

IL-8 RECEPTOR ANTAGONISTS

FIELD OF THE INVENTION

This invention relates to novel sulfonamide substituted diphenyl thiourea compounds, pharmaceutical compositions, processes for their preparation, and use thereof in treating IL-8, GRO α , GRO β , GRO γ , NAP-2, and ENA-78 mediated diseases.

BACKGROUND OF THE INVENTION

Many different names have been applied to Interleukin-8 (IL-8), such as neutrophil attractant/activation protein-1 (NAP-1), monocyte derived neutrophil chemotactic factor (MDNCF), neutrophil activating factor (NAF), and T-cell lymphocyte chemotactic factor. Interleukin-8 is a chemoattractant for neutrophils, basophils, and a subset of T-cells. It is produced by a majority of nucleated cells including macrophages, fibroblasts, endothelial and epithelial cells exposed to TNF, IL-1 α , IL-1 β or LPS, and by neutrophils themselves when exposed to LPS or chemotactic factors such as FMLP. M. Baggiolini et al., *J. Clin. Invest.* 84, 1045 (1989); J. Schroder et al, *J. Immunol.* 139, 3474 (1987) and *J. Immunol.* 144, 2223 (1990) ; Strieter, et al., *Science* 243, 1467 (1989) and *J. Biol. Chem.* 264, 10621 (1989); Cassatella et al., *J. Immunol.* 148, 3216 (1992).

GRO α , GRO β , GRO γ and NAP-2 also belong to the chemokine family. Like IL-8 these chemokines have also been referred to by different names. For instance GRO α , β , γ have been referred to as MGS α , β and γ respectively (Melanoma Growth Stimulating Activity), see Richmond et al., *J. Cell Physiology* 129, 375 (1986) and Chang et al., *J. Immunol* 148, 451 (1992). All of the chemokines of the α -family which possess the ELR motif directly preceding the CXC motif bind to the IL-8 B receptor (CXCR2).

IL-8, GRO α , GRO β , GRO γ , NAP-2, and ENA-78 stimulate a number of functions in vitro. They have all been shown to have chemoattractant properties for neutrophils, while IL-8 and GRO α have demonstrated T-lymphocytes, and basophilic chemotactic activity. In addition IL-8 can induce histamine release from basophils from both normal and atopic individuals. GRO- α and IL-8 can in addition, induce lysosomal enzyme release and respiratory burst from neutrophils. IL-8 has

also been shown to increase the surface expression of Mac-1 (CD11b/CD18) on neutrophils without de novo protein synthesis. This may contribute to increased adhesion of the neutrophils to vascular endothelial cells. Many known diseases are characterized by massive neutrophil infiltration. As IL-8, GRO α , GRO β , GRO γ and NAP-2 promote the accumulation and activation of neutrophils, these chemokines have been implicated in a wide range of acute and chronic inflammatory disorders including psoriasis and rheumatoid arthritis, Baggiolini et al., FEBS Lett. 307, 97 (1992); Miller et al., Crit. Rev. Immunol. 12, 17 (1992); Oppenheim et al., Annu. Rev. Immunol. 9, 617 (1991); Seitz et al., J. Clin. Invest. 87, 463 (1991); Miller et al., Am. Rev. Respir. Dis. 146, 427 (1992); Donnelly et al., Lancet 341, 643 (1993). In addition the ELR chemokines (those containing the amino acids ELR motif just prior to the CXC motif) have also been implicated in angiostasis, Strieter et al., Science 258, 1798 (1992).

In vitro, IL-8, GRO α , GRO β , GRO γ and NAP-2 induce neutrophil shape change, chemotaxis, granule release, and respiratory burst, by binding to and activating receptors of the seven-transmembrane, G-protein-linked family, in particular by binding to IL-8 receptors, most notably the IL-8 β receptor (CXCR2). Thomas et al., J. Biol. Chem. 266, 14839 (1991); and Holmes et al., Science 253, 1278 (1991). The development of non-peptide small molecule antagonists for members of this receptor family has precedent. For a review see R. Freidinger in: Progress in Drug Research, Vol. 40, pp. 33-98, Birkhauser Verlag, Basel 1993. Hence, the IL-8 receptor represents a promising target for the development of novel anti-inflammatory agents.

Two high affinity human IL-8 receptors (77% homology) have been characterized: IL-8R α , which binds only IL-8 with high affinity, and IL-8R β , which has high affinity for IL-8 as well as for GRO α , GRO β , GRO γ and NAP-2. See Holmes et al., supra; Murphy et al., Science 253, 1280 (1991); Lee et al., J. Biol. Chem. 267, 16283 (1992); LaRosa et al., J. Biol. Chem. 267, 25402 (1992); and Gayle et al., J. Biol. Chem. 268, 7283 (1993).

There remains a need for treatment, in this field, for compounds, which are capable of binding to the IL-8 α or β receptor. Therefore, conditions associated with an increase in IL-8 production (which is responsible for chemotaxis of neutrophil

and T-cells subsets into the inflammatory site) would benefit by compounds, which are inhibitors of IL-8 receptor binding.


SUMMARY OF THE INVENTION

This invention provides for a method of treating a chemokine mediated
 5 disease, wherein the chemokine is one which binds to an IL-8 a or b receptor and which method comprises administering an effective amount of a compound of Formula (I) or a pharmaceutically acceptable salt thereof. In particular the chemokine is IL-8.

This invention also relates to a method of inhibiting the binding of IL-8 to its
 10 receptors in a mammal in need thereof which comprises administering to said mammal an effective amount of a compound of Formula (I).

The present invention also provides for the novel compounds of Formula (I), and pharmaceutical compositions comprising a compound of Formula (I), and a pharmaceutical carrier or diluent.

15 Compounds of Formula (I) useful in the present invention are represented by the structure:



20 wherein:

R is selected from the group consisting of cyano, OR₁₁, C(O)NR₁₅R₁₆, R₁₈, C(O)OR₁₁, C(O)R₁₁, and S(O)₂R₁₇;

R_b is independently selected from the group consisting of hydrogen, NR₆R₇, OH, OR_a, C₁₋₅alkyl, aryl, arylC₁₋₄alkyl, aryl C₂₋₄alkenyl; cycloalkyl, cycloalkyl C₁₋₅ alkyl,
 25 heteroaryl, heteroarylC₁₋₄alkyl, heteroarylC₂₋₄ alkenyl, heterocyclic, heterocyclic C₁₋₄alkyl, and a heterocyclic C₂₋₄alkenyl moiety, all of which moieties are optionally substituted one to three times independently by a substituent selected

from the group consisting of halogen, nitro, halosubstituted C₁₋₄ alkyl, C₁₋₄ alkyl, amino, mono and di-C₁₋₄ alkyl substituted amine, OR_a,

C(O)R_a, NR_aC(O)OR_a, OC(O)NR₆R₇, hydroxy, NR₉C(O)R_a, S(O)_tR_a,

C(O)NR₆R₇, C(O)OH, C(O)OR_a, S(O)_tNR₆R₇, and NHS(O)_tR_a; or the two R_b

- 5 substituents join to form a 3-10 membered ring, and containing, in addition to optionally substituted C₁₋₉ alkyl, independently, 0 to 3 substituents selected from the group consisting of NR_a, C(O), O, S, SO, and SO₂ moieties which are unsaturated or saturated.

R_a is selected from the group consisting of alkyl, aryl, arylC₁₋₄alkyl, heteroaryl,

- 10 heteroaryl C₁₋₄alkyl, heterocyclic, COOR_a, and a heterocyclic C₁₋₄alkyl moiety, all of which moieties are optionally substituted;

m is an integer having a value of 1 to 3;

m' is 0, or an integer having a value of 1 or 2;

n is an integer having a value of 1 to 3;

- 15 q is 0, or an integer having a value of 1 to 10;

t is 0, or an integer having a value of 1 or 2;

s is an integer having a value of 1 to 3;

R₁ is independently selected from the group consisting of

hydrogen, halogen, nitro, cyano, C₁₋₁₀ alkyl, halosubstituted C₁₋₁₀ alkyl, C₂₋₁₀ alkenyl,

- 20 C₁₋₁₀ alkoxy, halosubstituted C₁₋₁₀alkoxy, azide, S(O)_tR₄, (CR₈R₈)_q S(O)_tR₄,

hydroxy, hydroxy substituted C₁₋₄alkyl, aryl, aryl C₁₋₄ alkyl, aryl C₂₋₁₀ alkenyl,

aryloxy, aryl C₁₋₄ alkyloxy, heteroaryl, heteroarylalkyl, heteroaryl C₂₋₁₀ alkenyl,

heteroaryl C₁₋₄ alkyloxy, heterocyclic, heterocyclic C₁₋₄alkyl,

heterocyclicC₁₋₄alkyloxy, heterocyclicC₂₋₁₀ alkenyl, (CR₈R₈)_q NR₄R₅,

- 25 (CR₈R₈)_qC(O)NR₄R₅, C₂₋₁₀ alkenyl C(O)NR₄R₅, (CR₈R₈)_q C(O)NR₄R₁₀,

S(O)₃R₈, (CR₈R₈)_q C(O)R₁₁, C₂₋₁₀ alkenyl C(O)R₁₁,

C₂₋₁₀ alkenyl C(O)OR₁₁, (CR₈R₈)_q C(O)OR₁₁, (CR₈R₈)_q OC(O)R₁₁,

(CR₈R₈)_qNR₄C(O)R₁₁, (CR₈R₈)_q C(NR₄)NR₄R₅, (CR₈R₈)_q NR₄C(NR₅)R₁₁,

(CR₈R₈)_q NHS(O)₂R₁₃, and (CR₈R₈)_q S(O)₂NR₄R₅, or two R₁ moieties together

- form O-(CH₂)_sO or a 5 to 6 membered saturated or unsaturated ring, wherein the alkyl, aryl, arylalkyl, heteroaryl, or heterocyclic moieties are optionally substituted; R₄ and R₅ are independently selected from the group consisting of hydrogen, optionally substituted C₁₋₄ alkyl, optionally substituted aryl, optionally substituted aryl C₁₋₄alkyl, optionally substituted heteroaryl, optionally substituted heteroaryl C₁₋₄alkyl, heterocyclic, and heterocyclicC₁₋₄ alkyl; or R₄ and R₅ together with the nitrogen to which they are attached form a 5 to 7 member ring which optionally comprises an additional heteroatom selected from the group consisting of O, N and S;
- R₆ and R₇ are independently selected from the group consisting of hydrogen, C₁₋₄ alkyl, heteroaryl, aryl, alkyl aryl, and alkyl C₁₋₄ heteroalkyl; or R₆ and R₇ together with the nitrogen to which they are attached form a 5 to 7 member ring which ring optionally contains an additional heteroatom selected from the group consisting of oxygen, nitrogen and sulfur, which ring is optionally substituted;
- Y is selected from the group consisting of hydrogen, halogen, nitro, cyano, halosubstituted C₁₋₁₀ alkyl, C₁₋₁₀ alkyl, C₂₋₁₀ alkenyl, C₁₋₁₀ alkoxy, halosubstituted C₁₋₁₀ alkoxy, azide, (CR₈R₈)_qS(O)_tR_a, (CR₈R₈)_qOR_a, hydroxy, hydroxy substituted C₁₋₄alkyl, aryl, aryl C₁₋₄ alkyl, aryloxy, arylC₁₋₄ alkyloxy, aryl C₂₋₁₀ alkenyl, heteroaryl, heteroarylalkyl, heteroaryl C₁₋₄ alkyloxy, heteroaryl C₂₋₁₀ alkenyl, heterocyclic, heterocyclic C₁₋₄alkyl, heterocyclicC₂₋₁₀ alkenyl, (CR₈R₈)_qNR₄R₅, C₂₋₁₀ alkenyl C(O)NR₄R₅, (CR₈R₈)_qC(O)NR₄R₅, (CR₈R₈)_qC(O)NR₄R₁₀, S(O)₃R₈, (CR₈R₈)_qC(O)R₁₁, C₂₋₁₀ alkenylC(O)R₁₁, (CR₈R₈)_qC(O)OR₁₁, C₂₋₁₀alkenylC(O)OR₁₁, (CR₈R₈)_qOC(O)R₁₁, (CR₈R₈)_qNR₄C(O)R₁₁, (CR₈R₈)_q NHS(O)_tR₁₃, (CR₈R₈)_q S(O)_tNR₄R₅, (CR₈R₈)_qC(NR₄)NR₄R₅, and (CR₈R₈)_q NR₄C(NR₅)R₁₁; or two Y moieties together form O-(CH₂)_s-O or a 5 to 6 membered saturated or unsaturated ring; wherein the alkyl, aryl, arylalkyl, heteroaryl, heteroaryl alkyl, heterocyclic, heterocyclicalkyl groups are optionally substituted;
- R₈ is hydrogen or C₁₋₄ alkyl;
- R₉ is hydrogen or a C₁₋₄ alkyl;
- R₁₀ is C₁₋₁₀ alkyl C(O)₂R₈;

- R₁₁ is selected from the group consisting of hydrogen, optionally substituted C₁₋₄ alkyl, optionally substituted aryl, optionally substituted aryl C₁₋₄alkyl, optionally substituted heteroaryl, optionally substituted heteroarylC₁₋₄alkyl, optionally substituted heterocyclic, and optionally substituted heterocyclicC₁₋₄alkyl; and
- 5 R₁₃ is selected from the group consisting of C₁₋₄ alkyl, aryl, aryl C₁₋₄alkyl, heteroaryl, heteroarylC₁₋₄alkyl, heterocyclic, and heterocyclicC₁₋₄alkyl; or a pharmaceutically acceptable salt thereof.

DETAILED DESCRIPTION OF THE INVENTION

The compounds of Formula (I), may also be used in association with the
 10 veterinary treatment of mammals, other than humans, in need of inhibition of IL-8 or other chemokines which bind to the IL-8 α and β receptors. Chemokine mediated diseases for treatment, therapeutically or prophylactically, in animals include disease states such as those noted herein in the Methods of Treatment section.

Suitably, R_b is independently hydrogen, NR₆R₇, OH, OR_a, C₁₋₄alkyl, aryl,
 15 arylC₁₋₄alkyl, aryl C₂₋₄alkenyl, heteroaryl, heteroarylC₁₋₄alkyl, heteroarylC₂₋₄alkenyl, heterocyclic, heterocyclic C₁₋₄alkyl, or a heterocyclic C₂₋₄alkenyl moiety, all of which moieties may be optionally substituted one to three times independently by halogen; nitro; halosubstituted C₁₋₄ alkyl; C₁₋₄ alkyl; amino, mono or di-C₁₋₄ alkyl substituted amine; cycloalkyl, cycloalkyl C₁₋₅ alkyl, OR_a; C(O)R_a;
 20 NR_aC(O)OR_a; OC(O)NR₆R₇; aryloxy; aryl C₁₋₄ oxy ; hydroxy; C₁₋₄ alkoxy; NR₉C(O)R_a; S(O)_mR_a; C(O)NR₆R₇; C(O)OH; C(O)OR_a; S(O)_tNR₆R₇; NHS(O)_tR_a; Alternatively, the two R_b substituents can join to form a 3-10 membered ring, optionally substituted and containing, in addition to carbon, independently, 1 to 3 NR₉, O, S, SO, or SO₂ moieties which can be optionally
 25 substituted.

Suitably, R_a is an alkyl, aryl, arylC₁₋₄alkyl, heteroaryl, heteroaryl C₁₋₄alkyl, heterocyclic, or a heterocyclic C₁₋₄alkyl moiety, all of which moieties may be optionally substituted.

Suitably R_1 is independently selected from hydrogen; halogen; nitro; cyano; halosubstituted C_{1-10} alkyl, such as CF_3 ; C_{1-10} alkyl, such as methyl, ethyl, isopropyl, or n-propyl; C_{2-10} alkenyl; C_{1-10} alkoxy, such as methoxy, or ethoxy; halosubstituted C_{1-10} alkoxy, such as trifluoromethoxy; azide; $(CR_8R_8)_q S(O)_t R_4$,
 5 wherein t is 0, 1 or 2; hydroxy; hydroxy C_{1-4} alkyl, such as methanol or ethanol; aryl, such as phenyl or naphthyl; aryl C_{1-4} alkyl, such as benzyl; aryloxy, such as phenoxy; aryl C_{1-4} alkyloxy, such as benzyloxy; heteroaryl; heteroarylalkyl; heteroaryl C_{1-4} alkyloxy; aryl C_{2-10} alkenyl; heteroaryl C_{2-10} alkenyl; heterocyclic C_{2-10} alkenyl; $(CR_8R_8)_q NR_4 R_5$; C_{2-10} alkenyl $C(O)NR_4 R_5$;
 10 $(CR_8R_8)_q C(O)NR_4 R_5$; $(CR_8R_8)_q C(O)NR_4 R_{10}$; $S(O)_3 H$; $S(O)_3 R_8$; $(CR_8R_8)_q C(O)R_{11}$; C_{2-10} alkenyl $C(O)R_{11}$; C_{2-10} alkenyl $C(O)OR_{11}$; $(CR_8R_8)_q C(O)R_{11}$; $(CR_8R_8)_q C(O)OR_{11}$; $(CR_8R_8)_q OC(O)R_{11}$; $(CR_8R_8)_q NR_4 C(O)R_{11}$; $(CR_8R_8)_q C(NR_4)NR_4 R_5$; $(CR_8R_8)_q NR_4 C(NR_5)R_{11}$; $(CR_8R_8)_q NHS(O)_t R_{13}$; $(CR_8R_8)_q S(O)_t NR_4 R_5$. All of the aryl, heteroaryl, and heterocyclic containing
 15 moieties may be optionally substituted as defined herein below.

For use herein the term "the aryl, heteroaryl, and heterocyclic containing moieties" refers to both the ring and the alkyl, or if included, the alkenyl rings, such as aryl, arylalkyl, and aryl alkenyl rings. The term "moieties" and "rings" may be interchangeably used throughout.

20 Suitably, R_4 and R_5 are independently hydrogen, optionally substituted C_{1-4} alkyl, optionally substituted aryl, optionally substituted aryl C_{1-4} alkyl, optionally substituted heteroaryl, optionally substituted heteroaryl C_{1-4} alkyl, heterocyclic, heterocyclic C_{1-4} alkyl, or R_4 and R_5 together with the nitrogen to which they are attached form a 5 to 7 member ring which may optionally comprise an additional
 25 heteroatom selected from O/N/S.

Suitably, R_8 is independently hydrogen or C_{1-4} alkyl.

Suitably, R_9 is hydrogen or a C_{1-4} alkyl;

Suitably, q is 0 or an integer having a value of 1 to 10.

Suitably, R_{10} is C_{1-10} alkyl $C(O)_2 R_8$, such as $CH_2 C(O)_2 H$ or
 30 $CH_2 C(O)_2 CH_3$.

Suitably, R_{11} is hydrogen, C_{1-4} alkyl, aryl, aryl C_{1-4} alkyl, heteroaryl, heteroaryl C_{1-4} alkyl, heterocyclic, or heterocyclic C_{1-4} alkyl.

Suitably, R_{12} is hydrogen, C_{1-10} alkyl, optionally substituted aryl or optionally substituted arylalkyl.

- 5 Suitably, R_{13} is C_{1-4} alkyl, aryl, arylalkyl, heteroaryl, heteroaryl C_{1-4} alkyl, heterocyclic, or heterocyclic C_{1-4} alkyl, wherein all of the aryl, heteroaryl and heterocyclic containing moieties may all be optionally substituted.

- Suitably, Y is independently selected from hydrogen; halogen; nitro; cyano; halosubstituted C_{1-10} alkyl; C_{1-10} alkyl; C_{2-10} alkenyl; C_{1-10} alkoxy; 10 halosubstituted C_{1-10} alkoxy; azide; $(CR_8R_8)_q S(O)_t R_a$; hydroxy; hydroxy C_{1-4} alkyl; aryl; aryl C_{1-4} alkyl; aryloxy; aryl C_{1-4} alkyloxy; heteroaryl; heteroarylalkyl; heteroaryl C_{1-4} alkyloxy; heterocyclic, heterocyclic C_{1-4} alkyl; aryl C_{2-10} alkenyl; heteroaryl C_{2-10} alkenyl; heterocyclic C_{2-10} alkenyl; $(CR_8R_8)_q$ NR_4R_5 ; C_{2-10} alkenyl $C(O)NR_4R_5$; $(CR_8R_8)_q C(O)NR_4R_5$; $(CR_8R_8)_q$ $C(O)NR_4R_{10}$; $S(O)_3H$; $S(O)_3R_8$; $(CR_8R_8)_q C(O)R_{11}$; C_{2-10} alkenyl $C(O)R_{11}$; C_{2-10} alkenyl $C(O)OR_{11}$; $(CR_8R_8)_q C(O)OR_{12}$; $(CR_8R_8)_q OC(O) R_{11}$; $(CR_8R_8)_q C(NR_4)NR_4R_5$; $(CR_8R_8)_q NR_4C(NR_5)R_{11}$; $(CR_8R_8)_q NR_4C(O)R_{11}$; $(CR_8R_8)_q NHS(O)_t R_{13}$; or $(CR_8R_8)_q S(O)_t NR_4R_5$; or two Y moieties together may form $O-(CH_2)_s-O$ or a 5 to 6 membered saturated or unsaturated ring. The aryl, 20 heteroaryl and heterocyclic containing moieties noted above may all be optionally substituted as defined herein.

Suitably s is an integer having a value of 1 to 3.

When Y forms a dioxybridge, s is preferably 1. When Y forms an additional unsaturated ring, it is preferably 6 membered resulting in a naphthylene ring system.

- 25 These ring systems may be substituted 1 to 3 times by other Y moieties as defined above.

Suitably, R_a is an alkyl, aryl C_{1-4} alkyl, heteroaryl, heteroaryl- C_{1-4} alkyl, heterocyclic, or a heterocyclic C_{1-4} alkyl, wherein all of these moieties may all be optionally substituted.

Y is preferably a halogen, C₁₋₄ alkoxy, optionally substituted aryl, optionally substituted aryloxy or arylalkoxy, methylene dioxy, NR₄R₅, thio C₁₋₄alkyl, thioaryl, halosubstituted alkoxy, optionally substituted C₁₋₄ alkyl, or hydroxy alkyl. Y is more preferably mono-substituted halogen, disubstituted
 5 halogen, mono-substituted alkoxy, disubstituted alkoxy, methylenedioxy, aryl, or alkyl, more preferably these groups are mono or di-substituted in the 2'- position or 2', 3'-position.

While Y may be substituted in any of the ring positions, n is preferably one. While both R₁ and Y can both be hydrogen, it is preferred that at least one of the
 10 rings is substituted, preferably both rings are substituted.

As used herein, "optionally substituted" unless specifically defined shall mean such groups as halogen, such as fluorine, chlorine, bromine or iodine; hydroxy; hydroxy substituted C₁₋₁₀alkyl; (C₈R₈)_qOR₄; C₁₋₁₀ alkoxy, such as methoxy or ethoxy; two substituents together may form O-(CH₂)_s-O; S(O)_{m'} C₁₋₁₀
 15 alkyl, wherein m' is 0, 1 or 2, such as methyl thio, methyl sulfinyl or methyl sulfonyl; amino, mono & di-substituted amino, such as in the NR₄R₅ group; NHC(O)R₄; C(O)NR₄R₅; C(O)OR₄; S(O)_tNR₄R₅; NHS(O)_tR₂₀, C₁₋₁₀ alkyl, such as methyl, ethyl, propyl, isopropyl, or t-butyl; halosubstituted C₁₋₁₀ alkyl, such as CF₃; an optionally substituted aryl, such as phenyl, or an optionally substituted
 20 arylalkyl, such as benzyl or phenethyl, optionally substituted heterocyclic, optionally substituted heterocyclicalkyl, optionally substituted heteroaryl, optionally substituted heteroaryl alkyl, wherein these aryl, heteroaryl, or heterocyclic moieties may be substituted one to two times by halogen; hydroxy; hydroxy substituted alkyl; C₁₋₁₀ alkoxy; S(O)_m C₁₋₁₀ alkyl; amino, mono & di-substituted alkyl amino, such
 25 as in the NR₄R₅ group; C₁₋₁₀ alkyl, or halosubstituted C₁₋₁₀ alkyl, such as CF₃.

R₂₀ is suitably C₁₋₄ alkyl, aryl, aryl C₁₋₄alkyl, heteroaryl, heteroarylC₁₋₄alkyl, heterocyclic, or heterocyclicC₁₋₄alkyl.

Suitable pharmaceutically acceptable salts are well known to those skilled in the art and include basic salts of inorganic and organic acids, such as hydrochloric
 30 acid, hydrobromic acid, sulphuric acid, phosphoric acid, methane sulphonic acid, ethane sulphonic acid, acetic acid, malic acid, tartaric acid, citric acid, lactic acid,

oxalic acid, succinic acid, fumaric acid, maleic acid, benzoic acid, salicylic acid, phenylacetic acid and mandelic acid. In addition, pharmaceutically acceptable salts of compounds of Formula (I) may also be formed with a pharmaceutically acceptable cation. Suitable pharmaceutically acceptable cations are well known to those skilled in the art and include alkaline, alkaline earth, ammonium and quaternary ammonium cations.

The following terms, as used herein, refer to:

- "halo" - all halogens, that is chloro, fluoro, bromo and iodo.
- "C₁₋₁₀alkyl" or "alkyl" - both straight and branched chain moieties of 1 to 10 carbon atoms, unless the chain length is otherwise limited, including, but not limited to, methyl, ethyl, *n*-propyl, *iso*-propyl, *n*-butyl, *sec*-butyl, *iso*-butyl, *tert*-butyl, *n*-pentyl and the like.
- "cycloalkyl" is used herein to mean cyclic moiety, preferably of 3 to 8 carbons, including but not limited to cyclopropyl, cyclopentyl, cyclohexyl, and the like.
- "alkenyl" is used herein at all occurrences to mean straight or branched chain moiety of 2-10 carbon atoms, unless the chain length is limited thereto, including, but not limited to ethenyl, 1-propenyl, 2-propenyl, 2-methyl-1-propenyl, 1-butenyl, 2-butenyl and the like.
- "aryl" - phenyl and naphthyl;
- "heteroaryl" (on its own or in any combination, such as "heteroaryloxy", or "heteroaryl alkyl") - a 5-10 membered aromatic ring system in which one or more rings contain one or more heteroatoms selected from the group consisting of N, O or S, such as, but not limited, to pyrrole, pyrazole, furan, thiophene, quinoline, isoquinoline, quinazolinyl, pyridine, pyrimidine, oxazole, tetrazole, thiazole, thiadiazole, triazole, imidazole, or benzimidazole.
- "heterocyclic" (on its own or in any combination, such as "heterocyclicalkyl") - a saturated or partially unsaturated 4-10 membered ring system in which one or more rings contain one or more heteroatoms selected from the group consisting of N, O, or S; such as, but not limited to, pyrrolidine, piperidine, piperazine, morpholine, tetrahydropyran, thiomorpholine, or

imidazolidine. Furthermore, sulfur may be optionally oxidized to the sulfone or the sulfoxide.

- "arylalkyl" or "heteroarylalkyl" or "heterocyclicalkyl" is used herein to mean C₁-10 alkyl, as defined above, attached to an aryl, heteroaryl or heterocyclic moiety, as also defined herein, unless otherwise indicated.
- "sulfinyl" - the oxide S (O) of the corresponding sulfide, the term "thio" refers to the sulfide, and the term "sulfonyl" refers to the fully oxidized S(O)₂ moiety.
- "wherein two R₁ moieties (or two Y moieties) may together form a 5 or 6 membered saturated or unsaturated ring" is used herein to mean the formation of an aromatic ring system, such as naphthalene, or is a phenyl moiety having attached a 6 membered partially saturated or unsaturated ring such as a C₆ cycloalkenyl, i.e. hexene, or a C₅ cycloalkenyl moiety, such as cyclopentene.

Illustrative compounds of Formula (I) include:

- 15 N-(2-bromophenyl)-N'-[4-chloro-2-hydroxy-3-(N'',N''dimethylaminosulfonyl)phenyl] cyanoguanidine;
- N-[4-chloro-2-hydroxy-3-(N'',N''-dimethylaminosulfonyl)phenyl]-N'-(2,3-dichlorophenyl) cyanoguanidine;
- N-(2-bromophenyl)-N'-[4-chloro-2-hydroxy-3-[S-(+)-(2-methoxymethyl)pyrrolidin-20 1-yl]aminosulfonylphenyl] cyanoguanidine;
- N-(2,3-dichlorophenyl)-N'-[4-chloro-2-hydroxy-3-[S-(+)-(2-methoxymethyl)pyrrolidin-1-yl]aminosulfonylphenyl] cyanoguanidine;
- N-phenyl-N'-[4-chloro-2-hydroxy-3-[S-(+)-(2-methoxymethyl)pyrrolidin-1-yl]aminosulfonylphenyl] cyanoguanidine;
- 25 N-(2-bromophenyl)-N'-[4-chloro-2-hydroxy-3-[R-(2-methoxymethyl)pyrrolidin-1-yl]aminosulfonylphenyl] cyanoguanidine;
- N-(2,3-dichlorophenyl)-N'-[4-chloro-2-hydroxy-3-[R-(2-methoxymethyl)pyrrolidin-1-yl]aminosulfonylphenyl] cyanoguanidine;
- N-(2-bromophenyl)-N'-[4-chloro-2-hydroxy-3-(N''-30 isoxazolidinylaminosulfonylphenyl] cyanoguanidine;

- N-(2,3-dichlorophenyl)-N'-[4-chloro-2-hydroxy-3-(N"-isoxazolidinylaminosulfonyl)phenyl] cyanoguanidine;
 N-(2-bromophenyl)-N'-[4-chloro-2-hydroxy-3-(N"-tetrahydroisoxazylaminosulfonyl)phenyl] cyanoguanidine;
 5 N-(2,3-dichlorophenyl)-N'-[4-chloro-2-hydroxy-3-(N"-tetrahydroisoxazylaminosulfonyl)phenyl] cyanoguanidine;
 N-(2,3-dichlorophenyl)-N'-[4-chloro-2-hydroxy-3-(4-thiomorpholinylaminosulfonyl)phenyl] cyanoguanidine;
 N-[4-chloro-2-hydroxy-3-[N",N"-dimethylaminosulfonyl]phenyl]-N'-(2-bromophenyl) propylguanidine;
 10 N-(2-bromophenyl)-N'-[4-chloro-2-hydroxy-3-(4-oxidithiomorpholino)amino sulfonylphenyl] cyanoguanidine;
 N-(2,3-chlorophenyl)-N'-[4-chloro-2-hydroxy-3-(4-oxidithiomorpholino)amino sulfonylphenyl] cyanoguanidine;
 15 N-(2-bromophenyl)-N'-[4-chloro-2-hydroxy-3-(N"-methylpiperazino)amino sulfonylphenyl] cyanoguanidine;
 N-(2,3-dichlorophenyl)-N'-[4-chloro-2-hydroxy-3-(N"-methylpiperazino)amino sulfonylphenyl] cyanoguanidine;
 N-(2-bromophenyl)-N'-[4-chloro-2-hydroxy-3-(N"-ethylmorpholino)amino
 20 sulfonylphenyl] cyanoguanidine;
 N-(2,3-dichlorophenyl)-N'-[4-chloro-2-hydroxy-3-(N"-ethylmorpholino)amino sulfonylphenyl] cyanoguanidine;
 N-(2-bromophenyl)-N'-{4-chloro-2-hydroxy-3-[N"-ethyl-2-(2-ethylpyrrolidino)]amino sulfonylphenyl} cyanoguanidine;
 25 N-(2,3-dichlorophenyl)-N'-{4-chloro-2-hydroxy-3-[N"-ethyl-2-(2-ethylpyrrolidino)]amino sulfonylphenyl} cyanoguanidine;
 N-(2-bromophenyl)-N'-{4-chloro-2-hydroxy-3-[S-(+)-(2-carboxy)pyrrolidin-1-yl]amino sulfonylphenyl} cyanoguanidine;
 N-(2,3-dichlorophenyl)-N'-{4-chloro-2-hydroxy-3-[S-(+)-(2-carboxy)pyrrolidin-1-yl]amino sulfonylphenyl} cyanoguanidine;
 30 N-(2-bromo-3-fluorophenyl)-N'-[4-chloro-2-hydroxy-3-[S-(+)-(2-methoxymethyl)pyrrolidin-1-yl]sulfonylphenyl] cyanoguanidine;

N-(2-phenoxyphenyl)-N'-[4-chloro-2-hydroxy-3-[S-(+)-(2-methoxymethyl)pyrrolidin-1-yl]sulfonylphenyl] cyanoguanidine; and

N-(2-benzoxypyphenyl)-N'-[4-chloro-2-hydroxy-3-[S-(+)-(2-methoxymethyl)pyrrolidin-1-yl]sulfonylphenyl] cyanoguanidine;


5 or a pharmaceutically acceptable salt thereof.

METHODS OF PREPARATION

The compounds of Formulas (I) may be obtained by applying synthetic procedures, some of which are illustrated in the Schemes below.

10

Scheme 1



Scheme 1

- a) i) NCS, AcOH, H₂O, ii) LiOH, MeOH b) H₂SO₄, HNO₃ c) KOH, MeOH d)
 PCl₅, POCl₃ e) NR'R''H, Et₃N f) NaH, H₂O g) Pd/C, H₂ h) RCNO, DMF i) TBSCl,
 5 Imidazole j) MsCl, Et₃N k) NH₂CN, Hunig's base l) CsF/TBAF, MeOH/THF

The desired 4-chloro-N-(3-sulfonamido-2-hydroxyphenyl)-N''-phenyl
 cyanoguanidine can be synthesized from the commercially available 2,6-dichloro
 thiophenol using procedure elaborated in scheme 1. The thiol can be oxidized to the
 10 sulfonyl halide using a halogenating agent such as NCS, NBS, chlorine or bromine
 in the presence of a protic solvent such as alcohol, acetic acid or water. The sulfonyl
 halide can be hydrolyzed by using a metal hydroxide such as lithium or potassium
 hydroxide to form the corresponding sulfonic acid salt.


The sulfonic acid salt can then be nitrated under nitration conditions such as
 15 nitric acid in a solvent of strong acid such as sulfuric acid to form the nitro phenyl
 sulfonic acid 3-scheme 1. The sulfonic acid 3-scheme 1 can be converted to the
 sulfonamide 5-scheme 1 using a three step procedure involving the formation of the
 metal salt using a base such as potassium hydroxide, sodium hydride or sodium
 carbonate to form 4-scheme 1. The sulfonic acid salt is then converted to the
 20 sulfonyl chloride using PCl₅ with POCl₃ as a solvent. The sulfonyl chloride can
 then be converted to the corresponding sulfonamide using the desired amine
 HNR'R'' in triethyl amine at temperatures ranging from -78 °C to 25 °C to form the
 corresponding sulfonamide 5-scheme 1.

The chlorine ortho to the nitro group can be selectively hydrolyzed using
 25 sodium hydride/ water in THF at room temperature to form the phenol. The nitro can
 be reduced by conditions well known in the art such as hydrogen and palladium on
 carbon to form the corresponding aniline 6-scheme 1. The aniline can then be
 coupled with a commercially available thioisocyanate to form the desired thiourea.

The phenol in thiourea can be protected with TBSCl to form the
 30 corresponding compounds 7-scheme 1. The protected thiourea can be converted to
 the corresponding carbodiimide 8-scheme 1 using methanesulfonyl chloride and
 triethyl amine at 0°C. The carbodiimide can then be converted to the corresponding

protected cyanoguanidine using cyanamide and Hunig's base, followed by desilylation with CsF or TBAF to form the desired the cyanoguanidine 9-scheme1.

Scheme 2



- 5 If the desired hydroxyaniline 6-Scheme 1 is not commercially available, it can be prepared as outlined in Scheme 2: Commercially available substituted 3-chloroanilines 1-scheme-2 can be converted to the amide 2-scheme-2 using standard conditions well known in the art such as pivavolyl chloride and triethylamine in a suitable organic solvent such as methylene chloride. The amide 2-scheme-2 can be
- 10 converted to the benzoxazole 3-scheme-2 using an excess amount of a strong base such as butyllithium in a suitable organic solvent such as THF under reduced reaction temperatures (-20 to -40°C) followed by quenching the reaction with sulfur dioxide gas and converting resulting sulfinic acid salt to the sulfonyl chloride 3-scheme-2 using standard conditions well known in the art such as sulfonyl chloride
- 15 in a suitable organic solvent such as methylene chloride. The sulfonyl chloride 3-scheme-2 can be transformed to the sulfonamide 4-scheme-2 using standard conditions well known in the art by reacting it with the amine HN(R_b)₂ in the presence of a suitable amine base such as triethylamine in a suitable organic solvent such as methylene chloride. The desired phenolaniline 5 can be obtained from the
- 20 benzoxazole 4-scheme-2 using standard hydrolysis conditions well known in the art such as sulfuric acid in water and heating at 85°C.

SYNTHETIC EXAMPLES

The invention will now be described by reference to the following examples, which are merely illustrative and are not to be construed as a limitation of the scope of the present invention. All temperatures are given in degrees centigrade, all solvents are highest available purity and all reactions run under anhydrous conditions in an argon atmosphere unless otherwise indicated.

In the Examples, all temperatures are in degrees Centigrade (°C). Mass spectra were performed upon a VG Zab mass spectrometer using fast atom bombardment, unless otherwise indicated. ¹H-NMR (hereinafter "NMR") spectra were recorded at 250 MHz using a Bruker AM 250 or Am 400 spectrometer. Multiplicities indicated are: s=singlet, d=doublet, t=triplet, q=quartet, m=multiplet and br indicates a broad signal. Sat. indicates a saturated solution, eq indicates the proportion of a molar equivalent of reagent relative to the principal reactant.

Example 1 & 2

Preparation of N-[4-chloro-2-hydroxy-3-(N'',N''-dimethylaminosulfonyl) phenyl]-N'-(2,3-dichlorophenyl) cyanoguanidine and N-(2-bromophenyl)-N'-[4-chloro-2-hydroxy-3-(N'',N''dimethylaminosulfonyl)phenyl] cyanoguanidine
2,6-Dichloro-3-nitrobenzenesulfonic acid

Lithium hydroxide hydrate (8.96 g, 0.214mol) was added to a solution of 2,6-dichlorobenzenesulfonyl chloride (35 g, 0.146mol) in MeOH (300mL) and the reaction was allowed to stir at room temperature for 16 hr. The reaction mixture was filtered to remove suspended solids and then concentrated. The resulting solid was dried *in vacuo* overnight to remove any residual MeOH. The solid was then dissolved in H₂SO₄ (300mL) and chilled in an ice bath. A solution of H₂SO₄ (35mL) and HNO₃ (70 %, 10.7 mL) was slowly added to the above reaction over 90 min. The reaction was allowed to warm up to room temperature overnight and then slowly poured into ice water (1200mL) and extracted with EtOAc. The combined organic layers were dried (MgSO₄) and concentrated to yield 2,6-dichloro-3-nitrobenzenesulfonic acid (37.38 g, 96%) as the dihydrate. EI-MS (m/z) 270 (M).

2,6-Dichloro-3-nitrobenzenesulfonyl chloride

Potassium hydroxide (11.54 g, 0.206mol) was added to a solution of 2,6-dichloro-3-nitrobenzenesulfonic acid dihydrate (37.38 g, 0.137mol) in MeOH (500mL) and the reaction was allowed to stir at room temperature for 14 hr. The reaction mixture was concentrated and the resulting solid was dried *in vacuo* overnight. To this was added PCl₅ (29.60 g, 0.142mol) followed by POCl₃ (400mL) and the mixture was refluxed overnight. The reaction was then cooled to room temperature and concentrated. The resulting mixture was taken up in EtOAc and chilled in an ice bath. Ice chunks were slowly added to the reaction mixture to quench any leftover PCl₅. When bubbling ceased, water was added and the reaction mix was extracted with EtOAc. The organic layer was dried (MgSO₄) and concentrated to yield 2,6-Dichloro-3-nitrobenzenesulfonyl chloride (31.4 g, 79%).
¹H NMR (DMSO-*d*₆) δ 7.88 (d, 1H), 7.75 (d, 1H).

The following is the general procedure for sulfonamide formation:

N,N-Dimethyl-2,6-dichloro-3-nitrobenzenesulfonamide

Into a solution of 2,6-dichloro-3-nitrobenzenesulfonyl chloride (200 mg, 0.69 mmol) in 15 mL of dichloromethane at -78°C was added dropwise a solution of dimethylamine (2.0 M in MeOH, 0.345mL, 0.69 mmol) and triethylamine (0.14 mL, 1.04 mmol) in 10 mL of dichloromethane. The mixture was warmed to room temperature and stirred for 16 hours. The mixture was acidified to pH >1 with 1N aq. HCl, then extracted with ethyl acetate. The combined organic layer was then concentrated to give the crude material. Column chromatography on silica gel, eluting with ethyl acetate/hexane (30/70, v/v/), gave the desired (240mg, 70 %). EI-MS (m/z) 298 (M⁺).

The following is the general procedure for the hydrolysis of dichlorosulfonamide to phenol:

N, N-Dimethyl-6-chloro-2-hydroxy-3-nitrobenzenesulfonamide

A mixture of N, N-dimethyl-2,6-dichloro-3-nitrobenzenesulfonamide (2.64 g, 8.83 mmol), 60% sodium hydride (1.06 g, 26.5 mmol) and water (191 mg, 10.6 mmol) was heated to 35 °C while kept under argon atmosphere for 16 hours. The solvent was evaporated. when The reaction was almost complete as indicated by ¹H NMR. The residue was diluted with ethyl acetate and washed with 1N aq. HCl. The solvent was concentrated to give the crude material. Column chromatography on silica gel, eluting with ethyl acetate/hexane/acetic acid (40/58/2, v/v/v), gave the desired product (2.3 g, 93%). EI-MS (m/z) 279.5 (M).

10 The following is the general procedure for the hydrogenation of nitro compound to aniline:

N, N-Dimethyl-3-amino-6-chloro-2-hydroxybenzenesulfonamide

To a solution of N, N-dimethyl-6-chloro-2-hydroxy-3-nitrobenzenesulfonamide (2.3 g, 8.2 mmol) in ethyl acetate, was added 10% Pd/C (2.0 g). The mixture was flushed with argon, and then stirred under a hydrogen atmosphere at balloon pressure for 3 hours at room temperature. The mixture was filtered through celite and the celite was washed with methanol. The solvent was evaporated to give the desired product (2.0 g, 97%). EI-MS (m/z) 249.5 (M).

20 Preparation of N-(3,4-dichlorophenyl)2,2-dimethyl-propionamide.

3,4-dichloroaniline (150g) in TBME (1L) was cooled to 10-15 °C. 30% aq NaOH (141 g, 1.14 equiv) was added, and the solution stirred vigorously via overhead mechanical stirrer. Trimethylacetyl chloride ("PivCl", 126mL) was added at such a rate as to keep the internal temperature below 30 °C. During this addition, the solution mixture becomes thick with white solid product. When the addition was complete (10-15 min), the mixture was heated to 30-35 °C for 1 hr, and then allowed to cool. The reaction mixture was held at -5 °C (overnight), and then filtered, rinsing first with 90:10 water/MeOH (600mL) and then water (900mL). Drying under vacuum yielded 195g (86%) product, as off-white crystals. LCMS m/z 246(M-H)⁺.

Preparation of 2-tert-Butyl-6-chloro-benzoxazole-7-sulfonyl chloride

The solution of N-(3,4-dichloro-phenyl)-2,2-dimethyl-propionamide (10g, 41mmol) in dry THF (100mL) was cooled to -72 °C under argon. n-Butyl lithium (1.6M in hexane, 64mL, 102mmol) was added dropwise. The solution warmed to ca. -50 °C over 45 minutes, and then was kept in the -25 - -10 °C range for 2 hrs.

- 5 The solution was then re-cooled to -78 °C, and sulfur dioxide was bubbled through the solution for 30min. The solution was then allowed to warm to room temperature for 2h, and a Ar stream was bubbled through the solution, with a gas outlet provided so that any excess sulfur dioxide could escape during the warming. The THF solution was cooled in an ice bath, and sulfuryl chloride (3.58mL, 44.9mmol) was
10 added dropwise. After a few minutes, the solution was warmed to room temperature for overnight. The mixture was concentrated, diluted with ethyl acetate and washed with water. Decolorizing carbon was added and the mixture was filtered. The resulting solution was dried (sodium sulfate), filtered and concentrated to afford the title compound (12.4g, 98%). ¹H NMR (CDCl₃) • 7.92 (d, 1H, J=8.5 Hz), 7.57 (d,
15 1H, J=8.4 Hz), 1.57 (s, 9H).

General procedure for the synthesis of 7-sulfoamidebenzoxazoles

2-tert-Butyl-6-chloro-7-(4-methyl-piperazine-1-sulfonyl)-benzooxazole

- To a solution of 2-tert-butyl-6-chloro-benzooxazole-7-sulfonyl chloride
20 (2.69g, 8.73mmol) and triethylamine (2.44mL, 17.5mmol) in THF (60mL) at 0 °C was added 1-methylpiperazine (0.98mL, 8.83mmol). The reaction was warmed to room temperature and allowed to stir overnight. The solution was concentrated and then diluted with water and extracted with ethyl acetate (3 times). The combined organic layers were dried with MgSO₄, filtered, and concentrated. Flash
25 chromatography (80% ethyl acetate/ 20% Ethanol) on silica gel gave the title compound (2.45g, 76%). EI-MS m/z 372(M+H)⁺.

General procedure for the hydrolysis of the benzooxazole to the desired aniline.

6-Amino-3-chloro-2-(4-methyl-piperazine-1-sulfonyl)-phenol

- 30 To a solution of 2-tert-Butyl-6-chloro-7-(4-methyl-piperazine-1-sulfonyl)-benzooxazole (2.44g, 6.56mmol) in 1,4-dioxane (20mL) was treated with water

(4mL) and conc. H₂SO₄ (4mL). The mixture was heated to 85 °C for 14h. The reaction was cooled to room temperature, and then basified to pH=14 with 25% aq NaOH. washed. The mixture was extracted with ethyl acetate (3 times), dried with MgSO₄, filtered, and concentrated to afford the title compound (1.35g, 68%). EI-MS

5 m/z 306(M+H)⁺.

The following is the general procedure for thiourea formation:

N-[4-Chloro-2-hydroxy-3-[N",N"-dimethylaminosulfonyl]phenyl]-N'-(2,3-dichlorophenyl) thiourea

10 A solution of N,N-dimethyl-3-amino-6-chloro-2-hydroxybenzenesulfonamide (356 mg, 1.42 mmol) and 2,3-dichlorophenylisothiocyanate (318 mg, 1.57 mmol) in 1.0 mL of N,N-dimethylformamide was stirred at room temperature for 3 hours. Purification by column chromatography on silica gel, eluting with ethyl acetate/hexane (30/70, v/v) to give the desired product (440 mg, 68 %). EI-MS (m/z) 455.5 (M⁺).

N-[4-chloro-2-hydroxy-3-[N",N"-dimethylaminosulfonyl]phenyl]-N'-(2-bromophenyl) thiourea

20 A solution of N,N-dimethyl-3-amino-6-chloro-2-hydroxybenzenesulfonamide (50 mg, 0.2 mmol) and 2-bromophenylisothiocyanate (43 mg, 0.2 mmol) in 0.5 mL of N,N-dimethylformamide was stirred at room temperature for 3 hours. Purification by column chromatography on silica gel, eluting with ethyl acetate/hexane (30/70, v/v) to give the desired product (66 mg, 74 %). EI-MS (m/z) 465.5 (M⁺).

25

The following is the general procedure for protected phenyl thiourea formation:

N-[4-Chloro-2-*tert*-butyldimethylsilyloxy-3-(N",N"-dimethylaminosulfonyl)phenyl]-N'-(2-bromophenyl) thiourea

30 To a solution of N-[4-chloro-2-hydroxy-3-(N",N"-dimethylaminosulfonyl)phenyl]-N'-(2-bromophenyl) thiourea (500 mg, 1.07 mmol) in THF (20 mL), *tert*-butyldimethylsilyl chloride (810 mg, 5.35 mmol) and imidazole (144 mg, 2.14 mmol) were added. The reaction mixture was stirred at

room temperature for 16 hours. Then it was partitioned between ethyl acetate and water. The combined organic phase was dried and concentrated. Chromatography of the residue on silica gel (30%Ethyl acetate/Hexane) gave desired product (240 mg, 40%) and recovered starting material (280 mg). EI-MS m/z 580 (M^+).

5

N-[4-chloro-2-*tert*-butyldimethylsilyloxy-3-(N",N"-dimethylaminosulfonyl)phenyl]-N'-(2,3-dichlorophenyl) thiourea

To a solution of N-[4-chloro-2-hydroxy-3-(N",N"-dimethylaminosulfonyl)phenyl]-N'-(2,3-dichlorophenyl) thiourea (440 mg, 1.07 mmol) in THF (20 mL), *tert*-butyldimethylsilyl chloride (730 mg, 4.85 mmol) and imidazole (132 mg, 1.94 mmol) were added. The reaction mixture was stirred at room temperature for 16 hours. Then it was partitioned between ethyl acetate and water. The combined organic phase was dried and concentrated. Chromatography of the residue on silica gel (30%Ethyl acetate/Hexane) gave desired product (270 mg, 50%). EI-MS m/z 570 (M^+).

15

The following is the general procedure for carbodiimide formation:

N-[4-chloro-2-*tert*-butyldimethylsilyloxy-3-(N",N"-dimethylaminosulfonyl)phenyl]-N'-(2-bromophenyl)carbodiimide

To a solution of N-[4-chloro-2-*tert*-butyldimethylsilyloxy-3-(N",N"-dimethylaminosulfonyl)phenyl]-N'-(2-bromophenyl) thiourea (240 mg, 0.42 mmol) in dichloromethane (10 mL) at 0°C, methanesulfonyl chloride (0.065 mL, 0.84 mmol) and triethylamine (0.12 mL, 0.84 mmol) were added. 4-Dimethylaminopyridine was added as a catalyst. The reaction mixture was stirred at 0°C for 1 hour, then it was partitioned between dichloromethane and water. The combined organic phase was dried and concentrated to give the desired product (266.2 mg, crude). ^1H NMR (CDCl_3) δ 0.4 (s, 6H), 1.09 (s, 9H), 2.86 (s, 6H), 7.06 (d, 2H), 7.1-7.37 (m, 4H).

25

N-[4-chloro-2-*tert*-butyldimethylsilyloxy-3-(N",N"-dimethylaminosulfonyl)phenyl]-N'-(2,3-dichlorophenyl)carbodiimide

30

To a solution of N-[4-chloro-2-*tert*-butyldimethylsilyloxy-3-(N'',N''-dimethylaminosulfonyl)phenyl]-N'-(2,3-dichlorophenyl) thiourea (270 mg, 0.45 mmol) in dichloromethane (10 mL) at 0°C, methanesulfonyl chloride (0.07 mL, 0.9 mmol) and triethylamine (0.13 mL, 0.9 mmol) were added. 4-

- 5 Dimethylaminopyridine was added as a catalyst. The reaction mixture was stirred at 0°C for 1 hour, then it was partitioned between dichloromethane and water. The combined organic phase was dried and concentrated to give the desired product (270 mg, crude). ¹ H NMR (CDCl₃) δ 0.47 (s, 6H), 1.05 (s, 9H), 2.9 (s, 6H), 7.08 (d, 1H), 7.12 (d, 1H), 7.2 (t, 1H), 7.31 (m, 2H).

10

The following is the general procedure for cyanoguanidine formation:

N-[4-chloro-2-hydroxy-3-(N'',N''-dimethylaminosulfonyl)phenyl]-N'-(2-bromophenyl) cyanoguanidine

- To a solution of N-[4-chloro-2-*tert*-butyldimethylsilyloxy-3-(N'',N''-dimethylaminosulfonyl)phenyl]-N'-(2-bromophenyl)carbodiimide (266 mg, 0.49 mmol) in acetonitrile (5 mL) at room temperature, cyanamide (83 mg, 1.96 mmol) and N,N-diisopropylethylamine (76 mg, 0.59 mmol) was added. The reaction mixture was stirred at room temperature for 1 hour. The reaction mixture was concentrated under reduced pressure. The residue was diluted with a mixture of THF (3mL) and methanol (1 mL). Cesium fluoride (90 mg, 0.59 mmol) was added at 0°C. 20 The reaction mixture was stirred at 0°C for 3 hours. The reaction mixture was concentrated under reduced pressure. The residue was purified by Gilson HPLC to give the desired product (70 mg, 30 %). EI-MS m/z 473.75(M⁺).

- 25 N-[4-chloro-2-hydroxy-3-(N'',N''-dimethylaminosulfonyl)phenyl]-N'-(2,3-dichlorophenyl) cyanoguanidine

- To a solution of N-[4-chloro-2-*tert*-butyldimethylsilyloxy-3-(N'',N''-dimethylaminosulfonyl)phenyl]-N'-(2,3-dichlorophenyl)carbodiimide (270 mg, 0.47 mmol) in acetonitrile (5 mL) at room temperature, cyanamide (79 mg, 1.88 mmol) and N,N-diisopropylethylamine (76 mg, 0.56 mmol) was added. The reaction mixture was stirred at room temperature for 1 hour. The reaction mixture was concentrated under reduced pressure. The residue was diluted with a mixture of 30

tetrahydrofuran (3mL) and methanol (1 mL). Cesium fluoride (86 mg, 0.56 mmol) was added at 0°C. The reaction mixture was stirred at 0°C for 3 hours. The reaction mixture was concentrated under reduced pressure. The residue was purified by Gilson HPLC to give the desired product (98 mg, 48 %). EI-MS m/z 473.75(M⁺).

5

Example 3, 4 & 5

Preparation of N-(2-bromophenyl)-N'-[4-chloro-2-hydroxy-3-[S-(+)-(2-methoxymethyl)pyrrolidin-1-yl]aminosulfonylphenyl] cyanoguanidine, N-(2,3-dichlorophenyl)-N'-[4-chloro-2-hydroxy-3-[S-(+)-(2-methoxymethyl)pyrrolidin-1-yl]aminosulfonylphenyl] cyanoguanidine and N-(2-bromophenyl)-N'-[4-chloro-2-hydroxy-3-[S-(+)-(2-methoxymethyl)pyrrolidin-1-yl]aminosulfonylphenyl] cyanoguanidine

10

S-(+)-N-(2-Methoxymethyl)pyrrolidin-1-yl -2,6-dichloro-3-nitrobenzenesulfonamide

15

Following the general procedure for sulfonamide formation outlined in example 1, 6-dichloro-3-nitrobenzenesulfonyl chloride (1.0 g, 3.44 mmol), (S)-(+)-2(methoxymethyl)pyrrolidine (0.476 mL, 4.13 mmol) and triethylamine (0.72 mL, 5.16 mmol) were reacted to form the desired product (1.15 g, 91 %). EI-MS m/z 368 (M).

20

S-(+)-N-(2-Methoxymethyl)pyrrolidin-1-yl-6-chloro-2-hydroxy-3-nitrobenzenesulfonamide

25

Following the general hydrolysis procedure outlined in example 1, S-(+)-N-(2-methoxymethyl)pyrrolidin-1-yl-2,6-dichloro-3-nitrobenzenesulfonamide (1.15 g, 3.12 mmol), 60% sodium hydride (374 mg, 9.36 mmol) and water (73 mg, 4.02 mmol) were reacted to form the desired product (1.0 g, 91 %). EI-MS m/z 349.1 (M).

30

S-(+)-N-(2-Methoxymethyl)pyrrolidin-1-yl-3-amino-6-chloro-2-hydroxybenzenesulfonamide

Following the general hydrogenation procedure outlined in example 1, S-(+)-N-(2-methoxymethyl)pyrrolidin-1-yl-6-chloro-2-hydroxy-3-

nitrobenzenesulfonamide (1.0 g, 2.86 mmol) was reduced with hydrogen and Pd/C (600 mg) to form the desired product (0.9 g, 98 %). EI-MS m/z 319.1 (M⁺).

5 N-(2-Bromophenyl)-N'-[4-chloro-2-hydroxy-3-[S-(+)-(2-methoxymethyl)pyrrolidin-1-yl]aminosulfonylphenyl] thiourea

Following the general procedure for thiourea formation outlined in example 1, S-(+)-N-(2-methoxymethyl)pyrrolidin-1-yl-3-amino-6-chloro-2-hydroxybenzenesulfonamide (260 mg, 0.85 mmol) and 2-bromophenylisothiocyanate (182 mg, 0.85 mmol) were coupled to form the desired thiourea (231 mg, 53%). EI-MS m/z 535.1 (M⁺).

N-(2,3-Dichlorophenyl)-N'-[4-chloro-2-hydroxy-3-[S-(+)-(2-methoxymethyl)pyrrolidin-1-yl]aminosulfonylphenyl] thiourea

15 Following the general procedure for thiourea formation outlined in example 1, S-(+)-N-(2-methoxymethyl)pyrrolidin-1-yl-3-amino-6-chloro-2-hydroxybenzenesulfonamide (407 mg, 1.27 mmol) and 2,3-dichlorophenylisothiocyanate (285 mg, 1.4 mmol) were coupled to form the desired thiourea (370 mg, 54%). EI-MS m/z 525.1 (M⁺).

20 N-Phenyl-N'-[4-chloro-2-hydroxy-3-[S-(+)-(2-methoxymethyl)pyrrolidin-1-yl]aminosulfonylphenyl] thiourea

Following the general procedure for thiourea formation outlined in example 1, S-(+)-N-(2-methoxymethyl)pyrrolidin-1-yl-3-amino-6-chloro-2-hydroxybenzenesulfonamide (310 mg, 0.97 mmol) and phenylisothiocyanate (144 mg, 1.07 mmol) were coupled to form the desired thiourea (210 mg, 54%). EI-MS m/z 456 (M⁺).

N-(2-Bromophenyl)-N'-[4-chloro-2-tert-butyl dimethylsilyloxy-3-[S-(+)-(2-methoxymethyl)pyrrolidin-1-yl]aminosulfonylphenyl] thiourea

30 Following the general procedure for protected phenyl thiourea formation outlined in example 1, N-(2-bromophenyl)-N'-[4-chloro-2-hydroxy-3-[S-(+)-(2-methoxymethyl)pyrrolidin-1-yl]aminosulfonylphenyl] thiourea (230 mg, 0.44

mmol), *tert*-butyldimethylsilyl chloride (332 mg, 2.2 mmol) and imidazole (60 mg, 0.88 mmol) were reacted to form the desired product (136 mg, 48%). EI-MS *m/z* 650 (M^+).

5 N-(2,3-Dichlorophenyl)-N'-[4-chloro-2-*tert*-butyldimethylsilyloxy-3-[S-(+)-(2-methoxymethyl)pyrrolidin-1-yl]aminosulfonylphenyl] thiourea

Following the general procedure for protected phenyl thiourea formation outlined in example 1, N-(2,3-dichlorophenyl)-N'-[4-chloro-2-hydroxy-3-[S-(+)-(2-methoxymethyl)pyrrolidin-1-yl]aminosulfonylphenyl] thiourea (370 mg, 0.71
10 mmol), *tert*-butyldimethylsilyl chloride (370 mg, 0.71 mmol) and imidazole (97 mg, 1.42 mmol) were reacted to form the desired product (187 mg, 41%). EI-MS *m/z* 640.2 (M^+).

15 N-Phenyl-N'-[4-chloro-2-*tert*-butyldimethylsilyloxy-3-[S-(+)-(2-methoxymethyl)pyrrolidin-1-yl]aminosulfonylphenyl] thiourea

Following the general procedure for protected phenyl thiourea formation outlined in example 1, N-phenyl-N'-[4-chloro-2-hydroxy-3-[S-(+)-(2-methoxymethyl)pyrrolidin-1-yl]aminosulfonylphenyl] thiourea (210 mg, 0.46
20 mmol), *tert*-butyldimethylsilyl chloride (349 mg, 2.3 mmol) and imidazole (63 mg, 0.92 mmol) were reacted to form the desired product (125 mg, 41%). EI-MS *m/z* 571 (M^+).

25 N-(2-Bromophenyl)-N'-[4-chloro-2-*tert*-butyldimethylsilyloxy-3-[S-(+)-(2-methoxymethyl)pyrrolidin-1-yl]aminosulfonylphenyl] carbodiimide

Following the general procedure for carbodiimide formation outlined in example 1, N-(2-bromophenyl)-N'-[4-chloro-2-*tert*-butyldimethylsilyloxy-3-[S-(+)-(2-methoxymethyl)pyrrolidin-1-yl]aminosulfonylphenyl] thiourea (136 mg, 0.21
30 mmol), methanesulfonyl chloride (0.03 mL, 0.42 mmol) and triethylamine (0.06 mL, 0.42 mmol) were reacted to form the desired product (120 mg, 93%). ^1H NMR (CDCl_3) δ 0.37 (s, 6H), 1.01 (s, 9H), 1.72 (m, 4H), 3.25 (s, 3H), 3.3 (m, 2H), 3.45 (m, 2H), 4.15 (m, 1H), 7.10 (d, 2H), 7.2 (d, 1H), 7.3 (m, 1H), 7.39 (d, 1H), 7.6 (d, 1H).

N-(2,3-Dichlorophenyl)-N'-[4-chloro-2-*tert*-butyldimethylsilyloxy-3-[S-(+)-(2-methoxymethyl)pyrrolidin-1-yl]aminosulfonylphenyl] carbodiimide

Following the general procedure for carbodiimide formation outlined in example 1, N-(2,3-dichlorophenyl)-N'-[4-chloro-2-*tert*-butyldimethylsilyloxy-3-[S-(+)-(2-methoxymethyl)pyrrolidin-1-yl]aminosulfonylphenyl] thiourea (187 mg, 0.3 mmol), methanesulfonyl chloride (0.05 mL, 0.6 mmol) and triethylamine (0.08 mL, 0.6 mmol) were reacted to form the desired product (160 mg, 90%). ¹H NMR (CDCl₃) δ 0.32 (s, 6H), 1.01 (s, 9H), 1.35 (m, 4H), 3.24 (s, 3H), 3.42 (m, 4H), 4.14 (m, 1H), 7.10 (d, 1H), 7.12 (d, 1H), 7.2 (t, 1H), 7.29 (d, 1H), 7.32 (d, 1H).

N-Phenyl-N'-[4-chloro-2-*tert*-butyldimethylsilyloxy-3-[S-(+)-(2-methoxymethyl)pyrrolidin-1-yl]aminosulfonylphenyl] carbodiimide

Following the general procedure for carbodiimide formation outlined in example 1, N-phenyl-N'-[4-chloro-2-*tert*-butyldimethylsilyloxy-3-[S-(+)-(2-methoxymethyl)pyrrolidin-1-yl]aminosulfonylphenyl] thiourea (125 mg, 0.22 mmol), methanesulfonyl chloride (0.04 mL, 0.44 mmol) and triethylamine (0.06 mL, 0.44 mmol) were reacted to form the desired product (125 mg, crude). ¹H NMR (CDCl₃) δ 0.37 (s, 6H), 1.04 (s, 9H), 1.35 (m, 4H), 3.24 (s, 3H), 3.42 (m, 4H), 4.14 (m, 1H), 7.06 (d, 1H), 7.16 (d, 1H), 7.19 (m, 2H), 7.25 (d, 1H), 7.35 (t, 2H).

N-(2-Bromophenyl)-N'-[4-chloro-2-hydroxy-3-[S-(+)-(2-methoxymethyl)pyrrolidin-1-yl]aminosulfonylphenyl] cyanoguanidine

Following the general procedure for cyanoguanidine formation outlined in example 1, N-(2-bromophenyl)-N'-[4-chloro-2-*tert*-butyldimethylsilyloxy-3-[S-(+)-(2-methoxymethyl)pyrrolidin-1-yl]aminosulfonylphenyl] carbodiimide (152 mg, 0.25 mmol), cyanamide (42 mg, 1.0 mmol) and N,N-diisopropylethylamine (39 mg, 0.3 mmol) were reacted, followed by desilylation with Cesium fluoride (45.6 mg, 0.3 mmol) to form the desired product (33 mg, 25%). EI-MS m/z 543.6 (M⁺).

N-(2,3-Dichlorophenyl)-N'-[4-chloro-2-hydroxy-3-[S-(+)-(2-methoxymethyl)pyrrolidin-1-yl]aminosulfonylphenyl] cyanoguanidine

Following the general procedure for carbodiimide formation outlined in example 1, N-(2,3-dichlorophenyl)-N'-[4-chloro-2-*tert*-butyldimethylsilyloxy-3-[S-(+)-(2-methoxymethyl)pyrrolidin-1-yl]sulfonylphenyl] carbodiimide (160 mg, 0.26 mmol), cyanamide (44 mg, 1.04 mmol) and N,N-diisopropylethylamine (41 mg, 0.31 mmol) were reacted, followed by desilylation with Cesium fluoride (48 mg, 0.31 mmol) to form the desired product (35 mg, 25%). EI-MS m/z 532 (M⁺).

N-Phenyl-N'-[4-chloro-2-hydroxy-3-[S-(+)-(2-methoxymethyl)pyrrolidin-1-yl]aminosulfonylphenyl] cyanoguanidine

Following the general procedure for carbodiimide formation outlined in example 1, N-phenyl-N'-[4-chloro-2-*tert*-butyldimethylsilyloxy-3-[S-(+)-(2-methoxymethyl)pyrrolidin-1-yl]sulfonylphenyl] carbodiimide (125 mg, 0.23 mmol), cyanamide (38.6 mg, 0.92 mmol) and N,N-diisopropylethylamine (36.2 mg, 0.28 mmol) were reacted, followed by desilylation with Cesium fluoride (42 mg, 0.28 mmol) to form the desired product (38 mg, 35%). LC-MS m/z 462.2.

Example 6 & 7

Preparation of N-(2-bromophenyl)-N'-[4-chloro-2-hydroxy-3-[R-(2-methoxymethyl)pyrrolidin-1-yl]aminosulfonylphenyl] cyanoguanidine and N-(2,3-dichlorophenyl)-N'-[4-chloro-2-hydroxy-3-[R-(2-methoxymethyl)pyrrolidin-1-yl]aminosulfonylphenyl] cyanoguanidine

R-N-(2-Methoxymethyl)pyrrolidin-1-yl-2,6-dichloro-3-nitrobenzenesulfonamide

Following the general procedure for sulfonamide formation outlined in example 1, 6-dichloro-3-nitrobenzenesulfonyl chloride (2.0 g, 6.89 mmol), (R)-2-(methoxymethyl)pyrrolidine (0.783 mL, 8.27 mmol) and triethylamine (1.44 mL, 10.34 mmol) were reacted to form the desired product (1.69 g, 66 %). EI-MS m/z 368 (M⁺).

R-N-(2-methoxymethyl)pyrrolidin-1-yl-6-chloro-2-hydroxy-3-nitrobenzenesulfonamide

Following the general hydrolysis procedure outlined in example (R)-N-(2-methoxymethyl)pyrrolidin-1-yl -2,6-dichloro-3-nitrobenzenesulfonamide (2.16 g,

5.85 mmol), 60% sodium hydride (702 mg, 17.55 mmol) and water (137 mg, 7.6 mmol) were reacted to form the desired product (2.0 g, 97 %). EI-MS m/z 349.1 (M⁺).

5 (R)-N-(2-Methoxymethyl)pyrrolidin-1-yl-3-amino-6-chloro-2-hydroxybenzenesulfonamide

Following the general hydrogenation procedure outlined in example 1, (R)-N-(2-methoxymethyl)pyrrolidin-1-yl-6-chloro-2-hydroxy-3-nitrobenzenesulfonamide (2.0 g, 4.44 mmol) was reduced with hydrogen and Pd/C (1.5 g) to form the desired product (1.75 g, 96 %). EI-MS m/z 319.1 (M⁺).

10

N-(2-bromophenyl)-N'-[4-chloro-2-hydroxy-3-[(R)-(2-methoxymethyl)pyrrolidin-1-yl]aminosulfonylphenyl] thiourea

Following the general procedure for thiourea formation outlined in example 1, (R)-N-(2-methoxymethyl)pyrrolidin-1-yl-3-amino-6-chloro-2-hydroxybenzenesulfonamide (390 mg, 1.21 mmol) and 2-bromophenylthioisocyanate (286 mg, 1.33 mmol) were coupled to form the desired thiourea (462 mg, 71%). EI-MS m/z 535.1 (M⁺).

20 N-(2,3-Dichlorophenyl)-N'-[4-chloro-2-hydroxy-3-[(R)-2-methoxymethyl]pyrrolidin-1-yl]aminosulfonylphenyl] thiourea

Following the general procedure for thiourea formation outlined in example 1, (R)-N-(2-methoxymethyl)pyrrolidin-1-yl-3-amino-6-chloro-2-hydroxybenzenesulfonamide (360 mg, 1.12 mmol) and 2,3-dichlorophenylisothiocyanate (251 mg, 1.23 mmol) were coupled to form the desired thiourea (410 mg, 70%). EI-MS m/z 525.1 (M⁺).

N-(2-Bromophenyl)-N'-[4-chloro-2-tert-butyl dimethylsilyloxy-3-[(R)-(2-methoxymethyl)pyrrolidin-1-yl]aminosulfonylphenyl] thiourea

Following the general procedure for protected phenyl thiourea formation outlined in example 1, N-(2-bromophenyl)-N'-[4-chloro-2-hydroxy-3-[(R)-(2-methoxymethyl)pyrrolidin-1-yl]aminosulfonylphenyl] thiourea (462 mg, 0.86 mmol), tert-butyl dimethylsilyl chloride (651 mg, 4.3 mmol) and imidazole (118

mg, 1.73 mmol) were reacted to form the desired product (216 mg, 39%). EI-MS m/z 650 (M^+).

N-(2,3-Dichlorophenyl)-N'-[4-chloro-2-*tert*-butyldimethylsilyloxy-3-[(R)-(2-methoxymethyl)pyrrolidin-1-yl]aminosulfonylphenyl] thiourea

Following the general procedure for protected phenyl thiourea formation outlined in example 1, N-(2,3-dichlorophenyl)-N'-[4-chloro-2-hydroxy-3-[(R)-(2-methoxymethyl)pyrrolidin-1-yl]aminosulfonylphenyl] thiourea (410 mg, 0.78 mmol), *tert*-butyldimethylsilyl chloride (589 mg, 3.9 mmol) and imidazole (106 mg, 1.56 mmol) were reacted to form the desired product (202 mg, 40%). EI-MS m/z 640.2 (M^+).

N-(2-Bromophenyl)-N'-[4-chloro-2-*tert*-butyldimethylsilyloxy-3-[(R)-(2-methoxymethyl)pyrrolidin-1-yl]aminosulfonylphenyl] carbodiimide

Following the general procedure for carbodiimide formation outlined in example 1, N-(2-bromophenyl)-N'-[4-chloro-2-*tert*-butyldimethylsilyloxy-3-[(R)-(2-methoxymethyl)pyrrolidin-1-yl]aminosulfonylphenyl] thiourea (216 mg, 0.33 mmol), methanesulfonyl chloride (0.05 mL, 0.66 mmol) and triethylamine (0.09 mL, 0.66 mmol) were reacted to form the desired product (200 mg, 97%). 1H NMR ($CDCl_3$) δ 0.37 (s, 6H), 1.02 (s, 9H), 1.9 (m, 4H), 3.23 (m, 4H), 3.42 (m, 3H), 4.13 (m, 1H), 7.08 (d, 2H), 7.20 (d, 1H), 7.3 (t, 1H), 7.38 (d, 1H), 7.6 (d, 1H).

N-(2,3-Dichlorophenyl)-N'-[4-chloro-2-*tert*-butyldimethylsilyloxy-3-[(R)-(2-methoxymethyl)pyrrolidin-1-yl]aminosulfonylphenyl] carbodiimide

Following the general procedure for carbodiimide formation outlined in example 1, N-(2,3-dichlorophenyl)-N'-[4-chloro-2-*tert*-butyldimethylsilyloxy-3-[(R)-(2-methoxymethyl)pyrrolidin-1-yl]aminosulfonylphenyl] thiourea (202 mg, 0.32 mmol), methanesulfonyl chloride (0.05 mL, 0.64 mmol) and triethylamine (0.09 mL, 0.64 mmol) were reacted to form the desired product (190 mg, 99%). 1H NMR ($CDCl_3$) δ 0.34 (s, 6H), 1.04 (s, 9H), 1.4 (m, 2H), 1.8 (m, 2H), 3.15 (s, 3H), 3.25 (m, 2H), 3.43 (m, 2H), 4.13 (m, 1H), 7.09 (d, 2H), 7.12 (d, 1H), 7.19 (t, 1H), 7.3 (d, 1H), 7.34 (d, 1H).

N-(2-Bromophenyl)-N'-[4-chloro-2-hydroxy-3-[(R)-(2-methoxymethyl)pyrrolidin-1-yl]aminosulfonylphenyl] cyanoguanidine

Following the general procedure for cyanoguanidine formation outlined in
 5 example 1, N-(2-bromophenyl)-N'-[4-chloro-2-*tert*-butyldimethylsilyloxy-3-[(R)-(2-methoxymethyl)pyrrolidin-1-yl]aminosulfonylphenyl] carbodiimide (240 mg, 0.39 mmol), cyanamide (66 mg, 1.56 mmol) and N,N-diisopropylethylamine (63 mg, 0.47 mmol) were reacted, followed by desilylation with Cesium fluoride (72 mg, 0.47 mmol) to form the desired product (75 mg, 35%). EI-MS m/z 543.6 (M⁺).

10

N-(2,3-Dichlorophenyl)-N'-[4-chloro-2-hydroxy-3-[(R)-(2-methoxymethyl)pyrrolidin-1-yl]aminosulfonylphenyl] cyanoguanidine

Following the general procedure for cyanoguanidine formation outlined in
 example 1, N-(2,3-dichlorophenyl)-N'-[4-chloro-2-*tert*-butyldimethylsilyloxy-3-
 15 [(R)-(2-methoxymethyl)pyrrolidin-1-yl]aminosulfonylphenyl] carbodiimide (224 mg, 0.37 mmol), cyanamide (63 mg, 1.48 mmol) and N,N-diisopropylethylamine (60 mg, 0.43 mmol) were reacted, followed by desilylation with Cesium fluoride (67 mg, 0.43 mmol) to form the desired product (70 mg, 36%). EI-MS m/z 533.5 (M⁺).

Example 8 & 9

20 Preparation of N-(2-bromophenyl)-N'-[4-chloro-2-hydroxy-3-(N"-isoxazolidinylaminosulfonylphenyl] cyanoguanidine and N-(2,3-dichlorophenyl)-N'-[4-chloro-2-hydroxy-3-(N"-isoxazolidinylaminosulfonylphenyl] cyanoguanidine

N-(ethoxycarbonyl)isoxazolidine

25 To a solution of KOH (6.4g, 0.11mol) and hydroxyurethane (12g, 0.11mol) in ethanol (50mL) was added 1,3-dibromopropane (5.8mL, 0.057mol). The resulting suspension was heated at reflux for 1 hour. After the mixture was cooled to room temperature, an additional portion of KOH (3.2g, 0.055mol) and of dibromopropane (2.9mL, 0.028mol) was added. The mixture was then refluxed for 1 hour, cooled to
 30 room temperature, and solvent was evaporated. The residue was suspended in boiling ether three times and filtered. The combined filtrates were dried over sodium sulfate, filtered, and evaporated. A portion of 3g of the crude product was purified

by flush column chromatography (EtOAc/Hexane, gradient elution), yielding 1.18g of N-(ethoxycarbonyl)isoxazolidine. ¹ H NMR (CDCl₃) δ 1.15(t, 3H), 2.15 (q, 2H), 3.55(t, 2H), 3.8 (t, 2H), 4.1(q, 2H).

5 Isoxazolidine hydrochloride

N-(ethoxycarbonyl)isoxazolidine (1.18g, 9.1mmol) was dissolved in aqueous HCl (6N, 7mL) and heated at reflux for 2 hours. After being cooled to room temperature, this solution was washed with ether (3x) and then evaporated affording crude isoxazolidine hydrochloride which was recrystallized from ethanol/ether yielding
10 0.79 g (80%) of isoxazolidine hydrochloride. ¹ H NMR (CDCl₃; CH₃OD), δ 2.5 (q, 2H), 3.55 (t, 2H), 4.2(t, 2H).

(N-Isoxazolidinyl)-2,6-dichloro-3-nitrobenzenesulfonamide

Following the general procedure for sulfonamide formation outlined in
15 example 1,
2,6-dichloro-3-nitrobenzenesulfonyl chloride (1.5 g, 5.2 mmol), isoxazolidine hydrochloride (0.56 g, 5.2 mmol) and triethylamine (2.2 mL, 15.5 mmol) were reacted to form the desired product (1.2 g, 71 %). EI-MS m/z 327 (M⁺).

20 (N-Isoxazolidinyl)-2-hydroxy-6-dichloro-3-nitrobenzenesulfonamide

Following the general hydrolysis procedure outlined in example 1, (N-isoxazolidinyl)-2,6-dichloro-3-nitrobenzenesulfonamide (1.08 g, 3.3 mmol), 80% sodium hydride (0.3 g, 10.0 mmol) and water (72 mg, 4.0 mmol) were reacted to form the desired product (0.79 g, 77 %). EI-MS m/z 309 (M⁺).

25

(N-Isoxazolidinyl)-2-hydroxy-3-amino-6-dichloro-benzenesulfonamide

Following the general hydrogenation procedure outlined in example 1, (N-isoxazolidinyl)-2-hydroxy-6-dichloro-3-nitrobenzenesulfonamide (0.84 g, 2.7 mmol) was reduced with hydrogen and Pd/C (840 mg) to form the desired product (0.75 g
30 crude). EI-MS m/z 279 (M⁺).

N-(2-Bromophenyl)-N'-[4-chloro-2-hydroxy-3-(N''-isoxazolidinylaminosulfonylphenyl)] thiourea

Following the general procedure for thiourea formation outlined in example 1, (N-isoxazolidinyl)-2-hydroxy-3-amino-6-dichloro-benzenesulfonamide (624 mg, 2.24 mmol) and 2-bromophenylisothiocyanate (528 mg, 2.46 mmol) were coupled to form the desired thiourea (620 mg, 56%). EI-MS m/z 493 (M⁺).

N-(2,3-Dichlorophenyl)-N'-[4-chloro-2-hydroxy-3-(N''-isoxazolidinylaminosulfonylphenyl)] thiourea

Following the general procedure for thiourea formation outlined in example 1, (N-isoxazolidinyl)-2-hydroxy-3-amino-6-dichloro-benzenesulfonamide (590 mg, 2.11 mmol) and 2,3-dichlorophenylisothiocyanate (474 mg, 2.53 mmol) were coupled to form the desired thiourea (753 mg, 74%). EI-MS m/z 481.75 (M).

N-(2,3-Dichlorophenyl)-N'-[4-chloro-2-*tert*-butyldimethylsilyloxy-3-(N''-isoxazolidinylaminosulfonylphenyl)] thiourea

Following the general procedure for protected phenyl thiourea formation outlined in example 1, N-(2,3-dichlorophenyl)-N'-[4-chloro-2-hydroxy-3-(N''-isoxazolidinylaminosulfonylphenyl)] thiourea (726 mg, 1.51 mmol), *tert*-butyldimethylsilyl chloride (1.14 mg, 7.55 mmol) and imidazole (205 mg, 3.02 mmol) were reacted to form the desired product (355 mg, 66%). EI-MS m/z 597.83 (M⁺).

N-(2-Bromophenyl)-N'-[4-chloro-2-*tert*-butyldimethylsilyloxy-3-(N''-isoxazolidinylaminosulfonylphenyl)] carbodiimide

Following the general procedure for carbodiimide formation outlined in example 1, N-(2-bromophenyl)-N'-[4-chloro-2-*tert*-butyldimethylsilyloxy-3-(N''-isoxazolidinylaminosulfonylphenyl)] thiourea (480 mg, 0.79 mmol), methanesulfonyl chloride (0.12 mL, 1.58 mmol) and triethylamine (0.22 mL, 1.58 mmol) were reacted to form the desired product (480 mg, crude). ¹H NMR (CDCl₃) δ 0.39 (s, 6H), 1.1 (s, 9H), 2.4 (m, 2H), 3.76 (t, 2H), 4.22 (t, 2H), 7.09 (m, 2H), 7.21 (d, 1H), 7.29 (m, 1H), 7.45 (d, 1H), 7.6 (d, 1H).

N-(2,3-Dichlorophenyl)-N'-[4-chloro-2-*tert*-butyldimethylsilyloxy-3-(N''-isoxazolidinylaminosulfonylphenyl)] carbodiimide

Following the general procedure for carbodiimide formation outlined in
5 example 1, N-(2,3-dichlorophenyl)-N'-[4-chloro-2-*tert*-butyldimethylsilyloxy-3-(N''-isoxazolidinylaminosulfonylphenyl)] thiourea (355 mg, 0.59 mmol), methanesulfonyl chloride (0.09 mL, 1.18 mmol) and triethylamine (0.16 mL, 1.18 mmol) were reacted to form the desired product (355 mg, crude). ¹H NMR (CDCl₃) δ 0.38 (s, 6H), 1.03 (s, 9H), 2.4 (m, 2H), 3.77 (t, 2H), 4.22 (t, 2H), 7.12 (d, 1H), 7.16 (t, 1H),
10 7.26 (d, 1H), 7.31 (d, 1H), 7.40 (d, 1H).

N-(2-Bromophenyl)-N'-[4-chloro-2-*tert*-butyldimethylsilyloxy-3-(N''-isoxazolidinylaminosulfonylphenyl)] cyanoguanidine

Following the general procedure for cyanoguanidine formation outlined in
15 example 1, N-(2-bromophenyl)-N'-[4-chloro-2-*tert*-butyldimethylsilyloxy-3-(N''-isoxazolidinylaminosulfonylphenyl)] carbodiimide (480 mg, 0.84 mmol), cyanamide (142 mg, 3.36 mmol) and N,N-diisopropylethylamine (130 mg, 1.0 mmol) were reacted, followed by desilylation with Cesium fluoride (153.2 mg, 1.0 mmol) to form the desired product (150 mg,
20 36%). LC-MS m/z 500.

N-(2-Bromophenyl)-N'-[4-chloro-2-*tert*-butyldimethylsilyloxy-3-(N''-isoxazolidinylaminosulfonylphenyl)] cyanoguanidine

Following the general procedure for cyanoguanidine formation outlined in
25 example 1, N-(2,3-dichlorophenyl)-N'-[4-chloro-2-*tert*-butyldimethylsilyloxy-3-(N''-isoxazolidinylaminosulfonylphenyl)] carbodiimide (355 mg, 0.63 mmol), cyanamide (105 mg, 2.52 mmol) and N,N-diisopropylethylamine (100 mg, 1.26 mmol) were reacted, followed by desilylation with Cesium fluoride (115 mg, 0.76 mmol) to form the desired product (28 mg,
30 10%). LC-MS m/z 490.

Example 10 & 11

Preparation of N-(2-bromophenyl)-N'-[4-chloro-2-hydroxy-3-(N"-
tetrahydroisoxazylaminosulfonyl)phenyl] cyanoguanidine and N-(2,3-
dichlorophenyl)-N'-[4-chloro-2-hydroxy-3-(N"-tetrahydroisoxazylaminosulfonyl
5 phenyl] cyanoguanidine
N-(Ethoxycarbonyl)tetrahydroisoxazine

To a solution of KOH (3.34g, 59.6mmol) and hydroxyurethane (6.1g, 58.5mmol) in ethanol (25mL) was added 1,4-dibromobutane (3.5mL, 29.3mmol). The resulting suspension was heated at reflux for 1 hour. After the mixture was
10 cooled to room temperature, an additional portion of KOH (1.65g, 29.4mmol) and of dibromopropane (1.8mL, 15mmol) was added. The mixture was then refluxed for 1 hour, cooled to room temperature, and solvent was evaporated. The residue was suspended in boiling ether three times and filtered. The combined filtrates were dried over sodium sulfate, filtered, and evaporated. A portion of 4g of the crude product
15 was purified by flash column chromatography (EtOAc/Hexane, gradient elution), yielding 1.85g of N-(ethoxycarbonyl)tetrahydroisoxazine. ¹H NMR (CDCl₃) δ 1.05 (q, 3H), 1.45 (dd, 2H), 1.55 (dd, 2H), 3.4 (t, 2H), 3.7 (t, 2H), 3.95 (q, 2H).

Tetrahydroisoxazine hydrochloride

20 N-(ethoxycarbonyl)tetrahydroisoxazine (1.85g, 11.6mmol) was dissolved in aqueous HCl (6N, 7.8mL) and heated at reflux for 7 hours. After being cooled to room temperature, this solution was washed with ether (3x) and then evaporated affording crude tetrahydroisoxazine hydrochloride which was recrystallized from ethanol/ether yielding 0.74 g (52%) of tetrahydroisoxazine hydrochloride. ¹H NMR (CH₃OD) δ
25 1.85 (dd, 2H), 1.95 (dd, 2H), 3.4 (t, 2H), 4.25 (t, 2H).

c) (N-Tetrahydroisoxazinyl)-2,6-dichloro-3-nitrobenzenesulfonamide

Following the general procedure for sulfonamide formation outlined in example 1, 2,6-dichloro-3-nitrobenzenesulfonyl chloride (1.75 g, 6.0 mmol),
30 Tetrahydroisoxazine hydrochloride hydrochloride (0.63 g, 5.1 mmol) and triethylamine (2.2 mL, 15.5 mmol) were reacted to form the desired product (1.32 g, 75%). EI-MS m/z 341 (M⁺).

(N-Tetrahydroisoxaziny)-2-hydroxy-6-chloro-3-nitrobenzenesulfonamide

Following the general hydrolysis procedure outlined in example 1, (N-Tetrahydroisoxaziny)-2,6-dichloro-3-nitrobenzenesulfonamide (0.1 g, 0.29 mmol),
5 80% sodium hydride (26 mg, 0.88 mmol) and water (6.3 mg, 0.35 mmol) were reacted to form the desired product (50 mg, 53 %). EI-MS m/z 323 (M⁺).

(N-Tetrahydroisoxaziny)-2-hydroxy-3-amino-6-chlorobenzenesulfonamide

Following the general hydrogenation procedure outlined in example 1, (N-tetrahydroisoxaziny)-2-hydroxy-6-chloro-3-nitrobenzenesulfonamide (0.76 g, 2.35
10 mmol) was reduced with hydrogen and Pd/C (760 mg) to form the desired product (890 mg, crude). EI-MS m/z 293 (M⁺).

N-(2-Bromophenyl)-N'-[4-chloro-2-hydroxy-3-(N"-tetrahydro isoxazyl amino
15 sulfonyl)phenyl] thiourea

Following the general procedure for thiourea formation outlined in example 1, (N-tetrahydroisoxaziny)-2-hydroxy-3-amino-6-chlorobenzenesulfonamide (620
mg, 2.12 mmol) and 2-bromophenylisothiocyanate (500 mg, 2.33 mmol) were coupled to form the desired thiourea (627 mg, 58%). EI-MS m/z 507 (M).

N-(2,3-Dichlorophenyl)-N'-[4-chloro-2-hydroxy-3-(N"-tetrahydroisoxazylamino
20 sulfonyl)phenyl] thiourea

Following the general procedure for thiourea formation outlined in example 1, (N-tetrahydroisoxaziny)-2-hydroxy-3-amino-6-chlorobenzenesulfonamide (888
25 mg, 3.04 mmol) and 2,3-dichlorophenylisothiocyanate (682 mg, 3.34 mmol) were coupled to form the desired thiourea (958 mg, 64%). EI-MS m/z 495.67 (M).

N-(2-Bromophenyl)-N'-[4-chloro-2-tert-butyl dimethylsilyloxy-3-(N"-tetra hydro
isoxazylaminosulfonyl)phenyl] thiourea

30 Following the general procedure for protected phenyl thiourea formation outlined in example 1, N-(2-bromophenyl)-N'-[4-chloro-2-hydroxy-3-(N"-tetrahydroisoxazylaminosulfonyl)phenyl] thiourea

(627 mg, 1.24 mmol), *tert*-butyldimethylsilyl chloride (934 mg, 6.2 mmol) and imidazole (169 mg, 2.48 mmol) were reacted to form the desired product (350 mg, 45%). EI-MS *m/z* 622 (M^+).

5 N-(2,3-Dichlorophenyl)-N'-[4-chloro-2-*tert*-butyldimethylsilyloxy-3-(N"-tetrahydroisoxazylaminosulfonyl)phenyl] thiourea

Following the general procedure for protected phenyl thiourea formation outlined in example 1, N-(2,3-dichlorophenyl)-N'-[4-chloro-2-hydroxy-3-(N"-tetrahydroisoxazylaminosulfonyl)phenyl] thiourea

10 (898 mg, 1.81 mmol), *tert*-butyldimethylsilyl chloride (1.4 g, 9.05 mmol) and imidazole (246 mg, 3.62 mmol) were reacted to form the desired product (546 mg, 49%). EI-MS *m/z* 611.81 (M^+).

15 N-(2-Bromophenyl)-N'-[4-chloro-2-*tert*-butyldimethylsilyloxy-3-(N"-tetrahydroisoxazylaminosulfonyl)phenyl] carbodiimide

Following the general procedure for carbodiimide formation outlined in example 1, N-(2-bromophenyl)-N'-[4-chloro-2-*tert*-butyldimethylsilyloxy-3-(N"-tetrahydroisoxazylaminosulfonyl)phenyl] thiourea (385 mg, 0.74 mmol), methanesulfonyl chloride (0.12 mL, 1.48 mmol) and triethylamine (0.21 mL, 1.48
20 mmol) were reacted to form the desired product (385 mg, crude). ^1H NMR (CDCl_3) δ 0.35 (s, 6H), 1.02 (s, 9H), 1.68 (m, 2H), 1.9 (m, 2H), 3.04 (t, 2H), 4.0 (t, 2H), 7.09 (d, 1H), 7.12 (d, 1H), 7.2 (d, 1H), 7.3 (t, 1H), 7.42 (d, 1H), 7.6 (d, 1H).

25 N-(2,3-Dichlorophenyl)-N'-[4-chloro-2-*tert*-butyldimethylsilyloxy-3-(N"-tetrahydroisoxazylaminosulfonyl)phenyl] carbodiimide

Following the general procedure for carbodiimide formation outlined in example 1, N-(2,3-dichlorophenyl)-N'-[4-chloro-2-*tert*-butyldimethylsilyloxy-3-(N"-tetrahydroisoxazylaminosulfonyl)phenyl] thiourea (547 mg, 0.897 mmol), methanesulfonyl chloride (0.14 mL, 1.79 mmol) and triethylamine (0.25 mL, 1.79
30 mmol) were reacted to form the desired product (547 mg, crude). ^1H NMR (CDCl_3) δ 0.36 (s, 6H), 1.03 (s, 9H), 1.68 (m, 2H), 1.9 (m, 2H), 3.53 (t, 2H), 4.0 (t, 2H), 7.11 (m, 2H), 7.14 (t, 1H), 7.2 (d, 1H), 7.3 (d, 1H), 7.39 (d, 1H).

N-(2-Bromophenyl)-N'-[4-chloro-2-*tert*-butyldimethylsilyloxy-3-(N"-tetrahydroisoxazylaminosulfonyl)phenyl] cyanoguanidine

Following the general procedure for cyanoguanidine formation outlined in example 1, N-(2-bromophenyl)-N'-[4-chloro-2-*tert*-butyldimethylsilyloxy-3-(N"-tetrahydroisoxazylaminosulfonyl)phenyl] carbodiimide (385 mg, 0.79 mmol), cyanamide (133 mg, 3.16 mmol) and N,N-diisopropylethylamine (127 mg, 0.95 mmol) were reacted, followed by desilylation with Cesium fluoride (144 mg, 0.95 mmol) to form the desired product (100 mg, 26%). EI-MS m/z 515 (M⁺).

N-(2,3-Dichlorophenyl)-N'-[4-chloro-2-*tert*-butyldimethylsilyloxy-3-(N"-tetrahydroisoxazylaminosulfonyl)phenyl] cyanoguanidine

Following the general procedure for cyanoguanidine formation outlined in example 1, N-(2,3-dichlorophenyl)-N'-[4-chloro-2-*tert*-butyldimethylsilyloxy-3-(N"-tetrahydroisoxazylaminosulfonyl)phenyl] carbodiimide (547 mg, 0.95 mmol), cyanamide (159 mg, 3.8 mmol) and N,N-diisopropylethylamine (150 mg, 1.14 mmol) were reacted, followed by desilylation with Cesium fluoride (174 mg, 1.14 mmol) to form the desired product (190 mg, 40%). LC-MS m/z 504.

Example 12

Preparation of N-[4-chloro-2-hydroxy-3-[N",N"-dimethylaminosulfonyl]phenyl]-N'-(2-bromophenyl) propylguanidine
N, N-Dimethyl-3-amino-6-chloro-2-hydroxybenzenesulfonamide

To a solution of N, N-Dimethyl-6-chloro-2-hydroxy-3-nitrobenzenesulfonamide (550 mg, 1.96 mmol) in ethyl acetate, was added 10% Pd/C (550 mg). The mixture was flushed with argon, and then stirred under a hydrogen atmosphere at balloon pressure for 3 hours at room temperature. TLC showed the reaction was not complete. The mixture was filtered through celite and the celite was washed with methanol. The mixture was retreated under the same conditions as mentioned above. After 3 hours, the solvent was evaporated to give the desired product (480 mg, 98%). EI-MS (m/z) 250.82, 252.87 (M⁺).

N-[4-chloro-2-hydroxy-3-[N",N"-dimethylaminosulfonyl]phenyl]-N'-(2-bromophenyl) thiourea

A solution of N,N-dimethyl-3-amino-6-chloro-2-hydroxybenzene sulfonamide (480 mg, 3.58 mmol) and 2-bromophenylisothiocyanate (0.53 μ L, 3.94 mmol) in 10 mL of ethanol was stirred at room temperature for overnight. Purification by column chromatography on silica gel, eluting with ethyl acetate/hexane (gradient elution) gave the desired product (368 mg, 22 %). EI-MS (m/z) 463.67, 465.66, 467.65, 468.64 (M^+).

10 N-[4-chloro-2-tert-butyldimethylsiloxy-3-[N",N"-dimethylaminosulfonyl]phenyl]-N'-(2-bromophenyl) thiourea

To a solution of N-[4-chloro-2-hydroxy-3-[N",N"-dimethylamino sulfonyl]phenyl]-N'-(2-bromophenyl) thiourea (350 mg, 0.755 mmol) in THF (10 mL), tert-butyldimethylsilyl chloride (171 mg, 1.13 mmol) and imidazole (106mg, 1.56 mmol) were added. The reaction mixture was stirred at room temperature for overnight. Then it was partitioned between ethyl acetate and water. The combined organic phase was dried and concentrated. Chromatography of the residue on silica gel gave desired product (150 mg, 46 %) and recovered starting material (90 mg). EI-MS m/z 577.81, 579.66, 580.65, 581.46, 582.77 (M^+).

20

N-[4-chloro-2-tert-butyldimethylsiloxy-3-[N",N"-dimethylaminosulfonyl]phenyl]-N'-(2-bromophenyl)carbodiimide

To a solution of N-[4-chloro-2-tert-butyldimethylsiloxy-3-[N",N"-dimethylaminosulfonyl]phenyl]-N'-(2-bromophenyl) thiourea (150 mg, 0.26mmol) in dichloromethane (5 mL) at 0°C, methanesulfonyl chloride (40 μ L, 0.52 mmol), 4-Dimethylaminopyridine (10 mg) and triethylamine (0.11 mL, 0.77 mmol) were added. The reaction mixture was stirred at 0°C for 3 hours. Then it was partitioned between ethyl acetate and water. The combined organic phase was dried and concentrated to give the desired product (120 mg, 85%). IR: 2140 cm^{-1}

N-[4-chloro-2-hydroxy-3-[N",N"-dimethylaminosulfonyl]phenyl]-N'-(2-bromophenyl) propylguanidine

To a solution of N-[4-chloro-2-tert-butyl dimethylsiloxy-3-[N",N"-dimethylaminosulfonyl]phenyl]-N'-(2-bromophenyl)carbodiimide (60 mg, 0.11 mmol) in tetrahydrofuran (2 mL) at room temperature, N,N-diisopropylethylamine (14 μ L, 0.12 mmol) and n-propylamine (10 μ L, 0.12 mmol) were added. The reaction mixture was stirred at room temperature for 15 minutes, then cooled to 0°C, and TBAF and methanol (1 mL) were added. After 30 minutes, the mixture was quenched with water, and extracted with ethylacetate. The organic layer was concentrated under reduced pressure. The residue was purified by Gilson HPLC to give the desired product (19 mg, 35 %). EI-MS m/z 488.7, 490.68, 492.67 (M^+).

Example 13

Preparation of N-[4-chloro-2-hydroxy-3-(N",N"-dimethylaminosulfonyl)phenyl]-N'-(2-chlorophenyl) cyanoguanidine

a) N-[4-Chloro-2-hydroxy-3-(N",N"-dimethylaminosulfonyl)phenyl]-N'-(2-chlorophenyl) thiourea

Following the general procedure for thiourea formation outlined in example 12, N,N-dimethyl-3-amino-6-chloro-2-hydroxybenzenesulfonamide (450 mg, 1.8 mmol) and 2-chlorophenylisothiocyanate (335.6 mg, 1.98 mmol) were coupled to form the desired thiourea (526 mg, 70%). EI-MS m/z 421.47 (M^+).

b) N-(2-Chlorophenyl)-N'-[4-chloro-2-tert-butyl dimethylsilyloxy-3-(N",N"-dimethylaminosulfonyl)phenyl] thiourea

Following the general procedure for protected phenyl thiourea formation outlined in example 12, N-[4-chloro-2-hydroxy-3-(N",N"-dimethylamino sulfonyl)phenyl]-N'-(2-chlorophenyl) thiourea (500 mg, 1.19 mmol), tert-butyl dimethylsilyl chloride (897 mg, 5.95 mmol) and imidazole (162 mg, 2.38 mmol) were reacted to form the desired product (311 mg, 49%). EI-MS m/z 535.32 (M^+).

c) N-(2-Chlorophenyl)-N'-[4-chloro-2-tert-butyldimethylsilyloxy-3-(N'',N''-dimethylaminosulfonyl)phenyl]carbodiimide

Following the general procedure for carbodiimide formation outlined in example 12, N-(2-chlorophenyl)-N'-[4-chloro-2-tert-butyldimethylsilyloxy-3-(N'',N''-dimethylaminosulfonyl)phenyl] thiourea (311 mg, 0.58 mmol), methanesulfonyl chloride (0.09 mL, 1.16 mmol) and triethylamine (0.15 mL, 1.16 mmol) were reacted to form the desired product (300 mg, crude). ¹H NMR (CDCl₃) δ 0.37 (s, 6H), 1.03 (s, 9H), 2.87 (s, 6H), 7.05 (d, 1H), 7.12 (t, 1H), 7.22 (m, 2H), 7.34 (d, 1H), 7.42 (d, 1H).

d) N-(2-Chlorophenyl)-N'-[4-chloro-3-(N'',N''-dimethylaminosulfonyl)phenyl] cyanoguanidine

Following the general procedure for cyanoguanidine formation outlined in example 12, N-(2-chlorophenyl)-N'-[4-chloro-2-tert-butyldimethylsilyloxy-3-(N'',N''-dimethylaminosulfonyl)phenyl] carbodiimide (300 mg, 0.6 mmol), cyanamide (100 mg, 2.4 mmol) and N,N-diisopropylethylamine (94 mg, 0.72 mmol) were reacted, followed by desilylation with Cesium fluoride (110 mg, 0.72 mmol) to form the desired product (129 mg, 50%). EI-MS m/z 428.0 (M⁺). ¹H NMR (DMSO-d₆) δ 2.86 (s, 6H), 7.2 (d, 1H), 7.32 (t, 1H), 7.36 (t, 1H), 7.43 (d, 1H), 7.51 (d, 1H), 7.57 (d, 1H), 8.98 (s, 1H), 9.18 (s, 1H), 10.51 (s, 1H).

Example 14

Preparation of N-[4-chloro-2-hydroxy-3-(N'',N''-dimethylaminosulfonyl)phenyl]-N'-(2-fluoro-3-chlorophenyl) cyanoguanidine

a) 2-Fluoro-3-chlorophenylisothiocyanate

Into a solution of 2-fluoro-3-chloroaniline (1.0 g, 6.87 mmol) in 30 mL of toluene at room temperature, thiophosgene (0.8 mL, 10.3 mmol) and triethylamine (1.12 mL, 8.24 mmol) were added. The mixture was stirred at room temperature for 16 hours. The mixture was partitioned between ethyl acetate and water. The combined organic layer was then concentrated to give the desired product (950 mg, 74 %). ¹H NMR (CDCl₃) δ 7.09 (m, 2H), 7.30 (m, 1H).

b) N-[4-Chloro-2-hydroxy-3-(N",N"-dimethylaminosulfonyl)phenyl]-N'-(2-fluoro-3-chlorophenyl) thiourea

Following the general procedure for thiourea formation outlined in example 12, N,N-dimethyl-3-amino-6-chloro-2-hydroxybenzenesulfonamide (500 mg, 2 mmol) and 2-fluoro-3-chlorophenylisothiocyanate (374 mg, 2 mmol) were coupled to form the desired thiourea (583 mg, 67%). EI-MS m/z 438.2 (M⁺).

c) N-(2-Fluoro-3-chlorophenyl)-N'-[4-chloro-2-tert-butyldimethylsilyloxy-3-(N",N"-dimethylaminosulfonyl)phenyl]-N'-(2-fluoro-3-chlorophenyl) thiourea

Following the general procedure for protected phenyl thiourea formation outlined in example 12, N-[4-chloro-2-hydroxy-3-(N",N"-dimethylaminosulfonyl)phenyl]-N'-(2-fluoro-3-chlorophenyl) thiourea (533 mg, 1.22 mmol), tert-butyldimethylsilyl chloride (913 mg, 6.1 mmol) and imidazole (166 mg, 2.44 mmol) were reacted to form the desired product (412 mg, 61%). EI-MS m/z 552.2 (M⁺).

d) N-(2-Fluoro-3-chlorophenyl)-N'-[4-chloro-2-tert-butyldimethylsilyloxy-3-(N",N"-dimethylaminosulfonyl)phenyl]-N'-(2-fluoro-3-chlorophenyl) carbodiimide

Following the general procedure for carbodiimide formation outlined in example 12, N-(2-fluoro-3-chlorophenyl)-N'-[4-chloro-2-tert-butyldimethylsilyloxy-3-(N",N"-dimethylaminosulfonyl)phenyl]-N'-(2-fluoro-3-chlorophenyl) thiourea (412 mg, 0.75 mmol), methanesulfonyl chloride (0.14 mL, 1.5 mmol) and triethylamine (0.24 mL, 1.5 mmol) were reacted to form the desired product (410 mg, crude). ¹H NMR (CDCl₃) δ 0.37 (s, 6H), 0.99 (s, 9H), 2.85 (s, 6H), 7.05 (m, 2H), 7.25 (m, 3H).

e) Preparation of N-[4-chloro-2-hydroxy-3-(N",N"-dimethylaminosulfonyl)phenyl]-N'-(2-fluoro-3-chlorophenyl) cyanoguanidine

Following the general procedure for cyanoguanidine formation outlined in example 12, N-[4-chloro-2-tert-butyldimethylsilyloxy-3-(N",N"-dimethylaminosulfonyl)phenyl]-N'-(2-fluoro-3-chlorophenyl) carbodiimide (410 mg,

0.79 mmol), cyanamide (133mg, 3.16 mmol) and N,N-diisopropylethylamine (122 mg, 0.95 mmol) were reacted, followed by desilylation with Cesium fluoride (144 mg, 0.95 mmol) to form the desired product (140 mg, 40%). EI-MS m/z 446.2 (M^+). 1H NMR (DMSO- d_6) δ 2.87 (s, 6H), 7.2 (d, 1H), 7.32 (t, 1H), 7.40 (m, 2H), 7.59 (d, 1H), 9.11 (s, 1H), 9.3 (s, 1H), 10.53 (s, 1H).

Example 15

Preparation of N-[4-chloro-2-hydroxy-3-(N",N"-dimethylaminosulfonyl)phenyl]-N'-(2-trifluoromethylphenyl) cyanoguanidine

10 a) 2-Trifluoromethylisothiocyanate

Into a solution of 2-trifluoromethylaniline (1.0 g, 6.21 mmol) in 30 mL of toluene at room temperature, thiophosgene (0.72 mL, 9.31 mmol) and triethylamine (1.01 mL, 7.45 mmol) was added. The mixture was stirred at room temperature for 16 hours. The mixture was parationed between ethyl acetate and water. The combined organic layer was then concentrated to give the desired product (1.01 g, 80 %). 1H NMR ($CDCl_3$) δ 7.30 (m, 2H), 7.54 (t, 1H), 7.64 (d, 1H).

b) N-[4-Chloro-2-hydroxy-3-(N",N"-dimethylaminosulfonyl)phenyl]-N'-(2-trifluoromethylphenyl) thiourea

20 Following the general procedure for thiourea formation outlined in example 12, N,N-dimethyl-3-amino-6-chloro-2-hydroxybenzenesulfonamide (500 mg, 2 mmol) and 2-trifluoromethylphenylisothiocyanate (548 mg, 2 mmol) were coupled to form the desired thiourea (469 mg, 52%). EI-MS m/z 454.0 (M^+).

25 c) N-(2-Trifluoromethylphenyl)-N'-[4-chloro-2-tert-butyldimethylsilyloxy-3-(N",N"-dimethylaminosulfonyl)phenyl] thiourea

Following the general procedure for protected phenyl thiourea formation outlined in example 12, N-[4-chloro-2-hydroxy-3-(N",N"-dimethylaminosulfonyl)phenyl]-N'-(2-trifluoromethylphenyl) thiourea (416 mg, 1.0 mmol), tert-butyldimethylsilyl chloride (750 mg, 5.0 mmol) and imidazole (136 mg, 2.0 mmol) were reacted to form the desired product (250 mg, 45%). EI-MS m/z 568.0 (M^+).

d) N-(2-Trifluoromethylphenyl)-N'-[4-chloro-2-tert-butyldimethylsilyloxy-3-(N'',N''-dimethylaminosulfonyl)phenyl] carbodiimide

Following the general procedure for carbodiimide formation outlined in
 5 example 12, N-(2-trifluoromethylphenyl)-N'-[4-chloro-2-tert-butyldimethylsilyloxy-3-(N'',N''-dimethylaminosulfonyl)phenyl]thiourea (250 mg, 0.44 mmol),
 methanesulfonyl chloride (0.1 mL, 0.88 mmol) and triethylamine (0.14 mL, 0.88 mmol) were reacted to form the desired product (250 mg, crude). ¹H NMR (CDCl₃)
 δ 0.38 (s, 6H), 1.04 (s, 9H), 2.87 (s, 6H), 7.07 (d, 1H), 7.2 (d, 1H), 7.29 (m, 2H),
 10 7.54 (t, 1H), 7.66 (d, 1H).

e) Preparation of N-[4-chloro-2-hydroxy-3-(N'',N''-dimethylaminosulfonyl)phenyl]-N'-(2-trifluoromethylphenyl) cyanoguanidine
 cyanoguanidine

Following the general procedure for cyanoguanidine formation outlined in
 15 example 12, N-[4-chloro-2-tert-butyldimethylsilyloxy-3-(N'',N''-dimethylaminosulfonyl)phenyl]-N'-(2-trifluoromethylphenyl) carbodiimide (250 mg, 0.47 mmol), cyanamide (79 mg, 1.88 mmol) and N,N-diisopropylethylamine (73 mg, 0.56 mmol) were reacted, followed by desilylation with Cesium fluoride (86
 20 mg, 0.56 mmol) to form the desired product (80 mg, 37%). EI-MS m/z 462.0 (M⁺). ¹H NMR (DMSO-d₆) δ 2.86 (s, 6H), 7.14 (d, 1H), 7.53 (m, 2H), 7.59 (d, 1H), 7.75 (t, 1H), 7.77 (d, 1H), 8.91 (s, 1H), 9.28 (s, 1H), 10.53 (s, 1H).

Example 16

Preparation of N-[4-chloro-2-hydroxy-3-(N'',N''-dimethylaminosulfonyl)phenyl]-N'-(2-methylphenyl) cyanoguanidine
 25

a) N-[4-Chloro-2-hydroxy-3-(N'',N''-dimethylaminosulfonyl)phenyl]-N'-(2-methylphenyl) thiourea

Following the general procedure for thiourea formation outlined in example
 30 12, N,N-dimethyl-3-amino-6-chloro-2-hydroxybenzenesulfonamide (500 mg, 2 mmol) and 2-methylphenylisothiocyanate (298.4 mg, 2 mmol) were coupled to form the desired thiourea (557 mg, 70%). EI-MS m/z 400.0 (M⁺).

b) N-(2-Methylphenyl)-N'-[4-chloro-2-tert-butyldimethylsilyloxy-3-(N'',N''-dimethylaminosulfonyl)phenyl] thiourea

Following the general procedure for protected phenyl thiourea formation outlined in example 12, N-[4-Chloro-2-hydroxy-3-[N'',N''-dimethylaminosulfonyl]phenyl]-N'-(2-methylphenyl) thiourea (557 mg, 1.39 mmol), tert-butyldimethylsilyl chloride (1.04 mg, 6.95 mmol) and imidazole (189 mg, 2.78 mmol) were reacted to form the desired product (410 mg, 57%). EI-MS m/z 514.2 (M⁺).

10

c) N-(2-Methylphenyl)-N'-[4-chloro-2-tert-butyldimethylsilyloxy-3-(N'',N''-dimethylaminosulfonyl)phenyl] carbodiimide

Following the general procedure for carbodiimide formation outlined in example 12, N-(2-methylphenyl)-N'-[4-chloro-2-tert-butyldimethylsilyloxy-3-(N'',N''-dimethylaminosulfonyl)phenyl]thiourea (410 mg, 0.8 mmol), methanesulfonyl chloride (0.14 mL, 1.6 mmol) and triethylamine (0.23 mL, 1.6 mmol) were reacted to form the desired product (400 mg, crude). ¹H NMR (CDCl₃) δ 0.39 (s, 6H), 1.05 (s, 9H), 1.72 (m, 4H), 2.37 (s, 3H), 2.87 (s, 6H), 7.04 (d, 1H), 7.18 (m, 4H), 7.29 (d, 1H).

20

d) Preparation of N-[4-chloro-2-hydroxy-3-(N'',N''-dimethylaminosulfonyl)phenyl]-N'-(2-methylphenyl) cyanoguanidine
cyanoguanidine

Following the general procedure for cyanoguanidine formation outlined in example 12, N-[4-chloro-2-tert-butyldimethylsilyloxy-3-(N'',N''-dimethylaminosulfonyl)phenyl]-N'-(2-methylphenyl) carbodiimide (400 mg, 0.83 mmol), cyanamide (139.4 mg, 3.32 mmol) and N,N-diisopropylethylamine (129 mg, 1.0 mmol) were reacted, followed by desilylation with Cesium fluoride (152 mg, 1.0 mmol) to form the desired product (110 mg, 32%). EI-MS m/z 408.2 (M⁺). ¹H NMR (DMSO-d₆) δ 2.22 (s, 3H), 2.86 (s, 6H), 7.20 (m, 5H), 7.60 (d, 1H), 8.63 (s, 1H), 8.96 (s, 1H), 10.50 (s, 1H).

30

Example 17Preparation of N-[4-chloro-2-hydroxy-3-(N",N"-dimethylaminosulfonyl)phenyl]-N'-(2-fluorophenyl) cyanoguanidine

5 a) 2-Fluorophenylisothiocyanate

Into a solution of 2-fluoroaniline (1.0 g, 9.0 mmol) in 30 mL of toluene at room temperature, thiophosgene (1.06 mL, 13.5 mmol) and triethylamine (1.55 mL, 13.5 mmol) was added. The mixture was stirred at room temperature for 16 hours. The mixture was partitioned between ethyl acetate and water. The combined organic
10 layer was then concentrated to give the desired product (1.05 g, 76 %). ¹ H NMR (CDCl₃) δ 7.1-7.3 (m, 4H).

b) N-[4-Chloro-2-hydroxy-3-(N",N"-dimethylaminosulfonyl)phenyl]-N'-(2-fluorolphenyl) thiourea

15 Following the general procedure for thiourea formation outlined in example 12, N,N-dimethyl-3-amino-6-chloro-2-hydroxybenzenesulfonamide (500 mg, 2 mmol) and 2-fluorophenylisothiocyanate (306 mg, 2 mmol) were coupled to form the desired thiourea (497 mg, 62%). EI-MS m/z 404.2 (M⁺).

20 c) N-(2-Fluorophenyl)-N'-[4-chloro-2-tert-butyldimethylsilyloxy-3-(N",N"-dimethylaminosulfonyl)phenyl] thiourea

Following the general procedure for protected phenyl thiourea formation outlined in example 12, N-[4-chloro-2-hydroxy-3-(N",N"-dimethylaminosulfonyl)phenyl]-N'-(2-fluorophenyl) thiourea (440 mg, 1.09 mmol),
25 tert-butyldimethylsilyl chloride (817 mg, 5.45 mmol) and imidazole (148 mg, 2.18 mmol) were reacted to form the desired product (387 mg, 69%). EI-MS m/z 518.2 (M⁺).

30 d) N-(2-Fluorophenyl)-N'-[4-chloro-2-tert-butyldimethylsilyloxy-3-(N",N"-dimethylaminosulfonyl)phenyl] carbodiimide

Following the general procedure for carbodiimide formation outlined in example 12, N-(2-fluorophenyl)-N'-[4-chloro-2-tert-butyldimethylsilyloxy-3-(N",N"-

dimethylaminosulfonyl)phenyl]thiourea (357 mg, 0.69 mmol), methanesulfonyl chloride (0.12 mL, 1.38 mmol) and triethylamine (0.2 mL, 1.38 mmol) were reacted to form the desired product (350 mg, crude). ¹H NMR (CDCl₃) δ 0.38 (s, 6H), 1.05 (s, 9H), 1.72 (m, 4H), 2.87 (s, 6H), 7.06 (d, 1H), 7.19 (m, 3H), 7.28 (d, 1H), 7.31 (d, 1H).

e) Preparation of N-[4-chloro-2-hydroxy-3-(N",N"-dimethylaminosulfonyl)phenyl]-N'-(2-fluorophenyl) cyanoguanidine

Following the general procedure for cyanoguanidine formation outlined in example 12, N-[4-chloro-2-tert-butyldimethylsilyloxy-3-(N",N"-dimethylaminosulfonyl)phenyl]-N'-(2-fluorophenyl) carbodiimide (350 mg, 0.72 mmol), cyanamide (121 mg, 2.88 mmol) and N,N-diisopropylethylamine (111 mg, 0.86 mmol) were reacted, followed by desilylation with Cesium fluoride (132 mg, 0.86 mmol) to form the desired product (120 mg, 40%). EI-MS m/z 412.2 (M⁺). ¹H NMR (DMSO-d₆) δ 2.86 (s, 6H), 7.20 (m, 2H), 7.27 (m, 2H), 7.40 (t, 1H), 7.52 (d, 1H), 9.06 (s, 1H), 9.20 (s, 1H), 10.51 (s, 1H).

Example 18

Preparation of N-[4-chloro-2-hydroxy-3-(N",N"-dimethylaminosulfonyl)phenyl]-N'-(2,3-difluorophenyl) cyanoguanidine

a) 2,3-Difluorophenylisothiocyanate

Into a solution of 2,3-difluoroaniline (1.0 g, 7.74 mmol) in 30 mL of toluene at room temperature, thiophosgene (0.91 mL, 11.6 mmol) and triethylamine (1.3 mL, 11.6 mmol) was added. The mixture was stirred at room temperature for 16 hours. The mixture was partitioned between ethyl acetate and water. The combined organic layer was then concentrated to give the desired product (910 mg, 68 %). ¹H NMR (CDCl₃) δ 6.98 (m, 2H), 7.11 (m, 1H).

b) N-[4-Chloro-2-hydroxy-3-(N",N"-dimethylaminosulfonyl)phenyl]-N'-(2,3-difluorophenyl) thiourea

Following the general procedure for thiourea formation outlined in example 12, N,N-dimethyl-3-amino-6-chloro-2-hydroxybenzenesulfonamide (500 mg, 2 mmol) and 2,3-difluorophenylisothiocyanate (342 mg, 2 mmol) were coupled to form the desired thiourea (467 mg, 54%). EI-MS m/z 422.2 (M^+).

5

c) N-(2,3-Difluorophenyl)-N'-[4-chloro-2-tert-butyldimethylsilyloxy-3-(N'',N''-dimethylaminosulfonyl)phenyl] thiourea

Following the general procedure for protected phenyl thiourea formation outlined in example 12, N-[4-chloro-2-hydroxy-3-(N'',N''-dimethylaminosulfonyl)phenyl]-N'-(2,3-difluorophenyl)thiourea (418 mg, 1.0 mmol), tert-butyldimethylsilyl chloride (745 mg, 5.0 mmol) and imidazole (136 mg, 2.0 mmol) were reacted to form the desired product (347 mg, 65%). EI-MS m/z 536.2 (M^+).

15 d) N-(2,3-Difluorophenyl)-N'-[4-chloro-2-tert-butyldimethylsilyloxy-3-(N'',N''-dimethylaminosulfonyl)phenyl] carbodiimide

Following the general procedure for carbodiimide formation outlined in example 12, N-(2,3-difluorophenyl)-N'-[4-chloro-2-tert-butyldimethylsilyloxy-3-(N'',N''-dimethylaminosulfonyl)phenyl]thiourea (347 mg, 0.69 mmol), methanesulfonyl chloride (0.12 mL, 1.38 mmol) and triethylamine (0.2 mL, 1.04 mmol) were reacted to form the desired product (340 mg, crude). 1H NMR ($CDCl_3$) δ 0.39 (s, 6H), 1.03 (s, 9H), 2.86 (s, 6H), 6.95 (t, 1H), 7.03 (m, 2H), 7.08 (d, 1H), 7.31 (d, 1H).

25 e) Preparation of N-[4-chloro-2-hydroxy-3-(N'',N''-dimethylaminosulfonyl)phenyl]-N'-(2,3-difluorophenyl) cyanoguanidine

Following the general procedure for cyanoguanidine formation outlined in example 12, N-[4-chloro-2-tert-butyldimethylsilyloxy-3-(N'',N''-dimethylaminosulfonyl)phenyl]-N'-(2,3-difluorophenyl) carbodiimide (340 mg, 0.68 mmol), cyanamide (114 mg, 2.72 mmol) and N,N-diisopropylethylamine (106 mg, 0.82 mmol) were reacted, followed by desilylation with Cesium fluoride (124 mg, 0.82 mmol) to form the desired product (10 mg, 3.4%). EI-MS m/z 430.0 (M^+). 1H

30

NMR (DMSO- d_6) δ 2.87 (s, 6H), 7.21 (m, 3H), 7.30 (m, 1H), 7.54 (d, 1H), 9.2 (s, 1H), 10.54 (s, 1H).

Example 19

Preparation of N-[4-chloro-2-hydroxy-3-(N",N"-dimethylaminosulfonyl)phenyl]-N'-(2-methyl-3-fluorophenyl) cyanoguanidine

a) 2-Methyl-3-fluorophenylisothiocyanate

Into a solution of 2-methyl-3-fluoroaniline (1.0 g, 8.0 mmol) in 30 mL of toluene at room temperature, thiophosgene (0.94 mL, 12 mmol) and triethylamine (1.34 mL, 12 mmol) was added. The mixture was stirred at room temperature for 16 hours. The mixture was partitioned between ethyl acetate and water. The combined organic layer was then concentrated to give the desired product (1.1 g, 82 %). EI-MS m/z 168.2 (M^+).

b) N-[4-Chloro-2-hydroxy-3-(N",N"-dimethylaminosulfonyl)phenyl]-N'-(2-methyl-3-fluorophenyl) thiourea

Following the general procedure for thiourea formation outlined in example 12, N,N-dimethyl-3-amino-6-chloro-2-hydroxybenzenesulfonamide (560 mg, 2.23 mmol) and 2-methyl-3-fluorophenylisothiocyanate (372 mg, 2.23 mmol) were coupled to form the desired thiourea (570 mg, 61%). EI-MS m/z 418.2 (M^+).

c) N-(2-Methyl-3-fluorophenyl)-N'-[4-chloro-2-tert-butyldimethylsilyloxy-3-(N",N"-dimethylaminosulfonyl)phenyl] thiourea

Following the general procedure for protected phenyl thiourea formation outlined in example 12, N-[4-chloro-2-hydroxy-3-(N",N"-dimethylaminosulfonyl)phenyl]-N'-(2-methyl-3-fluorophenyl)thiourea (530 mg, 1.27 mmol), tert-butyldimethylsilyl chloride (951 mg, 6.35 mmol) and imidazole (173 mg, 2.54 mmol) were reacted to form the desired product (331 mg, 49%). EI-MS m/z 532.2 (M^+).

d) N-(2-Methyl-3-fluorophenyl)-N'-[4-chloro-2-tert-butyldimethylsilyloxy-3-(N",N"-dimethylaminosulfonyl)phenyl] carbodiimide

Following the general procedure for carbodiimide formation outlined in example 12, N-(2-methyl-3-fluorophenyl)-N'-[4-chloro-2-tert-butyl dimethylsilyloxy-3-(N'',N''-dimethylaminosulfonyl)phenyl]thiourea (330 mg, 0.62 mmol), methanesulfonyl chloride (0.12 mL, 1.38 mmol) and triethylamine (0.2 mL, 1.04 mmol) were reacted to form the desired product (320 mg, crude). ¹H NMR (CDCl₃) δ 0.38 (s, 6H), 1.05 (s, 9H), 2.28 (s, 3H), 2.85 (s, 6H), 6.89 (t, 1H), 6.93 (d, 1H), 7.05 (d, 1H), 7.13 (m, 1H), 7.19 (d, 1H).

f) Preparation of N-[4-chloro-2-hydroxy-3-(N'',N''-dimethylaminosulfonyl)phenyl]-N'-(2-methyl-3-fluorophenyl) cyanoguanidine

Following the general procedure for cyanoguanidine formation outlined in example 12, N-[4-chloro-2-tert-butyl dimethylsilyloxy-3-(N'',N''-dimethylaminosulfonyl)phenyl]-N'-(2-methyl-3-fluorophenyl) carbodiimide (320 mg, 0.64 mmol), cyanamide (108 mg, 2.56 mmol) and N,N-diisopropylethylamine (99 mg, 0.77 mmol) were reacted, followed by desilylation with Cesium fluoride (117 mg, 0.77 mmol) to form the desired product (120 mg, 44%). EI-MS m/z 426.2 (M⁺). ¹H NMR (DMSO-d₆) δ 2.12 (s, 3H), 2.86 (s, 6H), 7.09 (m, 3H), 7.22 (m, 1H), 7.56 (d, 1H), 8.82 (s, 1H), 9.10 (s, 1H), 10.51 (s, 1H).

Example 20

20 Preparation of N-[4-chloro-2-hydroxy-3-(N'',N''-dimethylaminosulfonyl)phenyl]-N'-(2-chloro-3-fluorophenyl) cyanoguanidine

a) 2-Chloro-3-fluorobenzoic acid

A solution of 3-fluorobenzoic acid (8.0 g, 64.3 mmol) in 40 mL of THF was added dropwise to a solution of sec-butyllithium (90 mL, 128.6 mmol) and N,N,N',N'-tetramethylethylenediamine (20.0 mL, 147.9 mmol) in THF (100 mL) at -90°C. After addition, the reaction mixture was stirred at -90°C for 30 minutes. Hexachloroethane (54 g, 257.2 mmol) in THF (100 mL) was added to reaction mixture dropwise. Then the reaction mixture was stirred at -78°C to room temperature for 16 hours. The solvent was evaporated and the water was added to the residue. The reaction mixture was acidified to PH=1 by added conc. hydrochloric acid. Then the reaction mixture was extracted with ether (3x). The combined organic

phase was dried and conc. The crude product was washed with hexane for 3 times. Then it was filtered to get pure product (9.2 g, 93%). EI-MS m/z 172.89 (M^+).

b) 2-Chloro-3-fluoroaniline

5 To a solution of 2-chloro-3-fluorobenzoic acid (4 g, 23 mmol) in $CHCl_3$ (50 mL), sulfonic acid (64 mL) was added and then sodium azide (2.62 g, 1.75 mmol) was added by portion. After addition, the reaction mixture was stirred at 50°C for 16 hours. The solvent was evaporated and basicified with ammonium hydroxide at ice-bath. Then was extracted with ethyl acetate (3x). The combine organic phase was
10 dried and conc. to give the desired product (2.61 g, 78%). LC-MS m/z 146.2 (M^+).

c) 2-Chloro-3-fluorophenylisothiocyanate

Into a solution of 2-chloro-3-fluoroaniline (2.61 g, 17.94 mmol) in 50 mL of toluene at room temperature, thiophosgene (2.1 mL, 26.91 mmol) and triethylamine
15 (3.02 mL, 26.91 mmol) was added. The mixture was stirred at room temperature for 16 hours. The mixture was partitioned between ethyl acetate and water. The combined organic layer was then concentrated to give the desired product (2.99 g, 89 %). 1H NMR ($CDCl_3$) δ 7.10 (m, 1H), 7.22 (m, 1H).

20 d) N-[4-Chloro-2-hydroxy-3-(N,N' -dimethylaminosulfonyl)phenyl]- N' -(2-chloro-3-fluorophenyl) thiourea

Following the general procedure for thiourea formation outlined in example 12, N,N -dimethyl-3-amino-6-chloro-2-hydroxybenzenesulfonamide (500 mg, 2.0 mmol) and 2-chloro-3-fluorophenylisothiocyanate (374 mg, 2.0 mmol) were coupled
25 to form the desired thiourea (623 mg, 71%). LC-MS m/z 438.2 (M^+).

e) N-(2-Chloro-3-fluorophenyl)- N' -[4-chloro-2-tert-butyldimethylsilyloxy-3-(N,N' -dimethylaminosulfonyl)phenyl] thiourea

Following the general procedure for protected phenyl thiourea formation
30 outlined in example 12, N-[4-chloro-2-hydroxy-3-(N,N' -dimethylaminosulfonyl)phenyl]- N' -(2-chloro-3-fluorophenyl)thiourea (575 mg, 1.32 mmol), tert-butyldimethylsilyl chloride (991 mg, 6.6 mmol) and imidazole (179 mg,

2.64 mmol) were reacted to form the desired product (367 mg, 50%). LC-MS m/z 552.2 (M^+).

- 5 f) N-(2-Chloro-3-fluorophenyl)-N'-[4-chloro-2-tert-butyldimethylsilyloxy-3-(N'',N''-dimethylaminosulfonyl)phenyl] carbodiimide

Following the general procedure for carbodiimide formation outlined in example 12, N-(2-chloro-3-fluorophenyl)-N'-[4-chloro-2-tert-butyldimethylsilyloxy-3-(N'',N''-dimethylaminosulfonyl)phenyl]thiourea (367 mg, 0.67 mmol), methanesulfonyl chloride (0.12 mL, 1.38 mmol) and triethylamine (0.2 mL, 1.04 mmol) were reacted to form the desired product (360 mg, crude). 1H NMR ($CDCl_3$) δ 0.36 (s, 6H), 1.03 (s, 9H), 1.2.87 (s, 6H), 7.01 (m, 2H), 7.09 (d, 1H), 7.21 (m, 1H), 7.34 (d, 1H).

- 15 g) Preparation of N-[4-chloro-2-hydroxy-3-(N'',N''-dimethylaminosulfonyl)phenyl]-N'-(2-chloro-3-fluorophenyl) cyanoguanidine

Following the general procedure for cyanoguanidine formation outlined in example 12, N-[4-chloro-2-tert-butyldimethylsilyloxy-3-(N'',N''-dimethylaminosulfonyl)phenyl]-N'-(2-chloro-3-fluorophenyl) carbodiimide (360 mg, 0.7 mmol), cyanamide (118 mg, 2.8 mmol) and N,N-diisopropylethylamine (109 mg, 0.84 mmol) were reacted, followed by desilylation with Cesium fluoride (128 mg, 0.84 mmol) to form the desired product (120 mg, 39%). LC-MS m/z 446.2 (M^+). 1H NMR ($DMSO-d_6$) δ 2.86 (s, 6H), 7.20 (d, 1H), 7.33 (t, 1H), 7.40 (m, 2H), 7.58 (d, 1H), 9.11 (s, 1H), 9.31 (s, 1H), 10.52 (s, 1H).

Example 21

- 25 Preparation of N-[4-chloro-2-hydroxy-3-(N'',N''-dimethylaminosulfonyl)phenyl]-N'-(2-chloro-4-fluorophenyl) cyanoguanidine

- a) 2-Chloro-4-fluorophenylisothiocyanate

Into a solution of 2-chloro-4-fluoroaniline (500 mg, 3.44 mmol) in a mixture of chloroform and water (10 mL/10 mL) at room temperature, thiophosgene (0.53 mL, 6.88 mmol) and sodium bicarbonate (1.09 g, 10.32 mmol) were added. The mixture was stirred at room temperature for 16 hours. The mixture was partitioned

between chloroform and water. The combined organic layer was then concentrated to give the desired product (586 mg, 91%). ¹ H NMR (CDCl₃) δ 6.98 (m, 1H), 7.20 (m, 2H).

- 5 b) N-[4-Chloro-2-hydroxy-3-(N'',N''-dimethylaminosulfonyl)phenyl]-N'-(2-chloro-4-fluorophenyl) thiourea

Following the general procedure for thiourea formation outlined in example 12, N,N-dimethyl-3-amino-6-chloro-2-hydroxybenzenesulfonamide (526 mg, 2.10 mmol) and 2-chloro-4-fluorophenylisothiocyanate (586 mg, 3.1 mmol) were coupled
10 to form the desired thiourea (505 mg, 55%). EI-MS m/z 437.75 (M⁺).

- c) N-(2-Chloro-4-fluorophenyl)-N'-[4-chloro-2-tert-butyldimethylsilyloxy-3-(N'',N''-dimethylaminosulfonyl)phenyl] thiourea

Following the general procedure for protected phenyl thiourea formation
15 outlined in example 12, N-[4-chloro-2-hydroxy-3-(N'',N''-dimethylaminosulfonyl)phenyl]-N'-(2-chloro-4-fluorophenyl)thiourea (470 mg, 1.08 mmol), tert-butyldimethylsilyl chloride (810 mg, 5.4 mmol) and imidazole (147 mg, 2.16 mmol) were reacted to form the desired product (420 mg, 71%). EI-MS m/z 551.75 (M⁺).

20

- d) N-(2-Chloro-4-fluorophenyl)-N'-[4-chloro-2-tert-butyldimethylsilyloxy-3-(N'',N''-dimethylaminosulfonyl)phenyl] carbodiimide

Following the general procedure for carbodiimide formation outlined in example 12, N-(2-chloro-4-fluorophenyl)-N'-[4-chloro-2-tert-butyldimethylsilyloxy-
25 3-(N'',N''-dimethylaminosulfonyl)phenyl]thiourea (420 mg, 0.76 mmol), methanesulfonyl chloride (0.11 mL, 1.52 mmol) and triethylamine (0.22 mL, 1.52 mmol) were reacted to form the desired product (420 mg, crude). ¹ H NMR (CDCl₃) δ 0.36 (s, 6H), 1.02 (s, 9H), 2.89 (s, 6H), 6.99 (m, 1H), 7.08 (d, 1H), 7.19 (m, 2H), 7.33 (d, 1H).

30

- e) Preparation of N-[4-chloro-2-hydroxy-3- (N'',N''-dimethylaminosulfonyl)phenyl]-N'-(2-chloro-4-fluorophenyl) cyanoguanidine

Following the general procedure for cyanoguanidine formation outlined in example 12, N-[4-chloro-2-tert-butyldimethylsilyloxy-3-(N",N"-dimethylaminosulfonyl)phenyl]-N'-(2-chloro-4-fluorophenyl) carbodiimide (420 mg, 0.81 mmol), cyanamide (136 mg, 3.24 mmol) and N,N-diisopropylethylamine (126 mg, 1.62 mmol) were reacted, followed by desilylation with Cesium fluoride (148 mg, 1.62 mmol) to form the desired product (180 mg, 50%). EI-MS m/z 446.2 (M⁺). ¹H NMR (CDCl₃) δ 2.86 (s, 6H), 7.16 (d, 1H), 7.24 (t, 1H), 7.46 (m, 1H), 7.55 (m, 2H), 8.94 (s, 1H), 9.15 (s, 1H), 10.50 (s, 1H).

Example 22

10 Preparation of N-(2-chloro-4-fluorophenyl)-N'-[4-chloro-2-hydroxy-3-(piperidone-4-ketone)aminosulfonylphenyl] cyanoguanidine

a) N-(2-Chloro-4-fluorophenyl)-N'-[4-chloro-2-hydroxy-3-(piperidone-4-ketone)aminosulfonylphenyl] thiourea

15 Following the general procedure for thiourea formation outlined in example 12, (N-piperidone-4-ketone)-2-hydroxy-3-amino-6-dichloro-benzenesulfonamide (732 mg, 2.4 mmol) and 2-chloro-4-fluorophenylisothiocyanate (see Example 46, 500 mg, 2.67 mmol) were coupled to form the desired thiourea (704 mg, 54%). EI-MS m/z 491.96 (M⁺).

20

b) N-(2-Chloro-4-fluorophenyl)-N'-[4-chloro-2-tert-butyldimethylsilyloxy-3-(piperidone-4-ketone)aminosulfonylphenyl] thiourea

Following the general procedure for protected phenyl thiourea formation outlined in example 12, N-(2-chloro-4-fluorophenyl)-N'-[4-chloro-2-hydroxy-3-(piperidone-4-ketone)aminosulfonylphenyl] thiourea (704 mg, 1.44 mmol), tert-butyldimethylsilyl chloride (1.08 g, 7.2 mmol) and imidazole (196 mg, 2.88 mmol) were reacted to form the desired product (340 mg, 39%). LC-MS m/z 606.2 (M⁺).

25 c) N-(2-Chloro-4-fluorophenyl)-N'-[4-chloro-2-tert-butyldimethylsilyloxy-3-(piperidone-4-ketone)aminosulfonylphenyl] carbodiimide

30 Following the general procedure for carbodiimide formation outlined in example 12, N-(2-chloro-4-fluorophenyl)-N'-[4-chloro-2-tert-butyldimethylsilyloxy-

3- (piperidone-4-ketone)aminosulfonylphenyl] thiourea (340 mg, 0.56 mmol), methanesulfonyl chloride (0.08 mL, 1.12 mmol) and triethylamine (0.16 mL, 1.12 mmol) were reacted to form the desired product (330 mg, crude). LC-MS m/z 572.2 (M^+).

5

d) N-(2-Chloro-4-fluorophenyl)-N'-[4-chloro-2-hydroxy-3-(piperidone-4-ketone)aminosulfonylphenyl] cyanoguanidine

Following the general procedure for cyanoguanidine formation outlined in example 12, N-(2-chloro-4-fluorophenyl)-N'-[4-chloro-2-tert-butyldimethylsilyloxy-3-(piperidone-4-ketone)aminosulfonylphenyl] carbodiimide (330 mg, 0.58 mmol), cyanamide (104 mg, 2.48 mmol) and N,N-diisopropylethylamine (96 mg, 0.74 mmol) were reacted, followed by desilylation with Cesium fluoride (113 mg, 0.74 mmol) to form the desired product (76 mg, 26%). LC-MS m/z 500.2. ^1H NMR (DMSO- d_6) δ 2.44 (t, 4H), 3.64 (s, 4H), 7.19 (d, 1H), 7.25 (t, 1H), 7.46 (m, 1H), 7.58 (d, 2H), 8.93 (s, 1H), 9.20 (s, 1H), 10.53 (s, 1H).

15

Example 23

Preparation of N-(2,3-dichlorophenyl)-N'-[4-chloro-2-hydroxy-3-(piperidone-4-ketone)aminosulfonylphenyl] cyanoguanidine

20 a) N-(2,3-Dichlorophenyl)-N'-[4-chloro-2-hydroxy-3-(piperidone-4-ketone)aminosulfonylphenyl] thiourea

Following the general procedure for thiourea formation outlined in example 12, (N-piperidone-4-ketone)-2-hydroxy-3-amino-6-dichloro-benzenesulfonamide (1.0 g, 3.3 mmol) and 2,3-dichlorophenylisothiocyanate (669 mg, 3.3 mmol) were coupled to form the desired thiourea (500 mg, 30%). LC-MS m/z 508.2.

25

b) N-(2,3-Dichlorophenyl)-N'-[4-chloro-2-tert-butyldimethylsilyloxy-3-(piperidone-4-ketone)aminosulfonylphenyl] thiourea

Following the general procedure for protected phenyl thiourea formation outlined in example 12, N-(2,3-dichlorophenyl)-N'-[4-chloro-2-hydroxy-3-(piperidone-4-ketone)aminosulfonylphenyl] thiourea (500 mg, 0.98 mmol), tert-butyldimethylsilyl chloride (765 mg, 4.9 mmol) and imidazole (140 mg, 1.98

30

mmol) were reacted to form the desired product (289 mg, 39%). LC-MS m/z 622.2 (M^+).

5 c) N-(2,3-Dichlorophenyl)-N'-[4-chloro-2-tert-butyldimethylsilyloxy-3-(piperidone-4-ketone)aminosulfonylphenyl] carbodiimide

Following the general procedure for carbodiimide formation outlined in example 12, N-(2,3-dichlorophenyl)-N'-[4-chloro-2-tert-butyldimethylsilyloxy-3-(piperidone-4-ketone)aminosulfonylphenyl] thiourea (289 mg, 0.46 mmol), methanesulfonyl chloride (0.07 mL, 0.92 mmol) and triethylamine (0.13 mL, 0.92 mmol) were reacted to form the desired product (280 mg, crude). 1H NMR ($CDCl_3$) δ 0.36 (s, 6H), 1.03 (s, 9H), 2.54 (t, 4H), 3.63 (t, 4H), 7.09 (d, 1H), 7.1-7.27(m, 2H), 7.32 (d, 1H), 7.38 (d, 1H).

15 d) N-(2,3-Dichlorophenyl)-N'-[4-chloro-2-hydroxy-3-(piperidone-4-ketone)aminosulfonylphenyl] cyanoguanidine

Following the general procedure for cyanoguanidine formation outlined in example 12, N-(2,3-dichlorophenyl)-N'-[4-chloro-2-tert-butyldimethylsilyloxy-3-(piperidone-4-ketone)aminosulfonylphenyl] carbodiimide (280 mg, 0.48 mmol), cyanamide (80.6 mg, 1.92 mmol) and N,N-diisopropylethylamine (75 mg, 0.57 mmol) were reacted, followed by desilylation with Cesium fluoride (88 mg, 0.57 mmol) to form the desired product (40 mg, 16%). LC-MS m/z 500.2. 1H NMR ($DMSO-d_6$) δ 2.44 (t, 4H), 3.64 (s, 4H), 7.13 (m, 1H), 7.41 (m, 2H), 7.54 (m, 2H), 7.68 (d, 1H), 9.07 (s, 1H), 9.4 (s, 1H), 10.56 (s, 1H).

Example 24

25 Preparation of N-(2-chloro-3-fluorophenyl)-N'-[4-chloro-2-hydroxy-3-(piperidone-4-ketone)aminosulfonylphenyl] cyanoguanidine

a) N-(2-Chloro-3-fluorophenyl)-N'-[4-chloro-2-hydroxy-3-(piperidone-4-ketone)aminosulfonylphenyl] thiourea

30 Following the general procedure for thiourea formation outlined in example 12, (N-piperidone-4-ketone)-2-hydroxy-3-amino-6-dichloro-benzenesulfonamide (908 mg, 3.3 mmol, not clean) and 2-chloro-3-fluorophenylisothiocyanate (See

Example 45, 620 mg, 3.3 mmol) were coupled to form the desired thiourea (350 mg, 24%). LC-MS m/z 492.2.

5 b) N-(2-Chloro-3-fluorophenyl)-N'-[4-chloro-2-tert-butyldimethylsilyloxy-3-(piperidone-4-ketone)aminosulfonylphenyl] thiourea

Following the general procedure for protected phenyl thiourea formation outlined in example 12, N-(2-chloro-3-fluorophenyl)-N'-[4-chloro-2-hydroxy-3-(piperidone-4-ketone)aminosulfonylphenyl] thiourea (350 mg, 0.71 mmol), tert-butyldimethylsilyl chloride (535 mg, 3.55 mmol) and imidazole (98 mg, 1.42
10 mmol) were reacted to form the desired product (280 mg, 65%). LC-MS m/z 606.2 (M⁺).

c) N-(2-Chloro-3-fluorophenyl)-N'-[4-chloro-2-tert-butyldimethylsilyloxy-3-(piperidone-4-ketone)aminosulfonylphenyl] carbodiimide
15 Following the general procedure for carbodiimide formation outlined in example 12, N-(2-chloro-3-fluorophenyl)-N'-[4-chloro-2-tert-butyldimethylsilyloxy-3-(piperidone-4-ketone)aminosulfonylphenyl] thiourea (280 mg, 0.46 mmol), methanesulfonyl chloride (0.07 mL, 0.92 mmol) and triethylamine (0.13 mL, 0.92 mmol) were reacted to form the desired product (280 mg, crude). ¹ H NMR (CDCl₃)
20 δ 0.36 (s, 6H), 1.03 (s, 9H), 2.54 (t, 4H), 3.63 (t, 4H), 7.01 (d, 1H), 7.11 (d, 1H), 7.24 (m, 2H), 7.36 (d, 1H).

d) N-(2-Chloro-3-fluorophenyl)-N'-[4-chloro-2-hydroxy-3-(piperidone-4-ketone)aminosulfonylphenyl] cyanoguanidine
25 Following the general procedure for cyanoguanidine formation outlined in example 12, N-(2-chloro-3-fluorophenyl)-N'-[4-chloro-2-tert-butyldimethylsilyloxy-3-(piperidone-4-ketone)aminosulfonylphenyl] carbodiimide (280 mg, 0.49 mmol), cyanamide (82.3 mg, 1.96 mmol) and N,N-diisopropylethylamine (75 mg, 0.57 mmol) were reacted, followed by desilylation with Cesium fluoride (88 mg, 0.57
30 mmol) to form the desired product (42 mg, 16%). LC-MS m/z 500.2. ¹H NMR (DMSO-d₆) δ 2.44 (t, 4H), 3.64 (s, 4H), 7.2 (m, 1H), 7.22 (t, 1H), 7.34 (m, 3H), 7.58 (d, 1H), 9.1 (s, 1H), 9.3 (s, 1H), 10.57 (s, 1H).

Example 25**Preparation of N-(2-bromo-3-fluorophenyl)-N'-[4-chloro-2-hydroxy-3-(piperidone-4-ketone)aminosulfonylphenyl] cyanoguanidine**

- 5 a) N-(2-Bromo-3-fluorophenyl)-N'-[4-chloro-2-hydroxy-3-(piperidone-4-ketone)aminosulfonylphenyl] thiourea

Following the general procedure for thiourea formation outlined in example 12, (N-piperidone-4-ketone)-2-hydroxy-3-amino-6-dichloro-benzenesulfonamide (1.5 g, 4.92 mmol, not clean) and 2-bromo-3-fluorophenylisothiocyanate (See
10 Example 16, 1.0 g, 4.92 mmol) were coupled to form the desired thiourea (550 mg, 24%). EL-MS m/z 535.96 (M).

- b) N-(2-Bromo-3-fluorophenyl)-N'-[4-chloro-2-tert-butyldimethylsilyloxy-3-(piperidone-4-ketone)aminosulfonylphenyl] thiourea

15 Following the general procedure for protected phenyl thiourea formation outlined in example 12, N-(2-bromo-3-fluorophenyl)-N'-[4-chloro-2-hydroxy-3-(piperidone-4-ketone)aminosulfonylphenyl] thiourea (550 mg, 1.03 mmol), tert-butyldimethylsilyl chloride (773 mg, 5.15 mmol) and imidazole (142 mg, 2.06 mmol) were reacted to form the desired product (256 mg, 38%). LC-MS m/z 649.96
20 (M).

- c) N-(2-bromo-3-fluorophenyl)-N'-[4-chloro-2-tert-butyldimethylsilyloxy-3-(piperidone-4-ketone)aminosulfonylphenyl] carbodiimide

Following the general procedure for carbodiimide formation outlined in
25 example 12, N-(2-bromo-3-fluorophenyl)-N'-[4-chloro-2-tert-butyldimethylsilyloxy-3-(piperidone-4-ketone)aminosulfonylphenyl] thiourea (256 mg, 0.4 mmol), methanesulfonyl chloride (0.07 mL, 0.92 mmol) and triethylamine (0.13 mL, 0.92 mmol) were reacted to form the desired product (256 mg, crude).). ¹ H NMR (CDCl₃) δ 0.36 (s, 6H), 1.03 (s, 9H), 2.54 (t, 4H), 3.63 (t, 4H), 6.96 (t, 1H), 7.0 (d,
30 1H), 7.10 (d, 1H), 7.24 (m, 1H), 7.39 (d, 1H).

d) N-(2-Bromo-3-fluorophenyl)-N'-[4-chloro-2-hydroxy-3-(piperidone-4-ketone)aminosulfonylphenyl] cyanoguanidine

Following the general procedure for cyanoguanidine formation outlined in example 12, N-(2-bromo-3-fluorophenyl)-N'-[4-chloro-2-tert-butyldimethylsilyloxy-3-(piperidone-4-ketone)aminosulfonylphenyl] carbodiimide (256 mg, 0.42 mmol),
5 cyanamide (82.3 mg, 1.96 mmol) and N,N-diisopropylethylamine (75 mg, 0.57 mmol) were reacted, followed by desilylation with Cesium fluoride (88 mg, 0.57 mmol) to form the desired product (48 mg, 21%). LC-MS m/z 544.2. ¹H NMR (DMSO-d₆) δ 2.44 (t, 4H), 3.64 (s, 4H), 7.10 (m, 1H), 7.28 (m, 2H), 7.42 (m, 1H),
10 7.59 (m, 1H), 9.02 (s, 1H), 9.53 (s, 1H), 10.5 (s, 1H).

Example 26

Preparation of N-(2-bromophenyl)-N'-[4-chloro-2-hydroxy-3-(homopiperazine-aminosulfonylphenyl)phenyl]cyanoguanidine

15 a) N-(3,4-Dichlorophenyl)-2,2-dimethyl-propionamide

3,4-dichloroaniline (150g) in TBME (1L) was cooled to 10-15 °C. 30% aq NaOH (141 g, 1.14 equiv) was added, and the solution stirred vigorously via overhead mechanical stirrer. Trimethylacetyl chloride ("PivCl", 126mL) was added at such a rate as to keep the internal temperature below 30 °C. During this addition,
20 the solution mixture becomes thick with white solid product. When the addition was complete (10-15 min), the mixture was heated to 30-35 °C for 1 hr, and then allowed to cool. The reaction mixture was held at -5 °C (overnight), and then filtered, rinsing first with 90:10 water/MeOH (600mL) and then water (900mL). Drying under vacuum yielded 195g (86%) product, as off-white crystals. LCMS m/z
25 246(M-H)⁺.

b) 2-tert-Butyl-6-chloro-benzooxazole-7-sulfonyl chloride

The solution of N-(3,4-dichloro-phenyl)-2,2-dimethyl-propionamide (10g, 41mmol) in dry THF (100mL) was cooled to -72 °C under argon. n-Butyl lithium
30 (1.6M in hexane, 64mL, 102mmol) was added dropwise. The solution warmed to ca. -50 °C over 45 minutes, and then was kept in the -25 - -10 °C range for 2 hrs.

The solution was then re-cooled to -78°C , and sulfur dioxide was bubbled through the solution for 30min. The solution was then allowed to warm to room temperature for 2h, and a Ar stream was bubbled through the solution, with a gas outlet provided so that any excess sulfur dioxide could escape during the warming. The THF solution was cooled in an ice bath, and sulfuryl chloride (3.58mL, 44.9mmol) was added dropwise. After a few minutes, the solution was warmed to room temperature for overnight. The mixture was concentrated, diluted with ethyl acetate and washed with water. Decolorizing carbon was added and the mixture was filtered. The resulting solution was dried (sodium sulfate), filtered and concentrated to afford the title compound (12.4g, 98%). ^1H NMR (CDCl_3) • 7.92 (d, 1H, $J=8.5$ Hz), 7.57 (d, 1H, $J=8.4$ Hz), 1.57 (s, 9H).

c) Homopiperazine-carboxylic acid tert-butyl ester

To a solution of homopiperazine (5.0 g, 49.92 mmol) in dichloromethane (100 mL), Di-tert-butyl dicarbonate (3.63 g, 16.64 mmol) and triethylamine (6.96 mL, 49.92 mmol) were added at room temperature. Then the reaction mixture was stirred at room temperature for 16 hours. The solid was filtered and the organic phase was washed with water (3x). The organic phase was dried (sodium sulfate), filtered and concentrated to afford the title compound (6.5 g, 66%). EI-MS m/z 197.79(M).

General procedure for the synthesis of sulfonylamides.

d) 2-tert-Butyl-6-chloro-7-(homopiperazine-1-sulfonyl)-benzooxazole

To a solution of 2-tert-butyl-6-chloro-benzooxazole-7-sulfonyl chloride (5.54, 18mmol) and triethylamine (2.51mL, 18mmol) in THF (100mL) at 0°C was added homopiperazine-carboxylic acid tert-butyl ester (3.0 g, 15mmol). The reaction was warmed to room temperature and allowed to stir overnight. The solution was concentrated and then diluted with water and extracted with ethyl acetate (3 times). The combined organic layers were dried with MgSO_4 , filtered, and concentrated. Flash chromatography (80% ethyl acetate/ 20% Ethanol) on silica gel gave the title compound (4.0g, 61%). LC-MS m/z 473.2.

General procedure for the hydrolysis of the benzooxazole to the desired aniline.

e) 6-Amino-3-chloro-2-(homopiperazine-1-sulfonyl)-phenol

To a solution of 2-tert-Butyl-6-chloro-7-(homopiperazine-1-sulfonyl)-benzooxazole (2.0g, 4.24mmol) in 1,4-dioxane (56mL) was treated with water (3.5mL) and conc. H₂SO₄ (3.5mL). The mixture was heated to 100 °C for 16h. The reaction was cooled to room temperature, and then basified to pH=14 with 25% aq NaOH. washed. The mixture was extracted with ethyl acetate (3 times), dried with MgSO₄, filtered, and concentrated to afford the title compound (1.04 g, 80%).
EI-MS m/z 306.2(M⁺).

General procedure for the protected amine group

f) 6-Amino-3-chloro-2-[(4-N-9-fluorenylmethylformate)homopiperazine-1-sulfonyl]-phenol

To a solution of 6-Amino-3-chloro-2-(homopiperazine-1-sulfonyl)-phenol (1.0 g, 3.27 mmol) in 1,4-dioxane (30 mL), 10% of sodium carbonate (7.5 mL) was added. Then the reaction mixture was cooled down to 0°C, 9-fluorenylmethyl chloroformate (800 mg, 3.27mmol) was added. After addition, the reaction mixture was stirred at 0°C for 1 hour. The solvent was evaporated. The residue was partitioned between ethyl acetate and 10% of sodium carbonate. The combined organic phase was dried with MgSO₄, filtered, and concentrated to afford the title compound (2.0 g, crude). EI-MS m/z 528.2(M⁺).

g) N-(2-Bromophenyl)-N'-[4-chloro-2-hydroxy-3-[(4-N-9-fluorenylmethylformate)homopiperazine aminosulfonylphenyl] thiourea

Following the general procedure for thiourea formation outlined in example 12, 6-Amino-3-chloro-2-[(4-N-9-fluorenylmethylformate)homopiperazine-1-sulfonyl]-phenol (2.0 g, 3.79 mmol, not clean) and 2-bromophenylisothiocyanate (811 mg, 3.79 mmol) were coupled to form the desired thiourea (2.0 g, 71%). EL-MS m/z 740.61 (M).

h) N-(2-Bromophenyl)-N'-[4-chloro-2-tert-butyldimethylsilyloxy-3-[(4-N-9-fluorenylmethylformate)homopiperazine aminosulfonylphenyl] thiourea

Following the general procedure for protected phenyl thiourea formation outlined in example 12, N-(2-bromophenyl)-N'-[4-chloro-2-hydroxy-3-[(4-N-9-fluorenylmethylformate)homopiperazine aminosulfonylphenyl] thiourea (1.0 g, 1.35 mmol), tert-butyldimethylsilyl chloride (1.02 g, 6.75 mmol) and imidazole (184 mg, 2.7 mmol) were reacted to form the desired product (836 mg, 73%). EI-MS m/z 854.97 (M).

10 i) N-(2-bromophenyl)-N'-[4-chloro-2-tert-butyldimethylsilyloxy-3-[(4-N-9-fluorenylmethylformate)homopiperazine aminosulfonylphenyl] carbodiimide

Following the general procedure for carbodiimide formation outlined in example 12, N-(2-bromophenyl)-N'-[4-chloro-2-tert-butyldimethylsilyloxy-3-[(4-N-9-fluorenylmethylformate)homopiperazine aminosulfonylphenyl] thiourea (836 mg, 0.98 mmol), methanesulfonyl chloride (0.15 mL, 1.96 mmol) and triethylamine (0.27 mL, 1.96 mmol) were reacted to form the desired product (772 mg, crude). ¹H NMR (CDCl₃) δ 0.40 (s, 6H), 1.06 (s, 9H), 1.58 (t, 1H), 1.68 (m, 1H), 1.92 (t, 4H), 3.03 (q, 1H), 3.10 (t, 1H), 3.36 (m, 3H), 3.44 (m, 2H), 4.23 (t, 1H), 4.64 (d, 2H), 7.07 (d, 1H), 7.33 (m, 8H), 7.60 (m, 3H), 7.73 (m, 2H).

20

j) N-(2-Bromophenyl)-N'-[4-chloro-2-hydroxy-3-(homopiperazine aminosulfonylphenyl] cyanoguanidine

Following the general procedure for cyanoguanidine formation outlined in example 12, N-(2-bromophenyl)-N'-[4-chloro-2-tert-butyldimethylsilyloxy-3-[(4-N-9-fluorenylmethylformate)homopiperazine aminosulfonylphenyl] carbodiimide (256 mg, 0.42 mmol), cyanamide (82.3 mg, 1.96 mmol) and N,N-diisopropylethylamine (75 mg, 0.57 mmol) were reacted, followed by desilylation with Cesium fluoride (88 mg, 0.57 mmol) and deprotected amino with 20% piperidine (8 mL) in THF (40mL) at room temperature for 30 minutes to form the desired product. The desired product (250 mg) was purified by Gilson HPLC to give pure product (50 mg, 20%). LC-MS m/z 527.2. ¹H NMR (DMSO-d₆) δ 1.68 (m, 2H), 3.06 (m, 2H), 3.2 (m, 2H), 3.34 (m,

30

2H), 3.71 (m, 2H), 6.25 (m, 1H), 7.19 (m, 1H), 7.4 (m, 3H), 7.71 (m, 1H), 8.76 (s, 1H).

Example 27

5 Preparation of N-(2-bromophenyl)-N'-[4-chloro-2-hydroxy -3-(N-methylhomopiperazine-aminosulfonylphenyl)phenyl]cyanoguanidine

To a solution of N-(2-bromophenyl)-N'-[4-chloro-2-hydroxy -3-(homopiperazine-aminosulfonylphenyl)phenyl]cyanoguanidine (287 mg, 0.54 mmol) in diglyme (10 mL), paraformaldehyde (35 mg, 1.08 mmol) and titanium isopropoxide (0.17 mL, 0.54 mmol) were added at room temperature. The reaction mixture was stirred at 60 °C for 1 hour and stirred at room temperature for 30 minutes. Then sodium borohydride (22 mg, 0.65 mmol) was added and heated to 60 °C for 4 hours. The reaction mixture was partitioned between ethyl acetate and water. The combined organic phase was dried with MgSO₄, filtered, and concentrated to afford the product. The product was purified by Gilson HPLC to give pure product (10 mg, 3%). LC-MS m/z 541.2. ¹H NMR (DMSO-d₆) δ 1.7 (m, 2H), 2.65 (s, 3H), 3.09 (m, 3H), 3.27 (m, 3H), 3.7 (m, 2H), 6.33 (d, 1H), 7.21 (t, 1H), 7.45 (m, 2H), 7.58 (d, 1H), 7.7 (d, 1H), 8.61 (s, 1H), 10.34 (s, 1H).

Example 28

20 Preparation of N-(2-bromophenyl)-N'-[4-chloro-2-hydroxy -3-(piperazine-aminosulfonylphenyl)phenyl]cyanoguanidine

a) 4-(2-tert-Butyl-6-chloro-benzooxazole-7-sulfonyl)-piperazine-1-carboxylic acid tert-butyl ester

25 Following the general procedure for the synthesis of sulfonylamides outlined in example 51, 2-tert-butyl-6-chloro-benzooxazole-7-sulfonyl chloride (5.0g, 16.2mmol), triethylamine (2.4mL, 17.2mmol), and piperazine-carboxylic acid tert-butyl ester (3.62g, 19.4mmol) were reacted in THF (50mL) to afford the title compound (5.44g, 67%). LCMS m/z 402(M-H)⁺ (desired -Boc).

30

b) 6-Amino-3-chloro-2-(piperazine-1-sulfonyl)-phenol

Following the general procedure for the hydrolysis of the benzooxazole to the desired aniline outlined in example 51, 4-(2-tert-butyl-6-chloro-benzooxazole-7-sulfonyl)-piperazine-1-carboxylic acid tert-butyl ester (2.0g, 4.3mmol), water (3.65mL), and H₂SO₄ (3.65mL) in 1,4-dioxane (60mL) were reacted to afford the
5 title compound (1.22g, 96%). LCMS m/z 292(M-H)⁺.

c) 6-Amino-3-chloro-2-[(4-N-9-fluorenylmethylformate)piperazine-1-sulfonyl]-phenol

Following the general procedure for protection of amino outlined in example
10 51,
6-Amino-3-chloro-2-(piperazine-1-sulfonyl)-phenol
(660 mg, 2.26 mmol), 10% of sodium carbonate (6.0 mL) and 9-fluorenylmethyl
chloroformate (584.7 mg, 2.26mmol) in 1,4-dioxane (6.78 mL) were reacted to
afford the title compound (1.22 g, crude). LC-MS m/z 514(M⁺).

15
d) N-(2-Bromophenyl)-N'-[4-chloro-2-hydroxy-3-[(4-N-9-fluorenylmethylformate)piperazine aminosulfonylphenyl]] thiourea

Following the general procedure for thiourea formation outlined in example
12, 6-Amino-3-chloro-2-[(4-N-9-fluorenylmethylformate)piperazine-1-sulfonyl]-
20 phenol (1.22 g, 2.4 mmol, not clean) and 2-bromophenylisothiocyanate (508 mg,
2.4 mmol) were coupled to form the desired thiourea (490 mg, 28%). EL-MS m/z
727.07 (M⁺).

e) N-(2-Bromophenyl)-N'-[4-chloro-2-tert-butyldimethylsilyloxy-3-[(4-N-9-fluorenylmethylformate)piperazine aminosulfonylphenyl]] thiourea
25

Following the general procedure for protected phenyl thiourea formation
outlined in example 12, N-(2-bromophenyl)-N'-[4-chloro-2-hydroxy-3-[(4-N-9-fluorenylmethylformate)piperazine aminosulfonylphenyl]] thiourea (490 mg, 0.67
mmol), tert-butyldimethylsilyl chloride (505 mg, 3.35 mmol) and imidazole (93 mg,
30 1.34 mmol) were reacted to form the desired product (407 mg, 73%). EI-MS m/z
841.08 (M⁺).

f) N-(2-Bromophenyl)-N'-[4-chloro-2-tert-butyldimethylsilyloxy-3-[(4-N-9-fluorenylmethylformate)piperazine aminosulfonylphenyl] carbodiimide

Following the general procedure for carbodiimide formation outlined in example 12, N-(2-bromophenyl)-N'-[4-chloro-2-tert-butyldimethylsilyloxy-3-[(4-N-9-fluorenylmethylformate)piperazine aminosulfonylphenyl] thiourea (640 mg, 0.76 mmol), methanesulfonyl chloride (0.13 mL, 1.52 mmol) and triethylamine (0.25 mL, 1.52 mmol) were reacted to form the desired product (640 mg, crude). ¹H NMR (CDCl₃) δ 0.4 (s, 6H), 1.1 (s, 9H), 3.2 (m, 4H), 3.48 (m, 4H), 4.22 (t, 1H), 4.51 (d, 2H), 7.07-7.44 (m, 9H), 7.54 (d, 2H), 7.6 (d, 1H), 7.74 (d, 2H).

g) N-(2-Bromophenyl)-N'-[4-chloro-2-hydroxy-3-(piperazine aminosulfonylphenyl] cyanoguanidine

Following the general procedure for cyanoguanidine formation outlined in example 12, N-(2-bromophenyl)-N'-[4-chloro-2-tert-butyldimethylsilyloxy-3-[(4-N-9-fluorenylmethylformate)piperazine aminosulfonylphenyl] carbodiimide (640 mg, 0.79 mmol), cyanamide (133 mg, 3.16 mmol) and N,N-diisopropylethylamine (122 mg, 0.95 mmol) were reacted, followed by desilylation with Cesium fluoride (144 mg, 0.95 mmol) and deprotected amino with 20% piperidine (8 mL) in THF (40mL) at room temperature for 30 minutes to form the desired product (purified by Gilson HPLC, 346 mg, 76%). LC-MS m/z 513.2. ¹H NMR (DMSO-d₆) δ 3.02 (t, 4H), 3.43 (t, 4H), 6.08 (d, 1H), 7.2 (m, 1H), 7.38 (m, 3H), 7.67 (d, 1H), 8.84 (s, 1H).

Example 29

Preparation of N-(2-chloro-3-fluorophenyl)-N'-[4-chloro-2-hydroxy-3-(piperazine-aminosulfonylphenyl)phenyl]cyanoguanidine

a) N-(2-Chloro-3-fluorophenyl)-N'-[4-chloro-2-hydroxy-3-[(4-N-9-fluorenylmethylformate)piperazine aminosulfonylphenyl] thiourea

Following the general procedure for thiourea formation outlined in example 12, 6-Amino-3-chloro-2-[(4-N-9-fluorenylmethylformate)piperazine-1-sulfonyl]-phenol (1.0 g, 1.94 mmol, not clean) and 2-chloro-3-fluorophenylisothiocyanate (See Example 45, 400 mg, 1.94 mmol) were coupled to form the desired thiourea (713 mg, 48%). EL-MS m/z 700.70 (M).

b) N-(2-Chloro-3-fluorophenyl)-N'-[4-chloro-2-tert-butyldimethylsilyloxy-3-[(4-N-9-fluorenylmethylformate)piperazine aminosulfonylphenyl] thiourea

Following the general procedure for protected phenyl thiourea formation outlined in example 12, N-(2-chloro-3-fluorophenyl)-N'-[4-chloro-2-hydroxy-3-[(4-N-9-fluorenylmethylformate)piperazine aminosulfonylphenyl] thiourea (713 mg, 1.02 mmol), tert-butyldimethylsilyl chloride (765 mg, 5.1 mmol) and imidazole (139 mg, 2.04 mmol) were reacted to form the desired product (455 mg, 55%). EI-MS m/z 814.68 (M).

10

c) N-(2-Chloro-3-fluorophenyl)-N'-[4-chloro-2-tert-butyldimethylsilyloxy-3-[(4-N-9-fluorenylmethylformate)piperazine aminosulfonylphenyl] carbodiimide

Following the general procedure for carbodiimide formation outlined in example 12, N-(2-chloro-3-fluorophenyl)-N'-[4-chloro-2-tert-butyldimethylsilyloxy-3-[(4-N-9-fluorenylmethylformate)piperazine aminosulfonylphenyl] thiourea (455 mg, 0.56 mmol), methanesulfonyl chloride (0.1 mL, 1.12 mmol) and triethylamine (0.18 mL, 1.12 mmol) were reacted to form the desired product (537 mg, crude). ¹H NMR (CDCl₃) δ 0.39 (s, 6H), 1.04 (s, 9H), 3.19 (m, 4H), 3.49 (m, 4H), 4.23 (t, 1H), 4.5 (d, 1H), 6.96 (t, 1H), 7.01-7.41 (m, 9H), 7.55 (d, 1H), 7.74 (d, 1H).

20

d) N-(2-Chloro-3-fluorophenyl)-N'-[4-chloro-2-hydroxy-3-(piperazine aminosulfonylphenyl] cyanoguanidine

Following the general procedure for cyanoguanidine formation outlined in example 12, N-(2-chloro-3-fluorophenyl)-N'-[4-chloro-2-tert-butyldimethylsilyloxy-3-[(4-N-9-fluorenylmethylformate)piperazine aminosulfonylphenyl] carbodiimide (537 mg, 0.69 mmol), cyanamide (116 mg, 2.76 mmol) and N,N-diisopropylethylamine (107 mg, 0.83 mmol) were reacted, followed by desilylation with Cesium fluoride (126 mg, 0.83 mmol) and deprotected amino with 20% piperidine (6 mL) in THF (30mL) at room temperature for 30 minutes to form the desired product (purified by Gilson HPLC, 45 mg, 14%). LC-MS m/z 487.0. ¹H NMR (DMSO-d₆) δ 3.02 (t, 4H), 3.43 (t, 4H), 6.2 (d, 1H), 7.2-7.39 (m, 4H), 9.03 (s, 1H).

30

Example 30Preparation of N-(2-bromophenyl)-N'-[4-chloro-2-hydroxy-3-(4-amino-piperidine aminosulfonylphenyl)phenyl]cyanoguanidine

- 5 a) [1-(2-tert-Butyl-6-chloro-benzooxazole-7-sulfonyl)-piperidin-4-yl]-carbamic acid tert-butyl ester

Following the general procedure for the synthesis of sulfonylamides outlined in example 51, 2-tert-butyl-6-chloro-benzooxazole-7-sulfonyl chloride (5.05g, 16.4mmol), triethylamine (4.57mL, 32.8mmol), and 4-N-Boc-aminopiperidine
10 (3.288g, 16.4mmol) were reacted in THF (125mL) to afford the title compound (4.18g, 54%). ¹H NMR (DMSO-d₆) • 7.98 (d, 1H, J=8.48 Hz), 7.63 (d, 1H, J=8.47 Hz), 3.73 (d, 2H), 3.35 (bs, 2H), 2.92 (m, 2H), 1.75 (d, 2H), 1.35 (s, 10H).

- b) 6-Amino-2-(4-amino-piperidine-1-sulfonyl)-3-chloro-phenol

15 Following the general procedure for the hydrolysis of the benzooxazole to the desired aniline outlined in example 51, [1-(2-tert-butyl-6-chloro-benzooxazole-7-sulfonyl)-piperidin-4-yl]-carbamic acid tert-butyl ester (4.18g, 8.86mmol), water (5.5mL), and H₂SO₄ (5.5mL) in 1,4-dioxane (55mL) were reacted to afford the title compound (2.03g, 75%). LCMS m/z 306(M-H)⁺.

20

- c) 6-Amino-3-chloro-2-[(4-N-9-fluorenylmethylformate)piperidin-1-sulfonyl]-phenol

Following the general procedure for protection of amino outlined in example 51, 6-Amino-3-chloro-2-(piperidin-1-sulfonyl)-phenol
25 (1.05 g, 3.43 mmol), 10% of sodium carbonate (9.1 mL) and 9-fluorenylmethyl chloroformate (887.4 mg, 3.43mmol) in 1,4-dioxane (10.3 mL) were reacted to afford the title compound (1.3 g, crude). LC-MS m/z 528.04(M⁺).

- d) N-(2-Bromophenyl)-N'-[4-chloro-2-hydroxy-3-[(4-N-9-fluorenylmethylformate) piperidine aminosulfonylphenyl] thiourea
30

Following the general procedure for thiourea formation outlined in example 12, 6-Amino-3-chloro-2-[(4-N-9-fluorenylmethylformate)piperidine-1-sulfonyl]-

phenol (1.3 g, 2.42 mmol, not clean) and 2-bromophenylisothiocyanate (520 mg, 2.42 mmol) were coupled to form the desired thiourea (700 mg, 39%). EL-MS m/z 739.07 (M).

- 5 e) N-(2-Bromophenyl)-N'-[4-chloro-2-tert-butyldimethylsilyloxy-3-[(4-N-9-fluorenylmethylformate)piperidine aminosulfonylphenyl] thiourea

Following the general procedure for protected phenyl thiourea formation outlined in example 12, N-(2-bromophenyl)-N'-[4-chloro-2-hydroxy-3-[(4-N-9-fluorenylmethylformate)piperazine aminosulfonylphenyl] thiourea (700 mg, 0.95
10 mmol), tert-butyldimethylsilyl chloride (713 mg, 4.75 mmol) and imidazole (131 mg, 1.9 mmol) were reacted to form the desired product (400 mg, 50%). EI-MS m/z 854.52 (M).

- 15 f) N-(2-Bromophenyl)-N'-[4-chloro-2-tert-butyldimethylsilyloxy-3-[(4-N-9-fluorenylmethylformate)piperidine aminosulfonylphenyl] carbodiimide

Following the general procedure for carbodiimide formation outlined in example 12, N-(2-bromophenyl)-N'-[4-chloro-2-tert-butyldimethylsilyloxy-3-[(4-N-9-fluorenylmethylformate)piperidine aminosulfonylphenyl] thiourea (374 mg, 0.57 mmol), methanesulfonyl chloride (0.1 mL, 1.12 mmol) and triethylamine (0.18 mL,
20 1.12 mmol) were reacted to form the desired product (374 mg, crude). ¹H NMR (CDCl₃) δ 0.38 (s, 6H), 1.03 (s, 9H), 1.49 (m, 2H), 1.93 (m, 2H), 2.89 (t, 2H), 3.72 (m, 2H), 4.2 (t, 1H), 4.4 (d, 2H), 4.73 (d, 1H), 7.06-7.34 (m, 8H), 7.41 (t, 2H), 7.61 (t, 2H), 7.24 (m, 1H), 7.76 (d, 2H).

- 25 g) N-(2-Bromophenyl)-N'-[4-chloro-2-hydroxy-3-(piperidine aminosulfonyl)phenyl] cyanoguanidine

Following the general procedure for cyanoguanidine formation outlined in example 12, N-(2-bromophenyl)-N'-[4-chloro-2-tert-butyldimethylsilyloxy-3-[(4-N-9-fluorenylmethylformate)piperidine aminosulfonylphenyl] carbodiimide (400 mg,
30 0.65 mmol), cyanamide (108 mg, 2.6 mmol) and N,N-diisopropylethylamine (99 mg, 0.78 mmol) were reacted, followed by desilylation with Cesium fluoride (123 mg, 0.78 mmol) and deprotected amino with 20%piperidine (4 mL) in THF (20mL)

at room temperature for 30 minutes to form the desired product (purified by Gilson HPLC, 105 mg, 41%). LC-MS m/z 527.2. ¹H NMR (DMSO-d₆) δ 1.43 (m, 2H), 1.83 (m, 2H), 2.9 (t, 2H), 2.95 (m, 2H), 3.66 (d, 2H), 6.09 (d, 1H), 7.14 (t, 1H), 7.34 (m, 3H), 7.64 (d, 1H).

5

Example 31

Preparation of N-(2-bromophenyl)-N'-{4-chloro-2-hydroxy-3-[(R)-3-amino-pyrrolidine] aminosulfonylphenyl}phenyl}cyanoguanidine

10 a) [(R)-1-(2-tert-Butyl-6-chloro-benzooxazole-7-sulfonyl)-pyrrolidin-3-yl]-carbamic acid tert-butyl ester

Following the general procedure for the synthesis of sulfonylamides outlined in example 51, 2-tert-butyl-6-chloro-benzooxazole-7-sulfonyl chloride (3.0g, 9.74mmol), triethylamine (1.63mL, 11.7mmol), and (R)-pyrrolidin-3-yl-carbamic acid tert-butyl ester (2.18g, 11.7mmol) were reacted in THF (30mL) to afford the
15 title compound (3.0g, 67%). ¹H NMR (CDCl₃) • 7.77 (d, 1H, J=8.5 Hz), 7.48 (d, 1H, J=8.5 Hz), 4.67 (bm, 1H), 4.22 (bm, 1H), 3.66 (bm, 2H), 3.52 (bm, 1H), 3.41 (bm, 1H), 2.19 (bm, 1H), 1.90 (bm, 1H), 1.48 (s, 9H).

20 b) 6-Amino-2-((R)-3-amino-pyrrolidine-1-sulfonyl)-3-chloro-phenol

Following the general procedure for the hydrolysis of the benzooxazole to the desired aniline outlined in example 51, [(R)-1-(2-tert-butyl-6-chloro-benzooxazole-7-sulfonyl)-pyrrolidin-3-yl]-carbamic acid tert-butyl ester (2.0g, 4.31mmol), water (3.6mL), and H₂SO₄ (3.6mL) in 1,4-dioxane (60mL) were reacted to afford the title compound (1.2g, 94%). LCMS m/z 292(M-H)⁺.

25

c) 6-Amino-3-chloro-2-[(R)-3-N-9-fluorenylmethylformate) pyrrolidine -1-sulfonyl]-phenol

Following the general procedure for protection of amino outlined in example 51, 6-Amino-2-((R)-3-amino-pyrrolidine-1-sulfonyl)-3-chloro-phenol

(907 mg, 3.1 mmol), 10% of sodium carbonate (7.5 mL) and 9-fluorenylmethyl chloroformate (802 mg, 3.1 mmol) in 1,4-dioxane (20 mL) were reacted to afford the title compound (1.7 g, crude). LC-MS m/z 514.2 (M^+).

- 5 d) N-(2-Bromophenyl)-N'-{4-chloro-2-hydroxy-3-[(R)-3-N-9-fluorenylmethylformate) pyrrolidine] aminosulfonylphenyl} thiourea

Following the general procedure for thiourea formation outlined in example 12, 6-Amino-3-chloro-2-[(R)-3-N-9-fluorenylmethylformate) pyrrolidine]-1-sulfonyl]-phenol (800 mg, 1.56 mmol, not clean) and 2-bromophenylisothiocyanate
10 (333 mg, 1.56 mmol) were coupled to form the desired thiourea (405 mg, 36%). EL-MS m/z 727.01 (M).

- e) N-(2-Bromophenyl)-N'-{4-chloro-2-tert-butyldimethylsilyloxy-3-[(R)-3-N-9-fluorenylmethylformate) pyrrolidine] aminosulfonylphenyl} thiourea

15 Following the general procedure for protected phenyl thiourea formation outlined in example 12, N-(2-bromophenyl)-N'-{4-chloro-2-hydroxy-3-[(R)-3-N-9-fluorenylmethylformate) pyrrolidine] aminosulfonylphenyl} thiourea (405 mg, 0.56 mmol), tert-butyldimethylsilyl chloride (420 mg, 2.8 mmol) and imidazole (76 mg, 1.12 mmol) were reacted to form the desired product (293 mg, 63%). EI-MS m/z
20 841.2 (M^+).

- f) N-(2-Bromophenyl)-N'-{4-chloro-2-tert-butyldimethylsilyloxy-3-[(R)-3-N-9-fluorenylmethylformate) pyrrolidine] aminosulfonylphenyl} carbodiimide

Following the general procedure for carbodiimide formation outlined in
25 example 12, N-(2-bromophenyl)-N'-{4-chloro-2-tert-butyldimethylsilyloxy-3-[(R)-3-N-9-fluorenylmethylformate) pyrrolidine] aminosulfonylphenyl} thiourea (293 mg, 0.35 mmol), methanesulfonyl chloride (0.05 mL, 0.7 mmol) and triethylamine (0.1 mL, 0.7 mmol) were reacted to form the desired product (293 mg, crude). 1H NMR ($CDCl_3$) δ 0.39 (s, 6H), 1.05 (s, 9H), 1.90 (m, 1H), 2.18 (m, 1H), 3.14 (m, 1H), 3.39
30 (m, 3H), 3.48 (m, 1H), 4.2 (t, 1H), 4.4 (d, 2H), 5.06 (d, 1H), 7.16-7.43 (m, 10H), 7.59 (d, 1H), 7.7 (d, 1H).

g) N-(2-Bromophenyl)-N'-{4-chloro-2-hydroxy-3-[(R)-3-aminopyrrolidine]aminosulfonylphenyl}cyanoguanidine

Following the general procedure for cyanoguanidine formation outlined in example 12, N-(2-bromophenyl)-N'-{4-chloro-2-tert-butyldimethylsilyloxy-3-[(R)-3-N-9-fluorenylmethylformate) pyrrolidine]aminosulfonylphenyl} carbodiimide (293 mg, 0.36 mmol), cyanamide (61 mg, 1.44 mmol) and N,N-diisopropylethylamine (56 mg, 0.43 mmol) were reacted, followed by desilylation with Cesium fluoride (66 mg, 0.43 mmol) and deprotected amino with 20%piperidine (4 mL) in THF (20mL) at room temperature for 30 minutes to form the desired product (purified by Gilson HPLC, 53mg, 25%). LC-MS m/z 513.2. ¹H NMR (DMSO-d₆) δ 1.6 (m, 2H), 2.0 (m, 1H), 2.2 (m, 1H), 2.95 (t, 1H), 3.35 (m, 1H), 3.50 (m, 2H), 3.72 (m, 1H), 6.9 (d, 1H), 7.2 (t, 1H), 7.42 (m, 2H), 7.7 (t, 2H), 8.32 (s, 1H).

METHOD OF TREATMENT

The compounds of Formula (I), or a pharmaceutically acceptable salt thereof can be used in the manufacture of a medicine for the prophylactic or therapeutic treatment of any disease state in a human, or other mammal, which is exacerbated or caused by excessive or unregulated IL-8 cytokine production by such mammal's cell, such as but not limited to monocytes and/or macrophages, or other chemokines which bind to the IL-8 α or β receptor, also referred to as the type I or type II receptor.

Accordingly, the present invention provides a method of treating a chemokine mediated disease, wherein the chemokine is one which binds to an IL-8 α or β receptor and which method comprises administering an effective amount of a compound of Formula (I) or a pharmaceutically acceptable salt thereof. In particular, the chemokines are IL-8, GROα, GROβ, GROγ, NAP-2 or ENA-78.

The compounds of Formula (I) are administered in an amount sufficient to inhibit cytokine function, in particular IL-8, GROα, GROβ, GROγ, NAP-2 or ENA-78, such that they are biologically regulated down to normal levels of physiological function, or in some case to subnormal levels, so as to ameliorate the disease state. Abnormal levels of IL-8, GROα, GROβ, GROγ, NAP-2 or ENA-78

for instance in the context of the present invention, constitute: (i) levels of free IL-8 greater than or equal to 1 picogram per mL; (ii) any cell associated IL-8, GRO α , GRO β , GRO γ , NAP-2 or ENA-78 above normal physiological levels; or (iii) the presence of IL-8, GRO α , GRO β , GRO γ , NAP-2 or ENA-78 above basal levels in
5 cells or tissues in which IL-8, GRO α , GRO β , GRO γ , NAP-2 or ENA-78 respectively, is produced.

The compounds of Formula (I), in generally have been shown to have a longer $t_{1/2}$ and improved oral bioavailability over the compounds disclosed in WO 96/25157 and WO 97/29743 whose disclosures are incorporated herein by reference.

10 There are many disease states in which excessive or unregulated IL-8 production is implicated in exacerbating and/or causing the disease. Chemokine mediated diseases include psoriasis, atopic dermatitis, osteo arthritis, rheumatoid arthritis, asthma, chronic obstructive pulmonary disease, adult respiratory distress syndrome, inflammatory bowel disease, Crohn's disease, ulcerative colitis, stroke,
15 septic shock, multiple sclerosis, endotoxic shock, gram negative sepsis, toxic shock syndrome, cardiac and renal reperfusion injury, glomerulonephritis, thrombosis, graft vs. host reaction, Alzheimer's disease, allograft rejections, malaria, restenosis, angiogenesis, atherosclerosis, osteoporosis, gingivitis and undesired hematopoietic stem cells release and diseases caused by respiratory viruses, herpesviruses, and
20 hepatitis viruses, meningitis, cystic fibrosis, pre-term labor, cough, pruritus, multi-organ dysfunction, trauma, strains, sprains, contusions, psoriatic arthritis, herpes, encephalitis, CNS vasculitis, traumatic brain injury, CNS tumors, subarachnoid hemorrhage, post surgical trauma, interstitial pneumonitis, hypersensitivity, crystal induced arthritis, acute and chronic pancreatitis, acute alcoholic hepatitis,
25 necrotizing enterocolitis, chronic sinusitis, uveitis, polymyositis, vasculitis, acne, gastric and duodenal ulcers, celiac disease, esophagitis, glossitis, airflow obstruction, airway hyperresponsiveness, bronchiolitis obliterans organizing pneumonia, bronchiectasis, bronchiolitis, bronchiolitis obliterans, chronic bronchitis, cor pulmonae, dyspnea, emphysema, hypercapnea, hyperinflation, hypoxemia,
30 hyperoxia-induced inflammations, hypoxia, surgical lung volume reduction, pulmonary fibrosis, pulmonary hypertension, right ventricular hypertrophy,

sarcoidosis, small airway disease, ventilation-perfusion mismatching, wheeze, colds and lupus.

These diseases are primarily characterized by massive neutrophil infiltration, T-cell infiltration, or neovascular growth, and are associated with increased IL-8, GRO α , GRO β , GRO γ , NAP-2 or ENA-78 production which is responsible for the chemotaxis of neutrophils into the inflammatory site or the directional growth of endothelial cells. In contrast to other inflammatory cytokines (IL-1, TNF, and IL-6), IL-8, GRO α , GRO β , GRO γ , NAP-2 or ENA-78 have the unique property of promoting neutrophil chemotaxis, enzyme release including but not limited to elastase release as well as superoxide production and activation. The α -chemokines but particularly, GRO α , GRO β , GRO γ , NAP-2 or ENA-78, working through the IL-8 type I or II receptor can promote the neovascularization of tumors by promoting the directional growth of endothelial cells. Therefore, the inhibition of IL-8 induced chemotaxis or activation would lead to a direct reduction in the neutrophil infiltration.

Recent evidence also implicates the role of chemokines in the treatment of HIV infections, Littleman *et al.*, Nature 381, pp. 661 (1996) and Koup *et al.*, Nature 381, pp. 667 (1996).

Present evidence also indicates the use of IL-8 inhibitors in the treatment of atherosclerosis. The first reference, Boisvert *et al.*, J. Clin. Invest., 1998, 101:353-363 shows, through bone marrow transplantation, that the absence of IL-8 receptors on stem cells (and, therefore, on monocytes/macrophages) leads to a reduction in the development of atherosclerotic plaques in LDL receptor deficient mice. Additional supporting references are: Apostolopoulos, *et al.*, Arterioscler. Thromb. Vasc. Biol., 1996, 16:1007-1012; Liu, *et al.*, Arterioscler. Thromb. Vasc. Biol., 1997, 17:317-323; Rus, *et al.*, Atherosclerosis, 1996, 127:263-271.; Wang *et al.*, J. Biol. Chem., 1996, 271:8837-8842; Yue, *et al.*, Eur. J. Pharmacol., 1993, 240:81-84; Koch, *et al.*, Am. J. Pathol., 1993, 142:1423-1431.; Lee, *et al.*, Immunol. Lett., 1996, 53, 109-113.; and Terkeltaub *et al.*, Arterioscler. Thromb., 1994, 14:47-53.

The present invention also provides for a means of treating, in an acute setting, as well as preventing, in those individuals deemed susceptible to, CNS injuries by the chemokine receptor antagonist compounds of Formula (I).

CNS injuries as defined herein include both open or penetrating head trauma, such as by surgery, or a closed head trauma injury, such as by an injury to the head region. Also included within this definition is ischemic stroke, particularly to the brain area.

5 Ischemic stroke may be defined as a focal neurologic disorder that results from insufficient blood supply to a particular brain area, usually as a consequence of an embolus, thrombi, or local atheromatous closure of the blood vessel. The role of inflammatory cytokines in this area has been emerging and the present invention provides a mean for the potential treatment of these injuries. Relatively little
10 treatment, for an acute injury such as these has been available.

TNF- α is a cytokine with proinflammatory actions, including endothelial leukocyte adhesion molecule expression. Leukocytes infiltrate into ischemic brain lesions and hence compounds, which inhibit or decrease levels of TNF would be useful for treatment of ischemic brain injury. See Liu *et al.*, Stroke, Vol. 25., No. 7,
15 pp. 1481-88 (1994) whose disclosure is incorporated herein by reference.

Models of closed head injuries and treatment with mixed 5-LO/CO agents is discussed in Shohami *et al.*, J. of Vasc & Clinical Physiology and Pharmacology, Vol. 3, No. 2, pp. 99-107 (1992) whose disclosure is incorporated herein by reference. Treatment, which reduced edema formation, was found to improve
20 functional outcome in those animals treated.

The compounds of Formula (I) are administered in an amount sufficient to inhibit IL-8, binding to the IL-8 alpha or beta receptors, from binding to these receptors, such as evidenced by a reduction in neutrophil chemotaxis and activation. The discovery that the compounds of Formula (I) are inhibitors of IL-8 binding is
25 based upon the effects of the compounds of Formulas (I) in the *in vitro* receptor binding assays which are described herein. The compounds of Formula (I) have been shown to be inhibitors of type II IL-8 receptors.

As used herein, the term "IL-8 mediated disease or disease state" refers to any and all disease states in which IL-8, GRO α , GRO β , GRO γ , NAP-2 or ENA-78
30 plays a role, either by production of IL-8, GRO α , GRO β , GRO γ , NAP-2 or ENA-78 themselves, or by IL-8, GRO α , GRO β , GRO γ , NAP-2 or ENA-78 causing another monokine to be released, such as but not limited to IL-1, IL-6 or TNF. A disease

state in which, for instance, IL-1 is a major component, and whose production or action, is exacerbated or secreted in response to IL-8, would therefore be considered a disease state mediated by IL-8.

As used herein, the term "chemokine mediated disease or disease state" refers to any and all disease states in which a chemokine which binds to an IL-8 α or β receptor plays a role, such as but not limited to IL-8, GRO- α , GRO- β , GRO γ , NAP-2 or ENA-78. This would include a disease state in which, IL-8 plays a role, either by production of IL-8 itself, or by IL-8 causing another monokine to be released, such as but not limited to IL-1, IL-6 or TNF. A disease state in which, for instance, IL-1 is a major component, and whose production or action, is exacerbated or secreted in response to IL-8, would therefore be considered a disease state mediated by IL-8.

As used herein, the term "cytokine" refers to any secreted polypeptide that affects the functions of cells and is a molecule, which modulates interactions between cells in the immune, inflammatory or hematopoietic response. A cytokine includes, but is not limited to, monokines and lymphokines, regardless of which cells produce them. For instance, a monokine is generally referred to as being produced and secreted by a mononuclear cell, such as a macrophage and/or monocyte. Many other cells however also produce monokines, such as natural killer cells, fibroblasts, basophils, neutrophils, endothelial cells, brain astrocytes, bone marrow stromal cells, epidermal keratinocytes and B-lymphocytes. Lymphokines are generally referred to as being produced by lymphocyte cells. Examples of cytokines include, but are not limited to, Interleukin-1 (IL-1), Interleukin-6 (IL-6), Interleukin-8 (IL-8), Tumor Necrosis Factor-alpha (TNF- α) and Tumor Necrosis Factor beta (TNF- β).

As used herein, the term "chemokine" refers to any secreted polypeptide that affects the functions of cells and is a molecule which modulates interactions between cells in the immune, inflammatory or hematopoietic response, similar to the term "cytokine" above. A chemokine is primarily secreted through cell transmembranes and causes chemotaxis and activation of specific white blood cells and leukocytes, neutrophils, monocytes, macrophages, T-cells, B-cells, endothelial cells and smooth

muscle cells. Examples of chemokines include, but are not limited to IL-8, GRO- α , GRO- β , GRO- γ , NAP-2, ENA-78, IP-10, MIP-1 α , MIP- β , PF4, and MCP 1, 2, and 3.

The present compounds are also useful in normalizing leukocyte counts as well as normalizing levels of circulating chemokines.

5 In order to use a compound of Formula (I) or a pharmaceutically acceptable salt thereof in therapy, it will normally be formulated into a pharmaceutical composition in accordance with standard pharmaceutical practice. This invention, therefore, also relates to a pharmaceutical composition comprising an effective, non-toxic amount of a compound of Formula (I) and a pharmaceutically acceptable
10 carrier or diluent.

Compounds of Formula (I), pharmaceutically acceptable salts thereof and pharmaceutical compositions incorporating such may conveniently be administered by any of the routes conventionally used for drug administration, for instance, orally, topically, parenterally or by inhalation. The compounds of Formula (I) may be
15 administered in conventional dosage forms prepared by combining a compound of Formula (I) with standard pharmaceutical carriers according to conventional procedures. The compounds of Formula (I) may also be administered in conventional dosages in combination with a known, second therapeutically active compound. These procedures may involve mixing, granulating and compressing or
20 dissolving the ingredients as appropriate to the desired preparation. It will be appreciated that the form and character of the pharmaceutically acceptable character or diluent is dictated by the amount of active ingredient with which it is to be combined, the route of administration and other well-known variables. The carrier(s) must be "acceptable" in the sense of being compatible with the other
25 ingredients of the formulation and not deleterious to the recipient thereof.

The pharmaceutical carrier employed may be, for example, either a solid or liquid. Exemplary of solid carriers are lactose, terra alba, sucrose, talc, gelatin, agar, pectin, acacia, magnesium stearate, stearic acid and the like. Exemplary of liquid carriers are syrup, peanut oil, olive oil, water and the like. Similarly, the carrier or
30 diluent may include time delay material well known to the art, such as glyceryl mono-stearate or glyceryl distearate alone or with a wax.

A wide variety of pharmaceutical forms can be employed. Thus, if a solid carrier is used, the preparation can be tableted, placed in a hard gelatin capsule in powder or pellet form or in the form of a troche or lozenge. The amount of solid carrier will vary widely but preferably will be from about 25mg to about 1g. When a liquid carrier is used, the preparation will be in the form of a syrup, emulsion, soft gelatin capsule, sterile injectable liquid such as an ampule or nonaqueous liquid suspension.

Compounds of Formula (I) may be administered topically, that is by non-systemic administration. This includes the application of a compound of Formula (I) externally to the epidermis or the buccal cavity and the instillation of such a compound into the ear, eye and nose, such that the compound does not significantly enter the blood stream. In contrast, systemic administration refers to oral, intravenous, intraperitoneal and intramuscular administration.

Formulations suitable for topical administration include liquid or semi-liquid preparations suitable for penetration through the skin to the site of inflammation such as liniments, lotions, creams, ointments or pastes, and drops suitable for administration to the eye, ear or nose. The active ingredient may comprise, for topical administration, from 0.001% to 10% w/w, for instance from 1% to 2% by weight of the Formulation. It may however comprise as much as 10% w/w but preferably will comprise less than 5% w/w, more preferably from 0.1% to 1% w/w of the Formulation.

Lotions according to the present invention include those suitable for application to the skin or eye. An eye lotion may comprise a sterile aqueous solution optionally containing a bactericide and may be prepared by methods similar to those for the preparation of drops. Lotions or liniments for application to the skin may also include an agent to hasten drying and to cool the skin, such as an alcohol or acetone, and/or a moisturizer such as glycerol or an oil such as castor oil or arachis oil.

Creams, ointments or pastes according to the present invention are semi-solid formulations of the active ingredient for external application. They may be made by mixing the active ingredient in finely divided or powdered form, alone or in solution or suspension in an aqueous or non-aqueous fluid, with the aid of suitable machinery, with a greasy or non-greasy base. The base may comprise hydrocarbons

such as hard, soft or liquid paraffin, glycerol, beeswax, a metallic soap; a mucilage; an oil of natural origin such as almond, corn, arachis, castor or olive oil; wool fat or its derivatives or a fatty acid such as steric or oleic acid together with an alcohol such as propylene glycol or a macrogel. The formulation may incorporate any
5 suitable surface active agent such as an anionic, cationic or non-ionic surfactant such as a sorbitan ester or a polyoxyethylene derivative thereof. Suspending agents such as natural gums, cellulose derivatives or inorganic materials such as siliceous silicas, and other ingredients such as lanolin, may also be included.

Drops according to the present invention may comprise sterile aqueous or
10 oily solutions or suspensions and may be prepared by dissolving the active ingredient in a suitable aqueous solution of a bactericidal and/or fungicidal agent and/or any other suitable preservative, and preferably including a surface active agent. The resulting solution may then be clarified by filtration, transferred to a suitable container which is then sealed and sterilized by autoclaving or maintaining
15 at 98-100°C for half an hour. Alternatively, the solution may be sterilized by filtration and transferred to the container by an aseptic technique. Examples of bactericidal and fungicidal agents suitable for inclusion in the drops are phenylmercuric nitrate or acetate (0.002%), benzalkonium chloride (0.01%) and chlorhexidine acetate (0.01%). Suitable solvents for the preparation of an oily
20 solution include glycerol, diluted alcohol and propylene glycol.

Compounds of formula (I) may be administered parenterally, that is by intravenous, intramuscular, subcutaneous intranasal, intrarectal, intravaginal or intraperitoneal administration. The subcutaneous and intramuscular forms of parenteral administration are generally preferred. Appropriate dosage forms for such
25 administration may be prepared by conventional techniques. Compounds of Formula (I) may also be administered by inhalation that is by intranasal and oral inhalation administration. Appropriate dosage forms for such administration, such as an aerosol formulation or a metered dose inhaler, may be prepared by conventional techniques.

30 For all methods of use disclosed herein for the compounds of Formula (I) the daily oral dosage regimen will preferably be from about 0.01 to about 80 mg/kg of total body weight. The daily parenteral dosage regimen about 0.001 to about 80

mg/kg of total body weight. The daily topical dosage regimen will preferably be from 0.1 mg to 150 mg, administered one to four, preferably two or three times daily. The daily inhalation dosage regimen will preferably be from about 0.01 mg/kg to about 1 mg/kg per day. It will also be recognized by one of skill in the art that the optimal quantity and spacing of individual dosages of a compound of Formula (I) or a pharmaceutically acceptable salt thereof will be determined by the nature and extent of the condition being treated, the form, route and site of administration, and the particular patient being treated, and that such optimums can be determined by conventional techniques. It will also be appreciated by one of skill in the art that the optimal course of treatment, i.e., the number of doses of a compound of Formula (I) or a pharmaceutically acceptable salt thereof given per day for a defined number of days, can be ascertained by those skilled in the art using conventional course of treatment determination tests.

The invention will now be described by reference to the following biological examples which are merely illustrative and are not to be construed as a limitation of the scope of the present invention.

BIOLOGICAL EXAMPLES

The IL-8, and GRO- α chemokine inhibitory effects of compounds of the present invention are determined by the following *in vitro* assay:

Receptor Binding Assays:

[¹²⁵I] IL-8 (human recombinant) is obtained from Amersham Corp., Arlington Heights, IL, with specific activity 2000 Ci/mmol. GRO- α is obtained from NEN- New England Nuclear. All other chemicals are of analytical grade. High levels of recombinant human IL-8 type α and β receptors were individually expressed in Chinese hamster ovary cells as described previously (Holmes, *et al.*, Science, 1991, 253, 1278). The Chinese hamster ovary membranes were homogenized according to a previously described protocol (Haour, *et al.*, J. Biol. Chem., 249 pp 2195-2205 (1974)). Except that the homogenization buffer is changed to 10mM Tris-HCL, 1mM MgSO₄, 0.5mM EDTA (ethylene-diaminetetra-acetic acid), 1mM PMSF (α -toluenesulphonyl fluoride), 0.5 mg/L Leupeptin, pH 7.5. Membrane protein concentration is determined using Pierce Co. micro-assay kit

using bovine serum albumin as a standard. All assays are performed in a 96-well micro plate format. Each reaction mixture contains ^{125}I IL-8 (0.25 nM) or ^{125}I GRO- α and 0.5 $\mu\text{g/mL}$ of IL-8R α or 1.0 $\mu\text{g/mL}$ of IL-8R β membranes in 20 mM Bis-Trispropane and 0.4 mM Tris HCl buffers, pH 8.0, containing 1.2 mM MgSO_4 , 0.1 mM EDTA, 25 mM Na and 0.03% CHAPS. In addition, drug or compound of interest is added which has been pre-dissolved in DMSO so as to reach a final concentration of between 0.01nM and 100 μM . The assay is initiated by addition of ^{125}I -IL-8. After 1 hour at room temperature the plate is harvested using a Tomtec 96-well harvester onto a glass fiber filtermat blocked with 1% polyethylenimine/ 0.5% BSA and washed 3 times with 25 mM NaCl, 10 mM TrisHCl, 1 mM MgSO_4 , 0.5 mM EDTA, 0.03 % CHAPS, pH 7.4. The filter is then dried and counted on the Betaplate liquid scintillation counter. The recombinant IL-8 R α , or Type I, receptor is also referred to herein as the non-permissive receptor and the recombinant IL-8 R β , or Type II, receptor is referred to as the permissive receptor.

Representative compounds of Formula (I), Examples 1 to 106 have exhibited positive inhibitory activity in this assay at IC_{50} levels < 30 μM .

Chemotaxis Assay:

The *in vitro* inhibitory properties of these compounds are determined in the neutrophil chemotaxis assay as described in Current Protocols in Immunology, vol. I, Suppl 1, Unit 6.12.3., whose disclosure is incorporated herein by reference in its entirety. Neutrophils where isolated from human blood as described in Current Protocols in Immunology Vol. I, Suppl 1 Unit 7.23.1, whose disclosure is incorporated herein by reference in its entirety. The chemoattractants IL-8, GRO- α , GRO- β , GRO- γ and NAP-2 are placed in the bottom chamber of a 48 multiwell chamber (Neuro Probe, Cabin John, MD) at a concentration between 0.1 and 100 nM. The two chambers are separated by a 5 μM polycarbonate filter. When compounds of this invention are tested, they are mixed with the cells (0.001 - 1000 nM) just prior to the addition of the cells to the upper chamber. Incubation is allowed to proceed for between about 45 and 90 min at about 37°C in a humidified incubator with 5% CO_2 . At the end of the incubation period, the polycarbonate membrane is removed and the topside washed, the membrane then stained using the

Diff Quick staining protocol (Baxter Products, McGaw Park, IL, USA). Cells which have chemotaxed to the chemokine are visually counted using a microscope.

Generally, four fields are counted for each sample, these numbers are averaged to give the average number of cells which had migrated. Each sample is tested in triplicate and each compound repeated at least four times. To certain cells (positive control cells) no compound is added, these cells represent the maximum chemotactic response of the cells. In the case where a negative control (unstimulated) is desired, no chemokine is added to the bottom chamber. The difference between the positive control and the negative control represents the chemotactic activity of the cells.

10 **Elastase Release Assay:**

The compounds of this invention are tested for their ability to prevent Elastase release from human neutrophils. Neutrophils are isolated from human blood as described in Current Protocols in Immunology Vol. I, Suppl 1 Unit 7.23.1. PMNs 0.88×10^6 cells suspended in Ringer's Solution (NaCl 118, KCl 4.56, NaHCO_3 25, KH_2PO_4 1.03, Glucose 11.1, HEPES 5 mM, pH 7.4) are placed in each well of a 96 well plate in a volume of 50 ul. To this plate is added the test compound (0.001 - 1000 nM) in a volume of 50 ul, Cytochalasin B in a volume of 50 ul (20ug/ml) and Ringers buffer in a volume of 50 ul. These cells are allowed to warm (37 °C, 5% CO_2 , 95% RH) for 5 min before IL-8, $\text{GRO}\alpha$, $\text{GRO}\beta$, $\text{GRO}\gamma$ or NAP-2 at a final concentration of 0.01 - 1000 nM was added. The reaction is allowed to proceed for 45 min before the 96 well plate is centrifuged (800 xg 5 min.) and 100 ul of the supernatant removed. This supernatant is added to a second 96 well plate followed by an artificial elastase substrate (MeOSuc-Ala-Ala-Pro-Val-AMC, Nova Biochem, La Jolla, CA) to a final concentration of 6 ug/ml dissolved in phosphate buffered saline. Immediately, the plate is placed in a fluorescent 96 well plate reader (Cytofluor 2350, Millipore, Bedford, MA) and data collected at 3 min intervals according to the method of Nakajima et al *J. Biol. Chem.* 254 4027 (1979). The amount of Elastase released from the PMNs is calculated by measuring the rate of MeOSuc-Ala-Ala-Pro-Val-AMC degradation.

TNF- α in Traumatic Brain Injury Assay

The present assay provides for examination of the expression of tumor necrosis factor mRNA in specific brain regions, which follow experimentally, induced lateral fluid-percussion traumatic brain injury (TBI) in rats. Adult Sprague-Dawley rats (n=42) were anesthetized with sodium pentobarbital (60 mg/kg, i.p.) and subjected to lateral fluid-percussion brain injury of moderate severity (2.4 atm.) centered over the left temporoparietal cortex (n=18), or "sham" treatment (anesthesia and surgery without injury, n=18). Animals are sacrificed by decapitation at 1, 6 and 24 hr. post injury, brains removed, and tissue samples of left (injured) parietal cortex (LC), corresponding area in the contralateral right cortex (RC), cortex adjacent to injured parietal cortex (LA), corresponding adjacent area in the right cortex (RA), left hippocampus (LH) and right hippocampus (RH) are prepared. Total RNA are isolated and Northern blot hybridization is performed and quantitated relative to an TNF- α positive control RNA (macrophage = 100%). A marked increase of TNF- α mRNA expression is observed in LH (104 $\pm$ 17% of positive control, p < 0.05 compared with sham), LC (105 $\pm$ 21%, p < 0.05) and LA (69 $\pm$ 8%, p < 0.01) in the traumatized hemisphere 1 hr. following injury. An increased TNF- α mRNA expression is also observed in LH (46 $\pm$ 8%, p < 0.05), LC (30 $\pm$ 3%, p < 0.01) and LA (32 $\pm$ 3%, p < 0.01) at 6 hr which resolves by 24 hr following injury. In the contralateral hemisphere, expression of TNF- α mRNA is increased in RH (46 $\pm$ 2%, p < 0.01), RC (4 $\pm$ 3%) and RA (22 $\pm$ 8%) at 1 hr and in RH (28 $\pm$ 11%), RC (7 $\pm$ 5%) and RA (26 $\pm$ 6%, p < 0.05) at 6 hr but not at 24 hr following injury. In sham (surgery without injury) or naive animals, no consistent changes in expression of TNF- α mRNA are observed in any of the 6 brain areas in either hemisphere at any times. These results indicate that following parasagittal fluid-percussion brain injury, the temporal expression of TNF- α mRNA is altered in specific brain regions, including those of the non-traumatized hemisphere. Since TNF- α is able to induce nerve growth factor (NGF) and stimulate the release of other cytokines from activated astrocytes, this post-traumatic alteration in gene expression of TNF- α plays an important role in both the acute and regenerative response to CNS trauma.

CNS Injury model for IL-1 β mRNA

This assay characterizes the regional expression of interleukin-1 β (IL-1 β) mRNA in specific brain regions following experimental lateral fluid-percussion traumatic brain injury (TBI) in rats. Adult Sprague-Dawley rats (n=42) are

anesthetized with sodium pentobarbital (60 mg/kg, i.p.) and subjected to lateral fluid-percussion brain injury of moderate severity (2.4 atm.) centered over the left temporoparietal cortex (n=18), or "sham" treatment (anesthesia and surgery without injury). Animals are sacrificed at 1, 6 and 24 hr. post injury, brains removed, and tissue samples of left (injured) parietal cortex (LC), corresponding area in the contralateral right cortex (RC), cortex adjacent to injured parietal cortex (LA), corresponding adjacent area in the right cortex (RA), left hippocampus (LH) and right hippocampus (RH) are prepared. Total RNA is isolated and Northern blot hybridization was performed and the quantity of brain tissue IL-1 β mRNA is presented as percent relative radioactivity of IL-1 β positive macrophage RNA which was loaded on the same gel. At 1 hr following brain injury, a marked and significant increase in expression of IL-1 β mRNA is observed in LC (20.0 $\pm$ 0.7% of positive control, n=6, p < 0.05 compared with sham animal), LH (24.5 $\pm$ 0.9%, p < 0.05) and LA (21.5 $\pm$ 3.1%, p < 0.05) in the injured hemisphere, which remained elevated up to 6 hr. post injury in the LC (4.0 $\pm$ 0.4%, n=6, p < 0.05) and LH (5.0 $\pm$ 1.3%, p < 0.05). In sham or naive animals, no expression of IL-1 β mRNA is observed in any of the respective brain areas. These results indicate that following TBI, the temporal expression of IL-1 β mRNA is regionally stimulated in specific brain regions. These regional changes in cytokines, such as IL-1 β play a role in the post-traumatic.


All publications, including but not limited to patents and patent applications, cited in this specification are herein incorporated by reference as if each individual publication were specifically and individually indicated to be incorporated by reference herein as though fully set forth.

The above description fully discloses the invention including preferred embodiments thereof. Modifications and improvements of the embodiments specifically disclosed herein are within the scope of the following claims. Without further elaboration, it is believed that one skilled in the art can, using the preceding description, utilize the present invention to its fullest extent. Therefore the Examples herein are to be construed as merely illustrative and not a limitation of the scope of the present invention in any way. The embodiments of the invention in which an exclusive property or privilege is claimed are defined as follows.

What is Claimed Is:

1. A compound of the formula:

5



wherein:

R is selected from the group consisting of cyano, OR₁₁, C(O)NR₁₅R₁₆, R₁₈,

10 C(O)OR₁₁, C(O)R₁₁, and S(O)₂R₁₇;

R'R'' is independently selected from the group consisting of hydrogen, NR₆R₇, OH,

OR_a, C₁₋₅alkyl, aryl, arylC₁₋₄alkyl, aryl C₂₋₄alkenyl; cycloalkyl, cycloalkyl

C₁₋₅ alkyl, heteroaryl, heteroarylC₁₋₄alkyl, heteroarylC₂₋₄ alkenyl,

heterocyclic, heterocyclic C₁₋₄alkyl, and a heterocyclic C₂₋₄alkenyl moiety, all

15 of which moieties may be optionally substituted one to three times independently

by a substituent selected from the group consisting of halogen, nitro,

halosubstituted C₁₋₄ alkyl, C₁₋₄ alkyl, amino, mono or di-C₁₋₄ alkyl substituted

amine, OR_a, C(O)R_a, NR_aC(O)OR_a, OC(O)NR₆R₇, hydroxy, NR₉C(O)R_a,

S(O)_mR_a, C(O)NR₆R₇, C(O)OH, C(O)OR_a, S(O)_tNR₆R₇, and NHS(O)_tR_a; or

20 the two R_b substituents join to form a 3-10 membered ring, optionally

substituted and containing, in addition to optionally substituted C₁₋₄ alkyl,

independently, 1 to 3 NR_a, O, S, SO, or SO₂ moieties, which moieties can be

optionally unsaturated;

R''' is selected from the group consisting of Y hydrogen, halogen, nitro, cyano,

25 halosubstituted C₁₋₁₀ alkyl, C₁₋₁₀ alkyl, C₂₋₁₀ alkenyl, C₁₋₁₀ alkoxy,

halosubstituted C₁₋₁₀ alkoxy, azide, (CR₈R₈)_qS(O)_tR_a, (CR₈R₈)_qOR_a, hydroxy,

hydroxy substituted C₁₋₄alkyl, aryl, aryl C₁₋₄ alkyl, aryloxy, arylC₁₋₄ alkyloxy,

- aryl C₂₋₁₀ alkenyl, heteroaryl, heteroarylalkyl, heteroaryl C₁₋₄ alkyloxy, heteroaryl C₂₋₁₀ alkenyl, heterocyclic, heterocyclic C₁₋₄alkyl, heterocyclicC₂₋₁₀ alkenyl, (CR₈R₈)_qNR₄R₅, C₂₋₁₀ alkenyl C(O)NR₄R₅, (CR₈R₈)_qC(O)NR₄R₅, (CR₈R₈)_qC(O)NR₄R₁₀, S(O)₃R₈, (CR₈R₈)_qC(O)R₁₁, C₂₋₁₀ alkenylC(O)R₁₁,
 5 (CR₈R₈)_qC(O)OR₁₁, C₂₋₁₀alkenylC(O)OR₁₁, (CR₈R₈)_qOC(O)R₁₁, (CR₈R₈)_qNR₄C(O)R₁₁, (CR₈R₈)_q NHS(O)_tR₁₃, (CR₈R₈)_q S(O)_tNR₄R₅, (CR₈R₈)_qC(NR₄)NR₄R₅, and (CR₈R₈)_q NR₄C(NR₅)R₁₁; or two Y moieties together form O-(CH₂)_s-O or a 5 to 6 membered saturated or unsaturated ring, such that the alkyl, aryl, arylalkyl, heteroaryl, heteroaryl alkyl, heterocyclic, and
 10 heterocyclicalkyl groups may be optionally substituted;
 R₁ is independently selected from the group consisting of hydrogen, halogen, nitro, cyano, C₁₋₁₀ alkyl, halosubstituted C₁₋₁₀ alkyl, C₂₋₁₀ alkenyl, C₁₋₁₀ alkoxy, halosubstituted C₁₋₁₀alkoxy, azide, S(O)_tR₄, (CR₈R₈)_q S(O)_tR₄, hydroxy, hydroxy substituted C₁₋₄alkyl, aryl, aryl C₁₋₄ alkyl, aryl C₂₋₁₀ alkenyl, aryloxy, aryl C₁₋₄
 15 alkyloxy, heteroaryl, heteroarylalkyl, heteroaryl C₂₋₁₀ alkenyl, heteroaryl C₁₋₄ alkyloxy, heterocyclic, heterocyclic C₁₋₄alkyl, heterocyclicC₁₋₄alkyloxy, heterocyclicC₂₋₁₀ alkenyl, (CR₈R₈)_q NR₄R₅, (CR₈R₈)_qC(O)NR₄R₅, C₂₋₁₀ alkenyl C(O)NR₄R₅, (CR₈R₈)_q C(O)NR₄R₁₀, S(O)₃R₈, (CR₈R₈)_q C(O)R₁₁, C₂₋₁₀ alkenyl C(O)R₁₁, C₂₋₁₀ alkenyl C(O)OR₁₁, (CR₈R₈)_q C(O)OR₁₁, (CR₈R₈)_q OC(O)R₁₁,
 20 (CR₈R₈)_qNR₄C(O)R₁₁, (CR₈R₈)_q C(NR₄)NR₄R₅, (CR₈R₈)_q NR₄C(NR₅)R₁₁, (CR₈R₈)_q NHS(O)_tR₁₃, and (CR₈R₈)_q S(O)_tNR₄R₅; or two R₁ moieties together form O-(CH₂)_sO or a 5 to 6 membered saturated or unsaturated ring, wherein the alkyl, aryl, arylalkyl, heteroaryl, and heterocyclic moieties may be optionally substituted.
- 25 2. The compound according to Claim 1 wherein R₁ is substituted in the 4- position by an electron withdrawing moiety.
 3. The compound according to Claim 2 wherein R₁ is halogen, cyano or nitro.
 4. The compound according to Claim 3 wherein R₁ is halogen.

5. The compound according to Claim 4 wherein R₁ is independently fluorine, chlorine, or bromine.
6. The compound according to Claim 1 wherein Y is mono-substituted in the 2'-position or 3'-position, or is disubstituted in the 2'- or 3'-position of a monocyclic ring.
7. The compound according to Claim 6 wherein Y is halogen.
8. The compound according to Claim 4 wherein Y is independently fluorine, chlorine, or bromine.
9. The compound according to Claim 1 wherein R_b is hydrogen, C₁₋₄ alkyl, C₁₋₄ alkyl substituted with C(O)OH, or C(O)OR_a.
10. The compound according to Claim 1 wherein Y is halogen, n is 1 or 2, R₁ is halogen, m is 1 or 2, and R_b is independently hydrogen, C₁₋₄ alkyl, C₁₋₄ alkyl substituted with C(O)OH, or C(O)OR_a.
11. The compound according to Claim 1 which is selected from the group consisting of:
 - 15 N-(2-bromophenyl)-N'-[4-chloro-2-hydroxy-3-(N'',N''dimethylaminosulfonyl)phenyl] cyanoguanidine;
 - N-[4-chloro-2-hydroxy-3-(N'',N''dimethylaminosulfonyl)phenyl]-N''-(2,3-dichlorophenyl) cyanoguanidine;
 - N-(2-bromophenyl)-N'-[4-chloro-2-hydroxy-3-[S-(+)-(2-methoxymethyl)pyrrolidin-1-yl]aminosulfonylphenyl] cyanoguanidine;
 - 20 N-(2,3-dichlorophenyl)-N'-[4-chloro-2-hydroxy-3-[S-(+)-(2-methoxymethyl)pyrrolidin-1-yl]aminosulfonylphenyl] cyanoguanidine;
 - N-phenyl-N'-[4-chloro-2-hydroxy-3-[S-(+)-(2-methoxymethyl)pyrrolidin-1-yl]aminosulfonylphenyl] cyanoguanidine;
 - 25 N-(2-bromophenyl)-N'-[4-chloro-2-hydroxy-3-[R-(2-methoxymethyl)pyrrolidin-1-yl]aminosulfonylphenyl] cyanoguanidine;
 - N-(2,3-dichlorophenyl)-N'-[4-chloro-2-hydroxy-3-[R-(2-methoxymethyl)pyrrolidin-1-yl]aminosulfonylphenyl] cyanoguanidine;
 - N-(2-bromophenyl)-N'-[4-chloro-2-hydroxy-3-(N''-isoxazolidinylaminosulfonylphenyl] cyanoguanidine;
 - 30

- N-(2,3-dichlorophenyl)-N'-[4-chloro-2-hydroxy-3-(N"-isoxazolidinylaminosulfonylphenyl)] cyanoguanidine;
- N-(2-bromophenyl)-N'-[4-chloro-2-hydroxy-3-(N"-tetrahydroisoxazylaminosulfonyl)phenyl] cyanoguanidine;
- 5 N-(2,3-dichlorophenyl)-N'-[4-chloro-2-hydroxy-3-(N"-tetrahydroisoxazylaminosulfonyl)phenyl] cyanoguanidine;
- N-(2,3-dichlorophenyl)-N'-[4-chloro-2-hydroxy-3-(4-thiomorpholinylaminosulfonyl)phenyl] cyanoguanidine;
- N-[4-chloro-2-hydroxy-3-[N",N"-dimethylaminosulfonyl]phenyl]-N'-(2-bromophenyl)
- 10 propylguanidine;
- N-(2-bromophenyl)-N'-[4-chloro-2-hydroxy-3-(4-oxidothiomorpholino)amino sulfonylphenyl] cyanoguanidine;
- N-(2,3-chlorophenyl)-N'-[4-chloro-2-hydroxy-3-(4-oxidothiomorpholino)amino sulfonylphenyl] cyanoguanidine;
- 15 N-(2-bromophenyl)-N'-[4-chloro-2-hydroxy-3-(N"-methylpiperazino)amino sulfonylphenyl] cyanoguanidine;
- N-(2,3-dichlorophenyl)-N'-[4-chloro-2-hydroxy-3-(N"-methylpiperazino)amino sulfonylphenyl] cyanoguanidine;
- N-(2-bromophenyl)-N'-[4-chloro-2-hydroxy-3-(N"-ethylmorpholino)amino
- 20 sulfonylphenyl] cyanoguanidine;
- N-(2,3-dichlorophenyl)-N'-[4-chloro-2-hydroxy-3-(N"-ethylmorpholino)amino sulfonylphenyl] cyanoguanidine;
- N-(2-bromophenyl)-N'-{4-chloro-2-hydroxy-3-[N"-ethyl-2-(2-ethylpyrrolidino)]amino sulfonylpheny} cyanoguanidine;
- 25 N-(2,3-dichlorophenyl)-N'-{4-chloro-2-hydroxy-3-[N"-ethyl-2-(2-ethylpyrrolidino)]amino sulfonylpheny} cyanoguanidine;
- N-(2-bromophenyl)-N'-{4-chloro-2-hydroxy-3-[S-(+)-(2-carboxy)pyrrolidin-1-yl]amino sulfonylpheny} cyanoguanidine;
- N-(2,3-dichlorophenyl)-N'-{4-chloro-2-hydroxy-3-[S-(+)-(2-carboxy)pyrrolidin-1-
- 30 yl]amino sulfonylpheny} cyanoguanidine;
- N-(2-bromo-3-fluorophenyl)-N'-[4-chloro-2-hydroxy-3-[S-(+)-(2-methoxymethyl)pyrrolidin-1-yl]sulfonylphenyl] cyanoguanidine;

N-(2-phenoxyphenyl)-N'-[4-chloro-2-hydroxy-3-[S-(+)-(2-methoxymethyl)pyrrolidin-1-yl]sulfonylphenyl] cyanoguanidine; and

N-(2-benzoxypyphenyl)-N'-[4-chloro-2-hydroxy-3-[S-(+)-(2-methoxymethyl)pyrrolidin-1-yl]sulfonylphenyl] cyanoguanidine;

5 or a pharmaceutically acceptable salt thereof.

12. A pharmaceutical composition comprising a compound according to any of Claims 1 to 11 and a pharmaceutically acceptable carrier or diluent.

13. A method of treating a chemokine mediated disease, wherein the chemokine binds to an IL-8 a or b receptor in a mammal, which method comprises administering to
10 said mammal an effective amount of a compound of the formula according to any one of Claims 1 to 11.

14. The method according to Claim 13 wherein the mammal is afflicted with a chemokine mediated disease selected from the group consisting of psoriasis, atopic dermatitis, osteo arthritis, rheumatoid arthritis, asthma, chronic obstructive
15 pulmonary disease, adult respiratory distress syndrome, inflammatory bowel disease, Crohn's disease, ulcerative colitis, stroke, septic shock, multiple sclerosis, endotoxic shock, gram negative sepsis, toxic shock syndrome, cardiac and renal reperfusion injury, glomerulonephritis, thrombosis, graft vs. host reaction, alzheimers disease, allograft rejections, malaria, restenosis, angiogenesis, atherosclerosis, osteoporosis,
20 gingivitis and undesired hematopoietic stem cells release and diseases caused by respiratory viruses, herpesviruses, and hepatitis viruses, meningitis, cystic fibrosis, pre-term labor, cough, pruritus, multi-organ dysfuctions, trauma, strains, sprains, contusions, psoriatic arthritis, herpes, encephalitis, CNS vasculitis, traumatic brain injury, CNS tumors, subarachnoid hemorrhage, post surgical trauma, interstitial
25 pneumonitis, hypersensitivity, crystal induced arthritis, acute and chronic pancreatitis, acute alcoholic hepatitis, necrotizing enterocolitis, chronic sinusitis, uveitis, polymyositis, vasculitis, acne, gastric and duodenal ulcers, celiac disease, esophagitis, glossitis, airflow obstruction, airway hyperresponsiveness, bronchiolitis obliterans organizing pneumonia, bronchiectasis, bronchiolitis, bronchiolitis
30 obliterans, chronic bronchitis, cor pulmonae, dyspnea, emphysema, hypercapnea, hyperinflation, hypoxemia, hyperoxia-induced inflammations, hypoxia, surgical lung volume reduction, pulmonary fibrosis, pulmonary hypertension, right

ventricular hypertrophy, sarcoidosis, small airway disease, ventilation-perfusion mismatching, wheeze, colds and lupus.