


N-ACYLAMINO BENZENE DERIVATIVES AS SELECTIVE MONOAMINE OXIDASE B INHIBITORS

This invention relates to N-acylamino aryl derivatives of the general formula



wherein

- R^1 is halogen, halogen-(C_1 - C_6)-alkyl, cyano, C_1 - C_6 -alkoxy or
 5 halogen-(C_1 - C_6)-alkoxy;
- R^{21} , R^{22} , R^{23} and R^{24} independently from each other are selected from the group
 consisting of hydrogen, (C_1 - C_6)-alkyl, halogen, halogen-(C_1 - C_6)-alkyl,
 hydroxy, C_1 - C_6 -alkoxy or -CHO;
- R^3 is hydrogen or C_1 - C_3 -alkyl;
- 10 R^4 , R^5 independently from each other are selected from the group
 consisting of hydrogen, C_1 - C_6 -alkyl, C_1 - C_6 -alkoxy or -COO(C_1 - C_6)alkyl;
 or R^4 and R^5 form together with the C-atom to which they are attach a
 C_3 - C_7 -cycloalkyl ring;
- R^6 is -CO-NR⁷R⁸; -COO(C_1 - C_6)-alkyl, -CN, -NR₂ or -NHC(O)R;
- 15 R^7 and R^8 independently from each other are selected from the group
 consisting of hydrogen, C_1 - C_6 -alkyl, NH₂ or hydroxy;
- R is hydrogen or C_1 - C_6 -alkyl;
- n is 0, 1, 2 or 3.
- X is -CHRO, -OCHR-, -CH₂S-, -SCH₂-, -CH₂CH₂-, -CH=CH- or -C≡C-;

20 and to pharmaceutically active acid addition salts thereof.

It has been found that the compounds of general formula I are selective monoamine
 oxidase B inhibitors.


Monoamine oxidase (MAO, EC 1.4.3.4) is a flavin-containing enzyme responsible for the oxidative deamination of endogenous monoamine neurotransmitters such as dopamine, serotonin, adrenaline, or noradrenaline, and trace amines, e.g. phenylethylamine, as well as a number of amine xenobiotics. The enzyme exists in two forms, MAO-A and MAO-B, encoded by different genes (A. W. Bach et al., *Proc. Natl. Acad. Sci. USA* 1988, 85, 4934-4938) and differing in tissue distribution, structure and substrate specificity. MAO-A has higher affinity for serotonin, octopamine, adrenaline, and noradrenaline; whereas the natural substrates for MAO-B are phenylethylamine and tyramine. Dopamine is thought to be oxidised by both isoforms. MAO-B is widely distributed in several organs including brain (A.M. Cesura and A. Pletscher, *Prog. Drug Research* 1992, 38, 171-297). Brain MAO-B activity appears to increase with age. This increase has been attributed to the gliosis associated with aging (C.J. Fowler et al., *J. Neural. Transm.* 1980, 49, 1-20). Additionally, MAO-B activity is significantly higher in the brains of patients with Alzheimer's disease (P. Dostert et al., *Biochem. Pharmacol.* 1989, 38, 555-561) and it has been found to be highly expressed in astrocytes around senile plaques (Saura et al., *Neuroscience* 1994, 70, 755-774). In this context, since oxidative deamination of primary monoamines by MAO produces NH_3 , aldehydes and H_2O_2 , agents with established or potential toxicity, it is suggested that there is a rationale for the use of selective MAO-B inhibitors for the treatment of dementia and Parkinson's disease. Inhibition of MAO-B causes a reduction in the enzymatic inactivation of dopamine and thus prolongation of the availability of the neurotransmitter in dopaminergic neurons. The degeneration processes associated with age and Alzheimer's and Parkinson's diseases may also be attributed to oxidative stress due to increased MAO activity and consequent increased formation of H_2O_2 by MAO-B. Therefore, MAO-B inhibitors may act by both reducing the formation of oxygen radicals and elevating the levels of monoamines in the brain.

Given the implication of MAO-B in the neurological disorders mentioned above, there is considerable interest to obtain potent and selective inhibitors that would permit control over this enzymatic activity. The pharmacology of some known MAO-B inhibitors is for example discussed by D. Bentué-Ferrer et al. in *CNS Drugs* 1996, 6, 217-236. Whereas a major limitation of irreversible and non-selective MAO inhibitor activity is the need to observe dietary precautions due to the risk of inducing a hypertensive crisis when dietary tyramine is ingested, as well as the potential for interactions with other medications (D. M. Gardner et al., *J. Clin. Psychiatry* 1996, 57, 99-104), these adverse events are of less concern with reversible and selective MAO inhibitors, in particular of MAO-B. Thus, there is a need for MAO-B inhibitors with a high selectivity and without the adverse side-effects typical of irreversible MAO inhibitors with low selectivity for the enzyme.

Summary of Invention

Disclosed herein is to provide compounds which may have the criteria mentioned above. It has been found that the compounds of formula I disclosed herein show the potential to be highly selective MAO-B inhibitors. Further disclosed herein is a process for the manufacture of compounds of formula I as well as the use of the compounds of formula I in the control or prevention of diseases mediated by monoamine oxidase B inhibitors, and respectively, their use for the production of corresponding medicaments.

According to a first aspect, there is provided compounds of the general formula



wherein

R^1 is halogen, halogen-(C₁-C₆)-alkyl, cyano, C₁-C₆-alkoxy or halogen-(C₁-C₆)-alkoxy;

R^{21} , R^{22} , R^{23} and R^{24} independently from each other are selected from the group consisting of hydrogen, (C₁-C₆)-alkyl, halogen, halogen-(C₁-C₆)-alkyl, hydroxy, C₁-C₆-alkoxy or -CHO;

R^3 is hydrogen or C₁-C₃-alkyl;

R^4 , R^5 independently from each other are selected from the group consisting of hydrogen, C₁-C₆-alkyl, C₁-C₆-alkoxy or -COO(C₁-C₆)alkyl; or R^4 and R^5 form together with the C-atom to which they are attached a C₃-C₇-cycloalkyl ring;

R^6 is -CO-NR⁷R⁸; -COO(C₁-C₆)-alkyl, -CN, -NR₂ or -NHC(O)R;

3a

R^7 and R^8 independently from each other are selected from the group consisting of hydrogen, C_1 - C_6 -alkyl, NH_2 or hydroxy;

R is hydrogen or C_1 - C_6 -alkyl;

n is 0, 1, 2 or 3;

X is $-CHRO$, $-OCHR-$, $-CH_2S-$, $-SCH_2-$, $-CH_2CH_2-$, $-CH=CH-$ or $-C\equiv C-$;

and pharmaceutically active acid addition salts thereof,

with the proviso that

N -(4-benzyloxy-2,6-dimethyl-phenyl)-2-diethylamino-acetamide and hydrochloride salt thereof,

2-diethylamino- N -[4-(4-methoxy-benzyloxy)-2,6-dimethyl-phenyl]-acetamide,

N -[4-(4-chloro-benzyloxy)-2,6-dimethyl-phenyl]-2-diethylamino-acetamide,

N -(4-benzyloxy-2-methyl-phenyl)-2-diethylamino-acetamide,

N -(4-benzyloxy-5-isopropyl-2-methyl-phenyl)-2-diethylamino-acetamide,

N -(4-benzyloxy-phenyl)-2-diethylamino-acetamide,

N -(4-benzyloxy-2,6-dichloro-phenyl)-2-diethylamino-acetamide,

N -(4-benzyloxy-phenyl)-2-dimethylamino-acetamide hydrochloride,

N -(4-benzyloxy-phenyl)-2-dimethylamino-propionamide hydrochloride,

N -[4-(4-chloro-benzyloxy)-phenyl]-2-dimethylamino-acetamide hydrochloride,

N -[4-(4-chloro-benzyloxy)-phenyl]-2-dimethylamino-propionamide hydrochloride,

N -(4-benzyloxy-phenyl)-2-diethylamino-acetamide hydrochloride,

N -(4-benzyloxy-phenyl)-2-diethylamino-propionamide hydrochloride,

N -[4-(4-chloro-benzyloxy)-phenyl]-2-diethylamino-acetamide hydrochloride,

N -[4-(4-chloro-benzyloxy)-phenyl]-2-diethylamino-propionamide hydrochloride,

N -(4-benzyloxy-2-methyl-phenyl)-2-diethylamino-acetamide,

N -(4-benzyloxy-2-methyl-phenyl)-2-(butyl-methyl-amino)-acetamide,

N -(4-benzyloxy-2-methyl-phenyl)-2-dibutylamino-acetamide,

N -(4-benzyloxy-2-methyl-phenyl)-2-dipropylamino-acetamide,

N -(4-benzyloxy-2-methyl-phenyl)-2-propylamino-acetamide,

N -(4-benzyloxy-2-methyl-phenyl)-2-hexylamino-acetamide,

N -(4-benzyloxy-2-methyl-phenyl)-2-diethylamino-propionamide,

N -(4-benzyloxy-2-methyl-phenyl)-2-butylamino-propionamide,

N -(4-benzyloxy-2-methyl-phenyl)-2-propylamino-propionamide,

N -(4-benzyloxy-2,6-dimethyl-phenyl)-2-butylamino-acetamide hydrochloride salt thereof,

N -(4-benzyloxy-2,6-dimethyl-phenyl)-2-dibutylamino-acetamide hydrochloride salt thereof

N -(4-benzyloxy-2,6-dimethyl-phenyl)-2-dipropylamino-acetamide, and

N -(4-benzyloxy-2-methylphenyl)-2-(butylamino)acetamide;

and compounds of the formula I wherein

n is 0, 1, 2 or 3; and

R^1 is halogen, cyano, C_1 - C_6 -alkoxy, $-(CH_2)_m-C(halogen)_3$ with m ranging from 0 to 3 or $-O-(CH_2)_m-C(halogen)_3$ with m ranging from 0 to 3; and

R^{21} , R^{22} , R^{23} , and R^{24} are all hydrogen; and

X is $-(CH_2)_m-$ with $m=2$, $-CH=CH-$, or $-C\equiv C-$; and

R^3 is hydrogen or C_1 - C_3 -alkyl; and

R^6 is $-CN$; and

R^4 , R^5 is hydrogen;

are excluded.

The following definitions of general terms used herein apply irrespective of whether the terms in question appear alone or in combination. It must be noted that, as used in the specification and the appended claims, the singular forms "a", "an," and "the" include plural forms unless the context clearly dictates otherwise.

The term "C₁-C₆-alkyl" ("lower alkyl") used herein denotes straight-chain or branched saturated hydrocarbon residues with 1 to 6 carbon atoms, preferably with 1 to 3 carbon atoms, such as methyl, ethyl, n-propyl, i-propyl, n-butyl, sec-butyl, t-butyl, and the like. Accordingly, the term "C₁-C₃-alkyl" means a straight-chain or branched saturated hydrocarbon residue with 1 to 3 carbon atoms.

The term "halogen" denotes fluorine, chlorine, bromine and iodine.

"Halogen-(C₁-C₆)-alkyl" or "halogen-(C₁-C₆)-alkoxy" means the lower alkyl residue or lower alkoxy residue, respectively, as defined herein substituted in any position with one or more halogen atoms as defined herein. Examples of halogenalkyl residues include, but are not limited to, 1,2-difluoropropyl, 1,2-dichloropropyl, trifluoromethyl, 2,2,2-trifluoroethyl, 2,2,2-trichloroethyl, and 1,1,1-trifluoropropyl, and the like. "Halogenalkoxy" includes trifluoromethoxy.


"C₁-C₆-Alkoxy" means the residue -O-R, wherein R is a lower alkyl residue as defined herein. Examples of alkoxy radicals include, but are not limited to, methoxy, ethoxy, isopropoxy, and the like.

The term "C₃-C₇-cycloalkyl" denotes a saturated carbocyclic group, containing 3 to 7 carbon atoms. For example, a cycloalkyl group is cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and cycloheptyl, and these groups may optionally be substituted by one or two C₁-C₄-alkyl substituents, for example methyl or ethyl.

"Pharmaceutically acceptable salts" of a compound means salts that are pharmaceutically acceptable, which are generally safe, non-toxic, and neither biologically nor otherwise undesirable, and that possess the desired pharmacological activity of the

parent compound. These salts may be derived from an inorganic or organic acid or base. If possible, compounds of formula I may be converted into pharmaceutically salts. It should be understood that pharmaceutically acceptable salts may be included herein.

According to a second aspect, there are provided compounds of the general formula Ia



wherein

R^1 is halogen, halogen-(C_1 - C_6)-alkyl, cyano,
 C_1 - C_6 -alkoxy or halogen-(C_1 - C_6)-alkoxy;

R^{21} , R^{22} , R^{23} and R^{24} independently from each other are selected from the group
consisting of hydrogen and halogen;

R^3 is hydrogen or C_1 - C_3 -alkyl;

R^4 is hydrogen or C_1 - C_6 -alkyl;

R^5 is hydrogen or C_1 - C_6 -alkyl;

or R^4 and R^5 together form a C_3 - C_7 -cycloalkyl ring;

R^6 is $-CO-NR^7R^8$;
 $-COOR^9$; or
 $-CN$;

R^7 and R^8 independently from each other are hydrogen, methyl or ethyl;

R^9 is C_1 - C_6 -alkyl; and

n is 1, 2 or 3;

and pharmaceutically active acid addition salts thereof.

In a preferred embodiment, there are provided compounds of formula I, wherein X is CH_2O , R^1 is fluorine or trifluoromethyl and R^6 is $-\text{COOCH}_3$, for example the following compounds:

N-[4-(3-fluoro-benzyloxy)-phenyl]-malonamic acid methyl ester,
N-[3-fluoro-4-(3-fluoro-benzyloxy)-phenyl]-malonamic acid methyl ester,
N-[4-(4-fluoro-benzyloxy)-phenyl]-malonamic acid methyl ester,

N-[2-fluoro-4-(3-fluoro-benzyloxy)-phenyl]-malonamic acid methyl ester,
 N-[4-(2,4-difluoro-benzyloxy)-phenyl]-malonamic acid methyl ester,
 N-[4-(2-fluoro-benzyloxy)-phenyl]-malonamic acid methyl ester,
 N-[4-(2,4,5-trifluoro-benzyloxy)-phenyl]-malonamic acid methyl ester,
 N-[2-fluoro-4-(4-fluoro-benzyloxy)-phenyl]-malonamic acid methyl ester,
 N-[4-(3,5-bis-trifluoromethyl-benzyloxy)-2-fluoro-phenyl]-malonamic acid methyl ester,
 N-[4-(3-fluoro-benzyloxy)-3-methyl-phenyl]-malonamic acid methyl ester or
 N-[3-chloro-4-(3-fluoro-benzyloxy)-phenyl]-malonamic acid methyl ester.

In a further preferred embodiment, there are provided compounds of formula I, wherein X is CH_2O and R^6 is $-\text{CONH}_2$, for example the following compounds:

cyclopropane-1,1-dicarboxylic acid amide [4-(3-fluoro-benzyloxy)-phenyl]-amide,
 N-[4-(3-fluoro-benzyloxy)-phenyl]-malonamide,
 N-[4-(3-fluoro-benzyloxy)-phenyl]-2-methyl-malonamide,
 N-[3-fluoro-4-(3-fluoro-benzyloxy)-phenyl]-malonamide,
 N-[4-(4-fluoro-benzyloxy)-phenyl]-malonamide,
 N-[4-(2,4-difluoro-benzyloxy)-phenyl]-malonamide,
 N-[4-(2,4,5-trifluoro-benzyloxy)-phenyl]-malonamide,
 N-[4-(2-fluoro-benzyloxy)-phenyl]-malonamide,
 N-(4-benzyloxy-phenyl)-malonamide,
 N-[4-(4-chloro-benzyloxy)-phenyl]-malonamide,
 N-[4-(3-fluoro-benzyloxy)-2-hydroxy-phenyl]-malonamide,
 N-[2-fluoro-4-(4-fluoro-benzyloxy)-phenyl]-malonamide,
 N-[4-(3-fluoro-benzyloxy)-3-methyl-phenyl]-malonamide,
 N-[3-chloro-4-(3-fluoro-benzyloxy)-phenyl]-malonamide or
 cyclopropane-1,1-dicarboxylic acid amide [2-fluoro-4-(4-fluoro-benzyloxy)-phenyl]-amide.

In another preferred embodiment, there are provided compounds of formula I wherein X is $-\text{OCH}_2-$ and R^6 is $-\text{NHCOCH}_3$ or $-\text{NHCOH}$. Examples of such compounds are the following:

2-Acetyl-amino-N-[2-fluoro-4-(4-fluoro-benzyloxy)-phenyl]-acetamide,
 2-Acetyl-amino-N-[2-fluoro-4-(3-fluoro-benzyloxy)-phenyl]-acetamide,
 N-[2-Fluoro-4-(4-fluoro-benzyloxy)-phenyl]-2-formyl-amino-acetamide or
 N-[2-Fluoro-4-(3-fluoro-benzyloxy)-phenyl]-2-formyl-amino-acetamide.

5a

In yet another preferred embodiment, there are provided compounds of formula I wherein X is $-\text{OCH}_2-$ and R^6 is $-\text{NH}_2$.

The following compound is an example thereof:

2-amino-N-[2-fluoro-4-(4-fluoro-benzyloxy)-phenyl]-acetamide.

In yet another preferred embodiment, there is provided a group of compounds of formula I, wherein X is

$-\text{CH}=\text{CH}-$ and R^6 is $-\text{COOCH}_3$ or $-\text{CONH}_2$, for example the followings:


N-{4-[2-(4-fluoro-phenyl)-vinyl]-phenyl}-malonamic acid methyl ester,

N-{4-[2-(3-fluoro-phenyl)-vinyl]-phenyl}-malonamide,


N-{4-[2-(4-fluoro-phenyl)-vinyl]-phenyl}-malonamide or

N-{4-[2-(3-fluoro-phenyl)-vinyl]-phenyl}-malonamic acid methyl ester.


According to a third aspect, there is provided a process for the manufacture of a compound of general formula I as described above, which process comprises reacting a compound of formula



10 with a compound of formula



to obtain a compound of formula




and, if desired, converting a compound of formula I into a pharmaceutically

15 acceptable salt.

According to a fourth aspect, there is provided a medicament containing one or more compounds as described above and pharmaceutically acceptable excipients for use in the treatment and prevention of diseases which are mediated by monoamine oxidase B inhibitors.

According to a fifth aspect, there is provided a compound of formula I as described above as well as its pharmaceutically acceptable salts for the treatment or prevention of diseases.

According to a sixth aspect, there is provided the use of a compound of general formula I



wherein

R^1 is halogen, halogen-(C₁-C₆)-alkyl, cyano, C₁-C₆-alkoxy or halogen-(C₁-C₆)-alkoxy;

R^{21} , R^{22} , R^{23} and R^{24} independently from each other are selected from the group consisting of hydrogen, (C₁-C₆)-alkyl, halogen, halogen-(C₁-C₆)-alkyl, hydroxy, C₁-C₆-alkoxy or -CHO;

R^3 is hydrogen or C₁-C₃-alkyl;

R^4 , R^5 independently from each other are selected from the group consisting of hydrogen, C₁-C₆-alkyl, C₁-C₆-alkoxy or -COO(C₁-C₆)-alkyl; or R^4 and R^5 form together with the C-atom to which they are attached a C₃-C₇-cycloalkyl ring;

R^6 is -CO-NR⁷R⁸; -COO(C₁-C₆)-alkyl, -CN, -NR₂ or -NHC(O)R;

R^7 and R^8 independently from each other are selected from the group consisting of hydrogen, C₁-C₆-alkyl, NH₂ or hydroxy;

R is hydrogen or C₁-C₆-alkyl;


6b

n is 0, 1, 2 or 3;

X is -CHRO, -OCHR-, -CH₂S-, -SCH₂-, -CH₂CH₂-, -CH=CH- or -C≡C-;

as well as its pharmaceutically acceptable salts for the manufacture of medicaments for the treatment and prevention of diseases which are mediated by monoamine oxidase B inhibitors.

According to a seventh aspect, there is provided the use of a compound of general formula Ia



wherein

R¹ is halogen, halogen-(C₁-C₆)-alkyl, cyano, C₁-C₆-alkoxy or halogen-(C₁-C₆)-alkoxy;

R²¹, R²², R²³ and R²⁴ independently from each other are selected from the group consisting of hydrogen and halogen;

R³ is hydrogen or C₁-C₃-alkyl;

R⁴ is hydrogen or C₁-C₆-alkyl;

R⁵ is hydrogen or C₁-C₆-alkyl;

or R⁴ and R⁵ together form a C₃-C₇-cycloalkyl ring;

R⁶ is -CO-NR⁷R⁸;
-COOR⁹; or
-CN;

R⁷ and R⁸ independently from each other are hydrogen, methyl or ethyl;

R⁹ is C₁-C₆-alkyl; and

n is 1, 2 or 3;

as well as its pharmaceutically acceptable salts for the manufacture of medicaments for the treatment and prevention of diseases which are mediated by monoamine oxidase B inhibitors.

Detail Description of Embodiments

In accordance with the present invention, a possibility to prepare compounds of general formula I is shown in scheme 1: The key intermediates A are accessible through nucleophilic substitution of aromatic nitro compounds containing p-substituted leaving groups with benzylic alcohols or thiols. P- substituted leaving groups can be for example halogens (F, Cl, Br, I), tosylates, mesylates or triflates. These substitution reactions can be conducted neat or in inert solvents like for example toluene or xylene. Preferred reaction temperatures are between 50° and 150 °C. Alternatively, compounds A can be prepared

- 7 -

- by Williamson-ether synthesis, starting from p-nitrophenols and benzylic halides, tosylates, mesylates or triflates. Bases used can be for example alcoholates or carbonates (sodium, potassium or cesium carbonate). Preferred solvents are lower alcohols, acetonitrile or lower ketones at temperatures between 20 °C and reflux temperature.
- 5 Another approach is the Mitsunobu-coupling of benzylic alcohols with p-nitrophenols. The reaction is done as usual in inert solvents like for example diethyl ether or tetrahydrofuran, using dialkyl-azo-dicarboxylates in presence of phosphines (for example tributyl- or triphenyl-phosphine).

- The key intermediates A are reduced to the amino-compounds B using catalytic
- 10 hydrogenation (for example Platinum on charcoal in lower alcohols, ethyl acetate or tetrahydrofuran). An alternative is the reduction of the nitro-group by metals like iron, tin, or zinc in acidic media like diluted hydrochloric acid or acetic acid. Metals can also be replaced by metal salts (for example tin-(II)-chloride).


- Intermediates B can be acylated by known methods to give the desired compounds I-A. These reactions can be done with acid chlorides and bases (for example trialkylamines, sodium carbonate or potassium carbonate, sodium hydrogen carbonate or potassium
- 20 hydrogen carbonate), eventually in presence of an acylation catalyst (e.g. 1 to 10 mol% of N,N-dimethyl-4-aminopyridine) in solvents like dichloromethane, ethyl acetate or acetonitrile, preferentially at room temperature. An alternative is the well known coupling of an acid with the amine B using coupling reagents like N,N'-
- 25 dicyclohexylcarbodiimide (DCC), N-(3-dimethylaminopropyl)-N'-ethyl-carbodiimide hydrochloride (EDC) or 1,1'-carbonyl-diimidazole (CDI) in solvents like dichloromethane, diethyl ether or THF, preferentially at 0 to 40 °C. Intermediates B can also primarily be monoalkylated by known methods (see for example Johnstone et al, *J. Chem. Soc. (C)* 1969, 2233 or Krishnamurthy et al., *Tetrahedron Lett.* 1982, 23, 3315) to give compounds D. These are then acylated as previously described to lead to the desired compounds I-B.

- Compounds with inverted ether or thioether linkers can be obtained by similar reactions, as depicted in scheme 1a. The key-intermediates A' are then further
- 30 transformed into the final products by the reaction sequences already mentioned in scheme 1.


Additional functional group manipulations can be done by standard methods on the acylated compounds to obtain all compounds of formula I (for example : functionalising the malonic position by deprotonation and reaction with electrophiles).

- 8 -

Scheme 1



Scheme 1a




- Alternatively, in accordance with the present invention, compounds B can also be prepared by alkylation of N-protected p-hydroxyanilines with benzylic halides or by
- 5 Mitsunobu-coupling of N-protected p-hydroxyanilines with benzylic alcohols (scheme 2) by the methods described previously. Protective groups PG can be for example N-Boc (N-tert.-butoxycarbonyl) or N-acetyl. Deprotection of E leads to the intermediates B.

Obviously, compounds with inverted ether and thioether linkers can be prepared by similar reaction sequences, inverting substituent-patterns on the aromatic moieties.

10


Scheme 2



Another method to prepare compounds of the type D or I-B involves cross-coupling reactions of arylstannanes (Lam et al., *Tetrahedron Lett.* 2002, 43, 3091),

arylboronates (Lam et al., *Synlett* 2000, 5, 674); Chan et al., *Tetrahedron Lett.* 1998, 39, 2933) or aryl halides (Buchwald et al., *J. Amer. Chem. Soc.* 1996, 118, 7215) with the corresponding amines or amides (scheme 3).


Scheme 3



Once again, compounds with inverted ether and thioether linkers can be prepared by similar reaction sequences, inverting substituent-patterns on the aromatic moieties.

A possibility to prepare compounds **I** where X is $-CH_2CH_2-$ or $-CH=CH-$ is shown in scheme 4. Wittig- or Wittig-Horner reactions between phosphor-ylides and aromatic aldehydes under standard conditions leads to styrene-derivatives **G**. These intermediates can be reduced to intermediates **H** or **I**. Usually, reduction of **G** by Bechamp-type conditions leads to compounds **H**, which can then be further reduced to compounds **I** by catalytic hydrogenation. Using harsher hydrogenation conditions, compounds **G** can be reduced in one step to give compound **I**. Compounds **H** or **I** are then further processed by the methods depicted in scheme 1 to yield the final products **I**. Another method to prepare compounds of the type **G** involves nitration of styrene-derivatives.

Scheme 4



The compounds of formula I are, as already mentioned above, monoamine oxidase B inhibitors and can be used for the treatment or prevention of diseases in which MAO-B inhibitors might be beneficial. These include acute and chronic neurological disorders, cognitive disorders and memory deficits. Treatable neurological disorders are for instance traumatic or chronic degenerative processes of the nervous system, such as Alzheimer's disease, other types of dementia, minimal cognitive impairment or Parkinson's disease. Other indications include psychiatric diseases such as depression, anxiety, panic attack, social phobia, schizophrenia, eating and metabolic disorders such as obesity as well as the prevention and treatment of withdrawal syndromes induced by abuse of alcohol, nicotine and other addictive drugs. Other treatable indications may be reward deficiency syndrome (G.M. Sullivan, International patent application No. WO 01/34172 A2), peripheral neuropathy caused by cancer chemotherapy (G. Bobotas, International Patent Application No. WO 97/33572 A1), or the treatment of multiple sclerosis (R.Y. Harris, International patent application No. WO 96/40095 A1) and other neuroinflammatory diseases.

The compounds of formula I are especially useful for the treatment and prevention of Alzheimer's disease and senile dementia.

The pharmacological activity of the compounds was tested using the following method:

The cDNA's encoding human MAO-A and MAO-B were transiently transfected into EBNA cells using the procedure described by E.-J. Schlaeger and K. Christensen (Transient Gene Expression in Mammalian Cells Grown in Serum-free Suspension Culture; Cytotechnology, 15: 1-13, 1998). After transfection, cells were homogenised by means of a Polytron homogenizer in 20 mM Tris HCl buffer, pH 8.0, containing 0.5 mM EGTA and 0.5 mM phenylmethanesulfonyl fluoride. Cell membranes were obtained by centrifugation at 45,000 x g and, after two rinsing step with 20 mM Tris HCl buffer, pH 8.0, containing 0.5 mM EGTA, membranes were eventually re-suspended in the above buffer and aliquots stored at -80 °C until use.

MAO-A and MAO-B enzymatic activity was assayed in 96-well-plates using a spectrophotometric assay adapted from the method described by M. Zhou and N. Panchuk-Voloshina (A One-Step Fluorometric Method for the Continuous Measurement of Monoamine Oxidase Activity, Analytical Biochemistry, 253: 169-174, 1997). Briefly, membrane aliquots were incubated in 0.1 M potassium phosphate buffer, pH 7.4, for 30 min at 37 °C with or without various concentrations of the compounds. After this period, the enzymatic reaction was started by the addition of the MAO substrate tyramine together with 1 U/ml horse-radish peroxidase (Roche Biochemicals) and 80 µM N-acetyl-3,7,-dihydroxyphenoxazine (Amplex Red, Molecular Probes). The samples were further incubated for 30 min at 37 °C in a final volume of 200 µl and absorbance was then determined at a wavelength of 570 nm using a SpectraMax plate reader (Molecular Devices). Background (non-specific) absorbance was determined in the presence of 10 µM clorgyline for MAO-A or 10 µM L-deprenyl for MAO-B.

IC₅₀ values were determined from inhibition curves obtained using nine inhibitor concentrations in duplicate, by fitting data to a four parameter logistic equation using a computer program.

The compounds of the present invention are specific MAO-B inhibitors. The IC₅₀ values of preferred compounds of formula I as measured in the assay described above are in the range of 1000 nM or less, typically 100 nM or less, and ideally 50 nM or less.

In the table above are disclosed some IC₅₀ values (nM) of preferred compounds.

Example No.	IC ₅₀ (nM)	Example No.	IC ₅₀ (nM)
1	11	39	5
2	7.9	40	15
3	10.9	41	13
6	13	44	9
8	9.6	47	14
9	7.2	48	19
10	11.2	49	10
11	7.5	50	6
14	9	56	10
25	11	57	18
27	11	66	16
28	8	70	11
29	8	72	7
34	18	75	13
38	16	76	16

The compounds of formula I can be used as medicaments, e.g. in the form of pharmaceutical preparations. The pharmaceutical preparations can be administered orally, e.g. in the form of tablets, coated tablets, dragées, hard and soft gelatine capsules, solutions, emulsions or suspensions. However, the administration can also be effected rectally, e.g. in the form of suppositories, or parenterally, e.g. in the form of injection solutions.

The compounds of formula I can be processed with pharmaceutically inert, inorganic or organic carriers for the production of pharmaceutical preparations. Lactose, corn starch or derivatives thereof, talc, stearic acid or its salts and the like can be used, for

- example, as such carriers for tablets, coated tablets, dragées and hard gelatine capsules. Suitable carriers for soft gelatine capsules are, for example, vegetable oils, waxes, fats, semi-solid and liquid polyols and the like; depending on the nature of the active substance no carriers are, however, usually required in the case of soft gelatine capsules.
- 5 Suitable carriers for the production of solutions and syrups are, for example, water, polyols, sucrose, invert sugar, glucose and the like. Adjuvants, such as alcohols, polyols, glycerol, vegetable oils and the like, can be used for aqueous injection solutions of water-soluble salts of compounds of formula I, but as a rule are not necessary. Suitable carriers for suppositories are, for example, natural or hardened oils, waxes, fats, semi-liquid or
- 10 liquid polyols and the like.

In addition, the pharmaceutical preparations can contain preservatives, solubilizers, stabilizers, wetting agents, emulsifiers, sweeteners, colorants, flavorants, salts for varying the osmotic pressure, buffers, masking agents or antioxidants. They may also contain other therapeutically valuable substances.

- 15 As mentioned earlier, medicaments containing a compound of formula I and a therapeutically inert excipient are also an object of the present invention, as is a process for the production of such medicaments which comprises bringing one or more compounds of formula I and, if desired, one or more other therapeutically valuable substances into a galenical dosage form together with one or more therapeutically inert
- 20 carriers.

- The dosage can vary within wide limits and will, of course, be fitted to the individual requirements in each particular case. In general, the effective dosage for oral or parenteral administration is between 0.01-20 mg/kg/day, with a dosage of 0.1-10 mg/kg/day being preferred for all of the indications described. The daily dosage for an adult
- 25 human being weighing 70 kg accordingly lies between 0.7-1400 mg per day, preferably between 7 and 700 mg per day.

The following examples are provided for illustration of the invention. They should not be considered as limiting the scope of the invention, but merely as being representative thereof.

30

Example 1

N-[4-(3-Fluoro-benzyloxy)-phenyl]-malonic acid methyl ester

a) 1-(3-Fluorobenzyloxy)-4-nitro-benzene

A mixture of 5.04 g (40 mmol) 3-fluorobenzyl alcohol and 1.29 g (4mmol) tris-(dioxo-3,6-heptyl)amine is treated with 2.47 g (44 mmol) of potassium hydroxide. The

- 15 -

mixture is stirred at room temperature for 10 min, then 5.55 g (44 mmol) of 4-fluoro-nitrobenzene is slowly added through a dropping funnel. The mixture is kept for 45 min at 80 °C, cooled to room temperature and diluted with about 75 ml water. Extraction with ethyl acetate and washing with 2M aqueous hydrochloric acid yields a yellowish
5 organic phase, which is dried and evaporated. The residue is recrystallised from methanol to give 6.07 g (61 %) of the title compound. Yellow crystals, mp = 104-105 °C.

b) 4-(3-Fluoro-benzyloxy)-phenylamine

3 g (12.1 mmol) of 1-(3-fluorobenzyloxy)-4-nitro-benzene is dissolved in 125 ml of methanol. 150 mg of Pt 5% on charcoal is added and hydrogenation done under normal
10 pressure for about 17 h. The catalyst is filtered and the solution evaporated to yield 2.51 g (95%) of crude brownish material. MS: m/e = 218.4 ($M^+ + H$).

c) N-[4-(3-Fluoro-benzyloxy)-phenyl]-malonamic acid methyl ester

A solution of 242 mg (1.77 mmol) chlorocarbonyl-acetic acid methyl ester in 2ml dichloromethane is added dropwise to an ice-cooled solution of 350 mg (1.61 mmol)
15 4-(3-fluoro-benzyloxy)-phenylamine in 7ml of pyridine. The mixture is then stirred for 2h at room temperature, diluted with water, acidified with dil. hydrochloric acid and extracted with ethyl acetate. The organic phase is dried. Chromatography (silica gel, ethyl acetate / hexane 1:1) yields 99 mg (19%) of the title compound. Yellowish solid; MS: m/e = 418.1 ($M^+ + H$).

20

Example 2

Cyclopropane-1,1-dicarboxylic acid amide [4-(3-fluoro-benzyloxy)-phenyl]-amide

A solution of 126 mg (1 mmol) of 1-(aminocarbonyl)-1-cyclopropan-carboxylic acid and of 212 mg (1 mmol) 4-(3-fluoro-benzyloxy)-phenylamine (prepared as in
25 example 1b) in 10 ml dichloromethane is cooled to 0 °C and treated with 206 mg (1.1 mmol) N-(-3-dimethylaminopropyl)-N'-ethyl-carbodiimide-hydrochloride (EDCI). The mixture is stirred for additional 2h at room temperature, evaporated and acidified with dil. citric acid. Extraction with ethyl acetate and recrystallisation from methanol yields 178 mg (56%) of the title compound. Yellowish solid; MS: m/e = 329.3 ($M^+ + H$).

Example 3

30 N-[4-(3-Fluoro-benzyloxy)-phenyl]-malonamide

99 mg (0.31 mmol) N-[4-(3-fluoro-benzyloxy)-phenyl]-malonamic acid methyl ester (example 1) is dissolved in a mixture of 0.5ml tetrahydrofuran and 1.0ml of aqueous ammonium hydroxide (25%). The reaction vessel is capped and kept at 60 °C for 2.5 h. The reaction mixture is cooled, evaporated to dryness and diluted with water.

Filtration provides 48 mg (51%) of the title compound as a light yellow solid. MS: $m/e = 303.2$ ($M^+ + H$).

Example 4

2-Cyano-N-[4-(3-fluoro-benzyloxy)-phenyl]-acetamide

- 5 The title compound is prepared in analogy to example 2, starting from 4-(3-fluoro-benzyloxy)-phenylamine (prepared as in example 1b) and cyanoacetic acid. Yield after recrystallisation from methanol: 50%. Colourless crystals with mp = 164-166 °C.

Example 5

N-[4-(3-Fluoro-benzyloxy)-phenyl]-2-methyl-malonamic acid methyl ester

- 10 The title compound is prepared in analogy to example 2, starting from 4-(3-fluoro-benzyloxy)-phenylamine (as prepared in example 1b) and 2-methyl-malonic acid monomethyl ester (Nader et al., *Chem. Ber.* 1986, 119, 1196). Yield after flash-chromatography with dichloromethane / methanol 98:2: 40%. Colourless solid. Mp = 119 °C.

Example 6

N-[4-(3-Fluoro-benzyloxy)-phenyl]-2-methyl-malonamide

The title compound is prepared in analogy to example 3 from N-[4-(3-fluoro-benzyloxy)-phenyl]-2-methyl-malonamic acid methyl ester and ammonium hydroxide. Colourless solid. Mp = 206 °C.

Example 7

N-[4-(3-Fluoro-benzyloxy)-phenyl]-2,2-dimethyl-malonamide

a) N-[4-(3-Fluoro-benzyloxy)-phenyl]-2,2-dimethyl-malonamic acid ethyl ester

- 25 The title compound is prepared in analogy to example 2, starting from 4-(3-fluoro-benzyloxy)-phenylamine and 2,2-dimethyl-malonic acid monoethyl ester (Holmes et al., *J. Amer. Chem. Soc.* 1984, 106, 2353). Yield after flash-chromatography with cyclohexane / ethyl acetate 7:3 : 79%. Colourless oil. MS: $m/e = 359$ (M^+).

b) N-[4-(3-Fluoro-benzyloxy)-phenyl]-2,2-dimethyl-malonamide

- 30 The title compound is prepared in analogy to example 3 from N-[4-(3-fluoro-benzyloxy)-phenyl]-2,2-dimethyl-malonamic acid ethyl ester and ammonium hydroxide. Colourless solid. MS: $m/e = 331.4$ ($M^+ + H$).

Example 8

N-[3-Fluoro-4-(3-fluoro-benzyloxy)-phenyl]-malonamic acid methyl ester

a) 3-Fluoro-4-(3-fluoro-benzyloxy)-nitrobenzene

A mixture of 10.0g (64mmol) 2-fluoro-4-nitrophenol, 13.24g (70mmol) 3-fluorobenzyl bromide and 17.6g (127mmol) potassium carbonate in 200ml methyl-ethylketone is heated overnight at 80°. Water is added and the product is extracted with ethyl acetate, dried and evaporated. Recrystallisation from diethyl ether / hexane yields
5 12.68g (75%) of a slightly yellow solid. MS: m/e = 265.1 (M^+).

b) 3-Fluoro-4-(3-fluoro-benzyloxy)-phenylamine

12.68g (47.8mmol) of 3-fluoro-4-(3-fluoro-benzyloxy)-nitrobenzene in 150ml ethyl acetate is treated with 1.27g of 5% platinum on charcoal and hydrogenated for 6 h at room temperature and normal pressure. The catalyst is filtered and the solution
10 evaporated to yield 11.03g (98%) of dark brown oil. MS: m/e = 235.1 (M^+).

c) N-[3-Fluoro-4-(3-fluoro-benzyloxy)-phenyl]-malonamic acid methyl ester

The title compound is prepared in analogy to example 1 from 3-fluoro-4-(3-fluoro-benzyloxy)-phenylamine and chlorocarbonyl-acetic acid methyl ester. Yield : 65%. Light yellow solid. MS: m/e = 336.2 ($M^+ + H$).

15

Example 9

N-[3-Fluoro-4-(3-fluoro-benzyloxy)-phenyl]-malonamide

The title compound is prepared in analogy to example 3 from N-[3-fluoro-4-(3-fluoro-benzyloxy)-phenyl]-malonamic acid methyl ester and ammonium hydroxide. Light yellow solid. MS: m/e = 321.2 ($M^+ + H$).

20

Example 10

N-[4-(4-Fluoro-benzyloxy)-phenyl]-malonamic acid methyl ester

a) 4-(4-Fluoro-benzyloxy)-nitrobenzene

The title compound is prepared in analogy to 1-(3-fluorobenzyloxy)-4-nitrobenzene, starting from 4-fluoro nitrobenzene and 4-fluoro benzyl alcohol. Yield : 86% of
25 a slightly yellow solid. Mp. = 124-126°.

b) 4-(4-Fluoro-benzyloxy)-phenylamine

Prepared in analogy to 4-(3-fluoro-benzyloxy)-phenylamine, by hydrogenation of 4-(4-fluoro-benzyloxy)-nitrobenzene. Yield: 100% of a slightly red solid. MS: m/e = 218.2 ($M^+ + H$).

c) N-[4-(4-Fluoro-benzyloxy)-phenyl]-malonic acid methyl ester

The title compound is prepared in analogy to example 1 from 4-(4-fluoro-benzyloxy)-phenylamine and chlorocarbonyl-acetic acid methyl ester. Yield : 75%. Colorless solid. MS: $m/e = 318.2 (M^+ + H)$.

5

Example 11N-[4-(4-Fluoro-benzyloxy)-phenyl]-malonamide

The title compound is prepared in analogy to example 3 from N-[4-(4-fluoro-benzyloxy)-phenyl]-malonic acid methyl ester and ammonium hydroxide. Yield : 80%. Colorless solid. MS: $m/e = 303.2 (M^+ + H)$.

10

Example 12N-[4-(4-Trifluoromethyl-benzyloxy)-phenyl]-malonic acid methyl estera) 4-(4-Trifluoromethyl-benzyloxy)-nitrobenzene

Prepared in analogy to 1-(3-fluorobenzyloxy)-4-nitro-benzene, starting from 4-fluoro nitrobenzene and 4-trifluoromethyl benzyl alcohol. Yield after recrystallisation from methanol : 82% of a brownish solid. Mp = 80.5 – 81.5 °C.

15

b) 4-(4-Trifluoromethyl-benzyloxy)-phenylamine

Prepared in analogy to 4-(3-fluoro-benzyloxy)-phenylamine, by hydrogenation of 4-(4-trifluoromethyl-benzyloxy)-nitrobenzene. Yield: 98% of a slightly yellow solid. MS: $m/e = 268.3 (M^+ + H)$.

20

c) N-[4-(4-Trifluoromethyl-benzyloxy)-phenyl]-malonic acid methyl ester

In analogy to example 1, the title compound is prepared from 4-(4-trifluoromethyl-benzyloxy)-phenylamine and chlorocarbonyl-acetic acid methyl ester. Yield : 71%. Colorless solid. MS: $m/e = 368.1 (M^+ + H)$.

Example 1325 N-[4-(4-Trifluoromethyl-benzyloxy)-phenyl]-malonamide

The title compound is prepared in analogy to example 3 from N-[4-(4-trifluoromethyl-benzyloxy)-phenyl]-malonic acid methyl ester and ammonium hydroxide. Yield = 65%. Colorless solid. MS: $m/e = 353.1 (M^+ + H)$.

Example 14N-[2-Fluoro-4-(3-fluoro-benzyloxy)-phenyl]-malonamic acid methyl estera) 2-Fluoro-4-(3-fluoro-benzyloxy)-nitrobenzene

Prepared in analogy to 3-fluoro-4-(3-fluoro-benzyloxy)-nitrobenzene, starting
5 from 3-fluoro-4-nitrophenol and 3-fluorobenzyl bromide. Yield after recrystallisation
from diethyl ether / hexane : 100% of a white solid. MS: $m/e = 265.0$ (M^+).

b) 2-Fluoro-4-(3-fluoro-benzyloxy)-phenylamine

The title compound is prepared in analogy to 3-fluoro-4-(3-fluoro-benzyloxy)-
phenylamine by hydrogenation of 2-Fluoro-4-(3-fluoro-benzyloxy)-nitrobenzene. Yield :
10 98% of a dark brown oil. MS: $m/e = 235.0$ (M^+).

c) N-[2-Fluoro-4-(3-fluoro-benzyloxy)-phenyl]-malonamic acid methyl ester

In analogy to example 1, the title compound is prepared from 2-fluoro-4-(3-fluoro-
benzyloxy)-phenylamine and chlorocarbonyl-acetic acid methyl ester. Yield : 47%.
Brownish solid. MS: $m/e = 336.1$ ($M^+ + H$).

15

Example 15N-[2-Fluoro-4-(3-fluoro-benzyloxy)-phenyl]-malonamide

The title compound is prepared in analogy to example 3 from N-[2-fluoro-4-(3-fluoro-
benzyloxy)-phenyl]-malonamic acid methyl ester and ammonium hydroxide. Yield =
87%. Slightly yellow solid. MS: $m/e = 321.3$ ($M^+ + H$).

20

Example 16N-[4-(3-Fluoro-phenoxy)methyl]-phenyl]-malonamic acid methyl estera) 1-Nitro-4-(3-fluoro-phenoxy)methyl)-benzene

A solution of 2.0 g (13 mmol) 4-nitrobenzyl alcohol and 3.60 g (13.7mmol)
triphenylphosphine in 30ml tetrahydrofuran is treated with 1.54g (13.7mmol) of 3-
25 fluorophenol. The mixture is cooled in an ice-bath and 2.39g (13.7mmol) of diethyl-aza-
dicarboxylate is slowly added. The ice-bath is removed and the reaction mixture stirred
over night at room temperature. The tetrahydrofuran is evaporated, the oily residue
trituated with diethyl ether, filtered and concentrated. Chromatography (silica gel, ethyl
acetate / hexane 1:9) yields 2.52g (78%) of the title compound as a slightly yellow solid.
30 MS: $m/e = 247.1$ ($M^+ + H$).

b) 4-(3-Fluoro-phenoxy)methyl)-phenylamine

1.0 g (4mmol) of 1-Nitro-4-(3-fluoro-phenoxy-methyl)-benzene in 40ml methanol is treated with 0.25g of 5% platinum on charcoal and hydrogenated for 1 h at room temperature and normal pressure. The catalyst is filtered and the solution evaporated to yield 0.98g of the crude title compound as yellow oil. MS: $m/e = 218.3 (M^+ + H)$.

5 c) N-[4-(3-Fluoro-phenoxy-methyl)-phenyl]-malonamic acid methyl ester

211mg (1mmol) of the crude 4-(3-fluoro-phenoxy-methyl)-phenylamine is dissolved in 3ml dichloromethane. 110mg (1.1mmol) of triethylamine is added and the reaction mixture cooled in an ice-bath 132mg (1mmol) of methyl malonyl chloride is then added dropwise. The reaction mixture is stirred for additional 30 minutes in the ice-bath, then for 45 minutes at room temperature. The reaction mixture is treated with water, acidified with citric acid and extracted 3 times with dichloromethane. The organic phase is dried and evaporated. Chromatography (silica gel, ethyl acetate / hexane 1:1) yields 98mg (32%) of the title compound as a colorless solid. MS: $m/e = 318.2 (M^+ + H)$.

Example 17

15 N-[4-(3-Fluoro-phenoxy-methyl)-phenyl]-malonamide

The title compound is prepared in analogy to example 3 from N-[4-(3-Fluoro-phenoxy-methyl)-phenyl]-malonamic acid methyl ester and ammonium hydroxide. Yield = 37%. Colorless solid. MS: $m/e = 303.2 (M^+ + H)$.

Example 18

20 Cyclopropane-1,1-dicarboxylic acid amide [4-(3-fluoro-phenoxy-methyl)-phenyl]-amide

The title compound is prepared in analogy to example 2 from 4-(3-Fluoro-phenoxy-methyl)-phenylamine and 1-(aminocarbonyl)-1-cyclopropane-carboxylic acid. Yield = 42%. Colorless solid. MS: $m/e = 329.3 (M^+ + H)$.

Example 19

25 N-[4-(3-Fluoro-benzyloxy)-phenyl]-2-methoxy-malonamic acid methyl ester

The title compound is prepared in analogy to example 2 from 4-(3-Fluoro-benzyloxy)-phenylamine and 2-methoxy-malonic acid monomethyl ester (Swan et al.; J.Chem.Soc.Perkin Trans.1; 1985; 1757-1766). Yield = 41%. Colorless solid. MS: $m/e = 348.2 (M^+ + H)$.

Example 20

N-[4-(3-Fluoro-benzyloxy)-2-trifluoromethyl-phenyl]-malonamic acid methyl ester

a) 4-(3-Fluoro-benzyloxy)-1-nitro-2-trifluoromethyl-benzene

Prepared in analogy to example 8 from 4-nitro-3-(trifluoromethyl)-phenol and 3-Fluoro-benzylbromide. Yield = 85%. Yellowish solid. MS (ISP): $m/e = 333.2$ ($M^+ + NH_4$).

b) 4-(3-Fluoro-benzyloxy)-2-trifluoromethyl-phenylamine

- 5 Prepared in analogy to example 8b) by hydrogenation of 4-(3-Fluoro-benzyloxy)-1-nitro-2-trifluoromethyl-benzene. Yield = 93%. Slightly yellow oil. MS: $m/e = 286.1$ ($M^+ + H$).

c) N-[4-(3-Fluoro-benzyloxy)-2-trifluoromethyl-phenyl]-malonamic acid methyl ester

- 10 0.515g (1.8mmol) 4-(3-Fluoro-benzyloxy)-2-trifluoromethyl-phenylamine is dissolved in 20ml ethyl acetate. 0.38g (4.51mmol) sodium hydrogencarbonate is added and the mixture cooled in the ice-bath. 0.27g (2.0mmol) methyl malonyl chloride, dissolved in 5ml ethyl acetate is slowly added over a period of about 15min. The resulting mixture is stirred at room temperature overnight. 10ml 2M aqueous hydrochloric acid is added, the organic layer separated and dried over magnesium sulfate. Evaporation of the
15 solvent gives a slightly yellow solid which is recrystallised from methanol. Colorless solid, Yield = 58%. MS: $m/e = 386.1$ ($M^+ + H$).

Example 21

N-[2,5-Difluoro-4-(3-fluoro-benzyloxy)-phenyl]-malonamide

a) 1,4-Difluoro-2-(3-fluoro-benzyloxy)-5-nitro-benzene

- 20 Prepared in analogy to example 1a) from 2,4,5-trifluoronitrobenzene and 3-fluoro-benzyl alcohol. Colorless solid. Yield = 55%. MS: $m/e = 283.1$ (M^+).

b) 2,5-Difluoro-4-(3-fluoro-benzyloxy)-phenylamine

Prepared in analogy to example 8b) by hydrogenation of 1,4-difluoro-2-(3-fluoro-benzyloxy)-5-nitro-benzene. Yield = 96%. Brown solid. MS: $m/e = 253.2$ (M^+).

- 25 c) N-[2,5-Difluoro-4-(3-fluoro-benzyloxy)-phenyl]-malonamic acid methyl ester

Prepared in analogy to example 20c) from 2,5-difluoro-4-(3-fluoro-benzyloxy)-phenylamine and methyl malonyl chloride. Greyish solid. Yield = 73%. MS: $m/e = 354.1$ ($M^+ + H$).

d) N-[2,5-Difluoro-4-(3-fluoro-benzyloxy)-phenyl]-malonamide

- 30 The title compound is prepared in analogy to example 3 from N-[2,5-difluoro-4-(3-fluoro-benzyloxy)-phenyl]-malonamic acid methyl ester and ammonium hydroxide. Yield = 56%. Colorless solid. MS: $m/e = 339.3$ ($M^+ + H$).

Example 22N-[4-(3-Fluoro-benzyloxy)-phenyl]-N-methyl-malonamic acid methyl estera) [4-(3-Fluoro-benzyloxy)-phenyl]-methyl-amine

- 5 0.855g (8.4mmol) acetic anhydride is cooled to 0° and 0.475g (10.3mmol) acetic acid added dropwise. The reaction mixture is then heated to 55° for 2 hours. After cooling to room temperature the mixture is diluted with 0.5ml dry tetrahydrofuran. A solution of 0.7g (3.2mmol) 4-(3-fluoro-benzyloxy)-phenylamine in 2ml tetrahydrofuran is added dropwise at room temperature and the resulting mixture is stirred for 3 hours.
- 10 Evaporation leaves a solid which is dissolved again in 2ml tetrahydrofuran and cooled to 0°. 0.64g (8.1mmol) borane-methyl sulfide complex is slowly added under stirring and cooling. The mixture is stirred 1h at 0° and 2h at room temperature. After addition of 5ml methanol, stirring is pursued for 1h. The mixture is treated with 10ml 1M aqueous hydrochloric acid and stirred overnight. The organic solvents are removed under vacuum
- 15 and the aqueous phase adjusted to pH = 10 bei addition of potassium carbonate. Extraction with diethyl ether, drying and evaporation yields 0.57g (76%) of a colorless solid. MS: m/e = 232.2 (M⁺+H).

b) N-[4-(3-Fluoro-benzyloxy)-phenyl]-N-methyl-malonamic acid methyl ester

- c) Prepared in analogy to example 20c) from [4-(3-fluoro-benzyloxy)-phenyl]-methyl-amine and methyl malonyl chloride. Yellowish oil. Yield = 70%. MS: m/e = 332.3 (M⁺+H).
- 20

Example 23N-[4-(3-Fluoro-benzylsulfanyl)-phenyl]-malonamic acid methyl estera) 1-Fluoro-3-(4-nitrophenylsulfanyl-methyl)-benzene

- 25 Prepared in analogy to example 8 from 4-nitrothiophenol and 3-fluoro-benzylbromide. Yellowish solid. Yield = 54%.

b) 4-(3-Fluoro-benzylsulfanyl)-phenylamine

- 1.75g (6.7mmol) 1-Fluoro-3-(4-nitrophenylsulfanylmethyl)-benzene and 5.55g (100mmol) of iron-powder are suspended in 35ml water and 0.2ml acetic acid.
- 30 The mixture is heated overnight at 90°, then cooled and treated with saturated sodium carbonate solution. The suspension is filtered, the solid washed several times with ethyl acetate and the aqueous phase extracted 3 times with ethyl acetate. The combined organic

phases are dried and evaporated to give 1.18g (76%) of a yellowish waxy solid. MS: $m/e = 234.3 (M^+ + H)$.

c) N-[4-(3-Fluoro-benzylsulfanyl)-phenyl]-malonamic acid methyl ester

Prepared in analogy to example 20c) from 4-(3-fluoro-benzylsulfanyl)-phenylamine and methyl malonyl chloride. Yellowish solid. Yield = 64%. MS: $m/e = 334.2 (M^+ + H)$.

Example 24

N-[4-(3-Fluoro-benzylsulfanyl)-phenyl]-malonamide

Prepared in analogy to example 3 from N-[4-(3-fluoro-benzylsulfanyl)-phenyl]-malonamic acid methyl ester and ammonium hydroxide. Yield = 47%. Colorless solid. MS: $m/e = 319.3 (M^+ + H)$.

Example 25

N-[4-(2,4-Difluoro-benzyloxy)-phenyl]-malonamic acid methyl ester

a) 2,4-Difluoro-1-(4-nitro-phenoxy-methyl)-benzene

Prepared in analogy to example 8 from 4-nitrophenol and 2,4-difluorobenzylbromide. Colorless solid. Yield = 86%. MS: $m/e = 265.1 (M^+)$.

b) 4-(2,4-Difluoro-benzyloxy)-phenylamine

Prepared in analogy to example 8b) by hydrogenation of 2,4-difluoro-1-(4-nitro-phenoxy-methyl)-benzene. Yield = 99%. Brown solid. MS: $m/e = 235.1 (M^+)$.

c) N-[4-(2,4-Difluoro-benzyloxy)-phenyl]-malonamic acid methyl ester

Prepared in analogy to example 20c) from 4-(2,4-difluoro-benzyloxy)-phenylamine and methyl malonyl chloride. Colorless solid. Yield = 99%. MS: $m/e = 336.2 (M^+ + H)$.

Example 26

N-[4-(2-Fluoro-benzyloxy)-phenyl]-malonamic acid methyl ester

a) 1-Fluoro-2-(4-nitrophenoxy-methyl)-benzene

Prepared in analogy to example 8 from 4-nitrophenol and 2-fluorobenzylbromide. Colorless solid. Yield = 74%. MS: $m/e = 247.1 (M^+)$.

b) 4-(2-Fluoro-benzyloxy)-phenylamine

Prepared in analogy to example 8b) by hydrogenation of 1-fluoro-2-(4-nitrophenoxy-methyl)-benzene. Yield = 99%. Brown oil. MS: $m/e = 217.2 (M^+)$.

c) N-[4-(2-Fluoro-benzyloxy)-phenyl]-malonamic acid methyl ester

Prepared in analogy to example 20c) from 4-(2-fluoro-benzyloxy)-phenylamine
5 and methyl malonyl chloride. Colorless solid. Yield = 70%. MS: $m/e = 318.2 (M^+ + H)$.

Example 27

N-[4-(2,4,5-Trifluoro-benzyloxy)-phenyl]-malonamic acid methyl ester

a) 1,2,4-Trifluoro-5-(4-nitro-phenoxy-methyl)-benzene

Prepared in analogy to example 8 from 4-nitrophenol and 2,4,5-trifluoro-
10 benzylbromide. Colorless solid. Yield = 92%. MS: $m/e = 283.0 (M^+)$.

b) 4-(2,4,5-Trifluoro-benzyloxy)-phenylamine

Prepared in analogy to example 8b) by hydrogenation 1,2,4-trifluoro-5-(4-nitro-phenoxy-methyl)-benzene. Yield = 97%. Slightly red solid. MS: $m/e = 254.1 (M^+ + H)$.

c) N-[4-(2,4,5-Trifluoro-benzyloxy)-phenyl]-malonamic acid methyl ester

15 Prepared in analogy to example 20c) from 4-(2,4,5-trifluoro-benzyloxy)-phenylamine and methyl malonyl chloride. Colorless solid. Yield = 91%. MS: $m/e = 352.2 (M^+ - H)$.

Example 28

N-[4-(2,4-Difluoro-benzyloxy)-phenyl]-malonamide

20 Prepared in analogy to example 3 from N-[4-(2,4-difluoro-benzyloxy)-phenyl]-malonamic acid methyl ester and ammonium hydroxide. Yield = 49%. Colorless solid. MS: $m/e = 321.1 (M^+ + H)$.

Example 29

N-[4-(2,4,5-Trifluoro-benzyloxy)-phenyl]-malonamide

25 Prepared in analogy to example 3 from N-[4-(2,4,5-trifluoro-benzyloxy)-phenyl]-malonamic acid methyl ester and ammonium hydroxide. Yield = 52%. Colorless solid. MS: $m/e = 339.2 (M^+ + H)$.

- 25 -

Example 30N-[4-(2-Fluoro-benzyloxy)-phenyl]-malonamide

Prepared in analogy to example 3 from N-[4-(2-fluoro-benzyloxy)-phenyl]-malonamic acid methyl ester and ammonium hydroxide. Yield = 30%. Colorless solid.

5 MS: m/e = 303.2 ($M^+ + H$).

Example 31N-{4-[1-(3-Fluoro-phenyl)-ethoxy]-phenyl}-malonamidea) 1-Fluoro-3-(1-(4-nitrophenoxy)-ethyl)-benzene

Prepared in analogy to example 16a) from 1-(3-fluoro-phenyl)-ethanol
10 (Balasubramanian et al., Synth. Commun., 1994, 24 (8), 1049) and 4-nitrophenol. Yellow oil. Yield = 77%. MS: m/e = 261.2 (M^+).

b) 4-[1-(3-Fluoro-phenyl)-ethoxy]-phenylamine

Prepared in analogy to example 8b) by hydrogenation of 1-fluoro-3-(1-(4-nitrophenoxy)-ethyl)-benzene. Yield = 95%. Yellowish oil. MS: m/e = 232.2 ($M^+ + H$).

15 c) N-{4-[1-(3-Fluoro-phenyl)-ethoxy]-phenyl}-malonamic acid methyl ester

Prepared in analogy to example 20c) from 4-[1-(3-fluoro-phenyl)-ethoxy]-phenylamine and methyl malonyl chloride. Yellowish solid. Yield = 30%. MS: m/e = 332.4 ($M^+ + H$).

d) N-{4-[1-(3-Fluoro-phenyl)-ethoxy]-phenyl}-malonamide

20 Prepared in analogy to example 3 from N-{4-[1-(3-fluoro-phenyl)-ethoxy]-phenyl}-malonamic acid methyl ester and ammonium hydroxide. Yield = 55%. Colorless solid. MS: m/e = 317.3 ($M^+ + H$).

Example 32N-(4-Benzyloxy-phenyl)-malonamic acid methyl ester

25 Prepared in analogy to example 20c) from 4-benzyloxylaniline and methyl malonyl chloride. Yellowish solid. Yield = 62%. MS: m/e = 300.2 ($M^+ + H$).

Example 33N-(4-Benzyloxy-phenyl)-malonamide

30 Prepared in analogy to example 3 from N-(4-benzyloxy-phenyl)-malonamic acid methyl ester and ammonium hydroxide. Yield = 85%. Colorless solid. MS: m/e = 285.1 ($M^+ + H$).

Example 34N-[4-(4-Chloro-benzyloxy)-phenyl]-malonamidea) N-(4-Hydroxy-phenyl)-malonamide

A suspension of 6.0g (18.5mmol) of N-(4-benzyloxy-phenyl)-malonamide in
5 150ml methanol is treated with 0.8g of palladium 5% on charcoal. Hydrogenation is
performed at room temperature and normal pressure overnight. The reaction mixture is
then brought to reflux and filtered hot. Cooling of the filtrate yields 3.56g (99%) of the
title compound as a colorless solid. MS: $m/e = 285.1 (M^+ + H)$.

b) N-[4-(4-Chloro-benzyloxy)-phenyl]-malonamide

10 Prepared in analogy to example 8 from N-(4-hydroxy-phenyl)-malonamide and 4-
chloro-benzylbromide. Colorless solid. Yield = 18%. MS: $m/e = 319.1 (M^+ + H)$.

Example 35N-[4-(3-Fluoro-benzyloxy)-phenyl]-2-hydrazinocarbonyl-acetamide

A suspension of 0.35g (1.1mmol) of N-[4-(3-fluoro-benzyloxy)-phenyl]-
15 malonamic acid methyl ester in 7ml methanol is treated with 83mg (1.7mmol) of
hydrazine hydrate. The reaction mixture is refluxed for 2h, then another portion of
300mg (6mmol) of hydrazine hydrate is added and reflux is pursued for another 8h. The
mixture is cooled and hold overnight in the frig. The precipitate is filtered off and
recrystallised from methanol to yield 0.043g (12%) of filthy, colorless needles. MS: $m/e =$
20 318.3 ($M^+ + H$).

Example 36N-[4-(3-Fluoro-benzyloxy)-phenyl]-N'-hydroxy-malonamide

A solution of sodium methylate is prepared by dissolving 0.23g (10mmol) sodium
in 10ml methanol. This solution is cooled to 0° and 0.25g (5mmol) of hydroxylamine
25 hydrochloride is added, followed by 0.317g (1mmol) of N-[4-(3-fluoro-benzyloxy)-
phenyl]-malonamic acid methyl ester. The mixture is stirred for 2h at room temperature,
diluted with 25ml of water and acidified to pH = 3 by addition of citric acid. The
precipitate is filtered and recrystallised from ethyl acetate, giving 0.123mg (39%) of a
colorless solid. MS: $m/e = 319.3 (M^+ + H)$.

30

Example 372-[4-(3-Fluoro-benzyloxy)-phenylcarbamoyl]-malonic acid dimethyl ester

0.15g (0.47mmol) of N-[4-(3-fluoro-benzyloxy)-phenyl]-malonamic acid methyl
ester is dissolved in 7ml tetrahydrofuran and cooled to 0°. 38mg (1mmol) of sodium

hydride (55 to 65% in oil) is added and the mixture stirred for 45 minutes at room temperature. After cooling again to 0° a solution of 56mg (0.6mmol) methyl chloroformate in 0.5ml tetrahydrofuran is slowly added with a syringe. The yellowish solution is stirred at room temperature for 2h, treated with about 40ml of water, acidified to pH = 3 by addition of citric acid and extracted 3 times with ethyl acetate. The organic phase is dried over magnesium sulfate and concentrated in vacuo yielding 212mg of a yellow gum. Chromatography (silica gel, ethyl acetate / cyclohexane 1:1) gives 62mg (35%) of a slightly yellow solid. MS: $m/e = 376.3$ ($M^+ + H$).

Example 38

10 N-[4-(3-Fluoro-benzyloxy)-2-hydroxy-phenyl]-malonamide

a) 3-(3-Fluoro-benzyloxy)-phenol

Prepared in analogy to example 8 from resorcinol and 3-fluoro-benzylbromide. Colorless oil. Yield = 7%. MS: $m/e = 218.1$ (M^+).

b) 5-(3-Fluoro-benzyloxy)-2-nitro-phenol

15 1.0g (5mmol) of 3-(3-fluoro-benzyloxy)-phenol is dissolved in 10ml acetic acid and cooled to 10°. 0.44g (5mmol) of 65% nitric acid is dissolved in 10ml acetic acid and slowly added to the previous solution. The colored mixture is stirred for 15min. at 10° and for 2h at room temperature. 50ml of water is added and the product extracted 3 times with ethyl acetate. Chromatography (silica gel, dichloromethane / methanol 98:2) yields 0.38g (32%) of the desired product as a yellow solid. MS: $m/e = 262.1$ ($M^+ - H$).

c) 2-Amino-5-(3-fluoro-benzyloxy)-phenol

Prepared in analogy to example 8b) by hydrogenation of 5-(3-fluoro-benzyloxy)-2-nitro-phenol. Yield = 95%. Greyish solid. MS: $m/e = 234.3$ ($M^+ + H$).

d) N-[4-(3-Fluoro-benzyloxy)-2-hydroxy-phenyl]-malonamic acid methyl ester

25 Prepared in analogy to example 20c) from 2-amino-5-(3-fluoro-benzyloxy)-phenol and methyl malonyl chloride. Yellowish solid. Yield = 90%. MS: $m/e = 334.3$ ($M^+ + H$).

e) N-[4-(3-Fluoro-benzyloxy)-2-hydroxy-phenyl]-malonamide

30 Prepared in analogy to example 3 from N-[4-(3-fluoro-benzyloxy)-2-hydroxy-phenyl]-malonamic acid methyl ester and ammonium hydroxide. Yield = 23%. Colorless solid. MS: $m/e = 319.4$ ($M^+ + H$).

Example 39

N-[2-Fluoro-4-(4-fluoro-benzyloxy)-phenyl]-malonamic acid methyl ester

a) 1-(2-Fluoro-4-iodo-phenyl)-2,5-dimethyl-1H-pyrrole

A solution of 2-fluoro-4-iodoaniline (25.3 g, 107 mmol), acetonylacetone (14.9 g, 131 mmol) and para-toluenesulfonic acid (203 mg, 1 mmol) in dry toluene was heated under reflux for 1 h with a Dean-Stark tap. After cooling to room temperature the mixture was poured into sodium hydrogen carbonate (saturated). The organic layer was separated, washed with brine, dried over sodium sulfate, filtered and then boiled with charcoal. Filtering to remove the charcoal followed by evaporation afforded the title compound (33.2 g, 98%) as a light brown solid. MS: $m/e = 315.1$ ($M^+ + H$).

b) 1-[2-Fluoro-4-(4-fluoro-benzyloxy)-phenyl]-2,5-dimethyl-1H-pyrrole

A solution of 1-(2-fluoro-4-iodo-phenyl)-2,5-dimethyl-1H-pyrrole (33.18 g, 105 mmol), 4-fluorobenzylalcohol (26.6 g, 211 mmol), cesium carbonate (68.6 g, 211 mmol), cuprous iodide (2.0 g, 11 mmol) and 1,10-phenanthroline (3.8 g, 21 mmol) in toluene (52 mL) was heated at 100 °C in an autoclave for 48 h. After cooling to room temperature, the mixture was filtered and poured into sodium hydroxide (1N). The resulting mixture was extracted with toluene and the organic extracts washed with brine, dried and evaporated. The residue was triturated with boiling hexane to leave the title compound (24.7 g, 75%) as a light brown solid. MS: $m/e = 314.1$ ($M^+ + H$).

c) 2-Fluoro-4-(4-fluoro-benzyloxy)-phenylamine (1:1) hydrochloride

A solution of the 1-[2-fluoro-4-(4-fluoro-benzyloxy)-phenyl]-2,5-dimethyl-1H-pyrrole (23.7 g, 76 mmol), potassium hydroxide (8.5 g, 151 mmol), hydroxylamine HCl (62.5 g, 899 mmol) in water (125 mL) and propan-2-ol (250 mL) was heated at 100 °C for 36 h. After cooling to room temperature, the mixture was poured into water, and extracted with diethylether. The combined organic layers were then washed with brine, dried and the resulting solution was diluted with HCl in diethylether. The resulting precipitate was then filtered off to afford the title compound (15.2 g, 74%) as a light brown solid. MS: $m/e = 236.0$ ($M^+ - Cl$).

d) N-[2-Fluoro-4-(4-fluoro-benzyloxy)-phenyl]-malonamic acid methyl ester

The title compound is prepared in analogy to example 1 from 2-fluoro-4-(4-fluoro-benzyloxy)-phenylamine (1:1) hydrochloride. Light yellow solid. Yield = 95%. MS: $m/e = 236.2$ ($M^+ + H$).

Alternatively the 2-fluoro-4-(4-fluoro-benzyloxy)-phenylamine (1:1) hydrochloride can be prepared as follows.

e) 2-Fluoro-4-(4-fluoro-benzyloxy)-nitrobenzene

A solution of 3-fluoro-4-nitrophenol (5.0 g, 32 mmol) and 4-fluorobenzyl bromide (6.3 g, 33 mmol) in acetone (50 mL) containing potassium carbonate (5.3 g, 38 mmol) was heated under reflux for 2 h. After cooling to room temperature the mixture was filtered

- 29 -

and evaporated to leave the title compound (7.3 g, 86%) as a light yellow solid. MS: $m/e = 265.1 (M^+ + H)$.

f) 2-Fluoro-4-(4-fluoro-benzyloxy)-phenylamine (1:1) hydrochloride

A solution of 2-fluoro-4-(4-fluoro-benzyloxy)-nitrobenzene (7.3 g, 27 mmol) in ethyl acetate (160 mL) was hydrogenated in the presence of Pt/C (5%, 1.1 g) at room temperature for 12 h. The mixture was then filtered and HCl in diethylether was added. The resulting precipitate was filtered off and dried to afford the title compound (7.3 g, 99%) as a white solid. MS: $m/e = 236.2 (M^+ - Cl)$.

Example 40

10 N-[2-Fluoro-4-(4-fluoro-benzyloxy)-phenyl]-malonamide

A solution of N-[2-fluoro-4-(4-fluoro-benzyloxy)-phenyl]-malonamic acid methyl ester (150 mg, 0.45 mmol) and ammonium hydroxide (2.0 mL) in THF (1.0 mL) in a sealed tube was heated at 55 °C for 6 h. After cooling to room temperature the mixture was evaporated and the title compound (60 mg, 42%) was purified by trituration from diethylether. MS: $m/e = 321.2 (M^+ + H)$.

Example 41

N-[4-(3,5-Bis-trifluoromethyl-benzyloxy)-2-fluoro-phenyl]-malonamic acid methyl ester

a) 4-(3,5-Bis-trifluoromethyl-benzyloxy)-2-fluoro-nitrobenzene

20 As described for example 39e, 3-fluoro-4-nitrophenol (5 g, 32 mmol) was converted to the title compound (12.2 g, 99%) [using 3,5-bis(trifluoromethyl)benzyl bromide (10.3 g, 33 mmol) instead of 4-fluorobenzyl bromide] which was obtained as a light yellow solid. MS: $m/e = 383.0 (M^+)$.

b) 4-(3,5-Bis-trifluoromethyl-benzyloxy)-2-fluoro-phenylamine (1:1) hydrochloride

25 As described for example 39f, 4-(3,5-bis-trifluoromethyl-benzyloxy)-2-fluoro-nitrobenzene (12.2 g, 32 mmol) was converted to the title compound (8.3 g, 66%) which was obtained as light pink solid. MS: $m/e = 388.2 (M^+ - H)$.

c) N-[4-(3,5-Bis-trifluoromethyl-benzyloxy)-2-fluoro-phenyl]-malonamic acid methyl ester

30 As described for example 39d, 4-(3,5-bis-trifluoromethyl-benzyloxy)-2-fluoro-phenylamine (1:1) hydrochloride (500 mg, 1 mmol) was converted to the title compound (590 mg, 92%) which was obtained as an off-white solid. MS: $m/e = 454.2 (M^+ + H)$.

Example 42N-[4-(3,5-Bis-trifluoromethyl-benzyloxy)-2-fluoro-phenyl]-malonamide

- As described for example 40, N-[4-(3,5-bis-trifluoromethyl-benzyloxy)-2-fluoro-phenyl]-malonamic acid methyl ester (250 mg, 0.56 mmol) was converted to the title compound (84 mg, 35%) which was obtained as a white solid after purification by chromatography on silica gel eluting with DCM:MeOH (9:1). MS: m/e = 439.2 ($M^+ + H$).

Example 43N-[2-Fluoro-4-(4-trifluoromethyl-benzyloxy)-phenyl]-malonamic acid methyl ester

- 10 a) 2-Fluoro-4-(4-trifluoromethyl-benzyloxy)-nitrobenzene

As described for example 39e, 3-fluoro-4-nitrophenol (5 g, 32 mmol) was converted to the title compound (9.9 g, 99%) [using 3-(trifluoromethyl)benzyl bromide (8.0 g, 33 mmol) instead of 4-fluorobenzyl bromide] which was obtained as a light yellow solid. MS: m/e = 315.0 (M^+).

- 15 b) 2-Fluoro-4-(4-trifluoromethyl-benzyloxy)-phenylamine (1:1) hydrochloride

As described for example 39f, 2-fluoro-4-(4-trifluoromethyl-benzyloxy)-nitrobenzene (4.9 g, 15 mmol) was converted to the title compound (3.0 g, 60%) which was obtained as a grey solid.

- c) N-[2-Fluoro-4-(4-trifluoromethyl-benzyloxy)-phenyl]-malonamic acid methyl ester

- 20 As described for example 39d, 2-fluoro-4-(4-trifluoromethyl-benzyloxy)-phenylamine (1:1) hydrochloride (500 mg, 1 mmol) was converted to the title compound (700 mg, 98%) which was obtained as an off-white solid. MS: m/e = 386.2 ($M^+ + H$).

Example 44N-[2-Fluoro-4-(4-trifluoromethyl-benzyloxy)-phenyl]-malonamide

- 25 As described for example 40, N-[2-fluoro-4-(4-trifluoromethyl-benzyloxy)-phenyl]-malonamic acid methyl ester (250 mg, 0.65 mmol) was converted to the title compound (94 mg, 39%) which was obtained as a white solid after purification by chromatography on silica gel eluting with DCM:MeOH (9:1). MS: m/e = 371.2 ($M^+ + H$).

Example 45

- 30 N-[4-(3-Fluoro-benzyloxy)-3-methoxy-phenyl]-malonamic acid methyl ester

- 31 -

a) 1-(3-Fluoro-benzyloxy)-2-methoxy-4-nitro-benzene

Prepared in analogy to example 8 from 4-nitroguaiacol and 3-fluoro-benzylbromide. Colorless oil. Yield = 74 %. MS: $m/e = 277.1$ (M^+).

b) 4-(3-Fluoro-benzyloxy)-3-methoxy-phenylamine

5 Prepared in analogy to example 8b) by hydrogenation of 1-(3-fluoro-benzyloxy)-2-methoxy-4-nitro-benzene. The crude reaction mixture was used without purification in the next step.

c) N-[4-(3-Fluoro-benzyloxy)-3-methoxy-phenyl]-malonamic acid methyl ester

10 Prepared in analogy to example 20c) from 4-(3-fluoro-benzyloxy)-3-methoxy-phenylamine and methyl malonyl chloride. Yellowish solid. Yield = 89%. MS: $m/e = 348.2$ ($M^+ + H$).

Example 46N-[4-(3-Fluoro-benzyloxy)-3-methoxy-phenyl]-malonamide

15 Prepared in analogy to example 3 from N-[4-(3-fluoro-benzyloxy)-3-methoxy-phenyl]-malonamic acid methyl ester and 7M ammonia in methanol. Yield = 71 %. Colorless solid. MS: $m/e = 333.0$ ($M^+ + H$).

Example 47N-[4-(3-Fluoro-benzyloxy)-3-methyl-phenyl]-malonamic acid methyl estera) 1-(3-Fluoro-benzyloxy)-2-methyl-4-nitro-benzene

20 Prepared in analogy to example 8 from 2-methyl-4-nitrophenol and 3-fluoro-benzylbromide. Colorless oil. Yield = 62 %. MS: $m/e = 261.1$ (M^+).

b) 4-(3-Fluoro-benzyloxy)-3-methyl-phenylamine

25 Prepared in analogy to example 8b) by hydrogenation of 1-(3-fluoro-benzyloxy)-2-methyl-4-nitro-benzene. The crude reaction mixture was used without purification in the next step.

c) N-[4-(3-Fluoro-benzyloxy)-3-methyl-phenyl]-malonamic acid methyl ester

Prepared in analogy to example 20c) from 4-(3-fluoro-benzyloxy)-3-methyl-phenylamine and methyl malonyl chloride. Colorless solid. Yield = 98%. MS: $m/e = 332.0$ ($M^+ + H$).

30

Example 48N-[4-(3-Fluoro-benzyloxy)-3-methyl-phenyl]-malonamide

Prepared in analogy to example 3 from N-[4-(3-fluoro-benzyloxy)-3-methyl-phenyl]-malonamic acid methyl ester and 7M ammonia in methanol. Yield = 60 %. Colorless solid. MS: $m/e = 317.0$ ($M^+ + H$).

Example 49

5 N-[3-Chloro-4-(3-fluoro-benzyloxy)-phenyl]-malonamic acid methyl ester

a) 2-Chloro-1-(3-fluoro-benzyloxy)-4-nitro-benzene

Prepared in analogy to example 8 from 2-chloro-4-nitrophenol and 3-fluoro-benzylbromide. Colorless solid. Yield = 84 %. MS: $m/e = 281.0$ (M^+).

b) 3-Chloro-4-(3-fluoro-benzyloxy)-phenylamine

10 Prepared in analogy to example 8b) by hydrogenation of 2-chloro-1-(3-fluoro-benzyloxy)-4-nitro-benzene. The crude reaction mixture was used without purification in the next step.

c) N-[3-Chloro-4-(3-fluoro-benzyloxy)-phenyl]-malonamic acid methyl ester

15 Prepared in analogy to example 20c) from 3-chloro-4-(3-fluoro-benzyloxy)-phenylamine and methyl malonyl chloride. Colorless solid. Yield = 78%. MS: $m/e = 352.1$ ($M^+ + H$).

Example 50

N-[3-Chloro-4-(3-fluoro-benzyloxy)-phenyl]-malonamide

20 Prepared in analogy to example 3 from N-[3-chloro-4-(3-fluoro-benzyloxy)-phenyl]-malonamic acid methyl ester and 7M ammonia in methanol. Yield = 100 %. Colorless solid. MS: $m/e = 337.1$ ($M^+ + H$).

Example 51

Cyclopropane-1,1-dicarboxylic acid amide [2-fluoro-4-(4-fluoro-benzyloxy)-phenyl]-amide

25 A mixture of 2-fluoro-4-(4-fluoro-benzyloxy)-phenylamine (1:1) hydrochloride (272 mg, 1 mmol), 1-carboxycyclopropanecarboxamide (129 mg, 1 mmol), N-(3-dimethylaminopropyl)-N'-ethylcarbodiimide hydrochloride (EDCI, 211 mg, 1.1 mmol) and triethylamine (111 mg, 1.1 mmol) in DCM (7 mL) was stirred at rt for 2 h. Then the mixture was poured into water and the mixture extracted with DCM. The combined
30 organic extracts were then dried and evaporated to afford the title compound (130 mg, 38%) as a light brown solid after purification by chromatography on silica gel eluting with ethyl acetate : heptane (1:1 to 3:1). MS: $m/e = 347.0$ ($M^+ + H$).

Example 52N-[2-Fluoro-4-(4-fluoro-benzyloxy)-phenyl]-2,2-dimethyl-malonamic acid methyl ester

A mixture of N-[2-fluoro-4-(4-fluoro-benzyloxy)-phenyl]-malonamic acid methyl ester (335 mg, 1 mmol), sodium hydride (55%, 873 mg, 2 mmol) and iodomethane (213 mg, 1.5 mmol) were stirred at room temperature for 3 h. Then the mixture was poured into water and the mixture extracted with ethyl acetate. The combined organic extracts were then dried and evaporated to afford the title compound (83 mg, 23%) as a light yellow solid after purification by chromatography on silica gel eluting with ethyl acetate : heptane (1:1 to 3:1). MS: m/e = 364.1 ($M^+ + H$).

10

Example 53N-[2-Fluoro-4-(4-fluoro-benzyloxy)-phenyl]-malonamic acid ethyl ester

The title compound is prepared in analogy to example 1 from 2-fluoro-4-(4-fluoro-benzyloxy)-phenylamine (1:1) hydrochloride [using ethylmalonyl chloride instead of methylmalonyl chloride]. Light yellow solid. Yield = 84%. MS: m/e = 350.3 ($M^+ + H$).

15

Example 54N-[4-(3-Fluoro-benzyloxy)-3-formyl-phenyl]-malonamic acid methyl estera) 2-(3-Fluoro-benzyloxy)-5-nitro-benzaldehyde

Prepared in analogy to example 8 from 2-hydroxy-5-nitrobenzaldehyde and 3-fluoro-benzylbromide. Colorless solid. Yield = 95 %. MS: m/e = 275.1 (M^+).

20

b) 5-Amino-2-(3-fluoro-benzyloxy)-benzaldehyde

Prepared in analogy to example 8b) by hydrogenation of 2-(3-fluoro-benzyloxy)-5-nitro-benzaldehyde. The crude reaction mixture was used without purification in the next step.

c) N-[4-(3-Fluoro-benzyloxy)-3-formyl-phenyl]-malonamic acid methyl ester

25

Prepared in analogy to example 20c) from 5-amino-2-(3-fluoro-benzyloxy)-benzaldehyde and methyl malonyl chloride. Colorless solid. Yield = 3 %. MS: m/e = 344.1 ($M^+ - H$).

Example 552-Ethyl-N-[4-(3-fluoro-benzyloxy)-phenyl]-malonamide

30

A solution of sodium ethanolate is prepared by dissolving 46mg (2mmol) sodium in 5ml ethanol. 302mg (1mmol) of N-[4-(3-fluoro-benzyloxy)-phenyl]-malonamide is added, followed by 343mg (2mmol) of ethyl iodide. The reaction mixture is refluxed for

2.5h, hydrolysed and extracted with ethyl acetate. Chromatography (silica gel; dichloromethane / methanol) yields 45mg (14%) of a colorless solid. MS: $m/e = 331.3$ ($M^+ + H$).

Example 56

5 N-{4-[2-(4-Fluoro-phenyl)-vinyl]-phenyl}-malonamic acid methyl ester

a) (4-Nitro-benzyl)-phosphonic acid diethyl ester

10g (46.3mmol) of 4-nitrobenzylbromide and 10.03g (60.3mmol) of triethyl phosphite are dissolved in 25ml dimethylformamide. The reaction mixture is held at 155° C for 1.5h, diluted with 20ml water and extracted 3 times with ethyl acetate. The organic
10 phase is dried and concentrated to yield 8.35g (66%) of the crude title compound. MS: $m/e = 274.1$ ($M^+ + H$).

b) 4-[2-(4-Fluoro-phenyl)-vinyl]-1-nitro-benzene

2.63g of sodium hydride are added to 50ml of dimethylformamide. The mixture is cooled to 0° C and 16g (58.6mmol) of (4-nitro-benzyl)-phosphonic acid diethyl ester is
15 added portionwise. The solution is stirred at room temperature for 1.5 h, then cooled to -10°. A solution of 6.09g (50mmol) 4-fluoro-benzaldehyde in 10ml dimethylformamide is slowly added to the mixture at -10°. Stirring at room temperature for 45 min followed by addition of 250ml water yields a precipitate which is filtered and recrystallised from ethanol / water 85:15 to yield 10.81g (91%) of a yellow solid. MS: $m/e = 243.1$ (M^+).

20 c) 4-[2-(4-Fluoro-phenyl)-vinyl]-phenylamine

A solution of 7.3g (30mmol) of 4-[2-(4-fluoro-phenyl)-vinyl]-1-nitro-benzene in 250ml of ethanol is treated with 80ml of aqueous 25% hydrochloric acid. The mixture is heated at 110° C and 5g (42mmol) of tin is added portionwise. After stirring for 3h at 110° C the suspension is cooled to room temperature, neutralised by addition of aqueous
25 sodium hydroxide and extracted 3 times with dichloromethane. Drying over magnesium sulfate and evaporations yields 5.46 g (85%) of a colorless solid. MS: $m/e = 214.2$ ($M^+ + H$).

d) N-{4-[2-(4-Fluoro-phenyl)-vinyl]-phenyl}-malonamic acid methyl ester

Prepared in analogy to example 20c) from 4-[2-(4-fluoro-phenyl)-vinyl]-
30 phenylamine and methyl malonyl chloride. Colorless solid. Yield = 97 %. MS: $m/e = 314.0$ ($M^+ + H$).

Example 57

N-{4-[2-(4-Fluoro-phenyl)-vinyl]-phenyl}-malonamide

Prepared in analogy to example 3 from N-{4-[2-(4-fluoro-phenyl)-vinyl]-phenyl}-malonamic acid methyl ester and 7M ammonia in methanol. Yield = 89 %. Colorless solid. MS: $m/e = 299.2 (M^+ + H)$.

Example 58

5 N-{4-[2-(3-Fluoro-phenyl)-vinyl]-phenyl}-malonamic acid methyl ester

a) 4-[2-(3-Fluoro-phenyl)-vinyl]-1-nitro-benzene

Prepared in analogy to example 56b) from (4-nitro-benzyl)-phosphonic acid diethyl ester and 3-fluoro-benzaldehyde. Yellow solid. Yield = 82%. MS: $m/e = 243.0 (M^+)$.

10 b) 4-[2-(3-Fluoro-phenyl)-vinyl]-phenylamine

2.41g (10mmol) of 4-[2-(3-fluoro-phenyl)-vinyl]-1-nitro-benzene is dissolved in 25ml of ethyl acetate and treated with 240mg of platinum 5% on charcoal. Hydrogenation is performed at room temperature and normal pressure for 4h. The catalyst is filtered off and the filtrate concentrated. The residue is crystallised from diethyl ether / heptane to yield 1.32g (62%) of an orange solid. MS: $m/e = 213.1 (M^+)$.

c) N-{4-[2-(3-Fluoro-phenyl)-vinyl]-phenyl}-malonamic acid methyl ester

Prepared in analogy to example 20c) from 4-[2-(3-fluoro-phenyl)-vinyl]-phenylamine and methyl malonyl chloride. Colorless solid. Yield = 99 %. MS: $m/e = 313.9 (M^+ + H)$.

20

Example 59

N-{4-[2-(3-Fluoro-phenyl)-vinyl]-phenyl}-malonamide

Prepared in analogy to example 3 from N-{4-[2-(3-fluoro-phenyl)-vinyl]-phenyl}-malonamic acid methyl ester and 7M ammonia in methanol. Yield = 68 %. Colorless solid. MS: $m/e = 299.2 (M^+ + H)$.

25

Example 60

N-{4-[2-(4-Fluoro-phenyl)-ethyl]-phenyl}-malonamic acid methyl ester

a) 4-[2-(4-Fluoro-phenyl)-ethyl]-phenylamine

3.46g of 4-[2-(4-fluoro-phenyl)-vinyl]-phenylamine is dissolved in 100ml of tetrahydrofuran, treated with 350mg of palladium 10% on charcoal and hydrogenated at room temperature and normal pressure for about 4h. The catalyst is filtered off and the filtrate concentrated to yield 3.46g (99%) of a yellow solid. MS: $m/e = 216.9 (M^+ + H)$.

b) N-{4-[2-(4-Fluoro-phenyl)-ethyl]-phenyl}-malonamic acid methyl ester

Prepared in analogy to example 20c) from 4-[2-(3-fluoro-phenyl)-vinyl]-phenylamine and methyl malonyl chloride. Colorless solid. Yield = 99 %. MS: $m/e = 316.0$ ($M^+ + H$).

Example 61

5 N-{4-[2-(4-Fluoro-phenyl)-ethyl]-phenyl}-malonamide

Prepared in analogy to example 3 from N-{4-[2-(4-fluoro-phenyl)-ethyl]-phenyl}-malonamic acid methyl ester and 7M ammonia in methanol. Yield = 94 %. Colorless solid. MS: $m/e = 301.1$ ($M^+ + H$).

Example 62

10 N-{4-[2-(3-Fluoro-phenyl)-ethyl]-phenyl}-malonamic acid methyl ester

a) 4-[2-(3-Fluoro-phenyl)-ethyl]-phenylamine

Prepared in analogy to example 60a) by hydrogenation of 4-[2-(3-fluoro-phenyl)-vinyl]-phenylamine. Yellow solid. Yield = 100%. MS: $m/e = 215.1$ (M^+).

b) N-{4-[2-(3-Fluoro-phenyl)-ethyl]-phenyl}-malonamic acid methyl ester

15 Prepared in analogy to example 20c) from 4-[2-(3-fluoro-phenyl)-ethyl]-phenylamine and methyl malonyl chloride. Colorless solid. Yield = 95 %. MS: $m/e = 316.0$ ($M^+ + H$).

Example 63

N-{4-[2-(4-Methoxy-phenyl)-vinyl]-phenyl}-malonamic acid methyl ester

20 a) 4-[2-(4-Methoxy-phenyl)-vinyl]-phenylamine

Prepared in analogy to example 56b) from (4-nitro-benzyl)-phosphonic acid diethyl ester and 4-methoxy-benzaldehyde. Yellow solid. Yield = 69%. MS: $m/e = 225.9$ (M^+).

b) N-{4-[2-(4-Methoxy-phenyl)-vinyl]-phenyl}-malonamic acid methyl ester

25 Prepared in analogy to example 20c) from 4-[2-(4-methoxy-phenyl)-vinyl]-phenylamine and methyl malonyl chloride. Colorless solid. Yield = 66 %. MS: $m/e = 326.1$ ($M^+ + H$).

Example 64

N-{4-[2-(4-Chloro-phenyl)-ethyl]-phenyl}-malonamic acid methyl ester

a) 4-[2-(4-Chloro-phenyl)-vinyl]-1-nitro-benzene

Prepared in analogy to example 56b) from (4-nitro-benzyl)-phosphonic acid diethyl ester and 4-chloro-benzaldehyde. Yellow solid. Yield = 95%. MS: $m/e = 259.1$ (M^+).

b) 4-[2-(4-Chloro-phenyl)-ethyl]-phenylamine

A solution of 5.0g (19.2mmol) 4-[2-(4-chloro-phenyl)-vinyl]-1-nitro-benzene in
5 100ml ethyl acetate is treated with 200mg of platinum 5% on charcoal. Hydrogenation is performed at room temperature and normal pressure overnight. The catalyst is filtered off and the filtrate evaporated to yield 3.17g (71%) of a slightly yellow solid. MS: $m/e = 231.9$ ($M^+ + H$).

c) N-{4-[2-(4-Chloro-phenyl)-ethyl]-phenyl}-malonamic acid methyl ester

10 Prepared in analogy to example 20c) from 4-[2-(4-chloro-phenyl)-ethyl]-phenylamine and methyl malonyl chloride. Colorless solid. Yield = 99 %. MS: $m/e = 332.1$ ($M^+ + H$).

Example 65

N-{4-[2-(4-Chloro-phenyl)-ethyl]-phenyl}-malonamide

15 Prepared in analogy to example 3 from N-{4-[2-(4-chloro-phenyl)-ethyl]-phenyl}-malonamic acid methyl ester and 7M ammonia in methanol. Yield = 65 %. Colorless solid. MS: $m/e = 317.1$ ($M^+ + H$).

Example 66

2-Amino-N-[2-fluoro-4-(4-fluoro-benzyloxy)-phenyl]-acetamide (1:1) hydrochloride

20 a) {[2-Fluoro-4-(4-fluoro-benzyloxy)-phenyl]carbonyl}-methyl}-carbamic acid tert-butyl ester

A mixture of 2-fluoro-4-(4-fluoro-benzyloxy)-phenylamine (1:1) hydrochloride (272 mg, 1 mmol), Boc-glycine (263 mg, 1.5 mmol), N,N' -dicyclohexylcarbodiimide (206 mg, 1 mmol) and pyridine (127 mg, 1.6 mmol) in ethyl acetate (6 mL) was stirred at room
25 temperature for 3 h. After this time, the reaction mixture was poured into sodium hydrogen carbonate (saturated), and the mixture extracted with ethyl acetate. The combined organic layers were then washed with brine and dried. Filtration and evaporation afforded the title compound (180 mg, 46%) as a light brown solid after chromatography on silica gel eluting with ethyl acetate : heptane (1:1 to 4:1). MS: $m/e =$
30 393.1 ($M^+ + H$).

b) 2-Amino-N-[2-fluoro-4-(4-fluoro-benzyloxy)-phenyl]-acetamide (1:1) hydrochloride

A solution of {[2-fluoro-4-(4-fluoro-benzyloxy)-phenylcarbamoyl]-methyl}-carbamic acid tert-butyl ester (148 mg, 0.4 mmol) in HCl dioxane (4 M, 3 mL) was stirred at room temperature for 1 h. The resulting precipitate was filtered off to afford the title compound (112 mg, 90%) as a white solid. MS: $m/e = 293.1$ ($M^+ - Cl$).

5

Example 67

(R)-2-Amino-N-[2-fluoro-4-(4-fluoro-benzyloxy)-phenyl]-propionamide

As described for example 66a and 66b, 2-fluoro-4-(4-fluoro-benzyloxy)-phenylamine (1:1) hydrochloride (200 mg, 0.7 mmol) was converted to the title compound (127 mg, 39%) [using Boc-D-Ala-OH instead of Boc-glycine] which was obtained as a white solid.

10 MS: $m/e = 307.0$ ($M^+ - Cl$).

Example 68

2-Amino-N-[2-fluoro-4-(4-fluoro-benzyloxy)-phenyl]-propionamide

As described for example 66a and 66b, 2-fluoro-4-(4-fluoro-benzyloxy)-phenylamine (1:1) hydrochloride (200 mg, 0.7 mmol) was converted to the title compound (41 mg, 15%) [using Boc-DL-Ala-OH instead of Boc-glycine] which was obtained as a white solid. MS: $m/e = 306.9$ ($M^+ + H$).

15

Example 69

1-[2-Fluoro-4-(4-fluoro-benzyloxy)-phenylcarbamoyl]-2S-methyl-propyl-ammonium; chloride

As described for example 66a and 66b, 2-fluoro-4-(4-fluoro-benzyloxy)-phenylamine (1:1) hydrochloride (200 mg, 0.7 mmol) was converted to the title compound (47 mg, 12%) [using Boc-L-Valine instead of Boc-glycine] which was obtained as a white solid. MS: $m/e = 335.2$ ($M^+ + H$).

20

Example 70

25 2-Acetyl-amino-N-[2-fluoro-4-(4-fluoro-benzyloxy)-phenyl]-acetamide

A solution of 2-amino-N-[2-fluoro-4-(4-fluoro-benzyloxy)-phenyl]-acetamide (1:1) hydrochloride (50 mg, 0.15 mmol), acetyl chloride (14.4 mg, 0.18 mmol), triethylamine (31 mg, 0.30 mmol) was stirred at 0 °C for 4 h. Then the mixture was diluted with water and the mixture extracted with DCM. The organic layer was then washed with brine, dried, filtered and evaporated to leave the title compound (50 mg, 99%) as a white solid. MS: $m/e = 335.2$ ($M^+ + H$).

30

Example 71

(R)-2-Acetyl-amino-N-[2-fluoro-4-(4-fluoro-benzyloxy)-phenyl]-propionamide

- 39 -

As described for example 70, (R)-2-amino-N-[2-fluoro-4-(4-fluoro-benzyloxy)-phenyl]-propionamide (50 mg, 0.15 mmol) was converted to the title compound (43 mg, 76%) which was obtained as a white solid. MS: $m/e = 349.4$ ($M^+ + H$).

Example 72

5 N-[2-Fluoro-4-(4-fluoro-benzyloxy)-phenyl]-2-formylamino-acetamide

Step 1: A mixture of acetic anhydride (40 mg, 0.39 mmol) and formic acid (22 mg, 0.48 mmol) was stirred at 0 °C and then heated uat 60 °C for 1 h. Step 2: During this time, 2-amino-N-[2-fluoro-4-(4-fluoro-benzyloxy)-phenyl]-acetamide (1:1) hydrochloride (50 mg, 0.15 mmol) was extracted with DCM and sodium hydrogen carboante (saturated) and the organic layer evaporated. Then to a mixture of the acetic formic anhydride (Step 1) in dry THF (0.5 mL) was added a solution of the amine (Step 2) in dry THF (1 mL) at room temperature and the resulting mixture stirred for 10 min. Then the reaction mixture was evaporated leave the title compound (44.1 mg, 91%) as a white solid. MS: $m/e = 321.1$ ($M^+ + H$).

15

Example 73

(R)-N-[2-Fluoro-4-(4-fluoro-benzyloxy)-phenyl]-2-formylamino-propionamide

As described for example 72, (R)-2-amino-N-[2-fluoro-4-(4-fluoro-benzyloxy)-phenyl]-propionamide (50 mg, 0.15 mmol) was converted to the title compound (16 mg, 30%) which was obtained as a white solid. MS: $m/e = 333.1$ ($M^+ - H$).

20

Example 74

2-Amino-N-[2-fluoro-4-(3-fluoro-benzyloxy)-phenyl]-acetamide hydrochloride (1:1)

As described for example 66a and 66b, 2-fluoro-4-(3-fluoro-benzyloxy)-phenylamine (1:1) hydrochloride (500 mg, 2.1 mmol) was converted to the title compound (300 mg, 44%) which was obtained as a white solid. MS: $m/e = 293.1$ ($M^+ - Cl$).

25

Example 75

2-Acetylamino-N-[2-fluoro-4-(3-fluoro-benzyloxy)-phenyl]-acetamide

As described for example 70, 2-fluoro-4-(3-fluoro-benzyloxy)-phenylamine (1:1) hydrochloride (150 mg, 0.45 mmol) was converted to the title compound (127 mg, 84%) which was obtained as a white solid. MS: $m/e = 335.2$ ($M^+ + H$).

30

Example 76

N-[2-Fluoro-4-(3-fluoro-benzyloxy)-phenyl]-2-formylamino-acetamide

As described for example 72, 2-fluoro-4-(3-fluoro-benzyloxy)-phenylamine (1:1) hydrochloride (133 mg, 0.4 mmol) was converted to the title compound (115 mg, 89%) which was obtained as a white solid. MS: $m/e = 321.1$ ($M^+ + H$).

- 40 -

Example 77

2-Amino-N-[2-fluoro-4-(4-trifluoromethyl-benzyloxy)-phenyl]-acetamide (1:1) hydrochloride

As described for example 66a and 66b, 2-fluoro-4-(4-trifluoromethyl-benzyloxy)-phenylamine (1:1) hydrochloride (600 mg, 2.1 mmol) was converted to the title compound (155 mg, 20%) which was obtained as a white solid. MS: m/e = 343.1 (M^+ -Cl).

Example 78

2-Acetylamino-N-[2-fluoro-4-(4-trifluoromethyl-benzyloxy)-phenyl]-acetamide

As described for example 70, 2-fluoro-4-(4-trifluoromethyl-benzyloxy)-phenylamine (1:1) hydrochloride (120 mg, 0.32 mmol) was converted to the title compound (119 mg, 98%) which was obtained as a white solid. MS: m/e = 385.2 (M^+ +H).

Example 79

2-Amino-N-[4-(3,5-bis-trifluoromethyl-benzyloxy)-2-fluoro-phenyl]-acetamide (1:1) hydrochloride

As described for example 66a and 66b, 4-(3,5-bis-trifluoromethyl-benzyloxy)-2-fluoro-phenylamine (1:1) hydrochloride (500 mg, 1.4 mmol) was converted to the title compound (180 mg, 30%) which was obtained as a white solid. MS: m/e = 411.2 (M^+ -Cl).

Example 80

2-Acetylamino-N-[4-(3,5-bis-trifluoromethyl-benzyloxy)-2-fluoro-phenyl]-acetamide

As described for example 70, 4-(3,5-bis-trifluoromethyl-benzyloxy)-2-fluoro-phenylamine (1:1) hydrochloride (130 mg, 0.29 mmol) was converted to the title compound (107 mg, 81%) which was obtained as a white solid. MS: m/e = 453.2 (M^+ +H).

Example 81

2-Amino-N-[4-(3-fluoro-benzyloxy)-phenyl]-acetamide; hydrochloridea) {[4-(3-Fluoro-benzyloxy)-phenyl]carbamoyl}-methyl}-carbamic acid tert-butyl ester

A mixture of 0.345g (1.59mmol) 4-(3-fluoro-benzyloxy)-phenylamine and 0.278g (1.59mmol) of N-tert-Butyloxyglycine is dissolved in 10ml of ethyl acetate, cooled to 0° and 0.328g (1.59mmol) N,N'-dicyclohexylcarbodiimide is added at once. The resulting slurry is stirred for additional 3h at room temperature and filtered. The filtrate is washed 3 times with saturated sodium carbonate solution, dried and concentrated. The residue is

- 41 -

trituated with 20ml diethylether to give 0.353g (59%) of the title compound as a colorless solid. . MS: m/e = 375.4 ($M^+ + H$).

b) 2-Amino-N-[4-(3-fluoro-benzyloxy)-phenyl]-acetamide; hydrochloride

- A slurry of 0.166g (0.44mmol) of {[4-(3-fluoro-benzyloxy)-phenylcarbonyl]-methyl}-
 5 carbamic acid tert-butyl ester in 1ml diethyl ether is treated with 3ml of a saturated
 solution of gaseous hydrochloric acid in diethyl ether. The mixture is refluxed for 4h,
 poured on water and made basic by addition of a saturated sodium carbonate solution.
 Extraction with ethyl acetate yields a yellowish solid which is dissolved in about 1ml of
 diethyl ether and treated with a few drops of a saturated solution of gaseous hydrochloric
 10 acid in diethyl ether. The precipitate is filtered to give 0.051g (37%) of a colorless solid.
 MS: m/e = 275.2 ($M^+ + H$).

Example A

Tablets of the following composition are produced in a conventional manner:

	<u>mg/Tablet</u>
15 Active ingredient	100
Powdered lactose	95
White corn starch	35
Polyvinylpyrrolidone	8
Na carboxymethylstarch	10
20 Magnesium stearate	2
Tablet weight	<u>250</u>

Example B

Tablets of the following composition are produced in a conventional manner:

	<u>mg/Tablet</u>
25 Active ingredient	200
Powdered lactose	100
White corn starch	64
Polyvinylpyrrolidone	12
Na carboxymethylstarch	20
30 Magnesium stearate	4
Tablet weight	<u>400</u>

Example C

Capsules of the following composition are produced:

	<u>mg/Capsule</u>
Active ingredient	50
5 Crystalline lactose	60
Microcrystalline cellulose	34
Talc	5
Magnesium stearate	1
Capsule fill weight	<u>150</u>

- 10 The active ingredient having a suitable particle size, the crystalline lactose and the microcrystalline cellulose are homogeneously mixed with one another, sieved and thereafter talc and magnesium stearate are admixed. The final mixture is filled into hard gelatine capsules of suitable size.


Example D

- 15 An injection solution may have the following composition and is manufactured in usual manner:

Active substance	1.0 mg
1 N HCl	20.0 µl
acetic acid	0.5 mg
20 NaCl	8.0 mg
phenol	10.0 mg
1 N NaOH	q.s. ad pH 5
H ₂ O	q.s. ad 1 ml

Claims

1. Compounds of the general formula



wherein

R^1 is halogen, halogen-(C_1 - C_6)-alkyl, cyano, C_1 - C_6 -alkoxy or halogen-(C_1 - C_6)-alkoxy;

R^{21} , R^{22} , R^{23} and R^{24} independently from each other are selected from the group consisting of hydrogen, (C_1 - C_6)-alkyl, halogen, halogen-(C_1 - C_6)-alkyl, hydroxy, C_1 - C_6 -alkoxy or $-CHO$;

R^3 is hydrogen or C_1 - C_3 -alkyl;

R^4 , R^5 independently from each other are selected from the group consisting of hydrogen, C_1 - C_6 -alkyl, C_1 - C_6 -alkoxy or $-COO(C_1-C_6)alkyl$;
or R^4 and R^5 form together with the C-atom to which they are attached a C_3 - C_7 -cycloalkyl ring;

R^6 is $-CO-NR^7R^8$, $-COO(C_1-C_6)alkyl$, $-CN$, $-NR_2$ or $-NHC(O)R$;

R^7 and R^8 independently from each other are selected from the group consisting of hydrogen, C_1 - C_6 -alkyl, NH_2 or hydroxy;

R is hydrogen or C_1 - C_6 -alkyl;

n is 0, 1, 2 or 3;

X is $-CHRO$, $-OCHR-$, $-CH_2S-$, $-SCH_2-$, $-CH_2CH_2-$, $-CH=CH-$ or $-C\equiv C-$;

and pharmaceutically active acid addition salts thereof,

with the proviso that

N-(4-benzyloxy-2,6-dimethyl-phenyl)-2-diethylamino-acetamide and hydrochloride salt thereof,

2-diethylamino-N-[4-(4-methoxy-benzyloxy)-2,6-dimethyl-phenyl]-acetamide,
 N-[4-(4-chloro-benzyloxy)-2,6-dimethyl-phenyl]-2-diethylamino-acetamide,
 N-(4-benzyloxy-2-methyl-phenyl)-2-diethylamino-acetamide,
 N-(4-benzyloxy-5-isopropyl-2-methyl-phenyl)-2-diethylamino-acetamide,
 N-(4-benzyloxy-phenyl)-2-diethylamino-acetamide,
 N-(4-benzyloxy-2,6-dichloro-phenyl)-2-diethylamino-acetamide,
 N-(4-benzyloxy-phenyl)-2-dimethylamino-acetamide hydrochloride,
 N-(4-benzyloxy-phenyl)-2-dimethylamino-propionamide hydrochloride,
 N-[4-(4-chloro-benzyloxy)-phenyl]-2-dimethylamino-acetamide hydrochloride,
 N-[4-(4-chloro-benzyloxy)-phenyl]-2-dimethylamino-propionamide hydrochloride,
 N-(4-benzyloxy-phenyl)-2-diethylamino-acetamide hydrochloride,
 N-(4-benzyloxy-phenyl)-2-diethylamino-propionamide hydrochloride,
 N-[4-(4-chloro-benzyloxy)-phenyl]-2-diethylamino-acetamide hydrochloride,
 N-[4-(4-chloro-benzyloxy)-phenyl]-2-diethylamino-propionamide hydrochloride,
 N-(4-benzyloxy-2-methyl-phenyl)-2-diethylamino-acetamide,
 N-(4-benzyloxy-2-methyl-phenyl)-2-(butyl-methyl-amino)-acetamide,
 N-(4-benzyloxy-2-methyl-phenyl)-2-dibutylamino-acetamide,
 N-(4-benzyloxy-2-methyl-phenyl)-2-dipropylamino-acetamide,
 N-(4-benzyloxy-2-methyl-phenyl)-2-propylamino-acetamide,
 N-(4-benzyloxy-2-methyl-phenyl)-2-hexylamino-acetamide,
 N-(4-benzyloxy-2-methyl-phenyl)-2-diethylamino-propionamide,
 N-(4-benzyloxy-2-methyl-phenyl)-2-butylamino-propionamide,
 N-(4-benzyloxy-2-methyl-phenyl)-2-propylamino-propionamide,
 N-(4-benzyloxy-2,6-dimethyl-phenyl)-2-butylamino-acetamide hydrochloride salt thereof,
 N-(4-benzyloxy-2,6-dimethyl-phenyl)-2-dibutylamino-acetamide hydrochloride salt thereof,
 N-(4-benzyloxy-2,6-dimethyl-phenyl)-2-dipropylamino-acetamide, and
 N-(4-benzyloxy-2-methylphenyl)-2-(butylamino)acetamide;
 and compounds of the formula I wherein

n is 0, 1, 2 or 3; and

R¹ is halogen, cyano, C₁-C₆-alkoxy, -(CH₂)_m-C(halogen)₃ with m ranging from 0 to 3
 or -O-(CH₂)_m-C(halogen)₃ with m ranging from 0 to 3; and

R²¹, R²², R²³, and R²⁴ are all hydrogen; and

X is -(CH₂)_m- with m=2, -CH=CH-, or -C≡C-; and

R³ is hydrogen or C₁-C₃-alkyl; and


R⁶ is -CN; and

R⁴, R⁵ is hydrogen;

are excluded.

2. Compounds according to claim 1, of the general formula

- 45 -



wherein

R^1 is halogen, halogen-(C_1 - C_6)-alkyl, cyano, C_1 - C_6 -alkoxy or halogen-(C_1 - C_6)-alkoxy;

R^{21} , R^{22} , R^{23} and R^{24} independently from each other are selected from the group consisting of hydrogen, (C_1 - C_6)-alkyl, halogen, halogen-(C_1 - C_6)-alkyl, hydroxy, C_1 - C_6 -alkoxy or $-CHO$;

R^3 is hydrogen or C_1 - C_3 -alkyl;

R^4 , R^5 independently from each other are selected from the group consisting of hydrogen, C_1 - C_6 -alkyl, C_1 - C_6 -alkoxy or $-COO(C_1-C_6)alkyl$;
or R^4 and R^5 form together with the C-atom to which they are attached a C_3 - C_7 -cycloalkyl ring;

R^6 is $-CO-NR^7R^8$; $-COO(C_1-C_6)alkyl$, $-CN$, $-NR_2$ or $-NHC(O)R$;

R^7 and R^8 independently from each other are selected from the group consisting of hydrogen, C_1 - C_6 -alkyl, NH_2 or hydroxy;

R is hydrogen or C_1 - C_6 -alkyl;

n is 0, 1, 2 or 3;

X is $-CHRO$, $-OCHR-$, $-CH_2S-$ or $-SCH_2-$;


and pharmaceutically active acid addition salts thereof,

with the proviso that

N-(4-benzyloxy-2,6-dimethyl-phenyl)-2-diethylamino-acetamide and hydrochloride salt thereof,
2-diethylamino-N-[4-(4-methoxy-benzyloxy)-2,6-dimethyl-phenyl]-acetamide,
N-[4-(4-chloro-benzyloxy)-2,6-dimethyl-phenyl]-2-diethylamino-acetamide,
N-(4-benzyloxy-2-methyl-phenyl)-2-diethylamino-acetamide,
N-(4-benzyloxy-5-isopropyl-2-methyl-phenyl)-2-diethylamino-acetamide,
N-(4-benzyloxy-phenyl)-2-diethylamino-acetamide,

N-(4-benzyloxy-2,6-dichloro-phenyl)-2-diethylamino-acetamide,
 N-(4-benzyloxy-phenyl)-2-dimethylamino-acetamide hydrochloride,
 N-(4-benzyloxy-phenyl)-2-dimethylamino-propionamide hydrochloride,
 N-[4-(4-chloro-benzyloxy)-phenyl]-2-dimethylamino-acetamide hydrochloride,
 N-[4-(4-chloro-benzyloxy)-phenyl]-2-dimethylamino-propionamide hydrochloride,
 N-(4-benzyloxy-phenyl)-2-diethylamino-acetamide hydrochloride,
 N-(4-benzyloxy-phenyl)-2-diethylamino-propionamide hydrochloride,
 N-[4-(4-chloro-benzyloxy)-phenyl]-2-diethylamino-acetamide hydrochloride,
 N-[4-(4-chloro-benzyloxy)-phenyl]-2-diethylamino-propionamide hydrochloride,
 N-(4-benzyloxy-2-methyl-phenyl)-2-diethylamino-acetamide,
 N-(4-benzyloxy-2-methyl-phenyl)-2-(butyl-methyl-amino)-acetamide,
 N-(4-benzyloxy-2-methyl-phenyl)-2-dibutylamino-acetamide,
 N-(4-benzyloxy-2-methyl-phenyl)-2-dipropylamino-acetamide,
 N-(4-benzyloxy-2-methyl-phenyl)-2-propylamino-acteamide,
 N-(4-benzyloxy-2-methyl-phenyl)-2-hexylamino-acetamide,
 N-(4-benzyloxy-2-methyl-phenyl)-2-diethylamino-propionamide,
 N-(4-benzyloxy-2-methyl-phenyl)-2-butylamino-propionamide,
 N-(4-benzyloxy-2-methyl-phenyl)-2-propylamino-propionamide,
 N-(4-benzyloxy-2,6-dimethyl-phenyl)-2-butylamino-acetamide hydrochloride salt thereof,
 N-(4-benzyloxy-2,6-dimethyl-phenyl)-2-dibutylamino-acetamide hydrochloride salt thereof,
 N-(4-benzyloxy-2,6-dimethyl-phenyl)-2-dipropylamino-acetamide, and
 N-(4-benzyloxy-2-methylphenyl)-2-(butylamino)acetamide
 are excluded.

3. Compounds according to claim 1, of the general formula



wherein

R^1 is halogen, halogen-(C_1 - C_6)-alkyl, cyano, C_1 - C_6 -alkoxy or halogen-(C_1 - C_6)-alkoxy;

R^{21} , R^{22} , R^{23} and R^{24} independently from each other are selected from the group consisting of hydrogen, (C_1 - C_6)-alkyl, halogen, halogen-(C_1 - C_6)-alkyl, hydroxy, C_1 - C_6 -alkoxy or -CHO;

- 47 -

R^3 is hydrogen or C_1 - C_3 -alkyl;

R^4, R^5 independently from each other are selected from the group consisting of hydrogen, C_1 - C_6 -alkyl, C_1 - C_6 -alkoxy or $-COO(C_1-C_6)alkyl$; or R^4 and R^5 form together with the C-atom to which they are attached a C_3 - C_7 -cycloalkyl ring;

R^6 is $-CO-NR^7R^8$, $-COO(C_1-C_6)alkyl$, $-NR_2$ or $-NHC(O)R$;

R^7 and R^8 independently from each other are selected from the group consisting of hydrogen, C_1 - C_6 -alkyl, NH_2 or hydroxy;

R is hydrogen or C_1 - C_6 -alkyl;

n is 0, 1, 2 or 3;

X is $-CH_2CH_2-$, $-CH=CH-$ or $-C\equiv C-$;


and pharmaceutically active acid addition salts thereof,

with the proviso that

N-(4-benzyloxy-2,6-dimethyl-phenyl)-2-diethylamino-acetamide and hydrochloride salt thereof,
 2-diethylamino-N-[4-(4-methoxy-benzyloxy)-2,6-dimethyl-phenyl]-acetamide,
 N-[4-(4-chloro-benzyloxy)-2,6-dimethyl-phenyl]-2-diethylamino-acetamide,
 N-(4-benzyloxy-2-methyl-phenyl)-2-diethylamino-acetamide,
 N-(4-benzyloxy-5-isopropyl-2-methyl-phenyl)-2-diethylamino-acetamide,
 N-(4-benzyloxy-phenyl)-2-diethylamino-acetamide,
 N-(4-benzyloxy-2,6-dichloro-phenyl)-2-diethylamino-acetamide,
 N-(4-benzyloxy-phenyl)-2-dimethylamino-acetamide hydrochloride,
 N-(4-benzyloxy-phenyl)-2-dimethylamino-propionamide hydrochloride,
 N-[4-(4-chloro-benzyloxy)-phenyl]-2-dimethylamino-acetamide hydrochloride,
 N-[4-(4-chloro-benzyloxy)-phenyl]-2-dimethylamino-propionamide hydrochloride,
 N-(4-benzyloxy-phenyl)-2-diethylamino-acetamide hydrochloride,
 N-(4-benzyloxy-phenyl)-2-diethylamino-propionamide hydrochloride,
 N-[4-(4-chloro-benzyloxy)-phenyl]-2-diethylamino-acetamide hydrochloride,
 N-[4-(4-chloro-benzyloxy)-phenyl]-2-diethylamino-propionamide hydrochloride,
 N-(4-benzyloxy-2-methyl-phenyl)-2-diethylamino-acetamide,
 N-(4-benzyloxy-2-methyl-phenyl)-2-(butyl-methyl-amino)-acetamide,
 N-(4-benzyloxy-2-methyl-phenyl)-2-dibutylamino-acetamide,
 N-(4-benzyloxy-2-methyl-phenyl)-2-dipropylamino-acetamide,
 N-(4-benzyloxy-2-methyl-phenyl)-2-propylamino-acetamide,
 N-(4-benzyloxy-2-methyl-phenyl)-2-hexylamino-acetamide,
 N-(4-benzyloxy-2-methyl-phenyl)-2-diethylamino-propionamide,
 N-(4-benzyloxy-2-methyl-phenyl)-2-butylamino-propionamide,
 N-(4-benzyloxy-2-methyl-phenyl)-2-propylamino-propionamide,

N-(4-benzyloxy-2,6-dimethyl-phenyl)-2-butylamino-acetamide hydrochloride salt thereof, N-(4-benzyloxy-2,6-dimethyl-phenyl)-2-dibutylamino-acetamide hydrochloride salt thereof, N-(4-Benzyloxy-2,6-dimethyl-phenyl)-2-dipropylamino-acetamide, and N-(4-benzyloxy-2-methylphenyl)-2-(butylamino)acetamide are excluded.

4. Compounds according to any one of claims 1 to 3, of the general formula



wherein

R^1 is halogen, halogen-(C_1 - C_6)-alkyl, cyano, C_1 - C_6 -alkoxy or halogen-(C_1 - C_6)-alkoxy;

R^{21} , R^{22} , R^{23} and R^{24} independently from each other are selected from the group consisting of hydrogen, (C_1 - C_6)-alkyl, halogen, halogen-(C_1 - C_6)-alkyl, hydroxy, C_1 - C_6 -alkoxy or -CHO;

R^3 is hydrogen or C_1 - C_3 -alkyl;

R^4 , R^5 independently from each other are selected from the group consisting of hydrogen, C_1 - C_6 -alkyl, C_1 - C_6 -alkoxy or -COO(C_1 - C_6)alkyl; or R^4 and R^5 form together with the C-atom to which they are attached a C_3 - C_7 -cycloalkyl ring;

R^6 is -CO-NR⁷R⁸; -COO(C_1 - C_6)-alkyl or -NHC(O)R;

R^7 and R^8 independently from each other are selected from the group consisting of hydrogen, C_1 - C_6 -alkyl, NH₂ or hydroxy;


R is hydrogen or C_1 - C_6 -alkyl;

n is 0, 1, 2 or 3;

X is -CHRO, -OCHR-, -CH₂S-, -SCH₂-, -CH₂CH₂-, -CH=CH- or -C≡C-;

and pharmaceutically active acid addition salts thereof.

5. Compounds of the general formula Ia



wherein

R^1 is halogen, halogen-(C₁-C₆)-alkyl, cyano, C₁-C₆-alkoxy or halogen-(C₁-C₆)-alkoxy;

R^{21} , R^{22} , R^{23} and R^{24} independently from each other are selected from the group consisting of hydrogen and halogen;

R^3 is hydrogen or C₁-C₃-alkyl;

R^4 is hydrogen or C₁-C₆-alkyl;

R^5 is hydrogen or C₁-C₆-alkyl;

or R^4 and R^5 together form a C₃-C₇-cycloalkyl ring;

R^6 is -CO-NR⁷R⁸;
-COOR⁹; or
-CN;

R^7 and R^8 independently from each other are hydrogen, methyl or ethyl;

R^9 is C₁-C₆-alkyl; and

n is 1, 2 or 3;

and pharmaceutically active acid addition salts thereof.

6. Compounds of formula I according to claim 1, wherein X is CH₂O, R^1 is fluorine or trifluoromethyl and R^6 is -COOCH₃.

7. Compounds according to claim 6, which are

N-[4-(3-fluoro-benzyloxy)-phenyl]-malonamic acid methyl ester,
 N-[3-fluoro-4-(3-fluoro-benzyloxy)-phenyl]-malonamic acid methyl ester,
 N-[4-(4-fluoro-benzyloxy)-phenyl]-malonamic acid methyl ester,
 N-[2-fluoro-4-(3-fluoro-benzyloxy)-phenyl]-malonamic acid methyl ester,
 N-[4-(2,4-difluoro-benzyloxy)-phenyl]-malonamic acid methyl ester,
 N-[4-(2-fluoro-benzyloxy)-phenyl]-malonamic acid methyl ester,
 N-[4-(2,4,5-trifluoro-benzyloxy)-phenyl]-malonamic acid methyl ester,
 N-[2-fluoro-4-(4-fluoro-benzyloxy)-phenyl]-malonamic acid methyl ester,
 N-[4-(3,5-bis-trifluoromethyl-benzyloxy)-2-fluoro-phenyl]-malonamic acid methyl ester,
 N-[4-(3-fluoro-benzyloxy)-3-methyl-phenyl]-malonamic acid methyl ester or
 N-[3-chloro-4-(3-fluoro-benzyloxy)-phenyl]-malonamic acid methyl ester.

8. Compounds of formula I according to claim 1, wherein X is CH_2O and R^6 is $-\text{CONH}_2$.

9. Compounds according to claim 8, which compounds are

cyclopropane-1,1-dicarboxylic acid amide [4-(3-fluoro-benzyloxy)-phenyl]-amide,

N-[4-(3-fluoro-benzyloxy)-phenyl]-malonamide,

N-[4-(3-fluoro-benzyloxy)-phenyl]-2-methyl-malonamide,

N-[3-fluoro-4-(3-fluoro-benzyloxy)-phenyl]-malonamide,

N-[4-(4-fluoro-benzyloxy)-phenyl]-malonamide,

N-[4-(2,4-difluoro-benzyloxy)-phenyl]-malonamide,

N-[4-(2,4,5-trifluoro-benzyloxy)-phenyl]-malonamide,

N-[4-(2-fluoro-benzyloxy)-phenyl]-malonamide,

N-(4-benzyloxy-phenyl)-malonamide,

N-[4-(4-chloro-benzyloxy)-phenyl]-malonamide,

N-[4-(3-fluoro-benzyloxy)-2-hydroxy-phenyl]-malonamide,

N-[2-fluoro-4-(4-fluoro-benzyloxy)-phenyl]-malonamide,

N-[4-(3-fluoro-benzyloxy)-3-methyl-phenyl]-malonamide,

N-[3-chloro-4-(3-fluoro-benzyloxy)-phenyl]-malonamide or

cyclopropane-1,1-dicarboxylic acid amide [2-fluoro-4-(4-fluoro-benzyloxy)-phenyl]-amide.

10. Compounds of formula I according to claim 1, wherein X is $-\text{CH}_2\text{O}-$ and R^6 is $-\text{NHCOCH}_3$ or $-\text{NHCOH}$.

11. Compounds according to claim 10, wherein the compounds are
2-Acetylamino-N-[2-fluoro-4-(4-fluoro-benzyloxy)-phenyl]-acetamide,
2-Acetylamino-N-[2-fluoro-4-(3-fluoro-benzyloxy)-phenyl]-acetamide,
N-[2-Fluoro-4-(4-fluoro-benzyloxy)-phenyl]-2-formylamino-acetamide or
N-[2-Fluoro-4-(3-fluoro-benzyloxy)-phenyl]-2-formylamino-acetamide.


12. Compounds of formula I according to claim 1, wherein X is $-\text{CH}_2\text{O}-$ and R^6 is $-\text{NH}_2$.

13. A compounds according to claim 12, wherein the compound is
2-amino-N-[2-fluoro-4-(4-fluoro-benzyloxy)-phenyl]-acetamide.

14. Compounds of formula I according to claim 1, wherein X is $-\text{CH}=\text{CH}-$ and R^6 is $-\text{COOCH}_3$ or $-\text{CONH}_2$.


15. Compounds according to claim 14, wherein the compounds are
N-{4-[2-(4-fluoro-phenyl)-vinyl]-phenyl}-malonamic acid methyl ester,
N-{4-[2-(3-fluoro-phenyl)-vinyl]-phenyl}-malonamide,
N-{4-[2-(4-fluoro-phenyl)-vinyl]-phenyl}-malonamide or
N-{4-[2-(3-fluoro-phenyl)-vinyl]-phenyl}-malonamic acid methyl ester.

16. A process for the manufacture of a compound of formula I according to any one of claims 1 to 15, which process comprises
reacting a compound of formula




with a compound of formula

- 52 -



to obtain a compound of formula



and, if desired, converting a compound of formula I into a pharmaceutically acceptable salt.


17. A compound of formula I according to any one of claims 1 to 15, when manufactured by a process according to claim 16.

18. A medicament containing one or more compounds as claimed in any one of claims 1 to 15 and pharmaceutically acceptable excipients for use in the treatment and prevention of diseases which are mediated by monoamine oxidase B inhibitors.

19. The medicament according to claim 18 for use in the treatment and prevention of Alzheimer's disease and senile dementia.

20. A compound of formula I according to any one of claims 1 to 12 as well as its pharmaceutically acceptable salts for the treatment or prevention of diseases.

21. The use of a compound of general formula I



wherein

R^1 is halogen, halogen-(C₁-C₆)-alkyl, cyano, C₁-C₆-alkoxy or halogen-(C₁-C₆)-alkoxy;

R^{21} , R^{22} , R^{23} and R^{24} independently from each other are selected from the group consisting of hydrogen, (C₁-C₆)-alkyl, halogen, halogen-(C₁-C₆)-alkyl, hydroxy, C₁-C₆-alkoxy or -CHO;

R^3 is hydrogen or C₁-C₃-alkyl;

R^4 , R^5 independently from each other are selected from the group consisting of hydrogen, C₁-C₆-alkyl, C₁-C₆-alkoxy or -COO(C₁-C₆)-alkyl;

or R^4 and R^5 form together with the C-atom to which they are attached a C₃-C₇-cycloalkyl ring;

R^6 is -CO-NR⁷R⁸; -COO(C₁-C₆)-alkyl, -CN, -NR₂ or -NHC(O)R;

R^7 and R^8 independently from each other are selected from the group consisting of hydrogen, C₁-C₆-alkyl, NH₂ or hydroxy;


R is hydrogen or C₁-C₆-alkyl;

n is 0, 1, 2 or 3;

X is -CHRO, -OCHR-, -CH₂S-, -SCH₂-, -CH₂CH₂-, -CH=CH- or -C≡C-;

as well as its pharmaceutically acceptable salts for the manufacture of medicaments for the treatment and prevention of diseases which are mediated by monoamine oxidase B inhibitors.

22. The use of a compound of general formula Ia



wherein

R^1 is halogen, halogen-(C₁-C₆)-alkyl, cyano, C₁-C₆-alkoxy or halogen-(C₁-C₆)-alkoxy;

- 54 -

R^{21} , R^{22} , R^{23} and R^{24} independently from each other are selected from the group consisting of hydrogen and halogen;

R^3 is hydrogen or C_1 - C_3 -alkyl;

R^4 is hydrogen or C_1 - C_6 -alkyl;

R^5 is hydrogen or C_1 - C_6 -alkyl;

or R^4 and R^5 together form a C_3 - C_7 -cycloalkyl ring;

R^6 is $-\text{CO}-\text{NR}^7\text{R}^8$;
 $-\text{COOR}^9$; or
 $-\text{CN}$;

R^7 and R^8 independently from each other are hydrogen, methyl or ethyl;

R^9 is C_1 - C_6 -alkyl; and

n is 1, 2 or 3;

as well as its pharmaceutically acceptable salts for the manufacture of medicaments for the treatment and prevention of diseases which are mediated by monoamine oxidase B inhibitors.

23. The use according to claim 21 or 22, wherein the disease is Alzheimer's disease or senile dementia.
