

LTBP COMPLEX-SPECIFIC INHIBITORS OF TGF β AND USES THEREOF

RELATED APPLICATIONS

[001] This international application claims the benefit of and priority to U.S. Provisional Application No. 62/798,927, filed January 30, 2019, the contents of which are expressly incorporated herein by reference in entirety.

BACKGROUND

[002] Transforming growth factor beta (TGF β) superfamily of growth factors are involved in a number of signaling cascades that regulate diverse biological processes including, but not limited to: inhibition of cell growth, tissue homeostasis, extracellular matrix (ECM) remodeling, endothelial to mesenchymal transition, cell migration and invasion, and immune modulation/suppression, as well as mesenchymal to epithelial transition. In relation to ECM remodeling, TGF β signaling may increase fibroblast populations and ECM deposition (*e.g.*, collagen). In the immune system, TGF β ligand modulates T regulatory cell function and maintenance of immune precursor cell growth and homeostasis. In normal epithelial cells, TGF β is a potent growth inhibitor and promoter of cellular differentiation. However, as tumors develop and progress, they frequently lose their negative growth response to TGF β . In this setting, TGF β may become a promoter of tumor development due to its ability to stimulate angiogenesis, alter the stromal environment, and induce local and systemic immunosuppression. For these and other reasons, TGF β has been a therapeutic target for a number of clinical indications. Despite much effort made to date by a number of groups, clinical development of a TGF β therapeutic has been challenging.

[003] Observations from preclinical studies, including in rats and dogs, have revealed certain toxicities associated with inhibition of TGF β *in vivo*. Moreover, although several TGF β inhibitors have been developed to date, most clinical programs targeting TGF β have been discontinued due to side effects or risk of toxicity.

[004] For example, Anderton *et al.* (Toxicology Pathology, 39: 916-24, 2011) reported that small molecule inhibitors of TGF β type I (ALK5) receptor induced heart valve lesions characterized by hemorrhage, inflammation, degeneration and proliferation of valvular interstitial cells in a preclinical animal model. The toxicity was observed in all heart valves at all doses tested. Frazier *et al.* (Toxicology Pathology, 35: 284-295, 2007) reported that administration of the small molecule inhibitor of TGF β type I (ALK5) receptor GW788388 induced physeal dysplasia in rats.

[005] Stauber *et al.* (J. Clin. Practice 4:3, 2014) reported that a chronic (≥ 3 months) administration of the inhibitor of TGF β receptor I kinase, LY2157299, which is being investigated for certain cancer

treatments, caused multiple organ toxicities involving the cardiovascular, gastrointestinal, immune, bone/cartilage, reproductive, and renal systems, in rats and dogs.

[006] Fresolimumab (GC1008), a “pan” TGF β antibody capable of neutralizing all human isoforms of TGF β , has been reported to induce an epithelial hyperplasia of the gingiva, bladder, and of the nasal turbinate epithelium after multiple administrations in studies with cynomolgus macaques (Lonning *et al.*, Current Pharmaceutical Biotechnology 12: 2176-89, 2011). Similarly, a variety of skin rashes/lesions, gingival bleeding and fatigue have been reported in clinical trials after administration of multiple doses of the drug. The most notable adverse reaction to fresolimumab includes the induction of cutaneous keratoacanthomas and/or squamous cell carcinomas in human cancer patients (see, for example: Lacouture *et al.*, 2015, Cancer Immunol Immunother, 64: 437-46; Stevenson *et al.*, 2013, OncoImmunology, 2:8, e26218; and Lonning *et al.*, 2011). Additional evidence from a clinical trial suggests that in some cases this antibody may accelerate tumor progression (Stevenson *et al.*, 2013, OncoImmunology, 2:8, e26218).

[007] Thus, new methods and compositions for modulating TGF β signaling are necessary that can be used to effectively and safely treat diseases and disorders involving TGF β , including, for example, cancer, fibrosis and inflammation.

[008] With an increasing recognition of potentially dangerous adverse effects associated with broad inhibition of TGF β , a number of groups have more recently turned to identifying inhibitors that target a *subset* - but not all - of the isoforms and still retain sufficient efficacy. For example, WO 2016/161410 discloses neutralizing antibodies that bind both TGF β 1 and TGF β 2 (*i.e.*, TGF β 1/2 inhibitors). WO 2006/116002 provides neutralizing antibodies that bind both TGF β 1 and TGF β 3 (*i.e.*, TGF β 1/3 inhibitors), albeit preferentially to the former. In addition to traditional monoclonal antibodies, some groups are developing engineered fusion proteins that function as so-called “ligand traps” (see, for example, WO 2018/158727, WO 2018029367 and WO 2018129331), at least some of which may be selective for TGF β 1/3. Another class of TGF β 1/3 inhibitors include inhibitors of alpha-V (α v) integrins such as antibodies against α v β 6, which is an integrin known to activate both TGF β 1 and TGF β 3 (*i.e.*, TGF β 1/3). Yet others continue to pursue “better” pan-inhibitors that inhibit all three isoforms (*i.e.*, TGF β 1/2/3 or pan-inhibitors) (see, for example, WO 2018/134681).

[009] From an efficacy standpoint, however, the prevailing view of the field remains to be that it is advantageous to inhibit multiple isoforms of TGF β to achieve therapeutic effects, and to accommodate this, toxicity management by “careful dosing regimen” is suggested as a solution (Brennan *et al.* (2018) mAbs, 10:1, 1-17).

[010] Recently, Applicant described isoform-selective TGF β 1 inhibitors which were demonstrated to be both safe and efficacious in animal models (see, for example: WO 2017/156500 and WO 2018/129329, incorporated by reference), supporting the notion that selectively targeting the TGF β 1

isoform, as opposed to broadly antagonizing all TGF β isoforms, may provide an advantageous approach to achieving efficacy with acceptable toxicity.

[011] Whilst the observed safety profile achieved by selective inhibition of TGF β 1 at doses that were shown efficacious in vivo is a promising step towards developing a TGF β 1 inhibitor for clinical applications, identification of TGF β 1 inhibitors that are capable of selectively affecting a defined subset of TGF β 1 effects (*e.g.*, TGF β inhibitors that are selective to LTBP-presented complexes) remained elusive. More recently, Applicant demonstrated that such “LTBP context-specific” inhibitors can be generated (WO 2019/023661, incorporated herein by reference) using the methods previously described by Applicant (see, for example, WO 2014/074532 and WO 2014/182676). However, the LTBP-selective TGF β 1 inhibitors described in the aforementioned international publication showed modest affinities and inhibitory activities, coupled with suboptimal cross-species reactivity.

SUMMARY OF THE INVENTION

[012] The present disclosure provides improved TGF β inhibitors capable of selectively targeting matrix-associated proTGF β complexes, such as LTBP1-proTGF β 1 and LTBP3-proTGF β 1.

[013] These inhibitors bind and inhibit LTBP1- and/or LTBP3-presented proTGF β at high affinities (at least nanomolar range) but do not bind and inhibit immune cell-associated TGF β , *e.g.*, GARP- and/or LRRC33-presented proTGF β 1, or the binding is below meaningful levels (*e.g.*, at least 50 times affinities for the LTBP complexes over GARP or LRRC33 complex). Thus, these inhibitors can selectively inhibit activation of TGF β in a context-dependent manner, such that they selectively bind, thereby inhibiting the TGF β signaling axis associated with the ECM. In particular, the present disclosure includes selective inhibitors of matrix-associated (*e.g.*, LTBP1 and/or LTBP3-associated) TGF β activation. In some embodiments, such inhibitors specifically bind a particular isoform of TGF β (*e.g.*, proTGF β 1, proTGF β 2, and/or proTGF β 3) associated with LTBP1 and/or LTBP3, thus also providing TGF β isoform specificity. In a particular embodiment, such inhibitors specifically bind to LTBP1/3-proTGF β 1. In any of the embodiments of the present invention, such inhibitors do not inhibit activation of TGF β 1 associated with immune cell function, mediated by GARP and/or LRRC33. The improved antibodies encompassed by the present disclosure have affinities towards human LTBP1-proTGF β 1 and/or human LTBP3-proTGF β 1 in at least a nanomolar range (*i.e.*, 1×10^{-9} M to 10×10^{-9} M). In some embodiments, such antibodies also have affinities towards murine LTBP1-proTGF β 1 and/or murine LTBP3-proTGF β 1 in at least a nanomolar range (*i.e.*, 1×10^{-9} M to 10×10^{-9} M).

[014] Rationale for the therapeutic use of a TGF β 1 inhibitor that does not target the GARP-proTGF β 1 complex on regulatory T cells is at least threefold:

[015] First, regulatory T cells play a crucial role in maintaining immune tolerance to self-antigens and in preventing autoimmune disease. Since Tregs generally suppress, dampen or downregulate induction and proliferation of effector T cells, systemic inhibition of this function may lead to overactive or exaggerated immune responses in the host by disabling the “break” that is normally provided by Treg cells. Thus, the approach taken here (*e.g.*, TGFβ1 inhibition without disabling Treg function) is aimed to avoid the risk of eliciting autoimmunity. Furthermore, patients who already have a propensity for developing over-sensitive immune responses or autoimmunity may be particularly at risk of triggering or exacerbating such conditions, without the availability of normal Treg function; and therefore, the inhibitors that selectively target the matrix TGFβ1 may advantageously minimize such risk.

[016] Second, evidence suggests that an alteration in the Th17/Treg ratio leads to an imbalance in pro-fibrotic Th17 cytokines, which correlate with severity of fibrosis, such as liver fibrosis (see, for example, Shoukry *et al.* (2017) J Immunol 198 (1 Supplement): 197.12). The present inventors reasoned that perturbation of the GARP arm of TGFβ1 function may directly or indirectly exacerbate fibrotic conditions.

[017] Third, regulatory T cells are indispensable for immune homeostasis and the prevention of autoimmunity. It was reasoned that, particularly for a TGFβ1 inhibition therapy intended for a long-term or chronic administration, it would be desirable to avoid potential side effects stemming from perturbation of normal Treg function in maintaining immune homeostasis (reviewed in, for example, Richert-Spuhler and Lund (2015) Prog Mol Biol Transl Sci. 136: 217–243). This strategy is at least in part aimed to preserve normal immune function, which is required, *inter alia*, for combatting infections.

[018] To this end, the inventors of the present disclosure set out to generate isoform-specific, context-selective inhibitors of TGFβ1 that selectively target matrix-associated TGFβ1 activation but not immune cell-associated TGFβ1 activation.

[019] Technical challenges that exist to date include limited ability to discern and selectively modulate these subpools of TGFβ1 present in various contexts (or “niches”) *in vivo*.

[020] In an effort to address this challenge, the present inventors have identified isoform-specific monoclonal antibodies that bind the latent TGFβ1 prodomain, with no detectable binding to latent TGFβ2 or TGFβ3, and that inhibit integrin-mediated activation of latent TGFβ1 *in vitro* with the context-dependency as described herein. The discovery and characterization of such antibodies was made possible, at least in part, by the development of context-dependent cell-based assays of TGFβ1 activation. In the process of this novel assay development and validation, it was demonstrated that, like the αVβ6 integrin, αVβ8 can also activate LTBP1-proTGFβ1. It was further demonstrated that, similar to the LTBP1 complex, LTBP3-proTGFβ1 can be activated by αVβ6. Antibodies discovered

by screening in these assays revealed a class of antibodies that binds and inhibits TGF β 1 only when presented by LTBP1 or LTBP3. Such LTBP-specific antibodies do not inhibit TGF β 1 in the context of the immune-associated TGF β 1 presenters GARP and LRRC33. Such antibodies are therapeutic candidates for the treatment of disorders including, *e.g.*, fibrotic conditions, and could allow chronic dosing that would avoid TGF β -related immune system activation. Methods of selecting a context-specific or context-independent TGF β 1 inhibitor for various fibrotic conditions are also provided herein.

[021] Accordingly, in one aspect, the invention provides isoform-specific TGF β antibodies, or antigen-binding fragments thereof, characterized in that they bind selectively to an LTBP1-TGF β 1 complex and/or a LTBP3-TGF β 1 complex with a $K_D \leq 50$ nM. In one embodiment, the invention provides isoform-specific TGF β antibodies, or antigen-binding fragments thereof, characterized in that they bind selectively to an LTBP1-TGF β 1 complex and/or a LTBP3-TGF β 1 complex with a $K_D \leq 25$ nM. In one embodiment, the invention provides isoform-specific TGF β antibodies, or antigen-binding fragments thereof, characterized in that they bind selectively to an LTBP1-TGF β 1 complex and/or a LTBP3-TGF β 1 complex with a $K_D \leq 10$ nM. In one embodiment, the invention provides an isolated antibody, or antigen-binding portion thereof, that selectively binds to a LTBP1-proTGF β 1 complex and a LTBP3-proTGF β 1 complex, wherein the antibody, or antigen-binding portion thereof, does not bind to one or more of the following targets: (a) LTBP1 alone; (b) proTGF β 1 alone; (c) a GARP-proTGF β 1 complex; and (d) a LRRC33-proTGF β 1 complex. In further embodiments, the invention provides isoform-specific TGF β antibodies, or antigen-binding fragments thereof, characterized in that they bind selectively to an LTBP1-TGF β 1 complex and/or a LTBP3-TGF β 1 complex with a $K_D < 5$ nM.

[022] In one aspect, the invention provides inhibitors of extracellular matrix-associated TGF β activation, which selectively bind a LTBP1/3-presented proTGF β latent complex. In one embodiment, the inhibitor does not inhibit immune cell-associated TGF β 1 activation, for example, immune cell-associated TGF β 1 activation that results from activation of a GARP-presented proTGF β 1 latent complex. In exemplary embodiments, the inhibitor is an antibody, or antigen-binding portion thereof.

[023] In other aspects, the invention provides TGF β antibodies, or antigen-binding fragments thereof, characterized in that they bind selectively to an LTBP1-TGF β complex and/or a LTBP3-TGF β complex. In some embodiments, the antibodies, or antigen-binding fragments thereof, selectively bind to LTBP1-TGF β 1. In some embodiments, such antibodies bind both human and murine counterparts.

[024] In one aspect, the invention provides an isolated antibody, or antigen-binding portion thereof, that selectively binds an LTBP1-proTGF β latent complex and/or an LTBP3-proTGF β latent complex,

thereby modulating release of mature TGF β growth factor from the latent complex, wherein the antibody, or antigen-binding portion thereof, does not bind mature TGF β 1 alone or a GARP-proTGF β 1 latent complex. In one embodiment, the antibody, or antigen-binding portion thereof, does not bind an LRRC33-proTGF β 1 latent complex. Alternatively, in one embodiment, the antibody, or antigen-binding portion thereof, binds an LRRC33-proTGF β 1 latent complex.

[025] In some embodiments, the antibody, or antigen-binding portion thereof, is specific to an LTBP1-proTGF β 1 latent complex. In other embodiments, the antibody, or antigen-binding portion thereof, is specific to an LTBP3-proTGF β 1 latent complex. In one embodiment, the antibody, or antigen-binding portion thereof, binds an LTBP1-proTGF β 1 complex and/or a LTBP3-proTGF β 1 complex with a dissociation constant (K_D) of at least about 10^{-8} M. In one embodiment, the antibody, or antigen-binding portion thereof, binds a human LTBP1-proTGF β 1 complex and/or a human LTBP3-proTGF β 1 complex with a K_D of < 50 nM as measured in a suitable in vitro binding assay such as Bio-Layer Interferometry (BLI). In one embodiment, the antibody, or antigen-binding portion thereof, binds a human LTBP1-proTGF β 1 complex and/or a human LTBP3-proTGF β 1 complex with a K_D of < 10 nM as measured in a suitable in vitro binding assay such as Bio-Layer Interferometry (BLI). In one embodiment, the antibody, or antigen-binding portion thereof, binds a mouse LTBP1-proTGF β 1 complex and/or a mouse LTBP3-proTGF β 1 complex with a K_D of < 50 nM as measured in a suitable in vitro binding assay such as Bio-Layer Interferometry (BLI). In one embodiment, the antibody, or antigen-binding portion thereof, binds a mouse LTBP1-proTGF β 1 complex and/or a mouse LTBP3-proTGF β 1 complex with a K_D of < 10 nM as measured in a suitable in vitro binding assay such as Bio-Layer Interferometry (BLI).

[026] The present disclosure further provides antibodies and antigen binding fragments thereof, which selectively bind an LTBP1-proTGF β complex and/or an LTBP3-proTGF β complex and have one or more yet further advantageous properties. Indeed, the inventors surprisingly found that such antibodies could be provided which bind a human LTBP1-proTGF β complex and a human LTBP3-proTGF β complex with high affinity, and advantageously slow dissociation rates, while also being cross-reactive with mouse LTBP1-proTGF β complex and mouse LTBP3-proTGF β complex and displaying no significant binding to human GARP-proTGF β complex (or indeed to human LRRC33-proTGF β complex).

[027] Further still, antibodies disclosed herein (including antibodies having one or more, or even all of the aforementioned advantageous properties) exhibit potent inhibition of TGF β 1 signaling in cell-based assays, and significantly reduce markers of fibrosis and TGF β signaling in multiple animal models of fibrosis.

[028] Thus, in some embodiments, the antibody, or antigen-binding fragment thereof binds a human LTBP1-proTGF β 1 complex and/or a human LTBP3-TGF β 1 complex with a K_D of < 5 nM as measured by BLI, and has one or more of the following properties:

- i) is cross-reactive with mouse LTBP1-proTGFβ1 complex;
- ii) is cross-reactive with mouse LTBP3-proTGFβ1 complex;
- iii) binds a mouse LTBP1-proTGFβ1 complex with a K_D of < 10 nM as measured by BLI;
- iv) binds a mouse LTBP3-proTGFβ1 complex with a K_D of < 10 nM as measured by BLI;
- v) binds a human LTBP1-proTGFβ1 complex and/or a human LTBP3-TGFβ1 complex with a K_D that is at least 50 times lower than the K_D when binding to a human GARP-proTGFβ1 complex under the same assay conditions;
- vi) does not show detectable binding to a human GARP-proTGFβ1 complex, as measured by BLI, under the same assay conditions as used to measure binding to human LTBP1-proTGFβ1 complex and/or a human LTBP3-TGFβ1 complex;
- vii) does not show detectable binding to an LRRC33-proTGFβ1 complex (*e.g.*, a human LRRC33-proTGFβ1 complex) as measured by BLI, under the same assay conditions as used to measure binding to human LTBP1-proTGFβ1 complex and/or human LTBP3-TGFβ1 complex.

[029] In some embodiments, the antibody or antigen-binding fragment has at least properties (i)-(v) above, and optionally (vii). In some embodiments, the antibody or antigen-binding fragment has at least properties (i)-(iv) and (vi) above, and optionally (vii). In some embodiments, the antibody or antigen-binding fragment has at least properties (i), (iii) and (v) above, and optionally (vii). In some embodiments, the antibody or antigen-binding fragment has at least properties (ii), (iv) and (v) above, and optionally (vii). In some embodiments, the antibody or antigen-binding fragment has at least properties (i), (iii) and (vi) above, and optionally (vii). In some embodiments, the antibody or antigen-binding fragment has at least properties (ii), (iv) and (vi) above, and optionally (vii). In some embodiments, the antibody or antigen-binding fragment has at least properties (i)-(iii) and (v) above, and optionally (vii). In some embodiments, the antibody or antigen-binding fragment has at least properties (i)-(iii) and (vi) above, and optionally (vii).

[030] In some preferred embodiments, the antibody or antigen-binding fragment binds a human LTBP1-proTGFβ1 complex and a human LTBP3-TGFβ1 complex with a K_D of < 5 nM as measured by BLI, and has all of the above properties (i)-(vii).

[031] The antibody or antigen-binding fragment may selectively bind a LTBP1/3-presented proTGFβ latent complex and inhibit extracellular matrix-associated TGFβ activation.

[032] Further still, further advantageous isoform-selective inhibitors of TGFβ1 activation may include monoclonal antibodies (including immunoglobulins and antigen-binding fragments or portions thereof) that exhibit slow dissociation rates (*i.e.*, off-rates, k_{OFF}). Thus, the invention is further based on the recognition that treatment of chronic and progressive disease such as fibrosis may

require inhibitors with superior durability, which may be reflected on the dissociation rate of such antibody.

[033] The affinity of an antibody to its antigen is typically measured as the equilibrium dissociation constant, or K_D . The ratio of the experimentally measured off- and on-rates (k_{OFF} / k_{ON}) can be used to calculate the K_D value. The k_{OFF} value represents the antibody dissociation rate, which indicates how quickly it dissociates from its antigen, whilst the k_{ON} value represents the antibody association rate which provides how quickly it binds to its antigen. The latter is typically concentration-dependent, while the former is concentration-independent. The K_D value relates to the concentration of antibody (the amount of antibody needed for a particular experiment) and so the lower the K_D value (lower concentration) and thus the higher the affinity of the antibody. With respect to a reference antibody, a higher affinity antibody may have a lower k_{OFF} rate, a higher k_{ON} rate, or both.

[034] Both the k_{OFF} and k_{ON} rates contribute to the overall affinity of a particular antibody to its antigen, and relative importance or impact of each component may depend on the mechanism of action of the antibody. For example, neutralizing antibodies, which bind mature growth factors (*e.g.*, soluble, transient TGF β 1 ligand liberated from a latent complex), must compete with the endogenous high-affinity receptors for ligand binding *in vivo*. Because the ligand-receptor interaction is a local event and because the ligand is short-lived, such antibodies must be capable of rapidly targeting and sequestering the soluble growth factor before the ligand finds its cellular receptor – thereby activating the TGF β 1 signaling pathway – in the tissue. Therefore, for ligand-targeting neutralizing antibodies to be potent, the ability to bind the target growth factor fast, *i.e.*, high association rates (k_{ON}), may be especially important.

[035] By contrast, Applicant reasoned that antibodies that inhibit the TGF β 1 signaling by preventing the activation (*e.g.*, release) of mature growth factor from the latent complex (“activation inhibitors”) may preferentially benefit from having slow dissociation rates once the antibody is engaged with the target antigen (*e.g.*, proTGF β 1 complexes). Unlike neutralizing antibodies, such antibodies do not directly compete with cellular receptors; rather, they work upstream of the signaling by targeting inactive precursor forms (*e.g.*, latent proTGF β 1 complexes) that remain dormant within a tissue environment thereby preemptively preventing the activation of TGF β 1. Such antibodies may exert their inhibitory activity by preventing mature growth factor from being liberated from the latent complex. For example, such antibodies may function like a “clamp” to lock the active growth factor in the prodomain cage structure to keep it in an inactive (*e.g.*, “latent”) state. Indeed, structural analyses, including epitope mapping, provided insight into the molecular mechanism underlining the ability of these antibodies to block TGF β 1 activation. In this regard, the Latency Lasso region of the prodomain may be a particularly useful target.

[036] Upon target engagement, antibodies that are able to remain bound to the target (*e.g.*, dissociate very slowly from the latent complex) are expected to be advantageous in achieving superior *in vivo* potency, due to enhanced durability of effects and/or avidity. Based on this recognition,

Applicant of the present disclosure sought to identify isoform-selective activation inhibitors of TGFβ1 with particularly low k_{OFF} values as compared to previously described antibodies. Thus, according to the invention, preferred antibodies have high affinities primarily attributable to a slow dissociation rate (k_{OFF}), as opposed to fast association rate (k_{ON}). Accordingly, in some embodiments, the antibody, or antigen-binding fragment thereof binds a human LTBP1-proTGFβ1 complex and/or a human LTBP3-TGFβ1 complex with a K_D of < 5 nM as measured by BLI, and has one or more of the following properties (which may be in addition to one of properties (i)-(vii), or combinations thereof set out above):

(viii) low dissociation rates (k_{OFF}) of $\leq 5 \times 10^{-4}$ (1/s), when binding human LTBP1-proTGFβ1 complex and/or human LTBP3-TGFβ1 complex (*e.g.*, as measured by a suitable in vitro binding/kinetics assay, such as by BLI, *e.g.*, Octet-based systems); and/or

(ix) long half-binding time ($t_{1/2}$) of ≥ 45 minutes when bound to human LTBP1-proTGFβ1 and/or human LTBP3-proTGFβ1 complex (*e.g.*, as measured by SPR).

[037] In some preferred embodiments, the antibody or antigen binding fragment comprises the following six CDRs:

- a) CDR-H1 comprising the amino acid sequence FTFRSYVMH;
- b) CDR-H2 comprising the amino acid sequence VISHEGS(X₁)KYYADSVKG, wherein: X₁ is L or G; and
- c) CDR-H3 comprising the amino acid sequence A(X₁)PRIAARRGGFG(X₂), wherein: X₁ is V, R or L; and X₂ is Y, S or T;
- d) CDR-L1 comprising the amino acid sequence TRS(X₁)G(X₂)ID(X₃)NYVQ, wherein, X₁ is S or H; X₂ is N, L, S or A; and X₃ is N, D or Y;
- e) CDR-L2 comprising the amino acid sequence ED(X₁)(X₂)RPS, wherein: X₁ is N, F or A; and X₂ is Q, I or V; and
- f) CDR-L3 comprising the amino acid sequence Q(X₁)YD(X₂)(X₃)(X₄)Q(X₅)VV, wherein: X₁ is S or G; X₂ is S, F, Y, D, H or W; X₃ is N, D or S; X₄ is N, A, L, E or T; and X₅ is G, R, A or L.

[038] In some preferred embodiments, the antibody or antigen-binding fragment thereof competes or cross-competes with an antibody having a heavy chain variable region sequence as set forth in SEQ ID NO: 318 and light chain variable region sequence as set forth in SEQ ID NO: 319 (*e.g.*, Ab42). The antibody may comprise a heavy chain variable region having an amino acid sequence that is at least 90% identical to SEQ ID NO: 318 and a light chain variable region having an amino acid sequence that is at least 90% identical to SEQ ID NO: 319.

[039] The antibody or antigen-binding fragment thereof provided herein may, in some preferred embodiments, comprise the following six CDRs (*e.g.*, those of Ab42):

CDR-H1 comprising the amino acid sequence FTFRSYVMH (SEQ ID NO: 166);

CDR-H2 comprising the amino acid sequence VISHEGSLKYYADSVKG (SEQ ID NO: 167);

CDR-H3 comprising the amino acid sequence ARPRIAARRGGFGY (SEQ ID NO: 168);

CDR-L1 comprising the amino acid sequence TRSSGNIDNNYVQ (SEQ ID NO: 169);

CDR-L2 comprising the amino acid sequence EDNQRPS (SEQ ID NO: 170); and

CDR-L3 comprising the amino acid sequence QSYDYDTQGVV (SEQ ID NO: 171).

[040] The antibody or antigen-binding fragment may further comprise a heavy chain variable region having an amino acid sequence that is at least 95% identical (optionally at least 98% identical) to SEQ ID NO: 318 and a light chain variable region having an amino acid sequence that is at least 95% identical (optionally at least 98% identical) to SEQ ID NO: 319.

[041] In some alternative embodiments, the antibody, or antigen-binding fragment thereof, comprises the following six CDRs:

- a) CDR-H1 comprising the amino acid sequence GSIRSSSYWG;
- b) CDR-H2 comprising the amino acid sequence SISYSATTYY;
- c) CDR-H3 comprising the amino acid sequence A(X₁)DPSYDS(X₂)AGM(X₃)V, wherein: X₁ is S or G; X₂ is A or I; and X₃ is D or Q;
- d) CDR-L1 comprising the amino acid sequence RAS(X₁)(X₂)IS(X₃)YLN, wherein: X₁ is K or Q; X₂ is V or S; and X₃ is S or Y;
- e) CDR-L2 comprising the amino acid sequence (X₁)AS(X₂)(X₃)QS, wherein: X₁ is Y, A or S; X₂ is S or N; and X₃ is L or R;
- f) CDR-L3 comprising the amino acid sequence QQ(X₁)(X₂)D(X₃)P(X₄)T, wherein: X₁ is S or G; X₂ is F or N; X₃ is W or F; and X₄ is F or L.

[042] In some embodiments, the antibody or antigen-binding fragment thereof competes or cross-competes with an antibody having a heavy chain variable region sequence as set forth in SEQ ID NO: 360 and light chain variable region sequence as set forth in SEQ ID NO: 361 (*e.g.*, Ab63). The antibody may comprise a heavy chain variable region having an amino acid sequence that is at least 90% identical to SEQ ID NO: 360 and a light chain variable region having an amino acid sequence that is at least 90% identical to SEQ ID NO: 361.

[043] The antibody or antigen-binding fragment thereof provided herein may comprise the following six CDRs (*e.g.*, those of Ab63):

CDR-H1 comprising the amino acid sequence GSIRSSSYWG (SEQ ID NO: 292);

CDR-H2 comprising the amino acid sequence SISYSATTYY (SEQ ID NO: 293);

CDR-H3 comprising the amino acid sequence AGDPSYDSIAGMQV (SEQ ID NO: 294);

CDR-L1 comprising the amino acid sequence RASQSISSYLN (SEQ ID NO: 295);

CDR-L2 comprising the amino acid sequence AASNLQS (SEQ ID NO: 296); and

CDR-L3 comprising the amino acid sequence QQSFDWPLT (SEQ ID NO: 297).

[044] The antibody or antigen-binding fragment may further comprise a heavy chain variable region having an amino acid sequence that is at least 95% identical (optionally at least 98% identical) to SEQ ID NO: 360 and a light chain variable region having an amino acid sequence that is at least 95% identical (optionally at least 98% identical) to SEQ ID NO: 361.

[045] In one aspect, the invention provides an antibody, or antigen-binding fragment thereof, for use in a method for treating a fibrotic disorder in a subject, wherein the antibody, or antigen-binding fragment thereof, specifically binds a human LTBP1-proTGF β complex and/or a human LTBP3-proTGF β complex, and does not bind a human GARP-proTGF β 1 complex;, wherein: a) the fibrotic disorder comprises chronic inflammation; b) the subject benefits from immune suppression; c) the subject has or is at risk of developing an autoimmune disease; d) the subject is a candidate for or has received an allograft transplant; e) the subject has an elevated Th17/Treg ratio; and/or, f) the subject is in need of a long-term or chronic administration of the TGF β 1 inhibitor. In some embodiments, the the subject has or is at risk of developing a metabolic disorder (and the subject is optionally a subject according to one or more of a-f)). In some embodiments, the the antibody, or antigen-binding fragment thereof, is an isoform-specific LTBP1-proTGF β 1 inhibitor and/or LTBP3-proTGF β 1 inhibitor.

[046] The antibodies or antigen-binding fragments thereof provided herein may be used in a method for treating a fibrotic disorder in a subject. The fibrotic disorder may comprise chronic inflammation. The subject may benefit from immune suppression. The subject may have or be at risk of developing an autoimmune disease. The subject may be a candidate for or may have received an allograft transplant.

[047] Alternatively, or in addition, the subject may have an elevated Th17/Treg ratio. The subject may be in need of a long-term or chronic administration of the TGF β 1 inhibitor.

[048] Alternatively, or in addition, the subject may have or be at risk of developing a metabolic disorder.

[049] In another aspect, the invention provides a method for making a composition comprising an antibody, or antigen-binding fragment thereof, that specifically binds a human LTBP1-proTGF β complex and/or a human LTBP3-proTGF β complex, and does not bind a human GARP-proTGF β 1 complex; wherein the antibody, or antigen-binding fragment thereof, inhibits TGF β 1 but does not inhibit TGF β 2 or TGF β 3, the method comprising steps of i) providing at least one antigen comprising LTBP1-proTGF β 1 and/or LTBP3-proTGF β 1, ii) selecting a first pool of antibodies, or antigen-binding fragments thereof, that specifically bind the at least one antigen of step (i) so as to provide specific binders of LTBP1-proTGF β 1 and/or LTBP3-proTGF β 1; iii) selecting a second pool of antibodies, or antigen-binding fragments thereof, that inhibit activation of TGF β 1, so as to generate specific inhibitors of TGF β 1 activation; iv) formulating an antibody, or antigen-binding fragment

thereof, that is present in the first pool of antibodies and the second pool of antibodies into a pharmaceutical composition, thereby making the composition comprising the antibody, or antigen-binding fragment thereof.

[050] In one embodiment, the method further comprises a step of removing from the first pool of antibodies, or antigen-binding fragments thereof, any antibodies, or antigen-binding fragments thereof, that bind GARP-proTGF β 1, LRRC33-proTGF β 1, mature TGF β 1, GARP-proTGF β 2, LRRC33-proTGF β 2, mature TGF β 2, GARP-proTGF β 3, LRRC33-proTGF β 3, mature TGF β 3, or any combinations thereof. In one embodiment, the method further comprises a step of determining or confirming isoform-specificity of the antibodies, or antigen-binding fragments thereof, selected in steps (ii) and/or (iii). In one embodiment, the method further comprises a step of selecting for antibodies, or antigen-binding fragments thereof, that are cross-reactive to human and rodent antigens. In one embodiment, the method further comprises a step of generating a fully human or humanized antibody, or antigen-binding fragment thereof, of the antibody, or antigen-binding fragment thereof, that is present in the first pool of antibodies and the second pool of antibodies.

[051] In one embodiment, the method further comprises a step of subjecting the antibody, or antigen-binding fragment thereof, that is present in the first pool of antibodies and the second pool of antibodies to affinity maturation and/or optimization, so as to provide an affinity matured and/or optimized antibody or fragment thereof. In one embodiment, the affinity maturation/optimization comprises a step of subjecting the antibody, or antigen-binding fragment thereof, that is present in the first pool of antibodies and/or the second pool of antibodies to light chain shuffling as described herein. In one embodiment, the affinity maturation/optimization comprises the step of subjecting the antibody, or antigen-binding fragment thereof, that is present in the first, second, and/or third pool of antibodies to CDR H1/H2 diversification as described herein. In one embodiment, the affinity maturation/optimization comprises the step of subjecting the antibody, or antigen-binding fragment thereof, to CDR-H3 mutagenesis as described herein. In one embodiment, the affinity maturation/optimization comprises the step of subjecting the antibodies, or antigen-binding fragment thereof, to light chain CDR mutagenesis as described herein. In one embodiment, the affinity maturation/optimization comprises the step of subjecting the antibodies, or antigen-binding fragment thereof, to light chain CDR L1/L2 diversification as described herein.

[052] In one embodiment, the method further comprises a step of determining affinity of the antibodies, or antigen-binding fragments thereof, from the first and/or second pools of antibodies to human LTBP1-proTGF β 1 and/or human LTBP3-proTGF β 1. In some embodiments, the method further comprises a step of removing from the first and/or second pools of antibodies, or antigen-binding fragments thereof, any antibodies, or antigen-binding fragments thereof, that bind to human LTBP1-proTGF β 1 and/or human LTBP3-proTGF β 1 with a K_D of > 100 nM, > 50 nM, > 25 nM, or > 10 nM, as measured in a suitable in vitro binding assay such as Bio-Layer Interferometry (BLI).

[053] In one embodiment, the method further comprises a step of determining affinity of the antibodies, or antigen-binding fragments thereof, from the first and/or second pools to mouse LTBP1-proTGF β 1 and/or mouse LTBP3-proTGF β 1. In some embodiments, the method further comprises a step of removing from the first and/or second pools of antibodies, or antigen-binding fragments thereof, any antibodies, or antigen-binding fragments thereof, that bind to mouse LTBP1-proTGF β 1 and/or mouse LTBP3-proTGF β 1 with a K_D of > 100 nM, > 50 nM, or > 10 nM, as measured in a suitable in vitro binding assay such as Bio-Layer Interferometry (BLI).

[054] In one embodiment, the method further comprises a step of removing from the first and/or second pools of antibodies, or antigen-binding fragments thereof, any antibodies, or antigen-binding fragments thereof, that do not bind mouse LTBP1-proTGF β 1 and/or mouse LTBP3-proTGF β 1.

[055] In one embodiment, the method further comprises a step of determining the IC_{50} of the antibodies, or antigen-binding fragments thereof, from of the first and/or second pools of antibodies, or antigen-binding fragments thereof, as measured by a suitable functional in vitro cell-based assay such as a *caga* assay, as described herein. In some embodiments, the method comprises the step of removing antibodies, or antigen-binding fragments thereof, from the first and/or second pools of antibodies, or antigen-binding fragments thereof, that have an IC_{50} of greater than 100 nM, 50 nM, 25 nM, 10 nM, or 5 nM as measured by a cell-based assay (such as a *caga* assay) as described herein.

[056] In some embodiments, the method comprises the step of removing antibodies, or antigen-binding fragments thereof, from the first and/or second pools, antibodies, or antigen-binding fragments thereof, that have an IC_{50} of greater than 50 nM or 10 nM as measured by an endogenous LTBP *caga* assay as described herein.

[057] In some embodiments, the method comprises the step of removing antibodies, or antigen-binding fragments thereof, from the first and/or second pools, antibodies, or antigen-binding fragments thereof, that have an IC_{50} of greater than 50 nM, 25 nM, or 10 nM, as measured by a human LTBP overexpression *caga* assay as described herein.

[058] In some embodiments, the method comprises the step of removing antibodies, or antigen-binding fragments thereof, from the first and/or second pools, antibodies, or antigen-binding fragments thereof, that have an IC_{50} of greater than 50 nM, 25 nM, 10 nM, or 5 nM, as measured by a murine LTBP overexpression *caga* assay as described herein.

[059] Processes and methods for identifying or selecting TGF β 1-selective inhibitors suitable for therapeutic use are encompassed by the invention, as are methods for making a composition comprising a TGF β 1-selective inhibitor. In preferred embodiments, a TGF β 1 inhibitor (*e.g.*, a selected inhibitor) includes one or more antibodies or antigen-binding fragments with particularly advantageous kinetics criteria characterized by: i) high affinities to each of human LTBP1/3-proTGF β 1 complexes (*e.g.*, $K_D < 5$ nM), and, ii) low dissociation rates (k_{OFF}), *e.g.*, $\leq 5 \times 10^{-4}$ (1/s), as

measured by a suitable in vitro binding/kinetics assay, such as by BLI, *e.g.*, Octet-based systems. The low dissociation rate criterion may be reflected in long dissociation half-time ($t_{1/2}$), *e.g.*, ≥ 45 minutes from human LTBP1-proTGF β 1 and/or human LTBP3-proTGF β 1 complexes. Preferably, the long dissociation half-time of an antibody or antigen-binding fragment thereof for the matrix-associated complex(es) is coupled with short dissociation half-time with respect to cell-associated complexes, *e.g.*, human GARP-proTGF β 1 and/or human LRRC33-proTGF β 1 complexes. In particular, preferred antibodies or fragments dissociate from human GARP-proTGF β 1 complex with $t_{1/2}$ of no more than 10 minutes, more preferably no more than 5 minutes. Likewise, methods for making a composition comprising a TGF β 1-selective inhibitor as described herein may further include a step of selecting such antibodies. The selected antibody or the plurality of antibodies are evaluated in preclinical studies comprising an efficacy study and a toxicology/safety study, employing suitable preclinical models. Effective amounts of the antibody or the antibodies determined in the efficacy study are below the level that results in undesirable toxicities determined in the toxicology/safety study. Preferably, the antibody or antibodies are selected which has/have at least 3-fold, 6-fold, and more preferably 10-fold therapeutic window. Effective amounts of the antibodies according to the present disclosure may be between about 0.1 mg/kg and about 30 mg/kg when administered weekly. In preferred embodiments, the maximally tolerated dose (MTD) of the antibodies according to the present disclosure is >100 mg/kg when dosed weekly for at least 4 weeks. In some embodiments, in a preclinical toxicology study, the antibodies show a NOAEL of >100 mg/kg/week, >200 mg/kg/week or >300 mg/kg/week, wherein optionally the toxicology study is a 4-week study, 8-week study, or a 12-week study. For example, the NOAEL is >100 mg/kg/week in a 12-week sub-chronic dosing regimen in healthy mice or rats.

[060] The present disclosure also includes a surprising finding that inhibition of TGF β 3 with a TGF β 3-selective inhibitor produced *pro-fibrotic* effects in mice. Similarly, concurrent inhibition of both TGF β 1 and TGF β 3 in the same model with a combination of a TGF β 1-selective inhibitor and a TGF β 3-selective inhibitor resulted in attenuated anti-fibrotic effects of the TGF β 1 inhibitor. These observations raise the possibility that non-selective TGF β inhibitors (such as pan-inhibitors and TGF β 1/3 inhibitors) may in fact exacerbate fibrosis. Advantageously, the antibodies disclosed herein (*e.g.*, Ab42 and variants thereof, as described herein) are isoform-selective in that they specifically target the latent TGF β 1 complex and do so with low dissociation rates. Thus, the invention includes the recognition that when selecting a particular TGF β inhibitor for patients with a fibrotic condition (*e.g.*, disease involving ECM dysregulation), isoform selectivity should be carefully considered so as to avoid risk of exacerbating ECM dysregulation. Accordingly, the present disclosure includes therapeutic methods comprising selecting a TGF β inhibitor that does not inhibit TGF β 3 to treat a subject with a fibrotic condition, (including preferred fibrotic conditions, as described herein).

[061] The isoform-selective LTBP1/3- proTGF β 1 complex-selective inhibitor as used herein may in some embodiments be selected from Ab31, Ab34, Ab37, Ab38, Ab39, Ab40, Ab41, Ab42, Ab43, Ab44, Ab45, Ab62, Ab63, and Ab64 (optionally Ab42 or Ab63) (*i.e.*, an antibody or antigen-binding fragment having the heavy and light chain variable regions of the corresponding Ab, as provided herein), a variant/derivative or antigen-binding fragment thereof, or an engineered molecule comprising an antigen-binding fragment thereof. In some preferred embodiments, the LTBP1/3- proTGF β 1 complex-selective inhibitor is Ab42, a variant/derivative or antigen-binding fragment thereof, or an engineered molecule comprising an antigen-binding fragment thereof. In preferred embodiments, the LTBP1/3- proTGF β 1 complex-selective inhibitor is Ab42 or an antigen-binding fragment thereof.

BRIEF DESCRIPTION OF THE FIGURES

[062] **Fig. 1** graphically depicts that targeting of the latent form of TGF β 1 provides isoform and context specificity.

[063] **Figs. 2A-2B** demonstrate the identification of isoform-specific and LTBP complex-specific binders of latent TGF β 1. **Fig. 2A** demonstrates that SR-AB1 binds latent TGF β 1, independent of the presenting molecule. SR-AB1 is a human monoclonal antibody that was discovered by yeast display, which selectively binds latent TGF β 1, without detectable binding to latent TGF β 2, TGF β 3, or mature TGF β 1. SR-AB1 cross-reacts with mouse, rat, and cynomolgus monkey proteins and binds to all four latent TGF β 1 complexes. **Fig. 2B** demonstrates that SR-AB2, an anti-LTBP1-proTGF β 1 antibody, does not bind GARP-proTGF β 1 or mature TGF β 1. SR-AB2 cross-reacts with rodent LTBP1-proTGF β 1.

[064] **Figs. 3A-3B** demonstrate functional assays (potency assays) to detect the inhibition of activated recombinant latent TGF β 1. **Fig. 3A** depicts the activation of latent TGF β 1 deposited in the extracellular matrix (ECM). In this assay, presenting molecules are co-transfected with proTGF β 1 in integrin-expressing cells. Transiently transfected cells are seeded in assay plates in the presence of inhibitors. Latent LTBP-proTGF β 1 complex is embedded in the ECM. TGF β reporter cells are then added to the system; free growth factor (released by integrin) signals and is detected by luciferase assay. **Fig. 3B** depicts the activation of latent TGF β 1 presented on the cell surface. Presenting molecules are co-transfected with proTGF β 1 in integrin-expressing cells. Latent TGF β 1 is expressed on the cell surface by GARP or LRRC33. TGF β reporter cells and inhibitors are then added to the system; free growth factor (released by integrin) signals and is detected by luciferase assay.

[065] **Figs. 4A-4B** depict the optimization of the recombinant functional assays. **Fig. 4A** depicts the relative contribution of presenting molecule and/or proTGF β 1 activation upon co-transfection of presenting molecule and proTGF β 1. **Fig. 4B** depicts the optimization of co-transfection: the ratio of

plasmid DNAs for presenting molecule and proTGFβ1. Equivalent amounts of each plasmid were optimal for co-transfection.

[066] **Fig. 5** demonstrates that fibronectin promotes integrin activation of LTBP-presented latent TGFβ1. Assay plates were pre-coated with fibronectin purified from human plasma. Fibronectin increases integrin-mediated activation of latent TGFβ1 presented by LTBP1 and/or LTBP3.

[067] **Fig. 6** is a graph demonstrating that SR-AB1 is a context-independent inhibitor of TGFβ1 activation. SR-AB1 was shown to inhibit integrin-dependent activation of TGFβ1 independent of the presenting molecule.

[068] **Figs. 7A, 7B, and 7C** present data confirming LTBP-selective inhibition of TGFβ1 large latent complex (LLC). **Fig. 7A** demonstrates that SR-AB2 specifically binds LTBP-proTGFβ1 complex; it does not bind proTGFβ1 or LTBP1 alone. SR-AB2 also does not bind GARP-proTGFβ1. **Fig. 7B** depicts that SR-AB2 inhibits integrin activation of LTBP1-proTGFβ1 (human and mouse complexes). **Fig. 7C** depicts that SR-AB2 inhibits integrin activation of LTBP3-proTGFβ1.

[069] **Fig. 8** presents the heavy chain and light chain variable region sequences of SR-AB2 (SEQ ID NOs: 7-8, respectively, in order of appearance). Complementary determining regions (CDRs) are underlined.

[070] **Fig. 9** is a graph demonstrating the binding specificity of SR-AB2 to LTBP1-proTGFβ1 and LTBP3-proTGFβ1 complexes.

[071] **Figs. 10A-10B** provide data showing context-selective inhibition of matrix-associated TGFβ1 activation by SR-AB2. **Fig. 10A** demonstrates that SR-AB2 inhibits LTBP-proTGFβ, wherein the transfected proTGFβ1 is presented by endogenous LTBP1/3. **Fig. 10B** demonstrates that SR-AB2 does not inhibit GARP-presented TGFβ1 activation. These assays were performed in LN229 cells, which express high LTBP3 mRNA, low LTBP1 mRNA, undetectable GARP, and undetectable LRRC33. TGFβ activity, normalized to vehicle, is shown on the y-axis.

[072] **Fig. 11** presents binding profiles and affinity data for LTBP complex-specific antibodies SR-AB10, SR-AB2, and SR-13.

[073] **Figs. 12A and 12B** are graphs showing improved potency of optimized LTBP complex-specific antibodies. **Fig. 12A** provides a graph showing improved inhibitory potency of SR-AB14 (an optimized SR-AB10) as measured by cell-based TGFβ reporter assays. **Fig. 12B** provides a graph showing improved inhibitory potency of SR-AB15 (an optimized SR-AB13) as measured by cell-based TGFβ assays.

[074] **Figs. 13A and 13B** are graphs showing improved potency of optimized LTBP complex-specific antibodies after CDR-H3 mutagenesis (*i.e.*, SR-AB20, SR-AB21, SR-AB22, and SR-AB23), as measured by cell-based TGFβ reporter assays.

[075] **Figs. 14A and 14B** are graphs showing improved potency of optimized LTBP complex-specific antibodies after CDR-H3 mutagenesis (*i.e.*, SR-AB24, SR-AB25, SR-AB26, SR-AB27, SR-AB28, and SR-AB29), as measured by cell-based TGF β reporter assays.

[076] **Fig. 15** is a graph that shows affinity matured antibodies show specific binding to the LTBP-proTGF β 1 complex.

[077] **Fig. 16** is a graph showing improved potency of optimized LTBP complex-specific antibodies after cycles 1, 2 and 3 of antibody optimization as measured by cell-based TGF β reporter assays.

[078] **Fig. 17** depicts the results of an enzyme-linked immunosorbent assay (ELISA) showing antibody binding to baculovirus (BV) particles, which tests antibody polyspecificity.

[079] **Fig. 18** depicts the results of affinity-capture Self-interaction Nanoparticle Spectroscopy (AC-SINS) assay, which tests antibody self-interaction. Increased plasmon wavelength indicates self-interaction.

[080] **Fig. 19** is a graph showing treatment with SR-AB42 and SR-AB31 inhibited the increase in hydroxyproline (HYP) (μ g/mg tissue) in liver tissue in animals on a choline-deficient high fat diet (CDHFD).

[081] **Fig. 20A** is a graph showing relative ratios of phosphorylated versus total (phosphorylated and unphosphorylated) Smad2/3 (pSMAD2/3:tSMAD2/3) in an Alport mouse model. A single dose of SR-AB42 or SR-AB63 was sufficient to significantly inhibit pSmad2/3 signaling in whole kidney lysates. **Fig. 20B** is a graph showing the amount of phosphorylated SMAD2/3 (pSMAD2/3) as determined by ELISA, and **Fig. 20C** is a graph showing the amount of total SMAD2/3 (tSMAD2/3) protein as determined by ELISA. As shown by **Fig. 20B** and **Fig. 20C**, reduction of pSMAD is contributing to the change in ratio shown in **Fig. 20A**.

[082] **Fig. 21** is a graph showing that the lead cycle 3 antibodies show no inhibition in the LTBP-TGF β 3 assay.

[083] **Fig. 22** provides 5 representative PSR-stained images from controls, CDHFD mice treated with Reference Ab, a TGF β 3 inhibitor, or both (left). A graph showing picosirius red area (%) in liver sections of CDHFD mice treated with Reference Ab, a TGF β 3 inhibitor, or both, as compared to control, is also provided (right).

[084] **Fig. 23** shows LTBP-complex antibodies such as SR-AB63 are highly specific and have picomolar monovalent affinities.

[085] **Fig. 24** shows LTBP-complex antibodies such as SR-AB42 are highly specific and have picomolar monovalent affinities.

DETAILED DESCRIPTION OF CERTAIN EMBODIMENTS

[086] The present invention provides compositions that are useful for reducing activation of TGF β . Inhibitors that target latent proTGF β complexes, upstream of growth factor-receptor interaction, are generally referred to as activation inhibitors of TGF β .

[087] To date, four presenting molecules for TGF β have been identified: latent TGF beta-binding protein 1 (“LTBP1”), latent TGF beta-binding protein 3 (“LTBP3”), glycoprotein A repetitions predominant (“GARP”) and leucine-rich repeat-containing protein 33 (“LRRC33”). Each of these presenting molecules can form disulfide bonds with a homodimeric pro-protein complex of the TGF β precursor, *i.e.*, proTGF β . The proTGF β complex remains dormant (latent) in the respective extracellular niche (*e.g.*, ECM and immune cell surface) until activation events trigger the release of soluble growth factor from the complex.

[088] As compared to the TGF β growth factors and the receptors, which are expressed broadly, the presenting molecules show more restricted or selective (*e.g.*, tissue-specific) expression patterns, giving rise to functional compartmentalization of TGF β activities by virtue of association. The four presenting molecule-proTGF β complexes, namely, LTBP1-proTGF β , LTBP3-proTGF β , GARP-proTGF β and LRRC33-proTGF β , therefore, provide discrete “contexts” of TGF β signaling within the tissue in which the presenting molecules are expressed. These contexts may be divided into two broad categories: i) TGF β signaling associated with the ECM (*e.g.*, matrix-associated TGF β function); and ii) TGF β signaling associated with cells (particularly certain immune cell function). The LTBP1-proTGF β and LTBP3-proTGF β complexes fall under the first category, while GARP-proTGF β and LRRC33-proTGF β complexes fall under the second category. Thus, disclosed herein are inhibitors of TGF β that are capable of selectively inhibiting the activation of TGF β that is associated with the ECM. In some embodiments, the inhibitors are also selective for a particular TGF β isoform (*e.g.*, proTGF β 1, proTGF β 2, and/or proTGF β 3).

[089] In exemplary embodiments, the compositions described herein are useful for selectively reducing activation of TGF β 1 in the context of an LTBP protein, *e.g.*, a LTBP1 and/or a LTBP3 protein. Such compositions advantageously inhibit activation of extracellular matrix-associated TGF β 1, without inhibiting TGF β 1 in the context of the immune-associated TGF β 1 presenting molecules GARP and LRRC33. The compositions described herein are useful for treating disorders associated with TGF β 1 activation, *e.g.*, fibrotic disorders. Accordingly, in embodiments, the invention provides compositions for reducing activation of TGF β 1, methods of use thereof, methods of manufacture, and treatment methods. Methods of selecting a TGF β 1 inhibitor for subjects exhibiting symptoms of a fibrotic disorder are also provided.

Definitions

[090] In order that the disclosure may be more readily understood, certain terms are first defined. These definitions should be read in light of the remainder of the disclosure and as understood by a person of ordinary skill in the art. Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood by a person of ordinary skill in the art. Additional definitions are set forth throughout the detailed description.

[091] *Affinity*: Affinity is the strength of binding of a molecule (such as an antibody) to its ligand (such as an antigen). It is typically measured and reported by the equilibrium dissociation constant (K_D). K_D is the ratio of the antibody dissociation rate (“off rate” or K_{off}), how quickly it dissociates from its antigen, to the antibody association rate (“on rate” or K_{on}) of the antibody, how quickly it binds to its antigen. For example, an antibody with an affinity of $\leq 1 \mu\text{M}$ has a K_D value that is $1 \mu\text{M}$ or lower (*i.e.*, $1 \mu\text{M}$ or higher affinity) determined by a suitable in vitro binding assay. Suitable in vitro assays, such as Biolayer Interferometry (*e.g.*, Octet) or surface plasmon resonance (*e.g.*, Biacore System) can be used to assess affinities, as measured by K_D values based on well-known methods.

[092] *Affinity maturation*: Affinity maturation is a type of antibody optimization and is a process of improving the affinity of an antibody or a fragment to its antigen and typically involves making one or more changes to the amino acid sequence of the antibody or the fragment to achieve greater affinity. Typically, a parental antibody and an affinity-matured counterpart retain the same epitope. Affinity maturation may include diversification and/or mutagenesis of one or more CDR sequences.

[093] *Antibody*: The term “antibody” encompasses any naturally-occurring, recombinant, modified or engineered immunoglobulin or immunoglobulin-like structure or antigen-binding fragment or portion thereof, or derivative thereof, as further described elsewhere herein. Thus, the term refers to an immunoglobulin molecule that specifically binds to a target antigen, and includes, for instance, chimeric, humanized, fully human, and bispecific antibodies. Unless otherwise specified to the contrary, the term “antibody” as used herein shall encompass antigen-binding fragments and variants thereof. An intact antibody will generally comprise at least two full-length heavy chains and two full-length light chains, but in some instances can include fewer chains such as antibodies naturally occurring in camelids which can comprise only heavy chains. Antibodies can be derived solely from a single source, or can be “chimeric,” that is, different portions of the antibody can be derived from two different antibodies. Antibodies, or antigen-binding portions thereof, can be produced in hybridomas, by recombinant DNA techniques, or by enzymatic or chemical cleavage of intact antibodies. The term antibodies, as used herein, includes monoclonal antibodies, bispecific antibodies, minibodies, domain antibodies, synthetic antibodies (sometimes referred to herein as “antibody mimetics”), chimeric antibodies, humanized antibodies, human antibodies, antibody fusions

(sometimes referred to herein as “antibody conjugates”), respectively. In some embodiments, the term also encompasses peptibodies.

[094] *Antigen*: The term “antigen” broadly includes any molecules comprising an antigenic determinant within a binding region(s) to which an antibody or a fragment specifically binds. An antigen can be a single-unit molecule (such as a protein monomer or a fragment) or a complex comprised of multiple components. An antigen provides an epitope, *e.g.*, a molecule or a portion of a molecule, or a complex of molecules or portions of molecules, capable of being bound by a selective binding agent, such as an antigen-binding protein (including, *e.g.*, an antibody). Thus, a selective binding agent may specifically bind to an antigen that is formed by two or more components in a complex. In some embodiments, the antigen is capable of being used in an animal to produce antibodies capable of binding to that antigen. An antigen can possess one or more epitopes that are capable of interacting with different antigen-binding proteins, *e.g.*, antibodies. In the context of the present disclosure, a suitable antigen is a complex (*e.g.*, multimeric complex comprised of multiple components in association) containing a proTGF dimer (“small latent complex” or SLC) preferably in association with a presenting molecule (together “large latent complex” or LLC). Each monomer of the proTGF dimer comprises a prodomain and a growth factor domain, separated by a furin cleavage sequence. Two such monomers form the proTGF dimer complex. This in turn is covalently associated with a presenting molecule via disulfide bonds, which involve a cysteine residue present near the N-terminus of each of the proTGF monomer. This multi-complex formed by a proTGF dimer bound to a presenting molecule is generally referred to as a large latent complex. An antigen complex suitable for screening antibodies or antigen-binding fragments, for example, includes a presenting molecule component of a large latent complex. Such presenting molecule component may be a full-length presenting molecule or a fragment(s) thereof. Minimum required portions of the presenting molecule typically contain at least 50 amino acids, but more preferably at least 100 amino acids of the presenting molecule polypeptide, which comprises two cysteine residues capable of forming covalent bonds with the proTGFβ1 dimer.

[095] *Antigen-binding portion/fragment*: The terms “antigen-binding portion” or “antigen-binding fragment” of an antibody, as used herein, refers to one or more fragments of an antibody that retain the ability to specifically bind to an antigen (*e.g.*, LTBP1-proTGFβ1 and LTBP3-proTGFβ1). Antigen-binding portions include, but are not limited to, any naturally occurring, enzymatically obtainable, synthetic, or genetically engineered polypeptide or glycoprotein that specifically binds an antigen to form a complex. In some embodiments, an antigen-binding portion of an antibody may be derived, *e.g.*, from full antibody molecules using any suitable standard techniques such as proteolytic digestion or recombinant genetic engineering techniques involving the manipulation and expression of DNA encoding antibody variable and optionally constant domains. Non-limiting examples of antigen-binding portions include: (i) Fab fragments, a monovalent fragment consisting of the VL, VH,

CL and CH1 domains; (ii) F(ab')₂ fragments, a bivalent fragment comprising two Fab fragments linked by a disulfide bridge at the hinge region; (iii) Fd fragments consisting of the VH and CH1 domains; (iv) Fv fragments consisting of the VL and VH domains of a single arm of an antibody; (v) single-chain Fv (scFv) molecules (see, *e.g.*, Bird et al. (1988) SCIENCE 242:423-426; and Huston et al. (1988) PROC. NAT'L. ACAD. SCI. USA 85:5879-5883); (vi) dAb fragments (see, *e.g.*, Ward et al. (1989) NATURE 341: 544-546); and (vii) minimal recognition units consisting of the amino acid residues that mimic the hypervariable region of an antibody (*e.g.*, an isolated complementarity determining region (CDR)). Other forms of single chain antibodies, such as diabodies are also encompassed. The term antigen-binding portion of an antibody includes a "single chain Fab fragment" otherwise known as an "scFab," comprising an antibody heavy chain variable domain (VH), an antibody constant domain 1 (CH1), an antibody light chain variable domain (VL), an antibody light chain constant domain (CL) and a linker, wherein said antibody domains and said linker have one of the following orders in N-terminal to C-terminal direction: a) VH-CH1-linker-VL-CL, b) VL-CL-linker-VH-CH1, c) VH-CL-linker-VL-CH1 or d) VL-CH1-linker-VH-CL; and wherein said linker is a polypeptide of at least 30 amino acids, preferably between 32 and 50 amino acids.

[096] *Advanced fibrosis*: As used herein, a subject suffers from advanced fibrosis if s/he has an advanced stage of a fibrotic disorder, particularly organ fibrosis, which renders the patient a candidate for receiving, or in need of, an allograft transplant.

[097] *As needed*: In the context of dosing regimens, the term "as needed" refers to a dosing regimen that is not based on a predetermined dosing schedule but instead based on one or more parameters or markers measured or monitored periodically during treatment, which provides information or guidance as to whether additional doses should be beneficial to the subject/patient. For instance, a pharmaceutical composition comprising a TGFβ inhibitor such as TGFβ_{1/2/3} inhibitors ("pan" inhibitors), TGFβ_{1/2} inhibitors and TGFβ_{1/3} inhibitors, may be administered, intermittently, on an "as needed" basis in a therapeutically effective amount sufficient to achieve and/or maintain clinical benefit (*e.g.*, reduction of one or more clinical markers of fibrosis). In some embodiments, administration of a LTBP1/3-complex selective TGFβ inhibitor such as any one of the antibodies disclosed herein (*e.g.*, Ab31, Ab34, Ab37, Ab38, Ab39, Ab40, Ab41, Ab42, Ab43, Ab44, Ab45, Ab62, Ab63, or Ab64 (optionally Ab42)) may be used in combination with a method of determining or monitoring therapeutic efficacy. In some embodiments, the LTBP1/3-complex selective TGFβ inhibitor is administered in patients only when clinical benefit from additional doses of the TGFβ inhibitor is expected. It is contemplated that, in order to manage toxicities, intermittent or "as-needed" dosing regimen may be required more frequently with isoform-non-selective inhibitors of TGFβ, as compared to TGFβ₁-selective inhibitors, such as those disclosed herein.

[098] *Bias*: In the context of the present disclosure, the term “bias” refers to skewed or uneven affinity towards or against a subset of antigens to which an antibody is capable of specifically binding. For example, an antibody is said to have bias when the affinity for one antigen complex and the affinity for another antigen complex are not equivalent (*e.g.*, more than five-fold difference in affinity). Antibodies characterized as “unbiased” have approximately equivalent affinities towards such antigen complexes (*e.g.*, less than five-fold difference in affinity). Antibodies of the present disclosure “selectively” bind EMC-associated complexes (LTBP1-proTGFβ1 and LTBP3-proTGFβ). Such selective binding may in some embodiments comprise binding such that relative affinities between at least one of the matrix-associated complexes and at least one (preferably both) of the cell-associated complexes (GARP-proTGFβ1 and/or LRRC33-proTGFβ1 complexes) is greater than fifty-fold.

[099] *Biolayer Interferometry (BLI)*: BLI is a label-free technology for optically measuring biomolecular interactions, *e.g.*, between a ligand immobilized on the biosensor tip surface and an analyte in solution. BLI provides the ability to monitor binding specificity, rates of association and dissociation, or concentration, with precision and accuracy. BLI platform instruments are commercially available, for example, from ForteBio and are commonly referred to as the Octet® System. BLI can be employed in carrying out in vitro binding assays as described herein.

[100] *Autoimmune disease*: An autoimmune disease is a condition arising from an abnormal or overactive immune response to a normal body part. Immunostimulating agents administered to such patients with autoimmune conditions may exacerbate the condition.

[101] *Cell-associated proTGFβ1*: The term refers to TGFβ1 or its signaling complex (*e.g.*, pro/latent TGFβ1) that is membrane-bound (*e.g.*, tethered to cell surface). Typically, such cell is an immune cell. TGFβ1 that is presented by GARP or LRRC33 is a cell-associated TGFβ1. GARP and LRRC33 are transmembrane presenting molecules that are expressed on cell surface of certain cells. GARP-proTGFβ1 and LRRC33-proTGFβ1 may be collectively referred to as “cell-associated” (or “cell-surface”) proTGFβ1 complexes, that mediate cell-associated (*e.g.*, immune cell-associated) TGFβ1 activation/signaling.

[102] *Chronic inflammation*: In the context of the present disclosure, fibrotic disorders that involve chronic inflammation are characterized by continuous or persistent injury to a tissue such that it does not resolve in normal healing after an initial injury. Chronic inflammation refers to a prolonged inflammatory response that involves a progressive change in the type of cells present at the site of inflammation (*e.g.*, fibrotic tissues). It is characterized by the simultaneous destruction and repair of the tissue from the inflammatory process. It can follow an acute form of inflammation or be a prolonged low-grade form.

[103] *Clinical benefit*: As used herein, the term “clinical benefits” is intended to include both efficacy and safety of a therapy. Thus, therapeutic treatment that achieves a desirable clinical benefit is both efficacious and safe (*e.g.*, with tolerable or acceptable toxicities or adverse events).

[104] *Combinatory or combinatorial epitope*: A combinatorial epitope is an epitope that is recognized and bound by a combinatorial antibody at a site (*i.e.*, antigenic determinant) formed by non-contiguous portions of a component or components of an antigen, which, in a three-dimensional structure, come together in close proximity to form the epitope. Thus, antibodies of the invention may bind an epitope formed by two or more components (*e.g.*, portions or segments) of a pro/latent TGF β 1 complex. A combinatory epitope may comprise amino acid residue(s) from a first component of the complex, and amino acid residue(s) from a second component of the complex, and so on. Each component may be of a single protein or of two or more proteins of an antigenic complex. A combinatory epitope is formed with structural contributions from two or more components (*e.g.*, portions or segments, such as amino acid residues) of an antigen or antigen complex.

[105] *Complementary determining region*: As used herein, the term “CDR” refers to the complementarity determining region within antibody variable sequences. There are three CDRs in each of the variable regions of the heavy chain and the light chain, which are designated CDR1, CDR2 and CDR3, for each of the variable regions. The exact boundaries of these CDRs have been defined differently according to different systems. The system described by Kabat (Kabat et al. (1987; 1991) Sequences of Proteins of Immunological Interest (National Institutes of Health, Bethesda, Md.) not only provides an unambiguous residue numbering system applicable to any variable region of an antibody, but also provides precise residue boundaries defining the three CDRs on each of the heavy and light chains. These CDRs may be referred to as Kabat CDRs.

[106] *Conformational epitope*: A conformational epitope is an epitope that is recognized and bound by a conformational antibody in a three-dimensional conformation, but not in an unfolded peptide of the same amino acid sequence. A conformational epitope may be referred to as a conformation-specific epitope, conformation-dependent epitope, or conformation-sensitive epitope. A corresponding antibody or fragment thereof that specifically binds such an epitope may be referred to as conformation-specific antibody, conformation-selective antibody, or conformation-dependent antibody. Binding of an antigen to a conformational epitope depends on the three-dimensional structure (conformation) of the antigen or antigen complex.

[107] *Context-specific*: Context-specific (or context-selective) antibodies of the invention (as opposed to “context-independent” antibodies) are capable of binding selectively to a *subset*, but not all, of proTGF β 1 complexes associated with a particular biological context. For example, matrix-selective targeting enables specific inhibition of TGF β 1 function associated with the ECM. ECM-selective inhibition can be achieved by the use of antibodies or fragments thereof that selectively

target the ECM components, LTBP1-proTGFβ1 and/or LTBP3-proTGFβ1. Antibodies and fragments disclosed herein therefore represent a class of context-specific antibodies. LTBP1-specific and LTBP3-specific inhibitors of TGFβ1 activation are also context-specific antibodies.

[108] *Cross-block/cross-blocking*: a first antibody or antigen-binding portion thereof and a second antibody or antigen-binding portion thereof cross-block with each other with respect to the same antigen, for example, as assayed by as measured by Biolayer Interferometry (such as Octet) or surface plasmon resonance (such as Biacore System), using standard test conditions, *e.g.*, according to the manufacturer's instructions (*e.g.*, binding assayed at room temperature, ~20-25°C). The first antibody or fragment thereof and the second antibody or fragment thereof may have the same epitope; may have non-identical but overlapping epitopes; or, may have separate (different) epitopes which are in close proximity in a three-dimensional space, such that antibody binding is cross-blocked via steric hindrance. "Cross-block" means that binding of the first antibody to an antigen prevents binding of the second antibody to the same antigen, and similarly, binding of the second antibody to an antigen prevents binding of the first antibody to the same antigen.

[109] *Dissociation rate*: The term dissociation rate as used herein has the meaning understood by the skilled artisan in the pertinent art (*e.g.*, antibody technology) as refers to a kinetics parameter measured by how fast/slow a ligand (*e.g.*, antibody or fragment) dissociates from its binding target (*e.g.*, antigen). Dissociation rate is also referred to as the "off" rate (" k_{OFF} "). Relative on/off rates between an antibody and its antigen (*i.e.*, k_{ON} and k_{OFF}) determine the overall strength of the interaction, or affinity, typically expressed as a dissociation constant, or K_D . Therefore, equivalent affinities (*e.g.*, K_D values) may be achieved by having fast association (high k_{ON}), slow dissociation (low k_{OFF}), or contribution from both factors. Monovalent interactions may be measured by the use of monovalent antigen-binding molecules/fragments, such as fAb (Fab), whilst divalent interactions may be measured by the use of divalent antigen-binding molecules such as whole immunoglobulins (*e.g.*, IgGs). Dissociation kinetics may be expressed in terms of dissociation half-time (sometimes referred to as half binding time), or $t_{1/2}$, defined as a duration of time it takes for one half the number of antibody molecules (*e.g.*, mAb, Fab, etc.) to dissociate from bound antigen. Thus, antibodies with slow dissociation rates have long dissociation half-time, and antibodies with fast dissociation rates have short dissociation half-time.

[110] *Dosage*: As used herein, typical therapeutic dosage of an antibody of the present invention ranges between about 1-30 mg/kg per dose. A typical dosing regimen may include once a week, every 2 weeks, every 3 weeks, every 4 weeks, once a month, every 6 weeks, etc.

[111] *ECM-associated (or "matrix-associated") TGFβ1*: The term refers to TGFβ1 or its signaling complex (*e.g.*, pro/latent TGFβ1) that is a component of (*e.g.*, deposited into) the extracellular matrix. TGFβ1 that is presented by LTBP1 or LTBP3 is an ECM-associated TGFβ1.

[112] *Effective amount*: An “effective amount” (or therapeutically effective amount) is a dosage or dosing regimen that achieves statistically significant clinical benefits in a patient population.

[113] *Fibrotic disorder*: The term “fibrosis” or “fibrotic condition/disorder” refers to the process or manifestation characterized by the pathological accumulation of extracellular matrix (ECM) components, such as collagens, within a tissue or organ. Fibrosis can include primary fibrosis, as well as secondary fibrosis that are associated with a disease or disorder.

[114] *GARP-proTGFβ1*: As used herein, the term “GARP-proTGFβ1” refers to a protein complex comprising a pro-protein form or latent form of a transforming growth factor-β1 (TGFβ1) protein associated with a glycoprotein-A repetitions predominant protein (GARP) or fragment or variant thereof. The proTGFβ1 homodimer is capable of forming covalent association with a single molecule of GARP via disulfide bonds. The term “GARP-TGFβ1” may be used interchangeably. GARP-proTGFβ1 expression is limited to certain cell types, such as regulatory T cells (Treg).

[115] *Human antibody*: The term “human antibody,” as used herein, is intended to include antibodies having variable and constant regions derived from human germline immunoglobulin sequences. The human antibodies of the present disclosure may include amino acid residues not encoded by human germline immunoglobulin sequences (*e.g.*, mutations introduced by random or site-specific mutagenesis in vitro or by somatic mutation in vivo), for example in the CDRs and in particular CDR3 (*e.g.*, CDR-H3 or CDR-L3 mutagenesis).

[116] *Humanized antibody*: The term “humanized antibody” refers to antibodies, which comprise heavy and light chain variable region sequences from a non-human species (*e.g.*, a mouse) but in which at least a portion of the VH and/or VL sequence has been altered to be more “human-like,” *i.e.*, more similar to human germline variable sequences. One type of humanized antibody is a CDR-grafted antibody.

[117] *Immune suppression/immunosuppression*: The term immunosuppression refers to suppression or reduction of the strength of the body's immune system. Patients who “benefit from immunosuppression” include those who have advanced stages of organ fibrosis and are candidates for, being considered for, or have undergone transplantation.

[118] *Isoform-specific*: The term “isoform specificity” refers to an agent's ability to discriminate one isoform over other structurally related isoforms (*i.e.*, selectivity). An isoform-specific TGFβ inhibitor exerts its inhibitory activity towards one isoform of TGFβ but not the other isoforms of TGFβ at a given concentration. For example, an isoform-specific TGFβ1 antibody selectively binds TGFβ1. A TGFβ1-specific inhibitor (antibody) preferentially targets (binds thereby inhibits) the TGFβ1 isoform over TGFβ2 or TGFβ3 with substantially greater affinity. For example, the selectivity in this context may refer to at least a 500-1000-fold difference in respective affinities as measured by an in vitro binding assay such as Octet and Biacore. In some embodiments, the selectivity

is such that the inhibitor when used at a dosage effective to inhibit TGF β 1 in vivo does not inhibit TGF β 2 and TGF β 3. Context-specific inhibitors of the present disclosure are also isoform-specific.

[119] *Isolated*: An “isolated” antibody as used herein, refers to an antibody that is substantially free of other antibodies having different antigenic specificities. In some embodiments, an isolated antibody is substantially free of other unintended cellular material and/or chemicals.

[120] *Long-term or chronic administration*: As used herein, a therapeutic regimen that involves over six months of treatment is considered long-term. In some patient populations, long-term therapeutic regimens involve administration of a drug (such as context-selective TGF β 1 inhibitors) for an indefinite duration of time.

[121] *LRRC33-proTGF β 1*: As used herein, the term “LRRC33-TGF β 1 complex” refers to a complex between a pro-protein form or latent form of transforming growth factor- β 1 (TGF β 1) protein and a Leucine-Rich Repeat-Containing Protein 33 (LRRC33; also known as Negative Regulator Of Reactive Oxygen Species or NRROS) or fragment or variant thereof. In some embodiments, a LRRC33-TGF β 1 complex comprises LRRC33 covalently linked with pro/latent TGF β 1 via one or more disulfide bonds. In other embodiments, a LRRC33-TGF β 1 complex comprises LRRC33 non-covalently linked with pro/latent TGF β 1. In some embodiments, a LRRC33-TGF β 1 complex is a naturally-occurring complex, for example a LRRC33-TGF β 1 complex in a cell.

[122] *LTBP1-TGF β 1*: As used herein, the term “LTBP1-TGF β 1 complex” (or “LTBP1-proTGF β 1 complex”) refers to a protein complex comprising a pro-protein form or latent form of transforming growth factor- β 1 (TGF β 1) protein (may be referred to as “proTGF β 1” herein) and a latent TGF-beta binding protein 1 (LTBP1) or fragment or variant thereof. In some embodiments, a LTBP1-TGF β 1 complex comprises LTBP1 covalently linked with pro/latent TGF β 1 via one or more disulfide bonds. In other embodiments, a LTBP1-TGF β 1 complex comprises LTBP1 non-covalently linked with pro/latent TGF β 1. In some embodiments, a LTBP1-TGF β 1 complex is a naturally-occurring complex, for example a LTBP1-TGF β 1 complex in a cell. An exemplary LTBP1-TGF β 1 complex is shown in FIG. 3.

[123] *LTBP3-TGF β 1*: As used herein, the term “LTBP3-TGF β 1 complex” (or “LTBP3-proTGF β 1 complex”) refers to a protein complex comprising a pro-protein form or latent form of transforming growth factor- β 1 (TGF β 1) protein (may be referred to as “proTGF β 1” herein) and a latent TGF-beta binding protein 3 (LTBP3) or fragment or variant thereof. In some embodiments, a LTBP3-TGF β 1 complex comprises LTBP3 covalently linked with pro/latent TGF β 1 via one or more disulfide bonds. In other embodiments, a LTBP3-TGF β 1 complex comprises LTBP1 non-covalently linked with pro/latent TGF β 1. In some embodiments, a LTBP3-TGF β 1 complex is a naturally-occurring complex, for example a LTBP3-TGF β 1 complex in a cell. An exemplary LTBP3-TGF β 1 complex is shown in FIG. 3.

[124] *Macrophages*: Macrophages are a type of white blood cells of the immune system and includes heterogeneous, phenotypically diverse subpopulations of myeloid cells. Some macrophages differentiate from bone marrow-derived, circulating monocytes, while others are tissue-specific macrophages that reside within particular anatomical or tissue locations (“resident” macrophages). Tissue-specific macrophages include but are not limited to: Adipose tissue macrophages; Kupffer cells (Liver); Sinus histiocytes (Lymph nodes); Alveolar macrophages (or dust cells, Pulmonary alveoli of lungs); Tissue macrophages (histiocytes) leading to giant cells (Connective tissue); Langerhans cells (Skin and mucosa); Microglia (Central nervous system); Hofbauer cells (Placenta); Intraglomerular mesangial cells (Kidney); Osteoclasts (Bone); Epithelioid cells (Granulomas); Red pulp macrophages (or Sinusoidal lining cells, Red pulp of spleen); Peritoneal macrophages (Peritoneal cavity); and, LysoMac (Peyer's patch). Macrophages, *e.g.*, bone-marrow derived monocytes, can be activated by certain stimuli (such as cytokines) resulting in polarized phenotypes, *e.g.*, M1 and M2. M2-biased activated macrophages are further classified into several phenotypically distinct subtypes, such as M2a, M2b, M2c (*e.g.*, pro-fibrotic) and M2d (pro-tumor or TAM-like).

[125] *Matrix-associated proTGFβ1*: LTBP1 and LTBP3 are presenting molecules that are components of the extracellular matrix (ECM). LTBP1-proTGFβ1 and LTBP3-proTGFβ1 may be collectively referred to as “ECM-associated” (or “matrix-associated”) proTGFβ1 complexes, that mediate ECM-associated TGFβ1 activation/signaling.

[126] *Maximally tolerated dose (MTD)*: The term MTD generally refers to, in the context of safety/toxicology considerations, the highest amount of a test article (such as a TGFβ1 inhibitor) evaluated with no observed adverse effect level (NOAEL). For example, the NOAEL for Ab2 in rats was the highest dose evaluated (100 mg/kg), suggesting that the MTD for Ab2 is >100 mg/kg, based on a four-week toxicology study.

[127] *Myeloid-derived suppressor cell*: Myeloid-derived suppressor cells (MDSCs) are a heterogeneous population of cells generated during various pathologic conditions and thought to represent a pathologic state of activation of monocytes and relatively immature neutrophils. MDSCs include at least two categories of cells termed i) “granulocytic” (G-MDSC) or polymorphonuclear (PMN-MDSC), which are phenotypically and morphologically similar to neutrophils; and ii) monocytic (M-MDSC) which are phenotypically and morphologically similar to monocytes. MDSCs are characterized by a distinct set of genomic and biochemical features, and can be distinguished by specific surface molecules. For example, human G-MDSCs/PMN-MDSCs typically express the cell-surface markers CD11b, CD33, CD15 and CD66. In addition, human G-MDSCs/PMN-MDSCs may also express HLA-DR and/or Arginase. By comparison, human M-MDSCs typically express the cell surface markers CD11b, CD33 and CD14. The MDSCs may also express CD39 and CD73 to mediate adenosine signaling involved in organ fibrosis (such as liver fibrosis, and lung fibrosis), cancer and myelofibrosis). In addition, human M-MDSCs may also express HLA-DR. In addition to such cell-

surface markers, MDSCs are characterized by the ability to suppress immune cells, such as T cells, NK cells and B cells. Immune suppressive functions of MDSCs may include inhibition of antigen-non-specific function and inhibition of antigen-specific function. MDSCs can express cell surface LRRC33 and/or LRRC33-proTGF β 1.

[128] *Myofibroblast*: Myofibroblasts are cells with certain phenotypes of fibroblasts and smooth muscle cells and generally express vimentin, alpha-smooth muscle actin (α -SMA; human gene ACTA2) and paladin. In many disease conditions involving extracellular matrix dysregulations (such as increased matrix stiffness), normal fibroblast cells become de-differentiated into myofibroblasts in a TGF β -dependent manner.

[129] *Off rate (k_{OFF})*: The off rate is a kinetic parameter of how fast or how slowly an antibody (such as mAb) or antigen-binding fragment (such as fAb) dissociates from its antigen and may be also referred to as the dissociation rate. Dissociation rates can be experimentally measured in suitable in vitro binding assays, such as BLI (Octet®)- and/or SPR (Biacore)-based systems. In the context of antibody-antigen binding kinetics, the term "half-binding-time" ($T_{1/2}$) or "dissociation half-time" refers to the duration of time required for half the number of antibody molecules (*e.g.*, mAb, Fab) to dissociate from the bound antigen (*e.g.*, LTBP1-proTGF β 1, LTBP3-proTGF β 1). Thus, an antibody that dissociates slowly (*i.e.*, low off rates) from its antigen has a long $T_{1/2}$. Conversely, an antibody that dissociates rapidly (*i.e.*, high off rates) from its antigen has a short $T_{1/2}$.

[130] *Pan-TGF β inhibitor/pan-inhibition of TGF β* : The term "pan-TGF β inhibitor" refers to any agent that is capable of inhibiting or antagonizing all three isoforms of TGF β . Such an inhibitor may be a small molecule inhibitor of TGF β isoforms. The term includes pan-TGF β antibody which refers to any antibody capable of binding to each of TGF β isoforms, *i.e.*, TGF β 1, TGF β 2, and TGF β 3. In some embodiments, a pan-TGF β antibody binds and neutralizes activities of all three isoforms, *i.e.*, TGF β 1, TGF β 2, and TGF β 3 activities.

[131] *Potency*: The term "potency" as used herein refers to activity of a drug, such as a functional antibody (or fragment) having inhibitory activity, with respect to concentration or amount of the drug to produce a defined effect. For example, an antibody capable of producing certain effects at a given dosage is more potent than another antibody that requires twice the amount (dosage) to produce equivalent effects. Potency may be measured in cell-based assays, such as TGF β activation/inhibition assays. In some cases, the degree of TGF β activation, such as activation triggered by integrin binding, can be measured in the presence or absence of test article (*e.g.*, inhibitory antibodies) in a cell-based system. Typically, antibodies with higher affinities tend to show higher potency than antibodies with lower affinities.

[132] *Presenting molecule*: Presenting molecules are proteins that form covalent bonds with latent pro-proteins (*e.g.*, proTGF β 1) and "present" the inactive complex in an extracellular niche

(such as ECM or immune cell surface) thereby maintaining its latency until an activation event occurs. Known presenting molecules for proTGF β 1 include: LTBP1, LTBP3, GARP and LRRC33, which can form presenting molecule-proTGF β 1 complexes, namely, LTBP1-proTGF β 1, LTBP3-proTGF β 1, GARP-proTGF β 1 and LRRC33-proTGF β 1, respectively. LTBP1 and LTBP3 are components of the extracellular matrix (ECM); therefore, LTBP1-proTGF β 1 and LTBP3-proTGF β 1 may be collectively referred to as “ECM-associated” (or “matrix-associated”) proTGF β 1 complexes, that mediate ECM-associated TGF β 1 signaling/activities. GARP and LRRC33, on the other hand, are transmembrane proteins expressed on cell surface of certain cells; therefore, GARP-proTGF β 1 and LRRC33-proTGF β 1 may be collectively referred to as “cell-associated” (or “cell-surface”) proTGF β 1 complexes, that mediate cell-associated (*e.g.*, immune cell-associated) TGF β 1 signaling/activities.

[133] *ProTGF β 1*: The term “proTGF β 1” as used herein is intended to encompass precursor forms of inactive TGF β 1 complex that comprises a prodomain sequence of TGF β 1 within the complex. Thus, the term can include the pro-, as well as the latent-forms of TGF β 1. The expression “pro/latent TGF β 1” may be used interchangeably. The “pro” form of TGF β 1 exists prior to proteolytic cleavage at the furin site. Once cleaved, the resulting form is said to be the “latent” form of TGF β 1. The “latent” complex remains associated until further activation trigger, such as integrin-driven activation event. The proTGF β 1 complex is comprised of dimeric TGF β 1 pro-protein polypeptides, linked with disulfide bonds. The latent dimer complex is covalently linked to a single presenting molecule via the cysteine residue at position 4 (Cys4) of each of the proTGF β 1 polypeptides. The adjective “latent” may be used generally to describe the “inactive” state of TGF β 1, prior to integrin-mediated or other activation events. The proTGF β 1 polypeptide contains a prodomain (LAP) and a growth factor domain (SEQ ID NO: 12).

[134] *Regulatory T cell (Treg)*: “Regulatory T cells,” or Tregs, are a type of immune cells characterized by the expression of the biomarkers, CD4, forkhead box P3 (FOXP3), and CD25, as well as STAT5. Tregs are sometimes referred to as suppressor T cells and represent a subpopulation of T cells that modulate the immune system, maintain tolerance to self-antigens, and prevent autoimmune disease. Tregs are immunosuppressive and generally suppress or downregulate induction and proliferation of effector T (Teff) cells. Tregs can develop in the thymus (so-called CD4+ Foxp3+ “natural” Tregs) or differentiate in the periphery upon priming of naïve CD4+ T cells by antigen-presenting cells (APCs), for example, following exposure to TGF β or retinoic acid. Treg cells produce and secrete cytokines including IL-10 and TGF β 1. Generally, differentiation of Treg and Th17 cells is negatively correlated.

[135] *Specific binding*: As used herein, the term “specific binding” or “specifically binds” means that the interaction of the antibody, or antigen-binding portion thereof, with an antigen is dependent upon the presence of a particular structure (*e.g.*, an antigenic determinant or epitope). For example, the antibody, or antigen-binding portion thereof, binds to a specific protein rather than to proteins

generally. In some embodiments, an antibody, or antigen-binding portion thereof, specifically binds to a target, *e.g.*, TGF β 1, if the antibody has a K_D for the target of at least about 10^{-6} M. More preferably, the measured K_D values of such antibody range between 10-100 nM. More preferably, the measured K_D values of such antibody range between 0.1-10 nM.

[136] *Subject*: The term “subject” in the context of therapeutic applications refers to an individual who receives clinical care or intervention, such as treatment, diagnosis, etc. Suitable subjects include vertebrates, including but not limited to mammals (*e.g.*, human and non-human mammals). Where the subject is a human subject, the term “patient” may be used interchangeably. In a clinical context, the term “a patient population” or “patient subpopulation” is used to refer to a group of individuals that falls within a set of criteria, such as clinical criteria (*e.g.*, disease presentations, disease stages, susceptibility to certain conditions, responsiveness to therapy, etc.), medical history, health status, gender, age group, genetic criteria (*e.g.*, carrier of certain mutation, polymorphism, gene duplications, DNA sequence repeats, etc.) and lifestyle factors (*e.g.*, smoking, alcohol consumption, exercise, etc.).

[137] *TGF β inhibitor*: The term “TGF β inhibitor” refers to any agent capable of antagonizing biological activities or function of TGF β growth factor (*e.g.*, TGF β 1, TGF β 2 and/or TGF β 3). The term is not intended to limit its mechanism of action and includes, for example, neutralizing inhibitors, receptor antagonists, soluble ligand traps, and activation inhibitors of TGF β .

[138] *T helper 17 cell*: T helper 17 cells (Th17) are a subset of pro-inflammatory T helper cells characterized by the markers STAT3 and ROR γ t and the production of cytokines including interleukin 17 (IL-17A/F) and IL-22. Th17 cells are differentiated when naive T cells are exposed to TGF β and IL-6. Th17 cells are generally associated with tissue inflammation, autoimmunity and clearance of certain pathogens. The differentiation of Th17 cells and Treg cells is generally inversely related. Imbalance in Th17-to-Treg ratios (*e.g.*, “Th17/Treg”) has been implicated in a number of pathologies, such as fibrotic conditions and autoimmune conditions.

[139] *Th17/Treg ratio*: Th17-to-Treg ratios refer to measured ratios (relative proportions) of the number of Th17 cells versus the number of Treg cells in a tissue or sample of interest. Typically, known cell markers are used to identify, sort or isolate the cell types. Such markers include cell-surface molecules expressed on the particular cell type; a cytokine or a panel of cytokines produced (*e.g.*, secreted) by the particular cell type, and/or mRNA expression of certain gene markers that serve as a signature/profile of the particular cell type. For example, the Th17/Treg ratio of one (1) means that there is an equal or equivalent number of each of the cell types within the tissue or sample being evaluated. The Th17/Treg ratio of two (2) means that there is approximately twice the number of Th17 cells as compared to Treg cells in the tissue or sample. An elevated Th17/Treg ratio may arise from an increased number of Th17 cells, a decreased number of Treg cells, or combination thereof.

[140] *Therapeutic window*: The term “therapeutic window” refers to a range of doses / concentrations that produces therapeutic response without causing significant / observable / unacceptable adverse effect (*e.g.*, within adverse effects that are acceptable or tolerable) in subjects. Therapeutic window may be calculated as a ratio between minimum effective concentrations (MEC) to the minimum toxic concentrations (MTC). To illustrate, a TGF β 1 inhibitor that achieves in vivo efficacy at 10 mg/kg and shows tolerability or acceptable toxicities at 100 mg/kg provides at least a 10-fold (*e.g.*, 10x) therapeutic window. By contrast, a pan-inhibitor of TGF β that is efficacious at 10 mg/kg but causes adverse effects at 5 mg/kg is said to have “dose-limiting toxicities.” For example, the applicants have found that a context-independent TGF β 1 inhibitor antibody is efficacious at dosage ranging between about <3 and 30 mg/kg/week and is free of observable toxicities associated with pan-inhibition of TGF β at least 100 mg/kg/week for 4 weeks in preclinical models such as rats. Based on this, the context-independent TGF β 1 inhibitor antibody shows at minimum a 3.3-fold and up to 33-fold therapeutic window.

[141] *Toxicity*: As used herein, the term “toxicity” or “toxicities” refers to unwanted in vivo effects in patients associated with a therapy administered to the patients, such as undesirable side effects and adverse events. “Tolerability” refers to a level of toxicities associated with a therapy or therapeutic regimen, which can be reasonably tolerated by patients, without discontinuing the therapy due to the toxicities (*i.e.*, acceptable level of toxicities). Typically, toxicity/toxicology studies are carried out in one or more preclinical models prior to clinical development to assess safety profiles of a drug candidate (*e.g.*, monoclonal antibody therapy). Toxicity/toxicology studies may help determine the “no observed adverse effect level (NOAEL)” and the “maximally tolerated dose (MTD)” of a test article, based on which a therapeutic window may be deduced. Preferably, a species that is shown to be sensitive to the particular intervention should be chosen as a preclinical animal model in which safety/toxicity study is to be carried out. In case of TGF β inhibition, suitable species include rats, dogs, and cynos. Mice are reported to be less sensitive to pharmacological inhibition of TGF β and may not reveal toxicities that are potentially dangerous in other species, including human, although certain studies report toxicities observed with pan-inhibition of TGF β in mice. To illustrate, the NOAEL for a context-independent TGF β 1 inhibitor antibody in rats was the highest dose evaluated (100 mg/kg), suggesting that the MTD is >100 mg/kg per week, based on a four-week toxicology study.

[142] *Treat/treatment*: The term “treat” or “treatment” includes therapeutic treatments, prophylactic treatments, and applications in which one reduces the risk that a subject will develop a disorder or other risk factor. Thus the term is intended to broadly mean: causing therapeutic benefits in a patient by, for example, enhancing or boosting the body’s immunity; reducing or reversing immune suppression; reducing, removing or eradicating harmful cells or substances from the body; reducing disease burden (*e.g.*, tumor burden); preventing recurrence or relapse; prolonging a refractory period,

and/or otherwise improving survival. Treatment does not require the complete curing of a disorder and encompasses embodiments in which one reduces symptoms or underlying risk factors. In the context of combination therapy, the term may also refer to: i) the ability of a second therapeutic to reduce the effective dosage of a first therapeutic so as to reduce side effects and increase tolerability; ii) the ability of a second therapy to render the patient more responsive to a first therapy; and/or iii) the ability to effectuate additive or synergistic clinical benefits.

[143] *Variable region*: The term “variable region” or “variable domain” refers to a portion of the light and/or heavy chains of an antibody, typically including approximately the amino-terminal 120 to 130 amino acids in the heavy chain and about 100 to 110 amino terminal amino acids in the light chain. In certain embodiments, variable regions of different antibodies differ extensively in amino acid sequence even among antibodies of the same species. The variable region of an antibody typically determines specificity of a particular antibody for its target.

TGFβ1

[144] In mammals, the transforming growth factor-beta (TGFβ) superfamily is comprised of at least 33 gene products. These include the bone morphogenetic proteins (BMPs), activins, growth and differentiation factors (GDFs), and the three isoforms of the TGFβ family: TGFβ1, TGFβ2, and TGFβ3. The TGFβs are thought to play key roles in diverse processes, such as inhibition of cell proliferation, extracellular matrix (ECM) remodeling, and immune homeostasis. The importance of TGFβ1 for T cell homeostasis is demonstrated by the observation that TGFβ1^{-/-} mice survive only 3-4 weeks, succumbing to multiorgan failure due to massive immune activation (Kulkarni, A.B., *et al.*, Proc Natl Acad Sci U S A, 1993. 90(2): p. 770-4; Shull, M.M., *et al.*, Nature, 1992. 359(6397): p. 693-9). The roles of TGFβ2 and TGFβ3 are less clear. Whilst the three TGFβ isoforms have distinct temporal and spatial expression patterns, they signal through the same receptors, TGFβRI and TGFβRII, although in some cases, for example for TGFβ2 signaling, type III receptors such as betaglycan are also required (Feng, X.H. and R. Derynck, Annu Rev Cell Dev Biol, 2005. 21: p. 659-93; Massague, J., Annu Rev Biochem, 1998. 67: p. 753-91). Ligand-induced oligomerization of TGFβRI/II triggers the phosphorylation of SMAD transcription factors, resulting in the transcription of target genes, such as Col1a1, Col3a1, ACTA2, and SERPINE1 (Massague, J., J. Seoane, and D. Wotton, Genes Dev, 2005. 19(23): p. 2783-810). SMAD-independent TGFβ signaling pathways have also been described, for example in cancer or in the aortic lesions of Marfan mice (Derynck, R. and Y.E. Zhang, Nature, 2003. 425(6958): p. 577-84; Holm, T.M., *et al.*, Science, 2011. 332(6027): p. 358-61).

[145] The biological importance of the TGFβ pathway in humans has been validated by genetic diseases. Camurati-Engelman disease results in bone dysplasia due to an autosomal dominant mutation in the TGFB1 gene, leading to constitutive activation of TGFβ1 signaling (Janssens, K., *et*

al., J Med Genet, 2006. 43(1): p. 1-11). Patients with Loeys/Dietz syndrome carry autosomal dominant mutations in components of the TGF β signaling pathway, which cause aortic aneurism, hypertelorism, and bifid uvula (Van Laer, L., H. Dietz, and B. Loeys, Adv Exp Med Biol, 2014. 802: p. 95-105). As TGF β pathway dysregulation has been implicated in multiple diseases, several drugs that target the TGF β pathway have been developed and tested in patients, but with limited success. Most TGF β inhibitors described to date lack isoform specificity as briefly summarized below.

[146] Fresolimumab, a humanized monoclonal antibody that binds and inhibits all three isoforms of TGF β has been tested clinically in patients with focal segmental glomerulosclerosis, malignant melanoma, renal cell carcinoma, and systemic sclerosis (Rice, L.M., *et al.*, J Clin Invest, 2015. 125(7): p. 2795-807; Trachtman, H., *et al.*, Kidney Int, 2011. 79(11): p. 1236-43; Morris, J.C., *et al.*, PLoS One, 2014. 9(3): p. e90353). Additional companies have developed monoclonal antibodies against the TGF β growth factors with varying degrees of selectivity for TGF β isoforms. Such agents likely elicit toxicities *in vivo* through residual activity against other TGF β family members besides TGF β 1. This lack of isoform specificity may be due to the high degree of sequence identity between isoforms.

[147] Other approaches to target the TGF β pathway include ACE-1332, a soluble TGF β RII-Fc ligand trap from Acceleron (Yung, L.M., *et al.*, A Am J Respir Crit Care Med, 2016. 194(9): p. 1140-1151), or small molecule inhibitors of the ALK5 kinase, such as Eli Lilly's galunisertib. ACE-1332 binds TGF β 1 and TGF β 3 with equally high affinity (Yung, L.M., *et al.*, Am J Respir Crit Care Med, 2016. 194(9): p. 1140-1151), and ALK5 inhibitors block the activity of all growth factors that signal through TGF β R1. Substantial toxicities have been found in preclinical studies using ALK5 inhibitors (Anderton, M.J., *et al.*, Toxicol Pathol, 2011. 39(6): p. 916-24; Stauber, A., *et al.*, Clinical Toxicology, 2014. 4(3): p. 1-10), and sophisticated clinical dosing schemes are required to maintain efficacy while reducing adverse events (Herbertz, S., *et al.*, Drug Des Devel Ther, 2015. 9: p. 4479-99). In fact, the question of TGF β signaling specificity and its possible effect on toxicity observed with the known TGF β inhibitors has not been raised in most, if not all, of the candidate drugs that attempted to block TGF β . For example, how much of the toxicities are due to inhibition of TGF β 1 versus TGF β 2 and/or TGF β 3 has not been addressed. Similarly, modes of TGF β activation have not been taken into account in designing or developing ways to antagonize TGF β signaling.

[148] Recent structural insights into the activation mechanism of TGF β 1 (Shi, M., *et al.*, Nature, 2011. 474(7351): p. 343-9) have enabled more specific approaches to TGF β inhibition (see, *e.g.*, PCT/US2017/21972, the entire contents of which are incorporated herein by reference). Unlike other cytokines, TGF β superfamily members are not secreted as active growth factors, but as dimeric pro-proteins which consist of an N-terminal prodomain and a C-terminal growth factor domain. Cleavage of proTGF β 1 by furin proteases separates the homodimeric growth factor domain from its prodomain, also referred to as latency associated peptide (LAP). However, the growth factor and LAP remain

noncovalently associated, forming a latent complex which is unable to bind its receptors and induce signaling. During translation, latent TGF β 1, also called the small latent complex (SLC), becomes linked to “presenting molecules” via disulfide bridges, forming the large latent complex (LLC). These molecules allow proTGF β 1 to be presented in specific cellular or tissue contexts. Two cysteines near the N-terminus of the latent TGF β 1 link to appropriately positioned cysteines on the presenting molecule. The identity of the presenting molecule depends on the environment and cell type producing latent TGF β 1. For example, fibroblasts secrete latent TGF β 1 tethered to latent TGF β -binding proteins (LTBPs), which then associate with proteins in the extracellular matrix (ECM) (*i.e.*, fibronectin, fibrillin-1) to link latent TGF β to the ECM (Robertson *et al.* Matrix Biol 47: 44-53 (2015) (FIG. 2A). On the surface of activated regulatory T cells latent TGF β 1 is covalently linked to the transmembrane protein GARP (glycoprotein-A repetitions predominant protein (GARP), and a protein closely related to GARP, LRRC33 (leucine-rich repeat-containing protein 33), serves as a presenting molecule for TGF β 1 on the surface of monocytes, macrophages and microglia (Wang, R., *et al.*, Mol Biol Cell, 2012. 23(6): p. 1129-39 and T.A. Springer, Int. BMP Conference 2016).

[149] A number of studies have shed light on the mechanisms of TGF β 1 activation. Three integrins, α V β 6, α V β 8, and α V β 1 have been demonstrated to be key activators of latent TGF β 1 (Reed, N.I., *et al.*, Sci Transl Med, 2015. 7(288): p. 288ra79; Travis, M.A. and D. Sheppard, Annu Rev Immunol, 2014. 32: p. 51-82; Munger, J.S., *et al.*, Cell, 1999. 96(3): p. 319-28). α V integrins bind the RGD sequence present in TGF β 1 and TGF β 1 LAPs with high affinity (Dong, X., *et al.*, Nat Struct Mol Biol, 2014. 21(12): p. 1091-6). Transgenic mice with a mutation in the TGF β 1 RGD site that prevents integrin binding, but not secretion, phenocopy the TGF β 1 $^{-/-}$ mouse (Yang, Z., *et al.*, J Cell Biol, 2007. 176(6): p. 787-93). Mice that lack both β 6 and β 8 integrins recapitulate all essential phenotypes of TGF β 1 and TGF β 3 knockout mice, including multiorgan inflammation and cleft palate, confirming the essential role of these two integrins for TGF β 1 activation in development and homeostasis (Aluwihare, P., *et al.*, J Cell Sci, 2009. 122(Pt 2): p. 227-32). Key for integrin-dependent activation of latent TGF β 1 is the covalent tether to presenting molecules; disruption of the disulfide bonds between GARP and TGF β 1 LAP by mutagenesis does not impair complex formation, but completely abolishes TGF β 1 activation by α V β 6 (Wang, R., *et al.*, Mol Biol Cell, 2012. 23(6): p. 1129-39). The recent structure of latent TGF β 1 illuminates how integrins enable release of active TGF β 1 from the latent complex: the covalent link of latent TGF β 1 to its presenting molecule anchors latent TGF β 1, either to the ECM through LTBPs, or to the cytoskeleton through GARP or LRRC33. Integrin binding to the RGD sequence results in a force-dependent change in the structure of LAP, allowing active TGF β 1 to be released and bind nearby receptors (Shi, M., *et al.*, Nature, 2011. 474(7351): p. 343-9). The importance of integrin-dependent TGF β 1 activation in disease has also been well validated. A small molecular inhibitor of α V β 1 protects against bleomycin-induced lung fibrosis and carbon tetrachloride-induced liver fibrosis (Reed, N.I., *et al.*, Sci Transl Med, 2015).

7(288): p. 288ra79), and $\alpha V\beta 6$ blockade with an antibody or loss of integrin $\beta 6$ expression suppresses bleomycin-induced lung fibrosis and radiation-induced fibrosis (Munger, J.S., *et al.*, Cell, 1999. 96(3): p. 319-28); Horan, G.S., *et al.*, Am J Respir Crit Care Med, 2008. 177(1): p. 56-65). In addition to integrins, other mechanisms of TGF β 1 activation have been implicated, including thrombospondin-1 and activation by proteases such as matrix metalloproteinases (MMPs), cathepsin D or kallikrein. However, the majority of these studies were performed *in vitro* using purified proteins; there is less evidence for the role of these molecules from *in vivo* studies. Knockout of thrombospondin-1 recapitulates some aspects of the TGF β 1-/- phenotype in some tissues, but is not protective in bleomycin-induced lung fibrosis, known to be TGF β -dependent (Ezzie, M.E., *et al.*, Am J Respir Cell Mol Biol, 2011. 44(4): p. 556-61). Additionally, knockout of candidate proteases did not result in a TGF β 1 phenotype (Worthington, J.J., J.E. Klementowicz, and M.A. Travis, Trends Biochem Sci, 2011. 36(1): p. 47-54). This could be explained by redundancies or by these mechanisms being critical in specific diseases rather than development and homeostasis.

[150] TGF β has been implicated in a number of biological processes, including fibrosis, immune-modulation and cancer progression. TGF β 1 was the first identified member of the TGF β superfamily of proteins. Like other members of the TGF β superfamily, TGF β 1 and the isoforms TGF β 2 and TGF β 3, are initially expressed as inactive precursor pro-protein forms (termed proTGF β). TGF β proteins (*e.g.*, TGF β 1, TGF β 2 and TGF β 3) are proteolytically cleaved by proprotein convertases (*e.g.*, furin) to yield the latent form (termed latent TGF β). In some embodiments, a pro-protein form or latent form of a TGF β protein (*e.g.*, TGF β 1, TGF β 2 and TGF β 3) may be referred to as “pro/latent TGF β protein”. TGF β 1 may be presented to other molecules in complex with multiple molecules including, for example, GARP (to form a GARP-TGF β 1 complex), LRRC33 (to form a LRRC33-TGF β 1 complex), LTBP1 (to form a LTBP1-TGF β 1 complex), and/or LTBP3 (to form a LTBP3-TGF β 1 complex). The TGF β 1 present in these complexes may be in either latent form (latent TGF β 1) or in precursor form (proTGF β 1).

Isoform Selectivity and Mechanisms of Action of TGF inhibitors

[151] From a safety standpoint, there has been an increasing recognition that broad inhibition of TGF β across isoforms may be a cause of observed toxicities, which underscores the fact that no TGF β inhibitors have been successfully developed to this day. To circumvent potentially dangerous adverse effects, a number of groups have recently turned to identifying inhibitors that target a *subset* - but not all - of the isoforms and still retain efficacy. From an efficacy standpoint, however, the prevailing view of the field remains to be that it is advantageous to inhibit multiple isoforms of TGF β to achieve therapeutic effects, and to accommodate this, toxicity management by “careful dosing regimen” is suggested as a solution (Brennan et al. (2018) mAbs, 10:1, 1-17). Consistent with this premise, numerous groups are developing TGF β inhibitors that target more than one isoforms. These include low molecular weight antagonists of TGF β receptors, *e.g.*, ALK5 antagonists, such as Galunisertib

(LY2157299 monohydrate); monoclonal antibodies (such as neutralizing antibodies) that inhibit all three isoforms (“pan-inhibitor” antibodies) (see, for example, WO 2018/134681); monoclonal antibodies that preferentially inhibit two of the three isoforms (*e.g.*, antibodies against TGF β 1/2 (for example WO 2016/161410) and TGF β 1/3 (for example WO 2006/116002); and engineered molecules (*e.g.*, fusion proteins) such as ligand traps (for example, WO 2018/029367; WO 2018/129331 and WO 2018/158727). Similarly, inhibitors of integrins such as α V β 6 also block integrin-dependent activation of both TGF β 1 and TGF β 3 and therefore may be considered as isoform-non-selective inhibitors of TGF β signaling. In addition, examples of antibodies that selectively bind and neutralize both TGF β 1 and TGF β 2 (*i.e.*, TGF β 1/2 inhibitors) include XOMA 089 (or NIS793) and variants (see, for example, WO 2016/161410).

[152] Previously, Applicant demonstrated that inhibition of TGF β 1 alone was sufficient to sensitize immunosuppressive tumors to a checkpoint inhibitor therapy even in tumors where both TGF β 1/3 are co-expressed (PCT/US2019/041373). Similarly, TGF β 1-selective inhibitors are shown to mitigate fibrosis in preclinical models, including mouse liver fibrosis model where both the TGF β 1/3 isoforms are co-expressed in the fibrotic tissue, albeit in discrete cell types, as observed by immunohistochemistry (data now shown). Surprisingly, inhibition of TGF β 3 promoted *pro-fibrotic* phenotypes. The exacerbation of fibrosis is observed when the TGF β 3 inhibitor is used alone. In addition, when used in combination with a TGF β 1-selective inhibitor, the TGF β 3 inhibitor *attenuated* the anti-fibrotic effect of the TGF β 1-selective inhibitor, as evidenced by increased collagen accumulation in the fibrotic liver. These results raise the possibility that inhibitory potency against TGF β 3 may be an undesirable feature of TGF β inhibitors to be used as therapy in situations where fibrosis is a concern.

[153] Beyond the fibrosis context, there is a broader implication to this unexpected finding since the pro-fibrotic phenotype (*e.g.*, increased collagen deposit into the ECM) is associated not only with fibrosis, but also with aspects of cancer progression, such as tumor invasion and metastasis. See, for example, Chakravarthy et al. (Nature Communications, (2018) 9:4692. “TGF- β -associated extracellular matrix genes link cancer-associated fibroblasts to immune evasion and immunotherapy failure”). Diseased tissues with dysregulated ECM, including fibrotic tissues and stroma of various tumor types, can express both TGF β 1 and TGF β 3. As of today, multiple groups are making effort to develop TGF β inhibitors that target both of these isoforms, such as ligand traps, neutralizing antibodies and integrin inhibitors. However, the finding presented herein cautions that such approach may in fact exacerbate (*e.g.*, worsen) the disease.

[154] Accordingly, the present disclosure provides the teaching that for the treatment of a disorder involving ECM dysregulation, such as fibrosis and cancer, a TGF β inhibitor that does *not* specifically target TGF β 3 should be selected. Preferably, such inhibitor is an isoform-selective inhibitor of TGF β 1, such as inhibitors that selectively target LTBP1/3-associated TGF β 1 (*e.g.*, as disclosed

herein). Related methods include a method for selecting a TGF β inhibitor for use in the treatment of a fibrotic disorder in a subject, wherein the method includes the steps of: testing potency of one or more candidate inhibitors for the ability to inhibit TGF β 1, TGF β 2 and TGF β 3, and selecting an inhibitor that inhibits TGF β 1 but does not inhibit TGF β 3, for therapeutic use. Related treatment methods can further comprise a step of administering to the subject the inhibitor that inhibits TGF β 1 but does not inhibit TGF β 3 in an amount sufficient to treat the fibrotic disorder or treat a subject having or at risk of developing a fibrotic disorder. Preferably, the selected inhibitor is an antibody or fragment thereof that selectively inhibits LTBP1- and/or LTBP3-associated TGF β 1 signaling (*e.g.*, as disclosed herein). In some embodiments, subjects at risk of developing a fibrotic disorder may suffer from a metabolic disorder, such as diabetes, obesity and NASH. The proposed exclusion of the subpopulation of patients is aimed to reduce risk of triggering, facilitating or exacerbating a pro-fibrotic effect.

[155] In addition to the possible concerns of inhibiting TGF β 3 addressed above, Takahashi et al. (Nat Metab. 2019, 1(2): 291-303) recently reported a beneficial role of TGF β 2 in regulating metabolism. The authors identified TGF β 2 as an exercise-induced adipokine, which stimulated glucose and fatty acid uptake in vitro, as well as tissue glucose uptake in vivo; which improved metabolism in obese mice; and, which reduced high fat diet-induced inflammation. Moreover, the authors observed that lactate, a metabolite released from muscle during exercise, stimulated TGF β 2 expression in human adipocytes and that a lactate-lowering agent reduced circulating TGF β 2 levels and reduced exercise-stimulated improvements in glucose tolerance. These observations suggest that therapeutic use of a TGF β inhibitor with inhibitory activity towards the TGF β 2 isoform may be harmful at least in the metabolic aspect.

[156] Without being bound by particular theory, it is contemplated that it is advantageous to select a TGF β 1-selective inhibitor as a TGF β inhibitor for use in the treatment of a metabolic disease, such as liver fibrosis associated with NASH. In preferred embodiments, the TGF β 1-selective inhibitor selected for use in the treatment of the metabolic disease selectively inhibits LTBP1/3-associated TGF β 1, such as the antibodies and fragments disclosed herein. Accordingly, the invention includes a TGF β inhibitor for use in the treatment of a metabolic disease in a subject, wherein the treatment comprises selection of a TGF β inhibitor that inhibits TGF β 1 but does not inhibit TGF β 2, optionally wherein the inhibitor is TGF β 1-selective, and administration of the inhibitor to a subject suffering from a metabolic disease. The metabolic disease may be a liver disease, such as liver fibrosis, NASH, NAFLD, optionally accompanied by obesity and/or type 2 diabetes. In preferred embodiments, the TGF β 1-selective inhibitor is an antibody or antigen-binding fragment thereof that selectively targets matrix-associated TGF β 1 (*e.g.*, LTBP1-proTGF β 1 and LTBP3-proTGF β 1), such as those disclosed herein.

[157] In preferred embodiments, a TGF β inhibitor for use in the treatment of a fibrotic disorder is an isoform-selective activation inhibitor of TGF β 1 (such as the novel antibodies with low k_{OFF} or long

t½ disclosed herein) capable of targeting matrix-associated TGFβ1-containing latent complexes in vivo.

[158] The antibodies of the present disclosure work by preventing the step of TGFβ1 activation. In some embodiments, such inhibitors can inhibit integrin-dependent (*e.g.*, mechanical or force-driven) activation of TGFβ1. In some embodiments, such inhibitors can inhibit protease-dependent or protease-induced activation of TGFβ1. The latter includes inhibitors that inhibit the TGFβ1 activation step in an integrin-independent manner. In some embodiments, such inhibitors can inhibit TGFβ1 activation irrespective of the mode of activation, *e.g.*, inhibit both integrin-dependent activation and protease-dependent activation of TGFβ1. Non-limiting examples of proteases which may activate TGFβ1 include serine proteases, such as Kallikreins, Chymotrypsin, Trypsin, Elastases, Plasmin, thrombin, as well as zinc metalloproteases (MMP family) such as MMP-2, MMP-9, MMP-12, MMP-13 and ADAM proteases (*e.g.*, ADAM10 and ADAM17). Kallikreins include plasma-Kallikreins and tissue Kallikreins, such as KLK1, KLK2, KLK3, KLK4, KLK5, KLK6, KLK7, KLK8, KLK9, KLK10, KLK11, KLK12, KLK13, KLK14 and KLK15.

Latent TGFβ-Binding Proteins (LTBPs)

[159] In mammals there are four known LTBPs, LTBP1-4, each with multiple splice variants (Robertson, I.B., *et al.*, Matrix Biol, 2015. 47: p. 44-53). LTBP2 is the only LTBP that does not associate with latent TGFβ (Saharinen, J. and J. Keski-Oja, Mol Biol Cell, 2000. 11(8): p. 2691-704). While the association between LTBP1 or LTBP3 and latent TGFβ1 has been well validated, the role of LTBP4 in TGFβ presentation is less clear. The complex with LTBP4 and latent TGFβ1 appears to form much less efficiently, potentially due to the absence of several negatively charged residues in the TGFβ-binding domain of LTBP4 (Saharinen, J. and J. Keski-Oja, Mol Biol Cell, 2000. 11(8): p. 2691-704; Chen, Y., *et al.*, J Mol Biol, 2005. 345(1): p. 175-86). Both LTBP4S^{-/-} mice and Urban-Rifkin-Davis syndrome patients, who have null mutations in LTBP4, suffer from disrupted elastic fiber assembly (Urban, Z., *et al.*, Am J Hum Genet, 2009. 85(5): p. 593-605; Dabovic, B., *et al.*, J Cell Physiol, 2015. 230(1): p. 226-36). Additionally, while LTBP4S^{-/-} mice have a lung septation and an elastogenesis defect, transgenic mice with an LTBP4 that cannot form a complex with latent TGFβ1 have no obvious phenotype (Dabovic, B., *et al.*, J Cell Physiol, 2015. 230(1): p. 226-36). Whether LTBP4 is directly involved in regulation of latent TGFβ1 by functioning as a presenting molecule is unclear; LTBP4 may instead be required for proper formation of elastic fibrils in the ECM and its loss indirectly affect latent TGFβ1 activation through defects in the ECM.

[160] In one aspect, the present invention is directed to inhibitors, *e.g.*, immunoglobulins, *e.g.*, antibodies, or antigen-binding portions thereof, that selectively bind to a complex containing a TGFβ pro-protein and a LTBP protein (*e.g.*, LTBP1 or LTBP3). In a preferred embodiment, the TGFβ protein is TGFβ1. In some embodiments, the binding molecules disclosed herein bind selectively to a

complex containing pro/latent TGF β 1 and LTBP1 or LTBP3. Such binding molecules can allow TGF β 1 activity to be selectively modulated in a context-dependent manner, *i.e.*, by modulating TGF β 1 in the context of a LTPB protein, without modulating the activity of TGF β 1 complexed with other presenting molecules (*e.g.*, GARP and/or LRRC33).

Antibodies that Selectively Inhibit LTBP-Mediated TGF β activation

[161] The present invention provides novel, TGF β inhibitors that selectively target matrix- or ECM-associated TGF β activities. More specifically, such inhibitors include isoform-specific, context-selective inhibitors of TGF β 1 activation that specifically bind latent forms of TGF β 1 (*e.g.*, proTGF β 1 complex) within the ECM environment and prevent release of mature growth factor from the complex at the niche. Such matrix-targeting inhibitors are context-specific in that they selectively bind proTGF β 1 associated with ECM presenting molecules, namely, LTBP1 and/or LTBP3. Thus, disclosed herein are monoclonal antibodies and fragments thereof capable of binding an epitope present in an LTBP1-proTGF β 1 complex and/or LTBP3-proTGF β 1 complex, whereas the epitope is not present in a GARP-proTGF β 1 complex and/or LRRC33-proTGF β 1 complex.

[162] In some embodiments, the context-selective inhibitors of the present disclosure are capable of specifically binding both a human LTBP1-proTGF β 1 complex and a human LTBP3-proTGF β 1 complex, with affinities of < 5 nM each (measured K_D values) in a suitable *in vitro* binding assay, such as Octet. In preferred embodiments, such antibodies bind both a human LTBP1-proTGF β 1 complex and a human LTBP3-proTGF β 1 with affinities of < 5 nM each (measured K_D values) in a suitable *in vitro* binding assay, such as Octet. On the other hand, these context-specific antibodies do not show any detectable binding to a human GARP-proTGF β 1 complex or a human LRRC33-proTGF β 1 complex under the same assay conditions. Preferably, such antibody or the fragment binds each of the human LTBP1-proTGF β 1 complex and the human LTBP3-proTGF β 1 complex with K_D of less than 1 nM.

[163] In some embodiments, the context-selective inhibitors of the present disclosure are capable of specifically binding either a human LTBP1-proTGF β 1 complex or a human LTBP3-proTGF β 1 complex. Neither shows any detectable binding to a human GARP-proTGF β 1 complex or a human LRRC33-proTGF β 1 complex under the same assay conditions.

[164] The art is familiar with suitable *in vitro* binding assays, including, for example, BLI-based assays such as Octet and SPR-based assays such as Biacore, which can be used to measure antibody-antigen interactions (*e.g.*, binding kinetics). As used herein, “no binding” in these contexts may refer to no detectable binding by a particular assay, *e.g.*, the binding, if any, is below the sensitivity of the assay. In some embodiments, “no binding” may refer to “no meaningful binding”, set by a cutoff, which defines the minimum level required to be considered meaningful as measured by a particular assay system. For example, in a BLI (Octet) assay, 0.1 nm of optical shift measured at predetermined

analyte (*e.g.*, target antigen) concentrations (*e.g.*, 100 nM, 200 nM, etc.) may indicate meaningful binding, below which may be considered as no binding (see for example, Example 9). Similarly, in a typical SPR (Biacore) assay, the cutoff level may be 0.2 RU (resonance unit). In some embodiments, the signal at 0 nM antibody (*e.g.*, background noise) is subtracted from all the sensorgrams obtained at higher concentrations. Under these conditions, using SPR (Biacore), 0.2 RUs may be a suitable cutoff.

[165] In some embodiments, the context-selective inhibitors of the present disclosure are capable of specifically binding a human LTBP1-proTGFβ1 complex or a human LTBP3-proTGFβ1 complex without showing detectable binding to a human GARP-proTGFβ1 complex, as measured by BLI, under the same assay conditions as used to measure binding to human LTBP1-proTGFβ1 complex and/or human LTBP3-TGFβ1.

[166] In some embodiments, the context-selective inhibitors of the present disclosure bind a human LTBP1-proTGFβ1 complex or a human LTBP3-proTGFβ1 complex with a K_D that is at least 50 times lower (*e.g.*, at least 75 times lower, at least 100 times lower) than the K_D when binding to a human GARP-proTGFβ1 complex under the same assay conditions. In some embodiments, the K_D is as determined by BLI or SPR. In some embodiments, the K_D is as determined by SPR.

[167] In some embodiments, the context-selective inhibitors of the present disclosure are capable of specifically binding a human LTBP1-proTGFβ1 complex or a human LTBP3-proTGFβ1 complex without showing detectable binding to an LRRC33-proTGFβ1 latent complex, as measured by BLI, under the same assay conditions as used to measure binding to human LTBP1-proTGFβ1 complex and/or human LTBP3-TGFβ1.

[168] In some embodiments, the context-selective inhibitors of the present disclosure bind a human LTBP1-proTGFβ1 complex or a human LTBP3-TGFβ1 complex with a K_D that is at least 50 times lower (*e.g.*, at least 75 times lower, at least 100 times lower) than the K_D when binding to a human LRRC33-proTGFβ1 complex under the same assay conditions. In some embodiments, the K_D is as determined by BLI or SPR. In some embodiments, the K_D is as determined by SPR.

[169] The invention includes the recognition that preferred antibodies (*e.g.*, immunoglobulins and antigen-binding fragments such as Fabs, as well as engineered constructs incorporating such fragments), once bound to its target/antigen (*e.g.*, human LTBP1-proTGFβ1, human LTBP3-TGFβ1), dissociates slowly from the antigen. Thus, the novel antibodies of the instant invention are selected not only for their high overall affinities (such as K_D of no more than 5 nM) but especially for their low dissociation rates. According to the present disclosure, such antibodies have dissociation rates of $\leq 5 \times 10^{-4}$ (1/s), as measured by BLI (*e.g.*, when binding to human LTBP1-proTGFβ1 and/or human LTBP3-TGFβ1). Such dissociation rates of $\leq 5 \times 10^{-4}$ (1/s) of the antibodies or antigen-binding fragments may be monovalent dissociation rates or divalent dissociation rates. In some embodiments, the antibody or the fragment dissociates slowly from human LTBP1-proTGFβ1 and/or human

LTBP3-TGF β 1, preferably for both human LTBP1-proTGF β 1 and human LTBP3-TGF β 1, with a monovalent dissociation half-time ($t_{1/2}$) of at least 45 minutes (*e.g.*, $\geq$ 45, 60, 75, 90 minutes) as measured by SPR. On the other hand, should binding to a human GARP-proTGF β 1 and/or LRRC33-TGF β 1 complex be detectable, such antibody dissociates rapidly from a human GARP-proTGF β 1 and/or human LRRC33-TGF β 1 complex(es), particularly the human GARP-proTGF β 1. In some embodiments, the antibody dissociates from human GARP-proTGF β 1 with $t_{1/2}$ of less than 5 minutes as measured by SPR. In particularly preferred embodiments, the antibody or the fragments show species cross-reactivity such that they bind murine counterparts with equivalent affinities.

[170] The TGF β 1 present in these complexes may be in either latent form (latent TGF β 1) or in precursor form (proTGF β 1). In one embodiment, the inhibitors do not significantly bind to LTBP1 alone (*e.g.*, when not complexed with TGF β 1). In another embodiment, the inhibitors do not significantly bind to LTBP3 alone (*e.g.*, when not complexed with TGF β 1). In another embodiment, the inhibitors do not significantly bind to TGF β 1 alone (*e.g.*, pro or latent TGF β 1 not complexed with LTBP1 or LTBP3, or mature TGF β 1). In another embodiment, the inhibitors that selectively bind a LTBP1-TGF β 1 complex and/or a LTBP3-TGF β 1 complex do not significantly bind to a complex containing TGF β 1 and another presenting molecule, *e.g.*, a GARP-TGF β 1 complex (*e.g.*, GARP complexed to pro- or latent TGF β 1) and/or a LRRC33-TGF β 1 complex (*e.g.*, LRRC33 complexed to pro- or latent TGF β 1). In one embodiment, the inhibitors that selectively bind LTBP1/3-TGF β 1 do not significantly bind one or more (*e.g.*, two or more, three or more, or all four) of the following: LTBP1 alone, TGF β 1 alone, a GARP-TGF β 1 complex, and a LRRC33-TGF β 1 complex. In addition, in some embodiments, the inhibitors do not significantly bind LTBP3 alone.

[171] As used herein, the term “inhibitor” refers to any agent capable of blocking or antagonizing TGF β 1 signaling. Such agents may include small molecule antagonists of TGF β 1 and biologic antagonists of TGF β 1 (*e.g.*, protein fragments and antibodies). In some embodiments, the inhibitor may be an antibody (including fragments thereof, such as Domain Antibodies (dAbs) as described in, for example, U.S. Patent 6,291,158; 6,582,915; 6,593,081; 6,172,197; and 6,696,245), a small molecule inhibitor, an Adnectin, an Affibody, a DARPin, an Anticalin, an Avimer, a Versabody or a gene therapy. Use of inhibitors encompassed by the present invention also includes antibody mimetics, such as monobodies and single-domain antibodies. Monobodies are synthetic binding proteins that typically employ a fibronectin type III domain (FN3) as a molecular scaffold. Monobodies include AdnectinsTM which are based on the 10th fibronectin type III domain.

[172] In some aspects, the inhibitors, *e.g.*, antibodies, or antigen-binding portions thereof, selectively bind to an epitope present on a LTBP1/3-TGF β 1 complex, that is not present on a GARP-TGF β 1 complex and/or a LRRC33-TGF β 1 complex. In some embodiments, the epitope is available due to a conformational change in LTBP1/3 and/or TGF β 1 that occurs when LTBP1/3 and TGF β 1 form a complex. In this embodiment, the epitope is not present in LTBP1/3 or TGF β 1 when the

proteins are not associated in a complex. In one embodiment, the epitope is present on TGFβ1, when TGFβ1 is in a complex with LTBP1 or LTBP3. In another embodiment, the epitope is present on LTBP1, when LTBP1 is in a complex with TGFβ1. In another embodiment, the epitope is present on LTBP3, when LTBP3 is in a complex with TGFβ1. In another embodiment, the epitope comprises residues from both LTBP1 and TGFβ1. In another embodiment, the epitope comprises residues from both LTBP3 and TGFβ1.

[173] Surprisingly, some of the LTBP1/3 complex-selective antibodies disclosed herein (*e.g.*, Ab14, Ab20, Ab21-23, Ab17, and Ab24-29) are capable of binding to the small latent complex (proTGFβ1 C4S) in the absence of a presenting molecule (*e.g.*, LTBP1/3) and yet exert context-selectivity. Without wishing to be bound by theory, this finding suggests that the antibodies (and variants thereof, or cross-competing antibodies) may bind an epitope that is available in the LTBP1/3-proTGFβ1 complex and in proTGFβ1 alone, but which is not available when an LRRC-type of presenting molecule (GARP or LRRC33) is present. The epitope might be entirely on latent TGFβ1, but gets occluded (directly or indirectly) when GARP or LRRC33 is complexed.

[174] Alternatively, LTBP-selective inhibitors according to the present disclosure may bind a combinatorial epitope that comprises one or more amino acid residues of LTBP1 or LTBP3 and one or more amino acid residues of proTGFβ1, which confer the context-selectivity towards an LTBP-bound complex over GARP/LRRC33-bound complex. In these embodiments, selectivity towards the isoform (TGFβ1) as well as the context (ECM) is attributable to the combined contributions from both elements of the antigen complex.

[175] In some embodiments, the inhibitors, *e.g.*, antibodies, or antigen-binding portions thereof, are selective for the TGFβ1 isoform. In such embodiments, the inhibitors, *e.g.*, antibodies, or antigen-binding portions thereof, do not bind to TGFβ2 and/or TGFβ3. For example, in one embodiment, the inhibitors, *e.g.*, antibodies, or antigen-binding portions thereof, selectively bind a LTBP1/3-TGFβ1 complex, but do not bind TGFβ2, or a complex containing TGFβ2. In another embodiment, the inhibitors, *e.g.*, antibodies, or antigen-binding portions thereof, selectively bind a LTBP1/3-TGFβ1 complex, but do not bind TGFβ3, or a complex containing TGFβ3.

[176] In some embodiments, the inhibitors, *e.g.*, antibodies, or antigen-binding portions thereof, do not prevent TGFβ1 from binding to integrin. For example, in some embodiments, the inhibitors, *e.g.*, antibodies, or antigen-binding portions thereof, do not mask the integrin-binding site of TGFβ1.

[177] In one aspect, the invention provides functional inhibitors, *e.g.*, antibodies, that modulate TGFβ1 activity. In exemplary embodiments, the antibodies described herein are inhibitory antibodies, which inhibit the function or activity of TGFβ1. In some embodiments, the antibodies, or antigen-binding portions thereof, inhibit the activation (release) of TGFβ1 from a LTBP1-TGFβ1 complex and/or a LTBP3-TGFβ1 complex. The present disclosure provides, in exemplary embodiments,

“context-specific” or “context-selective” inhibitors of TGFβ1 activation. Such inhibitors can bind a LTBP1/3-TGFβ1 complex and inhibit activation of TGFβ1 that is presented by LTBP1 or LTBP3, without inhibiting the activation of TGFβ1 presented by GARP and/or LRRC33. Accordingly, in some embodiments, the antibodies, or antigen-binding portions thereof, described herein inhibit the release of mature TGFβ1 from a LTBP1-TGFβ1 complex and/or a LTBP3-TGFβ1 complex, but do not inhibit the release of mature TGFβ1 from a GARP-TGFβ1 complex and/or a LRRC33-TGFβ1 complex. Due to the differential localization of LTBP, GARP, and LRRC33, the context-specific inhibitors of TGFβ1 provided by the present invention can block a particular subset of TGFβ1 activity *in vivo*. In one embodiment, the context-specific antibodies provided herein that inhibit LTBP1/3-TGFβ1 but do not inhibit GARP-TGFβ1 or LRRC33-TGFβ1 can be used to inhibit TGFβ1 localized to the extracellular matrix. In another embodiment, the context-specific antibodies can inhibit TGFβ1 without modulating TGFβ1-associated immune activity or immune response. In another embodiment, the context-specific antibodies can be used to inhibit TGFβ1 activity associated with the extracellular matrix without modulating TGFβ1 activity associated with hematopoietic cells. Accordingly, the context-specific antibodies can be used to inhibit LTBP1/3-associated TGFβ1 activity in applications in which TGFβ1 activation in the context of GARP and/or LRRC33 is undesirable, as described herein.

[178] In some embodiments, the TGFβ1 comprises a naturally occurring mammalian amino acid sequence. In some embodiment, the TGFβ1 comprises a naturally occurring human amino acid sequence. In some embodiments, the TGFβ1 comprises a human, a monkey, a rat or a mouse amino acid sequence.

[179] In some embodiments, an antibody, or antigen-binding portion thereof, described herein selectively binds to a complex comprising a TGFβ1 protein comprising the amino acid sequence set forth in SEQ ID NO: 9, and LTBP1 or LTBP3. In some embodiments, an antibody, or antigen-binding portion thereof, described herein selectively binds to a LTBP1/3-TGFβ1 complex which comprises a non-naturally-occurring TGFβ1 amino acid sequence (otherwise referred to herein as a non-naturally-occurring TGFβ1). For example, a non-naturally-occurring TGFβ1 may comprise one or more recombinantly generated mutations relative to a naturally-occurring TGFβ1 amino acid sequence.

[180] In some embodiments, an antibody, or antigen-binding portion thereof, described herein does not bind TGFβ2 and/or TGFβ3, or to protein complexes containing TGFβ2 and/or TGFβ3. Exemplary TGFβ2 and TGFβ3 amino acid sequences are set forth in SEQ ID NOs: 10 and 11, respectively. In some embodiments, a TGFβ1, TGFβ2, or TGFβ3 amino acid sequence comprises an amino acid sequence as set forth in SEQ ID NOs: 12-23, as shown in Table 1. In some embodiments, a TGFβ1 amino acid sequence comprises an amino acid sequence as set forth in SEQ ID NOs: 24-31, as shown in Table 2.

[181] TGFβ1

LSTCKTIDMELVKKRIEAIKQILSKLRLASPPSQGEVPPGPLPEAVLALYNSTRDRVAGESA
 EPEPEPEADYYAKEVTRVLMVETHNEIYDKFKQSTHSIYMFFNTSELREAVPEPVLLSRAELR
 LLRLKLKVEQHVELYQKYSNNSWRYLSNRLLAPSDSPEWLSFDVTGVVRQWLSRGGEIEGF
 RLSAHCSCDSRDNTLQVDINGFTTGRRGDLATIHGMNRPFLLLMTPLERAQHLQSSRHRRALD
 TNYCFSSTEKNCCVRQLYIDFRKDLGWKWIHEPKGYHANFCLGPCPYIWSLDTQYSKVLAL
 LYNQHNP GASAAPCCVPQALEPLIVYYVGRKPKVEQLSNMIVRSCKCS (SEQ ID NO: 9)

[182] TGFβ2

SLSTCSTLMDQFMKRIEAIKQILSKLKLTSPPEDYPEPEEVPPEVISIYNSTRDLLQEKASR
 RAAACERERSDEEYYAKEVYKIDMPPFFPSENAIPPTFYRPFYRIVRFDVSAMEKNASNLVKA
 EFRVFRQLQNPKEARVPEQRIELYQILKSKDLTSPTQRYIDSKVVKTRAEGEWLSFDVTDVHE
 WLHHKDRNLGFKISLHCPCTFVPSNNYIIPNKSEELARFAGIDGTSTYTSGDQKTIKSTRKK
 NSGKTPHLLLMLLPSYRLESQQTNRKRRALDAAYCFRNVQDNCCLRPLYIDFKRDLGWKW
 IHEPKGYANFACAGAPYLWSSDTQHSRVLSLYNTINPEASASPCCVSQDLEPLTILYYIGKTP
 KIEQLSNMIVKSKCS (SEQ ID NO: 10)

[183] TGFβ3

SLSLSTCTTLDFGHIKKRVEAIKQILSKLRLTSPPEPTVMTHVPYQVLALYNSTRELLEEMH
 GEREECTQENTESEYYAKEIHKFDMIQGLAEHNELAVCPKGITSKVFRFNVSSEKNRNTLNF
 RAEFRVLRVPNPSSKRNEQRIELFQILRPDEHIAKQRYIGGKNLPTRGTAEWLSFDVTDVRE
 WLLRRESNLGLEISIHCPCHTFQPNGDILENIHEVMEIKFKGVDNEDDHGRGDLGRLKKQKD
 HHNPHLILMMIPPHRLDNPGQGGQRRKRALDTNYCFRNLEENCCVRPLYIDFRQDLGWKWV
 HEPKGYANFACGAPYLRSAADTTHSTVLGLYNTLNPEASASPCCVPQDLEPLTILYYVGRTP
 KVEQLSNMVVKSKCS (SEQ ID NO: 11)

[184] Table 1. Exemplary TGFβ1, TGFβ2, and TGFβ3 amino acid sequences

Protein	Sequence	SEQ ID NO
proTGFβ1	LSTCKTIDMELVKKRIEAIKQILSKLRLASPPSQGEVPPGPLPEAVLALYNSTRDRVAGESA EPEPEPEADYYAKEVTRVLMVETHNEIYDKFKQSTHSIYMFFNTSELREAVPEPVLLSRAELR LLRLKLKVEQHVELYQKYSNNSWRYLSNRLLAPSDSPEWLSFDVTGVVRQWLSRGGEIEGF RLSAHCSCDSRDNTLQVDINGFTTGRRGDLATIHGMNRPFLLLMTPLERAQHLQSSRHRRALD TNYCFSSTEKNCCVRQLYIDFRKDLGWKWIHEP	12

	KGYHANFCLGPCPYIWSLDTQYSKVLALYNQHNP GASAAP CCVPQALEPLPIVYYVGRKPKVEQLSNMIVRSCKCS	
proTGFβ1 C4S	LSTSKTIDMELVKRKRIEAI RQILSKLRLASPPSQGEVPPGP LPEAVLALYNSTRDRVAGESAEPEPEPEADYYAKEVTRVL MVETHNEIYDKFKQSTHSIYMFFNTSELREAVPEPVLLSRA ELRLLRLKLKVEQHVELYQKYSNNSWRYLSNRLLAPSDSP EWLSFDVTGVVRQWLSRGGEIEGFRLSAHCSCDSRDNTLQ VDINGFTTGRRGDLATIHGMNRPFLLL MATPLERAQHLQSS RHRRALDTNYCFSSTEKNCCVRQLYIDFRKDLGWKWIHEP KGYHANFCLGPCPYIWSLDTQYSKVLALYNQHNP GASAAP CCVPQALEPLPIVYYVGRKPKVEQLSNMIVRSCKCS	13
proTGFβ1 D2G	LSTCKTIDMELVKRKRIEAI RQILSKLRLASPPSQGEVPPGP LPEAVLALYNSTRDRVAGESAEPEPEPEADYYAKEVTRVL MVETHNEIYDKFKQSTHSIYMFFNTSELREAVPEPVLLSRA ELRLLRLKLKVEQHVELYQKYSNNSWRYLSNRLLAPSDSP EWLSFDVTGVVRQWLSRGGEIEGFRLSAHCSCDSRDNTLQ VDINGFTTGRRGDLATIHGMNRPFLLL MATPLERAQHLQSS RHGALDTNYCFSSTEKNCCVRQLYIDFRKDLGWKWIHEPK GYHANFCLGPCPYIWSLDTQYSKVLALYNQHNP GASAAPC CVPQALEPLPIVYYVGRKPKVEQLSNMIVRSCKCS	14
proTGFβ1 C4S D2G	LSTSKTIDMELVKRKRIEAI RQILSKLRLASPPSQGEVPPGP LPEAVLALYNSTRDRVAGESAEPEPEPEADYYAKEVTRVL MVETHNEIYDKFKQSTHSIYMFFNTSELREAVPEPVLLSRA ELRLLRLKLKVEQHVELYQKYSNNSWRYLSNRLLAPSDSP EWLSFDVTGVVRQWLSRGGEIEGFRLSAHCSCDSRDNTLQ VDINGFTTGRRGDLATIHGMNRPFLLL MATPLERAQHLQSS RHGALDTNYCFSSTEKNCCVRQLYIDFRKDLGWKWIHEPK GYHANFCLGPCPYIWSLDTQYSKVLALYNQHNP GASAAPC CVPQALEPLPIVYYVGRKPKVEQLSNMIVRSCKCS	15
proTGFβ2	SLSTCSTLDMQFMRKRIEAI RQILSKLKLTSPPEDYPEPE EVPPEVISIYNSTRDLLQE KASRRAAACERERSDEEYYAKE VYKIDMPFFPSENAIPPTFYRPFYFRIVRFDVSAMEKNASNL VKAEFRVFRLQNPKARVPEQRIEL YQILKSKDLTSPTQRYID SKVVKTRAEGEWLSFDVTD AVHEWLHHKDRNLGFKISLH CPCCTFVPSNNYIIPNKSEELE ARFAGIDGTSTYTSGDQKTIK STRKKNSGKTPHLLMLLP SYRLESQQTNRKKRALDAAY	16

	CFRNVQDNCCLRPLYIDFKRDLGWKWIHEPKGYNANFCA GACPYLWSSDTQHSRVLSLYNTINPEASASPCCVSQDLEPL TILYYIGKTPKIEQLSNMIVKSKCS	
proTGFβ2 C5S	SLSTSSTLMDMDQFMRKRIEAIHQILSKLKLTSPPEDYPEPEE VPPEVISIYNSTRDLLQEKASRRAAACERERSDEEYYAKEV YKIDMPPFFPSENAIPPTFYRPFYFRIVRFDVSAMEKNASNLV KAEFRVFRLQNPKEARVPEQRIELYQILKSKDLTSPTQRYIDS KVVKTRAEGEWLSFDVTDVHEWLHHKDRNLGFKISLHC PCCTFVPSNNYIIPNKSEELARFAGIDGTSTYTSQDQKTIKS TRKKNSGKTPHLLLMLLPSYRLESQQTNRRKKRALDAAYC FRNVQDNCCLRPLYIDFKRDLGWKWIHEPKGYNANFCA GACPYLWSSDTQHSRVLSLYNTINPEASASPCCVSQDLEPLTI LYYIGKTPKIEQLSNMIVKSKCS	17
proTGFβ2 C5S D2G	SLSTSSTLMDMDQFMRKRIEAIHQILSKLKLTSPPEDYPEPEE VPPEVISIYNSTRDLLQEKASRRAAACERERSDEEYYAKEV YKIDMPPFFPSENAIPPTFYRPFYFRIVRFDVSAMEKNASNLV KAEFRVFRLQNPKEARVPEQRIELYQILKSKDLTSPTQRYIDS KVVKTRAEGEWLSFDVTDVHEWLHHKDRNLGFKISLHC PCCTFVPSNNYIIPNKSEELARFAGIDGTSTYTSQDQKTIKS TRKKNSGKTPHLLLMLLPSYRLESQQTNRRKGALDAAYCF RNVQDNCCLRPLYIDFKRDLGWKWIHEPKGYNANFCAG CPYLWSSDTQHSRVLSLYNTINPEASASPCCVSQDLEPLTIL YYIGKTPKIEQLSNMIVKSKCS	18
proTGFβ2 D2G	SLSTCSTLMDMDQFMRKRIEAIHQILSKLKLTSPPEDYPEPE EVPPEVISIYNSTRDLLQEKASRRAAACERERSDEEYYAKE VYKIDMPPFFPSENAIPPTFYRPFYFRIVRFDVSAMEKNASNL VKAEFRVFRLQNPKEARVPEQRIELYQILKSKDLTSPTQRYID SKVVKTRAEGEWLSFDVTDVHEWLHHKDRNLGFKISLH CPCCTFVPSNNYIIPNKSEELARFAGIDGTSTYTSQDQKTIK STRKKNSGKTPHLLLMLLPSYRLESQQTNRRKGALDAAYC FRNVQDNCCLRPLYIDFKRDLGWKWIHEPKGYNANFCA GACPYLWSSDTQHSRVLSLYNTINPEASASPCCVSQDLEPLTI LYYIGKTPKIEQLSNMIVKSKCS	19
proTGFβ3	SLSLSTCTLDFGHKKRVEAIRGQILSKLRLTSPPEPTVMT HVPYQVLALYNSTRELLEEMHGEREEGCTQENTESEYYAK EIHKFDMIQGLAEHNELAVCPKGITSKVFRFNVSSVEKNRT	20

	NLFRAEFRVLRVPNPSSKRNEQRIELFQILRPDEHIAKQRYI GGKNLPTRGTAEWLSFDVTDTVREWLLRRESNLGLEISIH PCHTFQPNGDILENIHEVMEIKFKGVDNEDDHGRGDLGRL KKQKDHHNPHLILMMIPPHRLDNPQGQGGQRKKRALDTNY CFRNLEENCCVRPLYIDFRQDLGWKWWHEPKGYANFCS GPCPYLRSADTTHSTVLGLYNTLNPEASASPCCVPQDLEPL TILYYVGRTPKVEQLSNMVKSKCS	
proTGFβ3 C7S	SLSLSTSTLDFGHIKKRVEAIRGQILSKLRLTSPPEPTVMT HVPYQVLALYNSTRELLEEMHGEREEGCTQENTESEYYAK EIHKFDMIQGLAEHNELAVCPKGITSKVFRFNVSSVEKNRT NLFRAEFRVLRVPNPSSKRNEQRIELFQILRPDEHIAKQRYI GGKNLPTRGTAEWLSFDVTDTVREWLLRRESNLGLEISIH PCHTFQPNGDILENIHEVMEIKFKGVDNEDDHGRGDLGRL KKQKDHHNPHLILMMIPPHRLDNPQGQGGQRKKRALDTNY CFRNLEENCCVRPLYIDFRQDLGWKWWHEPKGYANFCS GPCPYLRSADTTHSTVLGLYNTLNPEASASPCCVPQDLEPL TILYYVGRTPKVEQLSNMVKSKCS	21
proTGFβ3 C7S D2G	SLSLSTSTLDFGHIKKRVEAIRGQILSKLRLTSPPEPTVMT HVPYQVLALYNSTRELLEEMHGEREEGCTQENTESEYYAK EIHKFDMIQGLAEHNELAVCPKGITSKVFRFNVSSVEKNRT NLFRAEFRVLRVPNPSSKRNEQRIELFQILRPDEHIAKQRYI GGKNLPTRGTAEWLSFDVTDTVREWLLRRESNLGLEISIH PCHTFQPNGDILENIHEVMEIKFKGVDNEDDHGRGDLGRL KKQKDHHNPHLILMMIPPHRLDNPQGQGGQRKGALDTNYC FRNLEENCCVRPLYIDFRQDLGWKWWHEPKGYANFCSGP CPYLRSADTTHSTVLGLYNTLNPEASASPCCVPQDLEPLTIL YYVGRTPKVEQLSNMVKSKCS	22
proTGFβ3 D2G	SLSLSTCTLDFGHIKKRVEAIRGQILSKLRLTSPPEPTVMT HVPYQVLALYNSTRELLEEMHGEREEGCTQENTESEYYAK EIHKFDMIQGLAEHNELAVCPKGITSKVFRFNVSSVEKNRT NLFRAEFRVLRVPNPSSKRNEQRIELFQILRPDEHIAKQRYI GGKNLPTRGTAEWLSFDVTDTVREWLLRRESNLGLEISIH PCHTFQPNGDILENIHEVMEIKFKGVDNEDDHGRGDLGRL KKQKDHHNPHLILMMIPPHRLDNPQGQGGQRKGALDTNYC FRNLEENCCVRPLYIDFRQDLGWKWWHEPKGYANFCSGP CPYLRSADTTHSTVLGLYNTLNPEASASPCCVPQDLEPLTIL	23

	YYVGRTPKVEQLSNMVKCKCS	
--	-----------------------	--

[185] Table 2. Exemplary non-human TGFβ1 amino acid sequences

Protein	Species	Sequence	SEQ ID NO
proTGFβ1	Mouse	LSTCKTIDMELVKKRRIEAIKQILSKLRLASPPSQGEVPP GPLPEAVLALYNSTRDRVAGESADPEPEPEADYYAKEV TRVLMVDRNNAIYEKTKDISHSIYMFNTSDIREAVPEPP LLSRAELRLQRLKSSVEQHVELYQKYSNNSWRYLGNRL LTPTDTPEWLSFDVTGVVRQWLNQGDGIQGRFSAHCS CDSKDNKLHVEINGISPKRRGDLGTIHDMMNRPFLLMAT PLERAQHLHSSRHRRALDTNYCFSSTEKNCCVRQLYIDF RKDLGWKWIHEPKGYHANFCLGPCPYIWSLDTQYSKVL ALYNQHNP GASASPCCVPQALEPLPIVYYVGRKPKVEQ LSNMIVRSCKCS	24
proTGFβ1	Cyno	LSTCKTIDMELVKKRRIEAIKQILSKLRLASPPSQGEVPP GPLPEAVLALYNSTRDRVAGESAEPEPEPEADYYAKEVT RVLMVETHNEIYDKFKQSTHSIYMFNTSELREAVPEPV LLSRAELRLRLKLVKVEQHVELYQKYSNNSWRYLSNRL LAPSDSPEWLSFDVTGVVRQWLSRGGEIEGFRLSAHCS DSKDNTLQVDINGFTTGRRGDLATIHGMNRPFLLMAT PLERAQHLQSSRHRRALDTNYCFSSTEKNCCVRQLYIDF RKDLGWKWIHEPKGYHANFCLGPCPYIWSLDTQYSKVL ALYNQHNP GASAAPCCVPQALEPLPIVYYVGRKPKVEQ LSNMIVRSCKCS	25
TGFβ1 LAP C4S	Mouse	LSTSKTIDMELVKKRRIEAIKQILSKLRLASPPSQGEVPP GPLPEAVLALYNSTRDRVAGESADPEPEPEADYYAKEV TRVLMVDRNNAIYEKTKDISHSIYMFNTSDIREAVPEPP LLSRAELRLQRLKSSVEQHVELYQKYSNNSWRYLGNRL LTPTDTPEWLSFDVTGVVRQWLNQGDGIQGRFSAHCS CDSKDNKLHVEINGISPKRRGDLGTIHDMMNRPFLLMAT PLERAQHLHSSRHRR	26
TGFβ1 LAP C4S	Cyno	LSTSKTIDMELVKKRRIEAIKQILSKLRLASPPSQGEVPP GPLPEAVLALYNSTRDRVAGESAEPEPEPEADYYAKEVT	27

		RVLMVETHNEIYDKFKQSTHSIYMFNTSELREAVPEPV LLSRAELRLRLKLKVEQHVELYQKYSNNSWRYLSNRL LAPSDSPEWLSFDVTGVVRQWLSRGGEIEGFRLSAHCSC DSKDNTLQVDINGFTTGRRGDLATIHGMNRPFLLLMAT PLERAQHLQSSRHRR	
proTGFβ1 C4S D2G	Mouse	LSTSKTIDMELVKKRIEAIKQILSKLRLASPPSQGEVPP GPLPEAVLALYNSTRDRVAGESADPEPEPEADYYAKEV TRVLMVDRNNAIYEKTKDISHSIYMFNTSDIREAVPEPP LLSRAELRLQRLKSSVEQHVELYQKYSNNSWRYLGNRL LTPTDTPEWLSFDVTGVVRQWLNQGDGIQGFRLSAHCS CDSKDNKLHVEINGISPKRRGDLGTIHD MNRPFLLLMAT PLERAQHLHSSRHGALDTNYCFSSTEKNCCVRQLYIDFR KDLGWKWIHEPKGYHANFCLGPCPYIWSLDTQYSKVL LYNQHNPGASASPCCVPQALEPLPIVYYVGRKPKVEQLS NMIVRSCKCS	28
proTGFβ1 C4S	Mouse	LSTSKTIDMELVKKRIEAIKQILSKLRLASPPSQGEVPP GPLPEAVLALYNSTRDRVAGESADPEPEPEADYYAKEV TRVLMVDRNNAIYEKTKDISHSIYMFNTSDIREAVPEPP LLSRAELRLQRLKSSVEQHVELYQKYSNNSWRYLGNRL LTPTDTPEWLSFDVTGVVRQWLNQGDGIQGFRLSAHCS CDSKDNKLHVEINGISPKRRGDLGTIHD MNRPFLLLMAT PLERAQHLHSSRHRRALDTNYCFSSTEKNCCVRQLYIDF RKDLGWKWIHEPKGYHANFCLGPCPYIWSLDTQYSKVL ALYNQHNPGASASPCCVPQALEPLPIVYYVGRKPKVEQ LSNMIVRSCKCS	29
proTGFβ1 C4S	Cyno	LSTSKTIDMELVKKRIEAIKQILSKLRLASPPSQGEVPP GPLPEAVLALYNSTRDRVAGESAEPEPEPEADYYAKEVT RVLMVETHNEIYDKFKQSTHSIYMFNTSELREAVPEPV LLSRAELRLRLKLKVEQHVELYQKYSNNSWRYLSNRL LAPSDSPEWLSFDVTGVVRQWLSRGGEIEGFRLSAHCSC DSKDNTLQVDINGFTTGRRGDLATIHGMNRPFLLLMAT PLERAQHLQSSRHRRALDTNYCFSSTEKNCCVRQLYIDF RKDLGWKWIHEPKGYHANFCLGPCPYIWSLDTQYSKVL ALYNQHNPGASAAPCCVPQALEPLPIVYYVGRKPKVEQ LSNMIVRSCKCS	30
proTGFβ1	Cyno	LSTSKTIDMELVKKRIEAIKQILSKLRLASPPSQGEVPP	31

C4S D2G		GPLPEAVLALYNSTRDRVAGESAEPEPEPEADYYAKEVT RVLMVETHNEIYDKFKQSTHSIYMFNTSELREAVPEPV LLSRAELRLLRLKLKVEQHVELYQKYSNNSWRYLSNRL LAPSDSPEWLSFDVTGVVRQWLSRGGEIEGFRLSAHCSC DSKDNTLQVDINGFTTGRRGDLATIHGMNRPFLLLMAT PLERAQHLQSSRHGALDTNYCFSSTEKNCCVRQLYIDFR KDLGWKWIHEPKGYHANFCLGPCPYIWSLDTQYSKVL LYNQHNPGLASAPCCVPQALEPLIVYYVGRKPKVEQL SNMIVRSCKCS	
---------	--	--	--

[186] In some embodiments, an antibody, or antigen-binding portion thereof, as described herein, is capable of selectively binding to an LTBP-TGF β 1 complex. In some embodiments, antigenic protein complexes (*e.g.*, a LTBP-TGF β 1 complex) may comprise an LTBP protein selected from the following: LTBP1, LTBP2, LTBP3, and LTBP4.

[187] In some embodiments, the antibody, or antigen-binding portion thereof, selectively binds an LTBP1-TGF β 1 complex. In some embodiments, the LTBP1 protein is a naturally-occurring protein. In some embodiments, the LTBP1 protein is a non-naturally occurring protein. In some embodiments, the LTBP1 protein is a recombinant protein. Such recombinant LTBP1 protein may comprise LTBP1, alternatively spliced variants thereof, and/or fragments thereof. Recombinant LTBP1 proteins may also be modified to comprise one or more detectable labels. In some embodiments, the LTBP1 protein comprises a leader sequence (*e.g.*, a native or non-native leader sequence). In some embodiments, the LTBP1 protein does not comprise a leader sequence (*i.e.*, the leader sequence has been processed or cleaved). Such detectable labels may include, but are not limited to biotin labels, polyhistidine tags, myc tags, HA tags and/or fluorescent tags. In some embodiments, the LTBP1 protein is a mammalian LTBP1 protein. In some embodiments, the LTBP1 protein is a human, a monkey, a mouse, or a rat LTBP1 protein. In some embodiments, the LTBP1 protein comprises an amino acid sequence as set forth in SEQ ID NO: 32 in Table 3. In some embodiments, the LTBP1 protein comprises an amino acid sequence as set forth in SEQ ID NOs: 33 or SEQ ID NO: 34 in Table 3.

[188] In some embodiments, an antibody, or antigen-binding portion thereof, as described herein, is capable of binding to a LTBP3-TGF β 1 complex. In some embodiments, the LTBP3 protein is a naturally-occurring protein. In some embodiments, the LTBP3 protein is a non-naturally occurring protein. In some embodiments, the LTBP3 protein is a recombinant protein. Such recombinant LTBP3 protein may comprise LTBP3, alternatively spliced variants thereof and/or fragments thereof. In some embodiments, the LTBP3 protein comprises a leader sequence (*e.g.*, a native or non-native leader sequence). In some embodiments, the LTBP3 protein does not comprise a leader sequence

(i.e., the leader sequence has been processed or cleaved). Recombinant LTBP3 proteins may also be modified to comprise one or more detectable labels. Such detectable labels may include, but are not limited to biotin labels, polyhistidine tags, myc tags, HA tags and/or fluorescent tags. In some embodiments, the LTBP3 protein is a mammalian LTBP3 protein. In some embodiments, the LTBP3 protein is a human, a monkey, a mouse, or a rat LTBP3 protein. In some embodiments, the LTBP3 protein comprises an amino acid sequence as set forth in SEQ ID NO: 35. In some embodiments, the LTBP3 protein comprises an amino acid sequence as set forth in SEQ ID NOs: 36 or 37.

[189] **Table 3. Exemplary LTBP amino acid sequences.**

Protein	Species	Sequence	SEQ ID NO
LTBP1S	Human	NHTGRIKVVFTPSICKVTCTKGSCQNSCEKGNTTTL ISENGHAADTLTATNFRVVICHLPCMNGGQCSSRD KCQCPPNFTGKLCQIPVHGASVPKLYQHSQQPGKA LGTHVIHSTHTLPLTVTSQQGVKVKFPPNIVNIHVK HPPEASVQIHQVSRIDGPTGQKTKEAQPQSQVSY QGLPVQKTQTIHSTYSHQQVIPHVYPVAAKTQLGR CFQETIGSQCGKALPGLSKQEDCCGTVGTSWGFNK CQKCPKKPSYHGYNQMMELPGYKRVNNTFCQDI NECQLQGVCPNGECLNTMGSYRCTCKIGFGPDPTF SSCVPDPVISEEKGPCYRLVSSGRQCMHPLSVHLT KQLCCCSVGKAWGPHCEKCPLPGTAAFKEICPGG MGYTVSGVHRRRPIHHHVKGKPVFVKPKNTQPVA KSTHPPPLPAKEEPVEALTFSREHGPGVAEPEVATA PPEKEIPSLDQEKTKLEPGQPQLSPGISTIHLPQFPV VIEKTSPVPVEVAPEASTSSASQVIAPTQVTEINEC TVNPDICGAGHCINLPVRYTCICYEGYRFSEQQRKC VDIDECTQVQHLC SQRCENTEGSFLCICPAGFMAS EEGTNCIDVDECLRPDVC GEGHCVNTVGAFRCEYC DSGYRMTQRGRCEDECLNPSTCPDEQCVNSPGS YQCVPCTEGFRGWNGQCLDVDECLEPNVCANGDC SNLEGSYMCSCHKGYTRTPDHKHC RDIDECQQGN LCVNGQCKNTEGSFRCTCGQGYQLSAAKDQCEDI DECQHRHLCAHGQCRNTEGSFQCVC DQGYRASGL GDHCEDINECLEDKSVCQRGDCINTAGSYDCTCPD GFQLDDNKTCQDINECEHPGLCGPQGECLNTEGSF HCVCQQGFSISADGRTCEDIDECVNNTVCD SHGFC	32

		DNTAGSFRCLCYQGFQAPQDGQGCVDVNECELLS GVCGEAFCENVEGSFLCVCADENQEYSPMTGQCR SRTSTDLDVDVDQPKEEKKECYYNLNDASLCDNV LAPNVTKQECCCTSGVGWGDNCEIFPCPVLGTAEF TEMCPKGKGFVPAGESSEAGGENYKDADECLLFG QEICKNGFCLNTRPGYECYCKQGTYYDPVKLQCFD MDECQDPSSCIDGQCVNTEGSYNCFCTHPMVLDS EKRCIRPAESNEQIEETDVYQDLCWEHLSDEYVCS RPLVGKQTTYTECCCLYGEAWGMQCALCPLKDS DYAQLCNIPVTGRRQPYGRDALVDFSEQYTPADP YFIQDRFLNSFEELQAEECGILNGCENGRCVRVQEG YTCDCFDGYHLDATAKMTCDVNECDELNNRMSL CKNAKCINTDGSYKCLCLPGYVPSDKPNYCTPLNT ALNLEKDSLE	
LTBPIS	Cyno	NHTGRIKVVFTPSICKVTCTKGSCQNSCEKGNTTTL ISENGHAADTLTATNFRVVLCHLPCMNGGQCSSRD KCQCPPNFTGKLCQIPVHGASVPKLYQHSQQPGKA LGTHVIHSTHTLPLTVTSQQGVKVKFPPNIVNIHV HPPEASVQIHQVSRIDGPTGQKTKEAQPQGSQVSY QGLPVQKTQTIHSTYSHQQVIPHVYPVAAKTQLGR CFQETIGSQCGKALPGLSKQEDCCGTGTSWGFNK CQKCPKKPSYHGYNQMMELPGYKRVNNTFCQDI NECQLQGVCNPGCECLNTMGSYRCTCKIGFGPDPTF SSCVPDPVISEEKGPCYRLVSSGRQCMHPLSVHLT KQLCCCSVGKAWGPHCEKCPLPGTAAFKEICPGG MGYTVSGVHRRRPIHHHVKGKPVFVKPKNTQPVA KSTHPPPLPAKEEPVEALTFSREHGPGVAEPEVATA PPEKEIPSLDQEKTKLEPGQPQLSPGISTIHLPQFPV VIEKTSPVPVEVAPEASTSSASQVIAPTQVTEINEC TVNPDICGAGHCINLPVRYTCICYEGYKFSEQQRKC VDIDECTQVQHLCSQGRCENTEGSFLCICPAGFMAS EEGTNCIDVDECLRPDVCGEHCVNTVGAFRCEYC DSGYRMTQRGRCEDIDECLNPSTCPDEQCVNSPGS YQCVPTGFRGWNGQCLDVDECLPNVCTNGDC SNLEGSYMCCHKGYTRTPDHKHCKDIDECQQGN LCVNGQCKNTEGSFRCTCGQGYQLSAAKDQCEDI	33

		<p> DECHHHHLCAHGQCRNTEGSFQCVCDQGYRASGL GDHCEDINECLEDKSVQQRGDCINTAGSYDCTCPD GFQLDDNKTCQDINECEHPGLCGPQGECLNTEGSF HCVCQQGFSISADGRTCEDIDECVNNTVCDSHGFC DNTAGSFRCLCYQGFQAPQDQGQCVDVNECELLS GVCGEAFCENVEGSFLCVCADENQEYSPMTGQCR SRTSTDLDVEQPKEEKKECYYNLNDASLCDNVLAP NVTKQECCCTSGAGWGDNCEIFPCPVLGTAEFTTEM CPKGGKFVPAGESSEAGGENYKDADECLLFGQEI CKNGFCLNTRPGYECYCKQGTYYDPVKLQCFDMD ECQDPSSCIDGQCVNTEGSYNCFCTHPMVLDASEK RCIRPAESNEQIEETDVYQDLCWEHLSDEYVCSRPL VGKQTTYTECCCLYGEAWGMQCALCPMKDSDDY AQLCNIPVTGRRQPYGRDALVDFSEQYAPEADPYFI QDRFLNSFEELQAEECGILNGCENGRCVRVQEGYT CDCFDGYHLDATAKMTCDVDNECDELNNRMSLCK NAKCINTEGSYKCLCLPGYVPSDKPNYCTPLNTAL NLEKDSLE </p>	
LTBPIS	mouse	<p> NHTGRIKVVFTPSICKVTCTKGNCQNSCQKGNTTT LIENGHAADTLTATNFRVVICHLPCMNGGQCSSR DKCQCPPNFTGKLCQIPVLGASMPKLYQHAQQQG KALGSHVIHSTHTLPLTMTSQQGVKVKFPPNIVNIH VKHPPEASVQIHQVSRIDSPGGQKVKEAQPGQSQV SYQGLPVQKTQTVHSTYSHQQLIPHVYPVAAKTQL GRCFQETIGSQCGKALPGLSKQEDCCGTVGTSWGF NKCQKCPKKQSYHGYTQMMECLQGYKRVNNTFC QDINECQLQGVCNNGECLNTMGSYRCCKMGFGP DPTFSSCVPDPPIVISEEKGPCYRLVSPGRHCMHPLS VHLTKQICCCSVGKAWGPHCEKCPLPGTAAFEIC PGGMGYTVSGVHRRRPIHQHIGKEAVYVKPKNTQ PVAKSTHPPPLPAKEEPVEALTSSWEHGPRGAPEV VTAPPEKEIPSLDQEKTRLEPGQPQLSPGVSTIHLHP QFPVVVEKTSPPVPVEVAPEASTSSASQVIAPTQVT EINECTVNPDICGAGHCINLPVRYTCICYEGYKFSE QLRKCDIDECAQVRHLCSQGRCENTEGSFLCVC AGFMASEEGTNCIDVDECLRPDMCRDGRGRCINTAGA </p>	34

		FRCEYCDSGYRMSRRGYCEDIDECLKPSTCPPEEQC VNTPGSYQCVPTTEGFRGWNGQCLDVDECLQPKV CTNGSCTNLEGSYMCSCHRGYSPTPDHRHCQDIDE CQQGNLCMNGQCRNTDGSFRCTCGQGYQLSAAK DQCEDIDECEHHHLCSHGQCRNTEGSFQCVCNQG YRASVLGDHCEINEDSSVCQGGDCINTAGSY DCTCPDGFQLNDNKGCDINECAQPGLCGSHGECL NTQGSFHCVCCEQGFSISADGRTCEDIDECVNNTVC DSHGFCDNAGSFRCLCYQGFQAPQDGGQGVVDVN ECELLSGVCGEAFCEVVEGSFLCVCADENQEYSPM TGQCRSRVTEDSGVDRQPREEKKECYYNLNDASL CDNVLAPNVTKQECCCTSGAGWGDNCEIFPCPVQ GTAEFTEMCPRGKGLVPAGESSYDTGGENYKDA ECLLFGEIICKNGYCLNTQPGYECYCKQGTYYDPV KLQCFDMDECQDPNSCIDGQCVNTEGSYNCFCTHP MVLDASEKRCVQPTESNEQIEETDVYQDLCWEHLS EEYVCSRPLVGKQTTYTECCCLYGEAWGMQCALC PMKDSDDYAQLCNIPVTGRRRPYGRDALVDFSEQ YGPETDPYFIQDRFLNSFEELQAECEGILNGCENGR CVRVQEGYTCDCFDGYHLDMAKMTCDVDNECSE LNNRMSLCKNAKCINTEGSYKCLCLPGYIPSDKPN YCTPLNSALNLDKESDLE	
LTBP3S	Human	ETDECRLNQNICGHGECVPGPPDYSCHCNPGYRSH PQHRYCVDVNECEAEPCGPGRGICMNTGGSYNCH CNRGYRLHVGAGGRSCVDLNECAKPHLCGDGGFC INFPGHYKCNCYPGYRLKASRPPVCEDIDECRDPS CPDGKCEKPGSFKCIACQPGYRSQGGGACRDVNE CAEGSPCSPGWCENLPGSFRCTCAQGYAPAPDGRS CLDVDECEAGDVCDNGICSNTPGSFQCQCLSGYHL SRDRSHCEDIDECDFPAACIGGDCINTNGSYRCLCP QGHRLVGGRKCQDIDECSDPSLCLPHGACKNLQ GSYVCVCDEGFTPTQDQHGCEEVEQPHHKKECYL NFDDTVFCDSVLATNVTQGECCSLGAGWGDHCE IYPCPVYSSAEFHSLCPDGKGYTQDNNIVNYGIPAH RDIDECMLFGSEICKEGKCVNTQPGYECYCKQGFY YDGNLLECVDVDECLDESNCNRNGVCENTRGGYRC	364

		ACTPPAEYSPAQRQCL	
LTBP3	Human	GPAGERGAGGGGALARERFKVVFAPVICKRTCLK GQCRDSCQQGSNMTLIGENGHSTDTLTGSGFRVVV CPLPCMNGGQCSSRNQCLCPPDFTGRFCQVPAGGA GGGTGGSGPGLSRTGALSTGALPPLAPEGDSVASK HAIYAVQVIADPPGPGEGPPAQHAAFLVPLGPGQIS AEVQAPPPVVNVRVHHPPEASVQVHRIESSNAESA APSQHLLPHPKPSHPRPPTQKPLGRCFQDTLPKQPC GSNPLPGLTKQEDCCGSIGTAWGQSKCHKCPQLQY TGVQKPGPVRGEVGADCPQGYKRLNSTHCQDINE CAMPGVCRHGDCLNPNPGSYRCVCPGHSLSGSR QCIADKPEEKSLCFRLVSPEHQCQHPLTTRLTRQLC CCSVGKAWGARCQRCPTDGTAAFKEICPAGKGYH ILTSHQTLTIQGESDFSLFLHPDGPPKPQQLPESPSQ APPPEDTEEERGVTDDSPVSEERSVQQSHPTATTTP ARPYPELISRSPPTMRWFLPDLPPSRSAVEIAPTQV TETDECRLNQNICGHGECVPGPPDYSCHCNPGYRS HPQHRYCVDVNECEAEPCGPGRGICMNTGGSYN HCNRGYRLHVGAGGRSCVDLNECAKPHLCGDGGF CINFPGHYKCNCYPGYRLKASRPPVCEDIDECDPS SCPDGKCENKPGSFKCIACQPGYRSQGGGACRDVN ECAEGSPCSPGWCENLPGSFRCTCAQGYAPAPDGR SCLDVDECEAGDVCDNGICSNTPGSFQCQCLSGYH LSRDRSHCEDIDECDFPAACIGGDCINTNGSYRCLC PQGHRLVGGKRCQDIDECSQDPSLCLPHGACKNLQ GSYVCVCDEGFTPTQDQHGCEEVEQPHHKKECYL NFDDTVFCDSVLATNVTQQECCCSLGAGWGDHCE IYPCPVYSSAEFHSLCPDGKGYTQDNNIVNYGIPAH RDIDECMFLGSEICKEGKCVNTQPGYECYCKQGFY YDGNLLECVDVDECLDESNCNRNGVCENTRGGYRC ACTPPAEYSPAQRQCLSPEEMDVDECQDPAACRPG RCVNLPGSYRCECRPPWVPGPSGRDCQLPESPAER APERRDVCWSQRGEDGMCAGPLAGPALTFDDCCC RQGRGWGAQCRPCPPRGAGSHCPTSQSESNSFWD TSPLLLGKPPRDEDSSEEDSDECRCVSGRCVPRPGG AVCECPGGFQLDASRARCVDIDECRELNQRGLLCK	35

		SERCVENTSGSFRCVCKAGFARSRPHGACVPQRRR	
LTBP3	CYNO	GPAGERGAGGGGALARERFKVVFAPVICKRTCLK GQCRDSCQQGSNMTLIGENGHSTDTLTGSGFRVVV CPLPCMNGGQCSSRNQCLCPPDFTGRFCQVPAGGA GGGTGGSGPGLSRAGALSTGALPPLAPEGDSVASK HAIYAVQVIADPPGPGEGPPAQHAAFLVPLGPGQIS AEVQAPPPVVNVRVHHPPEASVQVHRIESSNAEGA APSQHLLPHPKPSHPRPPTQKPLGRCFQDTLPKQPC GSNPLPGLTKQEDCCGSIGTAWGQSKCHKCPQLQY TGVQKPGPVRGEVGADCPQGYKRLNSTHCQDINE CAMPGVCRHGDCLNPNPSYRCVCPGHSLSGSRRT QCIADKPEEKSLCFRLVSPEHQCQHPLTTRLTRQLC CCSVGKAWGARCQRCPADGTA AFKEICPAGKGYH ILTSHQTLTIQGESDFSLFLHPDGPPKPQQLPESPSQ APPPEDTEEERGVTDDSPVSEERSVQQSHPTATTSP ARPYPELISRPSPTMRWFLPDLPPSRSAVEIAPTQV TETDECRLNQNICGHGECVPGPPDYSCHCNPGYRS HPQHRYCVDVNECEAEPCGPGRGICMNTGGSYNCR HCNRGYRLHVGAGGRSCVDLNECAKPHLCGDGGF CINFPGHYKCNCYPGYRLKASRPPVCEDIDECRDP SCPDGKCENKPGSFKCIACQPGYRSQGGGACRDVN ECAEGSPCSPGWCENLPGSFRCTCAQGYAPAPDGR SCVDVDECEAGDVCDNGICTNTPGSFQCQCLSGYH LSRDRSHCEDIDECDFPAACIGGDCINTNGSYRCLC PQGHRLVGGRCQDIDECTQDPGLCLPHGACKNL QGSYVCVCDEGFTPTQDQHGCEEVEQPHHKKECY LNFDDTVFCDSVLATNVTQQECCCSLGAGWGDHC EIYPCPVYSSAEFHSLCPDGKGYTQDNNIVNYGIPA HRDIDECMLFGAEICKEGKCVNTQPGYECYCKQGF YYDGNLLECVDVDECLDESNCNRNGVCENTRGGYR CACTPPAEYSPAQRQCLSPEEMDVDECQDPAACRP GRCVNLPGSYRCECRPPWVPGPSGRDCQLPESPAE RAPERRDVCWSQRGEDGMCAGPQAGPALTFDDCC CRQGRGWGAQCRPCPPRGAGSQCPTSQSESNSFW DTSPLLLGKPRRDEDSSEEDSDECRCVSGRCVPRPG GAVCECPGGFQLDASRARCVDIDECRELNQRGLLC	36

		KSERCVNTSGSFRCVCKAGFARSRPHGACVPQRRR	
LTBP3	Mouse	GPAGERGTGGGGALARERFKVVFAPVICKRTCLKG QCRDSCQQGSNMTLIGENGHSTDTLTGSAFRVVVC PLPCMNGGQCSSRNQCLCPPDFTGRFCQVPAAGTG AGTGSSGPGGLARTGAMSTGPLPLAPEGESVASKH AIYAVQVIADPPGPGEGPPAQHAAFLVPLGPGQISA EVQAPPPVVNVRVHHPPEASVQVHRIECPNAEGPA SSQHLLPHPKPPHPRPPTQKPLGRFCQDTLPKQPCG SNPLPGLTKQEDCCGSIGTAWGQSKCHKCPQLQYT GVQKPVPVRGEVGADCPQGYKRLNSTHCQDINEC AMPGNVCHGDCLNNPGSYRCVCPGHSGLPLAAQ CIADKPEEKSLCFRLVSTEHQCQHPLTTRLTRQLCC CSVGKAWGARCQRCPADGTAAFKEICPGKGYHILT SHQTLTIQGESDFSLFLHPDGPPKPQQLPESPSRAPP LEDTEEERGVTMDPPVSEERSVQQSHPTTTTSPPRP YPELISRPSPTFHRFLPDLPPSRSAVEIAPTQVTETD ECRLNQNICGHGQCVPGPSDYSCHCNAGYRSH PQH RYCVDVNECEAEPCGPGKGICMNTGGSYNCHCNR GYRLHVGAGGRSCVDLNECAKPHLCGDGGFCINFP GHYKCNCYPGYRLKASRPICEDIDECRDPSTCPDG KCENKPGSFKCIACQPGYRSQGGGACRDVNECSEG TPCSPGWCENLPGSYRCTCAQYEPAQDGLSCIDVD ECEAGKVCQDGICTNTPGSFQCQCLSGYHL SRDRS RCEDIDECDFPAACIGGDCINTNGSYRCLCPLGHRL VGGRKCKKDIDECSDPGLCLPHACENLQGSYVC VCDEGFTLTQDQHGCEEVEQPHHKKECYLNFFDDT VFCDSVLATNVTQQECCCSLGAGWGDHCEIYPCPV YSSAEFHS LVPD GKRLHSGQQHCELCIPAH RDIDEC ILFGAEICKEGKCVNTQPGYECYCKQGFYDGNLL ECVDVDECLDESNCRNGVCENTRGGYRC ACTPPA EYSPAQAQCLIPERWSTPQRDV KCAGASEERTACV WGPWAGPALTFDDCCCRQ PRLGTQCRPCPPRG TG SQCPTSQSESNSFWDTSPLLL GKSPREDSSEEDSD ECRCVSGRCVPRPGGAVCECPGGFQLDASRARC V DIDECRELNQRGLLCKSERCVNTSGSFRCVCKAGF TRSRPHGPACLSAAADDAIAHTSVIDHRGYFH	37

LTBP3S	Mouse	ETDECRLNQNICGHGQCVPGPSDYSCHCNAGYRSH PQHRYCVDVNECEAEPCPGKGICMNTGGSYNCH CNRGYRLHVGAGGRSCVDLNECTKPHLCGDGGFC INFPGHYKCNCYPGYRLKASRPPICEDIDECRDPST CPDGKCENKPGSFKCIACQPGYRSQGGGACRDVNE CSEGTPCSPGWCENLPGSYRCTCAQGYEPAQDGLS CIDVDECEAGKVCQDGICTNTPGSFQCQCLSGYHL SRDRSRCEDIDECDFPAACIGGDCINTNGSYRCLCP QGHRLVGGGRKCQDIDECSDPGLCLPHGACENLQ GSYVCVCDEGFTLTQDQHGCEEVEQPHHKKECYL NFDDTVFCDSVLATNVTQQECCCSLGAGWGDHCE IYPCPVYSSAEFHSLCPDGKGYTQDNNIVNYGIPAH RDIDECILFGAEICKEGKCVNTQPGYECYCKQGFY YDGNLLECVDVDECLDESNCNRNGVCENTRGGYRC ACTPPAEYSPAQRQCL	365
--------	-------	---	-----

[190] In an exemplary embodiment, inhibitors, *e.g.*, antibodies, and antigen-binding portions thereof, that selectively bind LTBP1-TGF β 1 and/or LTBP3-TGF β 1 do not bind to a complex containing TGF β 1 and GARP or LRRC33. In one embodiment, the antibodies, or antigen-binding portions thereof, do not bind a GARP protein having a sequence set forth in SEQ ID NO:38 or SEQ ID NO:39, and do not bind to a complex containing said GARP protein. In another embodiment, the inhibitors, *e.g.*, antibodies, or antigen-binding portions thereof, do not bind a GARP protein having a sequence set forth in SEQ ID NO:40 or SEQ ID NO:41, and do not bind to a complex containing said GARP protein. In one embodiment, the inhibitors, *e.g.*, antibodies, or antigen-binding portions thereof, do not bind a LRRC33 protein having a sequence set forth in SEQ ID NO:42 or SEQ ID NO:43, and do not bind a complex containing said LRRC33 protein. In one embodiment, the inhibitors, *e.g.*, antibodies, or antigen-binding portions thereof, do not bind a GARP/LRRC33 chimera, *e.g.*, the GARP/LRRC33 chimera set forth in SEQ ID NO:44.

[191] **Table 4 – Exemplary GARP and LRRC33 amino acid sequences.**

Protein	Sequence	SEQ ID NO
GARP	AQHQDKVPCKMVDKKVSCQVLGLLQVPSVLPPDTETLDLS GNQLRSILASPLGFYTALRHLDLSTNEISFLQPGAFQALTHL	38

	<p>EHLSLAHNRLAMATALSAGGLGPLPRVTSLDLSGNSLYSG LLERLLGEAPSLHTLSLAENSLTRLTRHTFRDMPALEQLDL HSNVLMDIEDGAFEGPLRLTHLNLSRNSLTCISDFSLQQLRV LDLSCNSIEAFQTASQPQAEFQLTWDLRENKLLHFPDLAA LPRLIYLNLSNNLIRLPTGPPQDSKGIHAPSEGWSALPLSAPS GNASGRPLSQLLNLDLSYNEIELIPDSFLEHLTSLCFLNLSRN CLRTFEARRLGSLPCLMLLDLSHNALETLELGARALGSLRT LLLQGNALRDLPPYTFANLASLQRLNLQGNRVSPCGGPDEP GPSGCVAFSGITSLRSLSLVDNEIELLRAGAFHTPLTELDLS SNPGLEVATGALGGLEASLEVLAQGNGLMVLQVDLPFCFI CLKRLNLAENRLSHLPAWTQAVSLEVLDLRNNSFSLLPGSA MGGLETSRRLYLQGNPLSCCGNGWLAAQLHQGRVDVDA TQDLICRFSSQEEVSLSHVRPEDCEKGGLKNINLIILTFILVS AILTTLAACCCVRRQKFNQQYKA</p>	
sGARP	<p>AQHQDKVPCKMVDKKVSCQVLGLLQVPSVLPPDTETLDLS GNQLRSILASPLGFYALRHLDLSTNEISFLQPGAFQALTHL EHLSLAHNRLAMATALSAGGLGPLPRVTSLDLSGNSLYSG LLERLLGEAPSLHTLSLAENSLTRLTRHTFRDMPALEQLDL HSNVLMDIEDGAFEGPLRLTHLNLSRNSLTCISDFSLQQLRV LDLSCNSIEAFQTASQPQAEFQLTWDLRENKLLHFPDLAA LPRLIYLNLSNNLIRLPTGPPQDSKGIHAPSEGWSALPLSAPS GNASGRPLSQLLNLDLSYNEIELIPDSFLEHLTSLCFLNLSRN CLRTFEARRLGSLPCLMLLDLSHNALETLELGARALGSLRT LLLQGNALRDLPPYTFANLASLQRLNLQGNRVSPCGGPDEP GPSGCVAFSGITSLRSLSLVDNEIELLRAGAFHTPLTELDLS SNPGLEVATGALGGLEASLEVLAQGNGLMVLQVDLPFCFI CLKRLNLAENRLSHLPAWTQAVSLEVLDLRNNSFSLLPGSA MGGLETSRRLYLQGNPLSCCGNGWLAAQLHQGRVDVDA TQDLICRFSSQEEVSLSHVRPEDCEKGGLKNIN</p>	39
GARP mouse	<p>ISQRREQVPCRTVNKEALCHGLGLLQVPSVLSLDIQALYLS GNQLQSILVSPLGFYALRHLDLSDNQISFLQAGVFQALPY LEHLNLAHNRLATGMALNSGGLGRLPLLVS LDLSGNSLHG NLVERLLGETPRLRTL SLAENSLTRLARHTFWGMPAVEQL DLHSNVLMDIEDGA FEALPHLTHLNLSRNSLTCISDFSLQQL QVLDLSCNSIEAFQTAPEPQAQFQLAWDLRENKLLHFPDL AVFPRLIYLVSNLIQLPAGLPRGSEDLHAPSEGWSASPLS</p>	40

	NPSRNASTHPLSQLLNLDLSYNEIELVPASFLEHLTSRFLN LSRNCLRSFEARQVDSLPCLVLLDLSHNVLEALELGTKVLG SLQTLLQDNALQELPPYTFASLASLQRLNLQGNQVSPCGG PAEPGPPGCVDFSGIPTLHVLMAGNSMGMLRAGSFLHTP LTELDLSTNPGLDVATGALVGLEASLEVLELQGNGLTVLR VDLPCFLRLKRLNLAENQLSHLPAWTRAVSLEVLDLRNNS FSLPGNAMGGLETSLRRLYLQGNPLSCCGNGWLAAQLHQ GRVDVDATQDLICRFGSQEELSLSLVRPEDCEKGGLKNVN LILLSSFTLVSAIVLTTLATICFLRRQKLSQQYKA	
sGARP mouse	ISQRREQVPCRTVNKEALCHGLGLLQVPSVLSLDIQALYLS GNQLQSILVSPLGFYTALRHLDLSDNQISFLQAGVFQALPY LEHLNLAHNRLATGMALNSGGLGRLPLLVSDDLSGNSLHG NLVERLLGETPRLRTLSEAENSLTRLARHTFWGMPAVEQL DLHSNVLMIEDGAFAEALPHLTHLNLSRNSLTCISDFSLQQL QVLDLSCNSIEAFQTAPEPQAQFQLAWLDLRENKLLHFPDL AVFPRLIYNVSNNLIQLPAGLPRGSEDLHAPSEGWSASPLS NPSRNASTHPLSQLLNLDLSYNEIELVPASFLEHLTSRFLN LSRNCLRSFEARQVDSLPCLVLLDLSHNVLEALELGTKVLG SLQTLLQDNALQELPPYTFASLASLQRLNLQGNQVSPCGG PAEPGPPGCVDFSGIPTLHVLMAGNSMGMLRAGSFLHTP LTELDLSTNPGLDVATGALVGLEASLEVLELQGNGLTVLR VDLPCFLRLKRLNLAENQLSHLPAWTRAVSLEVLDLRNNS FSLPGNAMGGLETSLRRLYLQGNPLSCCGNGWLAAQLHQ GRVDVDATQDLICRFGSQEELSLSLVRPEDCEKGGLKNVN	41
LRRC33 (also known as NRROS; Uniprot Accession No. Q86YC3)	MELLPLWLCGFHFLT VGWRNRSGTATAASQGVCKLVG GAADCRGQSLASVPSSLPPhARMLTLDANPLKTLWNHSLQ PYPLLESLSLHSCHLERISRGAFQEQGHLRSLVLGDNCLSEN YEETAAALHALPGLRRLDLSGNALTEDMAALMLQNSSLR SVSLAGNTIMRLDDSVFEGLERLRELDLQRNYIFEIEGGAFD GLAELRHLNLAFNNLPCIVDFGLTRLRVLNVSYNVLEWFL ATGGEAAFELETDLDSHNQLLFFPLLPQYSKLRTLLLRDNN MGFYRDLYNTSSPREMVAQFLLVDGNVTNITTVSLWEEFS SSDLADLRFLDMSQNQFQYLPDGFRLKMPSLSHLNLHQNC LMTLHIREHEPPGALTELDLSHNQLSELHLAPGLASCLGSL RLFNLSSNQLLGVPPGLFANARNITTLDMSHNQISLCPLPAA SDRVGPPSCVDFRNMASLRSLSEGCGLGALPDCPFQGTSL	42

	<p>TYLDLSSNWGVLNGLAPLQDVAPMLQVLSLRNMGLHSSF MALDFSGFGNLRDLDSLGNCLTTFPRFGGSLAETDLRRN SLTALPQKAVSEQLSRGLRTIYLSQNPYDCCGVDGWGALQ HGQTVADWAMVTCNLSSKIIRVTELPGGVPRDCKWERLDL GLLYLVILPSCLTLLVACTVIVLTFKKPLLQVIKSRCHWSS VY</p> <p>* Native signal peptide is depicted in bold font.</p>	
soluble LRRC33 (sLRRC33)	<p>MDMRVPAQLLGLLLLWFSGVLGWRNRSGTATAASQGV CKLVGGAADCRGQSLASVPSSLPPHARMLTLDANPLKTLW NHSLQPYPLLESLSLHSCHLERISRGAFQEQGHLRSLVLGD NCLSENYEETAAALHALPGLRRLDSLGNALTEDMAALML QNLSSLRSVSLAGNTIMRLDDSVFEGLERLRELDLQRNYIFE IEGGAFDGLAELRHLNLA FNNLPCIVDFGLTRLRVLNVSYN VLEWFLATGGEAAFELETDL SHNQLLFFPLLPQYSKLRTL LLRDNNMGFYRDLYNTSSPREMVAQFLLVDGNVTNITTVS LWEEFSSSDLADLRFLDMSQNQFQYLPDGF LRKMPSLSHL NLHQNCMLTLHIREHEPPGALTELD SHNQLSELHLAPGLA SCLGSLRFLNLSSNQLLGVPPGLFANARNITTLDMSHNQISL CPLPAASDRVGPPSCVDFRNMASLRSLSEGCGLGALPDCP FQGTSLTYLDLSSNWGVLNGLAPLQDVAPMLQVLSLRNM GLHSSFMALDFSGFGNLRDLDSLGNCLTTFPRFGGSLAET LDLRRNSLTALPQKAVSEQLSRGLRTIYLSQNPYDCCGVDG WGALQHGGTVADWAMVTCNLSSKIIRVTELPGGVPRDCK WERLDLGLHHHHH</p> <p>* Modified human kappa light chain signal peptide is depicted in bold font. ** Histidine tag is underlined.</p>	43
Human LRRC33- GARP chimera	<p>MDMRVPAQLLGLLLLWFSGVLGWRNRSGTATAASQGV <u>CKLVGGAADCRGQSLASVPSSLPPHARMLTLDANPLKTLW</u> <u>NHSLQPYPLLESLSLHSCHLERISRGAFQEQGHLRSLVLGD</u> <u>NCLSENYEETAAALHALPGLRRLDSLGNALTEDMAALML</u> <u>QNLSSLRSVSLAGNTIMRLDDSVFEGLERLRELDLQRNYIFE</u> <u>IEGGAFDGLAELRHLNLA FNNLPCIVDFGLTRLRVLNVSYN</u> <u>VLEWFLATGGEAAFELETDL SHNQLLFFPLLPQYSKLRTL</u> <u>LLRDNNMGFYRDLYNTSSPREMVAQFLLVDGNVTNITTVS</u> <u>LWEEFSSSDLADLRFLDMSQNQFQYLPDGF LRKMPSLSHL</u> <u>NLHQNCMLTLHIREHEPPGALTELD SHNQLSELHLAPGLA</u></p>	44

	<p> <u>SCLGSLRLFNLSNQLLGVPGLFANARNITTLDMSHNQISL</u> <u>CPLPAASDRVGPPSCVDFRNMASLRSLSLEGCGLGALPDCP</u> <u>FQGTSLTYLDLSSNWGVNLNGSLAPLQDVAPMLQVLSLRNM</u> <u>GLHSSFMALDFSGFGNLRDLDLSGNCLTTFPRFGGSLALET</u> <u>LDLRRNSLTALPQKAVSEQLSRGLRTIYLSQNPYDCCGV</u> <u>WDG</u> <u>WGALQHGQTVADWAMVTCNLSSKIIRVTELPGGVPRDCK</u> <u>WERLDLGLLIILTFILVSAILLTTLAACCCVRRQKFNQOYKA</u> </p> <p> * Modified human kappa light chain signal peptide is depicted in bold font. ** LRRC33 ectodomain is underlined. # GARP transmembrane domain is italicized. ## GARP intracellular tail is double underlined. </p>	
--	---	--

[192] In another aspect, the invention provides methods of inhibiting TGFβ1 activation in the context of LTBP1 and/or LTBP3. In one embodiment, the method comprises exposing a LTBP1-proTGFβ1 complex or a LTBP3-proTGFβ1 complex an inhibitor, an antibody or antigen-binding portion thereof, and/or a pharmaceutical composition described herein. For example, in one embodiment, the inhibitor is an inhibitor of extracellular matrix-associated TGFβ1 activation, which selectively binds a LTBP1/3-presented proTGFβ1 latent complex. In one embodiment, the inhibitor does not inhibit immune cell-associated TGFβ1 activation, for example, immune cell-associated TGFβ1 activation that results from activation of a GARP-presented proTGFβ1 latent complex. In another embodiment, the antibody, or antigen-binding portion thereof, selectively binds an LTBP1-proTGFβ1 latent complex and/or an LTBP3-proTGFβ1 latent complex, thereby modulating release of mature TGFβ1 growth factor from the latent complex, wherein the antibody, or antigen-binding portion thereof, does not bind mature TGFβ1 alone or a GARP-proTGFβ1 latent complex. In one embodiment, the antibody, or antigen-binding portion thereof, inhibits the release of mature TGFβ1 from the LTBP1-proTGFβ1 complex and/or the LTBP3-proTGFβ1 complex. In one embodiment, the antibody, or antigen-binding portion thereof, does not inhibit the release of mature TGFβ1 from a GARP-proTGFβ1 complex or a LRRC33-proTGFβ1 complex.

[193] In one embodiment, the method is performed *in vitro*. In another embodiment, the method is performed *in vivo*. In one embodiment, the LTBP1-proTGFβ1 complex or the LTBP3-proTGFβ1 complex is in an extracellular matrix. The extracellular matrix can comprise, for example, fibrillin and/or fibronectin. In some embodiments, the extracellular matrix comprises a protein comprising an RGD motif.

[194] In some embodiments of the foregoing aspects, the antibody, or antigen-binding portion thereof, does not stimulate immune effector cells. In one embodiment, the antibody, or antigen-binding portion thereof, inhibits the release of mature TGFβ1 from a LTBP1-proTGFβ1 complex

and/or a LTBP3-proTGFβ1 complex, and does not inhibit the release of mature TGFβ1 from a GARP-proTGFβ1 complex and/or an LRRC33-proTGFβ1 complex.

[195] In some embodiments, inhibitors, *e.g.*, antibodies, of the present disclosure that selectively bind to a LTBP1-TGFβ1 complex and/or a LTBP3-TGFβ1 complex can bind the complex with relatively high affinity, *e.g.*, with a dissociation constant (K_D) less than 10^{-6} M, 10^{-7} M, 10^{-8} M, 10^{-9} M, 10^{-10} M, 10^{-11} M or lower. In one embodiment, an antibody, or antigen-binding portion thereof, binds a LTBP1-TGFβ1 complex and/or a LTBP3-TGFβ1 complex with a dissociation constant (K_D) of about 10^{-8} M, about 10^{-9} M, about 10^{-10} M, about 10^{-11} M, about 10^{-12} M, or about 10^{-13} M. For example, antibodies that selectively bind to a LTBP1-TGFβ1 complex and/or a LTBP3-TGFβ1 complex can bind the complex with an affinity between 5 pM and 500 nM, *e.g.*, between 10 pM and 100 nM, *e.g.*, between 50 pM and 50 nM. In one embodiment, the antibody, or antigen-binding fragment thereof, can bind a LTBP1-TGFβ1 complex and/or a LTBP3-TGFβ1 complex with an affinity of less than about 300 nM, for example about 20 nM or lower, about 10 nM or lower, about 500 pM or lower, or about 5 pM or lower. For example, the antibody, or antigen-binding fragment thereof, can bind a LTBP1-TGFβ1 complex and/or a LTBP3-TGFβ1 complex with an affinity of about 1 nm to about 350 nm, from about 10 nm to about 200 nm, from about 15 nm to about 250 nm, from about 20 nm to about 200 nm, about 1 nm, about 20 nm, about 25 nm, about 50 nm, about 100 nm, about 150 nm, about 200 nm, about 250 nm, about 300 nm, or about 500 nm.

[196] The disclosure also includes antibodies or antigen-binding fragments that compete with any of the antibodies described herein for binding to a LTBP1-TGFβ1 complex and/or a LTBP3-TGFβ1 complex. In some embodiments, such antibodies have an affinity for the complex of 50 nM or lower (*e.g.*, 20 nM or lower, 10 nM or lower, 500 pM or lower, 50 pM or lower, or 5 pM or lower). The affinity and binding kinetics of antibodies (or antigen-binding fragments thereof) that selectively bind to a LTBP1-TGFβ1 complex and/or a LTBP3-TGFβ1 complex can be tested using any suitable method, including but not limited to biosensor technology (*e.g.*, OCTET or BIACORE).

[197] In one embodiment, the antibodies, or antigen-binding fragments thereof, of the present disclosure do not compete with antibody SR-Ab1 for binding to a human LTBP1-proTGFβ1 complex.

[198] Aspects of the disclosure relate to antibodies that compete or cross-compete with any of the antibodies provided herein. The term “compete”, as used herein with regard to an antibody, means that a first antibody binds to an epitope (*e.g.*, an epitope of a LTBP1-TGFβ1 complex and/or an epitope of a LTBP3-TGFβ1 complex) in a manner sufficiently similar to the binding of a second antibody, such that the result of binding of the first antibody with its epitope is detectably decreased in the presence of the second antibody compared to the binding of the first antibody in the absence of the second antibody. The alternative, where the binding of the second antibody to its epitope is also detectably decreased in the presence of the first antibody, can, but need not be the case. That is, a first

antibody can inhibit the binding of a second antibody to its epitope without that second antibody inhibiting the binding of the first antibody to its respective epitope. However, where each antibody detectably inhibits the binding of the other antibody with its epitope or ligand, whether to the same, greater, or lesser extent, the antibodies are said to “cross-compete” with each other for binding of their respective epitope(s). Both competing and cross-competing antibodies are within the scope of this disclosure. Regardless of the mechanism by which such competition or cross-competition occurs (*e.g.*, steric hindrance, conformational change, or binding to a common epitope, or portion thereof), the skilled artisan would appreciate that such competing and/or cross-competing antibodies are encompassed and can be useful for the methods and/or compositions provided herein.

[199] Aspects of the disclosure relate to antibodies that compete or cross-compete with any of the specific antibodies, or antigen-binding portions thereof, as provided herein, *e.g.*, an antibody having one or more CDR sequences (1, 2, 3, 4, 5, or 6 CDR sequences) set forth in Table 5. In one embodiment, the invention provides antibodies, and antigen-binding fragments thereof, that compete or cross-compete with an antibody having heavy chain CDR sequences comprising SEQ ID NO:1, SEQ ID NO:2, and SEQ ID NO:3 as set forth in Table 5, and/or light chain CDR sequences comprising SEQ ID NO:4, SEQ ID NO:5, and SEQ ID NO:6 as set forth in Table 5. In one embodiment, the invention provides antibodies that compete or cross-compete with an antibody, or antigen-binding portion thereof, having a heavy chain variable region sequence comprising SEQ ID NO:7, and/or a light chain variable region sequence comprising SEQ ID NO:8. In some embodiments, an antibody, or antigen-binding portion thereof, binds at or near the same epitope as any of the antibodies provided herein. In some embodiments, an antibody, or antigen-binding portion thereof, binds near an epitope if it binds within 15 or fewer amino acid residues of the epitope. In some embodiments, any of the antibody, or antigen-binding portion thereof, as provided herein, binds within 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14 or 15 amino acid residues of an epitope that is bound by any of the antibodies provided herein.

[200] In another embodiment, provided herein is an antibody, or antigen-binding portion thereof, competes or cross-competes for binding to any of the antigens provided herein (*e.g.*, a LTBP1-TGFβ1 complex and/or a LTBP3-TGFβ1 complex) with an equilibrium dissociation constant, K_D , between the antibody and the protein of less than 10^{-6} M. In other embodiments, an antibody competes or cross-competes for binding to any of the antigens provided herein with a K_D in a range from 10^{-11} M to 10^{-6} M. In other embodiments, an antibody competes or cross-competes for binding to a human LTBP1-TGFβ1 complex and/or a human LTBP3-TGFβ1 complex with a K_D of < 50 nM as determined by a suitable in vitro binding assay, *e.g.*, BLI, such as Octet®. In other embodiments, an antibody competes or cross-competes for binding to a human LTBP1-TGFβ1 complex and/or a human LTBP3-TGFβ1 complex with a K_D of < 10 nM as determined by a suitable in vitro binding assay, *e.g.*, BLI, such as Octet.

[201] In some embodiments, the antibody or antigen-binding portion competes or cross-competes with an antibody having a heavy chain variable region sequence and light chain variable region sequence of Ab42, as set forth in Table 6 (*e.g.*, SEQ ID NOs: 318 and 319, respectively). The antibody may compete or cross-compete for binding to a human LTBP1-TGF β 1 complex and/or a human LTBP3-TGF β 1 complex with a K_D of < 10 nM as determined by a suitable *in vitro* binding assay, *e.g.*, BLI, such as Octet. The antibody may compete or cross-compete for binding to a human LTBP1-TGF β 1 complex and/or a human LTBP3-TGF β 1 complex with a K_D of < 5 nM as determined by a suitable *in vitro* binding assay, *e.g.*, BLI, such as Octet. The antibody may bind to a human LTBP1-TGF β 1 complex and a human LTBP3-TGF β 1 complex with a K_D of < 5 nM as determined by a suitable *in vitro* binding assay, *e.g.*, BLI, such as Octet. The antibody may not show any detectable binding to a human GARP-proTGF β 1 complex in a suitable *in vitro* binding assay, such as BLI (*e.g.*, Octet). The antibody may not show detectable binding to a human GARP-proTGF β 1 complex, as measured by BLI, under the same assay conditions as used to measure binding to human LTBP1-proTGF β 1 complex and/or human LTBP3-TGF β 1 complex. Alternatively, or in addition, the antibody or antigen-binding portion may bind (*e.g.*, selectively bind) a human LTBP1-proTGF β 1 complex and/or a human LTBP3-TGF β 1 complex with a K_D that is at least 50 times lower than the K_D when binding to a human GARP-proTGF β 1 complex (and optionally at least 50 times lower than the K_D when binding to a human LRRC33-proTGF β 1 complex) under the same assay conditions.

[202] In some embodiments, the antibody competes or cross-competes with an antibody having a heavy chain variable region sequence and light chain variable region sequence of Ab63, as set forth in Table 6 (*e.g.*, SEQ ID NOs: 360 and 361, respectively). The antibody may compete or cross-compete for binding to a human LTBP1-TGF β 1 complex and/or a human LTBP3-TGF β 1 complex with a K_D of < 10 nM as determined by a suitable *in vitro* binding assay, *e.g.*, BLI, such as Octet. The antibody may compete or cross-compete for binding to a human LTBP1-TGF β 1 complex and/or a human LTBP3-TGF β 1 complex with a K_D of < 5 nM as determined by a suitable *in vitro* binding assay, *e.g.*, BLI, such as Octet. The antibody may bind to a human LTBP1-TGF β 1 complex and a human LTBP3-TGF β 1 complex with a K_D of < 5 nM as determined by a suitable *in vitro* binding assay, *e.g.*, BLI, such as Octet. The antibody may not show any detectable binding to a human GARP-proTGF β 1 complex in a suitable *in vitro* binding assay, such as BLI (*e.g.*, Octet). The antibody may not show detectable binding to a human GARP-proTGF β 1 complex, as measured by BLI, under the same assay conditions as used to measure binding to human LTBP1-proTGF β 1 complex and/or human LTBP3-TGF β 1 complex. Alternatively, or in addition, the antibody or antigen-binding portion may bind (*e.g.*, selectively bind) a human LTBP1-proTGF β 1 complex and/or a human LTBP3-TGF β 1 complex with a K_D that is at least 50 times lower than the K_D when binding to a human GARP-proTGF β 1 complex (and optionally at least 50 times lower than the K_D when binding to a human LRRC33-proTGF β 1 complex) under the same assay conditions.

[203] In further embodiments, the antibody which selectively binds a human LTBP1-TGFβ1 complex and/or a human LTBP3-TGFβ1 complex may not show meaningful binding (*e.g.*, may not show a response of more than 0.1 units (nm)) on exposure to a human GARP-proTGFβ1 complex in a BLI assay (*e.g.*, Octet) when the human GARP-proTGFβ1 complex is at a concentration of 200nM.

[204] In some embodiments, provided herein is an anti-TGFβ1 antibody, or antigen-binding portion thereof, that competes for binding with an antibody, or antigen-binding portion thereof, described herein. In some embodiments, provided herein is an anti-TGFβ1 antibody, or antigen-binding portion thereof, that binds to the same epitope as an antibody, or antigen-binding portion thereof, described herein.

[205] The antibodies provided herein can be characterized using any suitable methods. For example, one method is to identify the epitope to which the antigen binds, or “epitope mapping.” There are many suitable methods for mapping and characterizing the location of epitopes on proteins, including solving the crystal structure of an antibody-antigen complex, competition assays, gene fragment expression assays, and synthetic peptide-based assays, as described, for example, in Chapter 11 of Harlow and Lane, *Using Antibodies, a Laboratory Manual*, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., 1999. In an additional example, epitope mapping can be used to determine the sequence to which an antibody binds. The epitope can be a linear epitope, *i.e.*, contained in a single stretch of amino acids, or a conformational epitope formed by a three-dimensional interaction of amino acids that may not necessarily be contained in a single stretch (primary structure linear sequence). In some embodiments, the epitope is a TGFβ1 epitope that is only available for binding by the antibody, or antigen-binding portion thereof, described herein, when the TGFβ1 is in a LTBP1-TGFβ1 complex and/or a LTBP3-TGFβ1 complex. In some embodiments, the epitope is present on a LTBP1/3-TGFβ1 complex, and is not present on a GARP-TGFβ1 complex and/or a LRRC33-TGFβ1 complex. In some embodiments, the epitope is available due to a conformational change in LTBP1/3 and/or TGFβ1 that occurs when LTBP1/3 and TGFβ1 form a complex. In this embodiment, the epitope is not present in LTBP1/3 or TGFβ1 when the proteins are not associated in a complex. In one embodiment, the epitope is present on TGFβ1, when TGFβ1 is in a complex with LTBP1 or LTBP3. In another embodiment, the epitope is present on LTBP1, when LTBP1 is in a complex with TGFβ1. In another embodiment, the epitope is present on LTBP3, when LTBP3 is in a complex with TGFβ1. In another embodiment, the epitope comprises residues from both LTBP1 and TGFβ1. In another embodiment, the epitope comprises residues from both LTBP3 and TGFβ1. Peptides of varying lengths (*e.g.*, at least 4-6 amino acids long) can be isolated or synthesized (*e.g.*, recombinantly) and used for binding assays with an antibody. In another example, the epitope to which the antibody binds can be determined in a systematic screen by using overlapping peptides derived from the target antigen sequence and determining binding by the antibody. According to the gene fragment expression assays, the open reading frame encoding the

target antigen is fragmented either randomly or by specific genetic constructions and the reactivity of the expressed fragments of the antigen with the antibody to be tested is determined. The gene fragments may, for example, be produced by PCR and then transcribed and translated into protein *in vitro*, in the presence of radioactive amino acids. The binding of the antibody to the radioactively labeled antigen fragments is then determined by immunoprecipitation and gel electrophoresis. Certain epitopes can also be identified by using large libraries of random peptide sequences displayed on the surface of phage particles (phage libraries). Alternatively, a defined library of overlapping peptide fragments can be tested for binding to the test antibody in simple binding assays. In an additional example, mutagenesis of an antigen-binding domain, domain swapping experiments and alanine scanning mutagenesis can be performed to identify residues required, sufficient, and/or necessary for epitope binding. For example, domain swapping experiments can be performed using a mutant of a target antigen in which various fragments of the LTBP1-TGF β 1 complex or LTBP3-TGF β 1 complex have been replaced (swapped) with sequences from a closely related, but antigenically distinct protein, such as another member of the TGF β protein family (*e.g.*, GDF11). By assessing binding of the antibody to the mutant of the LTBP1-TGF β 1 complex and/or LTBP3-TGF β 1 complex, the importance of the particular antigen fragment to antibody binding can be assessed.

[206] Alternatively, competition assays can be performed using other antibodies known to bind to the same antigen to determine whether an antibody binds to the same epitope as the other antibodies. Competition assays are well known to those of skill in the art.

[207] Further, the interaction of the any of the antibodies provided herein with one or more residues in a LTBP1-TGF β 1 complex and/or a LTBP3-TGF β 1 complex can be determined by routine technology. For example, a crystal structure can be determined, and the distances between the residues in a LTBP1-TGF β 1 complex and/or a LTBP3-TGF β 1 complex, and one or more residues in the antibody, can be determined accordingly. Based on such distance, whether a specific residue in a LTBP1/3-TGF β 1 complex interacts with one or more residues in the antibody can be determined. Further, suitable methods, such as competition assays and target mutagenesis assays, can be applied to determine the preferential binding of a candidate antibody.

[208] In some embodiments, the antibodies, or antigen-binding portions thereof, of the present invention that selectively bind to a LTBP1-TGF β 1 complex and/or a LTBP3-TGF β 1 complex include one or more of complementary determining regions (CDRs) shown in Table 5. In some embodiments, the invention provides a nucleic acid molecule that encodes an antibody, or antigen-binding portion thereof, that selectively binds to a LTBP1-TGF β 1 complex and/or a LTBP3-TGF β 1 complex, as described herein. In one embodiment, the nucleic acid molecules encode one or more of the CDR sequences shown in Table 5.

[209] Table 5. Complementary determining regions of the heavy chain (CDRHs) and the light chain (CDRLs) of SR-AB2, SR-AB10, SR-AB13, SR-AB22, SR-AB23, SR-AB31, SR-AB34, SR-AB37, and SR-AB38 to SR-AB64 as determined using the Kabat numbering scheme.

Antibody	SR-AB2
CDRH1	GYTFTSYG (SEQ ID NO: 1)
CDRH2	ISAYNGNT (SEQ ID NO: 2)
CDRH3	ARAPLGNFDS (SEQ ID NO: 3)
CDRL1	SGSIASNY (SEQ ID NO: 4)
CDRL2	EDN (SEQ ID NO: 5)
CDRL3	QSYDSSNHPVV (SEQ ID NO: 6)
Antibody	SR-AB10
CDRH1	FTFNYPPIH (SEQ ID NO: 94)
CDRH2	VMSYDGINKYYADSVKG (SEQ ID NO: 95)
CDRH3	ARPRIAARRGGFDY (SEQ ID NO: 96)
CDRL1	TRSSGNIDNNYVQ (SEQ ID NO: 97)
CDRL2	EDNQRPS (SEQ ID NO: 98)
CDRL3	QSYDSNQGTVV (SEQ ID NO: 99)
Antibody	SR-AB13
CDRH1	GSISSSYYWG (SEQ ID NO: 100)
CDRH2	SISYSGSTYY (SEQ ID NO: 101)
CDRH3	ARDPSYDSIAGMDV (SEQ ID NO: 102)
CDRL1	RASQSISSYLN (SEQ ID NO: 103)
CDRL2	AASNLS (SEQ ID NO: 104)
CDRL3	QQSFDFPFT (SEQ ID NO: 105)

Antibody	SR-AB22
CDRH1	FTFRSYVMH (SEQ ID NO: 108)
CDRH2	VISHEGSLKYYADSVKG (SEQ ID NO: 109)
CDRH3	AVPRIAARRGGFGY (SEQ ID NO: 110)
CDRL1	TRSSGNIDNNYVQ (SEQ ID NO: 111)
CDRL2	EDNQRPS (SEQ ID NO: 112)
CDRL3	QSYDSNQGCVV (SEQ ID NO: 113)
Antibody	SR-AB23
CDRH1	FTFRSYVMH (SEQ ID NO: 116)
CDRH2	VISHEGSLKYYADSVKG (SEQ ID NO: 117)
CDRH3	ARPRIAARRGGFGY (SEQ ID NO: 118)
CDRL1	TRSSGNIDNNYVQ (SEQ ID NO: 119)
CDRL2	EDNQRPS (SEQ ID NO: 120)
CDRL3	QSYDSNQGCVV (SEQ ID NO: 121)
Antibody	SR-AB31
CDRH1	FTFRSYVMH (SEQ ID NO: 124)
CDRH2	VISHEGSLKYYADSVKG (SEQ ID NO: 125)
CDRH3	AVPRIAARRGGFGY (SEQ ID NO: 126)
CDRL1	TRSSGNIDNNYVQ (SEQ ID NO: 127)
CDRL2	EDNQRPS (SEQ ID NO: 128)
CDRL3	QSYDFNNQGCVV (SEQ ID NO: 129)
Antibody	SR-AB34

CDRH1	FTFRSYVMH (SEQ ID NO: 130)
CDRH2	VISHEGSLKYYADSVKG (SEQ ID NO: 131)
CDRH3	AVPRIAARRGGFGY (SEQ ID NO: 132)
CDRL1	TRSSGNIDNNYVQ (SEQ ID NO: 133)
CDRL2	EDNQRPS (SEQ ID NO: 134)
CDRL3	QSYDYDAQGVV (SEQ ID NO: 135)
Antibody	SR-AB37
CDRH1	FTFRSYVMH (SEQ ID NO: 136)
CDRH2	VISHEGSLKYYADSVKG (SEQ ID NO: 137)
CDRH3	AVPRIAARRGGFGY (SEQ ID NO: 138)
CDRL1	TRSSGLIDNNYVQ (SEQ ID NO: 139)
CDRL2	EDNQRPS (SEQ ID NO: 140)
CDRL3	QSYDSDLQRVV (SEQ ID NO: 141)
Antibody	SR-AB38
CDRH1	FTFRSYVMH (SEQ ID NO: 142)
CDRH2	VISHEGSLKYYADSVKG (SEQ ID NO: 143)
CDRH3	AVPRIAARRGGFGY (SEQ ID NO: 144)
CDRL1	TRSSGSIDNNYVQ (SEQ ID NO: 145)
CDRL2	EDFIRPS (SEQ ID NO: 146)
CDRL3	QSYDDDLQGVV (SEQ ID NO: 147)
Antibody	SR-AB39
CDRH1	FTFRSYVMH (SEQ ID NO: 148)

CDRH2	VISHEGSLKYYADSVKG (SEQ ID NO: 149)
CDRH3	AVPRIAARRGGFGY (SEQ ID NO: 150)
CDRL1	TRSSGLIDDNYVQ (SEQ ID NO: 151)
CDRL2	EDAQRPS (SEQ ID NO: 152)
CDRL3	QSYDHDEQGVV (SEQ ID NO: 153)
Antibody	SR-AB40
CDRH1	FTFRSYVMH (SEQ ID NO: 154)
CDRH2	VISHEGSLKYYADSVKG (SEQ ID NO: 155)
CDRH3	ARPRIAARRGGFGY (SEQ ID NO: 156)
CDRL1	TRSSGNIDNNYVQ (SEQ ID NO: 157)
CDRL2	EDNQRPS (SEQ ID NO: 158)
CDRL3	QSYDYDYNQGVV (SEQ ID NO: 159)
Antibody	SR-AB41
CDRH1	FTFRSYVMH (SEQ ID NO: 160)
CDRH2	VISHEGSLKYYADSVKG (SEQ ID NO: 161)
CDRH3	ARPRIAARRGGFGY (SEQ ID NO: 162)
CDRL1	TRSSGNIDNNYVQ (SEQ ID NO: 163)
CDRL2	EDNQRPS (SEQ ID NO: 164)
CDRL3	QSYDYDNQAVV (SEQ ID NO: 165)
Antibody	SR-AB42
CDRH1	FTFRSYVMH (SEQ ID NO: 166)
CDRH2	VISHEGSLKYYADSVKG (SEQ ID NO: 167)

CDRH3	ARPRIAARRGGFGY (SEQ ID NO: 168)
CDRL1	TRSSGNIDNNYVQ (SEQ ID NO: 169)
CDRL2	EDNQRPS (SEQ ID NO: 170)
CDRL3	QSYDYDTQGVV (SEQ ID NO: 171)
Antibody	SR-AB43
CDRH1	FTFRSYVMH (SEQ ID NO: 172)
CDRH2	VISHEGSLKYYADSVKG (SEQ ID NO: 173)
CDRH3	ARPRIAARRGGFGY (SEQ ID NO: 174)
CDRL1	TRSSGNIDYNYVQ (SEQ ID NO: 175)
CDRL2	EDNVRPS (SEQ ID NO: 176)
CDRL3	QSYDSDNQRVV (SEQ ID NO: 177)
Antibody	SR-AB44
CDRH1	FTFRSYVMH (SEQ ID NO: 178)
CDRH2	VISHEGSLKYYADSVKG (SEQ ID NO: 179)
CDRH3	ARPRIAARRGGFGY (SEQ ID NO: 180)
CDRL1	TRSHGNIDDNYVQ (SEQ ID NO: 181)
CDRL2	EDNVRPS (SEQ ID NO: 182)
CDRL3	QSYDSDNQLVV (SEQ ID NO: 183)
Antibody	SR-AB45
CDRH1	FTFRSYVMH (SEQ ID NO: 184)
CDRH2	VISHEGSLKYYADSVKG (SEQ ID NO: 185)
CDRH3	ARPRIAARRGGFGY (SEQ ID NO: 186)

CDRL1	TRSSGAIDNYVQ (SEQ ID NO: 187)
CDRL2	EDFQRPS (SEQ ID NO: 188)
CDRL3	QSYDDDLQGVV (SEQ ID NO: 189)
Antibody	SR-AB46
CDRH1	FTFRSYVMH (SEQ ID NO: 190)
CDRH2	VISHEGSGKYYADSVKG (SEQ ID NO: 191)
CDRH3	ARPRIAARRGGFGS (SEQ ID NO: 192)
CDRL1	TRSSGNIDYNYVQ (SEQ ID NO: 193)
CDRL2	EDNVRPS (SEQ ID NO: 194)
CDRL3	QSYDSNQRVV (SEQ ID NO: 195)
Antibody	SR-AB47
CDRH1	FTFRSYVMH (SEQ ID NO: 196)
CDRH2	VISHEGSGKYYADSVKG (SEQ ID NO: 197)
CDRH3	ARPRIAARRGGFGS (SEQ ID NO: 198)
CDRL1	TRSSGNIDYNYVQ (SEQ ID NO: 199)
CDRL2	EDNVRPS (SEQ ID NO: 200)
CDRL3	QSYDYDNQAVV (SEQ ID NO: 201)
Antibody	SR-AB48
CDRH1	FTFRSYVMH (SEQ ID NO: 202)
CDRH2	VISHEGSGKYYADSVKG (SEQ ID NO: 203)
CDRH3	ARPRIAARRGGFGS (SEQ ID NO: 204)
CDRL1	TRSSGNIDYNYVQ (SEQ ID NO: 205)

CDRL2	EDNVRPS (SEQ ID NO: 206)
CDRL3	QSYDYDTQGVV (SEQ ID NO: 207)
Antibody	SR-AB49
CDRH1	FTFRSYVMH (SEQ ID NO: 208)
CDRH2	VISHEGSGKYYADSVKG (SEQ ID NO: 209)
CDRH3	ARPRIAARRGGFGS (SEQ ID NO: 210)
CDRL1	TRSSGNIDYNYVQ (SEQ ID NO: 211)
CDRL2	EDNVRPS (SEQ ID NO: 212)
CDRL3	QGYDWDTQGVV (SEQ ID NO: 213)
Antibody	SR-AB50
CDRH1	FTFRSYVMH (SEQ ID NO: 214)
CDRH2	VISHEGSGKYYADSVKG (SEQ ID NO: 215)
CDRH3	ARPRIAARRGGFGT (SEQ ID NO: 216)
CDRL1	TRSSGNIDYNYVQ (SEQ ID NO: 217)
CDRL2	EDNVRPS (SEQ ID NO: 218)
CDRL3	QSYDSNQRVV (SEQ ID NO: 219)
Antibody	SR-AB51
CDRH1	FTFRSYVMH (SEQ ID NO: 220)
CDRH2	VISHEGSGKYYADSVKG (SEQ ID NO: 221)
CDRH3	ARPRIAARRGGFGT (SEQ ID NO: 222)
CDRL1	TRSSGNIDYNYVQ (SEQ ID NO: 223)
CDRL2	EDNVRPS (SEQ ID NO: 224)

CDRL3	QSYDYDNQAVV (SEQ ID NO: 225)
Antibody	SR-AB52
CDRH1	FTFRSYVMH (SEQ ID NO: 226)
CDRH2	VISHEGSGKYYADSVKG (SEQ ID NO: 227)
CDRH3	ARPRIAARRGGFGT (SEQ ID NO: 228)
CDRL1	TRSSGNIDYNYVQ (SEQ ID NO: 229)
CDRL2	EDNVRPS (SEQ ID NO: 230)
CDRL3	QSYDYDTQGVV (SEQ ID NO: 2312)
Antibody	SR-AB53
CDRH1	FTFRSYVMH (SEQ ID NO: 232)
CDRH2	VISHEGSGKYYADSVKG (SEQ ID NO: 233)
CDRH3	ARPRIAARRGGFGT (SEQ ID NO: 234)
CDRL1	TRSSGNIDYNYVQ (SEQ ID NO: 235)
CDRL2	EDNVRPS (SEQ ID NO: 236)
CDRL3	QGYDWDYQGVV (SEQ ID NO: 237)
Antibody	SR-AB54
CDRH1	FTFRSYVMH (SEQ ID NO: 238)
CDRH2	VISHEGSGKYYADSVKG (SEQ ID NO: 239)
CDRH3	ALPRIAARRGGFGS (SEQ ID NO: 240)
CDRL1	TRSSGNIDYNYVQ (SEQ ID NO: 241)
CDRL2	EDNVRPS (SEQ ID NO: 242)
CDRL3	QSYDSNQRVV (SEQ ID NO: 243)

Antibody	SR-AB55
CDRH1	FTFRSYVMH (SEQ ID NO: 244)
CDRH2	VISHEGSGKYYADSVKG (SEQ ID NO: 245)
CDRH3	ALPRIAARRGGFGS (SEQ ID NO: 246)
CDRL1	TRSSGNIDYNYVQ (SEQ ID NO: 247)
CDRL2	EDNVRPS (SEQ ID NO: 2489)
CDRL3	QSYDYDNQAVV (SEQ ID NO: 249)
Antibody	SR-AB56
CDRH1	FTFRSYVMH (SEQ ID NO: 250)
CDRH2	VISHEGSGKYYADSVKG (SEQ ID NO: 251)
CDRH3	ALPRIAARRGGFGS (SEQ ID NO: 252)
CDRL1	TRSSGNIDYNYVQ (SEQ ID NO: 253)
CDRL2	EDNVRPS (SEQ ID NO: 254)
CDRL3	QSYDYDTQGVV (SEQ ID NO: 255)
Antibody	SR-AB57
CDRH1	FTFRSYVMH (SEQ ID NO: 256)
CDRH2	VISHEGSGKYYADSVKG (SEQ ID NO: 257)
CDRH3	ALPRIAARRGGFGS (SEQ ID NO: 258)
CDRL1	TRSSGNIDYNYVQ (SEQ ID NO: 259)
CDRL2	EDNVRPS (SEQ ID NO: 260)
CDRL3	QGYDWDTQGVV (SEQ ID NO: 261)
Antibody	SR-AB58

CDRH1	FTFRSYVMH (SEQ ID NO: 262)
CDRH2	VISHEGSGKYYADSVKG (SEQ ID NO: 263)
CDRH3	ALPRIAARRGGFGT (SEQ ID NO: 264)
CDRL1	TRSSGNIDYNYVQ (SEQ ID NO: 265)
CDRL2	EDNVRPS (SEQ ID NO: 266)
CDRL3	QSYDSNQRVV (SEQ ID NO: 267)
Antibody	SR-AB59
CDRH1	FTFRSYVMH (SEQ ID NO: 268)
CDRH2	VISHEGSGKYYADSVKG (SEQ ID NO: 269)
CDRH3	ALPRIAARRGGFGT (SEQ ID NO: 270)
CDRL1	TRSSGNIDYNYVQ (SEQ ID NO: 271)
CDRL2	EDNVRPS (SEQ ID NO: 272)
CDRL3	QSYDYDNQAVV (SEQ ID NO: 273)
Antibody	SR-AB60
CDRH1	FTFRSYVMH (SEQ ID NO: 274)
CDRH2	VISHEGSGKYYADSVKG (SEQ ID NO: 275)
CDRH3	ALPRIAARRGGFGT (SEQ ID NO: 276)
CDRL1	TRSSGNIDYNYVQ (SEQ ID NO: 277)
CDRL2	EDNVRPS (SEQ ID NO: 278)
CDRL3	QSYDYDTQGVV (SEQ ID NO: 279)
Antibody	SR-AB61
CDRH1	FTFRSYVMH (SEQ ID NO: 280)

CDRH2	VISHEGSGKYYADSVKG (SEQ ID NO: 281)
CDRH3	ALPRIAARRGGFGT (SEQ ID NO: 282)
CDRL1	TRSSGNIDYNYVQ (SEQ ID NO: 283)
CDRL2	EDNVRPS (SEQ ID NO: 284)
CDRL3	QGYDWDQTQGVV (SEQ ID NO: 285)
Antibody	SR-AB62
CDRH1	GSIRSSSYWG (SEQ ID NO: 286)
CDRH2	SISYSATYY (SEQ ID NO: 287)
CDRH3	ASDPSYDSAAGMDV (SEQ ID NO: 288)
CDRL1	RASKVISSYLN (SEQ ID NO: 289)
CDRL2	YASSLQS (SEQ ID NO: 290)
CDRL3	QQSNDWPFT (SEQ ID NO: 291)
Antibody	SR-AB63
CDRH1	GSIRSSSYWG (SEQ ID NO: 292)
CDRH2	SISYSATYY (SEQ ID NO: 293)
CDRH3	AGDPSYDSIAGMQV (SEQ ID NO: 294)
CDRL1	RASQSISSYLN (SEQ ID NO: 295)
CDRL2	AASNLQS (SEQ ID NO: 296)
CDRL3	QQSFDWPLT (SEQ ID NO: 297)
Antibody	SR-AB64
CDRH1	GSIRSSSYWG (SEQ ID NO: 298)
CDRH2	SISYSATYY (SEQ ID NO: 299)

CDRH3	AGDPSYDSIAGMQV (SEQ ID NO: 300)
CDRL1	RASQSIYYLN (SEQ ID NO: 301)
CDRL2	SASSRQS (SEQ ID NO: 302)
CDRL3	QQGFDFPLT (SEQ ID NO: 303)

[210] In some embodiments, antibodies of the present invention that selectively bind to a LTBP1-TGF β complex and/or a LTBP3-TGF β complex include any antibody, or antigen-binding portion thereof, comprising a CDRH1, CDRH2, CDRH3, CDRL1, CDRL2, or CDRL3, or combinations thereof, as provided in Table 5. In some embodiments, antibodies that selectively bind to a LTBP1-TGF β complex and/or a LTBP3-TGF β complex include CDRH1, CDRH2, CDRH3, CDRL1, CDRL2, and CDRL3 as provided in Table 5.

[211] The present invention also provides a nucleic acid sequence that encodes a molecule comprising CDRH1, CDRH2, CDRH3, CDRL1, CDRL2, or CDRL3, or combinations thereof, as provided in Table 5.

[212] Antibody heavy and light chain CDR3 domains may play a particularly important role in the binding specificity/affinity of an antibody for an antigen. Accordingly, in some embodiments, the antibodies, or antigen-binding portions thereof, that selectively bind to a LTBP1-TGF β complex and/or a LTBP3-TGF β complex, or the nucleic acid molecules that encode these antibodies, or antigen-binding portions thereof, can include at least the heavy and/or light chain CDR3 of the antibody shown in Table 5.

[213] Aspects of the invention relate to a monoclonal antibody, or antigen-binding portion thereof, that binds selectively to a LTBP1-TGF β complex and/or a LTBP3-TGF β complex, and that comprises six complementarity determining regions (CDRs): CDRH1, CDRH2, CDRH3, CDRL1, CDRL2, and CDRL3. The antibody, or antigen-binding portion thereof may have the CDRH1, CDRH2, CDRH3, CDRL1, CDRL2, and CDRL3 of one of the antibodies (*e.g.*, Ab42) shown in Table 5.

[214] In some embodiments, CDRH1 comprises a sequence as set forth in SEQ ID NO: 1. In some embodiments, CDRH2 comprises a sequence as set forth in SEQ ID NO: 2. In some embodiments, CDRH3 comprises a sequence as set forth in SEQ ID NO: 3. In some embodiments, CDRL1 comprises a sequence as set forth in SEQ ID NO: 4. In some embodiments, CDRL2 comprises a sequence as set forth in SEQ ID NO: 5. In some embodiments, CDRL3 comprises a sequence as set forth in SEQ ID NO: 6.

[215] In one aspect, the invention provides an isolated antibody, or an antigen-binding fragment thereof, that specifically binds a human LTBP1-proTGF β complex and/or a human LTBP3-proTGF β

complex and does not bind a human GARP-proTGFβ1 complex; wherein the antibody or the antigen-binding fragment thereof does not bind mature TGFβ1, mature TGFβ2 or mature TGFβ3; wherein the antibody or the antigen-binding fragment thereof is a fully human or humanized antibody or antigen-binding fragment thereof, wherein the antibody or the antigen-binding fragment thereof comprises at least three CDRs selected from the following, optionally comprising up to one or more amino acid changes for each of the CDRs: CDR-H1: SEQ ID NO:1; CDR-H2: SEQ ID NO:2; CDR-H3: SEQ ID NO:3; CDR-L1: SEQ ID NO:4; CDR-L2: SEQ ID NO:5; and, CDR-L3: SEQ ID NO:6. In some embodiments, the one or more amino acid changes comprises up to 1, 2, 3, 4, 5, or 6 amino acid changes for each of the CDRs.

[216] In some embodiments (*e.g.*, as for antibody SR-AB2, shown in Table 5), the antibody, or antigen-binding portion thereof, that selectively binds to a LTBP1-TGFβ complex and/or a LTBP3-TGFβ complex comprises: a CDRH1 comprising an amino acid sequence as set forth in SEQ ID NO: 1, a CDRH2 comprising an amino acid sequence as set forth in SEQ ID NO: 2, a CDRH3 comprising an amino acid sequence as set forth in SEQ ID NO: 3, a CDRL1 comprising an amino acid sequence as set forth in SEQ ID NO: 4, a CDRL2 comprising an amino acid sequence as set forth in SEQ ID NO: 5, and a CDRL3 comprising an amino acid sequence as set forth in SEQ ID NO: 6.

[217] In some embodiments, the antibody, or antigen-binding portion thereof, comprises a heavy chain variable region comprising a complementarity determining region 3 (CDR3) having the amino acid sequence of SEQ ID NO: 3 and a light chain variable region comprising a CDR3 having the amino acid sequence of SEQ ID NO: 6. In some embodiments, the antibody, or antigen-binding portion thereof, comprises a heavy chain variable region comprising a complementarity determining region 2 (CDR2) having the amino acid sequence of SEQ ID NO: 2 and a light chain variable region comprising a CDR2 having the amino acid sequence of SEQ ID NO: 5. In some embodiments, the antibody, or antigen-binding portion thereof, comprises a heavy chain variable region comprising a complementarity determining region 1 (CDR1) having the amino acid sequence of SEQ ID NO: 1 and a light chain variable region comprising a CDR1 having the amino acid sequence of SEQ ID NO: 4.

[218] The amino acid sequences of the heavy chain variable region (HCVR) and the light chain variable region (LCVR) of the antibody set forth in Table 5 (*e.g.*, SR-AB2) are provided in Table 6.

[219] In some embodiments, CDRH1 comprises a sequence as set forth in SEQ ID NO: 94. In some embodiments, CDRH2 comprises a sequence as set forth in SEQ ID NO: 95. In some embodiments, CDRH3 comprises a sequence as set forth in SEQ ID NO: 96. In some embodiments, CDRL1 comprises a sequence as set forth in SEQ ID NO: 97. In some embodiments, CDRL2 comprises a sequence as set forth in SEQ ID NO: 98. In some embodiments, CDRL3 comprises a sequence as set forth in SEQ ID NO: 99.

[220] In one aspect, the invention provides an isolated antibody, or an antigen-binding fragment thereof, that specifically binds a human LTBP1-proTGF β complex and/or a human LTBP3-proTGF β complex and does not bind a human GARP-proTGF β 1 complex; wherein the antibody or the antigen-binding fragment thereof does not bind mature TGF β 1, mature TGF β 2 or mature TGF β 3; wherein the antibody or the antigen-binding fragment thereof is a fully human or humanized antibody or antigen-binding fragment thereof, wherein the antibody or the antigen-binding fragment thereof comprises at least three CDRs selected from the following, optionally comprising one or more amino acid changes for each of the CDRs: CDR-H1: SEQ ID NO:94; CDR-H2: SEQ ID NO:95; CDR-H3: SEQ ID NO:96; CDR-L1: SEQ ID NO:97; CDR-L2: SEQ ID NO:98; and, CDR-L3: SEQ ID NO:99. In some embodiments, the one or more amino acid changes comprises up to 1, 2, 3, 4, 5, or 6 amino acid changes for each of the CDRs.

[221] In some embodiments (*e.g.*, as for antibody SR-AB10, shown in Table 5), the antibody, or antigen-binding portion thereof, that selectively binds to a LTBP1-TGF β complex and/or a LTBP3-TGF β complex comprises: a CDRH1 comprising an amino acid sequence as set forth in SEQ ID NO: 94, a CDRH2 comprising an amino acid sequence as set forth in SEQ ID NO: 95, a CDRH3 comprising an amino acid sequence as set forth in SEQ ID NO: 96, a CDRL1 comprising an amino acid sequence as set forth in SEQ ID NO: 97, a CDRL2 comprising an amino acid sequence as set forth in SEQ ID NO: 98, and a CDRL3 comprising an amino acid sequence as set forth in SEQ ID NO: 99.

[222] In some embodiments, the antibody, or antigen-binding portion thereof, comprises a heavy chain variable region comprising a complementarity determining region 3 (CDR3) having the amino acid sequence of SEQ ID NO: 96 and a light chain variable region comprising a CDR3 having the amino acid sequence of SEQ ID NO: 99. In some embodiments, the antibody, or antigen-binding portion thereof, comprises a heavy chain variable region comprising a complementarity determining region 2 (CDR2) having the amino acid sequence of SEQ ID NO: 95 and a light chain variable region comprising a CDR2 having the amino acid sequence of SEQ ID NO: 98. In some embodiments, the antibody, or antigen-binding portion thereof, comprises a heavy chain variable region comprising a complementarity determining region 1 (CDR1) having the amino acid sequence of SEQ ID NO: 94 and a light chain variable region comprising a CDR1 having the amino acid sequence of SEQ ID NO: 97.

[223] The amino acid sequences of the heavy chain variable region (HCVR) and the light chain variable region (LCVR) of the antibody set forth in Table 5 (*e.g.*, SR-AB10) are provided in Table 6.

[224] In some embodiments, CDRH1 comprises a sequence as set forth in SEQ ID NO: 100. In some embodiments, CDRH2 comprises a sequence as set forth in SEQ ID NO: 101. In some embodiments, CDRH3 comprises a sequence as set forth in SEQ ID NO: 102. In some embodiments,

CDRL1 comprises a sequence as set forth in SEQ ID NO: 103. In some embodiments, CDRL2 comprises a sequence as set forth in SEQ ID NO: 104. In some embodiments, CDRL3 comprises a sequence as set forth in SEQ ID NO: 105.

[225] In one aspect, the invention provides an isolated antibody, or an antigen-binding fragment thereof, that specifically binds a human LTBP1-proTGF β complex and/or a human LTBP3-proTGF β complex and does not bind a human GARP-proTGF β 1 complex; wherein the antibody or the antigen-binding fragment thereof does not bind mature TGF β 1, mature TGF β 2 or mature TGF β 3; wherein the antibody or the antigen-binding fragment thereof is a fully human or humanized antibody or antigen-binding fragment thereof, wherein the antibody or the antigen-binding fragment thereof comprises at least three CDRs selected from the following, optionally comprising one or more amino acid changes for each of the CDRs: CDR-H1: SEQ ID NO:100; CDR-H2: SEQ ID NO:101; CDR-H3: SEQ ID NO:102; CDR-L1: SEQ ID NO:103; CDR-L2: SEQ ID NO:104; and, CDR-L3: SEQ ID NO:105. In some embodiments, the one or more amino acid changes comprises up to 1, 2, 3, 4, 5, or 6 amino acid changes for each of the CDRs.

[226] In some embodiments (*e.g.*, as for antibody SR-AB13, shown in Table 5), the antibody, or antigen-binding portion thereof, that selectively binds to a LTBP1-TGF β complex and/or a LTBP3-TGF β complex comprises: a CDRH1 comprising an amino acid sequence as set forth in SEQ ID NO: 100, a CDRH2 comprising an amino acid sequence as set forth in SEQ ID NO: 101, a CDRH3 comprising an amino acid sequence as set forth in SEQ ID NO: 102, a CDRL1 comprising an amino acid sequence as set forth in SEQ ID NO: 103, a CDRL2 comprising an amino acid sequence as set forth in SEQ ID NO: 104, and a CDRL3 comprising an amino acid sequence as set forth in SEQ ID NO: 105.

[227] In some embodiments, the antibody, or antigen-binding portion thereof, comprises a heavy chain variable region comprising a complementarity determining region 3 (CDR3) having the amino acid sequence of SEQ ID NO: 102 and a light chain variable region comprising a CDR3 having the amino acid sequence of SEQ ID NO: 105. In some embodiments, the antibody, or antigen-binding portion thereof, comprises a heavy chain variable region comprising a complementarity determining region 2 (CDR2) having the amino acid sequence of SEQ ID NO: 101 and a light chain variable region comprising a CDR2 having the amino acid sequence of SEQ ID NO: 104. In some embodiments, the antibody, or antigen-binding portion thereof, comprises a heavy chain variable region comprising a complementarity determining region 1 (CDR1) having the amino acid sequence of SEQ ID NO: 100 and a light chain variable region comprising a CDR1 having the amino acid sequence of SEQ ID NO: 103.

[228] The amino acid sequences of the heavy chain variable region (HCVR) and the light chain variable region (LCVR) of the antibody set forth in Table 5 (*e.g.*, SR-AB13) are provided in Table 6.

[229] In some embodiments, CDRH1 comprises a sequence as set forth in SEQ ID NO: 124. In some embodiments, CDRH2 comprises a sequence as set forth in SEQ ID NO: 125. In some embodiments, CDRH3 comprises a sequence as set forth in SEQ ID NO: 126. In some embodiments, CDRL1 comprises a sequence as set forth in SEQ ID NO: 127. In some embodiments, CDRL2 comprises a sequence as set forth in SEQ ID NO: 128. In some embodiments, CDRL3 comprises a sequence as set forth in SEQ ID NO: 129.

[230] In one aspect, the invention provides an isolated antibody, or an antigen-binding fragment thereof, that specifically binds a human LTBP1-proTGF β complex and/or a human LTBP3-proTGF β complex; wherein the antibody or the antigen-binding fragment thereof does not bind mature TGF β 1, mature TGF β 2 or mature TGF β 3; wherein the antibody or the antigen-binding fragment thereof is a fully human or humanized antibody or antigen-binding fragment thereof, wherein the antibody or the antigen-binding fragment thereof comprises at least three (optionally all six) CDRs selected from the following, optionally comprising up to one or more amino acid changes for each of the CDRs: CDR-H1: SEQ ID NO:124; CDR-H2: SEQ ID NO:125; CDR-H3: SEQ ID NO:126; CDR-L1: SEQ ID NO:127; CDR-L2: SEQ ID NO:128; and, CDR-L3: SEQ ID NO:129. In some embodiments, the one or more amino acid changes comprises up to 1, 2, 3, 4, 5, or 6 amino acid changes for each of the CDRs.

[231] In some embodiments (*e.g.*, as for antibody SR-AB31, shown in Table 5), the antibody, or antigen-binding portion thereof, that selectively binds to a LTBP1-TGF β complex and/or a LTBP3-TGF β complex comprises: a CDRH1 comprising an amino acid sequence as set forth in SEQ ID NO: 124, a CDRH2 comprising an amino acid sequence as set forth in SEQ ID NO: 125, a CDRH3 comprising an amino acid sequence as set forth in SEQ ID NO: 126, a CDRL1 comprising an amino acid sequence as set forth in SEQ ID NO: 127, a CDRL2 comprising an amino acid sequence as set forth in SEQ ID NO: 128, and a CDRL3 comprising an amino acid sequence as set forth in SEQ ID NO: 129.

[232] In some embodiments, the antibody, or antigen-binding portion thereof, comprises a heavy chain variable region comprising a complementarity determining region 3 (CDR3) having the amino acid sequence of SEQ ID NO: 126 and a light chain variable region comprising a CDR3 having the amino acid sequence of SEQ ID NO: 129. In some embodiments, the antibody, or antigen-binding portion thereof, comprises a heavy chain variable region comprising a complementarity determining region 2 (CDR2) having the amino acid sequence of SEQ ID NO: 125 and a light chain variable region comprising a CDR2 having the amino acid sequence of SEQ ID NO: 128. In some embodiments, the antibody, or antigen-binding portion thereof, comprises a heavy chain variable region comprising a complementarity determining region 1 (CDR1) having the amino acid sequence of SEQ ID NO: 124 and a light chain variable region comprising a CDR1 having the amino acid sequence of SEQ ID NO: 127.

[233] The amino acid sequences of the HCVR and the LCVR of the antibody set forth in Table 5 (*e.g.*, SR-AB31) are provided in Table 6.

[234] In some embodiments, CDRH1 comprises a sequence as set forth in SEQ ID NO: 166. In some embodiments, CDRH2 comprises a sequence as set forth in SEQ ID NO: 167. In some embodiments, CDRH3 comprises a sequence as set forth in SEQ ID NO: 168. In some embodiments, CDRL1 comprises a sequence as set forth in SEQ ID NO: 169. In some embodiments, CDRL2 comprises a sequence as set forth in SEQ ID NO: 170. In some embodiments, CDRL3 comprises a sequence as set forth in SEQ ID NO: 171.

[235] In one aspect, the invention provides an isolated antibody, or an antigen-binding fragment thereof, that specifically binds a human LTBP1-proTGF β complex and/or a human LTBP3-proTGF β complex; wherein the antibody or the antigen-binding fragment thereof does not bind mature TGF β 1, mature TGF β 2 or mature TGF β 3; wherein the antibody or the antigen-binding fragment thereof is a fully human or humanized antibody or antigen-binding fragment thereof, wherein the antibody or the antigen-binding fragment thereof comprises at least three (optionally all six) CDRs selected from the following, optionally comprising up to one or more amino acid changes for each of the CDRs: CDR-H1: SEQ ID NO:166; CDR-H2: SEQ ID NO:167; CDR-H3: SEQ ID NO:168; CDR-L1: SEQ ID NO:169; CDR-L2: SEQ ID NO:170; and, CDR-L3: SEQ ID NO:171. In some embodiments, the one or more amino acid changes comprises up to 1, 2, 3, 4, 5, or 6 amino acid changes for each of the CDRs.

[236] In some embodiments (*e.g.*, as for antibody SR-AB42, shown in Table 5), the antibody, or antigen-binding portion thereof, that selectively binds to a LTBP1-TGF β complex and/or a LTBP3-TGF β complex comprises: a CDRH1 comprising an amino acid sequence as set forth in SEQ ID NO: 166, a CDRH2 comprising an amino acid sequence as set forth in SEQ ID NO: 167, a CDRH3 comprising an amino acid sequence as set forth in SEQ ID NO: 168, a CDRL1 comprising an amino acid sequence as set forth in SEQ ID NO: 169, a CDRL2 comprising an amino acid sequence as set forth in SEQ ID NO: 170, and a CDRL3 comprising an amino acid sequence as set forth in SEQ ID NO: 171.

[237] In some embodiments, the antibody, or antigen-binding portion thereof, comprises a heavy chain variable region comprising a complementarity determining region 3 (CDR3) having the amino acid sequence of SEQ ID NO: 168 and a light chain variable region comprising a CDR3 having the amino acid sequence of SEQ ID NO: 171. In some embodiments, the antibody, or antigen-binding portion thereof, comprises a heavy chain variable region comprising a complementarity determining region 2 (CDR2) having the amino acid sequence of SEQ ID NO: 167 and a light chain variable region comprising a CDR2 having the amino acid sequence of SEQ ID NO: 170. In some embodiments, the antibody, or antigen-binding portion thereof, comprises a heavy chain variable

region comprising a complementarity determining region 1 (CDR1) having the amino acid sequence of SEQ ID NO: 166 and a light chain variable region comprising a CDR1 having the amino acid sequence of SEQ ID NO: 169.

[238] The amino acid sequences of the HCVR and the LCVR of the antibody set forth in Table 5 (*e.g.*, SR-AB42) are provided in Table 6.

[239] In some embodiments, CDRH1 comprises a sequence as set forth in SEQ ID NO: 292. In some embodiments, CDRH2 comprises a sequence as set forth in SEQ ID NO: 293. In some embodiments, CDRH3 comprises a sequence as set forth in SEQ ID NO: 294. In some embodiments, CDRL1 comprises a sequence as set forth in SEQ ID NO: 295. In some embodiments, CDRL2 comprises a sequence as set forth in SEQ ID NO: 296. In some embodiments, CDRL3 comprises a sequence as set forth in SEQ ID NO: 297.

[240] In one aspect, the invention provides an isolated antibody, or an antigen-binding fragment thereof, that specifically binds a human LTBP1-proTGF β complex and/or a human LTBP3-proTGF β complex; wherein the antibody or the antigen-binding fragment thereof does not bind mature TGF β 1, mature TGF β 2 or mature TGF β 3; wherein the antibody or the antigen-binding fragment thereof is a fully human or humanized antibody or antigen-binding fragment thereof, wherein the antibody or the antigen-binding fragment thereof comprises at least three (optionally all six) CDRs selected from the following, optionally comprising up to one or more amino acid changes for each of the CDRs: CDR-H1: SEQ ID NO:292; CDR-H2: SEQ ID NO:293; CDR-H3: SEQ ID NO:294; CDR-L1: SEQ ID NO:295; CDR-L2: SEQ ID NO:296; and, CDR-L3: SEQ ID NO:297. In some embodiments, the one or more amino acid changes comprises up to 1, 2, 3, 4, 5, or 6 amino acid changes for each of the CDRs.

[241] In some embodiments (*e.g.*, as for antibody SR-AB63, shown in Table 5), the antibody, or antigen-binding portion thereof, that selectively binds to a LTBP1-TGF β complex and/or a LTBP3-TGF β complex comprises: a CDRH1 comprising an amino acid sequence as set forth in SEQ ID NO: 292, a CDRH2 comprising an amino acid sequence as set forth in SEQ ID NO: 293, a CDRH3 comprising an amino acid sequence as set forth in SEQ ID NO: 294, a CDRL1 comprising an amino acid sequence as set forth in SEQ ID NO: 295, a CDRL2 comprising an amino acid sequence as set forth in SEQ ID NO: 296, and a CDRL3 comprising an amino acid sequence as set forth in SEQ ID NO: 297.

[242] In some embodiments, the antibody, or antigen-binding portion thereof, comprises a heavy chain variable region comprising a complementarity determining region 3 (CDR3) having the amino acid sequence of SEQ ID NO: 294 and a light chain variable region comprising a CDR3 having the amino acid sequence of SEQ ID NO: 297. In some embodiments, the antibody, or antigen-binding portion thereof, comprises a heavy chain variable region comprising a complementarity determining

region 2 (CDR2) having the amino acid sequence of SEQ ID NO: 293 and a light chain variable region comprising a CDR2 having the amino acid sequence of SEQ ID NO: 296. In some embodiments, the antibody, or antigen-binding portion thereof, comprises a heavy chain variable region comprising a complementarity determining region 1 (CDR1) having the amino acid sequence of SEQ ID NO: 292 and a light chain variable region comprising a CDR1 having the amino acid sequence of SEQ ID NO: 295.

[243] The amino acid sequences of the HCVR and the LCVR of the antibody set forth in Table 5 (*e.g.*, SR-AB63) are provided in Table 6.

[244] Ten additional antibodies (Ab3-Ab12) were developed that specifically bind to a LTBP1-TGF β complex and/or a LTBP3-TGF β complex, and inhibit release of mature TGF β presented in the context of LTBP1/3. Table 6 also provides the HCVR and LCVR amino acid sequences of these additional LTBP context-specific antibodies, in addition to the HCVR and LCVR amino acid sequences of the antibodies referred to in Table 5.

Table 6. Heavy Chain Variable Region Sequence and Light Chain Variable Region Sequence of Antibodies that Specifically Bind a LTBP1/3-TGF β 1 Complex

Antibody	HCVR Sequence	LCVR Sequence
SR-AB2	QVQLVQSGAEVKKPGASVKVSCKA SGYTFTSYGISWVRQAPGQGLEWM GWISAYNGNTNYAQKLQGRVTMTT DTSTSTAYMELRSLRSDDTAVYYCA RAPLGNFDSWGQGTMTVSS (SEQ ID NO: 7)	NFMLTQPHSVSESPGKTVTISCTRSS GSIASNYVQWYQQRPGSSPTTVIYE DNQRPSGVDPDRFSGSIDSSNSASLTI SGLKTEDEADYYCQSYDSSNHPVVF GGGTKLTVL (SEQ ID NO: 8)
SR-AB3	QMQLVQSGAEVKKPGASVKVSCKA SGYTFTSYGISWVRQAPGQGLEWM GWISAYNGNTNYAQKLQGRVTMTT NTSTSTAYMELRSLRSDDTAVYYCA RDDYYYYGMDVWGQGTTLTVSS (SEQ ID NO: 74)	QSGLTQPASVSGSPGQSVTISCTGTS SDVGGYNYASWYQHPGKAPKLM YDVSKRPSGVDPDRFSGSKSGNTASL TISGLQAEDEADYYCSTSSSTVYF GTGTKLTVL (SEQ ID NO: 75)
SR-AB4	QVQLQQWGAGLLKPSETLSLTCAV YGGSFSGYYWSWIRQPPGKGLEWI GEIHSNSTNYNPSLKSRTISVDTSK NQFSLKLSSVTAADTAVYYCARGV GLGRFDPWGQGTTLTVSS (SEQ ID NO: 76)	QSELTQSPSASGTPGQRVTISCSGSN SNIGTNTVNWYQQFPGTAPKLLIYY NDQRPSGVSDRFSGSRSGTSASLAIN GLQSEDEADYYCATWDDSLSGVVF GGGTKLTVL (SEQ ID NO: 77)
SR-AB5	QVQLQQWGAGLLKPSETLSLTCAV YGGSFSGYYWSWIRQPPGKGLEWI GEINHSNSTNYNPSLKSRTISVDTS KNQFSLKLSSVTAADTAVYYCARG VGLGRFDPWGQGTTLTVSS (SEQ ID NO: 78)	QSELTQSPSASGTPGQRVTISCSGSN SNIGTNTVNWYQQFPGTAPKLLIYY NDQRPSGVSDRFSGSRSGTSASLAIN GLQSEDEADYYCATWDDSLSGVVF GGGTKLTVL (SEQ ID NO: 79)
SR-AB6	QVQLQQSGPGLVRPSQTLSTCAISG DSVSSNGAAWNWIRQSPSRGLEWL GRYYRSKWYNDYAVSVKSRITNP	NFMLTQPHSVSESPGKTVTISCTRSS GSIASNYVQWYQQRPGSAPTTVIYD DKQRPSGIPDRFSGSIDSSNSASLTI

	DTSKNQFSLKLTSTVPEDTAVYYCA RGEDWGYAFDIWGQGLTVTVSS (SEQ ID NO: 80)	SGLKTEDEADYYCQSYDSSNVVFG GGTKVTVL (SEQ ID NO: 81)
SR-AB7	QVQLVQSGAEVKKPGASVKVSCKA SGYTFTSYGISWVRQAPGQGLEWM GWISAYDGNTNYAQKLQGRVTMTT DTSTSTAYMELSSLRSDDTAVYYCA RNPYYYYMDVWGQGTITVTVSS (SEQ ID NO: 82)	QSELTQAPSVSVAPGQTARITCGGN NIGGRSKSVHWYQHKLGQAPVLIV YDNTDRPSGISERFSGSSSVNAATLT ITTAEAGDEGDYYCQVWDVSTDHV VFGGGTKVTVL (SEQ ID NO: 83)
SR-AB8	QVQLVESGAEVKKPGASVKVSCKA SGYTFTGYMHVVRQAPGQGLEW MGWINPNGGGTNYAQKFQGRVTM TRDTSISTAYMELSLRSDDTAVYY CANRRRGSADFIDWGQGLTVTVSS (SEQ ID NO: 84)	NFMLTQPHSVSESPGKTVTISCTGSS GSIASNYVQWYQQRPGSSPTTVIYE DNQRPSGVPDRFSGSIDSSSNSASLTI SGLKTEDEADYYCQSYDDNYHVIF GGGKLTVL (SEQ ID NO: 85)
SR-AB9	QVQLVESGGALVQPGGSLRLSCAAS GFTFSSYAMHWVRQAPGKGLEWV AVISYDGSNKYYADSVKGRFTISR NSKNTLYLQMNSLRAEDTAVYYCA KETGYGFLFWGQGTMTVTVSS (SEQ ID NO: 86)	NFMLTQPHSVSESPGRTLTIPCFRSS GNIGDSYVHWYQQRPGSAPTTVIYR DSQRPSGVPDRFSGSIDSSSNSASLTI SGLKTEDEAAYYCQSYDRSNQWVF GGGKLTVL (SEQ ID NO: 87)
SR-AB10	QLQLQESGGGVVQGRSLRLSCAAS GFTFNYPPIHWVRQAPGKGLEWVA VMSYDGINKYYADSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYYCAR PRIAARRGGFDYWGQGLTVTVSS (SEQ ID NO: 88)	NFMLTQPHSVSESPGKTVTISCTRSS GNIDNNYVQWYQQRPGSSPTTVIYE DNQRPSGVPDRFSGSIDSSSNSASLTI SGLKTEDEADYYCQSYSDSNQGVV FGGGKLTVL (SEQ ID NO: 89)
SR-AB11	QVQLVQSGAEVKKPGASVKVSCKA SGYTFTSYGISWVRQAPGQGLEWM GWISAYNGNTDYAQKLQGRVTMTT DTSTSTAYMELRGLRSDDTAVYYC ARAPLGNFDSWGQGLTVTVSS (SEQ ID NO: 90)	NFMLTQPHSVSESPGKTVTISCTRSS GSIASNYVQWYQQRPGSAPTTVIYE DNQRPSGVPDRFSGSIDSSSNSASLTI SGLKTEDEADYYCQSYDSSNHVVF GGGKTVTVL (SEQ ID NO: 91)
SR-AB12	EVQLLESGGGVVQGRSLRLSCAAS GFTFPNYAMSWVRQAPGKGLEWVS AISGSGGSTYYADSVKGRFTISRDN KNTLYLQMNSLRAEDTAVYYCAKD LEGGYYWDYYYYGMDVWGQGLT VTVSS (SEQ ID NO: 92)	NFMLTQPHSVSESPGKTVTISCTRSS GSIASNYVQWYQQRPGSSPTTVIYE DNQRPSGVPDRFSGSIDSSSNSASLTI SGLKTEDEADYYCQSYDSSIVVFGG GTQLTVL (SEQ ID NO: 93)
SR-AB13	QLQLQESGPGLVKPSETLSLTCTVSG GSISSSSYYWGWRQPPGKGLEWIG SISYSGSTYYNPSLKSRTISVDTSK NQFSLKLSSVTAADTAVYYCARDPS YDSIAGMDVWGQGTITVTVSS (SEQ ID NO: 106)	DIQLTQSPSSLSASVGDRVITICRAS QSISSYLNWYQKPGKAPKLLIYAA SNLQSGVPSRFSGSGSGTDFTLTISSL QPEDFATYYCQSFDFPFTFGGGTK VEIK (SEQ ID NO: 107)
SR-AB22	QVQLVESGGGVVQGRSLRLSCAAS GFTFRSYVMHWVRQAPGKGLEWV AVISHEGSLKYYADSVKGRFTISR NSKNTLYLQMNSLRAEDTAVYYCA VPRIAARRGGFGYWGQGLTVTVSS (SEQ ID NO: 114)	NFMLTQPHSVSESPGKTVTISCTRSS GNIDNNYVQWYQQRPGSSPTTVIYE DNQRPSGVPDRFSGSIDSSSNSASLTI SGLKTEDEADYYCQSYSDSNQGVV FGGGKLTVL (SEQ ID NO: 115)
SR-AB23	QVQLVESGGGVVQGRSLRLSCAAS GFTFRSYVMHWVRQAPGKGLEWV AVISHEGSLKYYADSVKGRFTISR	NFMLTQPHSVSESPGKTVTISCTRSS GNIDNNYVQWYQQRPGSSPTTVIYE DNQRPSGVPDRFSGSIDSSSNSASLTI

	NSKNTLYLQMNSLRAEDTAVYYCA RPRIAARRGGFGYWGQGLTVTVSS (SEQ ID NO: 122)	SGLKTEDEADYYCQSYDSDNQGVV FGGGTKLTVL (SEQ ID NO: 123)
SR-AB31	QVQLVESGGGVVQPGRSLRLSCAAS GFTFRSYVMHWVRQAPGKGLEWV AVISHEGSLKYYADSVKGRFTISR NSKNTLYLQMNSLRAEDTAVYYCA VPRIAARRGGFGYWGQGLTVTVSS (SEQ ID NO: 304)	NFMLTQPHSVSESPGKTVTISCTRSS GNIDNNYVQWYQQRPGSSPTTVIYE DNQRPSGVPDRFSGSIDSSNSASLTI SGLKTEDEADYYCQSYDFNNQGVV FGGGTKLTVL (SEQ ID NO: 305)
SR-AB34	QVQLVESGGGVVQPGRSLRLSCAAS GFTFRSYVMHWVRQAPGKGLEWV AVISHEGSLKYYADSVKGRFTISR NSKNTLYLQMNSLRAEDTAVYYCA VPRIAARRGGFGYWGQGLTVTVSS (SEQ ID NO: 306)	NFMLTQPHSVSESPGKTVTISCTRSS GNIDNNYVQWYQQRPGSSPTTVIYE DNQRPSGVPDRFSGSIDSSNSASLTI SGLKTEDEADYYCQSYDYDAQGVV FGGGTKLTVL (SEQ ID NO: 307)
SR-AB37	QVQLVESGGGVVQPGRSLRLSCAAS GFTFRSYVMHWVRQAPGKGLEWV AVISHEGSLKYYADSVKGRFTISR NSKNTLYLQMNSLRAEDTAVYYCA VPRIAARRGGFGYWGQGLTVTVSS (SEQ ID NO: 308)	NFMLTQPHSVSESPGKTVTISCTRSS GLIDNNYVQWYQQRPGSSPTTVIYE DNQRPSGVPDRFSGSIDSSNSASLTI SGLKTEDEADYYCQSYDSDLQRVV FGGGTKLTVL (SEQ ID NO: 309)
SR-AB38	QVQLVESGGGVVQPGRSLRLSCAAS GFTFRSYVMHWVRQAPGKGLEWV AVISHEGSLKYYADSVKGRFTISR NSKNTLYLQMNSLRAEDTAVYYCA VPRIAARRGGFGYWGQGLTVTVSS (SEQ ID NO: 310)	NFMLTQPHSVSESPGKTVTISCTRSS GSIDNNYVQWYQQRPGSSPTTVIYE DFIRPSGVPDRFSGSIDSSNSASLTIS GLKTEDEADYYCQSYDDDLQGVVF GGGTKLTVL (SEQ ID NO: 311)
SR-AB39	QVQLVESGGGVVQPGRSLRLSCAAS GFTFRSYVMHWVRQAPGKGLEWV AVISHEGSLKYYADSVKGRFTISR NSKNTLYLQMNSLRAEDTAVYYCA VPRIAARRGGFGYWGQGLTVTVSS (SEQ ID NO: 312)	NFMLTQPHSVSESPGKTVTISCTRSS GLIDNNYVQWYQQRPGSSPTTVIYE DAQRPSGVPDRFSGSIDSSNSASLTI SGLKTEDEADYYCQSYDHDEQGVV FGGGTKLTVL (SEQ ID NO: 313)
SR-AB40	QVQLVESGGGVVQPGRSLRLSCAAS GFTFRSYVMHWVRQAPGKGLEWV AVISHEGSLKYYADSVKGRFTISR NSKNTLYLQMNSLRAEDTAVYYCA RPRIAARRGGFGYWGQGLTVTVSS (SEQ ID NO: 314)	NFMLTQPHSVSESPGKTVTISCTRSS GNIDNNYVQWYQQRPGSSPTTVIYE DNQRPSGVPDRFSGSIDSSNSASLTI SGLKTEDEADYYCQSYDYSNQGVV FGGGTKLTVL (SEQ ID NO: 315)
SR-AB41	QVQLVESGGGVVQPGRSLRLSCAAS GFTFRSYVMHWVRQAPGKGLEWV AVISHEGSLKYYADSVKGRFTISR NSKNTLYLQMNSLRAEDTAVYYCA RPRIAARRGGFGYWGQGLTVTVSS (SEQ ID NO: 316)	NFMLTQPHSVSESPGKTVTISCTRSS GNIDNNYVQWYQQRPGSSPTTVIYE DNQRPSGVPDRFSGSIDSSNSASLTI SGLKTEDEADYYCQSYDYDNQAVV FGGGTKLTVL (SEQ ID NO: 317)
SR-AB42	QVQLVESGGGVVQPGRSLRLSCAAS GFTFRSYVMHWVRQAPGKGLEWV AVISHEGSLKYYADSVKGRFTISR NSKNTLYLQMNSLRAEDTAVYYCA RPRIAARRGGFGYWGQGLTVTVSS (SEQ ID NO: 318)	NFMLTQPHSVSESPGKTVTISCTRSS GNIDNNYVQWYQQRPGSSPTTVIYE DNQRPSGVPDRFSGSIDSSNSASLTI SGLKTEDEADYYCQSYDYDTQGVV FGGGTKLTVL (SEQ ID NO: 319)
SR-AB43	QVQLVESGGGVVQPGRSLRLSCAAS GFTFRSYVMHWVRQAPGKGLEWV AVISHEGSLKYYADSVKGRFTISR	NFMLTQPHSVSESPGKTVTISCTRSS GNIDNNYVQWYQQRPGSSPTTVIYE DNVRPSGVPDRFSGSIDSSNSASLTI

	NSKNTLYLQMNSLRAEDTAVYYCA RPRIAARRGGFGYWGQGLTVTVSS (SEQ ID NO: 320)	SGLKTEDEADYYCQSYDSNDQRRV FGGGTKLTVL (SEQ ID NO: 321)
SR-AB44	QVQLVESGGGVVQPGRSLRLSCAAS GFTFRSYVMHWVRQAPGKGLEWV AVISHEGSLKYYADSVKGRFTISR NSKNTLYLQMNSLRAEDTAVYYCA RPRIAARRGGFGYWGQGLTVTVSS (SEQ ID NO: 322)	NFMLTQPHSVSESPGKTVTISCTRSH GNIDNYVQWYQQRPGSSPTTVIYE DNVRPSGVPDRFSGSIDSSNSASLTI SGLKTEDEADYYCQSYDSNDQLVV FGGGTKLTVL (SEQ ID NO: 323)
SR-AB45	QVQLVESGGGVVQPGRSLRLSCAAS GFTFRSYVMHWVRQAPGKGLEWV AVISHEGSLKYYADSVKGRFTISR NSKNTLYLQMNSLRAEDTAVYYCA RPRIAARRGGFGYWGQGLTVTVSS (SEQ ID NO: 324)	NFMLTQPHSVSESPGKTVTISCTRSS GAIDNYVQWYQQRPGSSPTTVIYE DFQRPSGVPDRFSGSIDSSNSASLTI SGLKTEDEADYYCQSYDDDLQGVV FGGGTKLTVL (SEQ ID NO: 325)
SR-AB46	QVQLVESGGGVVQPGRSLRLSCAAS GFTFRSYVMHWVRQAPGKGLEWV AVISHEGSGKYYADSVKGRFTISR NSKNTLYLQMNSLRAEDTAVYYCA RPRIAARRGGFGSWGQGLTVTVSS (SEQ ID NO: 326)	NFMLTQPHSVSESPGKTVTISCTRSS GNIDNYVQWYQQRPGSSPTTVIYE DNVRPSGVPDRFSGSIDSSNSASLTI SGLKTEDEADYYCQSYDSNDQRRV FGGGTKLTVL (SEQ ID NO: 327)
SR-AB47	QVQLVESGGGVVQPGRSLRLSCAAS GFTFRSYVMHWVRQAPGKGLEWV AVISHEGSGKYYADSVKGRFTISR NSKNTLYLQMNSLRAEDTAVYYCA RPRIAARRGGFGSWGQGLTVTVSS (SEQ ID NO: 328)	NFMLTQPHSVSESPGKTVTISCTRSS GNIDNYVQWYQQRPGSSPTTVIYE DNVRPSGVPDRFSGSIDSSNSASLTI SGLKTEDEADYYCQSYDYDNQAVV FGGGTKLTVL (SEQ ID NO: 329)
SR-AB48	QVQLVESGGGVVQPGRSLRLSCAAS GFTFRSYVMHWVRQAPGKGLEWV AVISHEGSGKYYADSVKGRFTISR NSKNTLYLQMNSLRAEDTAVYYCA RPRIAARRGGFGSWGQGLTVTVSS (SEQ ID NO: 330)	NFMLTQPHSVSESPGKTVTISCTRSS GNIDNYVQWYQQRPGSSPTTVIYE DNVRPSGVPDRFSGSIDSSNSASLTI SGLKTEDEADYYCQSYDYDTQGVV FGGGTKLTVL (SEQ ID NO: 331)
SR-AB49	QVQLVESGGGVVQPGRSLRLSCAAS GFTFRSYVMHWVRQAPGKGLEWV AVISHEGSGKYYADSVKGRFTISR NSKNTLYLQMNSLRAEDTAVYYCA RPRIAARRGGFGSWGQGLTVTVSS (SEQ ID NO: 332)	NFMLTQPHSVSESPGKTVTISCTRSS GNIDNYVQWYQQRPGSSPTTVIYE DNVRPSGVPDRFSGSIDSSNSASLTI SGLKTEDEADYYCQGYDWDQGV VFGGTKLTVL (SEQ ID NO: 333)
SR-AB50	QVQLVESGGGVVQPGRSLRLSCAAS GFTFRSYVMHWVRQAPGKGLEWV AVISHEGSGKYYADSVKGRFTISR NSKNTLYLQMNSLRAEDTAVYYCA RPRIAARRGGFGTWGQGLTVTVSS (SEQ ID NO: 334)	NFMLTQPHSVSESPGKTVTISCTRSS GNIDNYVQWYQQRPGSSPTTVIYE DNVRPSGVPDRFSGSIDSSNSASLTI SGLKTEDEADYYCQSYDSNDQRRV FGGGTKLTVL (SEQ ID NO: 335)
SR-AB51	QVQLVESGGGVVQPGRSLRLSCAAS GFTFRSYVMHWVRQAPGKGLEWV AVISHEGSGKYYADSVKGRFTISR NSKNTLYLQMNSLRAEDTAVYYCA RPRIAARRGGFGTWGQGLTVTVSS (SEQ ID NO: 336)	NFMLTQPHSVSESPGKTVTISCTRSS GNIDNYVQWYQQRPGSSPTTVIYE DNVRPSGVPDRFSGSIDSSNSASLTI SGLKTEDEADYYCQSYDYDNQAVV FGGGTKLTVL (SEQ ID NO: 337)
SR-AB52	QVQLVESGGGVVQPGRSLRLSCAAS GFTFRSYVMHWVRQAPGKGLEWV AVISHEGSGKYYADSVKGRFTISR	NFMLTQPHSVSESPGKTVTISCTRSS GNIDNYVQWYQQRPGSSPTTVIYE DNVRPSGVPDRFSGSIDSSNSASLTI

	NSKNTLYLQMNSLRAEDTAVYYCA RPRIAARRGGFGTWGQGLTVTVSS (SEQ ID NO: 338)	SGLKTEDEADYYCQSYDYDTQGVV FGGGTKLTVL (SEQ ID NO: 339)
SR-AB53	QVQLVESGGGVVQPGRSLRLSCAAS GFTFRSYVMHWVRQAPGKGLEWV AVISHEGSGKYYADSVKGRFTISR NSKNTLYLQMNSLRAEDTAVYYCA RPRIAARRGGFGTWGQGLTVTVSS (SEQ ID NO: 340)	NFMLTQPHSVSESPGKTVTISCTRSS GNIDYNYVQWYQQRPGSSPTTVIYE DNVRPSGVPDRFSGSIDSSNSASLTI SGLKTEDEADYYCQGYDWDQGV VFGGGTKLTVL (SEQ ID NO: 341)
SR-AB54	QVQLVESGGGVVQPGRSLRLSCAAS GFTFRSYVMHWVRQAPGKGLEWV AVISHEGSGKYYADSVKGRFTISR NSKNTLYLQMNSLRAEDTAVYYCA LPRIAARRGGFGSWGQGLTVTVSS (SEQ ID NO: 342)	NFMLTQPHSVSESPGKTVTISCTRSS GNIDYNYVQWYQQRPGSSPTTVIYE DNVRPSGVPDRFSGSIDSSNSASLTI SGLKTEDEADYYCQSYDSNDQVRV FGGGTKLTVL (SEQ ID NO: 343)
SR-AB55	QVQLVESGGGVVQPGRSLRLSCAAS GFTFRSYVMHWVRQAPGKGLEWV AVISHEGSGKYYADSVKGRFTISR NSKNTLYLQMNSLRAEDTAVYYCA LPRIAARRGGFGSWGQGLTVTVSS (SEQ ID NO: 344)	NFMLTQPHSVSESPGKTVTISCTRSS GNIDYNYVQWYQQRPGSSPTTVIYE DNVRPSGVPDRFSGSIDSSNSASLTI SGLKTEDEADYYCQSYDYNQAVV FGGGTKLTVL (SEQ ID NO: 345)
SR-AB56	QVQLVESGGGVVQPGRSLRLSCAAS GFTFRSYVMHWVRQAPGKGLEWV AVISHEGSGKYYADSVKGRFTISR NSKNTLYLQMNSLRAEDTAVYYCA LPRIAARRGGFGSWGQGLTVTVSS (SEQ ID NO: 346)	NFMLTQPHSVSESPGKTVTISCTRSS GNIDYNYVQWYQQRPGSSPTTVIYE DNVRPSGVPDRFSGSIDSSNSASLTI SGLKTEDEADYYCQSYDYDTQGVV FGGGTKLTVL (SEQ ID NO: 347)
SR-AB57	QVQLVESGGGVVQPGRSLRLSCAAS GFTFRSYVMHWVRQAPGKGLEWV AVISHEGSGKYYADSVKGRFTISR NSKNTLYLQMNSLRAEDTAVYYCA LPRIAARRGGFGSWGQGLTVTVSS (SEQ ID NO: 348)	NFMLTQPHSVSESPGKTVTISCTRSS GNIDYNYVQWYQQRPGSSPTTVIYE DNVRPSGVPDRFSGSIDSSNSASLTI SGLKTEDEADYYCQGYDWDQGV VFGGGTKLTVL (SEQ ID NO: 349)
SR-AB58	QVQLVESGGGVVQPGRSLRLSCAAS GFTFRSYVMHWVRQAPGKGLEWV AVISHEGSGKYYADSVKGRFTISR NSKNTLYLQMNSLRAEDTAVYYCA LPRIAARRGGFGTWGQGLTVTVSS (SEQ ID NO: 350)	NFMLTQPHSVSESPGKTVTISCTRSS GNIDYNYVQWYQQRPGSSPTTVIYE DNVRPSGVPDRFSGSIDSSNSASLTI SGLKTEDEADYYCQSYDSNDQVRV FGGGTKLTVL (SEQ ID NO: 351)
SR-AB59	QVQLVESGGGVVQPGRSLRLSCAAS GFTFRSYVMHWVRQAPGKGLEWV AVISHEGSGKYYADSVKGRFTISR NSKNTLYLQMNSLRAEDTAVYYCA LPRIAARRGGFGTWGQGLTVTVSS (SEQ ID NO: 352)	NFMLTQPHSVSESPGKTVTISCTRSS GNIDYNYVQWYQQRPGSSPTTVIYE DNVRPSGVPDRFSGSIDSSNSASLTI SGLKTEDEADYYCQSYDYNQAVV FGGGTKLTVL (SEQ ID NO: 353)
SR-AB60	QVQLVESGGGVVQPGRSLRLSCAAS GFTFRSYVMHWVRQAPGKGLEWV AVISHEGSGKYYADSVKGRFTISR NSKNTLYLQMNSLRAEDTAVYYCA LPRIAARRGGFGTWGQGLTVTVSS (SEQ ID NO: 354)	NFMLTQPHSVSESPGKTVTISCTRSS GNIDYNYVQWYQQRPGSSPTTVIYE DNVRPSGVPDRFSGSIDSSNSASLTI SGLKTEDEADYYCQSYDYDTQGVV FGGGTKLTVL (SEQ ID NO: 355)
SR-AB61	QVQLVESGGGVVQPGRSLRLSCAAS GFTFRSYVMHWVRQAPGKGLEWV AVISHEGSGKYYADSVKGRFTISR	NFMLTQPHSVSESPGKTVTISCTRSS GNIDYNYVQWYQQRPGSSPTTVIYE DNVRPSGVPDRFSGSIDSSNSASLTI

	NSKNTLYLQMNSLRAEDTAVYYCALPRIAARRGGFGTWGQGTTLVTVSS (SEQ ID NO: 356)	SGLKTEDEADYYCQGYDWDVTQGVVFGGGTKLTVL (SEQ ID NO: 357)
SR-AB62	QLQLQESGPGGLAKPSETLSLTCTVSG GSIRSSSYWGWIRQPPGKGLEWIG SISYSATTYYNPSLKSRTISVDTSK NQFSLKLSSVTAADTAVYYCASDPS YDSAAGMDVWGQGTTVTVSS (SEQ ID NO: 358)	DIQMTQSPSSLSASVGDRVTITCRAS KVISSYLNWYQQKPGKAPKLLIYYA SSLQSGVPSRFSGSGSGTDFTLTISL QPEDFATYYCQQSNDWPFTFGGGT KVEIK (SEQ ID NO: 359)
SR-AB63	QLQLQESGPGGLVKPSETLSLTCTVSG GSIRSSSYWGWIRQPPGKGLEWIG SISYSATTYYNPSLKSRTISVDTSK NQFSLKLSSVTAADTAVYYCAGDPS YDSIAGMQVWGQGTTVTVSS (SEQ ID NO: 360)	DIQLTQSPSSLSASVGDRVTITCRAS QSISSYLNWYQQKPGKAPKLLIYAA SNLQSGVPSRFSGSGSGTDFTLTISL QPEDFATYYCQQSFDWPLTFGGGT KVEIK (SEQ ID NO: 361)
SR-AB64	QLQLQESGPGGLVKPSETLSLTCTVSG GSIRSSSYWGWIRQPPGKGLEWIG SISYSATTYYNPSLKSRTISVDTSK NQFSLKLSSVTAADTAVYYCAGDPS YDSIAGMQVWGQGTTVTVSS (SEQ ID NO: 362)	DIQMTQSPSSLSASVGDRVTITCRAS QSISSYLNWYQQKPGKAPKLLIYSA SSRQSGVPSRFSGSGSGTDFTLTISL QPEDFATYYCQGFDFPLTFGGGT KVEIK (SEQ ID NO: 363)

[245] Aspects of the invention relate to a monoclonal antibody, or antigen-binding portion thereof, that binds selectively to a LTBP1-TGF β complex and/or a LTBP3-TGF β complex, and that comprises a heavy chain variable region sequence and a light chain variable region sequence.

[246] In one aspect, the invention provides an isolated antibody or an antigen-binding fragment thereof, that specifically binds a human LTBP1-proTGF β complex and/or a human LTBP3-proTGF β complex and does not bind a human GARP-proTGF β 1 complex; wherein the antibody or the antigen-binding fragment thereof does not bind mature TGF β 1, mature TGF β 2 or mature TGF β 3; wherein the antibody or the antigen-binding fragment thereof is a fully human or humanized antibody or an antigen-binding fragment thereof; wherein the antibody or the antigen-binding fragment thereof comprises a variable heavy chain having an amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to any one of the variable region amino acid sequences set forth in Table 6.

[247] In one embodiment, the antibody, or antigen-binding fragment thereof, comprises a heavy chain variable region having an amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to SEQ ID NO: 7, SEQ ID NO: 74, SEQ ID NO: 76, SEQ ID NO: 78, SEQ ID NO: 80, SEQ ID NO: 82, SEQ ID NO: 84, SEQ ID NO: 86, SEQ ID NO: 88, SEQ ID NO: 90, SEQ ID NO: 92, or SEQ ID NO: 106. In one embodiment, the antibody, or antigen-binding fragment thereof, comprises a light chain variable region having an amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to SEQ ID NO: 8, SEQ ID NO: 75, SEQ ID NO: 77, SEQ ID NO: 79, SEQ ID NO: 81, SEQ ID NO: 83, SEQ ID NO: 85, SEQ ID NO: 87, SEQ ID NO: 89, SEQ ID NO: 91, SEQ ID NO: 93, or SEQ ID NO: 107.

[248] In one embodiment, the antibody, or antigen-binding fragment thereof, comprises a heavy chain variable region having an amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to SEQ ID NO: 7 and/or a light chain variable region having an amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to SEQ ID NO: 8. In one embodiment, the antibody, or antigen-binding fragment thereof, comprises a heavy chain variable region having an amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to SEQ ID NO: 7 or a light chain variable region having an amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to SEQ ID NO: 8. In one embodiment, the antibody, or antigen-binding fragment thereof, comprises a heavy chain variable region having an amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to SEQ ID NO: 7 and a light chain variable region having an amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to SEQ ID NO: 8.

[249] In one embodiment, the antibody, or antigen-binding fragment thereof, comprises a heavy chain variable region having an amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to SEQ ID NO: 74 and/or a light chain variable region having an amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to SEQ ID NO: 75. In one embodiment, the antibody, or antigen-binding fragment thereof, comprises a heavy chain variable region having an amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to SEQ ID NO: 74 or a light chain variable region having an amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to SEQ ID NO: 75. In one embodiment, the antibody, or antigen-binding fragment thereof, comprises a heavy chain variable region having an amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to SEQ ID NO: 74 and a light chain variable region having an amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to SEQ ID NO: 75.

[250] In one embodiment, the antibody, or antigen-binding fragment thereof, comprises a heavy chain variable region having an amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to SEQ ID NO: 76 and/or a light chain variable region having an amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to SEQ ID NO: 77. In one embodiment, the antibody, or antigen-binding fragment thereof, comprises a heavy chain variable region having an amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to SEQ ID NO: 76 or a light chain variable region having an amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to SEQ ID NO: 77. In one embodiment, the antibody, or antigen-binding fragment thereof, comprises a heavy chain variable region having an amino acid sequence that is at

least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to SEQ ID NO: 76 and a light chain variable region having an amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to SEQ ID NO: 77.

[251] In one embodiment, the antibody, or antigen-binding fragment thereof, comprises a heavy chain variable region having an amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to SEQ ID NO: 78 and/or a light chain variable region having an amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to SEQ ID NO: 79. In one embodiment, the antibody, or antigen-binding fragment thereof, comprises a heavy chain variable region having an amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to SEQ ID NO: 78 or a light chain variable region having an amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to SEQ ID NO: 79. In one embodiment, the antibody, or antigen-binding fragment thereof, comprises a heavy chain variable region having an amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to SEQ ID NO: 78 and a light chain variable region having an amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to SEQ ID NO: 79.

[252] In one embodiment, the antibody, or antigen-binding fragment thereof, comprises a heavy chain variable region having an amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to SEQ ID NO: 80 and/or a light chain variable region having an amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to SEQ ID NO: 81. In one embodiment, the antibody, or antigen-binding fragment thereof, comprises a heavy chain variable region having an amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to SEQ ID NO: 80 or a light chain variable region having an amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to SEQ ID NO: 81. In one embodiment, the antibody, or antigen-binding fragment thereof, comprises a heavy chain variable region having an amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to SEQ ID NO: 80 and a light chain variable region having an amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to SEQ ID NO: 81.

[253] In one embodiment, the antibody, or antigen-binding fragment thereof, comprises a heavy chain variable region having an amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to SEQ ID NO: 82 and/or a light chain variable region having an amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to SEQ ID NO: 83. In one embodiment, the antibody, or antigen-binding fragment thereof, comprises a heavy chain variable region having an amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to SEQ ID NO: 82 or a light chain variable

region having an amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to SEQ ID NO: 83. In one embodiment, the antibody, or antigen-binding fragment thereof, comprises a heavy chain variable region having an amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to SEQ ID NO: 82 and a light chain variable region having an amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to SEQ ID NO: 83.

[254] In one embodiment, the antibody, or antigen-binding fragment thereof, comprises a heavy chain variable region having an amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to SEQ ID NO: 84 and/or a light chain variable region having an amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to SEQ ID NO: 85. In one embodiment, the antibody, or antigen-binding fragment thereof, comprises a heavy chain variable region having an amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to SEQ ID NO: 84 or a light chain variable region having an amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to SEQ ID NO: 85. In one embodiment, the antibody, or antigen-binding fragment thereof, comprises a heavy chain variable region having an amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to SEQ ID NO: 84 and a light chain variable region having an amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to SEQ ID NO: 85.

[255] In one embodiment, the antibody, or antigen-binding fragment thereof, comprises a heavy chain variable region having an amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to SEQ ID NO: 86 and/or a light chain variable region having an amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to SEQ ID NO: 87. In one embodiment, the antibody, or antigen-binding fragment thereof, comprises a heavy chain variable region having an amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to SEQ ID NO: 86 or a light chain variable region having an amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to SEQ ID NO: 87. In one embodiment, the antibody, or antigen-binding fragment thereof, comprises a heavy chain variable region having an amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to SEQ ID NO: 86 and a light chain variable region having an amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to SEQ ID NO: 87.

[256] In one embodiment, the antibody, or antigen-binding fragment thereof, comprises a heavy chain variable region having an amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to SEQ ID NO: 88 and/or a light chain variable region having an amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99%

identical to SEQ ID NO: 89. In one embodiment, the antibody, or antigen-binding fragment thereof, comprises a heavy chain variable region having an amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to SEQ ID NO: 88 or a light chain variable region having an amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to SEQ ID NO: 89. In one embodiment, the antibody, or antigen-binding fragment thereof, comprises a heavy chain variable region having an amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to SEQ ID NO: 88 and a light chain variable region having an amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to SEQ ID NO: 89.

[257] In one embodiment, the antibody, or antigen-binding fragment thereof, comprises a heavy chain variable region having an amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to SEQ ID NO: 90 and/or a light chain variable region having an amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to SEQ ID NO: 91. In one embodiment, the antibody, or antigen-binding fragment thereof, comprises a heavy chain variable region having an amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to SEQ ID NO: 90 or a light chain variable region having an amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to SEQ ID NO: 91. In one embodiment, the antibody, or antigen-binding fragment thereof, comprises a heavy chain variable region having an amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to SEQ ID NO: 90 and a light chain variable region having an amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to SEQ ID NO: 91.

[258] In one embodiment, the antibody, or antigen-binding fragment thereof, comprises a heavy chain variable region having an amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to SEQ ID NO: 92 and/or a light chain variable region having an amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to SEQ ID NO: 93. In one embodiment, the antibody, or antigen-binding fragment thereof, comprises a heavy chain variable region having an amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to SEQ ID NO: 92 or a light chain variable region having an amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to SEQ ID NO: 93. In one embodiment, the antibody, or antigen-binding fragment thereof, comprises a heavy chain variable region having an amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to SEQ ID NO: 92 and a light chain variable region having an amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to SEQ ID NO: 93.

[259] In one embodiment, the antibody, or antigen-binding fragment thereof, comprises a heavy chain variable region having an amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to SEQ ID NO: 106 and/or a light chain variable region having an amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to SEQ ID NO: 107. In one embodiment, the antibody, or antigen-binding fragment thereof, comprises a heavy chain variable region having an amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to SEQ ID NO: 106 or a light chain variable region having an amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to SEQ ID NO: 107. In one embodiment, the antibody, or antigen-binding fragment thereof, comprises a heavy chain variable region having an amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to SEQ ID NO: 106 and a light chain variable region having an amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to SEQ ID NO: 107.

[260] In one aspect, the invention provides an isolated antibody or an antigen-binding fragment thereof, that specifically binds a human LTBP1-proTGF β complex and/or a human LTBP3-proTGF β complex. The antibody may selectively bind a human LTBP1-proTGF β complex and/or a human LTBP3-proTGF β complex. The antibody or the antigen-binding fragment thereof may not bind mature TGF β 1, mature TGF β 2 or mature TGF β 3. The antibody or the antigen-binding fragment thereof may be a fully human or humanized antibody or an antigen-binding fragment thereof. The antibody or the antigen-binding fragment thereof may comprises a variable heavy chain having an amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to any one of the variable region amino acid sequences set forth in Table 6. In some embodiments, the level of identity is at least 95% (optionally at least 98%).

[261] Accordingly, in one embodiment, the antibody, or antigen-binding fragment thereof, comprises a heavy chain variable region having an amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to SEQ ID NO: 318 and/or a light chain variable region having an amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to SEQ ID NO: 319. The antibody, or antigen-binding fragment thereof, may comprise a heavy chain variable region having an amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to SEQ ID NO: 318 or a light chain variable region having an amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to SEQ ID NO: 319. The antibody, or antigen-binding fragment thereof, may comprise a heavy chain variable region having an amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to SEQ ID NO: 318 and a light chain variable region having an amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to SEQ ID NO: 319.

[262] In another embodiment, the antibody, or antigen-binding fragment thereof, comprises a heavy chain variable region having an amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to SEQ ID NO: 360 and/or a light chain variable region having an amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to SEQ ID NO: 361. The antibody, or antigen-binding fragment thereof, may comprise a heavy chain variable region having an amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to SEQ ID NO: 360 or a light chain variable region having an amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to SEQ ID NO: 361. The antibody, or antigen-binding fragment thereof, may comprise a heavy chain variable region having an amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to SEQ ID NO: 360 and a light chain variable region having an amino acid sequence that is at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to SEQ ID NO: 361.

[263] In some embodiments, the heavy chain variable region and/or the light chain variable region sequences do not vary within any of the CDR sequences provided herein. For example, in some embodiments, the degree of sequence variation (*e.g.*, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99%) may occur within a heavy chain variable and/or a light chain variable amino acid sequence excluding any of the CDR sequences provided herein. In some embodiments, the antibody, or antigen-binding portion thereof, comprises a heavy chain variable domain comprising an amino acid sequence set forth in SEQ ID NO: 7 and a light chain variable domain comprising an amino acid sequence set forth in SEQ ID NO: 8. In some embodiments, the antibody, or antigen-binding fragment thereof, comprising a heavy chain variable region having an amino acid sequence that is at least 90% identical to SEQ ID NO: 318 and/or a light chain variable region having an amino acid sequence that is at least 90% identical to SEQ ID NO: 319 does not vary within any of the CDR sequences of Ab42 provided herein. In some embodiments, the antibody, or antigen-binding fragment thereof, comprising a heavy chain variable region having an amino acid sequence that is at least 90% identical to SEQ ID NO: 360 and/or a light chain variable region having an amino acid sequence that is at least 90% identical to SEQ ID NO: 361 does not vary within any of the CDR sequences of Ab63 provided herein.

[264] In some embodiments, the antibody, or antigen-binding portion thereof, comprises a heavy chain variable domain comprising an amino acid sequence set forth in Table 6, and/or a light chain variable domain comprising an amino acid sequence set forth in Table 6. For example, in some embodiments, the antibody, or antigen-binding portion thereof, comprises a heavy chain variable domain comprising an amino acid sequence set forth in SEQ ID NO: 6 and/or a light chain variable domain comprising an amino acid sequence set forth in SEQ ID NO: 7. In some embodiments, the antibody, or antigen-binding portion thereof, comprises a heavy chain variable domain comprising an

amino acid sequence set forth in SEQ ID NO: 6 or a light chain variable domain comprising an amino acid sequence set forth in SEQ ID NO: 7. In some embodiments, the antibody, or antigen-binding portion thereof, comprises a heavy chain variable domain comprising an amino acid sequence set forth in SEQ ID NO: 6 and a light chain variable domain comprising an amino acid sequence set forth in SEQ ID NO: 7.

[265] In some embodiments, the antibody, or antigen-binding portion thereof, comprises a heavy chain variable domain comprising an amino acid sequence set forth in SEQ ID NO: 74 and/or a light chain variable domain comprising an amino acid sequence set forth in SEQ ID NO: 75. In some embodiments, the antibody, or antigen-binding portion thereof, comprises a heavy chain variable domain comprising an amino acid sequence set forth in SEQ ID NO: 74 or a light chain variable domain comprising an amino acid sequence set forth in SEQ ID NO: 75. In some embodiments, the antibody, or antigen-binding portion thereof, comprises a heavy chain variable domain comprising an amino acid sequence set forth in SEQ ID NO: 74 and a light chain variable domain comprising an amino acid sequence set forth in SEQ ID NO: 75.

[266] In some embodiments, the antibody, or antigen-binding portion thereof, comprises a heavy chain variable domain comprising an amino acid sequence set forth in SEQ ID NO: 76 and/or a light chain variable domain comprising an amino acid sequence set forth in SEQ ID NO: 77. In some embodiments, the antibody, or antigen-binding portion thereof, comprises a heavy chain variable domain comprising an amino acid sequence set forth in SEQ ID NO: 76 or a light chain variable domain comprising an amino acid sequence set forth in SEQ ID NO: 77. In some embodiments, the antibody, or antigen-binding portion thereof, comprises a heavy chain variable domain comprising an amino acid sequence set forth in SEQ ID NO: 76 and a light chain variable domain comprising an amino acid sequence set forth in SEQ ID NO: 77.

[267] In some embodiments, the antibody, or antigen-binding portion thereof, comprises a heavy chain variable domain comprising an amino acid sequence set forth in SEQ ID NO: 78 and/or a light chain variable domain comprising an amino acid sequence set forth in SEQ ID NO: 79. In some embodiments, the antibody, or antigen-binding portion thereof, comprises a heavy chain variable domain comprising an amino acid sequence set forth in SEQ ID NO: 78 or a light chain variable domain comprising an amino acid sequence set forth in SEQ ID NO: 79. In some embodiments, the antibody, or antigen-binding portion thereof, comprises a heavy chain variable domain comprising an amino acid sequence set forth in SEQ ID NO: 78 and a light chain variable domain comprising an amino acid sequence set forth in SEQ ID NO: 79.

[268] In some embodiments, the antibody, or antigen-binding portion thereof, comprises a heavy chain variable domain comprising an amino acid sequence set forth in SEQ ID NO: 80 and/or a light chain variable domain comprising an amino acid sequence set forth in SEQ ID NO: 81. In some

embodiments, the antibody, or antigen-binding portion thereof, comprises a heavy chain variable domain comprising an amino acid sequence set forth in SEQ ID NO: 80 or a light chain variable domain comprising an amino acid sequence set forth in SEQ ID NO: 81. In some embodiments, the antibody, or antigen-binding portion thereof, comprises a heavy chain variable domain comprising an amino acid sequence set forth in SEQ ID NO: 80 and a light chain variable domain comprising an amino acid sequence set forth in SEQ ID NO: 81.

[269] In some embodiments, the antibody, or antigen-binding portion thereof, comprises a heavy chain variable domain comprising an amino acid sequence set forth in SEQ ID NO: 82 and/or a light chain variable domain comprising an amino acid sequence set forth in SEQ ID NO: 83. In some embodiments, the antibody, or antigen-binding portion thereof, comprises a heavy chain variable domain comprising an amino acid sequence set forth in SEQ ID NO: 82 or a light chain variable domain comprising an amino acid sequence set forth in SEQ ID NO: 83. In some embodiments, the antibody, or antigen-binding portion thereof, comprises a heavy chain variable domain comprising an amino acid sequence set forth in SEQ ID NO: 82 and a light chain variable domain comprising an amino acid sequence set forth in SEQ ID NO: 83.

[270] In some embodiments, the antibody, or antigen-binding portion thereof, comprises a heavy chain variable domain comprising an amino acid sequence set forth in SEQ ID NO: 84 and/or a light chain variable domain comprising an amino acid sequence set forth in SEQ ID NO: 85. In some embodiments, the antibody, or antigen-binding portion thereof, comprises a heavy chain variable domain comprising an amino acid sequence set forth in SEQ ID NO: 84 or a light chain variable domain comprising an amino acid sequence set forth in SEQ ID NO: 85. In some embodiments, the antibody, or antigen-binding portion thereof, comprises a heavy chain variable domain comprising an amino acid sequence set forth in SEQ ID NO: 84 and a light chain variable domain comprising an amino acid sequence set forth in SEQ ID NO: 85.

[271] In some embodiments, the antibody, or antigen-binding portion thereof, comprises a heavy chain variable domain comprising an amino acid sequence set forth in SEQ ID NO: 86 and/or a light chain variable domain comprising an amino acid sequence set forth in SEQ ID NO: 87. In some embodiments, the antibody, or antigen-binding portion thereof, comprises a heavy chain variable domain comprising an amino acid sequence set forth in SEQ ID NO: 86 or a light chain variable domain comprising an amino acid sequence set forth in SEQ ID NO: 87. In some embodiments, the antibody, or antigen-binding portion thereof, comprises a heavy chain variable domain comprising an amino acid sequence set forth in SEQ ID NO: 86 and a light chain variable domain comprising an amino acid sequence set forth in SEQ ID NO: 87.

[272] In some embodiments, the antibody, or antigen-binding portion thereof, comprises a heavy chain variable domain comprising an amino acid sequence set forth in SEQ ID NO: 88 and/or a light

chain variable domain comprising an amino acid sequence set forth in SEQ ID NO: 89. In some embodiments, the antibody, or antigen-binding portion thereof, comprises a heavy chain variable domain comprising an amino acid sequence set forth in SEQ ID NO: 88 or a light chain variable domain comprising an amino acid sequence set forth in SEQ ID NO: 89. In some embodiments, the antibody, or antigen-binding portion thereof, comprises a heavy chain variable domain comprising an amino acid sequence set forth in SEQ ID NO: 88 and a light chain variable domain comprising an amino acid sequence set forth in SEQ ID NO: 89.

[273] In some embodiments, the antibody, or antigen-binding portion thereof, comprises a heavy chain variable domain comprising an amino acid sequence set forth in SEQ ID NO: 90 and/or a light chain variable domain comprising an amino acid sequence set forth in SEQ ID NO: 91. In some embodiments, the antibody, or antigen-binding portion thereof, comprises a heavy chain variable domain comprising an amino acid sequence set forth in SEQ ID NO: 90 or a light chain variable domain comprising an amino acid sequence set forth in SEQ ID NO: 91. In some embodiments, the antibody, or antigen-binding portion thereof, comprises a heavy chain variable domain comprising an amino acid sequence set forth in SEQ ID NO: 90 and a light chain variable domain comprising an amino acid sequence set forth in SEQ ID NO: 91.

[274] In some embodiments, the antibody, or antigen-binding portion thereof, comprises a heavy chain variable domain comprising an amino acid sequence set forth in SEQ ID NO: 92 and/or a light chain variable domain comprising an amino acid sequence set forth in SEQ ID NO: 93. In some embodiments, the antibody, or antigen-binding portion thereof, comprises a heavy chain variable domain comprising an amino acid sequence set forth in SEQ ID NO: 92 or a light chain variable domain comprising an amino acid sequence set forth in SEQ ID NO: 93. In some embodiments, the antibody, or antigen-binding portion thereof, comprises a heavy chain variable domain comprising an amino acid sequence set forth in SEQ ID NO: 92 and a light chain variable domain comprising an amino acid sequence set forth in SEQ ID NO: 93.

[275] In some embodiments, the antibody, or antigen-binding portion thereof, comprises a heavy chain variable domain comprising an amino acid sequence set forth in SEQ ID NO: 106 and/or a light chain variable domain comprising an amino acid sequence set forth in SEQ ID NO: 107. In some embodiments, the antibody, or antigen-binding portion thereof, comprises a heavy chain variable domain comprising an amino acid sequence set forth in SEQ ID NO: 106 or a light chain variable domain comprising an amino acid sequence set forth in SEQ ID NO: 107. In some embodiments, the antibody, or antigen-binding portion thereof, comprises a heavy chain variable domain comprising an amino acid sequence set forth in SEQ ID NO: 106 and a light chain variable domain comprising an amino acid sequence set forth in SEQ ID NO: 107.

[276] In some embodiments, the antibody, or antigen-binding portion thereof, comprises a heavy chain variable domain comprising an amino acid sequence set forth in SEQ ID NO: 318 and/or a light chain variable domain comprising an amino acid sequence set forth in SEQ ID NO: 319. In some embodiments, the antibody, or antigen-binding portion thereof, comprises a heavy chain variable domain comprising an amino acid sequence set forth in SEQ ID NO: 318 or a light chain variable domain comprising an amino acid sequence set forth in SEQ ID NO: 319. In some embodiments, the antibody, or antigen-binding portion thereof, comprises a heavy chain variable domain comprising an amino acid sequence set forth in SEQ ID NO: 318 and a light chain variable domain comprising an amino acid sequence set forth in SEQ ID NO: 319.

[277] In some embodiments, the antibody, or antigen-binding portion thereof, comprises a heavy chain variable domain comprising an amino acid sequence set forth in SEQ ID NO: 360 and/or a light chain variable domain comprising an amino acid sequence set forth in SEQ ID NO: 361. In some embodiments, the antibody, or antigen-binding portion thereof, comprises a heavy chain variable domain comprising an amino acid sequence set forth in SEQ ID NO: 360 or a light chain variable domain comprising an amino acid sequence set forth in SEQ ID NO: 361. In some embodiments, the antibody, or antigen-binding portion thereof, comprises a heavy chain variable domain comprising an amino acid sequence set forth in SEQ ID NO: 360 and a light chain variable domain comprising an amino acid sequence set forth in SEQ ID NO: 361.

[278] The amino acid sequences of the heavy chain variable region (HCVR) and the light chain variable region (LCVR) of the antibody SR-AB2 set forth in Table 5 are provided below.

[279] SR-AB2 – Heavy chain variable region amino acid sequence

QVQLVQSGAEVKKPGASVKVSCKASGYTFTSYGISWVRQAPGQGLEWMGWISAYNGNTNY
AQKLQGRVTMTTDTSTSTAYMELRSLRSDDTAVYYCARAPLGNFDSWGQGTMTVTSS
(SEQ ID NO: 7)

[280] SR-AB2 – Light chain variable region amino acid sequence

NFMLTQPHSVSESPGKTVTISCTRSSGSIASNYVQWYQQRPGSSPTTVIYEDNQRPSGVPDRFS
GSIDSSSNSASLTISGLKTEDEADYYCQSYDSSNHPVVFGGGTKLTVL (SEQ ID NO: 8)

[281] The amino acid sequences of the heavy chain variable region (HCVR) and the light chain variable region (LCVR) of the antibody SR-AB10 set forth in Table 5 are provided below.

[282] SR-AB10 – Heavy chain variable region amino acid sequence

QLQLQESGGGVVQPGRSLRLSCAASGFTFNYPHIVWRQAPGKGLEWVAVMSYDGINKYY
ADSVKGRFTISRDNKNTLYLQMNSLRAEDTAVYYCARPRIAARRGGFDYWGQGTTLTVSS
(SEQ ID NO: 88)

[283] SR-AB10 – Light chain variable region amino acid sequence

NFMLTQPHSVSESPGKTVTISCTRSSGNIDNNYVQWYQQRPGSSPTTVIYEDNQRPSPGVDRF
SGSIDSSSNSASLTISGLKTEDEADYYCQSYSDSNQGVVFGGGTKLTVL (SEQ ID NO: 89)

[284] The amino acid sequences of the heavy chain variable region (HCVR) and the light chain variable region (LCVR) of the antibody SR-AB13 set forth in Table 5 are provided below.

[285] SR-AB13 – Heavy chain variable region amino acid sequence

QLQLQESGPGLVKPSETLSLTCTVSGGSISSSSYYWGWIRQPPGKGLEWIGSISYSGSTYYNPS
LKSRVTISVDTSKNQFSLKLSSVTAADTAVYYCARDPSYDSIAGMDVWGQGTTVTVSS (SEQ
ID NO: 106)

[286] SR-AB13 – Light chain variable region amino acid sequence

DIQLTQSPSSLSASVGDRVTITCRASQSISSYLNWYQQKPGKAPKLLIYAASNLSQGVPSRFSG
SGSGTDFTLTISLQPEDFATYYCQQSFDFPFTFGGGTKVEIK (SEQ ID NO: 107)

[287] The amino acid sequences of the heavy chain variable region (HCVR) and the light chain variable region (LCVR) of the antibody SR-AB42 set forth in Table 5 are provided below.

[288] SR-AB42 – Heavy chain variable region amino acid sequence

QVQLVESGGGVVQPGRSLRLSCAASGFTFRSYVMHWVRQAPGKGLEWVAVISHEGSLKYY
ADSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYYCARPRIAARRGGFGYWGQGTTLTVSS
(SEQ ID NO: 318)

[289] 63 – Light chain variable region amino acid sequence

NFMLTQPHSVSESPGKTVTISCTRSSGNIDNNYVQWYQQRPGSSPTTVIYEDNQRPSPGVDRF
SGSIDSSSNSASLTISGLKTEDEADYYCQSYDYDTQGVVFGGGTKLTVL (SEQ ID NO: 319)

[290] The amino acid sequences of the heavy chain variable region (HCVR) and the light chain variable region (LCVR) of the antibody SR-AB63 set forth in Table 5 are provided below.

[291] SR-AB63 – Heavy chain variable region amino acid sequence

QLQLQESGPGLVKPSETLSLTCTVSGGSIRSSSYWGWIRQPPGKGLEWIGSISYSATTYYNPS
LKSRVTISVDTSKNQFSLKLSSVTAADTAVYYCAGDPSYDSIAGMQVWGQGTTVTVSS (SEQ
ID NO: 360)

[292] SR-AB63 – Light chain variable region amino acid sequence

DIQLTQSPSSLSASVGDRVTITCRASQSISSYLNWYQQKPGKAPKLLIYAASNLSQGVPSRFSG
SGSGTDFTLTISLQPEDFATYYCQQSFDWPLTFGGGTKEIK (SEQ ID NO: 361)

[293] In some embodiments, antibodies, or antigen-binding portions thereof, of the invention that selectively bind to a LTBP1-TGF β complex and/or a LTBP3-TGF β complex have one or more CDR sequences substantially similar to CDRH1, CDRH2, CDRH3, CDRL1, CDRL2, and/or CDRL3. For example, the antibodies may include one or more CDR sequences as shown in Table 5 (SEQ ID NOs: 1-6, SEQ ID NOs: 94-99 or SEQ ID NOs: 100-105) containing up to 6, 5, 4, 3, 2, or 1 amino acid residue variations as compared to the corresponding CDR region in any one of SEQ ID NOs: 1-6, SEQ ID NOs: 94-99 or SEQ ID NOs: 100-105.

[294] In one embodiment, the antibody, or antigen-binding fragment thereof, comprises at least three CDRs selected from the following, optionally comprising up to 6 amino acid changes, for example 1, 2, 3, 4, 5, or 6 amino acid changes, for each of the CDRs CDR-H1: SEQ ID NO: 1; CDR-H2: SEQ ID NO: 2; CDR-H3: SEQ ID NO: 3; CDR-L1: SEQ ID NO: 4; CDR-L2: SEQ ID NO: 5; and, CDR-L3: SEQ ID NO: 6. In one embodiment, the antibody, or antigen-binding fragment thereof, comprises at least three CDRs selected from the following, optionally comprising up to 6 amino acid changes, for example 1, 2, 3, 4, 5, or 6 amino acid changes, for each of the CDRs CDR-H1: SEQ ID NO: 94; CDR-H2: SEQ ID NO: 95; CDR-H3: SEQ ID NO: 96; CDR-L1: SEQ ID NO: 97; CDR-L2: SEQ ID NO: 98; and, CDR-L3: SEQ ID NO: 99. In one embodiment, the antibody, or antigen-binding fragment thereof, comprises at least three CDRs selected from the following, optionally comprising up to 6 amino acid changes, for example 1, 2, 3, 4, 5, or 6 amino acid changes, for each of the CDRs CDR-H1: SEQ ID NO: 100; CDR-H2: SEQ ID NO: 101; CDR-H3: SEQ ID NO: 102; CDR-L1: SEQ ID NO: 103; CDR-L2: SEQ ID NO: 104; and, CDR-L3: SEQ ID NO: 105. In one embodiment, the antibody, or antigen-binding fragment thereof, comprises at least three CDRs selected from the following, optionally comprising up to 6 amino acid changes, for example 1, 2, 3, 4, 5, or 6 amino acid changes, for each of the CDRs CDR-H1: SEQ ID NO: 166; CDR-H2: SEQ ID NO: 167; CDR-H3: SEQ ID NO: 168; CDR-L1: SEQ ID NO: 169; CDR-L2: SEQ ID NO: 170; and, CDR-L3: SEQ ID NO: 171. In one embodiment, the antibody, or antigen-binding fragment thereof, comprises at least three CDRs selected from the following, optionally comprising up to 6 amino acid changes, for example 1, 2, 3, 4, 5, or 6 amino acid changes, for each of the CDRs CDR-H1: SEQ ID NO: 292; CDR-H2: SEQ ID NO: 293; CDR-H3: SEQ ID NO: 294; CDR-L1: SEQ ID NO: 295; CDR-L2: SEQ ID NO: 296; and, CDR-L3: SEQ ID NO: 297.

[295] In one aspect, the invention provides an antibody, or antigen-binding fragment thereof, comprising a heavy chain variable region comprising CDR-H1: SEQ ID NO: 1; CDR-H2: SEQ ID NO: 2; and CDR-H3: SEQ ID NO: 3; and a light chain variable region comprising CDR-L1: SEQ ID NO: 4; CDR-L2: SEQ ID NO: 5; and CDR-L3: SEQ ID NO: 6, optionally comprising one or more amino acid changes, for example 1, 2, 3, 4, 5, or 6 amino acid changes, for each of the CDRs.

[296] In one aspect, the invention provides an antibody, or antigen-binding fragment thereof, comprising a heavy chain variable region comprising CDR-H1: SEQ ID NO: 94; CDR-H2: SEQ ID

NO: 95; and CDR-H3: SEQ ID NO: 96; and a light chain variable region comprising CDR-L1: SEQ ID NO: 97; CDR-L2: SEQ ID NO: 98; and CDR-L3: SEQ ID NO: 99, optionally comprising one or more amino acid changes, for example 1, 2, 3, 4, 5, or 6 amino acid changes, for each of the CDRs.

[297] In one aspect, the invention provides an antibody, or antigen-binding fragment thereof, comprising a heavy chain variable region comprising CDR-H1: SEQ ID NO: 100; CDR-H2: SEQ ID NO: 101; and CDR-H3: SEQ ID NO: 102; and a light chain variable region comprising CDR-L1: SEQ ID NO: 103; CDR-L2: SEQ ID NO: 104; and CDR-L3: SEQ ID NO: 105, optionally comprising one or more amino acid changes, for example 1, 2, 3, 4, 5, or 6 amino acid changes, for each of the CDRs.

[298] In one aspect, the invention provides an antibody, or antigen-binding fragment thereof, comprising a heavy chain variable region comprising CDR-H1: SEQ ID NO: 166; CDR-H2: SEQ ID NO: 167; and CDR-H3: SEQ ID NO: 168; and a light chain variable region comprising CDR-L1: SEQ ID NO: 169; CDR-L2: SEQ ID NO: 170; and CDR-L3: SEQ ID NO: 171, optionally comprising one or more amino acid changes, for example 1, 2, 3, 4, 5, or 6 amino acid changes, for each of the CDRs. For instance, if there are changes within the CDRs, there may be up to 1 change per CDR. There may be no more than 2 changes across all 6 CDRs.

[299] In one aspect, the invention provides an antibody, or antigen-binding fragment thereof, comprising a heavy chain variable region comprising CDR-H1: SEQ ID NO: 292; CDR-H2: SEQ ID NO: 293; and CDR-H3: SEQ ID NO: 294; and a light chain variable region comprising CDR-L1: SEQ ID NO: 295; CDR-L2: SEQ ID NO: 296; and CDR-L3: SEQ ID NO: 297, optionally comprising one or more amino acid changes, for example 1, 2, 3, 4, 5, or 6 amino acid changes (*e.g.*, up to 2), for each of the CDRs. For instance, if there are changes within the CDRs, there may be up to 1 change per CDR. There may be no more than 2 changes across all 6 CDRs.

[300] In one aspect, the invention provides an antibody, or antigen-binding fragment thereof comprising a heavy chain variable region comprising a CDR-H1, CDR-H2, CDR-H3, CDR-L1, CDR-L2, and/or CDR-L3 having particular amino acid changes. As used herein, the phrase “amino acid changes” or “changes in amino acid residues” includes amino acid substitutions, additions, and/or deletions. In some embodiments, there are one or more changes to the amino acid residues with any one of the CDRs and/or variable regions described herein. For example, in some embodiments, the one or more amino acid changes comprises one amino acid change. In some embodiments, the one or more amino acid changes comprises up to two amino acid changes. In some embodiments, the one or more amino acid changes comprises up to three amino acid changes. In some embodiments, the one or more amino acid changes comprises up to four amino acid changes. In some embodiments, the one or more amino acid changes comprises up to five amino acid changes. In some embodiments, the one

or more amino acid changes comprises up to six amino acid changes. In some embodiments, the one or more amino acid changes comprises up to seven amino acid changes.

[301] For example, in some embodiments, the antibody, or antigen-binding fragment thereof, comprises a CDR-H1: SEQ ID NO:1, with the proviso that the threonine residue at position 4 of SEQ ID NO:1 may be substituted with a histidine, lysine, phenylalanine, or glycine. In some embodiments, the antibody, or antigen-binding fragment thereof, comprises a CDR-H1: SEQ ID NO:1, with the proviso that the serine residue at position 5 of SEQ ID NO:1 may be substituted with a leucine. In some embodiments, the antibody, or antigen-binding fragment thereof, comprises a CDR-H1: SEQ ID NO:1, with the proviso that the serine residue at position 9 of SEQ ID NO:1 may be substituted with an alanine.

[302] In some embodiments, the antibody, or antigen-binding fragment thereof, comprises a CDR-H1: SEQ ID NO:1, with the proviso that (i) the threonine residue at position 4 of SEQ ID NO:1 may be substituted with a histidine, lysine, phenylalanine, or glycine; (ii) the serine residue at position 5 of SEQ ID NO:1 may be substituted with a leucine; and/or, (iii) the serine residue at position 9 of SEQ ID NO:1 may be substituted with an alanine.

[303] In some embodiments, the antibody, or antigen-binding fragment thereof, comprises a CDR-H2: SEQ ID NO:2, with the proviso that the serine residue at position 3 of SEQ ID NO:2 may be substituted with an aspartate or asparagine. In some embodiments, the antibody, or antigen-binding fragment thereof, comprises a CDR-H2: SEQ ID NO:2, with the proviso that the tyrosine residue at position 5 of SEQ ID NO:2 may be substituted with a histidine. In some embodiments, the antibody, or antigen-binding fragment thereof, comprises a CDR-H2: SEQ ID NO:2, with the proviso that the asparagine residue at position 6 of SEQ ID NO:2 may be substituted with a serine. In some embodiments, the antibody, or antigen-binding fragment thereof, comprises a CDR-H2: SEQ ID NO:2, with the proviso that the asparagine residue at position 8 of SEQ ID NO:2 may be substituted with a phenylalanine, leucine, alanine, tyrosine, aspartate, or serine. In some embodiments, the antibody, or antigen-binding fragment thereof, comprises a CDR-H2: SEQ ID NO:2, with the proviso that the asparagine residue at position 10 of SEQ ID NO:2 may be substituted with an aspartate or alanine.

[304] In some embodiments, the antibody, or antigen-binding fragment thereof, comprises a CDR-H2: SEQ ID NO:2, with the proviso that (i) the serine residue at position 3 of SEQ ID NO:2 may be substituted with an aspartate or asparagine; (ii) the tyrosine residue at position 5 of SEQ ID NO:2 may be substituted with a histidine; (iii) the asparagine residue at position 6 of SEQ ID NO:2 may be substituted with a serine; (iv) the asparagine residue at position 8 of SEQ ID NO:2 may be substituted with a phenylalanine, leucine, alanine, tyrosine, aspartate, or serine; and/or, (v) the asparagine residue at position 10 of SEQ ID NO:2 may be substituted with a aspartate or alanine.

[305] In some embodiments, the antibody, or antigen-binding fragment thereof, comprises three heavy chain CDRs and three light chain CDRs, wherein the heavy chain CDRs comprise:

- a) CDR-H1: SEQ ID NO:1, with the proviso that:
 - i. the threonine residue at position 4 of SEQ ID NO:1 may be substituted with a histidine, lysine, phenylalanine, or glycine;
 - ii. the serine residue at position 5 of SEQ ID NO:1 may be substituted with a leucine; and/or,
 - iii. the serine residue at position 9 of SEQ ID NO:1 may be substituted with an alanine;
- b) CDR-H2: SEQ ID NO:2, with the proviso that:
 - i. the serine residue at position 3 of SEQ ID NO:2 may be substituted with an aspartate or asparagine;
 - ii. the tyrosine residue at position 5 of SEQ ID NO:2 may be substituted with a histidine;
 - iii. the asparagine residue at position 6 of SEQ ID NO:2 may be substituted with a serine;
 - iv. the asparagine residue at position 8 of SEQ ID NO:2 may be substituted with a phenylalanine, leucine, alanine, tyrosine, aspartate, or serine; and/or,
 - v. the asparagine residue at position 10 of SEQ ID NO:2 may be substituted with a aspartate or alanine;
- c) CDR-H3: SEQ ID NO:3, optionally comprising one or more amino acid changes.

[306] In some embodiments, the antibody, or antigen-binding fragment thereof, further comprises a CDR-L1 as set forth in SEQ ID NO:4, optionally comprising one or more amino acid changes. In some embodiments, the antibody, or antigen-binding fragment thereof, further comprises a CDR-L2 as set forth in SEQ ID NO:5, optionally comprising one or more amino acid changes. In some embodiments, the antibody, or antigen-binding fragment thereof, further comprises a CDR-L3 as set forth in SEQ ID NO:6, optionally comprising one or more amino acid changes.

[307] In some embodiments, the antibody, or antigen-binding fragment thereof, comprises at least three of the following six CDRs:

- a) CDR-H1: SEQ ID NO:1, with the proviso that:
 - i. the threonine residue at position 4 of SEQ ID NO:1 may be substituted with a histidine, lysine, phenylalanine, or glycine;
 - ii. the serine residue at position 5 of SEQ ID NO:1 may be substituted with an leucine; and/or,
 - iii. the serine residue at position 9 of SEQ ID NO:1 may be substituted with an alanine;

- b) CDR-H2: SEQ ID NO:2, with the proviso that:
 - i. the serine residue at position 3 of SEQ ID NO:2 may be substituted with an aspartate or asparagine;
 - ii. the tyrosine residue at position 5 of SEQ ID NO:2 may be substituted with a histidine;
 - iii. the asparagine residue at position 6 of SEQ ID NO:2 may be substituted with a serine;
 - iv. the asparagine residue at position 8 of SEQ ID NO:2 may be substituted with a phenylalanine, leucine, alanine, tyrosine, aspartate, or serine; and/or,
 - v. the asparagine residue at position 10 of SEQ ID NO:2 may be substituted with a aspartate or alanine;
- c) CDR-H3: SEQ ID NO:3, optionally comprising one or more amino acid changes;
- d) CDR-L1: SEQ ID NO:4, optionally comprising one or more amino acid changes;
- e) CDR-L2: SEQ ID NO:5, optionally comprising one or more amino acid changes; and,
- f) CDR-L3: SEQ ID NO:6, optionally comprising one or more amino acid changes.

[308] In some embodiments, the antibody, or antigen-binding fragment thereof, specifically binds a human LTBP1-proTGFβ1 complex and/or a human LTBP3-proTGFβ1 complex, and does not bind a human GARP-proTGFβ1 complex. In some embodiments, the antibody, or antigen-binding fragment thereof, does not bind a human GARP-proTGFβ1 complex. In some embodiments, the antibody, or antigen-binding fragment thereof, does not bind mature TGFβ1, mature TGFβ2 or mature TGFβ3.

[309] In a particular embodiment, the invention provides an isolated antibody that specifically binds a human LTBP1-proTGFβ1 complex and/or a human LTBP3-proTGFβ1 complex, and does not bind a human GARP-proTGFβ1 complex or a human LRRC33-proTGFβ1 complex; wherein the antibody does not bind mature TGFβ1, mature TGFβ2 or mature TGFβ3; wherein the antibody is a fully human or humanized antibody or a fragment thereof, and wherein the antibody comprises at least three of the following six CDRs:

- a) CDR-H1: SEQ ID NO:1, with the proviso that:
 - i. the threonine residue at position 4 of SEQ ID NO:1 may be substituted with a histidine, lysine, phenylalanine, or glycine;
 - ii. the serine residue at position 5 of SEQ ID NO:1 may be substituted with an leucine; and/or,
 - iii. the serine residue at position 9 of SEQ ID NO:1 may be substituted with an alanine;
- b) CDR-H2: SEQ ID NO:2, with the proviso that:
 - i. the serine residue at position 3 of SEQ ID NO:2 may be substituted with an aspartate or asparagine;

- ii. the tyrosine residue at position 5 of SEQ ID NO:2 may be substituted with a histidine;
- iii. the asparagine residue at position 6 of SEQ ID NO:2 may be substituted with a serine;
- iv. the asparagine residue at position 8 of SEQ ID NO:2 may be substituted with a phenylalanine, leucine, alanine, tyrosine, aspartate, or serine; and/or,
- v. the asparagine residue at position 10 of SEQ ID NO:2 may be substituted with a aspartate or alanine;
- c) CDR-H3: SEQ ID NO:3, optionally comprising one or more amino acid changes;
- d) CDR-L1: SEQ ID NO:4, optionally comprising one or more amino acid changes;
- e) CDR-L2: SEQ ID NO:5, optionally comprising one or more amino acid changes; and,
- f) CDR-L3: SEQ ID NO:6, optionally comprising one or more amino acid changes.

[310] In some embodiments, such antibody binds a human LTBP1-proTGF β 1 complex with a K_D of < 100 nM as measured in a suitable in vitro binding assay such as Bio-Layer Interferometry (BLI); and/or, such antibody binds a human LTBP3-proTGF β 1 complex with a K_D of < 100 nM as measured in a suitable in vitro binding assay such as Bio-Layer Interferometry (BLI). In some embodiments, such antibody binds a human LTBP1-proTGF β 1 complex with a K_D of < 50 nM as measured in a suitable in vitro binding assay such as Bio-Layer Interferometry (BLI); and/or, such antibody binds a human LTBP3-proTGF β 1 complex with a K_D of < 50 nM as measured in a suitable in vitro binding assay such as Bio-Layer Interferometry (BLI). In some embodiments, such antibody binds a human LTBP1-proTGF β 1 complex with a K_D of < 25 nM as measured in a suitable in vitro binding assay such as Bio-Layer Interferometry (BLI); and/or, such antibody binds a human LTBP3-proTGF β 1 complex with a K_D of < 25 nM as measured in a suitable in vitro binding assay such as Bio-Layer Interferometry (BLI). In some embodiments, such antibody binds a human LTBP1-proTGF β 1 complex with a K_D of < 10 nM as measured in a suitable in vitro binding assay such as Bio-Layer Interferometry (BLI); and/or, such antibody binds a human LTBP3-proTGF β 1 complex with a K_D of < 10 nM as measured in a suitable in vitro binding assay such as Bio-Layer Interferometry (BLI).

[311] In another embodiment, such antibody is cross-reactive with mouse LTBP1-proTGF β 1. In some embodiments, such antibody is also cross-reactive with mouse LTBP3-proTGF β 1. In some embodiments, such antibody binds a mouse LTBP1-proTGF β 1 complex with a K_D of < 100 nM as measured in a suitable in vitro binding assay such as Bio-Layer Interferometry (BLI); and/or, the antibody binds a mouse LTBP3-proTGF β 1 complex with a K_D of < 100 nM as measured in a suitable in vitro binding assay such as Bio-Layer Interferometry (BLI). In some embodiments, such antibody binds a mouse LTBP1-proTGF β 1 complex with a K_D of < 50 nM as measured in a suitable in vitro binding assay such as Bio-Layer Interferometry (BLI); and/or, the antibody binds a mouse LTBP3-

proTGFβ1 complex with a K_D of < 50 nM as measured in a suitable in vitro binding assay such as Bio-Layer Interferometry (BLI). In some embodiments, such antibody binds a mouse LTBP1-proTGFβ1 complex with a K_D of < 25 nM as measured in a suitable in vitro binding assay such as Bio-Layer Interferometry (BLI); and/or, the antibody binds a mouse LTBP3-proTGFβ1 complex with a K_D of < 25 nM as measured in a suitable in vitro binding assay such as Bio-Layer Interferometry (BLI). In some embodiments, such antibody binds a mouse LTBP1-proTGFβ1 complex with a K_D of < 10 nM as measured in a suitable in vitro binding assay such as Bio-Layer Interferometry (BLI); and/or, the antibody binds a mouse LTBP3-proTGFβ1 complex with a K_D of < 10 nM as measured in a suitable in vitro binding assay such as Bio-Layer Interferometry (BLI).

[312] In another embodiment, such antibody does not bind to human GARP-proTGFβ1. In preferred embodiments, such context-selective antibody is isoform-specific in that it selectively binds and inhibits the activation of TGFβ1 associated with LTBP1/3 and does not bind to human GARP-proTGFβ1.

[313] In another aspect, the antibody, or antigen-binding fragment thereof, comprises a CDR-H1: SEQ ID NO:94, with the proviso that the threonine residue at position 2 of SEQ ID NO:94 may be substituted with an alanine. In some embodiments, the antibody, or antigen-binding fragment thereof, comprises a CDR-H1: SEQ ID NO:94, with the proviso that the asparagine residue at position 4 of SEQ ID NO:94 may be substituted with an alanine, tyrosine, aspartate, serine, arginine, or histidine. In some embodiments, the antibody, or antigen-binding fragment thereof, comprises a CDR-H1: SEQ ID NO:94, with the proviso that the asparagine residue at position 5 of SEQ ID NO:94 may be substituted with a glutamine, serine, glycine, lysine, glutamate, arginine, or histidine. In some embodiments, the antibody, or antigen-binding fragment thereof, comprises a CDR-H1: SEQ ID NO:94, with the proviso that the tyrosine residue at position 6 of SEQ ID NO:94 may be substituted with an arginine. In some embodiments, the antibody, or antigen-binding fragment thereof, comprises a CDR-H1: SEQ ID NO:94, with the proviso that the proline residue at position 7 of SEQ ID NO:94 may be substituted with a glycine, alanine, leucine, serine, asparagine, valine, aspartate, or glutamine. In some embodiments, the antibody, or antigen-binding fragment thereof, comprises a CDR-H1: SEQ ID NO:94, with the proviso that the isoleucine residue at position 8 of SEQ ID NO:94 may be substituted with a methionine or leucine. In some embodiments, the antibody, or antigen-binding fragment thereof, comprises a CDR-H1: SEQ ID NO:94, with the proviso that the histidine residue at position 9 of SEQ ID NO:94 may be substituted with a phenylalanine, tyrosine, asparagine, or serine.

[314] In some embodiments, the antibody, or antigen-binding fragment thereof, comprises a CDR-H1: SEQ ID NO:94, with the proviso that (i) the threonine residue at position 2 of SEQ ID NO:94 may be substituted with an alanine; (ii) the asparagine residue at position 4 of SEQ ID NO:94 may be substituted with an alanine, tyrosine, aspartate, serine, arginine, or histidine; (iii) the asparagine

residue at position 5 of SEQ ID NO:94 may be substituted with a glutamine, serine, glycine, lysine, glutamate, arginine, or histidine; (iv) the tyrosine residue at position 6 of SEQ ID NO:94 may be substituted with a arginine; (v) the proline residue at position 7 of SEQ ID NO:94 may be substituted with a glycine, alanine, leucine, serine, asparagine, valine, aspartate, or glutamine; (vi) the isoleucine residue at position 8 of SEQ ID NO:94 may be substituted with a methionine or leucine; and/or, (vii) the histidine residue at position 9 of SEQ ID NO:94 may be substituted with a phenylalanine, tyrosine, asparagine, or serine.

[315] In some embodiments, the antibody, or antigen-binding fragment thereof, comprises three heavy chain CDRs and three light chain CDRs, wherein the heavy chain CDRs comprise:

- a) CDR-H1: SEQ ID NO:94, with the proviso that:
 - i. the threonine residue at position 2 of SEQ ID NO:94 may be substituted with an alanine;
 - ii. the asparagine residue at position 4 of SEQ ID NO:94 may be substituted with an alanine, tyrosine, aspartate, serine, arginine, or histidine;
 - iii. the asparagine residue at position 5 of SEQ ID NO:94 may be substituted with a glutamine, serine, glycine, lysine, glutamate, arginine, or histidine;
 - iv. the tyrosine residue at position 6 of SEQ ID NO:94 may be substituted with a arginine;
 - v. the proline residue at position 7 of SEQ ID NO:94 may be substituted with a glycine, alanine, leucine, serine, asparagine, valine, aspartate, or glutamine;
 - vi. the isoleucine residue at position 8 of SEQ ID NO:94 may be substituted with a methionine or leucine; and/or,
 - vii. the histidine residue at position 9 of SEQ ID NO:94 may be substituted with a phenylalanine, tyrosine, asparagine, or serine;
- b) CDR-H2: SEQ ID NO:95, optionally comprising one or more amino acid changes; and
- c) CDR-H3: SEQ ID NO:96, optionally comprising one or more amino acid changes.

[316] In some embodiments, the antibody, or antigen-binding fragment thereof, further comprises a CDR-L1 as set forth in SEQ ID NO:97, optionally comprising one or more amino acid changes. In some embodiments, the antibody, or antigen-binding fragment thereof, further comprises a CDR-L2 as set forth in SEQ ID NO:98, optionally comprising one or more amino acid changes. In some embodiments, the antibody, or antigen-binding fragment thereof, further comprises a CDR-L3 as set forth in SEQ ID NO:99, optionally comprising one or more amino acid changes.

[317] In some embodiments, the antibody, or antigen-binding fragment thereof, comprises at least three of the following six CDRs:

- a) CDR-H1: SEQ ID NO:94, with the proviso that:

- i. the threonine residue at position 2 of SEQ ID NO:94 may be substituted with an alanine;
- ii. the asparagine residue at position 4 of SEQ ID NO:94 may be substituted with an alanine, tyrosine, aspartate, serine, arginine, or histidine;
- iii. the asparagine residue at position 5 of SEQ ID NO:94 may be substituted with a glutamine, serine, glycine, lysine, glutamate, arginine, or histidine;
- iv. the tyrosine residue at position 6 of SEQ ID NO:94 may be substituted with a arginine;
- v. the proline residue at position 7 of SEQ ID NO:94 may be substituted with a glycine, alanine, leucine, serine, asparagine, valine, aspartate, or glutamine;
- vi. the isoleucine residue at position 8 of SEQ ID NO:94 may be substituted with a methionine or leucine; and/or,
- vii. the histidine residue at position 9 of SEQ ID NO:94 may be substituted with a phenylalanine, tyrosine, asparagine, or serine;
- b) CDR-H2: SEQ ID NO:95, optionally comprising one or more amino acid changes;
- c) CDR-H3: SEQ ID NO:96, optionally comprising one or more amino acid changes;
- d) CDR-L1: SEQ ID NO:97, optionally comprising one or more amino acid changes;
- e) CDR-L2: SEQ ID NO:98, optionally comprising one or more amino acid changes;
- and,
- f) CDR-L3: SEQ ID NO:99, optionally comprising one or more amino acid changes.

[318] In some embodiments, the antibody, or antigen-binding fragment thereof, specifically binds a human LTBP1-proTGFβ1 complex and/or a human LTBP3-proTGFβ1 complex, and does not bind a human GARP-proTGFβ1 complex. In some embodiments, the antibody, or antigen-binding fragment thereof, does not bind a human GARP-proTGFβ1 complex. In some embodiments, the antibody, or antigen-binding fragment thereof, does not bind mature TGFβ1, mature TGFβ2 or mature TGFβ3.

[319] In a particular embodiment, the antibody, or antigen-binding fragment thereof, specifically binds a human LTBP1-proTGFβ1 complex and/or a human LTBP3-proTGFβ1 complex, and does not bind a human GARP-proTGFβ1 complex; wherein the antibody does not bind mature TGFβ1, mature TGFβ2 or mature TGFβ3; wherein the antibody is a fully human or humanized antibody or a fragment thereof, and wherein the antibody comprises at least three of the following six CDRs:

- a) CDR-H1: SEQ ID NO:94, with the proviso that:
 - i. the threonine residue at position 2 of SEQ ID NO:94 may be substituted with an alanine;
 - ii. the asparagine residue at position 4 of SEQ ID NO:94 may be substituted with an alanine, tyrosine, aspartate, serine, arginine, or histidine;

- iii. the asparagine residue at position 5 of SEQ ID NO:94 may be substituted with a glutamine, serine, glycine, lysine, glutamate, arginine, or histidine;
- iv. the tyrosine residue at position 6 of SEQ ID NO:94 may be substituted with a arginine;
- v. the proline residue at position 7 of SEQ ID NO:94 may be substituted with a glycine, alanine, leucine, serine, asparagine, valine, aspartate, or glutamine;
- vi. the isoleucine residue at position 8 of SEQ ID NO:94 may be substituted with a methionine or leucine; and/or,
- vii. the histidine residue at position 9 of SEQ ID NO:94 may be substituted with a phenylalanine, tyrosine, asparagine, or serine;
- b) CDR-H2: SEQ ID NO:95, optionally comprising one or more amino acid changes;
- c) CDR-H3: SEQ ID NO:96, optionally comprising one or more amino acid changes;
- d) CDR-L1: SEQ ID NO:97, optionally comprising one or more amino acid changes;
- e) CDR-L2: SEQ ID NO:98, optionally comprising one or more amino acid changes;
- and,
- f) CDR-L3: SEQ ID NO:99, optionally comprising one or more amino acid changes.

[320] In some embodiments, such antibody binds a human LTBP1-proTGFβ1 complex with a K_D of < 100 nM as measured in a suitable in vitro binding assay such as Bio-Layer Interferometry (BLI); and/or, such antibody binds a human LTBP3-proTGFβ1 complex with a K_D of < 100 nM as measured in a suitable in vitro binding assay such as Bio-Layer Interferometry (BLI). In some embodiments, such antibody binds a human LTBP1-proTGFβ1 complex with a K_D of < 50 nM as measured in a suitable in vitro binding assay such as Bio-Layer Interferometry (BLI); and/or, such antibody binds a human LTBP3-proTGFβ1 complex with a K_D of < 50 nM as measured in a suitable in vitro binding assay such as Bio-Layer Interferometry (BLI). In some embodiments, such antibody binds a human LTBP1-proTGFβ1 complex with a K_D of < 25 nM as measured in a suitable in vitro binding assay such as Bio-Layer Interferometry (BLI); and/or, such antibody binds a human LTBP3-proTGFβ1 complex with a K_D of < 25 nM as measured in a suitable in vitro binding assay such as Bio-Layer Interferometry (BLI). In some embodiments, such antibody binds a human LTBP1-proTGFβ1 complex with a K_D of < 10 nM as measured in a suitable in vitro binding assay such as Bio-Layer Interferometry (BLI); and/or, such antibody binds a human LTBP3-proTGFβ1 complex with a K_D of < 10 nM as measured in a suitable in vitro binding assay such as Bio-Layer Interferometry (BLI).

[321] In another embodiment, such antibody is cross-reactive with mouse LTBP1-proTGFβ1. In some embodiments, such antibody is also cross-reactive with mouse LTBP3-proTGFβ1. In some embodiments, such antibody binds a mouse LTBP1-proTGFβ1 complex with a K_D of < 100 nM as measured in a suitable in vitro binding assay such as Bio-Layer Interferometry (BLI); and/or, the

antibody binds a mouse LTBP3-proTGFβ1 complex with a K_D of < 100 nM as measured in a suitable in vitro binding assay such as Bio-Layer Interferometry (BLI). In some embodiments, such antibody binds a mouse LTBP1-proTGFβ1 complex with a K_D of < 50 nM as measured in a suitable in vitro binding assay such as Bio-Layer Interferometry (BLI); and/or, the antibody binds a mouse LTBP3-proTGFβ1 complex with a K_D of < 50 nM as measured in a suitable in vitro binding assay such as Bio-Layer Interferometry (BLI). In some embodiments, such antibody binds a mouse LTBP1-proTGFβ1 complex with a K_D of < 25 nM as measured in a suitable in vitro binding assay such as Bio-Layer Interferometry (BLI); and/or, the antibody binds a mouse LTBP3-proTGFβ1 complex with a K_D of < 25 nM as measured in a suitable in vitro binding assay such as Bio-Layer Interferometry (BLI). In some embodiments, such antibody binds a mouse LTBP1-proTGFβ1 complex with a K_D of < 10 nM as measured in a suitable in vitro binding assay such as Bio-Layer Interferometry (BLI); and/or, the antibody binds a mouse LTBP3-proTGFβ1 complex with a K_D of < 10 nM as measured in a suitable in vitro binding assay such as Bio-Layer Interferometry (BLI).

[322] In another embodiment, such antibody does not bind to human GARP-proTGFβ1. In preferred embodiments, such context-selective antibody is also isoform-specific in that it selectively binds and inhibits the activation of TGFβ1 associated with LTBP1/3.

[323] In another aspect, the antibody, or antigen-binding fragment thereof, comprises a CDR-H1 comprising the amino acid sequence FTF(X₁)(X₂)YVMH, wherein, optionally: X₁ is S or R; and X₂ is G or S. In some embodiments, X₁ is S. In some embodiments, X₁ is R. In some embodiments, X₂ is G. In some embodiments, X₂ is S.

[324] In some embodiments, the antibody, or antigen-binding fragment thereof, comprises a CDR-H2 comprising the amino acid sequence (X₁)ISHEG(X₂)(X₃)KYYADSVKG, wherein, optionally: X₁ is V or S; X₂ is S or G; and X₃ is F or L. In some embodiments, X₁ is a V. In some embodiment, X₁ is a S. In some embodiments, X₂ is S. In some embodiments, X₂ is G. In some embodiments, X₃ is F. In some embodiments, X₃ is L.

[325] In some embodiments, the antibody, or antigen-binding fragment thereof, comprises a CDR-H3 comprising the amino acid sequence (X₁)(X₂)P(X₃)(X₄)(X₅)(X₆)RRGG(X₇)(X₈)(X₉), wherein, optionally: X₁ is A or V; X₂ is R, V, G or K; X₃ is R, H or L; X₄ is I, V or G; X₅ is A, S, or L; X₆ is A or V; X₇ is F or Y; X₈ is D, G, R, or S; and, X₉ is Y, G, R, L, V, A or K. In some embodiments, X₁ is A. In some embodiments, X₁ is V. In some embodiments, X₂ is R. In some embodiments, X₂ is V. In some embodiments, X₂ is G. In some embodiments, X₂ is K. In some embodiments, X₃ is R. In some embodiments, X₃ is H. In some embodiments, X₃ is L. In some embodiments, X₄ is I. In some embodiments, X₄ is V. In some embodiments, X₄ is G. In some embodiments, X₅ is A. In some embodiments, X₅ is S. In some embodiments, X₅ is L. In some embodiments, X₆ is A. In some embodiments, X₆ is V. In some embodiments, X₇ is F. In some embodiments, X₇ is Y. In some

embodiments, X₈ is D. In some embodiments, X₈ is G. In some embodiments, X₈ is R. In some embodiments, X₈ is S. In some embodiments, X₉ is Y. In some embodiments, X₉ is G. In some embodiments, X₉ is R. In some embodiments, X₉ is L. In some embodiments, X₉ is V. In some embodiments, X₉ is A. In some embodiments, X₉ is K.

[326] In some embodiments, the antibody, or antigen-binding fragment thereof, comprises three heavy chain CDRs and three light chain CDRs, wherein the heavy chain CDRs comprise:

- a) CDR-H1 comprising the amino acid sequence FTF(X₁)(X₂)YVMH, wherein, optionally: X₁ is S or R; and X₂ is G or S;
- b) CDR-H2 comprising the amino acid sequence (X₁)ISHEG(X₂)(X₃)KYYADSVKG, wherein, optionally: X₁ is V or S; X₂ is S or G; and X₃ is F or L; and
- c) CDR-H3 comprising the amino acid sequence (X₁)(X₂)P(X₃)(X₄)(X₅)(X₆)RRGG(X₇)(X₈)(X₉), wherein, optionally: X₁ is A or V; X₂ is R, V, G or K; X₃ is R, H or L; X₄ is I, V or G; X₅ is A, S, or L; X₆ is A or V; X₇ is F or Y; X₈ is D, G, R, or S; and, X₉ is Y, G, R, L, V, A or K.

[327] In some embodiments, the antibody, or antigen-binding fragment thereof, further comprises a CDR-L1 as set forth in SEQ ID NO:97, optionally comprising one or more amino acid changes. In some embodiments, the antibody, or antigen-binding fragment thereof, further comprises a CDR-L2 as set forth in SEQ ID NO:98, optionally comprising one or more amino acid changes. In some embodiments, the antibody, or antigen-binding fragment thereof, further comprises a CDR-L3 as set forth in SEQ ID NO:99, optionally comprising one or more amino acid changes.

[328] In some embodiments, the antibody, or antigen-binding fragment, comprises at least three of the following six CDRs:

- a) CDR-H1 comprising the amino acid sequence FTF(X₁)(X₂)YVMH, wherein, optionally: X₁ is S or R; and X₂ is G or S;
- b) CDR-H2 comprising the amino acid sequence (X₁)ISHEG(X₂)(X₃)KYYADSVKG, wherein, optionally: X₁ is V or S; X₂ is S or G; and X₃ is F or L;
- c) CDR-H3 comprising the amino acid sequence (X₁)(X₂)P(X₃)(X₄)(X₅)(X₆)RRGG(X₇)(X₈)(X₉), wherein, optionally: X₁ is A or V; X₂ is R, V, G or K; X₃ is R, H or L; X₄ is I, V or G; X₅ is A, S, or L; X₆ is A or V; X₇ is F or Y; X₈ is D, G, R, or S; and, X₉ is Y, G, R, L, V, A or K;
- d) CDR-L1 as set forth in SEQ ID NO:97, optionally comprising one or more amino acid changes;
- e) CDR-L2 as set forth in SEQ ID NO:98, optionally comprising one or more amino acid changes; and
- f) CDR-L3 as set forth in SEQ ID NO:99, optionally comprising one or more amino acid changes.

[329] In some embodiments, the antibody, or antigen-binding fragment thereof, specifically binds a human LTBP1-proTGFβ1 complex and/or a human LTBP3-proTGFβ1 complex, and does not bind a human GARP-proTGFβ1 complex. In some embodiments, the antibody, or antigen-binding fragment thereof, does not bind a human GARP-proTGFβ1 complex. In some embodiments, the antibody, or antigen-binding fragment thereof, does not bind mature TGFβ1, mature TGFβ2 or mature TGFβ3.

[330] In a particular embodiment, the antibody, or antigen-binding fragment, specifically binds a human LTBP1-proTGFβ1 complex and/or a human LTBP3-proTGFβ1 complex, and does not bind a human GARP-proTGFβ1 complex; wherein the antibody does not bind mature TGFβ1, mature TGFβ2 or mature TGFβ3; wherein the antibody is a fully human or humanized antibody or a fragment thereof, wherein the antibody comprises at least three of the following six CDRs:

- a) CDR-H1 comprising the amino acid sequence FTF(X₁)(X₂)YVMH, wherein, optionally: X₁ is S or R; and X₂ is G or S;
- b) CDR-H2 comprising the amino acid sequence (X₁)ISHEG(X₂)(X₃)KYYADSVKG, wherein, optionally: X₁ is V or S; X₂ is S or G; and X₃ is F or L;
- c) CDR-H3 comprising the amino acid sequence (X₁)(X₂)P(X₃)(X₄)(X₅)(X₆)RRGG(X₇)(X₈)(X₉), wherein, optionally: X₁ is A or V; X₂ is R, V, G or K; X₃ is R, H or L; X₄ is I, V or G; X₅ is A, S, or L; X₆ is A or V; X₇ is F or Y; X₈ is D, G, R, or S; and, X₉ is Y, G, R, L, V, A or K;
- d) CDR-L1 as set forth in SEQ ID NO:97, optionally comprising one or more amino acid changes;
- e) CDR-L2 as set forth in SEQ ID NO:98, optionally comprising one or more amino acid changes; and
- f) CDR-L3 as set forth in SEQ ID NO:99, optionally comprising one or more amino acid changes.

[331] In some embodiments, such antibody binds a human LTBP1-proTGFβ1 complex with a K_D of < 100 nM as measured in a suitable in vitro binding assay such as Bio-Layer Interferometry (BLI); and/or, such antibody binds a human LTBP3-proTGFβ1 complex with a K_D of < 100 nM as measured in a suitable in vitro binding assay such as Bio-Layer Interferometry (BLI). In some embodiments, such antibody binds a human LTBP1-proTGFβ1 complex with a K_D of < 50 nM as measured in a suitable in vitro binding assay such as Bio-Layer Interferometry (BLI); and/or, such antibody binds a human LTBP3-proTGFβ1 complex with a K_D of < 50 nM as measured in a suitable in vitro binding assay such as Bio-Layer Interferometry (BLI). In some embodiments, such antibody binds a human LTBP1-proTGFβ1 complex with a K_D of < 25 nM as measured in a suitable in vitro binding assay such as Bio-Layer Interferometry (BLI); and/or, such antibody binds a human LTBP3-proTGFβ1 complex with a K_D of < 25 nM as measured in a suitable in vitro binding assay such as Bio-Layer Interferometry (BLI). In some embodiments, such antibody binds a human LTBP1-proTGFβ1

complex with a K_D of < 10 nM as measured in a suitable in vitro binding assay such as Bio-Layer Interferometry (BLI); and/or, such antibody binds a human LTBP3-proTGF β 1 complex with a K_D of < 10 nM as measured in a suitable in vitro binding assay such as Bio-Layer Interferometry (BLI).

[332] In another embodiment, such antibody is cross-reactive with mouse LTBP1-proTGF β 1. In some embodiments, such antibody is also cross-reactive with mouse LTBP3-proTGF β 1. In some embodiments, such antibody binds a mouse LTBP1-proTGF β 1 complex with a K_D of < 100 nM as measured in a suitable in vitro binding assay such as Bio-Layer Interferometry (BLI); and/or, the antibody binds a mouse LTBP3-proTGF β 1 complex with a K_D of < 100 nM as measured in a suitable in vitro binding assay such as Bio-Layer Interferometry (BLI). In some embodiments, such antibody binds a mouse LTBP1-proTGF β 1 complex with a K_D of < 50 nM as measured in a suitable in vitro binding assay such as Bio-Layer Interferometry (BLI); and/or, the antibody binds a mouse LTBP3-proTGF β 1 complex with a K_D of < 50 nM as measured in a suitable in vitro binding assay such as Bio-Layer Interferometry (BLI). In some embodiments, such antibody binds a mouse LTBP1-proTGF β 1 complex with a K_D of < 25 nM as measured in a suitable in vitro binding assay such as Bio-Layer Interferometry (BLI); and/or, the antibody binds a mouse LTBP3-proTGF β 1 complex with a K_D of < 25 nM as measured in a suitable in vitro binding assay such as Bio-Layer Interferometry (BLI). In some embodiments, such antibody binds a mouse LTBP1-proTGF β 1 complex with a K_D of < 10 nM as measured in a suitable in vitro binding assay such as Bio-Layer Interferometry (BLI); and/or, the antibody binds a mouse LTBP3-proTGF β 1 complex with a K_D of < 10 nM as measured in a suitable in vitro binding assay such as Bio-Layer Interferometry (BLI).

[333] In another embodiment, such antibody does not bind to human GARP-proTGF β 1. In preferred embodiments, such context-selective antibody is also isoform-specific in that it selectively binds and inhibits the activation of TGF β 1 associated with LTBP1/3.

[334] In another aspect, the antibody, or antigen-binding fragment thereof, comprises a CDR-H1 comprising the amino acid sequence FTF(X₁)(X₂)YVMH, wherein, optionally: X₁ is S or R; and X₂ is G or S. In some embodiments, X₁ is S. In some embodiments, X₁ is R. In some embodiments, X₂ is G. In some embodiments, X₂ is S.

[335] In some embodiments, the antibody, or antigen-binding fragment thereof, comprises a CDR-H2 comprising the amino acid sequence (X₁)ISHEGS(X₂)KYYADSVKG, wherein, optionally: X₁ is V or S; and, X₂ is F or L. In some embodiments, X₁ is a V. In some embodiment, X₁ is a S. In some embodiments, X₃ is F. In some embodiments, X₃ is L.

[336] In some embodiments, the antibody, or antigen-binding fragment thereof, comprises a CDR-H3 comprising the amino acid sequence A(X₁)PRI(X₂)ARRGGFGY, wherein, optionally: X₁ is R or

V; X₂ is A or L. In some embodiments, X₁ is R. In some embodiments, X₁ is V. In some embodiments, X₂ is A. In some embodiments, X₂ is L.

[337] In some embodiments, the antibody, or antigen-binding fragment thereof, comprises three heavy chain CDRs and three light chain CDRs, wherein the heavy chain CDRs comprise:

- a) CDR-H1 comprising the amino acid sequence FTF(X₁)(X₂)YVMH, wherein, optionally: X₁ is S or R; and X₂ is G or S;
- b) CDR-H2 comprising the amino acid sequence (X₁)ISHEGS(X₂)KYYADSVKG, wherein, optionally: X₁ is V or S; and, X₂ is F or L; and
- c) CDR-H3 comprising the amino acid sequence A(X₁)PRI(X₂)ARRGGFGY, wherein, optionally: X₁ is R or V; X₂ is A or L.

[338] In some embodiments, the antibody, or antigen-binding fragment thereof, further comprises a CDR-L1 as set forth in SEQ ID NO:97, optionally comprising one or more amino acid changes. In some embodiment, the antibody, or antigen-binding fragment thereof, further comprises a CDR-L2 as set forth in SEQ ID NO:98, optionally comprising one or more amino acid changes. In some embodiments, the antibody, or antigen-binding fragment thereof, further comprises a CDR-L3 as set forth in SEQ ID NO:99, optionally comprising one or more amino acid changes.

[339] In some embodiments, the antibody, or antigen-binding fragment, comprises at least three of the following six CDRs:

- a) CDR-H1 comprising the amino acid sequence FTF(X₁)(X₂)YVMH, wherein, optionally: X₁ is S or R; and X₂ is G or S;
- b) CDR-H2 comprising the amino acid sequence (X₁)ISHEGS(X₂)KYYADSVKG, wherein, optionally: X₁ is V or S; and, X₂ is F or L;
- c) CDR- comprising the amino acid sequence A(X₁)PRI(X₂)ARRGGFGY, wherein, optionally: X₁ is R or V; X₂ is A or L;
- d) CDR-L1 as set forth in SEQ ID NO:97, optionally comprising one or more amino acid changes;
- e) CDR-L2 as set forth in SEQ ID NO:98, optionally comprising one or more amino acid changes; and
- f) CDR-L3 as set forth in SEQ ID NO:99, optionally comprising one or more amino acid changes.

[340] In some embodiments, the antibody, or antigen-binding fragment thereof, specifically binds a human LTBP1-proTGFβ1 complex and/or a human LTBP3-proTGFβ1 complex, and does not bind a human GARP-proTGFβ1 complex. In some embodiments, the antibody, or antigen-binding fragment thereof, does not bind a human GARP-proTGFβ1 complex. In some embodiments, the antibody, or antigen-binding fragment thereof, does not bind mature TGFβ1, mature TGFβ2 or mature TGFβ3.

[341] In a particular embodiment, the antibody, or antigen-binding fragment thereof, specifically binds a human LTBP1-proTGF β 1 complex and/or a human LTBP3-proTGF β 1 complex, and does not bind a human GARP-proTGF β 1 complex; wherein the antibody does not bind mature TGF β 1, mature TGF β 2 or mature TGF β 3; wherein the antibody is a fully human or humanized antibody or a fragment thereof, wherein the antibody comprises at least three of the following six CDRs:

- a) CDR-H1 comprising the amino acid sequence FTF(X₁)(X₂)YVMH, wherein, optionally: X₁ is S or R; and X₂ is G or S;
- b) CDR-H2 comprising the amino acid sequence (X₁)ISHEGS(X₂)KYYADSVKG, wherein: X₁ is V or S; and, X₂ is F or L;
- c) CDR-H3 comprising the amino acid sequence A(X₁)PRI(X₂)ARRGGFGY, wherein, optionally: X₁ is R or V; X₂ is A or L;
- d) CDR-L1 as set forth in SEQ ID NO:97, optionally comprising one or more amino acid changes;
- e) CDR-L2 as set forth in SEQ ID NO:98, optionally comprising one or more amino acid changes; and
- f) CDR-L3 as set forth in SEQ ID NO:99, optionally comprising one or more amino acid changes.

[342] In some embodiments, such antibody binds a human LTBP1-proTGF β 1 complex with a K_D of < 100 nM as measured in a suitable in vitro binding assay such as Bio-Layer Interferometry (BLI); and/or, such antibody binds a human LTBP3-proTGF β 1 complex with a K_D of < 100 nM as measured in a suitable in vitro binding assay such as Bio-Layer Interferometry (BLI). In some embodiments, such antibody binds a human LTBP1-proTGF β 1 complex with a K_D of < 50 nM as measured in a suitable in vitro binding assay such as Bio-Layer Interferometry (BLI); and/or, such antibody binds a human LTBP3-proTGF β 1 complex with a K_D of < 50 nM as measured in a suitable in vitro binding assay such as Bio-Layer Interferometry (BLI). In some embodiments, such antibody binds a human LTBP1-proTGF β 1 complex with a K_D of < 25 nM as measured in a suitable in vitro binding assay such as Bio-Layer Interferometry (BLI); and/or, such antibody binds a human LTBP3-proTGF β 1 complex with a K_D of < 25 nM as measured in a suitable in vitro binding assay such as Bio-Layer Interferometry (BLI). In some embodiments, such antibody binds a human LTBP1-proTGF β 1 complex with a K_D of < 10 nM as measured in a suitable in vitro binding assay such as Bio-Layer Interferometry (BLI); and/or, such antibody binds a human LTBP3-proTGF β 1 complex with a K_D of < 10 nM as measured in a suitable in vitro binding assay such as Bio-Layer Interferometry (BLI).

[343] In another embodiment, such antibody is cross-reactive with mouse LTBP1-proTGF β 1. In some embodiments, such antibody is also cross-reactive with mouse LTBP3-proTGF β 1. In some embodiments, such antibody binds a mouse LTBP1-proTGF β 1 complex with a K_D of < 100 nM as

measured in a suitable in vitro binding assay such as Bio-Layer Interferometry (BLI); and/or, the antibody binds a mouse LTBP3-proTGF β 1 complex with a K_D of < 100 nM as measured in a suitable in vitro binding assay such as Bio-Layer Interferometry (BLI). In some embodiments, such antibody binds a mouse LTBP1-proTGF β 1 complex with a K_D of < 50 nM as measured in a suitable in vitro binding assay such as Bio-Layer Interferometry (BLI); and/or, the antibody binds a mouse LTBP3-proTGF β 1 complex with a K_D of < 50 nM as measured in a suitable in vitro binding assay such as Bio-Layer Interferometry (BLI). In some embodiments, such antibody binds a mouse LTBP1-proTGF β 1 complex with a K_D of < 25 nM as measured in a suitable in vitro binding assay such as Bio-Layer Interferometry (BLI); and/or, the antibody binds a mouse LTBP3-proTGF β 1 complex with a K_D of < 25 nM as measured in a suitable in vitro binding assay such as Bio-Layer Interferometry (BLI). In some embodiments, such antibody binds a mouse LTBP1-proTGF β 1 complex with a K_D of < 10 nM as measured in a suitable in vitro binding assay such as Bio-Layer Interferometry (BLI); and/or, the antibody binds a mouse LTBP3-proTGF β 1 complex with a K_D of < 10 nM as measured in a suitable in vitro binding assay such as Bio-Layer Interferometry (BLI).

[344] In another embodiment, such antibody does not bind to human GARP-proTGF β 1. In preferred embodiments, such context-selective antibody is also isoform-specific in that it selectively binds and inhibits the activation of TGF β 1 associated with LTBP1/3.

[345] In another aspect, the antibody, or antigen-binding fragment thereof, comprises a CDR-H1: SEQ ID NO:100, with the proviso that the serine residue at position 4 of SEQ ID NO:100 may be substituted with a histidine. In some embodiments, the antibody, or antigen-binding fragment thereof, comprises a CDR-H1: SEQ ID NO:100, with the proviso that the serine residue at position 7 of SEQ ID NO:100 may be substituted with an alanine or glycine. In some embodiments, the antibody, or antigen-binding fragment thereof, comprises a CDR-H1: SEQ ID NO:100, with the proviso that the glycine residue at position 11 of SEQ ID NO:100 may be substituted with a threonine, serine, histidine, leucine, isoleucine, asparagine, valine, or alanine.

[346] In some embodiments, the antibody, or antigen-binding fragment thereof, comprises a CDR-H1: SEQ ID NO:100, with the proviso that (i) the serine residue at position 4 of SEQ ID NO:100 may be substituted with a histidine; (ii) the serine residue at position 7 of SEQ ID NO:100 may be substituted with an alanine or glycine; and/or, (iii) the glycine residue at position 11 of SEQ ID NO:100 may be substituted with a threonine, serine, histidine, leucine, isoleucine, asparagine, valine, or alanine.

[347] In some embodiments, the antibody, or antigen-binding fragment thereof, comprises a CDR-H1: SEQ ID NO:101, with the proviso that the serine residue at position 3 of SEQ ID NO:101 may be substituted with an alanine. In some embodiments, the antibody, or antigen-binding fragment thereof, comprises a CDR-H1: SEQ ID NO:101, with the proviso that the glycine residue at position 6

of SEQ ID NO:101 may be substituted with an alanine or serine. In some embodiments, the antibody, or antigen-binding fragment thereof, comprises a CDR-H1: SEQ ID NO:101, with the proviso that the serine residue at position 7 of SEQ ID NO:101 may be substituted with a threonine.

[348] In some embodiments, the antibody, or antigen-binding fragment thereof, comprises a CDR-H1: SEQ ID NO:101, with the proviso that (i) the serine residue at position 3 of SEQ ID NO:101 may be substituted with an alanine; (ii) the glycine residue at position 6 of SEQ ID NO:101 may be substituted with an alanine or serine; and/or, (iii) the serine residue at position 7 of SEQ ID NO:101 may be substituted with a threonine.

[349] In some embodiments, the antibody, or antigen-binding fragment thereof, comprises three heavy chain CDRs and three light chain CDRs, wherein the heavy chain CDRs comprise:

- a) CDR-H1: SEQ ID NO:100, with the proviso that:
 - i. the serine residue at position 4 of SEQ ID NO:100 may be substituted with a histidine;
 - ii. the serine residue at position 7 of SEQ ID NO:100 may be substituted with an alanine or glycine; and/or,
 - iii. the glycine residue at position 11 of SEQ ID NO:100 may be substituted with a threonine, serine, histidine, leucine, isoleucine, asparagine, valine, or alanine;
- b) CDR-H2: SEQ ID NO:101, with the proviso that:
 - i. the serine residue at position 3 of SEQ ID NO:101 may be substituted with an alanine;
 - ii. the glycine residue at position 6 of SEQ ID NO:101 may be substituted with an alanine or serine; and/or,
 - iii. the serine residue at position 7 of SEQ ID NO:101 may be substituted with a threonine; and
- c) CDR-H3: SEQ ID NO:102, optionally comprising up to three, four, five or six amino acid changes.

[350] In some embodiments, the antibody, or antigen-binding fragment thereof, further comprises a CDR-L1 as set forth in SEQ ID NO:103, optionally comprising one or more amino acid changes. In some embodiments, the antibody, or antigen-binding fragment thereof, further comprises a CDR-L2 as set forth in SEQ ID NO:104, optionally comprising one or more amino acid changes. In some embodiments, the antibody, or antigen-binding fragment thereof, further comprises a CDR-L3 as set forth in SEQ ID NO:105, optionally comprising one or more amino acid changes.

[351] In some embodiments, the antibody, or antigen-binding fragment thereof, comprises at least three of the following six CDRs:

- a) CDR-H1: SEQ ID NO:100, with the proviso that:

- i. the serine residue at position 4 of SEQ ID NO:100 may be substituted with a histidine;
- ii. the serine residue at position 7 of SEQ ID NO:100 may be substituted with an alanine or glycine; and/or,
- iii. the glycine residue at position 11 of SEQ ID NO:100 may be substituted with a threonine, serine, histidine, leucine, isoleucine, asparagine, valine, or alanine;
- b) CDR-H2: SEQ ID NO:101, with the proviso that:
 - i. the serine residue at position 3 of SEQ ID NO:101 may be substituted with an alanine;
 - ii. the glycine residue at position 6 of SEQ ID NO:101 may be substituted with an alanine or serine; and/or,
 - iii. the serine residue at position 7 of SEQ ID NO:101 may be substituted with a threonine; and
- c) CDR-H3: SEQ ID NO:102, optionally comprising one or more amino acid changes.
- d) CDR-L1: SEQ ID NO:103, optionally comprising one or more amino acid changes;
- e) CDR-L2: SEQ ID NO:104, optionally comprising one or more amino acid changes; and,
- f) CDR-L3: SEQ ID NO:105, optionally comprising one or more amino acid changes.

[352] In some embodiments, the antibody, or antigen-binding fragment, specifically binds a human LTBP1-proTGFβ1 complex and/or a human LTBP3-proTGFβ1 complex, and does not bind a human GARP-proTGFβ1 complex. In some embodiments, the antibody, or antigen-binding fragment thereof, does not bind a human GARP-proTGFβ1 complex. In some embodiments, the antibody, or antigen-binding fragment thereof, does not bind mature TGFβ1, mature TGFβ2 or mature TGFβ3.

[353] In a particular embodiment, the antibody, or antigen-binding fragment, specifically binds a human LTBP1-proTGFβ1 complex and/or a human LTBP3-proTGFβ1 complex, and does not bind a human GARP-proTGFβ1 complex; wherein the antibody does not bind mature TGFβ1, mature TGFβ2 or mature TGFβ3; wherein the antibody is a fully human or humanized antibody or a fragment thereof, wherein the antibody comprises at least three of the following six CDRs:

- a) CDR-H1: SEQ ID NO:100, with the proviso that:
 - i. the serine residue at position 4 of SEQ ID NO:100 may be substituted with a histidine;
 - ii. the serine residue at position 7 of SEQ ID NO:100 may be substituted with an alanine or glycine; and/or,
 - iii. the glycine residue at position 11 of SEQ ID NO:100 may be substituted with a threonine, serine, histidine, leucine, isoleucine, asparagine, valine, or alanine;
- b) CDR-H2: SEQ ID NO:101, with the proviso that:

- i. the serine residue at position 3 of SEQ ID NO:101 may be substituted with an alanine;
- ii. the glycine residue at position 6 of SEQ ID NO:101 may be substituted with an alanine or serine; and/or,
- iii. the serine residue at position 7 of SEQ ID NO:101 may be substituted with a threonine;
- c) CDR-H3: SEQ ID NO:102, optionally comprising one or more amino acid changes;
- d) CDR-L1: SEQ ID NO:103, optionally comprising one or more amino acid changes;
- e) CDR-L2: SEQ ID NO:104, optionally comprising one or more amino acid changes; and,
- f) CDR-L3: SEQ ID NO:105, optionally comprising one or more amino acid changes.

[354] In some embodiments, such antibody binds a human LTBP1-proTGF β 1 complex with a KD of < 100 nM as measured in a suitable in vitro binding assay such as Bio-Layer Interferometry (BLI); and/or, such antibody binds a human LTBP3-proTGF β 1 complex with a KD of < 100 nM as measured in a suitable in vitro binding assay such as Bio-Layer Interferometry (BLI). In some embodiments, such antibody binds a human LTBP1-proTGF β 1 complex with a KD of < 50 nM as measured in a suitable in vitro binding assay such as Bio-Layer Interferometry (BLI); and/or, such antibody binds a human LTBP3-proTGF β 1 complex with a KD of < 50 nM as measured in a suitable in vitro binding assay such as Bio-Layer Interferometry (BLI). In some embodiments, such antibody binds a human LTBP1-proTGF β 1 complex with a KD of < 25 nM as measured in a suitable in vitro binding assay such as Bio-Layer Interferometry (BLI); and/or, such antibody binds a human LTBP3-proTGF β 1 complex with a KD of < 25 nM as measured in a suitable in vitro binding assay such as Bio-Layer Interferometry (BLI). In some embodiments, such antibody binds a human LTBP1-proTGF β 1 complex with a KD of < 10 nM as measured in a suitable in vitro binding assay such as Bio-Layer Interferometry (BLI); and/or, such antibody binds a human LTBP3-proTGF β 1 complex with a KD of < 10 nM as measured in a suitable in vitro binding assay such as Bio-Layer Interferometry (BLI).

[355] In another embodiment, such antibody is cross-reactive with mouse LTBP1-proTGF β 1. In some embodiments, such antibody is also cross-reactive with mouse LTBP3-proTGF β 1. In some embodiments, such antibody binds a mouse LTBP1-proTGF β 1 complex with a KD of < 100 nM as measured in a suitable in vitro binding assay such as Bio-Layer Interferometry (BLI); and/or, the antibody binds a mouse LTBP3-proTGF β 1 complex with a KD of < 100 nM as measured in a suitable in vitro binding assay such as Bio-Layer Interferometry (BLI). In some embodiments, such antibody binds a mouse LTBP1-proTGF β 1 complex with a KD of < 50 nM as measured in a suitable in vitro binding assay such as Bio-Layer Interferometry (BLI); and/or, the antibody binds a mouse LTBP3-proTGF β 1 complex with a KD of < 50 nM as measured in a suitable in vitro binding assay such as

Bio-Layer Interferometry (BLI). In some embodiments, such antibody binds a mouse LTBP1-proTGFβ1 complex with a KD of < 25 nM as measured in a suitable in vitro binding assay such as Bio-Layer Interferometry (BLI); and/or, the antibody binds a mouse LTBP3-proTGFβ1 complex with a KD of < 25 nM as measured in a suitable in vitro binding assay such as Bio-Layer Interferometry (BLI). In some embodiments, such antibody binds a mouse LTBP1-proTGFβ1 complex with a KD of < 10 nM as measured in a suitable in vitro binding assay such as Bio-Layer Interferometry (BLI); and/or, the antibody binds a mouse LTBP3-proTGFβ1 complex with a KD of < 10 nM as measured in a suitable in vitro binding assay such as Bio-Layer Interferometry (BLI).

[356] In another embodiment, such antibody does not bind to human GARP-proTGFβ1. In preferred embodiments, such context-selective antibody is also isoform-specific in that it selectively binds and inhibits the activation of TGFβ1 associated with LTBP1/3.

[357] In another aspect, the antibody, or antigen-binding fragment thereof, comprises a CDR-H1 comprising the amino acid sequence G(X₁)I(X₂)S(X₃)SYYW(X₄), wherein, optionally: X₁ is S or P; X₂ is S, H or R; X₃ is S or G; and, X₄ is G, I, N or V. In some embodiments, X₁ is a S. In some embodiments, X₁ is P. In some embodiments, X₂ is S. In some embodiments, X₂ is H. In some embodiments, X₂ is R. In some embodiments, X₃ is S. In some embodiments, X₃ is G. In some embodiments, X₄ is G. In some embodiments, X₄ is I. In some embodiments, X₄ is N. In some embodiments, X₄ is V.

[358] In some embodiments, the antibody, or antigen-binding fragment thereof, comprises a CDR-H2 comprising the amino acid sequence SISYSA(X₁)TYYNPSLKS, wherein, optionally, X₁ is S or T. In some embodiments, X₁ is a S. In some embodiment, X₁ is a T.

[359] In some embodiments, the antibody, or antigen-binding fragment thereof, comprises a CDR-H3 comprising the amino acid sequence (X₁)(X₂)D(X₃)(X₄)Y(X₅)(X₆)(X₇)(X₈)G(X₉)(X₁₀)(X₁₁), wherein, optionally: X₁ is A or V; X₂ is R, S or G; X₃ is P, Y, R, V, I, H, T or E; X₄ is S, D, E or N; X₅ is D, A or T; X₆ is S, G, T or A; X₇ is I, A, R, Q, or V; X₈ is A, E, K, G or T; X₉ is M or I; X₁₀ is D, L, Q, V, N or G; and, X₁₁ is V, R, N, E or K. In some embodiments, X₁ is A. In some embodiments, X₁ is V. In some embodiments, X₂ is R. In some embodiments, X₂ is S. In some embodiments, X₂ is G. In some embodiments, X₃ is P. In some embodiments, X₃ is Y. In some embodiments, X₃ is R. In some embodiments, X₃ is V. In some embodiments, X₃ is I. In some embodiments, X₃ is H. In some embodiments, X₃ is T. In some embodiments, X₃ is E. In some embodiments, X₄ is S. In some embodiments, X₄ is D. In some embodiments, X₄ is E. In some embodiments, X₄ is N. In some embodiments, X₅ is D. In some embodiments, X₅ is A. In some embodiments, X₅ is T. In some embodiments, X₆ is S. In some embodiments, X₆ is G. In some embodiments, X₆ is T. In some embodiments, X₆ is A. In some embodiments, X₇ is I. In some embodiments, X₇ is A. In some embodiments, X₇ is R. In some embodiments, X₇ is Q. In some embodiments, X₇ is V. In some embodiments, X₈ is A. In some embodiments, X₈ is E. In some

embodiments, X_8 is K. In some embodiments, X_8 is G. In some embodiments, X_8 is T. In some embodiments, X_9 is M. In some embodiments, X_9 is I. In some embodiments, X_{10} is D. In some embodiments, X_{10} is L. In some embodiments, X_{10} is Q. In some embodiments, X_{10} is V. In some embodiments, X_{10} is N. In some embodiments, X_{10} is G. In some embodiments, X_{11} is V. In some embodiments, X_{11} is R. In some embodiments, X_{11} is N. In some embodiments, X_{11} is E. In some embodiments, X_{11} is K.

[360] In some embodiments, the antibody, or antigen-binding fragment thereof, comprises three heavy chain CDRs and three light chain CDRs, wherein the heavy chain CDRs comprise:

- a) CDR-H1 comprising the amino acid sequence $G(X_1)I(X_2)S(X_3)SYYW(X_4)$, wherein, optionally: X_1 is S or P; X_2 is S, H or R; X_3 is S or G; and, X_4 is G, I, N or V;
- b) CDR-H2 comprising the amino acid sequence $SISYSA(X_1)TYYNPSLKS$, wherein, optionally, X_1 is S or T; and
- c) CDR-H3 comprising the amino acid sequence $(X_1)(X_2)D(X_3)(X_4)Y(X_5)(X_6)(X_7)(X_8)G(X_9)(X_{10})(X_{11})$, wherein, optionally: X_1 is A or V; X_2 is R, S or G; X_3 is P, Y, R, V, I, H, T or E; X_4 is S, D, E or N; X_5 is D, A or T; X_6 is S, G, T or A; X_7 is I, A, R, Q, or V; X_8 is A, E, K, G or T; X_9 is M or I; X_{10} is D, L, Q, V, N or G; and, X_{11} is V, R, N, E or K.

[361] In some embodiments, the antibody, or antigen-binding fragment thereof, further comprises a CDR-L1 as set forth in SEQ ID NO:103, optionally comprising one or more amino acid changes. In some embodiment, the antibody, or antigen-binding fragment thereof, further comprises a CDR-L2 as set forth in SEQ ID NO:104, optionally comprising one or more amino acid changes. In some embodiments, the antibody, or antigen-binding fragment thereof, further comprises a CDR-L3 as set forth in SEQ ID NO:105, optionally comprising one or more amino acid changes.

[362] In some embodiments, the antibody, or antigen-binding fragment, comprises at least three of the following six CDRs:

- a) CDR-H1 comprising the amino acid sequence $G(X_1)I(X_2)S(X_3)SYYW(X_4)$, wherein, optionally: X_1 is S or P; X_2 is S, H or R; X_3 is S or G; and, X_4 is G, I, N or V;
- b) CDR-H2 comprising the amino acid sequence $SISYSA(X_1)TYYNPSLKS$, wherein, optionally, X_1 is S or T;
- c) CDR-H3 comprising the amino acid sequence $(X_1)(X_2)D(X_3)(X_4)Y(X_5)(X_6)(X_7)(X_8)G(X_9)(X_{10})(X_{11})$, wherein, optionally: X_1 is A or V; X_2 is R, S or G; X_3 is P, Y, R, V, I, H, T or E; X_4 is S, D, E or N; X_5 is D, A or T; X_6 is S, G, T or A; X_7 is I, A, R, Q, or V; X_8 is A, E, K, G or T; X_9 is M or I; X_{10} is D, L, Q, V, N or G; and, X_{11} is V, R, N, E or K;
- d) CDR-L1: SEQ ID NO:103, optionally comprising one or more amino acid changes;
- e) CDR-L2: SEQ ID NO: 104, optionally comprising one or more amino acid changes; and

- f) CDR-L3: SEQ ID NO: 105, optionally comprising one or more amino acid changes.

[363] In some embodiments, the antibody, or antigen-binding fragment thereof, specifically binds a human LTBP1-proTGFβ1 complex and/or a human LTBP3-proTGFβ1 complex, and does not bind a human GARP-proTGFβ1 complex. In some embodiments, the antibody, or antigen-binding fragment thereof, does not bind a human GARP-proTGFβ1 complex. In some embodiments, the antibody, or antigen-binding fragment thereof, does not bind mature TGFβ1, mature TGFβ2 or mature TGFβ3.

[364] In a particular embodiment, the antibody, or antigen-binding fragment, specifically binds a human LTBP1-proTGFβ1 complex and/or a human LTBP3-proTGFβ1 complex, and does not bind a human GARP-proTGFβ1 complex; wherein the antibody does not bind mature TGFβ1, mature TGFβ2 or mature TGFβ3; wherein the antibody is a fully human or humanized antibody or a fragment thereof, wherein the antibody comprises at least three of the following six CDRs:

- a) CDR-H1 comprising the amino acid sequence G(X₁)I(X₂)S(X₃)SYYW(X₄), wherein, optionally: X₁ is S or P; X₂ is S, H or R; X₃ is S or G; and, X₄ is G, I, N or V;
- b) CDR-H2 comprising the amino acid sequence SISYSA(X₁)TYYNPSLKS, wherein, optionally, X₁ is S or T;
- c) CDR-H3 comprising the amino acid sequence (X₁)(X₂)D(X₃)(X₄)Y(X₅)(X₆)(X₇)(X₈)G(X₉)(X₁₀)(X₁₁), wherein, optionally: X₁ is A or V; X₂ is R, S or G; X₃ is P, Y, R, V, I, H, T or E; X₄ is S, D, E or N; X₅ is D, A or T; X₆ is S, G, T or A; X₇ is I, A, R, Q, or V; X₈ is A, E, K, G or T; X₉ is M or I; X₁₀ is D, L, Q, V, N or G; and, X₁₁ is V, R, N, E or K;
- d) CDR-L1 as set forth in SEQ ID NO:103, optionally comprising one or more amino acid changes;
- e) CDR-L2 as set forth in SEQ ID NO: 104, optionally comprising one or more amino acid changes; and
- f) CDR-L3 as set forth in SEQ ID NO: 105, optionally comprising one or more amino acid changes.

[365] In some embodiments, such antibody binds a human LTBP1-proTGFβ1 complex with a KD of < 100 nM as measured in a suitable in vitro binding assay such as Bio-Layer Interferometry (BLI); and/or, such antibody binds a human LTBP3-proTGFβ1 complex with a KD of < 100 nM as measured in a suitable in vitro binding assay such as Bio-Layer Interferometry (BLI). In some embodiments, such antibody binds a human LTBP1-proTGFβ1 complex with a KD of < 50 nM as measured in a suitable in vitro binding assay such as Bio-Layer Interferometry (BLI); and/or, such antibody binds a human LTBP3-proTGFβ1 complex with a KD of < 50 nM as measured in a suitable in vitro binding assay such as Bio-Layer Interferometry (BLI). In some embodiments, such antibody binds a human LTBP1-proTGFβ1 complex with a KD of < 25 nM as measured in a suitable in vitro

binding assay such as Bio-Layer Interferometry (BLI); and/or, such antibody binds a human LTBP3-proTGF β 1 complex with a KD of < 25 nM as measured in a suitable in vitro binding assay such as Bio-Layer Interferometry (BLI). In some embodiments, such antibody binds a human LTBP1-proTGF β 1 complex with a KD of < 10 nM as measured in a suitable in vitro binding assay such as Bio-Layer Interferometry (BLI); and/or, such antibody binds a human LTBP3-proTGF β 1 complex with a KD of < 10 nM as measured in a suitable in vitro binding assay such as Bio-Layer Interferometry (BLI).

[366] In another embodiment, such antibody is cross-reactive with mouse LTBP1-proTGF β 1. In some embodiments, such antibody is also cross-reactive with mouse LTBP3-proTGF β 1. In some embodiments, such antibody binds a mouse LTBP1-proTGF β 1 complex with a KD of < 100 nM as measured in a suitable in vitro binding assay such as Bio-Layer Interferometry (BLI); and/or, the antibody binds a mouse LTBP3-proTGF β 1 complex with a KD of < 100 nM as measured in a suitable in vitro binding assay such as Bio-Layer Interferometry (BLI). In some embodiments, such antibody binds a mouse LTBP1-proTGF β 1 complex with a KD of < 50 nM as measured in a suitable in vitro binding assay such as Bio-Layer Interferometry (BLI); and/or, the antibody binds a mouse LTBP3-proTGF β 1 complex with a KD of < 50 nM as measured in a suitable in vitro binding assay such as Bio-Layer Interferometry (BLI). In some embodiments, such antibody binds a mouse LTBP1-proTGF β 1 complex with a KD of < 25 nM as measured in a suitable in vitro binding assay such as Bio-Layer Interferometry (BLI); and/or, the antibody binds a mouse LTBP3-proTGF β 1 complex with a KD of < 25 nM as measured in a suitable in vitro binding assay such as Bio-Layer Interferometry (BLI). In some embodiments, such antibody binds a mouse LTBP1-proTGF β 1 complex with a KD of < 10 nM as measured in a suitable in vitro binding assay such as Bio-Layer Interferometry (BLI); and/or, the antibody binds a mouse LTBP3-proTGF β 1 complex with a KD of < 10 nM as measured in a suitable in vitro binding assay such as Bio-Layer Interferometry (BLI).

[367] In another embodiment, such antibody does not bind to human GARP-proTGF β 1. In preferred embodiments, such context-selective antibody is also isoform-specific in that it selectively binds and inhibits the activation of TGF β 1 associated with LTBP1/3.

[368] In another aspect, the antibody, or antigen-binding fragment thereof, comprises a CDR-H1 comprising the amino acid G(X₁)I(X₂)SSSYW(X₃), wherein, optionally: X₁ is S or P; X₂ is H or R; and, X₃ is G, I or N. In some embodiments, X₁ is a S. In some embodiments, X₁ is a P. In some embodiments, X₂ is a H. In some embodiments, X₂ is an R. In some embodiments, X₃ is a G. In some embodiments, X₃ is an I. In some embodiments, X₃ is an N.

[369] In some embodiments, the antibody, or antigen-binding fragment thereof, comprises a CDR-H2 comprising the amino acid sequence SISYA(X₁)TYNPSLKS, wherein, optionally, X₁ is S or T. In some embodiments, X₁ is a S. In some embodiment, X₁ is a T.

[370] In some embodiments, the antibody, or antigen-binding fragment thereof, comprises a CDR-H3 comprising the amino acid sequence A(X₁)D(X₂)SYD(X₃)(X₄)AGM(X₅)(X₆), wherein, optionally: X₁ is R, S or G; X₂ is P or V; X₃ is S or A; X₄ is A, R, I or V; X₅ D, Q, or G; and, X₆ is V or R. In some embodiments, X₁ is R. In some embodiments, X₁ is S. In some embodiments, X₁ is G. In some embodiments, X₂ is P. In some embodiments, X₂ is V. In some embodiments, X₃ is S. In some embodiments, X₃ is A. In some embodiments, X₄ is A. In some embodiments, X₄ is R. In some embodiments, X₄ is I. In some embodiments, X₄ is V. In some embodiments, X₅ is D. In some embodiments, X₅ is Q. In some embodiments, X₅ is G. In some embodiments, X₆ is V. In some embodiments, X₆ is R.

[371] In some embodiments, the antibody, or antigen-binding fragment thereof, comprises three heavy chain CDRs and three light chain CDRs, wherein the heavy chain CDRs comprise:

- a) CDR-H1 comprising the amino acid sequence G(X₁)I(X₂)SSSYW(X₃), wherein, optionally: X₁ is S or P; X₂ is H or R; and, X₃ is G, I or N;
- b) CDR-H2 comprising the amino acid sequence SISYSA(X₁)TYYNPSLKS, wherein, optionally, X₁ is S or T; and
- c) CDR-H3 comprising the amino acid sequence A(X₁)D(X₂)SYD(X₃)(X₄)AGM(X₅)(X₆), wherein, optionally: X₁ is R, S or G; X₂ is P or V; X₃ is S or A; X₄ is A, R, I or V; X₅ D, Q, or G; and, X₆ is V or R.

[372] In some embodiments, the antibody, or antigen-binding fragment thereof, further comprises a CDR-L1 as set forth in SEQ ID NO:103, optionally comprising one or more amino acid changes. In some embodiment, the antibody, or antigen-binding fragment thereof, further comprises a CDR-L2 as set forth in SEQ ID NO:104, optionally comprising one or more amino acid changes. In some embodiments, the antibody, or antigen-binding fragment thereof, further comprises a CDR-L3 as set forth in SEQ ID NO:105, optionally comprising one or more amino acid changes.

[373] In some embodiments, the antibody, or antigen-binding fragment, comprises at least three of the following six CDRs:

- a) CDR-H1 comprising the amino acid sequence G(X₁)I(X₂)SSSYW(X₃), wherein, optionally: X₁ is S or P; X₂ is H or R; and, X₃ is G, I or N;
- b) CDR-H2 comprising the amino acid sequence SISYSA(X₁)TYYNPSLKS, wherein, optionally, X₁ is S or T;
- c) CDR-H3 comprising the amino acid sequence A(X₁)D(X₂)SYD(X₃)(X₄)AGM(X₅)(X₆), wherein, optionally: X₁ is R, S or G; X₂ is P or V; X₃ is S or A; X₄ is A, R, I or V; X₅ D, Q, or G; and, X₆ is V or R;
- d) CDR-L1: SEQ ID NO:103, optionally comprising one or more amino acid changes;
- e) CDR-L2: SEQ ID NO: 104, optionally comprising one or more amino acid changes; and

- f) CDR-L3: SEQ ID NO: 105, optionally comprising one or more amino acid changes.

[374] In some embodiments, the antibody, or antigen-binding fragment thereof, specifically binds a human LTBP1-proTGFβ1 complex and/or a human LTBP3-proTGFβ1 complex, and does not bind a human GARP-proTGFβ1 complex. In some embodiments, the antibody, or antigen-binding fragment thereof, does not bind a human GARP-proTGFβ1 complex. In some embodiments, the antibody, or antigen-binding fragment thereof, does not bind mature TGFβ1, mature TGFβ2 or mature TGFβ3.

[375] In a particular embodiment, the antibody, or antigen-binding fragment thereof, specifically binds a human LTBP1-proTGFβ1 complex and/or a human LTBP3-proTGFβ1 complex, and does not bind a human GARP-proTGFβ1 complex; wherein the antibody does not bind mature TGFβ1, mature TGFβ2 or mature TGFβ3; wherein the antibody is a fully human or humanized antibody or a fragment thereof, wherein the antibody comprises at least three of the following six CDRs:

- a) CDR-H1 comprising the amino acid sequence G(X₁)I(X₂)SSSYYW(X₃), wherein, optionally: X₁ is S or P; X₂ is H or R; and, X₃ is G, I or N;
- b) CDR-H2 comprising the amino acid sequence SISYSA(X₁)TYYNPSLKS, wherein, optionally, X₁ is S or T;
- c) CDR-H3 comprising the amino acid sequence A(X₁)D(X₂)SYD(X₃)(X₄)AGM(X₅)(X₆), wherein, optionally: X₁ is R, S or G, X₂ is P or V; X₃ is S or A; X₄ is A, R, I or V; X₅ D, Q, or G; and, X₆ is V or R;
- d) CDR-L1: SEQ ID NO:103, optionally comprising one or more amino acid changes;
- e) CDR-L2: SEQ ID NO: 104, optionally comprising one or more amino acid changes; and
- f) CDR-L3: SEQ ID NO: 105, optionally comprising one or more amino acid changes.

[376] In some embodiments, such antibody binds a human LTBP1-proTGFβ1 complex with a KD of < 100 nM as measured in a suitable in vitro binding assay such as Bio-Layer Interferometry (BLI); and/or, such antibody binds a human LTBP3-proTGFβ1 complex with a KD of < 100 nM as measured in a suitable in vitro binding assay such as Bio-Layer Interferometry (BLI). In some embodiments, such antibody binds a human LTBP1-proTGFβ1 complex with a KD of < 50 nM as measured in a suitable in vitro binding assay such as Bio-Layer Interferometry (BLI); and/or, such antibody binds a human LTBP3-proTGFβ1 complex with a KD of < 50 nM as measured in a suitable in vitro binding assay such as Bio-Layer Interferometry (BLI). In some embodiments, such antibody binds a human LTBP1-proTGFβ1 complex with a KD of < 25 nM as measured in a suitable in vitro binding assay such as Bio-Layer Interferometry (BLI); and/or, such antibody binds a human LTBP3-proTGFβ1 complex with a KD of < 25 nM as measured in a suitable in vitro binding assay such as Bio-Layer Interferometry (BLI). In some embodiments, such antibody binds a human LTBP1-proTGFβ1 complex with a KD of < 10 nM as measured in a suitable in vitro binding assay such as Bio-Layer Interferometry (BLI); and/or, such antibody binds a human LTBP3-proTGFβ1 complex

with a KD of < 10 nM as measured in a suitable in vitro binding assay such as Bio-Layer Interferometry (BLI).

[377] In another embodiment, such antibody is cross-reactive with mouse LTBP1-proTGFβ1. In some embodiments, such antibody is also cross-reactive with mouse LTBP3-proTGFβ1. In some embodiments, such antibody binds a mouse LTBP1-proTGFβ1 complex with a KD of < 100 nM as measured in a suitable in vitro binding assay such as Bio-Layer Interferometry (BLI); and/or, the antibody binds a mouse LTBP3-proTGFβ1 complex with a KD of < 100 nM as measured in a suitable in vitro binding assay such as Bio-Layer Interferometry (BLI). In some embodiments, such antibody binds a mouse LTBP1-proTGFβ1 complex with a KD of < 50 nM as measured in a suitable in vitro binding assay such as Bio-Layer Interferometry (BLI); and/or, the antibody binds a mouse LTBP3-proTGFβ1 complex with a KD of < 50 nM as measured in a suitable in vitro binding assay such as Bio-Layer Interferometry (BLI). In some embodiments, such antibody binds a mouse LTBP1-proTGFβ1 complex with a KD of < 25 nM as measured in a suitable in vitro binding assay such as Bio-Layer Interferometry (BLI); and/or, the antibody binds a mouse LTBP3-proTGFβ1 complex with a KD of < 25 nM as measured in a suitable in vitro binding assay such as Bio-Layer Interferometry (BLI). In some embodiments, such antibody binds a mouse LTBP1-proTGFβ1 complex with a KD of < 10 nM as measured in a suitable in vitro binding assay such as Bio-Layer Interferometry (BLI); and/or, the antibody binds a mouse LTBP3-proTGFβ1 complex with a KD of < 10 nM as measured in a suitable in vitro binding assay such as Bio-Layer Interferometry (BLI).

[378] In another embodiment, such antibody does not bind to human GARP-proTGFβ1. In preferred embodiments, such context-selective antibody is also isoform-specific in that it selectively binds and inhibits the activation of TGFβ1 associated with LTBP1/3.

[379] Also provided herein are antibodies, or antigen-binding fragments thereof, which binds a human LTBP1-proTGFβ1 complex and/or a human LTBP3-TGFβ1 complex with yet higher affinities and further advantageous combinations of binding properties.

[380] Accordingly, in one aspect, the antibody, or antigen-binding fragment thereof, comprises the following six CDRs:

- a) CDR-H1 comprising the amino acid sequence FTFRSYVMH;
- b) CDR-H2 comprising the amino acid sequence VISHEGS(X₁)KYYADSVKG, wherein: X₁ is L or G; and
- c) CDR-H3 comprising the amino acid sequence A(X₁)PRIAARRGGFG(X₂), wherein: X₁ is V, R or L; and X₂ is Y, S or T;
- d) CDR-L1 comprising the amino acid sequence TRS(X₁)G(X₂)ID(X₃)NYVQ, wherein, X₁ is S or H; X₂ is N, L, S or A; and X₃ is N, D or Y;

e) CDR-L2 comprising the amino acid sequence ED(X₁)(X₂)RPS, wherein: X₁ is N, F or A; and X₂ is Q, I or V; and

f) CDR-L3 comprising the amino acid sequence Q(X₁)YD(X₂)(X₃)(X₄)Q(X₅)VV, wherein: X₁ is S or G; X₂ is S, F, Y, D, H or W; X₃ is N, D or S; X₄ is N, A, L, E or T; and X₅ is G, R, A or L.

[381] In some embodiments, within CDR-H3: X₁ is R or L. Within CDR-L3: X₂ may be Y. Within CDR-L3: X₃ may be D; and X₄ may be T. In some preferred embodiments, within CDR-H3: X₁ is R or L (optionally R), within CDR-L3: X₂ is Y; and within CDR-L3: X₃ is D; and X₄ is T.

[382] In alternative embodiments, within CDR-H3: X₁ is R or L (optionally R), within CDR-L3: X₂ is Y and within CDR-L3: X₃ is D; X₄ is N; and X₅ is A.

[383] In some embodiments within CDR-L1: X₁ is S or H; X₂ is N or A; and X₃ is N, D or Y; within CDR-L2: X₁ is N or F; and X₂ is Q or V; and within CDR-L3: X₁ is S or G; X₂ is S, Y, D or W; X₃ is D or S; X₄ is N, L or T; and X₅ is G, R, A or L. In some embodiments, within CDR-L1: X₁ is S; X₂ is N; and X₃ is N or Y; within CDR-L2: X₁ is N; and X₂ is Q or V; and within CDR-L3: X₁ is S or G; X₂ is S, Y or W; X₃ is D; X₄ is N or T; and X₅ is G, R or A. In some embodiments, within CDR-L3: X₁ is S; X₂ is S or Y; X₃ is D; X₄ is N or T; and X₅ is G, R or A. In some embodiments, within CDR-L3: X₁ is S; X₂ is Y; X₃ is D; X₄ is N or T; and X₅ is G or A. In some preferred embodiments, within CDR-L3: X₁ is S; X₂ is Y; X₃ is D; X₄ is T; and X₅ is G.

[384] In particularly preferred embodiments, the antibody or antigen-binding fragment has the CDRs of Ab42, *e.g.*: CDR-H1 comprising the amino acid sequence FTFRSYVMH (SEQ ID NO: 166); CDR-H2 comprising the amino acid sequence VISHEGSLKYYADSVKG (SEQ ID NO: 167); CDR-H3 comprising the amino acid sequence ARPRIAARRGGFGY (SEQ ID NO: 168); CDR-L1 comprising the amino acid sequence TRSSGNIDNNYVQ (SEQ ID NO: 169); CDR-L2 comprising the amino acid sequence EDNQRPS (SEQ ID NO: 170); and CDR-L3 comprising the amino acid sequence QSYDYDTQGVV (SEQ ID NO: 171).

[385] In some embodiments, the antibody or antigen-binding fragment comprises a heavy chain variable region having an amino acid sequence that is at least 90% identical to SEQ ID NO: 318; and a light chain variable region having an amino acid sequence that is at least 90% identical to SEQ ID NO: 319.

[386] In some embodiments, the antibody, or antigen-binding fragment, binds a human LTBP1-proTGFβ1 complex and/or a human LTBP3-TGFβ1 complex with a K_D of < 5 nM as measured by suitable in vitro binding assay, such as BLI. For example, the antibody, or antigen-binding fragment, may bind a human LTBP1-proTGFβ1 complex and a human LTBP3-TGFβ1 complex with a K_D of < 5 nM as measured by suitable in vitro binding assay, such as BLI. In some embodiments, the

antibody, or antigen-binding fragment, binds the human LTBP1- and/or LTBP3-proTGF β 1 complex with a K_D of < 1 nM as measured by suitable in vitro binding assay, such as BLI.

[387] In some embodiments, the antibody, or antigen-binding fragment thereof does not show detectable binding to a human GARP-proTGF β 1 complex, as measured by BLI, under the same assay conditions as used to measure binding to human LTBP1-proTGF β 1 complex and/or human LTBP3-TGF β 1. For example, the antibody, or antigen-binding fragment may not show detectable binding to a human GARP-proTGF β 1 complex, as measured by BLI, under the same assay conditions as used to measure binding to human LTBP1-proTGF β 1 complex and human LTBP3-TGF β 1 complex.

[388] In some embodiments, the antibody, or antigen-binding fragment thereof, binds a human LTBP1-proTGF β 1 complex and/or a human LTBP3-TGF β 1 complex with a K_D that is at least 50 times lower (*e.g.*, at least 75 times lower, at least 100 times lower) than the K_D when binding to a human GARP-proTGF β 1 complex under the same assay conditions. For example, the antibody, or antigen-binding fragment thereof, may bind a human LTBP1-proTGF β 1 complex and a human LTBP3-TGF β 1 complex with a K_D that is at least 50 times lower (*e.g.*, at least 75 times lower, at least 100 times lower) than the K_D when binding to a human GARP-proTGF β 1 complex under the same assay conditions. In some embodiments, K_D is as determined by BLI or SPR. In some embodiments, K_D is as determined by SPR.

[389] In some embodiments, the antibody, or antigen-binding fragment thereof does not show detectable binding to an LRRC33-proTGF β 1 latent complex, as measured by BLI, under the same assay conditions as used to measure binding to human LTBP1-proTGF β 1 complex and/or human LTBP3-TGF β 1. For example, the antibody, or antigen-binding fragment thereof may not show detectable binding to an LRRC33-proTGF β 1 latent complex, as measured by BLI, under the same assay conditions as used to measure binding to human LTBP1-proTGF β 1 complex and human LTBP3-TGF β 1.

[390] In some embodiments, the antibody, or antigen-binding fragment thereof, binds a human LTBP1-proTGF β 1 complex and/or a human LTBP3-TGF β 1 complex with a K_D that is at least 50 times lower (*e.g.*, at least 75 times lower, at least 100 times lower) than the K_D when binding to a human LRRC33-proTGF β 1 complex under the same assay conditions. For example, the antibody, or antigen-binding fragment thereof, may bind a human LTBP1-proTGF β 1 complex and a human LTBP3-TGF β 1 complex with a K_D that is at least 50 times lower (*e.g.*, at least 75 times lower, at least 100 times lower) than the K_D when binding to a human LRRC33-proTGF β 1 complex under the same assay conditions. In some embodiments, K_D is as determined by BLI or SPR. In some embodiments, K_D is as determined by SPR.

[391] In some embodiments, the antibody, or antigen-binding fragment thereof, is cross-reactive with mouse LTBP1-proTGFβ1. In some embodiments, the antibody, or antigen-binding fragment thereof, is cross-reactive with mouse LTBP3-proTGFβ1. In some embodiments, the antibody, or antigen-binding fragment thereof, binds a mouse LTBP1-proTGFβ1 complex with a K_D of < 10 nM as measured by BLI. In some embodiments, the antibody, or antigen-binding fragment thereof, binds a mouse LTBP3-proTGFβ1 complex with a K_D of < 10 nM as measured by BLI.

[392] In further embodiments, the antibody which selectively binds a human LTBP1-TGFβ1 complex and/or a human LTBP3-TGFβ1 complex may not show meaningful binding (*e.g.*, may not show a response of more than 0.1 units (nm)) on exposure to a human GARP-proTGFβ1 complex in a BLI assay (*e.g.*, Octet) when the human GARP-proTGFβ1 complex is at a concentration of 200nM.

[393] In another aspect, the antibody, or antigen-binding fragment thereof, comprises the following six CDRs:

- a) CDR-H1 comprising the amino acid sequence G(X₁)I(X₂)S(X₃)SYYW(X₄), wherein, optionally: X₁ is S; X₂ is S, H or R; X₃ is S or G; and, X₄ is G, I, N or V;
- b) CDR-H2 comprising the amino acid sequence SISYS(X₁)(X₂)TYY, wherein, optionally: X₁ is G or A; and X₂ is S or T;
- c) CDR-H3 comprising the amino acid sequence A(X₁)DPSYDS(X₂)AGM(X₃)V, wherein, optionally: X₁ is R, S or G; X₂ is A or I; and X₃ is D or Q;
- d) CDR-L1 comprising the amino acid sequence RAS(X₁)(X₂)IS(X₃)YLN, wherein, optionally: X₁ is K or Q; X₂ is V or S; and X₃ is S or Y;
- e) CDR-L2 comprising the amino acid sequence (X₁)AS(X₂)(X₃)QS, wherein, optionally: X₁ is Y, A or S; X₂ is S or N; and X₃ is L or R;
- f) CDR-L3 comprising the amino acid sequence QQ(X₁)(X₂)D(X₃)P(X₄)T, wherein, optionally: X₁ is S or G; X₂ is F or N; X₃ is W or F; and X₄ is F or L.

[394] In some embodiments: within CDR-H1: X₁ is S; X₂ is S or R; X₃ is S; and, X₄ is G; within CDR-H2: X₁ is G or A; and X₂ is S or T; within CDR-H3: X₁ is R, S or G; X₂ is A or I; and X₃ is D or Q; within CDR-L1: X₁ is K or Q; X₂ is V or S; and X₃ is S or Y; within CDR-L2: X₁ is Y, A or S; X₂ is S or N; and X₃ is L or R; and within CDR-L3: X₁ is S or G; X₂ is F or N; X₃ is W or F; and X₄ is F or L.

[395] In some embodiments: CDR-H1 comprises the amino acid sequence GSIRSSSYWYG; CDR-H2 comprises the amino acid sequence SISYSATTYY; within CDR-H3: X₁ is S or G; X₂ is A or I; and X₃ is D or Q; within CDR-L1: X₁ is K or Q; X₂ is V or S; and X₃ is S or Y; within CDR-L2: X₁ is Y, A or S; X₂ is S or N; and X₃ is L or R; and within CDR-L3: X₁ is S or G; X₂ is F or N; X₃ is W or F; and X₄ is F or L.

[396] In some embodiments: CDR-H1 comprises the amino acid sequence GSIRSSSYWYG (SEQ ID NO: 292); CDR-H2 comprises the amino acid sequence SISYSATTYY (SEQ ID NO: 293); CDR-H3 comprises the amino acid sequence AGDPSYDSIAGMQV (SEQ ID NO: 294); CDR-L1 comprises the amino acid sequence RASQSISSYLN (SEQ ID NO: 295); CDR-L2 comprises the amino acid sequence AASNLQS (SEQ ID NO: 296); and CDR-L3 comprises the amino acid sequence QQSFDWPLT (SEQ ID NO: 297).

[397] In some embodiments, the antibody or antigen-binding fragment comprises a heavy chain variable region having an amino acid sequence that is at least 90% identical to SEQ ID NO: 360; and a light chain variable region having an amino acid sequence that is at least 90% identical to SEQ ID NO: 361.

[398] In some embodiments, the antibody, or antigen-binding fragment, binds a human LTBP1-proTGF β 1 complex and/or a human LTBP3-TGF β 1 complex with a K_D of < 5 nM as measured by suitable in vitro binding assay, such as BLI. For example, the antibody, or antigen-binding fragment, may bind a human LTBP1-proTGF β 1 complex and a human LTBP3-TGF β 1 complex with a K_D of < 5 nM as measured by suitable in vitro binding assay, such as BLI. In some embodiments, the antibody, or antigen-binding fragment, binds the human LTBP1- and/or LTBP3-proTGF β 1 complex with a K_D of < 1 nM as measured by suitable in vitro binding assay, such as BLI.

[399] In some embodiments, the antibody, or antigen-binding fragment thereof does not show detectable binding to a human GARP-proTGF β 1 complex, as measured by BLI, under the same assay conditions as used to measure binding to human LTBP1-proTGF β 1 complex and/or human LTBP3-TGF β 1. For example, the antibody, or antigen-binding fragment may not show detectable binding to a human GARP-proTGF β 1 complex, as measured by BLI, under the same assay conditions as used to measure binding to human LTBP1-proTGF β 1 complex and human LTBP3-TGF β 1 complex.

[400] In some embodiments, the antibody, or antigen-binding fragment thereof, binds a human LTBP1-proTGF β 1 complex and/or a human LTBP3-TGF β 1 complex with a K_D that is at least 50 times lower (*e.g.*, at least 75 times lower, at least 100 times lower) than the K_D when binding to a human GARP-proTGF β 1 complex under the same assay conditions. For example, the antibody, or antigen-binding fragment thereof, may bind a human LTBP1-proTGF β 1 complex and a human LTBP3-TGF β 1 complex with a K_D that is at least 50 times lower (*e.g.*, at least 75 times lower, at least 100 times lower) than the K_D when binding to a human GARP-proTGF β 1 complex under the same assay conditions. In some embodiments, K_D is as determined by BLI or SPR. In some embodiments, K_D is as determined by SPR.

[401] In some embodiments, the antibody, or antigen-binding fragment thereof does not show detectable binding to an LRRC33-proTGF β 1 latent complex, as measured by BLI, under the same assay conditions as used to measure binding to human LTBP1-proTGF β 1 complex and/or human

LTBP3-TGF β 1. For example, the antibody, or antigen-binding fragment thereof may not show detectable binding to an LRRC33-proTGF β 1 latent complex, as measured by BLI, under the same assay conditions as used to measure binding to human LTBP1-proTGF β 1 complex and human LTBP3-TGF β 1.

[402] In some embodiments, the antibody, or antigen-binding fragment thereof, binds a human LTBP1-proTGF β 1 complex and/or a human LTBP3-TGF β 1 complex with a K_D that is at least 50 times lower (*e.g.*, at least 75 times lower, at least 100 times lower) than the K_D when binding to a human LRRC33-proTGF β 1 complex under the same assay conditions. For example, the antibody, or antigen-binding fragment thereof, may bind a human LTBP1-proTGF β 1 complex and a human LTBP3-TGF β 1 complex with a K_D that is at least 50 times lower (*e.g.*, at least 75 times lower, at least 100 times lower) than the K_D when binding to a human LRRC33-proTGF β 1 complex under the same assay conditions. In some embodiments, K_D is as determined by BLI or SPR. In some embodiments, K_D is as determined by SPR.

[403] In some embodiments, the antibody, or antigen-binding fragment thereof, is cross-reactive with mouse LTBP1-proTGF β 1. In some embodiments, the antibody, or antigen-binding fragment thereof, is cross-reactive with mouse LTBP3-proTGF β 1. In some embodiments, the antibody, or antigen-binding fragment thereof, binds a mouse LTBP1-proTGF β 1 complex with a K_D of < 10 nM as measured by BLI. In some embodiments, the antibody, or antigen-binding fragment thereof, binds a mouse LTBP3-proTGF β 1 complex with a K_D of < 10 nM as measured by BLI.

[404] In further embodiments, the antibody which selectively binds a human LTBP1-TGF β 1 complex and/or a human LTBP3-TGF β 1 complex may not show meaningful binding (*e.g.*, may not show a response of more than 0.1 units (nm)) on exposure to a human GARP-proTGF β 1 complex in a BLI assay (*e.g.*, Octet) when the human GARP-proTGF β 1 complex is at a concentration of 200nM.

[405] In some embodiments, the “percent identity” of two amino acid sequences is determined using the algorithm of Karlin and Altschul Proc. Natl. Acad. Sci. USA 87:2264-68, 1990, modified as in Karlin and Altschul Proc. Natl. Acad. Sci. USA 90:5873-77, 1993. Such an algorithm is incorporated into the NBLAST and XBLAST programs (version 2.0) of Altschul, *et al.* J. Mol. Biol. 215:403-10, 1990. BLAST protein searches can be performed with the XBLAST program, score=50, word length=3 to obtain amino acid sequences homologous to the protein molecules of interest. Where gaps exist between two sequences, Gapped BLAST can be utilized as described in Altschul *et al.*, Nucleic Acids Res. 25(17):3389-3402, 1997. When utilizing BLAST and Gapped BLAST programs, the default parameters of the respective programs (*e.g.*, XBLAST and NBLAST) can be used.

[406] In any of the antibodies or antigen-binding fragments described herein, one or more conservative mutations can be introduced into the CDRs or framework sequences at positions where

the residues are not likely to be involved in an antibody-antigen interaction. In some embodiments, such conservative mutation(s) can be introduced into the CDRs or framework sequences at position(s) where the residues are not likely to be involved in interacting with a LTBP1-TGF β complex and/or a LTBP3-TGF β complex, as determined based on the crystal structure. In some embodiments, the likely interface (*e.g.*, residues involved in an antigen-antibody interaction) may be deduced from known structural information on another antigens sharing structural similarities.

[407] As used herein, a “conservative amino acid substitution” refers to an amino acid substitution that does not alter the relative charge or size characteristics of the protein in which the amino acid substitution is made. Variants can be prepared according to methods for altering polypeptide sequence known to one of ordinary skill in the art such as are found in references which compile such methods, *e.g.*, *Molecular Cloning: A Laboratory Manual*, J. Sambrook, *et al.*, eds., Second Edition, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York, 1989, or *Current Protocols in Molecular Biology*, F.M. Ausubel, *et al.*, eds., John Wiley & Sons, Inc., New York. Conservative substitutions of amino acids include substitutions made amongst amino acids within the following groups: (a) M, I, L, V; (b) F, Y, W; (c) K, R, H; (d) A, G; (e) S, T; (f) Q, N; and (g) E, D.

[408] In some embodiments, the antibodies provided herein comprise mutations that confer desirable properties to the antibodies. For example, to avoid potential complications due to Fab-arm exchange, which is known to occur with native IgG4 mAbs, the antibodies provided herein may comprise a stabilizing ‘Adair’ mutation (Angal *et al.*, “A single amino acid substitution abolishes the heterogeneity of chimeric mouse/human (IgG4) antibody,” *Mol Immunol* 30, 105-108; 1993), where serine 228 (EU numbering; residue 241 Kabat numbering) is converted to proline resulting in an IgG1-like (CPPCP (SEQ ID NO: 45)) hinge sequence. Accordingly, any of the antibodies may include a stabilizing ‘Adair’ mutation or the amino acid sequence CPPCP (SEQ ID NO: 45). In one embodiment, an antibody described herein comprises a heavy chain immunoglobulin constant domain of a human IgG₄ having a backbone substitution of Ser to Pro, that produces an IgG₁-like hinge and permits formation of inter-chain disulfide bonds.

[409] Antibodies of this disclosure that selectively bind to a LTBP1-TGF β complex and/or a LTBP3-TGF β complex may optionally comprise antibody constant regions or parts thereof. For example, a V_L domain may be attached at its C-terminal end to a light chain constant domain such as C κ or C λ . Similarly, a V_H domain or portion thereof may be attached to all or part of a heavy chain such as IgA, IgD, IgE, IgG, and IgM, and any isotype subclass. Antibodies may include suitable constant regions (see, for example, Kabat *et al.*, *Sequences of Proteins of Immunological Interest*, No. 91-3242, National Institutes of Health Publications, Bethesda, Md. (1991)). Therefore, antibodies within the scope of this may disclosure include V_H and V_L domains, or antigen-binding portions thereof, combined with any suitable constant region. In exemplary embodiments, the antibodies, or antigen-binding portions thereof, comprise a heavy chain immunoglobulin constant domain

containing all or a portion of a human IgG₁ or a human IgG₄ constant domain. In some embodiments, the antibody, or antigen-binding portion thereof, comprises a light chain immunoglobulin constant domain containing all or a portion of a human Ig lambda constant domain or a human Ig kappa constant domain.

[410] In some embodiments, antibodies that selectively bind to a LTBP1-TGF β complex and/or a LTBP3-TGF β complex may or may not include the framework region of the antibodies of SEQ ID NOs: 7 and 8. In some embodiments, antibodies that selectively bind to a LTBP1-TGF β complex and/or a LTBP3-TGF β complex are murine antibodies and include murine framework region sequences. In other embodiments, the antibodies are chimeric antibodies, or antigen-binding fragments thereof. In another embodiment, the antibodies are humanized antibodies, or antigen-binding fragments thereof. In another embodiment, the antibodies are fully human antibodies, or antigen-binding fragments thereof. In one embodiment, the antibody comprises a framework region comprising a human germline amino acid sequence.

[411] The antibodies, and antigen-binding fragments thereof, described herein can have any configuration suitable for binding antigen. For example, in one embodiment, the antibody, or antigen-binding portion thereof, comprises four polypeptide chains, including two heavy chain variable regions and two light chain variable regions. In another embodiment, the antibody, or antigen-binding portion thereof, comprises one heavy chain variable region and one light chain variable region. In exemplary embodiments, the antibody, or antigen-binding portion thereof, is a Fab fragment, a F(ab')₂ fragment, a scFab fragment, an scFv, or a diabody.

[412] In one embodiment, the antibody, or antigen-binding portion thereof, comprises a heavy chain immunoglobulin constant domain of a human IgG₁ constant domain or a human IgG₄ constant domain. In an exemplary embodiment, the heavy chain immunoglobulin constant domain is a human IgG₄ constant domain. In one embodiment, the antibody, or antigen-binding portion thereof, binds a conformational epitope. In one embodiment, the antibody, or antigen-binding portion thereof, binds a combinatorial epitope.

[413] In one embodiment, the antibody, or antigen-binding portion thereof, comprises a heavy chain immunoglobulin constant domain of a human IgG₄ constant domain having a backbone substitution of Ser to Pro that produces an IgG₁-like hinge and permits formation of inter-chain disulfide bonds. In one embodiment, the antibody, or antigen-binding portion thereof, further comprises a light chain immunoglobulin constant domain comprising a human Ig lambda constant domain, or a human Ig kappa constant domain.

[414] In one embodiment, the antibody is an IgG having four polypeptide chains which are two heavy chains and two light chains. In exemplary embodiments, the antibody can be a humanized

antibody, a human antibody, or a chimeric antibody. In one embodiment, the antibody comprises a framework having a human germline amino acid sequence.

[415] In one embodiment, the invention provides an antibody, or antigen-binding portion thereof, that competes for binding with an antibody, or antigen-binding portion thereof, described herein. In one embodiment, the invention provides an antibody, or antigen-binding portion thereof, that binds to the same epitope as an antibody, or antigen-binding portion thereof, described herein. In one embodiment, the antibody, or antigen-binding fragment thereof, does not compete with antibody SR-Ab1 for binding to a human LTBP1-proTGFβ1 complex.

Binding kinetics of novel antibodies

[416] The novel antibodies and antigen-binding fragments thereof (*e.g.*, Fabs) disclosed herein are characterized by enhanced binding properties. The antibodies and the fragments are capable of selectively binding to a LTBP1-TGFβ1 complex and/or a LTBP3-TGFβ1 complex (also known as large latent complexes (LLCs)). Recombinantly produced, purified protein complexes may be used as antigens (*e.g.*, antigen complexes) to screen, evaluate or confirm the ability of an antibody to bind the antigen complexes in suitable in vitro binding assays. Such assays are well known in the art and include but are not limited to: BLI-based assays (such as Octet®) and SPR-based assays (such as Biacore).

[417] Previously, antibodies and fragments that exhibited high affinities (*e.g.*, sub-nanomolar K_D) to the LLCs were identified. Here, advantageously, antibodies and fragments with particularly slow dissociation rates were specifically selected, aimed to achieve particularly durable inhibitory effects.

[418] Accordingly, selection of suitable TGFβ inhibitors for carrying out the methods and therapeutic use in accordance with the present disclosure may include carrying out in vitro binding assays to measure binding kinetics. In preferred embodiments, the antibody or the antigen-binding fragment binds hLTBP1-pro TGFβ1 and/or hLTBP3-proTGFβ1 with high affinity and low dissociation rate k_{OFF} , as described herein. Preferably, the antibody or the fragment further binds the murine LLC counterparts, namely, mLTBP1-proTGFβ1 and/or mLTBP3-proTGFβ1, with equivalent affinities as the human LLCs. In vitro binding kinetics may be readily determined by measuring interactions of test antibodies (such as antigen-binding fragments) and suitable antigen, such as LLCs and small latent complexes (SLCs). Suitable methods for in vitro binding assays to determine the parameters of binding kinetics include BLI-based assays such as Octet, and surface plasmon resonance-based assays, such as Biacore systems. An example of an Octet-based in vitro binding assay is provided in Example 9 / Table 9. Several antibodies were shown in this experiment to have “OFF” rates (k_{OFF}) that are $\leq 5 \times 10^{-4}$ (1/s). These results are in stark contrast to the results shown for Ab10, for which binding to hLTBP1-pro TGFβ1 and/or hLTBP3-proTGFβ1 could not even be detected by the same Octet assay (Table 8). An example of SPR-based in vitro binding assay is

provided in Example 9. Fab fragments of Ab42, Ab63 and Ab43, which are activation inhibitors of TGF β 1, were used in this experiment. As illustrated in this example, these antibodies have sub-nanomolar K_D and “OFF” rates that are $\leq 5 \times 10^{-4}$ (1/s). Thus, for example, Ab42 is able to remain bound to the antigen for a much longer duration of time (*e.g.*, greater $t_{1/2}$) than an antibody with much higher “OFF” rate, which “falls off” (*e.g.*, dissociates from) the antigen relatively quickly. Thus, the difference in the dissociation kinetics predominantly attributes to the notable difference in their overall affinities (K_D), which may result in enhanced potency. Therefore, characterization of binding kinetics provides useful information as to potential durability of effects and resulting *in vivo* potency.

[419] Accordingly, the invention includes a method of selecting a TGF β activation inhibitor for therapeutic use, wherein the method comprises selection of an antibody or antigen-binding fragment thereof that has a dissociation rate of $\leq 5 \times 10^{-4}$ (1/s) as measured by SPR. In some embodiments, the antibody or the fragment binds antigen with an affinity of less than 1 nM (*i.e.*, sub-nanomolar), *e.g.*, less than 500 pM, 400 pM, 300 pM, 200 pM, 100 pM, or 50 pM.

[420] The selection method may include determining or measuring dissociation half-time (also referred to as half binding time or $t_{1/2}$) of test antibody by suitable means, such as BLI-based assays and SPR-based assays. Monovalent or multivalent (*e.g.*, divalent) test antibodies may be used. For example, Fab fragments are suitable monovalent antibodies that can be used to determine dissociation half time. Similarly, full length immunoglobulins (*e.g.*, IgGs) may be used to determine dissociation half time. In some embodiments, the method comprises selection of an antibody or antigen-binding fragment thereof that has $t_{1/2}$ of 45 minutes or longer to human LTBP1-proTGF β 1 and/or human LTBP3-proTGF β 1. Preferably, the antibody or the fragment has $t_{1/2}$ of 45 minutes or longer to each of human LTBP1-proTGF β 1 and human LTBP3-proTGF β 1. More preferably, the antibody or the fragment further has $t_{1/2}$ of 45 minutes or longer to murine LTBP1-proTGF β 1 and/or murine LTBP3-proTGF β 1. Most preferably, the antibody or the fragment has $t_{1/2}$ of 45 minutes or longer to each of murine LTBP1-proTGF β 1 and murine LTBP3-proTGF β 1. The method may further include selection of an antibody or an antigen-binding fragment that has $t_{1/2}$ of 5 minutes or less to human GARP-proTGF β 1. Any of these antibodies with advantageous dissociation half time should preferably have a K_D of less than 1 nM as measured by BLI (*e.g.*, Octet) or SPR (*e.g.*, Biacore). Preferably, SPR assays are used to determine dissociation half-time ($t_{1/2}$).

Polypeptides

[421] Some aspects of the disclosure relate to isolated polypeptides. For example, in one embodiment, the invention provides an isolated polypeptide comprising CDRH1, CDRH2, CDRH3, CDRL1, CDRL2, or CDRL3, or combinations thereof, as provided in Table 5. In an exemplary embodiment, the isolated polypeptide can contain CDRH1, CDRH2, and CDRH3 as provided in

Table 5. In other embodiments, the isolated polypeptide can contain CDRL1, CDRL2, and CDRL3 as provided in Table 5. In some embodiments, the polypeptide can contain up to 6, 5, 4, 3, 2, or 1 amino acid residue variations as compared to the corresponding CDR region in any one of CDRH1, CDRH2, CDRH3, CDRL1, CDRL2, or CDRL3, or combinations thereof, as provided in Table 5. In one embodiment, the invention provides an isolated polypeptide comprising SEQ ID NO: 7. In another embodiment, the invention provides an isolated polypeptide comprising SEQ ID NO: 8. In another embodiment, the invention provides an isolated polypeptide comprising SEQ ID NO:7 and SEQ ID NO:8. In this embodiment, SEQ ID NO:7 and SEQ ID NO:8 can optionally be connected by a linker peptide. In some embodiments, the polypeptide is a heavy chain variable domain. In some embodiments, the polypeptide is at least 75% (*e.g.*, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99%) identical to SEQ ID NO: 7. In some embodiments, the polypeptide is a light chain variable domain. In some embodiments, the polypeptide is at least 75% (*e.g.*, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99%) identical to SEQ ID NO: 8.

[422] In another embodiment, the invention provides an isolated polypeptide comprising a heavy chain variable region sequence set forth in Table 6. In one embodiment, the invention provides an isolated polypeptide comprising SEQ ID NO: 7, SEQ ID NO:74, SEQ ID NO:76, SEQ ID NO:78, SEQ ID NO:80, SEQ ID NO:82, SEQ ID NO:84, SEQ ID NO:86, SEQ ID NO:88, SEQ ID NO:90, SEQ ID NO:92, or SEQ ID NO: 106. In one embodiment, the invention provides an isolated polypeptide comprising SEQ ID NO: 8, SEQ ID NO:75, SEQ ID NO:77, SEQ ID NO:79, SEQ ID NO:81, SEQ ID NO:83, SEQ ID NO:85, SEQ ID NO:87, SEQ ID NO:89, SEQ ID NO:91, SEQ ID NO:93, or SEQ ID NO: 107. In one embodiment, the invention provides an isolated polypeptide comprising SEQ ID NO: 318. In one embodiment, the invention provides an isolated polypeptide comprising SEQ ID NO: 319. In one embodiment, the invention provides an isolated polypeptide comprising SEQ ID NO: 360. In one embodiment, the invention provides an isolated polypeptide comprising SEQ ID NO: 361.

[423] In another embodiment, the invention provides an isolated polypeptide comprising a light chain variable region set forth in Table 6. In another embodiment, the invention provides an isolated polypeptide comprising a heavy chain variable region sequence set forth in Table 6 (*e.g.*, SEQ ID NO: 7, SEQ ID NO:74, SEQ ID NO:76, SEQ ID NO:78, SEQ ID NO:80, SEQ ID NO:82, SEQ ID NO:84, SEQ ID NO:86, SEQ ID NO:88, SEQ ID NO:90, SEQ ID NO:92, or SEQ ID NO: 106) and a light chain variable region sequence set forth in Table 6 (*e.g.*, SEQ ID NO: 8, SEQ ID NO:75, SEQ ID NO:77, SEQ ID NO:79, SEQ ID NO:81, SEQ ID NO:83, SEQ ID NO:85, SEQ ID NO:87, SEQ ID NO:89, SEQ ID NO:91, SEQ ID NO:93, or SEQ ID NO: 107). In this embodiment, the heavy chain and light chain sequences (*e.g.*, SEQ ID NO:7 and SEQ ID NO:8) can optionally be connected by a linker peptide. In some embodiments, the polypeptide is a heavy chain variable domain. In some

embodiments, the polypeptide is at least 75% (*e.g.*, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99%) identical to SEQ ID NO: 7, SEQ ID NO:74, SEQ ID NO:76, SEQ ID NO:78, SEQ ID NO:80, SEQ ID NO:82, SEQ ID NO:84, SEQ ID NO:86, SEQ ID NO:88, SEQ ID NO:90, SEQ ID NO:92, or SEQ ID NO: 106. In some embodiments, the polypeptide is a light chain variable domain. In some embodiments, the polypeptide is at least 75% (*e.g.*, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99%) identical to SEQ ID NO: 8, SEQ ID NO:75, SEQ ID NO:77, SEQ ID NO:79, SEQ ID NO:81, SEQ ID NO:83, SEQ ID NO:85, SEQ ID NO:87, SEQ ID NO:89, SEQ ID NO:91, SEQ ID NO:93, or SEQ ID NO: 107.

[424] In another embodiment, the invention provides an isolated polypeptide comprising a heavy chain variable region sequence set forth in SEQ ID NO: 318 and a light chain variable region sequence set forth in SEQ ID NO: 319. In another embodiment, the invention provides an isolated polypeptide comprising a heavy chain variable region sequence set forth in SEQ ID NO: 360 and a light chain variable region sequence set forth in SEQ ID NO: 361. In this embodiment, the heavy chain and light chain sequences (*e.g.*, SEQ ID NO: 318 and SEQ ID NO: 319) can be connected by a linker peptide. In some embodiments, the polypeptide is a heavy chain variable domain. In some embodiments, the polypeptide is at least 85% (*e.g.*, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99%) identical to SEQ ID NO: 318 or SEQ ID NO: 360. In some embodiments, the polypeptide is a light chain variable domain. In some embodiments, the polypeptide is at least 85% (*e.g.*, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99%) identical to SEQ ID NO: 319 or SEQ ID NO: 361.

Nucleic Acids

[425] In some embodiments, antibodies, antigen-binding portions thereof, and/or compositions of the present disclosure may be encoded by nucleic acid molecules. Such nucleic acid molecules include, without limitation, DNA molecules, RNA molecules, polynucleotides, oligonucleotides, mRNA molecules, vectors, plasmids and the like. In some embodiments, the present disclosure may comprise cells programmed or generated to express nucleic acid molecules encoding compounds and/or compositions of the present disclosure.

[426] In some embodiments, the invention provides a nucleic acid molecule that encodes the foregoing antibodies, or an antigen-binding portion thereof. For example, in one embodiment, the invention provides a nucleic acid molecule that encodes a polypeptide comprising CDRH1, CDRH2, CDRH3, CDRL1, CDRL2, or CDRL3, or combinations thereof, as provided in Table 5. The nucleic acid molecule can, in some embodiments, encode a polypeptide comprising CDRH1, CDRH2, and CDRH3 as provided in Table 5. In some embodiments, the nucleic acid molecule can encode a

polypeptide comprising CDRL1, CDRL2, and CDRL3 as provided in Table 5. In some embodiments, the nucleic acid molecule encodes a polypeptide that can contain up to 6, 5, 4, 3, 2, or 1 amino acid residue variations as compared to the corresponding CDR region in any one of CDRH1, CDRH2, CDRH3, CDRL1, CDRL2, or CDRL3, or combinations thereof, as provided in Table 5. In an exemplary embodiment, the nucleic acid molecule encodes a polypeptide comprising a heavy chain variable domain having at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% identity to the amino acid sequence set forth in SEQ ID NO: 7, and/or a light chain variable domain having at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% identity to the amino acid sequence set forth in SEQ ID NO: 8. In one embodiment, the nucleic acid molecule encodes a polypeptide comprising a heavy chain variable domain having at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% identity to a heavy chain variable region sequence set forth in Table 6, and/or a light chain variable domain having at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% identity to a light chain variable region sequence set forth in Table 6. In an exemplary embodiment, the nucleic acid molecule encodes a polypeptide comprising a heavy chain variable domain having at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% identity to the amino acid sequence set forth in SEQ ID NO: 7, and/or a light chain variable domain having at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% identity to the amino acid sequence set forth in SEQ ID NO: 8. In some embodiments, the nucleic acid molecule encodes an antibody, or antigen-binding portion thereof, comprising a heavy chain variable domain amino acid sequence set forth in SEQ ID NO: 7, SEQ ID NO:74, SEQ ID NO:76, SEQ ID NO:78, SEQ ID NO:80, SEQ ID NO:82, SEQ ID NO:84, SEQ ID NO:86, SEQ ID NO:88, SEQ ID NO:90, SEQ ID NO:92, or SEQ ID NO: 106, and a light chain variable domain amino acid sequence set forth in SEQ ID NO: 8, SEQ ID NO:75, SEQ ID NO:77, SEQ ID NO:79, SEQ ID NO:81, SEQ ID NO:83, SEQ ID NO:85, SEQ ID NO:87, SEQ ID NO:89, SEQ ID NO:91, SEQ ID NO:93, or SEQ ID NO: 107. In some embodiments, the nucleic acid molecule encodes an antibody, or antigen-binding portion thereof, comprising a heavy chain variable domain amino acid sequence set forth in SEQ ID NO: 7, and a light chain variable domain amino acid sequence set forth in SEQ ID NO: 8.

[427] In another embodiment, the invention provides nucleic acid molecule that encodes the foregoing antibodies, or an antigen-binding portion thereof. In an exemplary embodiment, the nucleic acid molecule encodes a polypeptide comprising a heavy chain variable domain having at least 85%, 90%, 95%, 96%, 97%, 98%, or 99% identity to the amino acid sequence set forth in SEQ ID NO: 318, and/or a light chain variable domain having at least 85%, 90%, 95%, 96%, 97%, 98%, or 99% identity to the amino acid sequence set forth in SEQ ID NO: 319. In another embodiment, the nucleic acid molecule encodes a polypeptide comprising a heavy chain variable domain having at least 85%, 90%, 95%, 96%, 97%, 98%, or 99% identity to the amino acid sequence set forth in SEQ ID NO: 360,

and/or a light chain variable domain having at least 85%, 90%, 95%, 96%, 97%, 98%, or 99% identity to the amino acid sequence set forth in SEQ ID NO: 361.

[428] In another embodiment, the invention provides a nucleic acid molecule that encodes a polypeptide comprising a heavy chain variable region sequence set forth in SEQ ID NO: 318 and a light chain variable region sequence set forth in SEQ ID NO: 319. In another embodiment, the invention provides a nucleic acid molecule that encodes an isolated polypeptide comprising a heavy chain variable region sequence set forth in SEQ ID NO: 360 and a light chain variable region sequence set forth in SEQ ID NO: 361. In this embodiment, the heavy chain and light chain sequences (*e.g.*, SEQ ID NO: 318 and SEQ ID NO: 319) can be connected by a linker peptide. In some embodiments, the nucleic acid molecule encodes a polypeptide which is a heavy chain variable domain. In some embodiments, the nucleic acid molecule encodes a polypeptide which is at least 85% (*e.g.*, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99%) identical to SEQ ID NO: 318 or SEQ ID NO: 360. In some embodiments, the nucleic acid molecule encodes a polypeptide which is a light chain variable domain. In some embodiments, the nucleic acid molecule encodes a polypeptide which is at least 85% (*e.g.*, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99%) identical to SEQ ID NO: 319 or SEQ ID NO: 361. In some embodiments, the nucleic acid molecule encodes an antibody, or antigen-binding portion thereof, comprising a heavy chain variable domain amino acid sequence set forth in SEQ ID NO: 318, and a light chain variable domain amino acid sequence set forth in SEQ ID NO: 319. In some embodiments, the nucleic acid molecule encodes an antibody, or antigen-binding portion thereof, comprising a heavy chain variable domain amino acid sequence set forth in SEQ ID NO: 360, and a light chain variable domain amino acid sequence set forth in SEQ ID NO: 361.

[429] In some cases, nucleic acids of the disclosure include codon-optimized nucleic acids. Methods of generating codon-optimized nucleic acids are known in the art and may include, but are not limited to those described in US Patent Nos. 5,786,464 and 6,114,148, the contents of each of which are herein incorporated by reference in their entirety. Also provided herein are expression vectors comprising any of the aforementioned nucleic acid(s).

Production of Antibodies that Bind a LTBP1/3-TGF β 1 Complex

[430] The art is familiar with various techniques and methods that may be used for obtaining antibodies, or antigen-binding fragments thereof, of the disclosure. For example, antibodies can be produced using recombinant DNA methods, hybridoma techniques, phage or yeast display technologies, transgenic animals (*e.g.*, a XenoMouse®) or some combination thereof.

Immunization and Hybridomas

[431] In some methods described herein, the specified antigen (*e.g.*, an LTBP1-TGF β 1 complex and/or an LTBP3-TGF β 1 complex) can be used to immunize a non-human animal ("host"), *e.g.*, a rodent, *e.g.*, a mouse, hamster, or rat. In one embodiment, the non-human animal is a mouse. In another embodiment, the host may be a camelid.

[432] After immunization, which may include single or multiple steps of antigen exposures (*e.g.*, injections), splenocytes are harvested from the animal and the associated B cells are fused with immortalized myeloma cells to form hybridomas for antibody production. Hybridomas may be generated in accordance with known methods (see *e.g.*, Kohler and Milstein (1975) *Nature*, 256: 495-499). Hybridomas formed in this manner are then screened using standard methods, such as enzyme-linked immunosorbent assay (ELISA), Bio-Layer Interferometry (BLI) technology (*e.g.*, OCTET) and surface plasmon resonance (*e.g.*, BIACORE) analysis, to identify one or more hybridomas that produce an antibody that specifically binds to a specified antigen. Any form of the specified antigen may be used as the immunogen, *e.g.*, recombinant antigen, naturally occurring forms, any variants or fragments thereof, as well as antigenic peptide thereof (*e.g.*, any of the epitopes described herein as a linear epitope or within a scaffold as a conformational epitope).

Screening library/libraries

[433] In some embodiments, the method or process of making or identifying antibodies includes a step of screening protein expression libraries that express antibodies or fragments thereof (*e.g.*, scFv), *e.g.*, phage, yeast, or ribosome display libraries. For example, a library of human combinatorial antibodies or scFv fragments can be synthesized on phages or yeast, the library is then screened with the antigen of interest or an antibody-binding portion thereof, and the phage or yeast that binds the antigen is isolated, from which one may obtain the antibodies or immunoreactive fragments (Vaughan *et al.*, 1996, PMID: 9630891; Sheets *et al.*, 1998, PMID: 9600934; Boder *et al.*, 1997, PMID: 9181578; Pepper *et al.*, 2008, PMID: 18336206).

[434] Phage display is further described, for example, in Ladner *et al.*, U.S. Pat. No. 5,223,409; Smith (1985) *Science* 228:1315-1317; Clackson *et al.* (1991) *Nature*, 352: 624-628; Marks *et al.* (1991) *J. Mol. Biol.*, 222: 581-597; WO 92/18619; WO 91/17271; WO 92/20791; WO 92/15679; WO 93/01288; WO 92/01047; WO 92/09690; and WO 90/02809. Yeast display is further described, for example, in US 7,700,302 and US 8,877,688. In particular methods, the yeast display library expresses full-length antibodies (*e.g.*, Adimab, LLC).

[435] Kits for generating phage or yeast display libraries are commercially available. There also are other methods and reagents that can be used in generating and screening antibody display libraries (see US 5,223,409; WO 92/18619, WO 91/17271, WO 92/20791, WO 92/15679, WO 93/01288, WO

92/01047, WO 92/09690; and Barbas et al., 1991, PMID: 1896445). Such techniques advantageously allow for the screening of large numbers of candidate antibodies.

[436] No matter how obtained, antibody producing cells (*e.g.*, yeast colonies, hybridomas, etc.) may be selected, cloned and further screened for desirable characteristics including, for example, robust growth, high antibody production and desirable antibody characteristics such as high affinity for the antigen of interest. Methods of selecting, cloning and expanding colonies and/or hybridomas are well known to those of ordinary skill in the art. Once the desired antibodies are identified, the relevant genetic material may be isolated, manipulated, and expressed using common, art-recognized molecular biology and biochemical techniques.

Humanization

[437] The antibodies or fragments of the present invention are preferably fully human antibodies or humanized antibodies. Thus, whatever the source, it will be appreciated that the method may comprise humanizing one or more antibodies or fragments thereof, wherein the human antibody sequences may be fabricated using art-known molecular engineering techniques and introduced into expression systems and host cells as described herein. Such non-natural recombinantly produced human antibodies (and subject compositions) are entirely compatible with the teachings of this disclosure and are expressly held to be within the scope of the instant invention. In certain select aspects, the LTBP1-TGF β 1 complex-binding and/or a LTBP3-TGF β 1 complex-binding antibodies of the invention will comprise a recombinantly produced human antibody.

Affinity maturation

[438] In some embodiments, antibodies produced by the methods described-above may be of moderate affinity (K_a of about 10^6 to 10^7 M⁻¹). Accordingly, antibodies or fragments thereof may be subjected to a process of affinity maturation as part of optimization, if desired. The term “affinity maturation” shall have the meaning readily understood by the skilled artisan. Briefly, it refers to further modifying the amino acid sequence of candidate antibodies or fragments (often referred to as “parent”) to achieve improved binding profiles to the specific antigen. Typically, a parental antibody and an affinity-matured counterpart (sometimes referred to as “progeny” or “offspring”) retain the same epitope. Suitable *in vitro* binding assays may be carried out to screen for improved binders at appropriate step(s) during the affinity maturation process. Optionally, in some embodiments, functional assays (*e.g.*, cell-based potency assays) may also be performed to confirm desired functionality.

[439] Affinity maturation typically involves sequence diversification and/or mutagenesis, whilst the exact means of introducing or generating mutations is not limiting. In some embodiments, mutagenesis comprises introducing one or more changes (*e.g.*, substitutions or deletions) in amino acid residues of one or more CDRs. Accordingly, in some embodiments the VR or CDR sequences

described herein may comprise up to one, two, three, four, five, or six amino acid changes. In some embodiments, the VR or CDR sequences described herein may comprise up to one, two, three, four, five, or six amino acid substitutions. In some embodiments, the VR or CDR sequences described herein may comprise up to one, two, three, or four deletions. Additionally or alternatively, mutagenesis may comprise so-called oligo-walking of variable regions or CDRs.

[440] For example, affinity maturation of antibodies can be accomplished by a number of methods including random mutagenesis (Gram H., et al. Proc. Natl. Acad. Sci. U.S.A. (1992) 89, 3576–3580, and Hawkins R. E., et al. J. Mol. Biol. (1992) 226, 889–896), random mutagenesis of CDR sequences, *e.g.*, CDR walking (Yang W. P., et al., J. Mol. Biol. (1995) 254, 392–403), directed mutagenesis of residues (Ho M., et al., J. Biol. Chem. (2005) 280, 607–617 and Ho M., et al., Proc. Natl. Acad. Sci. U.S.A. (2006) 103, 9637–9642), and approaches that reproduce somatic hypermutation (SHM) in vitro (Bowers P. M., et al., Proc. Natl. Acad. Sci. U.S.A. (2011) 108, 20455–20460).

[441] In one embodiment, antibodies may be affinity matured by conducting mutagenesis on the variable heavy or variable light chain regions. In another embodiment, antibodies may be affinity matured by conducting mutagenesis on any one of the variable heavy chain CDRs or variable light chain CDRs. In another embodiment, antibodies may be affinity matured by conducting mutagenesis on the variable heavy chain CDR3 (*e.g.*, CDR-H3 mutagenesis). In another embodiment, antibodies may be affinity matured by conducting mutagenesis on the variable heavy chain CDR2 (*e.g.*, CDR-H2 mutagenesis). In another embodiment, antibodies may be affinity matured by conducting mutagenesis on the variable heavy chain CDR1 (*e.g.*, CDR-H1 mutagenesis). In another embodiment, antibodies may be affinity matured by conducting mutagenesis on the variable light chain CDR3 (*e.g.*, CDR-L3 mutagenesis). In another embodiment, antibodies may be affinity matured by conducting mutagenesis on the variable light chain CDR2 (*e.g.*, CDR-L2 mutagenesis). In another embodiment, antibodies may be affinity matured by conducting mutagenesis on the variable light chain CDR1 (*e.g.*, CDR-L1 mutagenesis).

[442] In some embodiments, an antibody may be affinity matured and/or optimized by separately conducting mutagenesis on some or all of the three light chain CDRs (*i.e.*, CDR-L1, CDR-L2, and CDR-L3) to generate up to three different libraries of antibodies, each having unique mutations in one of the three CDRs. The new antibodies can then be screened for improved properties (*e.g.*, antigen-binding). Then, if further affinity improvements are desired, unique CDRs from each of the libraries can be mixed and matched to generate a new library of antibodies and screened for improved properties (*e.g.*, antigen-binding). In some embodiments, affinity maturation involves screening an antibody library comprising variants of one or more CDRs (“repertoire”), which may be combined with at least one CDR of a parent antibody. This process is sometimes called CDR shuffling or CDR diversification. In some embodiments, a variable heavy chain CDR3 (*e.g.*, CDR-H3) may be used to screen a library that contains variable heavy chain CDR1 and CDR2 repertoires of variants (*a.k.a.*,

CDR-H1/H2 diversification). In some embodiments, variable light chain CDR3 (*e.g.*, CDR-L3) may be used to screen a library that contains variable light chain CDR1 and CDR2 repertoires of variants (*a.k.a.*, CDR-L1/L2 diversification).

[443] In some embodiments, affinity maturation involves screening an antibody library comprising light chain variants (“repertoire”), which may be combined with a heavy chain of a parent antibody (light-chain shuffling). For example, in some embodiments, selected heavy chains are introduced into an antibody library comprising light chain variants thereby producing a new library of antibodies that can be screened for improved affinity. In some embodiments, repertoires of naturally occurring variable region variants may be obtained from unimmunized donors. Examples of heavy or light-chain shuffling are described in the following documents: Marks et al., (1992) *Nature Biotech* 10: 779-78; Schier et al., (1996) *J. Mol. Biol.* 255, 28-43; Park et al., (2000) *BBRC* 275. 553-557; and Chames et al., (2002) *J. Immunol* 1110-1118. In some embodiments, the light chain library comprises lambda light chains variants. In some embodiments, the light chain library comprises kappa light chains variants. In some embodiments, the light chain library comprises both lambda and kappa light chains variants.

[444] It should be appreciated that the various methods for affinity maturation may be combined in any order. For example, in one embodiment, a select antibody may undergo heavy chain CDR-H1/H2 diversification, followed by CDR-H3 mutagenesis. In another embodiment, a select antibody may undergo heavy chain CDR-H1/H2 diversification, followed by CDR-H3 mutagenesis, followed by light chain shuffling. In another embodiment, a select antibody may undergo heavy chain CDR-H1/H2 diversification, followed by light chain shuffling, followed by CDR-H3 mutagenesis. In another embodiment, a select antibody may undergo light chain shuffling, followed by heavy chain CDR-H1/H2 diversification, followed by CDR-H3 mutagenesis. In another embodiment, a select antibody may undergo light chain shuffling, followed by heavy chain CDR-H1/H2 diversification, followed by CDR-H3 mutagenesis, followed by CDR-L3 mutagenesis.

[445] In some embodiment, a select antibody may undergo light chain shuffling, followed by heavy chain CDR-H1/H2 diversification, followed by CDR-H3 mutagenesis, followed by CDR-L1, CDR-L2, and/or CDRL3 mutagenesis. In some embodiments, a select antibody may undergo heavy chain CDR-H1/H2 diversification, followed by CDR-H3 mutagenesis, followed by CDR-L1, CDR-L2, and/or CDRL3 mutagenesis. In some embodiments, light chain CDR mutagenesis is separately conducted on some or all of the three light chain CDRs (CDR-L1, CDR-L2, and CDRL3) to produce up to three libraries of antibodies. In some embodiments, the light chain CDRs from each of the libraries are mixed and matched to generate a new library of antibodies having unique combinations of light chain CDR mutations.

[446] In any of the methods for affinity maturation described above, the resulting new antibodies may be selected for binding to the target antigen (*e.g.*, a LTBP1-TGF β 1 complex and/or a LTBP3-TGF β 1 complex) using known techniques (*e.g.*, FACS). Binding specificity and affinity using FACS may be tested by varying antigen concentration and/or competition for unlabeled (cold) antigen. Binding affinity can be further assessed using other techniques known in the art, such as ELISA, BLI (*e.g.*, OCTET), and SPR (*e.g.*, BIACORE).

[447] In addition to the affinity maturation process discussed above, further optimization may be performed to achieve desired product profiles. Thus, antibodies may be further subjected to a step of optimization and selected based on certain physicochemical properties that are advantageous. For therapeutic antibodies (biologics), physicochemical criteria for developability that may be evaluated include, but are not limited to: solubility, stability, immunogenicity, lack of self-association or aggregation, Fc functionality, internalization profiles, pH-sensitivity, glycosylation and manufacturability such as cell viability and/or gene expression. In some embodiments, the process of optimization involves mutagenesis of one or more amino acid sequences within the constant regions.

[448] In one aspect, the invention provides a method for making a composition comprising an antibody, or antigen-binding fragment thereof, that specifically binds a human LTBP1-proTGF β 1 complex and/or a human LTBP3-proTGF β 1 complex, and does not bind a human GARP-proTGF β 1 complex; wherein the antibody, or antigen-binding fragment thereof, inhibits TGF β 1 but does not inhibit TGF β 2 or TGF β 3, the method comprising steps of i) providing at least one antigen comprising LTBP1-proTGF β 1 and/or LTBP3-proTGF β 1, ii) selecting a first pool of antibodies, or antigen-binding fragments thereof, that specifically bind the at least one antigen of step (i) so as to provide specific binders of LTBP1-proTGF β 1 and/or LTBP3-proTGF β 1; iii) selecting a second pool of antibodies, or antigen-binding fragments thereof, that inhibit activation of TGF β 1, so as to generate specific inhibitors of TGF β 1 activation; iv) formulating an antibody, or antigen-binding fragment thereof, that is present in the first pool of antibodies and the second pool of antibodies into a pharmaceutical composition, thereby making the composition comprising the antibody, or antigen-binding fragment thereof.

[449] In one embodiment, the method further comprises a step of removing from the first pool of antibodies, or antigen-binding fragments thereof, any antibodies, or antigen-binding fragments thereof, that bind GARP-proTGF β 1, LRRC33-proTGF β 1, mature TGF β 1, GARP-proTGF β 2, LRRC33-proTGF β 2, mature TGF β 2, GARP-proTGF β 3, LRRC33-proTGF β 3, mature TGF β 3, or any combinations thereof. In one embodiment, the method further comprises a step of determining or confirming isoform-specificity of the antibodies, or antigen-binding fragments thereof, selected in steps (ii) and/or (iii). In one embodiment, the method further comprises a step of selecting for antibodies, or antigen-binding fragments thereof, that are cross-reactive to human and rodent antigens. In one embodiment, the method further comprises a step of generating a fully human or humanized

antibody, or antigen-binding fragment thereof, of the antibody, or antigen-binding fragment thereof, that is present in the first pool of antibodies and the second pool of antibodies.

[450] In one embodiment, the method further comprises a step of subjecting the antibody, or antigen-binding fragment thereof, that is present in the first pool of antibodies and/or the second pool of antibodies to affinity maturation and/or optimization, so as to provide an affinity matured and/or optimized antibody or fragment thereof. In one embodiment, the affinity maturation and/or optimization comprises a step of subjecting the antibody, or antigen-binding fragment thereof, to light chain shuffling as described herein. In one embodiment, the affinity maturation and/or optimization comprises the step of subjecting the antibody, or antigen-binding fragment thereof, to CDR H1/H2 diversification as described herein. In one embodiment, the affinity maturation and/or optimization comprises the step of subjecting the antibody, or antigen-binding fragment thereof, to CDR-H3 mutagenesis as described herein. In one embodiment, the affinity maturation and/or optimization comprises the step of subjecting antibody, or antigen-binding fragment thereof, to light chain CDR mutagenesis as described herein. In one embodiment, the affinity maturation and/or optimization comprises the step of subjecting the antibody, or antigen-binding fragment thereof, to light chain CDR L1/L2 diversification as described herein.

[451] Further optimization steps may be carried out to provide physicochemical properties that are advantageous for therapeutic compositions. Such steps may include, but are not limited to, mutagenesis or engineering to provide improved solubility, lack of self-aggregation, stability, pH sensitivity, Fc function, and so on.

[452] In one embodiment, the method further comprises a step of determining affinity of the antibodies, or antigen-binding fragments thereof, to human LTBP1-proTGF β 1 and/or human LTBP3-proTGF β 1. In some embodiments, the method further comprises a step of removing from the first and/or second pools of antibodies, or antigen-binding fragments thereof, any antibodies, or antigen-binding fragments thereof, that bind to human LTBP1-proTGF β 1 and/or human LTBP3-proTGF β 1 with a K_D of > 100 nM, as measured in a suitable in vitro binding assay such as Bio-Layer Interferometry (BLI). In some embodiments, the method further comprises a step of removing from the first and/or second pools of antibodies, or antigen-binding fragments thereof, any antibodies, or antigen-binding fragments thereof, that bind to human LTBP1-proTGF β 1 and/or human LTBP3-proTGF β 1 with a K_D of > 50 nM, as measured in a suitable in vitro binding assay such as Bio-Layer Interferometry (BLI). In some embodiments, the method further comprises a step of removing from the first and/or second pools of antibodies, or antigen-binding fragments thereof, any antibodies, or antigen-binding fragments thereof, that bind to human LTBP1-proTGF β 1 and/or human LTBP3-proTGF β 1 with a K_D of > 25 nM, as measured in a suitable in vitro binding assay such as Bio-Layer Interferometry (BLI). In some embodiments, the method further comprises a step of removing from the first and/or second pools of antibodies, or antigen-binding fragments thereof, any antibodies, or

antigen-binding fragments thereof, that bind to human LTBP1-proTGF β 1 and/or human LTBP3-proTGF β 1 with a K_D of > 10 nM, as measured in a suitable in vitro binding assay such as Bio-Layer Interferometry (BLI).

[453] In one embodiment, the method further comprises a step of determining affinity of the antibodies, or antigen-binding fragments thereof, from the first and/or second pools of antibodies to mouse LTBP1-proTGF β 1 and/or mouse LTBP3-proTGF β 1. In some embodiments, the method further comprises a step of removing from the first and/or second pools of antibodies, or antigen-binding fragments thereof, any antibodies, or antigen-binding fragments thereof, that bind to mouse LTBP1-proTGF β 1 and/or mouse LTBP3-proTGF β 1 with a K_D of > 100 nM, as measured in a suitable in vitro binding assay such as Bio-Layer Interferometry (BLI). In some embodiments, the method further comprises a step of removing from the first and/or second pools of antibodies, or antigen-binding fragments thereof, any antibodies, or antigen-binding fragments thereof, that bind to mouse LTBP1-proTGF β 1 and/or mouse LTBP3-proTGF β 1 with a K_D of > 50 nM, as measured in a suitable in vitro binding assay such as Bio-Layer Interferometry (BLI). In some embodiments, the method further comprises a step of removing from the first and/or second pools of antibodies, or antigen-binding fragments thereof, any antibodies, or antigen-binding fragments thereof, that bind to mouse LTBP1-proTGF β 1 and/or mouse LTBP3-proTGF β 1 with a K_D of > 25 nM, as measured in a suitable in vitro binding assay such as Bio-Layer Interferometry (BLI). In some embodiments, the method further comprises a step of removing from the first and/or second pools of antibodies, or antigen-binding fragments thereof, any antibodies, or antigen-binding fragments thereof, that bind to mouse LTBP1-proTGF β 1 and/or mouse LTBP3-proTGF β 1 with a K_D of > 10 nM, as measured in a suitable in vitro binding assay such as Bio-Layer Interferometry (BLI).

[454] In one embodiment, the method further comprises a step of removing from the first and/or second pool of antibodies, or antigen-binding fragments thereof, any antibodies, or antigen-binding fragments thereof, that do not bind mouse LTBP1-proTGF β 1 and/or mouse LTBP3-proTGF β 1.

[455] In one embodiment, the method further comprises a step of determining the IC_{50} of the antibodies, or antigen-binding fragments thereof, from the first and/or second pools of antibodies, or antigen-binding fragments thereof, as measured by a suitable functional in vitro cell-based assay such as a *caga* assay, as described herein. In some embodiments, the method comprises the step of removing antibodies, or antigen-binding fragments thereof, from the first and/or second pools of antibodies, or antigen-binding fragments thereof, that have an IC_{50} of greater than 50 nM as measured by a *caga* assay as described herein. In some embodiments, the method comprises the step of removing antibodies, or antigen-binding fragments thereof, from the first and/or second pools of antibodies, or antigen-binding fragments thereof, that have an IC_{50} of greater than 25 nM as measured by a *caga* assay as described herein. In some embodiments, the method comprises the step of removing antibodies, or antigen-binding fragments thereof, from the first and/or second pools of

antibodies, or antigen-binding fragments thereof, that have an IC_{50} of greater than 10 nM as measured by a caga assay as described herein. In some embodiments, the method comprises the step of removing antibodies, or antigen-binding fragments thereof, from the first and/or second pools of antibodies, or antigen-binding fragments thereof, that have an IC_{50} of greater than 5 nM as measured by a caga assay as described herein.

[456] In some embodiments, the method comprises the step of removing antibodies, or antigen-binding fragments thereof, from the first and/or second pools of antibodies, or antigen-binding fragments thereof, that have an IC_{50} of greater than 100 nM as measured by an endogenous LTBP caga assay as described herein. In some embodiments, the method comprises the step of removing antibodies, or antigen-binding fragments thereof, from the first and/or second pools of antibodies, or antigen-binding fragments thereof, that have an IC_{50} of greater than 50 nM as measured by an endogenous LTBP caga assay as described herein. In some embodiments, the method comprises the step of removing antibodies, or antigen-binding fragments thereof, from the first and/or second pools of antibodies, or antigen-binding fragments thereof, that have an IC_{50} of greater than 25 nM as measured by an endogenous LTBP caga assay as described herein. In some embodiments, the method comprises the step of removing antibodies, or antigen-binding fragments thereof, from the first and/or second pools of antibodies, or antigen-binding fragments thereof, that have an IC_{50} of greater than 10 nM as measured by an endogenous LTBP caga assay as described herein. In some embodiments, the method comprises the step of removing antibodies, or antigen-binding fragments thereof, from the first and/or second pools of antibodies, or antigen-binding fragments thereof, that have an IC_{50} of greater than 5 nM as measured by an endogenous LTBP caga assay as described herein.

[457] In some embodiments, the method comprises the step of removing antibodies, or antigen-binding fragments thereof, from the first and/or second pools of antibodies, or antigen-binding fragments thereof, that have an IC_{50} of greater than 100 nM as measured by a human LTBP overexpression caga assay as described herein. In some embodiments, the method comprises the step of removing antibodies, or antigen-binding fragments thereof, from the first and/or second pools of antibodies, or antigen-binding fragments thereof, that have an IC_{50} of greater than 50 nM as measured by a human LTBP overexpression caga assay as described herein. In some embodiments, the method comprises the step of removing antibodies, or antigen-binding fragments thereof, from first and/or second pools of antibodies, or antigen-binding fragments thereof, that have an IC_{50} of greater than 25 nM as measured by a human LTBP overexpression caga assay as described herein. In some embodiments, the method comprises the step of removing antibodies, or antigen-binding fragments thereof, from the first and/or second pools of antibodies, or antigen-binding fragments thereof, that have an IC_{50} of greater than 10 nM as measured by a human LTBP overexpression caga assay as described herein. In some embodiments, the method comprises the step of removing antibodies, or antigen-binding fragments thereof, from the first and/or second pools of antibodies, or antigen-

binding fragments thereof, that have an IC_{50} of greater than 5 nM as measured by a human LTBP overexpression caga assay as described herein.

[458] In some embodiments, the method comprises the step of removing antibodies, or antigen-binding fragments thereof, from the first and/or second pools of antibodies, or antigen-binding fragments thereof, that have an IC_{50} of greater than 100 nM as measured by a murine LTBP overexpression caga assay as described herein. In some embodiments, the method comprises the step of removing antibodies, or antigen-binding fragments thereof, from the first and/or second pools of antibodies, or antigen-binding fragments thereof, that have an IC_{50} of greater than 50 nM as measured by a murine LTBP overexpression caga assay as described herein. In some embodiments, the method comprises the step of removing antibodies, or antigen-binding fragments thereof, from first and/or second pools of antibodies, or antigen-binding fragments thereof, that have an IC_{50} of greater than 25 nM as measured by a murine LTBP overexpression caga assay as described herein. In some embodiments, the method comprises the step of removing antibodies, or antigen-binding fragments thereof, from first and/or second pools of antibodies, or antigen-binding fragments thereof, that have an IC_{50} of greater than 10 nM as measured by a murine LTBP overexpression caga assay as described herein. In some embodiments, the method comprises the step of removing antibodies, or antigen-binding fragments thereof from first and/or second pools of antibodies, or antigen-binding fragments thereof, that have an IC_{50} of greater than 5 nM as measured by a murine LTBP overexpression caga assay as described herein.

[459] In another embodiment, a monoclonal antibody is obtained from the non-human animal, and then modified, *e.g.*, made chimeric, using suitable recombinant DNA techniques. A variety of approaches for making chimeric antibodies have been described. See *e.g.*, Morrison *et al.*, Proc. Natl. Acad. Sci. U.S.A. 81:6851, 1985; Takeda *et al.*, Nature 314:452, 1985, Cabilly *et al.*, U.S. Pat. No. 4,816,567; Boss *et al.*, U.S. Pat. No. 4,816,397; Tanaguchi *et al.*, European Patent Publication EP171496; European Patent Publication 0173494, United Kingdom Patent GB 2177096B.

[460] For additional antibody production techniques, *see, e.g.*, Antibodies: A Laboratory Manual, eds. Harlow *et al.*, Cold Spring Harbor Laboratory, 1988. The present disclosure is not necessarily limited to any particular source, method of production, or other special characteristics of an antibody.

[461] Some aspects of the present disclosure relate to host cells transformed with a polynucleotide or vector. Host cells may be a prokaryotic or eukaryotic cell. The polynucleotide or vector which is present in the host cell may either be integrated into the genome of the host cell or it may be maintained extrachromosomally. The host cell can be any prokaryotic or eukaryotic cell, such as a bacterial, insect, fungal, plant, animal or human cell. In some embodiments, fungal cells are, for example, those of the genus *Saccharomyces*, in particular those of the species *S. cerevisiae*. The term "prokaryotic" includes all bacteria which can be transformed or transfected with a DNA or RNA

molecules for the expression of an antibody or the corresponding immunoglobulin chains.

Prokaryotic hosts may include gram negative as well as gram positive bacteria such as, for example, *E. coli*, *S. typhimurium*, *Serratia marcescens* and *Bacillus subtilis*. The term "eukaryotic" includes yeast, higher plants, insects and vertebrate cells, *e.g.*, mammalian cells, such as NSO and CHO cells. Depending upon the host employed in a recombinant production procedure, the antibodies or immunoglobulin chains encoded by the polynucleotide may be glycosylated or may be non-glycosylated. Antibodies or the corresponding immunoglobulin chains may also include an initial methionine amino acid residue.

[462] In some embodiments, once a vector has been incorporated into an appropriate host, the host may be maintained under conditions suitable for high level expression of the nucleotide sequences, and, as desired, the collection and purification of the immunoglobulin light chains, heavy chains, light/heavy chain dimers or intact antibodies, antigen-binding fragments or other immunoglobulin forms may follow; see, Beychok, Cells of Immunoglobulin Synthesis, Academic Press, N.Y., (1979). Thus, polynucleotides or vectors are introduced into the cells which in turn produce the antibody or antigen-binding fragments. Furthermore, transgenic animals, preferably mammals, comprising the aforementioned host cells may be used for the large scale production of the antibody or antibody fragments.

[463] The transformed host cells can be grown in fermenters and cultured using any suitable techniques to achieve optimal cell growth. Once expressed, the whole antibodies, their dimers, individual light and heavy chains, other immunoglobulin forms, or antigen-binding fragments, can be purified according to standard procedures of the art, including ammonium sulfate precipitation, affinity columns, column chromatography, gel electrophoresis and the like; see, Scopes, "Protein Purification", Springer Verlag, N.Y. (1982). The antibody or antigen-binding fragments can then be isolated from the growth medium, cellular lysates, or cellular membrane fractions. The isolation and purification of the, *e.g.*, microbially expressed antibodies or antigen-binding fragments may be by any conventional means such as, for example, preparative chromatographic separations and immunological separations such as those involving the use of monoclonal or polyclonal antibodies directed, *e.g.*, against the constant region of the antibody.

[464] Aspects of the disclosure relate to a hybridoma, which provides an indefinitely prolonged source of monoclonal antibodies. As an alternative to obtaining immunoglobulins directly from the culture of hybridomas, immortalized hybridoma cells can be used as a source of rearranged heavy chain and light chain loci for subsequent expression and/or genetic manipulation. Rearranged antibody genes can be reverse transcribed from appropriate mRNAs to produce cDNA. In some embodiments, heavy chain constant region can be exchanged for that of a different isotype or eliminated altogether. The variable regions can be linked to encode single chain Fv regions. Multiple Fv regions can be linked to confer binding ability to more than one target or chimeric heavy and light

chain combinations can be employed. Any appropriate method may be used for cloning of antibody variable regions and generation of recombinant antibodies.

[465] In some embodiments, an appropriate nucleic acid that encodes variable regions of a heavy and/or light chain is obtained and inserted into an expression vectors which can be transfected into standard recombinant host cells. A variety of such host cells may be used. In some embodiments, mammalian host cells may be advantageous for efficient processing and production. Typical mammalian cell lines useful for this purpose include CHO cells, 293 cells, or NSO cells. The production of the antibody or antigen-binding fragment may be undertaken by culturing a modified recombinant host under culture conditions appropriate for the growth of the host cells and the expression of the coding sequences. The antibodies or antigen-binding fragments may be recovered by isolating them from the culture. The expression systems may be designed to include signal peptides so that the resulting antibodies are secreted into the medium; however, intracellular production is also possible.

[466] The disclosure also includes a polynucleotide encoding at least a variable region of an immunoglobulin chain of the antibodies described herein. In some embodiments, the variable region encoded by the polynucleotide comprises at least one complementarity determining region (CDR) of the VH and/or VL of the variable region of the antibody produced by any one of the above described hybridomas.

[467] Polynucleotides encoding antibody or antigen-binding fragments may be, *e.g.*, DNA, cDNA, RNA or synthetically produced DNA or RNA or a recombinantly produced chimeric nucleic acid molecule comprising any of those polynucleotides either alone or in combination. In some embodiments, a polynucleotide is part of a vector. Such vectors may comprise further genes such as marker genes which allow for the β the vector in a suitable host cell and under suitable conditions.

[468] In some embodiments, a polynucleotide is operatively linked to expression control sequences allowing expression in prokaryotic or eukaryotic cells. Expression of the polynucleotide comprises transcription of the polynucleotide into a translatable mRNA. Regulatory elements ensuring expression in eukaryotic cells, preferably mammalian cells, are well known to those skilled in the art. They may include regulatory sequences that facilitate initiation of transcription and optionally poly-A signals that facilitate termination of transcription and stabilization of the transcript. Additional regulatory elements may include transcriptional as well as translational enhancers, and/or naturally associated or heterologous promoter regions. Possible regulatory elements permitting expression in prokaryotic host cells include, *e.g.*, the PL, Lac, Trp or Tac promoter in *E. coli*, and examples of regulatory elements permitting expression in eukaryotic host cells are the AOX1 or GAL1 promoter in yeast or the CMV-promoter, SV40-promoter, RSV-promoter (Rous sarcoma virus), CMV-enhancer, SV40-enhancer or a globin intron in mammalian and other animal cells.

[469] Beside elements which are responsible for the initiation of transcription such regulatory elements may also include transcription termination signals, such as the SV40-poly-A site or the tk-poly-A site, downstream of the polynucleotide. Furthermore, depending on the expression system employed, leader sequences capable of directing the polypeptide to a cellular compartment or secreting it into the medium may be added to the coding sequence of the polynucleotide and have been described previously. The leader sequence(s) is (are) assembled in appropriate phase with translation, initiation and termination sequences, and preferably, a leader sequence capable of directing secretion of translated protein, or a portion thereof, into, for example, the extracellular medium. Optionally, a heterologous polynucleotide sequence can be used that encode a fusion protein including a C- or N-terminal identification peptide imparting desired characteristics, *e.g.*, stabilization or simplified purification of expressed recombinant product.

[470] In some embodiments, polynucleotides encoding at least the variable domain of the light and/or heavy chain may encode the variable domains of both immunoglobulin chains or only one. Likewise, polynucleotides may be under the control of the same promoter or may be separately controlled for expression. Furthermore, some aspects relate to vectors, particularly plasmids, cosmids, viruses and bacteriophages used conventionally in genetic engineering that comprise a polynucleotide encoding a variable domain of an immunoglobulin chain of an antibody or antigen-binding fragment; optionally in combination with a polynucleotide that encodes the variable domain of the other immunoglobulin chain of the antibody.

[471] In some embodiments, expression control sequences are provided as eukaryotic promoter systems in vectors capable of transforming or transfecting eukaryotic host cells, but control sequences for prokaryotic hosts may also be used. Expression vectors derived from viruses such as retroviruses, vaccinia virus, adeno-associated virus, herpes viruses, or bovine papilloma virus, may be used for delivery of the polynucleotides or vector into targeted cell population (*e.g.*, to engineer a cell to express an antibody or antigen-binding fragment). A variety of appropriate methods can be used to construct recombinant viral vectors. In some embodiments, polynucleotides and vectors can be reconstituted into liposomes for delivery to target cells. The vectors containing the polynucleotides (*e.g.*, the heavy and/or light variable domain(s) of the immunoglobulin chains encoding sequences and expression control sequences) can be transferred into the host cell by suitable methods, which vary depending on the type of cellular host.

Modifications

[472] Antibodies, or antigen-binding portions thereof, of the disclosure may be modified with a detectable label or detectable moiety, including, but not limited to, an enzyme, prosthetic group, fluorescent material, luminescent material, bioluminescent material, radioactive material, positron emitting metal, nonradioactive paramagnetic metal ion, and affinity label for detection and/or

isolation of a LTBP1-TGFβ1 complex or a LTBP3-TGFβ1 complex. The detectable substance or moiety may be coupled or conjugated either directly to the polypeptides of the disclosure or indirectly, through an intermediate (such as, for example, a linker (*e.g.*, a cleavable linker)) using suitable techniques. Non-limiting examples of suitable enzymes include horseradish peroxidase, alkaline phosphatase, β-galactosidase, glucose oxidase, or acetylcholinesterase; non-limiting examples of suitable prosthetic group complexes include streptavidin/biotin and avidin/biotin; non-limiting examples of suitable fluorescent materials include biotin, umbelliferone, fluorescein, fluorescein isothiocyanate, rhodamine, dichlorotriazinylamine fluorescein, dansyl chloride, or phycoerythrin; an example of a luminescent material includes luminol; non-limiting examples of bioluminescent materials include luciferase, luciferin, and aequorin; and examples of suitable radioactive material include a radioactive metal ion, *e.g.*, alpha-emitters or other radioisotopes such as, for example, iodine (¹³¹I, ¹²⁵I, ¹²³I, ¹²¹I), carbon (¹⁴C), sulfur (³⁵S), tritium (³H), indium (¹¹⁵mIn, ¹¹³mIn, ¹¹²In, ¹¹¹In), and technetium (⁹⁹Tc, ^{99m}Tc), thallium (²⁰¹Tl), gallium (⁶⁸Ga, ⁶⁷Ga), palladium (¹⁰³Pd), molybdenum (⁹⁹Mo), xenon (¹³³Xe), fluorine (¹⁸F), ¹⁵³Sm, Lu, ¹⁵⁹Gd, ¹⁴⁹Pm, ¹⁴⁰La, ¹⁷⁵Yb, ¹⁶⁶Ho, ⁹⁰Y, ⁴⁷Sc, ⁸⁶R, ¹⁸⁸Re, ¹⁴²Pr, ¹⁰⁵Rh, ⁹⁷Ru, ⁶⁸Ge, ⁵⁷Co, ⁶⁵Zn, ⁸⁵Sr, ³²P, ¹⁵³Gd, ¹⁶⁹Yb, ⁵¹Cr, ⁵⁴Mn, ⁷⁵Se, and tin (¹¹³Sn, ¹¹⁷Sn). The detectable substance may be coupled or conjugated either directly to the antibodies of the disclosure that bind selectively to a LTBP1-TGFβ1 complex and/or a LTBP3-TGFβ1 complex, or indirectly, through an intermediate (such as, for example, a linker) using suitable techniques. Any of the antibodies provided herein that are conjugated to a detectable substance may be used for any suitable diagnostic assays, such as those described herein.

[473] In addition, antibodies, or antigen-binding portions thereof, of the disclosure may also be modified with a drug to form, *e.g.*, an antibody-drug conjugate. The drug may be coupled or conjugated either directly to the polypeptides of the disclosure, or indirectly, through an intermediate (such as, for example, a linker (*e.g.*, a cleavable linker)) using suitable techniques.

Targeting Agents

[474] In some embodiments methods of the present disclosure comprise the use of one or more targeting agents to target an antibody, or antigen-binding portion thereof, as disclosed herein, to a particular site in a subject for purposes of enriching or localizing such agent(s) to a niche of interest. In some embodiments, such targeting may achieve modulating mature TGFβ release from a LTBP1-TGFβ1 complex and/or a LTBP3-TGFβ1 complex. For example, LTBP1-TGFβ1 and LTBP3-TGFβ1 complexes are typically localized to extracellular matrix. Thus, in some embodiments, antibodies disclosed herein can be conjugated to extracellular matrix targeting agents for purposes of localizing the antibodies to sites where LTBP1-TGFβ1 and LTBP3-TGFβ1 complexes reside. In such embodiments, selective targeting of antibodies leads to selective modulation of LTBP1-TGFβ1 and/or LTBP3-TGFβ1 complexes. In some embodiments, selective targeting of antibodies leads to selective inhibition of LTBP1-TGFβ1 and/or LTBP3-TGFβ1 complexes (*e.g.*, for purposes of treating fibrosis).

In some embodiments, extracellular matrix targeting agents include heparin binding agents, matrix metalloproteinase binding agents, lysyl oxidase binding domains, fibrillin-binding agents, hyaluronic acid binding agents, and others.

[475] In some embodiments, bispecific antibodies may be used having a first portion that selectively binds a LTBP1/3-TGF β 1 complex and a second portion that selectively binds a component of a target site, *e.g.*, a component of the ECM (*e.g.*, fibrillin).

[476] In some embodiments, such a target agent may be coupled to another agent or therapeutics to carry or localize the complex to a niche of interest.

Safety/toxicity considerations

Histopathology, toxicology

[477] As mentioned above, known pan-inhibitors that antagonize all TGF β isoforms, namely, TGF β 1, TGF β 2 and TGF β 3, have been documented to cause various toxicities across multiple mammalian species. Most notable known toxicities include cardiovascular toxicities (such as valvulopathy) and epithelial hyperplasia, skin lesions, inflammation and bleeding. More specifically, some of the observed toxicities associated with pan-TGF β inhibitors (*e.g.*, small molecule antagonists of the TGF β R and non-selective neutralizing antibodies) reported in the literature include the following.

[478] Cardiovascular toxicities associated with TGF β inhibition include, hyperplasia in aortic valve, right AV valve, and left AV valve; inflammation in aortic valve, left AV valve, and ascending aorta; hemorrhage in ascending aorta, aortic valve and left AV valve; connective tissue degeneration in ascending aorta (see for example, Strauber et al. (2014) “Nonclinical safety evaluation of a Transforming Growth Factor β receptor I kinase inhibitor in Fischer 344 rats and beagle dogs” J. Clin. Pract 4(3): 1000196).

[479] In addition, neutralizing antibodies that bind all three TGF β isoforms have been associated with certain epithelial toxicities, which are summarized in the table below.

Epithelial and other toxicities across species for 1D11 and GC1008

	Mice	Cyno	Human
Toxicities	<ul style="list-style-type: none"> • Hyperplasia and inflammation of tongue, gingiva, and esophagus. • Findings not reversible (12wk recovery) 	<ul style="list-style-type: none"> • Hyperplasia of gingiva, nasal epithelium, and bladder • Anemia lead to cessation of treatment • Changes were reversible (except bladder) 	<ul style="list-style-type: none"> • Gingival bleeding • Epistaxis • Headache • Fatigue • Various skin disorders, including keratoacanthomas (KA), hyperkeratosis, cutaneous SCC, and basal cell carcinoma
Drug / Dose /	1D11	GC1008	GC1008

Duration	Dosing: 50 mg/kg(3x/week) Duration: 9-12 weeks	Dosing: 10 and 50 mg/kg Duration: 6 months	Dose: 0.1, 0.3, 1, 3, 10, 15 mg/kg Duration: 4 monthly doses
Exposure	Serum conc. = 1-2 mg/mL (over 4-12 weeks)	Not disclosed	Half-life: 21.7d DN Cmax ~(350 ng/mL)mg
*Vitsky et. Al. Am. J Pathology vol. 174, 2009; and Lonning et. al. Current Pharmaceutical Biotech, 2011			

[480] Applicant of the present disclosure previously demonstrated the improved safety profiles of monoclonal antibodies that selectively block the activation step of TGF β 1 by targeting latent proTGF β 1 complex (see, for example, WO 2017/156500 and WO 2018/129329). In rat toxicology studies described therein, there were no observable test article-related toxicities when the animals were dosed with the inhibitors up to 100 mg/kg per week for 4 weeks.

[481] Building upon the earlier recognition by the applicant of the present disclosure (see PCT/US2017/021972) that lack of isoform-specificity of conventional TGF β antagonists may underlie the source of toxicities associated with TGF β inhibition, the present inventors sought to further achieve context-selective TGF β 1 inhibition for treating various diseases that manifest TGF β 1 dysregulation, particularly fibrotic conditions, with enhanced safety/tolerability. The work presented herein therefore further provided, among high-affinity inhibitors, a subset of antibodies with particularly low dissociation rates (k_{OFF}) in order to improve durability.

[482] Thus, in some embodiments, the novel antibody according to the present disclosure has the maximally tolerated dose (MTD) of >100 mg/kg when dosed weekly for at least 4 weeks (*e.g.*, 4, 6, 8, 10, 12 weeks). In some embodiments, the novel antibody according to the present disclosure has the no observed adverse effect level (NOAEL) of up to 100 mg/kg when dosed weekly for at least 4 weeks in rats. In some embodiments, the antibody has a NOAEL of at least 100 mg/kg/week when dosed for 4 weeks or 12 weeks in mice. Suitable animal models to be used for conducting safety/toxicology studies for TGF β inhibitors and TGF β 1 inhibitors include, but are not limited to: rats, dogs, cynos, and mice. In preferred embodiments, the minimum effective amount of the antibody based on a suitable preclinical efficacy study is below the NOAEL. More preferably, the minimum effective amount of the antibody is about one-third or less of the NOAEL. In particularly preferred embodiments, the minimum effective amount of the antibody is about one-sixth or less of the NOAEL. In some embodiments, the minimum effective amount of the antibody is about one-tenth or less of the NOAEL.

[483] In some embodiments, the invention encompasses an isoform-selective antibody capable of inhibiting TGF β 1 signaling, which, when administered to a subject, does not cause cardiovascular or known epithelial toxicities at a dose effective to treat a TGF β 1-related indication. In some

embodiments, the antibody has a minimum effective amount of about 3-10 mg/kg administered weekly, biweekly or monthly. Preferably, the antibody causes no to minimum toxicities at a dose that is at least six-times the minimum effective amount (*e.g.*, a six-fold therapeutic window). More preferably, the antibody causes no to minimum toxicities at a dose that is at least ten-times the minimum effective amount (*e.g.*, a ten-fold therapeutic window). Even more preferably, the antibody causes no to minimum toxicities at a dose that is at least fifteen-times the minimum effective amount (*e.g.*, a fifteen-fold therapeutic window).

[484] Thus, selection of an antibody or an antigen-binding fragment thereof for therapeutic use may include: selecting an antibody or antigen-binding fragment that meets the criteria of one or more of TGF β inhibitors (such as monoclonal antibodies and antigen-binding fragments selected for example for having slow dissociation rates, *e.g.*, k_{OFF} of $< 5 \times 10^{-4}$ (1/s)); carrying out an in vivo efficacy study in a suitable preclinical model to determine an effective amount of the antibody or the fragment; carrying out an in vivo safety/toxicology study in a suitable model to determine an amount of the antibody that is safe or toxic (*e.g.*, MTD, NOAEL, or any art-recognized parameters for evaluating safety/toxicity); and, selecting the antibody or the fragment that provides at least a three-fold therapeutic window (preferably 6-fold, more preferably a 10-fold therapeutic window, even more preferably a 15-fold therapeutic window). The selected antibody or the fragment may be used in the manufacture of a pharmaceutical composition comprising the antibody or the fragment. Such pharmaceutical composition may be used in the treatment of a TGF β 1 indication in a subject as described herein. For example, the TGF β 1 indication may be a fibrotic disorder. Preferably, a TGF β inhibitor to be selected for therapeutic use or large-scale manufacture, does not produce observable adverse effects in the treated animals after at least 4 week, *e.g.*, 8 weeks, and 12 weeks, of sustained exposure. In some embodiments, certain toxicities observed in histopathological analyses are considered non-adverse.

Immune safety assessment

[485] Cytokines play an important role in normal immune responses, but when the immune system is triggered to become hyperactive, the positive feedback loop of cytokine production can lead to a “cytokine storm” or hypercytokinemia, a situation in which excessive cytokine production causes an immune response that can damage organs, especially the lungs and kidneys, and even lead to death. Such condition is characterized by markedly elevated proinflammatory cytokines in the serum. Historically, a Phase 1 Trial of the anti-CD28 monoclonal antibody TGN1412 in healthy volunteers led to a life-threatening “cytokine storm” response resulted from an unexpected systemic and rapid induction of proinflammatory cytokines (Suntharalingam G et al. N Engl J Med. 2006 Sep 7;355(10):1018-28). This incident prompted heightened awareness of the potential danger associated with pharmacologic stimulation of T cells.

[486] Whilst TGF β -directed therapies do not target a specific T cell receptor or its ligand, it is contemplated that it is prudent to carry out immune safety assessment, including, for example, *in vitro* cytokine release assays, *in vivo* cytokine measurements from plasma samples of non-human primate treated with a TGF β inhibitor, and platelet assays using human platelets.

[487] In some embodiments, selection of a TGF β inhibitor for therapeutic use and/or large-scale production thereof includes an assessment of the ability for the TGF β inhibitor to trigger cytokine release from cytokine-producing cells. In such an assessment, one or more of the cytokines (*e.g.*, inflammatory cytokines) IL-2, TNF α , IFN γ , IL-1 β , CCL2 (MCP-1), and IL-6 may be assayed. In some embodiments, the cytokine-producing cells may include peripheral blood mononuclear cell (PBMC) constituents from healthy donors. Cytokine response after exposure to the TGF β inhibitor (such as an antibody disclosed) herein may be compared to release after exposure to a control, *e.g.*, an IgG isotype negative control, or any other suitable control depending on the TGF β inhibitor being tested. Cytokine activation may be assessed in plate-bound (*e.g.*, immobilized) and/or soluble assay formats. Levels of IFN γ , IL-2, IL-1 β , TNF α , IL-6, and CCL2 (MCP-1) should not exceed 10-fold, *e.g.*, 8-, 6-, 4-, or 2-fold the activation in the negative control. In some embodiments, a positive control may also be used to confirm cytokine activation in the sample, *e.g.*, in the PBMCs. In some embodiments, these *in vitro* cytokine release results may be further confirmed *in vivo*, *e.g.*, in an animal model such as a monkey toxicology study, *e.g.*, a 4-week GLP or non-GLP repeat-dose monkey study.

[488] In some embodiments, selection of an antibody or an antigen-binding fragment thereof for therapeutic use may include: identifying an antibody or antigen-binding fragment that meets the criteria of one or more of those described herein; carrying out an *in vivo* efficacy study in a suitable preclinical model to determine an effective amount of the antibody or the fragment; carrying out an *in vivo* safety/toxicology study in a suitable model to determine an amount of the antibody that is safe or toxic (*e.g.*, MTD, NOAEL, or any art-recognized parameters for evaluating safety/toxicity); and, selecting the antibody or the fragment that provides at least a three-fold therapeutic window (preferably 6-fold, more preferably a 10-fold therapeutic window, even more preferably a 15-fold therapeutic window). In certain embodiments, the *in vivo* efficacy study is carried out in two or more suitable preclinical models that recapitulate human conditions. In some embodiments, such preclinical models comprise TGF β 1-positive fibrosis. In some embodiments, the preclinical models are selected from liver fibrosis model, kidney fibrosis model, lung fibrosis model, heart (cardiac) fibrosis model, skin fibrosis model.

[489] Identification of an antibody or antigen-binding fragment thereof for therapeutic use may further include carrying out an immune safety assay, which may include, but is not limited to, measuring cytokine release and/or determining the impact of the antibody or antigen-binding fragment on platelet binding, activation, and/or aggregation. In certain embodiments, cytokine release

may be measured in vitro using PBMCs or in vivo using a preclinical model such as non-human primates. In certain embodiments, the antibody or antigen-binding fragment thereof does not induce a greater than 10-fold release in IL-6, IFN γ , and/or TNF α levels as compared to levels in an IgG control sample in the immune safety assessment. In certain embodiments, assessment of platelet binding, activation, and aggregation may be carried out in vitro using PBMCs. In some embodiments, the antibody or antigen-binding fragment thereof does not induce a more than 10% increase in platelet binding, activation, and/or aggregation as compared to buffer or isotype control in the immune safety assessment.

[490] The selected antibody or the fragment may be used in the manufacture of a pharmaceutical composition comprising the antibody or the fragment. Such pharmaceutical composition may be used in the treatment of a TGF β indication in a subject as described herein. For example, the TGF β indication may be a fibrotic disorder, such as organ fibrosis, *e.g.*, liver fibrosis. Thus, the invention includes a method for manufacturing a pharmaceutical composition comprising a TGF β inhibitor, wherein the method includes the step of selecting a TGF β inhibitor which is tested for immune safety as assessed by immune safety assessment comprising a cytokine release assay and optionally further comprising a platelet assay. The TGF β inhibitor selected by the method does not trigger unacceptable levels of cytokine release, as compared to control (such as IgG control). Similarly, the TGF β inhibitor selected by the method does not cause unacceptable levels of platelet aggregation, platelet activation and/or platelet binding. Such TGF β inhibitor is then manufactured at large-scale, for example 250L or greater, *e.g.*, 1000L, 2000L, 3000L, 4000L or greater, for commercial production of the pharmaceutical composition comprising the TGF β inhibitor.

Pharmaceutical Compositions

[491] The invention further provides pharmaceutical compositions used as a medicament suitable for administration in human and non-human subjects. One or more antibodies that selectively binds an LTBP1-TGF β 1 complex and/or an LTBP3-TGF β 1 complex can be formulated or admixed with a pharmaceutically acceptable carrier (excipient), including, for example, a buffer, to form a pharmaceutical composition. Such formulations may be used for the treatment of a disease or disorder that involves TGF β signaling or dysregulation thereof. In some embodiments, such disease or disorder associated with TGF β signaling involves one or more contexts, *i.e.*, the TGF β is associated with a particular type or types of presenting molecules. In some embodiments, such context occurs in a cell type-specific and/or tissue-specific manner. In some embodiments, for example, such context-dependent action of TGF β signaling is mediated in part via GARP, LRRC33, LTBP1 and/or LTBP3.

[492] In some embodiments, the antibody of the present invention binds selectively to a single context of TGF β , such that the antibody binds TGF β in a complex with LTBP presenting molecules, *e.g.*, LTBP1 and/or LTBP3. Thus, such pharmaceutical compositions may be administered to patients

for alleviating a TGF β -related indication (*e.g.*, fibrosis) associated with TGF β 1 activation/release from LTBP1 and/or LTBP3.

[493] A pharmaceutically “acceptable” carrier (excipient) means that the carrier is compatible with the active ingredient of the composition (and preferably, capable of stabilizing the active ingredient) and not deleterious to the subject to be treated. Examples of pharmaceutically acceptable excipients (carriers), including buffers, would be apparent to the skilled artisan and have been described previously. See, *e.g.*, Remington: The Science and Practice of Pharmacy 20th Ed. (2000) Lippincott Williams and Wilkins, Ed. K. E. Hoover. In one example, a pharmaceutical composition described herein contains more than one antibody that selectively binds a LTBP1-TGF β 1 complex and/or a LTBP3-TGF β 1 complex, where the antibodies recognize different epitopes/residues of the LTBP1-TGF β 1 complex and/or LTBP3-TGF β 1 complex.

[494] The pharmaceutical compositions to be used in the present methods can comprise pharmaceutically acceptable carriers, excipients, or stabilizers in the form of lyophilized formulations or aqueous solutions (Remington: The Science and Practice of Pharmacy 20th Ed. (2000) Lippincott Williams and Wilkins, Ed. K. E. Hoover). Acceptable carriers, excipients, or stabilizers are nontoxic to recipients at the dosages and concentrations used, and may comprise buffers such as phosphate, citrate, and other organic acids; antioxidants including ascorbic acid and methionine; preservatives (such as octadecyldimethylbenzyl ammonium chloride; hexamethonium chloride; benzalkonium chloride, benzethonium chloride; phenol, butyl or benzyl alcohol; alkyl parabens such as methyl or propyl paraben; catechol; resorcinol; cyclohexanol; 3-pentanol; and m-cresol); low molecular weight (less than about 10 residues) polypeptides; proteins, such as serum albumin, gelatin, or immunoglobulins; hydrophilic polymers such as polyvinylpyrrolidone; amino acids such as glycine, glutamine, asparagine, histidine, arginine, or lysine; monosaccharides, disaccharides, and other carbohydrates including glucose, mannose, or dextrans; chelating agents such as EDTA; sugars such as sucrose, mannitol, trehalose or sorbitol; salt-forming counter-ions such as sodium; metal complexes (*e.g.*, Zn-protein complexes); and/or non-ionic surfactants such as TWEENTM, PLURONICSTM or polyethylene glycol (PEG). Pharmaceutically acceptable excipients are further described herein.

[495] The invention also includes pharmaceutical compositions that comprise an antibody or fragment thereof according to the present invention, and a pharmaceutically acceptable excipient.

[496] Thus, the antibody or a molecule comprising an antigen-binding fragment of such antibody can be formulated into a pharmaceutical composition suitable for human administration.

[497] The pharmaceutical formulation may include one or more excipients. In some embodiments, excipient(s) may be selected from the list provided in the following:

<https://www.accessdata.fda.gov/scripts/cder/iig/index.Cfm?event=browseByLetter.page&Letter=A>

[498] The pharmaceutical composition is typically formulated to a final concentration of the active biologic (*e.g.*, monoclonal antibody, engineered binding molecule comprising an antigen-binding fragment, etc.) to be between about 2 mg/mL and about 200 mg/mL. For example, the final concentration (wt/vol) of the formulations may range between about 2-200, 2-180, 2-160, 2-150, 2-120, 2-100, 2-80, 2-70, 2-60, 2-50, 2-40, 5-200, 5-180, 5-160, 5-150, 5-120, 5-100, 5-80, 5-70, 5-60, 5-50, 5-40, 10-200, 10-180, 10-160, 10-150, 10-120, 10-100, 10-80, 10-70, 10-60, 10-50, 10-40, 20-200, 20-180, 20-160, 20-150, 20-120, 20-100, 20-80, 20-70, 20-60, 20-50, 20-40, 30-200, 30-180, 30-160, 30-150, 30-120, 30-100, 30-80, 30-70, 30-60, 30-50, 30-40, 40-200, 40-180, 40-160, 40-150, 40-120, 40-100, 40-80, 40-70, 40-60, 40-50, 50-200, 50-180, 50-160, 50-150, 50-120, 50-100, 50-80, 50-70, 50-60, 60-200, 60-180, 60-160, 60-150, 60-120, 60-100, 60-80, 60-70, 70-200, 70-180, 70-160, 70-150, 70-120, 70-100, 70-80 mg/mL. In some embodiments, the final concentration of the biologic in the formulation is about 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 100, 110, 120, 130, 140, 150, 160, 170, 180, 190, or 200 mg/mL.

[499] According to some embodiments, the TGF β inhibitor is administered in an amount of about 3000 mg, 2400 mg, 1600 mg, 800 mg, 240 mg, 80 mg, or less.

[500] The pharmaceutical compositions of the present invention are preferably formulated with suitable buffers. Suitable buffers include but are not limited to: phosphate buffer, citric buffer, and histidine buffer.

[501] The final pH of the formulation is typically between pH 5.0 and 8.0. For example, the pH of the pharmaceutical composition may be about 5.0, 5.2, 5.5, 6.0, 6.2, 6.5, 6.8, 7.0, 7.2, 7.4, 7.5, 7.6, or 7.8.

[502] The pharmaceutical composition of the present disclosure may comprise a surfactant, such as nonionic detergent, approved for the use in pharmaceutical formulations. Such surfactants include, for example, polysorbates, such as Polysorbate 20 (Tween-20), Polysorbate 80 (Tween-80) and NP-40.

[503] The pharmaceutical composition of the present disclosure may comprise a stabilizer. For liquid-protein preparations, stability can be enhanced by selection of pH-buffering salts, and often amino acids can also be used. It is often interactions at the liquid/air interface or liquid/solid interface (with the packaging) that lead to aggregation following adsorption and unfolding of the protein. Suitable stabilizers include but are not limited to: sucrose, maltose, sorbitol, as well as certain amino acids such as histidine, glycine, methionine and arginine.

[504] The pharmaceutical composition of the present disclosure may contain one or any combinations of the following excipients: Sodium Phosphate, Arginine, Sucrose, Sodium Chloride, Tromethamine, Mannitol, Benzyl Alcohol, Histidine, Sucrose, Polysorbate 80, Sodium Citrate, Glycine, Polysorbate 20, Trehalose, Poloxamer 188, Methionine, Trehalose, rhHyaluronidase, Sodium

Succinate, Potassium Phosphate, Disodium Edetate, Sodium Chloride, Potassium Chloride, Maltose, Histidine Acetate, Sorbitol, Pentetic Acid, Human Serum Albumin, Pentetic Acid.

[505] In some embodiments, the pharmaceutical composition of the present disclosure may contain a preservative.

[506] The pharmaceutical composition of the present disclosure is typically presented as a liquid or a lyophilized form. Typically, the products can be presented in vial (*e.g.*, glass vial). Products available in syringes, pens, or autoinjectors may be presented as pre-filled liquids in these container/closure systems.

[507] In some examples, the pharmaceutical composition described herein comprises liposomes containing an antibody that selectively binds a LTBP1-TGF β 1 complex and/or a LTBP3-TGF β 1 complex, which can be prepared by any suitable method, such as described in Epstein *et al.*, *Proc. Natl. Acad. Sci. USA* 82:3688 (1985); Hwang *et al.* *Proc. Natl. Acad. Sci. USA* 77:4030 (1980); and U.S. Pat. Nos. 4,485,045 and 4,544,545. Liposomes with enhanced circulation time are disclosed in U.S. Pat. No. 5,013,556. Particularly useful liposomes can be generated by the reverse phase evaporation method with a lipid composition comprising phosphatidylcholine, cholesterol and PEG-derivatized phosphatidylethanolamine (PEG-PE). Liposomes are extruded through filters of defined pore size to yield liposomes with the desired diameter.

[508] In some embodiments, liposomes with targeting properties are selected to preferentially deliver or localize the pharmaceutical composition to certain tissues or cell types. For example, certain nanoparticle-based carriers with bone marrow-targeting properties may be employed, *e.g.*, lipid-based nanoparticles or liposomes. See, for example, Sou (2012) “Advanced drug carriers targeting bone marrow”, ResearchGate publication 232725109.

[509] The antibodies that selectively bind a LTBP1-TGF β 1 complex and/or a LTBP3-TGF β 1 complex may also be entrapped in microcapsules prepared, for example, by coacervation techniques or by interfacial polymerization, for example, hydroxymethylcellulose or gelatin-microcapsules and poly-(methylmethacrylate) microcapsules, respectively, in colloidal drug delivery systems (for example, liposomes, albumin microspheres, microemulsions, nano-particles and nanocapsules) or in macroemulsions. Exemplary techniques have been described previously, see, *e.g.*, Remington, *The Science and Practice of Pharmacy* 20th Ed. Mack Publishing (2000).

[510] In other examples, the pharmaceutical composition described herein can be formulated in sustained-release format. Suitable examples of sustained-release preparations include semipermeable matrices of solid hydrophobic polymers containing the antibody, which matrices are in the form of shaped articles, *e.g.*, films, or microcapsules. Examples of sustained-release matrices include polyesters, hydrogels (for example, poly(2-hydroxyethyl-methacrylate), or poly(vinylalcohol)), polylactides (U.S. Pat. No. 3,773,919), copolymers of L-glutamic acid and 7 ethyl-L-glutamate, non-

degradable ethylene-vinyl acetate, degradable lactic acid-glycolic acid copolymers such as the LUPRON DEPOT™ (injectable microspheres composed of lactic acid-glycolic acid copolymer and leuprolide acetate), sucrose acetate isobutyrate, and poly-D-(-)-3-hydroxybutyric acid.

[511] The pharmaceutical compositions to be used for *in vivo* administration must be sterile. This is readily accomplished by, for example, filtration through sterile filtration membranes. Therapeutic antibody compositions are generally placed into a container having a sterile access port, for example, an intravenous solution bag or vial having a stopper pierceable by a hypodermic injection needle.

[512] The pharmaceutical compositions described herein can be in unit dosage forms such as tablets, pills, capsules, powders, granules, solutions or suspensions, or suppositories, for oral, parenteral or rectal administration, or administration by inhalation or insufflation.

[513] Suitable surface-active agents include, in particular, non-ionic agents, such as polyoxyethylenesorbitans (*e.g.*, TWEEEN™ 20, 40, 60, 80 or 85) and other sorbitans (*e.g.*, SPAN™ 20, 40, 60, 80 or 85). Compositions with a surface-active agent will conveniently comprise between 0.05 and 5% surface-active agent, and can be between 0.1 and 2.5%. It will be appreciated that other ingredients may be added, for example mannitol or other pharmaceutically acceptable vehicles, if necessary.

[514] Suitable emulsions may be prepared using commercially available fat emulsions, such as INTRALIPID™, LIPSYN™, INFONUTROL™, LIPOFUNDIN™ and LIPIPHYSAN™. The active ingredient may be either dissolved in a pre-mixed emulsion composition or alternatively it may be dissolved in an oil (*e.g.*, soybean oil, safflower oil, cottonseed oil, sesame oil, corn oil or almond oil) and an emulsion formed upon mixing with a phospholipid (*e.g.*, egg phospholipids, soybean phospholipids or soybean lecithin) and water. It will be appreciated that other ingredients may be added, for example glycerol or glucose, to adjust the tonicity of the emulsion. Suitable emulsions will typically contain up to 20% oil, for example, between 5 and 20%.

[515] The emulsion compositions can be those prepared by mixing an antibody that selectively binds a LTBP1-TGFβ1 complex and/or a LTBP3-TGFβ1 complex with Intralipid™ or the components thereof (soybean oil, egg phospholipids, glycerol and water).

Use of Inhibitors that Selectively Bind a LTBP1/3-TGFβ1 Complex

[516] The inhibitors, *e.g.*, antibodies and antigen-binding portions thereof, described herein that selectively bind a LTBP1/3-TGFβ1 complex can be used in a wide variety of applications in which modulation of TGFβ1 activity associated with LTBP1 or LTBP3 is desired.

[517] In one embodiment, the invention provides a method of inhibiting TGFβ1 activation by exposing a LTBP1-TGFβ1 complex and/or a LTBP3-TGFβ1 complex to an inhibitor, *e.g.*, antibody, or antigen-binding portion thereof, which selectively binds a LTBP1/3-TGFβ1 complex. The

foregoing method can be performed *in vitro*, *e.g.*, to inhibit TGF β 1 activation in cultured cells. The foregoing method can also be performed *in vivo*, *e.g.*, in a subject in need of TGF β 1 inhibition, or in an animal model in which the effect of TGF β 1 inhibition is to be assessed.

[518] Any inhibitor, *e.g.*, antibody, or antigen-binding portion thereof, described herein which selectively binds a LTBP1-TGF β 1 complex and/or a LTBP3-TGF β 1 complex, and any pharmaceutical composition comprising such antibody, is suitable for use in the methods of the invention. For example, in one embodiment, the inhibitor, *e.g.*, antibody, or antigen-binding portion thereof, selectively binds to a LTBP1-TGF β 1 complex and a LTBP3-TGF β 1 complex, but does not bind to one or more targets selected from LTBP1 alone, mature TGF β 1 alone, a GARP-TGF β 1 complex, a LRRC33-TGF β 1 complex, and combinations thereof. Exemplary inhibitor, *e.g.*, antibodies, can inhibit the release of mature TGF β 1 from a LTBP1-proTGF β 1 complex and/or a LTBP3-proTGF β 1 complex, without inhibiting the release of mature TGF β 1 from a GARP-proTGF β 1 complex and/or a LRRC33-proTGF β 1 complex.

[519] The antibody, or antigen-binding portion thereof, can, in some embodiments, bind a LTBP1-proTGF β 1 complex and/or a LTBP3-proTGF β 1 complex with a dissociation constant (K_D) of about 10^{-8} M or less. In some embodiments, the antibody, or antigen-binding portion thereof has a K_D value of about 10^{-9} M or less. In some embodiments, the antibody, or antigen-binding portion thereof has a K_D value of about 10^{-10} M or less (*e.g.*, about 10^{-11} M or less). In some embodiments, the antibody, or antigen-binding portion thereof has a K_D value of < 10 nM, < 5 nM < 1 nM) towards a LTBP1-proTGF β 1 complex and/or a LTBP3-proTGF β 1 complex as measured in a suitable *in vitro* binding assay such as BLI (*e.g.*, Octet). In one embodiment, the antibody, or antigen-binding portion thereof, comprises at least one (*e.g.*, one, two, or three) heavy chain CDRs shown in Table 5, and/or at least one (*e.g.*, one, two, three) light chain CDRs shown in Table 5. In an exemplary embodiment, the antibody, or antigen-binding portion thereof, comprises a heavy chain variable region comprising SEQ ID NO:7, and/or a light chain variable region comprising SEQ ID NO:8. Antibodies and antigen-binding portions thereof which bind the same epitope as the foregoing antibodies, and/or which compete for binding with the foregoing antibodies to LTBP1/3-proTGF β 1, are also useful in the methods described herein. Additional features of the antibodies, or antigen-binding portions thereof, that are suitable for practicing the methods of the invention are described herein.

[520] In one embodiment, contacting a LTBP1-TGF β 1 complex and/or a LTBP3-TGF β 1 complex with the inhibitor, *e.g.*, antibody, or antigen-binding portion thereof, inhibits the release of mature TGF β 1 from the LTBP1-TGF β 1 complex and/or the LTBP3-TGF β 1 complex. In one embodiment, said contacting does not inhibit the release of mature TGF β 1 from presenting molecules other than LTBP1 and LTBP3. For example, exposing a GARP-TGF β 1 complex or a LRRC33-TGF β 1 complex to a context-specific inhibitor, *e.g.*, antibody, that selectively binds LTBP1/3-TGF β 1 but does not

bind TGFβ1 in the context of GARP or LRRC33 will not inhibit the release of mature TGFβ1 from the GARP-TGFβ1 complex or the LRRC33-TGFβ1 complex.

[521] LTBP1 and LTBP3 are generally deposited in the extracellular matrix. Accordingly, in one embodiment, complexes comprising LTBP1-TGFβ1 and/or LTBP3-TGFβ1 are associated with the extracellular matrix, *e.g.*, bound to the extracellular matrix. In some embodiments, the LTBP1/3-TGFβ1 complexes are bound to extracellular matrix comprising fibrillin, and/or a protein containing an RGD motif.

[522] The invention also provides a method of reducing TGFβ1 activation in a subject, by administering to the subject an inhibitor, *e.g.*, antibody, or antigen-binding portion thereof, which selectively binds a LTBP1/3-TGFβ1 complex, as described herein. Any antibody, or antigen-binding portion thereof, described herein which selectively binds a LTBP1-TGFβ1 complex and/or a LTBP3-TGFβ1 complex, and any pharmaceutical composition comprising such antibody, is suitable for use in the methods of the invention.

[523] Exemplary LTBP1/3 inhibitors, *e.g.*, antibodies, bind a LTBP1/3-TGFβ1 complex, and inhibit TGFβ1 activation in a context-specific manner, by inhibiting release of TGFβ1 presented by LTBP1 and LTBP3, without inhibiting release of TGFβ1 presented by GARP and/or LRRC33. Such antibodies are useful for blocking a particular subset of TGFβ1 activity *in vivo*. In one embodiment, the context-specific antibodies provided herein can be used to inhibit TGFβ1 localized to the extracellular matrix. In another embodiment, the context-specific antibodies can inhibit TGFβ1 without modulating TGFβ1-associated immune activity or immune response, which is primarily mediated by TGFβ1 presented by GARP and LRRC33. In another embodiment, the context-specific antibodies can be used to inhibit TGFβ1 activity associated with the extracellular matrix (*e.g.*, LTBP1-associated TGFβ1 activity and LTBP3-associated TGFβ1 activity) without modulating TGFβ1 activity associated with hematopoietic cells, *e.g.*, hematopoietic cells that express GARP and/or LRRC33.

Clinical Applications

[524] Applicant previously described so-called “context-independent” inhibitors of TGFβ1 (see, for example: PCT/US2017/021972 and PCT/US2018/012601) which may be useful for treating various diseases and disorders involving TGFβ1 dysregulation, including, but are not limited to, cancer and fibrosis. Unlike traditional TGFβ1 antagonists, these context-independent TGFβ1 inhibitors are capable of selectively targeting the TGFβ1 isoform. Within the multifaceted biological functions driven by the TGFβ1 isoform, however, the context-independent inhibitors do not discriminate tissue-specific (thus context-specific) proTGFβ1 complexes, such that such inhibitors are capable of binding and thereby inhibiting release or activation of mature growth factor from any of the presenting molecule-proTGFβ1 complexes.

[525] Based at least in part on the recognition that it may be advantageous to provide even greater selectivity in targeting only a subset of TGF β activities, context-selective inhibitors of the present disclosure have been generated. It is contemplated that by further narrowing particular biological contexts in which to inhibit TGF β function, greater safety may be achieved in a subset of disease conditions or patient populations. Specifically, the inventors of the present invention have recognized that in certain conditions, systemic perturbation of immune regulation may be particularly undesirable. Because TGF β plays an important role in mediating immune response and maintaining immune homeostasis, broad inhibition of TGF β activities effectuated in a context-independent manner may lead to unwanted side effects without justifiable benefits. In these circumstances, it is envisaged that it is advantageous to specifically target and inhibit matrix-associated TGF β function using a context-selective inhibitor, such as those encompassed herein, which does not inhibit the immune components of TGF β 1 function.

[526] Accordingly, the context-specific antibodies can be used to inhibit LTBP1/3-associated TGF β activity in applications in which TGF β activation in the context of LTBP1 or LTBP3 is desirable, and in which TGF β activation in the context of GARP and/or or LRRC33 is detrimental.

[527] The disease may involve dysregulation or impairment of ECM components or function and comprises increased collagen deposition. In some embodiments, the dysregulation or impairment of ECM components or function may further comprise increased stiffness and/or ECM reorganization. In some embodiments, the dysregulation or impairment of ECM components or function includes increased myofibroblast cells within the disease site. In some embodiments, the dysregulation of the ECM includes increased stiffness of the matrix, which is implicated in the pathogenesis and/or disease progression of a variety of fibrotic conditions and tumors. In some embodiments, the dysregulation of the ECM involves fibronectin and/or fibrillin.

Rationale for the development of matrix-targeted TGF β inhibitors that do not inhibit GARP-associated TGF β

[528] The invention includes context-specific inhibitors of LTBP1-associated and/or LTBP3-associated TGF β . Such inhibitors therefore are capable of specifically targeting the ECM-associated latent TGF β complexes (*e.g.*, LTBP1-proTGF β 1 and/or LTBP3-proTGF β 1) thereby inhibiting the release of mature TGF β growth factor from the latent complex at disease environments, *e.g.*, fibrotic tissues. Such inhibitors show no significant binding activities towards a GARP-proTGF β 1 complex, thereby minimizing unwanted systemic immune modulations. Such antibodies may be advantageous for use in the treatment of conditions with ECM dysregulation, such as abnormal remodeling and/or stiffness of the ECM.

[529] The context-selective antibodies provided herein may be used in the treatment of a condition where it is undesirable to stimulate the subject's immune response and/or in situations where the

subject is expected to benefit from a long-term TGF β inhibition therapy to manage a chronic condition, such as many types of fibrosis.

[530] At least three bases for supporting potential benefits of a TGF β inhibitor that does not target the GARP-proTGF β 1 complex expressed on regulatory T cells are discussed below.

[531] First, GARP-expressing T regulatory cells are a component of the immune system that suppress or dampen immune responses of other cells. This notion may be referred to as “tolerance.” This is an important “self-check” built into the immune system to prevent excessive reactions that in some situations can result in life-threatening conditions, such as sepsis, cytokine release syndrome and cytokine storm. TGF β inhibition therapies that exert Treg-inhibitory effects may, therefore, pose certain risk when the normal Treg function is impaired, particularly for a prolonged duration of time, *e.g.*, therapeutic regimen involving treatment of six months or longer, and chronic treatment that is administered for an indefinite period of time. For this reason, patients in need of TGF β inhibition therapies, particularly to avoid the risk of eliciting autoimmunity, may benefit from TGF β 1 inhibitors that do not directly perturb the normal Treg function. For example, patient populations in need of a long-term TGF β inhibition therapy may include those with genetic or congenital conditions, such as DMD, CF and others. In addition, patient populations that suffer from conditions that include inflammation may benefit from a context-specific inhibitor that does not perturb the GARP/Treg function so as to minimize the risk of exacerbating the existing inflammatory conditions.

[532] Second, increasing evidence points to a link between disproportionate Th17/Treg ratios and pathologies involving inflammation and/or fibrosis. It is generally accepted that the differentiation of the two cell types, Th17 and Treg, is negatively regulated with an inverse relationship. TGF β 1 appears to be a master gatekeeper of this process, such that, TGF β 1 exposure promotes naïve T cells to differentiate into Foxp3⁺ Tregs, whereas TGF β 1 in combination with IL-6, promotes naïve T cells to differentiate into ROR γ t⁺ Th17 cells instead. In addition, once differentiated, these cell populations negatively regulate each other.

[533] Lines of evidence suggest that an imbalance in Th17/Treg ratios correlates with the pathogenesis and/or progression of fibrotic conditions involving chronic inflammation, or severity thereof.

[534] For example, Shoukry *et al.* reported that Th17 cytokines drive liver fibrosis by regulating TGF β signaling. The authors examined *ex vivo* the frequency of Th17 and Treg populations in liver biopsy samples and found that increased Th17/Treg ratio correlated with advanced fibrosis, as compared to moderate fibrosis or healthy tissue samples. Consistent with the observation, a strong bias towards Th17 cytokines, IL-22 in particular, was also detected in fibrotic livers. These data suggest that increased Th17/Treg ratios lead to an imbalance in pro-fibrotic Th17 cytokines, which correlate with severity of liver fibrosis.

[535] Similar inverse correlations of Th17 and Treg populations are observed in other diseases.

[536] For example, increased muscle expression of IL-17 has been reported in patients with Duchenne muscular dystrophy (DMD), which is a condition that manifests chronic inflammation. De Pasquale *et al.* (Neurology 78(17): 1309-14) found that DMD muscle biopsy samples contained higher levels of IL-17 (a Th17 marker) and lower levels of Foxp3 (a Treg marker) mRNA compared to control. Elevations in other proinflammatory cytokines, such as TNF- α and MCP-1, were also observed and were found to be associated with worse clinical outcome of patients. The authors concluded that the data point to a possible pathogenic role of IL-17.

[537] Similarly, Jamshidian *et al.* (J Neuroimmunol 2013, 262(1-2): 106-12) reported biased Treg/Th17 balance away from regulatory toward inflammatory phenotype in patients with relapsed multiple sclerosis and its correlation with severity of clinical symptoms.

[538] A role of regulatory T cells is also implicated in the pathogenesis of cystic fibrosis (CF). In particular, CF lungs affected by the disease are associated with exaggerated Th17 and Th2 cell responses, indicative of a classic inflammatory phenotype, but also with a deficiency in numbers or function (*i.e.*, impairment) of Treg cells (McGuire (2015) Am J Respir Crit Care Med 191(8): 866-8).

[539] Furthermore, Zhuang *et al.* (Scientific Reports (2017) 7: 40141) found imbalance of Th17/Treg cells in patients with acute anterior uveitis (anterior segment intraocular inflammation with the positive of human class I major histocompatibility complex), in which both a marked increase in Th17 cells and a marked decrease in Treg cells were seen.

[540] Taken together, the inventors of the present disclosure recognized that what appears to be a common feature in these various diseases associated with elevated Th17/Treg ratios is that the patient suffers from a fibrotic condition accompanied by an inflammatory component.

[541] Thus, it is envisaged in the present disclosure that TGF β inhibition therapy that spares the Treg/GARP-arm of the TGF β function may be particularly advantageous for an effective treatment of diseases characterized by an elevated level of Th17/Treg ratios. In this way, the context-selective inhibitors of TGF β according to the invention are aimed to avoid more systemic effects of TGF β inhibition that may interfere with Treg function, which may lead to exacerbation of existing fibrotic/inflammatory conditions in patients. Thus, the matrix-targeted TGF β inhibitors described herein are used in a method for treating a patient who has or at risk of developing a fibrotic disorder that comprises inflammation. In some embodiments, the patient has an elevated Th17-to-Treg cell ratio. In some embodiments, the elevated Th17/Treg ratio may be predominantly caused by an increased number of Th17 cells, while in other embodiments, the elevated Th17/Treg ratio may be predominantly caused by a decreased number of Treg cells in the patient (or a biological sample collected from the patient). Yet in further embodiments, the elevated Th17/Treg ratio may be caused by a combination of an increased number of Th17 cells and a decreased number of Treg cells. In

some embodiments, elevated levels of IL-17 and/or IL-22 detected in patients (or measured in samples collected from the patients) are also indicative of fibrotic conditions accompanied by chronic inflammation. Such patients may be therefore selected as candidates for receiving a context-selective TGF β 1 inhibitor therapy disclosed herein.

[542] The third line of reasoning for keeping the GARP-TGF β 1 axis intact in a TGF β inhibition therapy relates to the benefit of maintaining normal Treg function. As mentioned, GARP is expressed on the cell surface of Tregs and are thought to play a role in TGF β -mediated immunomodulation. Because Tregs are indispensable for immune homeostasis and the prevention of autoimmunity, unnecessary perturbation of which may put certain patient populations at higher risk of, for example, infections (reviewed, for example, by: Richert-Spuhler and Lund (2015) *Prog Mol Biol Transl Sci*. 136:217-243).

[543] The third line of reasoning for keeping the GARP-TGF β 1 axis intact in a TGF β inhibition therapy is that regulatory T cells function as a "break" to modulate or dampen over-reactive immune response. The discovery of Foxp3 as the master regulator of Treg cell development and function was critical for the understanding of Treg cell biology. Inactivating mutations in Foxp3 result in the spontaneous development of severe autoimmunity with a scurfy phenotype in mice and IPEX syndrome ('immune dysregulation, polyendocrinopathy, enteropathy, X-linked') in humans (see Dominguez-Villaur and Haler, *Nature Immunology* 19, 665-673, 2018). Thus, it raises the possibility that TGF β 1 therapy that elicits inhibitory effects of the Treg/GARP arm of TGF β function, especially in a prolonged treatment, may cause or exacerbate autoimmune response.

[544] Increasing evidence suggests that Tregs not only act to dampen over exuberant effector immune responses, they also have the ability to potentiate appropriate immune responses to pathogens, by participating in pathogen clearance and protection of the host from collateral damage. Such diverse function of Treg cells is particularly apparent in delicate tissues such as the lung, which is constantly exposed to an external environment from which a variety of pathogens and other foreign components (*e.g.*, viral pathogens, bacterial pathogens, fungal pathogens, and allergens) may gain access to host cells.

[545] For example, influenza virus infection elicits a strong proinflammatory cytokine response with abundance immune cell infiltration. In acute and/or severe infections, such response can cause serious sequelae in susceptible individuals. Tregs provide a mechanism for dampening viral infection-associated pathology by controlling the magnitude of immune response in the host. Indeed, pathogen-exposed Tregs retain protective effects in adoptive transfer. Moreover, such adoptive transfer of primed Tregs have been shown to ameliorate influenza virus-associated morbidity and to prolong survival in severe immunocompromised animal models.

[546] Accordingly, the invention provides use of an ECM-targeted, context-selective TGF β inhibitor (*e.g.*, LTBP1-selective or LTBP1/3-selective inhibitors of TGF β 1 activation inhibitors) for the treatment of a disease that involves matrix-associated TGF β dysregulation in a subject. The subject is suffering from or at risk of an infection. The infection can be viral infections (*e.g.*, influenza virus, respiratory syncytial virus or RSV, human immunodeficiency virus or HIV, MARS, SARS, herpes simplex virus or HSV, hepatitis A virus or HAV, hepatitis B virus or HBV, hepatitis C virus or HCV, CMV, Dengue virus, lymphocytic choriomeningitis virus, and West Nile virus), bacterial infections (meningitis, *Mycobacterium tuberculosis*, *Listeria monocytogenes*, *Citrobacter rodentium*, *Salmonella*, and *E. coli*), and/or fungal infections (*e.g.*, *Candida*, *Pneumocystis*, *Aspergillus*, *Cryptococcus*, and *Coccidioides*).

[547] Typically, high-risk or at-risk populations (individuals that are considered particularly susceptible to severe infections or infection-triggered responses) include pediatric populations (infants, young children, *e.g.*, human individuals under the age of 7); elderly populations (those who are 65 years or older); those with compromised immune system due to medical condition, health status, life styles such as smoking, and/or medications with immunosuppressive effects, etc.

[548] For example, certain medications cause weakened immunity, such as chemotherapy, therapies that target hematopoietic cells such as CD33 therapy, steroids, immunosuppressants, and statins.

[549] In some embodiments, high-risk or at-risk populations are those with existing medical conditions, such as those with chronic infections such as HIV, those with bone marrow transplantation, pre-diabetic individuals, diabetic individuals, those with autoimmune disorders such as RA, asthma and allergy.

[550] Thus, matrix-targeted, context-selective TGF β inhibitors encompassed herein may be particularly advantageous for treating patients who require a long-term or chronic TGF β therapy since in these scenarios it is beneficial to avoid impairment of immune homeostasis and the normal immune function that provides the ability to respond effectively to possible infections caused by a variety of pathogens such as those listed above.

[551] Accordingly, antibodies that selectively bind LTBP-TGF β (*e.g.*, LTBP1-TGF β 1 and LTBP3-TGF β 1), and that do not inhibit TGF β in the context of the immune-associated TGF β presenters GARP and LRRC33, are therapeutic candidates for the treatment of fibrotic indications such as organ fibrosis, and are aimed to avoid TGF β -related global immune activation. In one embodiment, the context-specific antibodies can be used to inhibit LTBP1/3-associated TGF β activity in applications in which TGF β -mediated immune suppression is beneficial, *e.g.*, in a subject who has received a transplant, who is a candidate for receiving a transplant, or who is expected to receive a transplant. In some embodiments, the subject has an advanced stage fibrosis and/or a bone marrow disease. In some embodiments, the subject has or is at risk of developing an autoimmune disorder.

[552] The foregoing methods can be used to treat a subject having a condition for which inhibition or reduction in LTBP-associated TGF β activity is beneficial. For example, the subject may have or be at risk for developing a disorder in which extracellular matrix-associated TGF β activity has been implicated.

[553] Integrin-mediated activation of latent TGF β in the extracellular matrix is a key contributor to fibrosis. Without wishing to be bound by theory, it is presently understood that integrins, including α V β 6 and α V β 8, can trigger the release of TGF β from presenting molecules including LTBP1 and LTBP3. Inhibiting release or activation of TGF β in this context can reduce or eliminate fibrosis, and/or symptoms associated therewith.

[554] As described, LTBP1 and LTBP3 are produced and are deposited extracellularly as components of the ECM, where they can “present” a proTGF β complex (latent, inactive precursor of TGF β 1) within the ECM. Upon stimulation, the LTBP1/3-proTGF β complex releases the TGF β growth factor (the active, mature form of growth factor) which in turn is thought to be involved in the regulation of the local tissue microenvironment, such as ECM maintenance/remodeling and the process of fibrosis, possibly by responding to various cytokines, chemokines and growth factors, and by interacting with other ECM components, such as fibronectin, Fibrillin, collagen, elastin, and matrix metalloproteinases (MMPs).

[555] In the normal wound healing process that occurs in response to an injury, for example, TGF β is thought to facilitate granular tissue formation, angiogenesis, and collagen synthesis and production. TGF β signaling is also implicated in abnormal tissue fibrogenesis (*i.e.*, fibrosis), which results in formation of excess fibrous connective tissue in an organ or tissue in a reparative or reactive process characterized by the pathological accumulation of extracellular matrix (ECM) components, such as collagens. In these and other situations, the TGF β axis may affect further aspects (in addition to fibrotic aspect), such as inflammation, recruitment and phenotypic switch of various cell types, which may be mediated by its interaction with one or more of the other presenting molecules, such as GARP/LRRC32 and LRRC33. In certain instances, it is advantageous to preferentially inhibit the LTBP1/3-context of TGF β activation, without significantly inhibiting one or more of the other contexts of TGF β 1 activation, in situations where ECM-associated TGF β that drives fibrosis is to be selectively inhibited.

[556] Accordingly, in one embodiment, the invention provides a method of reducing TGF β activation in a subject having, or at risk of developing, a fibrotic disorder by administering to the subject an antibody, or antigen-binding portion thereof, which selectively binds a LTBP1/3-TGF β complex, as described herein. In another embodiment, the invention provides a method of treating a fibrotic disorder by administering to the subject an antibody, or antigen-binding portion thereof, which selectively binds a LTBP1/3-TGF β complex, as described herein.

[557] In one embodiment, the fibrotic disorder is an organ fibrosis, wherein optionally, the organ fibrosis is an advanced organ fibrosis. In a further embodiment, the organ fibrosis is selected from the group consisting of kidney fibrosis, liver fibrosis, lung fibrosis, cardiac fibrosis, pancreatic fibrosis, skin fibrosis, scleroderma, muscle fibrosis, uterine fibrosis and endometriosis. In another further embodiment, the fibrotic disorder comprising chronic inflammation is a muscular dystrophy, multiple sclerosis (MS), or Cystic Fibrosis (CF). In a further embodiment, the muscular dystrophy is Duchenne muscular dystrophy (DMD). In another further embodiment, the MS comprises perivascular fibrosis. In a further embodiment, the lung fibrosis is idiopathic pulmonary fibrosis (IPF). In another further embodiment, the subject has chronic kidney disease (CKD). In another embodiment, the subject has nonalcoholic steatohepatitis (NASH).

[558] In exemplary embodiments, the fibrotic disorder is fibrosis, Alport syndrome, fibroids, desmoplasia, amyotrophic lateral sclerosis (ALS), or Duchenne muscular dystrophy (DMD).

[559] In one embodiment, the subject has desmoplasia.

[560] In one embodiment, the subject has organ fibrosis, for example, kidney fibrosis (*e.g.*, fibrosis associated with chronic kidney disease (CKD)), liver fibrosis (*e.g.*, fibrosis associated with nonalcoholic steatohepatitis (NASH)), lung fibrosis (*e.g.*, idiopathic pulmonary fibrosis (IPF)), cardiac fibrosis, and/or skin fibrosis (*e.g.*, scleroderma). In some embodiments, the subject can have advanced organ fibrosis. For example, the subject may be in need of an organ transplant. In one embodiment, the subject may be in need of an organ transplant, and the compounds and compositions described herein are administered to prevent allograft fibrosis from developing in the subject following receipt of the transplant.

[561] A recent study examined whether inhibiting integrin $\alpha V\beta 6$ could prevent TGF β -mediated allograft fibrosis after kidney transplantation (Lo *et al.*, Am. J. Transplant. (2013), 13:3085-3093). Surprisingly, animals treated with an inhibitory anti- $\alpha V\beta 6$ antibody experienced a significant decrease in rejection-free survival compared to placebo animals. The authors conclude that this result cautions against TGF β inhibition in kidney transplantation, because the immunosuppressive properties of TGF β help prevent allograft rejection. The inhibitors, *e.g.*, antibodies, and antigen-binding portions thereof, described herein advantageously inhibit activation of TGF β presented by LTBP1 or LTBP3 in the extracellular matrix, but do not inhibit activation of TGF β presented by GARP or LRRC33 on immune cells. Accordingly, the context-specific LTBP1/3-TGF β inhibitors, *e.g.*, antibodies, described herein can prevent or reduce allograft fibrosis, without eliminating the immunosuppressive properties of TGF β that are useful for preventing allograft rejection. Accordingly, in one aspect, the invention provides a method for treating a fibrotic disorder in a subject, comprising administering to the subject a therapeutically effective amount of an inhibitor of TGF β signaling, wherein the inhibitor is a selective inhibitor of ECM-associated TGF β ; and, wherein the subject benefits from suppressed

immunity. In one embodiment, the subject has a fibrotic condition and would benefit from an allograft transplant, or has received an allograft transplant.

[562] Additional fibrotic conditions for which antibodies and/or compositions of the present disclosure may be used therapeutically include, but are not limited to, lung indications (*e.g.*, idiopathic pulmonary fibrosis (IPF), chronic obstructive pulmonary disorder (COPD), allergic asthma, cystic fibrosis (CF), acute lung injury, eosinophilic esophagitis, pulmonary arterial hypertension and chemical gas-injury), kidney indications (*e.g.*, diabetic glomerulosclerosis, focal segmental glomerulosclerosis (FSGS), chronic kidney disease, fibrosis associated with kidney transplantation and chronic rejection, IgA nephropathy, and hemolytic uremic syndrome), liver fibrosis (*e.g.*, non-alcoholic steatohepatitis (NASH), chronic viral hepatitis, parasitemia, inborn errors of metabolism, toxin-mediated fibrosis, such as alcohol fibrosis, non-alcoholic steatohepatitis-hepatocellular carcinoma (NASH-HCC), primary biliary cirrhosis, and sclerosing cholangitis), cardiovascular fibrosis (*e.g.*, cardiomyopathy, hypertrophic cardiomyopathy, atherosclerosis and restenosis,) systemic sclerosis, skin fibrosis (*e.g.*, skin fibrosis in systemic sclerosis, diffuse cutaneous systemic sclerosis, scleroderma, pathological skin scarring, keloid, post-surgical scarring, scar revision surgery, radiation-induced scarring and chronic wounds), eye-related conditions such as subretinal fibrosis, uveitis syndrome, uveitis associated with idiopathic retroperitoneal fibrosis, extraocular muscle fibrosis, eye diseases associated with the major histocompatibility complex (MHC class I) or histocompatibility antigens, subretinal fibrosis in macular degeneration (*e.g.*, age-related macular degeneration) and cancers or secondary fibrosis (*e.g.*, myelofibrosis, head and neck cancer, M7 acute megakaryoblastic leukemia and mucositis). Other diseases, disorders or conditions related to fibrosis that may be treated using compounds and/or compositions of the present disclosure, include, but are not limited to Marfan's syndrome, stiff skin syndrome, scleroderma, rheumatoid arthritis, bone marrow fibrosis, Crohn's disease, ulcerative colitis, systemic lupus erythematosus, muscular dystrophy, (such as DMD), Dupuytren's contracture, Camurati-Engelmann disease, neural scarring, dementia, proliferative vitreoretinopathy, corneal injury, complications after glaucoma drainage surgery, and multiple sclerosis (MS). Many such fibrotic indications are also associated with inflammation of the affected tissue(s), indicating involvement of an immune component. Such inflammation may be accompanied by aberrant immune cell populations, such as increased numbers of Th17 cells, reduced numbers of Treg cells, and/or both. In each case, the affected patient may exhibit increased Th17/Treg cell ratios. In some embodiments, diseases to be treated with an antibody according to the present disclosure include metabolic disorders, such as metabolic liver disorders. Non-limiting examples of metabolic disorders include NASH, NAFLD, type 2 diabetes and obesity. In some embodiments, the disease to be treated with an antibody according to the present disclosure is aortic stenosis.

[563] In another aspect, the invention provides a method of selecting an isoform-specific TGFβ1 inhibitor for the treatment of a fibrotic disorder in a subject, comprising: (a) determining whether the subject manifests clinical presentations including fibrosis and one or more of the following: (i) inflammation; (ii) immune suppression; (iii) proliferative dysregulation; (iv) need for an allograft transplant; (v) at risk of severe infection; (vi) in need of a long-term TGFβ1 inhibition therapy; and (vii) manifestation of an autoimmune conditions(s); and (b) selecting an isoform-specific, context-dependent TGFβ1 inhibitor or an isoform-specific, context-independent TGFβ1 inhibitor for treatment of the fibrotic disorder based on the clinical presentations determined in step (a).

[564] In another aspect, the invention provides a method of treating a subject having a fibrotic disorder, comprising (a) selecting a treatment regimen comprising an isoform-specific TGFβ1 inhibitor for the subject, said selection comprising (i) determining whether the fibrotic disorder manifests clinical presentations including fibrosis and one or more of the following: inflammation, immune suppression, proliferative dysregulation, and need for an allograft transplant; and (ii) selecting a treatment regimen comprising an isoform-specific, context-dependent TGFβ1 inhibitor or an isoform-specific, context-independent TGFβ1 inhibitor, based on the clinical presentations determined in step (i); and (b) administering the selected treatment regimen to the subject.

[565] In one embodiment of the foregoing aspects, the fibrotic disorder manifests clinical presentations comprising fibrosis, inflammation, immune suppression, and proliferative dysregulation. In an exemplary embodiment, the fibrotic disorder is myelofibrosis, and the selected isoform-specific TGFβ1 inhibitor is an isoform-specific, context-independent TGFβ1 inhibitor.

[566] In another embodiment, the fibrotic disorder manifests clinical presentations comprising fibrosis, inflammation, and need for an allograft transplant. In one embodiment, the fibrotic disorder manifests clinical presentations comprising fibrosis and inflammation. In another embodiment, the fibrotic disorder is a degenerative disease.

[567] In one embodiment, the fibrotic disorder manifests clinical presentations comprising immune suppression and proliferative dysregulation. In an exemplary embodiment, the fibrotic disorder is associated with a solid tumor, and the selected isoform-specific TGFβ1 inhibitor is an isoform-specific LTBP1/3-specific inhibitor and/or a GARP-selective inhibitor. In one embodiment, the solid tumor is a malignant tumor. In another embodiment, the tumor is a benign tumor. In one embodiment, the subject has desmoplasia, for example, pancreatic desmoplasia. In another embodiment, the subject has fibroids.

[568] In another aspect, the invention provides a method of treating a subject having a fibrotic disorder with an isoform-specific, LTBP1/3-specific TGFβ1 inhibitor, comprising determining whether the fibrotic disorder manifests clinical presentations including fibrosis and the need for an allograft transplant; and administering an effective amount of an isoform-specific, LTBP1/3-specific

TGFβ1 inhibitor to the subject if the fibrotic disorder manifests fibrosis and the need for an allograft transplant.

[569] In another aspect, the invention provides a method of treating a subject having a fibrotic disorder with an isoform-specific, context-independent TGFβ1 inhibitor, comprising determining whether the fibrotic disorder manifests clinical presentations including fibrosis, immune suppression and/or proliferative dysregulation; and administering an effective amount of an isoform-specific, context-independent TGFβ1 inhibitor to the subject if the fibrotic disorder manifests fibrosis in conjunction with immune suppression and/or proliferative dysregulation.

[570] The inhibitors, *e.g.*, antibodies, described herein can be administered to a subject in an amount effective to treat or reduce symptoms of fibrosis. The effective amount of such an inhibitor is an amount effective to achieve both therapeutic efficacy and clinical safety in the subject. In one embodiment, an effective amount is an amount effective to reduce TGFβ1 activity in the extracellular matrix. In another embodiment, an effective amount is an amount effective to reduce fibrosis in a subject. In another embodiment, the effective amount does not inhibit TGFβ1-mediated immune suppression. In some embodiments, such an inhibitor, *e.g.*, antibody, is a context-specific inhibitor that can block activation of TGFβ1 that is mediated by an LTBP-containing, ECM-associated TGFβ1. In some embodiments, the LTBP is LTBP1 and/or LTBP3. Assays useful for determining the efficacy of the inhibitors, *e.g.*, antibodies, and/or compositions of the present disclosure for the alteration of fibrosis include, but are not limited to, histological assays for counting fibroblasts and basic immunohistochemical analyses known in the art.

Diseases Involving Proteases:

[571] Activation of TGFβ from its latent complex may be triggered by integrin in a force-dependent manner, and/or by proteases. Evidence suggests that certain classes of proteases may be involved in the process, including but are not limited to Ser/Thr proteases such as thrombin, Kallikreins, chymotrypsin, elastases, plasmin, as well as zinc metalloproteases of ADAM family such as ADAM 10 and ADAM 17, as well as MMP family, such as MMP-2, MMP-9 and MMP-13. MMP-2 degrades the most abundant component of the basement membrane, Collagen IV, raising the possibility that it may play a role in ECM-associated TGFβ1 regulation. MMP-9 has been implicated to play a central role in tumor progression, angiogenesis, stromal remodeling and metastasis. Thus, protease-dependent activation of TGFβ1 in the ECM may be important for treating cancer.

[572] Kallikreins (KLKs) are trypsin- or chymotrypsin-like serine proteases that include plasma Kallikreins and tissue Kallikreins. The ECM plays a role in tissue homeostasis acting as a structural and signaling scaffold and barrier to suppress malignant outgrowth. KLKs may play a role in degrading ECM proteins and other components which may facilitate tumor expansion and invasion. For example, KLK1 is highly upregulated in certain breast cancers and can activate pro-MMP-2 and

pro-MMP-9. KLK2 activates latent TGFβ1, rendering prostate cancer adjacent to fibroblasts permissive to cancer growth. KLK3 has been widely studied as a diagnostic marker for prostate cancer (PSA). KLK3 may directly activate TGFβ1 by processing plasminogen into plasmin, which proteolytically cleaves LAP. KLK6 may be a potential marker for Alzheimer's disease.

[573] Known activators of TGFβ1, such as plasmin, TSP-1 and αVβ6 integrin, all interact directly with LAP. It is postulated that proteolytic cleavage of LAP may destabilize the LAP-TGFβ interaction, thereby releasing active TGFβ1. It has been suggested that the region containing 54-LSKLRL-59 is important for maintaining TGFβ1 latency. Thus, agents (*e.g.*, antibodies) that stabilize the interaction, or block the proteolytic cleavage of LAP may prevent TGFβ activation.

[574] Many of these proteases associated with pathological conditions (*e.g.*, cancer) function through distinct mechanisms of action. Thus, targeted inhibition of particular proteases, or combinations of proteases, may provide therapeutic benefits for the treatment of conditions involving the protease-TGFβ axis. Accordingly, it is contemplated that inhibitors (*e.g.*, TGFβ1 antibodies) that selectively inhibit protease-induced activation of TGFβ1 may be advantageous in the treatment of such diseases (*e.g.*, cancer). Similarly, selective inhibition of TGFβ1 activation by one protease over another protease may also be preferred, depending on the condition being treated.

[575] Plasmin is a serine protease produced as a precursor form called Plasminogen. Upon release, Plasmin enters circulation and therefore is detected in serum. Elevated levels of Plasmin appear to correlate with cancer progression, possibly through mechanisms involving disruption of the extracellular matrix (*e.g.*, basement membrane and stromal barriers) which facilitates tumor cell motility, invasion, and metastasis. Plasmin may also affect adhesion, proliferation, apoptosis, cancer nutrition, oxygen supply, formation of blood vessels, and activation of VEGF (Didiasova *et al.*, *Int. J. Mol. Sci.*, 2014, 15, 21229-21252). In addition, Plasmin may promote the migration of macrophages into the tumor microenvironment (Philips *et al.*, *Cancer Res.* 2011 Nov 1;71(21):6676-83 and Choong *et al.*, *Clin. Orthop. Relat. Res.* 2003, 415S, S46-S58). Indeed, tumor-associated macrophages (TAMs) are well characterized drivers of tumorigenesis through their ability to promote tumor growth, invasion, metastasis, and angiogenesis.

[576] Plasmin activities have been primarily tied to the disruption of the ECM. However, there is mounting evidence that Plasmin also regulate downstream MMP and TGF beta activation. Specifically, Plasmin has been suggested to cause activation of TGF beta through proteolytic cleavage of the Latency Associated Peptide (LAP), which is derived from the N-terminal region of the TGF beta gene product (Horiguchi *et al.*, *J Biochem.* 2012 Oct; 152(4):321-9), resulting in the release of active growth factor. Since TGFβ1 may promote cancer progression, this raises the possibility that plasmin-induced activation of TGFβ may at least in part mediate this process.

[577] TGF β 1 has also been shown to regulate expression of uPA, which is a critical player in the conversion of Plasminogen into Plasmin (Santibanez, Juan F., *ISRN Dermatology*, 2013: 597927). uPA has independently been shown to promote cancer progression (*e.g.*, adhesion, proliferation, and migration) by binding to its cell surface receptor (uPAR) and promoting conversion of Plasminogen into Plasmin. Moreover, studies have shown that expression of uPA and/or plasminogen activator inhibitor-1 (PAI-1) are predictors of poor prognosis in colorectal cancer (D. Q. Seetoo, *et al.*, *Journal of Surgical Oncology*, vol. 82, no. 3, pp. 184–193, 2003), breast cancer (N. Harbeck *et al.*, *Clinical Breast Cancer*, vol. 5, no. 5, pp. 348–352, 2004), and skin cancer (Santibanez, Juan F., *ISRN Dermatology*, 2013: 597927). Thus, without wishing to be bound by a particular theory, the interplay between Plasmin, TGF β 1, and uPA may create a positive feedback loop towards promoting cancer progression. Accordingly, inhibitors that selectively inhibit Plasmin-dependent TGF β 1 activation may be particularly suitable for the treatment of cancers reliant on the Plasmin/TGF β 1 signaling axis.

[578] In one aspect of the invention, the isoform-specific inhibitors of TGF β 1 described herein include inhibitors that can inhibit protease-dependent activation of TGF β 1. In some embodiments, the inhibitors can inhibit protease-dependent TGF β 1 activation in an integrin-independent manner. In some embodiments, such inhibitors can inhibit TGF β 1 activation irrespective of the mode of activation, *e.g.*, inhibit both integrin-dependent activation and protease-dependent activation of TGF β 1. In some embodiments, the protease is selected from the group consisting of: serine proteases, such as Kallikreins, Chymotrypsin, Trypsin, Elastases, Plasmin, as well as zinc metalloproteases (MMP family) such as MMP-2, MMP-9 and MMP-13.

[579] In some embodiments, the inhibitors can inhibit Plasmin-induced activation of TGF β 1. In some embodiments, the inhibitors can inhibit Plasmin- and integrin-induced TGF β 1 activation. In some embodiments, the inhibitors are monoclonal antibodies that specifically bind TGF β 1. In some embodiments, the antibody is a monoclonal antibody that specifically binds proTGF β 1. In some embodiments, the antibody binds latent proTGF β 1 thereby inhibiting release of mature growth factor from the latent complex. In some embodiments, the high-affinity, LTBP-complex specific inhibitor of TGF β 1 activation suitable for use in the method of inhibiting Plasmin-dependent activation of TGF β 1. In some embodiments, the LTBP-complex specific inhibitor of TGF β 1 activation is selected from Ab31, Ab34, Ab37, Ab38, Ab39, Ab40, Ab41, Ab42, Ab43, Ab44, Ab45, Ab62, Ab63, and Ab64 (optionally Ab42 or Ab63) (*i.e.*, an antibody or antigen-binding fragment having the heavy and light chain variable regions of the corresponding Ab, as provided herein) a variant/derivative or antigen-binding fragment thereof, or an engineered molecule comprising an antigen-binding fragment thereof. In some preferred embodiments, the LTBP-complex specific inhibitor of TGF β 1 activation is Ab42, a variant/derivative or antigen-binding fragment thereof, or an engineered molecule comprising an antigen-binding fragment thereof. In preferred embodiments, the LTBP-complex specific inhibitor of TGF β 1 activation is Ab42 or an antigen-binding fragment thereof.

[580] In some embodiments, the inhibitor (*e.g.*, TGF β 1 antibody) inhibits cancer cell migration. In some embodiments, the inhibitor inhibits macrophage migration. In some embodiments, the inhibitor inhibits accumulation of TAMs.

[581] In another aspect, provided herein is a method for treating cancer in a subject in need thereof, the method comprising administering to the subject an effective amount of an TGF β 1 inhibitor (*e.g.*, TGF β 1 antibody), wherein the inhibitor inhibits protease-induced activation of TGF β 1 (*e.g.*, Plasmin), thereby treating cancer in the subject.

[582] In another aspect, provided herein is a method of reducing tumor growth in a subject in need thereof, the method comprising administering to the subject an effective amount of an TGF β 1 inhibitor (*e.g.*, TGF β 1 antibody), wherein the inhibitor inhibits protease-induced activation of TGF β 1 (*e.g.*, Plasmin), thereby reducing tumor growth in the subject.

Disease involving ECM dysregulation

[583] The extracellular matrix is a cell-secreted network that surrounds cells and is primarily composed of proteoglycans and fibrous proteins, the most abundant of which is collagen. The novel antibodies disclosed herein may be used in the treatment of diseases associated with extracellular matrix dysregulation. The diseases associated with extracellular matrix dysregulation are typically myofibroblast-driven pathologies and include cancer, fibrosis, and cardiovascular disease (reviewed, for example, in: Lampi and Reinhart-King (2018) "Targeting extracellular matrix stiffness to attenuate disease: From molecular mechanisms to clinical trials" *Sci Transl Med* 10(422): eaao0475). Progression of fibrotic conditions involves increased levels of matrix components deposited into the ECM and/or maintenance/remodeling of the ECM. TGF β 1 at least in part contributes to this process. This is supported, for example, by the observation that increased deposition of ECM components such as collagens can alter the mechanophysical properties of the ECM (*e.g.*, the stiffness of the matrix/substrate) and this phenomenon is associated with TGF β 1 signaling. The inhibitors of TGF β 1, such as those described herein may be used to block this process to counter disease progression involving ECM alterations, such as fibrosis. The LTBP-arm of such inhibitors can directly block ECM-associated pro/latent TGF β complexes which are presented by LTBP1 and/or LTBP3, thereby preventing activation/release of the growth factor from the complex in the disease niche. In some embodiments, the isoform-specific TGF β 1 inhibitors such as those described herein may normalize ECM stiffness to treat a disease that involves integrin-dependent signaling. In some embodiments, the integrin comprises an α 11 chain, β 1 chain, or both.

[584] Thus, the antibody may be administered to a subject diagnosed with a disease with extracellular matrix dysregulation in an amount effective to treat the disease. Therapeutically effective amount of the antibody may be an amount sufficient to reduce expression of one or more markers of myofibroblasts, such as α -SMA. The amount may be an amount sufficient to reduce the

stiffness of the extracellular matrix of an affected tissue (*e.g.*, fibrotic tissues). The amount may be an amount sufficient to reduce TGF β 1 downstream effectors, such as phosphorylation of SMAD2 and/or SMAD3. In some embodiments, the isoform-selective activation inhibitor of TGF β 1 is selected from Ab31, Ab34, Ab37, Ab38, Ab39, Ab40, Ab41, Ab42, Ab43, Ab44, Ab45, Ab62, Ab63, and Ab64 (optionally Ab42 or Ab63) a variant/derivative or antigen-binding fragment thereof, or an engineered molecule comprising an antigen-binding fragment thereof. In some preferred embodiments, the isoform-selective activation inhibitor of TGF β 1 is Ab42, a variant/derivative or antigen-binding fragment thereof, or an engineered molecule comprising an antigen-binding fragment thereof. In preferred embodiments, the TGF β 1-selective inhibitor is Ab42 or an antigen-binding fragment thereof.

Diseases Involving Epithelial-to-Mesenchymal Transition (EMT):

[585] EMT (epithelial mesenchymal transition) is the process by which epithelial cells with tight junctions switch to mesenchymal properties (phenotypes) such as loose cell-cell contacts. The process is observed in a number of normal biological processes as well as pathological situations, including embryogenesis, wound healing, cancer metastasis and fibrosis (reviewed in, for example, Shiga et al. (2015) "Cancer-Associated Fibroblasts: Their Characteristics and Their Roles in Tumor Growth." *Cancers*, 7: 2443-2458). Generally, it is believed that EMT signals are induced mainly by TGF β . Many types of cancer, for example, appear to involve transdifferentiation of cells towards mesenchymal phenotype (such as CAFs) which correlate with poorer prognosis. Thus, LTBP-specific inhibitors of TGF β 1, such as those described herein, may be used to treat a disease that is initiated or driven by EMT. Indeed, data exemplified herein (*e.g.*, FIGs. 12 and 13) show that such inhibitors have the ability to suppress expression of CAF markers *in vivo*, such as α -SMA, Col1 (Type I collagen), and FN (fibronectin).

Diseases involving Matrix Stiffening and Remodeling

[586] Progression of fibrotic conditions involves increased levels of matrix components deposited into the ECM and/or maintenance/remodeling of the ECM. TGF β 1 at least in part contributes to this process. This is supported, for example, by the observation that increased deposition of ECM components such as collagens can alter the mechanophysical properties of the ECM (*e.g.*, the stiffness of the matrix/substrate) and this phenomenon is associated with TGF β 1 signaling. To confirm this notion, the present inventors have evaluated the role of matrix stiffness in affecting integrin-dependent activation of TGF β in primary fibroblasts transfected with proTGF β and LTBP1, and grown on silicon-based substrates with defined stiffness (*e.g.*, 5 kPa, 15 kPa or 100 kPa). Matrices with greater stiffness enhance TGF β 1 activation, and this can be suppressed by antibodies, and antigen-binding portions thereof, which are capable of binding and thereby inhibiting TGF β 1 activation associated with LTBP1/3. These observations suggest that TGF β 1 influences ECM

properties (such as stiffness), which in turn can further induce TGF β 1 activation, reflective of disease progression. Thus, antibodies, and antigen-binding portions thereof, that selectively bind complexes of LTBP1-TGF β 1 and/or LTBP3-TGF β 1, such as those described herein may be used to block this process to counter disease progression involving ECM alterations, such as fibrosis, tumor growth, invasion, metastasis and desmoplasia. Such inhibitors can directly block ECM-associated pro/latent TGF β complexes which are presented by LTBP1 and/or LTBP3, thereby preventing activation/release of the growth factor from the complex in the disease niche.

Fibrosis:

[587] In response to tissue injury or chronic insult due to physical damage/trauma, toxic substances, and/or infection, a natural reparative process begins which involves several cell types including fibroblasts, several different types of immune cells, and resident epithelial and endothelial cells. However, if left unchecked, this process can lead to excessive accumulation of extracellular matrix (ECM) and fibrosis, which in turn can lead to progressive loss of tissue function and organ failure (Caja et al., *Int. J. Mol. Sci.* 2018, 19, 1294).

[588] Fibrosis can occur in several different organs, including lung, kidney, liver, heart, and skin. Independent of the organ, the fibrotic response is characterized by inflammation, altered epithelial-mesenchymal interactions, and proliferation of fibroblasts. One of the hallmarks of fibrosis is the differentiation of fibroblasts into myofibroblasts, which greatly contribute to the dysregulation of the ECM. However, myofibroblasts have also been proposed to come from other cellular sources (*e.g.*, endothelial cells, epithelial cells, and mesenchymal stem cells (Kim, K.K. et al, Cold Spring Harb. Perspect. Biol., 2017; Okabe, H. *Histol. Histopathol.*, 2016, 31, 141-148; and Li, C et al, *Nat Commun.*, 2016, 7, 11455). Moreover, immune cells play an important role in the process by secreting cytokines and chemokines which promote differentiation of myofibroblasts, stimulate ECM deposition, and recruit additional immune cells to the damaged tissue (Caja et al., *Int. J. Mol. Sci.* 2018, 19, 1294).

[589] Similar to fibrotic tissue, activation of cancer-associated fibroblasts can occur in the tumor milieu, which produces excessive amounts of ECM. The ECM provides a scaffold for the infiltration of other cells (*e.g.*, pro-tumorigenic immune cells) and a substrate for cell migration. In other cases, excessive ECM may act as a barrier against anti-tumorigenic immune cells.

[590] TGF β is recognized as the central orchestrator of the fibrotic response. TGF β can promote myofibroblast differentiation, recruit immune cells, and affect epithelial and endothelial cell differentiation. Particularly, TGF β upregulates the production of ECM and basement membrane proteins, such as fibronectin, collagen, laminin, osteopontin, tenascin, elastin, decorin. TGF β -induced myofibroblast differentiation can lead to additional deposition of ECM proteins, secretion of matrix metalloproteinase (MMPs), and myofibroblast proliferation (Fabregat et al, *FEBS J.* 2016, 283, 2219-

2232; Meng et al, *Nat. Rev. Nephrol.* 2016, 12, 325-338; and Kulkarni et al., *Am. J. Respir. Cell Mol. Biol.*, 2016, 54, 751-760). Additionally, TGF β mediates phenotypic changes affecting contractile proteins and collagen I in vascular smooth muscle cells (VSCM), and can activate myofibroblasts and other stromal cells to enhance the synthesis of collagen cross-linking proteins, such as lysyl oxidase (LOX) family of matrix-remodeling enzymes (Busnadiago et al., *Mol. Cell. Biol.* 2013, 33, 2388-2401). Moreover, TGF β has been shown to regulate both EMT and EndMT, which contributes to the differentiation of pro-fibrotic cell types, such as myofibroblasts and CAFs. Moreover, TGF β has been shown to induce epithelial apoptosis, which can promote lung and liver fibrosis among other tissues (Barbas-Filho et al., *J. Clin. Pathol.* 2001, 54, 132-138; and Wang et al., *Dev. Dyn.* 2017, 247, 492-508).

[591] Whether innate or recruited, macrophages are thought to play an important role in responding to tissue damage and repair. However, upon certain signals they can become pro-fibrotic. TGF β , among other cytokines, has also been shown to activate M2 macrophages, which are pro-inflammatory. Upon activation, these macrophages secrete their own cytokines, including TGF β , ECM components, angiogenic factors, and chemotactic factors. M2 macrophages have been shown to be essential for TGF β -driven lung fibrosis (Murray et al., *Int. J. Biochem. Cell Biol.* 2011, 43, 154-162).

[592] In light of increasing evidence pointing to the importance of M2-type macrophages for disease progression in many types of fibrosis, a question remained as to whether context-selective inhibition of LTBP1/3-associated TGF β 1 alone (that is, without addressing the macrophage-associated, LRRC33-arm of TGF β 1 activity) might be sufficient to produce a potent anti-fibrotic effect in vivo. Surprisingly, however, data presented herein suggest that selectively targeting the matrix-associated TGF β 1 (*e.g.*, LTBP1/3-proTGF β 1) appears to be just as effective – if not better – in achieving anti-fibrotic effects in multiple preclinical models, as targeting all four known LLCs (*e.g.*, LTBP1/3-proTGF β 1, GARP-proTGF β 1 and LRRC33-proTGF β 1) with the use of a so-called context-independent inhibitor of TGF β 1 (see, for example, FIGs. 19 and 20) and do so without triggering T cell stimulation mediated via GARP-proTGF β 1 inhibition. Moreover, previously disclosed LTBP1/3 complex-selective antibodies lacked robust species cross-reactivity that would be advantageous for both preclinical (*e.g.*, rodent) and clinical (*e.g.*, human) use. It was not clear whether the rare epitopes being sought which would confer both isoform-selectivity and context-selectivity would also enable favorable species cross-reactivity profiles. Advantageously, novel antibodies disclosed herein possess all of these criteria.

[593] According to the invention, isoform-specific TGF β 1 such as those described herein are used in the treatment of fibrosis (*e.g.*, fibrotic indications, fibrotic conditions) in a subject. Suitable inhibitors to carry out the present invention include antibodies and/or compositions according to the

present disclosure which may be useful for altering or ameliorating fibrosis. More specifically, such antibodies and/or compositions are selective antagonists of TGF β 1 that are capable of targeting TGF β 1 presented by various types of presenting molecules. TGF β 1 is recognized as the central orchestrator of the fibrotic response. Antibodies targeting TGF β decrease fibrosis in numerous preclinical models. Such antibodies and/or antibody-based compounds include LY2382770 (Eli Lilly, Indianapolis, IN). Also included are those described in U.S. Patent Numbers US 6,492,497, US 7,151,169, US 7,723,486 and U.S. Appl. Publ. No. 2011/0008364, the contents of each of which are herein incorporated by reference in their entirety. Prior art TGF β antagonists include, for example, agents that target and block integrin-dependent activation of TGF β .

[594] However, evidence suggests that such prior art agents may not mediate isoform-specific inhibition and may cause unwanted effects by inadvertently blocking normal function of TGF β 2 and/or TGF β 3. Indeed, data presented herein support this notion. Normal (undiseased) lung tissues contain relatively low but measurable levels of TGF β 2 and TGF β 3, but notably less TGF β 1. In comparison, in certain disease conditions such as fibrosis, TGF β 1 becomes preferentially upregulated relative to the other isoforms. Preferably, TGF β antagonists for use in the treatment of such conditions exert their inhibitory activities only towards the disease-induced or disease-associated isoform, while preserving the function of the other isoforms that are normally expressed to mediate tonic signaling in the tissue. Prior art inhibitors (LY2109761, a small molecule TGF β receptor antagonist, and a monoclonal antibody that targets α V β 6 integrin) both are shown to inhibit TGF β downstream tonic signaling in non-diseased rat BAL, raising the possibility that these inhibitors may cause unwanted side effects. Alternatively or additionally, agents that target and block integrin-dependent activation of TGF β may be capable of blocking only a subset of integrins responsible for disease-associated TGF β 1 activation, among numerous integrin types that are expressed by various cell types and play a role in the pathogenesis. Furthermore, even where such antagonists may selectively block integrin-mediated activation of the TGF β 1 isoform, it may be ineffective in blocking TGF β 1 activation triggered by other modes, such as protease-dependent activation. Accordingly, the isoform-specific inhibitors of TGF β 1 such as those described herein are aimed to prevent the activation step of TGF β 1 regardless of the particular mode of activation, while maintaining isoform selectivity.

[595] It is further contemplated that isoform-specific TGF β 1 inhibitors that preferentially inhibit matrix-associated over cell-associated antigen complexes (*i.e.*, display context-bias) may offer a therapeutic advantage in certain clinical situations (*e.g.*, the LTBP-specific inhibitors described herein). For example, TGF β 1 context-independent inhibitors (which target all four antigen complexes), may increase immune activation through the targeting of cell-associated TGF β 1 (*e.g.*, GARP-TGF β 1 which is expressed on regulatory T cells). Immune activation may be disadvantageous for certain patients, *e.g.*, patients with autoimmune disease or who are at risk of sepsis. Accordingly,

context-bias antibodies may be useful for treating diseases associate with matrix-associated TGFβ1 complexes (*e.g.*, fibrosis), while minimizing immune activation.

[596] Previously, it was contemplated that isoform-specific TGFβ3 inhibitors might offer an added therapeutic benefit in particular disease states. For example, certain fibrotic diseases to be treated with a TGFβ1 inhibitor may also be TGFβ3-positive (*i.e.*, TGFβ1+/TGFβ3+ fibrotic tissue) characterized in that the disease tissue (*e.g.*, fibrotic tissue) expresses both the isoforms. Accordingly, the invention includes the use of isoform-selective TGFβ1 inhibitor in conjunction with an isoform-selective TGFβ3 inhibitor in the treatment of such conditions.

[597] Fibrotic indications for which antibodies and/or compositions of the present disclosure may be used therapeutically include, but are not limited to lung indications (*e.g.*, idiopathic pulmonary fibrosis (IPF), chronic obstructive pulmonary disorder (COPD), allergic asthma, acute lung injury, eosinophilic esophagitis, pulmonary arterial hypertension and chemical gas-injury), kidney indications (*e.g.*, diabetic glomerulosclerosis, focal segmental glomeruloclerosis (FSGS), chronic kidney disease (CKD), fibrosis associated with kidney transplantation and chronic rejection, IgA nephropathy, and hemolytic uremic syndrome), liver fibrosis (*e.g.*, non-alcoholic steatohepatitis (NASH), chronic viral hepatitis, parasitemia, inborn errors of metabolism, toxin-mediated fibrosis, such as alcohol fibrosis, non-alcoholic steatohepatitis-hepatocellular carcinoma (NASH-HCC), primary biliary cirrhosis, and sclerosing cholangitis), cardiovascular fibrosis (*e.g.*, cardiomyopathy, hypertrophic cardiomyopathy, atherosclerosis and restenosis,) systemic sclerosis, skin fibrosis (*e.g.*, skin fibrosis in systemic sclerosis, diffuse cutaneous systemic sclerosis, scleroderma, pathological skin scarring, keloid, post-surgical scarring, scar revision surgery, radiation-induced scarring and chronic wounds), eye-related conditions such as subretinal fibrosis, uveitis syndrome, uveitis associated with idiopathic retroperitoneal fibrosis, extraocular muscle fibrosis, eye diseases associated with the major histocompatibility complex (MHC class I) or histocompatibility antigens, subretinal fibrosis in macular degeneration (*e.g.*, age-related macular degeneration), and cancers or secondary fibrosis (*e.g.*, myelofibrosis, head and neck cancer, M7 acute megakaryoblastic leukemia and mucositis). Other diseases, disorders or conditions related to fibrosis (including degenerative disorders) that may be treated using compounds and/or compositions of the present disclosure, include, but are not limited to adenomyosis, endometriosis, Marfan's syndrome, stiff skin syndrome, scleroderma, rheumatoid arthritis, bone marrow fibrosis, Crohn's disease, ulcerative colitis, systemic lupus erythematosus, muscular dystrophy (such as DMD), Parkinson's disease, ALS, Dupuytren's contracture, Camurati-Engelmann disease, neural scarring, dementia, proliferative vitreoretinopathy, corneal injury, complications after glaucoma drainage surgery, and multiple sclerosis (MS). Many such fibrotic indications are also associated with inflammation of the affected tissue(s), indicating involvement of an immune component. Such inflammation may be accompanied by aberrant immune cell

populations, such as increased numbers of Th17 cells, reduced numbers of Treg cells, and/or both. In each case, the affected patient may exhibit increased Th17/Treg cell ratios.

[598] In some embodiments, fibrotic indications that may be treated with the compositions and/or methods described herein include organ fibrosis, such as fibrosis of the lung (*e.g.*, IPF), fibrosis of the kidney (*e.g.*, fibrosis associated with CKD), fibrosis of the liver, fibrosis of the heart or cardiac tissues, fibrosis of the skin (*e.g.*, scleroderma), fibrosis of the uterus (*e.g.*, endometrium, myometrium), and fibrosis of the bone marrow. In some embodiments, such therapy may reduce or delay the need for organ transplantation in patients. In some embodiments, such therapy may prolong the survival of the patients.

[599] To treat IPF, patients who may benefit from the treatment include those with familial IPF and those with sporadic IPF. Administration of a therapeutically effective amount of an isoform-specific inhibitor of TGF β 1 may reduce myofibroblast accumulation in the lung tissues, reduce collagen deposits, reduce IPF symptoms, improve or maintain lung function, and prolong survival. In some embodiments, the inhibitor blocks activation of ECM-associated TGF β 1 (*e.g.*, pro/latent TGF β 1 presented by LTBP1/3) within the fibrotic environment of IPF.

[600] Nonalcoholic fatty liver disease (NAFLD) includes a spectrum of histological changes that begin with simple fatty infiltration of the liver, also known as simple or isolated steatosis or nonalcoholic fatty liver (NAFL), which may gradually, sometimes over decades, progress to the development of chronic inflammation (steatohepatitis or NASH), fibrosis, and ultimately cirrhosis. Only a subgroup of patients with NAFL will progress to NASH and subsequent cirrhosis. Currently, there are no clear criteria to identify this group of patients. NAFLD is the most common cause of chronic liver disease in North America. Currently, there are no approved drugs for the treatment of NASH. Given the high prevalence of NASH, the associated morbidity, the growing burden of end-stage liver disease, and limited availability of livers for organ transplantation, identifications of therapies that will slow the progress of, halt, or reverse NASH and NAFLD will address an unmet medical need.

[601] There is a consensus that TGF β is a central player in liver fibrosis (reviewed in, for example, Dewidar et al., *Cells* 2019, 8, 1419, the contents of which are incorporated herein by reference). The isoform-specific TGF β 1 inhibitors such as those provided herein (*i.e.*, isoform-specific inhibitors of TGF β 1 that are selective for LTBP1/3-TGF β 1 complexes, or “matrix-targeted” inhibitors) may be used to treat fibrotic conditions of the liver, such as nonalcoholic fatty liver (NAFL) and fibrosis associated with fatty liver (*e.g.*, NASH). The fatty liver may or may not be inflamed. Inflammation of the liver due to fatty liver (*i.e.*, steatohepatitis) may develop into scarring (fibrosis), which then often progresses to cirrhosis (scarring that distorts the structure of the liver and impairs its function). The inhibitor may therefore be used to treat such conditions. In some embodiments, the inhibitor blocks

activation of ECM-associated TGF β 1 (*e.g.*, pro/latent TGF β 1 presented by LTBP1/3) within the fibrotic environment of the liver. Administration of the inhibitor in a subject with such conditions may reduce one or more symptoms, prevent or retard progression of the disease, reduce or stabilize fat accumulations in the liver, reduce disease-associated biomarkers (such as serum collagen fragments), reduce liver scarring, reduce liver stiffness, and/or otherwise produce clinically meaningful outcome in a patient population treated with the inhibitor, as compared to a control population not treated with the inhibitor. In some embodiments, an effective amount of the inhibitor may achieve both reduced liver fat and reduced fibrosis (*e.g.*, scarring) in NASH patients. In some embodiment, an effective amount of the inhibitor may achieve improvement in fibrosis by at least one stage with no worsening steatohepatitis in NASH patients. In some embodiments, an effective amount of the inhibitor may reduce the rate of occurrence of liver failure and/or liver cancer in NASH patients. In some embodiments, an effective amount of the inhibitor may normalize, as compared to control, the levels of multiple inflammatory or fibrotic serum biomarkers as assessed following the start of the therapy, at, for example, 12-36 weeks. In some embodiments in NASH patients, the isoform-specific TGF β 1 inhibitors may be administered in patients who receive one or more additional therapies, including, but are not limited to myostatin inhibitors, which may generally enhance metabolic regulation in patients with clinical manifestation of metabolic syndrome, including NASH.

[602] In some embodiments, in NASH or NAFLD patients, the isoform-specific, matrix-targeted, TGF β 1 inhibitors may be administered in patients who receive an Acetyl CoA Carboxylase inhibitor (ACCi) (*e.g.*, firsocostat (aka GS-0976) or PF-05221304). Other therapeutics which may be useful in combination with the improved isoform-specific TGF β 1 inhibitors described herein, include, but are not limited to: GLP-1 receptor agonists or analogues (*e.g.*, semaglutide), farnesoid X receptor (FXR) agonists (*e.g.*, GS-9674; aka Cilofexor), ASK1 inhibitors (*e.g.*, selonsertib); obeticholic acid, PPAR agonists (*e.g.*, GFT505; aka elafibranor); nitazoxanide, ketohexokinase (KHK) inhibitors (*e.g.*, PF-06835919); myostatin inhibitors and/or Diacylglycerol O-Acyltransferase 2 (DGAT2) inhibitors (*e.g.*, PF-06865571). In some embodiments, any one or more of the above-mentioned therapeutics can be used in combination with an isoform specific TGF β 1 inhibitor of the present disclosure, for example, an isoform-specific TGF β 1 inhibitor in combination with a FXR agonist, an ACC inhibitor, and/or a GLP-1 analogue. In some embodiments, TGF β inhibitors may be used in combination with a myostatin inhibitor in the treatment of a metabolic liver disease in a subject, such as NASH and NAFLD, and liver fibrosis associated therewith. The subject may also suffer from type 2 diabetes and/or obesity. The TGF β inhibitors used are preferably TGF β 1-selective inhibitors, more preferably context-selective TGF β 1-selective inhibitors that target LTBP1/2-associated TGF β 1, such as those disclosed herein. The myostatin inhibitor is preferably a myostatin-selective inhibitor, such as SRK-015 (*e.g.*, see WO2017/218592A1) and trevogrumab, or any variant thereof, or an antibody according to WO 2016/098357.

[603] In some embodiments, treatment with the isoform specific TGFβ1 inhibitors alone or in combination with one or more additional therapeutics reduces hepatic fat as measured by MRI-PDFF. In some embodiments, the reduction of hepatic fat is at least 20%, *e.g.*, ≥20%, ≥ 25%, ≥ 30%, ≥ 35%, ≥ 40%, ≥ 45%, or ≥ 50%. In some embodiments, treatment with the isoform specific TGFβ1 inhibitors alone or in combination with one or more additional therapeutics reduces serum ALT and/or GGT by at least 20%, *e.g.*, ≥20%, ≥ 25%, ≥ 30%, ≥ 35%, ≥ 40%, ≥ 45%, or ≥ 50%. In some embodiments, treatment with the isoform specific TGFβ1 inhibitors alone or in combination with one or more additional therapeutics reduces bile acid synthesis.

[604] In some embodiments, either as monotherapy or in conjunction with one or more additional therapy (*e.g.*, combination therapy), the TGFβ1 inhibitors of the present disclosure may be effective to treat NASH. “Effective treatment” may refer to improvements in hepatic steatosis, liver stiffness, liver biochemistry and serum fibrosis markers. In some embodiments, a 12-week treatment may result in significant decline of at least 30 percent in hepatic fat measured by magnetic resonance imaging-proton density fat fraction (MRI-PDFF) from baseline to 12 weeks in at least 50% percent of patients. Improvements in liver biochemistry tests including serum ALT of median relative reduction of at least 25% and GGT of at least 25% along with markers of reduced bile acid synthesis, may be achieved at 12 weeks.

[605] In some embodiments, the NASH patients may have advanced liver fibrosis (stage F3/F4). In some embodiments, such patients have stage F3 advanced liver fibrosis. In some embodiments, such patients have stage F4 liver fibrosis characterized by cirrhosis. In some embodiments, the NASH patients develop or at risk of developing hepatocellular carcinoma and/or esophageal varices.

[606] Fibrosis staging in non-alcoholic fatty liver disease according to the classification derived by the Nonalcoholic Steatohepatitis Clinical Research Network Pathology Committee is provided below:

Stages of fibrosis

Fibrotic manifestation	Fibrosis Stage
Perisinusoidal or periportal fibrosis	1
Mild perisinusoidal fibrosis (zone 3)	1A
Moderate perisinusoidal fibrosis (zone 3)	1B
Portal/periportal fibrosis	1C
Perisinusoidal and portal/periportal fibrosis	2
Bridging fibrosis	3
Cirrhosis	4

[607] Therapeutic benefits may be also evaluated by burden of disease and patient-reported outcomes. NASH also has an impact on quality of life for those living with the condition, measured through patient-reported outcomes (PROs). PROs may be assessed using tools such as the Chronic Liver Disease Questionnaire (CLDQ-NASH) prior to treatment (*e.g.*, baseline), particularly those related to physical health-related scores,

[608] Treatment with a TGF β 1 inhibitor (such as LTBP1/3 complex-selective inhibitors described herein) either alone (*e.g.*, monotherapy) or in combination with another therapy, may be effective to improve the PROs as compared to the baseline, or as compared to those of population norms. In some embodiments, diabetes mellitus may be associated with impairment in PROs including physical functioning, bodily pain, general health and vitality. Treatment with a TGF β 1 inhibitor (such as LTBP1/3 complex-selective inhibitors described herein) either alone (*e.g.*, monotherapy) or in combination with another therapy, may be effective to improve physical health-related scores (such as PROs) among subjects with diabetes (*e.g.*, type 2 diabetes) and/or obesity.

[609] Published studies in the literature suggest that regulatory T cells (Tregs) may play a role in the progression of liver disease into later-stage fibrosis with greater severity. For example, Zhang et al. reported that persistence of liver cirrhosis is maintained by intrahepatic regulatory T cells that inhibit the process of fibrosis resolution (Transl Res. 2016; 169: 67-79.e1-2). Kobayashi et al. suggested that Tregs are involved in the progression of liver fibrosis into hepatocellular carcinoma (HCC), a process referred to as hepatocarcinogenesis (Clin Cancer Res. 2007; 13(3): 902-911).

[610] Accordingly, it is contemplated that careful selection of suitable TGF β inhibitor tailored to the disease type and stage of the disease progression should be considered to maximize therapeutic benefit to a particular patient or patient population.

[611] In some embodiments, a TGF β 1-selective, context-selective inhibitor that targets matrix-associated TGF β 1 (*e.g.*, LTBP1-proTGF β 1 and/or LTBP3-proTGF β 1), such as those disclosed herein, is selected for use in the treatment of an early-stage liver disease such as nonalcoholic fatty liver ("NAFL") and noncirrhotic liver fibrosis associated with NASH. The noncirrhotic liver fibrosis includes liver fibrosis of stages 1-3. The TGF β 1-selective, context-selective inhibitor is administered to the subject in an amount effective to treat the disease, *e.g.*, slow the progress of, halt, or reverse NAFL and noncirrhotic NASH. Preferably, the effective amount is sufficient to prevent progression to cirrhosis and cirrhosis complications, reduce the need for liver transplantation, and/or improve survival. In some embodiments, efficacy may be shown by, for example, reduction of inflammatory changes, improvement in fibrosis, or both. In some embodiments, the subject has a metabolic condition, such as obesity, type 2 diabetes. The subject with noncirrhotic NASH may include those with a NASH activity score (NAS) greater than or equal to 4 with at least 1 point each in inflammation and ballooning along with a NASH Clinical Research Network (CRN) fibrosis score

greater than stage 1 fibrosis but less than stage 4 fibrosis. In some embodiments, the treatment achieves resolution of steatohepatitis on overall histopathological reading and no worsening of liver fibrosis on NASH CRN fibrosis score. Resolution of steatohepatitis is defined as absent fatty liver disease or isolated or simple steatosis without steatohepatitis and a NAS score of 0-1 for inflammation, 0 for ballooning, and any value for steatosis; or, improvement in liver fibrosis greater than or equal to one stage (NASH CRN fibrosis score) and no worsening of steatohepatitis (defined as no increase in NAS for ballooning, inflammation or steatosis); or, both resolution of steatohepatitis and improvement in fibrosis as defined above.

[612] In some embodiments, a TGF β 1-selective, context-selective inhibitor that targets matrix-associated TGF β 1 (*e.g.*, LTBP1-proTGF β 1 and/or LTBP3-proTGF β 1), such as those disclosed herein, is selected for use in the treatment of NASH with compensated cirrhosis. NASH-associated compensated cirrhosis is characterized by significant scar formation that is evident by histopathology, with hepatocytes clustered in nodules surrounded by dense extracellular matrix. The TGF β 1-selective, context-selective inhibitor is administered to the subject in an amount effective to halt or slow progression of fibrosis, prevent clinical decompensation, reduce the need for liver transplantation, and/or improve survival.

[613] NASH with decompensated cirrhosis may be characterized by one or more of the following criteria: portal hypertension (evidence of portal hypertension may include low platelet counts, esophageal varices, ascites, history of hepatic encephalopathy, splenomegaly); elevated bilirubin; or elevated international normalized ratio or prolonged prothrombin time. Thus, patients having NASH with compensated cirrhosis may be those not meeting one or more of the aforementioned decompensated cirrhosis criteria.

[614] In some embodiments, a TGF β 1-selective, context-selective inhibitor that targets matrix-associated TGF β 1 (*e.g.*, LTBP1-proTGF β 1 and/or LTBP3-proTGF β 1), such as those disclosed herein, is selected for use in the treatment of NASH with decompensated cirrhosis or HCC associated with liver fibrosis. In some embodiments, upon progression of the disease into a late-stage or end-stage liver disease characterized by manifestation of cirrhosis or HCC, the TGF β 1-selective, context-selective inhibitor is replaced with a TGF β 1-selective, context-independent inhibitor capable of targeting both matrix-associated and immune cell-associated TGF β 1, *e.g.*, LTBP1-proTGF β 1, LTBP3-proTGF β 1, GARP-proTGF β 1 and LRRC33-proTGF β 1 (see, for example, WO 2020/014473) in an amount effective to treat liver cirrhosis or HCC. In some embodiments, the TGF β 1-selective, context-selective inhibitor and/or the TGF β 1-selective, context-independent inhibitor may be used as monotherapy or in conjunction with one or more additional therapy.

[615] In some embodiments, an effective amount of the inhibitor may normalize, as compared to control, the levels of multiple inflammatory or fibrotic serum biomarkers as assessed following the

start of the therapy, at, for example, 12-36 weeks. In some embodiments, inflammatory or fibrotic biomarkers may be used to assess severity of NAFLD (by measure levels of hepatic steatosis), select patients for treatment, and/or monitor disease progression or treatment response. For example, blood biomarkers and panels may include, but are not limited to:

- i) the Fatty liver index (BMI, waist circumference, serum triglycerides, and gamma-glutamyltransferase (GGT);
- ii) the Hepatic steatosis index (serum aspartate aminotransferase (AST):alanine aminotransferase (ALT) ratio, BMI, gender, and presence of diabetes mellitus);
- i) the NAFLD liver fat score (serum ALT, HDL cholesterol, triglycerides, haemoglobin A_{1c} and leukocyte count);
- ii) the SteatoTest (BioPredictive) (serum levels of total bilirubin, GGT, α 2-macroglobin, haptoglobin, ALT, apolipoprotein AI, total cholesterol, triglycerides, glucose (adjusted for age and gender) and BMI); and
- iii) the NAFLD ridge score (serum levels of ALT, HDL cholesterol, triglycerides, haemoglobin A_{1c}, leukocyte count, and comorbidity data (and the presence of hypertension)).

[616] In some embodiments, imaging biomarkers can be used to assess levels of hepatic steatosis. For example, imaging biomarkers may include but are not limited to: ultrasonography, controlled attenuation parameter (CAP), MRI-estimated proton density fat fraction (MRI-PDFF), and magnetic resonance spectroscopy (MRS).

[617] Liver biopsies are the current standard for diagnosis NASH, however, variability among pathologists limits the effectiveness of such diagnostic method. Accordingly, use of the Fatty Liver Inhibition of Progression (FLIP) algorithm (comprising histological steatosis, activity and fibrosis scores) may be used to improve the consistency of NASH diagnosis by biopsy. Moreover, many noninvasive biomarkers may also be useful for diagnosing and monitoring disease. Accordingly, in some embodiments, inflammatory or fibrotic biomarkers may be used to assess severity of NASH, select patients for treatment, and/or monitor disease progression or treatment response. Blood biomarkers may include:

- i) apoptosis markers, such as CK18 fragments, total cytokeratin and sFAS;
- ii) inflammatory markers, such as CRP, TNF, IL-8, and CXCL10;
- iii) lipid oxidation products, such as 11-HETE, 9-HODE, 13-HODE, 12-oxo-ODE, LA-13-HODE (oxNASHscore), and 11,12-diHETrE;
- iv) lysosomal enzymes, such as cathepsin D; and

- v) combination panels, such as NASHTest (BioPredictive) and NASH Diagnostics Panel (comprising, presence of diabetes mellitus, sex, BMI, and serum levels of triglyceride, CK18 fragments, and total CK18).

[618] In some embodiments, biomarkers and related panels may be useful in diagnosis levels of fibrosis and/or cirrhosis, select patients for treatment, and/or monitor disease progression or treatment response. For example, noninvasive tests of liver fibrosis and cirrhosis include, but are not limited to: AST:ALT ratio, AST:platelet ratio index, fibrosis-4 index (age, AST, ALT, and platelet count), NAFLD fibrosis score (age, BMI, impaired fasting glucose and/or diabetes, AST ALT, platelet count, and albumin), BARD score (AST, ALT, BMI, and diabetes).

[619] Specific fibrosis markers and panels may also be useful, and include, but are not limited to: hyaluronic acid; PIIPNP; Pro-C3; TIMP1; Laminin; enhanced liver fibrosis (ELF) panel (PIIPNP, hyaluronic acid, TIMP1); FibroTest (GGT, total bilirubin, α_2 m, apolipoprotein AI, and haptoglobin); and FibroMeter NAFLD (body weight, prothrombin index, ALT, AST, ferritin, and fasting glucose). Imaging biomarkers for liver fibrosis may include, but are not limited to: FibroScan (TE), point shear wave elastography (pSWE) (aka acoustic radiation force impulse (ARFI)), 2D-3D SWE, magnetic resonance elastography (MRE), and multiparametric MRI.

[620] Any RGFb-related disease with an inflammatory or auto inflammatory element may benefit from the novel inhibitors of the present disclosure which spare the regulatory T cell function. Particularly in liver diseases, *e.g.*, metabolic liver conditions, it may be advantageous to select a TGFb inhibitor that selectively target the matrix-associated TGFb1 signaling.

[621] In one embodiment, the methods and compositions for use as described herein are useful for treating a subject having primary biliary cholangitis (PBC). In one embodiment, the subject having PBC has been nonresponsive to UDCA (ursodeoxycholic acid) treatment. In one embodiment, the subject has Barcelona, Paris-I, Toronto, Rotterdam, or Paris-II insufficient response to UDCA. In one embodiment, the subject having PBC has $ALP \geq 2 \times ULN$ (upper limit of normal), and bilirubin $> 1 \times ULN$ despite an at least 1 year therapy with UDCA at the standard recommended dose (10-15 mg/kg b.w./day).

[622] In another embodiment, the methods and compositions for use as described herein are useful for treating a subject having primary sclerosing cholangitis (PSC). In one embodiment, the subject having PSC has an elevated ALP, an abnormal cholangiography, endoscopic retrograde cholangiopancreatography, or percutaneous transhepatic cholangiography. In one embodiment, the subject having PSC has a model for end-stage liver disease (MELD) score of at least 14.

[623] In another embodiment, the methods and compositions for use as described herein are useful for treating a subject having NASH. In one embodiment, a subject has fibrosis stage 2 NASH, fibrosis stage 3 NASH, or fibrosis stage 4 NASH. In one embodiment, a subject has a NAS (NAFLD

activity score) of at least 4, or of at least 5. In one embodiment, a subject with NASH has a NAS>4 and fibrosis stage 2 or stage 3. In one embodiment, a subject with NASH has a NAS>5 and fibrosis stage 2 or stage 3. In one embodiment, a subject with NASH has a model for end-stage liver disease (MELD) score of at least 14.

[624] The isoform-specific TGF β 1 inhibitors such as those provided herein may be used to treat fibrotic conditions of the kidney, *e.g.*, diseases characterized by extracellular matrix accumulation (IgA nephropathy, focal and segmental glomerulosclerosis, crescentic glomerulonephritis, lupus nephritis and diabetic nephropathy) in which significantly increased expression of TGF β in glomeruli and the tubulointerstitium has been observed. While glomerular and tubulointerstitial deposition of two matrix components induced by TGF β , fibronectin EDA+ and PAI-1, was significantly elevated in all diseases with matrix accumulation, correlation analysis has revealed a close relationship primarily with the TGF β 1 isoform. Accordingly, the isoform-specific TGF β 1 inhibitors are useful as therapeutic for a spectrum of human glomerular disorders, in which TGF β is associated with pathological accumulation of extracellular matrix.

[625] In some embodiments, the fibrotic condition of the kidney is associated with chronic kidney disease (CKD). CKD is caused primarily by high blood pressure or diabetes and claims more than one million lives each year. CKD patients require lifetime medical care that ranges from strict diets and medications to dialysis and transplants. In some embodiments, the TGF β 1 inhibitor therapy described herein may reduce or delay the need for dialysis and/or transplantation. In some embodiments, such therapy may reduce the need (*e.g.*, dosage, frequency) for other treatments. In some embodiments, the isoform-specific TGF β 1 inhibitors may be administered in patients who receive one or more additional therapies, including, but are not limited to myostatin inhibitors, which may generally enhance metabolic regulation in patients with CKD.

[626] Fibrotic conditions that may be treated with the TGF β 1 inhibitor of the present disclosure include conditions involving fibrosis and/or chronic inflammation. Such conditions may be neuromuscular disorders, including but are not limited to Duchenne muscular dystrophy (DMD), and other genetic disorders such as multiple sclerosis (MS) and cystic fibrosis (CF). Through the inhibition of both the ECM- and immune cell-associated TGF β 1 arms, the TGF β 1 inhibitor such as those described herein is thought to suppress fibrotic progression and restore M1/M2 macrophage polarization.

[627] The organ fibrosis which may be treated with the methods provided herein includes cardiac (*e.g.*, cardiovascular) fibrosis. In some embodiments, the cardiac fibrosis is associated with heart failure, *e.g.*, chronic heart failure (CHF). In some embodiments, the heart failure may be associated with myocardial diseases and/or metabolic diseases. In some embodiments, the isoform-specific, TGF β 1 inhibitors may be administered in patients who receive one or more additional therapies,

including, but are not limited to myostatin inhibitors in patients with cardiac dysfunction that involves heart fibrosis and metabolic disorder.

[628] In some embodiments, fibrotic conditions that may be treated with the compositions and/or methods described herein include desmoplasia. Desmoplasia may occur around a neoplasm, causing dense fibrosis around the tumor (*e.g.*, desmoplastic stroma), or scar tissue within the abdomen after abdominal surgery. In some embodiments, desmoplasia is associated with malignant tumor. Due to its dense formation surrounding the malignancy, conventional anti-cancer therapeutics (*e.g.*, chemotherapy) may not effectively penetrate to reach cancerous cells for clinical effects. Isoform-specific, inhibitors of TGF β 1 such as those described herein may be used to disrupt the desmoplasia, such that the fibrotic formation can be loosened to aid effects of anti-cancer therapy. In some embodiments, the isoform-specific inhibitors of TGF β 1 can be used as monotherapy (more below).

[629] In some embodiments, a patient has a fibrotic solid tumor (*e.g.*, desmoplasia) and is or has been excluded from a surgical candidate pool, such that the fibrotic solid tumor is considered to be non-resectable or non-operable (*e.g.*, risk of surgical intervention outweighs potential benefit thereof). Such patient may be a candidate for receiving a TGF β 1 inhibition therapy of the present disclosure. The TGF β 1 inhibition therapy of the present invention administered to such patients may render the tumor become resectable or operable so that the patient may become a candidate for surgical resection.

[630] To treat patients with fibrotic conditions, TGF β 1 isoform-specific inhibitors are administered to a subject in an amount effective to treat the fibrosis. The effective amount of such an antibody is an amount effective to achieve both therapeutic efficacy and clinical safety in the subject. In some embodiments, the inhibitor is an antibody that can block activation of an LTBP-mediated TGF β 1 localized (*e.g.*, tethered) in the ECM. In some embodiments, the LTBP is LTBP1 and/or LTBP3. In some embodiments, a LTBP-specific inhibitor of TGF β 1 can be combined with an inhibitor of LRRC33-proTGF β .

[631] Assays useful in determining the efficacy of the antibodies and/or compositions of the present disclosure for the alteration of fibrosis include, but are not limited to, histological assays for counting fibroblasts and basic immunohistochemical analyses known in the art.

[632] In some embodiments, circulating LAP fragment(s) may be used as a serum marker of fibrogenesis. See for example, US patent 8,198,412, the contents of which are incorporated herein by reference.

[633] There are many animal models that have been developed to study fibrosis. For example, certain high fat diets in mice has been shown to mimic both the histopathology and pathogenesis of human NAFLD. Moreover, some genetic models also display features of human metabolic syndrome and NAFLD, such as *db/db* and *ob/ob* mouse models. There are also animal models for the study of

NASH, which mainly consist of various diet-induced models, including, but not limited to, methionine and choline-deficient diet (MCD), high-cholesterol diet (HCD), choline-deficient high fat diet (CDHFD), choline-deficient L-amino acid-deficient diet, choline-deficient L-amino acid-deficient diet + carbon tetrachloride, high-fat diet + streptozotocin, high fat + high cholesterol diet (HFHC), high-fructose diet (HFD), and high-fructose high fat diet (HFHF). Genetic mouse models for the study of NASH include, but are not limited to *foz/foz* mice, Hepatocyte-specific PTEN-deficient mice, *Db/db* mice + diethylnitrosamine (DEN), and *db/db* mice + MCD. The details of all of these models, including the pluses and minus of each, are outlined in Jennie Ka Ching Lau et al., *J Pathol* 2017; **241**: 36–44; the contents of which are incorporated herein by reference.

[634] Other models useful for testing the efficacy of TGF β inhibitors in fibrosis include the carbon tetrachloride (CCL₄)-induced liver fibrosis model and adenine-induced kidney fibrosis model. Another model useful for testing the efficacy of isoform-specific TGF β inhibitors in liver fibrosis include the bile duct ligation (BDL) model (see, *e.g.*, Tag et al., *J Vis Exp.* 2015; (96): 52438). A useful genetic model of kidney fibrosis includes Alport model (discussed elsewhere herein).

[635] In any of such preclinical models, efficacy in fibrosis (*e.g.*, pro-fibrotic or anti-fibrotic effects *in vivo*) may be assessed by any suitable methods, such as: percent of picosirius red (PSR)-positive area in tissue sections; contents (quantification) of hydroxyproline in tissue; and immunohistochemical detection and quantification of col1A staining on tissue sections. In some embodiments, histopathological appraisal may be carried out, in which fibrosis scoring system can be employed. Pharmacological effects of test articles (*e.g.*, TGF β inhibitors) may be examined by suitable pharmacodynamics (PD) measures, such as measuring downstream signal transduction events. In case of the TGF β pathway, for example, suitable PD measure includes relative phosphorylation of SMAD2/3.

Muscle Conditions Associated with Fibrosis

[636] Accumulating evidence indicates that TGF β plays an important role in muscle homeostasis, repair, and regeneration. Agents, such as monoclonal antibodies described herein, that selectively modulate LTBP-associated TGF β signaling may be effective for treating damaged muscle fibers, such as in chronic/genetic muscular dystrophies, congenital fibrosis of ocular/extraocular muscles, and acute muscle injuries, without the toxicities associated with more broadly-acting TGF β inhibitors.

[637] Accordingly, the present invention provides methods for treating damaged muscle fibers using an agent that preferentially modulates a subset, but not all, of TGF β effects *in vivo*. Such agents can selectively modulate TGF β 1 signaling (“isoform-specific modulation”) in a particular context, *i.e.*, when presented by LTBP1 or LTBP3.

[638] In skeletal muscle, TGF β plays a variety of roles including inhibition of proliferation and differentiation, induction of atrophy, and development of fibrosis. TGF β reduces satellite cell

proliferation and prevents differentiation (via inhibition of MyoD and myogenin) (Allen, R.E. and L.K. J Cell Physiol, 1987. 133(3): p. 567-72; Brennan, T.J., *et al.*, Proc Natl Acad Sci U S A, 1991. 88(9): p. 3822-6; Massague, J., *et al.*, Proc Natl Acad Sci U S A, 1986. 83(21): p. 8206-10; Olson, E.N., *et al.*, J Cell Biol, 1986. 103(5): p. 1799-805). The isoform of TGF β (*i.e.*, TGF β 1, 2, or 3) is not specified in these early papers, but is presumed to be TGF β 1. TGF β also contributes to muscle fibrosis; direct injection of recombinant TGF β 1 results in skeletal muscle fibrosis, and pan-TGF β inhibition decreases fibrosis in acute and chronically injured muscle (Li, Y., *et al.*, Am J Pathol, 2004. 164(3): p. 1007-19; Mendias, C.L., *et al.*, Muscle Nerve, 2012. 45(1): p. 55-9; Nelson, C.A., *et al.*, Am J Pathol, 2011. 178(6): p. 2611-21). TGF β 1 is expressed by myofibers, macrophages, regulatory T cells, fibroblasts, and fibrocytes within the skeletal muscle (Li, Y., *et al.*, Am J Pathol, 2004. 164(3): p. 1007-19; Lemos, D.R., *et al.*, Nat Med, 2015. 21(7): p. 786-94; Villalta, S.A., *et al.*, Sci Transl Med, 2014. 6(258): p. 258ra142; Wang, X., *et al.*, J Immunol, 2016. 197(12): p. 4750-4761); and expression is increased upon injury and in disease (Li, Y., *et al.*, Am J Pathol, 2004. 164(3): p. 1007-19; Nelson, C.A., *et al.*, Am J Pathol, 2011. 178(6): p. 2611-21; Bernasconi, P., *et al.*, J Clin Invest, 1995. 96(2): p. 1137-44; Ishitobi, M., *et al.*, Neuroreport, 2000. 11(18): p. 4033-5). TGF β 2 and TGF β 3 are also upregulated (at the mRNA level) in mdx muscle (a mouse model of Duchenne muscular dystrophy), although to a lesser extent than TGF β 1 (Nelson, C.A., *et al.*, Am J Pathol, 2011. 178(6): p. 2611-21; Zhou, L., *et al.*, Neuromuscul Disord, 2006. 16(1): p. 32-8). Pessina, *et al.*, recently used lineage tracing experiments to show that cells of multiple origins within dystrophic muscle adopt a fibrogenic fate via a TGF β -dependent pathway (Pessina, P., *et al.*, Stem Cell Reports, 2015. 4(6): p. 1046-60).

[639] TGF β 1 has been implicated in human muscular dystrophies. Duchenne muscular dystrophy (DMD) is a severe, progressive, and ultimately fatal disease caused by the absence of dystrophin (Bushby, K., *et al.*, Lancet Neurol, 2010. 9(1): p. 77-93). Lack of dystrophin results in increased susceptibility to contraction-induced injury, leading to continual muscle degeneration (Petrof, B.J., *et al.*, Proc Natl Acad Sci U S A, 1993. 90(8): p. 3710-4; Dellorusso, C., *et al.*, J Muscle Res Cell Motil, 2001. 22(5): p. 467-75; Pratt, S.J., *et al.*, Cell Mol Life Sci, 2015. 72(1): p. 153-64). Repeated rounds of repair contribute to chronic inflammation, fibrosis, exhaustion of the satellite cell pool, eventual loss of mobility and death (Bushby, K., *et al.*, Lancet Neurol, 2010. 9(1): p. 77-93; McDonald, C.M., *et al.*, Muscle Nerve, 2013. 48(3): p. 343-56). Expression of TGF β 1 is significantly increased in patients with DMD and correlates with the extent of fibrosis observed in these patients (Bernasconi, P., *et al.*, J Clin Invest, 1995. 96(2): p. 1137-44; Chen, Y.W., *et al.*, Neurology, 2005. 65(6): p. 826-34). Excessive ECM deposition has detrimental effects on the contractile properties of the muscle and can limit access to nutrition as the myofibers are isolated from their blood supply (Klingler, W., *et al.*, Acta Myol, 2012. 31(3): p. 184-95). Recently, additional data has further implicated TGF β 1 in muscular dystrophies. Variants in LTBP4 have been found to modify disease severity in mouse and

human. In mouse, a variant of LTBP4 is protective in mice lacking dystrophin or γ -sarcoglycan (Coley, W.D., *et al.*, Hum Mol Genet, 2016. 25(1): p. 130-45; Heydemann, A., *et al.*, J Clin Invest, 2009. 119(12): p. 3703-12). In humans, two groups independently identified a variant of LTBP4 as protective in DMD, delaying loss of ambulation by several years (Flanigan, K.M., *et al.*, Ann Neurol, 2013. 73(4): p. 481-8; van den Bergen, J.C., *et al.*, J Neurol Neurosurg Psychiatry, 2015. 86(10): p. 1060-5).

[640] Although the nature of the genetic variants in mouse and human differs, in both species the protective variant results in decreased TGF β signaling (Heydemann, A., *et al.*, J Clin Invest, 2009. 119(12): p. 3703-12); Ceco, E., *et al.*, Sci Transl Med, 2014. 6(259): p. 259ra144). Many of the functions of TGF β 1 in skeletal muscle biology have been inferred from experiments in which purified active growth factor is injected into animals or added to cells in culture (Massague, J., *et al.*, Proc Natl Acad Sci U S A, 1986. 83(21): p. 8206-10; Li, Y., *et al.*, Am J Pathol, 2004. 164(3): p. 1007-19; Mendias, C.L., *et al.*, Muscle Nerve, 2012. 45(1): p. 55-9). Given the importance of cellular context for specific functions of TGF β 1 (see, for example, Hinck *et al.*, Cold Spring Harb. Perspect. Biol, 2016. 8(12)) it is possible that some of the effects observed in these experiments do not reflect the endogenous role(s) of the cytokine *in vivo*. For example, treatment of human dermal fibroblasts with recombinant TGF β 1, myostatin, or GDF11 results in nearly identical changes in gene expression in these cells, although *in vivo* the roles of these proteins are quite different (Tanner, J.W., Khalil, A., Hill, J., Franti, M., MacDonnell, S.M., Growth Differentiation Factor 11 Potentiates Myofibroblast Activation, in Fibrosis: From Basic Mechanisms to Targeted therapies. 2016: Keystone, CO).

[641] Multiple investigators have used inhibitors of TGF β to clarify the role of the growth factor *in vivo*. Treatment of mdx mice with the pan-TGF β neutralizing antibody 1D11 clearly results in reduced fibrosis (by histology and hydroxyproline content), reduced muscle damage (reduced serum creatine kinase and greater myofiber density), and improved muscle function (by plethysmography, force generation of isolated EDL muscles, and increased forelimb grip strength) (Nelson, C.A., *et al.*, Am J Pathol, 2011. 178(6): p. 2611-21; Andreetta, F., *et al.*, J Neuroimmunol, 2006. 175(1-2): p. 77-86; Gumucio, J.P., *et al.*, J Appl Physiol (1985), 2013. 115(4): p. 539-45). In addition, myofiber-specific expression of a dominant negative TGF β type II receptor protects against muscle damage after cardiotoxin injury and in δ -sarcoglycan-/- mice (Accornero, F., *et al.*, Hum Mol Genet, 2014. 23(25): p. 6903-15). The proteoglycan decorin, which is abundant in skeletal muscle and inhibits TGF β activity, decreases muscle fibrosis in mdx mice and following laceration injury (Li, Y., *et al.*, Mol Ther, 2007. 15(9): p. 1616-22; Gosselin, L.E., *et al.*, Muscle Nerve, 2004. 30(5): p. 645-53). Other molecules with TGF β inhibitory activity, such as suramin (an anti-neoplastic agent) and losartan (an angiotensin receptor blocker) have been effective in improving muscle pathology and reducing fibrosis in mouse models of injury, Marfan's syndrome, and muscular dystrophy (Spurney, C.F., *et al.*, J Cardiovasc Pharmacol Ther, 2011. 16(1): p. 87-95; Taniguti, A.P., *et al.*, Muscle Nerve,

2011. 43(1): p. 82-7; Bedair, H.S., *et al.*, Am J Sports Med, 2008. 36(8): p. 1548-54; Cohn, R.D., *et al.*, Nat Med, 2007. 13(2): p. 204-10). While all of the therapeutic agents described above do inhibit TGF β 1 or its signaling, none of them is specific for the TGF β 1 isoform. For example, 1D11 binds to and inhibits the TGF β 1, 2, and 3 isoforms (Dasch, J.R., *et al.*, J Immunol, 1989. 142(5): p. 1536-41). Suramin inhibits the ability of multiple growth factors to bind to their receptors, including PDGF, FGF, and EGF, in addition to TGF β 1 (Hosang, M., J Cell Biochem, 1985. 29(3): p. 265-73; Olivier, S., *et al.*, Eur J Cancer, 1990. 26(8): p. 867-71; Scher, H.I. and W.D. Heston, Cancer Treat Res, 1992. 59: p. 131-51). Decorin also inhibits myostatin activity, both by direct binding and through upregulation of follistatin, a myostatin inhibitor (Miura, T., *et al.*, Biochem Biophys Res Commun, 2006. 340(2): p. 675-80; Brandan, E., C. Cabello-Verrugio, and C. Vial, Matrix Biol, 2008. 27(8): p. 700-8; Zhu, J., *et al.*, J Biol Chem, 2007. 282(35): p. 25852-63). Losartan affects additional signaling pathways through its effects on the renin-angiotensin-aldosterone system, including the IGF-1/AKT/mTOR pathway (Burks, T.N., *et al.*, Sci Transl Med, 2011. 3(82): p. 82ra37; Sabharwal, R. and M.W. Chapleau, Exp Physiol, 2014. 99(4): p. 627-31; McIntyre, M., *et al.*, Pharmacol Ther, 1997. 74(2): p. 181-94). Therefore, all of these therapies inhibit additional molecules which may contribute to their therapeutic effects, as well as toxicities.

[642] Apart from chronic inflammation, the hallmark of DMD is excessive, and progressive, fibrosis. In advanced disease the fibrosis is so severe that it can actually isolate individual muscle fibers from their blood supply. It also alters the contractile properties of the muscle. In human patients, there is a strong correlation between the extent of TGF β 1 upregulation and fibrosis, and a strong link between the extent of fibrosis and negative mobility outcomes. Therefore, in some embodiments, LTBP-proTGF β 1 inhibitors may be administered to dystrophic patients for the prevention and/or reduction of fibrosis to selectively target the ECM-associated TGF β 1 effects in the disease. In some embodiments, various isoform- and/or context-selective agents described herein can be employed to achieve inhibition of TGF β 1 signaling to prevent fibrosis and promote myogenesis, but without having unwanted effects on the immune system (*e.g.*, through GARP or LRRC33).

Administration

[643] To practice the method disclosed herein, an effective amount of the pharmaceutical composition described above can be administered to a subject (*e.g.*, a human) in need of the treatment via a suitable route, such as intravenous administration, *e.g.*, as a bolus or by continuous infusion over a period of time, by intramuscular, intraperitoneal, intracerebrospinal, subcutaneous, intra-articular, intrasynovial, intrathecal, oral, inhalation or topical routes. When used in the treatment of fibrosis and/or metabolic conditions, preferably the TGF β inhibitor is administered subcutaneously. Commercially available nebulizers for liquid formulations, including jet nebulizers and ultrasonic nebulizers are useful for administration. Liquid formulations can be directly nebulized and lyophilized powder can be nebulized after reconstitution. Alternatively, inhibitors, *e.g.*, antibodies, or antigen-

binding portions thereof, that selectively bind a LTBP1-TGF β complex and/or a LTBP3-TGF β complex can be aerosolized using a fluorocarbon formulation and a metered dose inhaler, or inhaled as a lyophilized and milled powder.

[644] The subject to be treated by the methods described herein can be a mammal, more preferably a human. Mammals include, but are not limited to, farm animals, sport animals, pets, primates, horses, dogs, cats, mice and rats. A human subject who needs the treatment may be a human patient having, at risk for, or suspected of having a TGF β -related indication, such as those noted above. A subject having a TGF β -related indication can be identified by routine medical examination, *e.g.*, laboratory tests, organ functional tests, CT scans, or ultrasounds. A subject suspected of having any of such indication might show one or more symptoms of the indication. A subject at risk for the indication can be a subject having one or more of the risk factors for that indication.

[645] As used herein, the terms "effective amount" and "effective dose" refer to any amount or dose of a compound or composition that is sufficient to fulfill its intended purpose(s), *i.e.*, a desired biological or medicinal response in a tissue or subject at an acceptable benefit/risk ratio. For example, in certain embodiments of the present invention, the intended purpose may be to inhibit TGF β -1 activation *in vivo*, to achieve clinically meaningful outcome associated with the TGF β -1 inhibition. Effective amounts vary, as recognized by those skilled in the art, depending on the particular condition being treated, the severity of the condition, the individual patient parameters including age, physical condition, size, gender and weight, the duration of the treatment, the nature of concurrent therapy (if any), the specific route of administration and like factors within the knowledge and expertise of the health practitioner. These factors are well known to those of ordinary skill in the art and can be addressed with no more than routine experimentation. It is generally preferred that a maximum dose of the individual components or combinations thereof be used, that is, the highest safe dose according to sound medical judgment. It will be understood by those of ordinary skill in the art, however, that a patient may insist upon a lower dose or tolerable dose for medical reasons, psychological reasons or for virtually any other reasons.

[646] Empirical considerations, such as the half-life, generally will contribute to the determination of the dosage. For example, antibodies that are compatible with the human immune system, such as humanized antibodies or fully human antibodies, may be used to prolong half-life of the antibody and to prevent the antibody being attacked by the host's immune system. Frequency of administration may be determined and adjusted over the course of therapy, and is generally, but not necessarily, based on treatment and/or suppression and/or amelioration and/or delay of a TGF β -related indication. Alternatively, sustained continuous release formulations of an antibody that selectively binds a LTBP1-TGF β 1 complex and/or a LTBP3-TGF β 1 complex may be appropriate. Various formulations and devices for achieving sustained release would be apparent to the skilled artisan and are within the scope of this disclosure.

[647] In one example, dosages for an inhibitor, *e.g.*, antibody, that selectively binds a LTBP1-TGF β 1 complex and/or a LTBP3-TGF β 1 complex as described herein may be determined empirically in individuals who have been given one or more administration(s) of the inhibitor. Individuals are given incremental dosages of the inhibitor. To assess efficacy, an indicator of the TGF β -related indication can be followed. For example, methods for measuring for myofiber damage, myofiber repair, inflammation levels in muscle, and/or fibrosis levels in muscle are well known to one of ordinary skill in the art.

[648] The present invention encompasses the recognition that agents capable of modulating the activation step of TGF β s in an isoform-specific manner, and a context-specific manner, may provide improved safety profiles when used as a medicament. Accordingly, the invention includes inhibitors, *e.g.*, antibodies and antigen-binding fragments thereof, that selectively bind and inhibit activation of TGF β 1, but not TGF β 2 or TGF β 3, thereby conferring specific inhibition of the TGF β 1 signaling *in vivo* while minimizing unwanted side effects from affecting TGF β 2 and/or TGF β 3 signaling. Likewise, the invention includes inhibitors, *e.g.*, antibodies and antigen-binding fragments thereof, that selectively inhibit activation of TGF β 1 presented by LTBP1 and/or LTBP3, but not TGF β 1 presented by GARP or LRRC33, thereby conferring specific inhibition of LTBP1/3-associated TGF β 1 signaling *in vivo* while minimizing unwanted side effects caused by modulation of GARP-associated TGF β 1 and/or LRRC33-associated TGF β 1.

[649] In some embodiments, the inhibitors, *e.g.*, antibodies, or antigen-binding portions thereof, as described herein, are not toxic when administered to a subject. In some embodiments, the inhibitors, *e.g.*, antibodies, or antigen-binding portions thereof, as described herein, exhibit reduced toxicity when administered to a subject as compared to an antibody that binds to both TGF β 1 and TGF β 2. In some embodiments, the inhibitors, *e.g.*, antibodies, or antigen-binding portions thereof, as described herein, exhibit reduced toxicity when administered to a subject as compared to an inhibitor that binds to both TGF β 1 and TGF β 3. In some embodiments, the inhibitors, *e.g.*, antibodies, or antigen-binding portions thereof, as described herein, exhibit reduced toxicity when administered to a subject as compared to an inhibitor that binds to TGF β 1, TGF β 2 and TGF β 3.

[650] Generally for administration of any of the inhibitors, *e.g.*, antibodies, described herein, an initial candidate dosage can be about 0.5-30 mg/kg per dose, *e.g.*, about 0.5 mg/kg, about 1 mg/kg, about 2 mg/kg, about 5 mg/kg, about 10 mg/kg, about 15 mg/kg, about 20 mg/kg, about 30 mg/kg per dose. Typically, the composition comprising an antibody or a fragment thereof encompassed by the present disclosure is administered to a human patient at the dosage at suitable intervals, such as once or twice weekly, every 1-8 weeks, etc. In some embodiments, frequency of administration may be adjusted to, for example, twice a week, once a week, every two weeks, every three weeks, every four weeks, every six weeks, every eight weeks, etc.. For the purpose of the present disclosure, a typical daily dosage might range from about any of 1 mg/kg to 20 mg/kg or more, depending on the factors

mentioned above. For repeated administrations over several days or longer, depending on the condition, the treatment is sustained until a desired suppression of symptoms occurs or until sufficient therapeutic levels are achieved to alleviate a TGF β -related indication, or a symptom thereof.

[651] In one embodiment, the antibody, or antigen-binding fragment thereof, is administered to the subject at a dosage of between 0.1 and 30 mg/kg, between 0.5 and 30 mg/kg, between 1 and 30 mg/kg, between 5 and 30 mg/kg, between 10 and 30 mg/kg, between 15 and 30 mg/kg, between 20 and 30 mg/kg, between 25 and 30 mg/kg, between 0.1 and 25 mg/kg, between 0.5 and 25 mg/kg, between 1 and 25 mg/kg, between 5 and 25 mg/kg, between 10 and 25 mg/kg, between 15 and 25 mg/kg, between 20 and 25 mg/kg, between 0.1 and 20 mg/kg, between 0.5 and 20 mg/kg, between 1 and 20 mg/kg, between 5.0 and 20 mg/kg, between 10 and 20 mg/kg, between 15 and 20 mg/kg, between 0.1 and 15 mg/kg, between 0.5 and 15 mg/kg, between 1 and 15 mg/kg, between 5 and 15 mg/kg, between 10 and 15 mg/kg, between 5.0 and 20 mg/kg, between 10 and 20 mg/kg, between 15 and 20 mg/kg, between 0.1 and 10 mg/kg, between 0.5 and 10 mg/kg, between 1 and 10 mg/kg, between 5 and 10 mg/kg, optionally, wherein the subject is administered the antibody, or antigen-binding portion thereof, twice a week, once a week, once every 2 weeks, once every 3 weeks, once a month, or every other month.

[652] An exemplary dosing regimen comprises administering an initial dose of about 2 mg/kg, followed by one or more maintenance doses. For example, an initial dose may be between about 2 and 30 mg/kg, for instance, once a week or twice a week. Thereafter, maintenance dose(s) may follow, for example, between about 0.1 and 20 mg/kg, for instance, once a week, every other week, once a month, etc. However, other dosage regimens may be useful, depending on the pattern of pharmacokinetic decay that the practitioner wishes to achieve. Pharmacokinetics experiments have shown that the serum concentration of an inhibitor, *e.g.*, antibody, disclosed herein (*e.g.*, SR-AB2) remains stable for at least 7 days after administration to a preclinical animal model (*e.g.*, a mouse model). Without wishing to be bound by any particular theory, this stability post-administration may be advantageous since the antibody may be administered less frequently while maintaining a clinically effective serum concentration in the subject to whom the antibody is administered (*e.g.*, a human subject). In some embodiments, dosing frequency is once every week, every 2 weeks, every 4 weeks, every 5 weeks, every 6 weeks, every 7 weeks, every 8 weeks, every 9 weeks, or every 10 weeks; or once every month, every 2 months, or every 3 months, or longer. The progress of this therapy is easily monitored by conventional techniques and assays. The dosing regimen (including the antibody used) can vary over time.

[653] According to some embodiments, serum concentrations of the LTBP context-selective TGF β 1 inhibitor that are therapeutically effective to treat a TGF β 1-related indication in accordance with the present disclosure may be at least about 10 μ g/mL, *e.g.*, between about 10 μ g/mL and 1.0 mg/mL. In some embodiments, effective amounts of the antibody as measured by serum concentrations are about

20-400 $\mu\text{g/mL}$. In some embodiments, effective amounts of the antibody as measured by serum concentrations are about 100-800 $\mu\text{g/mL}$. In some embodiments, effective amounts of the inhibitor as measured by serum concentrations are at least about 20 $\mu\text{g/mL}$, *e.g.*, at least about 50 $\mu\text{g/mL}$, 100 $\mu\text{g/mL}$, 150 $\mu\text{g/mL}$ or 200 $\mu\text{g/mL}$. In preferred embodiments, in non-human primates, there are no observed toxicities (for example: no cardiotoxicities, hyperplasia and inflammation, dental and gingival findings) associated with such inhibitor after maintaining serum concentration levels of about 2,000-3,000 $\mu\text{g/mL}$ for at least 4 weeks, *e.g.*, at least 4 weeks, preferably at least 8 weeks, more preferably at least 12 weeks. Therefore, about 10-100 fold therapeutic window may be achieved.

[654] In some embodiments, for an adult patient of normal weight, doses ranging from about 0.3 to 5.00 mg/kg may be administered. The particular dosage regimen, *e.g.*, dose, timing and repetition, will depend on the particular individual and that individual's medical history, as well as the properties of the individual agents (such as the half-life of the agent, and other relevant considerations).

[655] For the purpose of the present disclosure, the appropriate dosage of an inhibitor, *e.g.*, antibody or antigen-binding fragment thereof, that selectively binds a LTBP1-TGF β complex and/or a LTBP3-TGF β complex will depend on the specific antibody (or compositions thereof) employed, the type and severity of the indication, whether the antibody is administered for preventive or therapeutic purposes, previous therapy, the patient's clinical history and response to the inhibitor, and the discretion of the attending physician. In some embodiments, a clinician will administer an inhibitor, *e.g.*, antibody, that selectively binds a LTBP1-TGF β 1 complex and/or a LTBP3-TGF β 1 complex, until a dosage is reached that achieves the desired result. Administration of an inhibitor, *e.g.*, antibody, that selectively binds a LTBP1-TGF β 1 complex and/or a LTBP3-TGF β 1 complex can be continuous or intermittent, depending, for example, upon the recipient's physiological condition, whether the purpose of the administration is therapeutic or prophylactic, and other factors known to skilled practitioners. The administration of an inhibitor, *e.g.*, antibody, that selectively binds a LTBP1-TGF β 1 complex and/or a LTBP3-TGF β 1 complex may be essentially continuous over a preselected period of time or may be in a series of spaced dose, *e.g.*, either before, during, or after developing a TGF β -related indication.

[656] Based on the observation that inhibiting TGF β 3 can increase collagen deposition or accumulation in fibrosis, add-on therapy comprising a TGF β 1-selective inhibitor (such as the novel antibodies disclosed herein) may be considered for patients who are treated with a TGF β inhibitor with TGF β 3-inhibiting activity, *e.g.*, inhibitors of TGF β 1/2/3, TGF β 1/3 and TGF β 3. Examples of TGF β inhibitors with TGF β 3-inhibiting activity include but are not limited to: low molecular weight antagonists of TGF β receptors, *e.g.*, ALK5 antagonists, such as Galunisertib (LY2157299 monohydrate); monoclonal antibodies (such as neutralizing antibodies) that inhibit all three isoforms ("pan-inhibitor" antibodies) (see, for example, WO 2018/134681); monoclonal antibodies that preferentially inhibit two of the three isoforms (*e.g.*, TGF β 1/3 (for example WO 2006/116002); and engineered molecules (*e.g.*, fusion proteins) such as ligand traps (for example, WO 2018/029367; WO

2018/129331 and WO 2018/158727). In some embodiments, the ligand trap comprises the structure in accordance with the disclosure of WO/2018/15872. In some embodiments, the ligand trap comprises the structure in accordance with the disclosure of WO 2018/029367; WO 2018/129331. In some embodiments, the ligand trap is a construct known as M7824. In some embodiments, the ligand trap is a construct known as AVID200. In some embodiments, the neutralizing pan-TGF β antibody is GC1008 or a derivative thereof. In some embodiments, such antibody comprises the sequence in accordance with the disclosure of WO/2018/134681.

[657] In some embodiments, the antibody is a neutralizing antibody that specifically binds both TGF β 1 and TGF β 3. In some embodiments such antibody preferentially binds TGF β 1 over TGF β 3. For example, the antibody comprises the sequence in accordance with the disclosure of WO/2006/116002. In some embodiments, the antibody is 21D1.

[658] The add-on therapy is aimed to counter or overcome the pro-fibrotic effect of TGF β 3 inhibition in patients who have received or are receiving a TGF β inhibitor with TGF β 3 inhibitory activities. In some embodiments, the patient has a fibrotic disorder or is at risk of developing a fibrotic disorder. For example, the patient may suffer from a metabolic condition that is associated with higher risk of developing liver fibrosis. The metabolic conditions linked to such risk include obesity, type 2 diabetes and NASH. Accordingly, the invention includes a TGF β 1-selective inhibitor for use in an add-on therapy of a subject treated with a TGF β 3 inhibitor, in an amount sufficient to reduce pro-fibrotic effects of the TGF β 3 inhibitor. In some embodiments, the subject has fibrosis. In some embodiments, the subject has myelofibrosis. In some embodiments, the subject has advanced cancer, *e.g.*, metastatic or locally advanced tumor. In some embodiments, the TGF β 1-selective inhibitor is selected from Ab31, Ab34, Ab37, Ab38, Ab39, Ab40, Ab41, Ab42, Ab43, Ab44, Ab45, Ab62, Ab63, and Ab64 (optionally Ab42 or Ab63) a variant/derivative or antigen-binding fragment thereof thereof, or an engineered molecule comprising an antigen-binding fragment thereof. In some embodiments, the TGF β 1-selective inhibitor is Ab42, a variant/derivative or antigen-binding fragment thereof, or an engineered molecule comprising an antigen-binding fragment thereof. In preferred embodiments, the TGF β 1-selective inhibitor is Ab42 or an antigen-binding fragment thereof.

[659] Without being bound by theory, in some embodiments, sparing of TGF β inhibitors with anti-TGF β 3 activities may be especially useful for treating patients who are diagnosed with a type of cancer known to be highly metastatic, myelofibrotic, and/or those having or are at risk of developing a fibrotic condition. Accordingly, the disclosure herein includes a TGF β inhibitor for use in the treatment of cancer wherein the inhibitor does not inhibit TGF β 3 and wherein the patient has a metastatic cancer or myelofibrosis, or the patient has or is at risk of developing a fibrotic condition , wherein optionally the fibrotic condition is non-alcoholic steatohepatitis (NASH).

[660] As used herein, the term “treating” refers to the application or administration of a composition including one or more active agents to a subject, who has a TGF β -related indication, a symptom of the indication, or a predisposition toward the indication, with the purpose to cure, heal, alleviate, relieve, alter, remedy, ameliorate, improve, or affect the indication, the symptom of the indication, or the predisposition toward the indication.

[661] Alleviating a TGF β -related indication with an inhibitor, *e.g.*, antibody, that selectively binds a LTBP1-TGF β 1 complex and/or a LTBP3-TGF β 1 complex includes delaying the development or progression of the indication, or reducing indication's severity. Alleviating the indication does not necessarily require curative results. As used therein, “delaying” the development of an indication associated with a TGF β -related indication means to defer, hinder, slow, retard, stabilize, and/or postpone progression of the indication. This delay can be of varying lengths of time, depending on the history of the indication and/or individuals being treated. A method that “delays” or alleviates the development of an indication, or delays the onset of the indication, is a method that reduces probability of developing one or more symptoms of the indication in a given time frame and/or reduces extent of the symptoms in a given time frame, when compared to not using the method. Such comparisons are typically based on clinical studies, using a number of subjects sufficient to give a statistically significant result.

Selection of a TGF β Inhibitor for Treating a Fibrotic Disorder

[662] Inhibitors of TGF β include isoform-non-selective inhibitors and isoform-selective inhibitors, with the former being the majority of known TGF β inhibitors/antagonists. Among the isoform-non-selective inhibitors are pan-inhibitors (TGF β 1/2/3 inhibitors), TGF β 1/2 inhibitors and TGF β 1/3 inhibitors. Isoform-selective inhibitors include neutralizing antibodies that selectively bind one isoform and activation inhibitors that target latent proTGF β complexes in an isoform-selective manner. The class of activation inhibitors include context-independent and context-selective inhibitors. For example, isoform-specific, context-independent TGF β 1 inhibitors have also been described, which bind pro/latent TGF β 1 presented by LTBP1/3, GARP, or LRRC33, and inhibit the release of mature TGF β 1 from the presenting molecule complex (see, *e.g.*, WO 2017/156500, WO 2020/014473 and WO 2020/014460). The entire contents of each of the foregoing applications are incorporated herein by reference. Context-specific antibodies that selectively bind a GARP-TGF β 1 complex and inhibit activation of TGF β 1 presented in the context of GARP have recently been described in WO 2018/013939.

[663] The present invention includes specific inhibitors of ECM-associated TGF β 1 activation, *e.g.*, antibodies, and antigen-binding portions thereof, that selectively bind a LTBP1/3-TGF β 1 complex, and which inhibit activation of TGF β 1 presented in the context of LTBP1 or LTBP3. The present

disclosure further provides guidance as to both selection of a suitable TGF β inhibitor among those listed above, tailored to certain patient populations and related therapeutic regimen.

[664] The surprising observations demonstrated herein (see Example 17; FIG. 22) that showing that TGF β 3 inhibition may in fact aggravate ECM dysregulation are informative in a decision making process of selecting the right TGF β inhibitor. For example, for use in the treatment of a fibrotic condition, it may be advantageous to select a TGF β inhibitor that lacks inhibitory activities towards TGF β 3. The observed exacerbation of fibrosis (*e.g.*, profibrotic effects) in response to TGF β 3 inhibition raises the possibility that role of TGF β 3 expands beyond homeostasis. Even more importantly, this may be relevant not only to fibrotic conditions but also in other disease contexts. Ample evidence suggests that dysregulation of the ECM is found in a number of disease conditions, including fibrosis and cancer. Indeed, many of the key profibrotic genes are also recognized among markers of various cancers. These markers include, for example, *col1A1*, *col3A1*, *PAI-1*, *CCL2*, *ACTA2*, *FN-1*, *CTGF* and *TGFB1*. Therefore, the finding that blockade of TGF β 3 appears harmful in fibrosis (*e.g.*, having a profibrotic effect) may be applicable to a broader scope of conditions associated with ECM dysregulation.

[665] The present invention encompasses insights into selecting “the *right* TGF β inhibitor” for “the right patient” to treat a disease condition with certain criteria and/or clinical features. In one aspect, the present invention provides use of preferred TGF β 1 inhibitors suitable for a particular patient population with fibrotic conditions. Accordingly, the invention includes use of an LTBP1/LTBP3-proTGF β 1 inhibitor in the treatment of a fibrotic condition in a subject, wherein the subject benefits from immunosuppression. This is based on the notion that at least a subset of TGF β 1 activities involves immune regulation which is mediated by GARP-associated and/or LRRC33-associated TGF β 1. Thus, the invention includes the recognition that use of TGF β 1 inhibitors that also affect the immune aspect of TGF β 1 effects may be detrimental for treating patients with fibrotic conditions where immunostimulation may cause exacerbation of the disease. The invention therefore aims at least in part to provide means of selectively inhibiting TGF β 1 effects within the ECM context (*e.g.*, LTBP-associated) while sparing TGF β 1 effects associated with non-ECM contexts (*e.g.*, immune cells, leukocytes, etc. expressing GARP or LRRC33 on cell surface), so as to prevent unwanted immunostimulation. This approach may be particularly advantageous in early-stage fibrosis, such as noncirrhotic liver fibrosis.

[666] In some embodiments, patient populations who benefit from both: i) inhibition of TGF β 1 signaling, and, ii) immunosuppression, include those who suffer from a severe or late stage organ fibrosis and who are to receive an allograft organ transplant. The severe or late stage organ fibrosis may be associated with IPF, CKD, and/or NASH. Such patients may have already received other therapies for treating the fibrotic disease, yet which may have failed to sufficiently treat or manage the condition. Attending physicians may determine that remaining treatment options may include

allograft transplantation. Such patients may be placed in a wait list for an available organ for transplantation. Such patients may be treated with an immunosuppressant. A selective inhibitor of LTBP1/LTBP3-presented TGF β 1 activation, which does not inhibit GARP-presented TGF β 1 activation, can be used to treat such patients, without raising risk of triggering immunostimulation mediated by effector T cells. Similarly, following the transplantation, such patients may continue to receive the selective inhibitor of LTBP1/LTBP3-presented TGF β 1 activation to avoid risk of an organ rejection.

[667] In some embodiments, patient populations who benefit from both: i) inhibition of TGF β 1 signaling, and, ii) immunosuppression, include those who suffer from a fibrotic disorder and who have an inflammatory or autoimmune condition.

[668] In some embodiments, the patient or patient population has or is at risk of developing one or more autoimmune disorders, such as: Achalasia; Addison's disease; Adult Still's disease; Agammaglobulinemia; Alopecia areata; Amyloidosis; Ankylosing spondylitis; Anti-GBM/Anti-TBM nephritis; Antiphospholipid syndrome; Autoimmune angioedema; Autoimmune dysautonomia; Autoimmune encephalomyelitis; Autoimmune hepatitis; Autoimmune inner ear disease (AIED); Autoimmune myocarditis; Autoimmune oophoritis; Autoimmune orchitis; Autoimmune pancreatitis; Autoimmune retinopathy; Autoimmune urticaria; Axonal & neuronal neuropathy (AMAN); Baló disease; Behcet's disease; Benign mucosal pemphigoid; Bullous pemphigoid; Castleman disease (CD); Celiac disease; Chagas disease; Chronic inflammatory demyelinating polyneuropathy (CIPD); Chronic recurrent multifocal osteomyelitis (CRMO); Churg-Strauss Syndrome (CSS) or Eosinophilic Granulomatosis (EGPA); Cicatricial pemphigoid; Cogan's syndrome; Cold agglutinin disease; Congenital heart block; Coxsackie myocarditis; CREST syndrome; Crohn's disease; Dermatitis herpetiformis; Dermatomyositis; Devic's disease (neuromyelitis optica); Discoid lupus; Dressler's syndrome; Endometriosis; Eosinophilic esophagitis (EoE); Eosinophilic fasciitis; Erythema nodosum; Essential mixed cryoglobulinemia; Evans syndrome; Fibromyalgia; Fibrosing alveolitis; Giant cell arteritis (temporal arteritis); Giant cell myocarditis; Glomerulonephritis; Goodpasture's syndrome; Granulomatosis with Polyangiitis; Graves' disease; Guillain-Barre syndrome; Hashimoto's thyroiditis; Hemolytic anemia; Henoch-Schönlein purpura (HSP); Herpes gestationis or pemphigoid gestationis (PG); Hidradenitis Suppurativa (HS) (Acne Inversa); Hypogammaglobulinemia; IgA Nephropathy; IgG4-related sclerosing disease; Immune thrombocytopenic purpura (ITP); Inclusion body myositis (IBM); Interstitial cystitis (IC); Juvenile arthritis; Juvenile diabetes (Type 1 diabetes); Juvenile myositis (JM); Kawasaki disease; Lambert-Eaton syndrome; Leukocytoclastic vasculitis; Lichen planus; Lichen sclerosus; Ligneous conjunctivitis; Linear IgA disease (LAD); Lupus; Lyme disease chronic; Meniere's disease; Microscopic polyangiitis (MPA); Mixed connective tissue disease (MCTD); Mooren's ulcer; Mucha-Habermann disease; Multifocal Motor Neuropathy (MMN) or MMNCB; Multiple sclerosis; Myasthenia gravis; Myositis; Narcolepsy; Neonatal Lupus;

Neuromyelitis optica; Neutropenia; Ocular cicatricial pemphigoid; Optic neuritis; Palindromic rheumatism (PR); PANDAS; Paraneoplastic cerebellar degeneration (PCD); Paroxysmal nocturnal hemoglobinuria (PNH); Parry Romberg syndrome; Pars planitis (peripheral uveitis); Parsonage-Turner syndrome; Pemphigus; Peripheral neuropathy; Perivenous encephalomyelitis; Pernicious anemia (PA); POEMS syndrome; Polyarteritis nodosa; Polyglandular syndromes type I, II, III; Polymyalgia rheumatica; Polymyositis; Postmyocardial infarction syndrome; Postpericardiotomy syndrome; Primary biliary cirrhosis; Primary sclerosing cholangitis; Progesterone dermatitis; Psoriasis; Psoriatic arthritis; Pure red cell aplasia (PRCA); Pyoderma gangrenosum; Raynaud's phenomenon; Reactive Arthritis; Reflex sympathetic dystrophy; Relapsing polychondritis; Restless legs syndrome (RLS); Retroperitoneal fibrosis; Rheumatic fever; Rheumatoid arthritis; Sarcoidosis; Schmidt syndrome; Scleritis; Scleroderma; Sjögren's syndrome; Sperm & testicular autoimmunity; Stiff person syndrome (SPS); Subacute bacterial endocarditis (SBE); Susac's syndrome; Sympathetic ophthalmia (SO); Takayasu's arteritis; Temporal arteritis/Giant cell ; arteritisThrombocytopenic purpura (TTP); Tolosa-Hunt syndrome (THS); Transverse myelitis; Type 1 diabetes; Ulcerative colitis (UC); Undifferentiated connective tissue disease (UCTD); Uveitis; Vasculitis; Vitiligo; Vogt-Koyanagi-Harada Disease.

[669] In some embodiments, the inflammatory or autoimmune condition is associated with the fibrosis. Non-limiting examples of inflammatory or autoimmune conditions associated with fibrosis include muscular dystrophy, such as DMD.

[670] In other embodiments, where patient populations who benefit from both: i) inhibition of TGFβ1 signaling, and, ii) immunosuppression, include those who suffer from a fibrotic disease and who have an inflammatory or autoimmune condition that is not directly associated with the fibrosis, but rather a discrete disorder.

[671] Such inflammatory or autoimmune conditions, whether or not directly associated with the underlining fibrotic disease or separate condition(s), may be caused by or associated with imbalance of regulatory T cells (Treg) in human autoimmune diseases. For example, such disorders that are linked to Treg dysregulation include, but are not limited to: Juvenile idiopathic arthritis; Rheumatoid arthritis (RA); Spondyloarthritis; Psoriatic arthritis; HCV mixed cryoglobulinaemia; cryoglobulinaemia; Multiple sclerosis; Autoimmune liver disease; Systemic lupus erythematoses; Immune-mediated diabetes; Myasthenia gravis; Primary Sjögren syndrome; Kawasaki disease; and, Inflammatory bowel disease (IBD).

[672] Thus, LTBP1/3-sepective inhibitors of TGFβ1 signaling, such as those described herein, can be used to treat patients who suffer from a fibrotic condition and inflammatory or autoimmune condition such as one or more of the disorders listed above. The LTBP1/3-sepective inhibitors of

TGFβ1 signaling used accordingly can treat or alleviate TGFβ1-dependent fibrosis in the ECM, while sparing immune-associated TGFβ1 signaling.

[673] Accordingly, related methods of the invention include methods for selecting an appropriate TGFβ1 inhibitor for treating a fibrotic disorder, based on the clinical manifestations of the fibrotic disorder in a subject. In one embodiment, the invention provides a method of selecting an isoform-specific TGFβ1 inhibitor for treatment of a fibrotic disorder in a subject. The method comprises (a) determining whether the fibrotic disorder manifests clinical presentations including fibrosis and one or more of inflammation, immune suppression, proliferative dysregulation, and need for an allograft transplant, and (b) selecting an isoform-specific, context-dependent TGFβ1 inhibitor or an isoform-specific, context-independent TGFβ1 inhibitor for treatment of the fibrotic disorder based on the clinical presentations determined in step (a). In another embodiment, the invention provides a method of treating a subject having a fibrotic disorder, comprising selecting a treatment regimen including an isoform-specific TGFβ1 inhibitor for the subject, and administering the selected treatment regimen to the subject, wherein the selection comprises (a) determining whether the fibrotic disorder manifests clinical presentations including fibrosis and one or more of the following: inflammation, immune suppression, proliferative dysregulation, and need for an allograft transplant; and (b) selecting a treatment regimen comprising an isoform-specific, context-dependent TGFβ1 inhibitor or an isoform-specific, context-independent TGFβ1 inhibitor, based on the clinical presentations determined in step (a).

[674] Subjects afflicted with fibrotic disorders can display a wide range of symptoms, in addition to fibrosis. The specific combination of clinical manifestations in a subject can guide the selection of an appropriate TGFβ1-inhibitory treatment regimen. For example, a context-independent, isoform-specific TGFβ1 inhibitor can be used to treat the subject if the subject's clinical manifestations indicate a need for inhibition of TGFβ1, without modulating the activity of TGFβ2 or TGFβ3. A treatment regimen including a LTBP context-specific inhibitor can be used to treat the subject if the subject's clinical manifestations indicate that inhibition of TGFβ1 in the extracellular matrix would be beneficial. A LTBP context-specific inhibitor is also advantageous if the subject's clinical manifestations indicate that stimulation of immune effector cells is undesirable. A GARP context-specific inhibitor can be used to treat the subject if the subject's clinical manifestations indicate that blocking the activation/release of TGFβ1 on regulatory T cells (Treg cells) would be beneficial, *e.g.*, to prevent Treg cells from suppressing effector T cell activity. A LRRC33 context-specific inhibitor can be used to treat the subject if the subject's clinical manifestations indicate that blocking the activation/release of TGFβ1 on myeloid cells, monocytes, macrophages, dendritic cells and/or microglia would be beneficial, *e.g.*, to reverse or reduce immune suppression in the subject.

[675] By way of example, a subject having a fibrotic disorder may display clinical manifestations including fibrosis, inflammation, immune suppression, and proliferative dysregulation. Fibrotic

disorders which commonly present with the foregoing combination of symptoms include, *e.g.*, myelofibrosis. In this embodiment, an isoform-specific, context-independent TGF β 1 inhibitor can be selected for treating the subject.

[676] A subject having a fibrotic disorder may display clinical manifestations including fibrosis, inflammation, and need for an allograft transplant. Fibrotic disorders which commonly present with the foregoing combination of symptoms include, *e.g.*, organ fibrosis, such as kidney fibrosis (*e.g.*, fibrosis associated with chronic kidney disease), liver fibrosis (*e.g.*, fibrosis associated with nonalcoholic steatohepatitis (NASH)), or lung fibrosis (*e.g.*, fibrosis associated with idiopathic pulmonary fibrosis (IPF)). In this embodiment, a context-specific LTBP1/3-specific inhibitor is selected for treating the subject.

[677] In another example, a subject having a fibrotic disorder may display clinical manifestations including fibrosis and inflammation. Fibrotic disorders which commonly present with the foregoing combination of symptoms include, *e.g.*, scleroderma. In this embodiment, a context-specific LTBP1/3-specific inhibitor is selected for treating the subject. Additional fibrotic disorders which commonly present with the foregoing combination of symptoms include, *e.g.*, degenerative diseases, such as muscular dystrophy, *e.g.*, Duchenne muscular dystrophy (DMD). In this embodiment, a context-specific LTBP1/3-specific inhibitor is selected for treating the subject.

[678] A subject having a fibrotic disorder may display clinical manifestations including immune suppression and proliferative dysregulation. Fibrotic disorders which commonly present with the foregoing combination of symptoms include, *e.g.*, solid tumors. In some embodiments, the solid tumor is a malignant tumor. In other embodiments, the solid tumor is a benign tumor. In an exemplary embodiment, the subject has desmoplasia (*e.g.*, pancreatic desmoplasia). In some embodiments, patients may have a solid tumor that has been assessed as “inoperable” or not suitable for surgical resection. Thus, in some embodiments, patients are not candidates for surgical resection of the tumor. However, TGF β 1 inhibition therapy comprising a context-selective TGF β 1 inhibitor of the present invention may reverse such non-candidate patients to be more suited for receiving a surgery. In some embodiments, subjects having a solid tumor are poorly responsive to cancer therapy (*e.g.*, the tumor is resistant to the cancer therapy), such as chemotherapy, radiation therapy, CAR-T therapy and checkpoint inhibitor therapy. TGF β 1 inhibition therapy comprising a context-selective TGF β 1 inhibitor of the present invention may at least in part reverse the resistance to render the patient more responsive to the cancer therapy. In some embodiments, a combination therapy comprising both the context-selective TGF β 1 inhibition therapy and the cancer therapy may synergistically treat the cancer. In some embodiments, the context-selective TGF β 1 inhibition therapy administered in conjunction with the cancer therapy may reduce the required dosage of the cancer therapy to produce equivalent or improved clinical effects.

[679] In another exemplary embodiment, the subject has fibroids. In the foregoing embodiments, in which the fibrotic disorder displays clinical manifestations including immune suppression and proliferative dysregulation, a context-specific LTBP1/3-specific inhibitor and/or a context-specific GARP-specific inhibitor are selected for treating the subject.

[680] In another aspect, the invention provides a method of treating a subject having a fibrotic disorder with an isoform-specific, LTBP1/3 context-specific TGF β 1 inhibitor, by selecting a subject having a fibrotic disorder manifesting clinical presentations including fibrosis and the need for an allograft transplant, and administering an effective amount of an isoform-specific, LTBP1/3-specific TGF β 1 inhibitor to the subject. In one embodiment, the method comprises determining whether the fibrotic disorder manifests clinical presentations including fibrosis and the need for an allograft transplant. The LTBP1/3-specific TGF β 1 inhibitor is administered to the subject if the subject exhibits symptoms including fibrosis and the need for an allograft transplant.

[681] In another aspect, the invention provides a method of treating a subject having a fibrotic disorder with an isoform-specific, context-independent TGF β 1 inhibitor, by selecting a subject having a fibrotic disorder manifesting clinical presentations including fibrosis, immune suppression, and/or proliferative dysregulation, and administering an effective amount of an isoform-specific, context-independent TGF β 1 inhibitor to the subject. In one embodiment, the method comprises determining whether the fibrotic disorder manifests clinical presentations including fibrosis, immune suppression, and/or proliferative dysregulation. The isoform-specific, context-independent TGF β 1 inhibitor is administered to the subject if the subject exhibits symptoms including fibrosis, immune suppression, and/or proliferative dysregulation.

[682] Clinical manifestations including inflammation, immune suppression, proliferative dysregulation, and/or the need for an allograft transplant can be determined in a subject having a fibrotic disorder using methods and practices known in the art. Such methods include, for example, physical examination and standard diagnostic tests. In one embodiment, inflammation can be assessed by determining if a subject displays an elevated level of inflammatory biomarkers in plasma, blood, or serum. Such inflammatory biomarkers include, for example, C-reactive protein, interleukin 1 (IL-1), interleukin 6 (IL-6), tumor necrosis factor α (TNF- α), or combinations thereof. Blood tests including erythrocyte sedimentation rate (ESR) and plasma viscosity (PV) can also indicate the presence of inflammation in a subject with a fibrotic disorder. In another embodiment, immune suppression can be assessed by determining the number and composition of a subject's blood cells, *e.g.*, T cells, B cells, NK cells, monocytes, macrophages, etc. Immune suppression can also be assessed by determining if the subject is taking or has a history of taking immunosuppressant medications, or determining if the subject has a condition associated with immune suppression (*e.g.*, hematological malignancies, HIV/AIDS, etc.). In another embodiment, proliferative dysregulation can be assessed using standard tests including blood tests, biopsy, and/or imaging procedures such as CT scan,

ultrasound, and MRI. Other standard tests for diagnosing cancer (*e.g.*, biomarker tests, etc.) can also be used to assess proliferative dysregulation. The need for an allograft transplant can be determined by a clinician using standard procedures. In one embodiment, the loss or partial loss of organ function, or an increased likelihood of loss of organ function, indicates the need for a transplant.

[683] As mentioned, the present invention provides selective targeting of the ECM-associated TGF β 1 complexes enabled by the use of antibodies that are capable of specifically binding LTBP-presented TGF β 1 precursors. While some antibodies of the present invention are capable of binding and inhibiting both LTBP1- and LTBP3-associated proTGF β 1 complexes, others show even greater selectivity in that they only bind either LTBP1-proTGF β 1 or LTBP3-proTGF β 1.

[684] The invention therefore encompasses the recognition that certain patient populations may benefit from TGF β 1 inhibition therapy comprising a context-selective inhibitor that is specific to LTBP1/3-proTGF β 1, over TGF β inhibitors that also affect the immune components of TGF β signaling namely, TGF β associated with GARP. Accordingly, it is contemplated herein that to treat a TGF β -related condition in a subject who has or is at risk of developing an autoimmune condition, a TGF β inhibitor that selectively inhibits matrix-associated TGF β (such as LTBP1/3 context-selective inhibitors of TGF β 1 disclosed herein) may provide therapeutic benefits while minimizing risk of overstimulating the immune system. Such subject may suffer from or may be at risk of developing an autoimmune disorder, such as: Achalasia; Addison's disease; Adult Still's disease;

Agammaglobulinemia; Alopecia areata; Amyloidosis; Ankylosing spondylitis; Anti-GBM/Anti-TBM nephritis; Antiphospholipid syndrome; Autoimmune angioedema; Autoimmune dysautonomia; Autoimmune encephalomyelitis; Autoimmune hepatitis; Autoimmune inner ear disease (AIED); Autoimmune myocarditis; Autoimmune oophoritis; Autoimmune orchitis; Autoimmune pancreatitis; Autoimmune retinopathy; Autoimmune urticaria; Axonal & neuronal neuropathy (AMAN); Baló disease; Behcet's disease; Benign mucosal pemphigoid; Bullous pemphigoid; Castleman disease (CD); Celiac disease; Chagas disease; Chronic inflammatory demyelinating polyneuropathy (CIDP); Chronic recurrent multifocal osteomyelitis (CRMO); Churg-Strauss Syndrome (CSS) or Eosinophilic Granulomatosis (EGPA); Cicatricial pemphigoid; Cogan's syndrome; Cold agglutinin disease; Congenital heart block; Cocksackie myocarditis; CREST syndrome; Crohn's disease; Dermatitis herpetiformis; Dermatomyositis; Devic's disease (neuromyelitis optica); Discoid lupus; Dressler's syndrome; Endometriosis; Eosinophilic esophagitis (EoE); Eosinophilic fasciitis; Erythema nodosum; Essential mixed cryoglobulinemia; Evans syndrome; Fibromyalgia; Fibrosing alveolitis; Giant cell arteritis (temporal arteritis); Giant cell myocarditis; Glomerulonephritis; Goodpasture's syndrome; Granulomatosis with Polyangiitis; Graves' disease; Guillain-Barre syndrome; Hashimoto's thyroiditis; Hemolytic anemia; Henoch-Schonlein purpura (HSP); Herpes gestationis or pemphigoid gestationis (PG); Hidradenitis Suppurativa (HS) (Acne Inversa); Hypogammaglobulinemia; IgA Nephropathy; IgG4-related sclerosing disease; Immune thrombocytopenic purpura (ITP); Inclusion

body myositis (IBM); Interstitial cystitis (IC); Juvenile arthritis; Juvenile diabetes (Type 1 diabetes); Juvenile myositis (JM); Kawasaki disease; Lambert-Eaton syndrome; Leukocytoclastic vasculitis; Lichen planus; Lichen sclerosus; Ligneous conjunctivitis; Linear IgA disease (LAD); Lupus; Lyme disease chronic; Meniere's disease; Microscopic polyangiitis (MPA); Mixed connective tissue disease (MCTD); Mooren's ulcer; Mucha-Habermann disease; Multifocal Motor Neuropathy (MMN) or MMNCB; Multiple sclerosis; Myasthenia gravis; Myositis; Narcolepsy; Neonatal Lupus; Neuromyelitis optica; Neutropenia; Ocular cicatricial pemphigoid; Optic neuritis; Palindromic rheumatism (PR); PANDAS; Paraneoplastic cerebellar degeneration (PCD); Paroxysmal nocturnal hemoglobinuria (PNH); Parry Romberg syndrome; Pars planitis (peripheral uveitis); Parsonage-Turner syndrome; Pemphigus; Peripheral neuropathy; Perivenous encephalomyelitis; Pernicious anemia (PA); POEMS syndrome; Polyarteritis nodosa; Polyglandular syndromes type I, II, III; Polymyalgia rheumatica; Polymyositis; Postmyocardial infarction syndrome; Postpericardiotomy syndrome; Primary biliary cirrhosis; Primary sclerosing cholangitis; Progesterone dermatitis; Psoriasis; Psoriatic arthritis; Pure red cell aplasia (PRCA); Pyoderma gangrenosum; Raynaud's phenomenon; Reactive Arthritis; Reflex sympathetic dystrophy; Relapsing polychondritis; Restless legs syndrome (RLS); Retroperitoneal fibrosis; Rheumatic fever; Rheumatoid arthritis; Sarcoidosis; Schmidt syndrome; Scleritis; Scleroderma; Sjögren's syndrome; Sperm & testicular autoimmunity; Stiff person syndrome (SPS); Subacute bacterial endocarditis (SBE); Susac's syndrome; Sympathetic ophthalmia (SO); Takayasu's arteritis; Temporal arteritis/Giant cell ; arteritisThrombocytopenic purpura (TTP); Tolosa-Hunt syndrome (THS); Transverse myelitis; Type 1 diabetes; Ulcerative colitis (UC); Undifferentiated connective tissue disease (UCTD); Uveitis; Vasculitis; Vitiligo; Vogt-Koyanagi-Harada Disease.

[685] LTBP1 and LTBP3 are both components of the ECM, where they can display or "present" a latent TGF β precursor complex. Some observations from expression studies raise the possibility that deletion, ablation or functional inhibition of LTBP3 may cause certain toxicities. LTBP3^{-/-} mice (as well as some human mutations) have short stature, as well as bone and dental anomalies. These phenotypes are likely associated with disruptions in development, however, but it is possible that LTBP3 plays a role in homeostasis of these tissues in adults (expression in adult bone is reported). Based on these observations, in certain clinical situations (where the disease manifests in a tissue known to express LTBP3 and associated with toxicities) or in certain patient populations, such as pediatric patients who are still in active development, it may be advisable to avoid potential toxicities of LTBP3-related inhibition. Loss of LTBP1 function does appear to be sufficient to protect against at least some forms of fibrosis, as LTBP1^{-/-} KO mice are protected against liver fibrosis (induced by bile duct ligation). Taken together, these data raise the possibility that LTBP1-specific TGF β 1 inhibition could have a superior safety profile as compared to LTBP1/3-TGF β 1 inhibitors in certain situations.

[686] Accordingly, selection of patients or patient populations suitable or likely to benefit from the LTBP1/3-selective inhibitors of the present invention may involve evaluating or confirming expression profiles of LTBP1, LTBP2, LTBP3, LTBP4, GARP, LRRC33, pro- or mature TGF β 1, pro- or mature TGF β 2, pro- or mature TGF β 3, or any combinations thereof. Expression profiles may be obtained by measuring the presence/absence or levels of mRNA and/or proteins in suitable assays from biological samples collected from the subject (*e.g.*, patients). In some embodiments, a soluble circulating fragment(s) of LAP may be used as surrogate marker for the expression of the particular TGF β isoform. In some embodiments, TGF β 1 LAP fragments may be used as a marker of fibrogenesis. See for example, US patent 8,198,412, the contents of which are incorporated herein by reference.

Genetic Markers of Disease:

[687] It has been observed that abnormal activation of the TGF β 1 signal transduction pathway in various disease conditions is associated with altered gene expression of a number of markers. These gene expression markers (*e.g.*, as measured by mRNA) include, but are not limited to: Serpine 1 (encoding PAI-1), MCP-1 (also known as CCL2), Col1a1, Col3a1, FN1, TGF β 1, CTGF, ACTA2 (encoding α -SMA), SNAIL (drives EMT in fibrosis and metastasis by downregulating E-cadherin (Cdh1), MMP2 (matrix metalloprotease associated with EMT), MMP9 (matrix metalloprotease associated with EMT), TIMP1 (matrix metalloprotease associated with EMT), FOXP3 (marker of Treg induction), CDH1 (E cadherin (marker of epithelial cells) which is downregulated by TGF β), and, CDH2 (N cadherin (marker of mesenchymal cells) which is upregulated by TGF β). Interestingly, many of these genes are implicated to play a role in a diverse set of disease conditions, including various types of organ fibrosis, as well as in many cancers, which include myelofibrosis. Indeed, pathophysiological link between fibrotic conditions and abnormal cell proliferation, tumorigenesis and metastasis has been suggested. See for example, Cox and Erler (2014) *Clinical Cancer Research* 20(14): 3637-43 "Molecular pathways: connecting fibrosis and solid tumor metastasis"; Shiga et al. (2015) *Cancers* 7:2443-2458 "Cancer-associated fibroblasts: their characteristics and their roles in tumor growth"; Wynn and Barron (2010) *Semin. Liver Dis.* 30(3): 245-257 "Macrophages: master regulators of inflammation and fibrosis", contents of which are incorporated herein by reference. Without wishing to be bound by a particular theory, the inventors of the present disclosure contemplate that the TGF β 1 signaling pathway may in fact be a key link between these broad pathologies.

[688] The ability of chemotactic cytokines (or chemokines) to mediate leukocyte recruitment (*e.g.*, monocytes/macrophages) to injured or disease tissues has crucial consequences in disease progression. Members of the C-C chemokine family, such as monocyte chemoattractant protein 1

(MCP-1), also known as CCL2, macrophage inflammatory protein 1-alpha (MIP-1 α), also known as CCL3, and MIP-1 β , also known as CCL4, have been implicated in this process.

[689] For example, MCP-1/CCL2 is thought to play a role in both fibrosis and cancer. MCP-1/CCL2 is characterized as a profibrotic chemokine and is a monocyte chemoattractant, and evidence suggests that it may be involved in both initiation and progression of cancer. In fibrosis, MCP-1/CCL2 has been shown to play an important role in the inflammatory phase of fibrosis. For example, neutralization of MCP-1 resulted in a dramatic decrease in glomerular crescent formation and deposition of type I collagen. Similarly, passive immunotherapy with either anti-MCP-1 or anti-MIP-1 alpha antibodies is shown to significantly reduce mononuclear phagocyte accumulation in bleomycin-challenged mice, suggesting that MIP-1 alpha and MCP-1 contribute to the recruitment of leukocytes during the pulmonary inflammatory response (Smith, Biol Signals. 1996 Jul-Aug;5(4):223-31, "Chemotactic cytokines mediate leukocyte recruitment in fibrotic lung disease"). Elevated levels of MIP-1alpha in patients with cystic fibrosis and multiple myeloma have been reported (see, for example: Mrugacz et al., J Interferon Cytokine Res. 2007 Jun;27(6):491-5), supporting the notion that MIP-1 α is associated with localized or systemic inflammatory responses.

[690] Lines of evidence point the involvement of C-C chemokines in tumor progression. For example, tumor-derived MCP-1/CCL2 can promote "pro-cancer" phenotypes in macrophages. For example, in lung cancer, MCP-1/CCL2 has been shown to be produced by stromal cells and promote metastasis. In human pancreatic cancer, tumors secrete CCL2, and immunosuppressive CCR2-positive macrophages infiltrate these tumors. Patients with tumors that exhibit high CCL2 expression/low CD8 T-cell infiltrate have significantly decreased survival. Without wishing to be bound by particular theory, it is contemplated that monocytes that are recruited to an injured or diseased tissue environment may subsequently become polarized in response to local cues (such as in response to tumor-derived cytokines), thereby further contributing to disease progression. These M2-like macrophages are likely to contribute to immune evasion by suppressing effector cells, such as CD4+ and CD8+ T cells. In some embodiments, this process is in part mediated by LRRC33-TGF β 1 expressed by activated macrophages. In some embodiments, the process is in part mediated by GARP-TGF β 1 expressed by Tregs.

[691] Similarly, involvement of PAI-1/Serpine1 has been implicated in a variety of cancers, angiogenesis, inflammation, neurodegenerative diseases (*e.g.*, Alzheimer's Disease). Elevated expression of PAI-1 in tumor and/or serum is correlated with poor prognosis (*e.g.*, shorter survival, increased metastasis) in various cancers, such as breast cancer and bladder cancer (*e.g.*, transitional cell carcinoma) as well as myelofibrosis. In the context of fibrotic conditions, PAI-1 has been recognized as an important downstream effector of TGF β 1-induced fibrosis, and increased PAI-1 expression has been observed in various forms of tissue fibrosis, including lung fibrosis (such as Idiopathic Pulmonary Fibrosis (IPF)), kidney fibrosis, liver fibrosis and scleroderma. In some

embodiments, the process is in part mediated by ECM-associated TGFβ1, *e.g.*, via LTBP1 and/or LTBP3.

[692] Accordingly, in some embodiments, in vivo effects of the TGFβ1 inhibitor therapy may be assessed by measuring changes in gene markers. Suitable markers include TGFβ (*e.g.*, TGFβ1, TGFβ2, and TGFβ3). Suitable markers may also include one or more presenting molecules for TGFβ (*e.g.*, TGFβ1, TGFβ2, and TGFβ3), such as LTBP1, LTBP3, GARP (or LRRC32) and LRRC33. In some embodiments, suitable markers include mesenchymal transition genes (*e.g.*, AXL, ROR2, WNT5A, LOXL2, TWIST2, TAGLN, and/or FAP), immunosuppressive genes (*e.g.*, IL10, VEGFA, VEGFC), monocyte and macrophage chemotactic genes (*e.g.*, CCL2, CCL3, CCL4, CCL7, CCL8 and CCL13), and/or various fibrotic markers discussed herein. Preferred markers are plasma markers.

[693] In some embodiments, an LTBP complex inhibitor of TGFβ1 is used in the treatment of a disease associated with overexpression of one or more of the following: PAI-1 (encoded by Serpine1), MMP2, MMP9, MCP-1 (also known as CCL2), Col1a1, Col3a1, FN1, TGFβ1, CTGF, α-SMA, ITGA11, and ACTA2, wherein the treatment comprises administration of the inhibitor to a subject suffering from the disease in an amount effective to treat the disease. In some embodiments, the inhibitor is used to treat a disease associated with overexpression of PAI-1, MCP-1/CCL2, CTGF, and/or α-SMA. In some embodiments, the disease is myelofibrosis. In some embodiments, the disease is cancer, for example, cancer comprising a solid tumor. In some embodiments, the disease is organ fibrosis, *e.g.*, fibrosis of the liver, the kidney, the lung, the muscle, the skin and/or the cardiac or cardiovascular tissue. In some embodiments, the disease is Alport Syndrom. In some embodiments, the inhibitor reduces expression of one or more of the following: PAI-1 (encoded by Serpine1), MMP2, MMP9, MCP-1 (also known as CCL2), Col1a1, Col3a1, FN1, TGFβ1, CTGF, α-SMA, ITGA11, and ACTA2.

[694] Another biomarker which may be used to assess the in vivo effects of the TGFβ1 inhibitor therapy is blood urea nitrogen (BUN). Urea is naturally formed in the body as a by-product of protein breakdown. The urea travels from you liver to your kidneys where it is filtered/removed from the blood. Accordingly, BUN levels may increase in situations when a patient's kidneys are not functioning properly. For example, patients having kidney fibrosis may display increased BUN. Accordingly, in some embodiments, BUN is measured to assess the in vivo effects of the LTBP-specific inhibitors of TGFβ1 as described herein. In other embodiments, an LTBP-specific inhibitor of TGFβ1 is used in the treatment of a disease associated with increased BUN (*e.g.*, kidney fibrosis and/or acute or chronic kidney disease, damage, or failure). In a particular embodiment, the disease associated with increased BUN is Alport Syndrome.

[695] Accordingly, the present disclosure includes a method of selecting a candidate patient or patient population likely to respond to a TGFβ1 inhibition therapy. Such method may comprise a step

of testing a biological sample collected from the patient (or patient population), such as biopsy samples, for the expression of one or more of the markers discussed herein. Similarly, such genetic marker(s) may be used for purposes of monitoring the patient's responsiveness to a therapy. Monitoring may include testing two or more biological samples collected from the patient, for example, before and after administration of a therapy, and during the course of a therapeutic regimen over time, to evaluate changes in gene expression levels of one or more of the markers, indicative of therapeutic response or effectiveness.

[696] In some embodiments, a method of selecting a candidate patient or patient population likely to respond to a TGF β 1 inhibition therapy may comprise a step of identifying a patient or patient population previously tested for the genetic marker(s), such as those described herein, which showed aberrant expression thereof. In some embodiments, the aberrant marker expression includes elevated levels of at least one of the following: TGF β 1 (and/or TGFB1), LRRC33, GARP, LTBP1, LTBP3, CCL2, CCL3, PAI-1/Serpine1, MMP2, MMP9, Col1a1, Col3a1, FN1, CTGF, α -SMA, ITGA11, and ACTA2. In some embodiments, the patient or patient population (*e.g.*, biological samples collected therefrom) shows elevated TGF β 1 activation, phospho-Smad2/3, or combination thereof. In some embodiments, the patient or patient population shows elevated BUN.

Combination Therapies

[697] The disclosure further encompasses pharmaceutical compositions and related methods used as combination therapies for treating subjects who may benefit from TGF β 1 inhibition *in vivo*. In any of these embodiments, such subjects may receive combination therapies that include a first composition comprising at least one TGF β 1 inhibitor, *e.g.*, antibody or antigen-binding portion thereof, described herein, in conjunction with a second composition comprising at least one additional therapeutic intended to treat the same or overlapping disease or clinical condition. The first and second compositions may both act on the same cellular target, or discrete cellular targets. In some embodiments, the first and second compositions may treat or alleviate the same or overlapping set of symptoms or aspects of a disease or clinical condition. In some embodiments, the first and second compositions may treat or alleviate a separate set of symptoms or aspects of a disease or clinical condition. To give but one example, the first composition may treat a disease or condition associated with TGF β 1 signaling, while the second composition may treat inflammation or fibrosis associated with the same disease, etc. Such combination therapies may be administered in conjunction with each other. The phrase "in conjunction with," in the context of combination therapies, means that therapeutic effects of a first therapy overlaps temporarily and/or spatially with therapeutic effects of a second therapy in the subject receiving the combination therapy. Thus, the combination therapies may be formulated as a single formulation for concurrent administration, or as separate formulations, for sequential administration of the therapies.

[698] In preferred embodiments, combination therapies produce synergistic effects in the treatment of a disease. The term “synergistic” refers to effects that are greater than additive effects (*e.g.*, greater efficacy) of each monotherapy in aggregate.

[699] In some embodiments, combination therapies comprising a pharmaceutical composition described herein produce efficacy that is overall equivalent to that produced by another therapy (such as monotherapy of a second agent) but are associated with fewer unwanted adverse effect or less severe toxicity associated with the second agent, as compared to the monotherapy of the second agent. In some embodiments, such combination therapies allow lower dosage of the second agent but maintain overall efficacy. Such combination therapies may be particularly suitable for patient populations where a long-term treatment is warranted and/or involving pediatric patients.

[700] Accordingly, the invention provides pharmaceutical compositions and methods for use in combination therapies for the reduction of TGFβ1 protein activation and the treatment or prevention of diseases or conditions associated with TGFβ1 signaling, as described herein. Accordingly, the methods or the pharmaceutical compositions further comprise a second therapy. In some embodiments, the second therapy may be useful in treating or preventing diseases or conditions associated with TGFβ1 signaling. The second therapy may diminish or treat at least one symptom(s) associated with the targeted disease. The first and second therapies may exert their biological effects by similar or unrelated mechanisms of action; or either one or both of the first and second therapies may exert their biological effects by a multiplicity of mechanisms of action.

[701] It should be understood that the pharmaceutical compositions described herein may have the first and second therapies in the same pharmaceutically acceptable carrier or in a different pharmaceutically acceptable carrier for each described embodiment. It further should be understood that the first and second therapies may be administered simultaneously or sequentially within described embodiments.

[702] In one embodiment, the inhibitors, *e.g.*, antibodies, described herein that selectively binds a LTBP1-TGFβ1 complex and/or a LTBP3-TGFβ1 complex can be administered with another agent that inhibits TGFβ1 activity. For example, the second agent can be another context-specific TGFβ1 inhibitor. In one embodiment, the combination therapy comprises (i) an inhibitor, *e.g.*, antibody or antigen-binding portion thereof, that selectively binds a LTBP1-TGFβ1 complex and/or a LTBP3-TGFβ1 complex, and (ii) an inhibitor, *e.g.*, antibody or antigen-binding portion thereof, that selectively binds a GARP-TGFβ1 complex. In another embodiment, the combination therapy comprises (i) an inhibitor, *e.g.*, antibody or antigen-binding portion thereof, that selectively binds a LTBP1-TGFβ1 complex and/or a LTBP3-TGFβ1 complex, and (ii) an inhibitor, *e.g.*, antibody or antigen-binding portion thereof, that selectively binds a LRRC33-TGFβ1 complex. Context-specific antibodies that selectively bind LRRC33-TGFβ1 are described, for example, in US 62/503,785, and

context-specific antibodies that selectively bind GARP-TGF β 1 are described, above. The entire contents of the foregoing applications are incorporated by reference herein. In one embodiment, the combination therapy comprises (i) an inhibitor, *e.g.*, antibody or antigen-binding portion thereof, that selectively binds a LTBP1-TGF β 1 complex and/or a LTBP3-TGF β 1 complex, and (ii) an context-independent inhibitor, *e.g.*, antibody or antigen-binding portion thereof, that selectively binds pro/latent TGF β 1 in a complex with a presenting molecule (*e.g.*, LTBP1/3, GARP, and/or LRRC33). Context-independent inhibitors of TGF β 1 are described, for example, in WO 2017/156500, the entire contents of which are incorporated herein by reference.

[703] The one or more anti-TGF β 1 inhibitors, *e.g.*, antibodies, or antigen-binding portions thereof, of the invention may be used in combination with one or more additional therapeutic agents. Examples of the additional therapeutic agents which can be used with an anti-TGF β antibody of the invention include, but are not limited to, a myostatin inhibitor, a VEGF agonist, an IGF1 agonist, an FXR agonist, a CCR2 inhibitor, a CCR5 inhibitor, a dual CCR2/CCR5 inhibitor, a lysyl oxidase-like-2 inhibitor, an ASK1 inhibitor, an Acetyl-CoA Carboxylase (ACC) inhibitor, a p38 kinase inhibitor, Pirfenidone, Nintedanib, selonsertib, cilofexor, firsocostat, Pirfenidone, obeticholic acid, elafibranor, an anti-CD147 antibody, an anti-GP73 antibody, a Galactin-1 inhibitor, selonsertib, a caspase inhibitor (Emricasan, IDN-6556, PF-03491390), a GDF11 inhibitor, a GDF8/myostatin inhibitor, and the like. The GDF8/myostatin inhibitor is preferably a myostatin-selective inhibitor (*e.g.*, an antibody or antigen-binding fragment). The myostatin-selective inhibitor may bind latent myostatin. Non-limiting examples of myostatin-selective inhibitors include SRK-015 (*e.g.*, see WO2017/218592A1) and trevogrumab, or any variant thereof, or an antibody according to WO 2016/098357.

[704] In some embodiments, the additional agent is a checkpoint inhibitor. In some embodiments, the additional agent is selected from the group consisting of a PD-1 antagonist, a PDL1 antagonist, a PD-L1 or PDL2 fusion protein, a CTLA4 antagonist, a GITR agonist, an anti-ICOS antibody, an anti-ICOSL antibody, an anti-B7H3 antibody, an anti-B7H4 antibody, an anti-TIM3 antibody, an anti-LAG3 antibody, an anti-OX40 antibody, an anti-CD27 antibody, an anti-CD70 antibody, an anti-CD47 antibody, an anti-41BB antibody, an anti-PD-1 antibody, an oncolytic virus, and a PARP inhibitor. In some embodiments, the additional therapy is radiation. In some embodiments, the additional agent is a chemotherapeutic agent. In some embodiments, the chemotherapeutic agent is Taxol. In some embodiments, the additional agent is an anti-inflammatory agent. In some embodiments, the additional agent inhibits the process of monocyte/macrophage recruitment and/or tissue infiltration. In some embodiments, the additional agent is an inhibitor of hepatic stellate cell activation. In some embodiments, the additional agent is a chemokine receptor antagonist, *e.g.*, CCR2 antagonists and CCR5 antagonists. In some embodiments, such chemokine receptor antagonist is a dual specific antagonist, such as a CCR2/CCR5 antagonist. In some embodiments, the additional

agent to be administered as combination therapy is or comprises a member of the TGF β superfamily of growth factors or regulators thereof. In some embodiments, such agent is selected from modulators (*e.g.*, inhibitors and activators) of GDF8/myostatin and GDF11. In some embodiments, such agent is an inhibitor of GDF8/myostatin signaling. In some embodiments, such agent is a monoclonal antibody that binds a pro/latent myostatin complex and blocks activation of myostatin. In some embodiments, the monoclonal antibody that binds a pro/latent myostatin complex and blocks activation of myostatin does not bind free, mature myostatin.

[705] Combination therapy that includes a TGF β inhibitor (such as TGF β 1-selective inhibitors disclosed herein), in conjunction with one or more additional therapies, may be considered for treating a variety of liver diseases. Non-limiting examples of liver diseases include: non-alcoholic fatty liver disease (NAFLD), *e.g.*, non-alcoholic fatty liver (NAFL) and non-alcoholic steatohepatitis (NASH), which may include: noncirrhotic NASH with liver fibrosis, liver cirrhosis, NASH with compensated cirrhosis, NASH with decompensated cirrhosis, liver inflammation with fibrosis, liver inflammation without fibrosis; stage 2 and 3 liver fibrosis, stage 4 fibrosis (NASH cirrhosis or cirrhotic NASH with fibrosis), primary biliary cholangitis (PBC) (formerly known as primary biliary cirrhosis), and primary sclerosing cholangitis (PSC).

[706] One or more of the following therapies may be used in conjunction with the TGF β inhibitor (such as TGF β 1-selective inhibitors disclosed herein) for the treatment of a liver disease such as those listed above: Pioglitazone (PPAR γ agonist); Elafibranor (PPAR α/δ agonist); Saroglitazar (PPAR α/γ agonist); Obeticholic acid (FXR agonist); Liraglutide (GLP-1 receptor agonist); Aramchol (SCD inhibitor); Volixibat (SHP-626) (ASBT inhibitor); BMS-986036 (FGF-21 analogue); NGM-282 (FGF-19 analogue); Tesamorelin (GHRH analogue); NDI-010976 (ACC inhibitor); GS-9674 (FXR agonist); Dur-928 (Sulfated oxysterol); AZD4076 (miR-103/107 antagonist); Rosuvastatin (HMG-CoA reductase inhibitor); INT-767 (FXR/TGR5 agonist); Sevelamer (Bile acid sequestrant); Vitamin E (Antioxidant); Pentoxifylline (PDE inhibitor); Cenicriviroc (CCR2/CCR5 antagonist); Emricasan (Caspase inhibitors); GS-4997 (ASK1 inhibitor); Amlexanox (IKK ϵ /TBK1 inhibitor); PXS-4728A (VAP-1 inhibitor); Orlistat (Intestinal lipase inhibitor); IMM-124e (IgG-rich bovine colostrum); Solithromycin (Antibiotic); Faecal microbial transplant (Modulation of gut microbiome); Simtuzumab (LOXL2 antibody); GR-MD-02 (Galectin-3 inhibitor); Trevogrumab (myostatin inhibitor); Garetosmab (activin A inhibitor); and SRK-015 (myostatin inhibitor).

[707] Such combination therapies may advantageously utilize lower dosages of the administered therapeutic agents, thus avoiding possible toxicities or complications associated with the various monotherapies.

Assays for Detecting a LTBP1-TGF β 1 Complex and/or a LTBP3-TGF β 1 Complex

[708] In some embodiments, methods and compositions provided herein relate to a method for detecting a LTBP1-TGF β 1 complex and/or a LTBP3-TGF β 1 complex in a sample obtained from a subject. As used herein, a “subject” refers to an individual organism, for example, an individual mammal. In some embodiments, the subject is a human. In some embodiments, the subject is a non-human mammal. In some embodiments, the subject is a non-human primate. In some embodiments, the subject is a rodent. In some embodiments, the subject is a sheep, a goat, a cattle, poultry, a cat, or a dog. In some embodiments, the subject is a vertebrate, an amphibian, a reptile, a fish, an insect, a fly, or a nematode. In some embodiments, the subject is a research animal. In some embodiments, the subject is genetically engineered, *e.g.*, a genetically engineered non-human subject. The subject may be of either sex and at any stage of development. In some embodiments, the subject is a patient or a healthy volunteer.

[709] In some embodiments, a method for detecting a LTBP1-TGF β 1 complex and/or a LTBP3-TGF β 1 complex in a sample obtained from a subject involves (a) contacting the sample with an antibody that selectively binds a LTBP1-TGF β 1 complex and/or a LTBP3-TGF β 1 complex under conditions suitable for binding of the antibody to the antigen, if the antigen is present in the sample, thereby forming binding complexes; and (b) determining the level of the antibody bound to the antigen (*e.g.*, determining the level of the binding complexes).

[710] In one embodiment, a screening assay that utilizes biotinylated latent TGF β 1 complexes immobilized onto a surface is utilized, which allows for the activation of latent TGF β by integrins, *e.g.*, by providing a tether. Other, non-integrin activators could also be tested in that system. A readout can be measured through reporter cells or other TGF β -dependent cellular responses.

Cell-based assays for measuring TGF β activation

[711] Activation of TGF β (and inhibition thereof by a TGF β test inhibitor, such as an antibody) may be measured by any suitable method known in the art. For example, integrin-mediated activation of TGF β can be utilized in a cell-based assay, such as the “CAGA12” luciferase assay, described in more detail herein. Such an assay system may comprise the following components: i) a source of TGF β (recombinant, endogenous or transfected); ii) a source of integrin (recombinant, endogenous, or transfected); and iii) a reporter system that responds to TGF β activation, such as cells expressing TGF β receptors capable of responding to TGF β and translating the signal into a readable output (*e.g.*, luciferase activity in CAGA12 cells or other reporter cell lines). In some embodiments, the reporter cell line comprises a reporter gene (*e.g.*, a luciferase gene) under the control of a TGF β -responsive promoter (*e.g.*, a PAI-1 promoter). In some embodiments, certain promoter elements that confer sensitivity may be incorporated into the reporter system. In some embodiments, such promoter element is the CAGA12 element. Reporter cell lines that may be used in the assay have been

described, for example, in Abe *et al.* (1994) *Anal Biochem.* 216(2): 276-84, incorporated herein by reference. In some embodiments, each of the aforementioned assay components are provided from the same source (*e.g.*, the same cell). In some embodiments, two of the aforementioned assay components are provided from the same source, and a third assay component is provided from a different source. In some embodiments, all three assay components are provided from different sources. For example, in some embodiments, the integrin and the latent TGF β complex (proTGF β and a presenting molecule) are provided for the assay from the same source (*e.g.*, the same transfected cell line). In some embodiments, the integrin and the TGF are provided for the assay from separate sources (*e.g.*, two different cell lines, a combination of purified integrin and a transfected cell). When cells are used as the source of one or more of the assay components, such components of the assay may be endogenous to the cell, stably expressed in the cell, transiently transfected, or any combination thereof.

[712] A skilled artisan could readily adapt such assays to various suitable configurations. For instance, a variety of sources of TGF β may be considered. In some embodiments, the source of TGF β is a cell that expresses and deposits TGF β (*e.g.*, a primary cell, a propagated cell, an immortalized cell or cell line, etc.). In some embodiments, the source of TGF β is purified and/or recombinant TGF β immobilized in the assay system using suitable means. In some embodiments, TGF β immobilized in the assay system is presented within an extracellular matrix (ECM) composition on the assay plate, with or without de-cellularization, which mimics fibroblast-originated TGF β . In some embodiments, TGF β is presented on the cell surface of a cell used in the assay. Additionally, a presenting molecule of choice may be included in the assay system to provide suitable latent-TGF β complex. One of ordinary skill in the art can readily determine which presenting molecule(s) may be present or expressed in certain cells or cell types. Using such assay systems, relative changes in TGF β activation in the presence or absence of a test agent (such as an antibody) may be readily measured to evaluate the effects of the test agent on TGF β activation *in vitro*.

[713] Such cell-based assays may be modified or tailored in a number of ways depending on the TGF β isoform being studied, the type of latent complex (*e.g.*, presenting molecule), and the like. In some embodiments, a cell known to express integrin capable of activating TGF β may be used as the source of integrin in the assay. Such cells include SW480/ β 6 cells (*e.g.*, clone 1E7). In some embodiments, integrin-expressing cells may be co-transfected with a plasmid encoding a presenting molecule of interest (such as GARP, LRRC33, LTBP (*e.g.*, LTBP1 or LTBP3), etc.) and a plasmid encoding a pro-form of the TGF β isoform of interest (such as proTGF β 1). After transfection, the cells are incubated for sufficient time to allow for the expression of the transfected genes (*e.g.*, about 24 hours), cells are washed, and incubated with serial dilutions of a test agent (*e.g.*, an antibody). In some embodiments, tissue culture wells or plates may be coated with a substance that provides a favorable substrate upon which cells may adhere, grow, and/or deposit ECM components. This may

facilitate ECM architecture, organization or adhesion of the cells thereto. For example, charged substances such as poly-lysine may be used to pre-coat the tissue culture substrate. Additionally or alternatively, one or more components of ECM, such as laminins, fibronectins, etc., may be used as substrate for coating. In some embodiments, the cells are seeded on ECM protein-coated wells/plates prior to transfection. In some embodiments, the cells are seeded on ECM coated wells/plates after transfection. In some embodiments, the wells/plates are coated with fibronectin. In some embodiments, the wells/plates are coated with human fibronectin.

[714] After transfection (and optionally seeding on ECM coated well/plates), a reporter cell line (*e.g.*, CAGA12 cells) is added to the assay system, followed by appropriate incubation time to allow TGF β signaling. After an incubation period (*e.g.*, about 18-20 hours) following the addition of the test agent, signal/read-out (*e.g.*, luciferase activity) is detected using suitable means (*e.g.*, for luciferase-expressing reporter cell lines, the Bright-Glo reagent (Promega) can be used). In some embodiments, Luciferase fluorescence may be detected using a BioTek (Synergy H1) plate reader, with autogain settings.

Kits for Use in Alleviating Diseases/Disorders Associated with LTBP1/3-TGF β

[715] The present disclosure also provides kits for use in alleviating diseases/disorders associated with a TGF β -related indication. Such kits can include one or more containers comprising an inhibitor, *e.g.*, antibody, or antigen-binding portion thereof, that selectively binds to a LTBP1-TGF β 1 complex and/or a LTBP3-TGF β 1 complex, *e.g.*, any of those described herein.

[716] In some embodiments, the kit can comprise instructions for use in accordance with any of the methods described herein. The included instructions can comprise a description of administration of the inhibitor, *e.g.*, antibody, or antigen-binding portion thereof, that selectively binds a LTBP1-TGF β 1 complex and/or a LTBP3-TGF β 1 complex to treat, delay the onset, or alleviate a target disease as those described herein. The kit may further comprise a description of selecting an individual suitable for treatment based on identifying whether that individual has the target disease. In still other embodiments, the instructions comprise a description of administering an antibody, or antigen-binding portion thereof, to an individual at risk of the target disease.

[717] The instructions relating to the use of inhibitors, *e.g.*, antibodies, or antigen-binding portions thereof, that selectively bind a LTBP1-TGF β 1 complex and/or a LTBP3-TGF β 1 complex generally include information as to dosage, dosing schedule, and route of administration for the intended treatment. The containers may be unit doses, bulk packages (*e.g.*, multi-dose packages) or sub-unit doses. Instructions supplied in the kits of the disclosure are typically written instructions on a label or package insert (*e.g.*, a paper sheet included in the kit), but machine-readable instructions (*e.g.*, instructions carried on a magnetic or optical storage disk) are also acceptable. The label or package insert can indicate that the composition is used for treating, delaying the onset and/or alleviating a

disease or disorder associated with a TGF β -related indication. Instructions may be provided for practicing any of the methods described herein.

[718] The kits of this disclosure can be provided in suitable packaging. Suitable packaging includes, but is not limited to, vials, bottles, jars, flexible packaging (*e.g.*, sealed Mylar or plastic bags), and the like. Also contemplated are packages for use in combination with a specific device, such as an inhaler, nasal administration device (*e.g.*, an atomizer) or an infusion device such as a minipump. A kit may have a sterile access port (for example the container may be an intravenous solution bag or a vial having a stopper pierceable by a hypodermic injection needle). The container may also have a sterile access port (for example the container may be an intravenous solution bag or a vial having a stopper pierceable by a hypodermic injection needle). At least one active agent in the composition is an inhibitor, *e.g.*, antibody, or antigen-binding portion thereof, that selectively binds a LTBP1-TGF β 1 complex and/or a LTBP3-TGF β 1 complex, as described herein.

[719] Kits may optionally provide additional components such as buffers and interpretive information. Normally, the kit comprises a container and a label or package insert(s) on or associated with the container. In some embodiments, the disclosure provides articles of manufacture comprising contents of the kits described above.

Diagnostics, patient selection, monitoring

[720] Therapeutic methods that include TGF β 1 inhibition therapy may comprise diagnosis of a TGF β 1 indication and/or selection of patients likely to respond to such therapy. Additionally, patients who receive the TGF β 1 inhibitor may be monitored for therapeutic effects of the treatment, which typically involves measuring one or more suitable parameters which are indicative of the condition and which can be measured (*e.g.*, assayed) before and after the treatment and evaluating treatment-related changes in the parameters. For example, such parameters may include levels of biomarkers present in biological samples collected from the patients. Biomarkers may be RNA-based, protein-based, cell-based and/or tissue-based. For example, genes that are overexpressed in certain disease conditions may serve as the biomarkers to diagnose and/or monitor the disease or response to the therapy. Cell-surface proteins of disease-associated cell populations may serve as biomarkers. Such methods may include the direct measurements of disease parameters indicative of the extent of the particular disease. Any suitable sampling methods may be employed, such as serum/blood samples, biopsies, and imaging.

[721] While biopsies have traditionally been the standard for diagnosing and monitoring various diseases, such as fibrosis (*e.g.*, organ fibrosis) and proliferative disorders (*e.g.*, cancer), less invasive alternatives may be preferred. For example, many non-invasive *in vivo* imaging techniques may be used to diagnose, monitor, and select patients for treatment. Thus, the invention includes the use of *in vivo* imaging techniques to diagnose and/or monitor disease in a patient or subject. In some

embodiments, the patient or subject is receiving an isoform-specific TGF β 1 inhibitor as described herein. In some embodiments, the patient or subject is receiving an isoform-specific TGF β 1 inhibitor as described herein. In other embodiments, an in vivo imaging technique may be used to select patients for treatment with an isoform-specific TGF β 1 inhibitor. In some embodiments, such techniques may be used to determine if or how patients respond to a therapy, *e.g.*, TGF β 1 inhibition therapy.

[722] Exemplary in vivo imaging techniques used for the methods include, but are not limited to X-ray radiography, magnetic resonance imaging (MRI), medical ultrasonography or ultrasound, endoscopy, elastography, tactile imaging, thermography, medical photography. Other imaging techniques include nuclear medicine functional imaging, *e.g.*, positron emission tomography (PET) and Single-photon emission computed tomography (SPECT). Methods for conducting these techniques and analyzing the results are known in the art.

[723] Non-invasive imaging techniques commonly used to diagnose and monitor cancer include, but are not limited to: magnetic resonance imaging (MRI), computed tomography (CT), ultrasound, positron emission tomography (PET), single-photon emission computed tomography (SPECT), fluorescence reflectance imaging (FRI), and fluorescence mediated tomography (FMT). Hybrid imaging platforms may also be used to diagnose and monitor cancer. For example, hybrid techniques include, but are not limited to: PET-CT, FMT-CT, FMT-MRI, and PET-MRI. Dynamic contrast enhanced MRI (DCE-MRI) is another imaging technique commonly used to detect breast cancers. Methods for conducting these techniques and analyzing the results are known in the art.

[724] Non-invasive imaging techniques commonly used to diagnosis and monitor fibrosis include, but are not limited to: ultrasound (*e.g.*, conventional or contrast-enhanced ultrasound), ultrasound elastography (*e.g.*, transient elastography, point shear wave elastography and 2D-shear wave elastography), CT scan (*e.g.*, conventional CT or CT perfusion imaging), magnetic resonance imaging (MRI) (*e.g.*, conventional MRI, Magnetic resonance elastography, diffusion weighted magnetic resonance imaging, gadoxetic acid disodium, and magnetic resonance perfusion imaging).

[725] In some embodiments, non-invasive imaging techniques are used to assess levels of liver fibrosis or hepatic steatosis. For example, imaging techniques particularly useful to assess liver fibrosis may include but are not limited to: FibroScan (transient elastography; TE), point shear wave elastography (pSWE; a.k.a. acoustic radiation force impulse (ARFI)), 2D-3D SWE, magnetic resonance elastography (MRE), and multiparametric MRI. Imaging techniques particularly useful to assess hepatic steatosis may include but are not limited to: ultrasonography, controlled attenuation parameter (CAP) elastography, MRI-estimated proton density fat fraction (MRI-PDFF), and magnetic resonance spectroscopy (MRS). In some embodiments, the in vivo imaging technique is used to assess liver stiffness. In some embodiments, the in vivo imaging technique is used to detect and

assess intrahepatic triglyceride levels. In some embodiments, in vivo imaging technique is used to assess liver surface nodularity (LSN; a.k.a. “liver score”), liver stiffness, and/or liver segmental volume ratio (LSVR), which are all beneficial in the staging of hepatic fibrosis and sub-staging cirrhosis. Methods for conducting these techniques and analyzing the results are known in the art.

[726] More recently, non-invasive imaging methods are being developed which will allow the detection of cells of interest (*e.g.*, cytotoxic T cells, macrophages, and cancer cells) in vivo. See for example, www.imaginab.com/technology/; Tavare et al. (2014) PNAS, 111(3): 1108-1113; Tavare et al. (2015) J Nucl Med 56(8): 1258-1264; Rashidian et al. (2017) J Exp Med 214(8): 2243-2255; Beckford Vera et al. (2018) PLoS ONE 13(3): e0193832; and Tavare et al. (2015) Cancer Res 76(1): 73-82, each of which is incorporated herein by reference. So-called “T-cell tracking” is aimed to detect and localize anti-tumor effector T-cells in vivo. This may provide useful insights into understanding the immunosuppressive phenotype of solid tumors. Tumors that are well-infiltrated with cytotoxic T cells (“inflamed” or “hot” tumors) are likely to respond to cancer therapies such as checkpoint blockade therapy (CBT). On the other hand, tumors with immunosuppressive phenotypes tend to have poor T-cell infiltration even when there is an anti-tumor immune response. These so-called “immune excluded” tumors likely fail to respond to cancer therapies such as CBT. T-cell tracking techniques may reveal these different phenotypes and provide information to guide in therapeutic approach that would likely benefit the patients. For example, patients with an “immune excluded” tumor are likely benefit from a TGFβ1 inhibitor therapy to help reverse the immunosuppressive phenotype. It is contemplated that similar techniques may be used to diagnose and monitor other diseases, for example, fibrosis. Typically, antibodies or antibody-like molecules engineered with a detection moiety (*e.g.*, radiolabel, fluorescence, etc.) can be infused into a patient, which then will distribute and localize to sites of the particular marker (for instance CD8+ and M2 macrophages).

[727] Non-invasive in vivo imaging techniques may be applied in a variety of suitable methods for purposes of diagnosing patients; selecting or identifying patients who are likely to benefit from TGFβ1 inhibitor therapy; and/or, monitoring patients for therapeutic response upon treatment. Any cells with a known cell-surface marker may be detected/localized by virtue of employing an antibody or similar molecules that specifically bind to the cell marker. Typically, cells to be detected by the use of such techniques are immune cells, such as cytotoxic T lymphocytes, regulatory T cells, MDSCs, disease-associated macrophages, (M2 macrophages such as TAMs and FAMs), NK cells, dendritic cells, and neutrophils.

[728] Non-limiting examples of suitable immune cell markers include monocyte markers, macrophage markers (*e.g.*, M1 and/or M2 macrophage markers), CTL markers, suppressive immune cell markers, MDSC markers (*e.g.*, markers for G- and/or M-MDSCs), including but are not limited to: CD8, CD3, CD4, CD11b, CD163, CD206, CD68, CD14, CD15, CD66, CD34, CD25, and CD47.

[729] In some embodiments, the in vivo imaging technique measures hepatic steatosis, hepatic triglycerides, immune cells (*e.g.*, as described below), and/or myofibroblasts. In some embodiments, the treatment reduces triglycerides, steatosis, liver surface nodules, inflammation, and/or macrophages, in the diseased tissue. In some embodiments, the selected patient has an intrahepatic triglyceride content of >5.5% of liver volume, optionally wherein the intrahepatic triglyceride content is >10% of liver volume. In some embodiments, the treatment reduces intrahepatic triglyceride content to $\leq 5.5\%$ of liver volume. In some embodiments, the treatment reduces MDSCs in the diseased tissue. In some embodiments, the treatment reduces macrophages in the diseased tissue. In some embodiments, the effective amount is from 0.1 mg/kg to 30 mg/kg, optionally 3 mg/kg to 30 mg/kg. In some embodiments, the method further comprises monitoring the subject for a therapeutic response as described herein (*e.g.*, reduced triglycerides, reduced steatosis, reduced liver surface nodules, reduced inflammation, reduced macrophages, and/or reduced liver score).

Process of screening; manufacture

[730] The invention encompasses screening methods, production methods and manufacture processes of antibodies or fragments thereof which bind to and dissociates at slow rates from a hLTBP1-proTGF β 1 complex and/or a hLTBP3-proTGF β 1 complex, and pharmaceutical compositions and related kits comprising the same.

[731] Methods for making a pharmaceutical composition comprising the antibody (or an engineered construct comprising an antigen-binding fragment thereof) require identification and selection of such antibodies with desirable attributes. Here, the invention includes the recognition that antibodies with low k_{OFF} values may provide the durability that reflects the mechanism of action of these activation inhibitors, which do not rely on the ability to rapidly compete binding with endogenous receptors, but rather, exert inhibitory effects by latching onto inactive latent forms of TGF β 1 within the tissue. The ability to stay bound to the latent antigen complex (corresponding to low dissociation rates) may achieve durable potency in vivo.

[732] Accordingly, the invention provides a method for manufacturing a pharmaceutical composition comprising a TGF β 1-selective activation inhibitor, wherein the method comprises the steps of: selecting an antibody or antigen-binding fragment thereof that specifically binds a human LLC with a low dissociation rate (*e.g.*, $\leq 5 \times 10^{-4}$ (1/s)), and producing the antibody at large-scale.

[733] The selection of inhibitors with favorable off rates (low dissociation) may be determined with monovalent antibodies (*e.g.*, Fab fragments) or full-length antibodies (*e.g.*, IgGs).

[734] In some embodiments, the step of producing comprises a mammalian cell culture having a volume of 250L or greater, *e.g.*, 1000L, 2000L, 3000L, 4000L. The method may further comprise the step of purifying the antibody from the cell culture, and optionally formulating the purified antibody

into a pharmaceutical composition. In some embodiments, the method further comprises the step of testing the selected antibody in a suitable preclinical model for efficacy and safety and confirming that the antibody is efficacious at a NOAEL. The safety assessment may include in vivo toxicology study comprising histopathology and immune-directed safety assessment including, for example, in vitro cytokine release assays and platelet assays.

[735] In order to achieve durable inhibitory effects, antibodies with dissociation rates (*e.g.*, monovalent dissociation rates) of no greater than $10.0\text{E-}4$ (s^{-1}) (*e.g.*, $5.0\text{E-}4$ or less, $1.0\text{E-}4$ or less, $5.0\text{E-}5$ or less) may be selected for therapeutic use and/or large-scale manufacture in accordance with the present disclosure.

[736] While several embodiments of the present disclosure have been described and illustrated herein, those of ordinary skill in the art will readily envision a variety of other means and/or structures for performing the functions and/or obtaining the results and/or one or more of the advantages described herein, and each of such variations and/or modifications is deemed to be within the scope of the present disclosure. More generally, those skilled in the art will readily appreciate that all parameters, dimensions, materials, and configurations described herein are meant to be exemplary and that the actual parameters, dimensions, materials, and/or configurations will depend upon the specific application or applications for which the teachings of the present disclosure is/are used. Those skilled in the art will recognize, or be able to ascertain using no more than routine experimentation, many equivalents to the specific embodiments of the disclosure described herein. It is, therefore, to be understood that the foregoing embodiments are presented by way of example only and that, within the scope of the appended claims and equivalents thereto, the disclosure may be practiced otherwise than as specifically described and claimed. The present disclosure is directed to each individual feature, system, article, material, and/or method described herein. In addition, any combination of two or more such features, systems, articles, materials, and/or methods, if such features, systems, articles, materials, and/or methods are not mutually inconsistent, is included within the scope of the present disclosure.

[737] The present invention is further illustrated by the following examples, which are not intended to be limiting in any way. The entire contents of all references, patents and published patent applications cited throughout this application, as well as the Figures, are hereby incorporated herein by reference.

EXAMPLES

[738] Transforming growth factor beta 1 (TGF β 1) is expressed as a pro-protein that is proteolytically cleaved into a C-terminal growth factor and an N-terminal prodomain. After cleavage,

the prodomain remains noncovalently associated with the growth factor, preventing receptor binding. This latent TGF β 1 forms a large latent complex (LLC) through disulfide bonds that link the prodomain to presenting molecules, and these large latent complexes are then deposited into the extracellular matrix (ECM) or brought to the cell surface. These presenting molecules provide an anchor for specific α V β integrins to exert traction force on latent TGF β 1. Four TGF β 1 presenting proteins have been identified: Latent TGF β Binding Protein-1 (LTBP1) and LTBP3 are deposited in the extracellular matrix, while Glycoprotein-A Repetitions Predominant (GARP/LRRC32) and Leucine-Rich Repeat-Containing Protein 33 (LRRC33) present latent TGF β 1 on the surface of immune cells. TGF β 1 is involved in tissue homeostasis processes and regulation of immune responses, and dysregulation of its activation is a key driver of organ fibrosis, cancer, and autoimmunity.

[739] As compared to the TGF β growth factors and the receptors, which are expressed broadly, the four presenting molecule-proTGF β complexes, namely, LTBP1-proTGF β , LTBP3-proTGF β , GARP-proTGF β and LRRC33-proTGF β , show more restricted or selective (*e.g.*, tissue-specific) expression patterns, giving rise to functional compartmentalization of TGF β activities by virtue of association. The presenting molecule-proTGF β complexes therefore provide discrete “contexts” of TGF β signaling within the tissue in which the presenting molecules are expressed. These contexts may be divided into two broad categories: i) TGF β signaling associated with the ECM (*e.g.*, matrix-associated TGF β function); and ii) TGF β signaling associated with cells (particularly certain immune cell function). The LTBP1-proTGF β and LTBP3-proTGF β complexes fall under the first category, while GARP-proTGF β and LRRC33-proTGF β complexes fall under the second category.

[740] Non-selective targeting of TGF β activity for therapeutic purposes has been challenging due to dose-limiting toxicities reported for pan-TGF β pathway inhibitors, as well as immune system activation through chronic TGF β suppression. In an effort to address this therapeutic need for both isoform- and context-selectivity for TGF β 1 targeting, provided herein are inhibitors of TGF β that are capable of selectively inhibiting the activation of TGF β that is associated with the ECM. In some embodiments, the inhibitors are also selective for a particular TGF β isoform (*e.g.*, proTGF β 1, proTGF β 2, and/or proTGF β 3). The isoform-specific monoclonal antibodies bind the latent TGF β 1 prodomain, with no detectable binding to latent TGF β 2 or TGF β 3, and inhibit integrin-mediated activation of latent TGF β 1 *in vitro* with context-selectivity. In order to facilitate antibody discovery and characterization efforts, context-dependent cell-based assays of TGF β 1 activation were developed.

Example 1: Development of Context-Specific Inhibitors that Bind a LTBP1/3-TGF β 1 Complex

[741] SR-AB1 was used as a control. SR-AB1 binds latent TGF β 1 independent of the presenting molecule (see **Fig. 2A**).

[742] Antibodies that are selective for TGF β 1-containing large latent complexes were developed. SR-AB2 was selected for further analysis using the functional assays described in the below examples. The heavy and light chain variable regions of SR-AB2 were sequenced (**Fig. 8**); complementarity determining regions are underlined. It was demonstrated that SR-AB2 binds LTBP-presented latent TGF β 1 complexes but does not bind GARP-TGF β 1 or proTGF β 1 alone (**Fig. 2B**). However, as described below, the functional effect of such selective binding was unknown and could not be determined using currently known techniques without the further development of novel functional assays.

Example 2: Functional Assays to Detect Inhibition of Activated Recombinant Latent TGF β 1

[743] In order to identify isoform-specific inhibitors that bind the latent TGF β 1 prodomain with no detectable binding to latent TGF β 2 or TGF β 3 and that inhibit integrin-mediated activation of latent TGF β 1 *in vitro* with context-dependency, new functional assays were required. Prior to the instant invention, assays were not available which could detect isoform-specific TGF β 1 antibodies that bound only to LTBPs. Specifically, previous assay formats could not differentiate between the activation of proTGF β 1 presented by endogenous presenting molecules and the activation of proTGF β 1 presented by exogenous LTBPs. By directly transfecting integrin-expressing cells, the novel assays disclosed herein establish a window between endogenous presenter-proTGF β 1 activity and exogenous LTBP-proTGF β 1 activity. As LTBP-proTGF β 1 complexes are embedded in the extracellular matrix, the assay plate coating is also an important component of the assay. The use of high binding plates, coated with the ECM protein Fibronectin, made the LTBP assays more robust. In other words, prior to the instant disclosure, there was no assay window between proTGF β 1 transfection and co-transfection of LTBP1/3 + proTGF β 1. Prior to the instant invention, the only available assay format was a triple co-culture system: transfectants (latent TGF β presenting cells) + integrin expressing cells (activator) + CAGA cells (reporter). By combining the first two cell populations, and directly transfecting the integrin expressing cells with TGF β and presenting molecules, a window for LTBP-proTGF β 1 activation was established herein.

[744] The issue of ‘bulk transfection’ (*i.e.*, transfection in a separate well/plate/dish prior to seeding in the assay well) vs ‘direct transfection’ (*i.e.*, transfection in the assay well) protocol and whether an assay window is seen for LTBP complexes seems to be cell dependent in some situations. Thus, the discovery and characterization of LTBP1/3-TGF β 1 inhibitors, *e.g.*, antibodies and antigen-binding portions thereof, would not have been possible without the development of context-dependent cell-based assays of TGF β 1 activation described herein (see also **Fig. 3A** and **3B**).

[745] Specifically, to determine if the antibodies developed in Example 1 were functional, cell-based assays of α V β integrin activation of TGF β 1 large latent complex (LLC) were developed, which are specific for each known presenting molecule: LTBP1, LTBP3, GARP and LRRC33. Through the

process of assay development and optimization, it was determined that fibronectin is a critical ECM protein for the integrin-dependent *in vitro* activation of LTBP presented TGF β 1 LLCs. The context-independent and LTBP complex-specific TGF β 1 LLC antibodies were also validated as inhibitors of integrin-dependent activation using the below assays. Thus, the antibodies developed in Example 1 can be divided into 2 classes: antibodies which bind all TGF β 1 containing complexes (isoform-specific and context independent), and antibodies which only bind LTBP presented TGF β 1 LLC. As described in more detail herein, the development of an LTBP complex-specific class of inhibitor, which was not capable of being identified prior to the assays developed and described herein, enables a therapeutic approach for treating fibrotic indications, and could allow for chronic dosing while avoiding immune system activation due to TGF β 1 inhibition of immune suppressive cells.

Assay I. Activation of Latent TGF β 1 using SW480/ β 6 cells

[746] For the assay depicted in **Fig. 3A**, the following protocol was developed. This assay is optimal for extracellular matrix (LTBP presented) activation by integrin cells.

Materials:

- MvLu1-CAGA12 cells (Clone 4A4)
- SW480/ β 6 cells (Clone 1E7) (α V subunit is endogenously expressed at high levels; β 6 subunit is stably overexpressed)
- Costar white walled TC treated 96 well assay plate #3903
- Greiner Bio-One High Binding white μ clear 96 well assay plate #655094
- Human Fibronectin (Corning #354008)
- P200 multichannel pipet
- P20, P200, and P1000 pipets with sterile filter tips for each
- Sterile microfuge tubes and rack
- Sterile reagent reservoirs
- 0.4% trypan blue
- 2mL, 5mL, 10mL, and 25mL sterile pipets
- Tissue culture treated 100mm or 150mm plates
- 70%Ethanol
- Opti-MEM reduced serum media (Life Tech #31985-070)
- Lipofectamine 3000 (Life Tech #L3000015)
- Bright-Glo luciferase assay reagent (Promega #E2620)
- 0.25% Trypsin + 0.53mM EDTA
- proTGF β 1 expression plasmid, human (SR005)
- LTBP1S expression plasmid, human (SR044)
- LTBP3 expression plasmid, human (SR117)

- LRRC32 (GARP) expression plasmid, human (SR116)
- LRRC33 expression plasmid, human (SR386)

Equipment:

- BioTek Synergy H1 plate reader
- TC hood
- Bench top centrifuge
- CO2 incubator 37°C 5% CO2
- 37°C water/bead bath
- Platform shaker
- Microscope
- Hemocytometer/countess

Definitions:

- CAGA12 4A4 cells: Derivative of MvLu1 cells (Mink Lung Epithelial Cells), stably transfected with CAGA12 synthetic promoter, driving luciferase gene expression
- DMEM-0.1%BSA: Assay media; base media is DMEM (Gibco Cat# 11995-065), media also contains BSA diluted to 0.1% w/v, penicillin/streptomycin, and 4mM glutamine
- D10: DMEM 10% FBS, P/S, 4mM glutamine, 1% NEAA, 1X GlutaMAX (Gibco Cat# 35050061)
- SW480/β6 Media: D10 + 1000μg/mL G-418
- CAGA12 (4A4) media: D10 + 0.75μg/mL puromycin

Procedure:

[747] On Day 0, cells were seeded for transfection. SW480/β6 (clone 1E7) cells were detached with trypsin and pelleted (spin 5 min @ 200 x g). Cell pellet was resuspended in D10 media and viable cells per ml were counted. Cells were seeded at 5.0e6 cells/12ml/100mm TC dish. For CAGA12 cells, cells were passaged at a density of 1.0 million per T75 flask, to be used for the assay on Day 3. Cultures were incubated at 37°C and 5% CO₂.

[748] On Day 1, integrin-expressing cells were transfected. Manufacturer's protocol for transfection with Lipofectamine 3000 reagent was followed. Briefly, the following were diluted into OptiMEM I, for 125μl per well: 7.5μg DNA (presenting molecule) + 7.5μg DNA (proTGFβ1), 30μl P3000, and up to 125μl with OptiMEM I. The well was mixed by pipetting DNA together, then OptiMEM was added. P3000 was added, and everything was mixed well by pipetting. A master mix of Lipofectamine3000 was made, to be added to DNA mixes: for the LTBP1 assay: 15μl Lipofectamine3000, up to 125μl in OptiMEM I, per well; for the LTBP3 assay: 45μl Lipofectamine3000, up to 125μl in OptiMEM I, per well. Diluted Lipofectamine3000 was added to DNA, mixed well by pipetting, and incubated at room temp for 15min. After the incubation, the solution was mixed a few times by pipetting, and then 250μl of DNA:Lipofectamine3000 (2 x 125μl)

per dish was added dropwise. Each dish was gently swirled to mix and the dish was returned to the tissue culture incubator for ~ 24hrs.

[749] On Days 1-2, the assay plates were coated with human fibronectin. Specifically, lyophilized fibronectin was diluted to 1mg/ml in ultra-pure distilled water (sterile). 1mg/ml stock solution was diluted to 19.2µg/ml in PBS (sterile). 50µl/well was then added to the assay plate (high binding) and incubated O/N in tissue culture incubator (37°C and 5% CO₂). Final concentration was 3.0µg/cm².

[750] On Day 2, transfected cells were plated for assay and inhibitor addition. First, the fibronectin coating was washed by adding 200µl/well PBS to the fibronectin solution already in the assay plate. Removed wash manually with multichannel pipette. Wash was repeated for two washes total. The plate was allowed to dry at room temperature with lid off prior to cell addition. The cells were then plated by detaching with trypsin and pelleted (spin 5 min @ 200 x g.). The pellet was resuspended in assay media and viable cells were counted per ml. For the LTBP1 assay cells were diluted to 0.10e6cells/ml and seeded 50µl per well (5,000 cells per well). For the LTBP3 assay, cells were diluted to 0.05e6cells/ml and seeded 50µl per well (2,500 cells per well). To prepare functional antibody dilutions, antibodies were pre-diluted to a consistent working concentration in vehicle. Stock antibodies were serially diluted in vehicle (PBS is optimal, avoid sodium citrate buffer). Each point of serial dilution was diluted into assay media for a 4X final concentration of antibody. 25µl of 4X antibody was added per well and cultures were incubated at 37°C and 5% CO₂ for ~24 hours.

[751] On Day 3, the TGFβ reporter cells were added. CAGA12 (clone 4A4) cells for the assay were detached with trypsin and pelleted (spin 5 min @ 200 x g.). The pellet was resuspended in assay media and viable cells per ml were counted. Cells were diluted to 0.4e⁶cells/ml and seeded 50µl per well (20,000 cells per well). Cells were returned to incubator.

[752] On Day 4, the assay was read (16-20 hours after antibody and/or reporter cell addition). Bright-Glo reagent and test plate were allowed to come to room temperature before reading. Read settings on BioTek Synergy H1 were set using TMLC_std protocol – this method has an auto-gain setting. Selected positive control wells for autoscale (high). 100µL of Bright-Glo reagent was added per well. Incubated for 2min with shaking, at room temperature, protected plate from light. The plate was read on BioTek Synergy H1.

[753] In some embodiments, TGFβ activity associated with endogenous presenting molecules (*e.g.*, LTBP1/3) may be assessed by only transfecting proTGFβ1 (*i.e.*, without co-transfecting LTBP1/3). In some embodiments, the presenting molecule and proTGFβ DNA may be directly transfected into SW480/β6 cells seeded in an assay well (*i.e.*, “direct transfection”), rather than transfecting the cells in a separate well/dish/plate as essentially described above (*i.e.*, “bulk transfection”). See also Assay II below for direct transfection protocol. In some embodiments, SW480/β6 cells may be seeded in assay wells without fibronectin as essentially described in Assay II below.

[754] Results:

[755] Data generated from this assay reflected TGF β activity in cell supernatants (**Fig. 3A**). Specifically, SW480/ β 6 cells were bulk transfected with LTBP1/3 and proTGF β 1, and seeded on fibronectin as described above. Raw data units were relative light units (RLU). **Fig. 3A** demonstrates that transfection of LTBP1-proTGF β 1 and/or LTBP3-proTGF β 1, but not proTGF β 1 alone, induces a TGF β activation signal.

[756] The assay was further optimized as described in **Fig. 4**. Specifically, the relative contribution of presenting molecule and/or proTGF β 1 to latent TGF β 1 activation was determined. In this assay, SW480/ β 6 cells were bulk transfected with the indicated DNA molecules and seeded on pre-coated assay wells with fibronectin as essentially described above. As shown in **Fig. 4A**, a significant increase in latent TGF β 1 activation upon co-transfection of presenting molecule and proTGF β 1 was observed. **Fig. 4B** depicts the optimization of co-transfection by changing the ratio of plasmid DNAs for presenting molecule and proTGF β 1. Equivalent amounts of each plasmid were found to be optimal for co-transfection.

[757] **Fig. 5** demonstrates that fibronectin promotes integrin activation of LTBP-presented latent TGF β 1. In this assay, SW480/ β 6 cells were bulk transfected with the indicated DNA molecules and seeded on pre-coated assay well with varying concentrations of fibronectin purified from human plasma. Fibronectin increased activation of latent TGF β presented by LTBP1 and LTBP3.

Assay II. Activation of Latent TGF β 1 using LN229 cells

[758] For the assay depicted in **Fig. 3B**, the following protocol was developed. This assay, or “direct-transfection” protocol, is optimal for cell-surface presented TGF β 1 (GARP or LRRC33 presenter) activation by integrin cells. LN229 cells express integrin α V β 8 (as opposed to the SW480b6 cell line, which was engineered to express α V β 6).

[759] These two cell lines enable testing of antibodies on latent TGF β 1 activated by either of the two best validated TGF β -activating integrins.

Materials:

- MvLu1-CAGA12 cells (Clone 4A4)
- LN229 cell line (high levels of endogenous α V β 8 integrin)
- Costar white walled TC treated 96 well assay plate #3903
- Greiner Bio-One High Binding white μ clear 96 well assay plate #655094
- Human Fibronectin (Corning #354008)
- P200 multichannel pipet
- P20, P200, and P1000 pipets with sterile filter tips for each
- Sterile microfuge tubes and rack
- Sterile reagent reservoirs
- 0.4% trypan blue

- 2mL, 5mL, 10mL, and 25mL sterile pipets
- Tissue culture treated 100mm or 150mm plates
- 70%Ethanol
- Opti-MEM reduced serum media (Life Tech #31985-070)
- Lipofectamine 3000 (Life Tech #L3000015)
- Bright-Glo luciferase assay reagent (Promega #E2620)
- 0.25% Trypsin + 0.53mM EDTA
- proTGFb1 expression plasmid, human (SR005)
- LTBP1S expression plasmid, human (SR044)
- LTBP3 expression plasmid, human (SR117)
- LRRC32 (GARP) expression plasmid, human (SR116)
- LRRC33 expression plasmid, human (SR386)

Equipment:

- BioTek Synergy H1 plate reader
- TC hood
- Bench top centrifuge
- CO₂ incubator 37°C 5% CO₂
- 37°C water/bead bath
- Platform shaker
- Microscope
- Hemocytometer/countess

Definitions:

- CAGA12 4A4 cells: Derivative of MvLu1 cells (Mink Lung Epithelial Cells), stably transfected with CAGA12 synthetic promoter, driving luciferase gene expression
- DMEM-0.1%BSA: Assay media; base media is DMEM (Gibco Cat# 11995-065), media also contains BSA diluted to 0.1% w/v, penicillin/streptomycin, and 4mM glutamine
- D10: DMEM 10% FBS, P/S, 4mM glutamine, 1% NEAA, 1X GlutaMAX (Gibco Cat# 35050061)
- CAGA12 (4A4) media: D10 + 0.75ug/mL puromycin

Procedure:

[760] On Day 0, integrin expressing cells were seeded for transfection. Cells were detached with trypsin and pelleted (spin 5 min @ 200 x g). Cell pellet was resuspended in D10 media and count viable cells per ml. Cells were diluted to 0.1×10^6 cells/ml and seeded 100µl per well (10,000 cells per well) in an assay plate. For CAGA12 cells, passaged at a density of 1.5 million per T75 flask, to be used for the assay on Day 2. Cultures were incubated at 37°C and 5% CO₂.

[761] On Day 1, cells were transfected. The manufacturer's protocol was followed for transfection with Lipofectamine 3000 reagent. Briefly, the following was diluted into OptiMEM I, for 5µl per well: 0.1µg DNA (presenting molecule) + 0.1µg DNA (proTGFβ1), 0.4µl P3000, and up to 5µl with OptiMEM I. The well was mixed by pipetting DNA together, then add OptiMEM. Add P3000 and mix everything well by pipetting. A master mix was made with Lipofectamine3000, to be added to DNA mixes: 0.2µl Lipofectamine3000, up to 5µl in OptiMEM I, per well. Diluted Lipofectamine3000 was added to DNA, mixed well by pipetting, and incubated at room temp for 15min. After the incubation, the solution was mixed a few times by pipetting, and then 10µl per well of DNA:Lipofectamine3000 (2 x 5µl) was added. The cell plate was returned to the tissue culture incubator for ~ 24hrs.

[762] On Day 2, the antibody and TGFβ reporter cells were added. In order to prepare functional antibody dilutions, stock antibody in vehicle (PBS is optimal) was serially diluted. Then each point was diluted into assay media for 2X final concentration of antibody. After preparing antibodies, the cell plate was washed twice with assay media, by aspirating (vacuum aspirator) followed by the addition of 100µl per well assay media. After second wash, the assay media was replaced with 50µl per well of 2X antibody. The cell plate was returned to the incubator for ~ 15-20min.

[763] In order to prepare the CAGA12 (clone 4A4) cells for the assay, the cells were detached with trypsin and pelleted (spin 5 min @ 200 x g.). The pellet was resuspended in assay media and viable cells per ml were counted. Cells were diluted to 0.3×10^6 cells/ml and seeded 50µl per well (15,000 cells per well). Cells were returned to incubator.

[764] On Day 3, the assay was read about 16-20 hours after the antibody and/or reporter cell addition. Bright-Glo reagent and test plate were allowed to come to room temperature before reading. The read settings on BioTek Synergy H1 were set to use TMLC_std protocol – this method has an auto-gain setting. Positive control wells were set for autoscale (high). 100µL of Bright-Glo reagent was added per well. Incubated for 2min with shaking, at room temperature, protected plate from light. The plate was read on BioTek Synergy H1.

[765] In some embodiments, TGFβ activity associated with endogenous presenting molecules (*e.g.*, LTBP1/3) may be assessed by only transfecting proTGFβ1 (*i.e.*, without co-transfecting LTBP1/3). In some embodiments, the presenting molecule and proTGFβ DNA may be bulk transfected into LN229 cells in a separate well/dish/plate (*i.e.*, “bulk transfection”) as essentially described above in Assay I, rather than directly transfecting the cells (*i.e.*, “direct transfection”). In some embodiments, LN229 cells may be seeded in assay wells with fibronectin (see Assay I for fibronectin pre-coating protocol).

[766] Data generated from this assay reflects TGFβ1 activity in cell supernatants (**Fig. 3B**). Specifically, LN229 cells were seeded in assay wells without fibronectin and transfected with the indicated DNA molecules by “direct transfection”. Raw data units are relative light units (RLU).

Samples with high RLU values contained high amounts of free TGF β , samples with low RLU values contained low levels of TGF β .

Example 3. SR-AB1 Is A Context-Independent Inhibitor of TGF β LLC Activation by Integrin

[767] **Fig. 6** is a graph demonstrating that SR-AB1 is a context-independent inhibitor of TGF β 1 large latent complex (LLC) by integrin. In this assay, SW480/ β 6 cells were seeded in assay wells without fibronectin and directly transfected with GARP-proTGF β 1 or LRRC33-proTGF β 1, respectively, as essentially described in the above protocols in Example 2. To assess activation of TGF β 1 by LTBP1, LN229 cells were seeded in assay wells pre-coated with fibronectin and directly transfected with LTBP1-proTGF β 1, as essentially described in the above protocols in Example 2. SR-AB1 was shown to inhibit integrin activation of TGF β independent of the presenting molecule.

Example 4. SR-AB2 Is A Complex-Specific Inhibitor of LTBP-proTGF β 1

[768] SR-AB2 was selected for further analysis and testing for specificity for binding to different TGF β presenting molecules. Initially, an ELISA assay was conducted to test complex-specificity as follows.

[769] Materials:

- Solid white 96-well plates from the NeutrAvidin Coating of 96-Well Plates SOP
- Biotinylated antigen
- Maine Biotechnology Services Anti-His antibody MAB230P
- Jackson ImmunoResearch Laboratories Peroxidase Affinipure Goat α -human FC γ
- Fragment Specific. Catalogue number 109-035-008.
- Jackson ImmunoResearch Laboratories Affinipure Goat α -mouse FC γ Fragment Specific.
- Catalogue number 115-035-008
- QuantaBlu ELISA Substrate (Pierce Biotech catalog number 15162)

Equipment:

- Multi-channel pipette
- P200 tips
- Tabletop ultracentrifuge
- 1.5 mL centrifuge tubes
- 0.5 mL centrifuge tubes
- P1000
- P1000 tips
- P200
- P10
- P10 tips

- 15 mL Falcon tubes
- 50 mL Falcon tubes
- Biotek ELx 405 Select CW plate washer
- Multidrop
- Biotek Synergy H1 plate reader

Definitions:

- Wash buffer: TBS (Tris-Buffered Saline; 50 mM Tris-Cl, 150 mM NaCl, pH 7.6) with 0.05% Tween-20. For manual (hand) wash add 0.1% BSA as BSA is sticky and should not be used with the automated plate washer system.
- Sample Buffer: TBS (Tris-Buffered Saline; 50 mM Tris-Cl, 150 mM NaCl, pH 7.6) with 0.05% Tween-20 and 0.1% BSA.
- ELISA 3X protocol: Wash protocol on the Biotek ELx 405 Select CW plate washer. Washes with 200 μ L of wash buffer. Repeats the wash two additional times. Specifications: 3 cycles. No shaking. Dispenses 200 μ L per well. Dispense flow rate setting 7 (range 1 – 10). Dispense height 15.24 mm. Horizontal x dispense position 0mm. Horizontal y dispense position 0 mm. Aspirate height 3.048 mm. horizontal x aspirate position 1.372 mm. horizontal aspirate y position 0.452 mm. Aspiration rate 3.4 mm/second. Aspiration delay 0 milliseconds. Crosswire aspiration on final wash. Crosswire height 3.048 mm. Crosswire horizontal x position: -1.829 mm. Crosswire horizontal y position: -0.457 mm.

Procedure:

[770] Remove a pre-coated and pre-blocked 96-well plate from 4°C. Dump out the 1xPBS pH 7.4 1%BSA with 0.1% Tween-20 from the plate and forcefully hit the plate on a Styrofoam pad lined with paper towels. If a 96-well plate is not already prepared, prepare one with the NeutrAvidin Coating of 96-Well Plates protocol. Specifically, for one NeutrAvidin coated plate, remove 5 μ L from the top of the 1 mg/mL NeutrAvidin stock solution and dilute it into 10 mL of 1x carbonate buffer pH 9.4. Mix by inverting the falcon tube. Using a multi-channel pipette, place 100 μ L in each well of the Corning high binding 96-well assay plate and incubate the 96-well plate overnight at 4°C. Wash the 96-well plate with 200 μ L of wash buffer per well and repeat this step two additional times. The plate should be washed for a total of three times. Block the plate with 200 μ L per well of 1% BSA in PBS pH 7.4 and incubate the plate for 1 hour at 37°C or overnight at 4°C.

[771] Dilute biotinylated antigen in sample buffer. Optimal capture concentration should first be determined by titration for each individual protein. Add 50 μ L per well and incubate at room temperature for 1 hour.

[772] Wash plates using the plate washer with wash buffer using the ELISA 3X protocol.

[773] Dilute antibody in sample buffer. Screening of antibodies is performed at 1 μ g/mL. Prepare α -His coating control antibody (Maine Biotechnology Services catalogue number: MAB230P) at 1

μg/mL in sample buffer. Place 50 μL of diluted antibody on designated wells and incubate at room temperature for 1 hour.

[774] Wash plates using the plate washer with wash buffer using the ELISA 3X protocol.

[775] Dilute human secondary antibody (Jackson ImmunoResearch Laboratories Peroxidase

[776] Affinipure Goat α-human FCγ Fragment Specific. Catalogue number 109-035-008) 1:10,000 in sample buffer. For α-his wells, dilute mouse secondary antibody (Jackson ImmunoResearch Laboratories Affinipure Goat α-mouse FCγ Fragment Specific. Catalogue number 115-035-008) 1:10,000 in sample buffer.

[777] Place 50 μL of diluted antibody on designated wells. Incubate at room temperature for 1 hour. Wash plates using the plate washer with wash buffer using the ELISA 3X protocol.

[778] Prepare SuperSignal ELISA Femto Substrate (Pierce Biotech catalog number 15162) working solution according to manufacturer's protocol. 10 mL will be needed for one plate. Place 100 μL QuantBlu Substrate working solution per well. Incubate for 10 minutes at room temperature.

[779] Measure relative fluorescent units (RFU's) on a plate reader with an excitation of 325nm and emission of 420nm.

[780] Results:

[781] **Fig. 7A** demonstrates that SR-AB2 only binds LTBP-proTGFβ1 complex; it does not bind proTGFβ1 or LTBP1 alone by ELISA. **Fig. 7A** demonstrates that SR-AB2 does not bind GARP-proTGFβ1 by ELISA.

[782] Additionally, the ability of SR-AB2 to inhibit LTBP1/3-proTGFβ1 in a cell-based assay was also conducted. As essentially described above in Example 2, LN229 cells were seeded on fibronectin pre-coated assay wells and directly transfected with the indicated DNA molecules. **Fig. 7B** depicts that SR-AB2 inhibits integrin activation of LTBP1-proTGFβ1 (human and mouse complexes). **Fig. 7C** depicts that SR-AB2 inhibits integrin activation of LTBP3-proTGFβ1.

[783] It was demonstrated that SR-AB2 specifically binds to proTGFβ1:LTBP1 & 3 complexes, and not GARP-TGFβ1 or GARP-Lap complexes by ELISA as essentially described above (see **Fig. 9**).

[784] As discussed above, LTBP1 and LTBP3 are deposited in the extracellular matrix, while GARP/LRRC32 and LRRC33 present latent TGFβ1 on the surface of immune cells. It was demonstrated that SR-AB2 inhibits LTBP-proTGFβ1 signaling, but does not affect GARP-proTGFβ1 (**Fig. 10A and Fig. 10B**). **Fig. 10A** demonstrates that SR-AB2 inhibits LTBP-proTGFβ presented by endogenous LTBP1/3. This assay was performed in LN229 cells, which were seeded on fibronectin pre-coated wells and directly transfected with proTGFβ1. **Fig. 10B** demonstrates that SR-AB2 does not inhibit GARP-proTGFβ. SR-AB1 binds latent TGFβ1 independent of the presenting molecule. This assay was performed in LN229 cells, were seeded in assay wells without fibronectin and directly transfected with GARP-proTGFβ1.

[785] Inhibition of LTBP-proTGF β 1 by SR-AB2 was also shown in α V β 6 Integrin-dependent activation of LTBP1-presented TGF β 1 in cell-based assays (human and mouse, data not shown). SR-AB2 showed no inhibitory effects on overexpressed LRRC33-proTGF β 1.

Example 5: Octet binning of LTBP1-proTGF β 1 antibodies

[786] 500 nM of human LTBP1-proTGF β 1 complex was pre-incubated with 1 μ M of each test antibody. After an overnight incubation, the LTBP1-proTGF β 1+first antibody was tested for binding to a second antibody which was immobilized to an Anti-Human IgG Fc Capture (AHC) sensor tip at 67 nM. The sensor tip was blocked with a negative control antibody (HuNeg) before seeing the LTBP1-proTGF β 1+first antibody.

[787] Binding of the complex to a specific second antibody was normalized to the uninhibited interaction, that is the complex in the presence of a negative control antibody (HuNeg) that does not bind to the TGF β complex. Normalized responses less than 70% or less than 0.7 of the uninhibited interaction were considered antibodies that cross block. A response greater than 1 indicated that both antibodies were bound simultaneously. The results are shown in Table 7, below.

Table 7:

	SR-AB13	SR-AB10	SR-AB2	SR-AB1	Second Antibody
SR-AB13	0.47	1.38	1.29	1.08	
SR-AB10	1.27	0.51	1.53	0.82	
SR-AB2	1.31	1.54	0.47	1.11	
SR-AB1	1.43	0.76	1.38	0.69	
HuNeg	1	1	1	1	
First Antibody					

[788] As shown in Table 7, antibodies SR-AB13, SR-AB10, SR-AB2 and SR-AB1 do not cross block each other, and therefore each antibody occupies a distinct epitope on the surface of human LTBP1-proTGF β 1.

Example 6: In vitro Binding Profile and Affinity Data

[789] Suitable methods for in vitro binding assays to determine the parameters of binding kinetics include Bio-layer Interferometry (BLI)-based assays such as Octet, and surface plasmon resonance (SPR)-based assays, such as Biacore systems (see, for example:

http://www.biophysics.bioc.cam.ac.uk/wp-content/uploads/2011/02/Biacore_assay_handbook.pdf).

[790] The affinity of SR-AB10, SR-AB2 and SR-AB13 was measured by Octet assay. The protocol used to measure the affinity of the antibodies to the complexes provided herein is summarized below.

Materials:

- 96 well black polypropylene plates
- AHC Octet tips (anti-human IgG Fc capture tips) (FortéBio)
- 10x kinetics buffer (FortéBio) (diluted to 1x in PBS)
- reiner Bio-One 96-Well Half Area Microplates (VWR cat# 82050-044)

Procedure:

Notes: The volume within each well during an Octet experiment is 100 uL.

A shake speed of 1000 rpm is used for all assay steps.

Pre-Wet Tips:

- Biosensors must be pre-wet in 1x kinetics buffer (1x KB) for at least 10 minutes before starting an experiment. This can be done inside the Octet or on the bench.

Loading:

- Tips are baselined in 1x KB for 1 minute before loading.
- AHC tips are loaded with antibody at a concentration of 1 µg/mL (~7 nM) for 3 minutes. A limit is set so that loading will stop when any one sensor reaches a response of 1 nanometer.
- Tips are baselined in buffer for 1 minute after loading. The antibody should not dissociate from the tips during this time.

Antigen Association:

- TGFβ1 in complex with various presentation molecules was associated to the immobilized antibodies at a single concentration of 100 nM in 1x KB.

Antibody Dissociation:

- Dissociation in 1x KB was performed for 3 minutes.

Data Analysis using ForteBio Data analysis software 8.2:

- Processing: align Y axis to last 5 seconds of baseline, perform inter-step correction with align to dissociation, and perform Savitzky-Golay filtering.
- * Analysis: 1:1 fitting model is utilized. Fitting is local and full. (Local indicates each antibody is evaluated separately and full indicates that both association and dissociation are considered) Fit the curves and then save the report/export the data.

Results:

[791] **Fig. 11** presents the binding profile and affinity data for LTBP complex-specific antibodies SR-AB10, SR-AB2, and SR-AB13. Notably, SR-AB13 binds both human LTBP1 and human LTBP3 complexed with proTGFβ1, while SR-AB2 and SR-AB10 are specific to human LTBP1 complexed with TGFβ1.

Example 7: Improved potency of optimized LTBP-complex-specific antibodies

[792] LTBP complex-specific antibodies SR-AB10 and SR-AB13 were selected for an initial round of affinity maturation/optimization (*i.e.*, H1/H2 CDR shuffling/diversification as described herein) and particular progeny antibodies (SR-AB14 and SR-AB15) were assessed for their ability to inhibit TGF β activity using LN229 cells. For the assays depicted in **Fig. 12A and 12B**, the following protocol was used, which is a modified version of Assay II in Example 2.

Materials:

- MvLu1-CAGA12 cells (Clone 4A4)
- LN229 cell line (high levels of endogenous α V β 8 integrin)
- Costar white walled TC treated 96 well assay plate #3903
- Greiner Bio-One High Binding white μ clear 96 well assay plate #655094
- Human Fibronectin (Corning #354008)
- P200 multichannel pipet
- P20, P200, and P1000 pipets with sterile filter tips for each
- Sterile microfuge tubes and rack
- Sterile reagent reservoirs
- 0.4% trypan blue
- 2mL, 5mL, 10mL, and 25mL sterile pipets
- Tissue culture treated 100mm or 150mm plates
- 70%Ethanol
- Opti-MEM reduced serum media (Life Tech #31985-070)
- Lipofectamine 3000 (Life Tech #L3000015)
- Bright-Glo luciferase assay reagent (Promega #E2620)
- 0.25% Trypsin + 0.53mM EDTA
- proTGF β 1 expression plasmid, human
- LTBP1S expression plasmid, human

Equipment:

- PerkinElmer EnVision plate reader
- TC hood
- Bench top centrifuge
- CO2 incubator 37°C 5% CO2
- 37°C water/bead bath
- Platform shaker
- Microscope
- Hemocytometer/countess

Definitions:

- CAGA12 4A4 cells: Derivative of MvLu1 cells (Mink Lung Epithelial Cells), stably transfected with CAGA12 synthetic promoter, driving luciferase gene expression
- DMEM-0.1%BSA: Assay media; base media is DMEM (Gibco Cat# 11995-065), media also contains BSA diluted to 0.1% w/v, penicillin/streptomycin, and 4mM glutamine
- D10: DMEM 10% FBS, P/S, 4mM glutamine, 1% NEAA, 1X GlutaMAX (Gibco Cat# 35050061)
- CAGA12 (4A4) media: D10 + 0.75ug/mL puromycin

Procedure:

- Always work in sterile biosafety cabinet, and sterile technique should be used at all times.
- Prepare all media, sterilize all materials, and move materials into biosafety cabinet before starting.
- Growth medium should be warmed to 37°C.
- Use the reverse pipetting technique for almost all steps during this assay.
- Assays that rely on LTBP presentation of proTGFβ1 require pre-coating of assay plates with fibronectin, for 4hrs – overnight prior to seeding cells for transfection.

[793] On day -1, assay plates were coated with Fibronectin. A 1mg/ml stock solution of fibronectin was prepared in ultrapure water (sterile). Stock solution was diluted in PBS to 19.2μg/ml working solution and 50 μl was added to each well (3 μg/cm²). Plates were incubated at 37°C and 5% CO₂ overnight.

[794] On day 0, prior to cell seeding, assay plates were coated with Fibronectin (LTBP over expression assay only). A 1mg/ml stock solution of fibronectin was prepared in ultrapure water (sterile). Stock solution was diluted in PBS to 19.2μg/ml working solution and 50 μl was added to each well (3 μg/cm²). Plates were incubated at 37°C and 5% CO₂ for 4 hours. Following coating, assay plates were washed manually with multichannel pipette for 2 washes of 200 μl/well PBS. After final wash, plates were allowed to dry in the hood with lid off. LN229 cells were then detached with trypsin and pelleted (spun for 5 min at 200 x g). The cell pellet was resuspended in D10 media and viable cells per ml counted. Cells were diluted to 0.125e⁶ cells/ml and seeded 100μl per well (12,500 cells per well) in an assay plate. For CAGA12 cells to be used for assay on Day 2, cells passaged at a density of 1.5 million per T75 flask. Cultures were incubated at 37°C and 5% CO₂.

[795] On Day 1, LN229 cells were transfected. The manufacturer's protocol was followed for transfection with Lipofectamine 3000 reagent. Briefly, the following was diluted into OptiMEM I, for 5μl per well: 0.1μg DNA (proTGFβ1), optionally 0.1μg LTBP1 DNA, 0.4μl P3000, and up to 5μl with OptiMEM I. For **Fig. 12A**, 0.1μg proTGFβ1 (human) DNA was transfected alone, without LTBP DNA, to measure activation in the presence of endogenous presenting molecules. For **Fig. 12B**,

0.1µg LTBP1 (human) DNA was co-transfected with the proTGFβ1 (human) DNA to measure activation in the presence of overexpressed LTBP1.

[796] A master mix of Lipofectamine3000 was made by diluting 0.2µl Lipofectamine3000 in OptiMEM I, up to 5µl in OptiMEM I, per well. Diluted Lipofectamine3000 was added to the DNA mixture, mixed well by pipetting, and incubated at room temperature for 15min. After the incubation, 10µl of DNA:Lipofectamine3000 (2 x 5µl) mixture was added to each well. Plates were returned to the tissue culture incubator for ~24hrs.

[797] On day 2 the indicated antibodies and TGFβ reporter cells (CAGA12) were added to the wells. Antibodies were serially diluted into PBS, then further diluted into assay media until 2x final concentration. Plates were washed twice with assay media by aspirating (vacuum aspirator) followed by addition of 100µl per well assay media. After second wash, assay media was replaced with 50 µl per well of 2X antibody. Cell plate was returned to the incubator for ~ 15-20 min. CAGA12 (clone 4A4) cells were detached with trypsin and pelleted (spun 5 min at 200 x g.). Pellet was resuspended in assay media and viable cells counted. Viable cells were diluted to 0.3e⁶cells/ml and seeded 50µl per well (15,000 cells per well). Cells were returned to the incubator.

[798] On Day 3, the assay was read about 16-20 hours after the antibody and/or reporter cell addition. Bright-Glo reagent and test plate were allowed to come to room temperature before reading. The read settings on BioTek Synergy H1 were set to use TMLC_std protocol – this method has an auto-gain setting. Positive control wells were set for autoscale (high). 100µL of Bright-Glo reagent was added per well. Incubated for 2min with shaking, at room temperature, protected plate from light. The luminescence was then detected on a plate reader.

Results:

[799] Data generated from this assay reflected TGFβ activity in cell supernatants. **Fig. 12A** is a graph showing improved potency of SR-AB14 (optimized SR-AB10) as measured by TGFβ activity. Notably, SR-AB14 activity is similar to the context-independent antibody SR-AB1. This assay was performed without overexpressing LTBP1 and thus measures activation of TGFβ in the presence of endogenous presenting molecules. **Fig. 12B** is a graph showing improved potency of SR-AB15 (optimized SR-AB13) as measured by TGFβ activity. This assay was performed in the presence of overexpressed human LTBP1-pro TGFβ1. Notably, SR-AB14 activity is similar to the context-independent antibody SR-AB1.

[800] The assays depicted in **Figs. 12A** and **12B** were both performed in LN229 cells, which express low LTBP1 mRNA, high LTBP3 mRNA, undetectable GARP, and undetectable LRRC33.

Example 8: Improved affinity of optimized LTBP-proTGFβ1-specific antibodies after CDR-H3 mutagenesis

[801] LTBP complex-specific antibodies from the first round of affinity maturation/optimization (SR-AB14, SR-AB16, SR-AB17, SR-AB18, SR-AB19) were selected for a second round of affinity

maturation/optimization (*i.e.*, CDR-H3 mutagenesis as described herein) and particular progeny antibodies (SR-AB20, SR-AB21, SR-AB22, SR-AB23, SR-AB24, SR-AB25, SR-AB26, SR-AB27, SR-AB28, and SR-AB29) were assessed for their ability to bind various proTGFβ1 constructs. Binding affinities of the optimized antibodies were measured by Octet against human LTBP1-proTGFβ1, human LTBP3-proTGFβ1, mouse LTBP1-proTGFβ1, mouse LTBP3-proTGFβ1, and GARP-proTGFβ1 complexes, essentially as described in Example 6. In brief, test antibodies were immobilized to the surface of anti-human Fc capture biosensors (AHC) (ForteBio®) and binding was then tested against the various TGFβ1 complexes at a single concentration of 100nM to assess binding affinities. The antigens were allowed to associate for 3 minutes followed by a 5-minute dissociation. Kinetics buffer (ForteBio®) was used throughout the experiment, and K_D was determined using a 1:1 fitting model for each antibody antigen pair.

[802] Table 8 presents the binding profile and affinity data for the indicated LTBP complex-specific antibodies (IgG1-agly) determined by Octet. Notably, optimized antibodies SR-AB20, SR-AB21, SR-AB22, SR-AB23, SR-AB24, SR-AB25, SR-AB26, SR-AB27, SR-AB28, and SR-AB29 display single digit nM affinity for both human LTBP1 and human LTBP3 complexed with proTGFβ1. Additionally, none of the antibodies bound human GARP complexed with proTGFβ1 indicating that the antibodies are specific for LTBP complexes.

Table 8:

Ref	Lineage	ForteBio IgG K_D Hu LTBP1- proTGFβ1 (M) Avid	ForteBio IgG K_D Hu LTBP3- proTGFβ1 (M) Avid	ForteBio IgG K_D Mo LTBP1- proTGFβ1 (M) Avid	ForteBio IgG K_D Mo LTBP3- proTGFβ1 (M) Avid	ForteBio IgG K_D Hu GARP- proTGFβ1 (M) Avid
SR-AB10		P.F.	N.B.	1.42E-08	2.30E-08	N.B.
SR-AB16	SR-AB10	3.44E-08	1.69E-08	3.29E-08	2.12E-08	N.B.
SR-AB14	SR-AB10	2.44E-08	1.46E-08	2.04E-08	1.23E-08	N.B.
SR-AB20	SR-AB16	3.93E-09	2.08E-09	6.46E-09	2.23E-09	N.B.
SR-AB21	SR-AB16	6.78E-09	1.08E-09	1.06E-08	3.29E-08	N.B.
SR-AB22	SR-AB14	1.88E-09	P.F.	2.43E-09	1.56E-09	P.F.
SR-AB23	SR-AB14	5.98E-09	1.26E-09	8.38E-09	7.93E-09	N.B.
SR-AB13		3.92E-08	3.59E-08	5.95E-08	N.B.	N.B.
SR-AB17	SR-AB13	6.16E-09	5.28E-09	8.83E-09	5.36E-08	N.B.
SR-AB18	SR-AB13	9.27E-09	1.17E-08	1.45E-08	8.66E-08	N.B.
SR-AB19	SR-AB13	1.10E-08	1.64E-08	1.59E-08	8.40E-08	N.B.
SR-AB24	SR-AB17	1.74E-09	1.27E-09	2.07E-09	7.85E-08	N.B.
SR-AB25	SR-AB17	2.37E-09	1.05E-09	2.92E-09	3.88E-08	N.B.
SR-AB26	SR-AB17	2.73E-09	2.13E-09	3.73E-09	1.38E-08	N.B.
SR-AB27	SR-AB18	2.70E-09	2.33E-09	3.89E-09	5.11E-07	N.B.

SR-AB28	SR-AB18	2.84E-09	2.70E-09	4.52E-09	1.25E-08	N.B.
SR-AB29	SR-AB19	4.23E-09	4.08E-09	7.22E-09	7.91E-08	N.B.

P.F. = Poor Fit

N.B. = Non-binder under conditions of this assay

Example 9: Improved affinity of optimized LTBP-complex-specific antibodies after light chain optimization cycle 3

[803] LTBP complex-specific antibodies from the second round of affinity maturation/optimization were selected for a third round of affinity maturation/optimization (*i.e.*, light chain optimization as described herein) and particular progeny antibodies, were assessed for their ability to bind various proTGF β 1 constructs.

[804] Cycle 3 antibodies were generated by performing mutagenesis throughout the light chain CDR3 and performing a shuffle of premade sequences for the light chain CDR1 and CDR2. Antibodies of interest were identified through yeast display utilizing both positive and negative selections followed by sequencing.

[805] Binding affinities of the optimized antibodies were measured by Octet against human LTBP1-proTGF β 1, human LTBP3-proTGF β 1, mouse LTBP1-proTGF β 1, mouse LTBP3-proTGF β 1, and GARP-proTGF β 1 complexes, essentially as described in Example 6. In brief, test antibodies were immobilized to the surface of anti-human Fc capture biosensors (AHC) (ForteBio®) and binding was then tested against the various TGF β 1 complexes at a single concentration of 100nM to assess binding affinities. The antigens were allowed to associate for 3 minutes followed by a 5-minute dissociation. Kinetics buffer (ForteBio®) was used throughout the experiment, and KD was determined using a 1:1 fitting model for each antibody antigen pair.

[806] Table 9 presents the binding profile and affinity data for the indicated LTBP complex-specific antibodies (IgG1-agly) identified from the light chain optimization cycle 3, determined by Octet. Notably, several optimized antibodies display single digit nM affinity for both human LTBP1 and human LTBP3 complexed proTGF β 1. Additionally, as shown in Table 10, several antibodies did not bind human GARP complexed with proTGF β 1, under the same assay conditions, indicating that the antibodies are specific for LTBP complexes.

Table 9: Binding Profile and Affinity Data for Cycle 3 Optimized Antibodies (N.B. = non-binder under conditions of this assay)

Ref	Optimization Lineage	ForteBio IgG K _D Hu LTBp1-proTGFB1 (M) Avid	Hu LTBp1-proTGFB1 Kon (1/Ms)	Hu LTBp1-proTGFB1 Koff (1/s)	ForteBio IgG K _D Hu LTBp3-proTGFB1 (M) Avid	Hu LTBp3-proTGFB1 Kon (1/Ms)	Hu LTBp3-proTGFB1 Koff (1/s)	ForteBio IgG K _D Mo LTBp1-proTGFB1 (M) Avid	ForteBio IgG K _D Mo LTBp3-proTGFB1 (M) Avid	ForteBio IgG K _D Hu GARP-proTGFB1 (M) Avid
SR-AB30	SR-AB22	2.17E-10	2.77E+05	6.00E-05	1.56E-10	3.85E+05	6.00E-05	4.38E-10	4.21E-10	1.09E-08
SR-AB31	SR-AB22	4.12E-10	2.40E+05	9.89E-05	2.21E-10	3.08E+05	6.81E-05	6.82E-10	5.89E-10	1.12E-08
SR-AB32	SR-AB22	3.22E-10	2.83E+05	9.14E-05	1.49E-10	4.02E+05	6.00E-05	4.85E-10	4.18E-10	1.09E-08
SR-AB33	SR-AB22	5.61E-10	3.02E+05	1.69E-04	2.72E-10	4.45E+05	1.21E-04	8.67E-10	4.78E-10	1.61E-08
SR-AB34	SR-AB22	2.31E-10	2.59E+05	6.00E-05	1.79E-10	3.72E+05	6.68E-05	4.17E-10	4.50E-10	7.35E-08
SR-AB35	SR-AB22	3.15E-10	3.42E+05	1.08E-04	1.41E-10	4.85E+05	6.85E-05	3.89E-10	3.98E-10	6.66E-09
SR-AB36	SR-AB22	2.54E-10	2.89E+05	7.34E-05	1.46E-10	4.10E+05	6.00E-05	3.35E-10	4.29E-10	9.55E-09
SR-AB37	SR-AB22	3.28E-10	3.60E+05	1.18E-04	1.39E-10	5.45E+05	7.58E-05	3.84E-10	3.25E-10	3.61E-08
SR-AB38	SR-AB22	3.57E-10	3.27E+05	1.17E-04	1.45E-10	5.02E+05	7.26E-05	3.91E-10	3.31E-10	8.42E-09
SR-AB39	SR-AB22	2.05E-10	2.92E+05	6.00E-05	1.36E-10	4.41E+05	6.00E-05	2.32E-10	3.42E-10	3.89E-07
SR-AB40	SR-AB23	4.01E-10	1.50E+05	6.00E-05	3.33E-10	1.82E+05	6.07E-05	1.08E-09	1.16E-09	N.B.
SR-AB41	SR-AB23	2.96E-10	2.03E+05	6.00E-05	2.35E-10	2.55E+05	6.00E-05	4.04E-10	7.17E-10	7.64E-07
SR-AB42	SR-AB23	3.89E-10	1.54E+05	6.00E-05	2.59E-10	2.32E+05	6.00E-05	3.59E-10	1.05E-09	N.B.
SR-AB43	SR-AB23	6.49E-10	2.44E+05	1.59E-04	1.78E-10	3.37E+05	6.00E-05	7.31E-10	8.91E-10	N.B.
SR-AB44	SR-AB23	2.14E-10	2.80E+05	6.00E-05	1.46E-10	4.10E+05	6.00E-05	2.10E-10	4.07E-10	2.12E-07
SR-AB45	SR-AB23	1.93E-10	3.10E+05	6.00E-05	1.22E-10	4.91E+05	6.00E-05	1.93E-10	2.69E-10	1.95E-07
SR-AB62	SR-AB24	4.86E-10	1.40E+05	6.82E-05	3.68E-10	2.03E+05	7.47E-05	6.09E-10	3.37E-09	N.B.
SR-AB63	SR-AB26	6.11E-10	1.18E+05	7.24E-05	4.94E-10	1.59E+05	7.84E-05	1.09E-09	2.00E-09	N.B.
SR-AB64	SR-AB26	4.67E-10			4.71E-10	1.95E+05	9.18E-05	1.03E-09	2.85E-09	N.B.
Human IgG1 isotype control	N.A.	N.B.			N.B.			N.B.	N.B.	N.B.

Table 10:

Ref	VHCDR3 Lineage	ForteBio IgG K _D Hu GARP-proTGFβ1 in solution (M) Avid
SR-AB30	SR-AB10	1.09E-08
SR-AB31	SR-AB10	1.12E-08
SR-AB32	SR-AB10	1.09E-08
SR-AB33	SR-AB10	1.61E-08
SR-AB34	SR-AB10	7.35E-08
SR-AB35	SR-AB10	6.66E-09
SR-AB36	SR-AB10	9.55E-09
SR-AB37	SR-AB10	3.61E-08
SR-AB38	SR-AB10	8.42E-09
SR-AB39	SR-AB10	3.89E-07
SR-AB40	SR-AB10	N.B.
SR-AB41	SR-AB10	7.63909E-07
SR-AB42	SR-AB10	N.B.
SR-AB43	SR-AB10	N.B.
SR-AB44	SR-AB10	2.1153E-07
SR-AB45	SR-AB10	1.95124E-07
SR-AB62	SR-AB13	N.B.
SR-AB63	SR-AB13	N.B.
SR-AB64	SR-AB13	N.B.
Human IgG1 isotype control	N.A.	N.B.

N.B. = non-binder under conditions of this assay

[807] **Fig. 15** shows that affinity matured antibodies show specific binding to the LTBP-proTGFβ1 complex. This experiment was performed at 200, 100, 50, and 25 nM human GARP proTGFβ1 and human LTBP1 proTGFβ1. **Fig. 15** shows the 200 nM values, where the 0.1nm cutoff is used to determine what is and what is not meaningful binding. As shown in **Fig. 15**, for some antibodies, binding to GARP-proTGFβ1 is meaningful only at very high concentrations (200nM), and some antibodies show no binding to GARP-proTGFβ1 even at that high concentration.

Surface plasmon resonance (SPR)-based assays

[808] A Biacore 8K system was employed to determine the monovalent binding affinity and the kinetic parameters for antigen binding of test antibodies. Association and dissociation kinetics of Fab fragments SR-AB42-HuFab, SR-AB63-HuFab and SR-AB43-HuFab to antigen complexes were measured, and resulting k_a , k_d and K_D are provided below. Briefly, the binding kinetics were evaluated by surface plasmon resonance using Biacore 8K (GE Healthcare). Biotinylated capture antigens were immobilized to the chip (10nM, ~200 RU loading). A Biotin CAPture sensor chip was used to capture the biotinylated antigens. Fabs at 10, 5, 2.5, 1.25, and 0.6nM concentrations were injected over the captured antigens. 0 nM was used as a reference. Affinities of LTBP antibodies to

GARP and LRRC33 were confirmed using higher concentration of Fabs (100, 50, 25, 12.5, 6.25 nM). 0 nM was used as a reference. Multi-cycle kinetics was employed where each analyte concentration was injected in a separate cycle and the sensor chip surface was regenerated after each cycle. All the assays were carried out in freshly prepared 1x HBS-EP+ buffer (10 mM HEPES, 150 mM NaCl, 3 mM EDTA, 0.05% Tween20, pH 7.4). Data were fit globally to a 1:1 binding model to obtain the kinetic parameters. The sensorgram for 0 nM analyte concentration was used as reference.

[809] Results are shown in Tables 11-15 below. **Fig. 23** and **Fig. 24**, respectively, show that SR-AB63 and SR-AB42 human Fabs show context-selective binding of novel antibodies for huLTBP1 proTGFβ1 and huLTBP3 proTGFβ1.

Table 11

	huLTBP1 proTGFβ1		KD (M)
	ka (1/Ms)	kd (1/s)	
SR-AB42-HuFab*	5.43E+05	1.05E-04	1.94E-10
SR-AB63-HuFab**	3.71E+05	3.82E-08	1.03E-13
SR-AB43-HuFab	1.03E+06	2.84E-04	2.76E-10
SR-AB46-HuFab	2.47E+06	1.10E-04	4.44E-11
SR-AB47-HuFab	1.84E+06	1.36E-04	7.40E-11
SR-AB48-HuFab	1.76E+06	1.04E-04	5.93E-11
SR-AB49-HuFab	8.09E+05	1.38E-04	1.70E-10
SR-AB50-HuFab	3.31E+06	1.82E-04	5.50E-11
SR-AB51-HuFab	2.55E+06	1.50E-04	5.89E-11
SR-AB52-HuFab	2.50E+06	9.91E-05	3.96E-11
SR-AB53-HuFab	1.82E+06	1.28E-04	7.07E-11
SR-AB54-HuFab	1.80E+06	8.83E-05	4.91E-11
SR-AB55-HuFab	8.62E+05	1.13E-04	1.31E-10
SR-AB56-HuFab	2.58E+06	1.26E-04	4.89E-11
SR-AB57-HuFab	1.35E+06	1.81E-04	1.33E-10
SR-AB58-HuFab	3.11E+06	1.62E-04	5.21E-11
SR-AB59-HuFab	2.55E+06	1.30E-04	5.09E-11
SR-AB60-HuFab	2.35E+06	1.20E-04	5.10E-11
SR-AB61-HuFab	1.17E+06	1.55E-04	1.32E-10

*Average of five replicate experiments

**Average of four replicate experiments

Table 12

	huLTBP3 proTGFβ1		KD (M)
	ka (1/Ms)	kd (1/s)	
SR-AB42-HuFab	4.05E+05	1.42E-04	3.50E-10
SR-AB63-HuFab	3.94E+05	4.99E-05	1.27E-10

Table 13

	huGARP proTGFb1		
	ka (1/Ms)	kd (1/s)	KD (M)
SR-AB42-HuFab	1.55E+05	6.04E-03	3.89E-08
SR-AB63-HuFab	2.66E+03	9.30E-04	3.49E-07

Table 14

	huLRRC33 proTGFb1		
	ka (1/Ms)	kd (1/s)	KD (M)
SR-AB42-HuFab	8.09E+04	3.53E-03	4.36E-08
SR-AB63-HuFab	5.00E+03	1.08E-03	2.16E-07

Table 15

	Murine LTBP3 proTGFb1		
	ka (1/Ms)	kd (1/s)	KD (M)
SR-AB42-HuFab*	1.23E+08	3.97E-01	2.66E-09
SR-AB43-HuFab*	1.82E+09	6.37E+00	4.43E-09

*Average of two replicate experiments

Example 10: Improved potency of optimized LTBP complex-specific antibodies after CDR-H3 mutagenesis

[810] LTBP complex-specific antibodies from the first round of affinity maturation/optimization (*i.e.*, SR-AB14, SR-AB16, SR-AB17, SR-AB18, SR-AB19) were selected for a second round of affinity maturation/optimization (*i.e.*, CDR-H3 mutagenesis as described herein) and particular progeny antibodies (SR-AB20, SR-AB21, SR-AB22, SR-AB23, SR-AB24, SR-AB25, SR-AB26, SR-AB27, SR-AB28, and SR-AB29) were assessed for their ability to inhibit TGF β activity using LN229 cells. For the assays depicted in **Fig. 13A, 13B, 14A, and 14B** the following protocol was used, which is a modified version of Assay II in Example 2.

Materials:

- MvLu1-CAGA12 cells (Clone 4A4)
- LN229 cell line (high levels of endogenous α V β 8 integrin)
- Costar white walled TC treated 96 well assay plate #3903
- Greiner Bio-One High Binding white uclear 96 well assay plate #655094
- Human Fibronectin (Corning #354008)
- P200 multichannel pipet
- P20, P200, and P1000 pipets with sterile filter tips for each
- Sterile microfuge tubes and rack
- Sterile reagent reservoirs

- 0.4% trypan blue
- 2mL, 5mL, 10mL, and 25mL sterile pipets
- Tissue culture treated 100mm or 150mm plates
- 70%Ethanol
- Opti-MEM reduced serum media (Life Tech #31985-070)
- Lipofectamine 3000 (Life Tech #L3000015)
- Bright-Glo luciferase assay reagent (Promega #E2620)
- 0.25% Trypsin + 0.53mM EDTA
- proTGFb1 expression plasmid, human
- proTGFb1 expression plasmid, mouse
- LTBP1S expression plasmid, mouse

Equipment:

- PerkinElmer EnVision plate reader
- TC hood
- Bench top centrifuge
- CO2 incubator 37C 5% CO2
- 37C water/bead bath
- Platform shaker
- Microscope
- Hemocytometer/countess

Definitions:

- CAGA12 4A4 cells: Derivative of MvLu1 cells (Mink Lung Epithelial Cells), stably transfected with CAGA12 synthetic promoter, driving luciferase gene expression
- DMEM-0.1%BSA: Assay media; base media is DMEM (Gibco Cat# 11995-065), media also contains BSA diluted to 0.1% w/v, penicillin/streptomycin, and 4mM glutamine
- D10: DMEM 10% FBS, P/S, 4mM glutamine, 1% NEAA, 1X GlutaMAX (Gibco Cat# 35050061)
- CAGA12 (4A4) media: D10 + 0.75ug/mL puromycin

Procedure:

- Always work in sterile biosafety cabinet, and sterile technique should be used at all times.
- Prepare all media, sterilize all materials, and move materials into biosafety cabinet before starting. Growth medium should be warmed to 37°C.
- Use the reverse pipetting technique for almost all steps during this assay.

[811] On day 0, prior to cell seeding, assay plates were coated with Fibronectin (LTBP over expression assay only). A 1mg/ml stock solution of fibronectin was prepared in ultrapure water

(sterile). Stock solution was diluted in PBS to 19.2ug/ml working solution and 50 μ l was added to each well (3 ug/cm²). Plates were incubated at 37°C and 5% CO₂ for 4 hours. Following coating, assay plates were washed manually with multichannel pipette for 2 washes of 200 ul/well PBS. After final wash, plates were allowed to dry in the hood with lid off. LN229 cells were then detached with trypsin and pelleted (spun for 5 min at 200 x g). The cell pellet was resuspended in D10 media and viable cells per ml counted. Cells were diluted to 0.125e⁶ cells/ml and seeded 100 μ l per well (12,500 cells per well) in an assay plate. For CAGA12 cells to be used for assay on Day 2, cells passaged at a density of 1.5 million per T75 flask. Cultures were incubated at 37°C and 5% CO₂.

[812] On Day 1, LN229 cells were transfected. The manufacturer's protocol was followed for transfection with Lipofectamine 3000 reagent. Briefly, the following was diluted into OptiMEM I, for 5 μ l per well: 0.1 μ g proTGF β 1 DNA combined with 0.1 μ g presentation molecule DNA, and 0.4 μ l P3000, up to 5 μ l with OptiMEM I. For **Fig. 13A and Fig. 14A**, 0.1 μ g human proTGF β 1 DNA was transfected with 0.1ug empty vector, without LTBP DNA, to measure activation in the presence of endogenous presenting molecules. For **Fig. 13B and Fig. 14B**, 0.1 μ g mouse LTBP1 DNA was co-transfected with the mouse proTGF β 1 DNA to measure activation in the presence of overexpressed LTBP1.

[813] A master mix of Lipofectamine3000 was made by diluting 0.2 μ l Lipofectamine3000 in OptiMEM I, up to 5 μ l in OptiMEM I, per well. Diluted Lipofectamine3000 was added to the DNA mixture, mixed well by pipetting, and incubated at room temperature for 15min. After the incubation, 100 μ l of DNA:Lipofectamine3000 (2 x 5 μ l) mixture was added to each well. Plates were returned to the tissue culture incubator for ~24hrs.

[814] On day 2 the indicated antibodies and TGF β reporter cells (CAGA12) were added to the wells. Antibodies were serially diluted into PBS, then further diluted into assay media until 2x final concentration. Plates were washed twice with assay media by aspirating (vacuum aspirator) followed by addition of 100 μ l per well assay media. After second wash, assay media was replaced with 50 μ l per well of 2X antibody. Cell plate was returned to the incubator for ~ 15-20 min. CAGA12 (clone 4A4) cells were detached with trypsin and pelleted (spun 5 min at 200 x g.). Pellet was resuspended in assay media and viable cells counted. Viable cells were diluted to 0.3e⁶cells/ml and seeded 50 μ l per well (15,000 cells per well). Cells were returned to the incubator.

[815] On Day 3, the assay was read about 16-20 hours after the antibody and/or reporter cell addition. Bright-Glo reagent and test plate were allowed to come to room temperature before reading. 100 μ L of Bright-Glo reagent was added per well. Incubated for 2min with shaking, at room temperature, protected plate from light. The luminescence was then detected on a plate reader.

Results:

[816] Data generated from this assay reflected TGF β activity in cell supernatants. **Figs. 13A and 13B** are graphs showing improved potency of antibodies SR-AB20, SR-AB21, SR-AB22, and SR-

AB23 (SR-AB10 family antibodies) as measured by TGF β activity. The **Fig. 13A** assay was performed without overexpressing LTBP1 and thus measures activation of TGF β in the presence of endogenous presenting molecules. The **Fig. 13B** assay was performed with overexpressed murine LTBP1-proTGF β 1. Notably, antibodies SR-AB22 and SR-AB23 show activity that is similar to (or greater than) the context-independent antibody SR-AB1 in both assays.

[817] **Fig. 14A and Fig. 14B** are graphs showing improved potency of antibodies SR-AB24, SR-AB25, SR-AB26, SR-AB27, SR-AB28, and SR-AB29 (SR-AB13 family antibodies) as measured by TGF β activity. The **Fig. 14A** assay was performed without overexpressing LTBP1 and thus measures activation of TGF β in the presence of endogenous presenting molecules. The **Fig. 14B** assay was performed with overexpressed murine LTBP1-proTGF β 1. Notably, the optimized antibodies show activity that is greater than the context-independent antibody SR-AB1 in the LTBP1-proTGF β 1 overexpression cell assay (**Fig. 14B**).

[818] The assays depicted in **Fig. 13A**, **Fig. 13B**, **Fig. 14A**, and **Fig. 14B** were all performed in LN229 cells, which express low LTBP1 mRNA, high LTBP3 mRNA, undetectable GARP, and undetectable LRRC33.

Example 11: Improved potency of optimized LTBP complex-specific antibodies after light chain mutagenesis

[819] LTBP complex-specific antibodies from the third and fourth rounds of affinity maturation/optimization were assessed for their ability to inhibit TGF β activity using LN229 cells. The assays were carried out using the protocol described in Example 10.

[820] To calculate inhibitory potency (IC₅₀) values, raw luminescence values were normalized against wells treated with PBS only (vehicle). Normalized values were plotted using PRISM® software, and values were calculated using 3-point non-linear regression. The Tables below show IC₅₀ values (nM) for isoform-specific antibodies in two different assay formats: an endogenous human LTBP (hLTBP) assay and mouse LTBP assay (mLTBP). IC₅₀ values were determined in the two assay formats for 4 different antibody lineages: SR-AB22, SR-AB23, SR-AB24 and SR-AB26, shown in Tables 16-19.

Table 16: SR-AB22 lineage

	<i>Endogenous hLTBP assay</i>			<i>mLTBP1 assay</i>
Ref Ab	0.7682		SR-AB36 (cycle 3)	0.3
SR-AB34 (cycle 3)	0.9088		SR-AB30 (cycle 3)	0.4038
SR-AB39 (cycle 3)	1.015		SR-AB34 (cycle 3)	0.4231
SR-AB36 (cycle 3)	1.153		SR-AB35 (cycle 3)	0.4298
SR-AB31 (cycle 3)	1.261		SR-AB31 (cycle 3)	0.4313
SR-AB35 (cycle 3)	1.343		SR-AB33 (cycle 3)	0.4491

SR-AB30 (cycle 3)	1.344		SR-AB32 (cycle 3)	0.4744
SR-AB31 (cycle 3)	1.563		SR-AB39 (cycle 3)	0.4786
SR-AB37 (cycle 3)	1.93		SR-AB37 (cycle 3)	0.5426
SR-AB32 (cycle 3)	2.223		SR-AB22 (cycle 2)	1.394
SR-AB10 (parent)	3.64		Ref Ab	1.571
SR-AB1	4.446		SR-AB1	6.381
SR-AB22 (cycle 2)	4.618		SR-AB14 (cycle1)	12.59
SR-AB14 (cycle1)	51.65		SR-AB10 (parent)	118.9

Table 17: SR-AB23 lineage

	<i>Endogenous hLTBP</i>			<i>mLTBP1</i>
Ref Ab	0.7682		SR-AB45 (cycle 3)	0.2638
SR-AB42 (cycle 3)	1.434		SR-AB42 (cycle 3)	0.2708
SR-AB44 (cycle 3)	1.515		SR-AB44 (cycle 3)	0.3001
SR-AB41 (cycle 3)	1.542		SR-AB43 (cycle 3)	0.3189
SR-AB43 (cycle 3)	1.756		SR-AB40 (cycle 3)	0.333
SR-AB40 (cycle 3)	2.105		SR-AB41 (cycle 3)	0.4086
SR-AB45 (cycle 3)	2.247		SR-AB23 (cycle 2)	0.9458
SR-AB10 (parent)	3.64		Ref Ab	1.571
SR-AB1	4.446		SR-AB1	6.381
SR-AB23 (cycle 2)	6.043		SR-AB14 (cycle1)	12.59
SR-AB14 (cycle1)	51.65		SR-AB10 (parent)	118.9

Table 18: SR-AB24 lineage

	<i>Endogenous hLTBP</i>			<i>mLTBP1</i>
Ref Ab	0.7682		SR-AB62 (cycle 3)	0.3861
SR-AB1	4.446		SR-AB24 (cycle 2)	1.313
SR-AB13 (parent)	4.825		Ref Ab	1.571
SR-AB62 (cycle 3)	8.683		SR-AB17 (cycle 1)	4.038
SR-AB24 (cycle 2)	9.599		SR-AB1	6.381
SR-AB17 (cycle 1)	24.14		SR-AB13 (parent)	25.44

Table 19: SR-AB26 lineage

	<i>Endogenous hLTBP</i>			<i>mLTBP1</i>
SR-AB1	0.7682		SR-AB63 (cycle 3)	0.2075
SR-AB63 (cycle 3)	1.777		SR-AB64 (cycle 3)	0.2683
SR-AB64 (cycle 3)	3.12		SR-AB26 (cycle 2)	1.172
SR-AB1	4.446		Ref Ab	1.571
SR-AB13 (parent)	4.825		SR-AB17 (cycle 1)	4.038
SR-AB26 (cycle 2)	8.742		SR-AB1	6.381
SR-AB17 (cycle 1)	24.14		SR-AB13 (parent)	25.44

[821] Data generated from this assay reflected TGF β activity in cell supernatants. As shown in **Fig. 16**, the functional activity (potency) of LTBP complex specific antibodies improved in each cycle of affinity maturation. As shown in **Fig. 16**, improvement from each cycle of affinity maturation resulted in activity exceeding the context-independent reference antibody (Ref Ab), a potent context-independent inhibitor of latent TGF β 1. The assays described above were all performed in LN229 cells transiently transfected with murine LTBP1 and murine proTGF β 1.

[822] For the results shown in Table 20, the assay was carried out using the protocol described in Example 10, but in this assay Fabs were utilized.

Table 20

Antibody	IC50, nM
RefAb (hIgG4)	0.7676
RefAb (Fab)	3.607
SR-AB47 Fab	0.1861
SR-AB59 Fab	0.1947
SR-Ab56 Fab	0.4393
SR-AB52 Fab	0.4589
SR-AB61 Fab	0.4699
SR-AB41 Fab	0.4861
SR-AB46 Fab	0.4946
SR-AB48 Fab	0.5233
SR-AB51 Fab	0.5354
SR-AB55 Fab	0.5363
SR-AB42 Fab	0.6675
SR-AB60 Fab	0.742
SR-AB57 Fab	0.7754
SR-AB49 Fab	1.024
SR-AB43 Fab	1.077
SR-AB53 Fab	1.181
SR-AB58 Fab	4.856
SR-AB54 Fab	5.87
SR-AB23 Fab	8.533
SR-AB50 Fab	10.97

Example 12: Developability

[823] Antibodies that exhibit favorable biophysical properties have an increased success rate in development. LTBP complex-specific antibodies from the third and fourth rounds of affinity maturation/optimization were assessed for aggregation propensity and polyspecificity. **Fig. 17** shows the results of an enzyme-linked immunosorbent assay (ELISA) showing SR-AB binding to baculovirus (BV) particles. **Fig. 18** shows the results of affinity-capture Self-interaction Nanoparticle Spectroscopy (AC-SINS) Assay. This assay tests how likely an antibody is to interact with itself. It

uses gold nanoparticles that are coated with anti-Fc antibodies. When a dilute solution of antibodies is added, they rapidly become immobilized on the Fc-coated gold beads. If these antibodies subsequently interact with one another, it leads to shorter interatomic distances and an increase in the plasmon wavelength that can be detected by spectroscopy.

[824] These results show that the candidate antibodies demonstrate favorable developability properties, as evidenced by a plasmon shift less than 5 nM in the AC-SINS assay and a BV ELISA signal less than 5 fold of the negative control.

Example 13: Inhibition of TGF β signaling by LTBP complex-specific antibodies in choline-deficient high fat diet (CDHFD) model of mouse NASH

[825] The Choline deficient high fat diet (CDHFD) is an established diet-induced model of Non-Alcoholic steatohepatitis (NASH). In this model, male C57BL/6J mice are fed a choline deficient, 0.1% Methionine, high fat diet for 12 weeks. Three to six weeks after the start of the CDHFD, activation of pro-fibrotic processes can be detected through increased expression of α -smooth muscle actin (α -SMA) protein, a marker for activation of hepatic stellate cells, and elevated tissue hydroxyproline content reflecting increased collagen synthesis and deposition (Matsumoto et. al, *Int J Exp Pathol*. 2013 Apr;94(2):93-103).

[826] LTBP complex-specific antibodies were tested for their ability to inhibit and/or reduce the extent of liver fibrosis in mice in the CDHFD model as follows. An outline of the study is shown below in Table 21.

Table 21

Group	Diet	Lineage	Ab (mgG1)	Dose (mg/kg)	Frequency	Takedown	N
1	Regular Chow		NA	NA	NA	4 weeks	5
2	CDAA/HFD		NA	NA	NA	4 weeks	8
3	Regular Chow		HuNeg	15	2x weekly	12 weeks	5
4	CDAA/HFD		HuNeg	15	2x weekly	12 weeks	10
3	CDAA/HFD		Ref Ab	15	2x weekly	12 weeks	10
4	CDAA/HFD	SR-AB23	SR-AB42	15	2x weekly	12 weeks	10
5	CDAA/HFD	SR-AB22	SR-AB31	15	2x weekly	12 weeks	10

[827] Animals in the test cohorts were on the CDHFD for the duration of the study. A separate cohort of mice (Group 1) were administered a regular chow diet as a study control. Antibodies SR-AB42 and SR-AB31 were administered to mice by intraperitoneal (i.p.) injection beginning at 4 weeks post start of the CDHFD. Animals were dosed at antibody test concentrations 15mg/kg twice a week (*i.e.*, 30 mg/kg/week) for a test duration of 8 weeks (*i.e.*, weeks 4 through 12). A mouse IgG1

isotype antibody was used as a negative control at 15mg/kg twice weekly (*i.e.*, 30 mg/kg/week). Ref Ab was used as a reference antibody due to it being a potent context-independent inhibitor of TGF β 1. Following 8 weeks of dosing, animals were sacrificed and livers were collected for analysis.

Endpoint readout was hydroxyproline content.

Hydroxyproline Content in CDHFD Liver Fibrosis Model

[828] Hydroxyproline, produced by hydroxylation of the amino acid proline, is a signature amino acid for major component of fibrillar collagens, and comprises approximately 13.5% of the protein. Hydroxyproline acts as an important diagnostic indicator of the severity of fibrosis. Animals fed a CDHFD show an increase in hydroxyproline (HYP) in liver tissue. As shown in **Fig. 19**, treatment with SR-AB42 and SR-AB31 inhibited the increase in HYP (μ g/mg tissue) in liver tissue in animals on CDHFD.

Example 14: Inhibition of TGF β signaling by LTBP complex-specific antibodies in a genetic model of Alport syndrome

[829] The murine Col4a3 $-/-$ model is an established genetic model of autosomal recessive Alport syndrome. Alport mice lack a functional collagen 4A3 gene (Col4A3 $-/-$) and therefore cannot form normal type IV collagen trimers, which require α 3, α 4, and α 5 chains. Col4a3 $-/-$ mice develop fibrosis in the kidney consistent with renal fibrosis in human patients, including interstitial fibrosis and tubular atrophy, and Col4a3 $-/-$ mice develop end-stage renal disease (ESRD) between 8 and 30 weeks of age, depending on the genetic background of the mouse. The structural and functional manifestation of renal pathology in Col4a3 $-/-$ mice, combined with the progression to ESRD make Col4a3 $-/-$ mice an ideal model to understand kidney fibrosis. Previous reports point to the importance of the TGF β signaling pathway in this process, and treatment with either an inhibitor of α V β 6 integrin, a known activator of TGF β 1 and TGF β 3, or with a TGF β ligand trap has been reported to prevent renal fibrosis and inflammation in Alport mice (Hahm *et al.* (2007) *The American Journal of Pathology*, 170(1): 110-125).

[830] LTBP complex-specific antibodies were tested for their ability to inhibit and/or reduce the renal fibroses in Alport mice as follows. An outline of the study is shown below in Table 22.

Table 22

Group	Genotype	Ab (hIgG4)	Dose (mg/kg)	Frequency	N
1	Het	HuNeg	30	1	6
2	KO	HuNeg	30	1	10
3	KO	Ref Ab	30	1	10
4	KO	SR-AB42	30	1	10
5	KO	SR-AB63	30	1	10

[831] F1 offspring from Col4a3 +/- males on a 129/Sv genetic background crossed to Col4a3 +/- females on a C57BL/6 genetic background were employed for the study. These mice typically exhibit proteinuria by 4-5 weeks old and typically progress to ESRD by 13-15 weeks old, providing a good therapeutic window for testing efficacy of treatment. 11 week old Col4a3/- mice were dosed with 30 mg/kg SR-AB42 and SR-AB63 intraperitoneally (i.p.) 48 hours prior to animal sacrifice and kidney collection. Ref Ab was used as a reference antibody due to it being a potent context-independent inhibitor of TGFβ1 activation.

pSMAD Analysis in Alport Kidney Model

[832] It is well documented that TGFβ receptor activation leads to a downstream signaling cascade of intracellular events, including phosphorylation of Smad2/3. Therefore, the ability of SR-AB42 and SR-AB63 treatment to inhibit TGFβ signaling was assessed in kidney lysate samples by measuring relative phosphorylation levels of Smad2/3 as assayed by ELISA (Cell Signaling Technologies) according to the manufacturer's instructions. **Fig. 20A** is a graph showing relative ratios of phosphorylated versus total (phosphorylated and unphosphorylated) Smad2/3 (pSMAD2/3:tSMAD2/3) in an Alport mouse model. **Fig. 20B** is a graph showing the amount of phosphorylated SMAD2/3 (pSMAD2/3) as determined by ELISA and **Fig. 20C** is a graph showing the amount of total SMAD2/3 (tSMAD2/3) protein as determined by ELISA. As shown by **Fig. 20B** and **Fig. 20C**, reduction of pSMAD is contributing to the change in ratio shown in **Fig. 20A**. A single dose of SR-AB42 or SR-AB63 was sufficient to significantly inhibit pSmad2/3 signaling in whole kidney lysates, demonstrating efficient target engagement by SR-AB42 and SR-AB63 and that LTBP complex-specific inhibition in the Alport model reduces pSmad levels.

Example 15. Determination of proTGFβ1 C4S binding

[833] A modified proTGFβ1 complex (proTGFβ1 C4S) in which the cysteine residue at position 4 of the pro-domain has been substituted with a serine residue (described in WO 2014/182676) was used to assess antibody binding to C4S antigen by Octet. Cycle 2 antibodies were immobilized and tested against 3 antigens. As shown in the Tables below, robust antibody binding to LTBP-1 proTGFβ1 antigen (Table 23) and proTGFβ1C4S antigen was observed (Table 24), while binding to LRRC33 proTGFβ1 antigen was not observed (Table 25).

Table 23: Binding to human LTBP-1 proTGFβ1

Sample ID	Loading Sample ID	Response (nm)
human LTBP-1 proTGFβ1	SR-AB14	0.3197
human LTBP-1 proTGFβ1	SR-AB20	0.4132
human LTBP-1 proTGFβ1	SR-AB21	0.3196
human LTBP-1 proTGFβ1	SR-AB22	0.4317
human LTBP-1 proTGFβ1	SR-AB23	0.3449
human LTBP-1 proTGFβ1	SR-AB17	0.3036
human LTBP-1 proTGFβ1	SR-AB24	0.3796

human LTBP-1 proTGFB1	SR-AB25	0.3085
human LTBP-1 proTGFB1	SR-AB26	0.2953
human LTBP-1 proTGFB1	SR-AB27	0.3717
human LTBP-1 proTGFB1	SR-AB28	0.4046
human LTBP-1 proTGFB1	SR-AB29	0.3242
human LTBP-1 proTGFB1	HuNeg	-0.0007
human LTBP-1 proTGFB1	SR-AB16	0.2594
human LTBP-1 proTGFB1	SR-AB1	0.4311

Table 24: Binding to human TGFb1 C4S

Sample ID	Loading Sample ID	Response (nm)
human TGFb1 C4S	SR-AB14	0.2002
human TGFb1 C4S	SR-AB20	0.2891
human TGFb1 C4S	SR-AB21	0.1812
human TGFb1 C4S	SR-AB22	0.2937
human TGFb1 C4S	SR-AB23	0.2263
human TGFb1 C4S	SR-AB17	0.2835
human TGFb1 C4S	SR-AB24	0.3846
human TGFb1 C4S	SR-AB25	0.329
human TGFb1 C4S	SR-AB26	0.3764
human TGFb1 C4S	SR-AB27	0.3814
human TGFb1 C4S	SR-AB28	0.3605
human TGFb1 C4S	SR-AB29	0.3334
human TGFb1 C4S	HuNeg	0.001
human TGFb1 C4S	SR-AB16	0.1454
human TGFb1 C4S	SR-AB1	0.3604

Table 25: Binding to human LRRC33 proTGFB1

Sample ID	Loading Sample ID	Response (nm)
human LRRC33 proTGFB1	SR-AB14	0.03
human LRRC33 proTGFB1	SR-AB20	0.0894
human LRRC33 proTGFB1	SR-AB21	0.0141
human LRRC33 proTGFB1	SR-AB22	0.1329
human LRRC33 proTGFB1	SR-AB23	0.0233
human LRRC33 proTGFB1	SR-AB17	0.0043
human LRRC33 proTGFB1	SR-AB24	0.0394
human LRRC33 proTGFB1	SR-AB25	0.024
human LRRC33 proTGFB1	SR-AB26	0.0208
human LRRC33 proTGFB1	SR-AB27	0.0317
human LRRC33 proTGFB1	SR-AB28	0.037
human LRRC33 proTGFB1	SR-AB29	0.0288
human LRRC33 proTGFB1	HuNeg	0.0023

human LRRC33 proTGFβ1	SR-AB16	0.0161
human LRRC33 proTGFβ1	SR-AB1	0.4006

Example 16. Cycle 3 Lead Antibodies Show No Inhibition in TGFβ3 Assay

[834] **Fig. 21** is a graph showing that the lead cycle 3 antibodies show no inhibition in the LTBP-TGFβ3 assay. The TGFβ3 assay is performed similarly to the assay in Example 10, but proTGFβ3 is transfected instead of proTGFβ1. No LTBP construct is transfected as this assay relies on the endogenous presentation molecules in the LN229 cell line

Example 17: Pro-fibrotic effects of TGFβ3-selective activation inhibitor in liver fibrosis model

[835] Liver expresses both TGFβ1 and TGFβ3. To investigate whether inhibition of both isoforms in the CDHFD model would further mitigate fibrosis of the liver, CDHFD mice were treated with a potent TGFβ1-selective activation inhibitor (which inhibits activation of TGFβ1 in the context of human LTBP1 and LTBP3 complexes) and a TGFβ3-selective activation inhibitor, either alone or in combination. In mice treated with TGFβ3 inhibitor, exacerbation of the disease was observed, as evidenced by PSR analysis and additional histopathology analyses (**Fig. 22**). At 12 weeks, *e.g.*, end of the study, animals that received the TGFβ1-selective activation inhibitor showed significantly less fibrosis as compared to IgG-treated animals (negative control). By contrast, animals treated with TGFβ3-selective inhibitor showed significantly more fibrosis with approximately 12% PSR positive area. Animals that received a combination of both TGFβ3-selective inhibitor and TGFβ1-selective inhibitor showed an intermediate level of fibrosis, indicating that antifibrotic effects of the TGFβ1-selective activation inhibitor is being mitigated by TGFβ3 inhibition.

Example 18: Improved affinity of optimized LTBP-complex-specific antibodies after light chain optimization cycle 4

[836] LTBP complex-specific antibodies from the third round of affinity maturation/optimization were selected for a fourth round of affinity maturation/optimization (*i.e.*, light chain optimization as described herein) and particular progeny antibodies, were assessed for their ability to bind various proTGFβ1 constructs.

Materials:

- 96 well plates – Greinerbio-one REF 650101
- Microplate Foils-GE Healthcare – CAT# 28-9758-16
- Biotin CAPture Kit GE Healthcare product number 28920233
- Biacore running buffer – supplied as a 20x solution, diluted in Milli-Q water, TEKnova CAT# H8022

Method:

[837] These experiments were performed using the Biacore 8K using the method “multi-cycle kinetics/affinity using Biotin CAPture kit.”

[838] Data collection rate was set to 10 Hz and the Biotin CAPture step was performed for 300 seconds with a flow rate of 2 μ L/minute. The Biotin CAPture reagent was used after a 1:5 dilution in 1x Biacore running buffer. Biotinylated ligands (the various complexes of TGF β 1) were utilized at a concentration of 10 nM and were immobilized to the sensor chip for 180 seconds with a flow rate of 10 μ L/minute. The analytes, which are each of the Fabs, were tested at 10, 5, 2.5, 1.25, and 0.625 nM with a 0 nM control included. Analyte contact time was 120 seconds and the dissociation time was 900 seconds. The regeneration cycle utilized was 120 seconds with a flow rate of 10 μ L/minute.

[839] The evaluation method used was “multi-cycle affinity using capture” and a 1:1 binding model was used. For every Fab tested against a given biotinylated antigen, the Rmax was set to equal the Rmax of an affinity-matured version of Reference Ab with significantly slower binding off-rate and therefore higher affinity binding to that same antigen in the same experiment. Global fits were utilized for every KD determination. The KD values are shown in Table 26 below.

Table 26

	HuLTBP1-ProTGFβ1	HuLTBP3-ProTGFβ1	HuGARP-ProTGFβ1	HuLRRC33-ProTGFβ1
SR-AB42-HuFab	0.381	0.57	144.614	134.249
SR-AB47-HuFab	0.217	0.33	6.496	6.663
SR-AB49-HuFab	0.319	0.58	6.551	7.704
Ref Ab-HuFab	0.077	0.11	0.202	0.154
	MuLTBP1-ProTGFβ1	MuLTBP3-ProTGFβ1	MuGARP-ProTGFβ1	MuLRRC33-ProTGFβ1
SR-AB42-HuFab	0.611	0.68	29.102	109.989
SR-AB47-HuFab	0.198	0.40	1.950	9.164
SR-AB49HuFab	0.379	0.65	2.820	6.945
Ref Ab-HuFab	0.078	0.09	0.264	0.107

[840] **Table 27** shows the monovalent half-binding-times ($T_{1/2}$). The monovalent half-binding-time ($T_{1/2}$) value was at least 45 minutes for each of hLTBP1-proTGF β 1 and hLTBP3-proTGF β 1 complexes. The monovalent $T_{1/2}$ value was less than 5 minutes for each of hGARP-proTGF β 1 and hLRRC33-proTGF β 1 complexes, as measured by SPR.

Table 27: Monovalent Half-Binding Times ($T_{1/2}$)

LTBP-SR-AB42 HuFab	KD (nM)	$T_{1/2}$ Binding
Hu LTBP1-ProTGF β 1	0.38	1.7 hours
Hu LTBP3-ProTGF β 1	0.57	1.7 hours

Hu GARP-ProTGFβ1	144	1.9 min
Hu LRRC33-ProTGFβ1	134	4.3 min
Mu LTBP1-ProTGFβ1	0.6	1.1 hours
Mu LTBP3-ProTGFβ1	0.68	1.2 hours
Mu GARP-ProTGFβ1	29	5.1 min
Mu LRRC33-ProTGFβ1	110	7.4 min

Embodiments

The present invention encompasses various embodiments. Non-limiting examples are listed below:

1. An antibody, or antigen-binding fragment thereof, comprising at least three of the following six CDRs:

- a) CDR-H1: SEQ ID NO:94, with the proviso that, optionally:
 - i. the threonine residue at position 2 of SEQ ID NO:94 may be substituted with an alanine;
 - ii. the asparagine residue at position 4 of SEQ ID NO:94 may be substituted with an alanine, tyrosine, aspartate, serine, arginine, or histidine;
 - iii. the asparagine residue at position 5 of SEQ ID NO:94 may be substituted with a glutamine, serine, glycine, lysine, glutamate, arginine, or histidine;
 - iv. the tyrosine residue at position 6 of SEQ ID NO:94 may be substituted with a arginine;
 - v. the proline residue at position 7 of SEQ ID NO:94 may be substituted with a glycine, alanine, leucine, serine, asparagine, valine, aspartate, or glutamine;
 - vi. the isoleucine residue at position 8 of SEQ ID NO:94 may be substituted with a methionine or leucine; and/or,
 - vii. the histidine residue at position 9 of SEQ ID NO:94 may be substituted with a phenylalanine, tyrosine, asparagine, or serine;
- b) CDR-H2: SEQ ID NO:95, optionally comprising one or more amino acid changes;
- c) CDR-H3: SEQ ID NO:96, optionally comprising one or more amino acid changes;
- d) CDR-L1: SEQ ID NO:97, optionally comprising one or more amino acid changes;
- e) CDR-L2: SEQ ID NO:98, optionally comprising one or more amino acid changes;
- and,
- f) CDR-L3: SEQ ID NO:99, optionally comprising one or more amino acid changes.

2. The antibody, or antigen-binding fragment thereof, according to embodiment 1, comprising at least three of the following six CDRs:

- a) CDR-H1 comprising the amino acid sequence FTF(X₁)(X₂)YVMH, wherein, optionally: X₁ is S or R; and X₂ is G or S;
- b) CDR-H2 comprising the amino acid sequence (X₁)ISHEG(X₂)(X₃)KYYADSVKG, wherein, optionally: X₁ is V or S; X₂ is S or G; and X₃ is F or L; and
- c) CDR-H3 comprising the amino acid sequence (X₁)(X₂)P(X₃)(X₄)(X₅)(X₆)RRGG(X₇)(X₈)(X₉), wherein, optionally: X₁ is A or V; X₂ is R, V, G or K; X₃ is R, H or L; X₄ is I, V or G; X₅ is A, S, or L; X₆ is A or V; X₇ is F or Y; X₈ is D, G, R, or S; and, X₉ is Y, G, R, L, V, A or K.

- d) CDR-L1 as set forth in SEQ ID NO:97, optionally comprising one or more amino acid changes;
 - e) CDR-L2 as set forth in SEQ ID NO:98, optionally comprising one or more amino acid changes; and
 - f) CDR-L3 as set forth in SEQ ID NO:99, optionally comprising one or more amino acid changes.
3. The antibody, or antigen-binding fragment thereof, according to embodiment 2, wherein:
- a) within CDR-H2; X₂ is S; and
 - b) within CDR-H3; X₁ is A; X₂ is R or V; X₃ is R; X₄ is I; X₅ is A or L; X₆ is A; X₇ is F; X₈ is G; and X₉ and Y.
- 3.1 The antibody, or antigen-binding fragment thereof, according to embodiment 3, wherein one or more of the CDR-L1, CDR-L2, and CDR-L3 is/are affinity matured and/or optimized by CDR diversification and/or mutagenesis.
4. The antibody, or antigen-binding fragment thereof, according to any one of the preceding embodiments, wherein the antibody, or antigen-binding fragment thereof, comprises a heavy chain variable region having an amino acid sequence that is at least 90% identical to SEQ ID NO: SEQ ID NO: 88; and/or,
- wherein the antibody comprises a variable light chain variable region having an amino acid sequence that is at least 90% identical to SEQ ID NO: 89.
5. The antibody, or antigen-binding fragment thereof, according to any one of the preceding embodiments, comprising:
- a heavy chain variable region having an amino acid sequence that is at least 90% identical to SEQ ID NO: 88; and
 - a light chain variable region having an amino acid sequence that is at least 90% identical to SEQ ID NO: 89.
6. An antibody, or antigen-binding fragment thereof, comprising at least three of the following six CDRs:
- a) CDR-H1: SEQ ID NO:100, with the proviso that, optionally:
 - i. the serine residue at position 4 of SEQ ID NO:100 may be substituted with a histidine;

- ii. the serine residue at position 7 of SEQ ID NO:100 may be substituted with an alanine or glycine; and/or,
 - iii. the glycine residue at position 11 of SEQ ID NO:100 may be substituted with a threonine, serine, histidine, leucine, isoleucine, asparagine, valine, or alanine;
 - b) CDR-H2: SEQ ID NO:101, with the proviso that, optionally:
 - i. the serine residue at position 3 of SEQ ID NO:101 may be substituted with an alanine;
 - ii. the glycine residue at position 6 of SEQ ID NO:101 may be substituted with an alanine or serine; and/or,
 - iii. the serine residue at position 7 of SEQ ID NO:101 may be substituted with a threonine;
 - c) CDR-H3: SEQ ID NO:102, optionally comprising one or more amino acid changes;
 - d) CDR-L1: SEQ ID NO:103, optionally comprising one or more amino acid changes;
 - e) CDR-L2: SEQ ID NO:104, optionally comprising one or more amino acid changes; and,
 - f) CDR-L3: SEQ ID NO:105, optionally comprising one or more amino acid changes.
7. The antibody, or antigen-binding fragment thereof, according to embodiment 6, comprising at least three of the following six CDRs:
- g) CDR-H1 comprising the amino acid sequence G(X₁)I(X₂)S(X₃)SYYW(X₄), wherein, optionally: X₁ is S or P; X₂ is S, H or R; X₃ is S or G; and, X₄ is G, I, N or V;
 - h) CDR-H2 comprising the amino acid sequence SISYSA(X₁)TYYNPSLKS, wherein, optionally: X₁ is S or T;
 - i) CDR-H3 comprising the amino acid sequence (X₁)(X₂)D(X₃)(X₄)Y(X₅)(X₆)(X₇)(X₈)G(X₉)(X₁₀)(X₁₁), wherein, optionally: X₁ is A or V; X₂ is R, S or G; X₃ is P, Y, R, V, I, H, T or E; X₄ is S, D, E or N; X₅ is D, A or T; X₆ is S, G, T or A; X₇ is I, A, R, Q, or V; X₈ is A, E, K, G or T; X₉ is M or I; X₁₀ is D, L, Q, V, N or G; and, X₁₁ is V, R, N, E or K;
 - j) CDR-L1: SEQ ID NO:103, optionally comprising one or more amino acid changes;
 - k) CDR-L2: SEQ ID NO: 104, optionally comprising one or more amino acid changes; and
 - l) CDR-L3: SEQ ID NO: 105, optionally comprising one or more amino acid changes.
8. The antibody, or antigen-binding fragment thereof, according to embodiment 7, wherein:
- g) within CDR-H1: X₂ is H or R; X₃ is S; and, X₄ is G, I or N; and
 - h) within CDR-H3: X₁ is A; X₃ is P or V; X₄ is S; X₅ is D; X₆ is S or A; X₇ is A, R, I or V; X₈ is A; X₉ is M; X₁₀ is D, Q, or G; and, X₁₁ is V or R.

8.1 The antibody, or antigen-binding fragment thereof, according to embodiment 8, wherein one or more of the CDR-L1, CDR-L2, and CDR-L3 is/are affinity matured and/or optimized by CDR diversification and/or mutagenesis.

9. The antibody, or antigen-binding fragment thereof, according to any one of embodiments 6-8.1, comprising at least three CDRs selected from the following, optionally comprising one or more amino acid changes for each of the CDRs:

CDR-H1: SEQ ID NO:100;

CDR-H2: SEQ ID NO:101;

CDR-H3: SEQ ID NO:102;

CDR-L1: SEQ ID NO:103;

CDR-L2: SEQ ID NO:104; and,

CDR-L3: SEQ ID NO:105.

10. The antibody, or antigen-binding fragment thereof, according to one of embodiments 6-9, comprising a variable heavy chain region and a variable light chain region, wherein:

i) the variable heavy chain region comprises an amino acid sequence that is at least 90% identical to SEQ ID NO: 106; and/or,

ii) the variable light chain variable region comprises an amino acid sequence that is at least 90% identical to SEQ ID NO: 107.

11. The antibody, or antigen-binding fragment thereof, according to one of embodiments 6-9, wherein the antibody or antigen-binding portion thereof, comprises:

i) a variable heavy chain region having an amino acid sequence that is at least 90% identical to SEQ ID NO: 106; and,

ii) a light chain variable region having an amino acid sequence that is at least 90% identical to SEQ ID NO: 107.

12. The antibody, or antigen-binding fragment thereof, according to any one of the preceding embodiments, comprising all six CDRs.

12.1 The antibody, or antigen-binding fragment thereof, according to any one of the preceding embodiments, wherein the one or more amino acid changes comprises one amino acid change.

- 12.2 The antibody, or antigen-binding fragment thereof, according to any one of the preceding embodiments, wherein the one or more amino acid changes comprises up to two amino acid changes.
- 12.3 The antibody, or antigen-binding fragment thereof, according to any one of the preceding embodiments, wherein the one or more amino acid changes comprises up to three amino acid changes.
- 12.4 The antibody, or antigen-binding fragment thereof, according to any one of the preceding embodiments, wherein the one or more amino acid changes comprises up to four amino acid changes.
- 12.5 The antibody, or antigen-binding fragment thereof, according to any one of the preceding embodiments, wherein the one or more amino acid changes comprises up to five amino acid changes.
- 12.6 The antibody, or antigen-binding fragment thereof, according to any one of the preceding embodiments, wherein the one or more amino acid changes comprises up to six amino acid changes.
- 12.7 The antibody, or antigen-binding fragment thereof, according to any one of the preceding embodiments, wherein the one or more amino acid changes comprises up to seven amino acid changes.
- 12.8 The antibody, or antigen-binding fragment thereof, according to any one of the preceding embodiments, wherein any one of the CDRs comprise no amino acid changes.
13. The antibody, or antigen-binding fragment thereof, according to any one of the preceding embodiments, wherein the antibody, or antigen-binding fragment thereof, specifically binds a human LTBP1-proTGF β complex.
14. The antibody, or antigen-binding fragment thereof, according to any one of the preceding embodiments, wherein the antibody, or antigen-binding fragment thereof, specifically binds a human LTBP3-proTGF β complex.
15. The antibody, or antigen-binding fragment thereof, according to any one of the preceding embodiments, wherein the antibody, or antigen-binding fragment thereof, does not bind a human GARP-proTGF β complex.
16. The antibody, or antigen-binding fragment thereof, according to any one of the preceding embodiments, wherein the antibody, or antigen-binding fragment thereof, specifically binds a human

LTBP1-proTGF β complex and/or a human LTBP3-proTGF β complex, and does not bind a human GARP-proTGF β complex.

17. The antibody, or antigen-binding fragment thereof, according to any one of the preceding embodiments, wherein the antibody, or antigen-binding fragment thereof does not bind mature TGF β 1, mature TGF β 2 or mature TGF β 3.

18. The antibody, or antigen-binding fragment thereof, according to any one of the preceding embodiments, wherein the antibody, or antigen-binding fragment thereof, is a fully human or humanized antibody.

19. The antibody, or antigen-binding fragment thereof, according to any one of the preceding embodiments, wherein the antibody, or antigen-binding fragment thereof, is an isolated antibody, or antigen-binding fragment thereof.

20. The antibody, or antigen-binding fragment thereof, according to any one of the preceding embodiments, wherein the antibody, or antigen-binding fragment thereof, does not compete with any one of the antibodies SR-Ab1, SR-Ab2 or SR-Ab13 for binding to a human LTBP1-proTGF β 1 complex.

21. The antibody, or antigen-binding fragment thereof, according to any one of the preceding embodiments, wherein the antibody, or antigen-binding fragment thereof, does not compete with any one of the antibodies SR-Ab1, SR-Ab2 or SR-Ab10 for binding to a human LTBP1-proTGF β 1 complex.

22. The antibody, or antigen-binding fragment thereof, according to any one of the preceding embodiments, wherein the antibody is an IgG4 or IgG1 subtype.

23. The antibody, or antigen-binding fragment thereof, according to any one of the preceding embodiments, wherein the antibody is specific for human LTBP1-TGF β 1 complex.

24. The antibody, or antigen-binding fragment thereof, according to any one of the preceding embodiments, wherein the antibody is specific for human LTBP3-TGF β 1 complex.

25. The antibody, or antigen-binding fragment thereof, according to any one of the preceding embodiments, wherein the antibody is specific for human LTBP1-TGF β 1 complex and human LTBP3-TGF β 1 complex.
26. The antibody, or antigen-binding fragment thereof, according to any one of the preceding embodiments, wherein the antibody does not bind a human GARP-TGF β 1 complex or GARP-TGF β 3 complex.
27. The antibody, or antigen-binding fragment thereof, according to any one of the preceding embodiments, wherein the antibody, or antigen-binding fragment thereof, binds a human LTBP1-proTGF β 1 complex with a K_D of < 50 nM as measured by Bio-Layer Interferometry (BLI).
28. The antibody, or antigen-binding fragment thereof, according to any one of the preceding embodiments, wherein the antibody, or antigen-binding fragment thereof, binds a human LTBP3-proTGF β 1 complex with a K_D of < 50 nM as measured by Bio-Layer Interferometry (BLI).
29. The antibody, or antigen-binding fragment thereof, according to any one of the preceding embodiments, wherein the antibody, or antigen-binding fragment thereof, binds a human LTBP1-proTGF β 1 complex with a K_D of < 10 nM as measured by Bio-Layer Interferometry (BLI).
30. The antibody, or antigen-binding fragment thereof, according to any one of the preceding embodiments, wherein the antibody, or antigen-binding fragment thereof, binds a human LTBP3-proTGF β 1 complex with a K_D of < 10 nM as measured by Bio-Layer Interferometry (BLI).
31. The antibody, or antigen-binding fragment thereof, according to any one of embodiments 27-30, wherein the antibody, or antigen-binding fragment thereof, is cross-reactive with mouse LTBP1-proTGF β 1.
32. The antibody, or antigen-binding fragment thereof, according to any one of embodiments 27-30, wherein the antibody, or antigen-binding fragment thereof, is cross-reactive with mouse LTBP3-proTGF β 1.
33. The antibody, or antigen-binding fragment thereof, according to any one of embodiments 27-30, wherein the antibody, or antigen-binding fragment thereof, is cross-reactive with both mouse LTBP1-proTGF β 1 and mouse LTBP3-proTGF β 1.

34. The antibody, or antigen-binding fragment thereof, according to any one of the preceding embodiments, wherein the antibody, or antigen-binding fragment thereof, binds a mouse LTBP1-proTGF β 1 complex with a K_D of < 50 nM as measured by Bio-Layer Interferometry (BLI).
35. The antibody, or antigen-binding fragment thereof, according to any one of the preceding embodiments, wherein the antibody, or antigen-binding fragment thereof, binds a mouse LTBP3-proTGF β 1 complex with a K_D of < 50 nM as measured by Bio-Layer Interferometry (BLI).
36. The antibody, or antigen-binding fragment thereof, according to any one of the preceding embodiments, wherein the antibody, or antigen-binding fragment thereof, binds a mouse LTBP1-proTGF β 1 complex with a K_D of < 10 nM as measured by Bio-Layer Interferometry (BLI).
37. The antibody, or antigen-binding fragment thereof, according to any one of the preceding embodiments, wherein the antibody, or antigen-binding fragment thereof, binds a mouse LTBP3-proTGF β 1 complex with a K_D of < 10 nM as measured by Bio-Layer Interferometry (BLI).
38. The antibody, or antigen-binding fragment thereof, according to any one of the preceding embodiments, wherein the antibody, or antigen-binding fragment thereof, is isoform-specific in that it selectively binds and inhibits the activation of TGF β 1 associated with LTBP1/3.
39. The antibody, or antigen-binding fragment thereof, according to embodiment 38, wherein the antibody, or antigen-binding fragment thereof, does not bind GARP- TGF β 1.
40. An antibody, or antigen-binding fragment thereof, comprising the following six CDRs:
- a) CDR-H1 comprising the amino acid sequence FTFRSYVMH;
 - b) CDR-H2 comprising the amino acid sequence VISHEGS(X₁)KYYADSVKG, wherein: X₁ is L or G; and
 - c) CDR-H3 comprising the amino acid sequence A(X₁)PRIAARRGGFG(X₂), wherein: X₁ is V, R or L; and X₂ is Y, S or T;
 - d) CDR-L1 comprising the amino acid sequence TRS(X₁)G(X₂)ID(X₃)NYVQ, wherein, X₁ is S or H; X₂ is N, L, S or A; and X₃ is N, D or Y;
 - e) CDR-L2 comprising the amino acid sequence ED(X₁)(X₂)RPS, wherein: X₁ is N, F or A; and X₂ is Q, I or V; and
 - f) CDR-L3 comprising the amino acid sequence Q(X₁)YD(X₂)(X₃)(X₄)Q(X₅)VV, wherein: X₁ is S or G; X₂ is S, F, Y, D, H or W; X₃ is N, D or S; X₄ is N, A, L, E or T; and X₅ is G, R, A or L.

41. The antibody, or antigen-binding fragment thereof, according to embodiment 40, wherein:
within CDR-H3: X₁ is R or L.
42. The antibody, or antigen-binding fragment thereof, according to embodiment 41, wherein:
within CDR-L3: X₂ is Y.
43. The antibody, or antigen-binding fragment thereof, according to embodiment 42, wherein:
within CDR-L3: X₃ is D; and X₄ is T.
44. The antibody, or antigen-binding fragment thereof, according to embodiment 42, wherein:
within CDR-L3: X₃ is D; X₄ is N; and X₅ is A.
45. The antibody, or antigen-binding fragment thereof, according to embodiment 41, wherein:
within CDR-L1: X₁ is S or H; X₂ is N or A; and X₃ is N, D or Y;
within CDR-L2: X₁ is N or F; and X₂ is Q or V; and
within CDR-L3: X₁ is S or G; X₂ is S, Y, D or W; X₃ is D or S; X₄ is N, L or T; and X₅ is G,
R, A or L.
46. The antibody, or antigen-binding fragment thereof, according to embodiment 45, wherein:
within CDR-L1: X₁ is S; X₂ is N; and X₃ is N or Y;
within CDR-L2: X₁ is N; and X₂ is Q or V; and
within CDR-L3: X₁ is S or G; X₂ is S, Y or W; X₃ is D; X₄ is N or T; and X₅ is G, R or A.
47. The antibody, or antigen-binding fragment thereof, according to embodiment 46, wherein:
within CDR-L3: X₁ is S; X₂ is S or Y; X₃ is D; X₄ is N or T; and X₅ is G, R or A.
48. The antibody, or antigen-binding fragment thereof, according to embodiment 47, wherein:
within CDR-L3: X₁ is S; X₂ is Y; X₃ is D; X₄ is N or T; and X₅ is G or A.
49. The antibody, or antigen-binding fragment thereof, according to embodiment 48, wherein:
within CDR-L3: X₁ is S; X₂ is Y; X₃ is D; X₄ is T; and X₅ is G.
50. The antibody, or antigen-binding fragment thereof, according to any one of embodiments 45-49, wherein:
 - a) CDR-H1 comprises the amino acid sequence of SEQ ID NO: 166;

- b) CDR-H2 comprises the amino acid sequence of SEQ ID NO: 167;
- c) CDR-H3 comprises the amino acid sequence of SEQ ID NO: 168;
- d) CDR-L1 comprises the amino acid sequence of SEQ ID NO: 169;
- e) CDR-L2 comprises the amino acid sequence of SEQ ID NO: 170; and
- f) CDR-L3 comprises the amino acid sequence of SEQ ID NO: 171.

51. The antibody, or antigen-binding fragment thereof, according to any one of embodiments 40-50, which comprises:

a heavy chain variable region having an amino acid sequence that is at least 90% identical to SEQ ID NO: 318; and

a light chain variable region having an amino acid sequence that is at least 90% identical to SEQ ID NO: 319.

52. The antibody, or antigen-binding fragment thereof, according to any one of embodiments 40-51, which competes or cross-competes with an antibody having a heavy chain variable region sequence as set forth in SEQ ID NO: 318 and light chain variable region sequence as set forth in SEQ ID NO: 319.

53. An antibody, or antigen-binding fragment thereof, which selectively binds to human LTBP1-TGF β 1 complex and human LTBP3-TGF β 1 complex and competes or cross-competes with an antibody having a heavy chain variable region sequence as set forth in SEQ ID NO: 318 and light chain variable region sequence as set forth in SEQ ID NO: 319.

54. An antibody, or antigen-binding fragment thereof, comprising the following six CDRs:

- g) CDR-H1 comprising the amino acid sequence G(X₁)I(X₂)S(X₃)SYYW(X₄), wherein, optionally: X₁ is S; X₂ is S, H or R; X₃ is S or G; and, X₄ is G, I, N or V;
- h) CDR-H2 comprising the amino acid sequence SISYS(X₁)(X₂)TYY, wherein, optionally: X₁ is G or A; and X₂ is S or T;
- i) CDR-H3 comprising the amino acid sequence A(X₁)DPSYDS(X₂)AGM(X₃)V, wherein, optionally: X₁ is R, S or G; X₂ is A or I; and X₃ is D or Q;
- j) CDR-L1 comprising the amino acid sequence RAS(X₁)(X₂)IS(X₃)YLN, wherein, optionally: X₁ is K or Q; X₂ is V or S; and X₃ is S or Y;
- k) CDR-L2 comprising the amino acid sequence (X₁)AS(X₂)(X₃)QS, wherein, optionally: X₁ is Y, A or S; X₂ is S or N; and X₃ is L or R;
- l) CDR-L3 comprising the amino acid sequence QQ(X₁)(X₂)D(X₃)P(X₄)T, wherein, optionally: X₁ is S or G; X₂ is F or N; X₃ is W or F; and X₄ is F or L.

55. The antibody, or antigen-binding fragment thereof, according to embodiment 54, wherein:
- a) within CDR-H1: X₁ is S; X₂ is S or R; X₃ is S; and, X₄ is G;
 - b) within CDR-H2: X₁ is G or A; and X₂ is S or T;
 - c) within CDR-H3: X₁ is R, S or G; X₂ is A or I; and X₃ is D or Q;
 - d) within CDR-L1: X₁ is K or Q; X₂ is V or S; and X₃ is S or Y;
 - e) within CDR-L2: X₁ is Y, A or S; X₂ is S or N; and X₃ is L or R; and
 - f) within CDR-L3: X₁ is S or G; X₂ is F or N; X₃ is W or F; and X₄ is F or L.
56. The antibody, or antigen-binding fragment thereof, according to embodiment 55, wherein:
- a) CDR-H1 comprises the amino acid sequence GSIRSSSYWYG;
 - b) CDR-H2 comprises the amino acid sequence SISYSATTYY;
 - c) within CDR-H3: X₁ is S or G; X₂ is A or I; and X₃ is D or Q;
 - d) within CDR-L1: X₁ is K or Q; X₂ is V or S; and X₃ is S or Y;
 - e) within CDR-L2: X₁ is Y, A or S; X₂ is S or N; and X₃ is L or R; and
 - f) within CDR-L3: X₁ is S or G; X₂ is F or N; X₃ is W or F; and X₄ is F or L.
57. The antibody, or antigen-binding fragment thereof, according to embodiment 56, wherein:
- g) CDR-H1 comprises the amino acid sequence of SEQ ID NO: 292;
 - h) CDR-H2 comprises the amino acid sequence of SEQ ID NO: 293;
 - i) CDR-H3 comprises the amino acid sequence of SEQ ID NO: 294;
 - j) CDR-L1 comprises the amino acid sequence of SEQ ID NO: 295;
 - k) CDR-L2 comprises the amino acid sequence of SEQ ID NO: 296; and
 - l) CDR-L3 comprises the amino acid sequence of SEQ ID NO: 297.
58. The antibody, or antigen-binding fragment thereof, according to any one of embodiments 54-57, which is an antibody according to any one of claims 1-5.
59. The antibody, or antigen-binding fragment thereof, according to any one of embodiments 54-58, which comprises:
- a heavy chain variable region having an amino acid sequence that is at least 90% identical to SEQ ID NO: 360; and
 - a light chain variable region having an amino acid sequence that is at least 90% identical to SEQ ID NO: 361.

60. The antibody, or antigen-binding fragment thereof, according to any one of embodiments 54-59, which competes or cross-competes with an antibody having a heavy chain variable region sequence as set forth in SEQ ID NO: 360 and light chain variable region sequence as set forth in SEQ ID NO: 361.
61. An antibody, or antigen-binding fragment thereof, which selectively binds to human LTBP1-TGF β 1 complex and human LTBP3-TGF β 1 complex and competes or cross-competes with an antibody having a heavy chain variable region sequence as set forth in SEQ ID NO: 360 and light chain variable region sequence as set forth in SEQ ID NO: 361.
62. The antibody, or antigen-binding fragment thereof, according to any one of embodiments 40-61, which does not show detectable binding to a human GARP-proTGF β 1 complex, as measured by BLI, under the same assay conditions as used to measure binding to human LTBP1-proTGF β 1 complex and human LTBP3-TGF β 1 complex.
63. The antibody, or antigen-binding fragment thereof, according to any one of embodiments 40-62, which binds a human LTBP1-proTGF β 1 complex and/or a human LTBP3-TGF β 1 complex with a K_D that is at least 50 times lower than the K_D when binding to a human GARP-proTGF β 1 complex under the same assay conditions.
64. The antibody, or antigen-binding fragment thereof, according to any one of embodiments 40-63, which does not show detectable binding to an LRRC33-proTGF β 1 complex, as measured by BLI, under the same assay conditions as used to measure binding to human LTBP1-proTGF β 1 complex and human LTBP3-TGF β 1 complex.
65. The antibody, or antigen-binding fragment thereof, according to any one of embodiments 40-64, wherein the antibody, or antigen-binding fragment thereof has a monovalent half-binding-time ($t_{1/2}$) of at least 45 minutes for each of hLTBP1-proTGF β 1 and hLTBP3-proTGF β 1 complexes, as measured by SPR.
66. The antibody, or antigen-binding fragment thereof, according to embodiment 65 which has a monovalent $t_{1/2}$ of less than 5 minutes for each of hGARP-proTGF β 1 and hLRRC33-proTGF β 1 complexes, as measured by SPR.

67. The antibody, or antigen-binding fragment thereof, according to any one of the preceding embodiments, which binds a human LTBP1-proTGF β 1 complex and a human LTBP3-TGF β 1 complex with a K_D of < 5 nM as measured by Bio-Layer Interferometry (BLI), optionally < 1 nM.
68. The antibody, or antigen-binding fragment thereof, according to any one of the preceding embodiments, which is cross-reactive with mouse LTBP1-proTGF β 1.
69. The antibody, or antigen-binding fragment thereof, according to any one of the preceding embodiments, which is cross-reactive with mouse LTBP3-proTGF β 1.
70. The antibody, or antigen-binding fragment thereof, according to any one of the preceding embodiments, wherein the antibody, or antigen-binding fragment thereof, binds a mouse LTBP1-proTGF β 1 complex with a K_D of < 10 nM as measured by Bio-Layer Interferometry (BLI).
71. The antibody, or antigen-binding fragment thereof, according to any one of the preceding embodiments, wherein the antibody, or antigen-binding fragment thereof, binds a mouse LTBP3-proTGF β 1 complex with a K_D of < 10 nM as measured by Bio-Layer Interferometry (BLI).
72. The antibody, or antigen-binding fragment thereof, according to any one of the preceding embodiments, wherein the antibody, or antigen-binding fragment thereof cross-reacts with human and murine LTBP1-proTGF β 1 and LTBP3-proTGF β 1 complexes, each with a K_D of < 5 nM, optionally < 1 nM.
73. The antibody, or antigen-binding fragment thereof, according to any one of the preceding embodiments, wherein the antibody is an IgG4 or IgG1 subtype.
74. The antibody, or antigen-binding fragment thereof, according to embodiment 73, wherein the antibody is a human IgG4 subtype, wherein optionally the antibody comprises a backbone substitution of Ser to Pro that produces an IgG1-like hinge.
75. A pharmaceutical composition comprising the antibody, or antigen-binding fragment thereof, according to any one of the preceding embodiments and a pharmaceutically acceptable excipient.
76. The pharmaceutical composition according to embodiment 75, which is prepared for intravenous administration or subcutaneous administration.

77. A composition comprising a multi-dose vial containing the pharmaceutical composition of embodiments 75 or 76.
78. A composition comprising a single-dose syringe containing the pharmaceutical composition of embodiments 75 or 76, optionally wherein the syringe is a disposable syringe.
79. The composition according to any one of embodiments 75 or 78 for use in a method for the treatment of a fibrotic condition in a human subject, wherein the treatment comprises administration of the composition to the subject in an amount effective to treat the fibrotic disorder.
80. The composition for use according to embodiment 79, wherein the fibrotic disorder is an organ fibrosis.
81. The composition for use according to embodiment 80, wherein the organ fibrosis is an advanced organ fibrosis.
82. The composition for use according to embodiments 80 or 81, wherein the organ fibrosis is selected from the group consisting of:
kidney fibrosis, liver fibrosis, lung fibrosis, cardiac fibrosis, pancreatic fibrosis, skin fibrosis, scleroderma, muscle fibrosis, uterine fibrosis and endometriosis.
83. The composition for use according to embodiment 82, wherein:
- a) the fibrotic disorder comprises chronic inflammation;
 - b) the subject benefits from immune suppression;
 - c) the subject has or is at risk of developing an autoimmune disease;
 - d) the subject is a candidate for an allograft transplant; and/or,
 - e) the subject has received an allograft transplant.
84. The composition for use according to embodiment 79, wherein the fibrotic disorder comprising chronic inflammation is a muscular dystrophy, multiple sclerosis (MS), or Cystic Fibrosis (CF).
85. The composition for use according to embodiment 84, wherein the muscular dystrophy is Duchenne muscular dystrophy (DMD).

86. The composition for use according to embodiment 84, wherein the MS comprises perivascular fibrosis.
87. The composition for use according to embodiment 82, wherein the lung fibrosis is idiopathic pulmonary fibrosis (IPF).
88. The composition for use according to embodiment 82, wherein the subject has chronic kidney disease (CKD).
89. The composition for use according to embodiment 82, wherein the subject has liver fibrosis associated with nonalcoholic steatohepatitis (NASH) or nonalcoholic fatty liver disease (NAFLD).
90. The composition for use according to embodiment 89, wherein the subject has cirrhosis or hepatocellular carcinoma associated with NASH.
91. The composition for use according to embodiment 90, wherein the subject suffers from a metabolic condition (*e.g.*, obesity, type 2 diabetes).
92. The composition for use according to any one of embodiments 89-91, wherein the the subject is further treated with a myostatin inhibitor, wherein optionally the myostatin inhibitor is a myostatin-selective inhibitor, wherein further optionally the myostatin-selective inhibitor is SRK-015, trevogrumab, or any variant thereof, or an antibody according to WO 2016/098357.
93. The composition for use according to any one of embodiments 79-92, wherein the antibody is administered to the subject at a dosage of between 0.1 and 30 mg/kg.
94. The composition for use according to embodiment 93, wherein the antibody is administered twice a week, once a week, once every 2 weeks, once every 3 weeks, once every 4 weeks, once a month, once every 6 weeks, or every other month.
95. The composition for use according to embodiment 93, wherein a therapeutic regimen comprises an initial phase of a therapy and a subsequent phase of the therapy,
wherein the subject receives a loading dose during the initial phase followed by a maintenance dose during the subsequent phase.

96. The composition for use according to embodiment 95, wherein the loading dose is between 2-30 mg/kg, and the maintenance dose is between 0.1-20 mg/kg.
97. The composition for use according to embodiment 95 or 96, wherein the loading dose is administered to the subject twice a week, once a week, once every 2 weeks or every 3 weeks.
98. The composition for use according to any one of embodiments 95-97, wherein the maintenance dose is administered to the subject once every 2-12 weeks.
99. The composition for use according to any one of embodiments 95-97, wherein the maintenance dose is administered to the subject on an as-needed basis.
100. The composition for use according to any one of embodiments 79-99, wherein the method further comprises testing or confirming expression of TGF β 1, LTBP1 or LTBP3 in a biological sample collected from the subject.
101. A method for making a composition according to any one of embodiments 75-78, comprising an antibody, or antigen-binding fragment thereof, that specifically binds a human LTBP1-proTGF β complex and/or a human LTBP3-proTGF β complex, and does not bind a human GARP-proTGF β complex; wherein the antibody or fragment inhibits TGF β 1 but does not inhibit TGF β 2 or TGF β 3, the method comprising steps of:
- i) providing at least one antigen comprising LTBP1-proTGF β 1 and/or LTBP3-proTGF β 1,
 - ii) selecting a first pool of antibodies or fragments that specifically bind the at least one antigen of step (i) so as to provide specific binders of LTBP1-proTGF β 1 and/or LTBP3-proTGF β 1;
 - iii) selecting a second pool of antibodies or fragments that inhibit activation of TGF β 1, so as to generate specific inhibitors of TGF β 1 activation; and
 - iv) formulating an antibody or fragment that is present in the first pool of antibodies and the second pool of antibodies into a pharmaceutical composition, thereby making the composition comprising the antibody or fragment.
102. The method of embodiment 101, wherein the method further comprises a step of:
- removing from the first pool of antibodies, or fragments, any antibodies or fragments that bind GARP-proTGF β 1, LRRC33-proTGF β 1, mature TGF β 1, GARP-proTGF β 2, LRRC33-proTGF β 2, mature TGF β 2, GARP-proTGF β 3, LRRC33-proTGF β 3, mature TGF β 3, or any combinations thereof.

103. The method of embodiment 101 or 102, wherein the method further comprises a step of:
determining or confirming isoform-specificity of the antibodies or fragments selected in steps (ii) and/or (iii).
104. The method according to any one of embodiments 101-103, wherein the method further comprises a step of:
selecting antibodies or fragments that are cross-reactive to human and rodent antigens.
105. The method according to any one of embodiments 101-104, wherein the method further comprises a step of:
generating a fully human or humanized antibody or fragment, of the antibody or fragment that is presented in the first pool of antibodies and the second pool of antibodies.
106. The method according to any one of embodiments 101-105, wherein the method further comprises a step of:
subjecting the antibody or fragment that is present in the first pool of antibodies and the second pool of antibodies to affinity maturation and/or optimization, so as to provide an affinity matured and/or optimized antibody or fragment.
107. The method according to any one of embodiments 101-106, wherein the affinity maturation and/or optimization comprises a step of subjecting the antibody, or antigen-binding fragment thereof, to light chain shuffling.
108. The method according to any one of embodiments 101-107, wherein the affinity maturation and/or optimization comprises the step of subjecting the antibody, or antigen-binding fragment thereof, to CDR H1/H2 diversification.
109. The method according to any one of embodiments 101-108, wherein the affinity maturation and/or optimization comprises the step of subjecting the antibody, or antigen-binding fragment thereof, to CDR-H3 mutagenesis.
110. The method according to any one of embodiments 101-109, wherein the affinity maturation and/or optimization comprises the step of subjecting antibody, or antigen-binding fragment thereof, to light chain CDR mutagenesis.

111. The method according to any one of embodiments 101-110, wherein the affinity maturation and/or optimization comprises the step of subjecting the antibody, or antigen-binding fragment thereof, to light chain CDR L1/L2 diversification.
112. The method according to any one of embodiments 101-111, wherein the method further comprises a step of determining affinity of the antibodies, or antigen-binding fragments thereof, to human LTBP1-proTGF β 1 and/or human LTBP3-proTGF β 1.
113. The method according to embodiments 101-112, wherein the method further comprises a step of removing from the first and/or second pools of antibodies, or antigen-binding fragments thereof, any antibodies, or antigen-binding fragments thereof, that bind to human LTBP1-proTGF β 1 and/or human LTBP3-proTGF β 1 with a K_D of > 100 nM, as measured by Bio-Layer Interferometry (BLI).
114. The method according to embodiment 101-113, wherein the method further comprises a step of removing from the first and/or second pools of antibodies, or antigen-binding fragments thereof, any antibodies, or antigen-binding fragments thereof, that bind to human LTBP1-proTGF β 1 and/or human LTBP3-proTGF β 1 with a K_D of > 50 nM, as measured by Bio-Layer Interferometry (BLI).
115. The method according to embodiment 101-114, wherein the method further comprises a step of removing from the first and/or second pools of antibodies, or antigen-binding fragments thereof, any antibodies, or antigen-binding fragments thereof, that bind to human LTBP1-proTGF β 1 and/or human LTBP3-proTGF β 1 with a K_D of > 25 nM, as measured by Bio-Layer Interferometry (BLI).
116. The method according to embodiment 101-115, wherein the method further comprises a step of removing from the first and/or second pools of antibodies, or antigen-binding fragments thereof, any antibodies, or antigen-binding fragments thereof, that bind to human LTBP1-proTGF β 1 and/or human LTBP3-proTGF β 1 with a K_D of > 10 nM, as measured by Bio-Layer Interferometry (BLI).
117. The method according to any one of embodiments 101-116, wherein the method further comprises a step of determining affinity of the antibodies, or antigen-binding fragments thereof, from the first and/or second pools of antibodies to mouse LTBP1-proTGF β 1 and/or mouse LTBP3-proTGF β 1.
118. The method according to embodiment 117, wherein the method further comprises a step of removing from the first and/or second pools of antibodies, or antigen-binding fragments thereof, any

antibodies, or antigen-binding fragments thereof, that bind to mouse LTBP1-proTGF β 1 and/or mouse LTBP3-proTGF β 1 with a K_D of > 100 nM, as measured by Bio-Layer Interferometry (BLI).

119. The method according to embodiment 117, wherein the method further comprises a step of removing from the first and/or second pools of antibodies, or antigen-binding fragments thereof, any antibodies, or antigen-binding fragments thereof, that bind to mouse LTBP1-proTGF β 1 and/or mouse LTBP3-proTGF β 1 with a K_D of > 50 nM, as measured by Bio-Layer Interferometry (BLI).

120. The method according to embodiment 117, wherein the method further comprises a step of removing from the first and/or second pools of antibodies, or antigen-binding fragments thereof, any antibodies, or antigen-binding fragments thereof, that bind to mouse LTBP1-proTGF β 1 and/or mouse LTBP3-proTGF β 1 with a K_D of > 25 nM, as measured by Bio-Layer Interferometry (BLI).

121. The method according to embodiment 117, wherein the method further comprises a step of removing from the first and/or second pools of antibodies, or antigen-binding fragments thereof, any antibodies, or antigen-binding fragments thereof, that bind to mouse LTBP1-proTGF β 1 and/or mouse LTBP3-proTGF β 1 with a K_D of > 10 nM, as measured by Bio-Layer Interferometry (BLI).

122. The method according to any one of embodiments 101-121, wherein the method further comprises a step of removing from the first and/or second pool of antibodies, or antigen-binding fragments thereof, any antibodies, or antigen-binding fragments thereof, that do not bind mouse LTBP1-proTGF β 1 and/or mouse LTBP3-proTGF β 1.

123. The method according to any one of embodiments 101-122, wherein the method further comprises a step of determining the IC_{50} of the antibodies, or antigen-binding fragments thereof, from the first and/or second pools of antibodies, or antigen-binding fragments thereof, as measured by a suitable functional in vitro cell-based assay.

124. The method according to embodiments 123, wherein the suitable functional in vitro cell-based assay is a cell-based assay.

125. The method according to embodiments 123 or 124, wherein the method comprises the step of removing antibodies, or antigen-binding fragments thereof, from the first and/or second pools of antibodies, or antigen-binding fragments thereof, that have an IC_{50} of > 100 nM.

126. The method according to embodiments 123 or 124, wherein the method comprises the step of removing antibodies, or antigen-binding fragments thereof, from the first and/or second pools of antibodies, or antigen-binding fragments thereof, that have an IC_{50} of > 50 nM.
127. The method according to embodiments 123 or 124, wherein the method comprises the step of removing antibodies, or antigen-binding fragments thereof, from the first and/or second pools of antibodies, or antigen-binding fragments thereof, that have an IC_{50} of > 25 nM.
128. The method according to embodiments 123 or 124, wherein the method comprises the step of removing antibodies, or antigen-binding fragments thereof, from the first and/or second pools of antibodies, or antigen-binding fragments thereof, that have an IC_{50} of > 10 nM.
129. The method according to embodiments 123 or 124, wherein the method comprises the step of removing antibodies, or antigen-binding fragments thereof, from the first and/or second pools of antibodies, or antigen-binding fragments thereof, that have an IC_{50} of > 5 nM.
130. The method according to any one of embodiments 124-129, wherein the caga assay is an endogenous LTBP caga assay.
131. The method according to any one of embodiments 124-129, wherein the caga assay is a human LTBP overexpression caga assay.
132. The method according to any one of embodiments 124-129, wherein the caga assay is a murine LTBP overexpression caga assay.
133. A method for manufacturing a pharmaceutical composition comprising an antibody, or antigen-binding fragment thereof, that selectively binds a human LTBP1-proTGF β complex and/or a human LTBP3-proTGF β complex, the method comprising the steps of:
- selecting an antibody or antigen-binding fragment thereof that preferentially inhibits matrix-associated TGF β 1 over immune cell-associated TGF β 1;
 - producing the antibody in a cell culture comprising cells expressing the antibody, wherein the cell culture has a volume of 250L or greater,
 - optionally further comprising the step of purifying the antibody from the cell culture; and
 - further optionally comprising the step of formulating the purified antibody into a pharmaceutical composition.

134. The method according to 133, wherein the pharmaceutical composition is formulated for subcutaneous administration.

135. The method according to 133, wherein the selection step further comprises selecting an antibody or antigen-binding fragment that has $t_{1/2}$ of 45 minutes or longer for each of human LTBP1-proTGF β and human LTBP3-proTGF β complexes and optionally $t_{1/2}$ of 5 minutes or less for human GARP-proTGF β complex, as measured by SPR.

136. A method for making an antibody, or an antigen-binding fragment thereof, the method comprising steps of:

i) providing an antibody or a fragment that comprises at least three heavy chain CDR sequences of (CDR-H1) SEQ ID NO:94, (CDR-H2) SEQ ID NO:95, and (CDR-H3) SEQ ID NO:96; and

ii) subjecting the antibody or fragment of step (i) to affinity maturation and/or optimization, so as to provide an affinity matured and/or optimized antibody or fragment.

137. The method according to embodiment 136, wherein the antibody, or antigen-binding fragment thereof, further comprises light chain CDR sequences of (CDR-L1) SEQ ID NO:97, (CDR-L2) SEQ ID NO:98, and (CDR-L3) SEQ ID NO:99.

138. The method according to embodiments 136 or 137, wherein the antibody or fragment comprises a variable heavy chain region having an amino acid sequence as set forth in SEQ ID NO: 88.

139. The method according to any one of embodiments 136-138, wherein the antibody or fragment comprises a variable light chain region having an amino acid sequence as set forth in SEQ ID NO: 89.

140. A method for making an antibody, or an antigen-binding fragment thereof, the method comprising steps of:

i) providing an antibody or a fragment that comprises at least three CDR sequences of (CDR-H1) SEQ ID NO:100, (CDR-H2) SEQ ID NO:101, and (CDR-H3) SEQ ID NO:102; and

ii) subjecting the antibody or fragment of step (i) to affinity maturation and/or optimization, so as to provide an affinity matured and/or optimized antibody or fragment.

141. The method of embodiment 140, wherein the antibody, or antigen-binding fragment thereof, further comprises light chain CDR sequences of (CDR-L1) SEQ ID NO:103, (CDR-L2) SEQ ID NO:104, and (CDR-L3) SEQ ID NO:105.

142. The method according to embodiments 140 or 141, wherein the antibody or fragment comprises a variable heavy chain region having an amino acid sequence as set forth in SEQ ID NO: 106.

143. The method according to any one of embodiments 140-142, wherein the antibody or fragment comprises a variable light chain region having an amino acid sequence as set forth in SEQ ID NO: 107.

144. A method for making an antibody, or an antigen-binding fragment thereof, the method comprising steps of:

i) providing an antibody or a fragment that comprises at least three heavy chain CDR sequences of (CDR-H1) SEQ ID NO:108, (CDR-H2) SEQ ID NO:109, and (CDR-H3) SEQ ID NO:110 and

ii) subjecting the antibody or fragment of step (i) to affinity maturation and/or optimization, so as to provide an affinity matured and/or optimized antibody or fragment.

145. The method according to embodiment 144, wherein the antibody, or antigen-binding fragment thereof, further comprises light chain CDR sequences of (CDR-L1) SEQ ID NO:111, (CDR-L2) SEQ ID NO:112, and (CDR-L3) SEQ ID NO:113.

146. The method according to embodiments 144 or 145, wherein the antibody or fragment comprises a variable heavy chain region having an amino acid sequence as set forth in SEQ ID NO: 114.

147. The method according to any one of embodiments 144-146, wherein the antibody or fragment comprises a variable light chain region having an amino acid sequence as set forth in SEQ ID NO: 115.

148. A method for making an antibody, or an antigen-binding fragment thereof, the method comprising steps of:

i) providing an antibody or a fragment that comprises at least three heavy chain CDR sequences of (CDR-H1) SEQ ID NO:116, (CDR-H2) SEQ ID NO:117, and (CDR-H3) SEQ ID NO:118 and

ii) subjecting the antibody or fragment of step (i) to affinity maturation and/or optimization, so as to provide an affinity matured and/or optimized antibody or fragment.

149. The method according to embodiment 148, wherein the antibody, or antigen-binding fragment thereof, further comprises light chain CDR sequences of (CDR-L1) SEQ ID NO:119, (CDR-L2) SEQ ID NO:120, and (CDR-L3) SEQ ID NO:121.

150. The method according to embodiments 148 or 149, wherein the antibody or fragment comprises a variable heavy chain region having an amino acid sequence as set forth in SEQ ID NO: 122.

151. The method according to any one of embodiments 148-150, wherein the antibody or fragment comprises a variable light chain region having an amino acid sequence as set forth in SEQ ID NO: 123.

152. The method according to any one of embodiments 136-151, wherein step (ii) comprises light-chain shuffling.

153. The method according to any one of embodiments 136-152, wherein step (ii) comprises CDR-H1/H2 diversification.

154. The method according to any one of embodiments 136-153, wherein step (ii) comprises CDR-L1/L2 diversification.

155. The method according to any one of embodiments 136-154, wherein step (ii) comprises mutagenesis within any one of the CDRs, variable regions, and/or constant regions.

156. The method according to embodiment 155, wherein the mutagenesis is within a CDR.

157. The method according to any one of embodiments 155 and 156, wherein the mutagenesis is within a CDR-H3.

158. The method according to any one of embodiments 155-157, wherein the mutagenesis is within a CDR-L1.
159. The method according to any one of embodiments 155-158, wherein the mutagenesis is within a CDR-L2.
160. The method according to any one of embodiments 155-159, wherein the mutagenesis is within a CDR-L3.
161. The method according to embodiment 155, wherein the mutagenesis is within a variable region.
162. The method according to embodiment 155, wherein the mutagenesis is within a constant region.
163. The method of any one of embodiments 136-162, wherein the method further comprises a step of:
selecting affinity matured and/or optimized antibodies, or antigen-binding fragments thereof, that do not bind GARP-proTGF β 1, LRRC33-proTGF β 1, mature TGF β 1, GARP-proTGF β 2, LRRC33-proTGF β 2, mature TGF β 2, GARP-proTGF β 3, LRRC33-proTGF β 3, mature TGF β 3, or any combinations thereof.
164. The method of any one of embodiments 136-163, wherein the method further comprises a step of:
determining or confirming isoform-specificity of the affinity matured and/or optimized antibodies or fragments.
165. The method according to any one of embodiments 136-164, wherein the method further comprises a step of:
selecting antibodies or fragments that are cross-reactive to human and rodent antigens.
166. The method according to any one of embodiments 136-165, wherein the method further comprises a step of:
generating a fully human or humanized antibody or fragment, of the affinity matured and/or optimized antibody or fragment.

167. The method according to any one of embodiments 136-166, wherein the method further comprises a step of:
determining affinity of the affinity matured and/or optimized antibodies, or fragments, to human LTBP1-proTGF β 1 and/or human LTBP3-proTGF β 1.
168. The method according to embodiments 136-167, wherein the method further comprises a step of: selecting affinity matured and/or optimized antibodies, or fragments, that bind to human LTBP1-proTGF β 1 and/or human LTBP3-proTGF β 1 with a K_D of < 100 nM, as measured by Bio-Layer Interferometry (BLI).
169. The method according to embodiment 136-168, wherein the method further comprises a step of: selecting affinity matured and/or optimized antibodies, or fragments, that bind to human LTBP1-proTGF β 1 and/or human LTBP3-proTGF β 1 with a K_D of < 50 nM, as measured by Bio-Layer Interferometry (BLI).
170. The method according to embodiment 136-169, wherein the method further comprises a step of: selecting affinity matured and/or optimized antibodies, or fragments, that bind to human LTBP1-proTGF β 1 and/or human LTBP3-proTGF β 1 with a K_D of < 25 nM, as measured by Bio-Layer Interferometry (BLI).
171. The method according to embodiment 136-170, wherein the method further comprises a step of: selecting affinity matured and/or optimized antibodies, or fragments, that bind to human LTBP1-proTGF β 1 and/or human LTBP3-proTGF β 1 with a K_D of < 10 nM, as measured by Bio-Layer Interferometry (BLI).
172. The method according to any one of embodiments 136-171, wherein the method further comprises a step of:
determining affinity of the affinity matured and/or optimized antibodies or fragments, to mouse LTBP1-proTGF β 1 and/or mouse LTBP3-proTGF β 1.
173. The method according to embodiment 136-172, wherein the method further comprises a step of: selecting affinity matured and/or optimized antibodies, or fragments, that bind to mouse LTBP1-proTGF β 1 and/or mouse LTBP3-proTGF β 1 with a K_D of < 100 nM, as measured by Bio-Layer Interferometry (BLI).

174. The method according to embodiment 136-173, wherein the method further comprises a step of: selecting affinity matured and/or optimized antibodies, or fragments, that bind to mouse LTBP1-proTGF β 1 and/or mouse LTBP3-proTGF β 1 with a K_D of < 50 nM, as measured by Bio-Layer Interferometry (BLI).
175. The method according to embodiment 136-174, wherein the method further comprises a step of: selecting affinity matured and/or optimized antibodies, or fragments, that bind to mouse LTBP1-proTGF β 1 and/or mouse LTBP3-proTGF β 1 with a K_D of < 25 nM, as measured by Bio-Layer Interferometry (BLI).
176. The method according to embodiment 136-175, wherein the method further comprises a step of: selecting affinity matured and/or optimized antibodies, or fragments, that bind to mouse LTBP1-proTGF β 1 and/or mouse LTBP3-proTGF β 1 with a K_D of < 10 nM, as measured by Bio-Layer Interferometry (BLI).
177. The method according to any one of embodiments 136-176, wherein the method further comprises a step of:
selecting affinity matured and/or optimized antibodies, or fragments, that do not bind mouse LTBP1-proTGF β 1 and/or mouse LTBP3-proTGF β 1.
178. The method according to any one of embodiments 136-177, wherein the method further comprises a step of:
determining the IC_{50} of the affinity matured and/or optimized antibodies, or fragments, as measured by a suitable functional in vitro cell-based assay.
179. The method according to embodiments 178, wherein the suitable functional in vitro cell-based assay is a caga assay.
180. The method according to any one of embodiments 136-179, wherein the method comprises the step of:
selecting affinity matured and/or optimized antibodies, or fragments, that have an IC_{50} of < 100 nM.
181. The method according to any one of embodiments 136-179, wherein the method comprises the step of:

selecting affinity matured and/or optimized antibodies, or fragments, that have an IC_{50} of < 50 nM.

182. The method according to embodiments 136-179, wherein the method comprises the step of: selecting affinity matured and/or optimized antibodies, or fragments, that have an IC_{50} of < 25 nM.

183. The method according to embodiments 136-179, wherein the method comprises the step of: selecting affinity matured and/or optimized antibodies, or fragments, that have an IC_{50} of < 10 nM.

184. The method according to embodiments 136-179, wherein the method comprises the step of: selecting affinity matured and/or optimized antibodies, or fragments, that have an IC_{50} of < 5 nM.

185. The method according to any one of embodiments 179-184, wherein the caga assay is an endogenous LTBP caga assay.

186. The method according to any one of embodiments 179-184, wherein the caga assay is a human LTBP overexpression caga assay.

187. The method according to any one of embodiments 179-184, wherein the caga assay is a murine LTBP overexpression caga assay.

188. The antibody or the fragment, use thereof, or related methods thereof, according to any one of the preceding embodiments,

a) which comprises the following variable heavy and/or variable light chain sequences, or a variant thereof:

i) the variable heavy chain sequences

QVQLVESGGGVVQPGRSLRLSCAASGFTFRSYVMHWVRQAPGKGLEWVAVISHEGSLKYY
ADSVKGRFTISRDN SKNTLYQMNSLRAEDTAVYYCAVPRIAARRGGFGYWGQGLTLTVSS

or

QVQLVESGGGVVQPGRSLRLSCAASGFTFRSYVMHWVRQAPGKGLEWVAVISHEGSLKYY
ADSVKGRFTISRDN SKNTLYQMNSLRAEDTAVYYCARPRIAARRGGFGYWGQGLTLTVSS,

wherein optionally the variant is at least 85% identical, 90% identical, or 95% identical to the corresponding variable heavy chain sequence.

ii) the variable light chain sequences

NFMLTQPHSVSESPGKTVTISCTRSSGNIDNNYVQWYQQRPGSSPTTVIYEDNQRPSGVPDRF
SGSIDSSSNSASLTISGLKTEDEADYYCQSYDSDNQGVVFGGGTKLTVL or
NFMLTQPHSVSESPGKTVTISCTRSSGNIDNNYVQWYQQRPGSSPTTVIYEDNQRPSGVPDRF
SGSIDSSSNSASLTISGLKTEDEADYYCQSYDSDNQGVVFGGGTKLTVL, wherein optionally
the variant is at least 85% identical, 90% identical, or 95% identical to the corresponding variable
light chain sequence;

b) which has the following variable heavy chain CDR sequences, or variant thereof:

i) CDR-H1 of FTFRSYVMH optionally comprising one or more amino acid changes;

ii) CDR-H2 of VISHEGSLKYYADSVKG optionally comprising one or more amino acid
changes; and/or;

iii) CDR-H3 of AVPRIAARRGGFGY or ARPRIAARRGGFGY optionally comprising one
or more amino acid changes; and/or;

c) which has the following variable light chain CDR sequences, or variant thereof:

i) CDR1-L1 of TRSSGNIDNNYVQ optionally comprising one or more amino acid changes;

ii) CDR-L2 of EDNQRPS optionally comprising one or more amino acid changes; and/or;

iii) CDR-L3 of QSYDSDNQGVV optionally comprising one or more amino acid changes.

A1. An isolated antibody that specifically binds a human LTBP1-proTGF β complex and/or a
human LTBP3-proTGF β complex, and does not bind a human GARP-proTGF β complex;
wherein the antibody does not bind mature TGF β 1, mature TGF β 2 or mature TGF β 3;
wherein the antibody is a fully human or humanized antibody, or antigen-binding fragment
thereof,

wherein the antibody comprises at least three CDRs selected from the following, optionally
comprising up to 3 amino acid changes for each of the CDRs:

CDR-H1: SEQ ID NO:100;

CDR-H2: SEQ ID NO:101;

CDR-H3: SEQ ID NO:102;

CDR-L1: SEQ ID NO:103;

CDR-L2: SEQ ID NO:104; and,

CDR-L3: SEQ ID NO:105.

A1.1. An isolated antibody that specifically binds a human LTBP1-proTGF β complex and/or a
human LTBP3-proTGF β complex, and does not bind a human GARP-proTGF β complex;

wherein the antibody does not bind mature TGF β 1, mature TGF β 2 or mature TGF β 3;

wherein the antibody is a fully human or humanized antibody, or antigen-binding fragment thereof,

wherein the antibody comprises at least three of the following six CDRs:

- a) CDR-H1: SEQ ID NO:100, with the proviso that:
 - i. the serine residue at position 4 of SEQ ID NO:100 may be substituted with a histidine;
 - ii. the serine residue at position 7 of SEQ ID NO:100 may be substituted with an alanine or glycine; and/or,
 - iii. the glycine residue at position 11 of SEQ ID NO:100 may be substituted with a threonine, serine, histidine, leucine, isoleucine, asparagine, valine, or alanine;
- b) CDR-H2: SEQ ID NO:101, with the proviso that:
 - i. the serine residue at position 3 of SEQ ID NO:101 may be substituted with an alanine;
 - ii. the glycine residue at position 6 of SEQ ID NO:101 may be substituted with an alanine or serine; and/or,
 - iii. the serine residue at position 7 of SEQ ID NO:101 may be substituted with a threonine;
- c) CDR-H3: SEQ ID NO:102, optionally comprising up to three amino acid changes;
- d) CDR-L1: SEQ ID NO:103, optionally comprising up to three amino acid changes;
- e) CDR-L2: SEQ ID NO:104, optionally comprising up to three amino acid changes; and,
- f) CDR-L3: SEQ ID NO:105, optionally comprising up to three amino acid changes.

A2. An isolated antibody that specifically binds a human LTBP1-proTGF β complex and/or a human LTBP3-proTGF β complex, and does not bind a human GARP-proTGF β complex; wherein the antibody does not bind mature TGF β 1, mature TGF β 2 or mature TGF β 3; wherein the antibody is a fully human or humanized antibody, or antigen-binding fragment thereof;

wherein the antibody comprises a heavy chain variable region having an amino acid sequence that is at least 90% identical to SEQ ID NO: 106; and/or,

wherein the antibody comprises a variable light chain variable region having an amino acid sequence that is at least 90% identical to SEQ ID NO: 107.

A3. The antibody according to embodiment A2, wherein the antibody comprises:

a heavy chain variable region having an amino acid sequence that is at least 90% identical to SEQ ID NO: 106; and

a light chain variable region having an amino acid sequence that is at least 90% identical to SEQ ID NO: 107.

A3.1 An isolated antibody that specifically binds a human LTBP1-proTGF β complex and/or a human LTBP3-proTGF β complex, and does not bind a human GARP-proTGF β complex;

wherein the antibody does not bind mature TGF β 1, mature TGF β 2 or mature TGF β 3;

wherein the antibody is a fully human or humanized antibody, or antigen-binding fragment thereof,

wherein the antibody comprises at least three of the following six CDRs:

- a) CDR-H1: SEQ ID NO:94, with the proviso that:
 - i. the threonine residue at position 2 of SEQ ID NO:94 may be substituted with an alanine;
 - ii. the asparagine residue at position 4 of SEQ ID NO:94 may be substituted with an alanine, tyrosine, aspartate, serine, arginine, or histidine;
 - iii. the asparagine residue at position 5 of SEQ ID NO:94 may be substituted with a glutamine, serine, glycine, lysine, glutamate, arginine, or histidine;
 - iv. the tyrosine residue at position 6 of SEQ ID NO:94 may be substituted with a arginine;
 - v. the proline residue at position 7 of SEQ ID NO:94 may be substituted with a glycine, alanine, leucine, serine, asparagine, valine, aspartate, or glutamine;
 - vi. the isoleucine residue at position 8 of SEQ ID NO:94 may be substituted with a methionine or leucine; and/or,
 - vii. the histidine residue at position 9 of SEQ ID NO:94 may be substituted with a phenylalanine, tyrosine, asparagine, or serine;
- b) CDR-H2: SEQ ID NO:95, optionally comprising up to six amino acid changes;
- c) CDR-H3: SEQ ID NO:96, optionally comprising up to three amino acid changes;
- d) CDR-L1: SEQ ID NO:97, optionally comprising up to three amino acid changes;
- e) CDR-L2: SEQ ID NO:98, optionally comprising up to three amino acid changes; and,
- f) CDR-L3: SEQ ID NO:99, optionally comprising up to three amino acid changes.

A3.2. An isolated antibody that specifically binds a human LTBP1-proTGF β complex and/or a human LTBP3-proTGF β complex, and does not bind a human GARP-proTGF β complex;

wherein the antibody does not bind mature TGF β 1, mature TGF β 2 or mature TGF β 3;

wherein the antibody is a fully human or humanized antibody, or antigen-binding fragment thereof;

wherein the antibody comprises a heavy chain variable region having an amino acid sequence that is at least 90% identical to SEQ ID NO: 88; and/or,
wherein the antibody comprises a variable light chain variable region having an amino acid sequence that is at least 90% identical to SEQ ID NO: 89.

A3.3. The antibody according to embodiment A3.2, wherein the antibody comprises:

a heavy chain variable region having an amino acid sequence that is at least 90% identical to SEQ ID NO: 88; and

a light chain variable region having an amino acid sequence that is at least 90% identical to SEQ ID NO: 89.

A3.4. An isolated antibody that specifically binds a human LTBP1-proTGF β complex and/or a human LTBP3-proTGF β complex, and does not bind a human GARP-proTGF β complex;

wherein the antibody does not bind mature TGF β 1, mature TGF β 2 or mature TGF β 3;

wherein the antibody is a fully human or humanized antibody, or antigen-binding fragment thereof,

wherein the antibody comprises at least three of the following six CDRs:

- a) CDR-H1: SEQ ID NO:1, with the proviso that:
 - i. the threonine residue at position 4 of SEQ ID NO:1 may be substituted with a histidine, lysine, phenylalanine, or glycine;
 - ii. the serine residue at position 5 of SEQ ID NO:1 may be substituted with a leucine; and/or,
 - iii. the serine residue at position 9 of SEQ ID NO:1 may be substituted with an alanine;
- b) CDR-H2: SEQ ID NO:2, with the proviso that:
 - i. the serine residue at position 3 of SEQ ID NO:2 may be substituted with an aspartate or asparagine;
 - ii. the tyrosine residue at position 5 of SEQ ID NO:2 may be substituted with a histidine;
 - iii. the asparagine residue at position 6 of SEQ ID NO:2 may be substituted with a serine;
 - iv. the asparagine residue at position 8 of SEQ ID NO:2 may be substituted with a phenylalanine, leucine, alanine, tyrosine, aspartate, or serine; and/or,
 - v. the asparagine residue at position 10 of SEQ ID NO:2 may be substituted with a aspartate or alanine;
- c) CDR-H3: SEQ ID NO:3, optionally comprising up to three amino acid changes;

- d) CDR-L1: SEQ ID NO:4, optionally comprising up to three amino acid changes;
- e) CDR-L2: SEQ ID NO:5, optionally comprising up to three amino acid changes; and,
- f) CDR-L3: SEQ ID NO:6, optionally comprising up to three amino acid changes.

A3.5. An isolated antibody that specifically binds a human LTBP1-proTGF β complex and/or a human LTBP3-proTGF β complex, and does not bind a human GARP-proTGF β complex;
wherein the antibody does not bind mature TGF β 1, mature TGF β 2 or mature TGF β 3;
wherein the antibody is a fully human or humanized antibody, or antigen-binding fragment thereof;
wherein the antibody comprises a heavy chain variable region having an amino acid sequence that is at least 90% identical to SEQ ID NO: 7; and/or,
wherein the antibody comprises a variable light chain variable region having an amino acid sequence that is at least 90% identical to SEQ ID NO: 8.

A3.6. The antibody according to embodiment A3.5, wherein the antibody comprises:
a heavy chain variable region having an amino acid sequence that is at least 90% identical to SEQ ID NO: 7; and
a light chain variable region having an amino acid sequence that is at least 90% identical to SEQ ID NO: 8.

A4. The antibody according to any one of embodiments A1-A3.6, wherein the antibody does not compete with any one of the antibodies SR-Ab1, SR-Ab2 or SR-Ab10 for binding to a human LTBP1-proTGF β 1 complex.

A5. The antibody according to any one of embodiments A1-A4, wherein the antibody is an IgG4 or IgG1 subtype.

A5.1 The antibody according to any one of embodiments A1-A5, wherein the antibody is specific for human LTBP1-TGF β 1 complex.

A5.2 The antibody according to any one of embodiments A1-A5.1, wherein the antibody is specific for human LTBP3-TGF β 1 complex.

A5.3 The antibody according to any one of embodiments A1-A5.2, wherein the antibody is specific for human LTBP1-TGF β 1 complex and human LTBP3-TGF β 1 complex.

A5.4 The antibody according to any one of embodiments A1-A5.3, wherein the antibody does not bind a human GARP-TGF β 1 complex or GARP-TGF β 3 complex.

A6. A pharmaceutical composition comprising the antibody of any one of embodiments A1-A5.4 and a pharmaceutically acceptable excipient.

A7. The pharmaceutical composition according to embodiment A6, which is prepared for intravenous administration or subcutaneous administration.

A8. A composition comprising a multi-dose vial containing the pharmaceutical composition of embodiment A6 or A7.

A9. A composition comprising a single-dose syringe containing the pharmaceutical composition of embodiment A6 or A7, optionally wherein the syringe is a disposable syringe.

A10. The composition of any one of embodiments A6-A9 for use in a method for the treatment of a fibrotic condition in a human subject, wherein the treatment comprises administration of the composition to the subject in an amount effective to treat the fibrotic disorder.

A11. The composition for use according to embodiment A10, wherein the fibrotic disorder is an organ fibrosis.

A12. The composition for use according to embodiment A11, wherein the organ fibrosis is an advanced organ fibrosis.

A13. The composition for use according to embodiment A10 or A11, wherein the organ fibrosis is selected from the group consisting of:
kidney fibrosis, liver fibrosis, lung fibrosis, cardiac fibrosis, pancreatic fibrosis, skin fibrosis, scleroderma, muscle fibrosis, uterine fibrosis and endometriosis.

A14. The composition for use according to embodiment A10, wherein:
a) the fibrotic disorder comprises chronic inflammation;
b) the subject benefits from immune suppression;
c) the subject has or is at risk of developing an autoimmune disease;
d) the subject is a candidate for an allograft transplant; and/or,
e) the subject has received an allograft transplant.

A15. The composition for use according to embodiment A10, wherein the fibrotic disorder comprising chronic inflammation is a muscular dystrophy, multiple sclerosis (MS), or Cystic Fibrosis (CF).

A16. The composition for use according to embodiment A15, wherein the muscular dystrophy is Duchenne muscular dystrophy (DMD).

A17. The composition for use according to embodiment A15, wherein the MS comprises perivascular fibrosis.

A18. The composition for use according to embodiment A13, wherein the lung fibrosis is idiopathic pulmonary fibrosis (IPF).

A19. The composition for use according to embodiment A13, wherein the subject has chronic kidney disease (CKD).

A20. The composition for use according to embodiment A13, wherein the subject has nonalcoholic steatohepatitis (NASH).

A21. The composition for use according to any one of embodiments A10-A20, wherein the antibody is administered to the subject at a dosage of between 0.1 and 30 mg/kg.

A22. The composition for use according to embodiment A21, wherein the antibody is administered twice a week, once a week, once every 2 weeks, once every 3 weeks, once a month, or every other month.

A23. The composition for use according to embodiment A21, wherein a therapeutic regimen comprises an initial phase of a therapy and a subsequent phase of the therapy,
wherein the subject receives a loading dose during the initial phase followed by a maintenance dose during the subsequent phase.

A24. The composition for use according to embodiment A21, wherein the loading dose is between 2-30 mg/kg, and the maintenance dose is between 0.1-20 mg/kg.

A25. The composition for use according to embodiment A21, wherein the loading dose is administered to the subject twice a week or once a week.

A26. The composition for use according to embodiment A21, wherein the maintenance dose is administered to the subject once every 2-8 weeks.

A27. The composition for use according to any one of embodiments A10-A26, wherein the method further comprises testing or confirming expression of TGF β 1, LTBP1 or LTBP3 in a biological sample collected from the subject.

A28. A method for making a composition of any one of embodiments A6-A9, comprising an antibody, or antigen-binding fragment thereof, that specifically binds a human LTBP1-proTGF β complex and/or a human LTBP3-proTGF β complex, and does not bind a human GARP-proTGF β complex; wherein the antibody or fragment inhibits TGF β 1 but does not inhibit TGF β 2 or TGF β 3, the method comprising steps of:

- i) providing at least one antigen comprising LTBP1-proTGF β 1 and/or LTBP3-proTGF β 1,
- ii) selecting a first pool of antibodies or fragments that specifically bind the at least one antigen of step (i) so as to provide specific binders of LTBP1-proTGF β 1 and/or LTBP3-proTGF β 1;
- iii) selecting a second pool of antibodies or fragments that inhibit activation of TGF β 1, so as to generate specific inhibitors of TGF β 1 activation; and
- iv) formulating an antibody or fragment that is present in the first pool of antibodies and the second pool of antibodies into a pharmaceutical composition, thereby making the composition comprising the antibody or fragment.

A29. The method of embodiment A28, wherein the method further comprises a step of:
removing from the first pool of antibodies, or fragments, any antibodies or fragments that bind GARP-proTGF β 1, LRRC33-proTGF β 1, mature TGF β 1, GARP-proTGF β 2, LRRC33-proTGF β 2, mature TGF β 2, GARP-proTGF β 3, LRRC33-proTGF β 3, mature TGF β 3, or any combinations thereof.

A30. The method of embodiment A28 or A29, wherein the method further comprises a step of:
determining or confirming isoform-specificity of the antibodies or fragments selected in steps (ii) and/or (iii).

A31. The method of any one of embodiments A28-A30, wherein the method further comprises a step of:
selecting antibodies or fragments that are cross-reactive to human and rodent antigens.

A32. The method of any one of embodiments A28-A31, wherein the method further comprises a step of:

generating a fully human or humanized antibody or fragment, of the antibody or fragment that is presented in the first pool of antibodies and the second pool of antibodies.

A33. The method of any one of embodiments A28-A32, wherein the method further comprises a step of:

subjecting the antibody or fragment that is present in the first pool of antibodies and the second pool of antibodies to affinity maturation and/or optimization, so as to provide an affinity matured and/or optimized antibody or fragment.

A34. A method for making an antibody, or an antigen-binding fragment thereof, the method comprising steps of:

i) providing an antibody or a fragment that comprises at least three CDR sequences of (CDR-H1) SEQ ID NO:1, (CDR-H2) SEQ ID NO:2, and (CDR-H3) SEQ ID NO:3; and

ii) subjecting the antibody or fragment of step (i) to affinity maturation and/or optimization, so as to provide an affinity matured and/or optimized antibody or fragment.

A35. A method for making an antibody, or an antigen-binding fragment thereof, the method comprising steps of:

i) providing an antibody or a fragment that comprises at least three CDR sequences of (CDR-H1) SEQ ID NO:94, (CDR-H2) SEQ ID NO:95, and (CDR-H3) SEQ ID NO:96; and

ii) subjecting the antibody or fragment of step (i) to affinity maturation and/or optimization, so as to provide an affinity matured and/or optimized antibody or fragment.

A36. A method for making an antibody, or an antigen-binding fragment thereof, the method comprising steps of:

i) providing an antibody or a fragment that comprises at least three CDR sequences of (CDR-H1) SEQ ID NO:100, (CDR-H2) SEQ ID NO:101, and (CDR-H3) SEQ ID NO:102; and

ii) subjecting the antibody or fragment of step (i) to affinity maturation and/or optimization, so as to provide an affinity matured and/or optimized antibody or fragment.

A37. A method for making an antibody, or an antigen-binding fragment thereof, the method comprising steps of:

i) providing an antibody or a fragment that comprises a heavy chain variable region having an amino acid sequence set forth in SEQ ID NO: 7; and

ii) subjecting the antibody or fragment of step (i) to affinity maturation and/or optimization, so as to provide an affinity matured and/or optimized antibody, or fragment.

A38. A method for making antibody, or an antigen-binding fragment thereof, the method comprising steps of:

i) providing an antibody or a fragment that comprises a heavy chain variable region having an amino acid sequence set forth in SEQ ID NO: 88; and

ii) subjecting the antibody or fragment of step (i) to affinity maturation and/or optimization, so as to provide an affinity matured and/or optimized antibody or fragment.

A39. A method for making an antibody, or an antigen-binding fragment thereof, the method comprising steps of:

i) providing an antibody or a fragment that comprises a heavy chain variable region having an amino acid sequence set forth in SEQ ID NO: 106; and

ii) subjecting the antibody or fragment of step (i) to affinity maturation and/or optimization, so as to provide an affinity matured and/or optimized antibody or fragment.

A40. The method of any one of embodiments A34-A39, wherein step (ii) comprises mutagenesis.

A41. The method of embodiment A40, wherein the mutagenesis is within a CDR.

A42. The method of embodiment A40, wherein the mutagenesis is within a variable region.

A43. The method of embodiment A40, wherein the mutagenesis is within a constant region.

A44. The method of any one of embodiments A34-A43, wherein step (ii) comprises light-chain shuffling.

CLAIMS

What is claimed is:

1. An isolated antibody that specifically binds a human LTBP1-proTGF β complex and/or a human LTBP3-proTGF β complex, and does not bind a human GARP-proTGF β complex;
wherein the antibody does not bind mature TGF β 1, mature TGF β 2 or mature TGF β 3;
wherein the antibody is a fully human or humanized antibody, or antigen-binding fragment thereof,
wherein the antibody comprises at least three of the following six CDRs:
 - a) CDR-H1: SEQ ID NO:94, with the proviso that:
 - i. the threonine residue at position 2 of SEQ ID NO:94 may be substituted with an alanine;
 - ii. the asparagine residue at position 4 of SEQ ID NO:94 may be substituted with an alanine, tyrosine, aspartate, serine, arginine, or histidine;
 - iii. the asparagine residue at position 5 of SEQ ID NO:94 may be substituted with a glutamine, serine, glycine, lysine, glutamate, arginine, or histidine;
 - iv. the tyrosine residue at position 6 of SEQ ID NO:94 may be substituted with a arginine;
 - v. the proline residue at position 7 of SEQ ID NO:94 may be substituted with a glycine, alanine, leucine, serine, asparagine, valine, aspartate, or glutamine;
 - vi. the isoleucine residue at position 8 of SEQ ID NO:94 may be substituted with a methionine or leucine; and/or,
 - vii. the histidine residue at position 9 of SEQ ID NO:94 may be substituted with a phenylalanine, tyrosine, asparagine, or serine;
 - b) CDR-H2: SEQ ID NO:95, comprising up to six amino acid changes;
 - c) CDR-H3: SEQ ID NO:96, comprising up to three amino acid changes;
 - d) CDR-L1: SEQ ID NO:97, comprising up to three amino acid changes;
 - e) CDR-L2: SEQ ID NO:98, comprising up to three amino acid changes; and,
 - f) CDR-L3: SEQ ID NO:99, comprising up to three amino acid changes.
2. An isolated antibody that specifically binds a human LTBP1-proTGF β complex and/or a human LTBP3-proTGF β complex, and does not bind a human GARP-proTGF β complex;
wherein the antibody does not bind mature TGF β 1, mature TGF β 2 or mature TGF β 3;
wherein the antibody is a fully human or humanized antibody, or antigen-binding fragment thereof;

wherein the antibody comprises a heavy chain variable region having an amino acid sequence that is at least 90% identical to SEQ ID NO: 88; and/or,

wherein the antibody comprises a variable light chain variable region having an amino acid sequence that is at least 90% identical to SEQ ID NO: 89.

3. An antibody, or antigen-binding fragment thereof, comprising at least three of the following six CDRs:

a) CDR-H1 comprising the amino acid sequence FTF(X₁)(X₂)YVMH, wherein: X₁ is S or R; and X₂ is G or S;

b) CDR-H2 comprising the amino acid sequence (X₁)ISHEG(X₂)(X₃)KYYADSVKG, wherein: X₁ is V or S; X₂ is S or G; and X₃ is F or L; and

c) CDR-H3 comprising the amino acid sequence (X₁)(X₂)P(X₃)(X₄)(X₅)(X₆)RRGG(X₇)(X₈)(X₉), wherein: X₁ is A or V; X₂ is R, V, G or K; X₃ is R, H or L; X₄ is I, V or G; X₅ is A, S, or L; X₆ is A or V; X₇ is F or Y; X₈ is D, G, R, or S; and, X₉ is Y, G, R, L, V, A or K.

d) CDR-L1 as set forth in SEQ ID NO:97, comprising up to three amino acid changes;

e) CDR-L2 as set forth in SEQ ID NO:98, comprising up to three amino acid changes;

and

f) CDR-L3 as set forth in SEQ ID NO:99, comprising up to three amino acid changes.

4. The antibody, or antigen-binding fragment thereof, according to claim 3, wherein:

b) within CDR-H2; X₂ is S; and

c) within CDR-H3; X₁ is A; X₂ is R or V; X₃ is R; X₄ is I; X₅ is A or L; X₆ is A; X₇ is F; X₈ is G; and X₉ and Y.

5. The antibody according to any one of claims 1-4, wherein the antibody comprises:

a heavy chain variable region having an amino acid sequence that is at least 90% identical to SEQ ID NO: 88; and

a light chain variable region having an amino acid sequence that is at least 90% identical to SEQ ID NO: 89.

6. The antibody, or antigen-binding fragment thereof, according to any one of claims 3-5, wherein the up to three amino acid changes comprises up to two amino acid changes.

7. The antibody, or antigen-binding fragment thereof, according to any preceding claim, comprising all six CDRs.

8. The antibody according to any one of the preceding claims, wherein the antibody is specific for human LTBP1-TGF β 1 complex.
9. The antibody according to any one of the preceding claims, wherein the antibody is specific for human LTBP3-TGF β 1 complex.
10. The antibody according to any one of the preceding claims, wherein the antibody is specific for human LTBP1-TGF β 1 complex and human LTBP3-TGF β 1 complex.
11. An antibody, or antigen-binding fragment thereof, comprising the following six CDRs:
 - a) CDR-H1 comprising the amino acid sequence FTFRSYVMH;
 - b) CDR-H2 comprising the amino acid sequence VISHEGS(X₁)KYYADSVKG, wherein: X₁ is L or G; and
 - c) CDR-H3 comprising the amino acid sequence A(X₁)PRIAARRGGFG(X₂), wherein: X₁ is V, R or L; and X₂ is Y, S or T;
 - d) CDR-L1 comprising the amino acid sequence TRS(X₁)G(X₂)ID(X₃)NYVQ, wherein, X₁ is S or H; X₂ is N, L, S or A; and X₃ is N, D or Y;
 - e) CDR-L2 comprising the amino acid sequence ED(X₁)(X₂)RPS, wherein: X₁ is N, F or A; and X₂ is Q, I or V; and
 - f) CDR-L3 comprising the amino acid sequence Q(X₁)YD(X₂)(X₃)(X₄)Q(X₅)VV, wherein: X₁ is S or G; X₂ is S, F, Y, D, H or W; X₃ is N, D or S; X₄ is N, A, L, E or T; and X₅ is G, R, A or L.
12. The antibody, or antigen-binding fragment thereof, according to claim 11, wherein: within CDR-H3: X₁ is R or L.
13. The antibody, or antigen-binding fragment thereof, according to claim 12, wherein: within CDR-L3: X₂ is Y.
14. The antibody, or antigen-binding fragment thereof, according to claim 13, wherein: within CDR-L3: X₃ is D; and X₄ is T.
15. The antibody, or antigen-binding fragment thereof, according to claim 13, wherein: within CDR-L3: X₃ is D; X₄ is N; and X₅ is A.
16. The antibody, or antigen-binding fragment thereof, according claim 12, wherein:

within CDR-L1: X₁ is S or H; X₂ is N or A; and X₃ is N, D or Y;

within CDR-L2: X₁ is N or F; and X₂ is Q or V; and

within CDR-L3: X₁ is S or G; X₂ is S, Y, D or W; X₃ is D or S; X₄ is N, L or T; and X₅ is G, R, A or L.

17. The antibody, or antigen-binding fragment thereof, according to claim 16, wherein:
within CDR-L1: X₁ is S; X₂ is N; and X₃ is N or Y;
within CDR-L2: X₁ is N; and X₂ is Q or V; and
within CDR-L3: X₁ is S or G; X₂ is S, Y or W; X₃ is D; X₄ is N or T; and X₅ is G, R or A.
18. The antibody, or antigen-binding fragment thereof, according to claim 17, wherein:
within CDR-L3: X₁ is S; X₂ is S or Y; X₃ is D; X₄ is N or T; and X₅ is G, R or A.
19. The antibody, or antigen-binding fragment thereof, according to claim 18, wherein:
within CDR-L3: X₁ is S; X₂ is Y; X₃ is D; X₄ is N or T; and X₅ is G or A.
20. The antibody, or antigen-binding fragment thereof, according to claim 19, wherein:
within CDR-L3: X₁ is S; X₂ is Y; X₃ is D; X₄ is T; and X₅ is G.
21. The antibody, or antigen-binding fragment thereof, according to any one of claims 16-20, wherein:
 - a) CDR-H1 comprises the amino acid sequence of SEQ ID NO: 166;
 - b) CDR-H2 comprises the amino acid sequence of SEQ ID NO: 167;
 - c) CDR-H3 comprises the amino acid sequence of SEQ ID NO: 168;
 - d) CDR-L1 comprises the amino acid sequence of SEQ ID NO: 169;
 - e) CDR-L2 comprises the amino acid sequence of SEQ ID NO: 170; and
 - f) CDR-L3 comprises the amino acid sequence of SEQ ID NO: 171.
22. The antibody, or antigen-binding fragment thereof, according to any one of claims 11-21, which comprises:
 - a heavy chain variable region having an amino acid sequence that is at least 90% identical to SEQ ID NO: 318; and
 - a light chain variable region having an amino acid sequence that is at least 90% identical to SEQ ID NO: 319.

23. The antibody, or antigen-binding fragment thereof, according to any one of claims 11-22, which competes or cross-competes with an antibody having a heavy chain variable region sequence as set forth in SEQ ID NO: 318 and light chain variable region sequence as set forth in SEQ ID NO: 319.
24. An antibody, or antigen-binding fragment thereof, which selectively binds to human LTBP1-TGF β 1 complex and human LTBP3-TGF β 1 complex and competes or cross-competes with an antibody having a heavy chain variable region sequence as set forth in SEQ ID NO: 318 and light chain variable region sequence as set forth in SEQ ID NO: 319.
25. The antibody, or antigen-binding fragment thereof, according to any one of claims 11-24, which does not show detectable binding to a human GARP-proTGF β 1 complex, as measured by BLI, under the same assay conditions as used to measure binding to human LTBP1-proTGF β 1 complex and a human LTBP3-TGF β 1 complex.
26. The antibody, or antigen-binding fragment thereof, according to any one of claims 11-25, which binds a human LTBP1-proTGF β 1 complex and/or a human LTBP3-TGF β 1 complex with a K_D that is at least 50 times lower than the K_D when binding to a human GARP-proTGF β 1 complex under the same assay conditions.
27. The antibody, or antigen-binding fragment thereof, according to any one of claims 11-26, which does not show detectable binding to an LRRC33-proTGF β 1 complex, as measured by BLI, under the same assay conditions as used to measure binding to human LTBP1-proTGF β 1 complex and human LTBP3-TGF β 1 complex.
28. The antibody, or antigen-binding fragment thereof, according to any one of claims 11-27, wherein the antibody, or antigen-binding fragment thereof has a monovalent half-binding-time ($t_{1/2}$) of at least 45 minutes for each of hLTBP1-proTGF β 1 and hLTBP3-proTGF β 1 complexes, as measured by SPR.
29. The antibody, or antigen-binding fragment thereof, according to claim 28 which has a monovalent $t_{1/2}$ of less than 5 minutes for each of hGARP-proTGF β 1 and hLRRC33-proTGF β 1 complexes, as measured by SPR.
30. The antibody, or antigen-binding fragment thereof, according to any one of the preceding claims, which binds a human LTBP1-proTGF β 1 complex and a human LTBP3-TGF β 1 complex with a K_D of < 5 nM as measured by Bio-Layer Interferometry (BLI), optionally < 1 nM.

31. The antibody, or antigen-binding fragment thereof, according to any one of the preceding claims, which is cross-reactive with mouse LTBP1-proTGF β 1.
32. The antibody, or antigen-binding fragment thereof, according to any one of the preceding claims, which is cross-reactive with mouse LTBP3-proTGF β 1.
33. The antibody, or antigen-binding fragment thereof, according to any one of the preceding claims, wherein the antibody, or antigen-binding fragment thereof, binds a mouse LTBP1-proTGF β 1 complex with a K_D of < 10 nM as measured by Bio-Layer Interferometry (BLI).
34. The antibody, or antigen-binding fragment thereof, according to any one of the preceding claims, wherein the antibody, or antigen-binding fragment thereof, binds a mouse LTBP3-proTGF β 1 complex with a K_D of < 10 nM as measured by Bio-Layer Interferometry (BLI).
35. The antibody, or antigen-binding fragment thereof, according to any one of the preceding claims, wherein the antibody, or antigen-binding fragment thereof cross-reacts with human and murine LTBP1-proTGF β 1 and LTBP3-proTGF β 1 complexes, each with a K_D of < 5 nM or optionally < 1 nM.
36. The antibody, or antigen-binding fragment thereof, according to any one of the preceding claims, wherein the antibody is an IgG4 or IgG1 subtype, optionally wherein the antibody is a human IgG4 subtype and comprises a backbone substitution of Ser to Pro that produces an IgG1-like hinge.
37. A pharmaceutical composition comprising the antibody of any one of the preceding claims and a pharmaceutically acceptable excipient.
38. The pharmaceutical composition according to claim 37, which is prepared for intravenous administration or subcutaneous administration.
39. A composition comprising a multi-dose vial containing the pharmaceutical composition of claim 37 or 38.
40. A composition comprising a single-dose syringe containing the pharmaceutical composition of claim 37 or 38, optionally wherein the syringe is a disposable syringe.

41. The composition of any one of claims 37-40 for use in a method for the treatment of a fibrotic condition in a human subject, wherein the treatment comprises administration of the composition to the subject in an amount effective to treat the fibrotic disorder.
42. The composition for use according to claim 41, wherein the fibrotic disorder is an organ fibrosis.
43. The composition for use according to claim 42, wherein the organ fibrosis is an advanced organ fibrosis.
44. The composition for use according to claim 42 or 43, wherein the organ fibrosis is selected from the group consisting of: kidney fibrosis, liver fibrosis, lung fibrosis, cardiac fibrosis, pancreatic fibrosis, skin fibrosis, scleroderma, muscle fibrosis, uterine fibrosis and endometriosis.
45. The composition for use according to claim 44, wherein the lung fibrosis is idiopathic pulmonary fibrosis (IPF).
46. The composition for use according to claim 44, wherein the subject has chronic kidney disease (CKD).
47. The composition for use according to claim 44, wherein the liver fibrosis is associated with nonalcoholic steatohepatitis (NASH).
48. The composition for use according to any one of claims 41-47, wherein the antibody is administered to the subject at a dosage of between 0.1 and 30 mg/kg.
49. The composition for use according to claim 48, wherein the antibody is administered twice a week, once a week, once every 2 weeks, once every 3 weeks, once a month, or every other month.
50. The composition for use according to claim 48, wherein a therapeutic regimen comprises an initial phase of a therapy and a subsequent phase of the therapy, wherein the subject receives a loading dose during the initial phase followed by a maintenance dose during the subsequent phase.
51. The composition for use according to claim 50, wherein the loading dose is between 2-30 mg/kg, and the maintenance dose is between 0.1-20 mg/kg.

52. The composition for use according to claim 50 or 51, wherein the loading dose is administered to the subject twice a week or once a week.
53. The composition for use according to any one of claims 50-52, wherein the maintenance dose is administered to the subject once every 2-8 weeks.
54. The composition for use according to any one of claims 41-53, wherein the method further comprises testing or confirming expression of TGF β 1, LTBP1 or LTBP3 in a biological sample collected from the subject.
55. A method for making a composition of any one of claims 37-40, comprising an antibody, or antigen-binding fragment thereof, that specifically binds a human LTBP1-proTGF β complex and/or a human LTBP3-proTGF β complex, the method comprising steps of:
- i) selecting an antibody or an antigen-binding fragment thereof that dissociates from human LTBP1-proTGF β complex and/or a human LTBP3-proTGF β complex with $t_{1/2}$ of at least 45 minutes, and,
 - ii) formulating the antibody or fragment into a pharmaceutical composition, thereby making the composition comprising the antibody or fragment.
56. The method of 55, further comprising the step of: selecting an antibody or antigen-binding fragment with IC₅₀ of < 5 nM as measured by cell-based assays, optionally < 2 nM.
57. The method of 55 or 56, further comprising the step of: confirming efficacy in an in vivo preclinical model, wherein optionally the preclinical model is a liver fibrosis model, kidney fibrosis model, or cardiac fibrosis model.
58. The method of any one of claims 55-57, wherein the method further comprises a step of: selecting antibodies or fragments that are cross-reactive to human and rodent antigens.
59. The method of any one of claims 55-58, wherein the method further comprises a step of: subjecting the antibody or fragment that is present in the first pool of antibodies and the second pool of antibodies to affinity maturation and/or optimization, so as to provide an affinity matured and/or optimized antibody or fragment.