(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau

(43) International Publication Date
5 February 2004 (05.02.2004)

PCT

(10) International Publication Number
WO 2004/011900 A2

(51) International Patent Classification[7]: G01N

(21) International Application Number:
PCT/US2003/022611

(22) International Filing Date: 17 July 2003 (17.07.2003)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/398,724 25 July 2002 (25.07.2002) US

(71) Applicant: **ACLARA BIOSCIENCES, INC.** [US/US]; 1288 Pear Avenue, Mountain View, CA 94043 (US).

(72) Inventors: **CHAN-HUI, Po-Ying**; 13056 Parkhurst Drive, Oakland, CA 94619 (US). **SHI, Yining**; 7170 Bark Lane #2, San Jose, CA 95129 (US). **PIDAPARTHI, Sailaja**; 8055 Park Vill Circle, Cupertino, CA 95014 (US). **DUA, Rajiv**; 173 Whelan Way, Manteca, CA 95336 (US). **SINGH, Sharat**; 3420 Royal Meadow Lane, San Jose, CA 95135 (US).


(74) Agent: **MACEVICZ, Stephen, C.**; Aclara Biosciences, Inc., 1288 Pear Avenue, Mountain View, CA 94043 (US).

(81) Designated States *(national)*: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States *(regional)*: ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE,

[Continued on next page]

(54) Title: DETECTING RECEPTOR OLIGOMERIZATION



(57) Abstract: Methods are provided for detecting formation of oligomeric complexes of molecules on the surface of cell membranes. These methods employ pairs of tagged probes and cleaving probes, each of which binds specificly to a cell surface molecule. The tagged probe includes a molecular tag that is linked to a first binding compound through a cleavable linkage, and the cleaving probe includes a second binding agent and a cleavage-inducing moiety that can cleave the linkage when within a defined proximity thereto. Binding of the two probes to cell surface molecules that have formed an oligomeric complex results in release of the molecular tag from the binding compound, providing a measure of formation of the complex.

WO 2004/011900 A2

ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— *without international search report and to be republished upon receipt of that report*

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.