

PROLIFERATION SIGNATURE AND PROGNOSIS

FOR GASTROINTESTINAL CANCER

FIELD OF THE INVENTION

- 5 This invention relates to methods and compositions for determining the prognosis of cancer, particularly gastrointestinal cancer, in a patient. Specifically, this invention relates to the use of genetic markers for determining the prognosis of cancer, such as gastrointestinal cancer, based on cell proliferation signatures.

10 BACKGROUND OF THE INVENTION

Cellular proliferation is the most fundamental process in living organisms, and as such is precisely regulated by the expression level of proliferation-associated genes (1). Loss of proliferation control is a hallmark of cancer, and it is thus not surprising that growth-regulating genes are abnormally expressed in tumours relative to the neighbouring normal
15 tissue (2). Proliferative changes may accompany other changes in cellular properties, such as invasion and ability to metastasize, and therefore could affect patient outcome. This association has attracted substantial interest and many studies have been devoted to the exploration of tumour cell proliferation as a potential indicator of outcome.

20 Cell proliferation is usually assessed by flow cytometry or, more commonly, in tissues, by immunohistochemical evaluation of proliferation markers (3). The most widely used proliferation marker is Ki-67, a protein expressed in all cell cycle phases except for the resting phase G_0 (4). Using Ki-67, a clear association between the proportion of cycling cells and clinical outcome has been established in malignancies such as breast cancer,
25 lung cancer, soft tissue tumours, and astrocytoma (5). In breast cancer, this association has also been confirmed by microarray analysis, leading to a proliferative gene expression profile that has been employed for identifying patients at increased risk of recurrence (6).

However, in colorectal cancer (CRC), the proliferation index (PI) has produced conflicting
30 results as a prognostic factor and therefore cannot be applied in a clinical context (see below). Studies vary with respect to patient selection, sampling methods, cut-off point levels, antibody choices, staining techniques and the way data have been collected and interpreted. The methodological differences and heterogeneity of these studies may partly explain the contradictory results (7),(8). The use of Ki-67 as a proliferation marker also
35 has limitations. The Ki-67 PI estimates the fraction of actively cycling cells, but gives no indication of cell cycle length (3),(9). Thus, tumours with a similar PI may grow at dissimilar rates due to different cycling speeds. In addition, while Ki-67 mRNA is not

produced in resting cells, protein may still be detectable in a proportion of colorectal tumours leading to an overestimated proliferation rate (10).

5 Since the assessment of a prognosis using a single proliferation marker does not appear to be reliable in CRC (see below), there is a need for further tools to predict the prognosis of gastrointestinal cancer. This invention provides further methods and compositions based on prognostic cancer markers, specifically gastrointestinal cancer prognostic markers, to aid in the prognosis and treatment of cancer.

10 SUMMARY OF THE INVENTION

In certain aspects of the invention, microarray analysis is used to identify genes that provide a proliferation signature for cancer cells. These genes, and the proteins encoded by those genes, are herein termed gastrointestinal cancer proliferation markers (GCPMs). In one aspect of the invention, the cancer for prognosis is gastrointestinal cancer,
15 particularly gastric or colorectal cancer.

In particular aspects, the invention includes a method for determining the prognosis of a cancer by identifying the expression levels of at least one GCPM in a sample. Selected GCPMs encode proteins that associated with cell proliferation, e.g., cell cycle
20 components. These GCPMs have the added utility in methods for determining the best treatment regime for a particular cancer based on the prognosis. In particular aspects, GCPM levels are higher in non-recurring tumour tissue as compared to recurring tumour tissue. These markers can be used either alone or in combination with each other, or other known cancer markers.

25

In an additional aspect, this invention includes a method for determining the prognosis of a cancer, comprising: (a) providing a sample of the cancer; (b) detecting the expression level of at least one GCPM family member in the sample; and (c) determining the prognosis of the cancer.

30

In another aspect, the invention includes a step of detecting the expression level of at least one GCPM RNA, for example, at least one mRNA. In a further aspect, the invention includes a step of detecting the expression level of at least one GCPM protein. In yet a further aspect, the invention includes a step of detecting the level of at least one GCPM
35 peptide. In yet another aspect, the invention includes detecting the expression level of at least one GCPM family member in the sample. In an additional aspect, the GCPM is a

gene associated with cell proliferation, such as a cell cycle component. In other aspects, the at least one GCPM is selected from Table A, Table B, Table C or Table D, herein.

In a still further aspect, the invention includes a method for detecting the expression level
5 of at least one GCPM set forth in Table A, Table B, Table C or Table D, herein. In an even further aspect, the invention includes a method for detecting the expression level of at least one of CDC2, MCM6, RPA3, MCM7, PCNA, G22P1, KPNA2, ANLN, APG7L, TOPK, GMNN, RRM1, CDC45L, MAD2L1, RAN, DUT, RRM2, CDK7, MLH3, SMC4L1, CSPG6, POLD2, POLE2, BCCIP, Pfs2, TREX1, BUB3, FEN1, DRF1, PREI3, CCNE1, RPA1,
10 POLE3, RFC4, MCM3, CHEK1, CCND1, and CDC37. In yet a further aspect, the invention comprises detecting the expression level of at least one of CDC2, RFC4, PCNA, CCNE1, CCND1, CDK7, MCM genes, FEN1, MAD2L1, MYBL2, RRM2, and BUB3.

In additional aspects, the expression levels of at least two, or at least 5, or at least 10, at
15 least 15, at least 20, at least 25, at least 30, at least 35, at least 40, at least 45, at least 50, or at least 75 of the proliferation markers or their expression products are determined, for example, as selected from Table A, Table, B, Table C or Table D; as selected from CDC2, MCM6, RPA3, MCM7, PCNA, G22P1, KPNA2, ANLN, APG7L, TOPK, GMNN, RRM1, CDC45L, MAD2L1, RAN, DUT, RRM2, CDK7, MLH3, SMC4L1, CSPG6, POLD2,
20 POLE2, BCCIP, Pfs2, TREX1, BUB3, FEN1, DRF1, PREI3, CCNE1, RPA1, POLE3, RFC4, MCM3, CHEK1, CCND1, and CDC37; or as selected from CDC2, RFC4, PCNA, CCNE1, CCND1, CDK7, MCM genes (e.g., one or more of MCM3, MCM6, and MCM7), FEN1, MAD2L1, MYBL2, RRM2, and BUB3.

25 In other aspects, the expression levels of all proliferation markers or their expression products are determined, for example, as listed in Table A, Table, B, Table C or Table D; as listed for the group CDC2, MCM6, RPA3, MCM7, PCNA, G22P1, KPNA2, ANLN, APG7L, TOPK, GMNN, RRM1, CDC45L, MAD2L1, RAN, DUT, RRM2, CDK7, MLH3, SMC4L1, CSPG6, POLD2, POLE2, BCCIP, Pfs2, TREX1, BUB3, FEN1, DRF1, PREI3,
30 CCNE1, RPA1, POLE3, RFC4, MCM3, CHEK1, CCND1, and CDC37; or as listed for the group CDC2, RFC4, PCNA, CCNE1, CCND1, CDK7, MCM genes (e.g., one or more of MCM3, MCM6, and MCM7), FEN1, MAD2L1, MYBL2, RRM2, and BUB3.

In yet a further aspect, the invention includes a method of determining a treatment regime
35 for a cancer comprising: (a) providing a sample of the cancer; (b) detecting the expression level of at least one GCPM family member in the sample; (c) determining the

prognosis of the cancer based on the expression level of at least one GCPM family member; and (d) determining the treatment regime according to the prognosis.

5 In yet another aspect, the invention includes a device for detecting at least one GCPM, comprising: (a) a substrate having at least one GCPM capture reagent thereon; and (b) a detector capable of detecting the at least one captured GCPM, the capture reagent, or a complex thereof.

10 An additional aspect of the invention includes a kit for detecting cancer, comprising: (a) GCPM capture reagent; (b) a detector capable of detecting the captured GCPM, the capture reagent, or a complex thereof; and, optionally, (c) instructions for use. In certain aspects, the kit also includes a substrate for the GCPM as captured.

15 Yet a further aspect of the invention includes a method for detecting at least one GCPM using quantitative PCR, comprising: (a) a forward primer specific for the at least one GCPM; (b) a reverse primer specific for the at least one GCPM; (c) PCR reagents; and, optionally, at least one of: (d) a reaction vial; and (e) instructions for use.

20 Additional aspects of this invention include a kit for detecting the presence of at least one GCPM protein or peptide, comprising: (a) an antibody or antibody fragment specific for the at least one GCPM protein or peptide; and, optionally, at least one of: (b) a label for the antibody or antibody fragment; and (c) instructions for use. In certain aspects, the kit also includes a substrate having a capture agent for the at least one GCPM protein or peptide.

25 In specific aspects, this invention includes a method for determining the prognosis of gastrointestinal cancer, especially colorectal or gastric cancer, comprising the steps of: (a) providing a sample, e.g., tumour sample, from a patient suspected of having gastrointestinal cancer; (b) measuring the presence of a GCPM protein using an ELISA method.

30

In additional aspects of this invention, one or more GCPMs of the invention are selected from the group outlined in Table A, Table B, Table C or Table D, herein. Other aspects and embodiments of the invention are described herein below.

BRIEF DESCRIPTION OF THE DRAWINGS

This invention is described with reference to specific embodiments thereof and with reference to the figures.

5 FIG. 1: An overview of the approach used to derive and apply the gene proliferation signature (GPS) disclosed herein.

FIG. 2A: K-means clustering of 73 Cohort A tumours into two groups according to the expression level of the gene proliferation signature. FIG. 2B: Bar graph of Ki-67 PI (%);
10 vertical line represents the mean Ki-67 PI across all samples. Tumours with a proliferation index about and below the mean are shown in red and green, respectively. The results show that over-expression of the proliferation signature is not always associated with a higher Ki-67 PI.

FIG. 3: Kaplan-Meier survival curves according to the expression level of GPS (gene
15 proliferation signal) and Ki-67 PI. Both overall (OS) and recurrence-free survival (RFS) are significantly shorter in patients with low GPS expression in colorectal cancer Cohort A (a, b) and colorectal cancer Cohort B (c, d). No difference was observed in the survival rates of Cohort A patients according to Ki-67 PI (e, f). *P* values from Log rank test are indicated.

20 FIG. 4: Kaplan-Meier survival curves according to the expression level of GPS (gene proliferation signal) in gastric cancer patients. Overall survival is significantly shorter in patients with low GPS expression in this cohort of 38 gastric cancer patients of mixed stage. *P* values from Log rank test are indicated.

25 FIG. 5: A box-and-whisker plot showing differential expression between cycling cells in the exponential phase (EP) and growth-inhibited cells in the stationary phase (SP) of 11 QRT-PCR-validated genes. The box range includes the 25 to the 75 percentiles of the data. The horizontal line in the box represents the median value. The "whiskers" are the largest and smallest values (excluding outliers). Any points more than 3/2 times of the
30 interquartile range from the end of a box will be outliers and presented as a dot. The Y axis represents the log 2 fold change of the ratio between cell line RNA and reference RNA. Analysis was performed using SPSS software.

DETAILED DESCRIPTION OF THE INVENTION

35 Because a single proliferation marker is insufficient for obtaining reliable CRC prognosis, the simultaneous analysis of several growth-related genes by microarray was employed to provide a more quantitative and objective method to determine the proliferation state of a

gastrointestinal tumour. Table 1 (below) illustrates the previously published and conflicting results shown for use of the proliferation index (PI) as a prognostic factor for colorectal cancer.

5 **Table 1: Summary of studies on the association of proliferation indices with the CRC patients' survival**

Study	Number of patients	Dukes stage	Marker	Association with survival
Evans et al, 2006 ¹¹	40	A-C	Ki-67	No association was found between proliferation index and survival
Rosati et al, 2004 ¹²	103	B-C	Ki-67	
Ishida et al, 2004 ¹³	51	C	Ki-67	
Buglioni et al, 1999 ¹⁴	171	A-D	Ki-67	
Guerra et al, 1998 ¹⁵	108	A-C	PCNA	
Kyzer and Gordon, 1997 ¹⁶	30	B-D	Ki-67	
Jansson and Sun, 1997 ¹⁷	255	A-D	Ki-67	
Baretton et al, 1996 ¹⁸	95	A-B	Ki-67	
Sun et al, 1996 ¹⁹	293	A-C	PCNA	
Kubota et al, 1992 ²⁰	100	A-D	Ki-67	High proliferation index was associated with shorter survival
Valera et al, 2005 ²¹	106	A-D	Ki-67	
Dziegiel et al, 2003 ²²	81	NI	Ki-67	
Scopa et al, 2003 ²³	117	A-D	Ki-67	
Bhatavdekar et al, 2001 ²⁴	98	B-C	Ki-67	
Chen et al, 1997 ²⁵	70	B-C	Ki-67	
Choi et al, 1997 ²⁶	86	B-D	PCNA	Low proliferation index was associated with shorter survival
Hilska et al, 2005 ²⁷	363	A-D	Ki-67	
Salminen et al, 2005 ²⁸	146	A-D	Ki-67	
Garrity et al, 2004 ²⁹	366	B-C	Ki-67	
Allegra et al, 2003 ³⁰	706	B-C	Ki-67	
Palmqvist et al, 1999 ³¹	56	B	Ki-67	
Paradiso et al, 1996 ³²	71	NI	PCNA	
Neoptolemos et al, 1995 ³³	79	A-C	PCNA	
NI: No information available				

In contrast, the present disclosure has succeeded in (i) defining a CRC-specific gene proliferation signature (GPS) using a cell line model; and (ii) determining the prognostic significance of the GPS in the prediction of patient outcome and its association with clinico-pathologic variables in two independent cohorts of CRC patients.

Definitions

Before describing embodiments of the invention in detail, it will be useful to provide some definitions of terms used herein.

As used herein "antibodies" and like terms refer to immunoglobulin molecules and immunologically active portions of immunoglobulin (Ig) molecules, i.e., molecules that contain an antigen binding site that specifically binds (immunoreacts with) an antigen. These include, but are not limited to, polyclonal, monoclonal, chimeric, single chain, Fc, Fab, Fab', and Fab₂ fragments, and a Fab expression library. Antibody molecules relate to any of the classes IgG, IgM, IgA, IgE, and IgD, which differ from one another by the nature

of heavy chain present in the molecule. These include subclasses as well, such as IgG1, IgG2, and others. The light chain may be a kappa chain or a lambda chain. Reference herein to antibodies includes a reference to all classes, subclasses, and types. Also included are chimeric antibodies, for example, monoclonal antibodies or fragments thereof
5 that are specific to more than one source, e.g., a mouse or human sequence. Further included are camelid antibodies, shark antibodies or nanobodies.

The term "marker" refers to a molecule that is associated quantitatively or qualitatively with the presence of a biological phenomenon. Examples of "markers" include a
10 polynucleotide, such as a gene or gene fragment, RNA or RNA fragment; or a polypeptide such as a peptide, oligopeptide, protein, or protein fragment; or any related metabolites, by products, or any other identifying molecules, such as antibodies or antibody fragments, whether related directly or indirectly to a mechanism underlying the phenomenon. The markers of the invention include the nucleotide sequences (e.g., GenBank sequences) as
15 disclosed herein, in particular, the full-length sequences, any coding sequences, any fragments, or any complements thereof.

The terms "GCPM" or "gastrointestinal cancer proliferation marker" or "GCPM family member" refer to a marker with increased expression that is associated with a positive
20 prognosis, e.g., a lower likelihood of recurrence cancer, as described herein, but can exclude molecules that are known in the prior art to be associated with prognosis of gastrointestinal cancer. It is to be understood that the term GCPM does not require that the marker be specific only for gastrointestinal tumours. Rather, expression of GCPM can be altered in other types of tumours, including malignant tumours.

25 Non-limiting examples of GCPMs are included in Table A, Table B, Table C or Table D, herein below, and include, but are not limited to, the specific group CDC2, MCM6, RPA3, MCM7, PCNA, G22P1, KPNA2, ANLN, APG7L, TOPK, GMNN, RRM1, CDC45L, MAD2L1, RAN, DUT, RRM2, CDK7, MLH3, SMC4L1, CSPG6, POLD2, POLE2, BCCIP,
30 Pfs2, TREX1, BUB3, FEN1, DRF1, PREI3, CCNE1, RPA1, POLE3, RFC4, MCM3, CHEK1, CCND1, and CDC37; and the specific group CDC2, RFC4, PCNA, CCNE1, CCND1, CDK7, MCM genes (e.g., one or more of MCM3, MCM6, and MCM7), FEN1, MAD2L1, MYBL2, RRM2, and BUB3.

35 The terms "cancer" and "cancerous" refer to or describe the physiological condition in mammals that is typically characterized by abnormal or unregulated cell growth. Cancer and cancer pathology can be associated, for example, with metastasis, interference with

the normal functioning of neighbouring cells, release of cytokines or other secretory products at abnormal levels, suppression or aggravation of inflammatory or immunological response, neoplasia, premalignancy, malignancy, invasion of surrounding or distant tissues or organs, such as lymph nodes, etc. Specifically included are gastrointestinal
5 cancers, such as esophageal, stomach, small bowel, large bowel, anal, and rectal cancers, particularly included are gastric and colorectal cancers.

The term "colorectal cancer" includes cancer of the colon, rectum, and/or anus, and especially, adenocarcinomas, and may also include carcinomas (e.g., squamous
10 cloacogenic carcinomas), melanomas, lymphomas, and sarcomas. Epidermoid (nonkeratinizing squamous cell or basaloid) carcinomas are also included. The cancer may be associated with particular types of polyps or other lesions, for example, tubular adenomas, tubulovillous adenomas (e.g., villoglandular polyps), villous (e.g., papillary) adenomas (with or without adenocarcinoma), hyperplastic polyps, hamartomas, juvenile
15 polyps, polypoid carcinomas, pseudopolyps, lipomas, or leiomyomas. The cancer may be associated with familial polyposis and related conditions such as Gardner's syndrome or Peutz-Jeghers syndrome. The cancer may be associated, for example, with chronic fistulas, irradiated anal skin, leukoplakia, lymphogranuloma venereum, Bowen's disease (intraepithelial carcinoma), condyloma acuminatum, or human papillomavirus. In other
20 aspects, the cancer may be associated with basal cell carcinoma, extramammary Paget's disease, cloacogenic carcinoma, or malignant melanoma.

The terms "differentially expressed gene," "differential gene expression," and like phrases, refer to a gene whose expression is activated to a higher or lower level in a subject (e.g.,
25 test sample), specifically cancer, such as gastrointestinal cancer, relative to its expression in a control subject (e.g., control sample). The terms also include genes whose expression is activated to a higher or lower level at different stages of the same disease; in recurrent or non-recurrent disease; or in cells with higher or lower levels of proliferation. A differentially expressed gene may be either activated or inhibited at the polynucleotide
30 level or polypeptide level, or may be subject to alternative splicing to result in a different polypeptide product. Such differences may be evidenced by a change in mRNA levels, surface expression, secretion or other partitioning of a polypeptide, for example.

Differential gene expression may include a comparison of expression between two or
35 more genes or their gene products; or a comparison of the ratios of the expression between two or more genes or their gene products; or a comparison of two differently processed products of the same gene, which differ between normal subjects and diseased

subjects; or between various stages of the same disease; or between recurring and non-recurring disease; or between cells with higher and lower levels of proliferation; or between normal tissue and diseased tissue, specifically cancer, or gastrointestinal cancer. Differential expression includes both quantitative, as well as qualitative, differences in the temporal or cellular expression pattern in a gene or its expression products among, for example, normal and diseased cells, or among cells which have undergone different disease events or disease stages, or cells with different levels of proliferation.

The term "expression" includes production of polynucleotides and polypeptides, in particular, the production of RNA (e.g., mRNA) from a gene or portion of a gene, and includes the production of a protein encoded by an RNA or gene or portion of a gene, and the appearance of a detectable material associated with expression. For example, the formation of a complex, for example, from a protein-protein interaction, protein-nucleotide interaction, or the like, is included within the scope of the term "expression". Another example is the binding of a binding ligand, such as a hybridization probe or antibody, to a gene or other oligonucleotide, a protein or a protein fragment and the visualization of the binding ligand. Thus, increased intensity of a spot on a microarray, on a hybridization blot such as a Northern blot, or on an immunoblot such as a Western blot, or on a bead array, or by PCR analysis, is included within the term "expression" of the underlying biological molecule.

The term "gastric cancer" includes cancer of the stomach and surrounding tissue, especially adenocarcinomas, and may also include lymphomas and leiomyosarcomas. The cancer may be associated with gastric ulcers or gastric polyps, and may be classified as protruding, penetrating, spreading, or any combination of these categories, or, alternatively, classified as superficial (elevated, flat, or depressed) or excavated.

The term "long-term survival" is used herein to refer to survival for at least 5 years, more preferably for at least 8 years, most preferably for at least 10 years following surgery or other treatment

The term "microarray" refers to an ordered arrangement of capture agents, preferably polynucleotides (e.g., probes) or polypeptides on a substrate. See, e.g., *Microarray Analysis*, M. Schena, John Wiley & Sons, 2002; *Microarray Biochip Technology*, M. Schena, ed., Eaton Publishing, 2000; *Guide to Analysis of DNA Microarray Data*, S. Knudsen, John Wiley & Sons, 2004; and *Protein Microarray Technology*, D. Kambhampati, ed., John Wiley & Sons, 2004.

The term "oligonucleotide" refers to a polynucleotide, typically a probe or primer, including, without limitation, single-stranded deoxyribonucleotides, single- or double-stranded ribonucleotides, RNA:DNA hybrids, and double-stranded DNAs. Oligonucleotides, such as
5 single-stranded DNA probe oligonucleotides, are often synthesized by chemical methods, for example using automated oligonucleotide synthesizers that are commercially available, or by a variety of other methods, including *in vitro* expression systems, recombinant techniques, and expression in cells and organisms.

10 The term "polynucleotide," when used in the singular or plural, generally refers to any polyribonucleotide or polydeoxyribonucleotide, which may be unmodified RNA or DNA or modified RNA or DNA. This includes, without limitation, single- and double-stranded DNA, DNA including single- and double- stranded regions, single- and double-stranded RNA, and RNA including single- and double-stranded regions, hybrid molecules comprising
15 DNA and RNA that may be single-stranded or, more typically, double-stranded or include single- and double-stranded regions. Also included are triple-stranded regions comprising RNA or DNA or both RNA and DNA. Specifically included are mRNAs, cDNAs, and genomic DNAs. The term includes DNAs and RNAs that contain one or more modified bases, such as tritiated bases, or unusual bases, such as inosine. The polynucleotides of
20 the invention can encompass coding or non-coding sequences, or sense or antisense sequences.

"Polypeptide," as used herein, refers to an oligopeptide, peptide, or protein sequence, or fragment thereof, and to naturally occurring, recombinant, synthetic, or semi-synthetic
25 molecules. Where "polypeptide" is recited herein to refer to an amino acid sequence of a naturally occurring protein molecule, "polypeptide" and like terms, are not meant to limit the amino acid sequence to the complete, native amino acid sequence for the full-length molecule. It will be understood that each reference to a "polypeptide" or like term, herein, will include the full-length sequence, as well as any fragments, derivatives, or variants
30 thereof.

The term "prognosis" refers to a prediction of medical outcome (e.g., likelihood of long-term survival); a negative prognosis, or bad outcome, includes a prediction of relapse, disease progression (e.g., tumour growth or metastasis, or drug resistance), or mortality; a
35 positive prognosis, or good outcome, includes a prediction of disease remission, (e.g., disease-free status), amelioration (e.g., tumour regression), or stabilization.

The terms "prognostic signature," "signature," and the like refer to a set of two or more markers, for example GCPMs, that when analysed together as a set allow for the determination of or prediction of an event, for example the prognostic outcome of colorectal cancer. The use of a signature comprising two or more markers reduces the effect of individual variation and allows for a more robust prediction. Non-limiting examples of GCPMs are included in Table A, Table B, Table C or Table D, herein below, and include, but are not limited to, the specific group CDC2, MCM6, RPA3, MCM7, PCNA, G22P1, KPNA2, ANLN, APG7L, TOPK, GMNN, RRM1, CDC45L, MAD2L1, RAN, DUT, RRM2, CDK7, MLH3, SMC4L1, CSPG6, POLD2, POLE2, BCCIP, Pfs2, TREX1, BUB3, FEN1, DRF1, PREI3, CCNE1, RPA1, POLE3, RFC4, MCM3, CHEK1, CCND1, and CDC37; and the specific group CDC2, RFC4, PCNA, CCNE1, CCND1, CDK7, MCM genes (e.g., one or more of MCM3, MCM6, and MCM7), FEN1, MAD2L1, MYBL2, RRM2, and BUB3.

In the context of the present invention, reference to "at least one," "at least two," "at least five," etc., of the markers listed in any particular set (e.g., any signature) means any one or any and all combinations of the markers listed.

The term "prediction method" is defined to cover the broader genus of methods from the fields of statistics, machine learning, artificial intelligence, and data mining, which can be used to specify a prediction model. These are discussed further in the Detailed Description section.

The term "prediction model" refers to the specific mathematical model obtained by applying a prediction method to a collection of data. In the examples detailed herein, such data sets consist of measurements of gene activity in tissue samples taken from recurrent and non-recurrent colorectal cancer patients, for which the class (recurrent or non-recurrent) of each sample is known. Such models can be used to (1) classify a sample of unknown recurrence status as being one of recurrent or non-recurrent, or (2) make a probabilistic prediction (i.e., produce either a proportion or percentage to be interpreted as a probability) which represents the likelihood that the unknown sample is recurrent, based on the measurement of mRNA expression levels or expression products, of a specified collection of genes, in the unknown sample. The exact details of how these gene-specific measurements are combined to produce classifications and probabilistic predictions are dependent on the specific mechanisms of the prediction method used to construct the model.

The term "proliferation" refers to the processes leading to increased cell size or cell number, and can include one or more of: tumour or cell growth, angiogenesis, innervation, and metastasis.

- 5 The term "qPCR" or "QPCR" refers to quantitative polymerase chain reaction as described, for example, in PCR Technique: Quantitative PCR, J.W. Larrick, ed., Eaton Publishing, 1997, and A-Z of Quantitative PCR, S. Bustin, ed., IUL Press, 2004.

- 10 The term "tumour" refers to all neoplastic cell growth and proliferation, whether malignant or benign, and all pre-cancerous and cancerous cells and tissues.

Sensitivity", "specificity" (or "selectivity"), and "classification rate", when applied to the describing the effectiveness of prediction models mean the following:

- 15 "Sensitivity" means the proportion of truly positive samples that are also *predicted* (by the model) to be positive. In a test for cancer recurrence, that would be the proportion of recurrent tumours predicted by the model to be recurrent. "Specificity" or "selectivity" means the proportion of truly negative samples that are also *predicted* (by the model) to be negative. In a test for CRC recurrence, this equates to the proportion of non-recurrent samples that are predicted to by non-recurrent by the model. "Classification Rate" is the
20 proportion of all samples that are correctly classified by the prediction model (be that as positive or negative).

- "Stringent conditions" or "high stringency conditions", as defined herein, typically: (1) employ low ionic strength and high temperature for washing, for example 0.015 M sodium
25 chloride/0.0015 M sodium citrate/0.1% sodium dodecyl sulfate at 50°C; (2) employ a denaturing agent during hybridization, such as formamide, for example, 50% (v/v) formamide with 0.1% bovine serum albumin/0.1% Ficoll/0.1% polyvinylpyrrolidone/50 mM sodium phosphate buffer at pH 6.5 with 750 mM sodium chloride, 75 mM sodium citrate at 42°C; or (3) employ 50% formamide, 5X SSC (0.75 M NaCl, 0.075 M sodium citrate), 50
30 mM sodium phosphate (pH 6.8), 0.1% sodium pyrophosphate, 5X, Denhardt's solution, sonicated salmon sperm DNA (50 µg/ml), 0.1% SDS, and 10% dextran sulfate at 42°C, with washes at 42°C in 0.2X SSC (sodium chloride/sodium citrate) and 50% formamide at 55°C, followed by a high-stringency wash comprising 0.1X SSC containing EDTA at 55°C.

- 35 "Moderately stringent conditions" may be identified as described by Sambrook et al., Molecular Cloning: A Laboratory Manual, New York: Cold Spring Harbor Press, 1989, and include the use of washing solution and hybridization conditions (e. g., temperature, ionic

strength, and % SDS) less stringent than those described above. An example of moderately stringent conditions is overnight incubation at 37°C in a solution comprising: 20% formamide, 5X SSC (150 mM NaCl, 15 mM trisodium citrate), 50 mM sodium phosphate (pH 7.6), 5X Denhardt's solution, 10% dextran sulfate, and 20 mg/ml denatured
5 sheared salmon sperm DNA, followed by washing the filters in 1X SSC at about 37-50°C. The skilled artisan will recognize how to adjust the temperature, ionic strength, etc. as necessary to accommodate factors such as probe length and the like.

The practice of the present invention will employ, unless otherwise indicated, conventional
10 techniques of molecular biology (including recombinant techniques), microbiology, cell biology, and biochemistry, which are within the skill of the art. Such techniques are explained fully in the literature, such as, Molecular Cloning: A Laboratory Manual, 2nd edition, Sambrook et al., 1989; Oligonucleotide Synthesis, MJ Gait, ed., 1984; Animal Cell Culture, R.I. Freshney, ed., 1987; Methods in Enzymology, Academic Press, Inc.;
15 Handbook of Experimental Immunology, 4th edition, D.M. Weir & C.C. Blackwell, eds., Blackwell Science Inc., 1987; Gene Transfer Vectors for Mammalian Cells, J.M. Miller & M.P. Calos, eds., 1987; Current Protocols in Molecular Biology, F.M. Ausubel et al., eds., 1987; and PCR: The Polymerase Chain Reaction, Mullis et al., eds., 1994.

20 **Description of Embodiments of the Invention**

Cell proliferation is an indicator of outcome in some malignancies. In colorectal cancer, however, discordant results have been reported. As these results are based on a single proliferation marker, the present invention discloses the use of microarrays to overcome this limitation, to reach a firmer conclusion, and to determine the prognostic role of cell
25 proliferation in colorectal cancer. The microarray-based proliferation studies shown herein indicate that reduced rate of the proliferation signature in colorectal cancer is associated with poor outcome. The invention can therefore be used to identify patients at high risk of early death from cancer.

30 The present invention provides for markers for the determination of disease prognosis, for example, the likelihood of recurrence of tumours, including gastrointestinal tumours. Using the methods of the invention, it has been found that numerous markers are associated with the progression of gastrointestinal cancer, and can be used to determine the prognosis of cancer. Microarray analysis of samples taken from patients with various
35 stages of colorectal tumours has led to the surprising discovery that specific patterns of marker expression are associated with prognosis of the cancer.

An increase in certain GCPMs, for example, markers associated with cell proliferation, is indicative of positive prognosis. This can include decreased likelihood of cancer recurrence after standard treatment, especially for gastrointestinal cancer, such as gastric or colorectal cancer. Conversely, a decrease in these markers is indicative of a negative prognosis. This can include disease progression or the increased likelihood of cancer recurrence, especially for gastrointestinal cancer, such as gastric or colorectal cancer. A decrease in expression can be determined, for example, by comparison of a test sample (e.g., tumour sample) to samples associated with a positive prognosis. An increase in expression can be determined, for example, by comparison of a test sample (e.g., tumour samples) to samples associated with a negative prognosis.

For example, to obtain a prognosis, a patient's sample (e.g., tumour sample) can be compared to samples with known patient outcome. If the patient's sample shows increased expression of GCPMs that is comparable to samples with good outcome, and/or higher than samples with poor outcome, then a positive prognosis is implicated. If the patient's sample shows decreased expression of GCPMs that is comparable to samples with poor outcome, and/or lower than samples with good outcome, then a negative prognosis is implicated. Alternatively, a patient's sample can be compared to samples of actively proliferating/non-proliferating tumour cells. If the patient's sample shows increased expression of GCPMs that is comparable to actively proliferating cells, and/or higher than non-proliferating cells, then a positive prognosis is implicated. If the patient's sample shows decreased expression of GCPMs that is comparable to non-proliferating cells, and/or lower than actively proliferating cells, then a negative prognosis is implicated.

The invention provides for a set of genes, identified from cancer patients with various stages of tumours, outlined in Table C that are shown to be prognostic for colorectal cancer. These genes are all associated with cell proliferation and establish a relationship between cell proliferation genes and their utility in cancers prognosis. It has also been found that the genes in the prognostic signature listed in Table C are also correlated with additional cell proliferation genes. Based on these finding, the invention also provides for a set of cell cycle genes, shown in Table D, that are differentially expressed between high and low proliferation groups, for use as prognostic markers. Further, based on the surprising finding of the correlation between prognosis and cell proliferation-related genes, the invention also provides for a set of proliferation-related genes differentially expressed between cell lines in high and low proliferative states (Table A) and known proliferative-

related genes (Table B). The genes outlined in Table A, Table B, Table C and Table D provide for a set of gastrointestinal cancer prognostic markers (GCPMs).

5 As one approach, the expression of a panel of markers (e.g., GCPMs) can be analysed by techniques including Linear Discriminant Analysis (LDA) to work out a prognostic score. The marker panel selected and prognostic score calculation can be derived through extensive laboratory testing and multiple independent clinical development studies.

10 The disclosed GCPMs therefore provide a useful tool for determining the prognosis of cancer, and establishing a treatment regime specific for that tumour. In particular, a positive prognosis can be used by a patient to decide to pursue standard or less invasive treatment options. A negative prognosis can be used by a patient to decide to terminate treatment or to pursue highly aggressive or experimental treatments. In addition, a patient can chose treatments based on their impact on cell proliferation or the expression of cell proliferation markers (e.g., GCPMs). In accordance with the present invention, treatments that specifically target cells with high proliferation or specifically decrease expression of cell proliferation markers (e.g., GCPMs) would not be preferred for patients with gastrointestinal cancer, such as colorectal cancer or gastric cancer.

20 Levels of GCPMs can be detected in tumour tissue, tissue proximal to the tumour, lymph node samples, blood samples, serum samples, urine samples, or faecal samples, using any suitable technique, and can include, but is not limited to, oligonucleotide probes, quantitative PCR, or antibodies raised against the markers. The expression level of one GCPM in the sample will be indicative of the likelihood of recurrence in that subject. However, it will be appreciated that by analyzing the presence and amounts of expression of a plurality of GCPMs, and constructing a proliferation signature, the sensitivity and accuracy of prognosis will be increased. Therefore, multiple markers according to the present invention can be used to determine the prognosis of a cancer.

30 The present invention relates to a set of markers, in particular, GCPMs, the expression of which has prognostic value, specifically with respect to cancer-free survival. In specific aspects, the cancer is gastrointestinal cancer, particularly, gastric or colorectal cancer, and, in further aspects, the colorectal cancer is an adenocarcinoma.

35 In one aspect, the invention relates to a method of predicting the likelihood of long-term survival of a cancer patient without the recurrence of cancer, comprising determining the expression level of one or more proliferation markers or their expression products in a

sample obtained from the patient, normalized against the expression level of all RNA transcripts or their products in the sample, or of a reference set of RNA transcripts or their expression products, wherein the proliferation marker is the transcript of one or more markers listed in Table A, Table B, Table C or Table D, herein. In particular aspects, a decrease in expression levels of one or more GCPM indicates a decreased likelihood of long-term survival without cancer recurrence, while an increase in expression levels of one or more GCPM indicates an increased likelihood of long-term survival without cancer recurrence.

10 In a further aspect, the expression levels one or more, for example at least two, or at least 3, or at least 4, or at least 5, or at least 10, at least 15, at least 20, at least 25, at least 30, at least 35, at least 40, at least 45, at least 50, or at least 75 of the proliferation markers or their expression products are determined, e.g., as selected from Table A, Table B, Table C or Table D; as selected from CDC2, MCM6, RPA3, MCM7, PCNA, G22P1, KPNA2, ANLN, APG7L, TOPK, GMNN, RRM1, CDC45L, MAD2L1, RAN, DUT, RRM2, CDK7, 15 MLH3, SMC4L1, CSPG6, POLD2, POLE2, BCCIP, Pfs2, TREX1, BUB3, FEN1, DRF1, PREI3, CCNE1, RPA1, POLE3, RFC4, MCM3, CHEK1, CCND1, and CDC37; or as selected from CDC2, RFC4, PCNA, CCNE1, CCND1, CDK7, MCM genes (e.g., one or more of MCM3, MCM6, and MCM7), FEN1, MAD2L1, MYBL2, RRM2, and BUB3.

20

In another aspect, the method comprises the determination of the expression levels of all proliferation markers or their expression products, e.g., as listed in Table A, Table B, Table C or Table D; as listed for the group CDC2, MCM6, RPA3, MCM7, PCNA, G22P1, KPNA2, ANLN, APG7L, TOPK, GMNN, RRM1, CDC45L, MAD2L1, RAN, DUT, RRM2, 25 CDK7, MLH3, SMC4L1, CSPG6, POLD2, POLE2, BCCIP, Pfs2, TREX1, BUB3, FEN1, DRF1, PREI3, CCNE1, RPA1, POLE3, RFC4, MCM3, CHEK1, CCND1, and CDC37; or as listed for the group CDC2, RFC4, PCNA, CCNE1, CCND1, CDK7, MCM genes (e.g., one or more of MCM3, MCM6, and MCM7), FEN1, MAD2L1, MYBL2, RRM2, and BUB3.

30 The invention includes the use of archived paraffin-embedded biopsy material for assay of all markers in the set, and therefore is compatible with the most widely available type of biopsy material. It is also compatible with several different methods of tumour tissue harvest, for example, via core biopsy or fine needle aspiration. In a further aspect, RNA is isolated from a fixed, wax-embedded cancer tissue specimen of the patient. Isolation may 35 be performed by any technique known in the art, for example from core biopsy tissue or fine needle aspirate cells.

In another aspect, the invention relates to an array comprising polynucleotides hybridizing to two or more markers as selected from Table A, Table B, Table C or Table D; as selected from CDC2, MCM6, RPA3, MCM7, PCNA, G22P1, KPNA2, ANLN, APG7L, TOPK, GMNN, RRM1, CDC45L, MAD2L1, RAN, DUT, RRM2, CDK7, MLH3, SMC4L1, CSPG6, POLD2, POLE2, BCCIP, Pfs2, TREX1, BUB3, FEN1, DRF1, PREI3, CCNE1, RPA1, POLE3, RFC4, MCM3, CHEK1, CCND1, and CDC37; or as selected from CDC2, RFC4, PCNA, CCNE1, CCND1, CDK7, MCM genes (e.g., one or more of MCM3, MCM6, and MCM7), FEN1, MAD2L1, MYBL2, RRM2, and BUB3.

In particular aspects, the array comprises polynucleotides hybridizing to at least 3, or at least 5, or at least 10, or at least 15, or at least 20, at least 25, at least 30, at least 35, at least 40, at least 45, at least 50, or at least 75 or all of the markers listed in Table A, Table B, Table C or Table D; as listed in the group CDC2, MCM6, RPA3, MCM7, PCNA, G22P1, KPNA2, ANLN, APG7L, TOPK, GMNN, RRM1, CDC45L, MAD2L1, RAN, DUT, RRM2, CDK7, MLH3, SMC4L1, CSPG6, POLD2, POLE2, BCCIP, Pfs2, TREX1, BUB3, FEN1, DRF1, PREI3, CCNE1, RPA1, POLE3, RFC4, MCM3, CHEK1, CCND1, and CDC37; or as listed in the group CDC2, RFC4, PCNA, CCNE1, CCND1, CDK7, MCM genes (e.g., one or more of MCM3, MCM6, and MCM7), FEN1, MAD2L1, MYBL2, RRM2, and BUB3.

In another specific aspect, the array comprises polynucleotides hybridizing to the full set of markers listed in Table A, Table B, Table C or Table D; as listed for the group CDC2, MCM6, RPA3, MCM7, PCNA, G22P1, KPNA2, ANLN, APG7L, TOPK, GMNN, RRM1, CDC45L, MAD2L1, RAN, DUT, RRM2, CDK7, MLH3, SMC4L1, CSPG6, POLD2, POLE2, BCCIP, Pfs2, TREX1, BUB3, FEN1, DRF1, PREI3, CCNE1, RPA1, POLE3, RFC4, MCM3, CHEK1, CCND1, and CDC37; or as listed for the group CDC2, RFC4, PCNA, CCNE1, CCND1, CDK7, MCM genes (e.g., one or more of MCM3, MCM6, and MCM7), FEN1, MAD2L1, MYBL2, RRM2, and BUB3.

The polynucleotides can be cDNAs, or oligonucleotides, and the solid surface on which they are displayed can be glass, for example. The polynucleotides can hybridize to one or more of the markers as disclosed herein, for example, to the full-length sequences, any coding sequences, any fragments, or any complements thereof.

In still another aspect, the invention relates to a method of predicting the likelihood of long-term survival of a patient diagnosed with cancer, without the recurrence of cancer, comprising the steps of: (1) determining the expression levels of the RNA transcripts or the expression products of the full set or a subset of the markers listed in Table A, Table

B, Table C or Table D, herein, in a sample obtained from the patient, normalized against the expression levels of all RNA transcripts or their expression products in the sample, or of a reference set of RNA transcripts or their products; (2) subjecting the data obtained in step (1) to statistical analysis; and (3) determining whether the likelihood of the long-term survival has increased or decreased.

In yet another aspect, the invention concerns a method of preparing a personalized genomics profile for a patient, e.g., a cancer patient, comprising the steps of: (a) subjecting a sample obtained from the patient to expression analysis; (b) determining the expression level of one or more markers selected from the marker set listed in any one of Table A, Table B, Table C or Table D, wherein the expression level is normalized against a control gene or genes and optionally is compared to the amount found in a reference set; and (c) creating a report summarizing the data obtained by the expression analysis. The report may, for example, include prediction of the likelihood of long term survival of the patient and/or recommendation for a treatment modality of the patient.

In additional aspects, the invention relates to a prognostic method comprising: (a) subjecting a sample obtained from a patient to quantitative analysis of the expression level of the RNA transcript of at least one marker selected from Table A, Table B, Table C or Table D, herein, or its product, and (b) identifying the patient as likely to have an increased likelihood of long-term survival without cancer recurrence if the normalized expression levels of the marker or markers, or their products, are above defined expression threshold. In alternate aspects, step (b) comprises identifying the patient as likely to have a decreased likelihood of long-term survival without cancer recurrence if the normalized expression levels of the marker or markers, or their products, are decreased below a defined expression threshold.

In particular, the relatively low expression of proliferation markers is associated with poor outcome. This can include disease progression or the increased likelihood of cancer recurrence, especially for gastrointestinal cancer, such as gastric or colorectal cancer. By contrast, the relatively high expression of proliferation markers is associated with a good outcome. This can include decreased likelihood of cancer recurrence after standard treatment, especially for gastrointestinal cancer, such as gastric or colorectal cancer. Low expression can be determined, for example, by comparison of a test sample (e.g., tumour sample) to samples associated with a positive prognosis. High expression can be determined, for example, by comparison of a test sample (e.g., tumour sample) to samples associated with a negative prognosis.

For example, to obtain a prognosis, a patient's sample (e.g., tumour sample) can be compared to samples with known patient outcome. If the patient's sample shows high expression of GCPMs that is comparable to samples with good outcome, and/or higher than samples with poor outcome, then a positive prognosis is implicated. If the patient's sample shows low expression of GCPMs that is comparable to samples with poor outcome, and/or lower than samples with good outcome, then a negative prognosis is implicated. Alternatively, a patient's sample can be compared to samples of actively proliferating/non-proliferating tumour cells. If the patient's sample shows high expression of GCPMs that is comparable to actively proliferating cells, and/or higher than non-proliferating cells, then a positive prognosis is implicated. If the patient's sample shows low expression of GCPMs that is comparable to non-proliferating cells, and/or lower than actively proliferating cells, then a negative prognosis is implicated.

As further examples, the expression levels of a prognostic signature comprising two or more GCPMs from a patient's sample (e.g., tumour sample) can be compared to samples of recurrent/non-recurrent cancer. If the patient's sample shows increased or decreased expression of GCPMs by comparison to samples of non-recurrent cancer, and/or comparable expression to samples of recurrent cancer, then a negative prognosis is implicated. If the patient's sample shows expression of GCPMs that is comparable to samples of non-recurrent cancer, and/or lower or higher expression than samples of recurrent cancer, then a positive prognosis is implicated.

As one approach, a prediction method can be applied to a panel of markers, for example the panel of GCPMs outlined in Table A, Table B Table C or Table D, in order to generate a predictive model. This involves the generation of a prognostic signature, comprising two or more GCPMs.

The disclosed GCPMs in Table A, Table B, Table C or Table D therefore provide a useful set of markers to generate prediction signatures for determining the prognosis of cancer, and establishing a treatment regime, or treatment modality, specific for that tumour. In particular, a positive prognosis can be used by a patient to decide to pursue standard or less invasive treatment options. A negative prognosis can be used by a patient to decide to terminate treatment or to pursue highly aggressive or experimental treatments. In addition, a patient can chose treatments based on their impact on the expression of prognostic markers (e.g., GCPMs).

Levels of GCPMs can be detected in tumour tissue, tissue proximal to the tumour, lymph node samples, blood samples, serum samples, urine samples, or faecal samples, using any suitable technique, and can include, but is not limited to, oligonucleotide probes, quantitative PCR, or antibodies raised against the markers. It will be appreciated that by
5 analyzing the presence and amounts of expression of a plurality of GCPMs in the form of prediction signatures, and constructing a prognostic signature, the sensitivity and accuracy of prognosis will be increased. Therefore, multiple markers according to the present invention can be used to determine the prognosis of a cancer.

10 The invention includes the use of archived paraffin-embedded biopsy material for assay of the markers in the set, and therefore is compatible with the most widely available type of biopsy material. It is also compatible with several different methods of tumour tissue harvest, for example, via core biopsy or fine needle aspiration. In certain aspects, RNA is isolated from a fixed, wax-embedded cancer tissue specimen of the patient. Isolation may
15 be performed by any technique known in the art, for example from core biopsy tissue or fine needle aspirate cells.

In one aspect, the invention relates to a method of predicting a prognosis, e.g., the likelihood of long-term survival of a cancer patient without the recurrence of cancer,
20 comprising determining the expression level of one or more prognostic markers or their expression products in a sample obtained from the patient, normalized against the expression level of other RNA transcripts or their products in the sample, or of a reference set of RNA transcripts or their expression products. In specific aspects, the prognostic marker is one or more markers listed in Table A, Table B, Table C or Table D or is
25 included as one or more of the prognostic signatures derived from the markers listed in Table A, Table B, Table C or Table D.

In further aspects, the expression levels of the prognostic markers or their expression products are determined, e.g., for the markers listed in Table A, Table B, Table C or Table
30 D, a prognostic signature derived from the markers listed in Table A, Table B, Table C or Table D. In another aspect, the method comprises the determination of the expression levels of a full set of prognosis markers or their expression products, e.g., for the markers listed in Table A, Table B, Table C or Table D, or, a prognostic signature derived from the markers listed in Table A, Table B, Table C or Table D.

35

In an additional aspect, the invention relates to an array (e.g., microarray) comprising polynucleotides hybridizing to two or more markers, e.g., for the markers listed in Table A,

Table B, Table C or Table D, or a prognostic signature derived from the markers listed in Table A, Table B, Table C or Table D. In particular aspects, the array comprises polynucleotides hybridizing to prognostic signature derived from the markers listed in Table A, Table B, Table C or Table D, or e.g., for a prognostic signature. In another
5 specific aspect, the array comprises polynucleotides hybridizing to the full set of markers, e.g., for the markers listed in Table A, Table B, Table C or Table D, or, e.g., for a prognostic signature.

For these arrays, the polynucleotides can be cDNAs, or oligonucleotides, and the solid
10 surface on which they are displayed can be glass, for example. The polynucleotides can hybridize to one or more of the markers as disclosed herein, for example, to the full-length sequences, any coding sequences, any fragments, or any complements thereof. In particular aspects, an increase or decrease in expression levels of one or more GCPM indicates a decreased likelihood of long-term survival, e.g., due to cancer recurrence,
15 while a lack of an increase or decrease in expression levels of one or more GCPM indicates an increased likelihood of long-term survival without cancer recurrence.

In further aspects, the invention relates to a kit comprising one or more of: (1) extraction buffer/reagents and protocol; (2) reverse transcription buffer/reagents and protocol; and
20 (3) quantitative PCR buffer/reagents and protocol suitable for performing any of the foregoing methods. Other aspects and advantages of the invention are illustrated in the description and examples included herein.

Table A: GCPMs for cell proliferation signature

Unique ID	Gene Symbol	Gene Name	GenBank Acc. No.	Gene Aliases
A:09020	CCND1	cyclin D1	NM_053056	BCL1; PRAD1; U21B31; D11S287E
C:0921	CCNE1	cyclin E1	NM_001238, NM_057182	CCNE
A:05382	CDC2	cell division cycle 2, G1 to S and G2 to M	NM_001786, NM_033379	CDK1; MGC111195; DKFZp686L2022 2
A:09842	CDK7	cyclin-dependent kinase 7 (MO15 homolog, <i>Xenopus laevis</i> , cdk-activating kinase)	NM_001799	CAK1; STK1; CDKN7; p39MO15
B:7793	CHEK1	CHK1 checkpoint homolog (S. pombe)	NM_001274	CHK1
A:03447	CSE1L	CSE1 chromosome segregation 1-like (yeast)	NM_001316	CAS; CSE1; XPO2; MGC117283; MGC130036; MGC130037
A:05535	DKC1	dyskeratosis congenita 1, dyskerin	NM_001363	DKC; NAP57; NOLA4; XAP101; dyskerin
A:07296	DUT	dUTP pyrophosphatase	NM_001025248, NM_001025249, NM_001948	dUTPase; FLJ20622
C:2467	E4F1	E4F transcription factor 1	NM_004424	E4F; MGC99614
B:9065	FEN1	flap structure-specific endonuclease 1	NM_004111	MF1; RAD2; FEN-1
A:01437	FH	fumarate hydratase	NM_000143	MCL; LRCC; HLRCC; MCUL1
B:9714	XRCC6	X-ray repair complementing defective repair in Chinese hamster cells 6 (Ku autoantigen, 70kDa)	NM_001469	ML8; KU70; TLAA; CTC75; CTCBF; G22P1
B:3553_hk-r1	GPS1	G protein pathway suppressor 1	NM_004127; NM_212492	CSN1; COPS1; MGC71287
B:4036	KPNA2	karyopherin alpha 2 (RAG cohort 1, importin alpha 1)	NM_002266	QIP2; RCH1; IPOA1; SRP1alpha
A:06387	MAD2L1	MAD2 mitotic arrest deficient-like 1 (yeast)	NM_002358	MAD2; HSMAD2

A:08668	MCM3	MCM3 minichromosome maintenance deficient 3 (<i>S. cerevisiae</i>)	NM_002388	HCC5; P1.h; RLFb; MGC1157; P1-MCM3
B:8147	MCM6	MCM6 minichromosome maintenance deficient 6 (MIS5 homolog, <i>S. pombe</i>) (<i>S. cerevisiae</i>)	NM_005915	Mis5; P105MCM; MCG40308
B:7620	MCM7	MCM7 minichromosome maintenance deficient 7 (<i>S. cerevisiae</i>)	NM_005916, NM_182776	MCM2; CDC47; P85MCM; P1CDC47; PNAS-146; CDABP0042; P1.1-MCM3
A:10600	RAB8A	RAB8A, member RAS oncogene family	NM_005370	MEL; RAB8
A:09470	KITLG	KIT ligand	NM_000899, NM_003994	SF; MGF; SCF; KL-1; Klt; DKFZp686F2250
A:06037	MYBL2	v-myb myeloblastosis viral oncogene homolog (avian)-like 2	NM_002466	BMVB; MGC15600
A:01677	NME1	non-metastatic cellis 1, protein (NM23A) expressed in	NM_000269, NM_198175	AWD; GAAD; NM23; NDPKA; NM23-H1
A:03397	PRDX1	peroxiredoxin 1	NM_002574, NM_181696, NM_181697	PAG; PAGA; PAGB; MSP23; NKEFA; TDPX2
A:03715	PCNA	proliferating cell nuclear antigen	NM_002592, NM_182649	MGC8367
A:02929	POLD2	polymerase (DNA directed), delta 2, regulatory subunit 50kDa	NM_006230	None
A:04680	POLE2	polymerase (DNA directed), epsilon 2 (p59 subunit)	NM_002692	DPE2
A:09169	RAN	RAN, member RAS oncogene family	NM_006325	TC4; Gsp1; ARA24
A:09145	RBBP8	retinoblastoma binding protein 8	NM_002894, NM_203291, NM_203292	RIM; CTIP
A:09921	RFC4	replication factor C (activator 1) 4, 37kDa	NM_002916, NM_181573	A1; RFC37; MGC27291
A:10597	RPA1	replication protein A1, 70kDa	NM_002945	HSSB; RF-A; RP-A; REPA1; RPA70
A:00231	RPA3	replication protein A3, 14kDa	NM_002947	REPA3

A:09802	RRM1	ribonucleotide reductase M1 polypeptide	NM_001033	R1; RR1; RIR1
B:3501	RRM2	ribonucleotide reductase M2 polypeptide	NM_001034	R2; RR2M
A:08332	S100A5	S100 calcium binding protein A5	NM_002962	S100D
A:07314	FSCN1	fascin homolog 1, actin-bundling protein (Strongylocentrotus purpuratus)	NM_003088	SNL; p55; FLJ38511
A:03507	FOSL1	FOS-like antigen 1	NM_005438	FRA1; fra-1
A:09331	CDC45L	CDC45 cell division cycle 45-like (S. cerevisiae)	NM_003504	CDC45; CDC45L2; PORC-PI-1
A:09436	SMC3	structural maintenance of chromosomes 3	NM_005445	BAM; BMH; HCAP; CSPG6; SMC3L1
A:09747	BUB3	BUB3 budding uninhibited by benzimidazoles 3 homolog (yeast)	NM_001007793, NM_004725	BUB3L; hBUB3
A:00891	WDR39	WD repeat domain 39	NM_004804	CIAO1
A:05648	SMC4	structural maintenance of chromosomes 4	NM_001002799, NM_001002800, NM_005496	CAPC; SMC4L1; hCAP-C
B:7911	TOB1	transducer of ERBB2, 1	NM_005749	TOB; TROB; APRO6; PIG49; TROB1; MGC34446; MGC104792
A:04760	ATG7	ATG7 autophagy related 7 homolog (S. cerevisiae)	NM_006395	GSA7; APG7L; DKFZp434N0735
A:04950	CCT7	chaperonin containing TCP1, subunit 7 (eta)	NM_001009570, NM_006429	Ccth; Nip7-1; CCT-ETA; MGC110985; TCP-1-eta
A:09500	CCT2	chaperonin containing TCP1, subunit 2 (beta)	NM_006431	CCTB; 99D8.1; PRO1633; CCT-beta; MGC142074; MGC142076; TCP-1-beta
A:03486	CDC37	CDC37 cell division cycle 37 homolog (S. cerevisiae)	NM_007065	P50CDC37
B:7247	TREX1	three prime repair exonuclease 1	NM_016381, NM_032166, NM_033627, NM_033628, NM_033629, NM_130384	AGS1; DRN3; ATRIP; FLJ12343; DKFZp434J0310

A:01322	PARK7	Parkinson disease (autosomal recessive, early onset) 7	NM_007262	DJ1; DJ-1; FLJ27376
A:09401	PREI3	preimplantation protein 3	NM_015387, NM_199482	2C4D; MOB1; MOB3; CGI-95; MGC12264
A:09724	MLH3	mutL homolog 3 (E. coli)	NM_001040108, NM_014381	HNPCC7; MGC138372
A:02984	CACYBP	calcyclin binding protein	NM_001007214, NM_014412	SIP; GIG5; MGC87971; PNAS-107; S100A6BP; RP1-102G20.6
A:09821	MCTS1	malignant T cell amplified sequence 1	NM_014060	MCT1; MCT-1
A:03435	GMNN	geminin, DNA replication inhibitor	NM_015895	Gem; RP3-369A17.3
B:1035	GIN52	GIN5 complex subunit 2 (Psf2 homolog)	NM_016095	PSF2; Pfs2; HSPC037
A:02209	POLE3	polymerase (DNA directed), epsilon 3 (p17 subunit)	NM_017443	p17; YBL1; CHRAC17; CHARAC17
A:05280	ANLN	anillin, actin binding protein	NM_018685	scra; Scraps; ANILLIN; DKFZp779A055
A:07468	SEPT11	septin 11	NM_018243	None
A:03912	PBK	PDZ binding kinase	NM_018492	SPK; TOPK; Nori-3; FLJ14385
B:8449	BCCIP	BRCA2 and CDKN1A interacting protein	NM_016567, NM_078468, NM_078469	TOK-1
B:2392	DBF4B	DBF4 homolog B (S. cerevisiae)	NM_025104, NM_145663	DRF1; ASKL1; FLJ13087; MGC15009
B:6501	CD276	CD276 molecule	NM_001024736, NM_025240	B7H3; B7-H3
B:5467	LAMA1	laminin, alpha 1	NM_005559	LAMA

Table A: Proliferation-related genes differentially expressed between cell lines in high and low proliferative states. Genes that were differentially expressed between cell lines in confluent (low proliferation) and semi-confluent (high proliferation) states (see Figure 1) were identified by microarray analysis on 30K MWG Biotech arrays. Table A comprises the subset of these genes that were categorized by gene ontology analysis as cell proliferation-related.

Table B: GCPMs for cell proliferation signature

Unique ID	Gene Description	LocusLink	GenBank Accession
B:7560	v-abl Abelson murine leukaemia viral oncogene homolog 1 (ABL1), transcript variant a, mRNA	25	NM_005157
A:09071	acetylcholinesterase (YT blood group) (ACHE), transcript variant E4-E5, mRNA	43	NM_015831, NM_000665
A:04114	acid phosphatase 2, lysosomal (ACP2), mRNA	53	NM_001610
A:09146	acid phosphatase, prostate (ACPP), mRNA	55	NM_001099
A:09585	adrenergic, alpha-1D-, receptor (ADRA1D), mRNA	146	NM_000678
A:08793	adrenergic, alpha-1B-, receptor (ADRA1B), mRNA	147	NM_000679
C:0326	adrenergic, alpha-1A-, receptor (ADRA1A), transcript variant 4, mRNA	148	NM_033304
A:02272	adrenergic, alpha-2A-, receptor (ADRA2A), mRNA	150	NM_000681
A:05807	jagged 1 (Alagille syndrome) (JAG1), mRNA	182	NM_000214
A:02268	aryl hydrocarbon receptor (AHR), mRNA	196	NM_001621
A:00978	allograft inflammatory factor 1 (AIF1), transcript variant 2, mRNA	199	NM_004847
A:06335	adenylate kinase 1 (AK1), mRNA	203	NM_000476
A:07028	v-akt murine thymoma viral oncogene homolog 1 (AKT1), transcript variant 1, mRNA	207	NM_005163
A:05949	v-akt murine thymoma viral oncogene homolog 2 (AKT2), mRNA	208	NM_001626
B:9542	arachidonate 15-lipoxygenase, second type (ALOX15B), mRNA	247	NM_001141
A:02569	bridging integrator 1 (BIN1), transcript variant 8, mRNA	274	NM_004305
C:0393	amyloid beta (A4) precursor protein-binding, family B, member 1 (Fe65) (APBB1), transcript variant 1, mRNA	322	NM_001164
B:5288	amyloid beta (A4) precursor protein-binding, family B, member 2 (Fe65-like) (APBB2), mRNA	323	NM_173075
A:09151	adenomatosis polyposis coli (APC), mRNA	324	NM_000038
B:3616	baculoviral IAP repeat-containing 5 (survivin) (BIRC5), transcript variant 1, mRNA	332	NM_001168
C:2007	androgen receptor (dihydrotestosterone receptor; testicular feminization; spinal and bulbar muscular atrophy; Kennedy disease) (AR), transcript variant 2, mRNA	367	NM_001011645
A:04819	amphiregulin (schwannoma-derived growth factor) (AREG), mRNA	374	NM_001657

A:01709	ras homolog gene family, member G (rho G) (RHOG), mRNA	391	NM_001665
B:6554	ataxia telangiectasia mutated (includes complementation groups A, C and D) (ATM), transcript variant 1, mRNA	472	NM_000051
A:02418	ATPase, Cu++ transporting, beta polypeptide (ATP7B), transcript variant 1, mRNA	545	NM_000053
A:05997	AXL receptor tyrosine kinase (AXL), transcript variant 2, mRNA	558	NM_001699
B:0073	brain-specific angiogenesis inhibitor 1 (BAI1), mRNA	575	NM_001702
A:07209	BCL2-associated X protein (BAX), transcript variant beta, mRNA	581	NM_004324
B:1845	Bardet-Biedl syndrome 4 (BBS4), mRNA	586	NM_033028
A:00571	branched chain aminotransferase 2, mitochondrial (BCAT2), mRNA	588	NM_001190
A:09020	cyclin D1 (CCND1), mRNA	595	NM_053056
A:10775	B-cell CLL/lymphoma 2 (BCL2), nuclear gene encoding mitochondrial protein, transcript variant alpha, mRNA	596	NM_000633
A:09014	B-cell CLL/lymphoma 3 (BCL3), mRNA	602	NM_005178
C:2412	B-cell CLL/lymphoma 6 (zinc finger protein 51) (BCL6), transcript variant 1, mRNA	604	NM_001706
A:08794	tumour necrosis factor receptor superfamily, member 17 (TNFRSF17), mRNA	608	NM_001192
A:01162	Bloom syndrome (BLM), mRNA	641	NM_000057
B:5276	basonuclein 1 (BNC1), mRNA	646	NM_001717
B:3766	polymerase (RNA) III (DNA directed) polypeptide D, 44kDa (POLR3D), mRNA	661	NM_001722
C:2188	dystonin (DST), transcript variant 1, mRNA	667	NM_183380
B:5103	breast cancer 1, early onset (BRCA1), transcript variant BRCA1a, mRNA	672	NM_007294
A:03676	breast cancer 2, early onset (BRCA2), mRNA	675	NM_000059
A:07404	zinc finger protein 36, C3H type-like 1 (ZFP36L1), mRNA	677	NM_004926
B:5146	zinc finger protein 36, C3H type-like 2 (ZFP36L2), mRNA	678	NM_006887
B:4758	bone marrow stromal cell antigen 2 (BST2), mRNA	684	NM_004335
B:4642	betacellulin (BTC), mRNA	685	NM_001729
C:2483	B-cell translocation gene 1, anti-proliferative (BTG1), mRNA	694	NM_001731
B:0618	BUB1 budding uninhibited by benzimidazoles 1 homolog (yeast) (BUB1), mRNA	699	NM_004336
A:09398	BUB1 budding uninhibited by benzimidazoles 1 homolog beta (yeast) (BUB1B), mRNA	701	NM_001211

A:01104	chromosome 8 open reading frame 1 (C8orf1), mRNA	734	NM_004337
B:3828	calmodulin 2 (phosphorylase kinase, delta) (CALM2), mRNA	805	NM_001743
B:6851	calpain 1, (mu/I) large subunit (CAPN1), mRNA	823	NM_005186
A:09763	calpain, small subunit 1 (CAPNS1), transcript variant 1, mRNA	826	NM_001749
B:0205	core-binding factor, runt domain, alpha subunit 2; translocated to, 3 (CBFA2T3), transcript variant 2, mRNA	863	NM_175931
B:2901	runt-related transcription factor 3 (RUNX3), transcript variant 2, mRNA	864	NM_004350
A:01132	cholecystokinin B receptor (CCKBR), mRNA	887	NM_176875
A:04253	cyclin A2 (CCNA2), mRNA	890	NM_001237
A:04253	cyclin A2 (CCNA2), mRNA	891	NM_001237
A:09352	cyclin C (CCNC), transcript variant 1, mRNA	892	NM_005190
A:10559	cyclin D2 (CCND2), mRNA	894	NM_001759
A:02240	cyclin D3 (CCND3), mRNA	896	NM_001760
C:0921	cyclin E1 (CCNE1), transcript variant 1, mRNA	898	NM_001238
C:0921	cyclin E1 (CCNE1), transcript variant 1, mRNA	899	NM_001238
B:5261	cyclin G1 (CCNG1), transcript variant 1, mRNA	900	NM_004060
A:07154	cyclin G2 (CCNG2), mRNA	901	NM_004354
A:07930	cyclin H (CCNH), mRNA	902	NM_001239
A:01253	cyclin T1 (CCNT1), mRNA	904	NM_001240
B:0645	cyclin T2 (CCNT2), transcript variant b, mRNA	905	NM_058241
C:2676	CD3E antigen, epsilon polypeptide (TIT3 complex) (CD3E), mRNA	916	NM_000733
A:10068	CD5 antigen (p56-62) (CD5), mRNA	921	NM_014207
A:07504	tumour necrosis factor receptor superfamily, member 7 (TNFRSF7), mRNA	939	NM_001242
A:05558	CD28 antigen (Tp44) (CD28), mRNA	940	NM_006139
A:07387	CD86 antigen (CD28 antigen ligand 2, B7-2 antigen) (CD86), transcript variant 1, mRNA	942	NM_175862
A:06344	tumour necrosis factor receptor superfamily, member 8 (TNFRSF8), transcript variant 1, mRNA	943	NM_001243
A:03064	tumour necrosis factor (ligand) superfamily, member 8 (TNFSF8), mRNA	944	NM_001244
A:03802	CD33 antigen (gp67) (CD33), mRNA	945	NM_001772
A:07407	CD40 antigen (TNF receptor superfamily member 5) (CD40), transcript variant 1, mRNA	958	NM_001250

B:9757	CD40 ligand (TNF superfamily, member 5, hyper-IgM syndrome) (CD40LG), mRNA	959	NM_000074
A:07070	CD68 antigen (CD68), mRNA	968	NM_001251
A:04715	tumour necrosis factor (ligand) superfamily, member 7 (TNFSF7), mRNA	970	NM_001252
A:09638	CD81 antigen (target of antiproliferative antibody 1) (CD81), mRNA	975	NM_004356
A:05382	cell division cycle 2, G1 to S and G2 to M (CDC2), transcript variant 1, mRNA	983	NM_001786
A:00282	cell division cycle 2-like 1 (PITSLRE proteins) (CDC2L1), transcript variant 2, mRNA	984	NM_033486
A:00282	cell division cycle 2-like 1 (PITSLRE proteins) (CDC2L1), transcript variant 2, mRNA	985	NM_033486
A:07718	CDC5 cell division cycle 5-like (S. pombe) (CDC5L), mRNA	988	NM_001253
A:00843	septin 7 (SEPT7), transcript variant 1, mRNA	989	NM_001788
A:05789	CDC6 cell division cycle 6 homolog (S. cerevisiae) (CDC6), mRNA	990	NM_001254
A:03063	CDC20 cell division cycle 20 homolog (S. cerevisiae) (CDC20), mRNA	991	NM_001255
B:4185	cell division cycle 25A (CDC25A), transcript variant 1, mRNA	993	NM_001789
A:04022	cell division cycle 25B (CDC25B), transcript variant 3, mRNA	994	NM_021873
B:9539	cell division cycle 25C (CDC25C), transcript variant 1, mRNA	995	NM_001790
B:5590	cell division cycle 27 (CDC27)	996	NM_001256
B:9041	cell division cycle 34 (CDC34), mRNA	997	NM_004359
A:03518	cyclin-dependent kinase 2 (CDK2), transcript variant 2, mRNA	1017	NM_052827
A:02068	cyclin-dependent kinase 3 (CDK3), mRNA	1018	NM_001258
B:4838	cyclin-dependent kinase 4 (CDK4), mRNA	1019	NM_000075
A:10302	cyclin-dependent kinase 5 (CDK5), mRNA	1020	NM_004935
A:01923	cyclin-dependent kinase 6 (CDK6), mRNA	1021	NM_001259
A:09842	cyclin-dependent kinase 7 (MO15 homolog, Xenopus laevis, cdk-activating kinase) (CDK7), mRNA	1022	NM_001799
A:08302	cyclin-dependent kinase 8 (CDK8), mRNA	1024	NM_001260
A:05151	cyclin-dependent kinase 9 (CDC2-related kinase) (CDK9), mRNA	1025	NM_001261
A:09736	cyclin-dependent kinase inhibitor 1A (p21, Cip1) (CDKN1A), transcript variant 2, mRNA	1026	NM_078467

A:05571	cyclin-dependent kinase inhibitor 1B (p27, Kip1) (CDKN1B), mRNA	1027	NM_004064
A:08441	cyclin-dependent kinase inhibitor 1C (p57, Kip2) (CDKN1C), mRNA	1028	NM_000076
B:9782	cyclin-dependent kinase inhibitor 2A (melanoma, p16, inhibits CDK4) (CDKN2A), transcript variant 4, mRNA	1029	NM_058195
C:6459	cyclin-dependent kinase inhibitor 2B (p15, inhibits CDK4) (CDKN2B), transcript variant 1, mRNA	1030	NM_004936
B:0604	cyclin-dependent kinase inhibitor 2C (p18, inhibits CDK4) (CDKN2C), transcript variant 1, mRNA	1031	NM_001262
A:03310	cyclin-dependent kinase inhibitor 2D (p19, inhibits CDK4) (CDKN2D), transcript variant 2, mRNA	1032	NM_079421
A:05799	cyclin-dependent kinase inhibitor 3 (CDK2-associated dual specificity phosphatase) (CDKN3), mRNA	1033	NM_005192
B:9170	centromere protein B, 80kDa (CENPB), mRNA	1059	NM_001810
A:07769	centromere protein E, 312kDa (CENPE), mRNA	1062	NM_001813
A:06471	centromere protein F, 350/400ka (mitosin) (CENPF), mRNA	1063	NM_016343
A:03128	centrin, EF-hand protein, 1 (CETN1), mRNA	1068	NM_004066
A:05554	centrin, EF-hand protein, 2 (CETN2), mRNA	1069	NM_004344
B:4016	centrin, EF-hand protein, 3 (CDC31 homolog, yeast) (CETN3), mRNA	1070	NM_004365
B:5082	regulator of chromosome condensation 1 RCC1	1104	NM_001048194, NM_001048195, NM_001269
B:7793	CHK1 checkpoint homolog (S. pombe) (CHEK1), mRNA	1111	NM_001274
B:8504	checkpoint suppressor 1 (CHES1), mRNA	1112	NM_005197
A:00320	cholinergic receptor, muscarinic 1 (CHRM1), mRNA	1128	NM_000738
A:10168	cholinergic receptor, muscarinic 3 (CHRM3), mRNA	1131	NM_000740
A:06655	cholinergic receptor, muscarinic 4 (CHRM4), mRNA	1132	NM_000741
A:00869	cholinergic receptor, muscarinic 5 (CHRM5), mRNA	1133	NM_012125
C:0649	CDC28 protein kinase regulatory subunit 1B (CKS1B), mRNA	1163	NM_001826
B:6912	CDC28 protein kinase regulatory subunit 2 (CKS2), mRNA	1164	NM_001827
A:07840	CDC-like kinase 1 (CLK1), transcript variant 1, mRNA	1195	NM_004071
B:8665	polo-like kinase 3 (Drosophila) (PLK3), mRNA	1263	NM_004073
B:8651	collagen, type IV, alpha 3 (Goodpasture antigen) (COL4A3), transcript variant 1, mRNA	1285	NM_000091

B:4734	mitogen-activated protein kinase 8 (MAP3K8), mRNA	1326	NM_005204
B:3778	cysteine-rich protein 1 (intestinal) (CRIP1), mRNA	1396	NM_001311
B:3581	cysteine-rich protein 2 (CRIP2), mRNA	1397	NM_001312
B:5543	v-crk sarcoma virus CT10 oncogene homolog (avian) (CRK), transcript variant 1, mRNA	1398	NM_005206
B:6254	v-crk sarcoma virus CT10 oncogene homolog (avian)-like (CRKL), mRNA	1399	NM_005207
A:03447	CSE1 chromosome segregation 1-like (yeast) (CSE1L), transcript variant 2, mRNA	1434	NM_177436
A:10730	colony stimulating factor 1 (macrophage) (CSF1), transcript variant 2, mRNA	1435	NM_172210
A:05457	colony stimulating factor 1 receptor, formerly McDonough feline sarcoma viral (v-fms) oncogene homolog (CSF1R), mRNA	1436	NM_005211
B:1908	colony stimulating factor 3 (granulocyte) (CSF3), transcript variant 2, mRNA	1440	NM_172219
A:01629	c-src tyrosine kinase (CSK), mRNA	1445	NM_004383
A:07097	casein kinase 2, alpha prime polypeptide (CSNK2A2), mRNA	1459	NM_001896
B:3639	cysteine and glycine-rich protein 2 (CSRP2), mRNA	1466	NM_001321
B:8929	C-terminal binding protein 1 CTBP1	1487	NM_001012614, NM_001328
A:08689	C-terminal binding protein 2 (CTBP2), transcript variant 1, mRNA	1488	NM_001329
A:02604	cardiotrophin 1 (CTF1), mRNA	1489	NM_001330
A:05018	disabled homolog 2, mitogen-responsive phosphoprotein (Drosophila) (DAB2), mRNA	1601	NM_001343
A:09374	deleted in colorectal carcinoma (DCC), mRNA	1630	NM_005215
A:05576	dynactin 1 (p150, glued homolog, Drosophila) (DCTN1), transcript variant 1, mRNA	1639	NM_004082
A:04346	growth arrest and DNA-damage-inducible, alpha (GADD45A), mRNA	1647	NM_001924
B:9526	DNA-damage-inducible transcript 3 (DDIT3), mRNA	1649	NM_004083
B:6726	DEAD/H (Asp-Glu-Ala-Asp/His) box polypeptide 11 (CHL1-like helicase homolog, S. cerevisiae) (DDX11), transcript variant 1, mRNA	1663	NM_030653
B:1955	deoxyhypusine synthase (DHPS), transcript variant 1, mRNA	1725	NM_001930
A:09887	diaphanous homolog 2 (Drosophila) (DIAPH2), transcript variant 12C, mRNA	1730	NM_007309
B:4704	septin 1 (SEPT1), mRNA	1731	NM_052838

A:05535	dyskeratosis congenita 1, dyskerin (DKC1), mRNA	1736	NM_001363
A:06695	discs, large homolog 3 (neuroendocrine-dlg, Drosophila) (DLG3), mRNA	1741	NM_021120
B:9032	dystrophin myotonia-containing WD repeat motif (DMWD), mRNA	1762	NM_004943
B:4936	DNA2 DNA replication helicase 2-like (yeast) (DNA2L), mRNA	1763	XM_166103, XM_938629
B:5286	dynein, cytoplasmic 1, heavy chain 1 (DYNC1H1), mRNA	1778	NM_001376
B:9089	dynamitin 2 (DNM2), transcript variant 4, mRNA	1785	NM_001005362
A:05674	deoxynucleotidyltransferase, terminal (DNTT), transcript variant 1, mRNA	1791	NM_004088
A:00269	heparin-binding EGF-like growth factor (HBEGF), mRNA	1839	NM_001945
B:3724	deoxythymidylate kinase (thymidylate kinase) (DTYMK), mRNA	1841	NM_012145
A:01114	dual specificity phosphatase 1 (DUSP1), mRNA	1843	NM_004417
A:08044	dual specificity phosphatase 4 (DUSP4), transcript variant 2, mRNA	1846	NM_057158
B:0206	dual specificity phosphatase 6 (DUSP6), transcript variant 1, mRNA	1848	NM_001946
A:07296	dUTP pyrophosphatase (DUT), nuclear gene encoding mitochondrial protein, transcript variant 2, mRNA	1854	NM_001948
B:5540	E2F transcription factor 1 (E2F1), mRNA	1869	NM_005225
B:4216	E2F transcription factor 2 (E2F2), mRNA	1870	NM_004091
B:6451	E2F transcription factor 3 (E2F3), mRNA	1871	NM_001949
A:03567	E2F transcription factor 4, p107/p130-binding (E2F4), mRNA	1874	NM_001950
C:2484	E2F transcription factor 5, p130-binding (E2F5), mRNA	1875	NM_001951
B:9807	E2F transcription factor 6 (E2F6), transcript variant a, mRNA	1876	NM_001952
C:2467	E4F transcription factor 1 (E4F1), mRNA	1877	NM_004424
A:04592	endothelial cell growth factor 1 (platelet-derived) (ECGF1), mRNA	1890	NM_001953
A:00257	endothelial differentiation, lysophosphatidic acid G-protein-coupled receptor, 2 (EDG2), transcript variant 1, mRNA	1903	NM_001401
A:08155	endothelin 1 (EDN1), mRNA	1906	NM_001955
A:08447	endothelin receptor type A (EDNRA), mRNA	1909	NM_001957
A:09410	epidermal growth factor (beta-urogastrone) (EGF), mRNA	1950	NM_001963

A:10005	epidermal growth factor receptor (erythroblastic leukaemia viral (v-erb-b) oncogene homolog, avian) (EGFR), transcript variant 1, mRNA	1956	NM_005228
A:03312	early growth response 4 (EGR4), mRNA	1961	NM_001965
A:06719	eukaryotic translation initiation factor 4 gamma, 2 (EIF4G2), mRNA	1982	NM_001418
A:10651	E74-like factor 5 (ets domain transcription factor) (ELF5), transcript variant 2, mRNA	2001	NM_001422
A:07972	ELK3, ETS-domain protein (SRF accessory protein 2) (ELK3), mRNA	2004	NM_005230
A:06224	elastin (supravalvular aortic stenosis, Williams-Beuren syndrome) (ELN), mRNA	2006	NM_000501
A:10267	epithelial membrane protein 1 (EMP1), mRNA	2012	NM_001423
A:09610	epithelial membrane protein 2 (EMP2), mRNA	2013	NM_001424
A:00767	epithelial membrane protein 3 (EMP3), mRNA	2014	NM_001425
A:07219	glutamyl aminopeptidase (aminopeptidase A) (ENPEP), mRNA	2028	NM_001977
A:10199	E1A binding protein p300 (EP300), mRNA	2033	NM_001429
A:10325	EPH receptor B4 (EPHB4), mRNA	2050	NM_004444
A:04352	glutamyl-prolyl-tRNA synthetase (EPRS), mRNA	2059	NM_004446
A:04352	glutamyl-prolyl-tRNA synthetase (EPRS), mRNA	2060	NM_004446
A:08200	nuclear receptor subfamily 2, group F, member 6 (NR2F6), mRNA	2063	NM_005234
B:1429	v-erb-b2 erythroblastic leukaemia viral oncogene homolog 2, neuro/glioblastoma derived oncogene homolog (avian) ERBB2	2064	NM_001005862, NM_004448
A:02313	v-erb-a erythroblastic leukaemia viral oncogene homolog 4 (avian) (ERBB4), mRNA	2066	NM_005235
A:08898	epiregulin (EREG), mRNA	2069	NM_001432
A:07916	Ets2 repressor factor (ERF), mRNA	2077	NM_006494
B:9779	v-ets erythroblastosis virus E26 oncogene like (avian) (ERG), transcript variant 1, mRNA	2078	NM_182918
C:2388	enhancer of rudimentary homolog (Drosophila) (ERH), mRNA	2079	NM_004450
B:5360	endogenous retroviral sequence K(C4), 2 ERVK2	2087	U87595
C:2799	estrogen receptor 1 (ESR1), mRNA	2099	NM_000125
A:01596	v-ets erythroblastosis virus E26 oncogene homolog 1 (avian) (ETS1), mRNA	2113	NM_005238
A:07704	v-ets erythroblastosis virus E26 oncogene homolog 2 (avian) (ETS2), mRNA	2114	NM_005239

A:00924	ecotropic viral integration site 2A (EVI2A), transcript variant 2, mRNA	2123	NM_014210
A:07732	exostoses (multiple) 1 (EXT1), mRNA	2131	NM_000127
A:10493	exostoses (multiple) 2 (EXT2), transcript variant 1, mRNA	2132	NM_000401
A:07741	coagulation factor II (thrombin) (F2), mRNA	2147	NM_000506
A:06727	coagulation factor II (thrombin) receptor (F2R), mRNA	2149	NM_001992
A:10554	fatty acid binding protein 3, muscle and heart (mammary-derived growth inhibitor) (FABP3), mRNA	2170	NM_004102
A:10780	fatty acid binding protein 5 (psoriasis-associated) (FABP5), mRNA	2172	NM_001444
B:9700	fatty acid binding protein 7, brain FABP7	2173	NM_001446
C:2632	PTK2B protein tyrosine kinase 2 beta (PTK2B), transcript variant 1, mRNA	2185	NM_173174
A:07570	Fanconi anemia, complementation group G (FANCG), mRNA	2189	NM_004629
A:08248	membrane-spanning 4-domains, subfamily A, member 2 (Fc fragment of IgE, high affinity I, receptor for; beta polypeptide) (MS4A2), mRNA	2206	NM_000139
B:9065	flap structure-specific endonuclease 1 (FEN1), mRNA	2237	NM_004111
A:10689	glypican 4 (GPC4), mRNA	2239	NM_001448
B:7897	fer (fps/fes related) tyrosine kinase (phosphoprotein NCP94) (FER), mRNA	2242	NM_005246
B:1852	fibrinogen alpha chain (FGA), transcript variant alpha-E, mRNA	2243	NM_000508
B:1909	fibrinogen beta chain (FGB), mRNA	2244	NM_005141
A:07894	fibroblast growth factor 1 (acidic) (FGF1), transcript variant 1, mRNA	2246	NM_000800
B:7727	fibroblast growth factor 2 (basic) (FGF2), mRNA	2247	NM_002006
A:01551	fibroblast growth factor 3 (murine mammary tumour virus integration site (v-int-2) oncogene homolog) (FGF3), mRNA	2248	NM_005247
A:10568	fibroblast growth factor 4 (heparin secretory transforming protein 1, Kaposi sarcoma oncogene) (FGF4), mRNA	2249	NM_002007
C:2679	fibroblast growth factor 5 (FGF5), transcript variant 2, mRNA	2250	NM_033143
A:04438	fibroblast growth factor 6 (FGF6), mRNA	2251	NM_020996
C:2713	fibroblast growth factor 7 (keratinocyte growth factor) (FGF7), mRNA	2252	NM_002009
B:8151	fibroblast growth factor 8 (androgen-induced) (FGF8),	2253	NM_006119

	transcript variant B, mRNA		
A:10353	fibroblast growth factor 9 (glia-activating factor) (FGF9), mRNA	2254	NM_002010
A:10837	fibroblast growth factor 10 (FGF10), mRNA	2255	NM_004465
B:1815	fibrinogen gamma chain (FGG), transcript variant gamma-B, mRNA	2266	NM_021870
A:01437	fumarate hydratase (FH), nuclear gene encoding mitochondrial protein, mRNA	2271	NM_000143
A:04648	fragile histidine triad gene (FHIT), mRNA	2272	NM_002012
B:1938	c-fos induced growth factor (vascular endothelial growth factor D) (FIGF), mRNA	2277	NM_004469
B:5100	fms-related tyrosine kinase 1 (vascular endothelial growth factor/vascular permeability factor receptor) FLT1	2321	NM_002019
A:05859	fms-related tyrosine kinase 3 (FLT3), mRNA	2322	NM_004119
A:05362	fms-related tyrosine kinase 3 ligand (FLT3LG), mRNA	2323	NM_001459
A:05281	v-fos FBJ murine osteosarcoma viral oncogene homolog (FOS), mRNA	2353	NM_005252
A:01965	FBJ murine osteosarcoma viral oncogene homolog B (FOSB), mRNA	2354	NM_006732
A:01738	fyn-related kinase (FRK), mRNA	2444	NM_002031
A:03614	FK506 binding protein 12-rapamycin associated protein 1 (FRAP1), mRNA	2475	NM_004958
A:08973	ferritin, heavy polypeptide 1 (FTH1), mRNA	2495	NM_002032
A:03646	FYN oncogene related to SRC, FGR, YES (FYN), transcript variant 1, mRNA	2534	NM_002037
B:9714	X-ray repair complementing defective repair in Chinese hamster cells 6 (Ku autoantigen, 70kDa) (XRCC6), mRNA	2547	NM_001469
A:02378	GRB2-associated binding protein 1 (GAB1), transcript variant 2, mRNA	2549	NM_002039
A:07229	cyclin G associated kinase (GAK), mRNA	2580	NM_005255
B:9019	growth arrest-specific 1 (GAS1), mRNA	2619	NM_002048
B:9019	growth arrest-specific 1 (GAS1), mRNA	2620	NM_002048
B:9020	growth arrest-specific 6 (GAS6), mRNA	2621	NM_000820
A:10093	growth arrest-specific 8 (GAS8), mRNA	2622	NM_001481
A:09801	glucagon (GCG), mRNA	2641	NM_002054
A:09968	nuclear receptor subfamily 6, group A, member 1 (NR6A1), transcript variant 3, mRNA	2649	NM_033335

B:4833	growth factor, augmentor of liver regeneration (ERV1 homolog, <i>S. cerevisiae</i>) (GFER), mRNA	2671	NM_005262
A:08908	growth factor independent 1 (GFI1), mRNA	2672	NM_005263
A:02108	GPI anchored molecule like protein (GML), mRNA	2765	NM_002066
A:05004	gonadotropin-releasing hormone 1 (luteinizing-releasing hormone) (GNRH1), mRNA	2796	NM_000825
B:4823	stratifin (SFN), mRNA	2810	NM_006142
B:3553_hk-r1	G protein pathway suppressor 1 (GPS1), transcript variant 1, mRNA	2873	NM_212492
A:04124	G protein pathway suppressor 2 (GPS2), mRNA	2874	NM_004489
A:05918	granulin (GRN), transcript variant 1, mRNA	2896	NM_002087
C:0852	glucocorticoid receptor DNA binding factor 1 GRLF1	2909	NM_004491
A:04681	chemokine (C-X-C motif) ligand 1 (melanoma growth stimulating activity, alpha) (CXCL1), mRNA	2919	NM_001511
A:07763	gastrin-releasing peptide receptor (GRPR), mRNA	2925	NM_005314
B:9294	glycogen synthase kinase 3 beta (GSK3B), mRNA	2932	NM_002093
A:07312	G1 to S phase transition 1 (GSPT1), mRNA	2935	NM_002094
A:09859	mutS homolog 6 (<i>E. coli</i>) (MSH6), mRNA	2956	NM_000179
A:04525	general transcription factor IIH, polypeptide 1 (62kD subunit) (GTF2H1), mRNA	2965	NM_005316
B:9176	hepatoma-derived growth factor (high-mobility group protein 1-like) (HDGF), mRNA	3068	NM_004494
B:8961	hepatocyte growth factor (hepapoietin A; scatter factor) (HGF), transcript variant 3, mRNA	3082	NM_001010932
A:05880	hematopoietically expressed homeobox (HHEX), mRNA	3090	NM_002729
A:05673	hexokinase 2 (HK2), mRNA	3099	NM_000189
A:10377	high-mobility group box 1 (HMGB1), mRNA	3146	NM_002128
A:07252	solute carrier family 29 (nucleoside transporters), member 2 (SLC29A2), mRNA	3177	NM_001532
A:04416	heterogeneous nuclear ribonucleoprotein L (HNRPL), transcript variant 1, mRNA	3191	NM_001533
C:1926	homeo box C10 (HOXC10), mRNA	3226	NM_017409
A:08912	homeo box D13 (HOXD13), mRNA	3239	NM_000523
A:05637	v-Ha-ras Harvey rat sarcoma viral oncogene homolog (HRAS), transcript variant 1, mRNA	3265	NM_005343
A:08143	heat shock 70kDa protein 1A (HSPA1A), mRNA	3304	NM_005345

A:05469	heat shock 70kDa protein 2 (HSPA2), mRNA	3306	NM_021979
A:09246	5-hydroxytryptamine (serotonin) receptor 1A (HTR1A), mRNA	3350	NM_000524
A:07300	HUS1 checkpoint homolog (S. pombe) (HUS1), mRNA	3364	NM_004507
B:7639	interferon, gamma-inducible protein 16 IFI16	3428	NM_005531
A:04388	interferon, beta 1, fibroblast (IFNB1), mRNA	3456	NM_002176
A:02473	interferon, omega 1 (IFNW1), mRNA	3467	NM_002177
B:5220	insulin-like growth factor 1 (somatomedin C) IGF1	3479	NM_000618
C:0361	insulin-like growth factor 1 receptor IGF1R	3480	NM_000875
B:5688	insulin-like growth factor 2 (somatomedin A) (IGF2), mRNA	3481	NM_000612
A:09232	insulin-like growth factor binding protein 4 (IGFBP4), mRNA	3487	NM_001552
A:02232	insulin-like growth factor binding protein 6 (IGFBP6), mRNA	3489	NM_002178
A:03385	insulin-like growth factor binding protein 7 (IGFBP7), mRNA	3490	NM_001553
B:8268	cysteine-rich, angiogenic inducer, 61 CYR61	3491	NM_001554
C:2817	immunoglobulin mu binding protein 2 (IGHMBP2), mRNA	3508	NM_002180
A:07761	interleukin 1, alpha (IL1A), mRNA	3552	NM_000575
A:08500	interleukin 1, beta (IL1B), mRNA	3553	NM_000576
A:02668	interleukin 2 (IL2), mRNA	3558	NM_000586
A:03791	interleukin 2 receptor, alpha (IL2RA), mRNA	3559	NM_000417
B:4721	interleukin 2 receptor, gamma (severe combined immunodeficiency) (IL2RG), mRNA	3561	NM_000206
A:09679	interleukin 3 (colony-stimulating factor, multiple) (IL3), mRNA	3562	NM_000588
A:05115	interleukin 4 (IL4), transcript variant 1, mRNA	3565	NM_000589
A:04767	interleukin 5 (colony-stimulating factor, eosinophil) (IL5), mRNA	3567	NM_000879
A:00154	interleukin 5 receptor, alpha (IL5RA), transcript variant 1, mRNA	3568	NM_000564
A:00705	interleukin 6 (interferon, beta 2) (IL6), mRNA	3569	NM_000600
B:6258	interleukin 6 receptor (IL6R), transcript variant 1, mRNA	3570	NM_000565
A:04305	interleukin 7 (IL7), mRNA	3574	NM_000880
A:06269	interleukin 8 (IL8), mRNA	3576	NM_000584
A:10396	interleukin 9 (IL9), mRNA	3578	NM_000590
B:9037	interleukin 8 receptor, beta (IL8RB), mRNA	3579	NM_001557
A:07447	interleukin 9 receptor (IL9R), transcript variant 1, mRNA	3581	NM_002186
A:07424	interleukin 10 (IL10), mRNA	3586	NM_000572
C:2709	interleukin 11 (IL11), mRNA	3589	NM_000641

A:02631	interleukin 12A (natural killer cell stimulatory factor 1, cytotoxic lymphocyte maturation factor 1, p35) (IL12A), mRNA	3592	NM_000882
A:01248	interleukin 12B (natural killer cell stimulatory factor 2, cytotoxic lymphocyte maturation factor 2, p40) (IL12B), mRNA	3593	NM_002187
A:02885	interleukin 12 receptor, beta 1 (IL12RB1), transcript variant 1, mRNA	3594	NM_005535
B:4956	interleukin 12 receptor, beta 2 (IL12RB2), mRNA	3595	NM_001559
C:2230	interleukin 13 (IL13), mRNA	3596	NM_002188
A:02144	interleukin 13 receptor, alpha 2 (IL13RA2), mRNA	3599	NM_000640
A:05823	interleukin 15 (IL15), transcript variant 3, mRNA	3600	NM_000585
A:05507	interleukin 15 receptor, alpha (IL15RA), transcript variant 1, mRNA	3601	NM_002189
A:09902	tumour necrosis factor receptor superfamily, member 9 (TNFRSF9), mRNA	3604	NM_001561
A:01751	interleukin 18 (interferon-gamma-inducing factor) (IL18), mRNA	3606	NM_001562
B:1174	interleukin enhancer binding factor 3, 90kDa (ILF3), transcript variant 1, mRNA	3609	NM_012218
A:06560	integrin-linked kinase (ILK), transcript variant 1, mRNA	3611	NM_004517
A:04679	inner centromere protein antigens 135/155kDa (INCENP), mRNA	3619	NM_020238
B:8330	inhibitor of growth family, member 1 (ING1), transcript variant 4, mRNA	3621	NM_005537
A:05295	inhibin, alpha (INH A), mRNA	3623	NM_002191
A:02189	inhibin, beta A (activin A, activin AB alpha polypeptide) (INHBA), mRNA	3624	NM_002192
B:4601	chemokine (C-X-C motif) ligand 10 (CXCL10), mRNA	3627	NM_001565
B:3728	insulin induced gene 1 (INSIG1), transcript variant 1, mRNA	3638	NM_005542
A:08018	insulin-like 4 (placenta) (INSL4), mRNA	3641	NM_002195
A:02981	interferon regulatory factor 1 (IRF1), mRNA	3659	NM_002198
A:00655	interferon regulatory factor 2 (IRF2), mRNA	3660	NM_002199
B:4265	interferon stimulated exonuclease gene 20kDa (ISG20), mRNA	3669	NM_002201
C:0395	jagged 2 (JAG2), transcript variant 1, mRNA	3714	NM_002226
A:05470	Janus kinase 2 (a protein tyrosine kinase) (JAK2), mRNA	3717	NM_004972
A:04848	v-jun sarcoma virus 17 oncogene homolog (avian) (JUN), mRNA	3725	NM_002228
A:08730	jun B proto-oncogene (JUNB), mRNA	3726	NM_002229

A:06684	kinesin family member 11 (KIF11), mRNA	3832	NM_004523
B:4887	kinesin family member C1 (KIFC1), mRNA	3833	NM_002263
A:02390	kinesin family member 22 (KIF22), mRNA	3835	NM_007317
B:4036	karyopherin alpha 2 (RAG cohort 1, importin alpha 1) (KPNA2), mRNA	3838	NM_002266
B:8230	v-Ki-ras2 Kirsten rat sarcoma viral oncogene homolog (KRAS), transcript variant b, mRNA	3845	NM_004985
A:08264	keratin 16 (focal non-epidermolytic palmoplantar keratoderma) (KRT16), mRNA	3868	NM_005557
B:6112	lymphocyte-specific protein tyrosine kinase (LCK), mRNA	3932	NM_005356
A:02572	leukaemia inhibitory factor (cholinergic differentiation factor) (LIF), mRNA	3976	NM_002309
A:02207	ligase I, DNA, ATP-dependent (LIG1), mRNA	3978	NM_000234
A:08891	ligase III, DNA, ATP-dependent (LIG3), nuclear gene encoding mitochondrial protein, transcript variant alpha, mRNA	3980	NM_013975
A:05297	ligase IV, DNA, ATP-dependent (LIG4), mRNA	3981	NM_206937
B:8631	LIM domain only 1 (rhombotin 1) (LMO1), mRNA	4004	NM_002315
A:00504	LIM domain containing preferred translocation partner in lipoma (LPP), mRNA	4029	NM_005578
A:00504	LIM domain containing preferred translocation partner in lipoma (LPP), mRNA	4030	NM_005578
B:0707	low density lipoprotein-related protein 1 (alpha-2-macroglobulin receptor) (LRP1), mRNA	4035	NM_002332
A:09461	low density lipoprotein receptor-related protein 5 (LRP5), mRNA	4041	NM_002335
A:03776	low density lipoprotein receptor-related protein associated protein 1 (LRPAP1), mRNA	4043	NM_002337
B:7687	latent transforming growth factor beta binding protein 2 (LTBP2), mRNA	4053	NM_000428
C:2653	v-yes-1 Yamaguchi sarcoma viral related oncogene homolog (LYN), mRNA	4067	NM_002350
A:10613	tumour-associated calcium signal transducer 2 (TACSTD2), mRNA	4070	NM_002353
A:03716	MAX dimerization protein 1 (MXD1), mRNA	4084	NM_002357
A:06387	MAD2 mitotic arrest deficient-like 1 (yeast) (MAD2L1), mRNA	4085	NM_002358
B:5699	v-maf musculoaponeurotic fibrosarcoma oncogene homolog G (avian) (MAFG), transcript variant 1,	4097	NM_002359

	mRNA		
A:03848	MAS1 oncogene (MAS1), mRNA	4142	NM_002377
B:9275	megakaryocyte-associated tyrosine kinase (MATK), transcript variant 1, mRNA	4145	NM_139355
B:4426	mutated in colorectal cancers (MCC), mRNA	4163	NM_002387
A:08834	MCM2 minichromosome maintenance deficient 2, mitotin (<i>S. cerevisiae</i>) (MCM2), mRNA	4171	NM_004526
A:08668	MCM3 minichromosome maintenance deficient 3 (<i>S. cerevisiae</i>) (MCM3), mRNA	4172	NM_002388
B:7581	MCM4 minichromosome maintenance deficient 4 (<i>S. cerevisiae</i>) (MCM4), transcript variant 1, mRNA	4173	NM_005914
B:7805	MCM5 minichromosome maintenance deficient 5, cell division cycle 46 (<i>S. cerevisiae</i>) (MCM5), mRNA	4174	NM_006739
B:8147	MCM6 minichromosome maintenance deficient 6 (MIS5 homolog, <i>S. pombe</i>) (<i>S. cerevisiae</i>) (MCM6), mRNA	4175	NM_005915
B:7620	MCM7 minichromosome maintenance deficient 7 (<i>S. cerevisiae</i>) MCM7	4176	NM_005916
B:4650	midkine (neurite growth-promoting factor 2) (MDK), transcript variant 1, mRNA	4192	NM_001012334
B:8649	Mdm2, transformed 3T3 cell double minute 2, p53 binding protein (mouse) (MDM2), transcript variant MDM2a, mRNA	4193	NM_006878
A:03964	Mdm4, transformed 3T3 cell double minute 4, p53 binding protein (mouse) (MDM4), mRNA	4194	NM_002393
A:10600	RAB8A, member RAS oncogene family (RAB8A), mRNA	4218	NM_005370
B:8222	met proto-oncogene (hepatocyte growth factor receptor) MET	4233	NM_000245
A:09470	KIT ligand (KITLG), transcript variant b, mRNA	4254	NM_000899
A:01575	O-6-methylguanine-DNA methyltransferase (MGMT), mRNA	4255	NM_002412
A:10388	antigen identified by monoclonal antibody Ki-67 (MKI67), mRNA	4288	NM_002417
A:06073	mutL homolog 1, colon cancer, nonpolyposis type 2 (<i>E. coli</i>) (MLH1), mRNA	4292	NM_000249
B:7492	myeloid/lymphoid or mixed-lineage leukaemia (trithorax homolog, <i>Drosophila</i>); translocated to, 7 (MLLT7), mRNA	4303	NM_005938
A:09644	meningioma (disrupted in balanced translocation) 1 (MN1), mRNA	4330	NM_002430
A:08968	menage a trois 1 (CAK assembly factor) (MNAT1), mRNA	4331	NM_002431

A:02100	MAX binding protein (MNT), mRNA	4335	NM_020310
A:02282	v-mos Moloney murine sarcoma viral oncogene homolog (MOS), mRNA	4342	NM_005372
A:06141	myeloproliferative leukaemia virus oncogene (MPL), mRNA	4352	NM_005373
A:04072	MRE11 meiotic recombination 11 homolog A (S. cerevisiae) (MRE11A), transcript variant 1, mRNA	4361	NM_005591
A:04072	MRE11 meiotic recombination 11 homolog A (S. cerevisiae) (MRE11A), transcript variant 1, mRNA	4362	NM_005591
A:04514	mutS homolog 2, colon cancer, nonpolyposis type 1 (E. coli) (MSH2), mRNA	4436	NM_000251
A:06785	mutS homolog 3 (E. coli) (MSH3), mRNA	4437	NM_002439
A:02756	mutS homolog 4 (E. coli) (MSH4), mRNA	4438	NM_002440
A:09339	mutS homolog 5 (E. coli) (MSH5), transcript variant 1, mRNA	4439	NM_025259
A:04591	macrophage stimulating 1 receptor (c-met-related tyrosine kinase) (MST1R), mRNA	4486	NM_002447
A:05992	metallothionein 3 (growth inhibitory factor (neurotrophic)) (MT3), mRNA	4504	NM_005954
C:2393	mature T-cell proliferation 1 (MTCP1), nuclear gene encoding mitochondrial protein, transcript variant B1, mRNA	4515	NM_014221
A:01898	mutY homolog (E. coli) (MUTYH), mRNA	4595	NM_012222
A:10478	MAX interactor 1 (MXI1), transcript variant 1, mRNA	4601	NM_005962
B:5181	v-myb myeloblastosis viral oncogene homolog (avian) MYB	4602	NM_005375
B:5429	v-myb myeloblastosis viral oncogene homolog (avian)-like 1 (MYBL1), mRNA	4603	XM_034274, XM_933460, XM_938064
A:06037	v-myb myeloblastosis viral oncogene homolog (avian)-like 2 (MYBL2), mRNA	4605	NM_002466
A:02498	v-myc myelocytomatosis viral oncogene homolog (avian) (MYC), mRNA	4609	NM_002467
C:2723	myosin, heavy polypeptide 10, non-muscle (MYH10), mRNA	4628	NM_005964
B:4239	NGFI-A binding protein 2 (EGR1 binding protein 2) (NAB2), mRNA	4665	NM_005967
B:1584	nucleosome assembly protein 1-like 1 (NAP1L1), transcript variant 1, mRNA	4673	NM_139207
A:09960	neuroblastoma, suppression of tumourigenicity 1 (NBL1), transcript variant 1, mRNA	4681	NM_182744
A:02361	nucleotide binding protein 1 (MinD homolog, E. coli) (NUBP1), mRNA	4682	NM_002484

A:10519	nibrin (NBN), transcript variant 1, mRNA	4683	NM_002485
A:08868	NCK adaptor protein 1 (NCK1), mRNA	4690	NM_006153
A:07320	necdin homolog (mouse) (NDN), mRNA	4692	NM_002487
B:5481	Norrie disease (pseudoglioma) (NDP), mRNA	4693	NM_000266
B:4761	septin 2 (SEPT2), transcript variant 4, mRNA	4735	NM_004404
A:04128	neural precursor cell expressed, developmentally down-regulated 9 (NEDD9), transcript variant 1, mRNA	4739	NM_006403
B:7542	NIMA (never in mitosis gene a)-related kinase 1 (NEK1), mRNA	4750	NM_012224
A:00847	NIMA (never in mitosis gene a)-related kinase 2 (NEK2), mRNA	4751	NM_002497
B:7555	NIMA (never in mitosis gene a)-related kinase 3 (NEK3), transcript variant 1, mRNA	4752	NM_002498
B:9751	neurofibromin 1 (neurofibromatosis, von Recklinghausen disease, Watson disease) (NF1), mRNA	4763	NM_000267
B:7527	neurofibromin 2 (bilateral acoustic neuroma) (NF2), transcript variant 12, mRNA	4771	NM_181825
B:8431	nuclear factor I/A (NFIA), mRNA	4774	NM_005595
A:03729	nuclear factor I/B (NFIB), mRNA	4781	NM_005596
B:5428	nuclear factor I/C (CCAAT-binding transcription factor) (NFIC), transcript variant 1, mRNA	4782	NM_005597
C:5826	nuclear factor I/X (CCAAT-binding transcription factor) (NFIX), mRNA	4784	NM_002501
B:5078	nuclear transcription factor Y, gamma NFYC	4802	NM_014223
A:05462	NHP2 non-histone chromosome protein 2-like 1 (S. cerevisiae) (NHP2L1), transcript variant 1, mRNA	4809	NM_005008
A:01677	non-metastatic cells 1, protein (NM23A) expressed in (NME1), transcript variant 2, mRNA	4830	NM_000269
A:04306	non-metastatic cells 2, protein (NM23B) expressed in (NME2), transcript variant 1, mRNA	4831	NM_002512
C:1522	nucleolar protein 1, 120kDa (NOL1), transcript variant 2, mRNA	4839	NM_001033714
A:06565	neuropeptide Y (NPY), mRNA	4852	NM_000905
A:00579	Notch homolog 2 (Drosophila) (NOTCH2), mRNA	4853	NM_024408
A:02787	neuroblastoma RAS viral (v-ras) oncogene homolog (NRAS), mRNA	4893	NM_002524
B:6139	nuclear mitotic apparatus protein 1 (NUMA1), mRNA	4926	NM_006185
A:04432	opioid receptor, mu 1 (OPRM1), transcript variant MOR-1, mRNA	4988	NM_000914
A:02654	origin recognition complex, subunit 1-like (yeast) (ORC1L), mRNA	4998	NM_004153

A:01697	origin recognition complex, subunit 2-like (yeast) (ORC2L), mRNA	4999	NM_006190
A:06724	origin recognition complex, subunit 4-like (yeast) (ORC4L), transcript variant 2, mRNA	5000	NM_002552
C:0244	origin recognition complex, subunit 5-like (yeast) (ORC5L), transcript variant 2, mRNA	5001	NM_181747
A:09399	oncostatin M (OSM), mRNA	5008	NM_020530
A:07058	proliferation-associated 2G4, 38kDa (PA2G4), mRNA	5036	NM_006191
A:04710	platelet-activating factor acetylhydrolase, isoform Ib, alpha subunit 45kDa (PAFAH1B1), mRNA	5048	NM_000430
A:03397	peroxiredoxin 1 (PRDX1), transcript variant 1, mRNA	5052	NM_002574
B:4727	regenerating islet-derived 3 alpha (REG3A), transcript variant 1, mRNA	5068	NM_002580
A:03215	PRKC, apoptosis, WT1, regulator (PAWR), mRNA	5074	NM_002583
A:03715	proliferating cell nuclear antigen (PCNA), transcript variant 1, mRNA	5111	NM_002592
A:09486	PCTAIRE protein kinase 1 (PCTK1), transcript variant 1, mRNA	5127	NM_006201
A:09486	PCTAIRE protein kinase 1 (PCTK1), transcript variant 1, mRNA	5128	NM_006201
C:2666	platelet-derived growth factor alpha polypeptide (PDGFA), transcript variant 1, mRNA	5154	NM_002607
B:7519	platelet-derived growth factor beta polypeptide (simian sarcoma viral (v-sis) oncogene homolog) (PDGFB), transcript variant 1, mRNA	5155	NM_002608
A:02349	platelet-derived growth factor receptor, alpha polypeptide (PDGFRA), mRNA	5156	NM_006206
A:00876	PDZ domain containing 1 (PDZK1), mRNA	5174	NM_002614
A:04139	serpin peptidase inhibitor, clade F (alpha-2 antiplasmin, pigment epithelium derived factor), member 1 (SERPINF1), transcript variant 4, mRNA	5176	NM_002615
B:4669	prefoldin 1 (PFDN1), mRNA	5201	NM_002622
A:00156	placental growth factor, vascular endothelial growth factor-related protein (PGF), mRNA	5228	NM_002632
B:9242	phosphoinositide-3-kinase, catalytic, beta polypeptide (PIK3CB), mRNA	5291	NM_006219
A:09957	protein (peptidyl-prolyl cis/trans isomerase) NIMA-interacting 1 (PIN1), mRNA	5300	NM_006221
A:00888	pleiomorphic adenoma gene-like 1 (PLAGL1), transcript variant 2,	5325	NM_006718

	mRNA		
A:08398	plasminogen (PLG), mRNA	5340	NM_000301
B:3744	polo-like kinase 1 (<i>Drosophila</i>) (PLK1), mRNA	5347	NM_005030
B:4722	peripheral myelin protein 22 (PMP22), transcript variant 1, mRNA	5376	NM_000304
A:10286	PMS1 postmeiotic segregation increased 1 (<i>S. cerevisiae</i>) (PMS1), mRNA	5378	NM_000534
A:10286	PMS1 postmeiotic segregation increased 1 (<i>S. cerevisiae</i>) (PMS1), mRNA	5379	NM_000534
B:9336	postmeiotic segregation increased 2-like 2 (PMS2L2), mRNA	5380	NM_002679
B:9336	postmeiotic segregation increased 2-like 2 (PMS2L2), mRNA	5382	NM_002679
A:10467	postmeiotic segregation increased 2-like 5 (PMS2L5), mRNA	5383	NM_174930
A:10467	postmeiotic segregation increased 2-like 5 (PMS2L5), mRNA	5386	NM_174930
A:02096	PMS2 postmeiotic segregation increased 2 (<i>S. cerevisiae</i>) (PMS2), transcript variant 1, mRNA	5395	NM_000535
B:0731	septin 5 (SEPT5), transcript variant 1, mRNA	5413	NM_002688
A:09062	septin 4 (SEPT4), transcript variant 1, mRNA	5414	NM_004574
A:05543	polymerase (DNA directed), alpha (POLA), mRNA	5422	NM_016937
A:02852	polymerase (DNA directed), beta (POLB), mRNA	5423	NM_002690
A:09477	polymerase (DNA directed), delta 1, catalytic subunit 125kDa (POLD1), mRNA	5424	NM_002691
A:02929	polymerase (DNA directed), delta 2, regulatory subunit 50kDa (POLD2), mRNA	5425	NM_006230
B:3196	polymerase (DNA directed), epsilon POLE	5426	NM_006231
A:04680	polymerase (DNA directed), epsilon 2 (p59 subunit) (POLE2), mRNA	5427	NM_002692
A:08572	polymerase (DNA directed), gamma (POLG), mRNA	5428	NM_002693
A:08948	polymerase (RNA) mitochondrial (DNA directed) (POLRMT), nuclear gene encoding mitochondrial protein, mRNA	5442	NM_005035
A:00480	POU domain, class 1, transcription factor 1 (Pit1, growth hormone factor 1) (POU1F1), mRNA	5449	NM_000306
C:6960	peroxisome proliferative activated receptor, delta (PPARD), transcript variant 1, mRNA	5467	NM_006238
B:0695	PPAR binding protein (PPARBP), mRNA	5469	NM_004774
A:10622	pro-platelet basic protein (chemokine (C-X-C motif) ligand 7) (PPBP), mRNA	5473	NM_002704

A:08431	protein phosphatase 1G (formerly 2C), magnesium-dependent, gamma isoform (PPM1G), transcript variant 1, mRNA	5496	NM_177983
A:05348	protein phosphatase 1, catalytic subunit, alpha isoform (PPP1CA), transcript variant 1, mRNA	5499	NM_002708
B:0943	protein phosphatase 1, catalytic subunit, beta isoform (PPP1CB), transcript variant 1, mRNA	5500	NM_002709
A:02064	protein phosphatase 1, catalytic subunit, gamma isoform (PPP1CC), mRNA	5501	NM_002710
A:01231	protein phosphatase 2 (formerly 2A), catalytic subunit, alpha isoform (PPP2CA), mRNA	5515	NM_002715
A:03825	protein phosphatase 2 (formerly 2A), regulatory subunit A (PR 65), alpha isoform (PPP2R1A), mRNA	5518	NM_014225
A:01064	protein phosphatase 2 (formerly 2A), regulatory subunit A (PR 65), beta isoform (PPP2R1B), transcript variant 1, mRNA	5519	NM_002716
A:00874	protein phosphatase 2 (formerly 2A), regulatory subunit B", alpha (PPP2R3A), transcript variant 1, mRNA	5523	NM_002718
A:07683	protein phosphatase 3 (formerly 2B), catalytic subunit, beta isoform (calcineurin A beta) (PPP3CB), mRNA	5532	NM_021132
A:00032	protein phosphatase 5, catalytic subunit (PPP5C), mRNA	5536	NM_006247
A:02880	protein phosphatase 6, catalytic subunit (PPP6C), mRNA	5537	NM_002721
A:07833	primase, polypeptide 1, 49kDa (PRIM1), mRNA	5557	NM_000946
A:08706	primase, polypeptide 2A, 58kDa PRIM2A	5558	NM_000947
A:00953	protein kinase, cAMP-dependent, regulatory, type I, alpha (tissue specific extinguisher 1) (PRKAR1A), transcript variant 1, mRNA	5573	NM_002734
A:07305	protein kinase, cAMP-dependent, regulatory, type II, beta (PRKAR2B), mRNA	5578	NM_002736
A:08970	protein kinase D1 (PRKD1), mRNA	5587	NM_002742
A:05228	protein kinase, cGMP-dependent, type II (PRKG2), mRNA	5593	NM_006259
B:6263	mitogen-activated protein kinase 1 (MAPK1), transcript variant 1, mRNA	5594	NM_002745
B:5471	mitogen-activated protein kinase 3 (MAPK3), mRNA	5595	NM_002746
B:9088	mitogen-activated protein kinase 4 (MAPK4), mRNA	5596	NM_002747
A:03644	mitogen-activated protein kinase 6 (MAPK6), mRNA	5597	NM_002748

A:09951	mitogen-activated protein kinase 7 (MAPK7), transcript variant 1, mRNA	5598	NM_139033
A:00932	mitogen-activated protein kinase 13 (MAPK13), mRNA	5603	NM_002754
A:06747	mitogen-activated protein kinase 6 (MAP2K6), transcript variant 1, mRNA	5608	NM_002758
B:4014	mitogen-activated protein kinase 7 MAP2K7	5609	NM_145185
B:1372	eukaryotic translation initiation factor 2-alpha kinase 2 (EIF2AK2), mRNA	5610	NM_002759
B:5991	protein-kinase, interferon-inducible double stranded RNA dependent inhibitor, repressor of (P58 repressor) (PRKRIR), mRNA	5612	NM_004705
A:03959	prolactin (PRL), mRNA	5617	NM_000948
A:09385	protamine 1 (PRM1), mRNA	5619	NM_002761
A:02848	protamine 2 (PRM2), mRNA	5620	NM_002762
A:07907	kallikrein 10 (KLK10), transcript variant 1, mRNA	5655	NM_002776
A:01338	proteinase 3 (serine proteinase, neutrophil, Wegener granulomatosis autoantigen) (PRTN3), mRNA	5657	NM_002777
B:4949	presenilin 1 (Alzheimer disease 3) PSEN1	5663	NM_000021
A:00037	presenilin 2 (Alzheimer disease 4) (PSEN2), transcript variant 1, mRNA	5664	NM_000447
A:05430	peptide YY (PYY), mRNA	5697	NM_004160
A:05083	proteasome (prosome, macropain) 26S subunit, non-ATPase, 8 (PSMD8), mRNA	5714	NM_002812
A:10847	patched homolog (Drosophila) (PTCH), mRNA	5727	NM_000264
A:04029	phosphatase and tensin homolog (mutated in multiple advanced cancers 1) (PTEN), mRNA	5728	NM_000314
A:08708	parathyroid hormone-like hormone (PTH LH), transcript variant 2, mRNA	5744	NM_002820
B:4775	prothymosin, alpha (gene sequence 28) (PTMA), mRNA	5757	NM_002823
A:05250	parathymosin (PTMS), mRNA	5763	NM_002824
C:2316	pleiotrophin (heparin binding growth factor 8, neurite growth-promoting factor 1) (PTN), mRNA	5764	NM_002825
C:2627	quiescin Q6 (QSCN6), transcript variant 1, mRNA	5768	NM_002826
A:10310	protein tyrosine phosphatase, non-receptor type 6 (PTPN6), transcript variant 2, mRNA	5777	NM_080548
A:02619	RAD1 homolog (S. pombe) (RAD1), transcript variant 1, mRNA	5810	NM_002853
C:2196	purine-rich element binding protein A (PURA), mRNA	5813	NM_005859

B:1151	ras-related C3 botulinum toxin substrate 1 (rho family, small GTP binding protein Rac1) (RAC1), transcript variant Rac1b, mRNA	5879	NM_018890
A:05292	RAD9 homolog A (S. pombe) (RAD9A), mRNA	5883	NM_004584
A:10635	RAD17 homolog (S. pombe) (RAD17), transcript variant 8, mRNA	5884	NM_002873
A:07580	RAD21 homolog (S. pombe) (RAD21), mRNA	5885	NM_006265
A:07819	RAD51 homolog (RecA homolog, E. coli) (S. cerevisiae) (RAD51), transcript variant 1, mRNA	5888	NM_002875
A:09744	RAD51-like 1 (S. cerevisiae) (RAD51L1), transcript variant 1, mRNA	5890	NM_002877
B:0346	RAD51-like 3 (S. cerevisiae) RAD51L3	5892	NM_002878, NM_133629
B:1043	RAD52 homolog (S. cerevisiae) (RAD52), transcript variant beta, mRNA	5893	NM_134424
C:2457	v-raf-1 murine leukaemia viral oncogene homolog 1 (RAF1), mRNA	5894	NM_002880
B:8341	ral guanine nucleotide dissociation stimulator RALGDS	5900	NM_001042368, NM_006266
A:09169	RAN, member RAS oncogene family (RAN), mRNA	5901	NM_006325
C:0082	RAP1A, member of RAS oncogene family RAP1A	5906	NM_001010935, NM_002884
A:00423	RAP1B, member of RAS oncogene family (RAP1B), transcript variant 1, mRNA	5908	NM_015646
A:09690	retinoic acid receptor responder (tazarotene induced) 1 (RARRES1), transcript variant 2, mRNA	5918	NM_002888
A:08045	retinoic acid receptor responder (tazarotene induced) 3 (RARRES3), mRNA	5920	NM_004585
B:9011	retinoblastoma 1 (including osteosarcoma) (RB1), mRNA	5925	NM_000321
A:04888	retinoblastoma binding protein 4 (RBBP4), mRNA	5928	NM_005610
C:2267	retinoblastoma binding protein 6 (RBBP6), transcript variant 1, mRNA	5930	NM_006910
A:06741	retinoblastoma binding protein 7 (RBBP7), mRNA	5931	NM_002893
A:09145	retinoblastoma binding protein 8 (RBBP8), transcript variant 1, mRNA	5932	NM_002894
A:10222	retinoblastoma-like 1 (p107) (RBL1), transcript variant 1, mRNA	5933	NM_002895
A:08246	retinoblastoma-like 2 (p130) (RBL2), mRNA	5934	NM_005611
B:9795	RNA binding motif, single stranded interacting protein 1 (RBMS1),	5937	NM_016836

	transcript variant 1, mRNA		
B:1393	regenerating islet-derived 1 alpha (pancreatic stone protein, pancreatic thread protein) (REG1A), mRNA	5967	NM_002909
B:4741	regenerating islet-derived 1 beta (pancreatic stone protein, pancreatic thread protein) (REG1B), mRNA	5968	NM_006507
B:4741	regenerating islet-derived 1 beta (pancreatic stone protein, pancreatic thread protein) (REG1B), mRNA	5969	NM_006507
A:04164	REV3-like, catalytic subunit of DNA polymerase zeta (yeast) (REV3L), mRNA	5980	NM_002912
A:03348	replication factor C (activator 1) 1, 145kDa (RFC1), mRNA	5981	NM_002913
A:06693	replication factor C (activator 1) 2, 40kDa (RFC2), transcript variant 1, mRNA	5982	NM_181471
A:02491	replication factor C (activator 1) 3, 38kDa (RFC3), transcript variant 1, mRNA	5983	NM_002915
A:09921	replication factor C (activator 1) 4, 37kDa (RFC4), transcript variant 1, mRNA	5984	NM_002916
B:3726	replication factor C (activator 1) 5, 36kDa (RFC5), transcript variant 1, mRNA	5985	NM_007370
A:04896	ret finger protein (RFP), transcript variant alpha, mRNA	5987	NM_006510
A:04971	regulator of G-protein signalling 2, 24kDa (RGS2), mRNA	5997	NM_002923
B:8684	relaxin 2 (RLN2), transcript variant 2, mRNA	6024	NM_005059
A:10597	replication protein A1, 70kDa (RPA1), mRNA	6117	NM_002945
A:09203	replication protein A2, 32kDa (RPA2), mRNA	6118	NM_002946
A:00231	replication protein A3, 14kDa (RPA3), mRNA	6119	NM_002947
B:8856	ribosomal protein S4, X-linked (RPS4X), mRNA	6191	NM_001007
B:8856	ribosomal protein S4, X-linked (RPS4X), mRNA	6192	NM_001007
A:10444	ribosomal protein S6 kinase, 70kDa, polypeptide 2 (RPS6KB2), transcript variant 1, mRNA	6199	NM_003952
A:02188	ribosomal protein S25 (RPS25), mRNA	6232	NM_001028
A:08509	related RAS viral (r-ras) oncogene homolog (RRAS), mRNA	6237	NM_006270
A:09802	ribonucleotide reductase M1 polypeptide (RRM1), mRNA	6240	NM_001033
B:3501	ribonucleotide reductase M2 polypeptide (RRM2), mRNA	6241	NM_001034
A:08332	S100 calcium binding protein A5 (S100A5), mRNA	6276	NM_002962

C:1129	S100 calcium binding protein A6 (calcyclin) (S100A6), mRNA	6277	NM_014624
B:3690	S100 calcium binding protein A11 (calgizzarin) (S100A11), mRNA	6282	NM_005620
A:08910	S100 calcium binding protein, beta (neural) (S100B), mRNA	6285	NM_006272
A:05458	mitogen-activated protein kinase 12 (MAPK12), mRNA	6300	NM_002969
A:07786	tetraspanin 31 (TSPAN31), mRNA	6302	NM_005981
A:09884	C-type lectin domain family 11, member A (CLEC11A), mRNA	6320	NM_002975
A:00985	chemokine (C-C motif) ligand 3 (CCL3), mRNA	6348	NM_002983
A:00985	chemokine (C-C motif) ligand 3 (CCL3), mRNA	6349	NM_002983
B:0899	chemokine (C-C motif) ligand 14 (CCL14), transcript variant 2, mRNA	6358	NM_032962
B:0898	chemokine (C-C motif) ligand 23 (CCL23), transcript variant CKbeta8, mRNA	6368	NM_145898
B:5275	chemokine (C-X-C motif) ligand 11 (CXCL11), mRNA	6374	NM_005409
C:2038	SET translocation (myeloid leukaemia-associated) (SET), mRNA	6418	NM_003011
A:00679	SHC (Src homology 2 domain containing) transforming protein 1 (SHC1), transcript variant 1, mRNA	6464	NM_183001
B:9295	SCL/TAL1 interrupting locus (STIL), mRNA	6491	NM_003035
B:7410	signal-induced proliferation-associated gene 1 (SIPA1), transcript variant 1, mRNA	6494	NM_1532538
C:5435	S-phase kinase-associated protein 2 (p45) (SKP2), transcript variant 1, mRNA	6502	NM_005983
A:09017	signaling lymphocytic activation molecule family member 1 (SLAMF1), mRNA	6504	NM_003037
A:06456	solute carrier family 12 (potassium/chloride transporters), member 4 (SLC12A4), mRNA	6560	NM_005072
A:05730	SWI/SNF related, matrix associated, actin dependent regulator of chromatin, subfamily b, member 1 (SMARCB1), transcript variant 1, mRNA	6598	NM_003073
A:07314	fascin homolog 1, actin-bundling protein (Strongylocentrotus purpuratus) (FSCN1), mRNA	6624	NM_003088
A:04540	sparc/osteonectin, cwcv and kazal-like domains proteoglycan (testican) 1 (SPOCK1), mRNA	6695	NM_004598
A:09441	secreted phosphoprotein 1 (osteopontin, bone sialoprotein I, early T-lymphocyte activation 1) (SPP1), mRNA	6696	NM_000582

A:02264	v-src sarcoma (Schmidt-Ruppin A-2) viral oncogene homolog (avian) (SRC), transcript variant 1, mRNA	6714	NM_005417
A:04127	single-stranded DNA binding protein 1 (SSBP1), mRNA	6742	NM_003143
A:07245	signal sequence receptor, alpha (translocon-associated protein alpha) (SSR1), mRNA	6745	NM_003144
A:08350	somatostatin (SST), mRNA	6750	NM_001048
A:03956	somatostatin receptor 1 (SSTR1), mRNA	6751	NM_001049
C:1740	somatostatin receptor 2 (SSTR2), mRNA	6752	NM_001050
A:04237	somatostatin receptor 3 (SSTR3), mRNA	6753	NM_001051
A:04852	somatostatin receptor 4 (SSTR4), mRNA	6754	NM_001052
A:01484	somatostatin receptor 5 (SSTR5), mRNA	6755	NM_001053
A:03398	signal transducer and activator of transcription 1, 91kDa (STAT1), transcript variant alpha, mRNA	6772	NM_007315
A:05843	stromal interaction molecule 1 (STIM1), mRNA	6786	NM_003156
A:04562	NIMA (never in mitosis gene a)-related kinase 4 (NEK4), mRNA	6787	NM_003157
A:04814	serine/threonine kinase 6 (STK6), transcript variant 1, mRNA	6790	NM_198433
A:01764	aurora kinase C (AURKC), transcript variant 3, mRNA	6795	NM_003160
A:10309	suppressor of variegation 3-9 homolog 1 (Drosophila) (SUV39H1), mRNA	6839	NM_003173
A:01895	synaptonemal complex protein 1 (SYCP1), mRNA	6847	NM_003176
A:09854	spleen tyrosine kinase (SYK), mRNA	6850	NM_003177
A:02589	transcriptional adaptor 2 (ADA2 homolog; yeast)-like (TADA2L), transcript variant 1, mRNA	6871	NM_001488
A:01355	TAF1 RNA polymerase II, TATA box binding protein (TBP)-associated factor, 250kDa (TAF1), transcript variant 1, mRNA	6872	NM_004606
C:1960	T-cell acute lymphocytic leukaemia 1 (TAL1), mRNA	6886	NM_003189
C:2789	transcription factor 3 (E2A immunoglobulin enhancer binding factors E12/E47) (TCF3), mRNA	6930	NM_003200
B:4738	transcription factor 8 (represses interleukin 2 expression) (TCF8), mRNA	6935	NM_030751
A:03967	transcription factor 19 (SC1) (TCF19), mRNA	6941	NM_007109
A:05964	telomerase-associated protein 1 (TEP1), mRNA	7011	NM_007110
B:9167	telomeric repeat binding factor (NIMA-interacting) 1 (TERF1),	7013	NM_003218

	transcript variant 2, mRNA		
B:7401	telomeric repeat binding factor 2 (TERF2), mRNA	7014	NM_005652
C:0355	telomerase reverse transcriptase (TERT), transcript variant 1, mRNA	7015	NM_003219
A:07625	transcription factor A, mitochondrial (TFAM), mRNA	7019	NM_003201
A:06784	nuclear receptor subfamily 2, group F, member 1 (NR2F1), mRNA	7025	NM_005654
A:06784	nuclear receptor subfamily 2, group F, member 1 (NR2F1), mRNA	7027	NM_005654
B:5016	transcription factor Dp-2 (E2F dimerization partner 2) (TFDP2), mRNA	7029	NM_006286
B:5851	transforming growth factor, alpha (TGFA), mRNA	7039	NM_003236
A:07050	transforming growth factor, beta 1 (Camurati-Engelmann disease) (TGFB1), mRNA	7040	NM_000660
B:0094	transforming growth factor beta 1 induced transcript 1 (TGFB1I1), mRNA	7041	NM_015927
A:09824	transforming growth factor, beta 2 (TGFB2), mRNA	7042	NM_003238
B:7853	transforming growth factor, beta 3 (TGFB3), mRNA	7043	NM_003239
B:4156	transforming growth factor, beta-induced, 68kDa (TGFB1), mRNA	7045	NM_000358
A:03732	transforming growth factor, beta receptor II (70/80kDa) (TGFB2), transcript variant 2, mRNA	7048	NM_003242
B:0258	thrombopoietin (myeloproliferative leukaemia virus oncogene ligand, megakaryocyte growth and development factor) (THPO), transcript variant 3, mRNA	7066	NM_199356
B:4371	thyroid hormone receptor, alpha (erythroblastic leukaemia viral (v-erb-a) oncogene homolog, avian) (THRA), transcript variant 1, mRNA	7067	NM_199334
A:06139	Kruppel-like factor 10 (KLF10), transcript variant 1, mRNA	7071	NM_005655
A:08048	TIMP metalloproteinase inhibitor 1 (TIMP1), mRNA	7076	NM_003254
B:3686	transmembrane 4 L six family member 4 (TM4SF4), mRNA	7104	NM_004617
B:5451	topoisomerase (DNA) I (TOP1), mRNA	7150	NM_003286
B:7145	topoisomerase (DNA) II alpha 170kDa (TOP2A), mRNA	7153	NM_001067
A:04487	topoisomerase (DNA) II beta 180kDa (TOP2B), mRNA	7155	NM_001068
A:05345	topoisomerase (DNA) III alpha (TOP3A), mRNA	7156	NM_004618
A:07597	tumour protein p53 (Li-Fraumeni syndrome) (TP53), mRNA	7157	NM_000546
B:6951	tumour protein p53 binding protein, 2 (TP53BP2), transcript variant 1, mRNA	7159	NM_001031685

A:10089	tumour protein p73 (TP73), mRNA	7161	NM_005427
A:07179	tumour protein D52-like 1 (TPD52L1), transcript variant 4, mRNA	7165	NM_001003397
A:00700	tuberous sclerosis 1 (TSC1), transcript variant 1, mRNA	7248	NM_000368
C:2440	tuberous sclerosis 2 (TSC2), transcript variant 2, mRNA	7249	NM_021055
A:06571	thyroid stimulating hormone receptor (TSHR), transcript variant 1, mRNA	7253	NM_000369
A:02759	testis specific protein, Y-linked 1 (TSPY1), mRNA	7258	NM_003308
A:09121	tumour suppressing subtransferable candidate 1 (TSSC1), mRNA	7260	NM_003310
A:07936	TTK protein kinase (TTK), mRNA	7272	NM_003318
A:05365	tumour necrosis factor (ligand) superfamily, member 4 (tax-transcriptionally activated glycoprotein 1, 34kDa) (TNFSF4), mRNA	7292	NM_003326
B:0763	thioredoxin TXN	7295	NM_003329
B:4917	ubiquitin-activating enzyme E1 (A1S9T and BN75 temperature sensitivity complementing) (UBE1), transcript variant 1, mRNA	7317	NM_003334
A:08169	ubiquitin-conjugating enzyme E2D 1 (UBC4/5 homolog, yeast) (UBE2D1), mRNA	7321	NM_003338
A:07196	ubiquitin-conjugating enzyme E2D 3 (UBC4/5 homolog, yeast) (UBE2D3), transcript variant 1, mRNA	7323	NM_003340
A:04972	ubiquitin-conjugating enzyme E2 variant 1 (UBE2V1), transcript variant 1, mRNA	7335	NM_021988
B:0648	ubiquitin-conjugating enzyme E2 variant 2 (UBE2V2), mRNA	7336	NM_003350
C:2659	uromodulin (uromucoid, Tamm-Horsfall glycoprotein) (UMOD), transcript variant 2, mRNA	7369	NM_001008389
A:06855	vav 1 oncogene (VAV1), mRNA	7409	NM_005428
A:08040	vav 2 oncogene VAV2	7410	NM_003371
C:1128	vascular endothelial growth factor (VEGF), transcript variant 5, mRNA	7422	NM_001025369
B:5229	vascular endothelial growth factor B (VEGFB), mRNA	7423	NM_003377
A:06320	vascular endothelial growth factor C (VEGFC), mRNA	7424	NM_005429
A:06488	von Hippel-Lindau tumour suppressor (VHL), transcript variant 2, mRNA	7428	NM_198156
C:2407	vasoactive intestinal peptide (VIP), transcript variant 1, mRNA	7432	NM_003381
B:8107	vasoactive intestinal peptide receptor 1 (VIPR1), mRNA	7433	NM_004624
A:08324	tryptophanyl-tRNA synthetase (WARS), transcript variant 1, mRNA	7453	NM_004184

A:06953	WEE1 homolog (S. pombe) (WEE1), mRNA	7465	NM_003390
B:5487	Wilms tumour 1 (WT1), transcript variant D, mRNA	7490	NM_024426
C:0172	X-ray repair complementing defective repair in Chinese hamster cells 2 (XRCC2), mRNA	7516	NM_005431
A:02526	v-yes-1 Yamaguchi sarcoma viral oncogene homolog 1 (YES1), mRNA	7525	NM_005433
B:5702	ecotropic viral integration site 5 (EVI5), mRNA	7813	NM_005665
B:5523	BTG family, member 2 (BTG2), mRNA	7832	NM_006763
A:03788	interferon-related developmental regulator 2 (IFRD2), mRNA	7866	NM_006764
A:09614	v-maf musculoaponeurotic fibrosarcoma oncogene homolog K (avian) (MAFK), mRNA	7975	NM_002360
A:02920	frizzled homolog 3 (Drosophila) (FZD3), mRNA	7976	NM_017412
A:03507	FOS-like antigen 1 (FOSL1), mRNA	8061	NM_005438
A:00218	cullin 5 (CUL5), mRNA	8065	NM_003478
A:08128	CDK2-associated protein 1 (CDK2AP1), mRNA	8099	NM_004642
A:09843	melanoma inhibitory activity (MIA), mRNA	8190	NM_006533
A:09310	chromatin assembly factor 1, subunit B (p60) (CHAF1B), mRNA	8208	NM_005441
A:05798	SMC1 structural maintenance of chromosomes 1-like 1 (yeast) (SMC1L1), mRNA	8243	NM_006306
C:0317	axin 1 (AXIN1), transcript variant 1, mRNA	8312	NM_003502
B:0065	BRCA1 associated protein-1 (ubiquitin carboxy-terminal hydrolase) (BAP1), mRNA	8314	NM_004656
A:08801	CDC7 cell division cycle 7 (S. cerevisiae) (CDC7), mRNA	8317	NM_003503
A:09331	CDC45 cell division cycle 45-like (S. cerevisiae) (CDC45L), mRNA	8318	NM_003504
A:01727	growth factor independent 1B (potential regulator of CDKN1A, translocated in CML) (GFI1B), mRNA	8328	NM_004188
A:10009	MAD1 mitotic arrest deficient-like 1 (yeast) (MAD1L1), transcript variant 1, mRNA	8379	NM_003550
A:06561	breast cancer anti-estrogen resistance 3 (BCAR3), mRNA	8412	NM_003567
A:06461	reversion-inducing-cysteine-rich protein with kazal motifs (RECK), mRNA	8434	NM_021111
A:06991	RAD54-like (S. cerevisiae) (RAD54L), mRNA	8438	NM_003579
A:04140	NCK adaptor protein 2 (NCK2), transcript variant 1, mRNA	8440	NM_003581
B:6523	DEAH (Asp-Glu-Ala-His) box	8449	NM_003587

	polypeptide 16 DHX16		
A:09834	cullin 4B (CUL4B), mRNA	8450	NM_003588
A:06931	cullin 4A (CUL4A), transcript variant 1, mRNA	8451	NM_001008895
A:05012	cullin 3 (CUL3), mRNA	8452	NM_003590
A:05211	cullin 2 (CUL2), mRNA	8453	NM_003591
A:01673	cullin 1 (CUL1), mRNA	8454	NM_003592
C:0388	Kruppel-like factor 11 (KLF11), mRNA	8462	NM_003597
A:01318	suppressor of Ty 3 homolog (S. cerevisiae) (SUPT3H), transcript variant 2, mRNA	8464	NM_181356
A:01318	suppressor of Ty 3 homolog (S. cerevisiae) (SUPT3H), transcript variant 2, mRNA	8465	NM_181356
A:09841	protein phosphatase 1D magnesium-dependent, delta isoform (PPM1D), mRNA	8493	NM_003620
B:3627	interferon induced transmembrane protein 1 (9-27) (IFITM1), mRNA	8519	NM_003641
A:06665	growth arrest-specific 7 (GAS7), transcript variant a, mRNA	8522	NM_003644
A:10603	basic leucine zipper nuclear factor 1 (JEM-1) (BLZF1), mRNA	8548	NM_003666
A:10266	CDC14 cell division cycle 14 homolog A (S. cerevisiae) (CDC14A), transcript variant 2, mRNA	8556	NM_033312
A:09697	cyclin-dependent kinase (CDC2-like) 10 (CDK10), transcript variant 1, mRNA	8558	NM_003674
A:10520	protein kinase, interferon-inducible double stranded RNA dependent activator (PRKRA), mRNA	8575	NM_003690
A:00630	phosphatidic acid phosphatase type 2A (PPAP2A), transcript variant 2, mRNA	8611	NM_176895
B:9227	cell division cycle 2-like 5 (cholinesterase-related cell division controller) (CDC2L5), transcript variant 1, mRNA	8621	NM_003718
A:08282	tumour protein p73-like TP73L	8626	NM_003722
B:8989	aldo-keto reductase family 1, member C3 (3-alpha hydroxysteroid dehydrogenase, type II) (AKR1C3), mRNA	8644	NM_003739
B:1328	insulin receptor substrate 2 (IRS2), mRNA	8660	NM_003749
B:4001	CDC23 (cell division cycle 23, yeast, homolog) CDC23	8697	NM_004661
A:00144	tumour necrosis factor (ligand) superfamily, member 14 (TNFSF14), transcript variant 1, mRNA	8740	NM_003807
B:8481	tumour necrosis factor (ligand) superfamily, member 13 (TNFSF13), transcript variant alpha, mRNA	8741	NM_003808

A:09478	tumour necrosis factor (ligand) superfamily, member 9 (TNFSF9), mRNA	8744	NM_003811
B:8202	CD164 antigen, sialomucin (CD164), mRNA	8763	NM_006016
A:01775	RIO kinase 3 (yeast) (RIOK3), transcript variant 2, mRNA	8780	NM_145906
A:01775	RIO kinase 3 (yeast) (RIOK3), transcript variant 2, mRNA	8781	NM_145906
C:0356	tumour necrosis factor receptor superfamily, member 11a, NFKB activator (TNFRSF11A), mRNA	8792	NM_003839
A:03645	cellular repressor of E1A-stimulated genes 1 (CREG1), mRNA	8804	NM_003851
A:08261	galanin receptor 2 (GALR2), mRNA	8812	NM_003857
A:03558	cyclin-dependent kinase-like 1 (CDC2-related kinase) (CDKL1), mRNA	8814	NM_004196
B:0089	fibroblast growth factor 18 (FGF18), transcript variant 2, mRNA	8817	NM_033649
B:5592	sin3-associated polypeptide, 30kDa SAP30	8819	NM_003864
B:4763	IQ motif containing GTPase activating protein 1 (IQGAP1), mRNA	8827	NM_003870
C:0673	neuropilin 1 NRP1	8829	NM_001024628, NM_001024629, NM_003873
A:09407	histone deacetylase 3 (HDAC3), mRNA	8841	NM_003883
A:07011	alkB, alkylation repair homolog (E. coli) (ALKBH), mRNA	8847	NM_006020
A:06184	p300/CBP-associated factor (PCAF), mRNA	8850	NM_003884
A:06285	cyclin-dependent kinase 5, regulatory subunit 1 (p35) (CDK5R1), mRNA	8851	NM_003885
B:3696	chromosome 10 open reading frame 7 (C10orf7), mRNA	8872	NM_006023
C:2264	sphingosine kinase 1 (SPHK1), transcript variant 1, mRNA	8877	NM_021972
A:06721	CDC16 cell division cycle 16 homolog (S. cerevisiae) (CDC16), mRNA	8881	NM_003903
A:04142	zinc finger protein 259 (ZNF259), mRNA	8882	NM_003904
A:10737	MCM3 minichromosome maintenance deficient 3 (S. cerevisiae) associated protein (MCM3AP), mRNA	8888	NM_003906
A:03854	cyclin A1 (CCNA1), mRNA	8900	NM_003914
B:0704	B-cell CLL/lymphoma 10 (BCL10), mRNA	8915	NM_003921
A:03168	topoisomerase (DNA) III beta (TOP3B), mRNA	8940	NM_003935
B:9727	cyclin-dependent kinase 5, regulatory subunit 2 (p39) (CDK5R2), mRNA	8941	NM_003936

A:06189	protein regulator of cytokinesis 1 (PRC1), transcript variant 1, mRNA	9055	NM_003981
A:01168	DIRAS family, GTP-binding RAS-like 3 (DIRAS3), mRNA	9077	NM_004675
A:06043	protein kinase, membrane associated tyrosine/threonine 1 (PKMYT1), transcript variant 1, mRNA	9088	NM_004203
B:4778	ubiquitin specific peptidase 8 (USP8), mRNA	9101	NM_005154
B:8108	LATS, large tumour suppressor, homolog 1 (Drosophila) (LATS1), mRNA	9113	NM_004690
A:09436	chondroitin sulfate proteoglycan 6 (barnacan) (CSPG6), mRNA	9126	NM_005445
A:03606	cyclin B2 (CCNB2), mRNA	9133	NM_004701
A:10498	cyclin E2 (CCNE2), transcript variant 1, mRNA	9134	NM_057749
A:00971	Rho guanine nucleotide exchange factor (GEF) 1 (ARHGEF1), transcript variant 2, mRNA	9138	NM_004706
B:3843	hepatocyte growth factor-regulated tyrosine kinase substrate (HGS), mRNA	9146	NM_004712
A:03143	exonuclease 1 (EXO1), transcript variant 1, mRNA	9156	NM_006027
A:07881	oncostatin M receptor (OSMR), mRNA	9180	NM_003999
A:00335	ZW10, kinetochore associated, homolog (Drosophila) (ZW10), mRNA	9183	NM_004724
A:09747	BUB3 budding uninhibited by benzimidazoles 3 homolog (yeast) (BUB3), transcript variant 1, mRNA	9184	NM_004725
B:0692	leucine-rich, glioma inactivated 1 (LGI1), mRNA	9211	NM_005097
B:0692	leucine-rich, glioma inactivated 1 (LGI1), mRNA	9212	NM_005097
A:03609	nucleolar and coiled-body phosphoprotein 1 (NOLC1), mRNA	9221	NM_004741
A:04043	discs, large homolog 5 (Drosophila) (DLG5), mRNA	9231	NM_004747
A:05954	pituitary tumour-transforming 1 (PTTG1), mRNA	9232	NM_004219
B:0420	transforming growth factor beta regulator 4 (TBRG4), transcript variant 1, mRNA	9238	NM_004749
A:02479	endothelial differentiation, sphingolipid G-protein-coupled receptor, 5 (EDG5), mRNA	9294	NM_004230
A:06066	Kruppel-like factor 4 (gut) (KLF4), mRNA	9314	NM_004235
A:05541	glucagon-like peptide 2 receptor (GLP2R), mRNA	9340	NM_004246
A:00891	WD repeat domain 39 (WDR39), mRNA	9391	NM_004804
A:00519	lymphocyte antigen 86 (LY86), mRNA	9450	NM_004271

A:01180	Rho-associated, coiled-coil containing protein kinase 2 (ROCK2), mRNA	9475	NM_004850
A:01080	kinesin family member 23 (KIF23), transcript variant 2, mRNA	9493	NM_004856
A:04266	ADAM metalloproteinase with thrombospondin type 1 motif, 1 (ADAMTS1), mRNA	9510	NM_006988
B:9060	tumour protein p53 inducible protein 11 (TP53I11), mRNA	9537	NM_006034
A:04813	breast cancer anti-estrogen resistance 1 (BCAR1), mRNA	9564	NM_014567
A:09885	M-phase phosphoprotein 1 (MPHOSPH1), mRNA	9585	NM_016195
B:8184	mediator of DNA damage checkpoint 1 (MDC1), mRNA	9656	NM_014641
C:1135	extra spindle poles like 1 (S. cerevisiae) (ESPL1), mRNA	9700	NM_012291
C:0186	histone deacetylase 9 (HDAC9), transcript variant 4, mRNA	9734	NM_178423
A:05391	kinetochore associated 1 (KNTC1), mRNA	9735	NM_014708
B:0082	histone deacetylase 4 (HDAC4), mRNA	9759	NM_006037
B:0891	metastasis suppressor 1 (MTSS1), mRNA	9788	NM_014751
B:0062	Rho guanine nucleotide exchange factor (GEF) 11 (ARHGEF11), transcript variant 1, mRNA	9826	NM_014784
A:03269	tousled-like kinase 1 (TLK1), mRNA	9874	NM_012290
B:9335	RAB GTPase activating protein 1-like (RABGAP1L), transcript variant 1, mRNA	9910	NM_014857
A:08624	chromosome condensation-related SMC-associated protein 1 (CNAP1), mRNA	9918	NM_014865
B:8937	deleted in lung and esophageal cancer 1 (DLEC1), transcript variant DLEC1-L1, mRNA	9940	NM_007338
B:8656	major vault protein (MVP), transcript variant 1, mRNA	9961	NM_017458
A:02173	tumour necrosis factor (ligand) superfamily, member 15 (TNFSF15), mRNA	9966	NM_005118
A:05257	fibroblast growth factor binding protein 1 (FGFBP1), mRNA	9982	NM_005130
A:00752	REC8-like 1 (yeast) (REC8L1), mRNA	9985	NM_005132
A:01592	solute carrier family 12 (potassium/chloride transporters), member 6 (SLC12A6), mRNA	9990	NM_005135
A:04645	abl-interactor 1 (ABI1), transcript variant 1, mRNA	10006	NM_005470
A:10156	histone deacetylase 6 (HDAC6), mRNA	10013	NM_006044
B:2818	histone deacetylase 5 HDAC5	10014	NM_001015053, NM_005474

A:10510	chromatin assembly factor 1, subunit A (p150) (CHAF1A), mRNA	10036	NM_005483
A:05648	SMC4 structural maintenance of chromosomes 4-like 1 (yeast) (SMC4L1), transcript variant 3, mRNA	10051	NM_001002799
B:0675	tetraspanin 5 (TSPAN5), mRNA	10098	NM_005723
B:0685	tetraspanin 3 (TSPAN3), transcript variant 1, mRNA	10099	NM_005724
A:08229	tetraspanin 2 (TSPAN2), mRNA	10100	NM_005725
A:02634	tetraspanin 1 (TSPAN1), mRNA	10103	NM_005727
A:07852	RAD50 homolog (S. cerevisiae) (RAD50), transcript variant 1, mRNA	10111	NM_005732
B:4820	pre-B-cell colony enhancing factor 1 (PBEF1), transcript variant 1, mRNA	10135	NM_005746
B:7911	transducer of ERBB2, 1 (TOB1), mRNA	10140	NM_005749
B:0969	odt, odd Oz/ten-m homolog 1(Drosophila) (ODZ1), mRNA	10178	NM_014253
A:06242	RNA binding motif protein 7 (RBM7), mRNA	10179	NM_016090
A:03840	RNA binding motif protein 5 (RBM5), mRNA	10181	NM_005778
B:8194	M-phase phosphoprotein 9 (MPHOSPH9)	10198	NM_022782
A:09658	M-phase phosphoprotein 6 (MPHOSPH6), mRNA	10200	NM_005792
A:04009	ret finger protein 2 (RFP2), transcript variant 1, mRNA	10206	NM_005798
A:03270	proteoglycan 4 (PRG4), mRNA	10216	NM_005807
A:01614	A kinase (PRKA) anchor protein 8 (AKAP8), mRNA	10270	NM_005858
B:5575	stromal antigen 1 (STAG1), mRNA	10274	NM_005862
B:8332	aortic preferentially expressed gene 1 APEG1	10290	XM_001131579, XM_001128413
A:04828	DnaJ (Hsp40) homolog, subfamily A, member 2 (DNAJA2), mRNA	10294	NM_005880
B:0667	katanin p80 (WD repeat containing) subunit B 1 (KATNB1), mRNA	10300	NM_005886
A:04635	deleted in lymphocytic leukaemia, 1 (DLEU1) on chromosome 13	10301	NR_002605
B:2626	uracil-DNA glycosylase 2 (UNG2), transcript variant 1, mRNA	10309	NM_021147
A:09675	T-cell, immune regulator 1, ATPase, H ⁺ transporting, lysosomal V0 protein a isoform 3 (TCIRG1), transcript variant 1, mRNA	10312	NM_006019
A:09047	nucleophosmin/nucleoplasmin, 3 (NPM3), mRNA	10361	NM_006993
A:04517	synaptonemal complex protein 2 (SYCP2), mRNA	10388	NM_014258
A:06405	anaphase promoting complex subunit 10 (ANAPC10), mRNA	10393	NM_014885
A:04338	phosphatidylethanolamine N-methyltransferase (PEMT), nuclear gene encoding mitochondrial	10400	NM_007169

	protein, transcript variant 2, mRNA		
A:10053	kinetochore associated 2 (KNTC2), mRNA	10403	NM_006101
A:08539	Rap guanine nucleotide exchange factor (GEF) 3 (RAPGEF3), mRNA	10411	NM_006105
A:01717	SKB1 homolog (S. pombe) (SKB1), mRNA	10419	NM_006109
B:6182	RNA binding motif protein 14 (RBM14), mRNA	10432	NM_006328
B:4641	glycoprotein (transmembrane) nmb GPNMB	10457	NM_001005340, NM_002510
A:10829	MAD2 mitotic arrest deficient-like 2 (yeast) (MAD2L2), mRNA	10459	NM_006341
A:01067	transcriptional adaptor 3 (NGG1 homolog, yeast)-like (TADA3L), transcript variant 1, mRNA	10474	NM_006354
A:00010	vesicle transport through interaction with t-SNAREs homolog 1B (yeast) (VTI1B), mRNA	10490	NM_006370
B:1984	cartilage associated protein (CRTAP), mRNA	10491	NM_006371
A:07616	Sjogren's syndrome/scleroderma autoantigen 1 (SSSCA1), mRNA	10534	NM_006396
A:04760	ribonuclease H2, large subunit (RNASEH2A), mRNA	10535	NM_006397
A:10701	dynactin 2 (p50) (DCTN2), mRNA	10540	NM_006400
A:04950	chaperonin containing TCP1, subunit 7 (eta) (CCT7), transcript variant 1, mRNA	10574	NM_006429
A:04081	chaperonin containing TCP1, subunit 4 (delta) (CCT4), mRNA	10575	NM_006430
A:09500	chaperonin containing TCP1, subunit 2 (beta) (CCT2), mRNA	10576	NM_006431
A:09726	chromosome 6 open reading frame 108 (C6orf108), transcript variant 1, mRNA	10591	NM_006443
A:10196	SMC2 structural maintenance of chromosomes 2-like 1 (yeast) (SMC2L1), mRNA	10592	NM_006444
B:1048	ubiquitin specific peptidase 16 (USP16), transcript variant 1, mRNA	10600	NM_006447
A:08296	MAX dimerization protein 4 (MXD4), mRNA	10608	NM_006454
A:05163	synaptonemal complex protein SC65 (SC65), mRNA	10609	NM_006455
A:04356	STAM binding protein (STAMPB), transcript variant 1, mRNA	10617	NM_006463
B:3717	growth arrest-specific 2 like 1 (GAS2L1), transcript variant 1, mRNA	10634	NM_006478
A:01918	S-phase response (cyclin-related) (SPHAR), mRNA	10638	NM_006542
A:04374	KH domain containing, RNA binding, signal transduction associated 1 (KHDRBS1), mRNA	10657	NM_006559
A:08738	CCCTC-binding factor (zinc finger protein) (CTCF), mRNA	10664	NM_006565

A:08733	cell growth regulator with ring finger domain 1 (CGRRF1), mRNA	10668	NM_006568
A:07876	cell growth regulator with EF-hand domain 1 (CGREF1), mRNA	10669	NM_006569
A:05572	tumour necrosis factor (ligand) superfamily, member 13b (TNFSF13B), mRNA	10673	NM_006573
B:4752	polymerase (DNA-directed), delta 3, accessory subunit (POLD3), mRNA	10714	NM_006591
B:3500	polymerase (DNA directed), theta (POLQ), mRNA	10721	NM_199420
A:03035	nuclear distribution gene C homolog (A. nidulans) (NUDC), mRNA	10726	NM_006600
A:00069	transcription factor-like 5 (basic helix-loop-helix) (TCFL5), mRNA	10732	NM_006602
B:7543	polo-like kinase 4 (Drosophila) (PLK4), mRNA	10733	NM_014264
B:2404	stromal antigen 3 (STAG3), mRNA	10734	NM_012447
A:10760	stromal antigen 2 (STAG2), mRNA	10735	NM_006603
B:5933	transducer of ERBB2, 2 (TOB2), mRNA	10766	NM_016272
A:02195	polo-like kinase 2 (Drosophila) (PLK2), mRNA	10769	NM_006622
A:04982	zinc finger, MYND domain containing 11 (ZMYND11), transcript variant 1, mRNA	10771	NM_006624
B:2320	septin 9 (SEPT9), mRNA	10801	NM_006640
A:07660	thioredoxin-like 4A (TXNL4A), mRNA	10907	NM_006701
B:9218	SGT1, suppressor of G2 allele of SKP1 (S. cerevisiae) (SUGT1), mRNA	10910	NM_006704
A:08320	DBF4 homolog (S. cerevisiae) (DBF4), mRNA	10926	NM_006716
A:08852	spindlin (SPIN), mRNA	10927	NM_006717
A:00006	BTG family, member 3 (BTG3), mRNA	10950	NM_006806
A:01860	cytoskeleton-associated protein 4 (CKAP4), mRNA	10971	NM_006825
A:01595	microtubule-associated protein, RP/EB family, member 2 (MAPRE2), transcript variant 5, mRNA	10982	NM_014268
A:05220	cyclin I (CCNI), mRNA	10983	NM_006835
B:4359	kinesin family member 2C (KIF2C), mRNA	11004	NM_006845
A:09969	tousled-like kinase 2 (TLK2), mRNA	11011	NM_006852
A:04957	polymerase (DNA directed) sigma (POLS), mRNA	11044	NM_006999
A:01776	ubiquitin-conjugating enzyme E2C (UBE2C), transcript variant 1, mRNA	11065	NM_007019
A:09200	cytochrome b-561 domain containing 2 (CYB561D2), mRNA	11068	NM_007022
A:00904	topoisomerase (DNA) II binding protein 1 (TOPBP1), mRNA	11073	NM_007027

B:1407	ADAM metallopeptidase with thrombospondin type 1 motif, 8 (ADAMTS8), mRNA	11095	NM_007037
A:09918	katanin p60 (ATPase-containing) subunit A 1 (KATNA1), mRNA	11104	NM_007044
A:09825	PR domain containing 4 (PRDM4), mRNA	11108	NM_012406
B:7528	FGFR1 oncogene partner (FGFR1OP), transcript variant 1, mRNA	11116	NM_007045
A:04279	CD160 antigen (CD160), mRNA	11126	NM_007053
C:4275	TBC1 domain family, member 8 (with GRAM domain) (TBC1D8), mRNA	11138	NM_007063
A:03486	CDC37 cell division cycle 37 homolog (S. cerevisiae) (CDC37), mRNA	11140	NM_007065
A:06143	MYST histone acetyltransferase 2 (MYST2), mRNA	11143	NM_007067
A:06472	DMC1 dosage suppressor of mck1 homolog, meiosis-specific homologous recombination (yeast) (DMC1), mRNA	11144	NM_007068
A:07181	coronin, actin binding protein, 1A (CORO1A), mRNA	11151	NM_007074
A:04421	Huntingtin interacting protein E (HYPE), mRNA	11153	NM_007076
A:03200	PC4 and SFRS1 interacting protein 1 (PSIP1), transcript variant 2, mRNA	11168	NM_033222
C:0370	centrosomal protein 2 (CEP2), transcript variant 1, mRNA	11190	NM_007186
C:0370	centrosomal protein 2 (CEP2), transcript variant 1, mRNA	11191	NM_007186
A:02177	CHK2 checkpoint homolog (S. pombe) (CHEK2), transcript variant 1, mRNA	11200	NM_007194
A:09335	polymerase (DNA directed), gamma 2, accessory subunit (POLG2), mRNA	11232	NM_007215
A:08008	dynactin 3 (p22) (DCTN3), transcript variant 2, mRNA	11258	NM_024348
B:7247	three prime repair exonuclease 1 (TREX1), transcript variant 2, mRNA	11277	NM_033627
A:03276	polynucleotide kinase 3'-phosphatase (PNKP), mRNA	11284	NM_007254
A:01322	Parkinson disease (autosomal recessive, early onset) 7 (PARK7), mRNA	11315	NM_007262
B:5525	PDGFA associated protein 1 (PDAP1), mRNA	11333	NM_014891
A:05117	tumour suppressor candidate 2 (TUSC2), mRNA	11334	NM_007275
A:08584	activating transcription factor 5 (ATF5), mRNA	22809	NM_012068
A:10029	KIAA0971 (KIAA0971), mRNA	22868	NM_014929

C:4180	DENN/MADD domain containing 3 (DENND3), mRNA	22898	NM_014957
A:07655	microtubule-associated protein, RP/EB family, member 1 (MAPRE1), mRNA	22919	NM_012325
A:02013	sirtuin (silent mating type information regulation 2 homolog) 2 (<i>S. cerevisiae</i>) (SIRT2), transcript variant 2, mRNA	22933	NM_030593
A:07965	TPX2, microtubule-associated, homolog (<i>Xenopus laevis</i>) (TPX2), mRNA	22974	NM_012112
B:1032	apoptotic chromatin condensation inducer 1 ACIN1	22985	NM_014977
A:10375	androgen-induced proliferation inhibitor (APRIN), transcript variant 1, mRNA	23047	NM_015032
A:04696	nuclear receptor coactivator 6 (NCOA6), mRNA	23054	NM_014071
A:09165	KIAA0676 protein (KIAA0676), transcript variant 1, mRNA	23061	NM_198868
B:4976	KIAA0261 (KIAA0261), mRNA	23063	NM_015045
B:8950	KIAA0241 protein (KIAA0241), mRNA	23080	NM_015060
C:2458	p53-associated parkin-like cytoplasmic protein (PARC), mRNA	23113	NM_015089
B:9549	SMC5 structural maintenance of chromosomes 5-like 1 (yeast) (SMC5L1), mRNA	23137	NM_015110
B:4428	septin 6 (SEPT6), transcript variant 1, mRNA	23157	NM_145799
B:6278	KIAA0882 protein (KIAA0882), mRNA	23158	NM_015130
B:1443	septin 8 (SEPT8), mRNA	23176	XM_034872
B:8136	ankyrin repeat domain 15 (ANKRD15), transcript variant 1, mRNA	23189	NM_015158
B:4969	KIAA1086 (KIAA1086), mRNA	23217	XM_001130130, XM_001130674
A:10369	phospholipase C, beta 1 (phosphoinositide-specific) (PLCB1), transcript variant 2, mRNA	23236	NM_182734
B:0524	RAB6 interacting protein 1 (RAB6IP1), mRNA	23258	NM_015213
B:0230	inducible T-cell co-stimulator ligand ICOSLG	23308	NM_015259
B:0327	SAM and SH3 domain containing 1 (SASH1), mRNA	23328	NM_015278
B:5714	KIAA0650 protein (KIAA0650), mRNA	23347	XM_113962, XM_938891
B:8897	formin binding protein 4 (FNBP4), mRNA	23360	NM_015308
B:8228	barren homolog 1 (<i>Drosophila</i>) (BRRN1), mRNA	23397	NM_015341
B:9601	ATPase type 13A2 (ATP13A2), mRNA	23401	NM_022089

B:7418	TAR DNA binding protein (TARDBP), mRNA	23435	NM_007375
B:7878	microtubule-actin crosslinking factor 1 (MACF1), transcript variant 1, mRNA	23499	NM_012090
A:09105	RNA binding motif protein 9 (RBM9), transcript variant 2, mRNA	23543	NM_014309
B:1165	origin recognition complex, subunit 6 homolog-like (yeast) (ORC6L), mRNA	23594	NM_014321
B:3180	origin recognition complex, subunit 3-like (yeast) (ORC3L), transcript variant 2, mRNA	23595	NM_012381
A:00473	SPO11 meiotic protein covalently bound to DSB-like (S. cerevisiae) (SPO11), transcript variant 1, mRNA	23626	NM_012444
A:02179	RAB GTPase activating protein 1 (RABGAP1), mRNA	23637	NM_012197
A:06494	leucine zipper, down-regulated in cancer 1 (LDOC1), mRNA	23641	NM_012317
B:2198	protein phosphatase 1, regulatory (inhibitor) subunit 15A (PPP1R15A), mRNA	23645	NM_014330
C:3173	polymerase (DNA directed), alpha 2 (70kD subunit) (POLA2), mRNA	23649	NM_002689
A:03098	SH3-domain binding protein 4 (SH3BP4), mRNA	23677	NM_014521
C:1904	N-acetyltransferase 6 (NAT6), mRNA	24142	NM_012191
C:2118	unc-84 homolog B (C. elegans) (UNC84B), mRNA	25777	NM_015374
A:05344	RAD54 homolog B (S. cerevisiae) (RAD54B), transcript variant 1, mRNA	25788	NM_012415
A:06762	CDKN1A interacting zinc finger protein 1 (CIZ1), mRNA	25792	NM_012127
C:4297	Nipped-B homolog (Drosophila) (NIPBL), transcript variant B, mRNA	25836	NM_015384
A:09401	preimplantation protein 3 (PREI3), transcript variant 1, mRNA	25843	NM_015387
B:3103	breast cancer metastasis suppressor 1 (BRMS1), transcript variant 1, mRNA	25855	NM_015399
A:01151	protein kinase D2 (PRKD2), mRNA	25869	NM_016457
A:07688	EGF-like-domain, multiple 6 (EGFL6), mRNA	25975	NM_015507
B:6248	ankyrin repeat domain 17 (ANKRD17), transcript variant 1, mRNA	26057	NM_032217
A:02605	adaptor protein containing pH domain, PTB domain and leucine zipper motif 1 (APPL), mRNA	26060	NM_012096
A:02500	ets homologous factor (EHF), mRNA	26298	NM_012153
A:09724	mutL homolog 3 (E. coli) (MLH3), mRNA	27030	NM_014381

A:06200	lysosomal-associated membrane protein 3 (LAMP3), mRNA	27074	NM_014398
A:00686	tetraspanin 13 (TSPAN13), mRNA	27075	NM_014399
A:02984	calcyclin binding protein (CACYPB), transcript variant 1, mRNA	27101	NM_014412
A:00435	eukaryotic translation initiation factor 2-alpha kinase 1 (EIF2AK1), mRNA	27104	NM_014413
C:8169	SMC1 structural maintenance of chromosomes 1-like 2 (yeast) (SMC1L2), mRNA	27127	NM_148674
A:00927	sestrin 1 (SESN1), mRNA	27244	NM_014454
A:01831	RNA binding motif, single stranded interacting protein (RBMS3), transcript variant 2, mRNA	27303	NM_014483
A:06053	zinc finger protein 330 (ZNF330), mRNA	27309	NM_014487
A:03501	down-regulated in metastasis (DRIM), mRNA	27340	NM_014503
B:3842	polymerase (DNA directed), lambda (POLL), mRNA	27343	NM_013274
B:6569	polymerase (DNA directed), mu (POLM), mRNA	27434	NM_013284
B:4351	echinoderm microtubule associated protein like 4 (EML4), mRNA	27436	NM_019063
B:1612	cat eye syndrome chromosome region, candidate 4 CECR4	27443	AF307448
A:08058	protein phosphatase 2 (formerly 2A), regulatory subunit B", beta (PPP2R3B), transcript variant 1, mRNA	28227	NM_013239
A:09647	response gene to complement 32 (RGC32), mRNA	28984	NM_014059
A:09821	malignant T cell amplified sequence 1 (MCTS1), mRNA	28985	NM_014060
B:6485	HSPC135 protein (HSPC135), transcript variant 1, mRNA	29083	NM_014170
A:09945	PYD and CARD domain containing (PYCARD), transcript variant 1, mRNA	29108	NM_013258
C:1944	lectin, galactoside-binding, soluble, 13 (galectin 13) (LGALS13), mRNA	29124	NM_013268
A:02160	CD274 antigen (CD274), mRNA	29126	NM_014143
A:08075	replication initiator 1 (REPIN1), transcript variant 1, mRNA	29803	NM_013400
B:1479	anaphase promoting complex subunit 2 (ANAPC2), mRNA	29882	NM_013366
A:08657	protein predicted by clone 23882 (HSU79303), mRNA	29903	NM_013301
A:10453	replication protein A4, 34kDa (RPA4), mRNA	29935	NM_013347
A:02862	anaphase promoting complex subunit 4 (ANAPC4), mRNA	29945	NM_013367
A:10100	SERTA domain containing 1 (SERTAD1), mRNA	29950	NM_013376
A:05316	striatin, calmodulin binding protein 3 (STRN3), mRNA	29966	NM_014574
A:06440	G0/G1switch 2 (G0S2), mRNA	50486	NM_015714

A:08113	deleted in esophageal cancer 1 (DEC1), mRNA	50514	NM_017418
B:7919	hepatoma-derived growth factor, related protein 3 (HDGFRP3), mRNA	50810	NM_016073
A:07482	par-6 partitioning defective 6 homolog alpha (C. elegans) (PAR6A), transcript variant 1, mRNA	50855	NM_016948
A:03435	geminin, DNA replication inhibitor (GMNN), mRNA	51053	NM_015895
A:00171	ribosomal protein S27-like (RPS27L), mRNA	51065	NM_015920
B:1459	EGF-like-domain, multiple 7 (EGFL7), transcript variant 1, mRNA	51162	NM_016215
A:09081	tubulin, epsilon 1 (TUBE1), mRNA	51175	NM_016262
A:08522	hect domain and RLD 5 (HERC5), mRNA	51191	NM_016323
A:05174	phospholipase C, epsilon 1 (PLCE1), mRNA	51196	NM_016341
B:3533	dual specificity phosphatase 13 DUSP13	51207	NM_001007271, NM_001007272, NM_001007273, NM_001007274, NM_001007275, NM_016364
A:06537	Abi gene family, member 3 (ABI3), mRNA	51225	NM_016428
A:03107	transcription factor Dp family, member 3 (TFDP3), mRNA	51270	NM_016521
A:09430	SCAN domain containing 1 (SCAND1), transcript variant 1, mRNA	51282	NM_016558
B:9657	CD320 antigen (CD320), mRNA	51293	NM_016579
A:07215	fizzy/cell division cycle 20 related 1 (Drosophila) (FZR1), mRNA	51343	NM_016263
A:06101	Wilms tumour upstream neighbor 1 (WIT1), mRNA	51352	NM_015855
A:10614	E3 ubiquitin protein ligase, HECT domain containing, 1 (EDD1), mRNA	51366	NM_015902
B:9794	anaphase promoting complex subunit 5 (ANAPC5), mRNA	51433	NM_016237
B:1481	anaphase promoting complex subunit 7 (ANAPC7), mRNA	51434	NM_016238
A:08459	G-2 and S-phase expressed 1 (GTSE1), mRNA	51512	NM_016426
A:02842	APC11 anaphase promoting complex subunit 11 homolog (yeast) (ANAPC11), transcript variant 2, mRNA	51529	NM_0164760
B:2670	histone deacetylase 7A HDAC7A	51564	NM_015401,
A:07829	ubiquitin-conjugating enzyme E2D 4 (putative) (UBE2D4), mRNA	51619	NM_015983
A:09440	CDK5 regulatory subunit associated protein 1 (CDK5RAP1), transcript variant 2, mRNA	51654	NM_016082

B:1035	DNA replication complex GINS protein PSF2 (Pfs2), mRNA	51659	NM_016095
B:9464	sterile alpha motif and leucine zipper containing kinase AZK (ZAK), transcript variant 2, mRNA	51776	NM_133646
B:7871	ZW10 interactor antisense ZWINTAS	53588	X98261
B:3431	RNA binding motif protein 11 (RBM11), mRNA	54033	NM_144770
A:02209	polymerase (DNA directed), epsilon 3 (p17 subunit) (POLE3), mRNA	54107	NM_017443
A:04070	DKFZp434A0131 protein DKFZP434A0131	54441	NM_018991
A:05280	anillin, actin binding protein (scraps homolog, Drosophila) (ANLN), mRNA	54443	NM_018685
A:06475	spindlin family, member 2 (SPIN2), mRNA	54466	NM_019003
A:03960	cyclin J (CCNJ), mRNA	54619	NM_019084
B:3841	M-phase phosphoprotein, mpp8 (HSMPP8), mRNA	54737	NM_017520
B:8673	ropporin, raphilin associated protein 1 (ROPN1), mRNA	54763	NM_017578
A:02474	B-cell translocation gene 4 (BTG4), mRNA	54766	NM_017589
B:2084	G patch domain containing 4 (GPATC4), transcript variant 2, mRNA	54865	NM_182679
A:06639	hypothetical protein FLJ20422 (FLJ20422), mRNA	54929	NM_017814
C:2265	thioredoxin-like 4B (TXNL4B), mRNA	54957	NM_017853
B:7809	PIN2-interacting protein 1 (PINX1), mRNA	54984	NM_017884
B:8204	polybromo 1 (PB1), transcript variant 2, mRNA	55193	NM_018313
A:03321	hypothetical protein FLJ10781 (FLJ10781), mRNA	55228	NM_018215
B:2270	MOB1, Mps One Binder kinase activator-like 1B (yeast) MOBK1B	55233	NM_018221
A:08002	signal-regulatory protein beta 2 (SIRPB2), transcript variant 1, mRNA	55423	NM_018556
A:03524	tripartite motif-containing 36 (TRIM36), transcript variant 1, mRNA	55522	NM_018700
A:09474	chromosome 2 open reading frame 29 (C2orf29), mRNA	55571	NM_017546
A:05414	hypothetical protein H41 (H41), mRNA	55573	NM_017548
B:2133	CDC37 cell division cycle 37 homolog (S. cerevisiae)-like 1 (CDC37L1), mRNA	55664	NM_017913
B:8413	Nedd4 binding protein 2 (N4BP2), mRNA	55728	NM_018177
A:02898	checkpoint with forkhead and ring finger domains (CHFR), mRNA	55743	NM_018223
A:07468	sepin 11 (SEPT11), mRNA	55752	NM_018243

B:2252	chondroitin beta1,4 N-acetylgalactosaminyltransferase (ChGn), mRNA	55790	NM_018371
C:0033	B double prime 1, subunit of RNA polymerase III transcription initiation factor IIIB BDP1	55814	NM_018429
A:03912	PDZ binding kinase (PBK), mRNA	55872	NM_018492
A:10308	unc-45 homolog A (C. elegans) (UNC45A), transcript variant 1, mRNA	55898	NM_017979
A:02027	bridging integrator 3 (BIN3), mRNA	55909	NM_018688
C:0655	erbB2 interacting protein ERBB2IP	55914	NM_001006600, NM_018695
B:1503	septin 3 (SEPT3), transcript variant C, mRNA	55964	NM_145734
B:8446	gastrokine 1 (GKN1), mRNA	56287	NM_019617
A:00073	par-3 partitioning defective 3 homolog (C. elegans) (PARD3), mRNA	56288	NM_019619
A:03990	CTP synthase II (CTPS2), transcript variant 1, mRNA	56475	NM_019857
B:8449	BRCA2 and CDKN1A interacting protein (BCCIP), transcript variant B, mRNA	56647	NM_078468
B:1203	interferon, kappa (IFNK), mRNA	56832	NM_020124
B:1205	SLAM family member 8 (SLAMF8), mRNA	56833	NM_020125
A:00149	sphingosine kinase 2 (SPHK2), mRNA	56848	NM_020126
A:04220	Werner helicase interacting protein 1 (WRNIP1), transcript variant 1, mRNA	56897	NM_020135
A:09095	latexin (LXN), mRNA	56925	NM_020169
A:02450	dual specificity phosphatase 22 (DUSP22), mRNA	56940	NM_020185
C:0975	DC13 protein (DC13), mRNA	56942	NM_020188
A:04008	5',3'-nucleotidase, mitochondrial (NT5M), nuclear gene encoding mitochondrial protein, mRNA	56953	NM_020201
A:01586	kinesin family member 15 (KIF15), mRNA	56992	NM_020242
B:0396	catenin, beta interacting protein 1 (CTNNBIP1), transcript variant 1, mRNA	56998	NM_020248
B:3508	cyclin L1 (CCNL1), mRNA	57018	NM_020307
A:06501	cholinergic receptor, nicotinic, alpha polypeptide 10 (CHRNA10), mRNA	57053	NM_020402
B:7311	poly(rC) binding protein 4 (PCBP4), transcript variant 1, mRNA	57060	NM_020418
A:08184	chromosome 1 open reading frame 128 (C1orf128), mRNA	57095	NM_020362
B:3446	S100 calcium binding protein A14 (S100A14), mRNA	57402	NM_020672
C:5669	odz, odd Oz/ten-m homolog 2 (Drosophila) (ODZ2), mRNA	57451	XM_047995, XM_931456, XM_942208, XM_945786, XM_945788

B:8403	membrane-associated ring finger (C3HC4) 4 (MARCH4), mRNA	57574	NM_020814
B:1442	polymerase (DNA-directed), delta 4 (POLD4), mRNA	57804	NM_021173
B:1448	prokineticin 2 (PROK2), mRNA	60675	NM_021935
B:4091	CTF18, chromosome transmission fidelity factor 18 homolog (S. cerevisiae) (CHTF18), mRNA	63922	NM_022092
C:0644	TSPY-like 2 (TSPYL2), mRNA	64061	NM_022117
B:6809	chromosome 10 open reading frame 54 (C10orf54), mRNA	64115	NM_022153
A:10488	chromosome condensation protein G (HCAP-G), mRNA	64151	NM_022346
A:10186	spermatogenesis associated 1 (SPATA1), mRNA	64173	NM_022354
A:02978	DNA cross-link repair 1C (PSO2 homolog, S. cerevisiae) (DCLRE1C), transcript variant b, mRNA	64421	NM_022487
A:10112	anaphase promoting complex subunit 1 (ANAPC1), mRNA	64682	NM_022662
A:10470	FLJ20859 gene (FLJ20859), transcript variant 1, mRNA	64745	NM_001029991
B:3988	interferon stimulated exonuclease gene 20kDa-like 1 (ISG20L1), mRNA	64782	NM_022767
A:06358	DNA cross-link repair 1B (PSO2 homolog, S. cerevisiae) (DCLRE1B), mRNA	64858	NM_022836
A:10073	centromere protein H (CENPH), mRNA	64946	NM_022909
A:05903	chromosome 16 open reading frame 24 (C16orf24), mRNA	65990	NM_023933
A:07975	spermatogenesis associated 5-like 1 (SPATA5L1), mRNA	79029	NM_024063
A:01368	hypothetical protein MGC5297 (MGC5297), mRNA	79072	NM_024091
C:1382	basic helix-loop-helix domain containing, class B, 3 (BHLHB3), mRNA	79365	NM_030762
A:00699	NADPH oxidase, EF-hand calcium binding domain 5 (NOX5), mRNA	79400	NM_024505
A:05363	SMC6 structural maintenance of chromosomes 6-like 1 (yeast) (SMC6L1), mRNA	79677	NM_024624
A:09775	V-set domain containing T cell activation inhibitor 1 (VTCN1), mRNA	79679	NM_024626
B:6021	hypothetical protein FLJ21125 (FLJ21125), mRNA	79680	NM_024627
A:06447	Sin3A associated protein p30-like (SAP30L), mRNA	79685	NM_024632
A:08767	suppressor of variegation 3-9 homolog 2 (Drosophila) (SUV39H2), mRNA	79723	NM_024670
A:01156	chromosome 15 open reading frame 29 (C15orf29), mRNA	79768	NM_024713

A:03654	hypothetical protein FLJ13273 (FLJ13273), transcript variant 1, mRNA	79807	NM_001031720
A:10726	hypothetical protein FLJ13265 (FLJ13265), mRNA	79935	NM_024877
B:2392	Dbf4-related factor 1 (DRF1), transcript variant 2, mRNA	80174	NM_025104
B:2358	SMP3 mannosyltransferase (SMP3), mRNA	80235	NM_025163
A:02900	CDK5 regulatory subunit associated protein 3 (CDK5RAP3), transcript variant 2, mRNA	80279	NM_025197
C:0025	leucine rich repeat containing 27 (LRRC27), mRNA	80313	NM_030626
B:9631	ADAM metallopeptidase domain 33 (ADAM33), transcript variant 1, mRNA	80332	NM_025220
B:6501	CD276 antigen (CD276), transcript variant 2, mRNA	80381	NM_025240
A:05386	hypothetical protein MGC10334 (MGC10334), mRNA	80772	NM_001029885
A:08918	collagen, type XVIII, alpha 1 (COL18A1), transcript variant 1, mRNA	80781	NM_030582
C:0358	EGF-like-domain, multiple 8 (EGFL8), mRNA	80864	NM_030652
B:1020	C/EBP-induced protein (LOC81558), mRNA	81558	NM_030802
B:3550	DNA replication factor (CDT1), mRNA	81620	NM_030928
B:5661	cyclin L2 (CCNL2), mRNA	81669	NM_030937
B:1735	exonuclease NEF-sp (LOC81691), mRNA	81691	NM_030941
B:2768	ring finger protein 146 (RNF146), mRNA	81847	NM_030963
B:2350	interferon stimulated exonuclease gene 20kDa-like 2 (ISG20L2), mRNA	81875	NM_030980
B:3823	Cdk5 and Abl enzyme substrate 2 (CABLES2), mRNA	81928	NM_031215
B:8839	leucine rich repeat containing 48 (LRRC48), mRNA	83450	NM_031294
B:9709	katanin p60 subunit A-like 2 (KATNAL2), mRNA	83473	NM_031303
B:8709	sestrin 2 (SESN2), mRNA	83667	NM_031459
B:8721	CD99 antigen-like 2 (CD99L2), transcript variant 1, mRNA	83692	NM_031462
C:0565	regenerating islet-derived family, member 4 (REG4), mRNA	83998	NM_032044
B:3599	katanin p60 subunit A-like 1 (KATNAL1), transcript variant 1, mRNA	84056	NM_032116
B:3492	GAJ protein (GAJ), mRNA	84057	NM_032117
A:00224	IQ motif containing G (IQCG), mRNA	84223	NM_032263
C:1051	hypothetical protein MGC10911 (MGC10911), mRNA	84262	NM_032302
B:1756	prokineticin 1 (PROK1), mRNA	84432	NM_032414

B:3029	MCM8 minichromosome maintenance deficient 8 (<i>S. cerevisiae</i>) (MCM8), transcript variant 1, mRNA	84515	NM_032485
C:0555	RNA binding motif protein 13 (RBM13), mRNA	84552	NM_032509
C:1586	par-6 partitioning defective 6 homolog beta (<i>C. elegans</i>) (PARD6B), mRNA	84612	NM_032521
C:1872	resistin like beta (RETNLB), mRNA	84666	NM_032579
B:9569	protein phosphatase 1, regulatory subunit 9B, spinophilin (PPP1R9B), mRNA	84687	NM_032595
B:3610	hepatoma-derived growth factor-related protein 2 (HDGF2), transcript variant 2, mRNA	84717	NM_032631
B:4127	lamin B2 (LMNB2), mRNA	84823	NM_032737
B:2733	apoptosis-inducing factor (AIF)-like mitochondrion-associated inducer of death (AMID), mRNA	84883	NM_032797
B:4273	RAS-like, estrogen-regulated, growth inhibitor (REG), mRNA	85004	NM_032918
B:9560	cyclin B3 (CCNB3), transcript variant 1, mRNA	85417	NM_033670
C:0075	leucine rich repeat and coiled-coil domain containing 1 (LRRCC1), mRNA	85444	NM_033402
B:8110	tripartite motif-containing 4 (TRIM4), transcript variant alpha, mRNA	89765	NM_033017
B:6017	hypothetical gene CG018, CG018	90634	NM_052818
C:0238	NIMA (never in mitosis gene a)-related kinase 9 (NEK9), mRNA	91754	NM_033116
B:3862	Cdk5 and Abl enzyme substrate 1 (CABLES1), mRNA	91768	NM_138375
B:3802	chordin-like 1 (CHRD1), mRNA	91860	NM_145234
B:3730	family with sequence similarity 58, member A (FAM58A), mRNA	92002	NM_152274
B:6762	secretoglobin, family 3A, member 1 (SCGB3A1), mRNA	92304	NM_052863
B:4458	membrane-associated ring finger (C3HC4) 9 MARCH9	92979	NM_138396
B:9351	immunoglobulin superfamily, member 8 (IGSF8), mRNA	93185	NM_052868
B:1687	acid phosphatase, testicular (ACPT), transcript variant A, mRNA	93650	NM_033068
B:3540	RAS guanyl releasing protein 4 (RASGRP4), transcript variant 1, mRNA	115727	NM_170603
C:4836	topoisomerase (DNA) I, mitochondrial (TOP1MT), nuclear gene encoding mitochondrial protein, mRNA	116447	NM_052963
B:9435	mediator of RNA polymerase II transcription, subunit 12 homolog (yeast)-like (MED12L), mRNA	116931	NM_053002
C:3793	amyotrophic lateral sclerosis 2 (juvenile) chromosome region, candidate 19 (ALS2CR19), transcript variant b, mRNA	117583	NM_152526

C:3467	KIAA1977 protein (KIAA1977), mRNA	124404	NM_133450
C:3112	ubiquitin specific protease 43 (USP43), mRNA	124817	XM_945578
C:5265	hypothetical protein BC009732 (LOC133308), mRNA	133396	NM_178833
A:07401	myosin light chain 1 slow a (MLC1SA), mRNA	140466	NM_002475
C:1334	CCCTC-binding factor (zinc finger protein)-like (CTCFL), mRNA	140690	NM_080618
B:5293	chromosome 20 open reading frame 181 C20orf181	140849	U63828
B:9316	hypothetical protein MGC20470 (MGC20470), mRNA	143686	NM_145053
B:9599	septin 10 (SEPT10), transcript variant 1, mRNA	151011	NM_144710
C:0962	similar to hepatocellular carcinoma-associated antigen HCA557b (LOC151194), mRNA	151195	NM_145280
C:1752	connexin40 (CX40), mRNA	219771	NM_153368
B:3031	kinesin family member 6 (KIF6), mRNA	221527	NM_145027
B:1737	chromosome Y open reading frame 15A (CYorf15A), mRNA	246176	NM_001005852
B:8632	DNA directed RNA polymerase II polypeptide J-related gene (POLR2J2), transcript variant 3, mRNA	246778	NM_032959
A:08544	zinc finger, DHHC-type containing 24 (ZDHHC24), mRNA	254394	NM_207340
C:3659	growth arrest-specific 2 like 3 (GAS2L3), mRNA	283431	NM_174942
B:5467	laminin, alpha 1 (LAMA1), mRNA	284217	NM_005559
C:2399	hypothetical protein MGC26694 (MGC26694), mRNA	284439	NM_178526
C:5315	cation channel, sperm associated 3 (CATSPER3), mRNA	347733	NM_178019
B:0631	polymerase (DNA directed) nu (POLN), mRNA	353497	NM_181808

Table B: Known cell proliferation-related genes. All genes categorized as cell proliferation-related by gene ontology analysis and present on the Affymetrix HG-U133 platform.

5

General Approaches to Prognostic Marker Detection

The following approaches are non-limiting methods that can be used to detect the proliferation markers, including GCPM family members: microarray approaches using oligonucleotide probes selective for a GCPM; real-time qPCR on tumour samples using GCPM specific primers and probes; real-time qPCR on lymph node, blood, serum, faecal, or urine samples using GCPM specific primers and probes; enzyme-linked immunological

assays (ELISA); immunohistochemistry using anti-marker antibodies; and analysis of array or qPCR data using computers.

Other useful methods include northern blotting and *in situ* hybridization (Parker and Barnes, *Methods in Molecular Biology* 106: 247-283 (1999)); RNase protection assays (Hod, *BioTechniques* 13: 852-854 (1992)); reverse transcription polymerase chain reaction (RT-PCR; Weis et al., *Trends in Genetics* 8: 263-264 (1992)); serial analysis of gene expression (SAGE; Velculescu et al., *Science* 270: 484-487 (1995); and Velculescu et al., *Cell* 88: 243-51 (1997)), MassARRAY technology (Sequenom, San Diego, CA), and gene expression analysis by massively parallel signature sequencing (MPSS; Brenner et al., *Nature Biotechnology* 18: 630-634 (2000)). Alternatively, antibodies may be employed that can recognize specific complexes, including DNA duplexes, RNA duplexes, and DNA-RNA hybrid duplexes or DNA-protein duplexes.

Primary data can be collected and fold change analysis can be performed, for example, by comparison of marker expression levels in tumour tissue and non-tumour tissue; by comparison of marker expression levels to levels determined in recurring tumours and non-recurring tumours; by comparison of marker expression levels to levels determined in tumours with or without metastasis; by comparison of marker expression levels to levels determined in differently staged tumours; or by comparison of marker expression levels to levels determined in cells with different levels of proliferation. A negative or positive prognosis is determined based on this analysis. Further analysis of tumour marker expression includes matching those markers exhibiting increased or decreased expression with expression profiles of known gastrointestinal tumours to provide a prognosis.

A threshold for concluding that expression is increased is provided as, for example, at least a 1.5-fold or 2-fold increase, and in alternative embodiments, at least a 3-fold increase, 4-fold increase, or 5-fold increase. A threshold for concluding that expression is decreased is provided as, for example, at least a 1.5-fold or 2-fold decrease, and in alternative embodiments, at least a 3-fold decrease, 4-fold decrease, or 5-fold decrease. It can be appreciated that other thresholds for concluding that increased or decreased expression has occurred can be selected without departing from the scope of this invention.

It will also be appreciated that a threshold for concluding that expression is increased will be dependent on the particular marker and also the particular predictive model that is to

be applied. The threshold is generally set to achieve the highest sensitivity and selectivity with the lowest error rate, although variations may be desirable for a particular clinical situation. The desired threshold is determined by analysing a population of sufficient size taking into account the statistical variability of any predictive model and is calculated from the size of the sample used to produce the predictive model. The same applies for the determination of a threshold for concluding that expression is decreased. It can be appreciated that other thresholds, or methods for establishing a threshold, for concluding that increased or decreased expression has occurred can be selected without departing from the scope of this invention.

It is also possible that a prediction model may produce as its output a numerical value, for example a score, likelihood value or probability. In these instances, it is possible to apply thresholds to the results produced by prediction models, and in these cases similar principles apply as those used to set thresholds for expression values

Once the expression level of one or more proliferation markers in a tumour sample has been obtained the likelihood of the cancer recurring can then be determined. In accordance with the invention, a negative prognosis is associated with decreased expression of at least one proliferation marker, while a positive prognosis is associated with increased expression of at least one proliferation marker. In various aspects, an increase in expression is shown by at least 1, 2, 3, 4, 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, or 75 of the markers disclosed herein. In other aspects, a decrease in expression is shown by at least 1, 2, 3, 4, 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, or 75 of the markers disclosed herein

From the genes identified, proliferation signatures comprising one or more GCPMs can be used to determine the prognosis of a cancer, by comparing the expression level of the one or more genes to the disclosed proliferation signature. By comparing the expression of one or more of the GCPMs in a tumour sample with the disclosed proliferation signature, the likelihood of the cancer recurring can be determined. The comparison of expression levels of the prognostic signature to establish a prognosis can be done by applying a predictive model as described previously.

Determining the likelihood of the cancer recurring is of great value to the medical practitioner. A high likelihood of reoccurrence means that a longer or higher dose treatment should be given, and the patient should be more closely monitored for signs of recurrence of the cancer. An accurate prognosis is also of benefit to the patient. It allows

the patient, along with their partners, family, and friends to also make decisions about treatment, as well as decisions about their future and lifestyle changes. Therefore, the invention also provides for a method establishing a treatment regime for a particular cancer based on the prognosis established by matching the expression of the markers in
5 a tumour sample with the differential proliferation signature.

It will be appreciated that the marker selection, or construction of a proliferation signature, does not have to be restricted to the GCPMs disclosed in Table A, Table B, Table C or Table D, herein, but could involve the use of one or more GCPMs from the disclosed
10 signature, or a new signature may be established using GCPMs selected from the disclosed marker lists. The requirement of any signature is that it predicts the likelihood of recurrence with enough accuracy to assist a medical practitioner to establish a treatment regime.

15 Surprisingly, it was discovered that many of the GCPM were associated with increased levels of cell proliferation, and were also associated with a positive prognosis. It has similarly been found that there is a close correlation between the decreased expression level of GCPMs and a negative prognosis, e.g., an increased likelihood of gastrointestinal cancer recurring. Therefore, the present invention also provides for the use of a marker
20 associated with cell proliferation, e.g., a cell cycle component, as a GCPM.

As described herein, determination of the likelihood of a cancer recurring can be accomplished by measuring expression of one or more proliferation-specific markers. The methods provided herein also include assays of high sensitivity. In particular, qPCR is
25 extremely sensitive, and can be used to detect markers in very low copy number (e.g., 1 – 100) in a sample. With such sensitivity, prognosis of gastrointestinal cancer is made reliable, accurate, and easily tested.

Reverse Transcription PCR (RT-PCR)

30 Of the techniques listed above, the most sensitive and most flexible quantitative method is RT-PCR, which can be used to compare RNA levels in different sample populations, in normal and tumour tissues, with or without drug treatment, to characterize patterns of expression, to discriminate between closely related RNAs, and to analyze RNA structure.

35 For RT-PCR, the first step is the isolation of RNA from a target sample. The starting material is typically total RNA isolated from human tumours or tumour cell lines, and corresponding normal tissues or cell lines, respectively. RNA can be isolated from a

variety of samples, such as tumour samples from breast, lung, colon (e.g., large bowel or small bowel), colorectal, gastric, esophageal, anal, rectal, prostate, brain, liver, kidney, pancreas, spleen, thymus, testis, ovary, uterus, etc., tissues, from primary tumours, or tumour cell lines, and from pooled samples from healthy donors. If the source of RNA is a
5 tumour, RNA can be extracted, for example, from frozen or archived paraffin-embedded and fixed (e.g., formalin-fixed) tissue samples.

The first step in gene expression profiling by RT-PCR is the reverse transcription of the RNA template into cDNA, followed by its exponential amplification in a PCR reaction. The
10 two most commonly used reverse transcriptases are avian myeloblastosis virus reverse transcriptase (AMV-RT) and Moloney murine leukaemia virus reverse transcriptase (MMLV-RT). The reverse transcription step is typically primed using specific primers, random hexamers, or oligo-dT primers, depending on the circumstances and the goal of expression profiling. For example, extracted RNA can be reverse-transcribed using a
15 GeneAmp RNA PCR kit (Perkin Elmer, CA, USA), following the manufacturer's instructions. The derived cDNA can then be used as a template in the subsequent PCR reaction.

Although the PCR step can use a variety of thermostable DNA-dependent DNA
20 polymerases, it typically employs the Taq DNA polymerase, which has a 5'-3' nuclease activity but lacks a 3'-5' proofreading endonuclease activity. Thus, TaqMan (g) PCR typically utilizes the 5' nuclease activity of Taq or Tth polymerase to hydrolyze a hybridization probe bound to its target amplicon, but any enzyme with equivalent 5' nuclease activity can be used.

25 Two oligonucleotide primers are used to generate an amplicon typical of a PCR reaction. A third oligonucleotide, or probe, is designed to detect nucleotide sequence located between the two PCR primers. The probe is non-extendible by Taq DNA polymerase enzyme, and is labeled with a reporter fluorescent dye and a quencher fluorescent dye.
30 Any laser-induced emission from the reporter dye is quenched by the quenching dye when the two dyes are located close together as they are on the probe. During the amplification reaction, the Taq DNA polymerase enzyme cleaves the probe in a template-dependent manner. The resultant probe fragments disassociate in solution, and signal from the released reporter dye is free from the quenching effect of the second fluorophore.
35 One molecule of reporter dye is liberated for each new molecule synthesized, and detection of the unquenched reporter dye provides the basis for quantitative interpretation of the data.

TaqMan RT-PCR can be performed using commercially available equipment, such as, for example, ABI PRISM 7700tam Sequence Detection System (Perkin-Elmer-Applied Biosystems, Foster City, CA, USA), or Lightcycler (Roche Molecular Biochemicals, Mannheim, Germany). In a preferred embodiment, the 5' nuclease procedure is run on a real-time quantitative PCR device such as the ABI PRISM 7700tam Sequence Detection System. The system consists of a thermocycler, laser, charge-coupled device (CCD), camera, and computer. The system amplifies samples in a 96-well format on a thermocycler. During amplification, laser-induced fluorescent signal is collected in real-time through fibre optics cables for all 96 wells, and detected at the CCD. The system includes software for running the instrument and for analyzing the data.

5' nuclease assay data are initially expressed as Ct, or the threshold cycle. As discussed above, fluorescence values are recorded during every cycle and represent the amount of product amplified to that point in the amplification reaction. The point when the fluorescent signal is first recorded as statistically significant is the threshold cycle.

To minimize errors and the effect of sample-to-sample variation, RT-PCR is usually performed using an internal standard. The ideal internal standard is expressed at a constant level among different tissues, and is unaffected by the experimental treatment. RNAs most frequently used to normalize patterns of gene expression are mRNAs for the housekeeping genes glyceraldehyde-3-phosphate-dehydrogenase (GAPDH) and actin.

Real-time quantitative PCR (qPCR)

A more recent variation of the RT-PCR technique is the real time quantitative PCR, which measures PCR product accumulation through a dual-labeled fluorogenic probe (i.e., TaqMan@ probe). Real time PCR is compatible both with quantitative competitive PCR and with quantitative comparative PCR. The former uses an internal competitor for each target sequence for normalization, while the latter uses a normalization gene contained within the sample, or a housekeeping gene for RT-PCR. For further details see, e.g., Held et al., Genome Research 6: 986-994 (1996).

Expression levels can be determined using fixed, paraffin-embedded tissues as the RNA source. According to one aspect of the present invention, PCR primers and probes are designed based upon intron sequences present in the gene to be amplified. In this embodiment, the first step in the primer/probe design is the delineation of intron sequences within the genes. This can be done by publicly available software, such as the

DNA BLAT software developed by Kent, W. J., *Genome Res.* 12 (4): 656-64 (2002), or by the BLAST software including its variations. Subsequent steps follow well established methods of PCR primer and probe design.

- 5 In order to avoid non-specific signals, it is useful to mask repetitive sequences within the introns when designing the primers and probes. This can be easily accomplished by using the Repeat Masker program available on-line through the Baylor College of Medicine, which screens DNA sequences against a library of repetitive elements and returns a query sequence in which the repetitive elements are masked. The masked sequences can then
10 be used to design primer and probe sequences using any commercially or otherwise publicly available primer/probe design packages, such as Primer Express (Applied Biosystems); MGB assay-by-design (Applied Biosystems); Primer3 (Steve Rozen and Helen J. Skaletsky (2000) Primer3 on the WWW for general users and for biologist programmers in: Krawetz S, Misener S (eds) *Bioinformatics Methods and Protocols*:
15 *Methods in Molecular Biology*. Humana Press, Totowa, NJ, pp 365-386).

The most important factors considered in PCR primer design include primer length, melting temperature (T_m), and G/C content, specificity, complementary primer sequences, and 3' end sequence. In general, optimal PCR primers are generally 17-30 bases in
20 length, and contain about 20-80%, such as, for example, about 50-60% G+C bases. T_m s between 50 and 80°C, e.g., about 50 to 70°C are typically preferred. For further guidelines for PCR primer and probe design see, e.g., Dieffenbach, C. W. et al., *General Concepts for PCR Primer Design* in: *PCR Primer, A Laboratory Manual*, Cold Spring Harbor Laboratory Press, New York, 1995, pp. 133-155; Innis and Gelfand, *Optimization of PCRs*
25 in: *PCR Protocols, A Guide to Methods and Applications*, CRC Press, London, 1994, pp. 5-11; and Plasterer, T. N. *Primerselect: Primer and probe design*. *Methods Mol. Biol.* 70: 520-527 (1997), the entire disclosures of which are hereby expressly incorporated by reference.

30 **Microarray analysis**

Differential gene expression can also be identified, or confirmed using the microarray technique. Thus, the expression profile of GCPMs can be measured in either fresh or paraffin-embedded tumour tissue, using microarray technology. In this method, polynucleotide sequences of interest (including cDNAs and oligonucleotides) are plated,
35 or arrayed, on a microchip substrate. The arrayed sequences (i.e., capture probes) are then hybridized with specific polynucleotides from cells or tissues of interest (i.e., targets). Just as in the RT-PCR method, the source of RNA typically is total RNA isolated from

human tumours or tumour cell lines, and corresponding normal tissues or cell lines. Thus RNA can be isolated from a variety of primary tumours or tumour cell lines. If the source of RNA is a primary tumour, RNA can be extracted, for example, from frozen or archived paraffin-embedded and fixed (e.g., formalin-fixed) tissue samples, which are routinely prepared and preserved in everyday clinical practice.

In a specific embodiment of the microarray technique, PCR amplified inserts of cDNA clones are applied to a substrate. The substrate can include up to 1, 2, 3, 4, 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, or 75 nucleotide sequences. In other aspects, the substrate can include at least 10,000 nucleotide sequences. The microarrayed sequences, immobilized on the microchip, are suitable for hybridization under stringent conditions. As other embodiments, the targets for the microarrays can be at least 50, 100, 200, 400, 500, 1000, or 2000 bases in length; or 50-100, 100-200, 100-500, 100-1000, 100-2000, or 500-5000 bases in length. As further embodiments, the capture probes for the microarrays can be at least 10, 15, 20, 25, 50, 75, 80, or 100 bases in length; or 10-15, 10-20, 10-25, 10-50, 10-75, 10-80, or 20-80 bases in length.

Fluorescently labeled cDNA probes may be generated through incorporation of fluorescent nucleotides by reverse transcription of RNA extracted from tissues of interest. Labeled cDNA probes applied to the chip hybridize with specificity to each spot of DNA on the array. After stringent washing to remove non-specifically bound probes, the chip is scanned by confocal laser microscopy or by another detection method, such as a CCD camera. Quantitation of hybridization of each arrayed element allows for assessment of corresponding mRNA abundance. With dual colour fluorescence, separately labeled cDNA probes generated from two sources of RNA are hybridized pairwise to the array. The relative abundance of the transcripts from the two sources corresponding to each specified gene is thus determined simultaneously.

The miniaturized scale of the hybridization affords a convenient and rapid evaluation of the expression pattern for large numbers of genes. Such methods have been shown to have the sensitivity required to detect rare transcripts, which are expressed at a few copies per cell, and to reproducibly detect at least approximately two-fold differences in the expression levels (Scherena et al., Proc. Natl. Acad. Sci. USA 93 (2): 106-149 (1996)). Microarray analysis can be performed by commercially available equipment, following manufacturer's protocols, such as by using the Affymetrix GenChip technology, or Incyte's microarray technology. The development of microarray methods for large-scale analysis

of gene expression makes it possible to search systematically for molecular markers of cancer classification and outcome prediction in a variety of tumour types.

RNA isolation, purification, and amplification

5 General methods for mRNA extraction are well known in the art and are disclosed in standard textbooks of molecular biology, including Ausubel et al., *Current Protocols of Molecular Biology*, John Wiley and Sons (1997). Methods for RNA extraction from paraffin embedded tissues are disclosed, for example, in Rupp and Locker, *Lab Invest.* 56: A67 (1987), and De Sandres et al., *BioTechniques* 18: 42044 (1995). In particular, RNA
10 isolation can be performed using purification kit, buffer set, and protease from commercial manufacturers, such as Qiagen, according to the manufacturer's instructions. For example, total RNA from cells in culture can be isolated using Qiagen RNeasy mini-columns. Other commercially available RNA isolation kits include MasterPure Complete DNA and RNA Purification Kit (EPICENTRE (D, Madison, WI), and Paraffin Block RNA
15 Isolation Kit (Ambion, Inc.). Total RNA from tissue samples can be isolated using RNA Stat-60 (Tel-Test). RNA prepared from tumour can be isolated, for example, by cesium chloride density gradient centrifugation.

The steps of a representative protocol for profiling gene expression using fixed, paraffin-
20 embedded tissues as the RNA source, including mRNA isolation, purification, primer extension and amplification are given in various published journal articles (for example: T. E. Godfrey et al. *J. Molec. Diagnostics* 2: 84-91 (2000); K. Specht et al., *Am. J. Pathol.* 158: 419-29 (2001)). Briefly, a representative process starts with cutting about 10 μ m thick sections of paraffin-embedded tumour tissue samples. The RNA is then extracted, and
25 protein and DNA are removed. After analysis of the RNA concentration, RNA repair and/or amplification steps may be included, if necessary, and RNA is reverse transcribed using gene specific promoters followed by RT-PCR. Finally, the data are analyzed to identify the best treatment option(s) available to the patient on the basis of the characteristic gene expression pattern identified in the tumour sample examined.

30

Immunohistochemistry and proteomics

Immunohistochemistry methods are also suitable for detecting the expression levels of the proliferation markers of the present invention. Thus, antibodies or antisera, preferably polyclonal antisera, and most preferably monoclonal antibodies specific for each marker,
35 are used to detect expression. The antibodies can be detected by direct labeling of the antibodies themselves, for example, with radioactive labels, fluorescent labels, hapten labels such as, biotin, or an enzyme such as horse radish peroxidase or alkaline

phosphatase. Alternatively, unlabeled primary antibody is used in conjunction with a labeled secondary antibody, comprising antisera, polyclonal antisera or a monoclonal antibody specific for the primary antibody. Immunohistochemistry protocols and kits are well known in the art and are commercially available.

5

Proteomics can be used to analyze the polypeptides present in a sample (e.g., tissue, organism, or cell culture) at a certain point of time. In particular, proteomic techniques can be used to assess the global changes of protein expression in a sample (also referred to as expression proteomics). Proteomic analysis typically includes: (1) separation of individual
10 proteins in a sample by 2-D gel electrophoresis (2-D PAGE); (2) identification of the individual proteins recovered from the gel, e.g., by mass spectrometry or N-terminal sequencing, and (3) analysis of the data using bioinformatics. Proteomics methods are valuable supplements to other methods of gene expression profiling, and can be used, alone or in combination with other methods, to detect the products of the proliferation
15 markers of the present invention.

Selection of Differentially Expressed Genes.

An early approach to the selection of genes deemed significant involved simply looking at the "fold change" of a given gene between the two groups of interest. While this approach
20 hones in on genes that seem to change the most spectacularly, consideration of basic statistics leads one to realize that if the variance (or noise level) is quite high (as is often seen in microarray experiments), then seemingly large fold-change can happen frequently by chance alone.

25 Microarray experiments, such as those described here, typically involve the simultaneous measurement of thousands of genes. If one is comparing the expression levels for a particular gene between two groups (for example recurrent and non-recurrent tumours), the typical tests for significance (such as the t-test) are not adequate. This is because, in an ensemble of thousands of experiments (in this context each gene constitutes an
30 "experiment"), the probability of at least one experiment passing the usual criteria for significance by chance alone is essentially unity. In a test for significance, one typically calculates the probability that the "null hypothesis" is correct. In the case of comparing two groups, the null hypothesis is that there is no difference between the two groups. If a statistical test produces a probability for the null hypothesis below some threshold (usually
35 0.05 or 0.01), it is stated that we can reject the null hypothesis, and accept the hypothesis that the two groups are *significantly* different. Clearly, in such a test, a rejection of the null hypothesis by chance alone could be expected 1 in 20 times (or 1 in 100). The use of t-

tests, or other similar statistical tests for significance, fail in the context of microarrays, producing far too many false positives (or type I errors)

5 In this type of situation, where one is testing multiple hypotheses at the same time, one applies typical multiple comparison procedures, such as the Bonferroni Method (43). However such tests are too conservative for most microarray experiments, resulting in too many false negative (type II) errors.

10 A more recent approach is to do away with attempting to apply a probability for a given test being significant, and establish a means for selecting a subset of experiments, such that the expected proportion of Type I errors (or false discovery rate; 47) is controlled for. It is this approach that has been used in this investigation, through various implementations, namely the methods provided with BRB Array Tools (48), and the limma (11,42) package of Bioconductor (that uses the R statistical environment; 10,39).

15

General methodology for Data Mining: Generation of Prognostic Signatures

20 Data Mining is the term used to describe the extraction of "knowledge", in other words the "know-how", or predictive ability from (usually) large volumes of data (the dataset). This is the approach used in this study to generate prognostic signatures. In the case of this study the "know-how" is the ability to accurately predict prognosis from a given set of gene expression measurements, or "signature" (as described generally in this section and in more detail in the examples section).

25

The specific details used for the methods used in this study are described in Examples 17-20. However, application of any of the data mining methods (both those described in the Examples, and those described here) can follow this general protocol.

30 Data mining (49), and the related topic machine learning (40) is a complex, repetitive mathematical task that involves the use of one or more appropriate computer software packages (see below). The use of software is advantageous on the one hand, in that one does not need to be completely familiar with the intricacies of the theory behind each technique in order to successfully use data mining techniques, provided that one adheres
35 to the correct methodology. The disadvantage is that the application of data mining can often be viewed as a "black box": one inserts the data and receives the answer. How this is achieved is often masked from the end-user (this is the case for many of the techniques

described, and can often influence the statistical method chosen for data mining. For example, neural networks and support vector machines have a particularly complex implementation that makes it very difficult for the end user to extract out the "rules" used to produce the decision. On the other hand, k-nearest neighbours and linear discriminant analysis have a very transparent process for decision making that is not hidden from the user.

There are two types of approach used in data mining: supervised and unsupervised approaches. In the supervised approach, the information that is being linked to the data is known, such as categorical data (e.g. recurrent vs. non recurrent tumours). What is required is the ability to link the observed response (e.g. recurrence vs. non-recurrence) to the input variables. In the unsupervised approach, the classes within the dataset are not known in advance, and data mining methodology is employed to attempt to find the classes or structure within the dataset.

In the present example the supervised approach was used and is discussed in detail here, although it will be appreciated that any of the other techniques could be used.

The overall protocol involves the following steps:

- Data representation. This involves transformation of the data into a form that is most likely to work successfully with the chosen data mining technique. In where the data is numerical, such as in this study where the data being investigated represents relative levels of gene expression, this is fairly simple. If the data covers a large dynamic range (i.e. many orders of magnitude) often the log of the data is taken. If the data covers many measurements of separate samples on separate days by separate investigators, particular care has to be taken to ensure systematic error is minimised. The minimisation of systematic error (i.e. errors resulting from protocol differences, machine differences, operator differences and other quantifiable factors) is the process referred to here as "normalisation".
- Feature Selection. Typically the dataset contains many more data elements than would be practical to measure on a day-to-day basis, and additionally many elements that do not provide the information needed to produce a prediction model. The actual ability of a prediction model to describe a dataset is derived from some subset of the full dimensionality of the dataset. These dimensions the most important components (or features) of the dataset. Note in the context of

microarray data, the dimensions of the dataset are the individual genes. Feature selection, in the context described here, involves finding those genes which are most "differentially expressed". In a more general sense, it involves those groups which pass some statistical test for significance, i.e. is the level of a particular variable consistently higher or lower in one or other of the groups being investigated. Sometimes the features are those variables (or dimensions) which exhibit the greatest variance.

The application of feature selection is completely independent of the method used to create a prediction model, and involves a great deal of experimentation to achieve the desired results. Within this invention, the selection of significant genes, and those which correlated with the earlier successful model (the NZ classifier), entailed feature selection. In addition, methods of data reduction (such as principal component analysis) can be applied to the dataset.

- Training. Once the classes (e.g. recurrence/non-recurrence) and the features of the dataset have been established, and the data is represented in a form that is acceptable as input for data mining, the reduced dataset (as described by the features) is applied to the prediction model of choice. The input for this model is usually in the form a multi-dimensional numerical input, (known as a vector), with associated output information (a class label or a response). In the training process, selected data is input into the prediction model, either sequentially (in techniques such as neural networks) or as a whole (in techniques that apply some form of regression, such as linear models, linear discriminant analysis; support vector machines). In some instances (e.g. k-nearest neighbours) the dataset (or subset of the dataset obtained after feature selection) is itself the model. As discussed, effective models can be established with minimal understanding of the detailed mathematics, through the use of various software packages where the parameters of the model have been pre-determined by expert analysts as most likely to lead to successful results.

- Validation. This is a key component of the data-mining protocol, and the incorrect application of this frequently leads to errors. Portions of the dataset are to be set aside, apart from feature selection and training, to test the success of the prediction model. Furthermore, if the results of validation are used to effect feature selection and training of the model, then one obtains a further validation set to test the model before it is applied to real-life situations. If this process is not strictly

adhered to the model is likely to fail in real-world situations. The methods of validation are described in more detail below.

- Application. Once the model has been constructed, and validated, it must be packaged in some way as it is accessible to end users. This often involves implementation of some form a spreadsheet application, into which the model has been imbedded, scripting of a statistical software package, or refactoring of the model into a hard-coded application by information technology staff.

Examples of software packages that are frequently used are:

- Spreadsheet plugins, obtained from multiple vendors.
- The R statistical environment.
- The commercial packages MatLab, S-plus, SAS, SPSS, STATA.
- Free open-source software such as Octave (a MatLab clone)
- many and varied C++ libraries, which can be used to implement prediction models in a commercial, closed-source setting.

Examples of Data Mining Methods.

The methods can be by first performing the step of data mining process (above), and then applying the appropriate known software packages. Further description of the process of data mining is described in detail in many extremely well-written texts.(49)

- Linear models (49, 50): The data is treated as the input of a linear regression model, of which the class labels or responses variables are the output. Class labels, or other categorical data, must be transformed into numerical values (usually integer). In generalised linear models, the class labels or response variables are not themselves linearly related to the input data, but are transformed through the use of a "link function". Logistic regression is the most common form of generalized linear model.

- Linear Discriminant analysis (49, 51, 52). Provided the data is linearly separable (i.e. the groups or classes of data can be separated by a hyperplane, which is an n-dimensional extension of a threshold), this technique can be applied. A combination of variables is used to separate the classes, such that the between group variance is maximised, and the within-group variance is minimised. The byproduct of this is the formation of a classification rule. Application of this rule to samples of unknown class allows predictions or classification of class membership

to be made for that sample. There are variations of linear discriminant analysis such as nearest shrunken centroids which are commonly used for microarray analysis.

- Support vector machines (53): A collection of variables is used in conjunction with a collection of weights to determine a model that maximizes the separation between classes in terms of those weighted variables. Application of this model to a sample then produces a classification or prediction of class membership for that sample.

10

- Neural networks (52): The data is treated as input into a network of nodes, which superficially resemble biological neurons, which apply the input from all the nodes to which they are connected, and transform the input into an output. Commonly, neural networks use the "multiply and sum" algorithm, to transform the inputs from multiple connected input nodes into a single output. A node may not necessarily produce an output unless the inputs to that node exceed a certain threshold. Each node has as its input the output from several other nodes, with the final output node usually being linked to a categorical variable. The number of nodes, and the topology of the nodes can be varied in almost infinite ways, providing for the ability to classify extremely noisy data that may not be possible to categorize in other ways. The most common implementation of neural networks is the multi-layer perceptron.

15

20

- Classification and regression trees (54): In these, variables are used to define a hierarchy of rules that can be followed in a stepwise manner to determine the class of a sample. The typical process creates a set of rules which lead to a specific class output, or a specific statement of the inability to discriminate. A example classification tree is an implementation of an algorithm such as:

25

30

```
if gene A > x and gene Y > x and gene Z = z
then
  class A
else if geneA = q
then
  class B
```

35

- Nearest neighbour methods (51, 52). Predictions or classifications are made by comparing a sample (of unknown class) to those around it (or known class), with

5 closeness defined by a distance function. It is possible to define many different distance functions. Commonly used distance functions are the Euclidean distance (an extension of the Pythagorean distance, as in triangulation, to n-dimensions), various forms of correlation (including Pearson Correlation co-efficient). There are also transformation functions that convert data points that would not normally be interconnected by a meaningful distance metric into euclidean space, so that Euclidean distance can then be applied (e.g. Mahalanobis distance). Although the distance metric can be quite complex, the basic premise of k-nearest neighbours is quite simple, essentially being a restatement of "find the k-data vectors that are most similar to the unknown input, find out which class they correspond to, and vote as to which class the unknown input is".

- **Other methods:**

- Bayesian networks. A directed acyclic graph is used to represent a collection of variables in conjunction with their joint probability distribution, which is then used to determine the probability of class membership for a sample.
- Independent components analysis, in which independent signals (e.g., class membership) re isolated (into components) from a collection of variables. These components can then be used to produce a classification or prediction of class membership for a sample.

Ensemble learning methods in which a collection of prediction methods are combined to produce a joint classification or prediction of class membership for a sample

25 There are many variations of these methodologies that can be explored (49), and many new methodologies are constantly being defined and developed. It will be appreciated that any one of these methodologies can be applied in order to obtain an acceptable result. Particular care must be taken to avoid overfitting, by ensuring that all results are tested via a comprehensive validation scheme.

Validation

Application of any of the prediction methods described involves both training and cross-validation (43, 55) before the method can be applied to new datasets (such as data from a clinical trial). Training involves taking a subset of the dataset of interest (in this case gene expression measurements from colorectal tumours), such that it is stratified across the classes that are being tested for (in this case recurrent and non-recurrent

tumours). This training set is used to generate a prediction model (defined above), which is tested on the remainder of the data (the testing set).

It is possible to alter the parameters of the prediction model so as to obtain better performance in the testing set, however, this can lead to the situation known as overfitting, where the prediction model works on the training dataset but not on any external dataset. In order to circumvent this, the process of validation is followed. There are two major types of validation typically applied, the first (hold-out validation) involves partitioning the dataset into three groups: testing, training, and validation. The validation set has no input into the training process whatsoever, so that any adjustment of parameters or other refinements must take place during application to the testing set (but not the validation set). The second major type is cross-validation, which can be applied in several different ways, described below.

There are two main sub-types of cross-validation: K-fold cross-validation, and leave-one-out cross-validation

K-fold cross-validation: The dataset is divided into K subsamples, each subsample containing approximately the same proportions of the class groups as the original. In each round of validation, one of the K subsamples is set aside, and training is accomplished using the remainder of the dataset. The effectiveness of the training for that round is gauged by how correctly the classification of the left-out group is. This procedure is repeated K -times, and the overall effectiveness ascertained by comparison of the predicted class with the known class.

25

Leave-one-out cross-validation: A commonly used variation of K-fold cross validation, in which $K=n$, where n is the number of samples.

Combinations of CCPMS, such as those described above in Tables 1 and 2, can be used to construct predictive models for prognosis.

30

Prognostic Signatures

Prognostic signatures, comprising one or more of these markers, can be used to determine the outcome of a patient, through application of one or more predictive models derived from the signature. In particular, a clinician or researcher can determine the differential expression (e.g., increased or decreased expression) of the one or more markers in the signature, apply a predictive model, and thereby predict the negative

35

prognosis, e.g., likelihood of disease relapse, of a patient, or alternatively the likelihood of a positive prognosis (continued remission).

5 In still further aspects, the invention includes a method of determining a treatment regime for a cancer comprising: (a) providing a sample of the cancer; (b) detecting the expression level of a GgCPM family member in said sample; (c) determining the prognosis of the cancer based on the expression level of a CCPM family member; and (d) determining the treatment regime according to the prognosis.

10 In still further aspects, the invention includes a device for detecting a GCPM, comprising: a substrate having a GCPM capture reagent thereon; and a detector associated with said substrate, said detector capable of detecting a GCPM associated with said capture reagent. Additional aspects include kits for detecting cancer, comprising: a substrate; a GCPM capture reagent; and instructions for use. Yet further aspects of the invention
15 include method for detecting aGCPM using qPCR, comprising: a forward primer specific for said CCPM; a reverse primer specific for said GCPM; PCR reagents; a reaction vial; and instructions for use.

Additional aspects of this invention comprise a kit for detecting the presence of a GCPM
20 polypeptide or peptide, comprising: a substrate having a capture agent for said GCPM polypeptide or peptide; an antibody specific for said GCPM polypeptide or peptide; a reagent capable of labeling bound antibody for said GCPM polypeptide or peptide; and instructions for use.

25 In yet further aspects, this invention includes a method for determining the prognosis of colorectal cancer, comprising the steps of: providing a tumour sample from a patient suspected of having colorectal cancer; measuring the presence of a GCPM polypeptide using an ELISA method. In specific aspects of this invention the GCPM of the invention is selected from the markers set forth in Table A, Table B, Table C or Table D. In still further
30 aspects, the GCPM is included in a prognostic signature

While exemplified herein for gastrointestinal cancer, e.g., gastric and colorectal cancer, the GCPMs of the invention also find use for the prognosis of other cancers, e.g., breast
35 cancers, prostate cancers, ovarian cancers, lung cancers (such as adenocarcinoma and, particularly, small cell lung cancer), lymphomas, gliomas, blastomas (e.g., medulloblastomas), and mesothelioma, where decreased or low expression is associated

with a positive prognosis, while increased or high expression is associated with a negative prognosis.

EXAMPLES

5 The examples described herein are for purposes of illustrating embodiments of the invention. Other embodiments, methods, and types of analyses are within the scope of persons of ordinary skill in the molecular diagnostic arts and need not be described in detail hereon. Other embodiments within the scope of the art are considered to be part of this invention.

10

EXAMPLE 1: Cell cultures

The experimental scheme is shown in FIG. 1. Ten colorectal cell lines were cultured and harvested at semi- and full-confluence. Gene expression profiles of the two growth stages were analyzed on 30,000 oligonucleotide arrays and a gene proliferation signature (GPS; 15 Table C) was identified by gene ontology analysis of differentially expressed genes. Unsupervised clustering was then used to independently dichotomize two cohorts of clinical colorectal samples (Cohort A: 73 stage I-IV on oligo arrays, Cohort B: 55 stage II on Affymetrix chips) based on the similarities of the GPS expression. Ki-67 immunostaining was also performed on tissue sections from Cohort A tumours. Following 20 this, the correlation between proliferation activity and clinico-pathologic parameters was investigated.

Ten colorectal cancer cell lines derived from different disease stages were included in this study: DLD-1, HCT-8, HCT-116, HT-29, LoVo, Ls174T, SK-CO-1, SW48, SW480, and 25 SW620 (ATCC, Manassas, VA). Cells were cultivated in a 5% CO₂ humidified atmosphere at 37°C in alpha minimum essential medium supplemented with 10% fetal bovine serum, 100 IU/ml penicillin and 100 µg/ml streptomycin (GIBCO-Invitrogen, CA). Two cell cultures were established for each cell line. The first culture was harvested upon reaching semi-confluence (50-60%). When cells in the second culture reached full-confluence 30 (determined both microscopically and macroscopically), media was replaced, and cells were harvested twenty-four hours later to prepare RNA from the growth-inhibited cells. Array experiments were carried out on RNA extracted from each cell culture. In addition, a second culturing experiment was done following the same procedure and extracted RNA was used for dye-reversed hybridizations.

35

EXAMPLE 2: Patients

Two cohorts of patients were analysed. Cohort A included 73 New Zealand colorectal cancer patients who underwent surgery at Dunedin and Auckland hospitals between 1995 and 2000. These patients were part of a prospective cohort study and included all disease stages. Tumour samples were collected fresh from the operation theatre, snap frozen in liquid nitrogen and stored at -80°C. Specimens were reviewed by a single pathologist (H-S Y) and tumours were staged according to the TNM system (34). Of the 73 patients, 32 developed disease recurrence and 41 remained recurrence-free after a minimum of five years follow up. The median overall survival was 29.5 and 66 months for recurrent and recurrent-free patients, respectively. Twenty patients received 5-FU-based post-operative adjuvant chemotherapy and 12 patients received radiotherapy (7 pre- and 5 post-operative).

Cohort B included a group of 55 German colorectal patients who underwent surgery at the Technical University of Munich between 1995 and 2001 and had fresh frozen samples stored in a tissue bank. All 55 had stage II disease, 26 developed disease recurrence (median survival 47 months) and 29 remained recurrence-free (median survival 82 months). None of patients received chemotherapy or radiotherapy. Clinico-pathologic variables of both cohorts are summarised as part of Table 2.

20

Table 2: Clinico-pathologic parameters and their association with the GPS expression and Ki-67 PI

Parameters	Number of patients		GPS		Ki-67 PI*	
	cohort A	cohort B	cohort A (p-value) [§]	cohort B (p-value) [§]	Mean ± SD	p-value [§]
Age [†] < Mean	34	31	1	0.79	74.4±17.9	0.6
> Mean	39	24			77.9±17.3	
Sex Male	35	33	0.16	1	77.3±15.3	1
Female	38	22			75.3±19.5	
Site [‡] Right side	30	12	1	0.2	80.4±13.3	0.2
Left side	43	43			73.1±19.7	
Grade Well	9	0	0.22	0.2	75.6±18.1	
Moderate	50	33			73.9±18.9	0.98
Poor	14	22			84.3±9.3	
Dukes stage A	10	0	0.006	NA	78.8±17.3	0.73
B	27	55			75.7±18.4	
C	28	0			76±16.1	
D	8	0			75.9±22	
T stage T1	5	0	0.16	0.62	71.3±22.4	0.16
T2	11	11			85.4±7.4	
T3	50	41			76±17	
T4	7	3			66.2±26.3	
N stage N0	38	55	0.03	NA	76.5±17.9	1
N1+N2	35	0			76±17.4	
Vascular invasion Yes	5	1	0.67	NA	54.4±31.5	0.32
No	68	54			78±15	
Lymphatic invasion Yes	32	5	0.06	0.35	76.5±18.3	0.6
No	41	50			75.1±17.3	
Lymphocyte infiltration Mild	35	15	0.89	1	75±18.6	0.85
Moderate	27	25			79.4±16.5	
Prominent	11	15			73.5±18.3	
Margin Infiltrative	45	NA	0.47	NA	75.8±18.9	1
Expansive	28				77.1±15.7	
Recurrence Yes	32	26	0.03	<0.001	75.6±19	0.79
No	41	29			76.8±16.2	
Total	73	55			76.3±17.5	

§ A Fisher's Exact Test or Kruskal-Wallis Test were used for testing association between clinico-pathologic parameters and GPS expression or Ki-67 PI, as appropriate.
* Ki-67 immunostaining was performed on tumor sections from cohort A patients.
[‡] Proximal and distal to splenic flexure, respectively
[†] Average age 68 and 63 years for cohort A and B patients, respectively
NA: not applicable

5 EXAMPLE 3: Array preparation and gene expression analysis

Cohort A tumours and cell lines: Tissue samples and cell lines were homogenised and RNA was extracted using Tri-Reagent (Progenz, Auckland, NZ). The RNA was then purified using RNeasy mini column (Qiagen, Victoria, Australia) according to the manufacture's protocol. Ten micrograms of total RNA extracted from each culture or

10 tumour sample was oligo-dT primed and cDNA synthesis was carried out in the presence of aa-dUTP and Superscript II RNase H-Reverse Transcriptase (Invitrogen). Cy dyes were incorporated into cDNA using the indirect amino-allyl cDNA labelling method. cDNA derived from a pool of 12 different cell lines was used as the reference for all hybridizations. The Cy5-dUTP-tagged cDNA from an individual colorectal cell line or

15 tissue sample was combined with Cy3-dUTP-tagged cDNA from reference sample. The

mixture was then purified using a QiaQuick PCR purification Kit (Qiagen, Victoria, Australia) and co-hybridized to a microarray spotted with the MWG 30K Oligo Set (MWG Biotech, NC). cDNA samples from the second culturing experiment were additionally analysed on microarrays using reverse labelling.

5

Arrays were scanned with a GenePix 4000B Microarray Scanner and data were analysed using GenePix Pro 4.1 Microarray Acquisition and Analysis Software (Axon, CA). The foreground intensities from each channel were $\log_2$ transformed and normalised using the SNOMAD software (35). Normalised values were collated and filtered using BRB-Array Tools Version 3.2 (developed by Dr. Richard Simon and Amy Peng Lam, Biometric Research Branch, National Cancer Institute). Low intensity genes, and genes for which over 20% of measurements across tissue samples or cell lines were missing, were excluded from further analysis.

10

15 Cohort B tumours: Total RNA was extracted from each tumour using RNeasy Mini Kit and purified on RNeasy Columns (Qiagen, Hilden, Germany). Ten micrograms of total RNA was used to synthesize double-stranded cDNA with SuperScript II reverse transcriptase (GIBCO-Invitrogen, NY) and an oligo-dT-T7 primer (Eurogentec, Koeln, Germany). Biotinylated cRNA was synthesized from the double-stranded cDNA using the Promega RiboMax T7-kit (Promega, Madison, WI) and Biotin-NTP labelling mix (Loxo, Dossenheim, Germany). Then, the biotinylated cRNA was purified and fragmented. The fragmented cRNA was hybridized to Affymetrix HGU133A GeneChips (Affymetrix, Santa Clara, CA) and stained with streptavidin-phycoerythrin. The arrays were then scanned with a HP-argon-ion laser confocal microscope and the digitized image data were processed using the Affymetrix® Microarray Suite 5.0 Software. All Affymetrix U133A GeneChips passed quality control to eliminate scans with abnormal characteristics. Background correction and normalization were performed in the R computing environment using the robust multi-array average function implemented in the Bioconductor package affy.

20

25

30 **EXAMPLE 4: Quantitative real-time PCR (QPCR)**

The expression of eleven genes (MAD2L1, POLE2, CDC2, MCM6, MCM7, RANSEH2A, TOPK, KPNA2, G22P1, PCNA, and GMNN) was validated using the cDNA from the cell cultures. Total RNA (2 μ g) was reverse transcribed using Superscript II RNase H-Reverse Transcriptase kit (Invitrogen) and oligo dT primer (Invitrogen). QPCR was performed on an ABI Prism 7900HT Sequence Detection System (Applied Biosystems) using Taqman Gene Expression Assays (Applied Biosystems). Relative fold changes were calculated

35

using the $2^{-\Delta\Delta CT}$ method³⁶ with Topoisomerase 3A as the internal control. Reference RNA was used as the calibrator to enable comparison between different experiments.

EXAMPLE 5: Immunohistochemical analysis

5 Immunohistochemical expression of Ki-67 antigen (MIB-1; DakoCytomation, Denmark) was investigated on 4 μ m sections of 73 paraffin-embedded primary colorectal tumours from Cohort A. Endogenous peroxidase activity was blocked with 0.3% hydrogen peroxidase in methanol and antigens were retrieved in boiling citrate buffer (pH 6). Non-specific binding sites were blocked with 5% normal goat serum containing 1% BSA.
10 Primary antibody (dilution 1:50) was detected using the EnVision system (Dako EnVision, CA) and the DAB substrate kit (Vector laboratories, CA). Five high-power fields were selected using a 10 x 10 microscope grid and cell counts were performed manually in a blind fashion without knowledge of the clinico-pathologic data. The Ki-67 proliferation index (PI) was presented as the percentage of positively stained nuclei for each tumour.

15

EXAMPLE 6: Statistical analysis

Statistical analyses were performed using SPSS® version 14.0.0 (SPSS Inc., Chicago, IL). Ki-67 proliferation indices were presented as mean $\pm$ SD. A Fisher's Exact Test or Kruskal-Wallis Test was used to evaluate the differences between categorized groups
20 based on the expression of the GPS or the Ki-67 PI versus the clinico-pathologic parameters. A P value $\leq$ 0.05 was considered significant. Overall survival (OS) and recurrence-free survival (RFS) were plotted using the method of Kaplan and Meier (37). A log-rank test was used to test for differences in survival time between the categorized groups. Relative risk and associated confidence intervals were also estimated for each
25 variable using the Cox univariate model, and a multivariate Cox proportional hazard model was developed using forward stepwise regression with predictive variables that were significant in the univariate analysis. K-means clustering method was used to classify clinical samples based on the expression level of GPS.

30 EXAMPLE 7: Identification of a gene proliferation signature (GPS) using a colorectal cell line model

An overview of the approach used to derive and apply a gene proliferation signature (GPS) is summarised in FIG. 1. The GPS, including 38 mitotic cell cycle genes (Table C), was relatively over-expressed in cycling cells in semi-confluent cultures. Low proliferation,
35 defined by low GPS expression, was associated with unfavourable clinico-pathologic variables, shorter overall and recurrence-free survival ($p < 0.05$). No association was found between Ki-67 proliferation index and clinico-pathologic variables or clinical outcome.

Table C: GCPMs for cell proliferation signature

Unique ID	Average Fold change EP/SP	Gene Symbol	Gene Name	GenBank Acc. No.	Gene Aliases
A:05382	1.91	CDC2	cell division cycle 2, G1 to S and G2 to M	NM_001786, NM_033379	CDK1; MGC111195; DKFZp686L2022
B:8147	1.89	MCM6	MCM6 minichromosome maintenance deficient 6 (MIS5 homolog, S. pombe) (S. cerevisiae)	NM_005915	Mis5; P105MCM; MCG40308
A:00231	1.75	RPA3	replication protein A3, 14kDa	NM_002947	REPA3
B:7620	1.69	MCM7	MCM7 minichromosome maintenance deficient 7 (S. cerevisiae)	NM_005916, NM_182776	MCM2; CDC47; P85MCM; P1CDC47; PNAS-146; CDABP0042; P1.1-MCM3
A:03715	1.68	PCNA	proliferating cell nuclear antigen	NM_002592, NM_182649	MGC8367
B:9714	1.59	XRCC6	X-ray repair complementing defective repair in Chinese hamster cells 6 (Ku autoantigen, 70kDa)	NM_001469	ML8; KU70; TLAA; CTC75; CTCBF; G22P1
B:4036	1.56	KPNA2	karyopherin alpha 2 (RAG cohort 1, importin alpha 1)	NM_002266	QIP2; RCH1; IPOA1; SRP1alpha
A:05280	1.56	ANLN	anillin, actin binding protein	NM_018685	scra; Scraps; ANILLIN; DKFZp779A055
A:04760	1.52	APG7L	ATG7 autophagy related 7 homolog (S. cerevisiae)	NM_006395	GSA7; APG7L; DKFZp434N0735; ATG7
A:03912	1.52	PBK	PDZ binding kinase	NM_018492	SPK; TOPK; Nori-3; FLJ14385
A:03435	1.51	GMNN	geminin, DNA replication inhibitor	NM_015895	Gem; RP3-369A17.3
A:09802	1.51	RRM1	ribonucleotide reductase M1 polypeptide	NM_001033	R1; RR1; RIR1
A:09331	1.49	CDC45L	CDC45 cell division cycle 45-like (S.	NM_003504	CDC45; CDC45L2; PORC-PI-1

			cerevisiae)		
A:06387	1.46	MAD2L1	MAD2 mitotic arrest deficient-like 1 (yeast)	NM_002358	MAD2; HSMAD2
A:09169	1.45	RAN	RAN, member RAS oncogene family	NM_006325	TC4; Gsp1; ARA24
A:07296	1.43	DUT	dUTP pyrophosphatase	NM_001025248, NM_001025249, NM_001948	dUTPase; FLJ20622
B:3501	1.42	RRM2	ribonucleotide reductase M2 polypeptide	NM_001034	R2; RR2M
A:09842	1.41	CDK7	cyclin-dependent kinase 7 (MO15 homolog, Xenopus laevis, cdk-activating kinase)	NM_001799	CAK1; STK1; CDKN7; p39MO15
A:09724	1.40	MLH3	mutL homolog 3 (E. coli)	NM_001040108, NM_014381	HNPCC7; MGC138372
A:05648	1.39	SMC4	structural maintenance of chromosomes 4	NM_001002799, NM_001002800, NM_005496	CAPC; SMC4L1; hCAP-C
A:09436	1.39	SMC3	structural maintenance of chromosomes 3	NM_005445	BAM; BMH; HCAP; CSPG6; SMC3L1
A:02929	1.39	POLD2	polymerase (DNA directed), delta 2, regulatory subunit 50kDa	NM_006230	None
A:04680	1.38	POLE2	polymerase (DNA directed), epsilon 2 (p59 subunit)	NM_002692	DPE2
B:8449	1.38	BCCIP	BRCA2 and CDKN1A interacting protein	NM_016567, NM_078468, NM_078469	TOK-1
B:1035	1.37	GIN52	GIN5 complex subunit 2 (Psf2 homolog)	NM_016095	PSF2; Pfs2; HSPC037
B:7247	1.37	TREX1	three prime repair exonuclease 1	NM_016381, NM_032166, NM_033627, NM_033628, NM_033629, NM_130384	AGS1; DRN3; ATRIP; FLJ12343; DKFZp434J0310
A:09747	1.35	BUB3	BUB3 budding uninhibited by benzimidazoles 3 homolog (yeast)	NM_001007793, NM_004725	BUB3L; hBUB3
B:9065	1.32	FEN1	flap structure-	NM_004111	MF1; RAD2;

			specific endonuclease 1		FEN-1
B:2392	1.32	DBF4B	DBF4 homolog B (<i>S. cerevisiae</i>)	NM_025104, NM_145663	DRF1; ASKL1; FLJ13087; MGC15009
A:09401	1.31	PREI3	preimplantation protein 3	NM_015387, NM_199482	2C4D; MOB1; MOB3; CGI- 95; MGC12264
C:0921	1.30	CCNE1	cyclin E1	NM_001238, NM_057182	CCNE
A:10597	1.30	RPA1	replication protein A1, 70kDa	NM_002945	HSSB; RF-A; RP-A; REPA1; RPA70
A:02209	1.29	POLE3	polymerase (DNA directed), epsilon 3 (p17 subunit)	NM_017443	p17; YBL1; CHRA17; CHARAC17
A:09921	1.26	RFC4	replication factor C (activator 1) 4, 37kDa	NM_002916, NM_181573	A1; RFC37; MGC27291
A:08668	1.26	MCM3	MCM3 -minichromosome maintenance deficient 3 (<i>S.</i> <i>cerevisiae</i>)	NM_002388	HCC5; P1.h; RLFB; MGC1157; P1-MCM3
B:7793	1.25	CHEK1	CHK1 checkpoint homolog (<i>S.</i> <i>pombe</i>)	NM_001274	CHK1
A:09020	1.22	CCND1	cyclin D1	NM_053056	BCL1; PRAD1; U21B31; D11S287E
A:03486	1.22	CDC37	CDC37 cell division cycle 37 homolog (<i>S.</i> <i>cerevisiae</i>)	NM_007065	P50CDC37

The GPS was identified as a subset of genes whose expression correlates with CRC cell proliferation rate. Statistical Analysis of Microarray (SAM; Reference 38) was used to identify genes differentially expressed (DE) between exponentially growing (semi-confluent) and non-cycling (fully-confluent) CRC cell lines (FIG. 1, stage 1). To adjust for gene specific dye bias and other sources of variation, each culture set was analysed independently. Analyses were limited to 502 DE genes for which a significant expression difference was observed between two growth stages in both sets of cultures (false discovery rate < 1%). Gene Ontology (GO) analysis was carried out using EASE39 to identify the biological process categories that were significantly reflected in the DE genes.

Cell-proliferation related categories were over-represented mainly due to genes upregulated in exponentially growing cells. The mitotic cell cycle category (GO:0000278) was defined as the GPS because (i) this biological process was the most over-represented GO term (EASE score=5.5211); and (ii) all 38 mitotic cell cycle genes (Table C) were expressed at higher levels in rapidly growing compared to growth-inhibited cells. The expression of eleven genes from the GPS was assessed by QPCR and correlated with corresponding values obtained from the array data. Therefore, QPCR confirmed that elevated expression of the proliferation signature genes correlates with the increased proliferation in CRC cell lines (FIG. 5).

EXAMPLE 8: Classification of CRC samples according to the expression level of gene proliferation signature

In order to examine the relative proliferation state of CRC tumours and the utility of the GPS for clinical application, CRC tumours from two cohorts were stratified into two clusters based on the expression of GPS (FIG. 1, stage 2). Expression values of the 38 genes defining the GPS were first obtained from the microarray-generated expression profiles of tumours. Tumours from each cohort were then separately classified into two clusters (K=2) based on their GPS expression level similarities using K-means unsupervised clustering. Analysis of DE genes between two defined clusters using all filtered genes revealed that the GPS was contained within the list of genes upregulated in cluster 1 (FIG. 2A, upper panel) relative to cluster 2 (lower panel) in both cohorts. Thus, the tumours in cluster 1 are characterised by high GPS expression, while the tumours in cluster 2 are characterised by low GPS expression.

EXAMPLE 9: Low gene proliferation signature is associated with unfavourable clinico-pathologic variables

Table 2 summarises the association between GPS expression levels and clinico-pathologic variables. An association was observed between low proliferation activity, defined by low GPS expression, and an increased risk of recurrence in both cohorts (P=0.03 and <0.001 for Cohort A and B, respectively). In Cohort A, low GPS expression was also associated with a higher disease stage and lymph node metastasis (P=0.006 and 0.03 respectively). In addition, tumours with lymphatic invasion from Cohort A tended to be less proliferative than tumours without lymphatic invasion, albeit without reaching statistical significance (P=0.06). No association was found between the GPS expression level and tumour site, age, sex, degree of differentiation, T-stage, vascular invasion, degree of lymphocyte infiltration and tumour margin.

EXAMPLE 10: Gene proliferation signature predicts clinical outcome

To examine the performance of the GPS in predicting patient outcome, Kaplan-Meier survival analysis was used to compare RFS and OS between low and high GPS tumours (FIG. 3). All patients were censored at 60 months post-operation. In colorectal cancer Cohort A, OS and RFS were shorter in patients with low GPS expression (Log rank test $P=0.04$ and 0.01 , respectively). In colorectal cancer Cohort B, low GPS expression was also associated with decreased OS ($P=0.0004$) and RFS ($P=0.0002$). When the parameters predicting OS and RFS in univariate analysis were investigated in a multivariate model, disease stage was the only independent predictor of 5-year OS, while disease stage and T-stage were independent predictors of RFS in Cohort A. In Cohort B, low GPS expression and lymphatic invasion showed an independent contribution to both OS and RFS. If survival analysis was limited to Cohort B patients without lymphatic invasion, low GPS was still associated with shorter OS and RFS, confirming the independence of the GPS as a predictor. Analyses of single and multiple-variable associations with survival are summarized in Table 3.

Low GPS expression was also associated with decreased 5-year overall survival in patients with gastric cancer ($p=0.008$). A Kaplan-Meier survival plot comparing the overall survival of low and high GPS gastric tumours is shown in Fig. 4.

Table 3: Uni- and multivariate analysis of prognostic factors for OS and RFS in both cohorts

	Parameter s	Overall Survival				Recurrence-free Survival			
		Univariate analysis		Multivariate analysis §		Univariate analysis		Multivariate analysis §	
		Hazard ratio *	p-value	Hazard ratio *	p-value	Hazard ratio *	p-value	Hazard ratio *	p-value
Cohort A	Dukes stage	4.2 (2.4-7.4)	<0.001	4.2 (2.4-7.4)	<0.001	3.9 (2.1-7.2)	<0.001	3.5 (1.9-6.6)	<0.001
	T-stage	2.1 (1.2-3.8)	0.011	-	-	2.7 (1.4-5.2)	0.003	2.2 (1-5.1)	0.040
	N stage	4.4 (2-9.6)	<0.001	-	-	4.3 (1.8-10)	0.001	-	-
	Lymphatic invasion (+ vs. -)	0.16 (0.07-0.36)	<0.001	-	-	0.2 (0.09-0.43)	<0.001	-	-
	Margin (infiltrative vs. expansive)	4.3 (1.7-11.9)	0.002	-	-	3.7 (1.4-10.1)	0.008	-	-
	GPS expression (low vs. high)	0.46 (0.2-0.9)	0.037	-	-	0.33(0.14-0.78)	0.011	-	-
Cohort B	Lymphatic invasion (+ vs. -)	0.25 (0.08-0.78)	0.016	0.3 (0.09-0.9)	0.037	0.23 (0.08-0.63)	0.005	0.27 (0.1-0.77)	0.014
	GPS expression (low vs. high)	0.23 (0.06-0.81)	0.022	0.25 (0.07-0.89)	0.032	0.25 (0.09-0.67)	0.006	0.27 (0.1-0.73)	0.010
* Hazard ratio determined by Cox regression model; confidence interval=95%									
§ Final results of Cox regression analysis using a forward stepwise method (enter limit=0.05, remove limit=0.10)									

EXAMPLE 11: Ki-67 is not associated with clinico-pathologic variables or survival

Ki-67 immunostaining was performed on tissue sections from Cohort A tumours only as paraffin-embedded samples were unavailable for Cohort B (FIG. 1, stage 3). Nuclear staining was detected in all 73 CRC tumours. Ki-67 PI ranged from 25 to 96 %, with a mean value of 76.3 ± 17.5 . Using the mean Ki-67 value as a cut-off point, tumours were assigned into two groups with low or high PI. Ki-67 PI was neither associated with clinico-pathologic variables (Table 2) nor survival (FIG. 3). When the survival analysis was limited to the patients with the highest and lowest Ki-67 values, no statistical difference was observed (data not shown). The sum of these results indicates that the low expression of growth-related genes is associated with poor outcome in colorectal cancer, and Ki-67 was not sensitive enough to detect an association. These findings can be used as additional criteria for identifying patients at high risk of early death from cancer.

EXAMPLE 12: Selection of correlated cell proliferation genes

Cohort B (55 German CRC patients; Table 2) were first classified into low and high proliferation groups using the 38 gene cell proliferation signature (Table C) and the K-means clustering method (Pearson uncentered, 1000 permutations, threshold of occurrence in the same cluster sat at 80%). Statistical Analysis of Microarrays (SAM) was then applied to identify differentially expressed genes between low and high proliferation groups (FDR=0) when all filtered genes (16041 genes) were included for the analysis. 754 genes were found to be over-expressed in high proliferation group. The GATHER gene ontology program was then used to identify the most over-represented gene ontology categories within the list of differentially expressed genes. The cell cycle category was the most over-represented category within the list of differentially expressed genes. 102 cell cycle genes which are differentially expressed between the low and high proliferation groups (in addition to the original 38 gene signature) are shown in Table D.

Table D: Cell Cycle Genes that are Differentially Expressed in Low and High Proliferation

Gene Title	Gene Symbol	Chromosomal Location	Probe Set ID	Representative Public ID
asp (abnormal spindle) homolog, microcephaly associated (Drosophila)	ASPM	chr1q31	219918_s_at	NM_018123
aurora kinase A	AURKA	chr20q13.2-q13.3	204092_s_at 208079_s_at	NM_003600 NM_003158
aurora kinase B	AURKB	chr17p13.1	209464_at	AB011446
baculoviral IAP repeat-containing 5 (survivin)	BIRC5	chr17q25	202094_at 202095_s_at 210334_x_at	AA648913 NM_001168 AB028869
Bloom syndrome	BLM	chr15q26.1	205733_at	NM_000057
breast cancer 1, early onset	BRCA1	chr17q21	204531_s_at 211851_x_at	NM_007295 AF005068
BUB1 budding uninhibited by benzimidazoles 1 homolog (yeast)	BUB1	chr2q14	209642_at 215509_s_at	AF043294 AL137654
BUB1 budding uninhibited by benzimidazoles 1 homolog beta (yeast)	BUB1B	chr15q15	203755_at	NM_001211
cyclin A2	CCNA2	chr4q25-q31	203418_at 213226_at	NM_001237 AI346350
cyclin B1	CCNB1	chr5q12	214710_s_at	BE407516
cyclin B2	CCNB2	chr15q22.2	202705_at	NM_004701
cyclin E2	CCNE2	chr8q22.1	205034_at 211814_s_at	NM_004702 AF112857
cyclin F	CCNF	chr16p13.3	204826_at 204827_s_at	NM_001761 U17105
cyclin J	CCNJ	chr10pter-q26.12	219470_x_at	NM_019084

cyclin T2	CCNT2	chr2q21.3	204645_at	NM_001241
chaperonin containing TCP1, subunit 2 (beta)	CCT2	chr12q15	201946_s_at	AL545982
cell division cycle 20 homolog (S. cerevisiae)	CDC20	chr1p34.1	202870_s_at	NM_001255
cell division cycle 25 homolog A (S. pombe)	CDC25A	chr3p21	204695_at	AI343459
cell division cycle 25 homolog C (S. pombe)	CDC25C	chr5q31	205167_s_at	NM_001790
			217010_s_at	AF277724
cell division cycle 27 homolog (S. cerevisiae)	CDC27	chr17q12-q23.2	217879_at	AL566824
cell division cycle 6 homolog (S. cerevisiae)	CDC6	chr17q21.3	203968_s_at	NM_001254
cyclin-dependent kinase 2	CDK2	chr12q13	204252_at	M68520
			211804_s_at	AB012305
cyclin-dependent kinase 4	CDK4	chr12q14	202246_s_at	NM_000075
cyclin-dependent kinase inhibitor 3 (CDK2-associated dual specificity phosphatase)	CDKN3	chr14q22	209714_s_at	AF213033
chromatin licensing and DNA replication factor 1	CDT1	chr16q24.3	209832_s_at	AF321125
centromere protein E, 312kDa	CENPE	chr4q24-q25	205046_at	NM_001813
centromere protein F, 350/400ka (mitosin)	CENPF	chr1q32-q41	207828_s_at	NM_005196
			209172_s_at	U30872
chromatin assembly factor 1, subunit A (p150)	CHAF1A	chr19p13.3	203975_s_at	BF000239
			203976_s_at	NM_005483
			214426_x_at	BF062223
CHK2 checkpoint homolog (S. pombe)	CHEK2	chr22q11 22q12.1	210416_s_at	BC004207
CDC28 protein kinase regulatory subunit 1B	CKS1B	chr1q21.2	201897_s_at	NM_001826
CDC28 protein kinase regulatory subunit 2	CKS2	chr9q22	204170_s_at	NM_001827
DEAD/H (Asp-Glu-Ala-Asp/His) box polypeptide 11 (CHL1-like helicase homolog, S. cerevisiae)	DDX11	chr12p11	210206_s_at	U33833
extra spindle pole bodies homolog 1 (S. cerevisiae)	ESPL1	chr12q	38158_at	D79987
exonuclease 1	EXO1	chr1q42-q43	204603_at	NM_003686
fumarate hydratase	FH	chr1q42.1	203032_s_at	AI363836
fyn-related kinase	FRK	chr6q21-q22.3	207178_s_at	NM_002031
G-2 and S-phase expressed 1	GTSE1	chr22q13.2-q13.3	204318_s_at	NM_016426
			215942_s_at	BF973178

high mobility group AT-hook 1	HMGA1	chr6p21	206074_s_at	NM_002131
high-mobility group box 2	HMGB2	chr4q31	208808_s_at	BC000903
interleukin enhancer binding factor 3, 90kDa	ILF3	chr19p13.2	208931_s_at	AF147209
			211375_s_at	AF141870
kinesin family member 11	KIF11	chr10q24.1	204444_at	NM_004523
kinesin family member 22	KIF22	chr16p11.2	202183_s_at	NM_007317
			216969_s_at	AC002301
kinesin family member 23	KIF23	chr15q23	204709_s_at	NM_004856
kinesin family member 2C	KIF2C	chr1p34.1	209408_at	U63743
			211519_s_at	AY026505
kinesin family member C1	KIFC1	chr6p21.3	209680_s_at	BC000712
kinetochore associated 1	KNTC1	chr12q24.31	206316_s_at	NM_014708
ligase I, DNA, ATP-dependent	LIG1	chr19q13.2-q13.3	202726_at	NM_000234
mitogen-activated protein kinase 1	MAPK1	chr22q11.2 22q11.21	208351_s_at	NM_002745
minichromosome maintenance complex component 2	MCM2	chr3q21	202107_s_at	NM_004526
minichromosome maintenance complex component 4	MCM4	chr8q11.2	212141_at	AA604621
			212142_at	AI936566
			222036_s_at	AI859865
			222037_at	AI859865
minichromosome maintenance complex component 5	MCM5	chr22q13.1	201755_at	NM_006739
			216237_s_at	AA807529
antigen identified by monoclonal antibody, Ki-67	MKI67	chr10q25-qter	212020_s_at	AU152107
			212021_s_at	AU132185
			212022_s_at	BF001806
			212023_s_at	AU147044
M-phase phosphoprotein 1	MPHOSPH1	chr10q23.31	205235_s_at	NM_016195
M-phase phosphoprotein 9	MPHOSPH9	chr12q24.31	206205_at	NM_022782
mutS homolog 6 (E. coli)	MSH6	chr2p16	202911_at	NM_000179
			211450_s_at	D89646
non-SMC condensin I complex, subunit D2	NCAPD2	chr12p13.3	201774_s_at	AK022511
non-SMC condensin I complex, subunit G	NCAPG	chr4p15.33	218662_s_at	NM_022346
			218663_at	NM_022346
non-SMC condensin I complex, subunit H	NCAPH	chr2q11.2	212949_at	D38553

NDC80 homolog, kinetochore complex component (<i>S. cerevisiae</i>)	NDC80	chr18p11.32	204162_at	NM_006101
NIMA (never in mitosis gene a)-related kinase 2	NEK2	chr1q32.2-q41	204641_at	NM_002497
		chr1q32.2-q41	211080_s_at	Z25425
NIMA (never in mitosis gene a)-related kinase 4	NEK4	chr3p21.1	204634_at	NM_003157
non-metastatic cells 1, protein (NM23A) expressed in	NME1	chr17q21.3	201577_at	NM_000269
nucleolar and coiled-body phosphoprotein 1	NOLC1	chr10q24.32	205895_s_at	NM_004741
nucleophosmin (nucleolar phosphoprotein B23, numatrin)	NPM1	chr5q35	221691_x_at	AB042278
			221923_s_at	AA191576
nucleoporin 98kDa	NUP98	chr11p15.5	203194_s_at	AA527238
origin recognition complex, subunit 1-like (yeast)	ORC1L	chr1p32	205085_at	NM_004153
origin recognition complex, subunit 4-like (yeast)	ORC4L	chr2q22-q23	203351_s_at	AF047598
origin recognition complex, subunit 6 like (yeast)	ORC6L	chr16q12	219105_x_at	NM_014321
protein kinase, membrane associated tyrosine/threonine 1	PKMYT1	chr16p13.3	204267_x_at	NM_004203
polo-like kinase 1 (<i>Drosophila</i>)	PLK1	chr16p12.1	202240_at	NM_005030
polo-like kinase 4 (<i>Drosophila</i>)	PLK4	chr4q28	204886_at	AL043646
			204887_s_at	NM_014264
			211088_s_at	Z25433
PMS1 postmeiotic segregation increased 1 (<i>S. cerevisiae</i>)	PMS1	chr2q31-q33 2q31.1	213677_s_at	BG434893
polymerase (DNA directed), theta	POLQ	chr3q13.33	219510_at	NM_006596
protein phosphatase 1D magnesium-dependent, delta isoform	PPM1D	chr17q23.2	204566_at	NM_003620
protein phosphatase 2 (formerly 2A), regulatory subunit A, beta isoform	PPP2R1B	chr11q23.2	202886_s_at	M65254
protein phosphatase 6, catalytic subunit	PPP6C	chr9q33.3	206174_s_at	NM_002721
protein regulator of cytokinesis 1	PRC1	chr15q26.1	218009_s_at	NM_003981
primase, DNA, polypeptide 1 (49kDa)	PRIM1	chr12q13	205053_at	NM_000946

primase, DNA, polypeptide 2 (58kDa)	PRIM2	chr6p12-p11.1	205628_at	NM_000947
protein arginine methyltransferase 5	PRMT5	chr14q11.2-q21	217786_at	NM_006109
pituitary tumor-transforming 1	PTTG1	chr5q35.1	203554_x_at	NM_004219
pituitary tumor-transforming 3	PTTG3	chr8q13.1	208511_at	NM_021000
RAD51 homolog (RecA homolog, E. coli) (S. cerevisiae)	RAD51	chr15q15.1	205024_s_at	NM_002875
RAD54 homolog B (S. cerevisiae)	RAD54B	chr8q21.3-q22	219494_at	NM_012415
Ras association (RalGDS/AF-6) domain family member 1	RASSF1	chr3p21.3	204346_s_at	NM_007182
replication factor C (activator 1) 2, 40kDa	RFC2	chr7q11.23	1053_at	M87338
			203696_s_at	NM_002914
replication factor C (activator 1) 3, 38kDa	RFC3	chr13q12.3-q13	204128_s_at	NM_002915
replication factor C (activator 1) 5, 36.5kDa	RFC5	chr12q24.2-q24.3	203209_at	BC001866
			203210_s_at	NM_007370
ribonuclease H2, subunit A	RNASEH2A	chr19p13.13	203022_at	NM_006397
SET nuclear oncogene	SET	chr9q34	213047_x_at	A1278616
S-phase kinase-associated protein 2 (p45)	SKP2	chr5p13	210567_s_at	BC001441
structural maintenance of chromosomes 2	SMC2	chr9q31.1	204240_s_at	NM_006444
			213253_at	AU154486
sperm associated antigen 5	SPAG5	chr17q11.2	203145_at	NM_006461
SFRS protein kinase 1	SRPK1	chr6p21.3-p21.2	202199_s_at	AW082913
signal transducer and activator of transcription 1, 91kDa	STAT1	chr2q32.2	AFFX-HUMISGF3A/M97935_5_at	AFFX-HUMISGF3A/M97935_5
suppressor of variegation 3-9 homolog 2 (Drosophila)	SUV39H2	chr10p13	219262_at	NM_024670
TAR DNA binding protein	TARDBP	chr1p36.22	200020_at	NM_007375
transcription factor A, mitochondrial	TFAM	chr10q21	203177_x_at	NM_003201
topoisomerase (DNA) II binding protein 1	TOPBP1	chr3q22.1	202633_at	NM_007027
TPX2, microtubule-associated, homolog (Xenopus laevis)	TPX2	chr20q11.2	210052_s_at	AF098158
TTK protein kinase	TTK	chr6q13-q21	204822_at	NM_003318
tubulin, gamma 1	TUBG1	chr17q21	201714_at	NM_001070

Conclusions

The present invention is the first to report an association between a gene proliferation signature and major clinico-pathologic variables as well as outcome in colorectal cancer. The disclosed study investigated the proliferation state of tumours using an *in vitro*-
5 derived multi-gene proliferation signature and by Ki-67 immunostaining. According to the results herein, low expression of the GPS in tumours was associated with a higher risk of recurrence and shorter survival in two independent cohorts of patients. In contrast, Ki-67 proliferation index was not associated with any clinically relevant endpoints.

- 10 The colorectal GPS encompasses 38 mitotic cell cycle genes and includes a core set of genes (CDC2, RFC4, PCNA, CCNE1, CDK7, MCM genes, FEN1, MAD2L1, MYBL2, RRM2 and BUB3) that are part of proliferation signatures defined for tumours of the breast (40),(41), ovary (42), liver (43), acute lymphoblastic leukaemia (44), neuroblastoma (45), lung squamous cell carcinoma (46), head and neck (47), prostate (48), and stomach (49).
- 15 This represents a conserved pattern of expression, as most of these genes have been found to be highly overexpressed in fast-growing tumours and to reflect a high proportion of rapidly cycling cells (50). Therefore, the expression level of the colorectal GPS provides a measure for the proliferative state of a tumour.
- 20 In this study, several clinico-pathologic variables related to poor outcome (disease stage, lymph node metastasis and lymphatic invasion) were associated with low GPS expression in Cohort A patients. In Cohort B, consisting entirely of stage II tumours, the study assessed the association between the GPS and lymphatic invasion. The association failed to reach statistical significance due to the small number of tumours with lymphatic
25 invasion in this cohort (5/55). Without being bound by theory, the low GPS expression in more advanced tumours may indicate that CRC progression is not driven by enhanced proliferation. While accelerated proliferation may still be an important driving force during the initial phases of tumourigenesis, it is possible that more advanced disease is more dependent on processes such as genetic instability to allow continuous selection.
- 30 Consistent with our finding, two large-scale studies reported an association between decreased expression of CDK2, cyclin E and A, and advanced stage, deep infiltration and lymph node metastasis (51),(52).

The relationship between low GPS and unfavourable clinico-pathologic variables
35 suggested that the GPS should also predict patient outcome. Indeed, in both Cohort A and B, low GPS expression was associated with a higher risk of recurrence and shorter overall and recurrence-free survival. In Cohort B, where all patients had stage II tumours,

the association remained in multivariate analysis. However, in Cohort A, where patients had stage I-IV disease, the association was not independent of tumour stage. The number of patients with and without recurrence, within each stage of disease in Cohort A, was probably insufficient to demonstrate an independent association between the GPS and survival. In Cohort B, low GPS expression and lymphatic invasion remained independent predictors in multivariate analysis suggesting that the GPS may improve the prediction of CRC patient outcome within the same disease stage. Not surprisingly, the presence of lymph node and distant organ involvement were the most powerful predictors of outcome as these are direct manifestations of tumour metastasis.

Treatment with radiotherapy or chemotherapy, used in 18% and 27% of Cohort A patients respectively, was a possible confounding factor in this study. Theoretically, the improved survival associated with elevated GPS expression might reflect the better response of fast proliferating tumours to cancer treatment (53),(54). However, no correlation was found between treatment and GPS expression. Furthermore, no patients in Cohort B received adjuvant therapy indicating that the association between GPS and survival is independent of treatment. It should be noted that this study was not designed to investigate the relationship between tumour proliferation and response to chemotherapy or radiotherapy.

The sample size may also explain the lack of an association between clinico-pathologic variables and survival with Ki-67 PI in the present study. As mentioned above, other studies on Ki-67 and CRC outcome have reported inconsistent findings. However, in the three other CRC studies with the largest sample size a low Ki-67 PI was associated with a worse prognosis (27),(29),(30). We came to the same conclusion applying the GPS, but based on a much smaller sample size. The multi-gene expression analysis was therefore a more sensitive tool to assess the relationship between proliferation and prognosis than the Ki-67 PI.

The biological reason behind an unfavourable prognosis in tumours with a low GPS will involve further investigation. Mechanisms that could potentially contribute to worse clinical outcome in low GPS tumours include: (i) a more effective immune response to rapidly proliferating tumours; (ii) a higher level of genetic damage that may render cancer cells more resistant to apoptosis, and increase invasiveness, but also perturb smooth replication machinery; (iii) an increased number of cancer stem cells that divide slowly, similar to normal stem cells, but have a high metastatic potential; and (iv) a higher proportion of microsatellite unstable tumours which have a high proliferation rate but a relatively good prognosis.

In sum, the present invention has clarified the previous, conflicting results relating to the prognostic role of cell proliferation in colorectal cancer. A GPS has been developed using CRC cell lines and has been applied to two independent patient cohorts. It was found that
5 low expression of growth-related genes in CRC was associated with more advanced tumour stage (Cohort A) and poor clinical outcome within the same stage (Cohort B). Multi-gene expression analysis was shown as a more powerful indicator than the long-established proliferation marker, Ki-67, for predicting outcome. For future studies, it will be useful to determine the reasons that CRC differs from other common epithelia cancers,
10 such as breast and lung cancers (e.g., in reference to Ki-67). This will likely provide insights into important underlying biological mechanisms. From a practical viewpoint, the ability to stratify recurrence risk within a given pathological stage could enable adjuvant therapy to be targeted more accurately. Thus, GPS expression can be used as an adjunct to conventional staging for identifying patients at high risk of recurrence and death from
15 colorectal cancer.

All publications and patents mentioned in the above specification are herein incorporated by reference.

20 Wherein in the foregoing description reference has been made to integers or components having known equivalents, such equivalents are herein incorporated as if individually set forth.

Although the invention has been described by way of example and with reference to
25 possible embodiments thereof, it is to be appreciated that improvements and/or modifications may be made without departing from the scope or the spirit thereof.

References:

1. Evan GI, Vousden KH: Proliferation, cell cycle and apoptosis in cancer. Nature
30 411:342-8, 2001
2. Whitfield ML, George LK, Grant GD, et al: Common markers of proliferation. Nat Rev Cancer 6:99-106, 2006
3. Rew DA, Wilson GD: Cell production rates in human tissues and tumours and their significance. Part 1: an introduction to the techniques of measurement and their
35 limitations. Eur J Surg Oncol 26:227-38, 2000
4. Endle E, Gerdes J: The Ki-67 protein: fascinating forms and an unknown function. Exp Cell Res 257:231-7, 2000

5. Brown DC, Gatter KC: Ki67 protein: The immaculate deception. *Histopathology* 40:2-11, 2002
6. Paik S, Shak S, Tang G, et al: A multigene assay to predict recurrence of tamoxifen-treated, node-negative breast cancer. *N Engl J Med* 351:2817-26, 2004
- 5 7. Ofner D, Grothaus A, Riedmann B, et al: MIB1 in colorectal carcinomas: its evaluation by three different methods reveals lack of prognostic significance. *Anal Cell Pathol* 12:61-70, 1996
8. Ihmann T, Liu J, Schwabe W, et al: High-level mRNA quantification of proliferation marker pKi-67 is correlated with favorable prognosis in colorectal carcinoma. *J Cancer*
- 10 *Res Clin Oncol* 130:749-756, 2004
9. Van Oijen MG, Medema RH, Slootweg PJ, et al: Positivity of the proliferation marker pKi-67 in non-cycling cells. *Am J Clin Pathol* 110:24-31, 1998
10. Duchrow M, Ziemann T, Windhövel U, et al: Colorectal carcinomas with high MIB-1 labelling indices but low pKi67 mRNA levels correlate with better prognostic outcome.
- 15 *Histopathology* 42:566-574, 2003
11. Evans C, Morrison I, Heriot AG, et al: The correlation between colorectal cancer rates of proliferation and apoptosis and systemic cytokine levels; plus their influence upon survival. *Br J Cancer* 94:1412-9, 2006
12. Rosati G, Chiacchio R, Reggiardo G, et al: Thymidylate synthase expression, p53, bcl-
- 20 2, Ki-67 and p27 in colorectal cancer: relationships with tumour recurrence and survival. *Tumour Biol* 25:258-63, 2004
13. Ishida H, Miwa H, Tatsuta M, et al: Ki-67 and CEA expression as prognostic markers in Dukes' C colorectal cancer. *Cancer Lett* 207:109-115, 2004
14. Buglioni S, D'Agnano I, Cosimelli M, et al: Evaluation of multiple bio-pathological
- 25 factors in colorectal adenocarcinomas: independent prognostic role of p53 and bcl-2. *Int J Cancer* 84:545-52, 1999
15. Guerra A, Borda F, Javier Jimenez F, et al: Multivariate analysis of prognostic factors in resected colorectal cancer: a new prognostic index. *Eur J Gastroenterol Hepatol* 10:51-8, 1998
- 30 16. Kyzer S, Gordon PH: Determination of proliferative activity in colorectal carcinoma using monoclonal antibody Ki67. *Dis Colon Rectum* 40:322-5, 1997
17. Jansson A, Sun XF: Ki-67 expression in relation to clinicopathological variables and prognosis in colorectal adenocarcinomas. *APMIS* 105:730-4, 1997
18. Baretton GB, Diebold J, Christoforis G, et al: Apoptosis and immunohistochemical bcl-
- 35 2 expression in colorectal adenomas and carcinomas. Aspects of carcinogenesis and prognostic significance. *Cancer* 77:255-64, 1996

19. Sun XF, Carstensen JM, Stal O, et al: Proliferating cell nuclear antigen (PCNA) in relation to ras, c-erbB-2, p53, clinico-pathological variables and prognosis in colorectal adenocarcinoma. *Int J Cancer* 69:5-8, 1996
20. Kubota Y, Petras RE, Easley KA, et al: Ki-67-determined growth fraction versus
5 standard staging and grading parameters in colorectal carcinoma. A multivariate analysis. *Cancer* 70:2602-9, 1992
21. Valera V, Yokoyama N, Walter B, et al: Clinical significance of Ki-67 proliferation index in disease progression and prognosis of patients with resected colorectal carcinoma. *Br J Surg* 92:1002-7, 2005
- 10 22. Dziegiel P, Forgacz J, Suder E, et al: Prognostic significance of metallothionein expression in correlation with Ki-67 expression in adenocarcinomas of large intestine. *Histol Histopathol* 18:401-7, 2003
23. Scopa CD, Tsamandas AC, Zolata V, et al: Potential role of bcl-2 and Ki-67 expression and apoptosis in colorectal carcinoma: a clinicopathologic study. *Dig Dis Sci*
15 48:1990-7, 2003
24. Bhatavdekar JM, Patel DD, Chikhlikar PR, et al: Molecular markers are predictors of recurrence and survival in patients with Dukes B and Dukes C colorectal adenocarcinoma. *Dis Colon Rectum* 44:523-33, 2001
25. Chen YT, Henk MJ, Carney KJ, et al: Prognostic Significance of Tumor Markers in
20 Colorectal Cancer Patients: DNA Index, S-Phase Fraction, p53 Expression, and Ki-67 Index. *J Gastrointest Surg* 1:266-273, 1997
26. Choi HJ, Jung IK, Kim SS, et al: Proliferating cell nuclear antigen expression and its relationship to malignancy potential in invasive colorectal carcinomas. *Dis Colon Rectum* 40:51-9, 1997
- 25 27. Hilska M, Collan YU, O Laine VJ, et al: The significance of tumour markers for proliferation and apoptosis in predicting survival in colorectal cancer. *Dis Colon Rectum* 48:2197-208, 2005
28. Salminen E, Palmu S, Vahlberg T, et al: Increased proliferation activity measured by immunoreactive Ki67 is associated with survival improvement in rectal/recto sigmoid
30 cancer. *World J Gastroenterol* 11:3245-9, 2005
29. Garrity MM, Burgart LJ, Mahoney MR, et al: Prognostic value of proliferation, apoptosis, defective DNA mismatch repair, and p53 overexpression in patients with resected Dukes' B2 or C colon cancer: a North Central Cancer Treatment Group Study. *J Clin Oncol* 22:1572-82, 2004
- 35 30. Allegra CJ, Paik S, Colangelo LH, et al: Prognostic value of thymidylate synthase, Ki-67, and p53 in patients with Dukes' B and C colon cancer: a National Cancer Institute-

- National Surgical Adjuvant Breast and Bowel Project collaborative study. *J Clin Oncol* 21:241-50, 2003
31. Palmqvist R, Sellberg P, Öberg A, et al: Low tumour cell proliferation at the invasive margin is associated with a poor prognosis in Dukes' stage B colorectal cancers. *Br J Cancer* 79:577-81, 1999
32. Paradiso A, Rabinovich M, Vallejo C, et al: p53 and PCNA expression in advanced colorectal cancer: response to chemotherapy and long-term prognosis. *Int J Cancer* 69:437-41, 1996
33. Neoptolemos JP, Oates GD, Newbold KM, et al: Cyclin/proliferation cell nuclear antigen immunohistochemistry does not improve the prognostic power of Dukes' or Jass' classifications for colorectal cancer. *Br J Surg* 82:184-7, 1995
34. Compton C, Fenoglio-Preiser CM, Pettigrew N, et al: American joint committee on cancer prognostic factors consensus conference. Colorectal working group. *Cancer* 88:1739-1757, 2000
35. Colantuoni C, Henry G, Zeger S, et al: SNOMAD (Standardization and Normalization of MicroArray Data): web-accessible gene expression data analysis. *Bioinformatics* 18:1540-1541, 2002
36. Livak KJ, Schmittgen TD: Analysis of Relative Gene Expression Data Using Real-Time Quantitative PCR and the 2- $\Delta\Delta CT$ Method. *METHODS* 25:402-408, 2001
37. Pocock SJ, Clayton TC, Altman DG: Survival plots of time-to-event outcomes in clinical trials: good practice and pitfalls. *Lancet* 359:1686-89, 2002
38. Trushar VG, Tibshirani R, Chu G: Significance analysis of microarrays applied to the ionizing radiation response. *Proc Natl Acad Sci USA* 98:5116-21, 2001
39. Hosack DA, Dennis G, Sherman BT, et al: Identifying biological themes within lists of genes with EASE. *Genome biology* 4:R70, 2003
40. Perou CM, Jeffrey SS, DE Rijn MV: Distinctive gene expression patterns in human mammary epithelial cells and breast cancers. *Proc. Natl. Acad. Sci. USA* 96:9212-17, 1999
41. Perou CM: Molecular portraits of human breast tumours. *Nature* 406:747-752, 2000
42. Welsh JB, Zarrinkar PP, Sapinoso LM, et al: Analysis of gene expression profiles in normal and neoplastic ovarian tissue samples identifies candidate molecular markers of epithelial ovarian cancer. *Proc. Natl Acad. Sci. USA* 98:1176-1181, 2001
43. Chen X, Cheung ST, So S, et al: Gene expression patterns in human liver cancers. *Mol. Biol. Cell* 13:1929-1939, 2002
44. Kirschner-Schwabe R, Lottaz C, Todling J, et al: Expression of late cell cycle genes and an increased proliferative capacity characterize very early relapse of childhood acute lymphoblastic leukemia. *Clin Cancer Res* 12:4553-61, 2006

45. Krasnoselsky AL, Whiteford CC, Wei JS, et al: Altered expression of cell cycle genes distinguishes aggressive neuroblastoma. *Oncogene* 24:1533-1541, 2005
46. Inamura K, Fujiwara T, Hoshida Y, et al: Two subclasses of lung squamous cell carcinoma with different gene expression profiles and prognosis identified by hierarchical clustering and non-negative matrix factorization. *Oncogene* 24:7105-13, 2005
- 5 47. Chung CH, Parker JS, Karaca G, et al: Molecular classification of head and neck squamous cell carcinomas using patterns of gene expression. *Cancer Cell* 5:489-500, 2004
48. LaTulippe E, Satagopan J, Smith A, et al: Comprehensive gene expression analysis of prostate cancer reveals distinct transcriptional programs associated with metastatic disease. *Cancer Res* 62:4499-4506, 2002
- 10 49. Hippo Y, Taniguchi H, Tsutumi S, et al: Global gene expression analysis of gastric cancer by oligonucleotide microarrays. *Cancer Res* 62:233-40, 2002
50. Whitfield ML, Sherlock G, Saldanha AJ, et al: Identification of genes periodically expressed in the human cell cycle and their expression in tumours. *Mol Biol Cell* 13:1977-2000, 2002
- 15 51. Li JQ, Miki H, Ohmori M, et al: Expression of cyclin E and cyclin-dependent kinase 2 correlates with metastasis and prognosis in colorectal carcinoma. *Hum Pathol* 32:945-53, 2001
- 20 52. Li JQ, Miki H, Wu F, et al: Cyclin A correlates with carcinogenesis and metastasis, and p27 (kip1) correlates with lymphatic invasion, in colorectal neoplasms. *Hum Pathol* 33, 1006-15, 2002
53. Itamochi H, Kigawa J, Sugiyama T, et al: Low proliferation activity may be associated with chemoresistance in clear cell carcinoma of the ovary. *Obstet Gynecol* 100:281-287, 2002
- 25 54. Imdahl A, Jenkner J, Ihling C, et al: Is MIB-1 proliferation index a predictor for response to neoadjuvant therapy in patients with esophageal cancer? *Am J Surg* 179:514-520, 2000

CLAIMS

1. A prognostic signature for determining progression of gastrointestinal cancer in a patient, comprising one or more genes selected from Table A, Table B, Table C or Table D.

2. The signature of claim 1, wherein the signature comprises one or more genes selected from any one of CDC2, MCM6, RPA3, MCM7, PCNA, G22P1, KPNA2, ANLN, APG7L, TOPK, GMNN, RRM1, CDC45L, MAD2L1, RAN, DUT, RRM2, CDK7, MLH3, SMC4L1, CSPG6, POLD2, POLE2, BCCIP, Pfs2, TREX1, BUB3, FEN1, DRF1, PREI3, CCNE1, RPA1, POLE3, RFC4, MCM3, CHEK1, CCND1, and CDC37.

3. A method of predicting the likelihood of long-term survival of a gastrointestinal cancer patient without the recurrence of gastrointestinal cancer, comprising determining the expression level of one or more prognostic RNA transcripts or their expression products in a gastrointestinal sample obtained from the patient, normalized against the expression level of all RNA transcripts or their products in the gastrointestinal cancer tissue sample, or of a reference set of RNA transcripts or their expression products;

wherein the prognostic RNA transcript is the transcript of one or more genes selected from table A, Table B, Table C or Table D ; and

establishing likelihood of long-term survival without gastrointestinal cancer recurrence.

4. The method of claim 3, wherein at least one prognostic RNA transcripts or its expression products is selected from any one of CDC2, MCM6, RPA3, MCM7, PCNA, G22P1, KPNA2, ANLN, APG7L, TOPK, GMNN, RRM1, CDC45L, MAD2L1, RAN, DUT, RRM2, CDK7, MLH3, SMC4L1, CSPG6, POLD2, POLE2, BCCIP, Pfs2, TREX1, BUB3, FEN1, DRF1, PREI3, CCNE1, RPA1, POLE3, RFC4, MCM3, CHEK1, CCND1, and CDC37

5. The method of claim 3 or claim 4 comprising determining the expression level of at least two, at least five, at least 10, or at least 15 of the prognostic RNA transcripts or their expression products.

6. The method according to any one of claims 3 to 5, wherein increased expression of the one or more prognostic RNA transcripts or their expression products indicates an increased likelihood of long-term survival without gastrointestinal cancer recurrence.

7. The method according to any one of claims 3 to 5, wherein a predictive model is applied, established by applying a predictive method to expressions levels of the predictive signature in recurrent and non-recurrent tumour samples, to establishing likelihood of long-term survival without gastrointestinal cancer recurrence.

5

8. The method of claim 7, wherein said predictive method is selected from the group consisting of linear models, support vector machines, neural networks, classification and regression trees, ensemble learning methods, discriminant analysis, nearest neighbor method, bayesian networks, independent components analysis.

10

9. The method of any one of claims 3 to 8 wherein the gastrointestinal cancer is gastric cancer or colorectal cancer.

15

10. The method of any one of claims 3 to 9 wherein the expression level of one or more prognostic RNA transcripts is determined.

11. The method of any one of claims 3 to 10 wherein the RNA is isolated from a fixed, wax- embedded gastrointestinal cancer tissue specimen of the patient.

20

12. The method of any one of claims 3 to 10 wherein the RNA is isolated from core biopsy tissue or fine needle aspirate cells.

13. An array comprising polynucleotides hybridizing to two or more genes selected from table A, Table B, Table C or Table D.

25

14. An array of claim 13 comprising polynucleotides hybridizing to two or more of the following genes: CDC2, MCM6, RPA3, MCM7, PCNA, G22P1, KPNA2, ANLN, APG7L, TOPK, GMNN, RRM1, CDC45L, MAD2L1, RAN, DUT, RRM2, CDK7, MLH3, SMC4L1, CSPG6, POLD2, POLE2, BCCIP, Pfs2, TREX1, BUB3, FEN1, DRF1, PREI3, CCNE1, RPA1, POLE3, RFC4, MCM3, CHEK1, CCND1, and CDC37.

30

15. The array of claim 13 or claim 14 comprising polynucleotides hybridizing to at least 3, at least five, at least 10 or at least 15 of the genes.

35

16. The array of claim 13 comprising polynucleotides hybridizing to the following genes: CDC2, MCM6, RPA3, MCM7, PCNA, G22P1, KPNA2, ANLN, APG7L, TOPK, GMNN, RRM1, CDC45L, MAD2L1, RAN, DUT, RRM2, CDK7, MLH3, SMC4L1, CSPG6, POLD2,

POLE2, BCCIP, Pfs2, TREX1, BUB3, FEN1, DRF1, PREI3, CCNE1, RPA1, POLE3, RFC4, MCM3, CHEK1, CCND1, and CDC37.

17. The array of any one of claims 13 to 16 wherein the polynucleotides are cDNAs.

18. The array of claim 17 wherein the cDNAs are about 500 to 5000 bases long.

19. The array of claim any one of claims 13 to 16 wherein the polynucleotides are oligonucleotides.

20. The array of claim 19 wherein the oligonucleotides are about 20 to 80 bases long.

21. The array of any one of claims 13 to 20 wherein the solid surface is glass.

22. A method of predicting the likelihood of long-term survival of a patient diagnosed with gastrointestinal cancer, without the recurrence of gastrointestinal cancer, comprising the steps of:

(1) determining the expression levels of the RNA transcripts or the expression products of genes or a gene selected from table A, Table B, Table C or Table D, in a gastrointestinal cancer tissue sample obtained from the patient, normalized against the expression levels of all RNA transcripts or their expression products in the gastrointestinal cancer tissue sample, or of a reference set of RNA transcripts or their products;

(2) subjecting the data obtained in step (1) to statistical analysis; and

(3) determining whether the likelihood of the long-term survival has increased or decreased;

and establishing the likelihood of long-term survival without gastrointestinal cancer recurrence.

23. The method of claim 22, wherein at least one prognostic RNA transcripts or its expression products is selected from any one CDC2, MCM6, RPA3, MCM7, PCNA, G22P1, KPNA2, ANLN, APG7L, TOPK, GMNN, RRM1, CDC45L, MAD2L1, RAN, DUT, RRM2, CDK7, MLH3, SMC4L1, CSPG6, POLD2, POLE2, BCCIP, Pfs2, TREX1, BUB3, FEN1, DRF1, PREI3, CCNE1, RPA1, POLE3, RFC4, MCM3, CHEK1, CCND1, and CDC37.

24. The method of claim 22 or claim 23 wherein the statistical analysis is performed by using the Cox Proportional Hazards model.
25. A method of preparing a personalized genomics profile for a cancer patient, comprising the steps of: (a) subjecting RNA extracted from a gastrointestinal tissue obtained from the patient to gene expression analysis; (b) determining the expression level of one or more genes selected from the gastrointestinal cancer gene set listed in any one of Table A, Table B, Table C or Table D, wherein the expression level is normalized against a control gene or genes and optionally is compared to the amount found in a gastrointestinal cancer reference tissue set; and (c) creating a report summarizing the data obtained by the gene expression analysis.
26. The method of claim 25, wherein the gastrointestinal tissue comprises gastrointestinal cancer cells.
27. The method of claim 25 wherein the gastrointestinal tissue is obtained from a fixed, paraffin-embedded biopsy sample.
28. The method of claim 27 wherein the RNA is fragmented.
29. The method of any one of claims 22 to 28 wherein the report includes prediction of the likelihood of long term survival of the patient.
30. The method of any one of claims 22 to 29 wherein the report includes recommendation for a treatment modality of the patient.
31. A prognostic method comprising: (a) subjecting a sample comprising gastrointestinal cancer cells obtained from a patient to quantitative analysis of the levels of RNA transcripts of at least one gene selected from any one of Table A, Table B, Table C or table D, or its product, and (b) identifying the patient as likely to have an increased likelihood of long-term survival without gastrointestinal cancer recurrence if normalized expression levels of the gene or genes, or their products, are elevated above a defined expression threshold.
32. The method of claim 31, wherein at least one prognostic RNA transcripts or its expression products is selected from any one CDC2, MCM6, RPA3, MCM7, PCNA, G22P1, KPNA2, ANLN, APG7L, TOPK, GMNN, RRM1, CDC45L, MAD2L1, RAN, DUT,

RRM2, CDK7, MLH3, SMC4L1, CSPG6, POLD2, POLE2, BCCIP, Pfs2, TREX1, BUB3, FEN1, DRF1, PREI3, CCNE1, RPA1, POLE3, RFC4, MCM3, CHEK1, CCND1, and CDC37.

5 33. The method of claim 31 or claim 32, wherein the levels of the RNA transcripts of the genes are normalized relative to the mean level of the RNA transcript or the product of two or more housekeeping genes.

10 34. The method of claim 33 wherein the housekeeping genes are selected from the group consisting of glyceraldehyde-3-phosphate dehydrogenase (GAPDH), Cypl, albumin, actins, tubulins, cyclophilin hypoxanthine phosphoribosyltransferase (HRPT), L32, 28S, and 185.

15 35. The method of any one of claims 31 to 34 wherein the sample is subjected to global gene expression analysis of all genes present above the limit of detection.

20 36. The method of any one of claims 31 to 35 wherein the levels of RNA transcripts of the genes are normalized relative to the mean signal of the RNA transcripts or the products of all assayed genes or a subset thereof.

37. The method of any one of claims 31 to 36 wherein the levels of RNA transcripts are determined by quantitative RT-PCR, and the signal is a Ct value.

25 38. The method of claim 36 wherein the assayed genes include at least 50 or at least 100 cancer related genes.

39. The method of any one of claims 31 to 38 wherein the patient is human.

30 40. The method of any one of claims 31 to 39 wherein the sample is a fixed, paraffin-embedded tissue (FPET) sample, or fresh or frozen tissue sample.

41. The method of any one of claims 31 to 39 wherein the sample is a tissue sample from fine needle, core, or other types of biopsy.

35 42. The method of any one of claims 31 to 41 wherein the quantitative analysis is performed by quantitative RT-PCR.

43. The method of any one of claims 31 to 41 wherein the quantitative analysis is performed by quantifying the products of the genes.

44. The method of any one of claims 31 to 41 wherein the products are quantified by immunohistochemistry or by proteomics technology.

45. The method of any one of claims 31 to 44 further comprising the step of preparing a report indicating that the patient has an increased likelihood of long-term survival without gastrointestinal cancer recurrence.

46. A kit comprising one or more of (1) extraction buffer/reagents and protocol; (2) reverse transcription buffer/reagents and protocol; and (3) quantitative RT-PCR buffer/reagents and protocol suitable for performing the method of any one of claims 3, 25, or 30.