

Benzimidazole-proline derivatives

The present invention relates to novel benzimidazole-proline derivatives of formula (I) and their use as pharmaceuticals. The invention also concerns related aspects including processes for the preparation of the compounds, pharmaceutical compositions containing one or more compounds of formula (I), and their use as orexin receptor antagonists, especially as orexin-1 receptor antagonists.

Orexins (orexin A or OX-A and orexin B or OX-B) are neuropeptides found in 1998 by two research groups, orexin A is a 33 amino acid peptide and orexin B is a 28 amino acid peptide (Sakurai T. *et al.*, Cell, **1998**, 92, 573-585). Orexins are produced in discrete neurons of the lateral hypothalamus and bind to the G-protein-coupled receptors (OX₁ and OX₂ receptors). The orexin-1 receptor (OX₁) is selective for OX-A, and the orexin-2 receptor (OX₂) is capable to bind OX-A as well as OX-B. Orexin receptor antagonists are a novel type of nervous system or psychotropic drugs. Their mode of action in animals and humans involves either blockade of both orexin-1 and orexin-2 receptor (dual antagonists), or individual and selective blockade of either the orexin-1 or the orexin-2 receptor (selective antagonists) in the brain. Orexins were initially found to stimulate food consumption in rats suggesting a physiological role for these peptides as mediators in the central feedback mechanism that regulates feeding behaviour (Sakurai T. *et al.*, Cell, **1998**, 92, 573-585).

On the other hand, orexin neuropeptides and orexin receptors play an essential and central role in regulating circadian vigilance states. In the brain, orexin neurons collect sensory input about internal and external states and send short intrahypothalamic axonal projections as well as long projections to many other brain regions. The particular distribution of orexin fibers and receptors in basal forebrain, limbic structures and brainstem regions - areas related to the regulation of waking, sleep and emotional reactivity- suggests that orexins exert essential functions as regulators of behavioral arousal; by activating wake-promoting cell firing, orexins contribute to orchestrate all brain arousal systems that regulate circadian activity, energy balance and emotional reactivity. This role opens large therapeutic opportunities for medically addressing numerous mental health disorders possibly relating to orexinergic dysfunctions [see for example: Tsujino N and Sakurai T, "Orexin/hypocretin: a neuropeptide at the interface of sleep, energy homeostasis, and reward systems.", Pharmacol Rev. **2009**, 61:162-176; and Carter ME et al., "The brain hypocretins and their receptors: mediators of allostatic arousal.", Curr Op Pharmacol. **2009**, 9: 39-45] that are described in the following sections. It was also observed that orexins regulate states of sleep

and wakefulness opening potentially novel therapeutic approaches to insomnia and other sleep disorders (Chemelli R.M. *et al.*, Cell, **1999**, 98, 437-451).

Human memory is comprised of multiple systems that have different operating principles and different underlying neuronal substrates. The major distinction is between the capacity for conscious, declarative memory and a set of unconscious, non-declarative memory abilities. Declarative memory is further subdivided into semantic and episodic memory. Non-declarative memory is further subdivided into priming and perceptual learning, procedural memory for skills and habits, associative and non-associative learning, and some others. While semantic memory refers to the general knowledge about the world, episodic memory is autobiographical memory of events. Procedural memories refer to the ability to perform skill-based operations, as e.g. motor skills. Long-term memory is established during a multiple stage process through gradual changes involving diverse brain structures, beginning with learning, or memory acquisition, or formation. Subsequently, consolidation of what has been learned may stabilize memories. When long-term memories are retrieved, they may return to a labile state in which original content may be updated, modulated or disrupted. Subsequently, reconsolidation may again stabilize memories. At a late stage, long-term memory may be resistant to disruption. Long-term memory is conceptually and anatomically different from working memory, the latter of which is the capacity to maintain temporarily a limited amount of information in mind. Behavioural research has suggested that the human brain consolidates long-term memory at certain key time intervals. The initial phase of memory consolidation may occur in the first few minutes after we are exposed to a new idea or learning experience. The next, and possibly most important phase, may occur over a longer period of time, such as during sleep; in fact, certain consolidation processes have been suggested to be sleep-dependent [R. Stickgold *et al.*, Sleep-dependent memory consolidation; Nature **2005**, 437, 1272-1278]. Learning and memory processes are believed to be fundamentally affected in a variety of neurological and mental disorders, such as e.g. mental retardation, Alzheimer's disease or depression. Indeed, memory loss or impairment of memory acquisition is a significant feature of such diseases, and no effective therapy to prevent this detrimental process has emerged yet.

In addition, both anatomical and functional evidence from *in vitro* and *in vivo* studies suggest an important positive interaction of the endogenous orexin system with reward pathways of the brain [Aston-Jones G *et al.*, Brain Res **2010**, 1314, 74-90; Sharf R *et al.*, Brain Res **2010**, 1314, 130-138]. Selective pharmacological OXR-1 blockade reduced cue- and stress-induced reinstatement of cocaine seeking [Boutrel B, *et al.*, "Role for hypocretin in mediating stress-induced reinstatement of cocaine-seeking behavior." Proc Natl Acad Sci **2005**,

102(52), 19168-19173; Smith RJ et al., "Orexin/hypocretin signaling at the orexin 1 receptor regulates cue-elicited cocaine-seeking." *Eur J Neurosci* **2009**, 30(3), 493-503; Smith RJ et al., "Orexin/hypocretin is necessary for context-driven cocaine-seeking." *Neuropharmacology* 2010, 58(1), 179-184], cue-induced reinstatement of alcohol seeking [Lawrence AJ et al., *Br J Pharmacol* **2006**, 148(6), 752-759] and nicotine self-administration [Hollander JA et al., *Proc Natl Acad Sci* **2008**, 105(49), 19480-19485; LeSage MG et al., *Psychopharmacology* **2010**, 209(2), 203-212]. Orexin-1 receptor antagonism also attenuated the expression of amphetamine- and cocaine-induced CPP [Gozzi A et al., *PLoS One* **2011**, 6(1), e16406; Hutcheson DM et al., *Behav Pharmacol* **2011**, 22(2), 173-181], and reduced the expression or development of locomotor sensitization to amphetamine and cocaine [Borgland SL et al., *Neuron* **2006**, 49(4), 589-601; Quarta D et al., "The orexin-1 receptor antagonist SB-334867 reduces amphetamine-evoked dopamine outflow in the shell of the nucleus accumbens and decreases the expression of amphetamine sensitization." *Neurochem Int* **2010**, 56(1), 11-15].

The effect of a drug to diminish addictions may be modelled in normal or particularly sensitive mammals used as animal models [see for example Spealman et al, *Pharmacol. Biochem. Behav.* **1999**, 64, 327-336; or T.S. Shippenberg, G.F. Koob, "Recent advances in animal models of drug addiction" in *Neuropsychopharmacology: The fifth generation of progress*; K.L.Davis, D. Charney, J.T.Doyle, C. Nemeroff (eds.) **2002**; chapter 97, pages 1381-1397].

Several converging lines of evidence furthermore demonstrate a direct role of the orexin system as modulator of the acute stress response. For instance, stress (i.e. psychological stress or physical stress) is associated with increased arousal and vigilance which in turn is controlled by orexins [Sutcliffe, JG et al., *Nat Rev Neurosci* **2002**, 3(5), 339-349]. Orexin neurons are likely to be involved in the coordinated regulation of behavioral and physiological responses in stressful environments [Y. Kayaba et al., *Am. J. Physiol. Regul. Integr. Comp. Physiol.* **2003**, 285:R581-593]. Hypocretin/orexin contributes to the expression of some but not all forms of stress and arousal [Furlong T M et al., *Eur J Neurosci* **2009**, 30(8), 1603-1614]. Stress response may lead to dramatic, usually time-limited physiological, psychological and behavioural changes that may affect appetite, metabolism and feeding behavior [Chrousos, GP et al., *JAMA* **1992**, 267(9), 1244-1252]. The acute stress response may include behavioural, autonomic and endocrinological changes, such as promoting heightened vigilance, decreased libido, increased heart rate and blood pressure, or a redirection of blood flow to fuel the muscles, heart and the brain [Majzoub, JA et al., *European Journal of Endocrinology* **2006**, 155 (suppl_1) S71-S76].

As outlined above the orexin system regulates homeostatic functions such as sleep-wake cycle, energy balance, emotions and reward. Orexins are also involved in mediating the acute behavioral and autonomous nervous system response to stress [Zhang Wet al., "Multiple components of the defense response depend on orexin: evidence from orexin knockout mice and orexin neuron-ablated mice." *Auton Neurosci* **2006**, 126-127, 139-145].

Mood disorders including all types of depression and bipolar disorder are characterized by disturbed "mood" and feelings, as well as by sleeping problems (insomnia as well as hypersomnia), changes in appetite or weight and reduced pleasure and loss of interest in daily or once enjoyed activities [Liu X et al., *Sleep* **2007**, 30(1): 83-90]. Thus, there is a strong rationale that disturbances in the orexin system may contribute to the symptoms of mood disorders. Evidence in humans, for instance, exists that depressed patients show blunted diurnal variation in CSF orexin levels [Salomon RM et al., *Biol Psychiatry* **2003**, 54(2), 96-104]. In rodent models of depression, orexins were also shown to be involved. Pharmacological induction of a depressive behavioral state in rats, for instance, revealed an association with increased hypothalamic orexin levels [Feng P et al., *J Psychopharmacol* **2008**, 22(7): 784-791]. A chronic stress model of depression in mice also demonstrated an association of molecular orexin system disturbances with depressed behavioral states and a reversal of these molecular changes by antidepressant treatment [Nollet et al., *NeuroPharm* **2011**, 61(1-2):336-46].

The orexin system is also involved in stress-related appetitive/reward seeking behaviour (Berridge CW et al., *Brain Res* **2009**, 1314, 91-102). In certain instances, a modulatory effect on stress may be complementary to an effect on appetitive/reward seeking behaviour as such. For instance, an OX₁ selective orexin receptor antagonist was able to prevent footshock stress induced reinstatement of cocaine seeking behaviour [Boutrel, B et al., *Proc Natl Acad Sci* **2005**, 102(52), 19168-19173]. In addition, stress is also known to play an integral part in withdrawal which occurs during cessation of drug taking (Koob, GF et al., *Curr Opin Investig Drugs* **2010**, 11(1), 63-71).

Orexins have been found to increase food intake and appetite [Tsujino, N, Sakurai, T, *Pharmacol Rev* **2009**, 61(2) 162-176]. As an additional environmental factor, stress can contribute to binge eating behaviour, and lead to obesity [Adam, TC et al. *Physiol Behav* **2007**, 91(4) 449-458]. Animal models that are clinically relevant models of binge eating in humans are described for example in W. Foulds Mathes et al.; *Appetite* **2009**, 52, 545-553.

A number of recent studies report that orexins may play a role into several other important functions relating to arousal, especially when an organism must respond to unexpected

stressors and challenges in the environment [Tsujino N and Sakurai T. *Pharmacol Rev.* **2009**, 61:162-176; Carter ME, Borg JS and deLecea L., *Curr Op Pharmacol.* **2009**, 9: 39-45; C Boss, C Brisbare-Roch, F Jenck, *Journal of Medicinal Chemistry* **2009**, 52: 891-903]. The orexin system interacts with neural networks that regulate emotion, reward and energy homeostasis to maintain proper vigilance states. Dysfunctions in its function may thus relate to many mental health disorders in which vigilance, arousal, wakefulness or attention is disturbed.


The compound (2R)-2-((1S)-6,7-dimethoxy-1-[2-(4-trifluoromethyl-phenyl)-ethyl]-3,4-dihydro-1H-isoquinolin-2-yl)-N-methyl-2-phenyl-acetamide (WO2005/118548), a dual orexin receptor antagonist, showed clinical efficacy in humans when tested for the indication primary insomnia. In the rat, the compound has been shown to decrease alertness, characterized by decreases in both active wake and locomotion; and to dose-dependently increase the time spent in both REM and NREM sleep [Brisbare et al., *Nature Medicine* **2007**, 13, 150-155]. The compound further attenuated cardiovascular responses to conditioned fear and novelty exposure in rats [Furlong T M et al., *Eur J Neurosci* **2009**, 30(8), 1603-1614]. It is also active in an animal model of conditioned fear: the rat fear-potentiated startle paradigm (WO2009/047723) which relates to emotional states of fear and anxiety diseases such as anxieties including phobias and post traumatic stress disorders (PTSDs). In addition, intact declarative and non-declarative learning and memory has been demonstrated in rats treated with this compound [WO2007/105177, H Dietrich, F Jenck, *Psychopharmacology* **2010**, 212, 145-154]. Said compound furthermore decreased brain levels of amyloid-beta (A β) as well as A β plaque deposition after acute sleep restriction in amyloid precursor protein transgenic mice [JE Kang et al., "Amyloid-beta dynamics are regulated by orexin and the sleep-wake cycle.", *Science* **2009**, 326(5955): 1005-1007]. The accumulation of the A β in the brain extracellular space is hypothesized to be a critical event in the pathogenesis of Alzheimer's disease. The so-called and generally known "amyloid cascade hypothesis" links A β to Alzheimer's disease and, thus, to the cognitive dysfunction, expressed as impairment of learning and memory. The compound has also been shown to induce antidepressant-like activity in a mouse model of depression, when administered chronically [Nollet et al., *NeuroPharm* **2011**, 61(1-2):336-46]. Moreover, the compound has been shown to attenuate the natural activation induced by orexin A in fasted hungry rats exposed to food odors [MJ Prud'homme et al., *Neuroscience* **2009**, 162(4), 1287-1298]. The compound also displayed pharmacological activity in a rat model of nicotine self-administration [LeSage MG et al., *Psychopharmacology* **2010**, 209(2), 203-212]. Another dual orexin receptor antagonist, N-biphenyl-2-yl-1-[[1-methyl-1H-benzimidazol-2-yl)sulfanyl]acetyl]-L-prolinamide inhibited

nicotine-reinstatement for a conditioned reinforcer and reduced behavioral (locomotor sensitization) and molecular (transcriptional responses) changes induced by repeated amphetamine administration in rodents [Winrow et al., *Neuropharmacology* **2009**, 58(1),185-94].

5 Orexin receptor antagonists comprising a 2-substituted saturated cyclic amide derivatives (such as 2-substituted pyrrolidine-1-carboxamides) are known for example from WO2008/020405, WO2008/038251, WO2008/081399, WO2008/087611, WO2008/117241, WO2008/139416, WO2009/004584, WO2009/016560, WO2009/016564, WO2009/040730, WO2009/104155, WO2010/004507, WO2010/038200, WO2001/096302, WO2002/044172,
10 WO2002/089800, WO2002/090355, WO2003/002559, WO2003/032991, WO2003/041711, WO2003/051368, WO2003/051873, WO2004/026866, WO2004/041791, WO2004/041807, WO2004/041816, WO2009/003993, WO2009/003997, WO2009/124956, WO2010/060470, WO2010/060471, WO2010/060472, WO2010/063662, WO2010/063663, WO2010/072722, WO2010/122151, and WO2008/150364. A particular pyrrolidine derived compound is
15 disclosed in Langmead et. al, *Brit. J. Pharmacol.* **2004**, 141, 340-346 as being highly orexin-1 selective. WO2003/002561 discloses certain N-aroyl cyclic amine derivatives, encompassing benzimidazol-2-yl-methyl substituted pyrrolidine derivatives, as orexin receptor antagonists. Despite the great number of prior art compounds and their high structural variability, all compounds share a common structural feature, i.e. in position 2 of the saturated cyclic amide
20 a linker group such as at least a methylene group (or longer groups such as -CH₂-NH-CO-, -CH₂-NH-, -CH₂-O-, -CH₂-S-, etc.) link the cyclic amide to the respective aromatic ring system substituent. It has now surprisingly been found that, despite the substantial conformational changes that may be expected from the removal of a linker between two rigid structural elements, the present compounds, that have a benzimidazole ring directly attached to a
25 pyrrolidine amide in position 2, are potent orexin receptor antagonists.

The present invention, thus, provides novel benzimidazole-proline derivatives, which are non-peptide antagonists of human orexin receptors. These compounds are in particular of potential use in the treatment of disorders relating to orexinergic dysfunctions, comprising especially sleep disorders, anxiety disorders, addiction disorders, cognitive dysfunctions,
30 mood disorders, or appetite disorders.

1) A first aspect of the invention relates to compounds of the formula (I)



Formula (I)

wherein

5 **Ar¹** represents

- phenyl or 5- or 6-membered heteroaryl, wherein said phenyl or 5- or 6-membered heteroaryl independently is mono-, di-, or tri-substituted; wherein
 - one of said substituents is attached in *ortho*-position to the point of attachment of Ar¹ to the rest of the molecule; wherein said substituent is phenyl or 5- or 6-membered heteroaryl; wherein said phenyl or 5- or 6-membered heteroaryl substituent is independently unsubstituted, mono-, di-, or tri-substituted, wherein the substituents are independently selected from (C₁₋₄)alkyl, (C₁₋₄)alkoxy, halogen, cyano, (C₁₋₃)fluoroalkyl, and (C₁₋₃)fluoroalkoxy; or said *ortho* substituent is benzo[1,3]dioxolyl, or 2-(3-methoxy-phenyl)-ethynyl;
 - and the other of said substituents, if present, is/are independently selected from (C₁₋₄)alkyl; (C₁₋₄)alkoxy; (C₃₋₆)cycloalkyl; halogen; cyano; (C₁₋₃)fluoroalkyl; (C₁₋₃)fluoroalkoxy; -NR¹⁰R¹¹, wherein R¹⁰ and R¹¹ independently represent hydrogen or (C₁₋₄)alkyl, or R¹⁰ and R¹¹ together with the nitrogen to which they are attached to form a pyrrolidine ring; unsubstituted pyridinyl; and phenyl which is unsubstituted, or mono- or di-substituted, wherein the substituents are independently selected from (C₁₋₄)alkyl, (C₁₋₄)alkoxy, cyano, and halogen;

one of **R²** and **R³** represents hydrogen, and the other represents hydrogen or (C₁₋₄)alkyl (notably one of **R²** and **R³** represents hydrogen, and the other represents hydrogen or methyl); and

one of **R^{4a}** and **R^{4b}** represents hydrogen, and the other represents hydrogen, (C₁₋₄)alkoxy (especially methoxy), or halogen (especially fluorine); or **R^{4a}** and **R^{4b}** together represent a group H₂C=; or both **R^{4a}** and **R^{4b}** represent fluorine; wherein, in case **R³** is different from hydrogen, both **R^{4a}** and **R^{4b}** represent hydrogen;

R¹ represents hydrogen, (C₁₋₄)alkyl (especially methyl or ethyl), (C₃₋₆)cycloalkyl-(CH₂)- (especially cyclopropyl-methyl), (C₂₋₃)fluoroalkyl (especially 2-fluoro-ethyl), or (C₁₋₄)alkoxy-(C₂₋₄)alkyl (especially 2-methoxy-ethyl); and

(**R**⁵)_n represents one to three optional substituents (i.e. n represents the integer 0, 1, 2, or 3)

- 5 independently selected from (C₁₋₄)alkyl, (C₁₋₄)alkoxy (especially methoxy), halogen, (C₁₋₄)alkyl-thio- (especially H₃C-S-), (C₁₋₃)fluoroalkyl (especially trifluoromethyl), (C₁₋₃)fluoroalkoxy (especially trifluoromethoxy), (C₁₋₃)fluoroalkyl-thio- (especially F₃C-S-), hydroxy-(C₁₋₄)alkyl- (especially HO-CH₂-), (C₁₋₄)alkoxy-carbonyl- (especially H₃CO-CO-), nitro, hydroxy, and cyano; or (**R**⁵)_n represents a group -O-CH₂-CH₂-O-; or (**R**⁵)_n represents a
10 fused phenyl group which, together with the benzimidazole moiety to which it is fused to, forms a 1H-naphtho[2,3-d]imidazol-2-yl group;

with the exception of

[2-(1H-benzimidazol-2-yl)-1-pyrrolidinyl][2-(1H-pyrazol-1-yl)phenyl]-methanone (CAS Reg. No. 1293846-63-5);

[2-(1H-benzimidazol-2-yl)-1-pyrrolidinyl][5-(2,5-dimethyl-1H-pyrrol-1-yl)-1-methyl-1H-pyrazol-4-yl]-methanone

- 15 (CAS Reg. No. 1288543-08-7);

[2-(1H-benzimidazol-2-yl)-1-pyrrolidinyl][1,1'-biphenyl]-2-yl-methanone (CAS Reg. No. 1277849-61-2);

[2-(6-methyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl][3-phenyl-isoxazol-4-yl]-methanone (CAS Reg. No. 1413410-98-6);

- 20 [2-(1H-imidazol-2-yl)phenyl][2-(6-methyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone (CAS Reg. No. 1378205-71-0);


[2-(6-methyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl][3-(thien-2-yl)-1H-pyrazol-4-yl]-methanone (CAS Reg. No. 1377970-45-0);

[2-(6-methyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl][3-phenyl-1H-pyrazol-4-yl]-methanone (CAS Reg. No. 1377872-25-7);

- 25 [2-(1H-benzimidazol-2-yl)-pyrrolidin-1-yl][3-cyclopropyl-1-phenyl-1H-pyrazol-5-yl]-methanone (CAS Reg. No. 1331048-98-5); and

[2-(1H-benzimidazol-2-yl)-pyrrolidin-1-yl][1-(2-fluorophenyl)-5-(1H-pyrrol-1-yl)-1H-pyrazol-4-yl]-methanone (CAS Reg. No. 1290361-06-6).

2) A second aspect of the invention relates to compounds of the formula (I) according to embodiment 1), which are also compounds of the formula (II); wherein the absolute configuration is as depicted in formula (II):



Formula (II)

wherein

Ar¹ represents

- phenyl or 5- or 6-membered heteroaryl, wherein said phenyl or 5- or 6-membered heteroaryl independently is mono-, di-, or tri-substituted; wherein

➤ one of said substituents is attached in *ortho*-position to the point of attachment of Ar¹ to the rest of the molecule; wherein said substituent is phenyl or 5- or 6-membered heteroaryl; wherein said phenyl or 5- or 6-membered heteroaryl substituent is independently unsubstituted, mono-, di-, or tri-substituted, wherein the substituents are independently selected from (C₁₋₄)alkyl, (C₁₋₄)alkoxy, halogen, cyano, (C₁₋₃)fluoroalkyl, and (C₁₋₃)fluoroalkoxy; or said *ortho* substituent is benzo[1,3]dioxolyl, or 2-(3-methoxy-phenyl)-ethynyl;

➤ and the other of said substituents, if present, is/are independently selected from (C₁₋₄)alkyl; (C₁₋₄)alkoxy; (C₃₋₆)cycloalkyl; halogen; cyano; (C₁₋₃)fluoroalkyl; (C₁₋₃)fluoroalkoxy; -NR¹⁰R¹¹, wherein R¹⁰ and R¹¹ independently represent hydrogen or (C₁₋₄)alkyl, or R¹⁰ and R¹¹ together with the nitrogen to which they are attached to form a pyrrolidine ring; unsubstituted pyridinyl; and phenyl which is unsubstituted, or mono- or di-substituted, wherein the substituents are independently selected from (C₁₋₄)alkyl, (C₁₋₄)alkoxy, cyano, and halogen;

one of **R²** and **R³** represents hydrogen, and the other represents hydrogen or (C₁₋₄)alkyl (notably one of **R²** and **R³** represents hydrogen, and the other represents hydrogen or methyl); and

one of **R^{4a}** and **R^{4b}** represents hydrogen, and the other represents hydrogen, (C₁₋₄)alkoxy (especially methoxy), or halogen (especially fluorine); or **R^{4a}** and **R^{4b}** together represent a

group $\text{H}_2\text{C=}$; or both R^{4a} and R^{4b} represent fluorine; wherein, in case R^3 is different from hydrogen, both R^{4a} and R^{4b} represent hydrogen;


R^1 represents hydrogen, (C_{1-4}) alkyl (especially methyl or ethyl), (C_{3-6}) cycloalkyl- (CH_2) - (especially cyclopropyl-methyl), (C_{2-3}) fluoroalkyl (especially 2-fluoro-ethyl), or (C_{1-4}) alkoxy-
5 (C_{2-4}) alkyl (especially 2-methoxy-ethyl); and

$(\text{R}^5)_n$ represents one to three optional substituents (i.e. n represents the integer 0, 1, 2, or 3) independently selected from (C_{1-4}) alkyl (especially methyl), (C_{1-4}) alkoxy (especially methoxy), halogen (especially fluorine, chlorine or bromine), (C_{1-4}) alkyl-thio- (especially $\text{H}_3\text{C-S-}$), (C_{1-3}) fluoroalkyl (especially trifluoromethyl), (C_{1-3}) fluoroalkoxy (especially trifluoromethoxy),
10 (C_{1-3}) fluoroalkyl-thio- (especially $\text{F}_3\text{C-S-}$), hydroxy- (C_{1-4}) alkyl- (especially HO-CH_2 -), (C_{1-4}) alkoxy-carbonyl- (especially $\text{H}_3\text{CO-CO-}$), nitro, hydroxy, and cyano; or $(\text{R}^5)_n$ represents a group $-\text{O-CH}_2\text{-CH}_2\text{-O-}$; or $(\text{R}^5)_n$ represents a fused phenyl group which, together with the benzimidazole moiety to which it is fused to, forms a 1H-naphtho[2,3-d]imidazol-2-yl group;

with the exception of

15 $[(\text{S})\text{-2-(1H-benzimidazol-2-yl)-1-pyrrolidinyl}][2\text{-(1H-pyrazol-1-yl)phenyl}]\text{-methanone}$;
 $[(\text{S})\text{-2-(1H-benzimidazol-2-yl)-1-pyrrolidinyl}][5\text{-(2,5-dimethyl-1H-pyrrol-1-yl)-1-methyl-1H-pyrazol-4-yl}]\text{-methanone}$;
 $[(\text{S})\text{-2-(1H-benzimidazol-2-yl)-1-pyrrolidinyl}][1,1'\text{-biphenyl}]\text{-2-yl-methanone}$;
 $[(\text{S})\text{-2-(6-methyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl}][3\text{-phenyl-isoxazol-4-yl}]\text{-methanone}$;
 $[(\text{S})\text{-2-(1H-imidazol-2-yl)phenyl}][2\text{-(6-methyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl}]\text{-methanone}$;
20 $[(\text{S})\text{-2-(6-methyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl}][3\text{-(thien-2-yl)-1H-pyrazol-4-yl}]\text{-methanone}$;
 $[(\text{S})\text{-2-(6-methyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl}][3\text{-phenyl-1H-pyrazol-4-yl}]\text{-methanone}$;
 $[(\text{S})\text{-2-(1H-benzimidazol-2-yl)-pyrrolidin-1-yl}][3\text{-cyclopropyl-1-phenyl-1H-pyrazol-5-yl}]\text{-methanone}$; and
 $[(\text{S})\text{-2-(1H-benzimidazol-2-yl)-pyrrolidin-1-yl}][1\text{-(2-fluorophenyl)-5-(1H-pyrrol-1-yl)-1H-pyrazol-4-yl}]\text{-methanone}$.

3) A third aspect of the invention relates to compounds of the formula (II) according to embodiment 2), which are also compounds of the formula (III); wherein the absolute configuration is as depicted in formula (III):



Formula (III)

wherein

Ar¹ represents

- phenyl or 5- or 6-membered heteroaryl, wherein said phenyl or 5- or 6-membered heteroaryl independently is mono-, di-, or tri-substituted; wherein

➤ one of said substituents is attached in *ortho*-position to the point of attachment of Ar¹ to the rest of the molecule; wherein said substituent is phenyl or 5- or 6-membered heteroaryl; wherein said phenyl or 5- or 6-membered heteroaryl substituent is independently unsubstituted, mono-, di-, or tri-substituted, wherein the substituents are independently selected from (C₁₋₄)alkyl, (C₁₋₄)alkoxy, halogen, cyano, (C₁₋₃)fluoroalkyl, and (C₁₋₃)fluoroalkoxy; or said *ortho* substituent is benzo[1,3]dioxolyl;

➤ and the other of said substituents, if present, is/are independently selected from (C₁₋₄)alkyl; (C₁₋₄)alkoxy; (C₃₋₆)cycloalkyl; halogen; cyano; (C₁₋₃)fluoroalkyl; (C₁₋₃)fluoroalkoxy; and -NR¹⁰R¹¹, wherein R¹⁰ and R¹¹ independently represent hydrogen or (C₁₋₄)alkyl, or R¹⁰ and R¹¹ together with the nitrogen to which they are attached to form a pyrrolidine ring;

one of **R²** and **R³** represents hydrogen, and the other represents hydrogen or (C₁₋₄)alkyl (notably one of **R²** and **R³** represents hydrogen, and the other represents hydrogen or methyl); and

R^{4a} and **R^{4b}** independently represent hydrogen or halogen (especially fluorine); or **R^{4a}** represents (C₁₋₄)alkoxy (especially methoxy) and **R^{4b}** represents hydrogen; or **R^{4a}** and **R^{4b}** together represent a group H₂C=;

wherein, in case **R³** is different from hydrogen, both **R^{4a}** and **R^{4b}** represent hydrogen;

R^1 represents hydrogen, or (C_{1-4}) alkyl (especially methyl or ethyl); and

R^{14} , R^{15} , R^{16} and R^{17} together represent one to three optional substituents (i.e. at least one of R^{14} , R^{15} , R^{16} and R^{17} is hydrogen) [notably R^{14} , R^{15} , R^{16} and R^{17} together represent one to three substituents (i.e. at least one of R^{14} , R^{15} , R^{16} and R^{17} is hydrogen and at least one of R^{14} , R^{15} , R^{16} and R^{17} is different from hydrogen)], wherein

- R^{14} and R^{17} independently represent hydrogen, (C_{1-4}) alkyl, (C_{1-4}) alkoxy (especially methoxy), (C_{1-4}) alkyl-thio- (especially H_3C-S-), halogen, (C_{1-3}) fluoroalkyl (especially trifluoromethyl), (C_{1-4}) alkoxy-carbonyl- (especially $H_3CO-CO-$), hydroxy- (C_{1-4}) alkyl- (especially $HO-CH_2-$), hydroxy, or nitro; and
- R^{15} and R^{16} independently represent hydrogen, (C_{1-4}) alkyl, (C_{1-4}) alkoxy (especially methoxy), (C_{1-4}) alkyl-thio- (especially H_3C-S-), halogen, (C_{1-3}) fluoroalkyl (especially trifluoromethyl), (C_{1-3}) fluoroalkoxy (especially trifluoromethoxy), (C_{1-3}) fluoroalkyl-thio- (especially F_3C-S-), hydroxy- (C_{1-4}) alkyl- (especially $HO-CH_2-$), or cyano;

or R^{14} and R^{15} together, or R^{16} and R^{17} together, represent a group $-O-CH_2-CH_2-O-$;

or R^{15} and R^{16} together represent a fused phenyl group which, together with the benzimidazole moiety to which it is fused to, forms a 1H-naphtho[2,3-d]imidazol-2-yl group;

with the exception of

[(S)-2-(1H-benzimidazol-2-yl)-1-pyrrolidinyl][2-(1H-pyrazol-1-yl)phenyl]-methanone;

[(S)-2-(1H-benzimidazol-2-yl)-1-pyrrolidinyl][5-(2,5-dimethyl-1H-pyrrol-1-yl)-1-methyl-1H-pyrazol-4-yl]-methanone;

[(S)-2-(1H-benzimidazol-2-yl)-1-pyrrolidinyl][1,1'-biphenyl]-2-yl-methanone;

wherein, in a sub-embodiment, in addition to the above-listed three compounds also the following compounds are excluded from the scope of the compounds of embodiment 3):

[(S)-2-(6-methyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl][3-phenyl-isoxazol-4-yl]-methanone;

[(S)-2-(1H-imidazol-2-yl)phenyl][2-(6-methyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;

[(S)-2-(6-methyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl][3-(thien-2-yl)-1H-pyrazol-4-yl]-methanone;

[(S)-2-(6-methyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl][3-phenyl-1H-pyrazol-4-yl]-methanone;

[(S)-2-(1H-benzimidazol-2-yl)-pyrrolidin-1-yl][3-cyclopropyl-1-phenyl-1H-pyrazol-5-yl]-methanone; and

[(S)-2-(1H-benzimidazol-2-yl)-pyrrolidin-1-yl][1-(2-fluorophenyl)-5-(1H-pyrrol-1-yl)-1H-pyrazol-4-yl]-methanone.

The compounds of formula (I) contain at least one stereogenic center which is situated in position 2 of the pyrrolidine moiety. Preferably, the absolute configuration of the pyrrolidine moiety of the present compounds, especially the absolute configuration of said chiral center in position 2 of the pyrrolidine moiety, is as depicted in formula (II) and (III) of embodiments 2) or 3); i.e. for example for R^2 being the substituent of lowest priority (e.g. when R^2 is hydrogen or methyl), said chiral center is preferably in absolute (S) configuration. In addition,


the compounds of formulae (I), (II), (III), and the compounds of formulae (IV), (V), and (VI) below, may contain one or more further stereogenic or asymmetric centers, such as one or more asymmetric carbon atoms. The compounds of formulae (I), (II), (III), (IV), (V), and (VI) may thus be present as mixtures of stereoisomers or preferably as pure stereoisomers.

5 Mixtures of stereoisomers may be separated in a manner known to a person skilled in the art.

In addition, it is well understood that, in case the benzimidazole moiety of the present compounds is unsubstituted on the ring nitrogen having a free valency (i.e. R¹ represents hydrogen) such benzimidazole moiety represents tautomeric forms. Thus, substituents (R⁵)_n of the benzimidazole moiety may be attached in the position(s) *ortho* to the bridgehead atoms (i.e. attached in position(s) 4 and/or 7, corresponding to R¹⁴ and/or R¹⁷), and/or in the position(s) *meta* to the bridgehead atoms, (i.e. attached in position(s) 5 and/or 6, corresponding to R¹⁵ and/or R¹⁶). It is understood that the two *ortho*, and, respectively, the two *meta* positions are considered equivalent. For example, the group 4-methyl-1H-benzoimidazol-2-yl is understood to signify the same group as 7-methyl-1H-benzoimidazol-2-yl and 4-methyl-3H-benzoimidazol-2-yl and 7-methyl-3H-benzoimidazol-2-yl.

The present invention also includes isotopically labelled, especially ²H (deuterium) labelled compounds of formulae (I), (II), (III), (IV), (V), and (VI), which compounds are identical to the compounds of formulae (I), (II), (III), (IV), (V), and (VI) except that one or more atoms have each been replaced by an atom having the same atomic number but an atomic mass different from the atomic mass usually found in nature. Isotopically labelled, especially ²H (deuterium) labelled compounds of formulae (I), (II), (III), (IV), (V), and (VI) and salts thereof are within the scope of the present invention. Substitution of hydrogen with the heavier isotope ²H (deuterium) may lead to greater metabolic stability, resulting e.g. in increased *in-vivo* half-life or reduced dosage requirements, or may lead to reduced inhibition of cytochrome P450 enzymes, resulting e.g. in an improved safety profile. In one embodiment of the invention, the compounds of formulae (I), (II), (III), (IV), (V), and (VI) are not isotopically labelled, or they are labelled only with one or more deuterium atoms. In a sub-embodiment, the compounds of formulae (I), (II), (III), (IV), (V), and (VI) are not isotopically labelled at all. Isotopically labelled compounds of formulae (I), (II), (III), (IV), (V), and (VI) be prepared in analogy to the methods described hereinafter, but using the appropriate isotopic variation of suitable reagents or starting materials.

In this patent application, a dotted line shows the point of attachment of the radical drawn. For example, the radical drawn below



represents a 2-(2-triazolyl)-phenyl group.

Where the plural form is used for compounds, salts, pharmaceutical compositions, diseases and the like, this is intended to mean also a single compound, salt, or the like.

- 5 Any reference to compounds of formulae (I), (II), (III), (IV), (V), and (VI) according to any one of embodiments 1) to 57) is to be understood as referring also to the salts (and especially the pharmaceutically acceptable salts) of such compounds, as appropriate and expedient.

The term "pharmaceutically acceptable salts" refers to salts that retain the desired biological activity of the subject compound and exhibit minimal undesired toxicological effects. Such salts include inorganic or organic acid and/or base addition salts depending on the presence of basic and/or acidic groups in the subject compound. For reference see for example "Salt selection for basic drugs", *Int. J. Pharm.* (1986), **33**, 201-217; "Handbook of Pharmaceutical Salts. Properties, Selection and Use.", P. Heinrich Stahl, Camille G. Wermuth (Eds.), Wiley-VCH, 2008; and "Pharmaceutical Salts and Co-crystals", Johan Wouters and Luc Quéré (Eds.), RSC Publishing, 2012.

Definitions provided herein are intended to apply uniformly to the compounds of formulae (I), (II), (III), (IV), (V) and (VI) as defined in any one of embodiments 1) to 45), and, *mutatis mutandis*, throughout the description and the claims unless an otherwise expressly set out definition provides a broader or narrower definition. It is well understood that a definition or preferred definition of a term defines and may replace the respective term independently of (and in combination with) any definition or preferred definition of any or all other terms as defined herein.

The term "halogen" means fluorine, chlorine, or bromine, preferably fluorine or chlorine.

The term "alkyl", used alone or in combination, refers to a saturated straight or branched chain alkyl group containing one to six carbon atoms. The term "(C_{x-y})alkyl" (x and y each being an integer), refers to an alkyl group as defined before, containing x to y carbon atoms. For example a (C₁₋₄)alkyl group contains from one to four carbon atoms. Examples of alkyl groups are methyl, ethyl, propyl, isopropyl, n-butyl, isobutyl, sec.-butyl and tert.-butyl. Preferred are methyl and ethyl. Most preferred is methyl.

30 The term "alkoxy", used alone or in combination, refers to an alkyl-O- group wherein the alkyl group is as defined before. The term "(C_{x-y})alkoxy" (x and y each being an integer) refers to

an alkoxy group as defined before containing x to y carbon atoms. For example a (C₁₋₄)alkoxy group means a group of the formula (C₁₋₄)alkyl-O- in which the term "(C₁₋₄)alkyl" has the previously given significance. Examples of alkoxy groups are methoxy, ethoxy, n-propoxy, isopropoxy, n-butoxy, isobutoxy, sec.-butoxy and tert.-butoxy. Preferred are ethoxy and especially methoxy.

The term "(C₁₋₄)alkyl-thio-" used alone or in combination, refers to a group of the formula (C₁₋₄)alkyl-S- in which the term "(C₁₋₄)alkyl" has the previously given significance. An example is CH₃-S-.

The term "fluoroalkyl" refers to an alkyl group as defined before containing one to three carbon atoms in which one or more (and possibly all) hydrogen atoms have been replaced with fluorine. The term "(C_{x-y})fluoroalkyl" (x and y each being an integer) refers to a fluoroalkyl group as defined before containing x to y carbon atoms. For example a (C₁₋₃)fluoroalkyl group contains from one to three carbon atoms in which one to seven hydrogen atoms have been replaced with fluorine. Representative examples of fluoroalkyl groups include trifluoromethyl, 2-fluoroethyl, 2,2-difluoroethyl and 2,2,2-trifluoroethyl. Preferred are (C₁)fluoroalkyl groups such as trifluoromethyl.

The term "fluoroalkoxy" refers to an alkoxy group as defined before containing one to three carbon atoms in which one or more (and possibly all) hydrogen atoms have been replaced with fluorine. The term "(C_{x-y})fluoroalkoxy" (x and y each being an integer) refers to a fluoroalkoxy group as defined before containing x to y carbon atoms. For example a (C₁₋₃)fluoroalkoxy group contains from one to three carbon atoms in which one to seven hydrogen atoms have been replaced with fluorine. Representative examples of fluoroalkoxy groups include trifluoromethoxy, difluoromethoxy, 2-fluoroethoxy, 2,2-difluoroethoxy and 2,2,2-trifluoroethoxy. Preferred are (C₁)fluoroalkoxy groups such as trifluoromethoxy and difluoromethoxy.

The term "(C₁₋₃)fluoroalkyl-thio-" refers to (C₁₋₃)fluoroalkyl group as defined before, which is linked to the rest of the molecule through a sulfur atom. An example is CF₃-S-.

In case two of (R⁵)_n together form a group -O-CH₂-CH₂-O-, such group together with the benzimidazole moiety especially forms a 7,8-dihydro-3H-6,9-dioxo-1,3-diazacyclopenta[a]naphthalen-2-yl group.

Particular examples of Ar¹ representing a phenyl group, wherein said phenyl is mono-, di-, or tri-substituted; wherein one of said substituents is attached in *ortho*-position to the point of attachment of Ar¹ to the rest of the molecule; are such that the other of said substituents, if

present, is/are independently selected from (C₁₋₄)alkyl; (C₁₋₄)alkoxy; (C₃₋₆)cycloalkyl; halogen; cyano; (C₁₋₃)fluoroalkyl; (C₁₋₃)fluoroalkoxy; unsubstituted pyridinyl; and phenyl which is unsubstituted, or mono- or di-substituted, wherein the substituents are independently selected from (C₁₋₄)alkyl, (C₁₋₄)alkoxy, cyano, and halogen [notably the other of said substituents, if present, is/are independently selected from (C₁₋₄)alkyl, (C₁₋₄)alkoxy, halogen, cyano, (C₁₋₃)fluoroalkyl, and (C₁₋₃)fluoroalkoxy; especially from (C₁₋₄)alkyl, (C₁₋₄)alkoxy, and halogen]. Particular examples of such phenyl groups which are further substituted in *ortho* position as used for the group **Ar**¹ are 1,2-phenylene, 4-methyl-1,2-phenylene, 5-methyl-1,2-phenylene, 6-methyl-1,2-phenylene, 4,5-dimethyl-1,2-phenylene, 5-fluoro-1,2-phenylene, 6-fluoro-1,2-phenylene, 5-chloro-1,2-phenylene, 5-cyano-1,2-phenylene, 5-methoxy-1,2-phenylene, 4,5-dimethoxy-1,2-phenylene, 5-trifluoromethyl-1,2-phenylene, 5-trifluoromethoxy-1,2-phenylene, 6-fluoro-5-methyl-1,2-phenylene, and 6-fluoro-5-methoxy-1,2-phenylene; and in addition to the above listed groups: 5-bromo-1,2-phenylene, 3,4-dimethyl-1,2-phenylene, 5-(pyridine-3-yl)-1,2-phenylene, and 5-(3-cyano-phenyl)-1,2-phenylene; wherein in the above groups the carbonyl group is attached in position 1.

Examples of the particular phenyl groups which are substituents of the group **Ar**¹ are especially phenyl groups which are unsubstituted, mono-, or di-substituted, wherein the substituents are independently selected from (C₁₋₄)alkyl, (C₁₋₄)alkoxy, halogen, cyano, (C₁₋₃)fluoroalkyl, and (C₁₋₃)fluoroalkoxy [notably from (C₁₋₄)alkyl, (C₁₋₄)alkoxy, halogen, and (C₁₋₃)fluoroalkyl]. Particular examples are phenyl, 2-methyl-phenyl, 3-methyl-phenyl, 4-methyl-phenyl, 3,4-dimethyl-phenyl, 3-methoxy-phenyl, 2-fluoro-phenyl, 3-fluoro-phenyl, 4-fluoro-phenyl, 3,5-difluoro-phenyl, 4-fluoro-3-methyl-phenyl, 3-fluoro-4-methyl-phenyl, 3-chloro-phenyl, 4-chloro-phenyl, 2,3-dichloro-phenyl, 3,4-dichloro-phenyl, 4-bromo-phenyl, 3-trifluoromethyl-phenyl, 4-trifluoromethyl-phenyl; and in addition to the above-listed groups: 2-chloro-phenyl, 3,4-difluoro-phenyl, 2-methoxy-phenyl, 4-methoxy-phenyl, 3-cyano-phenyl, 4-cyano-phenyl, 3-trifluoromethoxy-phenyl, and 4-trifluoromethoxy-phenyl.

The term "heteroaryl", if not explicitly stated otherwise, means a 5- to 10-membered monocyclic, or bicyclic, aromatic ring containing 1 to a maximum of 3 heteroatoms independently selected from oxygen, nitrogen and sulfur. Examples of such heteroaryl groups are 5-membered monocyclic heteroaryl groups such as furanyl, oxazolyl, isoxazolyl, oxadiazolyl, thienyl, thiazolyl, isothiazolyl, thiadiazolyl, pyrrolyl, imidazolyl, pyrazolyl, and triazolyl; 6-membered monocyclic heteroaryl such as pyridyl, pyrimidyl, pyridazinyl, and pyrazinyl; and 8- to 10-membered bicyclic heteroaryl such as indolyl, isoindolyl, benzofuranyl, isobenzofuranyl, benzothiophenyl, indazolyl, benzimidazolyl, benzoxazolyl (or benzooxazolyl), benzisoxazolyl, benzothiazolyl, benzoisothiazolyl, benzotriazolyl,

benzoxadiazolyl, benzothiadiazolyl, quinolinyl, isoquinolinyl, naphthyridinyl, cinnolinyl, quinazolinyl, quinoxalinyl, phthalazinyl, pyrazolo[1,5-a]pyridyl, pyrazolo[1,5-a]pyrimidyl, imidazo[1,2-a]pyridyl, 1H-pyrrolo[3,2-b]pyridyl, 1H-pyrrolo[2,3-b]pyridyl, pyrrolo[3,2-d]pyrimidinyl, pyrrolo[2,3-d]pyrimidinyl, 4H-furo[3,2-b]pyrrolyl, pyrrolo[2,1-b]thiazolyl, imidazo[2,1-b]thiazolyl and purinyl.

Examples of the particular 5- or 6-membered heteroaryl groups which are further substituted in *ortho* position as used for the group **Ar**¹ are the above mentioned 5- or 6-membered heteroaryl groups, notably the 5-membered heteroaryl groups oxazolyl, isoxazolyl, thiophenyl, thiazolyl, isothiazolyl; and the 6-membered heteroaryl groups pyridyl, pyrimidyl and pyrazinyl. In a sub-embodiment, examples are oxazolyl (notably oxazol-4,5-diyl (in particular 2-methyl-oxazol-4,5-diyl), isoxazolyl (notably isoxazol-3,4-diyl, in particular 5-methyl-isoxazol-3,4-diyl), thiazolyl (notably thiazol-4,5-diyl, in particular 2-methyl-thiazol-4,5-diyl, 2-cyclopropyl-thiazol-4,5-diyl, 2-dimethylamino-thiazol-4,5-diyl, 2-(1-pyrrolidinyl)-thiazol-4,5-diyl) and thiophenyl (notably thiophen-2,3-diyl, in particular thiophen-2,3-diyl, 5-methyl-thiophen-2,3-diyl); as well as pyridyl (notably pyridin-2,3-diyl, in particular pyridin-2,3-diyl, 6-methyl-pyridin-2,3-diyl), pyrimidyl (notably pyrimidin-4,5-diyl, in particular pyrimidin-4,5-diyl, 2-methyl-pyrimidin-4,5-diyl), and pyrazinyl (notably pyrazin-2,3-diyl, in particular pyrazin-2,3-diyl). These groups are at least mono-substituted in ortho position, and preferably and independently carry no further substituent or one further substituent as explicitly defined. In particular such optional further substituent may independently be selected from (C₁₋₄)alkyl (especially methyl); (C₃₋₆)cycloalkyl (especially cyclopropyl); -NR¹⁰R¹¹, wherein R¹⁰ and R¹¹ independently represent (C₁₋₄)alkyl (especially methyl), or R¹⁰ and R¹¹ together with the nitrogen to which they are attached to form a pyrrolidine ring; and phenyl which is unsubstituted, or mono- or di-substituted, wherein the substituents are independently selected from (C₁₋₄)alkyl, (C₁₋₄)alkoxy, cyano, and halogen. In a sub-embodiment, such optional further substituent may independently be selected from (C₁₋₄)alkyl, (C₁₋₄)alkoxy, (C₃₋₆)cycloalkyl, halogen, cyano, (C₁₋₃)fluoroalkyl, (C₁₋₃)fluoroalkoxy, and -NR¹⁰R¹¹ [wherein especially oxazolyl groups carry no further substituent or one further substituent selected from (C₁₋₄)alkyl, (C₃₋₆)cycloalkyl, -NR¹⁰R¹¹; thiazolyl groups carry no further substituent or one further substituent selected from (C₁₋₄)alkyl, (C₃₋₆)cycloalkyl, -NR¹⁰R¹¹, and phenyl which is unsubstituted, or mono- or di-substituted, wherein the substituents are independently selected from (C₁₋₄)alkyl, (C₁₋₄)alkoxy, cyano, and halogen (notably methyl, cyclopropyl, and phenyl which is unsubstituted, or mono- or di-substituted, wherein the substituents are independently selected from (C₁₋₄)alkyl, (C₁₋₄)alkoxy, cyano, and halogen; especially methyl and cyclopropyl); thiophenyl, pyridinyl and pyrimidinyl groups carry no further substituent or

one further substituent selected from (C₁₋₄)alkyl (especially methyl); and pyrazinyl group carry no further substituent]. The above groups are preferably attached to the rest of the molecule (i.e. the carbonyl group) in position 4 of oxazolyl, isoxazolyl, or thiazolyl groups; in position 2 of pyridyl or pyrazinyl groups; in position 2 of thiophenyl groups; and in position 5 of pyrimidinyl groups. In a further sub-embodiment, particular examples of such groups are 2-methyl-thiazol-4,5-diyl, 2-cyclopropyl-thiazol-4,5-diyl, 2-dimethylamino-thiazol-4,5-diyl, thiophen-2,3-diyl; as well as 6-methyl-pyridin-2,3-diyl, 6-methyl-pyridin-3,4-diyl, 2-methyl-pyrimidin-4,5-diyl, pyrimidin-4,5-diyl, and pyrazin-2,3-diyl. Further particular examples are 2-(4-fluoro-2-methoxy-phenyl)-thiazol-4,5-diyl, 2-(2-methoxy-phenyl)-thiazol-4,5-diyl, 2-(3-methoxy-phenyl)-thiazol-4,5-diyl, 2-(4-methoxy-phenyl)-thiazol-4,5-diyl, 2-(3,4-dimethyl-phenyl)-thiazol-4,5-diyl, 2-(2,3-difluoro-phenyl)-thiazol-4,5-diyl, 2-(2,5-difluoro-phenyl)-thiazol-4,5-diyl, 2-(4-fluoro-2-methyl-phenyl)-thiazol-4,5-diyl, 2-(4-fluoro-3-methyl-phenyl)-thiazol-4,5-diyl, 2-(2-fluoro-4-methyl-phenyl)-thiazol-4,5-diyl, 2-(3-chloro-4-methyl-phenyl)-thiazol-4,5-diyl, 2-(3-cyano-phenyl)-thiazol-4,5-diyl, 2-(2-methyl-phenyl)-thiazol-4,5-diyl, 2-(3-methyl-phenyl)-thiazol-4,5-diyl, 2-(4-methyl-phenyl)-thiazol-4,5-diyl, 2-(2-fluoro-phenyl)-thiazol-4,5-diyl, 2-(3-fluoro-phenyl)-thiazol-4,5-diyl, 2-(4-fluoro-phenyl)-thiazol-4,5-diyl, 2-(2-chloro-phenyl)-thiazol-4,5-diyl, 2-(3-chloro-phenyl)-thiazol-4,5-diyl, and 2-phenyl)-thiazol-4,5-diyl.

Examples of the particular 5- or 6-membered heteroaryl groups which are substituents of the group **Ar**¹ are the above mentioned 5- or 6-membered heteroaryl groups; notably the 5-membered heteroaryl groups oxazolyl, isoxazolyl, oxadiazolyl, thienyl, thiazolyl, isothiazolyl, thiadiazolyl, imidazolyl, pyrazolyl, triazolyl, and the 6-membered heteroaryl groups pyridyl, pyrimidyl, pyrazinyl and pyridazinyl. In a sub-embodiment, such groups are especially pyrazolyl, triazolyl, pyridinyl and pyrimidinyl, notably pyrazol-1-yl, pyrimidin-2-yl, and [1,2,3]triazol-2-yl. The above mentioned groups may be unsubstituted or substituted as explicitly defined; wherein pyrazol-1-yl, and [1,2,3]triazol-2-yl groups are preferably unsubstituted. Particular examples are pyrazol-1-yl, [1,2,3]triazol-2-yl, 2-methyl-thiazol-4-yl, 3-methyl-[1,2,4]oxadiazol-5-yl, 6-methoxy-pyridin-3-yl, pyridin-2-yl, pyridin-3-yl, and pyrimidin-2-yl [notably the 5-membered heteroaryl group [1,2,3]triazol-2-yl; and the 6-membered heteroaryl group pyrimidin-2-yl].

Further embodiments of the invention are presented hereinafter:

4) Another embodiment relates to compounds according to any one of embodiments 1) to 3), wherein one of **R**² and **R**³ represents hydrogen, and the other represents hydrogen or methyl (wherein especially **R**³ represents hydrogen and **R**² represents hydrogen or methyl).

5) Another embodiment relates to compounds according to any one of embodiments 1) to 4), wherein R^{4a} and R^{4b} represent hydrogen; or R^{4a} represents methoxy or fluorine, and R^{4b} represents hydrogen; or R^{4a} and R^{4b} together represent a group $H_2C=$; wherein, in case R^3 is different from hydrogen, both R^{4a} and R^{4b} represent hydrogen.

5 6) Another embodiment relates to compounds according to any one of embodiments 1) to 3), wherein

- one of R^2 and R^3 represents hydrogen, and the other represents hydrogen or methyl (wherein especially R^3 represents hydrogen and R^2 represents hydrogen or methyl); and

10 • R^{4a} and R^{4b} represent hydrogen; or R^{4a} represents methoxy or fluorine, and R^{4b} represents hydrogen; or R^{4a} and R^{4b} together represent a group $H_2C=$; wherein, in case R^3 is methyl, both R^{4a} and R^{4b} represent hydrogen.

7) Another embodiment relates to compounds according to any one of embodiments 1) to 3), wherein

- 15
- R^3 represents hydrogen;
 - R^2 represents hydrogen or methyl; and
 - R^{4a} and R^{4b} represent hydrogen; or R^{4a} represents methoxy or fluorine, and R^{4b} represents hydrogen; or R^{4a} and R^{4b} together represent a group $H_2C=$.

8) Another embodiment relates to compounds according to any one of embodiments 1) to 3),
20 wherein R^2 , R^3 , R^{4a} and R^{4b} represent hydrogen.

9) Another embodiment relates to compounds according to any one of embodiments 1) to 3), wherein R^2 represents hydrogen or methyl; R^3 represents hydrogen; and R^{4a} and R^{4b} represent hydrogen, or R^{4a} and R^{4b} together represent a group $H_2C=$.

10) Another embodiment relates to compounds according to any one of embodiments 1) to
25 3), wherein R^2 represents hydrogen or methyl (especially methyl); and R^3 , R^{4a} and R^{4b} represent hydrogen.

11) Another embodiment relates to compounds according to any one of embodiments 1) to 3), wherein R^2 represents hydrogen or methyl; R^3 represents hydrogen; and R^{4a} and R^{4b} together represent a group $H_2C=$.

30 12) Another embodiment relates to compounds according to any one of embodiments 1) to 3), wherein R^2 represents hydrogen or methyl; R^3 represents hydrogen; R^{4a} represents methoxy or fluorine (especially R^{4a} represents methoxy); and R^{4b} represents hydrogen.

13) Another embodiment relates to compounds according to any one of embodiments 1) to 3), wherein R^3 represents methyl and R^2 , R^{4a} and R^{4b} represent hydrogen.

14) Another embodiment relates to compounds according to any one of embodiments 1) to 3), wherein both R^{4a} and R^{4b} represent fluorine and R^2 and R^3 represent hydrogen.

5 15) Another embodiment relates to compounds of formula (I) or (II) according to any one of embodiments 1), 2), or 4) to 14), wherein $(R^5)_n$ represents one or two optional substituents (i.e. n represents the integer 0, 1, or 2) (especially $(R^5)_n$ represents one or two substituents; i.e. n represents the integer 1 or 2) independently selected from (C₁₋₄)alkyl (especially methyl), (C₁₋₄)alkoxy (especially methoxy), halogen (especially fluorine, chlorine or bromine),
10 (C₁₋₄)alkyl-thio- (especially H₃C-S-), (C₁₋₃)fluoroalkyl (especially trifluoromethyl), (C₁₋₃)fluoroalkoxy (especially trifluoromethoxy), (C₁₋₃)fluoroalkyl-thio- (especially F₃C-S-), and cyano.

16) Another embodiment relates to compounds of formula (III) according to any one of embodiments 3) to 14); or to compounds of formula (I) or (II) according to any one of
15 embodiments 1) to 14), wherein the fragment



wherein

R^{14} , R^{15} , R^{16} and R^{17} together represent one or two optional substituents (i.e. at least two of R^{14} , R^{15} , R^{16} and R^{17} are hydrogen) [notably R^{14} , R^{15} , R^{16} and R^{17} together represent one or
20 two substituents (i.e. at least two of R^{14} , R^{15} , R^{16} and R^{17} are hydrogen and at least one of R^{14} , R^{15} , R^{16} and R^{17} is different from hydrogen)], wherein

- R^{14} and R^{17} independently represent hydrogen, (C₁₋₄)alkyl, (C₁₋₄)alkoxy (especially methoxy), (C₁₋₄)alkyl-thio- (especially H₃C-S-), halogen, (C₁₋₃)fluoroalkyl (especially trifluoromethyl), (C₁₋₄)alkoxy-carbonyl- (especially H₃CO-CO-), hydroxy-(C₁₋₄)alkyl-
25 (especially HO-CH₂-), hydroxy, or nitro; and
- R^{15} and R^{16} independently represent hydrogen, (C₁₋₄)alkyl, (C₁₋₄)alkoxy (especially methoxy), (C₁₋₄)alkyl-thio- (especially H₃C-S-), halogen, (C₁₋₃)fluoroalkyl (especially trifluoromethyl), (C₁₋₃)fluoroalkoxy (especially trifluoromethoxy), (C₁₋₃)fluoroalkyl-thio- (especially F₃C-S-), hydroxy-(C₁₋₄)alkyl- (especially HO-CH₂-), or cyano;
- 30 or R^{14} and R^{15} together, or R^{16} and R^{17} together, represent a group -O-CH₂-CH₂-O-;

or R^{15} and R^{16} together represent a fused phenyl group which, together with the benzimidazole moiety to which it is fused to, forms a 1H-naphtho[2,3-d]imidazol-2-yl group.

17) Another embodiment relates to compounds of formula (III) according to any one of embodiments 3) to 14); or to compounds of formula (I) or (II) according to any one of
 5 embodiments 1) to 14), wherein the fragment



wherein

R^{14} , R^{15} , R^{16} and R^{17} together represent one or two optional substituents [notably R^{14} , R^{15} , R^{16} and R^{17} together represent one or two substituents (i.e. at least two of R^{14} , R^{15} , R^{16} and
 10 R^{17} are hydrogen and at least one of R^{14} , R^{15} , R^{16} and R^{17} is different from hydrogen)],
 wherein

- R^{14} and R^{17} independently represent hydrogen, (C_{1-4}) alkyl, (C_{1-4}) alkoxy (especially methoxy), (C_{1-4}) alkyl-thio- (especially H_3C-S-), halogen, or (C_{1-3}) fluoroalkyl (especially trifluoromethyl); and
- 15 • R^{15} and R^{16} independently represent hydrogen, (C_{1-4}) alkyl, (C_{1-4}) alkoxy (especially methoxy), (C_{1-4}) alkyl-thio- (especially H_3C-S-), halogen, (C_{1-3}) fluoroalkyl (especially trifluoromethyl), (C_{1-3}) fluoroalkoxy (especially trifluoromethoxy), (C_{1-3}) fluoroalkyl-thio- (especially F_3C-S-), or cyano.

18) Another embodiment relates to compounds of formula (III) according to any one of
 20 embodiments 3) to 14); or to compounds of formula (I) or (II) according to any one of
 embodiments 1) to 14), wherein the fragment



wherein

R^{14} , R^{15} , R^{16} and R^{17} together represent one or two substituents (i.e. at least two of R^{14} , R^{15} , R^{16} and R^{17} are hydrogen and at least one of R^{14} , R^{15} , R^{16} and R^{17} is different from hydrogen), wherein

- one of R^{15} and R^{16} represents (C₁₋₄)alkyl, (C₁₋₄)alkoxy (especially methoxy), (C₁₋₄)alkyl-thio- (especially H₃C-S-), halogen, (C₁₋₃)fluoroalkyl (especially trifluoromethyl), (C₁₋₃)fluoroalkoxy (especially trifluoromethoxy), or cyano;
- or one of R^{14} and R^{17} represents (C₁₋₄)alkyl, (C₁₋₄)alkoxy (especially methoxy), (C₁₋₄)alkyl-thio- (especially H₃C-S-), halogen, or (C₁₋₃)fluoroalkyl (especially trifluoromethyl);

and the remaining substituent, if present, is one of R^{15} and R^{16} representing (C₁₋₄)alkyl, (C₁₋₄)alkoxy (especially methoxy), or halogen.

19) Another embodiment relates to compounds according to any one of embodiments 1) to 18), wherein R^1 represents hydrogen.

20) Another embodiment relates to compounds according to any one of embodiments 1) to 18), wherein R^1 represents (C₁₋₄)alkyl (especially methyl or ethyl, notably methyl).

21) Another embodiment relates to compounds according to any one of embodiments 1) to 20), wherein Ar^1 represents phenyl or 5- or 6-membered heteroaryl, wherein said phenyl or 5- or 6-membered heteroaryl independently is mono-, di-, or tri-substituted; wherein

- one of said substituents is attached in *ortho*-position to the point of attachment of Ar^1 to the rest of the molecule; wherein said substituent is phenyl or 5- or 6-membered heteroaryl; wherein said phenyl or 5- or 6-membered heteroaryl substituent is independently unsubstituted, mono-, di-, or tri-substituted (especially unsubstituted, mono-, or di-substituted), wherein the substituents are independently selected from (C₁₋₄)alkyl, (C₁₋₄)alkoxy, halogen, cyano, (C₁₋₃)fluoroalkyl, and (C₁₋₃)fluoroalkoxy; or said *ortho* substituent is benzo[1,3]dioxol-5-yl;
- and the other of said substituents, if present, is/are independently selected from (C₁₋₄)alkyl; (C₁₋₄)alkoxy; (C₃₋₆)cycloalkyl; halogen; cyano; (C₁₋₃)fluoroalkyl; (C₁₋₃)fluoroalkoxy; and -NR¹⁰R¹¹, wherein R¹⁰ and R¹¹ independently represent (C₁₋₄)alkyl, or R¹⁰ and R¹¹ together with the nitrogen to which they are attached to form a pyrrolidine ring.

22) Another embodiment relates to compounds of formula (I) or (II) according to any one of embodiments 1) to 20); or *mutatis mutandis* to compounds of formula (III) according to any one of embodiments 3) to 20); wherein Ar^1 represents phenyl or 5- or 6-membered

heteroaryl, wherein said phenyl or 5- or 6-membered heteroaryl independently is mono-, di-, or tri-substituted; wherein

- one of said substituents is attached in *ortho*-position to the point of attachment of Ar¹ to the rest of the molecule; wherein said substituent is phenyl or 5- or 6-membered heteroaryl; wherein said phenyl or 5- or 6-membered heteroaryl substituent is independently unsubstituted, or mono-, or di-substituted, wherein the substituents are independently selected from (C₁₋₂)alkyl, (C₁₋₂)alkoxy, halogen, cyano, trifluoromethyl, and trifluoromethoxy;
or said *ortho* substituent is benzo[1,3]dioxolyl;
or said *ortho* substituent is 2-(3-methoxy-phenyl)-ethynyl
- and the other of said substituents, if present, is/are independently selected from methyl; methoxy; cyclopropyl; halogen; cyano; trifluoromethyl; trifluoromethoxy; dimethylamino; pyrrolidin-1-yl; unsubstituted pyridinyl; and phenyl which is unsubstituted, or mono- or di-substituted, wherein the substituents are independently selected from methyl, methoxy, cyano, and halogen.

23) Another embodiment relates to compounds according to any one of embodiments 1) to 20), wherein

- **Ar¹** represents 5-membered heteroaryl, wherein the 5-membered heteroaryl is mono- or di-substituted; wherein
 - one of said substituents is attached in *ortho*-position to the point of attachment of Ar¹ to the rest of the molecule; wherein said *ortho*-substituent is phenyl, or 6-membered heteroaryl (especially pyridyl), which phenyl or 6-membered heteroaryl is independently unsubstituted, mono-, di-, or tri-substituted (especially unsubstituted, mono-, or di-substituted), wherein the substituents are independently selected from (C₁₋₄)alkyl, (C₁₋₄)alkoxy, halogen, cyano, (C₁₋₃)fluoroalkyl, and (C₁₋₃)fluoroalkoxy [wherein phenyl is especially unsubstituted, or mono-, or di-substituted with (C₁₋₄)alkyl, (C₁₋₄)alkoxy, halogen, (C₁₋₃)fluoroalkyl, or (C₁₋₃)fluoroalkoxy; and 6-membered heteroaryl is notably pyridyl which is mono-substituted with (C₁₋₄)alkoxy];
 - and the other of said substituents, if present, is/are independently selected from (C₁₋₄)alkyl (especially methyl); (C₃₋₆)cycloalkyl (especially cyclopropyl); -NR¹⁰R¹¹, wherein R¹⁰ and R¹¹ independently represent (C₁₋₄)alkyl (especially methyl), or R¹⁰ and R¹¹ together with the nitrogen to which they are attached to form a pyrrolidine ring; and phenyl which is unsubstituted, or mono- or di-substituted, wherein the substituents are independently selected from

(C₁₋₄)alkyl, (C₁₋₄)alkoxy, cyano, and halogen (especially the other of said substituents represents methyl or cyclopropyl);

- or **Ar**¹ represents 6-membered heteroaryl, wherein the 6-membered heteroaryl is mono-, di-, or tri-substituted (especially mono- or di-substituted); wherein

5 ➤ one of said substituents is attached in *ortho*-position to the point of attachment of **Ar**¹ to the rest of the molecule; wherein

 ○ said *ortho*-substituent is unsubstituted 5-membered heteroaryl (notably pyrazol-1-yl or [1,2,3]triazol-2-yl, especially in case **Ar**¹ represents pyridyl);

10 ○ or said *ortho*-substituent is phenyl which is unsubstituted, mono-, di-, or tri-substituted (especially unsubstituted, mono-, or di-substituted), wherein the substituents are independently selected from (C₁₋₄)alkyl, (C₁₋₄)alkoxy, halogen, cyano, (C₁₋₃)fluoroalkyl, and (C₁₋₃)fluoroalkoxy (especially (C₁₋₄)alkyl, (C₁₋₄)alkoxy, halogen and (C₁₋₃)fluoroalkyl);

15 ➤ and the other of said substituents, if present, is/are independently selected from (C₁₋₄)alkyl, (C₁₋₄)alkoxy, halogen, and (C₁₋₃)fluoroalkyl (especially (C₁₋₄)alkyl, and halogen; notably methyl);

- or **Ar**¹ represents phenyl which is mono-, di-, or tri-substituted; wherein

20 ➤ one of said substituents is attached in *ortho*-position to the point of attachment of **Ar**¹ to the rest of the molecule; wherein

 ○ said *ortho*-substituent is phenyl which is unsubstituted, mono-, di-, or tri-substituted (especially unsubstituted, mono-, or di-substituted), wherein the substituents are independently selected from (C₁₋₄)alkyl, (C₁₋₄)alkoxy, halogen, cyano, (C₁₋₃)fluoroalkyl, and (C₁₋₃)fluoroalkoxy;

25 ○ or said *ortho* substituent is benzo[1,3]dioxol-5-yl;

 ○ or said *ortho*-substituent is 6-membered heteroaryl which is unsubstituted, mono-, or di-substituted (especially unsubstituted or mono-substituted), wherein the substituents are independently selected from (C₁₋₄)alkyl, (C₁₋₄)alkoxy, halogen, and (C₁₋₃)fluoroalkyl (especially mono-substituted with (C₁₋₄)alkyl, or (C₁₋₄)alkoxy);

30 ○ or said *ortho*-substituent is 5-membered heteroaryl which is unsubstituted, mono-, or di-substituted (especially unsubstituted or mono-substituted), wherein the substituents are independently selected from (C₁₋₄)alkyl, (C₁₋₄)alkoxy, halogen, and (C₁₋₃)fluoroalkyl (especially (C₁₋₄)alkyl, notably methyl);

35

○ or said *ortho*-substituent is 2-(3-methoxy-phenyl)-ethynyl;

- and the other of said substituents, if present, is/are independently selected from (C₁₋₄)alkyl; (C₁₋₄)alkoxy; (C₃₋₆)cycloalkyl; halogen; cyano; (C₁₋₃)fluoroalkyl; (C₁₋₃)fluoroalkoxy; unsubstituted pyridinyl; and phenyl which is unsubstituted, or mono- or di-substituted, wherein the substituents are independently selected from (C₁₋₄)alkyl, (C₁₋₄)alkoxy, cyano, and halogen [notably selected from (C₁₋₄)alkyl, (C₁₋₄)alkoxy, halogen, cyano, (C₁₋₃)fluoroalkyl, and (C₁₋₃)fluoroalkoxy; especially from (C₁₋₄)alkyl, (C₁₋₄)alkoxy, and halogen].

24) Another embodiment relates to compounds according to any one of embodiments 1) to 23), wherein one or both of the following characteristics are present:

- in case **Ar**¹ represents a 5-membered heteroaryl group, such group is an oxazolyl, thienyl, or thiazolyl group (especially a thiazolyl group); and/or
- in case **Ar**¹ represents a 6-membered heteroaryl group, such group is a pyridinyl, pyrazinyl, or pyrimidinyl group (especially a pyridinyl group);

wherein said groups independently are substituted as defined in any one of the preceeding embodiments.


25) Another embodiment relates to compounds according to any one of embodiments 1) to 24), wherein one or both of the following characteristics are present:

- in case said *ortho* substituent of **Ar**¹ represents a 5-membered heteroaryl group, such group is a triazolyl (especially unsubstituted [1,2,3]triazol-2-yl), a pyrazolyl (especially unsubstituted pyrazol-1-yl), an oxadiazolyl (especially 3-methyl-[1,2,4]oxadiazol-5-yl), or a thiazolyl group (especially 2-methyl-thiazol-4-yl); [notably such group is [1,2,3]triazol-2-yl or pyrazol-1-yl]; and/or
- in case said *ortho* substituent of **Ar**¹ represents a 6-membered heteroaryl group, such group is a pyridinyl or a pyrimidinyl group (especially 6-methoxy-pyridin-3-yl, pyridin-2-yl, pyridin-3-yl, or pyrimidin-2-yl; preferably 6-methoxy-pyridin-3-yl, pyridin-2-yl or pyrimidin-2-yl);


wherein said groups independently are unsubstituted or substituted as defined in any one of the preceeding embodiments, or as explicitly defined herein.

26) Another embodiment relates to compounds according to any one of the embodiments 1) to 20) wherein **Ar¹** is a group independently selected from the following groups A, B, C, D, E, F, G, H, I, J, K, L, or M:


A.




B.




C.







10 D.




E.







J.




K.



5 L.



M.




10 wherein the groups listed in A and L are preferred groups, and wherein each of the groups A to M forms a particular sub-embodiment.

27) Another embodiment relates to compounds of formula (III) according to any one of embodiments 3) to 14), or to compounds of formula (I) or (II) according to embodiment 16), wherein

- 5
- R^2 , R^3 , R^{4a} , R^{4b} represent hydrogen; R^{15} represents chlorine and R^{14} represents methyl, or R^{16} represents chlorine and R^{17} represents methyl, and the remaining represent hydrogen;


and Ar^1 is as defined in any one of embodiments 21) to 26); wherein, in a sub-embodiment, a particular group for Ar^1 is:



- 10 28) Another embodiment relates to compounds of formula (III) according to any one of embodiments 3) to 14), or to compounds of formula (I) or (II) according to embodiment 16), wherein R^2 , R^3 , R^{4a} , R^{4b} , R^{14} , R^{15} , R^{16} and R^{17} are characterized by one or more of the following characteristics:

- 15
- R^2 , R^3 , R^{4a} , R^{4b} , R^{15} , and R^{16} represent hydrogen; one of R^{14} and R^{17} represents $-S-CH_3$, and the remaining represents hydrogen; or
 - R^2 , R^3 , R^{4a} , R^{4b} , R^{14} , and R^{17} represent hydrogen; one of R^{15} and R^{16} represents $-O-CH_3$ or $-S-CH_3$, and the remaining represents hydrogen;


and Ar^1 is as defined in any one of embodiments 21) to 26); wherein, in a sub-embodiment, a particular group for Ar^1 is:



29) Another embodiment relates to compounds of formula (III) according to any one of embodiments 3) to 14), or to compounds of formula (I) or (II) according to embodiment 16), wherein

- 25
- R^2 , R^3 , R^{4a} , R^{4b} , R^{15} , and R^{16} represent hydrogen; one of R^{14} and R^{17} represents $-S-CH_3$, and the remaining represents hydrogen;


and Ar¹ is as defined in any one of embodiments 21) to 26); wherein, in a sub-embodiment, particular groups for Ar¹ are:



30) Another embodiment relates to compounds of formula (III) according to any one of
 5 embodiments 3) to 14), or to compounds of formula (I) or (II) according to embodiment 16), wherein R², R³, R^{4a}, R^{4b}, R¹⁴, R¹⁵, R¹⁶ and R¹⁷ are characterized by one, or more (in any combination), or all of the following characteristics:

- R², R³, R^{4a}, R^{4b}, R¹⁷, and R¹⁴ represent hydrogen, one of R¹⁵ and R¹⁶ represents -O-CH₃ or -S-CH₃, and the remaining represents hydrogen; or
- 10 • R² represents methyl; R³, R^{4a}, and R^{4b} represent hydrogen; R¹⁵ represents chlorine and R¹⁴ represents methyl, or R¹⁶ represents chlorine and R¹⁷ represents methyl, and the remaining represent hydrogen; or
- R² represents methyl; R³, R^{4a}, R^{4b}, R¹⁴ and R¹⁷ represent hydrogen; R¹⁶ and R¹⁵ both represent methyl;


15 and Ar¹ is as defined in any one of embodiments 21) to 26); wherein, in a sub-embodiment, particular groups for Ar¹ are:



31) Another embodiment relates to compounds of formula (III) according to any one of
 20 embodiments 3) to 14), or to compounds of formula (I) or (II) according to embodiment 16), wherein

- R² represents methyl; R³, R^{4a}, R^{4b}, R¹⁴ and R¹⁷ represent hydrogen; R¹⁵ and R¹⁶ both represent methyl;


and Ar¹ is as defined in any one of embodiments 21) to 26); wherein, in a sub-embodiment, particular groups for Ar¹ are:



32) Another embodiment relates to compounds according of formual (III) according to any one of embodiments 3) to 14), or to compounds of formual (I) or (II) according to embodiment 16), wherein

- R² represents methyl; R³, R^{4a}, R^{4b}, R¹⁴ and R¹⁷ represent hydrogen; R¹⁵ and R¹⁶ both represent methoxy;


and Ar¹ is as defined in any one of embodiments 21) to 26); wherein, in a sub-embodiment, particular groups for Ar¹ are:



33) Another embodiment relates to compounds of formual (III) according to any one of embodiments 3) to 14), or to compounds of formual (I) or (II) according to embodiment 16), wherein R², R³, R^{4a}, R^{4b}, R¹⁴, R¹⁵, R¹⁶ and R¹⁷ are characterized by one or more of the following characteristics:

- R² represents methyl; R³, R^{4a}, R^{4b}, R¹⁴ and R¹⁷ represent hydrogen; one of R¹⁵ and R¹⁶ represents trifluoromethyl, and the remaining represents hydrogen, chlorine or fluorine;
- R² represents methyl; R³, R^{4a}, and R^{4b} represent hydrogen; R¹⁴ represents chlorine and R¹⁶ represents trifluoromethyl, or R¹⁷ represents chlorine and R¹⁵ represents trifluoromethyl, and the remaining represent hydrogen;


and Ar¹ is as defined in any one of embodiments 21) to 26); wherein, in a sub-embodiment, a particular group for Ar¹ is:



34) Another embodiment relates to compounds of formula (III) according to any one of embodiments 3) to 14), or to compounds of formula (I) or (II) according to embodiment 16), wherein R^2 , R^3 , R^{4a} , R^{4b} , R^{14} , R^{15} , R^{16} and R^{17} are characterized by one, or more (in any combination), or all of the following characteristics:

- 5
 - R^3 represents methyl; R^2 , R^{4a} , and R^{4b} , represent hydrogen; R^{14} represents methyl and R^{15} represents chlorine, or R^{17} represents methyl and R^{16} represents chlorine, and the remaining represent hydrogen; or
 - R^3 represents methyl; R^2 , R^{4a} , R^{4b} , R^{14} and R^{17} represent hydrogen; one of R^{15} and R^{16} represents methyl, and the remaining represents chlorine; or
- 10
 - R^3 represents methyl; R^2 , R^{4a} , R^{4b} , R^{14} and R^{17} represent hydrogen; R^{15} and R^{16} both represent methyl; or
 - R^3 represents methyl; R^2 , R^{4a} , and R^{4b} represent hydrogen; R^{14} represents methyl and R^{16} represents chlorine, or R^{17} represents methyl and R^{15} represents chlorine, and the remaining represent hydrogen; or
- 15
 - R^3 represents methyl; R^2 , R^{4a} , and R^{4b} represent hydrogen; R^{14} represents methyl and R^{16} represents bromine, or R^{17} represents methyl and R^{15} represents bromine, and the remaining represent hydrogen; or
 - R^3 represents methyl; R^2 , R^{4a} , R^{4b} , R^{14} and R^{17} represent hydrogen; R^{15} and R^{16} both represent methoxy;


20 and Ar^1 is as defined in any one of embodiments 21) to 26); wherein, in a sub-embodiment, particular groups for Ar^1 are:



35) Another embodiment relates to compounds of formula (III) according to any one of embodiments 3) to 14), or to compounds of formula (I) or (II) according to embodiment 16), wherein R^2 , R^3 , R^{4a} , R^{4b} , R^{14} , R^{15} , R^{16} and R^{17} are characterized by one, or more (in any combination), or all of the following characteristics:

- 5 • R^2 , R^3 , and R^{4b} represent hydrogen; R^{4a} represents methoxy; R^{14} represents methyl and R^{15} represents hydrogen or chlorine, or R^{17} represents methyl and R^{16} represents hydrogen or chlorine, and the remaining represent hydrogen; or
- R^2 , R^3 , and R^{4a} represent hydrogen; R^{4b} represents methoxy; R^{14} represents methyl and R^{15} represents hydrogen or chlorine, or R^{17} represents methyl and R^{16} represents hydrogen or chlorine, and the remaining represents hydrogen; or
- 10 • R^2 , R^3 , and R^{4b} represent hydrogen; R^{4a} represents fluorine; R^{14} represents methyl and R^{15} represents hydrogen or chlorine, or R^{17} represents methyl and R^{16} represents hydrogen or chlorine, and the remaining represents hydrogen; or
- R^2 , R^3 , and R^{4a} represent hydrogen; R^{4b} represents fluorine; R^{14} represents methyl and R^{15} represents hydrogen or chlorine, or R^{17} represents methyl and R^{16} represents hydrogen or chlorine, and the remaining represents hydrogen; or
- 15 • R^2 and R^3 represent hydrogen; R^{4a} and R^{4b} both represent fluorine; R^{14} represents methyl and R^{15} represents hydrogen or chlorine, or R^{17} represents methyl and R^{16} represents hydrogen or chlorine, and the remaining represents hydrogen;


20 and Ar^1 is as defined in any one of embodiments 21) to 26); wherein, in a sub-embodiment, particular groups for Ar^1 are:



36) Another embodiment relates to compounds of formula (III) according to any one of embodiments 3) to 14), or to compounds of formula (I) or (II) according to embodiment 16), wherein

- R^2 and R^3 represent hydrogen, R^{4a} and R^{4b} together represent $H_2C=$, and R^{14} , R^{15} , R^{16} and R^{17} together represent one or two substituents (i.e. two or three of R^{14} , R^{15} , R^{16} and R^{17} are hydrogen), wherein
 - R^{14} represents hydrogen or methoxy;
 - R^{15} represents hydrogen, chlorine, methyl or methoxy;
 - R^{16} represents hydrogen, methyl, chlorine, methoxy, trifluoromethoxy, tert.-butyl or $-S-CH_3$; and
 - R^{17} represents hydrogen, methyl, methoxy, bromine or $-S-CH_3$;


and Ar^1 is as defined in any one of embodiments 21) to 26); wherein particular groups for Ar^1 are:



37) Another embodiment relates to compounds according to any one of embodiments 27) to 36), wherein R^1 represents hydrogen.


38) Another embodiment relates to compounds according to any one of embodiments 27) to 36), wherein R^1 represents (C_{1-4}) alkyl (especially methyl or ethyl, notably methyl).

39) Another embodiment relates to compounds of formula (I) according to any one of embodiments 1) to 26); wherein the fragment




independently selected from the following groups A, B, or C:


A.



B.




5 C.



wherein it is understood that the benzimidazolyl moieties of groups A and B may be present in form of tautomers; and wherein the groups listed in group A are preferred groups.

40) Another embodiment relates to compounds of formula (I) according to any one of embodiments 1) to 14); wherein

- 5 • R^2 represents methyl; and R^3 , R^{4a} , and R^{4b} represent hydrogen (in which case the present embodiment is a particular embodiment for the compounds of formula (V) below); or
- R^3 represents methyl; and R^2 , R^{4a} , and R^{4b} represent hydrogen (in which case the present embodiment is a particular embodiment for the compounds of formula (VI) below);
- 10 • the fragment




R^1 represents hydrogen; and

R^{14} , R^{15} , R^{16} and R^{17} together represent one or two substituents [i.e. at least two of R^{14} , R^{15} , R^{16} and R^{17} are hydrogen and at least one of R^{14} , R^{15} , R^{16} and R^{17} is different from hydrogen]; wherein

- one of R^{15} and R^{16} represents methyl, methoxy, fluorine, chlorine, bromine, trifluoromethyl, or trifluoromethoxy;
- one of R^{14} and R^{17} , or the remaining of R^{15} and R^{16} , represents hydrogen, methyl, methoxy, fluorine, or chlorine;
- and the remaining of R^{14} , R^{15} , R^{16} and R^{17} represent hydrogen;

wherein, in a sub-embodiment, particular benzimidazole fragments are independently selected from the following groups:




wherein it is understood that said benzimidazolyl moieties may be present in form of tautomers;


and

- **Ar¹** is as defined in any one of embodiments 21) to 26);
wherein, in a sub-embodiment, particular groups **Ar¹** are independently selected from the following groups A, B, C, D, or E:


A.




B.




C.



D.



E.



wherein, in a further sub-embodiment, **Ar¹** is group is especially selected from the groups listed in group A.

- 5 41) Another embodiment relates to compounds of formula (I) according to any one of embodiments 1) to 14); wherein

- **R²** represents methyl; and **R³**, **R^{4a}**, and **R^{4b}** represent hydrogen (in which case the present embodiment is a particular embodiment for the compounds of formula (V) below); or

- 10 **R³** represents methyl; and **R²**, **R^{4a}**, and **R^{4b}** represent hydrogen (in which case the present embodiment is a particular embodiment for the compounds of formula (VI) below);

- the fragment




R^1 represents hydrogen; and

R^{14} , R^{15} , R^{16} and R^{17} are as follows:

- 5
 - R^{15} represents chlorine and R^{16} represents methyl, or R^{16} represents chlorine and R^{15} represents methyl, and R^{14} and R^{17} represent hydrogen;
 - or R^{15} represents chlorine and R^{14} represents methyl, or R^{16} represents chlorine and R^{17} represents methyl, and the remaining of R^{14} , R^{15} , R^{16} and R^{17} represent hydrogen;
- 10
 - or R^{16} represents chlorine and R^{14} represents methyl, or R^{15} represents chlorine and R^{17} represents methyl, and the remaining of R^{14} , R^{15} , R^{16} and R^{17} represent hydrogen;
 - R^{15} represents bromine and R^{16} represents fluorine, or R^{16} represents bromine and R^{15} represents fluorine, and R^{14} and R^{17} represent hydrogen;
- 15
 - or R^{16} represents bromine and R^{14} represents methyl, or R^{15} represents bromine and R^{17} represents methyl, and the remaining of R^{14} , R^{15} , R^{16} and R^{17} represent hydrogen;
 - or R^{15} and R^{16} both represent methoxy, and R^{14} and R^{17} represent hydrogen;
 - or one of R^{15} and R^{16} represents trifluoromethyl, the remaining of R^{15} and R^{16} represents hydrogen or chlorine; and R^{14} and R^{17} represent hydrogen;
- 20
 - or one of R^{15} and R^{16} represents trifluoromethoxy, and the remaining of R^{15} and R^{16} , and R^{14} and R^{17} represent hydrogen;

wherein, in a sub-embodiment, particular benzimidazole fragments are independently selected from the following groups:




wherein it is understood that said benzimidazolyl moieties may be present in form of tautomers;


and

- **Ar¹** is as defined in any one of embodiments 21) to 26);
wherein, in a sub-embodiment, particular groups for **Ar¹** are independently selected from the following groups A, B, C, D, or E:


A.




B.




C.



D.



E.



- 5 wherein, in a further sub-embodiment, **Ar¹** is group is especially selected from the groups listed in group A.

42) The invention, thus, relates to compounds of the formula (I) as defined in embodiment 1), compounds of the formula (II) as defined in embodiment 2), compounds of the formula (III) as defined in embodiment 3); or to such compounds further limited by the characteristics of any one of embodiments 6) to 41), under consideration of their respective dependencies; to pharmaceutically acceptable salts thereof; and to the use of such compounds as medicaments especially in the treatment of mental health disorders relating to orexinergic dysfunctions, which disorders are as defined below and which are especially selected from sleep disorders, anxiety disorders, addiction disorders, cognitive dysfunctions, mood disorders, or appetite disorders. For avoidance of any doubt, especially the following embodiments relating to the compounds of formula (I), (II), and (III) are thus possible and intended and herewith specifically disclosed in individualized form:


1, 6+1, 17+1, 17+6+1, 18+1, 18+6+1, 19+1, 19+6+1, 19+17+1, 19+17+6+1, 19+18+1, 19+18+6+1, 22+1, 22+6+1, 22+17+1, 22+17+6+1, 22+18+1, 22+18+6+1, 22+19+1, 22+19+6+1, 22+19+17+1, 22+19+17+6+1,

- 22+19+18+1, 22+19+18+6+1, 23+1, 23+6+1, 23+17+1, 23+17+6+1, 23+18+1, 23+18+6+1, 23+19+1, 23+19+6+1, 23+19+17+1, 23+19+17+6+1, 23+19+18+1, 23+19+18+6+1, 24+1, 24+6+1, 24+17+1, 24+17+6+1, 24+18+1, 24+18+6+1, 24+19+1, 24+19+6+1, 24+19+17+1, 24+19+17+6+1, 24+19+18+1, 24+19+18+6+1, 24+22+1, 24+22+6+1, 24+22+17+1, 24+22+17+6+1, 24+22+18+1, 24+22+18+6+1, 24+22+19+1, 24+22+19+6+1, 24+22+19+17+1, 24+22+19+17+6+1, 24+22+19+18+1, 24+22+19+18+6+1, 24+23+1, 24+23+6+1, 24+23+17+1, 24+23+17+6+1, 24+23+18+1, 24+23+18+6+1, 24+23+19+1, 24+23+19+6+1, 24+23+19+17+1, 24+23+19+17+6+1, 24+23+19+18+1, 24+23+19+18+6+1, 25+1, 25+6+1, 25+17+1, 25+17+6+1, 25+18+1, 25+18+6+1, 25+19+1, 25+19+6+1, 25+19+17+1, 25+19+17+6+1, 25+19+18+1, 25+19+18+6+1, 25+22+1, 25+22+6+1, 25+22+17+1, 25+22+17+6+1, 25+22+18+1, 25+22+18+6+1, 25+22+19+1, 25+22+19+6+1, 25+22+19+17+1, 25+22+19+17+6+1, 25+22+19+18+1, 25+22+19+18+6+1, 25+23+1, 25+23+6+1, 25+23+17+1, 25+23+17+6+1, 25+23+18+1, 25+23+18+6+1, 25+23+19+1, 25+23+19+6+1, 25+23+19+17+1, 25+23+19+17+6+1, 25+23+19+18+1, 25+23+19+18+6+1, 25+24+1, 25+24+6+1, 25+24+17+1, 25+24+17+6+1, 25+24+18+1, 25+24+18+6+1, 25+24+19+1, 25+24+19+6+1, 25+24+19+17+1, 25+24+19+17+6+1, 25+24+19+18+1, 25+24+19+18+6+1, 25+24+22+1, 25+24+22+6+1, 25+24+22+17+1, 25+24+22+17+6+1, 25+24+22+18+1, 25+24+22+18+6+1, 25+24+22+19+1, 25+24+22+19+6+1, 25+24+22+19+17+1, 25+24+22+19+17+6+1, 25+24+22+19+18+1, 25+24+22+19+18+6+1, 25+24+23+1, 25+24+23+6+1, 25+24+23+17+1, 25+24+23+17+6+1, 25+24+23+18+1, 25+24+23+18+6+1, 25+24+23+19+1, 25+24+23+19+6+1, 25+24+23+19+17+1, 25+24+23+19+17+6+1, 25+24+23+19+18+1, 25+24+23+19+18+6+1, 26+1, 26+6+1, 26+17+1, 26+17+6+1, 26+18+1, 26+18+6+1, 26+19+1, 26+19+6+1, 26+19+17+1, 26+19+17+6+1, 26+19+18+1, 26+19+18+6+1, 39+1, 39+6+1, 39+22+1, 39+22+6+1, 39+23+1, 39+23+6+1, 39+24+1, 39+24+6+1, 39+24+22+1, 39+24+22+6+1, 39+24+23+1, 39+24+23+6+1, 39+25+1, 39+25+6+1, 39+25+22+1, 39+25+22+6+1, 39+25+23+1, 39+25+23+6+1, 39+25+24+1, 39+25+24+6+1, 39+25+24+22+1, 39+25+24+22+6+1, 39+25+24+23+1, 39+25+24+23+6+1, 39+26+1, 39+26+6+1, 39+26+10+1, 39+26+13+1, 40+1, 41+1;
- 2, 6+2, 17+2, 17+6+2, 18+2, 18+6+2, 19+2, 19+6+2, 19+17+2, 19+17+6+2, 19+18+2, 19+18+6+2, 22+2, 22+6+2, 22+17+2, 22+17+6+2, 22+18+2, 22+18+6+2, 22+19+2, 22+19+6+2, 22+19+17+2, 22+19+17+6+2, 22+19+18+2, 22+19+18+6+2, 23+2, 23+6+2, 23+17+2, 23+17+6+2, 23+18+2, 23+18+6+2, 23+19+2, 23+19+6+2, 23+19+17+2, 23+19+17+6+2, 23+19+18+2, 23+19+18+6+2, 24+2, 24+6+2, 24+17+2, 24+17+6+2, 24+18+2, 24+18+6+2, 24+19+2, 24+19+6+2, 24+19+17+2, 24+19+17+6+2, 24+19+18+2, 24+19+18+6+2, 24+22+2, 24+22+6+2, 24+22+17+2, 24+22+17+6+2, 24+22+18+2, 24+22+18+6+2, 24+22+19+2, 24+22+19+6+2, 24+22+19+17+2, 24+22+19+17+6+2, 24+22+19+18+2, 24+22+19+18+6+2, 24+23+2, 24+23+6+2, 24+23+17+2, 24+23+17+6+2, 24+23+18+2, 24+23+18+6+2, 24+23+19+2, 24+23+19+6+2, 24+23+19+17+2, 24+23+19+17+6+2, 24+23+19+18+2, 24+23+19+18+6+2, 25+2, 25+6+2, 25+17+2, 25+17+6+2, 25+18+2, 25+18+6+2, 25+19+2, 25+19+6+2, 25+19+17+2, 25+19+17+6+2, 25+19+18+2, 25+19+18+6+2, 25+22+2, 25+22+6+2, 25+22+17+2, 25+22+17+6+2, 25+22+18+2, 25+22+18+6+2,

25+22+19+2, 25+22+19+6+2, 25+22+19+17+2, 25+22+19+17+6+2, 25+22+19+18+2, 25+22+19+18+6+2,
 25+23+2, 25+23+6+2, 25+23+17+2, 25+23+17+6+2, 25+23+18+2, 25+23+18+6+2, 25+23+19+2,
 25+23+19+6+2, 25+23+19+17+2, 25+23+19+17+6+2, 25+23+19+18+2, 25+23+19+18+6+2, 25+24+2,
 25+24+6+2, 25+24+17+2, 25+24+17+6+2, 25+24+18+2, 25+24+18+6+2, 25+24+19+2, 25+24+19+6+2,
 5 25+24+19+17+2, 25+24+19+17+6+2, 25+24+19+18+2, 25+24+19+18+6+2, 25+24+22+2, 25+24+22+6+2,
 25+24+22+17+2, 25+24+22+17+6+2, 25+24+22+18+2, 25+24+22+18+6+2, 25+24+22+19+2,
 25+24+22+19+6+2, 25+24+22+19+17+2, 25+24+22+19+17+6+2, 25+24+22+19+18+2, 25+24+22+19+18+6+2,
 25+24+23+2, 25+24+23+6+2, 25+24+23+17+2, 25+24+23+17+6+2, 25+24+23+18+2, 25+24+23+18+6+2,
 25+24+23+19+2, 25+24+23+19+6+2, 25+24+23+19+17+2, 25+24+23+19+17+6+2, 25+24+23+19+18+2,
 10 25+24+23+19+18+6+2, 26+2, 26+6+2, 26+17+2, 26+17+6+2, 26+18+2, 26+18+6+2, 26+19+2, 26+19+6+2,
 26+19+17+2, 26+19+17+6+2, 26+19+18+2, 26+19+18+6+2, 37+34+26+10+2, 37+34+26+13+2, 39+2, 39+6+2,
 39+22+2, 39+22+6+2, 39+23+2, 39+23+6+2, 39+24+2, 39+24+6+2, 39+24+22+2, 39+24+22+6+2, 39+24+23+2,
 39+24+23+6+2, 39+25+2, 39+25+6+2, 39+25+22+2, 39+25+22+6+2, 39+25+23+2, 39+25+23+6+2,
 39+25+24+2, 39+25+24+6+2, 39+25+24+22+2, 39+25+24+22+6+2, 39+25+24+23+2, 39+25+24+23+6+2,
 15 39+26+2, 39+26+6+2, 39+26+10+2, 39+26+13+2, 40+2, 41+2;
 3, 24+23+3, 25+23+3, 25+24+3, 25+24+23+3, 37+34+25+24+23+3.

In the list above the numbers refer to the embodiments according to their numbering provided hereinabove whereas “+” indicates the dependency from another embodiment. The different individualized embodiments are separated by commas. In other words, “39+26+6+2”
 20 for example refers to embodiment 39) depending on embodiment 26), depending on embodiment 6), depending on embodiment 2), i.e. embodiment “39+26+6+2” corresponds to the compounds of formula (II) according to embodiment 2) further limited by all the features of the embodiments 6), 26), and 39).

43) A further aspect of the invention relates to compounds of the formula (I) according to
 25 embodiment 1), which are also compounds of the formula (IV); wherein the absolute configuration is as depicted in formula (IV):



Formula (IV)

wherein

- **Ar¹** represents 5-membered heteroaryl selected from oxazolyl, thienyl, and thiazolyl (especially thiazolyl), wherein said 5-membered heteroaryl is mono- or di-substituted; wherein

- one of said substituents is attached in *ortho*-position to the point of attachment of **Ar¹** to the rest of the molecule; wherein said *ortho*-substituent is phenyl, or pyridyl, which phenyl or pyridyl is independently unsubstituted, mono-, or di-substituted, wherein the substituents are independently selected from (C₁₋₄)alkyl, (C₁₋₄)alkoxy, halogen, and (C₁₋₃)fluoroalkyl, [wherein phenyl is especially unsubstituted, mono-, or di-substituted with methyl, ethyl, methoxy, halogen or trifluoromethyl; and pyridyl is mono-substituted with methoxy];
- and the other of said substituents, if present, is independently selected from methyl; cyclopropyl; dimethylamino; pyrrolidin-1-yl; and phenyl which is unsubstituted, or mono- or di-substituted, wherein the substituents are independently selected from methyl, methoxy, cyano, and halogen (especially the other of said substituents represents methyl or cyclopropyl);

- or **Ar¹** represents 6-membered heteroaryl selected from pyridinyl, pyrazinyl, and pyrimidinyl (especially pyridinyl), wherein said 6-membered heteroaryl is mono- or di-substituted; wherein

- one of said substituents is attached in *ortho*-position to the point of attachment of **Ar¹** to the rest of the molecule; wherein
 - said *ortho*-substituent is pyrazol-1-yl or [1,2,3]triazol-2-yl;
 - or said *ortho*-substituent is phenyl which is unsubstituted, mono-, or di-substituted, wherein the substituents are independently selected from methyl, methoxy, halogen and trifluoromethyl;
- and the other of said substituents, if present, is methyl;

- or **Ar¹** represents phenyl which is mono-, di-, or tri-substituted; wherein

- one of said substituents is attached in *ortho*-position to the point of attachment of **Ar¹** to the rest of the molecule; wherein
 - said *ortho*-substituent is phenyl which is unsubstituted, mono-, or di-substituted, wherein the substituents are independently selected from methyl, methoxy, halogen, cyano, trifluoromethyl, and trifluoromethoxy;
 - or said *ortho* substituent is benzo[1,3]dioxol-5-yl;
 - or said *ortho* substituent is 2-(3-methoxy-phenyl)-ethynyl

- or said *ortho*-substituent is unsubstituted pyrimidinyl or pyridinyl;
- or said *ortho*-substituent is unsubstituted pyrazol-1-yl or [1,2,3]triazol-2-yl; or oxadiazolyl or thiazolyl, optionally mono-substituted with methyl (especially unsubstituted pyrazol-1-yl or [1,2,3]triazol-2-yl);

5 ➤ and the other of said substituents, if present, is/are independently selected from methyl, methoxy, halogen, cyano, trifluoromethyl, trifluoromethoxy, unsubstituted pyridinyl, and phenyl optionally mono-substituted with cyano [especially methyl, methoxy, and halogen];

10 one of R^2 and R^3 represents hydrogen, the other of R^2 and R^3 represents hydrogen or methyl; and

R^{4a} and R^{4b} independently represent hydrogen or fluorine; or R^{4a} represents methoxy and R^{4b} represents hydrogen; or R^{4a} and R^{4b} together represent a group $H_2C=$;

wherein, in case R^3 is different from hydrogen, both R^{4a} and R^{4b} represent hydrogen;

15 (notably one of R^2 and R^3 represents hydrogen, the other of R^2 and R^3 represents methyl, and R^{4a} and R^{4b} both represent hydrogen)

R^1 represents hydrogen, methyl, ethyl, cyclopropyl-methyl, 2-fluoro-ethyl or 2-methoxy-ethyl (especially hydrogen); and

20 R^{14} , R^{15} , R^{16} and R^{17} together represent one to three optional substituents (i.e. at least one of R^{14} , R^{15} , R^{16} and R^{17} is hydrogen) [notably R^{14} , R^{15} , R^{16} and R^{17} together represent one to three substituents (i.e. at least one of R^{14} , R^{15} , R^{16} and R^{17} is hydrogen and at least one of R^{14} , R^{15} , R^{16} and R^{17} is different from hydrogen)], wherein

- R^{14} and R^{17} independently represent hydrogen, (C_{1-4}) alkyl, (C_{1-4}) alkoxy (especially methoxy), (C_{1-4}) alkyl-thio- (especially H_3C-S-), halogen, (C_{1-3}) fluoroalkyl (especially trifluoromethyl), (C_{1-4}) alkoxy-carbonyl- (especially $H_3CO-CO-$), hydroxy- (C_{1-4}) alkyl- (especially $HO-CH_2-$), hydroxy, or nitro; and
- R^{15} and R^{16} independently represent hydrogen, (C_{1-4}) alkyl, (C_{1-4}) alkoxy (especially methoxy), (C_{1-4}) alkyl-thio- (especially H_3C-S-), halogen, (C_{1-3}) fluoroalkyl (especially trifluoromethyl), (C_{1-3}) fluoroalkoxy (especially trifluoromethoxy), (C_{1-3}) fluoroalkyl-thio- (especially F_3C-S-), hydroxy- (C_{1-4}) alkyl- (especially $HO-CH_2-$), or cyano;

30 or R^{14} and R^{15} together, or R^{16} and R^{17} together, represent a group $-O-CH_2-CH_2-O-$;

or R^{15} and R^{16} together represent a fused phenyl group which, together with the benzimidazole moiety to which it is fused to, forms a 1H-naphtho[2,3-d]imidazol-2-yl group;

with the exception of

[(S)-2-(1H-benzimidazol-2-yl)-1-pyrrolidinyl][2-(1H-pyrazol-1-yl)phenyl]-methanone; and


[(S)-2-(1H-benzimidazol-2-yl)-1-pyrrolidinyl][1,1'-biphenyl]-2-yl-methanone;

wherein all characteristics disclosed in embodiments 4) to 42) are intended to apply *mutatis mutandis* also to the compounds formula (IV) according to embodiment 43); wherein especially the following embodiments are thus possible and intended and herewith specifically disclosed in individualized form:

43, 10+43, 13+43, 18+10+43, 18+13+43, 18+43, 19+10+43, 19+13+43, 19+18+10+43, 19+18+13+43, 19+18+43, 19+43, 26+10+43, 26+13+43, 26+18+10+43, 26+18+13+43, 26+18+43, 26+19+10+43, 26+19+13+43, 26+19+18+10+43, 26+19+18+13+43, 26+19+18+43, 26+19+43, 26+43, 39+10+43, 39+13+43, 39+18+10+43, 39+18+13+43, 39+18+43, 39+19+10+43, 39+19+13+43, 39+19+18+10+43, 39+19+18+13+43, 39+19+18+43, 39+19+43, 39+26+10+43, 39+26+13+43, 39+26+18+10+43, 39+26+18+13+43, 39+26+18+43, 39+26+19+10+43, 39+26+19+13+43, 39+26+19+18+10+43, 39+26+19+18+13+43, 39+26+19+18+43, 39+26+19+43, 39+26+43, 39+43.

In the list above the numbers refer to the embodiments according to their numbering provided hereinabove whereas "+" indicates the dependency from another embodiment as outlined above.

44) A preferred aspect of the invention relates to compounds of the formula (I) according to embodiment 1), which are also compounds of the formula (V); wherein the absolute configuration is as depicted in formula (V):



Formula (V)

wherein

Ar¹ represents

- phenyl or 5- or 6-membered heteroaryl, wherein said phenyl or 5- or 6-membered heteroaryl independently is mono-, di-, or tri-substituted; wherein
 - one of said substituents is attached in *ortho*-position to the point of attachment of Ar¹ to the rest of the molecule; wherein said substituent is phenyl or 5- or 6-membered heteroaryl; wherein said phenyl or 5- or 6-membered heteroaryl substituent is independently unsubstituted, mono-, di-, or tri-substituted, wherein the substituents are independently selected from (C₁₋₄)alkyl,

(C₁₋₄)alkoxy, halogen, cyano, (C₁₋₃)fluoroalkyl, and (C₁₋₃)fluoroalkoxy; or said ortho substituent is benzo[1,3]dioxolyl;

- and the other of said substituents, if present, is/are independently selected from (C₁₋₄)alkyl; (C₁₋₄)alkoxy; (C₃₋₆)cycloalkyl; halogen; cyano; (C₁₋₃)fluoroalkyl; (C₁₋₃)fluoroalkoxy; -NR¹⁰R¹¹, wherein R¹⁰ and R¹¹ independently represent hydrogen or (C₁₋₄)alkyl, or R¹⁰ and R¹¹ together with the nitrogen to which they are attached to form a pyrrolidine ring; unsubstituted pyridinyl; and phenyl which is unsubstituted, or mono- or di-substituted, wherein the substituents are independently selected from (C₁₋₄)alkyl, (C₁₋₄)alkoxy, cyano, and halogen;

10 **R**¹ represents hydrogen, (C₁₋₄)alkyl (especially methyl or ethyl), (C₃₋₆)cycloalkyl-(CH₂)- (especially cyclopropyl-methyl), (C₂₋₃)fluoroalkyl (especially 2-fluoro-ethyl), or (C₁₋₄)alkoxy-(C₂₋₄)alkyl (especially 2-methoxy-ethyl); and

(**R**⁵)_n represents one to three optional substituents (i.e. n represents the integer 0, 1, 2, or 3) independently selected from (C₁₋₄)alkyl, (C₁₋₄)alkoxy (especially methoxy), halogen, (C₁₋₄)alkyl-thio- (especially H₃C-S-), (C₁₋₃)fluoroalkyl (especially trifluoromethyl), (C₁₋₃)fluoroalkoxy (especially trifluoromethoxy), (C₁₋₃)fluoroalkyl-thio- (especially F₃C-S-), hydroxy-(C₁₋₄)alkyl- (especially HO-CH₂-), (C₁₋₄)alkoxy-carbonyl- (especially H₃CO-CO-), nitro, hydroxy, and cyano; or (**R**⁵)_n represents a group -O-CH₂-CH₂-O-


wherein all characteristics disclosed in embodiments 15) to 42) are intended to apply *mutatis mutandis* also to the compounds formula (V) according to embodiment 44); wherein especially the following embodiments are thus possible and intended and herewith specifically disclosed in individualized form:

44, 17+44, 18+44, 19+17+44, 19+18+44, 19+44, 22+17+44, 22+18+44, 22+19+17+44, 22+19+18+44, 22+19+44, 22+44, 23+17+44, 23+18+44, 23+19+17+44, 23+19+18+44, 23+19+44, 23+44, 24+17+44, 24+18+44, 24+19+17+44, 24+19+18+44, 24+19+44, 24+22+17+44, 24+22+18+44, 24+22+19+17+44, 24+22+19+18+44, 24+22+19+44, 24+22+44, 24+23+17+44, 24+23+18+44, 24+23+19+17+44, 24+23+19+18+44, 24+23+19+44, 24+23+44, 24+44, 25+17+44, 25+18+44, 25+19+17+44, 25+19+18+44, 25+19+44, 25+22+17+44, 25+22+18+44, 25+22+19+17+44, 25+22+19+18+44, 25+22+19+44, 25+22+44, 25+23+17+44, 25+23+18+44, 25+23+19+17+44, 25+23+19+18+44, 25+23+19+44, 25+23+44, 25+24+17+44, 25+24+18+44, 25+24+19+17+44, 25+24+19+18+44, 25+24+19+44, 25+24+22+17+44, 25+24+22+18+44, 25+24+22+19+17+44, 25+24+22+19+18+44, 25+24+22+19+44, 25+24+22+44, 25+24+23+17+44, 25+24+23+18+44, 25+24+23+19+17+44, 25+24+23+19+18+44, 25+24+23+19+44, 25+24+23+44, 25+24+44, 25+44, 26+17+44, 26+18+44, 26+19+17+44, 26+19+18+44, 26+19+44, 26+44, 39+22+44, 39+23+44,

39+24+22+44, 39+24+23+44, 39+24+44, 39+25+22+44, 39+25+23+44, 39+25+24+22+44, 39+25+24+23+44, 39+25+24+44, 39+25+44, 39+26+44, 39+44, 40+44, 41+44.

In the list above the numbers refer to the embodiments according to their numbering provided hereinabove whereas "+" indicates the dependency from another embodiment as outlined above.

45) Another preferred aspect of the invention relates to compounds of the formula (I) according to embodiment 1), which are also compounds of the formula (VI); wherein the absolute configuration is as depicted in formula (VI):



Formula (VI)

wherein

Ar¹ represents

- phenyl or 5- or 6-membered heteroaryl, wherein said phenyl or 5- or 6-membered heteroaryl independently is mono-, di-, or tri-substituted; wherein
 - one of said substituents is attached in *ortho*-position to the point of attachment of Ar¹ to the rest of the molecule; wherein said substituent is phenyl or 5- or 6-membered heteroaryl; wherein said phenyl or 5- or 6-membered heteroaryl substituent is independently unsubstituted, mono-, di-, or tri-substituted, wherein the substituents are independently selected from (C₁₋₄)alkyl, (C₁₋₄)alkoxy, halogen, cyano, (C₁₋₃)fluoroalkyl, and (C₁₋₃)fluoroalkoxy; or said ortho substituent is benzo[1,3]dioxolyl;
 - and the other of said substituents, if present, is/are independently selected from (C₁₋₄)alkyl; (C₁₋₄)alkoxy; (C₃₋₆)cycloalkyl; halogen; cyano; (C₁₋₃)fluoroalkyl; (C₁₋₃)fluoroalkoxy; -NR¹⁰R¹¹, wherein R¹⁰ and R¹¹ independently represent hydrogen or (C₁₋₄)alkyl, or R¹⁰ and R¹¹ together with the nitrogen to which they are attached to form a pyrrolidine ring; unsubstituted pyridinyl; and phenyl which is unsubstituted, or mono- or di-substituted, wherein the substituents are independently selected from (C₁₋₄)alkyl, (C₁₋₄)alkoxy, cyano, and halogen;

R^1 represents hydrogen, (C_{1-4}) alkyl (especially methyl or ethyl), (C_{3-6}) cycloalkyl- (CH_2) - (especially cyclopropyl-methyl), (C_{2-3}) fluoroalkyl (especially 2-fluoro-ethyl), or (C_{1-4}) alkoxy- (C_{2-4}) alkyl (especially 2-methoxy-ethyl); and

$(R^5)_n$ represents one to three optional substituents (i.e. n represents the integer 0, 1, 2, or 3)

5 independently selected from (C_{1-4}) alkyl, (C_{1-4}) alkoxy (especially methoxy), halogen, (C_{1-4}) alkyl-thio- (especially H_3C-S-), (C_{1-3}) fluoroalkyl (especially trifluoromethyl), (C_{1-3}) fluoroalkoxy (especially trifluoromethoxy), (C_{1-3}) fluoroalkyl-thio- (especially F_3C-S-), hydroxy- (C_{1-4}) alkyl- (especially $HO-CH_2-$), (C_{1-4}) alkoxy-carbonyl- (especially $H_3CO-CO-$), nitro, hydroxy, and cyano; or $(R^5)_n$ represents a group $-O-CH_2-CH_2-O-$.

10 wherein all characteristics disclosed in embodiments 15) to 42) are intended to apply *mutatis mutandis* also to the compounds formula (VI) according to embodiment 45); wherein especially the following embodiments are thus possible and intended and herewith specifically disclosed in individualized form:

45, 17+45, 18+45, 19+17+45, 19+18+45, 19+45, 22+17+45, 22+18+45, 22+19+17+45, 22+19+18+45,
 15 22+19+45, 22+45, 23+17+45, 23+18+45, 23+19+17+45, 23+19+18+45, 23+19+45, 23+45, 24+17+45,
 24+18+45, 24+19+17+45, 24+19+18+45, 24+19+45, 24+22+17+45, 24+22+18+45, 24+22+19+17+45,
 24+22+19+18+45, 24+22+19+45, 24+22+45, 24+23+17+45, 24+23+18+45, 24+23+19+17+45,
 24+23+19+18+45, 24+23+19+45, 24+23+45, 24+45, 25+17+45, 25+18+45, 25+19+17+45, 25+19+18+45,
 25+19+45, 25+22+17+45, 25+22+18+45, 25+22+19+17+45, 25+22+19+18+45, 25+22+19+45, 25+22+45,
 20 25+23+17+45, 25+23+18+45, 25+23+19+17+45, 25+23+19+18+45, 25+23+19+45, 25+23+45, 25+24+17+45,
 25+24+18+45, 25+24+19+17+45, 25+24+19+18+45, 25+24+19+45, 25+24+22+17+45, 25+24+22+18+45,
 25+24+22+19+17+45, 25+24+22+19+18+45, 25+24+22+19+45, 25+24+22+45, 25+24+23+17+45,
 25+24+23+18+45, 25+24+23+19+17+45, 25+24+23+19+18+45, 25+24+23+19+45, 25+24+23+45, 25+24+45,
 25+45, 26+17+45, 26+18+45, 26+19+17+45, 26+19+18+45, 26+19+45, 26+45, 39+22+45, 39+23+45,
 25 39+24+22+45, 39+24+23+45, 39+24+45, 39+25+22+45, 39+25+23+45, 39+25+24+22+45, 39+25+24+23+45,
 39+25+24+45, 39+25+45, 39+26+45, 39+45, 40+45, 41+45.

In the list above the numbers refer to the embodiments according to their numbering provided hereinabove whereas "+" indicates the dependency from another embodiment as outlined above.

30 46) Another embodiment relates to compounds according to embodiment 1) selected from:

[2-(5-Chloro-6-methyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-[5-(3-chloro-phenyl)-2-methyl-thiazol-4-yl]-methanone;

[5-(3-Chloro-phenyl)-2-methyl-thiazol-4-yl]-[2-(4,6-dimethyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;

- [2-(4-Bromo-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-[5-(3-chloro-phenyl)-2-methyl-thiazol-4-yl]-methanone;
[2-(5-Bromo-7-methyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-[5-(3-chloro-phenyl)-2-methyl-thiazol-4-yl]-
methanone;
[2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-[5-(3-chloro-phenyl)-2-methyl-thiazol-4-yl]-
5 methanone;
[5-(3-Chloro-phenyl)-2-methyl-thiazol-4-yl]-[2-(6-chloro-5-trifluoromethyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-
methanone;
[2-(4-Bromo-6-fluoro-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-[5-(3-chloro-phenyl)-2-methyl-thiazol-4-yl]-
methanone;
10 [5-(3-Chloro-phenyl)-2-methyl-thiazol-4-yl]-[2-(4,5-dimethyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
[2-(5,6-Dichloro-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-(2-methyl-5-phenyl-thiazol-4-yl)-methanone;
[2-(5-Chloro-6-methyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-(2-methyl-5-phenyl-thiazol-4-yl)-methanone;
[2-(4,6-Dimethyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-(2-methyl-5-phenyl-thiazol-4-yl)-methanone;
[5-(3-Chloro-phenyl)-2-methyl-thiazol-4-yl]-[(S)-2-(7-methylsulfanyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-
15 methanone;
[2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-(2-methyl-5-phenyl-thiazol-4-yl)-methanone;
[2-(6-Chloro-5-trifluoromethyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-(2-methyl-5-phenyl-thiazol-4-yl)-methanone;
[(S)-2-(4,6-Dimethyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-(2-methyl-5-phenyl-thiazol-4-yl)-methanone;
[(S)-2-(5-Chloro-6-methyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-(2-methyl-5-phenyl-thiazol-4-yl)-methanone;
20 [(S)-2-(6-Bromo-4-methyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-(2-methyl-5-phenyl-thiazol-4-yl)-methanone;
[(S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-(2-methyl-5-phenyl-thiazol-4-yl)-methanone;
[(S)-2-(5-Chloro-6-trifluoromethyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-(2-methyl-5-phenyl-thiazol-4-yl)-
methanone;
[(S)-2-(6-Methylsulfanyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-(4-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
25 (4,5-Dimethoxy-2-[1,2,3]triazol-2-yl-phenyl)-[(S)-2-(6-methylsulfanyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-
methanone;
(5-Methoxy-2-[1,2,3]triazol-2-yl-phenyl)-[(S)-2-(7-methylsulfanyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-
methanone;
(2-Fluoro-3-methoxy-6-[1,2,3]triazol-2-yl-phenyl)-[(S)-2-(7-methylsulfanyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-
30 methanone;
[(S)-2-(7-Methylsulfanyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-(4-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
[(S)-2-(6-Methylsulfanyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-(2-methyl-5-p-tolyl-thiazol-4-yl)-methanone;
[(S)-2-(7-Methylsulfanyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
[4-(3-Chloro-phenyl)-2-methyl-pyrimidin-5-yl]-[(S)-2-(6-methylsulfanyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-
35 methanone;

- [4-(3,4-Dimethyl-phenyl)-2-methyl-pyrimidin-5-yl]-[(S)-2-(6-methylsulfanyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- (2-Methyl-4-phenyl-pyrimidin-5-yl)-[(S)-2-(6-methylsulfanyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- [4-(4-Bromo-3-chloro-phenyl)-2-methyl-pyrimidin-5-yl]-[(S)-2-(6-methylsulfanyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- 5 [2-Cyclopropyl-5-phenyl-thiazol-4-yl]-[(S)-2-(7-methylsulfanyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- [5-(3-Methoxy-phenyl)-2-methyl-thiazol-4-yl]-[(S)-2-(7-methylsulfanyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- [(S)-2-(7-Methylsulfanyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-[2-methyl-5-(3-trifluoromethyl-phenyl)-thiazol-4-yl]-methanone;
- 10 [(S)-2-(7-Methylsulfanyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-(2-methyl-5-p-tolyl-thiazol-4-yl)-methanone;
- (2-Cyclopropyl-5-p-tolyl-thiazol-4-yl)-[(S)-2-(7-methylsulfanyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- [2-Cyclopropyl-5-(4-fluoro-phenyl)-thiazol-4-yl]-[(S)-2-(7-methylsulfanyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- 15 (3',4'-Dimethyl-biphenyl-2-yl)-[(S)-2-(7-methylsulfanyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- [4-(3,4-Dichloro-phenyl)-2-methyl-pyrimidin-5-yl]-[(S)-2-(7-methylsulfanyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- [4-(3,4-Dichloro-phenyl)-pyrimidin-5-yl]-[(S)-2-(7-methylsulfanyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- 20 [4-(3,4-Dimethyl-phenyl)-pyrimidin-5-yl]-[(S)-2-(7-methylsulfanyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- [4-(4-Bromo-3-chloro-phenyl)-pyrimidin-5-yl]-[(S)-2-(7-methylsulfanyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- [4-(3,4-Dimethyl-phenyl)-2-methyl-pyrimidin-5-yl]-[(S)-2-(7-methylsulfanyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- 25 (2-Methyl-4-phenyl-pyrimidin-5-yl)-[(S)-2-(7-methylsulfanyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- [4-(4-Bromo-3-chloro-phenyl)-2-methyl-pyrimidin-5-yl]-[(S)-2-(7-methylsulfanyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- [4-(3,5-Dichloro-phenyl)-2-methyl-pyrimidin-5-yl]-[(S)-2-(7-methylsulfanyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- 30 [(2S,4R)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-4-fluoro-pyrrolidin-1-yl]-[5-(3-chloro-phenyl)-2-methyl-thiazol-4-yl]-methanone;

- [(2S,4R)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-4-fluoro-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
[5-(3-Chloro-phenyl)-2-methyl-thiazol-4-yl]-[(2S,4R)-4-fluoro-2-(4-methyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- 5 [(2S,4R)-2-(6-Chloro-7-methyl-1H-benzimidazol-2-yl)-4-methoxy-pyrrolidin-1-yl]-(5-chloro-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
[(2S,4R)-2-(6-Chloro-7-methyl-1H-benzimidazol-2-yl)-4-methoxy-pyrrolidin-1-yl]-(2-fluoro-3-methoxy-6-[1,2,3]triazol-2-yl-phenyl)-methanone;
[(2S,4R)-2-(6-Chloro-7-methyl-1H-benzimidazol-2-yl)-4-methoxy-pyrrolidin-1-yl]-(5-methoxy-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- 10 [(2S,4R)-2-(6-Chloro-7-methyl-1H-benzimidazol-2-yl)-4-methoxy-pyrrolidin-1-yl]-(2-fluoro-3-methyl-6-[1,2,3]triazol-2-yl-phenyl)-methanone;
[(2S,4R)-2-(6-Chloro-7-methyl-1H-benzimidazol-2-yl)-4-methoxy-pyrrolidin-1-yl]-(4,5-dimethyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- 15 [(2S,4R)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-4-methoxy-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
[(4R)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-4-methoxy-2-methyl-pyrrolidin-1-yl]-(2-methyl-5-phenyl-thiazol-4-yl)-methanone;
[(4R)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-4-methoxy-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- 20 [(4R)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-4-methoxy-2-methyl-pyrrolidin-1-yl]-[5-(3-chloro-phenyl)-2-methyl-thiazol-4-yl]-methanone;
[5-(3-Chloro-phenyl)-2-methyl-thiazol-4-yl]-[(4R)-4-methoxy-2-methyl-2-(4-methyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- 25 [(4R)-4-Methoxy-2-methyl-2-(4-methyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-(2-methyl-5-phenyl-thiazol-4-yl)-methanone;
[(2S,4R)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-4-methoxy-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- 30 [(2S,4R)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-4-methoxy-2-methyl-pyrrolidin-1-yl]-[5-(3-chloro-phenyl)-2-methyl-thiazol-4-yl]-methanone;
[(2S,4R)-2-(6-Chloro-7-methyl-1H-benzimidazol-2-yl)-4-methoxy-pyrrolidin-1-yl]-(2-methyl-5-m-tolyl-thiazol-4-yl)-methanone;
[(2S,4R)-2-(6-Chloro-7-methyl-1H-benzimidazol-2-yl)-4-methoxy-pyrrolidin-1-yl]-[2-dimethylamino-5-(3,4-dimethyl-phenyl)-thiazol-4-yl]-methanone;

- [(2S,4R)-2-(6-Chloro-7-methyl-1H-benzoimidazol-2-yl)-4-methoxy-pyrrolidin-1-yl]-[5-(3,4-dimethyl-phenyl)-2-methyl-thiazol-4-yl]-methanone;
- [(2S,4R)-2-(6-Chloro-7-methyl-1H-benzoimidazol-2-yl)-4-methoxy-pyrrolidin-1-yl]-[5-(4-chloro-phenyl)-2-methyl-thiazol-4-yl]-methanone;
- 5 [(2S,4R)-2-(6-Chloro-7-methyl-1H-benzoimidazol-2-yl)-4-methoxy-pyrrolidin-1-yl]-(2-methyl-5-p-tolyl-thiazol-4-yl)-methanone;
- [(2S,4R)-2-(6-Chloro-7-methyl-1H-benzoimidazol-2-yl)-4-methoxy-pyrrolidin-1-yl]-[5-(4-methoxy-phenyl)-2-methyl-thiazol-4-yl]-methanone;
- [(2S,4R)-2-(6-Chloro-7-methyl-1H-benzoimidazol-2-yl)-4-methoxy-pyrrolidin-1-yl]-(3',4'-dimethyl-biphenyl-2-yl)-
- 10 methanone;
- [(2S,4R)-4-Methoxy-2-(4-methyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-(2-methyl-5-phenyl-thiazol-4-yl)-methanone;
- [(2S,4R)-2-(6-Chloro-7-methyl-1H-benzoimidazol-2-yl)-4-methoxy-pyrrolidin-1-yl]-(4'-fluoro-3'-methyl-biphenyl-2-yl)-methanone;
- 15 [(2S,4R)-2-(6-Chloro-7-methyl-1H-benzoimidazol-2-yl)-4-methoxy-pyrrolidin-1-yl]-[5-(3-chloro-phenyl)-2-dimethylamino-thiazol-4-yl]-methanone;
- [(4R)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-4-methoxy-2-methyl-pyrrolidin-1-yl]-(5-fluoro-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- [(4R)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-4-methoxy-2-methyl-pyrrolidin-1-yl]-(5-chloro-2-[1,2,3]triazol-
- 20 2-yl-phenyl)-methanone;
- [(4R)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-4-methoxy-2-methyl-pyrrolidin-1-yl]-(2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- [(4R)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-4-methoxy-2-methyl-pyrrolidin-1-yl]-(4,5-dimethyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- 25 [(4R)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-4-methoxy-2-methyl-pyrrolidin-1-yl]-(4-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- [(4R)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-4-methoxy-2-methyl-pyrrolidin-1-yl]-(4,5-dimethoxy-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- [(4R)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-4-methoxy-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-pyrazol-1-yl-
- 30 phenyl)-methanone;
- [(4R)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-4-methoxy-2-methyl-pyrrolidin-1-yl]-(6-methyl-3-pyrazol-1-yl-pyridin-2-yl)-methanone;
- [(4R)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-4-methoxy-2-methyl-pyrrolidin-1-yl]-(6-methyl-3-[1,2,3]triazol-2-yl-pyridin-2-yl)-methanone;

- [(2S,4R)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-4-methoxy-pyrrolidin-1-yl]-(2-methyl-5-phenyl-thiazol-4-yl)-methanone;
- [5-(3-Chloro-phenyl)-2-methyl-thiazol-4-yl]-[(2S,4R)-4-methoxy-2-(4-methyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- 5 [(2S,4S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-4-methoxy-pyrrolidin-1-yl]-[5-(3-chloro-phenyl)-2-methyl-thiazol-4-yl]-methanone;
- [(2S,4R)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-4-methoxy-pyrrolidin-1-yl]-[5-(3-chloro-phenyl)-2-methyl-thiazol-4-yl]-methanone;
- (5-Chloro-2-[1,2,3]triazol-2-yl-phenyl)-[(2S,3S)-2-(5,6-dimethoxy-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-
- 10 methanone;
- [(2S,3S)-2-(5,6-Dimethoxy-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(4-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- [(2S,3S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(4-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- 15 [(2S,3S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(5-methoxy-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- [(2S,3S)-2-(5,6-Dimethyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(4-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- [(2S,3S)-2-(5,6-Dimethyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(5-methoxy-2-[1,2,3]triazol-2-yl-phenyl)-
- 20 methanone;
- [(2S,3S)-2-(5,6-Dimethyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- [(S)-2-Methyl-2-(4-methyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-(2-methyl-5-phenyl-thiazol-4-yl)-methanone;
- [(S)-2-(5,6-Dimethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-
- 25 methanone;
- [(S)-2-(5,6-Dimethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(4,4'-dimethyl-biphenyl-2-yl)-methanone;
- [(S)-2-(5,6-Dimethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(3'-fluoro-biphenyl-2-yl)-methanone;
- [(S)-2-(5,6-Dimethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(3'-methyl-biphenyl-2-yl)-methanone;
- [(S)-2-(4,5-Dimethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(3'-fluoro-4-methyl-biphenyl-2-yl)-
- 30 methanone;
- [(S)-2-(4,5-Dimethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(3'-methoxy-4-methyl-biphenyl-2-yl)-methanone;
- [(S)-2-(4,5-Dimethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(4,3'-dimethyl-biphenyl-2-yl)-methanone;
- [(S)-2-(4,5-Dimethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(4'-methoxy-4-methyl-biphenyl-2-yl)-
- 35 methanone;

- [(S)-2-(4,5-Dimethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(4,4'-dimethyl-biphenyl-2-yl)-methanone;
[(S)-2-(4,5-Dimethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(2'-fluoro-4-methyl-biphenyl-2-yl)-methanone;
[(S)-2-(4,5-Dimethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(4'-fluoro-4-methyl-biphenyl-2-yl)-methanone;
5 [(S)-2-(4,5-Dimethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(3'-fluoro-biphenyl-2-yl)-methanone;
[(S)-2-(4,5-Dimethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(3'-methyl-biphenyl-2-yl)-methanone;
[(S)-2-(4,5-Dimethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(4'-methoxy-biphenyl-2-yl)-methanone;
[(S)-2-(4,5-Dimethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(4'-methyl-biphenyl-2-yl)-methanone;
10 [(S)-2-(6-Bromo-5-fluoro-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(4,5-dimethyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
[(S)-2-(6-Bromo-5-fluoro-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-chloro-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
[(S)-2-(5-tert-Butyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
15 [(S)-2-(6-Bromo-5-fluoro-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(4-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
[(S)-2-(6-Chloro-5-trifluoromethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(2-fluoro-6-[1,2,3]triazol-2-yl-phenyl)-methanone;
20 [(S)-2-(6-Chloro-5-trifluoromethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(2-fluoro-3-methyl-6-[1,2,3]triazol-2-yl-phenyl)-methanone;
[(S)-2-(6-Chloro-5-trifluoromethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(4,5-dimethoxy-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
[(S)-2-(6-Chloro-5-trifluoromethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-fluoro-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
25 [(S)-2-(6-Chloro-5-trifluoromethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(4,5-dimethyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
[(S)-2-(6-Chloro-5-trifluoromethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-pyrazol-1-yl-phenyl)-methanone;
30 (5-Chloro-2-[1,2,3]triazol-2-yl-phenyl)-[(S)-2-(6-chloro-5-trifluoromethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-methanone;
[(S)-2-(4-Chloro-6-trifluoromethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
[(S)-2-(6-Chloro-5-trifluoromethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(4-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
35

- [(S)-2-(6-Chloro-5-trifluoromethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- [(S)-2-(6-Chloro-5-trifluoromethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(6-methyl-3-[1,2,3]triazol-2-yl-pyridin-2-yl)-methanone;
- 5 [(S)-2-(6-Chloro-5-trifluoromethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(2-fluoro-3-methoxy-6-[1,2,3]triazol-2-yl-phenyl)-methanone;
- [(S)-2-(6-Chloro-5-trifluoromethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(3'-methoxy-4-methyl-biphenyl-2-yl)-methanone;
- [(S)-2-(5-Chloro-6-methyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- 10 [(S)-2-(6-Chloro-5-trifluoromethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(2'-fluoro-4-methyl-biphenyl-2-yl)-methanone;
- [(S)-2-(6-Chloro-5-trifluoromethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(4'-fluoro-4-methyl-biphenyl-2-yl)-methanone;
- 15 [(S)-2-(6-Chloro-5-trifluoromethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(3'-fluoro-4-methyl-biphenyl-2-yl)-methanone;
- 2'-[(S)-2-(6-Chloro-5-trifluoromethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidine-1-carbonyl]-4'-methyl-biphenyl-3-carbonitrile;
- [(S)-2-(4,6-Dimethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- 20 [(5-Chloro-2-[1,2,3]triazol-2-yl-phenyl)-[(S)-2-methyl-2-(5-trifluoromethyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- (4,5-Dimethyl-2-[1,2,3]triazol-2-yl-phenyl)-[(S)-2-methyl-2-(5-trifluoromethyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- 25 (4-Methyl-2-[1,2,3]triazol-2-yl-phenyl)-[(S)-2-methyl-2-(5-trifluoromethyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- [(S)-2-(4-Bromo-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- (3'-Fluoro-4-methyl-biphenyl-2-yl)-[(S)-2-methyl-2-(5-trifluoromethyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- 30 (4,3'-Dimethyl-biphenyl-2-yl)-[(S)-2-methyl-2-(5-trifluoromethyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- [(S)-2-(5-Bromo-7-methyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- (4'-Methoxy-4-methyl-biphenyl-2-yl)-[(S)-2-methyl-2-(5-trifluoromethyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- 35

- (4,4'-Dimethyl-biphenyl-2-yl)-[(S)-2-methyl-2-(5-trifluoromethyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
(2-Benzo[1,3]dioxol-5-yl-5-methyl-phenyl)-[(S)-2-methyl-2-(5-trifluoromethyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
[(S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
5 [5-(3-Chloro-phenyl)-2-methyl-thiazol-4-yl]-[(S)-2-(4,5-difluoro-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-methanone;
[(S)-2-(5,6-Dimethoxy-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
10 [(S)-2-(4,5-Difluoro-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(2-methyl-5-m-tolyl-thiazol-4-yl)-methanone;
[(S)-2-(4,5-Difluoro-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-[5-(3-methoxy-phenyl)-2-methyl-thiazol-4-yl]-methanone;
[(S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-pyrimidin-2-yl-phenyl)-methanone;
15 [(S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-pyridin-2-yl-phenyl)-methanone;
[(S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-pyridin-3-yl-phenyl)-methanone;
[(S)-2-(6-Bromo-5-fluoro-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
20 [(S)-2-(6-Chloro-5-trifluoromethyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
[(S)-2-(4-Bromo-6-fluoro-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
25 [(S)-2-(4,5-Dimethyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
[(S)-2-(4-Isopropyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
[(S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(2-methyl-5-phenyl-thiazol-4-yl)-methanone;
30 [(S)-2-(7,8-Dihydro-3H-6,9-dioxo-1,3-diaza-cyclopenta[a]naphthalen-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
[(S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-chloro-2-[1,2,3]triazol-2-yl-phenyl)-methanone;

- [(S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methoxy-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- [(S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(2-fluoro-3-methoxy-6-[1,2,3]triazol-2-yl-phenyl)-methanone;
- 5 [(S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(4-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- [(S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(4,5-dimethoxy-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- [5-(3-Chloro-phenyl)-2-methyl-thiazol-4-yl]-[(S)-2-methyl-2-(4-methyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-
- 10 methanone;
- [(S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(4,5-dimethyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- [(S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-pyrazol-1-yl-phenyl)-methanone;
- 15 [(S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(6-methyl-3-pyrazol-1-yl-pyridin-2-yl)-methanone;
- [(S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-fluoro-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- [(S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(2-[1,2,3]triazol-2-yl-phenyl)-
- 20 methanone;
- [(S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(6-methyl-3-[1,2,3]triazol-2-yl-pyridin-2-yl)-methanone;
- [(S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(2-fluoro-6-[1,2,3]triazol-2-yl-phenyl)-methanone;
- 25 [(S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-[5-(3-chloro-phenyl)-2-methyl-thiazol-4-yl]-methanone;
- (4,5-Dimethyl-2-[1,2,3]triazol-2-yl-phenyl)-[(S)-2-methyl-2-(4-methyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- (5-Chloro-2-[1,2,3]triazol-2-yl-phenyl)-[(S)-2-(5,6-dimethyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-
- 30 methanone;
- [(S)-2-(5,6-Dimethyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methoxy-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- [(S)-2-(5,6-Dimethyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(4-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;

- [(S)-2-(5,6-Dimethyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(4,5-dimethyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- (5-Chloro-2-[1,2,3]triazol-2-yl-phenyl)-[(S)-2-(5,6-dimethoxy-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-methanone;
- 5 [(S)-2-(5,6-Dimethoxy-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methoxy-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- [(S)-2-(5,6-Dichloro-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- [(S)-2-(5,6-Dimethoxy-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(2-[1,2,3]triazol-2-yl-5-trifluoromethyl-phenyl)-methanone;
- 10 [(S)-2-(5,6-Dimethoxy-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(4,5-dimethyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- [(S)-2-(5,6-Dimethoxy-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-pyrazol-1-yl-phenyl)-methanone;
- 15 [(S)-2-(5,6-Dimethyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(3'-methoxy-4-methyl-biphenyl-2-yl)-methanone;
- [(S)-2-(5,6-Dimethyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(4'-methoxy-4-methyl-biphenyl-2-yl)-methanone;
- [(S)-2-(5,6-Dimethyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(2'-fluoro-4-methyl-biphenyl-2-yl)-methanone;
- 20 [(S)-2-(5,6-Dimethyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(4'-fluoro-4-methyl-biphenyl-2-yl)-methanone;
- [(S)-2-(5,6-Dimethyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(3'-fluoro-4-methyl-biphenyl-2-yl)-methanone;
- 25 [(S)-2-(5,6-Dimethyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(4,3'-dimethyl-biphenyl-2-yl)-methanone;
- [5-(4-Bromo-phenyl)-2-methyl-thiazol-4-yl]-[(S)-2-(7-methyl-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-methanone;
- [(S)-2-(7-Methyl-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(2-methyl-5-m-tolyl-thiazol-4-yl)-methanone;
- [5-(3-Bromo-phenyl)-2-methyl-thiazol-4-yl]-[(S)-2-(7-methyl-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-methanone;
- 30 [(S)-2-(7-Methyl-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(5-(3,5-Dimethyl-phenyl)-2-methyl-thiazol-4-yl)-methanone;
- [(S)-2-(7-methyl-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(5-(3,5-Dimethyl-phenyl)-2-methyl-thiazol-4-yl)-methanone;
- [5-(3-Bromo-4-fluoro-phenyl)-2-methyl-thiazol-4-yl]-[(S)-2-(7-methyl-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-methanone;

- [(S)-2-(7-Methyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-[2-methyl-5-(3-trifluoromethyl-phenyl)-thiazol-4-yl]-methanone;
- [5-(3,4-Dichloro-phenyl)-2-methyl-thiazol-4-yl]-[(S)-2-(7-methyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-methanone;
- 5 [5-(3-Methoxy-phenyl)-2-methyl-thiazol-4-yl]-[(S)-2-(7-methyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-methanone;
- [5-(4-Bromo-phenyl)-2-methyl-thiazol-4-yl]-[(S)-2-(6-chloro-7-methyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-methanone;
- 10 [(S)-2-(6-Chloro-7-methyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-[5-(6-methoxy-pyridin-3-yl)-2-methyl-thiazol-4-yl]-methanone;
- [(S)-2-(6-Chloro-7-methyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-[5-(4-fluoro-phenyl)-2-methyl-thiazol-4-yl]-methanone;
- [(S)-2-(6-Chloro-7-methyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-[5-(4-chloro-phenyl)-2-methyl-thiazol-4-yl]-methanone;
- 15 [(S)-2-(6-Chloro-7-methyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-[2-methyl-5-(3-trifluoromethyl-phenyl)-thiazol-4-yl]-methanone;
- [(S)-2-(6-Chloro-7-methyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-[5-(3-methoxy-phenyl)-2-methyl-thiazol-4-yl]-methanone;
- [(S)-2-(6-Chloro-7-methyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-[2-methyl-5-m-tolyl-thiazol-4-yl]-methanone;
- 20 [5-(3-Bromo-phenyl)-2-methyl-thiazol-4-yl]-[(S)-2-(6-chloro-7-methyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-methanone;
- [(S)-2-(6-Chloro-7-methyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-[5-(3,5-dimethyl-phenyl)-2-methyl-thiazol-4-yl]-methanone;
- 25 [5-(3-Bromo-4-fluoro-phenyl)-2-methyl-thiazol-4-yl]-[(S)-2-(6-chloro-7-methyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-methanone;
- [(S)-2-(6-Chloro-7-methyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-[5-(3,4-dichloro-phenyl)-2-methyl-thiazol-4-yl]-methanone;
- [(S)-2-(6-Chloro-7-methyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-[6-methyl-3-[1,2,3]triazol-2-yl-pyridin-2-yl]-methanone;
- 30 [(S)-2-(6-Chloro-7-methyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-[5-chloro-2-[1,2,3]triazol-2-yl-phenyl]-methanone;
- [(S)-2-(6-Chloro-7-methyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-[4,5-dimethyl-2-[1,2,3]triazol-2-yl-phenyl]-methanone;

- [(S)-2-(6-Chloro-7-methyl-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(5-methoxy-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- [(S)-2-(6-Chloro-7-methyl-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(2-fluoro-3-methyl-6-[1,2,3]triazol-2-yl-phenyl)-methanone;
- 5 [(S)-2-(6-Chloro-7-methyl-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(2-fluoro-3-methoxy-6-[1,2,3]triazol-2-yl-phenyl)-methanone;
- [(S)-2-(6-Chloro-5-trifluoromethyl-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-[5-(3,4-dimethyl-phenyl)-2-methyl-thiazol-4-yl]-methanone;
- [2-(5-Methoxy-1H-benzoimidazol-2-yl)-2-methyl-4-methylene-pyrrolidin-1-yl]-(2-methyl-5-m-tolyl-thiazol-4-yl)-
- 10 methanone;
- [2-Dimethylamino-5-(3,4-dimethyl-phenyl)-thiazol-4-yl]-[2-(5-methoxy-1H-benzoimidazol-2-yl)-2-methyl-4-methylene-pyrrolidin-1-yl]-methanone;
- [5-(3,4-Dimethyl-phenyl)-2-methyl-thiazol-4-yl]-[2-(5-methoxy-1H-benzoimidazol-2-yl)-2-methyl-4-methylene-pyrrolidin-1-yl]-methanone;
- 15 [5-(4-Chloro-phenyl)-2-methyl-thiazol-4-yl]-[2-(5-methoxy-1H-benzoimidazol-2-yl)-2-methyl-4-methylene-pyrrolidin-1-yl]-methanone;
- [2-(5-Methoxy-1H-benzoimidazol-2-yl)-2-methyl-4-methylene-pyrrolidin-1-yl]-[5-(4-methoxy-phenyl)-2-methyl-thiazol-4-yl]-methanone;
- [2-(5-Methoxy-1H-benzoimidazol-2-yl)-2-methyl-4-methylene-pyrrolidin-1-yl]-(2-methyl-5-p-tolyl-thiazol-4-yl)-
- 20 methanone;
- [5-(3-Chloro-phenyl)-2-dimethylamino-thiazol-4-yl]-[2-(5-methoxy-1H-benzoimidazol-2-yl)-2-methyl-4-methylene-pyrrolidin-1-yl]-methanone;
- [5-(3-Chloro-phenyl)-2-methyl-thiazol-4-yl]-[2-(5-methoxy-1H-benzoimidazol-2-yl)-2-methyl-4-methylene-pyrrolidin-1-yl]-methanone;
- 25 [2-(5-Methoxy-1H-benzoimidazol-2-yl)-2-methyl-4-methylene-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- (5-Chloro-2-[1,2,3]triazol-2-yl-phenyl)-[2-(5-methoxy-1H-benzoimidazol-2-yl)-2-methyl-4-methylene-pyrrolidin-1-yl]-methanone;
- (4,5-Dimethyl-2-[1,2,3]triazol-2-yl-phenyl)-[2-(5-methoxy-1H-benzoimidazol-2-yl)-2-methyl-4-methylene-pyrrolidin-1-yl]-methanone;
- 30 [(S)-2-(5-tert-Butyl-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(4,5-dimethyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- [(S)-2-(5-tert-Butyl-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(4-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;

- [(S)-2-(5-tert-Butyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(4,5-dimethoxy-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- [(S)-2-(5-Chloro-6-trifluoromethyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(2-fluoro-6-[1,2,3]triazol-2-yl-phenyl)-methanone;
- 5 [(S)-2-(5-Chloro-6-trifluoromethyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(5-fluoro-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- (5-Chloro-2-[1,2,3]triazol-2-yl-phenyl)-[(S)-2-(5-chloro-6-trifluoromethyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-methanone;
- [(S)-2-(5-Chloro-6-trifluoromethyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- 10 [(S)-2-(5-Chloro-6-trifluoromethyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(2-fluoro-3-methoxy-6-[1,2,3]triazol-2-yl-phenyl)-methanone;
- [(S)-2-(5-Chloro-6-trifluoromethyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(2-fluoro-3-methyl-6-[1,2,3]triazol-2-yl-phenyl)-methanone;
- 15 [(S)-2-(5-Chloro-6-trifluoromethyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(4,5-dimethyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- [(S)-2-(5-Chloro-6-trifluoromethyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(4-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- [(S)-2-(5-Chloro-6-trifluoromethyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(4,5-dimethoxy-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- 20 [(S)-2-(5-Chloro-6-trifluoromethyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(5-methyl-2-pyrazol-1-yl-phenyl)-methanone;
- [(S)-2-(5-Chloro-6-trifluoromethyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(6-methyl-3-pyrazol-1-yl-pyridin-2-yl)-methanone;
- 25 [(S)-2-(5-Chloro-6-trifluoromethyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(6-methyl-3-[1,2,3]triazol-2-yl-pyridin-2-yl)-methanone;
- [(S)-2-(5,6-Dimethyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(5-fluoro-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- (5-Chloro-2-[1,2,3]triazol-2-yl-phenyl)-[(S)-2-(5,6-dimethyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-methanone;
- 30 [(S)-2-(5,6-Dimethyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- [(S)-2-(5,6-Dimethyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(2-fluoro-3-methoxy-6-[1,2,3]triazol-2-yl-phenyl)-methanone;
- [(S)-2-(5,6-Dimethyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(2-fluoro-3-methyl-6-[1,2,3]triazol-2-yl-phenyl)-methanone;
- 35 [(S)-2-(5,6-Dimethyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(2-fluoro-3-methyl-6-[1,2,3]triazol-2-yl-phenyl)-methanone;

- [(S)-2-(5,6-Dimethyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(4,5-dimethyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
[5-(3-Bromo-4-fluoro-phenyl)-2-methyl-thiazol-4-yl]-[(S)-2-(5-methoxy-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-methanone;
- 5 (2-Cyclopropyl-5-m-tolyl-thiazol-4-yl)-[(S)-2-(5-methoxy-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-methanone;
[2-Dimethylamino-5-(3,4-dimethyl-phenyl)-thiazol-4-yl]-[(S)-2-(5-methoxy-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-methanone;
[5-(3-Bromo-phenyl)-2-cyclopropyl-thiazol-4-yl]-[(S)-2-(5-methoxy-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-
- 10 1-yl]-methanone;
[2-Cyclopropyl-5-(3-methoxy-phenyl)-thiazol-4-yl]-[(S)-2-(5-methoxy-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-methanone;
[2-(3,4-Dimethyl-phenyl)-5-methyl-thiophen-3-yl]-[(S)-2-(5-methoxy-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-methanone;
- 15 [(S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(2-methyl-5-phenyl-thiazol-4-yl)-methanone;
[(S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
[5-(3-Chloro-phenyl)-2-methyl-thiazol-4-yl]-[(S)-2-(4-methyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-
- 20 methanone;
[(S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-[5-(3-chloro-phenyl)-2-methyl-thiazol-4-yl]-methanone;
[5-(3,4-Dimethyl-phenyl)-2-methyl-thiazol-4-yl]-[(S)-2-(6-methoxy-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-methanone;
- 25 [(S)-2-(4,5-Dimethyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-[5-(3,4-dimethyl-phenyl)-2-methyl-thiazol-4-yl]-methanone;
[(S)-2-(4,7-Dimethoxy-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-[5-(3,4-dimethyl-phenyl)-2-methyl-thiazol-4-yl]-methanone;
[5-(3,4-Dimethyl-phenyl)-2-methyl-thiazol-4-yl]-[(S)-4-methylene-2-(4-methylsulfanyl-1H-benzimidazol-2-yl)-
- 30 pyrrolidin-1-yl]-methanone;
[(S)-2-(4-Bromo-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-[5-(3,4-dimethyl-phenyl)-2-methyl-thiazol-4-yl]-methanone;
[(S)-2-(4-Chloro-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-[5-(3,4-dimethyl-phenyl)-2-methyl-thiazol-4-yl]-methanone;

- [5-(3,4-Dimethyl-phenyl)-2-methyl-thiazol-4-yl]-[(S)-4-methylene-2-(5-methylsulfanyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- [5-(3,4-Dimethyl-phenyl)-2-methyl-thiazol-4-yl]-[(S)-2-(4-methyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-methanone;
- 5 [(S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-[5-(3,4-dimethyl-phenyl)-2-methyl-thiazol-4-yl]-methanone;
- [(S)-2-(5-Chloro-6-methyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-[5-(3,4-dimethyl-phenyl)-2-methyl-thiazol-4-yl]-methanone;
- [(S)-2-(5,6-Dimethyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-[5-(3,4-dimethyl-phenyl)-2-methyl-
- 10 thiazol-4-yl]-methanone;
- [(S)-2-(5-tert-Butyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-[5-(3,4-dimethyl-phenyl)-2-methyl-thiazol-4-yl]-methanone;
- [(S)-2-(4,6-Dimethyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-[5-(3,4-dimethyl-phenyl)-2-methyl-thiazol-4-yl]-methanone;
- 15 [(S)-2-(5,6-Dimethyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- [(S)-2-(4,6-Dimethyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- [(S)-2-(5-Chloro-6-methyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-
- 20 phenyl)-methanone;
- [(S)-2-(5-Chloro-6-trifluoromethyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- [(S)-2-(5-Methoxy-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(2-methyl-5-m-tolyl-thiazol-4-yl)-methanone;
- 25 [2-(3,4-Dimethyl-phenyl)-5-methyl-thiophen-3-yl]-[2-(5-methoxy-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-methanone;
- [2-(3,4-Dimethyl-phenyl)-5-methyl-thiophen-3-yl]-[4-methylene-2-(7-methylsulfanyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- [2-(3,4-Dimethyl-phenyl)-5-methyl-thiophen-3-yl]-[4-methylene-2-(5-methylsulfanyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- 30 [2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-[2-(3,4-dimethyl-phenyl)-5-methyl-thiophen-3-yl]-methanone;
- [2-(5,6-Dimethyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-[2-(3,4-dimethyl-phenyl)-5-methyl-thiophen-3-yl]-methanone;

- [2-(4,6-Dimethyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-[2-(3,4-dimethyl-phenyl)-5-methyl-thiophen-3-yl]-methanone;
- [2-(4,5-Dimethyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-[2-(3,4-dimethyl-phenyl)-5-methyl-thiophen-3-yl]-methanone;
- 5 [2-(4-Bromo-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-[2-(3,4-dimethyl-phenyl)-5-methyl-thiophen-3-yl]-methanone;
- [2-(3,4-Dimethyl-phenyl)-5-methyl-thiophen-3-yl]-[2-(4-methyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-methanone;
- [5-(3-Chloro-phenyl)-2-methyl-thiazol-4-yl]-[(S)-2-(5-methoxy-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-methanone;
- 10 [2-(5-Chloro-6-methyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-[2-(3,4-dimethyl-phenyl)-5-methyl-thiophen-3-yl]-methanone;
- [2-(5-tert-Butyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-[2-(3,4-dimethyl-phenyl)-5-methyl-thiophen-3-yl]-methanone;
- 15 [2-(4,7-Dimethoxy-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-[2-(3,4-dimethyl-phenyl)-5-methyl-thiophen-3-yl]-methanone;
- [2-(7-Chloro-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-[2-(3,4-dimethyl-phenyl)-5-methyl-thiophen-3-yl]-methanone;
- (2-Cyclopropyl-5-m-tolyl-thiazol-4-yl)-[2-(5-methoxy-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-methanone;
- 20 (2-Cyclopropyl-5-m-tolyl-thiazol-4-yl)-[4-methylene-2-(7-methylsulfanyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- (2-Cyclopropyl-5-m-tolyl-thiazol-4-yl)-[4-methylene-2-(5-methylsulfanyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- 25 [2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(2-cyclopropyl-5-m-tolyl-thiazol-4-yl)-methanone;
- [5-(3,4-Dimethyl-phenyl)-2-methyl-thiazol-4-yl]-[(S)-2-(5-methoxy-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-methanone;
- (2-Cyclopropyl-5-m-tolyl-thiazol-4-yl)-[2-(5,6-dimethyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-methanone;
- 30 (2-Cyclopropyl-5-m-tolyl-thiazol-4-yl)-[2-(4,6-dimethyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-methanone;
- (2-Cyclopropyl-5-m-tolyl-thiazol-4-yl)-[2-(4,5-dimethyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-methanone;
- 35 [2-(4-Bromo-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(2-cyclopropyl-5-m-tolyl-thiazol-4-yl)-methanone;

- (2-Cyclopropyl-5-m-tolyl-thiazol-4-yl)-[2-(4-methyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-methanone;
- [2-(5-Chloro-6-methyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(2-cyclopropyl-5-m-tolyl-thiazol-4-yl)-methanone;
- 5 [2-(5-Chloro-6-trifluoromethyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(2-cyclopropyl-5-m-tolyl-thiazol-4-yl)-methanone;
- (2-Cyclopropyl-5-m-tolyl-thiazol-4-yl)-[2-(4,7-dimethoxy-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-methanone;
- [2-(7-Chloro-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(2-cyclopropyl-5-m-tolyl-thiazol-4-yl)-methanone;
- 10 [(S)-2-(5-Chloro-1,4-dimethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- [(S)-2-(6-Chloro-1,7-dimethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- [(S)-2-(5-Bromo-1,7-dimethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- 15 [(S)-2-(6-Bromo-1,4-dimethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- (S)-(5-(3-chlorophenyl)-2-methylthiazol-4-yl)(2-(1,4-dimethyl-1H-benzo[d]imidazol-2-yl)-2-methylpyrrolidin-1-yl)methanone;
- 20 (S)-(5-(3-chlorophenyl)-2-methylthiazol-4-yl)(2-(1,7-dimethyl-1H-benzo[d]imidazol-2-yl)-2-methylpyrrolidin-1-yl)methanone;
- (S)-(5-(3-chlorophenyl)-2-methylthiazol-4-yl)(2-(1-ethyl-4-methyl-1H-benzo[d]imidazol-2-yl)-2-methylpyrrolidin-1-yl)methanone;
- (S)-(5-(3-chlorophenyl)-2-methylthiazol-4-yl)(2-(1-ethyl-7-methyl-1H-benzo[d]imidazol-2-yl)-2-methylpyrrolidin-1-yl)methanone;
- 25 (S)-(2-(5-chloro-1,4-dimethyl-1H-benzo[d]imidazol-2-yl)-2-methylpyrrolidin-1-yl)(2-methyl-5-phenylthiazol-4-yl)methanone;
- (S)-(2-(6-chloro-1,7-dimethyl-1H-benzo[d]imidazol-2-yl)-2-methylpyrrolidin-1-yl)(2-methyl-5-phenylthiazol-4-yl)methanone;
- 30 (S)-(2-(5-chloro-1-ethyl-4-methyl-1H-benzo[d]imidazol-2-yl)-2-methylpyrrolidin-1-yl)(2-methyl-5-phenylthiazol-4-yl)methanone; and
- (S)-(2-(6-chloro-1-ethyl-7-methyl-1H-benzo[d]imidazol-2-yl)-2-methylpyrrolidin-1-yl)(2-methyl-5-phenylthiazol-4-yl)methanone.

47) In addition to the above-listed compounds, further compounds according to embodiment
 35 1) are selected from:

- [5-(3-Chloro-phenyl)-2-methyl-thiazol-4-yl]-[2-(4-chloro-6-trifluoromethyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- [4-(3-Chloro-phenyl)-2-methyl-pyrimidin-5-yl]-[(S)-2-(7-methylsulfanyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- 5 [(2S,3S)-3-Methyl-2-(4-methyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- [5-(3-Chloro-phenyl)-2-methyl-thiazol-4-yl]-[(2S,3S)-2-(5,6-dimethyl-1H-benzoimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-methanone;
- 10 [(2S,3S)-2-(6-Chloro-4-methyl-1H-benzoimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-[5-(4-fluoro-phenyl)-2-methyl-thiazol-4-yl]-methanone;
- [(2S,3S)-2-(6-Chloro-4-methyl-1H-benzoimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- [(2S,3S)-2-(5-Chloro-1H-benzoimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-[5-(3-chloro-phenyl)-2-methyl-thiazol-4-yl]-methanone;
- 15 [5-(3-Chloro-phenyl)-2-methyl-thiazol-4-yl]-[(2S,3S)-2-(5,6-difluoro-1H-benzoimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-methanone;
- 2-[(2S,3S)-1-[5-(3-Chloro-phenyl)-2-methyl-thiazole-4-carbonyl]-3-methyl-pyrrolidin-2-yl]-1H-benzoimidazole-4-carboxylic acid methyl ester;
- [5-(3-Chloro-phenyl)-2-methyl-thiazol-4-yl]-[(2S,3S)-2-(4-methoxy-1H-benzoimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-methanone;
- 20 [(2S,3S)-2-(5-Chloro-6-methyl-1H-benzoimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-[5-(3-chloro-phenyl)-2-methyl-thiazol-4-yl]-methanone;
- [5-(3-Chloro-phenyl)-2-methyl-thiazol-4-yl]-[(2S,3S)-2-(4,6-dimethyl-1H-benzoimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-methanone;
- 25 [5-(3-Chloro-phenyl)-2-methyl-thiazol-4-yl]-[(2S,3S)-2-(4-hydroxymethyl-1H-benzoimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-methanone;
- [5-(3-Chloro-phenyl)-2-methyl-thiazol-4-yl]-[(2S,3S)-3-methyl-2-(4-trifluoromethyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- [5-(3-Chloro-phenyl)-2-methyl-thiazol-4-yl]-[(2S,3S)-2-(5-fluoro-1H-benzoimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-methanone;
- 30 [(2S,3S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-[5-(3-chloro-phenyl)-2-methyl-thiazol-4-yl]-methanone;
- [(2S,3S)-2-(4-Chloro-1H-benzoimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-[5-(3-chloro-phenyl)-2-methyl-thiazol-4-yl]-methanone;

- [5-(3-Chloro-phenyl)-2-methyl-thiazol-4-yl]-[(2S,3S)-2-(4,5-difluoro-1H-benzoimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-methanone;
- [5-(3-Chloro-phenyl)-2-methyl-thiazol-4-yl]-[(2S,3S)-2-(4,5-dimethyl-1H-benzoimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-methanone;
- 5 [5-(3-Chloro-phenyl)-2-methyl-thiazol-4-yl]-[(2S,3S)-2-(4-isopropyl-1H-benzoimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-methanone;
- [5-(3-Chloro-phenyl)-2-methyl-thiazol-4-yl]-[(2S,3S)-2-(5-methoxy-1H-benzoimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-methanone;
- 10 [(2S,3S)-2-(5-Chloro-7-methyl-1H-benzoimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-[5-(3-chloro-phenyl)-2-methyl-thiazol-4-yl]-methanone;
- (5-Methyl-2-[1,2,3]triazol-2-yl-phenyl)-[(2S,3S)-3-methyl-2-(5-trifluoromethyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- [5-(3-Chloro-phenyl)-2-methyl-thiazol-4-yl]-[(2S,3S)-3-methyl-2-(5-methyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- 15 [(2S,3S)-2-(5,6-Dimethyl-1H-benzoimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-[5-methyl-2-[1,2,3]triazol-2-yl-phenyl]-methanone;
- [(2S,3S)-2-(5-Chloro-6-methyl-1H-benzoimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-[5-methyl-2-[1,2,3]triazol-2-yl-phenyl]-methanone;
- 20 [(2S,3S)-2-(4,6-Dimethyl-1H-benzoimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-[5-methyl-2-[1,2,3]triazol-2-yl-phenyl]-methanone;
- [(2S,3S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-[5-methyl-2-[1,2,3]triazol-2-yl-phenyl]-methanone;
- [(2S,3S)-2-(4,5-Dimethyl-1H-benzoimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-[5-methyl-2-[1,2,3]triazol-2-yl-phenyl]-methanone;
- 25 [(2S,3S)-2-(5-Chloro-7-methyl-1H-benzoimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-[5-methyl-2-[1,2,3]triazol-2-yl-phenyl]-methanone;
- [(2S,3S)-3-Methyl-2-(4-methyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-[2-methyl-5-phenyl-thiazol-4-yl]-methanone;
- [(2S,3S)-3-Methyl-2-(5-methyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-[2-methyl-5-phenyl-thiazol-4-yl]-methanone;
- 30 [5-(3,5-Difluoro-phenyl)-2-methyl-thiazol-4-yl]-[(2S,3S)-2-(5,6-dimethoxy-1H-benzoimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-methanone;
- [(2S,3S)-2-(5,6-Dimethoxy-1H-benzoimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-[5-(4-fluoro-phenyl)-2-methyl-thiazol-4-yl]-methanone;
- [(2S,3S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-[2-[1,2,3]triazol-2-yl-phenyl]-methanone;

- [(2S,3S)-2-(1H-Benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-[5-(3-chloro-phenyl)-2-methyl-thiazol-4-yl]-methanone;
- [(2S,3S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(5-chloro-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- 5 [(2S,3S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(4'-methyl-biphenyl-2-yl)-methanone;
- [(2S,3S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-[5-(4-chloro-phenyl)-2-methyl-thiazol-4-yl]-methanone;
- [(2S,3S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(5-methyl-2-pyrazol-1-yl-phenyl)-methanone;
- 10 [(2S,3S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(2-methyl-4-p-tolyl-thiazol-5-yl)-methanone;
- [(2S,3S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-[5-(3,5-difluoro-phenyl)-2-methyl-thiazol-4-yl]-methanone;
- 15 [(2S,3S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(2-fluoro-3-methyl-6-[1,2,3]triazol-2-yl-phenyl)-methanone;
- [(2S,3S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-[5-(6-methoxy-pyridin-3-yl)-2-methyl-thiazol-4-yl]-methanone;
- [5-(3-Chloro-phenyl)-2-methyl-thiazol-4-yl]-[(2S,3S)-3-methyl-2-(4-methyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- 20 [(2S,3S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-[5-(4-fluoro-phenyl)-2-methyl-thiazol-4-yl]-methanone;
- [(2S,3S)-2-(5-Chloro-6-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(5-chloro-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- 25 [(2S,3S)-2-(5-Chloro-6-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(4-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- [(2S,3S)-2-(5-Chloro-6-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(5-methyl-2-pyrazol-1-yl-phenyl)-methanone;
- [(2S,3S)-2-(5-Chloro-6-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(5-methoxy-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- 30 [(2S,3S)-2-(5-Chloro-6-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(2-fluoro-3-methyl-6-[1,2,3]triazol-2-yl-phenyl)-methanone;
- [(2S,3S)-2-(5-Chloro-6-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(2-[1,2,3]triazol-2-yl-phenyl)-methanone;

- [(2S,3S)-2-(5-Chloro-6-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(4'-methyl-biphenyl-2-yl)-methanone;
- [(2S,3S)-2-(5-Chloro-6-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-[5-(4-chloro-phenyl)-2-methyl-thiazol-4-yl]-methanone;
- 5 [(2S,3S)-2-(5-Chloro-6-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(2-methyl-4-p-tolyl-thiazol-5-yl)-methanone;
- [(2S,3S)-2-(5-Chloro-6-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-[5-(3,5-difluoro-phenyl)-2-methyl-thiazol-4-yl]-methanone;
- (5-Chloro-2-[1,2,3]triazol-2-yl-phenyl)-[(2S,3S)-2-(5,6-dimethyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-methanone;
- 10 [(2S,3S)-2-(5,6-Dimethyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(5-methyl-2-pyrazol-1-yl-phenyl)-methanone;
- [(2S,3S)-2-(5,6-Dimethyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(4'-methyl-biphenyl-2-yl)-methanone;
- [5-(4-Chloro-phenyl)-2-methyl-thiazol-4-yl]-[(2S,3S)-2-(5,6-dimethyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-methanone;
- 15 [(2S,3S)-2-(5,6-Dimethyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(2-methyl-4-p-tolyl-thiazol-5-yl)-methanone;
- [5-(3,5-Difluoro-phenyl)-2-methyl-thiazol-4-yl]-[(2S,3S)-2-(5,6-dimethyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-methanone;
- 20 [5-(3-Chloro-phenyl)-2-methyl-thiazol-4-yl]-[(2S,3S)-3-methyl-2-(5-trifluoromethyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- [(2S,3S)-2-(5,6-Dimethyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-[5-(4-fluoro-phenyl)-2-methyl-thiazol-4-yl]-methanone;
- [(2S,3S)-2-(6-Chloro-4-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(5-chloro-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- 25 [(2S,3S)-2-(6-Chloro-4-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(4'-methyl-biphenyl-2-yl)-methanone;
- [(2S,3S)-2-(6-Chloro-4-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(4-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- 30 [(2S,3S)-2-(6-Chloro-4-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-[5-(4-chloro-phenyl)-2-methyl-thiazol-4-yl]-methanone;
- [(2S,3S)-2-(6-Chloro-4-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(5-methyl-2-pyrazol-1-yl-phenyl)-methanone;
- [(2S,3S)-2-(6-Chloro-4-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(2-methyl-4-p-tolyl-thiazol-5-yl)-methanone;
- 35

- [(2S,3S)-2-(6-Chloro-4-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(5-methoxy-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- [(2S,3S)-2-(6-Chloro-4-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-[5-(3,5-difluoro-phenyl)-2-methyl-thiazol-4-yl]-methanone;
- 5 [(2S,3S)-2-(6-Chloro-4-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(2-fluoro-3-methyl-6-[1,2,3]triazol-2-yl-phenyl)-methanone;
- [(S)-2-(5-Chloro-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- [(S)-2-(6-Chloro-5-fluoro-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- 10 [(S)-2-(6-Bromo-5-fluoro-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-pyrazol-1-yl-phenyl)-methanone;
- [(S)-2-(6-Bromo-5-fluoro-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(6-methyl-3-[1,2,3]triazol-2-yl-pyridin-2-yl)-methanone;
- 15 [(S)-2-(6-Chloro-5-trifluoromethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(2-pyrazol-1-yl-phenyl)-methanone;
- [(S)-2-(6-Chloro-5-trifluoromethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(6-methyl-3-pyrazol-1-yl-pyridin-2-yl)-methanone;
- [(S)-2-(6-Chloro-5-trifluoromethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(2-[1,2,3]triazol-2-yl-5-trifluoromethyl-phenyl)-methanone;
- 20 [(S)-2-(6-Chloro-5-trifluoromethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(2-[1,2,3]triazol-2-yl-5-trifluoromethoxy-phenyl)-methanone;
- (5-Methyl-2-[1,2,3]triazol-2-yl-phenyl)-[(S)-2-methyl-2-(4-trifluoromethyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- 25 (5-Methyl-2-pyrazol-1-yl-phenyl)-[(S)-2-methyl-2-(5-trifluoromethyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- (6-Methyl-3-[1,2,3]triazol-2-yl-pyridin-2-yl)-[(S)-2-methyl-2-(5-trifluoromethyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- (3'-Methoxy-4-methyl-biphenyl-2-yl)-[(S)-2-methyl-2-(5-trifluoromethyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- 30 4'-Methyl-2'-[(S)-2-methyl-2-(5-trifluoromethyl-1H-benzimidazol-2-yl)-pyrrolidine-1-carbonyl]-biphenyl-4-carbonitrile;
- (2'-Fluoro-4-methyl-biphenyl-2-yl)-[(S)-2-methyl-2-(5-trifluoromethyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;

(4'-Fluoro-4-methyl-biphenyl-2-yl)-[(S)-2-methyl-2-(5-trifluoromethyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;

[(S)-2-(4,5-Difluoro-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(4,5-dimethyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;

5 (5-Methyl-2-[1,2,3]triazol-2-yl-phenyl)-[(S)-2-methyl-2-(5-trifluoromethoxy-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;

[(S)-2-(4,5-Difluoro-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(2-methyl-5-phenyl-thiazol-4-yl)-methanone;

[(S)-2-(4,5-Difluoro-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(2-methyl-5-p-tolyl-thiazol-4-yl)-methanone;

[5-(4-Chloro-phenyl)-2-methyl-thiazol-4-yl]-[(S)-2-(4,5-difluoro-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-

10 methanone;

[(S)-2-(4,5-Difluoro-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-[5-(3-fluoro-phenyl)-2-methyl-thiazol-4-yl]-methanone;

[(S)-2-(6-Fluoro-5-trifluoromethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;

15 [(S)-2-(5-Bromo-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;

(5-Methyl-2-[1,2,3]triazol-2-yl-phenyl)-[(S)-2-methyl-2-(5-trifluoromethyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;

[(2S,4R)-2-(6-Chloro-1,7-dimethyl-1H-benzimidazol-2-yl)-4-methoxy-pyrrolidin-1-yl]-(5-methoxy-2-[1,2,3]triazol-

20 2-yl-phenyl)-methanone; and

[(2S,4R)-2-(5-Chloro-1,4-dimethyl-1H-benzimidazol-2-yl)-4-methoxy-pyrrolidin-1-yl]-(5-methoxy-2-[1,2,3]triazol-2-yl-phenyl)-methanone.

48) In addition to the above-listed compounds, further compounds according to embodiment 1) are selected from:

25 [2-(5,6-Dimethoxy-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-(2-methyl-5-phenyl-thiazol-4-yl)-methanone;

[2-(4,5-Dimethyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-(2-methyl-5-phenyl-thiazol-4-yl)-methanone;

[2-(5-Methoxy-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;

(5-Fluoro-2-[1,2,3]triazol-2-yl-phenyl)-[(S)-2-(6-methylsulfanyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;

(5-Chloro-2-[1,2,3]triazol-2-yl-phenyl)-[(S)-2-(6-methylsulfanyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;

30 (5-Methoxy-2-[1,2,3]triazol-2-yl-phenyl)-[(S)-2-(6-methylsulfanyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;

3-[[(S)-2-(6-Methylsulfanyl-1H-benzimidazol-2-yl)-pyrrolidine-1-carbonyl]-4-[1,2,3]triazol-2-yl-benzonitrile;

[(S)-2-(6-Methylsulfanyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-(2-methyl-6-[1,2,3]triazol-2-yl-phenyl)-methanone;

(5-Fluoro-2-[1,2,3]triazol-2-yl-phenyl)-[(S)-2-(6-methoxy-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;

35 (5-Chloro-2-[1,2,3]triazol-2-yl-phenyl)-[(S)-2-(6-methoxy-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;

- [(S)-2-(6-Methoxy-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-(5-methoxy-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
3-[(S)-2-(6-Methoxy-1H-benzimidazol-2-yl)-pyrrolidine-1-carbonyl]-4-[1,2,3]triazol-2-yl-benzonitrile;
[(S)-2-(6-Methoxy-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
[(S)-2-(6-Methoxy-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-(4-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
5 [(S)-2-(6-Methoxy-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-(2-methyl-6-[1,2,3]triazol-2-yl-phenyl)-methanone;
(4,5-Dimethoxy-2-[1,2,3]triazol-2-yl-phenyl)-[(S)-2-(6-methoxy-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
(5-Fluoro-2-[1,2,3]triazol-2-yl-phenyl)-[(S)-2-(7-methylsulfanyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
(2-Fluoro-6-[1,2,3]triazol-2-yl-phenyl)-[(S)-2-(7-methylsulfanyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
(5-Chloro-2-[1,2,3]triazol-2-yl-phenyl)-[(S)-2-(7-methylsulfanyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
10 3-[(S)-2-(7-Methylsulfanyl-1H-benzimidazol-2-yl)-pyrrolidine-1-carbonyl]-4-[1,2,3]triazol-2-yl-benzonitrile;
[(S)-2-(6-Methylsulfanyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
[(S)-2-(7-Methylsulfanyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-(2-methyl-6-[1,2,3]triazol-2-yl-phenyl)-methanone;
(2-Fluoro-3-methyl-6-[1,2,3]triazol-2-yl-phenyl)-[(S)-2-(5-methoxy-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-
methanone;
15 (5-Methyl-2-pyrazol-1-yl-phenyl)-[(S)-2-(5-methylsulfanyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
(2-Fluoro-3-methyl-6-[1,2,3]triazol-2-yl-phenyl)-[(S)-2-(5-methylsulfanyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-
methanone;
[(S)-2-(5-Methoxy-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
(3',4'-Dimethyl-biphenyl-2-yl)-[(S)-2-(6-methoxy-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
20 [(S)-2-(6-Methylsulfanyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
[(S)-2-(7-Methoxy-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
(3',4'-Dimethyl-biphenyl-2-yl)-[(S)-2-(6-methylsulfanyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
[(S)-2-(6-Methylsulfanyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-(2-[1,2,3]triazol-2-yl-phenyl)-methanone;
25 [(S)-2-(5-Methylsulfanyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
[(S)-2-(7-Methylsulfanyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-(2-[1,2,3]triazol-2-yl-phenyl)-methanone;
[(2S,4R)-2-(6-Chloro-7-methyl-1H-benzimidazol-2-yl)-4-methoxy-pyrrolidin-1-yl]-(2-methyl-6-[1,2,3]triazol-2-yl-
phenyl)-methanone;
[(2S,4R)-2-(6-Chloro-7-methyl-1H-benzimidazol-2-yl)-4-methoxy-pyrrolidin-1-yl]-(4-methyl-2-[1,2,3]triazol-2-yl-
30 phenyl)-methanone;
[(2S,4R)-2-(6-Chloro-7-methyl-1H-benzimidazol-2-yl)-4-methoxy-pyrrolidin-1-yl]-(5-fluoro-2-[1,2,3]triazol-2-yl-
phenyl)-methanone;
(5-Chloro-2-[1,2,3]triazol-2-yl-phenyl)-[(2S,4R)-4-methoxy-2-(6-methoxy-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-
methanone;

- (4,5-Dimethyl-2-[1,2,3]triazol-2-yl-phenyl)-[(2S,4R)-4-methoxy-2-(6-methoxy-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- [(S)-2-(5,6-Dimethyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- [5-(6-Methoxy-pyridin-3-yl)-2-methyl-thiazol-4-yl]-[(S)-2-(7-methyl-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-methanone;
- 5 [5-(4-Fluoro-phenyl)-2-methyl-thiazol-4-yl]-[(S)-2-(7-methyl-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-methanone;
- (4,5-Dimethyl-2-[1,2,3]triazol-2-yl-phenyl)-[(S)-2-(7-methyl-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-methanone;
- 10 [5-(4-Chloro-phenyl)-2-methyl-thiazol-4-yl]-[(S)-2-(7-methyl-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-methanone;
- (5-Methoxy-2-[1,2,3]triazol-2-yl-phenyl)-[(S)-2-(7-methyl-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-methanone;
- [(S)-2-(7-Methyl-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(6-methyl-3-[1,2,3]triazol-2-yl-pyridin-2-yl)-methanone;
- 15 [(S)-2-(5-Methoxy-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- [(S)-2-(5-Methoxy-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- [(S)-2-(6-Chloro-7-methyl-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(5-methyl-2-pyrazol-1-yl-phenyl)-methanone;
- 20 [(S)-2-(6-Chloro-7-methyl-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(6-methyl-3-pyrazol-1-yl-pyridin-2-yl)-methanone;
- (5-Chloro-2-[1,2,3]triazol-2-yl-phenyl)-[(S)-2-(6-methoxy-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-methanone;
- 25 [(S)-2-(6-Methoxy-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(5-methyl-2-pyrazol-1-yl-phenyl)-methanone;
- (4,5-Dimethyl-2-[1,2,3]triazol-2-yl-phenyl)-[(S)-2-(6-methoxy-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-methanone;
- [(S)-2-(5-Methoxy-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(3'-methyl-biphenyl-2-yl)-methanone;
- 30 [(S)-2-(6-Methoxy-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(5-methoxy-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- (2-Fluoro-3-methyl-6-[1,2,3]triazol-2-yl-phenyl)-[(S)-2-(6-methoxy-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-methanone;
- [(S)-2-(5-tert-Butyl-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(5-fluoro-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- 35

- [(S)-2-(5-tert-Butyl-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(5-chloro-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- [(S)-2-(5-tert-Butyl-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- [(S)-2-(5-tert-Butyl-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(2-fluoro-3-methyl-6-[1,2,3]triazol-2-yl-phenyl)-methanone;
- [(S)-2-(5-tert-Butyl-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(2-methyl-6-[1,2,3]triazol-2-yl-phenyl)-methanone;
- [(S)-2-(5-tert-Butyl-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(5-methyl-2-pyrazol-1-yl-phenyl)-methanone;
- [(S)-2-(5-tert-Butyl-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(6-methyl-3-[1,2,3]triazol-2-yl-pyridin-2-yl)-methanone;
- (5-Fluoro-2-[1,2,3]triazol-2-yl-phenyl)-[(S)-2-(5-methoxy-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-methanone;
- [(S)-2-(5-Chloro-6-trifluoromethyl-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(2-methyl-6-[1,2,3]triazol-2-yl-phenyl)-methanone;
- [(S)-2-(5,6-Dimethyl-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(2-fluoro-6-[1,2,3]triazol-2-yl-phenyl)-methanone;
- [(S)-2-(5-Methoxy-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-[5-(4-methoxy-phenyl)-2-methyl-thiazol-4-yl]-methanone;
- [5-(4-Bromo-phenyl)-2-methyl-thiazol-4-yl]-[(S)-2-(5-methoxy-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-methanone;
- [(S)-2-(5-Methoxy-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(2-methyl-5-phenyl-thiazol-4-yl)-methanone;
- [5-(3-Fluoro-phenyl)-2-methyl-thiazol-4-yl]-[(S)-2-(5-methoxy-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-methanone;
- [(S)-2-(5-Methoxy-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(2-m-tolyl-thiophen-3-yl)-methanone;
- [2-(3,4-Dimethyl-phenyl)-thiophen-3-yl]-[(S)-2-(5-methoxy-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-methanone;
- [(S)-2-(4-Methyl-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(2-methyl-5-phenyl-thiazol-4-yl)-methanone;
- [5-(4-Chloro-phenyl)-2-methyl-thiazol-4-yl]-[(S)-2-(5-methoxy-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-methanone;
- [(S)-4-Methylene-2-(7-methylsulfanyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- [(S)-4-Methylene-2-(5-methylsulfanyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;

[(S)-2-(4,5-Dimethyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone; and

[(S)-2-(5-tert-Butyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone.

- 5 49) In addition to the above-listed compounds, further compounds according to embodiment 1) are selected from:

[(2S,3S)-2-(4-Chloro-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-[2-(3,4-dimethyl-phenyl)-thiophen-3-yl]-methanone;

- 10 [5-(3,4-Dimethyl-phenyl)-2-methyl-thiazol-4-yl]-[(2S,3S)-3-methyl-2-(4-trifluoromethyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- [2-(3,4-Dimethyl-phenyl)-thiophen-3-yl]-[(2S,3S)-3-methyl-2-(4-trifluoromethyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;

[(2S,3S)-2-(5,6-Dimethoxy-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(5-methyl-2-pyrimidin-2-yl-phenyl)-methanone;

- 15 [(S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(4-fluoro-biphenyl-2-yl)-methanone;
- [(S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-fluoro-2-pyridin-2-yl-phenyl)-methanone;
- [(S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(2-fluoro-6-pyridin-2-yl-phenyl)-methanone;

- 20 [(S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(2-fluoro-6-pyrimidin-2-yl-phenyl)-methanone;

[(S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(4-methyl-biphenyl-2-yl)-methanone;

[(S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-fluoro-2-pyrimidin-2-yl-phenyl)-methanone;

- 25 [(S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(3-fluoro-biphenyl-2-yl)-methanone;
- (4,5-Dimethyl-2-[1,2,3]triazol-2-yl-phenyl)-[(S)-2-methyl-2-(5-trifluoromethoxy-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- (4-Methyl-2-[1,2,3]triazol-2-yl-phenyl)-[(S)-2-methyl-2-(5-trifluoromethoxy-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;

- 30 [5-(3,4-Dimethyl-phenyl)-2-methyl-thiazol-4-yl]-[(S)-2-methyl-2-(5-trifluoromethoxy-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;

[(S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(2-methyl-5-phenyl-pyridin-4-yl)-methanone;

[(S)-2-Methyl-2-(5-trifluoromethyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-(4-pyridin-3-yl-biphenyl-2-yl)-methanone;

- [(S)-2-(6-Bromo-4,5-dimethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- [(S)-2-(6-Bromo-4,5-dimethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(2-fluoro-6-[1,2,3]triazol-2-yl-phenyl)-methanone;
- 5 Biphenyl-2-yl-[(S)-2-(6-bromo-4,5-dimethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-methanone;
- [(S)-2-(6-Bromo-4,5-dimethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(4-methyl-biphenyl-2-yl)-methanone;
- [(S)-2-(6-Bromo-4,5-dimethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(2-methyl-5-o-tolyl-thiazol-4-yl)-methanone;
- 10 [(S)-2-(6-Bromo-4,5-dimethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-(2,3-dichloro-phenyl)-2-methyl-thiazol-4-yl)-methanone;
- [(S)-2-(6-Bromo-4,5-dimethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-(3,4-dimethyl-phenyl)-2-methyl-thiazol-4-yl)-methanone;
- [(S)-2-(6-Bromo-4,5-dimethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-(3,5-difluoro-phenyl)-2-methyl-thiazol-4-yl)-methanone;
- 15 [(S)-2-(6-Bromo-4,5-dimethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-(3-methoxy-phenyl)-2-methyl-thiazol-4-yl)-methanone;
- [(S)-2-(6-Bromo-4,5-dimethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-(4-fluoro-phenyl)-2-methyl-thiazol-4-yl)-methanone;
- 20 [(S)-2-(5-Bromo-7-methyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-pyrimidin-2-yl-phenyl)-methanone;
- [(S)-2-Methyl-2-(7-methyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-(5-methyl-2-pyrimidin-2-yl-phenyl)-methanone;
- [(S)-2-(6-Chloro-5-trifluoromethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-pyrimidin-2-yl-phenyl)-methanone;
- 25 [(S)-2-(5,6-Dimethoxy-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-pyrimidin-2-yl-phenyl)-methanone;
- [(S)-2-(5,6-Dimethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-pyrimidin-2-yl-phenyl)-methanone;
- [(S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(3-fluoro-4,3'-dimethoxy-biphenyl-2-yl)-methanone;
- 30 [(S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(3,2'-difluoro-4-methoxy-biphenyl-2-yl)-methanone;
- [(S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(3,4'-difluoro-4-methoxy-biphenyl-2-yl)-methanone;

- [(S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(3,3'-difluoro-4-methoxy-biphenyl-2-yl)-methanone;
- [(S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(3-fluoro-4-methoxy-3'-methyl-biphenyl-2-yl)-methanone;
- 5 [(S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(3-fluoro-4-methoxy-4'-methyl-biphenyl-2-yl)-methanone;
- [(S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(3-fluoro-4-methoxy-3'-trifluoromethyl-biphenyl-2-yl)-methanone;
- (6-Benzo[1,3]dioxol-5-yl-2-fluoro-3-methoxy-phenyl)-[(S)-2-(5-chloro-4-methyl-1H-benzoimidazol-2-yl)-2-methyl-
- 10 pyrrolidin-1-yl]-methanone;
- 2'-[(S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidine-1-carbonyl]-3'-fluoro-4'-methoxy-biphenyl-3-carbonitrile;
- [(S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(3-fluoro-4-methoxy-3'-trifluoromethoxy-biphenyl-2-yl)-methanone;
- 15 [(S)-2-(5,6-Dimethyl-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(4-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- (4,5-Dimethoxy-2-[1,2,3]triazol-2-yl-phenyl)-[(S)-2-(5,6-dimethyl-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-methanone;
- (4-Methyl-2-[1,2,3]triazol-2-yl-phenyl)-[(S)-2-methyl-2-(1,5,6-trimethyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-
- 20 methanone;
- [(S)-2-(1-Ethyl-5,6-dimethyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(4-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- [(S)-2-(5,6-Dimethoxy-1-methyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(4,5-dimethyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- 25 (4,5-Dimethyl-2-[1,2,3]triazol-2-yl-phenyl)-[(S)-2-(1-ethyl-5,6-dimethoxy-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-methanone;
- (4'-Fluoro-4-methyl-biphenyl-2-yl)-[(S)-2-methyl-2-(1,5,6-trimethyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- [(S)-2-(1-Ethyl-5,6-dimethyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(4'-fluoro-4-methyl-biphenyl-2-yl)-
- 30 methanone;
- (4,5-Dimethyl-2-[1,2,3]triazol-2-yl-phenyl)-[(S)-2-methyl-2-(1,5,6-trimethyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- [(S)-2-(5-Chloro-1,4-dimethyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(6-methyl-3-[1,2,3]triazol-2-yl-pyridin-2-yl)-methanone;

- [(S)-2-(5-Chloro-1,4-dimethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- [(S)-2-(6-Chloro-1-methyl-5-trifluoromethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- 5 [(S)-2-(5-Chloro-1-ethyl-4-methyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- [(S)-2-(5-Chloro-1,4-dimethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-chloro-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- [(S)-2-(5-Chloro-1,4-dimethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methoxy-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- 10 (4,3'-Dimethyl-biphenyl-2-yl)-[(S)-2-methyl-2-(1,5,6-trimethyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- (4,3'-Dimethyl-biphenyl-2-yl)-[(S)-2-(1-ethyl-5,6-dimethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-methanone;
- (4,3'-Dimethyl-biphenyl-2-yl)-{(S)-2-[1-(2-fluoro-ethyl)-5,6-dimethyl-1H-benzimidazol-2-yl]-2-methyl-pyrrolidin-1-yl}-methanone;
- 15 (4,3'-Dimethyl-biphenyl-2-yl)-{(S)-2-[1-(2-methoxy-ethyl)-5,6-dimethyl-1H-benzimidazol-2-yl]-2-methyl-pyrrolidin-1-yl}-methanone;
- (3'-Fluoro-4-methyl-biphenyl-2-yl)-[(S)-2-methyl-2-(1,5,6-trimethyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- 20 [(S)-2-(1-Ethyl-5,6-dimethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(3'-fluoro-4-methyl-biphenyl-2-yl)-methanone;
- [(S)-2-(5-Chloro-1,4-dimethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-[5-(3-chloro-phenyl)-2-methyl-thiazol-4-yl]-methanone;
- [(S)-2-(1-Cyclopropylmethyl-5,6-dimethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(3'-fluoro-4-methyl-biphenyl-2-yl)-methanone;
- 25 {(S)-2-[1-(2-Fluoro-ethyl)-5,6-dimethyl-1H-benzimidazol-2-yl]-2-methyl-pyrrolidin-1-yl}-(3'-fluoro-4-methyl-biphenyl-2-yl)-methanone;
- (4,4'-Dimethyl-biphenyl-2-yl)-[(S)-2-methyl-2-(1,5,6-trimethyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- (4,4'-Dimethyl-biphenyl-2-yl)-[(S)-2-(1-ethyl-5,6-dimethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-methanone;
- 30 (4,4'-Dimethyl-biphenyl-2-yl)-{(S)-2-[1-(2-fluoro-ethyl)-5,6-dimethyl-1H-benzimidazol-2-yl]-2-methyl-pyrrolidin-1-yl}-methanone;
- (3'-Methoxy-4-methyl-biphenyl-2-yl)-[(S)-2-methyl-2-(1,5,6-trimethyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;

(2'-Fluoro-4-methyl-biphenyl-2-yl)-[(S)-2-methyl-2-(1,5,6-trimethyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone;

[(S)-2-(1-Ethyl-5,6-dimethyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(3'-methoxy-4-methyl-biphenyl-2-yl)-methanone;

5 {[(S)-2-[1-(2-Fluoro-ethyl)-5,6-dimethyl-1H-benzoimidazol-2-yl]-2-methyl-pyrrolidin-1-yl]}-(3'-methoxy-4-methyl-biphenyl-2-yl)-methanone;

{[(S)-2-[1-(2-Fluoro-ethyl)-5,6-dimethyl-1H-benzoimidazol-2-yl]-2-methyl-pyrrolidin-1-yl]}-(2'-fluoro-4-methyl-biphenyl-2-yl)-methanone; and

10 [(S)-2-(5,6-Dimethoxy-1-methyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone.

50) In addition to the above-listed compounds, further compounds according to embodiment 1) are selected from:

[(2S,3S)-2-(5-Chloro-6-methyl-1H-benzoimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(5-methyl-2-pyridin-2-yl-phenyl)-methanone;

15 (4-Chloro-biphenyl-2-yl)-[(2S,3S)-2-(5-chloro-6-methyl-1H-benzoimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-methanone;

[(2S,3S)-2-(5-Chloro-6-methyl-1H-benzoimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(5-methyl-2-pyridin-3-yl-phenyl)-methanone;

20 [(2S,3S)-2-(5-Chloro-6-methyl-1H-benzoimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(4-fluoro-biphenyl-2-yl)-methanone;

[(2S,3S)-2-(5-Chloro-6-methyl-1H-benzoimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(5-fluoro-2-pyridin-2-yl-phenyl)-methanone;

2-[(2S,3S)-2-(5-Chloro-6-methyl-1H-benzoimidazol-2-yl)-3-methyl-pyrrolidine-1-carbonyl]-biphenyl-4-carbonitrile;

25 [(2S,3S)-2-(5-Chloro-6-methyl-1H-benzoimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(4-methyl-2-pyridin-2-yl-phenyl)-methanone;

[(2S,3S)-2-(5-Chloro-6-methyl-1H-benzoimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(4-methyl-2-pyridin-3-yl-phenyl)-methanone;

[(2S,3S)-2-(5-Chloro-6-methyl-1H-benzoimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(5-methyl-biphenyl-2-yl)-methanone;

30 [(2S,3S)-2-(5-Chloro-6-methyl-1H-benzoimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(3-fluoro-biphenyl-2-yl)-methanone;

[(2S,3S)-2-(5-Chloro-6-methyl-1H-benzoimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(2-fluoro-3-methyl-6-pyridin-2-yl-phenyl)-methanone;

35 [(2S,3S)-2-(5-Chloro-6-methyl-1H-benzoimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(3-fluoro-4-methyl-biphenyl-2-yl)-methanone;

- [(2S,3S)-2-(5-Chloro-6-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(4-methyl-biphenyl-2-yl)-methanone;
- [(2S,3S)-2-(5-Chloro-6-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(2-fluoro-3-methyl-6-pyridin-3-yl-phenyl)-methanone;
- 5 [(2S,3S)-2-(4-Chloro-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-[5-(3,5-difluoro-phenyl)-2-methyl-thiazol-4-yl]-methanone;
- [(2S,3S)-2-(4-Chloro-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-[5-(4-chloro-phenyl)-2-methyl-thiazol-4-yl]-methanone;
- [(2S,3S)-2-(4-Chloro-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-[5-(3-methoxy-phenyl)-2-methyl-thiazol-4-yl]-methanone;
- 10 [(2S,3S)-2-(4-Chloro-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(2-methyl-5-m-tolyl-thiazol-4-yl)-methanone;
- [5-(4-Bromo-phenyl)-2-methyl-thiazol-4-yl]-[(2S,3S)-2-(4-chloro-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-methanone;
- [(2S,3S)-2-(4-Chloro-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(2-methyl-5-p-tolyl-thiazol-4-yl)-methanone;
- 15 [(2S,3S)-2-(4-Chloro-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-[5-(3,4-dimethyl-phenyl)-2-methyl-thiazol-4-yl]-methanone;
- [(2S,3S)-2-(4-Chloro-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-[2-dimethylamino-5-(3,4-dimethyl-phenyl)-thiazol-4-yl]-methanone;
- [(2S,3S)-2-(5-Chloro-6-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(4-methyl-2-pyrimidin-2-yl-phenyl)-methanone;
- 20 [(2S,3S)-2-(5-Chloro-6-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(5-fluoro-2-pyrimidin-2-yl-phenyl)-methanone;
- [(2S,3S)-2-(5-Chloro-6-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(2-fluoro-6-pyrimidin-2-yl-phenyl)-methanone;
- 25 [(2S,3S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-[2-(3-methoxy-phenylethynyl)-phenyl]-methanone;
- [(2S,3S)-2-(5-Chloro-6-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-[2-(3-methoxy-phenylethynyl)-phenyl]-methanone;
- [5-(4-Fluoro-phenyl)-2-methyl-thiazol-4-yl]-[(2S,3S)-3-methyl-2-(5-trifluoromethyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- 30 [5-(3-Methoxy-phenyl)-2-methyl-thiazol-4-yl]-[(2S,3S)-3-methyl-2-(5-trifluoromethyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- (2-Methyl-5-m-tolyl-thiazol-4-yl)-[(2S,3S)-3-methyl-2-(5-trifluoromethyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;

- [5-(4-Chloro-phenyl)-2-methyl-thiazol-4-yl]-[(2S,3S)-3-methyl-2-(5-trifluoromethyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- (2-Methyl-5-p-tolyl-thiazol-4-yl)-[(2S,3S)-3-methyl-2-(5-trifluoromethyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- 5 [5-(4-Methoxy-phenyl)-2-methyl-thiazol-4-yl]-[(2S,3S)-3-methyl-2-(5-trifluoromethyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- [5-(4-Bromo-phenyl)-2-methyl-thiazol-4-yl]-[(2S,3S)-3-methyl-2-(5-trifluoromethyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- [5-(3,4-Difluoro-phenyl)-2-methyl-thiazol-4-yl]-[(2S,3S)-3-methyl-2-(5-trifluoromethyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- 10 [2-(3,4-Dimethyl-phenyl)-thiophen-3-yl]-[(2S,3S)-3-methyl-2-(5-trifluoromethyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- [5-(3,4-Dimethyl-phenyl)-2-methyl-thiazol-4-yl]-[(2S,3S)-3-methyl-2-(5-trifluoromethyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- 15 [5-(2-Chloro-phenyl)-2-methyl-thiazol-4-yl]-[(2S,3S)-3-methyl-2-(5-trifluoromethyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- [5-(2-Methoxy-phenyl)-2-methyl-thiazol-4-yl]-[(2S,3S)-3-methyl-2-(5-trifluoromethyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- [5-(3,5-Difluoro-phenyl)-2-methyl-thiazol-4-yl]-[(2S,3S)-3-methyl-2-(5-trifluoromethyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- 20 [5-(4-Fluoro-phenyl)-2-methyl-thiazol-4-yl]-[(2S,3S)-3-methyl-2-(4-trifluoromethyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- [5-(4-Chloro-phenyl)-2-methyl-thiazol-4-yl]-[(2S,3S)-3-methyl-2-(4-trifluoromethyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- 25 [5-(4-Bromo-phenyl)-2-methyl-thiazol-4-yl]-[(2S,3S)-3-methyl-2-(4-trifluoromethyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- (2-Methyl-5-p-tolyl-thiazol-4-yl)-[(2S,3S)-3-methyl-2-(4-trifluoromethyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- [5-(3,4-Difluoro-phenyl)-2-methyl-thiazol-4-yl]-[(2S,3S)-3-methyl-2-(4-trifluoromethyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- 30 [5-(2-Methoxy-phenyl)-2-methyl-thiazol-4-yl]-[(2S,3S)-3-methyl-2-(4-trifluoromethyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- [5-(3,5-Difluoro-phenyl)-2-methyl-thiazol-4-yl]-[(2S,3S)-3-methyl-2-(4-trifluoromethyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone;

- (2-Methyl-5-m-tolyl-thiazol-4-yl)-[(2S,3S)-3-methyl-2-(4-trifluoromethyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- [5-(4-Methoxy-phenyl)-2-methyl-thiazol-4-yl]-[(2S,3S)-3-methyl-2-(4-trifluoromethyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- 5 [(2S,3S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(5-methyl-2-pyridin-2-yl-phenyl)-methanone;
- (4-Chloro-biphenyl-2-yl)-[(2S,3S)-2-(5-chloro-4-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-methanone;
- 10 [(2S,3S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(5-methyl-2-pyridin-3-yl-phenyl)-methanone;
- [(2S,3S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(4-fluoro-biphenyl-2-yl)-methanone;
- [(2S,3S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(5-fluoro-2-pyridin-2-yl-phenyl)-methanone;
- 15 2-[(2S,3S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidine-1-carbonyl]-biphenyl-4-carbonitrile;
- [(2S,3S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(4-methyl-2-pyridin-2-yl-phenyl)-methanone;
- [(2S,3S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(5-methyl-biphenyl-2-yl)-methanone;
- 20 [(2S,3S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(3-fluoro-biphenyl-2-yl)-methanone;
- [(2S,3S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(4-trifluoromethyl-biphenyl-2-yl)-methanone;
- [(2S,3S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(3-fluoro-4-methyl-biphenyl-2-yl)-methanone;
- 25 [(2S,3S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(4-methyl-biphenyl-2-yl)-methanone;
- [(2S,3S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(2-fluoro-3-methyl-6-pyridin-3-yl-phenyl)-methanone;
- 30 [(2S,3S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(4-methyl-2-pyrimidin-2-yl-phenyl)-methanone;
- [(2S,3S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(5-fluoro-2-pyrimidin-2-yl-phenyl)-methanone;
- [(2S,3S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(2-fluoro-3-methyl-6-pyrimidin-2-yl-phenyl)-methanone;
- 35

- [(2S,3S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(2-fluoro-6-pyrimidin-2-yl-phenyl)-methanone;
- [(2S,3S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(5-methyl-2-pyrimidin-2-yl-phenyl)-methanone;
- 5 [(2S,3S)-2-(5-Chloro-6-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(5-methyl-2-pyrimidin-2-yl-phenyl)-methanone;
- [(2S,3S)-2-(6-Chloro-7-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(5-methyl-2-pyrimidin-2-yl-phenyl)-methanone;
- [(2S,3S)-2-(5-Chloro-7-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(5-methyl-2-pyrimidin-2-yl-phenyl)-methanone;
- 10 [(S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-[5-(4-fluoro-phenyl)-2-phenyl-thiazol-4-yl]-methanone;
- [2,5-Bis-(4-fluoro-phenyl)-thiazol-4-yl]-[(S)-2-(5-chloro-4-methyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-methanone;
- 15 [(S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-[5-(4-fluoro-phenyl)-2-m-tolyl-thiazol-4-yl]-methanone;
- [(S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-[5-(4-fluoro-phenyl)-2-(3-methoxy-phenyl)-thiazol-4-yl]-methanone;
- [(S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-[5-(4-fluoro-phenyl)-2-p-tolyl-thiazol-4-yl]-methanone;
- 20 [(S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-[5-(4-fluoro-phenyl)-2-(2-fluoro-phenyl)-thiazol-4-yl]-methanone;
- [(S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-[5-(4-fluoro-phenyl)-2-(3-fluoro-phenyl)-thiazol-4-yl]-methanone;
- 25 (4-Methyl-biphenyl-2-yl)-[(S)-2-methyl-2-(5-trifluoromethoxy-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- [5-(3,5-Difluoro-phenyl)-2-methyl-thiazol-4-yl]-[(S)-2-methyl-2-(5-trifluoromethoxy-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- [5-(3-Methoxy-phenyl)-2-methyl-thiazol-4-yl]-[(S)-2-methyl-2-(5-trifluoromethoxy-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- 30 [5-(4-Fluoro-phenyl)-2-methyl-thiazol-4-yl]-[(S)-2-methyl-2-(5-trifluoromethoxy-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- [(S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-[2-(2,3-difluoro-phenyl)-5-(4-fluoro-phenyl)-thiazol-4-yl]-methanone;
- [(S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-[2-(4-fluoro-3-methyl-phenyl)-5-(4-fluoro-phenyl)-thiazol-4-yl]-methanone;
- 35

- [(S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-[2-(2-fluoro-4-methyl-phenyl)-5-(4-fluoro-phenyl)-thiazol-4-yl]-methanone;
 3-[4-[(S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidine-1-carbonyl]-5-(4-fluoro-phenyl)-thiazol-2-yl]-benzonitrile;
- 5 [(S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-[2-(2-chloro-phenyl)-5-(4-fluoro-phenyl)-thiazol-4-yl]-methanone;
 [(S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-[2-(3-chloro-phenyl)-5-(4-fluoro-phenyl)-thiazol-4-yl]-methanone;
 [(S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-[2-(4-fluoro-2-methyl-phenyl)-5-(4-fluoro-phenyl)-thiazol-4-yl]-methanone;
- 10 [(S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-[2-(4-fluoro-2-methoxy-phenyl)-5-(4-fluoro-phenyl)-thiazol-4-yl]-methanone;
 [(S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-[2-(2,5-difluoro-phenyl)-5-(4-fluoro-phenyl)-thiazol-4-yl]-methanone;
- 15 [(S)-2-(6-Bromo-4,5-dimethyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(3,4-dimethyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
 3'-[(S)-2-Methyl-2-(5-trifluoromethyl-1H-benzoimidazol-2-yl)-pyrrolidine-1-carbonyl]-4'-[1,2,3]triazol-2-yl-biphenyl-3-carbonitrile;
 2'-[(S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidine-1-carbonyl]-3'-fluoro-4'-methoxy-
- 20 biphenyl-4-carbonitrile;
 [(S)-2-(6-Bromo-5-fluoro-1-methyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
 (3'-Fluoro-4-methyl-biphenyl-2-yl)-{(S)-2-[1-(2-methoxy-ethyl)-5,6-dimethyl-1H-benzoimidazol-2-yl]-2-methyl-pyrrolidin-1-yl}-methanone; and
- 25 [(2S,3S)-2-(5-Chloro-6-methyl-1H-benzoimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(2-fluoro-3-methyl-6-pyrimidin-2-yl-phenyl)-methanone.
- 51) In addition to the above-listed compounds, further compounds according to embodiment 1) are selected from:
- (4-Bromo-biphenyl-2-yl)-[(S)-2-methyl-2-(5-trifluoromethyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- 30 [(S)-2-(5,6-Dimethyl-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(2-methyl-6-[1,2,3]triazol-2-yl-phenyl)-methanone;
 [(S)-2-(5,6-Dimethyl-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(5-methyl-2-pyrazol-1-yl-phenyl)-methanone; and
 [(S)-2-(5,6-Dimethyl-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(6-methyl-3-[1,2,3]triazol-2-yl-pyridin-2-yl)-methanone.
- 35

52) In another embodiment, preferred compounds according to embodiment 1) are selected from:

[(2S,3S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(2-fluoro-6-pyrimidin-2-yl-phenyl)-methanone;

5 [(2S,3S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(2-fluoro-3-methyl-6-pyrimidin-2-yl-phenyl)-methanone;

[(2S,3S)-2-(5-Chloro-6-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(2-fluoro-3-methyl-6-pyrimidin-2-yl-phenyl)-methanone;

10 [(2S,3S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(5-chloro-2-[1,2,3]triazol-2-yl-phenyl)-methanone;

[(2S,3S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(2-fluoro-3-methyl-6-[1,2,3]triazol-2-yl-phenyl)-methanone;

3'-[(S)-2-Methyl-2-(5-trifluoromethyl-1H-benzimidazol-2-yl)-pyrrolidine-1-carbonyl]-4'-[1,2,3]triazol-2-yl-biphenyl-3-carbonitrile; and

15 [(S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-[5-(4-fluoro-phenyl)-2-phenyl-thiazol-4-yl]-methanone.

53) In another embodiment, preferred compounds according to embodiment 1) are selected from::

20 (5-Methyl-2-[1,2,3]triazol-2-yl-phenyl)-[(S)-2-methyl-2-(5-trifluoromethyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;

(5-Methyl-2-[1,2,3]triazol-2-yl-phenyl)-[(S)-2-methyl-2-(5-trifluoromethoxy-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;

[(S)-2-(5,6-Dimethoxy-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;

25 [(S)-2-(6-Bromo-5-fluoro-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;

[(S)-2-(6-Chloro-5-trifluoromethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;

30 [(S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-chloro-2-[1,2,3]triazol-2-yl-phenyl)-methanone;

[(S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(2-fluoro-3-methoxy-6-[1,2,3]triazol-2-yl-phenyl)-methanone;

[(S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(4-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;

[(S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-pyrazol-1-yl-phenyl)-methanone;

[(S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(6-methyl-3-pyrazol-1-yl-pyridin-2-yl)-methanone;

5 [(S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-fluoro-2-[1,2,3]triazol-2-yl-phenyl)-methanone;

[(S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(2-[1,2,3]triazol-2-yl-phenyl)-methanone;

10 (5-Chloro-2-[1,2,3]triazol-2-yl-phenyl)-[(S)-2-(5,6-dimethoxy-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-methanone;

[(S)-2-(5,6-Dimethoxy-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methoxy-2-[1,2,3]triazol-2-yl-phenyl)-methanone;

[(S)-2-(5,6-Dimethoxy-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-pyrazol-1-yl-phenyl)-methanone;

15 [(S)-2-(6-Chloro-5-trifluoromethyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(4,5-dimethyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone; and

(4,5-Dimethyl-2-[1,2,3]triazol-2-yl-phenyl)-[(S)-2-methyl-2-(5-trifluoromethyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone.

54) In another embodiment, further preferred compounds according to embodiment 1) are:

20 [(S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(6-methyl-3-[1,2,3]triazol-2-yl-pyridin-2-yl)-methanone; and

[(S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-pyrimidin-2-yl-phenyl)-methanone.

55) A particularly preferred compound according to embodiment 1) is:

25 [(S)-2-(5-Bromo-7-methyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone.

56) Another particularly preferred compound according to embodiment 1) is:

[(S)-2-(5-Bromo-7-methyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-pyrimidin-2-yl-phenyl)-methanone.

30 57) Another particularly preferred compound according to embodiment 1) is:

[(S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methoxy-2-[1,2,3]triazol-2-yl-phenyl)-methanone.

The compounds of formulae (I), (II), (III), (IV), (V), and (VI) according to any one of embodiments 1) to 57) and their pharmaceutically acceptable salts can be used as

medicaments, e.g. in the form of pharmaceutical compositions for enteral (such especially oral) or parenteral administration (including topical application or inhalation).

The production of the pharmaceutical compositions can be effected in a manner which will be familiar to any person skilled in the art (see for example Remington, *The Science and Practice of Pharmacy*, 21st Edition (2005), Part 5, "Pharmaceutical Manufacturing" [published by Lippincott Williams & Wilkins]) by bringing the described compounds of formulae (I), (II) and (III) or their pharmaceutically acceptable salts, optionally in combination with other therapeutically valuable substances, into a galenical administration form together with suitable, non-toxic, inert, therapeutically compatible solid or liquid carrier materials and, if desired, usual pharmaceutical adjuvants.

The present invention also relates to a method for the prevention or treatment of a disease or disorder mentioned herein comprising administering to a subject a pharmaceutically active amount of a compound of formulae (I), (II), (III), (IV), (V), and (VI) according to any one of embodiments 1) to 57).

In a preferred embodiment of the invention, the administered amount of such a compound of formulae (I), (II), (III), (IV), (V), and (VI) according to any one of embodiments 1) to 57) is comprised between 1 mg and 1000 mg per day, particularly between 5 mg and 500 mg per day, more particularly between 25 mg and 400 mg per day, especially between 50 mg and 200 mg per day.

For avoidance of any doubt, if compounds are described as useful for the prevention or treatment of certain diseases, such compounds are likewise suitable for use in the preparation of a medicament for the prevention or treatment of said diseases.

The compounds according to formulae (I), (II), (III), (IV), (V), and (VI) according to any one of embodiments 1) to 57) are useful for the prevention or treatment of disorders relating to orexinergic dysfunctions.

Such disorders relating to orexinergic dysfunctions are diseases or disorders where an antagonist of a human orexin receptor is required, notably mental health disorders relating to orexinergic dysfunctions. The above mentioned disorders may in particular be defined as comprising sleep disorders, anxiety disorders, addiction disorders, cognitive dysfunctions, mood disorders, or appetite disorders. In one sub-embodiment, the above mentioned disorders comprise especially anxiety disorders, addiction disorders and mood disorders, notably anxiety disorders and addiction disorders. In another sub-embodiment, the above mentioned disorders comprise especially sleep disorders.

In addition, further disorders relating to orexinergic dysfunctions are selected from treating, controlling, ameliorating or reducing the risk of epilepsy, including absence epilepsy; treating or controlling pain, including neuropathic pain; treating or controlling Parkinson's disease; treating or controlling psychosis including acute mania and bipolar disorder; treating or
5 controlling stroke, particularly ischemic or haemorrhagic stroke; blocking an emetic response i.e. nausea and vomiting; and treating or controlling agitation, in isolation or co-morbid with another medical condition.

Anxiety disorders can be distinguished by the primary object or specificity of threat, ranging from rather diffuse as in generalized anxiety disorder, to circumscribed as encountered in
10 phobic anxieties (PHOBs) or post-traumatic stress disorders (PTSDs). Anxiety disorders may, thus, be defined as comprising generalized anxiety disorders (GAD), obsessive compulsive disorders (OCDs), acute stress disorders, posttraumatic stress disorders (PTSDs), panic anxiety disorders (PADs) including panic attacks, phobic anxieties (PHOBs), specific phobia, social phobia (social anxiety disorder), avoidance, somatoform disorders
15 including hypochondriasis, separation anxiety disorder, anxiety disorders due to a general medical condition, and substance induced anxiety disorders. In a sub-embodiment, particular examples of circumscribed threat induced anxiety disorders are phobic anxieties or post-traumatic stress disorders. Anxiety disorders especially include post-traumatic stress disorders, obsessive compulsive disorders, panic attacks, phobic anxieties, and avoidance.

Addiction disorders may be defined as addictions to one or more rewarding stimuli, notably to one rewarding stimulus. Such rewarding stimuli may be of either natural or synthetic origin. Examples of such rewarding stimuli are substances / drugs {of either natural or synthetic origin; such as cocaine, amphetamines, opiates [of natural or (semi-)synthetic origin such as morphine or heroin], cannabis, ethanol, mescaline, nicotine, and the like}, which substances /
25 drugs may be consumed alone or in combination; or other rewarding stimuli {of either natural origin (such as food, sweet, fat, or sex, and the like), or synthetic origin [such as gambling, or internet/IT (such as immoderate gaming, or inappropriate involvement in online social networking sites or blogging), and the like]}. In a sub-embodiment, addiction disorders relating to psychoactive substance use, abuse, seeking and reinstatement are defined as all
30 types of psychological or physical addictions and their related tolerance and dependence components. Substance-related addiction disorders especially include substance use disorders such as substance dependence, substance craving and substance abuse; substance-induced disorders such as substance intoxication, substance withdrawal, and substance-induced delirium. The expression "prevention or treatment of addictions" (i.e.
35 preventive or curative treatment of patients who have been diagnosed as having an

addiction, or as being at risk of developing addictions) refers to diminishing addictions, notably diminishing the onset of addictions, to weakening their maintenance, to facilitating withdrawal, to facilitating abstinence, or to attenuating, decreasing or preventing the occurrence of reinstatement of addiction (especially to diminishing the onset of addictions, to
5 facilitating withdrawal, or to attenuating, decreasing or preventing the occurrence of reinstatement of addiction).

Mood disorders include major depressive episode, manic episode, mixed episode and hypomanic episode; depressive disorders including major depressive disorder, dysthymic disorders; bipolar disorders including bipolar I disorder, bipolar II disorder (recurrent major
10 depressive episodes with hypomanic episodes), cyclothymic disorder; mood disorders including mood disorder due to a general medical condition (including the subtypes with depressive features, with major depressive-like episode, with manic features, and with mixed features), substance-induced mood disorder (including the subtypes with depressive features, with manic features, and with mixed features). Such mood disorders are especially
15 major depressive episode, major depressive disorder, mood disorder due to a general medical condition; and substance-induced mood disorder.

Appetite disorders comprise eating disorders and drinking disorders. Eating disorders may be defined as comprising eating disorders associated with excessive food intake and complications associated therewith; anorexias; compulsive eating disorders; obesity (due to
20 any cause, whether genetic or environmental); obesity-related disorders including overeating and obesity observed in Type 2 (non-insulin-dependent) diabetes patients; bulimias including bulimia nervosa; cachexia; and binge eating disorder. Particular eating disorders comprise metabolic dysfunction; dysregulated appetite control; compulsive obesities; bulimia or anorexia nervosa. In a sub-embodiment, eating disorders may be defined as especially
25 comprising anorexia nervosa, bulimia, cachexia, binge eating disorder, or compulsive obesities. Drinking disorders include polydipsias in psychiatric disorders and all other types of excessive fluid intake. Pathologically modified food intake may result from disturbed appetite (attraction or aversion for food); altered energy balance (intake vs. expenditure); disturbed perception of food quality (high fat or carbohydrates, high palatability); disturbed
30 food availability (unrestricted diet or deprivation) or disrupted water balance.

Cognitive dysfunctions include deficits in attention, learning and especially memory functions occurring transiently or chronically in psychiatric, neurologic, neurodegenerative, cardiovascular and immune disorders, and also occurring transiently or chronically in the normal, healthy, young, adult, or especially aging population. Cognitive dysfunctions

especially relate to the enhancement or maintenance of memory in patients who have been diagnosed as having, or being at risk of developing, diseases or disorders in which diminished memory (notably declarative or procedural) is a symptom [in particular dementias such as frontotemporal dementia, or dementia with Lewy bodies, or (especially) Alzheimer's disease]. Especially, the term "prevention or treatment of cognitive dysfunctions" relates to the enhancement or maintenance of memory in patients who have a clinical manifestation of a cognitive dysfunction, especially expressed as a deficit of declarative memory, linked to dementias such as frontotemporal dementia, or dementia with Lewy bodies, or (especially) Alzheimer's disease. Furthermore, the term "prevention or treatment of cognitive dysfunctions" also relates to improving memory consolidation in any of the above mentioned patient populations.

Sleep disorders comprise dyssomnias, parasomnias, sleep disorders associated with a general medical condition and substance-induced sleep disorders. In particular, dyssomnias include intrinsic sleep disorders (especially insomnias, breathing-related sleep disorders, periodic limb movement disorder, and restless leg syndrome), extrinsic sleep disorders, and circadian-rhythm sleep disorders. Dyssomnias notably include insomnia, primary insomnia, idiopathic insomnia, insomnias associated with depression, emotional/mood disorders, aging, Alzheimer's disease or cognitive impairment; REM sleep interruptions; breathing-related sleep disorders; sleep apnea; periodic limb movement disorder (nocturnal myoclonus), restless leg syndrome, circadian rhythm sleep disorder; shift work sleep disorder; and jet-lag syndrome. Parasomnias include arousal disorders and sleep-wake transition disorders; notably parasomnias include nightmare disorder, sleep terror disorder, and sleepwalking disorder. Sleep disorders associated with a general medical condition are in particular sleep disorders associated with diseases such as mental disorders, neurological disorders, neuropathic pain, and heart and lung diseases. Substance-induced sleep disorders include especially the subtypes insomnia type, parasomnia type and mixed type, and notably include conditions due to drugs which cause reductions in REM sleep as a side effect. Sleep disorders especially include all types of insomnias, sleep-related dystonias; restless leg syndrome; sleep apneas; jet-lag syndrome; shift work sleep disorder, delayed or advanced sleep phase syndrome, or insomnias related to psychiatric disorders. In addition, sleep disorders further include sleep disorders associated with aging; intermittent treatment of chronic insomnia; situational transient insomnia (new environment, noise) or short-term insomnia due to stress; grief; pain or illness.

In the context of the present invention, it is to be understood that, in case certain environmental conditions such as stress or fear (wherein stress may be of social origin (e.g.

social stress) or of physical origin (e.g. physical stress), including stress caused by fear) facilitate or precipitate any of the disorders or diseases as defined before, the present compounds may be particularly useful for the treatment of such environmentally conditioned disorder or disease.

5 **Preparation of compounds of formula (I):**

The compounds of formulae (I), (II), (III), (IV), (V), and (VI) can be prepared by the methods given below, by the methods given in the experimental part below or by analogous methods. Optimum reaction conditions may vary with the particular reactants or solvents used, but such conditions can be determined by a person skilled in the art by routine optimisation procedures. In some cases the final product may be further modified, for example, by manipulation of substituents to give a new final product. These manipulations may include, but are not limited to, reduction, oxidation, alkylation, acylation, and hydrolysis reactions which are commonly known to those skilled in the art. In some cases the order of carrying out the following reaction schemes, and/or reaction steps, may be varied to facilitate the reaction or to avoid unwanted reaction products.


Compounds of formulae (I), (II), (III), (IV), (V), and (VI) of the present invention can be prepared according to the general sequence of reactions outlined below wherein Ar^1 , R^1 , R^2 , R^3 , R^{4a} , R^{4b} and $(\text{R}^5)_n$ are as defined for formula (I).

The synthesis of compound of formulae (I), wherein R^1 represents hydrogen, starts from proline-derivatives (**a**) which are commercially available or prepared as described below.

There are two general synthetic approaches of equal importance towards compounds of formulae (I).

Synthetic approach 1 starts with a Boc-protection of the respective proline derivative **a** under standard conditions by for example dissolving the proline **a** in a solvent such as DCM or THF and adding a base to the solution, for example DIPEA, TEA or aqueous Na_2CO_3 followed by the addition of Boc_2O . The reaction is performed at room temperature and is usually complete within a few hours and results in the Boc-protected proline derivative **b** which is then coupled with the appropriate phenylene-diamine- or pyridine-diamine derivative in solvents such as THF, DCM or DMF in the presence of a coupling agent such as HBTU or TBTU or the like and a base, for example DIPEA or TEA to give compound **c**. To obtain the benzimidazole derivative **d** the precursor **c** is dissolved in AcOH and heated to 100°C for 1 h. Compound **d** is Boc-deprotected under acidic conditions such as 4M HCl in dioxane (preferred method) or TFA in DCM to give precursor **e** which is converted into final compound **f** by an amide coupling reaction with $\text{Ar}^1\text{-COOH}$ in a solvent such as THF, DMF or


DCM in the presence of a coupling agent such as TBTU, HBTU, HATU, EDC or the like and a base such as DIPEA, TEA or N-methylmorpholine.



Synthetic approach 2 starts with an esterification (usually methyl ester formation) of the proline derivative **a** by dissolving the starting material in THF and adding 5 equivalents of the respective alcohol (usually MeOH) followed by the addition of EDC and DMAP. The reaction is run at RT and is usually complete within a few hours. The methyl-ester derivative **g** is acylated with $\text{Ar}^1\text{-COOH}$ under conditions described above to result in intermediate **h**. Esterhydrolysis under standard conditions by dissolving the ester derivative **h** in THF / MeOH = 1 / 1 followed by the addition of 2 equivalents of aq. 1M NaOH solution. The reaction runs


at RT and is usually complete after a few hours to result in the carboxylic acid derivative **i**. The final compounds **f** are obtained via precursor **j** by applying the same conditions as described for the amide-coupling and the cyclization in synthetic approach 1.

The following scheme depicts the methodology to prepare N-methylated benzimidazolo-derivatives of formula (I), wherein R¹ represents methyl:




Final compounds **g1** and **g2** are prepared by simple alkylation of compounds **f** (compounds of formula (I) wherein R¹ is H) with an alkyl halide (for example MeI) in the presence of a carbonate base (e.g. Cs₂CO₃) in a polar, aprotic solvent (e.g. DMF) at RT. In case the phenylring of the benzimidazole-system is unsymmetrically substituted with respect to R¹⁴, R¹⁵, R¹⁶, and R¹⁷ the alkylation (methylation) reaction results in two isomeric products **g1** and **g2** as depicted above.

The following proline derivatives are commercially available and were purchased and used as starting materials according to the methods described herein:



The following proline derivatives may be prepared according to the procedures outlined in the following scheme:



- 5 The diastereomeric 4-hydroxy-proline derivatives **k** and **n** are commercially available starting points. Both compounds may be fully deprotonated with NaH in DMF followed by the addition of an excess of MeI to give the 4-methoxy-proline-methylester derivatives **l** and **o** which can be transformed into the carboxylic acid derivatives **m** and **p** by standard ester hydrolysis (e.g. 1M NaOH, MeOH, THF, rt).
- 10 Whenever the compounds of formulae (I), (II), (III), (IV), (V), or (VI) are obtained in the form of mixtures of stereoisomers such as especially enantiomers, the stereoisomers can be separated using methods known to one skilled in the art: e.g. by formation and separation of diastereomeric salts or by HPLC over a chiral stationary phase such as a Daicel ChiralPak AD-H (5 μ m) column, a Daicel ChiralCel OD-H (5 μ m) column, a Daicel ChiralCel OD (10

μm) column, a Daicel ChiralPak IA (5 μm) column, a Daicel ChiralPak IB (5 μm) column, a Daicel ChiralPak IC (5 μm) column, or a (R,R)-Whelk-01 (5 μm) column. Typical conditions of chiral HPLC are an isocratic mixture of eluent A (EtOH, in presence or absence of a base like TEA and/or diethylamine or of an acid like TFA) and eluent B (heptane).

- 5 The following examples are provided to illustrate the invention. These examples are illustrative only and should not be construed as limiting the invention in any way.

Experimental Part

I. Chemistry

- 10 All temperatures are stated in °C. Commercially available starting materials were used as received without further purification. Unless otherwise specified, all reactions were carried out in oven-dried glassware under an atmosphere of nitrogen. Compounds were purified by flash column chromatography on silica gel or by preparative HPLC. Compounds described in the invention are characterised by LC-MS data (retention time t_R is given in min; molecular weight obtained from the mass spectrum is given in g/mol) using the conditions listed below.
- 15 In cases where compounds of the present invention appear as a mixture of conformational isomers, particularly visible in their LC-MS spectra, the retention time of the most abundant conformer is given. Racemates can be separated into their enantiomers by preparative HPLC (column: ChiralPaK IC 250x4.6 mm, 5 μm, 45% ethanol in heptane).

LC-MS with acidic conditions

- 20 **Method A:** Agilent 1100 series with mass spectrometry detection (MS: Finnigan single quadrupole). Column: Zorbax SB-aq (3.5 μm, 4.6 x 50 mm). Conditions: MeCN [eluent A]; water + 0.04% TFA [eluent B]. Gradient: 95% B → 5% B over 1.5 min (flow: 4.5 mL/min). Detection: UV/Vis + MS.

- 25 **Method B:** Agilent 1100 series with mass spectrometry detection (MS: Finnigan single quadrupole). Column: Waters XBridge C18 (2.5 μm, 4.6 x 30 mm). Conditions: MeCN [eluent A]; water + 0.04% TFA [eluent B]. Gradient: 95% B → 5% B over 1.5 min (flow: 4.5 mL/min). Detection: UV/Vis + MS.

LC-MS with basic conditions

- 30 **Method C:** Agilent 1100 series with mass spectrometry detection (MS: Finnigan single quadrupole). Column: Zorbax Extend C18 (5 μm, 4.6 x 50 mm). Conditions: MeCN [eluent A]; 13 mmol/L NH₃ in water [eluent B]. Gradient: 95% B → 5% B over 1.5 min (flow: 4.5 mL/min). Detection: UV/Vis + MS.

Method D: Agilent 1100 series with mass spectrometry detection (MS: Finnigan single quadrupole). Column: Waters XBridge C18 (5 μ m, 4.6 x 50 mm). Conditions: MeCN [eluent A]; 13 mmol/L NH₃ in water [eluent B]. Gradient: 95% B $\rightarrow$ 5% B over 1.5 min (flow: 4.5 mL/min). Detection: UV/Vis + MS.

5 Preparative HPLC with acidic conditions

Method E: Column: Waters XBridge (10 μ m, 75 x 30 mm). Conditions: MeCN [eluent A]; water + 0.5% HCOOH [eluent B]; Gradient: 90% B $\rightarrow$ 5% B over 6.4 min (flow: 75 mL/min). Detection: UV/Vis + MS.

Preparative HPLC with basic conditions

10 **Method F:** Column: Waters XBridge (10 μ m, 75 x 30 mm). Conditions: MeCN [eluent A]; water + 0.5% NH₄OH (25% aq.) [eluent B]; Gradient: 90% B $\rightarrow$ 5% B over 6.5 min (flow: 75 mL/min). Detection: UV/Vis + MS

Abbreviations (as used hereinbefore or hereinafter):

Abbreviations (as used herein and in the description above):

15	Ac	Acetyl (such as in OAc = acetate, AcOH = acetic acid)
	AcOH	Acetic acid
	anh.	Anhydrous
	aq.	aqueous
	atm	Atmosphere
20	tBME	tert-Butylmethylether
	Boc	<i>tert</i> -Butoxycarbonyl
	Boc ₂ O	di- <i>tert</i> -Butyl dicarbonate
	BSA	Bovine serum albumine
	Bu	Butyl such as in tBu = <i>tert</i> -butyl = tertiary butyl
25	CC	Column Chromatography on silica gel
	CHO	Chinese Hamster Ovary
	conc.	Concentrated
	DCE	1,2-Dichloroethane
	DCM	Dichloromethane
30	DEA	Diethylamine
	DIPEA	Diisopropylethylamine
	DMF	<i>N,N</i> -Dimethylformamide
	DMSO	Dimethyl sulfoxide

	EDC	
	ELSD	Evaporative Light-Scattering Detection
	eq	Equivalent(s)
	ES	Electron spray
5	Et	Ethyl
	Et ₂ O	Diethyl ether
	EtOAc	Ethyl acetate
	EtOH	Ethanol
	Ex.	Example
10	FC	Flash Chromatography on silica gel
	FCS	Foatal calf serum
	FLIPR	Fluorescent imaging plate reader
	h	Hour(s)
	HATU	
15	HBSS	Hank's balanced salt solution
	HBTU	
	HEPES	4-(2-Hydroxyethyl)-piperazine-1-ethanesulfonic acid
	¹ H-NMR	Nuclear magnetic resonance of the proton
	HPLC	High performance liquid chromatography
20	LC-MS	Liquid chromatography – Mass Spectroscopy
	Lit.	Literature
	M	Exact mass (as used for LC-MS)
	Me	Methyl
	MeCN	Acetonitrile
25	MeOH	Methanol
	Mel	Methyl iodide
	MHz	Megahertz
	μl	microliter
	min	Minute(s)
30	MS	Mass spectroscopy
	N	Normality
	Pd(OAc) ₂	Palladium diacetate
	Pd(PPh ₃) ₄	Tetrakis(triphenylphosphine)palladium(0)
	PL-HCO ₃	Polymer supported hydrogen carbonate
35	Ph	Phenyl

	PPh ₃	Triphenylphosphine
	prep.	Preparative
	RT	Room temperature
	sat.	Saturated
5	TBTU	O-(Benzotriazol-1-yl)-N,N,N',N'-tetramethyluronium tetrafluoroborate
	TEA	Triethylamine
	TFA	trifluoroacetic acid
	Tf	Trifluoromethansulfonyl
	THF	Tetrahydrofuran
10	t _R	Retention time
	UV	Ultra violet

I-Chemistry

All temperatures are stated in °C. The commercially available starting materials were used as received without further purification. Compounds are purified by flash column chromatography on silica gel (FC) or by preparative HPLC. Compounds described in the invention are characterized by LC-MS (retention time t_R is given in min.; molecular weight obtained from the mass spectrum is given in g/mol, using the conditions listed below). If the mass is not detectable the compounds are also characterized by ¹H-NMR (400 MHz: Bruker; chemical shifts are given in ppm relative to the solvent used; multiplicities: s = singlet, d = doublet, t = triplet; p = pentuplet, hex = hexet, hept = heptet, m = multiplet, br = broad, coupling constants are given in Hz).

LC-MS with acidic conditions (conditions A)

Apparatus: Agilent 1100 series with mass spectroscopy detection (MS : Finnigan single quadrupole). Column: Waters XBridge C18 (2.5 μm, 4.6 x 30 mm). Conditions: MeCN [eluent A]; water + 0.04% TFA [eluent B]. Gradient: 95% B → 5% B over 1.5 min. (flow: 4.5 ml/min.). Detection: UV/Vis + MS.

LC-MS with basic conditions (conditions B)

Apparatus: Agilent 1100 series with mass spectroscopy detection (MS : Finnigan single quadrupole). Column: Waters XBridge C18 (5 μm, 4.6 x 50 mm). Conditions: MeCN [eluent A]; 13 mmol/l NH₃ in water [eluent B]. Gradient: 95% B → 5% B over 1.5 min. (flow: 4.5 ml/min.). Detection: UV/Vis + MS.

Preparative HPLC for purification of compounds (conditions C)

Column: Waters XBridge (10 μ m, 75 x 30 mm). Conditions: MeCN [eluent A]; water + 0.5% NH₄OH (25% aq.) [eluent B]; Gradient: 90% B $\rightarrow$ 5% B over 6.5 min. (flow: 75 ml/min.).

Detection: UV + ELSD.

5 Preparative HPLC for purification of compounds (conditions D)

Column: Waters Atlantis T3 OBD (10 μ m, 75 x 30 mm). Conditions: MeCN [eluent A]; water + 0.5% HCOOH [eluent B]; Gradient: 90% B $\rightarrow$ 5% B over 6.4 min. (flow: 75 ml/min.).

Detection: UV + ELSD.

LC-MS with basic conditions (conditions E)

- 10 Apparatus: Agilent 1100 series with mass spectroscopy detection (MS : Finnigan single quadrupole). Column: Agilent Zorbax Extend-C18 (5 μ m, 4.6 x 50 mm). Conditions: MeCN [eluent A]; 13 mmol/l NH₃ in water [eluent B]. Gradient: 95% B $\rightarrow$ 5% B over 1.5 min. (flow: 4.5 ml/min.). Detection: UV + MS.

LC-MS with acidic conditions (conditions F)


- 15 Apparatus: Agilent 1100 series with mass spectroscopy detection (MS : Finnigan single quadrupole). Column: Agilent Zorbax SB-Aq, (3.5 μ m, 4.6 x 50mm). Conditions: MeCN [eluent A]; water + 0.04% TFA [eluent B]. Gradient: 95% B $\rightarrow$ 5% B over 1.5 min. (flow: 4.5 ml/min.). Detection: UV + MS.

- 20 The following examples illustrate the preparation of compounds of the invention but do not at all limit the scope thereof.

Preparation of precursors and intermediates:***A Preparation of building blocks of formula Ar¹-CO-OH:***

- 25 The preparation of these acids is described in detail for example in the following documents: WO 2008/020405; WO 2008/038251; WO 2008/081399; WO 2008/139416. All other carboxylic acids used in the experimental part which are not described in the following section are either commercially available or fully described in the literature listed above and in the introduction part.

In addition to commercially available building blocks, further particular building blocks of formula Ar¹-CO-OH are prepared as follows:



5 A.1 2-Fluoro-3-methyl-6-(2H-1,2,3-triazol-2-yl)benzoic acid 39

2-Fluoro-3-methyl-6-(2H-1,2,3-triazol-2-yl)benzoic acid **39** is synthesized in analogy to procedures reported in WO2008/069997.

In a dry Schlenk Tube at RT under nitrogen are successively charged 2-fluoro-6-iodo-3-methyl-benzoic acid (1.786 mmol, 1 eq), CuI (0.089 mmol, 0.05 eq), 1H-1,2,3-triazole (3.571 mmol, 2 eq), Cs₂CO₃ (3.571 mmol, 2 eq) and DMF (2.5 mL). The resulting blue suspension is stirred at 80°C overnight. The obtained reaction mixture is taken up in 1 M aq. HCl and extracted twice with EtOAc. The combined organic layers are dried over Na₂SO₄, filtered and concentrated under reduced pressure. Purification is achieved by preparative HPLC (conditions D) to give the titled compound (246 mg) as a pale yellow solid. LC-MS (conditions A): t_R = 0.55 min, [M + 1]⁺ = 222.19.

A.2 5-Methoxy-2-(2H-1,2,3-triazol-2-yl)benzoic acid 40

The title compound is prepared in analogy to compound **39** starting from 2-iodo-5-methoxybenzoic acid (1.798 mmol, 1 eq). **40** (313 mg) is obtained as a yellow solid. LC-MS (conditions A): t_R = 0.49 min, [M + 1]⁺ = 220.07.

A.3 2-Fluoro-3-methoxy-6-(2H-1,2,3-triazol-2-yl)benzoic acid 41

The title compound is prepared in analogy to compound **39** starting from 2-fluoro-6-iodo-3-methoxy-benzoic acid (1.689 mmol, 1 eq). **41** (221 mg) is obtained as a pale yellow solid. LC-MS (conditions A): $t_R = 0.48$ min, $[M + 1]^+ = 238.18$.

5 A.4 2-(2H-1,2,3-Triazol-2-yl)-5-(trifluoromethyl)benzoic acid 42

The title compound is prepared in analogy to compound **39** starting from 2-iodo-5-trifluorobenzoic acid (1.582 mmol, 1 eq). **42** (268 mg, 66%) is obtained as a white solid. LC-MS (conditions A): $t_R = 0.64$ min, $[M + 1]^+ = 257.91$.

A.5 2-(2H-1,2,3-Triazol-2-yl)-5-(trifluoromethoxy)benzoic acid 43

- 10 The title compound is prepared in analogy to compound **39** starting from 2-iodo-5-(trifluoromethoxy)benzoic acid (1.506 mmol, 1 eq). **43** (243 mg) is obtained as an off-white solid. LC-MS (conditions A): $t_R = 0.66$ min, $[M + 1]^+ = 273.69$.

A.6 5-Cyano-2-(2H-1,2,3-triazol-2-yl)benzoic acid 44

- 15 The title compound is prepared in analogy to compound **39** starting from 5-cyano-2-iodobenzoic acid (1.831 mmol, 1 eq). **44** (214 mg) is obtained as a grey solid. LC-MS (conditions A): $t_R = 0.46$ min, $[M + 1]^+ =$ not detectable. ^1H NMR (D_6 -DMSO): δ 13.49 (m, 1 H), 8.21 (m, 1 H), 8.18 (m, 2 H), 8.15 (m, 1 H), 8.03 (m, 1 H).

A.7 5-Methyl-2-(pyridin-2-yl)benzoic acid 45

- 20 a) In a dry Schlenk Tube at RT under nitrogen are successively charged 2-iodo-5-methylbenzoic acid methyl ester (13.765 mmol, 1 eq), CuI (2.753 mmol, 0.2 eq), CsF (27.529 mmol, 2 eq), 2-tributylstannylpyridine (20.647 mmol, 1.5 eq), Pd(PPh₃)₄ (1.376 mmol, 0.1 eq) and DMF (60 mL). The resulting suspension is stirred at 90°C overnight. The obtained reaction mixture is diluted with EtOAc and filtered through a short pad of Celite®. A solution of sat. aq. NaHCO₃ is then added to the filtrate and the aq. phase extracted with EtOAc (3
- 25 times). The combined organic layers are washed with H₂O and brine, dried over Na₂SO₄, filtered and concentrated under reduced pressure. Purification is achieved by FC (EtOAc/Heptane 1:4 to 3:7) to give methyl 5-methyl-2-(pyridin-2-yl)benzoate (2.64 g) as a brown oil. LC-MS (conditions A): $t_R = 0.67$ min, $[M + 1]^+ = 228.07$.

- 30 b) To a solution of methyl 5-methyl-2-(pyridin-2-yl)benzoate (11.617 mmol, 1 eq) in MeOH (15 mL) and THF (17 mL) is added 1 M NaOH (23.233 mL, 2 eq). The resulting mixture is stirred at RT overnight. The volatiles are evaporated under reduced pressure and the remaining aq. phase is acidified with 2 M HCl to pH = 1-2 and extracted with DCM (3 times).

The combined organic layers are dried over Na₂SO₄, filtered and concentrated under reduced pressure to give 5-methyl-2-(pyridin-2-yl)benzoic acid **45** (2.65 g) as a pale brown foam. LC-MS (conditions A): $t_R = 0.39$ min, $[M + 1]^+ = 214.25$.

A.8 2-Fluoro-3-methyl-6-(1H-pyrazol-1-yl)benzoic acid 46

- 5 The title compound is prepared in analogy to compound **39** replacing the 1*H*-1,2,3-triazole with 1*H*-pyrazole (25 mmol, 2 eq). **46** (1.86 g) is obtained as a light yellow solid. LC-MS (conditions F): $t_R = 0.63$ min, $[M + 1]^+ = 221.16$.

A.9 2-Fluoro-3-methyl-6-(pyridin-2-yl)benzoic acid 47

- 10 a) In a dry Schlenk Tube at RT under nitrogen are successively charged methyl 2-fluoro-6-iodo-3-methylbenzoate (9.18 mmol, 1 eq), CuI (1.84 mmol, 0.2 eq), CsF (18.4 mmol, 2 eq), 2-tributylstannylpyridine (9.18 mmol, 1 eq), Pd(PPh₃)₄ (0.918 mmol, 0.1 eq) and DMF (40 mL). The resulting suspension is stirred at 90°C overnight. The obtained reaction mixture is diluted with EtOAc and filtered through a short pad of Celite®. A solution of sat. aq. NaHCO₃ is then added to the filtrate and the aq. phase extracted with EtOAc (3 times). The combined
15 organic layers are washed with H₂O and brine, dried over Na₂SO₄, filtered and concentrated under reduced pressure. Purification is achieved by FC (Teledyne Isco Combiflash Rf, SiO₂ cartridge 120 g; Heptane to EtOAc/Heptane 3:7) to give methyl 2-fluoro-3-methyl-6-(pyridin-2-yl)benzoate as a brown oil. LC-MS (conditions F): $t_R = 0.74$ min, $[M + 1]^+ = 246.15$.

- 20 b) To a solution of methyl 2-fluoro-3-methyl-6-(pyridin-2-yl)benzoate (6.36 mmol, 1 eq) in MeOH (11.3 mL) is added NaOH 32% (6.27 mL). The resulting mixture is stirred at 60°C for 1 hour. The volatiles are evaporated under reduced pressure and the remaining aq. phase is acidified with 7 M HCl to pH = 1-2. The pink suspension is concentrated under reduced pressure and the obtained solid is purified by preparative HPLC (conditions D) to give 2-fluoro-3-methyl-6-(pyridin-2-yl)benzoic acid **47** (1.14 g) as a light pink solid. LC-MS
25 (conditions F): $t_R = 0.47$ min, $[M + 1]^+ = 232.17$.

A.10 3-fluoro-4-methyl-[1,1'-biphenyl]-2-carboxylic acid 48


- 30 a) In a dry Schlenk Tube at RT under nitrogen are successively charged methyl 2-fluoro-6-iodo-3-methylbenzoate (23.5 mmol, 1 eq), Pd(PPh₃)₄ (1.17 mmol, 0.05 eq) and toluene (60 mL). The resulting mixture is stirred at RT for 15 minutes before a solution of phenylboronic acid (25.8 mmol, 1.1 eq) in EtOH (26 mL) and 2 M Na₂CO₃ (54 mL) are successively added. The resulting mixture is stirred at reflux overnight. The obtained reaction mixture is diluted with Et₂O and the solvents are removed under reduced pressure. The residue is purified by FC (Teledyne Isco Combiflash Rf, SiO₂ cartridge 120 g; heptane to EtOAc/heptane 3:97) to

give methyl 3-fluoro-4-methyl-[1,1'-biphenyl]-2-carboxylate as a light yellow oil. LC-MS (conditions F): $t_R = 0.94$ min, $[M + 1]^+ = 245.19$.


b) To a solution of methyl 3-fluoro-4-methyl-[1,1'-biphenyl]-2-carboxylate (23 mmol, 1 eq) in MeOH (42 mL) is added NaOH 32% (24 mL). The resulting mixture is stirred at 65°C for 2 hour. The volatiles are evaporated under reduced pressure and the remaining aq. phase is acidified with 7 M HCl to pH = 1-2. The resulting suspension is filtered under vacuum and the obtained solid dried under high vacuum. 3-Fluoro-4-methyl-[1,1'-biphenyl]-2-carboxylic acid **48** (4.35 g) is obtained as a white solid. LC-MS (conditions F): $t_R = 0.81$ min, $[M + 1]^+ =$ not detectable.

5-Aryl-2-methyl-thiazole-4-carboxylic acids.

These acids were prepared by a three-step sequence depicted in the scheme below, e.g. Darzens condensation of 4-fluoro-benzaldehyde **49** with methyl dichloroacetate to give the methyl keto-ester **50** (Hamamoto H. *et al* *Tetrahedron Asymmetry* **2000**, 11, 4485-4497) which is converted to the desired 5-(4-fluoro-phenyl)-2-methyl-4-carboxylic acid **51** by reaction with thioamide (US 3,282,927) followed by ester hydrolysis under basic conditions.




The following compounds were prepared according to the same sequence:



Preparation of the Examples:

Synthesis of Example 1.1:



Step 1: 5-Methyl-2-(2H-1,2,3-triazol-2-yl)benzoic acid (**2**; 12.27 g, 60.4 mmol) and L-proline methylester hydrochloride (**1**; 10.20 g, 60.4 mmol) are dissolved in DCM (250 ml) followed by the addition of DIPEA (23.42 g, 31 ml, 181 mmol) and HATU (22.95 g, 60.4 mmol). Stirring at room temperature is continued for 1 h, the DCM is evaporated under reduced pressure and EtOAc (750 ml) is added. The reaction mixture is extracted with brine (3 x 300 ml). The organic layer is dried with MgSO₄, filtered and the solvent is evaporated under reduced pressure to give (S)-methyl 1-(5-methyl-2-(2H-1,2,3-triazol-2-yl)benzoyl)pyrrolidine-2-carboxylate (**3**) which is used in the next step without further purification. LC-MS: t_R = 0.66 min; $[M+H]^+$ = 315.09.


Step 2: (S)-Methyl 1-(5-methyl-2-(2H-1,2,3-triazol-2-yl)benzoyl)pyrrolidine-2-carboxylate (**3**, 18.81 g, 60.4 mmol) is dissolved in a mixture of MeOH (180 ml) and THF (210 ml) followed by the addition of aq. NaOH solution (1 M; 150 ml). Stirring is continued for 1 h. The reaction mixture is concentrated (removal of the organic solvents) under reduced pressure and acidified to pH 1 by the addition of aq. HCl (1M). The product is extracted with EtOAc (3 x 250 ml). The combined organic layers are dried with MgSO₄, filtered and the solvent is evaporated under reduced pressure to give 17.14 g (87%) (S)-1-(5-methyl-2-(2H-1,2,3-triazol-2-yl)benzoyl)pyrrolidine-2-carboxylic acid (**4**). LC-MS: t_R = 0.57 min; $[M+H]^+$ = 301.17.

Step 3: (S)-1-(5-Methyl-2-(2H-1,2,3-triazol-2-yl)benzoyl)pyrrolidine-2-carboxylic acid (**4**; 17.11 g, 51.3 mmol) is dissolved in DCM (250 ml) at RT followed by the addition of 3,4-diaminoanisole dihydrochloride (**5**; 10.83 g, 51.3 mmol). To this reaction mixture DIPEA (23.2 g, 180 mmol) is then slowly added followed by the addition of HATU (19.7 g, 51.8 mmol). Stirring is continued for 1 h at RT. The reaction mixture is concentrated under reduced pressure. The residue is taken up in EtOAc (750 ml) and washed with brine (2x 500 ml). The organic layer is dried with MgSO₄, filtered and concentrated under reduced pressure. The crude product is purified by flash master chromatography (Silicagel; EtOAc / MeOH = 95 / 5) to give (S)-N-(2-amino-4-methoxyphenyl)-1-(5-methyl-2-(2H-1,2,3-triazol-2-yl)benzoyl)pyrrolidine-2-carboxamide (**6**). LC-MS: t_R = 0.64 min; $[M+H]^+$ = 421.17.

Step 4: (S)-N-(2-amino-4-methoxyphenyl)-1-(5-methyl-2-(2H-1,2,3-triazol-2-yl)benzoyl)pyrrolidine-2-carboxamide (**6**; 21.57 g, 51.3 mmol) is dissolved in pure AcOH (390 ml) and heated to 100°C for 30 minutes. The reaction mixture is cooled to rt and the acetic acid is removed under reduced pressure. The residual material is carefully diluted with sat. aq. NaHCO₃ solution (600 ml). The product precipitated and is filtered off and purified by flash master chromatography (Silicagel, DCM / MeOH = 98 / 2) to give (S)-(2-(6-methoxy-1H-

benzo[d]imidazol-2-yl)pyrrolidin-1-yl)(5-methyl-2-(2H-1,2,3-triazol-2-yl)phenyl)methanone (Example 1.1) as a colorless powder. LC-MS: $t_R = 0.55$ min; $[M+H]^+ = 403.16$.

Synthesis of Example 2.1:



5

Step 1: (2S,4S)-1-(tert-Butoxycarbonyl)-4-fluoropyrrolidine-2-carboxylic acid (7; 144.27 mg, 0.6 mmol) is dissolved in a 1/1 mixture of DMF / DCM (1.2 ml) followed by the addition of DIPEA (0.44 ml, 2.52 mmol) and 4-chloro-3-methylbenzene-1,2-diamine (8; 121.95 mg, 0.6 mmol) dissolved in a 1/1 mixture of DMF / DCM (1.2 ml) and finally by the addition of HATU (240 mg, 0.63 mmol) dissolved in DMF (1 ml). The reaction mixture is stirred at RT for 15 h then passed through PL-HCO₃ packed filter syringe (1g) with a 1/1-mixture of DMF / DCM (4 ml). The solvent is removed under reduced pressure to give (2S,4S)-tert-butyl 2-((2-amino-4-chloro-3-methylphenyl)carbamoyl)-4-fluoropyrrolidine-1-carboxylate (9) which is used in the next step without further purification. LC-MS: $t_R = 0.79$ min; $[M+H]^+ = 372.26$.

10

Step 2: (2S,4S)-tert-Butyl 2-((2-amino-4-chloro-3-methylphenyl)carbamoyl)-4-fluoropyrrolidine-1-carboxylate (9; 220 mg, 0.6 mmol) is dissolved in 100 % AcOH (3 ml, 52.5 mmol) and heated to 60°C for 3 h. Toluene (3 ml) is added to the reaction mixture and the solvents are removed under reduced pressure to give (2S,4S)-tert-butyl 2-(6-chloro-7-methyl-1H-benzo[d]imidazol-2-yl)-4-fluoropyrrolidine-1-carboxylate (10), which is used in the next step without further purification. LC-MS: $t_R = 0.8$ min; $[M+H]^+ = 354.25$.


20

Step 3: (2S,4S)-tert-Butyl 2-(6-chloro-7-methyl-1H-benzo[d]imidazol-2-yl)-4-fluoropyrrolidine-1-carboxylate (10; 210 mg, 0.6 mmol) is dissolved in dioxane (0.5 ml) and a solution of HCl in dioxane (4M, 3 ml, 12 mmol) is carefully added. The reaction mixture turns into a

suspension. To solubilize the reaction mixture MeOH (1 ml) is added. Stirring is continued for 2 h. The solvents are removed under reduced pressure to give 6-chloro-2-((2S,4S)-4-fluoropyrrolidin-2-yl)-7-methyl-1H-benzo[d]imidazole hydrochloride (11) which is used in the next step without further purification. LC-MS: $t_R = 0.61$ min; $[M+H]^+ = 254.14$.

- 5 Step 4: 6-Chloro-2-((2S,4S)-4-fluoropyrrolidin-2-yl)-7-methyl-1H-benzo[d]imidazole hydrochloride (11; 30 mg, 0.1 mmol) is dissolved in DCM (0.2 ml) and DIPEA (0.072 ml, 0.42 mmol) is added, followed by the addition of a solution of 5-methyl-2-(2H-1,2,3-triazol-2-yl)benzoic acid (2; 20.3 mg, 0.1 mmol), HATU (40 mg, 0.105 mmol) and DIPEA (80 mg, 0.62 mmol) in 0.5 ml DMF. Stirring is continued at room temperature for 16 h. The reaction mixture is diluted with DCM / MeOH = 1/1 (1 ml) followed by the addition of PL-HCO₃-resin (213 mg, 0.4 mmol) and stirring is continued for 2 h. The resin is filtered off, the solvent is evaporated under reduced pressure and the product is purified by preparative HPLC to give ((2S,4S)-2-(6-chloro-7-methyl-1H-benzo[d]imidazol-2-yl)-4-fluoropyrrolidin-1-yl)(5-methyl-2-(2H-1,2,3-triazol-2-yl)phenyl)methanone (Example 2.1) as a colorless powder. LC-MS: $t_R =$
- 10
- 15 1.14 min; $[M+H]^+ = 439.25$.

Synthesis of Example 3.1:



Ex. 3.1

Step 1: (2S,4S)-1-(tert-Butoxycarbonyl)-4-hydroxypyrrolidine-2-carboxylic acid (**12**, 1.503 g, 6.5 mmol) is dissolved in DMF (9.06 ml) and added to a suspension of NaH (60% dispersion

in mineral oil, 676 mg, 16.9 mmol) in DMF (9.06 ml) in an argon atmosphere at RT and stirring is continued for 30 minutes followed by the addition of MeI (3.14 g, 22.14 mmol). Stirring is continued for 23 h. An aq., sat. solution of KH_2PO_4 (30 ml) is added followed by the addition of solid KH_2PO_4 (400 mg) and another portion of an aq., sat. solution of KH_2PO_4 (10 ml). The product is extracted with Et_2O (3 x 40 ml). The combined organic layers are dried with MgSO_4 , filtered and the solvent is removed under reduced pressure to give the 3-methoxy-proline methylester-intermediate, which is dissolved in MeOH (4.5 ml) followed by the addition of an aq. NaOH solution (3M, 3.76 ml, 11.3 mmol). Stirring is continued for 12 h. The reaction mixture is neutralized to pH = 9, by the addition of a sat. solution of KH_2PO_4 and extracted with Et_2O (3 x 8 ml). The combined ether layers are discarded and the aq. layer is acidified to pH = 4.5 with aq. HCl. The product is extracted with EtOAc (3 x 40 ml). The combined organic layers are dried with MgSO_4 , filtered and the solvent is removed under reduced pressure to give (2S,4S)-1-(tert-butoxycarbonyl)-4-methoxypyrrolidine-2-carboxylic acid (**13**).

Step 2: (2S,4S)-1-(tert-Butoxycarbonyl)-4-methoxypyrrolidine-2-carboxylic acid (**13**; 147.1 mg, 0.6 mmol) is dissolved in a 1/1 mixture of DMF / DCM (1.2 ml) followed by the addition of DIPEA (0.44 ml, 2.52 mmol) and 4-chloro-3-methylbenzene-1,2-diamine (**8**; 121.95 mg, 0.6 mmol) dissolved in a 1/1 mixture of DMF / DCM (1.2 ml) and finally by the addition of HATU (240 mg, 0.63 mmol) dissolved in DMF (1 ml). The reaction mixture is stirred at RT for 15 h then passed through PL- HCO_3 packed filter syringe (1g) with a 1/1-mixture of DMF / DCM (4 ml). The solvent is removed under reduced pressure to give (2S,4S)-tert-butyl 2-((2-amino-4-chloro-3-methylphenyl)carbamoyl)-4-methoxypyrrolidine-1-carboxylate (**14**) which is used in the next step without further purification. LC-MS: t_R = 0.8 min; $[\text{M}+\text{H}]^+$ = 384.28.


Step 3: (2S,4S)-tert-Butyl 2-((2-amino-4-chloro-3-methylphenyl)carbamoyl)-4-methoxypyrrolidine-1-carboxylate (**14**; 230 mg, 0.6 mmol) is dissolved in 100 % AcOH (3 ml, 52.5 mmol) and heated to 60°C for 3 h. Toluene (3 ml) is added to the reaction mixture and the solvents are removed under reduced pressure to give (2S,4S)-tert-butyl 2-(6-chloro-7-methyl-1H-benzo[d]imidazol-2-yl)-4-methoxypyrrolidine-1-carboxylate (**15**), which is used in the next step without further purification. LC-MS: t_R = 0.80 min; $[\text{M}+\text{H}]^+$ = 366.27.

Step 4: (2S,4S)-tert-Butyl 2-(6-chloro-7-methyl-1H-benzo[d]imidazol-2-yl)-4-methoxypyrrolidine-1-carboxylate (**15**; 220 mg, 0.6 mmol) is dissolved in dioxane (0.5 ml) and a solution of HCl in dioxane (4M, 3 ml, 12 mmol) is carefully added. The reaction mixture turns into a suspension. To solubilize the reaction mixture MeOH (1 ml) is added. Stirring is continued for 2 h. The solvents are removed under reduced pressure to give 6-chloro-2-

((2S,4S)-4-methoxypyrrolidin-2-yl)-7-methyl-1H-benzo[d]imidazole hydrochloride (**16**) which is used in the next step without further purification. LC-MS: $t_R = 0.61$ min; $[M+H]^+ = 266.16$.

Step 5: 6-Chloro-2-((2S,4S)-4-methoxypyrrolidin-2-yl)-7-methyl-1H-benzo[d]imidazole (**16**; 30 mg, 0.1 mmol) is dissolved in DCM (0.2 ml) and DIPEA (0.072 ml, 0.42 mmol) is added, followed by the addition of a solution of 5-methyl-2-(2H-1,2,3-triazol-2-yl)benzoic acid (**2**; 20.3 mg, 0.1 mmol), HATU (40 mg, 0.105 mmol) and DIPEA (80 mg, 0.62 mmol) in 0.5 ml DMF. Stirring is continued at room temperature for 16 h. The reaction mixture is diluted with DCM / MeOH = 1/1 (1 ml) followed by the addition of PL-HCO₃-resin (213 mg, 0.4 mmol) and stirring is continued for 2 h. The resin is filtered off, the solvent is evaporated under reduced pressure and the product is purified by preparative HPLC to give ((2S,4S)-2-(6-chloro-7-methyl-1H-benzo[d]imidazol-2-yl)-4-methoxypyrrolidin-1-yl)(5-methyl-2-(2H-1,2,3-triazol-2-yl)phenyl)methanone (Ex. 3.1) as a colorless powder. LC-MS: $t_R = 1.16$ min; $[M+H]^+ = 451.29$.

Synthesis of Example 4.1:



Step 1: (2S,3S)-3-Methylpyrrolidine-2-carboxylic acid (**17**; 2.45 g, 19 mmol) is dissolved in a 1 / 1-mixture of MeCN / water (66.5 ml) and TEA (13.6 ml; 69.9 mmol) is added followed by the addition of Boc₂O (6.22 g, 28.5 mmol) dissolved in MeCN (14.25 ml). The reaction mixture is stirred at RT for 90 minutes. The organic solvents are evaporated under reduced pressure and water (24 ml) is added to the residue followed by the addition of aq. NaOH solution (1M, 28.5 ml, 28.5 mmol). The product is extracted with Et₂O (5 x 60 ml). The

20


combined organic layers are dried with MgSO_4 , filtered and the solvent is evaporated under reduced pressure to give (2S,3S)-1-(tert-butoxycarbonyl)-3-methylpyrrolidine-2-carboxylic acid (**18**). LC-MS: $t_R = 0.64$ min; $[\text{M}+\text{H}]^+ = 230.16$.

Step 2: (2S,3S)-1-(tert-Butoxycarbonyl)-3-methylpyrrolidine-2-carboxylic acid (**18**; 69 mg, 0.3 mmol) is dissolved in a 1/1 mixture of DMF / DCM (0.6 ml) followed by the addition of DIPEA (0.22 ml, 1.26 mmol) and 3-methylbenzene-1,2-diamine (**19**; 36.9 mg, 0.3 mmol) dissolved in a 1/1 mixture of DMF / DCM (0.6 ml) and finally by the addition of HATU (120 mg, 0.315 mmol) dissolved in DMF (0.63 ml). The reaction mixture is stirred at RT for 17 h then passed through PL- HCO_3 packed filter syringe (640 mg) with a 1/1-mixture of DMF / DCM (6 ml). The solvent is removed under reduced pressure to give (2S,3S)-tert-butyl 2-((2-amino-3-methylphenyl)carbamoyl)-3-methylpyrrolidine-1-carboxylate (**20**) which is used in the next step without further purification. LC-MS: $t_R = 0.75$ min; $[\text{M}+\text{H}]^+ = 334.11$.

Step 3: (2S,3S)-tert-Butyl 2-((2-amino-3-methylphenyl)carbamoyl)-3-methylpyrrolidine-1-carboxylate (**20**; 100 mg, 0.3 mmol) is dissolved in 100 % AcOH (3 ml, 52.5 mmol) and heated to 60°C for 3 h. Toluene (3 ml) is added to the reaction mixture and the solvents are removed under reduced pressure to give (2S,3S)-tert-butyl 3-methyl-2-(7-methyl-1H-benzo[d]imidazol-2-yl)pyrrolidine-1-carboxylate (**21**), which is used in the next step without further purification. LC-MS: $t_R = 0.75$ min; $[\text{M}+\text{H}]^+ = 316.11$.

Step 4: (2S,3S)-tert-Butyl 3-methyl-2-(7-methyl-1H-benzo[d]imidazol-2-yl)pyrrolidine-1-carboxylate (**21**; 94.5 mg, 0.3 mmol) is dissolved in methanol (0.75 ml) and a solution of HCl in dioxane (4M, 1.5 ml, 6 mmol) is carefully added. Stirring is continued for 2 h. The solvents are removed under reduced pressure to give 7-methyl-2-((2S,3S)-3-methylpyrrolidin-2-yl)-1H-benzo[d]imidazole hydrochloride (**22**) which is used in the next step without further purification. LC-MS: $t_R = 0.58$ min; $[\text{M}+\text{H}]^+ = 216.14$.

Step 5: 7-Methyl-2-((2S,3S)-3-methylpyrrolidin-2-yl)-1H-benzo[d]imidazole hydrochloride (**22**; 25.2 mg, 0.1 mmol) is dissolved in DCM (0.2 ml) and DIPEA (0.072 ml, 0.42 mmol) is added, followed by the addition of a solution of 5-methyl-2-(2H-1,2,3-triazol-2-yl)benzoic acid (**2**; 20.3 mg, 0.1 mmol), HATU (40 mg, 0.105 mmol) and DIPEA (80 mg, 0.62 mmol) in 0.5 ml DMF. Stirring is continued at room temperature for 16 h. The reaction mixture is diluted with DCM / MeOH = 1/1 (1 ml) followed by the addition of PL- HCO_3 -resin (213 mg, 0.4 mmol) and stirring is continued for 2 h. The resin is filtered off, the solvent is evaporated under reduced pressure and the product is purified by preparative HPLC to give (5-methyl-2-(2H-1,2,3-triazol-2-yl)phenyl)((2S,3S)-3-methyl-2-(7-methyl-1H-benzo[d]imidazol-2-yl)pyrrolidin-1-yl)methanone (Ex. 4.1) as a colorless powder. LC-MS: $t_R = 0.96$ min; $[\text{M}+\text{H}]^+ = 401.16$.

Synthesis of Example 5.1:

Step 1: (S)-2-Methylpyrrolidine-2-carboxylic acid hydrochloride (**23**; 82.5 g, 498 mmol) is dissolved in a 1 / 1-mixture of MeCN / water (1000 ml) and TEA (210 ml; 1490 mmol) is added followed by the addition of Boc_2O (120 g, 548 mmol) dissolved in MeCN (200 ml). The reaction mixture is stirred at RT for 3 hours. The organic solvents are evaporated under reduced pressure and 2M aqueous sodium hydroxide (300 ml) is added to the residue. The aqueous phase is extracted with Et_2O (400 ml), and cooled to 10°C followed by slow addition of 25% aqueous hydrochloric acid until the pH = 5-6 followed by careful addition of aqueous 1M hydrochloric acid to pH = 2. The product precipitates and is filtered off, washed with water (300 ml) and dried at high vacuum to give 90.7 g of (S)-1-(tert-butoxycarbonyl)-2-methylpyrrolidine-2-carboxylic acid (**24**) as a slightly grey solid. LC-MS: $t_R = 0.63$ min; $[\text{M}+\text{H}]^+ = 230.24$.


Step 2: (S)-1-(tert-Butoxycarbonyl)-2-methylpyrrolidine-2-carboxylic acid (**24**; 137.6 mg, 0.6 mmol) is dissolved in a 1/1 mixture of DMF / DCM (1.2 ml) followed by the addition of DIPEA (0.44 ml, 2.52 mmol) and 3-methylbenzene-1,2-diamine (**19**; 73.8 mg, 0.6 mmol) dissolved in a 1/1 mixture of DMF / DCM (1.2 ml) and finally by the addition of HATU (240 mg, 0.63 mmol) dissolved in DMF (1.26 ml). The reaction mixture is stirred at RT for 17 h then passed through PL- HCO_3 packed filter syringe (1 g) with a 1/1-mixture of DMF / DCM (6 ml). The solvent is removed under reduced pressure to give (S)-tert-butyl 2-((2-amino-3-

methylphenyl)carbamoyl)-2-methylpyrrolidine-1-carboxylate (**25**) which is used in the next step without further purification. LC-MS: $t_R = 0.78$ min; $[M+H]^+ = 334.3$.

Step 3: (S)-tert-Butyl 2-((2-amino-3-methylphenyl)carbamoyl)-2-methylpyrrolidine-1-carboxylate (**25**; 200 mg, 0.6 mmol) is dissolved in 100 % AcOH (3 ml, 52.5 mmol) and
5 heated to 60°C for 3 h. Toluene (3 ml) is added to the reaction mixture and the solvents are removed under reduced pressure to give (S)-tert-butyl 2-methyl-2-(7-methyl-1H-benzo[d]imidazol-2-yl)pyrrolidine-1-carboxylate (**26**), which is used in the next step without further purification. LC-MS: $t_R = 0.76$ min; $[M+H]^+ = 316.24$.

Step 4: (S)-tert-Butyl 2-methyl-2-(7-methyl-1H-benzo[d]imidazol-2-yl)pyrrolidine-1-
10 carboxylate (**26**; 189.7 mg, 0.6 mmol) is dissolved in MeOH (1.5 ml) and a solution of HCl in dioxane (4M, 3 ml, 12 mmol) is carefully added. Stirring is continued for 2 h. The solvents are removed under reduced pressure to give (S)-7-methyl-2-(2-methylpyrrolidin-2-yl)-1H-benzo[d]imidazole hydrochloride (**27**) which is used in the next step without further purification. LC-MS: $t_R = 0.57$ min; $[M+H]^+ = 216.31$.

15 Step 5: (S)-7-Methyl-2-(2-methylpyrrolidin-2-yl)-1H-benzo[d]imidazole hydrochloride (**27**; 25.2 mg, 0.1 mmol) is dissolved in DCM (0.2 ml) and DIPEA (0.072 ml, 0.42 mmol) is added, followed by the addition of a solution of 2-methyl-5-phenylthiazole-4-carboxylic acid (**28**; 22 mg, 0.1 mmol), HATU (40 mg, 0.105 mmol) and DIPEA (80 mg, 0.62 mmol) in 0.5 ml DMF. Stirring is continued at RT for 16 h. The reaction mixture is diluted with DCM / MeOH = 1/1 (1
20 ml) followed by the addition of PL-HCO₃-resin (213 mg, 0.4 mmol) and stirring is continued for 2 h. The resin is filtered off, the solvent is evaporated under reduced pressure and the product is purified by preparative HPLC to give (S)-(2-methyl-2-(7-methyl-1H-benzo[d]imidazol-2-yl)pyrrolidin-1-yl)(2-methyl-5-phenylthiazol-4-yl)methanone (Ex. 5.1) as a colorless powder. LC-MS: $t_R = 1.19$ min; $[M+H]^+ = 417.31$.

Synthesis of Example 6.1:

Step 1: (S)-1-(tert-Butoxycarbonyl)-4-methylenepyrrolidine-2-carboxylic acid (**29**; 1.0 g, 4.4 mmol) and HATU (1.84 g, 4.84 mmol) is suspended in DCM (30 ml) followed by the addition of DIPEA (3.8 ml, 22 mmol) and 4-methoxybenzene-1,2-diamine hydrochloride (**5**; 960 mg, 4.4 mmol). The reaction mixture is stirred at RT for 16 h. The mixture is then poured onto brine (100 ml) and extracted with DCM (2x 100 ml). The combined organic layers are dried over MgSO_4 , filtered and the solvent is removed under reduced pressure. The crude product is purified by FC (ethyl acetate) to give (S)-tert-butyl 2-((2-amino-4-methoxyphenyl)carbamoyl)-4-methylenepyrrolidine-1-carboxylate (**30**) as a slightly yellow amorphous solid. LC-MS: $t_R = 0.65$ min; $[\text{M}+\text{H}]^+ = 348.03$.


Step 2: (S)-tert-Butyl 2-((2-amino-4-methoxyphenyl)carbamoyl)-4-methylenepyrrolidine-1-carboxylate (**30**; 1.52 g, 4.38 mmol) is dissolved in 100 % AcOH (43.8 ml, 766 mmol) and heated to 100°C for 30 minutes. The reaction mixture is concentrated under reduced pressure followed by the addition of DCM (150 ml) to the residue. The organic layer is washed with sat. aq. NaHCO_3 solution (150 ml) and brine (150 ml) and dried over MgSO_4 , filtered and the solvent is evaporated to give (S)-tert-butyl 2-(6-methoxy-1H-benzo[d]imidazol-2-yl)-4-methylenepyrrolidine-1-carboxylate (**31**) as a slightly yellow, amorphous solid. LC-MS: $t_R = 0.55$ min; $[\text{M}+\text{H}]^+ = 330.22$.

Step 3: (S)-tert-Butyl 2-(6-methoxy-1H-benzo[d]imidazol-2-yl)-4-methylenepyrrolidine-1-carboxylate (**31**; 1.21 g, 3.67 mmol) is dissolved in dioxane (19 ml) and a solution of HCl in dioxane (4M, 19 ml, 73.5 mmol) is carefully added. Stirring is continued for 2 h. The solvents

are removed under reduced pressure and the product is dried at HV to give (S)-6-methoxy-2-(4-methylenepyrrolidin-2-yl)-1H-benzo[d]imidazole hydrochloride (**32**), as a slightly beige solid. LC-MS: t_R = 0.37 min; $[M+H]^+$ = 230.22.

Step 4: 5-(4-Fluorophenyl)-2-methylthiazole-4-carboxylic acid (**33**; 12.4 mg, 0.207 mmol) is dissolved in MeCN (1 ml) and TBTU (66.5 mg, 0.207 mmol) and DIPEA (0.161 ml, 122 mg, 0.941 mmol) are added, followed by the addition of a solution of (S)-6-methoxy-2-(4-methylenepyrrolidin-2-yl)-1H-benzo[d]imidazole hydrochloride (**32**; 50 mg, 0.188 mmol) and DIPEA (8 mg, 0.6 mmol, 0.05ml) in 0.5 ml MeCN. Stirring is continued at RT for 16 h. The product is directly purified by preparative HPLC to give (S)-(5-(4-fluorophenyl)-2-methylthiazol-4-yl)(2-(6-methoxy-1H-benzo[d]imidazol-2-yl)-4-methylenepyrrolidin-1-yl)methanone (Ex. 6.1) as a colorless powder. LC-MS: t_R = 0.63 min; $[M+H]^+$ = 448.97.

Synthesis of Example 7.4:



(S)-(2-(5-bromo-7-methyl-1H-benzo[d]imidazol-2-yl)-2-methylpyrrolidin-1-yl)(5-methyl-2-(2H-1,2,3-triazol-2-yl)phenyl)methanone (**34**; 157 mg, 0.329 mmol) is dissolved in DMF (2 ml) and Cs_2CO_3 (161 mg, 0.493 mmol) is added followed by the slow addition of a solution of MeI (58.3 mg, 0.411 mmol) in DMF (1 ml) at 0°C. Stirring is continued for 60 min at 0°C. The reaction mixture is poured onto water (100 ml). The precipitated product is filtered off, washed with water (10 ml), dried at HV to give an isomeric mixture of Ex. 7.4 consisting of (S)-(2-(5-bromo-1,7-dimethyl-1H-benzo[d]imidazol-2-yl)-2-methylpyrrolidin-1-yl)(5-methyl-2-(2H-1,2,3-triazol-2-yl)phenyl)methanone (**35a**) and (S)-(2-(6-bromo-1,4-dimethyl-1H-benzo[d]imidazol-2-yl)-2-methylpyrrolidin-1-yl)(5-methyl-2-(2H-1,2,3-triazol-2-

yl)phenyl)methanone (**35b**) as a slightly grey powder. LC-MS: t_R = 0.79 min; $[M+H]^+$ = 495.11.

According to the procedures described herein before, the following examples are prepared:

Example	Compound name. LC-MS data
1.1	(5-Methyl-2-[1,2,3]triazol-2-yl-phenyl)-[(S)-2-(4-nitro-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: t_R = 0.84; $[M+H]^+$ = 418.2
1.2	[(S)-2-(5-Methoxy-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: t_R = 0.63; $[M+H]^+$ = 403.2
1.3	[(S)-2-(5-Chloro-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: t_R = 0.71; $[M+H]^+$ = 407.1
1.4	[(S)-2-(7-Methoxy-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: t_R = 0.64; $[M+H]^+$ = 403.2
1.5	[(S)-2-(7-Methylsulfanyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: t_R = 0.68; $[M+H]^+$ = 419.2
1.6	[(S)-2-(7-Chloro-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: t_R = 0.74; $[M+H]^+$ = 407.2
1.7	[(S)-2-(5-Methylsulfanyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: t_R = 0.68; $[M+H]^+$ = 419.2
1.9	[5-(3-Chloro-phenyl)-2-methyl-thiazol-4-yl]-[(S)-2-(5-methoxy-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: t_R = 0.76; $[M+H]^+$ = 453.2
1.10	[(S)-2-(5-Chloro-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-[5-(3-chloro-phenyl)-2-methyl-thiazol-4-yl]-methanone. LC-MS: t_R = 0.84; $[M+H]^+$ = 457.1
1.11	[5-(3-Chloro-phenyl)-2-methyl-thiazol-4-yl]-[(S)-2-(7-methoxy-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: t_R = 0.77; $[M+H]^+$ = 453.1
1.12	[5-(3-Chloro-phenyl)-2-methyl-thiazol-4-yl]-[(S)-2-(4-hydroxy-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: t_R = 0.71; $[M+H]^+$ = 439.1
1.13	[5-(3-Chloro-phenyl)-2-methyl-thiazol-4-yl]-[(S)-2-(7-methylsulfanyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: t_R = 0.82; $[M+H]^+$ = 469.1
1.14	[(S)-2-(7-Chloro-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-[5-(3-chloro-phenyl)-2-methyl-thiazol-4-yl]-methanone. LC-MS: t_R = 0.87; $[M+H]^+$ = 457.1
1.15	[5-(3-Chloro-phenyl)-2-methyl-thiazol-4-yl]-[(S)-2-(5-methylsulfanyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: t_R = 0.81; $[M+H]^+$ = 469.1

1.16	[(S)-2-(6-Methoxy-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.63$; $[M+H]^+ = 403.2$
1.17	[(S)-2-(6-Methylsulfanyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.69$; $[M+H]^+ = 419.2$
1.18	[5-(4-Bromo-phenyl)-2-methyl-thiazol-4-yl]-[(S)-2-(6-methoxy-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.77$; $[M+H]^+ = 497.1$
1.19	[5-(2,3-Dichloro-phenyl)-2-methyl-thiazol-4-yl]-[(S)-2-(6-methoxy-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.81$; $[M+H]^+ = 487.1$
1.20	[(S)-2-(6-Methoxy-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-[5-(3-methoxy-phenyl)-2-methyl-thiazol-4-yl]-methanone. LC-MS: $t_R = 0.70$; $[M+H]^+ = 449.2$
1.21	[5-(3-Fluoro-phenyl)-2-methyl-thiazol-4-yl]-[(S)-2-(6-methoxy-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.71$; $[M+H]^+ = 437.2$
1.22	[(S)-2-(6-Methoxy-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-(2-methyl-5-p-tolyl-thiazol-4-yl)-methanone. LC-MS: $t_R = 0.75$; $[M+H]^+ = 433.2$
1.23	(2-Cyclopropyl-5-p-tolyl-thiazol-4-yl)-[(S)-2-(6-methoxy-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.83$; $[M+H]^+ = 459.3$
1.24	[5-(4-Ethyl-phenyl)-2-methyl-thiazol-4-yl]-[(S)-2-(6-methoxy-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.81$; $[M+H]^+ = 447.2$
1.25	[5-(3-Chloro-phenyl)-2-methyl-thiazol-4-yl]-[(S)-2-(6-methoxy-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.76$; $[M+H]^+ = 453.2$
1.26	(3',4'-Dimethyl-biphenyl-2-yl)-[(S)-2-(6-methoxy-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.80$; $[M+H]^+ = 426.3$
1.27	[(S)-2-(6-Methoxy-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-(2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.58$; $[M+H]^+ = 389.2$
1.28	[4-(3,4-Dichloro-phenyl)-2-methyl-pyrimidin-5-yl]-[(S)-2-(6-methoxy-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.76$; $[M+H]^+ = 482.2$
1.29	[4-(3-Chloro-phenyl)-2-methyl-pyrimidin-5-yl]-[(S)-2-(6-methoxy-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.68$; $[M+H]^+ = 448.2$
1.30	[4-(4-Bromo-3-chloro-phenyl)-pyrimidin-5-yl]-[(S)-2-(6-methoxy-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.73$; $[M+H]^+ = 512.11$
1.31	[4-(3,4-Dimethyl-phenyl)-2-methyl-pyrimidin-5-yl]-[(S)-2-(6-methoxy-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.69$; $[M+H]^+ = 442.3$

1.32	[(S)-2-(6-Methoxy-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-(2-methyl-4-phenyl-pyrimidin-5-yl)-methanone. LC-MS: $t_R = 0.59$; $[M+H]^+ = 414.2$
1.33	[4-(4-Bromo-3-chloro-phenyl)-2-methyl-pyrimidin-5-yl]-[(S)-2-(6-methoxy-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.77$; $[M+H]^+ = 526.1$
1.34	[5-(4-Bromo-phenyl)-2-methyl-thiazol-4-yl]-[(S)-2-(6-methylsulfanyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.83$; $[M+H]^+ = 513.1$
1.35	(2-Cyclopropyl-5-phenyl-thiazol-4-yl)-[(S)-2-(6-methylsulfanyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.82$; $[M+H]^+ = 461.2$
1.36	[5-(2,3-Dichloro-phenyl)-2-methyl-thiazol-4-yl]-[(S)-2-(6-methylsulfanyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.86$; $[M+H]^+ = 503.1$
1.37	[5-(3-Methoxy-phenyl)-2-methyl-thiazol-4-yl]-[(S)-2-(6-methylsulfanyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.75$; $[M+H]^+ = 465.2$
1.38	[(S)-2-(6-Methylsulfanyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-[2-methyl-5-(3-trifluoromethyl-phenyl)-thiazol-4-yl]-methanone. LC-MS: $t_R = 0.85$; $[M+H]^+ = 503.2$
1.39	[5-(3-Fluoro-phenyl)-2-methyl-thiazol-4-yl]-[(S)-2-(6-methylsulfanyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.76$; $[M+H]^+ = 453.2$
1.40	[(S)-2-(6-Methylsulfanyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-(2-methyl-5-p-tolyl-thiazol-4-yl)-methanone. LC-MS: $t_R = 0.80$; $[M+H]^+ = 449.2$
1.41	(2-Cyclopropyl-5-p-tolyl-thiazol-4-yl)-[(S)-2-(6-methylsulfanyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.87$; $[M+H]^+ = 475.3$
1.42	[5-(4-Ethyl-phenyl)-2-methyl-thiazol-4-yl]-[(S)-2-(6-methylsulfanyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.85$; $[M+H]^+ = 463.2$
1.43	[5-(3-Chloro-phenyl)-2-methyl-thiazol-4-yl]-[(S)-2-(6-methylsulfanyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.81$; $[M+H]^+ = 469.2$
1.44	[5-(3,5-Difluoro-phenyl)-2-methyl-thiazol-4-yl]-[(S)-2-(6-methylsulfanyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.79$; $[M+H]^+ = 471.2$
1.45	[3-(3-Fluoro-5-methyl-phenyl)-pyrazin-2-yl]-[(S)-2-(6-methylsulfanyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.73$; $[M+H]^+ = 448.2$
1.46	(3',4'-Dimethyl-biphenyl-2-yl)-[(S)-2-(6-methylsulfanyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.85$; $[M+H]^+ = 442.2$
1.47	[(S)-2-(6-Methylsulfanyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-(2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.63$; $[M+H]^+ = 405.2$

1.49	[4-(3,4-Dichloro-phenyl)-2-methyl-pyrimidin-5-yl]-[(S)-2-(6-methylsulfanyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.82$; $[M+H]^+ = 498.1$
1.50	[4-(4-Fluoro-phenyl)-2-methyl-pyrimidin-5-yl]-[(S)-2-(6-methylsulfanyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.69$; $[M+H]^+ = 448.2$
1.51	[4-(3-Chloro-phenyl)-2-methyl-pyrimidin-5-yl]-[(S)-2-(6-methylsulfanyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.74$; $[M+H]^+ = 464.2$
1.52	[4-(3,4-Dichloro-phenyl)-pyrimidin-5-yl]-[(S)-2-(6-methylsulfanyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.78$; $[M+H]^+ = 484.1$
1.53	4-(3,4-Dimethyl-phenyl)-pyrimidin-5-yl]-[(S)-2-(6-methylsulfanyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.72$; $[M+H]^+ = 444.2$
1.54	[4-(4-Bromo-3-chloro-phenyl)-pyrimidin-5-yl]-[(S)-2-(6-methylsulfanyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.79$; $[M+H]^+ = 528.1$
1.55	[4-(3,4-Dimethyl-phenyl)-2-methyl-pyrimidin-5-yl]-[(S)-2-(6-methylsulfanyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.75$; $[M+H]^+ = 458.3$
1.56	[4-(3-Methoxy-phenyl)-2-methyl-pyrimidin-5-yl]-[(S)-2-(6-methylsulfanyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.67$; $[M+H]^+ = 460.3$
1.57	(2-Methyl-4-phenyl-pyrimidin-5-yl)-[(S)-2-(6-methylsulfanyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.66$; $[M+H]^+ = 430.2$
1.58	[4-(4-Bromo-3-chloro-phenyl)-2-methyl-pyrimidin-5-yl]-[(S)-2-(6-methylsulfanyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.83$; $[M+H]^+ = 542.0$
1.59	[4-(3,5-Dichloro-phenyl)-2-methyl-pyrimidin-5-yl]-[(S)-2-(6-methylsulfanyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.82$; $[M+H]^+ = 498.2$
1.60	[5-(4-Bromo-phenyl)-2-methyl-thiazol-4-yl]-[(S)-2-(7-methylsulfanyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.84$; $[M+H]^+ = 513.1$
1.61	(2-Cyclopropyl-5-phenyl-thiazol-4-yl)-[(S)-2-(7-methylsulfanyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.83$; $[M+H]^+ = 461.2$
1.62	[5-(2,3-Dichloro-phenyl)-2-methyl-thiazol-4-yl]-[(S)-2-(7-methylsulfanyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.86$; $[M+H]^+ = 503.1$
1.63	[5-(3-Methoxy-phenyl)-2-methyl-thiazol-4-yl]-[(S)-2-(7-methylsulfanyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.76$; $[M+H]^+ = 465.2$
1.64	[(S)-2-(7-Methylsulfanyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-[2-methyl-5-(3-trifluoromethyl-phenyl)-thiazol-4-yl]-methanone. LC-MS: $t_R = 0.86$; $[M+H]^+ = 503.2$

1.65	[5-(3-Fluoro-phenyl)-2-methyl-thiazol-4-yl]-[(S)-2-(7-methylsulfanyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.77$; $[M+H]^+ = 453.2$
1.66	[(S)-2-(7-Methylsulfanyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-(2-methyl-5-p-tolyl-thiazol-4-yl)-methanone. LC-MS: $t_R = 0.80$; $[M+H]^+ = 449.2$
1.67	[5-(3,4-Difluoro-phenyl)-2-methyl-thiazol-4-yl]-[(S)-2-(7-methylsulfanyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.80$; $[M+H]^+ = 471.1$
1.68	(2-Cyclopropyl-5-p-tolyl-thiazol-4-yl)-[(S)-2-(7-methylsulfanyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.88$; $[M+H]^+ = 475.3$
1.69	[5-(4-Ethyl-phenyl)-2-methyl-thiazol-4-yl]-[(S)-2-(7-methylsulfanyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.86$; $[M+H]^+ = 463.2$
1.70	[2-Cyclopropyl-5-(4-fluoro-phenyl)-thiazol-4-yl]-[(S)-2-(7-methylsulfanyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.84$; $[M+H]^+ = 479.2$
1.71	[5-(3,5-Difluoro-phenyl)-2-methyl-thiazol-4-yl]-[(S)-2-(7-methylsulfanyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.79$; $[M+H]^+ = 471.2$
1.72	[3-(4-Methoxy-phenyl)-pyrazin-2-yl]-[(S)-2-(7-methylsulfanyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.68$; $[M+H]^+ = 446.2$
1.73	[(S)-2-(7-Methylsulfanyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-[3-(3-trifluoromethyl-phenyl)-pyrazin-2-yl]-methanone. LC-MS: $t_R = 0.80$; $[M+H]^+ = 484.2$
1.74	[3-(3-Fluoro-5-methyl-phenyl)-pyrazin-2-yl]-[(S)-2-(7-methylsulfanyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.74$; $[M+H]^+ = 448.2$
1.75	(3',4'-Dimethyl-biphenyl-2-yl)-[(S)-2-(7-methylsulfanyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.87$; $[M+H]^+ = 442.2$
1.76	[(S)-2-(7-Methylsulfanyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-(2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.63$; $[M+H]^+ = 405.2$
1.78	[(S)-2-(7-Methylsulfanyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-(4-p-tolyl-pyrimidin-5-yl)-methanone. LC-MS: $t_R = 0.70$; $[M+H]^+ = 430.2$
1.79	[4-(3,4-Dichloro-phenyl)-2-methyl-pyrimidin-5-yl]-[(S)-2-(7-methylsulfanyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.86$; $[M+H]^+ = 498.2$
1.80	[4-(4-Fluoro-phenyl)-2-methyl-pyrimidin-5-yl]-[(S)-2-(7-methylsulfanyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.71$; $[M+H]^+ = 448.2$
1.81	[4-(3-Chloro-phenyl)-2-methyl-pyrimidin-5-yl]-[(S)-2-(7-methylsulfanyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.77$; $[M+H]^+ = 464.2$

1.82	[4-(3,4-Dichloro-phenyl)-pyrimidin-5-yl]-[(S)-2-(7-methylsulfanyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.74$; $[M+H]^+ = 444.3$
1.83	[4-(3,4-Dimethyl-phenyl)-pyrimidin-5-yl]-[(S)-2-(7-methylsulfanyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.83$; $[M+H]^+ = 528.0$
1.84	[4-(4-Bromo-3-chloro-phenyl)-pyrimidin-5-yl]-[(S)-2-(7-methylsulfanyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.77$; $[M+H]^+ = 458.2$
1.85	[4-(3,4-Dimethyl-phenyl)-2-methyl-pyrimidin-5-yl]-[(S)-2-(7-methylsulfanyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.68$; $[M+H]^+ = 460.2$
1.86	[4-(3-Methoxy-phenyl)-2-methyl-pyrimidin-5-yl]-[(S)-2-(7-methylsulfanyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.67$; $[M+H]^+ = 430.2$
1.87	(2-Methyl-4-phenyl-pyrimidin-5-yl)-[(S)-2-(7-methylsulfanyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.70$; $[M+H]^+ = 430.2$
1.88	[(S)-2-(7-Methylsulfanyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-(4-m-tolyl-pyrimidin-5-yl)-methanone. LC-MS: $t_R = 0.82$; $[M+H]^+ = 484.1$
1.89	[4-(3,5-Dichloro-phenyl)-pyrimidin-5-yl]-[(S)-2-(7-methylsulfanyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.87$; $[M+H]^+ = 542.1$
1.90	[4-(4-Bromo-3-chloro-phenyl)-2-methyl-pyrimidin-5-yl]-[(S)-2-(7-methylsulfanyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.86$; $[M+H]^+ = 498.2$
1.91	[4-(3,5-Dichloro-phenyl)-2-methyl-pyrimidin-5-yl]-[(S)-2-(7-methylsulfanyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.99$; $[M+H]^+ = 491.1$
1.92	rac-[5-(3-Chloro-phenyl)-2-methyl-thiazol-4-yl]-[2-(5,6-dichloro-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.86$; $[M+H]^+ = 501.1$
1.93	rac-[2-(5-Bromo-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-[5-(3-chloro-phenyl)-2-methyl-thiazol-4-yl]-methanone. LC-MS: $t_R = 0.94$; $[M+H]^+ = 491.1$
1.94	rac-[5-(3-Chloro-phenyl)-2-methyl-thiazol-4-yl]-[2-(5-trifluoromethyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.82$; $[M+H]^+ = 451.2$
1.95	rac-[5-(3-Chloro-phenyl)-2-methyl-thiazol-4-yl]-[2-(5,6-dimethyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.78$; $[M+H]^+ = 437.1$
1.96	rac-[5-(3-Chloro-phenyl)-2-methyl-thiazol-4-yl]-[2-(4-methyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.84$; $[M+H]^+ = 457.1$
1.97	rac-[2-(5-Chloro-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-[5-(3-chloro-phenyl)-2-methyl-thiazol-4-yl]-methanone. LC-MS: $t_R = 0.92$; $[M+H]^+ = 475.2$

1.98	rac-[2-(6-Chloro-5-fluoro-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-[5-(3-chloro-phenyl)-2-methyl-thiazol-4-yl]-methanone. LC-MS: $t_R = 0.91$; $[M+H]^+ = 479.2$
1.99	rac-[2-(5-tert-Butyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-[5-(3-chloro-phenyl)-2-methyl-thiazol-4-yl]-methanone. LC-MS: $t_R = 1.12$; $[M+H]^+ = 525.1$
1.100	rac-[5-(3-Chloro-phenyl)-2-methyl-thiazol-4-yl]-[2-(4-chloro-6-trifluoromethyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.86$; $[M+H]^+ = 471.1$
1.101	rac-[2-(5-Chloro-6-methyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-[5-(3-chloro-phenyl)-2-methyl-thiazol-4-yl]-methanone. LC-MS: $t_R = 0.83$; $[M+H]^+ = 451.2$
1.102	rac-[5-(3-Chloro-phenyl)-2-methyl-thiazol-4-yl]-[2-(4,6-dimethyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.71$; $[M+H]^+ = 453.2$
1.103	rac-[5-(3-Chloro-phenyl)-2-methyl-thiazol-4-yl]-[2-(4-hydroxymethyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.84$; $[M+H]^+ = 441.1$
1.104	rac-[5-(3-Chloro-phenyl)-2-methyl-thiazol-4-yl]-[2-(4-fluoro-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.97$; $[M+H]^+ = 491.2$
1.105	rac-[5-(3-Chloro-phenyl)-2-methyl-thiazol-4-yl]-[2-(4-trifluoromethyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.89$; $[M+H]^+ = 501.1$
1.106	rac-[2-(4-Bromo-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-[5-(3-chloro-phenyl)-2-methyl-thiazol-4-yl]-methanone. LC-MS: $t_R = 0.89$; $[M+H]^+ = 515.2$
1.107	rac-[2-(5-Bromo-7-methyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-[5-(3-chloro-phenyl)-2-methyl-thiazol-4-yl]-methanone. LC-MS: $t_R = 0.92$; $[M+H]^+ = 507.2$
1.108	rac-[5-(3-Chloro-phenyl)-2-methyl-thiazol-4-yl]-[2-(5-trifluoromethoxy-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.87$; $[M+H]^+ = 471.2$
1.109	rac-[2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-[5-(3-chloro-phenyl)-2-methyl-thiazol-4-yl]-methanone. LC-MS: $t_R = 0.73$; $[M+H]^+ = 483.2$
1.110	rac-[5-(3-Chloro-phenyl)-2-methyl-thiazol-4-yl]-[2-(5,6-dimethoxy-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 1.02$; $[M+H]^+ = 509.1$
1.111	rac-[5-(3-Chloro-phenyl)-2-methyl-thiazol-4-yl]-[2-(6-fluoro-5-trifluoromethyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.93$; $[M+H]^+ = 519.1$
1.112	rac-[2-(6-Bromo-5-fluoro-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-[5-(3-chloro-phenyl)-2-methyl-thiazol-4-yl]-methanone. LC-MS: $t_R = 1.08$; $[M+H]^+ = 525.3$
1.113	rac-[5-(3-Chloro-phenyl)-2-methyl-thiazol-4-yl]-[2-(6-chloro-5-trifluoromethyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.99$; $[M+H]^+ = 519.1$

1.114	rac-[2-(4-Bromo-6-fluoro-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-[5-(3-chloro-phenyl)-2-methyl-thiazol-4-yl]-methanone. LC-MS: $t_R = 0.82$; $[M+H]^+ = 451.2$
1.115	rac-[5-(3-Chloro-phenyl)-2-methyl-thiazol-4-yl]-[2-(4,5-dimethyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.87$; $[M+H]^+ = 465.2$
1.116	rac-[5-(3-Chloro-phenyl)-2-methyl-thiazol-4-yl]-[2-(4-isopropyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.75$; $[M+H]^+ = 481.2$
1.117	rac-[5-(3-Chloro-phenyl)-2-methyl-thiazol-4-yl]-[2-(7,8-dihydro-3H-6,9-dioxo-1,3-diaza-cyclopenta[]naphthalen-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 1.12$; $[M+H]^+ = 525.1$
1.118	rac-[2-(4,6-Bis-trifluoromethyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-(2-methyl-5-phenyl-thiazol-4-yl)-methanone. LC-MS: $t_R = 0.92$; $[M+H]^+ = 457.1$
1.119	rac-[2-(5,6-Dichloro-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-(2-methyl-5-phenyl-thiazol-4-yl)-methanone. LC-MS: $t_R = 0.79$; $[M+H]^+ = 467.1$
1.120	rac-[2-(5-Bromo-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-(2-methyl-5-phenyl-thiazol-4-yl)-methanone. LC-MS: $t_R = 0.87$; $[M+H]^+ = 457.2$
1.121	rac-(2-Methyl-5-phenyl-thiazol-4-yl)-[2-(5-trifluoromethyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.76$; $[M+H]^+ = 417.2$
1.122	rac-[2-(5,6-Dimethyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-(2-methyl-5-phenyl-thiazol-4-yl)-methanone. LC-MS: $t_R = 0.71$; $[M+H]^+ = 403.2$
1.123	rac-[2-(4-Methyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-(2-methyl-5-phenyl-thiazol-4-yl)-methanone. LC-MS: $t_R = 0.85$; $[M+H]^+ = 445.3$
1.124	rac-[2-(5-tert-Butyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-(2-methyl-5-phenyl-thiazol-4-yl)-methanone. LC-MS: $t_R = 1.06$; $[M+H]^+ = 491.1$
1.125	rac-[2-(4-Chloro-6-trifluoromethyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-(2-methyl-5-phenyl-thiazol-4-yl)-methanone. LC-MS: $t_R = 0.80$; $[M+H]^+ = 437.2$
1.126	rac-[2-(5-Chloro-6-methyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-(2-methyl-5-phenyl-thiazol-4-yl)-methanone. LC-MS: $t_R = 0.76$; $[M+H]^+ = 417.2$
1.127	rac-[2-(4,6-Dimethyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-(2-methyl-5-phenyl-thiazol-4-yl)-methanone. LC-MS: $t_R = 0.63$; $[M+H]^+ = 419.1$
1.128	rac-[2-(4-Hydroxymethyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-(2-methyl-5-phenyl-thiazol-4-yl)-methanone. LC-MS: $t_R = 0.90$; $[M+H]^+ = 457.2$
1.129	rac-(2-Methyl-5-phenyl-thiazol-4-yl)-[2-(4-trifluoromethyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.81$; $[M+H]^+ = 467.1$

1.130	rac-[2-(4-Bromo-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-(2-methyl-5-phenyl-thiazol-4-yl)-methanone. LC-MS: $t_R = 0.82$; $[M+H]^+ = 481.1$
1.131	rac-[2-(5-Bromo-7-methyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-(2-methyl-5-phenyl-thiazol-4-yl)-methanone. LC-MS: $t_R = 0.86$; $[M+H]^+ = 473.2$
1.132	rac-(2-Methyl-5-phenyl-thiazol-4-yl)-[2-(5-trifluoromethoxy-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.81$; $[M+H]^+ = 437.2$
1.133	rac-[2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-(2-methyl-5-phenyl-thiazol-4-yl)-methanone. LC-MS: $t_R = 0.67$; $[M+H]^+ = 449.2$
1.134	rac-[2-(5,6-Dimethoxy-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-(2-methyl-5-phenyl-thiazol-4-yl)-methanone. LC-MS: $t_R = 0.96$; $[M+H]^+ = 475.2$
1.135	rac-[2-(6-Fluoro-5-trifluoromethyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-(2-methyl-5-phenyl-thiazol-4-yl)-methanone. LC-MS: $t_R = 0.86$; $[M+H]^+ = 485.1$
1.136	rac-[2-(6-Bromo-5-fluoro-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-(2-methyl-5-phenyl-thiazol-4-yl)-methanone. LC-MS: $t_R = 1.01$; $[M+H]^+ = 491.2$
1.137	rac-[2-(6-Chloro-5-trifluoromethyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-(2-methyl-5-phenyl-thiazol-4-yl)-methanone. LC-MS: $t_R = 0.92$; $[M+H]^+ = 485.1$
1.138	rac-[2-(4-Bromo-6-fluoro-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-(2-methyl-5-phenyl-thiazol-4-yl)-methanone. LC-MS: $t_R = 0.75$; $[M+H]^+ = 417.2$
1.139	rac-[2-(4,5-Dimethyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-(2-methyl-5-phenyl-thiazol-4-yl)-methanone. LC-MS: $t_R = 0.69$; $[M+H]^+ = 447.2$
1.140	rac-[2-(7,8-Dihydro-3H-6,9-dioxo-1,3-diaza-cyclopenta[]naphthalen-2-yl)-pyrrolidin-1-yl]-(2-methyl-5-phenyl-thiazol-4-yl)-methanone. LC-MS: $t_R = 0.63$; $[M+H]^+ = 403.2$
1.141	rac-[2-(5-Methoxy-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.76$; $[M+H]^+ = 417.2$
1.142	[(S)-2-(4,6-Dimethyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-(2-methyl-5-phenyl-thiazol-4-yl)-methanone. LC-MS: $t_R = 0.80$; $[M+H]^+ = 437.2$
1.143	[(S)-2-(5-Chloro-6-methyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-(2-methyl-5-phenyl-thiazol-4-yl)-methanone. LC-MS: $t_R = 0.82$; $[M+H]^+ = 481.1$
1.144	[(S)-2-(6-Bromo-4-methyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-(2-methyl-5-phenyl-thiazol-4-yl)-methanone. LC-MS: $t_R = 0.81$; $[M+H]^+ = 437.1$
1.145	[(S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-(2-methyl-5-phenyl-thiazol-4-yl)-methanone. LC-MS: $t_R = 1.02$; $[M+H]^+ = 491.1$

1.146	[(S)-2-(5-Chloro-6-trifluoromethyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-(2-methyl-5-phenyl-thiazol-4-yl)-methanone. LC-MS: $t_R = 0.67$; $[M+H]^+ = 423.2$
1.147	(5-Fluoro-2-[1,2,3]triazol-2-yl-phenyl)-[(S)-2-(6-methylsulfanyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.65$; $[M+H]^+ = 423.2$
1.148	(2-Fluoro-6-[1,2,3]triazol-2-yl-phenyl)-[(S)-2-(6-methylsulfanyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.72$; $[M+H]^+ = 439.2$
1.149	(5-Chloro-2-[1,2,3]triazol-2-yl-phenyl)-[(S)-2-(6-methylsulfanyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.66$; $[M+H]^+ = 435.2$
1.150	(5-Methoxy-2-[1,2,3]triazol-2-yl-phenyl)-[(S)-2-(6-methylsulfanyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.77$; $[M+H]^+ = 473.2$
1.151	[(S)-2-(6-Methylsulfanyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-(2-[1,2,3]triazol-2-yl-5-trifluoromethyl-phenyl)-methanone. LC-MS: $t_R = 0.65$; $[M+H]^+ = 430.2$
1.152	3-[(S)-2-(6-Methylsulfanyl-1H-benzimidazol-2-yl)-pyrrolidine-1-carbonyl]-4-[1,2,3]triazol-2-yl-benzonitrile. LC-MS: $t_R = 0.69$; $[M+H]^+ = 419.2$
1.153	[(S)-2-(6-Methylsulfanyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-(4-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.69$; $[M+H]^+ = 419.2$
1.154	[(S)-2-(6-Methylsulfanyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-(2-methyl-6-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.63$; $[M+H]^+ = 465.2$
1.155	(4,5-Dimethoxy-2-[1,2,3]triazol-2-yl-phenyl)-[(S)-2-(6-methylsulfanyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.61$; $[M+H]^+ = 407.2$
1.156	(5-Fluoro-2-[1,2,3]triazol-2-yl-phenyl)-[(S)-2-(6-methoxy-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.66$; $[M+H]^+ = 423.1$
1.157	(5-Chloro-2-[1,2,3]triazol-2-yl-phenyl)-[(S)-2-(6-methoxy-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.61$; $[M+H]^+ = 419.2$
1.158	[(S)-2-(6-Methoxy-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-(5-methoxy-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.59$; $[M+H]^+ = 414.2$
1.159	3-[(S)-2-(6-Methoxy-1H-benzimidazol-2-yl)-pyrrolidine-1-carbonyl]-4-[1,2,3]triazol-2-yl-benzonitrile. LC-MS: $t_R = 0.64$; $[M+H]^+ = 403.2$
1.160	[(S)-2-(6-Methoxy-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-(4-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.63$; $[M+H]^+ = 403.2$
1.161	[(S)-2-(6-Methoxy-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-(2-methyl-6-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.58$; $[M+H]^+ = 449.2$

1.162	(4,5-Dimethoxy-2-[1,2,3]triazol-2-yl-phenyl)-[(S)-2-(6-methoxy-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.67$; $[M+H]^+ = 423.2$
1.163	(5-Fluoro-2-[1,2,3]triazol-2-yl-phenyl)-[(S)-2-(7-methylsulfanyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.66$; $[M+H]^+ = 423.2$
1.164	(2-Fluoro-6-[1,2,3]triazol-2-yl-phenyl)-[(S)-2-(7-methylsulfanyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.73$; $[M+H]^+ = 439.1$
1.165	(5-Chloro-2-[1,2,3]triazol-2-yl-phenyl)-[(S)-2-(7-methylsulfanyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.66$; $[M+H]^+ = 435.2$
1.166	(5-Methoxy-2-[1,2,3]triazol-2-yl-phenyl)-[(S)-2-(7-methylsulfanyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.66$; $[M+H]^+ = 430.1$
1.167	3-[(S)-2-(7-Methylsulfanyl-1H-benzoimidazol-2-yl)-pyrrolidine-1-carbonyl]-4-[1,2,3]triazol-2-yl-benzonitrile. LC-MS: $t_R = 0.68$; $[M+H]^+ = 453.2$
1.168	(2-Fluoro-3-methoxy-6-[1,2,3]triazol-2-yl-phenyl)-[(S)-2-(7-methylsulfanyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.69$; $[M+H]^+ = 419.2$
1.169	[(S)-2-(7-Methylsulfanyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-(4-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.70$; $[M+H]^+ = 419.3$
1.170	[(S)-2-(7-Methylsulfanyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-(2-methyl-6-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.63$; $[M+H]^+ = 402.2$
1.171	[(S)-2-(5-Methoxy-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-(5-methyl-2-pyrazol-1-yl-phenyl)-methanone. LC-MS: $t_R = 0.65$; $[M+H]^+ = 421.2$
1.172	(2-Fluoro-3-methyl-6-[1,2,3]triazol-2-yl-phenyl)-[(S)-2-(5-methoxy-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.59$; $[M+H]^+ = 404.2$
1.173	[(S)-2-(5-Methoxy-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-(6-methyl-3-[1,2,3]triazol-2-yl-pyridin-2-yl)-methanone. LC-MS: $t_R = 0.71$; $[M+H]^+ = 437.2$
1.174	(2-Fluoro-3-methyl-6-[1,2,3]triazol-2-yl-phenyl)-[(S)-2-(4-methylsulfanyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.64$; $[M+H]^+ = 420.2$
1.175	[(S)-2-(4-Methylsulfanyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-(6-methyl-3-[1,2,3]triazol-2-yl-pyridin-2-yl)-methanone. LC-MS: $t_R = 0.64$; $[M+H]^+ = 404.2$
1.176	[(S)-2-(5-Methylsulfanyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-(2-pyrazol-1-yl-phenyl)-methanone. LC-MS: $t_R = 0.69$; $[M+H]^+ = 418.2$
1.177	(5-Methyl-2-pyrazol-1-yl-phenyl)-[(S)-2-(5-methylsulfanyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.64$; $[M+H]^+ = 419.2$

1.178	(6-Methyl-3-pyrazol-1-yl-pyridin-2-yl)-[(S)-2-(5-methylsulfanyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.71$; $[M+H]^+ = 437.2$
1.179	(2-Fluoro-3-methyl-6-[1,2,3]triazol-2-yl-phenyl)-[(S)-2-(5-methylsulfanyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.65$; $[M+H]^+ = 420.2$
1.180	[(S)-2-(5-Methylsulfanyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-(6-methyl-3-[1,2,3]triazol-2-yl-pyridin-2-yl)-methanone. LC-MS: $t_R = 0.72$; $[M+H]^+ = 421.2$
2.1	[(2S,4S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-4-fluoro-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.75$; $[M+H]^+ = 439.2$
2.2	[(2S,4S)-4-Fluoro-2-(4-methyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-(2-methyl-5-phenyl-thiazol-4-yl)-methanone. LC-MS: $t_R = 0.72$; $[M+H]^+ = 421.2$
2.3	[(2S,4R)-4-Fluoro-2-(4-methyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-(2-methyl-5-phenyl-thiazol-4-yl)-methanone. LC-MS: $t_R = 0.73$; $[M+H]^+ = 421.2$
2.4	[(S)-4,4-Difluoro-2-(4-methyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-(2-methyl-5-phenyl-thiazol-4-yl)-methanone. LC-MS: $t_R = 0.78$; $[M+H]^+ = 439.2$
2.5	[(2S,4R)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-4-fluoro-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.76$; $[M+H]^+ = 439.2$
2.6	[(S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-4,4-difluoro-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.88$; $[M+H]^+ = 457.2$
2.7	[(2S,4S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-4-fluoro-pyrrolidin-1-yl]-(2-methyl-5-phenyl-thiazol-4-yl)-methanone. LC-MS: $t_R = 0.81$; $[M+H]^+ = 455.2$
2.8	[(S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-4,4-difluoro-pyrrolidin-1-yl]-(2-methyl-5-phenyl-thiazol-4-yl)-methanone. LC-MS: $t_R = 0.94$; $[M+H]^+ = 473.2$
2.9	[5-(3-Chloro-phenyl)-2-methyl-thiazol-4-yl]-[(2S,4R)-4-fluoro-2-(4-methyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.80$; $[M+H]^+ = 455.2$
2.10	[(2S,4R)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-4-fluoro-pyrrolidin-1-yl]-[5-(3-chloro-phenyl)-2-methyl-thiazol-4-yl]-methanone. LC-MS: $t_R = 0.91$; $[M+H]^+ = 489.1$
2.11	[(2S,4R)-4-Fluoro-2-(4-methyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.63$; $[M+H]^+ = 405.2$
3.1	[(2S,4S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-4-methoxy-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.75$; $[M+H]^+ = 451.2$
3.2	[(2S,4R)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-4-methoxy-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.74$; $[M+H]^+ = 451.3$

3.3	[(2S,4S)-4-Methoxy-2-(4-methyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-(2-methyl-5-phenyl-thiazol-4-yl)-methanone. LC-MS: $t_R = 0.73$; $[M+H]^+ = 433.2$
3.4	[(2S,4R)-4-Methoxy-2-(4-methyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-(2-methyl-5-phenyl-thiazol-4-yl)-methanone. LC-MS: $t_R = 0.73$; $[M+H]^+ = 433.2$
3.5	[(2S,4S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-4-methoxy-pyrrolidin-1-yl]-(2-methyl-5-phenyl-thiazol-4-yl)-methanone. LC-MS: $t_R = 0.82$; $[M+H]^+ = 467.2$
3.6	[(2S,4R)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-4-methoxy-pyrrolidin-1-yl]-(2-methyl-5-phenyl-thiazol-4-yl)-methanone. LC-MS: $t_R = 0.82$; $[M+H]^+ = 467.2$
3.7	[5-(3-Chloro-phenyl)-2-methyl-thiazol-4-yl]-[(2S,4R)-4-methoxy-2-(4-methyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.80$; $[M+H]^+ = 467.2$
3.8	[(2S,4S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-4-methoxy-pyrrolidin-1-yl]-[5-(3-chloro-phenyl)-2-methyl-thiazol-4-yl]-methanone. LC-MS: $t_R = 0.88$; $[M+H]^+ = 501.2$
3.9	[(2S,4R)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-4-methoxy-pyrrolidin-1-yl]-[5-(3-chloro-phenyl)-2-methyl-thiazol-4-yl]-methanone. LC-MS: $t_R = 0.89$; $[M+H]^+ = 501.1$
3.10	[(2S,4R)-2-(6-Chloro-7-methyl-1H-benzimidazol-2-yl)-4-methoxy-pyrrolidin-1-yl]-(5-chloro-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.79$; $[M+H]^+ = 471.2$
3.11	[(2S,4R)-2-(6-Chloro-7-methyl-1H-benzimidazol-2-yl)-4-methoxy-pyrrolidin-1-yl]-(2-fluoro-3-methoxy-6-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.74$; $[M+H]^+ = 485.2$
3.12	[(2S,4R)-2-(6-Chloro-7-methyl-1H-benzimidazol-2-yl)-4-methoxy-pyrrolidin-1-yl]-(5-methoxy-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.72$; $[M+H]^+ = 467.2$
3.13	[(2S,4R)-2-(6-Chloro-7-methyl-1H-benzimidazol-2-yl)-4-methoxy-pyrrolidin-1-yl]-(2-fluoro-3-methyl-6-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.72$; $[M+H]^+ = 469.3$
3.14	[(2S,4R)-2-(6-Chloro-7-methyl-1H-benzimidazol-2-yl)-4-methoxy-pyrrolidin-1-yl]-(2-methyl-6-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.75$; $[M+H]^+ = 451.2$
3.15	[(2S,4R)-2-(6-Chloro-7-methyl-1H-benzimidazol-2-yl)-4-methoxy-pyrrolidin-1-yl]-(4-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.75$; $[M+H]^+ = 451.2$
3.16	[(2S,4R)-2-(6-Chloro-7-methyl-1H-benzimidazol-2-yl)-4-methoxy-pyrrolidin-1-yl]-(4,5-dimethyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.77$; $[M+H]^+ = 465.2$
3.17	[(2S,4R)-2-(6-Chloro-7-methyl-1H-benzimidazol-2-yl)-4-methoxy-pyrrolidin-1-yl]-(2-[1,2,3]triazol-2-yl-5-trifluoromethyl-phenyl)-methanone. LC-MS: $t_R = 0.84$; $[M+H]^+ = 505.2$
3.18	[(2S,4R)-2-(6-Chloro-7-methyl-1H-benzimidazol-2-yl)-4-methoxy-pyrrolidin-1-yl]-(5-fluoro-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.74$; $[M+H]^+ = 455.2$

3.19	(5-Chloro-2-[1,2,3]triazol-2-yl-phenyl)-[(2S,4R)-4-methoxy-2-(6-methoxy-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.66$; $[M+H]^+ = 453.2$
3.20	[(2S,4R)-4-Methoxy-2-(6-methoxy-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-(5-methoxy-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.60$; $[M+H]^+ = 449.2$
3.21	[(2S,4R)-4-Methoxy-2-(6-methoxy-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-(4-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.63$; $[M+H]^+ = 433.3$
3.22	(4,5-Dimethyl-2-[1,2,3]triazol-2-yl-phenyl)-[(2S,4R)-4-methoxy-2-(6-methoxy-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.67$; $[M+H]^+ = 447.2$
3.23	(5-Fluoro-2-[1,2,3]triazol-2-yl-phenyl)-[(2S,4R)-4-methoxy-2-(6-methoxy-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.61$; $[M+H]^+ = 437.2$
3.24	[(2RS)-(4R)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-4-methoxy-2-methyl-pyrrolidin-1-yl]-(2-methyl-5-phenyl-thiazol-4-yl)-methanone. LC-MS: $t_R = 0.84$; $[M+H]^+ = 481.2$
3.25	[(2RS)-(4R)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-4-methoxy-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.79$; $[M+H]^+ = 465.2$
3.26	[(2RS)-(4R)-4-Methoxy-2-methyl-2-(4-methyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.72$; $[M+H]^+ = 431.2$
3.27	[(2RS)-(4R)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-4-methoxy-2-methyl-pyrrolidin-1-yl]-[5-(3-chloro-phenyl)-2-methyl-thiazol-4-yl]-methanone. LC-MS: $t_R = 0.90$; $[M+H]^+ = 515.2$
3.28	[5-(3-Chloro-phenyl)-2-methyl-thiazol-4-yl]-[(2RS)-(4R)-4-methoxy-2-methyl-2-(4-methyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.82$; $[M+H]^+ = 481.2$
3.29	[(2RS)-(4R)-4-Methoxy-2-methyl-2-(4-methyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-(2-methyl-5-phenyl-thiazol-4-yl)-methanone. LC-MS: $t_R = 0.76$; $[M+H]^+ = 447.2$
3.30	[(2S,4R)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-4-methoxy-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.82$; $[M+H]^+ = 465.3$
3.31	[(2S,4R)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-4-methoxy-2-methyl-pyrrolidin-1-yl]-[5-(3-chloro-phenyl)-2-methyl-thiazol-4-yl]-methanone. LC-MS: $t_R = 0.95$; $[M+H]^+ = 515.2$
3.32	[(2S,4R)-2-(6-Chloro-7-methyl-1H-benzimidazol-2-yl)-4-methoxy-pyrrolidin-1-yl]-(2-methyl-5-m-tolyl-thiazol-4-yl)-methanone. LC-MS: $t_R = 0.90$; $[M+H]^+ = 481.2$
3.33	[(2S,4R)-2-(6-Chloro-7-methyl-1H-benzimidazol-2-yl)-4-methoxy-pyrrolidin-1-yl]-[2-dimethylamino-5-(3,4-dimethyl-phenyl)-thiazol-4-yl]-methanone. LC-MS: $t_R = 0.97$; $[M+H]^+ = 524.3$
3.34	[(2S,4R)-2-(6-Chloro-7-methyl-1H-benzimidazol-2-yl)-4-methoxy-pyrrolidin-1-yl]-[5-(3,4-dimethyl-phenyl)-2-methyl-thiazol-4-yl]-methanone. LC-MS: $t_R = 0.95$; $[M+H]^+ = 495.3$

3.35	[(2S,4R)-2-(6-Chloro-7-methyl-1H-benzoimidazol-2-yl)-4-methoxy-pyrrolidin-1-yl]-[5-(4-chloro-phenyl)-2-methyl-thiazol-4-yl]-methanone. LC-MS: $t_R = 0.94$; $[M+H]^+ = 501.2$
3.36	[(2S,4R)-2-(6-Chloro-7-methyl-1H-benzoimidazol-2-yl)-4-methoxy-pyrrolidin-1-yl]-(2-methyl-5-p-tolyl-thiazol-4-yl)-methanone. LC-MS: $t_R = 0.91$; $[M+H]^+ = 481.2$
3.37	[(2S,4R)-2-(6-Chloro-7-methyl-1H-benzoimidazol-2-yl)-4-methoxy-pyrrolidin-1-yl]-[5-(4-methoxy-phenyl)-2-methyl-thiazol-4-yl]-methanone. LC-MS: $t_R = 0.86$; $[M+H]^+ = 497.2$
3.38	[(2S,4R)-2-(6-Chloro-7-methyl-1H-benzoimidazol-2-yl)-4-methoxy-pyrrolidin-1-yl]-(3',4'-dimethyl-biphenyl-2-yl)-methanone. LC-MS: $t_R = 0.97$; $[M+H]^+ = 474.3$
3.39	Biphenyl-2-yl-[(2S,4R)-2-(6-chloro-7-methyl-1H-benzoimidazol-2-yl)-4-methoxy-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.87$; $[M+H]^+ = 446.3$
3.40	[(2S,4R)-2-(6-Chloro-7-methyl-1H-benzoimidazol-2-yl)-4-methoxy-pyrrolidin-1-yl]-(4'-fluoro-3'-methyl-biphenyl-2-yl)-methanone. LC-MS: $t_R = 0.94$; $[M+H]^+ = 478.3$
3.41	[(2S,4R)-2-(6-Chloro-7-methyl-1H-benzoimidazol-2-yl)-4-methoxy-pyrrolidin-1-yl]-[5-(3-chloro-phenyl)-2-dimethylamino-thiazol-4-yl]-methanone. LC-MS: $t_R = 0.95$; $[M+H]^+ = 530.3$
3.42	[(2RS)-(4R)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-4-methoxy-2-methyl-pyrrolidin-1-yl]-(2-fluoro-6-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.80$; $[M+H]^+ = 469.2$
3.43	[(2RS)-(4R)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-4-methoxy-2-methyl-pyrrolidin-1-yl]-(5-fluoro-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.80$; $[M+H]^+ = 469.3$
3.44	[(2RS)-(4R)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-4-methoxy-2-methyl-pyrrolidin-1-yl]-(5-chloro-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.86$; $[M+H]^+ = 485.2$
3.45	[(2RS)-(4R)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-4-methoxy-2-methyl-pyrrolidin-1-yl]-(2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.76$; $[M+H]^+ = 451.3$
3.46	[(2RS)-(4R)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-4-methoxy-2-methyl-pyrrolidin-1-yl]-(2-fluoro-3-methoxy-6-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.81$; $[M+H]^+ = 499.2$
3.47	[(2RS)-(4R)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-4-methoxy-2-methyl-pyrrolidin-1-yl]-(2-fluoro-3-methyl-6-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.85$; $[M+H]^+ = 483.3$
3.48	[(2RS)-(4R)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-4-methoxy-2-methyl-pyrrolidin-1-yl]-(4,5-dimethyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.87$; $[M+H]^+ = 479.3$
3.49	[(2RS)-(4R)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-4-methoxy-2-methyl-pyrrolidin-1-yl]-(4-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.83$; $[M+H]^+ = 465.3$
3.50	[(2RS)-(4R)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-4-methoxy-2-methyl-pyrrolidin-1-yl]-(2-[1,2,3]triazol-2-yl-5-trifluoromethyl-phenyl)-methanone. LC-MS: $t_R = 0.90$; $[M+H]^+ = 519.3$

3.51	[(2RS)-(4R)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-4-methoxy-2-methyl-pyrrolidin-1-yl]-(2-[1,2,3]triazol-2-yl-5-trifluoromethoxy-phenyl)-methanone. LC-MS: $t_R = 0.93$; $[M+H]^+ = 535.2$
3.52	[(2RS)-(4R)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-4-methoxy-2-methyl-pyrrolidin-1-yl]-(4,5-dimethoxy-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.77$; $[M+H]^+ = 511.3$
3.53	[(2RS)-(4R)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-4-methoxy-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-pyrazol-1-yl-phenyl)-methanone. LC-MS: $t_R = 0.83$; $[M+H]^+ = 464.3$
3.54	[(2RS)-(4R)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-4-methoxy-2-methyl-pyrrolidin-1-yl]-(2-pyrazol-1-yl-phenyl)-methanone. LC-MS: $t_R = 0.77$; $[M+H]^+ = 450.3$
3.55	[(2RS)-(4R)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-4-methoxy-2-methyl-pyrrolidin-1-yl]-(6-methyl-3-pyrazol-1-yl-pyridin-2-yl)-methanone. LC-MS: $t_R = 0.74$; $[M+H]^+ = 465.3$
3.56	[(2RS)-(4R)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-4-methoxy-2-methyl-pyrrolidin-1-yl]-(6-methyl-3-[1,2,3]triazol-2-yl-pyridin-2-yl)-methanone. LC-MS: $t_R = 0.74$; $[M+H]^+ = 466.3$
4.1	[(2S,3S)-3-Methyl-2-(4-methyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.66$; $[M+H]^+ = 401.3$
4.2	[(2S,3S)-3-Methyl-2-(5-methyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.68$; $[M+H]^+ = 401.2$
4.3	[5-(3-Chloro-phenyl)-2-methyl-thiazol-4-yl]-[(2S,3S)-3-methyl-2-(5-methyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.81$; $[M+H]^+ = 451.2$
4.4	[(2S,3S)-3-Methyl-2-(4-methyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-(2-methyl-5-phenyl-thiazol-4-yl)-methanone. LC-MS: $t_R = 0.74$; $[M+H]^+ = 417.2$
4.5	[(2S,3S)-3-Methyl-2-(5-methyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-(2-methyl-5-phenyl-thiazol-4-yl)-methanone. LC-MS: $t_R = 0.75$; $[M+H]^+ = 417.2$
4.6	[(2S,3S)-2-(1H-Benzoimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-[5-(3-chloro-phenyl)-2-methyl-thiazol-4-yl]-methanone. LC-MS: $t_R = 0.77$; $[M+H]^+ = 437.2$
4.7	[5-(3-Chloro-phenyl)-2-methyl-thiazol-4-yl]-[(2S,3S)-3-methyl-2-(4-methyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.81$; $[M+H]^+ = 451.2$
4.8	2-[(2S,3S)-1-[5-(3-Chloro-phenyl)-2-methyl-thiazole-4-carbonyl]-3-methyl-pyrrolidin-2-yl]-1H-benzoimidazole-5-carbonitrile. LC-MS: $t_R = 0.88$; $[M+H]^+ = 462.2$
4.9	[5-(3-Chloro-phenyl)-2-methyl-thiazol-4-yl]-[(2S,3S)-3-methyl-2-(5-trifluoromethyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.97$; $[M+H]^+ = 505.2$
4.10	[5-(3-Chloro-phenyl)-2-methyl-thiazol-4-yl]-[(2S,3S)-2-(5,6-dimethyl-1H-benzoimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.84$; $[M+H]^+ = 465.2$

4.11	[(2S,3S)-2-(5-Chloro-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-[5-(3-chloro-phenyl)-2-methyl-thiazol-4-yl]-methanone. LC-MS: $t_R = 0.87$; $[M+H]^+ = 471.2$
4.12	[5-(3-Chloro-phenyl)-2-methyl-thiazol-4-yl]-[(2S,3S)-2-(5,6-difluoro-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.88$; $[M+H]^+ = 473.1$
4.13	2-[(2S,3S)-1-[5-(3-Chloro-phenyl)-2-methyl-thiazole-4-carbonyl]-3-methyl-pyrrolidin-2-yl]-1H-benzimidazole-4-carboxylic acid methyl ester. LC-MS: $t_R = 0.88$; $[M+H]^+ = 495.2$
4.14	[5-(3-Chloro-phenyl)-2-methyl-thiazol-4-yl]-[(2S,3S)-2-(4-methoxy-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.79$; $[M+H]^+ = 467.2$
4.15	[(2S,3S)-2-(5-Chloro-6-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-[5-(3-chloro-phenyl)-2-methyl-thiazol-4-yl]-methanone. LC-MS: $t_R = 0.89$; $[M+H]^+ = 485.2$
4.16	[5-(3-Chloro-phenyl)-2-methyl-thiazol-4-yl]-[(2S,3S)-2-(4,6-dimethyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.85$; $[M+H]^+ = 465.2$
4.17	[5-(3-Chloro-phenyl)-2-methyl-thiazol-4-yl]-[(2S,3S)-2-(4-hydroxymethyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.73$; $[M+H]^+ = 467.2$
4.18	[5-(3-Chloro-phenyl)-2-methyl-thiazol-4-yl]-[(2S,3S)-2-(4-fluoro-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.87$; $[M+H]^+ = 455.1$
4.19	[5-(3-Chloro-phenyl)-2-methyl-thiazol-4-yl]-[(2S,3S)-3-methyl-2-(4-trifluoromethyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 1.01$; $[M+H]^+ = 505.2$
4.20	[5-(3-Chloro-phenyl)-2-methyl-thiazol-4-yl]-[(2S,3S)-2-(5-fluoro-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.81$; $[M+H]^+ = 455.1$
4.21	[(2S,3S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-[5-(3-chloro-phenyl)-2-methyl-thiazol-4-yl]-methanone. LC-MS: $t_R = 0.9$; $[M+H]^+ = 485.1$
4.22	[(2S,3S)-2-(4-Chloro-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-[5-(3-chloro-phenyl)-2-methyl-thiazol-4-yl]-methanone. LC-MS: $t_R = 0.91$; $[M+H]^+ = 471.1$
4.23	[5-(3-Chloro-phenyl)-2-methyl-thiazol-4-yl]-[(2S,3S)-2-(4,5-difluoro-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.96$; $[M+H]^+ = 473.2$
4.24	[5-(3-Chloro-phenyl)-2-methyl-thiazol-4-yl]-[(2S,3S)-2-(4,5-dimethyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.84$; $[M+H]^+ = 465.2$
4.25	[5-(3-Chloro-phenyl)-2-methyl-thiazol-4-yl]-[(2S,3S)-2-(4-isopropyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.89$; $[M+H]^+ = 479.2$
4.26	[5-(3-Chloro-phenyl)-2-methyl-thiazol-4-yl]-[(2S,3S)-2-(5-methoxy-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.79$; $[M+H]^+ = 467.2$

4.27	[5-(3-Chloro-phenyl)-2-methyl-thiazol-4-yl]-[(2S,3S)-2-(5-hydroxymethyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.71$; $[M+H]^+ = 467.2$
4.28	[(2S,3S)-2-(5-Chloro-7-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-[5-(3-chloro-phenyl)-2-methyl-thiazol-4-yl]-methanone. LC-MS: $t_R = 0.91$; $[M+H]^+ = 485.1$
4.29	(5-Methyl-2-[1,2,3]triazol-2-yl-phenyl)-[(2S,3S)-3-methyl-2-(5-trifluoromethyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.85$; $[M+H]^+ = 455.2$
4.30	[(2S,3S)-2-(5,6-Dimethyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.72$; $[M+H]^+ = 415.3$
4.31	[(2S,3S)-2-(4-Methoxy-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.66$; $[M+H]^+ = 417.2$
4.32	[(2S,3S)-2-(5-Chloro-6-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.77$; $[M+H]^+ = 435.2$
4.33	[(2S,3S)-2-(4,6-Dimethyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.71$; $[M+H]^+ = 415.3$
4.34	(5-Methyl-2-[1,2,3]triazol-2-yl-phenyl)-[(2S,3S)-3-methyl-2-(4-trifluoromethyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.89$; $[M+H]^+ = 455.2$
4.35	[(2S,3S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.76$; $[M+H]^+ = 435.2$
4.36	[(2S,3S)-2-(4-Chloro-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.77$; $[M+H]^+ = 421.2$
4.37	[(2S,3S)-2-(4,5-Dimethyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.7$; $[M+H]^+ = 415.2$
4.38	[(2S,3S)-2-(4-Isopropyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.75$; $[M+H]^+ = 429.3$
4.39	[(2S,3S)-2-(5-Chloro-7-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.77$; $[M+H]^+ = 435.2$
4.40	(5-Chloro-2-[1,2,3]triazol-2-yl-phenyl)-[(2S,3S)-2-(5,6-dimethoxy-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.71$; $[M+H]^+ = 467.2$
4.41	[(2S,3S)-2-(5,6-Dimethoxy-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(5-fluoro-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.64$; $[M+H]^+ = 451.3$
4.42	[(2S,3S)-2-(5,6-Dimethoxy-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(2-fluoro-6-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.63$; $[M+H]^+ = 451.3$

4.43	[(2S,3S)-2-(5,6-Dimethoxy-1H-benzoimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(5-methoxy-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: t_R = 0.64; $[M+H]^+$ = 463.3
4.44	[(2S,3S)-2-(5,6-Dimethoxy-1H-benzoimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(2-fluoro-3-methyl-6-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: t_R = 0.69; $[M+H]^+$ = 465.3
4.45	[(2S,3S)-2-(5,6-Dimethoxy-1H-benzoimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(2-fluoro-3-methoxy-6-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: t_R = 0.65; $[M+H]^+$ = 481.3
4.46	[(2S,3S)-2-(5,6-Dimethoxy-1H-benzoimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(4-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: t_R = 0.67; $[M+H]^+$ = 447.3
4.47	[(2S,3S)-2-(5,6-Dimethoxy-1H-benzoimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(2-[1,2,3]triazol-2-yl-5-trifluoromethyl-phenyl)-methanone. LC-MS: t_R = 0.76; $[M+H]^+$ = 501.3
4.48	[(2S,3S)-2-(5,6-Dimethoxy-1H-benzoimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(2-methyl-6-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: t_R = 0.66; $[M+H]^+$ = 447.3
4.49	[(2S,3S)-2-(5,6-Dimethoxy-1H-benzoimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: t_R = 0.62; $[M+H]^+$ = 433.3
4.50	[(2S,3S)-2-(5,6-Dimethoxy-1H-benzoimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(2-methyl-5-(4-trifluoromethyl-phenyl)-thiazol-4-yl)-methanone. LC-MS: t_R = 0.83; $[M+H]^+$ = 531.3
4.51	[(2S,3S)-2-(5,6-Dimethoxy-1H-benzoimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(2-methyl-4-p-tolyl-thiazol-5-yl)-methanone. LC-MS: t_R = 0.7; $[M+H]^+$ = 477.2
4.52	[5-(3,5-Difluoro-phenyl)-2-methyl-thiazol-4-yl]-[(2S,3S)-2-(5,6-dimethoxy-1H-benzoimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-methanone. LC-MS: t_R = 0.75; $[M+H]^+$ = 499.3
4.53	[(2S,3S)-2-(5,6-Dimethoxy-1H-benzoimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-[5-(6-methoxy-pyridin-3-yl)-2-methyl-thiazol-4-yl]-methanone. LC-MS: t_R = 0.68; $[M+H]^+$ = 494.4
4.54	[(2S,3S)-2-(5,6-Dimethoxy-1H-benzoimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-[5-(4-fluoro-phenyl)-2-methyl-thiazol-4-yl]-methanone. LC-MS: t_R = 0.73; $[M+H]^+$ = 481.3
4.55	[5-(3-Chloro-phenyl)-thiazol-4-yl]-[(2S,3S)-2-(5,6-dimethoxy-1H-benzoimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-methanone. LC-MS: t_R = 0.72; $[M+H]^+$ = 483.2
4.56	[(2S,3S)-2-(5,6-Dimethoxy-1H-benzoimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(2-pyrazol-1-yl-phenyl)-methanone. LC-MS: t_R = 0.62; $[M+H]^+$ = 432.3
4.57	[(2S,3S)-2-(5,6-Dimethoxy-1H-benzoimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(5-methyl-2-pyrazol-1-yl-phenyl)-methanone. LC-MS: t_R = 0.66; $[M+H]^+$ = 446.3
4.58	[(2S,3S)-2-(5,6-Dimethoxy-1H-benzoimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(4'-methyl-biphenyl-2-yl)-methanone. LC-MS: t_R = 0.80; $[M+H]^+$ = 456.4

4.59	[(2S,3S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.74$; $[M+H]^+ = 421.3$
4.60	[(2S,3S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(5-chloro-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.84$; $[M+H]^+ = 455.2$
4.61	[(2S,3S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(4'-methyl-biphenyl-2-yl)-methanone. LC-MS: $t_R = 0.94$; $[M+H]^+ = 444.3$
4.62	[(2S,3S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(4-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.79$; $[M+H]^+ = 435.3$
4.63	[(2S,3S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(5-(4-chloro-phenyl)-2-methyl-thiazol-4-yl)-methanone. LC-MS: $t_R = 0.94$; $[M+H]^+ = 485.2$
4.64	[(2S,3S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(5-methyl-2-pyrazol-1-yl-phenyl)-methanone. LC-MS: $t_R = 0.80$; $[M+H]^+ = 434.3$
4.65	[(2S,3S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(2-methyl-4-p-tolyl-thiazol-5-yl)-methanone. LC-MS: $t_R = 0.88$; $[M+H]^+ = 465.2$
4.66	[(2S,3S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(5-methoxy-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.77$; $[M+H]^+ = 451.3$
4.67	[(2S,3S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(5-(3,5-difluoro-phenyl)-2-methyl-thiazol-4-yl)-methanone. LC-MS: $t_R = 0.92$; $[M+H]^+ = 487.2$
4.68	[(2S,3S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(2-fluoro-3-methyl-6-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.83$; $[M+H]^+ = 453.2$
4.69	[(2S,3S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(5-(6-methoxy-pyridin-3-yl)-2-methyl-thiazol-4-yl)-methanone. LC-MS: $t_R = 0.83$; $[M+H]^+ = 482.2$
4.70	[(2S,3S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(5-(4-fluoro-phenyl)-2-methyl-thiazol-4-yl)-methanone. LC-MS: $t_R = 0.89$; $[M+H]^+ = 469.2$
4.71	[(2S,3S)-2-(5-Chloro-6-methyl-1H-benzoimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(5-chloro-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.83$; $[M+H]^+ = 455.2$
4.72	[(2S,3S)-2-(5-Chloro-6-methyl-1H-benzoimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(4-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.80$; $[M+H]^+ = 435.3$
4.73	[(2S,3S)-2-(5-Chloro-6-methyl-1H-benzoimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(5-methyl-2-pyrazol-1-yl-phenyl)-methanone. LC-MS: $t_R = 0.80$; $[M+H]^+ = 434.3$
4.74	[(2S,3S)-2-(5-Chloro-6-methyl-1H-benzoimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(5-methoxy-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.78$; $[M+H]^+ = 451.9$

4.75	[(2S,3S)-2-(5-Chloro-6-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(2-fluoro-3-methyl-6-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.83$; $[M+H]^+ = 453.3$
4.76	[(2S,3S)-2-(5-Chloro-6-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.74$; $[M+H]^+ = 421.2$
4.77	[(2S,3S)-2-(5-Chloro-6-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(4'-methyl-biphenyl-2-yl)-methanone. LC-MS: $t_R = 0.93$; $[M+H]^+ = 444.3$
4.78	[(2S,3S)-2-(5-Chloro-6-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-[5-(4-chloro-phenyl)-2-methyl-thiazol-4-yl]-methanone. LC-MS: $t_R = 0.93$; $[M+H]^+ = 485.2$
4.79	[(2S,3S)-2-(5-Chloro-6-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(2-methyl-4-p-tolyl-thiazol-5-yl)-methanone. LC-MS: $t_R = 0.86$; $[M+H]^+ = 465.3$
4.80	[(2S,3S)-2-(5-Chloro-6-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-[5-(3,5-difluoro-phenyl)-2-methyl-thiazol-4-yl]-methanone. LC-MS: $t_R = 0.91$; $[M+H]^+ = 487.2$
4.81	(5-Chloro-2-[1,2,3]triazol-2-yl-phenyl)-[(2S,3S)-2-(5,6-dimethyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.78$; $[M+H]^+ = 435.3$
4.82	[(2S,3S)-2-(5,6-Dimethyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(4-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.75$; $[M+H]^+ = 415.3$
4.83	[(2S,3S)-2-(5,6-Dimethyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(5-methyl-2-pyrazol-1-yl-phenyl)-methanone. LC-MS: $t_R = 0.75$; $[M+H]^+ = 414.3$
4.84	[(2S,3S)-2-(5,6-Dimethyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(5-methoxy-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.72$; $[M+H]^+ = 431.3$
4.85	[(2S,3S)-2-(5,6-Dimethyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.69$; $[M+H]^+ = 401.3$
4.86	[(2S,3S)-2-(5,6-Dimethyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(4'-methyl-biphenyl-2-yl)-methanone. LC-MS: $t_R = 0.87$; $[M+H]^+ = 424.3$
4.87	[5-(4-Chloro-phenyl)-2-methyl-thiazol-4-yl]-[(2S,3S)-2-(5,6-dimethyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.87$; $[M+H]^+ = 465.2$
4.88	[(2S,3S)-2-(5,6-Dimethyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(2-methyl-4-p-tolyl-thiazol-5-yl)-methanone. LC-MS: $t_R = 0.78$; $[M+H]^+ = 445.3$
4.89	[5-(3,5-Difluoro-phenyl)-2-methyl-thiazol-4-yl]-[(2S,3S)-2-(5,6-dimethyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.85$; $[M+H]^+ = 467.3$
4.90	[(2S,3S)-2-(5,6-Dimethyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-[5-(4-fluoro-phenyl)-2-methyl-thiazol-4-yl]-methanone. LC-MS: $t_R = 0.82$; $[M+H]^+ = 449.3$

4.91	[(2S,3S)-2-(6-Chloro-4-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(5-chloro-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.84$; $[M+H]^+ = 455.2$
4.92	[(2S,3S)-2-(6-Chloro-4-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(4'-methyl-biphenyl-2-yl)-methanone. LC-MS: $t_R = 0.93$; $[M+H]^+ = 444.3$
4.93	[(2S,3S)-2-(6-Chloro-4-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(4-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.80$; $[M+H]^+ = 435.2$
4.94	[(2S,3S)-2-(6-Chloro-4-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(5-(4-chloro-phenyl)-2-methyl-thiazol-4-yl)-methanone. LC-MS: $t_R = 0.94$; $[M+H]^+ = 485.2$
4.95	[(2S,3S)-2-(6-Chloro-4-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(5-methyl-2-pyrazol-1-yl-phenyl)-methanone. LC-MS: $t_R = 0.80$; $[M+H]^+ = 434.3$
4.96	[(2S,3S)-2-(6-Chloro-4-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(2-methyl-4-p-tolyl-thiazol-5-yl)-methanone. LC-MS: $t_R = 0.88$; $[M+H]^+ = 465.2$
4.97	[(2S,3S)-2-(6-Chloro-4-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(5-methoxy-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.77$; $[M+H]^+ = 451.3$
4.98	[(2S,3S)-2-(6-Chloro-4-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(5-(3,5-difluoro-phenyl)-2-methyl-thiazol-4-yl)-methanone. LC-MS: $t_R = 0.92$; $[M+H]^+ = 487.2$
4.99	[(2S,3S)-2-(6-Chloro-4-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(2-fluoro-3-methyl-6-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.83$; $[M+H]^+ = 453.2$
4.100	[(2S,3S)-2-(6-Chloro-4-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(5-(4-fluoro-phenyl)-2-methyl-thiazol-4-yl)-methanone. LC-MS: $t_R = 0.89$; $[M+H]^+ = 469.2$
4.101	[(2S,3S)-2-(6-Chloro-4-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.74$; $[M+H]^+ = 421.3$
4.102	[(2S,3S)-2-(5-Chloro-6-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(5-methyl-2-pyridin-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.91$; $[M+H]^+ = 445.9$
4.103	(4-Chloro-biphenyl-2-yl)-[(2S,3S)-2-(5-chloro-6-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.99$; $[M+H]^+ = 464.3$
4.104	[(2S,3S)-2-(5-Chloro-6-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(5-methyl-2-pyridin-3-yl-phenyl)-methanone. LC-MS: $t_R = 0.72$; $[M+H]^+ = 445.3$
4.105	[(2S,3S)-2-(5-Chloro-6-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(4-fluoro-biphenyl-2-yl)-methanone. LC-MS: $t_R = 0.94$; $[M+H]^+ = 448.3$
4.106	[(2S,3S)-2-(5-Chloro-6-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(5-fluoro-2-pyridin-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.81$; $[M+H]^+ = 449.3$

4.107	[(2S,3S)-2-(5-Chloro-6-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(2-fluoro-6-pyridin-3-yl-phenyl)-methanone. LC-MS: $t_R = 0.71$; $[M+H]^+ = 449.3$
4.108	2-[(2S,3S)-2-(5-Chloro-6-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidine-1-carbonyl]-biphenyl-4-carbonitrile. LC-MS: $t_R = 0.91$; $[M+H]^+ = 455.3$
4.109	[(2S,3S)-2-(5-Chloro-6-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(4-methyl-2-pyridin-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.81$; $[M+H]^+ = 445.4$
4.110	[(2S,3S)-2-(5-Chloro-6-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(4-methyl-2-pyridin-3-yl-phenyl)-methanone. LC-MS: $t_R = 0.72$; $[M+H]^+ = 445.3$
4.111	[(2S,3S)-2-(5-Chloro-6-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(5-methyl-biphenyl-2-yl)-methanone. LC-MS: $t_R = 0.96$; $[M+H]^+ = 444.3$
4.112	[(2S,3S)-2-(5-Chloro-6-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(3-fluoro-biphenyl-2-yl)-methanone. LC-MS: $t_R = 0.8$; $[M+H]^+ = 449.3$
4.113	[(2S,3S)-2-(5-Chloro-6-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(2-fluoro-3-methyl-6-pyridin-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.83$; $[M+H]^+ = 463.3$
4.114	[(2S,3S)-2-(5-Chloro-6-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(4-trifluoromethyl-biphenyl-2-yl)-methanone. LC-MS: $t_R = 1.02$; $[M+H]^+ = 498.3$
4.115	[(2S,3S)-2-(5-Chloro-6-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(3-fluoro-4-methyl-biphenyl-2-yl)-methanone. LC-MS: $t_R = 0.99$; $[M+H]^+ = 462.3$
4.116	[(2S,3S)-2-(5-Chloro-6-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(4-methyl-biphenyl-2-yl)-methanone. LC-MS: $t_R = 0.95$; $[M+H]^+ = 444.3$
4.117	[(2S,3S)-2-(5-Chloro-6-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(2-fluoro-3-methyl-6-pyridin-3-yl-phenyl)-methanone. LC-MS: $t_R = 0.75$; $[M+H]^+ = 463.3$
4.118	[(2S,3S)-2-(4-Chloro-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-[5-(4-fluoro-phenyl)-2-methyl-thiazol-4-yl]-methanone. LC-MS: $t_R = 0.92$; $[M+H]^+ = 455.2$
4.119	[(2S,3S)-2-(4-Chloro-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-[5-(3,5-difluoro-phenyl)-2-methyl-thiazol-4-yl]-methanone. LC-MS: $t_R = 0.96$; $[M+H]^+ = 473.2$
4.120	[(2S,3S)-2-(4-Chloro-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-[5-(4-chloro-phenyl)-2-methyl-thiazol-4-yl]-methanone. LC-MS: $t_R = 0.98$; $[M+H]^+ = 471.2$
4.121	[(2S,3S)-2-(4-Chloro-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-[5-(3-methoxy-phenyl)-2-methyl-thiazol-4-yl]-methanone. LC-MS: $t_R = 0.9$; $[M+H]^+ = 467.2$
4.122	[(2S,3S)-2-(4-Chloro-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(2-methyl-5-m-tolyl-thiazol-4-yl)-methanone. LC-MS: $t_R = 0.95$; $[M+H]^+ = 451.3$

4.123	[5-(4-Bromo-phenyl)-2-methyl-thiazol-4-yl]-[(2S,3S)-2-(4-chloro-1H-benzoimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 1.0$; $[M+H]^+ = 515.2$
4.124	[(2S,3S)-2-(4-Chloro-1H-benzoimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(2-methyl-5-p-tolyl-thiazol-4-yl)-methanone. LC-MS: $t_R = 0.95$; $[M+H]^+ = 451.3$
4.125	[(2S,3S)-2-(4-Chloro-1H-benzoimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-[5-(3,4-dimethyl-phenyl)-2-methyl-thiazol-4-yl]-methanone. LC-MS: $t_R = 0.99$; $[M+H]^+ = 465.3$
4.126	[(2S,3S)-2-(4-Chloro-1H-benzoimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-[5-(3,4-difluoro-phenyl)-2-methyl-thiazol-4-yl]-methanone. LC-MS: $t_R = 0.95$; $[M+H]^+ = 473.2$
4.127	[(2S,3S)-2-(4-Chloro-1H-benzoimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-[2-(3,4-dimethyl-phenyl)-thiophen-3-yl]-methanone. LC-MS: $t_R = 1.02$; $[M+H]^+ = 450.3$
4.128	[(2S,3S)-2-(4-Chloro-1H-benzoimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-[2-dimethylamino-5-(3,4-dimethyl-phenyl)-thiazol-4-yl]-methanone. LC-MS: $t_R = 1.0$; $[M+H]^+ = 494.3$
4.129	[(2S,3S)-2-(4-Chloro-1H-benzoimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-[5-(2-methoxy-phenyl)-2-methyl-thiazol-4-yl]-methanone. LC-MS: $t_R = 0.89$; $[M+H]^+ = 467.3$
4.130	[(2S,3S)-2-(5-Chloro-6-methyl-1H-benzoimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(4-methyl-2-pyrimidin-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.81$; $[M+H]^+ = 446.3$
4.131	[(2S,3S)-2-(5-Chloro-6-methyl-1H-benzoimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(5-fluoro-2-pyrimidin-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.81$; $[M+H]^+ = 450.3$
4.132	[(2S,3S)-2-(5-Chloro-6-methyl-1H-benzoimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(2-fluoro-3-methyl-6-pyrimidin-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.83$; $[M+H]^+ = 464.3$
4.133	[(2S,3S)-2-(5-Chloro-6-methyl-1H-benzoimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(2-fluoro-6-pyrimidin-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.80$; $[M+H]^+ = 450.3$
4.134	[(2S,3S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-[2-(3-methoxy-phenylethynyl)-phenyl]-methanone. LC-MS: $t_R =$; $[M+H]^+ =$
4.135	[(2S,3S)-2-(5-Chloro-6-methyl-1H-benzoimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-[2-(3-methoxy-phenylethynyl)-phenyl]-methanone. LC-MS: $t_R =$; $[M+H]^+ =$
4.136	[5-(4-Fluoro-phenyl)-2-methyl-thiazol-4-yl]-[(2S,3S)-3-methyl-2-(5-trifluoromethyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.99$; $[M+H]^+ = 489.3$
4.137	[5-(3-Methoxy-phenyl)-2-methyl-thiazol-4-yl]-[(2S,3S)-3-methyl-2-(5-trifluoromethyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.98$; $[M+H]^+ = 501.3$
4.138	(2-Methyl-5-m-tolyl-thiazol-4-yl)-[(2S,3S)-3-methyl-2-(5-trifluoromethyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 1.02$; $[M+H]^+ = 485.3$

4.139	[5-(4-Chloro-phenyl)-2-methyl-thiazol-4-yl]-[(2S,3S)-3-methyl-2-(5-trifluoromethyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 1.05$; $[M+H]^+ = 505.3$
4.140	(2-Methyl-5-p-tolyl-thiazol-4-yl)-[(2S,3S)-3-methyl-2-(5-trifluoromethyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 1.02$; $[M+H]^+ = 485.3$
4.141	[5-(4-Methoxy-phenyl)-2-methyl-thiazol-4-yl]-[(2S,3S)-3-methyl-2-(5-trifluoromethyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.97$; $[M+H]^+ = 501.3$
4.142	[5-(4-Bromo-phenyl)-2-methyl-thiazol-4-yl]-[(2S,3S)-3-methyl-2-(5-trifluoromethyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 1.06$; $[M+H]^+ = 549.2$
4.143	[5-(3,4-Difluoro-phenyl)-2-methyl-thiazol-4-yl]-[(2S,3S)-3-methyl-2-(5-trifluoromethyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 1.02$; $[M+H]^+ = 507.3$
4.144	[2-(3,4-Dimethyl-phenyl)-thiophen-3-yl]-[(2S,3S)-3-methyl-2-(5-trifluoromethyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 1.09$; $[M+H]^+ = 484.3$
4.145	[5-(3,4-Dimethyl-phenyl)-2-methyl-thiazol-4-yl]-[(2S,3S)-3-methyl-2-(5-trifluoromethyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 1.06$; $[M+H]^+ = 499.3$
4.146	[5-(2-Chloro-phenyl)-2-methyl-thiazol-4-yl]-[(2S,3S)-3-methyl-2-(5-trifluoromethyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 1.00$; $[M+H]^+ = 505.2$
4.147	[5-(2-Methoxy-phenyl)-2-methyl-thiazol-4-yl]-[(2S,3S)-3-methyl-2-(5-trifluoromethyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.97$; $[M+H]^+ = 501.3$
4.148	[5-(3,5-Difluoro-phenyl)-2-methyl-thiazol-4-yl]-[(2S,3S)-3-methyl-2-(5-trifluoromethyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 1.02$; $[M+H]^+ = 507.3$
4.149	[5-(4-Fluoro-phenyl)-2-methyl-thiazol-4-yl]-[(2S,3S)-3-methyl-2-(4-trifluoromethyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 1.03$; $[M+H]^+ = 489.3$
4.150	[5-(4-Chloro-phenyl)-2-methyl-thiazol-4-yl]-[(2S,3S)-3-methyl-2-(4-trifluoromethyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 1.09$; $[M+H]^+ = 505.2$
4.151	[5-(4-Bromo-phenyl)-2-methyl-thiazol-4-yl]-[(2S,3S)-3-methyl-2-(4-trifluoromethyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 1.11$; $[M+H]^+ = 549.2$
4.152	[5-(3,4-Dimethyl-phenyl)-2-methyl-thiazol-4-yl]-[(2S,3S)-3-methyl-2-(4-trifluoromethyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 1.1$; $[M+H]^+ = 499.3$
4.153	(2-Methyl-5-p-tolyl-thiazol-4-yl)-[(2S,3S)-3-methyl-2-(4-trifluoromethyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 1.06$; $[M+H]^+ = 485.3$
4.154	[5-(3,4-Difluoro-phenyl)-2-methyl-thiazol-4-yl]-[(2S,3S)-3-methyl-2-(4-trifluoromethyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 1.06$; $[M+H]^+ = 507.3$

4.155	[5-(2-Chloro-phenyl)-2-methyl-thiazol-4-yl]-[(2S,3S)-3-methyl-2-(4-trifluoromethyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 1.04$; $[M+H]^+ = 505.2$
4.156	[5-(2-Methoxy-phenyl)-2-methyl-thiazol-4-yl]-[(2S,3S)-3-methyl-2-(4-trifluoromethyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.99$; $[M+H]^+ = 501.3$
4.157	[5-(3,5-Difluoro-phenyl)-2-methyl-thiazol-4-yl]-[(2S,3S)-3-methyl-2-(4-trifluoromethyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 1.06$; $[M+H]^+ = 507.3$
4.158	(2-Methyl-5-m-tolyl-thiazol-4-yl)-[(2S,3S)-3-methyl-2-(4-trifluoromethyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 1.06$; $[M+H]^+ = 485.3$
4.159	[5-(4-Methoxy-phenyl)-2-methyl-thiazol-4-yl]-[(2S,3S)-3-methyl-2-(4-trifluoromethyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 1.00$; $[M+H]^+ = 501.3$
4.160	[2-(3,4-Dimethyl-phenyl)-thiophen-3-yl]-[(2S,3S)-3-methyl-2-(4-trifluoromethyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 1.14$; $[M+H]^+ = 484.3$
4.161	[(2S,3S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(5-methyl-2-pyridin-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.81$; $[M+H]^+ = 445.3$
4.162	(4-Chloro-biphenyl-2-yl)-[(2S,3S)-2-(5-chloro-4-methyl-1H-benzoimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.99$; $[M+H]^+ = 464.3$
4.163	[(2S,3S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(5-methyl-2-pyridin-3-yl-phenyl)-methanone. LC-MS: $t_R = 0.73$; $[M+H]^+ = 445.4$
4.164	[(2S,3S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(4-fluoro-biphenyl-2-yl)-methanone. LC-MS: $t_R = 0.95$; $[M+H]^+ = 448.3$
4.165	[(2S,3S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(5-fluoro-2-pyridin-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.81$; $[M+H]^+ = 445.3$
4.166	2-[(2S,3S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-3-methyl-pyrrolidine-1-carbonyl]-biphenyl-4-carbonitrile. LC-MS: $t_R = 0.92$; $[M+H]^+ = 455.3$
4.167	[(2S,3S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(4-methyl-2-pyridin-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.81$; $[M+H]^+ = 445.3$
4.168	[(2S,3S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(5-methyl-biphenyl-2-yl)-methanone. LC-MS: $t_R = 0.96$; $[M+H]^+ = 444.3$
4.169	[(2S,3S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(3-fluoro-biphenyl-2-yl)-methanone. LC-MS: $t_R = 0.94$; $[M+H]^+ = 448.3$
4.170	[(2S,3S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(4-trifluoromethyl-biphenyl-2-yl)-methanone. LC-MS: $t_R = 1.03$; $[M+H]^+ = 498.3$

4.171	[(2S,3S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(3-fluoro-4-methyl-biphenyl-2-yl)-methanone. LC-MS: $t_R = 1.00$; $[M+H]^+ = 462.3$
4.172	[(2S,3S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(4-methyl-biphenyl-2-yl)-methanone. LC-MS: $t_R = 0.95$; $[M+H]^+ = 444.3$
4.173	[(2S,3S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(2-fluoro-3-methyl-6-pyridin-3-yl-phenyl)-methanone. LC-MS: $t_R = 0.76$; $[M+H]^+ = 463.3$
4.174	[(2S,3S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(4-methyl-2-pyrimidin-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.81$; $[M+H]^+ = 446.3$
4.175	[(2S,3S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(5-fluoro-2-pyrimidin-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.81$; $[M+H]^+ = 450.3$
4.176	[(2S,3S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(2-fluoro-3-methyl-6-pyrimidin-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.80$; $[M+H]^+ = 464.3$
4.177	[(2S,3S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(2-fluoro-6-pyrimidin-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.80$; $[M+H]^+ = 450.3$
4.178	[(2S,3S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(5-methyl-2-pyrimidin-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.80$; $[M+H]^+ = 446.3$
4.179	[(2S,3S)-2-(5,6-Dimethoxy-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(5-methyl-2-pyrimidin-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.68$; $[M+H]^+ = 458.4$
4.180	[(2S,3S)-2-(5-Chloro-6-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(5-methyl-2-pyrimidin-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.80$; $[M+H]^+ = 446.3$
4.181	[(2S,3S)-2-(6-Chloro-7-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(5-methyl-2-pyrimidin-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.80$; $[M+H]^+ = 446.3$
4.182	[(2S,3S)-2-(5-Chloro-7-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(5-methyl-2-pyrimidin-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.80$; $[M+H]^+ = 446.3$
4.183	[(2S,3S)-2-(5,6-Dimethyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(5-methyl-2-pyrimidin-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.76$; $[M+H]^+ = 426.4$
5.1	[(S)-2-Methyl-2-(4-methyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-(2-methyl-5-phenyl-thiazol-4-yl)-methanone. LC-MS: $t_R = 0.79$; $[M+H]^+ = 417.3$
5.2	[(S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.81$; $[M+H]^+ = 435.3$
5.3	[(S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(2-methyl-5-phenyl-thiazol-4-yl)-methanone. LC-MS: $t_R = 0.83$; $[M+H]^+ = 451.2$

5.4	[5-(3-Chloro-phenyl)-2-methyl-thiazol-4-yl]-[(S)-2-methyl-2-(4-methyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.82$; $[M+H]^+ = 451.2$
5.5	[(S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-[5-(3-chloro-phenyl)-2-methyl-thiazol-4-yl]-methanone. LC-MS: $t_R = 0.9$; $[M+H]^+ = 485.2$
5.6	[(S)-2-Methyl-2-(4-methyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.68$; $[M+H]^+ = 401.2$
5.7	[(S)-2-(5,6-Dichloro-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.92$; $[M+H]^+ = 455.2$
5.8	[(S)-2-(5-Bromo-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.74$; $[M+H]^+ = 465.2$
5.9	(5-Methyl-2-[1,2,3]triazol-2-yl-phenyl)-[(S)-2-methyl-2-(5-trifluoromethyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.83$; $[M+H]^+ = 455.3$
5.10	[(S)-2-(5,6-Dimethyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.75$; $[M+H]^+ = 415.4$
5.11	[(S)-2-(5-Chloro-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.73$; $[M+H]^+ = 421.2$
5.12	[(S)-2-(6-Chloro-5-fluoro-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.81$; $[M+H]^+ = 439.2$
5.13	[(S)-2-(5-tert-Butyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.81$; $[M+H]^+ = 443.3$
5.14	[(S)-2-(4-Chloro-6-trifluoromethyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 1.11$; $[M+H]^+ = 489.2$
5.15	[(S)-2-(5-Chloro-6-methyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.76$; $[M+H]^+ = 435.2$
5.16	[(S)-2-(4,6-Dimethyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.76$; $[M+H]^+ = 415.4$
5.17	(5-Methyl-2-[1,2,3]triazol-2-yl-phenyl)-[(S)-2-methyl-2-(4-trifluoromethyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.96$; $[M+H]^+ = 455.2$
5.18	[(S)-2-(4-Bromo-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.82$; $[M+H]^+ = 465.2$
5.19	[(S)-2-(5-Bromo-7-methyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.82$; $[M+H]^+ = 479.2$

5.20	(5-Methyl-2-[1,2,3]triazol-2-yl-phenyl)-[(S)-2-methyl-2-(5-trifluoromethoxy-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.82$; $[M+H]^+ = 471.2$
5.21	[(S)-2-(5,6-Dimethoxy-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.68$; $[M+H]^+ = 447.4$
5.22	[(S)-2-(6-Fluoro-5-trifluoromethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.95$; $[M+H]^+ = 473.3$
5.23	[(S)-2-(6-Bromo-5-fluoro-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.82$; $[M+H]^+ = 483.2$
5.24	[(S)-2-(6-Chloro-5-trifluoromethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 1.01$; $[M+H]^+ = 489.2$
5.25	[(S)-2-(4-Bromo-6-fluoro-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.99$; $[M+H]^+ = 483.2$
5.26	[(S)-2-(4-Chloro-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.79$; $[M+H]^+ = 421.2$
5.27	[(S)-2-(4,5-Difluoro-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.88$; $[M+H]^+ = 423.3$
5.28	[(S)-2-(4,5-Dimethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.73$; $[M+H]^+ = 415.3$
5.29	[(S)-2-(4-Isopropyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.81$; $[M+H]^+ = 429.4$
5.30	(5-Methyl-2-[1,2,3]triazol-2-yl-phenyl)-[(S)-2-methyl-2-(5-trifluoromethylsulfanyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.89$; $[M+H]^+ = 487.2$
5.31	[(S)-2-(4,7-Dimethoxy-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.72$; $[M+H]^+ = 447.3$
5.32	[(S)-2-(5-Methoxy-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.66$; $[M+H]^+ = 417.2$
5.33	[(S)-2-(7,8-Dihydro-3H-6,9-dioxo-1,3-diaza-cyclopenta[a]naphthalen-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.67$; $[M+H]^+ = 445.2$
5.34	[(S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-chloro-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.84$; $[M+H]^+ = 455.2$
5.35	[(S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(2-[1,2,3]triazol-2-yl-5-trifluoromethoxy-phenyl)-methanone. LC-MS: $t_R = 0.88$; $[M+H]^+ = 505.2$

5.36	[(S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methoxy-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.78$; $[M+H]^+ = 451.3$
5.37	[(S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(2-fluoro-3-methoxy-6-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.79$; $[M+H]^+ = 469.3$
5.38	[(S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(4-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.77$; $[M+H]^+ = 435.2$
5.39	[(S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(4,5-dimethoxy-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.72$; $[M+H]^+ = 481.2$
5.40	[(S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(2-[1,2,3]triazol-2-yl-5-trifluoromethyl-phenyl)-methanone. LC-MS: $t_R = 0.86$; $[M+H]^+ = 489.2$
5.41	[(S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(4,5-dimethyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.82$; $[M+H]^+ = 449.2$
5.42	[(S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(2-pyrazol-1-yl-phenyl)-methanone. LC-MS: $t_R = 0.73$; $[M+H]^+ = 420.2$
5.43	[(S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-pyrazol-1-yl-phenyl)-methanone. LC-MS: $t_R = 0.78$; $[M+H]^+ = 434.2$
5.44	[(S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(6-methyl-3-pyrazol-1-yl-pyridin-2-yl)-methanone. LC-MS: $t_R = 0.71$; $[M+H]^+ = 435.2$
5.45	[(S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-fluoro-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.75$; $[M+H]^+ = 439.1$
5.46	[(S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.72$; $[M+H]^+ = 421.2$
5.47	[(S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(6-methyl-3-[1,2,3]triazol-2-yl-pyridin-2-yl)-methanone. LC-MS: $t_R = 0.71$; $[M+H]^+ = 436.2$
5.48	[(S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(2-fluoro-6-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.77$; $[M+H]^+ = 439.3$
5.49	(5-Chloro-2-[1,2,3]triazol-2-yl-phenyl)-[(S)-2-methyl-2-(4-methyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.71$; $[M+H]^+ = 421.2$
5.50	(5-Methoxy-2-[1,2,3]triazol-2-yl-phenyl)-[(S)-2-methyl-2-(4-methyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.65$; $[M+H]^+ = 417.2$
5.51	(4,5-Dimethyl-2-[1,2,3]triazol-2-yl-phenyl)-[(S)-2-methyl-2-(4-methyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.76$; $[M+H]^+ = 415.4$

5.52	[(S)-2-Methyl-2-(4-methyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-(2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.65$; $[M+H]^+ = 387.3$
5.53	(5-Chloro-2-[1,2,3]triazol-2-yl-phenyl)-[(S)-2-(5,6-dimethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.75$; $[M+H]^+ = 435.2$
5.54	[(S)-2-(5,6-Dimethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methoxy-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.73$; $[M+H]^+ = 431.4$
5.55	[(S)-2-(5,6-Dimethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(2-fluoro-3-methoxy-6-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.7$; $[M+H]^+ = 449.3$
5.56	[(S)-2-(5,6-Dimethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(4-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.76$; $[M+H]^+ = 415.4$
5.57	(4,5-Dimethoxy-2-[1,2,3]triazol-2-yl-phenyl)-[(S)-2-(5,6-dimethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.67$; $[M+H]^+ = 461.3$
5.58	[(S)-2-(5,6-Dimethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(4,5-dimethyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.77$; $[M+H]^+ = 429.3$
5.59	[(S)-2-(5,6-Dimethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-pyrazol-1-yl-phenyl)-methanone. LC-MS: $t_R = 0.77$; $[M+H]^+ = 414.4$
5.60	[(S)-2-(5,6-Dimethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-fluoro-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.7$; $[M+H]^+ = 419.2$
5.61	[(S)-2-(5,6-Dimethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.67$; $[M+H]^+ = 401.2$
5.62	[(S)-2-(5,6-Dimethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(6-methyl-3-[1,2,3]triazol-2-yl-pyridin-2-yl)-methanone. LC-MS: $t_R = 0.68$; $[M+H]^+ = 416.2$
5.64	(5-Chloro-2-[1,2,3]triazol-2-yl-phenyl)-[(S)-2-(5,6-dimethoxy-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.69$; $[M+H]^+ = 467.2$
5.65	[(S)-2-(5,6-Dimethoxy-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-fluoro-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.63$; $[M+H]^+ = 451.3$
5.66	[(S)-2-(5,6-Dimethoxy-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(2-fluoro-6-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.61$; $[M+H]^+ = 451.3$
5.67	[(S)-2-(5,6-Dimethoxy-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methoxy-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.64$; $[M+H]^+ = 463.3$
5.68	[(S)-2-(5,6-Dimethoxy-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(2-fluoro-3-methyl-6-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.67$; $[M+H]^+ = 465.3$

5.69	[(S)-2-(5,6-Dimethoxy-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(2-fluoro-3-methoxy-6-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.63$; $[M+H]^+ = 481.3$
5.70	[(S)-2-(5,6-Dimethoxy-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(4-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.67$; $[M+H]^+ = 447.3$
5.71	[(S)-2-(5,6-Dimethoxy-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(2-[1,2,3]triazol-2-yl-5-trifluoromethyl-phenyl)-methanone. LC-MS: $t_R = 0.75$; $[M+H]^+ = 501.3$
5.72	[(S)-2-(5,6-Dimethoxy-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(4,5-dimethyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.72$; $[M+H]^+ = 461.3$
5.73	[(S)-2-(5,6-Dimethoxy-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.61$; $[M+H]^+ = 433.3$
5.74	[(S)-2-(5,6-Dimethoxy-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-[2-methyl-5-(4-trifluoromethyl-phenyl)-thiazol-4-yl]-methanone. LC-MS: $t_R = 0.85$; $[M+H]^+ = 531.3$
5.75	[(S)-2-(5,6-Dimethoxy-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(2-methyl-4-p-tolyl-thiazol-5-yl)-methanone. LC-MS: $t_R = 0.71$; $[M+H]^+ = 477.2$
5.76	[5-(3,5-Difluoro-phenyl)-2-methyl-thiazol-4-yl]-[(S)-2-(5,6-dimethoxy-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.77$; $[M+H]^+ = 499.3$
5.77	[(S)-2-(5,6-Dimethoxy-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-[5-(6-methoxy-pyridin-3-yl)-2-methyl-thiazol-4-yl]-methanone. LC-MS: $t_R = 0.69$; $[M+H]^+ = 494.3$
5.78	[(S)-2-(5,6-Dimethoxy-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-[5-(4-fluoro-phenyl)-2-methyl-thiazol-4-yl]-methanone. LC-MS: $t_R = 0.75$; $[M+H]^+ = 481.2$
5.79	[5-(4-Chloro-phenyl)-2-methyl-thiazol-4-yl]-[(S)-2-(5,6-dimethoxy-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.81$; $[M+H]^+ = 497.3$
5.80	[5-(3-Chloro-phenyl)-thiazol-4-yl]-[(S)-2-(5,6-dimethoxy-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.74$; $[M+H]^+ = 483.2$
5.81	[(S)-2-(5,6-Dimethoxy-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(2-pyrazol-1-yl-phenyl)-methanone. LC-MS: $t_R = 0.61$; $[M+H]^+ = 432.3$
5.82	[(S)-2-(5,6-Dimethoxy-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-pyrazol-1-yl-phenyl)-methanone. LC-MS: $t_R = 0.67$; $[M+H]^+ = 446.3$
5.83	[(S)-2-(5,6-Dimethoxy-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(6-methyl-3-pyrazol-1-yl-pyridin-2-yl)-methanone. LC-MS: $t_R = 0.61$; $[M+H]^+ = 447.3$
5.84	[(S)-2-(5,6-Dimethoxy-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(2-[1,2,3]triazol-2-yl-5-trifluoromethoxy-phenyl)-methanone. LC-MS: $t_R = 0.77$; $[M+H]^+ = 517.3$

5.85	[(S)-2-(5,6-Dimethoxy-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(6-methyl-3-[1,2,3]triazol-2-yl-pyridin-2-yl)-methanone. LC-MS: $t_R = 0.62$; $[M+H]^+ = 448.3$
5.86	[(S)-2-(5,6-Dimethoxy-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(4,5-dimethoxy-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.62$; $[M+H]^+ = 493.3$
5.87	[(S)-2-(5,6-Dimethoxy-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(3'-methyl-biphenyl-2-yl)-methanone. LC-MS: $t_R = 0.8$; $[M+H]^+ = 456.3$
5.88	[(S)-2-(5,6-Dimethoxy-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(4'-methyl-biphenyl-2-yl)-methanone. LC-MS: $t_R = 0.8$; $[M+H]^+ = 456.3$
5.89	[(S)-2-(5,6-Dimethoxy-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(2-methyl-4-m-tolyl-pyrimidin-5-yl)-methanone. LC-MS: $t_R = 0.68$; $[M+H]^+ = 472.3$
5.90	[(S)-2-(5,6-Dimethoxy-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(4-phenyl-pyrimidin-5-yl)-methanone. LC-MS: $t_R = 0.59$; $[M+H]^+ = 444.3$
5.91	[(S)-2-(5,6-Dimethyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(4'-fluoro-4,2'-dimethyl-biphenyl-2-yl)-methanone. LC-MS: $t_R = 0.95$; $[M+H]^+ = 456.4$
5.92	[(S)-2-(5,6-Dimethyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(2'-methoxy-4-methyl-biphenyl-2-yl)-methanone. LC-MS: $t_R = 0.89$; $[M+H]^+ = 454.4$
5.93	[(S)-2-(5,6-Dimethyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(3'-methoxy-4-methyl-biphenyl-2-yl)-methanone. LC-MS: $t_R = 0.89$; $[M+H]^+ = 454.3$
5.94	[(S)-2-(5,6-Dimethyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(4'-methoxy-4-methyl-biphenyl-2-yl)-methanone. LC-MS: $t_R = 0.89$; $[M+H]^+ = 454.4$
5.95	[(S)-2-(5,6-Dimethyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(2'-fluoro-4-methyl-biphenyl-2-yl)-methanone. LC-MS: $t_R = 0.88$; $[M+H]^+ = 442.3$
5.96	[(S)-2-(5,6-Dimethyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(4'-fluoro-4-methyl-biphenyl-2-yl)-methanone. LC-MS: $t_R = 0.89$; $[M+H]^+ = 442.3$
5.97	[(S)-2-(5,6-Dimethyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(3'-fluoro-4-methyl-biphenyl-2-yl)-methanone. LC-MS: $t_R = 0.9$; $[M+H]^+ = 442.3$
5.98	[(S)-2-(5,6-Dimethyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(4,2'-dimethyl-biphenyl-2-yl)-methanone. LC-MS: $t_R = 0.93$; $[M+H]^+ = 438.4$
5.99	[(S)-2-(5,6-Dimethyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(4,3'-dimethyl-biphenyl-2-yl)-methanone. LC-MS: $t_R = 0.93$; $[M+H]^+ = 438.4$
5.100	[(S)-2-(5,6-Dimethyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(4,4'-dimethyl-biphenyl-2-yl)-methanone. LC-MS: $t_R = 0.93$; $[M+H]^+ = 438.4$

5.101	[(S)-2-(5,6-Dimethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(3'-methoxy-biphenyl-2-yl)-methanone. LC-MS: $t_R = 0.83$; $[M+H]^+ = 440.3$
5.102	[(S)-2-(5,6-Dimethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(4'-methoxy-biphenyl-2-yl)-methanone. LC-MS: $t_R = 0.82$; $[M+H]^+ = 440.3$
5.103	[(S)-2-(5,6-Dimethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(2'-fluoro-biphenyl-2-yl)-methanone. LC-MS: $t_R = 0.82$; $[M+H]^+ = 428.3$
5.104	[(S)-2-(5,6-Dimethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(4'-fluoro-biphenyl-2-yl)-methanone. LC-MS: $t_R = 0.83$; $[M+H]^+ = 428.3$
5.105	[(S)-2-(5,6-Dimethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(3'-fluoro-biphenyl-2-yl)-methanone. LC-MS: $t_R = 0.83$; $[M+H]^+ = 428.3$
5.106	[(S)-2-(5,6-Dimethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(3'-methyl-biphenyl-2-yl)-methanone. LC-MS: $t_R = 0.87$; $[M+H]^+ = 424.4$
5.107	[(S)-2-(5,6-Dimethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(4'-methyl-biphenyl-2-yl)-methanone. LC-MS: $t_R = 0.87$; $[M+H]^+ = 424.3$
5.108	[(S)-2-(4,5-Dimethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(4'-fluoro-4,2'-dimethyl-biphenyl-2-yl)-methanone. LC-MS: $t_R = 0.95$; $[M+H]^+ = 456.4$
5.109	[(S)-2-(4,5-Dimethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(3'-fluoro-4-methyl-biphenyl-2-yl)-methanone. LC-MS: $t_R = 0.9$; $[M+H]^+ = 442.3$
5.110	[(S)-2-(4,5-Dimethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(3'-methoxy-4-methyl-biphenyl-2-yl)-methanone. LC-MS: $t_R = 0.89$; $[M+H]^+ = 454.4$
5.111	[(S)-2-(4,5-Dimethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(4,3'-dimethyl-biphenyl-2-yl)-methanone. LC-MS: $t_R = 0.93$; $[M+H]^+ = 438.4$
5.112	[(S)-2-(4,5-Dimethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(4'-methoxy-4-methyl-biphenyl-2-yl)-methanone. LC-MS: $t_R = 0.89$; $[M+H]^+ = 454.3$
5.113	[(S)-2-(4,5-Dimethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(4,4'-dimethyl-biphenyl-2-yl)-methanone. LC-MS: $t_R = 0.94$; $[M+H]^+ = 438.4$
5.114	[(S)-2-(4,5-Dimethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(2'-fluoro-4-methyl-biphenyl-2-yl)-methanone. LC-MS: $t_R = 0.89$; $[M+H]^+ = 442.3$
5.115	[(S)-2-(4,5-Dimethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(4'-fluoro-4-methyl-biphenyl-2-yl)-methanone. LC-MS: $t_R = 0.90$; $[M+H]^+ = 442.3$
5.116	[(S)-2-(4,5-Dimethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(3'-fluoro-biphenyl-2-yl)-methanone. LC-MS: $t_R = 0.84$; $[M+H]^+ = 428.3$

5.117	[(S)-2-(4,5-Dimethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(3'-methoxy-biphenyl-2-yl)-methanone. LC-MS: $t_R = 0.83$; $[M+H]^+ = 440.3$
5.118	[(S)-2-(4,5-Dimethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(3'-methyl-biphenyl-2-yl)-methanone. LC-MS: $t_R = 0.88$; $[M+H]^+ = 424.3$
5.119	[(S)-2-(4,5-Dimethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(4'-methoxy-biphenyl-2-yl)-methanone. LC-MS: $t_R = 0.83$; $[M+H]^+ = 440.3$
5.120	[(S)-2-(4,5-Dimethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(4'-methyl-biphenyl-2-yl)-methanone. LC-MS: $t_R = 0.88$; $[M+H]^+ = 424.4$
5.121	[(S)-2-(4,5-Dimethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(2'-fluoro-biphenyl-2-yl)-methanone. LC-MS: $t_R = 0.83$; $[M+H]^+ = 428.3$
5.122	[(S)-2-(4,5-Dimethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(4'-fluoro-biphenyl-2-yl)-methanone. LC-MS: $t_R = 0.84$; $[M+H]^+ = 428.3$
5.123	[(S)-2-(6-Bromo-5-fluoro-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(2-fluoro-6-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.83$; $[M+H]^+ = 487.1$
5.124	[(S)-2-(6-Bromo-5-fluoro-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(2-fluoro-3-methyl-6-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.90$; $[M+H]^+ = 501.1$
5.125	[(S)-2-(6-Bromo-5-fluoro-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(4,5-dimethoxy-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.79$; $[M+H]^+ = 529.2$
5.126	[(S)-2-(6-Bromo-5-fluoro-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-fluoro-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.83$; $[M+H]^+ = 487.2$
5.127	[(S)-2-(6-Bromo-5-fluoro-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(4,5-dimethyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.90$; $[M+H]^+ = 497.2$
5.128	[(S)-2-(6-Bromo-5-fluoro-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-pyrazol-1-yl-phenyl)-methanone. LC-MS: $t_R = 0.86$; $[M+H]^+ = 482.2$
5.129	[(S)-2-(6-Bromo-5-fluoro-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-chloro-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.90$; $[M+H]^+ = 503.1$
5.130	[(S)-2-(6-Bromo-5-fluoro-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(4-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.85$; $[M+H]^+ = 483.2$
5.131	[(S)-2-(6-Bromo-5-fluoro-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(6-methyl-3-[1,2,3]triazol-2-yl-pyridin-2-yl)-methanone. LC-MS: $t_R = 0.75$; $[M+H]^+ = 484.2$
5.132	[(S)-2-(6-Bromo-5-fluoro-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(2-fluoro-3-methoxy-6-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.85$; $[M+H]^+ = 517.2$

5.133	[(S)-2-(6-Chloro-5-trifluoromethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(2-fluoro-6-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: t_R = 1.04; $[M+H]^+$ = 493.2
5.134	[(S)-2-(6-Chloro-5-trifluoromethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(2-fluoro-3-methyl-6-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: t_R = 1.14; $[M+H]^+$ = 507.2
5.135	[(S)-2-(6-Chloro-5-trifluoromethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(4,5-dimethoxy-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: t_R = 0.98; $[M+H]^+$ = 535.2
5.136	[(S)-2-(6-Chloro-5-trifluoromethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-fluoro-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: t_R = 1.03; $[M+H]^+$ = 493.2
5.137	[(S)-2-(6-Chloro-5-trifluoromethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(4,5-dimethyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: t_R = 1.09.; $[M+H]^+$ = 503.3
5.138	[(S)-2-(6-Chloro-5-trifluoromethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-pyrazol-1-yl-phenyl)-methanone. LC-MS: t_R = 1.05.; $[M+H]^+$ = 488.2
5.139	(5-Chloro-2-[1,2,3]triazol-2-yl-phenyl)-[(S)-2-(6-chloro-5-trifluoromethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-methanone. LC-MS: t_R = 1.09; $[M+H]^+$ = 509.2
5.140	[(S)-2-(6-Chloro-5-trifluoromethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(4-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: t_R = 1.05; $[M+H]^+$ = 489.2
5.141	[(S)-2-(6-Chloro-5-trifluoromethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(2-pyrazol-1-yl-phenyl)-methanone. LC-MS: t_R = 1.00; $[M+H]^+$ = 474.2
5.142	[(S)-2-(6-Chloro-5-trifluoromethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: t_R = 0.99; $[M+H]^+$ = 475.2
5.143	[(S)-2-(6-Chloro-5-trifluoromethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(6-methyl-3-pyrazol-1-yl-pyridin-2-yl)-methanone. LC-MS: t_R = 0.94; $[M+H]^+$ = 489.2
5.144	[(S)-2-(6-Chloro-5-trifluoromethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(2-[1,2,3]triazol-2-yl-5-trifluoromethyl-phenyl)-methanone. LC-MS: t_R = 1.12; $[M+H]^+$ = 543.2
5.145	[(S)-2-(6-Chloro-5-trifluoromethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(6-methyl-3-[1,2,3]triazol-2-yl-pyridin-2-yl)-methanone. LC-MS: t_R = 0.94; $[M+H]^+$ = 490.2
5.146	[(S)-2-(6-Chloro-5-trifluoromethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(2-fluoro-3-methoxy-6-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: t_R = 1.05; $[M+H]^+$ = 523.2
5.147	[(S)-2-(6-Chloro-5-trifluoromethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(2-[1,2,3]triazol-2-yl-5-trifluoromethoxy-phenyl)-methanone. LC-MS: t_R = 1.14; $[M+H]^+$ = 559.2
5.148	[(S)-2-(6-Chloro-5-trifluoromethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(2'-methoxy-4-methyl-biphenyl-2-yl)-methanone. LC-MS: t_R = 1.18; $[M+H]^+$ = 528.2

5.149	[(S)-2-(6-Chloro-5-trifluoromethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(3'-methoxy-4-methyl-biphenyl-2-yl)-methanone. LC-MS: $t_R = 1.18$; $[M+H]^+ = 528.3$
5.150	[(S)-2-(6-Chloro-5-trifluoromethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(4'-methoxy-4-methyl-biphenyl-2-yl)-methanone. LC-MS: $t_R = 1.19$; $[M+H]^+ = 528.2$
5.151	[(S)-2-(6-Chloro-5-trifluoromethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(2'-fluoro-4-methyl-biphenyl-2-yl)-methanone. LC-MS: $t_R = 1.18$; $[M+H]^+ = 516.2$
5.152	[(S)-2-(6-Chloro-5-trifluoromethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(4'-fluoro-4-methyl-biphenyl-2-yl)-methanone. LC-MS: $t_R = 1.19$; $[M+H]^+ = 516.2$
5.153	[(S)-2-(6-Chloro-5-trifluoromethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(3'-fluoro-4-methyl-biphenyl-2-yl)-methanone. LC-MS: $t_R = 1.19$; $[M+H]^+ = 516.2$
5.154	[(S)-2-(6-Chloro-5-trifluoromethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(4,4'-dimethyl-biphenyl-2-yl)-methanone. LC-MS: $t_R = 1.24$; $[M+H]^+ = 512.3$
5.155	[(S)-2-(6-Chloro-5-trifluoromethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(4-methyl-3'-trifluoromethyl-biphenyl-2-yl)-methanone. LC-MS: $t_R = 1.25$; $[M+H]^+ = 566.2$
5.156	(2-Benzo[1,3]dioxol-5-yl-5-methyl-phenyl)-[(S)-2-(6-chloro-5-trifluoromethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 1.16$; $[M+H]^+ = 542.2$
5.157	[(S)-2-(6-Chloro-5-trifluoromethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(4-methyl-4'-trifluoromethyl-biphenyl-2-yl)-methanone. LC-MS: $t_R = 1.26$; $[M+H]^+ = 566.2$
5.158	[(S)-2-(6-Chloro-5-trifluoromethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(4-methyl-4'-trifluoromethoxy-biphenyl-2-yl)-methanone. LC-MS: $t_R = 1.28$; $[M+H]^+ = 582.3$
5.159	2'-[(S)-2-(6-Chloro-5-trifluoromethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidine-1-carbonyl]-4'-methyl-biphenyl-3-carbonitrile. LC-MS: $t_R = 1.13$; $[M+H]^+ = 523.2$
5.160	2'-[(S)-2-(6-Chloro-5-trifluoromethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidine-1-carbonyl]-4'-methyl-biphenyl-4-carbonitrile. LC-MS: $t_R = 1.13$; $[M+H]^+ = 523.3$
5.161	[(S)-2-(6-Chloro-5-trifluoromethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(4,2',3'-trimethyl-biphenyl-2-yl)-methanone. LC-MS: $t_R = 1.27$; $[M+H]^+ = 526.3$
5.162	[(S)-2-(6-Chloro-5-trifluoromethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(4'-ethoxy-4-methyl-biphenyl-2-yl)-methanone. LC-MS: $t_R = 1.24$; $[M+H]^+ = 542.3$
5.163	[(S)-2-(6-Chloro-5-trifluoromethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(4-methyl-3'-trifluoromethoxy-biphenyl-2-yl)-methanone. LC-MS: $t_R = 1.27$; $[M+H]^+ = 582.2$
5.164	(2-Fluoro-6-[1,2,3]triazol-2-yl-phenyl)-[(S)-2-methyl-2-(5-trifluoromethyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.82$; $[M+H]^+ = 459.2$

5.165	(5-Fluoro-2-[1,2,3]triazol-2-yl-phenyl)-[(S)-2-methyl-2-(5-trifluoromethyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.82$; $[M+H]^+ = 459.2$
5.166	(5-Chloro-2-[1,2,3]triazol-2-yl-phenyl)-[(S)-2-methyl-2-(5-trifluoromethyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.89$; $[M+H]^+ = 475.2$
5.167	[(S)-2-Methyl-2-(5-trifluoromethyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-(2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.78$; $[M+H]^+ = 441.3$
5.168	(2-Fluoro-3-methoxy-6-[1,2,3]triazol-2-yl-phenyl)-[(S)-2-methyl-2-(5-trifluoromethyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.84$; $[M+H]^+ = 489.3$
5.169	(2-Fluoro-3-methyl-6-[1,2,3]triazol-2-yl-phenyl)-[(S)-2-methyl-2-(5-trifluoromethyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.89$; $[M+H]^+ = 473.3$
5.170	(4,5-Dimethyl-2-[1,2,3]triazol-2-yl-phenyl)-[(S)-2-methyl-2-(5-trifluoromethyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.89$; $[M+H]^+ = 469.3$
5.171	(4-Methyl-2-[1,2,3]triazol-2-yl-phenyl)-[(S)-2-methyl-2-(5-trifluoromethyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.84$; $[M+H]^+ = 455.3$
5.172	[(S)-2-Methyl-2-(5-trifluoromethyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-(2-[1,2,3]triazol-2-yl-5-trifluoromethyl-phenyl)-methanone. LC-MS: $t_R = 0.93$; $[M+H]^+ = 509.2$
5.173	[(S)-2-Methyl-2-(5-trifluoromethyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-(2-[1,2,3]triazol-2-yl-5-trifluoromethoxy-phenyl)-methanone. LC-MS: $t_R = 0.95$; $[M+H]^+ = 525.3$
5.174	(4,5-Dimethoxy-2-[1,2,3]triazol-2-yl-phenyl)-[(S)-2-methyl-2-(5-trifluoromethyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.78$; $[M+H]^+ = 501.3$
5.175	(5-Methyl-2-pyrazol-1-yl-phenyl)-[(S)-2-methyl-2-(5-trifluoromethyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.85$; $[M+H]^+ = 454.3$
5.176	[(S)-2-Methyl-2-(5-trifluoromethyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-(2-pyrazol-1-yl-phenyl)-methanone. LC-MS: $t_R = 0.79$; $[M+H]^+ = 440.3$
5.177	(6-Methyl-3-pyrazol-1-yl-pyridin-2-yl)-[(S)-2-methyl-2-(5-trifluoromethyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.75$; $[M+H]^+ = 455.3$
5.178	(6-Methyl-3-[1,2,3]triazol-2-yl-pyridin-2-yl)-[(S)-2-methyl-2-(5-trifluoromethyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.75$; $[M+H]^+ = 456.3$
5.179	(3-Methyl-2-[1,2,3]triazol-2-yl-phenyl)-[(S)-2-methyl-2-(5-trifluoromethyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.80$; $[M+H]^+ = 455.3$
5.180	3-[(S)-2-Methyl-2-(5-trifluoromethyl-1H-benzoimidazol-2-yl)-pyrrolidine-1-carbonyl]-4-[1,2,3]triazol-2-yl-benzonitrile. LC-MS: $t_R = 0.80$; $[M+H]^+ = 466.2$

5.181	(4'-Fluoro-4,2'-dimethyl-biphenyl-2-yl)-[(S)-2-methyl-2-(5-trifluoromethyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 1.03$; $[M+H]^+ = 496.3$
5.182	(3'-Fluoro-4-methyl-biphenyl-2-yl)-[(S)-2-methyl-2-(5-trifluoromethyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.99$; $[M+H]^+ = 482.3$
5.183	[(S)-2-Methyl-2-(5-trifluoromethyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-(4-methyl-4'-trifluoromethyl-biphenyl-2-yl)-methanone. LC-MS: $t_R = 1.08$; $[M+H]^+ = 532.3$
5.184	(4-Methyl-3'-trifluoromethoxy-biphenyl-2-yl)-[(S)-2-methyl-2-(5-trifluoromethyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 1.09$; $[M+H]^+ = 548.3$
5.185	(2'-Methoxy-4-methyl-biphenyl-2-yl)-[(S)-2-methyl-2-(5-trifluoromethyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.98$; $[M+H]^+ = 494.3$
5.186	(4,2'-Dimethyl-biphenyl-2-yl)-[(S)-2-methyl-2-(5-trifluoromethyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 1.02$; $[M+H]^+ = 478.3$
5.187	(4-Methyl-4'-trifluoromethoxy-biphenyl-2-yl)-[(S)-2-methyl-2-(5-trifluoromethyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 1.09$; $[M+H]^+ = 548.3$
5.188	(3'-Methoxy-4-methyl-biphenyl-2-yl)-[(S)-2-methyl-2-(5-trifluoromethyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.98$; $[M+H]^+ = 494.3$
5.189	(4,3'-Dimethyl-biphenyl-2-yl)-[(S)-2-methyl-2-(5-trifluoromethyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 1.03$; $[M+H]^+ = 478.3$
5.190	4'-Methyl-2'-[(S)-2-methyl-2-(5-trifluoromethyl-1H-benzoimidazol-2-yl)-pyrrolidine-1-carbonyl]-biphenyl-3-carbonitrile. LC-MS: $t_R = 0.94$; $[M+H]^+ = 489.3$
5.191	(4'-Methoxy-4-methyl-biphenyl-2-yl)-[(S)-2-methyl-2-(5-trifluoromethyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.98$; $[M+H]^+ = 494.3$
5.192	(4,4'-Dimethyl-biphenyl-2-yl)-[(S)-2-methyl-2-(5-trifluoromethyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 1.04$; $[M+H]^+ = 478.3$
5.193	4'-Methyl-2'-[(S)-2-methyl-2-(5-trifluoromethyl-1H-benzoimidazol-2-yl)-pyrrolidine-1-carbonyl]-biphenyl-4-carbonitrile. LC-MS: $t_R = 0.95$; $[M+H]^+ = 489.3$
5.194	(2'-Fluoro-4-methyl-biphenyl-2-yl)-[(S)-2-methyl-2-(5-trifluoromethyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.98$; $[M+H]^+ = 482.3$
5.195	[(S)-2-Methyl-2-(5-trifluoromethyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-(4-methyl-3'-trifluoromethyl-biphenyl-2-yl)-methanone. LC-MS: $t_R = 1.07$; $[M+H]^+ = 532.3$
5.196	[(S)-2-Methyl-2-(5-trifluoromethyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-(4,2',3'-trimethyl-biphenyl-2-yl)-methanone. LC-MS: $t_R = 1.06$; $[M+H]^+ = 492.3$

5.197	(4'-Fluoro-4-methyl-biphenyl-2-yl)-[(S)-2-methyl-2-(5-trifluoromethyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.99$; $[M+H]^+ = 482.2$
5.198	(2-Benzo[1,3]dioxol-5-yl-5-methyl-phenyl)-[(S)-2-methyl-2-(5-trifluoromethyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.96$; $[M+H]^+ = 508.3$
5.199	[(S)-2-(4,5-Difluoro-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(4,5-dimethyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.92$; $[M+H]^+ = 437.3$
5.200	(5-Chloro-2-[1,2,3]triazol-2-yl-phenyl)-[(S)-2-(4,5-difluoro-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.90$; $[M+H]^+ = 443.2$
5.201	[(S)-2-(4,5-Difluoro-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-pyrazol-1-yl-phenyl)-methanone. LC-MS: $t_R = 0.86$; $[M+H]^+ = 422.3$
5.202	[(S)-2-(4,5-Difluoro-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(4-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.86$; $[M+H]^+ = 423.3$
5.203	[(S)-2-(4,5-Difluoro-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(2-fluoro-3-methoxy-6-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.85$; $[M+H]^+ = 457.3$
5.204	[(S)-2-(4,5-Difluoro-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-fluoro-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.83$; $[M+H]^+ = 427.3$
5.205	[(S)-2-(4,5-Difluoro-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(6-methyl-3-[1,2,3]triazol-2-yl-pyridin-2-yl)-methanone. LC-MS: $t_R = 0.73$; $[M+H]^+ = 424.2$
5.206	[(S)-2-(4,5-Difluoro-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(2-methyl-5-phenyl-thiazol-4-yl)-methanone. LC-MS: $t_R = 0.90$; $[M+H]^+ = 439.2$
5.207	[5-(3-Chloro-phenyl)-2-methyl-thiazol-4-yl]-[(S)-2-(4,5-difluoro-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.98$; $[M+H]^+ = 473.2$
5.208	[(S)-2-(4,5-Difluoro-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-[5-(4-fluoro-phenyl)-2-methyl-thiazol-4-yl]-methanone. LC-MS: $t_R = 0.92$; $[M+H]^+ = 457.3$
5.209	[(S)-2-(4,5-Difluoro-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(2-methyl-5-p-tolyl-thiazol-4-yl)-methanone. LC-MS: $t_R = 0.97$; $[M+H]^+ = 453.3$
5.210	[(S)-2-(4,5-Difluoro-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(2-methyl-5-m-tolyl-thiazol-4-yl)-methanone. LC-MS: $t_R = 0.96$; $[M+H]^+ = 453.3$
5.211	[5-(4-Chloro-phenyl)-2-methyl-thiazol-4-yl]-[(S)-2-(4,5-difluoro-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.99$; $[M+H]^+ = 473.2$
5.212	[(S)-2-(4,5-Difluoro-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-[5-(3-fluoro-phenyl)-2-methyl-thiazol-4-yl]-methanone. LC-MS: $t_R = 0.92$; $[M+H]^+ = 457.2$

5.213	[(S)-2-(4,5-Difluoro-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-[5-(3-methoxy-phenyl)-2-methyl-thiazol-4-yl]-methanone. LC-MS: $t_R = 0.91$; $[M+H]^+ = 469.3$
5.214	[(S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-pyrimidin-2-yl-phenyl)-methanone. LC-MS: $t_R = 77$; $[M+H]^+ = 446.2$
5.215	[(S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-pyridin-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.75$; $[M+H]^+ = 445.2$
5.216	[(S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-pyridin-3-yl-phenyl)-methanone. LC-MS: $t_R = 0.66$; $[M+H]^+ = 445.2$
5.217	[(S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-pyridin-3-yl-phenyl)-methanone. LC-MS: $t_R = 0.7$; $[M+H]^+ = 445.3$
5.218	[(S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(4-fluoro-biphenyl-2-yl)-methanone. LC-MS: $t_R = 0.95$; $[M+H]^+ = 448.3$
5.219	[(S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-fluoro-2-pyridin-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.8$; $[M+H]^+ = 449.3$
5.220	[(S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-fluoro-2-pyridin-3-yl-phenyl)-methanone. LC-MS: $t_R = 0.7$; $[M+H]^+ = 449.3$
5.221	[(S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(2-fluoro-6-pyridin-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.82$; $[M+H]^+ = 449.3$
5.222	[(S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(2-fluoro-6-pyrimidin-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.8$; $[M+H]^+ = 450.3$
5.223	[(S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(4-methyl-biphenyl-2-yl)-methanone. LC-MS: $t_R = 0.82$; $[M+H]^+ = 446.3$
5.224	[(S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-fluoro-2-pyrimidin-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.81$; $[M+H]^+ = 450.3$
5.225	[(S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(3-fluoro-biphenyl-2-yl)-methanone. LC-MS: $t_R = 0.94$; $[M+H]^+ = 448.3$
5.226	[(S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-[5-(4-fluoro-phenyl)-2-phenyl-thiazol-4-yl]-methanone. LC-MS: $t_R = 1.09$; $[M+H]^+ = 531.3$
5.227	[2,5-Bis-(4-fluoro-phenyl)-thiazol-4-yl]-[(S)-2-(5-chloro-4-methyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 1.09$; $[M+H]^+ = 549.4$
5.228	[(S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-[5-(4-fluoro-phenyl)-2-#o-tolyl-thiazol-4-yl]-methanone. LC-MS: $t_R = 1.12$; $[M+H]^+ = 545.3$

5.229	[(S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-[5-(4-fluoro-phenyl)-2-(2-methoxy-phenyl)-thiazol-4-yl]-methanone. LC-MS: $t_R = 1.09$; $[M+H]^+ = 561.3$
5.230	[(S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-[5-(4-fluoro-phenyl)-2-m-tolyl-thiazol-4-yl]-methanone. LC-MS: $t_R = 1.14$; $[M+H]^+ = 545.3$
5.231	[(S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-[5-(4-fluoro-phenyl)-2-(3-methoxy-phenyl)-thiazol-4-yl]-methanone. LC-MS: $t_R = 1.1$; $[M+H]^+ = 561.3$
5.232	[(S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-[5-(4-fluoro-phenyl)-2-p-tolyl-thiazol-4-yl]-methanone. LC-MS: $t_R = 1.14$; $[M+H]^+ = 545.3$
5.233	[(S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-[5-(4-fluoro-phenyl)-2-(2-fluoro-phenyl)-thiazol-4-yl]-methanone. LC-MS: $t_R = 1.1$; $[M+H]^+ = 549.3$
5.234	[(S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-[5-(4-fluoro-phenyl)-2-(3-fluoro-phenyl)-thiazol-4-yl]-methanone. LC-MS: $t_R = 1.1$; $[M+H]^+ = 549.3$
5.235	(4,5-Dimethyl-2-[1,2,3]triazol-2-yl-phenyl)-[(S)-2-methyl-2-(5-trifluoromethoxy-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.93$; $[M+H]^+ = 485.3$
5.236	(4-Methyl-biphenyl-2-yl)-[(S)-2-methyl-2-(5-trifluoromethoxy-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 1.01$; $[M+H]^+ = 480.3$
5.237	(5-Methoxy-2-[1,2,3]triazol-2-yl-phenyl)-[(S)-2-methyl-2-(5-trifluoromethoxy-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.85$; $[M+H]^+ = 487.3$
5.238	(4-Methyl-2-[1,2,3]triazol-2-yl-phenyl)-[(S)-2-methyl-2-(5-trifluoromethoxy-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.88$; $[M+H]^+ = 471.3$
5.239	(5-Methyl-2-pyridin-2-yl-phenyl)-[(S)-2-methyl-2-(5-trifluoromethoxy-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.83$; $[M+H]^+ = 481.3$
5.240	(4-Methyl-2-pyridin-2-yl-phenyl)-[(S)-2-methyl-2-(5-trifluoromethoxy-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.83$; $[M+H]^+ = 481.3$
5.241	(4-Methyl-2-pyridin-3-yl-phenyl)-[(S)-2-methyl-2-(5-trifluoromethoxy-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.74$; $[M+H]^+ = 481.3$
5.242	(5-Methyl-biphenyl-2-yl)-[(S)-2-methyl-2-(5-trifluoromethoxy-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 1.01$; $[M+H]^+ = 480.3$
5.243	(2-Methyl-5-o-tolyl-thiazol-4-yl)-[(S)-2-methyl-2-(5-trifluoromethoxy-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.98$; $[M+H]^+ = 501.3$
5.244	[5-(2,3-Dichloro-phenyl)-2-methyl-thiazol-4-yl]-[(S)-2-methyl-2-(5-trifluoromethoxy-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 1.03$; $[M+H]^+ = 555.2$

5.245	[5-(3,4-Dimethyl-phenyl)-2-methyl-thiazol-4-yl]-[(S)-2-methyl-2-(5-trifluoromethoxy-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 1.03$; $[M+H]^+ = 515.3$
5.246	[5-(3,5-Difluoro-phenyl)-2-methyl-thiazol-4-yl]-[(S)-2-methyl-2-(5-trifluoromethoxy-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.99$; $[M+H]^+ = 523.3$
5.247	[5-(3-Methoxy-phenyl)-2-methyl-thiazol-4-yl]-[(S)-2-methyl-2-(5-trifluoromethoxy-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.95$; $[M+H]^+ = 517.3$
5.248	[5-(4-Fluoro-phenyl)-2-methyl-thiazol-4-yl]-[(S)-2-methyl-2-(5-trifluoromethoxy-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.96$; $[M+H]^+ = 505.3$
5.249	[(S)-2-(6-Bromo-4-methyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-[2-(3-methoxy-phenylethynyl)-phenyl]-methanone. LC-MS: $t_R = 1.0$; $[M+H]^+ = 528.3$
5.250	[2-(3-Methoxy-phenylethynyl)-phenyl]-[(S)-2-methyl-2-(5-trifluoromethyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 1.05$; $[M+H]^+ = 504.4$
5.251	[(S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-[2-(2,3-difluoro-phenyl)-5-(4-fluoro-phenyl)-thiazol-4-yl]-methanone. LC-MS: $t_R = 1.12$; $[M+H]^+ = 567.3$
5.252	[(S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-[2-(4-fluoro-3-methyl-phenyl)-5-(4-fluoro-phenyl)-thiazol-4-yl]-methanone. LC-MS: $t_R = 1.15$; $[M+H]^+ = 563.3$
5.253	[(S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-[2-(2-fluoro-4-methyl-phenyl)-5-(4-fluoro-phenyl)-thiazol-4-yl]-methanone. LC-MS: $t_R = 1.15$; $[M+H]^+ = 563.3$
5.254	[(S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-[2-(3,4-dimethyl-phenyl)-5-(4-fluoro-phenyl)-thiazol-4-yl]-methanone. LC-MS: $t_R = 1.18$; $[M+H]^+ = 559.4$
5.255	3-[4-[(S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidine-1-carbonyl]-5-(4-fluoro-phenyl)-thiazol-2-yl]-benzonitrile. LC-MS: $t_R = 1.03$; $[M+H]^+ = 556.3$
5.256	[(S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-[2-(2-chloro-phenyl)-5-(4-fluoro-phenyl)-thiazol-4-yl]-methanone. LC-MS: $t_R = 1.13$; $[M+H]^+ = 565.3$
5.257	[(S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-[2-(3-chloro-phenyl)-5-(4-fluoro-phenyl)-thiazol-4-yl]-methanone. LC-MS: $t_R = 1.15$; $[M+H]^+ = 565.3$
5.258	[(S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-[2-(3-chloro-4-methyl-phenyl)-5-(4-fluoro-phenyl)-thiazol-4-yl]-methanone. LC-MS: $t_R = 1.2$; $[M+H]^+ = 579.3$
5.259	[(S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-[2-(4-fluoro-2-methyl-phenyl)-5-(4-fluoro-phenyl)-thiazol-4-yl]-methanone. LC-MS: $t_R = 1.13$; $[M+H]^+ = 563.3$
5.260	[(S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-[2-(4-fluoro-2-methoxy-phenyl)-5-(4-fluoro-phenyl)-thiazol-4-yl]-methanone. LC-MS: $t_R = 1.1$; $[M+H]^+ = 579.4$

5.261	[(S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-[2-(2,5-difluoro-phenyl)-5-(4-fluoro-phenyl)-thiazol-4-yl]-methanone. LC-MS: $t_R = 1.11$; $[M+H]^+ = 567.3$
5.262	(4-Bromo-biphenyl-2-yl)-[(S)-2-methyl-2-(5-trifluoromethyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 1.08$; $[M+H]^+ = 528.2$
5.263	[(S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(2-methyl-5-[1,2,3]triazol-2-yl-pyridin-4-yl)-methanone. LC-MS: $t_R = 0.73$; $[M+H]^+ = 436.3$
5.264	[(S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(2-methyl-5-phenyl-pyridin-4-yl)-methanone. LC-MS: $t_R = 0.73$; $[M+H]^+ = 445.3$
5.265	[(S)-2-Methyl-2-(5-trifluoromethyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-(4-pyridin-3-yl-biphenyl-2-yl)-methanone. LC-MS: $t_R = 0.82$; $[M+H]^+ = 527.3$
5.266	[(S)-2-(6-Bromo-4,5-dimethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.89$; $[M+H]^+ = 493.3$
5.267	[(S)-2-(6-Bromo-4,5-dimethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(3,4-dimethyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.88$; $[M+H]^+ = 507.3$
5.268	[(S)-2-(6-Bromo-4,5-dimethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(2-fluoro-6-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.86$; $[M+H]^+ = 497.3$
5.269	Biphenyl-2-yl-[(S)-2-(6-bromo-4,5-dimethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.96$; $[M+H]^+ = 488.3$
5.270	[(S)-2-(6-Bromo-4,5-dimethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(4-methyl-biphenyl-2-yl)-methanone. LC-MS: $t_R = 1.01$; $[M+H]^+ = 502.3$
5.271	[(S)-2-(6-Bromo-4,5-dimethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(2-methyl-5-o-tolyl-thiazol-4-yl)-methanone. LC-MS: $t_R = 1.0$; $[M+H]^+ = 523.3$
5.272	[(S)-2-(6-Bromo-4,5-dimethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-[5-(2,3-dichloro-phenyl)-2-methyl-thiazol-4-yl]-methanone. LC-MS: $t_R = 1.04$; $[M+H]^+ = 577.2$
5.273	[(S)-2-(6-Bromo-4,5-dimethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-[5-(3,4-dimethyl-phenyl)-2-methyl-thiazol-4-yl]-methanone. LC-MS: $t_R = 1.04$; $[M+H]^+ = 537.3$
5.274	[(S)-2-(6-Bromo-4,5-dimethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-[5-(3,5-difluoro-phenyl)-2-methyl-thiazol-4-yl]-methanone. LC-MS: $t_R = 0.99$; $[M+H]^+ = 545.3$
5.275	[(S)-2-(6-Bromo-4,5-dimethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-[5-(3-methoxy-phenyl)-2-methyl-thiazol-4-yl]-methanone. LC-MS: $t_R = 0.96$; $[M+H]^+ = 539.3$
5.276	[(S)-2-(6-Bromo-4,5-dimethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-[5-(4-fluoro-phenyl)-2-methyl-thiazol-4-yl]-methanone. LC-MS: $t_R = 0.97$; $[M+H]^+ = 527.3$

5.277	[(S)-2-(5-Bromo-7-methyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-pyrimidin-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.84$; $[M+H]^+ = 490.3$
5.278	[(S)-2-Methyl-2-(7-methyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-(5-methyl-2-pyrimidin-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.73$; $[M+H]^+ = 412.3$
5.279	[(S)-2-(6-Chloro-5-trifluoromethyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-pyrimidin-2-yl-phenyl)-methanone. LC-MS: $t_R = 1.07$; $[M+H]^+ = 500.3$
5.280	[(S)-2-(5,6-Dimethoxy-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-pyrimidin-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.7$; $[M+H]^+ = 458.4$
5.281	(5-Methyl-2-pyrimidin-2-yl-phenyl)-[(S)-2-methyl-2-(5-trifluoromethoxy-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.86$; $[M+H]^+ = 482.3$
5.282	[(S)-2-(6-Bromo-5-fluoro-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-pyrimidin-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.86$; $[M+H]^+ = 494.2$
5.283	[(S)-2-(6,7-Difluoro-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-pyrimidin-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.89$; $[M+H]^+ = 434.3$
5.284	[(S)-2-(5,6-Dimethyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-pyrimidin-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.77$; $[M+H]^+ = 426.4$
5.285	[(S)-2-(5-Chloro-6-trifluoromethyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-[5-(4-fluoro-phenyl)-2-phenyl-thiazol-4-yl]-methanone. LC-MS: $t_R = 1.29$; $[M+H]^+ = 585.3$
5.286	[(S)-2-(5-Chloro-6-trifluoromethyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-[5-(4-fluoro-phenyl)-2-o-tolyl-thiazol-4-yl]-methanone. LC-MS: $t_R = 1.32$; $[M+H]^+ = 599.3$
5.287	[(S)-2-(5-Chloro-6-trifluoromethyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-[5-(4-fluoro-phenyl)-2-m-tolyl-thiazol-4-yl]-methanone. LC-MS: $t_R = 1.34$; $[M+H]^+ = 599.3$
5.288	[(S)-2-(5-Chloro-6-trifluoromethyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-[5-(4-fluoro-phenyl)-2-p-tolyl-thiazol-4-yl]-methanone. LC-MS: $t_R = 1.34$; $[M+H]^+ = 599.3$
5.289	[(S)-2-(5-Chloro-6-trifluoromethyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-[5-(4-fluoro-phenyl)-2-(2-fluoro-phenyl)-thiazol-4-yl]-methanone. LC-MS: $t_R = 1.31$; $[M+H]^+ = 603.3$
5.290	[(S)-2-(5-Chloro-6-trifluoromethyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-[5-(4-fluoro-phenyl)-2-(3-fluoro-phenyl)-thiazol-4-yl]-methanone. LC-MS: $t_R = 1.30$; $[M+H]^+ = 603.3$
5.291	[2,5-Bis-(4-fluoro-phenyl)-thiazol-4-yl]-[(S)-2-(5-chloro-6-trifluoromethyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 1.30$; $[M+H]^+ = 603.3$
5.292	[(S)-2-(5-Chloro-6-trifluoromethyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-[5-(4-fluoro-phenyl)-2-(3-methoxy-phenyl)-thiazol-4-yl]-methanone. LC-MS: $t_R = 1.30$; $[M+H]^+ = 615.3$

5.293	[(S)-2-(5-Chloro-6-trifluoromethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-[5-(4-fluoro-phenyl)-2-(4-methoxy-phenyl)-thiazol-4-yl]-methanone. LC-MS: $t_R = 1.28$; $[M+H]^+ = 615.3$
5.294	[5-(4-Fluoro-phenyl)-2-(2-fluoro-phenyl)-thiazol-4-yl]-[(S)-2-methyl-2-(6-trifluoromethyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 1.15$; $[M+H]^+ = 569.3$
5.295	3'-[(S)-2-Methyl-2-(5-trifluoromethyl-1H-benzimidazol-2-yl)-pyrrolidine-1-carbonyl]-4'-[1,2,3]triazol-2-yl-biphenyl-3-carbonitrile. LC-MS: $t_R = 0.99$; $[M+H]^+ = 542.3$
5.296	[(S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(3-fluoro-4,3'-dimethoxy-biphenyl-2-yl)-methanone. LC-MS: $t_R = 0.95$; $[M+H]^+ = 508.3$
5.297	[(S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(3-fluoro-4,4'-dimethoxy-biphenyl-2-yl)-methanone. LC-MS: $t_R = 0.93$; $[M+H]^+ = 508.3$
5.298	[(S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(3,2'-difluoro-4-methoxy-biphenyl-2-yl)-methanone. LC-MS: $t_R = 0.95$; $[M+H]^+ = 496.3$
5.299	[(S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(3,4'-difluoro-4-methoxy-biphenyl-2-yl)-methanone. LC-MS: $t_R = 0.96$; $[M+H]^+ = 496.3$
5.300	[(S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(3,3'-difluoro-4-methoxy-biphenyl-2-yl)-methanone. LC-MS: $t_R = 0.96$; $[M+H]^+ = 496.3$
5.301	[(S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(3-fluoro-4-methoxy-3'-methyl-biphenyl-2-yl)-methanone. LC-MS: $t_R = 1.00$; $[M+H]^+ = 492.3$
5.302	[(S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(3-fluoro-4-methoxy-4'-methyl-biphenyl-2-yl)-methanone. LC-MS: $t_R = 1.01$; $[M+H]^+ = 492.3$
5.303	[(S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(3-fluoro-4-methoxy-3'-trifluoromethyl-biphenyl-2-yl)-methanone. LC-MS: $t_R = 1.05$; $[M+H]^+ = 546.3$
5.304	(6-Benzo[1,3]dioxol-5-yl-2-fluoro-3-methoxy-phenyl)-[(S)-2-(5-chloro-4-methyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.93$; $[M+H]^+ = 522.3$
5.305	[(S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(3-fluoro-4-methoxy-4'-trifluoromethyl-biphenyl-2-yl)-methanone. LC-MS: $t_R = 1.06$; $[M+H]^+ = 546.3$
5.306	[(S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(3-fluoro-4-methoxy-4'-trifluoromethoxy-biphenyl-2-yl)-methanone. LC-MS: $t_R = 1.07$; $[M+H]^+ = 562.4$
5.307	2'-[(S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidine-1-carbonyl]-3'-fluoro-4'-methoxy-biphenyl-3-carbonitrile. LC-MS: $t_R = 0.92$; $[M+H]^+ = 503.3$
5.308	2'-[(S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidine-1-carbonyl]-3'-fluoro-4'-methoxy-biphenyl-4-carbonitrile. LC-MS: $t_R = 0.93$; $[M+H]^+ = 503.3$

5.309	[(S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(3-fluoro-4-methoxy-3'-trifluoromethoxy-biphenyl-2-yl)-methanone. LC-MS: $t_R = 1.07$; $[M+H]^+ = 562.3$
5.310	[(S)-2-(6-Bromo-4-methyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(2-methyl-5-phenyl-pyridin-4-yl)-methanone. LC-MS: $t_R = 0.75$; $[M+H]^+ = 489.3$
6.1	[5-(4-Fluoro-phenyl)-2-methyl-thiazol-4-yl]-[(S)-2-(5-methoxy-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.74$; $[M+H]^+ = 449.2$
6.2	[(S)-2-(5-Methoxy-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-[5-(4-methoxy-phenyl)-2-methyl-thiazol-4-yl]-methanone. LC-MS: $t_R = 0.73$; $[M+H]^+ = 461.2$
6.3	[(S)-2-(5-Methoxy-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(2-methyl-5-phenyl-thiazol-4-yl)-methanone. LC-MS: $t_R = 0.72$; $[M+H]^+ = 431.2$
6.4	[(S)-2-(5-Methoxy-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(3-p-tolyl-pyrazin-2-yl)-methanone. LC-MS: $t_R = 0.67$; $[M+H]^+ = 426.2$
6.5	[5-(4-Chloro-phenyl)-2-methyl-thiazol-4-yl]-[(S)-2-(5-methoxy-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.79$; $[M+H]^+ = 465.2$
6.6	[(S)-2-(5-Methoxy-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(3-m-tolyl-pyrazin-2-yl)-methanone. LC-MS: $t_R = 0.67$; $[M+H]^+ = 426.2$
6.7	[(S)-2-(5-Methoxy-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(2-methyl-5-m-tolyl-thiazol-4-yl)-methanone. LC-MS: $t_R = 0.77$; $[M+H]^+ = 445.2$
6.8	[5-(3-Chloro-phenyl)-2-methyl-thiazol-4-yl]-[(S)-2-(5-methoxy-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.79$; $[M+H]^+ = 465.1$
6.9	[5-(3,4-Dimethyl-phenyl)-2-methyl-thiazol-4-yl]-[(S)-2-(5-methoxy-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.81$; $[M+H]^+ = 459.3$
6.10	(2-Cyclopropyl-5-phenyl-thiazol-4-yl)-[(S)-2-(5-methoxy-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.80$; $[M+H]^+ = 457.2$
6.11	[(S)-2-(5-Methoxy-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(2-methyl-4-m-tolyl-pyrimidin-5-yl)-methanone. LC-MS: $t_R = 0.68$; $[M+H]^+ = 440.2$
6.12	[(S)-2-(5-Methoxy-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.67$; $[M+H]^+ = 415.2$
6.13	[(S)-2-(5-Methoxy-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.61$; $[M+H]^+ = 401.2$
6.14	[(S)-2-(5-Methoxy-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(3'-methyl-biphenyl-2-yl)-methanone. LC-MS: $t_R = 0.79$; $[M+H]^+ = 424.2$

6.15	[(S)-2-(5-Methoxy-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(4'-methyl-biphenyl-2-yl)-methanone. LC-MS: $t_R = 0.79$; $[M+H]^+ = 424.2$
6.16	[(S)-2-(5-Methoxy-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(2-methyl-4-phenyl-pyrimidin-5-yl)-methanone. LC-MS: $t_R = 0.62$; $[M+H]^+ = 426.2$
6.17	[(S)-2-(5-Methoxy-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(4-m-tolyl-pyrimidin-5-yl)-methanone. LC-MS: $t_R = 0.64$; $[M+H]^+ = 426.3$
6.18	(5-Fluoro-2-[1,2,3]triazol-2-yl-phenyl)-[(S)-2-(5-methoxy-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.64$; $[M+H]^+ = 419.2$
6.19	(2-Fluoro-6-[1,2,3]triazol-2-yl-phenyl)-[(S)-2-(5-methoxy-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.63$; $[M+H]^+ = 419.2$
6.20	[(S)-2-(5-Methoxy-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-[2-(3-methyl-[1,2,4]oxadiazol-5-yl)-phenyl]-methanone. LC-MS: $t_R = 0.62$; $[M+H]^+ = 416.2$
6.21	[(S)-2-(5-Methoxy-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-[2-(2-methyl-thiazol-4-yl)-phenyl]-methanone. LC-MS: $t_R = 0.67$; $[M+H]^+ = 431.2$
6.22	[5-(4-Bromo-phenyl)-2-methyl-thiazol-4-yl]-[(S)-2-(5-methoxy-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.80$; $[M+H]^+ = 509.1$
6.23	[5-(2,3-Dichloro-phenyl)-2-methyl-thiazol-4-yl]-[(S)-2-(5-methoxy-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.84$; $[M+H]^+ = 499.2$
6.24	[5-(3-Bromo-4-fluoro-phenyl)-2-methyl-thiazol-4-yl]-[(S)-2-(5-methoxy-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.81$; $[M+H]^+ = 527.1$
6.25	[2-Cyclopropyl-5-(3-fluoro-4-methyl-phenyl)-thiazol-4-yl]-[(S)-2-(5-methoxy-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.87$; $[M+H]^+ = 489.2$
6.26	(2-Cyclopropyl-5-m-tolyl-thiazol-4-yl)-[(S)-2-(5-methoxy-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.85$; $[M+H]^+ = 471.3$
6.27	[2-Dimethylamino-5-(3,4-dimethyl-phenyl)-thiazol-4-yl]-[(S)-2-(5-methoxy-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.85$; $[M+H]^+ = 488.3$
6.28	[5-(3-Bromo-phenyl)-2-cyclopropyl-thiazol-4-yl]-[(S)-2-(5-methoxy-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.88$; $[M+H]^+ = 535.1$
6.29	[5-(3-Fluoro-phenyl)-2-pyrrolidin-1-yl-thiazol-4-yl]-[(S)-2-(5-methoxy-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.86$; $[M+H]^+ = 504.3$
6.30	[(S)-2-(5-Methoxy-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-[2-methyl-5-(3-trifluoromethyl-phenyl)-thiazol-4-yl]-methanone. LC-MS: $t_R = 0.83$; $[M+H]^+ = 499.2$

6.31	[2-Cyclopropyl-5-(3-methoxy-phenyl)-thiazol-4-yl]-[(S)-2-(5-methoxy-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.81$; $[M+H]^+ = 487.2$
6.32	[5-(3-Chloro-phenyl)-thiazol-4-yl]-[(S)-2-(5-methoxy-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.74$; $[M+H]^+ = 451.1$
6.33	[5-(3-Fluoro-phenyl)-2-methyl-thiazol-4-yl]-[(S)-2-(5-methoxy-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.74$; $[M+H]^+ = 449.2$
6.34	[5-(2-Chloro-phenyl)-2-methyl-thiazol-4-yl]-[(S)-2-(5-methoxy-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.79$; $[M+H]^+ = 465.2$
6.35	(2-Cyclopropyl-5-p-tolyl-thiazol-4-yl)-[(S)-2-(5-methoxy-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.85$; $[M+H]^+ = 471.3$
6.36	[2-(3,4-Dimethyl-phenyl)-5-methyl-thiophen-3-yl]-[(S)-2-(5-methoxy-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.86$; $[M+H]^+ = 458.2$
6.37	[5-(3,5-Difluoro-phenyl)-2-methyl-thiazol-4-yl]-[(S)-2-(5-methoxy-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.76$; $[M+H]^+ = 467.2$
6.38	[(S)-2-(5-Methoxy-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(2-m-tolyl-thiophen-3-yl)-methanone. LC-MS: $t_R = 0.77$; $[M+H]^+ = 430.1$
6.39	[2-(3,4-Dimethyl-phenyl)-thiophen-3-yl]-[(S)-2-(5-methoxy-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.81$; $[M+H]^+ = 444.2$
6.40	[(S)-2-(4-Methyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(2-methyl-5-phenyl-thiazol-4-yl)-methanone. LC-MS: $t_R = 0.74$; $[M+H]^+ = 415.2$
6.41	[(S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(2-methyl-5-phenyl-thiazol-4-yl)-methanone. LC-MS: $t_R = 0.86$; $[M+H]^+ = 449.2$
6.42	[(S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.80$; $[M+H]^+ = 433.2$
6.43	[(S)-2-(1H-Benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-[5-(3-chloro-phenyl)-2-methyl-thiazol-4-yl]-methanone. LC-MS: $t_R = 0.78$; $[M+H]^+ = 435.1$
6.44	[5-(3-Chloro-phenyl)-2-methyl-thiazol-4-yl]-[(S)-2-(4-methyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.81$; $[M+H]^+ = 449.2$
6.45	[(S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-[5-(3-chloro-phenyl)-2-methyl-thiazol-4-yl]-methanone. LC-MS: $t_R = 0.93$; $[M+H]^+ = 483.1$
6.46	[5-(3,4-Dimethyl-phenyl)-2-methyl-thiazol-4-yl]-[(S)-2-(6-methoxy-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.81$; $[M+H]^+ = 459.2$

6.47	[(S)-2-(4,5-Dimethyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-[5-(3,4-dimethyl-phenyl)-2-methyl-thiazol-4-yl]-methanone. LC-MS: $t_R = 0.86$; $[M+H]^+ = 457.2$
6.48	[(S)-2-(4,7-Dimethoxy-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-[5-(3,4-dimethyl-phenyl)-2-methyl-thiazol-4-yl]-methanone. LC-MS: $t_R = 0.85$; $[M+H]^+ = 489.3$
6.49	[5-(3,4-Dimethyl-phenyl)-2-methyl-thiazol-4-yl]-[(S)-4-methylene-2-(4-methylsulfanyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.88$; $[M+H]^+ = 475.3$
6.50	[(S)-2-(4-Bromo-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-[5-(3,4-dimethyl-phenyl)-2-methyl-thiazol-4-yl]-methanone. LC-MS: $t_R = 0.97$; $[M+H]^+ = 507.2$
6.51	[(S)-2-(4-Chloro-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-[5-(3,4-dimethyl-phenyl)-2-methyl-thiazol-4-yl]-methanone. LC-MS: $t_R = 0.96$; $[M+H]^+ = 463.2$
6.52	[5-(3,4-Dimethyl-phenyl)-2-methyl-thiazol-4-yl]-[(S)-4-methylene-2-(5-methylsulfanyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.86$; $[M+H]^+ = 475.3$
6.53	[5-(3,4-Dimethyl-phenyl)-2-methyl-thiazol-4-yl]-[(S)-2-(4-methyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.83$; $[M+H]^+ = 443.2$
6.54	[(S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-[5-(3,4-dimethyl-phenyl)-2-methyl-thiazol-4-yl]-methanone. LC-MS: $t_R = 0.94$; $[M+H]^+ = 477.2$
6.55	[(S)-2-(5-Chloro-6-methyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-[5-(3,4-dimethyl-phenyl)-2-methyl-thiazol-4-yl]-methanone. LC-MS: $t_R = 0.92$; $[M+H]^+ = 477.2$
6.56	[(S)-2-(5,6-Dimethyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-[5-(3,4-dimethyl-phenyl)-2-methyl-thiazol-4-yl]-methanone. LC-MS: $t_R = 0.86$; $[M+H]^+ = 457.3$
6.57	[(S)-2-(5-tert-Butyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-[5-(3,4-dimethyl-phenyl)-2-methyl-thiazol-4-yl]-methanone. LC-MS: $t_R = 0.95$; $[M+H]^+ = 485.3$
6.58	[(S)-2-(4,6-Dimethyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-[5-(3,4-dimethyl-phenyl)-2-methyl-thiazol-4-yl]-methanone. LC-MS: $t_R = 0.87$; $[M+H]^+ = 457.2$
6.59	[(S)-4-Methylene-2-(7-methylsulfanyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-[5-methyl-2-[1,2,3]triazol-2-yl-phenyl]-methanone. LC-MS: $t_R = 0.74$; $[M+H]^+ = 431.2$
6.60	[(S)-4-Methylene-2-(5-methylsulfanyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-[5-methyl-2-[1,2,3]triazol-2-yl-phenyl]-methanone. LC-MS: $t_R = 0.73$; $[M+H]^+ = 431.2$
6.61	[(S)-2-(5,6-Dimethyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-[5-methyl-2-[1,2,3]triazol-2-yl-phenyl]-methanone. LC-MS: $t_R = 0.73$; $[M+H]^+ = 413.2$
6.62	[(S)-2-(4,6-Dimethyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-[5-methyl-2-[1,2,3]triazol-2-yl-phenyl]-methanone. LC-MS: $t_R = 0.72$; $[M+H]^+ = 413.2$

6.63	[(S)-2-(4,5-Dimethyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.72$; $[M+H]^+ = 413.2$
6.64	[(S)-2-(4-Bromo-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.84$; $[M+H]^+ = 463.1$
6.65	[(S)-2-(5-Chloro-6-methyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.80$; $[M+H]^+ = 433.2$
6.66	[(S)-2-(5-tert-Butyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.82$; $[M+H]^+ = 441.3$
6.67	[(S)-2-(5-Chloro-6-trifluoromethyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 1.05$; $[M+H]^+ = 487.2$
6.68	[(S)-2-(4,7-Dimethoxy-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.73$; $[M+H]^+ = 445.2$
6.69	[(S)-2-(7-Chloro-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.82$; $[M+H]^+ = 419.1$
6.71	rac-[2-(3,4-Dimethyl-phenyl)-5-methyl-thiophen-3-yl]-[2-(5-methoxy-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.86$; $[M+H]^+ = 458.2$
6.72	rac-[2-(3,4-Dimethyl-phenyl)-5-methyl-thiophen-3-yl]-[4-methylene-2-(7-methylsulfanyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.94$; $[M+H]^+ = 474.2$
6.73	rac-[2-(3,4-Dimethyl-phenyl)-5-methyl-thiophen-3-yl]-[4-methylene-2-(5-methylsulfanyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.92$; $[M+H]^+ = 474.2$
6.74	rac-[2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-[2-(3,4-dimethyl-phenyl)-5-methyl-thiophen-3-yl]-methanone. LC-MS: $t_R = 0.99$; $[M+H]^+ = 476.2$
6.75	rac-[2-(5,6-Dimethyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-[2-(3,4-dimethyl-phenyl)-5-methyl-thiophen-3-yl]-methanone. LC-MS: $t_R = 0.76$; $[M+H]^+ = 456.1$
6.76	rac-[2-(4,6-Dimethyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-[2-(3,4-dimethyl-phenyl)-5-methyl-thiophen-3-yl]-methanone. LC-MS: $t_R = 0.91$; $[M+H]^+ = 456.3$
6.77	rac-[2-(4,5-Dimethyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-[2-(3,4-dimethyl-phenyl)-5-methyl-thiophen-3-yl]-methanone. LC-MS: $t_R = 0.90$; $[M+H]^+ = 456.2$
6.78	rac-[2-(4-Bromo-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-[2-(3,4-dimethyl-phenyl)-5-methyl-thiophen-3-yl]-methanone. LC-MS: $t_R = 1.06$; $[M+H]^+ = 506.2$
6.79	rac-[2-(3,4-Dimethyl-phenyl)-5-methyl-thiophen-3-yl]-[2-(4-methyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.88$; $[M+H]^+ = 442.2$

6.80	rac-[2-(5-Chloro-6-methyl-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-[2-(3,4-dimethyl-phenyl)-5-methyl-thiophen-3-yl]-methanone. LC-MS: $t_R = 1.00$; $[M+H]^+ = 476.2$
6.81	rac-[2-(5-tert-Butyl-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-[2-(3,4-dimethyl-phenyl)-5-methyl-thiophen-3-yl]-methanone. LC-MS: $t_R = 0.99$; $[M+H]^+ = 484.3$
6.82	rac-[2-(5-Chloro-6-trifluoromethyl-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-[2-(3,4-dimethyl-phenyl)-5-methyl-thiophen-3-yl]-methanone. LC-MS: $t_R = 1.24$; $[M+H]^+ = 530.2$
6.83	rac-[2-(4,7-Dimethoxy-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-[2-(3,4-dimethyl-phenyl)-5-methyl-thiophen-3-yl]-methanone. LC-MS: $t_R = 0.90$; $[M+H]^+ = 488.3$
6.84	rac-[2-(7-Chloro-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-[2-(3,4-dimethyl-phenyl)-5-methyl-thiophen-3-yl]-methanone. LC-MS: $t_R = 1.04$; $[M+H]^+ = 462.2$
6.85	rac-[2-(3,4-Dimethyl-phenyl)-5-methyl-thiophen-3-yl]-[4-methylene-2-(1H-naphtho[2,3-d]imidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.94$; $[M+H]^+ = 478.3$
6.86	rac-(2-Cyclopropyl-5-m-tolyl-thiazol-4-yl)-[2-(5-methoxy-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.85$; $[M+H]^+ = 471.3$
6.87	rac-(2-Cyclopropyl-5-m-tolyl-thiazol-4-yl)-[4-methylene-2-(7-methylsulfanyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.92$; $[M+H]^+ = 487.2$
6.88	rac-(2-Cyclopropyl-5-m-tolyl-thiazol-4-yl)-[4-methylene-2-(5-methylsulfanyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.90$; $[M+H]^+ = 487.2$
6.89	rac-[2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(2-cyclopropyl-5-m-tolyl-thiazol-4-yl)-methanone. LC-MS: $t_R = 0.97$; $[M+H]^+ = 489.2$
6.90	rac-(2-Cyclopropyl-5-m-tolyl-thiazol-4-yl)-[2-(5,6-dimethyl-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.90$; $[M+H]^+ = 469.3$
6.91	rac-(2-Cyclopropyl-5-m-tolyl-thiazol-4-yl)-[2-(4,6-dimethyl-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.91$; $[M+H]^+ = 469.3$
6.92	rac-(2-Cyclopropyl-5-m-tolyl-thiazol-4-yl)-[2-(4,5-dimethyl-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.90$; $[M+H]^+ = 469.3$
6.93	rac-[2-(4-Bromo-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(2-cyclopropyl-5-m-tolyl-thiazol-4-yl)-methanone. LC-MS: $t_R = 1.01$; $[M+H]^+ = 519.1$
6.94	rac-(2-Cyclopropyl-5-m-tolyl-thiazol-4-yl)-[2-(4-methyl-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.87$; $[M+H]^+ = 455.3$
6.95	rac-[2-(5-Chloro-6-methyl-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(2-cyclopropyl-5-m-tolyl-thiazol-4-yl)-methanone. LC-MS: $t_R = 0.96$; $[M+H]^+ = 489.2$

6.96	rac-[2-(5-tert-Butyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(2-cyclopropyl-5-m-tolyl-thiazol-4-yl)-methanone. LC-MS: $t_R = 0.98$; $[M+H]^+ = 497.3$
6.97	rac-[2-(5-Chloro-6-trifluoromethyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(2-cyclopropyl-5-m-tolyl-thiazol-4-yl)-methanone. LC-MS: $t_R = 1.19$; $[M+H]^+ = 543.1$
6.98	rac-(2-Cyclopropyl-5-m-tolyl-thiazol-4-yl)-[2-(4,7-dimethoxy-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.89$; $[M+H]^+ = 501.3$
6.99	rac-[2-(7-Chloro-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(2-cyclopropyl-5-m-tolyl-thiazol-4-yl)-methanone. LC-MS: $t_R = 1.00$; $[M+H]^+ = 475.2$
6.100	[5-(4-Bromo-phenyl)-2-methyl-thiazol-4-yl]-[(S)-2-(7-methyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.83$; $[M+H]^+ = 493.2$
6.101	[(S)-2-(7-Methyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(2-methyl-5-m-tolyl-thiazol-4-yl)-methanone. LC-MS: $t_R = 0.80$; $[M+H]^+ = 429.2$
6.102	(5-Chloro-2-[1,2,3]triazol-2-yl-phenyl)-[(S)-2-(7-methyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.71$; $[M+H]^+ = 419.2$
6.103	[5-(6-Methoxy-pyridin-3-yl)-2-methyl-thiazol-4-yl]-[(S)-2-(7-methyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.71$; $[M+H]^+ = 446.2$
6.104	[5-(3-Bromo-phenyl)-2-methyl-thiazol-4-yl]-[(S)-2-(7-methyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.83$; $[M+H]^+ = 493.1$
6.105	[(S)-2-(7-Methyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(6-methyl-3-pyrazol-1-yl-pyridin-2-yl)-methanone. LC-MS: $t_R = 0.65$; $[M+H]^+ = 399.2$
6.106	[5-(4-Fluoro-phenyl)-2-methyl-thiazol-4-yl]-[(S)-2-(7-methyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.76$; $[M+H]^+ = 433.2$
6.107	[5-(3,5-Dimethyl-phenyl)-2-methyl-thiazol-4-yl]-[(S)-2-(7-methyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.85$; $[M+H]^+ = 443.2$
6.108	(4,5-Dimethyl-2-[1,2,3]triazol-2-yl-phenyl)-[(S)-2-(7-methyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.72$; $[M+H]^+ = 413.2$
6.109	[5-(4-Chloro-phenyl)-2-methyl-thiazol-4-yl]-[(S)-2-(7-methyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.81$; $[M+H]^+ = 449.1$
6.110	[5-(3-Bromo-4-fluoro-phenyl)-2-methyl-thiazol-4-yl]-[(S)-2-(7-methyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.84$; $[M+H]^+ = 511.1$
6.111	(5-Methoxy-2-[1,2,3]triazol-2-yl-phenyl)-[(S)-2-(7-methyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.65$; $[M+H]^+ = 415.2$

6.112	[(S)-2-(7-Methyl-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-[2-methyl-5-(3-trifluoromethyl-phenyl)-thiazol-4-yl]-methanone. LC-MS: $t_R = 0.85$; $[M+H]^+ = 483.2$
6.113	[5-(3,4-Dichloro-phenyl)-2-methyl-thiazol-4-yl]-[(S)-2-(7-methyl-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.87$; $[M+H]^+ = 483.1$
6.114	[5-(3-Methoxy-phenyl)-2-methyl-thiazol-4-yl]-[(S)-2-(7-methyl-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.75$; $[M+H]^+ = 445.2$
6.115	[(S)-2-(7-Methyl-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(6-methyl-3-[1,2,3]triazol-2-yl-pyridin-2-yl)-methanone. LC-MS: $t_R = 0.65$; $[M+H]^+ = 400.2$
6.116	[5-(4-Bromo-phenyl)-2-methyl-thiazol-4-yl]-[(S)-2-(6-chloro-7-methyl-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.95$; $[M+H]^+ = 527.2$
6.117	[(S)-2-(6-Chloro-7-methyl-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-[5-(6-methoxy-pyridin-3-yl)-2-methyl-thiazol-4-yl]-methanone. LC-MS: $t_R = 0.82$; $[M+H]^+ = 480.2$
6.118	[(S)-2-(6-Chloro-7-methyl-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-[5-(4-fluoro-phenyl)-2-methyl-thiazol-4-yl]-methanone. LC-MS: $t_R = 0.88$; $[M+H]^+ = 467.2$
6.119	[(S)-2-(6-Chloro-7-methyl-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-[5-(4-chloro-phenyl)-2-methyl-thiazol-4-yl]-methanone. LC-MS: $t_R = 0.93$; $[M+H]^+ = 483.1$
6.120	[(S)-2-(6-Chloro-7-methyl-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-[2-methyl-5-(3-trifluoromethyl-phenyl)-thiazol-4-yl]-methanone. LC-MS: $t_R = 0.96$; $[M+H]^+ = 517.2$
6.121	[(S)-2-(6-Chloro-7-methyl-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-[5-(3-methoxy-phenyl)-2-methyl-thiazol-4-yl]-methanone. LC-MS: $t_R = 0.86$; $[M+H]^+ = 479.2$
6.122	[(S)-2-(6-Chloro-7-methyl-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(2-methyl-5-m-tolyl-thiazol-4-yl)-methanone. LC-MS: $t_R = 0.90$; $[M+H]^+ = 463.2$
6.123	[5-(3-Bromo-phenyl)-2-methyl-thiazol-4-yl]-[(S)-2-(6-chloro-7-methyl-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.94$; $[M+H]^+ = 527.1$
6.124	[(S)-2-(6-Chloro-7-methyl-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-[5-(3,5-dimethyl-phenyl)-2-methyl-thiazol-4-yl]-methanone. LC-MS: $t_R = 0.95$; $[M+H]^+ = 477.2$
6.125	[5-(3-Bromo-4-fluoro-phenyl)-2-methyl-thiazol-4-yl]-[(S)-2-(6-chloro-7-methyl-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.95$; $[M+H]^+ = 545.1$
6.126	[(S)-2-(6-Chloro-7-methyl-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-[5-(3,4-dichloro-phenyl)-2-methyl-thiazol-4-yl]-methanone. LC-MS: $t_R = 0.99$; $[M+H]^+ = 517$
6.127	[(S)-2-(6-Chloro-7-methyl-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(6-methyl-3-[1,2,3]triazol-2-yl-pyridin-2-yl)-methanone. LC-MS: $t_R = 0.74$; $[M+H]^+ = 434.2$

6.128	[(S)-2-(6-Chloro-7-methyl-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(5-chloro-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.84$; $[M+H]^+ = 453.1$
6.129	[(S)-2-(6-Chloro-7-methyl-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(2-[1,2,3]triazol-2-yl-5-trifluoromethyl-phenyl)-methanone. LC-MS: $t_R = 0.89$; $[M+H]^+ = 487.2$
6.130	[(S)-2-(6-Chloro-7-methyl-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(4,5-dimethyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.84$; $[M+H]^+ = 447.2$
6.131	[(S)-2-(6-Chloro-7-methyl-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(5-methoxy-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.78$; $[M+H]^+ = 449.2$
6.132	[(S)-2-(6-Chloro-7-methyl-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(2-fluoro-3-methyl-6-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.83$; $[M+H]^+ = 451.2$
6.133	[(S)-2-(6-Chloro-7-methyl-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(2-fluoro-3-methoxy-6-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.80$; $[M+H]^+ = 467.2$
6.134	[(S)-2-(6-Chloro-7-methyl-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(5-methyl-2-pyrazol-1-yl-phenyl)-methanone. LC-MS: $t_R = 0.81$; $[M+H]^+ = 432.2$
6.135	[(S)-2-(6-Chloro-7-methyl-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(6-methyl-3-pyrazol-1-yl-pyridin-2-yl)-methanone. LC-MS: $t_R = 0.75$; $[M+H]^+ = 433.2$
6.136	(5-Chloro-2-[1,2,3]triazol-2-yl-phenyl)-[(S)-2-(6-methoxy-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.70$; $[M+H]^+ = 435.1$
6.137	[(S)-2-(6-Methoxy-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(5-methyl-2-pyrazol-1-yl-phenyl)-methanone. LC-MS: $t_R = 0.67$; $[M+H]^+ = 414.2$
6.138	[(S)-2-(6-Methoxy-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(6-methyl-3-pyrazol-1-yl-pyridin-2-yl)-methanone. LC-MS: $t_R = 0.63$; $[M+H]^+ = 415.2$
6.139	(4,5-Dimethyl-2-[1,2,3]triazol-2-yl-phenyl)-[(S)-2-(6-methoxy-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.72$; $[M+H]^+ = 429.2$
6.140	[(S)-2-(6-Methoxy-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(5-methoxy-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.65$; $[M+H]^+ = 431.2$
6.141	(2-Fluoro-3-methyl-6-[1,2,3]triazol-2-yl-phenyl)-[(S)-2-(6-methoxy-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.69$; $[M+H]^+ = 433.2$
6.142	[(S)-2-(6-Chloro-5-trifluoromethyl-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(5-(3,4-dimethyl-phenyl)-2-methyl-thiazol-4-yl)-methanone. LC-MS: $t_R = 1.15$; $[M+H]^+ = 531.2$
6.143	rac-[2-(5-Methoxy-1H-benzoimidazol-2-yl)-2-methyl-4-methylene-pyrrolidin-1-yl]-(2-methyl-5-m-tolyl-thiazol-4-yl)-methanone. LC-MS: $t_R = 0.85$; $[M+H]^+ = 459.3$

6.144	rac-[2-Dimethylamino-5-(3,4-dimethyl-phenyl)-thiazol-4-yl]-[2-(5-methoxy-1H-benzimidazol-2-yl)-2-methyl-4-methylene-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.96$; $[M+H]^+ = 502.3$
6.145	rac-[5-(3,4-Dimethyl-phenyl)-2-methyl-thiazol-4-yl]-[2-(5-methoxy-1H-benzimidazol-2-yl)-2-methyl-4-methylene-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.90$; $[M+H]^+ = 473.3$
6.146	rac-[5-(4-Chloro-phenyl)-2-methyl-thiazol-4-yl]-[2-(5-methoxy-1H-benzimidazol-2-yl)-2-methyl-4-methylene-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.86$; $[M+H]^+ = 479.2$
6.147	rac-[2-(5-Methoxy-1H-benzimidazol-2-yl)-2-methyl-4-methylene-pyrrolidin-1-yl]-[5-(4-methoxy-phenyl)-2-methyl-thiazol-4-yl]-methanone. LC-MS: $t_R = 0.80$; $[M+H]^+ = 475.3$
6.148	rac-[2-(5-Methoxy-1H-benzimidazol-2-yl)-2-methyl-4-methylene-pyrrolidin-1-yl]-(2-methyl-5-p-tolyl-thiazol-4-yl)-methanone. LC-MS: $t_R = 0.85$; $[M+H]^+ = 459.3$
6.149	rac-(3',4'-Dimethyl-biphenyl-2-yl)-[2-(5-methoxy-1H-benzimidazol-2-yl)-2-methyl-4-methylene-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.90$; $[M+H]^+ = 452.3$
6.150	rac-Biphenyl-2-yl-[2-(5-methoxy-1H-benzimidazol-2-yl)-2-methyl-4-methylene-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.80$; $[M+H]^+ = 424.3$
6.151	rac-(4'-Fluoro-3'-methyl-biphenyl-2-yl)-[2-(5-methoxy-1H-benzimidazol-2-yl)-2-methyl-4-methylene-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.87$; $[M+H]^+ = 456.3$
6.152	rac-[5-(3-Chloro-phenyl)-2-dimethylamino-thiazol-4-yl]-[2-(5-methoxy-1H-benzimidazol-2-yl)-2-methyl-4-methylene-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.92$; $[M+H]^+ = 508.2$
6.153	rac-[5-(3-Chloro-phenyl)-2-methyl-thiazol-4-yl]-[2-(5-methoxy-1H-benzimidazol-2-yl)-2-methyl-4-methylene-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.86$; $[M+H]^+ = 479.2$
6.154	rac-[2-(5-Methoxy-1H-benzimidazol-2-yl)-2-methyl-4-methylene-pyrrolidin-1-yl]-(2-methyl-5-phenyl-thiazol-4-yl)-methanone. LC-MS: $t_R = 0.79$; $[M+H]^+ = 445.3$
6.155	rac-[2-(5-Methoxy-1H-benzimidazol-2-yl)-2-methyl-4-methylene-pyrrolidin-1-yl]-[2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.66$; $[M+H]^+ = 415.3$
6.156	rac-[2-(5-Methoxy-1H-benzimidazol-2-yl)-2-methyl-4-methylene-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.72$; $[M+H]^+ = 429.3$
6.157	rac-(5-Chloro-2-[1,2,3]triazol-2-yl-phenyl)-[2-(5-methoxy-1H-benzimidazol-2-yl)-2-methyl-4-methylene-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.75$; $[M+H]^+ = 449.2$
6.158	rac-[2-(5-Methoxy-1H-benzimidazol-2-yl)-2-methyl-4-methylene-pyrrolidin-1-yl]-(5-methoxy-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.70$; $[M+H]^+ = 445.3$
6.159	rac-(4,5-Dimethyl-2-[1,2,3]triazol-2-yl-phenyl)-[2-(5-methoxy-1H-benzimidazol-2-yl)-2-methyl-4-methylene-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.78$; $[M+H]^+ = 443.3$

6.160	[(S)-2-(5-tert-Butyl-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(5-fluoro-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.81$; $[M+H]^+ = 445.3$
6.161	[(S)-2-(5-tert-Butyl-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(5-chloro-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.86$; $[M+H]^+ = 461.3$
6.162	[(S)-2-(5-tert-Butyl-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.79$; $[M+H]^+ = 427.3$
6.163	[(S)-2-(5-tert-Butyl-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(2-fluoro-3-methoxy-6-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.81$; $[M+H]^+ = 475.3$
6.164	[(S)-2-(5-tert-Butyl-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(2-fluoro-3-methyl-6-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.84$; $[M+H]^+ = 459.3$
6.165	[(S)-2-(5-tert-Butyl-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(4,5-dimethyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.87$; $[M+H]^+ = 455.3$
6.166	[(S)-2-(5-tert-Butyl-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(4-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.84$; $[M+H]^+ = 441.4$
6.167	[(S)-2-(5-tert-Butyl-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(2-methyl-6-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.83$; $[M+H]^+ = 441.3$
6.168	[(S)-2-(5-tert-Butyl-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(2-[1,2,3]triazol-2-yl-5-trifluoromethyl-phenyl)-methanone. LC-MS: $t_R = 0.91$; $[M+H]^+ = 495.3$
6.169	[(S)-2-(5-tert-Butyl-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(2-[1,2,3]triazol-2-yl-5-trifluoromethoxy-phenyl)-methanone. LC-MS: $t_R = 0.93$; $[M+H]^+ = 511.3$
6.170	[(S)-2-(5-tert-Butyl-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(4,5-dimethoxy-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.78$; $[M+H]^+ = 487.3$
6.171	[(S)-2-(5-tert-Butyl-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(5-methyl-2-pyrazol-1-yl-phenyl)-methanone. LC-MS: $t_R = 0.84$; $[M+H]^+ = 440.3$
6.172	[(S)-2-(5-tert-Butyl-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(2-pyrazol-1-yl-phenyl)-methanone. LC-MS: $t_R = 0.8$; $[M+H]^+ = 426.3$
6.173	[(S)-2-(5-tert-Butyl-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(6-methyl-3-pyrazol-1-yl-pyridin-2-yl)-methanone. LC-MS: $t_R = 0.80$; $[M+H]^+ = 441.3$
6.174	[(S)-2-(5-tert-Butyl-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(6-methyl-3-[1,2,3]triazol-2-yl-pyridin-2-yl)-methanone. LC-MS: $t_R = 0.80$; $[M+H]^+ = 442.3$
6.175	[(S)-2-(5-Chloro-6-trifluoromethyl-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(2-fluoro-6-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 1.05$; $[M+H]^+ = 491.2$

6.176	[(S)-2-(5-Chloro-6-trifluoromethyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(5-fluoro-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 1.04$; $[M+H]^+ = 491.2$
6.177	(5-Chloro-2-[1,2,3]triazol-2-yl-phenyl)-[(S)-2-(5-chloro-6-trifluoromethyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 1.09$; $[M+H]^+ = 507.2$
6.178	[(S)-2-(5-Chloro-6-trifluoromethyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 1.01$; $[M+H]^+ = 473.2$
6.179	[(S)-2-(5-Chloro-6-trifluoromethyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(2-fluoro-3-methoxy-6-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 1.06$; $[M+H]^+ = 521.2$
6.180	[(S)-2-(5-Chloro-6-trifluoromethyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(2-fluoro-3-methyl-6-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 1.11$; $[M+H]^+ = 505.2$
6.181	[(S)-2-(5-Chloro-6-trifluoromethyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(4,5-dimethyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 1.1$; $[M+H]^+ = 501.2$
6.182	[(S)-2-(5-Chloro-6-trifluoromethyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(4-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 1.07$; $[M+H]^+ = 487.2$
6.183	[(S)-2-(5-Chloro-6-trifluoromethyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(2-methyl-6-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 1.09$; $[M+H]^+ = 487.3$
6.184	[(S)-2-(5-Chloro-6-trifluoromethyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(2-[1,2,3]triazol-2-yl-5-trifluoromethyl-phenyl)-methanone. LC-MS: $t_R = 1.12$; $[M+H]^+ = 541.1$
6.185	[(S)-2-(5-Chloro-6-trifluoromethyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(2-[1,2,3]triazol-2-yl-5-trifluoromethoxy-phenyl)-methanone. LC-MS: $t_R = 1.14$; $[M+H]^+ = 557.2$
6.186	[(S)-2-(5-Chloro-6-trifluoromethyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(4,5-dimethoxy-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.99$; $[M+H]^+ = 533.2$
6.187	[(S)-2-(5-Chloro-6-trifluoromethyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(5-methyl-2-pyrazol-1-yl-phenyl)-methanone. LC-MS: $t_R = 1.06$; $[M+H]^+ = 486.2$
6.188	[(S)-2-(5-Chloro-6-trifluoromethyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(2-pyrazol-1-yl-phenyl)-methanone. LC-MS: $t_R = 1.02$; $[M+H]^+ = 472.2$
6.189	[(S)-2-(5-Chloro-6-trifluoromethyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(6-methyl-3-pyrazol-1-yl-pyridin-2-yl)-methanone. LC-MS: $t_R = 0.98$; $[M+H]^+ = 487.3$
6.190	[(S)-2-(5-Chloro-6-trifluoromethyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(6-methyl-3-[1,2,3]triazol-2-yl-pyridin-2-yl)-methanone. LC-MS: $t_R = 0.98$; $[M+H]^+ = 488.2$
6.191	[(S)-2-(5,6-Dimethyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(2-fluoro-6-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.69$; $[M+H]^+ = 417.3$

6.192	[(S)-2-(5,6-Dimethyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(5-fluoro-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.70$; $[M+H]^+ = 417.3$
6.193	(5-Chloro-2-[1,2,3]triazol-2-yl-phenyl)-[(S)-2-(5,6-dimethyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.76$; $[M+H]^+ = 433.2$
6.194	[(S)-2-(5,6-Dimethyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.68$; $[M+H]^+ = 399.3$
6.195	[(S)-2-(5,6-Dimethyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(2-fluoro-3-methoxy-6-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.70$; $[M+H]^+ = 447.3$
6.196	[(S)-2-(5,6-Dimethyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(2-fluoro-3-methyl-6-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.74$; $[M+H]^+ = 431.3$
6.197	[(S)-2-(5,6-Dimethyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(4,5-dimethyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.78$; $[M+H]^+ = 427.3$
6.198	[(S)-2-(5,6-Dimethyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(4-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.8$; $[M+H]^+ = 413.4$
6.199	[(S)-2-(5,6-Dimethyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(2-methyl-6-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.79$; $[M+H]^+ = 413.4$
6.200	(4,5-Dimethoxy-2-[1,2,3]triazol-2-yl-phenyl)-[(S)-2-(5,6-dimethyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.74$; $[M+H]^+ = 459.4$
6.201	[(S)-2-(5,6-Dimethyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(5-methyl-2-pyrazol-1-yl-phenyl)-methanone. LC-MS: $t_R = 0.8$; $[M+H]^+ = 412.4$
6.202	[(S)-2-(5,6-Dimethyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(6-methyl-3-[1,2,3]triazol-2-yl-pyridin-2-yl)-methanone. LC-MS: $t_R = 0.75$; $[M+H]^+ = 414.4$
7.1	[(2S,4R)-2-(6-Chloro-1,7-dimethyl-1H-benzimidazol-2-yl)-4-methoxy-pyrrolidin-1-yl]-(5-methoxy-2-[1,2,3]triazol-2-yl-phenyl)-methanone and/or [(2S,4R)-2-(5-Chloro-1,4-dimethyl-1H-benzimidazol-2-yl)-4-methoxy-pyrrolidin-1-yl]-(5-methoxy-2-[1,2,3]triazol-2-yl-phenyl)-methanone (mixture of isomers). LC-MS: $t_R = 0.78$; $[M+H]^+ = 481.3$
7.2	[(S)-4-Methylene-2-(1,5,6-trimethyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.77$; $[M+H]^+ = 427.4$
7.3	[(S)-2-(5-Chloro-1,4-dimethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone and/or [(S)-2-(6-Chloro-1,7-dimethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone (mixture of isomers). LC-MS: $t_R = 0.84$; $[M+H]^+ = 449.3$

7.4	[(S)-2-(5-Bromo-1,7-dimethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone and/or [(S)-2-(6-Bromo-1,4-dimethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone (mixture of isomers). LC-MS: $t_R = 0.86$; $[M+H]^+ = 493.2$
7.5	(S)-(5-(3-chlorophenyl)-2-methylthiazol-4-yl)(2-(1,4-dimethyl-1H-benzo[d]imidazol-2-yl)-2-methylpyrrolidin-1-yl)methanone and/or (S)-(5-(3-chlorophenyl)-2-methylthiazol-4-yl)(2-(1,7-dimethyl-1H-benzo[d]imidazol-2-yl)-2-methylpyrrolidin-1-yl)methanone (mixture of isomers). LC-MS: $t_R = 0.88$; $[M+H]^+ = 465.1$
7.6	(S)-(5-(3-chlorophenyl)-2-methylthiazol-4-yl)(2-(1-ethyl-4-methyl-1H-benzo[d]imidazol-2-yl)-2-methylpyrrolidin-1-yl)methanone and/or (S)-(5-(3-chlorophenyl)-2-methylthiazol-4-yl)(2-(1-ethyl-7-methyl-1H-benzo[d]imidazol-2-yl)-2-methylpyrrolidin-1-yl)methanone (mixture of isomers). LC-MS: $t_R = 0.93$; $[M+H]^+ = 479.1$
7.7	(S)-(2-(5-chloro-1,4-dimethyl-1H-benzo[d]imidazol-2-yl)-2-methylpyrrolidin-1-yl)(2-methyl-5-phenylthiazol-4-yl)methanone and/or (S)-(2-(6-chloro-1,7-dimethyl-1H-benzo[d]imidazol-2-yl)-2-methylpyrrolidin-1-yl)(2-methyl-5-phenylthiazol-4-yl)methanone (mixture of isomers). LC-MS: $t_R = 0.93$; $[M+H]^+ = 465.1$
7.8	(S)-(2-(5-chloro-1-ethyl-4-methyl-1H-benzo[d]imidazol-2-yl)-2-methylpyrrolidin-1-yl)(2-methyl-5-phenylthiazol-4-yl)methanone and/or (S)-(2-(6-chloro-1-ethyl-7-methyl-1H-benzo[d]imidazol-2-yl)-2-methylpyrrolidin-1-yl)(2-methyl-5-phenylthiazol-4-yl)methanone (mixture of isomers). LC-MS: $t_R = 0.97$; $[M+H]^+ = 479.2$
7.9	[(S)-2-(5,6-Dimethoxy-1-methyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.74$; $[M+H]^+ = 461.4$
7.10	[(S)-2-(1-Ethyl-5,6-dimethoxy-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.78$; $[M+H]^+ = 475.4$
7.11	(4-Methyl-2-[1,2,3]triazol-2-yl-phenyl)-[(S)-2-methyl-2-(1,5,6-trimethyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.82$; $[M+H]^+ = 429.4$
7.12	[(S)-2-(1-Ethyl-5,6-dimethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(4-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.87$; $[M+H]^+ = 443.4$
7.13	[(S)-2-(5,6-Dimethoxy-1-methyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(4,5-dimethyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.80$; $[M+H]^+ = 475.4$
7.14	(4,5-Dimethyl-2-[1,2,3]triazol-2-yl-phenyl)-[(S)-2-(1-ethyl-5,6-dimethoxy-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.83$; $[M+H]^+ = 489.4$

7.15	(4'-Fluoro-4-methyl-biphenyl-2-yl)-[(S)-2-methyl-2-(1,5,6-trimethyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.97$; $[M+H]^+ = 456.4$
7.16	[(S)-2-(1-Ethyl-5,6-dimethyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(4'-fluoro-4-methyl-biphenyl-2-yl)-methanone. LC-MS: $t_R = 1.0$; $[M+H]^+ = 470.4$
7.17	(4,5-Dimethyl-2-[1,2,3]triazol-2-yl-phenyl)-[(S)-2-methyl-2-(1,5,6-trimethyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.87$; $[M+H]^+ = 443.4$
7.18	[(S)-2-(5-Chloro-1,4-dimethyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(6-methyl-3-[1,2,3]triazol-2-yl-pyridin-2-yl)-methanone and / or [(S)-2-(3-Chloro-1,2-dimethyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(6-methyl-3-[1,2,3]triazol-2-yl-pyridin-2-yl)-methanone (mixture of isomers). LC-MS: $t_R = 0.83$; $[M+H]^+ = 450.3$
7.19	[(S)-2-(5-Chloro-1,4-dimethyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(2-[1,2,3]triazol-2-yl-phenyl)-methanone and / or [(S)-2-(3-Chloro-1,2-dimethyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(2-[1,2,3]triazol-2-yl-phenyl)-methanone (mixtures of isomers). LC-MS: $t_R = 0.81$; $[M+H]^+ = 435.3$
7.20	[(S)-2-(6-Chloro-1-methyl-5-trifluoromethyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone and or [(S)-2-(5-Chloro-1-methyl-6-trifluoromethyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone (mixtures of isomers). LC-MS: $t_R = 1.09$; $[M+H]^+ = 531.3$
7.21	[(S)-2-(5-Chloro-1-ethyl-4-methyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone and / or [(S)-2-(3-Chloro-1-ethyl-2-methyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone (mixtures of isomers). LC-MS: $t_R = 0.93$; $[M+H]^+ = 463.3$
7.22	[(S)-2-(5-Chloro-1,4-dimethyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-chloro-2-[1,2,3]triazol-2-yl-phenyl)-methanone and / or [(S)-2-(3-Chloro-1,2-dimethyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-chloro-2-[1,2,3]triazol-2-yl-phenyl)-methanone (mixtures of isomers). LC-MS: $t_R = 0.93$; $[M+H]^+ = 469.3$
7.23	[(S)-2-(5-Chloro-1,4-dimethyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methoxy-2-[1,2,3]triazol-2-yl-phenyl)-methanone and / or [(S)-2-(3-Chloro-1,2-dimethyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methoxy-2-[1,2,3]triazol-2-yl-phenyl)-methanone (mixtures of isomers). LC-MS: $t_R = 0.86$; $[M+H]^+ = 465.4$
7.24	(5-Chloro-2-[1,2,3]triazol-2-yl-phenyl)-[(S)-2-(5,6-dimethoxy-1-methyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.77$; $[M+H]^+ = 481.3$

7.25	(4,3'-Dimethyl-biphenyl-2-yl)-[(S)-2-methyl-2-(1,5,6-trimethyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 1.01$; $[M+H]^+ = 452.4$
7.26	[(S)-2-(1-Cyclopropylmethyl-5,6-dimethyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(4,3'-dimethyl-biphenyl-2-yl)-methanone. LC-MS: $t_R = 1.1$; $[M+H]^+ = 492.4$
7.27	(4,3'-Dimethyl-biphenyl-2-yl)-[(S)-2-(1-ethyl-5,6-dimethyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 1.04$; $[M+H]^+ = 466.4$
7.28	(4,3'-Dimethyl-biphenyl-2-yl)-{(S)-2-[1-(2-fluoro-ethyl)-5,6-dimethyl-1H-benzoimidazol-2-yl]-2-methyl-pyrrolidin-1-yl}-methanone. LC-MS: $t_R = 1.03$; $[M+H]^+ = 484.4$
7.29	(4,3'-Dimethyl-biphenyl-2-yl)-{(S)-2-[1-(2-methoxy-ethyl)-5,6-dimethyl-1H-benzoimidazol-2-yl]-2-methyl-pyrrolidin-1-yl}-methanone. LC-MS: $t_R = 1.06$; $[M+H]^+ = 496.4$
7.30	(3'-Fluoro-4-methyl-biphenyl-2-yl)-[(S)-2-methyl-2-(1,5,6-trimethyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.97$; $[M+H]^+ = 456.4$
7.31	[(S)-2-(1-Ethyl-5,6-dimethyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(3'-fluoro-4-methyl-biphenyl-2-yl)-methanone. LC-MS: $t_R = 1.0$; $[M+H]^+ = 470.4$
7.32	[(S)-2-(6-Bromo-5-fluoro-1-methyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone and / or [(S)-2-(4-Bromo-3-fluoro-1-methyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone (mixtures of isomers). LC-MS: $t_R = 0.89$; $[M+H]^+ = 497.3$
7.33	[(S)-2-(5-Chloro-1,4-dimethyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-[5-(3-chloro-phenyl)-2-methyl-thiazol-4-yl]-methanone and / or [(S)-2-(3-Chloro-1,2-dimethyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-[5-(3-chloro-phenyl)-2-methyl-thiazol-4-yl]-methanone (mixtures of isomers). LC-MS: $t_R = 1.05$; $[M+H]^+ = 499.2$
7.34	[(S)-2-(1-Cyclopropylmethyl-5,6-dimethyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(3'-fluoro-4-methyl-biphenyl-2-yl)-methanone. LC-MS: $t_R = 1.06$; $[M+H]^+ = 496.4$
7.35	{(S)-2-[1-(2-Fluoro-ethyl)-5,6-dimethyl-1H-benzoimidazol-2-yl]-2-methyl-pyrrolidin-1-yl}-(3'-fluoro-4-methyl-biphenyl-2-yl)-methanone. LC-MS: $t_R = 0.99$; $[M+H]^+ = 488.4$
7.36	(3'-Fluoro-4-methyl-biphenyl-2-yl)-{(S)-2-[1-(2-methoxy-ethyl)-5,6-dimethyl-1H-benzoimidazol-2-yl]-2-methyl-pyrrolidin-1-yl}-methanone. LC-MS: $t_R = 1.02$; $[M+H]^+ = 500.4$
7.37	(4,4'-Dimethyl-biphenyl-2-yl)-[(S)-2-methyl-2-(1,5,6-trimethyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 1.02$; $[M+H]^+ = 452.5$
7.38	(4,4'-Dimethyl-biphenyl-2-yl)-[(S)-2-(1-ethyl-5,6-dimethyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 1.04$; $[M+H]^+ = 466.4$

7.39	[(S)-2-(1-Cyclopropylmethyl-5,6-dimethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(4,4'-dimethyl-biphenyl-2-yl)-methanone. LC-MS: $t_R = 1.1$; $[M+H]^+ = 492.4$
7.40	(4,4'-Dimethyl-biphenyl-2-yl)-{(S)-2-[1-(2-fluoro-ethyl)-5,6-dimethyl-1H-benzimidazol-2-yl]-2-methyl-pyrrolidin-1-yl}-methanone. LC-MS: $t_R = 1.03$; $[M+H]^+ = 484.4$
7.41	(4,4'-Dimethyl-biphenyl-2-yl)-{(S)-2-[1-(2-methoxy-ethyl)-5,6-dimethyl-1H-benzimidazol-2-yl]-2-methyl-pyrrolidin-1-yl}-methanone. LC-MS: $t_R = 1.06$; $[M+H]^+ = 496.4$
7.42	(3'-Methoxy-4-methyl-biphenyl-2-yl)-[(S)-2-methyl-2-(1,5,6-trimethyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.98$; $[M+H]^+ = 468.4$
7.43	(2'-Fluoro-4-methyl-biphenyl-2-yl)-[(S)-2-methyl-2-(1,5,6-trimethyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone. LC-MS: $t_R = 0.97$; $[M+H]^+ = 456.4$
7.44	[(S)-2-(1-Ethyl-5,6-dimethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(3'-methoxy-4-methyl-biphenyl-2-yl)-methanone. LC-MS: $t_R = 1.01$; $[M+H]^+ = 482.4$
7.45	[(S)-2-(1-Ethyl-5,6-dimethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(2'-fluoro-4-methyl-biphenyl-2-yl)-methanone. LC-MS: $t_R = 1.00$; $[M+H]^+ = 470.4$
7.46	{(S)-2-[5,6-Dimethoxy-1-(2-methoxy-ethyl)-1H-benzimidazol-2-yl]-2-methyl-pyrrolidin-1-yl}-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.81$; $[M+H]^+ = 505.4$
7.47	[(S)-2-(1-Cyclopropylmethyl-5,6-dimethoxy-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.86$; $[M+H]^+ = 501.4$
7.48	{(S)-2-[1-(2-Fluoro-ethyl)-5,6-dimethoxy-1H-benzimidazol-2-yl]-2-methyl-pyrrolidin-1-yl}-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone. LC-MS: $t_R = 0.78$; $[M+H]^+ = 493.4$
7.49	[(S)-2-(1-Cyclopropylmethyl-5,6-dimethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(3'-methoxy-4-methyl-biphenyl-2-yl)-methanone. LC-MS: $t_R = 1.07$; $[M+H]^+ = 508.4$
7.50	[(S)-2-(1-Cyclopropylmethyl-5,6-dimethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(2'-fluoro-4-methyl-biphenyl-2-yl)-methanone. LC-MS: $t_R = 1.06$; $[M+H]^+ = 496.4$
7.51	{(S)-2-[1-(2-Fluoro-ethyl)-5,6-dimethyl-1H-benzimidazol-2-yl]-2-methyl-pyrrolidin-1-yl}-(3'-methoxy-4-methyl-biphenyl-2-yl)-methanone. LC-MS: $t_R = 0.99$; $[M+H]^+ = 500.4$
7.52	{(S)-2-[1-(2-Fluoro-ethyl)-5,6-dimethyl-1H-benzimidazol-2-yl]-2-methyl-pyrrolidin-1-yl}-(2'-fluoro-4-methyl-biphenyl-2-yl)-methanone. LC-MS: $t_R = 0.99$; $[M+H]^+ = 488.4$
7.53	{(S)-2-[1-(2-Methoxy-ethyl)-5,6-dimethyl-1H-benzimidazol-2-yl]-2-methyl-pyrrolidin-1-yl}-(3'-methoxy-4-methyl-biphenyl-2-yl)-methanone. LC-MS: $t_R = 1.02$; $[M+H]^+ = 512.4$
7.54	(2'-Fluoro-4-methyl-biphenyl-2-yl)-{(S)-2-[1-(2-methoxy-ethyl)-5,6-dimethyl-1H-benzimidazol-2-yl]-2-methyl-pyrrolidin-1-yl}-methanone. LC-MS: $t_R = 1.02$; $[M+H]^+ = 500.4$

II-Biological assays

Antagonistic activities on both orexin receptors have been measured for each example compound using the following procedure:

In vitro assay: Intracellular calcium measurements:

- 5 Chinese hamster ovary (CHO) cells expressing the human orexin-1 receptor and the human orexin-2 receptor, respectively, are grown in culture medium (Ham F-12 with L-Glutamine) containing 300 µg/ml G418, 100 U/ml penicillin, 100 µg/ml streptomycin and 10 % heat inactivated fetal calf serum (FCS). The cells are seeded at 20'000 cells / well into 384-well black clear bottom sterile plates (Greiner). The seeded plates are incubated overnight at
10 37°C in 5% CO₂.

Human orexin-A as an agonist is prepared as 1 mM stock solution in MeOH: water (1:1), diluted in HBSS containing 0.1 % bovine serum albumin (BSA), NaHCO₃: 0.375g/l and 20 mM HEPES for use in the assay at a final concentration of 3 nM.

- Antagonists are prepared as 10 mM stock solution in DMSO, then diluted in 384-well plates using DMSO followed by a transfer of the dilutions into in HBSS containing 0.1 % bovine serum albumin (BSA), NaHCO₃: 0.375g/l and 20 mM HEPES. On the day of the assay, 50 µl of staining buffer (HBSS containing 1% FCS, 20 mM HEPES, NaHCO₃: 0.375g/l, 5 mM probenecid (Sigma) and 3 µM of the fluorescent calcium indicator fluo-4 AM (1 mM stock solution in DMSO, containing 10% pluronic) is added to each well. The 384-well cell-plates
15 are incubated for 50 min at 37° C in 5% CO₂ followed by equilibration at RT for 30 min before measurement.

- Within the Fluorescent Imaging Plate Reader (FLIPR Tetra, Molecular Devices), antagonists are added to the plate in a volume of 10 µl/well, incubated for 120 min and finally 10 µl/well of agonist is added. Fluorescence is measured for each well at 1 second intervals, and the
25 height of each fluorescence peak is compared to the height of the fluorescence peak induced by an approximate EC₇₀ (for example 5 nM) of orexin-A with vehicle in place of antagonist. The IC₅₀ value (the concentration of compound needed to inhibit 50 % of the agonistic response) is determined and may be normalized using the obtained IC₅₀ value of a on-plate reference compound. Optimized conditions are achieved by adjustment of pipetting speed and cell splitting regime. The calculated IC₅₀ values may fluctuate depending on the daily cellular assay performance. Fluctuations of this kind are known to those skilled in the art.
30 Average IC₅₀ values from several measurements are given as geometric mean values.

Antagonistic activities of example compounds with respect to the Ox₁ and the Ox₂ receptor are displayed in *Table 1*.

Table 1

Example Number	IC ₅₀ Ox1 [nM]	IC ₅₀ Ox2 [nM]	Example Number	IC ₅₀ Ox1 [nM]	IC ₅₀ Ox2 [nM]	Example Number	IC ₅₀ Ox1 [nM]	IC ₅₀ Ox2 [nM]
Ex. 1.1	1500	108	Ex. 1.10	98	248	Ex. 1.100	27	104
Ex. 1.101	21	50	Ex. 1.102	17	17	Ex. 1.103	364	112
Ex. 1.104	657	769	Ex. 1.105	45	89	Ex. 1.106	17	31
Ex. 1.107	28	58	Ex. 1.108	171	835	Ex. 1.109	5	10
Ex. 1.11	110	54	Ex. 1.110	164	46	Ex. 1.111	32	243
Ex. 1.112	69	289	Ex. 1.113	5	58	Ex. 1.114	30	74
Ex. 1.115	37	19	Ex. 1.116	86	248	Ex. 1.117	136	56
Ex. 1.118	191	364	Ex. 1.119	27	77	Ex. 1.12	273	344
Ex. 1.120	147	557	Ex. 1.121	76	287	Ex. 1.122	55	45
Ex. 1.123	147	70	Ex. 1.124	119	42	Ex. 1.125	39	73
Ex. 1.126	18	24	Ex. 1.127	33	22	Ex. 1.128	1480	331
Ex. 1.129	93	124	Ex. 1.13	15	24	Ex. 1.130	113	75
Ex. 1.131	37	48	Ex. 1.132	90	614	Ex. 1.133	4	11
Ex. 1.134	116	26	Ex. 1.135	32	194	Ex. 1.136	81	278
Ex. 1.137	4	33	Ex. 1.138	35	109	Ex. 1.139	103	26
Ex. 1.14	34	32	Ex. 1.140	192	46	Ex. 1.141	507	13
Ex. 1.142	24	13	Ex. 1.143	14	16	Ex. 1.144	33	27
Ex. 1.145	3	7	Ex. 1.146	7	23	Ex. 1.147	685	24
Ex. 1.148	1170	44	Ex. 1.149	143	9	Ex. 1.15	67	45
Ex. 1.150	301	11	Ex. 1.151	763	110	Ex. 1.152	282	23
Ex. 1.153	40	3	Ex. 1.154	483	21	Ex. 1.155	36	16
Ex. 1.156	1810	29	Ex. 1.157	419	10	Ex. 1.158	538	20
Ex. 1.159	903	23	Ex. 1.16	245	5	Ex. 1.160	245	5
Ex. 1.161	1450	22	Ex. 1.162	458	27	Ex. 1.163	167	17
Ex. 1.164	217	24	Ex. 1.165	76	9	Ex. 1.166	45	4
Ex. 1.167	97	15	Ex. 1.168	20	57	Ex. 1.169	37	3
Ex. 1.17	97	8	Ex. 1.170	258	20	Ex. 1.171	2210	36
Ex. 1.172	1160	19	Ex. 1.173	3930	54	Ex. 1.174	68	87
Ex. 1.175	528	76	Ex. 1.176	2190	99	Ex. 1.177	842	30
Ex. 1.178	2680	96	Ex. 1.179	486	28	Ex. 1.18	935	153
Ex. 1.180	1490	50				Ex. 1.19	660	76
Ex. 1.2	349	7	Ex. 1.20	692	121	Ex. 1.21	1180	149
Ex. 1.22	415	52	Ex. 1.23	523	212	Ex. 1.24	850	81
Ex. 1.25	364	90	Ex. 1.26	125	22	Ex. 1.27	1040	35

Ex. 1.28	114	110	Ex. 1.29	369	182	Ex. 1.3	265	51
Ex. 1.30	624	147	Ex. 1.31	195	78	Ex. 1.32	347	139
Ex. 1.33	76	48	Ex. 1.34	124	56	Ex. 1.35	117	99
Ex. 1.36	109	27	Ex. 1.37	154	62	Ex. 1.38	360	110
Ex. 1.39	157	73	Ex. 1.4	241	30	Ex. 1.40	74	23
Ex. 1.41	115	64	Ex. 1.42	263	65	Ex. 1.43	97	58
Ex. 1.44	404	120	Ex. 1.45	501	109	Ex. 1.46	128	13
Ex. 1.47	446	13	Ex. 1.49	32	42	Ex. 1.5	56	23
Ex. 1.50	394	107	Ex. 1.51	66	21	Ex. 1.52	737	121
Ex. 1.53	616	90	Ex. 1.54	726	80	Ex. 1.55	35	10
Ex. 1.56	252	88	Ex. 1.57	76	22	Ex. 1.58	32	24
Ex. 1.59	77	252	Ex. 1.6	202	46	Ex. 1.60	61	48
Ex. 1.61	21	21	Ex. 1.62	128	197	Ex. 1.63	12	9
Ex. 1.64	12	18	Ex. 1.65	63	147	Ex. 1.66	20	47
Ex. 1.67	127	84	Ex. 1.68	7	8	Ex. 1.69	37	31
Ex. 1.7	93	5	Ex. 1.70	26	26	Ex. 1.71	95	162
Ex. 1.72	112	180	Ex. 1.73	136	282	Ex. 1.74	52	100
Ex. 1.75	5	9	Ex. 1.76	279	17	Ex. 1.78	102	408
Ex. 1.79	5	46				Ex. 1.80	41	564
Ex. 1.81	20	116	Ex. 1.82	13	17	Ex. 1.83	17	36
Ex. 1.84	14	20	Ex. 1.85	5	23	Ex. 1.86	56	118
Ex. 1.87	30	77	Ex. 1.88	102	109	Ex. 1.89	39	180
Ex. 1.9	337	69	Ex. 1.90	2	14	Ex. 1.91	13	25
Ex. 1.92	23	69	Ex. 1.93	102	458	Ex. 1.94	42	414
Ex. 1.95	18	15	Ex. 1.96	54	43	Ex. 1.97	95	314
Ex. 1.98	105	387	Ex. 1.99	58	50	Ex. 2.1	56	53
Ex. 2.10	5	8	Ex. 2.11	69	124	Ex. 2.2	1300	317
Ex. 2.3	67	61	Ex. 2.4	425	131	Ex. 2.5	6	12
Ex. 2.6	52	59	Ex. 2.7	66	42	Ex. 2.8	70	51
Ex. 2.9	17	18	Ex. 3.1	102	69	Ex. 3.10	45	5
Ex. 3.11	6	9	Ex. 3.12	13	1	Ex. 3.13	16	7
Ex. 3.14	674	23	Ex. 3.15	96	4	Ex. 3.16	24	2
Ex. 3.17	368	51	Ex. 3.18	97	4	Ex. 3.19	1930	21
Ex. 3.2	13	6	Ex. 3.20	1860	40	Ex. 3.21	2510	33
Ex. 3.22	576	9	Ex. 3.23	3890	67	Ex. 3.24	5	12
Ex. 3.25	5	7	Ex. 3.26	50	66	Ex. 3.27	5	16
Ex. 3.28	10	19	Ex. 3.29	28	20	Ex. 3.3	1290	255
Ex. 3.30	2	2	Ex. 3.31	5	4	Ex. 3.32	0.9	5
Ex. 3.33	0.8	10	Ex. 3.34	0.9	8	Ex. 3.35	3	4

Ex. 3.36	5	9	Ex. 3.37	8	8	Ex. 3.38	6	13
Ex. 3.39	74	57	Ex. 3.4	44	14	Ex. 3.40	8	7
Ex. 3.41	0.7	7	Ex. 3.42	193	395	Ex. 3.43	10	13
Ex. 3.44	3	6	Ex. 3.45	16	19	Ex. 3.46	223	288
Ex. 3.47	225	471	Ex. 3.48	3	4	Ex. 3.49	13	18
Ex. 3.5	109	74	Ex. 3.50	232	252	Ex. 3.51	94	44
Ex. 3.52	28	27	Ex. 3.53	10	20	Ex. 3.54	52	44
Ex. 3.55	18	45	Ex. 3.56	18	37	Ex. 3.6	5	4
Ex. 3.7	9	5	Ex. 3.8	22	12	Ex. 3.9	1	2
Ex. 4.1	29	778	Ex. 4.10	2	833	Ex. 4.100	6	1280
Ex. 4.101	18	183	Ex. 4.11	10	4060	Ex. 4.12	23	4870
Ex. 4.13	26	4910	Ex. 4.14	15	4030	Ex. 4.15	2	670
Ex. 4.16	1	250	Ex. 4.17	14	1670	Ex. 4.18	43	2930
Ex. 4.19	2	422	Ex. 4.2	132	669	Ex. 4.20	28	2450
Ex. 4.21	1	133	Ex. 4.22	6	1075	Ex. 4.23	9	1940
Ex. 4.24	2	367	Ex. 4.25	4	441	Ex. 4.26	17	2280
Ex. 4.27	59	5590	Ex. 4.28	2	486	Ex. 4.29	30	353
Ex. 4.3	8	1350	Ex. 4.30	12	155	Ex. 4.31	77	774
Ex. 4.32	7	156	Ex. 4.33	9	294	Ex. 4.34	37	1880
Ex. 4.35	3	193	Ex. 4.36	70	2740	Ex. 4.37	24	988
Ex. 4.38	55	756	Ex. 4.39	14	315	Ex. 4.4	11	1170
Ex. 4.40	21	28	Ex. 4.41	182	248	Ex. 4.42	365	1130
Ex. 4.43	39	97	Ex. 4.44	38	246	Ex. 4.45	124	5780
Ex. 4.46	14	51	Ex. 4.47	176	281	Ex. 4.48	391	1320
Ex. 4.49	221	533	Ex. 4.5	29	3190	Ex. 4.50	373	4220
Ex. 4.51	99	677	Ex. 4.52	19	1067	Ex. 4.53	111	4440
Ex. 4.54	21	2322	Ex. 4.55	167	7970	Ex. 4.56	283	2620
Ex. 4.57	59	250	Ex. 4.58	66	2170	Ex. 4.59	6	67
Ex. 4.6	29	3890	Ex. 4.60	2	65	Ex. 4.61	0.9	72
Ex. 4.62	3	19	Ex. 4.63	0.8	122	Ex. 4.64	5	800
Ex. 4.65	8	207	Ex. 4.66	1	23	Ex. 4.67	0.9	251
Ex. 4.68	2	400	Ex. 4.69	2	191	Ex. 4.7	2	603
Ex. 4.70	0.6	404	Ex. 4.71	7	133	Ex. 4.72	4	51
Ex. 4.73	17	1290	Ex. 4.74	4	78	Ex. 4.75	8	636
Ex. 4.76	8	113	Ex. 4.77	5	790	Ex. 4.78	3	1630
Ex. 4.79	18	541	Ex. 4.8	68	9930	Ex. 4.80	3	2020
Ex. 4.81	4	79	Ex. 4.82	4	32	Ex. 4.83	159	2322
Ex. 4.84	8	58	Ex. 4.85	7	43	Ex. 4.86	7	352
Ex. 4.87	4	2076	Ex. 4.88	11	182	Ex. 4.89	3	2170

Ex. 4.9	2	2780	Ex. 4.90	6	2350	Ex. 4.91	17	183
Ex. 4.92	5	212	Ex. 4.93	4	74	Ex. 4.94	3	543
Ex. 4.95	28	1170	Ex. 4.96	31	996	Ex. 4.97	4	129
Ex. 4.98	4	1300	Ex. 4.99	13	981	Ex. 4.102	10	103
Ex. 4.103	4	178	Ex. 4.104	14	1660	Ex. 4.105	5	177
Ex. 4.106	23	529	Ex. 4.107	47	1760	Ex. 4.108	7	419
Ex. 4.109	3	108	Ex. 4.110	13	1520	Ex. 4.111	2	110
Ex. 4.112	3	1060	Ex. 4.113	9	2040	Ex. 4.114	33	1360
Ex. 4.115	7	754	Ex. 4.116	6	198	Ex. 4.117	16	2570
Ex. 4.118	50	4350	Ex. 4.119	21	2050	Ex. 4.120	14	725
Ex. 4.121	6	475	Ex. 4.122	3	610	Ex. 4.123	21	871
Ex. 4.124	8	591	Ex. 4.125	1	81	Ex. 4.126	43	2380
Ex. 4.127	1	19	Ex. 4.128	1	387	Ex. 4.129	35	2750
Ex. 4.130	11	128	Ex. 4.131	23	501	Ex. 4.132	87	1320
Ex. 4.133	25	359	Ex. 4.134	1	86	Ex. 4.135	4	203
Ex. 4.136	14	4360	Ex. 4.137	5	3090	Ex. 4.138	2	1340
Ex. 4.139	5	2260	Ex. 4.140	7	2170	Ex. 4.141	5	2760
Ex. 4.142	6	2160	Ex. 4.143	24	3680	Ex. 4.144	9	463
Ex. 4.145	3	635	Ex. 4.146	6	2110	Ex. 4.147	13	9650
Ex. 4.148	7	3180	Ex. 4.149	14	752	Ex. 4.150	6	180
Ex. 4.151	5	185	Ex. 4.152	2	40	Ex. 4.153	3	80
Ex. 4.154	26	4800	Ex. 4.155	42	891	Ex. 4.156	14	489
Ex. 4.157	13	1710	Ex. 4.158	2	308	Ex. 4.159	4	215
Ex. 4.160	2	27	Ex. 4.161	2	191	Ex. 4.162	2	299
Ex. 4.163	20	4400	Ex. 4.164	7	725	Ex. 4.165	20	968
Ex. 4.166	2	568	Ex. 4.167	13	144	Ex. 4.168	10	486
Ex. 4.169	4	892	Ex. 4.170	11	1430	Ex. 4.171	1.5	571
Ex. 4.172	2	457	Ex. 4.173	13	6330	Ex. 4.174	12	132
Ex. 4.175	14	249	Ex. 4.176	15	1067	Ex. 4.177	9	219
Ex. 4.178	7	350	Ex. 4.179	27	69	Ex. 4.180	8	304
Ex. 4.181	3	230	Ex. 4.182	7	694	Ex. 4.183	93	2256
						Ex. 5.1	26	13
Ex. 5.10	6	16	Ex. 5.100	17	27	Ex. 5.101	59	54
Ex. 5.102	34	82	Ex. 5.103	46	70	Ex. 5.104	92	97
Ex. 5.105	24	28	Ex. 5.106	25	37	Ex. 5.107	58	53
Ex. 5.108	265	161	Ex. 5.109	3	4	Ex. 5.11	29	413
Ex. 5.110	13	14	Ex. 5.111	5	7	Ex. 5.112	12	16
Ex. 5.113	12	15	Ex. 5.114	10	21	Ex. 5.115	22	18
Ex. 5.116	39	19	Ex. 5.117	57	38	Ex. 5.118	25	35

Ex. 5.119	30	29	Ex. 5.12	15	134	Ex. 5.120	48	26
Ex. 5.121	56	43	Ex. 5.122	99	74	Ex. 5.123	146	304
Ex. 5.124	113	411	Ex. 5.125	84	126	Ex. 5.126	117	396
Ex. 5.127	1	6	Ex. 5.128	20	219	Ex. 5.129	20	62
Ex. 5.13	6	31	Ex. 5.130	24	58	Ex. 5.131	15	178
Ex. 5.132	54	615	Ex. 5.133	11	45	Ex. 5.134	8	20
Ex. 5.135	10	49	Ex. 5.136	6	35	Ex. 5.137	3	6
Ex. 5.138	2	10	Ex. 5.139	2	21	Ex. 5.14	6	31
Ex. 5.140	5	28	Ex. 5.141	27	244	Ex. 5.142	8	24
Ex. 5.143	7	82	Ex. 5.144	20	210	Ex. 5.145	6	30
Ex. 5.146	15	43	Ex. 5.147	27	112	Ex. 5.148	35	159
Ex. 5.149	21	26	Ex. 5.15	5	22	Ex. 5.150	37	49
Ex. 5.151	20	40	Ex. 5.152	24	51	Ex. 5.153	22	35
Ex. 5.154	59	61	Ex. 5.155	134	249	Ex. 5.156	38	54
Ex. 5.157	367	538	Ex. 5.158	525	879	Ex. 5.159	23	37
Ex. 5.16	16	18	Ex. 5.160	38	113	Ex. 5.161	405	388
Ex. 5.162	252	250	Ex. 5.163	142	401	Ex. 5.164	104	419
Ex. 5.165	65	355	Ex. 5.166	13	59	Ex. 5.167	91	351
Ex. 5.168	64	570	Ex. 5.169	63	197	Ex. 5.17	11	76
Ex. 5.170	2	13	Ex. 5.171	12	58	Ex. 5.172	92	704
Ex. 5.173	290	1138	Ex. 5.174	59	133	Ex. 5.175	24	355
Ex. 5.176	389	1067	Ex. 5.177	76	860	Ex. 5.178	20	280
Ex. 5.179	368	1125	Ex. 5.18	23	84	Ex. 5.180	42	261
Ex. 5.181	385	556	Ex. 5.182	10	55	Ex. 5.183	339	1165
Ex. 5.184	300	413	Ex. 5.185	91	283	Ex. 5.186	120	294
Ex. 5.187	310	>2589	Ex. 5.188	23	119	Ex. 5.189	16	62
Ex. 5.19	9	20	Ex. 5.190	58	119	Ex. 5.191	18	69
Ex. 5.192	23	88	Ex. 5.193	24	203	Ex. 5.194	13	98
Ex. 5.195	73	102	Ex. 5.196	91	199	Ex. 5.197	18	166
Ex. 5.198	18	34	Ex. 5.199	12	83	Ex. 5.2	2	5
Ex. 5.20	19	161	Ex. 5.200	115	517	Ex. 5.201	265	861
Ex. 5.202	227	924	Ex. 5.203	195	1380	Ex. 5.204	360	900
Ex. 5.205	149	852	Ex. 5.206	19	117	Ex. 5.207	15	78
Ex. 5.208	53	284	Ex. 5.209	17	101	Ex. 5.21	22	18
Ex. 5.210	3	27	Ex. 5.211	27	159	Ex. 5.212	26	239
Ex. 5.213	16	51	Ex. 5.214	2	4	Ex. 5.215	4	11
Ex. 5.216	15	45	Ex. 5.22	7	86	Ex. 5.23	15	80
Ex. 5.24	2	9	Ex. 5.25	27	81	Ex. 5.26	39	197
Ex. 5.27	46	507	Ex. 5.28	4	19	Ex. 5.29	13	59

Ex. 5.3	2	4	Ex. 5.30	57	164	Ex. 5.31	44	740
Ex. 5.32	90	46	Ex. 5.33	25	29	Ex. 5.34	2	9
Ex. 5.35	102	99	Ex. 5.36	2	3	Ex. 5.37	10	16
Ex. 5.38	4	12	Ex. 5.39	36	20	Ex. 5.4	10	6
Ex. 5.40	44	91	Ex. 5.41	2	2	Ex. 5.42	49	75
Ex. 5.43	3	14	Ex. 5.44	8	29	Ex. 5.45	5	12
Ex. 5.46	6	14	Ex. 5.47	5	15	Ex. 5.48	12	25
Ex. 5.49	85	114	Ex. 5.5	3	5	Ex. 5.50	49	33
Ex. 5.51	10	18	Ex. 5.52	107	43	Ex. 5.53	24	23
Ex. 5.54	33	20	Ex. 5.55	113	202	Ex. 5.56	12	15
Ex. 5.57	57	55	Ex. 5.58	3	6	Ex. 5.59	42	84
Ex. 5.6	32	59	Ex. 5.60	156	54	Ex. 5.61	112	26
Ex. 5.62	35	130				Ex. 5.64	17	19
Ex. 5.65	200	154	Ex. 5.66	872	458	Ex. 5.67	27	23
Ex. 5.68	141	150	Ex. 5.69	414	1200	Ex. 5.7	5	29
Ex. 5.70	31	83	Ex. 5.71	56	22	Ex. 5.72	19	16
Ex. 5.73	98	57	Ex. 5.74	592	414	Ex. 5.75	300	126
Ex. 5.76	33	47	Ex. 5.77	64	153	Ex. 5.78	31	72
Ex. 5.79	43	97	Ex. 5.8	13	129	Ex. 5.80	548	360
Ex. 5.81	258	130	Ex. 5.82	27	15	Ex. 5.83	426	104
Ex. 5.84	138	76	Ex. 5.85	316	114	Ex. 5.86	94	85
Ex. 5.87	203	157	Ex. 5.88	192	166	Ex. 5.89	148	119
Ex. 5.9	8	81	Ex. 5.90	232	473	Ex. 5.91	217	1290
Ex. 5.92	62	177	Ex. 5.93	15	19	Ex. 5.94	14	37
Ex. 5.95	11	29	Ex. 5.96	13	32	Ex. 5.97	6	11
Ex. 5.98	120	264	Ex. 5.99	6	11	Ex. 5.217	98	274
Ex. 5.218	13	36	Ex. 5.219	19	72	Ex. 5.220	98	249
Ex. 5.221	19	83	Ex. 5.222	26	68	Ex. 5.223	3	12
Ex. 5.224	7	21	Ex. 5.225	16	44	Ex. 5.226	7	155
Ex. 5.227	20	521	Ex. 5.228	56	2030	Ex. 5.229	54	858
Ex. 5.230	21	530	Ex. 5.231	15	135	Ex. 5.232	20	2055
Ex. 5.233	5	263	Ex. 5.234	8	131	Ex. 5.235	5	16.5
Ex. 5.236	26	173	Ex. 5.237	58	223	Ex. 5.238	23	91
Ex. 5.239	35	594	Ex. 5.240	38	232	Ex. 5.241	54	1000
Ex. 5.242	56	401	Ex. 5.243	88	1120	Ex. 5.244	62	641
Ex. 5.245	14	80	Ex. 5.246	28	533	Ex. 5.247	10	136
Ex. 5.248	22	494	Ex. 5.249	38	53	Ex. 5.250	184	992
Ex. 5.251	13	774	Ex. 5.252	20	1640	Ex. 5.253	12	1187
Ex. 5.254	39	9270	Ex. 5.255	1	145	Ex. 5.256	13	402

Ex. 5.257	13	329	Ex. 5.258	36	6520	Ex. 5.259	29	4660
Ex. 5.260	20	2870	Ex. 5.261	4	466	Ex. 5.262	113	1
Ex. 5.263	56	304	Ex. 5.264	25	65	Ex. 5.265	9	32
Ex. 5.266	1	1	Ex. 5.267	12	105	Ex. 5.268	6	9
Ex. 5.269	15	22	Ex. 5.270	7	9	Ex. 5.271	5	6
Ex. 5.272	8	12	Ex. 5.273	7	12	Ex. 5.274	3	8
Ex. 5.275	5	6	Ex. 5.276	3	8	Ex. 5.277	6	21
Ex. 5.278	22	31	Ex. 5.279	3	8	Ex. 5.280	36	16
Ex. 5.281	34	380	Ex. 5.282	47	97	Ex. 5.283	87	397
Ex. 5.284	10	18	Ex. 5.285	32	452	Ex. 5.286	40	4592
Ex. 5.287	53	1165	Ex. 5.288	38	1340	Ex. 5.289	38	532
Ex. 5.290	33	2140	Ex. 5.291	48	347	Ex. 5.292	34	671
Ex. 5.293	77	4040	Ex. 5.294	39	714	Ex. 5.295	8	254
Ex. 5.296	25	31	Ex. 5.297	41	57	Ex. 5.298	15	21
Ex. 5.299	17	21	Ex. 5.300	10	18	Ex. 5.301	24	16
Ex. 5.302	28	42	Ex. 5.303	29	21	Ex. 5.304	19	86
Ex. 5.305	35	41	Ex. 5.306	33	69	Ex. 5.307	20	49
Ex. 5.308	21	122	Ex. 5.309	23	21	Ex. 5.310	32	70
						Ex. 6.1	367	33
Ex. 6.10	206	81	Ex. 6.100	55	4	Ex. 6.101	10	0.9
Ex. 6.102	364	32	Ex. 6.103	184	6	Ex. 6.104	13	0.8
Ex. 6.105	1060	77	Ex. 6.106	99	5	Ex. 6.107	74	16
Ex. 6.108	414	7	Ex. 6.109	112	7	Ex. 6.11	135	56
Ex. 6.110	15	1	Ex. 6.111	272	14	Ex. 6.112	15	0.7
Ex. 6.113	5	1	Ex. 6.114	16	0.7	Ex. 6.115	333	9
Ex. 6.116	14	3	Ex. 6.117	19	3	Ex. 6.118	21	4
Ex. 6.119	14	3	Ex. 6.12	317	6	Ex. 6.120	8	0.6
Ex. 6.121	1	0.5	Ex. 6.122	0.9	0.6	Ex. 6.123	2	0.7
Ex. 6.124	15	4	Ex. 6.125	2	0.8	Ex. 6.126	3	0.8
Ex. 6.127	17	4	Ex. 6.128	20	1	Ex. 6.129	215	39
Ex. 6.13	155	17	Ex. 6.130	11	0.6	Ex. 6.131	9	0.5
Ex. 6.132	13	4	Ex. 6.133	3	2	Ex. 6.134	75	10
Ex. 6.135	83	5	Ex. 6.136	765	4	Ex. 6.137	4950	23
Ex. 6.138	>6970	34	Ex. 6.139	137	0.6	Ex. 6.14	87	12
Ex. 6.140	667	10	Ex. 6.141	883	9	Ex. 6.142	17	22
Ex. 6.143	13	32	Ex. 6.144	15	56	Ex. 6.145	21	75
Ex. 6.146	16	33	Ex. 6.147	13	34	Ex. 6.148	19	40
Ex. 6.149	59	90	Ex. 6.15	298	37	Ex. 6.150	218	91
Ex. 6.151	46	44	Ex. 6.152	5	20	Ex. 6.153	8	16

Ex. 6.154	32	38	Ex. 6.155	649	105	Ex. 6.156	34	12
Ex. 6.157	70	21	Ex. 6.158	57	35	Ex. 6.159	11	10
Ex. 6.16	307	151	Ex. 6.160	326	8	Ex. 6.161	51	3
Ex. 6.162	330	7	Ex. 6.163	259	72	Ex. 6.164	100	9
Ex. 6.165	9	3	Ex. 6.166	33	4	Ex. 6.167	573	25
Ex. 6.168	261	54	Ex. 6.169	344	479	Ex. 6.17	308	47
Ex. 6.170	18	15	Ex. 6.171	162	19	Ex. 6.172	1260	67
Ex. 6.173	842	64	Ex. 6.174	233	15	Ex. 6.175	77	19
Ex. 6.176	58	18	Ex. 6.177	13	6	Ex. 6.178	44	10
Ex. 6.179	6	7	Ex. 6.18	1530	29	Ex. 6.180	5	4
Ex. 6.181	2	2	Ex. 6.182	12	6	Ex. 6.183	127	21
Ex. 6.184	67	73	Ex. 6.185	239	103	Ex. 6.186	27	6
Ex. 6.187	24	15	Ex. 6.188	115	66	Ex. 6.189	47	25
Ex. 6.19	1530	61	Ex. 6.190	10	9	Ex. 6.191	134	7
Ex. 6.192	48	7	Ex. 6.193	41	4	Ex. 6.194	51	5
Ex. 6.195	43	11	Ex. 6.196	59	12	Ex. 6.197	6	2
Ex. 6.2	223	17	Ex. 6.20	1750	49	Ex. 6.21	1970	76
Ex. 6.22	410	19	Ex. 6.23	452	67	Ex. 6.24	53	17
Ex. 6.25	154	52	Ex. 6.26	26	24	Ex. 6.27	5	9
Ex. 6.28	20	15	Ex. 6.29	69	92	Ex. 6.3	243	18
Ex. 6.30	203	31	Ex. 6.31	41	22	Ex. 6.32	814	103
Ex. 6.33	213	23	Ex. 6.34	398	66	Ex. 6.35	146	80
Ex. 6.36	15	2	Ex. 6.37	422	45	Ex. 6.38	269	19
Ex. 6.39	106	9	Ex. 6.4	2510	118	Ex. 6.40	70	9
Ex. 6.41	4	2	Ex. 6.42	12	2	Ex. 6.43	368	62
Ex. 6.44	10	2	Ex. 6.45	1	1	Ex. 6.46	30	4
Ex. 6.47	6	1	Ex. 6.48	14	7	Ex. 6.49	3	1
Ex. 6.5	216	19	Ex. 6.50	5	1	Ex. 6.51	8	3
Ex. 6.52	8	2	Ex. 6.53	4	1	Ex. 6.54	3	1
Ex. 6.55	5	1	Ex. 6.56	6	1	Ex. 6.57	11	11
Ex. 6.58	4	1	Ex. 6.59	76	10	Ex. 6.6	477	52
Ex. 6.60	83	3	Ex. 6.61	41	4	Ex. 6.62	44	4
Ex. 6.63	88	3	Ex. 6.64	142	33	Ex. 6.65	34	3
Ex. 6.66	57	2	Ex. 6.67	9	5	Ex. 6.68	767	62
Ex. 6.69	389	51	Ex. 6.7	15	3			
Ex. 6.71	24	2	Ex. 6.72	4	6	Ex. 6.73	10	4
Ex. 6.74	4	11	Ex. 6.75	8	4	Ex. 6.76	4	4
Ex. 6.77	3	9	Ex. 6.78	5	12	Ex. 6.79	2	4
Ex. 6.8	21	7	Ex. 6.80	13	19	Ex. 6.81	19	23

Ex. 6.82	36	208	Ex. 6.83	8	23	Ex. 6.84	8	14
Ex. 6.85	108	42	Ex. 6.86	22	20	Ex. 6.87	1	2
Ex. 6.88	20	15	Ex. 6.89	2	2	Ex. 6.9	7	4
Ex. 6.90	6	1	Ex. 6.91	2	1	Ex. 6.92	4	2
Ex. 6.93	0.9	3	Ex. 6.94	2	1	Ex. 6.95	9	2
Ex. 6.96	234	68	Ex. 6.97	14	10	Ex. 6.98	4	3
Ex. 6.99	5	1	Ex. 6.198	12	3	Ex. 6.199	143	6
Ex. 6.200	58	10	Ex. 6.201	148	25	Ex. 6.202	89	13
			Ex. 7.1	16	162	Ex. 7.2	420	69
Ex. 7.3	1	3	Ex. 7.4	7	6	Ex. 7.5	9	2
Ex. 7.6	5	6	Ex. 7.7	2	3	Ex. 7.8	2	4
Ex. 7.9	28	28	Ex. 7.10	33	42	Ex. 7.11	17	720
Ex. 7.12	15	52	Ex. 7.13	21	49	Ex. 7.14	23	50
Ex. 7.15	11	65	Ex. 7.16	8	21	Ex. 7.17	4	19
Ex. 7.18	1	4	Ex. 7.19	5	8	Ex. 7.20	2	14
Ex. 7.21	1	2	Ex. 7.22	1	4	Ex. 7.23	1	3
Ex. 7.24	33	33	Ex. 7.25	15	18	Ex. 7.26	42	54
Ex. 7.27	23	17	Ex. 7.28	14	19	Ex. 7.29	23	70
Ex. 7.30	9	12	Ex. 7.31	8	19	Ex. 7.32	7	124
Ex. 7.33	1	3	Ex. 7.34	16	84	Ex. 7.35	6	20
Ex. 7.36	16	115	Ex. 7.37	11	20	Ex. 7.38	8	30
Ex. 7.39	43	61	Ex. 7.40	23	39	Ex. 7.41	44	106
Ex. 7.42	38	28	Ex. 7.43	24	42	Ex. 7.44	21	35
Ex. 7.45	31	60	Ex. 7.46	78	77	Ex. 7.47	39	60
Ex. 7.48	33	40	Ex. 7.49	96	173	Ex. 7.50	66	171
Ex. 7.51	19	27	Ex. 7.52	27	60	Ex. 7.53	46	190
Ex. 7.54	87	270						

Compounds of the present invention may be further characterized with regard to their general pharmacokinetic and pharmacological properties using conventional assays well known in the art; for example relating to their bioavailability in different species (such as rat or dog); or relating to their ability to cross the blood-brain barrier, using for example a human P-glycoprotein 1 (MDR 1) substrate assay, or an in vivo assay to determine drug concentrations in the brain, e.g. in rats after oral dosing; or relating to their functional behavior in different disease related animal models {for example: the sedative effect of the compound using Electroencephalography (EEG) and Electromyography (EMG) signal measurments [F. Jenck et al., Nature Medicine **2007**, 13, 150-155]; the effect of the compound in the fear-potentiated startle paradigm [Fendt M et al., Neuroscience Biobehav

Rev. **1999**, 23, 743-760; WO2009/0047723]; the effect of the compound on stress-induced hyperthermia [Vinkers CH et al., European J Pharmacol. **2008**, 585, 407-425]; the effect of the compound on morphine-induced locomotor sensitization [Vanderschuren LJMJ et al., in Self DW, Staley JK (eds.) "Behavioral Neuroscience of Drug Addiction", Current Topics in Behavioral Neurosciences 3 (2009), 179-195] }; or for their properties with regard to drug safety and/or toxicological properties using conventional assays well known in the art, for example relating to cytochrome P450 enzyme inhibition and time dependent inhibition, pregnane X receptor (PXR) activation, glutathione binding, or phototoxic behavior.

Measurement of brain and systemic concentration after oral administration:

In order to assess brain penetration, the concentration of the compound is measured in plasma ([P]), and brain ([B]), sampled 3 h (or at different time points) following oral administration (e.g. 100 mg/kg) to male wistar rats. The compounds are formulated e.g. in 100% PEG 400. Samples are collected in the same animal at the same time point (+/- 5 min). Blood is sampled from the vena cava caudalis into containers with EDTA as anticoagulant and centrifuged to yield plasma. Brain is sampled after cardiac perfusion of 10 mL NaCl 0.9% and homogenized into one volume of cold phosphate buffer (pH 7.4). All samples are extracted with MeOH and analyzed by LC-MS/MS. Concentrations are determined with the help of calibration curves.

Results obtained for the compound of Example 5.19:

(3 h after oral administration (100 mg/kg), n = 3): [P] = 2095 ng / ml; [B] = 3880 ng / g.

Results obtained for the compound of Example 5.36:

(3 h after oral administration (100 mg/kg), n = 3): [P] = 1280 ng / ml; [B] = 1808 ng / g.

Results obtained for the compound of Example 5.277:

(3 h after oral administration (100 mg/kg), n = 3): [P] = 3560 ng / ml; [B] = 5880 ng / g.

Sedative effects: EEG, EMG and behavioural indices of alertness recorded by radiotelemetry in vivo in Wistar rats.

Electroencephalography (EEG) and Electromyography (EMG) signals were measured by telemetry using TL11M2-F20-EET miniature radiotelemetric implants (Data Science Int.) with two pairs of differential leads.

Surgical implantation was performed under general anesthesia with Ketamin/Xylazin, for cranial placement of one differential pair of EEG electrodes and one pair of EMG leads inserted in either side of the muscles of the neck. After surgery, rats recovered in a thermoregulated chamber and received analgesic treatment with subcutaneous

buprenorphine twice a day for 2 d. They were then housed individually and allowed to recover for a minimum of 2 weeks. Thereafter, rats—in their home cage—were placed in a ventilated sound-attenuating box, on a 12-h light / 12-h dark cycle, for acclimatization before continuous EEG / EMG recordings started. The telemetric technology that we used in this study allows accurate and stress-free acquisition of biosignals in rats placed in their familiar home cage environment, with no recording leads restricting their movements. Variables analyzed included four different stages of vigilance and sleep, spontaneous activity in the home cage and body temperature. Sleep and wake stages were evaluated using a rodent scoring software (Somnologica Science) directly processing electrical biosignals on 10 s contiguous epochs. The scoring is based on frequency estimation for EEG and amplitude discrimination for EMG and locomotor activity. Using these measurements, the software determines the probability that all components within each epoch best represent active waking (AW), quiet waking (QW), non-REM-sleep (NREM) or REM-sleep (REM). The percentage of total time spent in AW, QW, NREM- and REM-sleep was calculated per 12 h light or dark period. The latency to the onset of the first significant NREM- and REM-sleep episodes and the frequency and duration of those episodes were also calculated. AW, QW, NREM- and REM-sleep, home cage activity and body temperature were measured at baseline for at least one total circadian cycle (12 h-night, 12 h-day) before a test compound was administered. If baseline measurements indicated that animals were stable, test compound or vehicle was given in the evening by oral gavage at the end of the baseline 12-h day period, immediately before the nocturnal rise in orexin and activity in rats. All variables were subsequently recorded for 12 h following administration of the orexin receptor antagonist.


The compound of Example 5.19 has been tested in this assay (oral dosage: 30 mg/kg po; effects analyzed over 6 hours): Results are: -28% on active wake, -46% on home cage activity, +31% on NREM sleep, +69% on REM sleep; when compared to vehicle controls.

The compound of Example 5.36 has been tested in this assay (oral dosage: 30 mg/kg po; effects analyzed over 6 hours): Results are: -24% on active wake, -31% on home cage activity, +27% on NREM sleep, +53% on REM sleep; when compared to vehicle controls.

The compound of Example 5.277 has been tested in this assay (oral dosage: 30 mg/kg po; effects analyzed over 6 hours): Results are: -22% on active wake, -42% on home cage activity, +21% on NREM sleep, +52% on REM sleep; when compared to vehicle controls.

Claims

1. A compound of formula (I)



Formula (I)

5 wherein

Ar¹ represents

- phenyl or 5- or 6-membered heteroaryl, wherein said phenyl or 5- or 6-membered heteroaryl independently is mono-, di-, or tri-substituted; wherein

➤ one of said substituents is attached in *ortho*-position to the point of attachment of Ar¹ to the rest of the molecule; wherein said substituent is phenyl or 5- or 6-membered heteroaryl; wherein said phenyl or 5- or 6-membered heteroaryl substituent is independently unsubstituted, mono-, di-, or tri-substituted, wherein the substituents are independently selected from (C₁₋₄)alkyl, (C₁₋₄)alkoxy, halogen, cyano, (C₁₋₃)fluoroalkyl, and (C₁₋₃)fluoroalkoxy; or said *ortho* substituent is benzo[1,3]dioxolyl, or 2-(3-methoxy-phenyl)-ethynyl;

➤ and the other of said substituents, if present, is/are independently selected from (C₁₋₄)alkyl; (C₁₋₄)alkoxy; (C₃₋₆)cycloalkyl; halogen; cyano; (C₁₋₃)fluoroalkyl; (C₁₋₃)fluoroalkoxy; -NR¹⁰R¹¹, wherein R¹⁰ and R¹¹ independently represent hydrogen or (C₁₋₄)alkyl, or R¹⁰ and R¹¹ together with the nitrogen to which they are attached to form a pyrrolidine ring; unsubstituted pyridinyl; and phenyl which is unsubstituted, or mono- or di-substituted, wherein the substituents are independently selected from (C₁₋₄)alkyl, (C₁₋₄)alkoxy, cyano, and halogen;

one of **R²** and **R³** represents hydrogen, and the other represents hydrogen or (C₁₋₄)alkyl; and

one of **R^{4a}** and **R^{4b}** represents hydrogen, and the other represents hydrogen, (C₁₋₄)alkoxy, or halogen; or **R^{4a}** and **R^{4b}** together represent a group H₂C=; or both **R^{4a}** and **R^{4b}** represent fluorine; wherein, in case **R³** is different from hydrogen, both **R^{4a}** and **R^{4b}** represent hydrogen;

R¹ represents hydrogen, (C₁₋₄)alkyl, (C₃₋₆)cycloalkyl-(CH₂)-, (C₂₋₃)fluoroalkyl, or (C₁₋₄)alkoxy-(C₂₋₄)alkyl; and

(R⁵)_n represents one to three optional substituents independently selected from (C₁₋₄)alkyl, (C₁₋₄)alkoxy, halogen, (C₁₋₄)alkyl-thio-, (C₁₋₃)fluoroalkyl, (C₁₋₃)fluoroalkoxy, (C₁₋₃)fluoroalkyl-thio-, hydroxy-(C₁₋₄)alkyl-, (C₁₋₄)alkoxy-carbonyl-, nitro, hydroxy, and cyano; or (R⁵)_n represents a group -O-CH₂-CH₂-O-; or (R⁵)_n represents a fused phenyl group which, together with the benzimidazole moiety to which it is fused to, forms a 1H-naphtho[2,3-d]imidazol-2-yl group;

with the exception of

[2-(1H-benzimidazol-2-yl)-1-pyrrolidinyl][2-(1H-pyrazol-1-yl)phenyl]-methanone;

[2-(1H-benzimidazol-2-yl)-1-pyrrolidinyl][5-(2,5-dimethyl-1H-pyrrol-1-yl)-1-methyl-1H-pyrazol-4-yl]-methanone;

[2-(1H-benzimidazol-2-yl)-1-pyrrolidinyl][1,1'-biphenyl]-2-yl-methanone;

[2-(6-methyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl][3-phenyl-isoxazol-4-yl]-methanone;

[2-(1H-imidazol-2-yl)phenyl][2-(6-methyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;

[2-(6-methyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl][3-(thien-2-yl)-1H-pyrazol-4-yl]-methanone;


[2-(6-methyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl][3-phenyl-1H-pyrazol-4-yl]-methanone;

[2-(1H-benzimidazol-2-yl)-pyrrolidin-1-yl][3-cyclopropyl-1-phenyl-1H-pyrazol-5-yl]-methanone; and

[2-(1H-benzimidazol-2-yl)-pyrrolidin-1-yl][1-(2-fluorophenyl)-5-(1H-pyrrol-1-yl)-1H-pyrazol-4-yl]-methanone;

or a pharmaceutically acceptable salt thereof.

2. A compound according to claim 1; which is also a compound of the formula (II); wherein the absolute configuration is as depicted in formula (II):



Formula (II)

wherein

Ar¹ represents

- phenyl or 5- or 6-membered heteroaryl, wherein said phenyl or 5- or 6-membered heteroaryl independently is mono-, di-, or tri-substituted; wherein
 - one of said substituents is attached in *ortho*-position to the point of attachment of Ar¹ to the rest of the molecule; wherein said substituent is phenyl or 5- or 6-membered heteroaryl; wherein said phenyl or 5- or 6-membered heteroaryl substituent is independently unsubstituted, mono-, di-, or tri-substituted,

wherein the substituents are independently selected from (C₁₋₄)alkyl, (C₁₋₄)alkoxy, halogen, cyano, (C₁₋₃)fluoroalkyl, and (C₁₋₃)fluoroalkoxy; or said ortho substituent is benzo[1,3]dioxolyl, or 2-(3-methoxy-phenyl)-ethynyl;

- 5 ➤ and the other of said substituents, if present, is/are independently selected from (C₁₋₄)alkyl; (C₁₋₄)alkoxy; (C₃₋₆)cycloalkyl; halogen; cyano; (C₁₋₃)fluoroalkyl; (C₁₋₃)fluoroalkoxy; -NR¹⁰R¹¹, wherein R¹⁰ and R¹¹ independently represent hydrogen or (C₁₋₄)alkyl, or R¹⁰ and R¹¹ together with the nitrogen to which they are attached to form a pyrrolidine ring; unsubstituted pyridinyl; and phenyl
10 which is unsubstituted, or mono- or di-substituted, wherein the substituents are independently selected from (C₁₋₄)alkyl, (C₁₋₄)alkoxy, cyano, and halogen;

one of R² and R³ represents hydrogen, and the other represents hydrogen or (C₁₋₄)alkyl; and

one of R^{4a} and R^{4b} represents hydrogen, and the other represents hydrogen, (C₁₋₄)alkoxy, or halogen; or R^{4a} and R^{4b} together represent a group H₂C=; or both R^{4a} and R^{4b} represent

- 15 fluorine; wherein, in case R³ is different from hydrogen, both R^{4a} and R^{4b} represent hydrogen;

R¹ represents hydrogen, (C₁₋₄)alkyl, (C₃₋₆)cycloalkyl-(CH₂)-, (C₂₋₃)fluoroalkyl, or (C₁₋₄)alkoxy-(C₂₋₄)alkyl; and

(R⁵)_n represents one to three optional substituents independently selected from (C₁₋₄)alkyl, (C₁₋₄)alkoxy, halogen, (C₁₋₄)alkyl-thio-, (C₁₋₃)fluoroalkyl, (C₁₋₃)fluoroalkoxy, (C₁₋₃)fluoroalkyl-thio-, hydroxy-(C₁₋₄)alkyl-, (C₁₋₄)alkoxy-carbonyl-, nitro, hydroxy, and cyano; or (R⁵)_n
20 represents a group -O-CH₂-CH₂-O-; or (R⁵)_n represents a fused phenyl group which, together with the benzimidazole moiety to which it is fused to, forms a 1H-naphtho[2,3-d]imidazol-2-yl group;

with the exception of

- 25 [(S)-2-(1H-benzimidazol-2-yl)-1-pyrrolidinyl][2-(1H-pyrazol-1-yl)phenyl]-methanone;
 [(S)-2-(1H-benzimidazol-2-yl)-1-pyrrolidinyl][5-(2,5-dimethyl-1H-pyrrol-1-yl)-1-methyl-1H-pyrazol-4-yl]-methanone;
 [(S)-2-(1H-benzimidazol-2-yl)-1-pyrrolidinyl][1,1'-biphenyl]-2-yl-methanone;
 [(S)-2-(1H-imidazol-2-yl)phenyl][2-(6-methyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
 [(S)-2-(6-methyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl][3-(thien-2-yl)-1H-pyrazol-4-yl]-methanone;
30 [(S)-2-(6-methyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl][3-phenyl-1H-pyrazol-4-yl]-methanone;
 [(S)-2-(1H-benzimidazol-2-yl)-pyrrolidin-1-yl][3-cyclopropyl-1-phenyl-1H-pyrazol-5-yl]-methanone; and
 [(S)-2-(1H-benzimidazol-2-yl)-pyrrolidin-1-yl][1-(2-fluorophenyl)-5-(1H-pyrrol-1-yl)-1H-pyrazol-4-yl]-methanone;


or a pharmaceutically acceptable salt thereof.

3. A compound according to claim 2; wherein

- one of R^2 and R^3 represents hydrogen, and the other represents hydrogen or methyl; and
- R^{4a} and R^{4b} represent hydrogen; or R^{4a} represents methoxy or fluorine, and R^{4b} represents hydrogen; or R^{4a} and R^{4b} together represent a group $H_2C=$; wherein, in case R^3 is methyl, both R^{4a} and R^{4b} represent hydrogen;

or a pharmaceutically acceptable salt thereof.

4. A compound according to claim 1; which is also a compound of the formula (V); wherein the absolute configuration is as depicted in formula (V):



Formula (V)

wherein

Ar^1 represents

- phenyl or 5- or 6-membered heteroaryl, wherein said phenyl or 5- or 6-membered heteroaryl independently is mono-, di-, or tri-substituted; wherein
 - one of said substituents is attached in *ortho*-position to the point of attachment of Ar^1 to the rest of the molecule; wherein said substituent is phenyl or 5- or 6-membered heteroaryl; wherein said phenyl or 5- or 6-membered heteroaryl substituent is independently unsubstituted, mono-, di-, or tri-substituted, wherein the substituents are independently selected from (C_{1-4}) alkyl, (C_{1-4}) alkoxy, halogen, cyano, (C_{1-3}) fluoroalkyl, and (C_{1-3}) fluoroalkoxy; or said *ortho* substituent is benzo[1,3]dioxolyl;
 - and the other of said substituents, if present, is/are independently selected from (C_{1-4}) alkyl; (C_{1-4}) alkoxy; (C_{3-6}) cycloalkyl; halogen; cyano; (C_{1-3}) fluoroalkyl; (C_{1-3}) fluoroalkoxy; $-NR^{10}R^{11}$, wherein R^{10} and R^{11} independently represent hydrogen or (C_{1-4}) alkyl, or R^{10} and R^{11} together with the nitrogen to which they are attached to form a pyrrolidine ring; unsubstituted pyridinyl; and phenyl which is unsubstituted, or mono- or di-substituted, wherein the substituents are independently selected from (C_{1-4}) alkyl, (C_{1-4}) alkoxy, cyano, and halogen;

R^1 represents hydrogen, (C_{1-4}) alkyl, (C_{3-6}) cycloalkyl- $(CH_2)-$, (C_{2-3}) fluoroalkyl, or (C_{1-4}) alkoxy- (C_{2-4}) alkyl; and

$(R^5)_n$ represents one to three optional substituents independently selected from (C_{1-4}) alkyl, (C_{1-4}) alkoxy, halogen, (C_{1-4}) alkyl-thio-, (C_{1-3}) fluoroalkyl, (C_{1-3}) fluoroalkoxy, (C_{1-3}) fluoroalkyl-thio-, hydroxy- (C_{1-4}) alkyl-, (C_{1-4}) alkoxy-carbonyl-, nitro, hydroxy, and cyano;

or $(R^5)_n$ represents a group $-O-CH_2-CH_2-O-$;

or a pharmaceutically acceptable salt thereof.

5. A compound according to any one of claims 1 to 4; wherein the fragment



wherein R^{14} , R^{15} , R^{16} and R^{17} together represent one or two substituents, wherein

- one of R^{15} and R^{16} represents (C_{1-4}) alkyl, (C_{1-4}) alkoxy, (C_{1-4}) alkyl-thio-, halogen, (C_{1-3}) fluoroalkyl, (C_{1-3}) fluoroalkoxy, or cyano;
- or one of R^{14} and R^{17} represents (C_{1-4}) alkyl, (C_{1-4}) alkoxy, (C_{1-4}) alkyl-thio-, halogen, or (C_{1-3}) fluoroalkyl;


and the remaining substituent, if present, is one of R^{15} and R^{16} representing (C_{1-4}) alkyl, (C_{1-4}) alkoxy, or halogen;

or a pharmaceutically acceptable salt thereof.

6. A compound according to claim 5; wherein R^1 represents hydrogen;


or a pharmaceutically acceptable salt thereof.

7. A compound according to any one of claims 1 to 4; wherein the fragment:




represents a group independently selected from the following groups A, B, or C:


A.



B.



C.



or a pharmaceutically acceptable salt thereof.

5 8. A compound according to any one of claims 1 to 7; wherein

- **Ar¹** represents 5-membered heteroaryl, wherein the 5-membered heteroaryl is mono- or di-substituted; wherein

➤ one of said substituents is attached in *ortho*-position to the point of attachment of **Ar¹** to the rest of the molecule; wherein said *ortho*-substituent is phenyl, or 6-membered heteroaryl, which phenyl or 6-membered heteroaryl is independently unsubstituted, mono-, di-, or tri-substituted, wherein the substituents are independently selected from (C₁₋₄)alkyl, (C₁₋₄)alkoxy, halogen, cyano, (C₁₋₃)fluoroalkyl, and (C₁₋₃)fluoroalkoxy;

➤ and the other of said substituents, if present, is/are independently selected from (C₁₋₄)alkyl; (C₃₋₆)cycloalkyl; -NR¹⁰R¹¹, wherein R¹⁰ and R¹¹ independently represent (C₁₋₄)alkyl, or R¹⁰ and R¹¹ together with the nitrogen to which they are attached to form a pyrrolidine ring; and phenyl which is unsubstituted, or mono- or di-substituted, wherein the substituents are independently selected from (C₁₋₄)alkyl, (C₁₋₄)alkoxy, cyano, and halogen;

- or **Ar¹** represents 6-membered heteroaryl, wherein the 6-membered heteroaryl is mono-, di-, or tri-substituted; wherein

➤ one of said substituents is attached in *ortho*-position to the point of attachment of **Ar¹** to the rest of the molecule; wherein


- said *ortho*-substituent is unsubstituted 5-membered heteroaryl;

- or said *ortho*-substituent is phenyl which is unsubstituted, mono-, di-, or tri-substituted, wherein the substituents are independently selected from (C₁₋₄)alkyl, (C₁₋₄)alkoxy, halogen, cyano, (C₁₋₃)fluoroalkyl, and (C₁₋₃)fluoroalkoxy;
- 5 ➤ and the other of said substituents, if present, is/are independently selected from (C₁₋₄)alkyl, (C₁₋₄)alkoxy, halogen, and (C₁₋₃)fluoroalkyl;
- or Ar¹ represents phenyl which is mono-, di-, or tri-substituted; wherein
 - one of said substituents is attached in *ortho*-position to the point of attachment of Ar¹ to the rest of the molecule; wherein
 - 10 ○ said *ortho*-substituent is phenyl which is unsubstituted, mono-, di-, or tri-substituted, wherein the substituents are independently selected from (C₁₋₄)alkyl, (C₁₋₄)alkoxy, halogen, cyano, (C₁₋₃)fluoroalkyl, and (C₁₋₃)fluoroalkoxy;
 - or said *ortho* substituent is benzo[1,3]dioxol-5-yl;
 - 15 ○ or said *ortho*-substituent is 6-membered heteroaryl which is unsubstituted, mono-, or di-substituted, wherein the substituents are independently selected from (C₁₋₄)alkyl, (C₁₋₄)alkoxy, halogen, and (C₁₋₃)fluoroalkyl;
 - or said *ortho*-substituent is 5-membered heteroaryl which is unsubstituted, mono-, or di-substituted, wherein the substituents are independently
 - 20 selected from (C₁₋₄)alkyl, (C₁₋₄)alkoxy, halogen, and (C₁₋₃)fluoroalkyl;
 - or said *ortho*-substituent is 2-(3-methoxy-phenyl)-ethynyl;
 - and the other of said substituents, if present, is/are independently selected from (C₁₋₄)alkyl; (C₁₋₄)alkoxy; (C₃₋₆)cycloalkyl; halogen; cyano; (C₁₋₃)fluoroalkyl; (C₁₋₃)fluoroalkoxy; unsubstituted pyridinyl; and phenyl which is unsubstituted, or
 - 25 mono- or di-substituted, wherein the substituents are independently selected from (C₁₋₄)alkyl, (C₁₋₄)alkoxy, cyano, and halogen.


or a pharmaceutically acceptable salt thereof.

9. A compound according to any one of claims 1 to 7; wherein; **Ar¹** is a group independently selected from the following groups A, B, C, D, E, F, G, H, I, J, K, L, or M:


A.




5 B.



C.








D.




10

E.







J.




K.



5 L.



M.



10 or a pharmaceutically acceptable salt thereof.

10. A compound according to claim 1; selected from the group consisting of:

[2-(5-Chloro-6-methyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-[5-(3-chloro-phenyl)-2-methyl-thiazol-4-yl]-methanone;

[5-(3-Chloro-phenyl)-2-methyl-thiazol-4-yl]-[2-(4,6-dimethyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;

5 [2-(4-Bromo-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-[5-(3-chloro-phenyl)-2-methyl-thiazol-4-yl]-methanone;

[2-(5-Bromo-7-methyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-[5-(3-chloro-phenyl)-2-methyl-thiazol-4-yl]-methanone;

[2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-[5-(3-chloro-phenyl)-2-methyl-thiazol-4-yl]-methanone;

10 [5-(3-Chloro-phenyl)-2-methyl-thiazol-4-yl]-[2-(6-chloro-5-trifluoromethyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;

[2-(4-Bromo-6-fluoro-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-[5-(3-chloro-phenyl)-2-methyl-thiazol-4-yl]-methanone;

[5-(3-Chloro-phenyl)-2-methyl-thiazol-4-yl]-[2-(4,5-dimethyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;

15 [2-(5,6-Dichloro-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-(2-methyl-5-phenyl-thiazol-4-yl)-methanone;

[2-(5-Chloro-6-methyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-(2-methyl-5-phenyl-thiazol-4-yl)-methanone;

[2-(4,6-Dimethyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-(2-methyl-5-phenyl-thiazol-4-yl)-methanone;

[5-(3-Chloro-phenyl)-2-methyl-thiazol-4-yl]-[(S)-2-(7-methylsulfanyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;

20 [2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-(2-methyl-5-phenyl-thiazol-4-yl)-methanone;

[2-(6-Chloro-5-trifluoromethyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-(2-methyl-5-phenyl-thiazol-4-yl)-methanone;

[(S)-2-(4,6-Dimethyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-(2-methyl-5-phenyl-thiazol-4-yl)-methanone;

[(S)-2-(5-Chloro-6-methyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-(2-methyl-5-phenyl-thiazol-4-yl)-methanone;

[(S)-2-(6-Bromo-4-methyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-(2-methyl-5-phenyl-thiazol-4-yl)-methanone;

25 [(S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-(2-methyl-5-phenyl-thiazol-4-yl)-methanone;

[(S)-2-(5-Chloro-6-trifluoromethyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-(2-methyl-5-phenyl-thiazol-4-yl)-methanone;

[(S)-2-(6-Methylsulfanyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-(4-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;

(4,5-Dimethoxy-2-[1,2,3]triazol-2-yl-phenyl)-[(S)-2-(6-methylsulfanyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-

30 methanone;

(5-Methoxy-2-[1,2,3]triazol-2-yl-phenyl)-[(S)-2-(7-methylsulfanyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;

(2-Fluoro-3-methoxy-6-[1,2,3]triazol-2-yl-phenyl)-[(S)-2-(7-methylsulfanyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;

35 [(S)-2-(7-Methylsulfanyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-(4-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;

- [(S)-2-(6-Methylsulfanyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-(2-methyl-5-p-tolyl-thiazol-4-yl)-methanone;
[(S)-2-(7-Methylsulfanyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
[4-(3-Chloro-phenyl)-2-methyl-pyrimidin-5-yl]-[(S)-2-(6-methylsulfanyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- 5 [4-(3,4-Dimethyl-phenyl)-2-methyl-pyrimidin-5-yl]-[(S)-2-(6-methylsulfanyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
(2-Methyl-4-phenyl-pyrimidin-5-yl)-[(S)-2-(6-methylsulfanyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
[4-(4-Bromo-3-chloro-phenyl)-2-methyl-pyrimidin-5-yl]-[(S)-2-(6-methylsulfanyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- 10 (2-Cyclopropyl-5-phenyl-thiazol-4-yl)-[(S)-2-(7-methylsulfanyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
[5-(3-Methoxy-phenyl)-2-methyl-thiazol-4-yl]-[(S)-2-(7-methylsulfanyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
[(S)-2-(7-Methylsulfanyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-[2-methyl-5-(3-trifluoromethyl-phenyl)-thiazol-4-yl]-methanone;
- 15 [(S)-2-(7-Methylsulfanyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-(2-methyl-5-p-tolyl-thiazol-4-yl)-methanone;
(2-Cyclopropyl-5-p-tolyl-thiazol-4-yl)-[(S)-2-(7-methylsulfanyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
[2-Cyclopropyl-5-(4-fluoro-phenyl)-thiazol-4-yl]-[(S)-2-(7-methylsulfanyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
(3',4'-Dimethyl-biphenyl-2-yl)-[(S)-2-(7-methylsulfanyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- 20 [4-(3,4-Dichloro-phenyl)-2-methyl-pyrimidin-5-yl]-[(S)-2-(7-methylsulfanyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
[4-(3,4-Dichloro-phenyl)-pyrimidin-5-yl]-[(S)-2-(7-methylsulfanyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
[4-(3,4-Dimethyl-phenyl)-pyrimidin-5-yl]-[(S)-2-(7-methylsulfanyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- 25 [4-(4-Bromo-3-chloro-phenyl)-pyrimidin-5-yl]-[(S)-2-(7-methylsulfanyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
[4-(3,4-Dimethyl-phenyl)-2-methyl-pyrimidin-5-yl]-[(S)-2-(7-methylsulfanyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- 30 (2-Methyl-4-phenyl-pyrimidin-5-yl)-[(S)-2-(7-methylsulfanyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
[4-(4-Bromo-3-chloro-phenyl)-2-methyl-pyrimidin-5-yl]-[(S)-2-(7-methylsulfanyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
[4-(3,5-Dichloro-phenyl)-2-methyl-pyrimidin-5-yl]-[(S)-2-(7-methylsulfanyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- 35 [5-(3-Chloro-phenyl)-2-methyl-thiazol-4-yl]-[2-(5,6-dichloro-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;

- [5-(3-Chloro-phenyl)-2-methyl-thiazol-4-yl]-[2-(5,6-dimethyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
[(2S,4R)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-4-fluoro-pyrrolidin-1-yl]-[5-(3-chloro-phenyl)-2-methyl-thiazol-4-yl]-methanone;
[(2S,4R)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-4-fluoro-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
5 [5-(3-Chloro-phenyl)-2-methyl-thiazol-4-yl]-[(2S,4R)-4-fluoro-2-(4-methyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
[(2S,4R)-2-(6-Chloro-7-methyl-1H-benzoimidazol-2-yl)-4-methoxy-pyrrolidin-1-yl]-(5-chloro-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
10 [(2S,4R)-2-(6-Chloro-7-methyl-1H-benzoimidazol-2-yl)-4-methoxy-pyrrolidin-1-yl]-(2-fluoro-3-methoxy-6-[1,2,3]triazol-2-yl-phenyl)-methanone;
[(2S,4R)-2-(6-Chloro-7-methyl-1H-benzoimidazol-2-yl)-4-methoxy-pyrrolidin-1-yl]-(5-methoxy-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
[(2S,4R)-2-(6-Chloro-7-methyl-1H-benzoimidazol-2-yl)-4-methoxy-pyrrolidin-1-yl]-(2-fluoro-3-methyl-6-[1,2,3]triazol-2-yl-phenyl)-methanone;
15 [(2S,4R)-2-(6-Chloro-7-methyl-1H-benzoimidazol-2-yl)-4-methoxy-pyrrolidin-1-yl]-(4,5-dimethyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
[(2S,4R)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-4-methoxy-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
20 [(4R)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-4-methoxy-2-methyl-pyrrolidin-1-yl]-(2-methyl-5-phenyl-thiazol-4-yl)-methanone;
[(4R)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-4-methoxy-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
[(4R)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-4-methoxy-2-methyl-pyrrolidin-1-yl]-[5-(3-chloro-phenyl)-2-methyl-thiazol-4-yl]-methanone;
25 [5-(3-Chloro-phenyl)-2-methyl-thiazol-4-yl]-[(4R)-4-methoxy-2-methyl-2-(4-methyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
[(4R)-4-Methoxy-2-methyl-2-(4-methyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-(2-methyl-5-phenyl-thiazol-4-yl)-methanone;
30 [(2S,4R)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-4-methoxy-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
[(2S,4R)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-4-methoxy-2-methyl-pyrrolidin-1-yl]-[5-(3-chloro-phenyl)-2-methyl-thiazol-4-yl]-methanone;
[(2S,4R)-2-(6-Chloro-7-methyl-1H-benzoimidazol-2-yl)-4-methoxy-pyrrolidin-1-yl]-(2-methyl-5-m-tolyl-thiazol-4-yl)-methanone;
35

- [(2S,4R)-2-(6-Chloro-7-methyl-1H-benzimidazol-2-yl)-4-methoxy-pyrrolidin-1-yl]-[2-dimethylamino-5-(3,4-dimethyl-phenyl)-thiazol-4-yl]-methanone;
- [(2S,4R)-2-(6-Chloro-7-methyl-1H-benzimidazol-2-yl)-4-methoxy-pyrrolidin-1-yl]-[5-(3,4-dimethyl-phenyl)-2-methyl-thiazol-4-yl]-methanone;
- 5 [(2S,4R)-2-(6-Chloro-7-methyl-1H-benzimidazol-2-yl)-4-methoxy-pyrrolidin-1-yl]-[5-(4-chloro-phenyl)-2-methyl-thiazol-4-yl]-methanone;
- [(2S,4R)-2-(6-Chloro-7-methyl-1H-benzimidazol-2-yl)-4-methoxy-pyrrolidin-1-yl]-(2-methyl-5-p-tolyl-thiazol-4-yl)-methanone;
- [(2S,4R)-2-(6-Chloro-7-methyl-1H-benzimidazol-2-yl)-4-methoxy-pyrrolidin-1-yl]-[5-(4-methoxy-phenyl)-2-methyl-
- 10 thiazol-4-yl]-methanone;
- [(2S,4R)-2-(6-Chloro-7-methyl-1H-benzimidazol-2-yl)-4-methoxy-pyrrolidin-1-yl]-(3',4'-dimethyl-biphenyl-2-yl)-methanone;
- [(2S,4R)-4-Methoxy-2-(4-methyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-(2-methyl-5-phenyl-thiazol-4-yl)-methanone;
- 15 [(2S,4R)-2-(6-Chloro-7-methyl-1H-benzimidazol-2-yl)-4-methoxy-pyrrolidin-1-yl]-(4'-fluoro-3'-methyl-biphenyl-2-yl)-methanone;
- [(2S,4R)-2-(6-Chloro-7-methyl-1H-benzimidazol-2-yl)-4-methoxy-pyrrolidin-1-yl]-[5-(3-chloro-phenyl)-2-dimethylamino-thiazol-4-yl]-methanone;
- [(4R)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-4-methoxy-2-methyl-pyrrolidin-1-yl]-(5-fluoro-2-[1,2,3]triazol-2-
- 20 yl-phenyl)-methanone;
- [(4R)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-4-methoxy-2-methyl-pyrrolidin-1-yl]-(5-chloro-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- [(4R)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-4-methoxy-2-methyl-pyrrolidin-1-yl]-(2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- 25 [(4R)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-4-methoxy-2-methyl-pyrrolidin-1-yl]-(4,5-dimethyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- [(4R)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-4-methoxy-2-methyl-pyrrolidin-1-yl]-(4-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- [(4R)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-4-methoxy-2-methyl-pyrrolidin-1-yl]-(4,5-dimethoxy-2-
- 30 [1,2,3]triazol-2-yl-phenyl)-methanone;
- [(4R)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-4-methoxy-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-pyrazol-1-yl-phenyl)-methanone;
- [(4R)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-4-methoxy-2-methyl-pyrrolidin-1-yl]-(6-methyl-3-pyrazol-1-yl-pyridin-2-yl)-methanone;

- [(4R)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-4-methoxy-2-methyl-pyrrolidin-1-yl]-(6-methyl-3-[1,2,3]triazol-2-yl-pyridin-2-yl)-methanone;
- [(2S,4R)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-4-methoxy-pyrrolidin-1-yl]-(2-methyl-5-phenyl-thiazol-4-yl)-methanone;
- 5 [5-(3-Chloro-phenyl)-2-methyl-thiazol-4-yl]-[(2S,4R)-4-methoxy-2-(4-methyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- [(2S,4S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-4-methoxy-pyrrolidin-1-yl]-[5-(3-chloro-phenyl)-2-methyl-thiazol-4-yl]-methanone;
- [(2S,4R)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-4-methoxy-pyrrolidin-1-yl]-[5-(3-chloro-phenyl)-2-methyl-
- 10 thiazol-4-yl]-methanone;
- (5-Chloro-2-[1,2,3]triazol-2-yl-phenyl)-[(2S,3S)-2-(5,6-dimethoxy-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-methanone;
- [(2S,3S)-2-(5,6-Dimethoxy-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(4-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- 15 [(2S,3S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(4-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- [(2S,3S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(5-methoxy-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- [(2S,3S)-2-(5,6-Dimethyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(4-methyl-2-[1,2,3]triazol-2-yl-phenyl)-
- 20 methanone;
- [(2S,3S)-2-(5,6-Dimethyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(5-methoxy-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- [(2S,3S)-2-(5,6-Dimethyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- 25 [(S)-2-Methyl-2-(4-methyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-(2-methyl-5-phenyl-thiazol-4-yl)-methanone;
- [(S)-2-(5,6-Dimethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- [(S)-2-(5,6-Dimethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(4,4'-dimethyl-biphenyl-2-yl)-methanone;
- [(S)-2-(5,6-Dimethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(3'-fluoro-biphenyl-2-yl)-methanone;
- 30 [(S)-2-(5,6-Dimethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(3'-methyl-biphenyl-2-yl)-methanone;
- [(S)-2-(4,5-Dimethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(3'-fluoro-4-methyl-biphenyl-2-yl)-methanone;
- [(S)-2-(4,5-Dimethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(3'-methoxy-4-methyl-biphenyl-2-yl)-methanone;
- 35 [(S)-2-(4,5-Dimethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(4,3'-dimethyl-biphenyl-2-yl)-methanone;

- [(S)-2-(4,5-Dimethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(4'-methoxy-4-methyl-biphenyl-2-yl)-methanone;
- [(S)-2-(4,5-Dimethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(4,4'-dimethyl-biphenyl-2-yl)-methanone;
- [(S)-2-(4,5-Dimethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(2'-fluoro-4-methyl-biphenyl-2-yl)-methanone;
- 5 [(S)-2-(4,5-Dimethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(4'-fluoro-4-methyl-biphenyl-2-yl)-methanone;
- [(S)-2-(4,5-Dimethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(3'-fluoro-biphenyl-2-yl)-methanone;
- [(S)-2-(4,5-Dimethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(3'-methyl-biphenyl-2-yl)-methanone;
- 10 [(S)-2-(4,5-Dimethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(4'-methoxy-biphenyl-2-yl)-methanone;
- [(S)-2-(4,5-Dimethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(4'-methyl-biphenyl-2-yl)-methanone;
- [(S)-2-(6-Bromo-5-fluoro-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(4,5-dimethyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- [(S)-2-(6-Bromo-5-fluoro-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-chloro-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- 15 [(S)-2-(5-tert-Butyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- [(S)-2-(6-Bromo-5-fluoro-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(4-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- 20 [(S)-2-(6-Chloro-5-trifluoromethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(2-fluoro-6-[1,2,3]triazol-2-yl-phenyl)-methanone;
- [(S)-2-(6-Chloro-5-trifluoromethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(2-fluoro-3-methyl-6-[1,2,3]triazol-2-yl-phenyl)-methanone;
- [(S)-2-(6-Chloro-5-trifluoromethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(4,5-dimethoxy-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- 25 [(S)-2-(6-Chloro-5-trifluoromethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-fluoro-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- [(S)-2-(6-Chloro-5-trifluoromethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(4,5-dimethyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- 30 [(S)-2-(6-Chloro-5-trifluoromethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-pyrazol-1-yl-phenyl)-methanone;
- (5-Chloro-2-[1,2,3]triazol-2-yl-phenyl)-[(S)-2-(6-chloro-5-trifluoromethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-methanone;
- [(S)-2-(4-Chloro-6-trifluoromethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- 35

- [(S)-2-(6-Chloro-5-trifluoromethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(4-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- [(S)-2-(6-Chloro-5-trifluoromethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- 5 [(S)-2-(6-Chloro-5-trifluoromethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(6-methyl-3-[1,2,3]triazol-2-yl-pyridin-2-yl)-methanone;
- [(S)-2-(6-Chloro-5-trifluoromethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(2-fluoro-3-methoxy-6-[1,2,3]triazol-2-yl-phenyl)-methanone;
- [(S)-2-(6-Chloro-5-trifluoromethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(3'-methoxy-4-methyl-biphenyl-10 2-yl)-methanone;
- [(S)-2-(5-Chloro-6-methyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- [(S)-2-(6-Chloro-5-trifluoromethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(2'-fluoro-4-methyl-biphenyl-2-yl)-methanone;
- 15 [(S)-2-(6-Chloro-5-trifluoromethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(4'-fluoro-4-methyl-biphenyl-2-yl)-methanone;
- [(S)-2-(6-Chloro-5-trifluoromethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(3'-fluoro-4-methyl-biphenyl-2-yl)-methanone;
- 2'-[(S)-2-(6-Chloro-5-trifluoromethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidine-1-carbonyl]-4'-methyl-biphenyl-3-20 carbonitrile;
- [(S)-2-(4,6-Dimethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- (5-Chloro-2-[1,2,3]triazol-2-yl-phenyl)-[(S)-2-methyl-2-(5-trifluoromethyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- 25 (4,5-Dimethyl-2-[1,2,3]triazol-2-yl-phenyl)-[(S)-2-methyl-2-(5-trifluoromethyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- (4-Methyl-2-[1,2,3]triazol-2-yl-phenyl)-[(S)-2-methyl-2-(5-trifluoromethyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- [(S)-2-(4-Bromo-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-30 methanone;
- (3'-Fluoro-4-methyl-biphenyl-2-yl)-[(S)-2-methyl-2-(5-trifluoromethyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- (4,3'-Dimethyl-biphenyl-2-yl)-[(S)-2-methyl-2-(5-trifluoromethyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- [(S)-2-(5-Bromo-7-methyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-35 methanone;

- (4'-Methoxy-4-methyl-biphenyl-2-yl)-[(S)-2-methyl-2-(5-trifluoromethyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- (4,4'-Dimethyl-biphenyl-2-yl)-[(S)-2-methyl-2-(5-trifluoromethyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- (2-Benzo[1,3]dioxol-5-yl-5-methyl-phenyl)-[(S)-2-methyl-2-(5-trifluoromethyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- 5 [(S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- [5-(3-Chloro-phenyl)-2-methyl-thiazol-4-yl]-[(S)-2-(4,5-difluoro-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-methanone;
- 10 [(S)-2-(5,6-Dimethoxy-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- [(S)-2-(4,5-Difluoro-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(2-methyl-5-m-tolyl-thiazol-4-yl)-methanone;
- [(S)-2-(4,5-Difluoro-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-[5-(3-methoxy-phenyl)-2-methyl-thiazol-4-yl]-methanone;
- 15 [(S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-pyrimidin-2-yl-phenyl)-methanone;
- [(S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-pyridin-2-yl-phenyl)-methanone;
- [(S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-pyridin-3-yl-phenyl)-methanone;
- 20 [(S)-2-(6-Bromo-5-fluoro-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- [(S)-2-(6-Chloro-5-trifluoromethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- 25 [(S)-2-(4-Bromo-6-fluoro-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- [(S)-2-(4,5-Dimethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- [(S)-2-(4-Isopropyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- 30 [(S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(2-methyl-5-phenyl-thiazol-4-yl)-methanone;
- [(S)-2-(7,8-Dihydro-3H-6,9-dioxo-1,3-diaza-cyclopenta[a]naphthalen-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;

- [(S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-chloro-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- [(S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methoxy-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- 5 [(S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(2-fluoro-3-methoxy-6-[1,2,3]triazol-2-yl-phenyl)-methanone;
- [(S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(4-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- [(S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(4,5-dimethoxy-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- 10 [5-(3-Chloro-phenyl)-2-methyl-thiazol-4-yl]-[(S)-2-methyl-2-(4-methyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- [(S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(4,5-dimethyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- 15 [(S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-pyrazol-1-yl-phenyl)-methanone;
- [(S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(6-methyl-3-pyrazol-1-yl-pyridin-2-yl)-methanone;
- [(S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-fluoro-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- 20 [(S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- [(S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(6-methyl-3-[1,2,3]triazol-2-yl-pyridin-2-yl)-methanone;
- 25 [(S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(2-fluoro-6-[1,2,3]triazol-2-yl-phenyl)-methanone;
- [(S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-[5-(3-chloro-phenyl)-2-methyl-thiazol-4-yl]-methanone;
- (4,5-Dimethyl-2-[1,2,3]triazol-2-yl-phenyl)-[(S)-2-methyl-2-(4-methyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- 30 (5-Chloro-2-[1,2,3]triazol-2-yl-phenyl)-[(S)-2-(5,6-dimethyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-methanone;
- [(S)-2-(5,6-Dimethyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methoxy-2-[1,2,3]triazol-2-yl-phenyl)-methanone;

- [(S)-2-(5,6-Dimethyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(4-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- [(S)-2-(5,6-Dimethyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(4,5-dimethyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- 5 [(5-Chloro-2-[1,2,3]triazol-2-yl-phenyl)-[(S)-2-(5,6-dimethoxy-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-methanone;
- [(S)-2-(5,6-Dimethoxy-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methoxy-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- [(S)-2-(5,6-Dichloro-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- 10 [(S)-2-(5,6-Dimethoxy-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(2-[1,2,3]triazol-2-yl-5-trifluoromethyl-phenyl)-methanone;
- [(S)-2-(5,6-Dimethoxy-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(4,5-dimethyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- 15 [(S)-2-(5,6-Dimethoxy-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-pyrazol-1-yl-phenyl)-methanone;
- [(S)-2-(5,6-Dimethyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(3'-methoxy-4-methyl-biphenyl-2-yl)-methanone;
- [(S)-2-(5,6-Dimethyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(4'-methoxy-4-methyl-biphenyl-2-yl)-methanone;
- 20 [(S)-2-(5,6-Dimethyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(2'-fluoro-4-methyl-biphenyl-2-yl)-methanone;
- [(S)-2-(5,6-Dimethyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(4'-fluoro-4-methyl-biphenyl-2-yl)-methanone;
- 25 [(S)-2-(5,6-Dimethyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(3'-fluoro-4-methyl-biphenyl-2-yl)-methanone;
- [(S)-2-(5,6-Dimethyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(4,3'-dimethyl-biphenyl-2-yl)-methanone;
- [5-(4-Bromo-phenyl)-2-methyl-thiazol-4-yl]-[(S)-2-(7-methyl-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-methanone;
- 30 [(S)-2-(7-Methyl-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(2-methyl-5-m-tolyl-thiazol-4-yl)-methanone;
- [5-(3-Bromo-phenyl)-2-methyl-thiazol-4-yl]-[(S)-2-(7-methyl-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-methanone;
- [5-(3,5-Dimethyl-phenyl)-2-methyl-thiazol-4-yl]-[(S)-2-(7-methyl-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-methanone;

- [5-(3-Bromo-4-fluoro-phenyl)-2-methyl-thiazol-4-yl]-[(S)-2-(7-methyl-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-methanone;
- [(S)-2-(7-Methyl-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-[2-methyl-5-(3-trifluoromethyl-phenyl)-thiazol-4-yl]-methanone;
- 5 [5-(3,4-Dichloro-phenyl)-2-methyl-thiazol-4-yl]-[(S)-2-(7-methyl-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-methanone;
- [5-(3-Methoxy-phenyl)-2-methyl-thiazol-4-yl]-[(S)-2-(7-methyl-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-methanone;
- [5-(4-Bromo-phenyl)-2-methyl-thiazol-4-yl]-[(S)-2-(6-chloro-7-methyl-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-methanone;
- 10 [(S)-2-(6-Chloro-7-methyl-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-[5-(6-methoxy-pyridin-3-yl)-2-methyl-thiazol-4-yl]-methanone;
- [(S)-2-(6-Chloro-7-methyl-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-[5-(4-fluoro-phenyl)-2-methyl-thiazol-4-yl]-methanone;
- 15 [(S)-2-(6-Chloro-7-methyl-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-[5-(4-chloro-phenyl)-2-methyl-thiazol-4-yl]-methanone;
- [(S)-2-(6-Chloro-7-methyl-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-[2-methyl-5-(3-trifluoromethyl-phenyl)-thiazol-4-yl]-methanone;
- [(S)-2-(6-Chloro-7-methyl-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-[5-(3-methoxy-phenyl)-2-methyl-thiazol-4-yl]-methanone;
- 20 [(S)-2-(6-Chloro-7-methyl-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(2-methyl-5-m-tolyl-thiazol-4-yl)-methanone;
- [5-(3-Bromo-phenyl)-2-methyl-thiazol-4-yl]-[(S)-2-(6-chloro-7-methyl-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-methanone;
- 25 [(S)-2-(6-Chloro-7-methyl-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-[5-(3,5-dimethyl-phenyl)-2-methyl-thiazol-4-yl]-methanone;
- [5-(3-Bromo-4-fluoro-phenyl)-2-methyl-thiazol-4-yl]-[(S)-2-(6-chloro-7-methyl-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-methanone;
- [(S)-2-(6-Chloro-7-methyl-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-[5-(3,4-dichloro-phenyl)-2-methyl-thiazol-4-yl]-methanone;
- 30 [(S)-2-(6-Chloro-7-methyl-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(6-methyl-3-[1,2,3]triazol-2-yl-pyridin-2-yl)-methanone;
- [(S)-2-(6-Chloro-7-methyl-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(5-chloro-2-[1,2,3]triazol-2-yl-phenyl)-methanone;

- [(S)-2-(6-Chloro-7-methyl-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(4,5-dimethyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- [(S)-2-(6-Chloro-7-methyl-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(5-methoxy-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- 5 [(S)-2-(6-Chloro-7-methyl-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(2-fluoro-3-methyl-6-[1,2,3]triazol-2-yl-phenyl)-methanone;
- [(S)-2-(6-Chloro-7-methyl-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(2-fluoro-3-methoxy-6-[1,2,3]triazol-2-yl-phenyl)-methanone;
- [(S)-2-(6-Chloro-5-trifluoromethyl-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-[5-(3,4-dimethyl-phenyl)-2-
- 10 methyl-thiazol-4-yl]-methanone;
- [2-(5-Methoxy-1H-benzoimidazol-2-yl)-2-methyl-4-methylene-pyrrolidin-1-yl]-(2-methyl-5-m-tolyl-thiazol-4-yl)-methanone;
- [2-Dimethylamino-5-(3,4-dimethyl-phenyl)-thiazol-4-yl]-[2-(5-methoxy-1H-benzoimidazol-2-yl)-2-methyl-4-methylene-pyrrolidin-1-yl]-methanone;
- 15 [5-(3,4-Dimethyl-phenyl)-2-methyl-thiazol-4-yl]-[2-(5-methoxy-1H-benzoimidazol-2-yl)-2-methyl-4-methylene-pyrrolidin-1-yl]-methanone;
- [5-(4-Chloro-phenyl)-2-methyl-thiazol-4-yl]-[2-(5-methoxy-1H-benzoimidazol-2-yl)-2-methyl-4-methylene-pyrrolidin-1-yl]-methanone;
- [2-(5-Methoxy-1H-benzoimidazol-2-yl)-2-methyl-4-methylene-pyrrolidin-1-yl]-[5-(4-methoxy-phenyl)-2-methyl-
- 20 thiazol-4-yl]-methanone;
- [2-(5-Methoxy-1H-benzoimidazol-2-yl)-2-methyl-4-methylene-pyrrolidin-1-yl]-(2-methyl-5-p-tolyl-thiazol-4-yl)-methanone;
- [5-(3-Chloro-phenyl)-2-dimethylamino-thiazol-4-yl]-[2-(5-methoxy-1H-benzoimidazol-2-yl)-2-methyl-4-methylene-pyrrolidin-1-yl]-methanone;
- 25 [5-(3-Chloro-phenyl)-2-methyl-thiazol-4-yl]-[2-(5-methoxy-1H-benzoimidazol-2-yl)-2-methyl-4-methylene-pyrrolidin-1-yl]-methanone;
- [2-(5-Methoxy-1H-benzoimidazol-2-yl)-2-methyl-4-methylene-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- (5-Chloro-2-[1,2,3]triazol-2-yl-phenyl)-[2-(5-methoxy-1H-benzoimidazol-2-yl)-2-methyl-4-methylene-pyrrolidin-1-
- 30 yl]-methanone;
- (4,5-Dimethyl-2-[1,2,3]triazol-2-yl-phenyl)-[2-(5-methoxy-1H-benzoimidazol-2-yl)-2-methyl-4-methylene-pyrrolidin-1-yl]-methanone;
- [(S)-2-(5-tert-Butyl-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(4,5-dimethyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;

- [(S)-2-(5-tert-Butyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(4-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- [(S)-2-(5-tert-Butyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(4,5-dimethoxy-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- 5 [(S)-2-(5-Chloro-6-trifluoromethyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(2-fluoro-6-[1,2,3]triazol-2-yl-phenyl)-methanone;
- [(S)-2-(5-Chloro-6-trifluoromethyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(5-fluoro-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- (5-Chloro-2-[1,2,3]triazol-2-yl-phenyl)-[(S)-2-(5-chloro-6-trifluoromethyl-1H-benzimidazol-2-yl)-4-methylene-
- 10 pyrrolidin-1-yl]-methanone;
- [(S)-2-(5-Chloro-6-trifluoromethyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- [(S)-2-(5-Chloro-6-trifluoromethyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(2-fluoro-3-methoxy-6-[1,2,3]triazol-2-yl-phenyl)-methanone;
- 15 [(S)-2-(5-Chloro-6-trifluoromethyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(2-fluoro-3-methyl-6-[1,2,3]triazol-2-yl-phenyl)-methanone;
- [(S)-2-(5-Chloro-6-trifluoromethyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(4,5-dimethyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- [(S)-2-(5-Chloro-6-trifluoromethyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(4-methyl-2-[1,2,3]triazol-2-
- 20 yl-phenyl)-methanone;
- [(S)-2-(5-Chloro-6-trifluoromethyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(4,5-dimethoxy-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- [(S)-2-(5-Chloro-6-trifluoromethyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(5-methyl-2-pyrazol-1-yl-phenyl)-methanone;
- 25 [(S)-2-(5-Chloro-6-trifluoromethyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(6-methyl-3-pyrazol-1-yl-pyridin-2-yl)-methanone;
- [(S)-2-(5-Chloro-6-trifluoromethyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(6-methyl-3-[1,2,3]triazol-2-yl-pyridin-2-yl)-methanone;
- [(S)-2-(5,6-Dimethyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(5-fluoro-2-[1,2,3]triazol-2-yl-phenyl)-
- 30 methanone;
- (5-Chloro-2-[1,2,3]triazol-2-yl-phenyl)-[(S)-2-(5,6-dimethyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-methanone;
- [(S)-2-(5,6-Dimethyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- [(S)-2-(5,6-Dimethyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(2-fluoro-3-methoxy-6-[1,2,3]triazol-2-yl-
- 35 phenyl)-methanone;

- [(S)-2-(5,6-Dimethyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(2-fluoro-3-methyl-6-[1,2,3]triazol-2-yl-phenyl)-methanone;
- [(S)-2-(5,6-Dimethyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(4,5-dimethyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- 5 [5-(3-Bromo-4-fluoro-phenyl)-2-methyl-thiazol-4-yl]-[(S)-2-(5-methoxy-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-methanone;
- (2-Cyclopropyl-5-m-tolyl-thiazol-4-yl)-[(S)-2-(5-methoxy-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-methanone;
- [2-Dimethylamino-5-(3,4-dimethyl-phenyl)-thiazol-4-yl]-[(S)-2-(5-methoxy-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-methanone;
- 10 [5-(3-Bromo-phenyl)-2-cyclopropyl-thiazol-4-yl]-[(S)-2-(5-methoxy-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-methanone;
- [2-Cyclopropyl-5-(3-methoxy-phenyl)-thiazol-4-yl]-[(S)-2-(5-methoxy-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-methanone;
- 15 [2-(3,4-Dimethyl-phenyl)-5-methyl-thiophen-3-yl]-[(S)-2-(5-methoxy-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-methanone;
- [(S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(2-methyl-5-phenyl-thiazol-4-yl)-methanone;
- [(S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- 20 [5-(3-Chloro-phenyl)-2-methyl-thiazol-4-yl]-[(S)-2-(4-methyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-methanone;
- [(S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-[5-(3-chloro-phenyl)-2-methyl-thiazol-4-yl]-methanone;
- 25 [5-(3,4-Dimethyl-phenyl)-2-methyl-thiazol-4-yl]-[(S)-2-(6-methoxy-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-methanone;
- [(S)-2-(4,5-Dimethyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-[5-(3,4-dimethyl-phenyl)-2-methyl-thiazol-4-yl]-methanone;
- [(S)-2-(4,7-Dimethoxy-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-[5-(3,4-dimethyl-phenyl)-2-methyl-thiazol-4-yl]-methanone;
- 30 [5-(3,4-Dimethyl-phenyl)-2-methyl-thiazol-4-yl]-[(S)-4-methylene-2-(4-methylsulfanyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- [(S)-2-(4-Bromo-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-[5-(3,4-dimethyl-phenyl)-2-methyl-thiazol-4-yl]-methanone;

- [(S)-2-(4-Chloro-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-[5-(3,4-dimethyl-phenyl)-2-methyl-thiazol-4-yl]-methanone;
- [5-(3,4-Dimethyl-phenyl)-2-methyl-thiazol-4-yl]-[(S)-4-methylene-2-(5-methylsulfanyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- 5 [5-(3,4-Dimethyl-phenyl)-2-methyl-thiazol-4-yl]-[(S)-2-(4-methyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-methanone;
- [(S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-[5-(3,4-dimethyl-phenyl)-2-methyl-thiazol-4-yl]-methanone;
- [(S)-2-(5-Chloro-6-methyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-[5-(3,4-dimethyl-phenyl)-2-methyl-
- 10 thiazol-4-yl]-methanone;
- [(S)-2-(5,6-Dimethyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-[5-(3,4-dimethyl-phenyl)-2-methyl-thiazol-4-yl]-methanone;
- [(S)-2-(5-tert-Butyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-[5-(3,4-dimethyl-phenyl)-2-methyl-thiazol-4-yl]-methanone;
- 15 [(S)-2-(4,6-Dimethyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-[5-(3,4-dimethyl-phenyl)-2-methyl-thiazol-4-yl]-methanone;
- [(S)-2-(5,6-Dimethyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- [(S)-2-(4,6-Dimethyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-
- 20 methanone;
- [(S)-2-(5-Chloro-6-methyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- [(S)-2-(5-Chloro-6-trifluoromethyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- 25 [(S)-2-(5-Methoxy-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(2-methyl-5-m-tolyl-thiazol-4-yl)-methanone;
- [2-(3,4-Dimethyl-phenyl)-5-methyl-thiophen-3-yl]-[2-(5-methoxy-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-methanone;
- [2-(3,4-Dimethyl-phenyl)-5-methyl-thiophen-3-yl]-[4-methylene-2-(7-methylsulfanyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- 30 [2-(3,4-Dimethyl-phenyl)-5-methyl-thiophen-3-yl]-[4-methylene-2-(5-methylsulfanyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- [2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-[2-(3,4-dimethyl-phenyl)-5-methyl-thiophen-3-yl]-methanone;

- [2-(5,6-Dimethyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-[2-(3,4-dimethyl-phenyl)-5-methyl-thiophen-3-yl]-methanone;
- [2-(4,6-Dimethyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-[2-(3,4-dimethyl-phenyl)-5-methyl-thiophen-3-yl]-methanone;
- 5 [2-(4,5-Dimethyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-[2-(3,4-dimethyl-phenyl)-5-methyl-thiophen-3-yl]-methanone;
- [2-(4-Bromo-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-[2-(3,4-dimethyl-phenyl)-5-methyl-thiophen-3-yl]-methanone;
- [2-(3,4-Dimethyl-phenyl)-5-methyl-thiophen-3-yl]-[2-(4-methyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-methanone;
- 10 [5-(3-Chloro-phenyl)-2-methyl-thiazol-4-yl]-[(S)-2-(5-methoxy-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-methanone;
- [2-(5-Chloro-6-methyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-[2-(3,4-dimethyl-phenyl)-5-methyl-thiophen-3-yl]-methanone;
- 15 [2-(5-tert-Butyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-[2-(3,4-dimethyl-phenyl)-5-methyl-thiophen-3-yl]-methanone;
- [2-(4,7-Dimethoxy-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-[2-(3,4-dimethyl-phenyl)-5-methyl-thiophen-3-yl]-methanone;
- [2-(7-Chloro-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-[2-(3,4-dimethyl-phenyl)-5-methyl-thiophen-3-yl]-methanone;
- 20 (2-Cyclopropyl-5-m-tolyl-thiazol-4-yl)-[2-(5-methoxy-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-methanone;
- (2-Cyclopropyl-5-m-tolyl-thiazol-4-yl)-[4-methylene-2-(7-methylsulfanyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- 25 (2-Cyclopropyl-5-m-tolyl-thiazol-4-yl)-[4-methylene-2-(5-methylsulfanyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- [2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(2-cyclopropyl-5-m-tolyl-thiazol-4-yl)-methanone;
- [5-(3,4-Dimethyl-phenyl)-2-methyl-thiazol-4-yl]-[(S)-2-(5-methoxy-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-methanone;
- 30 (2-Cyclopropyl-5-m-tolyl-thiazol-4-yl)-[2-(5,6-dimethyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-methanone;
- (2-Cyclopropyl-5-m-tolyl-thiazol-4-yl)-[2-(4,6-dimethyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-methanone;

- (2-Cyclopropyl-5-m-tolyl-thiazol-4-yl)-[2-(4,5-dimethyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-methanone;
- [2-(4-Bromo-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(2-cyclopropyl-5-m-tolyl-thiazol-4-yl)-methanone;
- (2-Cyclopropyl-5-m-tolyl-thiazol-4-yl)-[2-(4-methyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-methanone;
- 5 [2-(5-Chloro-6-methyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(2-cyclopropyl-5-m-tolyl-thiazol-4-yl)-methanone;
- [2-(5-Chloro-6-trifluoromethyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(2-cyclopropyl-5-m-tolyl-thiazol-4-yl)-methanone;
- 10 (2-Cyclopropyl-5-m-tolyl-thiazol-4-yl)-[2-(4,7-dimethoxy-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-methanone;
- [2-(7-Chloro-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(2-cyclopropyl-5-m-tolyl-thiazol-4-yl)-methanone;
- [(S)-2-(5-Chloro-1,4-dimethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- 15 [(S)-2-(6-Chloro-1,7-dimethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- [(S)-2-(5-Bromo-1,7-dimethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- [(S)-2-(6-Bromo-1,4-dimethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- 20 (S)-(5-(3-chlorophenyl)-2-methylthiazol-4-yl)(2-(1,4-dimethyl-1H-benzo[d]imidazol-2-yl)-2-methylpyrrolidin-1-yl)methanone;
- (S)-(5-(3-chlorophenyl)-2-methylthiazol-4-yl)(2-(1,7-dimethyl-1H-benzo[d]imidazol-2-yl)-2-methylpyrrolidin-1-yl)methanone;
- 25 (S)-(5-(3-chlorophenyl)-2-methylthiazol-4-yl)(2-(1-ethyl-4-methyl-1H-benzo[d]imidazol-2-yl)-2-methylpyrrolidin-1-yl)methanone;
- (S)-(5-(3-chlorophenyl)-2-methylthiazol-4-yl)(2-(1-ethyl-7-methyl-1H-benzo[d]imidazol-2-yl)-2-methylpyrrolidin-1-yl)methanone;
- (S)-(2-(5-chloro-1,4-dimethyl-1H-benzo[d]imidazol-2-yl)-2-methylpyrrolidin-1-yl)(2-methyl-5-phenylthiazol-4-yl)methanone;
- 30 (S)-(2-(6-chloro-1,7-dimethyl-1H-benzo[d]imidazol-2-yl)-2-methylpyrrolidin-1-yl)(2-methyl-5-phenylthiazol-4-yl)methanone;
- (S)-(2-(5-chloro-1-ethyl-4-methyl-1H-benzo[d]imidazol-2-yl)-2-methylpyrrolidin-1-yl)(2-methyl-5-phenylthiazol-4-yl)methanone;

(S)-(2-(6-chloro-1-ethyl-7-methyl-1H-benzo[d]imidazol-2-yl)-2-methylpyrrolidin-1-yl)(2-methyl-5-phenylthiazol-4-yl)methanone;

[5-(3-Chloro-phenyl)-2-methyl-thiazol-4-yl]-[2-(4-chloro-6-trifluoromethyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone;

5 [4-(3-Chloro-phenyl)-2-methyl-pyrimidin-5-yl]-[(S)-2-(7-methylsulfanyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone;

[(2S,3S)-3-Methyl-2-(4-methyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;

10 [5-(3-Chloro-phenyl)-2-methyl-thiazol-4-yl]-[(2S,3S)-2-(5,6-dimethyl-1H-benzoimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-methanone;

[(2S,3S)-2-(6-Chloro-4-methyl-1H-benzoimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-[5-(4-fluoro-phenyl)-2-methyl-thiazol-4-yl]-methanone;

[(2S,3S)-2-(6-Chloro-4-methyl-1H-benzoimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(2-[1,2,3]triazol-2-yl-phenyl)-methanone;

15 [(2S,3S)-2-(5-Chloro-1H-benzoimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-[5-(3-chloro-phenyl)-2-methyl-thiazol-4-yl]-methanone;

[5-(3-Chloro-phenyl)-2-methyl-thiazol-4-yl]-[(2S,3S)-2-(5,6-difluoro-1H-benzoimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-methanone;

20 2-[(2S,3S)-1-[5-(3-Chloro-phenyl)-2-methyl-thiazole-4-carbonyl]-3-methyl-pyrrolidin-2-yl]-1H-benzoimidazole-4-carboxylic acid methyl ester;

[5-(3-Chloro-phenyl)-2-methyl-thiazol-4-yl]-[(2S,3S)-2-(4-methoxy-1H-benzoimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-methanone;

[(2S,3S)-2-(5-Chloro-6-methyl-1H-benzoimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-[5-(3-chloro-phenyl)-2-methyl-thiazol-4-yl]-methanone;

25 [5-(3-Chloro-phenyl)-2-methyl-thiazol-4-yl]-[(2S,3S)-2-(4,6-dimethyl-1H-benzoimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-methanone;

[5-(3-Chloro-phenyl)-2-methyl-thiazol-4-yl]-[(2S,3S)-2-(4-hydroxymethyl-1H-benzoimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-methanone;

30 [5-(3-Chloro-phenyl)-2-methyl-thiazol-4-yl]-[(2S,3S)-3-methyl-2-(4-trifluoromethyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone;

[5-(3-Chloro-phenyl)-2-methyl-thiazol-4-yl]-[(2S,3S)-2-(5-fluoro-1H-benzoimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-methanone;

[(2S,3S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-[5-(3-chloro-phenyl)-2-methyl-thiazol-4-yl]-methanone;

- [(2S,3S)-2-(4-Chloro-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-[5-(3-chloro-phenyl)-2-methyl-thiazol-4-yl]-methanone;
- [5-(3-Chloro-phenyl)-2-methyl-thiazol-4-yl]-[(2S,3S)-2-(4,5-difluoro-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-methanone;
- 5 [5-(3-Chloro-phenyl)-2-methyl-thiazol-4-yl]-[(2S,3S)-2-(4,5-dimethyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-methanone;
- [5-(3-Chloro-phenyl)-2-methyl-thiazol-4-yl]-[(2S,3S)-2-(4-isopropyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-methanone;
- [5-(3-Chloro-phenyl)-2-methyl-thiazol-4-yl]-[(2S,3S)-2-(5-methoxy-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-methanone;
- 10 [(2S,3S)-2-(5-Chloro-7-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-[5-(3-chloro-phenyl)-2-methyl-thiazol-4-yl]-methanone;
- (5-Methyl-2-[1,2,3]triazol-2-yl-phenyl)-[(2S,3S)-3-methyl-2-(5-trifluoromethyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- 15 [5-(3-Chloro-phenyl)-2-methyl-thiazol-4-yl]-[(2S,3S)-3-methyl-2-(5-methyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- [(2S,3S)-2-(5,6-Dimethyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- [(2S,3S)-2-(5-Chloro-6-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- 20 [(2S,3S)-2-(4,6-Dimethyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- [(2S,3S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- 25 [(2S,3S)-2-(4,5-Dimethyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- [(2S,3S)-2-(5-Chloro-7-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- [(2S,3S)-3-Methyl-2-(4-methyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-(2-methyl-5-phenyl-thiazol-4-yl)-methanone;
- 30 [(2S,3S)-3-Methyl-2-(5-methyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-(2-methyl-5-phenyl-thiazol-4-yl)-methanone;
- [5-(3,5-Difluoro-phenyl)-2-methyl-thiazol-4-yl]-[(2S,3S)-2-(5,6-dimethoxy-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-methanone;
- [(2S,3S)-2-(5,6-Dimethoxy-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-[5-(4-fluoro-phenyl)-2-methyl-thiazol-4-yl]-methanone;

- [(2S,3S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- [(2S,3S)-2-(1H-Benzoimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-[5-(3-chloro-phenyl)-2-methyl-thiazol-4-yl]-methanone;
- 5 [(2S,3S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(5-chloro-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- [(2S,3S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(4'-methyl-biphenyl-2-yl)-methanone;
- [(2S,3S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-[5-(4-chloro-phenyl)-2-methyl-
- 10 thiazol-4-yl]-methanone;
- [(2S,3S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(5-methyl-2-pyrazol-1-yl-phenyl)-methanone;
- [(2S,3S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(2-methyl-4-p-tolyl-thiazol-5-yl)-methanone;
- 15 [(2S,3S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-[5-(3,5-difluoro-phenyl)-2-methyl-thiazol-4-yl]-methanone;
- [(2S,3S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(2-fluoro-3-methyl-6-[1,2,3]triazol-2-yl-phenyl)-methanone;
- [(2S,3S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-[5-(6-methoxy-pyridin-3-yl)-2-
- 20 methyl-thiazol-4-yl]-methanone;
- [5-(3-Chloro-phenyl)-2-methyl-thiazol-4-yl]-[(2S,3S)-3-methyl-2-(4-methyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- [(2S,3S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-[5-(4-fluoro-phenyl)-2-methyl-thiazol-4-yl]-methanone;
- 25 [(2S,3S)-2-(5-Chloro-6-methyl-1H-benzoimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(5-chloro-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- [(2S,3S)-2-(5-Chloro-6-methyl-1H-benzoimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(4-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- [(2S,3S)-2-(5-Chloro-6-methyl-1H-benzoimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(5-methyl-2-pyrazol-1-yl-phenyl)-
- 30 methanone;
- [(2S,3S)-2-(5-Chloro-6-methyl-1H-benzoimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(5-methoxy-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- [(2S,3S)-2-(5-Chloro-6-methyl-1H-benzoimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(2-fluoro-3-methyl-6-[1,2,3]triazol-2-yl-phenyl)-methanone;

- [(2S,3S)-2-(5-Chloro-6-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- [(2S,3S)-2-(5-Chloro-6-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(4'-methyl-biphenyl-2-yl)-methanone;
- 5 [(2S,3S)-2-(5-Chloro-6-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-[5-(4-chloro-phenyl)-2-methyl-thiazol-4-yl]-methanone;
- [(2S,3S)-2-(5-Chloro-6-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(2-methyl-4-p-tolyl-thiazol-5-yl)-methanone;
- [(2S,3S)-2-(5-Chloro-6-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-[5-(3,5-difluoro-phenyl)-2-methyl-
- 10 thiazol-4-yl]-methanone;
- (5-Chloro-2-[1,2,3]triazol-2-yl-phenyl)-[(2S,3S)-2-(5,6-dimethyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-methanone;
- [(2S,3S)-2-(5,6-Dimethyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(5-methyl-2-pyrazol-1-yl-phenyl)-methanone;
- 15 [(2S,3S)-2-(5,6-Dimethyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(4'-methyl-biphenyl-2-yl)-methanone;
- [5-(4-Chloro-phenyl)-2-methyl-thiazol-4-yl]-[(2S,3S)-2-(5,6-dimethyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-methanone;
- [(2S,3S)-2-(5,6-Dimethyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(2-methyl-4-p-tolyl-thiazol-5-yl)-methanone;
- 20 [5-(3,5-Difluoro-phenyl)-2-methyl-thiazol-4-yl]-[(2S,3S)-2-(5,6-dimethyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-methanone;
- [5-(3-Chloro-phenyl)-2-methyl-thiazol-4-yl]-[(2S,3S)-3-methyl-2-(5-trifluoromethyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- [(2S,3S)-2-(5,6-Dimethyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-[5-(4-fluoro-phenyl)-2-methyl-thiazol-4-
- 25 yl]-methanone;
- [(2S,3S)-2-(6-Chloro-4-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(5-chloro-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- [(2S,3S)-2-(6-Chloro-4-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(4'-methyl-biphenyl-2-yl)-methanone;
- 30 [(2S,3S)-2-(6-Chloro-4-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(4-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- [(2S,3S)-2-(6-Chloro-4-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-[5-(4-chloro-phenyl)-2-methyl-thiazol-4-yl]-methanone;
- [(2S,3S)-2-(6-Chloro-4-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(5-methyl-2-pyrazol-1-yl-phenyl)-
- 35 methanone;

- [(2S,3S)-2-(6-Chloro-4-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(2-methyl-4-p-tolyl-thiazol-5-yl)-methanone;
- [(2S,3S)-2-(6-Chloro-4-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(5-methoxy-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- 5 [(2S,3S)-2-(6-Chloro-4-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-[5-(3,5-difluoro-phenyl)-2-methyl-thiazol-4-yl]-methanone;
- [(2S,3S)-2-(6-Chloro-4-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(2-fluoro-3-methyl-6-[1,2,3]triazol-2-yl-phenyl)-methanone;
- [(S)-2-(5-Chloro-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- 10 [(S)-2-(6-Chloro-5-fluoro-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- [(S)-2-(6-Bromo-5-fluoro-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-pyrazol-1-yl-phenyl)-methanone;
- 15 [(S)-2-(6-Bromo-5-fluoro-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(6-methyl-3-[1,2,3]triazol-2-yl-pyridin-2-yl)-methanone;
- [(S)-2-(6-Chloro-5-trifluoromethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(2-pyrazol-1-yl-phenyl)-methanone;
- [(S)-2-(6-Chloro-5-trifluoromethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(6-methyl-3-pyrazol-1-yl-pyridin-2-yl)-methanone;
- 20 [(S)-2-(6-Chloro-5-trifluoromethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(2-[1,2,3]triazol-2-yl-5-trifluoromethyl-phenyl)-methanone;
- [(S)-2-(6-Chloro-5-trifluoromethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(2-[1,2,3]triazol-2-yl-5-trifluoromethoxy-phenyl)-methanone;
- 25 (5-Methyl-2-[1,2,3]triazol-2-yl-phenyl)-[(S)-2-methyl-2-(4-trifluoromethyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- (5-Methyl-2-pyrazol-1-yl-phenyl)-[(S)-2-methyl-2-(5-trifluoromethyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- (6-Methyl-3-[1,2,3]triazol-2-yl-pyridin-2-yl)-[(S)-2-methyl-2-(5-trifluoromethyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- 30 (3'-Methoxy-4-methyl-biphenyl-2-yl)-[(S)-2-methyl-2-(5-trifluoromethyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- 4'-Methyl-2'-[(S)-2-methyl-2-(5-trifluoromethyl-1H-benzimidazol-2-yl)-pyrrolidine-1-carbonyl]-biphenyl-4-carbonitrile;

- (2'-Fluoro-4-methyl-biphenyl-2-yl)-[(S)-2-methyl-2-(5-trifluoromethyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- (4'-Fluoro-4-methyl-biphenyl-2-yl)-[(S)-2-methyl-2-(5-trifluoromethyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- 5 [(S)-2-(4,5-Difluoro-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(4,5-dimethyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- (5-Methyl-2-[1,2,3]triazol-2-yl-phenyl)-[(S)-2-methyl-2-(5-trifluoromethoxy-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- [(S)-2-(4,5-Difluoro-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(2-methyl-5-phenyl-thiazol-4-yl)-methanone;
- 10 [(S)-2-(4,5-Difluoro-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(2-methyl-5-p-tolyl-thiazol-4-yl)-methanone;
- [5-(4-Chloro-phenyl)-2-methyl-thiazol-4-yl]-[(S)-2-(4,5-difluoro-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-methanone;
- [(S)-2-(4,5-Difluoro-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-[5-(3-fluoro-phenyl)-2-methyl-thiazol-4-yl]-methanone;
- 15 [(S)-2-(6-Fluoro-5-trifluoromethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- [(S)-2-(5-Bromo-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- (5-Methyl-2-[1,2,3]triazol-2-yl-phenyl)-[(S)-2-methyl-2-(5-trifluoromethyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- 20 [(2S,4R)-2-(6-Chloro-1,7-dimethyl-1H-benzimidazol-2-yl)-4-methoxy-pyrrolidin-1-yl]-(5-methoxy-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- [(2S,4R)-2-(5-Chloro-1,4-dimethyl-1H-benzimidazol-2-yl)-4-methoxy-pyrrolidin-1-yl]-(5-methoxy-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- 25 [2-(5,6-Dimethoxy-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-(2-methyl-5-phenyl-thiazol-4-yl)-methanone;
- [2-(4,5-Dimethyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-(2-methyl-5-phenyl-thiazol-4-yl)-methanone;
- [2-(5-Methoxy-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- (5-Fluoro-2-[1,2,3]triazol-2-yl-phenyl)-[(S)-2-(6-methylsulfanyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- (5-Chloro-2-[1,2,3]triazol-2-yl-phenyl)-[(S)-2-(6-methylsulfanyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- 30 (5-Methoxy-2-[1,2,3]triazol-2-yl-phenyl)-[(S)-2-(6-methylsulfanyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- 3-[(S)-2-(6-Methylsulfanyl-1H-benzimidazol-2-yl)-pyrrolidine-1-carbonyl]-4-[1,2,3]triazol-2-yl-benzonitrile;
- [(S)-2-(6-Methylsulfanyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-(2-methyl-6-[1,2,3]triazol-2-yl-phenyl)-methanone;
- (5-Fluoro-2-[1,2,3]triazol-2-yl-phenyl)-[(S)-2-(6-methoxy-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- 35 (5-Chloro-2-[1,2,3]triazol-2-yl-phenyl)-[(S)-2-(6-methoxy-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;

- [(S)-2-(6-Methoxy-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-(5-methoxy-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
3-[(S)-2-(6-Methoxy-1H-benzimidazol-2-yl)-pyrrolidine-1-carbonyl]-4-[1,2,3]triazol-2-yl-benzonitrile;
[(S)-2-(6-Methoxy-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
[(S)-2-(6-Methoxy-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-(4-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
5 [(S)-2-(6-Methoxy-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-(2-methyl-6-[1,2,3]triazol-2-yl-phenyl)-methanone;
(4,5-Dimethoxy-2-[1,2,3]triazol-2-yl-phenyl)-[(S)-2-(6-methoxy-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
(5-Fluoro-2-[1,2,3]triazol-2-yl-phenyl)-[(S)-2-(7-methylsulfanyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
(2-Fluoro-6-[1,2,3]triazol-2-yl-phenyl)-[(S)-2-(7-methylsulfanyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
(5-Chloro-2-[1,2,3]triazol-2-yl-phenyl)-[(S)-2-(7-methylsulfanyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
10 3-[(S)-2-(7-Methylsulfanyl-1H-benzimidazol-2-yl)-pyrrolidine-1-carbonyl]-4-[1,2,3]triazol-2-yl-benzonitrile;
[(S)-2-(6-Methylsulfanyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
[(S)-2-(7-Methylsulfanyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-(2-methyl-6-[1,2,3]triazol-2-yl-phenyl)-methanone;
(2-Fluoro-3-methyl-6-[1,2,3]triazol-2-yl-phenyl)-[(S)-2-(5-methoxy-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-
methanone;
15 (5-Methyl-2-pyrazol-1-yl-phenyl)-[(S)-2-(5-methylsulfanyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
(2-Fluoro-3-methyl-6-[1,2,3]triazol-2-yl-phenyl)-[(S)-2-(5-methylsulfanyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-
methanone;
[(S)-2-(5-Methoxy-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
(3',4'-Dimethyl-biphenyl-2-yl)-[(S)-2-(6-methoxy-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
20 [(S)-2-(5-methoxy-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
(3',4'-Dimethyl-biphenyl-2-yl)-[(S)-2-(6-methylsulfanyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
[(S)-2-(6-Methylsulfanyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-(2-[1,2,3]triazol-2-yl-phenyl)-methanone;
25 [(S)-2-(5-Methylsulfanyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
[(S)-2-(7-Methylsulfanyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-(2-[1,2,3]triazol-2-yl-phenyl)-methanone;
[(2S,4R)-2-(6-Chloro-7-methyl-1H-benzimidazol-2-yl)-4-methoxy-pyrrolidin-1-yl]-(2-methyl-6-[1,2,3]triazol-2-yl-
phenyl)-methanone;
[(2S,4R)-2-(6-Chloro-7-methyl-1H-benzimidazol-2-yl)-4-methoxy-pyrrolidin-1-yl]-(4-methyl-2-[1,2,3]triazol-2-yl-
30 phenyl)-methanone;
[(2S,4R)-2-(6-Chloro-7-methyl-1H-benzimidazol-2-yl)-4-methoxy-pyrrolidin-1-yl]-(5-fluoro-2-[1,2,3]triazol-2-yl-
phenyl)-methanone;
(5-Chloro-2-[1,2,3]triazol-2-yl-phenyl)-[(2S,4R)-4-methoxy-2-(6-methoxy-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-
methanone;

- (4,5-Dimethyl-2-[1,2,3]triazol-2-yl-phenyl)-[(2S,4R)-4-methoxy-2-(6-methoxy-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- [(S)-2-(5,6-Dimethyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- [5-(6-Methoxy-pyridin-3-yl)-2-methyl-thiazol-4-yl]-[(S)-2-(7-methyl-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-methanone;
- 5 [5-(4-Fluoro-phenyl)-2-methyl-thiazol-4-yl]-[(S)-2-(7-methyl-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-methanone;
- (4,5-Dimethyl-2-[1,2,3]triazol-2-yl-phenyl)-[(S)-2-(7-methyl-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-methanone;
- 10 [5-(4-Chloro-phenyl)-2-methyl-thiazol-4-yl]-[(S)-2-(7-methyl-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-methanone;
- (5-Methoxy-2-[1,2,3]triazol-2-yl-phenyl)-[(S)-2-(7-methyl-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-methanone;
- [(S)-2-(7-Methyl-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(6-methyl-3-[1,2,3]triazol-2-yl-pyridin-2-yl)-methanone;
- 15 [(S)-2-(5-Methoxy-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- [(S)-2-(5-Methoxy-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- [(S)-2-(6-Chloro-7-methyl-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(5-methyl-2-pyrazol-1-yl-phenyl)-methanone;
- 20 [(S)-2-(6-Chloro-7-methyl-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(6-methyl-3-pyrazol-1-yl-pyridin-2-yl)-methanone;
- (5-Chloro-2-[1,2,3]triazol-2-yl-phenyl)-[(S)-2-(6-methoxy-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-methanone;
- 25 [(S)-2-(6-Methoxy-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(5-methyl-2-pyrazol-1-yl-phenyl)-methanone;
- (4,5-Dimethyl-2-[1,2,3]triazol-2-yl-phenyl)-[(S)-2-(6-methoxy-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-methanone;
- [(S)-2-(5-Methoxy-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(3'-methyl-biphenyl-2-yl)-methanone;
- 30 [(S)-2-(6-Methoxy-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(5-methoxy-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- (2-Fluoro-3-methyl-6-[1,2,3]triazol-2-yl-phenyl)-[(S)-2-(6-methoxy-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-methanone;
- [(S)-2-(5-tert-Butyl-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(5-fluoro-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- 35

- [(S)-2-(5-tert-Butyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(5-chloro-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- [(S)-2-(5-tert-Butyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- [(S)-2-(5-tert-Butyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(2-fluoro-3-methyl-6-[1,2,3]triazol-2-yl-phenyl)-methanone;
- 5 [(S)-2-(5-tert-Butyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(2-methyl-6-[1,2,3]triazol-2-yl-phenyl)-methanone;
- [(S)-2-(5-tert-Butyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(5-methyl-2-pyrazol-1-yl-phenyl)-methanone;
- 10 [(S)-2-(5-tert-Butyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(6-methyl-3-[1,2,3]triazol-2-yl-pyridin-2-yl)-methanone;
- (5-Fluoro-2-[1,2,3]triazol-2-yl-phenyl)-[(S)-2-(5-methoxy-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-methanone;
- [(S)-2-(5-Chloro-6-trifluoromethyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(2-methyl-6-[1,2,3]triazol-2-yl-phenyl)-methanone;
- 15 [(S)-2-(5,6-Dimethyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(2-fluoro-6-[1,2,3]triazol-2-yl-phenyl)-methanone;
- [(S)-2-(5-Methoxy-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(5-(4-methoxy-phenyl)-2-methyl-thiazol-4-yl)-methanone;
- 20 [5-(4-Bromo-phenyl)-2-methyl-thiazol-4-yl]-[(S)-2-(5-methoxy-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-methanone;
- [(S)-2-(5-Methoxy-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(2-methyl-5-phenyl-thiazol-4-yl)-methanone;
- [5-(3-Fluoro-phenyl)-2-methyl-thiazol-4-yl]-[(S)-2-(5-methoxy-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-methanone;
- 25 [(S)-2-(5-Methoxy-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(2-m-tolyl-thiophen-3-yl)-methanone;
- [2-(3,4-Dimethyl-phenyl)-thiophen-3-yl]-[(S)-2-(5-methoxy-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-methanone;
- [(S)-2-(4-Methyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(2-methyl-5-phenyl-thiazol-4-yl)-methanone;
- 30 [5-(4-Chloro-phenyl)-2-methyl-thiazol-4-yl]-[(S)-2-(5-methoxy-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-methanone;
- [(S)-4-Methylene-2-(7-methylsulfanyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- [(S)-4-Methylene-2-(5-methylsulfanyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- 35

[(S)-2-(4,5-Dimethyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone; and

[(S)-2-(5-tert-Butyl-1H-benzimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;

5 or a pharmaceutically acceptable salt thereof.

11. A compound according to claim 1; selected from the group consisting of:

[(2S,3S)-2-(4-Chloro-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-[2-(3,4-dimethyl-phenyl)-thiophen-3-yl]-methanone;

10 [5-(3,4-Dimethyl-phenyl)-2-methyl-thiazol-4-yl]-[(2S,3S)-3-methyl-2-(4-trifluoromethyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;

[2-(3,4-Dimethyl-phenyl)-thiophen-3-yl]-[(2S,3S)-3-methyl-2-(4-trifluoromethyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;

15 [(2S,3S)-2-(5,6-Dimethoxy-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(5-methyl-2-pyrimidin-2-yl-phenyl)-methanone;

[(S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(4-fluoro-biphenyl-2-yl)-methanone;

[(S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-fluoro-2-pyridin-2-yl-phenyl)-methanone;

20 [(S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(2-fluoro-6-pyridin-2-yl-phenyl)-methanone;

[(S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(2-fluoro-6-pyrimidin-2-yl-phenyl)-methanone;

[(S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(4-methyl-biphenyl-2-yl)-methanone;

25 [(S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-fluoro-2-pyrimidin-2-yl-phenyl)-methanone;

[(S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(3-fluoro-biphenyl-2-yl)-methanone;

(4,5-Dimethyl-2-[1,2,3]triazol-2-yl-phenyl)-[(S)-2-methyl-2-(5-trifluoromethoxy-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;

30 (4-Methyl-2-[1,2,3]triazol-2-yl-phenyl)-[(S)-2-methyl-2-(5-trifluoromethoxy-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;

[5-(3,4-Dimethyl-phenyl)-2-methyl-thiazol-4-yl]-[(S)-2-methyl-2-(5-trifluoromethoxy-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;

[(S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(2-methyl-5-phenyl-pyridin-4-yl)-methanone;

35 [(S)-2-Methyl-2-(5-trifluoromethyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-(4-pyridin-3-yl-biphenyl-2-yl)-methanone;

- [(S)-2-(6-Bromo-4,5-dimethyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- [(S)-2-(6-Bromo-4,5-dimethyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(2-fluoro-6-[1,2,3]triazol-2-yl-phenyl)-methanone;
- 5 Biphenyl-2-yl-[(S)-2-(6-bromo-4,5-dimethyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-methanone;
- [(S)-2-(6-Bromo-4,5-dimethyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(4-methyl-biphenyl-2-yl)-methanone;
- [(S)-2-(6-Bromo-4,5-dimethyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(2-methyl-5-o-tolyl-thiazol-4-yl)-methanone;
- 10 [(S)-2-(6-Bromo-4,5-dimethyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-(2,3-dichloro-phenyl)-2-methyl-thiazol-4-yl)-methanone;
- [(S)-2-(6-Bromo-4,5-dimethyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-(3,4-dimethyl-phenyl)-2-methyl-thiazol-4-yl)-methanone;
- [(S)-2-(6-Bromo-4,5-dimethyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-(3,5-difluoro-phenyl)-2-methyl-
- 15 thiazol-4-yl)-methanone;
- [(S)-2-(6-Bromo-4,5-dimethyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-(3-methoxy-phenyl)-2-methyl-thiazol-4-yl)-methanone;
- [(S)-2-(6-Bromo-4,5-dimethyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-(4-fluoro-phenyl)-2-methyl-thiazol-4-yl)-methanone;
- 20 [(S)-2-(5-Bromo-7-methyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-pyrimidin-2-yl-phenyl)-methanone;
- [(S)-2-Methyl-2-(7-methyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-(5-methyl-2-pyrimidin-2-yl-phenyl)-methanone;
- [(S)-2-(6-Chloro-5-trifluoromethyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-pyrimidin-2-yl-phenyl)-methanone;
- 25 [(S)-2-(5,6-Dimethoxy-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-pyrimidin-2-yl-phenyl)-methanone;
- [(S)-2-(5,6-Dimethyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-pyrimidin-2-yl-phenyl)-methanone;
- [(S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(3-fluoro-4,3'-dimethoxy-biphenyl-2-yl)-
- 30 methanone;
- [(S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(3,2'-difluoro-4-methoxy-biphenyl-2-yl)-methanone;
- [(S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(3,4'-difluoro-4-methoxy-biphenyl-2-yl)-methanone;

- [(S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(3,3'-difluoro-4-methoxy-biphenyl-2-yl)-methanone;
- [(S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(3-fluoro-4-methoxy-3'-methyl-biphenyl-2-yl)-methanone;
- 5 [(S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(3-fluoro-4-methoxy-4'-methyl-biphenyl-2-yl)-methanone;
- [(S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(3-fluoro-4-methoxy-3'-trifluoromethyl-biphenyl-2-yl)-methanone;
- (6-Benzo[1,3]dioxol-5-yl-2-fluoro-3-methoxy-phenyl)-[(S)-2-(5-chloro-4-methyl-1H-benzoimidazol-2-yl)-2-methyl-
- 10 pyrrolidin-1-yl]-methanone;
- 2'-[(S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidine-1-carbonyl]-3'-fluoro-4'-methoxy-biphenyl-3-carbonitrile;
- [(S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(3-fluoro-4-methoxy-3'-trifluoromethoxy-biphenyl-2-yl)-methanone;
- 15 [(S)-2-(5,6-Dimethyl-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(4-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- (4,5-Dimethoxy-2-[1,2,3]triazol-2-yl-phenyl)-[(S)-2-(5,6-dimethyl-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-methanone;
- (4-Methyl-2-[1,2,3]triazol-2-yl-phenyl)-[(S)-2-methyl-2-(1,5,6-trimethyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-
- 20 methanone;
- [(S)-2-(1-Ethyl-5,6-dimethyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(4-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- [(S)-2-(5,6-Dimethoxy-1-methyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(4,5-dimethyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- 25 (4,5-Dimethyl-2-[1,2,3]triazol-2-yl-phenyl)-[(S)-2-(1-ethyl-5,6-dimethoxy-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-methanone;
- (4'-Fluoro-4-methyl-biphenyl-2-yl)-[(S)-2-methyl-2-(1,5,6-trimethyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- [(S)-2-(1-Ethyl-5,6-dimethyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(4'-fluoro-4-methyl-biphenyl-2-yl)-
- 30 methanone;
- (4,5-Dimethyl-2-[1,2,3]triazol-2-yl-phenyl)-[(S)-2-methyl-2-(1,5,6-trimethyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- [(S)-2-(5-Chloro-1,4-dimethyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(6-methyl-3-[1,2,3]triazol-2-yl-pyridin-2-yl)-methanone;

- [(S)-2-(5-Chloro-1,4-dimethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- [(S)-2-(6-Chloro-1-methyl-5-trifluoromethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- 5 [(S)-2-(5-Chloro-1-ethyl-4-methyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- [(S)-2-(5-Chloro-1,4-dimethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-chloro-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- [(S)-2-(5-Chloro-1,4-dimethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methoxy-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- 10 (4,3'-Dimethyl-biphenyl-2-yl)-[(S)-2-methyl-2-(1,5,6-trimethyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- (4,3'-Dimethyl-biphenyl-2-yl)-[(S)-2-(1-ethyl-5,6-dimethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-methanone;
- (4,3'-Dimethyl-biphenyl-2-yl)-{(S)-2-[1-(2-fluoro-ethyl)-5,6-dimethyl-1H-benzimidazol-2-yl]-2-methyl-pyrrolidin-1-yl}-methanone;
- 15 (4,3'-Dimethyl-biphenyl-2-yl)-{(S)-2-[1-(2-methoxy-ethyl)-5,6-dimethyl-1H-benzimidazol-2-yl]-2-methyl-pyrrolidin-1-yl}-methanone;
- (3'-Fluoro-4-methyl-biphenyl-2-yl)-[(S)-2-methyl-2-(1,5,6-trimethyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- 20 [(S)-2-(1-Ethyl-5,6-dimethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(3'-fluoro-4-methyl-biphenyl-2-yl)-methanone;
- [(S)-2-(5-Chloro-1,4-dimethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-[5-(3-chloro-phenyl)-2-methyl-thiazol-4-yl]-methanone;
- [(S)-2-(1-Cyclopropylmethyl-5,6-dimethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(3'-fluoro-4-methyl-biphenyl-2-yl)-methanone;
- 25 {(S)-2-[1-(2-Fluoro-ethyl)-5,6-dimethyl-1H-benzimidazol-2-yl]-2-methyl-pyrrolidin-1-yl}-(3'-fluoro-4-methyl-biphenyl-2-yl)-methanone;
- (4,4'-Dimethyl-biphenyl-2-yl)-[(S)-2-methyl-2-(1,5,6-trimethyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- (4,4'-Dimethyl-biphenyl-2-yl)-[(S)-2-(1-ethyl-5,6-dimethyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-methanone;
- 30 (4,4'-Dimethyl-biphenyl-2-yl)-{(S)-2-[1-(2-fluoro-ethyl)-5,6-dimethyl-1H-benzimidazol-2-yl]-2-methyl-pyrrolidin-1-yl}-methanone;
- (3'-Methoxy-4-methyl-biphenyl-2-yl)-[(S)-2-methyl-2-(1,5,6-trimethyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;

- (2'-Fluoro-4-methyl-biphenyl-2-yl)-[(S)-2-methyl-2-(1,5,6-trimethyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- [(S)-2-(1-Ethyl-5,6-dimethyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(3'-methoxy-4-methyl-biphenyl-2-yl)-methanone;
- 5 {(S)-2-[1-(2-Fluoro-ethyl)-5,6-dimethyl-1H-benzoimidazol-2-yl]-2-methyl-pyrrolidin-1-yl}-(3'-methoxy-4-methyl-biphenyl-2-yl)-methanone;
- {(S)-2-[1-(2-Fluoro-ethyl)-5,6-dimethyl-1H-benzoimidazol-2-yl]-2-methyl-pyrrolidin-1-yl}-(2'-fluoro-4-methyl-biphenyl-2-yl)-methanone;
- [(S)-2-(5,6-Dimethoxy-1-methyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- 10 [(2S,3S)-2-(5-Chloro-6-methyl-1H-benzoimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(5-methyl-2-pyridin-2-yl-phenyl)-methanone;
- (4-Chloro-biphenyl-2-yl)-[(2S,3S)-2-(5-chloro-6-methyl-1H-benzoimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-methanone;
- 15 [(2S,3S)-2-(5-Chloro-6-methyl-1H-benzoimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(5-methyl-2-pyridin-3-yl-phenyl)-methanone;
- [(2S,3S)-2-(5-Chloro-6-methyl-1H-benzoimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(4-fluoro-biphenyl-2-yl)-methanone;
- [(2S,3S)-2-(5-Chloro-6-methyl-1H-benzoimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(5-fluoro-2-pyridin-2-yl-phenyl)-methanone;
- 20 2-[(2S,3S)-2-(5-Chloro-6-methyl-1H-benzoimidazol-2-yl)-3-methyl-pyrrolidine-1-carbonyl]-biphenyl-4-carbonitrile;
- [(2S,3S)-2-(5-Chloro-6-methyl-1H-benzoimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(4-methyl-2-pyridin-2-yl-phenyl)-methanone;
- [(2S,3S)-2-(5-Chloro-6-methyl-1H-benzoimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(4-methyl-2-pyridin-3-yl-phenyl)-methanone;
- 25 [(2S,3S)-2-(5-Chloro-6-methyl-1H-benzoimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(5-methyl-biphenyl-2-yl)-methanone;
- [(2S,3S)-2-(5-Chloro-6-methyl-1H-benzoimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(3-fluoro-biphenyl-2-yl)-methanone;
- 30 [(2S,3S)-2-(5-Chloro-6-methyl-1H-benzoimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(2-fluoro-3-methyl-6-pyridin-2-yl-phenyl)-methanone;
- [(2S,3S)-2-(5-Chloro-6-methyl-1H-benzoimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(3-fluoro-4-methyl-biphenyl-2-yl)-methanone;
- [(2S,3S)-2-(5-Chloro-6-methyl-1H-benzoimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(4-methyl-biphenyl-2-yl)-methanone;
- 35

- [(2S,3S)-2-(5-Chloro-6-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(2-fluoro-3-methyl-6-pyridin-3-yl-phenyl)-methanone;
- [(2S,3S)-2-(4-Chloro-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-[5-(3,5-difluoro-phenyl)-2-methyl-thiazol-4-yl]-methanone;
- 5 [(2S,3S)-2-(4-Chloro-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-[5-(4-chloro-phenyl)-2-methyl-thiazol-4-yl]-methanone;
- [(2S,3S)-2-(4-Chloro-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-[5-(3-methoxy-phenyl)-2-methyl-thiazol-4-yl]-methanone;
- [(2S,3S)-2-(4-Chloro-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(2-methyl-5-m-tolyl-thiazol-4-yl)-methanone;
- 10 [5-(4-Bromo-phenyl)-2-methyl-thiazol-4-yl]-[(2S,3S)-2-(4-chloro-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-methanone;
- [(2S,3S)-2-(4-Chloro-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(2-methyl-5-p-tolyl-thiazol-4-yl)-methanone;
- [(2S,3S)-2-(4-Chloro-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-[5-(3,4-dimethyl-phenyl)-2-methyl-thiazol-4-yl]-methanone;
- 15 [(2S,3S)-2-(4-Chloro-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-[2-dimethylamino-5-(3,4-dimethyl-phenyl)-thiazol-4-yl]-methanone;
- [(2S,3S)-2-(5-Chloro-6-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(4-methyl-2-pyrimidin-2-yl-phenyl)-methanone;
- [(2S,3S)-2-(5-Chloro-6-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(5-fluoro-2-pyrimidin-2-yl-phenyl)-methanone;
- 20 [(2S,3S)-2-(5-Chloro-6-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(2-fluoro-6-pyrimidin-2-yl-phenyl)-methanone;
- [(2S,3S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-[2-(3-methoxy-phenylethynyl)-phenyl]-methanone;
- 25 [(2S,3S)-2-(5-Chloro-6-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-[2-(3-methoxy-phenylethynyl)-phenyl]-methanone;
- [5-(4-Fluoro-phenyl)-2-methyl-thiazol-4-yl]-[(2S,3S)-3-methyl-2-(5-trifluoromethyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- [5-(3-Methoxy-phenyl)-2-methyl-thiazol-4-yl]-[(2S,3S)-3-methyl-2-(5-trifluoromethyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- 30 (2-Methyl-5-m-tolyl-thiazol-4-yl)-[(2S,3S)-3-methyl-2-(5-trifluoromethyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- [5-(4-Chloro-phenyl)-2-methyl-thiazol-4-yl]-[(2S,3S)-3-methyl-2-(5-trifluoromethyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;

- (2-Methyl-5-p-tolyl-thiazol-4-yl)-[(2S,3S)-3-methyl-2-(5-trifluoromethyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- [5-(4-Methoxy-phenyl)-2-methyl-thiazol-4-yl]-[(2S,3S)-3-methyl-2-(5-trifluoromethyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- 5 [5-(4-Bromo-phenyl)-2-methyl-thiazol-4-yl]-[(2S,3S)-3-methyl-2-(5-trifluoromethyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- [5-(3,4-Difluoro-phenyl)-2-methyl-thiazol-4-yl]-[(2S,3S)-3-methyl-2-(5-trifluoromethyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- [2-(3,4-Dimethyl-phenyl)-thiophen-3-yl]-[(2S,3S)-3-methyl-2-(5-trifluoromethyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- 10 [5-(3,4-Dimethyl-phenyl)-2-methyl-thiazol-4-yl]-[(2S,3S)-3-methyl-2-(5-trifluoromethyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- [5-(2-Chloro-phenyl)-2-methyl-thiazol-4-yl]-[(2S,3S)-3-methyl-2-(5-trifluoromethyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- 15 [5-(2-Methoxy-phenyl)-2-methyl-thiazol-4-yl]-[(2S,3S)-3-methyl-2-(5-trifluoromethyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- [5-(3,5-Difluoro-phenyl)-2-methyl-thiazol-4-yl]-[(2S,3S)-3-methyl-2-(5-trifluoromethyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- [5-(4-Fluoro-phenyl)-2-methyl-thiazol-4-yl]-[(2S,3S)-3-methyl-2-(4-trifluoromethyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- 20 [5-(4-Chloro-phenyl)-2-methyl-thiazol-4-yl]-[(2S,3S)-3-methyl-2-(4-trifluoromethyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- [5-(4-Bromo-phenyl)-2-methyl-thiazol-4-yl]-[(2S,3S)-3-methyl-2-(4-trifluoromethyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- 25 (2-Methyl-5-p-tolyl-thiazol-4-yl)-[(2S,3S)-3-methyl-2-(4-trifluoromethyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- [5-(3,4-Difluoro-phenyl)-2-methyl-thiazol-4-yl]-[(2S,3S)-3-methyl-2-(4-trifluoromethyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- [5-(2-Methoxy-phenyl)-2-methyl-thiazol-4-yl]-[(2S,3S)-3-methyl-2-(4-trifluoromethyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- 30 [5-(3,5-Difluoro-phenyl)-2-methyl-thiazol-4-yl]-[(2S,3S)-3-methyl-2-(4-trifluoromethyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- (2-Methyl-5-m-tolyl-thiazol-4-yl)-[(2S,3S)-3-methyl-2-(4-trifluoromethyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone;

- [5-(4-Methoxy-phenyl)-2-methyl-thiazol-4-yl]-[(2S,3S)-3-methyl-2-(4-trifluoromethyl-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- [(2S,3S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(5-methyl-2-pyridin-2-yl-phenyl)-methanone;
- 5 (4-Chloro-biphenyl-2-yl)-[(2S,3S)-2-(5-chloro-4-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-methanone;
- [(2S,3S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(5-methyl-2-pyridin-3-yl-phenyl)-methanone;
- [(2S,3S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(4-fluoro-biphenyl-2-yl)-
- 10 methanone;
- [(2S,3S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(5-fluoro-2-pyridin-2-yl-phenyl)-methanone;
- 2-[(2S,3S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidine-1-carbonyl]-biphenyl-4-carbonitrile;
- [(2S,3S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(4-methyl-2-pyridin-2-yl-phenyl)-
- 15 methanone;
- [(2S,3S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(5-methyl-biphenyl-2-yl)-methanone;
- [(2S,3S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(3-fluoro-biphenyl-2-yl)-methanone;
- 20 [(2S,3S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(4-trifluoromethyl-biphenyl-2-yl)-methanone;
- [(2S,3S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(3-fluoro-4-methyl-biphenyl-2-yl)-methanone;
- [(2S,3S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(4-methyl-biphenyl-2-yl)-
- 25 methanone;
- [(2S,3S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(2-fluoro-3-methyl-6-pyridin-3-yl-phenyl)-methanone;
- [(2S,3S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(4-methyl-2-pyrimidin-2-yl-phenyl)-methanone;
- 30 [(2S,3S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(5-fluoro-2-pyrimidin-2-yl-phenyl)-methanone;
- [(2S,3S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(2-fluoro-3-methyl-6-pyrimidin-2-yl-phenyl)-methanone;
- [(2S,3S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(2-fluoro-6-pyrimidin-2-yl-phenyl)-
- 35 methanone;

- [(2S,3S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(5-methyl-2-pyrimidin-2-yl-phenyl)-methanone;
- [(2S,3S)-2-(5-Chloro-6-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(5-methyl-2-pyrimidin-2-yl-phenyl)-methanone;
- 5 [(2S,3S)-2-(6-Chloro-7-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(5-methyl-2-pyrimidin-2-yl-phenyl)-methanone;
- [(2S,3S)-2-(5-Chloro-7-methyl-1H-benzimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(5-methyl-2-pyrimidin-2-yl-phenyl)-methanone;
- 10 [(S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-[5-(4-fluoro-phenyl)-2-phenyl-thiazol-4-yl]-methanone;
- [2,5-Bis-(4-fluoro-phenyl)-thiazol-4-yl]-[(S)-2-(5-chloro-4-methyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-methanone;
- [(S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-[5-(4-fluoro-phenyl)-2-m-tolyl-thiazol-4-yl]-methanone;
- 15 [(S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-[5-(4-fluoro-phenyl)-2-(3-methoxy-phenyl)-thiazol-4-yl]-methanone;
- [(S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-[5-(4-fluoro-phenyl)-2-p-tolyl-thiazol-4-yl]-methanone;
- [(S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-[5-(4-fluoro-phenyl)-2-(2-fluoro-phenyl)-thiazol-4-yl]-methanone;
- 20 [(S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-[5-(4-fluoro-phenyl)-2-(3-fluoro-phenyl)-thiazol-4-yl]-methanone;
- (4-Methyl-biphenyl-2-yl)-[(S)-2-methyl-2-(5-trifluoromethoxy-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- [5-(3,5-Difluoro-phenyl)-2-methyl-thiazol-4-yl]-[(S)-2-methyl-2-(5-trifluoromethoxy-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- 25 [5-(3-Methoxy-phenyl)-2-methyl-thiazol-4-yl]-[(S)-2-methyl-2-(5-trifluoromethoxy-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- [5-(4-Fluoro-phenyl)-2-methyl-thiazol-4-yl]-[(S)-2-methyl-2-(5-trifluoromethoxy-1H-benzimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- 30 [(S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-[2-(2,3-difluoro-phenyl)-5-(4-fluoro-phenyl)-thiazol-4-yl]-methanone;
- [(S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-[2-(4-fluoro-3-methyl-phenyl)-5-(4-fluoro-phenyl)-thiazol-4-yl]-methanone;
- [(S)-2-(5-Chloro-4-methyl-1H-benzimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-[2-(2-fluoro-4-methyl-phenyl)-5-(4-fluoro-phenyl)-thiazol-4-yl]-methanone;
- 35

- 3-[4-[(S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidine-1-carbonyl]-5-(4-fluoro-phenyl)-thiazol-2-yl]-benzonitrile;
- [(S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-[2-(2-chloro-phenyl)-5-(4-fluoro-phenyl)-thiazol-4-yl]-methanone;
- 5 [(S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-[2-(3-chloro-phenyl)-5-(4-fluoro-phenyl)-thiazol-4-yl]-methanone;
- [(S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-[2-(4-fluoro-2-methyl-phenyl)-5-(4-fluoro-phenyl)-thiazol-4-yl]-methanone;
- [(S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-[2-(4-fluoro-2-methoxy-phenyl)-5-(4-fluoro-phenyl)-thiazol-4-yl]-methanone;
- 10 [(S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-[2-(2,5-difluoro-phenyl)-5-(4-fluoro-phenyl)-thiazol-4-yl]-methanone;
- [(S)-2-(6-Bromo-4,5-dimethyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(3,4-dimethyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- 15 3'-[(S)-2-Methyl-2-(5-trifluoromethyl-1H-benzoimidazol-2-yl)-pyrrolidine-1-carbonyl]-4'-[1,2,3]triazol-2-yl-biphenyl-3-carbonitrile;
- 2'-[(S)-2-(5-Chloro-4-methyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidine-1-carbonyl]-3'-fluoro-4'-methoxy-biphenyl-4-carbonitrile;
- [(S)-2-(6-Bromo-5-fluoro-1-methyl-1H-benzoimidazol-2-yl)-2-methyl-pyrrolidin-1-yl]-(5-methyl-2-[1,2,3]triazol-2-yl-phenyl)-methanone;
- 20 (3'-Fluoro-4-methyl-biphenyl-2-yl)-{(S)-2-[1-(2-methoxy-ethyl)-5,6-dimethyl-1H-benzoimidazol-2-yl]-2-methyl-pyrrolidin-1-yl}-methanone;
- [(2S,3S)-2-(5-Chloro-6-methyl-1H-benzoimidazol-2-yl)-3-methyl-pyrrolidin-1-yl]-(2-fluoro-3-methyl-6-pyrimidin-2-yl-phenyl)-methanone;
- 25 (4-Bromo-biphenyl-2-yl)-[(S)-2-methyl-2-(5-trifluoromethyl-1H-benzoimidazol-2-yl)-pyrrolidin-1-yl]-methanone;
- [(S)-2-(5,6-Dimethyl-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(2-methyl-6-[1,2,3]triazol-2-yl-phenyl)-methanone;
- [(S)-2-(5,6-Dimethyl-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(5-methyl-2-pyrazol-1-yl-phenyl)-methanone; and
- 30 [(S)-2-(5,6-Dimethyl-1H-benzoimidazol-2-yl)-4-methylene-pyrrolidin-1-yl]-(6-methyl-3-[1,2,3]triazol-2-yl-pyridin-2-yl)-methanone;

or a pharmaceutically acceptable salt thereof.

12. A pharmaceutical composition comprising, as active principle, one or more compounds according to any one of claims 1 to 11, or a pharmaceutically acceptable salt thereof, and at least one therapeutically inert excipient.

13. A pharmaceutical composition according to claim 12 for use in the prevention or treatment of mental health diseases or disorders relating to orexinergic dysfunctions.

14. A compound according to any one of claims 1 to 11, or a pharmaceutically acceptable salt thereof, for use as a medicament.

15. A compound according to any one of claims 1 to 11, or a pharmaceutically acceptable salt thereof, for use in the prevention or treatment of diseases or disorders selected from sleep disorders, anxiety disorders, addiction disorders, cognitive dysfunctions, mood disorders, and appetite disorders.

16. A compound according to any one of claims 1 to 11, or a pharmaceutically acceptable salt thereof, for use in the preparation of a medicament for the prevention or treatment of diseases or disorders selected from sleep disorders, anxiety disorders, addiction disorders, cognitive dysfunctions, mood disorders, and appetite disorders.

17. A method to treat a disease or disorder selected from a sleep disorder, an anxiety disorder, an addiction disorder, a cognitive dysfunction, a mood disorder, or an appetite disorder; comprising administering to a patient in need thereof, the compound of claim 1 or 11 in free or pharmaceutically acceptable salt form.