

COMBINATION THERAPY OF T CELL ACTIVATING BISPECIFIC ANTIGEN BINDING MOLECULES
CD3 ABD FOLATE RECEPTOR 1 (FOLR1) AND PD-1 AXIS BINDING ANTAGONISTS

SEQUENCE LISTING

The instant application contains a Sequence Listing which has been submitted electronically in ASCII format and is hereby incorporated by reference in its entirety. Said ASCII copy, created on November 11, 2015, is named 32401_SL.txt and is 527,137 bytes in size.

FIELD OF THE INVENTION

The present invention relates to combination therapies employing T cell activating bispecific antigen binding molecule and a PD-1 axis binding antagonist, and, optionally, a TIM3 antagonist, and the use of these combination therapies for the treatment of cancer.

BACKGROUND

Monoclonal antibodies are powerful therapeutic agents for the treatment of cancer that selectively target antigens which are differentially expressed on cancer cells.

Bispecific antibodies designed to bind with one antigen binding moiety to a surface antigen on target cells, and with the second antigen binding moiety to an activating, invariant component of the T cell receptor (TCR) complex, have become of interest in recent years. The simultaneous binding of such an antibody to both of its targets will force a temporary interaction between target cell and T cell, causing activation of any cytotoxic T cell and subsequent lysis of the target cell. Hence, the immune response is re-directed to the target cells and is independent of peptide antigen presentation by the target cell or the specificity of the T cell as would be relevant for normal MHC-restricted activation of CTLs. In this context it is crucial that CTLs are only activated when a target cell is presenting the bispecific antibody to them, i.e., the immunological synapse is mimicked. Particularly desirable are bispecific antibodies that do not require lymphocyte preconditioning or co-stimulation in order to elicit efficient lysis of target cells. It is not well understood how TCBs affect the T cell itself beyond activation of certain effector function.

Activation of resting T lymphocytes, or T cells, by antigen-presenting cells (APCs) appears to require two signal inputs. Lafferty et al, Aust. J. Exp. Biol. Med. ScL 53: 27-42

(1975). The primary, or antigen specific, signal is transduced through the T-cell receptor (TCR) following recognition of foreign antigen peptide presented in the context of the major histocompatibility-complex (MHC). The second, or co-stimulatory, signal is delivered to T-cells by co-stimulatory molecules expressed on antigen-presenting cells (APCs), and promotes T-cell clonal expansion, cytokine secretion and effector function. 5 (APCs), and promotes T-cell clonal expansion, cytokine secretion and effector function. Lenschow et al., Ann. Rev. Immunol. 14:233 (1996). In the absence of co-stimulation, T cells can become refractory to antigen stimulation, do not mount an effective immune response, and may result in exhaustion or tolerance to foreign antigens.

T cells can receive both positive and negative secondary co- stimulatory signals. 10 The balance of positive and negative signals is important to elicit effective immune responses, while maintaining immune tolerance and preventing autoimmunity. Negative secondary signals appear necessary for induction of T-cell tolerance, while positive signals promote T cell activation.

Recently, it has been discovered that T cell dysfunction or anergy occurs 15 concurrently with an induced and sustained expression of the inhibitory receptor, programmed death 1 polypeptide (PD-1). One of its ligands, PD-L1 is overexpressed in many cancers and is often associated with poor prognosis (Okazaki T et al., Intern. Immun. 2007 19(7):813) (Thompson RH et al., Cancer Res 2006, 66(7):3381). Interestingly, the majority of tumor infiltrating T lymphocytes predominantly express PD- 20 1, in contrast to T lymphocytes in normal tissues and peripheral blood T lymphocytes indicating that up-regulation of PD-1 on tumor-reactive T cells can contribute to impaired antitumor immune responses (Blood 2009 1 14(8): 1537).

T cell Immunoglobulin- and Mucin domain-containing molecule 3 (TIM3), is important in immune regulation. This cell surface protein is expressed, preferentially, by 25 type 1 T helper cells and has been implicated in the regulation of macrophage activation, inflammatory conditions and cancer (Majeti R et al., PNAS, 106 (2009) 3396-3401 and WO2009/091547). Binding of TIM-3 to one of its ligands (e.g., galectin-9) can suppress the Th1 response by inducing programmed cell death, thereby supporting peripheral tolerance. Treatment with TIM-3 siRNA or with an anti-TIM-3 antagonist antibody 30 increases secretion of interferon alpha from CD4 positive T-cells, supporting the inhibitory role of TIM-3 in human T cells. Examples of the anti-TIM-3 monoclonal antibodies include are disclosed in WO2013/06490 and US2012/189617 (Ngiow et al., Cancer Res 7:6567 (2011)).

FOLR1 is expressed on tumor cells of various origins, e.g., ovarian and lung cancer. Several approaches to target FOLR1 with therapeutic antibodies, such as farletuzumab, antibody drug conjugates, or adoptive T cell therapy for imaging of tumors have been described (Kandalafi et al., *J Transl Med.* 2012 Aug 3;10:157. doi: 10.1186/1479-5876-
5 10-157; van Dam et al., *Nat Med.* 2011 Sep 18;17(10):1315-9. doi: 10.1038/nm.2472; Clifton et al., *Hum Vaccin.* 2011 Feb;7(2):183-90. Epub 2011 Feb 1; Kelemen et al., *Int J Cancer.* 2006 Jul 15;119(2):243-50; Vaitilingam et al., *J Nucl Med.* 2012 Jul;53(7); Teng et al., 2012 Aug;9(8):901-8. doi: 10.1517/17425247.2012.694863. Epub 2012 Jun 5.

Some attempts have been made to target folate receptor-positive tumors with constructs
10 that target the folate receptor and CD3 (Kranz et al., *Proc Natl Acad Sci U S A.* Sep 26, 1995; 92(20): 9057–9061; Roy et al., *Adv Drug Deliv Rev.* 2004 Apr 29;56(8):1219-31; Huiting Cui et al *Biol Chem.* Aug 17, 2012; 287(34): 28206–28214; Lamers et al., *Int. J. Cancer.* 60(4):450 (1995); Thompson et al., *MAbs.* 2009 Jul-Aug;1(4):348-56. Epub 2009 Jul 19; Mezzanzanica et al., *Int. J. Cancer,* 41, 609–615 (1988).

15 There remains a need for such an optimal therapy for treating, stabilizing, preventing, and/or delaying development of various cancers.

SUMMARY

Broadly, the present disclosure relates to bispecific antibodies combining a Folate Receptor 1 (FolR1) targeting antigen binding site with a second antigen binding site that
20 targets CD3 and their use in combination with a PD-1 axis binding antagonist, e.g., for the treatment of cancer. In one embodiment, the combination further comprises a TIM3 antagonist. The methods and combinations of the present disclosure enable enhanced immunotherapy. The advantage over conventional treatment is the specificity of inducing T cell activation only at the site where FolR1 is expressed as well as the reduction and/or
25 reversal of low T cell mediated activity also termed T cell exhaustion due to the combination with a PD-1 axis binding antagonist, and, optionally, a TIM3 antagonist.

Accordingly, the present disclosure provides a method for treating or delaying progression of a cancer in an individual comprising administering to the individual an effective amount of a T cell activating bispecific antigen binding molecule and a PD-1
30 axis binding antagonist. In one aspect, the present disclosure provides use of a T cell activating bispecific antigen binding molecule and a PD-1 axis binding antagonist in the manufacture of a medicament for combination therapy in the treatment of or delaying the

progression of a cancer, wherein the T cell activating bispecific antigen binding molecule comprises a first Fab fragment capable of specific binding to CD3, a second Fab fragment capable of specific binding to Folate Receptor 1 (FolR1), wherein the PD-1 axis binding antagonist inhibits the binding of PD-1 to its ligand binding partners, wherein the PD-1

- 5 axis binding antagonist is selected from the group consisting of an anti-PD-1 antibody, an anti-PD-L1 antibody and an anti-PD-L2 antibody, and wherein the cancer is a FolR1-positive cancer.

In one embodiment, the T cell activating bispecific antigen binding molecule comprises a first antigen binding moiety capable of specific binding to CD3 and a second antigen

- 10 binding moiety capable of specific binding to Folate Receptor 1 (FolR1). In one embodiment, the first antigen binding moiety comprises at

least one heavy chain complementarity determining region (CDR) amino acid sequence selected from the group consisting of SEQ ID NO: 37, SEQ ID NO: 38 and SEQ ID NO: 39 and at least one light chain CDR selected from the group of SEQ ID NO: 32, SEQ ID NO: 33, SEQ ID NO: 34. In one embodiment, the first antigen binding moiety comprises
5 a variable heavy chain comprising an amino acid sequence of SEQ ID NO: 36 and a variable light chain comprising an amino acid sequence of SEQ ID NO: 31. In one embodiment, the T cell activating bispecific antigen binding molecule further comprises a third antigen binding moiety capable of specific binding to FolR1. In one embodiment, the second and third antigen binding moiety capable of specific binding to FolR1
10 comprise identical heavy chain complementarity determining region (CDR) and light chain CDR sequences. In one embodiment, the third antigen binding moiety is identical to the second antigen binding moiety. In one embodiment, at least one of the first, second and third antigen binding moiety is a Fab molecule.

In one embodiment, the antigen binding moiety capable of specific binding to Folate
15 Receptor 1 (FolR1) comprises at least one heavy chain complementarity determining region (CDR) amino acid sequence selected from the group consisting of SEQ ID NO: 16, SEQ ID NO: 17 and SEQ ID NO: 18 and at least one light chain CDR selected from the group of SEQ ID NO: 32, SEQ ID NO: 33, SEQ ID NO: 34. In one embodiment, the antigen binding moiety capable of specific binding to Folate Receptor 1 (FolR1)
20 comprises a variable heavy chain comprising an amino acid sequence of SEQ ID NO: 15 and a variable light chain comprising an amino acid sequence of SEQ ID NO: 31. In one embodiment, the antigen binding moiety capable of specific binding to Folate Receptor 1 (FolR1) comprises at least one heavy chain complementarity determining region (CDR) amino acid sequence selected from the group consisting of SEQ ID NO: 8, SEQ ID NO: 56 and SEQ ID NO: 57 and at least one light chain CDR selected from the group of SEQ
25 ID NO: 59, SEQ ID NO: 60, SEQ ID NO: 65. In one embodiment, the antigen binding moiety capable of specific binding to Folate Receptor 1 (FolR1) comprises a variable heavy chain comprising an amino acid sequence of SEQ ID NO: 55 and a variable light chain comprising an amino acid sequence of SEQ ID NO: 64. In one embodiment, the
30 antigen binding moiety capable of specific binding to Folate Receptor 1 (FolR1) comprises at least one heavy chain complementarity determining region (CDR) amino acid sequence selected from the group consisting of SEQ ID NO: 8, SEQ ID NO: 9 and SEQ ID NO: 50 and at least one light chain CDR selected from the group of SEQ ID NO:

52, SEQ ID NO: 53, SEQ ID NO: 54. In one embodiment, the antigen binding moiety capable of specific binding to FolR1 comprises:

- a) a complementarity determining region heavy chain 1 (CDR-H1) amino acid sequences of SEQ ID NO: 8;
- 5 (b) a CDR-H2 amino acid sequence of SEQ ID NO: 9;
- (c) a CDR-H3 amino acid sequence of SEQ ID NO: 50;
- (d) a complementarity determining region light chain 1 (CDR-L1) amino acid sequence of SEQ ID NO: 52;
- (e) a CDR-L2 amino acid sequence of SEQ ID NO: 53, and
- 10 (f) a CDR-L3 amino acid sequence of SEQ ID NO: 54.

In one such embodiment, the antigen binding moiety capable of specific binding to FolR1 comprises a variable heavy chain comprising an amino acid sequence of SEQ ID NO: 49 and a variable light chain comprising an amino acid sequence of SEQ ID NO: 51.

15

In one embodiment, the T cell activating bispecific antigen binding molecule binds to a human FolR1, a cynomolgus monkey FolR1 and a murine FolR1.

In one embodiment, the T cell activating bispecific antigen binding molecule induces proliferation of a human CD3 positive T cell in vitro.

- 20 In one embodiment, the T cell activating bispecific antigen binding molecule induces human peripheral blood mononuclear cell mediated killing of a FolR1-expressing human tumor cell in vitro.

In one embodiment, the T cell activating bispecific antigen binding molecule induces T cell mediated killing of a FolR1-expressing human tumor cell in vitro. In one

- 25 embodiment, the T cell activating bispecific antigen binding molecule induces T cell mediated killing of the FolR1-expressing human tumor cell in vitro with an EC₅₀ of between about 36 pM and about 39573 pM after 24 hours. In one embodiment, the T cell activating bispecific antigen binding molecule induces upregulation of cell surface expression of at least one of CD25 and CD69 on the T cell as measured by flow
- 30 cytometry. In one embodiment, the T cell activating bispecific antigen binding molecule binds human FolR1 with an apparent K_D of about 5.36 pM to about 4 nM. In one embodiment, the T cell activating bispecific antigen binding molecule binds human and cynomolgus FolR1 with an apparent K_D of about 4 nM. In one embodiment, the T cell

activating bispecific antigen binding molecule binds murine FolR1 with an apparent K_D of about 1.5 nM. In one embodiment, the T cell activating bispecific antigen binding molecule binds human FolR1 with a monovalent binding K_D of at least about 1000 nM. In one embodiment, the T cell activating bispecific antigen binding molecule binds to

5 FolR1 expressed on a human tumor cell. In one embodiment, the T cell activating bispecific antigen binding molecule binds to a conformational epitope on human FolR1. In one embodiment, the T cell activating bispecific antigen binding molecule does not bind to human Folate Receptor 2 (FolR2) or to human Folate Receptor 3 (FolR3). In one embodiment, the antigen binding moiety binds to a FolR1 polypeptide comprising the

10 amino acids 25 to 234 of human FolR1 (SEQ ID NO:227). In one embodiment, the FolR1 antigen binding moiety binds to a FolR1 polypeptide comprising the amino acid sequence of SEQ ID NOs:227, 230 and 231, and wherein the FolR1 antigen binding moiety does not bind to a FolR polypeptide comprising the amino acid sequence of SEQ ID NOs:228 and 229. In one embodiment, the T cell activating bispecific antigen binding

15 molecule comprises a) a first antigen-binding site that competes for binding to human FolR1 with a reference antibody comprising a variable heavy chain domain (VH) of SEQ ID NO: 49 and a variable light chain domain of SEQ ID NO: 51; and b) a second antigen-binding site that competes for binding to human CD3 with a reference antibody comprising a variable heavy chain domain (VH) of SEQ ID NO: 36 and a variable light

20 chain domain of SEQ ID NO: 31, wherein binding competition is measured using a surface plasmon resonance assay.

In one embodiment, the T cell activating bispecific antigen binding molecule comprises a first, a second, a third, a fourth and a fifth polypeptide chain that form a first, a second and a third antigen binding moiety, wherein the first antigen binding moiety is capable of

25 binding CD3 and the second and the third antigen binding moiety each are capable of binding Folate Receptor 1 (FolR1), wherein a) the first and the second polypeptide chain comprise, in amino (N)-terminal to carboxyl (C)-terminal direction, VLD1 and CLD1; b) the third polypeptide chain comprises, in N-terminal to C-terminal direction, VLD2 and CH1D2; c) the fourth polypeptide chain comprises, in N-terminal to C-terminal direction,

30 VHD1, CH1D1, CH2D1 and CH3D1; d) the fifth polypeptide chain comprises VHD1, CH1D1, VHD2, CLD2, CH2D2 and CH3D2; wherein

VLD1 is a first light chain variable domain

VLD2 is a second light chain variable domain

CLD1 is a first light chain constant domain

CLD2 is a second light chain constant domain

VHD1 is a first heavy chain variable domain

VHD2 is a second heavy chain variable domain

5 CH1D1 is a first heavy chain constant domain 1

CH1D2 is a second heavy chain constant domain 1

CH2D1 is a first heavy chain constant domain 2

CH2D2 is a second heavy chain constant domain 2

CH3D1 is a first heavy chain constant domain 3

10 CH3D2 is a second heavy chain constant domain 3.

In one such embodiment,

- a. the third polypeptide chain and VHD2 and CLD2 of the fifth polypeptide chain form the first antigen binding moiety capable of binding CD3;
- b. the first polypeptide chain and VHD1 and CH1D1 of the fourth polypeptide chain form the second binding moiety capable of binding to FolR1; and
- c. the second polypeptide chain and VHD1 and CH1D1 of the fifth polypeptide chain form the third binding moiety capable of binding to FolR1.

20 In one such embodiment, the first and second polypeptide chain comprise the amino acid sequence of SEQ ID NO:399. In one such embodiment, the third polypeptide chain comprises the amino acid sequence of SEQ ID NO:86. In one such embodiment, the fourth polypeptide chain comprises the amino acid sequence of SEQ ID NO:394. In one such embodiment, the fifth polypeptide chain comprises the amino acid sequence of SEQ ID NO:397. In one embodiment,

- a. the first and second polypeptide chain comprise the amino acid sequence of SEQ ID NO:399;
- b. the third polypeptide chain comprises the amino acid sequence of SEQ ID NO:86;
- c. the fourth polypeptide chain comprises the amino acid sequence of SEQ ID NO:394; and
- d. the fifth polypeptide chain comprise the amino acid sequence of SEQ ID NO:397.

In some embodiments, the bispecific antibody is bivalent both for FolR1 and CD3.

In some embodiments, the bispecific antibody comprises one or more Fab fragment(s) comprising an antigen binding site specific for CD3, wherein the variable
5 regions or the constant regions of the heavy and light chain are exchanged.

In some embodiments, the bispecific antibody comprises an Fc domain, at least one Fab fragment comprising the antigen binding site specific for FolR1, and at least one Fab fragment comprising the antigen binding site specific for CD3 wherein either the
10 variable regions or the constant regions of the heavy and light chain of at least one Fab fragment are exchanged.

In some embodiments, the bispecific antibody comprises:

- a) an Fc domain,
- 15 b) a first and second Fab fragment each comprising an antigen binding site specific for FolR1,
- c) a third Fab fragment comprising an antigen binding site specific for CD3, wherein the third Fab fragment is connected at the C-terminus of the variable heavy chain (VH) to the second subunit of the Fc domain and wherein the third Fab fragment is
20 connected at the N-terminus of the variable heavy chain to the C-terminus of the second Fab fragment.

In one embodiment at least one of said Fab fragments is connected to the Fc domain via a peptide linker.

In one embodiment said bispecific antibody comprises an Fc domain, which comprises
25 one or more amino acid substitution that reduces binding to Fc receptors and/or effector function. In one embodiment said one or more amino acid substitution is at one or more positions selected from the group of L234, L235, and P329. In one embodiment each subunit of the Fc domain comprises three amino acid substitutions that abolish binding to an activating or inhibitory Fc receptor and/or effector function wherein said amino acid
30 substitutions are L234A, L235A and P329G.

In some embodiments, the PD-1 axis binding antagonist is selected from the group consisting of a PD-1 binding antagonist, a PDL1 binding antagonist and a PDL2 binding antagonist.

In some embodiments, the PD-1 axis binding antagonist is a PD-1 binding antagonist. In some embodiments, the PD-1 binding antagonist inhibits the binding of PD-1 to its ligand binding partners. In some embodiments, the PD-1 binding antagonist inhibits the binding of PD-1 to PDL1. In some embodiments, the PD-1 binding antagonist inhibits the binding of PD-1 to PDL2. In some embodiments, the PD-1 binding antagonist inhibits the binding of PD-1 to both PDL1 and PDL2. In some embodiments, PD-1 binding antagonist is an antibody. In some embodiments, the anti-PD-1 antibody is a monoclonal antibody. In some embodiments, the anti-PD-1 antibody is an antibody fragment selected from the group consisting of Fab, Fab'-SH, Fv, scFv, and (Fab')₂ fragments. In some embodiments, PD-1 binding antagonist is nivolumab, pembrolizumab, CT-011, or AMP-224.

In some embodiments, the PD-1 axis binding antagonist is a PDL1 binding antagonist. In some embodiments, the PDL1 binding antagonist inhibits the binding of PDL1 to PD-1. In some embodiments, the PDL1 binding antagonist inhibits the binding of PDL1 to B7-1. In some embodiments, the PDL1 binding antagonist inhibits the binding of PDL1 to both PD-1 and B7-1. In some embodiments, the PDL1 binding antagonist is an anti-PDL1 antibody. In some embodiments, the anti-PDL1 antibody is a monoclonal antibody. In some embodiments, the anti-PDL1 antibody is an antibody fragment selected from the group consisting of Fab, Fab'-SH, Fv, scFv, and (Fab')₂ fragments. In some embodiments, the anti-PDL1 antibody is a humanized antibody or a human antibody. In some embodiments, the PDL1 binding antagonist is selected from the group consisting of: YW243.55.S70, MPDL3280A, MDX-1105, and MEDI4736.

In some embodiments, the anti-PDL1 antibody comprises a heavy chain comprising HVR-H1 sequence of SEQ ID NO:289, HVR-H2 sequence of SEQ ID NO:290, and HVR-H3 sequence of SEQ ID NO:291; and a light chain comprising HVR-L1 sequence of SEQ ID NO:292, HVR-L2 sequence of SEQ ID NO:293, and HVR-L3 sequence of SEQ ID NO:294. In some embodiments, anti-PDL1 antibody comprises a heavy chain variable region comprising the amino acid sequence of SEQ ID NO:280 or SEQ ID NO:281 and a light chain variable region comprising the amino acid sequence of SEQ ID NO:383. In some embodiments, the anti-PDL1 antibody comprises a heavy chain comprising the amino acid sequence of SEQ ID NO:278 and/or a light chain comprising the amino acid sequence of SEQ ID NO:279.

In some embodiments, the PD-1 axis binding antagonist is a PDL2 binding antagonist. In some embodiments, PDL2 binding antagonist is an antibody. In some embodiments, the

anti-PDL2 antibody is a monoclonal antibody. In some embodiments, the anti-PDL2 antibody is an antibody fragment selected from the group consisting of Fab, Fab'-SH, Fv, scFv, and (Fab')₂ fragments. In some embodiments, PDL2 binding antagonist is an immunoadhesin.

- 5 In one embodiment, the method of any of the above embodiments further comprises administering to the individual a T cell immunoglobulin mucin 3 (TIM3) antagonist. In one embodiment, the TIM3 antagonist is an anti-TIM3 antibody. In one embodiment, the anti-TIM3 antibody induces internalization of TIM3 on a TIM3 expressing cell of at least 45% after 120 Minutes at 37 °C wherein internalization is measured by FACS analysis. In
- 10 one embodiment, the anti-TIM3 antibody has one or more of the following properties:
- a) competes for binding to TIM3 with an anti-Tim3 antibody comprising the VH of SEQ ID NO:7 and VL of of SEQ ID NO: 8
 - b) binds to a human and cynomolgous TIM3
 - c) shows as immunoconjugate a cytotoxic activity on TIM3 expressing cells
 - 15 d) induces interferon-gamma release.

In one embodiment, the anti-TIM3 antibody has one or more of the following properties:

- a. competes for binding to TIM3 with an anti-Tim3 antibody comprising the VH of SEQ ID NO:7 and VL of of SEQ ID NO: 8
- b. binds to a human and cynomolgous TIM3
- 20 c. shows as immunoconjugate a cytotoxic activity on TIM3 expressing cells
- d. induces interferon-gamma release.

- In one embodiment, the anti-TIM3 antibody is a monoclonal antibody. In one embodiment, the anti-TIM3 antibody is a human, humanized, or chimeric antibody. In one embodiment, the anti-TIM3 antibody is an antibody fragment that binds to TIM3. In
- 25 one embodiment, the anti-TIM3 antibody is Fab fragment. In one embodiment, the anti-TIM3 antibody comprises:

- A) (a) a VH domain comprising (i) HVR-H1 comprising the amino acid sequence of SEQ ID NO:304, (ii) HVR-H2 comprising the amino acid sequence of SEQ ID NO:305, and (iii) HVR-H3 comprising an amino acid sequence selected from SEQ
- 30 ID NO:306; and (b) a VL domain comprising (i) HVR-L1 comprising the amino acid sequence of SEQ ID NO:307; (ii) HVR-L2 comprising the amino acid sequence of SEQ ID NO:308 and (iii) HVR-L3 comprising the amino acid sequence of SEQ ID NO:309; or

B) (a) a VH domain comprising (i) HVR-H1 comprising the amino acid sequence of SEQ ID NO:304, (ii) HVR-H2 comprising the amino acid sequence of SEQ ID NO:305, and (iii) HVR-H3 comprising an amino acid sequence selected from SEQ ID NO:306; and (b) a VL domain comprising (i) HVR-L1 comprising the amino acid sequence of SEQ ID NO:314; (ii) HVR-L2 comprising the amino acid sequence of SEQ ID NO:308 and (iii) HVR-L3 comprising the amino acid sequence of SEQ ID NO:309; or

C) (a) a VH domain comprising (i) HVR-H1 comprising the amino acid sequence of SEQ ID NO:304, (ii) HVR-H2 comprising the amino acid sequence of SEQ ID NO:305, and (iii) HVR-H3 comprising an amino acid sequence selected from SEQ ID NO:306; and (b) a VL domain comprising (i) HVR-L1 comprising the amino acid sequence of SEQ ID NO:315; (ii) HVR-L2 comprising the amino acid sequence of SEQ ID NO:308 and (iii) HVR-L3 comprising the amino acid sequence of SEQ ID NO:309; or

D) (a) a VH domain comprising (i) HVR-H1 comprising the amino acid sequence of SEQ ID NO:316, (ii) HVR-H2 comprising the amino acid sequence of SEQ ID NO:317, and (iii) HVR-H3 comprising an amino acid sequence selected from SEQ ID NO:318; and (b) a VL domain comprising (i) HVR-L1 comprising the amino acid sequence of SEQ ID NO:319; (ii) HVR-L2 comprising the amino acid sequence of SEQ ID NO:320 and (iii) HVR-L3 comprising the amino acid sequence of SEQ ID NO:321; or

E) (a) a VH domain comprising (i) HVR-H1 comprising the amino acid sequence of SEQ ID NO:324, (ii) HVR-H2 comprising the amino acid sequence of SEQ ID NO:325, and (iii) HVR-H3 comprising an amino acid sequence selected from SEQ ID NO:326; and (b) a VL domain comprising (i) HVR-L1 comprising the amino acid sequence of SEQ ID NO:327; (ii) HVR-L2 comprising the amino acid sequence of SEQ ID NO:328 and (iii) HVR-L3 comprising the amino acid sequence of SEQ ID NO:329; or.

F) (a) a VH domain comprising (i) HVR-H1 comprising the amino acid sequence of SEQ ID NO:332, (ii) HVR-H2 comprising the amino acid sequence of SEQ ID NO:333, and (iii) HVR-H3 comprising an amino acid sequence selected from SEQ ID NO:334; and (b) a VL domain comprising (i) HVR-L1 comprising the amino acid sequence of SEQ ID NO:335; (ii) HVR-L2 comprising the amino acid sequence of

SEQ ID NO:336 and (iii) HVR-L3 comprising the amino acid sequence of SEQ ID NO:337; or

5 G) (a) a VH domain comprising (i) HVR-H1 comprising the amino acid sequence of SEQ ID NO:340, (ii) HVR-H2 comprising the amino acid sequence of SEQ ID NO:341, and (iii) HVR-H3 comprising an amino acid sequence selected from SEQ ID NO:342; and (b) a VL domain comprising (i) HVR-L1 comprising the amino acid sequence of SEQ ID NO:343; (ii) HVR-L2 comprising the amino acid sequence of SEQ ID NO:344 and (iii) HVR-L3 comprising the amino acid sequence of SEQ ID NO:345; or

10 H) (a) a VH domain comprising (i) HVR-H1 comprising the amino acid sequence of SEQ ID NO:348, (ii) HVR-H2 comprising the amino acid sequence of SEQ ID NO:349, and (iii) HVR-H3 comprising an amino acid sequence selected from SEQ ID NO:350; and (b) a VL domain comprising (i) HVR-L1 comprising the amino acid sequence of SEQ ID NO:351; (ii) HVR-L2 comprising the amino acid sequence of SEQ ID NO:352 and (iii) HVR-L3 comprising the amino acid sequence of SEQ ID NO:353; or

I) (a) a VH domain comprising (i) HVR-H1 comprising the amino acid sequence of SEQ ID NO:356, (ii) HVR-H2 comprising the amino acid sequence of SEQ ID NO:357, and (iii) HVR-H3 comprising an amino acid sequence selected from SEQ ID NO:358; and (b) a VL domain comprising (i) HVR-L1 comprising the amino acid sequence of SEQ ID NO:359; (ii) HVR-L2 comprising the amino acid sequence of SEQ ID NO:360 and (iii) HVR-L3 comprising the amino acid sequence of SEQ ID NO:361; or

20 J) (a) a VH domain comprising (i) HVR-H1 comprising the amino acid sequence of SEQ ID NO:364, (ii) HVR-H2 comprising the amino acid sequence of SEQ ID NO:365, and (iii) HVR-H3 comprising an amino acid sequence selected from SEQ ID NO:366; and (b) a VL domain comprising (i) HVR-L1 comprising the amino acid sequence of SEQ ID NO:367; (ii) HVR-L2 comprising the amino acid sequence of SEQ ID NO:368 and (iii) HVR-L3 comprising the amino acid sequence of SEQ ID NO:369.

30 In one embodiment, the anti-TIM3 antibody is a full length IgG1 antibody with mutations S228P, L235E and P329G according to the EU index of Kabat numbering. In one embodiment, the anti-TIM3 antibody is any one of the antibodies described in WO

2011/155607, WO 2013/006490, WO 03/063792, WO 2009/097394, and WO 2011/159877. In one embodiment, the anti-TIM3 antibody is F38-2E2.

In one embodiment, the cancer contains a KRAS wildtype. In one embodiment, the cancer contains an activating KRAS mutation.

- 5 In one embodiment, the treatment results in a sustained response in the individual after cessation of the treatment. In one embodiment, at least one of the T cell activating bispecific antigen binding molecule and the PD-1 axis binding antagonist is administered continuously. In one embodiment, at least one of the T cell activating bispecific antigen binding molecule and the PD-1 axis binding antagonist is administered intermittently. In
- 10 one embodiment, the PD-1 axis binding antagonist is administered before the FolR1 TCB. In one embodiment, the PD-1 axis binding antagonist is administered simultaneous with the FolR1 TCB. In one embodiment, the PD-1 axis binding antagonist is administered after the FolR1 TCB. In one embodiment, the cancer is selected from the group consisting of ovarian cancer, lung cancer, breast cancer, renal cancer, colorectal cancer,
- 15 endometrial cancer. In one embodiment, at least one of the T cell activating bispecific antigen binding molecule and the PD-1 axis binding antagonist is administered intravenously, intramuscularly, subcutaneously, topically, orally, transdermally, intraperitoneally, intraorbitally, by implantation, by inhalation, intrathecally, intraventricularly, or intranasally.
- 20 In one embodiment, T cells in the individual have enhanced activation, proliferation and/or effector function relative to prior to the administration of the combination. In one embodiment, T cells in the individual have enhanced activation, proliferation and/or effector function relative to administration of the T cell activating bispecific antigen binding molecule alone. In one embodiment, T cell effector function is secretion of at
- 25 least one of IL-2, IFN- γ and TNF- α . In one embodiment, the individual comprises less than about 15% PD-1^{hi} expressing tumor-infiltrating T cells.

The disclosure provides for a method of enhancing immune function in an individual having a FolR1 positive cancer comprising administering to the individual an effective amount of a combination of a T cell activating bispecific antigen binding molecule

- 30 specific for Folate Receptor 1 (FolR1) and CD3, and a PD-1 axis binding antagonist. In one aspect, the disclosure provides for use of a combination of a T cell activating bispecific antigen binding molecule specific for Folate Receptor 1 (FolR1) and CD3, and

a PD-1 axis binding antagonist in the manufacture of a medicament for combination therapy in enhancing immune function in an individual having a FolR1-positive cancer, wherein the PD-1 axis binding antagonist inhibits the binding of PD-1 to its ligand binding partners, and wherein the PD-1 axis binding antagonist is selected from the group

5 consisting of an anti-PD-1 antibody, an anti-PD-L1 antibody and an anti-PD-L2 antibody. In one embodiment, T cells in the individual have enhanced activation, proliferation and/or effector function relative to prior to the administration of the combination. In one embodiment, T cells in the individual have enhanced activation,

proliferation and/or effector function relative to administration of the T cell activating bispecific antigen binding molecule alone. In one embodiment, T cell effector function is secretion of at least one of IL-2, IFN- γ and TNF- α .

In one embodiment, the individual comprises less than about 15% PD-1^{hi} expressing tumor-infiltrating T cells.

5 The disclosure provides for a method for selecting a patient for treatment with a combination of a T cell activating bispecific antigen binding molecule specific for Folate Receptor 1 (FolR1) and CD3, and a PD-1 axis binding antagonist comprising measuring the level of PD-1 expression, wherein a patient having less than about 15% PD-1^{hi}
10 expressing T cells is selected for treatment with the combination.

In another aspect, the disclosure provides for a method for selecting a patient for treatment with a combination of a T cell activating bispecific antigen binding molecule specific for Folate Receptor 1 (FolR1) and CD3, and a PD-1 axis binding antagonist comprising measuring the level of PD-1 expression, wherein a patient having less than
15 about 15% PD-1^{hi} expressing T cells is selected for treatment with the combination, wherein the PD-1 axis binding antagonist inhibits the binding of PD-1 to its ligand binding partners, and wherein the PD-1 axis binding antagonist is selected from the group consisting of an anti-PD-1 antibody, an anti-PD-L1 antibody and an anti-PD-L2 antibody. The disclosure provides for a kit comprising a T cell activating bispecific antigen binding
20 molecule specific for Folate Receptor 1 (FolR1) and CD3, and a package insert comprising instructions for using the T cell activating bispecific antigen binding molecule with a PD-1 axis binding antagonist to treat or delay progression of cancer in an individual. In one embodiment, the kit further comprises instructions for using the T cell activating bispecific antigen binding molecule with a TIM3 antagonist.

25 The disclosure provides for a kit comprising a T cell activating bispecific antigen binding molecule specific for Folate Receptor 1 (FolR1) and CD3 and a PD-1 axis binding antagonist, and a package insert comprising instructions for using the T cell activating bispecific antigen binding molecule and the PD-1 axis binding antagonist to treat or delay progression of cancer in an individual.

30 In another aspect, the disclosure provides a kit comprising a T cell activating bispecific antigen binding molecule specific for Folate Receptor 1 (FolR1) and CD3 and a PD-1 axis binding antagonist as a combined pharmaceutical formulation for use in treating or delaying progression of cancer in an individual, and a package insert comprising instructions for using the T cell activating bispecific antigen binding molecule and the

PD-1 axis binding antagonist to treat or delay progression of cancer in an individual, wherein the cancer is a FolR1-positive cancer and wherein the PD-1 axis binding antagonist inhibits the binding of PD-1 to its ligand binding partners, wherein the PD-1 axis binding antagonist is selected from the group consisting of an anti-PD-1 antibody, an
5 anti-PD-L1 antibody and an anti-PD-L2 antibody.

In one embodiment, the kit further comprises a TIM3 antagonist. In one embodiment, the PD-1 axis binding antagonist is an anti-PD-1 antibody or an anti-PDL-1 antibody. In one embodiment, the PD-1 axis binding antagonist is an anti-PD-1 immunoadhesin.

The disclosure provides for a pharmaceutical composition comprising a T cell activating
10 bispecific antigen binding molecule specific for Folate Receptor 1 (FolR1) and CD3, a PD-1 axis binding antagonist and a pharmaceutically acceptable carrier. In another aspect, the disclosure provides for a pharmaceutical composition comprising a T cell activating bispecific antigen binding molecule specific for Folate Receptor 1 (FolR1) and CD3, a PD-1 axis binding antagonist and a pharmaceutically acceptable carrier, wherein the PD-1
15 axis binding antagonist inhibits the binding of PD-1 to its ligand binding partners, and wherein the PD-1 axis binding antagonist is selected from the group consisting of an anti-PD-1 antibody, an anti-PD-L1 antibody and an anti-PD-L2 antibody. In one embodiment, the pharmaceutical composition further comprises a TIM3 antagonist.

The disclosure provides for a use of a combination of a T cell activating bispecific antigen
20 binding molecule specific for Folate Receptor 1 (FolR1) and CD3 and a PD-1 axis binding antagonist in the manufacture of a medicament for the treatment of cancer.

In another aspect, the disclosure provides for a use of a T cell activating bispecific antigen binding molecule specific for Folate Receptor 1 (FolR1) and CD3, a PD-1 axis binding antagonist, and a TIM3 antagonist in the manufacture of a medicament for combination
25 therapy in the treatment of cancer, wherein the cancer is a FolR1-positive cancer and wherein the PD-1 axis binding antagonist inhibits the binding of PD-1 to its ligand binding partners, wherein the PD-1 axis binding antagonist is selected from the group consisting of an anti-PD-1 antibody, an anti-PD-L1 antibody and an anti-PD-L2 antibody.

In one embodiment, the medicament is for treatment of ovarian cancer, lung cancer, breast cancer, renal cancer, colorectal cancer, endometrial cancer.

In certain embodiments of the present disclosure, advantageously said T cell activating bispecific antigen binding molecule and/or PD-1 axis binding antagonist is human or
5 humanized.

In some embodiments, the bispecific antibody comprises an Fc domain, at least one Fab fragment comprising the antigen binding site specific for FolR1, and at least one Fab fragment comprising the antigen binding site specific for CD3.

The disclosure provides for a method for treating or delaying progression of a
10 cancer in an individual comprising administering to the individual an effective amount of a T cell activating bispecific antigen binding molecule and a TIM3 antagonist. In some embodiments, the T cell activating bispecific antigen binding molecule comprises an Fc domain, two Fab fragments comprising each an antigen binding site specific for FolR1, and one Fab fragment comprising an antigen binding site specific for CD3.

15 The present disclosure provides the use of a combination of a T cell activating bispecific antigen binding molecule that binds to FolR1 and CD3, and a PD-1 axis binding antagonist in the manufacture of a medicament for the treatment of cancer.

The present disclosure provides the use of a combination of a T cell activating bispecific antigen binding molecule that binds to FolR1 and CD3, a PD-1 axis binding
20 antagonist and a TIM3 antagonist in the manufacture of a medicament for the treatment of cancer.

Embodiments of the present disclosure will now be described by way of example and not limitation with reference to the accompanying figures. However various further aspects and embodiments of the present disclosure will be apparent to those skilled in the
25 art in view of the present disclosure.

“and/or” where used herein is to be taken as specific disclosure of each of the two specified features or components with or without the other. For example “A and/or B” is to be taken as specific disclosure of each of (i) A, (ii) B and (iii) A and B, just as if each is set out individually herein.

30 Unless context dictates otherwise, the descriptions and definitions of the features set out above are not limited to any particular aspect or embodiment of the disclosure and

apply equally to all aspects and embodiments which are described.

BRIEF DESCRIPTION OF THE FIGURES

Figures 1A-I illustrate exemplary configurations of the T cell activating bispecific antigen binding molecules (TCBs) of the invention. All constructs except the kappa-lambda format in (Fig. 1I) have P329G LALA mutations and comprise knob-into-hole Fc fragments with knob-into-hole modifications. (Fig. 1A) Illustration of the “FolR1 TCB 2+1 inverted (common light chain)”. The FolR1 binder is fused at the C-terminus of the Fab heavy chain to the N-terminus of the first subunit of the Fc domain comprising the knob modification. These constructs are not crossed and have three times the same VLCL light chain. (Fig. 1B) Illustration of the “FolR1 TCB 1+1 head-to-tail (common light chain)”. These constructs are not crossed and have two times the same VLCL light chain. (Fig. 1C) Illustration of the “FolR1 TCB 1+1 classical (common light chain)”. These constructs are not crossed and have two times the same VLCL light chain. (Fig. 1D) Illustration of the “FolR1 TCB 2+1 classical (common light chain)”. The CD3 binder is fused at the C-terminus of the Fab heavy chain to the N-terminus of the first subunit of the Fc domain comprising the knob modification. These constructs are not crossed and have three times the same VLCL light chain. (Fig. 1E) Illustration of the “FolR1 TCB 2+1 crossfab classical”. These constructs comprise a Ck-VH chain for the CD3 binder instead of the conventional CH1-VH chain. The CD3 binder is fused at the C-terminus of the Fab heavy chain to the N-terminus of the first subunit of the Fc domain comprising the knob modification. (Fig. 1F) Illustration of the “FolR1 TCB 2+1 crossfab inverted”. These constructs comprise a Ck-VH chain for the CD3 binder instead of the conventional CH1-VH chain. The FolR1 binder is fused at the C-terminus of the Fab heavy chain to the N-terminus of the first subunit of the Fc domain comprising the knob modification. (Fig. 1G) Illustration of the “FolR1 TCB 1+1 crossfab head-to-tail”. These constructs comprise a Ck-VH chain for the CD3 binder instead of the conventional CH1-VH chain. (Fig. 1H) Illustration of the “FolR1 TCB 1+1 crossfab classical”. These constructs comprise a Ck-VH chain for the CD3 binder instead of the conventional CH1-VH chain. Figure 1I illustrates the CD3/FolR1 kappa-lambda antibody format. These constructs comprise a crossed common light chain VLCH1 and one crossed VHCL chain specific for CD3 and one crossed VHCL chain specific for FolR1.

Figures 2A-C depict graphs summarizing Binding of FolR1 IgG binders to HeLa cells. Binding of newly generated FolR1 binders to FolR1 expressed on HeLa cells were determined by flow cytometry. Bound antibodies were detected with a fluorescently labeled anti-human secondary antibody.

5 **Figures 3A-B** depict graphs summarizing specificity of FolR1 binders for FolR1. Binding of FolR1 IgGs to HEK cells transiently transfected with either FolR1 or FolR2 was analyzed by flow cytometry to identify clones which bind specifically to FolR1 and not to FolR2. The antibodies were detected with a fluorescently labeled anti-human secondary antibody.

10 **Figures 4A-B** depict graphs summarizing cross-reactivity of FolR1 binders to cyFolR1. Cross-reactivity of the FolR1 antibodies to cyno FolR1 was addressed on HEK cells transiently transfected with cyFolR1 by flow cytometry. The antibodies were detected with a fluorescently labeled anti-human secondary antibody.

Figure 5 depicts a graph illustrating internalization of FolR1 TCBs after binding.

15 Internalization of the four FolR1 TCBs after binding to FolR1 was tested on HeLa cells. Remaining FolR1 TCBs on the surface were detected with a fluorescently labeled anti-human secondary antibody after indicated time points of incubation at 37°C. Percentage of internalization was calculated.

Figures 6A-E depict graphs summarizing binding of FolR1 IgGs to cells with different FolR1 expression levels. Binding of 9D11, 16D5 and Mov19 IgG to tumor cells with different FolR1 expression levels was analyzed by flow cytometry. DP47 IgG was included as isotype control and MKN-45 were included as FolR1 negative cell line. The antibodies were detected with a fluorescently labeled anti-human secondary antibody.

20 **Figures 7A-L** depict graphs summarizing T cell mediated killing of HT-29 and SKOV3 cells. FolR1 TCBs were used to test T cell mediated killing of HT-29 and SKOV3 tumor cells and upregulation of activation marker on T cells upon killing. (Figs. 7A-D) T cell mediated killing of HT-29 and SKOV3 cells in the presence of 9D11 FolR1 TCB and 16D5 FolR1 TCB was measured by LDH release after 24 h and 48 h. DP47 TCB was included as negative control. After 48 h incubation upregulation of the activation marker
25 CD25 and CD69 on CD8 T cells and CD4 T cells upon killing of SKOV3 (Figs. 7E-H) or HT-29 (Fig. 7I-L) tumor cells was assessed by flow cytometry.

Figure 8 depicts a graph showing absence of anti-FolR1 binding to erythrocytes. Erythrocytes were gated as CD235a positive population and binding of 9D11 IgG, 16D5

IgG, Mov19 IgG and DP47 IgG to this population was determined by flow cytometry. The antibodies were detected with a fluorescently labeled anti-human secondary antibody. **Figures 9A-D** depict graphs summarizing activation marker upregulation in whole blood. CD25 and CD69 activation marker upregulation of CD4 T cells and CD8 T cells 24 h after addition of 9D11 FolR1 TCB, 16D5 FolR1 TCB, Mov19 FolR1 TCB and DP47 TCB was analyzed by flow cytometry.

Figures 10A-C depict T-cell killing induced by 36F2 TCB, 16D5 TCB, 16D5 TCB classical, 16D5 TCB 1+1 and 16D5 TCB HT of Hela (high FolR1) (Fig. 24A), Skov-3 (medium FolR1) (Fig. 24B) and HT-29 (low FolR1) (Fig. 24C) human tumor cells (E:T = 10:1, effectors human PBMCs, incubation time 24 h). DP47 TCB was included as non-binding control.

Figures 11A-B show expression of inhibitory receptors on tumor-infiltrating T cells. CD8⁺ and CD4⁺ T cells in tumor samples were characterized by flow cytometry for their expression of inhibitory receptors.

Figures 12A-O show activation of CD8⁺ T cells in tumor digests and malignant effusions upon exposure to FolR1-TCB. Tumor digests or malignant effusions were cultured for 24h in the presence or absence of FolR1-TCB or the control TCB DP-47. The expression of activation markers or markers of T cell function on CD8⁺ T cells was determined by flow cytometry (Fig. 12A-M). Fig. 12J-K show representative FACS plots showing FolR1-TCB-induced T cell activation in a high responding (BS-269) or a low responding patient (BS-212). Fig. 12L depicts FACS plots showing FolR1-TCB-induced activation marker expression in T cells from a representative patient. The graphs in Fig. 12M depict the increase in marker expression after FolR1-TCB treatment with mean and standard deviations. As comparison, PBMC from healthy donors were co-cultured with the Skov3 tumor cell line and stimulated with FolR1-TCB. Fig. 12N depicts IFN- γ , IL-2, TNF and perforin in the cell culture supernatants as determined by Cytometric Bead Array or ELISA and normalized to the amount of 1×10^5 CD3⁺ T-cells (IFN- γ , TNF, IL-2) or CD3⁺ CD8⁺ T-cells (perforin) in the culture. Fig. 12O shows that FolR1-TCB-induced tumor cell killing varies largely in tumor digests and malignant effusions. FolR1 positive and negative tumor digests, malignant effusions or PBMCs from healthy donors were co-cultured with exogenously added fluorescently labeled FolR1⁺ Skov3 cells at an E:T ratio of 1:1 for 24 h in the presence or absence of FolR1-TCB. The FolR1-TCB-induced specific killing of the Skov3 cells was determined by flow cytometry by measuring

activated caspase 3 and the live/dead marker LIVE/DEAD[®]-near-IR. FolR1-TCB-mediated killing was calculated as follows: % specific killing = $100 - [(\% \text{ of Skov3 live cells in FolR1-TCB treated sample} / \% \text{ of Skov3 live cells in untreated sample}) \times 100]$. FACS plots show FolR1-TCB-induced killing in a representative patient. The p-values were calculated using the unpaired Mann-Whitney test.

Figures 13A-C show that FolR1-TCB-induced T cell activation shows no correlation with E:T ratio (Fig. 13A) or the amount of FolR1⁺ tumor cells (Fig. 13B). Tumor digests or malignant effusions were cultured for 24h in the presence or absence of FolR1-TCB. The FolR1-TCB induced expression of CD25 was correlated to E:T ratio or the amount of target cells. MFI: mean fluorescence intensity.

Figures 14A-L show FolR1-TCB induced T cell activation inversely correlates with expression of PD-1 and Tim-3. Tumor digests or malignant effusions were cultured for 24h in the presence or absence of FolR1-TCB. The expression of activation markers or markers of T cell function on CD8⁺ T cells was determined by flow cytometry. The FolR1-TCB induced expression of CD25 (Fig. 4A-C), CD137 (Fig. 14D-F), ICOS (Fig. 14G-I) and granzyme B (Fig. 14J-L) was correlated to baseline single- or co-expression of the inhibitory receptors PD-1 and Tim-3.

Figures 15A-C show FolR1-TCB induced IL-2 secretion inversely correlates with co-expression of PD-1 and Tim-3. Tumor digests or malignant effusions were cultured for 24h in the presence or absence of FolR1 TCB. IL-2 in the cell culture supernatants was determined by ELISA and normalized to the amount of T cells. The FolR1 TCB induced IL-2 secretion was correlated to baseline single- or co-expression of the inhibitory receptors PD-1 and Tim-3.

Figures 16A-F show FolR1-TCB induced tumor cell killing inversely correlates with co-expression of PD-1 and Tim-3. Tumor digests or malignant effusions were co-cultured with exogenously added fluorescence labelled Skov3 cells at a T cell to target cell ratio of 1:1 for 24h in the presence or absence of FolR1 TCB. The FolR1-TCB specific killing of the Skov3 cells was determined by flow cytometry by measuring activated caspase 3 and the live/dead marker Live/Dead-near-IR. The specific killing was correlated to baseline single or co-expression of the inhibitory receptors PD-1, Tim-3 and CTLA-4.

Figures 17A-H show activation of tumor-infiltrating CD8⁺ T cells upon exposure to catumaxomab. Tumor digests or malignant effusions were cultured for 24h in the presence or absence of catumaxomab. (Fig. 17A-D) The expression of activation markers

or markers of T cell function on CD8⁺ T cells was determined by flow cytometry. (Fig. 17E-H) Graphs showing the baseline expression of inhibitory receptors.

Figures 18A-R show Catumaxomab-induced T cell activation inversely correlates with co-expression of inhibitory receptors. Tumor digests or malignant effusions were cultured for 24h in the presence or absence of catumaxomab. T cell activation and effector functions were correlated to the expression of PD-1 (Fig. 18A-F), Tim-3 (Fig. 18G-L) or of the combination of PD-1 and Tim-3 (Fig. 18M-R).

Figures 19A-H show expression of inhibitory receptors on tumor-infiltrating T cells in Non-small cell lung cancer patients. CD8⁺ and CD4⁺ T cells in tumor samples were characterized by flow cytometry for their expression of inhibitory receptors (Fig. 19A-F). Fig. 19G shows the gating strategy for one representative donor. Fig. 19H shows results of analysis and heat mapping of indicated cell subsets based on the percentage of expression, with the use of an Excel conditional formatting program.

Figures 20A-E show T cell activation and effector functions upon polyclonal stimulation by CD3/CD28 antibodies. Expression of CD25 and Granzyme B (Fig. 20A-B) as well as IL-2, IFN- γ and TNF- α (Fig. 20C-E) as markers for T cell activation and effector function, respectively, was analyzed in T cells from digested tumor samples after stimulation of whole tumor digests with agonistic CD3 and CD28 antibodies.

Figures 21A-N show expression of inhibitory receptors and T cell dysfunction.

Expression of CD25 and Granzyme B (Fig. 21A-B) as well as IL-2, IFN- γ and TNF- α (Fig. 21C-E) upon polyclonal stimulation by an anti-CD3/ anti-CD28 antibodies correlates with the cumulative expression of inhibitory receptors indicated by the iR Score. Fig. 21F shows an exemplary calculation of iR scores. The percentage of expression of PD-1, Tim-3, CTLA-4, LAG-3 and BTLA was analyzed in all NSCLC samples and the median as well as interquartile ranges were determined. For the calculation of the iR score each patient received points for the expression of each of the determined inhibitory receptors based on the quartile within which the expression coincided. A maximum of 15 points could be reached; the calculated score of each sample was normalized to this maximum amount of points. Fig. 21G-K show expression of inhibitory receptors increases with tumor stage. Expression of inhibitory receptors on CD8⁺ tumor infiltrating T-cells was correlated to the TNM stage. Fig. 21L-N show increased cumulative expression of inhibitory receptors with tumor progression. The cumulative expression of the inhibitory receptors PD-1, Tim-3, CTLA-4, LAG-3 and

BTLA, as represented by the iR score, was correlated to the nodal status and the TNM stage.

Figures 22A-I show expression of PD-1 and Tim-3 correlates with T cell dysfunction. Expression of CD25 and Granzyme B (Fig. 22A-C) as well as IL-2, IFN- γ and TNF- α (Fig. 22D-F) upon polyclonal stimulation by CD3/CD28 correlates with the expression of PD-1 (Fig. 22A-C), Tim-3 (Fig. 22D-F) or PD-1/Tim-3 (Fig. 22G-I) on tumor-infiltrating T cells.

Figures 23A-E show that the effect of PD-1 or combined PD-1/Tim-3 blockade varies between patients. Digests were stimulated by agonistic anti-CD3/anti-CD28 antibodies with the addition of blocking antibodies to PD-1 alone or in combination with Tim-3. Secretion of IFN- γ , TNF- α and IL-2 was determined by ELISA and normalized to 1×10^6 T cells. Fig. 23A-C show T cells from a patient where T cell function can be rescued by addition of blocking Abs (BS-268) and T cells from a patient with no response to PD-1 or PD-1/Tim-3 blockade. The difference in expression ($[\% \text{ expression Ab treated}] - [\% \text{ expression untreated}]$) is shown. Fig. 23D shows respective flow cytometry plots with PD-1^{hi} and PD-1^{int} subsets. Fig. 23E shows a summary of IL-2, TNF- α and IFN- γ secretion by T cells from six patients, as determined by ELISA and normalized to 1×10^6 CD3⁺ T cells.

Figures 24A-F show that the effect of PD-1 or combined PD-1/Tim-3 blockade differs in PD-1^{hi} and PD-1^{int} subsets. Correlation of the increase in cytokine production by PD-1 or combined PD-1/Tim-3 blockade with PD-1^{hi} and PD-1^{int} subsets are indicated by PD-1^{hi}/PD-1^{int} ratio.

Figures 25A-I show activation of CD4⁺ T cells in tumor digests and malignant effusions upon exposure to FolR1-TCB. Tumor digests or malignant effusions were cultured for 24h in the presence or absence of FolR1-TCB or the control TCB DP-47. The expression of activation markers or markers of T cell function on CD8⁺ T cells was determined by flow cytometry.

Figures 26A-C show FolR1-TCB induced T cell activation is independent of CTLA-4, Lag-3 and BTLA expression. Tumor digests or malignant effusions were cultured for 24h in the presence or absence of FolR1-TCB. The expression of CD25 on CD8⁺ T cells was determined by flow cytometry. The FolR1-TCB induced expression of CD25 was correlated to baseline expression of CTLA-4, Lag-3 and BTLA.

Figures 27A-C show FolR1-TCB induces cytokine secretion only in patients with a low percentage of PD-1^{hi} expressing CD8⁺ T cells. Tumor digests or malignant effusions were cultured for 24h in the presence or absence of FolR1-TCB. IFN- γ , TNF and IL-2 in the cell culture supernatants was determined and normalized to the amount of 1×10^5 T cells in the culture. The FolR1-TCB induced cytokine secretion was correlated to baseline PD-1^{hi} expression.

Figures 28A-F show that treatment with a PD-1 blocking antibody fails to induce cytokine secretion in tumor digests or malignant effusions from patients with lung and ovarian cancer with a low percentage of PD-1^{hi} expressing cells. Tumor digests or malignant effusions were cultured for 24h with FolR1-TCB in the presence or absence of PD-1 blocking antibody (Fig. 28A-C) or the combination of PD-1 and Tim-3 blocking antibodies (Fig. 28D-F). IFN- γ , TNF and IL-2 in the cell culture supernatants was determined and normalized to the amount of 1×10^5 T cells in the culture. The cytokine secretion induced by the blocking antibodies compared to FolR1-TCB treatment alone was correlated to baseline PD-1^{hi} expression.

Figures 29A-B show results from a FACS based internalization assay. The data show that the Fab fragment (<TIM-3> Fab) of anti-TIM3 antibody Tim3_0022 (abbreviated as <TIM-3> Ab(022)) internalized into rec CHOK1 cells expressing huTIM-3 after incubation at 37°C with similar kinetic as the antibody in the full IgG format.

Figures 30A-B show binding of anti-TIM3 antibodies to RPMI-8226 cells (antibody designation clone 0016 refers to antibody Tim3_0016, clone 0016 refers to antibody Tim3_0016 variant (antibody Tim3_0018), clone 0022 refers to antibody Tim3_00122, etc.). Fig. 30B shows binding of anti-TIM3 antibodies to Pfeiffer cells (antibody designation clone 0016 refers to antibody Tim3_0016, clone 0016 refers to antibody Tim3_0016 variant (antibody Tim3_0018), clone 0022 refers to antibody Tim3_00122, etc.).

Figure 31 shows expression level of TIM-3 on different patient AML cell samples by FACS using anti-TIM-3 mAbs.

Figure 32 shows a heat map of expression of inhibitory receptors on NSCLC associated TILs. Co-expression of inhibitory receptors on tumor-infiltrating CD8⁺ T-cells positive for the indicated immune checkpoint is shown as a heat map displaying the percentage of expression for the additional receptors.

Figure 33 shows a radar plot of expression of inhibitory receptors on NSCLC associated TILs. Co-expression of inhibitory receptors on tumor-infiltrating CD8⁺ T-cells positive for the indicated immune checkpoint is shown as a radar plot indicating the mean expression and standard deviation of the four other receptors.

- 5 **Figures 34A-D** show the percentage of PD-1^{hi} or PD-1^{int} CD8⁺ T cells expressing additional immune checkpoints. Each dot represents one patient samples. The p values were calculated using the Wilcoxon rank sum test.

Figures 35A-F show intratumoral T cell inhibitory receptor expression and T cell function. Fig. 35A shows the gating strategy for identification of PD-1^{hi}, PD-1^{int}, and PD-1^{neg} CD8⁺ subsets of T-cells from two representative patients. Fig. 35B shows distribution of indicated T cell subsets in the tumor samples analyzed. Fig. 35C shows that T-cell functions induced by anti-CD3/-CD28 stimulation depend on the PD-1 expression level of CD8⁺ T-cells. Tumor digests and malignant effusions were cultured for 24 h in the presence or absence of agonistic anti-CD3/-CD28 antibodies. The increase in the expression of CD25 on CD8⁺ T-cells (Fig. 35C) and the increase in the effector cytokines IFN- γ , IL-2, and TNF (Fig. 35D) were determined in PD-1^{hi} scarce and abundant tumors. p-values were calculated using the unpaired Mann-Whitney test. Tumor samples were divided according to the percentage of PD-1^{hi} expressing CD8⁺ cells in two groups with PD-1^{hi} scarce and abundant expression, respectively (Fig. 35E). The expression level of the inhibitory receptors PD-1, Tim-3, CTLA-4, Lag-3, and BTLA was determined by flow cytometry on CD8⁺ T-cells from tumor digests or malignant effusions (Fig. 35F).

Figures 36A-E show patterns of inhibitory receptor expression and percentage of scarce and abundant CD8⁺ T-cells. Fig. 36A-D show co-expression of Tim-3, CTLA-4, Lag-3, and BTLA on PD-1^{hi}, PD-1^{int}, and PD-1^{neg} CD8⁺ T-cells. The p-values were calculated using one-way ANOVA with Bonferroni post-hoc-test. Fig. 36E: FolR1⁺ tumor samples were divided according to the percentage of PD-1^{hi} expressing CD8⁺ cells in two groups with PD-1^{hi} scarce and abundant expression, respectively.

Figures 37A-H show that FolR1-TCB-induced T-cell functions depend on the PD-1 expression level of CD8⁺ T-cells. FolR1⁺ tumor digests and malignant effusions were cultured for 24 h in the presence or absence of FolR1-TCB. The increase in the expression of activation markers on CD8⁺ T-cells (Fig. 37A-C) and the increase in the effector cytokines IFN- γ , IL-2, TNF, and perforin (Fig. 37D-G) was determined in PD-1^{hi}

scarce and abundant tumors. Fig. 37H shows target cell killing. Both FolR1 positive and negative tumor samples were adjusted by addition of the FolR1⁺ Skov3 cell line to an E:T ratio of 1:1 and killing was compared in PD-1^{hi} scarce and abundant tumors. p-values were calculated using the unpaired Mann-Whitney test.

- 5 **Figures 38A-E** show that PD-1 blockade increases cytokine production but not their cytolytic function in T-cells from PD-1^{hi} scarce tumors only. Fig. 38A-D: FolR1⁺ tumor digests or malignant effusions were cultured for 24 h with FolR1-TCB in the presence or absence of a PD-1 blocking antibody. IFN- γ , IL-2, TNF, and perforin in the cell culture supernatants were determined by Cytometric Bead Array or ELISA and normalized to the
- 10 amount of 1×10^5 CD3⁺ T-cells (IFN- γ , IL-2, TNF, Fig. 38A-C) or CD3⁺ CD8⁺ T-cells (perforin, Fig. 38D). The increase in cytokine secretion upon combined FolR1-TCB and anti-PD-1 treatment compared with FolR1-TCB alone was determined in PD-1^{hi} scarce and abundant tumors. Fig. 38E: Tumor digests or malignant effusions were co-cultured with exogenously added fluorescently labeled Skov3 cells at an E:T ratio of 1:1 for 24 h
- 15 in the presence or absence of a PD-1 blocking antibody and FolR1-TCB. The increase in specific killing by the anti-PD-1 antibody was compared in PD-1^{hi} scarce and abundant tumors. p-values were calculated using the unpaired Mann-Whitney test.

Figure 39 shows detailed patient characteristics.

- Figures 40A-C** show activation of CD8⁺ T-cells upon exposure to increasing
- 20 concentrations of FolR1-TCB. PBMCs were co-cultured with Skov3 cells for 24 h in the presence or absence of FolR1-TCB or the unspecific control DP-47-TCB. Fig. 40A shows the expression of FolR1 on Skov3. Shaded histogram: isotype control; open histogram: anti-FolR1-antibody. Fig. 40B: The expression of the activation markers CD25, CD137, and ICOS on CD8⁺ T-cells was determined by flow cytometry. Fig. 40C: IFN- γ , IL-2, and TNF in the cell culture supernatants were determined by ELISA and normalized to
- 25 the amount of 1×10^5 CD3⁺ T-cells.

DETAILED DESCRIPTION OF EMBODIMENTS OF THE INVENTION

I. DEFINITIONS

- 30 An “acceptor human framework” for the purposes herein is a framework comprising the amino acid sequence of a light chain variable domain (VL) framework or a heavy chain variable domain (VH) framework derived from a human immunoglobulin framework or a human consensus framework, as defined below. An acceptor human

framework “derived from” a human immunoglobulin framework or a human consensus framework may comprise the same amino acid sequence thereof, or it may contain amino acid sequence changes. In some embodiments, the number of amino acid changes are 10 or less, 9 or less, 8 or less, 7 or less, 6 or less, 5 or less, 4 or less, 3 or less, or 2 or less. In
5 some embodiments, the VL acceptor human framework is identical in sequence to the VL human immunoglobulin framework sequence or human consensus framework sequence.

“Affinity” refers to the strength of the sum total of noncovalent interactions between a single binding site of a molecule (e.g., an antibody) and its binding partner (e.g., an antigen). Unless indicated otherwise, as used herein, “binding affinity” refers to
10 intrinsic binding affinity which reflects a 1:1 interaction between members of a binding pair (e.g., antibody and antigen). The affinity of a molecule X for its partner Y can generally be represented by the dissociation constant (K_d). Affinity can be measured by common methods known in the art, including those described herein. Specific illustrative and exemplary embodiments for measuring binding affinity are described in the
15 following.

An “affinity matured” antibody refers to an antibody with one or more alterations in one or more hypervariable regions (HVRs), compared to a parent antibody which does not possess such alterations, such alterations resulting in an improvement in the affinity of the antibody for antigen.

20 The term “A bispecific antibody that specifically binds Folate Receptor1 (FolR1) and CD3,” “T cell activating bispecific antigen binding molecule specific for FolR1 and CD3” and “FolR1 TCB” are used interchangeably herein and refer to a bispecific antibody that is capable of binding FolR1 and CD3 with sufficient affinity such that the antibody is useful as a diagnostic and/or therapeutic agent in targeting CD3⁺ T cells to
25 FolR2⁺ target cells.

The terms “anti-TIM3 antibody” and “TIM3 antibody” are used synonymously herein to refer to an antibody that specifically binds to TIM3. An anti-TIM3 antibody described herein refers to an antibody that is capable of binding TIM3, especially a TIM3 polypeptide expressed on a cell surface, with sufficient affinity such that the antibody is
30 useful as a diagnostic and/or therapeutic agent. In one embodiment, the extent of binding of an antibody that specifically binds TIM3 to an unrelated non-TIM3 protein is less than about 10% of the binding of the antibody to TIM3 as measured, e.g., by a radioimmunoassay (RIA) or flow cytometry (FACS). In certain embodiments, an

antibody that specifically binds TIM3 has a dissociation constant (K_d) of $\leq 1\mu\text{M}$, $\leq 100\text{ nM}$, $\leq 10\text{ nM}$, $\leq 1\text{ nM}$, $\leq 0.1\text{ nM}$, $\leq 0.01\text{ nM}$, or $\leq 0.001\text{ nM}$ (e.g., 10^{-8} M or less, e.g. from 10^{-8} M to 10^{-13} M , e.g., from 10^{-9} M to 10^{-13} M). In certain embodiments, an antibody that specifically binds TIM3 binds to an epitope of TIM3 that is conserved among DR5 from different species. Preferably said antibody binds to human and cynomolgous monkey TIM3. The term "An antibody that specifically binds TIM3" also encompasses bispecific antibodies that are capable of binding TIM3 and a second antigen.

The term "antibody" herein is used in the broadest sense and encompasses various antibody structures, including but not limited to monoclonal antibodies, polyclonal antibodies, multispecific antibodies (e.g., bispecific antibodies), and antibody fragments so long as they exhibit the desired antigen-binding activity.

An "antibody fragment" refers to a molecule other than an intact antibody that comprises a portion of an intact antibody that binds the antigen to which the intact antibody binds. Examples of antibody fragments include but are not limited to Fv, Fab, Fab', Fab'-SH, F(ab')₂; diabodies, cross-Fab fragments; linear antibodies; single-chain antibody molecules (e.g. scFv); and multispecific antibodies formed from antibody fragments. scFv antibodies are, e.g. described in Houston, J.S., Methods in Enzymol. 203 (1991) 46-96). In addition, antibody fragments comprise single chain polypeptides having the characteristics of a VH domain, namely being able to assemble together with a VL domain, or of a VL domain, namely being able to assemble together with a VH domain to a functional antigen binding site and thereby providing the antigen binding property of full length antibodies.

As used herein, "Fab fragment" refers to an antibody fragment comprising a light chain fragment comprising a VL domain and a constant domain of a light chain (CL), and a VH domain and a first constant domain (CH1) of a heavy chain. In one embodiment the bispecific antibodies of the invention comprise at least one Fab fragment, wherein either the variable regions or the constant regions of the heavy and light chain are exchanged. Due to the exchange of either the variable regions or the constant regions, said Fab fragment is also referred to as "cross-Fab fragment" or "xFab fragment" or "crossover Fab fragment". Two different chain compositions of a crossover Fab molecule are possible and comprised in the bispecific antibodies of the invention: On the one hand, the variable regions of the Fab heavy and light chain are exchanged, i.e. the crossover Fab molecule comprises a peptide chain composed of the light chain variable region (VL) and

the heavy chain constant region (CH1), and a peptide chain composed of the heavy chain variable region (VH) and the light chain constant region (CL). This crossover Fab molecule is also referred to as CrossFab_(VLVH). On the other hand, when the constant regions of the Fab heavy and light chain are exchanged, the crossover Fab molecule comprises a peptide chain composed of the heavy chain variable region (VH) and the light chain constant region (CL), and a peptide chain composed of the light chain variable region (VL) and the heavy chain constant region (CH1). This crossover Fab molecule is also referred to as CrossFab_(CLCH1). Bispecific antibody formats comprising crossover Fab fragments have been described, for example, in WO 2009/080252, WO 2009/080253, WO 2009/080251, WO 2009/080254, WO 2010/136172, WO 2010/145792 and WO 2013/026831.

A “single chain Fab fragment” or “scFab” is a polypeptide consisting of an antibody heavy chain variable domain (VH), an antibody constant domain 1 (CH1), an antibody light chain variable domain (VL), an antibody light chain constant domain (CL) and a linker, wherein said antibody domains and said linker have one of the following orders in N-terminal to C-terminal direction:

a) VH-CH1-linker-VL-CL, b) VL-CL-linker-VH-CH1, c) VH-CL-linker-VL-CH1 or d) VL-CH1-linker-VH-CL; and wherein said linker is a polypeptide of at least 30 amino acids, preferably between 32 and 50 amino acids. Said single chain Fab fragments a) VH-CH1-linker-VL-CL, b) VL-CL-linker-VH-CH1, c) VH-CL-linker-VL-CH1 and d) VL-CH1-linker-VH-CL, are stabilized via the natural disulfide bond between the CL domain and the CH1 domain. In addition, these single chain Fab molecules might be further stabilized by generation of interchain disulfide bonds via insertion of cysteine residues (e.g. position 44 in the variable heavy chain and position 100 in the variable light chain according to Kabat numbering). The term “N-terminus denotes the last amino acid of the N-terminus. The term “C-terminus denotes the last amino acid of the C-terminus. By “fused” or “connected” is meant that the components (e.g. a Fab molecule and an Fc domain subunit) are linked by peptide bonds, either directly or via one or more peptide linkers.

The term “linker” as used herein refers to a peptide linker and is preferably a peptide with an amino acid sequence with a length of at least 5 amino acids, preferably with a length of 5 to 100, more preferably of 10 to 50 amino acids. In one embodiment said peptide linker is $(G_xS)_n$ (SEQ ID NOS 384 and 385) or $(G_xS)_nG_m$ (SEQ ID NOS 429

and 430) with G = glycine, S = serine, and ($x = 3, n = 3, 4, 5$ or 6 , and $m = 0, 1, 2$ or 3) or ($x = 4, n = 2, 3, 4$ or 5 and $m = 0, 1, 2$ or 3), preferably $x = 4$ and $n = 2$ or 3 , more preferably with $x = 4, n = 2$. In one embodiment said peptide linker is $(G_4S)_2$ (SEQ ID NO: 386).

The term "immunoglobulin molecule" refers to a protein having the structure of a naturally occurring antibody. For example, immunoglobulins of the IgG class are heterotetrameric glycoproteins of about 150,000 daltons, composed of two light chains and two heavy chains that are disulfide-bonded. From N- to C-terminus, each heavy chain has a variable region (VH), also called a variable heavy domain or a heavy chain variable domain, followed by three constant domains (CH1, CH2, and CH3), also called a heavy chain constant region. Similarly, from N- to C-terminus, each light chain has a variable region (VL), also called a variable light domain or a light chain variable domain, followed by a constant light (CL) domain, also called a light chain constant region. The heavy chain of an immunoglobulin may be assigned to one of five types, called α (IgA), δ (IgD), ϵ (IgE), γ (IgG), or μ (IgM), some of which may be further divided into subtypes, e.g. γ_1 (IgG₁), γ_2 (IgG₂), γ_3 (IgG₃), γ_4 (IgG₄), α_1 (IgA₁) and α_2 (IgA₂). The light chain of an immunoglobulin may be assigned to one of two types, called kappa (κ) and lambda (λ), based on the amino acid sequence of its constant domain. An immunoglobulin essentially consists of two Fab molecules and an Fc domain, linked via the immunoglobulin hinge region.

An "antibody that binds to the same epitope" as a reference antibody refers to an antibody that blocks binding of the reference antibody to its antigen in a competition assay by 50% or more, and conversely, the reference antibody blocks binding of the antibody to its antigen in a competition assay by 50% or more. An exemplary competition assay is provided herein.

The term "antigen binding domain" refers to the part of an antigen binding molecule that comprises the area which specifically binds to and is complementary to part or all of an antigen. Where an antigen is large, an antigen binding molecule may only bind to a particular part of the antigen, which part is termed an epitope. An antigen binding domain may be provided by, for example, one or more antibody variable domains (also called antibody variable regions). Preferably, an antigen binding domain comprises an antibody light chain variable region (VL) and an antibody heavy chain variable region (VH).

The term "chimeric" antibody refers to an antibody in which a portion of the heavy and/or light chain is derived from a particular source or species, while the remainder of the heavy and/or light chain is derived from a different source or species, usually prepared by recombinant DNA techniques. Chimeric antibodies comprising a rabbit variable region and a human constant region are preferred. Other preferred forms of "chimeric antibodies" encompassed by the present invention are those in which the constant region has been modified or changed from that of the original antibody to generate the properties according to the invention, especially in regard to C1q binding and/or Fc receptor (FcR) binding. Such chimeric antibodies are also referred to as "class-switched antibodies". Chimeric antibodies are the product of expressed immunoglobulin genes comprising DNA segments encoding immunoglobulin variable regions and DNA segments encoding immunoglobulin constant regions. Methods for producing chimeric antibodies involve conventional recombinant DNA and gene transfection techniques are well known in the art. See e.g. Morrison, S.L., et al., Proc. Natl. Acad. Sci. USA 81 (1984) 6851-6855; US Patent Nos. 5,202,238 and 5,204,244.

The term "cytotoxic agent" as used herein refers to a substance that inhibits or prevents a cellular function and/or causes cell death or destruction. Cytotoxic agents include, but are not limited to, radioactive isotopes (e.g., At²¹¹, I¹³¹, I¹²⁵, Y⁹⁰, Re¹⁸⁶, Re¹⁸⁸, Sm¹⁵³, Bi²¹², P³², Pb²¹² and radioactive isotopes of Lu); chemotherapeutic agents or drugs (e.g., methotrexate, adriamycin, vinca alkaloids (vincristine, vinblastine, etoposide), doxorubicin, melphalan, mitomycin C, chlorambucil, daunorubicin or other intercalating agents); growth inhibitory agents; enzymes and fragments thereof such as nucleolytic enzymes; antibiotics; toxins such as small molecule toxins or enzymatically active toxins of bacterial, fungal, plant or animal origin, including fragments and/or variants thereof; and the various antitumor or anticancer agents disclosed below.

"Effector functions" refer to those biological activities attributable to the Fc region of an antibody, which vary with the antibody isotype. Examples of antibody effector functions include: C1q binding and complement dependent cytotoxicity (CDC); Fc receptor binding; antibody-dependent cell-mediated cytotoxicity (ADCC); antibody-dependent cellular phagocytosis (ADCP), cytokine secretion, immune complex-mediated antigen uptake by antigen presenting cells; down regulation of cell surface receptors (e.g. B cell receptor); and B cell activation.

As used herein, the terms “engineer, engineered, engineering”, are considered to include any manipulation of the peptide backbone or the post-translational modifications of a naturally occurring or recombinant polypeptide or fragment thereof. Engineering includes modifications of the amino acid sequence, of the glycosylation pattern, or of the side chain group of individual amino acids, as well as combinations of these approaches.

The term “amino acid mutation” as used herein is meant to encompass amino acid substitutions, deletions, insertions, and modifications. Any combination of substitution, deletion, insertion, and modification can be made to arrive at the final construct, provided that the final construct possesses the desired characteristics, e.g., reduced binding to an Fc receptor, or increased association with another peptide. Amino acid sequence deletions and insertions include amino- and/or carboxy-terminal deletions and insertions of amino acids. Particular amino acid mutations are amino acid substitutions. For the purpose of altering e.g. the binding characteristics of an Fc region, non-conservative amino acid substitutions, i.e. replacing one amino acid with another amino acid having different structural and/or chemical properties, are particularly preferred. Amino acid substitutions include replacement by non-naturally occurring amino acids or by naturally occurring amino acid derivatives of the twenty standard amino acids (e.g. 4-hydroxyproline, 3-methylhistidine, ornithine, homoserine, 5-hydroxylysine). Amino acid mutations can be generated using genetic or chemical methods well known in the art. Genetic methods may include site-directed mutagenesis, PCR, gene synthesis and the like. It is contemplated that methods of altering the side chain group of an amino acid by methods other than genetic engineering, such as chemical modification, may also be useful. Various designations may be used herein to indicate the same amino acid mutation. For example, a substitution from proline at position 329 of the Fc domain to glycine can be indicated as 329G, G329, G₃₂₉, P329G, or Pro329Gly.

An “effective amount” of an agent, e.g., a pharmaceutical formulation, refers to an amount effective, at dosages and for periods of time necessary, to achieve the desired therapeutic or prophylactic result.

The term “Fc domain” or “Fc region” herein is used to define a C-terminal region of an immunoglobulin heavy chain that contains at least a portion of the constant region. The term includes native sequence Fc regions and variant Fc regions. Although the boundaries of the Fc region of an IgG heavy chain might vary slightly, the human IgG heavy chain Fc region is usually defined to extend from Cys226, or from Pro230, to the

carboxyl-terminus of the heavy chain. However, the C-terminal lysine (Lys447) of the Fc region may or may not be present. Unless otherwise specified herein, numbering of amino acid residues in the Fc region or constant region is according to the EU numbering system, also called the EU index, as described in Kabat et al., Sequences of Proteins of Immunological Interest, 5th Ed. Public Health Service, National Institutes of Health, Bethesda, MD, 1991. A "subunit" of an Fc domain as used herein refers to one of the two polypeptides forming the dimeric Fc domain, i.e. a polypeptide comprising C-terminal constant regions of an immunoglobulin heavy chain, capable of stable self-association. For example, a subunit of an IgG Fc domain comprises an IgG CH2 and an IgG CH3 constant domain.

A "modification promoting the association of the first and the second subunit of the Fc domain" is a manipulation of the peptide backbone or the post-translational modifications of an Fc domain subunit that reduces or prevents the association of a polypeptide comprising the Fc domain subunit with an identical polypeptide to form a homodimer. A modification promoting association as used herein particularly includes separate modifications made to each of the two Fc domain subunits desired to associate (i.e. the first and the second subunit of the Fc domain), wherein the modifications are complementary to each other so as to promote association of the two Fc domain subunits. For example, a modification promoting association may alter the structure or charge of one or both of the Fc domain subunits so as to make their association sterically or electrostatically favorable, respectively. Thus, (hetero)dimerization occurs between a polypeptide comprising the first Fc domain subunit and a polypeptide comprising the second Fc domain subunit, which might be non-identical in the sense that further components fused to each of the subunits (e.g. antigen binding moieties) are not the same. In some embodiments the modification promoting association comprises an amino acid mutation in the Fc domain, specifically an amino acid substitution. In a particular embodiment, the modification promoting association comprises a separate amino acid mutation, specifically an amino acid substitution, in each of the two subunits of the Fc domain.

"Framework" or "FR" refers to variable domain residues other than hypervariable region (HVR) residues. The FR of a variable domain generally consists of four FR domains: FR1, FR2, FR3, and FR4. Accordingly, the HVR and FR sequences generally

appear in the following sequence in VH (or VL): FR1-H1(L1)-FR2-H2(L2)-FR3-H3(L3)-FR4.

The terms "full length antibody," "intact antibody," and "whole antibody" are used herein interchangeably to refer to an antibody having a structure substantially similar to a native antibody structure or having heavy chains that contain an Fc region as defined herein.

The terms "host cell," "host cell line," and "host cell culture" are used interchangeably and refer to cells into which exogenous nucleic acid has been introduced, including the progeny of such cells. Host cells include "transformants" and "transformed cells," which include the primary transformed cell and progeny derived therefrom without regard to the number of passages. Progeny may not be completely identical in nucleic acid content to a parent cell, but may contain mutations. Mutant progeny that have the same function or biological activity as screened or selected for in the originally transformed cell are included herein.

A "human antibody" is one which possesses an amino acid sequence which corresponds to that of an antibody produced by a human or a human cell or derived from a non-human source that utilizes human antibody repertoires or other human antibody-encoding sequences. This definition of a human antibody specifically excludes a humanized antibody comprising non-human antigen-binding residues. As also mentioned for chimeric and humanized antibodies according to the invention the term "human antibody" as used herein also comprises such antibodies which are modified in the constant region to generate the properties according to the invention, especially in regard to C1q binding and/or FcR binding, e.g. by "class switching" i.e. change or mutation of Fc parts (e.g. from IgG1 to IgG4 and/or IgG1/IgG4 mutation.)

The term "recombinant human antibody", as used herein, is intended to include all human antibodies that are prepared, expressed, created or isolated by recombinant means, such as antibodies isolated from a host cell such as a NS0 or CHO cell or from an animal (e.g. a mouse) that is transgenic for human immunoglobulin genes or antibodies expressed using a recombinant expression vector transfected into a host cell. Such recombinant human antibodies have variable and constant regions in a rearranged form. The recombinant human antibodies according to the invention have been subjected to *in vivo* somatic hypermutation. Thus, the amino acid sequences of the VH and VL regions of the recombinant antibodies are sequences that, while derived from and related to

human germ line VH and VL sequences, may not naturally exist within the human antibody germ line repertoire *in vivo*.

A “human consensus framework” is a framework which represents the most commonly occurring amino acid residues in a selection of human immunoglobulin VL or VH framework sequences. Generally, the selection of human immunoglobulin VL or VH sequences is from a subgroup of variable domain sequences. Generally, the subgroup of sequences is a subgroup as in Kabat et al., *Sequences of Proteins of Immunological Interest*, Fifth Edition, NIH Publication 91-3242, Bethesda MD (1991), vols. 1-3. In one embodiment, for the VL, the subgroup is subgroup kappa I as in Kabat et al., *supra*. In one embodiment, for the VH, the subgroup is subgroup III as in Kabat et al., *supra*.

A “humanized” antibody refers to a chimeric antibody comprising amino acid residues from non-human HVRs and amino acid residues from human FRs. In certain embodiments, a humanized antibody will comprise substantially all of at least one, and typically two, variable domains, in which all or substantially all of the HVRs (e.g., CDRs) correspond to those of a non-human antibody, and all or substantially all of the FRs correspond to those of a human antibody. A humanized antibody optionally may comprise at least a portion of an antibody constant region derived from a human antibody. A “humanized form” of an antibody, e.g., a non-human antibody, refers to an antibody that has undergone humanization. Other forms of “humanized antibodies” encompassed by the present invention are those in which the constant region has been additionally modified or changed from that of the original antibody to generate the properties according to the invention, especially in regard to C1q binding and/or Fc receptor (FcR) binding.

The term “hypervariable region” or “HVR,” as used herein refers to each of the regions of an antibody variable domain which are hypervariable in sequence and/or form structurally defined loops (“hypervariable loops”). Generally, native four-chain antibodies comprise six HVRs; three in the VH (H1, H2, H3), and three in the VL (L1, L2, L3). HVRs generally comprise amino acid residues from the hypervariable loops and/or from the “complementarity determining regions” (CDRs), the latter being of highest sequence variability and/or involved in antigen recognition. Exemplary hypervariable loops occur at amino acid residues 26-32 (L1), 50-52 (L2), 91-96 (L3), 26-32 (H1), 53-55 (H2), and 96-101 (H3). (Chothia and Lesk, *J. Mol. Biol.* 196:901-917 (1987).) Exemplary CDRs (CDR-L1, CDR-L2, CDR-L3, CDR-H1, CDR-H2, and CDR-H3) occur at amino acid residues 24-34 of L1, 50-56 of L2, 89-97 of L3, 31-35B of H1, 50-65 of H2, and 95-102

of H3. (Kabat et al., *Sequences of Proteins of Immunological Interest*, 5th Ed. Public Health Service, National Institutes of Health, Bethesda, MD (1991).) Hypervariable regions (HVRs) are also referred to as complementarity determining regions (CDRs), and these terms are used herein interchangeably in reference to portions of the variable region that form the antigen binding regions. This particular region has been described by Kabat et al., U.S. Dept. of Health and Human Services, "Sequences of Proteins of Immunological Interest" (1983) and by Chothia *et al.*, *J. Mol. Biol.* 196:901-917 (1987), where the definitions include overlapping or subsets of amino acid residues when compared against each other. Nevertheless, application of either definition to refer to a CDR of an antibody or variants thereof is intended to be within the scope of the term as defined and used herein. The appropriate amino acid residues which encompass the CDRs as defined by each of the above cited references are set forth below in Table A as a comparison. The exact residue numbers which encompass a particular CDR will vary depending on the sequence and size of the CDR. Those skilled in the art can routinely determine which residues comprise a particular CDR given the variable region amino acid sequence of the antibody.

TABLE A. CDR Definitions¹

CDR	Kabat	Chothia	AbM ²
V _H CDR1	31-35	26-32	26-35
V _H CDR2	50-65	52-58	50-58
V _H CDR3	95-102	95-102	95-102
V _L CDR1	24-34	26-32	24-34
V _L CDR2	50-56	50-52	50-56
V _L CDR3	89-97	91-96	89-97

¹ Numbering of all CDR definitions in Table A is according to the numbering conventions set forth by Kabat et al. (see below).

² "AbM" with a lowercase "b" as used in Table A refers to the CDRs as defined by Oxford Molecular's "AbM" antibody modeling software.

Kabat *et al.* also defined a numbering system for variable region sequences that is applicable to any antibody. One of ordinary skill in the art can unambiguously assign this system of "Kabat numbering" to any variable region sequence, without reliance on any experimental data beyond the sequence itself. As used herein, "Kabat numbering" refers to the numbering system set forth by Kabat et al., U.S. Dept. of Health and Human Services, "Sequence of Proteins of Immunological Interest" (1983). Unless otherwise

specified, references to the numbering of specific amino acid residue positions in an antibody variable region are according to the Kabat numbering system.

With the exception of CDR1 in VH, CDRs generally comprise the amino acid residues that form the hypervariable loops. CDRs also comprise “specificity determining residues,” or “SDRs,” which are residues that contact antigen. SDRs are contained within regions of the CDRs called abbreviated-CDRs, or a-CDRs. Exemplary a-CDRs (a-CDR-L1, a-CDR-L2, a-CDR-L3, a-CDR-H1, a-CDR-H2, and a-CDR-H3) occur at amino acid residues 31-34 of L1, 50-55 of L2, 89-96 of L3, 31-35B of H1, 50-58 of H2, and 95-102 of H3. (See Almagro and Fransson, *Front. Biosci.* 13:1619-1633 (2008).) Unless otherwise indicated, HVR residues and other residues in the variable domain (e.g., FR residues) are numbered herein according to Kabat et al., *supra*.

An “immunoconjugate” is an antibody conjugated to one or more heterologous molecule(s), including but not limited to a cytotoxic agent.

An “individual” or “subject” is a mammal. Mammals include, but are not limited to, domesticated animals (e.g., cows, sheep, cats, dogs, and horses), primates (e.g., humans and non-human primates such as monkeys), rabbits, and rodents (e.g., mice and rats). In certain embodiments, the individual or subject is a human.

An “isolated” antibody is one which has been separated from a component of its natural environment. In some embodiments, an antibody is purified to greater than 95% or 99% purity as determined by, for example, electrophoretic (e.g., SDS-PAGE, isoelectric focusing (IEF), capillary electrophoresis) or chromatographic (e.g., ion exchange or reverse phase HPLC). For review of methods for assessment of antibody purity, see, e.g., Flatman et al., *J. Chromatogr. B* 848:79-87 (2007).

An “isolated” nucleic acid refers to a nucleic acid molecule that has been separated from a component of its natural environment. An isolated nucleic acid includes a nucleic acid molecule contained in cells that ordinarily contain the nucleic acid molecule, but the nucleic acid molecule is present extrachromosomally or at a chromosomal location that is different from its natural chromosomal location.

“Isolated nucleic acid encoding a bispecific antibody that specifically binds DR5 and FAP antibody” refers to one or more nucleic acid molecules encoding antibody heavy and light chains (or fragments thereof), including such nucleic acid molecule(s) in a single vector or separate vectors, and such nucleic acid molecule(s) present at one or more locations in a host cell.

The term "monoclonal antibody" as used herein refers to an antibody obtained from a population of substantially homogeneous antibodies, i.e., the individual antibodies comprising the population are identical and/or bind the same epitope, except for possible variant antibodies, e.g., containing naturally occurring mutations or arising during
5 production of a monoclonal antibody preparation, such variants generally being present in minor amounts. In contrast to polyclonal antibody preparations, which typically include different antibodies directed against different determinants (epitopes), each monoclonal antibody of a monoclonal antibody preparation is directed against a single determinant on an antigen. Thus, the modifier "monoclonal" indicates the character of the antibody as
10 being obtained from a substantially homogeneous population of antibodies, and is not to be construed as requiring production of the antibody by any particular method. For example, the monoclonal antibodies to be used in accordance with the present invention may be made by a variety of techniques, including but not limited to the hybridoma method, recombinant DNA methods, phage-display methods, and methods utilizing
15 transgenic animals containing all or part of the human immunoglobulin loci, such methods and other exemplary methods for making monoclonal antibodies being described herein.

A "naked antibody" refers to an antibody that is not conjugated to a heterologous moiety (e.g., a cytotoxic moiety) or radiolabel. The naked antibody may be present in a
20 pharmaceutical formulation.

"Native antibodies" refer to naturally occurring immunoglobulin molecules with varying structures. For example, native IgG antibodies are heterotetrameric glycoproteins of about 150,000 daltons, composed of two identical light chains and two identical heavy chains that are disulfide-bonded. From N- to C-terminus, each heavy chain has a variable
25 region (VH), also called a variable heavy domain or a heavy chain variable domain, followed by three constant domains (CH1, CH2, and CH3). Similarly, from N- to C-terminus, each light chain has a variable region (VL), also called a variable light domain or a light chain variable domain, followed by a constant light (CL) domain. The light chain of an antibody may be assigned to one of two types, called kappa (κ) and lambda (λ),
30 based on the amino acid sequence of its constant domain.

A "blocking" antibody or an "antagonist" antibody is one that inhibits or reduces a biological activity of the antigen it binds. In some embodiments, blocking antibodies or antagonist antibodies substantially or completely inhibit the biological activity of the

antigen. For example, the anti-PD-L1 antibodies of the invention block the signaling through PD-1 so as to restore a functional response by T-cells (e.g., proliferation, cytokine production, target cell killing) from a dysfunctional state to antigen stimulation.

5 An "agonist" or activating antibody is one that enhances or initiates signaling by the antigen to which it binds. In some embodiments, agonist antibodies cause or activate signaling without the presence of the natural ligand.

The term "package insert" is used to refer to instructions customarily included in commercial packages of therapeutic products, that contain information about the indications, usage, dosage, administration, combination therapy, contraindications and/or
10 warnings concerning the use of such therapeutic products.

"No substantial cross-reactivity" means that a molecule (e.g., an antibody) does not recognize or specifically bind an antigen different from the actual target antigen of the molecule (e.g. an antigen closely related to the target antigen), particularly when compared to that target antigen. For example, an antibody may bind less than about 10%
15 to less than about 5% to an antigen different from the actual target antigen, or may bind said antigen different from the actual target antigen at an amount consisting of less than about 10%, 9%, 8%, 7%, 6%, 5%, 4%, 3%, 2%, 1%, 0.5%, 0.2%, or 0.1%, preferably less than about 2%, 1%, or 0.5%, and most preferably less than about 0.2% or 0.1% antigen different from the actual target antigen.

20 "Percent (%) amino acid sequence identity" with respect to a reference polypeptide sequence is defined as the percentage of amino acid residues in a candidate sequence that are identical with the amino acid residues in the reference polypeptide sequence, after aligning the sequences and introducing gaps, if necessary, to achieve the maximum percent sequence identity, and not considering any conservative substitutions
25 as part of the sequence identity. Alignment for purposes of determining percent amino acid sequence identity can be achieved in various ways that are within the skill in the art, for instance, using publicly available computer software such as BLAST, BLAST-2, ALIGN or Megalign (DNASTAR) software. Those skilled in the art can determine appropriate parameters for aligning sequences, including any algorithms needed to
30 achieve maximal alignment over the full length of the sequences being compared. For purposes herein, however, % amino acid sequence identity values are generated using the sequence comparison computer program ALIGN-2. The ALIGN-2 sequence comparison computer program was authored by Genentech, Inc., and the source code has been filed

with user documentation in the U.S. Copyright Office, Washington D.C., 20559, where it is registered under U.S. Copyright Registration No. TXU510087. The ALIGN-2 program is publicly available from Genentech, Inc., South San Francisco, California, or may be compiled from the source code. The ALIGN-2 program should be compiled for use on a
5 UNIX operating system, including digital UNIX V4.0D. All sequence comparison parameters are set by the ALIGN-2 program and do not vary.

In situations where ALIGN-2 is employed for amino acid sequence comparisons, the % amino acid sequence identity of a given amino acid sequence A to, with, or against a given amino acid sequence B (which can alternatively be phrased as a given amino acid
10 sequence A that has or comprises a certain % amino acid sequence identity to, with, or against a given amino acid sequence B) is calculated as follows:

$$100 \text{ times the fraction } X/Y$$

15 where X is the number of amino acid residues scored as identical matches by the sequence alignment program ALIGN-2 in that program's alignment of A and B, and where Y is the total number of amino acid residues in B. It will be appreciated that where the length of amino acid sequence A is not equal to the length of amino acid sequence B, the % amino acid sequence identity of A to B will not equal the % amino acid sequence
20 identity of B to A. Unless specifically stated otherwise, all % amino acid sequence identity values used herein are obtained as described in the immediately preceding paragraph using the ALIGN-2 computer program.

The term "pharmaceutical formulation" refers to a preparation which is in such form as to permit the biological activity of an active ingredient contained therein to be
25 effective, and which contains no additional components which are unacceptably toxic to a subject to which the formulation would be administered.

A "pharmaceutically acceptable carrier" refers to an ingredient in a pharmaceutical formulation, other than an active ingredient, which is nontoxic to a subject. A pharmaceutically acceptable carrier includes, but is not limited to, a buffer,
30 excipient, stabilizer, or preservative.

The term "PD-1 axis binding antagonist" is a molecule that inhibits the interaction of a PD-1 axis binding partner with either one or more of its binding partner, so as to remove T-cell dysfunction resulting from signaling on the PD-1 signaling axis - with a

result being to restore or enhance T-cell function {e.g., proliferation, cytokine production, target cell killing). As used herein, a PD- 1 axis binding antagonist includes a PD- 1 binding antagonist, a PD-L1 binding antagonist and a PD-L2 binding antagonist.

The term "PD-1 binding antagonists" is a molecule that decreases, blocks, inhibits, abrogates or interferes with signal transduction resulting from the interaction of PD-1 with one or more of its binding partners, such as PD-L1, PD-L2. In some embodiments, the PD-1 binding antagonist is a molecule that inhibits the binding of PD-1 to its binding partners. In a specific aspect, the PD-1 binding antagonist inhibits the binding of PD- 1 to PD-L1 and/or PD-L2. For example, PD-1 binding antagonists include anti-PD-1 antibodies, antigen binding fragments thereof, immunoadhesins, fusion proteins, oligopeptides and other molecules that decrease, block, inhibit, abrogate or interfere with signal transduction resulting from the interaction of PD- 1 with PD-L1 and/or PD-L2. In one embodiment, a PD-1 binding antagonist reduces the negative co-stimulatory signal mediated by or through cell surface proteins expressed on T lymphocytes mediated signaling through PD-1 so as render a dysfunctional T-cell less dysfunctional (e.g. , enhancing effector responses to antigen recognition). In some embodiments, the PD-1 binding antagonist is an anti-PD-1 antibody. In a specific aspect, a PD- 1 binding antagonist is MDX- 1106 described herein. In another specific aspect, a PD-1 binding antagonist is Merck 3745 described herein. In another specific aspect, a PD-1 binding antagonist is CT-01 1 described herein.

The term "PD-L1 binding antagonists" is a molecule that decreases, blocks, inhibits, abrogates or interferes with signal transduction resulting from the interaction of PD-L1 with either one or more of its binding partners, such as PD-1 , B7-1 . In some embodiments, a PD-L1 binding antagonist is a molecule that inhibits the binding of PD-L 1 to its binding partners. In a specific aspect, the PD-L1 binding antagonist inhibits binding of PD-L1 to PD-1 and/or B7-1. In some embodiments, the PD-L1 binding antagonists include anti-PD-L1 antibodies, antigen binding fragments thereof, immunoadhesins, fusion proteins, oligopeptides and other molecules that decrease, block, inhibit, abrogate or interfere with signal transduction resulting from the interaction of PD-L1 with one or more of its binding partners, such as PD-1, B7- 1 . In one embodiment, a PD-L1 binding antagonist reduces the negative co-stimulatory signal mediated by or through cell surface proteins expressed on T lymphocytes mediated signaling through PD-L 1 so as to render a dysfunctional T-cell less dysfunctional (e.g. , enhancing effector

responses to antigen recognition). In some embodiments, a PD-L1 binding antagonist is an anti-PD-L1 antibody. In a specific aspect, an anti-PD-L1 antibody is YW243.55.S70 described herein. In another specific aspect, an anti-PD-L1 antibody is MDX- 1 105 described herein. In still another specific aspect, an anti-PD-L1 antibody is MPDL3280A 5 described herein.

The term "PD-L2 binding antagonists" is a molecule that decreases, blocks, inhibits, abrogates or interferes with signal transduction resulting from the interaction of PD-L2 with either one or more of its binding partners, such as PD- 1. In some 10 embodiments, a PD-L2 binding antagonist is a molecule that inhibits the binding of PD-L2 to its binding partners. In a specific aspect, the PD-L2 binding antagonist inhibits binding of PD-L2 to PD- 1. In some embodiments, the PD-L2 antagonists include anti-PD-L2 antibodies, antigen binding fragments thereof, immunoadhesins, fusion proteins, oligopeptides and other molecules that decrease, block, inhibit, abrogate or interfere with 15 signal transduction resulting from the interaction of PD- L2 with either one or more of its binding partners, such as PD-1. In one embodiment, a PD-L2 binding antagonist reduces the negative co-stimulatory signal mediated by or through cell surface proteins expressed on T lymphocytes mediated signaling through PD-L2 so as render a dysfunctional T-cell less dysfunctional (e.g. , enhancing effector responses to antigen recognition). In some 20 embodiments, a PD-L2 binding antagonist is an immunoadhesin.

A "PD-1 oligopeptide " "PD-L1 oligopeptide " or "PD-L2 oligopeptide" is an oligopeptide that binds, preferably specifically, to a PD-1 , PD-L1 or PD-L2 negative 25 costimulatory polypeptide, respectively, including a receptor, ligand or signaling component, respectively, as described herein. Such oligopeptides may be chemically synthesized using known oligopeptide synthesis methodology or may be prepared and purified using recombinant technology. Such oligopeptides are usually at least about 5 30 amino acids in length, alternatively at least about 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, or 100 amino acids in length or more. Such oligopeptides may be identified using well known techniques. In this regard, it is noted that techniques for screening oligopeptide libraries for oligopeptides that are capable of specifically binding to a polypeptide target are well known in the art (see, e.g., U.S.

Patent Nos. 5,556,762, 5,750,373, 4,708,871 , 4,833,092, 5,223,409, 5,403,484, 5,571,689, 5,663, 143; PCT Publication Nos. WO 84/03506 and WO84/03564; Geysen et al., Proc. Natl. Acad. Sci. U.S.A., 81:3998-4002 (1984); Geysen et al, Proc. Natl. Acad. Sci. U.S.A., 82: 178-182 (1985); Geysen et al, in Synthetic Peptides as Antigens, 130-149 (1986); Geysen et al., J. Immunol. Metk, 102:259-274 (1987); Schoofs et al., J. Immunol., 140:611-616 (1988), Cwirla, S. E. et al. Proc. Natl. Acad. Sci. USA, 87:6378 (1990); Lowman, H.B. et al. Biochemistry, 30: 10832 (1991); Clackson, T. et al. Nature, 352: 624 (1991); Marks, J. D. et al., J. Mol. Biol., 222:581 (1991); Kang, A.S. et al. Proc. Natl. Acad. Sci. USA, 88:8363 (1991), and Smith, G. P., Current Opin. Biotechnol, 2:668 (1991).

The term "anergy" refers to the state of unresponsiveness to antigen stimulation resulting from incomplete or insufficient signals delivered through the T-cell receptor (e.g. increase in intracellular Ca^{+2} in the absence of ras-activation). T cell anergy can also result upon stimulation with antigen in the absence of co-stimulation, resulting in the cell becoming refractory to subsequent activation by the antigen even in the context of costimulation. The unresponsive state can often be overridden by the presence of Interleukin-2. Anergic T-cells do not undergo clonal expansion and/or acquire effector functions.

The term "exhaustion" refers to T cell exhaustion as a state of T cell dysfunction that arises from sustained TCR signaling that occurs during many chronic infections and cancer. It is distinguished from anergy in that it arises not through incomplete or deficient signaling, but from sustained signaling. It is defined by poor effector function, sustained expression of inhibitory receptors and a transcriptional state distinct from that of functional effector or memory T cells. Exhaustion prevents optimal control of infection and tumors. Exhaustion can result from both extrinsic negative regulatory pathways (e.g., immunoregulatory cytokines) as well as cell intrinsic negative regulatory (costimulatory) pathways (PD-1, B7-H3, B7-H4, etc.).

"Enhancing T-cell function " means to induce, cause or stimulate a T-cell to have a sustained or amplified biological function, or renew or reactivate exhausted or inactive T-cells. Examples of enhancing T-cell function include: increased secretion of γ -interferon from CD8⁺ T-cells, increased proliferation, increased antigen responsiveness (e.g. , viral, pathogen, or tumor clearance) relative to such levels before the intervention. In one embodiment, the level of . enhancement is as least 50%, alternatively 60%, 70%,

80%, 90%, 100%, 120%, 150%, 200%. The manner of measuring this enhancement is known to one of ordinary skill in the art.

"Tumor immunity" refers to the process in which tumors evade immune recognition and clearance. Thus, as a therapeutic concept, tumor immunity is "treated" when such evasion is attenuated, and the tumors are recognized and attacked by the immune system. Examples of tumor recognition include tumor binding, tumor shrinkage and tumor clearance. [0046] "Immunogenicity" refers to the ability of a particular substance to provoke an immune response. Tumors are immunogenic and enhancing tumor immunogenicity aids in the clearance of the tumor cells by the immune response. Examples of enhancing tumor immunogenicity include treatment with anti-PDL antibodies and a ME inhibitor.

"Sustained response" refers to the sustained effect on reducing tumor growth after cessation of a treatment. For example, the tumor size may remain to be the same or smaller as compared to the size at the beginning of the administration phase. In some embodiments, the sustained response has a duration at least the same as the treatment duration, at least 1.5X, 2.0X, 2.5X, or 3.0X length of the treatment duration.

The term "Fibroblast activation protein (FAP)", as used herein, refers to any native FAP from any vertebrate source, including mammals such as primates (e.g. humans) and rodents (e.g., mice and rats), unless otherwise indicated. The term encompasses "full-length," unprocessed FAP as well as any form of FAP that results from processing in the cell. The term also encompasses naturally occurring variants of FAP, e.g., splice variants or allelic variants. Preferably, an anti-FAP antibody of the invention binds to the extracellular domain of FAP. The amino acid sequence of exemplary FAP polypeptide sequences, including the sequence of human FAP, are disclosed in WO 2012/020006.

As used herein, "treatment" (and grammatical variations thereof such as "treat" or "treating") refers to clinical intervention in an attempt to alter the natural course of the individual being treated, and can be performed either for prophylaxis or during the course of clinical pathology. Desirable effects of treatment include, but are not limited to, preventing occurrence or recurrence of disease, alleviation of symptoms, diminishment of any direct or indirect pathological consequences of the disease, preventing metastasis, decreasing the rate of disease progression, amelioration or palliation of the disease state,

and remission or improved prognosis. In some embodiments, antibodies of the invention are used to delay development of a disease or to slow the progression of a disease.

The term cancer as used herein refers to proliferative diseases, such as the cancer is colorectal cancer, sarcoma, head and neck cancer, squamous cell carcinoma, breast
5 cancer, pancreatic cancer, gastric cancer, non-small-cell lung carcinoma, small-cell lung cancer and mesothelioma, including refractory versions of any of the above cancers, or a combination of one or more of the above cancers. In one embodiment, the cancer is colorectal cancer and optionally the chemotherapeutic agent is Irinotecan. In
10 embodiments in which the cancer is sarcoma, optionally the sarcoma is chondrosarcoma, leiomyosarcoma, gastrointestinal stromal tumours, fibrosarcoma, osteosarcoma, liposarcoma or malignant fibrous histiocytoma.

The term "variable region" or "variable domain" refers to the domain of an antibody heavy or light chain that is involved in binding the antibody to antigen. The variable domains of the heavy chain and light chain (VH and VL, respectively) of a native
15 antibody generally have similar structures, with each domain comprising four conserved framework regions (FRs) and three hypervariable regions (HVRs). (See, e.g., Kindt et al. *Kuby Immunology*, 6th ed., W.H. Freeman and Co., page 91 (2007).) A single VH or VL domain may be sufficient to confer antigen-binding specificity. Furthermore, antibodies that bind a particular antigen may be isolated using a VH or VL domain from an antibody
20 that binds the antigen to screen a library of complementary VL or VH domains, respectively. See, e.g., Portolano et al., *J. Immunol.* 150:880-887 (1993); Clarkson et al., *Nature* 352:624-628 (1991).

As used herein, the term "antigen binding molecule" refers in its broadest sense to a molecule that specifically binds an antigenic determinant. Examples of antigen binding
25 molecules are immunoglobulins and derivatives, e.g. fragments, thereof.

The term "antigen-binding site of an antibody" when used herein refer to the amino acid residues of an antibody which are responsible for antigen-binding. The antigen-binding portion of an antibody comprises amino acid residues from the "complementary determining regions" or "CDRs". "Framework" or "FR" regions are
30 those variable domain regions other than the hypervariable region residues as herein defined. Therefore, the light and heavy chain variable domains of an antibody comprise from N- to C-terminus the domains FR1, CDR1, FR2, CDR2, FR3, CDR3, and FR4. Especially, CDR3 of the heavy chain is the region which contributes most to antigen

binding and defines the antibody's properties. CDR and FR regions are determined according to the standard definition of Kabat et al., Sequences of Proteins of Immunological Interest, 5th ed., Public Health Service, National Institutes of Health, Bethesda, MD (1991) and/or those residues from a "hypervariable loop".

5 Antibody specificity refers to selective recognition of the antibody for a particular epitope of an antigen. Natural antibodies, for example, are monospecific. "Bispecific antibodies" according to the invention are antibodies which have two different antigen-binding specificities. Antibodies of the present invention are specific for two different antigens, i.e. DR5 as first antigen and FAP as second antigen.

10 The term "monospecific" antibody as used herein denotes an antibody that has one or more binding sites each of which bind to the same epitope of the same antigen.

 The term "bispecific" means that the antigen binding molecule is able to specifically bind to at least two distinct antigenic determinants. Typically, a bispecific antigen binding molecule comprises at least two antigen binding sites, each of which is
15 specific for a different antigenic determinant. In certain embodiments the bispecific antigen binding molecule is capable of simultaneously binding two antigenic determinants, particularly two antigenic determinants expressed on two distinct cells.

 The antibody provided herein is a multispecific antibody, e.g. a bispecific antibody. Multispecific antibodies are monoclonal antibodies that have binding
20 specificities for at least two different sites. Provided herein is a bispecific antibody, with binding specificities for FAP and DR5. In certain embodiments, bispecific antibodies may bind to two different epitopes of DR5. Bispecific antibodies may also be used to localize cytotoxic agents to cells which express DR5. Bispecific antibodies can be prepared as full length antibodies or antibody fragments.

25 Techniques for making multispecific antibodies include, but are not limited to, recombinant co-expression of two immunoglobulin heavy chain-light chain pairs having different specificities (see Milstein and Cuello, *Nature* 305: 537 (1983)), WO 93/08829, and Traunecker et al., *EMBO J.* 10: 3655 (1991)), and "knob-in-hole" engineering (see, e.g., U.S. Patent No. 5,731,168). Multi-specific antibodies may also be made by
30 engineering electrostatic steering effects for making antibody Fc-heterodimeric molecules (WO 2009/089004); cross-linking two or more antibodies or fragments (see, e.g., US Patent No. 4,676,980, and Brennan et al., *Science*, 229: 81 (1985)); using leucine zippers to produce bi-specific antibodies (see, e.g., Kostelny et al., *J. Immunol.*, 148(5):1547-

1553 (1992)); using "diabody" technology for making bispecific antibody fragments (see, e.g., Hollinger et al., *Proc. Natl. Acad. Sci. USA*, 90:6444-6448 (1993)); and using single-chain Fv (sFv) dimers (see, e.g., Gruber et al., *J. Immunol.*, 152:5368 (1994)); and preparing trispecific antibodies as described, e.g., in Tutt et al. *J. Immunol.* 147: 60
5 (1991).

Engineered antibodies with three or more functional antigen binding sites, including "Octopus antibodies," are also included herein (see, e.g. US 2006/0025576A1).

The antibody or fragment herein also includes a "Dual Acting FAb" or "DAF" comprising at least one antigen binding site that binds to FAP or DR5 as well as another,
10 different antigen (see, US 2008/0069820, for example).

The term "valent" as used within the current application denotes the presence of a specified number of binding sites in an antibody molecule. As such, the terms "bivalent", "tetravalent", and "hexavalent" denote the presence of two binding sites, four binding sites, and six binding sites, respectively, in an antibody molecule. The bispecific
15 antibodies according to the invention are at least "bivalent" and may be "trivalent" or "multivalent" (e.g. "tetravalent" or "hexavalent").

Antibodies of the present invention have two or more binding sites and are bispecific. That is, the antibodies may be bispecific even in cases where there are more than two binding sites (i.e. that the antibody is trivalent or multivalent). Bispecific
20 antibodies of the invention include, for example, multivalent single chain antibodies, diabodies and triabodies, as well as antibodies having the constant domain structure of full length antibodies to which further antigen-binding sites (e.g., single chain Fv, a VH domain and/or a VL domain, Fab, or (Fab)₂) are linked via one or more peptide-linkers. The antibodies can be full length from a single species, or be chimerized or humanized.

25 The term "vector," as used herein, refers to a nucleic acid molecule capable of propagating another nucleic acid to which it is linked. The term includes the vector as a self-replicating nucleic acid structure as well as the vector incorporated into the genome of a host cell into which it has been introduced. Certain vectors are capable of directing the expression of nucleic acids to which they are operatively linked. Such vectors are
30 referred to herein as "expression vectors."

The term "amino acid" as used within this application denotes the group of naturally occurring carboxy α -amino acids comprising alanine (three letter code: ala, one letter code: A), arginine (arg, R), asparagine (asn, N), aspartic acid (asp, D), cysteine (cys,

C), glutamine (gln, Q), glutamic acid (glu, E), glycine (gly, G), histidine (his, H), isoleucine (ile, I), leucine (leu, L), lysine (lys, K), methionine (met, M), phenylalanine (phe, F), proline (pro, P), serine (ser, S), threonine (thr, T), tryptophan (trp, W), tyrosine (tyr, Y), and valine (val, V).

5 As used herein, the expressions “cell”, “cell line”, and “cell culture” are used interchangeably and all such designations include progeny. Thus, the words “transfectants” and “transfected cells” include the primary subject cell and cultures derived there from without regard for the number of transfers. It is also understood that all progeny may not be precisely identical in DNA content, due to deliberate or inadvertent
10 mutations. Variant progeny that have the same function or biological activity as screened for in the originally transformed cell are included.

“Affinity” refers to the strength of the sum total of noncovalent interactions between a single binding site of a molecule (e.g., an antibody) and its binding partner (e.g., an antigen). Unless indicated otherwise, as used herein, “binding affinity” refers to
15 intrinsic binding affinity which reflects a 1:1 interaction between members of a binding pair (e.g., antibody and antigen). The affinity of a molecule X for its partner Y can generally be represented by the dissociation constant (K_d). Affinity can be measured by common methods known in the art, including those described herein. Specific illustrative and exemplary embodiments for measuring binding affinity are described in the
20 following.

As used herein, the term “binding” or “specifically binding” refers to the binding of the antibody to an epitope of the antigen in an in-vitro assay, preferably in a surface plasmon resonance assay (SPR, BIAcore, GE-Healthcare Uppsala, Sweden). The affinity of the binding is defined by the terms k_a (rate constant for the association of the antibody
25 from the antibody/antigen complex), k_d (dissociation constant), and KD (kD/k_a). Binding or specifically binding means a binding affinity (KD) of 10^{-8} mol/l or less, preferably 10^{-9} M to 10^{-13} mol/l.

Binding of the antibody to the death receptor can be investigated by a BIAcore assay (GE-Healthcare Uppsala, Sweden). The affinity of the binding is defined by the
30 terms k_a (rate constant for the association of the antibody from the antibody/antigen complex), k_d (dissociation constant), and KD (kD/k_a)

“Reduced binding”, for example reduced binding to an Fc receptor, refers to a decrease in affinity for the respective interaction, as measured for example by SPR. For clarity the

term includes also reduction of the affinity to zero (or below the detection limit of the analytic method), i.e. complete abolishment of the interaction. Conversely, "increased binding" refers to an increase in binding affinity for the respective interaction.

"T cell activation" as used herein refers to one or more cellular response of a T lymphocyte, particularly a cytotoxic T lymphocyte, selected from: proliferation, differentiation, cytokine secretion, cytotoxic effector molecule release, cytotoxic activity, and expression of activation markers. The T cell activating bispecific antigen binding molecules of the invention are capable of inducing T cell activation. Suitable assays to measure T cell activation are known in the art described herein.

10 A "target cell antigen" as used herein refers to an antigenic determinant presented on the surface of a target cell, for example a cell in a tumor such as a cancer cell or a cell of the tumor stroma. In particular "target cell antigen" refers to Folate Receptor 1.

As used herein, the terms "first" and "second" with respect to antigen binding moieties etc., are used for convenience of distinguishing when there is more than one of each type of moiety. Use of these terms is not intended to confer a specific order or orientation of the T cell activating bispecific antigen binding molecule unless explicitly so stated.

The term "epitope" includes any polypeptide determinant capable of specific binding to an antibody. In certain embodiments, epitope determinant include chemically active surface groupings of molecules such as amino acids, sugar side chains, phosphoryl, or sulfonyl, and, in certain embodiments, may have specific three dimensional structural characteristics, and or specific charge characteristics. An epitope is a region of an antigen that is bound by an antibody.

As used herein, the term "antigenic determinant" is synonymous with "antigen" and "epitope," and refers to a site (e.g. a contiguous stretch of amino acids or a conformational configuration made up of different regions of non-contiguous amino acids) on a polypeptide macromolecule to which an antigen binding moiety binds, forming an antigen binding moiety-antigen complex. Useful antigenic determinants can be found, for example, on the surfaces of tumor cells, on the surfaces of virus-infected cells, on the surfaces of other diseased cells, on the surface of immune cells, free in blood serum, and/or in the extracellular matrix (ECM). The proteins referred to as antigens herein, e.g., FolR1 and CD3, can be any native form the proteins from any vertebrate source, including mammals such as primates (e.g. humans) and rodents (e.g. mice and rats), unless otherwise indicated. In a particular embodiment the antigen is a human

protein. Where reference is made to a specific protein herein, the term encompasses the “full-length”, unprocessed protein as well as any form of the protein that results from processing in the cell. The term also encompasses naturally occurring variants of the protein, e.g. splice variants or allelic variants. Exemplary human proteins useful as
5 antigens include, but are not limited to: FolR1 (Folate receptor alpha (FRA); Folate binding protein (FBP); human FolR1 UniProt no.: P15328; murine FolR1 UniProt no.: P35846; cynomolgus FolR1 UniProt no.: G7PR14) and CD3, particularly the epsilon subunit of CD3 (see UniProt no. P07766 (version 130), NCBI RefSeq no. NP_000724.1, SEQ ID NO:150 for the human sequence; or UniProt no. Q95LI5 (version 49), NCBI
10 GenBank no. BAB71849.1, for the cynomolgus [*Macaca fascicularis*] sequence). The T cell activating bispecific antigen binding molecule of the invention binds to an epitope of CD3 or a target cell antigen that is conserved among the CD3 or target antigen from different species. In certain embodiments the T cell activating bispecific antigen binding molecule of the invention binds to CD3 and FolR1, but does not bind to FolR2 (Folate
15 receptor beta; FRB; human FolR2 UniProt no.: P14207) or FolR3 (Folate receptor gamma; human FolR3 UniProt no.: P41439).

As used herein, the terms “engineer, engineered, engineering,” particularly with the prefix “glyco-,” as well as the term “glycosylation engineering” are considered to include any manipulation of the glycosylation pattern of a naturally occurring or
20 recombinant polypeptide or fragment thereof. Glycosylation engineering includes metabolic engineering of the glycosylation machinery of a cell, including genetic manipulations of the oligosaccharide synthesis pathways to achieve altered glycosylation of glycoproteins expressed in cells. Furthermore, glycosylation engineering includes the effects of mutations and cell environment on glycosylation. In one embodiment, the
25 glycosylation engineering is an alteration in glycosyltransferase activity. In a particular embodiment, the engineering results in altered glucosaminyltransferase activity and/or fucosyltransferase activity.

II. COMPOSITIONS AND METHODS

30 In one aspect, the invention is based on the use of a therapeutic combination of a T cell activating bispecific antigen binding molecule, e.g., a T cell activating bispecific antigen binding molecule comprising a first antigen binding site specific for Folate Receptor 1 (FolR1) and a second antigen binding site specific for CD3, and a PD-1 axis

binding antagonist, e.g., for the treatment of cancer. In some embodiments the therapeutic combination further includes a TIM3 antagonist.

A. Combination therapies of a T cell activating bispecific antigen binding molecule and a PD-1 axis binding antagonist

5 Broadly, the present invention relates to T cell activating bispecific antigen binding molecules and their use in combination with a PD-1 axis binding antagonists. The advantage of the combination over monotherapy is that the T cell activating bispecific antigen binding molecules used in the present invention enable re-direction and activation of T cells to the targeted cell while the PD-1 axis binding antagonist enhances
10 T cell function by reducing T cell exhaustion.

 In one aspect, provided herein is a method for treating or delaying progression of cancer in an individual comprising administering to the individual an effective amount of a T cell activating bispecific antigen binding molecules, e.g., a FolR1-TCB, and a PD-1 axis binding antagonist. In some embodiments, the treatment results in sustained response
15 in the individual after cessation of the treatment. The methods of this invention may find use in treating conditions where enhanced immunogenicity is desired such as increasing tumor immunogenicity for the treatment of cancer. A variety of cancers may be treated, or their progression may be delayed, including but are not limited to a cancer that may contain a BRAF V600E mutation, a cancer that may contain a BRAF wildtype, a cancer
20 that may contain a KRAS wildtype, or a cancer that may contain an activating KRAS mutation.

 In some embodiments, the individual has endometrial cancer. The endometrial cancer may be at early stage or late state. In some embodiments, the individual has melanoma. The melanoma may be at early stage or at late stage. In some embodiments,
25 the individual has colorectal cancer. The colorectal cancer may be at early stage or at late stage. In some embodiments, the individual has lung cancer, e.g., non-small cell lung cancer. The non-small cell lung cancer may be at early stage or at late stage. In some embodiments, the individual has pancreatic cancer. The pancreatic cancer may be at early stage or late state. In some embodiments, the individual has a hematological
30 malignancy. The hematological malignancy may be early stage or late stage. In some embodiments, the individual has ovarian cancer. The ovarian cancer may be at early stage or at late stage. In some embodiments, the individual has breast cancer. The breast cancer

may be at early stage or at late stage. In some embodiments, the individual has renal cell carcinoma. The renal cell carcinoma may be at early stage or at late stage.

In some embodiments, the individual is a mammal, such as domesticated animals (e.g., cows, sheep, cats, dogs, and horses), primates (e.g., humans and non-human
5 primates such as monkeys), rabbits, and rodents (e.g., mice and rats). In some embodiments, the individual treated is a human.

In another aspect, provided herein is a method of enhancing immune function in an individual having cancer comprising administering an effective amount of a T cell activating bispecific antigen binding molecules, specifically, a FolR1-TCB, and a PD-1
10 axis binding antagonist.

In some embodiments, the T cells in the individual have enhanced priming, activation, proliferation and/or effector function relative to prior to the administration of the T cell activating bispecific antigen binding molecules and the PD-1 pathway antagonist. In some embodiments, the T cell effector function is secretion of at least one
15 of IL-2, IFN- γ and TNF- α . In one embodiment, administering a FolR1-TCB and an anti-PDL-1 antibody results in increased T cell secretion of IL-2, IFN- γ and TNF- α . In some embodiments, the T cell is a CD8+ T cell. In some embodiments, the T cell priming is characterized by elevated CD44 expression and/or enhanced cytolytic activity in CD8 T cells. In some embodiments, the CD8 T cell activation is characterized by an elevated
20 frequency of γ -IFT⁺ CD8 T cells. In some embodiments, the CD8 T cell is an antigen-specific T-cell. In some embodiments, the immune evasion by signaling through PD-L1 surface expression is inhibited. In some embodiments, the cancer has elevated levels of T-cell infiltration.

In some embodiments, the combination therapy of the invention comprises
25 administration of a FolR1-TCB and a PD-1 axis binding antagonist. The FolR1-TCB and a PD-1 axis binding antagonist may be administered in any suitable manner known in the art. For example, FolR1-TCB and a PD-1 axis binding antagonist may be administered sequentially (at different times) or concurrently (at the same time). In some
embodiments, the FolR1-TCB is administered continuously. In some embodiments, the
30 FolR1-TCB is administered intermittently. In some embodiments, the FolR1-TCB is administered before administration of the PD-1 axis binding antagonist. In some
embodiments, the FolR1-TCB is administered simultaneously with administration of the

PD-1 axis binding antagonist. In some embodiments, the FolR1-TCB is administered after administration of the PD-1 axis binding antagonist.

In some embodiments, provided is a method for treating or delaying progression of cancer in an individual comprising administering to the individual an effective amount of a T cell activating bispecific antigen binding molecules, e.g., a FolR1-TCB, and a PD-1 axis binding antagonist, further comprising administering an additional therapy. Specifically contemplated is an embodiment in which the additional therapy comprises a TIM-3 antagonist. Accordingly, in one aspect, provided herein is a method for treating or delaying progression of cancer in an individual comprising administering to the individual an effective amount of a T cell activating bispecific antigen binding molecules, specifically, a FolR1-TCB, a PD-1 axis binding antagonist, and a TIM-3 antagonist. Any TIM3 antagonist, e.g., those described herein, can be used. The additional therapy may also be radiation therapy, surgery (e.g., lumpectomy and a mastectomy), chemotherapy, gene therapy, DNA therapy, viral therapy, R A therapy, immunotherapy, bone marrow transplantation, nanotherapy, monoclonal antibody therapy, or a combination of the foregoing. The additional therapy may be in the form of adjuvant or neoadjuvant therapy. In some embodiments, the additional therapy is the administration of small molecule enzymatic inhibitor or anti-metastatic agent. In some embodiments, the additional therapy is the administration of side-effect limiting agents (e.g., agents intended to lessen the occurrence and/or severity of side effects of treatment, such as anti-nausea agents, etc.). In some embodiments, the additional therapy is radiation therapy. In some embodiments, the additional therapy is surgery. In some embodiments, the additional therapy is a combination of radiation therapy and surgery. In some embodiments, the additional therapy is gamma irradiation. In some embodiments, the additional therapy is therapy targeting P13K/A T/mTOR pathway, HSP90 inhibitor, tubulin inhibitor, apoptosis inhibitor, and/or chemopreventative agent. The additional therapy may be one or more of the chemotherapeutic agents described hereabove.

T cell activating bispecific antigen binding molecules, e.g., a FolR1-TCB, and the PD-1 axis binding antagonist may be administered by the same route of administration or by different routes of administration. In some embodiments, T cell activating bispecific antigen binding molecules, e.g., a FolR1-TCB is administered intravenously, intramuscularly, subcutaneously, topically, orally, transdermally, intraperitoneally, intraprbitally, by implantation, by inhalation, intrathecally, intraventricularly, or

intranasally. In some embodiments, the PD-1 axis binding antagonist is administered intravenously, intramuscularly, subcutaneously, topically, orally, transdermally, intraperitoneally, intraorbitally, by implantation, by inhalation, intrathecally, intraventricularly, or intranasally. An effective amount of the T cell activating bispecific antigen binding molecules and the PD-1 axis binding antagonist may be administered for prevention or treatment of disease. The appropriate dosage of the T cell activating bispecific antigen binding molecules and/or the PD-1 axis binding antagonist may be determined based on the type of disease to be treated, the type of the T cell activating bispecific antigen binding molecules and the PD-1 axis binding antagonist, the severity and course of the disease, the clinical condition of the individual, the individual's clinical history and response to the treatment, and the discretion of the attending physician.

Any of the T cell activating bispecific antigen binding molecules, PD-1 axis binding antagonists and the TIM-3 antagonists known in the art or described below may be used in the methods.

In a further aspect, the present invention provides a pharmaceutical composition comprising a T cell activating bispecific antigen binding molecules as described herein, a PD-1 axis binding antagonists as described herein and a pharmaceutically acceptable carrier. In some embodiments, the pharmaceutical composition further comprises a TIM3 antagonist.

In a further aspect, the invention provides for a kit comprising a T cell activating bispecific antigen binding molecule specific for Folate Receptor 1 (FolR1) and CD3, and a package insert comprising instructions for using the T cell activating bispecific antigen binding molecule with a PD-1 axis binding antagonist to treat or delay progression of cancer in an individual. In some embodiments, the kit further comprises instructions for using the T cell activating bispecific antigen binding molecule with a TIM3 antagonist.

In a further aspect, the invention provides for a kit comprising a T cell activating bispecific antigen binding molecule specific for Folate Receptor 1 (FolR1) and CD3 and a PD-1 axis binding antagonist, and a package insert comprising instructions for using the T cell activating bispecific antigen binding molecule and the PD-1 axis binding antagonist to treat or delay progression of cancer in an individual. In one embodiment, the kit further comprises a TIM3 antagonist. In one of the embodiments, the PD-1 axis binding antagonist is an anti-PD-1 antibody or an anti-PDL-1 antibody. In one embodiment, the PD-1 axis binding antagonist is an anti-PD-1 immunoadhesin.

In a further aspect, the invention provides a kit comprising:

(i) a first container comprising a composition which comprises a T cell activating bispecific antigen binding molecule specific for Folate Receptor 1 (FolR1) and CD3 as described herein; and

5 (ii) a second container comprising a composition comprising a PD-1 axis binding antagonist.

In a further aspect, the invention provides a kit comprising:

(i) a first container comprising a composition which comprises a T cell activating bispecific antigen binding molecule specific for Folate Receptor 1 (FolR1) and CD3 as
10 described herein;

(ii) a second container comprising a composition comprising a PD-1 axis binding antagonist; and

(iii) a third container comprising a composition comprising a TIM3 antagonist.

15 **B. Exemplary T cell activating bispecific antigen binding molecule for use in the invention**

The T cell activating bispecific antigen binding molecule of the invention is bispecific, i.e. it comprises at least two antigen binding moieties capable of specific binding to two distinct antigenic determinants, i.e. to CD3 and to FolR1. According to the invention, the
20 antigen binding moieties are Fab molecules (i.e. antigen binding domains composed of a heavy and a light chain, each comprising a variable and a constant region). In one embodiment said Fab molecules are human. In another embodiment said Fab molecules are humanized. In yet another embodiment said Fab molecules comprise human heavy and light chain constant regions.

25 The T cell activating bispecific antigen binding molecule of the invention is capable of simultaneous binding to the target cell antigen FolR1 and CD3. In one embodiment, the T cell activating bispecific antigen binding molecule is capable of crosslinking a T cell and a FolR1 expressing target cell by simultaneous binding to the target cell antigen FolR1 and CD3. In an even more particular embodiment, such simultaneous binding results in
30 lysis of the FolR1 expressing target cell, particularly a FolR1 expressing tumor cell. In one embodiment, such simultaneous binding results in activation of the T cell. In other embodiments, such simultaneous binding results in a cellular response of a T lymphocyte, particularly a cytotoxic T lymphocyte, selected from the group of: proliferation,

differentiation, cytokine secretion, cytotoxic effector molecule release, cytotoxic activity, and expression of activation markers. In one embodiment, binding of the T cell activating bispecific antigen binding molecule to CD3 without simultaneous binding to the target cell antigen FolR1 does not result in T cell activation.

- 5 In one embodiment, the T cell activating bispecific antigen binding molecule is capable of re-directing cytotoxic activity of a T cell to a FolR1 expressing target cell. In a particular embodiment, said re-direction is independent of MHC-mediated peptide antigen presentation by the target cell and and/or specificity of the T cell.

Particularly, a T cell according to some of the embodiments of the invention is a cytotoxic
10 T cell. In some embodiments the T cell is a CD4⁺ or a CD8⁺ T cell, particularly a CD8⁺ T cell.

The T cell activating bispecific antigen binding molecule of the invention comprises at least one antigen binding moiety capable of binding to CD3 (also referred to herein as an “CD3 antigen binding moiety” or “first antigen binding moiety”). In a particular
15 embodiment, the T cell activating bispecific antigen binding molecule comprises not more than one antigen binding moiety capable of specific binding to CD3. In one embodiment the T cell activating bispecific antigen binding molecule provides monovalent binding to CD3. In a particular embodiment CD3 is human CD3 or cynomolgus CD3, most particularly human CD3. In a particular embodiment the CD3
20 antigen binding moiety is cross-reactive for (i.e. specifically binds to) human and cynomolgus CD3. In some embodiments, the first antigen binding moiety is capable of specific binding to the epsilon subunit of CD3 (see UniProt no. P07766 (version 130), NCBI RefSeq no. NP_000724.1, SEQ ID NO:150 for the human sequence; UniProt no. Q95LI5 (version 49), NCBI GenBank no. BAB71849.1, for the cynomolgus [Macaca
25 fascicularis] sequence).

In some embodiments, the CD3 antigen binding moiety comprises at least one heavy chain complementarity determining region (CDR) selected from the group consisting of SEQ ID NO: 37, SEQ ID NO: 38 and SEQ ID NO: 39 and at least one light chain CDR selected from the group of SEQ ID NO: 32, SEQ ID NO: 33, SEQ ID NO: 34.

- 30 In one embodiment the CD3 antigen binding moiety comprises the heavy chain CDR1 of SEQ ID NO: 37, the heavy chain CDR2 of SEQ ID NO: 38, the heavy chain CDR3 of SEQ ID NO:39, the light chain CDR1 of SEQ ID NO: 32, the light chain CDR2 of SEQ ID NO: 33, and the light chain CDR3 of SEQ ID NO:34.

In one embodiment the CD3 antigen binding moiety comprises a variable heavy chain comprising an amino acid sequence of: SEQ ID NO: 36 and a variable light chain comprising an amino acid sequence of: SEQ ID NO: 31.

5 In one embodiment the CD3 antigen binding moiety comprises a heavy chain variable region sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to SEQ ID NO: 36 and a light chain variable region sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to SEQ ID NO: 31.

The T cell activating bispecific antigen binding molecule of the invention comprises at
10 least one antigen binding moiety capable of binding to the target cell antigen FolR1 (also referred to herein as an "FolR1 binding moiety" or "second" or "third" antigen binding moiety). In one embodiment, the antigen binding moiety capable of binding to the target cell antigen FolR1 does not bind to FolR2 or FolR3. In a particular embodiment the FolR1 antigen binding moiety is cross-reactive for (i.e. specifically binds to) human and
15 cynomolgus FolR1. In certain embodiments, the T cell activating bispecific antigen binding molecule comprises two antigen binding moieties capable of binding to the target cell antigen FolR1. In a particular such embodiment, each of these antigen binding moieties specifically binds to the same antigenic determinant. In an even more particular embodiment, all of these antigen binding moieties are identical. In one embodiment the T
20 cell activating bispecific antigen binding molecule comprises not more than two antigen binding moieties capable of binding to FolR1.

The FolR1 binding moiety is generally a Fab molecule that specifically binds to FolR1 and is able to direct the T cell activating bispecific antigen binding molecule to which it is connected to a target site, for example to a specific type of tumor cell that expresses
25 FolR1.

In one aspect the present invention provides a T cell activating bispecific antigen binding molecule comprising

- (i) a first antigen binding moiety which is a Fab molecule capable of specific
30 binding to CD3, and which comprises at least one heavy chain complementarity determining region (CDR) selected from the group consisting SEQ ID NO: 37, SEQ ID NO: 38 and SEQ ID NO: 39 and at least one light chain CDR selected from the group of SEQ ID NO: 32, SEQ ID NO: 33, SEQ ID NO: 34; and

- (ii) a second antigen binding moiety which is a Fab molecule capable of specific binding to Folate Receptor 1 (FolR1).

In one embodiment the first antigen binding moiety which is a Fab molecule capable of specific binding to CD3 comprises a variable heavy chain comprising an amino acid
5 sequence of SEQ ID NO: 36 and a variable light chain comprising an amino acid sequence of SEQ ID NO: 31.

In one embodiment the T cell activating bispecific antigen binding molecule additionally comprises

- 10 (iii) a third antigen binding moiety which is a Fab molecule capable of specific binding to FolR1.

In one such embodiment the second and third antigen binding moiety capable of specific binding to FolR1 comprise identical heavy chain complementarity determining region
15 (CDR) and light chain CDR sequences. In one such embodiment the third antigen binding moiety is identical to the second antigen binding moiety.

In one embodiment the T cell activating bispecific antigen binding molecule of any of the above embodiments additionally comprises an Fc domain composed of a first and a
20 second subunit capable of stable association.

In one embodiment the first antigen binding moiety and the second antigen binding moiety are each fused at the C-terminus of the Fab heavy chain to the N-terminus of the first or second subunit of the Fc domain.

25 In one embodiment the third antigen binding moiety is fused at the C-terminus of the Fab heavy chain to the N-terminus of the Fab heavy chain of the first antigen binding moiety, optionally via a peptide linker.

In a further particular embodiment, not more than one antigen binding moiety capable of specific binding to CD3 is present in the T cell activating bispecific antigen binding
30 molecule (i.e. the T cell activating bispecific antigen binding molecule provides monovalent binding to CD3).

T cell activating bispecific antigen binding molecule with a common light chain

The inventors of the present invention generated a bispecific antibody wherein the binding moieties share a common light chain that retains the specificity and efficacy of the parent monospecific antibody for CD3 and can bind a second antigen (e.g., FolR1) using the same light chain. The generation of a bispecific molecule with a common light chain that retains the binding properties of the parent antibody is not straight-forward as the common CDRs of the hybrid light chain have to effectuate the binding specificity for both targets. In one aspect the present invention provides a T cell activating bispecific antigen binding molecule comprising a first and a second antigen binding moiety, one of which is a Fab molecule capable of specific binding to CD3 and the other one of which is a Fab molecule capable of specific binding to FolR1, wherein the first and the second Fab molecule have identical VLCL light chains. In one embodiment said identical light chain (VLCL) comprises the light chain CDRs of SEQ ID NO: 32, SEQ ID NO: 33 and SEQ ID NO: 34. In one embodiment said identical light chain (VLCL) comprises SEQ ID NO: 35.

15

In one embodiment the present invention provides a T cell activating bispecific antigen binding molecule comprising

(i) a first antigen binding moiety which is a Fab molecule capable of specific binding to CD3, and which comprises at least one heavy chain complementarity determining region (CDR) selected from the group consisting of SEQ ID NO: 37, SEQ ID NO: 38 and SEQ ID NO: 39 and at least one light chain CDR selected from the group of SEQ ID NO: 32, SEQ ID NO: 33, SEQ ID NO: 34;

(ii) a second antigen binding moiety which is a Fab molecule capable of specific binding to Folate Receptor 1 (FolR1) and which comprises at least one heavy chain complementarity determining region (CDR) selected from the group consisting of SEQ ID NO: 16, SEQ ID NO: 17 and SEQ ID NO: 18 and at least one light chain CDR selected from the group of SEQ ID NO: 32, SEQ ID NO: 33, SEQ ID NO: 34.

In one such embodiment the CD3 antigen binding moiety comprises the heavy chain CDR1 of SEQ ID NO: 37, the heavy chain CDR2 of SEQ ID NO: 38, the heavy chain CDR3 of SEQ ID NO: 39, the light chain CDR1 of SEQ ID NO: 32, the light chain CDR2 of SEQ ID NO: 33, and the light chain CDR3 of SEQ ID NO: 34 and the FolR1 antigen

30

binding moiety comprises the heavy chain CDR1 of SEQ ID NO: 16, the heavy chain CDR2 of SEQ ID NO: 17, the heavy chain CDR3 of SEQ ID NO:18, the light chain CDR1 of SEQ ID NO: 32, the light chain CDR2 of SEQ ID NO: 33, and the light chain CDR3 of SEQ ID NO:34.

5

In one embodiment the present invention provides a T cell activating bispecific antigen binding molecule comprising

- (i) a first antigen binding moiety which is a Fab molecule capable of specific binding to CD3 comprising a variable heavy chain comprising an amino acid sequence of SEQ ID NO: 36 and a variable light chain comprising an amino acid sequence of SEQ ID NO: 31.
- (ii) a second antigen binding moiety which is a Fab molecule capable of specific binding to Folate Receptor 1 (FolR1) comprising a variable heavy chain comprising an amino acid sequence of SEQ ID NO: 15 and a variable light chain comprising an amino acid sequence of SEQ ID NO: 31.

15

In a further embodiment, the antigen binding moiety that is specific for FolR1 comprises a heavy chain variable region sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to SEQ ID NO:15 and a light chain variable region sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to SEQ ID NO: 31 or variants thereof that retain functionality.

In one embodiment the T cell activating bispecific antigen binding molecule comprises a polypeptide sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to SEQ ID NO: 36, a polypeptide sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to SEQ ID NO:15, and a polypeptide sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to SEQ ID NO: 31.

In one embodiment the T cell activating bispecific antigen binding molecule additionally comprises

- (iii) a third antigen binding moiety (which is a Fab molecule) capable of specific binding to FolR1.

30

In one such embodiment the second and third antigen binding moiety capable of specific binding to FolR1 comprise identical heavy chain complementarity determining region

(CDR) and light chain CDR sequences. In one such embodiment the third antigen binding moiety is identical to the second antigen binding moiety.

Hence in one embodiment the present invention provides a T cell activating bispecific
5 antigen binding molecule comprising

- (i) a first antigen binding moiety which is a Fab molecule capable of specific binding to CD3, and which comprises at least one heavy chain complementarity determining region (CDR) selected from the group consisting of SEQ ID NO: 37, SEQ ID NO: 38 and SEQ ID NO: 39 and at least one light chain CDR selected from the group of SEQ ID NO: 32,
10 SEQ ID NO: 33, SEQ ID NO: 34;
- (ii) a second antigen binding moiety which is a Fab molecule capable of specific binding to Folate Receptor 1 (FolR1) and which comprises at least one heavy chain complementarity determining region (CDR) selected from the group consisting of SEQ ID NO: 16, SEQ ID NO: 17 and SEQ ID NO: 18 and at least one light chain CDR
15 selected from the group of SEQ ID NO: 32, SEQ ID NO: 33, SEQ ID NO: 34.
- (iii) a third antigen binding moiety which is a Fab molecule capable of specific binding to Folate Receptor 1 (FolR1) and which comprises at least one heavy chain complementarity determining region (CDR) selected from the group consisting of SEQ ID NO: 16, SEQ ID NO: 17 and SEQ ID NO: 18 and at least one light chain CDR selected from the group of
20 SEQ ID NO: 32, SEQ ID NO: 33, SEQ ID NO: 34.

In one such embodiment the CD3 antigen binding moiety comprises the heavy chain CDR1 of SEQ ID NO: 37, the heavy chain CDR2 of SEQ ID NO: 38, the heavy chain CDR3 of SEQ ID NO: 39, the light chain CDR1 of SEQ ID NO: 32, the light chain CDR2
25 of SEQ ID NO: 33, and the light chain CDR3 of SEQ ID NO: 34 and the FolR1 antigen binding moiety comprises the heavy chain CDR1 of SEQ ID NO: 16, the heavy chain CDR2 of SEQ ID NO: 17, the heavy chain CDR3 of SEQ ID NO: 18, the light chain CDR1 of SEQ ID NO: 32, the light chain CDR2 of SEQ ID NO: 33, and the light chain CDR3 of SEQ ID NO: 34.

30

In one embodiment the present invention provides a T cell activating bispecific antigen binding molecule comprising

- (i) a first antigen binding moiety which is a Fab molecule capable of specific binding to CD3 comprising a variable heavy chain comprising an amino acid sequence of SEQ ID NO: 36 and a variable light chain comprising an amino acid sequence of SEQ ID NO: 31.
- (ii) a second antigen binding moiety which is a Fab molecule capable of specific binding to Folate Receptor 1 (FolR1) comprising a variable heavy chain comprising an amino acid sequence of SEQ ID NO: 15 and a variable light chain comprising an amino acid sequence of SEQ ID NO: 31.
- (iii) a third antigen binding moiety which is a Fab molecule capable of specific binding to Folate Receptor 1 (FolR1) comprising a variable heavy chain comprising an amino acid sequence of SEQ ID NO: 15 and a variable light chain comprising an amino acid sequence of SEQ ID NO: 31.

In one embodiment the present invention provides a T cell activating bispecific antigen binding molecule comprising

- (i) a first antigen binding moiety which is a Fab molecule capable of specific binding to CD3, and which comprises at least one heavy chain complementarity determining region (CDR) selected from the group consisting of SEQ ID NO: 37, SEQ ID NO: 38 and SEQ ID NO: 39 and at least one light chain CDR selected from the group of SEQ ID NO: 32, SEQ ID NO: 33, SEQ ID NO: 34;
- (ii) a second antigen binding moiety which is a Fab molecule capable of specific binding to Folate Receptor 1 (FolR1) and which comprises at least one heavy chain complementarity determining region (CDR) selected from the group consisting of SEQ ID NO:16, SEQ ID NO:402 and SEQ ID NO:400 and at least one light chain CDR selected from the group of SEQ ID NO: 32, SEQ ID NO: 33, SEQ ID NO: 34.

25

- In one such embodiment the CD3 antigen binding moiety comprises the heavy chain CDR1 of SEQ ID NO: 37, the heavy chain CDR2 of SEQ ID NO: 38, the heavy chain CDR3 of SEQ ID NO:39, the light chain CDR1 of SEQ ID NO: 32, the light chain CDR2 of SEQ ID NO: 33, and the light chain CDR3 of SEQ ID NO:34, and the FolR1 antigen binding moiety comprises the heavy chain CDR1 of SEQ ID NO:16, the heavy chain CDR2 of SEQ ID NO:402, the heavy chain CDR3 of SEQ ID NO:400, the light chain CDR1 of SEQ ID NO: 32, the light chain CDR2 of SEQ ID NO: 33, and the light chain CDR3 of SEQ ID NO:34.

30

In one embodiment the present invention provides a T cell activating bispecific antigen binding molecule comprising

- (i) a first antigen binding moiety which is a Fab molecule capable of specific binding to CD3 comprising a variable heavy chain comprising an amino acid sequence of SEQ ID NO: 36 and a variable light chain comprising an amino acid sequence of SEQ ID NO: 31.
- (ii) a second antigen binding moiety which is a Fab molecule capable of specific binding to Folate Receptor 1 (FolR1) comprising a variable heavy chain comprising an amino acid sequence of SEQ ID NO:401 and a variable light chain comprising an amino acid sequence of SEQ ID NO: 31.

In a further embodiment, the antigen binding moiety that is specific for FolR1 comprises a heavy chain variable region sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to SEQ ID NO:401 and a light chain variable region sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to SEQ ID NO: 31 or variants thereof that retain functionality.

In one embodiment the T cell activating bispecific antigen binding molecule comprises a polypeptide sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to SEQ ID NO: 36, a polypeptide sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to SEQ ID NO:401, and a polypeptide sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to SEQ ID NO: 31.

In one embodiment the T cell activating bispecific antigen binding molecule additionally comprises

- (iii) a third antigen binding moiety (which is a Fab molecule) capable of specific binding to FolR1.

In one such embodiment the second and third antigen binding moiety capable of specific binding to FolR1 comprise identical heavy chain complementarity determining region (CDR) and light chain CDR sequences. In one such embodiment the third antigen binding moiety is identical to the second antigen binding moiety.

Hence in one embodiment the present invention provides a T cell activating bispecific antigen binding molecule comprising

- (i) a first antigen binding moiety which is a Fab molecule capable of specific binding to CD3, and which comprises at least one heavy chain complementarity determining region (CDR) selected from the group consisting of SEQ ID NO: 37, SEQ ID NO: 38 and SEQ ID NO: 39 and at least one light chain CDR selected from the group of SEQ ID NO: 32, SEQ ID NO: 33, SEQ ID NO: 34;
- (ii) a second antigen binding moiety which is a Fab molecule capable of specific binding to Folate Receptor 1 (FolR1) and which comprises at least one heavy chain complementarity determining region (CDR) selected from the group consisting of SEQ ID NO:16, SEQ ID NO:402 and SEQ ID NO:400 and at least one light chain CDR selected from the group of SEQ ID NO: 32, SEQ ID NO: 33, SEQ ID NO: 34.
- (iii) a third antigen binding moiety which is a Fab molecule capable of specific binding to Folate Receptor 1 (FolR1) and which comprises at least one heavy chain complementarity determining region (CDR) selected from the group consisting of SEQ ID NO:16, SEQ ID NO:402 and SEQ ID NO:400 and at least one light chain CDR selected from the group of SEQ ID NO: 32, SEQ ID NO: 33, SEQ ID NO: 34.

In one such embodiment the CD3 antigen binding moiety comprises the heavy chain CDR1 of SEQ ID NO: 37, the heavy chain CDR2 of SEQ ID NO: 38, the heavy chain CDR3 of SEQ ID NO:39, the light chain CDR1 of SEQ ID NO: 32, the light chain CDR2 of SEQ ID NO: 33, and the light chain CDR3 of SEQ ID NO:34 and the FolR1 antigen binding moiety comprises the heavy chain CDR1 of SEQ ID NO:16, the heavy chain CDR2 of SEQ ID NO:402, the heavy chain CDR3 of SEQ ID NO:400, the light chain CDR1 of SEQ ID NO: 32, the light chain CDR2 of SEQ ID NO: 33, and the light chain CDR3 of SEQ ID NO:34.

In one embodiment the present invention provides a T cell activating bispecific antigen binding molecule comprising

- (i) a first antigen binding moiety which is a Fab molecule capable of specific binding to CD3 comprising a variable heavy chain comprising an amino acid sequence of SEQ ID NO: 36 and a variable light chain comprising an amino acid sequence of SEQ ID NO: 31.

- (ii) a second antigen binding moiety which is a Fab molecule capable of specific binding to Folate Receptor 1 (FolR1) comprising a variable heavy chain comprising an amino acid sequence of SEQ ID NO:401 and a variable light chain comprising an amino acid sequence of SEQ ID NO: 31.
- 5 (iii) a third antigen binding moiety which is a Fab molecule capable of specific binding to Folate Receptor 1 (FolR1) comprising a variable heavy chain comprising an amino acid sequence of SEQ ID NO:401 and a variable light chain comprising an amino acid sequence of SEQ ID NO: 31.
- 10 Thus, in one embodiment, the invention relates to bispecific molecules wherein at least two binding moieties have identical light chains and corresponding remodeled heavy chains that confer the specific binding to the T cell activating antigen CD3 and the target cell antigen FolR1, respectively. The use of this so-called 'common light chain' principle, i.e. combining two binders that share one light chain but still have separate specificities,
- 15 prevents light chain mispairing. Thus, there are less side products during production, facilitating the homogenous preparation of T cell activating bispecific antigen binding molecules.

The components of the T cell activating bispecific antigen binding molecule can be fused
20 to each other in a variety of configurations. Exemplary configurations are depicted in Figures 1A-I and are further described below.

In some embodiments, said T cell activating bispecific antigen binding molecule further comprises an Fc domain composed of a first and a second subunit capable of stable association. Below exemplary embodiments of T cell activating bispecific antigen binding
25 molecule comprising an Fc domain are described.

T cell activating bispecific antigen binding molecule with a crossover Fab fragment

The inventors of the present invention generated a second bispecific antibody format wherein one of the binding moieties is a crossover Fab fragment. In one aspect of the invention a monovalent bispecific antibody is provided, wherein one of the Fab fragments
30 of an IgG molecule is replaced by a crossover Fab fragment. Crossover Fab fragments are Fab fragments wherein either the variable regions or the constant regions of the heavy and

- light chain are exchanged. Bispecific antibody formats comprising crossover Fab fragments have been described, for example, in WO2009080252, WO2009080253, WO2009080251, WO2009080254, WO2010/136172, WO2010/145792 and WO2013/026831. In a particular embodiment, the first antigen binding moiety is a crossover Fab molecule wherein either the variable or the constant regions of the Fab light chain and the Fab heavy chain are exchanged. Such modification prevent mispairing of heavy and light chains from different Fab molecules, thereby improving the yield and purity of the T cell activating bispecific antigen binding molecule of the invention in recombinant production. In a particular crossover Fab molecule useful for the T cell activating bispecific antigen binding molecule of the invention, the variable regions of the Fab light chain and the Fab heavy chain are exchanged. In another crossover Fab molecule useful for the T cell activating bispecific antigen binding molecule of the invention, the constant regions of the Fab light chain and the Fab heavy chain are exchanged.
- 15 In one embodiment the T cell activating bispecific antigen binding molecule comprises
- (i) a first antigen binding moiety which is a crossover Fab molecule capable of specific binding to CD3, comprising at least one heavy chain complementarity determining region (CDR) selected from the group consisting of SEQ ID NO: 37, SEQ ID NO: 38 and SEQ ID NO: 39 and at least one light chain CDR selected from the group of SEQ ID NO: 32, SEQ ID NO: 33, SEQ ID NO: 34;
- 20 (ii) a second antigen binding moiety capable of specific binding to Folate Receptor 1 (FolR1) comprising at least one heavy chain complementarity determining region (CDR) selected from the group consisting of SEQ ID NO: 8, SEQ ID NO: 56 and SEQ ID NO: 57 and at least one light chain CDR selected from the group of SEQ ID NO: 59, SEQ ID NO: 60, SEQ ID NO: 65.
- 25

In one such embodiment the CD3 antigen binding moiety comprises the heavy chain CDR1 of SEQ ID NO: 37, the heavy chain CDR2 of SEQ ID NO: 38, the heavy chain CDR3 of SEQ ID NO:39, the light chain CDR1 of SEQ ID NO: 32, the light chain CDR2 of SEQ ID NO: 33, and the light chain CDR3 of SEQ ID NO:34 and the FolR1 antigen binding moiety comprises the heavy chain CDR1 of SEQ ID NO: 8, the heavy chain CDR2 of SEQ ID NO: 56, the heavy chain CDR3 of SEQ ID NO:57, the light chain

30

CDR1 of SEQ ID NO: 59, the light chain CDR2 of SEQ ID NO: 60, and the light chain CDR3 of SEQ ID NO: 65.

In one embodiment, the second antigen binding moiety is a conventional Fab molecule.

5

In one embodiment the T cell activating bispecific antigen binding molecule comprises

(i) a first antigen binding moiety which is a crossover Fab molecule capable of specific binding to CD3 comprising a variable heavy chain comprising an amino acid sequence of SEQ ID NO: 36 and a variable light chain comprising an amino acid sequence of SEQ ID NO: 31.

10

(ii) a second antigen binding moiety which is a Fab molecule capable of specific binding to Folate Receptor 1 (FolR1) comprising a variable heavy chain comprising an amino acid sequence of SEQ ID NO: 55 and a variable light chain comprising an amino acid sequence of SEQ ID NO: 64.

15 In one embodiment, the second antigen binding moiety is a conventional Fab molecule.

In a further embodiment, the antigen binding moiety that is specific for FolR1 comprises a heavy chain variable region sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to SEQ ID NO: 55 and a light chain variable region sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to SEQ ID NO: 64 or variants thereof that retain functionality.

20

In one embodiment the T cell activating bispecific antigen binding molecule comprises a polypeptide sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to SEQ ID NO: 36, a polypeptide sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to SEQ ID NO: 31, a polypeptide sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to SEQ ID NO: 55, and a polypeptide sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to SEQ ID NO: 64.

25

30 In one embodiment the T cell activating bispecific antigen binding molecule additionally comprises

(iii) a third antigen binding moiety capable of specific binding to FolR1.

In one embodiment, the third antigen binding moiety is a conventional Fab molecule. In one embodiment, the third antigen binding moiety is a crossover Fab molecule.

- 5 In one such embodiment the second and third antigen binding moiety capable of specific binding to FolR1 comprise identical heavy chain complementarity determining region (CDR) and light chain CDR sequences. In one such embodiment the third antigen binding moiety is identical to the second antigen binding moiety.
- 10 In one embodiment the T cell activating bispecific antigen binding molecule comprises
- (i) a first antigen binding moiety which is a crossover Fab molecule capable of specific binding to CD3, comprising at least one heavy chain complementarity determining region (CDR) selected from the group consisting of SEQ ID NO: 37, SEQ ID NO: 38 and SEQ ID NO: 39 and at least one light chain CDR selected from the group of SEQ ID NO: 32, 15 SEQ ID NO: 33, SEQ ID NO: 34;
- (ii) a second antigen binding moiety capable of specific binding to Folate Receptor 1 (FolR1) comprising at least one heavy chain complementarity determining region (CDR) selected from the group consisting of SEQ ID NO: 8, SEQ ID NO: 56 and SEQ ID NO: 57 and at least one light chain CDR selected from the group of SEQ ID NO: 59, SEQ ID 20 NO: 60, SEQ ID NO: 65.
- (iii) a third antigen binding moiety capable of specific binding to Folate Receptor 1 (FolR1) comprising at least one heavy chain complementarity determining region (CDR) selected from the group consisting of SEQ ID NO: 8, SEQ ID NO: 56 and SEQ ID NO: 57 and at least one light chain CDR selected from the group of SEQ ID NO: 59, SEQ ID 25 NO: 60, SEQ ID NO: 65.

In one such embodiment the CD3 antigen binding moiety comprises the heavy chain CDR1 of SEQ ID NO: 37, the heavy chain CDR2 of SEQ ID NO: 38, the heavy chain CDR3 of SEQ ID NO: 39, the light chain CDR1 of SEQ ID NO: 32, the light chain CDR2 of SEQ ID NO: 33, and the light chain CDR3 of SEQ ID NO: 34 and the FolR1 antigen 30 binding moiety comprises the heavy chain CDR1 of SEQ ID NO: 8, the heavy chain CDR2 of SEQ ID NO: 56, the heavy chain CDR3 of SEQ ID NO: 57, the light chain

CDR1 of SEQ ID NO: 59, the light chain CDR2 of SEQ ID NO: 60, and the light chain CDR3 of SEQ ID NO: 65.

5 In one embodiment, the second antigen binding moiety and the third antigen binding moiety are both a conventional Fab molecule.

In one embodiment the T cell activating bispecific antigen binding molecule comprises
(i) a first antigen binding moiety which is a crossover Fab molecule capable of specific binding to CD3 comprising a variable heavy chain comprising an amino acid sequence of
10 SEQ ID NO: 36 and a variable light chain comprising an amino acid sequence of SEQ ID NO: 31.

(ii) a second antigen binding moiety which is a Fab molecule capable of specific binding to Folate Receptor 1 (FolR1) comprising a variable heavy chain comprising an amino acid sequence of SEQ ID NO: 55 and a variable light chain comprising an amino acid
15 sequence of SEQ ID NO: 64.

(iii) a third antigen binding moiety which is a Fab molecule capable of specific binding to Folate Receptor 1 (FolR1) comprising a variable heavy chain comprising an amino acid sequence of SEQ ID NO: 55 and a variable light chain comprising an amino acid sequence of SEQ ID NO: 64.

20 In one embodiment, the second antigen binding moiety and the third antigen binding moiety are both a conventional Fab molecule.

In one embodiment the T cell activating bispecific antigen binding molecule comprises

(i) a first antigen binding moiety which is a crossover Fab molecule capable of specific binding to CD3, comprising at least one heavy chain complementarity determining region
25 (CDR) selected from the group consisting of SEQ ID NO: 37, SEQ ID NO: 38 and SEQ ID NO: 39 and at least one light chain CDR selected from the group of SEQ ID NO: 32, SEQ ID NO: 33, SEQ ID NO: 34;

(ii) a second antigen binding moiety capable of specific binding to Folate Receptor 1 (FolR1) comprising at least one heavy chain complementarity determining region (CDR)
30 selected from the group consisting of SEQ ID NO: 8, SEQ ID NO: 9 and SEQ ID NO: 50

and at least one light chain CDR selected from the group of SEQ ID NO: 52, SEQ ID NO: 53, SEQ ID NO: 54.

In one such embodiment the CD3 antigen binding moiety comprises the heavy chain CDR1 of SEQ ID NO: 37, the heavy chain CDR2 of SEQ ID NO: 38, the heavy chain CDR3 of SEQ ID NO:39, the light chain CDR1 of SEQ ID NO: 32, the light chain CDR2 of SEQ ID NO: 33, and the light chain CDR3 of SEQ ID NO:34 and the FolR1 antigen binding moiety comprises the heavy chain CDR1 of SEQ ID NO: 8, the heavy chain CDR2 of SEQ ID NO: 9, the heavy chain CDR3 of SEQ ID NO:50, the light chain CDR1 of SEQ ID NO: 52, the light chain CDR2 of SEQ ID NO: 53, and the light chain CDR3 of SEQ ID NO:54.

In one embodiment, the second antigen binding moiety is a conventional Fab molecule.
In one embodiment, the second antigen binding moiety is a crossover Fab molecule.

- 15 In one embodiment the T cell activating bispecific antigen binding molecule comprises
- (i) a first antigen binding moiety which is a crossover Fab molecule capable of specific binding to CD3 comprising a variable heavy chain comprising an amino acid sequence of SEQ ID NO: 36 and a variable light chain comprising an amino acid sequence of SEQ ID NO: 31.
 - 20 (ii) a second antigen binding moiety which is a Fab molecule capable of specific binding to Folate Receptor 1 (FolR1) comprising a variable heavy chain comprising an amino acid sequence of SEQ ID NO: 49 and a variable light chain comprising an amino acid sequence of SEQ ID NO: 51.

In one embodiment, the second antigen binding moiety is a conventional Fab molecule.

- 25 In one embodiment, the second antigen binding moiety is a crossover Fab molecule.

In a further embodiment, the antigen binding moiety that is specific for FolR1 comprises a heavy chain variable region sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to SEQ ID NO:49 and a light chain variable region sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to SEQ ID NO: 51 or variants thereof that retain functionality.

In one embodiment the T cell activating bispecific antigen binding molecule comprises a polypeptide sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to SEQ ID NO: 36, a polypeptide sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to SEQ ID NO: 31, a polypeptide sequence that is at least about 5 95%, 96%, 97%, 98%, 99% or 100% identical to SEQ ID NO:49, and a polypeptide sequence that is at least about 95%, 96%, 97%, 98%, 99% or 100% identical to SEQ ID NO: 51.

In one embodiment the T cell activating bispecific antigen binding molecule additionally 10 comprises

- (iii) a third antigen binding moiety capable of specific binding to FolR1.

In one embodiment, the third antigen binding moiety is a conventional Fab molecule. In one embodiment, the second antigen binding moiety is a crossover Fab molecule. 15

In one such embodiment the second and third antigen binding moiety capable of specific binding to FolR1 comprise identical heavy chain complementarity determining region (CDR) and light chain CDR sequences. In one such embodiment the third antigen binding moiety is identical to the second antigen binding moiety. 20

In one embodiment the T cell activating bispecific antigen binding molecule comprises

- (i) a first antigen binding moiety which is a crossover Fab molecule capable of specific binding to CD3, comprising at least one heavy chain complementarity determining region (CDR) selected from the group consisting of SEQ ID NO: 37, SEQ ID NO: 38 and SEQ 25 ID NO: 39 and at least one light chain CDR selected from the group of SEQ ID NO: 32, SEQ ID NO: 33, SEQ ID NO: 34;
- (ii) a second antigen binding moiety capable of specific binding to Folate Receptor 1 (FolR1) comprising at least one heavy chain complementarity determining region (CDR) selected from the group consisting of SEQ ID NO: 8, SEQ ID NO: 9 and SEQ ID NO: 49 30 and at least one light chain CDR selected from the group of SEQ ID NO: 52, SEQ ID NO: 53, SEQ ID NO: 54.

(iii) a third antigen binding moiety capable of specific binding to Folate Receptor 1 (FolR1) comprising at least one heavy chain complementarity determining region (CDR) selected from the group consisting of SEQ ID NO: 8, SEQ ID NO: 9 and SEQ ID NO: 50 and at least one light chain CDR selected from the group of SEQ ID NO: 52, SEQ ID NO: 53, SEQ ID NO: 54.

In one such embodiment the CD3 antigen binding moiety comprises the heavy chain CDR1 of SEQ ID NO: 37, the heavy chain CDR2 of SEQ ID NO: 38, the heavy chain CDR3 of SEQ ID NO:39, the light chain CDR1 of SEQ ID NO: 32, the light chain CDR2 of SEQ ID NO: 33, and the light chain CDR3 of SEQ ID NO:34 and the FolR1 antigen binding moiety comprises the heavy chain CDR1 of SEQ ID NO: 8, the heavy chain CDR2 of SEQ ID NO: 9, the heavy chain CDR3 of SEQ ID NO:50, the light chain CDR1 of SEQ ID NO: 52, the light chain CDR2 of SEQ ID NO: 53, and the light chain CDR3 of SEQ ID NO:54.

In one embodiment, the second antigen binding moiety and the third antigen binding moiety are both a conventional Fab molecule.

In one embodiment the T cell activating bispecific antigen binding molecule comprises (i) a first antigen binding moiety which is a crossover Fab molecule capable of specific binding to CD3 comprising a variable heavy chain comprising an amino acid sequence of SEQ ID NO: 36 and a variable light chain comprising an amino acid sequence of SEQ ID NO: 31.

(ii) a second antigen binding moiety which is a Fab molecule capable of specific binding to Folate Receptor 1 (FolR1) comprising a variable heavy chain comprising an amino acid sequence of SEQ ID NO: 49 and a variable light chain comprising an amino acid sequence of SEQ ID NO: 51.

(iii) a third antigen binding moiety which is a Fab molecule capable of specific binding to Folate Receptor 1 (FolR1) comprising a variable heavy chain comprising an amino acid sequence of SEQ ID NO: 49 and a variable light chain comprising an amino acid sequence of SEQ ID NO: 51.

In one embodiment, the second antigen binding moiety and the third antigen binding moiety are both a conventional Fab molecule.

Thus, in one embodiment, the invention relates to bispecific molecules wherein two binding moieties confer specific binding to FolR1 and one binding moiety confers
5 specificity to the T cell activating antigen CD3. One of the heavy chains is modified to ensure proper pairing of the heavy and light chains, thus eliminating the need for a common light chain approach. The presence of two FolR1 binding sites enables appropriate engagement with the target antigen FolR1 and the activation of T cells.

The components of the T cell activating bispecific antigen binding molecule can be fused
10 to each other in a variety of configurations. Exemplary configurations are depicted in Figures 1A-I and are further described below.

In some embodiments, said T cell activating bispecific antigen binding molecule further comprises an Fc domain composed of a first and a second subunit capable of stable association. Below exemplary embodiments of T cell activating bispecific antigen binding
15 molecule comprising an Fc domain are described.

T cell activating bispecific antigen binding molecule formats

As depicted above and in Figures 1A-I, in one embodiment the T cell activating bispecific antigen binding molecules comprise at least two Fab fragments having identical light chains (VLCL) and having different heavy chains (VHCL) which confer the specificities
20 to two different antigens, i.e. one Fab fragment is capable of specific binding to a T cell activating antigen CD3 and the other Fab fragment is capable of specific binding to the target cell antigen FolR1.

In another embodiment the T cell activating bispecific antigen binding molecule comprises at least two antigen binding moieties (Fab molecules), one of which is a
25 crossover Fab molecule and one of which is a conventional Fab molecule. In one such embodiment the first antigen binding moiety capable of specific binding to CD3 is a crossover Fab molecule and the second antigen binding moiety capable of specific binding to FolR is a conventional Fab molecule.

These components of the T cell activating bispecific antigen binding molecule can be fused to each other in a variety of configurations. Exemplary configurations are depicted in Figures 1A-I.

In some embodiments, the first and second antigen binding moiety are each fused at the C-terminus of the Fab heavy chain to the N-terminus of the first or the second subunit of the Fc domain. In a specific such embodiment, the T cell activating bispecific antigen binding molecule essentially consists of a first and a second antigen binding moiety, an Fc domain composed of a first and a second subunit, and optionally one or more peptide linkers, wherein the first and second antigen binding moiety are each fused at the C-terminus of the Fab heavy chain to the N-terminus of the first or the second subunit of the Fc domain. In one such embodiment the first and second antigen binding moiety both are Fab fragments and have identical light chains (VLCL). In another such embodiment the first antigen binding moiety capable of specific binding to CD3 is a crossover Fab molecule and the second antigen binding moiety capable of specific binding to FolR is a conventional Fab molecule.

In one embodiment, the second antigen binding moiety is fused at the C-terminus of the Fab heavy chain to the N-terminus of the first or the second subunit of the Fc domain and the first antigen binding moiety is fused at the C-terminus of the Fab heavy chain to the N-terminus of the Fab heavy chain of the second antigen binding moiety. In a specific such embodiment, the T cell activating bispecific antigen binding molecule essentially consists of a first and a second antigen binding moiety, an Fc domain composed of a first and a second subunit, and optionally one or more peptide linkers, wherein the first antigen binding moiety is fused at the C-terminus of the Fab heavy chain to the N-terminus of the Fab heavy chain of the second antigen binding moiety, and the second antigen binding moiety is fused at the C-terminus of the Fab heavy chain to the N-terminus of the first or the second subunit of the Fc domain. In one such embodiment the first and second antigen binding moiety both are Fab fragments and have identical light chains (VLCL). In another such embodiment the first antigen binding moiety capable of specific binding to CD3 is a crossover Fab molecule and the second antigen binding moiety capable of specific binding to FolR is a conventional Fab molecule. Optionally, the Fab light chain of the first antigen binding moiety and the Fab light chain of the second antigen binding moiety may additionally be fused to each other.

In other embodiments, the first antigen binding moiety is fused at the C-terminus of the Fab heavy chain to the N-terminus of the first or second subunit of the Fc domain. In a particular such embodiment, the second antigen binding moiety is fused at the C-terminus of the Fab heavy chain to the N-terminus of the Fab heavy chain of the first antigen binding moiety. In a specific such embodiment, the T cell activating bispecific antigen binding molecule essentially consists of a first and a second antigen binding moiety, an Fc domain composed of a first and a second subunit, and optionally one or more peptide linkers, wherein the second antigen binding moiety is fused at the C-terminus of the Fab heavy chain to the N-terminus of the Fab heavy chain of the first antigen binding moiety, and the first antigen binding moiety is fused at the C-terminus of the Fab heavy chain to the N-terminus of the first or the second subunit of the Fc domain. In one such embodiment the first and second antigen binding moiety both are Fab fragments and have identical light chains (VLCL). In another such embodiment the first antigen binding moiety capable of specific binding to CD3 is a crossover Fab molecule and the second antigen binding moiety capable of specific binding to FolR is a conventional Fab molecule. Optionally, the Fab light chain of the first antigen binding moiety and the Fab light chain of the second antigen binding moiety may additionally be fused to each other.

The antigen binding moieties may be fused to the Fc domain or to each other directly or through a peptide linker, comprising one or more amino acids, typically about 2-20 amino acids. Peptide linkers are known in the art and are described herein. Suitable, non-immunogenic peptide linkers include, for example, $(G_4S)_n$ (SEQ ID NO: 387), $(SG_4)_n$ (SEQ ID NO: 388), $(G_4S)_n$ (SEQ ID NO: 387) or $G_4(SG_4)_n$ (SEQ ID NO: 389) peptide linkers. "n" is generally a number between 1 and 10, typically between 2 and 4. A particularly suitable peptide linker for fusing the Fab light chains of the first and the second antigen binding moiety to each other is $(G_4S)_2$ (SEQ ID NO: 386). An exemplary peptide linker suitable for connecting the Fab heavy chains of the first and the second antigen binding moiety is EPKSC(D)- $(G_4S)_2$ (SEQ ID NOS 390 and 391). Additionally, linkers may comprise (a portion of) an immunoglobulin hinge region. Particularly where an antigen binding moiety is fused to the N-terminus of an Fc domain subunit, it may be fused via an immunoglobulin hinge region or a portion thereof, with or without an additional peptide linker.

It has been found by the inventors of the present invention that T cell activating bispecific antigen binding molecule comprising two binding moieties specific for the target cell antigen FolR have superior characteristics compared to T cell activating bispecific antigen binding molecule comprising only one binding moiety specific for the target cell antigen FolR.

Accordingly, in certain embodiments, the T cell activating bispecific antigen binding molecule of the invention further comprises a third antigen binding moiety which is a Fab molecule capable of specific binding to FolR. In one such embodiment the second and third antigen binding moiety capable of specific binding to FolR1 comprise identical heavy chain complementarity determining region (CDR) and light chain CDR sequences, i.e., the heavy chain CDR sequences of the second antigen binding moiety are the same as the heavy chain CDR sequences of the third antigen binding moiety, and the light chain CDR sequences of the second antigen binding moiety are the same as the light chain CDR sequences of the third antigen binding moiety. In one such embodiment the third antigen binding moiety is identical to the second antigen binding moiety (i.e. they comprise the same amino acid sequences).

In one embodiment, the first and second antigen binding moiety are each fused at the C-terminus of the Fab heavy chain to the N-terminus of the first or second subunit of the Fc domain and the third antigen binding moiety is fused at the C-terminus of the Fab heavy chain to the N-terminus of the Fab heavy chain of the first antigen binding moiety. In a specific such embodiment, the T cell activating bispecific antigen binding molecule essentially consists of a first, a second and a third antigen binding moiety, an Fc domain composed of a first and a second subunit, and optionally one or more peptide linkers, wherein the first and second antigen binding moiety are each fused at the C-terminus of the Fab heavy chain to the N-terminus of the first subunit of the Fc domain and the third antigen binding moiety is fused at the C-terminus of the Fab heavy chain to the N-terminus of the Fab heavy chain of the first antigen binding moiety. In one such embodiment the first, second and third antigen binding moiety are conventional Fab fragments and have identical light chains (VLCL). In another such embodiment the first antigen binding moiety capable of specific binding to CD3 is a crossover Fab molecule and the second and third antigen binding moiety capable of specific binding to FolR is a

conventional Fab molecule. Optionally, the Fab light chain of the first antigen binding moiety and the Fab light chain of the third antigen binding moiety may additionally be fused to each other.

In another aspect, the invention provides for a bispecific antibody comprising a) a first antigen-binding site that competes for binding to human FolR1 with a reference antibody comprising a variable heavy chain domain (VH) of SEQ ID NO: 49 and a variable light chain domain of SEQ ID NO: 51; and b) a second antigen-binding site that competes for binding to human CD3 with a reference antibody comprising a variable heavy chain domain (VH) of SEQ ID NO: 36 and a variable light chain domain of SEQ ID NO: 31, wherein binding competition is measured using a surface plasmon resonance assay.

In another aspect, the invention provides for a T cell activating bispecific antigen binding molecule comprising a first antigen binding moiety capable of specific binding to CD3, and a second antigen binding moiety capable of specific binding to Folate Receptor 1 (FolR1), wherein the T cell activating bispecific antigen binding molecule binds to the same epitope on human FolR1 as a first reference antibody comprising a variable heavy chain domain (VH) of SEQ ID NO: 49 and a variable light chain domain of SEQ ID NO: 51; and wherein the T cell activating bispecific antigen binding molecule binds to the same epitope on human CD3 as a second reference antibody comprising a variable heavy chain domain (VH) of SEQ ID NO: 36 and a variable light chain domain of SEQ ID NO: 31.

In another aspect, the invention provides for a T cell activating bispecific antigen binding molecule that comprises a first, second, third, fourth and fifth polypeptide chain that form a first, a second and a third antigen binding moiety wherein the first antigen binding moiety is capable of binding CD3 and the second and the third antigen binding moiety each are capable of binding Folate Receptor 1 (FolR1). The first and the second polypeptide chain comprise, in amino (N)-terminal to carboxyl (C)-terminal direction, a first light chain variable domain (VLD1) and a first light chain constant domain (CLD1). The third polypeptide chain comprises, in N-terminal to C-terminal direction, second light chain variable domain (VLD2) and a second heavy chain constant domain 1 (CH1D2). The fourth polypeptide chain comprises, in N-terminal to C-terminal direction, a first heavy chain variable domain (VHD1), a first heavy chain constant domain 1 (CH1D1), a first heavy chain constant domain 2 (CH2D1) and a first heavy chain constant domain 3 (CH3D1). The fifth polypeptide chain comprises VHD1, CH1D1, a second heavy chain

variable domain (VHD2), a second light chain constant domain (CLD2), a second heavy chain constant domain 2 (CH2D2) and a second heavy chain constant domain 3 (CH3D2). The third polypeptide chain and VHD2 and CLD2 of the fifth polypeptide chain form the first antigen binding moiety capable of binding CD3. The second polypeptide chain and VHD1 and CH1D1 of the fifth polypeptide chain form the third binding moiety capable of binding to FolR1. The first polypeptide chain and VHD1 and CH1D1 of the fourth polypeptide chain form the second binding moiety capable of binding to FolR1.

In another embodiment, the second and the third antigen binding moiety are each fused at the C-terminus of the Fab heavy chain to the N-terminus of the first or second subunit of the Fc domain, and the first antigen binding moiety is fused at the C-terminus of the Fab heavy chain to the N-terminus of the Fab heavy chain of the second antigen binding moiety. In a specific such embodiment, the T cell activating bispecific antigen binding molecule essentially consists of a first, a second and a third antigen binding moiety, an Fc domain composed of a first and a second subunit, and optionally one or more peptide linkers, wherein the second and third antigen binding moiety are each fused at the C-terminus of the Fab heavy chain to the N-terminus of the first subunit of the Fc domain and the first antigen binding moiety is fused at the C-terminus of the Fab heavy chain to the N-terminus of the Fab heavy chain of the third antigen binding moiety. In one such embodiment the first, second and third antigen binding moiety are conventional Fab fragments and have identical light chains (VLCL). In another such embodiment the first antigen binding moiety capable of specific binding to CD3 is a crossover Fab molecule and the second and third antigen binding moiety capable of specific binding to FolR is a conventional Fab molecule. Optionally, the Fab light chain of the first antigen binding moiety and the Fab light chain of the second antigen binding moiety may additionally be fused to each other.

The antigen binding moieties may be fused to the Fc domain directly or through a peptide linker. In a particular embodiment the antigen binding moieties are each fused to the Fc domain through an immunoglobulin hinge region. In a specific embodiment, the immunoglobulin hinge region is a human IgG₁ hinge region.

In one embodiment the first and the second antigen binding moiety and the Fc domain are part of an immunoglobulin molecule. In a particular embodiment the immunoglobulin

molecule is an IgG class immunoglobulin. In an even more particular embodiment the immunoglobulin is an IgG₁ subclass immunoglobulin. In another embodiment the immunoglobulin is an IgG₄ subclass immunoglobulin. In a further particular embodiment the immunoglobulin is a human immunoglobulin. In other embodiments the immunoglobulin is a chimeric immunoglobulin or a humanized immunoglobulin.

In a particular embodiment said T cell activating bispecific antigen binding molecule the first and the second antigen binding moiety and the Fc domain are part of an immunoglobulin molecule, and the third antigen binding moiety is fused at the C-terminus of the Fab heavy chain to the N-terminus of the Fab heavy chain of the first antigen binding moiety, wherein the first, second and third antigen binding moiety are conventional Fab fragments and have identical light chains (VLCL), wherein the first antigen binding moiety capable of specific binding to CD3 comprises at least one heavy chain complementarity determining region (CDR) selected from the group consisting of SEQ ID NO: 37, SEQ ID NO: 38 and SEQ ID NO: 39 and at least one light chain CDR selected from the group of SEQ ID NO: 32, SEQ ID NO: 33 and SEQ ID NO: 34; and the second and the third antigen binding moiety capable of specific binding to FolR1 comprise at least one heavy chain complementarity determining region (CDR) selected from the group consisting of SEQ ID NO: 16, SEQ ID NO: 17 and SEQ ID NO: 18 and at least one light chain CDR selected from the group of SEQ ID NO: 32, SEQ ID NO: 33 and SEQ ID NO: 34.

In a particular embodiment said T cell activating bispecific antigen binding molecule the first and the second antigen binding moiety and the Fc domain are part of an immunoglobulin molecule, and the third antigen binding moiety is fused at the C-terminus of the Fab heavy chain to the N-terminus of the Fab heavy chain of the first antigen binding moiety, wherein the first, second and third antigen binding moiety are conventional Fab fragments and have identical light chains (VLCL), wherein the first antigen binding moiety capable of specific binding to CD3 comprises a variable heavy chain comprising a sequence of SEQ ID NO: 36, a variable light chain comprising a sequence of SEQ ID NO: 31; and the second and the third antigen binding moiety capable of specific binding to FolR1 comprise a variable heavy chain comprising a sequence of SEQ ID NO: 15, a variable light chain comprising a sequence of SEQ ID NO: 31.

In a particular embodiment said T cell activating bispecific antigen binding molecule the first and the second antigen binding moiety and the Fc domain are part of an immunoglobulin molecule, and the third antigen binding moiety is fused at the C-terminus of the Fab heavy chain to the N-terminus of the Fab heavy chain of the first antigen binding moiety and the first antigen binding moiety capable of specific binding to CD3 is a crossover Fab molecule wherein either the variable or the constant regions of the Fab light chain and the Fab heavy chain are exchanged, comprising at least one heavy chain complementarity determining region (CDR) selected from the group consisting of SEQ ID NO: 37, SEQ ID NO: 38 and SEQ ID NO: 39 and at least one light chain CDR selected from the group of SEQ ID NO: 32, SEQ ID NO: 33 and SEQ ID NO: 34; and the second and the third antigen binding moiety capable of specific binding to FolR1 comprise at least one heavy chain complementarity determining region (CDR) selected from the group consisting of SEQ ID NO: 8, SEQ ID NO: 56 and SEQ ID NO: 57 and at least one light chain CDR selected from the group of SEQ ID NO: 59, SEQ ID NO: 60 and SEQ ID NO: 65.

In a particular embodiment said T cell activating bispecific antigen binding molecule the first and the second antigen binding moiety and the Fc domain are part of an immunoglobulin molecule, and the third antigen binding moiety is fused at the C-terminus of the Fab heavy chain to the N-terminus of the Fab heavy chain of the first antigen binding moiety and the first antigen binding moiety capable of specific binding to CD3 is a crossover Fab molecule wherein either the variable or the constant regions of the Fab light chain and the Fab heavy chain are exchanged, wherein the first antigen binding moiety capable of specific binding to CD3 comprises a variable heavy chain comprising a sequence of SEQ ID NO: 36, a variable light chain comprising a sequence of SEQ ID NO: 31; and the second and the third antigen binding moiety capable of specific binding to FolR1 comprise a variable heavy chain comprising a sequence of SEQ ID NO: 55, a variable light chain comprising a sequence of SEQ ID NO: 65.

In one embodiment the T cell activating bispecific antigen binding molecule is monovalent for each antigen. In a particular embodiment the T cell activating bispecific antigen binding molecule can bind to human CD3 and human folate receptor alpha (FolR1) and was made without employing a hetero-dimerization approach, such as, e.g.,

knob-into-hole technology. For example, the molecule can be produced by employing a common light chain library and CrossMab technology. In a particular embodiment, The variable region of the CD3 binder is fused to the CH1 domain of a standard human IgG1 antibody to form the VLVH crossed molecule (fused to Fc) which is common for both specificities. To generate the crossed counterparts (VHCL), a CD3 specific variable heavy chain domain is fused to a constant human λ light chain whereas a variable heavy chain domain specific for human FolR1 (e.g., isolated from a common light chain library) is fused to a constant human κ light chain. The resulting desired molecule with correctly paired chains comprises both kappa and lambda light chains or fragments thereof. Consequently, this desired bispecific molecule species can be purified from mispaired or homodimeric species with sequential purification steps selecting for kappa and lambda light chain, in either sequence. In one particular embodiment, purification of the desired bispecific antibody employs subsequent purification steps with KappaSelect and LambdaFabSelect columns (GE Healthcare) to remove undesired homodimeric antibodies.

Fc domain

The Fc domain of the T cell activating bispecific antigen binding molecule consists of a pair of polypeptide chains comprising heavy chain domains of an immunoglobulin molecule. For example, the Fc domain of an immunoglobulin G (IgG) molecule is a dimer, each subunit of which comprises the CH2 and CH3 IgG heavy chain constant domains. The two subunits of the Fc domain are capable of stable association with each other. In one embodiment the T cell activating bispecific antigen binding molecule of the invention comprises not more than one Fc domain.

In one embodiment according the invention the Fc domain of the T cell activating bispecific antigen binding molecule is an IgG Fc domain. In a particular embodiment the Fc domain is an IgG₁ Fc domain. In another embodiment the Fc domain is an IgG₄ Fc domain. In a more specific embodiment, the Fc domain is an IgG₄ Fc domain comprising an amino acid substitution at position S228 (Kabat numbering), particularly the amino acid substitution S228P. This amino acid substitution reduces in vivo Fab arm exchange of IgG₄ antibodies (see Stubenrauch et al., Drug Metabolism and Disposition 38, 84-91 (2010)). In a further particular embodiment the Fc domain is human.

Fc domain modifications promoting heterodimerization

T cell activating bispecific antigen binding molecules according to the invention comprise different antigen binding moieties, fused to one or the other of the two subunits of the Fc domain, thus the two subunits of the Fc domain are typically comprised in two non-identical polypeptide chains. Recombinant co-expression of these polypeptides and subsequent dimerization leads to several possible combinations of the two polypeptides. To improve the yield and purity of T cell activating bispecific antigen binding molecules in recombinant production, it will thus be advantageous to introduce in the Fc domain of the T cell activating bispecific antigen binding molecule a modification promoting the association of the desired polypeptides.

Accordingly, in particular embodiments the Fc domain of the T cell activating bispecific antigen binding molecule according to the invention comprises a modification promoting the association of the first and the second subunit of the Fc domain. The site of most extensive protein-protein interaction between the two subunits of a human IgG Fc domain is in the CH3 domain of the Fc domain. Thus, in one embodiment said modification is in the CH3 domain of the Fc domain.

In a specific embodiment said modification is a so-called “knob-into-hole” modification, comprising a “knob” modification in one of the two subunits of the Fc domain and a “hole” modification in the other one of the two subunits of the Fc domain.

The knob-into-hole technology is described e.g. in US 5,731,168; US 7,695,936; Ridgway et al., *Prot Eng* 9, 617-621 (1996) and Carter, *J Immunol Meth* 248, 7-15 (2001). Generally, the method involves introducing a protuberance (“knob”) at the interface of a first polypeptide and a corresponding cavity (“hole”) in the interface of a second polypeptide, such that the protuberance can be positioned in the cavity so as to promote heterodimer formation and hinder homodimer formation. Protuberances are constructed by replacing small amino acid side chains from the interface of the first polypeptide with larger side chains (e.g. tyrosine or tryptophan). Compensatory cavities of identical or similar size to the protuberances are created in the interface of the second polypeptide by replacing large amino acid side chains with smaller ones (e.g. alanine or threonine).

Accordingly, in a particular embodiment, in the CH3 domain of the first subunit of the Fc domain of the T cell activating bispecific antigen binding molecule an amino acid residue is replaced with an amino acid residue having a larger side chain volume, thereby generating a protuberance within the CH3 domain of the first subunit which is positionable in a cavity within the CH3 domain of the second subunit, and in the CH3

domain of the second subunit of the Fc domain an amino acid residue is replaced with an amino acid residue having a smaller side chain volume, thereby generating a cavity within the CH3 domain of the second subunit within which the protuberance within the CH3 domain of the first subunit is positionable.

- 5 The protuberance and cavity can be made by altering the nucleic acid encoding the polypeptides, e.g. by site-specific mutagenesis, or by peptide synthesis.

In a specific embodiment, in the CH3 domain of the first subunit of the Fc domain the threonine residue at position 366 is replaced with a tryptophan residue (T366W), and in the CH3 domain of the second subunit of the Fc domain the tyrosine residue at position
10 407 is replaced with a valine residue (Y407V). In one embodiment, in the second subunit of the Fc domain additionally the threonine residue at position 366 is replaced with a serine residue (T366S) and the leucine residue at position 368 is replaced with an alanine residue (L368A).

In yet a further embodiment, in the first subunit of the Fc domain additionally the serine
15 residue at position 354 is replaced with a cysteine residue (S354C), and in the second subunit of the Fc domain additionally the tyrosine residue at position 349 is replaced by a cysteine residue (Y349C). Introduction of these two cysteine residues results in formation of a disulfide bridge between the two subunits of the Fc domain, thus further stabilizing the dimer (Carter, J Immunol Methods 248, 7-15 (2001)).

- 20 In a particular embodiment the antigen binding moiety capable of binding to CD3 is fused (optionally via the antigen binding moiety capable of binding to a target cell antigen) to the first subunit of the Fc domain (comprising the “knob” modification). Without wishing to be bound by theory, fusion of the antigen binding moiety capable of binding to CD3 to the knob-containing subunit of the Fc domain will (further) minimize the generation of
25 antigen binding molecules comprising two antigen binding moieties capable of binding to CD3 (steric clash of two knob-containing polypeptides).

In an alternative embodiment a modification promoting association of the first and the second subunit of the Fc domain comprises a modification mediating electrostatic steering effects, e.g. as described in PCT publication WO 2009/089004. Generally, this method
30 involves replacement of one or more amino acid residues at the interface of the two Fc domain subunits by charged amino acid residues so that homodimer formation becomes electrostatically unfavorable but heterodimerization electrostatically favorable.

Fc domain modifications abolishing Fc receptor binding and/or effector function

The Fc domain confers to the T cell activating bispecific antigen binding molecule favorable pharmacokinetic properties, including a long serum half-life which contributes to good accumulation in the target tissue and a favorable tissue-blood distribution ratio.

- 5 At the same time it may, however, lead to undesirable targeting of the T cell activating bispecific antigen binding molecule to cells expressing Fc receptors rather than to the preferred antigen-bearing cells. Moreover, the co-activation of Fc receptor signaling pathways may lead to cytokine release which, in combination with the T cell activating properties and the long half-life of the antigen binding molecule, results in excessive
- 10 activation of cytokine receptors and severe side effects upon systemic administration. Activation of (Fc receptor-bearing) immune cells other than T cells may even reduce efficacy of the T cell activating bispecific antigen binding molecule due to the potential destruction of T cells e.g. by NK cells.

- Accordingly, in particular embodiments the Fc domain of the T cell activating bispecific
- 15 antigen binding molecules according to the invention exhibits reduced binding affinity to an Fc receptor and/or reduced effector function, as compared to a native IgG₁ Fc domain. In one such embodiment the Fc domain (or the T cell activating bispecific antigen binding molecule comprising said Fc domain) exhibits less than 50%, preferably less than 20%, more preferably less than 10% and most preferably less than 5% of the binding affinity to
- 20 an Fc receptor, as compared to a native IgG₁ Fc domain (or a T cell activating bispecific antigen binding molecule comprising a native IgG₁ Fc domain), and/or less than 50%, preferably less than 20%, more preferably less than 10% and most preferably less than 5% of the effector function, as compared to a native IgG₁ Fc domain domain (or a T cell activating bispecific antigen binding molecule comprising a native IgG₁ Fc domain). In
- 25 one embodiment, the Fc domain domain (or the T cell activating bispecific antigen binding molecule comprising said Fc domain) does not substantially bind to an Fc receptor and/or induce effector function. In a particular embodiment the Fc receptor is an Fcγ receptor. In one embodiment the Fc receptor is a human Fc receptor. In one embodiment the Fc receptor is an activating Fc receptor. In a specific embodiment the Fc
- 30 receptor is an activating human Fcγ receptor, more specifically human FcγRIIIa, FcγRI or FcγRIIa, most specifically human FcγRIIIa. In one embodiment the effector function is one or more selected from the group of CDC, ADCC, ADCP, and cytokine secretion. In a particular embodiment the effector function is ADCC. In one embodiment the Fc domain

domain exhibits substantially similar binding affinity to neonatal Fc receptor (FcRn), as compared to a native IgG₁ Fc domain domain. Substantially similar binding to FcRn is achieved when the Fc domain (or the T cell activating bispecific antigen binding molecule comprising said Fc domain) exhibits greater than about 70%, particularly greater than
5 about 80%, more particularly greater than about 90% of the binding affinity of a native IgG₁ Fc domain (or the T cell activating bispecific antigen binding molecule comprising a native IgG₁ Fc domain) to FcRn.

In certain embodiments the Fc domain is engineered to have reduced binding affinity to an Fc receptor and/or reduced effector function, as compared to a non-engineered Fc
10 domain. In particular embodiments, the Fc domain of the T cell activating bispecific antigen binding molecule comprises one or more amino acid mutation that reduces the binding affinity of the Fc domain to an Fc receptor and/or effector function. Typically, the same one or more amino acid mutation is present in each of the two subunits of the Fc domain. In one embodiment the amino acid mutation reduces the binding affinity of the
15 Fc domain to an Fc receptor. In one embodiment the amino acid mutation reduces the binding affinity of the Fc domain to an Fc receptor by at least 2-fold, at least 5-fold, or at least 10-fold. In embodiments where there is more than one amino acid mutation that reduces the binding affinity of the Fc domain to the Fc receptor, the combination of these amino acid mutations may reduce the binding affinity of the Fc domain to an Fc receptor
20 by at least 10-fold, at least 20-fold, or even at least 50-fold. In one embodiment the T cell activating bispecific antigen binding molecule comprising an engineered Fc domain exhibits less than 20%, particularly less than 10%, more particularly less than 5% of the binding affinity to an Fc receptor as compared to a T cell activating bispecific antigen binding molecule comprising a non-engineered Fc domain. In a particular embodiment
25 the Fc receptor is an Fcγ receptor. In some embodiments the Fc receptor is a human Fc receptor. In some embodiments the Fc receptor is an activating Fc receptor. In a specific embodiment the Fc receptor is an activating human Fcγ receptor, more specifically human FcγRIIIa, FcγRI or FcγRIIa, most specifically human FcγRIIIa. Preferably, binding to each of these receptors is reduced. In some embodiments binding affinity to a
30 complement component, specifically binding affinity to C1q, is also reduced. In one embodiment binding affinity to neonatal Fc receptor (FcRn) is not reduced. Substantially similar binding to FcRn, i.e. preservation of the binding affinity of the Fc domain to said receptor, is achieved when the Fc domain (or the T cell activating bispecific antigen

binding molecule comprising said Fc domain) exhibits greater than about 70% of the binding affinity of a non-engineered form of the Fc domain (or the T cell activating bispecific antigen binding molecule comprising said non-engineered form of the Fc domain) to FcRn. The Fc domain, or T cell activating bispecific antigen binding molecules of the invention comprising said Fc domain, may exhibit greater than about 80% and even greater than about 90% of such affinity. In certain embodiments the Fc domain of the T cell activating bispecific antigen binding molecule is engineered to have reduced effector function, as compared to a non-engineered Fc domain. The reduced effector function can include, but is not limited to, one or more of the following: reduced complement dependent cytotoxicity (CDC), reduced antibody-dependent cell-mediated cytotoxicity (ADCC), reduced antibody-dependent cellular phagocytosis (ADCP), reduced cytokine secretion, reduced immune complex-mediated antigen uptake by antigen-presenting cells, reduced binding to NK cells, reduced binding to macrophages, reduced binding to monocytes, reduced binding to polymorphonuclear cells, reduced direct signaling inducing apoptosis, reduced crosslinking of target-bound antibodies, reduced dendritic cell maturation, or reduced T cell priming. In one embodiment the reduced effector function is one or more selected from the group of reduced CDC, reduced ADCC, reduced ADCP, and reduced cytokine secretion. In a particular embodiment the reduced effector function is reduced ADCC. In one embodiment the reduced ADCC is less than 20% of the ADCC induced by a non-engineered Fc domain (or a T cell activating bispecific antigen binding molecule comprising a non-engineered Fc domain).

In one embodiment the amino acid mutation that reduces the binding affinity of the Fc domain to an Fc receptor and/or effector function is an amino acid substitution. In one embodiment the Fc domain comprises an amino acid substitution at a position selected from the group of E233, L234, L235, N297, P331 and P329. In a more specific embodiment the Fc domain comprises an amino acid substitution at a position selected from the group of L234, L235 and P329. In some embodiments the Fc domain comprises the amino acid substitutions L234A and L235A. In one such embodiment, the Fc domain is an IgG₁ Fc domain, particularly a human IgG₁ Fc domain. In one embodiment the Fc domain comprises an amino acid substitution at position P329. In a more specific embodiment the amino acid substitution is P329A or P329G, particularly P329G. In one embodiment the Fc domain comprises an amino acid substitution at position P329 and a

further amino acid substitution at a position selected from E233, L234, L235, N297 and P331. In a more specific embodiment the further amino acid substitution is E233P, L234A, L235A, L235E, N297A, N297D or P331S. In particular embodiments the Fc domain comprises amino acid substitutions at positions P329, L234 and L235. In more particular embodiments the Fc domain comprises the amino acid mutations L234A, L235A and P329G ("P329G LALA"). In one such embodiment, the Fc domain is an IgG₁ Fc domain, particularly a human IgG₁ Fc domain. The "P329G LALA" combination of amino acid substitutions almost completely abolishes Fcγ receptor binding of a human IgG₁ Fc domain, as described in PCT publication no. WO 2012/130831, incorporated herein by reference in its entirety. WO 2012/130831 also describes methods of preparing such mutant Fc domains and methods for determining its properties such as Fc receptor binding or effector functions.

IgG₄ antibodies exhibit reduced binding affinity to Fc receptors and reduced effector functions as compared to IgG₁ antibodies. Hence, in some embodiments the Fc domain of the T cell activating bispecific antigen binding molecules of the invention is an IgG₄ Fc domain, particularly a human IgG₄ Fc domain. In one embodiment the IgG₄ Fc domain comprises amino acid substitutions at position S228, specifically the amino acid substitution S228P. To further reduce its binding affinity to an Fc receptor and/or its effector function, in one embodiment the IgG₄ Fc domain comprises an amino acid substitution at position L235, specifically the amino acid substitution L235E. In another embodiment, the IgG₄ Fc domain comprises an amino acid substitution at position P329, specifically the amino acid substitution P329G. In a particular embodiment, the IgG₄ Fc domain comprises amino acid substitutions at positions S228, L235 and P329, specifically amino acid substitutions S228P, L235E and P329G. Such IgG₄ Fc domain mutants and their Fcγ receptor binding properties are described in PCT publication no. WO 2012/130831, incorporated herein by reference in its entirety.

In a particular embodiment the Fc domain exhibiting reduced binding affinity to an Fc receptor and/or reduced effector function, as compared to a native IgG₁ Fc domain, is a human IgG₁ Fc domain comprising the amino acid substitutions L234A, L235A and optionally P329G, or a human IgG₄ Fc domain comprising the amino acid substitutions S228P, L235E and optionally P329G.

In certain embodiments N-glycosylation of the Fc domain has been eliminated. In one such embodiment the Fc domain comprises an amino acid mutation at position N297,

particularly an amino acid substitution replacing asparagine by alanine (N297A) or aspartic acid (N297D).

In addition to the Fc domains described hereinabove and in PCT publication no. WO 2012/130831, Fc domains with reduced Fc receptor binding and/or effector function also include those with substitution of one or more of Fc domain residues 238, 265, 269, 270, 297, 327 and 329 (U.S. Patent No. 6,737,056). Such Fc mutants include Fc mutants with substitutions at two or more of amino acid positions 265, 269, 270, 297 and 327, including the so-called "DANA" Fc mutant with substitution of residues 265 and 297 to alanine (US Patent No. 7,332,581).

10 Mutant Fc domains can be prepared by amino acid deletion, substitution, insertion or modification using genetic or chemical methods well known in the art. Genetic methods may include site-specific mutagenesis of the encoding DNA sequence, PCR, gene synthesis, and the like. The correct nucleotide changes can be verified for example by sequencing.

15 Binding to Fc receptors can be easily determined e.g. by ELISA, or by Surface Plasmon Resonance (SPR) using standard instrumentation such as a BIAcore instrument (GE Healthcare), and Fc receptors such as may be obtained by recombinant expression. A suitable such binding assay is described herein. Alternatively, binding affinity of Fc domains or cell activating bispecific antigen binding molecules comprising an Fc domain for Fc receptors may be evaluated using cell lines known to express particular Fc receptors, such as human NK cells expressing FcγIIIa receptor.

Effector function of an Fc domain, or a T cell activating bispecific antigen binding molecule comprising an Fc domain, can be measured by methods known in the art. A suitable assay for measuring ADCC is described herein. Other examples of *in vitro* assays to assess ADCC activity of a molecule of interest are described in U.S. Patent No. 25 5,500,362; Hellstrom et al. Proc Natl Acad Sci USA 83, 7059-7063 (1986) and Hellstrom et al., Proc Natl Acad Sci USA 82, 1499-1502 (1985); U.S. Patent No. 5,821,337; Bruggemann et al., J Exp Med 166, 1351-1361 (1987). Alternatively, non-radioactive assays methods may be employed (see, for example, ACTITM non-radioactive cytotoxicity assay for flow cytometry (CellTechnology, Inc. Mountain View, CA); and CytoTox 96® 30 non-radioactive cytotoxicity assay (Promega, Madison, WI)). Useful effector cells for such assays include peripheral blood mononuclear cells (PBMC) and Natural Killer (NK) cells. Alternatively, or additionally, ADCC activity of the molecule of interest may be

assessed *in vivo*, e.g. in a animal model such as that disclosed in Clynes et al., Proc Natl Acad Sci USA 95, 652-656 (1998).

In some embodiments, binding of the Fc domain to a complement component, specifically to C1q, is reduced. Accordingly, in some embodiments wherein the Fc domain is engineered to have reduced effector function, said reduced effector function includes reduced CDC. C1q binding assays may be carried out to determine whether the T cell activating bispecific antigen binding molecule is able to bind C1q and hence has CDC activity. See e.g., C1q and C3c binding ELISA in WO 2006/029879 and WO 2005/100402. To assess complement activation, a CDC assay may be performed (see, for example, Gazzano-Santoro et al., J Immunol Methods 202, 163 (1996); Cragg et al., Blood 101, 1045-1052 (2003); and Cragg and Glennie, Blood 103, 2738-2743 (2004)).

Fc domain modifications promoting heterodimerization

The T cell activating bispecific antigen binding molecule of the invention comprise different antigen binding moieties, some of which are fused to one or the other of the two subunits of the Fc domain, thus the two subunits of the Fc domain are typically comprised in two non-identical polypeptide chains. Recombinant co-expression of these polypeptides and subsequent dimerization leads to several possible combinations of the two polypeptides. To improve the yield and purity of the bispecific antibodies of the invention in recombinant production, it will thus be advantageous to introduce in the Fc domain of the bispecific antibodies of the invention a modification promoting the association of the desired polypeptides.

Accordingly, in particular embodiments, the Fc domain of the bispecific antibodies of the invention comprises a modification promoting the association of the first and the second subunit of the Fc domain. The site of most extensive protein-protein interaction between the two subunits of a human IgG Fc domain is in the CH3 domain of the Fc domain. Thus, in one embodiment said modification is in the CH3 domain of the Fc domain.

In a specific embodiment, said modification is a so-called “knob-into-hole” modification, comprising a “knob” modification in one of the two subunits of the Fc domain and a “hole” modification in the other one of the two subunits of the Fc domain. The knob-into-hole technology is described e.g. in US 5,731,168; US 7,695,936; Ridgway et al., Prot Eng 9, 617-621 (1996) and Carter, J Immunol Meth 248, 7-15 (2001).

Generally, the method involves introducing a protuberance (“knob”) at the

interface of a first polypeptide and a corresponding cavity ("hole") in the interface of a second polypeptide, such that the protuberance can be positioned in the cavity so as to promote heterodimer formation and hinder homodimer formation. Protuberances are constructed by replacing small amino acid side chains from the interface of the first
5 polypeptide with larger side chains (e.g. tyrosine or tryptophan). Compensatory cavities of identical or similar size to the protuberances are created in the interface of the second polypeptide by replacing large amino acid side chains with smaller ones (e.g. alanine or threonine).

Accordingly, in a particular embodiment, in the CH3 domain of the first subunit of
10 the Fc domain of the bispecific antibodies of the invention an amino acid residue is replaced with an amino acid residue having a larger side chain volume, thereby generating a protuberance within the CH3 domain of the first subunit which is positionable in a cavity within the CH3 domain of the second subunit, and in the CH3 domain of the second subunit of the Fc domain an amino acid residue is replaced with an
15 amino acid residue having a smaller side chain volume, thereby generating a cavity within the CH3 domain of the second subunit within which the protuberance within the CH3 domain of the first subunit is positionable.

The protuberance and cavity can be made by altering the nucleic acid encoding the polypeptides, e.g. by site-specific mutagenesis, or by peptide synthesis.

20 In a specific embodiment, in the CH3 domain of the first subunit of the Fc domain the threonine residue at position 366 is replaced with a tryptophan residue (T366W), and in the CH3 domain of the second subunit of the Fc domain the tyrosine residue at position 407 is replaced with a valine residue (Y407V). In one embodiment, in the second subunit of the Fc domain additionally the threonine residue at position 366 is replaced with a
25 serine residue (T366S) and the leucine residue at position 368 is replaced with an alanine residue (L368A).

In yet a further embodiment, in the first subunit of the Fc domain additionally the serine residue at position 354 is replaced with a cysteine residue (S354C), and in the second subunit of the Fc domain additionally the tyrosine residue at position 349 is
30 replaced by a cysteine residue (Y349C). Introduction of these two cysteine residues results in formation of a disulfide bridge between the two subunits of the Fc domain, further stabilizing the dimer (Carter, J Immunol Methods 248, 7-15 (2001)).

In an alternative embodiment a modification promoting association of the first and the second subunit of the Fc domain comprises a modification mediating electrostatic steering effects, e.g. as described in WO 2009/089004. Generally, this method involves replacement of one or more amino acid residues at the interface of the two Fc domain subunits by charged amino acid residues so that homodimer formation becomes electrostatically unfavorable but heterodimerization electrostatically favorable.

In one embodiment, a T cell activating bispecific antigen binding molecule that binds to FolR1 and CD3 according to any of the above embodiments comprises an Immunoglobulin G (IgG) molecule with two binding sites specific for FolR1, wherein the Fc part of the first heavy chain comprises a first dimerization module and the Fc part of the second heavy chain comprises a second dimerization module allowing a heterodimerization of the two heavy chains of the IgG molecule.

In a further preferred embodiment, the first dimerization module comprises knobs and the second dimerization module comprises holes according to the knobs into holes strategy (see Carter P.; Ridgway J.B.B.; Presta L.G.: Immunotechnology, Volume 2, Number 1, February 1996 , pp. 73-73(1)).

Biological properties and functional characteristics of T cell activating bispecific antigen binding molecules

One of skill in the art can appreciate the advantageous efficiency of a molecule that selectively distinguishes between cancerous and non-cancerous, healthy cells. One way to accomplish this goal is by appropriate target selection. Markers expressed exclusively on tumor cells can be employed to selectively target effector molecules or cells to tumor cells while sparing normal cells that do not express such marker. However, in some instances, so called tumor cell markers are also expressed in normal tissue, albeit at lower levels. This expression in normal tissue raises the possibility of toxicity. Thus, there was a need in the art for molecules that can more selectively target tumor cells. The invention described herein provides for T cell activating bispecific antigen binding molecules that selectively target FolR1-positive tumor cells and not normal, non-cancerous cells that express FolR1 at low levels or not at all. In one embodiment, the T cell activating bispecific antigen binding molecule comprises at least two, preferably two, FolR1 binding moieties of relatively low affinity that confer an avidity effect which allows for differentiation between high and low FolR1 expressing cells. Because tumor cells express FolR1 at high or intermediate levels, this embodiment of the invention selectively

binds to, and/or induces killing of, tumor cells and not normal, non-cancerous cells that express FolR1 at low levels or not at all. In one embodiment, the T cell activating bispecific antigen binding molecule is in the 2+1 inverted format. In one embodiment, the T cell activating bispecific antigen binding molecule induces T cell mediated killing of FolR1-positive tumor cells and not non-tumor cells and comprises a CD3 antigen binding moiety that comprises the heavy chain CDR1 of SEQ ID NO: 37, the heavy chain CDR2 of SEQ ID NO: 38, the heavy chain CDR3 of SEQ ID NO:39, the light chain CDR1 of SEQ ID NO: 32, the light chain CDR2 of SEQ ID NO: 33, and the light chain CDR3 of SEQ ID NO:34 and two FolR1 antigen binding moieties that each comprise the heavy chain CDR1 of SEQ ID NO: 8, the heavy chain CDR2 of SEQ ID NO: 9, the heavy chain CDR3 of SEQ ID NO:50, the light chain CDR1 of SEQ ID NO: 52, the light chain CDR2 of SEQ ID NO: 53, and the light chain CDR3 of SEQ ID NO:54.

In one specific embodiment, the T cell activating bispecific antigen binding molecule does not induce killing of a normal cells having less than about 1000 copies of FolR1 its surface.

In addition to the above advantageous characteristics, one embodiment of the invention does not require chemical cross linking or a hybrid approach to be produced. Accordingly, in one embodiment, the invention provides for T cell activating bispecific antigen binding molecule capable of production in CHO cells. In one embodiment, the T cell activating bispecific antigen binding molecule comprises humanized and human polypeptides. In one embodiment, the T cell activating bispecific antigen binding molecule does not cause FcγR crosslinking. In one such embodiment, the T cell activating bispecific antigen binding molecule is capable of production in CHO cells and comprises a CD3 antigen binding moiety that comprises the heavy chain CDR1 of SEQ ID NO: 37, the heavy chain CDR2 of SEQ ID NO: 38, the heavy chain CDR3 of SEQ ID NO:39, the light chain CDR1 of SEQ ID NO: 32, the light chain CDR2 of SEQ ID NO: 33, and the light chain CDR3 of SEQ ID NO:34 and two FolR1 antigen binding moieties that each comprise the heavy chain CDR1 of SEQ ID NO: 8, the heavy chain CDR2 of SEQ ID NO: 9, the heavy chain CDR3 of SEQ ID NO:50, the light chain CDR1 of SEQ ID NO: 52, the light chain CDR2 of SEQ ID NO: 53, and the light chain CDR3 of SEQ ID NO:54.

As noted above, some embodiments contemplated herein include T cell activating bispecific antigen binding molecules having two binding moieties that confer specific

binding to FolR1 and one binding moiety that confers specificity to the T cell activating antigen CD3, wherein each individual FolR1 binding moiety engages the antigen with low affinity. Because the molecule comprises two antigen binding moieties that confer binding to FolR1, the overall avidity of the molecule, nevertheless, provides effective
5 binding to FolR1-expressing target cells and activation of T cells to induce T cell effector function. Considering that while FolR1 is expressed at various level on tumor cells, it is also expressed at very low levels (e.g., less than about 1000 copies on the cell surface) in certain normal cells, one of skill in the art can readily recognize the advantageous efficiency of such a molecule for use as a therapeutic agent. Such molecule selectively
10 targets tumor cells over normal cells. Such molecule, thus, can be administered to an individual in need thereof with significantly less concern about toxicity resulting from FolR1 positive normal cells compared to molecules that bind to FolR1 with high affinity to induce effector function.

In one embodiment, the T cell activating bispecific antigen binding molecule binds
15 human FolR1 with an apparent K_D of about 5.36 pM to about 4 nM. In one embodiment, the T cell activating bispecific antigen binding molecule binds human and cynomolgus FolR1 with an apparent K_D of about 4 nM. In one embodiment, the T cell activating bispecific antigen binding molecule binds murine FolR1 with an apparent K_D of about 1.5 nM. In one embodiment, the T cell activating bispecific antigen binding molecule binds
20 human FolR1 with a monovalent binding K_D of at least about 1000 nM. In a specific embodiment, the T cell activating bispecific antigen binding molecule binds human and cynomolgus FolR1 with an apparent K_D of about 4 nM, binds murine FolR1 with an apparent K_D of about 1.5 nM, and comprises a CD3 antigen binding moiety that comprises the heavy chain CDR1 of SEQ ID NO: 37, the heavy chain CDR2 of SEQ ID
25 NO: 38, the heavy chain CDR3 of SEQ ID NO:39, the light chain CDR1 of SEQ ID NO: 32, the light chain CDR2 of SEQ ID NO: 33, and the light chain CDR3 of SEQ ID NO:34 and two FolR1 antigen binding moieties that each comprise the heavy chain CDR1 of SEQ ID NO: 8, the heavy chain CDR2 of SEQ ID NO: 9, the heavy chain CDR3 of SEQ ID NO:50, the light chain CDR1 of SEQ ID NO: 52, the light chain CDR2 of SEQ ID
30 NO: 53, and the light chain CDR3 of SEQ ID NO:54. In one embodiment, the T cell activating bispecific antigen binding molecule binds human FolR1 with a monovalent binding K_D of at least about 1000 nM and comprises a CD3 antigen binding moiety that comprises the heavy chain CDR1 of SEQ ID NO: 37, the heavy chain CDR2 of SEQ ID

NO: 38, the heavy chain CDR3 of SEQ ID NO:39, the light chain CDR1 of SEQ ID NO: 32, the light chain CDR2 of SEQ ID NO: 33, and the light chain CDR3 of SEQ ID NO:34 and two FolR1 antigen binding moieties that each comprise the heavy chain CDR1 of SEQ ID NO: 8, the heavy chain CDR2 of SEQ ID NO: 9, the heavy chain CDR3 of SEQ ID NO:50, the light chain CDR1 of SEQ ID NO: 52, the light chain CDR2 of SEQ ID NO: 53, and the light chain CDR3 of SEQ ID NO:54.

As described above, the T cell activating bispecific antigen binding molecules contemplated herein can induce T cell effector function, e.g., cell surface marker expression, cytokine production, T cell mediated killing. In one embodiment, the T cell activating bispecific antigen binding molecule induces T cell mediated killing of the FolR1-expressing target cell, such as a human tumor cell, in vitro. In one embodiment, the T cell is a CD8 positive T cell. Examples of FolR1-expressing human tumor cells include but are not limited to Hela, Skov-3, HT-29, and HRCEpiC cells. Other FolR1 positive human cancer cells that can be used for in vitro testing are readily available to the skilled artisan. In one embodiment, the T cell activating bispecific antigen binding molecule induces T cell mediated killing of the FolR1-expressing human tumor cell in vitro with an EC50 of between about 36 pM and about 39573 pM after 24 hours. Specifically contemplated are T cell activating bispecific antigen binding molecules that induce T cell mediated killing of the FolR1-expressing tumor cell in vitro with an EC50 of about 36 pM after 24 hours. In one embodiment, the T cell activating bispecific antigen binding molecule induces T cell mediated killing of the FolR1-expressing tumor cell in vitro with an EC50 of about 178.4 pM after 24 hours. In one embodiment, the T cell activating bispecific antigen binding molecule induces T cell mediated killing of the FolR1-expressing tumor cell in vitro with an EC50 of about 134.5 pM or greater after 48 hours. The EC50 can be measure by methods known in the art, for example by methods disclosed herein by the examples.

In one embodiment, the T cell activating bispecific antigen binding molecule of any of the above embodiments induces upregulation of cell surface expression of at least one of CD25 and CD69 on the T cell as measured by flow cytometry. In one embodiment, the T cell is a CD4 positive T cell or a CD8 positive T cell.

In one embodiment, the T cell activating bispecific antigen binding molecule of any of the above embodiments binds to FolR1 expressed on a human tumor cell. In one embodiment, the T cell activating bispecific antigen binding molecule of any of the above

embodiments binds to a conformational epitope on human FolR1. In one embodiment, the T cell activating bispecific antigen binding molecule of any of the above embodiments does not bind to human Folate Receptor 2 (FolR2) or to human Folate Receptor 3 (FolR3). In one embodiment of the T cell activating bispecific antigen binding molecule of any of the above embodiments, the antigen binding moiety binds to a FolR1 polypeptide comprising the amino acids 25 to 234 of human FolR1 (SEQ ID NO:227). In one embodiment of the T cell activating bispecific antigen binding molecule of any of the above embodiments, the FolR1 antigen binding moiety binds to a FolR1 polypeptide comprising the amino acid sequence of SEQ ID NOs:227, 230 and 231, and wherein the FolR1 antigen binding moiety does not bind to a FolR polypeptide comprising the amino acid sequence of SEQ ID NOs:228 and 229. In one specific embodiment, the T cell activating bispecific antigen binding molecule comprises a FolR1 antigen binding moiety that binds to a FolR1 polypeptide comprising the amino acid sequence of SEQ ID NOs:227, 230 and 231, and wherein the FolR1 antigen binding moiety does not bind to a FolR polypeptide comprising the amino acid sequence of SEQ ID NOs:228 and 229, and comprises a CD3 antigen binding moiety that comprises the heavy chain CDR1 of SEQ ID NO: 37, the heavy chain CDR2 of SEQ ID NO: 38, the heavy chain CDR3 of SEQ ID NO:39, the light chain CDR1 of SEQ ID NO: 32, the light chain CDR2 of SEQ ID NO: 33, and the light chain CDR3 of SEQ ID NO:34 and two FolR1 antigen binding moieties that each comprise the heavy chain CDR1 of SEQ ID NO: 8, the heavy chain CDR2 of SEQ ID NO: 9, the heavy chain CDR3 of SEQ ID NO:50, the light chain CDR1 of SEQ ID NO: 52, the light chain CDR2 of SEQ ID NO: 53, and the light chain CDR3 of SEQ ID NO:54.

With respect to the FolR1, the T cell activating bispecific antigen binding molecules contemplated herein can have agonist, antagonist or neutral effect. Examples of agonist effect include induction or enhancement of signaling through the FolR1 upon engagement by the FolR1 binding moiety with the FolR1 receptor on the target cell. Examples of antagonist activity include abrogation or reduction of signaling through the FolR1 upon engagement by the FolR1 binding moiety with the FolR1 receptor on the target cell. This can, for example, occur by blocking or reducing the interaction between folate with FolR1.

Exemplary PD-1 Axis Binding Antagonists for use in the invention

Provided herein are methods for treating or delaying progression of cancer in an individual comprising administering to the individual an effective amount of a T cell activating bispecific antigen binding molecule and a PD-1 axis binding antagonist. For

5 example, a PD-1 axis binding antagonist includes a PD-1 binding antagonist, a PDL1 binding antagonist and a PDL2 binding antagonist. Alternative names for “PD-1” include CD279 and SLEB2. Alternative names for “PDL1” include B7-H1, B7-4, CD274, and B7-H. Alternative names for “PDL2” include B7-DC, Btdc, and CD273. In some embodiments, PD-1, PDL1, and PDL2 are human PD-1, PDL1 and PDL2.

10 In some embodiments, the PD-1 binding antagonist is a molecule that inhibits the binding of PD-1 to its ligand binding partners. In a specific aspect the PD-1 ligand binding partners are PDL1 and/or PDL2. In another embodiment, a PDL1 binding antagonist is a molecule that inhibits the binding of PDL1 to its binding partners. In a specific aspect, PDL1 binding partners are PD-1 and/or B7-1. In another embodiment, the PDL2 binding

15 antagonist is a molecule that inhibits the binding of PDL2 to its binding partners. In a specific aspect, a PDL2 binding partner is PD-1. The antagonist may be an antibody, an antigen binding fragment thereof, an immunoadhesin, a fusion protein, or oligopeptide. In some embodiments, the PD-1 binding antagonist is an anti-PD-1 antibody (e.g., a human antibody, a humanized antibody, or a chimeric antibody). In some embodiments,

20 the anti-PD-1 antibody is selected from the group consisting of nivolumab, pembrolizumab, and CT-011. In some embodiments, the PD-1 binding antagonist is an immunoadhesin (e.g., an immunoadhesin comprising an extracellular or PD-1 binding portion of PDL1 or PDL2 fused to a constant region (e.g., an Fc region of an immunoglobulin sequence). In some embodiments, the PD-1 binding antagonist is AMP-

25 224. Nivolumab, also known as MDX-1106-04, MDX-1106, ONO-4538, BMS-936558, and OPDIVO®, is an anti-PD-1 antibody described in WO2006/121168. Pembrolizumab, also known as MK-3475, Merck 3475, lambrolizumab, KEYTRUDA®, and SCH-900475, is an anti-PD-1 antibody described in WO2009/114335. CT-011, also known as hBAT or hBAT-1, is an anti-PD-1 antibody described in WO2009/101611. AMP-224, also known

30 as B7-DCIg, is a PDL2-Fc fusion soluble receptor described in WO2010/027827 and WO2011/066342.

In some embodiments, the anti-PD-1 antibody is nivolumab (CAS Registry Number:946414-94-4). In a still further embodiment, provided is an isolated anti-PD-1 antibody comprising a heavy chain variable region comprising the heavy chain variable region amino acid sequence from SEQ ID NO:274 and/or a light chain variable region comprising the light chain variable region amino acid sequence from SEQ ID NO:275. In a still further embodiment, provided is an isolated anti-PD-1 antibody comprising a heavy chain and/or a light chain sequence, wherein:

(a) the heavy chain sequence has at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99% or 100% sequence identity to the heavy chain sequence:

QVQLVESGGGVVQPGRSLRLDCKASGITFSNSGMHWVRQAPGKGLEWVAVIWY
DGSKRYYADSVKGRFTISRDN SKNTLFLQMNSLRAEDTAVYYCATNDDYWGQG
TLVTVSSASTKGPSVFPLAPCSRSTSESTAALGCLVKDYFPEPVTVSWNSGALTSG
VHTFPAVLQSSGLYSLSSVVTVPSSSLGKTKTYTCNVDPHKPSNTKVDKRVESKYGP
PCPPCPAPEFLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSQEDPEVQFNWYVD
GVEVHNAKTKPREEQFNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKGLPSSIEK
TISKAKGQPREPQVYTLPPSQQEEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPEN
NYKTTTPVLDSDGSFFLYSRLTVDKSRWQEGNVFSCSVMHEALHNHYTQKSLSL
SLGK (SEQ ID NO:274), or

(b) the light chain sequences has at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99% or 100% sequence identity to the light chain sequence:

EIVLTQSPATLSLSPGERATLSCRASQSVSSYLAWYQQKPGQAPRLLIYDASNRAT
GIPARFSGSGSGTDFTLTISSELPEDFAVYYCQQSSNWPRFTFGQGTKVEIKRTVAA
PSVFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQWKVDNALQSGNSQESVTEQD
SKDSTYSLSSSTLTLSKADYEKHKVYACEVTHQGLSPVTKSFNRGEC (SEQ ID
NO:275).

In some embodiments, the anti-PD-1 antibody is pembrolizumab (CAS Registry Number: 1374853-91-4). In a still further embodiment, provided is an isolated anti-PD-1 antibody comprising a heavy chain variable region comprising the heavy chain variable region amino acid sequence from SEQ ID NO:276 and/or a light chain variable region comprising the light chain variable region amino acid sequence from SEQ ID NO:277. In

a still further embodiment, provided is an isolated anti-PD-1 antibody comprising a heavy chain and/or a light chain sequence, wherein:

(a) the heavy chain sequence has at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least

5 99% or 100% sequence identity to the heavy chain sequence: QVQLVQSGVE
VKKPGASVKVSKASGYTFT NYMYWVRQA PGQGLEWMGG INPSNGGTNF
NEKFKNRVTLTTDSSTTTAY MELKSLQFDD TAVYYCARRDYRFDMGFDYW
GGGTTVTVSSASTKGPSVFP LAPCSRSTSE STAALGCLVKDYFPEPVTVS
WNSGALTSGVHTFPAVLQSS GLYSLSSVVT VPSSSLGKTYTCNVDPKPS
10 NTKVDKRVESKYGPPCPPCP APEFLGGPSV FLPPKPKDTLMISRTPEVT
CVVVDVSQEDPEVQFNWYVD GVEVHNAKTK PREEQFNSTYRVVSVLTVLH
QDWLNGKEYKCKVSNKGLPS SIEKTISKAK GQPREPQVYTLPPSQEEMTK
NQVSLTCLVKGFYPSDIAVE WESNGQPENN YKTTTPVLDSDGSFFLYSRL
TVDKSRWQEGNVFSCSVMHE ALHNHYTQKS LSLSLGK (SEQ ID NO:276), or

15 (b) the light chain sequences has at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99% or 100% sequence identity to the light chain sequence:

EIVLTQSPATLSLSPGERATLSCRASKGVSTSGYSYLHWYQQKPGQAPRLLIYLAS
YLESQVPAFSGSGSGTDFTLTISSELPEDFAVYYCQHSRDLP LTFGGGTKEIKR
20 TVAAPS VFIFPPSDEQLKSGTASVCLLN NFYPREAKVQWKVDNALQSGNSQESV
TEQDSKDSSTLSSTLTLSKADYEKHKVYACEVTHQGLSSPVT KSFNRGEC (SEQ
ID NO:277).

In some embodiments, the PDL1 binding antagonist is anti-PDL1 antibody. In some
embodiments, the anti-PDL1 binding antagonist is selected from the group consisting of
25 YW243.55.S70, MPDL3280A, MDX-1105, and MEDI4736. MDX-1105, also known as
BMS-936559, is an anti-PDL1 antibody described in WO2007/005874. Antibody
YW243.55.S70 (heavy and light chain variable region sequences shown in SEQ ID Nos.
20 and 21, respectively) is an anti-PDL1 described in WO 2010/077634 A1. MEDI4736
is an anti-PDL1 antibody described in WO2011/066389 and US2013/034559, each
30 incorporated herein by reference as if set forth in their entirety.

Examples of anti-PDL1 antibodies useful for the methods of this invention, and methods
for making thereof are described in PCT patent application WO 2010/077634 A1 and US
Patent No. 8,217,149, each incorporated herein by reference as if set forth in their entirety.

- In some embodiments, the PD-1 axis binding antagonist is an anti-PDL1 antibody. In some embodiments, the anti-PDL1 antibody is capable of inhibiting binding between PDL1 and PD-1 and/or between PDL1 and B7-1. In some embodiments, the anti-PDL1 antibody is a monoclonal antibody. In some embodiments, the anti-PDL1 antibody is an antibody fragment selected from the group consisting of Fab, Fab'-SH, Fv, scFv, and (Fab')₂ fragments. In some embodiments, the anti-PDL1 antibody is a humanized antibody. In some embodiments, the anti-PDL1 antibody is a human antibody. The anti-PDL1 antibodies useful in this invention, including compositions containing such antibodies, such as those described in WO 2010/077634 A1, may be used in combination with a T cell activating antigen binding molecule, and, optionally an anti-TIM3 antagonist antibody, to treat cancer. In some embodiments, the anti-PDL1 antibody comprises a heavy chain variable region comprising the amino acid sequence of SEQ ID NO:382 and a light chain variable region comprising the amino acid sequence of SEQ ID NO:383.
- 15 In one embodiment, the anti-PDL1 antibody contains a heavy chain variable region polypeptide comprising an HVR-H1, HVR-H2 and HVR-H3 sequence, wherein:
- (a) the HVR-H1 sequence is GFTFSX1SWIH (SEQ ID NO:283);
 - (b) the HVR-H2 sequence is AWIX2PYGGSX3YYADSVKG (SEQ ID NO:284);
 - (c) the HVR-H3 sequence is RHWPGGFDY (SEQ ID NO:285);
- 20 further wherein: X1 is D or G; X2 is S or L; X3 is T or S.
- In one specific aspect, X1 is D; X2 is S and X3 is T. In another aspect, the polypeptide further comprises variable region heavy chain framework sequences juxtaposed between the HVRs according to the formula: (HC-FR1)-(HVR-H1)-(HC-FR2)-(HVR-H2)-(HC-FR3)-(HVRH3)-(HC-FR4). In yet another aspect, the framework sequences are derived from human consensus framework sequences. In a further aspect, the framework sequences are VH subgroup III consensus framework. In a still further aspect, at least one of the framework sequences is the following:
- HC-FR1 is EVQLVESGGGLVQPGGSLRLSCAAS (SEQ ID NO:295)
- HC-FR2 is WVRQAPGKGLEWV (SEQ ID NO:296)
- 30 HC-FR3 is RFTISADTSKNTAYLQMNSLRAEDTAVYYCAR (SEQ ID NO:297)
- HC-FR4 is WGQGTLVTVSA (SEQ ID NO:298).
- In a still further aspect, the heavy chain polypeptide is further combined with a variable region light chain comprising an HVR-L1, HVR-L2 and HVR-L3, wherein:

(a) the HVR-L1 sequence is RASQX4X5X6TX7X8A (SEQ ID NO:286);

(b) the HVR-L2 sequence is SASX9LX10S, (SEQ ID NO:287);

(c) the HVR-L3 sequence is QQX11X12X13X14PX15T (SEQ ID NO:288);

further wherein: X4 is D or V; X5 is V or I; X6 is S or N; X7 is A or F; X8 is V or L; X9 is F or T; X10 is Y or A; X11 is Y, G, F, or S; X12 is L, Y, F or W; X13 is Y, N, A, T, G, F or I; X14 is H, V, P, T or I; X15 is A, W, R, P or T.

In a still further aspect, X4 is D; X5 is V; X6 is S; X7 is A; X8 is V; X9 is F; X10 is Y; X11 is Y; X12 is L; X13 is Y; X14 is H; X15 is A. In a still further aspect, the light chain further comprises variable region light chain framework sequences juxtaposed between the HVRs according to the formula: (LC-FR1)-(HVR-L1)-(LC-FR2)-(HVR-L2)-(LC-FR3)-(HVR-L3)-(LCFR4).

In a still further aspect, the framework sequences are derived from human consensus framework sequences. In a still further aspect, the framework sequences are VL kappa I consensus framework. In a still further aspect, at least one of the framework sequence is the following:

LC-FR1 is DIQMTQSPSSLSASVGDRVTITC (SEQ ID NO:300)

LC-FR2 is WYQQKPGKAPKLLIY (SEQ ID NO:301)

LC-FR3 is GVPSRFSGSGSGTDFTLTISLQPEDFATYYC (SEQ ID NO:302)

LC-FR4 is FGQGTKVEIKR (SEQ ID NO:303).

In another embodiment, provided is an isolated anti-PDL1 antibody or antigen binding fragment comprising a heavy chain and a light chain variable region sequence, wherein:

(a) the heavy chain comprises and HVR-H1, HVR-H2 and HVR-H3, wherein further:

(i) the HVR-H1 sequence is GFTFSX1SWIH (SEQ ID NO:283)

(ii) the HVR-H2 sequence is AWIX2PYGGSX3YYADSVKG (SEQ ID NO:284)

(iii) the HVR-H3 sequence is RHWPGGFDY (SEQ ID NO:285)

(b) the light chain comprises and HVR-L1, HVR-L2 and HVR-L3, wherein further:

(i) the HVR-L1 sequence is RASQX4X5X6TX7X8A (SEQ ID NO:286)

(ii) the HVR-L2 sequence is SASX9LX10S (SEQ ID NO:287)

(iii) the HVR-L3 sequence is QQX11X12X13X14PX15T (SEQ ID NO:288)

Further wherein: X1 is D or G; X2 is S or L; X3 is T or S; X4 is D or V; X5 is V or I; X6 is S or N; X7 is A or F; X8 is V or L; X9 is F or T; X10 is Y or A; X11 is Y, G, F, or S;

X12 is L, Y, F or W; X13 is Y, N, A, T, G, F or I; X14 is H, V, P, T or I; X15 is A, W, R, P or T.

In a specific aspect, X1 is D; X2 is S and X3 is T. In another aspect, X4 is D; X5 is V; X6 is S; X7 is A; X8 is V; X9 is F; X10 is Y; X11 is Y; X12 is L; X13 is Y; X14 is H;

5 X15 is A. In yet another aspect, X1 is D; X2 is S and X3 is T, X4 is D; X5 is V; X6 is S; X7 is A; X8 is V; X9 is F; X10 is Y; X11 is Y; X12 is L; X13 is Y; X14 is H and X15 is A.

In a further aspect, the heavy chain variable region comprises one or more framework sequences juxtaposed between the HVRs as: (HC-FR1)-(HVR-H1)-(HC-FR2)-(HVR-

10 H2)-(HCFR3)-(HVR-H3)-(HC-FR4), and the light chain variable regions comprises one

or more framework sequences juxtaposed between the HVRs as: (LC-FR1)-(HVR-L1)-(LC-FR2)-(HVR-L2)-(LC-FR3)-(HVR-L3)-(LC-FR4). In a still further aspect, the

framework sequences are derived from human consensus framework sequences. In a still

further aspect, the heavy chain framework sequences are derived from a Kabat subgroup I,

15 II, or III sequence. In a still further aspect, the heavy chain framework sequence is a VH

subgroup III consensus framework. In a still further aspect, one or more of the heavy chain framework sequences is the following:

HC-FR1 EVQLVESGGGLVQPGGSLRLSCAAS (SEQ ID NO:295)

HC-FR2 WVRQAPGKGLEWV (SEQ ID NO:296)

20 HC-FR3 RFTISADTSKNTAYLQMNSLRAEDTAVYYCAR (SEQ ID NO:297)

HC-FR4 WGQGTLVTVSA (SEQ ID NO:298).

In a still further aspect, the light chain framework sequences are derived from a Kabat

kappa I, II, II or IV subgroup sequence. In a still further aspect, the light chain framework

sequences are VL kappa I consensus framework. In a still further aspect, one or more of

25 the light chain framework sequences is the following:

LC-FR1 DIQMTQSPSSLSASVGDRVTITC (SEQ ID NO:300)

LC-FR2 WYQQKPGKAPKLLIY (SEQ ID NO:301)

LC-FR3 GVPSRFSGSGSGTDFTLTISSLQPEDFATYYC (SEQ ID NO:302)

LC-FR4 FGQGTVKVEIKR (SEQ ID NO:303).

30 In a still further specific aspect, the antibody further comprises a human or murine

constant region. In a still further aspect, the human constant region is selected from the

group consisting of IgG1, IgG2, IgG2, IgG3, IgG4. In a still further specific aspect, the

human constant region is IgG1. In a still further aspect, the murine constant region is

selected from the group consisting of IgG1, IgG2A, IgG2B, IgG3. In a still further aspect, the murine constant region is IgG2A. In a still further specific aspect, the antibody has reduced or minimal effector function. In a still further specific aspect the minimal effector function results from an “effectorless Fc mutation” or aglycosylation. In still a further embodiment, the effector-less Fc mutation is an N297A or D265A/N297A substitution in the constant region.

In yet another embodiment, provided is an anti-PDL1 antibody comprising a heavy chain and a light chain variable region sequence, wherein:

(a) the heavy chain further comprises an HVR-H1, HVR-H2 and an HVRH3 sequence having at least 85% sequence identity to GFTFSDSWIH (SEQ ID NO:289), AWISPYGGSTYYADSVKG (SEQ ID NO:290), and RHWPGGFDY (SEQ ID NO:291), respectively, or

(b) the light chain further comprises an HVR-L1, HVR-L2 and an HVR-L3 sequence having at least 85% sequence identity to RASQDVSTAVA (SEQ ID NO:292), SASFLYS (SEQ ID NO:293) and QQYLYHPAT (SEQ ID NO:294), respectively.

In a specific aspect, the sequence identity is 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or 100%. In another aspect, the heavy chain variable region comprises one or more framework sequences juxtaposed between the HVRs as: (HCFR1)-(HVR-H1)-(HC-FR2)-(HVR-H2)-(HC-FR3)-(HVR-H3)-(HC-FR4), and the light chain variable regions comprises one or more framework sequences juxtaposed between the HVRs as: (LC-FR1)-(HVR-L1)-(LC-FR2)-(HVR-L2)-(LC-FR3)-(HVR-L3)-(LC-FR4). In yet another aspect, the framework sequences are derived from human consensus framework sequences. In a still further aspect, the heavy chain framework sequences are derived from a Kabat subgroup I, II, or III sequence. In a still further aspect, the heavy chain framework sequence is a VH subgroup III consensus framework. In a still further aspect, one or more of the heavy chain framework sequences is the following:

HC-FR1 EVQLVESGGGLVQPGGSLRLSCAAS (SEQ ID NO:295)

HC-FR2 WVRQAPGKGGLEWV (SEQ ID NO:296)

HC-FR3 RFTISADTSKNTAYLQMNSLRAEDTAVYYCAR (SEQ ID NO:297)

HC-FR4 WGQGTLVTVSA (SEQ ID NO:298).

In a still further aspect, the light chain framework sequences are derived from a Kabat kappa I, II, III or IV subgroup sequence. In a still further aspect, the light chain framework

sequences are VL kappa I consensus framework. In a still further aspect, one or more of the light chain framework sequences is the following:

LC-FR1 DIQMTQSPSSLSASVGDRVITC (SEQ ID NO:300)

LC-FR2 WYQQKPGKAPKLLIY (SEQ ID NO:301)

5 LC-FR3 GVPSRFSGSGSGTDFTLTISLQPEDFATYYC (SEQ ID NO:302)

LC-FR4 FGQGTKVEIKR (SEQ ID NO:303).

In a still further specific aspect, the antibody further comprises a human or murine constant region. In a still further aspect, the human constant region is selected from the group consisting of IgG1, IgG2, IgG2, IgG3, IgG4. In a still further specific aspect, the human constant region is IgG1. In a still further aspect, the murine constant region is selected from the group consisting of IgG1, IgG2A, IgG2B, IgG3. In a still further aspect, the murine constant region is IgG2A. In a still further specific aspect, the antibody has reduced or minimal effector function. In a still further specific aspect the minimal effector function results from an "effectorless Fc mutation" or aglycosylation. In still a further embodiment, the effector-less Fc mutation is an N297A or D265A/N297A substitution in the constant region.

In a still further embodiment, provided is an isolated anti-PDL1 antibody comprising a heavy chain and a light chain variable region sequence, wherein:

(a) the heavy chain sequence has at least 85% sequence identity to the heavy chain sequence:

EVQLVESGGGLVQPGGSLRLSCAASGFTFSDSWIHWVRQAPGKGLEWVAWIS
PYGGSTYYADSVKGRFTISADTSKNTAYLQMNSLRAEDTAVYYCARRHWPGGF
DYWGQGTLLVTVSA (SEQ ID NO:382), or

(b) the light chain sequence has at least 85% sequence identity to the light chain sequence:

DIQMTQSPSSLSASVGDRVITCRASQDVSTAVAWYQQKPGKAPKLLIY
SASFLYSGVPSRFSGSGSGTDFTLTISLQPEDFATYYCQQYLYHPATFGQGTKVEI
KR (SEQ ID NO:383).

In a specific aspect, the sequence identity is 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or 100%. In another aspect, the heavy chain variable region comprises one or more framework sequences juxtaposed between the HVRs as: (HCFR1)-(HVR-H1)-(HC-FR2)-(HVR-H2)-(HC-FR3)-(HVR-H3)-(HC-FR4), and the light chain variable regions comprises one or more framework sequences

juxtaposed between the HVRs as: (LC-FR1)-(HVR-L1)-(LC-FR2)-(HVR-L2)-(LC-FR3)-(HVR-L3)-(LC-FR4). In yet another aspect, the framework sequences are derived from human consensus framework sequences. In a further aspect, the heavy chain framework sequences are derived from a Kabat subgroup I, II, or III sequence. In a still further aspect,

5 the heavy chain framework sequence is a VH subgroup III consensus framework. In a still further aspect, one or more of the heavy chain framework sequences is the following:

HC-FR1 EVQLVESGGGLVQPGGSLRLSCAAS (SEQ ID NO:295)

HC-FR2 WVRQAPGKGGLEWV (SEQ ID NO:296)

HC-FR3 RFTISADTSKNTAYLQMNSLRAEDTAVYYCAR (SEQ ID NO:297)

10 HC-FR4 WGQGTLVTVSA (SEQ ID NO:298).

In a still further aspect, the light chain framework sequences are derived from a Kabat kappa I, II, II or IV subgroup sequence. In a still further aspect, the light chain framework sequences are VL kappa I consensus framework. In a still further aspect, one or more of the light chain framework sequences is the following:

15 LC-FR1 DIQMTQSPSSLSASVGDRVTITC (SEQ ID NO:300)

LC-FR2 WYQQKPGKAPKLLIY (SEQ ID NO:301)

LC-FR3 GVPSRFSGSGSGTDFTLTISLQPEDFATYYC (SEQ ID NO:302)

LC-FR4 FGQGTEKVEIKR (SEQ ID NO:303).

In a still further specific aspect, the antibody further comprises a human or murine

20 constant region. In a still further aspect, the human constant region is selected from the group consisting of IgG1, IgG2, IgG2, IgG3, IgG4. In a still further specific aspect, the human constant region is IgG1. In a still further aspect, the murine constant region is selected from the group consisting of IgG1, IgG2A, IgG2B, IgG3. In a still further aspect, the murine constant region is IgG2A. In a still further specific aspect, the antibody has

25 reduced or minimal effector function. In a still further specific aspect, the minimal effector function results from production in prokaryotic cells. In a still further specific aspect the minimal effector function results from an "effector-less Fc mutation" or aglycosylation. In still a further embodiment, the effector-less Fc mutation is an N297A or D265A/N297A substitution in the constant region.

30 In another further embodiment, provided is an isolated anti-PDL1 antibody comprising a heavy chain and a light chain variable region sequence, wherein:

(a) the heavy chain sequence has at least 85% sequence identity to the heavy chain

sequence:EVQLVESGGGLVQPGGSLRLSCAASGFTFSDSWIHWVRQAPGKGLEWV
AWISPYGGSTYYADSVKGRFTISADTSKNTAYLQMNSLRAEDTAVYYCARRHW
PGGFDYWGQGTLLTVSS (SEQ ID NO:280), or

(b) the light chain sequence has at least 85% sequence identity to the light chain sequence:

- 5 DIQMTQSPSSLSASVGDRVTITCRASQDVSTAVAWYQQKPGKAPKLLIY
SASFLYSGVPSRFSGSGSGTDFTLTISLQPEDFATYYCQQYLYHPATFGQGTKVEI
KR (SEQ ID NO:383).

In a still further embodiment, provided is an isolated anti-PDL1 antibody comprising a heavy chain and a light chain variable region sequence, wherein:

- 10 (a) the heavy chain sequence has at least 85% sequence identity to the heavy chain sequence:

EVQLVESGGGLVQPGGSLRLSCAASGFTFSDSWIHWVRQAPGKGLEWVAVISPY
GGSTYYADSVKGRFTISADTSKNTAYLQMNSLRAEDTAVYYCARRHWPGGFDY
WGQGTLLTVSSASTK (SEQ ID NO:281), or

- 15 (b) the light chain sequences has at least 85% sequence identity to the light chain sequence:

DIQMTQSPSSLSASVGDRVTITCRASQDVSTAVAWYQQKPGKAPKLLIYSASF
LYSGVPSRFSGSGSGTDFTLTISLQPEDFATYYCQQYLYHPATFGQGTKVEIKR
(SEQ ID NO:282).

- 20 In a specific aspect, the sequence identity is 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or 100%. In another aspect, the heavy chain variable region comprises one or more framework sequences juxtaposed between the HVRs as: (HCFR1)-(HVR-H1)-(HC-FR2)-(HVR-H2)-(HC-FR3)-(HVR-H3)-(HC-FR4), and the light chain variable regions comprises one or more framework sequences juxtaposed
25 between the HVRs as: (LC-FR1)-(HVR-L1)-(LC-FR2)-(HVR-L2)-(LC-FR3)-(HVR-L3)-(LC-FR4). In yet another aspect, the framework sequences are derived from human consensus framework sequences. In a further aspect, the heavy chain framework sequences are derived from a Kabat subgroup I, II, or III sequence. In a still further aspect, the heavy chain framework sequence is a VH subgroup III consensus framework. In a still
30 further aspect, one or more of the heavy chain framework sequences is the following:

HC-FR1 EVQLVESGGGLVQPGGSLRLSCAAS (SEQ ID NO:295)

HC-FR2 WVRQAPGKGLEWV (SEQ ID NO:296)

HC-FR3 RFTISADTSKNTAYLQMNSLRAEDTAVYYCAR (SEQ ID NO:297)

HC-FR4 WGQGTLVTVSS (SEQ ID NO:299).

In a still further aspect, the light chain framework sequences are derived from a Kabat kappa I, II, III or IV subgroup sequence. In a still further aspect, the light chain framework sequences are VL kappa I consensus framework. In a still further aspect, one or more of the light chain framework sequences is the following:

LC-FR1 DIQMTQSPSSLSASVGDRVTITC (SEQ ID NO:300)

LC-FR2 WYQQKPGKAPKLLIY (SEQ ID NO:301)

LC-FR3 GVPSRFSGSGSGTDFTLTISSLQPEDFATYYC (SEQ ID NO:302)

LC-FR4 FGQGTKVEIKR (SEQ ID NO:303).

- 10 In a still further specific aspect, the antibody further comprises a human or murine constant region. In a still further aspect, the human constant region is selected from the group consisting of IgG1, IgG2, IgG2, IgG3, IgG4. In a still further specific aspect, the human constant region is IgG1. In a still further aspect, the murine constant region is selected from the group consisting of IgG1, IgG2A, IgG2B, IgG3. In a still further aspect, the murine constant region is IgG2A. In a still further specific aspect, the antibody has reduced or minimal effector function. In a still further specific aspect, the minimal effector function results from production in prokaryotic cells. In a still further specific aspect the minimal effector function results from an "effector-less Fc mutation" or aglycosylation. In still a further embodiment, the effector-less Fc mutation is an N297A or D265A/N297A substitution in the constant region.

- 20 In yet another embodiment, the anti-PDL1 antibody is MPDL3280A (CAS Registry Number: 1422185-06-5). In a still further embodiment, provided is an isolated anti-PDL1 antibody comprising a heavy chain variable region comprising the heavy chain variable region amino acid sequence from SEQ ID NO:24 or SEQ ID NO:28 and/or a light chain variable region comprising the light chain variable region amino acid sequence from SEQ ID NO:21. In a still further embodiment, provided is an isolated anti-PDL1 antibody comprising a heavy chain and/or a light chain sequence, wherein:

- 25 (a) the heavy chain sequence has at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99% or 100% sequence identity to the heavy chain sequence:

EVQLVESGGGLVQPGGSLRLSCAASGFTFSDSWIHWVRQAPGKGLEWVAWISPY
GGSTYYADSVKGRFTISADTSKNTAYLQMNSLRAEDTAVYYCARRHWPGGFDY
WGQGTLVTVSSASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSG

ALTSGVHTFPAVLQSSGLYSLSSVVTVPSSSLGTQTYICNVNHKPSNTKVDKKVE
 PKSCDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVDVSHEDPE
 VKFNWYVDGVEVHNAKTKPREEQYASTYRVVSVLTVLHQDWLNGKEYKCKV
 NKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVE
 5 WESNGQPENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQQGNVFCFSVMHEALH
 NHYTQKSLSLSPG (SEQ ID NO:278), or

(b) the light chain sequences has at least 85%, at least 90%, at least 91%, at least 92%, at
 least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least
 99% or 100% sequence identity to the light chain sequence:

10 DIQMTQSPSSLSASVGDRVTITCRASQDVSTAVAWYQQKPGKAPKLLIYSASFLY
 SGVPSRFSGSGSGTDFTLTISLQPEDFATYYCQQYLYHPATFGQGTKVEIKRTVA
 APSVFIFPPSDEQLKSGTASVCLLNNFYPRKAKVQWKVDNALQSGNSQESVTEQ
 DSKDSTYLSSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC (SEQ ID
 NO:279).

15 In a still further embodiment, the invention provides for compositions comprising any
 of the above described anti-PDL1 antibodies in combination with at least one
 pharmaceutically acceptable carrier.

In a still further embodiment, provided is an isolated nucleic acid encoding a light
 chain or a heavy chain variable region sequence of an anti-PDL1 antibody, wherein:

20 (a) the heavy chain further comprises an HVR-H1, HVR-H2 and an HVR-H3 sequence
 having at least 85% sequence identity to GFTFSDSWIH (SEQ ID NO:289),
 AWISPYGGSTYYADSVKG (SEQ ID NO:290) and RHWPGGFDY (SEQ ID NO:291),
 respectively, and

(b) the light chain further comprises an HVR-L1, HVR-L2 and an HVR-L3 sequence
 25 having at least 85% sequence identity to RASQDVSTAVA (SEQ ID NO:292),
 SASFLYS (SEQ ID NO:293) and QQYLYHPAT (SEQ ID NO:294), respectively.

In a specific aspect, the sequence identity is 86%, 87%, 88%, 89%, 90%, 91%, 92%,
 93%, 94%, 95%, 96%, 97%, 98%, 99% or 100%. In another aspect, the heavy chain variable
 region comprises one or more framework sequences juxtaposed between the HVRs as:

30 (HC-FR1)-(HVR-H1)-(HC-FR2)-(HVR-H2)-(HC-FR3)-(HVR-H3)-(HC-FR4), and the
 light chain variable regions comprises one or more framework sequences juxtaposed
 between the HVRs as: (LC-FR1)-(HVR-L1)-(LC-FR2)-(HVR-L2)-(LC-FR3)-(HVR-L3)-
 (LC-FR4). In yet another aspect, the framework sequences are derived from human

consensus framework sequences. In a further aspect, the heavy chain framework sequences are derived from a Kabat subgroup I, II, or III sequence. In a still further aspect, the heavy chain framework sequence is a VH subgroup III consensus framework. In a still further aspect, one or more of the heavy chain framework sequences is the following:

- 5 HC-FR1 EVQLVESGGGLVQPGGSLRLSCAAS (SEQ ID NO:295)
- HC-FR2 WVRQAPGKGGLEWV (SEQ ID NO:296)
- HC-FR3 RFTISADTSKNTAYLQMNSLRAEDTAVYYCAR (SEQ ID NO:297)
- HC-FR4 WGQGTLVTVSA (SEQ ID NO:298).

In a still further aspect, the light chain framework sequences are derived from a Kabat
 10 kappa I, II, II or IV subgroup sequence. In a still further aspect, the light chain framework sequences are VL kappa I consensus framework. In a still further aspect, one or more of the light chain framework sequences is the following:

- LC-FR1 DIQMTQSPSSLSASVGDRVTITC (SEQ ID NO:300)
- LC-FR2 WYQQKPGKAPKLLIY (SEQ ID NO:301)
- 15 LC-FR3 GVPSRFSGSGSGTDFLTISLQPEDFATYYC (SEQ ID NO:302)
- LC-FR4 FGQGTKVEIKR (SEQ ID NO:303).

In a still further specific aspect, the antibody described herein (such as an anti-PD-1 antibody, an anti-PDL1 antibody, or an anti-PDL2 antibody) further comprises a human or murine constant region. In a still further aspect, the human constant region is selected
 20 from the group consisting of IgG1, IgG2, IgG2, IgG3, IgG4. In a still further specific aspect, the human constant region is IgG1. In a still further aspect, the murine constant region is selected from the group consisting of IgG1, IgG2A, IgG2B, IgG3. In a still further aspect, the murine constant region is IgG2A. In a still further specific aspect, the antibody has reduced or minimal effector function. In a still further specific aspect, the
 25 minimal effector function results from production in prokaryotic cells. In a still further specific aspect the minimal effector function results from an "effector-less Fc mutation" or aglycosylation. In still a further aspect, the effector-less Fc mutation is an N297A or D265A/N297A substitution in the constant region.

In a still further aspect, provided herein are nucleic acids encoding any of the
 30 antibodies described herein. In some embodiments, the nucleic acid further comprises a vector suitable for expression of the nucleic acid encoding any of the previously described anti-PDL1, anti-PD-1, or anti-PDL2 antibodies. In a still further specific aspect, the vector further comprises a host cell suitable for expression of the nucleic acid. In a still

further specific aspect, the host cell is a eukaryotic cell or a prokaryotic cell. In a still further specific aspect, the eukaryotic cell is a mammalian cell, such as Chinese Hamster Ovary (CHO).

The antibody or antigen binding fragment thereof, may be made using methods

- 5 known in the art, for example, by a process comprising culturing a host cell containing nucleic acid encoding any of the previously described anti-PDL1, anti-PD-1, or anti-PDL2 antibodies or antigen-binding fragment in a form suitable for expression, under conditions suitable to produce such antibody or fragment, and recovering the antibody or fragment.

- 10 In some embodiments, the isolated anti-PDL1 antibody is aglycosylated.

Glycosylation of antibodies is typically either N-linked or O-linked. N-linked refers to the attachment of the carbohydrate moiety to the side chain of an asparagine residue. The tripeptide sequences asparagine-X-serine and asparagine-X-threonine, where X is any amino acid except proline, are the recognition sequences for enzymatic attachment of the
15 carbohydrate moiety to the asparagine side chain. Thus, the presence of either of these tripeptide sequences in a polypeptide creates a potential glycosylation site. O-linked glycosylation refers to the attachment of one of the sugars N-aceylgalactosamine, galactose, or xylose to a hydroxyamino acid, most commonly serine or threonine, although 5-hydroxyproline or 5-hydroxylysine may also be used. Removal of
20 glycosylation sites from an antibody is conveniently accomplished by altering the amino acid sequence such that one of the above-described tripeptide sequences (for N-linked glycosylation sites) is removed. The alteration may be made by substitution of an asparagine, serine or threonine residue within the glycosylation site another amino acid residue (e.g., glycine, alanine or a conservative substitution).

- 25 In any of the embodiments herein, the isolated anti-PDL1 antibody can bind to a human PDL1, for example a human PDL1 as shown in UniProtKB/Swiss-Prot Accession No.Q9NZQ7.1, or a variant thereof.

In a still further embodiment, the invention provides for a composition comprising an anti-PDL1, an anti-PD-1, or an anti-PDL2 antibody or antigen binding fragment thereof
30 as provided herein and at least one pharmaceutically acceptable carrier. In some embodiments, the anti-PDL1, anti-PD-1, or anti-PDL2 antibody or antigen binding fragment thereof administered to the individual is a composition comprising one or more pharmaceutically acceptable carrier.

Any of the pharmaceutically acceptable carriers described herein or known in the art may be used.

In some embodiments, the anti-PDL1 antibody described herein is in a formulation comprising the antibody at an amount of about 60 mg/mL, histidine acetate in a concentration of about 20 mM, sucrose in a concentration of about 120 mM, and polysorbate (e.g., polysorbate 20) in a concentration of 0.04% (w/v), and the formulation has a pH of about 5.8. In some embodiments, the anti-PDL1 antibody described herein is in a formulation comprising the antibody in an amount of about 125 mg/mL, histidine acetate in a concentration of about 20 mM, sucrose is in a concentration of about 240 mM, and polysorbate (e.g., polysorbate 20) in a concentration of 0.02% (w/v), and the formulation has a pH of about 5.5.

Exemplary TIM3 Antagonists for use in the invention

Provided herein are methods for treating or delaying progression of cancer in an individual comprising administering to the individual an effective amount of a T cell activating bispecific antigen binding molecule, a PD-1 axis binding antagonist, and a TIM-3 antagonist. In one embodiment, the TIM-3 antagonist is an anti-TIM-3 antibody. In some embodiments, the anti-TIM3 induces internalization of TIM3 expressed on a cell of at least 45% after 120 Minutes at 37 °C as determined by FACS analysis. The cell is, e.g., a RPMI8226 cells (ATCC ® CCL-155™). In one embodiment, the antibody induces internalization of TIM3 on TIM3 expressing RPMI8226 cells (ATCC ® CCL-155™) of at least 55% after 120 Minutes at 37 °C as determined by FACS analysis. In one embodiment, the antibody induces internalization of TIM3 on TIM3 expressing RPMI8226 cells (ATCC ® CCL-155™) of at least 60% after 240 Minutes at 37 °C as determined by FACS analysis. In one embodiment, the antibody induces internalization of TIM3 on TIM3 expressing RPMI8226 cells (ATCC ® CCL-155™) of at least 65% after 240 Minutes at 37 °C as determined by FACS analysis.

In some embodiments, the anti-TIM3 antibody competes for binding to TIM3 with an anti-Tim3 antibody comprising the VH and VL of Tim3_0016. In some embodiments, the anti-TIM3 antibody binds to a human and cynomolgous TIM3. In some embodiments, the anti-TIM3 antibody shows as an immunoconjugate a cytotoxic activity on TIM3 expressing cells. In one such embodiment, the immunoconjugate has a relative

IC50 value of the cytotoxic activity as *Pseudomonas* exotoxin A conjugate on RPMI-8226 cells of 0.1 or lower. In one embodiment, the anti-TIM3 antibody induces interferon-gamma release as determined by MLR assay.

In certain embodiments, the anti-TIM3 antibody binds to a human and cynomolgous

5 TIM3 and induces interferon-gamma release as determined by a MLR assay.

In one embodiment, the anti-TIM3 antibody comprises at least one, two, three, four, five, or six HVRs selected from (a) HVR-H1 comprising the amino acid sequence of SEQ ID NO:304; (b) HVR-H2 comprising the amino acid sequence of SEQ ID NO:305; (c) HVR-H3 comprising the amino acid sequence of SEQ ID NO:306; (d) HVR-L1 comprising the
10 amino acid sequence of SEQ ID NO:307; or HVR-L1 comprising the amino acid sequence of SEQ ID NO:314; HVR-L1 comprising the amino acid sequence of SEQ ID NO:315; (e) HVR-L2 comprising the amino acid sequence of SEQ ID NO:308; and (f) HVR-L3 comprising the amino acid sequence of SEQ ID NO:309.

In one embodiment, the anti-TIM3 antibody comprises (a) HVR-H1 comprising the
15 amino acid sequence of SEQ ID NO:304; (b) HVR-H2 comprising the amino acid sequence of SEQ ID NO:305; (c) HVR-H3 comprising the amino acid sequence of SEQ ID NO:306; (d) HVR-L1 comprising the amino acid sequence of SEQ ID NO:307; or HVR-L1 comprising the amino acid sequence of SEQ ID NO:314; or HVR-L1 comprising the amino acid sequence of SEQ ID NO:315; (e) HVR-L2 comprising the
20 amino acid sequence of SEQ ID NO:308; and (f) HVR-L3 comprising the amino acid sequence of SEQ ID NO:309.

In one embodiment, the anti-TIM3 antibody comprises (a) HVR-H1 comprising the amino acid sequence of SEQ ID NO:304; (b) HVR-H2 comprising the amino acid
25 sequence of SEQ ID NO:305; (c) HVR-H3 comprising the amino acid sequence of SEQ ID NO:306; (d) HVR-L1 comprising the amino acid sequence of SEQ ID NO:307; (e) HVR-L2 comprising the amino acid sequence of SEQ ID NO:308; and (f) HVR-L3 comprising the amino acid sequence of SEQ ID NO:309.

In one embodiment, the anti-TIM3 antibody comprises (a) HVR-H1 comprising the
30 amino acid sequence of SEQ ID NO:304; (b) HVR-H2 comprising the amino acid sequence of SEQ ID NO:305; (c) HVR-H3 comprising the amino acid sequence of SEQ ID NO:306; (d) HVR-L1 comprising the amino acid sequence of SEQ ID NO:314; (e)

HVR-L2 comprising the amino acid sequence of SEQ ID NO:308; and (f) HVR-L3 comprising the amino acid sequence of SEQ ID NO:309.

In one embodiment, the anti-TIM3 antibody comprises (a) HVR-H1 comprising the amino acid sequence of SEQ ID NO:304; (b) HVR-H2 comprising the amino acid

- 5 sequence of SEQ ID NO:305; (c) HVR-H3 comprising the amino acid sequence of SEQ ID NO:306; (d) HVR-L1 comprising the amino acid sequence of SEQ ID NO:315; (e) HVR-L2 comprising the amino acid sequence of SEQ ID NO:308; and (f) HVR-L3 comprising the amino acid sequence of SEQ ID NO:309.

- 10 In one embodiment, the anti-TIM3 antibody comprises (a) a VH domain comprising at least one, at least two, or all three VH HVR sequences selected from (i) HVR-H1 comprising the amino acid sequence of SEQ ID NO:304, (ii) HVR-H2 comprising the amino acid sequence of SEQ ID NO:305, and (iii) HVR-H3 comprising an amino acid sequence selected from SEQ ID NO:306; and (b) a VL domain comprising at least one, at
15 least two, or all three VL HVR sequences selected from (i) HVR-L1 comprising the amino acid sequence of SEQ ID NO:307; or HVR-L1 comprising the amino acid sequence of SEQ ID NO:314; or HVR-L1 comprising the amino acid sequence of SEQ ID NO:315; (ii) HVR-L2 comprising the amino acid sequence of SEQ ID NO:308 and (c) HVR-L3 comprising the amino acid sequence of SEQ ID NO:309.

- 20 In one embodiment, the anti-TIM3 antibody comprises (a) a VH domain comprising (i) HVR-H1 comprising the amino acid sequence of SEQ ID NO:304, (ii) HVR-H2 comprising the amino acid sequence of SEQ ID NO:305, and (iii) HVR-H3 comprising an amino acid sequence selected from SEQ ID NO:306; and (b) a VL domain comprising (i) HVR-L1 comprising the amino acid sequence of SEQ ID NO:307; or HVR-L1
25 comprising the amino acid sequence of SEQ ID NO:314; or HVR-L1 comprising the amino acid sequence of SEQ ID NO:315; (ii) HVR-L2 comprising the amino acid sequence of SEQ ID NO:308 and (iii) HVR-L3 comprising the amino acid sequence of SEQ ID NO:309.

In one embodiment, the anti-TIM3 antibody comprises (a) a VH domain comprising (i)

- 30 HVR-H1 comprising the amino acid sequence of SEQ ID NO:304, (ii) HVR-H2 comprising the amino acid sequence of SEQ ID NO:305, and (iii) HVR-H3 comprising an amino acid sequence selected from SEQ ID NO:306; and (b) a VL domain comprising (i) HVR-L1 comprising the amino acid sequence of SEQ ID NO:307; (ii) HVR-L2

comprising the amino acid sequence of SEQ ID NO:308 and (iii) HVR-L3 comprising the amino acid sequence of SEQ ID NO:309.

In one embodiment, the anti-TIM3 antibody comprises (a) a VH domain comprising (i) HVR-H1 comprising the amino acid sequence of SEQ ID NO:304, (ii) HVR-H2

- 5 comprising the amino acid sequence of SEQ ID NO:305, and (iii) HVR-H3 comprising an amino acid sequence selected from SEQ ID NO:306; and (b) a VL domain comprising (i) HVR-L1 comprising the amino acid sequence of SEQ ID NO:314; (ii) HVR-L2

comprising the amino acid sequence of SEQ ID NO:308 and (iii) HVR-L3 comprising the amino acid sequence of SEQ ID NO:309.

- 10 In one embodiment, the anti-TIM3 antibody comprises (a) a VH domain comprising (i) HVR-H1 comprising the amino acid sequence of SEQ ID NO:304, (ii) HVR-H2 comprising the amino acid sequence of SEQ ID NO:305, and (iii) HVR-H3 comprising an amino acid sequence selected from SEQ ID NO:306; and (b) a VL domain comprising (i) HVR-L1 comprising the amino acid sequence of SEQ ID NO:315; (ii) HVR-L2

- 15 comprising the amino acid sequence of SEQ ID NO:308 and (iii) HVR-L3 comprising the amino acid sequence of SEQ ID NO:309.

In one embodiment such anti-TIM3 antibody comprises

- i) comprises a VH sequence of SEQ ID NO:310 and a VL sequence of SEQ ID NO:311;
- 20 ii) comprises a VH sequence of SEQ ID NO:312 and a VL sequence of SEQ ID NO:313;
- iii) or humanized variant of the VH and VL of the antibody under i) or ii).

In one embodiment, the anti-TIM3 antibody comprises at least one, two, three, four, five, or six HVRs selected from (a) HVR-H1 comprising the amino acid sequence of SEQ ID

- 25 NO:316; (b) HVR-H2 comprising the amino acid sequence of SEQ ID NO:317; (c) HVR-H3 comprising the amino acid sequence of SEQ ID NO:318; (d) HVR-L1 comprising the amino acid sequence of SEQ ID NO:319; (e) HVR-L2 comprising the amino acid sequence of SEQ ID NO:320; and (f) HVR-L3 comprising the amino acid sequence of SEQ ID NO:321.

- 30 In one embodiment, the anti-TIM3 antibody comprises (a) HVR-H1 comprising the amino acid sequence of SEQ ID NO:316; (b) HVR-H2 comprising the amino acid sequence of SEQ ID NO:317; (c) HVR-H3 comprising the amino acid sequence of SEQ ID NO:318; (d) HVR-L1 comprising the amino acid sequence of SEQ ID NO:319; (e)

HVR-L2 comprising the amino acid sequence of SEQ ID NO:320; and (f) HVR-L3 comprising the amino acid sequence of SEQ ID NO:321.

In one embodiment, the anti-TIM3 antibody comprises (a) a VH domain comprising at least one, at least two, or all three VH HVR sequences selected from (i) HVR-H1 comprising the amino acid sequence of SEQ ID NO:316, (ii) HVR-H2 comprising the amino acid sequence of SEQ ID NO:317, and (iii) HVR-H3 comprising an amino acid sequence selected from SEQ ID NO:318; and (b) a VL domain comprising at least one, at least two, or all three VL HVR sequences selected from (i) HVR-L1 comprising the amino acid sequence of SEQ ID NO:319; (ii) HVR-L2 comprising the amino acid sequence of SEQ ID NO:320 and (c) HVR-L3 comprising the amino acid sequence of SEQ ID NO:321.

In one embodiment, the anti-TIM3 antibody comprises (a) a VH domain comprising (i) HVR-H1 comprising the amino acid sequence of SEQ ID NO:316, (ii) HVR-H2 comprising the amino acid sequence of SEQ ID NO:317, and (iii) HVR-H3 comprising an amino acid sequence selected from SEQ ID NO:318; and (b) a VL domain comprising (i) HVR-L1 comprising the amino acid sequence of SEQ ID NO:319; (ii) HVR-L2 comprising the amino acid sequence of SEQ ID NO:320 and (iii) HVR-L3 comprising the amino acid sequence of SEQ ID NO:321.

In one embodiment such anti-TIM3 antibody comprises

- i) comprises a VH sequence of SEQ ID NO:322 and a VL sequence of SEQ ID NO:323;
- ii) or humanized variant of the VH and VL of the antibody under i).

In one embodiment, the anti-TIM3 antibody comprises at least one, two, three, four, five, or six HVRs selected from (a) HVR-H1 comprising the amino acid sequence of SEQ ID NO:324; (b) HVR-H2 comprising the amino acid sequence of SEQ ID NO:325; (c) HVR-H3 comprising the amino acid sequence of SEQ ID NO:326; (d) HVR-L1 comprising the amino acid sequence of SEQ ID NO:327; (e) HVR-L2 comprising the amino acid sequence of SEQ ID NO:328; and (f) HVR-L3 comprising the amino acid sequence of SEQ ID NO:329.

In one embodiment, the anti-TIM3 antibody comprises (a) HVR-H1 comprising the amino acid sequence of SEQ ID NO:324; (b) HVR-H2 comprising the amino acid sequence of SEQ ID NO:325; (c) HVR-H3 comprising the amino acid sequence of SEQ ID NO:326; (d) HVR-L1 comprising the amino acid sequence of SEQ ID NO:327; (e)

HVR-L2 comprising the amino acid sequence of SEQ ID NO:328; and (f) HVR-L3 comprising the amino acid sequence of SEQ ID NO:329.

In one embodiment, the anti-TIM3 antibody comprises (a) a VH domain comprising at least one, at least two, or all three VH HVR sequences selected from (i) HVR-H1
5 comprising the amino acid sequence of SEQ ID NO:324, (ii) HVR-H2 comprising the amino acid sequence of SEQ ID NO:325, and (iii) HVR-H3 comprising an amino acid sequence selected from SEQ ID NO:326; and (b) a VL domain comprising at least one, at least two, or all three VL HVR sequences selected from (i) HVR-L1 comprising the amino acid sequence of SEQ ID NO:327; (ii) HVR-L2 comprising the amino acid
10 sequence of SEQ ID NO:328 and (c) HVR-L3 comprising the amino acid sequence of SEQ ID NO:329.

In one embodiment, the anti-TIM3 antibody comprises (a) a VH domain comprising (i) HVR-H1 comprising the amino acid sequence of SEQ ID NO:324, (ii) HVR-H2 comprising the amino acid sequence of SEQ ID NO:325, and (iii) HVR-H3 comprising an
15 amino acid sequence selected from SEQ ID NO:326; and (b) a VL domain comprising (i) HVR-L1 comprising the amino acid sequence of SEQ ID NO:327; (ii) HVR-L2 comprising the amino acid sequence of SEQ ID NO:328 and (iii) HVR-L3 comprising the amino acid sequence of SEQ ID NO:329.

In one embodiment such anti-TIM3 antibody comprises
20 i) comprises a VH sequence of SEQ ID NO:330 and a VL sequence of SEQ ID NO:331;
ii) or humanized variant of the VH and VL of the antibody under i).

In one embodiment, the anti-TIM3 antibody comprises at least one, two, three, four, five, or six HVRs selected from (a) HVR-H1 comprising the amino acid sequence of SEQ ID
25 NO:332; (b) HVR-H2 comprising the amino acid sequence of SEQ ID NO:333; (c) HVR-H3 comprising the amino acid sequence of SEQ ID NO:334; (d) HVR-L1 comprising the amino acid sequence of SEQ ID NO:335; (e) HVR-L2 comprising the amino acid sequence of SEQ ID NO:336; and (f) HVR-L3 comprising the amino acid sequence of SEQ ID NO:337.

30 In one embodiment, the anti-TIM3 antibody comprises (a) HVR-H1 comprising the amino acid sequence of SEQ ID NO:332; (b) HVR-H2 comprising the amino acid sequence of SEQ ID NO:333; (c) HVR-H3 comprising the amino acid sequence of SEQ ID NO:334; (d) HVR-L1 comprising the amino acid sequence of SEQ ID NO:335; (e)

HVR-L2 comprising the amino acid sequence of SEQ ID NO:336; and (f) HVR-L3 comprising the amino acid sequence of SEQ ID NO:337.

In one embodiment, the anti-TIM3 antibody comprises (a) a VH domain comprising at least one, at least two, or all three VH HVR sequences selected from (i) HVR-H1
5 comprising the amino acid sequence of SEQ ID NO:332, (ii) HVR-H2 comprising the amino acid sequence of SEQ ID NO:333, and (iii) HVR-H3 comprising an amino acid sequence selected from SEQ ID NO:334; and (b) a VL domain comprising at least one, at least two, or all three VL HVR sequences selected from (i) HVR-L1 comprising the amino acid sequence of SEQ ID NO:335; (ii) HVR-L2 comprising the amino acid
10 sequence of SEQ ID NO:336 and (c) HVR-L3 comprising the amino acid sequence of SEQ ID NO:337.

In one embodiment, the anti-TIM3 antibody comprises (a) a VH domain comprising (i) HVR-H1 comprising the amino acid sequence of SEQ ID NO:332, (ii) HVR-H2 comprising the amino acid sequence of SEQ ID NO:333, and (iii) HVR-H3 comprising an
15 amino acid sequence selected from SEQ ID NO:334; and (b) a VL domain comprising (i) HVR-L1 comprising the amino acid sequence of SEQ ID NO:335; (ii) HVR-L2 comprising the amino acid sequence of SEQ ID NO:336 and (iii) HVR-L3 comprising the amino acid sequence of SEQ ID NO:337.

In one embodiment such anti-TIM3 antibody comprises
20 i) comprises a VH sequence of SEQ ID NO:338 and a VL sequence of SEQ ID NO:339;
ii) or humanized variant of the VH and VL of the antibody under i).

In one aspect, the invention provides an anti-TIM3 antibody comprising at least one, two, three, four, five, or six HVRs selected from (a) HVR-H1 comprising the amino acid
25 sequence of SEQ ID NO:340; (b) HVR-H2 comprising the amino acid sequence of SEQ ID NO:341; (c) HVR-H3 comprising the amino acid sequence of SEQ ID NO:342; (d) HVR-L1 comprising the amino acid sequence of SEQ ID NO:343; (e) HVR-L2 comprising the amino acid sequence of SEQ ID NO:344; and (f) HVR-L3 comprising the amino acid sequence of SEQ ID NO:345.

30 In one embodiment, the anti-TIM3 antibody comprises (a) HVR-H1 comprising the amino acid sequence of SEQ ID NO:340; (b) HVR-H2 comprising the amino acid sequence of SEQ ID NO:341; (c) HVR-H3 comprising the amino acid sequence of SEQ ID NO:342; (d) HVR-L1 comprising the amino acid sequence of SEQ ID NO:343; (e)

HVR-L2 comprising the amino acid sequence of SEQ ID NO:344; and (f) HVR-L3 comprising the amino acid sequence of SEQ ID NO:345.

In one embodiment, the anti-TIM3 antibody comprises (a) a VH domain comprising at least one, at least two, or all three VH HVR sequences selected from (i) HVR-H1 comprising the amino acid sequence of SEQ ID NO:340, (ii) HVR-H2 comprising the amino acid sequence of SEQ ID NO:341, and (iii) HVR-H3 comprising an amino acid sequence selected from SEQ ID NO:342; and (b) a VL domain comprising at least one, at least two, or all three VL HVR sequences selected from (i) HVR-L1 comprising the amino acid sequence of SEQ ID NO:343; (ii) HVR-L2 comprising the amino acid sequence of SEQ ID NO:344 and (c) HVR-L3 comprising the amino acid sequence of SEQ ID NO:345.

In one embodiment, the anti-TIM3 antibody comprises (a) a VH domain comprising (i) HVR-H1 comprising the amino acid sequence of SEQ ID NO:340, (ii) HVR-H2 comprising the amino acid sequence of SEQ ID NO:341, and (iii) HVR-H3 comprising an amino acid sequence selected from SEQ ID NO:342; and (b) a VL domain comprising (i) HVR-L1 comprising the amino acid sequence of SEQ ID NO:343; (ii) HVR-L2 comprising the amino acid sequence of SEQ ID NO:344 and (iii) HVR-L3 comprising the amino acid sequence of SEQ ID NO:345.

In one embodiment such anti-TIM3 antibody comprises

- i) comprises a VH sequence of SEQ ID NO:346 and a VL sequence of SEQ ID NO:347;
- ii) or humanized variant of the VH and VL of the antibody under i).

In one embodiment, the anti-TIM3 antibody comprises at least one, two, three, four, five, or six HVRs selected from (a) HVR-H1 comprising the amino acid sequence of SEQ ID NO:348; (b) HVR-H2 comprising the amino acid sequence of SEQ ID NO:349; (c) HVR-H3 comprising the amino acid sequence of SEQ ID NO:350; (d) HVR-L1 comprising the amino acid sequence of SEQ ID NO:351; (e) HVR-L2 comprising the amino acid sequence of SEQ ID NO:352; and (f) HVR-L3 comprising the amino acid sequence of SEQ ID NO:353.

In one aspect, the invention provides an anti-TIM3 antibody comprising (a) HVR-H1 comprising the amino acid sequence of SEQ ID NO:348; (b) HVR-H2 comprising the amino acid sequence of SEQ ID NO:349; (c) HVR-H3 comprising the amino acid sequence of SEQ ID NO:350; (d) HVR-L1 comprising the amino acid sequence of SEQ

ID NO:351; (e) HVR-L2 comprising the amino acid sequence of SEQ ID NO:352; and (f) HVR-L3 comprising the amino acid sequence of SEQ ID NO:353.

In one embodiment, the anti-TIM3 antibody comprises (a) a VH domain comprising at least one, at least two, or all three VH HVR sequences selected from (i) HVR-H1
5 comprising the amino acid sequence of SEQ ID NO:348, (ii) HVR-H2 comprising the amino acid sequence of SEQ ID NO:349, and (iii) HVR-H3 comprising an amino acid sequence selected from SEQ ID NO:350; and (b) a VL domain comprising at least one, at least two, or all three VL HVR sequences selected from (i) HVR-L1 comprising the amino acid sequence of SEQ ID NO:351; (ii) HVR-L2 comprising the amino acid
10 sequence of SEQ ID NO:352 and (c) HVR-L3 comprising the amino acid sequence of SEQ ID NO:353.

In one embodiment, the anti-TIM3 antibody comprises (a) a VH domain comprising (i) HVR-H1 comprising the amino acid sequence of SEQ ID NO:348, (ii) HVR-H2 comprising the amino acid sequence of SEQ ID NO:349, and (iii) HVR-H3 comprising an
15 amino acid sequence selected from SEQ ID NO:350; and (b) a VL domain comprising (i) HVR-L1 comprising the amino acid sequence of SEQ ID NO:351; (ii) HVR-L2 comprising the amino acid sequence of SEQ ID NO:352 and (iii) HVR-L3 comprising the amino acid sequence of SEQ ID NO:353.

In one embodiment such anti-TIM3 antibody comprises
20 i) comprises a VH sequence of SEQ ID NO:354 and a VL sequence of SEQ ID NO:355;
ii) or humanized variant of the VH and VL of the antibody under i).

In one embodiment, the anti-TIM3 antibody comprises at least one, two, three, four, five, or six HVRs selected from (a) HVR-H1 comprising the amino acid sequence of SEQ ID
25 NO:356; (b) HVR-H2 comprising the amino acid sequence of SEQ ID NO:357; (c) HVR-H3 comprising the amino acid sequence of SEQ ID NO:358; (d) HVR-L1 comprising the amino acid sequence of SEQ ID NO:359; (e) HVR-L2 comprising the amino acid sequence of SEQ ID NO:360; and (f) HVR-L3 comprising the amino acid sequence of SEQ ID NO:361.

30 In one embodiment, the anti-TIM3 antibody comprises (a) HVR-H1 comprising the amino acid sequence of SEQ ID NO:356; (b) HVR-H2 comprising the amino acid sequence of SEQ ID NO:357; (c) HVR-H3 comprising the amino acid sequence of SEQ ID NO:358; (d) HVR-L1 comprising the amino acid sequence of SEQ ID NO:359; (e)

HVR-L2 comprising the amino acid sequence of SEQ ID NO:360; and (f) HVR-L3 comprising the amino acid sequence of SEQ ID NO:361.

In one embodiment, the anti-TIM3 antibody comprises (a) a VH domain comprising at least one, at least two, or all three VH HVR sequences selected from (i) HVR-H1 comprising the amino acid sequence of SEQ ID NO:356, (ii) HVR-H2 comprising the amino acid sequence of SEQ ID NO: 357, and (iii) HVR-H3 comprising an amino acid sequence selected from SEQ ID NO: 358; and (b) a VL domain comprising at least one, at least two, or all three VL HVR sequences selected from (i) HVR-L1 comprising the amino acid sequence of SEQ ID NO: 359; (ii) HVR-L2 comprising the amino acid sequence of SEQ ID NO:360 and (c) HVR-L3 comprising the amino acid sequence of SEQ ID NO:361.

In one embodiment, the anti-TIM3 antibody comprises (a) a VH domain comprising (i) HVR-H1 comprising the amino acid sequence of SEQ ID NO: 356, (ii) HVR-H2 comprising the amino acid sequence of SEQ ID NO: 357, and (iii) HVR-H3 comprising an amino acid sequence selected from SEQ ID NO: 358; and (b) a VL domain comprising (i) HVR-L1 comprising the amino acid sequence of SEQ ID NO: 359; (ii) HVR-L2 comprising the amino acid sequence of SEQ ID NO:360 and (iii) HVR-L3 comprising the amino acid sequence of SEQ ID NO:361.

In one embodiment such anti-TIM3 antibody comprises

- i) comprises a VH sequence of SEQ ID NO:362 and a VL sequence of SEQ ID NO:363;
- ii) or humanized variant of the VH and VL of the antibody under i).

In one embodiment, the anti-TIM3 antibody comprises at least one, two, three, four, five, or six HVRs selected from (a) HVR-H1 comprising the amino acid sequence of SEQ ID NO:364; (b) HVR-H2 comprising the amino acid sequence of SEQ ID NO:365; (c) HVR-H3 comprising the amino acid sequence of SEQ ID NO:366; (d) HVR-L1 comprising the amino acid sequence of SEQ ID NO:367; (e) HVR-L2 comprising the amino acid sequence of SEQ ID NO:368; and (f) HVR-L3 comprising the amino acid sequence of SEQ ID NO:369.

In one embodiment, the anti-TIM3 antibody comprises (a) HVR-H1 comprising the amino acid sequence of SEQ ID NO:364; (b) HVR-H2 comprising the amino acid sequence of SEQ ID NO:365; (c) HVR-H3 comprising the amino acid sequence of SEQ ID NO:366; (d) HVR-L1 comprising the amino acid sequence of SEQ ID NO:367; (e)

HVR-L2 comprising the amino acid sequence of SEQ ID NO:368; and (f) HVR-L3 comprising the amino acid sequence of SEQ ID NO:369.

In one embodiment, the anti-TIM3 antibody comprises (a) a VH domain comprising at least one, at least two, or all three VH HVR sequences selected from (i) HVR-H1 comprising the amino acid sequence of SEQ ID NO:364, (ii) HVR-H2 comprising the amino acid sequence of SEQ ID NO:365, and (iii) HVR-H3 comprising an amino acid sequence selected from SEQ ID NO:366; and (b) a VL domain comprising at least one, at least two, or all three VL HVR sequences selected from (i) HVR-L1 comprising the amino acid sequence of SEQ ID NO:367; (ii) HVR-L2 comprising the amino acid sequence of SEQ ID NO:368 and (c) HVR-L3 comprising the amino acid sequence of SEQ ID NO:369.

In one embodiment, the anti-TIM3 antibody comprises (a) a VH domain comprising (i) HVR-H1 comprising the amino acid sequence of SEQ ID NO:364, (ii) HVR-H2 comprising the amino acid sequence of SEQ ID NO:365, and (iii) HVR-H3 comprising an amino acid sequence selected from SEQ ID NO:366; and (b) a VL domain comprising (i) HVR-L1 comprising the amino acid sequence of SEQ ID NO:367; (ii) HVR-L2 comprising the amino acid sequence of SEQ ID NO:368 and (iii) HVR-L3 comprising the amino acid sequence of SEQ ID NO:369.

In one embodiment such anti-TIM3 antibody comprises

- i) comprises a VH sequence of SEQ ID NO:370 and a VL sequence of SEQ ID NO:371;
- ii) or humanized variant of the VH and VL of the antibody under i).

In any of the above embodiments, an anti-TIM3 antibody is humanized. In one embodiment, an anti-TIM3 antibody comprises HVRs as in any of the above embodiments, and further comprises an acceptor human framework, e.g. a human immunoglobulin framework or a human consensus framework. In another embodiment, an anti-TIM3 antibody comprises HVRs as in any of the above embodiments, and further comprises a VH and VL comprising such HVRs. In a further aspect, the anti-TIM3 antibody binds to the same epitope as an anti-TIM3 antibody provided herein. For example, in certain embodiments, anti-TIM3 antibody binds to the same epitope as anti-TIM3 antibody comprising a VH sequence of SEQ ID NO:310 and a VL sequence of SEQ ID NO:311, or anti-TIM3 antibody binds to the same epitope as anti-TIM3 antibody comprising a VH sequence of SEQ ID NO:312 and a VL sequence of SEQ ID NO:313, or

an antibody is provided that binds to the same epitope as anti-TIM3 antibody comprising a VH sequence of SEQ ID NO:322 and a VL sequence of SEQ ID NO:323, or an antibody is provided that binds to the same epitope as anti-TIM3 antibody comprising a VH sequence of SEQ ID NO:330 and a VL sequence of SEQ ID NO:331, or an antibody is provided that binds to the same epitope as anti-TIM3 antibody comprising a VH sequence of SEQ ID NO:338 and a VL sequence of SEQ ID NO:339, or an antibody is provided that binds to the same epitope as anti-TIM3 antibody comprising a VH sequence of SEQ ID NO:346 and a VL sequence of SEQ ID NO:347, or an antibody is provided that binds to the same epitope as anti-TIM3 antibody comprising a VH sequence of SEQ ID NO:354 and a VL sequence of SEQ ID NO:355, or an antibody is provided that binds to the same epitope as anti-TIM3 antibody comprising a VH sequence of SEQ ID NO:362 and a VL sequence of SEQ ID NO:363, or an antibody is provided that binds to the same epitope as anti-TIM3 antibody comprising a VH sequence of SEQ ID NO:370 and a VL sequence of SEQ ID NO:371. In one preferred embodiment an antibody is provided that binds to the same epitope as an anti-TIM3 antibody comprising a VH sequence of SEQ ID NO:310 and a VL sequence of SEQ ID NO:311.

In one embodiment, the anti-TIM3 competes for binding to human TIM3 with an anti-TIM3 antibody comprising a VH sequence of SEQ ID NO:310 and a VL sequence of SEQ ID NO:311 as determined in a competition assay using TIM3 expressing RPMI-8226 cells (ATCC ® CCL-155™).

In one embodiment, the anti-TIM3 antibody according to any of the above embodiments is a monoclonal antibody, including a chimeric, humanized or human antibody. In one embodiment, an anti-TIM3 antibody is an antibody fragment, e.g., a Fv, Fab, Fab', scFv, diabody, or F(ab')₂ fragment. In another embodiment, the antibody is a full length antibody, e.g., an intact IgG1 or IgG4 antibody or other antibody class or isotype as defined herein.

In a further aspect, an anti-TIM3 antibody according to any of the above embodiments may incorporate any of the features, singly or in combination, as described herein.

In one embodiment, the anti-TIM3 antibody is any of the antibodies described in WO 2011/155607, WO 2013/006490, WO 03/063792, WO 2009/097394, or WO 2011/159877. In one embodiment, the anti-TIM3 antibody is F38-2E2. In some embodiments, the anti-TIM-3 antibodies are antibodies from hybridomas 8B.2C12 and

25F.1D6 and prepared as disclosed in U. S. Patent application Nos: 2004/0005322 and 2005/0191721, Sabatos, C. A. et al., Nature Immunol. 4:1102-1110, 2003, and Sanchez-Fueyo, A. et al., Nature Immunol. 4:1093-101 2003, all of which are hereby incorporated by reference as if set forth in their entirety. Other antibodies to TIM-3 are specifically contemplated and can be produced, e.g., with the methods disclosed herein. The nucleotide and protein sequences of TIM3 human sequences can be found at Genbank accession number AF251707.1 and Uniprot accession number Q8TDQ0. An exemplary human TIM3 amino acid sequence is set forth at SEQ ID NO:380; an exemplary human TIM3 extracellular domain amino acid sequence is set forth at SEQ ID NO:381.

10 ***Antibody Preparation***

As described above, in some embodiments, the PD-1 binding antagonist is an antibody (e.g., an anti-PD-1 antibody, an anti-PDL1 antibody, or an anti-PDL2 antibody). In some embodiments, the TIM3 antagonist is an antibody (e.g., an anti-TIM3 antagonist antibody). The antibodies described herein may be prepared using techniques available in the art for generating antibodies, exemplary methods of which are described in more detail in the following sections.

The antibody is directed against an antigen of interest. For example, the antibody may be directed against PD-1 (such as human PD-1), PDL1 (such as human PDL1), PDL2 (such as human PDL2), an TIM3 (such as human TIM3). Preferably, the antigen is a biologically important polypeptide and administration of the antibody to a mammal suffering from a disorder can result in a therapeutic benefit in that mammal.

In certain embodiments, an antibody described herein has a dissociation constant (Kd) of 1 μ M, 150 nM, 100 nM, 50 nM, 10 nM, 1 nM, 0.1 nM, 0.01 nM, or 0.001 nM (e.g. 10⁻⁸ M or less, e.g. from 10⁻⁸ M to 10⁻¹³ M, e.g., from 10⁻⁹ M to 10⁻¹³ M).

In one embodiment, Kd is measured by a radiolabeled antigen binding assay (RIA) performed with the Fab version of an antibody of interest and its antigen as described by the following assay. Solution binding affinity of Fabs for antigen is measured by equilibrating Fab with a minimal concentration of (125I)-labeled antigen in the presence of a titration series of unlabeled antigen, then capturing bound antigen with an anti-Fab antibody-coated plate (see, e.g., Chen et al., *J. Mol. Biol.* 293:865-881(1999)). To establish conditions for the assay, MICROTITER® multi-well plates (Thermo Scientific) are coated overnight with 5 μ g/ml of a capturing anti-Fab antibody (Cappel Labs) in 50 mM sodium carbonate (pH 9.6), and subsequently blocked with 2% (w/v) bovine serum

albumin in PBS for two to five hours at room temperature (approximately 23°C). In a non-adsorbent plate (Nunc #269620), 100 pM or 26 pM [¹²⁵I]-antigen are mixed with serial dilutions of a Fab of interest. The Fab of interest is then incubated overnight; however, the incubation may continue for a longer period (*e.g.*, about 65 hours) to ensure that equilibrium is reached. Thereafter, the mixtures are transferred to the capture plate for incubation at room temperature (*e.g.*, for one hour). The solution is then removed and the plate washed eight times with 0.1% polysorbate 20 (TWEEN-20[®]) in PBS. When the plates have dried, 150 µl/well of scintillant (MICROSCINT-20 TM; Packard) is added, and the plates are counted on a TOPCOUNT TM gamma counter (Packard) for ten minutes. Concentrations of each Fab that give less than or equal to 20% of maximal binding are chosen for use in competitive binding assays.

According to another embodiment, K_d is measured using surface plasmon resonance assays using a BIACORE[®]-2000 or a BIACORE[®]-3000 (BIAcore, Inc., Piscataway, NJ) at 25°C with immobilized antigen CM5 chips at ~10 response units (RU). Briefly, carboxymethylated dextran biosensor chips (CM5, BIACORE, Inc.) are activated with *N*-ethyl-*N*'-(3-dimethylaminopropyl)-carbodiimide hydrochloride (EDC) and *N*-hydroxysuccinimide (NHS) according to the supplier's instructions. Antigen is diluted with 10 mM sodium acetate, pH 4.8, to 5 µg/ml (~0.2 µM) before injection at a flow rate of 5 µl/minute to achieve approximately 10 response units (RU) of coupled protein. Following the injection of antigen, 1 M ethanolamine is injected to block unreacted groups. For kinetics measurements, two-fold serial dilutions of Fab (0.78 nM to 500 nM) are injected in PBS with 0.05% polysorbate 20 (TWEEN-20TM) surfactant (PBST) at 25°C at a flow rate of approximately 25 µl/min. Association rates (k_{on}) and dissociation rates (k_{off}) are calculated using a simple one-to-one Langmuir binding model (BIACORE[®] Evaluation Software version 3.2) by simultaneously fitting the association and dissociation sensorgrams. The equilibrium dissociation constant (K_d) is calculated as the ratio k_{off}/k_{on}. See, *e.g.*, Chen et al., *J. Mol. Biol.* 293:865-881 (1999). If the on-rate exceeds 10⁶ M⁻¹ s⁻¹ by the surface plasmon resonance assay above, then the on-rate can be determined by using a fluorescent quenching technique that measures the increase or decrease in fluorescence emission intensity (excitation = 295 nm; emission = 340 nm, 16 nm band-pass) at 25°C of a 20 nM anti-antigen antibody (Fab form) in PBS, pH 7.2, in the presence of increasing concentrations of antigen as measured in a spectrometer, such

as a stop-flow equipped spectrophotometer (Aviv Instruments) or a 8000-series SLM-AMINCO TM spectrophotometer (ThermoSpectronic) with a stirred cuvette.

In some embodiments, an anti-TIM3 antibody as described herein exhibits a binding affinity of at least 100 pM or less against human TIM3, a binding affinity of at least 300 pM or less against human TIM3, a binding affinity of at least 400 pM or less against human TIM3, a neutralizing ability of at least 40 nM or less against the human TIM3, a neutralizing ability of at least 120 nM or less against the human TIM3, and a neutralizing ability of at least 31 nM or less against the human TIM3. In these embodiments, binding affinity may be measured by surface plasmon resonance as described in U.S. Patent No. 8,771,697,

Antibody Fragments

In certain embodiments, an antibody described herein is an antibody fragment. Antibody fragments include, but are not limited to, Fab, Fab', Fab'-SH, F(ab')₂, Fv, and scFv fragments, and other fragments described below. For a review of certain antibody fragments, see Hudson et al. *Nat. Med.* 9:129-134 (2003). For a review of scFv fragments, see, e.g., Pluckthün, in *The Pharmacology of Monoclonal Antibodies*, vol. 113, Rosenberg and Moore eds., (Springer-Verlag, New York), pp. 269-315 (1994); see also WO 93/16185; and U.S. Patent Nos. 5,571,894 and 5,587,458. For discussion of Fab and F(ab')₂ fragments comprising salvage receptor binding epitope residues and having increased in vivo half-life, see U.S. Patent No. 5,869,046.

Diabodies are antibody fragments with two antigen-binding sites that may be bivalent or bispecific. See, for example, EP 404,097; WO 1993/01161; Hudson et al., *Nat. Med.* 9:129-134 (2003); and Hollinger et al., *Proc. Natl. Acad. Sci. USA* 90: 6444-6448 (1993).

Triabodies and tetrabodies are also described in Hudson et al., *Nat. Med.* 9:129-134 (2003). Single-domain antibodies are antibody fragments comprising all or a portion of the heavy chain variable domain or all or a portion of the light chain variable domain of an antibody. In certain embodiments, a single-domain antibody is a human single-domain antibody (Domantis, Inc., Waltham, MA; see, e.g., U.S. Patent No. 6,248,516 B1).

Antibody fragments can be made by various techniques, including but not limited to proteolytic digestion of an intact antibody as well as production by recombinant host cells (e.g. *E. coli* or phage), as described herein.

Chimeric and Humanized Antibodies

In certain embodiments, an antibody described herein is a chimeric antibody. Certain chimeric antibodies are described, e.g., in U.S. Patent No. 4,816,567; and Morrison et al., *Proc. Natl. Acad. Sci. USA*, 81:6851-6855 (1984)). In one example, a chimeric antibody

5 comprises a non-human variable region (e.g., a variable region derived from a mouse, rat, hamster, rabbit, or non-human primate, such as a monkey) and a human constant region. In a further example, a chimeric antibody is a “class switched” antibody in which the class or subclass has been changed from that of the parent antibody. Chimeric antibodies include antigen-binding fragments thereof. In certain embodiments, a chimeric antibody

10 is a humanized antibody. Typically, a non-human antibody is humanized to reduce immunogenicity to humans, while retaining the specificity and affinity of the parental non-human antibody. Generally, a humanized antibody comprises one or more variable domains in which HVRs, e.g., CDRs, (or portions thereof) are derived from a non-human antibody, and FRs (or portions thereof) are derived from human antibody sequences. A

15 humanized antibody optionally will also comprise at least a portion of a human constant region. In some embodiments, some FR residues in a humanized antibody are substituted with corresponding residues from a non-human antibody (e.g., the antibody from which the HVR residues are derived), e.g., to restore or improve antibody specificity or affinity. Humanized antibodies and methods of making them are reviewed, e.g., in Almagro and

20 Fransson, *Front. Biosci.* 13:1619-1633 (2008), and are further described, e.g., in Riechmann et al., *Nature* 332:323-329 (1988); Queen et al., *Proc. Nat'l Acad. Sci. USA* 86:10029-10033 (1989); US Patent Nos. 5, 821,337, 7,527,791, 6,982,321, and 7,087,409; Kashmiri et al., *Methods* 36:25-34 (2005) (describing SDR (a-CDR) grafting); Padlan, *Mol. Immunol.* 28:489-498 (1991) (describing “resurfacing”); Dall’Acqua et al., *Methods*

25 36:43-60 (2005) (describing “FR shuffling”); and Osbourn et al., *Methods* 36:61-68 (2005) and Klimka et al., *Br. J. Cancer*, 83:252-260 (2000) (describing the “guided selection” approach to FR shuffling).

Human framework regions that may be used for humanization include but are not limited to: framework regions selected using the “best-fit” method (see, e.g., Sims et al. *J.*

30 *Immunol.* 151:2296 (1993)); framework regions derived from the consensus sequence of human antibodies of a particular subgroup of light or heavy chain variable regions (see, e.g., Carter et al. *Proc. Natl. Acad. Sci. USA*, 89:4285 (1992); and Presta et al. *J. Immunol.*, 151:2623 (1993)); human mature (somatically mutated) framework regions or

human germline framework regions (see, e.g., Almagro and Fransson, *Front. Biosci.* 13:1619-1633 (2008)); and framework regions derived from screening FR libraries (see, e.g., Baca et al., *J. Biol. Chem.* 272:10678-10684 (1997) and Rosok et al., *J. Biol. Chem.* 271:22611-22618 (1996)).

5

Human Antibodies

In certain embodiments, an antibody described herein is a human antibody. Human antibodies can be produced using various techniques known in the art. Human antibodies are described generally in van Dijk and van de Winkel, *Curr. Opin. Pharmacol.* 5: 368-74
10 (2001) and Lonberg, *Curr. Opin. Immunol.* 20:450-459 (2008).

Human antibodies may be prepared by administering an immunogen to a transgenic animal that has been modified to produce intact human antibodies or intact antibodies with human variable regions in response to antigenic challenge. Such animals typically contain all or a portion of the human immunoglobulin loci, which replace the endogenous
15 immunoglobulin loci, or which are present extrachromosomally or integrated randomly into the animal's chromosomes. In such transgenic mice, the endogenous immunoglobulin loci have generally been inactivated. For review of methods for obtaining human antibodies from transgenic animals, see Lonberg, *Nat. Biotech.* 23:1117-1125 (2005). See also, e.g., U.S. Patent Nos. 6,075,181 and 6,150,584 describing
20 XENOMOUSE™ technology; U.S. Patent No. 5,770,429 describing HUMAB® technology; U.S. Patent No. 7,041,870 describing K-M MOUSE® technology, and U.S. Patent Application Publication No. US 2007/0061900, describing VELOCIMOUSE® technology). Human variable regions from intact antibodies generated by such animals may be further modified, e.g., by combining with a different human constant region.

25 Human antibodies can also be made by hybridoma-based methods. Human myeloma and mouse-human heteromyeloma cell lines for the production of human monoclonal antibodies have been described. (See, e.g., Kozbor *J. Immunol.*, 133: 3001 (1984); Brodeur et al., *Monoclonal Antibody Production Techniques and Applications*, pp. 51-63 (Marcel Dekker, Inc., New York, 1987); and Boerner et al., *J. Immunol.*, 147: 86 (1991).)
30 Human antibodies generated via human B-cell hybridoma technology are also described in Li et al., *Proc. Natl. Acad. Sci. USA*, 103:3557-3562 (2006). Additional methods include those described, for example, in U.S. Patent No. 7,189,826 (describing production of monoclonal human IgM antibodies from hybridoma cell lines) and Ni, *Xiandai*

Mianyxue, 26(4):265-268 (2006) (describing humanhuman hybridomas). Human hybridoma technology (Trioma technology) is also described in Vollmers and Brandlein, *Histology and Histopathology*, 20(3):927-937 (2005) and Vollmers and Brandlein, *Methods and Findings in Experimental and Clinical Pharmacology*, 27(3):185-91 (2005).

- 5 Human antibodies may also be generated by isolating Fv clone variable domain sequences selected from human-derived phage display libraries. Such variable domain sequences may then be combined with a desired human constant domain. Techniques for selecting human antibodies from antibody libraries are described below.

10 ***Library-Derived Antibodies***

- Antibodies may be isolated by screening combinatorial libraries for antibodies with the desired activity or activities. For example, a variety of methods are known in the art for generating phage display libraries and screening such libraries for antibodies possessing the desired binding characteristics. Such methods are reviewed, e.g., in Hoogenboom et al. in *Methods in Molecular Biology* 178:1-37 (O'Brien et al., ed., Human Press, Totowa, NJ, 2001) and further described, e.g., in the McCafferty et al., *Nature* 348:552-554; Clackson et al., *Nature* 352: 624-628 (1991); Marks et al., *J. Mol. Biol.* 222: 581-597 (1992); Marks and Bradbury, in *Methods in Molecular Biology* 248:161-175 (Lo, ed., Human Press, Totowa, NJ, 2003); Sidhu et al., *J. Mol. Biol.* 338(2): 299-310 (2004); Lee et al., *J. Mol. Biol.* 340(5): 1073-1093 (2004); Fellouse, *Proc. Natl. Acad. Sci. USA* 101(34): 12467-12472 (2004); and Lee et al., *J. Immunol. Methods* 284(1-2): 119-132(2004).

- In certain phage display methods, repertoires of VH and VL genes are separately cloned by polymerase chain reaction (PCR) and recombined randomly in phage libraries, which can then be screened for antigen-binding phage as described in Winter et al., *Ann. Rev. Immunol.*, 12: 433-455 (1994). Phage typically display antibody fragments, either as singlechain Fv (scFv) fragments or as Fab fragments. Libraries from immunized sources provide highaffinity antibodies to the immunogen without the requirement of constructing hybridomas. Alternatively, the naive repertoire can be cloned (e.g., from human) to provide a single source of antibodies to a wide range of non-self and also self antigens without any immunization as described by Griffiths et al., *EMBO J*, 12: 725-734 (1993). Finally, naive libraries can also be made synthetically by cloning unrearranged V-gene segments from stem cells, and using PCR primers containing random sequence to encode the highly variable CDR3 regions and to accomplish rearrangement *in vitro*, as described

by Hoogenboom and Winter, *J. Mol. Biol.*, 227:381-388 (1992). Patent publications describing human antibody phage libraries include, for example: US Patent No. 5,750,373, and US Patent Publication Nos. 2005/0079574, 2005/0119455, 2005/0266000, 2007/0117126, 2007/0160598, 2007/0237764, 2007/0292936, and 2009/0002360.

- 5 Antibodies or antibody fragments isolated from human antibody libraries are considered human antibodies or human antibody fragments herein.

Multispecific Antibodies

- In certain embodiments, an antibody described herein is a multispecific antibody, e.g. a
10 bispecific antibody. Multispecific antibodies are monoclonal antibodies that have binding specificities for at least two different sites. Examples of T cell activating bispecific antigen binding molecules specific for FolR1 and CD3 are described herein. In some embodiments, the PD1 axis component antagonist is multispecific. In one of the binding specificities is for a PD-1 axis component (e.g., PD-1, PDL1, or PDL2) and the other is
15 for any other antigen. In some embodiments, one of the binding specificities is for IL-17 or IL-17R and the other is for any other antigen. In certain embodiments, bispecific antibodies may bind to two different epitopes of a PD-1 axis component (e.g., PD-1, PDL1, or PDL2), IL-17, or IL-17R. Bispecific antibodies can be prepared as full length antibodies or antibody fragments.
- 20 In some embodiments, one of the binding specificities is for a PD-1 axis component (e.g., PD-1, PDL1, or PDL2) and the other is for IL-17 or IL-17R. Provided herein are methods for treating or delaying progression of cancer in an individual comprising administering to the individual an effective amount of a multispecific antibody, wherein the multispecific antibody comprises a first binding specificity for a PD-1 axis component
25 (e.g., PD-1, PDL1, or PDL2) and a second binding specificity for IL-17 or IL-17R. In some embodiments, a multispecific antibody may be made by any of the techniques described herein and below.

- Techniques for making multispecific antibodies include, but are not limited to, recombinant co-expression of two immunoglobulin heavy chain-light chain pairs having
30 different specificities (see Milstein and Cuello, *Nature* 305: 537 (1983)), WO 93/08829, and Traunecker et al., *EMBO J.* 10: 3655 (1991)), and “knob-in-hole” engineering (see, e.g., U.S. Patent No. 5,731,168). Multi-specific antibodies may also be made by engineering electrostatic steering effects for making antibody Fc-heterodimeric molecules

(WO 2009/089004A1); crosslinking two or more antibodies or fragments (see, e.g., US Patent No. 4,676,980, and Brennan et al., *Science*, 229: 81 (1985)); using leucine zippers to produce bi-specific antibodies (see, e.g., Kostelny et al., *J. Immunol.*, 148(5):1547-1553 (1992)); using “diabody” technology for making bispecific antibody fragments (see, e.g., Hollinger et al., *Proc. Natl. Acad. Sci. USA*, 90:6444-6448 (1993)); and using single-chain Fv (sFv) dimers (see, e.g. Gruber et al., *J. Immunol.*, 152:5368 (1994)); and preparing trispecific antibodies as described, e.g., in Tutt et al. *J. Immunol.* 147: 60 (1991). Engineered antibodies with three or more functional antigen binding sites, including “Octopus antibodies,” are also included herein (see, e.g. US 2006/0025576A1).

- 10 The antibody or fragment herein also includes a “Dual Acting FAb” or “DAF” comprising an antigen binding site that binds to a PD-1 axis component (e.g., PD-1, PDL1, or PDL2), IL-17, or IL-17R as well as another, different antigen (see, US 2008/0069820, for example).

15 **C. Nucleic Acid sequences, vectors and methods of production**

- Polynucleotides encoding a T cell activating bispecific antigen binding molecule, e.g., a T cell activating bispecific antigen binding molecule comprising a first antigen binding site specific for Folate Receptor 1 (FolR1) and a second antigen binding site specific for CD3, and antibodies may be used for production of the T cell activating
- 20 bispecific antigen binding molecule and antibodies described herein. The T cell activating bispecific antigen binding molecule and antibodies of the invention may be expressed as a single polynucleotide that encodes the entire bispecific antigen binding molecule or as multiple (e.g., two or more) polynucleotides that are co-expressed. Polypeptides encoded by polynucleotides that are co-expressed may associate through,
- 25 e.g., disulfide bonds or other means to form a functional T cell activating bispecific antigen binding molecule and antibody. For example, the light chain portion of a Fab fragment may be encoded by a separate polynucleotide from the portion of the bispecific antibody or the antibody binding to FolR1 comprising the heavy chain portion of the Fab fragment, an Fc domain subunit and optionally (part of) another Fab fragment. When co-
- 30 expressed, the heavy chain polypeptides will associate with the light chain polypeptides to form the Fab fragment. In another example, the portion of the T cell activating bispecific antigen binding molecule or the FolR1 antigen binding portion provided therein comprising one of the two Fc domain subunits and optionally (part of) one or more Fab

fragments could be encoded by a separate polynucleotide from the portion of the bispecific antibody or the antibody binding to FolR1 provided therein comprising the other of the two Fc domain subunits and optionally (part of) a Fab fragment. When co-expressed, the Fc domain subunits will associate to form the Fc domain.

5 In certain embodiments the polynucleotide or nucleic acid is DNA. In other embodiments, a polynucleotide of the present invention is RNA, for example, in the form of messenger RNA (mRNA). RNA of the present invention may be single stranded or double stranded.

10 **D. Antibody Variants**

In certain embodiments, amino acid sequence variants of the T cell activating bispecific antigen binding molecule specific for FolR1 and CD3 provided herein and antibodies are contemplated, in addition to those described above. For example, it may be desirable to improve the binding affinity and/or other biological properties of the T cell

15 activating bispecific antigen binding molecule. Amino acid sequence variants of a T cell activating bispecific antigen binding molecule and antibody may be prepared by introducing appropriate modifications into the nucleotide sequence encoding the T cell activating bispecific antigen binding molecule or antibody, or by peptide synthesis. Such modifications include, for example, deletions from, and/or insertions into and/or

20 substitutions of residues within the amino acid sequences of the antibody. Any combination of deletion, insertion, and substitution can be made to arrive at the final construct, provided that the final construct possesses the desired characteristics, e.g., antigen-binding.

25 **1. Substitution, Insertion, and Deletion Variants**

In certain embodiments, variants having one or more amino acid substitutions are provided. Sites of interest for substitutional mutagenesis include the HVRs and FRs. Conservative substitutions are shown in Table B under the heading of "conservative substitutions." More substantial changes are provided in Table B under the heading of

30 "exemplary substitutions," and as further described below in reference to amino acid side chain classes. Amino acid substitutions may be introduced into an antibody of interest and the products screened for a desired activity, e.g., retained/improved antigen binding, decreased immunogenicity, or improved ADCC or CDC.

TABLE B

Original Residue	Exemplary Substitutions	Preferred Substitutions
Ala (A)	Val; Leu; Ile	Val
Arg (R)	Lys; Gln; Asn	Lys
Asn (N)	Gln; His; Asp, Lys; Arg	Gln
Asp (D)	Glu; Asn	Glu
Cys (C)	Ser; Ala	Ser
Gln (Q)	Asn; Glu	Asn
Glu (E)	Asp; Gln	Asp
Gly (G)	Ala	Ala
His (H)	Asn; Gln; Lys; Arg	Arg
Ile (I)	Leu; Val; Met; Ala; Phe; Norleucine	Leu
Leu (L)	Norleucine; Ile; Val; Met; Ala; Phe	Ile
Lys (K)	Arg; Gln; Asn	Arg
Met (M)	Leu; Phe; Ile	Leu
Phe (F)	Trp; Leu; Val; Ile; Ala; Tyr	Tyr
Pro (P)	Ala	Ala
Ser (S)	Thr	Thr
Thr (T)	Val; Ser	Ser
Trp (W)	Tyr; Phe	Tyr
Tyr (Y)	Trp; Phe; Thr; Ser	Phe
Val (V)	Ile; Leu; Met; Phe; Ala; Norleucine	Leu

5 Amino acids may be grouped according to common side-chain properties:

(1) hydrophobic: Norleucine, Met, Ala, Val, Leu, Ile;

(2) neutral hydrophilic: Cys, Ser, Thr, Asn, Gln;

(3) acidic: Asp, Glu;

(4) basic: His, Lys, Arg;

10 (5) residues that influence chain orientation: Gly, Pro;

(6) aromatic: Trp, Tyr, Phe.

Non-conservative substitutions will entail exchanging a member of one of these classes for another class.

One type of substitutional variant involves substituting one or more hypervariable region residues of a parent antibody (e.g. a humanized or human antibody). Generally, the resulting variant(s) selected for further study will have modifications (e.g., improvements) in certain biological properties (e.g., increased affinity, reduced immunogenicity) relative to the parent antibody and/or will have substantially retained certain biological properties of the parent antibody. An exemplary substitutional variant is an affinity matured antibody, which may be conveniently generated, e.g., using phage display-based affinity maturation techniques such as those described herein. Briefly, one or more HVR residues are mutated and the variant antibodies displayed on phage and screened for a particular biological activity (e.g. binding affinity).

Alterations (e.g., substitutions) may be made in HVRs, e.g., to improve antibody affinity. Such alterations may be made in HVR “hotspots,” i.e., residues encoded by codons that undergo mutation at high frequency during the somatic maturation process (see, e.g., Chowdhury, *Methods Mol. Biol.* 207:179-196 (2008)), and/or SDRs (a-CDRs), with the resulting variant VH or VL being tested for binding affinity. Affinity maturation by constructing and reselecting from secondary libraries has been described, e.g., in Hoogenboom et al. in *Methods in Molecular Biology* 178:1-37 (O’Brien et al., ed., Human Press, Totowa, NJ, (2001).) In some embodiments of affinity maturation, diversity is introduced into the variable genes chosen for maturation by any of a variety of methods (e.g., error-prone PCR, chain shuffling, or oligonucleotide-directed mutagenesis). A secondary library is then created. The library is then screened to identify any antibody variants with the desired affinity. Another method to introduce diversity involves HVR-directed approaches, in which several HVR residues (e.g., 4-6 residues at a time) are randomized. HVR residues involved in antigen binding may be specifically identified, e.g., using alanine scanning mutagenesis or modeling. CDR-H3 and CDR-L3 in particular are often targeted.

In certain embodiments, substitutions, insertions, or deletions may occur within one or more HVRs so long as such alterations do not substantially reduce the ability of the antibody to bind antigen. For example, conservative alterations (e.g., conservative substitutions as provided herein) that do not substantially reduce binding affinity may be made in HVRs. Such alterations may be outside of HVR “hotspots” or SDRs. In certain

embodiments of the variant VH and VL sequences provided above, each HVR either is unaltered, or contains no more than one, two or three amino acid substitutions.

A useful method for identification of residues or regions of an antibody that may be targeted for mutagenesis is called "alanine scanning mutagenesis" as described by
5 Cunningham and Wells (1989) *Science*, 244:1081-1085. In this method, a residue or group of target residues (e.g., charged residues such as arg, asp, his, lys, and glu) are identified and replaced by a neutral or negatively charged amino acid (e.g., alanine or polyalanine) to determine whether the interaction of the antibody with antigen is affected. Further substitutions may be introduced at the amino acid locations demonstrating
10 functional sensitivity to the initial substitutions. Alternatively, or additionally, a crystal structure of an antigen-antibody complex to identify contact points between the antibody and antigen. Such contact residues and neighboring residues may be targeted or eliminated as candidates for substitution. Variants may be screened to determine whether they contain the desired properties.

15 Amino acid sequence insertions include amino- and/or carboxyl-terminal fusions ranging in length from one residue to polypeptides containing a hundred or more residues, as well as intrasequence insertions of single or multiple amino acid residues. Examples of terminal insertions include an antibody with an N-terminal methionyl residue. Other insertional variants of the antibody molecule include the fusion to the N- or C-terminus of
20 the antibody to an enzyme (e.g. for ADEPT) or a polypeptide which increases the serum half-life of the antibody.

2. Glycosylation variants

In certain embodiments, a T cell activating bispecific antigen binding molecule or
25 an antibody provided herein is altered to increase or decrease the extent to which the antibody is glycosylated. Addition or deletion of glycosylation sites to an antibody may be conveniently accomplished by altering the amino acid sequence such that one or more glycosylation sites is created or removed.

Where the T cell activating bispecific antigen binding molecule or the antibody
30 used with the invention comprises an Fc region, the carbohydrate attached thereto may be altered. Native antibodies produced by mammalian cells typically comprise a branched, biantennary oligosaccharide that is generally attached by an N-linkage to Asn297 of the CH2 domain of the Fc region. See, e.g., Wright et al. *TIBTECH* 15:26-32 (1997). The

oligosaccharide may include various carbohydrates, e.g., mannose, N-acetyl glucosamine (GlcNAc), galactose, and sialic acid, as well as a fucose attached to a GlcNAc in the “stem” of the biantennary oligosaccharide structure. In some embodiments, modifications of the oligosaccharide in a bispecific antibody or an antibody binding to DR5 of the invention may be made in order to create antibody variants with certain improved properties.

In one embodiment, bispecific antibody variants or variants of antibodies are provided having a carbohydrate structure that lacks fucose attached (directly or indirectly) to an Fc region. For example, the amount of fucose in such antibody may be from 1% to 80%, from 1% to 65%, from 5% to 65% or from 20% to 40%. The amount of fucose is determined by calculating the average amount of fucose within the sugar chain at Asn297, relative to the sum of all glycostructures attached to Asn 297 (e. g. complex, hybrid and high mannose structures) as measured by MALDI-TOF mass spectrometry, as described in WO 2008/077546, for example. Asn297 refers to the asparagine residue located at about position 297 in the Fc region (Eu numbering of Fc region residues); however, Asn297 may also be located about ± 3 amino acids upstream or downstream of position 297, i.e., between positions 294 and 300, due to minor sequence variations in antibodies. Such fucosylation variants may have improved ADCC function. See, e.g., US Patent Publication Nos. US 2003/0157108 (Presta, L.); US 2004/0093621 (Kyowa Hakko Kogyo Co., Ltd). Examples of publications related to “defucosylated” or “fucose-deficient” antibody variants include: US 2003/0157108; WO 2000/61739; WO 2001/29246; US 2003/0115614; US 2002/0164328; US 2004/0093621; US 2004/0132140; US 2004/0110704; US 2004/0110282; US 2004/0109865; WO 2003/085119; WO 2003/084570; WO 2005/035586; WO 2005/035778; WO2005/053742; WO2002/031140; Okazaki et al. *J. Mol. Biol.* 336:1239-1249 (2004); Yamane-Ohnuki et al. *Biotech. Bioeng.* 87: 614 (2004). Examples of cell lines capable of producing defucosylated antibodies include Lec13 CHO cells deficient in protein fucosylation (Ripka et al. *Arch. Biochem. Biophys.* 249:533-545 (1986); US Pat Appl No US 2003/0157108 A1, Presta, L; and WO 2004/056312 A1, Adams *et al.*, especially at Example 11), and knockout cell lines, such as alpha-1,6-fucosyltransferase gene, *FUT8*, knockout CHO cells (see, e.g., Yamane-Ohnuki et al. *Biotech. Bioeng.* 87: 614 (2004); Kanda, Y. et al., *Biotechnol. Bioeng.*, 94(4):680-688 (2006); and WO2003/085107).

T cell activating bispecific antigen binding molecule variants and antibody variants are further provided with bisected oligosaccharides, e.g., in which a biantennary oligosaccharide attached to the Fc region of the T cell activating bispecific antigen binding molecule binding to FolR1 is bisected by GlcNAc. Such T cell activating
5 bispecific antigen binding molecule variants may have reduced fucosylation and/or improved ADCC function. Examples of such antibody variants are described, e.g., in WO 2003/011878 (Jean-Mairet et al.); US Patent No. 6,602,684 (Umana et al.); and US 2005/0123546 (Umana *et al.*). Antibody variants with at least one galactose residue in the oligosaccharide attached to the Fc region are also provided. Such antibody variants may
10 have improved CDC function. Such antibody variants are described, e.g., in WO 1997/30087 (Patel et al.); WO 1998/58964 (Raju, S.); and WO 1999/22764 (Raju, S.).

3. Cysteine engineered antibody variants

In certain embodiments, it may be desirable to create cysteine engineered T cell
15 activating bispecific antigen binding molecule and antibodies, e.g., THIOMABS, in which one or more residues of the T cell activating bispecific antigen binding molecule are substituted with cysteine residues. In particular embodiments, the substituted residues occur at accessible sites of the T cell activating bispecific antigen binding molecule. By substituting those residues with cysteine, reactive thiol groups are thereby positioned at
20 accessible sites of the antibody and may be used to conjugate the antibody to other moieties, such as drug moieties or linker-drug moieties, to create an immunoconjugate. In certain embodiments, any one or more of the following residues may be substituted with cysteine: V205 (Kabat numbering) of the light chain; A118 (EU numbering) of the heavy chain; and S400 (EU numbering) of the heavy chain Fc region. Cysteine engineered
25 antibodies may be generated as described, e.g., in U.S. Patent No. 7,521,541.

E. Recombinant Methods and Compositions

T cell activating bispecific antigen binding molecule and antibodies of the invention may be obtained, for example, by solid-state peptide synthesis (e.g. Merrifield
30 solid phase synthesis) or recombinant production. For recombinant production one or more polynucleotide encoding the T cell activating bispecific antigen binding molecule or antibodies (or fragments), e.g., as described above, is isolated and inserted into one or more vectors for further cloning and/or expression in a host cell. Such polynucleotide

may be readily isolated and sequenced using conventional procedures. In one embodiment a vector, preferably an expression vector, comprising one or more of the polynucleotides of the invention is provided. Methods which are well known to those skilled in the art can be used to construct expression vectors containing the coding

5 sequence of a T cell activating bispecific antigen binding molecule or an antibody along with appropriate transcriptional/translational control signals. These methods include *in vitro* recombinant DNA techniques, synthetic techniques and *in vivo* recombination/genetic recombination. See, for example, the techniques described in Maniatis et al., MOLECULAR CLONING: A LABORATORY MANUAL, Cold Spring Harbor

10 Laboratory, N.Y. (1989); and Ausubel et al., CURRENT PROTOCOLS IN MOLECULAR BIOLOGY, Greene Publishing Associates and Wiley Interscience, N.Y (1989). The expression vector can be part of a plasmid, virus, or may be a nucleic acid fragment. The expression vector includes an expression cassette into which the polynucleotide encoding T cell activating bispecific antigen binding molecule (fragment) or an antibody (fragment)

15 (i.e. the coding region) is cloned in operable association with a promoter and/or other transcription or translation control elements. As used herein, a "coding region" is a portion of nucleic acid which consists of codons translated into amino acids. Although a "stop codon" (TAG, TGA, or TAA) is not translated into an amino acid, it may be considered to be part of a coding region, if present, but any flanking sequences, for

20 example promoters, ribosome binding sites, transcriptional terminators, introns, 5' and 3' untranslated regions, and the like, are not part of a coding region. Two or more coding regions can be present in a single polynucleotide construct, e.g. on a single vector, or in separate polynucleotide constructs, e.g. on separate (different) vectors. Furthermore, any vector may contain a single coding region, or may comprise two or more coding regions,

25 e.g. a vector of the present invention may encode one or more polypeptides, which are post- or co-translationally separated into the final proteins via proteolytic cleavage. In addition, a vector, polynucleotide, or nucleic acid of the invention may encode heterologous coding regions, either fused or unfused to a polynucleotide encoding the T cell activating bispecific antigen binding molecule (fragment) or an antibody, or variant

30 or derivative thereof. Heterologous coding regions include without limitation specialized elements or motifs, such as a secretory signal peptide or a heterologous functional domain. An operable association is when a coding region for a gene product, e.g. a polypeptide, is associated with one or more regulatory sequences in such a way as to

place expression of the gene product under the influence or control of the regulatory sequence(s). Two DNA fragments (such as a polypeptide coding region and a promoter associated therewith) are "operably associated" if induction of promoter function results in the transcription of mRNA encoding the desired gene product and if the nature of the linkage between the two DNA fragments does not interfere with the ability of the expression regulatory sequences to direct the expression of the gene product or interfere with the ability of the DNA template to be transcribed. Thus, a promoter region would be operably associated with a nucleic acid encoding a polypeptide if the promoter was capable of effecting transcription of that nucleic acid. The promoter may be a cell-specific promoter that directs substantial transcription of the DNA only in predetermined cells.

Other transcription control elements, besides a promoter, for example enhancers, operators, repressors, and transcription termination signals, can be operably associated with the polynucleotide to direct cell-specific transcription. Suitable promoters and other transcription control regions are disclosed herein. A variety of transcription control regions are known to those skilled in the art. These include, without limitation, transcription control regions, which function in vertebrate cells, such as, but not limited to, promoter and enhancer segments from cytomegaloviruses (e.g. the immediate early promoter, in conjunction with intron-A), simian virus 40 (e.g. the early promoter), and retroviruses (such as, e.g. Rous sarcoma virus). Other transcription control regions include those derived from vertebrate genes such as actin, heat shock protein, bovine growth hormone and rabbit α -globin, as well as other sequences capable of controlling gene expression in eukaryotic cells. Additional suitable transcription control regions include tissue-specific promoters and enhancers as well as inducible promoters (e.g. promoters inducible tetracyclins). Similarly, a variety of translation control elements are known to those of ordinary skill in the art. These include, but are not limited to ribosome binding sites, translation initiation and termination codons, and elements derived from viral systems (particularly an internal ribosome entry site, or IRES, also referred to as a CITE sequence). The expression cassette may also include other features such as an origin of replication, and/or chromosome integration elements such as retroviral long terminal repeats (LTRs), or adeno-associated viral (AAV) inverted terminal repeats (ITRs).

Polynucleotide and nucleic acid coding regions of the present invention may be associated with additional coding regions which encode secretory or signal peptides,

which direct the secretion of a polypeptide encoded by a polynucleotide of the present invention. For example, if secretion of the T cell activating bispecific antigen binding molecule or the antibody is desired, DNA encoding a signal sequence may be placed upstream of the nucleic acid encoding a bispecific antibody of the invention or the

5 antibody binding to DR5 of the invention or a fragment thereof. According to the signal hypothesis, proteins secreted by mammalian cells have a signal peptide or secretory leader sequence which is cleaved from the mature protein once export of the growing protein chain across the rough endoplasmic reticulum has been initiated. Those of ordinary skill in the art are aware that polypeptides secreted by vertebrate cells generally

10 have a signal peptide fused to the N-terminus of the polypeptide, which is cleaved from the translated polypeptide to produce a secreted or "mature" form of the polypeptide. In certain embodiments, the native signal peptide, *e.g.* an immunoglobulin heavy chain or light chain signal peptide is used, or a functional derivative of that sequence that retains the ability to direct the secretion of the polypeptide that is operably associated with it.

15 Alternatively, a heterologous mammalian signal peptide, or a functional derivative thereof, may be used. For example, the wild-type leader sequence may be substituted with the leader sequence of human tissue plasminogen activator (TPA) or mouse β -glucuronidase.

DNA encoding a short protein sequence that could be used to facilitate later

20 purification (*e.g.* a histidine tag) or assist in labeling the T cell activating bispecific antigen binding molecule may be included within or at the ends of the T cell activating bispecific antigen binding molecule (fragment) or the antibody (fragment) encoding polynucleotide.

In a further embodiment, a host cell comprising one or more polynucleotides of the

25 invention is provided. In certain embodiments a host cell comprising one or more vectors of the invention is provided. The polynucleotides and vectors may incorporate any of the features, singly or in combination, described herein in relation to polynucleotides and vectors, respectively. In one such embodiment a host cell comprises (*e.g.* has been transformed or transfected with) a vector comprising a polynucleotide that encodes a T

30 cell activating bispecific antigen binding molecule or an antibody of the invention or a part thereof. As used herein, the term "host cell" refers to any kind of cellular system which can be engineered to generate the T cell activating bispecific antigen binding molecule, *e.g.*, the FolR1 T cell activating bispecific antigen binding molecules disclosed

herein, or antibody, e.g., anti-PD-1 antibodies, anti-PD-L1 antibodies, and anti-TIM3 antibodies of the invention or fragments thereof. Host cells suitable for replicating and for supporting expression of T cell activating bispecific antigen binding molecule and antibodies of the invention are well known in the art. Such cells may be transfected or
5 transduced as appropriate with the particular expression vector and large quantities of vector containing cells can be grown for seeding large scale fermenters to obtain sufficient quantities of the T cell activating bispecific antigen binding molecule and antibodies for clinical applications. Suitable host cells include prokaryotic microorganisms, such as *E. coli*, or various eukaryotic cells, such as Chinese hamster
10 ovary cells (CHO), insect cells, or the like. For example, polypeptides may be produced in bacteria in particular when glycosylation is not needed. After expression, the polypeptide may be isolated from the bacterial cell paste in a soluble fraction and can be further purified. In addition to prokaryotes, eukaryotic microbes such as filamentous fungi or yeast are suitable cloning or expression hosts for polypeptide-encoding vectors,
15 including fungi and yeast strains whose glycosylation pathways have been “humanized”, resulting in the production of a polypeptide with a partially or fully human glycosylation pattern. See Gerngross, *Nat Biotech* 22, 1409-1414 (2004), and Li et al., *Nat Biotech* 24, 210-215 (2006). Suitable host cells for the expression of (glycosylated) polypeptides are also derived from multicellular organisms (invertebrates and vertebrates). Examples of
20 invertebrate cells include plant and insect cells. Numerous baculoviral strains have been identified which may be used in conjunction with insect cells, particularly for transfection of *Spodoptera frugiperda* cells. Plant cell cultures can also be utilized as hosts. See e.g. US Patent Nos. 5,959,177, 6,040,498, 6,420,548, 7,125,978, and 6,417,429 (describing PLANTIBODIESTM technology for producing antibodies in transgenic plants). Vertebrate
25 cells may also be used as hosts. For example, mammalian cell lines that are adapted to grow in suspension may be useful. Other examples of useful mammalian host cell lines are monkey kidney CV1 line transformed by SV40 (COS-7); human embryonic kidney line (293 or 293T cells as described, e.g., in Graham et al., *J Gen Virol* 36, 59 (1977)), baby hamster kidney cells (BHK), mouse sertoli cells (TM4 cells as described, e.g., in
30 Mather, *Biol Reprod* 23, 243-251 (1980)), monkey kidney cells (CV1), African green monkey kidney cells (VERO-76), human cervical carcinoma cells (HELA), canine kidney cells (MDCK), buffalo rat liver cells (BRL 3A), human lung cells (W138), human liver cells (Hep G2), mouse mammary tumor cells (MMT 060562), TRI cells (as described,

e.g., in Mather et al., *Annals N.Y. Acad Sci* 383, 44-68 (1982)), MRC 5 cells, and FS4 cells. Other useful mammalian host cell lines include Chinese hamster ovary (CHO) cells, including dhfr⁻ CHO cells (Urlaub et al., *Proc Natl Acad Sci USA* 77, 4216 (1980)); and myeloma cell lines such as YO, NS0, P3X63 and Sp2/0. For a review of certain

- 5 mammalian host cell lines suitable for protein production, see, e.g., Yazaki and Wu, *Methods in Molecular Biology*, Vol. 248 (B.K.C. Lo, ed., Humana Press, Totowa, NJ), pp. 255-268 (2003). Host cells include cultured cells, e.g., mammalian cultured cells, yeast cells, insect cells, bacterial cells and plant cells, to name only a few, but also cells comprised within a transgenic animal, transgenic plant or cultured plant or animal tissue.
- 10 In one embodiment, the host cell is a eukaryotic cell, preferably a mammalian cell, such as a Chinese Hamster Ovary (CHO) cell, a human embryonic kidney (HEK) cell or a lymphoid cell (e.g., YO, NS0, Sp20 cell).

- Standard technologies are known in the art to express foreign genes in these systems. Cells expressing a polypeptide comprising either the heavy or the light chain of
- 15 an antigen binding domain such as an antibody, may be engineered so as to also express the other of the antibody chains such that the expressed product is an antibody that has both a heavy and a light chain.

- Any animal species of antibody, antibody fragment, antigen binding domain or variable region can be used in the T cell activating bispecific antigen binding molecules of the
- 20 invention. Non-limiting antibodies, antibody fragments, antigen binding domains or variable regions useful in the present invention can be of murine, primate, or human origin. If the T cell activating bispecific antigen binding molecule is intended for human use, a chimeric form of antibody may be used wherein the constant regions of the antibody are from a human. A humanized or fully human form of the antibody can also be
- 25 prepared in accordance with methods well known in the art (see e. g. U.S. Patent No. 5,565,332 to Winter). Humanization may be achieved by various methods including, but not limited to (a) grafting the non-human (e.g., donor antibody) CDRs onto human (e.g. recipient antibody) framework and constant regions with or without retention of critical framework residues (e.g. those that are important for retaining good antigen binding
- 30 affinity or antibody functions), (b) grafting only the non-human specificity-determining regions (SDRs or a-CDRs; the residues critical for the antibody-antigen interaction) onto human framework and constant regions, or (c) transplanting the entire non-human variable domains, but "cloaking" them with a human-like section by replacement of

surface residues. Humanized antibodies and methods of making them are reviewed, e.g., in Almagro and Fransson, *Front Biosci* 13, 1619-1633 (2008), and are further described, e.g., in Riechmann et al., *Nature* 332, 323-329 (1988); Queen et al., *Proc Natl Acad Sci USA* 86, 10029-10033 (1989); US Patent Nos. 5,821,337, 7,527,791, 6,982,321, and 7,087,409; Jones et al., *Nature* 321, 522-525 (1986); Morrison et al., *Proc Natl Acad Sci* 81, 6851-6855 (1984); Morrison and Oi, *Adv Immunol* 44, 65-92 (1988); Verhoeyen et al., *Science* 239, 1534-1536 (1988); Padlan, *Molec Immun* 31(3), 169-217 (1994); Kashmiri et al., *Methods* 36, 25-34 (2005) (describing SDR (a-CDR) grafting); Padlan, *Mol Immunol* 28, 489-498 (1991) (describing "resurfacing"); Dall'Acqua et al., *Methods* 36, 43-60 (2005) (describing "FR shuffling"); and Osbourn et al., *Methods* 36, 61-68 (2005) and Klimka et al., *Br J Cancer* 83, 252-260 (2000) (describing the "guided selection" approach to FR shuffling). Human antibodies and human variable regions can be produced using various techniques known in the art. Human antibodies are described generally in van Dijk and van de Winkel, *Curr Opin Pharmacol* 5, 368-74 (2001) and Lonberg, *Curr Opin Immunol* 20, 450-459 (2008). Human variable regions can form part of and be derived from human monoclonal antibodies made by the hybridoma method (see e.g. *Monoclonal Antibody Production Techniques and Applications*, pp. 51-63 (Marcel Dekker, Inc., New York, 1987)). Human antibodies and human variable regions may also be prepared by administering an immunogen to a transgenic animal that has been modified to produce intact human antibodies or intact antibodies with human variable regions in response to antigenic challenge (see e.g. Lonberg, *Nat Biotech* 23, 1117-1125 (2005)). Human antibodies and human variable regions may also be generated by isolating Fv clone variable region sequences selected from human-derived phage display libraries (see e.g., Hoogenboom et al. in *Methods in Molecular Biology* 178, 1-37 (O'Brien et al., ed., Human Press, Totowa, NJ, 2001); and McCafferty et al., *Nature* 348, 552-554; Clackson et al., *Nature* 352, 624-628 (1991)). Phage typically display antibody fragments, either as single-chain Fv (scFv) fragments or as Fab fragments.

In certain embodiments, the antigen binding moieties useful in the present invention are engineered to have enhanced binding affinity according to, for example, the methods disclosed in U.S. Pat. Appl. Publ. No. 2004/0132066, the entire contents of which are hereby incorporated by reference. The ability of the T cell activating bispecific antigen binding molecule of the invention to bind to a specific antigenic determinant can be measured either through an enzyme-linked immunosorbent assay (ELISA) or other

techniques familiar to one of skill in the art, e.g. surface plasmon resonance technique (analyzed on a BIACORE T100 system) (Liljeblad, et al., Glyco J 17, 323-329 (2000)), and traditional binding assays (Heeley, Endocr Res 28, 217-229 (2002)). Competition assays may be used to identify an antibody, antibody fragment, antigen binding domain or variable domain that competes with a reference antibody for binding to a particular antigen, e.g. an antibody that competes with the V9 antibody for binding to CD3. In certain embodiments, such a competing antibody binds to the same epitope (e.g. a linear or a conformational epitope) that is bound by the reference antibody. Detailed exemplary methods for mapping an epitope to which an antibody binds are provided in Morris (1996) "Epitope Mapping Protocols," in Methods in Molecular Biology vol. 66 (Humana Press, Totowa, NJ). In an exemplary competition assay, immobilized antigen (e.g. CD3) is incubated in a solution comprising a first labeled antibody that binds to the antigen (e.g. V9 antibody, described in US 6,054,297) and a second unlabeled antibody that is being tested for its ability to compete with the first antibody for binding to the antigen. The second antibody may be present in a hybridoma supernatant. As a control, immobilized antigen is incubated in a solution comprising the first labeled antibody but not the second unlabeled antibody. After incubation under conditions permissive for binding of the first antibody to the antigen, excess unbound antibody is removed, and the amount of label associated with immobilized antigen is measured. If the amount of label associated with immobilized antigen is substantially reduced in the test sample relative to the control sample, then that indicates that the second antibody is competing with the first antibody for binding to the antigen. See Harlow and Lane (1988) Antibodies: A Laboratory Manual ch.14 (Cold Spring Harbor Laboratory, Cold Spring Harbor, NY).

In certain embodiments, the antigen binding moieties useful in the present invention are engineered to have enhanced binding affinity according to, for example, the methods disclosed in U.S. Pat. Appl. Publ. No. 2004/0132066, the entire contents of which are hereby incorporated by reference. The ability of the T cell activating bispecific antigen binding molecule or the antibody of the invention to bind to a specific antigenic determinant can be measured either through an enzyme-linked immunosorbent assay (ELISA) or other techniques familiar to one of skill in the art, e.g. surface plasmon resonance technique (analyzed on a BIACORE T100 system) (Liljeblad, et al., Glyco J 17, 323-329 (2000)), and traditional binding assays (Heeley, Endocr Res 28, 217-229 (2002)). Competition assays may be used to identify an antibody, antibody fragment, antigen

binding domain or variable domain that competes with a reference antibody for binding to a particular antigen. In certain embodiments, such a competing antibody binds to the same epitope (e.g. a linear or a conformational epitope) that is bound by the reference antibody. Detailed exemplary methods for mapping an epitope to which an antibody binds are
5 provided in Morris (1996) "Epitope Mapping Protocols," in *Methods in Molecular Biology* vol. 66 (Humana Press, Totowa, NJ). In an exemplary competition assay, immobilized antigen is incubated in a solution comprising a first labeled antibody that binds to the antigen and a second unlabeled antibody that is being tested for its ability to compete with the first antibody for binding to the antigen. The second antibody may be
10 present in a hybridoma supernatant. As a control, immobilized antigen is incubated in a solution comprising the first labeled antibody but not the second unlabeled antibody.

After incubation under conditions permissive for binding of the first antibody to the antigen, excess unbound antibody is removed, and the amount of label associated with immobilized antigen is measured. If the amount of label associated with immobilized
15 antigen is substantially reduced in the test sample relative to the control sample, then that indicates that the second antibody is competing with the first antibody for binding to the antigen. See Harlow and Lane (1988) *Antibodies: A Laboratory Manual* ch.14 (Cold Spring Harbor Laboratory, Cold Spring Harbor, NY).

T cell activating bispecific antigen binding molecules and antibodies prepared as
20 described herein may be purified by art-known techniques such as high performance liquid chromatography, ion exchange chromatography, gel electrophoresis, affinity chromatography, size exclusion chromatography, and the like. The actual conditions used to purify a particular protein will depend, in part, on factors such as net charge, hydrophobicity, hydrophilicity etc., and will be apparent to those having skill in the art.
25 For affinity chromatography purification an antibody, ligand, receptor or antigen can be used to which the bispecific antibody or the antibody binding to DR5 binds. For example, for affinity chromatography purification of bispecific antibodies of the invention, a matrix with protein A or protein G may be used. Sequential Protein A or G affinity chromatography and size exclusion chromatography can be used to isolate a bispecific
30 antibody essentially as described in the Examples. The purity of the bispecific antibody or the antibody binding to DR5 can be determined by any of a variety of well-known analytical methods including gel electrophoresis, high pressure liquid chromatography, and the like.

F. Assays

T cell activating bispecific antigen binding molecules, e.g., a T cell activating bispecific antigen binding molecules comprising a first antigen binding site specific for Folate Receptor 1 (FolR1) and a second antigen binding site specific for CD3, and antibodies, e.g., anti-PD-1 axis binding antagonist antibodies and anti-TIM3 antagonist antibodies provided herein may be identified, screened for, or characterized for their physical/chemical properties and/or biological activities by various assays known in the art.

1. Affinity assays

The affinity of the T cell activating bispecific antigen binding molecules, e.g., a T cell activating bispecific antigen binding molecules comprising a first antigen binding site specific for Folate Receptor 1 (FolR1) and a second antigen binding site specific for CD3, and antibodies, e.g., anti-PD-1 axis binding antagonist antibodies and anti-TIM3 antagonist antibodies provided herein for their respective antigen, e.g., FolR1, PD-1, PD-L1, TIM3, can be determined in accordance with the methods set forth in the Examples by surface plasmon resonance (SPR), using standard instrumentation such as a BIAcore instrument (GE Healthcare), and receptors or target proteins such as may be obtained by recombinant expression. Alternatively, binding of T cell activating bispecific antigen binding molecules and antibodies provided therein to their respective antigen may be evaluated using cell lines expressing the particular receptor or target antigen, for example by flow cytometry (FACS).

K_D may be measured by surface plasmon resonance using a BIAcore® T100 machine (GE Healthcare) at 25°C. To analyze the interaction between the Fc-portion and Fc receptors, His-tagged recombinant Fc-receptor is captured by an anti-Penta His antibody (Qiagen) ("Penta His" disclosed as SEQ ID NO: 392) immobilized on CM5 chips and the bispecific constructs are used as analytes. Briefly, carboxymethylated dextran biosensor chips (CM5, GE Healthcare) are activated with N-ethyl-N'-(3-dimethylaminopropyl)-carbodiimide hydrochloride (EDC) and N-hydroxysuccinimide (NHS) according to the supplier's instructions. Anti Penta-His antibody ("Penta His" disclosed as SEQ ID NO: 392) is diluted with 10 mM sodium acetate, pH 5.0, to 40 µg/ml before injection at a flow rate of 5 µl/min to achieve approximately 6500 response units (RU) of coupled protein. Following the injection of the ligand, 1 M ethanolamine is injected to block unreacted groups. Subsequently the Fc-receptor is captured for 60 s at 4

or 10 nM. For kinetic measurements, four-fold serial dilutions of the bispecific construct (range between 500 nM and 4000 nM) are injected in HBS-EP (GE Healthcare, 10 mM HEPES, 150 mM NaCl, 3 mM EDTA, 0.05 % Surfactant P20, pH 7.4) at 25 °C at a flow rate of 30 µl/min for 120 s.

5 To determine the affinity to the target antigen, bispecific constructs are captured by an anti human Fab specific antibody (GE Healthcare) that is immobilized on an activated CM5-sensor chip surface as described for the anti Penta-His antibody ("Penta His" disclosed as SEQ ID NO: 392). The final amount of coupled protein is is approximately 12000 RU. The bispecific constructs are captured for 90 s at 300 nM. The
10 target antigens are passed through the flow cells for 180 s at a concentration range from 250 to 1000 nM with a flowrate of 30 µl/min. The dissociation is monitored for 180 s.

Bulk refractive index differences are corrected for by subtracting the response obtained on reference flow cell. The steady state response was used to derive the dissociation constant K_D by non-linear curve fitting of the Langmuir binding isotherm.
15 Association rates (k_{on}) and dissociation rates (k_{off}) are calculated using a simple one-to-one Langmuir binding model (BIAcore® T100 Evaluation Software version 1.1.1) by simultaneously fitting the association and dissociation sensorgrams. The equilibrium dissociation constant (K_D) is calculated as the ratio k_{off}/k_{on} . See, e.g., Chen et al., J Mol Biol 293, 865-881 (1999).

20

2. Binding assays and other assays

In one aspect, a T cell activating bispecific antigen binding molecules, e.g., a T cell activating bispecific antigen binding molecules comprising a first antigen binding site specific for Folate Receptor 1 (FolR1) and a second antigen binding site specific for CD3,
25 and antibodies, e.g., anti-PD-1 axis binding antagonist antibodies and anti-TIM3 antagonist antibodies of the invention is tested for its antigen binding activity, e.g., by known methods such as ELISA, Western blot, etc.

In another aspect, competition assays may be used to identify an antibody or fragment that competes with a specific reference antibody for binding to the respective
30 antigens. In certain embodiments, such a competing antibody binds to the same epitope (e.g., a linear or a conformational epitope) that is bound by a specific reference antibody. Detailed exemplary methods for mapping an epitope to which an antibody binds are provided in Morris (1996) "Epitope Mapping Protocols," in *Methods in Molecular*

Biology vol. 66 (Humana Press, Totowa, NJ). Further methods are described in the example section.

3. Activity assays

- 5 In one aspect, assays are provided for identifying T cell activating bispecific antigen binding molecules, e.g., a T cell activating bispecific antigen binding molecules comprising a first antigen binding site specific for Folate Receptor 1 (FolR1) and a second antigen binding site specific for CD3, and antibodies, e.g., anti-PD-1 axis binding antagonist antibodies and anti-TIM3 antagonist antibodies provided herein having
- 10 biological activity. Biological activity may include, e.g., inducing DNA fragmentation, induction of apoptosis and lysis of targeted cells. Antibodies having such biological activity *in vivo* and/or *in vitro* are also provided.

In certain embodiments, T cell activating antigen binding molecule and antibody of the invention is tested for such biological activity. Assays for detecting cell lysis (e.g.

15 by measurement of LDH release) or apoptosis (e.g. using the TUNEL assay) are well known in the art. Assays for measuring ADCC or CDC are also described in WO 2004/065540 (see Example 1 therein), the entire content of which is incorporated herein by reference.

20 G. Pharmaceutical Formulations

Pharmaceutical formulations of a T cell activating bispecific antigen binding molecules, e.g., a T cell activating bispecific antigen binding molecule comprising a first antigen binding site specific for Folate Receptor 1 (FolR1) and a second antigen binding site specific for CD3, and antibodies, e.g., anti-PD-1 axis binding antagonist antibodies

25 and anti-TIM3 antagonist antibodies as described herein are prepared by mixing such T cell activating bispecific antigen binding molecules or antibody having the desired degree of purity with one or more optional pharmaceutically acceptable carriers (*Remington's Pharmaceutical Sciences* 16th edition, Osol, A. Ed. (1980)), in the form of lyophilized formulations or aqueous solutions. Pharmaceutically acceptable carriers are generally

30 nontoxic to recipients at the dosages and concentrations employed, and include, but are not limited to: buffers such as phosphate, citrate, and other organic acids; antioxidants including ascorbic acid and methionine; preservatives (such as octadecyldimethylbenzyl ammonium chloride; hexamethonium chloride; benzalkonium chloride; benzethonium

chloride; phenol, butyl or benzyl alcohol; alkyl parabens such as methyl or propyl paraben; catechol; resorcinol; cyclohexanol; 3-pentanol; and m-cresol); low molecular weight (less than about 10 residues) polypeptides; proteins, such as serum albumin, gelatin, or immunoglobulins; hydrophilic polymers such as polyvinylpyrrolidone; amino acids such as glycine, glutamine, asparagine, histidine, arginine, or lysine; monosaccharides, disaccharides, and other carbohydrates including glucose, mannose, or dextrins; chelating agents such as EDTA; sugars such as sucrose, mannitol, trehalose or sorbitol; salt-forming counter-ions such as sodium; metal complexes (*e.g.* Zn-protein complexes); and/or non-ionic surfactants such as polyethylene glycol (PEG). Exemplary pharmaceutically acceptable carriers herein further include interstitial drug dispersion agents such as soluble neutral-active hyaluronidase glycoproteins (sHASEGP), for example, human soluble PH-20 hyaluronidase glycoproteins, such as rHuPH20 (HYLENEX[®], Baxter International, Inc.). Certain exemplary sHASEGPs and methods of use, including rHuPH20, are described in US Patent Publication Nos. 2005/0260186 and 2006/0104968. In one aspect, a sHASEGP is combined with one or more additional glycosaminoglycanases such as chondroitinases.

Exemplary lyophilized antibody formulations are described in US Patent No. 6,267,958. Aqueous antibody formulations include those described in US Patent No. 6,171,586 and WO2006/044908, the latter formulations including a histidine-acetate buffer.

The formulation herein may also contain more than one active ingredients as necessary for the particular indication being treated, preferably those with complementary activities that do not adversely affect each other. Such active ingredients are suitably present in combination in amounts that are effective for the purpose intended.

Active ingredients may be entrapped in microcapsules prepared, for example, by coacervation techniques or by interfacial polymerization, for example, hydroxymethylcellulose or gelatin-microcapsules and poly-(methylmethacrylate) microcapsules, respectively, in colloidal drug delivery systems (for example, liposomes, albumin microspheres, microemulsions, nano-particles and nanocapsules) or in macroemulsions. Such techniques are disclosed in *Remington's Pharmaceutical Sciences* 16th edition, Osol, A. Ed. (1980).

Sustained-release preparations may be prepared. Suitable examples of sustained-release preparations include semipermeable matrices of solid hydrophobic polymers

containing the antibody, which matrices are in the form of shaped articles, *e.g.* films, or microcapsules.

The formulations to be used for *in vivo* administration are generally sterile. Sterility may be readily accomplished, *e.g.*, by filtration through sterile filtration
5 membranes.

H. Therapeutic Methods and Compositions

The therapeutic combinations comprising one or more of the T cell activating bispecific antigen binding molecules and the anti-PD-1 axis binding antagonist antibody and, optionally, the TIM3 antagonist provided herein may be used in therapeutic methods.

10 In one aspect, a T cell activating bispecific antigen binding molecules that binds to Folate Receptor 1 (FolR1) and CD3 for use as a medicament is provided for use in combination with an anti-PD-1 axis binding antagonist antibody. In certain embodiments, a T cell activating bispecific antigen binding molecules that binds to FolR1 and CD3 for use in combination with an anti-PD-1 axis binding antagonist antibody is provided for use
15 in a method of treatment. In certain embodiments, the combination further comprises a TIM3 antagonist, *e.g.*, an anti-TIM3 antagonist antibody. In certain embodiments, the invention provides a T cell activating bispecific antigen binding molecules that binds to FolR1 and CD3 and an anti-PD-1 axis binding antagonist antibody for use in a method of treating an individual having cancer comprising administering to the individual an
20 effective amount of the T cell activating bispecific antigen binding molecules that binds to FolR1 and CD3 and the anti-PD-1 axis binding antagonist antibody. In one such embodiment, the method further comprises administering to the individual an effective amount of at least one TIM3 antagonist, *e.g.*, as described below. An “individual” according to any of the above embodiments is preferably a human. In one preferred
25 embodiment, said cancer is pancreatic cancer, sarcoma or colorectal carcinoma. In other embodiments, the cancer is colorectal cancer, sarcoma, head and neck cancers, squamous cell carcinomas, breast cancer, pancreatic cancer, gastric cancer, non-small-cell lung carcinoma, small-cell lung cancer or mesothelioma. In embodiments in which the cancer is breast cancer, the breast cancer may be triple negative breast cancer.

30 In a further aspect, the invention provides the use of a therapeutic combination comprising a T cell activating bispecific antigen binding molecules that binds to FolR1 and CD3 and an anti-PD-1 axis binding antagonist antibody in the manufacture or preparation of a medicament. In one embodiment, the combination further comprises a

TIM3 antagonist. In one embodiment, the medicament is for treatment of cancer. In a further embodiment, the medicament is for use in a method of treating cancer comprising administering to an individual having cancer an effective amount of the medicament. In one such embodiment, the method further comprises administering to the individual an effective amount of at least one additional therapeutic agent, e.g., as described below. An “individual” according to any of the above embodiments may be a human.

In a further aspect, the invention provides a method for treating cancer. In one embodiment, the method comprises administering to an individual having cancer an effective amount of a therapeutic combination comprising a T cell activating bispecific antigen binding molecules that binds to FolR1 and CD3 and an anti-PD-1 axis binding antagonist antibody. In one such embodiment, the method further comprises administering to the individual an effective amount of at least one additional therapeutic agent, as described below. In one such embodiment, the at least one additional therapeutic agent is an anti-TIM3 antagonist antibody. An “individual” according to any of the above embodiments may be a human. In one preferred embodiment said cancer is pancreatic cancer, sarcoma or colorectal carcinoma. In other embodiments, the cancer is colorectal cancer, sarcoma, head and neck cancers, squamous cell carcinomas, breast cancer, pancreatic cancer, gastric cancer, non-small-cell lung carcinoma, small-cell lung cancer or mesothelioma.

In a further aspect, the invention provides pharmaceutical formulations comprising any of the T cell activating bispecific antigen binding molecules that binds to FolR1 and CD3 provided herein, e.g., for use in any of the above therapeutic methods, and an anti-PD-1 axis binding antagonist antibody. In one embodiment, a pharmaceutical formulation comprises any of the T cell activating bispecific antigen binding molecules that binds to FolR1 provided herein and a pharmaceutically acceptable carrier. In another embodiment, a pharmaceutical formulation comprises any of T cell activating bispecific antigen binding molecules that binds to FolR1 and CD3 and an anti-PD-1 axis binding antagonist antibody provided herein and at least one additional therapeutic agent, e.g., as described below.

A bispecific antibody can be administered by any suitable means, including parenteral, intrapulmonary, and intranasal, and, if desired for local treatment, intralesional administration. Parenteral infusions include intramuscular, intravenous, intraarterial, intraperitoneal, or subcutaneous administration. Dosing can be by any suitable route, e.g.

by injections, such as intravenous or subcutaneous injections, depending in part on whether the administration is brief or chronic. Various dosing schedules including but not limited to single or multiple administrations over various time-points, bolus administration, and pulse infusion are contemplated herein.

5 Bispecific antibodies may be formulated, dosed, and administered in a fashion consistent with good medical practice. Factors for consideration in this context include the particular disorder being treated, the particular mammal being treated, the clinical condition of the individual patient, the cause of the disorder, the site of delivery of the agent, the method of administration, the scheduling of administration, and other factors
10 known to medical practitioners. The bispecific antibody need not be, but is optionally formulated with one or more agents currently used to prevent or treat the disorder in question. The effective amount of such other agents depends on the amount of antibody present in the formulation, the type of disorder or treatment, and other factors discussed above. These are generally used in the same dosages and with administration routes as
15 described herein, or about from 1 to 99% of the dosages described herein, or in any dosage and by any route that is empirically/clinically determined to be appropriate.

For the prevention or treatment of disease, the appropriate dosage of a bispecific antibody will depend on the type of disease to be treated, the type of antibody, the severity and course of the disease, whether the bispecific antibody is administered for
20 preventive or therapeutic purposes, previous therapy, the patient's clinical history and response to the bispecific antibody and the discretion of the attending physician. The bispecific antibody is suitably administered to the patient at one time or over a series of treatments. Depending on the type and severity of the disease, about 1 μ g/kg to 15 mg/kg (e.g. 0.1mg/kg-10mg/kg) of the bispecific antibody or the novel antibody binding to DR5
25 can be an initial candidate dosage for administration to the patient, whether, for example, by one or more separate administrations, or by continuous infusion. One typical daily dosage might range from about 1 μ g/kg to 100 mg/kg or more, depending on the factors mentioned above. For repeated administrations over several days or longer, depending on the condition, the treatment would generally be sustained until a desired suppression of
30 disease symptoms occurs. One exemplary dosage of the bispecific would be in the range from about 0.05 mg/kg to about 10 mg/kg. Thus, one or more doses of about 0.5 mg/kg, 2.0 mg/kg, 4.0 mg/kg or 10 mg/kg may be administered to the patient. Such doses may be administered intermittently, e.g. every week or every three weeks (e.g. such that the

patient receives from about two to about twenty, or *e.g.* about six doses of the bispecific antibody). An initial higher loading dose, followed by one or more lower doses may be administered. However, other dosage regimens may be useful. The progress of this therapy is easily monitored by conventional techniques and assays.

5 It is understood that any of the above formulations or therapeutic methods may be carried out using an immunoconjugate of the invention in place of or in addition to the T cell activating bispecific antigen binding molecules that binds to FolR1 and CD3 and the anti-PD-1 axis binding antagonist antibody, and, optionally, the anti-TIM3 antagonist antibody.

10

I. Articles of Manufacture

In another aspect of the invention, an article of manufacture containing materials useful for the treatment, prevention and/or diagnosis of the disorders described above is provided. The article of manufacture comprises a container and a label or package insert
15 on or associated with the container. Suitable containers include, for example, bottles, vials, syringes, IV solution bags, etc. The containers may be formed from a variety of materials such as glass or plastic. The container holds a composition which is by itself or combined with another composition effective for treating, preventing and/or diagnosing the condition and may have a sterile access port (for example the container may be an
20 intravenous solution bag or a vial having a stopper pierceable by a hypodermic injection needle). At least one active agent in the composition is a bispecific antibody and an additional active agent is the further chemotherapeutic agent as described herein. The label or package insert indicates that the composition is used for treating the condition of choice. Moreover, the article of manufacture may comprise (a) a first container with a
25 composition contained therein, wherein the composition comprises a bispecific antibody; and (b) a second container with a composition contained therein, wherein the composition comprises a further cytotoxic or otherwise therapeutic agent. The article of manufacture in this embodiment of the invention may further comprise a package insert indicating that the compositions can be used to treat a particular condition. Alternatively, or
30 additionally, the article of manufacture may further comprise a second (or third) container comprising a pharmaceutically-acceptable buffer, such as bacteriostatic water for injection (BWI), phosphate-buffered saline, Ringer's solution and dextrose solution. It

may further include other materials desirable from a commercial and user standpoint, including other buffers, diluents, filters, needles, and syringes.

It is understood that any of the above articles of manufacture may include an immunoconjugate of the invention in place of or in addition to the T cell activating
5 bispecific antigen binding molecules that binds to FolR1 and CD3 and the anti-PD-1 axis binding antagonist antibody and, optionally, the anti-TIM3 antagonist antibody.

III. EXAMPLES

The following are examples of methods and compositions of the invention. It is
10 understood that various other embodiments may be practiced, given the general description provided above.

The following are examples of methods and compositions of the invention. It is understood that various other embodiments may be practiced, given the general description provided above.

15

General methods

Recombinant DNA Techniques

Standard methods were used to manipulate DNA as described in Sambrook et al., Molecular cloning: A laboratory manual; Cold Spring Harbor Laboratory Press, Cold
20 Spring Harbor, New York, 1989. The molecular biological reagents were used according to the manufacturers' instructions. General information regarding the nucleotide sequences of human immunoglobulins light and heavy chains is given in: Kabat, E.A. et al., (1991) Sequences of Proteins of Immunological Interest, 5th ed., NIH Publication No. 91-3242.

25

DNA Sequencing

DNA sequences were determined by double strand sequencing.

Gene Synthesis

30 Desired gene segments where required were either generated by PCR using appropriate templates or were synthesized by Genearth AG (Regensburg, Germany) from synthetic oligonucleotides and PCR products by automated gene synthesis. In cases where no exact

gene sequence was available, oligonucleotide primers were designed based on sequences from closest homologues and the genes were isolated by RT-PCR from RNA originating from the appropriate tissue. The gene segments flanked by singular restriction endonuclease cleavage sites were cloned into standard cloning / sequencing vectors. The plasmid DNA was purified from transformed bacteria and concentration determined by UV spectroscopy. The DNA sequence of the subcloned gene fragments was confirmed by DNA sequencing. Gene segments were designed with suitable restriction sites to allow sub-cloning into the respective expression vectors. All constructs were designed with a 5'-end DNA sequence coding for a leader peptide which targets proteins for secretion in eukaryotic cells.

Isolation of primary human pan T cells from PBMCs

Peripheral blood mononuclear cells (PBMCs) were prepared by Histopaque density centrifugation from enriched lymphocyte preparations (buffy coats) obtained from local blood banks or from fresh blood from healthy human donors. Briefly, blood was diluted with sterile PBS and carefully layered over a Histopaque gradient (Sigma, H8889). After centrifugation for 30 minutes at 450 x g at room temperature (brake switched off), part of the plasma above the PBMC containing interphase was discarded. The PBMCs were transferred into new 50 ml Falcon tubes and tubes were filled up with PBS to a total volume of 50 ml. The mixture was centrifuged at room temperature for 10 minutes at 400 x g (brake switched on). The supernatant was discarded and the PBMC pellet washed twice with sterile PBS (centrifugation steps at 4°C for 10 minutes at 350 x g). The resulting PBMC population was counted automatically (ViCell) and stored in RPMI1640 medium, containing 10% FCS and 1% L-alanyl-L-glutamine (Biochrom, K0302) at 37°C, 5% CO₂ in the incubator until assay start.

T cell enrichment from PBMCs was performed using the Pan T Cell Isolation Kit II (Miltenyi Biotec #130-091-156), according to the manufacturer's instructions. Briefly, the cell pellets were diluted in 40 µl cold buffer per 10 million cells (PBS with 0.5% BSA, 2 mM EDTA, sterile filtered) and incubated with 10 µl Biotin-Antibody Cocktail per 10 million cells for 10 min at 4°C. 30 µl cold buffer and 20 µl Anti-Biotin magnetic beads per 10 million cells were added, and the mixture incubated for another 15 min at 4°C. Cells were washed by adding 10-20x the current volume and a subsequent centrifugation step at 300 x g for 10 min. Up to 100 million cells were resuspended in 500 µl buffer.

Magnetic separation of unlabeled human pan T cells was performed using LS columns (Miltenyi Biotec #130-042-401) according to the manufacturer's instructions. The resulting T cell population was counted automatically (ViCell) and stored in AIM-V medium at 37°C, 5% CO₂ in the incubator until assay start (not longer than 24 h).

5

Isolation of primary human naive T cells from PBMCs

Peripheral blood mononuclear cells (PBMCs) were prepared by Histopaque density centrifugation from enriched lymphocyte preparations (buffy coats) obtained from local blood banks or from fresh blood from healthy human donors. T-cell enrichment from
10 PBMCs was performed using the Naive CD8⁺ T cell isolation Kit from Miltenyi Biotec (#130-093-244), according to the manufacturer's instructions, but skipping the last isolation step of CD8⁺ T cells (also see description for the isolation of primary human pan T cells).

15 Isolation of murine pan T cells from splenocytes

Spleens were isolated from C57BL/6 mice, transferred into a GentleMACS C-tube (Miltenyi Biotec #130-093-237) containing MACS buffer (PBS + 0.5% BSA + 2 mM EDTA) and dissociated with the GentleMACS Dissociator to obtain single-cell suspensions according to the manufacturer's instructions. The cell suspension was passed
20 through a pre-separation filter to remove remaining undissociated tissue particles. After centrifugation at 400 x g for 4 min at 4°C, ACK Lysis Buffer was added to lyse red blood cells (incubation for 5 min at room temperature). The remaining cells were washed with MACS buffer twice, counted and used for the isolation of murine pan T cells. The negative (magnetic) selection was performed using the Pan T Cell Isolation Kit from
25 Miltenyi Biotec (#130-090-861), following the manufacturer's instructions. The resulting T cell population was automatically counted (ViCell) and immediately used for further assays.

Isolation of primary cynomolgus PBMCs from heparinized blood

30 Peripheral blood mononuclear cells (PBMCs) were prepared by density centrifugation from fresh blood from healthy cynomolgus donors, as follows: Heparinized blood was diluted 1:3 with sterile PBS, and Lymphoprep medium (Axon Lab #1114545) was diluted to 90% with sterile PBS. Two volumes of the diluted blood were layered over one volume

of the diluted density gradient and the PBMC fraction was separated by centrifugation for 30 min at 520 x g, without brake, at room temperature. The PBMC band was transferred into a fresh 50 ml Falcon tube and washed with sterile PBS by centrifugation for 10 min at 400 x g at 4°C. One low-speed centrifugation was performed to remove the platelets
5 (15 min at 150 x g, 4°C), and the resulting PBMC population was automatically counted (ViCell) and immediately used for further assays.

Example 1

Purification of biotinylated Folate receptor-Fc fusions

- 10 To generate new antibodies against human FolR1 the following antigens and screening tools were generated as monovalent Fc fusion proteins (the extracellular domain of the antigen linked to the hinge region of Fc-knob which is co-expressed with an Fc-hole molecule). The antigen genes were synthesized (Geneart, Regensburg, Germany) based on sequences obtained from GenBank or SwissProt and inserted into expression vectors
15 to generate fusion proteins with Fc-knob with a C-terminal Avi-tag for in vivo or in vitro biotinylation. In vivo biotinylation was achieved by co-expression of the bacterial birA gene encoding a bacterial biotin ligase during production. Expression of all genes was under control of a chimeric MPSV promoter on a plasmid containing an oriP element for stable maintenance of the plasmids in EBNA containing cell lines.
- 20 For preparation of the biotinylated monomeric antigen/Fc fusion molecules, exponentially growing suspension HEK293 EBNA cells were co-transfected with three vectors encoding the two components of fusion protein (knob and hole chains) as well as BirA, an enzyme necessary for the biotinylation reaction. The corresponding vectors were used at a 9.5 : 9.5 : 1 ratio ("antigen ECD- Fc knob-avi tag" : "Fc hole" : "BirA").
- 25 For protein production in 500 ml shake flasks, 400 million HEK293 EBNA cells were seeded 24 hours before transfection. For transfection cells were centrifuged for 5 minutes at 210 g, and supernatant was replaced by pre-warmed CD CHO medium. Expression vectors were resuspended in 20 mL of CD CHO medium containing 200 µg of vector DNA. After addition of 540 µL of polyethylenimine (PEI), the solution was mixed for 15
30 seconds and incubated for 10 minutes at room temperature. Afterwards, cells were mixed with the DNA/PEI solution, transferred to a 500 mL shake flask and incubated for 3 hours at 37°C in an incubator with a 5% CO₂ atmosphere. After the incubation, 160 mL of F17

medium was added and cells were cultured for 24 hours. One day after transfection, 1 mM valproic acid and 7% Feed 1 (Lonza) were added to the culture. The production medium was also supplemented with 100 μ M biotin. After 7 days of culturing, the cell supernatant was collected by spinning down cells for 15 min at 210 g. The solution was
5 sterile filtered (0.22 μ m filter), supplemented with sodium azide to a final concentration of 0.01 % (w/v), and kept at 4°C.

Secreted proteins were purified from cell culture supernatants by affinity chromatography using Protein A, followed by size exclusion chromatography. For affinity chromatography, the supernatant was loaded on a HiTrap ProteinA HP column (CV = 5
10 mL, GE Healthcare) equilibrated with 40 mL 20 mM sodium phosphate, 20 mM sodium citrate pH 7.5. Unbound protein was removed by washing with at least 10 column volumes of 20 mM sodium phosphate, 20 mM sodium citrate pH 7.5. The bound protein was eluted using a linear pH-gradient created over 20 column volumes of 20 mM sodium citrate, 100 mM sodium chloride, 100 mM glycine, pH 3.0 . The column was then washed
15 with 10 column volumes of 20 mM sodium citrate, 100 mM sodium chloride, 100 mM glycine, pH 3.0. pH of collected fractions was adjusted by adding 1/10 (v/v) of 0.5 M sodium phosphate, pH 8.0. The protein was concentrated and filtered prior to loading on a HiLoad Superdex 200 column (GE Healthcare) equilibrated with 20 mM histidine, 140 mM sodium chloride, pH 6.0.

20 The protein concentration was determined by measuring the optical density (OD) at 280 nm, using the molar extinction coefficient calculated on the basis of the amino acid sequence. Purity and molecular weight of the FolR1-Fc-fusion was analyzed by SDS capillary electrophoresis in the presence and absence of a reducing agent following the manufacturer instructions (instrument Caliper LabChipGX, Perkin Elmer). The aggregate
25 content of samples was analyzed using a TSKgel G3000 SW XL analytical size-exclusion column (Tosoh) equilibrated in 25 mM K₂HPO₄, 125 mM NaCl, 200 mM L-arginine monohydrochloride, 0.02 % (w/v) NaN₃, pH 6.7 running buffer at 25°C.

Purified antigen-Fc-fusion proteins were analyzed by surface plasmon resonance assays using commercially available antibodies to confirm correct and natural conformation of
30 the antigens (data not shown).

Table 1: Antigens produced for isolation, selection and counter selection of human FolR1 antibodies

-155-

Antigen	ECD (aa)	Accession number	Sequence	Seq ID No
human FolR1	25 - 234	P15328	RIAWARTELLNVCMAKHHKEKPGPEDKLHEQCRPWR KNACCSTNTSQEAHKDVSYLRFNWNHCGEMAPACKR HFIQDTCLYECSPNLGPWQQVDQSWRKERVNLNPLC KEDCEQWWEDCRTSYTCKSNWHKGWNWTSGFNKA AACQPFHFYFPTPTVLCNEIWTSHSYKVSNSYRSGS GRCIQMWFDPAQGNPNNEEVARFYAAAM	227
human FolR2	17 - 230	P14207	TMCSAQDRTDLLNVCMDAKHHKTKPGPEDKLHDQCSP WKKNACCTASTSQELHKDTSRLYNFNWDHCGKMEPAC KRHF IQDTCLYECSPNLGPWQQVQSWRKERFLDVP LCKEDCQRWWEDCHTSHTCKSNWHRGWDWTSGVNKC PAGALCRTFESYFPTPAALCEGLWSHSYKVSNSYRSGS GRCIQMWFDPAQGNPNNEEVARFYAAAMHVN	228
human FolR3	24 - 243	P41439	SARARTDLLNVCMAKHHKTPSPEDELYGCSPWKK NACCTASTSQELHKDTSRLYNFNWDHCGKMEPTCKRH FIQDSCLYECSPNLGPWIRQVQSWRKERILNPLCK EDCERWWEDCRTSYTCKSNWHKGWNWTSGINECPAGA LCSTFESYFPTPAALCEGLWSHSFKVSNSYRSGSGRCI QMWFDPAQGNPNNEEVAKFYAAAMNAGAPSRGIIDS	229
murine FolR1	25 - 232	P35846	TRARTELLNVCMDAKHHKEKPGPEDNLHDQCSPWKT N SCCSTNTSQEAHKDISYLRFNWNHCGTMTSECKRHF IQDTCLYECSPNLGPWQQVDQSWRKERILDVPLCKE DCQQWWEDCQSSFTCKSNWHKGWNWSSGHNECPVGAS CHPFTFYFPTSAALCEEIWSHSYKLSNSYRSGSGRCI QMWFDPAQGNPNNEEVARFYAEAMS	230
cynomolg us FolR1	25 - 234	G7PR14	EAQTRTARARTELLNVCMAKHHKEKPGPEDKLHEQ CRPWKNACCSTNTSQEAHKDVSYLRFNWNHCGEMAP ACKRHFIQDTCLYECSPNLGPWQQVDQSWRKERVNL N VPLCKEDCERWWEDCRTSYCKSNWHKGWNWTSGFNKC PVGAAACQPFHFYFPTPTVLCNEIWTYSYKVSNSYRSGS GRCIQMWFDPAQGNPNNEEVARFYAAAMS	231

Table 2: Summary of the yield and final monomer content of the FolR- Fc- fusions.

Antigen	Monomer [%] (SEC)	Yield
huFolR1	100	30 mg/L
cyFolR1	100	32 mg/L
muFolR1	100	31 mg/L
huFolR2	100	16 mg/L
huFolR3	95	38 mg/L

5

Example 2**Generation of common light chain with CD3ε specificity**

The T cell activating bispecific molecules described herein comprise at least one CD3 binding moiety. This moiety can be generated by immunizing laboratory animals,

screening phage library or using known anti-CD3 antibodies. The common light chain with CD3 ϵ specificity was generated by humanizing the light chain of a murine parental anti-CD3 ϵ antibody (CH2527). For humanization of an antibody of non-human origin, the CDR residues from the non-human antibody (donor) have to be transplanted onto the framework of a human (acceptor) antibody. Generally, acceptor framework sequences are selected by aligning the sequence of the donor to a collection of potential acceptor sequences and choosing one that has either reasonable homology to the donor, or shows similar amino acids at some positions critical for structure and activity. In the present case, the search for the antibody acceptor framework was performed by aligning the mouse VL-domain sequence of the parental antibody to a collection of human germline sequences and choosing the human sequence that showed high sequence identity. Surprisingly, a good match in terms of framework sequence homology was found in a rather infrequent human light chain belonging to the V-domain family 7 of the lambda type, more precisely, hVL_7_46 (IMGT nomenclature, GenBank Acc No. Z73674). This infrequent human light chain was subsequently chosen as acceptor framework for humanization of the light chain of CH2527. The three complementarity determining regions (CDRs) of the mouse light chain variable domain were grafted onto this acceptor framework. Since the framework 4 region is not part of the variable region of the germline V-gene, the alignment for this region (J-element) was done individually. Hence the IGLJ3-02 sequence was chosen for humanization of this light chain.

Thirteen humanized variants were generated (CH2527-VL7_46-1 to VL7_46-10, VL7_46-12 to VL7_46-14). These differ in framework residues (and combinations thereof) that were back-mutated to the murine V-domain sequence or in CDR-residues (Kabat definition) that could be kept identical to the human germline sequence. The following framework residues outside the CDRs were back-mutated to the murine residues in the final humanized VL-domain variant VL7_46-13 (murine residues listed): V36, E38, F44, G46, G49, and G57, respectively. The human J-element IGLJ3-02 was 100% identical to the J-element of the murine parental antibody.

Example 3

SPR assessment of humanized variants with CD3 ϵ specificity

Humanized VL variants were assessed as chimera in a 2+1 TCB format, i.e. humanized light chain V-domains were paired with murine heavy chain V-domains. SPR assessment

was carried out on a ProteOn XPR36 instrument (Bio-Rad). More precisely, the variants were captured directly from the culture supernatant on an anti-Fab derivatized GLM sensorchip (Goat Anti-Human IgG, F(ab')₂ Fragment Specific, Jackson ImmunoResearch) in vertical orientation. The following analytes were subsequently injected horizontally as single concentrations to assess binding to human and cynomolgus CD3ε: 3μM hu CD3ε(-1-26)-Fc(knob)-avi (ID807) and 2.5μM cy CD3ε(-1-26)-Fc(knob)-Avi-Fc(hole) (ID873), respectively. Binding responses were qualitatively compared to binding of the murine control construct and graded + (comparable binding observed), +/- (reduced binding observed) and – (no binding observed). The capture antibody was regenerated after each cycle of ligand capture and analyte binding and the murine construct was re-injected at the end of the study to confirm the activity of the capture surface. The results are summarized in Table 3.

humanized VL variant	binding to CD3ε
murine_CH2527-VL	+
CH2527-VL7_46-1	-
CH2527-VL7_46-2	-
CH2527-VL7_46-3	-
CH2527-VL7_46-4	-
CH2527-VL7_46-5	-
CH2527-VL7_46-6	-
CH2527-VL7_46-7	-
CH2527-VL7_46-8	-
CH2527-VL7_46-9	-
CH2527-VL7_46-10	-
CH2527-VL7_46-12	+/-
CH2527-VL7_46-13	+
CH2527-VL7_46-14	-

Table 3 Qualitative binding assessment based on SPR for the humanized light chain variants combined with the murine heavy chain of CH2527. Only the humanized light chain variant that was finally chosen, CH2527-VL7_46-13, highlighted in bold letters, exhibited comparable binding to human and cynomolgus CD3ε.

Example 4

Properties of humanized common light chain with CD3 ϵ specificity

The light chain V-domain variant that was chosen for the humanized lead molecule is VL7_46-13. The degree of humanness, i.e. the sequence homology of the humanized V-domain to the human germline V-domain sequence was determined. For VL7_46-13, the overall sequence identity with the closest human germline homolog is 65% before humanization and 80% afterwards. Omitting the CDR regions, the sequence identity is 92% to the closest human germline homolog. As can be seen from Table 3, VL7_46-13 is the only humanized VL variant out of a panel of 13 variants that showed comparable binding to the parental murine antibody and also retained its cross-reactivity to cynomolgus CD3 ϵ . This result indicates that it was not trivial to humanize the murine VL-domain without losing binding affinity to CD3 ϵ which required several back-mutations to murine framework residues (in particular G46) while retaining G24 in CDR1. In addition, this result shows that the VL-domain plays a crucial role in target recognition. Importantly, the humanized VL-domain VL7_46-13 based on an infrequent human germline belonging to the V-domain family 7 of the lambda type and retaining affinity and specificity for CD3 ϵ , is also suitable to be used as a common light chain in phage-displayed antibody libraries of the Fab-format and enables successful selection for novel specificities which greatly facilitates the generation and production of bispecific molecules binding to CD3 ϵ and e.g. a tumor target and sharing the same 'common' light chain.

Example 5

Generation of a phage displayed antibody library using a human germ-line Common Light Chain derived from HVK1-39

Several approaches to generate bispecific antibodies that resemble full length human IgG utilize modifications in the Fc region that induce heterodimerization of two distinct heavy chains. Such examples include knobs-into-holes (Merchant et al., Nat Biotechnol. 1998 Jul;16(7):677-81) SEED (Davis et al., Protein Eng Des Sel. 2010 Apr;23(4):195-202) and electrostatic steering technologies (Gunasekaran et al., J Biol Chem. 2010 Jun 18;285(25):19637-46). Although these approaches enable effective heterodimerization of two distinct heavy chains, appropriate pairing of cognate light and heavy chains remains a

problem. Usage of a common light chain (LC) can solve this issue (Merchant, et al. Nat Biotech 16, 677–681 (1998)).

Here, we describe the generation of an antibody library for the display on a M13 phage. Essentially, we designed a multi framework library for the heavy chain with one constant
5 (or “common”) light chain. This library is designed for generating multispecific antibodies without the need to use sophisticated technologies to avoid light chain mispairing.

By using a common light chain the production of these molecules can be facilitated as no mispairing occurs any longer and the isolation of a highly pure bispecific antibody is
10 facilitated. As compared to other formats the use of Fab fragments as building blocks as opposed to e.g. the use of scFv fragments results in higher thermal stability and the lack of scFv aggregation and intermolecular scFv formation.

Library generation

15 In the following the generation of an antibody library for the display on M13 phage is described. Essentially, we designed a multi framework library for the heavy chain with one constant (or “common”) light chain.

We used these heavy chains in the library (GenBank Accession Numbers in brackets):

IGHV1-46*01 (X92343) (SEQ ID NO:104) ,
20 IGHV1-69*06 (L22583), (SEQ ID NO:105)
IGHV3-15*01 (X92216), (SEQ ID NO:106)
IGHV3-23*01 (M99660), (SEQ ID NO:107)
IGHV4-59*01 (AB019438), (SEQ ID NO:108)
IGHV5-51*01 (M99686), (SEQ ID NO:109)

25 All heavy chains use the IGHJ2 as J-element, except the IGHV1-69*06 which uses IGHJ6 sequence. The design of the randomization included the CDR-H1, CDR-H2, and CDR-H3. For CDR-H1 and CDR-H2 a “soft” randomization strategy was chosen, and the randomization oligonucleotides were such that the codon for the amino acid of the germ-line sequence was present at 50%. All other amino acids, except cysteine, were summing
30 up for the remaining 50%. In CDR-H3, where no germ-line amino acid is present due to the presence of the genetic D-element, oligonucleotides were designed that allow for the usage of randomized inserts between the V-element and the J-element of 4 to 9 amino acids in length. Those oligonucleotides contained in their randomized part e.g. The three

amino acids G/Y/S are present to 15% each, those amino acids

A/D/T/R/P/L/V/N/W/F/I/E are present to 4,6% each.

Exemplary methods for generation of antibody libraries are described in Hoogenboom et al., Nucleic Acids Res. 1991, 19, 4133–413; Lee et., al J. Mol. Biol. (2004) 340, 1073–

5 1093.

The light chain is derived from the human sequence hVK1-39, and is used in an unmodified and non-randomized fashion. This will ensure that the same light chain can be used for other projects without additional modifications.

Exemplary Library selection:

10 Selections with all affinity maturation libraries are carried out in solution according to the following procedure using a monomeric and biotinylated extracellular domain of a target antigen X.

1. 10^{12} phagemid particles of each library are bound to 100nM biotinylated soluble antigen for 0.5 h in a total volume of 1ml. 2. Biotinylated antigen is captured and
15 specifically bound phage particles are isolated by addition of $\sim 5 \times 10^7$ streptavidin-coated magnetic beads for 10 min. 3. Beads are washed using 5-10x 1ml PBS/Tween20 and 5-10x 1ml PBS. 4. Elution of phage particles is done by addition of 1ml 100mM TEA (triethylamine) for 10 min and neutralization by addition of 500ul 1M Tris/HCl pH 7.4 and 5. Re-infection of exponentially growing E. coli TG1 bacteria, infection with helper
20 phage VCSM13 and subsequent PEG/NaCl precipitation of phagemid particles is applied in subsequent selection rounds. Selections are carried out over 3-5 rounds using either constant or decreasing (from 10^{-7} M to 2×10^{-9} M) antigen concentrations. In round 2, capture of antigen/phage complexes is performed using neutravidin plates instead of streptavidin beads. All binding reactions are supplemented either with 100 nM bovine
25 serum albumin, or with non-fat milk powder in order to compete for unwanted clones arising from mere sticky binding of the antibodies to the plastic support.

Selections are being carried out over three or four rounds using decreasing antigen concentrations of the antigen starting from 100nM and going down to 5nM in the final selection round. Specific binders are defined as signals ca. $5 \times$ higher than background
30 and are identified by ELISA. Specific binders are identified by ELISA as follows: 100 μ l of 10nM biotinylated antigen per well are coated on neutravidin plates. Fab-containing bacterial supernatants are added and binding Fabs are detected via their Flag-tags by using an anti-Flag/HRP secondary antibody. ELISA-positive clones are bacterially

expressed as soluble Fab fragments in 96-well format and supernatants are subjected to a kinetic screening experiment by SPR-analysis using ProteOn XPR36 (BioRad). Clones expressing Fabs with the highest affinity constants are identified and the corresponding phagemids are sequenced. For further characterization, the Fab sequences are amplified
5 via PCR from the phagemid and cloned via appropriate restriction sites into human IgG1 expression vectors for mammalian production.

Generation of a phage displayed antibody library using a humanized CD3ε specific Common Light Chain

Here, the generation of an antibody library for the display on M13 phage is described.

10 Essentially, we designed a multi framework library for the heavy chain with one constant (or “common”) light chain. This library was designed for the generation of Fc-containing, but FcγR binding inactive T cell bispecific antibodies of IgG1 P329G LALA or IgG4 SPLE PG isotype in which one or two Fab recognize a tumor surface antigen expressed on a tumor cell whereas the remaining Fab arm of the antibody recognizes CD3ε on a T
15 cell.

Library generation

In the following the generation of an antibody library for the display on M13 phage is described. Essentially, we designed a multi framework library for the heavy chain with one constant (or “common”) light chain. This library is designed solely for the generation
20 of Fc-containing, but FcγR binding inactive T cell bispecific antibodies of IgG1 P329G LALA or IgG4 SPLE PG isotype.

Diversity was introduced via randomization oligonucleotides only in the CDR3 of the different heavy chains. Methods for generation of antibody libraries are well known in the art and are described in (Hoogenboom et al., Nucleic Acids Res. 1991, 19, 4133–413; or
25 in: Lee et., al J. Mol. Biol. (2004) 340, 1073–1093).

We used these heavy chains in the library:

IGHV1-46*01 (X92343), (SEQ ID NO:104)
IGHV1-69*06 (L22583), (SEQ ID NO:105)
IGHV3-15*01 (X92216), (SEQ ID NO:106)
30 IGHV3-23*01 (M99660), (SEQ ID NO:107)
IGHV4-59*01 (AB019438), (SEQ ID NO:108)
IGHV5-51*01 (M99686), (SEQ ID NO:109)

We used the light chain derived from the humanized human and Cynomolgus CD3 ϵ specific antibody CH2527 in the library: (VL7_46-13; SEQ ID NO:112). This light chain was not randomized and used without any further modifications in order to ensure compatibility with different bispecific binders.

- 5 All heavy chains use the IGHJ2 as J-element, except the IGHV1-69*06 which uses IGHJ6 sequence. The design of the randomization focused on the CDR-H3 only, and PCR oligonucleotides were designed that allow for the usage of randomized inserts between the V-element and the J-element of 4 to 9 amino acids in length.

10

Example 6

Selection of antibody fragments from common light chain libraries (comprising light chain with CD3 ϵ specificity) to FolR1

The antibodies 16A3, 15A1, 18D3, 19E5, 19A4, 15H7, 15B6, 16D5, 15E12, 21D1,
15 16F12, 21A5, 21G8, 19H3, 20G6, and 20H7 comprising the common light chain VL7_46-13 with CD3 ϵ specificity were obtained by phage display selections against different species (human, cynomolgus and murine) of FolR1. Clones 16A3, 15A1, 18D3, 19E5, 19A4, 15H7, 15B6, 21D1, 16F12, 19H3, 20G6, and 20H7 were selected from a sub-library in which the common light chain was paired with a heavy chain repertoire
20 based on the human germline VH1_46. In this sub-library, CDR3 of VH1_46 has been randomized based on 6 different CDR3 lengths. Clones 16D5, 15E12, 21A5, and 21G8 were selected from a sub-library in which the common light chain was paired with a heavy chain repertoire based on the human germline VH3_15. In this sub-library, CDR3 of VH3_15 has been randomized based on 6 different CDR3 lengths. In order to obtain
25 species cross-reactive (or murine FolR1-reactive) antibodies, the different species of FolR1 were alternated (or kept constant) in different ways over 3 rounds of biopanning: 16A3 and 15A1 (human – cynomolgus – human FolR1); 18D3 (cynomolgus – human – murine FolR1); 19E5 and 19A4 (3 rounds against murine FolR1); 15H7, 15B6, 16D5, 15E12, 21D1, 16F12, 21A5, 21G8 (human – cynomolgus – human FolR1); 19H3, 20G6,
30 and 20H7 (3 rounds against murine FolR1).

Human, murine and cynomolgus FolR1 as antigens for the phage display selections as well as ELISA- and SPR-based screenings were transiently expressed as N-terminal

monomeric Fc-fusion in HEK EBNA cells and *in vivo* site-specifically biotinylated via co-expression of BirA biotin ligase at the avi-tag recognition sequence located at the C-terminus of the Fc portion carrying the receptor chain (Fc knob chain). In order to assess the specificity to FolR1, two related receptors, human FolR2 and FolR3 were generated in
5 the same way.

Selection rounds (biopanning) were performed in solution according to the following pattern:

1. Pre-clearing of $\sim 10^{12}$ phagemid particles on maxisorp plates coated with 10 ug/ml of an unrelated human IgG to deplete the libraries of antibodies recognizing the Fc-portion
10 of the antigen.
2. Incubating the non-Fc-binding phagemid particles with 100nM biotinylated human, cynomolgus, or murine FolR1 for 0.5h in the presence of 100nM unrelated non-biotinylated Fc knob-into-hole construct for further depletion of Fc-binders in a total volume of 1ml.
- 15 3. Capturing the biotinylated FolR1 and attached specifically binding phage by transfer to 4 wells of a neutravidin pre-coated microtiter plate for 10 min (in rounds 1 & 3).
4. Washing the respective wells using 5x PBS/Tween20 and 5x PBS.
5. Eluting the phage particles by addition of 250 ul 100 mM TEA (triethylamine) per well for 10 min and neutralization by addition of 500 ul 1 M Tris/HCl pH 7.4 to the
20 pooled eluates from 4 wells.
6. Post-clearing of neutralized eluates by incubation on neutravidin pre-coated microtiter plate with 100 nM biotin-captured FolR2 or FolR3 for final removal of Fc- and unspecific binders.
7. Re-infection of log-phase *E. coli* TG1 cells with the supernatant of eluted phage
25 particles, infection with helperphage VCSM13, incubation on a shaker at 30 °C over night and subsequent PEG/NaCl precipitation of phagemid particles to be used in the next selection round.

Selections were carried out over 3 rounds using constant antigen concentrations of 100nM. In round 2, in order to avoid enrichment of binders to neutravidin, capture of antigen : phage complexes was performed by addition of 5.4×10^7 streptavidin-coated magnetic beads. Specific binders were identified by ELISA as follows: 100ul of 25 nM biotinylated human, cynomolgus, or murine FolR1 and 10 ug/ml of human IgG were coated on neutravidin plates and maxisorp plates, respectively. Fab-containing bacterial supernatants were added and binding Fabs were detected via their Flag-tags using an anti-Flag/HRP secondary antibody. Clones exhibiting signals on human FolR1 and being negative on human IgG were short-listed for further analyses and were also tested in a similar fashion against the remaining two species of FolR1. They were bacterially expressed in a 0.5 liter culture volume, affinity purified and further characterized by SPR-analysis using BioRad's ProteOn XPR36 biosensor.

Affinities (K_D) of selected clones were measured by surface plasmon resonance (SPR) using a ProteOn XPR36 instrument (Biorad) at 25 °C with biotinylated human, cynomolgus, and murine FolR1 as well as human FolR2 and FolR3 (negative controls) immobilized on NLC chips by neutravidin capture. Immobilization of antigens (ligand): Recombinant antigens were diluted with PBST (10 mM phosphate, 150 mM sodium chloride pH 7.4, 0.005% Tween 20) to 10 µg/ml, then injected at 30 µl/minute in vertical orientation. Injection of analytes: For 'one-shot kinetics' measurements, injection direction was changed to horizontal orientation, two-fold dilution series of purified Fab (varying concentration ranges) were injected simultaneously along separate channels 1-5, with association times of 200 s, and dissociation times of 600 s. Buffer (PBST) was injected along the sixth channel to provide an "in-line" blank for referencing. Association rate constants (k_{on}) and dissociation rate constants (k_{off}) were calculated using a simple one-to-one Langmuir binding model in ProteOn Manager v3.1 software by simultaneously fitting the association and dissociation sensorgrams. The equilibrium dissociation constant (K_D) was calculated as the ratio k_{off}/k_{on} . Table 4 lists the equilibrium dissociation constants (K_D) of the selected clones specific for FolR1.

Table 4: Equilibrium dissociation constants (K_D) for anti-FolR1 antibodies (Fab-format) selected by phage display from common light chain sub-libraries comprising VL7_46-13, a humanized light chain specific for CD3ε. K_D in nM.

Clone	huFolR1 [nM]	cyFolR1 [nM]	muFolR1 [nM]	huFolR2 [nM]	huFolR3 [nM]
16A3	21.7	18	very weak	no binding	no binding
15A1	30.9	17.3	very weak	no binding	no binding
18D3	93.6	40.2	very weak	no binding	no binding
19E5	522	276	19.4	no binding	no binding
19A4	2050	4250	43.1	no binding	no binding
15H7	13.4	72.5	no binding	no binding	no binding
15B6	19.1	13.9	no binding	no binding	no binding
16D5	39.5	114	no binding	no binding	no binding
15E12	55.7	137	no binding	no binding	no binding
21D1	62.6	32.1	no binding	no binding	no binding
16F12	68	90.9	no binding	no binding	no binding
21A5	68.8	131	no binding	no binding	no binding
21G8	130	261	no binding	no binding	no binding
19H3	no binding	no binding	89.7	no binding	no binding
20G6	no binding	no binding	78.5	no binding	no binding

Example 7

Selection of antibody fragments from generic multi-framework libraries to FolR1

5

The antibodies 11F8, 36F2, 9D11, 5D9, 6B6, and 14E4 were obtained by phage display selections based on generic multi-framework sub-libraries against different species (human, cynomolgus and murine) of FolR1. In these multi-framework sub-libraries, different VL-domains with randomized CDR3 (3 different lengths) are paired with different VH-domains with randomized CDR3 (6 different lengths). The selected clones are of the following VL/VH pairings: 11F8 (Vk_{1_5}/VH_{1_69}), 36F2 (Vk_{3_20}/VH_{1_46}), 9D11 (Vk_{2D_28}/VH_{1_46}), 5D9 (Vk_{3_20}/VH_{1_46}), 6B6 (Vk_{3_20}/VH_{1_46}), and 14E4 (Vk_{3_20}/VH_{3_23}). In order to obtain species cross-reactive (or murine FolR1-reactive) antibodies, the different species of FolR1 were alternated (or kept constant) in different ways over 3 or 4 rounds of biopanning: 11F8 (cynomolgus – murine – human FolR1); 36F2 (human – murine – cynomolgus – murine FolR1); 9D11 (cynomolgus – human – cynomolgus FolR1); 5D9 (human – cynomolgus – human FolR1); 6B6 (human – cynomolgus – human FolR1) and 14E4 (3 rounds against murine FolR1).

20

Human, murine and cynomolgus FolR1 as antigens for the phage display selections as well as ELISA- and SPR-based screenings were transiently expressed as N-terminal

monomeric Fc-fusion in HEK EBNA cells and *in vivo* site-specifically biotinylated via co-expression of BirA biotin ligase at the avi-tag recognition sequence located at the C-terminus of the Fc portion carrying the receptor chain (Fc knob chain). In order to assess the specificity to FolR1, two related receptors, human FolR2 and FolR3 were generated in
5 the same way.

Selection rounds (biopanning) were performed in solution according to the following pattern:

1. Pre-clearing of $\sim 10^{12}$ phagemid particles on maxisorp plates coated with 10 ug/ml of an unrelated human IgG to deplete the libraries of antibodies recognizing the Fc-portion
10 of the antigen.
2. Incubating the non-Fc-binding phagemid particles with 100nM biotinylated human, cynomolgus, or murine FolR1 for 0.5h in the presence of 100nM unrelated non-biotinylated Fc knob-into-hole construct for further depletion of Fc-binders in a total volume of 1ml.
- 15 3. Capturing the biotinylated FolR1 and attached specifically binding phage by transfer to 4 wells of a neutravidin pre-coated microtiter plate for 10 min (in rounds 1 & 3).
4. Washing the respective wells using 5x PBS/Tween20 and 5x PBS.
5. Eluting the phage particles by addition of 250 ul 100 mM TEA (triethylamine) per well for 10 min and neutralization by addition of 500 ul 1 M Tris/HCl pH 7.4 to the pooled
20 eluates from 4 wells.
6. Post-clearing of neutralized eluates by incubation on neutravidin pre-coated microtiter plate with 100 nM biotin-captured FolR2 or FolR3 for final removal of Fc- and unspecific binders.
7. Re-infection of log-phase *E. coli* TG1 cells with the supernatant of eluted phage
25 particles, infection with helperphage VCSM13, incubation on a shaker at 30 °C over night and subsequent PEG/NaCl precipitation of phagemid particles to be used in the next selection round.

Selections were carried out over 3 rounds using constant antigen concentrations of 100nM. In round 2 and 4, in order to avoid enrichment of binders to neutravidin, capture of antigen : phage complexes was performed by addition of 5.4×10^7 streptavidin-coated magnetic beads. Specific binders were identified by ELISA as follows: 100ul of 25 nM
 5 biotinylated human, cynomolgus, or murine FolR1 and 10 ug/ml of human IgG were coated on neutravidin plates and maxisorp plates, respectively. Fab-containing bacterial supernatants were added and binding Fabs were detected via their Flag-tags using an anti-Flag/HRP secondary antibody. Clones exhibiting signals on human FolR1 and being negative on human IgG were short-listed for further analyses and were also tested in a
 10 similar fashion against the remaining two species of FolR1. They were bacterially expressed in a 0.5 liter culture volume, affinity purified and further characterized by SPR-analysis using BioRad's ProteOn XPR36 biosensor.

Affinities (K_D) of selected clones were measured by surface plasmon resonance (SPR) using a ProteOn XPR36 instrument (Biorad) at 25 °C with biotinylated human,
 15 cynomolgus, and murine FolR1 as well as human FolR2 and FolR3 (negative controls) immobilized on NLC chips by neutravidin capture. Immobilization of antigens (ligand): Recombinant antigens were diluted with PBST (10 mM phosphate, 150 mM sodium chloride pH 7.4, 0.005% Tween 20) to 10 µg/ml, then injected at 30 µl/minute in vertical orientation. Injection of analytes: For 'one-shot kinetics' measurements, injection
 20 direction was changed to horizontal orientation, two-fold dilution series of purified Fab (varying concentration ranges) were injected simultaneously along separate channels 1-5, with association times of 150 or 200 s, and dissociation times of 200 or 600 s, respectively. Buffer (PBST) was injected along the sixth channel to provide an "in-line" blank for referencing. Association rate constants (k_{on}) and dissociation rate constants (k_{off})
 25 were calculated using a simple one-to-one Langmuir binding model in ProteOn Manager v3.1 software by simultaneously fitting the association and dissociation sensorgrams. The equilibrium dissociation constant (K_D) was calculated as the ratio k_{off}/k_{on} . Table 5 lists the equilibrium dissociation constants (K_D) of the selected clones specific for FolR1.

Table 5: Equilibrium dissociation constants (K_D) for anti-FolR1 antibodies (Fab-format)
 30 selected by phage display from generic multi-framework sub-libraries. K_D in nM.

Clone	K _D (nM)				
	huFolR1	cyFolR1	muFolR1	huFolR2	huFolR3
11F8	632	794	1200	no binding	no binding
36F2	1810	1640	737	no binding	no binding
9D11	8.64	5.29	no binding	no binding	no binding
5D9	8.6	5.9	no binding	no binding	no binding
6B6	14.5	9.4	no binding	no binding	no binding
14E4	no binding	no binding	6.09	no binding	no binding

Example 8

Production and purification of novel FolR1 binders in IgG and T-cell bispecific formats

5 To identify FolR1 binders which are able to induce T-cell dependent killing of selected target cells the antibodies isolated from a common light chain- or Fab-library were converted into the corresponding human IgG1 format. In brief, the variable heavy and variable light chains of unique FolR1 binders from phage display were amplified by standard PCR reactions using the Fab clones as the template. The PCR products were

10 purified and inserted (either by restriction endonuclease and ligase based cloning, or by 'recombineering' using the InFusion kit from Invitrogen) into suitable expression vectors in which they are fused to the appropriate human constant heavy or human constant light chain. The expression cassettes in these vectors consist of a chimeric MPSV promoter and a synthetic polyadenylation site. In addition, the plasmids contain the oriP region from the

15 Epstein Barr virus for the stable maintenance of the plasmids in HEK293 cells harboring the EBV nuclear antigen (EBNA). After PEI mediated transfection the antibodies were transiently produced in HEK293 EBNA cells and purified by standard ProteinA affinity chromatography followed by size exclusion chromatography as described:

Transient transfection and production

20 All (bispecific) antibodies (if not obtained from a commercial source) used herein were transiently produced in HEK293 EBNA cells using a PEI mediated transfection procedure for the required vectors as described below. HEK293 EBNA cells are cultivated in suspension serum free in CD CHO culture medium. For the production in 500 ml shake flask 400 million HEK293 EBNA cells are seeded 24 hours before transfection (for

25 alternative scales all amounts were adjusted accordingly). For transfection cells are centrifuged for 5 min by 210 x g, supernatant is replaced by pre-warmed 20 ml CD CHO medium. Expression vectors are mixed in 20 ml CD CHO medium to a final amount of 200 µg DNA. After addition of 540 µl PEI solution is vortexed for 15 s and subsequently

incubated for 10 min at room temperature. Afterwards cells are mixed with the DNA/PEI solution, transferred to a 500 ml shake flask and incubated for 3 hours by 37 °C in an incubator with a 5 % CO₂ atmosphere. After incubation time 160 ml F17 medium is added and cell are cultivated for 24 hours. One day after transfection 1 mM valproic acid and 7 % Feed 1 is added. After 7 days cultivation supernatant is collected for purification by centrifugation for 15 min at 210 x g, the solution is sterile filtered (0.22 µm filter) and sodium azide in a final concentration of 0.01 % w/v is added, and kept at 4°C. After production the supernatants were harvested and the antibody containing supernatants were filtered through 0.22 µm sterile filters and stored at 4 °C until purification.

10

Antibody purification

All molecules were purified in two steps using standard procedures, such as protein A affinity purification (Äkta Explorer) and size exclusion chromatography. The supernatant obtained from transient production was adjusted to pH 8.0 (using 2 M TRIS pH 8.0) and applied to HiTrap PA FF (GE Healthcare, column volume (cv) = 5 ml) equilibrated with 8 column volumes (cv) *buffer A* (20 mM sodium phosphate, 20 mM sodium citrate, pH 7.5). After washing with 10 cv of *buffer A*, the protein was eluted using a pH gradient to *buffer B* (20 mM sodium citrate pH 3, 100 mM NaCl, 100 mM glycine) over 12 cv. Fractions containing the protein of interest were pooled and the pH of the solution was gently adjusted to pH 6.0 (using 0.5 M Na₂HPO₄ pH 8.0). Samples were concentrated to 2 ml using ultra-concentrators (Vivaspin 15R 30.000 MWCO HY, Sartorius) and subsequently applied to a HiLoad™ 16/60 Superdex™ 200 preparative grade (GE Healthcare) equilibrated with 20 mM Histidine, pH 6.0, 140 mM NaCl, 0.01% Tween-20. The aggregate content of eluted fractions was analyzed by analytical size exclusion chromatography. Therefore, 30 µl of each fraction was applied to a TSKgel G3000 SW XL analytical size-exclusion column (Tosoh) equilibrated in 25 mM K₂HPO₄, 125 mM NaCl, 200 mM L-arginine monohydrochloride, 0.02 % (w/v) NaN₃, pH 6.7 running buffer at 25°C. Fractions containing less than 2 % oligomers were pooled and concentrated to final concentration of 1 - 1.5 mg/ml using ultra concentrators (Vivaspin 15R 30.000 MWCO HY, Sartorius). The protein concentration was determined by measuring the optical density (OD) at 280 nm, using the molar extinction coefficient calculated on the basis of the amino acid sequence. Purity and molecular weight of the constructs were analyzed by SDS capillary electrophoresis in the presence and absence of a reducing

30

agent following the manufacturer instructions (instrument Caliper LabChipGX, Perkin Elmer). Purified proteins were frozen in liquid N₂ and stored at -80 °C.

Based on *in vitro* characterization results selected binders were converted into a T-cell bispecific format. In these molecules the FolR1:CD3 binding moieties are arranged in a 2:1 order with the FolR1 Fabs being located at the N-terminus. For clones isolated from the standard Fab library the CD3 binding part was generated as a CrossFab (CH1Cκ crossing) while for the clones from the common light chain library no crossing was necessary. These bispecific molecules were produced and purified analogously to the IgGs.

- 10 Table 6: Yield and monomer content of novel FolR1 binders in IgG and TCB format, respectively.

#	Clone	Library	IgG		TCB	
			Yield [mg/L]	Monomer [%]	Yield [mg/L]	Monomer [%]
1	11F8	Fab	8.03	96.26	--	--
2	14E4	Fab	8.90	98.12	--	--
3	15B6	CLC	7.72	100.00	--	--
4	15E12	CLC	6.19	100.00	--	--
5	15H7	CLC	8.94	100.00	--	--
6	16A3	CLC	0.60	n.d.	--	--
7	16D5	CLC	36.50	96.96	4.36	97.19
8	16F12	CLC	5.73	97.17	--	--
9	18D3	CLC	0.90	n.d.	--	--
10	19A4	CLC	38.32	100.00	37.50	100.00
11	19E5	CLC	46.09	100.00	--	--
12	19H3	CLC	7.64	100.00	--	--
13	20G6	CLC	24.00	100.00	--	--
14	20H7	CLC	45.39	100.00	--	--
15	21A5	CLC	1.38	98.56	47.31	95.08
16	21D1	CLC	5.47	100.00	--	--
17	21G8	CLC	6.14	97.28	9.27	100.00
18	36F2	Fab	11.22	100.00	18.00	100.00

19	5D9	Fab	20.50	100.00	0.93	97.32
20	6B6	Fab	3.83	100.00	4.17	91.53
21	9D11	Fab	14.61	100.00	2.63	100.00

CLC: Common light chain

Example 9

2+1 and 1+1 T-cell bispecific formats

- 5 Four different T-cell bispecific formats were prepared for one common light chain binder (16D5) and three formats for one binder from the Fab library (9D11) to compare their killing properties in vitro.

The standard format is the 2+1 inverted format as already described (FolR1:CD3 binding moieties arranged in a 2:1 order with the FolR1 Fabs located at the N-terminus). In the
 10 2+1 classical format the FolR1:CD3 binding moieties are arranged in a 2:1 order with the CD3 Fab being located at the N-terminus. Two monovalent formats were also prepared. The 1+1 head-to-tail has the FolR1:CD3 binding moieties arranged in a 1:1 order on the same arm of the molecule with the FolR1 Fab located at the N-terminus. In the 1+1
 15 classical format the FolR1:CD3 binding moieties are present once, each on one arm of the molecule. For the 9D11 clone isolated from the standard Fab library the CD3 binding part was generated as a CrossFab (CH1Cκ crossing) while for the 16D5 from the common light chain library no crossing was necessary. These bispecific molecules were produced and purified analogously to the standard inverted T-cell bispecific format.

- 20 Table 7: Summary of the yield and final monomer content of the different T-cell bispecific formats.

Construct	Monomer [%] (SEC)	Yield
16D5 FolR1 TCB 2+1 (inverted)	96%	5.4 mg/L
16D5 FolR1 TCB 2+1 (classical)	90%	4.6 mg/L
16D5 FolR1 TCB 1+1 (head-to-	100 %	5.4 mg/L

tail)		
16D5 FolR1 TCB 1+1 (classical)	100%	0.7 mg/L
9D11 FolR1 TCB 2+1 (inverted)	100%	2.6 mg/L
9D11 FolR1 TCB 1+1 (head-to-tail)	100%	6.1 mg/L
9D11 FolR1 TCB 1+1 (classical)	96%	1.3 mg/L
Mov19 FolR1 TCB 2+1 (inverted)	98%	3 mg/L
Mov19 FolR1 TCB 1+1 (head-to-tail)	100%	5.2 mg/L

Example 10

Biochemical characterization of FolR1 binders by surface plasmon resonance

Binding of FolR1 binders as IgG or in the T-cell bispecific format to different recombinant folate receptors (human FolR1, 2 and 3, murine FolR1 and cynomolgus FolR1; all as Fc fusions) was assessed by surface plasmon resonance (SPR). All SPR experiments were performed on a Biacore T200 at 25 °C with HBS-EP as running buffer (0.01 M HEPES pH 7.4, 0.15 M NaCl, 3 mM EDTA, 0.005% Surfactant P20, Biacore, Freiburg/Germany).

10 Single injections

First the anti-FolR1 IgGs were analyzed by single injections (Table 1) to characterize their crossreactivity (to human, murine and cyno FolR1) and specificity (to human FolR1, human FolR2, human FolR3). Recombinant biotinylated monomeric Fc fusions of human, cynomolgus and murine Folate Receptor 1 (FolR1-Fc) or human Folate Receptor 2 and 3 (FolR2-Fc, FolR3-Fc) were directly coupled on a SA chip using the standard coupling instruction (Biacore, Freiburg/Germany). The immobilization level was about 300-400 RU. The IgGs were injected for 60 seconds at a concentration of 500 nM. IgGs binding to huFolR2 and huFolR3 were rejected for lack of specificity. Most of the binders are only crossreactive between human and cyno FolR1, additional crossreactivity to murine FolR1 went most of the time hand in hand with loss of specificity.

Table 8: Crossreactivity and specificity of 25 new folate receptor 1 binders (as IgGs) as well as of two control IgGs (Mov19 and Farletuzumab). + means binding, - means no binding, +/- means weak binding.

Clone name	Binding to huFolR1	Binding to cyFolR1	Binding to muFolR1	Binding to huFolR2	Binding to huFolR3
------------	--------------------	--------------------	--------------------	--------------------	--------------------

Mov19	+	+	-	-	-
Farletuzumab	+	+	-	-	-
16A3	+	+	+/-	-	-
18D3	+	+	-	-	-
19E5	+	+	+	+	+
19A4	-	-	+	+	+
15H7	+	+	+	-	-
15B6	+	+	-	-	-
16D5	+	+	-	-	-
15E12	+	+	+/-	+	+
21D1	+	+	+/-	-	-
16F12	+	+	-	-	-
21A5	+	+	-	-	+/-
21G8	+	+	-	+	+
19H3	-	-	+	-	-
20G6	-	-	+	-	-
20H7	-	-	+	-	-
9D11	+	+	-	-	-
5D9	+	+	-	+	+
6B6	+	+	-	+	+
11F8	+	+	+	+	+
36F2	+	+	+	-	-
14E4	-	-	+	-	-

Avidity to Folate Receptor 1

The avidity of the interaction between the anti-FolR1 IgGs or T cell bispecifics and the recombinant folate receptors was determined as described below (Table 9).

Recombinant biotinylated monomeric Fc fusions of human, cynomolgus and murine Folate Receptor 1 (FolR1-Fc) were directly coupled on a SA chip using the standard coupling instruction (Biacore, Freiburg/Germany). The immobilization level was about 300-400 RU. The anti-FolR1 IgGs or T cell bispecifics were passed at a concentration range from 2.1 to 500 nM with a flow of 30 μ L/minutes through the flow cells over 180 seconds. The dissociation was monitored for 600 seconds. Bulk refractive index differences were corrected for by subtracting the response obtained on reference flow cell immobilized with recombinant biotinylated IL2 receptor Fc fusion. For the analysis of the interaction of 19H3 IgG and murine folate receptor 1, folate (Sigma F7876) was added in the HBS-EP running buffer at a concentration of 2.3 μ M. The binding curves resulting from the bivalent binding of the IgGs or T cell bispecifics were approximated to a 1:1 Langmuir binding and fitted with that model (which is not correct, but gives an idea of the

avidity). The apparent avidity constants for the interactions were derived from the rate constants of the fitting using the Bia Evaluation software (GE Healthcare).

Table 9: Bivalent binding (avidity with apparent KD) of selected FolR1 binders as IgGs or as T-cell bispecifics (TCB) on human and cyno FolR1.

Analyte	Ligand	ka (1/Ms)	kd (1/s)	Apparent KD (M)
16D5 TCB	huFolR1	8.31E+04	3.53E-04	4.24E-09
	cyFolR1	1.07E+05	3.70E-04	3.45E-09
9D11 TCB	huFolR1	1.83E+05	9.83E-05	5.36E-10
	cyFolR1	2.90E+05	6.80E-05	2.35E-10
21A5 TCB	huFolR1	2.43E+05	2.64E-04	1.09E-09
	cyFolR1	2.96E+05	2.76E-04	9.32E-10
36F2 IgG	huFolR1	2.62E+06	1.51E-02	5.74E-9
	cyFolR1	3.02E+06	1.60E-02	5.31E-9
	muFolR1	3.7E+05	6.03E-04	1.63E-9
Mov19 IgG	huFolR1	8.61E+05	1.21E-04	1.4E-10
	cyFolR1	1.29E+06	1.39E-04	1.08E-10
Farletuzumab	huFolR1	1.23E+06	9E-04	7.3E-10
	cyFolR1	1.33E+06	8.68E-04	6.5E-10
19H3 IgG	muFolR1	7.1E+05	1.1E-03	1.55E-09

5

1. Affinity to Folate Receptor 1

The affinity of the interaction between the anti-FolR1 IgGs or the T cell bispecifics and the recombinant folate receptors was determined as described below (Table 10).

For affinity measurement, direct coupling of around 6000-7000 resonance units (RU) of the anti-human Fab specific antibody (Fab capture kit, GE Healthcare) was performed on a CM5 chip at pH 5.0 using the standard amine coupling kit (GE Healthcare). Anti-FolR1 IgGs or T cell bispecifics were captured at 20 nM with a flow rate of 10 µl/min for 20 or 40 sec, the reference flow cell was left without capture. Dilution series (6.17 to 500 nM or 12.35 to 3000 nM) of human or cyno Folate Receptor 1 Fc fusion were passed on all flow cells at 30 µl/min for 120 or 240 sec to record the association phase. The dissociation phase was monitored for 240 s and triggered by switching from the sample solution to HBS-EP. The chip surface was regenerated after every cycle using a double injection of 60 sec 10 mM Glycine-HCl pH 2.1 or pH 1.5. Bulk refractive index differences were corrected for by subtracting the response obtained on the reference flow cell 1. The affinity constants for the interactions were derived from the rate constants by fitting to a 1:1 Langmuir binding using the Bia Evaluation software (GE Healthcare).

Table 10: Monovalent binding (affinity) of selected FolR1 binders as IgGs or as T-cell bispecifics (TCB) on human and cyno FolR1.

Ligand	Analyte	ka (1/Ms)	kd (1/s)	KD (M)
16D5 TCB	huFolR1	1.53E+04	6.88E-04	4.49E-08
	cyFolR1	1.32E+04	1.59E-03	1.21E-07
9D11 TCB	huFolR1	3.69E+04	3.00E-04	8.13E-09
	cyFolR1	3.54E+04	2.06E-04	5.82E-09
21A5 TCB	huFolR1	1.79E+04	1.1E-03	6.16E-08
	cyFolR1	1.48E+04	2.06E-03	1.4E-07
Mov19 IgG	huFolR1	2.89E+05	1.59E-04	5.5E-10
	cyFolR1	2.97E+05	1.93E-04	6.5E-10
Farletuzumab	huFolR1	4.17E+05	2.30E-02	5.53E-08
	cyFolR1	5.53E+05	3.73E-02	6.73E-08

2. Affinity to CD3

- 5 The affinity of the interaction between the anti-FolR1 T cell bispecifics and the recombinant human CD3 $\epsilon\delta$ -Fc was determined as described below (Table 11).

For affinity measurement, direct coupling of around 9000 resonance units (RU) of the anti-human Fab specific antibody (Fab capture kit, GE Healthcare) was performed on a CM5 chip at pH 5.0 using the standard amine coupling kit (GE Healthcare). Anti-FolR1 T

10 cell bispecifics were captured at 20 nM with a flow rate of 10 μ l/min for 40 sec, the reference flow cell was left without capture. Dilution series (6.17 to 500 nM) of human CD3 $\epsilon\delta$ -Fc fusion were passed on all flow cells at 30 μ l/min for 240 sec to record the association phase. The dissociation phase was monitored for 240 s and triggered by switching from the sample solution to HBS-EP. The chip surface was regenerated after

15 every cycle using a double injection of 60 sec 10 mM Glycine-HCl pH 2.1. Bulk refractive index differences were corrected for by subtracting the response obtained on the reference flow cell 1. The affinity constants for the interactions were derived from the rate constants by fitting to a 1:1 Langmuir binding using the Bia Evaluation software (GE Healthcare).

- 20 Table 11: Monovalent binding (affinity) of selected FolR1 T-cell bispecifics (TCB) on human CD3-Fc.

Ligand	Analyte	ka (1/Ms)	kd (1/s)	KD (M)
16D5 TCB	huCD3	4.25E+04	3.46E-03	8.14E-08

21A5 TCB	huCD3	3.72E+04	3.29E-03	8.8E-08
-----------------	--------------	----------	----------	---------

The CD3 binding part is identical for all constructs and the affinity is similar for the measured T cell bispecifics (KD range between 60 and 90 nM).

Example 11

Simultaneous binding T cell bispecifics on Folate Receptor 1 and CD3

- 5 Simultaneous binding of the anti-FolR1 T cell bispecifics on recombinant Folate Receptor 1 and recombinant human CD3 $\epsilon\delta$ -Fc was determined as described below.

Recombinant biotinylated monomeric Fc fusions of human, cynomolgus and murine Folate Receptor 1 (FolR1-Fc) were directly coupled on a SA chip using the standard coupling instruction (Biacore, Freiburg/Germany). The immobilization level was about
 10 300-400 RU. The anti-FolR1 T cell bispecifics were injected for 60 s at 500 nM with a flow of 30 μ L/minutes through the flow cells, followed by an injection of hu CD $\epsilon\delta$ -Fc for 60 s at 500 nM. Bulk refractive index differences were corrected for by subtracting the response obtained on reference flow cell immobilized with recombinant biotinylated IL2 receptor Fc fusion. The four T cell bispecifics tested (16D5 TCB, 21A5 TCB, 51C7 TCB
 15 and 45D2 TCB) were able to bind simultaneously to Folate Receptor 1 and human CD3 as expected.

Example 12

Epitope binning

For epitope binning, the anti-FolR1 IgGs or T cell bispecifics were directly immobilized
 20 on a CM5 chip at pH 5.0 using the standard amine coupling kit (GE Healthcare), with a final response around 700 RU. 500 nM huFolR1-Fc was then captured for 60 s, followed by 500 nM of the different binders for 30 s. The surface was regenerated with two injections of 10 mM glycine pH 2 for 30 s each. It is assessed if the different binders can bind to huFolR1 captured on immobilized binders (Table 12).

- 25 Table 12: Epitope characterization of selected FolR1 binders as IgGs or as T-cell bispecifics (TCB) on human FolR1. + means binding, - means no binding, +/- means weak binding

		Analytes in solution					
	On huFolR1	16D5 TCB	21A5 TCB	9D11 TCB	36F2 IgG	Mov19 IgG	Farletuzumab
Immobilized	16D5 TCB	-	-	-	+	+	+
	21A5 TCB	-	-	-	+	+	+
	9D11 TCB	No additional binding on FolR1 possible once captured on 9D11					
	36F2 IgG	Measure not possible, huFolR1 dissociates too rapidly					
	Mov19 IgG	+	+	+/-	-	-	-

Based on these results and additional data with simultaneous binding on immobilized huFolR1, the binders were separated in three groups. It is not clear if 9D11 has a separate epitope because it displaces all the other binders. 16D5 and 21A5 seem to be in the same group and Mov19, Farletuzumab (Coney et al., Cancer Res. 1991 Nov 15;51(22):6125-32; Kalli et al., Curr Opin Investig Drugs. 2007 Dec;8(12):1067-73) and 36F2 in another (Table 13). However, 36F2 binds to a different epitope than Mov 19 and Farletuzumab as it binds to human, cynomous and murine FolR1.

Table 13: Epitope grouping of selected FolR1 binders as IgGs or as T-cell bispecifics (TCB) on human FolR1

Epitope 1	Epitope 2	Epitope 3
16D5	9D11	Mov19
21A5		Farletuzumab
		36F2

Example 13

Selection of binders

FolR1 binders in the IgG formats were screened by surface plasmon resonance (SPR) and
5 by in vitro assay on cells to select the best candidates.

The anti-FolR1 IgGs were analyzed by SPR to characterize their crossreactivity (to
human, murine and cynomolgus FolR1) and specificity (to human FolR1, human FolR2,
human FolR3). Unspecific binding to human FolR2 and 3 was considered an exclusion
factor. Binding and specificity to human FolR1 was confirmed on cells. Some binders did
10 not bind on cells expressing FolR1 even though they recognized the recombinant human
FolR1 in SPR. Aggregation temperature was determined but was not an exclusion factor
because the selected binders were all stable. Selected binders were tested in a
polyreactivity ELISA to check for unspecific binding, which led to the exclusion of four
binders. This process resulted in an initial selection of three binders: 36F2 (Fab library),
15 9D11 (Fab library) and 16D5 (common light chain). 36F2 dissociated rapidly from
huFolR1 and was, therefore, initially not favored.

Example 14

Specific binding of newly generated FolR1 binders to human FolR1 positive 20 tumor cells

New FolR1 binders were generated via Phage Display using either a Fab library or a
common light chain library using the CD3 light chain. The identified binders were
converted into a human IgG1 format and binding to FolR1 high expressing HeLa cells
was addressed. As reference molecule the human FolR1 binder Mov19 was included.

25 Most of the binders tested in this assay showed intermediate to good binding to FolR1
with some clones binding equally well as Mov19 (see Figure 2). The clones 16A3, 18D3,
15H7, 15B6, 21D1, 14E4 and 16F12 were excluded because binding to FolR1 on cells
could not be confirmed by flow cytometry. In a next step the selected clones were tested
for specificity to human FolR1 by excluding binding to the closely related human FolR2.
30 HEK cells were transiently transfected with either human FolR1 or human FolR2 to
address specificity. The clones 36F2 and 9D11 derived from the Fab library and the
clones 16D5 and 21A5 derived from the CLC library bind specifically to human FolR1

and not to human FolR2 (see Figures 3A-B). All the other tested clones showed at least some binding to human FolR2 (see Figures 3A-B). Therefore these clones were excluded from further characterization. In parallel cross-reactivity of the FolR1 clones to cyno FolR1 was addressed by performing binding studies to HEK cells transiently transfected with cyno FolR1. All tested clones were able to bind cyno FolR1 and the four selected human FolR1 specific clones 36F2, 9D11, 16D5 and 21A5 bind comparably well human and cyno FolR1 (Figure 4). Subsequently three human FolR1 specific cyno cross-reactive binders were converted into TCB format and tested for induction of T cell killing and T cell activation. These clones were 9D11 from the Fab library and 16D5 and 21A5 from the CLC library. As reference molecule Mov19 FolR1 TCB was included in all studies. These FolR1 TCBs were then used to compare induction of internalization after binding to FolR1 on HeLa cells. All three tested clones are internalized upon binding to FolR1 comparable to internalization upon binding of Mov19 FolR1 TCB (Figure 5). 21A5 FolR1 TCB was discontinued due to signs of polyreactivity.

15

Example 15

T cell-mediated killing of FolR1-expressing tumor target cells induced by FolR1 TCB antibodies

The FolR1 TCBs were used to determine T cell mediated killing of tumor cells expressing FolR1. A panel of potential target cell lines was used to determine FolR1 binding sites by Qifikit analysis.

20

The used panel of tumor cells contains FolR1 high, intermediate and low expressing tumor cells and a FolR1 negative cell line.

Table 14: FolR1 binding sites on tumor cells

Cell line	Origin	FolR1 binding sites
Hela	Cervix adenocarcinoma	2'240'716
Skov3	Ovarian adenocarcinoma	91'510
OVCAR5	Ovarian adenocarcinoma	22'077
HT29	Colorectal adenocarcinoma	10'135
MKN45	Gastric adenocarcinoma	54

Binding of the three different FoLR1 TCBs (containing 9D11, 16D5 and Mov19 binders) to this panel of tumor cell lines was determined showing that the FoLR1 TCBs bind specifically to FoLR1 expressing tumor cells and not to a FoLR1 negative tumor cell line.

- 5 The amount of bound construct is proportional to the FoLR1 expression level and there is still good binding of the constructs to the FoLR1 low cell line HT-29 detectable. In addition there is no binding of the negative control DP47 TCB to any of the used cell lines (Figures 6A-E). DP47 TCB is an untargeted TCB and was prepared as described in WO2014/131712.
- 10 The intermediate expressing cell line SKOV3 and the low expressing cell line HT-29 were further on used to test T cell mediated killing and T cell activation using 16D5 TCB and 9D11 TCB; DP47 TCB was included as negative control. Both cell lines were killed in the presence of already very low levels of 16D5 TCB and 9D11 TCB and there was no difference in activity between both TCBs even though 9D11 TCB binds stronger to FoLR1
- 15 than 16D5 TCB. Overall killing of SKOV3 cells was higher compared to HT-29 which reflects the higher expression levels of FoLR1 on SKOV3 cells (Figures 7A-D). In line with this, a strong upregulation of the activation marker CD25 and CD69 on CD4⁺ T cells and CD8⁺ T cells was detected. Activation of T cells was very similar in the presence of SKOV3 cells and HT-29 cells. The negative control DP47 TCB does not induce any
- 20 killing at the used concentrations and there was no significant upregulation of CD25 and CD69 on T cells.

Table 15: EC50 values of tumor cell killing and T cell activation with SKOV3 cells

Construct	Killing 24 h (pM)	Killing 48 h (pM)	CD4+ CD69+ (%)	CD4+ CD25+ (%)	CD8+ CD69+ (%)	CD8+ CD25+ (%)
9D11 FoLR1 TCB	1.1	0.03	0.51	0.46	0.019	0.03
16D5 FoLR1	0.7	0.04	0.34	0.33	0.025	0.031

-181-

TCB						
-----	--	--	--	--	--	--

Table 16: EC50 values of tumor cell killing and T cell activation with HT-29 cells

Construct	Killing 24 h (pM)	Killing 48 h (pM)	CD4+ CD69+ (%)	CD4+ CD25+ (%)	CD8+ CD69+ (%)	CD8+ CD25+ (%)
9D11 FolR1 TCB	2.3	0.1	1.22	1.11	0.071	0.084
16D5 FolR1 TCB	2.8	0.1	0.69	0.62	0.021	0.028

Example 16

5

**Binding of FolR1 TCB antibodies to erythrocytes
and T cell activation in whole blood**

To prove that there is no spontaneous activation in the absence of FoLR1 expressing tumor cells we tested if there is binding of the FolR1 clones to erythrocytes which might potentially express FolR1. We could not observe any specific binding of 9D11 IgG, 16D5 IgG and Mov19 IgG to erythrocytes, as negative control DP47 IgG was included (Figure 8).

To exclude any further unspecific binding to blood cells or unspecific activation via FoLR1 TCB, 9D11 TCB, 16D5 TCB and Mov19 TCB were added into whole blood and upregulation of CD25 and CD69 on CD4⁺ T cells and CD8⁺ T cells was analyzed by flow cytometry. DP47 TCB was included as negative control. No activation of T cells with any of the tested constructs could be observed by analyzing upregulation of CD25 and CD69 on CD4⁺ T cells and CD8⁺ T cells (Figure 9).

Example 17**T-cell killing induced by 36F2 TCB and 16D5 TCB****in different monovalent and bivalent T-cell bispecific formats**

T-cell killing mediated by 36F2 TCB, 16D5 TCB, 16D5 TCB classical, 16D5 TCB 1+1
5 and 16D5 TCB HT antibodies of Hela, Skov-3 (medium FolR1, about 70000-90000
copies) and HT-29 (low FolR1, about 10000) human tumor cells was assessed. DP47
TCB antibody was included as negative control. Human PBMCs were used as effectors
and the killing was detected at 24 h of incubation with the bispecific antibody. Briefly,
target cells were harvested with Trypsin/EDTA, washed, and plated at density of 25 000
10 cells/well using flat-bottom 96-well plates. Cells were left to adhere overnight. Peripheral
blood mononuclear cells (PBMCs) were prepared by Histopaque density centrifugation of
enriched lymphocyte preparations (buffy coats) obtained from healthy human donors.
Fresh blood was diluted with sterile PBS and layered over Histopaque gradient (Sigma,
#H8889). After centrifugation (450 x g, 30 minutes, room temperature), the plasma above
15 the PBMC-containing interphase was discarded and PBMCs transferred in a new falcon
tube subsequently filled with 50 ml of PBS. The mixture was centrifuged (400 x g, 10
minutes, room temperature), the supernatant discarded and the PBMC pellet washed twice
with sterile PBS (centrifugation steps 350 x g, 10 minutes). The resulting PBMC
population was counted automatically (ViCell) and stored in RPMI1640 medium
20 containing 10% FCS and 1% L-alanyl-L-glutamine (Biochrom, K0302) at 37°C, 5% CO₂
in cell incubator until further use (no longer than 24 h). For the killing assay, the antibody
was added at the indicated concentrations (range of 0.01 pM – 100 nM in triplicates).
PBMCs were added to target cells at final E:T ratio of 10:1. Target cell killing was
assessed after 24 h of incubation at 37°C, 5% CO₂ by quantification of LDH released into
25 cell supernatants by apoptotic/necrotic cells (LDH detection kit, Roche Applied Science,
#11 644 793 001). Maximal lysis of the target cells (= 100%) was achieved by incubation
of target cells with 1% Triton X-100. Minimal lysis (= 0%) refers to target cells co-
incubated with effector cells without bispecific construct. The results show target-
specific killing of all three FolR1+ target cell lines induced by 36F2 TCB and 16D5 TCB
30 (Figure 10).

Example 18

Generation of anti-TIM3 antibodies

Immunization of mice NMRI mice were immunized genetically, using a plasmid expression vector coding for full-length human Tim-3 by intradermal application of 100
5 ug vector DNA (plasmid 15304_hTIM3-fl), followed by Electroporation (2 square pulses of 1000 V/cm, duration 0.1 ms, interval 0.125 s; followed by 4 square pulses of 287.5 V/cm, duration 10 ms, interval 0.125 s. Mice received either 6 consecutive immunizations at days 0, 14, 28, 42, 56, 70, and 84. Blood was taken at days 36, 78 and 92 and serum prepared, which was used for titer determination by ELISA (see below). Animals with
10 highest titers were selected for boosting at day 96, by intravenous injection of 50 ug of recombinant human Tim-3 human Fc chimera, and monoclonal antibodies were isolated by hybridoma technology, by fusion of splenocytes to myeloma cell line 3 days after boost.

Determination of serum titers (ELISA) Human recombinant Tim-3 human Fc chimera was
15 immobilized on a 96-well NUNC Maxisorp plate at 0.3 ug/ml, 100 ul/well, in PBS, followed by: blocking of the plate with 2% Crotein C in PBS, 200 ul/well; application of serial dilutions of antisera, in duplicates, in 0.5% Crotein C in PBS, 100 ul/well; detection with HRP-conjugated goat anti-mouse antibody (Jackson ImmunoResearch/Dianova 115-036-071; 1/16 000). For all steps, plates were incubated for 1 h at 37⁰ C. Between all
20 steps, plates were washed 3 times with 0.05% Tween 20 in PBS. Signal was developed by addition of BM Blue POD Substrate soluble (Roche), 100 ul/well; and stopped by addition of 1 M HCl, 100 ul/well. Absorbance was read out at 450 nm, against 690 nm as reference. Titer was defined as dilution of antisera resulting in half-maximal signal.

Example 19

25 Characterization anti-Tim3 antibodies

ELISA for Tim3 Nunc-Maxi Sorp Streptavidine plates (MicroCoat #11974998/MC1099) were coated by 25 µl/well with Tim3-ECD-His-Biotin (biotinylated with BirA Ligase) and incubated at RT for 1 h while shaking at 400 rpm rotation. After washing (3x90 µl/well with PBST-buffer) 25 µl aTim3 samples or diluted (1:2 steps) reference antibody
30 aTim3 F38-2E2 (Biolegend) was added and incubated 1h at RT. After washing (3x90 µl/well with PBST-buffer) 25µl/well sheep-anti-mouse-POD (GE NA9310V) was added in 1:9000 dilution and incubated at RT for 1 h while shaking at 400 rpm rotation. After washing (4x90 µl/well with PBST-buffer) 25 µl/well TMB substrate (Calbiochem, #CL07)

was added and incubated until OD 1.5 – 2.5. Then the reaction was stopped by addition of 25 µl/well 1N HCL-solution. Measurement took place at 370/492 nm. ELISA results are listed as EC50-values [ng/ml] in summary Table 17 below.

Cell ELISA for Tim3 Adherent CHO-K1 cell line stably transfected with plasmid

- 5 15312_hTIM3-fl_pUC_Neo coding for full-length human Tim3 and selection with G418 (Neomycin resistance marker on plasmid) were seeded at a concentration of 1.2x10E6 cells/ml into 384-well flat bottom plates and grown over night.

At the next day 25 µl/well Tim3 sample or aTim3 reference antibody F38-2E2 Azide free (Biolegend, 354004) was added and incubated for 2h at 4°C (to avoid internalization).

- 10 After washing (3x90µl/well PBST (BIOTEK Washer: Prog. 29, 1 x 90) cells were fixed by flicking out residual buffer and addition of 50µl/well 0,05% Glutaraldehyde: Dilution 1:500 of 25% Glutaraldehyde (Sigma Cat.No: G5882) in 1xPBS-buffer and incubated for 1h at RT. After washing (3x90µl/well PBST (BIOTEK Washer: Prog. 21, 3x90 GreinLysin) 25 µl/well secondary antibody was added for detection (Sheep-anti-mouse-
15 POD; Horseradish POD linked F(ab')₂ Fragment ; GE NA9310) followed by 2h incubation at RT while shaking at 400 rpm. After washing (3x90µl/well PBST (BIOTEK Washer: Prog. 21, 3x90 GreinLysin) 25 µl/well TMB substrate solution (Roche 11835033001) was added and incubated until OD 1.5 – 2.5. Then the reaction was stopped by addition of 25 µl/well 1N HCL-solution. Measurement took place at 370/492
20 nm. Cell ELISA results are listed as “EC50 CHO-Tim3”-values [ng/ml] in summary table Table 17 below.

Table 17: Binding Affinities of exemplary antibodies (ELISA and BIACORE)

Assay	Tim3_0 018	Tim3 _0021	Tim3 _0028	Tim3 _0026	Tim3 _0033	Tim3 _0038
Affinity KD [nM]	3.4 / 1.1	204 / 4.1	173 / 2.8	6.2 / 1.5	n.f. / 3.1	7.6 / 0.6
mono / bivalent Tim3						
EC50 ELISA [nM]	0.56		0.22			0.501
EC50 ELISA [ng/ml]	94	47	37	47	1321	83
EC50 CHO-Tim3 [nM]	0.52		0.32			0.17

EC50 CHO-Tim3 [ng/ml]	87	73	53	69	3710	29
-----------------------	----	----	----	----	------	----

Biacore characterization of the Tim3 ABs A surface plasmon resonance (SPR) based assay has been used to determine the kinetic parameters of the binding between several murine Tim3 binders as well as commercial human Tim3 binding references. Therefore, an anti-mouse IgG was immobilized by amine coupling to the surface of a (Biacore) CM5 sensor chip. The samples were then captured and hu/cy Tim3-ECD was bound to them. The sensor chip surface was regenerated after each analysis cycle. The equilibrium constant K_D was finally gained by fitting the data to a 1:1 langmuir interaction model. About 12000 response units (RU) of 30 μ g/ml anti-mouse IgG (GE Healthcare #BR-1008-38) were coupled onto the spots 1,2,4 and 5 of the flow cells 1-4 (spots 1,5 are active and spots 2,4 are reference spots) of a CM5 sensor chip in a Biacore B4000 at pH 5.0 by using an amine coupling kit supplied by GE Healthcare.

The sample and running buffer was HBS-EP+ (0.01 M HEPES, 0.15 M NaCl, 3 mM EDTA, 0.05% v/v Surfactant P20, pH 7.4). Flow cell temperature was set to 25 °C and sample compartment temperature to 12 °C. The system was primed with running buffer. The samples were injected for 30 seconds with a concentration of 200 μ g/ml and bound to the spots 1 and 5 of each flow cell, allowing the measurement of eight samples in parallel. Then a complete set of different (monomeric cyno, monomeric human and huFc fused dimeric human Tim3-ECD) concentrations (s. Table X) was injected over each sample for 240 s followed by a dissociation time of 30/1800 s (s. Table 1). Each analysis cycle (sample capture, spot 1 and 5 – Tim3 ECD injection) was then regenerated with a 30 seconds long injection of Glycine-HCl pH 1.7. The flow rate was set to 30 μ l/min for the whole run. Finally, the double referenced data was fitted to a 1:1 langmuir interaction model with the Biacore B4000 Evaluation Software. Resulting K_D values are shown in Table 17 and 18.

Table 18: Binding affinities determined by Biacore-KD values gained by a kinetic SPR measurement.-n.f. means no fit possible, most likely due to no or weak binding.

Sample	huTim3 K_D (25 °C) [M]	huTim3Fc K_D (25 °C) [M]	cyTim3 K_D (25 °C) [M]
TIM3-0016	3.29E-09	1.09E-09	2.16E-08
TIM3-0016 variant (0018)	3.40E-09	1.11E-09	4.19E-08
TIM3-0021	2.04E-07	4.07E-09	n.f.

TIM3-0022	1.26E-07	1.52E-09	2.84E-08
TIM3-0026	6.23E-09	1.52E-09	n.f.
TIM3-0028	1.73E-07	2.77E-09	n.f.
TIM3-0030	3.11E-09	1.28E-09	n.f.
TIM3-0033	n.f.	3.05E-09	n.f.
TIM3-0038	7.56E-09	5.69E-10	n.f.
Reference antibody Biolegend F38-2E2	1.36E-08	7.50E-09	1.68E-07
Reference antibody USB 11E365	1.34E-08	7.73E-09	1.41E-07

Example 20

Generation of anti-Tim3 antibody derivatives

Chimeric antibodies derivatives Chimeric Tim3 antibodies were generated by amplifying the variable heavy and light chain regions of the anti-TIM3 mouse antibodies Tim3-0016, Tim3-0016 variant (0018), Tim3-0021, Tim3-0022, Tim3-0026, Tim3-0028, Tim3-0030, and Tim3-0033, Tim3-0038 from via PCR and cloning them into heavy chain expression vectors as fusion proteins with human IgG1 backbones / human CH1-Hinge-CH2-CH3 with LALA and PG mutations (Leucine 234 to Alanine, Leucine 235 to Alanine, Proline 329 to Glycine) abrogating effector functions and light chain expression vectors as fusion proteins to human C-kappa. LC and HC Plasmids were then cotransfected into HEK293 and purified after 7 days from supernatants by standard methods for antibody purification.

Removal of glycosylation site NYT: Modifying 1 HVR-L1 position in Tim3-0016, Tim3_0016 variant (named 0018 or Tim3_0018) by substitution of N by Q or S Mutations within the variable light vchain region of Tim3_0016 and Tim3_0016 variant (0018) were generated by in vitro mutagenesis using Agilent "Quick Change Lightning Site-directed Mutagenesis Kit" according manufacturer's instructions. By this method the asparagine (N) of the glycosylation site motif NYT in the light chain HVR-L1 (SEQ ID NO: 4) was replaced by glutamine (Q) (resulting in SEQ ID NO: 11 = Tim3_0016_HVR-L1 variant 1_ NQ) or, alternatively, the asparagine (N) was replaced by serine (S) (resulting in SEQ ID NO: 12 = Tim3_0016_HVR-L1 variant 2_ NS). In both glycosylation site motif NYT was successfully modified. LC and HC Plasmids coding for the variants were then cotransfected into HEK293 and purified after 7 days from supernatants by standard methods for antibody purification. The generated mutants

were tested by ELISA on human Tim3, ELISA on cynomolgus Tim3 and cellular ELISA on adherent CHO-K1 cells expressing full-length human Tim3. All mutants generated were found to show even more functional binding to human TIM3 (human) , cyno TIM3 (cyno) or human TIMR on CHO cells than the parental antibodies Tim3_0016 or the

5 Tim3_0016 antibody variant Tim3_0018 respectively.

Table 19:

Mutants tested	Biochem Human		Biochem Cyno		cellular bindg. CHO-Tim3	
	EC50 [ng/ml] values in relation to the sample's max value	inflexion point [ng/ml]	EC50 [ng/ml] values in relation to the sample's max value	inflexion point [ng/ml]	EC50 [ng/ml] values in relation to the sample's max value	inflexion point [ng/ml]
aTim3 F38 2E2	73,2	88,3	423,0	209871,3	150,2	224,3
aTim3 0018	15,1	15,3	14,6	14,6	26,4	29,4
aTim3 0018MutNQ	12,0	10,8	13,2	10,8	13,4	12,8
aTim3 0018MutNS	10,3	6,5	11,9	6,5	11,2	11,1
aTim3 0016 MutNQ	7,6	5,7	8,3	5,7	6,3	5,4
aTim3 0016MutNS	8,5	5,5	9,7	5,5	9,1	8,5

Example 21

Fluorescent Labeling of Purified Monoclonal Antibody

10 The fluorescent labeling of the hybridoma derived monoclonal antibody was carried out by using Alexa Fluor 488 Monoclonal Antibody Labeling Kit (manufactured by Invitrogen) according to the manufacturer's instructions. After the labeling, each antibody was confirmed to be positively labeled with Alexa Fluor 488 (hereinafter referred to as "Alexa-488") by FACSCalibur (manufactured by BD Biosciences) analysis for TIM-3

15 expressing RPMI-8226 and Pfeiffer cells.

Example 22

Classification of Binding Epitope Groups using FACS based Competition Assay

The relation of epitopes between generated anti-TIM3 antibodies and six anti-TIM3 reference antibodies was analyzed by a FACS based binding competition assay. The

20 TIM3 reference antibodies were the following: antibodies 4177 and 8213 as described in US2012/189617 , antibodies 1.7E10 and 27.12E12 as described in WO2013/06490; antibody 344823 (Clone 344823, manufactured by R&D Systems) and antibody F38-2E2 (Clone F38-2E2, manufactured by BioLegend and R&D Systems). In brief, the test antibody was allowed to interact and bind with the TIM-3 expressing RPMI-8226 cells

25 (ATCC ® CCL-155™) and then it was evaluated by flow cytometry method whether another anti-TIM-3 antibody could also bind to TIM-3 expressing cells.

In short human TIM3 expressing RPMI-8226 cells were incubated with BD human Fc Block for 10 min at RT and stained in two different experimental setups to exclude the impact of the difference in the affinity of the tested antibodies on the binding:

-188-

1) with disclosed purified anti-TIM3 (10µg/ml in BD staining buffer for 0.5h at 4oC), which were conjugated with Alexa*488 according to the manufacturer's instructions (Molecular Probes A-20181) with an average of 2.7 fluorophores per antibody. Then a) unlabeled (1-4) reference recombinant anti-TIM3 antibodies or Isotype control were added (10µg/ml) for 0.5h at 4oC in BD SB and after washing with BD SB stained with PE-labeled anti-huFcγ Abs (JIR, 109-116-098, 1:200, 0.5h at 4oC in BD SB) or b) PE labeled (5-6) available reference anti-TIM3 antibodies or appropriate Isotype controls were added (10µg/ml) for 0.5h at 4oC in BD SB. After washing and centrifugation MFI signals of stained RPMI-8226 cells were analyzed by BD Biosciences FACSCanto flow cytometer.

Table 20: Summary of epitope characterization.

	Max % Inhibition of Binding					
	Epitope group 1			Epitope group 3		
	1a		1b	3a	3b	
	TIM3-0016	TIM3-0018	TIM3-0026	TIM3-0022	TIM3-0028	TIM3-0038
clone 4177	1	-9	29	79	-3	0
clone 8213	-2	9	9	9	38	29
clone 1-7E10	-5	15	24	0	20	7
clone 27-12E12	-1	4	22	40	82	94
clone 344823	0	0	3	102	107	99
clone F38-2E2	-7	-6	2	77	75	94

100	>90
100	>50
100	>30
100	>20

Results from the FACS based epitope groups mapping show that Tim3_0016 and Tim3_0016 variant Tim3_0018 show no binding competition with any tested anti-TIM-3 reference antibodies and it was suggested that these Abs recognized the new epitope different from the epitopes to which all previous described TIM3 reference antibodies recognized whereas Tim3_0022, Tim3_0026, Tim3_0028 and Tim3_0038 compete to different extend for binding to surface expressed TIM3 on JRPMI-8226 cells with various competitors.

Example 23

Effect of human anti-TIM-3 Antibodies on Cytokine Production in a Mixed Lymphocyte Reaction (MLR)

A mixed lymphocyte reaction was used to demonstrate the effect of blocking the TIM-3 pathway to lymphocyte effector cells. T cells in the assay were tested for activation and their IFN-gamma secretion in the presence or absence of an anti-TIM-3 mAbs.

Human Lymphocytes were isolated from peripheral blood of healthy donor by density gradient centrifugation using Leukosep (Greiner Bio One, 227 288). Briefly, heparinized blood were diluted with the three fold volume of PBS and 25 ml aliquots of the diluted blood were layered in 50 ml Leukosep tubes. After centrifugation at 800 x g for 15 min at room temperature (w/o break) the lymphocyte containing fractions were harvested, washed in PBS and used directly in functional assay or resuspended in freezing medium (10% DMSO, 90 %FCS) at 1.0×10^7 cells/ml and stored in liquid nitrogen. 1:1 target/responder cell ratio was used in MLR assay (i.e. each MLR culture contained - 2.0E+05 PBMCs from each donor in a total volume of 200 μ l. Anti-TIM3 monoclonal antibodies Tim3_0016, Tim3_0016 variant (Tim3_0018), Tim3_0021, Tim3_0022, Tim3_0026, Tim3_0028, Tim3_0030, Tim3_0033, Tim3_0038 and F38-2E2 (BioLegend), were added to each culture at different antibody concentrations. Either no antibody or an isotype control antibody was used as a negative control and recombinant human IL-2 (20 EU/ml) was used as positive control. The cells were cultured for 6 days at 37°C. After day 6 100 μ l of medium was taken from each culture for cytokine measurement. The levels of IFN-gamma were measured using OptEIA ELISA kit (BD Biosciences).

The results are shown in Table 21 (IFN-g secretion/release). The anti-TIM-3 monoclonal antibodies promoted T cell activation and IFN-gamma secretion in concentration dependent manner. The anti-TIM3 antibodies Tim3_0021, Tim3_0022, Tim3_0028, and Tim3_0038 reduce release of the inflammatory cytokine IFN-gamma more than the F38-2E2 antibody. Tim3_0016, Tim3_0016 variant (Tim3_0018), Tim3_0033 and Tim3_0038 showed a similar release when compared the F38-2E2 antibody. In contrast, cultures containing the isotype control antibody did not show an increase in IFN-gamma secretion.

Table 21: Percentage anti-Tim3 antibody induced IFN-gamma release in comparison to recombinant human IL-2 (20 EU/ml) (= 100%) as positive control and no antibody as negative control (Donors

Compound concentration	MLR+IL-2 20U/ml	Isotype IgG2a	F38-2E2	Tim3 0016	Tim3 0018	Tim3 0021	Tim3 0022	Tim3 0026	Tim3 0028	Tim3 0030	Tim3 0033	Tim3 0038	Isotype hIgG1
40 µg/ml		2	36	33	36	112	58	25	40	14	35	51	0
10 µg/ml	100	0	26	22	30	108	38	16	38	4	30	38	5
1 µg/ml		0	7	7	12	101	18	18	12	3	0	1	0

Example 24

Internalization of anti-TIM-3 Antibodies into TIM-3 expressing cells

TIM-3-specific antibodies described herein can be internalized into TIM-3-expressing cells, including TIM-3 expressing lymphoma, multiple myeloma and AML cells. For example, the disclosed TIM-3 specific antibodies and fragments thereof are shown to be internalized into rec TIM3 CHO cells stably expressing human TIM-3 as evaluated by cell based ELISA, flow cytometry (FACS) and confocal microscopy.

Stable Tim3-transfected CHO-K1 cells (clone 8) (4x10⁴ cells/well/100µl) were seeded into 96 well-MTP using fresh culture medium. After overnight cell attachment, cell culture medium was removed and the test antibodies were added to the cells (10µg/ml in cell culture medium) and incubated for 0.5 hour at 4°C. As reference, a commercial mouse-anti-human antibody (TIM3 MAB 11E365 (US Biological, T5469-92P) was used. After washing (2x with cell culture medium) and centrifugation cells were incubated for 3 h at a) 4°C or b) 37°C in 200 µl cell culture medium. Internalization typically occurs at 37°C, but not at 4°C, which provides another control for the reaction. Then cells were fixated with 100 µl/well 0.05 % glutaraldehyde (Sigma Cat.No: G5882) in 1xPBS for 10 min at room temperature (RT). This was followed by three washing steps with 200 µl PBS-T and secondary antibody sheep-anti-mouse-POD (Horseradish POD linked F(ab')₂ Fragment ; GE NA9310)) were added for 1 hour at RT. After the final washing steps (3xPBS-T), TMB substrate was added (Roche order no. 11835033001) for 15 min and color development was stopped using 1N HCl. Final ODs were determined by measurement at 450/620 nm in an ELISA reader. This cellular ELISA procedure was used for medium throughput evaluation of the internalizing capacity of the testing antibodies which were purified from hybridoma supernatants.

The percentage of internalization was calculated as follow:

$$\text{Internalization [\%]} = (1 - \text{OD}_{\text{sample}_{37^{\circ}\text{C}}} / \text{OD}_{\text{sample}_{4^{\circ}\text{C}}}) * 100$$

The results are shown in Fig. 29A and B for (Internalization). Almost all tested anti-TIM-3 monoclonal antibodies were similar well internalized into stable Tim3-transfected CHO-K1 cells after 3h incubation at 37°C (not all data shown).

5 The determination of EC50 internalizing values (time dependency) as well as comparison of the kinetics of the internalization depending on mono- vs. bivalency was estimated by FACS for selected candidates.

In short, human TIM3 stable expressing CHO-K1 cells were seeded (4×10^5 cells/well/50µl) into 96 well-v bottom MTP using fresh culture medium and incubated with Redimune® NF Liquid for 10 min at RT to block unspecific binding. Then 50 µl/well of selected purified anti-TIM3 (10µg/ml in cell culture medium) were added and incubated for 1h at 4°C. After washing (with cell culture medium) and centrifugation cells were incubated for 0.25, 0.5, 1, 2, 3, 4, 6 and 24 h at a) 4°C or b) 37°C in 200 µl cell culture medium. Then cells were washed with PBS/1%BSA and secondary antibody Alexa Fluor 488 Goat-anti-mouseIgG, F(ab)2 were added for 1 hour at 4°C. After washing and centrifugation 125 µl of CellFix (BD Bioscience, 1:1000) were added and MFI signals of stained cells were analyzed by BD Biosciences FACSCanto flow cytometer.

The percentage of internalization was calculated as follow:

$$\text{Internalization [\%]} = (1 - \text{MFI}_{\text{sample}_{37^\circ\text{C}}} / \text{MFI}_{\text{sample}_{4^\circ\text{C}}}) * 100$$

20 Example for the evaluation of time dependent internalization of anti-TIM3 antibodies Tim3_0016, Tim3_0016 variant (Tim3_0018), Tim3_0021, Tim3_0028, Tim3_0030, Tim3_0033, Tim3_0038 on RPMI-8226 cells (ATCC ® CCL-155™):

The presently disclosed anti-TIM3 antibodies are internalized rapidly into TIM3 expressing RPMI-8226 cells (ATCC ® CCL-155™) at a high level. The experiments were conducted as described above with TIM3 expressing RPMI-8226 cells (ATCC ® CCL-155™) instead of rec CHO-K1 cells expressing huTIM-3. Results are shown in the Table 22. The following antibodies were used as TIM3 reference antibodies: antibody 8213 as described in US2012/189617, antibody 27.12E12 as described in WO2013/06490. Tim3_0016, Tim3_0016 variant (Tim3_0018), Tim3_0038 were used as human IgG1 chimeric versions.

Table 22: Percentage internalization at the indicated time point (0 min set as 0 percent).

5		Percentage internalization of anti-TIM3 antibodies				
	Antibody	30 Min	60 Min	120 Min	240 Min	26h
10	8213	22	22	43	52	72
	27.12E12	19	22	25	46	59
	Tim3_0016	33	52	55	66	87
	Tim3_0018	39	41	80	70	88
	Tim3_0021	70	75	74	78	77
	Tim3_0028	50	59	67	68	83
	Tim3_0033	75	81	82	82	80
	Tim3_0038	22	20	45	46	63

The results show that the tested antibodies are rapidly internalized at high percentage compared to reference antibodies on RPMI-8226 cells (ATCC ® CCL-155™).

Example 25

20 Binding of anti-TIM-3 Antibodies to isolated human Monocytes expressing TIM-3

CD14+ Monocytes were isolated from anticoagulated peripheral blood of healthy donors by density gradient centrifugation using Ficoll-Paque (GE Healthcare) (see General Protocols in the User Manuals or visit www.miltenyibiotec.com/protocols) and subsequent positive selection via CD14 MicroBeads. First the CD14+ cells are
 25 magnetically labeled with CD14 MicroBeads. Then the cell suspension is loaded onto a MACS® Column which is placed in the magnetic field of a MACS Separator. The magnetically labeled CD14+ cells are retained in the column. The unlabeled cells run through, this cell fraction is depleted of CD14+ cells. After removal of the column from the magnetic field, the magnetically retained CD14+ cells can be eluted as the positively
 30 selected cell fraction. After centrifugation at 200 x g for 10 min at room temperature the monocytes were harvested and used directly in binding assay or resuspended in freezing medium (10% DMSO, 90 %FCS) at 1.0E+07 cells/ml and stored in liquid nitrogen.

As shown in the literature Monocytes express constitutively TIM3 on their surface. 1x10⁵
 35 CD14+ isolated human monocytes (50µl/well) were put into 96 well-v bottom MTP in fresh culture medium and incubated with Redimune® NF Liquid for 15 min at RT to block unspecific binding. Then 50 µl/well of disclosed anti-TIM3 mAbs or reference anti-TIM-3 mAbs 344823 (R&D) and F38-2E2 (BioLegend) (10µg/ml in cell culture medium)

were added and incubated for 1h at 4°C. Then cells were washed with PBS/1%BSA and secondary antibody PE-labeled Goat-anti-mouse F(ab')₂ were (Jackson Lab 115-006-072) added for 1 hour at 4°C. After washing and centrifugation MFI signals of stained cells were analyzed by BD Biosciences FACSCanto flow cytometer.

- 5 The specific binding was calculated as follow:

Specific Binding [MFI] = Geom. Mean MFI_{sample} - Geom. Mean MFI_{isotype control}

The results are shown in Table 8: (Binding to human Monocytes). TIM3 clones

Tim3_0016, Tim3_0018, Tim3_0020, Tim3_0028 and Tim3_0038 bind to human monocytes of different donors even better than the reference anti- TIM-3 Abs .

- 10 Table 23: Binding to human Monocytes.

	donor1 (CD14+)	donor2 (CD14+)	donor3 (CD14+)
Tim3 0016	2122	1634	1690
Tim3 0018	2326	1818	1943
Tim3 0020	1917	1377	1462
Tim3 0021	1134	951	1197
Tim3 0022	1468	1111	1235
Tim3 0026	1665	1016	900
Tim3 0030	1411	419	466
Tim3 0038	1637	1368	1401
Tim3 0028	1351	950	1607
Tim3 0033	480	328	595
M-IgG2b	0	13	0
M-IgG1	144	55	213
<TIM-3>PE Mab, M-IgG1 (Cl F38-2E2; Biolegend)	516	493	460
<TIM-3> PE Mab , Rat IgG2A (Clone 344823, R&D)	1010	917	814
Rat-IgG2A-PE	71	68	70

Example 26

Binding of anti-TIM-3 Antibodies to isolated cyno Monocytes expressing TIM-3

- 15 CD14⁺ Monocytes were isolated from cynomolgus monkey anticoagulated peripheral blood (Covance) by density gradient centrifugation using Ficoll-Paque (GE Healthcare) (see General Protocols in the User Manuals or visit www.miltenyibiotec.com/protocols) and subsequent positiv selection via NHP CD14 MicroBeads. First the CD14⁺ cells are magnetically labeled with CD14 MicroBeads. Then the cell suspension is loaded onto a MACS® Column which is placed in the magnetic field of a MACS Separator. The

magnetically labeled CD14+ cells are retained in the column. The unlabeled cells run through, this cell fraction is depleted of CD14+ cells. After removal of the column from the magnetic field, the magnetically retained CD14+ cells can be eluted as the positively selected cell fraction. After centrifugation at 200 x g for 10 min at room temperature the monocytes were harvested and used directly in binding assay or resuspended in freezing medium (10% DMSO, 90 %FCS) at 1.0E+07 cells/ml and stored in liquid nitrogen.

As shown in the literature Monocytes express constitutively TIM3 on their surface. 1x10⁵ CD14+ isolated cyno monocytes (50µl/well) were put into 96 well-v bottom MTP in fresh culture medium and incubated with Redimune® NF Liquid for 15 min at RT to block unspecific binding. Then 50 µl/well of Alexa488 labeled anti-TIM3 (10µg/ml in cell culture medium) were added and incubated for 1h at 4°C. After washing and centrifugation MFI signals of stained cells were analyzed by BD Biosciences FACSCanto flow cytometer.

The specific binding was calculated as follow:

Specific Binding [MFI] = Geom. Mean MFI_{sample} - Geom. Mean MFI_{isotype control}

The results are shown in Table 9 (Binding to Cyno Monocytes). TIM3 clones Tim3_0016, Tim3_0018, Tim3_0026, Tim3_0028 and , Tim3_0030 bind to cyno monocytes of different cyno donors.

Table 24: Binding to Cyno Monocytes.

	cyno1 (16719M) CD14+	cyno2 (17435M) CD14+	cyno3 (30085F) CD14+
AF +PI	75	83	84
HumTIM-3 Alexa488 R&D (34482)	158	121	143
Rat-IgG2A-Alexa488	84	86	91
hum TIM-3 A488 F38-2E2 (NOVUS Biol)	135	136	124
M-IgG1-Alexa 488	72	82	83
Tim3 _0016- A488	157	177	187
Tim3 _0016 variant 0018- A488	301	480	417
Tim3 0022- A488	115	134	138
Tim3 0026- A488	137	184	197
Tim3 0028- A488	3936	2996	4090
Tim3 0038- A488	97	107	120
Tim3 _0020- A488	274	378	354
Tim3 0021 A488	348	473	399

-195-

Tim3 0030 A488	119	163	144
Tim3 0033 A488	71	81	83
TIM-3 (4177) A488	78	83	85
TIM-3 (8213) A488	75	83	87

Example 27

Binding of anti-TIM-3 Antibodies to NHL and MM cell lines expressing TIM-3

- 5 The binding capacity of disclosed anti-TIM3 antibodies and two anti-TIM3 reference antibodies clones (1) 4177 and (2) 8213 (Kyowa) was analyzed by a FACS. In short human TIM3 expressing B cell lymphoma cells (exemplified as Pfeiffer cells) and multiple myeloma cells (exemplified as RPMI-8226 cells) were incubated with BD human Fc Block for 10 min at RT to block unspecific binding. Then 2×10^5 cells
- 10 (50 μ l/well) were put into 96 well-v bottom MTP and 50 μ l/well of Alexa488 labeled anti-TIM3 (10 μ g/ml in BD Staining buffer) were added and incubated for 1h at 4°C. After washing and centrifugation MFI signals of stained cells were analyzed by BD Biosciences FACSCanto flow cytometer.

The specific binding was calculated as follow:

- 15 Specific Binding [MFI] = Geom. Mean MFI_{sample} - Geom. Mean MFI_{isotype control}
- The results are shown in FIG. 2A and 2B (Binding to RPMI-8226 and Pfeiffer cells).

Example 10: Cytotoxic activity of anti-TIM-3 Antibodies on TIM-3 expressing NHL and MM cells

- TIM3-specific antibodies conjugated with pseudomonas exotoxin (PE 24) effectively kill
- 20 TIM3-expressing cells. The cytotoxic activity of disclosed anti-TIM3 antibodies and one commercial available anti-TIM3 reference antibody clone 11E365 (available from US Biological) was analysed with Promega CellTiter-Glo Luminescent Cell Viability Assay. In short to 5×10^3 (50 μ l/well in 96 well MTP, in triplicate) recombinant CHO K1 stable expressing human TIM-3 or 2×10^4 cells (50 μ l/well in 96 well MTP, in triplicate)
- 25 human TIM3 expressing B cell lymphoma cells (exemplified as Pfeiffer cells) or multiple myeloma cells (exemplified as RPMI-8226 cells) were added 25 μ l/well 1:5 serial dilution of disclosed anti-TIM-3 antibodies with the highest concentration of 10 μ g/ml or appropriate media to untreated cells or Isotype control to untargeted treated cells. Treatment ranges from 10 μ g/ml to 1 ng/ml in triplicate. All antibodies were used as full

length mouse Fcγ versions . For conjugation of the conjugation of the Pseudomonas exotoxin 10μg/ml of mouse Fcγ fragment specific Fabs conjugated with PE 24 were added and incubated for 3 days at 37°C. Cycloheximide as a known protein synthesis inhibitor in eukaryotes was used as positive control. Viability of treated cells were measured with Promega CellTiter-Glo Luminescent Cell Viability Assay.

The cytotoxic activity was calculated as follow:

$$\text{Rel. Inhibition [\%]} = (1 - (\text{Esampel} - \text{E negative control}) / (\text{E positive control} - \text{E negative control})) * 100$$

The results are shown in Table 25.

10 Table 25: Cytotoxic activity of anti-TIM3 mAbs on TIM-3 expressing recombinant, NHL and MM cell lines in sandwich format.

Antibodies and references (all anti TIM3 antibodies conjugated to a deimmunized Pseudomonas exotoxin A)	IC50 [nM]		
	recTIM-3 CHO cells	Pfeiffer cells	RPMI-8226
Tim3_0016	0,04	0,09	0,55
Tim3_0016 variant 1 (Tim3_0018)	0,05	0,10	0,66
Tim3_0020	0,07	0,11	>64
Tim3_0021	0,04	0,10	5,9
Tim3_0022	0,02	0,07	0,36
Tim3_0023	0,03	0,08	>64
Tim3_0026	0,03	0,08	>64
Tim3_0030	0,03	0,10	>64
Tim3_0033	0,11	0,20	0,79
Tim3_0038	0,01	< 0.002	0,16
US Biol. Clone 11E365	0,7	1,2	1,1
Cells w/o Ab	-	-	-
Cells + <mFc> Fab PE	-	-	-
IgG2A + <mFc> Fab PE	-	-	-
Cycloheximide	135	181	245

All tested TIM3 clones are highly potent (IC50 range 0,01-0,2 nM) on recombinant CHO K1 stable expressing human TIM-3 and Pfeiffer cells expressing high and moderate levels of TIM-3 and even more potent in their cytotoxic activity than the strong internalizing reference anti-TIM-3 Ab clone 11E365, US Biological. TIM3 clones 0016,

0018, 0021, 0022, 0033 and 0038 are also potent on RPMI-8226 cells expressing 5 fold lower TIM-3 level compare to recombinant CHO TIM-3 cells .

Example 28: Comparison of the cytotoxic activity of disclosed anti-TIM3 antibodies vs. two anti-TIM3 reference antibodies 1.7.E10 and 27-12E12 (as described in

5 WO2013/06490).

The cytotoxic activity of disclosed anti-TIM3 antibodies and two anti-TIM3 reference antibodies the TIM3 reference antibodies 1.7E10 and 27.12E12 as described in

WO2013/06490 was analysed with Promega CellTiter-Glo Luminescent Cell Viability Assay as described above. All antibodies were used as full lenght human IgG1 format

10 includimng the human Fcgamma part. In this experiment conjugation of the

Pseudomonas exotoxin was achieved via human Fc γ fragment specific Fabs conjugated with PE 24 (10 μ g/ml) which were added and incubated for 5 days at 37°C.

The results are shown in Table 26.

Table 26: Comparison of cytotoxic activity of anti-TIM3 mAbs on TIM-3 expressing

15 NHL and MM cell lines.

	Pfeiffer cells		RPMI-8226 cells	
Antibodies and references (all anti TIM3 antibodies conjugated to a deimmunized Pseudomonas exotoxin A)	Max. killing	Rel. IC50 [nM]	Max. killing	Rel. IC50 [nM]
Cycloheximide	100 [%]	271	100 [%]	111
1.7E10	60.3 [%]	0,68	65.7 [%]	2,544
27-12E12	75.7 [%]	0,02	86.6 [%]	0,111
Tim3_0016	84.9 [%]	0,05	86.6 [%]	0,063
Tim3_0016 variant (Tim3_0018)	82.9 [%]	0,06	88.1 [%]	0,081
Tim3_0026	78.3 [%]	<0.02	83.1 [%]	0,067
Tim3_002	82.6 [%]	<0.02	83.8 [%]	0,047
Isotype Control hIgG1	3.2 [%]	N.A	0.4 [%]	N.A

All disclosed TIM3 clones are highly active (IC50 range 0,02-0,08 nM) on Pfeiffer and RPMI-8226 cells expressing TIM-3 and even more potent in their cytotoxic activity than

20 the strong internalizing reference anti-TIM-3 Ab clone 27-12E12. All antibodies were compared as Pseudomonas exotoxin (PE24) conjugates using the same Pseudomonas exotoxin under the same conditions.

Example 28**Cytotoxic activity of Fab-PE24 constructs of disclosed anti-TIM3 antibodies on MM, NHL and AML cell lines (expressing TIM3, but not PSMA).**

The cytotoxic activity was analysed with Promega CellTiter-Glo Luminescent Cell

- 5 Viability Assay as described above. 1:5 serial dilutions of Fab-fragments of disclosed anti-TIM3 antibodies directly conjugated to PE24 with the highest concentration of 50µg/ml or appropriate media to untreated cells or non-binding anti-PSMA Fab-PE24 control to untargated treated cells were incubated with 7,5 x10³ Pfeiffer cells or 2 x10³ RPMI-8226 cells (50µl/well in 98 well MTP) for 4 days at 37°C. Treatment ranges from
- 10 50µg/ml to 8ng/ml in triplicate. Cycloheximide was used as positive control.

The results are shown in Table 27.

Table 27: Cytotoxic activity of Fab-PE24 constructs of disclosed anti-TIM3 antibodies on MM, NHL and AML cell lines.

	RPMI-8226		Karpas-299		CMK		TF-1		MOLM-13	
Antibodies and references (all anti TIM3 antibodies conjugated to a deimmunized Pseudomonas exotoxin A)	Max. killing	IC50 [nM]	Max. killing	IC50 [nM]	Max. killin g	IC50 [nM]	Max. killin g	IC50 [nM]	Max. killin g	IC50 [nM]
Cycloheximide	100 [%]	281	100 [%]	113	100 [%]	149,0	100 [%]	207	100 [%]	156
Anti_PSMA	10.5 [%]	N.A.	40.1 [%]	N.A.	8.98 [%]	N.A.	5.27 [%]	N.A.	18.9 [%]	N.A.
Tim3_0022	99.1 [%]	1,9	98.8 [%]	10	67.1 [%]	255	58.6 [%]	299	58.5 [%]	579
Tim3_0016	99.3 [%]	1,1	99.2 [%]	4	64.8 [%]	225	54.2 [%]	534	62.7 [%]	459

- All tested Fab-PE24 constructs of disclosed anti-TIM3 antibodies are highly potent (IC50
- 15 range 1-10 nM) on MM (RPMI-8226) and NHL (Karpas-299) cells expressing moderate level of TIM-3 and demonstrate significant cytotoxic activity on AML cell lines (CMK, TF-1, MOLM-13) expressing very low levels of TIM-3.

Example 29

Cytotoxic activity of Immuno conjugates (Pseudomonas Exotoxin A conjugates (Fab-PE24 constructs) of disclosed anti-TIM3 on primary leukemic stem/progenitor AML cells from relapsed/refractory patients

CD34+ cells from peripheral blood of relapsed/refractory patients were obtained from AllCells, LLC, Alameda, CA. After confirmation of purity and viability of all samples (purity range 84-94% and viability range 95-99%) the expression level of TIM-3 was evaluated by FACS as described in Example 7 using anti-TIM-3 mAbs 344823 (R&D).

(see Fig. 31). All tested (4/4) primary leukemic stem/progenitor (CD34+) AML samples from relapsed/refractory patients demonstrate homogeneous expression of TIM-3 at different levels.

For the evaluation of cytotoxic activity of Fab-PE24 constructs of disclosed anti-TIM3 clones 0016 and 0022 on primary CD34+ AML cells 1×10^4 cells (50 μ l/well in 96 well MTP, in triplicate) were incubated with 1:5 serial dilutions of Fab-fragments with the highest concentration of 50 μ g/ml or appropriate media to untreated cells or non-binding anti-PSMA Fab-PE24 control to untargeted treated cells for 3 days at 37°C. Cycloheximide was used as positive control. Cytotoxic activity was analysed with Promega CellTiter-Glo Luminescent Cell Viability Assay as described above in Example 28.

The results are shown in Table 28. (Cytotoxic activity of Fab-PE24 constructs of disclosed anti-TIM3 antibodies on primary CD34+ AML cells).

Table 28: Cytotoxic activity of Fab-PE24 constructs of disclosed anti-TIM3 antibodies on primary CD34+ AML cells).

	D1; AML CD34+ PB0136 cells		D2; AML CD34+ PB0142 cells		D3; AML CD34+ PB0135 cells		D4; AML CD34+ PB0193 cells	
Antibodies and references (all anti TIM3 antibodies conjugated to a deimmunized Pseudomonas exotoxin A)	Max. killing	IC50 [nM]	Max. killing	IC50 [nM]	Max. killing	IC50 [nM]	Max. killing	IC50 [nM]

-200-

Cycloheximide	100 [%]	212	100 [%]	262	100 [%]	121	100 [%]	208
anti-PSMA	2 [%]	N.A.	8 [%]	N.A.	18 [%]	N.A.	12 [%]	N.A.
TIM-3 0022-cFP	38 [%]	>691	75 [%]	107	31 [%]	>691	57 [%]	375
TIM-3 0016-cFP	48 [%]	>691	79 [%]	30	44 [%]	>691	69 [%]	116

Fab-PE24 constructs of anti-TIM3 antibodies Tim3_0016 and Tim3_0022 are highly potent on (2/4) primary AML samples (PB0142 and PB0135) (IC50 range 30-116 nM) and demonstrate significant cytotoxic activity on all (4/4) primary leukemic stem/progenitor (CD34+) AML cells expressing different levels of TIM-3.

5

Example 30

Comparison of potency of Fab-PE24 constructs of selected anti-TIM3 antibodies on NHL and MM cell lines

The evaluation of cytotoxic activity of sortase coupled Fab-PE24 constructs of selected disclosed anti-TIM3 antibodies was analysed with Promega CellTiter-Glo Luminescent Cell Viability Assay as described above in Example 28.

The results are shown in Table 29.

Table 29: Cytotoxic activity of Fab-PE24 constructs of selected anti-TIM3 antibodies on NHL and MM cells.

15

Antibodies and references (all anti TIM3 antibodies conjugated to a deimmunized Pseudomonas exotoxin A)	Pfeiffer cells		RPMI-8226 cells	
	Max. killing	IC50 [nM]	Max. killing	IC50 [nM]
Cycloheximide	100 [%]	271,1	100 [%]	153
anti-PSMA	25.2 [%]	N.A.	21.5 [%]	N.A.
TIM-3 0022	99.9 [%]	1,58	99.6 [%]	2,14
TIM-3 0016	99.6 [%]	0,77	99.2 [%]	0,61
TIM-3 0021	98.4 [%]	2,15	99.1 [%]	3,61
TIM-3 0033	99.8 [%]	5,30	99.7 [%]	5,73
TIM-3 0038	99.6 [%]	0,47	98.3 [%]	0,32

-201-

High cytotoxic potency was demonstrated with Fab-PE24 constructs of all selected disclosed anti-TIM3 antibodies (IC₅₀ range 0,3-5 nM) on NHL (Pfeiffer) and MM (RPMI-8226) cells expressing moderate level of TIM-3.

The highest cytotoxic activity was observed with Fab-PE24 constructs of disclosed anti-TIM3 antibodies Tim3_ 0016 and Tim3_ 0038.

Example 31

Comparison of cytotoxic activity of Fab-PE24 construct vs. total-IgG-Amatoxin conjugate of the same clone of disclosed anti-TIM-3 antibody on Pfeiffer cells

The evaluation of cytotoxic activity of conjugated Fab-PE24 construct of disclosed anti-TIM3 clone 0016 vs. total IgG of the same clone conjugated with Amatoxin (according to th procedures described in WO2012/041504 (conjugated via the 6' C-atom of amatoxin amino acid 4, particularly via an oxygen atom bound to the 6' C-atom of amatoxin amino acid, and wherein the TIM3 antibody is connected by a linker via a urea moiety) was analysed with Promega CellTiter-Glo Luminescent Cell Viability Assay as described above in Example 12. The results are shown in Table 30.

Table 30: Cytotoxic activity of Fab-PE24 construct vs. total IgG- Amatoxin conjugate of anti-TIM3 clone 0016 on NHL cells

Pfeiffer cells	Max. killing	IC ₅₀ [nM]
Cycloheximide	100 [%]	163
Isotype hIgG1 Amatoxin	28 [%]	N.A.
TIM-3 0016- Amatoxin	93.3 [%]	0,81
TIM-3 0016-PE24	99.8 [%]	0,25

Cytotoxic activity of Amanitin-conugated anti-TIM-3 clone 0016 (IC₅₀ 0,8 nM) is comparable with cytotoxic activity of Fab-PE24 construct of the same clone (IC₅₀ 0,3 nM) on NHL (Pfeiffer) cells expressing moderate level of TIM-3.

Example 32

Patients and tumor sample processing

Freshly excised solid tumor lesions and malignant effusions were collected from 34 patients with non-small cell lung cancer, 7 patients with ovarian cancer and 1 patient with renal cell carcinoma (RCC) between. The solid tumor lesions were dissociated mechanically and digested using accutase (PAA), collagenase IV (Worthington),

hyaluronidase (Sigma), and DNase type IV (Sigma) directly after excision. Single-cell suspensions were prepared. The cellular fraction of malignant effusions was isolated by density gradient centrifugation using Histopaque-1119 (Sigma). All samples were stored in liquid nitrogen until further usage. The study was approved by the local Ethical Review Board (Ethikkommission Nordwestschweiz).

Example 33

Tumor sample characterization

All tumor samples were comprehensively characterized by multicolor flow cytometry. The following antibodies were used for flow cytometric analysis: α -CD4-PE, α -CD8-PE-Cy7, α -CD11b-PerCP-eFluor710, α -CD45-PE-Cy7, α -CD45-PerCP-Cy5.5, α -CD137-FITC, α -BTLA-Biotin, α -CTLA-4-PE, α -ICOS-FITC, α -IFN- γ -FITC, α -Lag-3-APC (all eBioscience), α -CD3-PECF594, α -CD25-BV605, α -CD69-FITC, α -Epcam-FITC, α -granzyme B-PE, α -active caspase 3-PE, α -PD-1-BV605, Steptavidin-BV711 (all BD Bioscience), α -CD45RA-BV421, α -CCR7-AlexaFluor647, α -FoxP3-AlexaFluor647, α -Tim-3-BV421, α -Tim-3-BV605 (all Biolegend). Dead cells were stained with LIVE/DEAD® Fixable *Near-IR* Dead Cell Stain Kit or LIVE/DEAD® Fixable Blue Dead Cell Stain Kit (Invitrogen). For intracellular stainings Fixation and Permabilization Buffers from eBioscience were used. Samples were acquired for flow cytometric analysis on a BD LSR Fortessa. The human IL-2, IFN- γ and TNF ELISA sets were all obtained from BD Bioscience.

CD8⁺ and CD4⁺ T cells (CD45⁺CD3⁺) were characterized for the expression of the surface markers PD-1, Tim-3, CTLA-4, Lag-3, BTLA, CD25, CD69, CD137, ICOS, CD45RA and CCR7. Tumor cells (CD45⁺Epcam⁺) were characterized for the expression of FolR1 comparing the binding of a FolR1 specific antibody with its matched isotype control. Only samples that were positive for FolR1 expression were used for treatment with FolR1-TCB, and samples expressing EpCAM for treatment with catumaxomab, respectively.

Example 34

Ex vivo treatment of tumor samples with FolR1-TCB

FolR1 positive tumor digests or malignant effusions were thawed, washed and plated in 96-well flat bottom cell culture plates (BD Falcon) with a density of 3×10^5 cells/200 μ l/well in complete medium (DMEM + Sodium Pyruvate (1mM) + MEM non essential

AA (1x) + L-Glutamin (2mM) + Penicillin/Streptomycin (100ng/ml) + 2-Mercaptoethanol (50nM) + Ciproxin (1mg/ml) + 10% human Serum). The samples were cultured in the presence or absence of FolR1-TCB or DP47 TCB at a concentration of 2 nM for 24h. Activation of CD8⁺ and CD4⁺ T cells (CD45⁺CD3⁺) upon FolR1-TCB treatment was determined by multicolor flow cytometry by measuring the expression of the cell surface markers CD25, CD69, CD137, ICOS, PD-1 and Tim-3. Furthermore the expression of granzyme B and IFN- γ was determined by intracellular staining. The concentration of IL-2 in the cell culture supernatants was measured by ELISA (human IL-2 ELISA set, BD OptEIA) following the instructions of the manufacturer.

10

Example 35

Ex vivo treatment of tumor samples with catumaxomab

The trifunctional TCB catumaxomab (Removab®) was obtained from Fresenius. The experimental conditions were similar as indicated above for FolR1-TCB. Briefly,

15 EpCAM positive tumor digests or malignant effusions were cultured in the presence or absence of catumaxomab at a concentration of 10 ng/ml for 24h. Analysis of CD8⁺ and CD4⁺ positive T cells (CD45⁺CD3⁺) was performed as described above.

Example 36

Killing assay

20

To determine the FolR1-TCB induced tumor cell killing, 3x10⁴ CFSE-labelled Skov3 cells were cocultured with tumor samples in the presence or absence of FolR1-TCB at a concentration of 2 nM for 24h in 96-well flat bottom cell culture plates. The E:T ratio (E: effector CD45⁺CD3⁺ cells; T: target FolR1⁺ cells from tumor and added Skov3 cells) was adjusted to 1:1 in each well and the cell number of the added tumor samples was calculated for each sample according to prior characterization by flow cytometry. Cell death of Skov3 cells was determined by flow cytometry by measuring activated caspase 3 and the live/dead marker Live/Dead-near-IR. The assay was performed in triplicates. The FolR1-TCB mediated killing was calculated according to the following equation: % of specific killing = 100 – [(% of Skov3 live cells in FolR1-TCB treated sample / % of Skov3 live cells in untreated sample) x 100].

30

To compare the FolR1-TCB-induced killing capacity of T-cells between tumor samples, and to exclude additional factors suppressing T-cell functionality, such as expression of

PD-L1 on the tumor cells, we exogenously added CFSE-labeled FolR1⁺ Skov3 cells to the tumor digests and adjusted the E:T ratio to 1:1, essentially as described above. We then measured the FolR1-TCB-induced killing of CFSE-labeled Skov3 cells, which allowed us to also include FolR1⁻ tumor samples into the analysis. As some tumors from the initial cohort could not be used to characterize TCB-mediated tumor cell killing due to a very low amount of effector cells, a separate cohort of 12 tumor digests and 5 malignant effusions from 15 non-small cell lung cancer (NSCLC) and two epithelial ovarian carcinoma (EOC) patients was analyzed. All samples were characterized for their CD3⁺ effector and FolR1⁺ target cell content (Fig. 39). Tumor cell killing of CD3⁺ T-cells from patients was compared with PBMC-derived T-cells from healthy donors. A substantial heterogeneity in tumor cell killing between individual patients was observed ($26 \pm 11.8\%$) after 24 h (Fig. 12O). Of note, CD3⁺ T-cells from healthy donors induced a significantly better killing than TILs ($42.8 \pm 9.7\%$, $p=0.013$). Exposure to a control TCB with no binding to a tumor antigen (DP47-TCB) did not induce any tumor cell killing.

15

Example 37

Polyclonal stimulation with anti-CD3/CD28 antibodies

A 96-well flat-bottom plate was precoated with 0.5 $\mu\text{g/ml}$ anti-CD3 ϵ (clone OKT3, Biolegend) for 2 hrs at 37°C. Afterwards, the antibody solution was removed and the plate washed extensively. Frozen tumor suspensions were thawed, washed and cultured at 3×10^5 cells/200 μL /well in complete medium with 2 $\mu\text{g/ml}$ anti-CD28 antibody (clone 28.2, eBioscience) for 24 hrs. After 24 hrs of incubation cells were collected, washed and analyzed by flow cytometry for expression of activation markers e.g. CD25 and T cell effector functions e.g. granzyme B and IFN- γ on CD8⁺ T cells. Supernatants were collected for IL-2, IFN- γ and TNF- α ELISA which was performed according to the manufacturer's instructions.

25

Example 38

Restoring of T cell function by PD-1 blockade

Tumor digests were stimulated by agonistic anti-CD3 and anti-CD28 antibodies as described above in the presence or absence of 10 $\mu\text{g/ml}$ anti-PD-1 antibody (MDX5C4) per well and incubated for 24 hrs. After 24 hrs cells were collected, washed and analyzed by flow cytometry. Supernatants were collected for IL-2, IFN- γ and TNF- α ELISA which was performed according to the manufacturer's instructions.

30

Example 39

Activation of T cells in tumor digests and malignant effusions by FolR1 TCB

The T cell bispecific antibodies engaging CD3 and folate receptor 1 (Mov19 based FolR1-TCB and the control antibody DP47-TCB were provided by Roche Glycart. The anti-PD-1 antibody 5C4 is described in US Pat. No. 8,008,449. The anti-Tim3 antibody F38-2EL was used. For flow cytometric characterization of FolR1 expression the antibody anti-FolR1-APC (aa25-233) from LifeSpanBiosciences and its matched isotype control (Biolegend) were used. Tumor lesions from 15 patients with FolR1⁺ tumors were characterized for T cell activation induced by FolR1 TCB. The samples consisted of 9 single cell suspensions and 6 malignant effusions derived from patients with NSCLC (n=7), ovarian cancer (n=7), and renal cell cancer (n=1). The amount of CD3⁺ T cells and of FolR1⁺ tumor cells was highly variable between patients (CD3⁺: mean 33.9% ± standard deviation of 16.6%, FolR1⁺: 17.1% ± 16.8%). Characterization of the expression of the inhibitory receptors PD-1, Tim-3, CTLA-4, Lag- and BTLA on T cells revealed a large heterogeneity among patients (Fig. 11A-B). While the tumor-infiltrating CD8⁺ T cells showed high levels of PD-1, Tim-3 and CTLA-4 (31.6% ± 25%; 22.2% ± 20.8% and 18.7% ± 14.4%, respectively), Lag-3 and BTLA were only expressed on a minority of cells in all patients of this cohort (3.5% ± 4.9% and 2.3% ± 1.7%, respectively). Inhibitory receptors on CD4⁺ T cells were distributed similarly, with a slightly more prominent expression of CTLA-4.

To determine FolR1-TCB induced T cell activation, tumor samples were cultured in the presence or absence of FolR1-TCB or the control TCB DP-47. Then, T cells were characterized by multicolor flow cytometry for expression of activation markers and T cell effector functions, as described above. Fig. 12A-O reveals a large heterogeneity in FolR1-TCB induced T cell activation between patients. In particular, while the vast majority of patients expressed CD69 already at baseline, upregulation of CD25, CD137, and ICOS, varying from 9 - 80%, 2.5 - 50% and 3.5 - 71%, respectively was observed. Acquisition of effector functions such as IFN- γ secretion, CD107 degranulation and expression of granzyme B was observed, ranging from 3.7 - 59%, a fold change of 1 - 7 or 1.3 - 64, respectively (Fig. 12A-I). The inhibitory receptors PD-1 and Tim-3 were further upregulated as a marker of activation upon FolR1-TCB treatment, irrespective of their baseline expression. Exposure to TCB DP-47 did not induce any T cell activation.

The upregulation of CD25 and ICOS induced by FolR1-TCB stimulation was significantly stronger in peripheral CD8⁺ T-cells from healthy donors than for tumor-derived CD8⁺ cells ($p=0.002$ and $p<0.001$, respectively; Fig. 12J, Fig. 12L, Fig. 12M). The secretion of T-cell effector cytokines IFN- γ , IL-2, and TNF upon FolR1-TCB stimulation was largely diminished amongst TILs in the majority of tumors compared with PBMCs from healthy donors ($p=0.0047$, $p<0.001$, and $p=0.006$, respectively; Fig. 12N). FolR1-TCB-induced perforin secretion was highly variable in TILs, and severely impaired in a subset of patients (Fig. 12N).

Similarly, despite a lower upregulation of granzyme B, FolR1-TCB induced activation and acquisition of effector functions of CD4⁺ T cells (Fig. 25A-I). To assess whether the abundance of intra-tumoral T cells or FolR1 expression impacts on T cell activation upon TCB exposure, the upregulation of activation markers was correlated to the E:T ratio (E: effector CD45⁺CD3⁺ T cells; T: FolR1⁺ cells) and to the percentage and to the level of tumor antigen expression of FolR1⁺ cells (Fig. 13A-C). The latter was determined by the mean fluorescence intensity of FolR1 on tumor cells (CD45⁻EpCAM⁺) using flow cytometry (Fig. 13C). However, neither of these parameters did influence T cell activation, i.e., even low amounts of FolR1⁺ cells, high E:T ratios, or poor T-cell infiltration have been sufficient for an efficient upregulation of activation and functional markers. In addition, the presence of potentially immune-suppressive cell populations such as regulatory T-cells or immature myeloid cells did not influence T-cell activation or T-cell function.

Example 40

FolR1-TCB induced T cell activation inversely correlates

with expression of PD-1 and Tim-3

High expression of inhibitory receptors has been described as a hallmark of exhausted T cells. Therefore, a dysfunctional state of tumor-infiltrating T cells may impact efficacy of the FolR1 TCB and may be responsible, at least in part, for heterogeneous T cell activation upon TCB exposure. To this end, the co-expression of inhibitory receptors, as determined at baseline, was correlated to FolR1 TCB induced upregulation of activation markers and T cell effector functions. Both PD-1 and Tim-3 expression on CD8⁺ T cells thereby negatively correlated with T cell activation determined by expression of CD25, CD137 and ICOS. CD8⁺ T cells with a high expression of PD-1 or Tim-3 showed a

marginal effect upon FolR1-TCB treatment, while T cells with a low expression of these inhibitory receptors could be strongly activated upon treatment with FolR1-TCB (Fig. 14A-I). Measurement of FolR1-TCB induced IL-2 secretion normalized to the content of T cells in the samples revealed the same dependencies on PD-1 and Tim-3 expression (Fig. 15A-C), while FolR1-TCB induced upregulation of granzyme B was less dependent on prior expression of these inhibitory receptors (Fig. 14J-L). Interestingly, the baseline expression of CTLA-4, Lag-3 and BTLA on CD8⁺ T cells did not correlate with FolR1-TCB induced T cell activation (Fig. 26A-C). Expression of inhibitory receptors on CD4⁺ T cells was much less predictive for FolR1-TCB induced CD4⁺ T cell activation compared to the expression of the same receptors on CD8⁺ T cells.

Example 41

FolR1-TCB induced tumor cell killing inversely correlates with expression of PD-1 and Tim-3

To investigate FolR1-TCB induced killing of tumor cells at an adjusted E:T ratio of 1:1, CFSE-labelled Skov3 cells were exogenously added to the tumor digests which contain a previously determined amount of CD3⁺ T cells using multicolor flow cytometry. FolR1-TCB induced killing of Skov3 cells was determined by measuring activated caspase 3 and a live/dead marker. In line with the FolR1-TCB induced T cell activation as measured by CD25 up-regulation, the specific killing upon FolR1-TCB exposure negatively correlated with single or co-expression of PD-1 and Tim-3 on CD8⁺ T cells. Furthermore, FolR1-TCB induced killing was also influenced by the baseline expression of CTLA-4 and the co-expression of PD-1 and CTLA-4. However, the impact of CTLA-4 expression on FolR1-TCB induced tumor cell killing was less pronounced compared to PD-1 and Tim-3 expression.

Example 42

Treatment of fresh tumor lesions with catumaxomab -

Activation of tumor-infiltrating T cells using catumaxomab and correlation with expression of inhibitory receptors

To determine to which extent catumaxomab induces T cell activation and to confirm the findings described above using a second, independent T cell bispecific molecule, 4 tumor digests from patients with NSCLC were exposed to catumaxomab, a trifunctional bispecific antibody recognizing CD3 on T cells and EpCAM on tumor cells. Then, T cells

were characterized by flow cytometry for expression of activation markers and T cell effector functions (Fig. 17A-D). Validating our data above for FolR1-TCB, we observed a striking heterogeneity in catumaxomab induced T cell activation. Accordingly, the baseline expression of inhibitory receptors differed between the patients (Fig. 17E-H).

- 5 Analysis of T cell activation and effector function upon treatment with catumaxomab revealed two groups of patients according to PD-1 and/or Tim-3 expression on CD8⁺ T cells confirming our findings with FolR1-TCB (Fig. 18A-R). PD-1^{low}, Tim-3^{low}, and, even more pronounced, both PD-1^{low} /Tim-3^{low} expressing cells, failed to be activated by catumaxomab, whereas PD-1^{high}, Tim-3^{high}, and PD-1^{high} /Tim-3^{high} T cells substantially
10 upregulated CD25, CD69, CD137, ICOS, granzyme B and IFN- γ .

Example 43

Polyclonal stimulation of tumor-infiltrating T cells by CD3/CD28 -

Immune phenotyping of tumor-infiltrating T cell subsets in non-small cell lung

15 cancer samples

We investigated the expression of co-inhibitory T cell receptors and differentiation markers on tumor-infiltrating CD3⁺CD8⁺ and CD3⁺CD4⁺ T cell subsets from 34 patients NSCLC using multicolor flow cytometry. The majority of tumors showed a high expression of the inhibitory receptor PD-1 (Fig. 19A-B), a major regulator of T cell
20 exhaustion. Of note, expression of other checkpoint inhibitors such as Tim-3, CTLA-4, LAG-3 or BTLA showed substantial variation between T cells obtained from different tumors (Fig. 19A-B).

Example 44

Cumulative expression of inhibitory receptors defines T cell dysfunction

- 25 In this Example, polyclonal stimulation was used in a sub-optimal dose to assess the impact of inhibitory receptors on T cell function. The effect of stimulation with agonistic anti-CD3 and anti-CD28 antibodies on T cell activation, as exemplified by CD25 expression, and on T cell effector function as analyzed by IFN- γ , TNF- α and IL-2 production as well as granzyme B expression varied substantially between patients as
30 determined by flow cytometry (Figure 20A-B) and ELISA (Fig. 20C-E). Of note, we observed different levels of T cell function, varying from T cell populations that exhibit a largely preserved T cell function (i.e., sustained CD25 and granzyme B expression, as

well as IL-2, IFN- γ and TNF- α production) to those with abrogated T cell function (loss of CD25 and granzyme B expression and of cytokine production).

To analyze the impact of multiple inhibitory receptors on T cell functionality we defined the inhibitory receptor (iR) score as a marker for the cumulative expression of inhibitory
5 receptors on T cells. To this end, the percentage of expression of PD-1, Tim-3, CTLA-4, Lag-3 and BTLA was analysed in all NSCLC samples and a score based on the median and interquartile ranges of each expressed receptor was defined and calculated for each sample (e.g., Fig. 21F). Tumor-infiltrating CD8⁺ T cells expressing a high iR score indicating expression of multiple inhibitory receptors showed a marginal effect upon
10 polyclonal stimulation, correlating with their highly dysfunctional state, whereas T cells with a low iR score could be strongly activated upon polyclonal stimulation (Fig. 21A-E). Upregulation of T cell effector functions, indicated by IL-2, IFN- γ and TNF- α production, not only correlated with the cumulative expression of inhibitory receptors but similarly with PD-1 and Tim-3 expression as well with the co-expression of both receptors (Fig.
15 22A-I), indicating a significant contribution of PD-1 and Tim-3 to T cell dysfunction.

Example 45

Inhibitory Receptor Expression

Single and cumulative expression of inhibitory receptors increases with tumor
20 progression. The expression of inhibitory receptors correlated with tumor stage and tumor progression. The number of PD-1, Tim-3 and LAG-3 positive cells was clearly increased in advanced tumor stages (Fig. 21G-K). No clear correlation was observed for the expression of CTLA-4, which may indicate that this receptor acts via a different inhibitory mechanism. BTLA was generally expressed at a low level and only a small
25 increase was found in advanced tumor stages (Fig. 21K). A significant increase in the cumulative expression of inhibitory receptors, as reflected by the *iR score*, was observed in patients with nodal positive cancers and advanced tumor stages whereas primary tumor size did not significantly correlate with the *iR score* (Fig. 21L-M). These data suggest a gradual and continuous upregulation of inhibitory receptors, during tumor progression,
30 which are most likely involved in T cell exhaustion in NSCLC.

Inhibitory receptors are gradually expressed on tumor-infiltrating T cells. To explore the role of simultaneous expression of distinct inhibitory receptors on single T cells, the concomitant expression of these receptors in CD8⁺ T cells (Figs. 32, 33) relative

to the expression of any of the five analyzed receptors was analyzed. Expression is shown as heat map, displaying the percentage of expression for the individual patients (Fig. 32) or as a radar plot, which shows the expression as mean and standard deviation of the four respective receptors on CD8⁺ T cells, pregated for the fifth, indicated immune checkpoint (Fig. 33). CD8⁺PD-1⁺ T cells on average expressed the lowest percentages of other inhibitory receptors, whereas CD8⁺BTLA⁺ T cells expressed all of the four other inhibitory receptors at high levels, indicating that BTLA marks a particularly exhausted T cell subset (Figs. 32, 33). An increase in the number of co-expressed inhibitory receptors was observed from CD8⁺Tim-3⁺ T cells over CD8⁺CTLA-4⁺ T cells to CD8⁺LAG-3⁺ T cells (Figs. 32, 33). These findings suggest a gradual acquisition of inhibitory receptors with PD-1 as a broadly expressed, early marker, while BTLA is upregulated rather late during T cell exhaustion.

Example 46

Blockade of PD-1 can partially restore T cell function

Rescue of T cell function by PD-1 blocking antibodies depends on the level of PD-1 expression. As we found a clear correlation between the expression of inhibitory receptors, particularly PD-1 and Tim-3, and T cell activation upon polyclonal stimulation, blockade of the PD-1 or PD-1/Tim-3 pathways might restore T cell function. However, addition of a blocking antibody to PD-1 (5C4) or combined blockade of PD-1 and Tim-3 upon stimulation with agonistic anti-CD3 and anti-CD28 antibodies could restore T cell effector function such as production and secretion of IL-2, IFN- γ and TNF- α only in some patients whereas in other patients only a marginal effect was seen (Fig. 23A-D). As observed in a chronic murine LCMV infection model (Blackburn et al., PNAS 105(39):15016 (2008)), we identified PD-1^{hi} and PD-1^{int} subsets in tumor-infiltrating CD8⁺ T cells from NSCLC patients. In brief, PD-1^{hi}, PD-1^{int}, and PD-1^{neg} subsets could be identified based on their measured fluorescence intensity. Cells from 33 patients were analysed for PD-1 expression to define uniform parameters for reproducible discernment of the three subsets. The analysis covered the whole spectrum of PD-1 expression levels and included tumor samples with clearly distinguished PD-1^{neg} or PD-1^{hi} populations. This allowed to set the gates for this analysis, which was then applied to all samples. Only PD-1^{int} expressing T cell subsets appeared to be rescued in activation upon PD-1 or combined PD-1/Tim-3 blockade, while no effect in T cell activation was observed upon

blockade in PD-1^{hi} cells (Fig. 24). The latter may exhibit a more exhausted phenotype which appears to be resistant to PD-1 blockade alone.

This finding was confirmed in T cells activated by FolR1 TCB. T cells were stimulated with FolR1 as described above. Blockade of PD-1 further strengthened FolR1-TCB induced T cell activation of T cells from a subset of patients.

Measurement of FolR1-TCB induced IFN- γ , TNF and IL-2 secretion normalized to the content of T cells in the samples revealed that in patient cell populations with a substantial amount of PD-1^{hi} expressing (approximately > 15%) cells were not able to secrete these cytokines. In contrast, cytokine secretion could be induced in most patient cell populations with a lower amount of PD-1^{hi} expressing (approximately < 15%) cells (Figure 27A-C). In the latter group, addition of a blocking antibody to PD-1 or combined blockade of PD-1 and Tim-3 upon stimulation by FolR1-TCB stimulation increased production of IL-2, IFN- γ and TNF- α (Figure 28A-F). The PD-1^{hi} expressing subset therefore may exhibit a more exhausted phenotype which appears to be resistant to PD-1 blockade alone.

Thus, T cell effector functions such as production of IL-2, IFN- γ and TNF- α could be restored in TILs from some NSCLC patients, whereas in other patients only a marginal recovery of T cell functions could be achieved. The increase in cytokine production upon exposure to anti-CD3/CD28 stimulation in combination with the PD-1 blocking antibody was compared to the percentage of PD-1^{hi} CD8⁺ T cells from the PD-1 positive population per patient. The increase in cytokine expression upon PD-1 blockade inversely correlated with the percentage of PD-1^{hi} T cells, indicating that patients expressing larger numbers of PD-1^{hi} T cells respond poorly to PD-1 blockade alone (FIG. 24A-C). As T cell dysfunction correlates with the expression of multiple inhibitory receptors (i.e., patients with a high *iR score*) and response to a PD-1 directed therapy correlates with the expression levels of PD-1 on CD8⁺ T cells, we further analyzed the expression of Tim-3, CTLA-4, LAG-3 and BTLA in PD-1^{hi} and PD-1^{int} CD8⁺ T cells. Remarkably, PD-1^{hi} T cells expressed significantly higher levels of additional receptors compared to PD-1^{int} subsets (Fig. 34). Thus, PD-1^{hi} and PD-1^{int} may identify two distinct T cell populations where PD-1^{hi} T cells may exhibit a more exhausted phenotype, which cannot be recovered by PD-1 blockade alone.

The data presented herein for the first time provides a comprehensive phenotypical and functional analysis of tumor-infiltrating CD8⁺ T cells from patients with NSCLC. The

data shows that these cells mainly possess an effector memory phenotype (CCR7-CD45RA^{low}) and show large heterogeneity in expression of inhibitory receptors such as PD-1, Tim-3, CTLA-4, LAG-3 and BTLA. Nevertheless, a clear increase in the number of receptors expressed on tumor-infiltrating lymphocytes (TILs) from late stage tumors was observed, which reflects the progress of T cell dysfunction during tumor development. The data presented herein shows that the effector functions of TILs were impaired in the vast majority of patients, and that impairment correlated with the expression of inhibitory receptors. To recover T cell function in a clinically relevant setting we combined polyclonal T cell stimulation with antibody-mediated inhibition of PD-1. The effect of PD-1 blockade on T cell functionality varied between TILs from different patients, but could be predicted by assessing the percentage of CD8⁺ T cells expressing PD-1 at high levels.

Here, we could demonstrate that the functionality of TILs can be correlated with and is largely affected by the number and expression level of inhibitory receptors. Of note, even T cells expressing low levels of inhibitory receptors showed some degree of impaired functionality, as the secretion of IL-2 was impaired in the vast majority of patients. Overall the activation and effector function of CD8⁺ T cells inversely correlated with the cumulative expression of inhibitory receptors, indicating a direct contribution of different inhibitory pathways to T cell dysfunction in NSCLC.

Our analysis of five inhibitory receptors on tumor infiltrating CD8⁺ T cells showed a clear increase of the single and cumulative expression of these inhibitory receptors in tumor tissues from NSCLC patients presenting with tumor-positive lymph nodes and advanced tumor stages. Expression of CTLA-4 differed from the other four receptors with the highest percentage of positive cells at early stages, which may indicate a distinct role of CTLA-4 in regulating T cell immunity (Topalian *et al.*, Safety, activity, and immune correlates of anti-PD-1 antibody in cancer. N. Engl. J. Med. 366, 2443 (Jun 28, 2012)). Co-expression analysis of additional inhibitory receptors on single cells, relative to the expression of one given receptor, showed a gradual expression, with early and late upregulation of PD-1 and BTLA, respectively. This may reflect the dynamic process of T cell exhaustion.

The findings presented herein underscore the clinical relevance of inhibitory receptor expression during NSCLC tumor progression, associated with progressive failure of immune control of tumor growth. We document here two populations of CD8⁺ tumor-

infiltrating T cells characterized by different levels of PD-1 expression (PD-1^{hi} and PD-1^{int} subsets). The occurrence of PD-1^{hi} T cells did not correlate with the percentage of PD-1 expression. Interestingly, we observed that the effect of PD-1 blockade correlated with the levels of PD-1 expression, with minimal effects on responsiveness of TILs with high proportions of PD-1^{hi} subpopulations. These findings are in line with experiments in a murine, chronic LCMV infection model where the subset of PD-1^{int} DbGP33-specific CD8⁺ T cells could be restored upon PD-1 blockade. In contrast, the PD-1^{hi} subset appeared more “exhausted,” i.e., exhibited signs of functional exhaustion, and responded poorly to PD-1 blockade. Thus, the level of PD-1 expression may represent a novel marker to define distinct T cell subsets in human cancers and, may serve as a predictor of responses to treatment with PD-1 blocking antibodies.

Example 47

Activation of T-cells from healthy donors and cancer patients by FolR1-TCB

To assess the effect of FolR1-TCB on T-cell activation peripheral blood mononuclear cells (PBMCs) from healthy donors were co-cultured with the FolR1⁺ ovarian cancer cell line Skov3 (Fig. 40A). Upon exposure to increasing concentrations of FolR1-TCB ranging from 0.6 pM to 2 nM for 24 h we observed a strong activation of CD8⁺ T-cells with upregulation of CD25, CD137, and ICOS. In addition, T-cells secreted IL-2, IFN- γ , and TNF. Exposure to DP47-TCB, a TCB directed against an irrelevant antigen, did not induce any T-cell activation (Fig. 40B and C).

Example 48

Inhibitory receptor expression is highly diverse in tumor-infiltrating CD8⁺ T-cells

As tumor-resident T-cells frequently display a highly dysfunctional phenotype, the observed heterogeneity in T-cell activation among different patients after FolR1-TCB stimulation may be due to an impaired TIL functionality. A hallmark of dysfunctional T-cells in both chronic viral infections and in tumors is the overexpression of inhibitory receptors. To this end, we determined the expression of the immune checkpoints PD-1, Tim-3, CTLA-4, Lag-3, and BTLA on tumor-infiltrating CD8⁺ T-cells in all tumor samples. We observed a high diversity in frequency and combined expression of these receptors amongst different tumors; PD-1 was found to be the most prominent inhibitory receptor with the highest percentage of expression (60.2 $\pm$ 30%), followed by Tim-3 (29.5 $\pm$ 24.4%), CTLA-4 (24.6 $\pm$ 17.6%), Lag-3 (7.0 $\pm$ 5.9%), and BTLA (3.9 $\pm$ 2.6%) (Fig. 35F). As described previously in a murine chronic viral infection model (Blackburn et al., Proc

Natl Acad Sci U S A 2008;105(39):15016-21) and, as shown herein, in human tumors, the PD-1⁺ population could be divided into a PD-1^{hi} and a PD-1^{int} expressing subpopulation (Fig. 35A). Analysis of additional inhibitory receptors expressed on these particular subsets showed a significantly higher expression of all other inhibitory receptors, including Tim-3, CTLA-4, Lag-3, and BTLA, in the PD-1^{hi} subpopulation as compared with the expression of these receptors in the PD-1^{int} and PD-1^{neg} subsets (Fig. 36A-D). Therefore, we used the percentage of PD-1^{hi} T-cells in the CD8⁺ subset as a surrogate marker for the cumulative expression of inhibitory receptors. The tumor samples were divided according to the frequency of PD-1^{hi} cells into two groups with high (PD-1^{hi} abundant tumors) and low frequencies of PD-1^{hi} expressing T-cells (PD-1^{hi} scarce tumors), respectively. A cut-off value of 30% PD-1^{hi} expression was chosen to separate the two groups. The percentage of PD-1^{hi} cells ranged from 39.1–60.5% in the PD-1^{hi} abundant (49.5 ± 7.9%) and from 2.65–19.5% in the PD-1^{hi} scarce group (8.4 ± 5.7%; Fig. 36E). The cut-off value was validated in a second cohort of 14 NSCLC and 2 ovarian cancer patients with a similar distribution in the frequency of PD-1^{hi} cells, where we observed comparable results upon polyclonal stimulation by anti-CD3/anti-CD28 antibodies (Fig. 39).

Example 49

FolR1-TCB-induced T-cell activation largely depends on

the level of PD-1 expression on CD8⁺ T-cells

We analyzed whether the expression of inhibitory receptors could be correlated with a diminished T-cell functionality upon FolR1-TCB treatment. Consistent with the results described in Example 41 above, FolR1-TCB-induced T-cell activation, as exemplified by CD25, CD137, and ICOS expression (p=0.028; p<0.001, and p=0.008, respectively), and T-cell effector functions, indicated by IFN-γ, IL-2, TNF, as well as perforin secretion, were significantly impaired in PD-1^{hi} abundant tumors compared with PD-1^{hi} scarce tumors (p=0.019; p=0.007; p=0.028, and p=0.029, respectively; Fig. 37A-G). Similarly, PD-1^{hi} abundant tumors displayed a significantly reduced cytotoxicity upon FolR1-TCB stimulation whereas a strong tumor cell killing could be observed in the majority of PD-1^{hi} scarce tumors (p=0.021; Fig. 37H).

Example 50**PD-1 blockade restores FolR1-TCB-induced T-cell
function only in PD-1^{hi} scarce tumors**

As the level of PD-1 expression on TILs correlates with the efficacy of FolR1-TCB, we
5 analyzed whether blockade of the PD-1/PD-L1 axis in combination with FolR1-TCB
treatment might be able to restore T-cell function. We found that upon combined
treatment with FolR1-TCB and the PD-1 blocking antibody nivolumab (MDX5C4)
secretion of the effector cytokines IFN- γ , TNF, and IL-2 as well as perforin could be
increased only in some of the PD-1^{hi} scarce tumors. In contrast, in PD-1^{hi} abundant
10 tumors PD-1 blockade failed to elicit any response (Fig. 38A-D). Of note, cytotoxic
tumor cell killing could neither be improved in T-cells from PD-1^{hi} scarce nor from PD-
1^{hi} abundant tumors by the additional PD-1 blockade (Fig. 38E).

The examples set forth herein describe the immuno-modulatory capacity of a CD3 x
FolR1-specific TCB in primary cancer lesions from patients with non-small cell lung
15 cancer (NSCLC), epithelial ovarian carcinoma (EOC) and renal cell carcinoma (RCC).
Compared with fully functional peripheral T-cells from healthy donors, we observed a
substantial heterogeneity in FolR1-TCB-induced tumor cell killing and T-cell activation
among different human tumor samples, resulting in partial or complete impairment of T-
cell function in the majority of patients. Comprehensive analysis of inhibitory receptor
20 expression on the cell surface of intratumoral T-cells revealed that the efficacy of T-cell
activation by FolR1-TCB inversely correlated with the expression levels of PD-1.
Patients with PD-1^{hi} abundant tumors displayed impaired T-cell activation and effector
function upon FolR1-TCB treatment. Additionally, these patients did not respond to PD-
1 blockade in contrast to their PD-1^{hi} scarce expressing counterparts. Thus, the bioactivity
25 of bispecific antibodies is considerably hampered by T-cell dysfunction, which is
orchestrated, at least in part, by the sustained and highly diverse expression of inhibitory
receptors.

We observed a strong upregulation of T-cell activation markers, effector cytokine
secretion and tumor cell killing upon FolR1-TCB stimulation in PBMCs from healthy
30 donors (Fig. 40). In stark contrast, however, T-cell effector functions largely varied and
were generally diminished in intratumoral T-cells. Particularly, killing capacity and
effector cytokine production was significantly lower in TILs with complete loss of IL-2
production and severely impaired TNF and IFN- γ secretion in the majority of tumors.

We documented the expression of the inhibitory receptors PD-1, Tim-3, CTLA-4, Lag-3, and BTLA on intratumoral CD8⁺ T-cells. PD-1 displayed the broadest expression of all analyzed inhibitory receptors. Observations from chronic murine LCMV infections by Blackburn suggest the presence of functionally distinct PD-1 positive T-cell subsets, which can be separated on the basis of MFI levels, using flow cytometry (Blackburn et al., PNAS 105(39):15016 (2008)). Of note, PD-1^{hi} T-cell subsets displayed a high co-expression of Tim-3 and CTLA-4 and to a lesser extent of Lag-3 and BTLA, while their PD-1^{int} counterparts expressed only low levels of other inhibitory receptors, comparable to PD-1^{neg} T-cells. The frequency of PD-1^{hi} CD8⁺ T-cells differed largely between patients and allowed us to discriminate between PD-1^{hi} abundant and scarce tumors. In contrast to patients with a PD-1^{hi} scarce phenotype, FolR1-TCB-mediated T-cell activation and tumor cell killing was significantly impaired in tumors displaying a PD-1^{hi} abundant phenotype. These data extend and confirm previous observations that the activation and effector function of CD8⁺ T-cells correlates with the co-expression of multiple immune checkpoints (Sakuishi et al., J Exp Med 2010;207(10):2187-94; Fourcade et al., J Exp Med 2010;207(10):2175-86; Grosso et al., J Immunol 2009;182(11):6659-69; Matsuzaki et al., Proc Natl Acad Sci U S A 2010;107(17):7875-80; Fourcade et al., Cancer Res 2012;72(4):887-96). The frequency of PD-1^{hi} T-cells may therefore be useful as a surrogate marker for the functionality of TILs upon TCB activation as well as serve as a predictive marker for the therapeutic responses to TCB treatment. This immune profile could guide the selection of patients who are likely to respond to immunotherapy such as TCBs. Its correlation with clinical benefits remains to be determined in prospective clinical interventions.

A promising avenue to improve the therapeutic efficacy of TCBs lies in the blockade of inhibitory signals on T-cells. As PD-1 was the most prominently expressed inhibitory receptor in all tumors analyzed we assessed whether PD-1 blockade could enhance T-cell effector functions upon TCB activation. Of note, we observed increased secretion of effector cytokines upon combined FolR1-TCB and anti-PD-1 treatment, though only in PD-1^{hi} scarce tumors. Thus, novel therapeutic strategies, exploring the transformation of PD-1^{hi} into PD-1^{int} T-cells to increase the susceptibility to PD-1/PD-L1 blockade, are clearly needed.

Remarkably, we observed no improvement on tumor cell killing upon concomitant PD-1 blockade in all of the tumor samples. Thus, blockade of a single immune checkpoint may

not be sufficient to restore the cytolytic capacity of TILs. In a mouse tumor model, however, blockade of the PD-1/PD-L1 axis has been shown to increase T-cell infiltration into tumors (Curran et al., Proc Natl Acad Sci U S A 2010;107(9):4275-80), a characteristic of this treatment, which could not be addressed by our *in vitro* approach.

- 5 Thus, the therapeutic effect of PD-1 blockade *in vivo* might not only result from improving T-cell cytotoxicity of residual intratumoral T-cells, but from the sustained functionality of newly infiltrating T-cells. TCB-induced T-cell activation has been shown to upregulate PD-1 expression, which may lead to secondary resistance in the presence of PD-L1 expressed on both tumor cells and infiltrating immune cells as recently
- 10 demonstrated both with a Her2-specific TCB and with a carcinoembryonic antigen- (CEA) specific TCB (Junttila et al., Cancer Res 2014;74(19):5561-71; Osada et al., Cancer Immunol Immunother 2015). Importantly, blockade of the PD-1/PD-L1 axis could completely restore TCB-induced T-cell function both *in vitro* and in a mouse tumor model. These observations indicate that co-administration of checkpoint inhibitors is
- 15 capable of preventing secondary resistance, which may add to the dysfunctional state of TILs and limit the therapeutic efficacy of TCBs. Further work is clearly needed to determine optimal combination regimens of checkpoint inhibitors and TCBs. It will also be crucial to identify inhibitory and activating T-cell-receptors with non-redundant functions as potential therapeutic targets.
- 20 Our findings clearly indicate that bispecific antibodies such as FolR1-TCB are capable of causing T-cells to upregulate co-stimulatory molecules, produce inflammatory cytokines, and acquire cytolytic function. We have observed different states of T-cell dysfunction, which are orchestrated, at least in part, by the expression of inhibitory receptors and, in some instances, reduce the effectiveness of the TCB. As FolR1-TCB-induced effector
- 25 functions could only be partially restored by PD-1 blockade, our results suggest a rather complex immune regulation, which utilizes multiple and eventually non-redundant pathways to maintain T-cell dysfunction within the tumor environment.

Sequences

Amino acid sequences of exemplary embodiments

- 5 1) FolR binders useful in common light chain format, variable heavy chain

Description	Sequence	Seq ID No
16A3	QVQLVQSGAEVKKPGASVKVSCKASGYTFTSYMHWVRQAPGGGLE WMGIINPSGGSTSYAQKFQGRVTMTRDTSTSTVYMELSSLRSED VYYCARNYYAGVTPFDYWGGTGLTVSS	1
18D3	QVQLVQSGAEVKKPGASVKVSCKASGYTFTSYMHWVRQAPGGGLE WMGIINPSGGSTSYAQKFQGRVTMTRDTSTSTVYMELSSLRSED VYYCARNYYTGGSSAFDYWGQGLTVSS	2
15H7	QVQLVQSGAEVKKPGASVKVSCKASGYTFTSYMHWVRQAPGGGLE WMGIINPSGGSTSYAQKFQGRVTMTRDTSTSTVYMELSSLRSED VYYCARNYYLFTSFQDYWGQGLTVSS	3
15B6	QVQLVQSGAEVKKPGASVKVSCKASGYTFTSYMHWVRQAPGGGLE WMGIINPSGGSTSYAQKFQGRVTMTRDTSTSTVYMELSSLRSED VYYCARNYYIGIVPFDYWGGTGLTVSS	4
21D1	QVQLVQSGAEVKKPGASVKVSCKASGYTFTSYMHWVRQAPGGGLE WMGIINPSGGSTSYAQKFQGRVTMTRDTSTSTVYMELSSLRSED VYYCARNYYVGVSPFDYWGGTGLTVSS	5
16F12	QVQLVQSGAEVKKPGASVKVSCKASGYTFTSYMHWVRQAPGGGLE WMGIINPSGGSTSYAQKFQGRVTMTRDTSTSTVYMELSSLRSED VYYCARNFTVLRVPFDYWGGTGLTVSS	6
15A1	QVQLVQSGAEVKKPGASVKVSCKASGYTFTSYMHWVRQAPGGGLE WMGIINPSGGSTSYAQKFQGRVTMTRDTSTSTVYMELSSLRSED VYYCARNYYIGVVTFDYWGGTGLTVSS	7
15A1_CDR 1	SYMH	8
15A1_CDR 2	IINPSGGSTSYAQKFQG	9
15A1_CDR 3	NYYIGVVTFDY	10
19E5	QVQLVQSGAEVKKPGASVKVSCKASGYTFTSYMHWVRQAPGGGLE WMGIINPSGGSTSYAQKFQGRVTMTRDTSTSTVYMELSSLRSED VYYCARGEWRRYTSFDYWGGTGLTVSS	11
19E5_CDR 1	SYMH	8
19E5_CDR 2	IINPSGGSTSYAQKFQG	9
19E5_CDR 3	GEWRRYTSFDY	12
19A4	QVQLVQSGAEVKKPGASVKVSCKASGYTFTSYMHWVRQAPGGGLE WMGIINPSGGSTSYAQKFQGRVTMTRDTSTSTVYMELSSLRSED VYYCARGGWIRWEHFDYWGGTGLTVSS	13

19A4_CDR 1	SYMH	8
19A4_CDR 2	IINPSGGSTSYAQKFQG	9
19A4_CDR 3	GGWIRWEHFDY	14
16D5	EVQLVESGGGLVKPGGSLRLSCAASGFTFSNAWMSWVRQAPGKGLE WVGRIKSKTDGGTTDYAAPVKGRFTISRDDSKNTLYLQMNSLKTED TAVYYCTTPWEWSWYDYWGQGTLTVSS	15
16D5_CDR 1	NAWMS	16
16D5_CDR 2	RIKSKTDGGTTDYAAPVKG	17
16D5_CDR 3	PWEWSWYDY	18
15E12	EVQLVESGGGLVKPGGSLRLSCAASGFTFSNAWMSWVRQAPGKGLE WVGRIKSKTDGGTTDYAAPVKGRFTISRDDSKNTLYLQMNSLKTED TAVYYCTTPWEWSYFDYWGQGTLTVSS	19
15E12_CD R1	NAWMS	16
15E12_CD R2	RIKSKTDGGTTDYAAPVKG	17
15E12_CD R3	PWEWSYFDY	20
21A5	EVQLVESGGGLVKPGGSLRLSCAASGFTFSNAWMSWVRQAPGKGLE WVGRIKSKTDGGTTDYAAPVKGRFTISRDDSKNTLYLQMNSLKTED TAVYYCTTPWEWAWFDYWGQGTLTVSS	21
21A5_CDR 1	NAWMS	16
21A5_CDR 2	RIKSKTDGGTTDYAAPVKG	17
21A5_CDR 3	PWEWAWFDY	22
21G8	EVQLVESGGGLVKPGGSLRLSCAASGFTFSNAWMSWVRQAPGKGLE WVGRIKSKTDGGTTDYAAPVKGRFTISRDDSKNTLYLQMNSLKTED TAVYYCTTPWEWAYFDYWGQGTLTVSS	23
21G8_CDR 1	NAWMS	16
21G8_CDR 2	RIKSKTDGGTTDYAAPVKG	17
21G8_CDR 3	PWEWAYFDY	24
19H3	QVQLVQSGAEVKKPGASVKVSKASGYTFTSYMHWVRQAPGGGLE WMGIINPSGGSTSYAQKFQGRVTMTRDTSTSTVYMELSSLRSEDTA	25

-220-

	VYYCARTGWSRWGYMDYWGQGTLTVSS	
19H3_CDR 1	SYMH	8
19H3_CDR 2	IINPSGGSTSYAQKFQG	9
19H3_CDR 3	TGWSRWGYMDY	26
20G6	QVQLVQSGAEVKKPGASVKVSCASGYTFTSYMHVVRQAPGGGLE WMGIINPSGGSTSYAQKFQGRVTMTRDTSTSTVYMELSSLRSED TA VYYCARGEWIRYYHFDYWGQGTLTVSS	27
20G6_CDR 1	SYMH	8
20G6_CDR 2	IINPSGGSTSYAQKFQG	9
20G6_CDR 3	GEWIRYYHFDY	28
20H7	QVQLVQSGAEVKKPGASVKVSCASGYTFTSYMHVVRQAPGGGLE WMGIINPSGGSTSYAQKFQGRVTMTRDTSTSTVYMELSSLRSED TA VYYCARVGWYRWGYMDYWGQGTLTVSS	29
20H7_CDR 1	SYMH	8
20H7_CDR 2	IINPSGGSTSYAQKFQG	9
20H7_CDR 3	VGWYRWGYMDY	30

2) CD3 binder common light chain (CLC)

Description	Sequence	Seq ID No
common CD3 light chain (VL)	QAVVTQEPSTLVSPGGTVTLTCGSSTGAVTTSNYANWVQEK P GQAFRGLIGGTNKRAPGTPARFSGSLLGGKAALTLSGAQPED EAEYYCALWYSNLWVFGGGTKLTVL	31
common CD3 light chain_CDR 1	GSSTGAVTTSNYAN	32
common CD3 light chain_CDR 2	GTNKRAP	33
common CD3 light chain_CDR 3	ALWYSNLWV	34

-221-

common CD3 light chain (VLCL)	QAVVTQEPSTLVSPGGTVTLTCGSSTGAVTTSNYANWVQEKP GQAFRGLIGGTNKRAPGTPARFSGSLLGGKAALTLSGAQPED EAEYYCALWYSNLWVFGGGTKLTVLGQPKAAPSVTLFPPSSE ELQANKATLVCLISDFYPGAVTVAWKADSSPVKAGVETTTPS KQSNNKYAASSYLSLTPEQWKSHRSYSCQVTHEGSTVEKTVA PTECS	35
--	---	----

3) CD3 binder, heavy chain

Description	Sequence	Seq ID No
CD3 variable heavy chain (VH)	EVQLLES GGGLVQPGGSLRLSCAASGFTFSTYAMNWVRQAPGKG LEWVSRIRSKYNNYATYYADSVKGRFTISRDDSKNTLYLQMNSL RAEDTAVYYCVRHGNFGNSYVSWFAYWGQGTLLTVSS	36
CD3 heavy chain (VH)_CDR 1	TYAMN	37
CD3 heavy chain (VH)_CDR 2	RIRSKYNNYATYYADSVKG	38
CD3 heavy chain (VH)_CDR 3	HGNFGNSYVSWFAY	39
CD3 full heavy chain (VHCH1)_	EVQLLES GGGLVQPGGSLRLSCAASGFTFSTYAMNWVRQAPGKG LEWVSRIRSKYNNYATYYADSVKGRFTISRDDSKNTLYLQMNSL RAEDTAVYYCVRHGNFGNSYVSWFAYWGQGTLLTVSSASTKGPS VFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVH TFPAVLQSSGLYSLSSVVTVPSSSLGTQTYICNVNHKPSNTKVD KKVEPKSC	40
CD3 constant heavy chain CH1	ASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSG ALTSGVHTFPAVLQSSGLYSLSSVVTVPSSSLGTQTYICNVNHK PSNTKVDKKVEPKSC	84

4) FolR binders useful for crossfab Format

Description	Sequence	Seq ID No
11F8_VH	QVQLVQSGAEVKKPGSSVKVSKASGGTFSSYAI SWVRQAPGQGLE WMGGIIPIFGTANYAQKFQGRVTITADKSTSTAYMELSSLRSEDTA VYYCARAVFYRAWYSFDYWGQGTITVTVSS	41

11F8_VH_CD R1	SYAIS	42
11F8_VH_CD R2	GIIPIFGTANYAQKFQG	43
11F8_VH_CD R3	AVFYRAWYSFDY	44
11F8_VL	DIQMTQSPSTLSASVGDRVTITCRASQSISSWLAWYQQKPGKAPKL LIYDASSLES ¹ GVPSRFSGSGSGTEFTLTIS ² SLQPDDFATYYCQQYT SPPPTFGQGTKVEIK	45
11F8_VL_CD R1	RASQSISSWLA	46
11F8_VL_CD R2	DASSLES	47
11F8_VL_CD R3	QQYTSPPT	48
36F2_VH	QVQLVQSGAEVKKPGASVKVSCKASGYTFTSY ³ YMHWVRQAPGQGLE WMGIINPSGGSTSYAQKFQGRVTMTHTDSTSTVYME ⁴ LSLRSEDTA VYYCARSFFTG ⁵ FHLDYWGQGT ⁶ LVTVSS	49
36F2_VH_CD R1	SYMH	8
36F2_VH_CD R2	IINPSGGSTSYAQKFQG	9
36F2_VH_CD R3	SFFTG ⁵ FHLDY	50
36F2_VL	EIVLTQSPGTLSLSPGERATLSCRASQSVSSSYLAWYQQKPGQAPR LLIYGASSRATGIPDRFSGSGSGTDFTLTISRLEPEDFAVYYCQQY TNEHY ⁷ YTFGQGTKVEIK	51
36F2_VL_CD R1	RASQSVSSSYLA	52
36F2_VL_CD R2	GASSRAT	53
36F2_VL_CD R3	QQYTNEHY ⁷ T	54
9D11_VH	QVQLVQSGAEVKKPGASVKVSCKASGYTFTSY ³ YMHWVRQAPGQGLE WMGIINPSGGPTSYAQKFQGRVTMT ⁸ TRDTSTSTVYME ⁴ LSLRSEDTA VYYCARGDFAWLDYWGQGT ⁶ LVTVSS	55
9D11_VH_CD R1	SYMH	8
9D11_VH_CD R2	IINPSGGPTSYAQKFQG	56
9D11_VH_CD R3	GDFAWLDY	57
9D11_VL	DIVMTQSP ⁹ LSLPVTPGEPASISCRSSQSL ¹⁰ LHSNGYNYLDWYLQKPG QSPQLLIYLG ¹¹ SNRASGV ¹² PDRFSGSGSGTDFTLKISRVEAEDVGVYY	58

-223-

	CMQASIMNRTFGQGTKVEIK	
9D11_VL_CD R1	RSSQSLLSHNGYNYLD	59
9D11_VL_CD R2	LGSNRAS	60
9D11_VL_CD R3	MQASIMNRT	61
9D11_VL N95S	DIVMTQSPLSLPVTTPGEPASISCRSSQSLLSHNGYNYLDWYLQKPG QSPQLLIYLGSNRASGVPDRFSGSGSGTDFTLKISRVEAEDVGVYY CMQASIMSRTFGQGTKVEIK	62
9D11_VL N95S_CDR3	MQASIMSRT	63
9D11_VL N95Q	DIVMTQSPLSLPVTTPGEPASISCRSSQSLLSHNGYNYLDWYLQKPG QSPQLLIYLGSNRASGVPDRFSGSGSGTDFTLKISRVEAEDVGVYY CMQASIMQRTFGQGTKVEIK	64
9D11_VL N95Q_CDR3	MQASIMQRT	65
9D11_VL T97A	DIVMTQSPLSLPVTTPGEPASISCRSSQSLLSHNGYNYLDWYLQKPG QSPQLLIYLGSNRASGVPDRFSGSGSGTDFTLKISRVEAEDVGVYY CMQASIMNRAFGQGTKVEIK	66
9D11_VL T97A	MQASIMNRA	67
9D11_VL T97N	DIVMTQSPLSLPVTTPGEPASISCRSSQSLLSHNGYNYLDWYLQKPG QSPQLLIYLGSNRASGVPDRFSGSGSGTDFTLKISRVEAEDVGVYY CMQASIMNRNFGQGTKVEIK	68
9D11_VL T97N_CDR3	MQASIMNRN	69
5D9_VH	QVQLVQSGAEVKKPGASVKVSCKASGYTFTSYIMHWVRQAPGQGLE WMGIINPSGGSTSYAQKFQGRVTMTRDTSTSTVYMELSSLRSEDTA VYYCARSYIDMDYWGQGLTVTVSS	70
5D9_VH_CDR 1	SYIMH	8
5D9_VH_CDR 2	IINPSGGSTSYAQKFQG	9
5D9_VH_CDR 3	SYIDMDY	71
5D9_VL	EIVLTQSPGTLSPGERATLSCRASQSVSSSYLAWYQQKPGQAPR LLIYGASSRATGIPDRFSGSGSGTDFTLTISRLEPEDFAVYYCQQD NWSPTFGQGTKVEIK	72
5D9_VL_CDR 1	RASQSVSSSYLA	52
5D9_VL_CDR 2	GASSRAT	53
5D9_VL_CDR 3	QQDNWSPT	73

-224-

6B6_VH	QVQLVQSGAEVKKPGASVKVSCKASGYTFTSYMHVWRQAPGQGLE WMGIINPSSGGSTSYAQKFQGRVTMTTRDTSTSTVYMELSSLRSEDTA VYYCARSYVDMDYWGQGTLVTVSS	74
6B6_VH_CDR 1	SYMH	8
6B6_VH_CDR 2	IINPSSGGSTSYAQKFQG	9
6B6_VH_CDR 3	SYVDMDY	75
6B6_VL	EIVLTQSPGTLSPGERATLSCRASQSVSSSYLAWYQQKPGQAPR LLIYGASSRATGIPDRFSGSGSGTDFTLTISRLEPEDFAVYYCQQD IWSPTFGQGTKVEIK	76
6B6_VL_CDR 1	RASQSVSSSYLA	52
6B6_VL_CDR 2	GASSRAT	53
6B6_VL_CDR 3	QQDIWSPT	77
14E4_VH	EVQLLES GGGLVQPGGSLRLSCAASGFTFSSYAMSWVRQAPGKGLE WVSAISGSGGSTYYADSVKGRFTISRDN SKNTLYLQMNSLRAEDTA VYYCAKDSSYVEWYAFDYWGQGTLVTVSS	78
14E4_VH_CD R1	SYAMS	79
14E4_VH_CD R2	AISGSGGSTYYADSVKG	80
14E4_VH_CD R3	DSSYVEWYAFDY	81
14E4_VL	EIVLTQSPGTLSPGERATLSCRASQSVSSSYLAWYQQKPGQAPR LLIYGASSRATGIPDRFSGSGSGTDSTLTISRLEPEDFAVYYCQQP TSSPITFG QGTKVEIK	82
14E4_VL_CD R1	RASQSVSSSYLA	52
14E4_VL_CD R2	GASSRAT	53
14E4_VL_CD R3	QQPTSSPIT	83

5) CD3 binder useful in crossfab Format

Description	Sequence	Seq ID No
CD3 heavy chain (VH)	EVQLLES GGGLVQPGGSLRLSCAASGFTFSTYAMN WVRQAPG KGLEWVSRIRSKYNNYATYYADSVKGRFTISRDD SKNTLYLQ MNSLRAEDTAVYYCVRHGNFGNSYVSWFAYWGQGTLVTVSS	36
CD3 heavy	TYAMN	37

chain (VH)_CDR 1		
CD3 heavy chain (VH)_CDR 2	RIRSKYNNYATYYADSVKG	38
CD3 heavy chain (VH)_CDR 3	HGNFGNSYVSWFAY	39
CD3 light chain (VL)	QAVVTQEP SLTVSPGGTVTLTCGSSTGAVTTSNYANWVQEK GQAFRGLIGGTNKRAPGTPARFSGSLLGGKAALTLGAQPED EAEYYCALWYSNLWVFGGGTKLTVL	31
CD3 light chain_CDR 1	GSSTGAVTTSNYAN	32
CD3 light chain_CDR 2	GTNKRAP	33
CD3 light chain_CDR 3	ALWYSNLWV	34
pETR12940: crossed common CD3 light chain (VLCH1)	QAVVTQEP SLTVSPGGTVTLTCGSSTGAVTTSNYANWVQEK GQAFRGLIGGTNKRAPGTPARFSGSLLGGKAALTLGAQPED EAEYYCALWYSNLWVFGGGTKLTVLSSASTKGPSVFPLAPSS KSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVL QSSGLYSLSSVTVPSSSLGTQTYICNVNHKPSNTKVDKKVE PKSC	86
Crossed CD3 heavy chain (VHCκ); e.g. in pCON1057	EVQLLES GGGVLVQPGGSLRLSCAASGFTFSTYAMNWVRQAPG KGLEWVSRIRSKYNNYATYYADSVKGRFTISRDDSKNTLYLQ MNSLRAEDTAVYYCVRHGNFGNSYVSWFAYWGQGLTVTVSSA SVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQWKV DNALQSGNSQESVTEQDSKDYSLSSLTLSKADYEKHKVY ACEVTHQGLSSPVTKSFNRGEC	87
CD3- CH1	ASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWN SGALTSGVHTFPAVLQSSGLYSLSSVTVPSSSLGTQTYICN VNHKPSNTKVDKKVEPKSC	85
CD3- ckappa	VAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQWKVD NALQSGNSQESVTEQDSKDYSLSSLTLSKADYEKHKVYA CEVTHQGLSSPVTKSFNRGEC	88

6) Exemplary amino acid sequences of CD3-FcR bispecific antibodies 2+1 inverted crossmab format

Description	Sequence	Seq ID No
VHCH1[9D11]_VHCL[CD3]_Fcknob_PGLALA pCON1057	QVQLVQSGAEVKKPGASVKVSCKASGYTFTSYMHWVRQAPGQGLEWMGIINPSSGGPTSYAQKFQGRVTMTTRDTSTSTVYMELSSLRSEDTAVYYCARGDFAWLDYWGQGLTVTVSSASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVVTVPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDGGGGSGGGGSEVQLLESGLLVQPGGSLRLSCLASGFTFTSYAMNWVRQAPGKGLIEWVSRIRSKYNNYATYYADSVKGRFTISRDDSKNTLYLQMNSLRAEDTAVYYCVRHGNFGNSYVSWFAYWGQGLTVTVSSASVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQWKVDNALQSGNSQESVTEQDSKDSSTYSLSSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGECGCDKTHTCPPCPAPEAAGGPSVFLFPPKPKDTLMISRTPEVTCVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALGAPIEKTIISKAKGQPREPQVYTLPPCRDELTKNQVSLWCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGSFFFLYSKLTVDKSRWQQGNVFCSCVMHEALHNHYTQKSLSLSPGK	94
9D11_Fchole_PGLALA_HYRF	QVQLVQSGAEVKKPGASVKVSCKASGYTFTSYMHWVRQAPGQGLEWMGIINPSSGGPTSYAQKFQGRVTMTTRDTSTSTVYMELSSLRSEDTAVYYCARGDFAWLDYWGQGLTVTVSSASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVVTVPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHTCPPCPAPEAAGGPSVFLFPPKPKDTLMISRTPEVTCVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALGAPIEKTIISKAKGQPREPQVCTLPSPRDELTKNQVSLSCAVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGSFFFLVSKLTVDKSRWQQGNVFCSCVMHEALHNHYTQKSLSLSPGK	95
9D11_LC pCON1063	DIVMTQSPLSLPVTPGEPASISCRSSQSLHLSNGYNYLDWYLQKPGQSPQQLLIYLGSNRASGVDPDRFSGSGSDFTLKI SRVEAEDVGVYYCMQASIMNRTFGQGTKVEIKRTVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQWKVDNALQSGNSQESVTEQDSKDSSTYSLSSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC	96
VLCH1[CD3] pETR12940	QAVVTQEPSTLVSPGGTVTLTCGSSTGAVTTSNYANWVQEKPGQAFRGLIGGTNKRAPGTPTARFSGSLGGAALTLGAQPEDEAEYYCALWYSNLWVFGGGTKLTVLSSASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVVTVPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSC	86
CH1	ASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVVTVPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCD	428
VHCH1[36F2]_VHCL[CD3]_Fcknob_PGLALA pCON1056	QVQLVQSGAEVKKPGASVKVSCKASGYTFTSYMHWVRQAPGQGLEWMGIINPSSGGSTSYAQKFQGRVTMTHTDTSTSTVYMELSSLRSEDTAVYYCARSFFTGFHLDYWGQGLTVTVSSASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVVTVPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDGGGGSGGGSEVQLLESGLLVQPGGSLRLSCLASGFTFTSYAMNWVRQAPGKGLIEWVSRIRSKYNNYATYYADSVKGRFTISRDDSKNTLYLQMNSLR AEDTAVYYCVRHGNFGNSYVSWFAYWGQGLTVTVSSASVAAPSVFI FPPSDEQLKSGTASVVCLLNNFYPREAKVQWKVDNALQSGNSQESVTEQDSKDSSTYSLSSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGECGCDKTHTCPPCPAPEAAGGPSVFLFPPKPKDTLMISRTPEVTCVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLT	393

	VLHQDWLNGKEYKCKVSNKALGAPIEKTISKAKGQPREPQVYTLPP CRDELTKNQVSLWCLVKGFYPSDIAVEWESNGQPENNYKTTPVLDS SDGSFFLYSKLTVDKSRWQQGNVSCSVMHEALHNHYTQKSLSLSP GK	
36F2-Fc hole PGLALA pCON1050	QVQLVQSGAEVKKPGASVKVSCKASGYTFTSYMHWRQAPGQGLE WMGIINPSGGSTSYAQKFQGRVTMTHTSTSTVYMEISSLRSEDTA VYYCARSFFTGFHLDYWGQGTSLVTVSSASTKGPSVFPLAPSSKSTS GGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSL SSVVTVPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHTCPP CPAPEAAGGPSVFLLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVK FNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYK CKVSNKALGAPIEKTISKAKGQPREPQVCTLPSPRDELTKNQVSL CAVKGFYPSDIAVEWESNGQPENNYKTTPVLDSGDSFFLVSKLTV DKSRWQQGNVSCSVMHEALHNHYTQKSLSLSPGK	394
36F2 LC pCON1062	EIVLTQSPGTLSLSPGERATLSCRASQSVSSSYLAWYQQKPGQAPR LLIYGASSRATGIPDRFSGSGSGTDFTLTISRLEPEDFAVYYCQQY TNEHYTFTGQGTKVEIKRTVAAPSVFIFPPSDEQLKSGTASVVCLL NNFYPREAKVQWKVDNALQSGNSQESVTEQDSKSTYSLSSLTLS KADYEEKHKVYACEVTHXGLSSPVTKSFNRGEC	395
CD3 VLCH1 pETR12940	QAVVTQEPSTLVSPGGTVTLTCGSSTGAVTTSNYANWVQEKPGQAF RGLIGGTNKRAPGTPARFSGSLGGKAALTLGAQPEDEAEYYCAL WYSNLWVFGGGTKLTVLSSASTKGPSVFPLAPSSKSTSGGTAALGC LVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVVTVP SSSLGTQTYICNVNHKPSNTKVDKKVEPKSC	86

7) Exemplary amino acid sequences of CD3-FcR bispecific antibodies with common light chain

VHCH1[1 6D5]_VH CH1[CD3] _Fcknob pCON999	EVQLVESGGGLVKPGGSLRLSCAASGFTFSNAWMSWVRQAPGKG LEWVGRIKSKTDGGTTDYAAPVKGRFTISRDDSKNTLYLQMNSL KTEDTAVYYCTTPWEWSWYDYWGQGTSLVTVSSASTKGPSVFPLA PSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAV LQSSGLYSLSSVVTVPSSSLGTQTYICNVNHKPSNTKVDKKVEP KSCDGGGGSGGGSEVQLLESGLLVQPGGSLRLSCAASGFTFS TYAMNWVRQAPGKGLEWVSRIRSKYNNYATYYADSVKGRFTISR DDSKNTLYLQMNSLRAEDTAVYYCVRHGNFGNSYVSWFAYWGQ TLTVSSASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPV TVSWNSGALTSGVHTFPAVLQSSGLYSLSSVVTVPSSSLGTQTY ICNVNHKPSNTKVDKKVEPKSCDKTHTCPPCPAPEAAGGPSVFL FPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHN AKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALGA PIEKTISKAKGQPREPQVYTLPPCRDELTKNQVSLWCLVKGFYPS DIAVEWESNGQPENNYKTTPVLDSGDSFFLYSKLTVDKSRWQ QGNVSCSVMHEALHNHYTQKSLSLSPGK	89
VHCH1[1 6D5]_Fcho le pCON983	EVQLVESGGGLVKPGGSLRLSCAASGFTFSNAWMSWVRQAPGKG LEWVGRIKSKTDGGTTDYAAPVKGRFTISRDDSKNTLYLQMNSL KTEDTAVYYCTTPWEWSWYDYWGQGTSLVTVSSASTKGPSVFPLA PSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAV LQSSGLYSLSSVVTVPSSSLGTQTYICNVNHKPSNTKVDKKVEP KSCDKTHTCPPCPAPEAAGGPSVFLLFPPKPKDTLMISRTPEVTC VVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSV LTVLHQDWLNGKEYKCKVSNKALGAPIEKTISKAKGQPREPQVC TLPPSRDELTKNQVSLSCAVKGFYPSDIAVEWESNGQPENNYKT	90

	TPPVLDSDGSFFLVSKLTVDKSRWQQGNVFSCSVMEALHNRT QKSLSLSPGK	
CD3_com mon light chain pETR1319 7	QAVVTQEPSTLVSPGGTVTLTCGSSTGAVTTSNYANWVQEKPGQ AFRGLIGGTNKRAPGTPARFSGSLLGGKAALTLSGAQPEDEAEY YCALWYSNLWVFGGGTKLTVLGQPKAAPSVTLFPPSSEELQANK ATLVCLISDFYPGAVTVAWKADSSPVKAGVETTTTPSKQSNNKYA ASSYLSLTPEQWKSHRSYSCQVTHEGSTVEKTVAPTECS	35
VHCH1[C D3]_VHC H1[16D5] _Fcknob_P GLALA pETR1393 2	EVQLLESGGGLVQPGGSLRLSCAASGFTTFSTYAMNWVRQAPGKG LEWVSRIIRSKYNNYATYYADSVKGRFTISRDDSKNTLYLQMNSL RAEDTAVYYCVRHGNFGNSYVSWFAYWGQGTLLTVVSSASTKGPS VFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVH TFPAVLQSSGLYSLSSVVTVPSSSLGTQTYICNVNHKPSNTKVD KKVEPKSCDGGGGSGGGGSEVQLVESGGGLVKPGGSLRLSCAAS GFTFSNAWMSWVRQAPGKGLEWVGRIKSKTDGGTTDYAAPVKGR FTISRDDSKNTLYLQMNSLKTEDTAVYYCTTPWEWSWYDYWGQ TLTVVSSASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPV TVSWNSGALTSGVHTFPAVLQSSGLYSLSSVVTVPSSSLGTQTY ICNVNHKPSNTKVDKKVEPKSCDKTHTCPPCPAPEAAGGPSVFL FPPKPKDTLMISRTPEVTCVVDVSHEDPEVKFNWYVDGVEVHN AKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALGA PIEKTISKAKGQPREPQVYTLPPCRDELTKNQVSLWCLVKGFYP SDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQ QGNVFSCSVMEALHNHYTQKSLSLSPGK	91
CD3_Fckn ob_PGLAL A pETR1391 7	EVQLLESGGGLVQPGGSLRLSCAASGFTTFSTYAMNWVRQAPGKG LEWVSRIIRSKYNNYATYYADSVKGRFTISRDDSKNTLYLQMNSL RAEDTAVYYCVRHGNFGNSYVSWFAYWGQGTLLTVVSSASTKGPS VFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVH TFPAVLQSSGLYSLSSVVTVPSSSLGTQTYICNVNHKPSNTKVD KKVEPKSCDKTHTCPPCPAPEAAGGPSVFLFPPKPKDTLMISRT PEVTCVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTY RVVSVLTVLHQDWLNGKEYKCKVSNKALGAPIEKTISKAKGQPR EPQVYTLPPCRDELTKNQVSLWCLVKGFYPSPDIAVEWESNGQPE NNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVFSCSVMEAL HNHYTQKSLSLSPGK	92
Fc_hole_P GLALA_H YRF pETR1075 5	DKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVV DVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTV LHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVCTLP PSRDELTKNQVSLSCAVKGFYPSPDIAVEWESNGQPENNYKTTP VLDSDGSFFLVSKLTVDKSRWQQGNVFSCSVMEALHNRTQKS LSLSPGK	93
VHCL[CD 3]_Fcknob _PGLALA pETR1337 8	EVQLLESGGGLVQPGGSLRLSCAASGFTTFSTYAMNWVRQAPGKG LEWVSRIIRSKYNNYATYYADSVKGRFTISRDDSKNTLYLQMNSL RAEDTAVYYCVRHGNFGNSYVSWFAYWGQGTLLTVVSSASVAAPS VFIFPPSDEQLKSGTASVVCCLNNFYPREAKVQWKVDNALQSGN SQESVTEQDSKDYSLSSSTLTLSKADYEKHKVYACEVTHQGLS SPVTKSFNRGECDKTHTCPPCPAPEAAGGPSVFLFPPKPKDTLM ISRTPEVTCVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQY NSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALGAPIEKTISKAK GQPREPQVYTLPPCRDELTKNQVSLWCLVKGFYPSPDIAVEWESN GQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVFSCSV MEALHNHYTQKSLSLSPGK	98

16D5 inverted 2+1 with N100A in CDR H3 pETR1409 6	EVQLVESGGGLVKPGGSLRLSCAASGFTFSNAWMSWVRQAPGKG LEWVGRIKSKTDGGTTDYAAPVKGRFTISRDDSKNTLYLQMNSL KTEDTAVYYCTTPWEWSWYDYWGQGTTLTVSSASTKGPSVFPLA PSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAV LQSSGLYSLSSVVTVPSSSLGTQTYICNVNHKPSNTKVDKKVEP KSCDGGGSGGGGSEVQLLESGLLVQPGGSLRLSCAASGFTFS TYAMNWVRQAPGKGLEWVSRIRSKYNNYATYYADSVKGRFTISR DDSKNTLYLQMNSLRAEDTAVYYCVRHGNFGASYVSWFAYWGQG TLTVSSASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPV TVSWNSGALTSGVHTFPAVLQSSGLYSLSSVVTVPSSSLGTQTY ICNVNHKPSNTKVDKKVEPKSCDKTHTCPPCPAPEAAGGPSVFL FPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHN AKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALGA PIEKTISKAKGQPREPQVYTLPPCRDELTKNQVSLWCLVKGFYP SDIAVEWESNGQPENNYKTPPVLDSDGSFFLYSKLTVDKSRWQ QGNVFSCSVMHEALHNHYTQKSLSLSPGK	99
16D5 inverted 2+1 with S100aA in CDR H3 pETR1409 7	EVQLVESGGGLVKPGGSLRLSCAASGFTFSNAWMSWVRQAPGKG LEWVGRIKSKTDGGTTDYAAPVKGRFTISRDDSKNTLYLQMNSL KTEDTAVYYCTTPWEWSWYDYWGQGTTLTVSSASTKGPSVFPLA PSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAV LQSSGLYSLSSVVTVPSSSLGTQTYICNVNHKPSNTKVDKKVEP KSCDGGGSGGGGSEVQLLESGLLVQPGGSLRLSCAASGFTFS TYAMNWVRQAPGKGLEWVSRIRSKYNNYATYYADSVKGRFTISR DDSKNTLYLQMNSLRAEDTAVYYCVRHGNFGNAYVSWFAYWGQG TLTVSSASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPV TVSWNSGALTSGVHTFPAVLQSSGLYSLSSVVTVPSSSLGTQTY ICNVNHKPSNTKVDKKVEPKSCDKTHTCPPCPAPEAAGGPSVFL FPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHN AKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALGA PIEKTISKAKGQPREPQVYTLPPCRDELTKNQVSLWCLVKGFYP SDIAVEWESNGQPENNYKTPPVLDSDGSFFLYSKLTVDKSRWQ QGNVFSCSVMHEALHNHYTQKSLSLSPGK	100
CD3 light chain fused to CH1; Fc _γ PGLAL A; pETR1386 2	QAVVTQEPSTLVSPGGTVTLTCGSSTGAVTTSNYANWVQEKPGQ AFRGLIGGTNKRAPGTPARFSGSLLGGKAALTLSGAQPEDEAEY YCALWYSNLWVFGGGTKLTVLSSASTKGPSVFPLAPSSKSTSGG TAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSL SSVVTVPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHTC PPCPAPEAAGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHED PEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWL NGKEYKCKVSNKALGAPIEKTISKAKGQPREPQVYTLPPSRDEL TKNQVSLTCLVKGFYPSTIAVEWESNGQPENNYKTPPVLDSDG SFFLYSKLTVDKSRWQQGNVFSCSVMHEALHNHYTQKSLSLSPG K	101
16D5 VH fused to constant kappa chain; pETR1385 9	EVQLVESGGGLVKPGGSLRLSCAASGFTFSNAWMSWVRQAPGKG LEWVGRIKSKTDGGTTDYAAPVKGRFTISRDDSKNTLYLQMNSL KTEDTAVYYCTTPWEWSWYDYWGQGTTLTVSSASVAAPSVFIFP PSDEQLKSGTASVVCLLNNFYPREAKVQWKVDNALQSGNSQESV TEQDSKDSSTYSLSSTLTLSKADYEKHKVYACEVTHQGLSSPVTK SFNRGEC	102
CD3 VH	EVQLLESGLLVQPGGSLRLSCAASGFTFSTYAMNWVRQAPGKG LEWVSRIRSKYNNYATYYADSVKGRFTISRDDSKNTLYLQMNSL	103

fused to constant lambda chain; pETR13860	RAEDTAVYYCVRHGNFGNSYVSWFAYWGQGLTVTVSSASPKAAP SVTLFPSPSEELQANKATLVCLISDFYPGAVTVAWKADSSPVKA GVETTTSPSKQSNNKYAASSYLSLTPEQWKSHRSYSCQVTHEGST VEKTVAPTECS	
IGHV1-46*01 (X92343), plus JH4 element	QVQLVQSGAEVKKPGASVKVSCKASGYTFTSYMH WVRQAPGQGLEWMGIINPSGGSTSYAQKFQGRVTM TRDTSTSTVYMELSSLRSEDNAVYYCARGGSGGSFD YWGQGLTVTVSS	104
IGHV1-69*06 (L22583), plus JH6 element	QVQLVQSGAEVKKPGSSVKVSCKASGGTFSSYAISW VRQAPGQGLEWMGGIPIFGTANYAQKFQGRVTITA DKSTSTAYMELSSLRSEDNAVYYCARGGSGGSMDA WGQGTTVTVSS	105
IGHV3-15*01 (X92216), plus JH4 element	EVQLVESGGGLVKPGGSLRLSCAASGFTFSNAWMS WVRQAPGKGLEWVGRIKSKTDGGTTDYAAPVKGRF TISRDDSKNTLYLQMNSLKTEDNAVYYCTTGSGGS FDYWGQGLTVTVSS	106
IGHV3-23*01 (M99660), plus JH4 element	EVQLLES GGGLVQPGGSLRLSCAASGFTFSSYAMSW VRQAPGKGLEWVSAISGSGGSTYYADSVKGRFTISR DNSKNTLYLQMNSLRAEDNAVYYCAKGGSGGSFDY WGQGLTVTVSS	107
IGHV4-59*01 (AB019438), plus JH4 element	QVQLQESGPGLVKPSETLSLTCTVSGGSISSYYWSWI RQPPGKGLEWIGYIYYSGSTNYNPSLKSRVTISVDT KNQFSLKLSSVTAADNAVYYCARGGSGGSFDYWGQ GTLTVTVSS	108
IGHV5-51*01 (M99686), plus JH4 element	EVQLVQSGAEVKKPGESLKISCKGSGYSFTSYWIGW VRQMPGKGLEWMGIIPGDS DTRYSPSFQGGV TISA DKSISTAYLQWSSLKASDTAMY CARGGSGGSFDY WGQGLTVTVSS	109
CD3 specific antibody based on humanized CH2527 light chain	QTVVTQEPSLTVSPGGTVTLTCGSSTGAVTTSNYAN WVQEKPGQA FRGLIGGTNKRAPGTPARFSGSLLGGK AALTLSGAQPEDEAEYYCALWYSNLWVFGGGTKLT VL	110
hVK1-39 (JK4 J-element)	DIQMTQSPSSLSASVGDRVTITCRASQSISSYLNWYQ QKPGKAPKLLIYAASSLQSGVPSRFSGSGSGTDFTLT SSLQPEDFATYYCQQSYSTPLTFGGGKVEIK	111

-231-

VL7_46-13 (humanized anti-CD3 antibody light chain)	QAVVTQEPSLTVSPGGTVTLTCGSSTGAVTTSNYAN WVQEKPQAFRGLIGGTNKRAPGTPARFSGSLLGGK AALTLSGAQPEDEAEYYCALWYSNLWVFGGGTKLT VL	112
--	--	-----

8) Exemplary 16D5 variants with reduced affinity

a. Exemplary light chain variants with reduced affinity

Name	Sequence	Seq ID No
K53A aa	QTVVTQEPSLTVSPGGTVTLTC <u>GSSTGAVTTSNYAN</u> WVQQKPGQAPRGLIGG <u>TNARAP</u> GTPARFSGSLLGGKAALTLSGVQPEDEAEYYC <u>ALWYSNLWV</u> FGGGT KLTVL	113
K53A_VL _CDR1	GSSTGAVTTSNYAN	32
K53A_VL _CDR2	GTNARAP	396
K53A_VL _CDR3	ALWYSNLWV	34
S93A aa	QTVVTQEPSLTVSPGGTVTLTC <u>GSSTGAVTTSNYAN</u> WVQQKPGQAPRGLIGG <u>TNKRAP</u> GTPARFSGSLLGGKAALTLSGVQPEDEAEYYC <u>ALWYANLWV</u> FGGGT KLTVL	114
S93A_VL CDR1	GSSTGAVTTSNYAN	32
S93A_VL CDR2	GTNKRAP	33
S93A_VL CDR3	ALWYANLWV	397

b. Exemplary heavy chain variants with reduced affinity

Name	Sequence	Seq ID No
S35H aa	EVQLVESGGGLVKPGGSLRLSCAASGFTFS <u>NAWMH</u> WVRQAPGKGLEWVGRIK <u>SKTDGGTTDYAAPVKG</u> RFTISRDDSKNTLYLQMNSLKTEDTAVYYCTT <u>PWEW</u> <u>SWYDY</u> WGQGTLVTVSSAS	115
S35H_VH _CDR1	NAWMH	398
S35H_VH _CDR2	RIKSKTDGGTTDYAAPVKG	17
S35H_VH _CDR3	PWEWSWYDY	18
G49S aa	EVQLVESGGGLVKPGGSLRLSCAASGFTFS <u>NAWMS</u> WVRQAPGKGLEWVSRIK <u>SKTDGGTTDYAAPVKG</u> RFTISRDDSKNTLYLQMNSLKTEDTAVYYCTT <u>PWEW</u> <u>SWYDY</u> WGQGTLVTVSSAS	116
G49S_VH _CDR1	NAWMS	16
G49S_VH	RIKSKTDGGTTDYAAPVKG	17

-232-

_CDR2		
G49S_VH _CDR3	PWEWSWYDY	18
R50S aa	EVQLVESGGGLVKPGGSLRLSCAASGFTFS <u>NAWMS</u> WVRQAPGKGLEWVG <u>SIK</u> <u>SKTDGGTTDYAAPVKG</u> RFTISRDDSKNTLYLQMNSLKTEDTAVYYCTT <u>PWEW</u> <u>SWYDY</u> WGQGTLVTVSSAS	117
R50S_VH _CDR1	NAWMS	16
R50S_VH _CDR2	SIKSKTDGGTTDYAAPVKG	399
R50S_VH _CDR3	PWEWSWYDY	18
W96Y aa	EVQLVESGGGLVKPGGSLRLSCAASGFTFS <u>NAWMS</u> WVRQAPGKGLEWVG <u>RIK</u> <u>SKTDGGTTDYAAPVKG</u> RFTISRDDSKNTLYLQMNSLKTEDTAVYYCTT <u>PYEW</u> <u>SWYDY</u> WGQGTLVTVSSAS	118
W96Y_V H_CDR1	NAWMS	16
W96Y_V H_CDR2	RIKSKTDGGTTDYAAPVKG	17
W96Y_V H_CDR3	PYEWWSWYDY	400
W98Y aa	EVQLVESGGGLVKPGGSLRLSCAASGFTFS <u>NAWMS</u> WVRQAPGKGLEWVG <u>RIK</u> <u>SKTDGGTTDYAAPVKG</u> RFTISRDDSKNTLYLQMNSLKTEDTAVYYCTT <u>PWEY</u> <u>SWYDY</u> WGQGTLVTVSSAS	119
W98Y_V H_CDR1	NAWMS	16
W98Y_V H_CDR2	RIKSKTDGGTTDYAAPVKG	17
W98Y_V H_CDR3	PWEYSWYDY	232

9) Additional exemplary embodiments generated from a phage display library (CDRs underlined)

Name	Sequence	Seq ID No
90D7 aa	QVQLVQSGAEVKKPGASVKVSCKASGYTFT <u>SYMH</u> WVRQAPGQGLEWMG <u>IIN</u> <u>PSGGSTSYAQKFQG</u> RVMTMRDTSSTVYMELSSLRSED TAVYYCARN <u>NYTIVV</u> <u>SPFDY</u> WGQGTLVTVSSAS	120
90D7_VH _CDR1	SYMH	8
90D7_VH _CDR2	IINPSGGSTSYAQKFQG	9
90D7_VH _CDR3	NYTIVVSPFDY	233
90C1	QVQLVQSGAEVKKPGASVKVSCKASGYTFT <u>SYMH</u> WVRQAPGQGLEWMG <u>IIN</u> <u>PSGGSTSYAQKFQG</u> RVMTMRDTSSTVYMELSSLRSED TAVYYCARN <u>NYFIGS</u>	121

-233-

aa	<u>VAMDY</u> WGQGTILVTVSSAS	
90C1_VH _CDR1	SYMH	8
90C1_VH _CDR2	IINPSGGSTSYAQKFQG	9
90C1_VH _CDR3	NYFIGSVAMDY	234
5E8 VH aa	QVQLVQSGAEVKKPGASVKVSCKASGYTFT SYMH WVRQAPGGGLEWMGI IIN PSGGSTSYAQKFQG RVMTMTRDTSTSTVYMEISSLRSED TAVYYCARG GLTYSM DY WGQGTILVTVSSAS	122
5E8_VH_ CDR1	SYMH	8
5E8_VH_ CDR2	IINPSGGSTSYAQKFQG	9
5E8_VH_ CDR3	GLTYSMDY	235
5E8 VL aa	DIVMTQSPLSLPVTPGEPASISCR RSSQSLLS NGYNYLDWYLQKPGQSPQLL IY LGSNRAS GVPDFRSGSGSGTDFTLKISRVEAEDVGVYYC MQALQIPNT FG QGTKVEIKRT	123
5E8_VL_ CDR1	RSSQSLLSNGYNYLD	59
5E8_VL_ CDR2	LGSNRAS	60
5E8_VL_ CDR3	MQALQIPNT	236
12A4 VH aa	EVQLLESGLLVQPGGSLRLSCAASGFTFS SYAMS WVRQAPGKGLEWVS AI SGSGSTYYADSVKGR FTISRDNKNTLYLQMNSLR AEDTAVYYCAK YAYALD Y WGQGTILVTVSSAS	124
12A4_VH _CDR1	SYAMS	79
12A4_VH _CDR2	AISGSGGSTYYADSVKG	80
12A4_VH _CDR3	YAYALDY	237
12A4 VL aa	EIVLTQSPGTLISLSPGERATLSC RASQSVSSSYLA WYQQKPGQAPRLLIY GA SSRAT GIPDRFSGSGSGTDFTLTISRLEPEDFAVYYC QQHGSSST FGQGTKV EIKRT	125
12A4_VL_ CDR1	RASQSVSSSYLA	52
12A4_VL_ CDR2	GASSRAT	53
12A4_VL_ CDR3	QQHGSSST	238
7A3 VH	QVQLVQSGAEVKKPGASVKVSCKASGYTFT SYMH WVRQAPGGGLEWMGI IIN PSGGSTSYAQKFQG RVMTMTRDTSTSTVYMEISSLRSED TAVYYCARG GDFSAG	126

-234-

aa	<u>RLMDY</u> WGQGTILVTVSSAS	
7A3_VH_ CDR1	SYMH	8
7A3_VH_ CDR2	IINPSGGSTSYAQKFQG	9
7A3_VH_ CDR3	GDFSAGRLMDY	239
7A3 VL aa	DIVMTQSPLSLPVTGPGEPAISCR <u>RSSQSLLS</u> NGYNYLDWYLQKPGQSPQLL IY <u>LGSNRAS</u> GVPDRFSGSGGTDFTLKISRVEAEDVGYYC <u>MQALQTPPI</u> TF GQGTKVEIKRT	127
7A3_VL_ CDR1	RSSQSLLSNGYNYLD	59
7A3_VL_ CDR2	LGSNRAS	60
7A3_VL_ CDR3	MQALQTPPI	240
6E10 VH aa	QVQLVQSGAEVKKPGASVKVSCKASGYTFT <u>SYMH</u> WVRQAPGQGLEWMGI <u>IIN</u> <u>PSGGSTSYAQKFQG</u> RVTMTRDTSTSTVYME <u>LS</u> SLRSEDYAVYYCARG <u>GDYNAF</u> <u>DY</u> WGHGTLVTVSSAS	128
6E10_VH _CDR1	SYMH	8
6E10_VH _CDR2	IINPSGGSTSYAQKFQG	9
6E10_VH _CDR3	GDYNAFDY	241
6E10 VL aa	DIVMTQSPLSLPVTGPGEPAISCR <u>RSSQSLLS</u> NGYNYLDWYLQKPGQSPQLL IY <u>LGSNRAS</u> GVPDRFSGSGGTDFTLKISRVEAEDVGYYC <u>MQAWHSPT</u> TFGQ GTKVEIKRT	129
6E10_VL_ CDR1	RSSQSLLSNGYNYLD	59
6E10_VL_ CDR2	LGSNRAS	60
6E10_VL_ CDR3	MQAWHSPT	242
12F9 VH aa	QVQLVQSGAEVKKPGASVKVSCKASGYTFT <u>SYMH</u> WVRQAPGQGLEWMGI <u>IIN</u> <u>PSGGSTSYAQKFQG</u> RVTMTRDTSTSTVYME <u>LS</u> SLRSEDYAVYYCARG <u>GATYTM</u> <u>DY</u> WGQGTILVTVSSAS	130
12F9_VH _CDR1	SYMH	8
12F9_VH _CDR2	IINPSGGSTSYAQKFQG	9
12F9_VH _CDR3	GATYTM	243
12F9 VL	DIVMTQSPLSLPVTGPGEPAISCR <u>RSSQSLLS</u> NGYNYLDWYLQKPGQSPQLL IY <u>LGSNRAS</u> GVPDRFSGSGGTDFTLKISRVEAEDVGYYC <u>MQALQTPPI</u> TFG	131

-235-

aa	QGTKVEIKRT	
12F9_VL_ CDR1	RSSQSLHLSNGYNYLD	59
12F9_VL_ CDR2	LGSNRAS	60
12F9_VL_ CDR3	MQALQTPIT	244

10) 9D11 Glycosite variants: variable light chain of exemplary embodiments
(CDRs underlined)

Variant	Sequence	Seq ID No
N95S	DIVMTQSPLSLPVTPGEPASISCRSSQSLHLSNGYNYLDWYLQKPGQSPQLL IYLGSNRASGVPDRFSGSGSGTDFTLKISRVEAEDVGVYYCMQASIM <u>S</u> RTFG QGTKVEIK	132
12F9_VL_ CDR1	RSSQSLHLSNGYNYLD	59
12F9_VL_ CDR2	LGSNRAS	60
12F9_VL_ CDR3	MQASIMSRT	63
N95Q	DIVMTQSPLSLPVTPGEPASISCRSSQSLHLSNGYNYLDWYLQKPGQSPQLL IYLGSNRASGVPDRFSGSGSGTDFTLKISRVEAEDVGVYYCMQASIM <u>Q</u> RTFG QGTKVEIK	133
N95Q_VL_ _CDR1	RSSQSLHLSNGYNYLD	59
N95Q_VL_ _CDR2	LGSNRAS	60
N95Q_VL_ _CDR3	MQASIMQRT	65
T97A	DIVMTQSPLSLPVTPGEPASISCRSSQSLHLSNGYNYLDWYLQKPGQSPQLL IYLGSNRASGVPDRFSGSGSGTDFTLKISRVEAEDVGVYYCMQASIMN <u>R</u> AFG QGTKVEIK	134
T97A_VL_ _CDR1	RSSQSLHLSNGYNYLD	59
T97A_VL_ _CDR2	LGSNRAS	60
T97A_VL_ _CDR3	MQASIMNRA	67
T97N	DIVMTQSPLSLPVTPGEPASISCRSSQSLHLSNGYNYLDWYLQKPGQSPQLL IYLGSNRASGVPDRFSGSGSGTDFTLKISRVEAEDVGVYYCMQASIMN <u>R</u> NFG QGTKVEIK	135
T97N_VL_ _CDR1	RSSQSLHLSNGYNYLD	59
T97N_VL	LGSNRAS	60

-236-

_CDR2		
T97N_VL	MQASIMNRN	69
_CDR3		

11) Deamination Variants

Variant	Sequence	Seq ID No
16D5 VH_D52d E	EVQLVESGGGLVQPGGSLRLSCAASGFTFSNAWMSWVRQAPGKGLEWVGRIK SKT E GGTTDYAAPVKGRFTISRDDSKNTLYLQMNSLKTEDTAVYYCTTPWEW SWYDYWGQGTLLTVSS	248
16D5 VH_D52d Q	EVQLVESGGGLVQPGGSLRLSCAASGFTFSNAWMSWVRQAPGKGLEWVGRIK SKT Q GGTTDYAAPVKGRFTISRDDSKNTLYLQMNSLKTEDTAVYYCTTPWEW SWYDYWGQGTLLTVSS	249
CD3_VH N100A	EVQLLESGGGLVQPGGSLRLSCAASGFTFSTYAMNWVRQAPGKGLEWVSRIR SKYNNYATYYADSVKGRFTISRDDSKNTLYLQMNSLRAEDTAVYYCVRHGNF G ASYVSWFAYWGQGTLLTVSS	250
CD3_VH S100aA	EVQLLESGGGLVQPGGSLRLSCAASGFTFSTYAMNWVRQAPGKGLEWVSRIR SKYNNYATYYADSVKGRFTISRDDSKNTLYLQMNSLRAEDTAVYYCVRHGNF GN A YVSWFAYWGQGTLLTVSS	251
16D5 [VHCH1] - CD3[VHC H1- N100A]- Fcknob_P GLALA	EVQLVESGGGLVQPGGSLRLSCAASGFTFSNAWMSWVRQAPGKGLEWVGRIK SKTDGGTTDYAAPVKGRFTISRDDSKNTLYLQMNSLKTEDTAVYYCTTPWEW SWYDYWGQGTLLTVSSASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEP VTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVTVTPSSSLGTQTYICNVNHK PSNTKVDKKVEPKSCDGGGGSGGGGSEVQLLESGGGLVQPGGSLRLSCAASG FTFSTYAMNWVRQAPGKGLEWVSRIRSKYNNYATYYADSVKGRFTISRDDSK NTLYLQMNSLRAEDTAVYYCVRHGNF G ASYVSWFAYWGQGTLLTVSSASTKG PSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVL QSSGLYSLSSVTVTPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHTC PPCPAPEAAGGPSVFLFPPKPKDTLMISRTPEVTCVVDVSHEDPEVKFNWY VDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALGA PIEKTISKAKGQPREPQVYTLPPCRDELTKNQVSLVCLVKGFYPSDIAVEWE SNGQPENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQQGNVSCSVMHEALHN HYTQKSLSLSPGK	252
16D5- Fchole- PGLALA	EVQLVESGGGLVQPGGSLRLSCAASGFTFSNAWMSWVRQAPGKGLEWVGRIK SKTDGGTTDYAAPVKGRFTISRDDSKNTLYLQMNSLKTEDTAVYYCTTPWEW SWYDYWGQGTLLTVSSASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEP VTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVTVTPSSSLGTQTYICNVNHK PSNTKVDKKVEPKSCDKTHTCPPCPAPEAAGGPSVFLFPPKPKDTLMISRT PEVTCVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVL HQDWLNGKEYKCKVSNKALGAPIEKTISKAKGQPREPQVCTLPSSRDELTKN QVSLSCAVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGSFFLVSKLTVD KSRWQQGNVSCSVMHEALHN R FTQKSLSLSPGK	253
CD3-CLC	QAVVTQEPSTLVSPGGTVTLTCGSSTGAVTTSNYANWVQEKPGQAFRGLIGG TNKRAPGTPARFSGSLLGGKAALTLGAQPEDEAEYYCALWYSNLWVFGGGT KLTVLGQPKAAPSVTLFPPSSEELQANKATLVCLISDFYPGAVTVAWKADSS PVKAGVETTTTPSKQSNKYAASSYLSLTPEQWKSHRSYSCQVTHEGSTVEKT VAPTECS	254
16D5 [VHCH1] - CD3[VHC	EVQLVESGGGLVQPGGSLRLSCAASGFTFSNAWMSWVRQAPGKGLEWVGRIK SKTDGGTTDYAAPVKGRFTISRDDSKNTLYLQMNSLKTEDTAVYYCTTPWEW SWYDYWGQGTLLTVSSASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEP VTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVTVTPSSSLGTQTYICNVNHK PSNTKVDKKVEPKSCDGGGGSGG	255

H1-S100aA]- Fcknob_P GLALA	GGSEVQLLESGGGLVQPGGSLRLSCAASGFTTFSTYAMNWVRQAPGKGLEWVS RIRSKYNNYATYYADSVKGRFTISRDDSKNTLYLQMNSLRAEDTAVYYCVRH GNFGNAYVSWFAYWGQGLTLTVSSASTKGPSVFPLAPSSKSTSGGTAALGCL VKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVVTVPSSSLGTQT YICNVNHKPSNTKVDKKVEPKSCDKTHTCPPCPAPEAAGGPSVFLFPPKPKD TLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYR VVSVLTVLHQDWLNGKEYKCKVSNKALGAPIEKTISKAKGQPREPQVYTLPP CRDELTKNQVSLWCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSF LYSKLTVDKSRWQQGNVFCSCVMHEALHNHYTQKSLSLSPGK	
9D11 [VHCH1] - CD3[VHC L- N100A]- Fcknob_P GLALA	QVQLVQSGAEVKKPGASVKVSCKASGYTFTSYMHWVRQAPGQGLEWMGIIN PSGGPTSYAQKFQGRVTMTRDTSTSTVYMELSSLRSEDVAVYYCARGDFAWL DYWGQGLTLTVSSASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVT VSWNSGALTSGVHTFPAVLQSSGLYSLSSVVTVPSSSLGTQTYICNVNHKPSN TKVDKKVEPKSCDGGGSGGGGSEVQLLESGGGLVQPGGSLRLSCAASGFTTF STYAMNWVRQAPGKGLEWVSRIIRSKYNNYATYYADSVKGRFTISRDDSKNTL YLQMNSLRAEDTAVYYCVRHGNFGASVSWFAYWGQGLTLTVSSASVAAPSV FIFPPSDEQLKSGTASVVCLLNNFYPREAKVQWKVDNALQSGNSQESVTEQD SKDSTYLSSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGECDKTHT CPPCPAPEAAGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNW YVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALG APIEKTISKAKGQPREPQVYTLPPCRDELTKNQVSLWCLVKGFYPSDIAVEW ESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVFCSCVMHEALH NHYTQKSLSLSPGK	256
9D11- Fchole	QVQLVQSGAEVKKPGASVKVSCKASGYTFTSYMHWVRQAPGQGLEWMGIIN PSGGPTSYAQKFQGRVTMTRDTSTSTVYMELSSLRSEDVAVYYCARGDFAWL DYWGQGLTLTVSSASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVT VSWNSGALTSGVHTFPAVLQSSGLYSLSSVVTVPSSSLGTQTYICNVNHKPSN TKVDKKVEPKSCDKTHTCPPCPAPEAAGGPSVFLFPPKPKDTLMISRTPEVT CVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQD WLNGKEYKCKVSNKALGAPIEKTISKAKGQPREPQVCTLPPSRDELTKNQVS LSCAVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLVSKLTVDKSR WQQGNVFCSCVMHEALHNHYTQKSLSLSPGK	257
9D11_LC [N95Q]	DIVMTQSPLSLPVTPGEPASISCRSSQSLLSNGYNYLDWYLQKPGQSPQLL IYLGSNRASGVPDFRSGSGSGTDFTLKISRVEAEDVGVYYCMQASIMQRTFG QGKVEIKRTVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQWKVD NALQSGNSQESVTEQDSKDSTYLSSTLTLSKADYEKHKVYACEVTHQGLSS PVTKSFNRGEC	258
CD3_VLC H1	QAVVTQEPSTLTVSPGGTVTLTCGSSTGAVTTSNYANWVQEKPGQAFRGLIGG TNKRAPGTTPARFSGSLLGGKAALTLGAQPEDEAEYYCALWYSNLWVFGGGT KLTVLSSASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGA LTSGVHTFPAVLQSSGLYSLSSVVTVPSSSLGTQTYICNVNHKPSNTKVDKK VEPKSC	259
9D11 [VHCH1] - CD3[VHC H1- S100aA]- Fcknob_P GLALA	QVQLVQSGAEVKKPGASVKVSCKASGYTFTSYMHWVRQAPGQGLEWMGIIN PSGGPTSYAQKFQGRVTMTRDTSTSTVYMELSSLRSEDVAVYYCARGDFAWL DYWGQGLTLTVSSASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVT VSWNSGALTSGVHTFPAVLQSSGLYSLSSVVTVPSSSLGTQTYICNVNHKPSN TKVDKKVEPKSCDGGGSGGGGSEVQLLESGGGLVQPGGSLRLSCAASGFTTF STYAMNWVRQAPGKGLEWVSRIIRSKYNNYATYYADSVKGRFTISRDDSKNTL YLQMNSLRAEDTAVYYCVRHGNFGNAYVSWFAYWGQGLTLTVSSASVAAPSV FIFPPSDEQLKSGTASVVCLLNNFYPREAKVQWKVDNALQSGNSQESVTEQD SKDSTYLSSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGECDKTHT CPPCPAPEAAGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNW YVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALG APIEKTISKAKGQPREPQVYTLPPCRDELTKNQVSLWCLVKGFYPSDIAVEW ESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVFCSCVMHEALH NHYTQKSLSLSPGK	260

12) Mov19 based FolR1 TCBs of exemplary embodiments (CDRs underlined)

Name	Sequence	Seq ID No
pETR1164 6 Mov19 VH-CH1- Fchole PG/LALA	QVQLQQSGAELVKPGASVKISCKASGYSTF <u>GYFMN</u> WVKQSHGKSLEWIGRIH <u>PYDGD</u> <u>TFYNQNF</u> <u>KD</u> KATLTVDKSSNTAHMELLSLTSEDFAVYYCTR <u>YDGSRA</u> <u>MDY</u> WGQGTITVTVSSASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVT VSWNSGALTSGVHTFPAVLQSSGLYSLSSVTVTPSSSLGTQTYICNVNHKPS NTKVDKKVEPKSCDKTHTCPPCPAPEAAGGPSVFLFPPKPKDTLMISRTPEV TCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQ DWLNGKEYKCKVSNKALGAPIEKTISKAKGQPREPQVCTLPPSRDELTKNQV SLSCAVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGSFFLVSKLTVDKS RWQQGNVFCFSVMHEALHNHYTQKSLSLSPGK	136
pETR1164 7 Mov19 VH-CH1- CD3 VH- CL- Fcknob PG/LALA	QVQLQQSGAELVKPGASVKISCKASGYSTF <u>GYFMN</u> WVKQSHGKSLEWIGRIH <u>PYDGD</u> <u>TFYNQNF</u> <u>KD</u> KATLTVDKSSNTAHMELLSLTSEDFAVYYCTR <u>YDGSRA</u> <u>MDY</u> WGQGTITVTVSSASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVT VSWNSGALTSGVHTFPAVLQSSGLYSLSSVTVTPSSSLGTQTYICNVNHKPS NTKVDKKVEPKSCDGGGGSGGGGSEVQLVESGGGLVQPKGSLKLSCAAS <u>GFT</u> <u>FNTYAMN</u> WVRQAPGKGLEWVAR <u>IRSKYNNYATYYADSVKD</u> RFTISRDDSQSI LYLQMNNLKTEDTAMYYCVR <u>HGNFGNSYVSWFAY</u> WGQGTITVTVSAASVAAPS VFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQWKVDNALQSGNSQESVTEQ DSKSTYLSLSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGECDKTH TCPPCPAPEAAGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFN WYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKAL GAPIEKTISKAKGQPREPQVYTLPPCRDELTKNQVSLWCLVKGFYPSDIAVE WESNGQPENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQQGNVFCFSVMHEAL HNHYTQKSLSLSPGK	137
pETR1164 4 Mov19 LC	DIETQSPASLAVSLGQRAII SC <u>KASQSVSFACTSLMH</u> WYHQKPGQPKLLI <u>YRASNLEA</u> GVPTRFSGSGSKTDFTLNIHPVEEEDAATYYC <u>QQSREYPYTF</u> GG GTKLEIKRTVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQWKVDN ALQSGNSQESVTEQDSKSTYLSLSTLTLSKADYEKHKVYACEVTHQGLSSP VTKSFNRGEC	138

Name	Sequence	Seq ID No
Hu IgG1 Fc	DKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPE VKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVS NKALPAPIEKTISKAKGQPREPQVYTLPPSRDELTKNQVSLTCLVKGFYPSD IAVEWESNGQPENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQQGNVFCFSVM HEALHNHYTQKSLSLSPGK	245

5 13) Additional FolR1 TCBs with intermediate affinity binders (CDRs according to Kabat, underlined)

Name	Sequence	Seq ID No
16D5 variant W96Y/D52 E VH	EVQLVESGGGLVKPGGSLRLSCAASGFTFS <u>NAWMS</u> WVRQAPGKGLEWV <u>GRIKSKTEGGTTDYAAPVKGR</u> FTISRDDSKNTLYLQMNSLKTEDTAVY YCTTPY <u>EW</u> SWYDYWGQGTITVTVSS	401
W96Y/D52 E_VH CDR1	NAWMS	16
W96Y/D52	RIKSKTEGGTTDYAAPVKG	402

E_VH CDR2		
W96Y/D52 E_VH CDR3	PYEWSWYDY	400
16D5 variant W96Y/D52 E VL	QAVVTQEP SLTVSPGGTVTLTCGSSTGAVTTSNYANWVQE KPGQA FRGLIGGTNKRAPGTPARFSGSLLGGKAALTLSGA QPEDEAEYYCALWYSNLWVFGGGTKLTVL	31
W96Y/D5 2E_CD3- VHCH1_F c- knob_PGL ALA pETR1494 5	EVQLVESGGGLVKPGGSLRLS CAASGFTFSNAWMSWVRQAPGKGLEWV GRIKSKTEGGTTDYAAPVKGRFTISRDDSKNTLYLQMNSLKTEDTAVY YCTTPYEWSWYDYWGQGLTVTVSSASTKGPSVFPLAPSSKSTSGGTAA LGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVTVTP SSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDGGGSGGGGSEVQILLE SGGGLVQPGGSLRLS CAASGFTFSTYAMNWVRQAPGKGLEWVSRIRSK YNNYATYYADSVKGRFTISRDDSKNTLYLQMNSLRAEDTAVYYCVRHG NFGNSYVSWFAYWGQGLTVTVSSASTKGPSVFPLAPSSKSTSGGTAA GCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVTVTPS SSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHTCPPCPAPEAAGGP SVFLFPPKPKDTLMI SRTPEVTCVVDVSHEDPEVKFNWYVDGVEVHN AKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALGAP IEK TISKAKGQPREPQVYTLPPCRDELTKNQVSLWCLVKGFYPSDIAVEWE SNGQPENNYKTTTPVLDSDGSFFFLYSKLTVDKSRWQQGNVFSCSVMHE ALHNHYTQKSLSLSPGK	403
W96Y/D5 2E_Fc- hole_PGL ALA_HYR F pETR1494 6	EVQLVESGGGLVKPGGSLRLS CAASGFTFSNAWMSWVRQAPGKGLEWV GRIKSKTEGGTTDYAAPVKGRFTISRDDSKNTLYLQMNSLKTEDTAVY YCTTPYEWSWYDYWGQGLTVTVSSASTKGPSVFPLAPSSKSTSGGTAA LGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVTVTP SSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHTCPPCPAPEAAGG PSVFLFPPKPKDTLMI SRTPEVTCVVDVSHEDPEVKFNWYVDGVEVH NAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALGAP IE KTISKAKGQPREPQVCTLPPSRDELTKNQVSLSCAVKGFYPSDIAVEW ESNGQPENNYKTTTPVLDSDGSFFFLYSKLTVDKSRWQQGNVFSCSVMH EALHNRFTQKSLSLSPGK	404
14B1 VH	EVQLLES GGGLVQPGGSLRLS CAASGFTFS <u>SYAMS</u> WVRQAPGKGLEWV <u>SAISGSGGSTYYADSVKGR</u> FTISRDN SKNTLYLQMNSLRAEDTAVYYC <u>ARGDYRYRYFDY</u> WGQGLTVTVSS	405
14B1 VL	<u>SSELTQDPAVSVALGQTVRITC</u> <u>OGDSLRSYYAS</u> WYQQKPGQAPVLVIY <u>GKNRNP</u> SGIPDRFS GSSSGNTASLTITGAQAEDEADYYC <u>NSRESPTG</u> <u>LVV</u> FGGGTKLTVL	406
14B1[EE]_ CD3[VLC H1]_Fc- knob_PGL ALA pETR1497 6	EVQLLES GGGLVQPGGSLRLS CAASGFTFSSYAMSWVRQAPGKGLEWV SAISGSGGSTYYADSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYYC ARGDYRYRYFDYWGQGLTVTVSSASTKGPSVFPLAPSSKSTSGGTAA GCLVEDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVTVTPS SSSLGTQTYICNVNHKPSNTKVDKVEPKSCDGGGSGGGGSGQAVVTQE PSLTVSPGGTVTLTCGSSTGAVTTSNYANWVQEKPGQA FRGLIGGTNK RAPGTPARFSGSLLGGKAALTLSGAQPEDEAEYYCALWYSNLWVFGGG TKLTVLSSASTKGPSVFPLAPSSKSTSGGTAA LGCLVKDYFPEPVTVS WNSGALTSGVHTFPAVLQSSGLYSLSSVTVTPSSSLGTQTYICNVNHK PSNTKVDKKVEPKSCDKTHTCPPCPAPEAAGGPSVFLFPPKPKDTLMI SRTPEVTCVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYR VVSVLTVLHQDWLNGKEYKCKVSNKALGAP IEKTISKAKGQPREPQVY TLPPCRDELTKNQVSLWCLVKGFYPSDIAVEWESNGQPENNYKTTTPV LDSDGSFFFLYSKLTVDKSRWQQGNVFSCSVMHEALHNHYTQKSLSLSP GK	407

14B1[EE]_ Fc- hole_PGL ALA pETR1497 7	EVQLLESGGGLVQPGGSLRLSCAASGFTFSYAMSWVRQAPGKGLEWV SAISGSGGSTYYADSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYYC ARGDYRYRYFDYWGQGTILVTVSSASTKGPSVFPLAPSSKSTSGGTAAL GCLVEDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVTVPS SSLGTQTYICNVNHKPSNTKVDEKVEPKSCDKTHTCPPCPAPEAAGGP SVFLFPPKPKDTLMI SRTPEVTCVVDVSHEDPEVKFNWYVDGVEVHN AKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALGAPIEK TISKAKGQPREPQVCTLPSPRDELTKNQVSLSCAVKGFYPSDIAVEWE SNGQPENNYKTPPVLDSDGSFFLVSKLTVDKSRWQQGNVFSCSVMHE ALHNHYTQKSLSLSPGK	408
14B1 LC [KK] Constant lambda pETR1497 9	SSELTQDPAVSVALGQTVRITCQGDSLRSYYASWYQQKPGQAPVLIY GKNNRPSGIPDRFSGSSSGNTASLTITGAQAEDEADYYCNSRESPTG LVVFGGGTKLTVLGQPKAAPSVTLFPPSSKKLQANKATLVCLISDFYP GAVTVAWKADSSPVKAGVETTTPSKQSNKNYAASSYLSLTPEQWKSHR SYSCQVTHEGSTVEKTVAPTECS	409
9C7 VH	QVQLVQSGAEVKKPGASVKVSCKASGYTFTSYMHWVRQAPGQGLEWM GIINPSGGSTSYAQKFQGRVTMTRDTSTSTVYME LSSLRSED TAVYYC ARGDWSYMDYWGQGTILVTVSS	410
9C7 VL	DIVMTQSPLSLPVTPGEPASISCRSSQSL LHSNGYNYLDWYLQKPGQS PQLLIYLGSNRASGVDPDRFSGSGSGTDFTLKI SRVEAEDVG VYYCMQA RQTPTFGQGTKVEIK	411
9C7[EE]_ CD3[VLC H1]_Fc- knob_PGL ALA pETR1497 4	QVQLVQSGAEVKKPGASVKVSCKASGYTFTSYMHWVRQAPGQGLEWM GIINPSGGSTSYAQKFQGRVTMTRDTSTSTVYME LSSLRSED TAVYYC ARGDWSYMDYWGQGTILVTVSSASTKGPSVFPLAPSSKSTSGGTAALG CLVEDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVTVPS SLGTQTYICNVNHKPSNTKVDEKVEPKSCDGGGSGGGGSQAVVTQEP SLTVSPGGTVLTTCGSSTGAVTTSNYANWVQEKPGQAFRGLIGGTNKR APGTPARFSGSLGGAALTLGSAQPEDEAEYYCALWYSNLWVFGGGT KLTVLSSASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSW NSGALTSGVHTFPAVLQSSGLYSLSSVTVPSSSLGTQTYICNVNHKPS NTKVDEKVEPKSCDKTHTCPPCPAPEAAGGP SVFLFPPKPKDTLMI S RTPETCVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRV VSVLTVLHQDWLNGKEYKCKVSNKALGAPIEKTISKAKGQPREPQVYT LPPCRDELTKNQVSLWCLVKGFYPSDIAVEWESNGQPENNYKTPPVLD SDGSFFLYSKLTVDKSRWQQGNVFSCSVMHEALHNHYTQKSLSLSPG K	412
9C7[EE]_ Fc- hole_PGL ALA pETR1497 5	QVQLVQSGAEVKKPGASVKVSCKASGYTFTSYMHWVRQAPGQGLEWM GIINPSGGSTSYAQKFQGRVTMTRDTSTSTVYME LSSLRSED TAVYYC ARGDWSYMDYWGQGTILVTVSSASTKGPSVFPLAPSSKSTSGGTAALG CLVEDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVTVPS SLGTQTYICNVNHKPSNTKVDEKVEPKSCDKTHTCPPCPAPEAAGGPS VFLFPPKPKDTLMI SRTPEVTCVVDVSHEDPEVKFNWYVDGVEVHNA KTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALGAPIEKT ISKAKGQPREPQVCTLPSPRDELTKNQVSLSCAVKGFYPSDIAVEWES NGQPENNYKTPPVLDSDGSFFLVSKLTVDKSRWQQGNVFSCSVMHEA LHNHYTQKSLSLSPGK	413
9C7 LC [RK] pETR1498 0	DIVMTQSPLSLPVTPGEPASISCRSSQSL LHSNGYNYLDWYLQKPGQS PQLLIYLGSNRASGVDPDRFSGSGSGTDFTLKI SRVEAEDVG VYYCMQA RQTPTFGQGTKVEIKRTVAAPSVFIFPPSDRKLKSGTASVVCLLNNFY PREAKVQWKVDNALQSGNSQESVTEQDSKDSYSTLSSTLTLSKADYEK HKVYACEVTHQGLSSPVTKSFNRGEC	414

14) Antigen Sequences

Antigen	Sequence	Seq ID No
hu FolR1	MAQRMTTQLLLLLLVWVAVVGEAQTRIAWARTELLNVCMNAAKHHKEKPGPEDKL HEQCRPWKRNACCSTNTSQAHKDVSYLRFNWNHCGEMAPACKRHF IQDTCL YECSPNLGPWIIQQVDQSWRKERVNLNPLCKEDCEQWWEDCRTSYTCKSNWHKG WNWTSGFNKCAGAACQPFHFYFPTPTVLCNEIWTHSYKVSNSYRSGSGRCIQM WFDPAQGNPNEEVARFYAAAMSGAGPWAAPFLLSLALMLLWLLS	139
huFolR1 ECD- AcTev- Fcknob- Avi tag	RIAWARTELLNVCMNAAKHHKEKPGPEDKLHEQCRPWKRNACCSTNTSQAHKD VSYLRFNWNHCGEMAPACKRHF IQDTCLYECSPNLGPWIIQQVDQSWRKERV NLNPLCKEDCEQWWEDCRTSYTCKSNWHKGWNWTSGFNKCAGAACQPFHFYF PTPTVLCNEIWTHSYKVSNSYRSGSGRCIQMWFDPAQGNPNEEVARFYAAAMVDE <u>QLYFQGGSPKSADKTHTCPPCPAPPELLGGPSVFLFPPKPKDTLMISRTPEVTC</u> VVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWL NGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPCRDELTKNQVSLWC LVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQQG NVFSCSVMHEALHNHYTQKSLSLSPGKSGGLNDIFEAQKIEWHE	140
Fchole	DKTHTCPPCPAPPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEV KFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNK ALPAPIEKTISKAKGQPREPQVCTLPSPRDELTKNQVSLSCAVKGFYPSDIAV EWESNGQPENNYKTTTPVLDSDGSFFLVSKLTVDKSRWQQGNVFCSSVMHEAL HNRTQKSLSLSPGK	141
mu FolR1	MAHLMTVQLLLLLVMWMAECAQSRATRARTTELLNVCMDAKHHKEKPGPEDNLHD QCSPWKTNSSCSTNTSQAHKDISYLRFNWNHCGTMTSECKRHF IQDTCLYE CSPNLGPWIIQQVDQSWRKERILDVPLCKEDCQWWEDCQSSFTCKSNWHKGWN WSSGHNECPVGASCHPFTFYFPTSAALCEEIWSHSYKLSNSYRSGSGRCIQMWF DPAQGNPNEEVARFYAEAMSGAGLHGTWPLLCSLSLVLLWVIS	142
mu FolR1 ECD- AcTev- Fcknob- Avitag	TRARTELLNVCMDAKHHKEKPGPEDNLHDQCSPWKTNSSCSTNTSQAHKDIS YLRFNWNHCGTMTSECKRHF IQDTCLYECSPNLGPWIIQQVDQSWRKERILDV PLCKEDCQWWEDCQSSFTCKSNWHKGWNWSSGHNECPVGASCHPFTFYFPTSA ALCEEIWSHSYKLSNSYRSGSGRCIQMWFDPAQGNPNEEVARFYAEAMVDE <u>QL</u> <u>YFQGGSPKSADKTHTCPPCPAPPELLGGPSVFLFPPKPKDTLMISRTPEVTCV</u> VDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNG KEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPCRDELTKNQVSLWCLV KGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQQGNV FSCSVMHEALHNHYTQKSLSLSPGKSGGLNDIFEAQKIEWHE	143
cy FolR1	MAQRMTTQLLLLLLVWVAVVGEAQTRTARARTELLNVCMNAAKHHKEKPGPEDKL HEQCRPWKKNACCSTNTSQAHKDVSYLRFNWNHCGEMAPACKRHF IQDTCL YECSPNLGPWIIQQVDQSWRKERVNLNPLCKEDCERWWEDCRTSYTCKSNWHKG WNWTSGFNKCVPGAACQPFHFYFPTPTVLCNEIWTSYKVSNSYRSGSGRCIQM WFDPAQGNPNEEVARFYAAAMSGAGPWAAPLALLSLALTLWLLS	144
cy FolR1 ECD- AcTev- Fcknob- Avi tag	RTARARTELLNVCMNAAKHHKEKPGPEDKLHEQCRPWKKNACCSTNTSQAHKD VSYLRFNWNHCGEMAPACKRHF IQDTCLYECSPNLGPWIIQQVDQSWRKERV NLNPLCKEDCEQWWEDCRTSYTCKSNWHKGWNWTSGFNKCVPGAACQPFHFYF PTPTVLCNEIWTSYKVSNSYRSGSGRCIQMWFDPAQGNPNEEVARFYAAAMVDE <u>QLYFQGGSPKSADKTHTCPPCPAPPELLGGPSVFLFPPKPKDTLMISRTPEVTC</u> VVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWL NGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPCRDELTKNQVSLWC LVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQQG NVFSCSVMHEALHNHYTQKSLSLSPGKSGGLNDIFEAQKIEWHE	145
hu FolR2	MVWKWMPLLLLLVCVATMCSAQDRDRLNVCMDAKHHKTKPGPEDKLHDQCSP WKKNACCTASTSQELHKDTSRLYNFNWDHCGKMEPACKRHF IQDTCLYECSPN LGPWIIQQVNQSWRKERFLDVPLCKEDCQRWWEDCHTSHTCKSNWHRGWDWTSG VNKCPAGALCRTFESYFPTPAALCEGLWSHSYKVSNSYRSGSGRCIQMWFDQAQ GNPNEEVARFYAAAMHVNAGEMHLHGTGGLLLSLALMLQLWLLG	146
hu FolR2 ECD- AcTev-	TMCSAQDRDRLNVCMDAKHHKTKPGPEDKLHDQCSPWKKNACCTASTSQELH KDTSRLYNFNWDHCGKMEPACKRHF IQDTCLYECSPNLGPWIIQQVNQSWRKER FLDVPLCKEDCQRWWEDCHTSHTCKSNWHRGWDWTSGVNKCPAGALCRTFESY FPTPAALCEGLWSHSYKVSNSYRSGSGRCIQMWFDQAQGNPNEEVARFYAAAMH	147

Fcknob-Avi tag	VVDEQLYFQGGSPKSADKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPCRDELTKNQVSLWCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVSCSVMHEALHNHYTQKSLSLSPGKSGGLNDIFEAQKIEWHE	
hu FolR3	MAWQMMQLLLLLALVTAAGSAQPR SARARTDLLNVCMAKHHKTQPSPEDELYGQCSPWKKNACCTASTSQELHKDTSRLYNFNWDHCGKMEPTCKRHF IQDSCLYE CSPNLGPWIRQVNQSWRKERILNVPLCKEDCERWWEDCRTSYTCKSNWHKGWNWTSGINECPAGALCSTFESYFPTPAALCEGLWSHSFKVSNYSRSGSGRCIQMWFDSAQGNPNEEVAKFYAAAMNAGAPSRGIIDS	148
hu FolR3 ECD-AcTev-Fcknob-Avi tag	SARARTDLLNVCMAKHHKTQPSPEDELYGQCSPWKKNACCTASTSQELHKDTSRLYNFNWDHCGKMEPTCKRHF IQDSCLYECSPNLGPWIRQVNQSWRKERILNVPLCKEDCERWWEDCRTSYTCKSNWHKGWNWTSGINECPAGALCSTFESYFPTPAALCEGLWSHSFKVSNYSRSGSGRCIQMWFDSAQGNPNEEVAKFYAAAMNAGAPSRGIIDSVDEQLYFQGGSPKSADKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPCRDELTKNQVSLWCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVSCSVMHEALHNHYTQKSLSLSPGKSGGLNDIFEAQKIEWHE	149
hu CD3e	MQSGTHWRVLGLCLLSVGWVGQDGNEEMGGITQTPYKVSISGTTVILTCPQYPGSEILWQHNDKNIGGEDDKNIGSDEHDLSLKEFSELEQSGYYVCYPRGSKPE DANFYLYLRARVCENCMEMDVMSVATIVIVDICITGGLLLL VYYWSKNRKAKA KPVTRGAGAGGRQGRQNKERPPVPNPDEYPIRKGQRDLYSGLNQRRRI	150

2) Nucleotide sequences of exemplary embodiments

Description	Sequence	Seq ID No
16A3	CAGGTGCAATTGGTTCAATCTGGTGCTGAAGTAAAAAACC GGGCGCTTCCGTTAAAGTGAGCTGCAAAGCATCCGGATACACCTTCACTTCCTATTACATGCACTGGGTTTCGTCAAGCCCCGGGCCAGGGTCTGGAA TGGATGGGCATCATTAAACCCAAGCGGTGGCTCTACCTCCTACGCGCAGAAATTCCAGGGTCGCGTCACGATGACCCGTGACACTAGCACCTCTACCGTTTATATGGAGCTGTCCAGCCTGCGTTCTGAAGATACTGCA GTGTACTACTGTGCACGCAACTACTACGCTGGTGTACTCCGTTTCGACTATTGGGGTCAAGGCACCCTCGTAACGGTTTCTTCT	151
15A1	CAGGTGCAATTGGTTCAATCTGGTGCTGAAGTAAAAAACC GGGCGCTTCCGTTAAAGTGAGCTGCAAAGCATCCGGATACACCTTCACTTCCTATTACATGCACTGGGTTTCGTCAAGCCCCGGGCCAGGGTCTGGAA TGGATGGGCATCATTAAACCCAAGCGGTGGCTCTACCTCCTACGCGCAGAAATTCCAGGGTCGCGTCACGATGACCCGTGACACTAGCACCTCTACCGTTTATATGGAGCTGTCCAGCCTGCGTTCTGAAGATACTGCA GTGTACTACTGTGCACGCAACTACTACATCGGTGTTGTTACTTTTCGACTATTGGGGTCAAGGCACCCTCGTAACGGTTTCTTCT	152
18D3	CAGGTGCAATTGGTTCAATCTGGTGCTGAAGTAAAAAACC GGGCGCTTCCGTTAAAGTGAGCTGCAAAGCATCCGGATACACCTTCACTTCCTATTACATGCACTGGGTTTCGTCAAGCCCCGGGCCAGGGTCTGGAA TGGATGGGCATCATTAAACCCAAGCGGTGGCTCTACCTCCTACGCGCAGAAATTCCAGGGTCGCGTCACGATGACCCGTGACACTAGCACCTCTACCGTTTATATGGAGCTGTCCAGCCTGCGTTCTGAAGATACTGCA GTGTACTACTGTGCACGCAACTACTACACTGGTGGTTCTTCTGCTTTCGACTATTGGGGTCAAGGCACCCTCGTAACGGTTTCTTCT	153
19E5	CAGGTGCAATTGGTTCAATCTGGTGCTGAAGTAAAAAACC GGGCG	154

	NTTCCGTTAAAGTGAGCTGCAAAGCATCCGGATACACCTTCACTTC CTATTACATGCACTGGGTTTCGTCAAGCCCCGGGCCAGGGTCTGGAA TGGATGGGCATCATTAAACCCAAGCGGTGGCTCTACCTCCTACGCGC AGAAATTCCAGGGTCGCGTCACGATGACCCGTGACACTAGCACCTC TACCGTTTATATGGAGCTGTCCAGCCTGCGTTCTGAAGATACTGCA GTGTACTACTGTGCACGCGGTGAATGGCGTCGTTACACTTCTTTTCG ACTATTGGGGTCAAGGCACCCCTCGTAACGGTTTCTTCT	
19A4	CAGGTGCAATTGGTTCAATCTGGTGCTGAAGTAAAAAACC GGCG CTTCCGTTAAAGTGAGCTGCAAAGCATCCGGATACACCTTCACTTC CTATTACATGCACTGGGTTTCGTCAAGCCCCGGGCCAGGGTCTGGAA TGGATGGGCATCATTAAACCCAAGCGGTGGCTCTACCTCCTACGCGC AGAAATTCCAGGGTCGCGTCACGATGACCCGTGACACTAGCACCTC TACCGTTTATATGGAGCTGTCCAGCCTGCGTTCTGAAGATACTGCA GTGTACTACTGTGCACGCGGTGGTTGGATCCGTTGGGAACATTTTCG ACTATTGGGGTCAAGGCACCCCTCGTAACGGTTTCTTCT	155
15H7	CAGGTGCAATTGGTTCAATCTGGTGCTGAAGTAAAAAACC GGCG CTTCCGTTAAAGTGAGCTGCAAAGCATCCGGATACACCTTCACTTC CTATTACATGCACTGGGTTTCGTCAAGCCCCGGGCCAGGGTCTGGAA TGGATGGGCATCATTAAACCCAAGCGGTGGCTCTACCTCCTACGCGC AGAAATTCCAGGGTCGCGTCACGATGACCCGTGACACTAGCACCTC TACCGTTTATATGGAGCTGTCCAGCCTGCGTTCTGAAGATACTGCA GTGTACTACTGTGCACGCAACTACTACCTGTTCTCTACTTCTTTTCG ACTATTGGGGTCAAGGCACCCCTCGTAACGGTTTCTTCT	156
15B6	CAGGTGCAATTGGTTCAATCTGGTGCTGAGGTAAAAAACC GGCG CTTCCGTTAAAGTGAGCTGCAAAGCATCCGGATACACCTTCACTTC CTATTACATGCACTGGGTTTCGTCAAGCCCCGGGCCAGGGTCTGGAA TGGATGGGCATCATTAAACCCAAGCGGTGGCTCTACCTCCTACGCGC AGAAATTCCAGGGTCGCGTCACGATGACCCGTGACACTAGCACCTC TACCGTTTATATGGAGCTGTCCAGCCTGCGTTCTGAAGATACTGCA GTGTACTACTGTGCACGCAACTACTACATCGGTATCGTTCCGTTTCG ACTATTGGGGTCAAGGCACCCCTCGTAACGGTTTCTTCT	157
16D5	GAGGTGCAATTGGTTGAATCTGGTGCTGGTGGTCTGGTAAAACC GGCG GTTCCCTGCGTCTGAGCTGCGCGGCTTCCGGATTACCTTCTCCAA CGCGTGGATGAGCTGGGTTTCGCCAGGCCCCGGGCAAAGGCCTCGAG TGGGTTGGTTCGTATCAAGTCTAAAACTGACGGTGGCACCACGGATT ACGCGGCTCCAGTTAAAGGTCGTTTTACCATTTCGCGACGATAG CAAAAACACTCTGTATCTGCAGATGAACTCTCTGAAAACCTGAAGAC ACCGCAGTCTACTACTGTACTACCCCGTGGGAATGGTCTTGGTACG ATTATTGGGGCCAGGGCACGCTGGTTACGGTGTCTTCC	158
15E12	GAGGTGCAATTGGTTGAATCTGGTGCTGGTGGTCTGGTAAAACC GGCG GTTCCNCGCTCTGAGCTGCGCGGCTTCCGGATTACCTTCTCCAA CGCGTGGATGAGCTGGGTTTCGCCAGGCCCCGGGCAAAGGCCTCGAG TGGGTTGGTTCGTATCAAGTCTAAAACTGACGGTGGCACCACGGATT ACGCGGCTCCAGTTAAAGGTCGTTTTACCATTTCGCGACGATAG CAAAAACACTCTGTATCTGCAGATGAACTCTCTGAAAACCTGAAGAC ACCGCAGTCTACTACTGTACTACCCCGTGGGAATGGTCTTACTTCG ATTATTGGGGCCAGGGCACGCTGGTTACGGTGTCTTCC	159
21D1	CAGGTGCAATTGGTTCAATCTGGTGCTGAAGTAAAAAACC GGCG CTTCCGTTAAAGTGAGCTGCAAAGCATCCGGATACACCTTCACTTC CTATTACATGCACTGGGTTTCGTCAAGCCCCGGGCCAGGGTCTGGAA TGGATGGGCATCATTAAACCCAAGCGGTGGCTCTACCTCCTACGCGC AGAAATTCCAGGGTCGCGTCACGATGACCCGTGACACTAGCACCTC TACCGTTTATATGGAGCTGTCCAGCCTGCGTTCTGAAGATACTGCA GTGTACTACTGTGCACGCAACTACTACGTTGGTGTCTTCCGTTTCG	160

	ACTATTGGGGTCAAGGCACCCTCGTAACGGTTTCTTCT	
16F12	CAGGTGCAATTGGTTCAATCTGGTGCTGAAGTAAAAAACC GGCG NTTCCGTTAAAGTGAGCTGCAAAGCATCCGGATACACCTTCACTTC CTATTACATGCACTGGGTTTCGTCAAGCCCCGGGCCAGGGTCTGGAA TGGATGGGCATCATTAAACCAAGCGGTGGCTCTACCTCNTACGCGC AGAAATTCCAGGGTCGCGTCACGATGACCCGTGACACTAGCACCTC TACCGTTTATATGGAGCTGTCCAGCCTGCGTTCTGAAGATACTGCA GTGTACTACTGTGCACGCAACTTCACTGTTCTGCGTGTTCCGTTTCG ACTATTGGGGTCAAGGCACCCTCGTAACGGTTTCTTCT	161
21A5	GAGGTGCAATTGGTTGAATCTGGTGGTGGTCTGGTAAAACCGGGCG GTTCCCTGCGTCTGAGCTGCGCGGCTTCCGGATTACCTTCTCCAA CGCGTGGATGAGCTGGGTTTCGCCAGGCCCCGGGCAAAGGCCTCGAG TGGGTTGGTTCGTATCAAGTCTAAACTGACGGTGGCACCACGGATT ACGCGGCTCCAGTTAAAGGTCGTTTTACCATTTCCTCGCAGCATAG CAAAAACACTCTGTATCTGCAGATGAACTCTCTGAAAACCTGAAGAC ACCGCAGTCTACTACTGTACTACCCCGTGGGAATGGGCTTGTTTCG ATTATTGGGGCCAGGGCACGCTGGTTACGGTGTCTTCC	162
21G8	GAGGTGCAATTGGTTGAATCTGGTGGTGGTCTGGTAAAACCGGGCG GTTCCCTGCGTCTGAGCTGCGCGGCTTCCGGATTACCTTCTCCAA CGCGTGGATGAGCTGGGTTTCGCCAGGCCCCGGGCAAAGGCCTCGAG TGGGTTGGTTCGTATCAAGTCTAAACTGACGGTGGCACCACGGATT ACGCGGCTCCAGTTAAAGGTCGTTTTACCATTTCCTCGCAGCATAG CAAAAACACTCTGTATCTGCAGATGAACTCTCTGAAAACCTGAAGAC ACCGCAGTCTACTACTGTACTACCCCGTGGGAATGGGCTTACTTCG ATTATTGGGGCCAGGGCACGCTGGTTACGGTGTCTTCC	163
19H3	CAGGTGCAATTGGTTCAATCTGGTGCTGAAGTAAAAAACC GGCG CTTCCGTTAAAGTGAGCTGCAAAGCATCCGGATACACCTTCACTTC CTATTACATGCACTGGGTTTCGTCAAGCCCCGGGCCAGGGTCTGGAA TGGATGGGCATCATTAAACCAAGCGGTGGCTCTACCTCCTACGCGC AGAAATTCCAGGGTCGCGTCACGATGACCCGTGACACTAGCACCTC TACCGTTTATATGGAGCTGTCCAGCCTGCGTTCTGAAGATACTGCA GTGTACTACTGTGCACGCACTGGTTGGTCTCGTTGGGGTTACATGG ACTATTGGGGCCAAGGCACCCTCGTAACGGTTTCTTCT	164
20G6	CAGGTGCAATTGGTTCAATCTGGTGCTGAAGTAAAAAACC GGCG CTTCCGTTAAAGTGAGCTGCAAAGCATCCGGATACACCTTCACTTC CTATTACATGCACTGGGTTTCGTCAAGCCCCGGGCCAGGGTCTGGAA TGGATGGGCATCATTAAACCAAGCGGTGGCTCTACCTCCTACGCGC AGAAATTCCAGGGTCGCGTCACGATGACCCGTGACACTAGCACCTC TACCGTTTATATGGAGCTGTCCAGCCTGCGTTCTGAAGATACTGCA GTGTACTACTGTGCACGCGGTGAATGGATCCGTTACTACCATTTCG ACTATTGGGGTCAAGGCACCCTCGTAACGGTTTCTTCT	165
20H7	CAGGTGCAATTGGTTCAATCTGGTGCTGAAGTAAAAAACC GGCG CTTCCGTTAAAGTGAGCTGCAAAGCATCCGGATACACCTTCACTTC CTATTACATGCACTGGGTTTCGTCAAGCCCCGGGCCAGGGTCTGGAA TGGATGGGCATCATTAAACCAAGCGGTGGCTCTACCTCCTACGCGC AGAAATTCCAGGGTCGCGTCACGATGACCCGTGACACTAGCACCTC TACCGTTTATATGGAGCTGTCCAGCCTGCGTTCTGAAGATACTGCA GTGTACTACTGTGCACGCGTTGGTTGGTACCGTTGGGGTTACATGG ACTATTGGGGTCAAGGCACCCTCGTAACGGTTTCTTCT	166
11F8_VH	CAGGTGCAATTGGTGCACTCTGGGGCTGAGGTGAAGAAGCCTGGGT CCTCGGTGAAGGTCTCCTGCAAGGCCTCCGGAGGCACATTCAGCAG CTACGCTATAAGCTGGGTGCGACAGGCCCTGGACAAGGGCTCGAG TGGATGGGAGGGATCATCCCTATCTTTGGTACAGCAAACCTACGCAC AGAAGTTCAGGGCAGGGTAACCATTACTGCAGACAAATCCACGAG	167

	CACAGCCTACATGGAGCTGAGCAGCCTGAGATCTGAGGACACCGCC GTGTATTACTGTGCGAGAGCTGTTTTCTACCGTGCTTGGTACTCTT TCGACTACTGGGGCCAAGGGACCACCGTGACCGTCTCCTCA	
11F8_VL	GACATCCAGATGACCCAGTCTCCTTCCACCTGTCTGCATCTGTAG GAGACCGTGTCAACCATCACTTGCCGTGCCAGTCAGAGTATTAGTAG CTGGTTGGCCTGGTATCAGCAGAAACCAGGGAAAGCCCCCTAAGCTC CTGATCTATGATGCCTCCAGTTTGGAAGTGGGGTCCCATCACGTT TCAGCGGCAGTGGATCCGGGACAGAATTCACCTCTCACCATCAGCAG CTTGCAGCCTGATGATTTTGCAACTTATTACTGCCAACAGTATAACC AGCCCACCACCAACGTTTGGCCAGGGCACCAAAGTCGAGATCAAG	168
36F2_VH	CAGGTGCAATTGGTTCAATCTGGTGCTGAAGTAAAAAAACCGGGCG CTTCCGTTAAAGTGAGCTGCAAAGCATCCGGATACACCTTCACTTC CTATTACATGCACTGGGTTTCGTCAAGCCCCGGGCCAGGGTCTGGAA TGGATGGGCATCATTAAACCAAGCGGTGGCTCTACCTCCTACGCGC AGAAATTCCAGGGTCGCGTCACGATGACCCATGACACTAGCACCTC TACCGTTTATATGGAGCTGTCCAGCCTGCGTTCTGAAGATACTGCA GTGTACTACTGTGCACGCTCTTTCTTCACTGGTTTCCATCTGGACT ATTGGGGTCAAGGCACCTCGTAACGGTTTCTTCT	169
36F2_VL	GAAATCGTGTTAACGCAGTCTCCAGGCACCTGTCTTTGTCTCCAG GGGAAAGAGCCACCCTCTCTTGCCAGGGCCAGTCAGAGTGTTAGCAG CAGCTACTTAGCCTGGTACCAGCAGAAACCTGGCCAGGCTCCCAGG CTCCTCATCTATGGAGCATCCAGCAGGGCCACTGGCATCCCAGACA GGTTCAGTGGCAGTGGATCCGGGACAGACTTCACTCTCACCATCAG CAGACTGGAGCCTGAAGATTTTGCAGTGTATTACTGTCAGCAGTAT ACCAACGAACATTATTATACGTTCCGGCCAGGGGACCAAAGTGGAAT TCAAA	170
9D11_VH	CAGGTGCAATTGGTTCAATCTGGTGCTGAAGTAAAAAAACCGGGCG CTTCCGTTAAAGTGAGCTGCAAAGCATCCGGATACACCTTCACTTC CTATTACATGCACTGGGTTTCGTCAAGCCCCGGGCCAGGGTCTGGAA TGGATGGGCATCATTAAACCAAGCGGTGGCCCTACCTCCTACGCGC AGAAATTCCAGGGTCGCGTCACGATGACCCGTGACACTAGCACCTC TACCGTTTATATGGAGCTGTCCAGCCTGCGTTCTGAAGATACTGCA GTGTACTACTGTGCACGCGGTGACTTCGCTTGGCTGGACTATTGGG GTCAAGGCACCTCGTAACGGTTTCTTCT	171
9D11_VL	GATATTGTTATGACTCAATCTCCACTGTCTCTGCCGGTGACTCCAG GCGAACCGGCGAGCATTTCTTGCCGTTCCAGCCAGTCTCTGCTGCA CTCCAACGGCTACAACCTATCTCGATTGGTACCTGCAAAAACCGGGT CAGAGCCCTCAGCTGCTGATCTACCTGGGCTCTAACCGCGCTTCCG GTGTACCGGACCGTTTCAGCGGCTCTGGATCCGGCACCGATTTCAC GTTGAAAATCAGCCGTGTTGAAGCAGAAGACGTGGGCGTTTATTAC TGTATGCAGGCAAGCATTATGAACCGGACTTTTGGTCAAGGCACCA AGGTCGAAATTAAA	172
9D11_VL N95S	GATATTGTTATGACTCAATCTCCACTGTCTCTGCCGGTGACTCCAG GCGAACCGGCGAGCATTTCTTGCCGTTCCAGCCAGTCTCTGCTGCA CTCCAACGGCTACAACCTATCTCGATTGGTACCTGCAAAAACCGGGT CAGAGCCCTCAGCTGCTGATCTACCTGGGCTCTAACCGCGCTTCCG GTGTACCGGACCGTTTCAGCGGCTCTGGATCCGGCACCGATTTCAC GTTGAAAATCAGCCGTGTTGAAGCAGAAGACGTGGGCGTTTATTAC TGTATGCAGGCAAGCATTATGAACCGGACTTTTGGTCAAGGCACCA AGGTCGAAATTAAA	173
9D11_VL N95Q	GATATTGTTATGACTCAATCTCCACTGTCTCTGCCGGTGACTCCAG GCGAACCGGCGAGCATTTCTTGCCGTTCCAGCCAGTCTCTGCTGCA CTCCAACGGCTACAACCTATCTCGATTGGTACCTGCAAAAACCGGGT CAGAGCCCTCAGCTGCTGATCTACCTGGGCTCTAACCGCGCTTCCG GTGTACCGGACCGTTTCAGCGGCTCTGGATCCGGCACCGATTTCAC GTTGAAAATCAGCCGTGTTGAAGCAGAAGACGTGGGCGTTTATTAC TGTATGCAGGCAAGCATTATGAACCGGACTTTTGGTCAAGGCACCA AGGTCGAAATTAAA	174

	GTGTACCGGACCGTTTTCAGCGGCTCTGGATCCGGCACCGATTTCAC GTTGAAAATCAGCCGTGTTGAAGCAGAAGACGTGGGCGTTTATTAC TGTATGCAGGCAAGCATTATGCAGCGGACTTTTGGTCAAGGCACCA AGGTCGAAATTAAA	
9D11_VL T97A	GATATTGTTATGACTCAATCTCCACTGTCTCTGCCGGTGACTCCAG GCGAACCGGCGAGCATTCTTGCCGTTCCAGCCAGTCTCTGTCTGCA CTCCAACGGCTACAACCTATCTCGATTGGTACCTGCAAAAACCGGGT CAGAGCCCTCAGCTGCTGATCTACCTGGGCTCTAACCGCGCTTCCG GTGTACCGGACCGTTTTCAGCGGCTCTGGATCCGGCACCGATTTCAC GTTGAAAATCAGCCGTGTTGAAGCAGAAGACGTGGGCGTTTATTAC TGTATGCAGGCAAGCATTATGAACCGGGCTTTTGGTCAAGGCACCA AGGTCGAAATTAAA	175
9D11_VL T97N	GATATTGTTATGACTCAATCTCCACTGTCTCTGCCGGTGACTCCAG GCGAACCGGCGAGCATTCTTGCCGTTCCAGCCAGTCTCTGTCTGCA CTCCAACGGCTACAACCTATCTCGATTGGTACCTGCAAAAACCGGGT CAGAGCCCTCAGCTGCTGATCTACCTGGGCTCTAACCGCGCTTCCG GTGTACCGGACCGTTTTCAGCGGCTCTGGATCCGGCACCGATTTCAC GTTGAAAATCAGCCGTGTTGAAGCAGAAGACGTGGGCGTTTATTAC TGTATGCAGGCAAGCATTATGAACCGGAATTTTGGTCAAGGCACCA AGGTCGAAATTAAA	176
5D9_VH	CAGGTGCAATTGGTTCAATCTGGTGCTGAAGTAAAAAAACCGGGCG CTTCCGTTAAAGTGAGCTGCAAAGCATCCGGATACACCTTCACTTC CTATTACATGCACTGGGTTTCGTCAAGCCCCGGGCCAGGGTCTGGAA TGGATGGGCATCATTAAACCAAGCGGTGGCTCTACCTCCTACGCGC AGAAATTCCAGGGTCGCGTACGATGACCCGTGACACTAGCACCTC TACCGTTTATATGGAGCTGTCCAGCCTGCGTTCTGAAGATACTGCA GTGTACTACTGTGCACGCTCTTACATCGACATGGACTATTGGGGTC AAGGCACCCTCGTAACGGTTTCTTCT	177
5D9_VL	GAAATCGTGTTAACGCAGTCTCCAGGCACCCTGTCTTTGTCTCCAG GGGAAAGAGCCACCCTCTCTTGCAAGGGCCAGTCAGAGTGTTAGCAG CAGTACTTAGCCTGGTACCAGCAGAAACCTGGCCAGGCTCCCAGG CTCCTCATCTATGGAGCATCCAGCAGGGCCACTGGCATCCCAGACA GGTTCAGTGGCAGTGGATCCGGGACAGACTTCACTCTCACCATCAG CAGACTGGAGCCTGAAGATTTTGCAGTGTATTACTGTCAGCAGGAT AACTGGAGCCCAACGTTTCGGCCAGGGGACCAAGTGGAATCAAA	178
6B6_VH	CAGGTGCAATTGGTTCAATCTGGTGCTGAAGTAAAAAAACCGGGCG CTTCCGTTAAAGTGAGCTGCAAAGCATCCGGATACACCTTCACTTC CTATTACATGCACTGGGTTTCGTCAAGCCCCGGGCCAGGGTCTGGAA TGGATGGGCATCATTAAACCAAGCGGTGGCTCTACCTCCTACGCGC AGAAATTCCAGGGTCGCGTACGATGACCCGTGACACTAGCACCTC TACCGTTTATATGGAGCTGTCCAGCCTGCGTTCTGAAGATACTGCA GTGTACTACTGTGCACGCTCTTACGTTGACATGGACTATTGGGGTC AAGGCACCCTCGTAACGGTTTCTTCT	179
6B6_VL	GAAATCGTGTTAACGCAGTCTCCAGGCACCCTGTCTTTGTCTCCAG GGGAAAGAGCCACCCTCTCTTGCAAGGGCCAGTCAGAGTGTTAGCAG CAGTACCTAGCCTGGTACCAGCAGAAACCTGGCCAGGCTCCCAGG CTCCTCATCTATGGAGCATCCAGCAGGGCCACTGGCATCCCAGACA GGTTCAGTGGCAGTGGATCCGGGACAGACTTCACTCTCACCATCAG CAGACTGGAGCCTGAAGATTTTGCAGTGTATTACTGTCAGCAGGAT ATTTGGAGCCCAACGTTTCGGCCAGGGGACCAAGTGGAATCAAA	180
14E4_VH	GAGGTGCAATTGTTGGAGTCTGGGGGAGGCTTGGTACAGCCTGGGG GGTCCCCTGAGACTCTCCTGTGCAGCCTCCGGATTACCTTTAGCAG TTATGCCATGAGCTGGGTCCGCCAGGCTCCAGGGAAGGGGCTGGAG TGGGTCTCAGCTATTAGTGGTAGTGGTGGTAGCACATACTACGCAG	181

	ACTCCGTGAAGGGCCGGTTACCATCTCCAGAGACAATTCCAAGAA CACGCTGTATCTGCAGATGAACAGCCTGAGAGCCGAGGACACGGCC GTATATTACTGTGCGAAAGACTCTTCTTACGTTGAATGGTACGCTT TCGACTACTGGGGCCAAGGAACCCCTGGTCACCGTCTCGAGT	
14E4_VL	GAAATCGTGTTAACGCAGTCTCCAGGCACCCCTGTCTTTGTCTCCAG GGGAAAGAGCCACCCTCTCTTGCAGGGCCAGTCAGAGTGTTAGCAG CAGCTACTTAGCCTGGTACCAGCAGAAACCTGGCCAGGCTCCCAGG CTCCTCATCTATGGAGCATCCAGCAGGGCCACTGGCATCCCAGACA GGTTCAGTGGCAGTGGATCCGGGACAGACTCCACTCTCACCATCAG CAGACTGGAGCCTGAAGATTTTGCAGTGTATTACTGTCAGCAGCCA ACCAGCAGCCCAATTACGTTTCGGCCAGGGGACCAAAGTGGAATCA AA	182
CD3 heavy chain (VHCH1)	GAGGTGCAGCTGCTGGAATCTGGCGGCGGACTGGTGCAGCCT GGCGGATCTCTGAGACTGAGCTGTGCCGCCAGCGGCTTCACC TTCAGCACCTACGCCATGAACTGGGTGCGCCAGGCCCCCTGGC AAAGGCCTGGAATGGGTGTCCCGGATCAGAAGCAAGTACAAC AACTACGCCACCTACTACGCCGACAGCGTGAAGGGCCGGTTC ACCATCAGCCGGGACGACAGCAAGAACACCCTGTACCTGCAG ATGAACAGCCTGCGGGCCGAGGACACCGCCGTGTACTATTGT GTGCGGCACGGCAACTTCGGCAACAGCTATGTGTCTTGGTTT GCCTACTGGGGCCAGGGCACCCCTCGTGACCGTGTCAAGCGCT AGTACCAAGGGCCCCAGCGTGTTCCTCCCTGGCACCCAGCAGC AAGAGCACATCTGGCGGAACAGCCGCTCTGGGCTGTCTGGTG AAAGACTACTTCCCCGAGCCCGTGACCGTGTCTTGGAATCT GGCGCCCTGACCAGCGGCGTGCACACCTTTCAGCCGTGCTG CAGAGCAGCGGCCTGTACTCCCTGTCCTCCGTGGTACCGTG CCCTCTAGCTCCCTGGGAACACAGACATATATCTGTAATGTC AATCACAAGCCTTCCAACACCAAAGTCGATAAGAAAGTCGAG CCCAAGAGCTGC	183
Crossed CD3 heavy chain (VHCκ)	GAGGTGCAGCTGCTGGAATCTGGCGGCGGACTGGTGCAGCCTGGCG GATCTCTGAGACTGAGCTGTGCCGCCAGCGGCTTCACCTTCAGCAC CTACGCCATGAACTGGGTGCGCCAGGCCCCCTGGCAAAGGCCTGGAA TGGGTGTCCCGGATCAGAAGCAAGTACAACAACCTACGCCACCTACT ACGCCGACAGCGTGAAGGGCCGGTTCACCATCAGCCGGGACGACAG CAAGAACACCCTGTACCTGCAGATGAACAGCCTGCGGGCCGAGGAC ACCGCCGTGTACTATTGTGTGCGGCACGGCAACTTCGGCAACAGCT ATGTGTCTTGGTTTGCCTACTGGGGCCAGGGCACCCCTCGTGACCGT GTCAAGCGCTAGTGTGGCCGCTCCCTCCGTGTTTATCTTTCCCCCA TCCGATGAACAGCTGAAAAGCGGCACCGCCTCCGTGCTGTGTCTGC TGAACAATTTTTACCCTAGGGAAGCTAAAGTGCAGTGGAAAGTGGA TAACGCACTGCAGTCCGGCAACTCCCAGGAATCTGTGACAGAACAG GACTCCAAGGACAGCACCTACTCCCTGTCTCCACCCTGACACTGT CTAAGGCTGATTATGAGAAACACAAAGTCTACGCCTGCGAAGTCAC CCATCAGGGCCTGAGCTCGCCCGTCACAAAGAGCTTCAACAGGGGA GAGTGT	184
Mutagenesi s primer GAB7734 N95Q	GCAGGCAAGCATTATGCAGCGGACTTTTGGTCAAGG	185
Mutagenesi	CAGGCAAGCATTATGAGCCGGACTTTTGGTCAAGG	186

s primer GAB7735 N95S		
Mutagenesi s primer GAB7736 T97A	CATTATGAACCGGGCTTTTGGTCAAGGCACCAAGGTC	187
Mutagenesi s primer GAB7737 T97N	CATTATGAACCGGAATTTTGGTCAAGGCACCAAGGTC	188
VHCH1[16 D5]_VHCH 1[CD3]_Fck nob_PGLA LA pCON999 (Inverted TCB with 16D5 2+1: pCON999 + pCON983 + pETR13197)	GAGGTGCAATTGGTTGAATCTGGTGGTGGTCTGGTAAAACCGGGCG GTTCCCTGCGTCTGAGCTGCGCGGCTTCCGGATTACCTTCTCCAA CGCGTGGATGAGCTGGGTTCGCCAGGCCCCGGGCAAAGGCCTCGAG TGGGTTGGTTCGTATCAAGTCTAAACTGACGGTGGCACCACGGATT ACGCGGCTCCAGTTAAAGGTCGTTTTACCATTTCCTCGCAGCATAG CAAAAACACTCTGTATCTGCAGATGAACTCTCTGAAAACCTGAAGAC ACCGCAGTCTACTACTGTACTACCCCGTGGGAATGGTCTTGGTACG ATTATTGGGGCCAGGGCACGCTGGTTACGGTGTCTTCCGCTAGCAC AAAGGGCCCTAGCGTGTTCCTCTGGCCCCCAGCAGCAAGAGCACA AGCGGCGGAACAGCCGCCCTGGGCTGCCTCGTGAAGGACTACTTCC CCGAGCCCGTGACAGTGTCTTGGAACAGCGGAGCCCTGACAAGCGG CGTGACACTTTCCCTGCCGTGCTGCAGAGCAGCGGCCTGTACTCC CTGAGCAGCGTGGTCACCGTGCCTAGCAGCAGCCTGGGCACCCAGA CCTACATCTGCAACGTGAACCACAAGCCCAGCAACACCAAAGTGGA CAAGAAGGTGGAGCCCCAAGAGCTGTGATGGCGGAGGAGGGTCCGGA GGCGGAGGATCCGAGGTGCAGCTGCTGGAATCTGGCGGCGGACTGG TGCAGCCTGGCGGATCTCTGAGACTGAGCTGTGCCGCCAGCGGCTT CACCTTCAGCACCTACGCCATGAACTGGGTGCGCCAGGCCCCCTGGC AAAGGCCTGGAATGGGTGTCCCGGATCAGAAGCAAGTACAACAAC ACGCCACCTACTACGCCGACAGCGTGAAGGGCCGGTTACCATCAG CCGGGACGACAGCAAGAACACCCTGTACCTGCAGATGAACAGCCTG CGGGCCGAGGACACCGCCGTGTACTATTGTGTGCGGCACGGCAACT TCGGCAACAGCTATGTGTCTTGGTTTGCCTACTGGGGCCAGGGCAC CCTCGTGACCGTGTCAAGCGCTAGTACCAAGGGCCCCAGCGTGTTC CCCCTGGCACCCAGCAGCAAGAGCACATCTGGCGGAACAGCCGCTC TGGGCTGTCTGGTGAAAGACTACTTCCCCGAGCCCGTGACCGTGTCT TTGGAACCTCTGGCGCCCTGACCAGCGGCGTGCACACCTTCCAGCC GTGCTGCAGAGCAGCGGCCTGTACTCCCTGTCCTCCGTGGTCACCG TGCCCTCTAGCTCCCTGGGAACACAGACATATATCTGTAATGTCAA TCACAAGCCTTCCAACACCAAAGTCGATAAGAAAGTCGAGCCCCAAG AGCTGCGACAAAACCTACACATGCCCACCGTGCCCAGCACCTGAAG CTGCAGGGGGACCGTCAGTCTTCCTCTTCCCCCAAACCCAAGGA CACCTCATGATCTCCCGGACCCCTGAGGTACATGCGTGGTGGTG GACGTGAGCCACGAAGACCCTGAGGTCAAGTTCAACTGGTACGTGG ACGGCGTGGAGGTGCATAATGCCAAGACAAAGCCGCGGGAGGAGCA GTACAACAGCACGTACCGTGTGGTCAGCGTCCTACCGTCCTGCAC CAGGACTGGCTGAATGGCAAGGAGTACAAGTGCAAGGTCTCCAACA AAGCCCTCGGCGCCCCCATCGAGAAAACCATCTCCAAGCCAAAGG GCAGCCCCGAGAACCACAGGTGTACACCCTGCCCCCATGCCGGGAT GAGCTGACCAAGAACCAGGTGAGCCTGTGGTGCCTGGTCAAAGGCT	189

	TCTATCCCAGCGACATCGCCGTGGAGTGGGAGAGCAATGGGCAGCC GGAGAACAACTACAAGACCACGCCTCCCGTGCTGGACTCCGACGGC TCCTTCTTCTCTACAGCAAGCTCACCGTGGACAAGAGCAGGTGGC AGCAGGGGAACGTCTTCTCATGCTCCGTGATGCATGAGGCTCTGCA CAACCACTACACGCAGAAGAGCCTCTCCCTGTCTCCGGGTAA	
VHCH1[16 D5]_Fchole _PGLALA_ HYRF pCON983	GAGGTGCAATTGGTTGAATCTGGTGGTGGTCTGGTAAAACCGGGCG GTTCCCTGCGTCTGAGCTGCGCGGCTTCCGGATTACCTTCTCCAA CGCGTGGATGAGCTGGGTTTCGCCAGGCCCCGGGCAAAGGCCTCGAG TGGGTTGGTTCGTATCAAGTCTAAACTGACGGTGGCACCACGGATT ACGCGGCTCCAGTTAAAGGTCGTTTTACCATTTCCTCGCAGCATAG CAAAAACACTCTGTATCTGCAGATGAACTCTCTGAAAACCTGAAGAC ACCGCAGTCTACTACTGTACTACCCCGTGGGAATGGTCTTGGTACG ATTATTGGGGCCAGGGCACGCTGGTTACGGTGTCTTCCGCTAGCAC CAAGGGCCCCCTCCGTGTTCCCCCTGGCCCCCAGCAGCAAGAGCACC AGCGGCGGCACAGCCGCTCTGGGCTGCCTGGTCAAGGACTACTTCC CCGAGCCCGTGACCGTGTCTTGGAAACAGCGGAGCCCTGACCTCCGG CGTGACACCTTCCCCGCGGTGCTGCAGAGTTCTGGCCTGTATAGC CTGAGCAGCGTGGTCACCGTGCCTTCTAGCAGCCTGGGCACCCAGA CCTACATCTGCAACGTGAACCACAAGCCCAGCAACACCAAGGTGGA CAAGAAGGTGGAGCCCCAAGAGCTGCGACAAAACCTCACACATGCCCA CCGTGCCCAGCACCTGAAGCTGCAGGGGGACCGTCAGTCTTCCTCT TCCCCC AAAACCCAAGGACACCCTCATGATCTCCCGGACCCCTGA GGTCACATGCGTGGTGGTGGACGTGAGCCACGAAGACCCTGAGGTC AAGTTCAACTGGTACGTGGACGGCGTGGAGGTGCATAATGCCAAGA CAAAGCCGCGGGAGGAGCAGTACAACAGCACGTACCGTGTGGTCAG CGTCTCACCGTCTTGCACCAGGACTGGCTGAATGGCAAGGAGTAC AAGTGCAAGGTCTCCAACAAAGCCCTCGGCGCCCCCATCGAGAAAA CCATCTCCAAGGCCAAAGGGCAGCCCCGAGAACCACAGGTGTGCAC CCTGCCCCCATCCCGGGATGAGCTGACCAAGAACCAGGTGAGCCTC TCGTGCGCAGTCAAAGGCTTCTATCCCAGCGACATCGCCGTGGAGT GGGAGAGCAATGGGCAGCCGGAGAACAACCTACAAGACCACGCCTCC CGTGCTGGACTCCGACGGCTCCTTCTTCTCTCGTGAGCAAGCTCACC GTGGACAAGAGCAGGTGGCAGCAGGGGAACGTCTTCTCATGCTCCG TGATGCATGAGGCTCTGCACAACCGCTTCACGCAGAAGAGCCTCTC CCTGTCTCCGGGTAA	190
CD3_comm on light chain pETR13197	CAGGCCGTCTGTGACCCAGGAACCCAGCCTGACAGTGTCTCCTGGCG GCACCGTGACCCTGACATGTGGCAGTTCTACAGGCGCCGTGACCAC CAGCAACTACGCCAACTGGGTGCAGGAAAAGCCCGGCCAGGCCTTC AGAGGACTGATCGGCGGCACCAACAAGAGAGCCCCTGGCACCCCTG CCAGATTGAGCGGATCTCTGCTGGGAGGAAAAGCCCGCCCTGACACT GTCTGGCGCCCAGCCAGAAGATGAGGCCGAGTACTACTGCGCCCTG TGGTACAGCAACCTGTGGGTGTTTCGGCGGAGGCACCAAGCTGACAG TCCTAGGTCAACCCAAGGCTGCCCCAGCGTGACCCTGTTCCCCC CAGCAGCGAGGAACTGCAGGCCAACAAGGCCACCCTGGTCTGCCTG ATCAGCGACTTCTACCCAGGCGCCGTGACCGTGGCCTGGAAGGCCG ACAGCAGCCCCGTGAAGGCCGCGTGGAGACCACCACCCCCAGCAA GCAGAGCAACAACAAGTACGCCGCCAGCAGCTACCTGAGCCTGACC CCCGAGCAGTGGAAGAGCCACAGGTCCTACAGCTGCCAGGTGACCC ACGAGGGCAGCACCGTGGAGAAAACCGTGGCCCCCACCAGGTGCAG C	191
VHCH1[C D3]_VHCH	GAGGTGCAGCTGCTGGAATCTGGCGGCGGACTGGTGCAGCCTGGCG GATCTCTGAGACTGAGCTGTGCCGCCAGCGGCTTCACCTTCAGCAC CTACGCCATGAACTGGGTGCGCCAGGCCCCCTGGCAAAGGCCTGGAA	192

<p>I[16D5]_Fc knob_PGL ALA pETR13932 (Classical TCB with 16D5; 2+1: pETR13932+ pCON983 + pETR13197)</p>	<p>TGGGTGTCCCGGATCAGAAGCAAGTACAACAACACTACGCCACCTACT ACGCCGACAGCGTGAAGGGCCGGTTTACCATCAGCCGGGACGACAG CAAGAACACCCTGTACCTGCAGATGAACAGCCTGCGGGCCGAGGAC ACCGCCGTGTACTATTGTGTGCGGCACGGCAACTTCGGCAACAGCT ATGTGTCTTGGTTTGCCTACTGGGGCCAGGGCACCCCTCGTGACCGT GTCATCTGCTAGCACAAAGGGCCCTAGCGTGTTCCCTCTGGCCCCC AGCAGCAAGAGCACAAAGCGGCGGAACAGCCGCCCTGGGCTGCCTCG TGAAGGACTACTTCCCCGAGCCCGTGACAGTGTCTTGGAACAGCGG AGCCCTGACAAGCGGCGTGACACACCTTCCCTGCCGTGCTGCAGAGC AGCGGCCTGTACTCCCTGAGCAGCGTGGTCACCGTGCTAGCAGCA GCCTGGGCACCCAGACCTACATCTGCAACGTGAACCACAAGCCAG CAACACCAAAGTGGACAAGAAGGTGGAGCCCAAGAGCTGTGATGGC GGAGGAGGGTCCGGAGGCGGAGGATCCGAGGTGCAATTGGTTGAAT CTGGTGGTGGTCTGGTAAAACCGGGCGGTTCCCTGCGTCTGAGCTG CGCGGCTTCCGGATTACCTTCTCCAACGCGTGGATGAGCTGGGTT CGCCAGGCCCCGGGCAAAGGCCCTCGAGTGGGTTGGTTCGTATCAAGT CTAAACTGACGGTGGCACCACGGATTACGCGGCTCCAGTTAAAGG TCGTTTTACCATTTCGCGACGATAGCAAAAACACTCTGTATCTG CAGATGAACTCTCTGAAACTGAAGACACCGCAGTCTACTACTGTA CTACCCCGTGGAATGGTCTTGGTACGATTATTGGGGCCAGGGCAC GCTGGTTACGGTGTCTAGCGCTAGTACCAAGGGCCCCAGCGTGTTT CCCCCTGGCACCCAGCAGCAAGAGCACATCTGGCGGAACAGCCGCTC TGGGCTGTCTGGTGAAAGACTACTTCCCCGAGCCCGTGACCGTGTC TTGGAACCTCTGGCGCCCTGACCAGCGGCGTGACACCTTTCAGCC GTGCTGCAGAGCAGCGGCCTGTACTCCCTGTCTTCCGTGGTCACCG TGCCCTCTAGCTCCCTGGGAACACAGACATATCTGTAATGTCAA TCACAAGCCTTCCAACACCAAAGTCGATAAGAAAGTCGAGCCCAAG AGCTGCGACAAAACCTCACACATGCCCACCGTGCCCAGCACCTGAAG CTGCAGGGGGACCGTCAGTCTTCTCTTCCCCCAAAACCCAAGGA CACCTCATGATCTCCCGGACCCCTGAGGTACATGCGTGGTGGTG GACGTGAGCCACGAAGACCCTGAGGTCAAGTTCAACTGGTACGTGG ACGGCGTGGAGGTGCATAATGCCAAGACAAAGCCGCGGGAGGAGCA GTACAACAGCACGTACCGTGTGGTACGCGTCTCACCGTCTGCAC CAGGACTGGCTGAATGGCAAGGAGTACAAGTGCAAGGTCTCCAACA AAGCCCTCGGCGCCCCCATCGAGAAAACCATCTCCAAGCCAAAGG GCAGCCCCGAGAACCACAGGTGTACACCCTGCCCCCATGCCGGGAT GAGCTGACCAAGAACCAGGTGACGCTGTGGTGCCTGGTCAAAGGCT TCTATCCCAGCGACATCGCCGTGGAGTGGGAGAGCAATGGGCAGCC GGAGAACAACCTACAAGACCACGCCTCCCGTGCTGGACTCCGACGGC TCCTTCTTCTCTACAGCAAGCTCACCGTGGACAAGAGCAGGTGGC AGCAGGGGAACGTCTTCTCATGCTCCGTGATGCATGAGGCTCTGCA CAACCACTACACGCAGAAGAGCCTCTCCCTGTCTCCGGGTAAA</p>	
<p>VHCH1[C D3]_Fckno b_PGLALA pETR13719 (16D5 IgG format, 1+1: pETR13719 + pCON983 + pETR13197)</p>	<p>GAGGTGCAGCTGCTGGAATCTGGCGGCGGACTGGTGCAGCCTGGCG GATCTCTGAGACTGAGCTGTGCCGCCAGCGGCTTCACCTTCAGCAC CTACGCCATGAACTGGGTGCGCCAGGCCCCCTGGCAAAGGCCTGGAA TGGGTGTCCCGGATCAGAAGCAAGTACAACAACACTACGCCACCTACT ACGCCGACAGCGTGAAGGGCCGGTTTACCATCAGCCGGGACGACAG CAAGAACACCCTGTACCTGCAGATGAACAGCCTGCGGGCCGAGGAC ACCGCCGTGTACTATTGTGTGCGGCACGGCAACTTCGGCAACAGCT ATGTGTCTTGGTTTGCCTACTGGGGCCAGGGCACCCCTCGTGACCGT GTCATCTGCTAGCACCAAGGGCCCATCGGTCTTCCCCCTGGCACCC TCCTCCAAGAGCACCTCTGGGGGCACAGCGGCCCTGGGCTGCCTGG TCAAGGACTACTTCCCCGAACCGGTGACGGTGTCTGTGGAACCTCAGG CGCCCTGACCAGCGGCGTGACACACCTTCCCGGCTGTCTACAGTCC</p>	<p>193</p>

	TCAGGACTCTACTCCCTCAGCAGCGTGGTGACCGTGCCCTCCAGCA GCTTGGGCACCCAGACCTACATCTGCAACGTGAATCACAAGCCCAG CAACACCAAGGTGGACAAGAAAGTTGAGCCCAAATCTTGTGACAAA ACTCACACATGCCACCGTGCCAGCACCTGAAGCTGCAGGGGGAC CGTCAGTCTTCTCTTCCCCCAAAACCCAAAGGACACCCCTCATGAT CTCCCGGACCCCTGAGGTACATGCGTGGTGGTGGACGTGAGCCAC GAAGACCCTGAGGTCAAGTTCAACTGGTACGTGGACGGCGTGGAGG TGCATAATGCCAAGACAAAGCCGCGGGAGGAGCAGTACAACAGCAC GTACCGTGTGGTCAGCGTCCTCACCGTCCTGCACCAGGACTGGCTG AATGGCAAGGAGTACAAGTGCAAGGTCTCCAACAAAGCCCTCGGCG CCCCATCGAGAAAACCATCTCCAAGCCAAAGGGCAGCCCCGAGA ACCACAGGTGTACACCCTGCCCCATGCCGGGATGAGCTGACCAAG AACCAGGTGAGCCTGTGGTGCCTGGTCAAAGGCTTCTATCCCAGCG ACATCGCCGTGGAGTGGGAGAGCAATGGGCAGCCGGAGAACAACTA CAAGACCACGCCTCCCGTGCTGGACTCCGACGGCTCCTTCTTCTC TACAGCAAGCTCACCGTGGACAAGAGCAGGTGGCAGCAGGGGAACG TCTTCTCATGCTCCGTGATGCATGAGGCTCTGCACAACCACTACAC GCAGAAGAGCCTCTCCCTGTCTCCGGGTAAA	
Fc_hole_PG LALA_HYR F pETR10755 (16D5 Head- to-tail, 1+1: pCON999 + pETR10755 + pETR13197)	GACAAAACCTCACACATGCCACCGTGCCAGCACCTGAATCCTGG GGGACCGTCAGTCTTCTCTTCCCCCAAAACCCAAAGGACACCCCT CATGATCTCCCGGACCCCTGAGGTACATGCGTGGTGGTGGACGTG AGCCACGAAGACCCTGAGGTCAAGTTCAACTGGTACGTGGACGGCG TGGAGGTGCATAATGCCAAGACAAAGCCGCGGGAGGAGCAGTACAA CAGCACGTACCGTGTGGTCAGCGTCCTCACCGTCCTGCACCAGGAC TGGCTGAATGGCAAGGAGTACAAGTGCAAGGTCTCCAACAAAGCCC TCCCAGCCCCCATCGAGAAAACCATCTCCAAGCCAAAGGGCAGCC CCGAGAACCACAGGTGTGCACCCTGCCCCCATCCCGGGATGAGCTG ACCAAGAACCAGGTGAGCCTCTCGTGCGCAGTCAAAGGCTTCTATC CCAGCGACATCGCCGTGGAGTGGGAGAGCAATGGGCAGCCGGAGAA CAACTACAAGACCACGCCTCCCGTGCTGGACTCCGACGGCTCCTTC TTCTCGTGAGCAAGCTCACCGTGGACAAGAGCAGGTGGCAGCAGG GGAACGTCTTCTCATGCTCCGTGATGCATGAGGCTCTGCACAACCG CTTCACGCAGAAGAGCCTCTCCCTGTCTCCGGGTAAA	194
VHCH1[9D 11]_VHCL[CD3]_Fckn ob_PGLAL A pCON1057 (9D11 inverted format, 2+1: pCON1057 + pCON1051 + pCON1063 + pETR12940)	CAGGTGCAATTGGTTCAATCTGGTGCTGAAGTAAAAAAACCGGGCG CTTCCGTTAAAGTGAGCTGCAAAGCATCCGGATACACCTTCACTTC CTATTACATGCACTGGGTTTCGTCAAGCCCCGGGCCAGGGTCTGGAA TGGATGGGCATCATTAACCCAAGCGGTGGCCCTACCTCCTACGCGC AGAAATTCCAGGGTCGCGTCACGATGACCCGTGACACTAGCACCTC TACCGTTTATATGGAGCTGTCCAGCCTGCGTTCTGAAGATACTGCA GTGTACTACTGTGCACGCGGTGACTTCGCTTGGCTGGACTATTGGG GTCAAGGCACCCCTCGTAACGGTTTCTTCTGCTAGCACAAAGGGCCC CAGCGTGTTCCCTCTGGCCCCCTAGCAGCAAGAGCACATCTGGCGGA ACAGCCGCCCTGGGCTGCCTCGTGAAGGACTACTTTCCCGAGCCTG TGACCGTGTCTCTGGAACCTCTGGCGCCCTGACAAGCGGCGTGCACAC CTTTCCAGCCGTGCTGCAGAGCAGCGGCCCTGTACTCTCTGAGCAGC GTGGTCACCGTGCCTAGCAGCAGCCTGGGCACCCAGACCTACATCT GCAACGTGAACCACAAGCCCAGCAACACCAAAGTGGACAAGAAGGT GGAGCCCAAGAGCTGTGATGGCGGAGGAGGGTCCGGAGGCGGAGGA TCCGAGGTGCAGCTGCTGGAATCTGGCGGCGGACTGGTGCAGCCTG GCGGATCTCTGAGACTGAGCTGTGCCGCCAGCGGCTTACCTTCAG CACCTACGCCATGAACTGGGTGCGCCAGGCCCTGGCAAAGGCCTG GAATGGGTGTCCCGGATCAGAAGCAAGTACAACAACCTACGCCACCT ACTACGCCGACAGCGTGAAGGGCCGGTTACCATCAGCCGGGACGA CAGCAAGAACACCCTGTACCTGCAGATGAACAGCCTGCGGGCCGAG	195

	<p>GACACCGCCGTGTACTATTGTGTGCGGCACGGCAACTTCGGCAACA GCTATGTGTCTTGGTTTGCTACTGGGGCCAGGGCACCTCTGTGAC CGTGTCAAGCGCTAGTGTGGCCGCTCCCTCCGTGTTTATCTTTCCC CCATCCGATGAACAGCTGAAAAGCGGCACCGCCTCCGTCTGTGTCTC TGCTGAACAATTTTTACCCTAGGGAAGCTAAAGTGCAGTGGAAGT GGATAACGCACTGCAGTCCGGCAACTCCCAGGAATCTGTGACAGAA CAGGACTCCAAGGACAGCACCTACTCCCTGTCCTCCACCCTGACAC TGTCTAAGGCTGATTATGAGAAACACAAAGTCTACGCCTGCGAAGT CACCCATCAGGGCCTGAGCTCGCCCGTCACAAAGAGCTTCAACAGG GGAGAGTGTGACAAGACCCACACCTGTCCCCCTTGTCTGCCCCCTG AAGCTGCTGGCGGCCCTTCTGTGTTCTGTTCCTGTTCCTCCCAAGCCAA GGACACCCTGATGATCAGCCGGACCCCCGAAGTGACCTGCGTGGTG GTGGATGTGTCCACGAGGACCCTGAAGTGAAGTTCAATTGGTACG TGGACGGCGTGGAAGTGCACAACGCCAAGACAAAGCCGCGGGAGGA GCAGTACAACAGCACGTACCGTGTGGTCAGCGTCCTCACCGTCCTG CACCAGGACTGGCTGAATGGCAAGGAGTACAAGTGAAGTCTCCA ACAAAGCCCTCGGCGCCCCCATCGAGAAAACCATCTCCAAAGCCAA AGGGCAGCCCCGAGAACCACAGGTGTACACCCTGCCCCCATGCCGG GATGAGCTGACCAAGAACCAGGTGAGCCTGTGGTGCCTGGTCAAAG GCTTCTATCCCAGCGACATCGCCGTGGAGTGGGAGAGCAATGGGCA GCCGGAGAACAACACTACAAGACCACGCCTCCCGTGCTGGACTCCGAC GGCTCCTTCTTCTCTACAGCAAGCTCACCGTGGACAAGAGCAGGT GGCAGCAGGGGAACGTCTTCTCATGCTCCGTGATGCATGAGGCTCT GCACAACCACTACACGCAGAAGAGCCTCTCCCTGTCTCCGGGTAAA</p>	
9D11_Fchol e_PGLALA _HYRF pCON1051	<p>CAGGTGCAATTGGTTCAATCTGGTGTCTGAAGTAAAAAACCAGGGCG CTTCCGTAAAGTGAGCTGCAAAGCATCCGGATACACCTTCACTTC CTATTACATGCACTGGGTTTCGTCAAGCCCCGGGCCAGGGTCTGGAA TGGATGGGCATCATTAAACCAAGCGGTGGCCCTACCTCCTACGCGC AGAAATTCCAGGGTCGCGTCACGATGACCCGTGACACTAGCACCTC TACCGTTTATATGGAGCTGTCCAGCCTGCGTTCTGAAGATACTGCA GTGTACTACTGTGCACGCGGTGACTTCGCTTGGCTGGACTATTGGG GTCAAGGCACCCCTCGTAACGGTTTCTTCTGCTAGCACCAAGGGCCC CTCCGTGTTCCCCCTGGCCCCCAGCAGCAAGAGCACCCAGCGCGGC ACAGCCGCTCTGGGCTGCCTGGTCAAGGACTACTTCCCCGAGCCCG TGACCGTGTCTTGGAACAGCGGAGCCCTGACCTCCGGCGTGCACAC CTTCCCCGCCGTGCTGCAGAGTTCTGGCCTGTATAGCCTGAGCAGC GTGGTCACCGTGCCTTCTAGCAGCCTGGGCACCCAGACCTACATCT GCAACGTGAACCACAAGCCCAGCAACACCAAGGTGGACAAGAAGGT GGAGCCCCAAGAGCTGCGACAAAACCTCACACATGCCACCGTGCCCA GCACCTGAAGCTGCAGGGGGACCGTCAGTCTTCTCTTCCCCCAA AACCAAGGACACCCTCATGATCTCCCGGACCCCTGAGGTACATG CGTGGTGGTGGACGTGAGCCACGAAGACCCTGAGGTCAAGTTCAAC TGGTACGTGGACGGCGTGGAGGTGCATAATGCCAAGACAAAGCCGC GGGAGGAGCAGTACAACAGCACGTACCGTGTGGTCAGCGTCCTCAC CGTCCTGCACCAGGACTGGCTGAATGGCAAGGAGTACAAGTGCAAG GTCTCCAACAAAGCCCTCGGCGCCCCCATCGAGAAAACCATCTCCA AAGCCAAAGGGCAGCCCCGAGAACCACAGGTGTGCACCCTGCCCCC ATCCCGGGATGAGCTGACCAAGAACCAGGTGAGCCTCTCGTGCGCA GTCAAAGGCTTCTATCCCAGCGACATCGCCGTGGAGTGGGAGAGCA ATGGGCAGCCGGAGAACAACACTACAAGACCACGCCTCCCGTGCTGGA CTCCGACGGCTCCTTCTTCTCTGAGCAAGCTCACCGTGGACAAG AGCAGGTGGCAGCAGGGGAACGTCTTCTCATGCTCCGTGATGCATG AGGCTCTGCACAACCACTACACGCAGAAGAGCCTCTCCCTGTCTCC GGGTAAA</p>	196

9D11_LC pCON1063	GATATTGTTATGACTCAATCTCCACTGTCTCTGCCGGTGACTCCAG GCGAACCGGCGAGCATTCTTGCCGTTCCAGCCAGTCTCTGCTGCA CTCCAACGGCTACAACCTATCTCGATTGGTACCTGCAAAAACCGGT CAGAGCCCTCAGCTGCTGATCTACCTGGGCTCTAACCGCGCTTCCG GTGTACCGGACCGTTTCAGCGGCTCTGGATCCGGCACCGATTTCAC GTTGAAAATCAGCCGTGTTGAAGCAGAAGACGTGGGCGTTTATTAC TGTATGCAGGCAAGCATTATGAACCGGACTTTTGGTCAAGGCACCA AGGTGCGAAATTAAACGTACGGTGGCTGCACCATCTGTCTTCATCTT CCCGCCATCTGATGAGCAGTTGAAATCTGGAAGTGCCTCTGTTGTG TGCTGCTGAATAACTTCTATCCCAGAGAGGCCAAAGTACAGTGGA AGGTGGATAACGCCCTCCAATCGGGTAACTCCCAGGAGAGTGTAC AGAGCAGGACAGCAAGGACAGCACCTACAGCCTCAGCAGCACCTG ACGCTGAGCAAAGCAGACTACGAGAAACACAAAGTCTACGCCTGCG AAGTCACCCATCAGGGCCTGAGCTCGCCCGTCACAAAGAGCTTCAA CAGGGGAGAGTGT	197
VLCH1[CD 3] pETR12940	CAGGCCGTCGTGACCCAGGAACCCAGCCTGACAGTGTCTCCTGGCG GCACCGTGACCTGACATGTGGCAGTTCTACAGGCGCCGTGACCAC CAGCAACTACGCCAACTGGGTGCAGGAAAAGCCCGGCCAGGCCTTC AGAGGACTGATCGGCGGCACCAACAAGAGAGCCCCCTGGCACCCCTG CCAGATTGAGCGGATCTCTGCTGGGAGGAAAGGCCGCCCTGACACT GTCTGGCGCCAGCCAGAAGATGAGGCCGAGTACTACTGCGCCCTG TGGTACAGCAACCTGTGGGTGTTTCGGCGGAGGCACCAAGCTGACAG TGCTGAGCAGCGCTTCCACCAAAGGCCCTTCCGTGTTTCTCTGGC TCCTAGCTCCAAGTCCACCTCTGGAGGCACCGCTGCTCTCGGATGC CTCGTGAAGGATTATTTTCTGAGCCTGTGACAGTGTCTTGGGAATA GCGGAGCACTGACCTCTGGAGTGCATACTTTCCCCGCTGTGCTGCA GTCTCTGGACTGTACAGCCTGAGCAGCGTGGTGACAGTGCCCAGC AGCAGCCTGGGCACCCAGACCTACATCTGCAACGTGAACCACAAGC CCAGCAACACCAAGGTGGACAAGAAGGTGGAACCCAAGTCTTGT	198
VHCL[CD3]_Fcknob_P GLALA pETR13378 (9D11 CrossMab format, 1+1: pETR13378 + pCON1051 + pCON1063 + pETR12940)	GAGGTGCAGCTGCTGGAATCTGGCGGCGGACTGGTGCAGCCTGGCG GATCTCTGAGACTGAGCTGTGCCGCCAGCGGCTTCACCTTCAGCAC CTACGCCATGAACTGGGTGCGCCAGGCCCCCTGGCAAAGGCCTGGAA TGGGTGTCCCGGATCAGAAGCAAGTACAACAACCTACGCCACCTACT ACGCCGACAGCGTGAAGGGCCGGTTCACCATCAGCCGGGACGACAG CAAGAACACCCCTGTACCTGCAGATGAACAGCCTGCGGGCCGAGGAC ACCGCCGTGTACTATTGTGTGCGGCACGGCAACTTCGGCAACAGCT ATGTGTCTTGGTTTGCCTACTGGGGCCAGGGCACCCCTCGTGACCGT GTCATCTGCTAGCGTGGCCGCTCCCTCCGTGTTTATCTTTCCCCCA TCCGATGAACAGCTGAAAAGCGGCACCGCCTCCGTGCTGTGTCTGC TGAACAATTTTTACCCTAGGGAAGCTAAAGTGCAGTGGAAGTGGA TAACGCACTGCAGTCCGGCAACTCCCAGGAATCTGTGACAGAACAG GACTCCAAGGACAGCACCTACTCCCTGTCTCCACCCTGACACTGT CTAAGGCTGATTATGAGAAACACAAAGTCTACGCCTGCGAAGTCAC CCATCAGGGCCTGAGCTCGCCCGTCACAAAGAGCTTCAACAGGGGA GAGTGTGACAAGACCCACACCTGTCCCCCTTGTCTGCCCCCTGAAG CTGCTGGCGGCCCTTCTGTGTTTCTGTTCCCCCAAGCCCAAGGA CACCTGATGATCAGCCGGACCCCCGAAGTGACCTGCGTGGTGGTG GATGTGTCCACGAGGACCCTGAAGTGAAGTTCAATTGGTACGTGG ACGGCGTGGAAGTGACAACGCCAAGACAAAGCCGCGGGAGGAGCA GTACAACAGCACGTACCGTGTGGTCAGCGTCTCACCCTCCTGCAC CAGGACTGGCTGAATGGCAAGGAGTACAAGTGCAAGGTCTCCAACA AAGCCCTCGGCGCCCCCATCGAGAAAACCATCTCCAAGCCAAAGG GCAGCCCCGAGAACCACAGGTGTACACCCTGCCCCCATGCCGGGAT GAGCTGACCAAGAACCAGGTGAGCCTGTGGTGCCTGGTCAAAGGCT	199

	TCTATCCCAGCGACATCGCCGTGGAGTGGGAGAGCAATGGGCAGCC GGAGAACAACACTACAAGACCACGCCTCCCGTGCTGGACTCCGACGGC TCCTTCTTCTCTACAGCAAGCTCACCGTGGACAAGAGCAGGTGGC AGCAGGGGAACGTCTTCTCATGCTCCGTGATGCATGAGGCTCTGCA CAACCACTACACGCAGAAGAGCCTCTCCCTGTCTCCGGGTAAA	
16D5 inverted 2+1 with N100A in CDR H3 pETR14096 (pETR14096 + pCON983 + pETR13197)	GAGGTGCAATTGGTTGAATCTGGTGGTGGTCTGGTAAAACCGGGCG GTTCCCTGCGTCTGAGCTGCGCGGCTTCCGGATTACCTTCTCCAA CGCGTGGATGAGCTGGGTTTCGCCAGGCCCGGGCAAAGGCCTCGAG TGGGTTGGTTCGTATCAAGTCTAAAACCTGACGGTGGCACCACGGATT ACGCGGCTCCAGTTAAAGGTCGTTTTACCATTTCCCGCGACGATAG CAAAAACACTCTGTATCTGCAGATGAACTCTCTGAAAACCTGAAGAC ACCGCAGTCTACTACTGTACTACCCCGTGGGAATGGTCTTGGTACG ATTATTGGGGCCAGGGCACGCTGGTTACGGTGTCTTCCGCTAGCAC AAAGGGCCCTAGCGTGTTCCCTCTGGCCCCCAGCAGCAAGAGCACA AGCGGCGGAACAGCCGCCCTGGGCTGCCTCGTGAAGGACTACTTCC CCGAGCCCGTGACAGTGTCTTGGAACAGCGGAGCCCTGACAAGCGG CGTGACACTTTCCCTGCCGTGCTGCAGAGCAGCGGCCTGTACTCC CTGAGCAGCGTGGTCACCGTGCCTAGCAGCAGCCTGGGCACCCAGA CCTACATCTGCAACGTGAACCACAAGCCCAGCAACACCAAAGTGGA CAAGAAGGTGGAGCCCAAGAGCTGTGATGGCGGAGGAGGGTCCGGA GGCGGAGGATCCGAGGTGCAGCTGCTGGAATCTGGCGGCGGACTGG TGCAGCCTGGCGGATCTCTGAGACTGAGCTGTGCCGCCAGCGGCTT CACCTTCAGCACCTACGCCATGAACTGGGTGCGCCAGGCCCTGGC AAAGGCCTGGAATGGGTGTCCCGGATCAGAAGCAAGTACAACAAC ACGCCACCTACTACGCCGACAGCGTGAAGGGCCGGTTACCATCAG CCGGGACGACAGCAAGAACACCCTGTACCTGCAGATGAACAGCCTG CGGGCCGAGGACACCGCCGTGTACTATTGTGTGCGGCACGGCAACT TCGGCGCCAGCTATGTGTCTTGGTTTGCCTACTGGGGCCAGGGCAC CCTCGTGACCGTGTCAAGCGCTAGTACCAAGGGCCCCAGCGTGTTC CCCCTGGCACCCAGCAGCAAGAGCACATCTGGCGGAACAGCCGCTC TGGGCTGTCTGGTGAAGACTACTTCCCCGAGCCCGTGACCGTGTCT TTGGAACCTCTGGCGCCCTGACCAGCGGCGTGCACACCTTTCCAGCC GTGCTGCAGAGCAGCGGCCTGTACTCCCTGTCTCCGTGGTACCG TGCCCTCTAGCTCCCTGGGAACACAGACATATATCTGTAATGTCAA TCACAAGCCTTCCAACACCAAAGTCGATAAGAAAGTCGAGCCCAAG AGCTGCGACAAAACCTCACACATGCCACCGTGCCAGCACCTGAAG CTGCAGGGGGACCGTCAGTCTTCTCTTCCCCCAAACCCAAGGA CACCTCATGATCTCCCGGACCCCTGAGGTACATGCGTGGTGGTG GACGTGAGCCACGAAGACCCTGAGGTCAAGTTCAACTGGTACGTGG ACGGCGTGGAGGTGCATAATGCCAAGACAAAGCCGCGGGAGGAGCA GTACAACAGCACGTACCGTGTGGTCAGCGTCCTACCGTCTCTGCAC CAGGACTGGCTGAATGGCAAGGAGTACAAGTGCAAGGTCTCCAACA AAGCCCTCGGCGCCCCCATCGAGAAAACCATCTCCAAAGCCAAAGG GCAGCCCCGAGAACCACAGGTGTACACCCTGCCCCCATGCCGGGAT GAGCTGACCAAGAACCAGGTACGCCTGTGGTGCCTGGTCAAAGGCT TCTATCCCAGCGACATCGCCGTGGAGTGGGAGAGCAATGGGCAGCC GGAGAACAACACTACAAGACCACGCCTCCCGTGCTGGACTCCGACGGC TCCTTCTTCTCTACAGCAAGCTCACCGTGGACAAGAGCAGGTGGC AGCAGGGGAACGTCTTCTCATGCTCCGTGATGCATGAGGCTCTGCA CAACCACTACACGCAGAAGAGCCTCTCCCTGTCTCCGGGTAAA	200
16D5 inverted 2+1 with	GAGGTGCAATTGGTTGAATCTGGTGGTGGTCTGGTAAAACCGGGCG GTTCCCTGCGTCTGAGCTGCGCGGCTTCCGGATTACCTTCTCCAA CGCGTGGATGAGCTGGGTTTCGCCAGGCCCGGGCAAAGGCCTCGAG TGGGTTGGTTCGTATCAAGTCTAAAACCTGACGGTGGCACCACGGATT	201

<p>S100aA in CDR H3 pETR14097 (pETR14097 + pCON983 + pETR13197)</p>	<p>ACGCGGCTCCAGTTAAAGGTCGTTTTACCATTTCCCGCGACGATAG CAAAAACACTCTGTATCTGCAGATGAACTCTCTGAAAACCTGAAGAC ACCGCAGTCTACTACTGTACTACCCCGTGGGAATGGTCTTGGTACG ATTATTGGGGCCAGGGCACGCTGGTTACGGTGTCTTCCGCTAGCAC AAAGGGCCCTAGCGTGTTCCCTCTGGCCCCCAGCAGCAAGAGCACA AGCGGCGGAACAGCCGCCCTGGGCTGCCTCGTGAAGGACTACTTCC CCGAGCCCGTGACAGTGTCTTGGAACAGCGGAGCCCTGACAAGCGG CGTGACACTTTCCCTGCCGTGCTGCAGAGCAGCGGCCTGTACTCC CTGAGCAGCGTGGTCACCGTGCCTAGCAGCAGCCTGGGCACCCAGA CCTACATCTGCAACGTGAACCACAAGCCCAGCAACACCAAAGTGGA CAAGAAGGTGGAGCCCAAGAGCTGTGATGGCGGAGGAGGGTCCGGA GGCGGAGGATCCGAGGTGCAGCTGCTGGAATCTGGCGGCGGACTGG TGCAGCCTGGCGGATCTCTGAGACTGAGCTGTGCCGCCAGCGGCTT CACCTTCAGCACCTACGCCATGAACTGGGTGCGCCAGGCCCCCTGGC AAAGGCCTGGAATGGGTGTCCCGGATCAGAAGCAAGTACAACAAC ACGCCACCTACTACGCCGACAGCGTGAAGGGCCGGTTCACCATCAG CCGGGACGACAGCAAGAACACCCTGTACCTGCAGATGAACAGCCTG CGGGCCGAGGACACCGCCGTGTACTATTGTGTGCGGCACGGCAACT TCGGCAACGCCTATGTGTCTTGGTTTGCTACTGGGGCCAGGGCAC CCTCGTGACCGTGTCAAGCGCTAGTACCAAGGGCCCCAGCGTGTTT CCCCCTGGCACCCAGCAGCAAGAGCACATCTGGCGGAACAGCCGCTC TGGGCTGTCTGGTGAAAGACTACTTCCCCGAGCCCGTGACCGTGT TTGGAACCTCTGGCGCCCTGACCAGCGGCGTGCACACCTTTCAGCC GTGCTGCAGAGCAGCGGCCTGTACTCCCTGTCCTCCGTGGTCACCG TGCCCTCTAGCTCCCTGGGAACACAGACATATATCTGTAATGTCAA TCACAAGCCTTCCAACACCAAAGTCGATAAGAAAGTCGAGCCCAAG AGCTGCGACAAAACCTCACACATGCCCACCGTGCCAGCACCTGAAG CTGCAGGGGGACCGTCAGTCTTCTCTTCCCCCAGGACCAAGGA CACCTCATGATCTCCCGGACCCCTGAGGTACATGCGTGGTGGTG GACGTGAGCCACGAAGACCCTGAGGTCAAGTTCAACTGGTACGTGG ACGGCGTGGAGGTGCATAATGCCAAGACAAAGCCGCGGGAGGAGCA GTACAACAGCACGTACCGTGTGGTCAGCGTCTCACCCTCCTGCAC CAGGACTGGCTGAATGGCAAGGAGTACAAGTGCAAGGTCTCCAACA AAGCCCTCGGCGCCCCCATCGAGAAAACCATCTCCAAAGCCAAAGG GCAGCCCCGAGAACCACAGGTGTACACCCTGCCCCCATGCCGGGAT GAGCTGACCAAGAACCAGGTGAGCCTGTGGTGCCTGGTCAAAGGCT TCTATCCCAGCGACATCGCCGTGGAGTGGGAGAGCAATGGGCAGCC GGAGAACAACCTACAAGACCACGCCTCCCGTGCTGGACTCCGACGGC TCCTTCTTCTCTACAGCAAGCTCACCGTGGACAAGAGCAGGTGGC AGCAGGGGAACGTCTTCTCATGCTCCGTGATGCATGAGGCTCTGCA CAACCACTACACGCAGAAGAGCCTCTCCCTGTCTCCGGGTAAA</p>	
<p>CD3 light chain fused to CH1; Fc_PGLAL A; pETR13862 (Kappa- lambda antibody with CD3 common</p>	<p>CAGGCCGTGCTGACCCAGGAACCCAGCCTGACAGTGTCTCCTGGCG GCACCGTGACCCTGACATGTGGCAGTTCTACAGGCGCCGTGACCAC CAGCAACTACGCCAAGTGGGTGCAGGAAAAGCCCGGCCAGGCCTTC AGAGGACTGATCGGCGGCACCAACAAGAGAGCCCTGGCACCCCTG CCAGATTGAGCGGATCTCTGCTGGGAGGAAAAGGCCGCCCTGACACT GTCTGGCGCCCAGCCAGAAGATGAGGCCGAGTACTACTGCGCCCTG TGGTACAGCAACCTGTGGGTGTTTCGGCGGAGGCACCAAGCTGACAG TGCTGAGCAGCGCTAGCACCAAGGGCCCATCGGTCTTCCCCCTGGC ACCTCCTCCAAGAGCACCTCTGGGGGCACAGCGGCCCTGGGCTGC CTGGTCAAGGACTACTTCCCCGAACCGGTGACGGTGTCTGGAAC CAGGCGCCCTGACCAGCGCGTGCACACCTTCCCGGCTGTCTTACA GTCTCAGGACTCTACTCCCTCAGCAGCGTGGTGACCGTGCCCTCC AGCAGCTTGGGCACCCAGACCTACATCTGCAACGTGAATCACAAGC</p>	<p>202</p>

light chain fused to CH1 + Fc_PGLALA. VHs fused to kappa or lambda constant chain pETR13859 + pETR13860 + pETR13862)	CCAGCAACACCAAGGTGGACAAGAAAGTTGAGCCCCAAATCTTGTGA CAAAACTCACACATGCCCCACCGTGCCCAGCACCTGAAGCTGCAGGG GGACCGTCAGTCTTCCCTCTTCCCCCAAACCCAAGGACACCCTCA TGATCTCCCGGACCCCTGAGGTCACATGCGTGGTGGTGGACGTGAG CCACGAAGACCCCTGAGGTCAAGTTCAACTGGTACGTGGACGGCGTG GAGGTGCATAATGCCAAGACAAAGCCGCGGGAGGAGCAGTACAACA GCACGTACCGTGTGGTCAGCGTCTCTACCGTCTCTGCACCAGGACTG GCTGAATGGCAAGGAGTACAAGTGCAAGGTCTCCAACAAAGCCCTC GGCGCCCCCATCGAGAAAACCATCTCCAAGCCAAAGGGCAGCCCC GAGAACCA CAGGTGTACACCCTGCCCCCATCCCGGGATGAGCTGACCAAGAACC AGGTCAGCCTGACCTGCCTGGTCAAAGGCTTCTATCCCAGCGACAT CGCCGTGGAGTGGGAGAGCAATGGGCAGCCGGAGAACAACATAAG ACCACGCCTCCCGTGTGGACTCCGACGGCTCCTTCTTCTCTACA GCAAGCTCACCGTGGACAAGAGCAGGTGGCAGCAGGGGAACGTCTT CTCATGCTCCGTGATGCATGAGGCTCTGCACAACCACTACACGCAG AAGAGCCTCTCCCTGTCTCCGGGTAAA	
16D5 VH fused to constant kappa chain; pETR13859	GAGGTGCAATTGGTTGAATCTGGTGGTGGTCTGGTAAAACCGGGCG GTTCCCTGCGTCTGAGCTGCGCGGCTTCCGGATTACCTTCTCCAA CGCGTGGATGAGCTGGGTTTCGCCAGGCCCGGGCAAAGGCCTCGAG TGGGTTGGTTCGTATCAAGTCTAAAAGTACGGTGGCACCACGGATT ACGCGGCTCCAGTTAAAGGTCGTTTTACCATTTCGCGACGATAG CAAAAACACTCTGTATCTGCAGATGAACTCTCTGAAAAGTGAAGAC ACCGCAGTCTACTACTGTACTACCCCGTGGGAATGGTCTTGGTACG ATTATTGGGGCCAGGGCACGCTGGTTACGGTGTCTTCCGCTAGCGT GGCCGCTCCCTCCGTGTTTCATCTTCCACCTTCCGACGAGCAGCTG AAGTCCGGCACCGCTTCTGTCTGTGCCTGCTGAACAACCTTCTACC CCCGCGAGGCCAAGGTGCAGTGGAAGGTGGACAACGCCCTGCAGTC CGGCAACAGCCAGGAATCCGTGACCGAGCAGGACTCCAAGGACAGC ACCTACTCCCTGTCTTCCACCTGACCCTGTCCAAGGCCGACTACG AGAAGCACAAGGTGTACGCTGCGAAGTGACCCACCAGGGCCTGTC TAGCCCCGTGACCAAGTCTTTCAACCGGGGCGAGTGC	203
CD3 VH fused to constant lambda chain; pETR13860	GAAGTGCAGCTGCTGGAATCCGGCGGAGGACTGGTGCAGCCTGGCG GATCTCTGAGACTGTCTTGTGCCGCTCCGGCTTACCTTCTCCAC CTACGCCATGAACTGGGTGCGACAGGCTCCTGGCAAGGGCCTGGAA TGGGTGTCCCGGATCAGATCCAAGTACAACAACCTACGCCACCTACT ACGCCGACTCCGTGAAGGGCCGGTTCACCATCTCTCGGGACGACTC CAAGAACACCCTGTACCTGCAGATGAACTCCCTGCGGGCCGAGGAC ACCGCCGTGTACTATTGTGTGCGGCACGGCAACTTCGGCAACTCCT ATGTGTCTTGGTTTGCCTACTGGGGCCAGGGCACCCCTCGTGACCGT GTCATCTGCTAGCCCCAAGGCTGCCCCAGCGTGACCCTGTTTCCC CCCAGCAGCGAGGAACTGCAGGCCAACAAGGCCACCCTGGTCTGCC TGATCAGCGACTTCTACCCAGGCGCCGTGACCGTGGCCTGGAAGGC CGACAGCAGCCCCGTGAAGGCCGGCGTGGAGACCACCACCCCCAGC AAGCAGAGCAACAACAAGTACGCCGCCAGCAGCTACCTGAGCCTGA CCCCGAGCAGTGGAAGAGCCACAGGTCTTACAGCTGCCAGGTGAC CCACGAGGGCAGCACC GTGGAGAAAACCGTGGCCCCCACCAGTGCAGC	204
VHCH1[36 F2]_VHCL[CD3]_Fc knob_PGL	CAGGTGCAATTGGTTCAATCTGGTGTCTGAAGTAAAAAACCGGGCGCTTCC GTTAAAGTGAGCTGCAAAGCATCCGGATACACCTTCACTTCTATTACATG CACTGGGTTTCGTCAAGCCCCGGGCCAGGGTCTGGAATGGATGGGCATCATT AACCCAAGCGGTGGCTCTACCTCTACGCGCAGAAATTCCAGGGTCGCGTC ACGATGACCCATGACACTAGCACCTTACCGTTTTATATGGAGCTGTCCAGC CTGCGTTCTGAAGATACTGCAGTGTACTACTGTGCACGCTCTTTCTTCACT GGTTTCCATCTGGACTATTGGGGTCAAGGCACCCTCGTAACGGTTTCTTCT	246

<p>ALA pCON1056</p>	<p>GCTAGCACAAAGGGCCCCAGCGTGTTCCCTCTGGCCCCCTAGCAGCAAGAGC ACATCTGGCGGAACAGCCGCCCTGGGCTGCCTCGTGAAGGACTACTTTCCC GAGCCTGTGACCGTGTCTTGGAACTCTGGCGCCCTGACAAGCGGCGTGAC ACCTTTCCAGCCGTGCTGCAGAGCAGCGCCCTGTACTCTCTGAGCAGCGTG GTCACCGTGCCTAGCAGCAGCCTGGGCACCCAGACCTACATCTGCAACGTG AACCACAAGCCCAGCAACACCAAGTGGACAAGAAGGTGGAGCCCCAAG AGCTGTGATGGCGGAGGAGGGTCCGGAGGCGGAGGATCCGAGGTGCAGCTG CTGGAATCTGGCGGCGGACTGGTGCAGCCTGGCGGATCTCTGAGACTGAGC TGTGCCGCCAGCGGCTTCACCTTCAGCACCTACGCCATGAACTGGGTGCGC CAGGCCCCCTGGCAAAGGCCTGGAATGGGTGTCCCGGATCAGAAGCAAGTAC AACAACCTACGCCACCTACTACGCCGACAGCGTGAAGGGCCGGTTACCATC AGCCGGGACGACAGCAAGAACACCCTGTACCTGCAGATGAACAGCCTGCGG GCCGAGGACACCGCCGTGTACTATTGTGTGCGGCACGGCAACTTCGGCAAC AGCTATGTGTCTTGGTTTGCCTACTGGGGCCAGGGCACCCCTCGTGACCGTG TCAAGCGCTAGTGTGGCCGCTCCCTCCGTGTTTATCTTTCCCCCATCCGAT GAACAGCTGAAAAGCGGCACCGCCTCCGTGCTGTGTCTGCTGAACAATTTT TACCCTAGGGAAGCTAAAGTGCAGTGGAAAGTGGATAACGCACTGCAGTCC GGCAACTCCCAGGAATCTGTGACAGAACAGGACTCCAAGGACAGCACCTAC TCCCTGTCTCTCCACCCTGACACTGTCTAAGGCTGATTATGAGAAACAC AAAGTCTACGCCTGCGAAGTCACCCATCAGGGCCTGAGCTCGCCCGTCACA AAGAGCTTCAACAGGGGAGAGTGTGACAAGACCCACACCTGTCCCCCTTGT CCTGCCCCCTGAAGCTGTGCGCGCCCTTCTGTGTTCTGTCTCCCCCAAG CCCAAGGACACCCTGATGATCAGCCGGACCCCCGAAGTGACCTGCGTGGTG GTGGATGTGTCCACGAGGACCCCTGAAGTGAAGTTCAATTGGTACGTGGAC GGCGTGGAAGTGCAACGCCAAGACAAAGCCGCGGGAGGAGCAGTACAAC AGCACGTACCGTGTGGTCAGCGTCTCACCCTCCTGCACCAGGACTGGCTG AATGGCAAGGAGTACAAGTGCAAGGTCTCCAACAAAGCCCTCGGCGCCCCC ATCGAGAAAACCATCTCCAAAGCCAAAGGGCAGCCCCGAGAACCACAGGTG TACACCCTGCCCCCATGCCGGGATGAGCTGACCAAGAACCAGGTACGCCTG TGGTGCCTGGTCAAAGGCTTCTATCCAGCGACATCGCCGTGGAGTGGGAG AGCAATGGGCAGCCGGAGAACAACATAAGACCACGCTCCCGTGTGTGGAC TCCGACGGCTCCTTCTTCTCTACAGCAAGCTCACCGTGGACAAGAGC AGGTGGCAGCAGGGGAACGTCTTCTCATGCTCCGTGATGCATGAGGCTCTG CACAACCACTACACGCAGAAGAGCCTCTCCCTGTCTCCGGGTAAA</p>	
<p>36F2-Fc hole PGLALA pCON1050</p>	<p>CAGGTGCAATTGGTTCAATCTGGTGCTGAAGTAAAAAAACCGGGCGCTTCC GTTAAAGTGAGCTGCAAAGCATCCGGATACACCTTCACTTCTATTACATG CACTGGGTTTCGTCAAGCCCCGGGCCAGGGTCTGGAATGGATGGGCATCATT AACCCAAGCGGTGGCTCTACCTCTACGCGCAGAAATTCAGGGTCGCGTC ACGATGACCCATGACACTAGCACCTCTACCGTTTATATGGAGCTGTCCAGC CTGCGTTCTGAAGATACTGCAGTGTACTACTGTGCACGCTCTTTCTTCACT GGTTTCCATCTGGACTATTGGGGTCAAGGCACCCCTCGTAACGGTTTCTTCT GCTAGCACCAAGGGCCCCCTCCGTGTTCCCCCTGGCCCCCAGCAGCAAGAGC ACCAGCGGCGGCACAGCCGCTCTGGGCTGCCTGGTCAAGGACTACTTCCCC GAGCCCGTGACCGTGTCTTGAACAGCGGAGCCCTGACCTCCGGCGTGCAC ACCTTCCCCGCGGTGCTGCAGAGTTCTGGCCTGTATAGCCTGAGCAGCGTG GTCACCGTGCCTTCTAGCAGCCTGGGCACCCAGACCTACATCTGCAACGTG AACCACAAGCCCAGCAACACCAAGGTGGACAAGAAGGTGGAGCCCCAAG AGCTGCGACAAAACCTACACATGCCCACCGTGCCCAGCACCTGAAGCTGCA GGGGGACCGTCAGTCTTCTCTTCCCCCAAAACCAAGGACACCCCTCATG ATCTCCCGGACCCCTGAGGTACATGCGTGGTGGTGGACGTGAGCCACGAA GACCCTGAGGTCAAAGTTCAACTGGTACGTGGACGGCGTGGAGGTGCATAAT GCCAAGACAAAGCCGCGGGAGGAGCAGTACAACAGCACGTACCGTGTGGTC AGCGTCTCACCCTCCTGCACCAGGACTGGCTGAATGGCAAGGAGTACAAG TGCAAGGTCTCCAACAAAGCCCTCGGCGCCCCCATCGAGAAAACCATCTCC AAAGCCAAAGGGCAGCCCCGAGAACCACAGGTGTGCACCCTGCCCCCATCC CGGGATGAGCTGACCAAGAACCAGGTGAGCCTCTCGTGCGCAGTCAAAGGC TTCTATCCCAGCAGATCGCCGTGGAGTGGGAGAGCAATGGGCAGCCGGAG AACAACATAAGACCACGCTCCCGTGTGACTCCGACGGCTCCTTCTTCTC CTCGTGAGCAAGCTCACCGTGGACAAGAGCAGGTGGCAGCAGGGGAACGTC</p>	<p>247</p>

-258-

	TTCTCATGCTCCGTGATGCATGAGGCTCTGCACAACCACTACACGCAG AAGAGCCTCTCCCTGTCTCCGGGTAAG	
36F2 LC pCON1062	GAAATCGTGTTAACGCAGTCTCCAGGCACCCCTGTCTTTGTCTCCAGGGGAA AGAGCCACCCCTCTCTTGCAGGGCCAGTCAGAGTGTTAGCAGCAGCTACTTA GCCTGGTACCAGCAGAAACCTGGCCAGGCTCCCAGGCTCCTCATCTATGGA GCATCCAGCAGGGCCACTGGCATCCCAGACAGGTTTCAGTGGCAGTGGATCC GGGACAGACTTCACTCTCACCATCAGCAGACTGGAGCCTGAAGATTTTGCA GTGTATTACTGTGACAGTATACCAACGAACATTATTATACGTTCCGGCCAG GGGACCAAAGTGGAATCAAACGTACGGTGGCTGCACCATCTGTCTTCATC TTCCCGCCATCTGATGAGCAGTTGAAATCTGGAAGTGCCTCTGTTGTGTGC CTGCTGAATAACTTCTATCCCAGAGAGGCCAAAGTACAGTGGAAAGGTGGAT AACGCCCTCCAATCGGGTAACCTCCCAGGAGAGTGTACAGAGCAGGACAGC AAGGACAGCACCTACAGCCTCAGCAGCACCCCTGACGCTGAGCAAAGCAGAC TACGAGAAACACAAAGTCTACGCTGCGAAGTCAACCATCANGGCCTGAGC TCGCCCCGTACAAAGAGCTTCAACAGGGGAGAGTGT	97
CD3 VLCH1 pETR12940	CAGGCCGTCGTGACCCAGGAACCCAGCCTGACAGTGTCTCCTGGCGGCACC GTGACCCTGACATGTGGCAGTTCTACAGGCGCCGTGACCACCAGCAACTAC GCCAACTGGGTGCAGGAAAAGCCCCGGCCAGGCCTTCAGAGGACTGATCGGC GGCACCAACAAGAGAGCCCCCTGGCACCCCTGCCAGATTACGCGGATCTCTG CTGGGAGGAAAGGCCGCCCTGACACTGTCTGGCGCCAGCCAGAAGATGAG GCCGAGTACTACTGCGCCCTGTGGTACAGCAACCTGTGGGTGTTTCGGCGGA GGCACCAAGCTGACAGTGTGAGCAGCGCTTCCACCAAAGGCCCTTCCGTG TTTCCTCTGGCTCCTAGCTCCAAGTCCACCTCTGGAGGCACCGCTGCTCTC GGATGCCTCGTGAAGGATTATTTTCCTGAGCCTGTGACAGTGTCTCGGAAT AGCGGAGCACTGACCTCTGGAGTGCATACTTTCCCCGCTGTGCTGCAGTCC TCTGGACTGTACAGCCTGAGCAGCGTGGTGACAGTGCCAGCAGCAGCCTG GGCACCCAGACCTACATCTGCAACGTGAACCACAAGCCCAGCAACACCAAG GTGGACAAGAAGGTGGAACCCAAGTCTTGT	198

Name	Sequence	Seq ID No
K53A nt	CAGACCGTCGTGACCCAGGAACCCAGCCTGACAGTGTCTCCTGGCGGCACC GTGACCCTGACATGTGGCAGTTCTACAGGCGCCGTGACCACCAGCAACTAC GCCAACTGGGTGCAGCAGAAGCCAGGCCAGGCTCCCAGAGGACTGATCGGC GGCACCAACGCCAGAGCCCCCTGGCACCCCTGCCAGATTACGCGGATCTCTG CTGGGAGGAAAGGCCGCCCTGACACTGTCTGGCGTGCAGCCTGAAGATGAG GCCGAGTACTACTGCGCCCTGTGGTACAGCAACCTGTGGGTGTTTCGGCGGA GGCACCAAGCTGACAGTCCTA	205
S93A nt	CAGACCGTCGTGACCCAGGAACCCAGCCTGACAGTGTCTCCTGGCGGCACC GTGACCCTGACATGTGGCAGTTCTACAGGCGCCGTGACCACCAGCAACTAC GCCAACTGGGTGCAGCAGAAGCCAGGCCAGGCTCCCAGAGGACTGATCGGC GGCACCAACAAGAGAGCCCCCTGGCACCCCTGCCAGATTACGCGGATCTCTG CTGGGAGGAAAGGCCGCCCTGACACTGTCTGGCGTGCAGCCTGAAGATGAG GCCGAGTACTACTGCGCCCTGTGGTACGCCAACCTGTGGGTGTTTCGGCGGA GGCACCAAGCTGACAGTCCTA	206

Name	Sequence	Seq ID No
S35H nt	GAGGTGCAATTGGTGGAAAGCGGAGGCGGCCTCGTGAAGCCTGGCGGATCT CTGAGACTGAGCTGTGCCGCCAGCGGCTTACCTTCAGCAACGCCTGGATG CACTGGGTGCGCCAGGCCCCCTGGAAAAGGACTCGAGTGGGTGGGACGGATC AAGAGCAAGACCGATGGCGGCACCACCGACTATGCCGCCCTGTGAAGGGC CGGTTACCATCAGCAGGGACGACAGCAAGAACACCCTGTACCTGCAGATG AACAGCCTGAAAACCGAGGACACCGCCGTGTACTACTGCACCACCCCTGG GAGTGGTCTTGGTACGACTATTGGGGCCAGGGCACCCCTCGTGACCCTGTCC TCTGCTAGC	207
G49S	GAGGTGCAATTGGTGGAAAGCGGAGGCGGCCTCGTGAAGCCTGGCGGATCT CTGAGACTGAGCTGTGCCGCCAGCGGCTTACCTTCAGCAACGCCTGGATG	208

-259-

nt	AGCTGGGTGCGCCAGGCCCTGGAAAAGGACTCGAGTGGGTGTCCCGGATC AAGAGCAAGACCGATGGCGGCACCACCGACTATGCCGCCCTGTGAAGGGC CGGTTACCATCAGCAGGGACGACAGCAAGAACACCCTGTACCTGCAGATG AACAGCCTGAAAACCGAGGACACCGCCGTGTACTACTGCACCACCCCTGG GAGTGGTCTTGGTACGACTATTGGGGCCAGGGCACCCCTCGTGACCGTGTCC TCTGCTAGC	
R50S nt	GAGGTGCAATTGGTGGAAAGCGGAGGCGGCCTCGTGAAAGCCTGGCGGATCT CTGAGACTGAGCTGTGCCGCCAGCGGCTTACCTTCAGCAACGCCTGGATG AGCTGGGTGCGCCAGGCCCTGGAAAAGGACTCGAGTGGGTGGGATCTATC AAGAGCAAGACCGACGCGGCACCACCGACTATGCCGCCCTGTGAAGGGC CGGTTACCATCAGCAGGGACGACAGCAAGAACACCCTGTACCTGCAGATG AACAGCCTGAAAACCGAGGACACCGCCGTGTACTACTGCACCACCCCTGG GAGTGGTCTTGGTACGACTATTGGGGCCAGGGCACCCCTCGTGACCGTGTCC TCT GCTAGC	209
W96Y nt	GAGGTGCAATTGGTGGAAAGCGGAGGCGGCCTCGTGAAAGCCTGGCGGATCT CTGAGACTGAGCTGTGCCGCCAGCGGCTTACCTTCAGCAACGCCTGGATG AGCTGGGTGCGCCAGGCCCTGGAAAAGGACTCGAGTGGGTGGGACGGATC AAGAGCAAGACCGATGGCGGCACCACCGACTATGCCGCCCTGTGAAGGGC CGGTTACCATCAGCAGGGACGACAGCAAGAACACCCTGTACCTGCAGATG AACAGCCTGAAAACCGAGGACACCGCCGTGTACTACTGCACCACCCCTAC GAGTGGTCTTGGTACGACTACTGGGGCCAGGGCACCCCTCGTGACCGTGTCA TCT GCTAGC	210
W98Y nt	GAGGTGCAATTGGTGGAAAGCGGAGGCGGCCTCGTGAAAGCCTGGCGGATCT CTGAGACTGAGCTGTGCCGCCAGCGGCTTACCTTCAGCAACGCCTGGATG AGCTGGGTGCGCCAGGCCCTGGAAAAGGACTCGAGTGGGTGGGACGGATC AAGAGCAAGACCGATGGCGGCACCACCGACTATGCCGCCCTGTGAAGGGC CGGTTACCATCAGCAGGGACGACAGCAAGAACACCCTGTACCTGCAGATG AACAGCCTGAAAACCGAGGACACCGCCGTGTACTACTGCACCACCCCTGG GAGTACTCTTGGTACGACTACTGGGGCCAGGGCACCCCTCGTGACCGTGTCA TCT GCTAGC	211

Name	Sequence	Seq ID No
90D7 nt	CAGGTGCAATTGGTTCAATCTGGTGCTGAAGTAAAAAAACCGGGCGCTTCC GTTAAAGTGAGCTGCAAAGCATCCGGATACACCTTCACTTCTATTACATG CACTGGGTTCGTCAAGCCCCGGGCCAGGGTCTGGAATGGATGGGCATCATT AACCCAAGCGGTGGCTCTACCTCCTACGCGCAGAAATTCAGGGTCGCGTC ACGATGACCCGTGACACTAGCACCTCTACCGTTTATATGGAGCTGTCCAGC CTGCGTTCTGAAGATACTGCAGTGTACTACTGTGCACGCAACTACTATC GTTGTTTCTCCGTTTCGACTATTGGGGTCAAGGCACCCCTCGTAACGGTTTCT TCTGCTAGC	212
90C1 nt	CAGGTGCAATTGGTTCAATCTGGTGCTGAAGTAAAAAAACCGGGCGCTTCC GTTAAAGTGAGCTGCAAAGCATCCGGATACACCTTCACTTCTATTACATG CACTGGGTTCGTCAAGCCCCGGGCCAGGGTCTGGAATGGATGGGCATCATT AACCCAAGCGGTGGCTCTACCTCCTACGCGCAGAAATTCAGGGTCGCGTC ACGATGACCCGTGACACTAGCACCTCTACCGTTTATATGGAGCTGTCCAGC CTGCGTTCTGAAGATACTGCAGTGTACTACTGTGCACGCAACTACTTCATC GGTTCTGTTGCTATGGACTATTGGGGTCAAGGCACCCCTCGTAACGGTTTCT TCTGCTAGC	213
5E8 VH nt	CAGGTGCAATTGGTTCAATCTGGTGCTGAAGTAAAAAAACCGGGCGCTTCC GTTAAAGTGAGCTGCAAAGCATCCGGATACACCTTCACTTCTATTACATG CACTGGGTTCGTCAAGCCCCGGGCCAGGGTCTGGAATGGATGGGCATCATT AACCCAAGCGGTGGCTCTACCTCCTACGCGCAGAAATTCAGGGTCGCGTC ACGATGACCCGTGACACTAGCACCTCTACCGTTTATATGGAGCTGTCCAGC CTGCGTTCTGAAGATACTGCAGTGTACTACTGTGCACGCGGTCGACTTAC TCTATGGACTATTGGGGTCAAGGCACCCCTCGTAACGGTTTCTTCTGCTAGC	214

5E8 VL nt	GATATTGTTATGACTCAATCTCCACTGTCTCTGCCGGTGACTCCAGGCGAA CCGGCGAGCATTTCTTGCCGTTCCAGCCAGTCTCTGCTGCACTCCAACGGC TACAACATATCTCGATTGGTACCTGCAAAAACCGGGTCAGAGCCCTCAGCTG CTGATCTACCTGGGCTCTAACCGCGCTTCCGGTGTACCGGACCGTTTCAGC GGCTCTGGATCCGGCACCATTTCACGTTGAAAAATCAGCCGTGTTGAAGCA GAAGACGTGGGCGTTTATTACTGTATGCAGGCACTGCAGATTCCAAACACT TTTGGTCAAGGCACCAAGGTCGAAATTAAACGTACG	215
12A4 VH nt	GAGGTGCAATTGTTGGAGTCTGGGGGAGGCTTGGTACAGCCTGGGGGGTCC CTGAGACTCTCCTGTGCAGCCTCCGGATTACCTTTAGCAGTTATGCCATG AGCTGGGTCCGCCAGGCTCCAGGGAAGGGGCTGGAGTGGGTCTCAGCTATT AGTGGTAGTGGTGGTAGCACATACTACGCAGACTCCGTGAAGGGCCGGTTC ACCATCTCCAGAGACAATTCCAAGAACACGCTGTATCTGCAGATGAACAGC CTGAGAGCCGAGGACACGGCCGTATATTACTGTGCGAAATACGCTTACGCT CTGGACTACTGGGGCCAAGGAACCCCTGGTCACCGTCTCGAGTGTCTAGC	216
12A4 VL nt	GAAATCGTGTTAACGCAGTCTCCAGGCACCCTGTCTTTGTCTCCAGGGGAA AGAGCCACCCTCTCTTGCAGGGCCAGTCAGAGTGTTAGCAGCAGCTACTTA GCCTGGTACCAGCAGAAACCTGGCCAGGCTCCCAGGCTCCTCATCTATGGA GCATCCAGCAGGGCCACTGGCATCCCAGACAGGTTTCACTGGCAGTGGAATCC GGGACAGACTTCACTCTCACCATCAGCAGACTGGAGCCTGAAGATTTTGCA GTGTATTACTGTGCAGCAGATGGCAGCAGCAGCACGTTTCGGCCAGGGGACC AAAGTGGAATCAAACGTACG	217
7A3 VH nt	CAGGTGCAATTGGTTCAATCTGGTGCTGAAGTAAAAAAACCGGGCGCTTCC GTTAAAGTGAGCTGCAAAGCATCCGGATACACCTTCACTTCTTATTACATG CACTGGGTTTCGTCAAGCCCCGGGCCAGGGTCTGGAATGGATGGGCATCATT AACCCAAGCGGTGGCTCTACCTCCTACGCGCAGAAATTCAGGGTTCGCGTC ACGATGACCCGTGACACTAGCACCTCTACCGTTTATATGGAGCTGTCCAGC CTGCGTTCTGAAGATACTGCAGTGTACTACTGTGCACGCGGTGACTTCTCT GCTGGTCTGTGATGGACTATTGGGGTCAAGGCACCCTCGTAACGGTTTCT TCTGCTAGC	218
7A3 VL nt	GATATTGTTATGACTCAATCTCCACTGTCTCTGCCGGTGACTCCAGGCGAA CCGGCGAGCATTTCTTGCCGTTCCAGCCAGTCTCTGCTGCACTCCAACGGC TACAACATATCTCGATTGGTACCTGCAAAAACCGGGTCAGAGCCCTCAGCTG CTGATCTACCTGGGCTCTAACCGCGCTTCCGGTGTACCGGACCGTTTCAGC GGCTCTGGATCCGGCACCATTTCACGTTGAAAAATCAGCCGTGTTGAAGCA GAAGACGTGGGCGTTTATTACTGTATGCAGGCACTGCAGACCCCACTTA ACCTTTGGTCAAGGCACCAAGGTCGAAATTAAACGTACG	219
6E10 VH nt	CAGGTGCAATTGGTTCAATCTGGTGCTGAAGTAAAAAAACCGGGCGCTTCC GTTAAAGTGAGCTGCAAAGCATCCGGATACACCTTCACTTCTTATTACATG CACTGGGTTTCGTCAAGCCCCGGGCCAGGGTCTGGAATGGATGGGCATCATT AACCCAAGCGGTGGCTCTACCTCCTACGCGCAGAAATTCAGGGTTCGCGTC ACGATGACCCGTGACACTAGCACCTCTACCGTTTATATGGAGCTGTCCAGC CTGCGTTCTGAAGATACTGCAGTGTACTACTGTGCACGCGGTGACTACAAC GCTTTCGACTATTGGGGTCAAGGCACCCTCGTAACGGTTTCTTCTGCTAGC	220
6E10 VL nt	GATATTGTTATGACTCAATCTCCACTGTCTCTGCCGGTGACTCCAGGCGAA CCGGCGAGCATTTCTTGCCGTTCCAGCCAGTCTCTGCTGCACTCCAACGGC TACAACATATCTCGATTGGTACCTGCAAAAACCGGGTCAGAGCCCTCAGCTG CTGATCTACCTGGGCTCTAACCGCGCTTCCGGTGTACCGGACCGTTTCAGC GGCTCTGGATCCGGCACCATTTCACGTTGAAAAATCAGCCGTGTTGAAGCA GAAGACGTGGGCGTTTATTACTGTATGCAGGCATGGCATAGCCCACTTTT GGTCAAGGCACCAAGGTCGAAATTAAACGTACG	221
12F9 VH nt	CAGGTGCAATTGGTTCAATCTGGTGCTGAAGTAAAAAAACCGGGCGCTTCC GTTAAAGTGAGCTGCAAAGCATCCGGATACACCTTCACTTCTTATTACATG CACTGGGTTTCGTCAAGCCCCGGGCCAGGGTCTGGAATGGATGGGCATCATT AACCCAAGCGGTGGCTCTACCTCCTACGCGCAGAAATTCAGGGTTCGCGTC ACGATGACCCGTGACACTAGCACCTCTACCGTTTATATGGAGCTGTCCAGC CTGCGTTCTGAAGATACTGCAGTGTACTACTGTGCACGCGGTGCTACTTAC ACTATGGACTATTGGGGTCAAGGCACCCTCGTAACGGTTTCTTCTGCTAGC	222
12F9 VL	GATATTGTTATGACTCAATCTCCACTGTCTCTGCCGGTGACTCCAGGCGAA CCGGCGAGCATTTCTTGCCGTTCCAGCCAGTCTCTGCTGCACTCCAACGGC	223

-261-

nt	TACAACTATCTCGATTGGTACCTGCAAAAACCGGGTCAGAGCCCTCAGCTG CTGATCTACCTGGGCTCTAACCGCGCTTCCGGTGTACCGGACCGTTTCAGC GGCTCTGGATCCGGCACCGATTTCACGTTGAAAATCAGCCGTGTTGAAGCA GAAGACGTGGGCGTTTATTACTGTATGCAGGCACTGCAGACCCCAATTACT TTTGGTCAAGGCACCAAGGTGCAAATTAAACGTACG	
----	--	--

Name	Sequence	Seq ID No
pETR11646 Mov19 VH- CH1-Fchole PG/LALA	CAGGTGCAGCTGCAGCAGTCTGGCGCCGAGCTCGTGAAAACCTGGCGCCTCC GTGAAGATCAGCTGCAAGGCCAGCGGCTACAGCTTACCGGCTACTTCATG AACTGGGTCAAGCAGAGCCACGGCAAGAGCCTGGAATGGATCGGCAGAATC CACCCCTACGACGGCGACACCTTCTACAACCAGAACTTCAAGGACAAGGCC ACCCTGACCGTGGACAAGAGCAGCAACACCGCCACATGGAAGTGTGAGC CTGACCAGCGAGGACTTCGCCGTGTACTACTGCACCAGATACGACGGCAGC CGGGCCATGGATTATTGGGGCCAGGGCACCACCGTGACAGTGTCCAGCGCT AGCACCAAGGGCCCCCTCCGTGTTCCCCCTGGCCCCCAGCAGCAAGAGCACC AGCGGGCGGCACAGCCGCTCTGGGCTGCCTGGTCAAGGACTACTTCCCCGAG CCCGTGACCGTGTCTTGAACAGCGGAGCCCTGACCTCCGGCGTGCACACC TTCCCCGCCGTGCTGCAGAGTTCTGGCCTGTATAGCCTGAGCAGCGTGGTC ACCGTGCCTTCTAGCAGCCTGGGCACCCAGACCTACATCTGCAACGTGAAC CACAAGCCCAGCAACACCAAGGTGGACAAGAAGGTGGAGCCCAAGAGCTGC GACAAAACCTACACATGCCCCACCGTGCCACAGCACTGAAGCTGCAGGGGGA CCGTGAGTCTTCTCTTCCCCCAAAACCAAGGACACCCCTCATGATCTCC CGGACCCCTGAGGTACATGCGTGGTGGTGGACGTGAGCCACGAAAGCCCT GAGGTCAAGTTCAACTGGTACGTGGACGGCGTGGAGGTGCATAATGCCAAG ACAAAGCCGCGGGAGGAGCAGTACAACAGCACGTACCGTGTGGTCAAGCGTC CTCACCGTCTTGACACAGGACTGGCTGAATGGCAAGGAGTACAAGTGAAG GTCTCCAACAAAGCCCTCGGCGCCCCATCGAGAAAACCATCTCCAAAGCC AAAGGGCAGCCCCGAGAACCACAGGTGTGCACCCTGCCCCCATCCCGGGAT GAGCTGACCAAGAACCAGGTGAGCCTCTCGTGCGCAGTCAAAGGCTTCTAT CCCAGCGACATCGCCGTGGAGTGGGAGAGCAATGGGCAGCCGGAGAACAAC TACAAGACCACGCTCCCGTGTGACTCCGACGGCTCCTTCTTCCCTCGTG AGCAAGCTCACCGTGGACAAGAGCAGGTGGCAGCAGGGGAACGTCTTCTCA TGCTCCGTGATGCATGAGGCTCTGCACAACCACTACACGCAGAAGAGCCCTC TCCCTGTCTCCGGGTAAA	224
pETR11647 Mov19 VH- CH1-CD3 VH-CL- Fcknob PG/LALA	CAGGTGCAGCTGCAGCAGTCTGGCGCCGAGCTCGTGAAAACCTGGCGCCTCC GTGAAGATCAGCTGCAAGGCCAGCGGCTACAGCTTACCGGCTACTTCATG AACTGGGTCAAGCAGAGCCACGGCAAGAGCCTGGAATGGATCGGCAGAATC CACCCCTACGACGGCGACACCTTCTACAACCAGAACTTCAAGGACAAGGCC ACCCTGACCGTGGACAAGAGCAGCAACACCGCCACATGGAAGTGTGAGC CTGACCAGCGAGGACTTCGCCGTGTACTACTGCACCAGATACGACGGCAGC CGGGCCATGGATTATTGGGGCCAGGGCACCACCGTGACAGTGTCCAGCGCT AGCACAAAGGGCCCCAGCGTGTTCCTCTGGCCCCCTAGCAGCAAGAGCACA TCTGGCGGAACAGCCGCCCTGGGCTGCCTCGTGAAGGACTACTTCCCCGAG CCTGTGACCGTGTCTTGAACCTCTGGCGCCCTGACAAGCGGCGTGCACACC TTTCCAGCCGTGCTGCAGAGCAGCGGCTGTACTCTCTGAGCAGCGTGGTC ACCGTGCCTAGCAGCAGCCTGGGCACCCAGACCTACATCTGCAACGTGAAC CACAAGCCCAGCAACACCAAGTGGACAAGAAGGTGGAGCCCAAGAGCTGT GATGGCGGAGGAGGGTCCGGAGGCGGAGGATCCGAAGTGCAGCTGGTGGAA AGCGGCGGAGGCCTGGTGCAGCCTAAGGGCTCTCTGAAGCTGAGCTGTGCC GCCAGCGGCTTACCTTCAACACCTACGCCATGAACTGGGTGCGCCAGGCC CCTGGCAAAGGCCTGGAATGGGTGGCCCCGATCAGAAGCAAGTACAACAAT TACGCCACCTACTACGCCGACAGCGTGAAGGACCGGTTACCATCAGCCGG GACGACAGCCAGAGCATCCTGTACCTGCAGATGAACAACCTGAAAACCGAG GACACCGCCATGTACTACTGCGTGCAGGACGGCAACTTCGGCAACAGCTAT GTGTCTTGGTTTGCCTACTGGGGCCAGGGCACCCCTCGTGACAGTGTCTGCT GCTAGCGTGGCCGCTCCCTCCGTGTTTATCTTTCCCCCATCCGATGAACAG CTGAAAAGCGGCACCGCCTCCGTGCTGTGTCTGCTGAACAATTTTTACCCCT AGGGAAGCTAAAGTGCAGTGGAAAGTGGATAACGCACCTGCAGTCCGGCAAC	225

-262-

	TCCCAGGAATCTGTGACAGAACAGGACTCCAAGGACAGCACCTACTCCCTG TCCTCCACCCTGACACTGTCTAAGGCTGATTATGAGAAACACAAAGTCTAC GCCTGCGAAGTCACCCATCAGGGCCTGAGCTCGCCCGTCACAAAGAGCTTC AACAGGGGAGAGTGTGACAAGACCCACACCTGTCCCCCTTGTCTGCCCCCT GAAGCTGCTGGCGGCCCTTCTGTGTTCTGTTCCTGTTCCTCCCAAGGCCAAGGAC ACCCTGATGATCAGCCGACCCCCGAAGTGACCTGCGTGTTGGTGGATGTG TCCCACGAGGACCCTGAAGTGAAAGTTCAATTGGTACGTGGACGGCGTGGAA GTGCACAACGCCAAGACAAAGCCGCGGGAGGAGCAGTACAACAGCACGTAC CGTGTGGTCAGCGTCTCACCCTCCTGCACCAGGACTGGCTGAATGGCAAG GAGTACAAGTGCAAGGTCTCCAACAAAGCCCTCGGCGCCCCCATCGAGAAA ACCATCTCCAAAGCCAAAGGGCAGCCCCGAGAACCACAGGTGTACACCCTG CCCCCATGCCGGGATGAGCTGACCAAGAACCAGGTGAGCCTGTGGTGCCTG GTCAAAGGCTTCTATCCCAGCGACATCGCCGTGGAGTGGGAGAGCAATGGG CAGCCGGAGAACAACACTACAAGACCACGCCTCCCGTGCTGGACTCCGACGGC TCCTTCTTCTCTACAGCAAGCTCACCCTGGACAAGAGCAGGTGGCAGCAG GGGAACGTCTTCTCATGTCCGTGATGCATGAGGCTCTGCACAACCACTAC ACGCAGAAGAGCCTCTCCCTGTCTCCGGGTAAA	
pETR11644 Mov19 LC	GACATCGAGCTGACCCAGAGCCCTGCCTCTCTGGCCGTGTCTCTGGGACAG AGAGCCATCATCAGCTGCAAGGCCAGCCAGAGCGTGTCTTTGCCGGCACC TCTCTGATGCACTGGTATCACCAGAAGCCCGGCCAGCAGCCCAAGCTGCTG ATCTACAGAGCCAGCAACCTGGAAGCCGGCGTGCCCAAGATTTTCCGGC AGCGGCAGCAAGACCGACTTCACCTGAACATCCACCCGTGGAAGAAGAG GACGCCGCCACCTACTACTGCCAGCAGAGCAGAGAGTACCCCTACACCTTC GGCGGAGGCACCAAGCTGGAATCAAGCGTACGGTGGCTGCACCATCTGTC TTCATCTTCCCGCCATCTGATGAGCAGTTGAAATCTGGAAGTGCCTCTGTT GTGTGCCTGCTGAATAACTTCTATCCCAGAGAGGCCAAAGTACAGTGGAAAG GTGGATAACGCCCTCCAATCGGGTAACTCCCAGGAGAGTGTACAGAGCAG GACAGCAAGGACAGCACCTACAGCCTCAGCAGCACCCCTGACGCTGAGCAAA GCAGACTACGAGAAACACAAAGTCTACGCCTGCGAAGTACCCATCAGGGC CTGAGCTCGCCCGTCACAAAGAGCTTCAACAGGGGAGAGTGT	226

Variant	Sequence	Seq ID No
16D5 VH_D52d E	GAGGTGCAATTGGTTGAATCTGGTGGTGGTCTGGTAAACCGGGCGGTTCCC TGCGTCTGAGCTGCGCGGCTTCCGGATTACCTTCTCCAACGCGTGGATGAG CTGGGTTTCGCCAGGCCCGGGCAAAGGCCTCGAGTGGGTTGGTCTGATCAAG TCTAAAACTGAGGGTGGCACCACGGATTACGCGGCTCCAGTTAAAGGTCGTT TTACCATTTCGCCGACGATAGCAAAAACACTCTGTATCTGCAGATGAATC TCTGAAAACCTGAAGACACCGCAGTCTACTACTGTACTACCCCGTGGGAATGG TCTTGGTACGATTATTGGGGCCAGGGCAGCTGGTTACGGTGTCTTCC	261
16D5 VH_D52d Q	GAGGTGCAATTGGTTGAATCTGGTGGTGGTCTGGTAAACCGGGCGGTTCCC TGCGTCTGAGCTGCGCGGCTTCCGGATTACCTTCTCCAACGCGTGGATGAG CTGGGTTTCGCCAGGCCCGGGCAAAGGCCTCGAGTGGGTTGGTCTGATCAAG TCTAAAACTCAGGGTGGCACCACGGATTACGCGGCTCCAGTTAAAGGTCGTT TTACCATTTCGCCGACGATAGCAAAAACACTCTGTATCTGCAGATGAATC TCTGAAAACCTGAAGACACCGCAGTCTACTACTGTACTACCCCGTGGGAATGG TCTTGGTACGATTATTGGGGCCAGGGCAGCTGGTTACGGTGTCTTCC	262
CD3_VH N100A	GAGGTGCAGCTGCTGGAATCTGGCGGCGGACTGGTGCAGCCTGGCGGATCTC TGAGACTGAGCTGTGCCGCCAGCGGCTTACCTTACAGCACCTACGCCATGAA CTGGGTGCGCCAGGCCCTGGCAAAGGCCTGGAATGGGTGTCCCGGATCAGA AGCAAGTACAACAACCTACGCCACCTACTACGCCGACAGCGTGAAGGGCCGGT TCACCATCAGCCGGGACGACAGCAAGAACACCCTGTACCTGCAGATGAACAG CCTGCGGGCCGAGGACACCGCCGTGTACTATTGTGTGCGGCACGGCAACTTC GGCGCCAGCTATGTGTCTTGGTTTGCCTACTGGGGCCAGGGCACCCCTCGTGA CCGTGTCAAGC	263
CD3_VH S100aA	GAGGTGCAGCTGCTGGAATCTGGCGGCGGACTGGTGCAGCCTGGCGGATCTC TGAGACTGAGCTGTGCCGCCAGCGGCTTACCTTACAGCACCTACGCCATGAA CTGGGTGCGCCAGGCCCTGGCAAAGGCCTGGAATGGGTGTCCCGGATCAGA AGCAAGTACAACAACCTACGCCACCTACTACGCCGACAGCGTGAAGGGCCGGT	264

	TCACCATCAGCCGGGACGACAGCAAGAACACCCTGTACCTGCAGATGAACAG CCTGCGGGCCGAGGACACCGCCGTGTACTATTGTGTGCGGCACGGCAACTTC GGCAACGCCCTATGTGTCTTGGTTTGCCTACTGGGGCCAGGGCACCCCTCGTGA CCGTGTCAAGC	
16D5 [VHCH1] - CD3[VHC H1- N100A]- Fcknob_P GLALA	GAGGTGCAATTGGTTGAATCTGGTGGTGGTCTGGTAAAACCGGGCGGTTCCC TGCGTCTGAGCTGCGCGGCTTCCGGATTACCTTCTCCAACGCGTGGATGAG CTGGGTTTCGCCAGGCCCGGGCAAAGGCCTCGAGTGGGTTGGTTCGTATCAAG TCTAAAACTGACGGTGGCACCACGGATTACGCGGCTCCAGTTAAAGGTCGTT TTACCATTTCGCGACGATAGCAAAAACACTCTGTATCTGCAGATGAACTC TCTGAAAACCTGAAGACACCGCAGTCTACTACTGTACTACCCCGTGGGAATGG TCTTGGTACGATTATTGGGGCCAGGGCACGCTGGTTACGGTGTCTTCCGCTA GCACAAAGGGCCCTAGCGTGTTCCTCTGGCCCCCAGCAGCAAGAGCACAAG CGGCGGAACAGCCGCCCTGGGCTGCCTCGTGAAGGACTACTTCCCCGAGCCC GTGACAGTGTCTTGAACAGCGGAGCCCTGACAAGCGGCGTGCACACTTTCC CTGCCGTGTGCAGAGCAGCGGCCTGTACTCCCTGAGCAGCGTGGTCACCGT GCCTAGCAGCAGCCTGGGCACCCAGACCTACATCTGCAACGTGAACCACAAG CCCAGCAACACCAAAGTGGACAAGAAGGTGGAGCCCAAGAGCTGTGATGGCG GAGGAGGGTCCGGAGGCGGAGGATCCGAGGTGCAGCTGTGGAATCTGGCGG CGGACTGGTGCAGCCTGGCGGATCTCTGAGACTGAGCTGTGCCGCCAGCGGC TTCACCTTCAGCACCTACGCCATGAACTGGGTGCGCCAGGCCCTGGCAAAG GCCTGGAATGGGTGTCCCGGATCAGAAGCAAGTACAACAACCTACGCCACCTA CTACGCCGACAGCGTGAAGGGCCGGTTCACCATCAGCCGGGACGACGACAAG AACACCCTGTACCTGCAGATGAACAGCCTGCGGGCCGAGGACACCGCGTGT ACTATTGTGTGCGGCACGGCAACTTCGGCGCCAGCTATGTGTCTTGGTTTGC CTACTGGGGCCAGGGCACCCCTCGTGACCGTGTCAAGCGCTAGTACCAAGGGC CCCAGCGTGTTCCTCCCTGGCACCCAGCAGCAAGAGCACATCTGGCGGAACAG CCGCTCTGGGCTGTCTGGTGAAAGACTACTTCCCCGAGCCCGTGACCGTGTCT TTGGAACCTCTGGCGCCCTGACCAGCGGCGTGCACACCTTTCCAGCCGTGCTG CAGAGCAGCGGCCTGTACTCCCTGTCTCCGTGGTCAACCGTGCCCTCTAGCT CCCTGGGAACACAGACATATATCTGTAATGTCAATCACAAGCCTTCCAACAC CAAAGTCGATAAGAAAGTCGAGCCCAAGAGCTGCGACAAAACCTCACACATGC CCACCGTGCCGAGCACCTGAAGCTGCAGGGGGACCGTCAGTCTTCCTCTTCC CCCCAAAACCCAAGGACACCCTCATGATCTCCCGGACCCCTGAGGTACACATG CGTGGTGGTGGACGTGAGCCACGAAGACCCTGAGGTCAAGTTCAACTGGTAC GTGGACGGCGTGGAGGTGCATAATGCCAAGACAAAGCCGCGGGAGGAGCAGT ACAACAGCACGTACCGTGTGGTCAAGCTCCTCACCCTCCTGCACCAGGACTG GCTGAATGGCAAGGAGTACAAGTGAAGGTCTCCAACAAAGCCCTCGGCGCC CCCATCGAGAAAACCATCTCCAAAGCCAAAGGGCAGCCCCGAGAACCACAGG TGTACACCCTGCCCCCATGCCGGGATGAGCTGACCAAGAACCAGGATGACCT GTGGTGCCTGGTCAAAGGCTTCTATCCAGCGACATCGCCGTGGAGTGGGAG AGCAATGGGCAGCCGGAGAACAACCTACAAGACCACGCTCCCGTGTGGACT CCGACGGCTCCTTCTTCTCTACAGCAAGCTCACCGTGGACAAGAGCAGGTG GCAGCAGGGGAACGTCTTCTCATGTCCGTGATGCATGAGGCTCTGCACAAC CACTACACGCAGAAGAGCCTCTCCCTGTCTCCGGGTAAA	265
16D5- Fchole- PGLALA	GAGGTGCAATTGGTTGAATCTGGTGGTGGTCTGGTAAAACCGGGCGGTTCCC TGCGTCTGAGCTGCGCGGCTTCCGGATTACCTTCTCCAACGCGTGGATGAG CTGGGTTTCGCCAGGCCCGGGCAAAGGCCTCGAGTGGGTTGGTTCGTATCAAG TCTAAAACTGACGGTGGCACCACGGATTACGCGGCTCCAGTTAAAGGTCGTT TTACCATTTCGCGACGATAGCAAAAACACTCTGTATCTGCAGATGAACTC TCTGAAAACCTGAAGACACCGCAGTCTACTACTGTACTACCCCGTGGGAATGG TCTTGGTACGATTATTGGGGCCAGGGCACGCTGGTTACGGTGTCTTCCGCTA GCACCAAGGGCCCTCCGTGTTCCCCCTGGCCCCCAGCAGCAAGAGCACCAG CGGCGGCACAGCCGTCTGGGCTGCCTGGTCAAGGACTACTTCCCCGAGCCC GTGACCGTGTCTTGAACAGCGGAGCCCTGACCTCCGGCGTGCACACCTTCC CCGCCGTGTGCAGAGTTCTGGCCTGTATAGCCTGAGCAGCGTGGTCACCGT GCCTTCTAGCAGCCTGGGCACCCAGACCTACATCTGCAACGTGAACCACAAG CCCAGCAACACCAAGGTGGACAAGAAGGTGGAGCCCAAGAGCTGCGACAAAA CTCACACATGCCACCGTGCCAGCCTGAAGCTGCAGGGGGACCGTCAGT CTTCTCTTCCCCCAAACCCAAGGACACCCTCATGATCTCCCGGACCCCT	266

	GAGGTCACATGCGTGGTGGTGGACGTGAGCCACGAAGACCCTGAGGTCAAGT TCAACTGGTACGTGGACGGCGTGGAGGTGCATAATGCCAAGACAAAGCCGCG GGAGGAGCAGTACAACAGCACGTACCGTGTGGTCAGCGTCCTCACCCTCCTG CACCAGGACTGGCTGAATGGCAAGGAGTACAAGTGCAAGGTCTCCAACAAAG CCCTCGGCGCCCCATCGAGAAAACCATCTCCAAAGCCAAAGGGCAGCCCCG AGAACCACAGGTGTGCACCCTGCCCCCATCCCGGGATGAGCTGACCAAGAAC CAGGTCAGCCTCTCGTGCAGTCAAAGGCTTCTATCCCAGCGACATCGCCG TGGAGTGGGAGAGCAATGGGCAGCCGGAGAACAACATAAGACCACGCTCC CGTGCTGGACTCCGACGGCTCCTTCTTCTCGTGAGCAAGCTCACCCTGGAC AAGAGCAGGTGGCAGCAGGGGAACGTCTTCTCATGCTCCGTGATGCATGAGG CTCTGCACAACCGCTTCACGCAGAAGAGCCTCTCCCTGTCTCCGGGTAA	
CD3-CLC	CAGGCCGTCTGACCCAGGAACCCAGCCTGACAGTGTCTCTGCGGCACCG TGACCCTGACATGTGGCAGTTCTACAGGCGCCGTGACCACCAGCAACTACGC CAACTGGGTGCAGGAAAAGCCCGGCCAGGCCTTCAGAGGACTGATCGGCGGC ACCAACAAGAGAGCCCCCTGGCACCCCTGCCAGATTTCAGCGGATCTCTGCTGG GAGGAAAGGCCGCCCTGACACTGTCTGGCGCCCAGCCAGAAGATGAGGCCGA GTACTACTGCGCCCTGTGGTACAGCAACCTGTGGGTGTTTCGGCGGAGGCACC AAGCTGACAGTCTTAGGTCAACCCAAGGCTGCCCCAGCGTGACCCTGTTCC CCCCCAGCAGCGAGGAACCTGCAGGCCAACAAGGCCACCCTGGTCTGCCTGAT CAGCGACTTCTACCCAGGCGCCGTGACCGTGGCCTGGAAGGCCGACAGCAGC CCCGTGAAGGCCGGCGTGGAGACCACCACCCCGAGCAAGCAGAGCAACAACA AGTACGCCGCCAGCAGCTACCTGAGCCTGACCCCGAGCAGTGGAAGAGCCA CAGGTCTACAGCTGCCAGGTGACCCACGAGGGCAGCACCGTGAGAAAACC GTGGCCCCCACCAGGTGCAGC	267
16D5 [VHCH1] - CD3[VHC H1- S100aA]- Fcknob_P GLALA	GAGGTGCAATTGGTTGAATCTGGTGGTGGTCTGGTAAACCGGGCGGTTCCT TGCGTCTGAGCTGCGCGGCTTCCGGATTACCTTCTCCAACGCGTGATGAG CTGGGTTCGCCAGGCCCCGGGCAAAGGCCTCGAGTGGGTGGTCTGATCAAG TCTAAACTGACGGTGGCACCACGGATTACGCGGCTCCAGTTAAAGGTGCTT TTACCATTTCCGCGACGATAGCAAAAACACTCTGTATCTGCAGATGAACCTC TCTGAAAACCTGAAGACACCGCAGTCTACTACTGTACTACCCCGTGGGAATGG TCTTGGTACGATTATTGGGGCCAGGGCAGCTGGTTACGGTGTCTTCCGCTA GCACAAAGGGCCCTAGCGTGTTCCTCTGGCCCCCAGCAGCAAGAGCACAAAG CGGCGGAACAGCCGCCCTGGGCTGCCTCGTGAAGGACTACTTCCCCGAGCCC GTGACAGTGTCTTGAACAGCGGAGCCCTGACAAGCGGCGTGACACTTTCC CTGCCGTGCTGCAGAGCAGCGGCCTGTACTCCCTGAGCAGCGTGGTCACCGT GCCTAGCAGCAGCCTGGGCACCCAGACCTACATCTGCAACGTGAACCACAAG CCCAGCAACACCAAAGTGGACAAGAAGGTGGAGCCCAAGAGCTGTGATGGCG GAGGAGGGTCCGGAGGCGGAGGATCCGAGGTGCAGCTGCTGGAATCTGGCGG CGGACTGGTGCAGCCTGGCGGATCTCTGAGACTGAGCTGTGCCGCCAGCGGC TTCACCTTCAGCACCTAGCCATGAACCTGGGTGCGCCAGGCCCTGGCAAAAG GCTTGAATGGGTGTCCCGGATCAGAAGCAAGTACAACAACCTACGCCACTA CTACGCCGACAGCGTGAAGGGCCGGTTACCATCAGCCGGGACGACAGCAAG AACACCCTGTACCTGCAGATGAACAGCCTGCGGGCCGAGGACACCGCCGTGT ACTATTGTGTGCGGCACGGCAACTTCGGCAACGCCTATGTGTCTTGGTTTGC CTACTGGGGCCAGGGCACCTCTGTGACCGTGTCAAGCGCTAGTACCAAGGGC CCCAGCGTGTTCCTTGGCACCCAGCAGCAAGAGCACATCTGGCGGAACAG CCGCTCTGGGCTGTCTGGTGAAAGACTACTTCCCCGAGCCCGTGACCGTGTCT TTGGAACCTCTGGCGCCCTGACCAGCGGCGTGACACCTTTCCAGCCGTGCTG CAGAGCAGCGGCCTGTACTCCCTGTCTCCGTGGTACCGTGCCCTCTAGCT CCCTGGGAACACAGACATATATCTGTAATGTCAATCACAAGCCTTCCAACAC CAAAGTCGATAAGAAAGTCGAGCCCCAAGAGCTGCGACAAAACCTCACACATGC CCACCGTGCCAGCACCTGAAGCTGCAGGGGGACCGTCAGTCTTCTCTTCC CCCCAAAACCCAAGGACACCCTCATGATCTCCCGGACCCCTGAGGTACATG CGTGGTGGTGGACGTGAGCCACGAAGACCCTGAGGTCAAGTTCAACTGGTAC GTGGACGGCGTGGAGGTGCATAATGCCAAGACAAAGCCGCGGGAGGAGCAGT ACAACAGCACGTACCGTGTGGTCAGCGTCTTACCCTCTGACACAGGACTG GCTGAATGGCAAGGAGTACAAGTGCAAGGTCTCCAACAAGCCCTCGGCGCC CCCATCGAGAAAACCATCTCCAAAGCCAAAGGGCAGCCCCGAGAACCACAGG TGTACACCCTGCCCCCATGCCGGGATGAGCTGACCAAGAACCAGGTACGCT	268

	GTGGTGCCTGGTCAAAGGCTTCTATCCCAGCGACATCGCCGTGGAGTGGGAG AGCAATGGGCAGCCGGAGAACAACCTACAAGACCACGCCTCCCGTGCTGGACT CCGACGGCTCCTTCTTCTCTACAGCAAGCTCACC GTGGACAAGAGCAGGTG GCAGCAGGGGAACGTCTTCTCATGCTCCGTGATGCATGAGGCTCTGCACAAC CACTACACGCAGAAGAGCCTCTCCCTGTCTCCGGGTAAA	
9D11 [VHCH1] - CD3[VHC L- N100A]- Fcknob_P GLALA	CAGGTGCAATTGGTTCAATCTGGTGCTGAAGTAAAAAACC GGCGCTTCCG TTAAAGTGAGCTGCAAAGCATCCGGATACACCTTCACTTCTATTACATGCA CTGGGTTCGTCAAGCCCCGGGCCAGGGTCTGGAATGGATGGGCATCATTAAC CCAAGCGGTGGCCCTACCTCTACGCGCAGAAATTCCAGGGTCGCGTCACGA TGACCCGTGACACTAGCACCTCTACCGTTTATATGGAGCTGTCCAGCCTGCG TTCTGAAGATACTGCAGTGTACTACTGTGCACGCGGTGACTTCGCTTGGCTG GACTATTGGGGTCAAGGCACCCTCGTAACGGTTTCTTCTGCTAGCACAAAGG GCCCCAGCGTGTTCCCTCTGGCCCCCTAGCAGCAAGAGCACATCTGGCGGAAC AGCCGCCCTGGGCTGCCTCGTGAAGGACTACTTTCCCGAGCCTGTGACCGTG TCCTGGAACCTCTGGCGCCCTGACAAGCGGCGTGACACCTTTCCAGCCGTGC TGCAGAGCAGCGGCCTGTACTCTCTGAGCAGCGTGGTACCCGTGCCTAGCAG CAGCCTGGGCACCCAGACCTACATCTGCAACGTGAACCACAAGCCCAGCAAC ACCAAAGTGGACAAGAAGGTGGAGCCCCAAGAGCTGTGATGGCGGAGGAGGGT CCGGAGGCGGAGGATCCGAGGTGCAGCTGCTGGAATCTGGCGGCGGACTGGT GCAGCCTGGCGGATCTCTGAGACTGAGCTGTGCCGCCAGCGGCTTCACCTTC AGCACCTACGCCATGAACTGGGTGCGCCAGGCCCCCTGGCAAAGGCCTGGAAT GGGTGTCCCGGATCAGAAGCAAGTACAACAACCTACGCCACCTACTACGCCGA CAGCGTGAAGGGCCGGTTACCATCAGCCGGGACGACAGCAAGAACACCCCTG TACCTGCAGATGAACAGCCTGCGGGCCGAGGACACCGCCGTGTACTATTGTG TGCGGCACGGCAACTTCGGCGCCAGCTATGTGTCTTGGTTTGCCTACTGGGG CCAGGGCACCCCTCGTGACCGTGTCAAGCGCTAGTGTGGCCGCTCCCTCCGTG TTTATCTTTCCCCATCCGATGAACAGCTGAAAAGCGGCACCGCCTCCGTGCG TGTGTCTGCTGAACAATTTTACCCTAGGGAAGCTAAAGTGCAGTGGAAAGT GGATAACGCACTGCAGTCCGGCAACTCCCAGGAATCTGTGACAGAACAGGAC TCCAAGGACAGCACCTACTCCCTGTCTCCACCCTGACACTGTCTAAGGCTG ATTATGAGAAACACAAAGTCTACGCCTGCGAAGTCACCCATCAGGGCCTGAG CTCGCCCGTCACAAAGAGCTTCAACAGGGGAGAGTGTGACAAGACCCACACC TGTCCTCCCTTGTCTGCCCCCTGAAGCTGCTGGCGGCCCTTCTGTGTTCTCTGT TCCCCCAAAGCCCAAGGACACCCTGATGATCAGCCGGACCCCCGAAGTGAC CTGCGTGGTGGTGGATGTGTCCCACGAGGACCCTGAAGTGAAGTTCAATTGG TACGTGGACGGCGTGGAAGTGACAACGCCAAGACAAAGCCGCGGGAGGAGC AGTACAACAGCACGTACCGTGTGGTCAGCGTCCTCACCCTCCTGCACCAGGA CTGGCTGAATGGCAAGGAGTACAAGTGCAAGGTCTCCAACAAAGCCCTCGGC GCCCCATCGAGAAAACCATCTCCAAAGCCAAAGGGCAGCCCCGAGAACCCAC AGGTGTACACCCTGCCCCATGCCGGGATGAGCTGACCAAGAAGCAGGTGAG CCTGTGGTGCCTGGTCAAAGGCTTCTATCCCAGCGACATCGCCGTGGAGTGG GAGAGCAATGGGCAGCCGGAGAACAACCTACAAGACCACGCCTCCCGTGCTGG ACTCCGACGGCTCCTTCTTCTCTACAGCAAGCTCACCCTGGACAAGAGCAG GTGGCAGCAGGGGAACGTCTTCTCATGCTCCGTGATGCATGAGGCTCTGCAC AACCCTACACGCAGAAGAGCCTCTCCCTGTCTCCGGGTAAA	269
9D11- Fchole	CAGGTGCAATTGGTTCAATCTGGTGCTGAAGTAAAAAACC GGCGCTTCCG TTAAAGTGAGCTGCAAAGCATCCGGATACACCTTCACTTCTATTACATGCA CTGGGTTCGTCAAGCCCCGGGCCAGGGTCTGGAATGGATGGGCATCATTAAC CCAAGCGGTGGCCCTACCTCTACGCGCAGAAATTCCAGGGTCGCGTCACGA TGACCCGTGACACTAGCACCTCTACCGTTTATATGGAGCTGTCCAGCCTGCG TTCTGAAGATACTGCAGTGTACTACTGTGCACGCGGTGACTTCGCTTGGCTG GACTATTGGGGTCAAGGCACCCTCGTAACGGTTTCTTCTGCTAGCACCAAGG GCCCCCTCCGTGTTCCCCCTGGCCCCCAGCAGCAAGAGCACACGCGGGCAC AGCCGCTCTGGGCTGCCTGGTCAAGGACTACTTCCCCGAGCCCGTGACCGTG TCCTGGAACAGCGGAGCCCTGACCTCCGGCGTGACACCTTCCCCGCCGTGC TGCAGAGTTCTGGCCTGTATAGCCTGAGCAGCGTGGTACCCGTGCCTTCTAG CAGCCTGGGCACCCAGACCTACATCTGCAACGTGAACCACAAGCCCAGCAAC ACCAAGGTGGACAAGAAGGTGGAGCCCAAGAGCTGCGACAAAACCTCACACAT GCCCACCGTGCCAGCACCTGAAGCTGCAGGGGGACCGTCAGTCTTCTCTT	270

	CCCCC AAAACCCAAGGACACCCTCATGATCTCCCGGACCCCTGAGGTCACA TGCGTGGTGGTGGACGTGAGCCACGAAGACCCTGAGGTCAAGTTCAACTGGT ACGTGGACGGCGTGGAGGTGCATAATGCCAAGACAAAGCCGCGGGAGGAGCA GTACAACAGCACGTACCGTGTGGTCAGCGTCCTCACCCTCCTGCACCAGGAC TGGCTGAATGGCAAGGAGTACAAGTGCAAGGTCTCCAACAAAGCCCTCGGCG CCCCCATCGAGAAAACCATCTCCAAAGCCAAAGGGCAGCCCCGAGAACCACA GGTGTGCACCCTGCCCCCATCCCGGGATGAGCTGACCAAGAACCAGGTGAGC CTCTCGTGCAGTCAAAGGCTTCTATCCAGCGACATCGCCGTGGAGTGGG AGAGCAATGGGCAGCCGGAGAACAACACTACAAGACCACGCCTCCCGTGCTGGA CTCCGACGGCTCCTTCTTCTCGTGAGCAAGCTCACCCTGGACAAGAGCAGG TGGCAGCAGGGGAACGTCTTCTCATGCTCCGTGATGCATGAGGCTCTGCACA ACCACTACACGCAGAAGAGCCTCTCCCTGTCTCCGGGTAAA	
9D11_LC [N95Q]	GATATTGTTATGACTCAATCTCCACTGTCTCTGCCGGTGACTCCAGGCGAAC CGGCGAGCATTTCTTGCCGTTCAGCCAGTCTCTGCTGCACTCCAACGGCTA CAACTATCTCGATTGGTACCTGCAAAAACCGGGTCAGAGCCCTCAGCTGCTG ATCTACCTGGGCTCTAACCGCGCTTCCGGTGTACCGGACCGTTTCAGCGGCT CTGGATCCGGCACCGATTTACGTTGAAAATCAGCCGTGTTGAAGCAGAAGA CGTGGGCGTTTATTACTGTATGCAGGCAAGCATTATGCAGCGGACTTTTGGT CAAGGCACCAAGGTCGAAATTAAACGTACGGTGGCTGCACCATCTGTCTTCA TCTTCCCGCCATCTGATGAGCAGTTGAAATCTGGAAGTGCCTCTGTTGTGTG CCTGCTGAATAACTTCTATCCAGAGAGGCCAAAGTACAGTGGAAGGTGGAT AAGCCCTCCAATCGGGTAACTCCAGGAGAGTGTACAGAGCAGGACAGCA AGGACAGCACCTACAGCCTCAGCAGCACCCCTGACGCTGAGCAAAGCAGACTA CGAGAAACACAAAGTCTACGCCTGCGAAGTCACCCATCAGGGCCTGAGCTCG CCCGTCACAAAGAGCTTCAACAGGGGAGAGTGT	271
CD3_VLC H1	CAGGCCGTGCTGACCCAGGAACCCAGCCTGACAGTGTCTCCTGGCGGCACCG TGACCCTGACATGTGGCAGTTCTACAGGCGCCGTGACCACCAGCAACTACGC CAACTGGGTGCAGGAAAAGCCCGGCCAGGCCTTCAGAGGACTGATCGGCGGC ACCAACAAGAGAGCCCTGGCACCCCTGCCAGATTCAGCGGATCTCTGCTGG GAGGAAAGGCCGCCCTGACACTGTCTGGCGCCAGCCAGAAGATGAGGCCGA GTACTACTGCGCCCTGTGGTACAGCAACCTGTGGGTGTTCCGGCGGAGGCACC AAGCTGACAGTGTGAGCAGCGCTTCCACCAAAGGCCCTTCCGTGTTTCCCTC TGGCTCCTAGCTCCAAGTCCACCTCTGGAGGCACCGCTGCTCTCGGATGCCT CGTGAAGGATTATTTTCTGAGCCTGTGACAGTGTCTTGAATAGCGGAGCA CTGACCTCTGGAGTGACATACTTTCCCGCTGTGCTGCAGTCTCTGGACTGT ACAGCCTGAGCAGCGTGGTGACAGTGCCAGCAGCAGCCTGGGCACCCAGAC CTACATCTGCAACGTGAACCACAAGCCAGCAACACCAAGGTGGACAAGAAG GTGGAACCAAGTCTTGT	272
9D11 [VHCH1] - CD3[VHC H1- S100aA]- Fcknob_P GLALA	CAGGTGCAATTGGTTCAATCTGGTGCTGAAGTAAAAAAACCGGGCGCTTCCG TTAAAGTGAGCTGCAAAGCATCCGGATACACCTTCACTTCTATTACATGCA CTGGGTTTCGTCAAGCCCGGGCAGGGTCTGGAATGGATGGGCATATTAAC CCAAGCGGTGGCCCTACCTCCTACGCGCAGAAATTCCAGGGTCGCGTCACGA TGACCCGTGACACTAGCACCTCTACCGTTTATATGGAGCTGTCCAGCCTGCG TTCTGAAGATACTGCAGTGTACTACTGTGCACGCGGTGACTTTCGCTTGGCTG GACTATTGGGGTCAAGGCACCCTCGTAACGGTTTCTTCTGCTAGCACAAAGG GCCCCAGCGTGTTCCTCTGGCCCCCTAGCAGCAAGAGCACATCTGGCGGAAC AGCCGCCCTGGGCTGCCTCGTGAAGGACTACTTTCCCGAGCCTGTGACCGTG TCCTGGAACCTCTGGCGCCCTGACAAGCGGCGTGACACCTTTCCAGCCGTGC TGCAGAGCAGCGGCTGTACTCTCTGAGCAGCGTGGTCACCGTGCCTAGCAG CAGCCTGGGCACCCAGACCTACATCTGCAACGTGAACCACAAGCCAGCAAC ACCAAAGTGGACAAGAAGGTGGAGCCCAAGAGCTGTGATGGCGGAGGAGGGT CCGGAGGCGGAGGATCCGAGGTGCAGCTGCTGGAATCTGGCGGCGGACTGGT GCAGCCTGGCGGATCTCTGAGACTGAGCTGTGCCGCCAGCGGCTTACCTTC AGCACCTACGCCATGAACTGGGTGCGCCAGGCCCCCTGGCAAAGGCCTGGAAT GGGTGTCCCGGATCAGAAGCAAGTACAACAACACTACGCCACCTACTACGCCGA CAGCGTGAAGGGCCGGTTTACCATCAGCCGGGACGACAGCAAGAACACCCCTG TACCTGCAGATGAACAGCCTGCGGGCCGAGGACACCGCCGTGTACTATTGTG TGCGGCACGGCAACTTCGGCAACGCCTATGTGTCTTGGTTTGCCTACTGGG CCAGGGCACCCCTCGTGACCGTGTCAAGCGCTAGTGTGGCCGCTCCCTCCGTG	273

-267-

	<p>TTTATCTTTCCCCATCCGATGAACAGCTGAAAAGCGGCACCGCCTCCGTCTG TGTGTCTGCTGAACAATTTTTACCCTAGGGAAGCTAAAGTGCAGTGGAAAGT GGATAACGCACTGCAGTCCGGCAACTCCCAGGAATCTGTGACAGAACAGGAC TCCAAGGACAGCACCTACTCCCTGTCTCCACCTGACACTGTCTAAGGCTG ATTATGAGAAACACAAAGTCTACGCCTGCGAAGTCACCCATCAGGGCCTGAG CTCGCCCGTACAAAGAGCTTCAACAGGGGAGAGTGTGACAAGACCCACACC TGTCCCCCTTGTCTGCCCCCTGAAGCTGCTGGCGGCCCTTCTGTGTTCTGT TCCCCCAAAGCCCAAGGACACCCTGATGATCAGCCGGACCCCCGAAGTGAC CTGCGTGGTGGTGGATGTGTCCCACGAGGACCCTGAAGTGAAGTTCAATTGG TACGTGGACGGCGTGGAAGTGCACAACGCCAAGACAAAGCCGCGGGAGGAGC AGTACAACAGCACGTACCGTGTGGTTCAGCGTCTCACCCTCCTGCACCAGGA CTGGCTGAATGGCAAGGAGTACAAGTGCAAGGTCTCCAACAAAGCCCTCGGC GCCCCCATCGAGAAAACCATCTCCAAAGCCAAAGGGCAGCCCCGAGAACCAC AGGTGTACACCCTGCCCCATGCCGGGATGAGCTGACCAAGAACCAGGTCAG CCTGTGGTGCCTGGTCAAAGGCTTCTATCCCAGCGACATCGCCGTGGAGTGG GAGAGCAATGGGCAGCCGGAGAACAATAACAAGACCACGCCTCCCGTGTCTGG ACTCCGACGGCTCCTTCTTCTCTACAGCAAGCTCACCCTGGACAAGAGCAG GTGGCAGCAGGGGAACGTCTTCTCATGCTCCGTGATGCATGAGGCTCTGCAC AACCCTACACGCAGAAGAGCCTCTCCCTGTCTCCGGGTAA</p>	
--	---	--

Name	Sequence	Seq ID No
16D5 variant W96Y/D5 2E VH	<p>GAGGTGCAATTGGTGGAAAGCGGAGGCGGCCTCGTGAAGCCTGGCGGATCTCT GAGACTGAGCTGTGCCGCCAGCGGCTTACCTTCAGCAACGCCTGGATGAGCT GGGTGCGCCAGGCCCCCTGGAAAAGGACTCGAGTGGGTGGGACGGATCAAGAGC AAGACCGAGGGCGGCACCACCGACTATGCCGCCCTGTGAAGGGCCGGTTCAC CATCAGCAGGGACGACAGCAAGAACACCCTGTACCTGCAGATGAACAGCCTGA AAACCGAGGACACCGCCGTGTACTACTGCACCACCCCTACGAGTGGTCTTGG TACGACTACTGGGGCCAGGGCACCCCTCGTGACCGTGTCTATCT</p>	415
W96Y/D 52E- _CD3- VHCH1 _Fc- knob_PG LALA pETR149 45	<p>GAGGTGCAATTGGTGGAAAGCGGAGGCGGCCTCGTGAAGCCTGGCGGATCTCT GAGACTGAGCTGTGCCGCCAGCGGCTTACCTTCAGCAACGCCTGGATGAGCT GGGTGCGCCAGGCCCCCTGGAAAAGGACTCGAGTGGGTGGGACGGATCAAGAGC AAGACCGAGGGCGGCACCACCGACTATGCCGCCCTGTGAAGGGCCGGTTCAC CATCAGCAGGGACGACAGCAAGAACACCCTGTACCTGCAGATGAACAGCCTGA AAACCGAGGACACCGCCGTGTACTACTGCACCACCCCTACGAGTGGTCTTGG TACGACTACTGGGGCCAGGGCACCCCTCGTGACCGTGTCTATCTGCTAGCACAAA GGGCCCTAGCGTGTTCCTCTGGCCCCCAGCAGCAAGAGCACAAAGCGGCGGAA CAGCCGCCCTGGGCTGCCTCGTGAAGGACTACTTCCCCGAGCCCGTGACAGTG TCTTGGAACAGCGGAGCCCTGACAAGCGGCGTGCACACCTTCCCTGCCGTGCT GCAGAGCAGC GGCCTGTACTCCCTGAGCAGCGTGGTACCCTGCCTAGCAGCAGCCTGGGCAC CCAGACCTACATCTGCAACGTGAACCACAAGCCCAGCAACACCAAAGTGGACA AGAAGGTGGAGCCCAAGAGCTGTGATGGCGGAGGAGGTCCGGAGGCGGAGGA TCCGAGGTGCAGCTGCTGGAATCTGGCGGCGGACTGGTGCAGCCTGGCGGATC TCTGAGACTGAGCTGTGCCGCCAGCGGCTTACCTTCAGCACCTACGCCATGA ACTGGGTGCGCCAGGCCCCCTGGCAAAGGCCTGGAATGGGTGTCCCGGATCAGA AGCAAGTACAACAATACTACGCCACCTACTACGCCGACAGCGTGAAGGGCCGGTT CACCATCAGCCGGGACGACAGCAAGAACACCCTGTACCTGCAGATGAACAGCC TGCGGGCCGAGGACACCGCCGTGTACTATTGTGTGCGGCACGGCAACTTCGGC AACAGCTATGTGTCTTGGTTTGCCTACTGGGGCCAGGGCACCCCTCGTGACCGT GTCAAGCGCT AGTACCAAGGGCCCCAGCGTGTTCCTTGGCACCCAGCAGCAAGAGCACATC TGGCGGAACAGCCGCTCTGGGCTGTCTGGTGAAAGACTACTTCCCCGAGCCCG TGACCGTGTCTTGGAATCTGGCGCCCTGACCAGCGGCGTGCACACCTTTCCA GCCGTGCTGCAGAGCAGCGGCCTGTACTCCCTGTCTCCGTGGTACCGTGGC CTCTAGCTCCCTGGGAACACAGACATATCTGTAATGTCAATCACAAGCCTT CCAACACCAAAGTCGATAAGAAAGTCGAGCCCAAGAGCTGCGACAAAACCTCAC ACATGCCACCGTGCCAGCACCTGAAGCTGCAGGGGACCGTCAGTCTTCT</p>	416

	<p>CTTCCCCCAAACCCAAGGACACCCTCATGATCTCCCGGACCCCTGAGGTCA CATGCGTGGTGGTGGACGTGAGCCACGAAGACCCTGAGGTCAAGTTCAACTGG TACGTGGACGGCGTGGAGGTGCATAATGCCAAGACAAAGCCGCGGGAGGAGCA GTACAACAGC</p> <p>ACGTACCGTGTGGTTCAGCGTCTCACCCTGCTGCACCAGGACTGGCTGAATGG CAAGGAGTACAAGTGCAAGGTCTCCAACAAAGCCCTCGGCGCCCCCATCGAGA AAACCATCTCCAAAGCCAAAGGGCAGCCCCGAGAACCACAGGTGTACACCCTG CCCCCATGCCGGGATGAGCTGACCAAGAACCAGGTGAGCCTGTGGTGCCTGGT CAAAGGCTTCTATCCCAGCGACATCGCCGTGGAGTGGGAGAGCAATGGGCAGC CGGAGAACAACACTACAAGACCACGCCTCCCGTGTGGACTCCGACGGCTCCTTC TTCCTCTACAGCAAGCTCACCCTGGACAAGAGCAGGTGGCAGCAGGGGAACGT CTTCTCATGCTCCGTGATGCATGAGGCTCTGCACAACCACTACACGCAGAAGA GCCTCTCCCTGTCTCCGGGTAAA</p>	
W96Y/D 52E_Fc- hole_PG LALA_H YRF pETR149 46	<p>GAGGTGCAATTGGTGGAAAGCGGAGGCGGCCTCGTGAAGCCTGGCGGATCTCT GAGACTGAGCTGTGCCGCCAGCGGCTTACCTTCAGCAACGCCTGGATGAGCT GGGTGCGCCAGGCCCCCTGGAAAAGGACTCGAGTGGGTGGGACGGATCAAGAGC AAGACCGAGGGCGGCACCACCGACTATGCCGCCCTGTGAAGGGCCGGTTAC CATCAGCAGGGACGACAGCAAGAACACCCTGTACCTGCAGATGAACAGCCTGA AAACCGAGGACACCGCCGTGTACTACTGCACCACCCCTACGAGTGGTCTTGG TACGACTACTGGGGCCAGGGCACCTCTGTGACCGTGTCTATCTGTAGACCAA GGGCCCCCTCCGTGTTCCCCCTGGCCCCCAGCAGCAAGAGCACCAGCGGCGGCA CAGCCGCTCTGGGCTGCCTGGTCAAGGACTACTTCCCCGAGCCCGTGACCGTG TCCTGGAACAGCGGAGCCCTGACCTCCGGCGTGCACACCTTCCCCGCCGTGCT GCAGAGTTCT</p> <p>GGCCTGTATAGCCTGAGCAGCGTGGTACCCTGCCTTCTAGCAGCCTGGGCAC CCAGACCTACATCTGCAACGTGAACCACAAGCCCAGCAACACCAAGGTGGACA AGAAGGTGGAGCCCCAAGAGCTGCGACAAAACCTACACATGCCACCGTGCCCA GCACCTGAAGCTGCAGGGGGACCGTCAGTCTTCTCTTCCCCCAAACCCAA GGACACCCTCATGATCTCCCGGACCCCTGAGGTACATGCGTGGTGGTGGACG TGAGCCACGAAGACCCTGAGGTCAAGTTCAACTGGTACGTGGACGGCGTGGAG GTGCATAATGCCAAGACAAAGCCGCGGGAGGAGCAGTACAACAGCACGTACCG TGTTGGTCAGCGTCTCACCCTGCTGCACCAGGACTGGCTGAATGGCAAGGAGT ACAAGTGCAAGGTCTCCAACAAAGCCCTCGGCGCCCCCATCGAGAAAACCATC TCCAAAGCCAAAGGGCAGCCCCGAGAACCACAGGTGTGCACCCTGCCCCCATC CCGGGATGAG</p> <p>CTGACCAAGAACCAGGTGAGCCTCTCGTGCAGTCAAAGGCTTCTATCCCAG CGACATCGCCGTGGAGTGGGAGAGCAATGGGCAGCCGGAGAACAACACTACAAGA CCACGCCCTCCCGTGTGGACTCCGACGGCTCCTTCTTCTCTCGTGAGCAAGCTC ACCGTGGACAAGAGCAGGTGGCAGCAGGGGAACGTCTTCTCATGCTCCGTGAT GCATGAGGCTCTGCACAACCGCTTACGCAGAAGAGCCTCTCCCTGTCTCCGG GTAAA</p>	417
14B1 VH	<p>GAGGTGCAATTGTTGGAGTCTGGGGGAGGCTTGGTACAGCCTGGGGGTCCCT GAGACTCTCCTGTGCAGCCTCCGGATTACCTTTAGCAGTTATGCCATGAGCT GGGTCCGCCAGGCTCCAGGGAAGGGGCTGGAGTGGGTCTCAGCTATTAGTGGT AGTGGTGGTAGCACATACTACGCAGACTCCGTGAAGGGCCGGTTACCATCTC CAGAGACAATTCCAAGAACACGCTGTATCTGCAGATGAACAGCCTGAGAGCCG AGGACACGGCCGTATATTACTGTGCGCTGGTGAAGTACCCTTACCGTTACTTC GACTACTGGGGCCAAGGAACCCCTGGTACCCTCTCGAGT</p>	418
14B1 VL	<p>TCTTCTGAAGTACTCAAGATCCAGCTGTTAGCGTGGCTCTGGGTGAGACTGT ACGTATCACCTGCCAAGGCGATTCTCTGCGCTCCTACTACGCAAGCTGGTACC AGCAGAAACCGGGTCAGGCCCAAGTTCTGGTGATTTACGGCAAAAACAACCGT CCGTCTGGGATCCCGGACCGTTTCTCCGGCAGCTCTTCCGGTAACACGGCGAG CCTCACCATCACTGGCGCTCAAGCAGAAGACGAGGCCGACTATTACTGTAAGT CTCGGGAAGCCCAACCGGCCTGGTTGTCTTCGGTGGCGGTACCAAGCTG ACCGTCTTA</p>	419
14B1[EE] _CD3[V LCH1]_F	<p>GAGGTGCAATTGTTGGAGTCTGGGGGAGGCTTGGTACAGCCTGGGGGTCCCT GAGACTCTCCTGTGCAGCCTCCGGATTACCTTTAGCAGTTATGCCATGAGCT GGGTCCGCCAGGCTCCAGGGAAGGGGCTGGAGTGGGTCTCAGCTATTAGTGGT AGTGGTGGTAGCACATACTACGCAGACTCCGTGAAGGGCCGGTTACCATCTC CAGAGACAATTCCAAGAACACGCTGTATCTGCAGATGAACAGCCTGAGAGCCG AGGACACGGCCGTATATTACTGTGCGCTGGTGAAGTACCCTTACCGTTACTTC GACTACTGGGGCCAAGGAACCCCTGGTACCCTCTCGAGT</p>	420

c- knob_PG LALA pETR149 76	<p>CAGAGACAATTCCAAGAACACGCTGTATCTGCAGATGAACAGCCTGAGAGCCG AGGACACGGCCGTATATTACTGTGCGCGTGGTGACTACCGTTACCGTTACTTC GACTACTGGGGCCAAGGAACCCCTGGTCACCGTCTCGAGTGCTAGCACCAAGGG CCCCCTCCGTGTTTCCTCTGGCCCCCTTCCAGCAAGTCCACCTCTGGCGGAACTG CCGCTCTGGGCTGCCTGGTGGAAGATTACTTCCCCGAGCCCGTGACCGTGTCC TGGAATTCTGGCGCTCTGACCTCCGGCGTGCACACCTTTCCAGCTGTGCTGCA GTCCTCCGGCCTGTACTCCCTGTCTCCGTCTGACAGTGCCCTCCAGCTCTC TGGGCACCCAGACCTACATCTGCAACGTGAACCACAAGCCCTCCAACACCAAG GTGGACGAGAAGGTGGAACCCAAGTCTTGCACGGTGGCGGAGGTTCCGGAGG CGGAGGATCCCAGGCTGTCTGACCCAGGAACCCCTCCCTGACAGTGTCTCCTG GCGGCACCGTGACCCCTGACCTGTGGATCTTCTACCGGCGCTGTGACCACCTCC AACTACGCCAATTGGGTGCAGGAAAAGCCCGGCCAGGCCTTCAGAGGACTGAT CGGCGGCACCAACAAGAGAGCCCTGGCACCCCTGCCAGATTCTCCGGTTCTC TGCTGGGCGGCAAGGCTGCCCTGACTCTGTCTGGTGCTCAGCCTGAGGACGAG GCCGAGTACTACTGCGCCCTGTGGTACTCCAACCTGTGGGTGTTCCGGCGGAGG CACCAAGCTGACCGTGCTGTCCAGCGCTTCCACCAAGGGACCCAGTGTGTTCC CCCTGGCCCCCAGCTCCAAGTCTACATCCGGTGGCACAGCTGCCCTGGGATGT CTCGTGAAGGACTACTTTCTTGAGCCTGTGACAGTGTCTTGGAACAGCGGAGC CCTGACCAGCGGAGTGCACACATTCCTTGCAGTGTCTGCAGAGCAGCGGCCTGT ATAGCCTGAGCAGCGTCTGACCGTGCCCTTCTCTAGCCTGGGAACACAGAGA TATATCTGTAATGTGAATCATAAGCCCCAGTAATACCAAGTGGATAAGAAAGT GGAACCTAAGAGCTGCGATAAGACCCACACCTGTCCCCCTGCCCTGCTCCTG AAGCTGCTGGTGGCCCTAGCGTGTTCTGTTCCTTCCCCCAAAGCCCAAGGACACC CTGATGATCTCCCGGACCCCCGAAGTGACCTGCGTGGTGGTGGATGTGTCCCA CGAGGACCCTGAAGTGAAGTTCAATTGGTACGTGGACGGCGTGGAAGTGCACA ACGCCAAGACCAAGCCTAGAGAGGAACAGTACAACCTCCACCTACCGGGTGGTG TCCGTGCTGACAGTGTCTGCACCAGGACTGGCTGAACGGCAAAGAGTACAAGTG CAAGGTGTCCAACAAGGCCCTGGGCGCTCCCATCGAAAAGACCATCTCCAAGG CCAAGGGCCAGCCCCGGGAACCCCAAGGTGTACACCCTGCCCCCATGCCGGGAT GAGCTGACCAAGAACCAGGTGAGCCTGTGGTGCCTGGTCAAAGGCTTCTATCC CAGCGACATCGCCGTGGAGTGGGAGAGCAATGGGCAGCCGGAGAACAACCTACA AGACCACGCCTCCCGTGTCTGGACTCCGACGGCTCCTTCTTCTCTACAGCAAG CTCACCGTGGACAAGAGCAGGTGGCAGCAGGGGAACGTCTTCTCATGTCCGT GATGCATGAGGCTCTGCACAACCACTACACGCAGAAGAGCCTCTCCCTGTCTC CGGGTAAA</p>	
14B1[EE] _Fc- hole_PG LALA pETR149 77	<p>GAGGTGCAATTGTTGGAGTCTGGGGGAGGCTTGGTACAGCCTGGGGGGTCCCT GAGACTCTCCTGTGCAGCCTCCGGATTACCTTTAGCAGTTATGCCATGAGCT GGGTCCGCCAGGCTCCAGGGAAGGGGCTGGAGTGGGTCTCAGCTATTAGTGGT AGTGGTGGTAGCACATACTACGACAGTCCGTGAAGGGCCGGTTACCATCTC CAGAGACAATTCCAAGAACACGCTGTATCTGCAGATGAACAGCCTGAGAGCCG AGGACACGGCCGTATATTACTGTGCGCGTGGTGACTACCGTTACCGTTACTTC GACTACTGGGGCCAAGGAACCCCTGGTCACCGTCTCGAGTGCTAGCACCAAGGG CCCCCTCCGTGTTCCCCCTGGCCCCCAGCAGCAAGAGCACCAGCGCGGCACAG CCGCTCTGGGCTGCCTGGTGCAGGACTACTTCCCCGAGCCCGTGACCGTGTCC TGGAACAGCGGAGCCCTGACCTCCGGCGTGCACACCTTCCCCGCCGTGCTGCA GAGTTCTGGCCTGTATAGCCTGAGCAGCGTGGTACCGTGCCCTTCTAGCAGCC TGGGCACCCAGACCTACATCTGCAACGTGAACCACAAGCCCAGCAACACCAAG GTGGACGAGAAGGTGGAGCCCAAGAGCTGCGACAAAACCTCACACATGCCACC GTGCCCAGCACCTGAAGCTGCAGGGGGACCGTCAGTCTTCTCTTCCCCCAA AACCCAAGGACACCCTCATGATCTCCCGGACCCCTGAGGTACATGCGTGGTG GTGGACGTGAGCCACGAAGACCCTGAGGTCAAGTTCAACTGGTACGTGGACGG CGTGGAGGTGCATAATGCCAAGACAAAGCCGCGGGAGGAGCAGTACAACAGCA CGTACCGTGTGGTACAGCTCCTACCGTCTCTGCACCAGGACTGGCTGAATGGC AAGGAGTACAAGTGCAAGGTCTCCAACAAAGCCCTCGGCGCCCCCATCGAGAA AACCATCTCCAAGCCAAAGGGCAGCCCCGAGAACCACAGGTGTGACCCCTGC CCCCATCCCGGGATGAGCTGACCAAGAACCAGGTGAGCCTCTCGTGGCAGTC AAAGGCTTCTATCCCAGCGACATCGCCGTGGAGTGGGAGAGCAATGGGACCC GGAGAACAACATAAGACCACGCTCCCGTGGTGGACTCCGACGGCTCCTTCT TCCTCGTGAGCAAGCTCACCGTGGACAAGAGCAGGTGGCAGCAGGGGAACGTC</p>	421

	TTCTCATGCTCCGTGATGCATGAGGCTCTGCACAACCACTACACGCAGAAGAG CCTCTCCCTGTCTCCGGGTAAA	
14B1 LC [KK] Constant lambda pETR149 79	TCTTCTGAACTGACTCAAGATCCAGCTGTTAGCGTGGCTCTGGGTGCACTGT ACGTATCACCTGCCAAGGCGATTCTCTGCGCTCCTACTACGCAAGCTGGTACC AGCAGAAACCGGGTCAGGCCCCAGTTCTGGTGATTTACGGCAAAAACAACCGT CCGTCTGGGATCCCGGACCGTTTCTCCGGCAGCTCTTCCGGTAACACGGCGAG CCTCACCATCACTGGCGCTCAAGCAGAAGACGAGGCCGACTATTACTGTAACT CTCGGGAAGCCACCAACCGGCCTGGTTGTCTTCGGTGGCGGTACCAAGCTG ACCGTCTTAGGTCAACCCAAGGCTGCCCCAGCGTGACCCTGTTCCCCCCAG CAGCAAGAACTGCAGGCCAACAAGGCCACCCTGGTCTGCCTGATCAGCGACT TCTACCCAGGCGCCGTGACCGTGGCCTGGAAGGCCGACAGCAGCCCCGTGAAG GCCGGCGTGGAGACCACCACCCCAAGCAAGCAGAGCAACAACAAGTACGCCGC CAGCAGCTACCTGAGCCTGACCCCGAGCAGTGGAAGAGCCACAGGTCCTACA GCTGCCAGGTGACCCACGAGGGCAGCACCGTGGAGAAAACCGTGGCCCCCACC GAGTGCAGC	422
9C7 VH	CAGGTGCAATTGGTTCAATCTGGTGCTGAAGTAAAAAACCAGGGCGCTTCCGT TAAAGTGAGCTGCAAAGCATCCGGATACACCTTCACTTCCCTATTACATGCACT GGGTTTCGTCAAGCCCCGGGCCAGGGTCTGGAATGGATGGGCATCATTAACCCA AGCGGTGGCTCTACCTCCTACGCGCAGAAATTCAGGGTTCGCGTCACGATGAC CCGTGACACTAGCACCTCTACCGTTTATATGGAGCTGTCCAGCCTGCGTTCTG AAGATACTGCAGTGTACTACTGTGCACGCGGTGACTGGTCTTACTACATGGAC TATTGGGGTCAAGGCACCCTCGTAACGGTTTCTTCT	423
9C7 VL	GATATTGTTATGACTCAATCTCCACTGTCTCTGCCGGTGAAGTCCAGGCGAACC GGCAGCATTTCTTGCCGTTCCAGCCAGTCTCTGCTGCACTCCAACGGCTACA ACTATCTCGATTGGTACCTGCAAAAACCGGGTCAGAGCCCTCAGCTGCTGATC TACCTGGGCTCTAACCAGCGCTTCCGGTGTACCGGACCGTTTCAGCGGCTCTGG ATCCGGCACCGATTTACGTTGAAAATCAGCCGTGTTGAAGCAGAAGACGTGG GCGTTTATTACTGTAIGCAGGCACGGCAGACCCCAACTTTTGGTCAAGGCACC AAGGTCGAAATTAAA	424
9C7[EE] _CD3[V LCH1]_F c- knob_PG LALA pETR149 74	CAGGTGCAATTGGTTCAATCTGGTGCTGAAGTAAAAAACCAGGGCGCTTCCGT TAAAGTGAGCTGCAAAGCATCCGGATACACCTTCACTTCCCTATTACATGCACT GGGTTTCGTCAAGCCCCGGGCCAGGGTCTGGAATGGATGGGCATCATTAACCCA AGCGGTGGCTCTACCTCCTACGCGCAGAAATTCAGGGTTCGCGTCACGATGAC CCGTGACACTAGCACCTCTACCGTTTATATGGAGCTGTCCAGCCTGCGTTCTG AAGATACTGCAGTGTACTACTGTGCACGCGGTGACTGGTCTTACTACATGGAC TATTGGGGTCAAGGCACCCTCGTAACGGTTTCTTCTGCTAGCACCAAGGGCCC CTCCGTGTTTCTCTGGCCCCCTTCCAGCAAGTCCACCTCTGGCGGAAGTCCG CTCTGGGCTGCCTGGTGGAAGATTACTTCCCCGAGCCCGTGACCGTGTCTTG AATTCTGGCGCTCTGACCTCCGGCGTGCACACCTTTCAGCTGTGCTGCAGTC CTCCGGCCTGTACTCCCTGTCTCCGTCTGTGACAGTGCCCTCCAGCTCTCTGG GCACCCAGACCTACATCTGCAACGTGAACCACAAGCCCTCCAACCAAGGTG GACGAGAAGGTGGAACCAAGTCTGCGACGGTGGCGGAGGTTCCGGAGGCGG AGGATCCCAGGCTGTCTGACCCAGGAACCCCTCCCTGACAGTGTCTCTGGCG GCACCGTGACCTGACCTGTGGATCTTCTACCGGCGCTGTGACCACCTCCAAC TACGCCAATTGGGTGCAGGAAAAGCCCGGCCAGGCCTTCAGAGGACTGATCGG CGGCACCAACAAGAGAGCCCCCTGGCACCCCTGCCAGATTCTCCGGTTCTCTGC TGGGCGGCAAGGCTGCCCTGACTCTGTCTGGTGCTCAGCCTGAGGACGAGGCC GAGTACTACTGCGCCCTGTGGTACTCCAACCTGTGGGTGTTTCGGCGGAGGCAC CAAGCTGACCGTGTCTGTCAGCGCTTCCACCAAGGGACCCAGTGTGTTCCCC TGGCCCCCAGCTCCAAGTCTACATCCGGTGGCACAGCTGCCCTGGGATGTCTC GTGAAGGACTACTTTCCCTGAGCCTGTGACAGTGTCTTGGAACAGCGGAGCCCT GACCAGCGGAGTGCACACATTCCCTGCACTGTGACAGCAGCGGCCTGTATA GCCTGAGCAGCGTCTGTGACCGTGCCTTCCCTCTAGCCTGGGAACACAGACATAT ATCTGTAATGTGAATCATAAGCCCAGTAATACCAAGTGGATAAGAAAGTGGGA ACCTAAGAGCTGCGATAAGACCCACACCTGTCCCCCCTGCCCTGCTCCTGAAG CTGCTGGTGGCCCTAGCGTGTTCCTGTTCCCCCAAGCCCAAGGACACCCTG ATGATCTCCCGGACCCCGAAGTGACCTGCGTGGTGGTGGATGTGTCCACGA GGACCTGAAGTGAAGTTCAATTGGTACGTGGACGGCGTGGAAGTGCACAACG CCAAGACCAAGCCTAGAGAGGAACAGTACAACCTCCACCTACCGGGTGGTGTCC	425

	GTGCTGACAGTGCTGCACCAGGACTGGCTGAACGGCAAAGAGTACAAGTGCAA GGTGTCCAACAAGGCCCTGGGCGCTCCCATCGAAAAGACCATCTCCAAGGCCA AGGGCCAGCCCCGGGAACCCAGGTGTACACCCTGCCCCCATGCCGGGATGAG CTGACCAAGAACCAGGTACGCTGTGGTGCCTGGTCAAAGGCTTCTATCCCAG CGACATCGCCGTGGAGTGGGAGAGCAATGGGCAGCCGGAGAACAACATAAGA CCACGCCTCCCGTGCTGGACTCCGACGGCTCCTTCTTCTCTACAGCAAGCTC ACCGTGGACAAGAGCAGGTGGCAGCAGGGGAACGTCTTCTCATGCTCCGTGAT GCATGAGGCTCTGCACAACCACTACACGCAGAAGAGCCTCTCCCTGTCTCCGG GTAAA	
9C7[EE] _Fc- hole_PG LALA pETR149 75	CAGGTGCAATTGGTTCAATCTGGTGTGAAGTAAAAAACCAGGGCGCTTCCGT TAAAGTGAGCTGCAAAGCATCCGGATACACCTTCACTTCTATTACATGCACT GGGTTTCGTCAAGCCCCGGGCCAGGGTCTGGAATGGATGGGCATCATTAACCCA AGCGGTGGCTCTACCTCTACGCGCAGAAATTCCAGGGTCGCGTCACGATGAC CCGTGACACTAGCACCTCTACCGTTTATATGGAGCTGTCCAGCCTGCGTTCTG AAGATACTGCAGTGTACTACTGTGCACGCGGTGACTGGTCTTACTACATGGAC TATTGGGGTCAAGGCACCCCTCGTAACGGTTTCTTCTGCTAGCACCAAGGGCCC CTCCGTGTTCCCCCTGGCCCCCAGCAGCAAGAGCACCAGCGGCGGCACAGCCG CTCTGGGCTGCCTGGTCGAGGACTACTTCCCCGAGCCCGTGACCGTGTCTTGG AACAGCGGAGCCCTGACCTCCGGCGTGCACACCTTCCCCGCCGTGCTGCAGAG TTCTGGCCTGTATAGCCTGAGCAGCGTGGTCACCGTGCCTTCTAGCAGCCTGG GCACCCAGACCTACATCTGCAACGTGAACCACAAGCCCAGCAACACCAAGGTG GACGAGAAGGTGGAGCCCAAGAGCTGCGACAAAACACACATGCCCAACCGTG CCCAGACCTGAAGCTGCAGGGGGACCGTCAGTCTTCTCTTCCCCCAAAAC CCAAGGACACCCCTCATGATCTCCCGGACCCCTGAGGTACATGCGTGGTGGTG GACGTGAGCCACGAAGACCCCTGAGGTCAAGTTCAACTGGTACGTGGACGGCGT GGAGGTGCATAATGCCAAGACAAAGCCGCGGGAGGAGCAGTACAACAGCACGT ACCGTGTGGTCAGCGTCTTACCGTCTTGCACCAGGACTGGCTGAATGGCAAG GAGTACAAGTGCAAGGTCTCCAACAAAGCCCTCGGCGCCCCCATCGAGAAAAC CATCTCCAAAGCCAAAGGGCAGCCCCGAGAACCACAGGTGTGCACCCCTGCCCC CATCCCGGGATGAGCTGACCAAGAACCAGGTACGCTCTCGTGCGCAGTCAAA GGCTTCTATCCCAGCGACATCGCCGTGGAGTGGGAGAGCAATGGGCAGCCGGA GAACAACATAAGACCACGCCTCCCGTGCTGGACTCCGACGGCTCCTTCTTCC TCGTGAGCAAGCTCACCGTGGACAAGAGCAGGTGGCAGCAGGGGAACGTCTTC TCATGCTCCGTGATGCATGAGGCTCTGCACAACCACTACACGCAGAAGAGCCT CTCCCTGTCTCCGGGTAAA	426
9C7 LC [RK] pETR149 80	GATATTGTTATGACTCAATCTCCACTGTCTCTGCCGGTGACTCCAGGCGAACC GGCGAGCATTTCTTGCCGTTCCAGCCAGTCTCTGCTGCACTCCAACGGCTACA ACTATCTCGATTGGTACCTGCAAAAACCGGGTCAGAGCCCTCAGCTGCTGATC TACCTGGGCTCTAACCAGCGCTTCCGGTGTACCGGACCGTTTCAGCGGCTCTGG ATCCGGCACCGATTTCACGTTGAAAATCAGCCGTGTTGAAGCAGAAGACGTGG GCGTTTATTACTGTAATGCAGGCACGGCAGACCCCAACTTTTGGTCAAGGCACC AAGGTGCAAAATTAAACGTACGGTGGCTGCACCATCTGTCTTCATCTTCCCCGCC ATCTGATCGGAAGTTGAAATCTGGAACCTGCCTCTGTTGTGTGCTGCTGAATA ACTTCTATCCCAGAGAGGCCAAAGTACAGTGGAAGGTGGATAACGCCCTCCAA TCGGGTAACCTCCAGGAGAGTGTACAGAGCAGGACAGCAAGGACAGCACCTA CAGCCTCAGCAGCACCTGACGCTGAGCAAAGCAGACTACGAGAAACACAAAG TCTACGCCTGCGAAGTCACCCATCAGGGCCTGAGCTCGCCCGTCACAAAGAGC TTCAACAGGGGAGAGTGT	427

Exemplary anti-PD1 antagonist sequences

Description	Sequence	Seq ID No
anti-PDL1 antibody	QVQLVESGGGVVQPGRSLRLDCKASGITFSNSGMHWVRQAPGKGLE WVAVIWDGSKRYYADSVKGRFTISRDNKNTLFLQMNSLRAEDTA VYYCATNDDYWGQGLTVTVSSASTKGPSVFPLAPCSRSTSESTAAL GCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVTV	274

-272-

	PSSSLGTKTYTCNVDHKPSNTKVDKRVESKYGPCCPPCPAPEFLGG PSVFLFPPKPKDTLMISRTPEVTCVVVDVSQEDPEVQFNWYVDGVE VHNAKTKPREEQFNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKGLP SSIEKTISKAKGQPREPQVYTLPPSQEEMTKNQVSLTCLVKGFYPS DIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSRLTVDKSRWQEGN VFSCSVMHEALHNHYTQKSLSLSPG	
anti-PDL1 antibody	EIVLTQSPATLSLSPGERATLSCRASQSVSSYLAWYQQKPGQAPRL LIYDASNRAITGIPARFSGSGSGTDFTLTISILEPEDFAVYYCQQSS NWPRTFGQGTKVEIKRTVAAPSVFIFPPSDEQLKSGTASVVCLLNN FYPREAKVQWKVDNALQSGNSQESVTEQDSKDSSTYSLSSTLTLSKA DYEKHKVYACEVTHQGLSSPVTKSFNRGEC	275
anti-PDL1 antibody	QVQLVQSGVEVKKPGASVKVSCASGYTFTNYYMYWVRQAPGQGLE WMGGINPSNGGTNFNEKFKNRVTLTDSSTTTAYMELKSLQFDDTA VYYCARRDYRFDMGFDYWGQGTITVTVSSASTKGPSVFPLAPCSRST SESTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYS LSSVTVTPSSSLGKTYTCNVDHKPSNTKVDKRVESKYGPCCPPCP APEFLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSQEDPEVQFN WYVDGVEVHNAKTKPREEQFNSTYRVVSVLTVLHQDWLNGKEYKCK VSNKGLPSSIEKTISKAKGQPREPQVYTLPPSQEEMTKNQVSLTCL VKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSRLTVDK SRWQEGNVFSCSVMHEALHNHYTQKSLSLSPG	276
anti-PDL1 antibody	EIVLTQSPATLSLSPGERATLSCRASKGVSTSGYSYLHWYQQKPGQ APRLLIYLAAYLESQVPAFSGSGSGTDFTLTISILEPEDFAVYYC QHSRDPLTFGGGTKVEIKRTVAAPSVFIFPPSDEQLKSGTASVVC LLNFFYPREAKVQWKVDNALQSGNSQESVTEQDSKDSSTYSLSSTLT LSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC	277
heavy	EVQLVESGGGLVQPGGSLRLSCAASGFTTFSDSWIHWVRQAPGKGLE WVAWISPYGGSTYYADSVKGRFTISADTSKNTAYLQMNSLRAEDTA VYYCARRHWPGGFDYWGQGTITVTVSSASTKGPSVFPLAPSSKSTSG GTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLS SVVTVTPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHTCPPC PAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKF NRYVDGVEVHNAKTKPREEQYASTYRVVSVLTVLHQDWLNGKEYKC KVSNAKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTC LVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVD KSRWQQGNVVFSCSVMHEALHNHYTQKSLSLSPG	278
light	DIQMTQSPSSLSASVGDRVTITCRASQDVSTAVAWYQQKPGKAPKL LIYSASFLYSGVPSRFSGSGSGTDFTLTISLQPEDFATYYCQQYL YHPATFGQGTKVEIKRTVAAPSVFIFPPSDEQLKSGTASVVCLLNN FYPREAKVQWKVDNALQSGNSQESVTEQDSKDSSTYSLSSTLTLSKA DYEKHKVYACEVTHQGLSSPVTKSFNRGEC	279
anti-PDL1 antibody VH	EVQLVESGGGLVQPGGSLRLSCAASGFTTFSDSWIHWVRQAPGKGLE WVAWISPYGGSTYYADSVKGRFTISADTSKNTAYLQMNSLRAEDTA VYYCARRHWPGGFDYWGQGTITVTVSS	280
anti-PDL1 antibody VH	EVQLVESGGGLVQPGGSLRLSCAASGFTTFSDSWIHWVRQAPGKGLE WVAWISPYGGSTYYADSVKGRFTISADTSKNTAYLQMNSLRAEDTA VYYCARRHWPGGFDYWGQGTITVTVSSASTK	281
anti-PDL1 antibody VL	DIQMTQSPSSLSASVGDRVTITCRASQDVSTAVAWYQQKPGKAPKL LIYSASFLYSGVPSRFSGSGSGTDFTLTISLQPEDFATYYCQQYL YHPATFGQGTKVEIKR	282
HVR-H1	GFTFSX1SWIH	283
HVR-H2	AWIX2PYGGSX3YYADSVKG	284
HVR-H3	RHWPGGFDY	285
HVR-L1	RASQX4X5X6TX7X8A	286

-273-

HVR-L2	SASX9LX10S	287
HVR-L3	QQX11X12X13X14PX15T	288
HVR-H1	GFTFSDSWIH	289
HVR-H2	AWISPYGGSTYYADSVKG	290
HVR-H3	RHWPGGFDY	291
HVR-L1	RASQDVSTAVA	292
HVR-L2	SASFLYS	293
HVR-L3	QQYLYHPAT	294
anti-PDL1 antibody HC-FR1	EVQLVESGGGLVQPGGSLRLSCAAS	295
anti-PDL1 antibody HC-FR2	HC-FR2 is WVRQAPGKGLEWV	296
anti-PDL1 antibody HC-FR3	RFTISADTSKNTAYLQMNSLRAEDTAVYYCAR	297
anti-PDL1 antibody HC-FR4	WGQGTLVTVSA	298
anti-PDL1 antibody HC-FR4	WGQGTLVTVSS	299
LC-FR1	DIQMTQSPSSLSASVGDRVTITC	300
LC-FR2	WYQQKPGKAPKLLIY	301
LC-FR3	GVPSRFSGSGSGTDFTLTISLQPEDFATYYC	302
LC-FR4	FGQGTKVEIKR	303
anti-PDL1 antibody VH	EVQLVESGGGLVQPGGSLRLSCAASGFTFSDSWIHWVRQAPGKGLE WVAWISPYGGSTYYADSVKGRFTISADTSKNTAYLQMNSLRAEDTA VYYCARRHWPGGFDYWGQGTLVTVSA	382
anti-PDL1 antibody VL	DIQMTQSPSSLSASVGDRVTITCRASQDVSTAVAWYQQKPGKAPKL LIYSASFLYSGVPSRFSGSGSGTDFTLTISLQPEDFATYYCQQYL YHPATFGQGTKVEIKR	383

Exemplary anti-TIM3 antibody sequences

Sequences of exemplary anti-TIM3 antibody amino acid sequences and exemplary TIM3 sequences are set forth in the sequence listing below as follows:

- | | | |
|----|---------------|-------------------------------|
| 5 | SEQ ID NO:304 | heavy chain HVR-H1, Tim3_0016 |
| | SEQ ID NO:305 | heavy chain HVR-H2, Tim3_0016 |
| | SEQ ID NO:306 | heavy chain HVR-H3, Tim3_0016 |
| | SEQ ID NO:307 | light chain HVR-L1, Tim3_0016 |
| | SEQ ID NO:308 | light chain HVR-L2, Tim3_0016 |
| 10 | SEQ ID NO:309 | light chain HVR-L3, Tim3_0016 |

	SEQ ID NO:310	heavy chain variable domain VH, Tim3_0016
	SEQ ID NO:311	light chain variable domain VL, Tim3_0016
	SEQ ID NO:312	heavy chain variable domain VH, Tim3_0016 variant (0018)
	SEQ ID NO:313	light chain variable domain VL, Tim3_0016 variant (0018)
5	SEQ ID NO:314	light chain HVR-L1, Tim3_0016_HVR-L1 variant 1_ NQ (removal of glycosylation site by N to Q mutation)
	SEQ ID NO:315	light chain HVR-L1, Tim3_0016_HVR-L1 variant 2_ NS (removal of glycosylation site by N to S mutation)
	SEQ ID NO:316	heavy chain HVR-H1, Tim3_0021
10	SEQ ID NO:317	heavy chain HVR-H2, Tim3_0021
	SEQ ID NO: 318	heavy chain HVR-H3, Tim3_0021
	SEQ ID NO:319	light chain HVR-L1, Tim3_0021
	SEQ ID NO:320	light chain HVR-L2, Tim3_0021
	SEQ ID NO:321	light chain HVR-L3, Tim3_0021
15	SEQ ID NO:322	heavy chain variable domain VH, Tim3_0021
	SEQ ID NO:323	light chain variable domain VL, Tim3_0021
	SEQ ID NO:324	heavy chain HVR-H1, Tim3_0022
	SEQ ID NO:325	heavy chain HVR-H2, Tim3_0022
	SEQ ID NO:326	heavy chain HVR-H3, Tim3_0022
20	SEQ ID NO:327	light chain HVR-L1, Tim3_0022
	SEQ ID NO:328	light chain HVR-L2, Tim3_0022
	SEQ ID NO:329	light chain HVR-L3, Tim3_0022
	SEQ ID NO:330	heavy chain variable domain VH, Tim3_0022
	SEQ ID NO:331	light chain variable domain VL, Tim3_0022
25	SEQ ID NO:332	heavy chain HVR-H1, Tim3_0026
	SEQ ID NO:333	heavy chain HVR-H2, Tim3_0026
	SEQ ID NO:334	heavy chain HVR-H3, Tim3_0026
	SEQ ID NO:335	light chain HVR-L1, Tim3_0026
	SEQ ID NO:336	light chain HVR-L2, Tim3_0026
30	SEQ ID NO:337	light chain HVR-L3, Tim3_0026
	SEQ ID NO:338	heavy chain variable domain VH, Tim3_0026
	SEQ ID NO:339	light chain variable domain VL, Tim3_0026
	SEQ ID NO:340	heavy chain HVR-H1, Tim3_0028
	SEQ ID NO:341	heavy chain HVR-H2, Tim3_0028
35	SEQ ID NO:342	heavy chain HVR-H3, Tim3_0028
	SEQ ID NO:343	light chain HVR-L1, Tim3_0028
	SEQ ID NO:344	light chain HVR-L2, Tim3_0028
	SEQ ID NO:345	light chain HVR-L3, Tim3_0028
	SEQ ID NO:346	heavy chain variable domain VH, Tim3_0028
40	SEQ ID NO:347	light chain variable domain VL, Tim3_0028
	SEQ ID NO:348	heavy chain HVR-H1, Tim3_0030
	SEQ ID NO:349	heavy chain HVR-H2, Tim3_0030
	SEQ ID NO:350	heavy chain HVR-H3, Tim3_0030
	SEQ ID NO:351	light chain HVR-L1, Tim3_0030
45	SEQ ID NO:352	light chain HVR-L2, Tim3_0030
	SEQ ID NO:353	light chain HVR-L3, Tim3_0030
	SEQ ID NO:354	heavy chain variable domain VH, Tim3_0030
	SEQ ID NO:355	light chain variable domain VL, Tim3_0030
	SEQ ID NO:356	heavy chain HVR-H1, Tim3_0033

-275-

- SEQ ID NO:357 heavy chain HVR-H2, Tim3_0033
 SEQ ID NO:358 heavy chain HVR-H3, Tim3_0033
 SEQ ID NO:359 light chain HVR-L1, Tim3_0033
 SEQ ID NO:360 light chain HVR-L2, Tim3_0033
 5 SEQ ID NO:361 light chain HVR-L3, Tim3_0033
 SEQ ID NO:362 heavy chain variable domain VH, Tim3_0033
 SEQ ID NO:363 light chain variable domain VL, Tim3_0033
 SEQ ID NO:364 heavy chain HVR-H1, Tim3_0038
 SEQ ID NO:365 heavy chain HVR-H2, Tim3_0038
 10 SEQ ID NO:366 heavy chain HVR-H3, Tim3_0038
 SEQ ID NO:367 light chain HVR-L1, Tim3_0038
 SEQ ID NO:368 light chain HVR-L2, Tim3_0038
 SEQ ID NO:369 light chain HVR-L3, Tim3_0038
 SEQ ID NO:370 heavy chain variable domain VH, Tim3_0038
 15 SEQ ID NO:371 light chain variable domain VL, Tim3_0038
 SEQ ID NO:372 an exemplary Pseudomonas exotoxin A variant 1 (deimmunized PE24 example)
 SEQ ID NO:373 an exemplary Pseudomonas exotoxin A variant 2 (deimmunized PE24 example)
 20 SEQ ID NO:374 human kappa light chain constant region
 SEQ ID NO:375 human lambda light chain constant region
 SEQ ID NO:376 human heavy chain constant region derived from IgG1
 SEQ ID NO:377 human heavy chain constant region derived from IgG1 with mutations L234A and L235A
 25 SEQ ID NO:378 human heavy chain constant region derived from IgG1 with mutations L234A, L235A and P329G
 SEQ ID NO:379 human heavy chain constant region derived from IgG4
 SEQ ID NO:380 exemplary human TIM3 sequences
 SEQ ID NO:381 human TIM3 Extracellular Domain (ECD)

30

<210> 304<211> 9<212> PRT<213> Mus musculus

35

<400> 304Gly Phe Ser Leu Ser Thr Ser Gly Met1 5

40

<210> 305<211> 3<212> PRT45 <213> Mus musculus<400> 305

-276-

Leu Asn Asp15 <210> 306<211> 8<212> PRT<213> Mus musculus10 <400> 306Asn Gly Tyr Leu Tyr Ala Leu Asp1 5

15

<210> 307<211> 6<212> PRT<213> Mus musculus

20

<400> 307Ser Ser Ser Val Asn Tyr1 5

25

<210> 308<211> 3<212> PRT30 <213> Mus musculus<400> 308Asp Ala Phe35 1<210> 309<211> 740 <212> PRT<213> Mus musculus<400> 30945 Trp Ser Ser Tyr Pro Trp Thr1 5<210> 310

-277-

<211> 120<212> PRT<213> Mus musculus5 <400> 310

Gln Val Thr Leu Lys Glu Ser Gly Pro Gly Ile Leu Gln Pro Ser Gln
1 5 10 15

10

Thr Leu Arg Leu Thr Cys Ser Phe Ser Gly Phe Ser Leu Ser Thr Ser
20 25 30

15

Gly Met Ser Val Gly Trp Ile Arg Gln Pro Ser Gly Lys Gly Leu Glu
35 40 45

20

Trp Leu Ala His Ile Trp Leu Asn Asp Asp Val Phe Phe Asn Pro Ala
50 55 60

25

Leu Lys Ser Arg Leu Thr Ile Ser Lys Asp Thr Ser Asn Asn Gln Val
65 70 75 80

Phe Leu Gln Ile Ala Ser Val Val Thr Ala Asp Thr Ala Thr Tyr Tyr
85 90 95

30

Cys Val Arg Ala Asn Gly Tyr Leu Tyr Ala Leu Asp Tyr Trp Gly Gln
100 105 110

35

Gly Thr Ser Val Thr Val Ser Ser
115 120

40

<210> 311<211> 106<212> PRT<213> Mus musculus

45

<400> 311

Gln Ile Val Leu Thr Gln Ser Pro Ala Ile Met Ser Ala Ser Pro Gly
1 5 10 15

-278-

- Gln Lys Val Thr Ile Thr Cys Ser Ala Ser Ser Ser Val Asn Tyr Thr
20 25 30
- 5 Gln Trp Tyr Gln Gln Lys Leu Gly Ser Ser Pro Lys Leu Trp Ile Tyr
35 40 45
- 10 Asp Ala Phe Lys Leu Ala Pro Gly Val Pro Ala Arg Phe Ser Gly Ser
50 55 60
- 15 Gly Thr Gly Thr Ser Tyr Ser Leu Thr Ile Ser Ser Met Glu Ala Glu
65 70 75 80
- Asp Ala Ala Ser Tyr Phe Cys His Gln Trp Ser Ser Tyr Pro Trp Thr
85 90 95
- 20 Phe Gly Gly Gly Thr Lys Leu Glu Ile Lys
100 105
- 25 <210> 312
<211> 120
<212> PRT
<213> Mus musculus
- 30 <400> 312
Gln Val Thr Leu Lys Glu Ser Gly Pro Gly Ile Leu Gln Pro Ser Gln
1 5 10 15
- 35 Thr Leu Ser Leu Thr Cys Ser Phe Ser Gly Phe Ser Leu Ser Thr Ser
20 25 30
- 40 Gly Met Ser Val Gly Trp Ile Arg Gln Pro Ser Gly Lys Gly Leu Glu
35 40 45
- 45 Trp Leu Ala His Ile Trp Leu Asn Asp Asp Val Phe Phe Asn Pro Ala
50 55 60
- Leu Lys Arg Arg Leu Thr Ile Ser Lys Asp Thr Ser Asn Asn Gln Val
65 70 75 80

-279-

5 Phe Leu Gln Ile Ala Ser Val Val Thr Ala Asp Thr Ala Thr Tyr Tyr
 85 90 95

10 Cys Val Arg Ala Asn Gly Tyr Leu Tyr Ala Leu Asp Tyr Trp Gly Gln
 100 105 110

10 Gly Ile Ser Val Thr Val Ser Ser
 115 120

15 <210> 313
 <211> 106
 <212> PRT
 <213> Mus musculus

20 <400> 313

25 Gln Ile Val Leu Thr Gln Ser Pro Ala Ile Met Ser Ala Ser Pro Gly
 1 5 10 15

30 Gln Lys Val Thr Ile Thr Cys Ser Ala Ser Ser Ser Val Asn Tyr Thr
 20 25 30

30 Gln Trp Tyr Gln Gln Lys Leu Gly Ser Ser Pro Lys Leu Trp Ile Tyr
 35 40 45

35 Asp Ala Phe Lys Leu Ala Pro Gly Val Pro Ala Arg Phe Ser Gly Ser
 50 55 60

40 Gly Thr Gly Thr Ser Tyr Ser Leu Thr Ile Ser Ser Met Glu Ala Glu
 65 70 75 80

45 Asp Ala Ala Ser Tyr Phe Cys His Gln Trp Ser Ser Tyr Pro Trp Thr
 85 90 95

45 Phe Gly Gly Gly Thr Lys Leu Glu Ile Lys
 100 105

-280-

5 <210> 314
 <211> 6
 <212> PRT
 <213> Mus musculus

 <400> 314

 Ser Ser Ser Val Gln Tyr
 1 5

10

<210> 315
 <211> 6
 <212> PRT
15 <213> Mus musculus

 <400> 315

 Ser Ser Ser Val Ser Tyr
20 1 5

<210> 316
 <211> 7
25 <212> PRT
 <213> Mus musculus

 <400> 316

30 Gly Tyr Ser Phe Thr Ser Tyr
 1 5

<210> 317
35 <211> 3
 <212> PRT
 <213> Mus musculus

 <400> 317
40 Ser Asp Ser
 1

45 <210> 318
 <211> 9
 <212> PRT
 <213> Mus musculus

-281-

<400> 318Gly Tyr Tyr Ala Trp Tyr Tyr Phe Asp1 5

5

<210> 319<211> 7<212> PRT10 <213> Mus musculus<400> 319Ser Gln Ser Ile Gly Asn Asn15 1 5<210> 320<211> 320 <212> PRT<213> Mus musculus<400> 32025 Tyr Ala Ser1<210> 32130 <211> 6<212> PRT<213> Mus musculus<400> 321

35

Ser Asn Ser Trp Pro Leu1 540 <210> 322<211> 120<212> PRT<213> Mus musculus45 <400> 322Gln Val Gln Leu Gln Gln Ser Gly Pro Gln Leu Val Arg Pro Gly Ala1 5 10 15

-282-

- Ser Val Gln Ile Ser Cys Lys Ala Ser Gly Tyr Ser Phe Thr Ser Tyr
20 25 30
- 5
- Leu Leu His Trp Leu Lys Gln Arg Pro Gly Gln Gly Leu Glu Trp Ile
35 40 45
- 10
- Gly Met Ile Asp Pro Ser Asp Ser Glu Thr Arg Leu Asn Gln Lys Phe
50 55 60
- 15
- Lys Asp Lys Ala Thr Leu Thr Val Asp Lys Ser Ser Ser Thr Ala Tyr
65 70 75 80
- 20
- Met Gln Leu Ser Ser Pro Thr Ser Glu Asp Ser Ala Val Tyr Tyr Cys
85 90 95
- 25
- Ala Arg Asp Gly Tyr Tyr Ala Trp Tyr Tyr Phe Asp Cys Trp Gly Gln
100 105 110
- 30
- Gly Thr Thr Leu Thr Val Ser Ser
115 120
- 35
- <210> 323
<211> 107
<212> PRT
<213> Mus musculus
- 40
- Asp Ile Val Leu Thr Gln Ser Pro Ala Thr Leu Ser Val Thr Pro Gly
1 5 10 15
- 45
- Asp Arg Val Ser Leu Ser Cys Arg Ala Ser Gln Ser Ile Gly Asn Asn
20 25 30
- Leu His Trp Tyr Gln Gln Lys Ser His Glu Ser Pro Arg Leu Leu Ile
35 40 45
- Lys Tyr Ala Ser His Ser Ile Ser Gly Ile Pro Ser Lys Phe Ser Gly

-283-

50 55 60

5 Thr Gly Ser Gly Thr Asp Phe Thr Leu Ser Phe Asn Ser Val Glu Thr
65 70 75 80

Glu Asp Phe Gly Met Tyr Phe Cys Gln Gln Ser Asn Ser Trp Pro Leu
85 90 95

10

Thr Phe Gly Ala Gly Thr Lys Leu Glu Leu Lys
100 105

15

<210> 324
<211> 5
<212> PRT
<213> Mus musculus

20

<400> 324

Gly Asp Ser Ile Ala
1 5

25

<210> 325
<211> 3
<212> PRT
30 <213> Mus musculus

<400> 325

Tyr Ser Gly
35 1

<210> 326
<211> 4
40 <212> PRT
<213> Mus musculus

<400> 326

45 Asp Tyr Phe Asp
1

<210> 327

-284-

<211> 7<212> PRT<213> Mus musculus5 <400> 327Arg Gln Asp Val Arg Lys Asn1 5

10

<210> 328<211> 3<212> PRT<213> Mus musculus

15

<400> 328Tyr Thr Ser1

20

<210> 329<211> 6<212> PRT25 <213> Mus musculus<400> 329Tyr Asp Asn Leu Pro Phe30 1 5<210> 330<211> 11435 <212> PRT<213> Mus musculus<400> 33040 Glu Val Gln Leu Gln Glu Ser Gly Pro Ser Leu Val Lys Pro Ser Gln1 5 10 15Thr Leu Ser Leu Thr Cys Ser Val Thr Gly Asp Ser Ile Ala Ser Ala45 20 25 30Tyr Trp Asn Trp Ile Arg Lys Phe Pro Gly Asn Lys Leu Glu Tyr Met35 40 45

-285-

5 Gly Tyr Ile Asn Tyr Ser Gly Ser Thr Tyr Tyr Asn Pro Ser Leu Lys
 50 55 60

10 Ser Arg Ile Ser Ile Thr Arg Asp Thr Ser Gln Asn Gln Tyr Tyr Leu
 65 70 75 80

15 Gln Leu Asn Ser Val Thr Thr Glu Asp Thr Ala Thr Tyr Tyr Cys Val
 85 90 95

20 Thr Gly Asp Tyr Phe Asp Tyr Trp Gly Arg Gly Thr Thr Leu Thr Val
 100 105 110

25 Ser Ser

30 <210> 331
 <211> 107
 <212> PRT
 <213> Mus musculus

35 <400> 331

40 Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Tyr Leu Gly
 1 5 10 15

45 Gly Lys Val Thr Ile Thr Cys Lys Ala Arg Gln Asp Val Arg Lys Asn
 20 25 30

50 Ile Gly Trp Tyr Gln His Lys Pro Gly Lys Gly Pro Arg Leu Leu Ile
 35 40 45

55 Trp Tyr Thr Ser Thr Leu Gln Ser Gly Ile Pro Ser Arg Phe Ser Gly
 50 55 60

60 Ser Gly Ser Gly Arg Asp Tyr Ser Phe Asn Ile Asn Asn Leu Glu Pro
 65 70 75 80

-286-

Glu Asp Ile Ala Thr Tyr Tyr Cys Leu Gln Tyr Asp Asn Leu Pro Phe
85 90 95

5 Thr Phe Gly Thr Gly Thr Lys Leu Glu Ile Arg
100 105

10 <210> 332
<211> 5
<212> PRT
<213> Mus musculus

15 <400> 332
Gly Tyr Thr Phe Thr
1 5

20 <210> 333
<211> 3
<212> PRT
<213> Mus musculus

25 <400> 333
Glu Thr Tyr
1

30 <210> 334
<211> 4
<212> PRT
<213> Mus musculus

35 <400> 334

Gly Tyr Pro Ala
1

40 <210> 335
<211> 12
<212> PRT
45 <213> Mus musculus

<400> 335

Ser Arg Thr Ile Leu His Ser Ser Gly Asn Thr Tyr

-287-

1 5 10

<210> 336

5 <211> 3

<212> PRT

<213> Mus musculus

<400> 336

10

Lys Val Ser

1

15 <210> 337

<211> 6

<212> PRT

<213> Mus musculus

20 <400> 337

Asp Ser His Val Pro Phe

1 5

25

<210> 338

<211> 115

<212> PRT

<213> Mus musculus

30

<400> 338

Gln Ile Gln Leu Val Gln Ser Gly Pro Glu Leu Lys Lys Pro Gly Glu

1 5 10 15

35

Thr Val Lys Ile Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Asp Tyr

 20 25 30

40

Ser Met His Trp Val Lys Gln Ala Pro Gly Arg Gly Leu Lys Trp Met

 35 40 45

45 Gly Tyr Ile Asn Thr Glu Thr Tyr Glu Pro Thr Phe Gly Ala Asp Phe

 50 55 60

Lys Gly Arg Phe Ala Phe Ser Leu Asp Thr Ser Ala Thr Thr Ala Tyr

-288-

65 70 75 80

5 Leu Gln Ile Asn Ser Leu Lys Thr Glu Asp Thr Ala Thr Phe Phe Cys
 85 90 95

10 Gly Gly Gly Gly Tyr Pro Ala Tyr Trp Gly Gln Gly Thr Val Val Ile
 100 105 110

15 Val Ser Ala
 115

15 <210> 339
 <211> 112
 <212> PRT
 <213> Mus musculus
 <400> 339

20 Asp Val Leu Met Thr Gln Thr Pro Leu Ser Leu Pro Val Ser Leu Gly
 1 5 10 15

25 Asp Gln Ala Ser Ile Ser Cys Arg Ser Ser Arg Thr Ile Leu His Ser
 20 25 30

30 Ser Gly Asn Thr Tyr Leu Glu Trp Tyr Leu Gln Lys Pro Gly Gln Ser
 35 40 45

35 Pro Lys Leu Leu Ile Tyr Lys Val Ser Asn Arg Phe Ser Gly Val Pro
 50 55 60

40 Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Asn Ile
 65 70 75 80

45 Ser Arg Val Glu Ala Glu Asp Leu Gly Val Tyr Tyr Cys Phe Gln Asp
 85 90 95

Ser His Val Pro Phe Thr Phe Gly Thr Gly Thr Lys Leu Glu Ile Lys
 100 105 110

-289-

- <210> 340
<211> 7
<212> PRT
5 <213> Mus musculus
- <400> 340
- Gly Phe Asn Ile Lys Thr Thr
10 1 5
- <210> 341
<211> 3
15 <212> PRT
<213> Mus musculus
- <400> 341
- 20 Ala Asp Asp
1
- <210> 342
25 <211> 8
<212> PRT
<213> Mus musculus
- <400> 342
- 30 Phe Gly Tyr Val Ala Trp Phe Ala
1 5
- 35 <210> 343
<211> 7
<212> PRT
<213> Mus musculus
- 40 <400> 343
- Ser Gln Ser Val Asp Asn Tyr
1 5
- 45 <210> 344
<211> 3
<212> PRT
<213> Mus musculus

-290-

<400> 344

5 Tyr Ala Ser
1

<210> 345<211> 610 <212> PRT<213> Mus musculus<400> 345

15 His Tyr Ser Ser Pro Tyr
1 5

<210> 34620 <211> 119<212> PRT<213> Mus musculus<400> 346

25 Glu Val Gln Leu Gln Gln Ser Val Ala Glu Leu Val Arg Pro Gly Ala
1 5 10 15

30 Ser Val Lys Leu Ser Cys Thr Ala Ser Gly Phe Asn Ile Lys Thr Thr
20 25 30

35 Tyr Met His Trp Val Lys Gln Arg Pro Glu Gln Gly Leu Glu Trp Ile
35 40 45

40 Gly Arg Ile Asp Pro Ala Asp Asp Asn Thr Lys Tyr Ala Pro Lys Phe
50 55 60

Gln Gly Lys Ala Thr Ile Thr Ala Asp Thr Ser Ser Asn Thr Ala Tyr
65 70 75 80

45 Leu Gln Leu Ser Ser Leu Thr Ser Glu Asp Ala Ala Ile Tyr Tyr Cys
85 90 95

-291-

Val Arg Asp Phe Gly Tyr Val Ala Trp Phe Ala Tyr Trp Gly Gln Gly
100 105 110

5 Thr Leu Val Thr Phe Ser Ala
115

<210> 347
 10 <211> 107
<212> PRT
<213> Mus musculus

<400> 347
 15 Asn Ile Val Met Thr Pro Thr Pro Lys Phe Leu Pro Val Ser Ser Gly
1 5 10 15

20 Asp Arg Val Thr Met Thr Cys Arg Ala Ser Gln Ser Val Asp Asn Tyr
20 25 30

Val Ala Trp Tyr Gln Gln Lys Pro Gly Gln Ser Pro Lys Leu Leu Ile
 25 35 40 45

Tyr Tyr Ala Ser Asn Arg Tyr Ile Gly Val Pro Asp Arg Phe Thr Gly
50 55 60
 30

Ser Gly Ser Gly Thr Asp Phe Thr Phe Thr Ile Ser Ser Val Gln Val
65 70 75 80

35 Glu Asp Leu Ala Val Tyr Phe Cys Gln Gln His Tyr Ser Ser Pro Tyr
85 90 95

40 Thr Phe Gly Ser Gly Thr Lys Leu Glu Ile Lys
100 105

<210> 348
 45 <211> 7
<212> PRT
<213> Mus musculus

<400> 348

-292-

Gly Tyr Pro Phe Ser Glu Tyr1 5

5

<210> 349<211> 3<212> PRT<213> Mus musculus

10

<400> 349Glu Thr Gly1

15

<210> 350<211> 4<212> PRT20 <213> Mus musculus<400> 350Gly Tyr Pro Ala25 1<210> 351<211> 1230 <212> PRT<213> Mus musculus<400> 35135 Ser Arg Ser Ile Val His Ser Ser Gly Asn Thr Tyr1 5 10<210> 35240 <211> 3<212> PRT<213> Mus musculus<400> 352

45

Lys Val Ser1

-293-

<210> 353<211> 5<212> PRT<213> Mus musculus

5

<400> 353Asp Ser His Val Pro1 5

10

<210> 354<211> 115<212> PRT15 <213> Mus musculus<400> 354Gln Ile Gln Leu Val Gln Ser Gly Pro Glu Leu Lys Lys Pro Gly Glu20 1 5 10 15Thr Val Lys Ile Ser Cys Lys Ala Ser Gly Tyr Pro Phe Ser Glu Tyr20 25 30

25

Ser Ile His Trp Val Lys Gln Ala Pro Gly Lys Gly Leu Lys Trp Met35 40 45

30

Val Tyr Val Asn Thr Glu Thr Gly Gln Pro Ile Val Gly Asp Asp Phe50 55 6035 Arg Gly Arg Phe Val Leu Ser Leu Glu Thr Ser Ala Ser Thr Ala Tyr65 70 75 80Leu Gln Ile Asn Asn Leu Lys Asn Glu Asp Thr Ala Thr Tyr Phe Cys40 85 90 95Gly Gly Gly Gly Tyr Pro Ala Tyr Trp Gly Gln Gly Thr Leu Val Thr100 105 110

45

Val Ser Ala115

-294-

<210> 355<211> 112<212> PRT5 <213> Mus musculus<400> 355

10 Asp Val Leu Met Thr Gln Thr Pro Leu Ser Leu Pro Val Ser Leu Gly
1 5 10 15

15 Asp Gln Ala Ser Ile Ser Cys Arg Ser Ser Arg Ser Ile Val His Ser
20 25 30

20 Ser Gly Asn Thr Tyr Leu Glu Trp Tyr Leu Gln Lys Pro Gly Gln Ser
35 40 45

20 Pro Lys Leu Leu Ile Tyr Lys Val Ser Asn Arg Phe Ser Gly Val Pro
50 55 60

25 Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Asn Ile
65 70 75 80

30 Ser Arg Val Glu Ala Glu Asp Leu Gly Val Tyr Tyr Cys Phe Gln Asp
85 90 95

35 Ser His Val Pro Phe Thr Phe Gly Thr Gly Thr Lys Leu Glu Ile Lys
100 105 110

<210> 356<211> 7<212> PRT40 <213> Mus musculus<400> 356

45 Gly Phe Thr Phe Ser Ser Ser
1 5

<210> 357<211> 3

-295-

<212> PRT<213> Mus musculus<400> 357

5

Ala Thr Gly110 <210> 358<211> 8<212> PRT<213> Mus musculus15 <400> 358Tyr Pro His Tyr Tyr Ala Met Asp1 5

20

<210> 359<211> 7<212> PRT<213> Mus musculus

25

<400> 359Ser Glu Asn Ile Phe Ser Asn1 5

30

<210> 360<211> 3<212> PRT35 <213> Mus musculus<400> 360Ser Ala Thr40 1<210> 361<211> 645 <212> PRT<213> Mus musculus<400> 361

-296-

Phe Tyr Lys Ile Pro Phe1 55 <210> 362<211> 121<212> PRT<213> Mus musculus10 <400> 362Gln Gly Gln Met His Gln Ser Gly Ala Glu Leu Val Lys Pro Gly Ser1 5 10 15

15

Ser Val Lys Leu Ser Cys Lys Thr Ser Gly Phe Thr Phe Ser Ser Ser20 25 3020 Phe Ile Ser Trp Leu Lys Gln Lys Pro Gly Gln Ser Leu Glu Trp Ile35 40 45Ala Trp Ile Tyr Ala Ala Thr Gly Ser Thr Ser Tyr Asn Gln Lys Phe25 50 55 60Thr Asn Lys Ala Gln Leu Thr Val Asp Thr Ser Ser Ser Ala Ala Tyr65 70 75 80

30

Met Gln Phe Ser Ser Leu Thr Thr Glu Asp Ser Ala Ile Tyr Tyr Cys85 90 95

35

Ala Arg His Ala Gly Tyr Pro His Tyr Tyr Ala Met Asp Tyr Trp Gly100 105 11040 Gln Gly Thr Ser Val Thr Val Ser Ser115 120<210> 36345 <211> 107<212> PRT<213> Mus musculus<400> 363

-297-

Asp Ile Gln Met Thr Gln Ser Pro Ala Ser Leu Ser Ala Ser Val Gly
1 5 10 15

5

Glu Thr Val Thr Ile Thr Cys Arg Ala Ser Glu Asn Ile Phe Ser Asn
20 25 30

10

Leu Ala Trp Tyr Gln Gln Lys Gln Gly Lys Ser Pro Gln Leu Leu Val
35 40 45

15

Tyr Ser Ala Thr Asn Leu Gly Asp Gly Val Pro Ser Arg Phe Ser Gly
50 55 60

20

Ser Gly Ser Gly Thr Gln Phe Ser Leu Lys Ile Asn Ser Leu Gln Pro
65 70 75 80

25

Glu Asp Phe Gly Asn Tyr Tyr Cys Gln His Phe Tyr Lys Ile Pro Phe
85 90 95

30

Thr Phe Gly Thr Gly Thr Lys Leu Glu Ile Lys
100 105

35

<210> 364
<211> 7
<212> PRT
<213> Mus musculus

40

<400> 364
Gly Phe Asn Ile Lys Asp Tyr
1 5

45

<210> 365
<211> 3
<212> PRT
<213> Mus musculus
<400> 365
Glu Asp Gly
1

-298-

- <210> 366
<211> 8
5 <212> PRT
<213> Mus musculus

<400> 366
10 His Gly Tyr Val Gly Trp Phe Ala
1 5

<210> 367
15 <211> 8
<212> PRT
<213> Mus musculus

<400> 367
20 Ala Ser Glu Asn Val Asp Thr Tyr
1 5

<210> 368
25 <211> 3
<212> PRT
<213> Mus musculus

30 <400> 368

Gly Ala Ser
1

35 <210> 369
<211> 6
<212> PRT
<213> Mus musculus
40 <400> 369

Ser Tyr Ser Tyr Pro Trp
1 5
45 <210> 370
<211> 119
<212> PRT

-299-

<213> Mus musculus<400> 370

5 Glu Val Gln Leu Gln Gln Ser Gly Ala Glu Pro Leu Lys Pro Gly Ala
1 5 10 15

10 Ser Val Lys Leu Thr Cys Thr Thr Ser Gly Phe Asn Ile Lys Asp Tyr
20 25 30

15 Tyr Ile His Trp Val Lys Gln Arg Ser Asp Gln Gly Leu Glu Trp Ile
35 40 45

Gly Arg Ile Asp Pro Glu Asp Gly Glu Leu Ile Tyr Ala Pro Lys Phe
50 55 60

20 Gln Asp Lys Ala Thr Ile Thr Val Asp Thr Ser Ser Asn Ile Ala Tyr
65 70 75 80

25 Leu Gln Leu Asn Ser Leu Thr Ser Glu Asp Thr Ala Val Tyr Tyr Cys
85 90 95

30 Ser Arg Asp His Gly Tyr Val Gly Trp Phe Ala Tyr Trp Gly Gln Gly
100 105 110

35 Thr Leu Val Thr Val Ser Ala
115

<210> 371<211> 107<212> PRT40 <213> Mus musculus<400> 371

45 Asn Val Val Met Thr Gln Ser Pro Lys Ser Met Ile Met Ser Val Gly
1 5 10 15

Gln Arg Val Thr Leu Asn Cys Lys Ala Ser Glu Asn Val Asp Thr Tyr
20 25 30

-300-

5 Val Ser Trp Tyr Gln Gln Lys Pro Glu Gln Ser Pro Lys Leu Leu Ile
 35 40 45

10 Tyr Gly Ala Ser Asn Arg Tyr Thr Gly Val Pro Asp Arg Phe Thr Gly
 50 55 60

15 Ser Arg Ser Ala Thr Asp Phe Thr Leu Thr Ile Ser Ser Val Gln Ala
 65 70 75 80

20 Glu Asp Leu Ala Val Tyr Tyr Cys Gly Gln Ser Tyr Ser Tyr Pro Trp
 85 90 95

25 Thr Phe Gly Gly Gly Thr Lys Leu Glu Phe Arg
 100 105

30 <210> 372
 <211> 219
 <212> PRT
 <213> Artificial

35 <220>
 <223> an exemplary Pseudomonas exotoxin A variant 1(deimmunized PE24
 example)

40 <400> 372

45 Pro Thr Gly Ala Glu Phe Leu Gly Asp Gly Gly Asp Val Ser Phe Ser
 1 5 10 15

50 Thr Arg Gly Thr Gln Asn Trp Thr Val Glu Arg Leu Leu Gln Ala His
 20 25 30

55 Ala Gln Leu Glu Glu Arg Gly Tyr Val Phe Val Gly Tyr His Gly Thr
 35 40 45

60 Phe Leu Glu Ala Ala Gln Ser Ile Val Phe Gly Gly Val Ala Ala Arg
 50 55 60

-301-

Ser Gln Asp Leu Ala Ala Ile Trp Ala Gly Phe Tyr Ile Ala Gly Asp
65 70 75 80

5 Pro Ala Leu Ala Tyr Gly Tyr Ala Gln Asp Gln Glu Pro Asp Ala Ala
 85 90 95

10 Gly Arg Ile Arg Asn Gly Ala Leu Leu Arg Val Tyr Val Pro Ala Ser
 100 105 110

15 Ser Leu Pro Gly Phe Tyr Arg Thr Ser Leu Thr Leu Ala Ala Pro Glu
 115 120 125

20 Ala Ala Gly Glu Val Glu Arg Leu Ile Gly His Pro Leu Pro Leu Ala
 130 135 140

25 Leu Asp Ala Ile Thr Gly Pro Glu Glu Glu Gly Gly Arg Leu Glu Thr
 145 150 155 160

30 Ile Leu Gly Trp Pro Leu Ala Glu Arg Thr Val Val Ile Pro Ser Ala
 165 170 175

35 Ile Pro Thr Asp Pro Arg Asn Val Gly Gly Asp Leu Asp Pro Ser Ser
 180 185 190

40 Ile Pro Asp Lys Glu Gln Ala Ile Ser Ala Leu Pro Asp Tyr Ala Ser
 195 200 205

45 Gln Pro Gly Lys Pro Pro Arg Glu Asp Leu Lys
 210 215

<210> 373
<211> 219
<212> PRT
<213> Artificial

<220>
<223> an exemplary Pseudomonas exotoxin A variant 2(deimmunized PE24
 example)

-302-

<400> 373

Pro Thr Gly Ala Glu Phe Leu Gly Asp Gly Gly Asp Val Ser Phe Ser
 1 5 10 15

5

Thr Arg Gly Thr Gln Asn Trp Thr Val Glu Arg Leu Leu Gln Ala His
 20 25 30

10

Ala Gln Leu Glu Glu Arg Gly Tyr Val Phe Val Gly Tyr His Gly Thr
 35 40 45

15

Ala Leu Glu Ala Ala Gln Ser Ile Val Phe Gly Gly Val Arg Ala Arg
 50 55 60

20

Ser Gln Asp Leu Arg Ala Ile Trp Arg Gly Phe Tyr Ile Ala Gly Asp
 65 70 75 80

25

Pro Ala His Ala Tyr Gly Tyr Ala Gln Asp Gln Glu Pro Asp Ala Arg
 85 90 95

30

Gly Arg Ile Ala Asn Gly Ala Leu Leu Arg Val Tyr Val Pro Ala Ser
 100 105 110

Ser Leu Pro Gly Phe Tyr Arg Thr Ser Leu Thr Leu Ala Ala Pro Glu
 115 120 125

35

Ala Ala Gly Glu Val Glu Arg Leu Ile Gly His Pro Leu Pro Leu Arg
 130 135 140

40

Leu Asp Ala Ile Thr Gly Pro Glu Glu Glu Gly Gly Arg Glu Glu Thr
 145 150 155 160

45

Ile Leu Gly Trp Pro Leu Ala Glu Arg Thr Val Val Ile Pro Ser Ala
 165 170 175

Ile Pro Thr Asp Pro Arg Asn Val Gly Gly Asp Leu Asp Pro Ser Ser
 180 185 190

-303-

- Ile Pro Asp Lys Glu Gln Ala Ile Ser Ala Leu Pro Asp Tyr Ala Ser
195 200 205
- 5 Gln Pro Gly Lys Pro Pro Arg Glu Asp Leu Lys
210 215
- 10 <210> 374
<211> 107
<212> PRT
<213> Homo Sapiens
- 15 <400> 374
- Arg Thr Val Ala Ala Pro Ser Val Phe Ile Phe Pro Pro Ser Asp Glu
1 5 10 15
- 20 Gln Leu Lys Ser Gly Thr Ala Ser Val Val Cys Leu Leu Asn Asn Phe
20 25 30
- 25 Tyr Pro Arg Glu Ala Lys Val Gln Trp Lys Val Asp Asn Ala Leu Gln
35 40 45
- 30 Ser Gly Asn Ser Gln Glu Ser Val Thr Glu Gln Asp Ser Lys Asp Ser
50 55 60
- 35 Thr Tyr Ser Leu Ser Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu
65 70 75 80
- Lys His Lys Val Tyr Ala Cys Glu Val Thr His Gln Gly Leu Ser Ser
85 90 95
- 40 Pro Val Thr Lys Ser Phe Asn Arg Gly Glu Cys
100 105
- 45 <210> 375
<211> 105
<212> PRT
<213> Homo Sapiens

-304-

<400> 375

Gln Pro Lys Ala Ala Pro Ser Val Thr Leu Phe Pro Pro Ser Ser Glu
1 5 10 15

5

Glu Leu Gln Ala Asn Lys Ala Thr Leu Val Cys Leu Ile Ser Asp Phe
20 25 30

10

Tyr Pro Gly Ala Val Thr Val Ala Trp Lys Ala Asp Ser Ser Pro Val
35 40 45

15

Lys Ala Gly Val Glu Thr Thr Thr Pro Ser Lys Gln Ser Asn Asn Lys
50 55 60

20

Tyr Ala Ala Ser Ser Tyr Leu Ser Leu Thr Pro Glu Gln Trp Lys Ser
65 70 75 80

25

His Arg Ser Tyr Ser Cys Gln Val Thr His Glu Gly Ser Thr Val Glu
85 90 95

30

Lys Thr Val Ala Pro Thr Glu Cys Ser
100 105

<210> 376

<211> 330

<212> PRT

<213> Homo Sapiens

35

<400> 376

Ala Ser Thr Lys Gly Pro Ser Val Phe Pro Leu Ala Pro Ser Ser Lys
1 5 10 15

40

Ser Thr Ser Gly Gly Thr Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr
20 25 30

45

Phe Pro Glu Pro Val Thr Val Ser Trp Asn Ser Gly Ala Leu Thr Ser
35 40 45

-305-

	<u>Gly Val His Thr Phe Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser</u>
	<u>50 55 60</u>
5	<u>Leu Ser Ser Val Val Thr Val Pro Ser Ser Ser Leu Gly Thr Gln Thr</u>
	<u>65 70 75 80</u>
10	<u>Tyr Ile Cys Asn Val Asn His Lys Pro Ser Asn Thr Lys Val Asp Lys</u>
	<u>85 90 95</u>
15	<u>Lys Val Glu Pro Lys Ser Cys Asp Lys Thr His Thr Cys Pro Pro Cys</u>
	<u>100 105 110</u>
20	<u>Pro Ala Pro Glu Leu Leu Gly Gly Pro Ser Val Phe Leu Phe Pro Pro</u>
	<u>115 120 125</u>
	<u>Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys</u>
	<u>130 135 140</u>
25	<u>Val Val Val Asp Val Ser His Glu Asp Pro Glu Val Lys Phe Asn Trp</u>
	<u>145 150 155 160</u>
30	<u>Tyr Val Asp Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu</u>
	<u>165 170 175</u>
35	<u>Glu Gln Tyr Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu</u>
	<u>180 185 190</u>
40	<u>His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn</u>
	<u>195 200 205</u>
	<u>Lys Ala Leu Pro Ala Pro Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly</u>
	<u>210 215 220</u>
45	<u>Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Asp Glu</u>
	<u>225 230 235 240</u>
	<u>Leu Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr</u>

-306-

245 250 255
 Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn
 5 260 265 270
 Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe
 10 275 280 285
 Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn
 15 290 295 300
 Val Phe Ser Cys Ser Val Met His Glu Ala Leu His Asn His Tyr Thr
 305 310 315 320
 Gln Lys Ser Leu Ser Leu Ser Pro Gly Lys
 20 325 330
 <210> 377
 25 <211> 330
 <212> PRT
 <213> homo sapiens
 <400> 377
 30 Ala Ser Thr Lys Gly Pro Ser Val Phe Pro Leu Ala Pro Ser Ser Lys
 1 5 10 15
 Ser Thr Ser Gly Gly Thr Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr
 35 20 25 30
 Phe Pro Glu Pro Val Thr Val Ser Trp Asn Ser Gly Ala Leu Thr Ser
 40 35 40 45
 Gly Val His Thr Phe Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser
 45 50 55 60
 Leu Ser Ser Val Val Thr Val Pro Ser Ser Ser Leu Gly Thr Gln Thr
 65 70 75 80

-307-

	<u>Tyr Ile Cys Asn Val Asn His Lys Pro Ser Asn Thr Lys Val Asp Lys</u>
	<u>85 90 95</u>
5	<u>Lys Val Glu Pro Lys Ser Cys Asp Lys Thr His Thr Cys Pro Pro Cys</u>
	<u>100 105 110</u>
10	<u>Pro Ala Pro Glu Ala Ala Gly Gly Pro Ser Val Phe Leu Phe Pro Pro</u>
	<u>115 120 125</u>
15	<u>Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys</u>
	<u>130 135 140</u>
20	<u>Val Val Val Asp Val Ser His Glu Asp Pro Glu Val Lys Phe Asn Trp</u>
	<u>145 150 155 160</u>
25	<u>Tyr Val Asp Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu</u>
	<u>165 170 175</u>
30	<u>Glu Gln Tyr Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu</u>
	<u>180 185 190</u>
35	<u>His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn</u>
	<u>195 200 205</u>
40	<u>Lys Ala Leu Pro Ala Pro Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly</u>
	<u>210 215 220</u>
45	<u>Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Asp Glu</u>
	<u>225 230 235 240</u>
50	<u>Leu Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr</u>
	<u>245 250 255</u>
55	<u>Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn</u>
	<u>260 265 270</u>

-308-

Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe
275 280 285

5 Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn
290 295 300

10 Val Phe Ser Cys Ser Val Met His Glu Ala Leu His Asn His Tyr Thr
305 310 315 320

15 Gln Lys Ser Leu Ser Leu Ser Pro Gly Lys
325 330

<210> 378
<211> 330
<212> PRT
 20 <213> homo sapiens

<400> 378

25 Ala Ser Thr Lys Gly Pro Ser Val Phe Pro Leu Ala Pro Ser Ser Lys
1 5 10 15

30 Ser Thr Ser Gly Gly Thr Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr
20 25 30

35 Phe Pro Glu Pro Val Thr Val Ser Trp Asn Ser Gly Ala Leu Thr Ser
35 40 45

40 Gly Val His Thr Phe Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser
50 55 60

45 Leu Ser Ser Val Val Thr Val Pro Ser Ser Ser Leu Gly Thr Gln Thr
65 70 75 80

45 Tyr Ile Cys Asn Val Asn His Lys Pro Ser Asn Thr Lys Val Asp Lys
85 90 95

Lys Val Glu Pro Lys Ser Cys Asp Lys Thr His Thr Cys Pro Pro Cys
100 105 110

-309-

5	<u>Pro Ala Pro Glu Ala Ala Gly Gly Pro Ser Val Phe Leu Phe Pro Pro</u> <u>115 120 125</u>
10	<u>Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys</u> <u>130 135 140</u>
15	<u>Val Val Val Asp Val Ser His Glu Asp Pro Glu Val Lys Phe Asn Trp</u> <u>145 150 155 160</u>
20	<u>Tyr Val Asp Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu</u> <u>165 170 175</u>
25	<u>Glu Gln Tyr Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu</u> <u>180 185 190</u>
30	<u>His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn</u> <u>195 200 205</u>
35	<u>Lys Ala Leu Gly Ala Pro Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly</u> <u>210 215 220</u>
40	<u>Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Asp Glu</u> <u>225 230 235 240</u>
45	<u>Leu Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr</u> <u>245 250 255</u>
	<u>Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn</u> <u>260 265 270</u>
	<u>Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe</u> <u>275 280 285</u>
	<u>Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn</u> <u>290 295 300</u>

-310-

- Val Phe Ser Cys Ser Val Met His Glu Ala Leu His Asn His Tyr Thr
305 310 315 320
- 5 Gln Lys Ser Leu Ser Leu Ser Pro Gly Lys
 325 330
- 10 <210> 379
<211> 327
<212> PRT
<213> Homo Sapiens
- 15 <400> 379
- Ala Ser Thr Lys Gly Pro Ser Val Phe Pro Leu Ala Pro Cys Ser Arg
1 5 10 15
- 20 Ser Thr Ser Glu Ser Thr Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr
 20 25 30
- 25 Phe Pro Glu Pro Val Thr Val Ser Trp Asn Ser Gly Ala Leu Thr Ser
 35 40 45
- 30 Gly Val His Thr Phe Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser
 50 55 60
- 35 Leu Ser Ser Val Val Thr Val Pro Ser Ser Ser Leu Gly Thr Lys Thr
65 70 75 80
- Tyr Thr Cys Asn Val Asp His Lys Pro Ser Asn Thr Lys Val Asp Lys
 85 90 95
- 40 Arg Val Glu Ser Lys Tyr Gly Pro Pro Cys Pro Ser Cys Pro Ala Pro
 100 105 110
- 45 Glu Phe Leu Gly Gly Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys
 115 120 125
- Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val Val

-311-

	<u>130</u>	<u>135</u>	<u>140</u>
5	<u>Asp Val Ser Gln Glu Asp Pro Glu Val Gln Phe Asn Trp Tyr Val Asp</u>		
	<u>145</u>	<u>150</u>	<u>155</u>
			<u>160</u>
10	<u>Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Phe</u>		
	<u>165</u>	<u>170</u>	<u>175</u>
15	<u>Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu His Gln Asp</u>		
	<u>180</u>	<u>185</u>	<u>190</u>
	<u>Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Gly Leu</u>		
	<u>195</u>	<u>200</u>	<u>205</u>
20	<u>Pro Ser Ser Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg</u>		
	<u>210</u>	<u>215</u>	<u>220</u>
25	<u>Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Gln Glu Glu Met Thr Lys</u>		
	<u>225</u>	<u>230</u>	<u>235</u>
			<u>240</u>
30	<u>Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp</u>		
	<u>245</u>	<u>250</u>	<u>255</u>
35	<u>Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys</u>		
	<u>260</u>	<u>265</u>	<u>270</u>
	<u>Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser</u>		
	<u>275</u>	<u>280</u>	<u>285</u>
40	<u>Arg Leu Thr Val Asp Lys Ser Arg Trp Gln Glu Gly Asn Val Phe Ser</u>		
	<u>290</u>	<u>295</u>	<u>300</u>
45	<u>Cys Ser Val Met His Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser</u>		
	<u>305</u>	<u>310</u>	<u>315</u>
			<u>320</u>
	<u>Leu Ser Leu Ser Leu Gly Lys</u>		
	<u>325</u>		

-312-

<210> 380<211> 2805 <212> PRT<213> homo sapiens<400> 380

10 Ser Glu Val Glu Tyr Arg Ala Glu Val Gly Gln Asn Ala Tyr Leu Pro
1 5 10 15

15 Cys Phe Tyr Thr Pro Ala Ala Pro Gly Asn Leu Val Pro Val Cys Trp
20 25 30

20 Gly Lys Gly Ala Cys Pro Val Phe Glu Cys Gly Asn Val Val Leu Arg
35 40 45

Thr Asp Glu Arg Asp Val Asn Tyr Trp Thr Ser Arg Tyr Trp Leu Asn
50 55 60

25 Gly Asp Phe Arg Lys Gly Asp Val Ser Leu Thr Ile Glu Asn Val Thr
65 70 75 80

30 Leu Ala Asp Ser Gly Ile Tyr Cys Cys Arg Ile Gln Ile Pro Gly Ile
85 90 95

35 Met Asn Asp Glu Lys Phe Asn Leu Lys Leu Val Ile Lys Pro Ala Lys
100 105 110

40 Val Thr Pro Ala Pro Thr Arg Gln Arg Asp Phe Thr Ala Ala Phe Pro
115 120 125

Arg Met Leu Thr Thr Arg Gly His Gly Pro Ala Glu Thr Gln Thr Leu
130 135 140

45 Gly Ser Leu Pro Asp Ile Asn Leu Thr Gln Ile Ser Thr Leu Ala Asn
145 150 155 160

-313-

	<u>Glu Leu Arg Asp Ser Arg Leu Ala Asn Asp Leu Arg Asp Ser Gly Ala</u>
	<u>165 170 175</u>
5	<u>Thr Ile Arg Ile Gly Ile Tyr Ile Gly Ala Gly Ile Cys Ala Gly Leu</u>
	<u>180 185 190</u>
10	<u>Ala Leu Ala Leu Ile Phe Gly Ala Leu Ile Phe Lys Trp Tyr Ser His</u>
	<u>195 200 205</u>
15	<u>Ser Lys Glu Lys Ile Gln Asn Leu Ser Leu Ile Ser Leu Ala Asn Leu</u>
	<u>210 215 220</u>
20	<u>Pro Pro Ser Gly Leu Ala Asn Ala Val Ala Glu Gly Ile Arg Ser Glu</u>
	<u>225 230 235 240</u>
25	<u>Glu Asn Ile Tyr Thr Ile Glu Glu Asn Val Tyr Glu Val Glu Glu Pro</u>
	<u>245 250 255</u>
30	<u>Asn Glu Tyr Tyr Cys Tyr Val Ser Ser Arg Gln Gln Pro Ser Gln Pro</u>
	<u>260 265 270</u>
35	<u>Leu Gly Cys Arg Phe Ala Met Pro</u>
	<u>275 280</u>
	<u><210> 381</u>
	<u><211> 181</u>
	<u><212> PRT</u>
	<u><213> homo sapiens</u>
	<u><400> 381</u>
40	<u>Ser Glu Val Glu Tyr Arg Ala Glu Val Gly Gln Asn Ala Tyr Leu Pro</u>
	<u>1 5 10 15</u>
45	<u>Cys Phe Tyr Thr Pro Ala Ala Pro Gly Asn Leu Val Pro Val Cys Trp</u>
	<u>20 25 30</u>
	<u>Gly Lys Gly Ala Cys Pro Val Phe Glu Cys Gly Asn Val Val Leu Arg</u>
	<u>35 40 45</u>

-314-

5	<u>Thr Asp Glu Arg Asp Val Asn Tyr Trp Thr Ser Arg Tyr Trp Leu Asn</u>
	<u>50 55 60</u>
10	<u>Gly Asp Phe Arg Lys Gly Asp Val Ser Leu Thr Ile Glu Asn Val Thr</u>
	<u>65 70 75 80</u>
15	<u>Leu Ala Asp Ser Gly Ile Tyr Cys Cys Arg Ile Gln Ile Pro Gly Ile</u>
	<u>85 90 95</u>
20	<u>Met Asn Asp Glu Lys Phe Asn Leu Lys Leu Val Ile Lys Pro Ala Lys</u>
	<u>100 105 110</u>
25	<u>Val Thr Pro Ala Pro Thr Arg Gln Arg Asp Phe Thr Ala Ala Phe Pro</u>
	<u>115 120 125</u>
30	<u>Arg Met Leu Thr Thr Arg Gly His Gly Pro Ala Glu Thr Gln Thr Leu</u>
	<u>130 135 140</u>
35	<u>Gly Ser Leu Pro Asp Ile Asn Leu Thr Gln Ile Ser Thr Leu Ala Asn</u>
	<u>145 150 155 160</u>
40	<u>Glu Leu Arg Asp Ser Arg Leu Ala Asn Asp Leu Arg Asp Ser Gly Ala</u>
	<u>165 170 175</u>
45	<u>Thr Ile Arg Ile Gly</u>
	<u>180</u>

* * *

Although the foregoing invention has been described in some detail by way of illustration and example for purposes of clarity of understanding, the descriptions and examples should not be construed as limiting the scope of the invention. The disclosures of all patent and scientific literature cited herein are expressly incorporated in their entirety by reference.

Claims

1. Use of a T cell activating bispecific antigen binding molecule and a PD-1 axis binding antagonist in the manufacture of a medicament for combination therapy in the treatment of or delaying the progression of a cancer, wherein the T cell activating bispecific antigen binding molecule comprises a first Fab fragment capable of specific binding to CD3, a second Fab fragment capable of specific binding to Folate Receptor 1 (FolR1), wherein the PD-1 axis binding antagonist inhibits the binding of PD-1 to its ligand binding partners, wherein the PD-1 axis binding antagonist is selected from the group consisting of an anti-PD-1 antibody, an anti-PD-L1 antibody and an anti-PD-L2 antibody, and wherein the cancer is a FolR1-positive cancer.
2. The use of claim 1, wherein the first Fab fragment comprises the heavy chain CDR1 SEQ ID NO: 37, the heavy chain CDR2 of SEQ ID NO: 38, the heavy chain CDR3 of SEQ ID NO: 39, the light chain CDR1 of SEQ ID NO: 32, the light chain CDR2 of SEQ ID NO: 33, and the light chain CDR3 of SEQ ID NO: 34.
3. The use of claim 1, wherein the first Fab fragment comprises a variable heavy chain comprising an amino acid sequence of SEQ ID NO: 36 and a variable light chain comprising an amino acid sequence of SEQ ID NO: 31.
4. The use of any one of claims 1-3, wherein the T cell activating bispecific antigen binding molecule further comprises a third Fab fragment capable of specific binding to FolR1.
5. The use of claim 4, wherein the second and third Fab fragment capable of specific binding to FolR1 comprise identical heavy chain complementarity determining region (CDR) and light chain CDR sequences.
6. The use of claim 5, wherein the third Fab fragment is identical to the second Fab fragment.

7. The use of any one of the preceding claims, wherein the Fab fragment capable of specific binding to Folate Receptor 1 (FolR1) comprises at least one heavy chain complementarity determining region (CDR) amino acid sequence selected from the group consisting of SEQ ID NO: 16, SEQ ID NO: 17 and SEQ ID NO: 18 and at least one light chain CDR selected from the group of SEQ ID NO: 32, SEQ ID NO: 33, SEQ ID NO: 34.
8. The use of claim 7, wherein the Fab fragment capable of specific binding to Folate Receptor 1 (FolR1) comprises a variable heavy chain comprising an amino acid sequence of SEQ ID NO: 15 and a variable light chain comprising an amino acid sequence of SEQ ID NO: 31.
9. The use of any one of claims 1-6, wherein the Fab fragment capable of specific binding to Folate Receptor 1 (FolR1) comprises at least one heavy chain complementarity determining region (CDR) amino acid sequence selected from the group consisting of SEQ ID NO: 8, SEQ ID NO: 56 and SEQ ID NO: 57 and at least one light chain CDR selected from the group of SEQ ID NO: 59, SEQ ID NO: 60, SEQ ID NO: 65.
10. The use of claim 9, wherein the Fab fragment capable of specific binding to Folate Receptor 1 (FolR1) comprises a variable heavy chain comprising an amino acid sequence of SEQ ID NO: 55 and a variable light chain comprising an amino acid sequence of SEQ ID NO: 64.
11. The use of any one of claims 1-6, wherein the Fab fragment capable of specific binding to Folate Receptor 1 (FolR1) comprises at least one heavy chain complementarity determining region (CDR) amino acid sequence selected from the group consisting of SEQ ID NO: 16, SEQ ID NO: 402 and SEQ ID NO: 400 and at least one light chain CDR selected from the group of SEQ ID NO: 32, SEQ ID NO: 33, SEQ ID NO: 34.

12. The use of claim 11, wherein the Fab fragment capable of specific binding to FolR1 comprises:
 - a) a complementarity determining region heavy chain 1 (CDR-H1) amino acid sequences of SEQ ID NO: 16;
 - (b) a CDR-H2 amino acid sequence of SEQ ID NO: 402;
 - (c) a CDR-H3 amino acid sequence of SEQ ID NO: 400;
 - (d) a complementarity determining region light chain 1 (CDR-L1) amino acid sequence of SEQ ID NO: 32;
 - (e) a CDR-L2 amino acid sequence of SEQ ID NO: 33, and
 - (f) a CDR-L3 amino acid sequence of SEQ ID NO: 34.
13. The use of claim 12, wherein the Fab fragment capable of specific binding to FolR1 comprises a variable heavy chain comprising an amino acid sequence of SEQ ID NO: 401 and a variable light chain comprising an amino acid sequence of SEQ ID NO: 31.
14. The use of any one of the preceding claims, wherein the anti-PD-L1 antibody comprises a heavy chain comprising HVR-H1 sequence of SEQ ID NO:289, HVR-H2 sequence of SEQ ID NO:290, and HVR-H3 sequence of SEQ ID NO:291; and a light chain comprising HVR-L1 sequence of SEQ ID NO: 292, HVR-L2 sequence of SEQ ID NO: 293, and HVR-L3 sequence of SEQ ID NO: 294.
15. The use of any one of the preceding claims, wherein the antibody comprises a heavy chain variable region comprising the amino acid sequence of SEQ ID NO:280 or SEQ ID NO:281 and a light chain variable region comprising the amino acid sequence of SEQ ID NO:383.
16. The use of any one of Claims 1-15, wherein the medicament further comprises a T cell immunoglobulin mucin 3 (TIM3) antagonist.
17. The use of Claim 16, wherein the TIM3 antagonist is an anti-TIM3 antibody.

18. The use of Claim 17, wherein the anti-TIM3 antibody is F38-2E2.
19. The use of any one of Claims 1-18, wherein the cancer is selected from the group consisting of ovarian cancer, lung cancer, breast cancer, renal cancer, colorectal cancer, endometrial cancer.
20. Use of a T cell activating bispecific antigen binding molecule specific for Folate Receptor 1 (FolR1) and CD3, and a PD-1 axis binding antagonist in the manufacture of a medicament for combination therapy in enhancing immune function in an individual having a FolR1-positive cancer, wherein the PD-1 axis binding antagonist inhibits the binding of PD-1 to its ligand binding partners, and wherein the PD-1 axis binding antagonist is selected from the group consisting of an anti-PD-1 antibody, an anti-PD-L1 antibody and an anti-PD-L2 antibody.
21. A method for selecting a patient for treatment with a combination of a T cell activating bispecific antigen binding molecule specific for Folate Receptor 1 (FolR1) and CD3, and a PD-1 axis binding antagonist comprising measuring the level of PD-1 expression, wherein a patient having less than about 15% PD-1^{hi} expressing T cells is selected for treatment with the combination, wherein the PD-1 axis binding antagonist inhibits the binding of PD-1 to its ligand binding partners, and wherein the PD-1 axis binding antagonist is selected from the group consisting of an anti-PD-1 antibody, an anti-PD-L1 antibody and an anti-PD-L2 antibody.
22. A kit comprising a T cell activating bispecific antigen binding molecule specific for Folate Receptor 1 (FolR1) and CD3 and a PD-1 axis binding antagonist as a combined pharmaceutical formulation for use in treating or delaying progression of cancer in an individual, and a package insert comprising instructions for using the T cell activating bispecific antigen binding molecule and the PD-1 axis binding antagonist to treat or delay progression of cancer in an individual, wherein the cancer is a FolR1-positive cancer and wherein the PD-1 axis binding antagonist inhibits the binding of PD-1 to its ligand binding partners, wherein the PD-1 axis binding antagonist is selected from

the group consisting of an anti-PD-1 antibody, an anti-PD-L1 antibody and an anti-PD-L2 antibody.

23. The kit of claim 22, further comprising a TIM3 antagonist.
24. A pharmaceutical composition comprising a T cell activating bispecific antigen binding molecule specific for Folate Receptor 1 (FolR1) and CD3, a PD-1 axis binding antagonist and a pharmaceutically acceptable carrier, wherein the PD-1 axis binding antagonist inhibits the binding of PD-1 to its ligand binding partners, and wherein the PD-1 axis binding antagonist is selected from the group consisting of an anti-PD-1 antibody, an anti-PD-L1 antibody and an anti-PD-L2 antibody.
25. The pharmaceutical composition of claim 24, further comprising a TIM3 antagonist.
26. Use of a T cell activating bispecific antigen binding molecule specific for Folate Receptor 1 (FolR1) and CD3, a PD-1 axis binding antagonist, and a TIM3 antagonist in the manufacture of a medicament for combination therapy in the treatment of cancer, wherein the cancer is a FolR1-positive cancer and wherein the PD-1 axis binding antagonist inhibits the binding of PD-1 to its ligand binding partners, wherein the PD-1 axis binding antagonist is selected from the group consisting of an anti-PD-1 antibody, an anti-PD-L1 antibody and an anti-PD-L2 antibody.
27. The use of claim 26, wherein the medicament is for treatment of ovarian cancer, lung cancer, breast cancer, renal cancer, colorectal cancer, endometrial cancer.