

Average Daily Stool frequency
Trial 1
Trial 2
Neg Con
Frequency (Number of Episodes / Day)
0
2
4
6
8
10
12
1
2
3
4
5
Days

FIG. 1



Average Daily Stool Consistency
Trial 1
Trial 2
Neg Con
Consistency (1 = Normal, 5 = Fully Liquid)
0
1
2
3
4
5
6
1
2
3
4
5
Days

FIG. 2



Average Doctor Assessed Well-Being
Trial 1
Trial 2
Neg Con
Wellbeing (1 = Normal, 5 = Most Sick)
0
1
2
3
4
5
6
1
2
3
4
5
Days

FIG. 3

FIG. 4




Stool Frequency (count)
11.00
10.00
9.00
8.00
7.00
6.00
5.00
4.00
3.00
2.00
1.00
1
2
3
4
5
Trial 1
Trial 2
Trial 3
Negative Controls

FIG. 5


Stool Consistency
5.00
4.50
4.00
3.50
3.00
2.50
2.00
1.50
1.00
1
2
3
4
5
Trial 1
Trial 2
Trial 3
Negative Controls



Doctor Reported Well Being
5.00
4.50
4.00
3.50
3.00
2.50
2.00
1.50
1.00
1
2
3
4
5
Trial 1
Trial 2
Trial 3
Negative Controls

FIG. 6

FIG. 7



Stool Frequency (count)
13.00
11.00
9.00
7.00
5.00
3.00
1.00
1
2
3
4
5
Trial 1
Trial 2
Trial 3 ES204A
Trial 3 ES204B
Trial 3 MT204A D
Trial 3 MT304A
Trial 3 MS204A
Trial 3 MS304A
Negative Controls

FIG. 8




Stool Consistency
5.00
4.50
4.00
3.50
3.00
2.50
2.00
1.50
1.00
1
2
3
4
5
Trial 1
Trial 2
Trial 3 ES204A
Trial 3 ES204B
Trial 3 MT204A D
Trial 3 MT304A
Trial 3 MS204A
Trial 3 MS304A
Negative Controls

FIG. 9


Doctor Reported Well Being
5.00
4.50
4.00
3.50
3.00
2.50
2.00
1.50
1.00
1
2
3
4
5
Trial 1
Trial 2
Trial 3 ES204A
Trial 3 ES204B
Trial 3 MT204A D
Trial 3 MT304A
Trial 3 MS204A
Trial 3 MS304A
Negative Controls