

TITLE OF THE INVENTION

OPTIMIZED EXPRESSION OF HPV 45 L1 IN YEAST

FIELD OF THE INVENTION

5 The present invention relates generally to the therapy of human papillomavirus (HPV). More specifically, the present invention relates to synthetic polynucleotides encoding HPV45 L1 protein, and to recombinant vectors and hosts comprising said polynucleotides. This invention also relates to HPV45 virus-like particles (VLPs), wherein the VLPs are produced by expressing recombinant HPV 45 L1 or L1 + L2 in yeast cells and to their use in vaccines and pharmaceutical compositions for preventing and treating HPV.

BACKGROUND OF THE INVENTION

 There are more than 80 types of human papillomavirus (HPV), many of which have been associated with a wide variety of biological phenotypes, from benign proliferative warts to malignant carcinomas (for review, *see* McMurray et al., *Int. J. Exp. Pathol.* 82(1): 15-33 (2001)). HPV6 and HPV11 are the types most commonly associated with benign warts and/or nonmalignant condyloma acuminata of the genital or respiratory mucosa. HPV16 and HPV18 are the high-risk types most frequently associated with *in situ* and invasive carcinomas of the cervix, vagina, vulva and anal canal. More than 90% of cervical carcinomas are associated with infections of HPV16, HPV18 or the less prevalent oncogenic types HPV31, -33, -45, -52 and -58 (Schiffman et al., *J. Natl. Cancer Inst.* 85(12): 958-64 (1993)). The observation that HPV DNA is detected in more than 90% of cervical cancers provides strong epidemiological evidence that HPVs cause cervical carcinoma (*see* Bosch et al., *J. Natl. Cancer Inst.* 87(11): 796-802 (1995)).

 Papillomaviruses are small (50-60 nm), nonenveloped, icosahedral DNA viruses that encode up to eight early and two late genes. The open reading frames (ORFs) of the viral genomes are designated E1 to E7, and L1 and L2, where "E" denotes early and "L" denotes late. L1 and L2 code for virus capsid proteins, while the E genes are associated with functions such as viral replication and cellular transformation.

 The L1 protein is the major capsid protein and has a molecular weight of 55-60 kDa. The L2 protein is the minor capsid protein. Immunological data suggest that most of the L2 protein is internal to the L1 protein in the viral capsid. Both the L1 and L2 proteins are highly conserved among different papillomaviruses.

 Expression of the L1 protein or a combination of the L1 and L2 proteins in yeast, insect cells, mammalian cells or bacteria leads to self-assembly of virus-like particles (VLPs) (for review, *see* Schiller and Roden, in *Papillomavirus Reviews: Current Research on Papillomaviruses*; Lacey, ed.

Leeds, UK: Leeds Medical Information, pp 101-12 (1996)). VLPs are morphologically similar to authentic virions and are capable of inducing high titres of neutralizing antibodies upon administration into an animal. Because VLPs do not contain the potentially oncogenic viral genome, they present a safe alternative to the use of live virus in HPV vaccine development (for review, *see* Schiller and Hidesheim, *J. Clin. Virol* 19: 67-74 (2000)). For this reason, the L1 and L2 genes have been identified as immunological targets for the development of prophylactic and therapeutic vaccines for HPV infection and disease.

HPV vaccine development and commercialization have been hindered by difficulties associated with obtaining high expression levels of exogenous genes in successfully transformed host organisms, limiting the production of purified protein. Therefore, despite the identification of wild-type nucleotide sequences encoding HPV L1 proteins such as HPV45 L1 protein, it would be highly desirable to develop a readily renewable source of crude HPV L1 protein that utilizes HPV45 L1-encoding nucleotide sequences that are optimized for expression in the intended host cell. Additionally, it would be useful to produce large quantities of HPV45 L1 VLPs having the immunity-conferring properties of the native proteins for use in vaccine development.

SUMMARY OF THE INVENTION

According to a first aspect of the present invention, there is provided a nucleic acid molecule comprising a sequence of nucleotides that encodes an HPV45 L1 protein as set forth in SEQ ID NO:2, the nucleic acid sequence being codon-optimized for high-level expression in a yeast cell.

According to a second aspect of the present invention a vector comprising the nucleic acid molecule as defined above.

According to a third aspect of the present invention, there is provided a host cell comprising the vector as defined above.

According to a fourth aspect of the present invention, there is provided virus-like particles (VLPs) comprised of recombinant L1 protein or recombinant L1 + L2 proteins of

HPV45, wherein the recombinant L1 protein or the recombinant L1 + L2 proteins are produced in yeast.

According to a fifth aspect of the present invention, there is provided a method of producing the VLPs as defined above comprising:

- (a) transforming yeast with a codon-optimized DNA molecule encoding HPV45 L1 protein or HPV45 L1 + L2 proteins;
- (b) cultivating the transformed yeast under conditions that permit expression of the codon-optimized DNA molecule to produce a recombinant papillomavirus protein; and
- (c) isolating the recombinant papillomavirus protein to produce the VLPs as defined above.

According to a sixth aspect of the present invention, there is provided a vaccine comprising the VLPs as defined above.

According to a seventh aspect of the present invention, there is provided pharmaceutical compositions comprising the VLPs as defined above.

According to an eighth aspect of the present invention, there is provided the use of vaccine as defined above in the preparation of a medicament for preventing HPV infection.

According to a ninth aspect of the present invention, there is provided the use of VLPs as defined above in the preparation of a medicament for inducing an immune response in an animal.

Disclosed herein are compositions and methods to elicit or enhance immunity to the protein products expressed by HPV45 L1 genes, which have been associated with cervical cancer. Specifically, the present invention provides polynucleotides encoding HPV45 L1 protein, wherein the polynucleotides have been codon-optimized for high level expression in a yeast cell. Also disclosed herein are HPV45 virus-like particles (VLPs), wherein said VLPs are produced by expressing recombinant HPV45 L1 or L1 + L2 in yeast cells and discloses use of HPV45 VLPs in pharmaceutical compositions and vaccines for the prevention and/or treatment of HPV-associated cancer.

Disclosed herein are synthetic DNA molecules encoding the HPV45 L1 protein. The codons of the synthetic molecules are designed so as to use the codons preferred by a yeast cell. The synthetic molecules may be used as a source of HPV45 L1 protein, which may self-assemble into VLPs. Said VLPs may be used

in a VLP-based vaccine.

An exemplary embodiment comprises a synthetic nucleic acid molecule which encodes the HPV45 L1 protein as set forth in SEQ ID NO:2, said nucleic acid molecule comprising a sequence of nucleotides as set forth in SEQ ID NO: 1.

Also disclosed are recombinant vectors and recombinant host cells, both prokaryotic and eukaryotic, which may contain the nucleic acid molecules disclosed throughout this specification.

Also disclosed herein is a process for expressing an HPV45 L1 protein in a recombinant host cell, comprising: (a) introducing a vector comprising a nucleic acid encoding an

HPV45 L1 protein into a yeast host cell; and (b) culturing the yeast host cell under conditions which allow expression of said HPV45 L1 protein.

Further disclosed herein is a process for expressing an HPV45 L1 protein in a recombinant host cell, comprising: (a) introducing a vector comprising a nucleic acid encoding an HPV45 L1 protein into a yeast host cell; wherein the nucleic acid molecule is codon-optimized for optimal expression in the yeast host cell and; (b) culturing the yeast host cell under conditions which allow expression of said HPV45 L1 protein.

In preferred embodiments, the nucleic acid comprises a sequence of nucleotides as set forth in SEQ ID NO: 1 (45 L1 R sequence).

Also disclosed herein are HPV45 virus-like particles (VLPs) which are produced in yeast cells, methods of producing HPV45 VLPs, and methods of using HPV45 VLPs.

In a preferred embodiment of the invention, the yeast is selected from the group consisting of: *Saccharomyces cerevisiae*, *Hansenula polymorpha*, *Pichia pastoris*, *Kluyveromyces fragilis*, *Kluyveromyces lactis*, and *Schizosaccharomyces pombe*.

Disclosed herein is an HPV45 VLP, wherein the VLP is produced by recombinant expression of HPV45 L1 or HPV45 L1 + L2 in a yeast cell.

Disclosed herein is an HPV45 VLP which comprises an HPV45 L1 protein encoded by a codon-optimized HPV45 L1 gene. In an exemplary embodiment of this aspect of the invention, the codon-optimized HPV45 L1 gene comprises a sequence of nucleotides as set forth in SEQ ID NO:1.

Disclosed herein is a method for inducing an immune response in an animal comprising administering HPV45 virus-like particles to the animal. In a preferred embodiment, the HPV45 VLPs are produced by a codon-optimized gene.

Disclosed herein is a method of preventing or treating HPV- associated cervical cancer comprising administering to a mammal a vaccine comprising HPV45 VLPs. In a preferred embodiment of this aspect of the invention, the HPV45 VLPs are produced in yeast.

Disclosed herein is a vaccine comprising HPV45 virus-like particles (VLPs). In an alternative embodiment of this aspect of the invention, the

vaccine further comprises VLPs of at least one additional HPV type. The at least one additional HPV type may be any HPV type of interest, including any HPV type described in the art or those subsequently identified. In a preferred embodiment, the HPV type is a type that is associated with a clinical phenotype such as warts or cervical cancer. In a further preferred embodiment, the at least one additional HPV type is selected from the group consisting of: HPV6, HPV11, HPV16, HPV18, HPV31, HPV33, HPV35, HPV39, HPV51, HPV52, HPV55, HPV56, HPV58, HPV59 and HPV68.

Disclosed herein are pharmaceutical compositions comprising HPV 45 virus-like particles. Further, this invention relates to pharmaceutical compositions comprising HPV45 VLPs and VLPs of at least one additional HPV type. In a preferred embodiment, the at least one additional HPV type is selected from the group consisting of: HPV6, HPV11, HPV16, HPV18, HPV31, HPV33, HPV35, HPV39, HPV51, HPV52, HPV55, HPV56, HPV58, HPV59, and HPV68.

As used throughout the specification and in the appended claims, the singular forms "a," "an," and "the" include the plural reference unless the context clearly dictates otherwise.

As used throughout the specification and appended claims, the following definitions and abbreviations apply:

The term "promoter" refers to a recognition site on a DNA strand to which the RNA polymerase binds. The promoter forms an initiation complex with RNA polymerase to initiate and drive transcriptional activity. The complex can be modified by activating sequences termed "enhancers" or "upstream activating sequences" or inhibiting sequences termed "silencers".

The term "vector" refers to some means by which DNA fragments can be introduced into a host organism or host tissue. There are various types of vectors including plasmids, viruses (including adenovirus), bacteriophages and cosmids.

The designation "45 L1 wild-type sequence" refers to the HPV45 L1 DNA sequence disclosed herein as SEQ ID NO:3 (45 L1 wt). Although the HPV 45 L1 wild-type DNA sequence has been described previously, it is not uncommon to find minor sequence variations between DNAs obtained from clinical isolates. Therefore, a representative 45 L1 wild-type DNA sequence was isolated from clinical samples previously shown to contain HPV 45 DNA (see EXAMPLE 1). The 45 L1 wild-type sequence was used as a reference sequence to compare the codon-optimized 45 L1 sequences disclosed herein (see FIGURE 1).

The designation "HPV 45 L1 R" or "45 L1 R" refers to an exemplary synthetic HPV45 L1 nucleotide sequence (SEQ ID NO: 1), disclosed herein, wherein the sequence was rebuilt so that it comprises codons that are preferred for high-level expression by a yeast cell.

The term "effective amount" means sufficient vaccine composition is introduced to produce the adequate levels of the polypeptide, so that an immune response results. One skilled in the art recognizes that this level may vary.

A "conservative amino acid substitution" refers to the replacement of one amino acid residue by another, chemically similar, amino acid residue. Examples of such conservative substitutions are: substitution of one hydrophobic residue (isoleucine, leucine, valine, or methionine) for another; substitution of one polar residue for another polar residue of the same charge (e.g., arginine for lysine; glutamic acid for aspartic acid).

The term "mammalian" refers to any mammal, including a human being.

"VLP" or "VLPs" mean(s) virus-like particle or virus-like particles.

"Synthetic" means that the HPV45 L1 gene was created so that it contains a sequence of nucleotides that is not the same as the sequence of nucleotides present in the designated naturally occurring wild-type HPV45 L1 gene (45 L1 wt, SEQ ID NO:3). As stated above, synthetic molecules are provided herein comprising a sequence of nucleotides comprising codons that are preferred for expression by yeast cells. The synthetic molecules provided herein encode the same amino acid sequence as the wild-type HPV45 L1 gene (SEQ ID NO:2).

10 BRIEF DESCRIPTION OF THE DRAWINGS

FIGURE 1 is a DNA sequence alignment showing nucleotides that were altered in the synthetic HPV45 L1 gene of the present invention (SEQ ID NO:1, indicated as "45 L1 R") (See EXAMPLE 2). The reference sequence is the 45 L1 wild-type sequence (SEQ ID NO:3, indicated as "45 L1 wt"; see EXAMPLE 1). Altered nucleotides are indicated at their corresponding location. Nucleotide number is contained within the parentheses. Identical nucleotides in the 45 L1 rebuilt sequence are indicated with dots.

FIGURE 2 shows the rebuilt synthetic HPV 45 L1 nucleotide and single-code amino acid sequence. Nucleotide number is indicated to the left.

FIGURE 3 shows a Northern blot probed specifically for HPV 45 L1 RNA under high stringency conditions (see EXAMPLE 4). Lane (1) contains 5 μ g of HPV 45 L1 R RNA and lane (2) contains 10 μ g of HPV 45 L1 R RNA. A single full-length RNA transcript is apparent on the blot. Arrow on left indicates the predicted position of a full length HPV 45 L1 transcript.

FIGURE 4 shows a Western Blot of HPV 45 L1 wt and three HPV 45 L1 R isolates. Contents of the lanes are: 45 L1 wt (lane 1), 45 L1 R #4 (lane 2), 45 L1 R #7 (lane 3), 45 L1 R #11 (lane 4), HPV 16 L1 control (lane 5). Fifteen micrograms of total yeast protein extract were loaded into each lane of a 10% SDS-PAGE gel. A goat polyclonal anti-serum against a TrpE-HPV 45 L1 fusion protein was used to specifically identify the HPV 45 L1 protein. The arrow at the left indicates the 55 kDa position.

FIGURE 5 shows a portion of the data from two ELISA experiments in ng VLP / μ g total protein (see EXAMPLE 7). A comparison between 45 L1 wt and two separate clones of 45 L1 R, is shown. Rebuilt HPV 45 L1 VLP expression was approximately 2 fold higher than the 45 L1 wt.

FIGURE 6 shows a representative sample of HPV 45 VLPs composed of HPV 45 L1 R protein molecules, described herein, as visualized by transmission electron microscopy (see EXAMPLE 8). The bar represents approximately 100 nm.

DETAILED DESCRIPTION OF THE INVENTION

The majority of cervical carcinomas are associated with infections of specific oncogenic types of human papillomavirus (HPV). The present invention relates to compositions and methods to elicit or enhance immunity to the protein products expressed by genes of oncogenic HPV types.

5 Specifically, the present invention provides polynucleotides encoding HPV45 L1 protein, which self-assemble into HPV45 virus-like particles (VLPs) and discloses use of said polynucleotides and VLPs in pharmaceutical compositions and vaccines for the prevention and/or treatment of HPV-associated cancer.

The wild-type HPV45 L1 nucleotide sequence has been reported (Genbank Accession # NC_001590; see also Delius and Hofmann, *Curr. Top. Microbiol. Immunol.* 186: 13-31 (1994)). The
10 present invention provides synthetic DNA molecules encoding the HPV45 L1 protein. The synthetic molecules of the present invention comprise a sequence of codons, wherein at least a portion of the codons have been altered to use the codons preferred by a yeast cell for high-level expression. The synthetic molecules may be used as a coding sequence for expression of HPV45 L1 protein, which may self-assemble into VLPs. Said VLPs may be used in a VLP-based vaccine to provide effective
15 immunoprophylaxis against papillomavirus infection through neutralizing antibody and cell-mediated immunity.

Expression of HPV VLPs in yeast cells offers the advantages of being cost-effective and easily adapted to large-scale growth in fermenters. However, many HPV L1 proteins, including HPV45 L1 are expressed at levels in yeast cells which are lower than what is desirable for commercial scale-up.

20 Accordingly, the present invention relates to HPV45 L1 gene sequences that are "optimized" for high level expression in a yeast cellular environment.

A "triplet" codon of four possible nucleotide bases can exist in over 60 variant forms. Because these codons provide the message for only 20 different amino acids (as well as transcription initiation and termination), some amino acids can be coded for by more than one codon, a phenomenon
25 known as codon redundancy. For reasons not completely understood, alternative codons are not uniformly present in the endogenous DNA of differing types of cells. Indeed, there appears to exist a variable natural hierarchy or "preference" for certain codons in certain types of cells. As one example, the amino acid leucine is specified by any of six DNA codons including CTA, CTC, CTG, CTT, TTA, and TTG. Exhaustive analysis of genome codon use frequencies for microorganisms has revealed
30 endogenous DNA of *E. coli* most commonly contains the CTG leucine-specifying codon, while the DNA of yeasts and slime molds most commonly includes a TTA leucine-specifying codon. In view of this hierarchy, it is generally believed that the likelihood of obtaining high levels of expression of a leucine-rich polypeptide by an *E. coli* host will depend to some extent on the frequency of codon use. For example, it is likely that a gene rich in TTA codons will be poorly expressed in *E. coli*, whereas a CTG

rich gene will probably be highly expressed in this host. Similarly, a preferred codon for expression of a leucine-rich polypeptide in yeast host cells would be TTA.

The implications of codon preference phenomena on recombinant DNA techniques are manifest, and the phenomenon may serve to explain many prior failures to achieve high expression levels of exogenous genes in successfully transformed host organisms—a less "preferred" codon may be repeatedly present in the inserted gene and the host cell machinery for expression may not operate as efficiently. This phenomenon suggests that synthetic genes which have been designed to include a projected host cell's preferred codons provide an optimal form of foreign genetic material for practice of recombinant protein expression. Thus, one aspect of this invention is an HPV45 L1 gene that is codon-optimized for high-level expression in a yeast cell. In a preferred embodiment of this invention, it has been found that the use of alternative codons encoding the same protein sequence may remove the constraints on expression of HPV45 L1 proteins by yeast cells.

In accordance with this invention, HPV45 L1 gene segments were converted to sequences having identical translated sequences but with alternative codon usage as described by Sharp and Cowe (Synonymous Codon Usage in *Saccharomyces cerevisiae*. *Yeast* 7: 657-678 (1991)), which is hereby incorporated by reference. The methodology generally consists of identifying codons in the wild-type sequence that are not commonly associated with highly expressed yeast genes and replacing them with optimal codons for high expression in yeast cells. The new gene sequence is then inspected for undesired sequences generated by these codon replacements (e.g., "ATTTA" sequences, inadvertent creation of intron splice recognition sites, unwanted restriction enzyme sites, high GC content, presence of transcription termination signals that are recognized by yeast, etc.). Undesirable sequences are eliminated by substitution of the existing codons with different codons coding for the same amino acid. The synthetic gene segments are then tested for improved expression.

The methods described above were used to create synthetic gene segments for HPV45 L1, resulting in a gene comprising codons optimized for high level expression in a yeast cellular environment. While the above procedure provides a summary of our methodology for designing codon-optimized genes for use in HPV vaccines, it is understood by one skilled in the art that similar vaccine efficacy or increased expression of genes may be achieved by minor variations in the procedure or by minor variations in the sequence.

Accordingly, the present invention relates to a synthetic polynucleotide comprising a sequence of nucleotides encoding an HPV45 L1 protein, or a biologically active fragment or mutant form of an HPV45 L1 protein, the polynucleotide sequence comprising codons optimized for expression in a yeast host cell. Said mutant forms of the HPV45 L1 protein include, but are not limited to: conservative amino acid substitutions; amino-terminal truncations, carboxy-terminal truncations, deletions, or additions. Any such biologically active fragment and/or mutant will encode either a protein or protein

fragment which at least substantially mimics the immunological properties of the HPV45 L1 protein as set forth in SEQ ID NO:2. The synthetic polynucleotides of the present invention encode mRNA molecules that express a functional HPV45 L1 protein so as to be useful in the development of a therapeutic or prophylactic HPV vaccine.

5 One aspect of this invention is a codon-optimized nucleic acid molecule which encodes the HPV45 L1 protein as set forth in SEQ ID NO:2, said nucleic acid molecule comprising a sequence of nucleotides as set forth in SEQ ID NO:1.

The present invention also relates to recombinant vectors and recombinant host cells, both prokaryotic and eukaryotic, which contain the nucleic acid molecules disclosed throughout this
10 specification.

The synthetic HPV45 DNA or fragments thereof constructed through the methods described herein may be recombinantly expressed by molecular cloning into an expression vector containing a suitable promoter and other appropriate transcription regulatory elements, and transferred into prokaryotic or eukaryotic host cells to produce recombinant HPV45 L1. Techniques for such
15 manipulations are fully described by Sambrook et al. (*Molecular Cloning: A Laboratory Manual*; Cold Spring Harbor Laboratory, Cold Spring Harbor, New York, (1989); *Current Protocols in Molecular Biology*, Ausubel et al., Green Pub. Associates and Wiley-Interscience, New York (1988); *Yeast Genetics: A Laboratory Course Manual*, Rose et al., Cold Spring Harbor Laboratory, Cold Spring Harbor, New York, (1990)), which are hereby incorporated by reference in their entirety.

20 Thus, the present invention relates to a process for expressing an HPV45 L1 protein in a recombinant host cell, comprising: (a) introducing a vector comprising a nucleic acid encoding an HPV45 L1 protein into a yeast host cell; and (b) culturing the yeast host cell under conditions which allow expression of said HPV45 L1 protein.

The present invention further relates to a process for expressing an HPV45 L1 protein in
25 a recombinant host cell, comprising: (a) introducing a vector comprising a nucleic acid encoding an HPV45 L1 protein into a yeast host cell; wherein the nucleic acid molecule is codon-optimized for optimal expression in the yeast host cell and; (b) culturing the yeast host cell under conditions which allow expression of said HPV45 L1 protein.

This invention further relates to a process for expressing an HPV45 L1 protein in a
30 recombinant host cell, comprising: (a) introducing a vector comprising a nucleic acid as set forth in SEQ ID NO:1 into a yeast host cell; and, (b) culturing the yeast host cell under conditions which allow expression of said HPV45 L1 protein.

The synthetic genes of the present invention can be assembled into an expression
35 cassette that comprises sequences designed to provide efficient expression of the HPV45 L1 protein in the host cell. The cassette preferably contains the synthetic gene, with related transcriptional and

translations control sequences operatively linked to it, such as a promoter, and termination sequences. In a preferred embodiment, the promoter is the *S.cerevisiae GAL1* promoter, although those skilled in the art will recognize that any of a number of other known yeast promoters such as the *GAL10*, *GAL7*, *ADHI*, *TDH3*, or *PGK* promoters, or other eukaryotic gene promoters may be used. A preferred transcriptional terminator is the *S.cerevisiae ADHI* terminator, although other known transcriptional terminators may also be used. The combination of *GAL1* promoter – *ADHI* terminator is particularly preferred.

Another aspect of this invention is an HPV45 virus-like particle (VLP) produced by recombinantly expressing the HPV45 L1 or L1 + L2 genes in a yeast cell, methods of producing HPV45 VLPs, and methods of using HPV45 VLPs. VLPs can self-assemble when L1, the major capsid protein of human and animal papillomaviruses, is expressed in yeast, insect cells, mammalian cells or bacteria (for review, see Schiller and Roden, in *Papillomavirus Reviews: Current Research on Papillomaviruses*; Lacey, ed. Leeds, UK: Leeds Medical Information, pp 101-12 (1996)). Morphologically indistinct HPV VLPs can also be produced by expressing a combination of the L1 and L2 capsid proteins. VLPs are composed of 72 pentamers of L1 in a T=7 icosahedral structure (Baker et al., *Biophys. J.* 60(6): 1445-56 (1991)).

VLPs are morphologically similar to authentic virions and are capable of inducing high titres of neutralizing antibodies upon administration into an animal. Immunization of rabbits (Breitburd et al., *J. Virol.* 69(6): 3959-63 (1995)) and dogs (Suzich et al., *Proc. Natl. Acad. Sci. USA* 92(25): 11553-57 (1995)) with VLPs was shown to both induce neutralizing antibodies and protect against experimental papillomavirus infection. However, because the VLPs do not contain the potentially oncogenic viral genome and can self-assemble when expressed from a single gene, they present a safe alternative to the use of live virus in HPV vaccine development (for review, see Schiller and Hidesheim, *J. Clin. Virol.* 19: 67-74 (2000)).

Marais and colleagues (*J. Med. Virol.* 60: 331-336 (2000)) disclose HPV 45 VLPs, which were produced in insect cells from an HPV45 L1 gene expressed from baculovirus. Expression of VLPs in insect cells is not advantageous for vaccine development because of the high associated costs. Additionally, it is often difficult to scale-up expression of HPV L1 genes in insect cell culture to the large volumes necessary for commercial product development. HPV 45 VLPs produced in yeast cells have not been described. Expression of HPV VLPs in yeast cells offers the advantages of being cost-effective and easily adapted to large-scale growth in fermenters. In addition, the yeast genome can be readily altered to ensure selection of recombinant, transformed yeast with increased growth and expression potential.

Thus, the present invention relates to virus-like particles comprised of recombinant L1 protein or recombinant L1 + L2 proteins of HPV45, wherein the recombinant protein is expressed in a yeast cell.

5 In a preferred embodiment of the invention, the HPV45 VLPs are produced by expressing HPV 45 L1 or HPV 45 L1 + L2 in a yeast selected from the group consisting of: *Saccharomyces cerevisiae*, *Hansenula polymorpha*, *Pichia pastoris*, *Kluyveromyces fragilis*, *Kluyveromyces lactis*, and *Schizosaccharomyces pombe*.

10 Another aspect of this invention is an HPV45 VLP which comprises an HPV45 L1 protein produced by expressing a codon-optimized HPV45 L1 gene. In a preferred embodiment of this aspect of the invention, the codon-optimized HPV45 L1 gene comprises a sequence of nucleotides as set forth in SEQ ID NO:1.

Yet another aspect of this invention is a method of producing HPV45 VLPs, comprising: (a) transforming yeast with a recombinant DNA molecule encoding HPV45 L1 protein or HPV45 L1 + L2 proteins; (b) cultivating the transformed yeast under conditions that permit expression of the recombinant DNA molecule to produce the recombinant HPV45 protein; and (c) isolating the recombinant HPV45 protein to produce HPV45 VLPs.

20 In a preferred embodiment of this aspect of the invention, the yeast is transformed with a codon-optimized HPV45 L1 gene to produce HPV45 VLPs. In a particularly preferred embodiment, the codon-optimized HPV45 L1 gene comprises a sequence of nucleotides as set forth in SEQ ID NO:1.

This invention also provides a method for inducing an immune response in an animal comprising administering HPV45 virus-like particles to the animal. In a preferred embodiment, the HPV45 VLPs are produced by a codon-optimized gene.

Yet another aspect of this invention is a method of preventing or treating HPV-associated cervical cancer comprising administering to a mammal a vaccine comprising HPV45 VLPs. 25 In a preferred embodiment of this aspect of the invention, the HPV45 VLPs are produced in yeast.

This invention also relates to a vaccine comprising HPV45 virus-like particles (VLPs).

30 In an alternative embodiment of this aspect of the invention, the vaccine further comprises VLPs of at least one additional HPV type. In a preferred embodiment, the at least one additional HPV type is selected from the group consisting of: HPV6, HPV11, HPV16, HPV18, HPV31, HPV33, HPV35, HPV39, HPV51, HPV52, HPV55, HPV56, HPV58, HPV59, and HPV68.

In a preferred embodiment of this aspect of the invention, the vaccine further comprises HPV16 VLPs.

In another preferred embodiment of the invention, the vaccine further comprises HPV16 VLPs and HPV18 VLPs.

In yet another preferred embodiment of the invention, the vaccine further comprises HPV6 VLPs, HPV11 VLPs, HPV16 VLPs and HPV18 VLPs.

This invention also relates to pharmaceutical compositions comprising HPV 45 virus-like particles. Further, this invention relates to pharmaceutical compositions comprising HPV45 VLPs and VLPs of at least one additional HPV type. In a preferred embodiment, the at least one additional HPV type is selected from the group consisting of: HPV6, HPV11, HPV16, HPV18, HPV31, HPV33, HPV35, HPV39, HPV51, HPV52, HPV55, HPV56, HPV58, HPV59, and HPV68.

Vaccine compositions of the present invention may be used alone at appropriate dosages defined by routine testing in order to obtain optimal inhibition of HPV45 infection while minimizing any potential toxicity. In addition, co-administration or sequential administration of other agents may be desirable.

The amount of virus-like particles to be introduced into a vaccine recipient will depend on the immunogenicity of the expressed gene product. In general, an immunologically or prophylactically effective dose of about 10 μ g to 100 μ g, and preferably about 20 μ g to 60 μ g of VLPs is administered directly into muscle tissue. Subcutaneous injection, intradermal introduction, impression through the skin, and other modes of administration such as intraperitoneal, intravenous, or inhalation delivery are also contemplated. It is also contemplated that booster vaccinations may be provided. Parenteral administration, such as intravenous, intramuscular, subcutaneous or other means of administration with adjuvants such as alum or Merck aluminum adjuvant, concurrently with or subsequent to parenteral introduction of the vaccine of this invention is also advantageous.

All publications mentioned herein are incorporated by reference for the purpose of describing and disclosing methodologies and materials that might be used in connection with the present invention. Nothing herein is to be construed as an admission that the invention is not entitled to antedate such disclosure by virtue of prior invention.

Having described preferred embodiments of the invention with reference to the accompanying drawings, it is to be understood that the invention is not limited to those precise embodiments, and that various changes and modifications may be effected therein by one skilled in the art without departing from the scope or spirit of the invention as defined in the appended claims.

The following examples illustrate, but do not limit the invention.

EXAMPLE 1

Determination of a Representative HPV 45 L1 Sequence

The HPV 45 L1 sequence has been described previously (Genbank Accession # NC_001590; *see* Delius and Hofmann *Curr. Top. Microbiol. Immunol.* 186: 13-31 (1994)). It is not

uncommon, however, to find minor sequence variations between DNAs obtained from clinical isolates. To determine a representative HPV45 L1 wild-type sequence, DNA was isolated from four clinical samples previously shown to contain HPV 45 DNA. HPV 45 L1 sequences were amplified in a polymerase chain reaction (PCR) using *Taq* DNA polymerase and the following primers: HPV 45 L1 F 5' - CCACCAACCACTATATAGGTATTC - 3' (SEQ ID NO:4) and HPV 45 L1 B 5' - CAAACATACATATATGTGCTAACA - 3' (SEQ ID NO:5). The amplified products were electrophoresed on agarose gels and visualized by ethidium bromide staining. The ~ 1500 bp L1 bands were excised and DNA purified using the QIA quick PCR purification kit (Qiagen, Hilden, Germany). The DNA was then ligated to a TA cloning vector (Invitrogen Corp., Carlsbad, CA), *E. Coli* was transformed with the ligation mixture and plated on LB agar with ampicillin plus IPTG and X-gal for blue/white colony selection. The plates were inverted and incubated for 16 hours at 37°C.

Colony PCR was performed on eight white colonies originating from each of the four clinical isolates that had been PCR amplified. HPV 45 L1 DNA was PCR-amplified using primers HPV 45 L1 F and HPV 45 L1 B. The specific PCR protocol consisted of 25 cycles of 15 seconds at 98°C (denaturation), 30 seconds at 50°C (annealing) and 2 minutes at 68°C (extension). PCR products were visualized by ethidium bromide staining after agarose gel electrophoresis. Several colonies from each isolate contained L1 bands. The second colony from each isolate was cultured in LB medium with ampicillin, shaking at 37°C for 16 hours. Minipreps were performed to extract plasmid DNA from the colonies.

DNA sequencing was performed on the plasmids containing amplified and cloned HPV 45 L1 DNA from the four clinical isolates. DNA and translated amino acid sequences were compared with one another and the Genbank HPV 45 L1 sequences. Sequence analysis of the four clinical isolates revealed that no sequence was identical to the Genbank sequence. The HPV 45 L1 plasmid from isolate #33 was chosen to be the representative 45 L1 wild-type (wt) sequence and is referred to herein as the "45 L1 wild-type sequence" (SEQ ID NO:3, see FIGURE 1). This sequence contained a nine nucleotide, three amino acid, deletion which maintained the subsequent sequence reading frame, five point mutations resulting in five amino acids changes and eight additional silent point mutations. Amino acids 495-497 in the Genbank sequence are deleted in the 45 L1 wt. Genbank amino acids numbers 23 (S->N), 55 (N->S), 303 (I->T), 357 (S->N), and 356 (Q->H) represent the five amino acids changes (Genbank aa -> 45 L1 wt aa). The 8 silent mutations were distributed throughout the sequence. See FIGURE 1.

The 45 L1 wild-type sequence was amplified using the LS-112 5' - CTCAGATCTCACAAACAATAATGGCTTTGTGGCGGCCTAGTGAC - 3' (SEQ ID NO:6) and HPV 45 L1 EAS 5' - GACAGATCTTATTTTTTACTACGTATACGTACACG - 3' (SEQ ID NO:7) primers to add *Bgl*III extensions. PCR was performed using *Vent* polymerase. The PCR product was visualized by ethidium bromide staining of an agarose gel. The ~ 1500 bp band was

excised and DNA purified using the QIAEX II gel extraction kit (Qiagen). The PCR product was then digested with *Bgl*III at 37°C for 2 hours and purified using the QIA quick PCR purification kit. The *Bgl*III-digested HPV 45 L1 PCR product was ligated to *Bam*HI-digested pGAL110 (described in Hofmann et al., *Virology* 209: 506-18 (1995)) and *E.coli* DH5 was transformed with the ligation mixture.

5 Colonies were screened by PCR for the HPV 45 L1 insert in the correct orientation. Sequence and orientation were confirmed by DNA sequencing. The selected clone was named pGAL110-HPV 45 L1 #6. This clone was digested with *Pst*I, *Eco*RI and *Hind*III to determine the restriction fragment profile of the HPV 45 L1 gene within the pGAL110 vector. DNA fragments were electrophoresed on agarose gels. The resulting restriction fragment profile was visualized by ethidium
10 bromide staining and viewed with UV light.

Maxiprep DNA was prepared. *Saccharomyces cerevisiae* cells were made competent by spheroplasting with Glusulase and transformed with pGAL110-HPV 45 L1 #6. The yeast transformation mixture was plated in Leu(-) sorbitol top-agar onto Leu(-) sorbitol plates and incubated inverted for 5-7 days at 30°C. Colonies were picked and streaked for clonal isolation on Leu(-) sorbitol plates. Isolated
15 colonies were subsequently grown in 5 ml of 5 X Leu(-) Ade(-) sorbitol with 1.6% glucose and 4% galactose in rotating tube cultures at 30°C to induce L1 transcription and protein expression.

EXAMPLE 2

Yeast codon optimization

20 Yeast-preferred codons have been described (Sharp, Paul M and Cowe, Elizabeth 1991 Synonymous Codon Usage in *Saccharomyces cerevisiae* YEAST 7: 657-678). Expression of the HPV 45 L1 wt protein was detectable; however, to obtain the increased expression necessary for commercial product development, the HPV 45 L1 gene was rebuilt utilizing yeast-preferred codons. Said rebuilt sequence would provide increased HPV45 L1 expression, which would be a significant advance over the
25 wild-type for use in vaccine development. When the sequence was rebuilt utilizing yeast-optimized codon sequences, the nucleotide sequence of 45 L1 wt was altered at 392 positions to produce 45 L1 R (R = rebuild). The amino acid sequence, however, was not altered. The nucleotide (SEQ ID NO:1) and amino acid (SEQ ID NO:2) sequences of HPV 45 L1 R are shown in FIGURE 2.

The strategy employed for the gene rebuild was to design long overlapping sense and
30 antisense oligomers that span the gene, substituting nucleotides with yeast-preferred codon sequences while maintaining the original amino acid sequence. These oligomers were used in place of template DNA in a PCR reaction. Additional amplification primers were designed and used to amplify the rebuilt sequences from the template oligomers. *Pfu* DNA polymerase was used in the PCR of 25 cycles.

The optimal conditions for amplification were section-specific; however most employed
35 a program resembling 94°C for 5 minutes (denaturing) followed by 25 cycles of 95°C for 30 sec

Following rebuild, the 1533 nt band was gel purified, ligated to pCR4 Blunt (Invitrogen) and *E. coli* TOP10 cells were transformed with the ligation mixture. Colonies were grown in 4 ml LB with ampicillin and plasmid DNA was extracted by standard miniprep techniques. The plasmid DNA was sequenced to confirm the presence of the desired changes for the 45 L1 rebuild. The 45 L1 R (rebuild) was re-amplified from pCR4Blunt-45 L1 R to add *Bam*HI extensions to both ends plus a yeast 5'-non-translated leader sequence upstream of the ATG initiation codon. The amplified fragment was cloned as above and plasmid DNA was sequenced. The plasmid, pCR4 Blunt-45 L1 R (*Bam*), was digested with *Bam*HI and DNA fragments were electrophoresed on an agarose gel. The ~1550 bp 45 L1 R (*Bam*) fragment was gel purified, ligated to *Bam*HI-digested pGAL110 and transformed into TOP10F' *E. coli* (Invitrogen).

Colonies were screened by PCR for the HPV 45 L1 R insert in the correct orientation in pGAL110. Sequence and orientation were confirmed by DNA sequencing. Maxiprep plasmid DNA was prepared and used for the transformation of *S. cerevisiae* cells which had been made competent by spheroplasting. The transformed yeast were plated in Leu(-) sorbitol top-agar onto Leu(-) sorbitol plates, which were incubated inverted for 7 days. Colonies were streaked for clonal isolation on Leu(-) sorbitol plates. Isolated colonies were subsequently grown in 5 ml of 5 X Leu(-) Ade(-) sorbitol with 1.6% glucose and 4% galactose in rotating tube cultures at 30°C to induce L1 transcription and protein expression. After 48 hours, a culture volume equivalent to an OD₆₀₀ = 10 was pelleted, the supernatant was removed and the pellets were frozen and stored at -70°C.

RNA preparation

Cell pellets of transformed yeast induced to express HPV 45 L1 by galactose induction were thawed on ice, suspended in 0.8 ml of Trizol reagent (Life Technologies, Gibco BRL) and incubated at room temperature for 5 minutes. One fifth volume of chloroform was added to the vial. It was then shaken vigorously for 15 seconds to mix and incubated at room temperature for 3 minutes. After a 5 minute centrifugation at 13 k rpms, the upper phase was collected and transferred to a new vial. 0.4 ml isopropanol was added and incubated at room temperature for 10 minutes. Centrifugation to pellet the RNA was performed at 13 k rpms for 10 minutes. The supernatant was decanted, the RNA pellet washed with 75% EtOH and centrifugation repeated. The supernatant was decanted and the RNA pellet allowed to air-dry for 15 minutes followed by suspension in RNase-free water. Spectrophotometry

was performed to determine the concentration of RNA in the sample using the assumption that an A₂₆₀ reading of 1 = 40 µg/ml RNA when the A₂₆₀/A₂₈₀ is 1.7-2.0.

EXAMPLE 4

5 Northern Blot Analysis

A 1.1% agarose formaldehyde gel was cast. Five and ten micrograms of RNA was combined with denaturing buffer (final concentrations: 6% formaldehyde, 50% formamide and 0.1 x MOPS) and heated to 65°C for 10 minutes. A one-tenth volume of gel loading buffer was added and the sample loaded onto the gel. Electrophoresis was performed at 75 volts in 1 x MOPS buffer for ~ 3 hours.
10 The gel was washed for 60 minutes in 10 x SSC.

The RNA was transferred to a Hybond-N+ nylon membrane (Amersham Biosciences, Piscataway, NJ) by capillary action over 16 hours in 10 x SSC. The RNA was then fixed to the nylon membrane by cross-linking using the Stratagene UV-Stratalinker (Stratagene, La Jolla, CA) auto-crosslink function. After fixing, the nylon membrane was allowed to air dry.

15 The Roche DIG High Prime DNA Labeling and Detection Kit I (F. Hoffmann-La Roche Ltd., Basel, Switzerland) was used to label 45 L1 R DNA sequences with DIG to be used as a probe to detect 45 L1 R RNA on the Northern blot. The pre-hybridization, hybridization and immunological development using an anti-DIG alkaline phosphatase conjugated antibody were performed per the manufacturer's recommendations. Briefly, the blot was pre-hybridized at 37°C for 30 minutes with
20 gentle shaking. The probe was denatured by heating to 95°C for 5 minutes and quenching on ice. The probe was added to the hybridization solution and applied to the membrane for 4 hours at 44.6°C with gentle shaking. The hybridization solution was then removed and the blot washed twice for 5 minutes in 2 x SSC with 0.1% SDS at room temperature, followed by an additional wash at 65°C with 0.5 x SSC and 0.1% SDS. The blot was then blocked for 30 minutes and anti-DIG alkaline phosphatase conjugated
25 antibody applied at a 1:5000 dilution for 30 minutes. The blot was washed and the presence of probe bound RNA determined by NBT/BCIP substrate detection of the alkaline phosphatase conjugated anti-DIG bound antibody.

Initial analysis of yeast expressing 45 L1 wt suggested that there was functional HPV 45 L1 transcription and translation; however, the sequence was rebuilt with yeast-preferred codon sequences
30 to obtain an increased level of expression, useful for vaccine development. The rebuilt 45 L1 sequence was engineered to omit any possible premature transcription termination sites to ensure robust transcription. Northern blot analysis of the 45 L1 R transcript revealed that full-length transcripts were generated (FIGURE 3).

35 EXAMPLE 5

HPV 45 L1 Protein Expression

Frozen yeast cell pellets of galactose-induced cultures equivalent to OD₆₀₀= 10, were thawed on ice and suspended in 300 µl of PC buffer (100 mM Na₂HPO₄ and 0.5 M NaCl, pH 7.0) with 2mM PMSF. Acid-washed 0.5mm glass beads, were added, ~ 0.5g/tube. The tubes were vortexed for 3 cycles of 5 minutes at 4°C with a 1 minute break. 7.5 µl of 20% TritonX100 was added and vortex repeated for 5 minutes at 4°C. The tubes were placed on ice for 15 minutes, and then centrifuged for 10 minutes at 4°C. The supernate was transferred to a sterile microfuge tube, labeled as total yeast protein extract, dated and stored at -70°C.

EXAMPLE 6

Western Blot Analysis

Total yeast protein extract from twenty isolated yeast colonies for each 45 L1 construct were analyzed by Western blot to confirm expression of 45 L1 protein after galactose induction.

Ten micrograms of total yeast protein extract was combined with SDS-PAGE loading buffer and heated to 95°C for 10 minutes. The proteins were loaded onto a 10% SDS-PAGE gel and electrophoresed in Tris-Glycine buffer. After protein separation, the proteins were Western transferred from the gel to nitrocellulose and the blot blocked in 1 x diluent buffer (Kirkegaard and Perry Laboratories) for 1 hour at room temperature with rocking. The blot was washed three times and then incubated with a 1:2500 dilution of goat anti-trpE-HPV 45 L1 serum at room temperature for 16 hours. The blot was then washed three times and incubated with a 1:2500 dilution of anti-goat-HRP conjugated antibody for 1 hr. The blot was again washed three times and NBT/BCIP detection substrate applied (Kirkegaard and Perry Laboratories). Immunoreactive proteins were detected as purple bands on the blot.

The results demonstrate that in all cases, the 45 L1 protein was detected as a distinct immunoreactive band on the nitrocellulose corresponding to approximately 57 kDa (FIGURE 4). The 16 L1 protein, which is approximately 55 kDa, was included as a positive control, along with HPV L1-free total yeast protein extract as a negative control (data not shown).

EXAMPLE 7

ELISA Assay

The yeast cells expressing HPV 45 L1 were grown by a variety of methods including rotating tube cultures, shake flasks and fermenters. The yeast were lysed and protein extracts were made to determine the amount of HPV 45 L1 virus-like particles (VLPs) produced per microgram of total protein. A sandwich ELISA was designed to demonstrate HPV 45 L1 VLP expression.

The H18R.5 monoclonal antibody (mAb), which recognizes both HPV type 18 and 45 L1 VLPs, was used to bind the 45 L1 VLPs found in the yeast protein extracts. The unbound proteins were washed away and H45.10B11.A5, an HPV 45 L1 VLP type and conformational specific mAb, was applied as a detection antibody. True, conformationally correct, 45 L1 VLPs were bound and detection was facilitated by the use of an anti-mouse IgG1 HRP-conjugated antibody and TMB substrate.

Specifically, H18R.5 was used to coat the bottom of Immulon 4 HBX 96 well plates overnight at 4°C. The plates were washed three times with PBS and 0.05% Tween 20, followed by blocking with blocking solution (PBS + 0.05% Tween 20 + 1% BSA). The plates were washed three times and antigens (total yeast cell lysates diluted in blocking solution to 100 µg/ml), were applied to row A in duplicate. Reference standards of purified HPV 45 L1 VLPs were applied to row A columns 1 and 2 at 3300ng/ml in 100 µg/ml total yeast protein. The reference and test samples were then serially diluted two-fold down each column. After three hours at room temperature the excess antigen was removed by aspiration and the plates washed 3 times. Cell supernatant containing HPV 45 L1 VLP conformational specific mAb H45.10B11.A5 was diluted 1:80 in blocking solution and applied to each well for an hour at room temperature. The plates were washed three times and an anti-mouse IgG1 HRP-conjugated antibody diluted 1:8000 in blocking solution was applied for 1 hour at room temperature. The plates were washed and TMB (Pierce) applied for 5 minutes to detect HRP-conjugated antibody complexes. The detection reaction was stopped by the addition of 2M H₂SO₄. Plates were read at 450 nm wavelength and the concentration of HPV 45 L1 VLP was determined by comparison to the reference standards in ng VLP / µg total protein.

A comparison between 45 L1 wt and two separate clones of 45 L1 R, all assayed in duplicate, is shown in FIGURE 5. Rebuilt HPV 45 L1 VLP expression was approximately 2 fold higher than the results observed for 45 L1 wt (FIGURE 6).

EXAMPLE 8

Transmission Electron Microscopy

To demonstrate that the 45 L1 protein was in fact self-assembling to form pentameric-L1 capsomers, which in turn self-assemble into virus-like particles, a partially purified 45 L1 R protein extract was analyzed by transmission EM.

Yeast cells were grown under small scale fermentation conditions, pelleted and the pellets were subjected to purification treatments. Pellet and clarified yeast extracts were analyzed by immunoblot to demonstrate L1 protein expression and retention through the purification procedure. Clarified yeast extracts were then subjected to centrifugation over a 45%-sucrose cushion and the resulting pellet suspended in buffer for analysis by transmission electron microscopy (EM).

A representative sample of the 45 L1 R VLPs produced is shown in FIGURE 6. The diameter of the spherical particles in this crude sample ranged from between 40 and 60 nm with some particles displaying a regular array of capsomers.

WHAT IS CLAIMED IS:

1. A nucleic acid molecule comprising a sequence of nucleotides that encodes an HPV45 L1 protein as set forth in SEQ ID NO:2, the nucleic acid sequence being codon-optimized for high-level expression in a yeast cell.
2. A vector comprising the nucleic acid molecule of claim 1.
3. A host cell comprising the vector of claim 3.
4. The host cell of claim 3, wherein the host cell is selected from the group consisting of: *Saccharomyces cerevisiae*, *Hansenula polymorpha*, *Pichia pastoris*, *Kluyveromyces fragilis*, *Kluyveromyces lactis*, and *Schizosaccharomyces pombe*.
5. The host cell of claim 4, wherein the host cell is *Saccharomyces cerevisiae*.
6. The nucleic acid molecule of claim 1, wherein the sequence of nucleotides comprises a sequence of nucleotides as set forth in SEQ ID NO:1.
7. A vector comprising the nucleic acid molecule of claim 6.
8. A host cell comprising the vector of claim 7.
9. Virus-like particles (VLPs) comprised of recombinant L1 protein or recombinant L1 + L2 proteins of HPV45, wherein the recombinant L1 protein or the recombinant L1 + L2 proteins are produced in yeast.
10. The VLPs of claim 9, wherein the recombinant L1 protein or recombinant L1 + L2 proteins are encoded by a codon-optimized HPV45 L1 nucleic acid molecule.
11. The VLPs of claim 10, wherein the codon-optimized nucleic acid molecule consists essentially of a sequence of nucleotides as set forth in SEQ ID NO:1.
12. A method of producing the VLPs of Claim 10, comprising:

- (a) transforming yeast with a codon-optimized DNA molecule encoding HPV45 L1 protein or HPV45 L1 + L2 proteins;
- (b) cultivating the transformed yeast under conditions that permit expression of the codon-optimized DNA molecule to produce a recombinant papillomavirus protein; and
- (c) isolating the recombinant papillomavirus protein to produce the VLPs of Claim 10.

- 13. A vaccine comprising the VLPs of Claim 10.
- 14. Pharmaceutical compositions comprising the VLPs of claim 10.
- 15. Use of the vaccine of Claim 13 in the preparation of a medicament for preventing HPV infection.
- 16. Use of VLPs of Claim 10 in the preparation of a medicament for inducing an immune response in an animal.
- 17. The virus-like particles of Claim 10 wherein the yeast is selected from the group consisting of *Saccharomyces cerevisiae*, *Hansenula polymorpha*, *Pichia pastoris*, *Kluyveromyces fragilis*, *Kluyveromyces lactis*, and *Schizosaccharomyces pombe*.
- 18. The virus-like particles of claim 17, wherein the yeast is *Saccharomyces cerevisiae*.
- 19. The vaccine of claim 13, further comprising VLPs of at least one additional HPV type.
- 20. The vaccine of claim 19, wherein the at least one additional HPV type is selected from the group consisting of: HPV6, HPV11, HPV16, HPV18, HPV31, HPV33, HPV35, HPV39, HPV51, HPV52, HPV55, HPV56, HPV58, HPV59, and HPV68.
- 21. The vaccine of claim 20, wherein the at least one HPV type comprises HPV16.
- 22. The vaccine of claim 21, further comprising HPV18 VLPs.

23. The vaccine of claim 22, further comprising HPV6 VLPs and HPV11 VLPs.
24. The vaccine of claim 23, further comprising HPV31 VLPs.
25. The vaccine of claim 22, further comprising HPV31 VLPs.

5

WO 2005/032586

PCT/US2004/031326

SEQUENCE LISTING

<110> Merck & Co., Inc.
 Bryan, Janine T.
 Brownlow, Michelle K.
 Schultz, Loren D.
 Jansen, Kathrin U.

<120> OPTIMIZED EXPRESSION OF HPV 45 L1 IN
 YEAST

<130> 21500-PCT

<150> 60/506,812

<151> 2003-09-29

<160> 8

<170> FastSEQ for Windows Version 4.0

<210> 1

<211> 1533

<212> DNA

<213> Artificial Sequence

<220>

<223> HPV45 L1R

<400> 1

```

atggccttgg ggagaccatc tgactctact gtctacttgc caccaccatc tgtcgctaga 60
gtcgtcaaca ctgacgacta cgtctccaga acctccatct tctaccacgc tggttcttcc 120
agattgttga ctgtcggtaa cccatacttc agagtcgtcc catccgggtgc tggtaacaag 180
caagctgttc caaagggtctc tgcttaccaa tacagagtct tcagagtcgc tttgccagac 240
ccaaacaagt tcgggttggc agactctact atctacaacc cagaaactca aagattgggtc 300
tgggcatgcg tcgggtatgga aatcggtaga ggtaaccat tgggtatcgg tttgtctgggt 360
caccattctt acaacaagtt ggacgacacc gaatccgctc acgctgctac tgcgtgctac 420
actcaagacg tcagagacaa cgtctctgtc gactacaagc aaacccaatt gtgtatcttg 480
ggttgtgtcc cagctatcgg tgaacactgg gctaagggtg ccttgtgtaa gccagctcaa 540
ttgcaaccag gtgactgtcc accattggaa ttgaagaaca ctatcatcga agacggtgac 600
atgggttgaca ctggttacgg tgctatggac ttctccaccc tgcaggacac taagtgtgaa 660
gttccatttg acatctgtca atctatctgt aagtacccag actacttgca aatgtccgct 720
gaccatacag gtgactctat gttcttctgt ttgagaagag aacaattggt cgctagacac 780
ttctggaaca gagctgggtg catgggtgac actgttccaa ctgacttgta catcaaggggt 840
acctctgcta acatgagaga aactccaggt tctgtgtct actctccatc tccatctggt 900
tctatcacta cttccgactc tcaattgttc aacaagccat actggttgca caaggctcaa 960
ggtcacaaca acggtatctg ttggcacaac caattgttcg tcaccgtcgt tgacactacc 1020
agatctacta acttgacctt gtgtgcttct actcaaaacc cagttccaaa cacttacgac 1080
ccaaccaagt tcaagcacta ctccagacac gtcgaggaat acgacttgca attcatcttc 1140
caattgtgta ctatcacctt gaccgtgaa gtcattgtct acattcactc tatgaactcc 1200
tctatcttgg aaaactggaa cttcgggtgt ccaccaccac caaccacctc cttgggtgac 1260
acttacagat tcttccaatc tgcgctgtc acttgtcaaa aggacaccac tccaccagaa 1320
aagcaagacc catacgacaa gttgaagttc tggactgttg acttgaagga aaagttctct 1380
tccgacttgg accaataccc attgggtaga aagttcttgg ttcaagctgg tttgagacgt 1440
agaccaacta tcgggtccacg taagagacca gctgcttcca cttccactgc ttctagacca 1500
gctaagcgtg tcagaatcag atccaagaag taa 1533

```

<210> 2

<211> 510

<212> PRT

<213> Human Papillomavirus Type 45

<400> 2

Met Ala Leu Trp Arg Pro Ser Asp Ser Thr Val Tyr Leu Pro Pro Pro
 1 5 10 15

WO 2005/032586

PCT/US2004/031326

Ser Val Ala Arg Val Val Asn Thr Asp Asp Tyr Val Ser Arg Thr Ser
 20 25 30
 Ile Phe Tyr His Ala Gly Ser Ser Arg Leu Leu Thr Val Gly Asn Pro
 35 40 45
 Tyr Phe Arg Val Val Pro Ser Gly Ala Gly Asn Lys Gln Ala Val Pro
 50 55 60
 Lys Val Ser Ala Tyr Gln Tyr Arg Val Phe Arg Val Ala Leu Pro Asp
 65 70 75 80
 Pro Asn Lys Phe Gly Leu Pro Asp Ser Thr Ile Tyr Asn Pro Glu Thr
 85 90 95
 Gln Arg Leu Val Trp Ala Cys Val Gly Met Glu Ile Gly Arg Gly Gln
 100 105 110
 Pro Leu Gly Ile Gly Leu Ser Gly His Pro Phe Tyr Asn Lys Leu Asp
 115 120 125
 Asp Thr Glu Ser Ala His Ala Ala Thr Ala Val Ile Thr Gln Asp Val
 130 135 140
 Arg Asp Asn Val Ser Val Asp Tyr Lys Gln Thr Gln Leu Cys Ile Leu
 145 150 155 160
 Gly Cys Val Pro Ala Ile Gly Glu His Trp Ala Lys Gly Thr Leu Cys
 165 170 175
 Lys Pro Ala Gln Leu Gln Pro Gly Asp Cys Pro Pro Leu Glu Leu Lys
 180 185 190
 Asn Thr Ile Ile Glu Asp Gly Asp Met Val Asp Thr Gly Tyr Gly Ala
 195 200 205
 Met Asp Phe Ser Thr Leu Gln Asp Thr Lys Cys Glu Val Pro Leu Asp
 210 215 220
 Ile Cys Gln Ser Ile Cys Lys Tyr Pro Asp Tyr Leu Gln Met Ser Ala
 225 230 235 240
 Asp Pro Tyr Gly Asp Ser Met Phe Phe Cys Leu Arg Arg Glu Gln Leu
 245 250 255
 Phe Ala Arg His Phe Trp Asn Arg Ala Gly Val Met Gly Asp Thr Val
 260 265 270
 Pro Thr Asp Leu Tyr Ile Lys Gly Thr Ser Ala Asn Met Arg Glu Thr
 275 280 285
 Pro Gly Ser Cys Val Tyr Ser Pro Ser Pro Ser Gly Ser Ile Thr Thr
 290 295 300
 Ser Asp Ser Gln Leu Phe Asn Lys Pro Tyr Trp Leu His Lys Ala Gln
 305 310 315 320
 Gly His Asn Asn Gly Ile Cys Trp His Asn Gln Leu Phe Val Thr Val
 325 330 335
 Val Asp Thr Thr Arg Ser Thr Asn Leu Thr Leu Cys Ala Ser Thr Gln
 340 345 350
 Asn Pro Val Pro Asn Thr Tyr Asp Pro Thr Lys Phe Lys His Tyr Ser
 355 360 365
 Arg His Val Glu Glu Tyr Asp Leu Gln Phe Ile Phe Gln Leu Cys Thr
 370 375 380
 Ile Thr Leu Thr Ala Glu Val Met Ser Tyr Ile His Ser Met Asn Ser
 385 390 395 400
 Ser Ile Leu Glu Asn Trp Asn Phe Gly Val Pro Pro Pro Pro Thr Thr
 405 410 415
 Ser Leu Val Asp Thr Tyr Arg Phe Val Gln Ser Val Ala Val Thr Cys
 420 425 430
 Gln Lys Asp Thr Thr Pro Pro Glu Lys Gln Asp Pro Tyr Asp Lys Leu
 435 440 445
 Lys Phe Trp Thr Val Asp Leu Lys Glu Lys Phe Ser Ser Asp Leu Asp
 450 455 460
 Gln Tyr Pro Leu Gly Arg Lys Phe Leu Val Gln Ala Gly Leu Arg Arg
 465 470 475 480
 Arg Pro Thr Ile Gly Pro Arg Lys Arg Pro Ala Ala Ser Thr Ser Thr
 485 490 495
 Ala Ser Arg Pro Ala Lys Arg Val Arg Ile Arg Ser Lys Lys
 500 505 510

WO 2005/032586

PCT/US2004/031326

<210> 3
 <211> 1533
 <212> DNA
 <213> Human Papillomavirus Type 45

<400> 3
 atggcctttgt ggcggcctag tgacagtacy gtatatcttc caccaccttc tgtggccaga 60
 gttgtcaaca ctgatgatta tgtgtctcgc acaagcatat tttaccatgc aggcagttcc 120
 cgattattaa ctgtaggcaa tccatatttt aggggtgtac ctagtgggtgc aggtaataaa 180
 caggctgttc ctaagggtatc cgcatatcag tatagggtgt ttagagtagc tttgcccgat 240
 cctaataaat ttggattacc tgattctact atatatataatc ctgaaacaca acgtttgggt 300
 tgggcatgtg taggtatgga aattgggtcgt gggcagcctt taggtattgg cctaagtggc 360
 catccatttt ataataaatt ggatgataca gaaagtgtc atgcagctac agctgttatt 420
 acgcaggatg ttagggataa tgtgtcagtt gattataagc aaacacagct gtgtatttta 480
 gggtgtgtac ctgctattgg tgagcactgg gccaaagggca cactttgtaa acctgcacaa 540
 ttgcaacctg gtgactgtcc tcctttggaa cttaaaaaaca ccattattga ggatgggtgat 600
 atgggtggata cagggttatgg ggcaatggat tttagtacat tgcaggatac aaagtgcgag 660
 gttccattag acattttgtca atccatctgt aaatatccag attatttgca aatgtctgct 720
 gatccctatg gggattctat gtttttttgc ctacgcctgt aacaactggt tgcaagacat 780
 ttttggaata gggcaggtgt tatgggtgac acagtaccta cagacctata tattaagggc 840
 actagcgcta atatgcgtga aacccttggc agtgtgtgtgt attccccttc tcccagtggc 900
 tctattacta cttctgattc tcaattattt aataagccat attgggttaca taaggcccag 960
 ggccataaca atgggtatttg ttggcataat cagttgtttg ttactgtagt ggacactacc 1020
 cgcagtacta atttaacatt atgtgcctct acacaaaatc ctgtgccaaa tacatatgat 1080
 cctactaagt ttaagcacta tagtagacat gtggagggaat atgatttaca gtttattttt 1140
 cagttgtgca ctattacttt aactgcagag gttatgtcat atatccatag tatgaatagt 1200
 agtatattgg aaaattggaa ttttggtgta cctccaccac ctactacaag tttagtggat 1260
 acatatcggt ttgtgcaatc agttgctgtt acctgtcaaa aggatactac acctccagaa 1320
 aagcaggatc catatgataa attaaagttt tggactgttg acctaaagga aaaattttcc 1380
 tccgatttgg atcaatatcc ccttgggtcga aagttttttag ttcaggctgg gttacgtcgt 1440
 aggcctacca taggacctcg taagcgtcct gctgcttcca cgtctactgc atctaggcct 1500
 gccaaacgtg tacgtatacg tagtaaaaaa taa 1533

<210> 4
 <211> 24
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> PCR Primer

<400> 4
 ccaccaccac ctatataggt attc

24

<210> 5
 <211> 24
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> PCR Primer

<400> 5
 caaacatata tatatgtgct aaca

24

<210> 6
 <211> 44
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> PCR Primer

WO 2005/032586

PCT/US2004/031326

<400> 6
ctcagatctc acaaaacaaa atggctttgt ggcggcctag tgac

44

<210> 7
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> PCR Primer

<400> 7
gacagatctt attttttact acgtatacgt acacg

35

<210> 8
<211> 1533
<212> DNA
<213> Artificial Sequence

<220>
<223> HPV45 L1R - Antisense

<400> 8
taccgaaaca cctctggtag actgagatga cagatgaacg gtggtggtag acagcgatct 60
cagcagttgt gactgctgat gcagaggtct tggaggtaga agatggtgcg accaagaagg 120
tctaacaact gacagccatt ggggtatgaag tctcagcagg gtaggccacg accattgttc 180
gttcgacaag gtttcagag acgaatgggtt atgtctcaga agtctcagcg aaacggtctg 240
ggtttggttca agccaaacgg tctgagatga tagatgttgg gtctttgagt ttctaaccag 300
acccgtacgc agccatacct ttagccatct ccagttggta acccatagcc aaacagacca 360
tggggtaaga tgttgttcaa cctgctgtgg cttagggcag tgccgacgat acgacagtag 420
tgagttctgc agtctctgtt gcagagacag ctgatgttcg tttgggttaa cacatagaac 480
ccaacacagg gtcgatagcc acttgtgacc cgattcccat ggaacacatt cggtcgagtt 540
aacgttggtc cactgacagg tggtaacctt aacttcttgt gatagtagct tctgccactg 600
taccaactgt gaccaatgcc acgatacctg aagaggtggg acgtcctgtg attcacactt 660
caaggtaacc tgtagacagt tagatagaca ttcattgggtc tgatgaacgt ttacaggcga 720
ctgggtatgc cactgagata caagaagaca aactcttctc ttgttaacaa gcgatctgtg 780
aagaccttgt ctcgaccaca gtaccactg tgacaagggt gactgaacat gtagttccca 840
tgagacgat tgtactctct ttgaggtcca aggacacaga tgagaggtag aggtagacca 900
agatagtgat gaaggctgag agttaacaag ttgttcggta tgaccaacgt gttccgagtt 960
ccagtgttgt tgccatagac aaccgtgttg gttaacaagc agtggcagca actgtgatgg 1020
tctagatgat tgaactggaa cacacgaaga tgagttttgg gtcaagggtt gtgaatgctg 1080
ggttggttca agttcgtgat gaggtctgtg cagctcctta tgctgaacgt taagtagaag 1140
gttaacacat gatagtggaa ctggcgactt cagtacagga tgtaagtgag atacttgagg 1200
agatagaacc ttttgacctt gaagccacaa ggtggtggtg gttggtggag gaaccaactg 1260
tgaatgtcta agcaggttag acagcgacag tgaacagttt tcctgtggtg aggtggtctt 1320
ttcgttctgg gtatgctgtt caacttcaag acctgacaac tgaacttcct tttcaagaga 1380
aggctgaacc tggttatggg taacctatct ttcaagaacc aagttcgacc aaactctgca 1440
tctggttgat agccagggtc attctctggt cgacgaaggt gaaggtgacg aagatctggt 1500
cgattcgcac agtcttagtc taggttcttc att 1533