

HIGH ALCOHOL CONTENT GEL-LIKE AND FOAMING COMPOSITIONS

CROSS REFERENCE TO RELATED U.S. APPLICATIONS

This patent application relates to, and claims the priority benefit from, United States Provisional Patent Application Serial No. 60/506,172 filed on September 29, 2003, and United States Provisional Patent Application Serial No. 60/591,601 filed on July 28, 2004 which is incorporated herein by reference in its entirety.

FIELD OF THE INVENTION

The present invention relates to compositions with high contents of lower alcohol (C₁₋₄) that could be a gel-like composition or a solution able to be dispensed as a foam. The compositions to be dispensed as foams contain a fluorosurfactant and when mixed with air provide a stable alcohol foam which can be used for personal cleaning or for disinfecting purposes. The gel-like compositions may or may not contain a fluorosurfactant.

BACKGROUND ART

Ethanol and/or Isopropyl alcohol compositions with at least 60% percent v/v (approximately 52% by weight) are well known to be antibacterial, therefore widely accepted for disinfecting purposes. Nonetheless due to the inherent characteristics of alcohol, it is perceived that the higher the content the better the product and a higher than 60% by volume alcohol content solution is more desirable.

Alcohol disinfectant solutions are generally thickened in order to eliminate the waste and facilitate spreading the composition throughout the desired area. It

is also known that other than gelling agents one can use paraffin or waxes to achieve thickening of a solution with high alcohol concentration. Such a composition added with lanolin to reduce the melting point closer to body temperature is described in United States Patent No. 2,054,989. One of the disadvantages of gels and such type thick alcohol containing compositions is that if they do not leave a tacky feeling on the hands after one use (although some do), the effect builds up after repetitive use during the day, making it necessary to eventually wash off the thickeners before continuing the usage of an alcohol antiseptic solution. The present invention if formulated for such type of product does not leave such a feel, and does not need to be washed off after having been used repeatedly.

Another way of thickening high alcohol content solutions has also been taught in United States Patent No. 6,090,395 and 6,623,744 where they use emulsifiers and surfactants as the thickening system to produce a hydroalcoholic lotion with a viscosity of at least 4,000cps. Also, United States Patent No. 4,956,170 discloses polyethoxylated non-ionic surfactants/emulsifiers to stabilize the added emollient oils in addition to a fatty alcohol although with the addition of a polymeric thickening agent to prepare a hydroalcoholic skin moisturizing/conditioning antimicrobial gel. The disinfecting compositions of the present invention that are gel-like have a viscosity lower than 4,000 cps and no polymeric thickening agent is added.

Generally speaking a high alcohol content disinfectant solution disinfects but does not clean. In order to make them disinfect and clean, so much soap would be needed that the skin would feel soapy and disagreeable, unacceptable for rubbing alcohol purposes. Nonetheless, a non-irritant skin disinfecting high

lower alcohol content formulation for use as a skin-washing agent is successfully attained by combining emulsifiers, surfactants and skin emollients to be used as a gel or ointment as described in United States Patent No. 5,629,006.

Surfactants other than for cleaning purposes are also used for spreading an aqueous composition containing one or more active substances rapidly and evenly over a surface due to their wetting properties. The use of good wetting agents definitely improves the efficient use of active substances in different compositions as described in United States Patent No. 5,928,993. Hence, the composition described in the present invention includes the addition of surfactants, specifically fluorosurfactants which are well known for their unparalleled wetting power and which are also surface-active in the lower alcohols used as disinfectant and solvent system in levels which make it acceptable even for rubbing alcohol purposes, providing cleaning, wetting and foaming properties to the composition.

Although a high alcohol content disinfectant solution has good disinfectant characteristics, it has a sharp smell and is generally perceived to cause drying of the skin, characteristics which can also be diminished to a desirable level in the present invention.

A greater than 40% v/v alcohol foam product, easy and safe to use, is desirable over conventional gel or ointment type composition products. The concentration of alcohol already poses a hazard in itself, and there are many applications in which the perceived risk may be diminished if it could be dispensed as a foam without the use of pressurized aerosol containers. A foam intended to be useful as a skin disinfecting agent must have a uniform consistency, spreadability, cleansing ability, and have a pleasant feel, i.e. have

rapid breaking power when pressure is applied; all of which present a challenge for a high lower alcohol content composition.

The description of an aqueous foaming skin disinfecting composition using 15% w/w alcohol as a co-solvent, which requires no pressurized container or added propellant to produce the foam, is described in United States Patent No. 3,962,150.

The foam-forming agents utilized heretofore, have been incapable of forming stable foams when the liquid phase has high alcohol content without using other ingredients. Furthermore, lower alcohols have been considered to be defoamers rather than foam-promoting chemicals. According to Klausner, in United States Patent No. 3,131,153, if more than 64% alcohol is used non-homogeneous compositions are obtained. The compositions in the patent required propellant to foam and the foams produced were of limited stability.

Prior to this invention, when a greater than 40% v/v alcohol concentration is required in a product, it is generally accepted that the product will be either liquid or gel, and that if a foam is desired then the concentration of alcohol would need to be reduced or the use of a propellant and a pressurized system would be required.

Surprisingly, in the few "foamable" high alcohol content products disclosed, the types of foam obtained were not similar to those expected from aqueous solutions. The foams obtained are described as fast or aerated foam, quick breaking, with low or limited stability, which would not last for more than one minute, being generally gone within seconds.

It has been disclosed that fluorosurfactants and alcohol can be combined to produce a "stable" foam by a process using high-pressurized means to

generate the foam. Highly stable pressurized foams containing high lower alcohol contents and methods of forming and using such pressurized foams in the oil industry using a non-ionic surfactant or mixture of non-ionic surfactants of a specific group of fluorosurfactants are provided in United States Patent No. 4,440,653. The compositions in this patent require the use of a pressurized gas system to generate the foam.

Various examples of compositions with a high lower alcohol content that are dispensed as a foam have been described, although for the purpose of the present invention the characteristics of the foam are not of the desired outcome, since they are fast breaking, of low stability and the foam is produced by means of propellants and aerosol containers only, as the one described in United States Patent No. 5,906,808, which discloses a product that uses an emulsifying wax NF, and a combination of stearyl and cetyl alcohol, or other wax combinations, which improve the foaming performance of the composition, in combination with cetyl lactate, to produce a 0.8% chlorhexidine gluconate alcohol product.

United States Patent No. 5,167,950 issued to Lins discloses a foam product which requires a propellant and no surfactant is added as a cleaning agent. The composition disclosed in this patent is based upon using an emulsifier system (fatty alcohol ROH 16-22 carbons) in combination with the use of a thickening agent (carbomer, klucel, etc.).

United States Patent No. 5,167,950 to Lins discloses an antimicrobial aerosol mousse having a high alcohol content. The mousse comprises alcohol, water, a polymeric gelling agent and a surfactant system comprising a C16-C22 alcohol, aerosol propellant and a non-ionic polyethoxylated surfactant. Despite the work done to date it has been shown that there is little specific knowledge on

how foams react and are formed, and surprisingly formulations that might seem not foamable result in the best foam producing ones while other formulations which seemed to have been producing foam even while being prepared did not perform well at all in some non-aerosol foam dispensers. The behaviour of aqueous foams is not the same of that of an alcohol foam.

The traditional ways of forming a gel using polymeric thickeners presents undesirable characteristics and similarly little has been done in forming emulsion-like thickened gels.

It would be very advantageous to have alcohol based disinfecting formulations which may be dispensed as either a gel or a foam. Further, it would be very advantageous and desirable to find a foaming agent that could be used in concentrations that would allow it to be used in products that can remain in the area on which they have been applied and do not need to be rinsed or wiped off due to small amounts of residue remaining after evaporation. Thus it would also be very advantageous to provide foams or gels that do not leave an unpleasant sticky after-feel as most commercial alcohol gel products are known to, or which clog up the dispensing equipment used to dispense the foams and gels.

SUMMARY OF THE INVENTION

It is an object of this invention to provide high alcohol content liquid compositions, which contain a surfactant/cleaning agent as well as a disinfectant/cleaning/solvent/carrier and that causes very little drying to the skin or the hands of the user and is able to be dispensed either as a gel or as a foam from both pressurized and non-pressurized systems.

The present invention provides high alcohol content compositions that are either gels or able to be dispensed as a foam, which are readily spread over the desired surface. Amongst the different applications where such compositions might be of use, it is another object to also provide an antimicrobial alcohol foam and an antimicrobial alcohol gel. The foamable compositions when dispensed from a suitable dispenser are stable and do not require the use of propellants and pressurized containers. The gels disclosed herein with a viscosity of less than 4,000cps do not use the gelling or thickening agents typically used in commercial gels and therefore after single or multiple applications of the gel there is not the usual tacky or sticky after-feel and the gel does not clog the dispensers from which the gels are dispensed.

These and other objects and advantages will be apparent from the following description of the invention. All percentages provided herein are based on the total weight unless otherwise indicated.

Accordingly, the present invention provides compositions for personal hygiene, as follows.

Foamable Compositions

The present invention provides a foamable composition, comprising:

- a) an alcohol C₁₋₄, or mixtures thereof, present in an amount greater than about 40% percent v/v of the total composition;
- b) an effective fluorinated surface active agent for wetting and foaming present in an amount of at least 0.001% weight percent of the total composition; and
- c) water present in an amount to balance the total composition to 100% weight percent.

In this aspect of the invention the effective fluorinated surface active agent is present in an amount from about 0.001% to about 10.0% weight percent of the total composition which is physiologically acceptable so it can be used in personal care type products.

In a preferred embodiment of the invention the fluorosurfactant may be an amphoteric polytetrafluoroethylene acetoxypentyl betaine of the following formula, $(CF_3CF_2(CF_2CF_2)_nCH_2CH_2(OAc)CH_2N^+(CH_3)_2CH_2COO^-)$ where $n = 2$ to 4 , an ethoxylated nonionic fluorosurfactant of the following structure: $R_fCH_2CH_2O(CH_2CH_2O)_xH$ where $R_f = F(CF_2CF_2)_y$, $x = 0$ to about 15 and $y = 1$ to about 7 ; or an anionic phosphate fluorosurfactant of the following structure: $(R_fCH_2CH_2O)_xP(O)(ONH_4)_y$ where $R_f = F(CF_2CF_2)_z$, $x = 1$ or 2 , $y = 2$ or 1 and $z = 1$ to about 7 , or mixtures thereof.

In another aspect of the invention there is provided an alcohol disinfecting composition, comprising;

- a) ethanol present in an amount between about 60% to 70% percent v/v of the total composition;
- b) a physiologically acceptable fluorosurfactant in an amount from about 0.01% to about 2.0% weight percent of the total composition;
- c) at least one nonionic surfactant selected from the group consisting of polyethoxylated fatty alcohols present in an amount from about 0.01 to about 10.0 % weight percent;
- d) a foam stabilizing agent;
- e) water in an amount to balance the total composition to 100% weight percent.

The present invention also provides an alcohol disinfecting composition comprising:

- a) ethanol present in an amount between about 60% to 70% percent v/v of the total composition;
- b) a physiologically acceptable anionic phosphate fluorosurfactant in an amount from about 0.01% to about 2.0% weight percent of the total composition;
- c) at least 1% n-propanol;
- d) foam stabilizing agents that at least include 1,3-Butyleneglycol % 2, Butoxyethanol in 0.001-3% ea;
- e) a lipid layer enhancer such as a mixture of alkylglucoside and glyceryl oleate; and
- f) water in an amount to balance the total composition to 100% weight percent.

Gel-Like Compositions

In this aspect of the invention there is provided an alcohol gel-like composition, comprising;

- a) an alcohol C₁₋₄, or mixtures thereof, present in an amount greater than about 40% percent v/v of the total composition;
- b) at least one nonionic surfactant selected from the group consisting of a poly(ethoxylated and/or propoxylated)alcohol, a poly(ethoxylated and/or propoxylated)ester, a derivative of a poly(ethoxylated and/or propoxylated)alcohol, a derivative of a poly(ethoxylated and/or propoxylated)ester, an alkyl alcohol, an alkenyl alcohol, an ester of a polyhydric alcohol, an ether of a polyhydric alcohol, an ester of a polyalkoxylated derivative

of a polyhydric alcohol, an ether of a polyalkoxylated derivative of a polyhydric alcohol, a sorbitan fatty acid ester, a polyalkoxylated derivative of a sorbitan fatty acid ester, and mixtures thereof, present in an amount between about 0.10 % to about 5% weight percent;

c) an emulsifier present in an amount between about 0.10 % to about 3.0% weight percent; and

d) water in an amount sufficient to form a stable gel-like disinfecting composition

The compositions disclosed herein provide a wide variety and range of high alcohol content disinfecting products. According to the percentage of alcohol used in the compositions and by varying the proportions of the other constituents in the formulation, foams with differing properties can be achieved thereby allowing the production of foams that are either coarse or wet which quickly flatten, or foams that are soft which contain fine bubbles and which are relatively dry having long foam stability, or creamy thick foams that are gel-like. Also, the compositions may or may not be disinfecting according to the percentage of alcohol.

It was surprisingly found that by varying the percentages of the ingredients an alcohol gel-like composition was obtained which did not dry the hands or leave a sticky after-feel and that did not clog the gel dispensers, having the desired consistency and showing a viscosity of less than 4,000 cps.

Some of the compositions can conveniently be manufactured in a two step process such that most of the alcohol can be added at a later time and/or location making it the first part a desirable concentrate suitable for shipping less hazardous goods and weight. Warming the first part from 30 to 80 degrees

Celsius, (depending on the particular composition) before adding the major portion of alcohol improves the long term stability of the compositions. This warming can either take place the same day in the same location where the finished composition is prepared or the concentrate first part can be stored or shipped elsewhere and the warming can take place either when the first part is mixing or right before adding the major portion of alcohol.

It should be evident that the described embodiment can be subjected to adjustment and/or improvement for specific applications either as a gel or a foam or to contain a desired active ingredient, without departing from the scope of the present invention. Different materials and/or ingredients will be then needed to compensate for the composition and/or foam stability disruption that might be generated by the change (i.e. introducing a more compatible secondary or even primary surfactant, adjusting the compatible foam stabilizer percentage and/or varying the relative amount of emulsifier and/or alcohol or water) or to compensate for shifts in desired viscosity and foam characteristics to obtain the desired gel (i.e. reduce the amount of fluorosurfactant or increase the polyethoxylated surfactants, or add an emulsifier and/or increase or decrease alcohol and/or water). These and other changes may be made in the details within the spirit of the invention, which is to be broadly construed and not to be limited except by the character of the claims appended hereto.

For example, the alcohol based compositions may contain up to 10 % by weight of other active ingredients or additives or combinations thereof commonly added to aerosol compositions or to toiletries, cosmetics, pharmaceuticals, etc. Materials that may be added may include organic gums and colloids, lower alkanolamides of higher fatty acids, short chain diols and/or triols,

alkylglucosides, fragrance, coloring matter, additional emollients, ultraviolet absorbers, solvents, emulsifiers, foam stabilizers or mixture of such stabilizers, suspending agents, buffers, conditioning agents, antioxidants, bactericides, medicinal active ingredient, and the like.

The present invention provides a composition, comprising;

a) an alcohol C_{1-4} , or mixtures thereof, present in an amount greater than about 40% v/v of the total composition;

b) at least one nonionic surfactant selected from the group consisting of a poly(ethoxylated and/or propoxylated)alcohol, a poly(ethoxylated and/or propoxylated)ester, a derivative of a poly(ethoxylated and/or propoxylated)alcohol, a derivative of a poly(ethoxylated and/or propoxylated)ester, an alkyl alcohol; an alkenyl alcohol, an ester of a polyhydric alcohol, an ether of a polyhydric alcohol, an ester of a polyalkoxylated derivative of a polyhydric alcohol, an ether of a polyalkoxylated derivative of a polyhydric alcohol, a sorbitan fatty acid ester, a polyalkoxylated derivative of a sorbitan fatty acid ester, and mixtures thereof, present in an amount between about 0.10 % to about 5% weight percent

c) an emulsifier present in an amount between about 0.10 % to about 3.0% weight percent; and

d) water in an amount sufficient to form a stable gel-like composition with a viscosity less than 4,000cps.

The invention will be described in connection with various specific examples, which are intended to be illustrative rather than limiting. Nevertheless, the present invention lends itself to the preparation of a wide variety of products,

and it is to be understood that other embodiments may be utilized and that changes may be made without departing from the scope of the invention.

DETAILED DESCRIPTION OF THE INVENTION

Definitions

The term “emollient” as used herein refers broadly to materials which are capable of maintaining or improving the moisture level, compliance, or appearance of the skin when used repeatedly.

The term “emulsifier” as used herein refers to surfactants or other materials added in small quantities to a mixture of two miscible liquids for the purpose of aiding in the formation and stabilization of an emulsion.

The phrase “emulsifying ingredients” as used herein is synonymous with emulsifier defined above.

The term “emulsion” as used herein refers to a colloidal dispersion of one liquid in another.

The term “surfactant” as used herein is the widely employed contraction for “surface active agents” which is the descriptive generic term for materials that preferentially adsorb at interfaces as a result of the presence of both lyophilic and lyophobic structural units, the adsorption generally resulting in the alteration of the surface or interfacial properties of the system.

The term “fluorosurfactant” as used herein refers to a fluorinated surface active agent which enables the composition in which it is contained to clean, wet and foam.

The phrase “foam stabilizer” as used herein refers to an additive that increases the amount or persistence of foam produced by a surfactant system.

The phrase "gel-like composition" as used herein refers to a hydroalcoholic solution with at least 40% v/v alcohol content, which is thickened by the use of emulsifiers and surfactant to have a viscosity greater than water and less than 4,000cps.

The term "disinfect" as used herein means to destroy or reduce harmful microorganisms.

The present invention provides compositions with high contents of lower alcohol (C_{1-4}) able to be dispensed as a gel or a foam. The foamable compositions when mixed with air deliver a stable foam to provide an alcoholic liquid solution which can be used for personal cleaning or for disinfecting purposes and which breaks on pressure application such as when a user rubs their hands or when applied over a surface. The gel composition delivers a liquid of the appropriate consistency to be readily spread on the hands, yet without dripping off. This gel composition with at least 60% v/v alcohol provides an effective disinfectant that does not leave a tacky after-feel once the alcohol has evaporated and that is common to such alcohol gels which use thickeners and gelling agents that have been used commonly in the past during single or multiple applications. The gel composition does not easily clog the dispensers as common gel products do.

The alcohol used in the present invention is a lower hydrocarbon chain alcohol such as a C_{1-4} alcohol. The preferred alcohol is chosen from ethanol, 2-propanol, or n-propanol, most preferably ethanol, well accepted by Health Care personnel as an adequate disinfectant at the right percentages. The invention anticipates that a single alcohol may be used or that a blend of two or more

alcohols may comprise the alcohol content of the composition either for a gel-like or foamable product.

FOAMABLE COMPOSITIONS

One of the main achievements of the present invention is making compositions with a greater than 40% v/v alcohol content able to be dispensed as a cosmetically appealing foam. The other important achievement is to obtain an alcohol gel without using the typical gelling agents known to those skilled in the art which would not clog dispensers or leave a tacky after-feel.

The use of a fluorosurfactant is the key ingredient as the primary foaming agent in the compositions designed to foam disclosed herein. Fluorosurfactants have various interesting properties such as leaving little residue, being able to function in harsh chemical and thermal environments; they have an unparalleled wetting power, etc. Unlike traditional surfactants, they show unusual surface-active properties in organic solvents that are known to those skilled in the art, and that have made them widely used for applications in coatings, oilfield, material finishes, cleaning, paints, etc.

The fluorosurfactants suitable for these types of compositions may include, but are not limited to, ethoxylates, glycerol esters, amine oxides, acetylenic alcohol derivatives, carboxylates, phosphates, carbohydrate derivatives, sulfonates, betaines, esters, polyamides, silicones, and hydrocarbon surfactants that have been fluorinated and are compatible with the other components being used for a particular formulation.

A preferred fluorosurfactant is polytetrafluoroethylene acetoxypropyl betaine $\text{CF}_3\text{CF}_2(\text{CF}_2\text{CF}_2)_n\text{CH}_2\text{CH}_2(\text{OAc})\text{CH}_2\text{N}^+(\text{CH}_3)_2\text{CH}_2\text{COO}^-$, where $n=2-4$. However, it is contemplated that other fluorosurfactants may be used including as

non-limiting examples for use in the present invention an ethoxylated nonionic fluorosurfactant of the following structure: $\text{RfCH}_2\text{CH}_2\text{O}(\text{CH}_2\text{CH}_2\text{O})_x\text{H}$ where $\text{Rf} = \text{F}(\text{CF}_2\text{CF}_2)_y$, $x = 0$ to about 15 and $y = 1$ to about 7; an anionic phosphate fluorosurfactant of the following structure: $(\text{RfCH}_2\text{CH}_2\text{O})_x\text{P}(\text{O})(\text{ONH}_4)_y$ where $\text{Rf} = \text{F}(\text{CF}_2\text{CF}_2)_z$, $x = 1$ or 2, $y = 2$ or 1, $x + y = 3$, and $z = 1$ to about 7 or mixtures thereof.

It was surprisingly found that despite the characteristics of fluorosurfactants, there was little or no information on their use to produce a foamable product with high alcohol content other than the one using non-ionic fluorosurfactants and pressurized gas as a foamer as taught in United States Patent No. 4,440,653 incorporated herein by reference.

Furthermore, in order to obtain a high alcohol content product able to produce a foam even if no pressurized containers or propellants are used would require surface tension values as low as possible so that the pressure required to produce such foam by hand pumps and mechanical means would be sufficient. Hence, the lower than 20 dynes/cm (0.01% DW 25°C) surface tension values achievable with these surfactants made them suitable for the application.

During the development of the present invention, it was unexpectedly found that a quick breaking aerated foam could even be obtained when using just ethanol and the fluorosurfactant, while using traditional surfactants at even double the percentage bore results that could not be even slightly similar and no foam at all could be obtained.

In order to achieve a commercially suitable formulation, reducing the amount of fluorosurfactant used while using the assistance of other ingredients such as secondary surfactants, emulsifiers, foam stabilizers, fragrances, and the

like ingredients employed in cosmetics, aerosols, toiletries, personal care, etc. is one of the approaches followed. One of the commercial products obtained uses emulsifiers and polyethoxylated fatty acid surfactants disclosed in United States Patent Nos. 5,167,950 and 6,090,395, both incorporated herein by reference, while other examples use a combination of different foam stabilizers to achieve a similar result.

Examples of secondary surfactants that may be used in the present compositions include alkylglucosides, a poly(ethoxylated and/or propoxylated)alcohol, a poly(ethoxylated and/or propoxylated)ester, a derivative of a poly(ethoxylated and/or propoxylated)alcohol, a derivative of a poly(ethoxylated and/or propoxylated)ester, an alkyl alcohol, an alkenyl alcohol, an ester of a polyhydric alcohol, an ether of a polyhydric alcohol, an ester of a polyalkoxylated derivative of a polyhydric alcohol, an ether of a polyalkoxylated derivative of a polyhydric alcohol, a sorbitan fatty acid ester, a polyalkoxylated derivative of a sorbitan fatty acid ester, a betaine, a sulfobetaines, imidazoline derivatives, aminoacid derivatives, lecithins, phosphatides, some amine oxides and sulfoxides and mixtures thereof, present in an amount between about 0.10 % to about 5% weight percent.

A preferred betaine is cocamidopropyl betaine. A preferred alkylglucoside is cocoglucoside. Preferred polyethoxylated fatty alcohols are polyethoxylated stearyl alcohol (21 moles ethylene oxide) and polyethoxylated stearyl alcohol (2 moles ethylene oxide), and a combination of these two.

The compositions may include an antimicrobial agent. The following antimicrobials are offered as non-limiting examples of suitable antimicrobials for use in the present invention and may include chlorhexidine salt, iodine, a

complexed form of iodine, parachlorometaxilenol, triclosan, hexachlorophene, a phenol, a surfactant having a long chain hydrophobic group and a quaternary group, hydrogen peroxide, silver, a silver salt, silver oxide, and mixtures thereof.

A preferred antimicrobial agent in the present compositions is chlorhexidine gluconate (CHG) present in an amount between about 0.50 % to about 4.0 % weight percent. Another preferred antimicrobial agent is didecyl dimethyl diamonium chloride in an amount between about 0.05% to 5% weight percent.

If the amount of ingredients employed is little enough not to leave a tacky feeling after the composition evaporates after single or multiple uses, and this is achieved while maintaining at least 60% v/v ethanol or n-propanol concentration or 70% v/v isopropanol, then the composition would be ideal for use as an alcohol hand sanitizer/disinfectant foamable composition.

The addition of water to the alcohol produces a more stable foam while allowing to reduce the amount of fluorosurfactant required to foam the product. For instance, using 0.5 to 1.0% fluorosurfactant with a 50 to 60% v/v alcohol water solution produces a stable foam that does not readily collapse and that produces a stable puff that does not fall even when inverted and does not collapse until pressure is applied (such as when rubbed in hands or on over a surface) to provide an alcoholic liquid solution.

The use of a mild non-irritant surfactant widely used in the cosmetic industry such as cocamidopropyl betaine as a secondary surfactant is more suitable to prepare the foamable hydroalcoholic composition of the present invention depending on the fluorosurfactant being used.

In order to stabilize the foam, foam stabilizers, as well as emulsifying ingredients have been tried with good results in allowing the product to be dispensed as a foam even when no propellant and/or pressurized container systems are used.

Examples of compatible foam stabilizers that can optionally be employed include lactic acid esters of monoglycerides, cationic emulsifiers, triquaternized stearic phospholipid complex, hydroxystearamide propyltriamine salts, lactic acid monoglycerides, food emulsifiers such as glyceryl monostearate, propylene glycol monostearate, sodium stearyl lactylate, silicone wax, an encapsulated oil, Microcapsule Mineral Oil.

A preferred foam stabilizer used in the present foamable compositions is cetyl betaine. A preferred combination of foam stabilizers is that of butyleneglycol, butoxyethanol and n-propanol.

Examples of moisturizers and/or emollients which may be used in the present formulations include lanolin, vinyl alcohol, polyvinyl pyrrolidone and polyols selected from the group consisting of glycerol, propylene glycol, glyceryl oleate and sorbitol, cocoglucoside or a fatty alcohol selected from the group consisting of cetyl alcohol, stearyl alcohol, lauryl alcohol, myristyl alcohol and palmityl alcohol, cetyl alcohol, cetareth 20, and combinations thereof, present in an amount up to about 5%.

The compositions formulated to be dispensed as a foam may be stored in an unpressurized dispenser having a dispenser pump for mixing the composition with air and dispensing foam therefrom. The composition may include an aerosol propellant in an amount from about 3 to about 20 weight percent of the total composition for pressurized discharge of the foam. The aerosol propellant may

include propane, carbon dioxide, butane, dichloro difluoro methane, dichloro tetra fluoro ethane, octafluorocyclo butane; 1,1,1,2-tetrafluoroethane; 1,1,1,2,3,3,3 heptafluoropropane, and 1,1,1,3,3,3,-hexafluoropropane.. When stored in a metal container with propellant, the formulation may include a corrosion inhibitor such as sorbic acid, benzoic acid, potassium sorbate and sodium benzoate, in an amount from about 0.1 to about 5 weight percent of the total composition.

GEL-LIKE COMPOSITIONS

Some of compositions studied had some gel-like properties. This characteristic led to the second most important achievement of the present invention; that is an alcohol gel with viscosities less than 4,000cps that do not use the conventional polymeric thickeners (i.e, cellulose derivatives, carbomers, etc) that are known to leave a sticky residue on surfaces on single and multiple applications that builds up. This discourages users and tends to clog the dispensers.

In order to prepare a gel-like composition, a fluorosurfactant is not required to form the gel, however, using a small amount improves the after-feel, it also allows one to reduce the usage of other surfactants required, therefore improving the performance of the composition. The use of the fluorosurfactant also noticeably improves the spreadability of the gel disinfecting compositions on the hands or a surface.

The following is a basic formulation of the gel-like compositions. An alcohol C₁₋₄, or mixtures thereof, present in an amount greater than about 40% v/v of the total composition, one or more nonionic surfactants present in an amount between about 0.10 % to about 5% weight percent, an emulsifier present

in an amount between about 0.10 % to about 3.0% weight percent, and water in an amount sufficient to form a stable gel-like composition.

Non-limiting examples of non-ionic surfactants include poly(ethoxylated and/or propoxylated)alcohol, a poly(ethoxylated and/or propoxylated)ester, a derivative of a poly(ethoxylated and/or propoxylated)alcohol, a derivative of a poly(ethoxylated and/or propoxylated)ester, an alkyl alcohol, an alkenyl alcohol, an ester of a polyhydric alcohol, an ether of a polyhydric alcohol, an ester of a polyalkoxylated derivative of a polyhydric alcohol, an ether of a polyalkoxylated derivative of a polyhydric alcohol, a sorbitan fatty acid ester, a polyalkoxylated derivative of a sorbitan fatty acid ester, and mixtures thereof.

A preferred non-ionic surfactant includes polyethoxylated fatty alcohols such as polyethoxylated stearyl alcohol (21 moles ethylene oxide) and polyethoxylated stearyl alcohol (2 moles ethylene oxide), and/or a combination of polyethoxylated fatty alcohols.

To make the gel-like compositions for personal hygiene applications, the compositions may include a physiologically acceptable fluorinated surface active agent up to about 2.5 %. A preferred fluorinated surface active agent is that used in the gel-like compositions, namely polytetrafluoroethylene acetoxypentyl betaine $\text{CF}_3\text{CF}_2(\text{CF}_2\text{CF}_2)_n\text{CH}_2\text{CH}_2(\text{OAc})\text{CH}_2\text{N}^+(\text{CH}_3)_2\text{CH}_2\text{COO}^-$, where $n = 2-4$.

Another fluorinated surface active agent also preferred is an ethoxylated nonionic fluorosurfactant of the following structure: $\text{RfCH}_2\text{CH}_2\text{O}(\text{CH}_2\text{CH}_2\text{O})_x\text{H}$ where $\text{Rf} = \text{F}(\text{CF}_2\text{CF}_2)_y$, $x = 0$ to about 15 and $y = 1$ to about 7 and yet another one is an anionic phosphate fluorosurfactant of the following structure:

$(\text{RfCH}_2\text{CH}_2\text{O})_x\text{P}(\text{O})(\text{ONH}_4)_y$ where $\text{Rf} = \text{F}(\text{CF}_2\text{CF}_2)_z$, $x = 1$ or 2, $y = 2$ or 1, $x + y = 3$, and $z = 1$ to about 7 or mixtures thereof.

The following non-limiting examples are set forth to show for the various preferred embodiments and are not in any way to limit the scope of the present invention.

EXAMPLES

Examples 1 through 12 were prepared to illustrate the ability to produce alcohol-based formulations which can be dispensed as foams using different surfactants and a solution of water and 50% ethanol. Examples 13 through 18 show increasing concentrations of ethanol and fluorosurfactant to produce foam. Examples 19 through 30 illustrate the ability to produce foam using different surfactants and a solution of 70%v/v Isopropanol. All parts and percentages are expressed by weight unless otherwise indicated.

Amount Ingredients	Ex. 1	Ex. 2	Ex. 3	Ex. 4	Ex. 5	Ex. 6
S.D. Alcohol 3-A	50.00	50.00	50.00	50.00	50.00	50.00
Fluorosurfactant	0.50	-----	-----	-----	-----	-----
Cocamidopropyl betaine (1)	-----	8.00	-----	-----	-----	-----
Alkylglucoside (2)	-----	-----	8.00	-----	-----	-----
Alkylglucoside (3)	-----	-----	-----	8.00	-----	-----
Glycomul L	-----	-----	-----	-----	8.00	-----
Sorbitan Sesquioleate	-----	-----	-----	-----	-----	8.00
Deionized Water	49.50	42.00	42.00	42.00	42.00	42.00
Total %	100.00	100.00	100.00	100.00	100.00	100.00

(1) Amphoteric, (2) Nonionic, (3) Anionic

Amount Ingredients	Ex. 7	Ex. 8	Ex. 9	Ex. 10	Ex. 11	Ex. 12
S.D. Alcohol 3-A	50.00	50.00	50.00	50.00	50.00	50.00
Polysorbate 20	8.00	-----	-----	-----	-----	-----
Polyoxyethylene Sorbitan Monooleate	-----	8.00	-----	-----	-----	-----
Sorbitan Monooleate	-----	-----	8.00	-----	-----	-----
Cocamidopropyl betaine & sodium caproyl lactate	-----	-----	-----	8.00	-----	-----
Cocamidopropyl hydroxysultaine	-----	-----	-----	-----	8.00	-----
Sodium Cocoamphoacetate	-----	-----	-----	-----	-----	8.00
Deionized Water	42.00	42.00	42.00	42.00	42.00	42.00
Total %	100.00	100.00	100.00	100.00	100.00	100.00

Amount Ingredients	Ex. 13	Ex. 14	Ex. 15	Ex. 16	Ex. 17	Ex. 18
S.D. Alcohol 3-A	50.00	60.00	65.00	70.00	80.00	92.50
Fluorosurfactant	0.10	0.75	0.80	1.50	2.00	7.5
Deionized Water	49.90	39.25	34.20	28.50	18.00	-----
Total %	100.00	100.00	100.00	100.00	100.00	100.00

Amount Ingredients	Ex. 19	Ex. 20	Ex. 21	Ex. 22	Ex. 23	Ex. 24
70%v/v Isopropanol	99.90	92.00	92.00	92.00	92.00	92.00
Fluorosurfactant	0.10	-----	-----	-----	-----	-----
Cocamidopropyl betaine (1)	-----	8.00	-----	-----	-----	-----
Alkylglucoside (2)	-----	-----	8.00	-----	-----	-----
Alkylglucoside (3)	-----	-----	-----	8.00	-----	-----
Glycomul L	-----	-----	-----	-----	8.00	-----
Sorbitan Sesquioleate	-----	-----	-----	-----	-----	8.00
Total %	100.00	100.00	100.00	100.00	100.00	100.00

(1) Amphoteric, (2) Nonionic, (3) Anionic

Amount Ingredients	Ex. 25	Ex. 26	Ex. 27	Ex. 28	Ex. 29	Ex. 30
70%v/v Isopropanol	92.00	92.00	92.00	92.00	92.00	92.00
Polysorbate 20	8.00	-----	-----	-----	-----	-----
Polyoxyethylene Sorbitan Monooleate	-----	8.00	-----	-----	-----	-----
Sorbitan Monooleate	-----	-----	8.00	-----	-----	-----
Cocamidopropylbetaine & sodium caproyl lactate	-----	-----	-----	8.00	-----	-----
Cocamidopropyl hydroxysultaine	-----	-----	-----	-----	8.00	-----
Sodium Cocoamphoacetate	-----	-----	-----	-----	-----	8.00
Total %	100.00	100.00	100.00	100.00	100.00	100.00

The solutions prepared, were evaluated as to whether foam was produced or not and if so, then the foam produced was described as follows:

Example	Foam Produced	Foam Evaluation/Description/Characteristics
Ex. 1	Yes	Very good stable stiff puff creamy and soft lasts minutes
Ex. 2	No	Just Very Wet Bubbles produced lasting <10 seconds
Ex. 3	No	-----
Ex. 4	No	-----
Ex. 5	No	-----
Ex. 6	No	Just Very Wet Bubbles produced lasting <7 seconds
Ex. 7	No	Just Very Wet Bubbles produced lasting <10 seconds
Ex. 8	No	-----
Ex. 9	No	Just Very Wet Bubbles produced lasting <10 seconds
Ex. 10	No	-----
Ex. 11	No	-----
Ex. 12	No	-----
Ex. 13	Yes	Very good stable stiff puff creamy and soft lasts minutes
Ex. 14	Yes	Very good creamy and soft lasts more than a minute
Ex. 15	Yes	Very good creamy and soft lasts more than a minute
Ex. 16	Yes	Very good creamy and soft lasts more than a minute
Ex. 17	Yes	Quick fast breaking foam lasts more than a 15secs
Ex. 18	Yes	Quick fast breaking foam lasts more than a 10secs
Ex. 19	Yes	Quick fast breaking foam lasts more than a 20secs
Ex. 20	No	-----
Ex. 21	No	-----
Ex. 22	No	-----
Ex. 23	No	-----
Ex. 24	No	-----
Ex. 25	No	-----
Ex. 26	No	-----
Ex. 27	No	-----
Ex. 28	No	-----
Ex. 29	No	-----
Ex. 30	No	-----

Comparatively, it was also found that for instance, Cocamidopropyl betaine (CAPB) alone even at 40% ethanol and at 3% CAPB, was unable to produce as good results as those with 60%v/v ethanol, and fluorosurfactants using much less percentage (less than 1.0%). Cocamidopropyl betaine does not give any acceptable foam above that percentage of alcohol and the lower than 60% v/v alcohol content makes it inadequate for a sanitizing solution. Also the solution left an unacceptable feeling on the skin after the alcohol evaporated (i.e. a soapy sticky feeling) indicating high levels of surfactant.

Very interestingly fluorosurfactants seemed to be a likely way to achieve a foaming composition that contains more than 40% v/v alcohol. The fact that foam could be achieved even when no added water or ingredients are used other than

95%v/v alcohol and the fluorosurfactant as shown in example 18 makes the present invention suitable for many different applications.

Below are some specific examples for compositions following the above formulation to produce alcohol hand sanitizing solutions; more than one being a foamable composition with alcohol being the only disinfectant ingredient, while other foamable compositions use an added antimicrobial such as Chlorhexidine Digluconate or Didecyl Dimethyl Diammonium Chloride and the third group being alcohol gel-like hand sanitizing solutions.

EXAMPLE 31

Alcohol hand sanitizing foamable disinfecting composition

0.01- 1.0 % * amphoteric, anionic or non-ionic fluorosurfactant (primary surfactant)

0.01 – 1.0% cocoamidopropylbetaine (secondary surfactant)

0.05 – 1.0% cetyl betaine (foam stabilizing agent)

0.10 – 1.5% emulsifier fatty alcohol ROH 16-22 carbons or combination that works well in a final formulation containing

60-70% v/v ethanol

Q.S. water

- Preferably Polytetrafluoroethylene Acetoxypentyl Betaine

$\text{CF}_3\text{CF}_2(\text{CF}_2\text{CF}_2)_n \text{CH}_2\text{CH}_2(\text{OAc})\text{CH}_2\text{N}^+(\text{CH}_3)_2\text{CH}_2\text{COO}^-$, where $n = 2-4$ or

an ethoxylated nonionic fluorosurfactant of the following structure:

$\text{RfCH}_2\text{CH}_2\text{O}(\text{CH}_2\text{CH}_2\text{O})_x\text{H}$ where $\text{Rf} = \text{F}(\text{CF}_2\text{CF}_2)_y$, $x = 0$ to about 15

and $y = 1$ to about 7; or an anionic phosphate fluorosurfactant of the

following structure: $(\text{RfCH}_2\text{CH}_2\text{O})_x \text{P}(\text{O})(\text{ONH}_4)_y$ where $\text{Rf} = \text{F}(\text{CF}_2$

$\text{CF}_2)_z$, $x = 1$ or 2 , $y = 2$ or 1 , $x + y = 3$, and $z = 1$ to about 7 , or mixtures thereof.

EXAMPLE 32

Alcohol hand sanitizing foamable disinfecting composition Concentrate

0.01- 1.0 % * amphoteric, anionic or non-ionic fluorosurfactant (primary surfactant)

0.01 – 12.0% 1,3 Butyleneglycol, 2-Butoxyethanol, n-propanol (foam stabilizing agents)

0.05 – 5.0% cocoglucoside, glycerin, glyceryl oleate (moisturizers, emollients and the like)

60-70% v/v ethanol, n-propanol, isopropanol or a combination thereof

Q.S. water

- Preferably an anionic phosphate fluorosurfactant of the following structure: $(\text{RfCH}_2\text{CH}_2\text{O})_x \text{P}(\text{O})(\text{ONH}_4)_y$ where $\text{Rf} = \text{F}(\text{CF}_2 \text{CF}_2)_z$, $x = 1$ or 2 , $y = 2$ or 1 , $x + y = 3$, and $z = 1$ to about 7 or an ethoxylated nonionic fluorosurfactant of the following structure: $\text{RfCH}_2\text{CH}_2\text{O}(\text{CH}_2\text{CH}_2\text{O})_x\text{H}$ where $\text{Rf} = \text{F}(\text{CF}_2 \text{CF}_2)_y$, $x = 0$ to about 15 and $y = 1$ to about 7 ; or

Polytetrafluoroethylene Acetoxypropyl Betaine

$\text{CF}_3\text{CF}_2(\text{CF}_2\text{CF}_2)_n \text{CH}_2\text{CH}_2(\text{OAc})\text{CH}_2\text{N}^+(\text{CH}_3)_2\text{CH}_2\text{COO}^-$, where $n = 2-4$, or mixtures thereof.

EXAMPLE 33**Alcohol hand sanitizing foamable disinfecting composition**

- a) ethanol present in an amount between about 60% to 70% percent v/v of the total composition;
- b) a physiologically acceptable anionic phosphate fluorosurfactant in an amount from about 0.01% to about 2.0% weight percent of the total composition;
- c) at least 1% n-propanol
- d) foam stabilizing agents that at least include 1,3-Butyleneglycol % 2, Butoxyethanol in 0.001-3% ea.
- e) a lipid layer enhancer such as a mixture of alkylglucoside and glyceryl oleate
- f) water in an amount to balance the total composition to 100% weight percent.

EXAMPLE 34**Chlorhexidine gluconate (CHG) & alcohol hand sanitizing foamable disinfecting composition**

Formulations 31 or 32 added with

0.50 – 4.0% Chlorhexidine Gluconate (CHG)

EXAMPLE 35

Formulations 31 or 32 added with

0.01 – 5.0% Didecyl Dimethyl Diammonium Chloride

EXAMPLE 36**Alcohol hand sanitizing gel-like disinfecting composition with a viscosity less than 4,000cps**

- 0.0- 1.0 % * amphoteric, anionic or non-ionic fluorosurfactant (primary surfactant)
- 0.10 – 2.0 % an emulsifier moisturizer and/or emollient preferably a non-ionic surfactant and/or a combination of cetearyl alcohol and ceteareth 20 or a combination thereof to give a composition with a viscosity of less than 4,000cps;
- 0.50 – 4.0% a combination of nonionic surfactants specifically from the group of the polyethoxylated fatty alcohols
- 60-70%v/v ethanol
- Q.S. water

- * Preferably Polytetrafluoroethylene Acetoxypentyl Betaine $\text{CF}_3\text{CF}_2(\text{CF}_2\text{CF}_2)_n\text{CH}_2\text{CH}_2(\text{OAc})\text{CH}_2\text{N}^+(\text{CH}_3)_2\text{CH}_2\text{COO}^-$, where $n = 2-4$ or an ethoxylated nonionic fluorosurfactant of the following structure: $\text{RfCH}_2\text{CH}_2\text{O}(\text{CH}_2\text{CH}_2\text{O})_x\text{H}$ where $\text{Rf} = \text{F}(\text{CF}_2\text{CF}_2)_y$, $x = 0$ to about 15 and $y = 1$ to about 7; or an anionic phosphate fluorosurfactant of the following structure: $(\text{RfCH}_2\text{CH}_2\text{O})_x\text{P}(\text{O})(\text{ONH}_4)_y$ where $\text{Rf} = \text{F}(\text{CF}_2\text{CF}_2)_z$, $x = 1$ or 2, $y = 2$ or 1, $x + y = 3$, and $z = 1$ to about 7, or mixtures thereof.

The process to prepare the compositions of the present invention described herein is straightforward since most of the ingredients are liquid. When

wax type ingredients are to be used, they can be incorporated by warming up to 40-45°C preferably to the alcohol portion while mixing and then allowing it to cool down or they could be added in "cold", at room temperature to the alcohol before any other ingredient and mixed until completely incorporated before adding the rest of the ingredients according to the composition. Whether all ingredients are liquid or not, warming from 30 to 80 degrees Celsius, (depending on the particular composition) increases the long term stability of the compositions. Active ingredients could be pre-dissolved into the water first. A process that anyone knowledgeable enough of the art would have no problem implementing. If a specific formulation cannot be adjusted for the foamable composition in the percentages of the ingredients, then there is still the option of modifying the characteristics of the foaming pump, such as changing pressures, screen sizes, etc.

The compositions described within the present invention improve over prior similar products commercially available in the high concentrations of alcohol, as well as in the fact of being able to foam even with no propellants or pressurized containers (using propellants would improve considerably the quality of the foam) and being able to produce alcohol gel-like compositions that do not leave a sticky after-feel that builds up and do not clog the dispensers after single or multiple applications.

Depending on the alcohol concentration and the application of the particular composition the foam produced can widely vary, being at the high end of a relatively fast breaking variety stable enough to be thoroughly spread onto the skin without waste in a unique way and the gel-like composition viscosity varies with the alcohol concentration. The gel-like composition obtained is a

unique approach that does not follow the traditional ways of making alcohol gels. In summary it could be said that the stated invention has exceeded expectations.

Due to the nature of the base composition with respect to the alcohol concentration and the quality of the ingredients, one of the logical first applications for the present invention would be as an alcohol hand disinfectant composition either for a foamable product or an alcohol gel-like product, examples of which are described above. Nevertheless, the present invention lends itself to the preparation of a wide variety of products, and it is to be understood that other embodiments may be utilized and that changes may be made without departing from the scope of the invention.

Consequently, it is intended that the claims be interpreted to cover such modifications and equivalents. To note a few, the following may be mentioned: medicated foams and gels, sunscreen foams and gels, hand cream foams, brush-less shaving cream foams, shower or bath oil foams, dry hair shampoo foams, make-up remover foams, analgesic foam rubs and gels, hair grooming foams and antiperspirants hair cleaning foam, antiperspirant foam, hair conditioner foams.

As used herein, the terms "comprises", "comprising", "includes" and "including" are to be construed as being inclusive and open ended, and not exclusive. Specifically, when used in this specification including claims, the terms "comprises", "comprising", "includes" and "including" and variations thereof mean the specified features, steps or components are included. These terms are not to be interpreted to exclude the presence of other features, steps or components.

The foregoing description of the preferred embodiments of the invention has been presented to illustrate the principles of the invention and not to limit the

invention to the particular embodiment illustrated. It is intended that the scope of the invention be defined by all of the embodiments encompassed within the following claims and their equivalents.

References Cited U.S. PATENT DOCUMENTS

2,054,989/1936	Moore	167/58
3,131,153/1964	Klausner	252/305
3,962,150/1976	Leonard et al.	252/542
4,440,653/1984	James et al.	252/8.55
5,167,950 12/1992	Lins	424/47
4,956,170 09/1990	Lee.....	514/772.1
5,629,006/1997	Minh et al.	424/405
5,906,808/1999	Osborne, et al.....	424/43
5,928,993/1999	Ingegärd	504/116
5,951,993 09/1999	Scholz et al.....	424/405
6,090,395 07/2000	Asmus et al.....	424/401
6,610,315 08/2003	Scholz et al.....	424/415
6,623,744 09/2003	Asmus et al.....	424/401
6,562,360 05/2003	Scholz et al.....	424/405

OTHER PUBLICATIONS

Myers, Drew; "Surfactant Science and Technology", second edition, Drew Myers, VCH Publishers, New York, 1992

Reduce Tension Dupont Zonyl® Fluorosurfactants Field Manual published by Dupont Co on May 2001

WHAT IS CLAIMED IS:

1. A composition, comprising;
 - a) an alcohol C₁₋₄, or mixtures thereof, present in an amount greater than about 40% v/v of the total composition;
 - b) an effective fluorinated surface active agent for wetting and foaming present in an amount of at least 0.001% weight percent of the total composition; and
 - c) water present in an amount to balance the total composition to 100% weight percent.
2. The composition according to claim 1 including air, wherein when the composition is mixed with air, the mixture of the composition and air forms a foam.
3. The composition according to claims 1 or 2 wherein the effective fluorinated surface active agent is a fluorosurfactant present in an amount from about 0.001% to about 10.0% weight percent of the total composition.
4. The composition according to claim 1, 2 or 3 wherein the effective fluorinated surface active agent is a physiologically acceptable fluorosurfactant, and wherein the fluorosurfactant is selected from the group consisting of fluorinated ethoxylates, glycerol esters, amine oxides, acetylenic alcohol derivatives, carboxylates, phosphates, carbohydrate derivatives,

sulfonates, betaines, esters, polyamides, silicones, and hydrocarbon surfactants.

5. The composition according to claim 1, 2, 3 or 4 wherein the alcohol C₁₋₄ is an aliphatic alcohol selected from the group consisting of methanol, ethanol, isopropanol, n-propanol, butanol and combinations thereof.

6. The composition according to claim 1, 2, 3, 4 or 5 wherein the fluorosurfactant is an amphoteric polytetrafluoroethylene acetoxypopyl betaine $(\text{CF}_3\text{CF}_2(\text{CF}_2\text{CF}_2)_n\text{CH}_2\text{CH}_2(\text{OAc})\text{CH}_2\text{N}^+(\text{CH}_3)_2\text{CH}_2\text{COO}^-)$ where $n = 2$ to 4.

7. The composition according to claim 1, 2, 3, 4 or 5 wherein the fluorosurfactant is an ethoxylated nonionic fluorosurfactant of the following structure: $\text{RfCH}_2\text{CH}_2\text{O}(\text{CH}_2\text{CH}_2\text{O})_x\text{H}$ where $\text{Rf} = \text{F}(\text{CF}_2\text{CF}_2)_y$, $x = 0$ to about 15, and $y = 1$ to about 7.

8. The composition according to claim 1, 2, 3, 4 or 5 wherein the fluorosurfactant is an anionic phosphate fluorosurfactant of the following structure: $(\text{RfCH}_2\text{CH}_2\text{O})_x\text{P}(\text{O})(\text{ONH}_4)_y$ where $\text{Rf} = \text{F}(\text{CF}_2\text{CF}_2)_z$, $x = 1$ or 2, $y = 2$ or 1, $x + y = 3$, and $z = 1$ to about 7 or mixtures thereof.

9. The composition according to claim 1, 2, 3, 4 or 5 wherein the fluorosurfactant is a mixture of two or more of the fluorosurfactants in claims 6, 7 and 8.

10. The composition according to claim 1, 2, 3, 4, 5, 6, 7, 8 or 9 wherein the alcohol is present in a range from about 40 % to about 90 % v/v.

11. The composition according to claim 1, 2, 3, 4, 5, 6, 7, 8 or 9 wherein the alcohol is ethanol present in an amount of at least 60% v/v, and wherein the composition is for use as an alcohol foam for personal hygiene.

12. The composition according to claim 1, 2, 3, 4, 5, 6, 7, 8 or 9 wherein the alcohol is a mixture of n-propanol and ethanol present in a combined amount of at least 60% v/v, and wherein the composition is for use as an alcohol foam for personal hygiene.

13. The composition according to claim 1, 2, 3, 4, 5, 6, 7, 8 or 9 wherein the alcohol is a mixture of isopropanol and ethanol present in a combined amount of at least 60% v/v, and wherein the composition is for use as as an alcohol foam for personal hygiene.

14. The composition according to claim 1, 2, 3, 4, 5, 6, 7, 8 or 9 wherein the alcohol is isopropanol present in an amount of at least 70% v/v, and wherein the composition is for use as an alcohol foam for personal hygiene.

15. The composition according to claim 1, 2, 3, 4, 5, 6, 7, 8 or 9 wherein alcohol is n-propanol present in an amount of at least 60% v/v, and wherein the composition is for use as an alcohol foam for personal hygiene.

16. The composition according to claim 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14 or 15 further including at least one additional surfactant for adjusting properties of the foam produced from the composition.
17. The composition according to claim 16 wherein the additional surfactant is selected from the group consisting of a poly(ethoxylated and/or propoxylated)alcohol, a poly(ethoxylated and/or propoxylated)ester, a derivative of a poly(ethoxylated and/or propoxylated)alcohol, a derivative of a poly(ethoxylated and/or propoxylated)ester, an alkyl alcohol, an alkenyl alcohol, an ester of a polyhydric alcohol, an ether of a polyhydric alcohol, an ester of a polyalkoxylated derivative of a polyhydric alcohol, an ether of a polyalkoxylated derivative of a polyhydric alcohol, a sorbitan fatty acid ester, a polyalkoxylated derivative of a sorbitan fatty acid ester, an alkylglucoside, a betaine, a sulfobetaine, an imidazoline derivative, an aminoacid derivative, a lecithin, a phosphatide, an amine oxide, a sulfoxide and mixtures thereof, present in an amount between about 0.10 % to about 5% weight percent.
18. The composition according to claim 17 wherein the betaine is cocamidopropyl betaine.
19. The composition according to claim 17 wherein the alkylglucoside is cocoglucoside.
20. The composition according to claim 17 wherein the

polyethoxylated fatty alcohol is polyethoxylated stearyl alcohol (21 moles ethylene oxide).

21. The composition according to claim 17 wherein the polyethoxylated fatty alcohol is polyethoxylated stearyl alcohol (2 moles ethylene oxide).
22. The composition according to claim 17 wherein the polyethoxylated fatty alcohol is a combination of polyethoxylated stearyl alcohol (21 moles ethylene oxide) and polyethoxylated stearyl alcohol (2 moles ethylene oxide).
23. The composition according to claim 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21 or 22 including a foam stabilizing agent present in an amount up to 5%.
24. The composition according to claim 23 wherein the foam stabilizing agent is selected from the group consisting of lactic acid esters of monoglycerides, cationic emulsifiers, triquaternized stearic phospholipid complex, hydroxystearamide propyltriamine salts, lactic acid monoglycerides, food emulsifiers such as glyceryl monostearate, propylene glycol monostearate, sodium stearyl lactylate, cetyl betaine, glycolether, n-propanol, butyleneglycol, silicone wax, an encapsulated oil, Microcapsule Mineral Oil, and combinations thereof.

25. The composition according to claim 23 wherein the foam stabilizing agent is selected from the group consisting of glycolether, n-propanol, butyleneglycol, and combinations thereof.

26. The composition according to any one of claims 1 to 25 including any one of a moisturizer, emollient and combinations thereof selected from the group consisting of lanolin, vinyl alcohol, polyvinyl pyrrolidone and polyols selected from the group consisting of glycerol, propylene glycol, butyleneglycol and sorbitol, or a fatty alcohol selected from the group consisting of cetyl alcohol, stearyl alcohol, lauryl alcohol, myristyl alcohol and palmityl alcohol, cetyl alcohol, cetareth 20, or an alkylglucoside and combinations thereof, present in an amount up to 5%.

27. The composition according to any one of claims 1 to 26 further comprising an acid or a base to adjust a pH of the composition to a pre-selected pH present in an amount from about 0.05 to about 0.5 % weight percent of the total composition.

28. The composition according to claim 27 wherein when an acid is used to adjust the pH the acid is selected from the group consisting of hydrochloric acid, citric acid and phosphoric acid, and wherein when a base is used to adjust the pH the base is sodium sesquicarbonate.

29. The composition according to any one of claims 1 to 28 including a preservative in an amount from about 0.01 to about 5 % weight percent of the total composition.

30. The composition according to any one of claims 1 to 29 including an antimicrobial agent.

31. The composition according to claim 30 wherein the antimicrobial agent is chlorhexidine gluconate present in an amount between about 0.50 % to about 4.0 % weight percent.

32. The composition according to claim 30 wherein the antimicrobial agent is didecyl dymethyl diamonium chloride present in an amount between about 0.05 % to about 5.0 % weight percent.

33. The composition according to claim 30 wherein the antimicrobial agent is selected from the group of a chlorhexidine salt, iodine, a complexed form of iodine, parachlorometaxylenol, triclosan, hexachlorophene, a phenol, a surfactant having a long chain hydrophobic group and a quaternary group, hydrogen peroxide, silver, a silver salt, silver oxide, and mixtures thereof.

34. The composition according to any one of claims 1 to 33 further comprising constituents selected from the group consisting of organic gums and colloids, lower alkanolamides of higher fatty acids, short chain diols and/or triols, fragrance, coloring matter, ultraviolet absorbers, solvents,

suspending agents, buffers, conditioning agents, antioxidants, bactericides and medicinally active ingredients, and combinations thereof.

35. The composition according to any one of claims 1 to 34 stored in an unpressurized dispenser having a dispenser pump for mixing the composition with air and dispensing foam therefrom.

36. The composition according to any one of claim 1 to 34 stored in a pressurized dispenser having a dispenser pump for mixing the composition with air or a propellant and dispensing foam therefrom, the composition including an aerosol propellant in an amount from about 3 to about 20 weight percent of the total composition.

37. The composition according to claim 36 wherein the aerosol propellant is selected from the group consisting of propane, carbon dioxide butane, dichloro difluoro methane, dichloro tetra fluoro ethane octafluorocyclo butane; 1,1,1,2-tetrafluoroethane; 1,1,1,2,3,3,3 heptafluoropropane, and 1,1,1,3,3,3,-hexafluoropropane.

38. The composition according to claim 36 or 37 including a corrosion inhibitor is selected from the group consisting of sorbic acid, benzoic acid, potassium sorbate and sodium benzoate, in an amount from about 0.1 to about 5 weight percent of the total composition.

39. A composition concentrate, comprising;

- a) an amphoteric, anionic or non-ionic fluorosurfactant present in an amount from about 0.01 to about 1.0 %;
- b) a foam stabilizing agent including at least from about 0.01 to about 12.0% of n-propanol;
- c) one of moisturizers, emollients and combinations thereof present in a range from about 0.05% to about 5.0%; and
- d) water.

40. The composition concentrate according to claim 39 wherein the amphoteric, anionic or non-ionic fluorosurfactant is anionic phosphate fluorosurfactant of the following structure: $(RfCH_2CH_2O)_xP(O)(ONH_4)_y$ where $Rf = F(CF_2CF_2)_z$, $x = 1$ or 2 , $y = 2$ or 1 , $x + y = 3$, and $z = 1$ to about 7 or an ethoxylated nonionic fluorosurfactant of the following structure: $RfCH_2CH_2O(CH_2CH_2O)_xH$ where $Rf = F(CF_2CF_2)_y$, $x = 0$ to about 15 and $y = 1$ to about 7 ; or Polytetrafluoroethylene Acetoxypopyl Betaine $CF_3CF_2(CF_2CF_2)_nCH_2CH_2(OAc)CH_2N^+(CH_3)_2CH_2COO^-$, where $n = 2-4$, or mixtures thereof

41. The composition concentrate according to claim 39 or 40 wherein the foam stabilizing agent includes 1,3 Butyleneglycol and 2-Butoxyethanol.

42. The composition concentrate according to claim 39, 40 or 41 wherein the moisturizers and emollients include cocoglucoside, glycerine and glyceryl oleate.

43. The composition concentrate according to claim 39, 40, 41 or 42 which is constituted as an alcohol disinfecting composition by adding

- a) an alcohol C₁₋₄, or mixtures thereof, present in an amount between about 60 to about 80% v/v of the total composition;
- c) water present in an amount to balance the total composition to 100% weight percent.

44. The composition concentrate according to claim 43 wherein the alcohol C₁₋₄ is selected from the group consisting of ethanol, n-propanol, isopropanol and combinations thereof.

45. The composition concentrate according to any one of claims 39 to 44 made by a process of mixing the constituents and then warming the concentrate to a temperature between about 30 to about 80 degrees Celsius prior to shipping the concentrate.

46. An alcohol disinfecting composition, comprising;

- a) a) an alcohol C₁₋₄, or mixtures thereof, present in an amount between about 60% to about 80% v/v of the total composition;
- b) a physiologically acceptable fluorosurfactant present in an amount from about 0.01% to about 2.0% weight percent of the total composition;
- c) a foam stabilizing agent present in an amount from about 0.01 to about 12.0 % weight percent;
- d) any one of moisturizers, emollients and combinations thereof present in an amount from about 0.05 to about 5.0 % weight percent; and

e) water in an amount to balance the total composition to 100% weight percent.

47. The alcohol disinfecting composition according to claim 46 including air, wherein when the disinfecting composition is mixed with air, the mixture of disinfecting composition and air forms a foam.

48. The alcohol disinfecting composition according to claims 46 or 47 wherein the fluorosurfactant is an amphoteric polytetrafluoroethylene acetoxypopyl betaine
 $(\text{CF}_3\text{CF}_2(\text{CF}_2\text{CF}_2)_n\text{CH}_2\text{CH}_2(\text{OAc})\text{CH}_2\text{N}^+(\text{CH}_3)_2\text{CH}_2\text{COO}^-)$ where $n = 2$ to 4 .

49. The alcohol disinfecting composition according to claim 46 or 47 wherein the fluorosurfactant is an ethoxylated nonionic fluorosurfactant of the following structure: $\text{RfCH}_2\text{CH}_2\text{O}(\text{CH}_2\text{CH}_2\text{O})_x\text{H}$ where $\text{Rf} = \text{F}(\text{CF}_2\text{CF}_2)_y$, $x = 0$ to about 15 and $y = 1$ to about 7.

50. The alcohol disinfecting composition according to claim 46 or 47 wherein the fluorosurfactant is an anionic phosphate fluorosurfactant of the following structure: $(\text{RfCH}_2\text{CH}_2\text{O})_x\text{P}(\text{O})(\text{ONH}_4)_y$ where $\text{Rf} = \text{F}(\text{CF}_2\text{CF}_2)_z$, $x = 1$ or 2 , $y = 2$ or 1 , $x + y = 3$, and $z = 1$ to about 7.

51. The alcohol disinfecting composition according to claim 46 or 47 wherein the fluorosurfactant is selected from the group consisting of fluorinated ethoxylates, glycerol esters, amine oxides, acetylenic alcohol

derivatives, carboxylates, phosphates, carbohydrate derivatives, sulfonates, betaines, esters, polyamides, silicones, hydrocarbon surfactants, and a mixture of two or more of the fluorosurfactants in claims 47, 48 and 49.

52. The alcohol disinfecting composition according to claim 46, 47, 48, 49, 50 or 51 wherein the foam stabilizing agent is selected from the group consisting of lactic acid esters of monoglycerides, cationic emulsifiers, triquaternized stearic phospholipid complex, hydroxystearamide propyltriamine salts, lactic acid monoglycerides, food emulsifiers such as glyceryl monostearate, propylene glycol monostearate, sodium stearyl lactylate, silicone wax, an encapsulated oil, Microcapsule Mineral Oil, butyleneglycol, butoxyethanol and or n-propanol and mixtures thereof.

53. The alcohol disinfecting composition according to any one of claims 46, 47, 48, 49, 50 or 51 wherein the foam stabilizing agent is selected from the group consisting of butoxyethanol, glycolether, n-propanol, butyleneglycol, and combinations thereof.

54. The alcohol disinfecting composition according to claims 46, 47, 48, 49, 50, 51, 52 or 53 including any one of a moisturizer, emollient and combinations thereof selected from the group consisting of lanolin, vinyl alcohol, polyvinyl pyrrolidone and polyols selected from the group consisting of glycerol, propylene glycol, butyleneglycol and sorbitol, or a fatty alcohol selected from the group consisting of cetyl alcohol, stearyl alcohol, lauryl

alcohol, myristyl alcohol and palmityl alcohol, cetyl alcohol, ceteareth 20, or an alkylglucoside and combinations thereof, present in an amount up to 5%.

55. The alcohol disinfecting composition according to claim 46, 47, 48, 49, 50, 51, 52 or 53 including a moisturizer and an emollient selected from the group consisting of glyceryl oleate, glycerine, cocoglucoside and combinations thereof.

56. The alcohol disinfecting composition according to claim 46, 47, 48, 49, 50, 51, 52, 53, 54 or 55 including an antimicrobial agent.

57. The alcohol disinfecting composition according to claim 56 wherein the antimicrobial agent is chlorhexidine gluconate present in an amount between about 0.50 % to about 4.0 % weight percent.

58. The alcohol disinfecting composition according to claim 56 wherein the antimicrobial agent is Didecyl Dymethyl Diamonium Chloride present in an amount between about 0.50 % to about 5.0 % weight percent.

59. The alcohol disinfecting composition according to claim 56 wherein the antimicrobial agent is selected from the group of a chlorhexidine salt, iodine, a complexed form of iodine, parachlorometaxlenol, triclosan, hexachlorophene, a phenol, a surfactant having a long chain hydrophobic group and a quaternary group, hydrogen peroxide, silver, a silver salt, silver oxide, and mixtures thereof.

60. The alcohol disinfecting composition according to any one of claims 46 to 59 including a preservative in an amount from about 0.01 to about 5 % weight percent of the total composition.

61. The alcohol disinfecting composition according to any one of claims 46 to 59 including a preservative in an amount from about 0.01 to about 5 % weight percent of the total composition.

62. The alcohol disinfecting composition according to any one of claims 46 to 61 further comprising an acid or a base to adjust a pH of the disinfecting composition to a pre-selected pH present in an amount from about 0.05 to about 0.5 % weight percent of the total composition.

63. The alcohol disinfecting composition according to claim 62 wherein the acid is selected from the group consisting of hydrochloric acid, citric acid and phosphoric acid, and the base is sodium sesquicarbonate.

64. The alcohol disinfecting composition according to any one of claims 46 to 63 stored in an unpressurized dispenser having a dispenser pump for mixing the disinfecting composition with air and dispensing foam therefrom.

65. The alcohol disinfecting composition according to any one of claims 46 to 63 stored in a pressurized dispenser having a dispenser pump for mixing the disinfecting composition with air or a propellant and dispensing foam

therefrom, the disinfecting composition including an aerosol propellant in an amount from about 3 to about 20 weight percent of the total composition.

66. The alcohol disinfecting composition according to claim 65 wherein the aerosol propellant is selected from the group consisting of propane, carbon dioxide, butane, dichloro difluoro methane, dichloro tetra fluoro ethane octafluorocyclo butane; 1,1,1,2-tetrafluoroethane; 1,1,1,2,3,3,3 heptafluoropropane, and 1,1,1,3,3,3,-hexafluoropropane.

67. The alcohol disinfecting composition according to claim 65 or 66 including a corrosion inhibitor is selected from the group consisting of sorbic acid, benzoic acid, potassium sorbate and sodium benzoate, in an amount from about 0.1 to about 5 weight percent of the total composition.

68. The alcohol disinfecting composition according to any one of claims 46 to 67 including any one of a moisturizer, emollient and combinations thereof selected from the group consisting of lanolin, vinyl alcohol, polyvinyl pyrrolidone and polyols selected from the group consisting of glycerol, propylene glycol, butyleneglycol and sorbitol, or a fatty alcohol selected from the group consisting of cetyl alcohol, stearyl alcohol, lauryl alcohol, myristyl alcohol and palmityl alcohol, cetyl alcohol, ceteareth 20, or an alkylglucoside and combinations thereof, present in an amount up to 5%.

69. The alcohol disinfecting composition according to any one of claims 46 to 68 further comprising other constituents or materials including organic

gums and colloids, lower alkanolamides of higher fatty acids, short chain diols and/or triols, fragrance, coloring matter, additional emollients, ultraviolet absorbers, solvents, emulsifiers, foam stabilizers or mixture of such stabilizers, suspending agents, buffers, conditioning agents, antioxidants, bactericides, medicinal active ingredient, and the like) that may be commonly added to aerosol compositions, toiletries, cosmetics and pharmaceuticals.

70. An alcohol disinfecting composition, comprising;

a) ethanol present in an amount between about 60% to 70% v/v of the total composition;

b) a physiologically acceptable fluorosurfactant or mixtures thereof in an amount from about 0.01% to about 2.0% weight percent of the total composition;

c) at least one secondary surfactant present in an amount from about 0.01 to about 10.0 % weight percent;

d) a foam stabilizing agent; and

e) water in an amount to balance the total composition to 100% weight percent.

71. The alcohol disinfecting composition according to claim 70 wherein the fluorosurfactant is an amphoteric polytetrafluoroethylene acetoxypyril betaine $(\text{CF}_3\text{CF}_2(\text{CF}_2\text{CF}_2)_n\text{CH}_2\text{CH}_2(\text{OAc})\text{CH}_2\text{N}^+(\text{CH}_3)_2\text{CH}_2\text{COO}^-)$ where $n = 2$ to 4.

72. The alcohol disinfecting composition according to claim 70 wherein the fluorosurfactant is an ethoxylated nonionic fluorosurfactant of the following structure: $R_fCH_2CH_2O(CH_2CH_2O)_xH$ where $R_f = F(CF_2CF_2)_y$, $x = 0$ to about 15 and $y = 1$ to about 7.

73. The alcohol disinfecting composition according to claim 70 wherein the fluorosurfactant is an anionic phosphate fluorosurfactant of the following structure: $(R_fCH_2CH_2O)_xP(O)(ONH_4)_y$ where $R_f = F(CF_2CF_2)_z$, $x = 1$ or 2, $y = 2$ or 1, $x + y = 3$, and $z = 1$ to about 7.

74. The alcohol disinfecting composition according to claim 70 wherein the fluorosurfactant is selected from the group consisting of fluorinated ethoxylates, glycerol esters, amine oxides, acetylenic alcohol derivatives, carboxylates, phosphates, carbohydrate derivatives, sulfonates, betaines, esters, polyamides, silicones, hydrocarbon surfactants, and mixtures of two or more of the fluorosurfactants in claims 71, 72 or 73.

75. The alcohol disinfecting composition according to claim 70, 71, 72, 73 or 74 wherein the secondary surfactant is at least one selected from the group consisting of a poly(ethoxylated and/or propoxylated)alcohol, a poly(ethoxylated and/or propoxylated)ester, a derivative of a poly(ethoxylated and/or propoxylated)alcohol, a derivative of a poly(ethoxylated and/or propoxylated)ester, an alkyl alcohol, an alkenyl alcohol, an ester of a polyhydric alcohol, an ether of a polyhydric alcohol, an ester of a polyalkoxylated derivative of a polyhydric alcohol, an ether of a

polyalkoxylated derivative of a polyhydric alcohol, a sorbitan fatty acid ester, a polyalkoxylated derivative of a sorbitan fatty acid ester, a betaine, a sulfobetaine, an imidazoline derivative, an aminoacid derivative, a lecithin, a phosphatide, an amine oxide, a sulfoxide and mixtures thereof, present in an amount between about 0.10 % to about 5% weight percent.

76. The alcohol disinfecting composition according to claim 70, 71, 72, 73 or 74 wherein the secondary surfactant is cocoamidopropyl betaine.

77. The alcohol disinfecting composition according to claim 70, 71, 72, 73, 74, 75 or 76 including an emulsifier or mixture of emulsifiers present in an amount between about 0.10 % to about 1.5% weight percent.

78. The alcohol disinfecting composition according to claim 77 wherein the emulsifier is a mixture of cetyl ethers and stearyl ethers of polyethylene oxide having an average of about 20 oxide units.

79. The alcohol disinfecting composition according to claim 70, 71, 72, 73, 74, 75, 76, 77 or 78 wherein the foam stabilizing agent is cetyl betaine.

80. The alcohol disinfecting composition according to claim 70, 71, 72, 73, 74, 75, 76, 77, 78 or 79 including an antimicrobial agent.

81. The alcohol disinfecting composition according to claim 80 wherein the antimicrobial agent is chlorhexidine gluconate present in an amount between about 0.50 % to about 4.0 % weight percent.
82. The alcohol disinfecting composition according to claim 80 wherein the antimicrobial agent is Didecyl Dymethyl Diamonium Chloride present in an amount between about 0.50 % to about 5.0 % weight percent.
83. The alcohol disinfecting composition according to claim 82 wherein the antimicrobial agent is selected from the group of a chlorhexidine salt, iodine, a complexed form of iodine, parachlorometaxylenol, triclosan, hexachlorophene, a phenol, a surfactant having a long chain hydrophobic group and a quaternary group, hydrogen peroxide, silver, a silver salt, silver oxide, and mixtures thereof.
84. The alcohol disinfecting composition according to claims 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82 or 83 including a preservative in an amount from about 0.01 to about 5 % weight percent of the total composition.
85. The alcohol disinfecting composition according to claims 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82 or 83 including a preservative in an amount from about 0.01 to about 5 % weight percent of the total composition.
86. The alcohol disinfecting composition according to any one of claims 70 to 85 further comprising an acid or a base to adjust a pH of the disinfecting

composition to a pre-selected pH present in an amount from about 0.05 to about 0.5 % weight percent of the total composition.

87. The alcohol disinfecting composition according to claim 86 wherein the acid is selected from the group consisting of hydrochloric acid, citric acid and phosphoric acid, and the base is sodium sesquicarbonate.

88. The alcohol disinfecting composition according to any one of claims 70 to 87 stored in an unpressurized dispenser having a dispenser pump for mixing the disinfecting composition with air and dispensing foam therefrom.

89. The alcohol disinfecting composition according to any one of claims 70 to 87 stored in a pressurized dispenser having a dispenser pump for mixing the disinfecting composition with air or a propellant and dispensing foam therefrom, the disinfecting composition including an aerosol propellant in an amount from about 3 to about 20 weight percent of the total composition.

90. The alcohol disinfecting composition according to claim 89 wherein the aerosol propellant is selected from the group consisting of propane, carbon dioxide, butane, dichloro difluoro methane, dichloro tetra fluoro ethane octafluorocyclo butane; 1,1,1,2-tetrafluoroethane; 1,1,1,2,3,3,3 heptafluoropropane, and 1,1,1,3,3,3,-hexafluoropropane.

91. The alcohol disinfecting composition according to claim 89 or 90 including a corrosion inhibitor is selected from the group consisting of sorbic

acid, benzoic acid, potassium sorbate and sodium benzoate, in an amount from about 0.1 to about 5 weight percent of the total composition.

92. The alcohol disinfecting composition according to any one of claims 70 to 91 including a moisturizer and/or emollient selected from the group consisting of lanolin, alkylglucoside, vinyl alcohol, polyvinyl pyrrolidone and polyols selected from the group consisting of glycerol, propylene glycol, glyceryl oleate and sorbitol, or a fatty alcohol selected from the group consisting of cetyl alcohol, stearyl alcohol, lauryl alcohol, myristyl alcohol and palmityl alcohol, cetyl alcohol, ceteareth 20, and combinations thereof, present in an amount up to 5% weight percent.

93. The alcohol disinfecting composition according to any one of claims 70 to 92 further comprising other constituents or materials including organic gums and colloids, lower alkanolamides of higher fatty acids, short chain diols and/or triols, fragrance, coloring matter, additional emollients, ultraviolet absorbers, solvents, emulsifiers, foam stabilizers or mixture of such stabilizers, suspending agents, buffers, conditioning agents, antioxidants, bactericides, medicinal active ingredient, and the like) that may be commonly added to aerosol compositions, toiletries, cosmetics and pharmaceuticals.

94. The alcohol disinfecting composition concentrate according to claim 43 or 44 made by a process of mixing the constituents and then shipping the concentrate, and just prior to reconstituting the concentrate with alcohol, warming the concentrate to a temperature between about 30 to 80 degrees.

95. An alcohol gel-like composition, comprising;
- a) an alcohol C₁₋₄, or mixtures thereof, present in an amount greater than about 40% v/v of the total composition;
 - b) at least one nonionic surfactant selected from the group consisting of a poly(ethoxylated and/or propoxylated)alcohol, a poly(ethoxylated and/or propoxylated)ester, a derivative of a poly(ethoxylated and/or propoxylated)alcohol, a derivative of a poly(ethoxylated and/or propoxylated)ester, an alkyl alcohol, an alkenyl alcohol, an ester of a polyhydric alcohol, an ether of a polyhydric alcohol, an ester of a polyalkoxylated derivative of a polyhydric alcohol, an ether of a polyalkoxylated derivative of a polyhydric alcohol, a sorbitan fatty acid ester, a polyalkoxylated derivative of a sorbitan fatty acid ester, and mixtures thereof, present in an amount between about 0.10 % to about 5% weight percent
 - c) an emulsifier present in an amount between about 0.10 % to about 3.0% weight percent ; and
 - d) water in an amount sufficient to form a stable gel composition with a viscosity less than 4,000 cps.
96. The alcohol gel-like composition according to claim 95 including an effective fluorinated surface active agent for wetting.
97. The alcohol gel-like composition according to claim 96 wherein the effective fluorinated surface active agent is a fluorosurfactant selected from the group consisting of fluorinated ethoxylates, glycerol esters, amine oxides, acetylenic alcohol derivatives, carboxylates, phosphates, carbohydrate

derivatives, sulfonates, betaines, esters, polyamides, silicones, hydrocarbon surfactants.

98. The alcohol gel-like composition according to claim 96 wherein the effective fluorinated surface active agent is an amphoteric polytetrafluoroethylene acetoxypentyl betaine

$(\text{CF}_3\text{CF}_2(\text{CF}_2\text{CF}_2)_n\text{CH}_2\text{CH}_2(\text{OAc})\text{CH}_2\text{N}^+(\text{CH}_3)_2\text{CH}_2\text{COO}^-)$ where $n = 2$ to 4 , present in an amount from about 0.001 % weight percent to about 2.5.

99. The alcohol gel-like composition according to claim 96 wherein the effective fluorinated surface active agent is an ethoxylated nonionic fluorosurfactant of the following structure: $\text{RfCH}_2\text{CH}_2\text{O}(\text{CH}_2\text{CH}_2\text{O})_x\text{H}$ where $\text{Rf} = \text{F}(\text{CF}_2\text{CF}_2)_y$, $x = 0$ to about 15 and $y = 1$ to about 7 present in an amount from about 0.001 % weight percent to about 2.5.

100. The alcohol gel-like composition according to claim 96 wherein the effective fluorinated surface active agent is an anionic phosphate fluorosurfactant of the following structure: $(\text{RfCH}_2\text{CH}_2\text{O})_x\text{P}(\text{O})(\text{ONH}_4)_y$ where $\text{Rf} = \text{F}(\text{CF}_2\text{CF}_2)_z$, $x = 1$ or 2 , $y = 2$ or 1 , $x + y = 3$, and $z = 1$ to about 7 present in an amount from about 0.001 % weight percent to about 2.5.

101. The alcohol gel-like composition according to claim 96 wherein the effective fluorinated surface active agent is a mixture of two or more of the fluorosurfactants in claims 96, 97 or 98 present in an amount from about 0.001 % weight percent to about 2.5.

102. The alcohol gel-like composition according to any one of claims 95 to 101 wherein the alcohol C₁₋₄ is an aliphatic alcohol selected from the group consisting of methanol, ethanol, 2-propanol and butanol.
103. The alcohol gel-like composition according to any one of claims 95 to 101 wherein the at least one nonionic surfactant is a polyethoxylated fatty alcohol which is polyethoxylated stearyl alcohol (21 moles ethylene oxide) in an amount of 0.1% to about 5% so that the final disinfecting composition has a final viscosity of less than 4,000cps.
104. The alcohol gel-like composition according to any one of claims 95 to 101 wherein the at least one nonionic surfactant is a polyethoxylated fatty alcohol which is polyethoxylated stearyl alcohol (2 moles ethylene oxide) present in an amount of 0.1% to about 5% so that the final disinfecting composition has a final viscosity of less than 4,000cps.
105. The alcohol gel-like composition according to any one of claims 95 to 101 wherein the at least one nonionic surfactant is a combination of polyethoxylated stearyl alcohol (21 moles ethylene oxide) and polyethoxylated stearyl alcohol (2 moles ethylene oxide) present in an amount of 0.1% to about 5% so that the final disinfecting composition has a final viscosity of less than 4,000cps.

106. The alcohol gel-like composition according to any one of claims 95 to 105 including an emulsifier moisturizer and emollient ingredient or mixture of ingredients present in an amount from about 0.10 to about 2.0 % weight percent so that the final disinfecting composition has a final viscosity of less than 4,000cps.

107. The alcohol gel-like composition according to claim 106 wherein the emulsifier moisturizer and emollient ingredient or mixture of ingredients is comprised of a mixture of cetearyl alcohol and ceteareth 20.

108. The alcohol gel-like composition according to any one of claims 95 to 107 including an antimicrobial agent.

109. The alcohol gel-like composition according to claim 108 wherein the antimicrobial agent is chlorhexidine gluconate present in an amount between about 0.50 % to about 4.0 % weight percent.

110. The alcohol gel-like composition according to claim 108 wherein the antimicrobial agent is selected from the group of a chlorhexidine salt, iodine, a complexed form of iodine, parachlorometaxyleneol, triclosan, hexachlorophene, a phenol, a surfactant having a long chain hydrophobic group and a quaternary group, hydrogen peroxide, silver, a silver salt, silver oxide, and mixtures thereof.

111. The alcohol gel-like composition according to any one of claims 95 to 110 including a preservative in an amount from about 0.01 to about 5 % weight percent of the total composition.

112. The alcohol gel-like composition according to any one of claims 95 to 111 further comprising an acid or a base to adjust a pH of the composition to a pre-selected pH present in an amount from about 0.05 to about 0.5 % weight percent of the total composition.

113. The alcohol gel-like composition according to claim 112 wherein the acid is selected from the group consisting of hydrochloric acid, citric acid and phosphoric acid, and the base is sodium sesquicarbonate.

114. The alcohol gel-like composition according to any one of claims 95 to 113 stored in an unpressurized dispenser having a dispenser pump for dispensing the composition therefrom in a gel-like form.

115. The alcohol gel-like composition according to any one of claims 95 to 113 stored in a pressurized dispenser having a dispenser pump for dispensing the disinfecting composition therefrom, including an aerosol propellant in an amount from about 3 to about 20 weight percent of the total composition.

116. The alcohol gel-like composition according to claim 115 wherein the aerosol propellant is selected from the group consisting of propane, carbon

dioxide butane, dichloro difluoro methane, dichloro tetra fluoro ethane and octafluorocyclo butane; 1,1,1,2-tetrafluoroethane; 1,1,1,2,3,3,3 heptafluoropropane, and 1,1,1,3,3,3,-hexafluoropropane.

117. The alcohol gel-like composition according to claim 115 or 116 including a corrosion inhibitor is selected from the group consisting of sorbic acid, benzoic acid, potassium sorbate and sodium benzoate, in an amount from about 0.1 to about 5 weight percent of the total composition.