

COMPOSITIONS AND METHODS FOR MINIMIZING PROTEIN LOSS AT LOW PROTEIN CONCENTRATIONS

CROSS REFERENCE OF RELATED APPLICATION

[0001] This application claims the benefit of U. S. Provisional Application No. 62/926,089 filed October. 25, 2019, which is incorporated in its entirety by reference herein.

SUBMISSION OF SEQUENCE LISTING ON ASCII TEXT FILE

[0002] The content of the following submission on ASCII text file is incorporated herein by reference in its entirety: a computer readable form (CRF) of the Sequence Listing (file name: A-2429-WO-PCT_ST25, date created: October 23, 2020 size: 451,608 bytes).

FIELD OF THE INVENTION

[0001] Inventions disclosed herein generally relate to the field of compositions comprising proteins, in particular, pharmaceutical compositions comprising therapeutic proteins at low protein concentrations. Inventions disclosed herein also relate to methods of administering the composition to a subject in need thereof.

BACKGROUND OF THE INVENTION

[0002] Therapeutic proteins are an important class of therapeutics for treating patients. Protein molecules are surface active and subject to potential adsorption to solid surfaces. Therapeutic proteins in pharmaceutical compositions could be adsorbed to solid surfaces that the proteins come into contact with (e.g., the surfaces of a container containing a pharmaceutical composition), which could lead to protein loss during storage and use. Generally, the concentration of therapeutic proteins in those compositions is high (e.g., 1 mg/mL or higher) such that protein adsorption to solid surfaces does not result in insufficient amount of drug available for administration to patients. However, when the concentration of proteins in compositions is low (e.g., less than 0.1 mg/mL, such as when a composition is diluted before administration to patients), the risk of protein loss can be more pronounced, which could potentially lead to insufficient amount of drug available for patient administration.

[0003] Surfactants are generally used in pharmaceutical compositions comprising therapeutic proteins, e.g., to prevent protein aggregation and stabilize proteins. It is unclear whether surfactants can be used to effectively prevent protein loss due to surface adsorption when proteins are present at low concentrations in pharmaceutical compositions (e.g., 0.1 mg/mL or less).

[0004] There is a need for protein compositions and methods that minimize protein loss due to adsorption to solid surfaces, especially when the compositions comprising protein at low protein concentrations.

SUMMARY OF THE INVENTION

[0005] Disclosed herein are compositions comprising proteins at low protein concentrations as well as methods of administering the compositions to a subject in need thereof. The compositions and methods disclosed herein have the advantage of minimizing or eliminating protein loss due to protein adsorption to solid surfaces and ensure accurate dosing of therapeutic proteins to patients.

[0006] In certain embodiments, disclosed herein is an aqueous composition comprising a protein and a surfactant, wherein the protein is present in the composition at a concentration of between about 0.001 µg/ml and about 100 µg/ml, and the surfactant is present in the composition at a concentration of at least about 0.25 x of the critical micelle concentration (CMC) of the surfactant.

[0007] In certain embodiments, the protein is a bispecific antibody construct comprising a first binding domain that binds to a target cell surface antigen, a second binding domain that binds to human CD3 on the surface of a T cell, and optionally, a third domain comprising, in an amino to carboxyl order, hinge-CH2 domain-CH3 domain-linker-hinge-CH2 domain-CH3 domain. In certain embodiments, the second binding domain comprises a polypeptide having the sequence of SEQ ID NO: 201. In certain embodiments, the bispecific antibody construct is present at a concentration of between about 0.001 µg/ml and about 50 µg/ml, or between about 0.01 µg/ml to about 50 µg/ml, or between 0.1 µg/ml to about 50 µg/ml, or 0.1 µg/ml to about 10 µg/ml, or 1 µg/ml to about 10 µg/ml.

[0008] In certain embodiments, the surfactant is a polysorbate, a poloxamer or triton x-100. In certain embodiments, the surfactant is polysorbate 80, polysorbate 60, polysorbate 40, polysorbate 20, or Triton X-100. In certain embodiments, the surfactant is poloxamer 188 or

poloxamer 407. In certain embodiments, the surfactant is present at a concentration of between about 0.25x and about 20x of the CMC, or between about 0.25x and about 10x of the CMC of the surfactant.

[0009] In certain embodiments, wherein the composition further comprising a salt, an amino acid, a saccharide or saccharide derivative, or combinations thereof. In certain embodiments, the salt is NaCl. In certain embodiments, the saccharide or saccharide derivative is a monosaccharide, a disaccharide, a cyclic polysaccharide or a sugar alcohol. In certain embodiments, the saccharide is sucrose, trehalose, mannitol or sorbitol. In certain embodiments, the amino acid is lysine.

[0010] In certain embodiments, wherein the pH of the composition is between about 3.5 and about 7.5. In certain embodiments, the pH of the composition is between about 4.2 and about 7.0.

[0011] In certain embodiments, the composition further comprises a buffer or a preservative. In certain embodiments, the buffer is an acetate buffer, a glutamate buffer, a citrate buffer, a succinate buffer, a tartrate buffer, a fumarate buffer, a maleate buffer, a histidine buffer, or phosphate buffer.

[0012] In certain embodiments, wherein each of the first and second binding domains of the bispecific antibody construct comprises a VH region and a VL region. In certain embodiments, the bispecific antibody construct is a single chain antibody construct. In certain embodiments, the bispecific antibody construct comprises a polypeptide having the amino acid sequence selected from SEQ ID NO: 11, SEQ ID NO: 13, SEQ ID NO: 15, SEQ ID NO: 24, SEQ ID NO: 25, SEQ ID NO: 47, SEQ ID NO: 48, SEQ ID NO: 66, SEQ ID NO: 67, SEQ ID NO: 76, SEQ ID NO: 78, SEQ ID NO: 87, SEQ ID NO: 88, SEQ ID NO: 108, SEQ ID NO: 109, SEQ ID NO: 110, SEQ ID NO: 165, SEQ ID NO: 166, SEQ ID NO: 167, SEQ ID NO: 176, SEQ ID NO: 178, and SEQ ID NO: 192. In certain embodiments, the bispecific antibody construct comprises a polypeptide comprising the amino acid sequence of SEQ ID NO: 13, SEQ ID NO: 15, SEQ ID NO: 25, SEQ ID NO: 48, SEQ ID NO: 67, SEQ ID NO: 78, SEQ ID NO: 88, SEQ ID NO: 109, SEQ ID NO: 110, SEQ ID NO: 165, SEQ ID NO: 166, SEQ ID NO: 167, SEQ ID NO: 178 or SEQ ID NO: 192.

[0013] In certain embodiments, the composition is contained in a plastic container such as an IV bag or IV tube. In certain embodiments, the container is made of a material comprising polyolefin, polyvinyl chloride (PVC), ethyl vinyl acetate (EVA), or polyurethane. In certain

embodiments, the container is made of a material comprising PVC and wherein the PVC is substantially free of di-2-ethylhexyl phthalate(DEHP) or tri-2-ethylhexyltrimellitate (TOTM)

[0014] In certain embodiments, disclosed herein is a pharmaceutical preparation comprising an aqueous pharmaceutical composition contained inside a container, wherein the aqueous pharmaceutical composition comprising: a bispecific antibody construct at a concentration of between about 0.001 µg/ml and about 100 µg/ml, and a surfactant at a concentration of at least about 0.25 x of CMC of the surfactant, wherein the surfactant has an HLB value of at least 20. In certain embodiments, the surfactant is poloxamer 188 or poloxamer 407. In certain embodiments, the aqueous pharmaceutical composition comprises the bispecific antibody construct at a concentration of between about 0.001 µg/ml and about 50 µg/ml. In certain embodiments, the aqueous pharmaceutical composition comprises the surfactant is at a concentration of between about 0.25x and about 20x of the CMC or between about 0.25x and about 10x of the CMC, of the surfactant.

[0015] In certain embodiments, the aqueous pharmaceutical composition further comprising a salt, a buffer, an amino acid, a saccharide or saccharide derivative, or combinations thereof. In certain embodiments, the aqueous pharmaceutical composition has a pH of between about 4.2 and about 7.0.

[0016] In certain embodiments, the container is made of a material comprising polyolefin, PVC, EVA or polyurethane (e.g., polyester and polyether).

[0017] In certain embodiments, the bispecific antibody construct comprises a polypeptide having the amino acid sequences selected from SEQ ID NO: 11, SEQ ID NO: 13, SEQ ID NO: 15, SEQ ID NO: 24, SEQ ID NO: 25, SEQ ID NO: 47, SEQ ID NO: 48, SEQ ID NO: 66, SEQ ID NO: 67, SEQ ID NO: 76, SEQ ID NO: 78, SEQ ID NO: 87, SEQ ID NO: 88, SEQ ID NO: 108, SEQ ID NO: 109, SEQ ID NO: 110, SEQ ID NO: 165, SEQ ID NO: 166, SEQ ID NO: 167, SEQ ID NO: 176, SEQ ID NO: 178, and SEQ ID NO: 192. In certain embodiments, the bispecific antibody construct comprises a polypeptide comprising the amino acid sequence of SEQ ID NO: 13, SEQ ID NO: 15, SEQ ID NO: 25, SEQ ID NO: 48, SEQ ID NO: 67, SEQ ID NO: 78, SEQ ID NO: 88, SEQ ID NO: 109, SEQ ID NO: 110, SEQ ID NO: 165, SEQ ID NO: 166, SEQ ID NO: 167, SEQ ID NO: 178 or SEQ ID NO: 192.

[0018] Also disclosed herein is a method of administering a bispecific antibody construct to a patient comprising: preparing an aqueous pharmaceutical composition in a container, wherein the aqueous pharmaceutical composition comprises the bispecific antibody construct at a concentration of between about 0.001 µg/ml and about 100 µg/ml and a surfactant at a

concentration of at least about 0.25 x of CMC of the surfactant, and administering the aqueous pharmaceutical composition to the patient, wherein the bispecific antibody construct comprises a polypeptide having the amino acid sequence selected from SEQ ID NO: 11, SEQ ID NO: 13, SEQ ID NO: 15, SEQ ID NO: 24, SEQ ID NO: 25, SEQ ID NO: 47, SEQ ID NO: 48, SEQ ID NO: 66, SEQ ID NO: 67, SEQ ID NO: 76, SEQ ID NO: 78, SEQ ID NO: 87, SEQ ID NO: 88, SEQ ID NO: 108, SEQ ID NO: 109, SEQ ID NO: 110, SEQ ID NO: 165, SEQ ID NO: 166, SEQ ID NO: 167, SEQ ID NO: 176, SEQ ID NO: 178, and SEQ ID NO: 192. In certain embodiments, the bispecific antibody construct comprises a polypeptide comprising the amino acid sequence of SEQ ID NO: 13, SEQ ID NO: 15, SEQ ID NO: 25, SEQ ID NO: 48, SEQ ID NO: 67, SEQ ID NO: 78, SEQ ID NO: 88, SEQ ID NO: 109, SEQ ID NO: 110, SEQ ID NO: 165, SEQ ID NO: 166, SEQ ID NO: 167, SEQ ID NO: 178 or SEQ ID NO: 192. In certain embodiments, the aqueous pharmaceutical composition comprises the bispecific antibody construct is at a concentration of between about 0.001 µg/ml and about 50 µg/ml.

[0019] In certain embodiments, the aqueous pharmaceutical composition comprises the surfactant is at a concentration of between about 0.25x and about 20x of the CMC or between about 0.25x and about 10x of the CMC, of the surfactant. In certain embodiments, the surfactant is polysorbate 80, polysorbate 60, polysorbate 40, polysorbate 20, poloxamer 188, poloxamer 407, or Triton X-100.

[0020] In certain embodiments, the aqueous pharmaceutical composition further comprising one or more selected from a salt, a buffer, an amino acid, a saccharide or a saccharide derivative, and a preservative. In certain embodiments, the aqueous pharmaceutical composition has a pH of between about 4.2 and about 7.0.

[0021] In certain embodiments, the container is made of a material comprising polyolefin, PVC, EVA, polyurethane. In certain embodiments, the surfactant is polysorbate 80, polysorbate 60, polysorbate 40, or polysorbate 20, and wherein the container is made of a material comprising PVC that is substantially free of DEHP or TOTM.

[0022] In certain embodiments, the aqueous pharmaceutical composition is prepared by diluting a first composition comprising the bispecific antibody construct with a suitable aqueous solution. In certain embodiments, the first composition is a liquid composition comprising the bispecific antibody construct. In certain embodiments, the first composition is a liquid composition reconstituted from a lyophilized composition comprising the bispecific antibody construct. In certain embodiments, the suitable solution comprises the surfactant at a concentration of at least about 0.25 x of CMC of the surfactant. In certain embodiments, the

aqueous pharmaceutical composition is prepared by adding the suitable aqueous solution into the container followed by adding an appropriate amount of the first composition into the container.

[0023] In certain embodiments, the patient is a cancer patient. In certain embodiments, the administration is IV administration.

BRIEF DESCRIPTION OF THE DRAWINGS

[0024] Figure 1 shows a diagram of the assay for measuring protein binding to a solid surface.

[0025] Figure 2 shows protein binding to a solid surface in the absence of surfactants.

[0026] Figure 3 shows different the addition of various surfactants prevented protein binding to solid surfaces.

[0027] Figure 4 shows that adding surfactants to the solid surface before adding protein prevents protein binding to surfaces more efficiently.

[0028] Figure 5 shows the impacts of different surfactants at the same concentration on the leaching of di-2-ethylhexyl phthalate (DEHP) from DEHP-containing PVC.

[0029] Figure 6 shows the impacts of different surfactants at same folds of CMC on leaching of DEHP from DEHP-containing PVC.

DETAILED DESCRIPTION

[0030] Inventions disclosed herein are based on the surprising finding that surfactants, when used at concentrations lower than their critical micelle concentration, can stabilize proteins at low concentrations in liquid compositions and effectively prevent protein loss due to adsorption to solid surfaces.

[0031] In some embodiments, disclosed herein is an aqueous composition comprising a protein and a surfactant, wherein the protein is present in the composition at a concentration of between about 0.001 µg/ml and about 100 µg/ml, and the surfactant is present in the composition at a concentration of at least about 0.25x of the critical micelle concentration (CMC) of the surfactant.

[0032] Surfactants that can be used in the composition can be any surfactant typically used in pharmaceutical compositions. In some embodiments, the surfactant is a non-ionic surfactant. In some embodiments, the surfactant is a polysorbate such as polysorbate 20, polysorbate 40, polysorbate 60, polysorbate 65, polysorbate 80 or polysorbate 85. In some

embodiments, the surfactant is polysorbate 20, in other embodiments, the surfactant is polysorbate 80. In some embodiments, the surfactant is a poloxamer such as poloxamer 124, poloxamer 188, poloxamer 237, poloxamer 338, and poloxamer 407. In some embodiments, the surfactant is Triton X-100. Various surfactants are available commercially (e.g., Tween 20, Tween 80, Pluronic F68, and Pluronic F127 etc.).

[0033] The surfactant can be present in the composition at a concentration that is at least about 0.25 times (0.25x) the critical micelle concentration (CMC) of that surfactant. In some embodiments, the surfactant is present in the composition at a concentration that is between about 0.25x and about 20x of the CMC of the surfactant. In some embodiments, the surfactant is present in the composition at a concentration that is between about 0.25x and about 15x, or between about 0.25x and about 10x, or between about 0.25x and about 8x, or between about 0.25x and about 6x, or between about 0.25x and about 4, or between about 0.25x and about 2x, or between about 0.25x and about 1x of the CMC of the surfactant.

[0034] In some embodiments, the surfactant is present in the composition at a concentration that is between about 0.5x and about 20x of the CMC of the surfactant. In some embodiments, the surfactant is present in the composition at a concentration that is between about 0.5x and about 15x, or between about 0.5x and about 10x, or between about 0.5x and about 8x, or between about 0.5x and about 6x, or between about 0.5x and about 4x, or between about 0.5x and about 2x, or between about 0.5x and about 1x of the CMC of the surfactant.

[0035] In some embodiments, the surfactant is present in the composition at a concentration of about 0.25x, about 0.5x, about 1x, about 2x, about 3x, about 4x, about 5x, about 6x, about 7x, about 8x, about 9x, about 10x, about 12x, about 14x, about 16x, about 18x, or about 20x of the CMC of the surfactant. In some embodiments, the surfactant is present in the composition at a concentration of about 1.25x, about 2.5x, about 3.5x, about 4.5x, about 5.5x, about 6.5x, about 7.5x, about 8.5x or about 9.5x of the CMC of the surfactant.

[0036] As used herein, the term “about,” when used to modify a particular value or a range, is understood to mean that there can be variations in a given value or range, including within 20 percent, e.g., within 10 percent, 5 percent, 4 percent, 3 percent, 2 percent, or 1 percent of the stated value or range.

[0037] Critical micelle concentration refers to the concentration of a surfactant above which micelles form, it is a property of a surfactant. The CMC value of a surfactant can be measured by experimental methods well known in the art such as fluorometry, surface tension, conductometry and dynamic light scattering. See e.g., Norman Scholz, Thomas Behnke, Ute

Resch-Genger. *Journal of Fluorescence* 28:465–476 (2018) and Önder Topel, Burçin Acar Çakır, Leyla Budama, Numan Hoda. *Journal of Molecular Liquids* 177 40–43 (2013). The CMC value of a surfactant can also be measured automatically using devices such as Attention® Sigma 700 or 701.

[0038] In some embodiments, the CMC value of each of the surfactants is listed in table 1 below.

[0039] Table 1. CMC of commonly used surfactants

Surfactant	CMC* (w/v%)
Polysorbate 20	0.007
Polysorbate 80	0.002
Poloxamer 188	0.4
Poloxamer 407	0.004
Triton X-100	0.014

* For the CMC values listed in the table, see e.g., le Maire M, Champeil P, Moller JV. 2000.

Interaction of membrane proteins and lipids with solubilizing detergents. *Biochim Biophys Acta* 1508:86–111; Suksiriworapong J, Rungvimolsin T, A-gomol A, Junyaprasert VB, Chantasart D. Development and characterization of lyophilized diazepam-loaded polymeric micelles. 2014. *AAPS PharmSciTech*. 15(1):52–64; https://www.sigmaaldrich.com/content/dam/sigma-aldrich/docs/Sigma/Product_Information_Sheet/1/t8532pis.pdf

[0040] The protein can be present in the composition at a concentration that is between about 0.001 µg/ml and about 100 µg/ml. In some embodiments, the protein is present in the composition at a concentration that is between about 0.001 µg/ml and about 90 µg/ml, or between about 0.001 µg/ml and about 80 µg/ml, or between about 0.001 µg/ml and about 70 µg/ml, or between about 0.001 µg/ml and about 60 µg/ml, or between about 0.001 µg/ml and about 50 µg/ml, or between about 0.001 µg/ml and about 40 µg/ml, between about 0.001 µg/ml and about 30 µg/ml, or between about 0.001 µg/ml and about 20 µg/ml, or between about 0.001 µg/ml and about 10 µg/ml, or between about 0.001 µg/ml and about 5 µg/ml, or between about 0.001 µg/ml and about 1 µg/ml, or between about 0.001 µg/ml and about 0.01 µg/ml.

[0041] In some embodiments, the protein is present in the composition at a concentration that is between about 0.01 µg/ml and about 100 µg/ml, or between about 0.01 µg/ml and about 80 µg/ml, or between about 0.01 µg/ml and about 70 µg/ml, or between about 0.01 µg/ml and about 60 µg/ml, or between about 0.01 µg/ml and about 50 µg/ml, or between about 0.01 µg/ml

and about 40 µg/ml, between about 0.01 µg/ml and about 30 µg/ml, or between about 0.01 µg/ml and about 20 µg/ml, or between about 0.01 µg/ml and about 10 µg/ml, or between about 0.01 µg/ml and about 5 µg/ml, or between about 0.01 µg/ml and about 1 µg/ml, or between about 0.01 µg/ml and about 0.1 µg/ml, or between about 0.1 µg/ml and about 1 µg/ml, or between about 0.1 µg/ml and about 5 µg/ml.

[0042] In some embodiments, the protein is present in the composition at a concentration of about 0.001 µg/ml, about 0.005 µg/ml, about 0.01 µg/ml, about 0.05 µg/ml, about 0.1 µg/ml, about 1 µg/ml, about 4 µg/ml, about 8 µg/ml, about 10 µg/ml, about 15 µg/ml, about 20 µg/ml, about 25 µg/ml, about 30 µg/ml, about 35 µg/ml, about 40 µg/ml, about 45 µg/ml, about 50 µg/ml, about 55 µg/ml, about 60 µg/ml, about 65 µg/ml, about 70 µg/ml, about 75 µg/ml, about 80 µg/ml, about 85 µg/ml, about 90 µg/ml, about 95 µg/ml, or about 100 µg/ml.

[0043] In the composition disclosed herein, any of the concentration or concentration range for the protein can be combined with any of the concentration or concentration range for the surfactant.

[0044] Any protein can be the protein in the composition. In some embodiments, the protein in the composition is a therapeutic protein such as an antigen binding protein or a fusion protein.

[0045] As used herein, the term "antigen binding protein" refers to a protein that specifically binds to one or more target antigens. An antigen binding protein includes, but not limited to an antibody (e.g., a monoclonal antibody). An antigen binding protein typically comprises an antigen-binding fragment that specifically binds to an antigen and, optionally, a scaffold or framework portion that allows the antigen-binding fragment to adopt a conformation that promotes binding of the antigen binding protein to the antigen. An "antigen binding fragment" refers to a portion of an antibody that lacks at least some of the amino acids present in a full-length heavy chain and/or light chain, but which is still capable of specifically binding to an antigen. An antigen-binding fragment includes, but is not limited to, a single-chain variable fragment (scFv), a nanobody (e.g. VH domain of camelid heavy chain antibodies; VHH fragment, see Cortez-Retamozo et al., Cancer Research, Vol.64:2853-57, 2004), a Fab fragment, a Fab' fragment, a F(ab')₂ fragment, a Fv fragment, a Fd fragment, and a complementarity determining region (CDR) fragment, and can be derived from any mammalian source, such as human, mouse, rat, rabbit, or camelid. Antigen-binding fragments may compete for binding of a target antigen with an intact antibody and the fragments may be produced by the modification of

intact antibodies (e.g. enzymatic or chemical cleavage) or synthesized de novo using recombinant DNA technologies or peptide synthesis known in the art.

[0046] In some embodiments, the protein is an antigen binding protein that is bispecific. As used herein, the term “bispecific” refers to an antigen binding protein capable of specifically binding to two different antigens or targets or epitopes. As used herein, an “epitope” refers to any determinant capable of being specifically bound by an antigen binding protein, such as an antibody or fragment thereof. In some embodiments, the bispecific antigen binding protein comprises a first domain specifically binds to one antigen or target and a second domain specifically binds to another antigen or target. In some embodiments, the first domain of the bispecific antigen binding protein specifically binds to a target cell surface antigen and the second binding domain of the bispecific antigen binding protein specifically binds to the human CD3, a subunit of the T cell receptor complex on T cells. In some preferred embodiments, the bispecific antigen binding protein is a bispecific T cell engager (BiTE®) antibody construct as described in, e.g., WO2008119567 and WO2017134140.

[0047] As used herein, the term “antibody construct” refers to a molecule in which the structure and/or function is/are based on the structure and/or function of an antibody, e.g., of a full-length or whole immunoglobulin molecule and/or is/are drawn from the variable heavy chain (VH) and/or variable light chain (VL) domains of an antibody or fragment thereof. An antibody construct is hence capable of binding to its specific target or antigen. Antibody construct also includes modified fragments of antibodies, such as scFv, di-scFv or bi(s)-scFv, scFv-Fc, scFv-zipper, scFab, Fab₂, Fab₃, diabodies, single chain diabodies, tandem diabodies (Tandab’s), tandem di-scFv, tandem tri-scFv, “multibodies” such as triabodies or tetrabodies, and single domain antibodies such as nanobodies or single variable domain antibodies comprising merely one variable domain, which might be VHH, VH or VL, that specifically bind an antigen or epitope independently of other V regions or domains.

[0048] In some embodiments, the bispecific antibody construct comprises a first binding domain and a second binding domain, wherein the first binding domain specifically binds to a first cell surface antigen and the second binding domain specifically binds to human CD3. In some embodiments, the first and the second domain of the bispecific antibody construct is a “bispecific single chain antibody construct”, more preferably a bispecific “single chain Fv” (scFv). In a scFv, VL and VH of an antibody are joined, e.g., by a synthetic linker, as a single protein chain in which the VL and VH regions pair to form a monovalent molecule; see e.g., Huston et al. (1988) Proc. Natl. Acad. Sci USA 85:5879-5883). The linker can be a short peptide

of about ten to about 25 amino acids, preferably about 15 to 20 amino acids. The linker is usually rich in glycine for flexibility, as well as serine or threonine for solubility, and can either connect the N-terminus of the VH with the C-terminus of the VL, or vice versa. The scFv retains the specificity of the original immunoglobulin, despite removal of the constant regions and introduction of the linker. The VH and VL regions are arranged in the order VH-VL or VL-VH. It is preferred that the VH-region is positioned N-terminally of a linker sequence, and the VL-region is positioned C-terminally of the linker sequence. In certain embodiments, the first and second domain of the bispecific antibody construct are in a format selected from (scFv)₂, scFv-single domain mAb, diabody and oligomers of any of those formats.

[0049] In some embodiments, the bispecific antibody construct further comprises a third domain. In some embodiments, the third domain is a single-chain Fc (scFc) domain. In some embodiments, the scFc domain is a scFc half-life extended (HLE) domain. In some preferred embodiments, the third domain of the bispecific antibody construct is an HLE domain with an amino to carboxyl order: hinge-CH2-CH3-linker-hinge-CH2-CH3.

[0050] In some embodiments, the first binding domain of the bispecific antibody construct binds to a first cell surface antigen. In some embodiment, the first cell surface antigen is CD70. CD70 (also known as CD27L or TNFSF7) is a type II integral membrane protein whose normal expression is restricted to a subset of activated T and B cells, mature dendritic cells and thymic medullar epithelial cells.

[0051] In some embodiments, the first cell surface antigen is a tumor antigen. The term "tumor antigen" as used herein is understood to refer to those antigens that are presented on tumor cells. These antigens can be presented on the cell surface with an extracellular part, which is often combined with a transmembrane and cytoplasmic part of the molecule. These antigens can sometimes be presented only by tumor cells and not by the normal ones. Tumor antigens can be exclusively expressed on tumor cells or might represent a tumor specific mutation compared to normal cells. In this case, they are called tumor-specific antigens. More common are antigens that are presented by tumor cells and normal cells, and they are called tumor-associated antigens. These tumor-associated antigens can be overexpressed compared to normal cells or are accessible for antibody binding in tumor cells due to the less compact structure of the tumor tissue compared to normal tissue. In some embodiments, the first binding domain binds to tumor antigens selected from CD19, CD33, epidermal growth factor receptor variant iii (EGFRvIII), mesothelin (MSLN), cadherin 19 (CDH19), FMS-like tyrosine kinase 3 (FLT3), delta-like ligand 3 (DLL3), Placental-Cadherin (CDH3), B-cell maturation antigen (BCMA), prostate-specific

membrane antigen (PSMA), human mucin 17 (MUC17), and claudin-18 isoform 2 (CLDN18.2). In some embodiments, the tumor antigens are human tumor antigens.

[0052] In some preferred embodiments, the bispecific antibody construct comprises a first domain, a second domain and optionally a third domain, wherein the first domain binds to CD70 and the second domain binds to human CD3, and the third domain (if present) is an HLE domain with an amino to carboxyl order: hinge-CH2-CH3-linker-hinge-CH2-CH3. In other preferred embodiments, the bispecific antibody construct comprises a first domain, a second domain and optionally a third domain, wherein the first domain binds to a tumor antigen selected from CD19, CD33, EGFRvIII, MSLN, CDH19, FLT3, DLL3, CDH3, BCMA, PSMA, MUC17 and C:DN18.2, and the second domain binds to human CD3, and the third domain (if present) is a HLE domain with an amino to carboxyl order: hinge-CH2-CH3-linker-hinge-CH2-CH3. In a preferred embodiment, the first and second domain are linked together via a peptide linker, and are linked to the third domain (if present) via a peptide linker. Preferred peptide linker have been described herein above and are characterized by the amino acid sequence Gly-Gly-Gly-Gly-Ser, i.e. Gly4Ser, or polymers thereof, i.e. (Gly4Ser)_x, where x is an integer of 1 or greater (e.g. 2, 3, 4, 5, 6, or 7). In some of the preferred embodiments, the second binding domain comprises a polypeptide having the amino acid sequence of SEQ ID NO: 201.

[0053] In some embodiments, the first binding domain specifically binds to CD33. In some embodiments, the first binding domain comprises a polypeptide having the amino acid sequence of any one of SEQ ID NOs: 1-15.

[0054] In some embodiments, the first binding domain specifically binds to EGFRvIII. In some embodiments, the first binding domain comprises a polypeptide having the amino acid sequence of any one of SEQ ID NOs: 16-26.

[0055] In some embodiments, the first binding domain specifically binds to MSLN. In some embodiments, the first binding domain comprises a polypeptide having the amino acid sequence of any one of SEQ ID NOs: 27-38 and 165.

[0056] In some embodiments, the first binding domain specifically binds to CDH19. In some embodiments, the first binding domain comprises a polypeptide having the amino acid sequence of any one of SEQ ID NOs: 39-56.

[0057] In some embodiments, the first binding domain specifically binds to DLL3. In some embodiments, the first binding domain comprises a polypeptide having the amino acid sequence of SEQ ID NOs: 68-78.

[0058] In some embodiments, the first binding domain specifically binds to CD19. In some embodiments, the first binding domain comprises a polypeptide having the amino acid sequence of any one of SEQ ID NOs: 79-88.

[0059] In some embodiments, the first binding domain specifically binds to FLT3. In some embodiments, the first binding domain comprises a polypeptide having the amino acid sequence of any one of SEQ ID NOs: 57-67.

[0060] In some embodiments, the first binding domain specifically binds to CDH3. In some embodiments, the first binding domain comprises a polypeptide having the amino acid sequence of any one of SEQ ID NOs: 89-99.

[0061] In some embodiments, the first binding domain specifically binds to BCMA. In some embodiments, the first binding domain comprises a polypeptide having the amino acid sequence of any one of SEQ ID NOs: 100-110.

[0062] In some embodiments, the first binding domain specifically binds to PSMA. In some embodiments, the first binding domain comprises a polypeptide having the amino acid sequence of any one of SEQ ID NOs: 111-155, 166 and 167.

[0063] In some embodiments, the first binding domain specifically binds to CD70. In some embodiments, the first binding domain comprises a polypeptide having the amino acid sequence of any one of SEQ ID NOs: 156-164.

[0064] In some embodiments, the second binding domain of the bispecific antibody construct specifically binds to the human CD3 epsilon on the surface of a T cell. In some embodiments, the second domain of the bispecific antibody construct specifically binds to an extracellular epitope of the human CD3ε chain. In some embodiments, the second domain of the bispecific antibody construct that specifically binds to the human CD3 comprises a VL region comprising CDR-L1 having the amino acid sequence of sequence of SEQ ID NO: 193, CDR-L2 having the amino acid of SEQ ID NO: 194, and CDR-L3 having the amino acid sequence of sequence of SEQ ID NO: 195, and a VH region comprising CDR-H1 having the amino acid sequence of sequence of SEQ ID NO: 196, CDR-H2 having the amino acid sequence of sequence of SEQ ID NO: 197, and CDR-H3 having the amino acid sequence of sequence of SEQ ID NO: 198.

[0065] In some embodiments, the second domain of the bispecific antibody construct comprises a VH having the amino acid sequence of SEQ ID NO: 199 and a VL having the amino acid of SEQ ID NO: 200. In some embodiments, the second domain of the bispecific antibody construct comprises a polypeptide having the amino acid sequence of SEQ ID NO: 201.

[0066] In some embodiments, the protein is a CD70xCD3 bispecific antibody construct, which comprises a first domain that specifically binds to CD70 and a second domain that specifically binds to CD3. In one embodiment, the first domain specifically binds to CD70 and comprises the CDRs as depicted in SEQ ID NOs: 156 to 161, the second domain specifically binds to CD3 and has the CDRs as depicted in SEQ ID NOs: 193-198. In some embodiments, the bispecific antibody construct further comprises an HLE domain (third domain). In one embodiment, the bispecific antibody construct comprises a VH and a VL, wherein the VH comprises a polypeptide comprising, consisting essentially of, or consisting of the amino acid sequence of SEQ ID NO: 162 and the VL comprises a polypeptide comprising, consisting essentially of, or consisting of the amino acid sequence of SEQ ID NO: 163. In one embodiment, the CD70xCD3 bispecific antibody construct comprises a polypeptide that comprises, consists essentially of, or consists of the amino acid sequence of SEQ ID NO: 164.

[0067] In some embodiments, the protein is a BCMAxCD3 bispecific antibody construct, which comprises a first domain that specifically binds to BCMA and a second domain that specifically binds to CD3. In some embodiments, the bispecific antibody further comprises an HLE domain (third domain). In one embodiment, the first domain specifically binds to BCMA and has the CDRs as depicted in SEQ ID NOs: 100 to 105, the second domain specifically binds to CD3 and has the CDRs as depicted in SEQ ID NOs: 193-198. In one embodiment, the BCMAxCD3 bispecific antibody construct comprises a VH and a VL, wherein the VH comprises a polypeptide comprising, consisting essentially of, or consisting of the amino acid sequence of SEQ ID NO: 106 and the VL comprises a polypeptide comprising, consisting essentially of, or consisting of the amino acid sequence of SEQ ID NO: 107. In one embodiment, the BCMAxCD3 bispecific antibody construct comprises a polypeptide that comprises, consists essentially of, or consists of the amino acid sequence of SEQ ID NO: 108. In one embodiment, the BCMAxCD3 bispecific antibody construct comprises a polypeptide that comprises, consists essentially of, or consists of the amino acid sequence of SEQ ID NO: 109. In another embodiment, the BCMAxCD3 bispecific antibody construct comprises a polypeptide that comprises, consists essentially of, or consists of the amino acid sequence of SEQ ID NO: 110.

[0068] In some embodiments, the bispecific antibody is a CD33xCD3 bispecific antibody construct, which comprises a first domain that specifically binds to CD33 and a second domain that specifically binds to CD3. In some embodiments, the bispecific antibody construct further comprises an HLE domain (third domain). In one embodiment, the first domain specifically binds to CD33 and has the CDRs as depicted in SEQ ID NOs: 3 to 5 and 8 to 10, the second

domain specifically binds to CD3 and has the CDRs as depicted in SEQ ID NOs: 193-198. In one embodiment, the CD33xCD3 bispecific antibody construct comprises a VH and a VL, wherein the VH comprises a polypeptide comprising, consisting essentially of, or consisting of the amino acid sequence of SEQ ID NO: 1 or 2 and the VL comprises a polypeptide comprising, consisting essentially of, or consisting of the amino acid sequence of SEQ ID NO: 6 or 7. In another embodiment, the CD33xCD3 bispecific antibody construct comprises a polypeptide that comprises, consists essentially of, or consists of the amino acid sequence of any one of SEQ ID NO: 11 or 12. In another embodiment, the BCMAxCD3 bispecific antibody construct comprises a polypeptide that comprises, consists essentially of, or consists of the amino acid sequence of SEQ ID NO: 13 or 15.

[0069] In some embodiments, the protein is an EGFRvIIIxCD3 bispecific antibody construct, which comprises a first domain that specifically binds to EGFRvIII and a second domain that specifically binds to CD3. In some embodiments, the bispecific antibody further comprises an HLE domain (third domain). In one embodiment, the first domain specifically binds to EGFRvIII and has the CDRs as depicted in SEQ ID NOs: 16 to 21, the second domain specifically binds to CD3 and has the CDRs as depicted in SEQ ID NOs: 193-198. In one embodiment, the EGFRvIIIxCD3 bispecific antibody construct comprises a VH and a VL, wherein the VH comprises a polypeptide comprising, consisting essentially of, or consisting of the amino acid sequence of SEQ ID NO: 22 and the VL comprises a polypeptide comprising, consisting essentially of, or consisting of the amino acid sequence of SEQ ID NO: 23. In one embodiment, the EGFRvIIIxCD3 bispecific antibody construct comprises a polypeptide that comprises, consists essentially of, or consists of the amino acid sequence of SEQ ID NO: 24 or 25. In one embodiment, the EGFRvIIIxCD3 bispecific antibody construct comprises a polypeptide that comprises, consists essentially of, or consists of the amino acid sequence of SEQ ID NO: 24 or 26.

[0070] In some embodiments, the protein is a MSLNxCD3 bispecific antibody construct, which comprises a first domain that specifically binds to MSLN and a second domain that specifically binds to CD3. In some embodiments, the bispecific antibody construct further comprises an HLE domain (third domain). In one embodiment, the first domain specifically binds to MSLN and has the CDRs as depicted in SEQ ID NOs: 27 to 32, the second domain specifically binds to CD3 and has the CDRs as depicted in SEQ ID NOs: 193 to 198. In one embodiment, the MSLNxCD3 bispecific antibody construct comprises a VH and a VL, wherein the VH comprises a polypeptide comprising, consisting essentially of, or consisting of the amino acid sequence of SEQ ID NO: 33 and the VL comprises a polypeptide comprising, consisting essentially of, or consisting of the amino acid

sequence of SEQ ID NO: 34. In one embodiment, the MSLNxCD3 bispecific antibody construct comprises a polypeptide that comprises, consists essentially of, or consists of the amino acid sequence of any one of SEQ ID NOs: 35-38. In one embodiment, the MSLNxCD3 bispecific antibody construct comprises a polypeptide that comprises, consists essentially of, or consists of the amino acid sequence of SEQ ID NO: 165.

[0071] In some embodiments, the protein is a CDH19xCD3 bispecific antibody construct, which comprises a first domain that specifically binds to CDH19 and a second domain that specifically binds to CD3. In some embodiments, the bispecific antibody further comprises an HLE domain (third domain). In one embodiment, the first domain specifically binds to CDH19 and has the CDRs as depicted in SEQ ID NOs: 39 to 44, the second domain specifically binds to CD3 and has the CDRs as depicted in SEQ ID NOs: 193 to 198. In one embodiment, the CDH19xCD3 bispecific antibody construct comprises a VH and a VL, wherein the VH comprises a polypeptide comprising, consisting essentially of, or consisting of the amino acid sequence of SEQ ID NO: 45 or 51 and the VL comprises a polypeptide comprising, consisting essentially of, or consisting of the amino acid sequence of SEQ ID NO: 46 or 52. In one embodiment, the CDH19xCD3 bispecific antibody construct comprises a polypeptide that comprises, consists essentially of, or consists of the amino acid sequence of any one of SEQ ID NOs: 47, 48-50, and 53-56, . In one embodiment, the CDH19xCD3 bispecific antibody construct comprises a polypeptide that comprises, consists essentially of, or consists of the amino acid sequence of SEQ ID NO: 48.

[0072] In some embodiments, the protein is a DLL3xCD3 bispecific antibody, which comprises a first domain that specifically binds to DLL3 and a second domain that specifically binds to CD3. In some embodiments, the bispecific antibody further comprises an HLE domain (third domain). In one embodiment, the first domain specifically binds to DLL3 and has the CDRs as depicted in SEQ ID NOs: 68 to 73, the second domain specifically binds to CD3 and has the CDRs as depicted in SEQ ID NOs: 193 to 198. In one embodiment, the DLL3xCD3 bispecific antibody construct comprises a VH and a VL, wherein the VH comprises a polypeptide comprising, consisting essentially of, or consisting of the amino acid sequence of SEQ ID NO: 74 and the VL comprises a polypeptide comprising, consisting essentially of, or consisting of the amino acid sequence of SEQ ID NO: 75. In one embodiment, the DLL3xCD3 bispecific antibody construct comprises a polypeptide that comprises, consists essentially of, or consists of the amino acid sequence of SEQ ID NO: 76-78. In one embodiment, the DLL3xCD3 bispecific antibody construct comprises a polypeptide that comprises, consists essentially of, or consists of the amino acid sequence of SEQ ID NO: 78.

[0073] In some embodiments, the protein is a FLT3xCD3 bispecific antibody construct, which comprises a first domain that specifically binds to FLT3 and a second domain that specifically binds to CD3. In some embodiments, the bispecific antibody further comprises an HLE domain (third domain). In one embodiment, the first domain specifically binds to FLT3 and has the CDRs as depicted in SEQ ID NOs: 57 to 62, the second domain specifically binds to CD3 and has the CDRs as depicted in SEQ ID NOs 193 to 198. In one embodiment, the FLT3xCD3 bispecific antibody construct comprises a VH and a VL, wherein the VH comprises a polypeptide comprising, consisting essentially of, or consisting of the amino acid sequence of SEQ ID NO: 63 and the VL comprises a polypeptide comprising, consisting essentially of, or consisting of the amino acid sequence of SEQ ID NO: 64. In one embodiment, the FTL3xCD3 bispecific antibody construct comprises a polypeptide that comprises, consists essentially of, or consists of the amino acid sequence of any of SEQ ID NO: 65-67. In one embodiment, the FTL3xCD3 bispecific antibody construct comprises a polypeptide that comprises, consists essentially of, or consists of the amino acid sequence of SEQ ID NO: 67.

[0074] In some embodiments, the protein is a CDH3xCD3 bispecific antibody construct, which comprises a first domain that specifically binds to CDH3 and a second domain that specifically binds to CD3. In some embodiments, the bispecific antibody construct further comprises an HLE domain (third domain). In one embodiment, the first domain specifically binds to CDH3 and has the CDRs as depicted in SEQ ID NOs: 89 to 94, the second domain specifically binds to CD3 and has the CDRs as depicted in SEQ ID NOs: 193 to 198. In one embodiment, the CDH3xCD3 bispecific antibody construct comprises a VH and a VL, wherein the VH comprises a polypeptide comprising, consisting essentially of, or consisting of the amino acid sequence of SEQ ID NO: 95 and the VL comprises a polypeptide comprising, consisting essentially of, or consisting of the amino acid sequence of SEQ ID NO: 96. In one embodiment, the CDH3xCD3 bispecific antibody construct comprises a polypeptide that comprises, consists essentially of, or consists of the amino acid sequence of any of SEQ ID NO: 97-99. In one embodiment, the CDH3xCD3 bispecific antibody construct comprises a polypeptide that comprises, consists essentially of, or consists of the amino acid sequence of SEQ ID NO: 99.

[0075] In some embodiments, the protein is a PSMAxCD3 bispecific antibody construct, which comprises a first domain that specifically binds to PSMA and a second domain that specifically binds to CD3. In some embodiments, the bispecific antibody construct further comprises an HLE domain (third domain). In one embodiment, the first domain specifically binds to PSMA and has the CDRs as depicted in any of SEQ ID NOs: 111-116, the second domain specifically binds to

CD3 and has the CDRs as depicted in SEQ ID NOs: 193 to 198. In one embodiment, the first domain binds to PSMA and has the CDRs as depicted in any of SEQ ID NOs: 126-131 and 141-146, the second domain binds to CD3 and has the CDRs as depicted in SEQ ID NOs: 193 to 198. In one embodiment, the PSMAxCD3 bispecific antibody comprises a VH and a VL, wherein the VH comprises a polypeptide comprising, consisting essentially of, or consisting of the amino acid sequence of SEQ ID NO: 117, 132 or 147 and the VL comprises a polypeptide comprising, consisting essentially of, or consisting of the amino acid sequence of SEQ ID NO: 118, 133, or 148. In one embodiment, the PSMAxCD3 bispecific antibody construct comprises a polypeptide that comprises, consists essentially of, or consists of the amino acid sequence of any one of SEQ ID NOs: 119-125, 134-140, and 149-155. In one embodiment, the PSMAxCD3 bispecific antibody construct comprises a polypeptide that comprises, consists essentially of, or consists of the amino acid sequence of any of SEQ ID NOs: 121, 122, 124, 125, 136, 137, 139, 140, 151, 152, 154 and 155. In one embodiment, the PSMAxCD3 bispecific antibody construct comprises a polypeptide that comprises, consists essentially of, or consists of the amino acid sequence of SEQ ID NO: 166 or 167.

[0076] In some embodiments, the protein is a Cldn18.2xCD3 bispecific antibody construct, which comprises a first domain that specifically binds to Cldn18.2 and a second domain that specifically binds to CD3. In some embodiments, the bispecific antibody further comprises an HLE domain (third domain). In one embodiment, the first domain specifically binds to Cldn18.2 and has the CDRs as depicted in SEQ ID NOs: 168 to 173, the second domain specifically binds to CD3 and has the CDRs as depicted in SEQ ID NOs 193 to 198. In one embodiment, the Cldn18.2xCD3 bispecific antibody construct comprises a VH and a VL, wherein the VH comprises a polypeptide comprising, consisting essentially of, or consisting of the amino acid sequence of SEQ ID NO: 174 and the VL comprises a polypeptide comprising, consisting essentially of, or consisting of the amino acid sequence of SEQ ID NO: 175. In one embodiment, the Cldn18.2xCD3 bispecific antibody construct comprises a polypeptide that comprises, consists essentially of, or consists of the amino acid sequence of SEQ ID NO: 176 or 178. In one embodiment, the Cldn18.2xCD3 bispecific antibody construct comprises a polypeptide that comprises, consists essentially of, or consists of the amino acid sequence of SEQ ID NO: 178.

[0077] In some embodiments, the protein is a MUC17xCD3 bispecific antibody construct, which comprises a first domain that specifically binds to MUC17 and a second domain that specifically binds to CD3. In some embodiments, the bispecific antibody further comprises an HLE domain (third domain). In one embodiment, the first domain specifically binds to MUC17 and has the CDRs

as depicted in SEQ ID NOs: 184 to 189, the second domain specifically binds to CD3 and has the CDRs as depicted in SEQ ID NOs 193 to 198. In one embodiment, the MUC17xCD3 bispecific antibody construct comprises a VH and a VL, wherein the VH comprises a polypeptide comprising, consisting essentially of, or consisting of the amino acid sequence of SEQ ID NO: 190 and the VL comprises a polypeptide comprising, consisting essentially of, or consisting of the amino acid sequence of SEQ ID NO: 191. In one embodiment, the MUC17xCD3 bispecific antibody construct comprises a polypeptide that comprises, consists essentially of, or consists of the amino acid sequence of SEQ ID NO: 192.

[0078] In certain embodiments, the bispecific antibody construct comprises a polypeptide that comprises, consists essentially or consists of the amino acid sequence of SEQ ID NO: 13, SEQ ID NO: 15, SEQ ID NO: 25, SEQ ID NO: 48, SEQ ID NO: 67, SEQ ID NO: 78, SEQ ID NO: 88, SEQ ID NO: 109, SEQ ID NO: 110, SEQ ID NO: 165, SEQ ID NO: 166, SEQ ID NO: 167, SEQ ID NO: 178 or SEQ ID NO: 192.

[0079] The bispecific antibody disclosed herein can be prepared by methods known in the art. For example, the bispecific antibody can be prepared by methods disclosed in WO2008/119657 and WO2017/134140.

Additional Excipients that May be Used in the Composition

[0080] In some embodiments, the composition further comprises one or more excipients suitable for pharmaceutical compositions. In some embodiments, the composition further comprises a salt, a buffer, a saccharide or a saccharide derivative, an amino acid, or a preservative, or combinations of two or more of the foregoing. In some embodiments, the composition further comprises a salt, a saccharide or a saccharide derivative, an amino acid, and optionally a preservative. In some embodiments, the composition further comprises a salt, a buffer, a saccharide or a saccharide derivative, and an amino acid. In some embodiments, the composition further comprises a salt, a buffer, a saccharide or a saccharide derivative, an amino acid, and a preservative.

[0081] Exemplary salts that may be used in the composition include salts that are suitable to be used in pharmaceutical compositions (e.g., NaCl). Exemplary buffers that may be used in the composition include acetate buffer, glutamate buffer, citrate buffer, succinate buffer, tartrate buffer, fumarate buffer, maleate buffer, histidine buffer, and phosphate buffer. Exemplary saccharides or saccharide derivatives include monosaccharides, disaccharides, cyclic polysaccharides and sugar alcohols, such as sugars (e.g., sucrose and trehalose) and sugar alcohol (e.g., mannitol and sorbitol). Exemplary amino acids include lysine, histidine, arginine,

glycine, methionine, and alanine. Exemplary preservatives include benzoates (e.g., benzyl alcohol and sodium benzoate) and sorbates (e.g., potassium sorbate).

[0082] The pH of the composition can be in the range of from about 3.5 to 7.5. In some embodiments, the composition has a pH of from about 4.0 to about 7.0. In some embodiments, the composition has a pH of from about 4.2 to about 7.0, or from about 5.0 to about 7.0, or from about 5.5 to about 7.0, or from about 6.0 to about 7.0, or from about 6.5 to about 7.0, or from about 4.2 to about 6.5, or from about 4.2 to about 6.0, or from about 4.2 to about 5.5, or from about 4.2 to about 5.0, or from about 5.0 to about 6.0, or from about 5.0 to about 6.5, or from about 5.0 to about 6.0, or from about 5.0 to about 5.5, or from about 5.5 to about 6.0, or from about 5.5 to about 6.5, or from about 6.0 to about 6.5. In some embodiments, the pH of the composition is about 3.5, about 4.0, about 4.5, about 5.0, about 5.5, about 6.0, about 6.5, about 7.0, or about 7.5.

[0083] The pH of the composition may be achieved by using one or more buffers listed above. In some embodiments, the composition comprising a buffer selected from an acetate buffer, a glutamate buffer, a citrate buffer, a succinate buffer, a tartrate buffer, a fumarate buffer, a maleate buffer, a histidine buffer, and a phosphate buffer. In some embodiments, the composition comprising a combination of two buffers selected from the above list of buffers, e.g., a glutamate buffer and a citrate buffer.

[0084] In some embodiments, the composition comprises substantially no additional buffer. As used herein, the phrase “substantially no additional buffer” refers that the composition contains no buffering agent added therein for the purpose of adjusting and/or maintaining the pH of the composition. For example, saline solution (0.9% NaCl solution) contains no additional buffer and has a pH of about 5.5. It is believed that the pH of saline solution is achieved and maintained in part by atmospheric CO₂ dissolved in water. See e.g., Reddi, B. *AJ Int. J. Med. Sci.* 10: 747-750 (2013). Such a composition may contain residual buffer that does not contribute to the buffering capacity of the composition.

[0085] In some embodiments, the composition comprises a protein at any of the concentration disclosed above, a surfactant at any of the concentration disclosed above, and further comprises a salt (e.g., NaCl), a sugar (e.g., sucrose), an amino acid (e.g., lysine), and optionally a preservative (e.g., benzyl alcohol). In some embodiments, the composition comprises a protein at any of the concentration disclosed above, a surfactant at any of the concentration disclosed above, and further comprises a salt (e.g., NaCl), a sugar (e.g., sucrose), an amino acid (e.g., lysine), a buffer (e.g., a glutamate and/or citrate buffer), and optionally a

preservative (e.g., benzyl alcohol). The pH of the composition can be any of the pH value disclosed above. In some embodiments, the pH of the composition is about 5.5.

[0086] In some embodiments, the composition disclosed herein is a pharmaceutical composition. As used herein, the term "pharmaceutical composition" is understood to refer to a formulation comprising a protein (e.g., a bispecific antibody construct) suitable for injection and/or administration into a patient (e.g., a human) in need thereof. More particularly, a pharmaceutical composition is substantially sterile and does not contain any agents that are unduly toxic or infectious to the recipient.

Containers for the Composition

[0087] In some embodiments, the composition disclosed herein is contained in a container. Containers that may be used herein include those suitable for pharmaceutical use, e.g., containers made of materials that are nontoxic and maintain physical integrity. In some embodiments, the container is a component for intravenous (IV) administration. As used herein, the phrase "component for IV administration" is understood to refer to a container or a part of a system that may contact the composition during IV administration. In some embodiments, the container is an IV bag or IV tubing.

[0088] Materials that may be used for making containers include those typically used for making pharmaceutical containers such as glass and plastic.

[0089] In some embodiments, the container is a plastic container. In some embodiments, the container is made of a material comprising polyolefin (e.g., polypropylene (PP) and polyethylene (PE)), polyvinyl chloride (PVC), ethyl vinyl acetate (EVA), or polyurethane (e.g., polyester and polyether). In some embodiments, the container is made of a material comprising PVC. In some embodiments, the PVC is substantially free of di-2-ethylhexyl phthalate (DEHP) or tri-2-ethylhexyltrimellitate (TOTM). As used herein, the term "substantially free of" is understood to refer to PVC in which DEHP or TOTM is not used and/or detected. DEHP and TOTM are plasticizers that may be used in making PVC softer therefore could be made into different shapes. In some embodiments, the container is made of a material that does not comprise PVC. In some embodiments, the container is made of a material comprising polyolefin, EVA, or polyurethane (e.g., polyester and polyether). In some embodiments, the container is made of a material comprising PP and/or PE.

[0090] Under certain conditions, DEHP or TOTM contained in certain types of PVC plastic can leach from the plastic in the presence of certain concentrations of certain surfactants (e.g., polysorbates, see Example 3). Not wish to be bound by any particular theory, it is believed

that the more lipophilic a surfactant, the easier DEHP or TOTM can leach out in the presence of the surfactant. The hydrophilic-lipophilic balance (HLB) of a surfactant is a measure of the degree to which it is hydrophilic or lipophilic, and can be calculated by methods known in the art. See e.g., Griffin, William C. Calculation of HLB Values of Non-Ionic Surfactants, *Journal of the Society of Cosmetic Chemists*, 5 (4): 249–56 (1954). The HLB value can be used to predict properties of a surfactant, e.g., $HLB > 10$ indicates the surfactant is more water-soluble (lipid-insoluble), while $HLB < 10$ indicates the surfactant is more lipid-soluble (water-insoluble). The HLB value of exemplary surfactants that can be used in the composition disclosed herein include 17 (polysorbate 20), 15 (polysorbate 80) (<https://pharmlabs.unc.edu/labs/emulsions/hlb.htm>) and 29 (poloxamer 188)

(http://www.rumapel.com.ar/cosmetica_miscelaneos/ficha_tecnica/Pluracare%20L-%20F%20Grades.pdf). In embodiments where the container is made of a PVC plastic comprising DEHP or TOTM, the surfactant in the composition preferably have an HLB value of at least 20 (e.g., a poloxamer, see Example 3), more preferably an HLB value of between 20 and 30.

[0091] Containers such as IV components (e.g., IV bags and tubes used for IV administration) that are made of the above listed materials are commercially available. For example, various suitable containers are available from manufactures such as Baxter Healthcare Corporation and B. Braun Medical Inc.

Pharmaceutical Preparation

[0092] Also disclosed herein are pharmaceutical preparations comprising the aqueous composition disclosed above. As used herein, the term “pharmaceutical preparation” is understood to refer to a preparation comprising the aqueous composition disclosed herein in a suitable pharmaceutical container prior to administration to a patient (e.g., a human). In some embodiments, disclosed herein is a pharmaceutical preparation comprising an aqueous pharmaceutical composition in a container, wherein the aqueous pharmaceutical composition comprising: a) a protein at a concentration of between about 0.001 µg/ml and about 100 µg/ml, and b) a surfactant at a concentration of at least about 0.25 x of the CMC of the surfactant, and wherein the protein is not blinatumomab.

[0093] In some embodiments, the surfactant comprised in the composition is a polysorbate. In some embodiments, the surfactant is a polysorbate selected from polysorbate 20, polysorbate 40, polysorbate 60, polysorbate 65, polysorbate 80 and polysorbate 85. In some embodiments, the surfactant is polysorbate 20 or polysorbate 80. In some embodiments, the

surfactant is a poloxamer. In some embodiments, the surfactant is poloxamer 188 or poloxamer 407. In some embodiments, the surfactant is Triton X-100.

[0094] In some embodiments, disclosed herein is a pharmaceutical preparation comprising an aqueous pharmaceutical composition in a container, wherein the aqueous pharmaceutical composition comprising: a) a bispecific antibody construct at a concentration of between about 0.001 µg/ml and about 100 µg/ml, and b) a surfactant at a concentration of at least about 0.25 x of CMC of the surfactant, wherein the surfactant is a poloxamer. In some embodiments, the poloxamer is poloxamer 188 or poloxamer 407.

[0095] In some embodiments, the surfactant is present in the composition at a concentration that is between about 0.25x and about 20x of the CMC of the surfactant. In some embodiments, the surfactant is present in the composition at a concentration that is between about 0.25x and about 10x of the CMC of the surfactant. In some embodiments, the surfactant is present in the composition at a concentration that is within any of the concentration ranges disclosed above. In some embodiments, the surfactant is present in the composition at a concentration that is any of the concentration disclosed above. The CMC value of a surfactant can be determined by the methods disclosed above. In some embodiments, the CMC value of certain commonly used surfactants are listed in Table 1.

[0096] In some embodiments, the aqueous pharmaceutical composition comprising the protein at a concentration of between about 0.001 µg/ml and about 50 µg/ml. In some embodiments, the composition comprising the protein (e.g., a bispecific antibody construct) at a concentration that is within any of the ranges disclosed above. In some embodiments, the protein (e.g., a bispecific antibody construct) is present in the composition at any of the concentrations disclosed above. Any of the concentration or concentration range for the protein disclosed above can be combined with any of the concentration or concentration range for the surfactant disclosed above.

[0097] As disclosed above, the protein can be a therapeutic protein such as an antigen binding protein and a fusion protein. In some embodiments, the protein is a bispecific antigen binding protein. In some embodiments, the protein is a bispecific antibody construct disclosed above. In some embodiments, the protein is a CD70xCD3 bispecific antibody construct. In some embodiments, the protein is a BCMAxCD3 bispecific antibody construct. In some embodiments, the protein is a CD33xCD3 bispecific antibody construct. In some embodiments, the protein is an EGFRvIIIxCD3 bispecific antibody construct. In some embodiments, the protein is a MSLNxCD3 bispecific antibody construct. In some embodiments, the protein is a CDH19xCD3

bispecific antibody construct. In some embodiments, the protein is a DLL3xCD3 bispecific antibody construct. In some embodiments, the protein is a FLT3xCD3 bispecific antibody construct. In some embodiments, the protein is a CDH3xCD3 bispecific antibody construct. In some embodiments, the protein is a PSMAxCD3 bispecific antibody construct. In some embodiments, the protein is a Cldn18.2xCD3 bispecific antibody construct. In some embodiments, the protein is a MUC17xCD3 bispecific antibody construct. Each of these bispecific antibody constructs is disclosed above. In certain embodiments, the protein comprises, consists essentially or consists of the amino acid sequence of SEQ ID NO: 13, SEQ ID NO: 15, SEQ ID NO: 25, SEQ ID NO: 48, SEQ ID NO: 67, SEQ ID NO: 78, SEQ ID NO: 88, SEQ ID NO: 109, SEQ ID NO: 110, SEQ ID NO: 165, SEQ ID NO: 166, SEQ ID NO: 167, SEQ ID NO: 178 or SEQ ID NO: 192.

[0098] The pH of the composition can be within any of the pH ranges disclosed above. In some embodiments, the pH of the composition can be any pH disclosed above.

[0099] In some embodiments, the aqueous pharmaceutical composition further comprises one or more excipients suitable for use in a pharmaceutical composition. In some embodiments, the aqueous pharmaceutical composition further comprises a salt, a buffer, an amino acid, a saccharide or a saccharide derivative, a preservative or combinations thereof. In some embodiments, the aqueous pharmaceutical composition further comprises a salt, an amino acid, a saccharide or a saccharide derivative, a preservative or combinations thereof. Exemplary salts, buffers, amino acids, saccharides or derivatives thereof, and preservatives that may be used in the composition are disclosed above.

[00100] In some embodiments, the container is a plastic container. In some embodiments, the container is made of a material comprising polyolefin (e.g., PP and PE), PVC, EVA, or polyurethane (e.g., polyester and polyether). In some embodiments, the container is made of a material comprising PVC. In some embodiments, the PVC is substantially free of DEHP or TOTM. In some embodiments, the container is made of a material that does not comprise PVC.

[00101] In some embodiments, the surfactant comprised in the composition is a poloxamer, and the container can be made of a material comprising polyolefin, PVC, EVA, or polyurethane (e.g., polyester and polyether). In some embodiments, the surfactant comprised in the composition is a polysorbate or Triton X-100, and the container can be made of a material comprising polyolefin, PVC that is substantially free of DEHP or TOTM, EVA, or polyurethane (e.g., polyester and polyether).

[00102] In some embodiments, the container is a component for intravenous (IV) administration. In some embodiments, the container is an IV bag or IV tubing.

[00103] In some embodiments, the pharmaceutical preparation comprises an aqueous pharmaceutical composition in a container, wherein the aqueous pharmaceutical composition comprising a bispecific antibody construct (e.g., any of the bispecific antibody construct disclosed herein), a surfactant (e.g., any of the surfactant disclosed herein), and further comprises a salt (e.g., NaCl), an amino acid (e.g., lysine), a saccharide or a saccharide derivative (e.g., sucrose or mannitol), optionally a preservative (e.g., and benzyl alcohol), and wherein the container is made of a material comprising polyolefin, PVC, EVA, or polyurethane (e.g., polyester and polyether). In some embodiments, the pharmaceutical preparation comprises an aqueous pharmaceutical composition in a container, wherein the aqueous pharmaceutical composition comprising a bispecific antibody construct (e.g., any of the bispecific antibody construct disclosed herein), a surfactant (e.g., any of the surfactant disclosed herein), and further comprises a salt (e.g., NaCl), a buffer (e.g., a glutamate buffer and/or a citrate buffer), an amino acid (e.g., lysine), a saccharide or a saccharide derivative (e.g., sucrose or mannitol), optionally a preservative (e.g., benzyl alcohol), and wherein the container is made of a material comprising polyolefin, PVC, EVA, or polyurethane (e.g., polyester and polyether). The concentration of the bispecific antibody construct can be any of the concentration disclosed above for the protein. The concentration of the surfactant can be any of the concentration disclosed above. The pH of the composition is in the range of between about 3.5 and about 7.0. In some embodiments, the pH of the composition is about 5.5. In some embodiments, the container is an IV bag.

Methods of Administering the Composition

[00104] Also disclosed herein are methods of administering the composition disclosed herein to a patient. In some embodiments, disclosed herein is a method of administering a protein to a patient, the method comprises a) preparing an aqueous pharmaceutical composition in a container, wherein the aqueous pharmaceutical composition comprises a protein (e.g., a bispecific antibody construct disclosed herein) at a concentration of between about 0.001 µg/ml and about 100 µg/ml and a surfactant at a concentration of at least about 0.25 x of CMC of the surfactant, and b) administering the aqueous pharmaceutical composition to the patient, wherein the protein is not blinatumomab.

[00105] In some embodiments, the aqueous pharmaceutical composition is prepared by diluting a first composition comprising the protein (e.g., a bispecific antibody construct) with a suitable aqueous solution. In some embodiments, the aqueous pharmaceutical composition is

prepared by adding the suitable solution into the container followed by adding an appropriate amount of a first composition into the container thereby diluting the first composition comprising the protein.

[00106] In some embodiments, the first composition is a liquid composition comprising the protein. In some embodiments, the first composition is a liquid composition reconstituted from a lyophilized composition comprising the protein. In some embodiments, the first composition is a liquid composition reconstituted from a lyophilized composition comprising the protein using sterile water.

[00107] In some embodiments, the suitable aqueous solution that is used for diluting the first composition comprises the surfactant at a concentration of at least about 0.25x of the CMC of the surfactant. In some embodiments, the suitable aqueous solution that is used for diluting the first composition comprises the surfactant at a concentration of between about 0.25x and 20x, or between about 0.25x and about 10x of the CMC of the surfactant. In some embodiments, the suitable aqueous solution that is used for diluting the first composition comprises the surfactant at a concentration of 0.25x, about 0.5 x, about 1x, about 2x, about 3x, about 4x, about 5x, about 6x, about 7x, about 8x, about 9x, about 10x, about 15x or about 20x of the CMC of the surfactant. In some embodiments, the suitable aqueous solution that is used for diluting the first composition further comprises NaCl. In some embodiments, the suitable aqueous solution that is used for diluting the first composition has a pH of between about 3.5 and about 7.0, or between about 4.0 and about 6.5. In some embodiments, the suitable aqueous solution that is used for diluting the first composition has a pH of about 5.5. In some embodiments, the method further comprising rinsing the container with the suitable aqueous solution that is used for diluting the first composition before the preparing step.

[00108] In some embodiments, the surfactant in the aqueous pharmaceutical composition is a polysorbate. In some embodiments, the surfactant is a polysorbate selected from polysorbate 20, polysorbate 40, polysorbate 60, polysorbate 65, polysorbate 80 and polysorbate 85. In some embodiments, the surfactant is polysorbate 20 or polysorbate 80. In some embodiments, the surfactant is a poloxamer. In some embodiments, the surfactant is poloxamer 188 or poloxamer 407. In some embodiments, the surfactant is Triton X-100.

[00109] In some embodiments, the surfactant in the aqueous pharmaceutical composition is present in the composition at a concentration that is between about 0.25x and about 20x of the CMC of the surfactant. In some embodiments, the surfactant is present in the aqueous pharmaceutical composition at a concentration that is between about 0.25x and about 10x of the

CMC of the surfactant. In some embodiments, the surfactant is present in the aqueous pharmaceutical composition at a concentration that is within any of the concentration ranges disclosed above. In some embodiments, the surfactant is present in the aqueous pharmaceutical composition at a concentration that is any of the concentrations disclosed above. The CMC value of a surfactant can be determined by the methods disclosed above. In some embodiments, the CMC value of certain commonly used surfactants are listed in Table 1.

[00110] In some embodiments, the aqueous pharmaceutical composition comprising the protein at a concentration of between about 0.001 µg/ml and about 50 µg/ml. In some embodiments, the aqueous pharmaceutical composition comprising the protein at a concentration that is within any of the ranges disclosed above. In some embodiments, the protein is present in the composition at any of the concentrations disclosed above.

[00111] As disclosed above, the protein can be a therapeutic protein such as an antigen binding protein, a mAb or a fusion protein. In some embodiments, the protein is a bispecific antigen binding protein. In some embodiments, the protein is a bispecific antibody construct disclosed above. In some embodiments, the protein is a CD70xCD3 bispecific antibody construct. In some embodiments, the protein is a BCMAxCD3 bispecific antibody construct. In some embodiments, the protein is a CD33xCD3 bispecific antibody construct. In some embodiments, the protein is an EGFRvIIIxCD3 bispecific antibody construct. In some embodiments, the protein is a MSLNxCD3 bispecific antibody construct. In some embodiments, the protein is a CDH19xCD3 bispecific antibody construct. In some embodiments, the protein is a DLL3xCD3 bispecific antibody construct. In some embodiments, the protein is a FLT3xCD3 bispecific antibody construct. In some embodiments, the protein is a CDH3xCD3 bispecific antibody construct. In some embodiments, the protein is a PSMAxCD3 bispecific antibody construct. In some embodiments, the protein is a Cldn18.2xCD3 bispecific antibody construct. In some embodiments, the protein is a MUC17xCD3 bispecific antibody construct. Each of these bispecific antibody constructs is disclosed above. In certain embodiments, the protein comprises, consists essentially or consists of the amino acid sequence of SEQ ID NO: 13, SEQ ID NO: 15, SEQ ID NO: 25, SEQ ID NO: 48, SEQ ID NO: 67, SEQ ID NO: 78, SEQ ID NO: 88, SEQ ID NO: 109, SEQ ID NO: 110, SEQ ID NO: 165, SEQ ID NO: 166, SEQ ID NO: 167, SEQ ID NO: 178 or SEQ ID NO: 192.

[00112] The pH of the composition can be within any of the pH ranges disclosed above. In some embodiments, the pH of the composition can be any pH disclosed above.

[00113] In some embodiments, the aqueous pharmaceutical composition further comprises one or more excipients suitable for use in a pharmaceutical composition. In some embodiments, the aqueous pharmaceutical composition further comprises a salt, a buffer, an amino acid, a saccharide or a saccharide derivative, optionally a preservative, or combinations of two or more of the foregoing. In some embodiments, the aqueous pharmaceutical composition further comprises a salt, an amino acid, a saccharide or a saccharide derivative, optionally a preservative, or combinations of two or more of the foregoing. Exemplary salts, buffers, amino acids, saccharides or derivatives thereof, and preservatives that may be used in the composition are disclosed above.

[00114] In some embodiments, the container is a plastic container or component. In some embodiments, the container is made of a material comprising polyolefin (e.g., PP and PE), PVC, EVA, or polyurethane (e.g., polyester and polyether). In some embodiments, the container is made of a material comprising PVC. In some embodiments, the PVC is substantially free of DEHP or TOTM. In some embodiments, the container is made of a material that does not comprise PVC. In some embodiments, the container is an IV bag or IV tube.

[00115] In some embodiments, the aqueous pharmaceutical composition is administered to the patient via IV administration. In some embodiments, the patient is a cancer patient. In some embodiments, the patient is a human.

Method for Accessing Binding of Protein to a Solid Surface

[00116] Also disclosed herein is a method for accessing binding of proteins to solid surfaces. In some embodiments, disclosed herein is of accessing binding of a protein to a solid surface, comprising: a) incubating an aqueous solution comprising the protein with the solid surface, wherein the protein is labeled with a fluorophore, b) removing the aqueous solution from the solid surface and rinse the surface, and c) imaging the solid surface using confocal microscopy.

Table 2 below lists the sequences disclosed herein

	Designation			AMINO ACID SEQUENCE
1.	CD33 ccVH E11	artificial	aa	QVQLVQSGAEVKKPGESVKVSKASGYFTNYGM NWVKQAPGQCLEWMGWINTYTGEPTYADKFQG RVTMTTDTSTSTAYMEIRNLGGDDTAVYYCARWS WSDGYVYFDYWGGQTSVTVSS

2.	CD33 VH E11	artificial	aa	QVQLVQSGAEVKKPGESVKVSKASGYFTFTNYGM NWVKQAPGQGLEWMGWINTYTGEPTYADKFQG RVTMTTDTSTSTAYMEIRNLGGDDTAVYYCARWS WSDGYVYFDYWGQGTSTVTVSS
3.	CD33 HCDR1 E11	artificial	aa	NYGMN
4.	CD33 HCDR2 E11	artificial	aa	WINTYTGEPTYADKFQG
5.	CD33 HCDR3 E11	artificial	aa	WSWSDGYVYFDY
6.	CD33 CC VL E11	artificial	aa	DIVMTQSPDSLTVSLGERTTINCKSSQSVLDSSTNK NSLAWYQQKPGQPPKLLLSWASTRESGIPDRFSGS GSGTDFTLTIDSPQPEDSATYYCQSAHFPITFGCG TRLEIK
7.	CD33 VL E11	artificial	aa	DIVMTQSPDSLTVSLGERTTINCKSSQSVLDSSTNK NSLAWYQQKPGQPPKLLLSWASTRESGIPDRFSGS GSGTDFTLTIDSPQPEDSATYYCQSAHFPITFGQG TRLEIK
8.	CD33 LCDR1 E11	artificial	aa	KSSQSVLDSSTNKNSLA
9.	CD33 LCDR2 E11	artificial	aa	WASTRES
10.	CD33 LCDR3 E11	artificial	aa	QSAHFPIT
11.	CD33 HL CC E11	artificial	aa	QVQLVQSGAEVKKPGESVKVSKASGYFTFTNYGM NWVKQAPGQCLEWMGWINTYTGEPTYADKFQG RVTMTTDTSTSTAYMEIRNLGGDDTAVYYCARWS WSDGYVYFDYWGQGTSTVTVSSGGGGSGGGGSG GGGSDIVMTQSPDSLTVSLGERTTINCKSSQSVLDS STNKNSLAWYQQKPGQPPKLLLSWASTRESGIPDR FSGSGSGTDFTLTIDSPQPEDSATYYCQSAHFPITF GCGTRLEIK
12.	CD33 HL E11	artificial	aa	QVQLVQSGAEVKKPGESVKVSKASGYFTFTNYGM NWVKQAPGQGLEWMGWINTYTGEPTYADKFQG RVTMTTDTSTSTAYMEIRNLGGDDTAVYYCARWS WSDGYVYFDYWGQGTSTVTVSSGGGGSGGGGSG GGGSDIVMTQSPDSLTVSLGERTTINCKSSQSVLDS STNKNSLAWYQQKPGQPPKLLLSWASTRESGIPDR FSGSGSGTDFTLTIDSPQPEDSATYYCQSAHFPITF GQGTRLEIK
13.	CD33 CC E11 HL x I2C HL Bispecific molecule	artificial	aa	QVQLVQSGAEVKKPGESVKVSKASGYFTFTNYGM NWVKQAPGQCLEWMGWINTYTGEPTYADKFQG RVTMTTDTSTSTAYMEIRNLGGDDTAVYYCARWS WSDGYVYFDYWGQGTSTVTVSSGGGGSGGGGSG

				GGGSDIVMTQSPDSLTVSLGERTTINCKSSQSVLDS STNKNSLAWYQQKPGQPPKLLSWASTRESGIPDR FSGSGSGTDFTLTIDSPQPEDSATYYCQSAHPITF GCGTRLEIKSGGGGSEVQLVESGGGLVQPGGSLKL SCAASGFTFNKYAMNWVRQAPGKGLEWVARIRS KYNYYATYYADSVKDRFTISRDDSKNTAYLQMNNL KTEDTAVYYCVRHGNFGNSYISYWAYWGQGLVT VSSGGGGSGGGGSGGGGSQTVVTQEPSLTVSPGG TVTLTCGSSTGAVTSGNYPNWVQKPGQAPRGLI GGTKFLAPGTPARFSGSLLGGKAALTLSGVQPEDEA EYYCVLWYSNRWVFGGGTKLTVL
14.	CD33 E11 HL x I2C HL	artificial	aa	MGWSCIIILFVATATGVHSQVQLVQSGAEVKKPGE SVKVSCASGYFTNYGMNWVKQAPGQGLEWM GWINTYTGEPTYADKFQGRVTMTTDTSTSTAYMEI RNLGGDDTAVYYCARWSWSDGYVYFDYWGQG TSVTSSGGGGSGGGGSGGGGSDIVMTQSPDSL VSLGERTTINCKSSQSVLDSSTNKNSLAWYQQKPG QPPKLLSWASTRESGIPDRFSGSGSGTDFTLTIDSP QPEDSATYYCQSAHPITFGQTRLEIKSGGGGSE VQLVESGGGLVQPGGSLKLSAASGFTFNKYAMN WVRQAPGKGLEWVARIRSKYNYYATYYADSVKDR FTISRDDSKNTAYLQMNNLKTEDTAVYYCVRHGNF GNSYISYWAYWGQGLVTVSSGGGGSGGGSGG GGSQTVVTQEPSLTVSPGGTVTLTCGSSTGAVTSG NYPNWVQKPGQAPRGLIGGTKFLAPGTPARFSG SLLGGKAALTLSGVQPEDEAEYYCVLWYSNRWVFG GGTKLTVLHHHHH
15.	CD33 CC x I2C- scFc Bispecific HLE molecule	artificial	aa	QVQLVQSGAEVKKPGESVKVSCASGYFTNYGMN WVKQAPGQCLEWMGWINTYTGEPTYADKFQGRVT MTTDTSTSTAYMEIRNLGGDDTAVYYCARWSWSDG YVYFDYWGQGTSTVTSSGGGGSGGGGSGGGGSDI VMTQSPDSLTVSLGERTTINCKSSQSVLDSSTNKNSL AWYQQKPGQPPKLLSWASTRESGIPDRFSGSGSGT DFTLTIDSPQPEDSATYYCQSAHPITFGCGTRLEIKS GGGGSEVQLVESGGGLVQPGGSLKLSAASGFTFNK YAMNWVRQAPGKGLEWVARIRSKYNYYATYYADS VKDRFTISRDDSKNTAYLQMNNLKTEDTAVYYCVRH GNFGNSYISYWAYWGQGLVTVSSGGGGSGGGGS GGGGSQTVVTQEPSLTVSPGGTVTLTCGSSTGAVTS GNYPNWVQKPGQAPRGLIGGTKFLAPGTPARFSG SLLGGKAALTLSGVQPEDEAEYYCVLWYSNRWVFGG GTKLTVLGGGGDKHTCPPCPAPELLGGPSVFLFPPK PKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGV EVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKE YKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRE EMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYK

				TTPPVLDSDGSFFLYSKLTVDKSRWQQGNVFSCSVM HEALHNHYTQKSLSLSPGKGGGGSGGGSGGGSGG GGGGSGGGSGGGSGDKTHTCPPCPAPPELLGGPSVFL FPPKPKDTLMISRTPEVTCVVDVSHEDPEVKFNWY VDGVEVHNAKTKPCEEQYGSTYRCVSVLTVLHQDW LNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTL PPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQP ENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVF SCSVMHEALHNHYTQKSLSLSPGK
16.	EGFRvIIIxCD3- scFc VH CDR1	artificial	aa	NYGMH
17.	EGFRvIIIxCD3- scFc VH CDR2	artificial	aa	VIWYDGSDKYYADSVRG
18.	EGFRvIIIxCD3- scFc VH CDR3	artificial	aa	DGYDILTGNPRDFDY
19.	EGFRvIIIxCD3- scFc VL CDR1	artificial	aa	RSSQSLVHSDGNTYLS
20.	EGFRvIIIxCD3- scFc VL CDR2	artificial	aa	RISRRFS
21.	EGFRvIIIxCD3- scFc VL CDR3	artificial	aa	MQSTHVPRT
22.	EGFRvIII_CCxC D3-scFc VH	artificial	aa	QVQLVESGGGVVQSGRSLRLSCAASGFTFRNYGMH WVRQAPGKCLEWVAVIWDGSDKYYADSVRGFTI SRDNSKNTLYLQMNSLRAEDTAVYYCARDGYDILT NPRDFDYWGQGLTVTVSS
23.	EGFRvIII_CCxC D3-scFc VL	artificial	aa	DTVMTQTPLSSHVTLGQPASISCRSSQSLVHSDGNTY LSWLQQRPGQPPRLIYRISRRFSGVPDRFSGSAGT DFTLEISRVEAEDVGVYYCMQSTHVPRTFGCGTKVEI K
24.	EGFRvIII_CCxC D3-scFc scFv	artificial	aa	QVQLVESGGGVVQSGRSLRLSCAASGFTFRNYGMH WVRQAPGKCLEWVAVIWDGSDKYYADSVRGFTI SRDNSKNTLYLQMNSLRAEDTAVYYCARDGYDILT NPRDFDYWGQGLTVTVSSGGGGSGGGSGGGSGD TVMTQTPLSSHVTLGQPASISCRSSQSLVHSDGNTYL SWLQQRPGQPPRLIYRISRRFSGVPDRFSGSAGTD FTLEISRVEAEDVGVYYCMQSTHVPRTFGCGTKVEIK
25.	EGFRvIII_CCxC D3-scFc Bispecific molecule	artificial	aa	QVQLVESGGGVVQSGRSLRLSCAASGFTFRNYGMH WVRQAPGKCLEWVAVIWDGSDKYYADSVRGFTI SRDNSKNTLYLQMNSLRAEDTAVYYCARDGYDILT NPRDFDYWGQGLTVTVSSGGGGSGGGSGGGSGD TVMTQTPLSSHVTLGQPASISCRSSQSLVHSDGNTYL SWLQQRPGQPPRLIYRISRRFSGVPDRFSGSAGTD FTLEISRVEAEDVGVYYCMQSTHVPRTFGCGTKVEIK SGGGGSEVQLVESGGGLVQPGGSLKLSAASGFTFN KYAMNWVRQAPGKLEWVARIRSKYNNYATYYAD SVKDRFTISRDDSKNTAYLQMNNLKTEDTAVYYCVR

				HGNFGNSYISYWAYWGQGLTVTVSSGGGSGGGG SGGGGSQTVVTQEPLTVSPGGTVTLTCGSSTGAVTS GNYPNWVQKPGQAPRGLIGGTKFLAPGTPARFSG SLLGGKAALTLSGVQPEDEAEYYCVLWYSNRWVFGG GTKLTVL
26.	EGFRvIII_CCxC D3-scFc Bispecific HLE molecule	artificial	aa	QVQLVESGGGVVQSGRSLRLSCAASGFTFRNYGMH WVRQAPGKCLEWVAVIWDGSDKYADSVRGRFTI SRDNSKNTLYLQMNSLRAEDTAVYCARDGYDILT NPRDFDYWGQGLTVTVSSGGGSGGGGSGGGGSD TVMTQTPLSSHVTLGQPASISCRSSQSLVHSDGNTYL SWLQQRPGQPPRLIYRISRRFSGVPDRFSGSAGTD FTLEISRVEADVGVYCMQSTHVPRTFGCGTKVEIK SGGGGSEVQLVESGGGLVQPGGSLKLSAASGFTFN KYAMNWVRQAPGKLEWVARIRSKYNNYATYYAD SVKDRFTISRDDSKNTAYLQMNNLKTEDTAVYCVR HGNFGNSYISYWAYWGQGLTVTVSSGGGSGGGG SGGGGSQTVVTQEPLTVSPGGTVTLTCGSSTGAVTS GNYPNWVQKPGQAPRGLIGGTKFLAPGTPARFSG SLLGGKAALTLSGVQPEDEAEYYCVLWYSNRWVFGG GTKLTVLGGGGDKHTCPPCPAPELLGGPSVFLFPPK PKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGV EVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKE YKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRE EMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYK TTPPVLDSDGSFFLYSKLTVDKSRWQQGNVFCFSVM HEALHNHYTQKSLSLSPGKGGGGSGGGGSGGGGSG GGGSGGGGSGGGGSDKHTCPPCPAPELLGGPSVFL FPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWY VDGVEVHNAKTKPCEEQYGSTYRCVSVLTVLHQDW LNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTL PPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQP ENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNV FCFSVMHEALHNHYTQKSLSLSPGK
27.	MSLN_5 VH CDR1	artificial	aa	DYYMT
28.	MSLN_5 VH CDR2	artificial	aa	YISSSGSTIYYADSVKG
29.	MSLN_5 VH CDR3	artificial	aa	DRNSHFDY
30.	MSLN_5 VL CDR1	artificial	aa	RASQGINTWLA
31.	MSLN_5 VL CDR2	artificial	aa	GASGLQS
32.	MSLN_5 VL CDR3	artificial	aa	QQAKSFPRT

33.	MSLN_5 VH	artificial	aa	QVQLVESGGGLVKPGGSLRLSCAASGFTFSDYYMT WIRQAPGKGLEWLSYISSSGSTIYYADSVKGRFTISR DNAKNSLFLQMNSLRAEDTAVYYCARDNRNSHFDY WGQGTLLTVSS
34.	MSLN_5 VL	artificial	aa	DIQMTQSPSSVSASVGDRVITITCRASQGINTWLA WYQQKPGKAPKLLIYGASGLQSGVPSRFSGSGSGT DFTLTISLQPEDFATYYCQAKSFPRFTGQGTKVEI K
35.	MSLN_5 scFv	artificial	aa	QVQLVESGGGLVKPGGSLRLSCAASGFTFSDYYMT WIRQAPGKGLEWLSYISSSGSTIYYADSVKGRFTISR DNAKNSLFLQMNSLRAEDTAVYYCARDNRNSHFDY WGQGTLLTVSSGGGGSGGGSGGGGSDIQMTQ SPSSVSASVGDRVITITCRASQGINTWLAWYQQK GKAPKLLIYGASGLQSGVPSRFSGSGSGTDFTLTIS LQPEDFATYYCQAKSFPRFTGQGTKVEIK
36.	MSLN_5x12C0 bispecific molecule	artificial	aa	QVQLVESGGGLVKPGGSLRLSCAASGFTFSDYYMT WIRQAPGKGLEWLSYISSSGSTIYYADSVKGRFTISR DNAKNSLFLQMNSLRAEDTAVYYCARDNRNSHFDY WGQGTLLTVSSGGGGSGGGSGGGGSDIQMTQ SPSSVSASVGDRVITITCRASQGINTWLAWYQQK GKAPKLLIYGASGLQSGVPSRFSGSGSGTDFTLTIS LQPEDFATYYCQAKSFPRFTGQGTKVEIKSGGGG SEVQLVESGGGLVQPGGSLKLSAASGFTFNKYAM NWVRQAPGKGLEWVARIRSKYNNYATYYADSVKD RFTISRDDSKNTAYLQMNNLKTEDTAVYYCVRHGN FGNSYISYWAYWGQGTLLTVSSGGGGSGGGGSG GGGSQTVVTQEPSTVSPGGTVTLTCSSTGAVTS GNYPNWVQQKPGQAPRGLIGGTKFLAPGTPARFS GSLGGAALTLSGVQPEDEAEYCVLWYSNRWVF GGGKLTVL
37.	MSLN_5xCD3- scFc Bispecific HLE molecule	artificial	aa	QVQLVESGGGLVKPGGSLRLSCAASGFTFSDYYMT WIRQAPGKGLEWLSYISSSGSTIYYADSVKGRFTISR DNAKNSLFLQMNSLRAEDTAVYYCARDNRNSHFDY WGQGTLLTVSSGGGGSGGGSGGGGSDIQMTQ SPSSVSASVGDRVITITCRASQGINTWLAWYQQK GKAPKLLIYGASGLQSGVPSRFSGSGSGTDFTLTIS LQPEDFATYYCQAKSFPRFTGQGTKVEIKSGGGG SEVQLVESGGGLVQPGGSLKLSAASGFTFNKYAM NWVRQAPGKGLEWVARIRSKYNNYATYYADSVKD RFTISRDDSKNTAYLQMNNLKTEDTAVYYCVRHGN FGNSYISYWAYWGQGTLLTVSSGGGGSGGGGSG GGGSQTVVTQEPSTVSPGGTVTLTCSSTGAVTS GNYPNWVQQKPGQAPRGLIGGTKFLAPGTPARFS GSLGGAALTLSGVQPEDEAEYCVLWYSNRWVF GGGKLTVLGGGGDKHTCPPCPAPELLGGPSVFL FPPKPKDTLMISRTPEVTCVVDVSHEDPEVKFNW

				YVDGVEVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQGQGNVFSCSVMHEALHNHYTQKSLSLSPGKGGGGSGGGGGSGGGGGSGGGGSDKHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQGQGNVFSCSVMHEALHNHYTQKSLSLSPGK
38.	MSLN_5_CCxC D3-scFc Bispecific HLE molecule	artificial	aa	QVQLVESGGGLVPGGSLRLSCAASGFTFSDHYMSWIRQAPGKCLEWFYSYISSGGIYYADSVKGRFTISRDNAKNSLYLQMNSLRAEDTAVYYCARDVGSFHDYWGQGTLTIVSSGGGGSGGGGSGGGGSDIQMTQSPSSVSASVGDRTITCRASQDISRWLAWYQQKPGKAPKLLISAASRLQSGVPSRFSGSGSGTDFLTISLQPEDFAIYYCQQAQKSFPRTFGCGTKVEIKSGGGGSEVQLVESGGGLVQPGGSLKLSCAASGFTFNKYAMNWRVQAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSKNTAYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQGTLTIVSSGGGGSGGGGSGGGGGSQTVVTQEPSTVSPGGTVTLTCSSTGAVTSGNYPNWWVQQKPGQAPRGLIGGTKFLAPGTPARFSGSLGGKAALTLSGVQPEDEAEYCVLWYSNRWVFGGGTKLTVLGGGGDKHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQGQGNVFSCSVMHEALHNHYTQKSLSLSPGK
39.	CDR-H1 CDH19 65254.007	artificial	aa	SYGMH

40.	CDR-H2 CDH19 65254.007	artificial	aa	FIWYEGSNKYAESVKD
41.	CDR-H3 CDH19 65254.007	artificial	aa	RAGIIGTIGYYYGMDV
42.	CDR-L1 CDH19 65254.007	artificial	aa	SGDRLGEKYTS
43.	CDR-L2 CDH19 65254.007	artificial	aa	QDTRPS
44.	CDR-L3 CDH19 65254.007	artificial	aa	QAWESSTVV
45.	VH CDH19 65254.007	artificial	aa	QVQLVESGGGVVQPGGSLRLSCAASGFTFSSYGM HWVRQAPGKGLEWVAFIWIWYEGSNKYAESVKDRF TISRDNKNTLYLQMNSLRAEDTAVYYCARRAGIIG TIGYYYGMDVWGQGTITVTVSS
46.	VL CDH19 65254.007	artificial	aa	SYELTQPPSVSVSPGQTASITCSGDRLGEKYTSWYQ QRPQGSPLLVIYQDTRPSGIPERFSGSNSGNTATL TISGTQAMDEADYYCQAWESSTVVFSGGKLTVL
47.	VH-VL CDH19 65254.007	artificial	aa	QVQLVESGGGVVQPGGSLRLSCAASGFTFSSYGM HWVRQAPGKGLEWVAFIWIWYEGSNKYAESVKDRF TISRDNKNTLYLQMNSLRAEDTAVYYCARRAGIIG TIGYYYGMDVWGQGTITVTVSSGGGGSGGGSGG GGSSYELTQPPSVSVSPGQTASITCSGDRLGEKYTS WYQQRPGQSPLLVIYQDTRPSGIPERFSGNSGN TATLTISGTQAMDEADYYCQAWESSTVVFSGGKLT TVLS
48.	CDH19 65254.007 x I2C	artificial	aa	QVQLVESGGGVVQPGGSLRLSCAASGFTFSSYGM HWVRQAPGKGLEWVAFIWIWYEGSNKYAESVKDRF TISRDNKNTLYLQMNSLRAEDTAVYYCARRAGIIG TIGYYYGMDVWGQGTITVTVSSGGGGSGGGSGG GGSSYELTQPPSVSVSPGQTASITCSGDRLGEKYTS WYQQRPGQSPLLVIYQDTRPSGIPERFSGNSGN TATLTISGTQAMDEADYYCQAWESSTVVFSGGKLT TVLSGGGGSEVQLVESGGGLVQPGGSLKLSAASG FTFNKYAMNWVRQAPGKLEWVARIRSKYNNYA TYYADSVKDRFTISRDDSKNTAYLQMNNLKTEDTA VYYCVRHGNFGNSYISYWAYWGQGLTVTVSSGGG SGGGGGSGGGGSQTVVTQEPSTVSPGGTVTLTC GSSTGAVTSGNYPNWVQKPGQAPRGLIGGKFL APGTPARFSGSLLGGKAALTLSGVQPEDEAEYCVL WYSNRWVFSGGKLTVLHHHHHH
49.	CDH19 65254.007 x I2C -scFc	artificial	aa	QVQLVESGGGVVQPGGSLRLSCAASGFTFSSYGMH WVRQAPGKGLEWVAFIWIWYEGSNKYAESVKDRFTIS RDNKNTLYLQMNSLRAEDTAVYYCARRAGIIGTIGY

	Bispecific HLE molecule			YYGMDVWGQGTTVTVSSGGGGSGGGGSGGGGSS YELTQPPSVSVSPGQTASITCSGDLGEKYTSWYQQR PGQSPLLVIYQDTRPSGIPERFSGSNSGNTATLTISG TQAMDEADYYCQAWESSTVVFGGGTCLTVLSGGGG SEVQLVESGGGLVQPGGSLKLSAASGFTFNKYAMN WVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRF TISRDDSKNTAYLQMNNLKTEDTAVYYCVRHGNFGN SYISYWAYWGQGLTVTVSSGGGGSGGGGSGGGGSS QTVVTQEPSLTVSPGGTVTLTCSSTGAVTSGNYPN WVQQKPGQAPRGLIGGTKFLAPGTPARFSGSLLGGK AALTLSGVQPEDEAEYYCVLWYSNRWVFGGGTKLTV LGGGGDKHTHTCPPCPAPELLGGPSVFLFPPKPKDTL MISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHN AKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCK VSNKALPAIEKTISKAKGQPREPQVYTLPPSREEMTK NQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTPP VLDSGDSFFLYSKLTVDKSRWQQGNVFSCSVMHEAL HNHYTQKSLSPGKGGGGSGGGGSGGGGSGGGG SGGGGSGGGGSDKHTHTCPPCPAPELLGGPSVFLFPP KPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDG VEVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNG KEYKCKVSNKALPAIEKTISKAKGQPREPQVYTLPPS REEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENN YKTPPVLDSDGSFFLYSKLTVDKSRWQQGNVFSCSV MHEALHNHYTQKSLSPGK
50.	CDH19 65254.007 x I2C - scFc_delGK Bispecific HLE molecule	artificial	aa	QVQLVESGGGVVQPGGSLRLSASGFTSSYGMH WVRQAPGKGLEWVAFIWEYSNKKYAESVKDRFTIS RDNSKNTLYLQMNSLRAEDTAVYYCARRAGIIGTIGY YYGMDVWGQGTTVTVSSGGGGSGGGGSGGGGSS YELTQPPSVSVSPGQTASITCSGDLGEKYTSWYQQR PGQSPLLVIYQDTRPSGIPERFSGSNSGNTATLTISG TQAMDEADYYCQAWESSTVVFGGGTCLTVLSGGGG SEVQLVESGGGLVQPGGSLKLSAASGFTFNKYAMN WVRQAPGKGLEWVARIRSKYNNYATYYADSVKDRF TISRDDSKNTAYLQMNNLKTEDTAVYYCVRHGNFGN SYISYWAYWGQGLTVTVSSGGGGSGGGGSGGGGSS QTVVTQEPSLTVSPGGTVTLTCSSTGAVTSGNYPN WVQQKPGQAPRGLIGGTKFLAPGTPARFSGSLLGGK AALTLSGVQPEDEAEYYCVLWYSNRWVFGGGTKLTV LGGGGDKHTHTCPPCPAPELLGGPSVFLFPPKPKDTL MISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHN AKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCK VSNKALPAIEKTISKAKGQPREPQVYTLPPSREEMTK NQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTPP VLDSGDSFFLYSKLTVDKSRWQQGNVFSCSVMHEAL HNHYTQKSLSPGKGGGGSGGGGSGGGGSGGGGSG

				GGGSGGGGSDKHTHTCPPCPAPPELLGGPSVFLFPPKP KDTLMISRTPEVTCVVDVSHEDPEVKFNWYVDGVE VHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEY KCKVSNKALPAPIEKTISKAKGQPREPVYTLPPSREE MTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKT TPPVLDSDGSFFLYSKLTVDKSRWQQGNVFCSSVMH EALHNHYTQKSLSLSPGK
51.	CDH19 65254.007_CC x I2C -scFc VH	artificial	aa	QVQLVESGGGVVQPGGSLRLSCAASGFTFSSYGMH WVRQAPGKCLEWVAFIWIYEGSNKYAESVKDRFTIS RDNSKNTLYLQMNSLRAEDTAVYYCARRAGIIGTIGY YYGMDVWGQGTITVTVSS
52.	CDH19 65254.007_CC x I2C -scFc VL	artificial	aa	SYELTQPPSVSVSPGQTASITCSGDRLEKYTSWYQQ RPGQSPLLVIYQDTRKPSGIPERFSGSNSGNTATLTIS GTQAMDEADYYCQAWESSTVVFSGCTKLTVL
53.	CDH19 65254.007_CC x I2C -scFc scFv	artificial	aa	QVQLVESGGGVVQPGGSLRLSCAASGFTFSSYGMH WVRQAPGKCLEWVAFIWIYEGSNKYAESVKDRFTIS RDNSKNTLYLQMNSLRAEDTAVYYCARRAGIIGTIGY YYGMDVWGQGTITVTVSSGGGGSGGGSGGGSS YELTQPPSVSVSPGQTASITCSGDRLEKYTSWYQQR PGQSPLLVIYQDTRKPSGIPERFSGSNSGNTATLTISG TQAMDEADYYCQAWESSTVVFSGCTKLTVL
54.	CDH19 65254.007_CC x I2C -scFc Bispecific molecule	artificial	aa	QVQLVESGGGVVQPGGSLRLSCAASGFTFSSYGMH WVRQAPGKCLEWVAFIWIYEGSNKYAESVKDRFTIS RDNSKNTLYLQMNSLRAEDTAVYYCARRAGIIGTIGY YYGMDVWGQGTITVTVSSGGGGSGGGSGGGSS YELTQPPSVSVSPGQTASITCSGDRLEKYTSWYQQR PGQSPLLVIYQDTRKPSGIPERFSGSNSGNTATLTISG TQAMDEADYYCQAWESSTVVFSGCTKLTVLSSGGGG SEVQLVESGGGLVQPGGSLKLSAASGFTFNKYAMN WVRQAPGKLEWVARIRSKYNNYATYYADSVKDRF TISRDDSKNTAYLQMNNLKTEDTAVYYCVRHGNFGN SYISYWAYWGQGLTVTVSSGGGGSGGGSGGGSS QTVVTQEPSTVSPGGTVTLTCSSTGAVTSGNYPN WVQQKPGQAPRGLIGGTFKFLAPGTPARFSGSLLGGK AALTLSGVQPEDEAEYYCVLWYSNRWVFGGGTKLT V
55.	CDH19 65254.007_CC x I2C -scFc Bispecific HLE molecule	artificial	aa	QVQLVESGGGVVQPGGSLRLSCAASGFTFSSYGMH WVRQAPGKCLEWVAFIWIYEGSNKYAESVKDRFTIS RDNSKNTLYLQMNSLRAEDTAVYYCARRAGIIGTIGY YYGMDVWGQGTITVTVSSGGGGSGGGSGGGSS YELTQPPSVSVSPGQTASITCSGDRLEKYTSWYQQR PGQSPLLVIYQDTRKPSGIPERFSGSNSGNTATLTISG TQAMDEADYYCQAWESSTVVFSGCTKLTVLSSGGGG SEVQLVESGGGLVQPGGSLKLSAASGFTFNKYAMN WVRQAPGKLEWVARIRSKYNNYATYYADSVKDRF TISRDDSKNTAYLQMNNLKTEDTAVYYCVRHGNFGN

				SYISYWAYWGQGLTVTVSSGGGGSGGGGSGGGGS QTVVTQEPSLTVSPGGTVTLTCGSSTGAVTSGNYPN WVQQKPGQAPRGLIGGTKFLAPGTPARFSGSLLGGK AALTLSGVQPEDEAEYYCVLWYSNRWVFGGGTKLTV LGGGGDKTHTCPPCPAPELLGGPSVFLFPPKPKDTL MISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHN AKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCK VSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTK NQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPP VLDSGDSFFLYSKLTVDKSRWQQGNVFSCSVMHEAL HNHYTQKSLSLSPGKGGGGSGGGGSGGGGSGGGG SGGGSGGGGSDKTHTCPPCPAPELLGGPSVFLFPP KPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDG VEVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNG KEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPS REEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENN YKTPPVLDSDGSFFLYSKLTVDKSRWQQGNVFSCSV MHEALHNHYTQKSLSLSPGK
56.	CDH19 65254.007_CC x I2C - scFc_delGK Bispecific HLE molecule	artificial	aa	QVQLVESGGGVVQPGGSLRLSCAASGFTSSYGMH WVRQAPGKCLEWVAFIWIYEGSNKYAESVKDRFTIS RDNSKNTLYLQMNSLRAEDTAVYYCARRAGIIGTIGY YYGMDVWGQGTITVTVSSGGGGSGGGGSGGGGSS YELTQPPSVSPGQTASITCSGDRLEKYSWYQQR PGQSPLLVIYQDTRKPSGIPERFSGSNSGNTATLTISG TQAMDEADYYCAWESSTVVFSGTKLTVLSSGGG SEVQLVESGGGLVQPGGSLKLSAASGFTFNKYAMN WVRQAPGKLEWVARIRSKYNNYATYYADSVKDRF TISRDDSKNTAYLQMNNLKTEDTAVYYCVRHGNFGN SYISYWAYWGQGLTVTVSSGGGGSGGGGSGGGGS QTVVTQEPSLTVSPGGTVTLTCGSSTGAVTSGNYPN WVQQKPGQAPRGLIGGTKFLAPGTPARFSGSLLGGK AALTLSGVQPEDEAEYYCVLWYSNRWVFGGGTKLTV LGGGGDKTHTCPPCPAPELLGGPSVFLFPPKPKDTL MISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHN AKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCK VSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTK NQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPP VLDSGDSFFLYSKLTVDKSRWQQGNVFSCSVMHEAL HNHYTQKSLSLSPGGGGSGGGGSGGGGSGGGGSG GGGSGGGGSDKTHTCPPCPAPELLGGPSVFLFPPK KDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVE VHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEY KCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREE MTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKT TPPVLDSDGSFFLYSKLTVDKSRWQQGNVFSCSVMH EALHNHYTQKSLSLSPGK

57.	FLT3_7 A8xCD3-scFc VH CDR1	artificial	aa	NARMGVS
58.	FLT3_7 A8xCD3-scFc VH CDR2	artificial	aa	HIFSNDEKSYSTSLKN
59.	FLT3_7 A8xCD3-scFc VH CDR3	artificial	aa	IVGYGSGWYGFFDY
60.	FLT3_7 A8xCD3-scFc VL CDR1	artificial	aa	RASQGIRNDLG
61.	FLT3_7 A8xCD3-scFc VL CDR2	artificial	aa	AASTLQS
62.	FLT3_7 A8xCD3-scFc VL CDR3	artificial	aa	LQHNSYPLT
63.	FLT3_7 A8xCD3-scFc VH	artificial	aa	QVTLKESGPTLVKPTETLTLTCTLSGFSLNNARMGVS WIRQPPGKCLEWLAHIFSNDEKSYSTSLKNRLTISKDS SKTQVVLMTNVDPVDTATYYCARIVGYGSGWYGFF DYWGQGLTVTVSS
64.	FLT3_ A8-scFc VL	artificial	aa	DIQMTQSPSSLSASVGDRTITCRASQGIRNDLGWY QQKPGKAPKRLIYAASTLQSGVPSRFSGSGSGTEFTLT ISSLPEDFATYYCLQHNSYPLTFGCGTKVEIK
65.	FLT3_7 A8xCD3- scFv	artificial	aa	QVTLKESGPTLVKPTETLTLTCTLSGFSLNNARMGVS WIRQPPGKCLEWLAHIFSNDEKSYSTSLKNRLTISKDS SKTQVVLMTNVDPVDTATYYCARIVGYGSGWYGFF DYWGQGLTVTVSSGGGGSGGGSGGGGSDIQMT QSPSSLSASVGDRTITCRASQGIRNDLGWYQQKPG KAPKRLIYAASTLQSGVPSRFSGSGSGTEFTLTISSLP EDFATYYCLQHNSYPLTFGCGTKVEIK
66.	FLT3_7 A8xCD3 Bispecific molecule	artificial	aa	QVTLKESGPTLVKPTETLTLTCTLSGFSLNNARMGVS WIRQPPGKCLEWLAHIFSNDEKSYSTSLKNRLTISKDS SKTQVVLMTNVDPVDTATYYCARIVGYGSGWYGFF DYWGQGLTVTVSSGGGGSGGGSGGGGSDIQMT QSPSSLSASVGDRTITCRASQGIRNDLGWYQQKPG KAPKRLIYAASTLQSGVPSRFSGSGSGTEFTLTISSLP EDFATYYCLQHNSYPLTFGCGTKVEIKSGGGGSEVQL VESGGGLVQPGGSLKLSAASGFTFNKYAMNWRQ APGKLEWVARIRSKYNNYATYYADSVKDRFTISRDD SKNTAYLQMNNLKTEDAVYYCVRHGNFGNSYISY WAYWGQGLTVTVSSGGGGSGGGSGGGGSGTQTVV TQEPSLTVPGGTVTLTCSSTGAVTSGNYPNWWVQQ KPGQAPRGLIGGTKFLAPGTPARFSGSLLGGKAALT SGVQPEDEAEYYCVLWYSNRWVFGGGTKLTVL

67.	FLT3_7 A8xCD3-scFc Bispecific HLE molecule	artificial	aa	QVTLKESGPTLVKPTETLTCTLSGFSLNNARMGVS WIRQPPGKCLEWLAHIFSNDEKSYSTSLKNRLTISKDS SKTQVVLMTNVDVDTATYYCARIVGYGSGWYGFF DYWGQGTTLTVSSGGGGSGGGGSGGGGSDIQMT QSPSSLSASVGDRVTITCRASQGIRNDLGWYQQKPG KAPKRLIYAASLTQSGVPSRFSGSGSGTEFTLTISLQP EDFATYYCLQHNSYPLTFGCGTKVEIKSGGGGSEVQL VESGGGLVQPGGSLKLSAASGFTFNKYAMNWVRQ APGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDD SKNTAYLQMNNLKTEDTAVYYCVRHGNFGNSYISY WAYWGQGTTLTVSSGGGGSGGGGSGGGGSGQTVV TQEPSLTVSPGGTVTLTCSSTGAVTSGNYPNWVQQ KPGQAPRGLIGGTKFLAPGTPARFSGSLLGGKAALT SGVQPEDEAEYYCVLWYSNRWVFGGGTKLTVLGGG GDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRT EVTQVVDVSHEDPEVKFNWYVDGVEVHNAKTKPC EEQYGSTYRCVSVTLVHQQDWLNGKEYCKVSNKAL PAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLT CLVKGFYPSDIAVEWESNGQPENNYKTPPVLDSDG SFFLYSKLTVDKSRWQQGNVFSCSVMHEALHNHYT QKSLSLSPGKGGGGSGGGGSGGGGSGGGGSGGGG SGGGGSDKTHTCPPCPAPELLGGPSVFLFPPKPKDTL MISRTPEVTQVVDVSHEDPEVKFNWYVDGVEVHN AKTKPCEEQYGSTYRCVSVTLVHQQDWLNGKEYCK VSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTK NQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTPP VLDSDGSFFLYSKLTVDKSRWQQGNVFSCSVMHEAL HNHYTQKSLSLSPGK
68	VH CDR1 DLL3_1_CC_del GK	artificial	aa	SYYS
69	VH CDR2 DLL3_1_CC_del GK	artificial	aa	YVYSGTTNYPNPSLKS
70	VH CDR3 DLL3_1_CC_del GK	artificial	aa	IAVTGFYFDY
71	VL CDR1 DLL3_1_CC_del GK	artificial	aa	RASQRVNNNYLA
72	VL CDR2 DLL3_1_CC_del GK	artificial	aa	GASSRAT

73	VL CDR3 DLL3_1_CC_del GK	artificial	aa	QQYDRSPLT
74.	VH DLL3_1_CC_del GK	artificial	aa	QVQLQESGPGLVKPSSETLSLTCTVSGGSISSYYWSWI RQPPGKCLEWIGYVYSGTTNYPNPSLKSRTISVDTSK NQFSLKLSSVTAADTAVYYCASIATGIFYFDYWGQG TLVTVSS
75.	VL DLL3_1_CC_del GK	artificial	aa	EIVLTQSPGTLSLSPGERVTLSRASQRVNNNYLAWY QQRPGQAPRLIYGASSRATGIPDRFSGSGSGTDFTL TISRLEPEDFAVYYCQQYDRSPLTFGCGTKLEIK
76.	DLL3_1_CC_del GK	artificial	aa	QVQLQESGPGLVKPSSETLSLTCTVSGGSISSYYWSWI RQPPGKCLEWIGYVYSGTTNYPNPSLKSRTISVDTSK NQFSLKLSSVTAADTAVYYCASIATGIFYFDYWGQG TLVTVSSGGGGSGGGSGGGGSEIVLTQSPGTLSLSP GERVTLSRASQRVNNNYLAWYQQRPGQAPRLIY GASSRATGIPDRFSGSGSGTDFTLTISRLEPEDFAVYY CQQYDRSPLTFGCGTKLEIK
77.	DLL3_1_CCxCD 3_delGK Bispecific molecule	artificial	aa	QVQLQESGPGLVKPSSETLSLTCTVSGGSISSYYWSWI RQPPGKCLEWIGYVYSGTTNYPNPSLKSRTISVDTSK NQFSLKLSSVTAADTAVYYCASIATGIFYFDYWGQG TLVTVSSGGGGSGGGSGGGGSEIVLTQSPGTLSLSP GERVTLSRASQRVNNNYLAWYQQRPGQAPRLIY GASSRATGIPDRFSGSGSGTDFTLTISRLEPEDFAVYY CQQYDRSPLTFGCGTKLEIKSGGGGSEVQLVESGGGL VQPGGSLKLSAASGFTFNKYAMNWRQAPGKGLE WVARIRSKYNNYATYYADSVKDRFTISRDDSKNTAYL QMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQ GTLVTVSSGGGGSGGGSGGGGSGTQVVTQEPSTV SPGGTVTLTCSSTGAVTSGNYPNWWQKPGQAPR GLIGGKFLAPGTPARFSGSLLGGKAALTLSGVQPED EAEYYCVLWYSNRWVFGGGTKLTVL
78.	DLL3_1_CCxCD 3-scFc_delGK Bispecific HLE molecule	artificial	aa	QVQLQESGPGLVKPSSETLSLTCTVSGGSISSYYWSWI RQPPGKCLEWIGYVYSGTTNYPNPSLKSRTISVDTSK NQFSLKLSSVTAADTAVYYCASIATGIFYFDYWGQG TLVTVSSGGGGSGGGSGGGGSEIVLTQSPGTLSLSP GERVTLSRASQRVNNNYLAWYQQRPGQAPRLIY GASSRATGIPDRFSGSGSGTDFTLTISRLEPEDFAVYY CQQYDRSPLTFGCGTKLEIKSGGGGSEVQLVESGGGL VQPGGSLKLSAASGFTFNKYAMNWRQAPGKGLE WVARIRSKYNNYATYYADSVKDRFTISRDDSKNTAYL QMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQ GTLVTVSSGGGGSGGGSGGGGSGTQVVTQEPSTV SPGGTVTLTCSSTGAVTSGNYPNWWQKPGQAPR GLIGGKFLAPGTPARFSGSLLGGKAALTLSGVQPED EAEYYCVLWYSNRWVFGGGTKLTVLGGGGDKHTC PPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVV

				DVSHEDPEVKFNWYVDGVEVHNAKTKPCEEQYGST YRCVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTI SKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGF YPSDIAVEWESNGQPENNYKTTTPVLDSDGSFFLYSK LTVDKSRWQQGNVFCFSVMHEALHNHYTQKSLSLS PGGGGSGGGGSGGGGSGGGGSGGGGSGGGGSGDK THTCPPCPAPELLGGSVFLFPPKPKDTLMISRTPEVT CVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEEQ YGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALPAPI EKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLV KGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGSFFL YSKLTVDKSRWQQGNVFCFSVMHEALHNHYTQKSL SLSPGK
79.	VH CDR1 CD19 97-G1RE-C2	artificial	aa	SYGMH
80.	VH CDR2 CD19 97-G1RE-C2	artificial	aa	VISYEGSNKYAESVKG
81.	VH CDR3 CD19 97-G1RE-C2	artificial	aa	DRGTIFGNYLEV
82.	VH CD19 97- G1RE-C2 CC	artificial	aa	QVQLVESGGGVVQPGRSLRLSCAASGFTFSSYGMH WVRQAPGKCLEWVAVISYEGSNKYAESVKGRTIS RDNSKNTLYLQMNSLRDEDTAVYYCARDRTIFGNY GLEVWGQGTTVTVSS
83.	VL CDR1 CD19 97-G1RE-C2	artificial	aa	RSSQSLHKNFNYLD
84.	VL CDR2 CD19 97-G1RE-C2	artificial	aa	LGSNRAS
85.	VL CDR3 CD19 97-G1RE-C2	artificial	aa	MQALQTPFT
86.	VL CD19 97- G1RE-C2 CC	artificial	aa	DIVMTQSPLSLPVISGEPASISCRSSQSLHKNFNYL DWYLQKPGQSPQLLIYLGSNRASGVPDRFSGSGSGT DFTLKISRVEAEDVGVYYCMQALQTPFTFGCGTKVDI K
87.	CD19 97-G1RE- C2 CC x I2C0	artificial	aa	MDMRVPAQLLGLLLWLGRGCDIVMTQSPLSLPVI SGEPASISCRSSQSLHKNFNYLDWYLQKPGQSPQL LIYLGSNRASGVPDRFSGSGSGTDFTLKISRVEAEDVG VYYCMQALQTPFTFGCGTKVDIKGGGGSGGGGSGG GGSQVQLVESGGGVVQPGRSLRLSCAASGFTFSSYG MHWVRQAPGKCLEWVAVISYEGSNKYAESVKGRTIS RDNSKNTLYLQMNSLRDEDTAVYYCARDRTIFG NYGLEVWGQGTTVTVSSGGGGSEVQLVESGGGLVQ PGGSLKLSAASGFTFNKYAMNWVRQAPGKGLEW VARIRSKYNNYATYYADSVKDRFTISRDDSKNTAYLQ MNNLKTEDAVYYCVRHGNFGNSYISYWAYWGQG TLVTVSSGGGGSGGGGSGGGGSGTQVTVQEPSTVSP GGTVTLTCSSTGAVTSGNYPNWWQQKPGQAPRGL

				IGGTKFLAPGTPARFSGSLLGGKAALTLSGVQPEDEA EYYCVLWYSNRWVFSGGKLTVL
88.	CD19 97-G1RE- C2 CC x I2C0- scFc	artificial	aa	MDMRVPAQLLGLLLWLRGARCDIVMTQSPLSLPVI SGEPASISCRSSQSLHKNFNYLDWYLQKPGQSPQL LIYLGSNRASGVDPDRFSGSGSGTDFTLKISRVEAEDVG VYYCMQALQTPFTFGCGTKVDIKGGGGSGGGGSGG GGSQVQLVESGGGVVQPGRSLRLSCAASGFTFSSYG MHWVRQAPGKCLEWVAVISYEGSNKYAESVKGRF TISRDNSKNTLYLQMNSLRDEDTAVYYCARDRTIFG NYGLEVWGQGTTVTVSSGGGGSEVQLVESGGGLVQ PGGSLKLSAASGFTFNKYAMNWVRQAPGKGLEW VARIRSKYNNYATYYADSVKDRFTISRDDSKNTAYLQ MNNLKTEDTAVYYCVRHGNFGNSYISYWAYWGQG TLVTVSSGGGGSGGGGSGGGGSGTQVTVQEPSLTVSP GGTVTLTCGSSTGAVTSGNYPNWVQKPGQAPRGL IGGTKFLAPGTPARFSGSLLGGKAALTLSGVQPEDEA EYYCVLWYSNRWVFSGGKLTVLGGGGDKTHTCPP CPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDV SHEDPEVKFNWYVDGVEVHNAKTKPCEEQYGSTYR CVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISK AKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYP SDIAVEWESNGQPENNYKTTTPVLDSDGSFFLYSKLT VDKSRWQQGNVFSCSVMHEALHNHYTQKSLSLSPG KGGGGSGGGGSGGGGSGGGGSGGGGSGGGGSGDK THTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVT CVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEEQ YGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALPAPI EKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLV KGFYP SDIAVEWESNGQPENNYKTTTPVLDSDGSFFL YSKLTVDKSRWQQGNVFSCSVMHEALHNHYTQKS LSLSPGK
89.	VH CDR1 CDH3 G8A 6-B12	artificial	aa	SYPIN
90.	VH CDR2 CDH3 G8A 6-B12	artificial	aa	VIWTGGGTNYASSVKG
91.	VH CDR3 CDH3 G8A 6-B12	artificial	aa	SRGVYDFDGRGAMDY
92.	VL CDR1 CDH3 G8A 6-B12	artificial	aa	KSSQSLLYSSNQKNYFA
93.	VL CDR2 CDH3 G8A 6-B12	artificial	aa	WASTRES
94.	VL CDR3 CDH3 G8A 6-B12	artificial	aa	QQYYSYPYT
95.	VH CDH3 G8A 6-B12	artificial	aa	EVQLLESGGGLVQPGGSLRLSCAASGFSFSSYPINWV RQAPGKGLEWVGVIWTGGGTNYASSVKGRTISR

				NSKNTVYLQMNSLRAEDTAVYYCAKSRGVYDFDGR GAMDYWGQGTLLTVSS
96.	VL CDH3 G8A 6-B12	artificial	aa	DIVMTQSPDSLAVSLGERATINCKSSQSLLYSSNQKN YFAWYQQKPGQPPKLLIYWASTRESGVPDRFSGSGS GTDFTLTISLQAEDVAVYYCQQYYSYPYTFGQGTKL EIK
97.	CDH3 G8A 6- B12 scFv	artificial	aa	EVQLLESGGGLVQPGGSLRLSCAASGFSFSSYPINWV RQAPGKGLEWVGVIWTGGGTNYASSVKGRFTISRD NSKNTVYLQMNSLRAEDTAVYYCAKSRGVYDFDGR GAMDYWGQGTLLTVSSGGGGSGGGSGGGGSDI VMTQSPDSLAVSLGERATINCKSSQSLLYSSNQKNYF AWYQQKPGQPPKLLIYWASTRESGVPDRFSGSGSGT DFTLTISLQAEDVAVYYCQQYYSYPYTFGQGTKLEIK
98.	CDH3 G8A 6- B12 x I2C0 bispecific molecule	artificial	aa	EVQLLESGGGLVQPGGSLRLSCAASGFSFSSYPINWV RQAPGKGLEWVGVIWTGGGTNYASSVKGRFTISRD NSKNTVYLQMNSLRAEDTAVYYCAKSRGVYDFDGR GAMDYWGQGTLLTVSSGGGGSGGGSGGGGSDI VMTQSPDSLAVSLGERATINCKSSQSLLYSSNQKNYF AWYQQKPGQPPKLLIYWASTRESGVPDRFSGSGSGT DFTLTISLQAEDVAVYYCQQYYSYPYTFGQGTKLEIK SGGGGSEVQLVESGGGLVQPGGSLKLSCAASGFTFN KYAMNWRQAPGKGLEWVARIRSKYNNYATYYAD SVKDRFTISRDDSKNTAYLQMNNLKTEDTAVYYCVR HGNFGNSYISYWAYWGQGTLLTVSSGGGGSGGGG SGGGGSQTVVTQEPSTVSPGGTVTLTCSSTGAVTS GNYPNWWVQQKPGQAPRGLIGGTFKFLAPGTPARFSG SLLGGKAALTLSGVQPEDEAEYCVLWYSNRWVFGG GKLTVL
99.	CDH3 G8A 6- B12 x I2C0 bispecific molecule HLE	artificial	aa	EVQLLESGGGLVQPGGSLRLSCAASGFSFSSYPINWV RQAPGKGLEWVGVIWTGGGTNYASSVKGRFTISRD NSKNTVYLQMNSLRAEDTAVYYCAKSRGVYDFDGR GAMDYWGQGTLLTVSSGGGGSGGGSGGGGSDI VMTQSPDSLAVSLGERATINCKSSQSLLYSSNQKNYF AWYQQKPGQPPKLLIYWASTRESGVPDRFSGSGSGT DFTLTISLQAEDVAVYYCQQYYSYPYTFGQGTKLEIK SGGGGSEVQLVESGGGLVQPGGSLKLSCAASGFTFN KYAMNWRQAPGKGLEWVARIRSKYNNYATYYAD SVKDRFTISRDDSKNTAYLQMNNLKTEDTAVYYCVR HGNFGNSYISYWAYWGQGTLLTVSSGGGGSGGGG SGGGGSQTVVTQEPSTVSPGGTVTLTCSSTGAVTS GNYPNWWVQQKPGQAPRGLIGGTFKFLAPGTPARFSG SLLGGKAALTLSGVQPEDEAEYCVLWYSNRWVFGG GKLTVLGGGGDKHTCPPCPAPELLGGPSVFLFPPK PKDTLMISRTPEVTCVVDVSHEDPEVKFNWYVDGV EVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKE YKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRE

				EMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYK TTPPVLDSDGSFFLYSKLTVDKSRWQQGNVFCFSVM HEALHNHYTQKSLSLSPGKGGGGSGGGSGGGSGG GGGGSGGGSGGGSGDKHTCPPCPAPELLGGPSVFL FPPKPKDTLMISRTPEVTCVVDVSHEDPEVKFNWY VDGVEVHNAKTKPCEEQYGSTYRCVSVLTVLHQDW LNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTL PPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQP ENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNV SCFSVMHEALHNHYTQKSLSLSPGK
100.	BCMA A7 27- C4-G7 CDR1 VH	artificial	aa	NHIIH
101.	BCMA A7 27- C4-G7 CDR2 VH	artificial	aa	YINPYPGYHAYNEKFQG
102.	BCMA A7 27- C4-G7 CDR3 VH	artificial	aa	DGYRDTDLVDY
103.	BCMA A7 27- C4-G7 CDR1 VL	artificial	aa	QASQDISNYLN
104.	BCMA A7 27- C4-G7 CDR2 VL	artificial	aa	YTSRLHT
105.	BCMA A7 27- C4-G7 CDR3 VL	artificial	aa	QQGNTLPWT
106.	BCMA A7 27- C4-G7 CC (44/100) VH	artificial	aa	QVQLVQSGAEVKKPGASVKVSCASGYFTNHHIHW VRQAPGQCLEWMGYINPYPGYHAYNEKFQGRATM TSDTSTVYMELSSLRSEDTAVYYCARDGYRDTDV LDYWGQGTLLVTVSS
107.	BCMA A7 27- C4-G7 CC (44/100) VL	artificial	aa	DIQMTQSPSSLSASVGDRVTITCQASQDISNYLNWY QQKPGKAPKLLIYYTSRLHTGVPSRFSGSGSGDFTFT ISSLEPEDIATYYCQQGNTLPWTFGCGTKLEIK
108.	BCMA A7 27- C4-G7 CC (44/100) scFv	artificial	aa	QVQLVQSGAEVKKPGASVKVSCASGYFTNHHIHW VRQAPGQCLEWMGYINPYPGYHAYNEKFQGRATM TSDTSTVYMELSSLRSEDTAVYYCARDGYRDTDV LDYWGQGTLLVTVSSGGGGSGGGSGGGSGDIQMT QSPSSLSASVGDRVTITCQASQDISNYLNWYQQKPG KAPKLLIYYTSRLHTGVPSRFSGSGSGDFTFTISSLEPE DIATYYCQQGNTLPWTFGCGTKLEIK
109.	BCMA A7 27- C4-G7 CC (44/100) x I2C0 bispecific molecule	artificial	aa	QVQLVQSGAEVKKPGASVKVSCASGYFTNHHIHW VRQAPGQCLEWMGYINPYPGYHAYNEKFQGRATM TSDTSTVYMELSSLRSEDTAVYYCARDGYRDTDV LDYWGQGTLLVTVSSGGGGSGGGSGGGSGDIQMT QSPSSLSASVGDRVTITCQASQDISNYLNWYQQKPG KAPKLLIYYTSRLHTGVPSRFSGSGSGDFTFTISSLEPE DIATYYCQQGNTLPWTFGCGTKLEIKSGGGGSEVQL VESGGGLVQPGGSLKLSCAASGFTFNKYAMNWVRQ APGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDD

				SKNTAYLQMNNLKTEDTAVYYCVRHGNFGNSYISY WAYWGQGTTLTVSSGGGGSGGGGSGGGGSQTVV TQEPSLTVSPGGTVTLTCSSTGAVTSGNYPNWWVQQ KPGQAPRGLIGGTKFLAPGTPARFSGSLLGGKAALT SGVQPEDEAEYYCVLWYSNRWVFGGGTKLTVL
110.	BCMA A7 27- C4-G7 CC (44/100) x I2C0-scFc bispesific molecule HLE	artificial	aa	QVQLVQSGAEVKKPGASVKVSCASGYTFTNHHHW VRQAPGQCLEWMGYINPYPGYHAYNEKFQGRATM TSDTSTSTVYMESSLRSEDTAVYYCARDGYRDTDV LDYWGQGTTLTVSSGGGGSGGGGSGGGGSDIQMT QSPSSLSASVGDRTITCQASQDISNYLNWYQQKPG KAPKLLIYYTSLRHTGVPSRFSGSGSGTDFTFTISSELEPE DIATYYCQQGNTLPWTFGCGTKVEIKSGGGGSEVQL VESGGGLVQPGGSLKLSAASGFTFNKYAMNWWVRQ APGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDD SKNTAYLQMNNLKTEDTAVYYCVRHGNFGNSYISY WAYWGQGTTLTVSSGGGGSGGGGSGGGGSQTVV TQEPSLTVSPGGTVTLTCSSTGAVTSGNYPNWWVQQ KPGQAPRGLIGGTKFLAPGTPARFSGSLLGGKAALT SGVQPEDEAEYYCVLWYSNRWVFGGGTKLTVLGGG GDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRT EVTCTVVDVSHEDPEVKFNWYVDGVEVHNAKTKPC EEQYGYSTYRCVSVTLVHLDWLNKEYCKVSNKAL PAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLT CLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDG SFFLYSKLTVDKSRWQQGNVFSCVMHEALHNHYT QKSLSLSPGKGGGGSGGGGSGGGGSGGGGSGGGG SGGGGSDKTHTCPPCPAPELLGGPSVFLFPPKPKDTL MISRTPEVTCTVVDVSHEDPEVKFNWYVDGVEVHN AKTKPCEEQYGYSTYRCVSVTLVHLDWLNKEYCK VSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTK NQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTP VLDSDGSFFLYSKLTVDKSRWQQGNVFSCVMHEAL HNHYTQKSLSLSPGK
111.	PM 76-B10.17 CC VH CDR1	artificial	aa	DYYMY
112.	PM 76-B10.17 CC VH CDR2	artificial	aa	IISDAGYYTYSDIIG
113.	PM 76-B10.17 CC VH CDR3	artificial	aa	GFPLLRHGAMDY
114.	PM 76-B10.17 CC VL CDR1	artificial	aa	KASQNV DANVA
145.	PM 76-B10.17 CC VL CDR2	artificial	aa	SASYVYW

116.	PM 76-B10.17 CC VL CDR3	artificial	aa	QQYDQQLIT
117.	PM 76-B10.17 CC VH	artificial	aa	QVQLVESGGGLVKPGESLRLSCAASGFTFSDYYMYW VRQAPGKCLEWVAIISDAGYYTYSDIIKGRFTISRDN AKNSLYLQMNSLKAEDTAVYYCARGFPLLRHGAMD YWGGGTLTVSS
118.	PM 76-B10.17 CC VL	artificial	aa	DIQMTQSPSSLSASVGDRTITCKASQNV DANVAW YQQKPGQAPKSLIYSASYVYWDVPSRFGSGSASGTDFT LTISVQSEDFATYYCQQYDQQLITFGCGTKLEIK
119.	PM 76-B10.17 CC scFv	artificial	aa	QVQLVESGGGLVKPGESLRLSCAASGFTFSDYYMYW VRQAPGKCLEWVAIISDAGYYTYSDIIKGRFTISRDN AKNSLYLQMNSLKAEDTAVYYCARGFPLLRHGAMD YWGGGTLTVSSGGGGSGGGSGGGSGGGSDIQMTQS PSSLSASVGDRTITCKASQNV DANVAWYQQKPGQ APKSLIYSASYVYWDVPSRFGSGSASGTDFTLTISVQS EDFATYYCQQYDQQLITFGCGTKLEIK
120.	PM 76-B10.17 CC x I2C0 bispecific molecule	artificial	aa	QVQLVESGGGLVKPGESLRLSCAASGFTFSDYYMYW VRQAPGKCLEWVAIISDAGYYTYSDIIKGRFTISRDN AKNSLYLQMNSLKAEDTAVYYCARGFPLLRHGAMD YWGGGTLTVSSGGGGSGGGSGGGSGGGSDIQMTQS PSSLSASVGDRTITCKASQNV DANVAWYQQKPGQ APKSLIYSASYVYWDVPSRFGSGSASGTDFTLTISVQS EDFATYYCQQYDQQLITFGCGTKLEIKSGGGGSEVQL VESGGGLVQPGGSLKLSAASGFTFNKYAMNWVRQ APGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDD SKNTAYLQMNNLKTEDTAVYYCVRHGNFGNSYISY WAYWGGGTLTVSSGGGGSGGGSGGGSGGGGQTIVV TQEPSLTVSPGGTVTLTCSSTGAVTSGNYPNWVQQ KPGQAPRGLIGGTKFLAPGTPARFSGSLLGGKAALT SGVQPEDEAEYCVLWYSNRWVFGGGTKLTVL
121.	PM 76-B10.17 CC x I2C0-scFc bispecific HLE molecule	artificial	aa	QVQLVESGGGLVKPGESLRLSCAASGFTFSDYYMYW VRQAPGKCLEWVAIISDAGYYTYSDIIKGRFTISRDN AKNSLYLQMNSLKAEDTAVYYCARGFPLLRHGAMD YWGGGTLTVSSGGGGSGGGSGGGSGGGSDIQMTQS PSSLSASVGDRTITCKASQNV DANVAWYQQKPGQ APKSLIYSASYVYWDVPSRFGSGSASGTDFTLTISVQS EDFATYYCQQYDQQLITFGCGTKLEIKSGGGGSEVQL VESGGGLVQPGGSLKLSAASGFTFNKYAMNWVRQ APGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDD SKNTAYLQMNNLKTEDTAVYYCVRHGNFGNSYISY WAYWGGGTLTVSSGGGGSGGGSGGGSGGGGQTIVV TQEPSLTVSPGGTVTLTCSSTGAVTSGNYPNWVQQ KPGQAPRGLIGGTKFLAPGTPARFSGSLLGGKAALT SGVQPEDEAEYCVLWYSNRWVFGGGTKLTVLGGG GDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTP

				EVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPC EEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKAL PAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLT CLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDG SFFLYSKLTVDKSRWQQGNVFSCSVMHEALHNHYT QKSLSLSPGKGGGGSGGGGSGGGGSGGGGSGGGG SGGGGSDKTHTCPPCPAPELLGGPSVFLFPPKPKDTL MISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHN AKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCK VSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTK NQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTP VLDSDGSFFLYSKLTVDKSRWQQGNVFSCSVMHEAL HNHYTQKSLSLSPGK
122.	PM 76-B10.17 CC x I2C0- scFc_delGK bispesific HLE molecule	artificial	aa	QVQLVESGGGLVKPGESLRSCAASGFTFSDYYMYW VRQAPGKCLEWVAIISDAGYYTYSDIIKGRFTISRDN AKNSLYLQMNSLKAEDTAVYYCARGFLLRHGAMD YWGQGTLLTVSSGGGGSGGGGSGGGGSDIQMTQS PSSLSASVGDRVITITCKASQNV DANVAWYQQKPGQ APKSLIYSASYVYWDVPSRFGSASGTDFTLTISVQS EDFATYYCQQYDQQLITFGCGTKLEIKSGGGGSEVQL VESGGGLVQPGGSLKLSAASGFTFNKYAMNWWVRQ APGKLEWVARIRSKYNNYATYYADSVKDRFTISRDD SKNTAYLQMNNLKTEDTAVYYCVRHGNFGNSYISY WAYWGQGTLLTVSSGGGGSGGGGSGGGGSQTVV TQEPSLTVSPGGTVTLTCSSTGAVTSGNYPNWWVQQ KPGQAPRGLIGGKFLAPGTPARFSGSLLGGKAALT SGVQPEDEAEYCVLWYSNRWVFGGGTKLTVLGGG GDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRT EVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPC EEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKAL PAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLT CLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDG SFFLYSKLTVDKSRWQQGNVFSCSVMHEALHNHYT QKSLSLSPGKGGGGSGGGGSGGGGSGGGGSGGGG GGGSDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMI SRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAK TKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVS NKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKN QVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVL DSDGSFFLYSKLTVDKSRWQQGNVFSCSVMHEALH NHYTQKSLSLSPGK
123.	PM 76-B10.17 CC x I2C0 CC (103/43)-scFc	artificial	aa	QVQLVESGGGLVKPGESLRSCAASGFTFSDYYMYW VRQAPGKCLEWVAIISDAGYYTYSDIIKGRFTISRDN AKNSLYLQMNSLKAEDTAVYYCARGFLLRHGAMD YWGQGTLLTVSSGGGGSGGGGSGGGGSDIQMTQS PSSLSASVGDRVITITCKASQNV DANVAWYQQKPGQ

	bispecific molecule			APKSLIYSASYVYWDVPSRFGSASGTDFTLTSSVQS EDFATYYCQYDQQLITFGCGTKLEIKSGGGGSEVQL VESGGGLVQPGGSLKLSAASGFTFNKYAMNWVRQ APGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDD SKNTAYLQMNNLKTEDTAVYYCVRHGNFGNSYISY WAYCGQGT LVT VSSGGGSGGGGSGGGGSGT VVT QEPSLTVSPGGTVTLTCSSTGAVTSGNYPNWVQQ KPGQCPRGLIGGTKFLAPGTPARFSGSLLGGKAALTS GVQPEDEAEYYCVLWYSNRWVFGGGTKLTVL
124.	PM 76-B10.17 CC x I2C0 CC (103/43)-scFc bispecific HLE molecule	artificial	aa	QVQLVESGGGLVKPGESLRLSAASGFTFSDYYMYW VRQAPGKCLEWVAIISDAGYYTYSDIIGRFTISRDN AKNSLYLQMNSLKAEDTAVYYCARGFPLLRHGAMD YWGGQGT LVT VSSGGGSGGGGSGGGGSDIQMTQS PSSLSASVGDRTITCKASQNV DANVAWYQQKPGQ APKSLIYSASYVYWDVPSRFGSASGTDFTLTSSVQS EDFATYYCQYDQQLITFGCGTKLEIKSGGGGSEVQL VESGGGLVQPGGSLKLSAASGFTFNKYAMNWVRQ APGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDD SKNTAYLQMNNLKTEDTAVYYCVRHGNFGNSYISY WAYCGQGT LVT VSSGGGSGGGGSGGGGSGT VVT QEPSLTVSPGGTVTLTCSSTGAVTSGNYPNWVQQ KPGQCPRGLIGGTKFLAPGTPARFSGSLLGGKAALTS GVQPEDEAEYYCVLWYSNRWVFGGGTKLTVLGGGG DKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPE VTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPCE EQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALP APIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLT CLVKGFYPSDIAVEWESNGQPENNYKTPPVLDSDG SFFLYSKLTVDKSRWQQGNVFSCSVMHEALHNHYT QKSLSLSPGKGGGGSGGGGSGGGGSGGGGSGGGG SGGGGSDKTHTCPPCPAPELLGGPSVFLFPPKPKDTL MISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHN AKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCK VSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTK NQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTPPV VLDSDGSFFLYSKLTVDKSRWQQGNVFSCSVMHEAL HNHYTQKSLSLSPGK
125.	PM 76-B10.17 CC x I2C0 CC (103/43)- scFc_delGK bispecific HLE molecule	artificial	aa	QVQLVESGGGLVKPGESLRLSAASGFTFSDYYMYW VRQAPGKCLEWVAIISDAGYYTYSDIIGRFTISRDN AKNSLYLQMNSLKAEDTAVYYCARGFPLLRHGAMD YWGGQGT LVT VSSGGGSGGGGSGGGGSDIQMTQS PSSLSASVGDRTITCKASQNV DANVAWYQQKPGQ APKSLIYSASYVYWDVPSRFGSASGTDFTLTSSVQS EDFATYYCQYDQQLITFGCGTKLEIKSGGGGSEVQL VESGGGLVQPGGSLKLSAASGFTFNKYAMNWVRQ APGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDD

				SKNTAYLQMNNLKTEDTAVYYCVRHGNFGNSYISY WAYCGQGTTLTVSSGGGGSGGGGSGGGGSGTQVVT QEPSLTVSPGGTVTLTCGSSTGAVTSGNYPNWWQQ KPGQCPRGLIGGTKFLAPGTPARFSGSLLGGKAALTS GVQPEDEAEYYCVLWYSNRWVFGGGTKLTVLGGGG DKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPE VTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPCE EQYGSTYRCVSVLTVLHQDWLNGKEYCKVSNKALP APIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLT CLVKGFYPSDIAVEWESNGQPENNYKTPPVLDSDG SFFLYSKLTVDKSRWQQGNVFCFSVMHEALHNHYT QKSLSLSPGGGGSGGGGSGGGGSGGGGSGGGGSG GGGSDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMI SRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAK TKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYCKVS NKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKN QVSLTCLVKGFYPSDIAVEWESNGQPENNYKTPPVLD SDGSFFLYSKLTVDKSRWQQGNVFCFSVMHEALH NHYTQKSLSLSPGK
126.	PM 76-B10.11 CC VH CDR1	artificial	aa	DYYMY
127.	PM 76-B10.11 CC VH CDR2	artificial	aa	IISDGGYYTYSDIIG
128.	PM 76-B10.11 CC VH CDR3	artificial	aa	GFLLRHGAMDY
129.	PM 76-B10.11 CC VL CDR1	artificial	aa	KASQNVDTNVA
130.	PM 76-B10.11 CC VL CDR2	artificial	aa	SASYVYW
131.	PM 76-B10.11 CC VL CDR3	artificial	aa	QQYDQQLIT
132.	PM 76-B10.11 CC VH	artificial	aa	QVQLVESGGGLVKPGESLRLSCAASGFTFSDYYMYW VRQAPGKGLEWVAIISDGGYYTYSDIIGRFTISRDN AKNSLYLQMNSLKAEDTAVYYCARGFLLRHGAMD YWGGQGTTLTVSS
133.	PM 76-B10.11 CC VL	artificial	aa	DIQMTQSPSSLSASVGDRVTITCKASQNVDTNVAWY QQKPGQAPKSLIYSASYVYWDVPSRFSGSASGTDFTL TISSVQSEDFATYYCQQYDQQLITFGGGTKLEIK
134.	PM 76-B10.11 CC scFv	artificial	aa	QVQLVESGGGLVKPGESLRLSCAASGFTFSDYYMYW VRQAPGKGLEWVAIISDGGYYTYSDIIGRFTISRDN AKNSLYLQMNSLKAEDTAVYYCARGFLLRHGAMD YWGGQGTTLTVSSGGGGSGGGGSGGGGSDIQMTQS

				PSSLSASVGDRVTITCKASQNVDTNVAWYQQKPGQ APKSLIYSASYVYWDVPSRFGSASGTDFTLTISVQS EDFATYYCQYDQQLITFGGGTKLEIK
135.	PM 76-B10.11 CC x I2C0 bispecific molecule	artificial	aa	QVQLVESGGGLVKPGESLRSCAASGFTFSDYYMYW VRQAPGKGLEWVAIISDGGYYTYSDIIKGRFTISRDN AKNSLYLQMNSLKAEDTAVYYCARGFLLRHGAMD YWGGQGLTVTVSSGGGGSGGGSGGGGSDIQTQS PSSLSASVGDRVTITCKASQNVDTNVAWYQQKPGQ APKSLIYSASYVYWDVPSRFGSASGTDFTLTISVQS EDFATYYCQYDQQLITFGGGTKLEIKSGGGGSEVQL VESGGGLVQPGGSLKLSAASGFTFNKYAMNWVRQ APGKLEWVARIRSKYNNYATYYADSVKDRFTISRDD SKNTAYLQMNNLKTEDTAVYYCVRHGNFGNSYISY WAYWGQGLTVTVSSGGGGSGGGSGGGGSQTQV TQEPSLTVSPGGTVTLTCSSTGAVTSGNYPNWVQQ KPGQAPRGLIGGTKFLAPGTPARFSGSLLGGKAALT SGVQPEDEAEYCVLWYSNRWVFGGGTKLTVL
136.	PM 76-B10.11 CC x I2C0-scFc bispecific HLE molecule	artificial	aa	QVQLVESGGGLVKPGESLRSCAASGFTFSDYYMYW VRQAPGKGLEWVAIISDGGYYTYSDIIKGRFTISRDN AKNSLYLQMNSLKAEDTAVYYCARGFLLRHGAMD YWGGQGLTVTVSSGGGGSGGGSGGGGSDIQTQS PSSLSASVGDRVTITCKASQNVDTNVAWYQQKPGQ APKSLIYSASYVYWDVPSRFGSASGTDFTLTISVQS EDFATYYCQYDQQLITFGGGTKLEIKSGGGGSEVQL VESGGGLVQPGGSLKLSAASGFTFNKYAMNWVRQ APGKLEWVARIRSKYNNYATYYADSVKDRFTISRDD SKNTAYLQMNNLKTEDTAVYYCVRHGNFGNSYISY WAYWGQGLTVTVSSGGGGSGGGSGGGGSQTQV TQEPSLTVSPGGTVTLTCSSTGAVTSGNYPNWVQQ KPGQAPRGLIGGTKFLAPGTPARFSGSLLGGKAALT SGVQPEDEAEYCVLWYSNRWVFGGGTKLTVLGGG GDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRT EVTCTVVDVSHEDPEVKFNWYVDGVEVHNAKTKPC EEQYGSTYRCVSVTLVHLDWLNKEYCKVSNKAL PAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLT CLVKGFYPSDIAVEWESNGQPENNYKTPPVLDSDG SFFLYSKLTVDKSRWQQGNVFSCSVMHEALHNHYT QKSLSLSPGKGGGGSGGGSGGGSGGGSGGGG SGGGGSDKTHTCPPCPAPELLGGPSVFLFPPKPKDTL MISRTPEVTCTVVDVSHEDPEVKFNWYVDGVEVHN AKTKPCEEQYGSTYRCVSVTLVHLDWLNKEYCK VSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTK NQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTPP VLDSDGSFFLYSKLTVDKSRWQQGNVFSCSVMHEAL HNHYTQKSLSLSPGK

137.	PM 76-B10.11 CC x I2C0- scFc_delGK bispecific HLE molecule	artificial	aa	QVQLVESGGGLVKPGESLRSCAASGFTFSDYYMYW VRQAPGKGLEWVAIISDGGYYTYSDIIKGRFTISRDN AKNSLYLQMNSLKAEDTAVYYCARGFLLRHGAMD YWGQGTLLTVSSGGGGSGGGGSGGGGSDIQMTQS PSSLSASVGDRTITCKASQNVDTNVAWYQQKPGQ APKSLIYSASYVYWDVPSRFGSASGTDFTLTSSVQS EDFATYYCQYDQQLITFGGGTKLEIKSGGGGSEVQL VESGGGLVQPGGSLKLSAASGFTFNKYAMNWVRQ APGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDD SKNTAYLQMNNLKTEDTAVYYCVRHGNFGNSYISY WAYWGQGTLLTVSSGGGGSGGGGSGGGGSGQTVV TQEPSLTVSPGGTVTLTCSSTGAVTSGNYPNWVQQ KPGQAPRGLIGGTKFLAPGTPARFSGSLLGGKAALT SGVQPEDEAEYYCVLWYSNRWVFGGGTKLTVLGGG GDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRT EVTCTVVDVSHEDPEVKFNWYVDGVEVHNAKTKPC EEQYGSTYRCVSLTVLHQDWLNGKEYCKVSNKAL PAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLT CLVKGFYPSDIAVEWESNGQPENNYKTPPVLDSDG SFFLYSKLTVDKSRWQQGNVFSCSVMHEALHNHYT QKSLSLSPGGGGSGGGGSGGGGSGGGGSGGGGSG GGGSDKHTCPPCPAPELLGGPSVFLFPPKPKDTLMI SRTPEVTCVVDVSHEDPEVKFNWYVDGVEVHNAK TKPCEEQYGSTYRCVSLTVLHQDWLNGKEYCKVS NKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKN QVSLTCLVKGFYPSDIAVEWESNGQPENNYKTPPV LSDGGSFFLYSKLTVDKSRWQQGNVFSCSVMHEALH NHYTQKSLSLSPGK
138.	PM 76-B10.11 CC x I2C0 CC (103/43)-scFc bispecific molecule	artificial	aa	QVQLVESGGGLVKPGESLRSCAASGFTFSDYYMYW VRQAPGKGLEWVAIISDGGYYTYSDIIKGRFTISRDN AKNSLYLQMNSLKAEDTAVYYCARGFLLRHGAMD YWGQGTLLTVSSGGGGSGGGGSGGGGSDIQMTQS PSSLSASVGDRTITCKASQNVDTNVAWYQQKPGQ APKSLIYSASYVYWDVPSRFGSASGTDFTLTSSVQS EDFATYYCQYDQQLITFGGGTKLEIKSGGGGSEVQL VESGGGLVQPGGSLKLSAASGFTFNKYAMNWVRQ APGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDD SKNTAYLQMNNLKTEDTAVYYCVRHGNFGNSYISY WAYCGQGTLLTVSSGGGGSGGGGSGGGGSGQTVVT QEPSLTVSPGGTVTLTCSSTGAVTSGNYPNWVQQ KPGQCPRGLIGGTKFLAPGTPARFSGSLLGGKAALT LSGVQPEDEAEYYCVLWYSNRWVFGGGTKLTVL
139.	PM 76-B10.11 CC x I2C0 CC (103/43)-scFc bispecific	artificial	aa	QVQLVESGGGLVKPGESLRSCAASGFTFSDYYMYW VRQAPGKGLEWVAIISDGGYYTYSDIIKGRFTISRDN AKNSLYLQMNSLKAEDTAVYYCARGFLLRHGAMD YWGQGTLLTVSSGGGGSGGGGSGGGGSDIQMTQS

	HLE molecule			PSSLSASVGDRTITCKASQNVDTNVAWYQQKPGQ APKSLIYSASYVYWDVPSRFGSASGTDFTLTSSVQS EDFATYYCQYDQQLITFGGGTKLEIKSGGGGSEVQL VESGGGLVQPGGSLKLSAASGFTFNKYAMNWVRQ APGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDD SKNTAYLQMNNLKTEDTAVYYCVRHGNFGNSYISY WAYCGQGTLLTVSSGGGGSGGGGSGGGGSQTVVT QEPSLTVSPGGTVTLTCSSTGAVTSGNYPNWVQQ KPGQCPRGLIGGTKFLAPGTPARFSGSLLGGKAALTS GVQPEDEAEYYCVLWYSNRWVFGGGTKLTVLGGGG DKHTCPCPAPELLGGPSVFLFPPKPKDTLMISRTPE VTCVVDVSHEDPEVKFNWYVDGVEVHNAKTKPCE EQYGSTYRCVSVLTVLHQDWLNGKEYCKVSNKALP APIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLT CLVKGFYPSDIAVEWESNGQPENNYKTPPVLDSDG SFFLYSKLTVDKSRWQQGNVFSCSVMHEALHNHYT QKSLSLSPGKGGGGSGGGGSGGGGSGGGGSGGGG SGGGGSDKHTCPCPAPELLGGPSVFLFPPKPKDTL MISRTPEVTCVVDVSHEDPEVKFNWYVDGVEVHN AKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYCK VSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTK NQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTPP VLDSDGSFFLYSKLTVDKSRWQQGNVFSCSVMHEAL HNHYTQKSLSLSPGK
140.	PM 76-B10.11 CC x 12C0 CC (103/43)- scFc_delGK bispecific HLE molecule	artificial	aa	QVQLVESGGGLVKPGESLRLSAASGFTFSDYYMYW VRQAPGKGLEWVAIISDGGYYTYYSIIKGRFTISRDN AKNSLYLQMNSLKAEDTAVYYCARGFPLLRHGAMD YWGQGTLLTVSSGGGGSGGGGSGGGGSDIQMTQS PSSLSASVGDRTITCKASQNVDTNVAWYQQKPGQ APKSLIYSASYVYWDVPSRFGSASGTDFTLTSSVQS EDFATYYCQYDQQLITFGGGTKLEIKSGGGGSEVQL VESGGGLVQPGGSLKLSAASGFTFNKYAMNWVRQ APGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDD SKNTAYLQMNNLKTEDTAVYYCVRHGNFGNSYISY WAYCGQGTLLTVSSGGGGSGGGGSGGGGSQTVVT QEPSLTVSPGGTVTLTCSSTGAVTSGNYPNWVQQ KPGQCPRGLIGGTKFLAPGTPARFSGSLLGGKAALTS GVQPEDEAEYYCVLWYSNRWVFGGGTKLTVLGGGG DKHTCPCPAPELLGGPSVFLFPPKPKDTLMISRTPE VTCVVDVSHEDPEVKFNWYVDGVEVHNAKTKPCE EQYGSTYRCVSVLTVLHQDWLNGKEYCKVSNKALP APIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLT CLVKGFYPSDIAVEWESNGQPENNYKTPPVLDSDG SFFLYSKLTVDKSRWQQGNVFSCSVMHEALHNHYT QKSLSLSPGKGGGGSGGGGSGGGGSGGGGSGGGGSG GGGSDKHTCPCPAPELLGGPSVFLFPPKPKDTLMI

				SRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAK TKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKV NKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKN QVSLTCLVKGFYPSDIAVEWESNGQPENNYKTPPV LDSGSEFLYSKLTVDKSRWQQGNVFCFSVMHEALH NHYTQKSLSPGK
141.	PM 76-B10.11 CC x I2C0-scFc VH CDR1	artificial	aa	DYYMY
142.	PM 76-B10.11 CC x I2C0-scFc VH CDR2	artificial	aa	IISDGGYYTYYSIIKG
143.	PM 76-B10.11 CC x I2C0-scFc VH CDR3	artificial	aa	GFPLLRHGAMDY
144.	PM 76-B10.11 CC x I2C0-scFc VL CDR1	artificial	aa	KASQNVDTNVA
145.	PM 76-B10.11 CC x I2C0-scFc VL CDR2	artificial	aa	SASYVYW
146.	PM 76-B10.11 CC x I2C0-scFc VL CDR3	artificial	aa	QQYDQQLIT
147.	PM 76-B10.11 CC x I2C0-scFc VH	artificial	aa	QVQLVESGGGLVKPGESLRSLCAASGFTFSDYYMYW VRQAPGKCLEWVAIISDGGYYTYYSIIKGRFTISRDN AKNSLYLQMNSLKAEDTAVYYCARGFPLLRHGAMD YWGQGTSLTVSS
148.	PM 76-B10.11 CC x I2C0-scFc VL	artificial	aa	DIQMTQSPSSLSASVGDRVTITCKASQNVDTNVAWY QQKPGQAPKSLIYSASYVYWDVPSRFGSGSASGTDFTL TISSVQSEDFATYYCQQYDQQLITFGCGTKLEIK
149.	PM 76-B10.11 CC x I2C0-scFc scFv	artificial	aa	QVQLVESGGGLVKPGESLRSLCAASGFTFSDYYMYW VRQAPGKCLEWVAIISDGGYYTYYSIIKGRFTISRDN AKNSLYLQMNSLKAEDTAVYYCARGFPLLRHGAMD YWGQGTSLTVSSGGGGSGGGSGGGGSDIQMTQS PSSLSASVGDRVTITCKASQNVDTNVAWYQQKPGQ APKSLIYSASYVYWDVPSRFGSGSASGTDFTLTISSVQS EDFATYYCQQYDQQLITFGCGTKLEIK
150.	PM 76-B10.11 CC x I2C0-scFc bispecific molecule	artificial	aa	QVQLVESGGGLVKPGESLRSLCAASGFTFSDYYMYW VRQAPGKCLEWVAIISDGGYYTYYSIIKGRFTISRDN AKNSLYLQMNSLKAEDTAVYYCARGFPLLRHGAMD YWGQGTSLTVSSGGGGSGGGSGGGGSDIQMTQS PSSLSASVGDRVTITCKASQNVDTNVAWYQQKPGQ APKSLIYSASYVYWDVPSRFGSGSASGTDFTLTISSVQS EDFATYYCQQYDQQLITFGCGTKLEIKSGGGGSEVQL

				VESGGGLVQPGGSLKLSAASGFTFNKYAMNWVRQ APGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDD SKNTAYLQMNNLKTEDTAVYYCVRHGNFGNSYISY WAYWGQGTTLTVSSGGGGSGGGGSGGGGSQTQV TQEPSLTVSPGGTVTLTCGSSTGAVTSGNYPNWWVQ KPGQAPRGLIGGTKFLAPGTPARFSGSLLGGKAALT SGVQPEDEAEYCVLWYSNRWVFGGGTKLTVL
151.	PM 76-B10.11 CC x I2C0-scFc bispecific HLE molecule	artificial	aa	QVQLVESGGGLVKPGESLRLSAASGFTFSDYYMYW VRQAPGKCLEWVAIISDGGYYTYSDIIKGRFTISRDN AKNSLYLQMNSLKAEDTAVYYCARGFLLRHGAMD YWGGQGTTLTVSSGGGGSGGGGSGGGGSDIQMTQS PSSLSASVGDRVTITCKASQNVDTNVAWYQQKPGQ APKSLIYSASYVYWDVPSRFGSASGTDFTLTSSVQS EDFATYYCQYDQQLITFGCGTKLEIKSGGGGSEVQL VESGGGLVQPGGSLKLSAASGFTFNKYAMNWVRQ APGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDD SKNTAYLQMNNLKTEDTAVYYCVRHGNFGNSYISY WAYWGQGTTLTVSSGGGGSGGGGSGGGGSQTQV TQEPSLTVSPGGTVTLTCGSSTGAVTSGNYPNWWVQ KPGQAPRGLIGGTKFLAPGTPARFSGSLLGGKAALT SGVQPEDEAEYCVLWYSNRWVFGGGTKLTVLGGG GDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRT EVTCTVVDVSHEDPEVKFNWYVDGVEVHNAKTKPC EEQYGSTYRCVSVTLVHLDWLNKEYCKVSNKAL PAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSL TLVKGFPYSDIAVEWESNGQPENNYKTTTPVLDSDG SFFLYSKLTVDKSRWQQGNVFSCSVMHEALHNHYT QKSLSLSPGKGGGGSGGGGSGGGGSGGGGSGGGG SGGGGSDKTHTCPPCPAPELLGGPSVFLFPPKPKDT LMISRTPEVTCTVVDVSHEDPEVKFNWYVDGVEVHN AKTKPCEEQYGSTYRCVSVTLVHLDWLNKEYCK VSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTK NQVSLTCLVKGFPYSDIAVEWESNGQPENNYKTTTP VLDSDGSFFLYSKLTVDKSRWQQGNVFSCSVMHEAL HNHYTQKSLSLSPGK
152.	PM 76-B10.11 CC x I2C0- scFc_delGK bispecific HLE molecule	artificial	aa	QVQLVESGGGLVKPGESLRLSAASGFTFSDYYMYW VRQAPGKCLEWVAIISDGGYYTYSDIIKGRFTISRDN AKNSLYLQMNSLKAEDTAVYYCARGFLLRHGAMD YWGGQGTTLTVSSGGGGSGGGGSGGGGSDIQMTQS PSSLSASVGDRVTITCKASQNVDTNVAWYQQKPGQ APKSLIYSASYVYWDVPSRFGSASGTDFTLTSSVQS EDFATYYCQYDQQLITFGCGTKLEIKSGGGGSEVQL VESGGGLVQPGGSLKLSAASGFTFNKYAMNWVRQ APGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDD SKNTAYLQMNNLKTEDTAVYYCVRHGNFGNSYISY WAYWGQGTTLTVSSGGGGSGGGGSGGGGSQTQV

				TQEPSLTVSPGGTVTLTCSSTGAVTSGNYPNWVQQ KPGQAPRGLIGGTKFLAPGTPARFSGSLLGGKAALT SGVQPEDEAEYYCVLWYSNRWVFGGGTKLTVLGGG GDKHTHTCPPCPAPPELLGGPSVFLFPPKPKDTLMISRT EVTCTVVDVSHEDPEVKFNWYVDGVEVHNAKTKPC EEQYGSTYRCVSVLTVLHQDWLNGKEYCKVSNKAL PAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLT CLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDG SFFLYSKLTVDKSRWQQGNVFCFSVMHEALHNHYT QKSLSLSPGGGGSGGGGSGGGGSGGGGSGGGGSGG GGGSDKHTHTCPPCPAPPELLGGPSVFLFPPKPKDTLMI SRTPEVTCTVVDVSHEDPEVKFNWYVDGVEVHNAK TKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYCKVSN KALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKN QVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPV LSDGGSFFLYSKLTVDKSRWQQGNVFCFSVMHEALH NHYTQKSLSLSPGK
153.	PM 76-B10.11 CC x 12C0 CC (103/43)-scFc bispecific molecule	artificial	aa	QVQLVESGGGLVQPGESLRSLCAASGFTFSDYYMYW VRQAPGKCLEWVAIISDGGYYTYYSIIKGRFTISRDN AKNSLYLQMNSLKAEDTAVYYCARGFPLLRHGAMD YWGQGTLLTVSSGGGGSGGGGSGGGGSDIQMTQS PSSLSASVGDRTITCKASQNVDTNVAWYQQKPGQ APKSLIYSASYVYWDVPSRFGSGASGTDFTLTSSVQS EDFATYYCQYDQQLITFGCGTKLEIKSGGGGSEVQL VESGGGLVQPGGSLKSLCAASGFTFNKYAMNWVRQ APGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDD SKNTAYLQMNNLKTEDTAVYYCVRHGNFGNSYISY WAYCGQGTLLTVSSGGGGSGGGGSGGGGSGTQVVT QEPSLTVSPGGTVTLTCSSTGAVTSGNYPNWVQQ KPGQCPRLIGGTKFLAPGTPARFSGSLLGGKAALTLS GVQPEDEAEYYCVLWYSNRWVFGGGTKLTVL
154.	PM 76-B10.11 CC x 12C0 CC (103/43)-scFc bispecific HLE molecule	artificial	aa	QVQLVESGGGLVQPGESLRSLCAASGFTFSDYYMYW VRQAPGKCLEWVAIISDGGYYTYYSIIKGRFTISRDN AKNSLYLQMNSLKAEDTAVYYCARGFPLLRHGAMD YWGQGTLLTVSSGGGGSGGGGSGGGGSDIQMTQS PSSLSASVGDRTITCKASQNVDTNVAWYQQKPGQ APKSLIYSASYVYWDVPSRFGSGASGTDFTLTSSVQS EDFATYYCQYDQQLITFGCGTKLEIKSGGGGSEVQL VESGGGLVQPGGSLKSLCAASGFTFNKYAMNWVRQ APGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDD SKNTAYLQMNNLKTEDTAVYYCVRHGNFGNSYISY WAYCGQGTLLTVSSGGGGSGGGGSGGGGSGTQVVT QEPSLTVSPGGTVTLTCSSTGAVTSGNYPNWVQQ KPGQCPRLIGGTKFLAPGTPARFSGSLLGGKAALTLS GVQPEDEAEYYCVLWYSNRWVFGGGTKLTVLGGGG DKHTHTCPPCPAPPELLGGPSVFLFPPKPKDTLMISRTPE

				VTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPCE EQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALP APIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLT CLVKGFYPSDIAVEWESNGQPENNYKTPPVLDSDG SFFLYSKLTVDKSRWQQGNVFSCSVMHEALHNHYT QKSLSLSPGKGGGGSGGGSGGGSGGGSGGGSGGGG SGGGSDKTHTCPPCPAPELLGGPSVFLFPPKPKDTL MISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNA AKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCK VSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTK NQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTPPV LDSDGSFFLYSKLTVDKSRWQQGNVFSCSVMHEAL HNHYTQKSLSLSPGK
155.	PM 76-B10.11 CC x 12C0 CC (103/43)- scFc_delGK bispecific HLE molecule	artificial	aa	QVQLVESGGGLVKPGESLRSCAASGFTFSDYYMYW VRQAPGKCLEWVAIISDGGYYTYSDIIKGRFTISRDN AKNSLYLQMNSLKAEDTAVYYCARGFLLRHGAMD YWGQGTLLTVSSGGGGSGGGSGGGSGGGSDIQMTQS PSSLSASVGDRTITCKASQNVDTNVAWYQQKPGQ APKSLIYSASYVYWDVPSRFGSASGTDFTLTSSVQS EDFATYYCQYDQQLITFGCGTKLEIKSGGGGSEVQL VESGGGLVQPGGSLKLSAASGFTFNKYAMNWWVRQ APGKLEWVARIRSKYNNYATYYADSVKDRFTISRDD SKNTAYLQMNNLKTEDTAVYYCVRHGNFGNSYISY WAYCGQGTLLTVSSGGGGSGGGSGGGSGGGSTVVT QEPSLTVSPGGTVTLTCSSTGAVTSGNYPNWVQQ KPGQCPRGLIGGKFLAPGTPARFSGSLLGGKAALTLS GVQPEDEAEYYCVLWYSNRWVFGGKTLTVLGGGG DKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPE VTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPCE EQYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALP APIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLT CLVKGFYPSDIAVEWESNGQPENNYKTPPVLDSDG SFFLYSKLTVDKSRWQQGNVFSCSVMHEALHNHYT QKSLSLSPGKGGGGSGGGSGGGSGGGSGGGSGGGG GGGSDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMI SRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAK TKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVS NKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKN QVSLTCLVKGFYPSDIAVEWESNGQPENNYKTPPVLD SDGSFFLYSKLTVDKSRWQQGNVFSCSVMHEALH NHYTQKSLSLSPGK
156.	CD70_21D_CC VH CDR1	artificial	aa	TYAMS
157.	CD70_21D_CC VH CDR2	artificial	aa	AISGSGGRTFYAESVEG

158.	CD70_21D_CC VH CDR3	artificial	aa	HDYSNYPYFDY
1659 .	CD70_21D_CC VL CDR1	artificial	aa	RASQSVRSTYLA
160.	CD70_21D_CC VL CDR2	artificial	aa	GASSRAT
161.	CD70_21D_CC VL CDR3	artificial	aa	QQYGDLPFT
162.	CD70_21D_CC VH	artificial	aa	EVQLLESGGGMVQPGGSLRLSCAASGFTFSTYAMS WVRQAPGKCLEWWSAISGSGGRTFYAESVEGRFTIS RDNSKNTLYLQMNSLRAEDTAVYYCAKHDYSNYPYF DYWGQGTLLTVSS
163.	CD70_21D_CC VL	artificial	aa	EIVLTQSPGTLSPGERATLSCRASQSVRSTYLAWYQ QK PGQAPRLLIYGASSRATGIPDRFSGSGSGTDFTLTISRL E PEDFAVYSCQQYGDLPFTFGCGTKLEIK
164.	CD70_21D_CCx 12C scFc	artificial	aa	EVQLLESGGGMVQPGGSLRLSCAASGFTFSTYAMS WVRQAPGKCLEWWSAISGSGGRTFYAESVEGRFTIS RDNSKNTLYLQMNSLRAEDTAVYYCAKHDYSNYPYF DYWGQGTLLTVSSGGGGSGGGGSGGGGSEIVLTQS PGTSLSPGERATLSCRASQSVRSTYLAWYQQKPGQ APRLLIYGASSRATGIPDRFSGSGSGTDFTLTISRLEPE DFAVYSCQQYGDLPFTFGCGTKLEIKSGGGGSEVQLV ESGGGLVQPGGSLKLSAASGFTFNKYAMNWVRQA PGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDS KNTAYLQMNNLKTEDTAVYYCVRHGNFGNSYISYW AYWGQGTLLTVSSGGGGSGGGGSGGGGSGTQTVVTQ EPSLTVSPGGTVTLTCSSTGAVTSGNYPNWVQQKP GQAPRGLIGGTFKFLAPGTPARFSGSLLGGKAALTLSG VQPEDEAEYYCVLWYSNRWVFGGGTKLTVLGGGGD KTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEV TCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEE QYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALPA PIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCL VKGFYPSDIAVEWESNGQPENNYKTPPVLDSDGSF FLYSKLTVDKSRWQQGNVFCFSVMHEALHNHYTQK SLSLSPGKGGGGSGGGGSGGGGSGGGGSGGGGSGG GGGSDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMI SRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAK TKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCKVS NKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKN QVSLTCLVKGFYPSDIAVEWESNGQPENNYKTPPVLD SDGSFFLYSKLTVDKSRWQQGNVFCFSVMHEALH NHYTQKSLSLSPGK

165	MSLN_5_CCxC D3-scFc Bispecific HLE molecule		MGWSCILFLVATATGVHSQVQLVESGGGLVKPGGS LRLSCAASGFTFSDYYMTWIRQAPGKGLEWLSYISS GSTIYYADSVKGRFTISRDNKNSLFLQMNSLRAEDT AVYYCARDNRNSHFDYWGGQTLTVSSGGGGSGGG GSGGGGSDIQMTQSPSSVSASVGDRVITICRASQGI NTWLAWYQQKPGKAPKLLIYGASGLQSGVPSRFSGS GSGTDFTLTISSLQPEDFATYYCQQAQSFPRTFGQGT KVEIKSGGGGSEVQLVESGGGLVQPGGSLKLSAAS GFTFNKYAMNWVRQAPGKGLEWVARIRSKYNNYA TYYADSVKDRFTISRDDSKNTAYLQMNNLKTEDTAV YYCVRHGNFGNSYISYWAYWGGQTLTVSSGGGGG GGGGSGGGGSQTQVTVQEPSTVSPGGTVTLTCSST GAVTSGNYPNWVQQKPGQAPRGLIGGTFKFLAPGTP ARFSGSLLGGKAALTLSGVQPEDEAEYCVLWYSNR WVFGGGTKLTVLGGGGDKTHTCPPCPAPELLGGPSV FLFPPKPKDTLMISRTPEVTCVVDVSHEDPEVKFNW YVDGVEVHNAKTKPCEEQYGYSTYRCVSVLTVLHQD WLNGKEYCKVSNKALPAPIEKTISKAKGQPREPQVY TLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNG QPENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQQGN VFSCVMHEALHNHYTQKSLSLSPGKGGGGSGGGG SGGGSGGGSGGGSGGGSGGGSDKTHTCPPCPAPELL GGPSVFLFPPKPKDTLMISRTPEVTCVVDVSHEDP EVKFNWYVDGVEVHNAKTKPCEEQYGYSTYRCVSVLT VLHQDWLNGKEYCKVSNKALPAPIEKTISKAKGQPR EPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEW ESNGQPENNYKTTTPVLDSDGSFFLYSKLTVDKSRW QQGNVFSCVMHEALHNHYTQKSLSLSPGK
166	Anti-PSMA x12C0 with cys- clamp, scFc Bispecific molecule HLE PM76-B10.11		QVQLVESGGGLVKPGESLRLSCAASGFTFSDYYMYW VRQAPGKCLEWVAIISDGGYYTYYSIIKGRFTISRDN AKNSLYLQMNSLKAEDTAVYYCARGFPLLRHGAMD YWGGQTLTVTVSSGGGGSGGGSGGGSDIQMTQS PSSLSASVGDRVITICKASQNVDTNVAWYQQKPGQ APKSLIYSASYVYWDVPSRFSGSASGTDFTLTISSVQS EDFATYYCQYDQQLITFGCGTKLEIKSGGGGSEVQL VESGGGLVQPGGSLKLSAASGFTFNKYAMNWVRQ APGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDD SKNTAYLQMNNLKTEDTAVYYCVRHGNFGNSYISY WAYWGGQTLTVTVSSGGGGSGGGSGGGSGGGGSQTQV TQEPSTVSPGGTVTLTCSSTGAVTSGNYPNWVQQ KPGQAPRGLIGGTFKFLAPGTPARFSGSLLGGKAALT SGVQPEDEAEYCVLWYSNRWVFGGGTKLTVLGGG GDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRT PEVTCVVDVSHEDPEVKFNWYVDGVEVHNAKTKPC EEQYGYSTYRCVSVLTVLHQDWLNGKEYCKVSNKAL PAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLT

				CLVKGFYPSDIAVEWESNGQPENNYKTTTPVLSDG SFFLYSKLTVDKSRWQQGNVFSCSVMHEALHNHYT QKSLSLSPGKGGGGSGGGSGGGSGGGSGGGG SGGGGSDKTHTCPPCPAPELLGGPSVFLFPPKPKDTL MISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHN AKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYCK VSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTK NQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTP VLSDGSGFFLYSKLTVDKSRWQQGNVFSCSVMHEAL HNHYTQKSLSLSPGK
167	Anti-PSMA IC20 bispecific molecule PM76-B10.17			QVQLVESGGGLVKPGESLRSCAASGFTFSDYYMYW VRQAPGKGLEWVAIISDGGYYTYSDIIKGRFTISRDN AKNSLYLQMNSLKAEDTAVYYCARGFLLRHGAMD YWGQGTLLTVSSGGGGSGGGSGGGSGGGSDIQTQS PSSLSASVGDRTITCKASQNVDTNVAWYQQKPGQ APKSLIYSASYRSDVPSRFGSASGTDFTLTISVQSE DFATYYCQQYDSYPYTFGGGTKLEIKSGGGGSEVQLV ESGGGLVQPGGSLKLSAASGFTFNKYAMNWWVRQA PGKLEWVARIRSKYNNYATYYADSVKDRFTISRDDS KNTAYLQMNNLKTEDTAVYYCVRHGNFGNSYISYW AYWGQGTLLTVSSGGGGSGGGSGGGSGGGSTVVTQ EPSLTVSPGGTVTLTCSSTGAVTSGNYPNWWQQKP GQAPRGLIGGTFKFLAPGTPARFSGSLLGGKAALTLSG VQPEDEAEYYCVLWYSNRWVFGGGTKLTVL
168	Anti-Cldn 18.2 VH CDR1 CL-1 and CL-2	artificial	aa	GYMH
169	Anti-Cldn18.2 VH CDR2	artificial	aa	WINPNSGGTKYAQKFQG
170	Anti-Cldn18.2 VH CDR3	artificial	aa	DRITVAGTYYYGMDV
171	Anti-Cldn18.2 VL CDR1	artificial	aa	RASQGVNNWLA
172	Anti-Cldn18.2 VL CDR2	artificial	aa	TASSLQS
173	Anti-Cldn18.2 VL CDR3	artificial	aa	QQANSFPIT
174	Anti-Cldn18.2 VH anti- CL-1	artificial	aa	QVQLVQSGAEVKKPGASVKVSCASGYTFTGYMHWVR QAPGQCLEWMGWINPNSGGTKYAQKFQGRVTMTTRDT SISTAYMELSRLRSDDTAVYYCARDITVAGTYYYGMDV WGQGTITVTVSS
175	Anti-Cldn18.2 VL CL-1	artificial	aa	DIQMTQSPSSVSASVGDRTITCRASQGVNNWLAWYQQ KPGKAPKLLIYTASSLQSGVPSRFGSGSGTDFTLTIRSLQP EDFATYYCQQANSFPITFGCGTRLEIK
176	Anti-Cldn18.2 scFv	artificial	aa	QVQLVQSGAEVKKPGASVKVSCASGYTFTGYMHWVR QAPGQCLEWMGWINPNSGGTKYAQKFQGRVTMTTRDT

	CL-1			SISTAYMELSRRLRSDDTAVYYCARDRITVAGTYYYYGMDV WGQGTTVTVSSGGGGSGGGSGGGSDIQMTQSPSSV SASVGDRVTITCRASQGVNNWLAWYQQKPGKAPKLLIYT ASSLQSGVPSRFSGSGSGTDFTLTIRSLQPEDFATYYCQQA NSFPITFGCGTRLEIK
177	Anti-Cldn 18.2xCD3 bispecific molecule CL-1 x12C	artificial	aa	QVQLVQSGAEVKKPGASVKVSCASGYFTGYMHVWR QAPGQCLEWMGWNPNSGGTKYAQKFQGRVTMTRDT SISTAYMELSRRLRSDDTAVYYCARDRITVAGTYYYYGMDV WGQGTTVTVSSGGGGSGGGSGGGSDIQMTQSPSSV SASVGDRVTITCRASQGVNNWLAWYQQKPGKAPKLLIYT ASSLQSGVPSRFSGSGSGTDFTLTIRSLQPEDFATYYCQQA NSFPITFGCGTRLEIKSGGGGSEVQLVESGGGLVQPGGSLK LSCAASGFTFNKYAMNWRQAPGKGLEWVARIRSKYNN YATYYADSVKDRFTISRDDSKNTAYLQMNNLKTEDAVYY CVRHGNFGNSYISYWAYWGQGTTLTVSSGGGGSGGGGS GGGGSQTVVTQEPSLTVSPGGTVTLTCSSTGAVTSGNY PNWVQQKPGQAPRGLIGGTFKFLAPGTPARFSGSLLGGKA ALTLGSGVQPEDEAEYCVLWYSNRWVFGGGTKLTVL
178	Anti-Cldn 18.2xCD3 Bispecific scFc molecule CL-1 x12C-scFc	artificial	aa	QVQLVQSGAEVKKPGASVKVSCASGYFTGYMHVWR QAPGQCLEWMGWNPNSGGTKYAQKFQGRVTMTRDT SISTAYMELSRRLRSDDTAVYYCARDRITVAGTYYYYGMDV WGQGTTVTVSSGGGGSGGGSGGGSDIQMTQSPSSV SASVGDRVTITCRASQGVNNWLAWYQQKPGKAPKLLIYT ASSLQSGVPSRFSGSGSGTDFTLTIRSLQPEDFATYYCQQA NSFPITFGCGTRLEIKSGGGGSEVQLVESGGGLVQPGGSLK LSCAASGFTFNKYAMNWRQAPGKGLEWVARIRSKYNN YATYYADSVKDRFTISRDDSKNTAYLQMNNLKTEDAVYY CVRHGNFGNSYISYWAYWGQGTTLTVSSGGGGSGGGGS GGGGSQTVVTQEPSLTVSPGGTVTLTCSSTGAVTSGNY PNWVQQKPGQAPRGLIGGTFKFLAPGTPARFSGSLLGGKA ALTLGSGVQPEDEAEYCVLWYSNRWVFGGGTKLTVLGGG GDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVT CVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEEQYGS TYRCVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKA KGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAV EWESNGQPENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQ QGNVFSCSVMHEALHNHYTQKSLSLSPGKGGGGSGGGG SGGGSGGGSGGGSGGGSGGGSDKHTCPPCPAPELLGG PSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNW YVDGVEVHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNG KEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREE MTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTP VLDSDGSFFLYSKLTVDKSRWQQGNVFSCSVMHEALHNH YTQKSLSLSPGK
179	Anti-Cldn18.2 VH CL-2	artificial	aa	QVQMVSQSGAEVKKHGASVKVSCASGYFTGYMHVW RQAPGQCLEWMGWNPNSGGTKYAQKFQGRVTMTRD TSISTAYMELSRRLRSDDTAVYYCARDRITVAGTYYYYGMD VWGQGTTVTVSS

180	Anti-Cldn18.2 VL CL-2	artificial	aa	DIQMTQSPSSVSASVGDRTITCRASQGVNNWLAWYQQ KPGKAPKLLIYTASSLQSGVPSRFSGSGSGTDFTLTIRSLQP EDFATYYCQQANSFPITFGCGTRLEIK
181	Anti-Cldn18.2 scFv CL-2	artificial	aa	QVQMVSQSGAEVKKHGASVKVSCASGYFTGYMHWW RQAPGQCLEWMGWINPNSGGTKYAQKFQGRVTMTRD TSISTAYMELSRRLSDDTAVYYCARDRITVAGTYYYGMD VWGQGTITVTVSSGGGGSGGGSGGGGSDIQMTQSPSS VSASVGDRTITCRASQGVNNWLAWYQQKPGKAPKLLIY TASSLQSGVPSRFSGSGSGTDFTLTIRSLQPEDFATYYCQQ ANSFPITFGCGTRLEIK
182	Anti-Cldn18.2xCD3 bispecific molecule CL-2xI2C	artificial	aa	QVQMVSQSGAEVKKHGASVKVSCASGYFTGYMHWW RQAPGQCLEWMGWINPNSGGTKYAQKFQGRVTMTRD TSISTAYMELSRRLSDDTAVYYCARDRITVAGTYYYGMD VWGQGTITVTVSSGGGGSGGGSGGGGSDIQMTQSPSS VSASVGDRTITCRASQGVNNWLAWYQQKPGKAPKLLIY TASSLQSGVPSRFSGSGSGTDFTLTIRSLQPEDFATYYCQQ ANSFPITFGCGTRLEIKSGGGGSEVQLVESGGGLVQPGGS LKLSCAASGFTFNKYAMNWRQAPGKLEWVARIRSKY NNYATYYADSVKDRFTISRDDSKNTAYLQMNNLKTEDTA VYYCVRHGNFGNSYISYWAYWGQGLTVTVSSGGGGSGG GGSGGGGSQTVTQEPSTVSPGGTVTLTCSSTGAVTS GNYPNWWVQKPGQAPRGLIGGTKFLAPGTPARFSGSLLG GKAALTLSGVQPEDEAEYCVLWYSNRWVFGGKTCLTVL
183	Anti-Cldn18.2xCD3 Bispecific scFc molecule CL-2xI2C-scFc	artificial	aa	QVQMVSQSGAEVKKHGASVKVSCASGYFTGYMHWW RQAPGQCLEWMGWINPNSGGTKYAQKFQGRVTMTRD TSISTAYMELSRRLSDDTAVYYCARDRITVAGTYYYGMD VWGQGTITVTVSSGGGGSGGGSGGGGSDIQMTQSPSS VSASVGDRTITCRASQGVNNWLAWYQQKPGKAPKLLIY TASSLQSGVPSRFSGSGSGTDFTLTIRSLQPEDFATYYCQQ ANSFPITFGCGTRLEIKSGGGGSEVQLVESGGGLVQPGGS LKLSCAASGFTFNKYAMNWRQAPGKLEWVARIRSKY NNYATYYADSVKDRFTISRDDSKNTAYLQMNNLKTEDTA VYYCVRHGNFGNSYISYWAYWGQGLTVTVSSGGGGSGG GGSGGGGSQTVTQEPSTVSPGGTVTLTCSSTGAVTS GNYPNWWVQKPGQAPRGLIGGTKFLAPGTPARFSGSLLG GKAALTLSGVQPEDEAEYCVLWYSNRWVFGGKTCLTVL GGGGDKTHTCPPCPAPPELLGGPSVFLFPPKPKDTLMISRT PEVTCVVDVSHEDPEVKFNWYVDGVEVHNAKTKPCEE QYGSTYRCVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEK TISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPS DIAVEWESNGQPENNYKTTTPVLDSDGSFFLYSKLTVDKS RWQQGNVFSCSVMHEALHNHYTQKSLSLSPGKGGGGS GGGGSGGGSGGGSGGGSGGGSGGGGSDKTHTCPPCPAP ELLGGPSVFLFPPKPKDTLMISRTPEVTCVVDVSHEDPEV KFNWYVDGVEVHNAKTKPCEEQYGSTYRCVSVLTVLHQD WLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPP SREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYK TTPVLDSDGSFFLYSKLTVDKSRWQQGNVFSCSVMHEA LHNHYTQKSLSLSPGK
184	Anti-MUC17	artificial	aa	GYYS

	VH CDR1 MU 8-B7			
185	Anti-MUC17 VH CDR2 MU 8-B7	artificial	aa	DIDASGSTKYNPSLKS
186	Anti-MUC17 VH CDR3 MU 8-B7	artificial	aa	KKYSTVWSYFDN
187	Anti-MUC17 VL CDR1 MU 8-B7	artificial	aa	SGDKLGDKYAS
188	Anti-MUC17 VL CDR2 MU 8-B7	artificial	aa	QDRKRPS
189	Anti-MUC17 VL CDR3 MU 8-B7	artificial	aa	QAWGSSTAV
190	Anti-MUC17 VH MU 8-B7	artificial	aa	QVQLQQWAGAGLLKPSETLSLTCAVYGGSFSGYYWSWIRQPPG KCLEWIGDIDASGSTKYNPSLKSRTISLDTSKNQFSLKLNSTAA DTAVYFCARKKYSTVWSYFDNWGQGLTVTVSS
191	Anti-MUC17 VL MU 8-B7	artificial	aa	SYELTQPSSVSPPGQTASITCSGDKLGDKYASWYQQKPG QSPVLVIYQDRKRPSGVPERFSGSNSGNTATLTISGTQAM DEADYYCQAWGSSTAVFGCGTKLTVL
192	bispecific molecule MU 8-B7 x I2C0scFc	artificial	aa	QVQLQQWAGAGLLKPSETLSLTCAVYGGSFSGYYWSWIR QPPGKCLEWIGDIDASGSTKYNPSLKSRTISLDTSKNQF LKLNSVTAADTAVYFCARKKYSTVWSYFDNWGQGLTVTV SSGGGGSGGGGGSGGGSSYELTQPSSVSPPGQTASITCS GDKLGDKYASWYQQKPGQSPVLVIYQDRKRPSGVPERFS GSNSGNTATLTISGTQAMDEADYYCQAWGSSTAVFGCG TKLTVLSGGGGSEVQLVESGGGLVQPGGSLKLSAASGFT FNKYAMNWVRQAPGKLEWVARIRSKYNNYATYYADSV KDRFTISRDDSKNTAYLQMNNLKTEDTAVYYCVRHGNFG NSYISYWAYWGQGLTVTVSSGGGGSGGGGGSGGGGSQT VVTQEPSTLTVSPGGTVTLTCSSTGAVTSGNYPNWVQQK PGQAPRGLIGGTKFLAPGTPARFSGSLLGGKAALTLSGVQ PEDEAEYYCVLWYSNRWVFSGGGLTVLGGGDKTHTC PPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVS HEDPEVKFNWYVDGVEVHNAKTKPCEEQYGSTYRCVSVL TVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREP QVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNG QPENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQQGNVFS CSVMHEALHNHYTQKSLSLSPGKGGGGSGGGGGSGGGGS GGGGSGGGGGSGGGGSDKTHTCPPCPAPELLGGPSVFLFP PKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVE VHNAKTKPCEEQYGSTYRCVSVLTVLHQDWLNGKEYKCK VSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQ VSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDG

				SFFLYSKLTVDKSRWQQGNVFSCSVMHEALHNHYTQKSL SLSPGK
193	CDR-L1 of I2C	artificial	aa	GSSTGAVTSGNYPN
194	CDR-L2 of I2C	artificial	aa	GTKFLAP
195	CDR-L3 of I2C	artificial	aa	VLWYSNRWV
196	CDR-H1 of I2C	artificial	aa	KYAMN
197	CDR-H2 of I2C	artificial	aa	RIRSKYNNYATYYADSVKD
198	CDR-H3 of I2C	artificial	aa	HGNFGNSYISYWAY
199	VH of I2C	artificial	aa	EVQLVESGGGLVQPGGSLKLSAASGFTFNKYAMNWVR QAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSK NTAYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWG QGTLVTVSS
200	VL of I2C	artificial	aa	QTVVTQEPSTLVSPGGTVTLTCSSTGAVTSGNYPNWVQ QKPGQAPRGLIGGKFLAPGTPARFSGSLLGGKAALTLSG VQPEDEAEYYCVLWYSNRWVFGGGTKLTVL
201	VH-VL of I2C	artificial	aa	EVQLVESGGGLVQPGGSLKLSAASGFTFNKYAMNWVR QAPGKGLEWVARIRSKYNNYATYYADSVKDRFTISRDDSK NTAYLQMNNLKTEDTAVYYCVRHGNFGNSYISYWAYWG QGTLVTVSSGGGGSGGGSGGGSGGGSGTQVVTQEPSTLVSP GGTVTLTCSSTGAVTSGNYPNWVQKPGQAPRGLIGG TKFLAPGTPARFSGSLLGGKAALTLSGVQPEDEAEYYCVL WYSNRWVFGGGTKLTVL

[00117] The invention will be more fully understood by reference to the following examples. The examples should not, however, be construed as limiting the scope of the invention.

EXAMPLES

EXAMPLE 1. METHOD OF MEASURING PROTEIN BINDING TO SOLID SURFACES

[00118] Bispecific antibody constructs were internally sourced. They were labeled with a fluorophore and purified after the labeling procedure.

[00119] Each measurement chamber contains a plastic coverslip. To measure the protein binding to solid surfaces, the measurement chambers with plastic coverslips on the bottom were incubated with a solution containing a fluorophore-protein (e.g., a fluorophore labeled antibody construct) first. Then the sample solution was aspirated out, the from the coverslips were rinsed and filled with buffer for imaging on a confocal microscope later. Fluorescence intensity as measured by the confocal microscope shows the binding of the bispecific antibody constructs to the coverslips. **Figure 1** shows the diagram of the experimental set-up.

[00120] **Figure 2** shows titrations of two fluorophore-labeled antibody constructs binding to the solid surfaces (e.g., coverslips), separately, in the absence of surfactant. The fluorescence intensities of bound fluorophore-labeled antibody constructs were measured by the confocal xy scans of the surface.

EXAMPLE 2. TREATMENT OF SOLID SURFACES WITH SURFACTANTS PREVENTING PROTEIN BINDING TO THE SURFACES

[00121] Several surfactants were used at different folds of their respective CMC to determine the effectiveness of each surfactant at preventing bispecific antibody construct binding to solid surface. In this study, a solution containing surfactants was incubated with the surface first, then the fluorophore labeled antibody constructs were added and incubated. After that, the solution was aspirated out, the surface was rinsed and filled with buffer for imaging on a confocal microscope. The results are shown in **Figure 3**. In the figure, the first group of bars are the bispecific antibody binding to the surface without any surfactants. These were served as benchmark and all the following groups of data were compared to those. From the 2nd to the last group are the relative percentages of protein bound to the surfaces pre-treated with different surfactants at different folds of their distinguish CMCs. The insert is a zoom-in for the lower region of the graph. PS 80, PS 20, P188, P407 and Triton X-100 were investigated.

[00122] The order of adding the surfactants and the antibodies to the surface was tested. For bispecific antibodies 1 & 2, two orders were tested: in the first one, a surfactant-containing solution was added to the surface before adding the antibodies to the surface; while the other one was vice versa. The results were shown in **Figure 4** (left is antibody 1 and right is antibody 2). For both graphs, the first group of bars are the bispecific antibody binding to the surface without any surfactants. These were served as benchmark and all the following groups of data were compared to those. From the 2nd to the last group are the relative percentages of the antibodies bound to the surfaces pre-treated with PS 80 at different folds of its CMC. The surfactants effectively prevented the proteins binding to the surfaces.

EXAMPLE 3. INCOMPATIBILITY OF CERTAIN SURFACTANTS AND THE PLASTICIZERS USED IN PLASTIC IV COMPONENTS

[00123] Baxter Viaflex PVC-DEHP IV bags pre-filled with saline diluent were used for the study. Surfactants polysorbate 80 (PS80), polysorbate 20 (PS20), poloxamer 188 (P188), poloxamer 407 (P407) and Triton X-100 were used in the study. Different amounts of different

surfactants were incubated with the bags at 25 °C for 24 hrs or 48 hrs. Then the bags were sampled and analyzed by reversed-phase ultra-high pressure liquid chromatography (RP-UHPLC) and detected by an UV detector. The mobile phase A & B are 0.1% trifluoroacetic acid (TFA) in DI-water and 0.1% TFA in acetonitrile. The gradient is listed in the Table 3 below. Flow rate is 0.6 ml/min. For quantification, a standard curve of DEHP was established under the same conditions.

Table 3. RP-UPLC gradient

Time (min)	Flow (mL/min)	%A	%B
0.0	0.6	95	5
1.5	0.6	5	95
3.1	0.6	5	95
3.2	0.6	95	5
4.0	0.6	95	5

[00124] PS 80, PS 20 and P188 were compared at 0.3 wt% in saline in PVC-DEHP IV bags for 24 hrs and 48 hrs at 25 °C. The results are shown in **Figure 5**. PS80 & PS20 caused significant leaching of DEHP from PVC-DEHP IV bags, while P188 didn't cause any leaching (Figure 5). Saline only was used as a control.

[00125] PS 80, PS 20, P188, P407 and Triton X-100 were compared at different folds of their respective CMC by incubating in saline in PVC-DEHP IV bags for 24 hrs at 25 °C. The amount of leached DEHP was plotted as a function of the folds of CMC in **Figure 6**. Clearly, polysorbates extract certain amounts of DEHP while poloxamers don't. And the amounts of DEHP leached are correlated with the amounts of surfactants used.