

COMPOUNDS USEFUL AS RAF KINASE INHIBITORS

CROSS-REFERENCE TO RELATED APPLICATIONS

[0001] The present invention claims priority to United States provisional patent application serial number 60/947,291, filed June 29, 2007, the entirety of which is hereby incorporated herein by reference.

TECHNICAL FIELD OF THE INVENTION

[0002] The present invention relates to compounds useful as inhibitors of protein kinases. The invention also provides pharmaceutically acceptable compositions comprising compounds of the present invention and methods of using said compositions in the treatment of various disorders.

BACKGROUND OF THE INVENTION

[0003] Cancer results from the deregulation of the normal processes that control cell division, differentiation and apoptotic cell death. Protein kinases play a critical role in this regulatory process. A partial non-limiting list of such kinases includes ab1, ATK, bcr-ab1, Blk, Brk, Btk, c-kit, c-met, c-src, CDK1, CDK2, CDK4, CDK6, cRaf1, CSF1R, CSK, EGFR, ErbB2, ErbB3, ErbB4, ERK, Fak, fes, FGFR1, FGFR2, FGFR3, FGFR4, FGFR5, Fgr, FLK4, flt-1, Fps, Frk, Fyn, Hck, IGF-1R, INS-R, Jak, KDR, Lck, Lyn, MEK, p38, PDGFR, PIK, PKC, PYK2, ros, tie₁, tie₂, TRK, Yes and Zap70. In mammalian biology, such protein kinases comprise mitogen activated protein kinase (MAPK) signalling pathways. MAPK signalling pathways are inappropriately activated by a variety of common disease-associated mechanisms such as mutation of ras genes and deregulation of growth factor receptors (Magnuson *et al.*, Seminars in Cancer Biology; 1994 (5), 247-252).

[0004] Additionally, protein kinases have been implicated as targets in central nervous system disorders (such as Alzheimer's), inflammatory disorders (such as psoriasis, arthritis), bone diseases (such as osteoporosis), atherosclerosis, restenosis, thrombosis, metabolic disorders (such as diabetes) and infectious diseases (such as viral and fungal infections).

[0005] One of the most commonly studied pathways involving kinase regulation is intracellular signalling from cell surface receptors to the nucleus. One example of this pathway includes a cascade of kinases in which members of the Growth Factor receptor Tyrosine Kinases (such as EGF-R, PDGF-R, VEGF-R, IGF1-R, the Insulin receptor) deliver signals through phosphorylation to other kinases such as Src Tyrosine kinase, and the Raf, Mek and Erk serine/threonine kinase families. Each of these kinases is represented by several family members, which play related, but functionally distinct roles. The loss of regulation of the growth factor signalling pathway is a frequent occurrence in cancer as well as other disease states.

[0006] The signals mediated by kinases have also been shown to control growth, death and differentiation in the cell by regulating the processes of the cell cycle. Progression through the eukaryotic cell cycle is controlled by a family of kinases called cyclin dependent kinases (CDKs). The regulation of CDK activation is complex, but requires the association of the CDK with a member of the cyclin family of regulatory subunits. A further level of regulation occurs through both activating and inactivating phosphorylations of the CDK subunit. The coordinate activation and inactivation of different cyclin/CDK complexes is necessary for normal progression through the cell cycle. Both the critical G1-S and G2-M transitions are controlled by the activation of different cyclin/CDK activities. In G1, both cyclin D/CDK4 and cyclin E/CDK2 are thought to mediate the onset of S-phase. Progression through S-phase requires the activity of cyclin A/CDK2 whereas the activation of cyclin A/cdc2 (CDK1) and cyclin B/cdc2 are required for the onset of metaphase. It is not surprising, therefore, that the loss of control of CDK regulation is a frequent event in hyperproliferative diseases and cancer.


[0007] Raf protein kinases are key components of signal transduction pathways by which specific extracellular stimuli elicit precise cellular responses in mammalian cells. Activated cell surface receptors activate ras/rap proteins at the inner aspect of the plasma membrane which in turn recruit and activate Raf proteins. Activated Raf proteins phosphorylate and activate the intracellular protein kinases MEK1 and MEK2. In turn, activated MEKs catalyze phosphorylation and activation of p42/p44 mitogen-activated protein kinase (MAPK). Various cytoplasmic and nuclear substrates of activated MAPK are known which directly or indirectly contribute to the cellular response to environmental change. Three distinct genes have been identified in mammals that encode Raf proteins; A-Raf, B-Raf and C-Raf (also known as Raf-1) and isoformic variants that result from differential splicing of mRNA are known.

[0008] Inhibitors of Raf kinases have been suggested for use in disruption of tumor cell growth and hence in the treatment of cancers, e.g., histiocytic lymphoma, lung adenocarcinoma, small cell lung cancer, and pancreatic and breast carcinoma; and also in the treatment and/or prophylaxis of disorders associated with neuronal degeneration resulting from ischemic events, including cerebral ischemia after cardiac arrest, stroke and multi-infarct dementia and also after cerebral ischemic events such as those resulting from head injury, surgery, and/or during childbirth.

[0009] Accordingly, there is a great need to develop compounds useful as inhibitors of protein kinases. In particular, it would be desirable to develop compounds that are useful as Raf inhibitors.

SUMMARY OF THE INVENTION

[0010] It has now been found that compounds of this invention, and pharmaceutically acceptable compositions thereof, are effective as inhibitors of one or more protein kinases. Such compounds are of formula **I**:



or a pharmaceutically acceptable salt thereof, wherein each of R^x , R^y , R^1 , L^1 , L^2 , Cy^1 , and Cy^2 are as defined in classes and subclasses herein, and pharmaceutical compositions thereof, as described generally and in subclasses herein, which compounds are useful as inhibitors of protein kinase (*e.g.*, Raf), and thus are useful, for example, for the treatment of Raf-mediated diseases.

[0011] In certain other embodiments, the invention provides pharmaceutical compositions comprising a compound of the invention, wherein the compound is present in an amount effective to inhibit Raf activity. In certain other embodiments, the invention provides pharmaceutical compositions comprising a compound of the invention and optionally further


comprising an additional therapeutic agent. In yet other embodiments, the additional therapeutic agent is an agent for the treatment of cancer.

[0012] In yet another aspect, the present invention provides methods for inhibiting kinase (e.g., Raf) activity in a patient or a biological sample, comprising administering to said patient, or contacting said biological sample with, an effective inhibitory amount of a compound of the invention. In still another aspect, the present invention provides methods for treating any disorder involving Raf activity, comprising administering to a subject in need thereof a therapeutically effective amount of a compound of the invention.

DETAILED DESCRIPTION OF CERTAIN EMBODIMENTS OF THE INVENTION

1. General Description of Compounds of the Invention:

[0013] In certain embodiments, the present invention provides a compound of formula I:



or a pharmaceutically acceptable salt thereof, wherein:

Cy¹ is an optionally substituted phenyl or 5–6 membered saturated, partially unsaturated, or aromatic ring having 1–3 heteroatoms independently selected from nitrogen, oxygen, or sulfur;

Cy² is an optionally substituted 5–14 membered saturated, partially unsaturated, or aromatic monocyclic, bicyclic, or tricyclic ring having 0–4 heteroatoms, independently selected from nitrogen, oxygen, or sulfur;

L¹ is a direct bond or an optionally substituted, straight or branched C_{1–6} alkylene chain;

L² is a direct bond, or is an optionally substituted, straight or branched C_{1–6} alkylene chain wherein 1 or 2 methylene units of L² are optionally and independently replaced by –O–, –S–, –N(R)–, –C(O)–, –C(O)N(R)–, –N(R)C(O)N(R)–, –N(R)C(O)–, –N(R)C(O)O–, –OC(O)N(R)–, –SO₂–, –SO₂N(R)–, –N(R)SO₂–, –OC(O)–, –C(O)O–, or a 3–6 membered cycloalkylene;

each R is independently hydrogen or an optionally substituted C₁₋₆ aliphatic group;

R¹ is hydrogen or an optionally substituted C₁₋₆ aliphatic group;

each of R^x and R^y is independently selected from -R², -halo, -NO₂, -CN, -OR², -SR², -N(R²)₂, -C(O)R², -CO₂R², -C(O)C(O)R², -C(O)CH₂C(O)R², -S(O)R², -S(O)₂R², -C(O)N(R²)₂, -SO₂N(R²)₂, -OC(O)R², -N(R²)C(O)R², -N(R²)N(R²)₂, -N(R²)-C(=NR²)N(R²)₂, -C(=NR²)N(R²)₂, -C=NOR², -N(R²)C(O)N(R²)₂, -N(R²)SO₂N(R²)₂, -N(R²)SO₂R², or -OC(O)N(R²)₂; and

each R² is independently hydrogen or an optionally substituted group selected from C₁₋₆ aliphatic, a C₆₋₁₀ monocyclic or bicyclic aryl ring, or a 5-10 membered saturated, partially unsaturated, or aromatic monocyclic or bicyclic ring having 1-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur, or

two R² on the same nitrogen are taken together with the nitrogen to form an optionally substituted 5-8 membered saturated, partially unsaturated, or aromatic ring having 1-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur.

[0014] Compounds of this invention include those generally set forth above and described specifically herein, and are illustrated in part by the various classes, subgenera and species disclosed herein. Additionally, the present invention provides pharmaceutically acceptable derivatives of the compounds of the invention, and methods of treating a subject using these compounds, pharmaceutical compositions thereof, or either of these in combination with one or more additional therapeutic agents.

2. Compounds and Definitions:

[0015] Definitions of specific functional groups and chemical terms are described in more detail below. For purposes of this invention, the chemical elements are identified in accordance with the Periodic Table of the Elements, CAS version, *Handbook of Chemistry and Physics*, 75th Ed., inside cover, and specific functional groups are generally defined as described therein. Additionally, general principles of organic chemistry, as well as specific functional moieties and reactivity, are described in *Organic Chemistry*, Thomas Sorrell, University Science Books, Sausalito, 1999; Smith and March *March's Advanced Organic Chemistry*, 5th Edition, John Wiley & Sons, Inc., New York, 2001; Larock, *Comprehensive Organic Transformations*, VCH Publishers, Inc., New York, 1989; Carruthers, *Some Modern Methods of Organic Synthesis*, 3rd

Edition, Cambridge University Press, Cambridge, 1987; the entire contents of each of which are incorporated herein by reference.

[0016] Certain compounds of the present invention can comprise one or more asymmetric centers, and thus can exist in various isomeric forms, e.g., stereoisomers and/or diastereomers. Thus, compounds of the invention and pharmaceutical compositions thereof may be in the form of an individual enantiomer, diastereomer or geometric isomer, or may be in the form of a mixture of stereoisomers. In certain embodiments, the compounds of the invention are enantiopure compounds. In certain other embodiments, mixtures of stereoisomers or diastereomers are provided.

[0017] Furthermore, certain compounds, as described herein, may have one or more double bonds that can exist as either the Z or E isomer, unless otherwise indicated. The invention additionally encompasses the compounds as individual isomers substantially free of other isomers and alternatively, as mixtures of various isomers, e.g., racemic mixtures of stereoisomers. In addition to the above-mentioned compounds *per se*, this invention also encompasses pharmaceutically acceptable derivatives of these compounds and compositions comprising one or more compounds.

[0018] Where a particular enantiomer is preferred, it may, in some embodiments be provided substantially free of the corresponding enantiomer, and may also be referred to as “optically enriched.” “Optically-enriched,” as used herein, means that the compound is made up of a significantly greater proportion of one enantiomer. In certain embodiments the compound is made up of at least about 90% by weight of a preferred enantiomer. In other embodiments the compound is made up of at least about 95%, 98%, or 99% by weight of a preferred enantiomer. Preferred enantiomers may be isolated from racemic mixtures by any method known to those skilled in the art, including chiral high pressure liquid chromatography (HPLC) and the formation and crystallization of chiral salts or prepared by asymmetric syntheses. See, for example, Jacques, et al., *Enantiomers, Racemates and Resolutions* (Wiley Interscience, New York, 1981); Wilen, et al., *Tetrahedron* 33:2725 (1977); Eliel, E.L. *Stereochemistry of Carbon Compounds* (McGraw-Hill, NY, 1962); Wilen, S.H. *Tables of Resolving Agents and Optical Resolutions* p. 268 (E.L. Eliel, Ed., Univ. of Notre Dame Press, Notre Dame, IN 1972).

[0019] The term “heteroatom” means one or more of oxygen, sulfur, nitrogen, phosphorus, or silicon (including, any oxidized form of nitrogen, sulfur, phosphorus, or silicon; the

quaternized form of any basic nitrogen or; a substitutable nitrogen of a heterocyclic ring, for example N (as in 3,4-dihydro-2*H*-pyrrolyl), NH (as in pyrrolidinyl) or NR⁺ (as in N-substituted pyrrolidinyl)).

[0020] As used herein a “direct bond” or “covalent bond” refers to a single, double or triple bond. In certain embodiments, a “direct bond” refers to a single bond.


[0021] The terms “halo” and “halogen” as used herein refer to an atom selected from fluorine (fluoro, -F), chlorine (chloro, -Cl), bromine (bromo, -Br), and iodine (iodo, -I).



[0022] The term “aliphatic” or “aliphatic group”, as used herein, denotes a hydrocarbon moiety that may be straight-chain (i.e., unbranched), branched, or cyclic (including fused, bridging, and spiro-fused polycyclic) and may be completely saturated or may contain one or more units of unsaturation, but which is not aromatic. Unless otherwise specified, aliphatic groups contain 1–6 carbon atoms. In some embodiments, aliphatic groups contain 1–4 carbon atoms, and in yet other embodiments aliphatic groups contain 1–3 carbon atoms. Suitable aliphatic groups include, but are not limited to, linear or branched, alkyl, alkenyl, and alkynyl groups, and hybrids thereof such as (cycloalkyl)alkyl, (cycloalkenyl)alkyl or (cycloalkyl)alkenyl.

[0023] The term “unsaturated”, as used herein, means that a moiety has one or more units of unsaturation.

[0024] The terms “cycloaliphatic”, “carbocycle”, “carbocyclyl”, “carbocyclo”, or “carbocyclic”, used alone or as part of a larger moiety, refer to a saturated or partially unsaturated cyclic aliphatic monocyclic or bicyclic ring systems, as described herein, having from 3 to 10 members, wherein the aliphatic ring system is optionally substituted as defined above and described herein. Cycloaliphatic groups include, without limitation, cyclopropyl, cyclobutyl, cyclopentyl, cyclopentenyl, cyclohexyl, cyclohexenyl, cycloheptyl, cycloheptenyl, cyclooctyl, cyclooctenyl, and cyclooctadienyl. In some embodiments, the cycloalkyl has 3–6 carbons. The terms “cycloaliphatic”, “carbocycle”, “carbocyclyl”, “carbocyclo”, or “carbocyclic” also include aliphatic rings that are fused to one or more aromatic or nonaromatic rings, such as decahydronaphthyl or tetrahydronaphthyl, where the radical or point of attachment is on the aliphatic ring.

[0025] As used herein, the term “cycloalkylene” refers to a bivalent cycloalkyl group. In certain embodiments, a cycloalkylene group is a 1,1-cycloalkylene group (i.e., a spiro-fused

ring). Exemplary 1,1-cycloalkylene groups include . In other embodiments, a cycloalkylene group is a 1,2-cycloalkylene group or a 1,3-cycloalkylene group. Exemplary 1,2-

cycloalkylene groups include  and .

[0026] The term “alkyl,” as used herein, refers to saturated, straight- or branched-chain hydrocarbon radicals derived from an aliphatic moiety containing between one and six carbon atoms by removal of a single hydrogen atom. In some embodiments, the alkyl group employed in the invention contains 1–5 carbon atoms. In another embodiment, the alkyl group employed contains 1–4 carbon atoms. In still other embodiments, the alkyl group contains 1–3 carbon atoms. In yet another embodiment, the alkyl group contains 1–2 carbons. Examples of alkyl radicals include, but are not limited to, methyl, ethyl, n-propyl, isopropyl, n-butyl, iso-butyl, sec-butyl, sec-pentyl, iso-pentyl, tert-butyl, n-pentyl, neopentyl, n-hexyl, sec-hexyl, n-heptyl, n-octyl, n-decyl, n-undecyl, dodecyl, and the like.

[0027] The term “alkenyl,” as used herein, denotes a monovalent group derived from a straight- or branched-chain aliphatic moiety having at least one carbon-carbon double bond by the removal of a single hydrogen atom. In certain embodiments, the alkenyl group employed in the invention contains 2–6 carbon atoms. In certain embodiments, the alkenyl group employed in the invention contains 2–5 carbon atoms. In some embodiments, the alkenyl group employed in the invention contains 2–4 carbon atoms. In another embodiment, the alkenyl group employed contains 2–3 carbon atoms. Alkenyl groups include, for example, ethenyl, propenyl, butenyl, 1-methyl-2-buten-1-yl, and the like.

[0028] The term “alkynyl,” as used herein, refers to a monovalent group derived from a straight- or branched-chain aliphatic moiety having at least one carbon-carbon triple bond by the removal of a single hydrogen atom. In certain embodiments, the alkynyl group employed in the invention contains 2–6 carbon atoms. In certain embodiments, the alkynyl group employed in the invention contains 2–5 carbon atoms. In some embodiments, the alkynyl group employed in the invention contains 2–4 carbon atoms. In another embodiment, the alkynyl group employed contains 2–3 carbon atoms. Representative alkynyl groups include, but are not limited to, ethynyl, 2-propynyl (propargyl), 1-propynyl, and the like.

[0029] The term “aryl” used alone or as part of a larger moiety as in “aralkyl”, “aralkoxy”, or “aryloxyalkyl”, refers to monocyclic and bicyclic ring systems having a total of five to 10 ring members, wherein at least one ring in the system is aromatic and wherein each ring in the system contains three to seven ring members. The term “aryl” may be used interchangeably with the term “aryl ring”. In certain embodiments of the present invention, “aryl” refers to an aromatic ring system which includes, but not limited to, phenyl, biphenyl, naphthyl, anthracyl and the like, which may bear one or more substituents. Also included within the scope of the term “aryl”, as it is used herein, is a group in which an aromatic ring is fused to one or more non-aromatic rings, such as indanyl, phthalimidyl, naphthimidyl, phenanthridinyl, or tetrahydronaphthyl, and the like.

[0030] The terms “heteroaryl” and “heteroar-”, used alone or as part of a larger moiety, e.g., “heteroaralkyl”, or “heteroaralkoxy”, refer to groups having 5 to 10 ring atoms, preferably 5, 6, or 9 ring atoms; having 6, 10, or 14 π electrons shared in a cyclic array; and having, in addition to carbon atoms, from one to five heteroatoms. The term “heteroatom” refers to nitrogen, oxygen, or sulfur, and includes any oxidized form of nitrogen or sulfur, and any quaternized form of a basic nitrogen. Heteroaryl groups include, without limitation, thienyl, furanyl, pyrrolyl, imidazolyl, pyrazolyl, triazolyl, tetrazolyl, oxazolyl, isoxazolyl, oxadiazolyl, thiazolyl, isothiazolyl, thiadiazolyl, pyridyl, pyridazinyl, pyrimidinyl, pyrazinyl, indoliziny, purinyl, naphthyridinyl, and pteridinyl. The terms “heteroaryl” and “heteroar-”, as used herein, also include groups in which a heteroaromatic ring is fused to one or more aryl, cycloaliphatic, or heterocyclyl rings, where the radical or point of attachment is on the heteroaromatic ring. Nonlimiting examples include indolyl, isoindolyl, benzothienyl, benzofuranyl, dibenzofuranyl, indazolyl, benzimidazolyl, benzthiazolyl, quinolyl, isoquinolyl, cinnoliny, phthalazinyl, quinazolinyl, quinoxalinyl, 4*H*-quinoliziny, carbazolyl, acridinyl, phenazinyl, phenothiazinyl, phenoxazinyl, tetrahydroquinoliny, tetrahydroisoquinoliny, and pyrido[2,3-*b*]-1,4-oxazin-3(4*H*)-one. A heteroaryl group may be mono- or bicyclic. The term “heteroaryl” may be used interchangeably with the terms “heteroaryl ring”, “heteroaryl group”, or “heteroaromatic”, any of which terms include rings that are optionally substituted. The term “heteroaralkyl” refers to an alkyl group substituted by a heteroaryl, wherein the alkyl and heteroaryl portions independently are optionally substituted.

[0031] As used herein, the terms “heterocycle”, “heterocyclyl”, “heterocyclic radical”, and “heterocyclic ring” are used interchangeably and refer to a stable 4- to 7-membered monocyclic

or 7–10-membered bicyclic heterocyclic moiety that is either saturated or partially unsaturated, and having, in addition to carbon atoms, one or more, preferably one to four, heteroatoms, as defined above. When used in reference to a ring atom of a heterocycle, the term "nitrogen" includes a substituted nitrogen. As an example, in a saturated or partially unsaturated ring having 0–3 heteroatoms selected from oxygen, sulfur or nitrogen, the nitrogen may be N (as in 3,4-dihydro-2*H*-pyrrolyl), NH (as in pyrrolidinyl), or ⁺NR (as in *N*-substituted pyrrolidinyl).

[0032] A heterocyclic ring can be attached to its pendant group at any heteroatom or carbon atom that results in a stable structure and any of the ring atoms can be optionally substituted. Examples of such saturated or partially unsaturated heterocyclic radicals include, without limitation, tetrahydrofuranyl, tetrahydrothienyl, pyrrolidinyl, pyrrolidonyl, piperidinyl, pyrrolinyl, tetrahydroquinolinyl, tetrahydroisoquinolinyl, decahydroquinolinyl, oxazolidinyl, piperazinyl, dioxanyl, dioxolanyl, diazepinyl, oxazepinyl, thiazepinyl, morpholinyl, and quinuclidinyl. The terms "heterocycle", "heterocyclyl", "heterocyclyl ring", "heterocyclic group", "heterocyclic moiety", and "heterocyclic radical", are used interchangeably herein, and also include groups in which a heterocyclyl ring is fused to one or more aryl, heteroaryl, or cycloaliphatic rings, such as indolinyl, 3*H*-indolyl, chromanyl, phenanthridinyl, or tetrahydroquinolinyl, where the radical or point of attachment is on the heterocyclyl ring. A heterocyclyl group may be mono- or bicyclic. The term "heterocyclylalkyl" refers to an alkyl group substituted by a heterocyclyl, wherein the alkyl and heterocyclyl portions independently are optionally substituted.

[0033] As used herein, the term "partially unsaturated" refers to a ring moiety that includes at least one double or triple bond between ring atoms. The term "partially unsaturated" is intended to encompass rings having multiple sites of unsaturation, but is not intended to include aryl or heteroaryl moieties, as herein defined.

[0034] The term "alkylene" refers to a bivalent alkyl group. An "alkylene chain" is a polymethylene group, i.e., $-(CH_2)_n-$, wherein *n* is a positive integer, preferably from 1 to 6, from 1 to 4, from 1 to 3, from 1 to 2, or from 2 to 3. A substituted alkylene chain is a polymethylene group in which one or more methylene hydrogen atoms are replaced with a substituent. Suitable substituents include those described below for a substituted aliphatic group.

[0035] As defined herein, an alkylene chain also can be optionally replaced by a functional group. An alkylene chain is "replaced" by a functional group when an internal methylene unit is

replaced with the functional group. Examples of suitable “interrupting functional groups” are described in the specification and claims herein.

[0036] As described herein, compounds of the invention may contain “optionally substituted” moieties. In general, the term “substituted”, whether preceded by the term “optionally” or not, means that one or more hydrogens of the designated moiety are replaced with a suitable substituent. Unless otherwise indicated, an “optionally substituted” group may have a suitable substituent at each substitutable position of the group, and when more than one position in any given structure may be substituted with more than one substituent selected from a specified group, the substituent may be either the same or different at every position. Combinations of substituents envisioned under this invention are preferably those that result in the formation of stable or chemically feasible compounds. The term “stable”, as used herein, refers to compounds that are not substantially altered when subjected to conditions to allow for their production, detection, and, in certain embodiments, their recovery, purification, and use for one or more of the purposes disclosed herein.

[0037] Suitable monovalent substituents on a substitutable carbon atom of an “optionally substituted” group are independently halogen; $-(CH_2)_{0-4}R^\circ$; $-(CH_2)_{0-4}OR^\circ$; $-O-(CH_2)_{0-4}C(O)OR^\circ$; $-(CH_2)_{0-4}CH(OR^\circ)_2$; $-(CH_2)_{0-4}SR^\circ$; $-(CH_2)_{0-4}Ph$, which may be substituted with R° ; $-(CH_2)_{0-4}O(CH_2)_{0-1}Ph$ which may be substituted with R° ; $-CH=CHPh$, which may be substituted with R° ; $-NO_2$; $-CN$; $-N_3$; $-(CH_2)_{0-4}N(R^\circ)_2$; $-(CH_2)_{0-4}N(R^\circ)C(O)R^\circ$; $-N(R^\circ)C(S)R^\circ$; $-(CH_2)_{0-4}N(R^\circ)C(O)NR^\circ_2$; $-N(R^\circ)C(S)NR^\circ_2$; $-(CH_2)_{0-4}N(R^\circ)C(O)OR^\circ$; $-N(R^\circ)N(R^\circ)C(O)R^\circ$; $-N(R^\circ)N(R^\circ)C(O)NR^\circ_2$; $-N(R^\circ)N(R^\circ)C(O)OR^\circ$; $-(CH_2)_{0-4}C(O)R^\circ$; $-C(S)R^\circ$; $-(CH_2)_{0-4}C(O)OR^\circ$; $-(CH_2)_{0-4}C(O)SR^\circ$; $-(CH_2)_{0-4}C(O)OSiR^\circ_3$; $-(CH_2)_{0-4}OC(O)R^\circ$; $-OC(O)(CH_2)_{0-4}SR^\circ$, $SC(S)SR^\circ$; $-(CH_2)_{0-4}SC(O)R^\circ$; $-(CH_2)_{0-4}C(O)NR^\circ_2$; $-C(S)NR^\circ_2$; $-C(S)SR^\circ$; $-SC(S)SR^\circ$, $-(CH_2)_{0-4}OC(O)NR^\circ_2$; $-C(O)N(OR^\circ)R^\circ$; $-C(O)C(O)R^\circ$; $-C(O)CH_2C(O)R^\circ$; $-C(NOR^\circ)R^\circ$; $-(CH_2)_{0-4}SSR^\circ$; $-(CH_2)_{0-4}S(O)_2R^\circ$; $-(CH_2)_{0-4}S(O)_2OR^\circ$; $-(CH_2)_{0-4}OS(O)_2R^\circ$; $-S(O)_2NR^\circ_2$; $-(CH_2)_{0-4}S(O)R^\circ$; $-N(R^\circ)S(O)_2NR^\circ_2$; $-N(R^\circ)S(O)_2R^\circ$; $-N(OR^\circ)R^\circ$; $-C(NH)NR^\circ_2$; $-P(O)_2R^\circ$; $-P(O)R^\circ_2$; $-OP(O)R^\circ_2$; $-OP(O)(OR^\circ)_2$; SiR°_3 ; $-(C_{1-4}$ straight or branched alkylene) $O-N(R^\circ)_2$; or $-(C_{1-4}$ straight or branched alkylene) $C(O)O-N(R^\circ)_2$, wherein each R° may be substituted as defined below and is independently hydrogen, C_{1-6} aliphatic, $-CH_2Ph$, $-O(CH_2)_{0-1}Ph$, or a 4–6-membered saturated, partially unsaturated, or aryl ring having 0–4 heteroatoms independently selected from nitrogen,

oxygen, or sulfur, or, notwithstanding the definition above, two independent occurrences of R[°], taken together with their intervening atom(s), form a 3–12-membered saturated, partially unsaturated, or aryl mono- or bicyclic ring having 0–4 heteroatoms independently selected from nitrogen, oxygen, or sulfur, which may be substituted as defined below.

[0038] Suitable monovalent substituents on R[°] (or the ring formed by taking two independent occurrences of R[°] together with their intervening atoms), are independently halogen, $-(CH_2)_{0-2}R^\bullet$, $-(haloR^\bullet)$, $-(CH_2)_{0-2}OH$, $-(CH_2)_{0-2}OR^\bullet$, $-(CH_2)_{0-2}CH(OR^\bullet)_2$, $-O(haloR^\bullet)$, $-CN$, $-N_3$, $-(CH_2)_{0-2}C(O)R^\bullet$, $-(CH_2)_{0-2}C(O)OH$, $-(CH_2)_{0-2}C(O)OR^\bullet$, $-(CH_2)_{0-2}SR^\bullet$, $-(CH_2)_{0-2}SH$, $-(CH_2)_{0-2}NH_2$, $-(CH_2)_{0-2}NHR^\bullet$, $-(CH_2)_{0-2}NR^\bullet_2$, $-NO_2$, $-SiR^\bullet_3$, $-OSiR^\bullet_3$, $-C(O)SR^\bullet$, $-(C_{1-4}$ straight or branched alkylene) $C(O)OR^\bullet$, or $-SSR^\bullet$ wherein each R[•] is unsubstituted or where preceded by “halo” is substituted only with one or more halogens, and is independently selected from C_{1–4} aliphatic, $-CH_2Ph$, $-O(CH_2)_{0-1}Ph$, or a 5–6-membered saturated, partially unsaturated, or aryl ring having 0–4 heteroatoms independently selected from nitrogen, oxygen, or sulfur. Suitable divalent substituents on a saturated carbon atom of R[°] include $=O$ and $=S$.

[0039] Suitable divalent substituents on a saturated carbon atom of an “optionally substituted” group include the following: $=O$, $=S$, $=NNR^*_2$, $=NNHC(O)R^*$, $=NNHC(O)OR^*$, $=NNHS(O)_2R^*$, $=NR^*$, $=NOR^*$, $-O(C(R^*_2))_{2-3}O-$, or $-S(C(R^*_2))_{2-3}S-$, wherein each independent occurrence of R^{*} is selected from hydrogen, C_{1–6} aliphatic which may be substituted as defined below, or an unsubstituted 5–6-membered saturated, partially unsaturated, or aryl ring having 0–4 heteroatoms independently selected from nitrogen, oxygen, or sulfur. Suitable divalent substituents that are bound to vicinal substitutable carbons of an “optionally substituted” group include: $-O(CR^*_2)_{2-3}O-$, wherein each independent occurrence of R^{*} is selected from hydrogen, C_{1–6} aliphatic which may be substituted as defined below, or an unsubstituted 5–6-membered saturated, partially unsaturated, or aryl ring having 0–4 heteroatoms independently selected from nitrogen, oxygen, or sulfur.

[0040] Suitable substituents on the aliphatic group of R^{*} include halogen, $-R^\bullet$, $-(haloR^\bullet)$, $-OH$, $-OR^\bullet$, $-O(haloR^\bullet)$, $-CN$, $-C(O)OH$, $-C(O)OR^\bullet$, $-NH_2$, $-NHR^\bullet$, $-NR^\bullet_2$, or $-NO_2$, wherein each R[•] is unsubstituted or where preceded by “halo” is substituted only with one or more halogens, and is independently C_{1–4} aliphatic, $-CH_2Ph$, $-O(CH_2)_{0-1}Ph$, or a 5–6-membered

saturated, partially unsaturated, or aryl ring having 0–4 heteroatoms independently selected from nitrogen, oxygen, or sulfur.

[0041] Suitable substituents on a substitutable nitrogen of an “optionally substituted” group include $-R^\dagger$, $-NR^\dagger_2$, $-C(O)R^\dagger$, $-C(O)OR^\dagger$, $-C(O)C(O)R^\dagger$, $-C(O)CH_2C(O)R^\dagger$, $-S(O)_2R^\dagger$, $-S(O)_2NR^\dagger_2$, $-C(S)NR^\dagger_2$, $-C(NH)NR^\dagger_2$, or $-N(R^\dagger)S(O)_2R^\dagger$; wherein each $R^\dagger$ is independently hydrogen, C_{1-6} aliphatic which may be substituted as defined below, unsubstituted $-OPh$, or an unsubstituted 5–6-membered saturated, partially unsaturated, or aryl ring having 0–4 heteroatoms independently selected from nitrogen, oxygen, or sulfur, or, notwithstanding the definition above, two independent occurrences of $R^\dagger$, taken together with their intervening atom(s) form an unsubstituted 3–12-membered saturated, partially unsaturated, or aryl mono- or bicyclic ring having 0–4 heteroatoms independently selected from nitrogen, oxygen, or sulfur.

[0042] Suitable substituents on the aliphatic group of $R^\dagger$ are independently halogen, $-R^\bullet$, $-(haloR^\bullet)$, $-OH$, $-OR^\bullet$, $-O(haloR^\bullet)$, $-CN$, $-C(O)OH$, $-C(O)OR^\bullet$, $-NH_2$, $-NHR^\bullet$, $-NR^\bullet_2$, or $-NO_2$, wherein each $R^\bullet$ is unsubstituted or where preceded by “halo” is substituted only with one or more halogens, and is independently C_{1-4} aliphatic, $-CH_2Ph$, $-O(CH_2)_{0-1}Ph$, or a 5–6-membered saturated, partially unsaturated, or aryl ring having 0–4 heteroatoms independently selected from nitrogen, oxygen, or sulfur.

3. Description of Exemplary Compounds:

[0043] As defined generally above, each of R^x and R^y is independently selected from $-R^2$, $-halo$, $-NO_2$, $-CN$, $-OR^2$, $-SR^2$, $-N(R^2)_2$, $-C(O)R^2$, $-CO_2R^2$, $-C(O)C(O)R^2$, $-C(O)CH_2C(O)R^2$, $-S(O)R^2$, $-S(O)_2R^2$, $-C(O)N(R^2)_2$, $-SO_2N(R^2)_2$, $-OC(O)R^2$, $-N(R^2)C(O)R^2$, $-N(R^2)N(R^2)_2$, $-N(R^2)-C(=NR^2)N(R^2)_2$, $-C(=NR^2)N(R^2)_2$, $-C=NOR^2$, $-N(R^2)C(O)N(R^2)_2$, $-N(R^2)SO_2N(R^2)_2$, $-N(R^2)SO_2R^2$, or $-OC(O)N(R^2)_2$, wherein R^2 is as defined above and described herein.

[0044] In certain embodiments, each of R^x and R^y is independently selected from $-R^2$, $halo$, $-OR^2$, $-N(R^2)_2$, $-OC(O)R^2$, $-N(R^2)C(O)R^2$, $-N(R^2)N(R^2)_2$, $-N(R^2)C(O)N(R^2)_2$, $-N(R^2)SO_2N(R^2)_2$, $-N(R^2)SO_2R^2$, or $-OC(O)N(R^2)_2$; wherein R^2 is as defined above and described herein. In some embodiments, each of R^x and R^y is independently selected from $-R^2$, $halo$, $-OR^2$, and $-N(R^2)_2$. In other embodiments, each of R^x and R^y is independently hydrogen, $halo$, $-OR^2$, $-N(R^2)_2$, or an optionally substituted group selected from C_{1-6} aliphatic or a 5–10

membered saturated, partially unsaturated, or aromatic monocyclic or bicyclic ring having 1–4 heteroatoms independently selected from nitrogen, oxygen, or sulfur.

[0045] In certain embodiments, R^x is selected from $-R^2$, $-\text{halo}$, $-\text{CN}$, or $-\text{CO}_2R^2$.

[0046] In certain embodiments, R^x is R^2 or halo . In some embodiments, R^x is hydrogen, CN , an optionally substituted C_{1-6} aliphatic group, or halo . In certain embodiments, R^x is hydrogen. In certain embodiments, R^x is fluoro, chloro or bromo. In other embodiments, R^x is chloro.

[0047] In certain embodiments, R^x is an optionally substituted C_{1-6} aliphatic group. In some embodiments, R^x is an optionally substituted C_{1-6} alkyl group. In other embodiments, R^x is an optionally substituted C_{1-3} alkyl group. In certain embodiments, R^x is an optionally substituted methyl, ethyl, n -propyl or isopropyl group. According to one embodiment, R^x is an optionally substituted methyl group. According to another embodiment, one or more substituents present on the C_{1-6} aliphatic, C_{1-6} alkyl, C_{1-3} alkyl, n -propyl, isopropyl, ethyl or methyl group include $-\text{N}(\text{R}^2)_2$, wherein R^2 is as defined above and described herein. In certain embodiments, R^x is $-\text{CF}_3$.

[0048] Exemplary R^x groups include those set forth in **Tables 1, 3, 4, and 5** in the Examples section, *infra*.

[0049] In certain embodiments, R^y is selected from $-R^2$, $-\text{OR}^2$, or $-\text{N}(\text{R}^2)_2$. In certain embodiments, R^y is independently selected from hydrogen, $-\text{OR}^2$, $-\text{N}(\text{R}^2)_2$, or an optionally substituted group selected from C_{1-6} aliphatic or a 5–10 membered saturated, partially unsaturated, or aromatic monocyclic or bicyclic ring having 1–4 heteroatoms independently selected from nitrogen, oxygen, or sulfur.

[0050] In certain embodiments, R^y is hydrogen.

[0051] In some embodiments, R^y is an optionally substituted C_{1-6} aliphatic group. In other embodiments, R^y is an optionally substituted C_{2-6} aliphatic group. In certain embodiments, R^y is an optionally substituted C_{2-6} alkenyl group. In certain embodiments, R^y is an optionally substituted C_{2-6} alkynyl group. According to one embodiment, R^y is an optionally substituted C_{2-5} alkynyl group. According to another embodiment, substituents present on the C_{1-6} aliphatic, C_{2-6} aliphatic, C_{2-6} alkenyl, C_{2-6} alkynyl or C_{2-5} alkynyl R^y group include $-(\text{CH}_2)_{0-4}\text{OR}^o$ or $-(\text{CH}_2)_{0-4}\text{N}(\text{R}^o)_2$ groups, wherein R^o is as defined above and herein.

[0052] In certain embodiments, R^y is an optionally substituted C₆₋₁₀ monocyclic or bicyclic aryl ring. In certain embodiments, R^y is an optionally substituted C₈₋₁₀ bicyclic aryl ring. In some embodiments, R^y is an optionally substituted phenyl ring.

[0053] According to one embodiment, R^y is an optionally substituted 5–10 membered saturated monocyclic or bicyclic ring having 1–4 heteroatoms independently selected from nitrogen, oxygen, or sulfur. In certain embodiments, R^y is an optionally substituted 5,6– or 6,6– fused saturated bicyclic ring having 1–4 heteroatoms independently selected from nitrogen, oxygen, or sulfur. In some embodiments, R^y is an optionally substituted 5–6 membered saturated monocyclic ring having 1–4 heteroatoms independently selected from nitrogen, oxygen, or sulfur. In other embodiments, R^y is an optionally substituted 5–6 membered saturated monocyclic ring having 1–2 heteroatoms independently selected from nitrogen, oxygen, or sulfur.

[0054] In certain embodiments, R^y is an optionally substituted 5–membered saturated monocyclic ring having 1–2 heteroatoms independently selected from nitrogen, oxygen, or sulfur. In some embodiments, R^y is an optionally substituted 5–membered saturated monocyclic ring having 2 heteroatoms independently selected from nitrogen, oxygen, or sulfur. In other embodiments, R^y is an optionally substituted 5–membered saturated monocyclic ring having 2 heteroatoms independently selected from nitrogen or oxygen.

[0055] In certain embodiments, R^y is an optionally substituted 6–membered saturated monocyclic ring having 1–2 heteroatoms independently selected from nitrogen, oxygen, or sulfur. In some embodiments, R^y is an optionally substituted 6–membered saturated monocyclic ring having 2 heteroatoms independently selected from nitrogen, oxygen, or sulfur. In other embodiments, R^y is an optionally substituted 6–membered saturated monocyclic ring having 2 heteroatoms independently selected from nitrogen or oxygen.

[0056] Exemplary R^y groups include optionally substituted octahydroazocinyl, thiocyclopentanyl, thiocyclohexanyl, pyrrolidinyl, piperidinyl, piperazinyl, tetrahydrothiopyranyl, tetrahydrothiophenyl, dithiolanyl, tetrahydrofuranyl, tetrahydropyranyl, dioxanyl, thioxanyl, morpholinyl, oxathiolanyl, imidazolidinyl, oxathiolanyl, oxazolidinyl, or thiazolidinyl groups. In certain embodiments, R^y is an optionally substituted imidazolidinyl, oxathiolanyl, oxazolidinyl, or thiazolidinyl group. In some embodiments, R^y is an optionally

substituted piperidinyl, piperazinyl, morpholinyl, or pyrrolidinyl group. In other embodiments, R^y is an optionally substituted morpholinyl group.

[0057] In certain embodiments, R^y is an optionally substituted 5-membered heteroaryl ring having 1–3 heteroatoms selected from nitrogen, oxygen, or sulfur. In some embodiments, R^y is an optionally substituted 5-membered heteroaryl ring having 1–2 heteroatoms selected from nitrogen, oxygen, or sulfur. In other embodiments, R^y is an optionally substituted 5-membered heteroaryl ring having 2 heteroatoms selected from nitrogen, oxygen, or sulfur. According to one aspect, R^y is an optionally substituted 5-membered heteroaryl ring having 1 heteroatom selected from nitrogen, oxygen, or sulfur. In certain embodiments, R^y is an optionally substituted 5-membered heteroaryl ring having 1 nitrogen atom, and an additional heteroatom selected from sulfur or oxygen. Exemplary R^y groups include optionally substituted pyrrolyl, pyrazolyl, imidazolyl, triazolyl, tetrazolyl, thiophenyl, furanyl, thiazolyl, isothiazolyl, thiadiazolyl, oxazolyl, isoxazolyl, or oxadiazolyl group.

[0058] In certain embodiments, R^y is an optionally substituted 6-membered heteroaryl ring having 1–4 heteroatoms independently selected from nitrogen, oxygen, or sulfur. In some embodiments, R^y is an optionally substituted 6-membered heteroaryl ring having 1–3 nitrogen atoms. In other embodiments, R^y is an optionally substituted 6-membered heteroaryl ring having 1–2 nitrogen atoms. According to one aspect, R^y is an optionally substituted 6-membered heteroaryl ring having 2 heteroatoms nitrogen atoms. Exemplary R^y groups include an optionally substituted pyridinyl, pyrimidinyl, pyrazolyl, pyrazinyl, pyridazinyl, triazinyl, or tetrazinyl group. In certain embodiments, R^y is an optionally substituted pyridinyl group.

[0059] In certain embodiments, R^y is an optionally substituted 5–10 membered partially unsaturated monocyclic or bicyclic ring having 1–3 heteroatoms independently selected from nitrogen, oxygen, or sulfur. In some embodiments, R^y is an optionally substituted 5–6 membered partially unsaturated monocyclic ring having 1–2 heteroatoms independently selected from nitrogen, oxygen, or sulfur. In other embodiments, R^y is an optionally substituted tetrahydropyridinyl group.

[0060] In certain embodiments, R^y is an optionally substituted 8–10 membered aromatic bicyclic heteroaryl ring having 1–4 heteroatoms independently selected from nitrogen, oxygen, or sulfur. In some embodiments, R^y is an optionally substituted 5,6-fused heteroaryl ring having 1–4 heteroatoms independently selected from nitrogen, oxygen, or sulfur. In other embodiments,

R^y is an optionally substituted 5,6-fused heteroaryl ring having 1–2 heteroatoms independently selected from nitrogen, oxygen, or sulfur. In certain embodiments, R^y is an optionally substituted 5,6-fused heteroaryl ring having 1 heteroatom independently selected from nitrogen, oxygen, or sulfur.

[0061] In certain embodiments, R^y is an optionally substituted 6,6-fused heteroaryl ring having 1–4 heteroatoms independently selected from nitrogen, oxygen, or sulfur. In some embodiments, R^y is an optionally substituted 6,6-fused heteroaryl ring having 1–2 heteroatoms independently selected from nitrogen, oxygen, or sulfur. In other embodiments, R^y is an optionally substituted 6,6-fused heteroaryl ring having 1 heteroatom independently selected from nitrogen, oxygen, or sulfur. According to one aspect, R^y is an optionally substituted 6,6-fused heteroaryl ring having 2 heteroatoms independently selected from nitrogen, oxygen, or sulfur. Exemplary R^y groups include an optionally substituted benzofuranyl, thianaphthenyl, pyrroliziny, indolyl, quinoliny, isoquinoliny, benzimidazolyl, imidazopyridiny, puriny, indazolyl, pyrrolopyridiny, cinnoliny, quinazoliny, phthalazinyl, naphthyridiny, or quinoxaliny group. In some embodiments, R^y is a pyrrolylpyridiny, imidazopyridiny, or puriny group. In other embodiments, R^y is a pyrrolylpyridiny group.

[0062] In certain embodiments, R^y is $-OR^2$, wherein R^2 is defined above and described herein. In certain embodiments, R^y is $-OR^2$, wherein R^2 is hydrogen or an optionally substituted C_{1-6} aliphatic group. In some embodiments, R^y is $-OR^2$, wherein R^2 is an optionally substituted C_{1-6} aliphatic group. In other embodiments, R^y is $-OR^2$, wherein R^2 is an optionally substituted C_{1-6} alkyl group. According to one aspect, R^y is $-OR^2$, wherein R^2 is an optionally substituted C_{1-3} alkyl group. In other embodiments, R^y is $-OR^2$, wherein R^2 is an optionally substituted C_{1-2} alkyl group. In some embodiments, R^y is $-OCH_3$. In other embodiments, R^y is $-OH$. In yet other embodiments, R^y is $-OR^2$, wherein R^2 is $-(CH_2)_{0-3}CH_2N(R^o)_2$, and wherein each R^o is defined and described herein.

[0063] In certain embodiments, R^y is $-N(R^2)_2$, wherein R^2 is defined above and described herein. In other embodiments, R^y is $-N(R^2)_2$, wherein each R^2 is independently hydrogen or an optionally substituted C_{1-6} aliphatic group.

[0064] In certain embodiments, R^y is $-NH_2$.

[0065] In certain embodiments, R^y is $-NHR^2$, wherein R^2 is an optionally substituted C_{1-6} aliphatic group. In some embodiments, R^y is $-NHR^2$, wherein R^2 is an optionally substituted C_{1-6}

₆ alkyl group. In other embodiments, R^y is -NHR², wherein R² is an optionally substituted C₁₋₃ alkyl group. According to one aspect, R^y is -NHR², wherein R² is an optionally substituted methyl or ethyl. Exemplary R^y groups include -NHCH₃, -NHCH₂CH₃, -NHCH₂CH₂CH₃, -NHCH(CH₃)₂, or -NH(C₃H₅), NHCH₂CH₂CH₂OH, and -N(CH₂CH₂)₂O, -NHCH₂CH₂CH₂NH(CH₃)₂.

[0066] In certain embodiments, R^y is -N(R²)₂, wherein each R² is independently hydrogen or an optionally substituted C₆₋₁₀ monocyclic or bicyclic aryl ring. In certain embodiments, R^y is -NHR², wherein R² is an optionally substituted C₆₋₁₀ monocyclic or bicyclic aryl ring. In certain embodiments, R^y is -NHR², wherein R² is an optionally substituted C₆ monocyclic aryl ring. In certain embodiments, R^y is -NHR², wherein R² is an optionally substituted C₈₋₁₀ bicyclic aryl ring.

[0067] In certain embodiments, R^y is -NHR², wherein R² is an optionally substituted 5-10 membered monocyclic or bicyclic heteroaryl ring having 1-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur. In some embodiments, R^y is -NHR², wherein R² is an optionally substituted 5-6 membered heteroaryl ring having 1-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur. In other embodiments, R^y is -NHR², wherein R² is an optionally substituted 5-6 membered heteroaryl ring having 1-2 heteroatoms independently selected from nitrogen, oxygen, or sulfur. In certain aspects, R^y is -NHR², wherein R² is an optionally substituted 5-6 membered heteroaryl ring having 1-2 nitrogen atoms.

[0068] In certain embodiments, R^y is -NHR², wherein R² is an optionally substituted 5 membered heteroaryl ring having 1-2 heteroatoms selected from nitrogen, oxygen, or sulfur. In some embodiments, R^y is -NHR², wherein R² is an optionally substituted 5 membered heteroaryl ring having 2 heteroatoms selected from nitrogen, oxygen, or sulfur. In other embodiments, R^y is -NHR², wherein R² is an optionally substituted 5 membered heteroaryl ring having 1 heteroatom selected from nitrogen, oxygen, or sulfur. In certain embodiments, R^y is -NHR², wherein R² is an optionally substituted 5 membered heteroaryl ring having a nitrogen atom, and another heteroatom selected from sulfur or oxygen.

[0069] In certain embodiments, R^y is -NHR², wherein R² is an optionally substituted 6 membered heteroaryl ring having 1-2 heteroatoms independently selected from nitrogen, oxygen, or sulfur. In certain embodiments, R^y is -NHR², wherein R² is an optionally substituted 6 membered heteroaryl ring having 1 heteroatom independently selected from nitrogen, oxygen,

or sulfur. In certain embodiments, R^y is $-NHR^2$ wherein R^2 is an optionally substituted 6 membered heteroaryl ring having 2 heteroatoms independently selected from nitrogen, oxygen, or sulfur. In certain embodiments, R^y is $-NHR^2$, wherein R^2 is an optionally substituted 6 membered heteroaryl ring having 1 heteroatom selected from nitrogen, and 1 heteroatom selected from sulfur or oxygen.

[0070] In certain embodiments, R^y is $-NHR^2$, wherein R^2 is an optionally substituted 5,6-fused heteroaryl ring having 1–4 heteroatoms independently selected from nitrogen, oxygen, or sulfur. In some embodiments, R^y is $-NHR^2$, wherein R^2 is an optionally substituted 5,6-fused heteroaryl ring having 1–2 heteroatoms independently selected from nitrogen, oxygen, or sulfur. In other embodiments, R^y is $-NHR^2$, wherein R^2 is an optionally substituted 5,6-fused heteroaryl ring having 1 heteroatom independently selected from nitrogen, oxygen, or sulfur. In certain aspects, R^y is $-NHR^2$, wherein R^2 is an optionally substituted 5,6-fused heteroaryl ring having 2 heteroatoms independently selected from nitrogen, oxygen, or sulfur.

[0071] In certain embodiments, R^y is $-NHR^2$, wherein R^2 is an optionally substituted 6,6-fused heteroaryl ring having 1–4 heteroatoms independently selected from nitrogen, oxygen, or sulfur. In some embodiments, R^y is $-NHR^2$, wherein R^2 is an optionally substituted 6,6-fused heteroaryl ring having 1–2 heteroatoms independently selected from nitrogen, oxygen, or sulfur. In other embodiments, R^y is $-NHR^2$, wherein R^2 is an optionally substituted 6,6-fused heteroaryl ring having 1 heteroatom independently selected from nitrogen, oxygen, or sulfur. In certain aspects, R^y is $-NHR^2$, wherein R^2 is an optionally substituted 6,6-fused heteroaryl ring having 2 heteroatoms independently selected from nitrogen, oxygen, or sulfur.

[0072] In certain embodiments, R^y is $-NHR^2$, wherein R^2 is an optionally substituted pyrrolyl, pyrazolyl, imidazolyl, pyridinyl, pyrimidinyl, pyrazinyl, pyridazinyl, triazinyl, tetrazinyl, tetrazolyl, pyrolyl, indolyl, quinolyl, isoquinolyl, benzimidazolyl, pyrrolylpyridinyl, indazolyl, cinnolyl, quinazolinyl, phthalazinyl, naphthyridinyl, quinoxalinyl, thiophenyl, thienyl, thianaphthenyl, furanyl, benzofuranyl, thiazolyl, isothiazolyl, thiadiazolyl, oxazolyl, isoxazolyl, or oxadiazolyl group. In other embodiments, R^y is $-NHR^2$, wherein R^2 is an optionally substituted pyridinyl, thiazolyl, isothiazolyl, oxazolyl, or isoxazolyl group. In certain embodiments, R^y is $-NHR^2$, wherein R^2 is an optionally substituted pyridinyl, thiazolyl or isoxazolyl group.

[0073] In certain embodiments, R^y is $-N(R^2)_2$, wherein two R^2 groups on the same nitrogen are taken together with the nitrogen to form a 5–8 membered saturated, partially unsaturated, or aromatic mono- or bicyclic ring having 1–4 heteroatoms independently selected from nitrogen, oxygen, or sulfur. In some embodiments, R^y is $-N(R^2)_2$, wherein two R^2 on the same nitrogen are taken together with the nitrogen to form an optionally substituted piperidinyl, piperazinyl, pyrrolidinyl, octahydroazocinyl or morpholinyl group. In other embodiments, R^y is $-N(R^2)_2$, wherein two R^2 on the same nitrogen are taken together with the nitrogen to form an optionally substituted piperidinyl, piperazinyl, morpholinyl, or pyrrolidinyl group. In certain aspects, R^y is $-N(R^2)_2$, wherein two R^2 on the same nitrogen are taken together with the nitrogen to form an optionally substituted morpholinyl group.

[0074] Exemplary R^y groups include those set forth in the Examples section, *infra*.

[0075] In certain embodiments, R^1 is hydrogen. In other embodiments, R^1 is an optionally substituted C_{1-6} aliphatic group. In certain embodiments, R^1 is an optionally substituted C_{1-6} alkyl group. In some embodiments, R^1 is an optionally substituted C_{1-3} alkyl group. In certain aspects, R^1 is an optionally substituted methyl or ethyl group. In certain embodiments, R^1 is an optionally substituted methyl group.

[0076] As defined above, L^1 is a direct bond or an optionally substituted, straight or branched C_{1-6} alkylene chain. In some embodiments, L^1 is a direct bond. In certain embodiments, L^1 is an optionally substituted, straight or branched C_{1-5} alkylene chain. In some embodiments, L^1 is an optionally substituted, straight or branched C_{1-4} alkylene chain. In other embodiments, L^1 is an optionally substituted, straight or branched C_{1-3} alkylene chain. According to some embodiments, L^1 is an optionally substituted, straight or branched C_{1-2} alkylene chain.

[0077] In certain embodiments, L^1 is an optionally substituted, straight or branched C_1 alkylene chain. In some embodiments, L^1 is an optionally substituted, straight or branched C_2 alkylene chain. In other embodiments, L^1 is an optionally substituted, straight or branched C_3 alkylene chain. According to some embodiments, L^1 is an optionally substituted, straight or branched C_4 alkylene chain. In certain aspects, L^1 is an optionally substituted, straight or branched C_5 alkylene chain. In other aspects, L^1 is an optionally substituted, straight or branched C_6 alkylene chain.

[0078] In certain embodiments, L^1 is an optionally substituted, straight C_{1-6} alkylene chain. In some embodiments, L^1 is a straight C_{1-6} alkylene chain. In other embodiments, L^1 is an

optionally substituted, branched C₁₋₆ alkylene chain. In certain aspects, L¹ is a branched C₁₋₆ alkylene chain. In certain embodiments, L¹ is -CH(C₁₋₆alkyl)-, -CH(C₁₋₅alkyl)-, -CH(C₁₋₄alkyl)-, -CH(C₁₋₃alkyl)-, or -CH(C₁₋₂alkyl)-. In certain embodiments, L¹ is -CH(CH₃)-. Exemplary L¹ groups include -CH₂-, -C(CH₃)₂-, -CH(CF₃)-, -CH(CHF₂)-, -CH(CH₂F)-, -CH(CH₂OH)-, -CH(CH₂NH₂)-, -CH(OCH₃)-, -CH(NHCH₃)-, -CH(N(CH₃)₂)-, -CH(SCH₃)-, -CH(=O)-, and -C(=CH₂)-.

[0079] Exemplary L¹ groups include those set forth in **Tables 2, 3, 4, and 5** in the Examples section, *infra*.

[0080] As defined generally above, Cy¹ is an optionally substituted phenyl or an optionally substituted 5-6 membered saturated, partially unsaturated, or aromatic ring having 1-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur. In some embodiments, Cy¹ is optionally substituted phenyl. In certain embodiments, Cy¹ is an optionally substituted 6 membered saturated, partially unsaturated, or aromatic ring having 1-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur. In other embodiments, Cy¹ is an optionally substituted 5-membered saturated, partially unsaturated, or aromatic ring having 1-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur. In certain aspects, Cy¹ is an optionally substituted 5-membered heteroaryl ring having 2 heteroatoms independently selected from nitrogen, oxygen, or sulfur. In other embodiments, Cy¹ is an optionally substituted 5-membered heteroaryl ring having 2 heteroatoms independently selected from nitrogen and oxygen. In some embodiments, Cy¹ is an optionally substituted 5-membered heteroaryl ring having 2 heteroatoms independently selected from nitrogen and sulfur.

[0081] Exemplary Cy¹ groups include an optionally substituted pyrrolyl, pyrazolyl, imidazolyl, triazolyl, tetrazolyl, thiophenyl, furanyl, thiazolyl, isothiazolyl, thiadiazolyl, oxazolyl, isoxazolyl, or oxadiazolyl group. In certain embodiments, Cy¹ is an optionally substituted thiazolyl or isoxazolyl group. In other embodiments, Cy¹ is an optionally substituted thiazolyl group. In some embodiments, Cy¹ is an unsubstituted thiazolyl group. In certain aspects, Cy¹ is an optionally substituted isoxazolyl group. According to another aspect, Cy¹ is an unsubstituted isoxazolyl group.

[0082] In other embodiments, Cy¹ is an optionally substituted 5-6 membered saturated ring having 1-2 heteroatoms independently selected from nitrogen and oxygen. In certain embodiments, Cy¹ is optionally substituted piperidinyl or pyrrolidinyl.

[0083] In other embodiments, Cy¹ is an optionally substituted 6-membered saturated, partially unsaturated or aryl ring having 1-2 nitrogens. In certain embodiments, Cy¹ is an optionally substituted pyridine or pyrimidine ring.

[0084] Exemplary Cy¹ groups include those set forth in the Examples section, *infra*.

[0085] As defined generally above, L² is a direct bond, or is an optionally substituted, straight or branched C₁₋₆ alkylene chain wherein 1 or 2 methylene units of L² are optionally and independently replaced by -O-, -S-, -N(R)-, -C(O)-, -C(O)N(R)-, -N(R)C(O)N(R)-, -N(R)C(O)-, -N(R)C(O)O-, -OC(O)N(R)-, -SO₂-, -SO₂N(R)-, -N(R)SO₂-, -OC(O)-, -C(O)O-, or a 3-6 membered cycloalkylene. In certain embodiments, L² is a direct bond.

[0086] In certain embodiments, L² is an optionally substituted, straight or branched C₁₋₆ alkylene chain wherein 1 or 2 methylene units of L² are replaced by -O-, -S-, -N(R)-, -C(O)-, -C(O)N(R)-, -N(R)C(O)N(R)-, -N(R)C(O)-, -N(R)C(O)O-, -OC(O)N(R)-, -SO₂-, -SO₂N(R)-, -N(R)SO₂-, -OC(O)-, or -C(O)O-; wherein each R is as defined above and described herein. In some embodiments, L² is an optionally substituted, straight or branched C₁₋₄ alkylene chain wherein 1 or 2 methylene units of L² are replaced by -O-, -S-, -N(R)-, -C(O)-, -C(O)N(R)-, -N(R)C(O)N(R)-, -N(R)C(O)-, -N(R)C(O)O-, -OC(O)N(R)-, -SO₂-, -SO₂N(R)-, -N(R)SO₂-, -OC(O)-, or -C(O)O-. In other embodiments, L² is an optionally substituted, straight or branched C₁₋₂ alkylene chain wherein 1 methylene unit of L² is replaced by -O-, -S-, -N(R)-, -C(O)-, -C(O)N(R)-, -N(R)C(O)N(R)-, -N(R)C(O)-, -N(R)C(O)O-, -OC(O)N(R)-, -SO₂-, -SO₂N(R)-, -N(R)SO₂-, -OC(O)-, or -C(O)O-. In certain aspects, L² is -O-, -S-, -N(R)-, -C(O)-, -C(O)N(R)-, -N(R)C(O)-, -SO₂-, -SO₂N(R)-, -N(R)SO₂-, -OC(O)-, or -C(O)O-. In other embodiments, L² is -C(O)N(R)-, -N(R)C(O)-, -SO₂N(R)-, -N(R)SO₂-, -OC(O)-, or -C(O)O-. In certain aspects, L² is -C(O)N(R)- or -N(R)C(O)-. In certain embodiments, L² is -C(O)N(H)- or -N(H)C(O)-. In certain embodiments, L² is -C(O)N(H)-.

[0087] Exemplary L² groups include those set forth in the Examples section, *infra*.

[0088] As defined generally above, Cy² is an optionally substituted 5-14 membered saturated, partially unsaturated, or aromatic monocyclic, bicyclic, or tricyclic ring having 0-4 heteroatoms, independently selected from nitrogen, oxygen, or sulfur.

[0089] In some embodiments, Cy² is optionally substituted phenyl.

[0090] In certain embodiments, Cy² is an optionally substituted 5-10 membered saturated, partially unsaturated, or aromatic monocyclic ring having 1-4 heteroatoms, independently

selected from nitrogen, oxygen, or sulfur. In other embodiments, Cy² is an optionally substituted 5–6 membered saturated, partially unsaturated, or aromatic monocyclic ring having 1–4 heteroatoms, independently selected from nitrogen, oxygen, or sulfur.

[0091] In certain embodiments, Cy² is an optionally substituted 5-membered saturated, partially unsaturated, or aromatic monocyclic ring having 1–3 heteroatoms, independently selected from nitrogen, oxygen, or sulfur. In some embodiments, Cy² is an optionally substituted 5-membered saturated, partially unsaturated, or aromatic monocyclic ring having 1–2 heteroatoms, independently selected from nitrogen, oxygen, or sulfur. In other embodiments, Cy² is an optionally substituted 5-membered heteroaryl ring having 1–3 heteroatoms, independently selected from nitrogen, oxygen, or sulfur. In still other embodiments, Cy² is an optionally substituted 5-membered heteroaryl ring having 1–2 heteroatoms, independently selected from nitrogen. Exemplary Cy² groups include an optionally substituted pyrrolyl, pyrazolyl, imidazolyl, triazolyl, tetrazolyl, thiophenyl, furanyl, thiazolyl, isothiazolyl, thiadiazolyl, oxazolyl, isoxazolyl, or oxadiazolyl group.

[0092] In certain embodiments, Cy² is an optionally substituted 6-membered saturated, partially unsaturated, or aromatic monocyclic ring having 1–4 heteroatoms, independently selected from nitrogen, oxygen, or sulfur. In some embodiments, Cy² is an optionally substituted 6-membered saturated, partially unsaturated, or aromatic monocyclic ring having 1–2 heteroatoms, independently selected from nitrogen, oxygen, or sulfur. In other embodiments, Cy² is an optionally substituted 6-membered heteroaryl ring having 1–4 nitrogen atoms. In certain aspects, Cy² is an optionally substituted 6-membered heteroaryl ring having 1–3 nitrogen atoms. In some embodiments, Cy² is an optionally substituted 6-membered heteroaryl ring having 1–2 nitrogen atoms. Exemplary Cy² groups include an optionally substituted pyridinyl, pyrimidinyl, pyrazinyl, pyridazinyl, triazinyl, or tetrazinyl group. In some embodiments, Cy² is an optionally substituted pyridinyl, pyrimidinyl or pyridazinyl group.

[0093] In certain embodiments, Cy² is an optionally substituted 8–10 membered saturated, partially unsaturated, or aromatic bicyclic ring having 1–4 heteroatoms, independently selected from nitrogen, oxygen, or sulfur. In some embodiments, Cy² is an optionally substituted 5,5–fused, 5,6–fused, or 6,6–fused saturated, partially unsaturated, or aromatic bicyclic ring having 1–4 heteroatoms, independently selected from nitrogen, oxygen, or sulfur. In other embodiments, Cy² is an optionally substituted 5,5–fused, 5,6–fused, or 6,6–fused heteroaryl ring

having 1–4 heteroatoms, independently selected from nitrogen, oxygen, or sulfur. In certain aspects, Cy^2 is an optionally substituted 5,5–fused, 5,6–fused, or 6,6–fused heteroaryl ring having 1–4 nitrogen atoms. In other embodiments, Cy^2 is an optionally substituted 5,6–fused heteroaryl ring having 1–4 nitrogen atoms. Exemplary Cy^2 groups include an optionally substituted pyrroliziny, indolyl, quinoliny, isoquinoliny, benzimidazolyl, imidazopyridiny, indazolyl, puriny, cinnoliny, quinazoliny, phthalaziny, naphthridiny, quinoxaliny, thianaphtheneyl, or benzofuranyl group. In certain aspects, Cy^2 is an optionally substituted benzimidazolyl, imidazopyridiny or puriny group.

[0094] In certain embodiments, Cy^2 is an optionally substituted 5–10 membered saturated, partially unsaturated, or aromatic monocyclic or bicyclic carbocyclic ring. In some embodiments, Cy^2 is an optionally substituted 5–10 membered saturated, partially unsaturated, or aromatic monocyclic or bicyclic carbocyclic ring. In other embodiments, Cy^2 is an optionally substituted 5–6 membered saturated, partially unsaturated, or aromatic monocyclic carbocyclic ring. In certain aspects, Cy^2 is an optionally substituted 5-membered saturated or partially unsaturated carbocyclic ring. According to one embodiment, Cy^2 is an optionally substituted 6 membered saturated, partially unsaturated, or aromatic ring. In still other embodiments, Cy^2 is an optionally substituted phenyl group.


[0095] In certain embodiments, Cy^2 is an optionally substituted 5,5–fused–, 5,6–fused, or 6,6–fused saturated, partially unsaturated, or aromatic bicyclic ring. In some embodiments, Cy^2 is an optionally substituted 5,5–fused, 5,6–fused, or 6,6–fused aromatic bicyclic ring. In other embodiments, Cy^2 is optionally substituted naphthalenyl, indanyl or indenyl group.

[0096] In certain embodiments, Cy^2 , as described above and herein, is optionally substituted with one or more groups selected from $-R^\circ$, $-\text{halo}$, $-\text{NO}_2$, $-\text{CN}$, $-\text{OR}^\circ$, $-\text{SR}^\circ$, $-\text{N}(\text{R}^\circ)_2$, $-\text{C}(\text{O})\text{R}^\circ$, $-\text{CO}_2\text{R}^\circ$, $-\text{C}(\text{O})\text{C}(\text{O})\text{R}^\circ$, $-\text{C}(\text{O})\text{CH}_2\text{C}(\text{O})\text{R}^\circ$, $-\text{S}(\text{O})\text{R}^\circ$, $-\text{S}(\text{O})_2\text{R}^\circ$, $-\text{C}(\text{O})\text{N}(\text{R}^\circ)_2$, $-\text{SO}_2\text{N}(\text{R}^\circ)_2$, $-\text{OC}(\text{O})\text{R}^\circ$, $-\text{N}(\text{R}^\circ)\text{C}(\text{O})\text{R}^\circ$, $-\text{N}(\text{R}^\circ)\text{N}(\text{R}^\circ)_2$, $-\text{C}=\text{NN}(\text{R}^\circ)_2$, $-\text{C}=\text{NOR}^\circ$, $-\text{N}(\text{R}^\circ)\text{C}(\text{O})\text{N}(\text{R}^\circ)_2$, $-\text{N}(\text{R}^\circ)\text{SO}_2\text{N}(\text{R}^\circ)_2$, $-\text{N}(\text{R}^\circ)\text{SO}_2\text{R}^\circ$, or $-\text{OC}(\text{O})\text{N}(\text{R}^\circ)_2$; wherein R° is as defined above and described herein. In other embodiments, Cy^2 is optionally substituted with C_{1-6} aliphatic or halogen. In some embodiments, Cy^2 is optionally substituted with Cl, F, CF_3 , or C_{1-4} alkyl. Exemplary substituents on Cy^2 include methyl, tert-butyl, and 1-methylcyclopropyl. In other embodiments, Cy^2 is mono- or di-substituted. In certain aspects, Cy^2 is optionally substituted at the *meta* or the *para* position with any one of the above-mentioned substituents. In some

embodiments, Cy^2 is substituted with R^o , wherein R^o is a 4–6-membered saturated, partially unsaturated, or aryl ring having 0–4 heteroatoms independently selected from nitrogen, oxygen, or sulfur,

[0097] Exemplary Cy^2 groups include those set forth in **Tables 2, 3, 4, and 5** in the Examples section, *infra*.

[0098] According to one aspect, the present invention provides a compound of formula **II**:



II


or a pharmaceutically acceptable salt thereof, wherein:

each of R^1 , R^x , and R^y is as defined above and described in classes and subclasses herein;

Cy^1 is an optionally substituted 5-membered saturated, partially unsaturated, or aromatic ring having 1–3 heteroatoms independently selected from nitrogen, oxygen, or sulfur; and

Cy^2 is optionally substituted phenyl or an optionally substituted 6-membered aromatic ring having 1–3 nitrogen atoms.

[0099] According to another aspect, the present invention provides a compound of formula **II'**:



II'



or a pharmaceutically acceptable salt thereof, wherein:

each of R^1 , R^x , and R^y is as defined above and described in classes and subclasses herein;

each of X, Y, and Z is independently -CH-, nitrogen, oxygen, or sulfur, wherein at least one of X, Y, or Z is a heteroatom and the circle depicted within the ring containing X, Y, and Z indicates that said ring is aromatic; and

Cy² is optionally substituted phenyl or an optionally substituted 6-membered aromatic ring having 1–3 nitrogen atoms.

[00100] Yet another aspect of the present invention provides a compound of formulae **II-a** and **II-b**:

**II-a****II-b**



or a pharmaceutically acceptable salt thereof, wherein:

each of R¹, R^x, and R^y is as defined above and described in classes and subclasses herein;

Cy¹ is an optionally substituted 5-membered saturated, partially unsaturated, or aromatic ring having 1–3 heteroatoms independently selected from nitrogen, oxygen, or sulfur; and

Cy² is optionally substituted phenyl or an optionally substituted 6-membered aromatic ring having 1–3 nitrogen atoms.

[00101] In certain embodiments, the present invention provides a compound of formulae **II-a'** and **II-b'**:

**II-a'****II-b'**

or a pharmaceutically acceptable salt thereof, wherein:


each of R^x, and R^y is as defined above and described in classes and subclasses herein;

Cy¹ is an optionally substituted 5-membered saturated, partially unsaturated, or aromatic ring having 1–3 heteroatoms independently selected from nitrogen, oxygen, or sulfur; and

Cy² is optionally substituted phenyl or an optionally substituted 6-membered aromatic ring having 1–3 nitrogen atoms.

[00102] In certain embodiments, the present invention provides a compound of formula **II-a** or **II-b** wherein Cy¹ is a 5-membered heteroaryl ring having 1-3 heteroatoms independently

selected from nitrogen, oxygen, or sulfur. Such compounds are represented by formulae **II-c** and **II-d**:



or a pharmaceutically acceptable salt thereof, wherein:


each of R^1 , R^x , and R^y is as defined above and described in classes and subclasses herein;

each of X, Y, and Z is independently -CH-, nitrogen, oxygen, or sulfur, wherein at least one of

X, Y, or Z is a heteroatom and the circle depicted within the ring containing X, Y, and Z indicates that said ring is aromatic; and

Cy^2 is optionally substituted phenyl or an optionally substituted 6-membered aromatic ring having 1–3 nitrogen atoms.

[00103] According to another embodiment, the present invention provides a method for preparing a compound of formula **II-a'**:



or a pharmaceutically acceptable salt thereof, wherein:


each of R^x and R^y is as defined above and described in classes and subclasses herein;

Cy^1 is an optionally substituted 5-membered saturated, partially unsaturated, or aromatic ring having 1–3 heteroatoms independently selected from nitrogen, oxygen, or sulfur; and

Cy^2 is optionally substituted phenyl or an optionally substituted 6-membered aromatic ring having 1–3 nitrogen atoms,

wherein said method comprises the steps depicted in Scheme **II**, below.

Scheme II



wherein each Cy^1 and Cy^2 is as defined above and described in classes and subclasses herein and A^- is a suitable chiral anion.

[0100] At step S-1, above, a compound of formula **II-i** is coupled to a compound of formula **II-ii**. Such coupling of a carboxylic acid group with an amine can be performed using methods well known to one of ordinary skill in the art. In certain embodiments, the carboxylic acid moiety of formula **II-i** is activated prior to coupling. In some embodiments, the carboxylic acid moiety is converted to an acyl halide group prior to coupling. In another embodiment, the carboxylic acid moiety is treated with a suitable reagent to form the acyl chloride thereof which is then coupled to the amine moiety of compound **II-ii** to form a compound of formula **II-iii**. Such reagents for forming acyl halides are well known to one of ordinary skill in the art and include oxalyl chloride and thionyl chloride, to name a few. In certain embodiments, the acyl halide of formula **II-iii** can be used directly in step S-2 without isolation or purification.

[00104] At step S-2, the ketone moiety of formula **II-iii** is converted to the oxime moiety of formula **II-iv**. In some embodiments, the compound of formula **II-iii** is treated with hydroxylamine to form a compound of formula **II-iv**. In certain embodiments, the compound of formula **II-iv** is about 1:1 E:Z configuration with respect to the -C=N- bond. In some embodiments, the present invention provides a compound of formula **II-iv** that is at least about

50%, 60%, 70%, 80%, 85%, 90%, 95%, 97%, 98%, 99%, or 100% in the E configuration with respect to the --C=N-- bond. In certain embodiments, the present invention provides a compound of formula **II-iv** that is at least about 50%, 60%, 70%, 80%, 85%, 90%, 95%, 97%, 98%, 99%, or 100% in the Z configuration with respect to the --C=N-- bond.

[00105] At step **S-3**, the oxime moiety of formula **II-iv** is converted to the amine group of formula **II-v**. In certain embodiments, the compound of formula **II-iv** is treated with zinc dust and acetic acid in an alcohol to form a compound of formula **II-v**. In certain embodiments, the alcohol is a C_{4-6} alkanol. In some embodiments, the alcohol is 1-butanol or pentanol.

[00106] At step **S-4 (a)**, the racemic compound **II-v** is treated with a chiral agent to form a diastereomeric salt of formula **II-vi-a**. In certain embodiments, the chiral acid has two carboxylate moieties as with, for example, tartaric acid or a derivative thereof. In some embodiments, the chiral acid is ditoluoyl tartaric acid. The term “chiral agent” means an enantiomerically enriched group which may be ionically or covalently bonded to the nitrogen of a compound of formula **II-v** to form **II-vi-a**. As used herein, the term “enantiomerically enriched”, as used herein means that one enantiomer makes up at least 85% of the preparation. In certain embodiments, the term enantiomerically enriched means that at least 90% of the preparation is one of the enantiomers. In other embodiments, the term means that at least 95% of the preparation is one of the enantiomers.

[00107] Chiral agents that are ionically bonded to said nitrogen include, for example, chiral acids. When the chiral agent is a chiral acid, the acid forms a diastereomeric salt with the nitrogen. The resulting diastereomers are then separated by suitable physical means. Examples of chiral acids include, but are not limited to, tartaric acid and tartaric acid derivatives, mandelic acid, malic acid, camphorsulfonic acid, and Mosher's acid, among others. In certain embodiments, the chiral acid is ditoluoyl-D-tartaric acid. In other embodiments, the chiral acid is ditoluoyl-L-tartaric acid. Other chiral agents that may be covalently bonded to the nitrogen are known in the art. Exemplary chiral acids include camphorsulfonic acid (-); tartaric acid (+); malic acid (-); N-acetyl-L-leucine (-); di-toluloyl-L-tartaric acid (-); deoxycholic acid (+); quinic acid (-); camphoric acid (+); N-BOC-alanine (-); tartaric acid (-); di-toluloyl-D-tartaric acid (+); camphorsulfonic acid (+); dibenzoyl-D-tartaric acid (+); L(+)citramalic; S-acetyl mandelic acid (+); and BOC-isoleucine(+).

[00108] At step S-4 (b), a diastereomeric salt of formula **II-vi-b** is obtained *via* suitable physical means. In some embodiments, “suitable physical means” refers to preferential crystallization, trituration, or slurry of a diastereomeric salt formed at step S-4 (a) above. In certain embodiments, a diastereomeric salt of formula **II-vi-b** is obtained *via* slurry. In other embodiments, the crystallization is achieved from a protic solvent. In still other embodiments, the protic solvent is an alcohol. It will be appreciated that the crystallization may be achieved using a single protic solvent or a combination of one or more protic solvents. Such solvents and solvent mixtures are well known to one of ordinary skill in the art and include, for example, one or more straight or branched alkyl alcohols. In certain embodiments, the crystallization is achieved from isopropyl alcohol and water.

[00109] At step S-4(a), a chiral acid is added to a compound of formula **II-v** to form a compound of formula **II-vi-a**. In certain embodiments, an equimolar amount of chiral acid is added. In other embodiments, a substoichiometric amount of chiral acid is added. In some embodiments, about 0.5 to about 0.75 molar equivalents of chiral acid are added. As used herein, the term “substoichiometric amount” denotes that the chiral acid is used in less than 1 mole equivalent relative to the compound of formula **II-v**.

[00110] In certain embodiments, the diastereomeric salt of formula **II-vi** comprises an equimolar amount of chiral acid and amine. In other embodiments, the diastereomeric salt of formula **II-vi** comprises a substoichiometric amount of chiral acid. In some embodiments, the diastereomeric salt of formula **II-vi** is a dihydrate.


[00111] It should be readily apparent to those skilled in the art that enantiomeric enrichment of one enantiomer in compound **II-vi-b** (ie resulting from preferential crystallization, trituration, or reslurry) causes an enantiomeric enrichment in the mother liquor of the other enantiomeric form. Therefore, according to another embodiment, the invention relates to a method of enhancing the percent enantiomeric excess (“%ee”) of a racemic compound of formula **II-vi-a** or enantiomerically enriched compound of formula **II-vi-b**.

[00112] At step S-5, the diastereomeric salt of formula **VI-vi-b** is treated with a suitable base obtain a compound of formula **II-vii**. Free bases according to the invention are also prepared, for example, by contacting a compound of formula **VI-vi-b** with a suitable base in the presence of a solvent suitable for free base formation. In certain embodiments, the suitable solvent is one or more polar aprotic solvent optionally mixed with a protic solvent. In some embodiments, the

suitable solvent is an ether mixed with an alcohol. In other embodiments, the suitable solvent is tert-butylmethyl ether and methanol or tert-butylmethyl ether and acetone. Such suitable bases include strong inorganic bases, *i.e.*, those that completely dissociate in water under formation of hydroxide anion. Exemplary suitable bases include metal hydroxides, including sodium hydroxide and potassium hydroxide. In some embodiments, the base is a carbonate base, *e.g.* sodium bicarbonate.

[00113] At step S-6, a compound of formula **II-vii** is coupled with a compound of formula **II-viii** to form a compound of formula **II-a**. Such coupling reactions are well known in the art. In certain embodiments, the coupling is achieved with a suitable coupling reagent. Such reagents are well known in the art and include, for example, DCC, HATU, and EDC, among others. In other embodiments, the carboxylic acid moiety is activated for use in the coupling reaction. Such activation includes formation of an acyl halide, use of a Mukaiyama reagent, and the like. These methods, and others, are known to one of ordinary skill in the art, *e.g.*, see, "Advanced Organic Chemistry," Jerry March, 5th Ed., pp. 351-357, John Wiley and Sons, N.Y.

[00114] According to another embodiment, the present invention provides a method for preparing a compound of formula **II-a'**:



II-a'


or a pharmaceutically acceptable salt thereof, wherein:

each of R^x and R^y is as defined above and described in classes and subclasses herein;

Cy¹ is an optionally substituted 5-membered saturated, partially unsaturated, or aromatic ring having 1–3 heteroatoms independently selected from nitrogen, oxygen, or sulfur; and


Cy² is optionally substituted phenyl or an optionally substituted 6-membered aromatic ring having 1–3 nitrogen atoms,

comprising the step of coupling a compound of formula **II-viii**:

**II-viii**

wherein each of R^x and R^y is as defined above and described in classes and subclasses herein;

with a compound of formula **II-vii**:

**II-vii**


wherein:

Cy^1 is an optionally substituted 5-membered saturated, partially unsaturated, or aromatic ring having 1–3 heteroatoms independently selected from nitrogen, oxygen, or sulfur; and

Cy^2 is optionally substituted phenyl or an optionally substituted 6-membered aromatic ring having 1–3 nitrogen atoms,

to form the compound of formula **II-a'**.

[00115] In certain embodiments, the compound of formula **II-vii**:


**II-vii**

wherein:

Cy^1 is an optionally substituted 5-membered saturated, partially unsaturated, or aromatic ring having 1–3 heteroatoms independently selected from nitrogen, oxygen, or sulfur; and


Cy^2 is optionally substituted phenyl or an optionally substituted 6-membered aromatic ring having 1–3 nitrogen atoms,

is prepared from a compound of formula **II-vi-b**:

**II-vi-b**

wherein A^- is a suitable chiral anion,
 comprising the step of treating the compound of formula **II-vi-b** with a suitable base to form a compound of formula **II-vii**.

[00116] In certain embodiments, the compound of formula **II-vi-b**:

**II-vi-b**


wherein:

A^- is a suitable chiral anion;

Cy^1 is an optionally substituted 5-membered saturated, partially unsaturated, or aromatic ring having 1–3 heteroatoms independently selected from nitrogen, oxygen, or sulfur;
 and


Cy^2 is optionally substituted phenyl or an optionally substituted 6-membered aromatic ring having 1–3 nitrogen atoms,

is prepared from a compound of formula **II-v**:

**II-v,**

comprising the steps of:


(a) treating the compound of formula **II-v** with a chiral agent to form a compound of formula **II-vi-a**:

**II-vi-a**

and

(b) separating the resulting diastereomers by suitable physical means to obtain a compound of formula **II-vi-b**.

[00117] In certain embodiments, the compound of formula **II-v**:


**II-v**

wherein:

Cy¹ is an optionally substituted 5-membered saturated, partially unsaturated, or aromatic ring having 1–3 heteroatoms independently selected from nitrogen, oxygen, or sulfur; and


Cy² is optionally substituted phenyl or an optionally substituted 6-membered aromatic ring having 1–3 nitrogen atoms,

is prepared from a compound of formula **II-iv**:

**II-iv**

comprising the step of converting the oxime moiety of formula **II-iv** to the amine group of formula **II-v**.

[00118] In some embodiments, the present invention provides a method for preparing a compound of formula **II-iv**:


**II-iv**

wherein:

Cy¹ is an optionally substituted 5-membered saturated, partially unsaturated, or aromatic ring having 1–3 heteroatoms independently selected from nitrogen, oxygen, or sulfur;
and


Cy² is optionally substituted phenyl or an optionally substituted 6-membered aromatic ring having 1–3 nitrogen atoms,

comprising the step of treating a compound of formula **II-iii**:



with hydroxylamine to form the compound of formula **II-iv**.

[00119] In certain embodiments, the compound of formula **II-iii**:




wherein:

Cy¹ is an optionally substituted 5-membered saturated, partially unsaturated, or aromatic ring having 1–3 heteroatoms independently selected from nitrogen, oxygen, or sulfur;
and


Cy² is optionally substituted phenyl or an optionally substituted 6-membered aromatic ring having 1–3 nitrogen atoms,

is prepared by coupling a compound of formula **II-i**:




wherein Cy¹ is an optionally substituted 5-membered saturated, partially unsaturated, or aromatic ring having 1–3 heteroatoms independently selected from nitrogen, oxygen, or sulfur,

with a compound of formula **II-ii**:

**II-ii**


wherein Cy^2 is optionally substituted phenyl or an optionally substituted 6-membered aromatic ring having 1–3 nitrogen atoms.

[00120] In certain embodiments, the present invention provides a compound of formula **II-vi-a** or **II-vi-b**:

**II-vi-a****II-vi-b**


wherein each of Cy^1 , Cy^2 and A^- is as defined herein.

[00121] In some embodiments, the present invention provides a compound of formula **II-iv**:

**II-iv**

wherein each of Cy^1 and Cy^2 is as defined herein.

[00122] According to another aspect, the present invention provides a compound of formula **III**:

**III**


or a pharmaceutically acceptable salt thereof, wherein:

each of R^1 , R^x , and R^y is as defined above and described in classes and subclasses herein;

Cy^1 is an optionally substituted 5-membered saturated, partially unsaturated, or aromatic ring having 1–3 heteroatoms independently selected from nitrogen, oxygen, or sulfur; and

Cy² is an optionally substituted 8–10 membered saturated, partially unsaturated, or aromatic bicyclic ring having 1–4 heteroatoms, independently selected from nitrogen, oxygen, or sulfur.

[00123] According to certain embodiments, the present invention provides a compound of formula **III'**:



III'


or a pharmaceutically acceptable salt thereof, wherein:

each of R¹, R^x, and R^y is as defined above and described in classes and subclasses herein;


each of X, Y, and Z is independently -CH-, nitrogen, oxygen, or sulfur, wherein at least one of X, Y, or Z is a heteroatom and the circle depicted within the ring containing X, Y, and Z indicates that said ring is aromatic; and

Cy² is optionally substituted phenyl or an optionally substituted 6-membered aromatic ring having 1–3 nitrogen atoms.

[00124] In certain aspects, the present invention provides a compound of formulae **III-a** and **III-b**:



III-a



III-b



or a pharmaceutically acceptable salt thereof, wherein:

each of R¹, R^x, and R^y is as defined above and described in classes and subclasses herein;

Cy¹ is an optionally substituted 5-membered saturated, partially unsaturated, or aromatic ring having 1–3 heteroatoms independently selected from nitrogen, oxygen, or sulfur; and

Cy² is an optionally substituted 8–10 membered saturated, partially unsaturated, or aromatic bicyclic ring having 1–4 heteroatoms, independently selected from nitrogen, oxygen, or sulfur.

[00125] In certain embodiments, the present invention provides a compound of formula **III-a** or **III-b** wherein Cy¹ is a 5-membered heteroaryl ring having 1-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur. Such compounds are represented by formulae **III-c** and **III-d**:

**III-c****III-d**

or a pharmaceutically acceptable salt thereof, wherein:


each of R¹, R^x, and R^y is as defined above and described in classes and subclasses herein;

each of X, Y, and Z is independently -CH-, nitrogen, oxygen, or sulfur, wherein at least one of X, Y, or Z is a heteroatom and the circle depicted within the ring containing X, Y, and Z indicates that said ring is aromatic; and

Cy² is optionally substituted phenyl or an optionally substituted 6-membered aromatic ring having 1–3 nitrogen atoms.

[00126] In certain embodiments, each of R¹, R^x, R^y, L¹, L², Cy¹, and Cy² is selected from those groups depicted in **Tables 1-5**, *infra*.

[00127] According to one aspect, the present invention provides a compound of formula **IV**:


**IV**

or a pharmaceutically acceptable salt thereof, wherein:

each of R¹, R^x, and R^y is as defined above and described in classes and subclasses herein;

Cy¹ is an optionally substituted 5-6 membered saturated, partially unsaturated, or aromatic ring having 1–3 heteroatoms independently selected from nitrogen, oxygen, or sulfur; and Cy² is optionally substituted phenyl or an optionally substituted 6-membered aromatic ring having 1–3 nitrogen atoms.

[00128] Yet another aspect of the present invention provides a compound of formulae **IV-a** and **IV-b**:



or a pharmaceutically acceptable salt thereof, wherein:

each of R¹, R^x, and R^y is as defined above and described in classes and subclasses herein;

Cy¹ is an optionally substituted 5-6 membered saturated, partially unsaturated, or aromatic ring having 1–3 heteroatoms independently selected from nitrogen, oxygen, or sulfur; and Cy² is optionally substituted phenyl or an optionally substituted 6-membered aromatic ring having 1–3 nitrogen atoms.

[00129] In certain embodiments, the present invention provides a compound of formula **IV**, **IV-a**, or **IV-b** wherein Cy¹ is a 5-membered heteroaryl ring having 1-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur.

[00130] Exemplary compounds of the present invention are set forth in the Examples at **Tables 3, 4, and 5, *infra***. In certain embodiments, the present invention provides a compound selected from those set forth in **Table 3**, or a pharmaceutically acceptable salt thereof. In some embodiments, the present invention provides a compound selected from those set forth in **Table 4**, or a pharmaceutically acceptable salt thereof. In other embodiments, the present invention provides a compound selected from those set forth in **Table 5**, or a pharmaceutically acceptable salt thereof.

4. Uses, Formulation and Administration

Pharmaceutically acceptable compositions

[0101] As discussed above, the present invention provides compounds that are inhibitors of protein kinases (*e.g.*, Raf kinase), and thus the present compounds are useful for the treatment of diseases, disorders, and conditions mediated by Raf kinase. In certain embodiments, the present invention provides a method for treating a Raf-mediated disorder. As used herein, the term “Raf-mediated disorder” includes diseases, disorders, and conditions mediated by Raf kinase. Such Raf-mediated disorders include melanoma, leukemia, or cancers such as colon, breast, gastric, ovarian, lung, brain, larynx, cervical, renal, lymphatic system, genitourinary tract (including bladder and prostate), stomach, bone, lymphoma, melanoma, glioma, papillary thyroid, neuroblastoma, and pancreatic cancer.

[0102] Raf-mediated disorders further include diseases afflicting mammals which are characterized by cellular proliferation. Such diseases include, for example, blood vessel proliferative disorders, fibrotic disorders, mesangial cell proliferative disorders, and metabolic diseases. Blood vessel proliferative disorders include, for example, arthritis and restenosis. Fibrotic disorders include, for example, hepatic cirrhosis and atherosclerosis. Mesangial cell proliferative disorders include, for example, glomerulonephritis, diabetic nephropathy, malignant nephrosclerosis, thrombotic microangiopathy syndromes, organ transplant rejection, and glomerulopathies. Metabolic disorders include, for example, psoriasis, diabetes mellitus, chronic wound healing, inflammation, and neurodegenerative diseases.

[0103] In another aspect of the present invention, pharmaceutically acceptable compositions are provided, wherein these compositions comprise any of the compounds as described herein, and optionally comprise a pharmaceutically acceptable carrier, adjuvant or vehicle. In certain embodiments, these compositions optionally further comprise one or more additional therapeutic agents.

[0104] It will also be appreciated that certain of the compounds of present invention can exist in free form for treatment, or where appropriate, as a pharmaceutically acceptable derivative thereof. According to the present invention, pharmaceutically acceptable derivatives include, but are not limited to, pharmaceutically acceptable salts, esters, salts of such esters, or any other adducts or derivatives that, upon administration to a patient in need, are capable of providing,

directly or indirectly, a compound as otherwise described herein, or a metabolite or residue thereof.

[0105] As used herein, the term "pharmaceutically acceptable salt" refers to those salts that are, within the scope of sound medical judgement, suitable for use in contact with the tissues of humans or animals without undue toxicity, irritation, allergic response, or the like, and are offer with a reasonable benefit/risk ratio. A "pharmaceutically acceptable salt" means any at least substantially non-toxic salt or salt of an ester of a compound of this invention that, upon administration to a recipient, is capable of providing, either directly or indirectly, a compound of this invention or an inhibitorily active metabolite or residue thereof. As used herein, the term "inhibitory metabolite or residue thereof" means that a metabolite or residue thereof is also an inhibitor of a Raf kinase.

[0106] Pharmaceutically acceptable salts are well known in the art. For example, S. M. Berge *et al.* describe pharmaceutically acceptable salts in detail in *J. Pharmaceutical Sciences*, **1977**, 66, 1–19, incorporated herein by reference. Pharmaceutically acceptable salts of the compounds of this invention include those derived from suitable inorganic and organic acids and bases. Examples of pharmaceutically acceptable, nontoxic acid addition salts are salts of an amino group formed with inorganic acids such as hydrochloric acid, hydrobromic acid, phosphoric acid, sulfuric acid and perchloric acid or with organic acids such as acetic acid, oxalic acid, maleic acid, tartaric acid, citric acid, succinic acid or malonic acid or by using other methods used in the art such as ion exchange. Other pharmaceutically acceptable salts include adipate, alginate, ascorbate, aspartate, benzenesulfonate, benzoate, bisulfate, borate, butyrate, camphorate, camphorsulfonate, citrate, cyclopentanepropionate, digluconate, dodecylsulfate, ethanesulfonate, formate, fumarate, glucoheptonate, glycerophosphate, gluconate, hemisulfate, heptanoate, hexanoate, hydroiodide, 2-hydroxy-ethanesulfonate, lactobionate, lactate, laurate, lauryl sulfate, malate, maleate, malonate, methanesulfonate, 2-naphthalenesulfonate, nicotinate, nitrate, oleate, oxalate, palmitate, pamoate, pectinate, persulfate, 3-phenylpropionate, phosphate, picrate, pivalate, propionate, stearate, succinate, sulfate, tartrate, thiocyanate, p-toluenesulfonate, undecanoate, valerate salts, and the like. Salts derived from appropriate bases include alkali metal, alkaline earth metal, ammonium and $N^+(C_{1-4}alkyl)_4$ salts. This invention also envisions the quaternization of any basic nitrogen-containing groups of the compounds disclosed herein. Water or oil-soluble or dispersible products may be obtained by such quaternization.

Representative alkali or alkaline earth metal salts include sodium, lithium, potassium, calcium, magnesium, and the like. Further pharmaceutically acceptable salts include, when appropriate, nontoxic ammonium, quaternary ammonium, and amine cations formed using counterions such as halide, hydroxide, carboxylate, sulfate, phosphate, nitrate, loweralkyl sulfonate and aryl sulfonate.

[0107] As described above, the pharmaceutically acceptable compositions of the present invention additionally comprise a pharmaceutically acceptable carrier, adjuvant, or vehicle, which, as used herein, includes any and all solvents, diluents, or other liquid vehicle, dispersion or suspension aids, surface active agents, isotonic agents, thickening or emulsifying agents, preservatives, solid binders, lubricants and the like, as suited to the particular dosage form desired. *Remington's Pharmaceutical Sciences*, Sixteenth Edition, E. W. Martin (Mack Publishing Co., Easton, Pa., 1980) discloses various carriers used in formulating pharmaceutically acceptable compositions and known techniques for the preparation thereof. Except insofar as any conventional carrier medium is incompatible with the compounds of the invention, such as by producing any undesirable biological effect or otherwise interacting in a deleterious manner with any other component(s) of the pharmaceutically acceptable composition, its use is contemplated to be within the scope of this invention. Some examples of materials which can serve as pharmaceutically acceptable carriers include, but are not limited to, ion exchangers, alumina, aluminum stearate, lecithin, serum proteins, such as human serum albumin, buffer substances such as phosphates, glycine, sorbic acid, or potassium sorbate, partial glyceride mixtures of saturated vegetable fatty acids, water, salts or electrolytes, such as protamine sulfate, disodium hydrogen phosphate, potassium hydrogen phosphate, sodium chloride, zinc salts, colloidal silica, magnesium trisilicate, polyvinyl pyrrolidone, polyacrylates, waxes, polyethylene–polyoxypropylene–block polymers, wool fat, sugars such as lactose, glucose and sucrose; starches such as corn starch and potato starch; cellulose and its derivatives such as sodium carboxymethyl cellulose, ethyl cellulose and cellulose acetate; powdered tragacanth; malt; gelatin; talc; excipients such as cocoa butter and suppository waxes; oils such as peanut oil, cottonseed oil; safflower oil; sesame oil; olive oil; corn oil and soybean oil; glycols; such a propylene glycol or polyethylene glycol; esters such as ethyl oleate and ethyl laurate; agar; buffering agents such as magnesium hydroxide and aluminum hydroxide; alginic acid; pyrogen-free water; isotonic saline; Ringer's solution; ethyl alcohol, and phosphate buffer

solutions, as well as other non-toxic compatible lubricants such as sodium lauryl sulfate and magnesium stearate, as well as coloring agents, releasing agents, coating agents, sweetening, flavoring and perfuming agents, preservatives and antioxidants can also be present in the composition, according to the judgment of the formulator.

Uses of Compounds and Pharmaceutically acceptable compositions

[0108] According to the present invention, provided compounds may be assayed in any of the available assays known in the art for identifying compounds having kinase inhibitory activity. For example, the assay may be cellular or non-cellular, *in vivo* or *in vitro*, high- or low-throughput format, etc.

[0109] In certain exemplary embodiments, compounds of this invention were assayed for their ability to inhibit protein kinases, more specifically Raf.

[0110] Thus, in one aspect, compounds of this invention which are of particular interest include those which:

- are inhibitors of protein kinases;
- exhibit the ability to inhibit Raf kinase;
- are useful for treating mammals (*e.g.*, humans) or animals suffering from an Raf-mediated disease or condition, and for helping to prevent or delay the onset of such a disease or condition;
- exhibit a favorable therapeutic profile (*e.g.*, safety, efficacy, and stability).

[0111] In certain embodiments, compounds of the invention are Raf kinase inhibitors. In certain exemplary embodiments, compounds of the invention are Raf inhibitors. In certain exemplary embodiments, compounds of the invention have $^{Cell}IC_{50}$ values ≤ 100 μM . In certain other embodiments, compounds of the invention have $^{Cell}IC_{50}$ values ≤ 75 μM . In certain other embodiments, compounds of the invention have $^{Cell}IC_{50}$ values ≤ 50 μM . In certain other embodiments, compounds of the invention have $^{Cell}IC_{50}$ values ≤ 25 μM . In certain other embodiments, compounds of the invention have $^{Cell}IC_{50}$ values ≤ 10 μM . In certain other embodiments, compounds of the invention have $^{Cell}IC_{50}$ values ≤ 7.5 μM . In certain other embodiments, of the invention compounds have $^{Cell}IC_{50}$ values ≤ 5 μM . In certain other embodiments, of the invention compounds have $^{Cell}IC_{50}$ values ≤ 2.5 μM . In certain other embodiments, of the invention compounds have $^{Cell}IC_{50}$ values ≤ 1 μM . In certain other embodiments, of the invention compounds have $^{Cell}IC_{50}$ values ≤ 800 nM. In certain other

embodiments, of the invention compounds have $^{Cell}IC_{50}$ values ≤ 600 nM. In certain other embodiments, inventive compounds have $^{Cell}IC_{50}$ values ≤ 500 nM. In certain other embodiments, compounds of the invention have $^{Cell}IC_{50}$ values ≤ 300 nM. In certain other embodiments, compounds of the invention have $^{Cell}IC_{50}$ values ≤ 200 nM. In certain other embodiments, of the invention compounds have $^{Cell}IC_{50}$ values ≤ 100 nM.

[0112] In yet another aspect, a method for the treatment or lessening the severity of an Raf-mediated disease or condition is provided comprising administering an effective amount of a compound, or a pharmaceutically acceptable composition comprising a compound to a subject in need thereof. In certain embodiments of the present invention an "effective amount" of the compound or pharmaceutically acceptable composition is that amount effective for treating or lessening the severity of a Raf-mediated disease or condition. The compounds and compositions, according to the method of the present invention, may be administered using any amount and any route of administration effective for treating or lessening the severity of a Raf-mediated disease or condition. The exact amount required will vary from subject to subject, depending on the species, age, and general condition of the subject, the severity of the infection, the particular agent, its mode of administration, and the like. In certain embodiments, compounds of the invention are formulated in dosage unit form for ease of administration and uniformity of dosage. The expression "dosage unit form" as used herein refers to a physically discrete unit of agent appropriate for the patient to be treated. It will be understood, however, that the total daily usage of the compounds and compositions of the present invention will be decided by the attending physician within the scope of sound medical judgment. The specific effective dose level for any particular patient or organism will depend upon a variety of factors including the disorder being treated and the severity of the disorder; the activity of the specific compound employed; the specific composition employed; the age, body weight, general health, sex and diet of the patient; the time of administration, route of administration, and rate of excretion of the specific compound employed; the duration of the treatment; drugs used in combination or coincidental with the specific compound employed, and like factors well known in the medical arts. The term "patient", as used herein, means an animal, preferably a mammal, and most preferably a human.

[0113] The pharmaceutically acceptable compositions of this invention can be administered to humans and other animals orally, rectally, parenterally, intracisternally, intravaginally,

intraperitoneally, topically (as by powders, ointments, or drops), buccally, as an oral or nasal spray, or the like, depending on the severity of the infection being treated. In certain embodiments, the compounds of the invention may be administered orally or parenterally at dosage levels of about 0.01 mg/kg to about 50 mg/kg and preferably from about 1 mg/kg to about 25 mg/kg, of subject body weight per day, one or more times a day, to obtain the desired therapeutic effect.

[0114] Liquid dosage forms for oral administration include, but are not limited to, pharmaceutically acceptable emulsions, microemulsions, solutions, suspensions, syrups and elixirs. In addition to the active compounds, the liquid dosage forms may contain inert diluents commonly used in the art such as, for example, water or other solvents, solubilizing agents and emulsifiers such as ethyl alcohol, isopropyl alcohol, ethyl carbonate, ethyl acetate, benzyl alcohol, benzyl benzoate, propylene glycol, 1,3-buthylene glycol, dimethylformamide, oils (in particular, cottonseed, groundnut, corn, germ, olive, castor, and sesame oils), glycerol, tetrahydrofurfuryl alcohol, polyethylene glycols and fatty acid esters of sorbitan, and mixtures thereof. Besides inert diluents, the oral compositions can also include adjuvants such as wetting agents, emulsifying and suspending agents, sweetening, flavoring, and perfuming agents.

[0115] Injectable preparations, for example, sterile injectable aqueous or oleaginous suspensions may be formulated according to the known art using suitable dispersing or wetting agents and suspending agents. The sterile injectable preparation may also be a sterile injectable solution, suspension or emulsion in a nontoxic parenterally acceptable diluent or solvent, for example, as a solution in 1,3-butanediol. Among the acceptable vehicles and solvents that may be employed are water, Ringer's solution, U.S.P. and isotonic sodium chloride solution. In addition, sterile, fixed oils are conventionally employed as a solvent or suspending medium. For this purpose any bland fixed oil can be employed including synthetic mono- or diglycerides. In addition, fatty acids such as oleic acid are used in the preparation of injectables.

[0116] The injectable formulations can be sterilized, for example, by filtration through a bacterial-retaining filter, or by incorporating sterilizing agents in the form of sterile solid compositions which can be dissolved or dispersed in sterile water or other sterile injectable medium prior to use.

[0117] In order to prolong the effect of a compound of the present invention, it is often desirable to slow the absorption of the compound from subcutaneous or intramuscular injection.

This may be accomplished by the use of a liquid suspension of crystalline or amorphous material with poor water solubility. The rate of absorption of the compound then depends upon its rate of dissolution that, in turn, may depend upon crystal size and crystalline form. Alternatively, delayed absorption of a parenterally administered compound form is accomplished by dissolving or suspending the compound in an oil vehicle. Injectable depot forms are made by forming microencapsule matrices of the compound in biodegradable polymers such as polylactide–polyglycolide. Depending upon the ratio of compound to polymer and the nature of the particular polymer employed, the rate of compound release can be controlled. Examples of other biodegradable polymers include poly(orthoesters) and poly(anhydrides). Depot injectable formulations are also prepared by entrapping the compound in liposomes or microemulsions that are compatible with body tissues.

[0118] Compositions for rectal or vaginal administration are preferably suppositories which can be prepared by mixing the compounds of this invention with suitable non-irritating excipients or carriers such as cocoa butter, polyethylene glycol or a suppository wax which are solid at ambient temperature but liquid at body temperature and therefore melt in the rectum or vaginal cavity and release the active compound.

[0119] Solid dosage forms for oral administration include capsules, tablets, pills, powders, and granules. In such solid dosage forms, the active compound is mixed with at least one inert, pharmaceutically acceptable excipient or carrier such as sodium citrate or dicalcium phosphate and/or a) fillers or extenders such as starches, lactose, sucrose, glucose, mannitol, and silicic acid, b) binders such as, for example, carboxymethylcellulose, alginates, gelatin, polyvinylpyrrolidone, sucrose, and acacia, c) humectants such as glycerol, d) disintegrating agents such as agar—agar, calcium carbonate, potato or tapioca starch, alginic acid, certain silicates, and sodium carbonate, e) solution retarding agents such as paraffin, f) absorption accelerators such as quaternary ammonium compounds, g) wetting agents such as, for example, cetyl alcohol and glycerol monostearate, h) absorbents such as kaolin and bentonite clay, and i) lubricants such as talc, calcium stearate, magnesium stearate, solid polyethylene glycols, sodium lauryl sulfate, and mixtures thereof. In the case of capsules, tablets and pills, the dosage form may also comprise buffering agents.

[0120] Solid compositions of a similar type may also be employed as fillers in soft and hard-filled gelatin capsules using such excipients as lactose or milk sugar as well as high molecular

weight polyethylene glycols and the like. The solid dosage forms of tablets, dragees, capsules, pills, and granules can be prepared with coatings and shells such as enteric coatings and other coatings well known in the pharmaceutical formulating art. They may optionally contain opacifying agents and can also be of a composition that they release the active ingredient(s) only, or preferentially, in a certain part of the intestinal tract, optionally, in a delayed manner. Examples of embedding compositions that can be used include polymeric substances and waxes. Solid compositions of a similar type may also be employed as fillers in soft and hard-filled gelatin capsules using such excipients as lactose or milk sugar as well as high molecular weight polyethylene glycols and the like.

[0121] The active compounds can also be in micro-encapsulated form with one or more excipients as noted above. The solid dosage forms of tablets, dragees, capsules, pills, and granules can be prepared with coatings and shells such as enteric coatings, release controlling coatings and other coatings well known in the pharmaceutical formulating art. In such solid dosage forms the active compound may be admixed with at least one inert diluent such as sucrose, lactose or starch. Such dosage forms may also comprise, as is normal practice, additional substances other than inert diluents, *e.g.*, tableting lubricants and other tableting aids such as a magnesium stearate and microcrystalline cellulose. In the case of capsules, tablets and pills, the dosage forms may also comprise buffering agents. They may optionally contain opacifying agents and can also be of a composition that they release the active ingredient(s) only, or preferentially, in a certain part of the intestinal tract, optionally, in a delayed manner. Examples of embedding compositions that can be used include polymeric substances and waxes.

[0122] Dosage forms for topical or transdermal administration of a compound of this invention include ointments, pastes, creams, lotions, gels, powders, solutions, sprays, inhalants or patches. The active component is admixed under sterile conditions with a pharmaceutically acceptable carrier and any needed preservatives or buffers as may be required. Ophthalmic formulation, ear drops, and eye drops are also contemplated as being within the scope of this invention. Additionally, the present invention contemplates the use of transdermal patches, which have the added advantage of providing controlled delivery of a compound to the body. Such dosage forms can be made by dissolving or dispensing the compound in the proper medium. Absorption enhancers can also be used to increase the flux of the compound across the

skin. The rate can be controlled by either providing a rate controlling membrane or by dispersing the compound in a polymer matrix or gel.

[0123] As described generally above, the compounds of the invention are useful as inhibitors of protein kinases. In one embodiment, the compounds of the invention are Raf kinase inhibitors, and thus, without wishing to be bound by any particular theory, the compounds and compositions are particularly useful for treating or lessening the severity of a disease, condition, or disorder where activation of Raf kinase is implicated in the disease, condition, or disorder. When activation of Raf kinase is implicated in a particular disease, condition, or disorder, the disease, condition, or disorder may also be referred to as a “Raf-mediated disease”. Accordingly, in another aspect, the present invention provides a method for treating or lessening the severity of a disease, condition, or disorder where activation of Raf kinase is implicated in the disease state.

[0124] The activity of a compound utilized in this invention as an Raf kinase inhibitor, may be assayed *in vitro*, *in vivo*, *ex vivo*, or in a cell line. *In vitro* assays include assays that determine inhibition of either the phosphorylation activity or ATPase activity of activated Raf. Alternate *in vitro* assays quantitate the ability of the inhibitor to bind to Raf. Inhibitor binding may be measured by radiolabelling the inhibitor (e.g., synthesizing the inhibitor to include a radioisotope) prior to binding, isolating the inhibitor/Raf, complex and determining the amount of radiolabel bound. Alternatively, inhibitor binding may be determined by running a competition experiment where new inhibitors are incubated with Raf bound to known radioligands.

[0125] The term “measurably inhibit”, as used herein means a measurable change in Raf activity between a sample comprising said composition and a Raf kinase and an equivalent sample comprising Raf kinase in the absence of said composition.

[0126] It will also be appreciated that the compounds and pharmaceutically acceptable compositions of the present invention can be employed in combination therapies, that is, the compounds and pharmaceutically acceptable compositions can be administered concurrently with, prior to, or subsequent to, one or more other desired therapeutics or medical procedures. The particular combination of therapies (therapeutics or procedures) to employ in a combination regimen will take into account compatibility of the desired therapeutics and/or procedures and the desired therapeutic effect to be achieved. It will also be appreciated that the therapies

employed may achieve a desired effect for the same disorder (for example, compound of the invention may be administered concurrently with another agent used to treat the same disorder), or they may achieve different effects (e.g., control of any adverse effects). As used herein, additional therapeutic agents that are normally administered to treat or prevent a particular disease, or condition, are known as "appropriate for the disease, or condition, being treated".

[0127] For example, other therapies, chemotherapeutic agents, or other anti-proliferative agents may be combined with the compounds of this invention to treat proliferative diseases and cancer. Examples of therapies or anticancer agents that may be used in combination with the inventive anticancer agents of the present invention include surgery, radiotherapy (e.g., gamma-radiation, neutron beam radiotherapy, electron beam radiotherapy, proton therapy, brachytherapy, and systemic radioactive isotopes), endocrine therapy, biologic response modifiers (e.g., interferons, interleukins, and tumor necrosis factor (TNF)), hyperthermia and cryotherapy, agents to attenuate any adverse effects (e.g., antiemetics), and other approved chemotherapeutic drugs.

Examples of chemotherapeutic anticancer agents that may be used as second active agents in combination with compounds of the invention include, but are not limited to, alkylating agents (e.g. mechlorethamine, chlorambucil, cyclophosphamide, melphalan, ifosfamide), antimetabolites (e.g., methotrexate), purine antagonists and pyrimidine antagonists (e.g. 6-mercaptopurine, 5-fluorouracil, cytarabine, gemcitabine), spindle poisons (e.g., vinblastine, vincristine, vinorelbine, paclitaxel), podophyllotoxins (e.g., etoposide, irinotecan, topotecan), antibiotics (e.g., doxorubicin, daunorubicin, bleomycin, mitomycin), nitrosoureas (e.g., carmustine, lomustine), inorganic ions (e.g., platinum complexes such as cisplatin, carboplatin), enzymes (e.g., asparaginase), hormones (e.g., tamoxifen, leuprolide, flutamide, and megestrol), topoisomerase II inhibitors or poisons, EGFR (Her1, ErbB-1) inhibitors (e.g., gefitinib), antibodies (e.g., rituximab), IMiDs (e.g., thalidomide, lenalidomide), various targeted agents (e.g., HDAC inhibitors such as vorinostat, Bcl-2 inhibitors, VEGF inhibitors); proteasome inhibitors (e.g., bortezomib), cyclin-dependent kinase inhibitors, and dexamethasone.

[0128] For a more comprehensive discussion of updated cancer therapies see, *The Merck Manual*, Seventeenth Ed. 1999, the entire contents of which are hereby incorporated by reference. See also the National Cancer Institute (NCI) website (www.nci.nih.gov) and the Food

and Drug Administration (FDA) website for a list of the FDA approved oncology drugs (www.fda.gov/cder/cancer/druglistframe – See Appendix).

[0129] Other examples of agents the inhibitors of this invention may also be combined with include, without limitation: treatments for Alzheimer's Disease such as Aricept[®] and Exelon[®]; treatments for Parkinson's Disease such as L-DOPA/carbidopa, entacapone, ropinrole, pramipexole, bromocriptine, pergolide, trihexephendyl, and amantadine; agents for treating Multiple Sclerosis (MS) such as beta interferon (*e.g.*, Avonex[®] and Rebif[®]), Copaxone[®], and mitoxantrone; treatments for asthma such as albuterol and Singulair[®]; agents for treating schizophrenia such as zyprexa, risperdal, seroquel, and haloperidol; anti-inflammatory agents such as corticosteroids, TNF blockers, IL-1 RA, azathioprine, cyclophosphamide, and sulfasalazine; immunomodulatory agents, including immunosuppressive agents, such as cyclosporin, tacrolimus, rapamycin, mycophenolate mofetil, interferons, corticosteroids, cyclophosphamide, azathioprine, and sulfasalazine; neurotrophic factors such as acetylcholinesterase inhibitors, MAO inhibitors, interferons, anti-convulsants, ion channel blockers, riluzole, and anti-Parkinson's agents; agents for treating cardiovascular disease such as beta-blockers, ACE inhibitors, diuretics, nitrates, calcium channel blockers, and statins; agents for treating liver disease such as corticosteroids, cholestyramine, interferons, and anti-viral agents; agents for treating blood disorders such as corticosteroids, anti-leukemic agents, and growth factors; and agents for treating immunodeficiency disorders such as gamma globulin.

[0130] Those additional agents may be administered separately from composition containing a compound of the invention, as part of a multiple dosage regimen. Alternatively, those agents may be part of a single dosage form, mixed together with a compound of this invention in a single composition. If administered as part of a multiple dosage regime, the two active agents may be submitted simultaneously, sequentially or within a period of time from one another normally within five hours from one another.

[0131] The amount of additional therapeutic agent present in the compositions of this invention will be no more than the amount that would normally be administered in a composition comprising that therapeutic agent as the only active agent. Preferably the amount of additional therapeutic agent in the presently disclosed compositions will range from about 50% to 100% of the amount normally present in a composition comprising that agent as the only therapeutically active agent.

[0132] The compounds of this invention or pharmaceutically acceptable compositions thereof may also be incorporated into compositions for coating implantable medical devices, such as prostheses, artificial valves, vascular grafts, stents and catheters. Accordingly, the present invention, in another aspect, includes a composition for coating an implantable device comprising a compound of the present invention as described generally above, and in classes and subclasses herein, and a carrier suitable for coating said implantable device. In still another aspect, the present invention includes an implantable device coated with a composition comprising a compound of the present invention as described generally above, and in classes and subclasses herein, and a carrier suitable for coating said implantable device.

[0133] Vascular stents, for example, have been used to overcome restenosis (re-narrowing of the vessel wall after injury). However, patients using stents or other implantable devices risk clot formation or platelet activation. These unwanted effects may be prevented or mitigated by pre-coating the device with a pharmaceutically acceptable composition comprising a kinase inhibitor. Suitable coatings and the general preparation of coated implantable devices are described in US Patents 6,099,562; 5,886,026; and 5,304,121. The coatings are typically biocompatible polymeric materials such as a hydrogel polymer, polymethyldisiloxane, polycaprolactone, polyethylene glycol, polylactic acid, ethylene vinyl acetate, and mixtures thereof. The coatings may optionally be further covered by a suitable topcoat of fluorosilicone, polysaccharides, polyethylene glycol, phospholipids or combinations thereof to impart controlled release characteristics in the composition.

[0134] Another aspect of the invention relates to inhibiting Raf activity in a biological sample or a patient, which method comprises administering to the patient, or contacting said biological sample with a compound of the present invention or a composition comprising said compound. The term "biological sample", as used herein, includes, without limitation, cell cultures or extracts thereof; biopsied material obtained from a mammal or extracts thereof; and blood, saliva, urine, feces, semen, tears, or other body fluids or extracts thereof.

[0135] Inhibition of Raf kinase activity in a biological sample is useful for a variety of purposes that are known to one of skill in the art. Examples of such purposes include, but are not limited to, blood transfusion, organ-transplantation, biological specimen storage, and biological assays.

TREATMENT KIT

[0136] In other embodiments, the present invention relates to a kit for conveniently and effectively carrying out the methods in accordance with the present invention. In general, the pharmaceutical pack or kit comprises one or more containers filled with one or more of the ingredients of the pharmaceutical compositions of the invention. Such kits are especially suited for the delivery of solid oral forms such as tablets or capsules. Such a kit preferably includes a number of unit dosages, and may also include a card having the dosages oriented in the order of their intended use. If desired, a memory aid can be provided, for example in the form of numbers, letters, or other markings or with a calendar insert, designating the days in the treatment schedule in which the dosages can be administered. Alternatively, placebo dosages, or calcium dietary supplements, either in a form similar to or distinct from the dosages of the pharmaceutical compositions, can be included to provide a kit in which a dosage is taken every day. Optionally associated with such container(s) can be a notice in the form prescribed by a governmental agency regulating the manufacture, use or sale of pharmaceutical products, which notice reflects approval by the agency of manufacture, use or sale for human administration.

EQUIVALENTS

[0137] The representative examples that follow are intended to help illustrate the invention, and are not intended to, nor should they be construed to, limit the scope of the invention. Indeed, various modifications of the invention and many further embodiments thereof, in addition to those shown and described herein, will become apparent to those skilled in the art from the full contents of this document, including the examples which follow and the references to the scientific and patent literature cited herein. It should further be appreciated that the contents of those cited references are incorporated herein by reference to help illustrate the state of the art.

[0138] The following examples contain important additional information, exemplification and guidance that can be adapted to the practice of this invention in its various embodiments and the equivalents thereof.


EXAMPLES

[0139] As depicted in the Examples below, in certain exemplary embodiments, compounds are prepared according to the following general procedures. It will be appreciated that, although the synthetic methods and Schemes depict the synthesis of certain compounds of the present

invention, the following methods and other methods known to one of ordinary skill in the art can be applied to all compounds and subclasses and species of each of these compounds, as described herein.

Synthesis of the Pyrimidine (“Left-Side”) Groups

Scheme 1.



[0140] Synthesis of Compound 1.1. To a stirred solution of diethyl acetylenedicarboxylate (20 g, 0.117 mol) and formamidine hydrochloride (9.4 g, 0.117 mol) in acetonitrile (400 mL) was added triethylamine (16.3 mL, 0.117 mol) dropwise at room temperature (RT) and the reaction mixture was heated at reflux for 16 hours (hr). The reaction mixture was cooled to 0 °C and the obtained solid was filtered and purified by silica gel column chromatography to furnish compound **1.1** (11g, 55.6%). ¹H NMR (200 MHz, DMSO-*d*₆): δ 12.7 (bs, 1H), 8.25 (s, 1H), 6.85 (s, 1H), 4.28 (q, *J* = 7 Hz, 2H), 1.25 (t, *J* = 7 Hz, 3H); LCMS: *m/z* 169 [*M*+1]⁺.


[0141] Synthesis of Compound 1.2. To a stirred solution of compound **1.1** (8 g, 0.047 mol) in DMF (22 mL) was added 1,3-dichloro-5,5-dimethyl hydantoin (NDDH; 5.6 g, 0.028 mol) in DMF (14.7 mL) and the reaction mixture stirred at RT for 1 hr. After complete consumption of the starting material was observed by TLC analysis, the reaction mixture was cooled to 0 °C and SOCl₂ (5.3 mL, 0.062) was added dropwise. After warming to RT and stirring for 1 hr, the reaction mixture was diluted with water (120 mL) and extracted with ether (3 x 200 mL). The combined organic layers were dried (Na₂SO₄), concentrated under reduced pressure and purified by column chromatography to give compound **1.2** (4.6g, 40.7%). ¹H NMR (200 MHz, DMSO-*d*₆): δ 9.15 (s, 1H), 4.42 (q, *J* = 7 Hz, 2H), 1.41 (t, *J* = 7 Hz, 3H). ¹³C NMR (125 MHz, DMSO-



d₆): 162.293, 159.731, 156.087, 155.993, 126.329, 62.962 and 13.803. LCMS: *m/z*: *m/z* 221 [M+1]⁺.

[0142] Synthesis of Compound 1.3. To a stirred solution of compound **1.2** (500 mg, 0.0022 mol) in 1,4-dioxane (5 mL) was added ethanolamine (152 mg, 0.0024 mol) and the reaction mixture was stirred at RT overnight. The progress of the reaction was monitored by TLC. After consumption of the starting material, the reaction mixture was concentrated under reduced pressure and purified by column chromatography (5% MeOH/ DCM) to give compound **1.3** (220 mg, 40%). ¹H NMR (200 MHz, DMSO-d₆): δ 8.40 (s, 1H), 7.70 (bs, N-H), 4.78 (bs, O-H), 4.28 (q, *J* = 7.4 Hz, 2H), 3.58–3.42 (m, 4H), 1.25 (t, *J* = 7.4 Hz, 3H); LCMS: *m/z* 246 [M+1]⁺.

[0143] Synthesis of Compound 1a. To a solution of ester **1.3** in THF (10 equiv.) and water (30 equiv.) was added LiOH (2.0 equiv.). The reaction mixture was stirred at RT for 1–3 hr and monitored by LCMS. THF was removed under reduced pressure and the resulting aqueous solution was neutralized with 2 N HCl. Precipitates were collected and dried to give the corresponding acid. In cases where precipitation did not occur, the mixture was lyophilized to give a crude product which was used for coupling without further purification.

[0144] Compounds 1a–1t. Using different amines and compound **1.2**, the following acids can be synthesized by the general method depicted in Scheme 1:




**Scheme 2.**


[0145] Synthesis of Compound 2.1. A mixture of compound 1.2 (250 mg, 0.0011 mol) and 4-amino pyridine (106 mg, 0.0011 mol) in acetonitrile (2.5 mL) was stirred in a sealed tube at 95 °C for 3 hr. After the reaction was judged to be complete by TLC analysis, the reaction mixture was cooled to 0 °C. The obtained solid was filtered and purified by column chromatography (50% ethyl acetate / hexane) to give compound 2.1 (100 mg, 33%). ¹H NMR (500 MHz, CDCl₃): δ 8.78 (s, 1H), 8.57 (d, *J* = 7.0 Hz, 2H), 7.70 (d, *J* = 6.0 Hz, 2H), 7.60 (bs, N-H), 4.50 (q, *J* = 7.0 Hz, 4H), 1.43 (t, *J* = 7.0 Hz, 3H); LCMS: *m/z* 279 [M+1]⁺.

[0146] Synthesis of Compound 2a. Compound 2.1 was hydrolyzed as described for compound 1 to afford 2 which was used without further purification. ¹H NMR (500 MHz, DMSO-*d*₆): δ 10.50 (bs, 1H), 8.88–8.36 (m, 5H). LCMS: 251 [M+1]⁺.

[0147] Compounds 2a–2g. Using different anilines and compound 1.2, the following acids can be synthesized as by the general method depicted in Scheme 2:




Scheme 3.




[0148] **Synthesis of Compound 3.1.** A solution of compound **1.2** (250 mg, 0.00113 mol) in MeOH (5 mL) in a sealed tube was stirred at 60 °C overnight. After consumption of the starting material, MeOH was removed under reduced pressure. The obtained crude material was purified by column chromatography (30% ethyl acetate / hexanes) to give compound **3.1** (78 mg, 31%). ¹H NMR (200 MHz, CD₃OD): δ 8.69 (s, 1H), 4.51 (s, 3H), 3.99 (s, 3H); LCMS: *m/z* 203 [M+1]⁺.

[0149] **Synthesis of Compound 3a.** Compound **3.1** was hydrolyzed as described for compound **1** to afford **3a** as a crude product which was used without further purification. ¹H NMR (200 MHz, DMSO-*d*₆): δ 8.58 (s, 1H), 3.98 (s, 3H); LCMS: 188 [M+1]⁺.

[0150] **Compounds 3a–3c.** Using different alcohols and compound **1.2**, the following acids can be synthesized as exemplified in Scheme 3:
























Scheme 4.

[0151] Synthesis of Compound 4.1 was synthesized using the approach shown in Scheme 1, except omitting the chlorination step using NDDH.


[0152] Synthesis of Compound 4a. To a microwave vial was added 6-chloro-pyrimidine-4-carboxylic acid ethyl ester (250 mg, 0.0013 mol), 4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-pyridine (275 mg, 0.00134 mol), 1,2-dimethoxyethane (5.0 mL, 0.048 mol), saturated sodium bicarbonate solution (0.9 mL, 0.009 mol), and tetrakis(triphenylphosphine)palladium(0) (150 mg, 0.00013 mol). The vial was purged with nitrogen, and sealed with a rubber septum. The reaction mixture was heated in the microwave (300 watts, 110°C) for 2 hr. LCMS indicated consumption of starting material. The major product was found to be the hydrolysis product [by LCMS ($M+1 = 202$)]. The reaction mixture was diluted with 50% MeOH/ CH_2Cl_2 (20 mL), and filtered through Celite[®]. The filtrate was concentrated under reduced pressure and the resultant residue was triturated with water (3 x 20 mL). The aqueous mixture was collected and washed with ethyl acetate EtOAc (3 x 10 mL) to remove residual ligand. The solution was neutralized with 1N HCl and lyophilized to give acid **4** as a light purple powder (160 mg, 50%) which was used without further purification. LCMS: m/z 202 [$M+1$]⁺.

[0153] Compounds 4a–4z. Using different boronic acids or esters, and compound **1.4**, the following acids can be synthesized by the general method depicted in Scheme 4:








Scheme 5.
















[0154] Compounds **5a–5eee**. Using different amines and compound **4.1**, the following acids can be synthesized as exemplified in Scheme 5:













Scheme 6.



[0155] Compounds **6a–6q**. Using different anilines and compound **4.1**, the following acids can be synthesized as exemplified in Scheme 6:




**Scheme 7.**

[0156] Synthesis of Compound 7.1. A solution of THP-protected homopropargyl alcohol (500 mg, 0.00324 mol) and triethylamine (0.4 mL, 0.00324 mol) in DMF (5 mL) was degassed for 30 min. Compound **4.1** (600 mg, 0.00324 mol), Pd(PPh₃)₄ (260 mg, 0.0002 mol) and CuI (20 mg) were added and the reaction mixture was stirred at RT for 16 hr. The reaction mixture was diluted with water (100 mL) and extracted with EtOAc (3 x 50 mL). The combined organic layers were washed with cold water (100 mL), dried over Na₂SO₄, concentrated under reduced pressure and purified by column chromatography to give **7.1** (350 mg, 42%). ¹H NMR (200 MHz, CDCl₃): δ 9.30 (s, 1H), 8.00 (s, 1H), 4.70 (t, *J* = 2.2 Hz, 1H), 4.52 (q, *J* = 7.2 Hz, 2H), 4.02–3.75 (m, 2H), 3.75–3.50 (m, 2H), 2.82 (t, *J* = 6.8 Hz, 2H), 1.82–1.41 (m, 4H); LCMS: *m/z* 304 [M+1]⁺.

[0157] Synthesis of Compound 7.2. Compound **7.1** was hydrolyzed as described for compound **1** to afford **7.2** which was used without further purification. ¹H NMR (200 MHz, DMSO-*d*₆): δ 9.07 (s, 1H), 7.67 (s, 1H), 4.66 (s, 1H), 3.79–3.56 (m, 4H). LCMS: *m/z* 276 [M+1]⁺.

Scheme 8.




[0158] Synthesis of Compound 8.1. A solution of 4-pentyn-1-ol (573 mg, 0.0068 mol) and triethylamine (689 mg, 0.0068 mol) in DMF (5 mL) was degassed for 30 min. Compound **1.2** (1g, 0.0045 mol), Pd(PPh₃)₄ (367 mg, 0.0003 mol), and CuI (50 mg) were added and the reaction mixture was stirred for 20 hr. After consumption of the starting material, the reaction mixture was diluted with water (100 mL) and extracted with EtOAc (3x 50 mL). The combined organic layers were washed with cold water (100 mL), dried over Na₂SO₄, concentrated under reduced pressure and purified by column chromatography (20% ethyl acetate/ hexane) to give compound **8.1** (848 mg, 69%). ¹H NMR (500 MHz, DMSO-d₆): δ 9.18 (s, 1H), 4.60 (t, *J* = 5.5 Hz, O-H), 4.43 (q, *J* = 7.5 Hz, 2H), 3.53 (t, *J* = 6.5 Hz, 2H), 2.65 (t, *J* = 6.5 Hz, 2H), 1.76–1.71 (m, 2H), 1.32 (t, *J* = 7.5 Hz, 3H); LCMS: *m/z* 268.9 [M+1]⁺.


[0159] Synthesis of Compound 8b. To a suspension of compound **8.1** (50 mg, 0.0011 mol) in water (2 mL) was added triethylamine (56 mg, 0.0005 mol) and the reaction mixture was stirred at RT for 16h. After completion of the starting material (by TLC), water was removed under reduced pressure and co-distilled with toluene (2 x 5 mL) to afford compound **8** (200 mg), which was used without any further purification.

[0160] Compounds 8a–8g. Using different propargyl alcohols and compound **4.1**, the following acids can be synthesized as exemplified in Scheme 8.

Scheme 8



Scheme 9.




[0161] **Synthesis of Compound 9.1.** To a solution of compound 1.2 (1250 mg, 0.00566 mol) in DMF (4 mL) was added potassium cyanide (520 mg, 0.0079 mol). The reaction mixture was stirred for 3 days. Additional KCN (360 mg) was added and the reaction mixture was stirred for another 24 hr. The mixture was diluted with EtOAc (150 mL) and washed with water (100 mL). The aqueous phase was extracted with EtOAc (100 mL). The organic phases were combined and washed with brine, dried over sodium sulfate and concentrated to give compound 9.1 (650 mg, 54%) as a dark brown oil. LCMS: m/z 212 $[M+1]^+$.

[0162] **Synthesis of Compound 9.2.** A vial was charged with compound 9.1 (35 mg, 0.00016 mol), acetic acid (0.7 mL, 0.01 mol), and aminoacetaldehyde dimethyl acetal (50 mg, 0.00047 mol). The reaction mixture was purged with nitrogen and stirred at 110°C overnight.

The solvent was removed and the crude product was purified on Gilson reverse phase HPLC to give compound **9.2** (15 mg, 38%) as a light brown oil. LCMS: m/z 253/255 $[M+1/M+3]^+$.

[0163] Synthesis of Compound 9. Compound **9.2** was hydrolyzed as described for compound **1** to give **9** which was used without further purification.


Scheme 10.




[0164] Synthesis of Compound 10.1. To a solution of compound **1.2** (1.0 g, 0.0045 mol) in methylene chloride (6 mL) was added trimethoxybenzylamine hydrochloride (1.0g, 0.0043 mol,) and diisopropylethylamine (1.5 mL, 0.0086 mol). The resulting mixture was stirred at RT for 3 hr. The reaction mixture was diluted with methylene chloride (80 mL) and washed 1 N HCl (2X) and brine (1 x). The organic layer was dried over MgSO₄ and concentrated to give compound **10.1** (1.6 g, 99%) as a yellow solid which was used without further purification. LCMS: m/z 382 $[M+1]^+$.

[0165] Synthesis of Compound 10.2. To a stirred solution of compound **10.1** (1.6g, 0.0042 mol) in DCM (5 mL) was added TFA (15 mL). The mixture was stirred at RT for 24 hr whereupon the solvent was removed under reduced pressure. Saturated aqueous NaHCO₃ was added to the residue and the resultant neutral aqueous mixture was extracted with EtOAc (3 x 50 mL). The combined organic layers were dried over MgSO₄ and concentrated under reduced pressure. The crude material was purified by column chromatography (0–60% ethyl acetate/hexanes) to give compound **10.2** (0.6 g, 70%) as off–white crystals.

[0166] Synthesis of Compound 10. Compound **10.2** was hydrolyzed as described for compound **1** to give acid **10**, which was used without further purification. ¹H NMR (400 MHz, Methanol–d₄): δ 8.21 (s, 1H); LCMS: 174 $[M+1]^+$.

Scheme 11.

[0167] **Synthesis of Compound 11.** Using ester **4.1** as the starting material, compound **11** was synthesized according to scheme **11**, following the procedure used for the synthesis of compound **10** (Scheme **10**).


Scheme 12.

[0168] **Synthesis of Compound 12.1:** Compound **12.1** was synthesized using a similar approach as for compound **4.1** (Scheme **4**).


[0169] **Synthesis of Compound 12.2.** To compound **12.1** (0.16 g, 0.91 mmol, 1.0 equiv), isoxazol-3-ylamine (92 mg, 1.1 mmol, 1.2 equiv), tris(dibenzylideneacetone)-dipalladium (21 mg, 0.023 mmol, 0.025 equiv), xantphos (39 mg, 0.068 mmol, 0.075 equiv), and Na_2CO_3 (133 mg, 1.4 mmol, 1.4 equiv) in toluene (3 mL) was added H_2O (16 μL , 0.91 mmol, 1.0 equiv). The reaction mixture was heated to 100 $^\circ\text{C}$ and stirred for 3 hr, whereupon it was cooled to RT. The mixture was filtered through Celite[®] and adsorbed onto SiO_2 gel. Purification by flash column chromatography (50–75–100% EtOAc/hexanes) afforded **12.2** (0.79 mg, 40%). LCMS: m/z : 221 $[\text{M}+1]^+$.

[0170] **Synthesis of Compound 12a.** To a solution of ester **12.2** (79 mg, 0.36 mmol) in THF (1.5 mL) was added a solution of LiOH (17 mg, 0.72 mmol, 2.0 equiv) in H₂O (0.50 mL). The reaction mixture was stirred at RT for 18 hr. The reaction mixture was concentrated, and the residue was dissolved in MeOH (5 mL) and water (10 mL). The solution was frozen and lyophilized for 2 days to provide **12a** (0.74 g, 100%, Li salt) as a white solid. LC-MS: *m/z*: 207 [M+1]⁺.

[0171] Compounds **12a–12c**. Using different aromatic amines and compound **12.1**, the following acids can be synthesized as exemplified in Scheme 12:




Scheme 13.




[0172] Compounds **13a** and **13b** can be synthesized by hydrolysis of compound **1.1** (Scheme 1) and by chlorination of compound **1.1** followed by hydrolysis.

Scheme 14-1.



[0173] **Synthesis of Compound 14a.** Using ester **1.1** as the starting material and 1,3-dibromo-5,5-N,N-dimethylhydantoin, compound **14a** was synthesized according to Scheme 14-1 by following the procedure used for the synthesis of compound **10** (Scheme 10).

Scheme 14-2.




[0174] **Compound 14.1.** Compound **14.1** was synthesized using a approach similar to that described in Scheme 10, except replacing the dichlorohydantoin reagent with a dibromohydantoin reagent for the first halogenation. ^1H NMR (500 MHz, CDCl_3): δ 8.93 (s, 1H), 4.51 (q, $J=7$ Hz, 2H), 1.49 (t, $J=7$ Hz, 3H). LCMS: m/z 265 $[\text{M}+1]^+$.

[0175] **Compound 14.4.** Aqueous methylamine (0.25 mL, 0.003 mol) was added to a solution of **13.1** (500 mg, 0.002 mol) in 1,4-dioxane (10 mL, 0.1 mol). The reaction mixture was stirred at RT for 18 hr. The solvent was removed *in vacuo* and the crude reaction mixture purified by reverse phase column chromatography to provide **14.4**. (350 mg, 60%). ^1H NMR (400 MHz, CDCl_3): δ 8.56 (s, 1H), 5.89 (bs, N-H), 4.47 (q, $J=7.3$ Hz, 2H), 3.12 (d, $J=4.8$ Hz, 3H), 4.43 (t, $J=4.8$ Hz, 3H; LCMS: m/z 261 $[\text{M}+1]^+$.


[0176] **Compound 14b.** Compound **14.4** (500 mg, 0.002 mol) was added to a mixture of tetrahydrofuran (2.22 mL, 0.0274 mol) and water (1.06 mL, 0.0592 mol) and the suspension stirred. Lithium hydroxide (130 mg, 0.0053 mol) was added and the reaction was stirred for 1.5 hr. The reaction mixture was then adjusted to pH 5 with 1N HCl. Solvent was removed *in vacuo* and the aqueous solution lyophilized to give the crude product **14.4** which was used without further purification. LCMS: m/z 233 $[\text{M}^+ + 1]$. ^1H NMR (400 MHz, $\text{DMSO}-d_6$): δ 8.23 (s, 1H), 6.97 (m, 1H, NH), 2.85 (d, $J=4.3$ Hz, 3H). LCMS: m/z 233 $[\text{M}+1]^+$.

Scheme 14-3.




[0177] **Compound 14c.** To a suspension of formamidinium hydrochloride (30 g, 0.252 mole) in ethanol (150ml) at 45 °C was added sodium ethoxide (prepared by dissolving Na (6.4 g, 0.282 mol) in ethanol (100 mL)) and mucobromic acid (25 g, 0.097 mol) in ethanol (50 mL). The two solutions were added simultaneously over 1 hr. After stirring the reaction mixture at 45-50 °C for 3 hr, the solvent was evaporated *in vacuo* and the residue was dissolved in ice water (100 mL). Decolorizing charcoal (2 g) was added and filtered. The filtrate was washed with ethyl acetate and aqueous layer was acidified with 12 N HCl. The aqueous layer was extracted with EtOAc (3X) and the combined organic layer was dried over anhydrous sodium sulfate. The solvent was evaporated and the residue was washed several times with ether to obtain **14c** as a light brown solid (3.8 g, 9.25%). ¹H NMR: (DMSO-d₆, 200MHz) δ: 9.22 (s, 1H), 9.18 (s, 1H).

[0178] **Compound 14d.** Using different amines and compound **14.1**, compound **14d** is prepared as exemplified in **Scheme 14.1**.



Scheme 15.



[0179] **Synthesis of Compound 15.1.** To cooled (0 °C) phosphorus oxychloride (20.0 mL, 215 mmol, 4.8 equiv.) was added DMF (6.4 mL, 83 mmol, 1.9 equiv) dropwise over 3 min. The reaction mixture was stirred for fifteen min and the ice bath was removed. 4,6-Dihydroxypyrimidine (5.0 g, 44.6 mmol, 1.0 equiv.) was added and the reaction mixture was heated to 130 °C and stirred for 3.5 hr. The mixture was cooled to RT and concentrated. Ice was slowly added to the dark brown residue, followed by 600 mL of ice water. The aqueous mixture was extracted with diethyl ether (5 x 100 mL), and the organic extracts were washed with aqueous saturated NaHCO₃ (2 x 100 mL) and brine (100 mL), and dried over anhydrous sodium sulfate and concentrated *in vacuo* to provide Compound **15** (4.42 g, 57%) as a crude orange solid, which was used without further purification.

[0180] **Synthesis of Compound 15.2.** To a solution of aldehyde **14.1** (1.50 g, 8.48 mmol, 1.0 equiv.) in toluene (18 mL) was added 7 M NH₃ in MeOH (1.8 mL, 12.7 mmol, 1.5 equiv.) and the reaction mixture was heated to 55 °C. Additional NH₃ was added (7 M in MeOH, 3.5 mL, 24.5 mmol) over the next 4 hr, and then the reaction mixture was cooled to RT. Water (2 mL) was added and the resultant mixture was concentrated. The residue was dissolve in MeOH

and adsorbed onto SiO₂ gel. Purification by flash column chromatography (20–25–33–40% EtOAc/hexanes) afforded **15.2** (0.88 g, 66%) as a beige solid. LCMS: *m/z*: 158 [M+1]⁺.


[0181] Synthesis of Compound 15.3. To a mixture of trimethoxybenzylamine (469 mg, 2.38 mmol, 1.0 equiv., HCl salt free based prior to use), 4 angstrom molecular sieves (290 mg), and aldehyde **15.2** (375 mg, 2.38 mmol, 1.0 equiv.) in dichloromethane (5 mL) was added acetic acid (0.14 mL, 2.43 mmol, 1.02 equiv.). After stirring for 3 hr at RT, sodium triacetoxyborohydride (757 mg, 3.57 mmol, 1.5 equiv.) was added and the reaction mixture was stirred at RT for 21.5 hr. The reaction mixture was diluted with dichloromethane (20 mL) and aqueous saturated NaHCO₃ (20 mL). The aqueous layer was extracted with dichloromethane (4 x 20 mL), and the combined organic extracts were washed with brine, dried over anhydrous sodium sulfate, and concentrated *in vacuo*. The resultant crude residue and Boc₂O (524 mg, 2.38 mmol, 1 equiv.) were dissolved in THF (10 mL), and pyridine (0.59 mL, 5.95 mmol, 2.5 equiv.) was added. After stirring at RT for 16.5 hr, the reaction mixture was diluted with water (25 mL), EtOAc (25 mL), and 1 N aqueous HCl (25 mL). The aqueous layer was extracted with EtOAc (4 x 30 mL). The combined organic extracts were washed with water (50 mL), 1 N aqueous HCl (50 mL), and brine (50 mL), dried over anhydrous sodium sulfate, and concentrated. Purification by flash column chromatography (50–60–66% EtOAc/hexanes) afforded compound **15.3** (403 mg, 39% over 2 steps) as a beige foam. LCMS: *m/z*: 439 [M+1]⁺.

[0182] Synthesis of Compound 15.4. A bomb was charged with chloride **15.3** (0.202 g, 0.46 mmol, 1.0 equiv.), bis(acetonitrile)dichloropalladium II (6 mg, 0.023 mmol, 0.05 equiv.), *rac*-BINAP (15 mg, 0.023 mmol, 0.05 equiv.), methanol (25 mL), and triethylamine (0.88 mL, 0.60 mmol, 1.3 equiv.). After purging and back-filling the bomb with CO (g) (3X, 50 psi), the bomb was pressurized to 50 psi CO. The reaction mixture was stirred at 100 °C for 22 hr, and then cooled to RT and the bomb was carefully vented. LC–MS analysis indicated incomplete conversion, so additional bis(acetonitrile)dichloropalladium II (18 mg, 0.069 mmol, 0.15 equiv.), *rac*-BINAP (44 mg, 0.069 mmol, 0.15 equiv.) and triethylamine (0.10 mL, 0.7 mmol) were added and the bomb was repressurized to 60 psi CO and heated to 105 °C. The reaction mixture was stirred at 105 °C for 23 hr, then cooled to RT and the bomb was carefully vented. The reaction mixture was filtered through Celite and adsorbed onto SiO₂ gel. Purification by flash


column chromatography (10–20–40–50–75–100% EtOAc/hexanes) afforded **15.4** (0.109 g, 51%) as a yellow foam. LCMS: m/z : 463 $[M+1]^+$.

[0183] Synthesis of Compound 15. To a solution of ester **15.4** (0.103 g, 0.22 mmol) in THF (0.85 mL) was added a solution of LiOH (6 mg, 0.27 mmol, 1.2 equiv.) in H₂O (0.27 mL). The reaction mixture was stirred at RT for 18 hr. The reaction mixture was concentrated, and the residue was dissolved in MeOH (5 mL) and water (10 mL). The solution was frozen and lyophilized for 2 days to provide **15** (0.101 g, 100%, Li salt) as a pale yellow solid. LC–MS: m/z : 449 $[M+1]^+$.


[0184] Compounds 15a–15e. Using different amines and compound **15.2**, the following acids can be synthesized as exemplified in Scheme 15:



Scheme 16.




[0185] Compound **16** is commercially available and was used without additional purification.

Scheme 17.

[0186] **Compound 17.1.** A 5 mL microwave vial was flushed with nitrogen gas. To the vial was added compound **14.3** (500 mg, 0.20 mmol), zinc cyanide (130 mg, 0.11 mmol), tris(dibenzylideneacetone)dipalladium(0) (20 mg, 0.002 mmol), 1,1'-bis-(diphenylphosphino)ferrocene (30 mg, 0.11 mmol), zinc acetate (20 mg, 0.009 mmol) and zinc (6 mg, 0.009 mmol). N,N-Dimethylformamide (2.3 mL, 2.9 mmol) was added and the reaction capped and flushed with nitrogen gas (3X). The reaction mixture was heated in a microwave reactor at 130 °C for 1 hr. The solvent was removed *in vacuo* and the residue added to 5 mL of 5% NaHCO₃ and extracted with EtOAc (3X). The organic layers were combined and washed with brine. The solvent was removed *in vacuo* and the crude product was used in subsequent reactions without further purification. LCMS: *m/z* 193.07 [M+1]⁺.


[0187] **Compound 17.** Compound **17** was obtained from **17.1** using the hydrolysis procedure outlined in Scheme 1 (**1.3** to **1a**). LCMS: *m/z* 165.16 [M+1]⁺.

Scheme 18.


[0188] **Synthesis of Compound 18.1.** To the suspension of NaH (70 mg, 60% NaH in paraffin oil, 0.00295 mol) in THF (5 mL) was added imidazole (201 mg, 0.00295 mol) at 0 °C and stirred for 30 min. Compound **4.1** (500 mg, 0.0026 mol) was added at 0 °C and the reaction mixture was heated at 60 °C for 18 hr. The reaction mixture was quenched with ice water (2 mL) and extracted with EtOAc (3 x 20 mL). The combined organic layers were dried over Na₂SO₄ and concentrated under reduced pressure. The crude material was purified by flash column chromatography to give **18.1** (300 mg, 52%).

[0189] **Synthesis of Compound 18a.** Compound **18.1** was hydrolyzed as described for compound **1** to give **18a** which was used without further purification


[0190] **Compounds 18a-18l.** Using different heterocycles or alcohols and compound **4.1**, the following acids can be synthesized as exemplified in Scheme **18**:




Scheme 19.1



[0191] **Synthesis of Compounds 19IID-19rrD.** Compounds **19IID-19rrD** were prepared by TFA deprotection of the corresponding Boc protected amines **5IID-5rrD** under standard TFA deprotection conditions.



Scheme 20.



[0192] **Synthesis of Compound 20.1.** To a stirred solution of NaOEt (2.7 g, 0.04 mol) in EtOH (40 mL) was added formamidine acetate (4.2 g, 0.04 mol), followed by addition of diethylfluoromalonate in ethanol (10 mL) at 0°C . The reaction mixture was stirred at 90°C overnight. Ethanol was removed under reduced pressure and the reaction mixture was acidified with conc. HCl to pH 1. The resulting solid was filtered and dried under vacuum to afford **20.1** (crude, 750 mg, 52%). $^1\text{H-NMR}$ (DMSO-d_6 200 MHz): δ 12.40 (bs, 2H), 7.89 (s, 1H).


[0193] **Synthesis of Compound 20.2.** A mixture of **20.1** (800 mg, 0.0062 mol) and N,N-diethylaniline in POCl₃ (3 mL) was refluxed at 100 °C overnight. The reaction mixture was poured into ice water and extracted with hexane (3 x 100 mL). The combined organic layers were washed with saturated NaHCO₃ and dried over Na₂SO₄. Hexane was removed under reduced pressure and the obtained crude material was purified by column chromatography to give 300 mg of **20.2** (30%). ¹H-NMR (CDCl₃ 200 MHz): δ 8.61 (s, 1H); m/z: 167 [M+1]⁺.

[0194] **Synthesis of Compound 20.3.** To a stirred solution of **20.2** (120 mg, 0.000722 mol) in n-butanol (0.5 mL) was added NH₄OH (1 mL). The reaction mixture was heated at 90 °C for 2.5 hr in a sealed tube. The reaction mixture was cooled to 0 °C, and the resulting solid was filtered and dried under vacuum to give **20.3** (60 mg, 57%). ¹H-NMR (DMSO-d₆ 500 MHz): δ 8.03 (s, 1H), 7.60 (s, 2H); m/z: 148 [M+1]⁺.


[0195] **Synthesis of Compound 20.4.** To a stirred solution of **20.3** (150 mg, 0.00102 mol) in n-butanol (2 mL) and acetonitrile (2 mL) was added DIPEA (0.2 mL, 0.0013 mol), [2,2'-bis(diphenylphosphino)-1,1'-binaphthyl]palladium (II) chloride (41 mg, 0.000051 mol) in a steel bomb was stirred at 100 °C under CO (100 psi) overnight. The progress of the reaction was monitored by TLC. After completion of the reaction, solvents were removed under reduced pressure and the obtained crude material was purified by column chromatography to give **20.4** (95 mg, 44%). ¹H-NMR (DMSO-d₆ 500 MHz): δ 8.21 (s, 1H), 7.64 (s, 2H), 4.29 (t, J= 6.5 Hz, 2H), 1.67 (m, 2H), 1.41 (m, 2H), 0.923 (t, J= 7.5 Hz, 3H); m/z: 214 [M+1]⁺.

[0196] **Synthesis of Compound 20a.** To the stirred solution of **20.4** (120 mg, 0.000563 mol) in THF (1 mL) and water (1 mL) was added LiOH (25 mg, 0.000619 mol) at 0 °C. The reaction mixture was stirred at RT for 2 hr. The reaction mixture was concentrated under reduced pressure to give 110 mg of **20a** (crude) as a Li salt. ¹H-NMR (DMSO-d₆ 500 MHz): δ 7.96 (s, 1H), 6.91 (s, 2H); m/z: 158 [M+1]⁺.

[0197] **Compounds 20a-20b.** Using different amines and compound **20.2**, the following acids can be synthesized as exemplified in **Scheme 20**.




Scheme 21.



[0198] Synthesis of Compound 21. To a solution of **5vDa** (56 mg, 0.10 mmol) in acetonitrile (1 mL, 20 mmol) in a sealed microwave tube under nitrogen atmosphere was added phosphoryl chloride (37 μ L, 0.40 mmol), followed by pyridine (8.1 μ L, 0.10 mmol). The reaction mixture was stirred at RT overnight. Next morning, the tube was heated at 80 °C under microwave irradiation for 10 min. The reaction was quenched by addition of water. The mixture was diluted with DMSO, purified by reverse phase preparatory HPLC (acetonitrile 10-90%, buffered with TFA), and lyophilized to afford 36 mg (yield 57%) of **21** as a white solid.

Scheme 22.




[0199] Synthesis of Compound 22.1. To a mixture of pyrimidine-4,6-dicarboxylic acid (34 mg, 0.20 mmol) in DMF (3 mL, 40 mmol), cooled with ice-bath, was added 1-hydroxybenzotriazole (200 mg, 0.15 mmol), N-(3-dimethylaminopropyl)-N'-ethylcarbodiimide hydrochloride (48 mg, 0.25 mmol) and 4-methylmorpholine (16 μ L, 0.15 mmol). The mixture was stirred at 0 °C for 30 min. Compound **Da** (70 mg, 0.20 mmol) was added, and stirred in the cold bath (allow the ice to melt) for 2 hr. LC-MS showed the desired mono-amide intermediate, along with di-amide byproduct. To the reaction mixture was added acetohydrazide (30 mg, 0.40

mmol), followed by an additional portion of N-(3-dimethylaminopropyl)-N'-ethylcarbodiimide hydrochloride. The reaction mixture was stirred at RT over the weekend. The reaction mixture was concentrated *in vacuo* to remove most of DMF solvent, re-dissolved in DMSO/MeOH, purified by reverse phase HPLC (20-100%), and lyophilized to obtain 20 mg (20%) of **22.1** as a white solid

[0200] Synthesis of Compound 22. A mixture of **22.1** (11 mg, 0.020 mmol) in phosphoryl chloride (250 μ L, 2.7 mmol) was heated in a sealed tube in a 100 °C oil bath for 30 min. The reaction mixture was partitioned between EtOAc and aqueous saturated NaHCO₃. The organic phase was dried, filtered, and concentrated. The residue was purified by reverse phase HPLC (20-100% acetonitrile, TFA) to provide 5 mg (50%) of **22** as a pink solid.

Scheme 23.



[0201] Synthesis of Compound 23.1. In a 100 mL round-bottom flask **D.3** (5.00 g, 0.0163 mole) and oxalyl chloride (1.52 mL, 0.0180 mole) were dissolved in acetonitrile (50.0 mL). The resulting solution evolved gas for 5 min. After 5 min, N,N-dimethylformamide (0.100 mL) was added dropwise at RT with much gas evolution. The reaction was allowed to stir at RT for 3 hr. Methanol (50.0 mL) was added in one portion and allowed to stir for an additional 2 hr. The solvent was then removed *in vacuo*. The resulting residue was then diluted with 250 mL of EtOAc and washed with 2 x 200 mL of sat NaHCO₃ 1 x 100 mL brine. The EtOAc layer was then dried over Na₂SO₄ and removed *in vacuo*. Yielded 5.00 g of **23.1**, which was used without further purification. LCMS *m/z* 321 [M+1]⁺:

[0202] **Synthesis of Compound 23.2.** In a 50 mL round-bottom flask **23.1** (5.00 g, 0.0156 mole) was taken up in HBr/AcOH (4.0 M, 10 mL). The resulting brown reaction mixture was allowed to stir at RT for 18 hr. After 18 hr the HBr/AcOH was removed *in vacuo* to yield a brown solid. The brown solid (HBr salt) was then titrated with CH₂Cl₂, which removed most of the brown color, yielding an off-white solid. The resulting solid was then taken up in 300 mL of EtOAc and washed with 75 mL of sat NaHCO₃ (2x) and 75 mL brine. The EtOAc was then dried over Na₂SO₄. The EtOAc was removed in vacuo to yield 1.70 g (0.0156 mol 59%) of the desired **23.2**, which was used without further purification. LCMS *m/z* 187 [M+1]⁺.

[0203] **Synthesis of Compound 23.3.** In a 100 mL round-bottom flask, **23.2** (1.21 g, 0.00697 mole), methyl 2-(1-aminoethyl)thiazole-5-carboxylate (1.30 g, 0.00697 mole), and TBTU (2.69 g, 0.00837 mole) were dissolved in N,N-dimethylformamide (25.0 mL, 0.323 mole) to which was added N,N-diisopropylethylamine (3.64 mL, 0.0209 mole). The resulting yellow brown solution was allowed to stir at RT for 3 hr. The reaction mixture was diluted with 250 mL EtOAc, washed with 75 mL satd. NaHCO₃ (2x), 75 mL of water, and 50 mL brine. The organic layer was dried over Na₂SO₄ and concentrated to a brown oil. The residue was purified by flash column chromatography (50% EtOAc/Hexanes gradient to 100% EtOAc) to yield 1.12 g (0.0070 47%) of desired product **23.3**. LCMS *m/z* 342 [M+1]⁺.


[0204] **Synthesis of Compound 23.4** In a 50 mL round-bottom flask, **23.3** (1.12 g, 0.00328 mol) was taken up in THF (20 mL, 0.2 mole) to which was added a solution of LiOH (0.08633 g, 0.003605 mole) in water (4 mL, 0.2 mole). The resulting reaction mixture was then allowed to stir at RT for 6 hr. The solvent was removed *in vacuo* and the resulting residue was taken up in 200 mL CH₂Cl₂ and then washed 50 mL sat NH₄Cl (2x) and brine to yield 0.653 g (0.0038 mole, 60%) of **23.4**. LCMS *m/z* 327 [M+1]⁺.

[0205] **Synthesis of Compound 23.5.** In a 25 mL round-bottom flask **23.4** (0.382 g, 0.00116 mole), 1-hydroxybenzotriazole (0.157 g, 0.00116 mole) and 2-amino-1-phenylethanone (0.200 g, 0.00116 mole) were taken up in DMF (5 mL, 0.06 mole), and to this was added N-(3-dimethylaminopropyl)-N'-ethylcarbodiimide hydrochloride (0.268 g, 0.00140 mole) and then N,N-diisopropylethylamine (0.203 mL, 0.00116 mole). The resulting cloudy solution was allowed to stir at RT for 3 hr. The reaction mixture was diluted with 100 mL EtOAc and washed with 50 mL Sat NaHCO₃ (2x) and brine. The organic layer was dried over Na₂SO₄, filtered, and

concentration. The residue was dissolved in CH_2Cl_2 and eluted through a silica gel column with 90-100% EtOAc/hexanes to afford 270 mg of **23.5**. LCMS m/z 446 $[\text{M}+1]^+$.


[0206] Synthesis of Compound 23. In a 5 mL microwave reaction vial **23.5** (0.100 g, 0.000225 mole) and ammonium acetate (0.173 g, 0.00225 mole) were taken up in acetic acid (4.0 mL, 0.070 mole). The vial was sealed and allowed to stir at RT for 5 min. The reaction was heated at 175 °C under microwave radiation for 15 min. The acetic acid was removed in vacuo to yield a slightly yellow oil. The oil was taken up in 100 mL of CH_2Cl_2 and washed with 50 mL of sat NaHCO_3 . The organic layer was then washed with 50 mL NaHCO_3 , 50 mL H_2O , and 35 mL brine. The organic layer was then dried over Na_2SO_4 and the solvent was removed *in vacuo*. The resulting slightly yellow oil was taken up in DMSO and purified by preparative HPLC (10%-90% CH_3CN /water 0.1% TFA acidic method) to yield 42 mg (0.00022 mol, 35%) of **23** as a TFA salt. LCMS m/z 427 $[\text{M}+1]^+$.

Scheme 24.



[0207] Synthesis of Compound 24a and 24b. A solution of **14aD** (200 mg, 0.0004 mole) and [1,1'-bis(diphenylphosphino)ferrocene]dichloropalladium(II) (10 mg, 0.00002 mol) in methanol (30 mL) was treated with *N,N*-diisopropylethylamine (75.90 μL , 0.0004358 mole). The mixture was then placed in the parr autoclave and flushed with CO. The autoclave was then charged with CO to 10 bar and heated to 100 °C for 16 hr. After cooling to RT, the mixture was filtered to remove any solid, and the resulting reddish brown solution was concentrated to a brown solid. The residue was purified via preparatory HPLC to afford **24a** (22 mg, 11%) and **24b** (2.8 mg, 1%). **24a**: m/z 530 $[\text{M}+1]$. **24b**: m/z 554 $[\text{M}+1]^+$.


Scheme 25.




[0208] Synthesis of Compound 25aDa. To a stirred solution of **25.1** (600 mg, 3.569 mmol) in DMF (50 mL) at 0 °C were added HOBT (361 mg, 2.676 mmol), EDCI (855 mg, 4.46 mmol) and NMM (288 mg, 2.854 mmol). After stirring for 30 min., **Da** (626 mg, 1.784 mmol) was added, and the reaction was stirred for 4 hr at 10-15 °C. After completion, the reaction mixture was diluted with water (50 mL) and extracted twice with DCM (2 x 50mL). The combined organic layer was dried over anhydrous Na_2SO_4 and concentrated. The residue was dissolved in water and acidified with 6 N HCl (pH = 2-3). The solid obtained was filtered, washed with water, EtOAc (5 mL), and hexane, and dried under vacuum. The white solid obtained was codistilled twice with CCl_4 to obtain **25aDa** (710 mg, 79.41%). ^1H NMR (DMSO- d_6 , 500 MHz) δ : 14.0-14.4 (s, 1H, D_2O exchangeable), 11.7 (s, 1H, D_2O exchangeable), 9.95 (d, 1H, D_2O exchangeable), 9.5 (s, 1H), 8.8 (2s, 2H), 8.5 (s, 1H), 8.4 (s, 1H), 5.5 (m, 1H), 1.7 (d, 3H); m/z 501 $[\text{M}+1]^+$.

[0209] Synthesis of Compound 25bDa. To a solution of compound **25aDa** (100.0 mg, 0.1997 mmol) in DMF (4.0 mL) were added HOBT (200 mg, 0.15 mmol), EDCI (47.8 mg, 0.250 mmol), and 4-methylmorpholine (60 μL , 0.6 mmol). This resulting brown solution was then treated with ammonium chloride (21 mg, 0.40 mmol). After stirring for 4 hr, this mixture was purified via preparatory Gilson HPLC (flow rate 20, from 10% B (MeCN with 0.1% formic acid) to 95% B in 10 min), affording **25bDa** as a white solid (19 mg, 19%). ^1H NMR (400MHz, DMSO- d_6) δ = 11.72 (br. s., 1 H), 9.93 (d, J = 8.6 Hz, 1 H), 9.50 (s, 1 H), 8.77 (s, 1 H), 8.75 (s, 1 H), 8.55 (s, 1 H), 8.47 (s, 1 H), 8.46 - 8.41 (m, 1 H), 8.10 (s, 1 H), 5.54 - 5.45 (m, 1 H), 1.70 (d, J = 7.1 Hz, 3 H); m/z 500 $[\text{M}+1]^+$.


[0210] **Compounds 25aD-25lD.** Using different amines and compound **D**, the following compounds were synthesized as exemplified in Scheme 25.




[0211] **Compounds 25cDa, 25iDa, 25kDa, 25lDa, 25mDa, and 25nDa.** Using different amines and compound **Da**, the following compounds were synthesized as exemplified in Scheme 25.




Scheme 26.




[0212] **Synthesis of Compounds 26aD-26eD.** Compounds 26aD-26eD were prepared by coupling of the corresponding carboxylic acid or acid chloride with amine 19lID under standard acid coupling conditions.




Scheme 27.
















[0213] **Synthesis of Compounds 27aD-26hD.** Compounds 27aD-27hD were prepared by hydrogenation of the corresponding alkynes under standard alkyne hydrogenation conditions.




Scheme 28.



[0214] **Synthesis of Compounds 28a-28m.** Compounds **28a-28m** were prepared by reductive amination of different amines and compound **28.2** as exemplified in Scheme **28**.




Scheme 29.




[0215] Synthesis of Compound 29a. To a solution of **10D** (30.0 mg, 0.059 mmol) in THF (5.00 mL) were added 1 M of potassium tert-butoxide in THF (0.071 mL, 0.07 mmol), 4-methylmorpholine (0.020 mL, 0.178 mmol) and iodoacetic acid (12.1 mg, 0.065 mmol). After stirring at 25 °C for 18 hr, the crude mixture was purified *via* preparatory reverse-phase HPLC, affording **29a** as a white solid (20 mg, 60%). ¹H NMR (400MHz, DMSO-d₆) δ = 11.77 (s, 1 H), 9.58 (d, *J* = 7.8 Hz, 1 H), 8.78 (s, 1 H), 8.76 (s, 1 H), 8.58 (s, 1 H), 8.47 (s, 1 H), 8.06 (t, *J* = 6.0 Hz, 1 H), 5.40 - 5.31 (m, 1 H), 4.07 (d, *J* = 6.1 Hz, 2 H), 1.59 (d, *J* = 7.1 Hz, 3 H); m/z 564 [M+1]⁺.

[0216] Synthesis of Compound 29b. Compound **29b** was synthesized as exemplified in **Scheme 29** substituting ethyl 2-bromoacetate for the iodoacetic acid.



Scheme 30.




[0217] **Synthesis of Compound 30.1.** A solution of tert-butyl 4-aminopiperidine-1-carboxylate (1g, 0.0049 mol), **10** (868 mg, 0.0049 mol), EDCI (2.3 g, 0.0124 mol) and HOBT (269 mg, 0.0019 mol) in DMF (10 ml) was stirred at RT for 16h. The reaction mixture was diluted with water (50 ml) and extracted with ethyl acetate (3x 50 ml). The combined organic layers was washed with water (3x30 ml), dried over Na₂SO₄ and concentrated under reduced pressure. The resulting mixture was purified by column chromatography to give **30.1** (1.7 mg, 58%). ¹H-NMR (CDCl₃ 500 MHz): δ 8.41 (s, 1H), 7.89 (bs, 1H), 5.65 (bs, 2N-H), 4.15 (m, 4H), 2.93 (m, 2H), 1.94 (m, 2H), 1.42 (s, 9H); m/z 356 [M+1]⁺.

[0218] **Synthesis of Compound 30.2.** To a stirred solution of **30.1** (600 mg, 0.0016 mol) in DCM (4 ml) and cooled to 0°C, then added TFA (4ml). The reaction mixture was stirred at RT for 2 hr and then volatiles were removed under reduced pressure. The residue was co-distilled with toluene (2x10ml) to afford **30.2** as light yellow solid (400 mg, 93%). ¹H-NMR (CD₃OD, 500 MHz) δ 8.28 (s, 1H), 4.18 (t, 1H), 3.70 (dd, 2H), 3.21 (m, 2H), 2.25 (d, 2H), 1.85 (m, 2H); m/z 255.9 [M+1]⁺.


[0219] **Synthesis of Compound 30a.** A solution of **30.2** (100mg, 0.00039 mol), **30.3** (86 mg, 0.00039 mol), EDCI (188 mg, 0.00098 mol), HOBT (23 mg, 0.00017 mol) and DIPEA (152mg, 0.0011) in DMF (3 ml) was stirred at RT for 16 hr. The reaction mixture was diluted with water (20 ml) and extracted with ethyl acetate (3x 20 ml). The combined organic layers were washed with (2x 10 ml), dried over Na₂SO₄ and concentrated under reduced pressure. The resulting mixture was purified by column chromatography to give **30a** (60 mg, 33%). ¹H-NMR (DMSO-d₆, 500 MHz): δ 8.62(d, J=7.5 Hz, 1H), 8.32 (s, 1H), 7.35 (dd, J=8, 1H), 6.95 (d, J= 25

Hz, 2H), 6.82 (d, $J=7.5\text{Hz}$, 1H); 6.18(d, 1H), 4.25(d, 1H), 4.0-.8(br, 4H), 3.15(t, 1H), 2.93(t, 1H), 1.92(br, 2H), 1.6-1.49 (br, 2H); m/z 456.8 $[M+1]^+$.

[0220] Synthesis of Compound 30b-30c. Compound **30b** was prepared as exemplified in **Scheme 30** utilizing tert-butyl 3-aminopyrrolidine-1-carboxylate. Compound **30c** was prepared as exemplified in **Scheme 31** utilizing tert-butyl 3-aminopiperidine-1-carboxylate.



Scheme 31.



[0221] Synthesis of Compound 31.1. The solution of tert-butyl 3-aminopiperidine-1-carboxylate (500 mg, 0.0024 mol), **10** (434 mg, 0.0024 mol), EDCI (1.2g, 0.0062 mol) and HOBT (136 mg, 0.0009 mol) in DMF (5 ml) was stirred at RT for 16 hr. The reaction mixture was diluted with water (25 ml) and extracted with ethyl acetate (3x 25 ml). The combined organic layers was washed with water (3x 15 ml), dried over Na_2SO_4 and concentrated under reduced pressure. The resulting mixture was purified by column chromatography to give **31.1** (560 mg, 63%). $^1\text{H-NMR}$ (DMSO- d_6 , 500MHz): δ 8.45 (s, 1H), 8.25 (s, 1H), 7.90-7.20 (bs, 2N-H), 3.89-3.65 (m, 3H), 2.95-2.78 (m, 2H), 1.94 -1.82 (m, 2H), 1.75-1.68 (m, 2H), 1.42 (s, 9H); m/z 355.9 $[M+1]^+$.

[0222] Synthesis of Compound 31.2. To the stirred solution of **31.1** (560 mg, 0.0015 mol) in DCM (4 ml) at 0°C was added TFA (3 ml). The reaction mixture was stirred at RT for 2 hr, DCM was removed under reduced pressure and the resulting crude material was co-distilled with toluene (2x10 ml) to obtain **31.2** (360 mg, 58%). $^1\text{H-NMR}$ (DMSO- d_6 , 500 MHz): δ 8.45 (s,

1H), 8.25 (s, 1H), 7.9-7.2 (bs, 2N-H), 3.89-3.65 (m, 3H), 2.95-2.78 (m, 2H), 1.94 -1.82 (m, 1H), 1.75-1.68 (m, 1H); m/z: 255.9 [M+1]⁺.



[0223] Synthesis of Compound 31.3. To a solution of **31.2** (100 mg, 0.0039 mol) in acetonitrile (5 ml) was added DIPEA (151 mg, 0.0011 mol), followed by tert-butyl bromoacetate (0.7 ml, 0.00047 mol) and stirred at RT for 6 hr. The reaction mixture was concentrated under reduced pressure and the resulting crude material was purified by column chromatography to give **31.3** (100 mg, 69%). ¹H-NMR (DMSO-d₆, 500 MHz): δ 8.41 (d, J= 8.0 Hz, 1H), 8.28 (s, 1H), 7.60-7.50 (bs, 2N-H), 5.74 (s, 1H), 3.12 (s, 2H), 2.81 (d, J= 8.5 Hz, 2H), 2.62-2.61 (m, 2H), 2.21-2.19 (m, 2H), 1.68-1.59 (m, 2H), 1.42 (s, 9H); m/z: 370 [M+1]⁺.

[0224] Synthesis of Compound 31.4. To the stirred solution of **31.3** (100 mg, 0.00022 mol) in DCM (2 ml) at 0°C was added TFA (2 ml). The reaction mixture was stirred at RT for 2 hr and DCM was removed under reduced pressure. The resulting crude material was co-distilled with toluene (2 x 10 ml) to obtain **31.4** (crude 70 mg). ¹H-NMR (CD₃OD, 500 MHz): δ 8.50 (s, 1H), 4.38-4.20 (m, 1H), 4.18 (s, 2H), 3.64-3.61 (m, 2H), 3.21-3.01 (m, 2H), 2.19-2.01 (m, 2H), 1.20-1.16 (m, 2H); m/z: 314 [M+1]⁺.

[0225] Synthesis of Compound 31a. The solution of **31.4** (100 mg, 0.00031 mol), 3-(trifluoromethyl) aniline (51 mg, 0.00031 mol), EDCI (152 mg, 0.00079 mol), HOBT (17 mg, 0.00012 mol) and DIPEA (50 mg, 0.00031 mol) in DMF (2 ml) was stirred at RT for 16 hr. The reaction mixture was diluted with water (20 ml) and extracted with ethyl acetate (3x 20 ml). The combined organic layers were washed with water (2x 10 ml), dried over Na₂SO₄ and concentrated under reduced pressure. The resulting mixture was purified by column chromatography to give **31a** (29 mg, 25%). ¹H-NMR (DMSO-d₆, 500 MHz): δ 9.99 (s, 1H), 8.57 (d, J=8.0 Hz, 1H), 8.28 (s, 1H), 8.10 (s, 1H), 7.82 (d, J=7.5 Hz, 1H), 7.54-7.51 (m, 1H), 7.39 (d, J=7.5 Hz, 1H), 4.03-3.98 (m, 1H), 3.20-3.06 (m, 2H), 2.69 (d, J=9.5 Hz, 1H), 2.49-2.30 (m, 2H), 1.68-1.59 (m, 3H), 1.42-1.34 (m, 1H); m/z: 456.9 [M+1]⁺.

[0226] Synthesis of Compound 31b. Compound **31b** was prepared as exemplified in **Scheme 31** utilizing tert-butyl 3-aminopyrrolidine-1-carboxylate. Compound **31c** was prepared as exemplified in **Scheme 31** utilizing tert-butyl 4-aminopiperidine-1-carboxylate.

Scheme 31.


**Scheme 32.**

[0227] Synthesis of compound 32.2. The solution of **31.4** (100mg, 0.00032 mol), 2-tert-butylpyrimidine-4,5-diamine **32.1** (63 mg, 0.00032 mol), EDCI (152 g, 0.00079 mol), HOBT (16 mg, 0.00011 mol) and DIPEA (124 mg, 0.00095 mol) in DMF (5 ml) was stirred at RT for 16 hr. The reaction mixture was diluted with water (20 ml) and extracted with ethyl acetate (3x 20 ml). The combined organic layers was washed with water (3x 20 ml), dried over Na₂SO₄ and concentrated under reduced pressure. The resulting mixture was purified by column chromatography to give **32.2** (100 mg, 68%). ¹H-NMR (DMSO-d₆, 500 MHz): δ 9.08 (s, 1H), 8.56 (bs, 1N-H), 8.25 (s, 1H), 8.00 (s, 1H), 6.46-6.38 (bs, 2H), 4.62-4.55 (bs, 2H), 4.02-3.99 (m, 1H), 3.18-3.09 (m, 2H), 2.39-2.36 (m, 2H), 2.22-2.15 (m, 2H), 2.02-1.98 (m, 2H), 1.69-1.58 (m, 2H), 1.22 (s, 9H); m/z 461.8 [M+1]⁺.


[0228] Synthesis of Compound 32a. To the stirred solution of **32.2** (100 mg, 0.0002 mol) in acetic acid (5 ml) was stirred at 130 °C for 24 hr. After completion of the starting material, acetic acid was completely removed under reduced pressure. The resulting reaction mixture was co-distilled with toluene (2x10ml) and the obtained crude was purified by preparative reverse-phase HPLC to give **32** (22mg, 23%). ¹H-NMR (DMSO-d₆, 500 MHz): δ 9.08(s, 1N-H), 8.56 (bs, 1N-H), 8.25 (s, 1H), 8.00 (s, 1H), 4.62-4.55 (bs, 2H), 4.02-3.99 (m, 1H), 3.18-3.09 (m, 2H),

2.39-2.36 (m, 2H), 2.22-2.15 (m, 2H), 2.02-1.98 (m, 2H), 1.69-1.58 (m, 2H), 1.22 (s, 9H); m/z 443.9 $[M+1]^+$.

[0229] Synthesis of Compound 32b. Compound 32b was synthesized as described in **Scheme 32** utilizing 2-trifluoromethylpyrimidine-4,5-diamine.



Scheme 33.



[0230] Synthesis of Compound 33.2. Compound 33.1 (30 g, 178.5 mmol) was treated with ammonium hydroxide solution (300 mL) at 0 °C. The reaction mixture was warmed to RT and stirred for 10 hr. After completion of the starting material (by TLC), the precipitated solid was filtered and dried under vacuum. The crude material was co-distilled with toluene to provide **33.2** (15 g, 60.43%) as brown color solid. ^1H NMR (200 MHz, DMSO- d_6) δ 11.6-11.0 (brs, 1H, D₂O exchangeable), 8.25 (s, 1H), 8.0-7.9 (brs, 1H, D₂O exchangeable), 7.9-7.8 (brs, 1H, D₂O exchangeable), 6.8 (s, 1H); m/z 140.0 $[M+1]^+$.

[0231] Synthesis of Compound 33.3. A mixture of **33.2** (15 g, 107.9 mmol) in POCl₃ (105 mL, 7 volumes) was heated at reflux for 16 hr. After completion of the starting material (by TLC), the reaction mixture was cooled to RT, poured into ice cold water and neutralized with

aqueous ammonium hydroxide solution. Aqueous layer was extracted with ethyl acetate (3 × 200 mL), and combined organic layers were dried over anhydrous Na₂SO₄ and evaporated under vacuum to afford crude compound. The crude compound was purified over silica gel column chromatography eluting with 10% ethyl acetate/hexane to afford **33.3** (9.8 g, 65.33%) as pale yellow syrup. ¹H NMR (1H, 200 MHz, CDCl₃): δ 9.15 (s, 1H), 7.75 (s, 1H).

[0232] **Synthesis of Compound 33.4.** To a stirred solution of **33.3** (4 g, 28.77 mmol) in absolute ethanol (40 mL) was added freshly prepared NaOEt (5.86 g, 86.33 mmol) at RT and stirred for 3 hr. After complete consumption of the starting material (by TLC), the reaction mixture was diluted with absolute ethanol (40 mL), and treated with NH₄OAc (8.87 g, 115.08 mmol) at RT and continued stirring for overnight at RT. The reaction mixture was filtered and the filtrate was evaporated under reduced pressure and the residue was dissolved in absolute ethanol (120 mL). To this chloro-acetone (6.93 mL, 86.33 mmol) was added at RT and the reaction mixture was heated at reflux temperature for 16 hr. After completion of the starting material (by TLC), the volatiles were evaporated under reduced pressure. The resulting residue was dissolved in water and neutralized with saturated NaHCO₃ solution. The aqueous layer was extracted with ethyl acetate (2 × 100 mL). The combined organic extracts were dried over anhydrous Na₂SO₄ and concentrated under vacuum. The crude material was purified by column chromatography eluting with 40% EtOAc/hexane to obtain **33.4** (300 mg, 6.97 %) as brown color solid. ¹H NMR (200 MHz, CDCl₃) δ 10.3-10.0 (brs, 1H, D₂O exchangeable), 8.65 (s, 1H), 7.41 (2s, 1H), 6.95 (2s, 1H), 4.6-4.4 (q, 2H), 2.4 (d, 3H), 1.41 (t, 3H); m/z 205.0 [M+1]⁺.

[0233] **Synthesis of Compound 33.5.** A mixture of **33.4** (300 mg, 1.47 mmol) in HBr-acetic acid (10 mL) was stirred at reflux temperature for 4 hr under inert atmosphere. After completion of the starting material (by TLC), the solvent was evaporated under reduced pressure to afford crude compound. The crude compound was dissolved in water; aqueous layer was washed with ethyl acetate (30 mL). The aqueous layer was evaporated under reduced pressure and to the crude compound was dried with toluene (co-distilled) to obtain **33.5** (200 mg, 77.51%) as brown colored solid. ¹H NMR (1H, 200 MHz, DMSO-*d*₆) : δ 8.15 (s, 1H), 6.80 (s, 1H), 6.50 (s, 1H), 2.18 (s, 3H); m/z 177.0 [M+1]⁺.


[0234] **Synthesis of Compound 33.6.** A mixture of **33.5** (0.2 g, 1.13 mmol) in POCl₃ (10 mL) was heated at reflux temperature for 4 hr under inert atmosphere. After completion of the starting precursor (by TLC), reaction mixture was poured into ice water and neutralized to pH ~

7 using NaHCO_3 . The aqueous layer was extracted with EtOAc (2×50 mL). The combined organic extracts were dried over anhydrous Na_2SO_4 and evaporated under reduced pressure to provide **33.6** (0.15 g, 68%) as a brown colored solid. ^1H NMR (200 MHz, $\text{DMSO}-d_6$) δ 9.18 (s, 1H), 8.60 (s, 1H), 7.41 (s, 1H), 2.31 (s, 3H); m/z 194.9 $[\text{M}+1]^+$.


[0235] Synthesis of Compound 33.7. To a stirred solution of **33.6** (0.3 g, 1.54 mmol) in acetonitrile (9.0 mL) and *n*-BuOH (9.0 mL) in steel bomb was added dppf-PdCl_2 (0.15 g) followed by *N*-ethyl-diisopropylamine (0.4 mL, 2.3 mmol) at RT under inert atmosphere. The steel bomb was filled with carbon monoxide (120 psi) and heated at 65 °C for 16 hr. After consumption of the starting material (by TLC), the reaction mixture was filtered through a pad of celite. The filtrate was evaporated under reduced pressure to obtain a crude material that was purified over silica gel column chromatography eluting with 30% EtOAc/hexane to afford **33.7** (mixture of two isomers) (0.2 g, 49%) as light brown color solid. ^1H NMR (200 MHz, CDCl_3) δ 10.4-10.2 (brs, 1H), 9.12 (2s, 1H), 8.62 (2s, 1H), 7.01 (d, 1H), 4.42 (t, 2H), 2.38 (d, 3H), 1.81-1.72 (m, 2H), 1.61-1.40 (m, 2H), 1.02 (t, 3H); m/z 260.9 $[\text{M}+1]^+$.

[0236] Synthesis of Compound 33a. To a stirred solution of **33.7** (0.2 g, 0.76 mmol) in THF (1.5 mL) was added LiOH solution (1M in H_2O) (0.769 mL, 0.76 mmol) at RT under inert atmosphere and the resulting mixture was stirred for 2 hr at RT. After complete consumption of starting precursor (by TLC), the volatiles were evaporated under vacuum and crude material was dissolved in water (10 mL). Aqueous layer was washed with EtOAc (10 mL) and acidified with 2N HCl at 0 °C. The precipitated solid was filtered, washed with hexane (10 mL) and dried under vacuum to provide **33.7** (0.15 g, 96%) as a yellow colored solid. ^1H NMR (500 MHz, $\text{DMSO}-d_6$) δ 14.20-13.91 (brs, 1H, D_2O exchangeable), 9.38 (s, 1H), 8.45 (s, 1H), 7.22 (s, 1H), 2.21 (s, 3H); m/z 205.0 $[\text{M}+1]^+$.

[0237] Compounds 33a-33b. Using 4-chloro-6-(5-(trifluoromethyl)-1H-imidazol-2-yl)pyrimidine (See WO2007076473 and WO2007076474), compound **33b** was synthesized as exemplified in Scheme 33.



Scheme 34.




[0238] **Synthesis of Compound 34.1.** To a stirred solution of 4-methyl-3-nitrobenzoic acid (500mg, 2.76 mmol) in DMF (10 mL), HOBT (560mg, 4.14 mmol), EDCI (794mg, 4.14 mmol) and 4-chloro-3-trifluoromethyl-aniline (540mg, 2.76 mmol) were added at 0 °C, and the reaction mixture was stirred for 6 hr at room temperature. The reaction mixture was diluted with water (50 mL) and extracted with EtOAc (2x50mL). The combined organic layers was washed with water, dried over anhydrous Na₂SO₄ and concentrated under reduced pressure. The residue was purified by column chromatography (SiO₂, 100% Hexane then gradient to 12% EtOAc/hexane) to afford compound **34.2** (700 mg, 70.7%) as light yellow solid. ¹H NMR (DMSO-D₆, 200 MHz) δ 10.8 (s, 1H, D₂O exchangeable), 8.6 (s, 1H), 8.5 (s, 1H), 8.2-8.3 (d, 1H), 8.1-8.2 (d, 1H), 7.6-7.8 (m, 2H), 2.6 (s, 3H); LCMS *m/z* 358.9 [M+1]⁺.

[0239] **Synthesis of Compound 34.2.** To a stirred solution of **34.1** (550mg, 1.553 mmol) in ethanol (50 mL) at room temperature, tin(II)chloride (1.38g, 6.133 mmol) was added and the reaction mixture was refluxed for 2 hr. The solvent was concentrated under vacuum. The residue was dissolved in EtOAc, washed with 2N NaOH and brine, dried over anhydrous Na₂SO₄, and concentrated to afford **34.2** (400mg, 79.36%) as a yellow solid. ¹H NMR (DMSO-D₆, 200 MHz) δ 10.2 (s, 1H, D₂O exchangeable), 8.4 (s, 1H), 8.05-8.15 (d, 1H), 7.6-7.65 (d, 1H), 7.2 (s, 1H), 7.1 (s, 1H), 5.05 (s, 2H, D₂O exchangeable), 2.1 (s, 3H); LCMS *m/z* 328.9 [M+1]⁺.


[0240] **Synthesis of Compound 34a.** To a stirred solution of **34.2** (250 mg, 0.76 mmol) in DMF (10 mL), HOBT (154mg, 1.14 mmol), EDCI (218mg, 1.14 mmol) and compound **10** (132mg, 0.76 mmol) were added at 0 °C and stirred for 8 hr at room temperature. After completion, the reaction mixture was diluted with water (50 mL) and extracted with twice with

EtOAc (2x50mL). The combined organic layers were washed with water, dried over anhydrous Na_2SO_4 , and concentrated. The crude compound obtained was purified by column chromatography (SiO_2 , 100% DCM then gradient to 2% MeOH/DCM) to afford **34a** (300 mg, 81.52%) as off white solid. ^1H NMR (DMSO- D_6 , 500 MHz) δ 10.6 (s, 1H, D_2O exchangeable), 10.3 (s, 1H, D_2O exchangeable), 8.4 (2s, 2H), 8.15 (m, 2H), 7.8 (m, 1H), 7.75 (m, 1H), 7.5 (m, 1H), 2.3 (s, 3H); LCMS m/z 484.26 $[\text{M}+1]^+$.


[0241] Compounds 34a-34h. Using different amines, the following compounds can be synthesized as exemplified in Scheme 34:



[00131] Compounds 34i. Using 4-methyl-3-nitro-aniline and 4-chloro-3-trifluoromethylbenzoic acid, the following compounds can be synthesized as exemplified in Scheme 34.
























[00132] **Compounds 34j-34k.** Using compound **O.7** or **N.6**, the following compounds can be synthesized as exemplified in Scheme **34** and Schemes **O** and **N**.
























[00133] In certain embodiments, the compound of formula for use in preparing compounds of the present invention is selected from those set forth in **Table 1**, below.


Table 1. Exemplary Compounds





















































































**12a****12b****12c****13a****13b****14a****14b****14c****14d****15a****15b****15c****15d****15e****16****17****18a****18b****18c****18d****18e****18f****18g****18h****18i****18j****18k****18l**



Synthesis of -L¹-Cy¹-L²-Cy² Moieties

(1) Thiazole condensation

Scheme A-1.



[0242] **Synthesis of Compound A.1.** To an ice cold solution of 2-aminoacetamide (100 g, 0.90 mol) in water/ dioxane (1200 mL, 1: 1), CbzCl (130 mL, 0.90 mol) was added slowly. The reaction was brought to RT and stirred at RT for 12 hr. Dioxane was removed under reduced pressure and the reaction mixture was filtered and air-dried to obtain compound A.1 as a white solid (167.0 g, 88%). ¹H NMR: (CDCl₃-DMSO-d₆, 200 MHz) δ: 7.4 (s, 5H), 6.8 (1H, D₂O exchangeable), 6.2 (1 H, D₂O exchangeable), 6.1 (1 H, D₂O exchangeable), 5.1 (s, 2 H), 3.8 (d, 2 H, *J* = 5 Hz); LCMS: *m/z* 209.3 [M+]⁺.


[0243] **Synthesis of Compound A.2.** To a solution of compound A.1 (0.5 g, 0.0024 mol) in dioxane (7 mL) was added Lawesson's reagent (0.5 g, 0.0013 mol). The reaction was heated at 60 °C for 30–45 min. The reaction was brought to RT and stirred for an additional 4 hr. Dioxane was removed under reduced pressure. The reaction mixture was diluted with EtOAc (3 mL) and the organic layer was washed with sat. NaHCO₃ (2 mL). The aqueous layer was again extracted with EtOAc (2 x 5 mL). The combined organic extracts were again washed with sat.

NaHCO₃ (3 x 5 mL), dried (Na₂SO₄) and concentrated under reduced pressure to furnish compound **A.2** as a light yellow solid (0.42 g, 79%). ¹H NMR: (CDCl₃-DMSO-d₆, 200 MHz) δ: 7.4 (s, 5 H), 6.4 (1 H, D₂O exchangeable), 5.2 (s, 2 H), 4.2 (d, 2 H, *J* = 5 Hz); LCMS: *m/z* 224.9 [M+1]⁺.

[0244] Synthesis of Compound A.3. Ethyl chloroacetate (50 g, 0.409 mol) and ethyl formate (30.3g, 0.409 mol) were taken in anhydrous toluene (500 mL) and cooled to 0 °C. NaOEt (33g, 0.485 mol) was added portion wise. The reaction mixture was stirred at 0 °C for 5 hr and then at RT for 12 hr. The reaction mixture was quenched with water (250 mL) and washed with Et₂O (2 x 250 mL). The aqueous layer was cooled to 0 °C and acidified to pH 4 using 5N HCl. The aqueous layer was extracted with Et₂O (3 x 300 mL). The combined organic layers were dried (Na₂SO₄) and concentrated under reduced pressure to obtain compound **A.3** as light brown oil (54 g, 88%), which was used without further purification.

[0245] Synthesis of Compound A.4. To a solution of aldehyde **A.3** (54 g, 0.36 mol) in anhydrous DMF (42 mL), was added a solution of compound **A.2** (40.3 g, 0.18 mol) in anhydrous DMF (320 mL). The reaction was heated at 50 °C for 3 days. The mixture was cooled to 0°C, and Et₂O (390 mL) followed by sat. NaHCO₃ solution (200 mL) were added slowly. After separation of the phases, the aqueous layer was extracted with Et₂O (2 x 300 mL). The combined organic extracts were washed with sat. NaHCO₃ (3 x 500 mL), dried (Na₂SO₄) and concentrated under reduced pressure to give crude material as thick brown oil, which was purified by column chromatography (EtOAc/hexanes) to give compound **A.4** as a brown solid (22 g, 19 %). ¹H NMR: (CDCl₃, 200 MHz) δ: 8.3 (s, 1 H), 7.4 (s, 5 H), 5.6 (brs, 1 H), 5.2 (s, 2H), 4.7 (d, 2 H, *J* = 5 Hz), 4.4 (m, 2 H), 1.4 (m, 3 H); LCMS: *m/z* 320.9 [M+1]⁺.

[0246] Synthesis of Compound A.5. To an ice-cold solution of compound **A.4** (10 g, 0.0311 mol) in THF/H₂O (80 mL, 1: 1) was added LiOH (2.6 g, 0.062 mol). The reaction was stirred for 3 hr, whereupon THF was removed under reduced pressure and the aqueous layer was extracted with Et₂O (2 x 50 mL). The aqueous layer was cooled to 0 °C and acidified with 3N HCl (20 mL) during which solid precipitated out. The solid was filtered, washed with water (2 x 100 mL) and dried to give compound **A.5** as a white solid (7 g, 77%). ¹H NMR: (CDCl₃-DMSO-d₆) δ 8.2 (s, 1 H), 7.4 (s, 5 H), (brs, 1 H), 5.2 (s, 2 H), 4.8 (d, 2 H, *J* = 4 Hz); ¹³C NMR: (DMSO-d₆, 60 MHz): 176.33, 162.04, 156.39, 147.62, 136.78, 130.25, 128.3, 127.7, 65.9, 42.71, 40.34; LCMS: *m/z* 292.8 [M+1]⁺.

(2) Oxalyl chloride coupling**Scheme A-2.**


[0247] Synthesis of Compound A.6. To a solution of 2-amino-4-trifluoropyridine (2.00 g, 0.0123 mol) in DMF (4 mL, 0.05 mol) was added a solution of 1,3-dichloro-5,5-dimethylhydantoin (1.4 g, 0.0074 mol) in DMF (4 mL) dropwise. The reaction was stirred at RT for 2 hr, whereupon the reaction mixture was diluted with ether (80 mL) and washed with water (10 mL). The organic phase was dried and concentrated to give the crude product, which was purified on combiflash (0–20% EtOAc/Hexanes) to give compound **A.6** as light yellow oil. (65% yield); ^1H NMR: (DMSO- d_6) δ 8.16 (s, 1 H), 6.87 (s, 1 H), 6.76 (brs, 1 H); LCMS: m/z 197 $[\text{M}+1]^+$.

[0248] Synthesis of Compound A.7. A 20 mL vial was charged with compound **A.5** (191.8 mg, 0.0006561 mol), methylene chloride (3.0 mL), a 2.0 M solution of oxalyl chloride in methylene chloride (390 μL) and DMF (10.0 μL , 0.000129 mol). The reaction mixture was stirred for 15 minutes at rt, then concentrated *in vacuo* and the resultant residue was taken up in acetonitrile (3.0 mL). To this solution was added a solution of compound **A.6** (129 mg, 0.000656 mol) and pyridine (0.5 mL, 0.006 mol) in acetonitrile (1.5 mL). The reaction mixture was stirred at RT overnight. The solvent was removed under reduced pressure, and the residue was purified by combiflash (0–30% EtOAc/ CH_2Cl_2) to give compound **A.7** in 49% yield. LCMS: m/z 471 $[\text{M}+1]^+$.

[0249] Synthesis of Compound A. A vial was charged with compound **A.7** (1.0E2 mg, 0.00021 mol), acetic acid (1.0 mL, 0.018 mol) and hydrogen bromide (300 μL , 4 M/ acetic acid). The reaction mixture was stirred at RT for 2h. The reaction mixture was diluted with methanol and concentrated under reduced pressure. The residue was diluted with aqueous NaHCO_3 and


ethyl acetate. After separation of the phases, the organic layer was washed with aqueous NaHCO₃ and brine, dried over sodium sulfate, and concentrated to give compound **A** as a light brown solid (73% yield), which was used without further purification. ¹H NMR (300 MHz, DMSO-d₆): δ 8.85 (s, 1 H), 8.79 (s, 1 H), 8.57 (s, 1 H), 4.48 (brs, 2 H). LCMS: *m/z* 337 [M+1]⁺.

Scheme B.



[0250] Synthesis of Compound B. Compound **A.5** was coupled to 4-chloro-3-trifluoromethyl-phenylamine and deprotected according to procedures described in Scheme **A.2**. ¹H NMR (400 MHz, CDCl₃): δ 8.40 (s, 1 H), 8.21 (d, *J* = 2.6 Hz, 1 H), 7.96 (dd, *J*¹ = 8.7 Hz, *J*² = 2.6, 1 H), 7.60 (d, *J* = 8.7 Hz, 1 H), 4.48 (brs, 2H); LCMS: *m/z* 336 [M+1]⁺.


Scheme C.



[0251] Synthesis of Compound C: Compound A.5 was coupled to 4-methyl-3-trifluoromethyl-phenylamine and deprotected according to procedures described in Scheme A.2.

Compound C.1. ¹H NMR: (MeOD-d₄, 400 MHz) δ: 8.3 (s, 1 H), 7.9(s, 1 H), 7.7 (d, 1 H, *J* = 8 Hz), 7.3–7.2 (m, 8 H), 5.0 (s, 2 H), 4.5 (s, 2 H), 2.4 (s, 3 H); LCMS: *m/z* 450.1 [M+1]⁺; R_f = 0.2 (50% EtOAc/hexanes). **Compound C.** LCMS: *m/z* 316.1 [M+1]⁺.

Scheme D.



[0252] As shown in Scheme **D**, using Z-alanine-NH₂ as starting material, compound **D** was synthesized following the same procedures as previously detailed in Methods 3 and 4, Schemes **A-1** and **A-2**.

[0253] **Synthesis of Compound D.1.** To a solution of Z-alanine-NH₂ (5 g, 22.5 mmol) in dioxane (100 mL) was added Lawesson's reagent (5.4 g, 13.5 mmol). The reaction was heated at 60 °C overnight. The solvent was removed under reduced pressure, the resulting residue was diluted with a 1:1 mixture of saturated aqueous NaHCO₃: H₂O (100 mL), and extracted with ethyl acetate (3 x 100 mL). The combined extracts were washed with brine (100 mL), dried over anhydrous sodium sulfate, and concentrated *in vacuo*. Purification by flash column chromatography (10–60% EtOAc / hexanes) afforded compound **D.1** (4.7 g, 90%) as a white solid. LCMS: *m/z*: 239 [M+1]⁺.

[0254] **Synthesis of Compound D.2.** Compound **D.1** was condensed with compound **A.3** according to the procedure described previously (Scheme **A-1**) to afford compound **D.2** (50% yield) as a light yellow solid. ¹H NMR (CDCl₃, 200 MHz): δ 8.3 (s, 1 H), 7.3–7.5 (m, 5 H), 5.4–5.5 (m, 1 H), 5.1 (m, 2 H), 4.3–4.4 (m, 2 H), 1.6–1.7 (d, 2 H), 1.3–1.4 (t, 3 H); LCMS: *m/z* 335 [M+1]⁺.


[0255] **Synthesis of Compound D.3.** Hydrolysis of compound **D.2** according to the procedure described previously (Scheme A-1) afford compound **D.3** (83.5% yield) as a white solid. ^1H NMR (CDCl_3 , 200 MHz): δ 8.2 (s, 1 H), 7.2–7.4 (m, 5 H), 5.1 (m, 2 H), 4.8–4.9 (m, 1 H), 1.3–1.5 (d, 2 H); ^{13}C NMR (75 MHz, $\text{DMSO}-d_6$): δ 181.12, 162.22, 155.81, 147.85, 136.89, 130.05, 128.46, 128.0, 127.89, 65.86, 20.47; LCMS: m/z 307 $[\text{M}+1]^+$.

[0256] **Synthesis of Compound D.4.** Compound **D.3** was coupled to compound **A.6** according to the procedure described previously (Scheme A-2) to afford compound **D.4** (60% yield). ^1H NMR (CDCl_3 , 200 MHz): δ 8.6 (s, 1 H), 8.4 (s, 2 H, 1 H D_2O exchangeable), 8.2 (s, 1 H), 7.2 (s, 5 H), 5.4–5.5 (m, 1 H), 5.1 (s, 2 H), 5.1 (s, 2 H), 1.7 (d, $J = 7$ Hz, 3 H); LCMS: m/z 484.9 $[\text{M}+1]^+$.

[0257] **Synthesis of Compound D.** Compound **D.4** was deprotected according to the procedure described previously (Scheme A-2) to afford compound **D** (85% yield). ^1H NMR (400 MHz, $\text{DMSO}-d_6$): δ 8.77 (s, 1 H), 8.70 (s, 1 H), 8.59 (s, 1 H), 4.22 (q, $J = 7.0$ Hz, 1 H), 1.39 (d, $J = 7.0$ Hz, 2 H); LCMS: m/z 351 $[\text{M}+1]^+$.

[0258] **Synthesis of Compound Da and Compound Db.** Compound **D** was separated by preparative chiral HPLC, using CHIRALCEL OJ column and hexane/IPA/EtOH (80:15:5) as the mobile phase to afford compound **Da** and compound **Db**.

Scheme D'.



[0259] Alternatively, compound **Da** (R) was prepared as shown in Scheme **D'**, above.

[0260] **Synthesis of Compound Diii.** To a clean dry flask was charged 21.83 g (127.5 mmols, 1.06 eq) of 2-acetylthiazole-5-carboxylic acid (Compound **Di**), 40.5 mL of 1,2-dimethoxyethane, and 42.8 mg (5 mol %) of N,N-dimethylformamide under a nitrogen atmosphere. The resulting mixture was allowed to stir at 20-30 °C while 15.85 g (123.8 mmols, 1.03eq) of oxalyl chloride was charged dropwise over 30 minutes. The resulting reaction solution was allowed to stir for at least 3 hr at 25 °C. In a separate flask was charged 28.07 g (120.5 mmols, 1 eq) of 5-chloro-4-(trifluoromethyl)pyridine-2-amine hydrochloride (Compound **Dii**), 87 mL of acetonitrile, and 29.1 mL of (360.3 mmols, 2.99 eq) pyridine under a nitrogen atmosphere. The resulting solution was cooled to 10 °C with stirring. To the cooled **Dii** solution was added the activated **Di** solution dropwise over 30 minutes. The final combined solution was allowed to warm to RT, and the stirring was continued for an additional 2 hr. This solution may be used in the next step without isolation. However, Compound **Diii** can be isolated from the solution at this point by adding water dropwise until a thick slurry is obtained.

[0261] **Synthesis of Compound Div.** The solution of **Diii**, from the procedure described above, was heated to 45 °C while maintaining stirring and a nitrogen atmosphere. To the heated solution was added 9.30 g of NH₂OH dropwise over 5 minutes. After the addition was complete, stirring was continued at 45 °C for an additional 4 hr. The reaction solution was then heated to 60 °C and 215 mL of water was added over the course of 1 hr. The resulting slurry was cooled to RT and filtered to collect the solids. The filter cake was washed with 25% v/v acetonitrile/water, then water, and dried to constant weight at ambient temperature. A total of 44.26 g of compound **Div** was produced in 98% yield. Mass spectra showed a molecular ion (M+1) of 365.01.

[0262] **Synthesis of Compound D.** To a clean dry flask was charged 11.5 g (31.5 mmols, 1 eq) of compound **Div**, 4.6 g (70.3 mmols, 2.23 eq) of zinc dust, 35 mL of water, and 57 mL of 1-butanol under a nitrogen atmosphere. While stirring vigorously, the resulting mixture was cooled to 0-5 °C. To the cold mixture was charged 10.8 mL (188.7 mmols, 6 eq) of acetic acid dropwise, while maintaining the internal reaction temperature of <10 °C. Once the addition is complete, the reaction was allowed to warm to 30 °C, and the stirring was continued for an additional 3-4 hr. After aging the reaction solution, the contents of the flask were cooled to ~5 °C, and 56 mL of NH₄OH was added dropwise while maintaining an internal temperature <10

°C. The biphasic mixture was warmed to 35 °C and the aqueous phase was removed. The organic layer was washed once more with a mixture of 24 mL of NH₄OH and 24 mL of water at 35 °C. The aqueous phase was removed and the 16 mL of heptane was added to the organic layer. The organic solution was then washed with a solution of 1.15 g of EDTA in 50 mL of water at 35 °C. The aqueous phase was removed, and the organic phase, at 35 °C, was filtered through a 4-5.5 micron filter funnel into a separate clean dry flask. To the filtered solution was added 215 mL of heptane at ambient temperature with stirring over the course of 1 hr. The slurry was cooled to 0-5 °C and held with stirring for an additional 3 hr. The solids were collected by filtration and washed with 35 mL of heptane in 2 portions. The wet solids were dried at 50 °C under high vacuum for 30 hr. Compound **D**, 8.52 g, was isolated as a pale pink solid in a 77% yield. The mass spectrum showed a molecular ion of 351.35 [M+1]⁺.


[0263] Synthesis of Compound Dv. To a clean dry flask was charged 80 g (228 mmoles, 1 eq) of Compound **D**, 263 g of 2-propanol, and 263 mL of water under a nitrogen atmosphere. The resulting mixture was heated to 53 °C and stirred until all the solids dissolved. In a separate clean dry flask was charged 59.2 g (153 mmoles, 0.67 eq) of *D*-ditoluoyl tartaric acid, 481 g of 2-propanol, and 206 g of water under a nitrogen atmosphere. The tartaric acid solution was stirred until all the solids dissolved at ambient temperature, and then added to the Compound **D** solution through a coarse filter funnel at such a rate to maintain the internal temperature of the Compound **D** solution at 45-53 °C. The coarse filter funnel was washed with an additional 40 mL of a 3:1 2-propanol : water solution. Immediately following the funnel wash, the stirring of combined solutions was stopped, and the contents of the flask were held at 45 °C for 9 hr. After aging, the reaction mixture was cooled to 20 °C, and the stirring was resumed. The contents of the flask were held at 20 °C with stirring for approximately 12 hr. The solids were then collected by filtration, and the wet solids were washed with 80 mL of a cold 2-propanol :water (3:1) solution in 2 portions. The wet solids were then dried at 50 °C under vacuum to constant weight. A total of 74.2 g of Compound **Dv** was obtained in a 88% yield.

[0264] The stereochemical purity of Compound **Dv** was further enhanced by the following procedure. To a clean dry flask was charged 66.5 g (90 mmoles, 1 eq) of Compound **Dv**, 335 g of water, and 1330 g of 2-propanol under a nitrogen atmosphere. With stirring, the contents of the flask were heated to 60 °C, and held at that temperature for 1 hr. After aging, the stirring was stopped, and the contents of the flask were cooled to 0 °C over 4 hr. During this cooling period,

the stirring was started and stopped after approximately 20 seconds 5 times over evenly spaced intervals. The contents of the flask were held at 0 °C for 2 hr without stirring. After aging, the solids were collected by filtration. The wet solids were dried at 50 °C under vacuum to constant weight. A total of 53.8 g of Compound **Dv** was obtained in a 81% yield. Mass spectral analysis (positive mode) showed a molecular ion of 351.43 [M+1]⁺.

[0265] Synthesis of Compound Da (R). To a clean dry flask was charged 156 g (217 mmoles, 1 eq) of Compound **Dv**, 1560 mL of methyl *tert*-butyl ether, and 780 mL of methanol under a nitrogen atmosphere. The contents of the flask were then stirred at ambient temperature, and a solution of 250 g (1110 mmoles, 5.26 eq) of sodium bicarbonate in 2340 mL of water was added slowly to maintain the internal temperature of ≤30 °C. The resulting mixture was stirred for an additional hr at 30 °C. After aging, the stirring was stopped and the organic and aqueous layers were allowed to separate. The aqueous layer was removed, and the organic layer was concentrated under vacuum to obtain a thick slurry. To the slurry was added 1000 mL of heptane, and the resulting mixture was cooled to 0-5 °C. The solids were collected from the cold solution by filtration. The wet solids were then dried at 50 °C under vacuum to constant weight. A total of 68.7 g of Compound **Da** was obtained in a 92% yield. Mass spectral analysis showed a molecular ion of 351.35 [M+1]⁺.

Scheme E.




[0266] Synthesis of Compound E. Compound **D.3** was coupled to 4-chloro-3-trifluoromethyl-phenylamine and deprotected according to procedures described in Scheme A-2. ¹H NMR (400 MHz, DMSO-*d*₆): δ 11.54 (s, 1 H), 9.06 (s, 1 H), 8.92 (br. s, 3 H), 8.30 (d, *J* =

Hz, 1 H), 8.05 (dd, $J = 8.8$, 2 Hz, 1 H), 7.86 (d, $J = 8.8$ Hz, 1 H), 4.91 (quintet, $J = 6$ Hz, 1 H), 1.65 (d, $J = 6.8$ Hz, 3 H). LCMS: m/z 350 $[M+1]^+$.

[0267] Synthesis of Compound E.2. To a flask containing compound **E** (10.3 mg, 0.0294 mmol) was added a solution of carbonic acid di-tert-butyl ester (17.6 mg, 0.0799 mmol) in CH_2Cl_2 (0.6 mL) at RT. Triethylamine (8 μL) was added and the reaction was stirred at RT overnight. Water and ethyl acetate were added to the reaction mixtures and the layers were separated. The aqueous layer was extracted once more with ethyl acetate. The combined organic layers were dried over anhydrous sodium sulfate and concentrated *in vacuo*. Purification by column chromatography (EtOAc/Hexanes) afforded compound **E.2** as a white solid (8.2 mg, 62%). $R_f = 0.1$ (100% EtOAc); LCMS: m/z : 450 $[M+1]^+$.


[0268] Synthesis of Compound Ea and Eb. Compound **E.2** was separated by preparative chiral HPLC, using CHIRALPAK AD column and hexanes/EtOH (85:15) as the mobile phase. The compounds were deprotected by treatment with 4M-hydrochloric acid in dioxane at RT to afford compound **Ea** and compound **Eb**. LCMS: m/z : 350 $[M+1]^+$.

Scheme F.




[0269] Synthesis of Compound F. Compound **D.3** was coupled to 3-chloro-4-trifluoromethyl-phenylamine and deprotected according to the procedures described in Scheme A.2. ^1H NMR (400 MHz, $\text{DMSO}-d_6$): δ 11.38 (s, 1 H), 8.96 (s, 1 H), 8.87 (br. s, 3 H), 8.42 (d, $J = 2.4$ Hz, 1 H), 8.18 (dd, $J = 9$, 2.6 Hz, 1 H), 7.73 (d, $J = 9$ Hz, 1 H), 4.91 (br. s, 1H), 1.65 (d, $J = 6.8$ Hz, 3 H); LCMS: m/z 350 $[M+1]^+$.

Scheme G.



[0270] Synthesis of Compound G: Compound **D.3** was coupled to 3-methyl-4-trifluoromethyl-phenylamine and deprotected according to the procedures described in Scheme A-2. **Compound G.1.** ^1H NMR: (MeOD- d_4 , 400 MHz) δ : 8.3 (s, 1 H), 7.9 (s, 1 H), 7.7 (d, 1 H, $J = 8$ Hz), 7.3–7.2 (m, 8 H), 5.0 (s, 2 H), 5.0–4.9 (m, 1 H), 2.4 (s, 3 H), 1.49 (d, 1 H, $J = 4$ Hz); LCMS: m/z 464.1 $[\text{M}+1]^+$; $R_f = 0.5$ (50% EtOAc/hexanes). **Compound G.** LCMS: m/z 330.1 $[\text{M}+1]^+$.

Scheme H-1.




[0271] Synthesis of Compound H.1. In a 50 mL round-bottomed flask, pinacolone (6.2 mL, 50.0 mmol) and methoxy-bis(dimethylamino)methane (10 mL) were heated at 110 °C under nitrogen. After 18 hr, the solvent was removed under reduced pressure. The crude product was purified by flash chromatography (hexanes/EtOAc = 1:1 $\rightarrow$ 1:3) to afford compound **H.1** (5.94 g, 77%) as a yellow oil which solidified upon standing. ^1H NMR (400 MHz, CDCl_3): δ 7.56 (d, $J = 12.7$ Hz, 1 H), 5.20 (d, $J = 12.7$ Hz, 1 H), 2.92 (br s, 6 H), 1.11 (s, 9 H); LCMS: m/z 156 $[\text{M}+1]^+$.

[0272] Synthesis of Compound H.2. To a solution of Na (74 mg, 3.22 mmol) in EtOH (21 mL) was added guanidine hydrochloride (308 mg, 3.22 mmol). The resultant suspension was stirred at RT, and after 30 min, a solution of compound **H.1** (500 mg, 3.22 mmol) in EtOH (2.1 mL) was added. The reaction was refluxed overnight under nitrogen. After 20 hr, the solvent was removed under reduced pressure. To the residue was added Et_2O and H_2O . The aqueous layer was extracted three times with Et_2O . The combined organic layers were washed with brine, dried over anhydrous sodium sulfate, filtered, and concentrated. The crude product was

purified by flash chromatography (hexanes/EtOAc = 1:1→1:3) to afford 379 mg (78%) of compound **H.2**. R_f = 0.3 (50% EtOAc/hexanes); ^1H NMR (400 MHz, MeOD-d_4): δ 8.11 (d, J = 5.38 Hz, 1 H), 6.69 (d, J = 5.38 Hz, 1 H), 1.27 (s, 9 H); LCMS: m/z 152 $[\text{M}+1]^+$.


[0273] Synthesis of Compound H.3. A solution of compound **H.2** (200 mg, 1.32 mmol) and N-chlorosuccinimide (185 mg, 1.39 mmol) in chloroform (3.4 mL) was refluxed. After 1.5 hr, sat. aq. NaHCO_3 and EtOAc were added. The aqueous layer was extracted three times with EtOAc. The combined organic layers were washed with brine, dried over anhydrous sodium sulfate, filtered, and concentrated. The crude product was purified by flash chromatography (hexanes/EtOAc = 5:1→3:1) to afford 200 mg (81%) of compound **H.3** as a white solid. ^1H NMR (400 MHz, MeOD-d_4): δ 8.02 (s, 1 H), 1.40 (s, 9 H); LCMS: m/z 186 $[\text{M}+1]^+$.

Scheme H-2.




[0274] Synthesis of Compound H. Compound **D.3** was coupled to 4-tert-butyl-5-chloropyrimidin-2-ylamine and deprotected according to procedures described in Scheme A-2. R_f = 0.2 (5% MeOH/EtOAc); LCMS: m/z 340 $[\text{M}+1]^+$.

Scheme I.



[0275] Synthesis of Compound I. Compound **D.3** was coupled to 6-tert-butylpyrimidin-4-ylamine and deprotected according to procedures described in Scheme A-2. R_f = 0.1 (5% MeOH/EtOAc); LCMS: m/z 306 $[\text{M}+1]^+$.


Scheme J-1.



[0276] Synthesis of Compound J.1. A flask was charged with 3,6-dichloropyridazine (1.49 g, 0.01 mol, 1.0 equiv), silver nitrate (0.17 g, 0.001 mol, 0.1 equiv), water (30 mL), pivalic acid (3.57 g, 0.035 mol, 3.5 equiv), and sulfuric acid (1.6 mL, 0.03 mol, 3.0 equiv). The mixture was heated to 70 °C and a solution of ammonium persulfate (2.28 g, 0.01 mol, 1.0 equiv) in water (10 mL) was added dropwise over ten minutes. The reaction was stirred at 70 °C for one hr and then cooled to RT. The reaction mixture was poured into ice water and then adjusted to pH 8 with aqueous ammonium hydroxide. The aqueous mixture was extracted with CH₂Cl₂ (2 x 250 mL). The combined organic extracts were filtered through a cotton plug, washed with aqueous 1 N NaOH (70 mL), dried over anhydrous MgSO₄ and concentrated under reduced pressure. Purification by flash column chromatography (20% EtOAc/hexanes) afforded the title compound (1.32 g, 64%) as a white solid. ¹H NMR: (CDCl₃, 400 MHz) δ: 7.5 (s, 1 H), 1.5 (s, 9 H); R_f = 0.5 (80% EtOAc/hexanes).


[0277] **Synthesis of Compound J.2.** To a solution of compound **J.1** (1.32 g, 0.006 mol) in EtOH (1 mL) was added 50% aqueous ammonium hydroxide (10 mL). The reaction mixture was stirred at 140 °C for 19 hr, then additional aqueous ammonium hydroxide (10 mL) was added and the mixture was stirred at 130 °C for one hr. After cooling to rt, the reaction mixture was concentrated under reduced pressure and the resultant residue was suspended in water. The solid was filtered, washed with water and Et₂O, and dried to afford compound **J.2** as a peach solid (0.27 g, 23%). ¹H NMR: (CDCl₃) δ 7.01 (s, 1 H), 1.5 (s, 9 H); LCMS: *m/z* 186.1 [M+1]⁺.

Scheme J-2.



[0278] **Synthesis of Compound J:** Compound **D.3** was coupled to compound **J.2**, 5-tert-butyl-6-chloro-pyridazin-3-ylamine, and deprotected according to procedures described in Scheme A-2. **Compound J.3.** LCMS: m/z 474.1 $[M+1]^+$; R_f = 0.4 (50% EtOAc/hexanes). **Compound J.** LCMS: m/z 340.1 $[M+1]^+$.


Scheme K.



[0279] **Synthesis of Compound K:** Compound **D.3** was coupled to compound **K.1**, 4-fluoro-3-trifluoromethyl-phenylamine, and deprotected according to procedures described in Scheme A.2. **Compound K.2.** R_f = 0.2 (50% EtOAc/hexanes); LCMS: m/z 468 $[M+1]^+$. **Compound K.** R_f = 0.1 (100% EtOAc); LCMS: m/z 334 $[M+1]^+$.

(3) Isoxazole synthesis


Scheme L-1.



[0280] **Synthesis of Compound L.1.** (2-Oxo-ethyl)-carbamic acid tert-butyl ester (1.0 g, 6.28 mmol), hydroxylamine hydrochloride (647 mg, 9.31 mmol) and pyridine (5 mL) were dissolved in methanol (40 mL) and the reaction was stirred at RT overnight. Solvent was removed at reduced pressure and the reaction was partitioned between chloroform and water. The aqueous layer was extracted with chloroform (2 x). The combined organic layers were dried over anhydrous sodium sulfate. Removal of solvent under reduced pressure afforded crude **L.1** which was used without further purification.

[0281] **Synthesis of Compound L.2.** To a solution of **L.1** (~1.2 g, ~6.28 mmol) in DMF (35 mL) was added *N*-chlorosuccinimide (1.05 g, 7.86 mmol) at RT. The reaction mixture was heated at 60 °C for one hr. The reaction mixture was cooled to 0°C and propynoic acid ethyl ester (1.8 mL, 17.8 mmol) was added. Triethylamine (1.06 mL, 7.61 mmol) in DMF (8 mL) was added dropwise over 30 minutes. The reaction mixture was slowly allowed to warm to RT. The reaction mixture was diluted with ethyl acetate and water. The layers were separated and the aqueous layer was extracted with ethyl acetate (2 x). The combined organic layers were washed with water followed by brine and dried over anhydrous sodium sulfate. After removal of the solvent under reduced pressure the crude material was purified by silica gel column chromatography (ethyl acetate/hexane) to afford **L.2** (1.68 g, 86%). ¹H NMR (400 MHz, CDCl₃): δ 6.93 (s, 1 H), 5.02 (br, 1 H), 4.42 (s, 2 H), 4.41 (q, 2 H, *J* = 6.9 Hz), 1.45 (s, 9 H), 1.39 (t, 3 H, *J* = 6.9 Hz); LCMS: *m/z* 271 [M+1]⁺.


[0282] **Synthesis of Compound L.3.** Compound **L.2** (1.68 g, 6.22 mmol) was dissolved in THF (20 mL) at 0 °C. Aqueous lithium hydroxide (1M-solution, 6.5 mL, 6.5 mmol) was added and the reaction was stirred for one hr. THF was removed under reduced pressure and the reaction mixture was washed with hexanes. The reaction mixture was acidified using 3N-hydrochloric acid and extracted with chloroform (3 x). The combined organic layers were dried over anhydrous sodium sulfate. Upon removal of solvent under reduced pressure, crude **L.3** was obtained (743 mg, 49%) which was used without further purification. LCMS: *m/z* 243 [M+1]⁺.

(4) HATU coupling**Scheme L-2.**

[0283] **Synthesis of Compound L.4.** Compound L.3 (51.0 mg, 0.211 mmol) and 4-methyl-3-trifluoromethyl-phenylamine (33 μ L, 0.230 mmol) were dissolved in DMF (1 mL) at RT. HATU (98.0 mg, 0.258 mmol) and triethylamine (74 μ L, 0.531 mmol) were added and the reaction mixture was stirred at RT overnight. Ethyl acetate and water were added to the reaction mixture and the layers were separated. The aqueous layer was extracted with ethyl acetate (2 x) and the combined layers were dried over anhydrous sodium sulfate. Upon removal of the solvent under reduced pressure, the crude L.4 was obtained as a white solid, which was used without further purification. LCMS: m/z 400 [M+1]⁺.


[0284] **Synthesis of Compound L.** Compound L.4 (<0.211 mmol) was dissolved in 20% TFA in dichloromethane (1 mL) at 0 °C. The reaction was allowed to warm to RT over one hr. Benzene was added and the solvents were removed under reduced pressure. The reaction mixture was dissolved in dichloromethane and saturated sodium bicarbonate solution was added. After separation of the phases, the aqueous layer was extracted with dichloromethane (2 x). The combined organic layers were dried over anhydrous sodium sulfate. The solvent was removed under reduced pressure and the crude L obtained was used without further purification. LCMS: m/z 300 [M+1]⁺.

Scheme M-1.




[0285] Synthesis of Compound M.2 and Compound M.3: As shown in Scheme M-1, using (1*R*)-(1-methyl-2-oxo-ethyl)-carbamic acid tert-butyl ester as starting material, compounds **M.2** and **M.3** were synthesized following the same procedures as previously detailed in Schemes L-1 and L-2. **Compound M.2.** This compound was prepared using a procedure described for compound L.2. ^1H NMR (400 MHz, CDCl_3): δ 6.88 (s, 1 H), 4.97 (br, 1 H), 4.41 (q, 2 H, $J = 7.4$ Hz), 1.53 (d, 3 H, $J = 4.9$ Hz), 1.44 (s, 9 H), 1.39 (t, 3 H, $J = 7.4$ Hz); LCMS: m/z 285 $[\text{M}+1]^+$. **Compound M.3.** This compound was prepared using a procedure described for compound L.3 in scheme L-1 and the product was used without further purification. LCMS: m/z 225 $[\text{M}+1]^+$.

Scheme M-2.




[0286] Synthesis of Compound Ma. Compound **M.3** was coupled to 4-methyl-3-trifluoromethyl-phenylamine and deprotected according to procedures described in Scheme L-2. LCMS: m/z 314 $[\text{M}+1]^+$.

Scheme M-3.




[0287] **Synthesis of Compound M.6 and M.7:** As shown in Scheme M-3, using (1*S*)-(1-Methyl-2-oxo-ethyl)-carbamic acid tert-butyl ester as starting material, compound **Mb** was synthesized following the same procedures as previously detailed in Schemes L-1 and L-2. **Compound M.6.** This compound was prepared using the procedure described for compound L.2. $^1\text{H NMR}$ (400 MHz, CDCl_3): δ 6.88 (s, 1 H), 4.97 (br, 1 H), 4.41 (q, 2 H, $J = 7.4$ Hz), 1.53 (d, 3 H, $J = 4.9$ Hz), 1.44 (s, 9 H), 1.39 (t, 3 H, $J = 7.4$ Hz); LCMS: m/z 285 $[\text{M}+1]^+$. **Compound M.7.** This compound was prepared using the procedure described for compound L.3 in scheme L-1 and the product was used without further purification. LCMS: m/z 225 $[\text{M}+1]^+$.

Scheme M-4.



[0288] **Synthesis of Compound Mb.** Compound **M.7** was coupled to 4-methyl-3-trifluoromethyl-phenylamine and deprotected according to procedures described in Scheme L-2. LCMS: m/z 314 $[\text{M}+1]^+$.


(5) Isoxazole regioisomer synthesis**Scheme N-1.**

[0289] Synthesis of Compound N.1. To a cooled ($-78\text{ }^{\circ}\text{C}$) solution of oxalyl chloride (90 mL, 1.03 mol) in CH_2Cl_2 was added dropwise a solution of DMSO (100 mL, 1.41 mol) in CH_2Cl_2 . The mixture was stirred at $-78\text{ }^{\circ}\text{C}$ for 1 hr, and a solution of (*R*)-tert-butyl 1-hydroxypropan-2-ylcarbamate (90 g, 0.51 mol) in CH_2Cl_2 was added. After stirring for 3 hr, 500 mL of triethylamine was added and the reaction mixture was stirred for another 3 h at -78°C . The reaction was quenched with 1% HCl and the reaction mixture was warmed to RT. The organic layer was separated and the aqueous layer was extracted with CH_2Cl_2 . The organic layer was washed with water, dried over MgSO_4 , and evaporated to provide crude **N.1**, (*R*)-tert-butyl 1-oxopropan-2-ylcarbamate (76.0 g, 85.4%). ^1H NMR (CDCl_3) δ 9.56 (s, 1 H), 4.23 (br s, 1 H), 1.45 (s, 9 H), 1.32 (s, 3 H).

[0290] Synthesis of Compound N.2. A solution of zinc (135 g, 2.08 mol), PPh_3 (545 g, 2.08 mol) and CBr_4 (682 g, 4.08 mol) in CH_2Cl_2 (2 L) was stirred at $0\text{ }^{\circ}\text{C}$ for 1.5 hr. A solution of (*R*)-tert-butyl 1-oxopropan-2-ylcarbamate (114 g, 0.66 mol) in DCM was added in one portion, and the reaction mixture was stirred at $0\text{ }^{\circ}\text{C}$ for another 3 hr. The mixture was quickly passed through a silica gel, and the solvent was evaporated to give the crude (*R*)-tert-butyl 4,4-dibromobut-3-en-2-ylcarbamate. To a cooled ($-78\text{ }^{\circ}\text{C}$) solution of the crude compound (*R*)-tert-butyl 4,4-dibromobut-3-en-2-ylcarbamate in THF (2 L) was added dropwise 2.5 M BuLi (0.75 L, 1.88 mol) under nitrogen. The reaction was quenched with water and the organic layer was separated. The aqueous layer was extracted with ethyl acetate. The organic layers were combined, washed with water, dried over MgSO_4 , filtered and concentrated to give the crude compound **N.2**, (*R*)-tert-butyl but-3-yn-2-ylcarbamate, which was used without further purification. ^1H NMR (CDCl_3) δ 4.47 (br s, 1 H), 2.24 (s, 1 H), 1.49 (s, 9 H), 1.27 (s, 3 H).

[0291] Synthesis of Compound N.4. To a stirred solution of (*R*)-tert-butyl but-3-yn-2-ylcarbamate (262.5 g, 1.56 mol) and (*Z*)-ethyl 2-chloro-2-(hydroxyimino)acetate (78.2 g, 0.52 mol) in DMF (1 L) was added dropwise Et₃N (216 mL, 1.56 mol) at 90 °C. The mixture was stirred for 5 hr, and then concentrated *in vacuo*. The residue was re-dissolved in ethyl acetate. The ethyl acetate solution was washed with water, dried over Na₂SO₄, and evaporated to provide the crude compound (*R*)-ethyl 5-(1-(tert-butoxycarbonylamino)ethyl)isoxazole-3-carboxylate. To a solution of (*R*)-ethyl 5-(1-(tert-butoxycarbonylamino)ethyl)isoxazole-3-carboxylate in THF (2 L) was added aqueous 2.5 N LiOH (1 L) at RT. The mixture was stirred for 1 hr, and then evaporated under reduced pressure to remove THF. The residue was partitioned between water (1 L) and ethyl acetate (0.5 L). The organic layer was separated and the aqueous layer was extracted with ethyl acetate twice. The aqueous layer was adjusted to pH 2 with 10% HCl and extracted with ethyl acetate (2 x 1 L). All the organic layers were combined, washed with water, dried over Na₂SO₄, filtered and concentrated under reduced pressure. The residue was dried under vacuum to give the crude product **N.4**, (*R*)-5-(1-(tert-butoxycarbonylamino)ethyl)isoxazole-3-carboxylic acid (55.2 g, 44.8%), which was used without further purification. ¹H NMR (CDCl₃) δ 6.57 (s, 1 H), 4.12 (q, 1 H), 1.56 (d, 3 H), 1.37 (s, 9 H).

Scheme N-2.




[0292] Synthesis of Compound N.5. To a suspension of glycine (300 g, 4 mol) in ethanol (1500 mL) was added dropwise SOCl₂ at -5 °C. After the addition was complete, the mixture was heated to reflux and stirred for 3 hr. The reaction mixture was cooled to 0 °C, and methyl *t*-butyl ether (500 mL) was added. The resultant suspension was filtered and the filter cake was washed with methyl *t*-butyl ether and dried under vacuum to provide the pure compound **N.5**, ethyl 2-aminoacetate (482 g, 86.7%) as a white solid. ¹H NMR (D₂O) δ 4.21 (q, 2 H), 3.84 (s, 2 H), 1.21 (t, 3 H).

[0293] Synthesis of Compound N.3. To a solution of compound ethyl 2-aminoacetate (30.0 g, 0.24 mol) in water (50 mL) and 36% HCl (36 mL) was added dropwise a solution of

NaNO₂ in water (100 mL) at -5 °C. The reaction mixture was extracted with ethyl acetate. The organic layer was dried over MgSO₄, filtered and concentrated to give compound **N.3**, (*Z*)-ethyl 2-chloro-2-(hydroxyimino)acetate (17.4 g, 42.1%). ¹H NMR (DMSO-d₆) δ 13.41 (s, 1 H), 4.25 (q, 2 H), 1.24 (t, 3 H).

Scheme N-3.




[0294] Synthesis of Compound N.6. A mixture of 2-nitro-4-(trifluoromethyl)phenylamine (240 g, 1.16 mol) and Raney Ni (10 g) in methanol (2400 mL) was stirred at RT under hydrogen (50 psi) overnight. The reaction mixture was filtered and concentrated to provide the compound **N.6** (197.7 g, 96.4%). ¹H NMR (CDCl₃) δ 6.98 (d, 1 H), 6.93 (s, 1 H), 6.71 (d, 2 H).

[0295] Synthesis of Compound N.7. To a solution of (*R*)-5-(1-(tert-butoxycarbonylamino)ethyl)-isoxazole-3-carboxylic acid (55 g, 0.215 mol) and Et₃N (36 mL, 0.26 mol) in THF (2 L) was added dropwise isobutyl chloroformate (33 mL, 0.26 mol) at -20 °C. The reaction mixture was stirred for 1 hr, and a solution of 4-(trifluoromethyl)benzene-1,2-diamine (45.4 g, 0.26 mol) in THF was added. After stirring for 2 h at -20°C, the mixture was allowed to warm up to RT and stirred for another 2 hr. Water was added to quench the reaction and the reaction mixture was evaporated under reduced pressure to remove THF. The aqueous layer was extracted with ethyl acetate (2 x). The combined organic layers were washed with water, dried over Na₂SO₄, filtered and concentrated. The residue was re-dissolved in acetic acid (250 mL) and stirred for 2 hr at 90°C. The solution was concentrated under vacuum and partitioned with ethyl acetate and water. The organic layer was separated, washed with water, Na₂CO₃ solution and brine, dried over Na₂SO₄, filtered and concentrated. The crude product was purified by column chromatography to afford compound **N.7**, (*R*)-tert-butyl 1-(3-(6-

(trifluoromethyl)-1H-benzo[d]imidazol-2-yl)isoxazol-5-yl)ethylcarbamate (75.7 g, 88.8%). ¹H NMR (DMSO-d₆) δ 7.8 (m, 4 H), 6.9 (s, 1H), 4.91 (m, 1 H), 1.46 (d, 3 H), 1.39 (s, 9 H).


[0296] Synthesis of Compound Na. A mixture of (*R*)-tert-butyl 1-(3-(6-(trifluoromethyl)-1H-benzo[d]imidazol-2-yl)isoxazol-5-yl)ethylcarbamate (86.5 g, 0.22 mol) in TFA (300 mL) was stirred at RT for 2 hr. The reaction mixture was concentrated *in vacuo* and re-dissolved in ethyl acetate. The ethyl acetate solution was washed with K₂CO₃ and water, dried over Na₂SO₄, and concentrated. The crude product was purified by column chromatography to afford compound **Na**, (*R*)-1-(3-(6-(trifluoromethyl)-1H-benzo[d]imidazol-2-yl)isoxazol-5-yl)ethanamine (30.2 g, 46.7%). ¹H NMR (DMSO-d₆) δ 7.98 (s, 1 H), 7.78 (d, 1 H), 7.56 (d, 1 H), 6.94 (s, 1 H), 4.16 (q, 1 H), 1.36 (d, 3 H).

Scheme N-4.



[0297] Synthesis of Compound Nb. This compound was synthesized in the same manner as described for compound **Na** in schemes **N-1** ~ **N-3** starting from (1*S*)-(1-methyl-2-oxo-ethyl)-carbamic acid tert-butyl ester. ¹H NMR (DMSO-d₆) δ 7.98 (s, 1 H), 7.78 (d, 1 H), 7.56 (d, 1 H), 6.94 (s, 1 H), 4.16 (q, 1 H), 1.36 (d, 3 H).

Scheme O.



[0298] Synthesis of Compound O.1. Pivalonitrile (13 g, 157 mmol) was dissolved in absolute ethanol (50 mL) and cooled in a salt-ice bath. HCl gas was bubbled through this solution for 1 h to saturate the solution. The reaction was warmed to RT. After 3 hr, the solvent was removed *in vacuo* to afford ethyl pivalimidate (16 g, 62%) as white solid. The crude ethyl pivalimidate (16 g, 97 mmol) was taken up in absolute ethanol (20 mL) and absolute ethanol saturated with ammonia (30 mL) was added. The reaction mixture was stirred at RT for 3 hr, whereupon ammonium chloride was filtered off and the salt washed with ethanol. The filtrate was concentrated *in vacuo* and the solid obtained was dried under vacuum to afford compound **O.1**, pivalimidamide (10 g, 76%). ¹H NMR (DMSO-*d*₆, 200 MHz): δ 8.6 (br s, 1 H), 1.2 (s, 9 H); LCMS *m/z* 101 [M+1]⁺.

[0299] Synthesis of Compound O.2. Sodium metal (15g, 0.65 moles) was added to dry toluene and the mixture was heated to 120 °C. Ethanol (38 mL, 0.847g) was added dropwise through an addition funnel, and the mixture was refluxed for 3 hr after the addition. The reaction was cooled to RT and dry ether (400 mL) was added. To the resultant suspension, a mixture of ethyl formate (45 mL, 75 mmol) and ethyl acetate (54.7 mL, 88 mmol) were added dropwise.

The reaction was stirred at RT for 3 days. Solvent was evaporated and the obtained solid **O.2**, sodium (*E*)-3-ethoxy-3-oxoprop-1-en-1-olate (60 g, 67%), was used without further purification.

[0300] **Synthesis of Compound O.3.** A mixture of **O.1** (25 g, 182 mmol), **O.2** (50 g, 363 mmol) and 5% aqueous sodium hydroxide (320 mL) was stirred at RT overnight. The reaction mixture was brought to pH ~5.0 with conc. HCl and the product was extracted with DCM (3 x). The combined organic layers were dried (Na₂SO₄) and concentrated *in vacuo*. The resultant crude residue was purified by column chromatography to obtain compound **O.3**, 2-tert-butylpyrimidin-4(3H)-one, as a yellow solid (15 g, 54%). ¹H NMR (CDCl₃, 200 MHz) δ: 12.2 (brs, D₂O exchangeable, 1 H), 8.0 (d, *J* = 6.9 Hz, 1 H), 6.3 (d, *J* = 6.9 Hz, 1 H), 1.4 (s, 9 H); LCMS: *m/z* 153 [M+1]⁺.

[0301] **Synthesis of Compound O.4.** Compound **O.3** (10 g, 66 mmol) was taken up in concentrated sulfuric acid (64 mL) and heated to 110 °C. To the reaction mixture at 110 °C, concentrated nitric acid (64 mL) was added dropwise in four equal portions. After 70% conversion, the reaction mixture was poured into ice water and extracted (DCM). The organic layer was dried (Na₂SO₄) and concentrated *in vacuo* to afford compound **O.4**, 2-tert-butyl-5-nitropyrimidin-4(3H)-one, as a white solid (5.0 g, 39%). ¹H NMR (CDCl₃, 200 MHz) δ: 12.0 (br s, 1 H), 9.0 (s, 1 H), 1.4 (s, 9 H); LCMS *m/z* 198 [M+1]⁺.

[0302] **Synthesis of Compound O.5.** A solution of compound **O.4** (12 g, 60.9 mmol) in phosphorus oxychloride (96 mL) was stirred at reflux for 5 hr. The reaction mixture was cooled to RT and the excess phosphorus oxychloride was concentrated *in vacuo*. The residue was added to ice-water and extracted into DCM. The organic layer was dried (Na₂SO₄) and removed *in vacuo* to afford compound **O.5**, 2-tert-butyl-4-chloro-5-nitropyrimidine, as a brown liquid (12 g, 92%) which was used without further purification.

[0303] **Synthesis of Compound O.6.** To a stirred solution of compound **O.5** (12 g, 55.7 mmol) in methanol (96 mL) was added ammonium hydroxide solution (156 mL) at 0–5 °C. The reaction was warmed to RT and stirred overnight. The mixture was concentrated *in vacuo*, and the residue was dissolved in water and extracted with DCM. The organic layer was dried (Na₂SO₄) and concentrated *in vacuo* to afford compound **O.6**, 2-tert-butyl-5-nitropyrimidin-4-

amine, as a light green solid (8.4 g, 77%). ¹H NMR (CDCl₃, 200 MHz) δ 9.2 (s, 1 H), 7.8 (br. s, 1 H), 6.0 (br. s, 1 H), 1.38 (s, 9 H); LCMS: *m/z* 197.0 [M+1]⁺.


[0304] Synthesis of Compound O.7. To a stirred solution of compound **O.6** (8.0 g, 40 mmol) in methanol (200 mL) was added 10% palladium carbon (1.0g). The reaction was stirred under an atmospheric pressure of hydrogen for 6 h at RT. The mixture was filtered through celite and the solution was concentrated *in vacuo* to afford compound **O.7**, 2-tert-butylpyrimidine-4,5-diamine, as an off-white solid (6.7 g, 98.96%). ¹H NMR: (CDCl₃, 200 MHz) δ 7.8 (s, 1 H), 4.7 (br. s, 2 H), 3.0 (br. s, 2 H), 1.35 (s, 9 H); ¹³C NMR: (CDCl₃, 60 MHz) δ 167.9, 155.9, 138.4, 125.2, 38.9, 30.2; LCMS: *m/z* 167.1 [M+1]⁺.

[0305] Synthesis of Compound O.8. To a three-neck round-bottom flask equipped with a thermometer, a magnetic stirrer and a nitrogen inlet was added ethyl acetate (50.0 mL), and CDI (9.7 g, 59.9 mmol) at RT. To the resultant slurry was added a solution of compound **N.4**, 5-(1-tert-butoxycarbonylamino-ethyl)-isoxazole-3-carboxylic acid (15.7 g, 60 mmol) in ethyl acetate (80.0 mL) at RT over 1 hr. The clear solution was heated to 40 °C for additional 10 min. The reaction was cooled to RT and to it was added a solution of compound **O.7** (10.0 g, 59.9 mmol) in DMF (20 mL) over 30 min. The reaction mixture was stirred at RT for an additional 5 hr, whereupon ethyl acetate (150 mL) was added. The mixture was washed with water (3 x 110 mL) and the organic layer was concentrated under reduced pressure to give compound **O.8**, (*R*)-tert-butyl 1-(3-(4-amino-2-tert-butylpyrimidin-5-ylcarbamoyl)isoxazol-5-yl)ethylcarbamate, as a glassy solid (25.7 g, 91.2%). ¹H NMR (CDCl₃, 200 MHz) δ: 8.3 (s, 1 H), 8.2 (s, 1 H), 6.65 (s, 1 H), 5.1–5.2 (m, 1 H), 1.6 (d, 3 H), 1.4 (s, 9 H), 1.3 (s, 9 H); LCMS: *m/z* 405.2 [M+1]⁺.

[0306] Synthesis of Compound O.9. To a three-neck round-bottom flask equipped with a thermometer, a magnetic stirrer and a nitrogen inlet was added compound **O.8** (17.6 g, 37.4 mmol) and methanol (60.0 mL) at RT. To the resultant clear solution was then added acetyl chloride (16.5 mL, 232 mmol) while maintaining the reaction temperature below 40 °C. The solution was stirred at RT for an additional 1 to 2 hr, whereupon ethyl acetate (95 mL) was added. The product started to crystallize from the reaction mixture and additional ethyl acetate (265 mL) was added over 1 hr. The resultant slurry was stirred for additional 1 h and filtered. The wet cake was washed with ethyl acetate (3 x 50 mL) and dried under vacuum to give compound **O.9** (13.11 g, 92 %) as a white solid. ¹H NMR (DMSO-d₆, 400 MHz) δ 10.64 (s, 1

H), 9.19 (br s, 3 H), 8.83 (s, 1 H), 7.17 (s, 1 H), 4.83 (br. s, 1 H), 1.64 (d, $J = 7$ Hz, 3 H), 1.41 (s, 9 H); LCMS: m/z 305.3 $[M+1]^+$.

Scheme P-1.



[0307] Synthesis of Compound P.1. 1-(1-methylcyclopropyl)ethanone (8 g, 81.5 mmol) and methoxybis(N,N-dimethyl)methane (14 g, 16.2 ml, 106.0 mmol) were heated at 110 °C for 18 hr. Excess methoxybis(N,N-dimethyl)methane was removed by concentration *in vacuo* to obtain compound **P.1** as yellow crystals (11.1g, 88.2%). ^1H NMR (CDCl_3 , 200 MHz) δ : 7.60 (d, $J = 11.3$ Hz, 1 H), 5.20 (d, $J = 11.3$ Hz, 1 H), 1.4 (s, 3 H), 1.1–1.2 (m, 2 H), 0.7–0.8 (m, 2 H); LCMS: m/z 154.2 $[M+1]^+$.


[0308] Synthesis of Compound P.2. In a 350 mL sealed flask (2-nitroethene-1,1-diyl)bis(methylsulfane) (15 g, 90 mmol) was dissolved in 7M ammonia in methanol (150 mL) and stirred at 50 °C overnight. After 18 hr, solvent was removed *in vacuo* and the solid obtained was washed with DCM to afford **P.2** as an orange solid (7.2g, 76.9%). ^1H NMR ($\text{DMSO}-d_6$, 200 MHz) δ : 6.6 (s, 1 H).

[0309] Synthesis of Compound P.3. Compound **P.1** (8.0 g, 52.3 mmol) and compound **P.2** (5.38 g, 52.3 mmol) were dissolved in AcOH:EtOH (1:4). The reaction mixture was heated at 100 °C for 16 hr, then cooled to RT and concentrated *in vacuo*. The resultant residue was dissolved in 1 M NaOH and extracted with ethyl acetate (3 x). The combined organic layers were washed with brine, dried over anhydrous sodium sulfate and concentrated *in vacuo*. The crude product was purified by column chromatography (50–100% DCM/hexane) to afford

compound **P.3** (4.8 g, 47.6%). ^1H NMR (CDCl_3 , 200 MHz): δ 8.25 (d, $J = 8.5$ Hz, 1 H), 6.6–6.7 (d, $J = 8.5$ Hz, 1 H), 1.5 (s, 3 H), 1.2–1.3 (m, 1 H), 0.8–0.9 (m, 1 H); LCMS: m/z 194.1 $[\text{M}+1]^+$.

[0310] Synthesis of Compound P.4. Compound **P.3** (5.0 g, 25.9 mmol) was dissolved in methanol (200 mL) and palladium/C (1.0 g) was added. The reaction mixture was stirred under an atmospheric pressure of hydrogen for 4 hr and filtered through Celite[®]. The filtrate was concentrated *in vacuo* to provide a residue which was purified by column chromatography (2% methanol/DCM) to obtain compound **P.4** (2 g, 47.4%). ^1H NMR: (CDCl_3 , 200 MHz) δ 6.85 (d, $J = 8.5$ Hz, 1 H), 6.7–6.8 (brs, $J = 8.5$ Hz, 1 H), 4.1–4.3 (br s, 2 H, NH), 3.1–3.3 (brs, 2 H, NH), 1.4 (s, 3 H), 1.0–1.1 (m, 2 H), 0.6–0.8 (m, 2 H); ^{13}C NMR (CDCl_3 , 60 MHz): δ 154.03, 148.50, 125.75, 123.08, 111.17, 23.24, 19.65, 15.80; LCMS: m/z 164.2 $[\text{M}+1]^+$.


Scheme P-2.



[0311] Synthesis of Compound P.5. Compound **N.4** (1 g, 0.004 mol) was dissolved in DMF (30 mL). Compound **P.4** (0.64 g, 0.004 mol), HATU (2.4 g, 0.006 mol), and diisopropylethylamine (3.0 mL, 0.02 mol) were added and the reaction mixture was stirred at RT for 1 hr. Solvent was removed *in vacuo* and the crude reaction mixture was dissolved in EtOAc and washed with saturated aqueous NaHCO_3 (3 x) and brine (1 x). The organic layer was dried over anhydrous sodium sulfate and concentrated *in vacuo*. The crude product was purified by column chromatography (0–5% MeOH/DCM) to afford compound **P.5** (1.28 g, 80%). ^1H NMR ($\text{DMSO}-d_6$, 200MHz): δ 9.89 (s, 1 H, NH), 7.64 (d, $J = 7.6$ Hz, 1 H, NH), 7.39 (d, $J = 6.6$ Hz, 1 H), 6.62 (s, 1 H), 6.59 (d, $J = 7.6$ Hz, 1 H), 5.64 (br s, 1 H), 4.91–4.84 (m, 1 H), 1.44 (s, 3 H), 1.49–1.39 (m, 12 H), 1.08 (dd, $J = 3.4$ Hz, $J = 2.6$ Hz, 2 H), 0.68 (dd, $J = 3.4$ Hz, $J = 2.6$ Hz, 2 H); LCMS: m/z 402.5 $[\text{M}+1]^+$.


[0312] **Synthesis of Compound P.6.** A solution of compound **P.5** (1.0 g, 0.0025 mol) in 4 N HCl/dioxane (5 mL) was stirred for 3 hr and concentrated *in vacuo*. The resultant residue (0.65 g, 86%) was used without further purification. LCMS: m/z 302.5 $[M+1]^+$.

Scheme Q.




[0313] **Synthesis of Compound Q.1.** Compound **L.3** (73.8 mg, 0.305 mmol), compound **N.6** (59.5 mg, 0.338 mmol) and HATU (139.7 mg, 0.367 mmol) were dissolved in DMF (1.5 mL) at rt. Triethylamine (106 μ L, 0.761 mmol) was added and the reaction was stirred at RT overnight. The reaction mixture was diluted with ethyl acetate and water was added. The layers were separated and the aqueous layer was extracted twice more with ethyl acetate. The combined organic layers were dried over anhydrous sodium sulfate and concentrated under reduced pressure. The crude material was purified using silica gel column chromatography (ethyl acetate/hexanes) to afford the coupled product in quantitative yield. This compound was dissolved in acetic acid (1 mL) and the reaction was stirred at 80 °C for one hr. After cooling, acetic acid was removed under vacuum and the crude product was purified using silica gel column chromatography (ethyl acetate/hexanes) to afford compound **Q.1** (85.4 mg, 73%). LCMS: m/z 383 $[M+1]^+$.

[0314] **Synthesis of Compound Q.** Compound **Q.1** (85.4 mg, 0.223 mmol) was dissolved in 20% TFA in dichloromethane (1 mL) at 0 °C and the reaction mixture was gradually warmed to RT over one hr. Benzene was added and the solvents were removed under reduced pressure. The resultant residue was dissolved in dichloromethane and saturated sodium bicarbonate solution was added. The layers were separated and the aqueous layer was extracted twice more with dichloromethane. The combined organic layers were dried over anhydrous sodium sulfate and concentrated under reduced pressure to afford compound **Q** which was used without further purification. LCMS: m/z 283 $[M+1]^+$.


Scheme R-1.

[0315] Synthesis of Compound R. This compound was synthesized in a similar manner as compound **Q** following Scheme **Q** using compound **M.3** instead of **L.3**. LCMS: m/z 297 [M+1]⁺.

Scheme R-2.

[0316] Synthesis of Compound Rb. This compound was synthesized in a similar manner as compound **Q** following scheme **Q** using compound **M.7** instead of **L.3**. LCMS: m/z 297 [M+1]⁺.

Scheme S.



[0317] Synthesis of Compound S.2. To **S.1** (10g, 0.0969 mol) in THF (60 ml) and water (60 mL) at 0°C was added sodium bicarbonate (16.27g, 0.193 mole) followed by N-(benzyloxy carbonyloxy) succinimide (60.37g, 0.242 mol). The reaction mixture was stirred at RT for 12 hr. The THF was removed under vacuum and the aqueous phase was washed with ether (2x100 mL). The aqueous phase was cooled to 0 °C and acidified to pH=2 with 5N HCL (50 mL). The reaction mixture was extracted with ethyl acetate (2x 100 mL); the combined organic layer was dried over sodium sulfate and concentrated under reduced pressure. The crude material was purified by column chromatography (1% MeOH in dichloromethane) to give **S.2** (16g, 72%). ^1H NMR (CDCl_3 , 200 MHz) δ 7.45-7.32 (m, 5H), 5.40 (bs, 1H,) 5.12 (s, 2H), 1.82 (s, 6H); LCMS: m/z 238 $[\text{M}+1]^+$.

[0318] Synthesis of Compound S.3. To a suspension of **S.2** (20g, 0.0843 mol) in acetonitrile were added (400 mL), di-tert-butyl-dicarbonate (24 mL, 0.107 mol), ammonium bicarbonate (8 g, 0.101 mol) and pyridine (5.2 ml). The reaction mixture was stirred at RT for 3h and then the acetonitrile was removed under reduced pressure. The reaction mixture was diluted with water (50 mL) and the resulting solid was removed by filtration. The solid was washed with water and dried to afford **S.3** (12 g, 63%) as a off-white solid. This material was used for the

next reaction with out any further purification. ^1H NMR (CDCl_3 , 200 MHz) δ 7.41-7.38 (m, 5H), 6.30 (bs, 1H), 5.40 (bs, 2H), 5.15 (s, 2H), 1.78 (s, 6H); LCMS: m/z 236 $[\text{M}+1]^+$.

[0319] Synthesis of Compound S.4. Lawessons reagent (10.28g, 0.0254 mol) was added to a suspension of **S.3** (10g, 0.04237 mol) in dioxane (58 mL) at RT. The reaction mixture was heated at 60°C for 30 minutes, cooled to RT and stirred for additional 1.5 hr. The resulting solution was concentrated under reduced pressure and the residue was diluted with saturated sodium bicarbonate (50 mL). The solid obtained was filtered, washed with water and dried under vacuum to afford an off-white solid **S.4** (8.0g, 75%) which was for the next step without further purification. ^1H NMR (CDCl_3 , 200 MHz) δ 7.90 (bs, 1H) 7.72 (bs, 1H) 7.41-7.7.38 (m, 5H), 5.58 (bs, 1H), 5.12 (s, 2H), 1.72 (s, 6H). LCMS: m/z 253 $[\text{M}+1]^+$.

[0320] Synthesis of Compound S.5. A solution of **A.3** (9.5 g, 0.0635 mol) in DMF (64 mL) was added to thioamide **S.4** (8 g, 0.031 mol). The reaction mixture was stirred at 50 °C under nitrogen atmosphere overnight. After cooling to rt, ether (70 mL) was added. The solution was cooled to 0°C and saturated sodium bicarbonate (30 mL) was added slowly. The reaction mixture was extracted with ether (2 x 50 mL); the combined organic layer was washed with saturated sodium bicarbonate (1 x 50 mL), dried over sodium sulfate and concentrated under vacuum to give a brown oil. Purification by column chromatography (20% ethyl acetate / hexane) provided compound **S.5** (6g, 54%) as a brown solid. ^1H NMR (CDCl_3 200 MHz) δ 8.13 (s, 1H) 7.40-7.35 (m, 5H) 5.70 (bs, 1H), 5.10 (s, 2H), 4.35 (q, $J=7.2$ Hz, 2H) 1.80 (s, 6H), 1.37 (t, $J=7.2$ Hz, 3H). LCMS m/z : 349 $[\text{M}+1]^+$.


[0321] Synthesis of Compound S.6. To a 0 °C solution of **S.5** (300 mg, 0.86 mmol) in THF (4 mL) and water (4 mL) was added lithium hydroxide (200 mg, 0.0258 mol) in water (1 mL). The reaction mixture was stirred at RT for 2.5 hr and then the solvent was removed under reduced pressure. The aq. layer was washed with ether (2x 15 ml), cooled to 0 °C and acidified to pH 2 with 5N HCl. The obtained precipitate was filtered and dried to give **S.6** (180 mg, 66%). ^1H NMR (DMSO-d_6 , 200 MHz) δ 13.45 (bs, 1H), 8.20 (bs, 1H), 8.18 (s, 1H), 7.40-7.38 (m, 5H), 5.02 (s, 2H), 1.60 (s, 6H). LCMS m/z : 320.9 $[\text{M}+1]^+$.

[0322] Synthesis of Compound S.7. To a solution of **S.6** (205 mg, 0.64 mmol) in methylene chloride (4 mL) at RT was added oxalyl chloride (160 μL , 0.0019 mol) followed by the addition of DMF (50 μL) and stirred at RT for 1 hr. Separately a solution of **A.6** (132 mg,

0.000672 mol), acetonitrile (2 mL) and pyridine (520 μ L, 0.0065 mol) was stirred at RT followed by the addition of chlorotrimethylsilane (100 μ L, 0.0008 mol). The acid chloride was concentrated under reduced pressure to a tan solid and redissolved in acetonitrile (2 mL). To the acid chloride solution was added the activated aniline. After 3 hr, the reaction mixture was diluted with ethyl acetate (75 mL) and washed with dilute citric acid (50 mL), aqueous sodium bicarbonate (50 mL) and water. The organic layer was dried over sodium sulfate and concentrated to a residue which was purified by to give compound **S.7**. LCMS m/z : 498.95 $[M+1]^+$.


[0323] Synthesis of Compound S. To a solution of **S.7** (80 mg, 0.16 mmol) in acetic acid (3 mL) was added 4M hydrogen bromide in acetic acid (1 mL, 0.004 mol) and stirred at RT for 4 hr. The reaction mixture was concentrated to a residue which was triturated with saturated sodium bicarbonate. The residue was dissolved in ethyl acetate and washed with saturated sodium bicarbonate. The organic layer was dried over sodium sulfate and concentrated to provide **S**. LCMS m/z : 364.97 $[M+1]^+$.

Scheme T



[0324] Synthesis of Compound T. The synthesis of **T** was accomplished following Scheme S substituting 1-amino-cyclopropanecarboxylic acid for 2-amino-2-methyl-propionic acid (**S.1**).

Scheme U.



[0325] Synthesis of Compound U.2. To a 2 mL reaction vial was charged with **U.1** (50 mg, 0.2 mmol), 4-trifluoromethylbenzenamine (30 μ L, 0.24 mmol), MeOH (500 μ L) and 4 M of HCl


in 1,4-dioxane (5 μ L, 0.02 mmol). The mixture was heated in microwave oven for 20 min at 120 °C. This crude mixture was purified via preparatory reverse-phase HPLC, affording **U.2** (30 mg, 50%). ¹H NMR (DMSO-d₆, 400 MHz) δ : 11.2 (br s, 1H), 8.2 (s, 1H), 7.8-7.9 (d, 2H), 7.7-7.8 (d, 2H), 2.4 (s, 3H); m/z 287 [M+1]⁺.

[0326] Synthesis of Compound U.3. To a solution of **U.2** (1.0 g, 3.49 mmol) in methanol (20 mL) at 0 °C, were added pyridine (1.17 mL, 13.98 mmol) and hydroxylamine hydrochloride (485 mg, 6.99 mmol). After stirring at RT overnight, methanol was removed and the residue was diluted with water. The formed solid was collected via filtration, affording compound **U.3** (800mg, 80%). ¹H NMR (mixture of cis, trans isomers, DMSO-d₆ 200 MHz) δ : 11.4 and 11.1 (1H, -OH), 10.7-10.8 (br s, 1H), 7.8-7.9 (d, 2H), 7.8 and 7.6 (s, 1H), 7.6-7.7 (d, 2H), 2.1 and 2.2 (s, 3H); m/z 302 [M+1]⁺.

[0327] Synthesis of Compound U. To a mixture of **U.3** (800mg, 2.65 mmol) in 1:1 ethanol and acetic acid (30 mL) was added Zn powder (1g, 15.9 mmol). After stirring overnight at RT, solvents were distilled off and residue was taken in water. The solution was basified with NH₄OH, extracted into EtOAc and concentrated. Crude compound was purified by column chromatography using DCM to 2-4% MeOH in DCM as elute to afford **U** as a brown color solid (500mg, 65.61%). ¹H NMR (DMSO-d₆, 200 MHz) δ : 10.4-10.6 (br s, 1H), 7.8-7.9 (d, 2H), 7.6-7.7 (d, 2H), 7.1 (s, 1H), 4.2-4.3 (m, 1H), 1.3-1.4 (d, 3H); m/z 288 [M+1]⁺.

[0328] Synthesis of Compound Ua and Ub. Preparatory chiral SFC of compound **U** (440 mg) on a Chiralpak AS-H (2 x 25cm) with an eluant of 30% isopropanol(0.1% Et₂NH)/CO₂ at 100 bar at 60 mL/min and monitoring at 220 nM afforded and 206 mg of **Ub** (ee >99%) as the first eluting peak and 186 mg of **Ua** (ee >99%) as the second eluting peak.

Scheme V.



[0329] **Synthesis of Compound V.3.** A RT solution of V.1 (10 g, 45.45 mmol) in ethanol (100 mL) was treated with V.2 (10.26 g, 68.18 mmol, Plouvier, B.; Bailly, C.; Houssin, R.; Henichart, J. P. *Heterocycles* **1991**, 32, 693-701), and the reaction mixture was heated at reflux for 16 hr. The ethanol solvent was distilled off and the residue was dissolved in EtOAc. The organic layer was washed with sodium bicarbonate solution, water, and brine, dried over anhydrous Na₂SO₄, filtered, and concentrated under vacuum. Purification by flash column chromatography (SiO₂, 100% hexane to 12% EtOAc-Hexane) afforded V.3 as a yellow solid (10g, 69.63%). ¹HNMR (CDCl₃, 200 MHz) δ 9.3-9.4 (br s, 1H, D₂O exchangeable), 8.0 (s, 1H), 7.6-7.7 (d, 2H), 7.3-7.4 (d, 2H), 4.2-4.4 (q, 2H), 1.3-1.4 (t, 3H); m/z: 317 [M+1].


[0330] **Synthesis of Compound V.4.** A solution of V.3 (4g, 12.65 mmol) in dry DCM (60 mL) was cooled to -78 °C under a N₂ atmosphere, and treated with DIBAL-H (38 mL, 1M solution in toluene, 38 mmol). The reaction was stirred at -78 °C for 2 hr, then quenched by addition of saturated NH₄Cl solution, and slowly warmed to RT. The reaction mixture was filtered through celite, and the filter cake was washed with DCM. The organic layer was separated and dried over anhydrous Na₂SO₄, filtered, and concentrated under vacuum. Purification by flash column chromatography (SiO₂, 100% hexanes to 25% ethyl acetate-Hexane) afforded V.4 as white solid (1.8g, 52%). ¹HNMR (DMSO-D₆, 200 MHz) δ: 10.5 (s, 1H, D₂O exchangeable), 7.7-7.8 (d, 2H), 7.5-7.6 (d, 2H), 7.1 (s, 1H), 5.3 (t, 1H, D₂O exchangeable), 4.5 (s, 2H); m/z: 274.9 [M+1]⁺.

[0331] **Synthesis of Compound V.5.** A solution of V.4 (1.8g, 6.57 mmol) in toluene (30 mL) and THF (10 mL) was cooled in an ice bath at 0 °C, and treated with diphenylphosphonic azide (2.835g, 13.139 mmol) and DBU (2g, 13.139 mmol). The reaction mixture was stirred

overnight at RT. The mixture was concentrated under vacuum, and the residue was purified by flash column chromatography to obtain **V.5** (1g, 51%) as yellow solid. ^1H NMR (1H, CDCl_3 , 200 MHz) δ : 7.6-7.7 (d, 2H), 7.5-7.6 (d, 2H), 7.3 (s, 1H), 4.4 (s, 2H); m/z : 300 $[\text{M}+1]^+$.


[0332] Synthesis of Compound V. A solution of SBN-69-5 (500mg, 1.672 mmol) in THF (20 mL) and water (1 mL) was treated with triphenylphosphine (657mg, 2.508 mmol). The mixture was stirred overnight at RT. Solvents were evaporated and the residue was purified by column chromatography (SiO_2 , 100% DCM to 2.5% MeOH/DCM) to obtain the product as brown colour solid. (300mg, 65.78%). ^1H NMR: (1H, DMSO- D_6 , 200 MHz) δ : 10.4-10.6 (br s, 1H), 7.7-7.9 (d, 2H), 7.6-7.7 (d, 2H), 7.1 (s, 1H), 3.9 (s, 2H); m/z : 274 $[\text{M}+1]^+$.

Scheme W.



[0333] Synthesis of Compound W. The synthesis of **W** was accomplished following **Scheme U** substituting 3-trifluoromethylaniline for 4-trifluoromethylaniline.

Scheme X.



[0334] Synthesis of Compound X.1. The synthesis of **X.1** was accomplished following **Scheme U** substituting 1-(6-chloro-3-pyridinyl)-1-ethanone for 1-(2-chlorothiazol-5-yl)ethanone (**U.1**).


[0335] Synthesis of Compound X.2. A suspension of **X.1** (804 mg, 2.87 mmole) in 30 mL of ethanol was treated with sodium borohydride (0.217 g, 5.74 mmol), and the reaction mixture was stirred at RT for 16 hr. The mixture was concentrated to dryness and the residue was dissolved in EtOAc and H_2O . The organic layer was separated, dried over MgSO_4 , filtered, and concentrated, absorbing onto 10 g SiO_2 . Purification by flash column chromatography (40 g

SiO₂, 10% EtOAc/hexane for 5 min then gradient to 60% EtOAc/hexanes over 15 min) afforded 738 mg (91%) of **X.2** as a clear oil that slowly solidified to a white solid. LCMS, $m/z = 284$ $[M+1]^+$.


[0336] **Synthesis of Compound X.3.** A solution of **X.2** (738 mg, 2.61 mmol) in anhydrous DCM (10 mL) was cooled in an ice bath, treated with diphenylphosphonic azide (0.817 mL, 3.79 mmol) in a dropwise fashion, and stirred for 15 min. 1,8-Diazabicyclo[5.4.0]undec-7-ene (0.567 mL, 3.79 mmol) was added in a dropwise fashion. The reaction mixture was stirred in the ice bath for 1 hr, warmed to RT and stirred for 16 hr. The reaction mixture was partitioned between EtOAc and H₂O. The organic layer was dried over MgSO₄, filtered, and concentrated, absorbing onto 5 g SiO₂. Purification by flash column chromatography (40 g SiO₂, 5% EtOAc/hexane then gradient to 40% EtOAc/hexanes) yielded **X.3** (464 mg, 58%) as a yellow viscous oil. LCMS $m/z = 292$ $[M+H]^+$.

[0337] **Synthesis of Compound X.** A solution of **X.3** (463 mg, 1.51 mmol) in THF (10 mL) and H₂O (3 mL) was treated with triphenylphosphine (0.593 g, 2.26 mmol) and was heated at 60 °C for 16 hr. The reaction mixture was cooled to RT, diluted with EtOAc and extracted with 1 N HCl (2x10mL). The aqueous layer was made basic by addition of 10% NaOH and extracted with EtOAc (2x). The combined organic layers were dried over MgSO₄, filtered, and concentrated to obtain **X** (316 mg, 75%) as a viscous oil that solidified to a white solid upon standing. LCMS $m/z = 282$ $[M+H]^+$.


Scheme Y.




[0338] **Synthesis of Compound Y.** The synthesis of **Y** was accomplished following **Scheme X** substituting 4-*t*-butyl-aniline for 4-trifluoromethylaniline.

Scheme Z.


[0339] Synthesis of Compound Z. The synthesis of **Z** was accomplished following **Scheme U** and **X** substituting 1-(2-chloropyrimidin-5-yl)ethanone (*Bioorg. Med. Chem.* **2005**, 13, 3707) for 1-(2-chlorothiazol-5-yl)ethanone (**U.1**).

Scheme AA.

[0340] Synthesis of Compound AA. The synthesis of **AA** was accomplished following **Scheme U** and **X** substituting 1-(2-chloropyrazin-5-yl)ethanone (*Bioorg. Med. Chem.* **2005**, 13, 3707) for 1-(2-chlorothiazol-5-yl)ethanone (**U.1**).


Scheme BB.

[0341] Synthesis of Compound BB. The synthesis of **BB** was accomplished following **Scheme U** substituting 1-(2-chloropyridazin-5-yl)ethanone (*Bioorg. Med. Chem.* **2005**, 13, 3707) for 1-(2-chlorothiazol-5-yl)ethanone (**U.1**).

Scheme CC.

[0342] **Synthesis of Compound CC.** The synthesis of **CC** was accomplished following **Scheme U** substituting 1-(2-chloropyridazin-5-yl)ethanone (*Bioorg. Med. Chem.* **2005**, 13, 3707) for 1-(2-chlorothiazol-5-yl)ethanone (**U.1**) and 4-*t*-butylaniline for 4-trifluoromethylaniline.


Scheme DD.



[0343] **Synthesis of Compound DD.2.** Compound **DD.2** was synthesized as described in **Scheme U**. m/z 270 $[M+1]^+$.

[0344] **Synthesis of Compound DD.** To a mixture of **DD.2** (200 mg, 0.7 mmol) in THF (5 mL) was added lithium tetrahydroaluminate (90 mg, 2.0 mmol) and heated it at 70 °C for 2 hr. After cooling down to 25 °C., the mixture was quenched with ice water, followed by added 1 N NaOH. The formed solid was removed via filtration, and the filtrate was concentrated and further purified *via* preparatory reverse-phase HPLC, affording **DD** (40 mg, 20%). m/z 256 $[M+1]^+$.

Scheme EE.




[0345] **Synthesis of Compound EE.2.** To a solution (in a flame dried vial) of ethanamine, 2,2'-oxybis[N,N-dimethyl- (0.50 mL, 2.6 mmol) in tetrahydrofuran (7.0 mL) at 0 °C, was added 1.0 M of ethylmagnesium bromide in tetrahydrofuran (2.6 mL, 2.6 mmol). After stirring at 0-5 °C for 15 min, this mixture was slowly added to a solution (in a flame dried vial) of **EE.1** (350 mg, 2.0 mmol) in tetrahydrofuran (4.0 mL) at -60 °C over 10 min and the resulted mixture was further stirred at -60 °C for 8 min. The mixture was then quenched with aqueous ammonium

chloride. The aqueous layer was extracted with EtOAc. The organic layer was concentrated to afford **EE.2** as a white solid (250 mg, 74%). m/z 170 $[M+1]^+$.

[0346] Synthesis of Compound EE. Compound **EE** was synthesized as described in **Scheme U**. m/z 284 $[M+1]^+$.

Scheme FF.




[0347] Synthesis of Compound FF.2. In a 50 mL round-bottom flask, **FF.1** (0.949 g, 0.641 mmole), 2-amino-1-phenylethanone (1.10 g, 0.00641 mole), and 1-hydroxybenzotriazole (0.866 g, 0.641 mmole) were dissolved in DMF (20 mL). The mixture was treated with N-(3-dimethylaminopropyl)-N'-ethylcarbodiimide hydrochloride (1.474 g, 0.7691 mmole) and N,N-diisopropylethylamine (1.12 mL, 0.641 mmole). The yellow reaction mixture was allowed to stir at RT for 18 hr and then diluted with 200 mL of EtOAc. The organic layer was washed 2x 50 mL of water. **FF.2** precipitated as a white solid which was collected by filtration. The filtrate was washed with 50 mL brine, dried over Na_2SO_4 , and concentrated. The combined solids were titrated with Et_2O to yield 1.55 g (0.0064 mol, 91%) of **FF.2**.

[0348] Synthesis of Compound FF.3. In a 20 mL microwave reaction vial **FF.2** (1.5 g, 0.0565 mole) and ammonium acetate (0.262 g, 0.023 mole) were suspended in acetic acid (10.0 mL). The mixture was then stirred at RT for 1 hr before then heated at 175 °C for 15 min under microwave irradiation. The acetic acid was then removed *in vacuo* and the resulting residue was neutralized to pH 7 with NaHCO_3 sat (aq) 100 mL and solid in the presence of 200 mL of EtOAc. The aqueous layer was washed 2x 75 mL EtOAc. The combined organic layers were dried over Na_2SO_4 , filtered, and concentrated to yield an orange tar. Purification by flash column chromatography (SiO_2 , 50% EtOAc/Hexanes gradient to 100% EtOAc) yielded 250 mg (18%) of **FF.3**.


[0349] **Synthesis of Compound FF.** In a 5 mL microwave reaction vial **FF.3** (0.250 g, 1.02 mmole) and 5% Pd/C (0.2 g) were taken up in methanol (4 mL). The reaction was stirred under a H₂ balloon at RT for 24 hr. The mixture was filtered through celite and concentration to yield 250 mg of **FF**.

Scheme GG.




[0350] **Synthesis of Compound GG.** The synthesis of **GG** was accomplished following **Scheme U** and **Scheme X** substituting 1-(2-chloropyrimidin-5-yl)ethanone (*Bioorg. Med. Chem.* **2005**, *13*, 3707) for 1-(2-chlorothiazol-5-yl)ethanone (**U.1**) and 4-*t*-butylaniline for 4-trifluoromethylaniline.

Scheme HH.




[0351] **Synthesis of Compound HH.** The synthesis of **HH** was accomplished following **Scheme X** substituting 4-chloro-3-trifluoromethylaniline for 4-trifluoromethylaniline. LCMS $m/z = 316 [M+1]^+$.

Scheme II.




[0352] **Synthesis of Compound II.** The synthesis of **II** was accomplished following **Scheme X** substituting 3-trifluoromethylaniline for 4-trifluoromethylaniline.


[0353] **Synthesis of Compounds JJ – TT.** Compounds **JJ – TT** could be synthesized following Scheme D using the appropriately substituted aniline for compound **A.6**.



[0354] **Scheme UUa.** Compound UUa can be synthesized following Scheme M substituting 3-trifluoromethylaniline for 4-methyl-3-trifluoromethyl-phenylamine..



Scheme VV.




[0355] **Synthesis of VV.2.** A solution of **VV.1** (2 g, 0.0085 mol), dimethylamine hydrochloride (1g, 0.0127 mol), EDCI (4.0 g, 0.0212 mol), HOBT (574 mg, 0.0042 mol) and DIPEA (1.4g, 0.0110 mol) in DMF (20 ml) was stirred at 80 °C for 16 hr. The reaction mixture was diluted with water (50 ml) and extracted with ethyl acetate (3x 100 ml). The combined organic layers was washed with water (3x 50 ml), dried over Na₂SO₄ and concentrated under reduced pressure. The resulting crude material was purified by column chromatography to give **VV.2** as a brown liquid (1.4g, 63%): ¹H-NMR (CDCl₃, 200 MHz): d 8.61(s, 1H), 8.58 (s, 1H); 8.11 (s, 1H), 3.23 (s, 3H), 3.13 (s, 3H); m/z: 263 [M+1]⁺.

[0356] **Synthesis of VV.3** A solution of **VV.2** (1.3g, 0.0049 mol), sodium dithionite (3.4g, 0.0198 mol), sodium carbonate (1g, 0.0099 mol) in MeOH (13 ml) and water (13ml) was stirred at RT for 2 hr. The volatiles were removed under reduced pressure and extracted with ethyl acetate (3x 100 ml). The combined organic layers was dried over Na₂SO₄ and concentrated under reduced pressure to obtain **VV.3** as a light yellow solid (600 mg, 54.5%). ¹H-NMR (CDCl₃, 200 MHz) δ 7.0 (s, 1H), 6.90 (s, 1H), 6.80 (s, 1H), 3.23 (s, 3H), 3.13 (s, 3H); m/z: 233 [M+1]⁺.


[0357] **Synthesis of VV.4** Compound **VV.4** was synthesized as described in **Scheme D** for compound **D.4**. m/z: 521 [M+1]⁺.

[0358] **Synthesis of VV.** Compound **VV** was synthesized as described in **Scheme D** for compound **D**. ¹H-NMR (CD₃OD, 200 MHz): δ 8.58 (s, 1H), 8.21 (s, 1H), 8.0 (s, 1H), 7.56 (s, 1H), 5.40-5.38 (m, 1H), 3.23 (s, 3H), 3.13 (s, 3H), 1.80 (d, J=7.0 Hz, 2H); m/z: 387 [M+1]⁺.


[0359] **Compounds WW-YY.** Using the appropriate amine, the following amines could be synthesized as exemplified in **Scheme VV**.



[0360] **Synthesis of Compound ZZ.** The synthesis of compound **ZZ** was accomplished following **Scheme X** substituting 4-chloroaniline for 4-trifluoromethylaniline. MS m/z 248.1 $[M+1]^+$.


























[0361] **Synthesis of Compound AAA.** The synthesis of compound **AAA** was accomplished following **Scheme DD** substituting 2-chloroisonicotinamide for compound **DD.1** and 3-trifluoromethylaniline for 4-*t*-butylaniline. MS m/z 268 $[M+1]^+$.




[0362] **Synthesis of Compound BBB.** The synthesis of compound **BBB** was accomplished following **Scheme U** substituting 4-chloro-3-(trifluoromethyl)aniline for 4-trifluoromethylaniline. MS m/z 322 $[M+1]^+$.


[0363] In certain embodiments, the compound of formula $-NH_2-L^1-Cy^1-L^2-Cy^2$ for use in preparing compounds of the present invention is selected from those set forth in **Table 2**, below.

Table 2. Exemplary -NH₂-L¹-Cy¹-L²-Cy² Moieties

**U****Ua****Ub****V****W****X****Y****Z****AA****BB****CC****DD****EE****FF****GG****HH****II****JJ****KK****LL****MM****NN**



General Coupling of the Pyrimidine (“Left-Side”) and -L¹-Cy¹-L²-Cy² Moieties
Scheme ZZ.



[0364] To a solution of the acid (1.3–1.6 equiv), the amine (1 equiv), and HOBT (0.3 equiv) in DMF (50 equiv) was added N-(3-dimethylaminopropyl)-N'-ethylcarbodiimide hydrochloride (1.5 eq.) and diisopropylethylamine (1.0 equiv). If the amine was used as a salt at least one additional equivalent of diisopropylamine was added. The reaction mixture was stirred at RT for 3–16 hr, monitored by LCMS. After the reaction is completed, the solution was diluted with EtOAc, washed with water and brine. The solvent was removed from the organic phase, and the residue purified on flash column chromatography (EtOAc/Hexanes or MeOH/CH₂Cl₂ as eluents) or reverse phase preparative HPLC (mobile phase: acetonitrile/water, buffered with 0.1% TFA or 0.1% formic acid) to give the desired product. In the case of a chiral final product, the chiral purity was monitored by chiral HPLC using Chiralcel OC or OJ-H column (mobile phase: ethanol/hexane buffered with 0.1% diethylamine).

[0365] In an alternative method, a clean dry flask was charged with the acid (1.05 equiv), the amine (1.00 equiv), and HOBT (0.20 equiv) under a nitrogen atmosphere. To the flask was then added DMF (22.65 equiv) and the mixture was stirred at 25 °C until all solids dissolved, or 30 minutes. To the solution/slurry was then added 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide hydrochloride (EDC) (1.05-1.15 equiv) as a solid in portions to keep the internal temperature of the flask below 35 °C. The reaction mixture was stirred at 25 °C for 2-3 hr, and monitored by LCMS. After the reaction is completed, the solution was diluted with 1-butanol (9.59 equiv) and the contents of the flask were heated to 60 °C. To the hot solution was then added water (486.7 eq) dropwise to initiate crystallization. The solids were then collected by filtration and washed 3 times with water. The wet cake was then charged back to a clean dry flask under nitrogen. To the solids was added water (194 to 292 equiv) with stirring. The solids were slurried for 3 hr, and then collected by filtration. The wet cake was washed with water 3 times, and dried at 50 °C under vacuum to constant weight. (In the case of a chiral final product, the chiral purity was monitored by chiral HPLC using Chiralcel OC, OC-H or OJ-H column (mobile phase: ethanol/hexane buffered with 0.1% diethylamine).

[0366] In some instances, an additional chemical transformation(s) was performed after amide formation. In those instances the following procedures were utilized.





[0367] **General THP deprotection conditions.** To a 0 °C solution of the THP protected alcohol in MeOH was added catalytic p-toluenesulfonic acid and the reaction mixture was stirred for 1 hr. Solid NaHCO₃ was added and MeOH was removed under reduced pressure. The

reaction mixture was diluted with water and extracted with CH₂Cl₂. The combined organic layers were dried over Na₂SO₄, concentrated under reduced pressure to provide the desired alcohol.

[0368] General azole cyclization conditions. Procedure used in the preparation of benzimidazoles and similar derivatives. A solution of the amino amide (0.1 mmol) and acetic acid (2 mL, 40.0 mmol) was heated in the microwave for 30 minutes at 170 °C. The solvent was removed and the solid was triturated with MeOH to afford the desired azole which could be purified by crystallization or column chromatography.

[0369] The following compounds of the present invention, set forth in **Table 3**, below, were prepared by general coupling Scheme **ZZ** described above.

Table 3. Exemplary Compounds of Formula I

#	Structure	Characterization Data
1aD		<i>m/z</i> 550 [M+1] ⁺ ; ¹ H NMR (200 MHz, DMSO- <i>d</i> ₆): δ 11.78 (bs, N-H), 9.53 (s, 1H), 9.49 (s, 1H), 8.77 (s, 1H), 8.74 (s, 1H) 8.56 (s, 1H), 8.43 (s, 1H), 7.63 (bs, 1H), 5.33 (q, <i>J</i> = 7.6 Hz, 1H), 3.50–3.41 (m, 4H), 1.57 (d, <i>J</i> = 6.8 Hz, 3H).
1bD		<i>m/z</i> 564 [M+1] ⁺
1cD		<i>m/z</i> 619 [M+1] ⁺
1dD		<i>m/z</i> 604 [M+1] ⁺







#	Structure	Characterization Data
1eB		m/z 620 $[M+1]^+$
1fB		m/z 596 $[M+1]^+$
1gB		m/z 604 $[M+1]^+$; 1H NMR (400 MHz, methanol- d_4): δ 8.53 (s, 1H), 8.45 (s, 1H), 8.22 (d, J = 2.6 Hz, 1H), 7.95 (dd, J = 9.0, 2.6 Hz, 1H), 7.60 (d, J = 9.0 Hz, 1H), 4.08 (brs, 2H), 4.00-3.96 (m, 2H), 3.82-3.67 (m, 6H), 3.51-3.47 (m, 2H), 3.28-3.20 (m, 2H).
1hB		m/z 618 $[M+1]^+$
1iD		m/z 548 $[M+1]^+$
1jD		m/z 520 $[M+1]^+$
1jDa		m/z 520 $[M^+ + 1]$; 1H NMR: (DMSO- d_6 , 400MHz) δ : 11.8 (s, 1H, NH), 9.5 (d, J = 7.9 Hz, 1H), 8.79 (s, 1H), 8.77 (s, 1H), 8.59 (s, 1H), 8.48 (s, 1H), 7.79-7.75 (m, 1H), 5.40-5.32 (m, 1H), 2.94 (d, J = 4.9 Hz, 3H), 1.61 (d, J = 6.9 Hz, 3H).







#	Structure	Characterization Data
1kD		m/z 534 [M+1] ⁺
1lB		m/z 559 [M+1] ⁺
1mB		m/z 545 [M+1] ⁺
1nB		m/z 561 [M+1] ⁺
1oD		m/z 589 [M+1] ⁺
1pD		m/z 577 [M+1] ⁺
1qD		m/z 591 [M+1] ⁺

#	Structure	Characterization Data
1rB		m/z 534 $[M+1]^+$
1rD		m/z 549 $[M+1]^+$
1rNa		m/z 595 $[M+1]^+$
2aD		m/z 583 $[M+1]^+$; 1H NMR δ 9.66 (d, NH), 8.77 (s, 1H), 8.76 (d, J = 9.0 Hz, 2H), 8.57 (s, 1H), 8.48 (s, 1H), 8.47 (s, 1H), 7.83 (s, 1H), 7.82 (s, 1H), 5.39 (q, J = 7.5 Hz, 1H), 1.60 (d, J = 7.0 Hz, 3H)
2bD		m/z 583 $[M+1]^+$
2cD		m/z 601 $[M+1]^+$
2dD		m/z 601 $[M+1]^+$
2eD		m/z 613 $[M+1]^+$








#	Structure	Characterization Data
2fD		m/z 613 [M+1] ⁺
3aD		m/z 521 [M+1] ⁺ ; ¹ H NMR δ 8.72 (s, 1H), 8.64 (s, 1H), 8.59 (s, 1H), 8.54 (s, 1H), 5.50 (q, J = 7.0 Hz, 1H) 4.15 (s, 3H), 1.70 (d, J = 7.0 Hz, 2H).
3bD		m/z 578 [M+1] ⁺
3cD		m/z 592 [M+1] ⁺
4aD		m/z 534 [M+1] ⁺
4aDa		m/z 534 [M+1] ⁺ ; ¹ H NMR (400 MHz, methanol-d ₄): δ 9.37 (bs, 1H), 8.70 (d, J = 8 Hz, 2H), 8.61 (s, 1H), 8.51 (s, 1H), 8.47 (s, 1H), 8.43 (s, 1H), 8.21 (d, J = 8 Hz, 2H), 5.53 (q, J = 8 Hz, 1H), 1.71 (d, J = 8 Hz, 1H).
4aNa		m/z 480 [M+1] ⁺ ; ¹ H NMR (400 MHz, Methanol-d ₄): δ 9.52 (s, 1H), 8.94 (brs, 2H), 8.72 (s, 1H), 8.62 (brs, 2H), 7.98 (s, 1H), 7.82 (d, J = 8.5 Hz, 1H), 7.62 (dd, J = 8.5, 1.5 Hz, 1H), 7.02 (s, 1H), 5.65 (q, J = 7.0 Hz, 1H), 1.82 (d, J = 7.0 Hz, 3H).








#	Structure	Characterization Data
4bD		m/z 534 [M+1] ⁺
4bDa		m/z 534 [M+1] ⁺ ; ¹ H NMR (400 MHz, methanol-d ₄): δ 9.55 (s, 1H), 9.47 (s, 1H), 8.96 (d, <i>J</i> = 8.0 Hz, 1H), 8.86 (d, <i>J</i> = 8.0 Hz, 1H), 8.73 (s, 1H), 8.62 (s, 1H), 8.59 (s, 1H), 8.55 (s, 1H), 7.88 (m, 1H), 5.64 (q, <i>J</i> = 8.0 Hz, 1H), 1.83 (d, <i>J</i> = 8.0 Hz, 1H)
4bO		m/z 470 [M+1] ⁺ ; ¹ H NMR (400 MHz, DMSO-d ₆): δ 9.80 (s, 1H), 9.49 (s, 1H), 9.48 (s, 1H), 9.16 (s, 1H), 8.84 (d, <i>J</i> = 4.0 Hz, 1H), 8.75 (dd, <i>J</i> = 8.0, 1.5 Hz, 1H), 8.66 (s, 1H), 7.70 (dd, <i>J</i> = 8.0, 4.0 Hz, 1H), 7.08 (s, 1H), 5.65 (q, <i>J</i> = 7.0 Hz, 1H), 1.82 (d, <i>J</i> = 7.0 Hz, 3H), 1.42 (s, 9H)
4cD		m/z 552 [M+1] ⁺
4cDa		m/z 551 [M+1] ⁺ ; ¹ H NMR (400 MHz, CDCl ₃ -d ₄): δ 9.43 (s, 1H), 8.74 (m, 2H), 8.71 (s, 1H), 8.67 (s, 1H), 8.41 (s, 1H), 8.34 (s, 1H), 7.44 (m, 1H), 5.67 (m, 1H), 1.84 (d, <i>J</i> = 8.0 Hz, 1H)
4dD		m/z 568 [M+1] ⁺
4cD		m/z 523 [M+1] ⁺







#	Structure	Characterization Data
4fD		m/z 619 [M+1] ⁺
4gD		m/z 551 [M+1] ⁺
4hD		m/z 552 [M+1] ⁺
4iD		m/z 564 [M+1] ⁺
4jD		m/z 564 [M+1] ⁺
4kD		m/z 552 [M+1] ⁺








#	Structure	Characterization Data
4lD		m/z 570 [M+1] ⁺
4mD		m/z 564 [M+1] ⁺
4nD		m/z 569 [M+1] ⁺
4oD		m/z 552 [M+1] ⁺
4pD		m/z 550 [M+1] ⁺
4qD		m/z 535 [M+1] ⁺








#	Structure	Characterization Data
4qDa		m/z 535 [M+1] ⁺
4rD		m/z 550 [M+1] ⁺
4sD		m/z 549 [M+1] ⁺
4tD		m/z 573 [M+1] ⁺
5aD		m/z 585 [M+1] ⁺ ;
5aDa		m/z 585 [M+1] ⁺ ; ¹ H NMR (400 MHz, Methanol-d ₄): δ 8.62 (s, 1H), 8.58 (s, 1H), 8.58 (s, 1H), 8.51 (s, 1H), 7.39 (s, 1H), 5.53 (q, J = 7.4 Hz, 1H), 3.80 (brs, 4H), 3.72 (t, J = 6.0 Hz, 2H), 2.64 (t, J = 6.0 Hz, 4H), 2.60 (t, J = 6.0 Hz, 2H), 1.75 (d, J = 7.4 Hz, 3H).
5bD		m/z 516 [M+1] ⁺

#	Structure	Characterization Data
5cD		m/z 530 [M+1] ⁺
5dD		m/z 542 [M+1] ⁺
5dDa		m/z 542 [M+1] ⁺
5dB		m/z 527 [M+1] ⁺
5eD		m/z 570 [M+1] ⁺
5fD		m/z 599 [M+1] ⁺
5gD		m/z 555 [M+1] ⁺

#	Structure	Characterization Data
5hD		m/z 569 [M+1] ⁺
5iA		m/z 486 [M+1] ⁺
5jD		m/z 486 [M+1] ⁺
5jDa		m/z 486 [M+1] ⁺ ; ¹ HNMR: (DMSO-d ₆ , 400MHz) δ: 11.8 (s, 1H, NH), 9.5 (d, J = 7.9 Hz, 1H), 8.79 (s, 1H), 8.75 (s, 1H), 8.59-8.54 (m, 2H), 8.06-7.99 (m, 1H), 7.11 (brs, 1H), 5.42-5.38 (m, 1H), 2.89 (brs, 3H), 1.61 (d, J = 6.9 Hz, 3H).
5kD		m/z 514 [M+1] ⁺
5lA		m/z 529 [M+1] ⁺
5lD		m/z 543 [M+1] ⁺









#	Structure	Characterization Data
5mD		m/z 557 [M+1] ⁺
5nD		m/z 515 [M+1] ⁺
5oD		m/z 555 [M+1] ⁺
5qD		m/z 581 [M+1] ⁺
5pDa		m/z 569 [M+1] ⁺
5rD		m/z 595 [M+1] ⁺

#	Structure	Characterization Data
5sD		m/z 625 [M+1] ⁺
5tD		m/z 583 [M+1] ⁺
5uD		m/z 597 [M+1] ⁺
6aD		m/z 567 [M+1] ⁺
6bD		m/z 567 [M+1] ⁺
6cD		m/z 574 [M+1] ⁺
6dD		m/z 579 [M+1] ⁺








#	Structure	Characterization Data
6cD		m/z 579 [M+1] ⁺
6fB		m/z 525 [M+1] ⁺
6gD		m/z 549 [M+1] ⁺
6hD		m/z 549 [M+1] ⁺
6iD		m/z 567 [M+1] ⁺
6jD		m/z 567 [M+1] ⁺
7aD		m/z 525 [M+1] ⁺







#	Structure	Characterization Data
8aD		m/z 559 [M+1] ⁺
8bD		m/z 573 [M+1] ⁺
9A		m/z 544 [M+1] ⁺ ; ¹ H NMR (300 MHz, Methanol-d ₄): δ 9.20 (s, 1H), 8.54 (s, 1H), 8.50 (s, 1H), 8.43 (s, 1H), 7.59 (s, 2H), 4.85 (brs, 2H);
9D		m/z 558 [M+1] ⁺ ; ¹ H NMR (400 MHz, DMSO-d ₆): δ 9.76 (d, J = 8.0 Hz, 1H), 9.22 (s, 1H), 8.79 (s, 2H), 8.59 (s, 1H), 7.45 (s, 2H), 5.44 (q, J = 8.0 Hz, 1H), 1.62 (d, J = 8.0 Hz, 3H).
10A		m/z 493 [M+1] ⁺ ; ¹ H NMR (400 MHz, methanol-d ₄): δ 8.64 (s, 1H), 8.59 (s, 1H), 8.53 (s, 1H), 8.37 (s, 1H), 4.89 (d, J = 4.0 Hz).
10D		m/z 506 [M+1] ⁺
10Da		m/z 506 [M+1] ⁺ ; ¹ H NMR (400 MHz, methanol-d ₄): δ 8.64 (s, 1H), 8.59 (s, 1H), 8.54 (s, 1H), 8.36 (s, 1H), 5.52 (q, J = 8.0 Hz), 1.74 (d, J = 8.0 Hz).
10Db		m/z 506 [M+1] ⁺ ; ¹ H NMR (400 MHz, methanol-d ₄): δ 8.64 (s, 1H), 8.59 (s, 1H), 8.54 (s, 1H), 8.36 (s, 1H), 5.52 (q, J = 8.0 Hz), 1.74 (d, J = 8.0 Hz).







#	Structure	Characterization Data
10E		m/z 505 [M+1] ⁺ ; ¹ H NMR (400 MHz, Methanol-d ₄): δ 8.44 (s, 1H), 8.34 (s, 1H), 8.20 (s, 1H), 7.96 (dd, J = 8.0, 2.0 Hz, 1H), 7.60 (d, J = 8.0 Hz, 1H), 5.50 (q, J = 7.4 Hz, 1H), 1.73 (d, J = 7.4 Hz, 3H).
10F		m/z 505 [M+1] ⁺ ; ¹ H NMR (400 MHz, Methanol-d ₄): δ 8.47 (s, 1H), 8.36 (s, 1H), 8.10 (s, 1H), 7.80 (dd, J = 8.0, 2.0 Hz, 1H), 7.77 (d, J = 8.0 Hz, 1H), 5.51 (q, J = 7.4 Hz, 1H), 1.74 (d, J = 7.4 Hz, 3H).
10Na		m/z 452 [M+1] ⁺
10O		m/z 442 [M+1] ⁺
10P		m/z 439 [M+1] ⁺ ; ¹ H NMR: (DMSO-d ₆ , 400MHz) δ: 13.8 (brs, 1H, NH), 9.4 (d, J = 8.1 Hz, 1H), 8.40 (s, 1H), 8.07 (brs, 1H, NH), 7.35 (d, J = 8.1 Hz, 1H), 7.02 (d, J = 0.7 Hz, 1H), 5.44 (m, 1H), 1.63 (d, J = 7.0 Hz, 3H), 1.59 (s, 3H), 1.26 (brs, 2H), 0.90 (brs, 2H).
11D		m/z 472 [M+1] ⁺
11Da		m/z 472 [M+1] ⁺ ¹ H NMR (400 MHz, CDCl ₃): δ 8.64 (s, 1H), 8.59 (s, 1H), 8.43 (s, 1H), 8.27 (s, 1H), 7.25 (s, 1H), 5.58 (q, J = 7.4 Hz, 1H), 5.16 (brs, 1H), 1.79 (d, J = 7.4 Hz, 3H).
12aDa		m/z 540 [M+1] ⁺

#	Structure	Characterization Data
12bDa		m/z 556 [M+1] ⁺
12cDa		m/z 538 [M+1] ⁺
13aD		m/z 473 [M+1] ⁺
13bD		m/z 507 [M+1] ⁺
14aD		m/z 552 [M+1] ⁺
14aDa		m/z 552 [M+1] ⁺ ; ¹ HNMR: (DMSO-d ₆ , 200MHz) δ: 11.76 (s, 1H), 9.48 (d, J = 7.8 Hz, 1H), 8.79 (s, 1H), 8.76 (s, 1H), 8.58 (s, 1H), 8.38 (s, 1H), 5.40-5.30 (m, 1H), 1.60 (d, J = 7.4 Hz, 3H).
14aDb		m/z 552 [M+1] ⁺
14aE		m/z 551 [M+1] ⁺

#	Structure	Characterization Data
14aO		m/z 486 [M ⁺ + 1]; ¹ H NMR (300 MHz, CDCl ₃): δ 9.34 (d, <i>J</i> = 7.4 Hz, 1H), 9.15 (s, 1H), 8.35 (s, 2H, NH), 7.04 (s, 1H), 5.43-5.33 (m, 1H), 1.57 (d, <i>J</i> = 7.4 Hz, 1H), 1.41 (s, 9H).
14bD		m/z 565 [M+1]; ¹ H NMR (500 MHz, CD ₃ OD): δ 8.64 (s, 1H), 8.60 (s, 1H), 8.54 (s, 1H), 8.49 (s, 1H), 5.51 (q, <i>J</i> = 8.0 Hz, 1H), 3.08 (s, 3H) 1.74 (d, <i>J</i> = 8.0 Hz, 3H).
14cD		m/z 537 [M+1]; ¹ H NMR: (DMSO-d ₆ , 200MHz) δ: 11.79 (s, 1H), 9.75 (d, <i>J</i> = 7.8 Hz, 1H), 9.28 (s, 1H), 9.21 (s, 1H), 8.79 (s, 1H), 8.78 (s, 1H), 8.58 (s, 1H), 5.45-5.38 (m, 1H), 1.63 (d, <i>J</i> = 7.1 HZ, 3H).
15aDa		m/z 502 [M+1] ⁺
15bDa		m/z 516 [M+1] ⁺
15cDa		m/z 530 [M+1] ⁺
15dDa		m/z 556 [M+1] ⁺
16D		m/z 443 [M+1] ⁺ ; ¹ H NMR (300 MHz, DMSO-d ₆): δ 11.9 (brs, 1H), 10.19 (t, <i>J</i> = 6.5 Hz, 1H), 9.47 (s, 1H), 9.19 (d, <i>J</i> = 5.1 Hz, 1H), 8.89 (s, 1H), 8.83 (s, 1H), 8.62 (s, 1H), 8.15 (d, <i>J</i> = 5.0 Hz, 1H), 4.88(d, <i>J</i> = 6.5 Hz, 2H).







#	Structure	Characterization Data
16Na		m/z 403 [M+1] ⁺ ; ¹ H NMR (300 MHz, DMSO-d ₆): δ 9.68 (d, J = 8.4 Hz, 1H), 9.33 (s, 1H), 9.05 (d, J = 5.1 Hz, 1H), 8.03 (d, J = 5.1 Hz, 1H), 7.95 (s, 1H), 7.76 (d, J = 8.4 Hz, 1H), 7.55 (d, J = 8.4 Hz, 1H), 7.00 (s, 1H), 5.48 (m, 1H), 1.63 (d, J = 8.0 Hz, 1H).
17D		m/z 497 [M+1] ⁺
18aDa		m/z 523 [M+1] ⁺
5vDa		¹ H NMR (400MHz, DMSO-d ₆) δ 11.74 (s, 1 H), 9.49 (d, J = 8.1 Hz, 1 H), 8.78 (s, 1 H), 8.74 (s, 1 H), 8.60 (s, 1 H), 8.55 (s, 1 H), 7.32 (s, 1 H), 5.42 (quin, J = 7.3 Hz, 1 H), 4.79 (d, J = 4.5 Hz, 1 H), 4.08 (br. s., 1 H), 3.78 (td, J = 4.0, 8.1 Hz, 1 H), 3.37 (dd, J = 3.3, 9.3 Hz, 2 H), 1.86 - 1.75 (m, 2 H), 1.66 (d, J = 7.1 Hz, 3 H), 1.43 - 1.31 (m, 2 H); m/z 556 [M+1] ⁺
5wDa		¹ H NMR (400MHz, DMSO-d ₆) δ 11.75 (s, 1 H), 9.49 (d, J = 8.1 Hz, 1 H), 8.78 (s, 1 H), 8.74 (s, 1 H), 8.59 (s, 1 H), 8.55 (s, 1 H), 7.06 - 6.93 (m, 1 H), 5.42 (quin, J = 7.3 Hz, 1 H), 5.14 - 4.98 (m, 1 H), 4.51 - 4.32 (m, 1 H), 3.78 - 3.44 (m, 3 H), 2.16 - 1.84 (m, 2 H), 1.66 (d, J = 7.1 Hz, 3 H); m/z 542 [M+1] ⁺
5xDa		m/z 542 [M+1] ⁺
18jDa		m/z 550 [M+1] ⁺







#	Structure	Characterization Data
21		¹ H NMR (400MHz, DMSO-d ₆) δ 11.75 (s, 1 H), 9.54 (d, J = 7.6 Hz, 1 H), 8.78 (s, 1 H), 8.74 (s, 1 H), 8.63 (s, 1 H), 8.56 (s, 1 H), 7.39 (br. s., 1 H), 5.43 (quin, J = 7.2 Hz, 1 H), 4.44 (dd, J = 3.5, 7.6 Hz, 1 H), 3.98 (br. s., 2 H), 3.60 (br. s., 2 H), 1.93 (br. s., 2 H), 1.73 - 1.57 (m, 5 H); m/z 636 [M+1] ⁺
18cDa		¹ H NMR (400MHz, DMSO-d ₆) δ 11.76 (s, 1 H), 9.86 (d, J = 8.1 Hz, 1 H), 9.22 (s, 1 H), 8.78 (s, 1 H), 8.75 (s, 1 H), 8.73 (s, 1 H), 8.55 (s, 1 H), 8.34 (s, 1 H), 7.89 (s, 1 H), 5.49 (quin, J = 7.3 Hz, 1 H), 2.19 (s, 3 H), 1.70 (d, J = 7.1 Hz, 3 H); m/z 537 [M+1] ⁺
20aDa		¹ H NMR (400MHz, DMSO-d ₆) δ 11.76 (s, 1 H), 9.48 (d, J = 8.1 Hz, 1 H), 8.78 (s, 1 H), 8.75 (s, 1 H), 8.57 (s, 1 H), 8.25 (d, J = 2.0 Hz, 1 H), 7.64 (br. s., 2 H), 5.37 (quin, J = 7.2 Hz, 1 H), 1.62 (d, J = 7.1 Hz, 3 H); m/z 490[M+1] ⁺
22.1		¹ H NMR (400MHz, DMSO-d ₆) δ 11.76 (s, 1 H), 9.48 (d, J = 8.1 Hz, 1 H), 8.78 (s, 1 H), 8.75 (s, 1 H), 8.57 (s, 1 H), 8.25 (d, J = 2.0 Hz, 1 H), 7.64 (br. s., 2 H), 5.37 (quin, J = 7.2 Hz, 1 H), 1.62 (d, J = 7.1 Hz, 3 H); m/z 557[M+1] ⁺
22		¹ H NMR (400MHz, DMSO-d ₆) δ 11.76 (s, 1 H), 10.02 (d, J = 8.1 Hz, 1 H), 9.59 (s, 1 H), 8.78 (s, 1 H), 8.76 (s, 1 H), 8.55 (s, 2 H), 5.51 (quin, J = 7.2 Hz, 1 H), 2.68 (s, 3 H), 1.71 (d, J = 7.1 Hz, 3 H); m/z 539[M+1] ⁺
25mDa		¹ H NMR (400MHz, DMSO-d ₆) δ 11.75 (s, 1 H), 9.94 (d, J = 8.1 Hz, 1 H), 9.80 (t, J = 6.3 Hz, 1 H), 9.51 (s, 1 H), 8.78 (s, 1 H), 8.75 (s, 1 H), 8.57 (s, 1 H), 8.55 (s, 1 H), 8.50 - 8.43 (m, 2 H), 7.75 (d, J = 7.6 Hz, 1 H), 7.35 (dd, J = 4.8, 7.8 Hz, 1 H), 5.50 (quin, J = 7.2 Hz, 1 H), 4.56 (d, J = 6.1 Hz, 2 H), 1.70 (d, J = 7.1 Hz, 3 H); m/z 591 [M+1] ⁺








#	Structure	Characterization Data
10MM		m/z 459 [M+1] ⁺
10G		m/z 485 [M+1] ⁺
10K		m/z 489 [M+1] ⁺
10NN		m/z 417 [M+1] ⁺
10V		m/z 429 [M+1] ⁺
33aDa		m/z 537 [M+1] ⁺







#	Structure	Characterization Data
10FF		^1H NMR (DMSO- <i>d</i> ₆) δ 9.30 - 9.39 (m, 1H), 9.17 (d, <i>J</i> = 1.8 Hz, 1H), 8.41 (dd, <i>J</i> = 8.3, 2.3 Hz, 1H), 8.37 (s, 1H), 8.15 (s, 1H), 7.86 - 7.93 (m, 2H), 7.56 - 7.62 (m, 1H), 7.48 - 7.55 (m, 2H), 7.37 - 7.43 (m, 1H), 4.62 (d, <i>J</i> = 6.0 Hz, 2H); <i>m/z</i> = 406 [M+1] ⁺
23.5		^1H NMR (MeOH- <i>d</i> ₄) δ 8.34 (d, <i>J</i> = 8.1 Hz, 2H), 8.07 (d, <i>J</i> = 7.1 Hz, 2H), 8.00 (s, 1H), 7.63 - 7.72 (m, 1H), 7.52 - 7.60 (m, 2H), 5.46 - 5.56 (m, 1H), 3.01 (s, 2H), 2.88 (s, 2H), 1.73 (d, <i>J</i> = 7.1 Hz, 3H); <i>m/z</i> = 445 [M+1] ⁺
23		^1H NMR (MeOH- <i>d</i> ₄) δ 8.43 (br. s., 1H), 8.36 (br. s., 1H), 7.93 (s, 1H), 7.76 - 7.82 (m, 2H), 7.49 - 7.56 (m, 2H), 7.43 - 7.49 (m, 1H), 5.54 (q, <i>J</i> = 7.1 Hz, 1H), 2.66 (br. s., 1H), 1.77 (d, <i>J</i> = 7.1 Hz, 3H); <i>m/z</i> = 425 [M+1] ⁺
35		<i>m/z</i> 549 [M+1] ⁺
5tDa		<i>m/z</i> 584 [M+1] ⁺
36		^1H NMR (400MHz, MeOD) δ = 8.93 (s, 1 H), 8.76 (s, 1 H), 8.61 (s, 1 H), 8.57 (s, 1 H), 8.51 (s, 1 H), 5.55 (q, <i>J</i> = 6.9 Hz, 1 H), 2.21 (s, 3 H), 1.76 (d, <i>J</i> = 7.1 Hz, 2 H); <i>m/z</i> 472 [M+1-Ac] ⁺

#	Structure	Characterization Data
5yDa		^1H NMR (400MHz, DMSO- d_6) δ = 11.74 (s, 1 H), 9.50 (d, J = 7.6 Hz, 1 H), 8.78 (s, 1 H), 8.74 (d, J = 2.5 Hz, 1 H), 8.62 (s, 1 H), 8.55 (s, 1 H), 7.31 (s, 1 H), 5.41 (m, 1 H), 3.76 - 3.62 (m, 4 H), 3.50 (t, J = 5.6 Hz, 2 H), 3.42 (q, J = 7.1 Hz, 2 H), 3.36-3.30 (m, 4 H), 1.66 (d, J = 6.6 Hz, 3 H), 1.11 (t, J = 7.1 Hz, 3 H); m/z 614 $[\text{M}+1]^+$
5zDa		^1H NMR (400MHz, DMSO- d_6) δ = 11.75 (s, 1 H), 9.58 (d, J = 8.6 Hz, 1 H), 8.78 (s, 1 H), 8.74 (d, J = 2.5 Hz, 1 H), 8.72 (s, 1 H), 8.55 (s, 1 H), 7.52 (s, 1 H), 5.48 - 5.39 (m, 1 H), 4.18 (br. s., 4 H), 3.21 (br. s., 4 H), 1.67 (d, 3 H); m/z 591 $[\text{M}+1]^+$
5aaDa		m/z 584 $[\text{M}+1]^+$
3bB		m/z 564 $[\text{M}+1]^+$
3bC		m/z 578 $[\text{M}+1]^+$
4dB		m/z 554 $[\text{M}+1]^+$

#	Structure	Characterization Data
4cB		m/z 508 [M+1] ⁺
1rA		m/z 536 [M+1] ⁺
14dA		m/z 608 [M+1] ⁺
18dDa		¹ H NMR (400MHz, DMSO-d ₆) δ = 11.76 (s, 1 H), 9.91 (d, J = 8.1 Hz, 1 H), 9.32 (s, 1 H), 8.78 (s, 1 H), 8.76 (d, J = 2.5 Hz, 1 H), 8.55 (s, 1 H), 8.21 (s, 1 H), 7.98 (s, 1 H), 7.00 (s, 1 H), 5.54 - 5.44 (m, 1 H), 2.69 (s, 3 H), 1.71 (d, J = 7.1 Hz, 3 H); m/z 537 [M+1] ⁺
33bDa		¹ H NMR (400MHz, DMSO-d ₆) δ = 14.11 (br. s., 1 H), 11.76 (s, 1 H), 9.90 (d, J = 8.1 Hz, 1 H), 9.44 (s, 1 H), 8.78 (s, 1 H), 8.76 (d, J = 2.5 Hz, 1 H), 8.56 (s, 1 H), 8.53 (s, 1 H), 8.11 (s, 1 H), 5.63 - 5.41 (m, 1 H), 1.71 (d, J = 7.1 Hz, 3 H); m/z 591 [M+1] ⁺
10GG		m/z: 426 [M+1] ⁺







#	Structure	Characterization Data
10Z		m/z: 438 [M+1] ⁺
10X		m/z: 437 [M+1] ⁺
10Y		m/z: 425 [M+1] ⁺
10HH		m/z: 471 [M+1] ⁺
17Da		¹ H NMR (400 MHz, CHLOROFORM-d) δ 8.68 (s, 1H), 8.66 (s, 1H), 8.55 - 8.59 (m, 1H), 8.52 (br. s., 1H), 8.45 (s, 1H), 8.30 (s, 1H), 5.97 (br. s., 2H), 5.55 - 5.65 (m, 1H), 1.80 (d, J = 6.95 Hz, 3H); LCMS: m/z: 497 [M+1] ⁺
14aOO		¹ H NMR (400 MHz, CHLOROFORM-d) δ 8.79 (d, J = 2.02 Hz, 1H), 8.63 (d, J = 8.08 Hz, 1H), 8.50 (s, 1H), 8.44 (d, J = 3.41 Hz, 2H), 7.35 (ddd, J = 0.76, 2.18, 8.56 Hz, 1H), 6.94 (d, J = 8.59 Hz, 1H), 5.96 (br. s., 2H), 5.51 - 5.62 (m, 1H), 4.59 (dt, J = 2.04, 6.79 Hz, 2H), 4.31 (t, J = 4.42 Hz, 2H), 2.10 (s, 3H), 1.78 (d, 3H)

#	Structure	Characterization Data
14aPP		LCMS: m/z: 573 [M+1] ⁺
14aQQ		LCMS: m/z: 614 [M+1] ⁺
14aRR		LCMS: m/z: 589 [M+1] ⁺
10SS		LCMS: m/z: 496 [M+1] ⁺
14aK		LCMS: m/z: 534 [M+1] ⁺
14aTT		LCMS: m/z: 559 [M+1] ⁺
4uDa		LCMS: m/z: 533 [M+1] ⁺








#	Structure	Characterization Data
25aD		m/z 501 [M+1] ⁺
25cD		m/z 514 [M+1] ⁺
6kD		m/z 565 [M+1] ⁺
29b		m/z 592 [M+1] ⁺
29a		m/z 564 [M+1] ⁺
24a		m/z 530 [M+1] ⁺







#	Structure	Characterization Data
24b		m/z 554 [M+1] ⁺
25dD		m/z 528 [M+1] ⁺
37		m/z 833 [M+1] ⁺
25eD		m/z 542 [M+1] ⁺
1tD		m/z 577 [M+1] ⁺
25fD		¹ H NMR (400MHz, DMSO-d ₆) δ = 11.77 (s, 1 H), 9.96 (d, J = 8.1 Hz, 1 H), 9.54 (d, J = 1.3 Hz, 1 H), 9.33 (t, J = 6.1 Hz, 1 H), 8.78 (s, 1 H), 8.76 (s, 1 H), 8.55 (d, J = 0.6 Hz, 1 H), 8.49 (d, J = 1.3 Hz, 1 H), 8.02 - 7.66 (m, 3 H), 7.78 (br. s., 2 H), 5.80 - 5.30 (m, 1 H), 3.74 - 3.40 (m, 2 H), 3.15 - 2.91 (m, 2 H), 1.70 (d, J = 7.0 Hz, 3 H); m/z 543 [M+1] ⁺







#	Structure	Characterization Data
25gD		m/z 544 [M+1] ⁺
25hD		m/z 585 [M+1] ⁺
25iD		¹ H NMR (400MHz ,DMSO-d ₆) δ = 11.72 (br. s., 1 H), 9.93 (d, J = 8.6 Hz, 1 H), 9.50 (s, 1 H), 8.77 (s, 1 H), 8.75 (s, 1 H), 8.55 (s, 1 H), 8.47 (s, 1 H), 8.46 - 8.41 (m, 1 H), 8.10 (s, 1 H), 5.54 - 5.45 (m, 1 H), 1.70 (d, J = 7.1 Hz, 3 H); m/z 500 [M+1] ⁺
25jD		m/z 611 [M+1] ⁺
25kD		m/z 583 [M+1] ⁺
25lD		¹ H NMR (400MHz ,DMSO-d ₆) δ = 9.94 (d, 1 H), 9.76 (d, 1 H), 9.53 (s, 1 H), 8.77 (s, 1 H), 8.75 (s, 1 H), 8.55 (s, 1 H), 8.45 (s, 1 H), 8.37 (s, 1 H), 5.54 - 5.46 (m, 1 H), 4.90 - 4.80 (m, 1 H), 3.98 - 3.89 (m, 4 H), 1.70 (d, J = 7.1 Hz, 3 H); m/z 555 [M+1] ⁺

#	Structure	Characterization Data
38		m/z 591 [M+1] ⁺
25lDa		¹ H NMR (400MHz, DMSO-d ₆) δ = 9.94 (d, 1 H), 9.76 (d, 1 H), 9.53 (s, 1 H), 8.77 (s, 1 H), 8.75 (s, 1 H), 8.55 (s, 1 H), 8.45 (s, 1 H), 8.37 (s, 1 H), 5.54 - 5.46 (m, 1 H), 4.90 - 4.80 (m, 1 H), 3.98 - 3.89 (m, 4 H), 1.70 (d, J = 7.1 Hz, 3 H); m/z 555 [M+1] ⁺
25bDa		¹ H NMR (400MHz, DMSO-d ₆) δ = 11.72 (br. s., 1 H), 9.93 (d, J = 8.6 Hz, 1 H), 9.50 (s, 1 H), 8.77 (s, 1 H), 8.75 (s, 1 H), 8.55 (s, 1 H), 8.47 (s, 1 H), 8.46 - 8.41 (m, 1 H), 8.10 (s, 1 H), 5.54 - 5.45 (m, 1 H), 1.70 (d, J = 7.1 Hz, 3 H); m/z 500 [M+1] ⁺
25nDa		m/z 528 [M+1] ⁺
1tDa		m/z 577 [M+1] ⁺
25kDa		m/z 583 [M+1] ⁺

#	Structure	Characterization Data
25cDa		m/z 514 [M+1] ⁺
10LL		m/z 584 [M+1] ⁺
10EE		m/z 439 [M+1] ⁺
10EEa		m/z 594 [M+1] ⁺
10II		m/z 437 [M+1] ⁺
10DD		m/z 411 [M+1] ⁺







#	Structure	Characterization Data
10DDa		m/z 566 [M+1] ⁺
10W		m/z 443 [M+1] ⁺
15eNa		m/z 574 [M+1] ⁺
15eDa		m/z 628 [M+1] ⁺
39aNa		m/z 490 [M+1] ⁺
39aEa		m/z 544 [M+1] ⁺
39bNa		m/z 476 [M+1] ⁺







#	Structure	Characterization Data
39bEa		m/z 530 [M+1] ⁺
2eD		¹ H-NMR (DMSO-D ₆ , 500 MHz): δ 11.85 (s, 1H), 9.78 (d, <i>J</i> = 8.5 Hz, 2H), 8.86 (d, <i>J</i> = 8.0 Hz, 3H), 8.58 (s, 2H), 8.38 (s, 1H), 8.01 (d, <i>J</i> = 8.0 Hz, 1H), 7.4 (d, <i>J</i> = 8.0 Hz, 1H), 5.35- 5.25 (m, 1H), 1.67 (d, <i>J</i> = 6.0 Hz, 3H); m/z 582.7 [M+1] ⁺
18iD		¹ H-NMR (CD ₃ OD, 200 MHz): δ 8.95 (s, 1H), 8.64 (d, <i>J</i> = 8.5 Hz, 2H), 8.46 (s, 1H), 7.43 (s, 1H), 5.55- 5.45 (m, 1H), 4.01 (s, 3H), 1.77 (d, <i>J</i> = 6.0 Hz, 3H); m/z 486.8 [M+1] ⁺
6iD		¹ H-NMR (DMSO-D ₆ , 500 MHz): δ 11.78 (s, 1H), 10.62 (s, 1H), 9.78 (d, <i>J</i> = 8.5 Hz, 1H), 9.02 (s, 1H), 8.95 (s, 1H), 8.76 (d, <i>J</i> = 14.0 Hz, 2H), 8.58 (s, 2H), 7.92 (d, <i>J</i> = 6.0 Hz, 1H), 7.56 (s, 1H), 5.45- 5.43 (m, 1H), 1.72 (d, <i>J</i> = 6.0 Hz, 3H); m/z 616.5 [M+1] ⁺
18kD		¹ H-NMR (CD ₃ OD, 500 MHz): δ 8.82 (s, 1H), 8.62 (s, 1H), 8.59 (s, 1H), 8.52 (s, 1H), 7.42 (s, 1H), 5.55- 5.51 (m, 1H), 4.45 (t, <i>J</i> = 7.5 Hz, 2H), 2.52 (t, <i>J</i> = 7.5 Hz, 2H), 2.25 (s, 6H), 2.04- 2.01 (m, 2H), 1.76 (d, <i>J</i> = 6.0 Hz, 3H); m/z 557.7 [M+1] ⁺
4zD		¹ H-NMR (CDCl ₃ , 500 MHz): δ 9.18 (s, 1H), 8.62 (s, 2H), 8.5 (s, 1H), 8.42 (s, 1H), 8.35 (s, 1H), 8.18 (s, 1H), 7.02 (bs, 1H), 5.63- 5.61 (m, 1H), 4.20 (bs, 2H), 3.68- 3.65 (m, 2H), 2.65- 2.62 (m, 2H), 1.82 (d, <i>J</i> = 6.0 Hz, 3H), 1.55 (s, 9H); m/z 637.8 [M+1] ⁺







#	Structure	Characterization Data
8eD		¹ H-NMR (DMSO-D ₆ , 500 MHz): δ 11.78 (s, 1H), 9.82 (d, <i>J</i> = 7.0 Hz, 1H), 9.39 (s, 1H), 8.78 (s, 1H), 8.75 (s, 1H), 8.56 (s, 1H), 7.92 (s, 1H), 5.63- 5.61 (m, 1H), 5.49- 5.42 (m, 1H), 4.39 (s, 2H), 1.63 (d, <i>J</i> = 7.0 Hz, 3H); <i>m/z</i> 510.7 [M+1] ⁺
18ID		¹ H-NMR (CD ₃ OD, 500 MHz): δ 8.82 (s, 1H), 8.62 (s, 1H), 8.59 (s, 1H), 8.52 (s, 1H), 7.42 (s, 1H), 5.59- 5.56 (m, 1H), 4.60 (t, <i>J</i> = 7.5 Hz, 2H), 2.82 (t, <i>J</i> = 7.5 Hz, 2H), 2.39 (s, 6H), 1.76 (d, <i>J</i> = 6.0 Hz, 3H); <i>m/z</i> 543.9 [M+1] ⁺
8dD		¹ H-NMR (DMSO-D ₆ , 500 MHz): δ 11.76 (s, 1H), 9.84 (d, <i>J</i> = 7.0 Hz, 1H), 9.38 (s, 1H), 8.79 (s, 1H), 8.75 (s, 1H), 8.56 (s, 1H), 7.92 (s, 1H), 5.45- 5.41 (m, 1H), 4.58 (s, 1H), 3.54- 3.51 (m, 2H), 2.62- 2.59 (m, 2H), 1.78- 1.74 (m, 2H), 1.62 (d, <i>J</i> = 7.0 Hz, 3H); <i>m/z</i> 538.8 [M+1] ⁺
10XX		¹ H-NMR (CD ₃ OD, 500 MHz): δ 8.45 (s, 1H), 8.39 (s, 1H), 8.12 (s, 1H), 7.92 (s, 1H), 7.45 (s, 1H), 5.48- 5.41 (m, 1H), 3.78-3.73 (m, 4H), 3.64 (s, 2H), 2.49-2.45 (m, 4H), 1.77 (d, <i>J</i> = 7.0 Hz, 3H); <i>m/z</i> 569.9 [M+1] ⁺
10WW		¹ H-NMR (CD ₃ OD, 500 MHz): δ 8.45 (s, 1H), 8.39 (s, 1H), 8.12 (s, 1H), 8.09 (s, 1H), 7.45 (s, 1H), 5.48- 5.41 (m, 1H), 3.85-3.60 (m, 6H), 3.55-3.40 (m, 2H), 1.77 (d, <i>J</i> = 7.5 Hz, 3H); <i>m/z</i> 583.7 [M+1] ⁺
5bbD		¹ H-NMR (CD ₃ OD, 500 MHz): δ 8.61 (s, 1H), 8.58 (s, 1H), 8.52 (s, 1H), 7.68 (s, 1H), 7.17 (s, 1H), 7.14 (s, 1H), 6.98 (s, 1H), 5.52- 5.51 (m, 1H), 4.16 (t, <i>J</i> = 7.0 Hz, 2H), 3.46- 3.35 (m, 2H), 2.15- 2.12 (m, 2H), 1.77 (d, <i>J</i> = 6.8 Hz, 3H); <i>m/z</i> 579.7 [M+1] ⁺

#	Structure	Characterization Data
8fD		$^1\text{H-NMR}$ (CDCl_3 , 500 MHz): δ 9.20 (s, 1H), 8.63 (s, 1H), 8.57 (d, J = 7.0 Hz, 1H), 8.55 (s, 1H), 8.43 (s, 1H), 8.27 (s, 1H), 8.15 (s, 1H), 5.61- 5.58 (m, 1H), 3.77- 3.36 (m, 4H), 3.61 (s, 2H), 2.62 (bs, 4H), 1.81 (d, J = 7.0 Hz, 3H); m/z 579.9 $[\text{M}+1]^+$
27fD		$^1\text{H-NMR}$ ($\text{CDCl}_3+\text{D}_2\text{O}$, 200 MHz): δ 9.12 (s, 1H), 8.67 (s, 1H), 8.50 (s, 1H), 8.38 (s, 1H), 8.01 (s, 1H), 5.63- 5.61 (m, 1H), 3.88 (bs, 4H), 2.99- 2.97 (m, 2H), 2.75- 2.72 (bs, 5H), 2.15- 2.12 (m, 3H), 1.81 (d, J = 7.0 Hz, 3H); m/z 583.6 $[\text{M}+1]^+$
5ccD		$^1\text{H NMR}$ (400MHz, DMSO-d_6) δ = 11.74 (s, 1 H), 9.43 (d, 1 H), 8.77 (s, 1 H), 8.73 (s, 1 H), 8.55 (s, 1 H), 8.53 (br. s., 1 H), 7.97 - 7.47 (m, 1 H), 7.09 (s, 1 H), 5.73 - 5.14 (m, 1 H), 3.45 - 3.36 (m, 6 H), 1.80 - 1.71 (m, 2 H), 1.65 (d, J = 7.0 Hz, 3 H), 1.10 (t, J = 7.0 Hz, 3 H); m/z 558 $[\text{M}+1]^+$
5ddD		$^1\text{H-NMR}$ (DMSO-D_6 , 500 MHz): δ 11.72 (s, 1H), 9.40 (d, J = 8 Hz, 1H), 8.76 (s, 1H), 8.72 (s, 1H), 8.54 (s, 1H), 8.50 (s, 1H), 7.86 (bs, 1H), 7.17 (s, 1H), 5.40- 5.37 (m, 1H), 4.85 (d, J = 4.5 Hz, 1H), 4.59 (s, 1H), 3.62- 3.60 (m, 3H), 1.64 (d, J = 7.0 Hz, 3H); m/z 545.6 $[\text{M}+1]^+$
5eeD		$^1\text{H-NMR}$ (DMSO-D_6 , 500 MHz): δ 11.76 (s, 1H), 9.39 (d, J = 7.0 Hz, 1H), 8.78 (s, 1H), 8.75 (s, 1H), 8.56 (s, 1H), 8.52 (s, 1H), 7.92 (bs, 1H), 7.12 (s, 1H), 5.39- 5.35 (m, 1H), 3.39- 3.35 (m, 4H), 2.45- 2.41(m, 4H), 1.73- 1.65 (m, 9H); m/z 582.7 $[\text{M}+1]^+$
8gD		$^1\text{H-NMR}$ (CDCl_3 , 200 MHz): δ 9.20 (s, 1H), 8.63- 8.55 (s, 1H), 8.57 (d, J = 7.0 Hz, 1H), 8.43 (s, 2H), 8.28 (s, 1H), 8.14 (s, 1H), 5.65- 5.58 (m, 1H), 3.65 (s, 2H), 3.50 (bs, 4H), 2.61 (bs, 4H), 1.82 (d, J = 8.0 Hz, 3H), 1.46 (s, 9H); m/z 678.5 $[\text{M}+1]^+$
5ffD		$^1\text{H-NMR}$ (CD_3OD , 200 MHz): δ 8.62 (s, 1H), 8.59 (s, 1H), 8.52 (s, 1H), 7.18 (s, 1H), 5.52- 5.49 (m, 1H), 3.48- 3.42 (m, 2H), 2.78- 2.60 (m, 6H), 1.90-1.82(m, 2H), 1.76 (d, J = 8.0 Hz, 3H), 1.02 (t, J = 7.5 Hz, 6H); m/z 584.9 $[\text{M}+1]^+$

#	Structure	Characterization Data
27eD		¹ H-NMR (DMSO-D ₆ , 200 MHz): δ 11.78 (s, 1H), 9.75 (d, <i>J</i> = 7.0 Hz, 1H), 9.05 (s, 1H), 8.78 (s, 1H), 8.75 (s, 1H), 8.57 (s, 1H), 5.42- 5.39 (m, 1H), 4.61-4.55 (m, 1H), 3.55- 3.49 (m, 2H), 2.98 (t, <i>J</i> = 7.5 Hz, 2H), 1.95- 1.92 (m, 2H), 1.62 (d, <i>J</i> = 8.0 Hz, 3H); <i>m/z</i> 548.8 [M+1] ⁺
27gD		¹ H-NMR (CDCl ₃ , 200 MHz): δ 9.15 (s, 1H), 8.64 (bs, 2H), 8.57 (s, 1H), 8.43(s, 1H), 8.28 (s, 1H), 8.00 (s, 1H), 5.65- 5.58 (m, 1H), 3.39 (bs, 4H), 2.93 (t, <i>J</i> = 8.0 Hz, 2H), 2.38 (bs, 6H), 2.01 (bs, 2H), 1.82 (d, <i>J</i> = 8.0 Hz, 3H), 1.46 (s, 9H); <i>m/z</i> 682.9 [M+1] ⁺
10VV		¹ H-NMR (CD ₃ OD, 500 MHz): □ 8.47 (s, 1H), 8.35 (s, 1H), 8.18 (s, 1H), 8.05 (s, 1H), 7.51 (s, 1H), 5.40-5.38 (m, 1H), 3.23 (s, 3H), 3.13 (s, 3H), 1.80 (d, <i>J</i> =7.0 Hz, 2H); <i>m/z</i> 542 [M+1] ⁺
18aD		¹ H-NMR (DMSO-D ₆ , 200 MHz): δ 11.78 (s, 1H), 9.78 (d, <i>J</i> = 8.5 Hz, 1H), 9.25 (s, 1H), 8.85 (s, 1H), 8.76 (d, <i>J</i> = 12.0 Hz, 2H), 8.38 (s, 1H), 8.21 (s, 1H), 8.17 (s, 1H), 7.18 (s, 1H), 5.45- 5.43 (m, 1H), 1.72 (d, <i>J</i> = 6.0 Hz, 3H); <i>m/z</i> 523.1 [M+1] ⁺
18dD		¹ H-NMR (DMSO-D ₆ , 500 MHz): δ 11.73 (s, 1H), 9.89 (d, <i>J</i> = 8.0 Hz, 1H), 9.31 (s, 1H), 8.76 (s, 1H), 8.74 (s, 1H), 8.54 (s, 1H), 8.20 (s, 1H), 7.96 (s, 1H), 6.99 (s, 1H), 5.50- 5.47 (m, 1H), 2.68 (s, 3H), 1.70 (d, <i>J</i> = 7.0 Hz, 3H); <i>m/z</i> 536.9 [M+1] ⁺
5ggD		¹ H-NMR (CD ₃ OD, 200 MHz): δ 8.62 (s, 1H), 8.58 (s, 1H), 8.52 (s, 2H), 7.12 (s, 1H), 5.52- 5.51 (m, 1H), 3.59- 3.31 (m, 3H), 2.65- 2.4 (m, 9H), 2.29 (s, 3H), 1.77 (d, <i>J</i> = 8.0 Hz, 3H); <i>m/z</i> 611.6 [M+1] ⁺
30b		¹ H-NMR (DMSO-D ₆ , 500MHz): □ 8.87 (bs, NH), 8.25 (s, 1H), 7.25 (t, <i>J</i> = 8 Hz, 2H), 6.95 (s, 1H), 6.88 (d, <i>J</i> = 8 Hz, 1H), 6.83 (d, <i>J</i> = 8 Hz, 1H), 6.12 (s, 1H), 4.48- 4.39 (m, 1H), 3.92-3.78 (m, 2H), 3.61-3.52 (m, 2H), 3.45-3.36 (m, 2H), 2.35-1.89 (m, 4H); <i>m/z</i> : 442 [M+1] ⁺

#	Structure	Characterization Data
5hhD		¹ H-NMR (DMSO-D ₆ , 500 MHz): δ 11.78 (s, 1H), 9.46 (d, <i>J</i> = 8.5 Hz, 1H), 8.82 (s, 1H), 8.78 (s, 1H), 8.58 (bs, 2H), 7.95 (bs, 2H), 7.12 (s, 1H), 5.43- 5.38 (m, 1H), 3.45 (bs, 2H), 3.12 (bs, 2H), 1.82 (s, 3H), 1.65 (d, <i>J</i> = 7.5 Hz, 3H); <i>m/z</i> 556.8 [M+1] ⁺
27hD		¹ H-NMR (CDCl ₃ , 500 MHz): δ 9.14 (s, 1H), 8.80 (s, 1H), 8.62-8.61 (m, 2H), 8.40 (s, 1H), 8.31 (s, 1H), 7.96 (s, 1H), 5.60-5.59 (m, 1H), 3.57-3.47 (m, 8H), 3.18-10 (m, 2H), 2.98-2.97 (m, 2H), 2.26-2.25 (m, 2H), 1.79 (d, <i>J</i> = 7 Hz, 3H); <i>m/z</i> 582.8 [M+1] ⁺
18eD		¹ H-NMR (DMSO-D ₆ , 500 MHz): δ 11.71 (s, 1H), 9.76 (d, <i>J</i> = 8 Hz, 1H), 9.13 (s, 1H), 8.75-8.73 (m, 2H), 8.54 (s, 1H), 8.23 (s, 1H), 7.89 (m, 2H), 6.41 (s, 2H), 5.49-5.47 (m, 1H), 1.69 (d, <i>J</i> = 7 Hz, 3H); <i>m/z</i> 521.8 [M+1] ⁺
18fD		¹ H-NMR (DMSO-D ₆ , 500 MHz): δ 11.72 (s, 1H), 9.85 (s, 1H), 9.23 (s, 1H), 8.75-8.73 (m, 3H), 8.53 (s, 1H), 8.37 (s, 1H), 8.01 (s, 1H), 6.71 (s, 1H), 5.50-5.49 (m, 1H), 1.69 (d, <i>J</i> = 7 Hz, 3H); <i>m/z</i> 522.8 [M+1] ⁺
5jjD		¹ H-NMR (DMSO-D ₆ , 500 MHz): δ 11.70 (s, 1H), 9.59 (d, <i>J</i> = 8.5 Hz, 1H), 8.75 (s, 1H), 8.72 (s, 1H), 8.58 (s, 1H), 8.54 (s, 1H), 7.22 (s, 1H), 6.97 (d, N-H), 5.43-5.42 (m, 1H), 4.15-4.09 (m, 1H), 3.79-3.29 (m, 4H), 2.15-2.08 (m, 1H), 1.98-1.90 (m, 1H), 1.65 (d, <i>J</i> = 7 Hz, 3H), 1.38 (s, 9H); <i>m/z</i> 640.7 [M+1] ⁺
5iiD		¹ H-NMR (DMSO-D ₆ , 500 MHz): δ 11.73 (s, 1H), 9.41 (bs, 1H), 8.78 (s, 1H), 8.76 (s, 1H), 8.54 (s, 1H), 8.52 (s, 1H), 7.81 (bs, 2H), 7.1 (s, 1H), 5.42- 5.39 (m, 1H), 3.51- 3.42 (m, 2H), 3.19- 3.12 (m, 2H), 1.81 (s, 3H), 1.76 (d, <i>J</i> = 7.0 Hz, 3H); <i>m/z</i> 570.9 [M+1] ⁺







#	Structure	Characterization Data
8hD		$^1\text{H-NMR}$ (CDCl_3 , 500 MHz): δ 9.70 (bs, 1H), 9.20 (s, 1H), 8.63-8.57 (m, 3H), 8.55 (s, 1H), 8.28 (s, 1H), 8.15 (s, 1H), 5.62-5.61 (m, 1H), 3.66 (s, 2H), 3.31-3.26 (m, 4H), 2.90-2.89 (m, 4H), 1.80 (d, J = 6.5 Hz, 3H); m/z 578.7 $[\text{M}+1]^+$
19iID		$^1\text{H-NMR}$ (DMSO-D_6 , 200 MHz): δ 11.73 (s, 1H), 9.52 (d, N-H), 8.76 (s, 1H), 8.72 (s, 1H), 8.64 (s, 1H), 8.53 (s, 1H), 7.33 (s, 1H), 5.44-5.36 (m, 1H), 3.78-3.58 (m, 4H), 3.54-3.35 (m, 4H), 2.03 (s, 3H), 1.67 (d, J = 7 Hz, 3H); m/z 582.8 $[\text{M}+1]^+$
5iID		$^1\text{H-NMR}$ (CDCl_3 , 200 MHz): δ 8.63-8.58 (m, 3H), 8.43 (s, 1H), 8.39 (s, 1H), 8.27 (s, 1H), 7.35 (s, 1H), 5.63-5.52 (m, 1H), 3.80-3.72 (m, 4H), 3.60-3.46 (m, 4H), 1.78 (d, J = 7 Hz, 3H), 1.49 (s, 9H); m/z 640.7 $[\text{M}+1]^+$
18cD		$^1\text{H-NMR}$ (DMSO-D_6 , 200 MHz): δ 11.78 (s, 1H), 9.82 (d, J = 8.5 Hz, 1H), 9.12 (s, 1H), 8.78 (t, J = 12.5 Hz, 3H), 8.48 (s, 1H), 8.34 (s, 1H), 7.18 (s, 1H), 5.45-5.43 (m, 1H), 2.21 (s, 3H), 1.68 (d, J = 7.5 Hz, 3H); m/z 536.8 $[\text{M}+1]^+$
18gD		$^1\text{H-NMR}$ (DMSO-D_6 , 500 MHz): δ 11.72 (s, 1H), 9.71 (d, J = 8.5 Hz, 1H), 9.10 (s, 1H), 8.86 (s, 1H), 8.76 (s, 1H), 8.73 (s, 1H), 8.54 (s, 1H), 5.46-5.45 (m, 1H), 4.01 (t, J = 7.5 Hz, 2H), 2.64 (t, J = 7.5 Hz, 2H), 2.10 (t, J = 7.5 Hz, 2H), 1.68 (d, J = 6.5 Hz, 3H); m/z 539.7 $[\text{M}+1]^+$
19jjD		$^1\text{H-NMR}$ (CD_3OD , 500 MHz): δ 8.64 (s, 1H), 8.61 (s, 1H), 8.59 (s, 1H), 8.53 (s, 1H), 7.21 (s, 1H), 5.56-5.52 (m, 1H), 4.10-4.08 (m, 1H), 3.94-3.84 (m, 1H), 3.86-3.66 (m, 3H), 2.64-2.44 (m, 1H), 2.35-2.25 (m, 1H), 1.77 (d, J = 7 Hz, 3H); m/z 540.7 $[\text{M}+1]^+$

#	Structure	Characterization Data
18hD		¹ H-NMR (DMSO-D ₆ , 500 MHz): δ 11.74 (s, 1H), 9.85 (d, J = 8 Hz, 1H), 9.10 (s, 1H), 8.76 (s, 1H), 8.74 (s, 1H), 8.62 (s, 1H), 8.54 (s, 1H), 8.29 (s, 1H), 6.53 (d, J = 8 Hz, 1H), 5.52-5.46 (m, 1H), 2.32 (s, 3H), 1.68 (d, J = 6.5 Hz, 3H); m/z 536.7 [M+1] ⁺
5ttD		¹ H-NMR (DMSO-D ₆ , 500 MHz): δ 11.71 (s, 1H), 9.40 (bs, 1H), 8.76 (s, 1H), 8.72 (s, 1H), 8.56 (s, 1H), 8.53 (s, 1H), 7.91 (bs, 1H), 7.32 (bs, 1H), 7.10 (s, 1H), 6.82 (bs, 1H), 5.42-5.36 (m, 1H), 3.62-3.48 (m, 2H), 2.44-2.32 (m, 2H), 1.64 (d, J = 6.5 Hz, 3H); m/z 542.7 [M+1] ⁺
26aD		¹ H-NMR (DMSO-D ₆ , 200 MHz): δ 11.73 (s, 1H), 9.52 (d, N-H), 8.76 (s, 1H), 8.72 (s, 1H), 8.64 (s, 1H), 8.53 (s, 1H), 7.33 (s, 1H), 5.44-5.36 (m, 1H), 3.78-3.58 (m, 4H), 3.54-3.35 (m, 4H), 2.49 (s, 3H), 1.64 (d, J = 7 Hz, 3H); m/z 582.8 [M+1] ⁺
5ssD		¹ H-NMR (DMSO-D ₆ , 500 MHz): δ 11.71 (s, 1H), 9.45 (d, N-H), 8.75 (s, 1H), 8.72 (s, 1H), 8.57 (s, 1H), 8.54 (s, 1H), 7.01 (s, 0.5H), 6.96 (s, 0.5H), 5.42-5.39 (m, 1H), 5.06 (s, 0.5H), 4.99 (s, 0.5H), 4.43 (s, 0.5H), 4.37 (s, 0.5H), 3.68-3.50 (m, 3H), 2.04-1.90 (m, 2H), 1.65 (d, J = 7 Hz, 3H); m/z 541.7 [M+1] ⁺
5mmD		¹ H-NMR (DMSO-D ₆ , 500 MHz): δ 11.70 (s, 1H), 9.45 (d, N-H), 8.75 (s, 1H), 8.72 (s, 1H), 8.56 (s, 1H), 8.54 (s, 1H), 7.00 (s, 1H), 6.97 (s, 1H), 5.42-5.39 (m, 1H), 3.65-2.99 (m, 6H), 2.45-2.09 (m, 3H), 1.65 (d, J = 7 Hz, 3H), 1.36 (s, 9H); m/z 654.8 [M+1] ⁺
30a		¹ H-NMR (DMSO-D ₆ , 500 MHz): δ 8.59 (d, J = 7.5 Hz, 1H), 8.29 (s, 1H), 7.35 (t, J = 8 Hz, 1H), 6.96 (s, 1H), 6.93 (d, J = 8 Hz, 1H), 6.82 (d, J = 7.5 Hz, 1H), 6.07 (t, J = 5 Hz, 1H), 4.25 (d, N-H), 4.02-3.89 (m, 4H), 3.20-3.15 (m, 2H), 2.86 (t, J = 11 Hz, 2H), 1.85-1.80 (m, 2H), 1.60-1.49 (m, 2H); m/z 456.8 [M+1] ⁺

#	Structure	Characterization Data
5nnD		¹ H-NMR (DMSO-D ₆ , 200 MHz): δ 11.71 (s, 1H), 9.45 (d, N-H), 8.75 (s, 1H), 8.72 (s, 1H), 8.56 (s, 1H), 8.54 (s, 1H), 7.30 (s, 1H), 6.95 (t, N-H), 5.42-5.39 (m, 1H), 4.45-4.39 (m, 2H), 3.13-2.84 (m, 4H), 1.92-0.9 (m, 4H), 1.65 (d, J = 7 Hz, 3H), 1.36 (s, 9H); m/z 668.8 [M+1] ⁺
28a		¹ H-NMR (DMSO-D ₆ , 200 MHz): δ 11.74 (s, 1H), 9.45 (d, N-H), 8.76 (s, 1H), 8.72 (s, 1H), 8.56-8.53 (m, 2H), 7.97-7.95 (m, 2H), 7.36 (t, J = 6.8 Hz, 1H), 7.08 (s, 1H), 6.61-6.43 (m, 3H), 5.37-5.35 (m, 1H), 3.49-3.23 (m, 4H), 1.65 (d, J = 7 Hz, 3H); m/z 591.8 [M+1] ⁺
28b		¹ H -NMR (DMSO-D ₆ , 200 MHz): δ 11.70 (s, 1H), 9.39 (s, N-H), 8.76 (s, 1H), 8.72 (s, 1H), 8.56-8.53 (m, 2H), 8.00-7.75 (m, 3H), 7.11-6.97 (m, 3H), 5.91 (s, 1H), 5.40-5.35 (m, 1H), 3.51-3.28 (m, 4H), 1.65 (d, J = 7 Hz, 3H); m/z 591.8 [M+1] ⁺
5ooD		¹ H -NMR (CD ₃ OD, 500 MHz): δ 8.62 (s, 1H), 8.58 (s, 1H), 8.57 (s, 1H), 8.52 (s, 1H), 7.42 (s, 1H), 5.55-5.51 (m, 1H), 4.57 (s, 1H), 4.51-4.46 (m, 1H), 3.68 (s, 1H), 3.20-3.15 (m, 2H), 1.98-1.96 (m, 2H), 1.76 (d, J = 7 Hz, 3H), 1.37-1.30 (m, 2H), 1.45 (s, 9H); m/z 654.7 [M+1] ⁺
19nnD		¹ H -NMR (DMSO-D ₆ , 200 MHz): δ 11.75 (s, 1H), 9.55 (d, N-H), 8.77 (s, 1H), 8.73 (s, 1H), 8.61 (s, 1H), 8.53 (s, 1H), 7.73 (bs, 2H), 7.36 (s, 1H), 5.44-5.37 (m, 1H), 4.47-4.45 (m, 2H), 3.06-2.93 (m, 2H), 2.73-2.60 (m, 2H), 1.97-1.12 (m, 5H), 1.65 (d, J = 7 Hz, 3H); m/z 568.7 [M+1] ⁺
19mmD		¹ H-NMR (DMSO-D ₆ , 500 MHz): δ 11.71 (s, 1H), 9.48 (s, 1H), 8.76 (s, 1H), 8.73 (s, 1H), 8.59 (s, 1H), 8.53 (s, 1H), 7.78 (bs, 2H), 7.01 (bs, 1H), 5.42- 5.39 (m, 1H), 3.02-2.85 (m, 6H), 2.16- 2.02 (m, 3H), 1.66 (d, J = 6.0 Hz, 3H); m/z 554.8 [M+1] ⁺

#	Structure	Characterization Data
5ppD		¹ H-NMR (DMSO-D ₆ , 500 MHz): δ 11.70 (s, 1H), 9.38 (d, J = 7.5 Hz, 1H), 8.76 (s, 1H), 8.72 (s, 1H), 8.54 (s, 1H), 8.50 (s, 1H), 7.88 (s, 1H), 7.11 (s, 1H), 5.41- 5.37 (m, 1H), 3.983.90 (m, 2H), 2.70-2.60 (m, 3H), 1.79-1.68 (m, 4H), 1.61 (d, J = 6.5 Hz, 3H), 1.38 (s, 9H), 1.03 (d, J = 9.5 Hz, 2H); m/z 568.9 [M-Boc].
19rrD		¹ H-NMR (DMSO-D ₆ , 500 MHz): δ 11.71 (s, 1H), 9.42 (s, 1H), 8.76 (s, 1H), 8.72 (s, 1H), 8.54 (s, 1H), 8.47 (s, 1H), 7.93 (bs, 2H), 7.11 (s, 1H), 5.40- 5.37 (m, 1H), 4.20 (bs, 1H), 3.05- 3.03 (m, 4H), 2.04- 2.02 (m, 4H), 1.65 (d, J = 6.0 Hz, 3H); m/z 554.8 [M+1] ⁺
31b		¹ H-NMR (DMSO-D ₆ , 500 MHz): δ 10.01 (s, 1H), 8.75 (d, J = 6.0 Hz, 1H), 8.29 (s, 1H), 8.10 (s, 1H), 7.86 (d, J = 8.0 Hz, 1H), 7.55-7.52 (m, 1H), 7.40 (d, J = 7.0 Hz, 1H), 4.42-4.19 (bs, 1H), 2.90-2.79 (m, 2H), 2.61- 2.58 (m, 2H), 2.25-2.18 (m, 1H), 1.82-1.75 (m, 1H); 443 [M+1] ⁺
19qqD		¹ H-NMR (DMSO-d ₆ , 500 MHz): δ 11.72 (s, 1H), 9.48 (s, 1H), 8.76 (s, 1H), 8.73 (s, 1H), 8.60 (s, 1H), 8.53 (s, 1H), 7.78 (bs, 2H), 7.01 (bs, 1H), 5.42- 5.39 (m, 1H), 3.84-3.78 (m, 2H), 3.34-3.30 (m, 2H), 2.95-2.90 (m, 3H), 2.16- 2.02 (m, 1H), 1.88-1.70 (m, 1H), 1.66 (d, J = 7.0 Hz, 3H); m/z 554.8 [M+1] ⁺
5qqD		¹ H-NMR (DMSO-D ₆ , 500 MHz): δ 11.71 (s, 1H), 9.46 (s, 1H), 8.75 (s, 1H), 8.72 (s, 1H), 8.56 (s, 1H), 8.54 (s, 1H), 7.00 (s, 1H), 6.97 (s, 1H), 5.41- 5.39 (m, 1H), 3.65- 3.51 (m, 1H), 3.50-3.40 (m, 2H), 3.00-2.82 (m, 3H), 2.42-2.37 (m, 1H), 2.10-1.98 (m, 1H), 1.69-1.66 (m, 1H), 1.64 (d, J = 6.0 Hz, 3H), 1.36 (s, 9H); m/z 654.7 [M+1] ⁺
19ooD		¹ H-NMR (CD ₃ OD-D ₄ , 500 MHz): δ 8.60 (s, 2H), 8.58 (s, 1H), 8.52 (s, 1H), 7.44 (s, 1H), 5.54- 5.51 (m, 1H), 4.66-4.62 (m, 2H), 4.53 (bs, 1H), 3.13- 3.08 (m, 2H), 2.09- 1.92 (m, 2H), 1.79 (d, J = 5.0 Hz, 3H), 1.52- 1.50 (m, 2H); m/z 555.0 [M+1] ⁺
19uuD		¹ H-NMR (CD ₃ OD-D ₄ , 500 MHz): δ 8.60 (s, 1H), 8.58 (s, 1H), 8.56 (s, 1H), 8.52 (s, 1H), 7.18 (s, 1H), 5.53- 5.50 (m, 1H), 3.57 (bs, 2H), 3.03- 3.00 (m, 2H), 2.04- 1.99 (m, 2H), 1.77 (d, J = 5.0 Hz, 3H); m/z 528.8 [M+1] ⁺

#	Structure	Characterization Data
5vvD		¹ H-NMR (DMSO-D ₆ , 500 MHz): δ 11.68 (s, 1H), 9.38 (s, 1H), 8.75 (s, 1H), 8.71 (s, 1H), 8.53 (s, 1H), 8.52 (s, 1H), 8.03 (s, 1H), 7.39 (bs, 1H), 7.22 (bs, 1H), 7.00 (bs, 1H), 5.41- 5.37 (m, 1H), 3.93 (bs, 2H), 1.65 (d, <i>J</i> = 5.0 Hz, 3H); <i>m/z</i> 528.7 [M+1] ⁺
5rrD		¹ H-NMR (DMSO-D ₆ , 500 MHz): δ 11.81 (bs, 1H), 9.39 (s, 1H), 8.75 (s, 1H), 8.71 (s, 1H), 8.53 (s, 1H), 8.52 (s, 1H), 7.80 (bs, 1H), 7.08 (s, 1H), 5.40- 5.37 (m, 1H), 4.06 (bs, 1H), 3.87 (bs, 2H), 2.92 (bs, 2H), 1.86 (bs, 2H), 1.64 (d, <i>J</i> = 5.0 Hz, 3H), 1.40 (s, 9H), 1.29- 1.22 (m, 2H); <i>m/z</i> 654.9 [M+1] ⁺
28c		¹ H-NMR (DMSO-D ₆ , 200 MHz): δ 11.74 (s, 1H), 9.47 (d, <i>J</i> = 8.2 Hz, 1H), 8.77 (s, 1H), 8.73 (s, 1H), 8.53 (s, 1H), 8.50 (s, 1H), 8.24 (s, 1H), 7.99-7.97(m, 1H), 7.11 (s, 1H), 5.41-5.33 (m, 1H), 3.24-3.16 (m, 4H), 1.63 (d, <i>J</i> = 7 Hz, 3H); <i>m/z</i> 581.96 [M+1] ⁺
28d		¹ H-NMR (DMSO-D ₆ , 500 MHz): δ 11.73 (bs, 1H), 9.46 (d, <i>J</i> = 8.0 Hz, 1H), 8.76 (s, 1H), 8.72 (s, 1H), 8.54 (s, 1H), 8.52 (s, 1H), 7.96 (bs, 1H), 7.08 (s, 1H), 7.05 (s, 1H), 5.42- 5.34 (m, 1H), 3.53- 3.23 (m, 4H), 1.65 (d, <i>J</i> = 5.0 Hz, 3H); <i>m/z</i> 582.8 [M+1] ⁺
28e		¹ H-NMR (DMSO-D ₆ , 500 MHz): δ 11.74 (bs, 1H), 9.46 (d, <i>J</i> = 8.0 Hz, 1H), 8.76 (s, 1H), 8.72 (s, 1H), 8.54 (s, 2H), 7.98 (s, 1H), 7.90 (s, 2H), 7.64 (s, 1H), 7.18- 7.10 (bs, 2H), 5.42- 5.35 (m, 1H), 3.46- 3.41 (m, 4H), 1.65 (d, <i>J</i> = 5.0 Hz, 3H); <i>m/z</i> 592.8 [M+1] ⁺
20aD		¹ H-NMR (DMSO-D ₆ , 500 MHz): δ 11.71 (bs, N-H), 9.44 (bs, N-H), 8.76 (s, 1H), 8.73 (s, 1H), 8.55 (s, 1H), 8.24 (s, 1H), 7.59 (s, 2H), 5.36 (m, 1H), 1.62 (d, <i>J</i> = 7.0 Hz, 3H); <i>m/z</i> 489.9 [M+1] ⁺

#	Structure	Characterization Data
30c		$^1\text{H-NMR}$ (DMSO- D_6 , 500 MHz): δ 8.71 (d, $J = 7.0$ Hz, 1H), 8.58 (d, $J = 8.0$ Hz, 1H), 8.27 (s, 2H), 7.85-7.35 (bs, 2N-H), 6.97-6.81 (m, 2H), 6.08 (d, $J = 4.0$ Hz, 1H), 6.18 (d, $J = 4.0$ Hz, 1H), 4.22 (d, $J = 10$ Hz, 1H), 4.01-3.85 (m, 2H), 3.82-3.66 (m, 2H), 3.07-3.00 (m, 2H), 2.78 (q, $J = 11$ Hz, 1H), 1.92-1.85 (m, 3H); m/z : 456.8 $[\text{M}+1]^+$
28f		$^1\text{H-NMR}$ (DMSO- D_6 , 200 MHz, Rotamers): δ 12.41 (s, 1H), 11.83 (s, 1H), 11.73 (s, 2H), 9.44 (d, $J = 8.6$ Hz, 2H), 8.76 (s, 2H), 8.76 (s, 2H), 8.72 (s, 2H), 8.54 (s, 4H), 7.96 (bs, 2H), 7.50 (s, 1H), 7.18 (s, 1H), 7.09 (m, 2H), 6.57 (bs, 1H), 6.34 (bs, 1H), 5.41-5.32 (m, 2H), 3.51-3.20 (m, 8H), 1.64 (d, $J = 6.8$ Hz, 6H); m/z 605.8 $[\text{M}+1]^+$
26cD		$^1\text{H-NMR}$ (DMSO- D_6 , 200 MHz): δ 11.73 (s, 1H), 9.53 (d, $J = 8.0$ Hz, 1H), 8.76 (s, 1H), 8.72 (s, 1H), 8.64 (s, 1H), 8.53 (s, 1H), 7.33 (s, 1H), 5.45-5.37 (m, 1H), 4.66 (t, $J = 5.4$ Hz, 1H), 4.12 (d, $J = 5.4$ Hz, 1H), 3.80-3.40 (m, 8H), 1.64 (d, $J = 6.8$ Hz, 3H); m/z 598.7 $[\text{M}+1]^+$
5wwD		$^1\text{H-NMR}$ (DMSO- D_6 , 500 MHz): δ 11.70 (s, 1H), 9.44 (d, $J = 7.0$ Hz, 1H), 8.75 (s, 1H), 8.72 (s, 1H), 8.59 (s, 1H), 8.54 (s, 1H), 7.30 (s, 1H), 7.27 (s, 1H), 6.77 (s, 1H), 5.40-5.35 (m, 1H), 4.43 (bs, 2H), 3.04-3.02 (m, 2H), 2.46-2.42 (m, 1H), 1.80-1.76 (m, 2H), 1.64 (d, $J = 7.5$ Hz, 3H), 1.49-1.46 (m, 2H); m/z 582.7 $[\text{M}+1]^+$
39a		$^1\text{H-NMR}$ (DMSO- D_6 , 500 MHz): δ 11.20 (bs, 1H), 9.39 (d, $J = 7.5$ Hz, 1H), 8.74 (s, 1H), 8.70 (s, 1H), 8.54 (s, 1H), 8.52 (s, 1H), 7.91 (bs, 2H), 7.07 (s, 1H), 5.40-5.36 (m, 1H), 3.40-3.39 (m, 2H), 3.22-3.18 (m, 2H), 1.78 (s, 3H), 1.64 (d, $J = 7.5$ Hz, 3H); m/z 556.8 $[\text{M}+1]^+$
5xxD		$^1\text{H-NMR}$ (DMSO- D_6 , 500 MHz): δ 11.70 (s, 1H), 9.37 (d, $J = 7.5$ Hz, 1H), 8.75 (s, 1H), 8.71 (s, 1H), 8.54 (s, 1H), 8.51 (s, 1H), 7.76 (d, $J = 6.5$ Hz, 1H), 7.08 (s, 1H), 5.40-5.35 (m, 1H), 3.83-3.82 (m, 1H), 2.73-2.70 (m, 2H), 2.15 (s, 3H), 2.00-1.95 (m, 4H), 1.86-1.84 (m, 2H), 1.64 (d, $J = 7.5$ Hz, 3H); m/z 568.8 $[\text{M}+1]^+$

#	Structure	Characterization Data
26bD		¹ H-NMR (CD ₃ OD, 500 MHz): δ 8.63 (s, 1H), 8.61 (s, 1H), 8.58 (s, 1H), 8.52 (s, 1H), 7.43 (s, 1H), 5.56-5.52 (m, 1H), 4.55 (s, 1H), 3.93-3.82 (m, 4H), 3.81-3.70 (m, 4H), 2.03-2.00 (m, 1H), 1.78 (d, <i>J</i> = 7.5 Hz, 3H) 0.94-0.85 (m, 4H); <i>m/z</i> 608.7 [M+1] ⁺
5aaaD		¹ H-NMR (DMSO-D ₆ , 500 MHz): δ 11.71 (s, 1H), 9.47 (d, <i>J</i> = 8.0 Hz, 1H), 8.75 (s, 1H), 8.72 (s, 1H), 8.61 (s, 1H), 8.54 (s, 1H), 7.34 (s, 1H), 5.41-5.38 (m, 1H), 4.00-3.98 (m, 2H), 3.54-3.52 (m, 2H), 3.16-3.12 (m, 2H), 1.97-1.93 (m, 2H), 1.72 (d, <i>J</i> = 7.5 Hz, 3H); <i>m/z</i> 564.8 [M+1] ⁺
26eD		¹ H-NMR (Acetone-D ₆ , 500 MHz): δ 10.34 (s, 1H), 8.90 (d, <i>J</i> = 8.0 Hz, 1H), 8.68 (s, 1H), 8.63 (s, 1H), 8.55 (s, 1H), 8.53 (s, 1H), 7.34 (s, 1H), 5.53-5.49 (m, 1H), 3.85-3.81 (m, 2H), 3.77-3.60 (m, 9H), 2.60-2.57 (m, 2H), 1.75 (d, <i>J</i> = 7.5 Hz, 3H); <i>m/z</i> 613.3 [M+1] ⁺
5yyD		¹ H-NMR (DMSO-D ₆ , 500 MHz): δ 11.71 (s, 1H), 9.38 (d, <i>J</i> = 8.0 Hz, 1H), 8.76 (s, 1H), 8.72 (s, 1H), 8.54 (s, 1H), 8.51 (s, 1H), 7.90 (s, 1H), 7.11 (s, 1H), 5.40-5.37 (m, 1H), 1.63-1.47 (m, 6H), 1.34 (d, <i>J</i> = 7.5 Hz, 3H), 1.33-1.22 (m, 4H); <i>m/z</i> 582.9 [M+1] ⁺
31c		¹ H-NMR (DMSO-D ₆ , 500 MHz): δ 10.04 (s, 1H), 8.49 (d, <i>J</i> = 8.0 Hz, 1H), 8.28 (s, 1H), 8.15 (s, 1H), 7.87 (d, <i>J</i> = 9.0 Hz, 1H), 7.53 (t, <i>J</i> = 8.0 Hz, 1H), 7.40 (d, <i>J</i> = 8.0 Hz, 1H), 3.71 (bs, 1H), 3.14 (s, 2H), 2.88-2.83 (m, 2H), 2.28-2.24 (m, 2H), 1.80-1.78 (m, 2H), 1.63-1.61 (m, 2H); <i>m/z</i> 457 [M+1] ⁺
31a		¹ H-NMR (DMSO-D ₆ , 500 MHz): δ 9.99 (s, 1H), 8.57 (d, <i>J</i> = 8.0 Hz, 1H), 8.28 (s, 1H), 8.10 (s, 1H), 7.82 (d, <i>J</i> = 7.5 Hz, 1H), 7.54-7.51 (m, 1H), 7.39 (d, <i>J</i> = 7.5 Hz, 1H), 4.03-3.98 (m, 1H), 3.20-3.06 (m, 2H), 2.69 (d, <i>J</i> = 9.5 Hz, 1H), 2.49-2.30 (m, 2H), 1.68-1.59 (m, 3H), 1.42-1.34 (m, 1H); <i>m/z</i> 456.9 [M+1] ⁺







#	Structure	Characterization Data
40b		¹ H-NMR (DMSO-D ₆ , 500 MHz): δ 11.71 (s, 1H), 9.40 (s, 1H), 8.75 (s, 1H), 8.72 (s, 1H), 8.54 (s, 1H), 8.50 (s, 1H), 7.89 (s, 1H), 7.21 (s, 1H), 5.40-5.37 (m, 1H), 3.69-3.67 (m, 1H), 3.48-3.47 (m, 1H), 1.64 (d, <i>J</i> = 6.5 Hz, 3H), 1.24 (s, 6H); <i>m/z</i> 570.9 [M+1] ⁺
20bDa		¹ H-NMR (DMSO-D ₆ , 500 MHz): δ 11.75 (s, 1H), 9.46 (d, <i>J</i> = 8.5 Hz, 1H), 8.75 (s, 1H), 8.72 (s, 1H), 8.48 (bs, 1H), 8.12 (s, 1H), 5.41-5.39 (m, 1H), 3.78-3.65 (m, 8H), 1.67 (d, <i>J</i> = 8.0 Hz, 3H); <i>m/z</i> 559.6 [M+1] ⁺
26dD		¹ H-NMR (CD ₃ OD, 500 MHz): δ 8.65 (s, 1H), 8.61 (s, 1H), 8.59 (s, 1H), 8.53 (s, 1H), 7.45 (s, 1H), 5.55-5.53 (m, 1H), 4.02 (s, 2H), 3.92-3.84 (m, 4H), 3.80-3.76 (m, 2H), 3.63-3.59 (m, 2H), 1.78 (d, <i>J</i> = 7.0 Hz, 3H); <i>m/z</i> 598.2 [M+1] ⁺
18mD		¹ H-NMR (DMSO-D ₆ , 200 MHz): δ 11.77 (s, 1H), 9.81 (d, <i>J</i> = 8.5 Hz, 1H), 9.13 (s, 1H), 8.78 (s, 1H), 8.76 (s, 1H), 8.75 (s, 1H), 8.46 (s, 1H), 5.47-5.43 (m, 1H), 4.23-4.13 (m, 2H), 3.98 (s, 2H), 3.57-3.54 (m, 2H), 1.64 (d, <i>J</i> = 6.5 Hz, 3H); <i>m/z</i> 554.8 [M+1] ⁺
28g		¹ H-NMR (Acetone-D ₆ , 500 MHz): δ 10.74 (bs, 1H), 8.89 (s, 1H), 8.67 (s, 1H), 8.63 (s, 1H), 8.55 (s, 1H), 8.47 (s, 1H), 7.20-7.16 (m, 3H), 5.49-5.40 (m, 1H), 3.57-3.48 (m, 6H), 1.74 (d, <i>J</i> = 6.5 Hz, 3H); <i>m/z</i> 580.7 [M+1] ⁺
28h		¹ H-NMR (DMSO-D ₆ , 500 MHz): δ 11.74 (s, 1H), 9.45 (s, 1H), 8.76 (s, 1H), 8.72 (s, 1H), 8.61 (s, 1H), 8.54 (s, 1H), 8.02 (s, 1H), 7.86 (s, 1H), 7.09 (s, 1H), 5.40-5.37 (m, 1H), 3.80-3.60 (m, 1H), 1.64 (d, <i>J</i> = 7 Hz, 3H); <i>m/z</i> 598.7 [M+1] ⁺







#	Structure	Characterization Data
5bbbDa		$^1\text{H-NMR}$ (DMSO- D_6 , 500 MHz): δ 11.74 (s, 1H), 9.49 (s, 1H), 8.76 (s, 1H), 8.72 (s, 1H), 8.57 (s, 1H), 8.54 (s, 1H), 7.20 (s, 0.5H), 6.97 (s, 0.5H), 5.41-5.38 (m, 1H), 4.99 (s, 0.5H), 4.80 (s, 0.5H), 4.25 (s, 0.5H), 3.95 (s, 0.5H), 3.63-3.49 (m, 3H), 2.04-1.93 (m, 4H), 1.65 (d, J = 6.5 Hz, 3H); m/z 555.7 $[\text{M}+1]^+$
41		$^1\text{H-NMR}$ (DMSO- D_6 , 500 MHz): δ 11.74 (s, 1H), 10.50 (s, 1H), 9.52 (s, 1H), 8.76 (s, 1H), 8.72 (s, 1H), 8.63 (s, 1H), 8.54 (s, 1H), 7.31 (s, 1H), 5.42-5.39 (m, 1H), 3.81-3.70 (m, 4H), 2.49-2.41 (m, 4H), 1.66 (d, J = 7 Hz, 3H), m/z 568.7 $[\text{M}+1]^+$
28i		$^1\text{H-NMR}$ (DMSO- D_6 , 500 MHz): δ 11.9 (s, 1H), 9.45 (s, 1H), 8.76 (s, 1H), 8.72 (s, 1H), 8.54 (s, 2H), 7.97 (s, 1H), 7.09 (s, 1H), 5.40-5.37 (m, 1H), 3.85-3.82 (m, 2H), 3.50 (bs, 2H), 3.36-3.31 (m, 2H), 1.64 (d, J = 7 Hz, 3H), m/z 599.7 $[\text{M}+1]^+$
28j		$^1\text{H-NMR}$ (DMSO- D_6 , 500 MHz): δ 11.74 (s, 1H), 9.44 (d, J = 7.5 Hz, 1H), 8.76 (s, 1H), 8.72 (s, 1H), 8.54 (s, 2H), 8.26 (s, 2H), 7.98 (s, 1H), 7.20 (s, 1H), 7.07 (s, 1H), 6.56 (s, 1H), 5.42-5.39 (m, 1H), 3.53-3.44 (m, 4H), 1.64 (d, J = 6.5 Hz, 3H), m/z 592.6 $[\text{M}+1]^+$
28k		$^1\text{H-NMR}$ (DMSO- D_6 , 500 MHz): δ 11.82 (s, 1H), 9.44 (d, NH), 8.75 (s, 1H), 8.70 (s, 1H), 8.55 (s, 1H), 8.53 (s, 1H), 8.39 (s, 1H), 8.00 (bs, 2H), 7.49 (s, 1H), 7.08 (s, 1H), 6.45 (s, 1H), 5.39-5.37 (m, 1H), 3.52-3.46 (m, 4H), 1.64 (d, J = 7.5 Hz, 3H), m/z 592.6 $[\text{M}+1]^+$
5dddDa		$^1\text{H-NMR}$ (DMSO- D_6 , 500 MHz): δ 11.74 (s, 1H), 9.55 (s, 1H), 8.76 (s, 1H), 8.73 (s, 1H), 8.67 (s, 1H), 8.54 (s, 1H), 7.39 (s, 1H), 5.43-5.40 (m, 1H), 4.10-4.00 (m, 4H), 3.31-3.29 (m, 4H), 1.66 (d, J = 7 Hz, 3H); m/z 553.7 $[\text{M}+1]^+$

#	Structure	Characterization Data
5vvDa		¹ H-NMR (DMSO-D ₆ , 500 MHz): δ 11.74 (s, 1H), 9.45 (d, <i>J</i> = 7 Hz, 1H), 8.76 (s, 1H), 8.72 (s, 1H), 8.54 (s, 1H), 8.53 (s, 1H), 8.08 (s, 1H), 7.43 (s, 1H), 7.22 (s, 1H), 7.05 (s, 1H), 5.40-5.37 (m, 1H), 3.93 (s, 2H), 1.64 (d, <i>J</i> = 7 Hz, 3H); <i>m/z</i> 528.7 [M+1] ⁺
26cDa		¹ H-NMR (DMSO-D ₆ , 500 MHz): δ 11.73 (s, 1H), 9.53 (d, <i>J</i> = 9 Hz, 1H), 8.76 (s, 1H), 8.71 (s, 1H), 8.64 (s, 1H), 8.54 (s, 1H), 7.34 (s, 1H), 5.42-5.39 (m, 1H), 4.69-4.68 (m, 1H), 4.12 (d, <i>J</i> = 5.5 Hz, 2H), 3.80-3.64 (m, 4H), 3.60-3.55 (m, 2H), 3.52-3.46 (m, 2H), 1.65 (d, <i>J</i> = 6.5 Hz, 3H); <i>m/z</i> 598.7 [M+1] ⁺
18fDa		¹ H-NMR (DMSO-D ₆ , 500 MHz): δ 11.74 (s, 1H), 9.88 (d, <i>J</i> = 7.5 Hz, 1H), 9.24 (s, 1H), 8.76-8.74 (m, 3H), 8.54 (s, 1H), 8.37 (s, 1H), 8.02 (s, 1H), 6.72 (s, 1H), 5.52-5.47 (m, 1H), 1.69 (d, <i>J</i> = 6.5 Hz, 3H); <i>m/z</i> 522.8 [M+1] ⁺
42a		¹ H-NMR (DMSO-D ₆ , 500 MHz): δ 11.75 (s, 1H), 9.58 (s, 1H), 8.77 (s, 1H), 8.73 (s, 1H), 8.60 (s, 1H), 8.54 (s, 1H), 7.38 (s, 1H), 5.43-5.40 (m, 1H), 4.60-4.40 (m, 2H), 3.08-2.97 (m, 2H), 2.18 (d, <i>J</i> = 6.5 Hz, 2H), 2.10-1.98 (m, 1H), 1.78 (d, <i>J</i> = 11 Hz, 2H), 1.65 (d, <i>J</i> = 7.5 Hz, 3H), 1.22-1.14 (m, 2H); <i>m/z</i> 597.7 [M+1] ⁺
42b		¹ H-NMR (DMSO-D ₆ , 500 MHz): δ 11.75 (s, 1H), 9.64 (bs, 1H), 8.77 (s, 1H), 8.74 (s, 1H), 8.63 (s, 1H), 8.54 (s, 1H), 7.45 (bs, 1H), 5.44-5.41 (m, 1H), 4.44-4.24 (m, 2H), 3.24-3.18 (m, 2H), 2.65-2.60 (m, 1H), 1.93 (d, <i>J</i> = 11 Hz, 2H), 1.66 (d, <i>J</i> = 7.5 Hz, 3H), 1.53 (d, <i>J</i> = 11.5 Hz, 2H); <i>m/z</i> 583.7 [M+1] ⁺
32b		¹ H-NMR (Acetone-D ₆ , 500 MHz): □ 11.95-11.89 (bs, 1N-H), 8.56 (bs, 1N-H), 8.25 (s, 1H), 8.00 (s, 1H), 7.52 (d, <i>J</i> = 8.0 Hz, 1H), 7.23 (d, <i>J</i> = 8.0 Hz, 1H), 4.62-4.55 (bs, 2H), 4.02-3.99 (m, 1H), 3.18-3.09 (m, 2H), 2.39-2.36 (m, 2H), 2.22-2.15 (m, 2H), 2.02-1.98 (m, 2H), 1.69-1.58 (m, 2H); <i>m/z</i> 453.8 [M+1] ⁺








#	Structure	Characterization Data
5eeeDa		¹ H-NMR (DMSO-D ₆ , 500 MHz): δ 11.74 (s, 1H), 9.51 (d, NH), 8.76 (s, 1H), 8.72 (s, 1H), 8.60 (s, 1H), 8.54 (s, 1H), 7.30 (s, 1H), 5.41-5.38 (m, 1H), 3.67- 3.58 (m, 4H), 3.48- 3.43 (m, 4H), 3.31 (s, 3H), 3.29- 3.22 (m, 4H), 1.65 (d, <i>J</i> = 7 Hz, 3H); <i>m/z</i> 598.7 [M+1] ⁺
32a		¹ H-NMR (DMSO-D ₆ , 500 MHz): δ 9.08 (s, 1N-H), 8.56 (bs, 1N-H), 8.25 (s, 1H), 8.00 (s, 1H), 4.62-4.55 (bs, 2H), 4.02-3.99 (m, 1H), 3.18-3.09 (m, 2H), 2.39-2.36 (m, 2H), 2.22-2.15 (m, 2H), 2.02-1.98 (m, 2H), 1.69-1.58 (m, 2H), 1.22 (s, 9H); <i>m/z</i> 443.9 [M+1] ⁺
4vDa		¹ H-NMR (DMSO-D ₆ , 200 MHz): δ 11.78 (s, 1H), 9.82 (d, <i>J</i> = 8.5 Hz, 1H), 9.12 (s, 1H), 8.78 (s, 1H), 8.76 (s, 1H), 8.68 (s, 1H), 8.54 (s, 1H), 8.25 (s, 1H), 8.21 (s, 1H), 5.46- 5.43 (m, 1H), 3.92 (s, 3H), 1.67 (d, <i>J</i> = 7.5 Hz, 3H); <i>m/z</i> 536.8 [M+1] ⁺
4vD		¹ H-NMR (DMSO-D ₆ , 500 MHz): δ 11.74 (s, 1H), 9.77 (d, NH), 9.19 (s, 1H), 8.76 (s, 1H), 8.74 (s, 1H), 8.65 (s, 1H), 8.54 (s, 1H), 8.25 (s, 1H), 8.20 (s, 1H), 5.48-5.45 (m, 1H), 3.92 (s, 3H), 1.68 (d, <i>J</i> = 6.5 Hz, 3H), <i>m/z</i> 536.7 [M+1] ⁺
5ttDa		¹ H-NMR (DMSO-D ₆ , 500 MHz): δ 11.74 (bs, 1H), 9.43 (bs, 1H), 8.73 (s, 1H), 8.71 (s, 1H), 8.39 (s, 2H), 7.92 (s, 1H), 7.39 (s, 1H), 7.15 (s, 1H), 6.99 (s, 1H), 5.39-5.34 (m, 1H), 3.47-3.44 (m, 2H), 2.41-2.38 (m, 2H), 1.69 (d, <i>J</i> = 6.5 Hz, 3H); <i>m/z</i> 542.6 [M+1] ⁺
28l		¹ H-NMR (DMSO-D ₆ , 500 MHz): δ 11.74 (s, 1H), 9.45 (s, 1H), 8.77 (s, 1H), 8.72 (s, 1H), 8.56 (s, 1H), 8.54 (s, 1H), 8.02 (bs, 1H), 7.92 (s, 1H), 7.90 (s, 1H), 7.09 (s, 1H), 5.42-5.39 (m, 1H), 3.57- 3.45 (m, 4H), 1.64 (d, <i>J</i> = 6.5 Hz, 3H); <i>m/z</i> 598.6 [M+1] ⁺

#	Structure	Characterization Data
4wD		¹ H-NMR (CD ₃ OD, 500 MHz): δ 9.28 (s, 1H), 8.65 (s, 1H), 8.62 (s, 1H), 8.59 (s, 1H), 8.54 (s, 1H), 7.83 (s, 1H), 7.13 (s, 1H), 5.62- 5.61 (m, 1H), 1.82 (d, <i>J</i> = 7 Hz, 3H); <i>m/z</i> 522.6 [M+1] ⁺
4cD		¹ H-NMR (DMSO-D ₆ , 500 MHz): δ 13.41 (s, 1H), 11.74 (s, 1H), 9.76 (d, <i>J</i> = 7.0 Hz, 1H), 9.20 (s, 1H), 8.76 (s, 1H), 8.74 (s, 1H), 8.72 (s, 1H), 8.54 (s, 1H), 8.29 (s, 1H), 8.27 (s, 1H), 5.48-5.45 (m, 1H), 1.69 (d, <i>J</i> = 7 Hz, 3H), <i>m/z</i> 522.9 [M+1] ⁺
28m		¹ H-NMR (DMSO-D ₆ , 500 MHz): δ 12.12 (s, 1H), 11.74 (s, 1H), 9.54 (s, 1H), 8.77 (s, 1H), 8.73 (s, 1H), 8.54 (s, 1H), 8.51 (s, 1H), 7.98 (bs, 1H), 7.14 (s, 1H), 6.97 (s, 1H), 5.45- 5.37 (m, 1H), 3.49 (bs, 4H), 2.82 (bs, 2H), 1.65 (d, <i>J</i> = 7.0 Hz, 3H); <i>m/z</i> 580.8 [M+1] ⁺
1nDa		¹ H-NMR (DMSO-D ₆ , 500 MHz): δ 11.76 (s, 1H), 9.56 (d, <i>J</i> = 7.0, 1H), 8.77 (s, 1H), 8.75 (s, 1H), 8.63 (s, 1H), 8.57 (s, 1H), 5.36- 5.33 (m, 1H), 3.71-3.68 (m, 8H), 1.58 (d, <i>J</i> = 7.5 Hz, 3H); <i>m/z</i> 575.7 [M+1] ⁺
5zzD		¹ H-NMR (DMSO-D ₆ , 500 MHz): δ 11.78 (s, 1H), 9.46 (d, <i>J</i> = 8.5 Hz, 1H), 8.78 (s, 1H), 8.76 (s, 1H), 8.58 (s, 1H), 8.56 (s, 1H), 7.25 (s, 1H), 5.43- 5.38 (m, 1H), 3.65 (bs, 4H), 1.67 (d, <i>J</i> = 7.5 Hz, 3H), 1.57 (bs, 6H); <i>m/z</i> 539.7 [M+1] ⁺
10U		¹ H-NMR: (DMSO-D ₆ , 500 MHz) δ: 10.50 (s, 1H), 9.10 (d, NH), 8.31 (s, 1H), 7.82 (d, <i>J</i> = 8 Hz, 2H), 7.62 (d, <i>J</i> = 8 Hz, 2H), 7.21 (s, 1H), 5.29-5.31 (m, 1H), 1.52 (d, <i>J</i> = 7 Hz, 3H); <i>m/z</i> 442.7 [M+1] ⁺







#	Structure	Characterization Data
4qU		$^1\text{H-NMR}$ (DMSO- D_6 , 500 MHz): δ 10.48 (s, 1H), 9.60 (s, 2H), 9.52 (d, J = 7.0 Hz, 2H), 9.37 (s, 1H), 8.70 (s, 1H), 7.77 (d, J = 8.0 Hz, 2H), 7.62 (d, J = 8.0 Hz, 2H), 7.22 (s, 1H), 5.40-5.37 (m, 1H), 1.64 (d, J = 7.0 Hz, 3H); m/z 471.7 $[\text{M}+1]^+$
4qV		$^1\text{H-NMR}$ (DMSO- D_6 , 500 MHz): δ 10.48 (s, 1H), 9.60 (s, 2H), 9.52 (d, J = 7.0 Hz, 2H), 9.37 (s, 1H), 8.70 (s, 1H), 7.77 (d, J = 8.0 Hz, 2H), 7.62 (d, J = 8.0 Hz, 2H), 7.22 (s, 1H), 4.51 (d, J = 3.0 Hz, 2H); m/z 457.9 $[\text{M}+1]^+$
5dV		$^1\text{H-NMR}$ (DMSO- D_6 , 500 MHz): δ 10.45 (s, 1H), 9.31 (s, 1H), 8.57 (s, 1H), 7.77 (d, J = 8 Hz, 2H), 7.62 (d, J = 8.0 Hz, 2H), 7.30 (s, 1H), 7.16 (s, 1H), 4.50 (d, J = 3.0 Hz, 2H), 3.63-3.60 (m, 8H); m/z 464.9 $[\text{M}+1]^+$
18dV		$^1\text{H-NMR}$ (DMSO- D_6 , 500 MHz): δ 10.49 (s, 1H), 9.69 (s, 1H), 9.25 (s, 1H), 8.17 (s, 1H), 7.95 (s, 1H), 7.77 (d, J = 8.0 Hz, 2H), 7.62 (d, J = 8.0 Hz, 2H), 7.21 (s, 1H), 6.99 (s, 1H), 4.56 (d, J = 6.0 Hz, 2H), 2.66 (s, 3H); m/z 459.9 $[\text{M}+1]^+$
5vV		$^1\text{H-NMR}$ (DMSO- D_6 , 500 MHz): δ 10.47 (s, 1H), 9.25 (d, J = 8.5 Hz, 1H), 8.46 (s, 1H), 7.69 (d, J = 8.5 Hz, 2H), 7.59 (d, J = 8.0 Hz, 2H), 7.24 (s, 1H), 7.16 (s, 1H), 4.47 (s, 2H), 3.77-3.73 (m, 1H), 3.25- 3.21 (m, 2H), 1.79- 1.77 (m, 2H), 1.40- 1.39 (m, 2H); m/z 478.9 $[\text{M}+1]^+$
25iU		$^1\text{H-NMR}$ (DMSO- D_6 , 500 MHz): δ 10.48 (s, 1H), 9.57 (d, J = 8.0 Hz, 1H), 9.48 (s, 1H), 8.39 (s, 1H), 7.77 (d, J = 8.5 Hz, 2H), 7.62 (d, J = 8.0 Hz, 2H), 7.20 (s, 1H), 5.37- 5.36 (m, 1H), 1.62 (d, J = 7.0 Hz, 3H); m/z 437.7 $[\text{M}+1]^+$

#	Structure	Characterization Data
10CC		¹ H-NMR (DMSO-D6- 500 MHz) δ 9.12 (s, 1H), 9.10 (d, NH), 8.25 (s, 1H), 7.62 (d, <i>J</i> = 8.5 Hz, 2H), 7.42 (d, <i>J</i> = 6.5 Hz, 1H), 7.32 (d, <i>J</i> = 8.5 Hz, 2H), 7.12 (d, <i>J</i> = 6.5 Hz, 1H), 5.23-5.19 (m, 1H), 1.45 (d, <i>J</i> = 8 Hz, 3H), 1.21 (s, 9H); <i>m/z</i> 425.9 [M+1] ⁺
5vU		¹ H-NMR (DMSO-D6, 500 MHz): δ 10.47 (s, 1H), 9.05 (d, <i>J</i> = 8.5 Hz, 1H), 8.53 (s, 1H), 7.77 (d, <i>J</i> = 8.5 Hz, 2H), 7.62 (d, <i>J</i> = 8.0 Hz, 2H), 7.28 (s, 1H), 7.17 (s, 1H), 5.30-5.27 (m, 1H), 4.79 (bs, 1H), 4.02 (bs, 1H), 3.77- 3.75 (m, 2H), 3.25- 3.21 (m, 2H), 1.78 (bs, 2H), 1.59 (d, <i>J</i> = 7.0 Hz, 3H), 1.35- 1.21 (m, 2H); <i>m/z</i> 493.2 [M+1] ⁺
18dU		¹ H-NMR (DMSO-D6, 500 MHz): δ 10.49 (s, 1H), 9.52 (d, <i>J</i> = 8.0 Hz, 1H), 9.25 (s, 1H), 8.17 (s, 1H), 7.95 (s, 1H), 7.77 (d, <i>J</i> = 8.0 Hz, 2H), 7.63 (d, <i>J</i> = 8.0 Hz, 2H), 7.21 (s, 1H), 6.99 (s, 1H), 5.37-5.35 (m, 1H), 2.66 (s, 3H), 1.63 (d, <i>J</i> = 7.0 Hz, 3H); <i>m/z</i> 474.1 [M+1] ⁺
25iV		¹ H-NMR (DMSO-D6, 500 MHz): δ 10.48 (s, 1H), 9.70 (s, 1H), 9.48 (s, 1H), 8.44 (s, 1H), 8.41 (s, 1H), 7.77 (d, <i>J</i> = 8.5 Hz, 2H), 7.62 (d, <i>J</i> = 8.0 Hz, 2H), 7.20 (s, 1H), 4.56 (d, <i>J</i> = 7.0 Hz, 3H); <i>m/z</i> 423.0 [M+1] ⁺
10BB		¹ H NMR (DMSO-D6, 500 MHz) δ 8.3 (s, 1H), 7.9 (d, <i>J</i> = 10 Hz, 2H), 7.6 (d, <i>J</i> = 10 Hz, 2H), 7.5 (d, <i>J</i> = 10 Hz, 1H), 7.2 (d, <i>J</i> = 10 Hz, 1H), 5.19-5.23 (m, 1H), 1.30 (s, 3H); <i>m/z</i> 438 [M+1] ⁺
10AA		¹ H-NMR (DMSO-D6, 500 MHz): δ 9.92 (s, 1H), 9.03 (d, <i>J</i> = 4 Hz, 1H), 8.31 (s, 1H), 8.27 (d, <i>J</i> = 4 Hz, 2H), 7.89 (d, <i>J</i> = 4 Hz, 2H), 7.64 (d, <i>J</i> = 4 Hz, 2H), 5.10 (m, 1H), 1.47 (d, <i>J</i> = 4 Hz, 3H); <i>m/z</i> 438 [M+1] ⁺

#	Structure	Characterization Data
5dJJ		¹ H-NMR (DMSO-D ₆ , 500 MHz): δ 10.88 (s, 1H), 9.53 (d, <i>J</i> = 8.0 Hz, 2H), 8.64 (s, 1H), 8.53 (s, 2H), 8.31 (d, <i>J</i> = 7.0 Hz, 1H), 7.85 (d, <i>J</i> = 7.0 Hz, 2H), 7.32 (s, 1H), 5.44-5.41 (m, 1H), 3.67 (s, 8H), 3.30 (s, 3H), 1.64 (d, <i>J</i> = 7.0, 3H); <i>m/z</i> 570.9 [M+1] ⁺
5vJJ		¹ H-NMR (DMSO-D ₆ , 500 MHz): δ 10.87 (s, 1H), 9.50 (d, <i>J</i> = 8.0 Hz, 2H), 8.59 (s, 1H), 8.52 (s, 2H), 8.31 (d, <i>J</i> = 5 Hz, 1H), 7.85 (d, <i>J</i> = 5 Hz, 2H), 7.31 (s, 1H), 5.42-5.34 (m, 1H), 4.78 (d, <i>J</i> = 5 Hz, 1H), 4.05 (bs, 1H), 3.8 (bs, 1H), 1.77 (s, 2H), 1.67 (d, <i>J</i> = 7.0, 3H), 1.35 (m, 4H); <i>m/z</i> 584.8 [M+1] ⁺
33aJJ		¹ H-NMR (DMSO-D ₆ , 500 MHz): δ 13.17 (s, 1H), 10.89 (s, 1H), 9.82 (s, 1H), 9.28 (s, 1H), 8.54 (d, <i>J</i> = 10 Hz, 2H), 8.42 (s, 1H), 8.31 (d, <i>J</i> = 5 Hz, 1H), 7.85 (d, <i>J</i> = 5 Hz, 2H), 6.98 (s, 1H), 5.45-5.41 (m, 1H), 2.25 (d, <i>J</i> = 10 Hz, 3H), 1.69 (d, <i>J</i> = 7.0, 3H); <i>m/z</i> 565.8 [M+1] ⁺
4qJJ		¹ H-NMR (DMSO-D ₆ , 500 MHz): δ 10.90 (s, 1H), 9.95 (d, <i>J</i> = 10 Hz, 1H), 9.62 (d, <i>J</i> = 10 Hz, 2H), 9.52 (s, 1H), 9.37 (s, 1H), 8.74 (s, 1H), 8.55 (d, <i>J</i> = 15 Hz, 3H), 8.31 (d, <i>J</i> = 5 Hz, 1H), 7.85 (d, <i>J</i> = 5 Hz, 2H), 5.51-5.55 (m, 1H), 3.28 (s, 3H), 1.72 (d, <i>J</i> = 5 Hz, 3H); <i>m/z</i> 563.9 [M+1] ⁺
4bU		¹ H-NMR (DMSO-D ₆ , 500 MHz): δ 10.46 (s, 1H), 9.44-9.38 (m, 3H), 8.76 (d, <i>J</i> = 4 Hz, 1H), 8.63 (d, <i>J</i> = 3 Hz, 1H), 8.58 (s, 1H), 7.78-7.75 (m, 2H), 7.62-7.59 (m, 3H), 7.22 (s, 1H), 5.40-5.37 (m, 1H), 1.54 (d, <i>J</i> = 7 Hz, 3H); <i>m/z</i> 470.7 [M+1] ⁺
4eU		¹ H-NMR (DMSO-D ₆ , 500 MHz): δ 12.32 (s, 1H), 10.47 (s, 1H), 9.34 (d, <i>J</i> = 9 Hz, 1H), 9.27 (s, 1H), 9.25 (s, 1H), 8.70-8.68 (m, 1H), 7.76 (d, <i>J</i> = 9 Hz, 2H), 7.71 (d, <i>J</i> = 5 Hz, 1H), 7.61 (d, <i>J</i> = 9 Hz, 2H), 6.47 (t, <i>J</i> = 13.5 Hz, 1H), 5.36-5.33 (m, 1H), 1.62 (d, <i>J</i> = 10 Hz, 3H); <i>m/z</i> 486.9 [M+1] ⁺

#	Structure	Characterization Data
6gU		¹ H-NMR (DMSO-D ₆ , 500 MHz): δ 10.54 (s, 1H), 10.47 (s, 1H), 9.16 (d, <i>J</i> = 8.5 Hz, 1H), 8.78 (s, 1H), 8.34 (d, <i>J</i> = 4.5 Hz, 1H), 8.29 (s, 1H), 7.78-7.75 (m, 4H), 7.64-7.61 (m, 2H), 7.19 (s, 1H), 7.05 (s, 1H), 5.34-5.31 (m, 1H), 1.61 (d, <i>J</i> = 4.2 Hz, 3H); <i>m/z</i> 486 [M+1] ⁺
5dKK		¹ H-NMR (CD ₃ OD, 500 MHz): δ 8.62 (s, 1H), 8.45 (s, 1H), 7.40 (s, 1H), 6.62 (s, 1H), 5.58-5.52 (m, 1H), 3.84-3.75 (m, 8H), 1.77 (d, <i>J</i> = 7.0 Hz, 3H), 1.38 (s, 9H); <i>m/z</i> 485.9 [M+1] ⁺
10KK		¹ H-NMR (CD ₃ OD, 500 MHz): δ 8.45 (s, 1H), 8.35 (s, 1H), 6.63 (s, 1H), 5.50-5.48 (m, 1H), 1.73 (d, <i>J</i> = 7.0 Hz, 3H), 1.38 (s, 9H); <i>m/z</i> 449.8 [M+1] ⁺
5dU		¹ H-NMR (DMSO-D ₆ , 500 MHz): δ 10.44 (s, 1H), 9.08 (d, <i>J</i> = 7.0, 1H), 8.61 (s, 1H), 7.78 (d, <i>J</i> = 8.5 Hz, 2H), 7.58 (d, <i>J</i> = 8.5 Hz, 2H), 7.31 (s, 1H), 7.21 (s, 1H), 5.23-5.22 (m, 1H), 3.65-3.60 (m, 8H), 1.65 (d, <i>J</i> = 7.0 Hz, 3H); <i>m/z</i> 479 [M+1] ⁺
5vvU		¹ H-NMR (DMSO-D ₆ , 500 MHz): δ 10.45 (s, 1H), 8.98 (d, <i>J</i> = 7.0, 1H), 8.45 (s, 1H), 8.04 (bs, 1H), 7.78 (d, <i>J</i> = 8.5 Hz, 2H), 7.59 (d, <i>J</i> = 8.5 Hz, 2H), 7.43 (bs, 1H), 7.15 (s, 2H), 7.07 (s, 1H), 5.24-5.22 (m, 1H), 3.89 (s, 2H), 1.56 (d, <i>J</i> = 7.0 Hz, 3H); <i>m/z</i> 465.7 [M+1] ⁺
5vKK		¹ H-NMR (CD ₃ OD, 500 MHz): δ 8.55 (s, 1H), 8.43 (s, 1H), 7.39 (s, 1H), 6.61 (s, 1H), 5.52-5.49 (m, 1H), 4.26-4.18 (m, 2H), 3.98-3.87 (m, 1H), 3.41-3.36 (m, 2H), 1.95-1.93 (m, 2H), 1.74 (d, <i>J</i> = 7.0 Hz, 3H), 1.52-1.50 (m, 2H), 1.36 (s, 9H); <i>m/z</i> 499.8 [M+1] ⁺
26cU		¹ H-NMR (DMSO-D ₆ , 500 MHz): δ 10.46 (s, 1H), 9.07 (d, <i>J</i> = 8.5 Hz, 1H), 8.58 (s, 1H), 7.76 (d, <i>J</i> = 9 Hz, 2H), 7.61 (d, <i>J</i> = 9 Hz, 2H), 7.31 (s, 1H), 7.17 (s, 1H), 5.31-5.28 (m, 1H), 4.66 (t, <i>J</i> = 11.5 Hz, 1H), 4.12 (d, <i>J</i> = 6 Hz, 2H), 3.78-3.65 (m, 4H), 3.60-3.50 (m, 2H), 3.48-3.40 (m, 2H), 1.59 (d, <i>J</i> = 7 Hz, 3H); <i>m/z</i> 536 [M+1] ⁺

#	Structure	Characterization Data
5ccU		¹ H-NMR (DMSO-D ₆ , 500 MHz): δ 10.44 (s, 1H), 8.95 (d, <i>J</i> = 7.0, 1H), 8.40 (s, 1H), 7.80 (s, 1H), 7.68 (d, <i>J</i> = 8.5 Hz, 2H), 7.58 (d, <i>J</i> = 8 Hz ⁺ , 2H), 7.14 (s, 1H), 6.99 (s, 1H), 5.25-5.11 (m, 1H), 3.40-3.38 (m, 6H), 1.72-1.64 (m, 2H), 1.54 (d, <i>J</i> = 7.5 Hz, 3H), 1.04 (t, <i>J</i> = 7 Hz, 3H); m/z 495.1 [M+1] ⁺
43		¹ H-NMR (CDCl ₃) δ = 8.25 (s, 1H), 8.0 (s, 1H), 7.78 (d, 1H), 7.58 (d, 1H), 6.22 (q, 1H), 2.56 (s, 3H), 1.88 (d, 3H)-
34a		¹ H NMR (DMSO-D ₆ , 500 MHz) δ 10.6 (s, 1H, D ₂ O exchangeable), 10.3 (s, 1H, D ₂ O exchangeable), 8.4 (2s, 2H), 8.15 (m, 2H), 7.8 (m, 1H), 7.75 (m, 1H), 7.5 (m, 1H), 2.3 (s, 3H); MS: m/z 484.26 [M+1] ⁺ .
34b		¹ H-NMR (ACETONE-D ₆ , 500 MHz): δ 9.95 (bs, 1H), 9.60 (bs, 1H), 8.54 (s, 1H), 8.38 (s, 1H), 7.82 (d, <i>J</i> = 8.5 Hz, 2H), 7.72 (d, 1H), 7.38-7.28 (m, 3H), 7.07-7.044 (m, 2H), 2.39 (s, 3H); m/z 381.9 [M+1] ⁺ .
34c		¹ H-NMR (DMSO-D ₆ , 500 MHz): δ 10.77 (s, 1H), 10.26 (s, 1H), 8.40-8.38 (m, 2H), 8.18-8.15 (m, 2H), 7.87-7.82 (m, 2H), 7.40 (d, <i>J</i> = 8.5 Hz, 1H), 7.17-7.15 (m, 1H), 2.33 (s, 3H); m/z 382.9 [M+1] ⁺ .
34d		¹ H-NMR (DMSO-D ₆ , 500 MHz): δ 10.48 (s, 1H), 10.33 (s, 1H), 8.94 (s, 1H), 8.42 (s, 1H), 8.32 (s, 1H), 8.21-8.12 (m, 2H), 7.84 (d, <i>J</i> = 8.5 Hz, 1H), 7.48-7.41 (m, 3H), 2.36 (s, 3H); m/z 382.8 [M+1] ⁺ .
34e		¹ H-NMR (DMSO-D ₆ , 500 MHz): δ 10.40 (s, 1H), 10.30 (s, 1H), 8.40 (s, 1H), 8.07 (s, 1H), 7.96 (s, 1H), 7.79 (d, <i>J</i> = 8.5 Hz, 1H), 7.72 (d, <i>J</i> = 8.5 Hz, 1H), 7.44-7.36 (m, 3H), 7.16 (d, <i>J</i> = 8.5 Hz, 1H), 2.33 (s, 3H); m/z 415.7 [M+1] ⁺ .

#	Structure	Characterization Data
34f		¹ H-NMR (DMSO-D ₆ , 500 MHz): δ 10.37 (s, 1H), 10.30 (s, 1H), 8.40 (s, 1H), 8.07 (s, 1H), 7.82-7.77 (m, 3H), 7.44-7.39 (m, 3H), 2.33 (s, 3H); m/z 415.8 [M+ 1] ⁺ .
34g		¹ H-NMR (DMSO-D ₆ , 500 MHz): δ 10.55 (s, 1H), 10.31 (s, 1H), 8.40 (s, 1H), 8.24 (s, 1H), 8.10-8.05 (m, 2H), 7.82 (d, J = 8.5 Hz, 1H), 7.61-7.58 (m, 1H), 7.45 (d, J = 8.5 Hz, 3H), 2.34 (s, 3H); m/z 449.8 [M+ 1] ⁺ .
34h		¹ H-NMR (DMSO-D ₆ , 500 MHz): δ 10.22 (bs, 1H), 8.43 (s, 1H), 8.39 (s, 1H), 7.94 (s, 1H), 7.62 (d, J = 8.5 Hz, 1H), 7.34 (d, J = 8.5 Hz, 1H), 2.76 (s, 3H), 2.28 (s, 3H); m/z 319.9 [M+ 1] ⁺ .
34i		¹ H NMR (DMSO-D ₆ , 500 MHz) δ 10.6 (s, 1H, D ₂ O exchangeable), 10.1 (s, 1H, D ₂ O exchangeable), 8.4 (s, 2H), 8.2-8.25 (d, 1H), 7.9-8.0 (d, 2H), 7.6-7.7 (d, 1H), 7.2-7.3 (d, 1H), 2.3 (s, 3H); MS: m/z 483.7 [M+1] ⁺ .
34j		¹ H-NMR (CD ₃ OD, 500 MHz): δ 8.83 (s, 1H), 8.33 (s, 1H), 8.32 (s, 1H), 7.89 (bs, 1H), 7.43 (d, J = 8.5 Hz, 1H), 2.36 (s, 3H), 1.38 (s, 9H); m/z 436.8 [M+1] ⁺ .
34k		¹ H-NMR (CD ₃ OD, 500 MHz): δ 8.32 (s, 2H), 7.88 (d, J = 8.0 Hz, 1H), 7.83 (s, 1H), 7.67 (d, J = 8.0 Hz, 1H), 7.47 (d, J = 8.0 Hz, 1H), 2.36 (s, 3H); m/z 446.8 [M+1] ⁺ .

[0370] Additional compounds of the present invention may be prepared according to general Scheme ZZ. Such compounds are set forth in **Table 4**, below.

Table 4. Exemplary Compounds of Formula I

Additional compounds			
1aA		1aB	
1aC		1aDa	
1aDb		1aE	
1aEa		1aEb	
1aF		1aG	
1aH		1aI	

Additional compounds			
1aJ		1aK	
1aL		1aMa	
1aMb		1aNa	
1aNb		1aO	
1aQ		1aRa	
1aRb		2aA	











Additional compounds			
2aB		2aC	
2aDa		2aDb	
2aE		2aEa	
2aEb		2aF	
2aG		2aH	
2aI		2aJ	
2aK		2aL	

Additional compounds			
2aMa		2aMb	
2aNa		2aNb	
2aO		2aP	
2aQ		2aRa	
2aRb		3aA	
3aB		3aC	

Additional compounds			
3aD		3aDa	
3aDb		3aE	
3aEa		3aEb	
3aF		3aG	
3aH		3aI	
3aJ		3aK	
3aL		3aMa	

Additional compounds			
3aMb		3aNa	
3aNb		3aO	
3aP		3aQ	
3aRa		3aRb	
4aA		4aB	
4aC		4aDb	

Additional compounds			
4aE		4aEa	
4aEb		4aF	
4aG		4aH	
4aI		4aJ	
4aK		4aL	
4aMa		4aMb	

Additional compounds			
4aNb		4aO	
4aP		4aQ	
4aRa		4aRb	
4ba		4bB	
4bC		4bDb	

Additional compounds			
4bE		4bEa	
4bEb		4bF	
4bG		4bH	
4bI		4bJ	
4bK		4bL	

Additional compounds			
4bMa		4bMb	
4bNa		4bNb	
4bP		4bQ	
4bRa		4bRb	
4eA		4eB	

Additional compounds			
4eC		4eDb	
4eE		4eEa	
4eEb		4eF	
4eG		4eH	
4eI		4eJ	

Additional compounds			
4eK		4eL	
4eMa		4eMb	
4eNa		4eNb	
4eO		4eP	
4eQ		4eRa	

Additional compounds			
4cRb		7aA	
7aB		7aC	
7aDa		7aDb	
7aE		7aEa	
7aEb		7aF	
7aG		7aH	

Additional compounds			
7aI		7aJ	
7aK		7aL	
7aMa		7aMb	
7aNa		7aNb	
7aO		7aP	
7aQ		7aRa	

Additional compounds			
7aRb		8bA	
8bB		8bC	
8bDa		8bDb	
8bE		8bEa	
8bEb		8bF	
8bG		8bH	















Additional compounds			
8bI		8bJ	
8bK		8bL	
8bMa		8bMb	
8bNa		8bNb	
8bO		8bP	
8bQ		8bRa	

Additional compounds			
8bRb		9B	
9C		9Da	
9Db		9E	
9Ea		9Eb	
9F		9G	
9H		9I	

Additional compounds			
9J		9K	
9L		9Ma	
9Mb		9Na	
9Nb		9O	
9P		9Q	

Additional compounds			
9Ra		9Rb	
10B		10C	
10Ea		10Eb	
10G		10H	
10I		10J	
10K		10L	

Additional compounds			
10Ma		10Mb	
10Na		10Nb	
10O		10Q	
10Ra		10Rb	
11A		11B	
11C		11Db	

Additional compounds			
11E		11Ea	
11Eb		11F	
11G		11H	
11I		11J	
11K		11L	
11Ma		11Mb	
11Na		11Nb	

Additional compounds			
11O		11P	
11Q		11Ra	
11Rb		12aA	
12aB		12aC	
12aD		12aDb	
12aE		12aEa	

Additional compounds			
12aEb		12aF	
12aG		12aH	
12aI		12aJ	
12aK		12aL	
12aMa		12aMb	
12aNa		12aNb	

Additional compounds			
12aO		12aP	
12aQ		12aRa	
12aRb		12bA	
12bB		12bC	
12bD		12bDb	
12bE		12bEa	

Additional compounds			
12bEb		12bF	
12bG		12bH	
12bI		12bJ	
12bK		12bL	
12bMa		12bMb	
12bNa		12bNb	

Additional compounds			
12bO		12bP	
12bQ		12bRa	
12bRb		5dA	
5dC		5dDa	
5dDb		5dE	
5dEa		5dEb	

Additional compounds			
5dF		5dG	
5dH		5dI	
5dJ		5dK	
5dL		5dMa	
5dMb		5dNa	

Additional compounds			
5dNb		5dO	
5dP		5dQ	
5dRa		5dRb	
15aA		15aB	
15aC		15aD	
15aDb		15aE	

Additional compounds			
15aEa		15aEb	
15aF		15aG	
15aH		15aI	
15aJ		15aK	
15aL		15aMa	
15aMb		15aNa	

Additional compounds			
15aNb		15aO	
15aP		15aQ	
15aRa		15aRb	
15bA		15bB	
15bC		15bD	
15bDb		15bE	

Additional compounds			
15bEa		15bEb	
15bF		15bG	
15bH		15bI	
15bJ		15bK	
15bL		15bM a	
15bMb		15bNa	















Additional compounds			
15bNb		15bO	
15bP		15bQ	
15bRa		15bRb	
15cA		15cB	
15cC		15cD	
15cDb		15cE	

Additional compounds			
15cEa		15cEb	
15cF		15cG	
15cH		15cI	
15cJ		15cK	
15cL		15cMa	
15cMb		15cNa	

Additional compounds			
15cNb		15cO	
15cP		15cQ	
15cRa		15cRb	
15dA		15dB	
15dC		15dD	
15dDb		15dE	
15dEa		15dEb	

Additional compounds			
15dF		15dG	
15dH		15dI	
15dJ		15dK	
15dL		15dM a	
15dMb		15dNa	
15dNb		15dO	
15dP		15dQ	

Additional compounds			
15dRa		15dRb	
10WW		8tD	
I-1		5tD	
I-2		I-3	
I-4		5cD	
I-5		I-6	
5ccD		I-7	

Additional compounds			
I-8		5ggD	
5ffD		5ccD	
5ddD		I-9	
I-10		I-11	
I-12		I-13	
I-14		5ttD	
5iiD		5vvD	

Additional compounds			
5hhD		I-15	
6mD		6oD	
6nD		33bD	
I-16		20aD	
I-17		I-18	
10S		10T	

[0371] In certain embodiments, the present invention provides a compound selected from those set forth in **Table 5**, below, where each compound # corresponds to a compound number as recited in **Table 3** or **Table 4**, *supra*.

Table 5. Selected Compounds of Formula I

#	Structure	Name
33bDa		(R)-N-(5-chloro-4-(trifluoromethyl)pyridin-2-yl)-2-(1-(6-(4-(trifluoromethyl)-1H-imidazol-2-yl)pyrimidine-4-carboxamido)ethyl)thiazole-5-carboxamide
26cD		N-(5-chloro-4-(trifluoromethyl)pyridin-2-yl)-2-(1-(6-(4-(2-hydroxyacetyl)piperazin-1-yl)pyrimidine-4-carboxamido)ethyl)thiazole-5-carboxamide
5vvD		2-(1-(6-(2-amino-2-oxoethylamino)pyrimidine-4-carboxamido)ethyl)-N-(5-chloro-4-(trifluoromethyl)pyridin-2-yl)thiazole-5-carboxamide
18cDa		(R)-N-(5-chloro-4-(trifluoromethyl)pyridin-2-yl)-2-(1-(6-(4-methyl-1H-imidazol-1-yl)pyrimidine-4-carboxamido)ethyl)thiazole-5-carboxamide
25bDa		(R)-N4-(1-(5-(5-chloro-4-(trifluoromethyl)pyridin-2-yl)carbamoyl)thiazol-2-yl)ethyl)pyrimidine-4,6-dicarboxamide

#	Structure	Name
25lDa		(R)-N4-(azetidin-3-yl)-N6-(1-(5-(5-chloro-4-(trifluoromethyl)pyridin-2-yl)carbamoyl)thiazol-2-yl)ethyl)pyrimidine-4,6-dicarboxamide
5vDa		(R)-N-(5-chloro-4-(trifluoromethyl)pyridin-2-yl)-2-(1-(6-(4-hydroxypiperidin-1-yl)pyrimidine-4-carboxamido)ethyl)thiazole-5-carboxamide
21		(R)-1-(6-(1-(5-(5-chloro-4-(trifluoromethyl)pyridin-2-yl)carbamoyl)thiazol-2-yl)ethyl)carbamoyl)pyrimidin-4-yl)piperidin-4-yl dihydrogen phosphate
19mmD		2-(1-(6-((R)-3-(aminomethyl)pyrrolidin-1-yl)pyrimidine-4-carboxamido)ethyl)-N-(5-chloro-4-(trifluoromethyl)pyridin-2-yl)thiazole-5-carboxamide
5wDa		N-(5-chloro-4-(trifluoromethyl)pyridin-2-yl)-2-((R)-1-(6-((R)-3-hydroxypyrrolidin-1-yl)pyrimidine-4-carboxamido)ethyl)thiazole-5-carboxamide
18fD		2-(1-(6-(1H-pyrazol-1-yl)pyrimidine-4-carboxamido)ethyl)-N-(5-chloro-4-(trifluoromethyl)pyridin-2-yl)thiazole-5-carboxamide

#	Structure	Name
25fD		N4-(2-aminoethyl)-N6-(1-(5-(5-chloro-4-(trifluoromethyl)pyridin-2-yl)carbamoyl)thiazol-2-yl)ethyl)pyrimidine-4,6-dicarboxamide
18dDa		(R)-N-(5-chloro-4-(trifluoromethyl)pyridin-2-yl)-2-(1-(6-(2-methyl-1H-imidazol-1-yl)pyrimidine-4-carboxamido)ethyl)thiazole-5-carboxamide
5dDa		(R)-N-(5-chloro-4-(trifluoromethyl)pyridin-2-yl)-2-(1-(6-morpholinopyrimidine-4-carboxamido)ethyl)thiazole-5-carboxamide
4qDa		(R)-2-(1-(4,5'-bipyrimidine-6-carboxamido)ethyl)-N-(5-chloro-4-(trifluoromethyl)pyridin-2-yl)thiazole-5-carboxamide
5aDa		(R)-N-(5-chloro-4-(trifluoromethyl)pyridin-2-yl)-2-(1-(6-(4-(2-hydroxyethyl)piperazin-1-yl)pyrimidine-4-carboxamido)ethyl)thiazole-5-carboxamide
15dDa		(R)-2-(1-(6-amino-5-(pyrrolidin-1-yl)methyl)pyrimidine-4-carboxamido)ethyl)-N-(5-chloro-4-(trifluoromethyl)pyridin-2-yl)thiazole-5-carboxamide

#	Structure	Name
5zDa		(R)-N-(5-chloro-4-(trifluoromethyl)pyridin-2-yl)-2-(1-(6-(4,4-Dioxothiomorpholin-1-yl)-pyrimidine-4-carboxamido)ethyl)thiazole-5-carboxamide
35		2-(1-(6-acetamidopyrimidine-4-carboxamido)ethyl)-N-(5-chloro-4-(trifluoromethyl)pyridin-2-yl)thiazole-5-carboxamide
18aDa		(R)-2-(1-(6-(1H-imidazol-1-yl)pyrimidine-4-carboxamido)ethyl)-N-(5-chloro-4-(trifluoromethyl)pyridin-2-yl)thiazole-5-carboxamide
5yDa		(R)-N-(5-chloro-4-(trifluoromethyl)pyridin-2-yl)-2-(1-(6-(4-(2-ethoxyethyl)piperazin-1-yl)pyrimidine-4-carboxamido)ethyl)thiazole-5-carboxamide
10E		2-(1-(6-amino-5-chloropyrimidine-4-carboxamido)ethyl)-N-(4-chloro-3-(trifluoromethyl)phenyl)thiazole-5-carboxamide
10Da		(R)-2-(1-(6-amino-5-chloropyrimidine-4-carboxamido)ethyl)-N-(5-chloro-4-(trifluoromethyl)pyridin-2-yl)thiazole-5-carboxamide

#	Structure	Name
10Db		(S)-2-(1-(6-amino-5-chloropyrimidine-4-carboxamido)ethyl)-N-(5-chloro-4-(trifluoromethyl)pyridin-2-yl)thiazole-5-carboxamide
1jDa		(R)-N-(5-chloro-4-(trifluoromethyl)pyridin-2-yl)-2-(1-(5-chloro-6-(methylamino)pyrimidine-4-carboxamido)ethyl)thiazole-5-carboxamide
5jDa		(R)-N-(5-chloro-4-(trifluoromethyl)pyridin-2-yl)-2-(1-(6-(methylamino)pyrimidine-4-carboxamido)ethyl)thiazole-5-carboxamide
17D		(R)-2-(1-(6-amino-5-cyanopyrimidine-4-carboxamido)ethyl)-N-(5-chloro-4-(trifluoromethyl)pyridin-2-yl)thiazole-5-carboxamide
11Da		(R)-2-(1-(6-amino-pyrimidine-4-carboxamido)ethyl)-N-(5-chloro-4-(trifluoromethyl)pyridin-2-yl)thiazole-5-carboxamide
4eDa		(R)-N-(5-chloro-4-(trifluoromethyl)pyridin-2-yl)-2-(1-(6-(2-fluoropyridin-3-yl)pyrimidine-4-carboxamido)ethyl)thiazole-5-carboxamide

#	Structure	Name
4bD		N-(5-chloro-4-(trifluoromethyl)pyridin-2-yl)-2-(1-(6-(pyridin-3-yl)pyrimidine-4-carboxamido)ethyl)thiazole-5-carboxamide
10D		2-(1-(6-amino-5-chloropyrimidine-4-carboxamido)ethyl)-N-(5-chloro-4-(trifluoromethyl)pyridin-2-yl)thiazole-5-carboxamide
10U		6-amino-5-chloro-N-(1-(2-(4-(trifluoromethyl)phenylamino)thiazol-5-yl)ethyl)pyrimidine-4-carboxamide
5aD		N-(5-chloro-4-(trifluoromethyl)pyridin-2-yl)-2-(1-(6-(4-hydroxypiperidin-1-yl)pyrimidine-4-carboxamido)ethyl)thiazole-5-carboxamide

Biological Assays

(1) Biochemical FRET assay

[0372] Method utilized for measuring the phosphorylation of MEK by wild-type (WT) B-Raf as a method for quantifying the ability of molecules to inhibit the enzymatic activity of WT-B-Raf.

[0373] In the assay methods described below, the following definitions apply:

“HEPES” refers to 4-(2-hydroxyethyl)-1-piperazineethanesulfonic acid;

“MEK” refers to mitogen activated extracellular signal-related kinase kinase;

“DTT” refers to dithiothreitol;

“APC” refers to allophycocyanin;

“TR-FRET” refers to time resolved fluorescence energy transfer;

“PBS” refers to phosphate buffered saline;

“PMSF” refers to phenyl methyl sulfonyl fluoride; and

“BSA” refers to bovine serum albumin.

Table 6. Reagents

Name	Units/Amount	Source	Catalog Number	Storage
Biotin-MEK1 (15:1)	DB021505 767 µg/mL (10.8 µM)	Biogen Idec.	In house	-80 °C
ATP	10mM, 500µl	Gibco BRL	8330-019	-20 °C
B-Raf (WT)	12µg/480µl 54% Pure (2.1µM)	Upstate	14-530M	-80 °C
DMSO	100%	Fisher	D128-500	RT
Streptavidin Allophycocyanin (SA-APC)	14.8uM SA (2.20 mg/ml)	Prozyme	PJ25S	4 °C, in the dark
Polyclonal Antiphospho MEK1/2(Ser 217/221) Antibody	265µg/ml (1.8uM)	Cell Signaling Technologies Inc.	9121	-20 °C
Lance Eu- W1024 Anti Rabbit IgG	880µg/ml (5.5µM)	Perkin Elmer	AD083	4 °C
LANCE 10X Detection Buffer	N/A	Perkin Elmer	CR97-100	4 °C
SuperBlock in TBS	N/A	Pierce	37535	4 °C

Table 7. Buffers

Master Buffer	Storage
50 mM HEPES, 60 mM NaCl, 3 mM MgCl ₂	4 °C
1M Dithiothreitol(DTT)	-20 °C in aliquots of 150µl
1M MnCl ₂	4 °C
20% BSA, 0.002% Sodium Azide.	4 °C
20% Tween-20	room temperature (~25 °C)
1M EDTA in dH ₂ O	room temperature (~25 °C)

[0374] **Equipment and Materials:** Analyst AD, LJI BioSystems, ID1615; 96 well $\frac{1}{2}$ Area Black Polystyrene plates. Costar 3694.

Table 8. Reagents

Reagents used for Kinase reaction:

50 μ M ATP

0.125 nM B-Raf (WT)

12.5 nM Biotin-MEK (15:1)

1% DMSO

50 mM Hepes, 60 mM NaCl, 3 mM MgCl₂, 2mM DTT, 0.25 mM MnCl₂, 0.01%BSA, 0.01% Tween-20

Reagents used for Detection Reaction

20nM SA-APC

2.5nM Polyclonal Anti p-MEK1/2 (Ser217/221)

2.5nM Eu-AntiRabbit IgG

1X Lance Detection Buffer

10% Superblock in TBS

WT Raf

[0375] Inhibitors were diluted 4-fold in 100% DMSO and added to a final concentration of 10 μ M to 40 pM to a solution containing 12.5 nM biotin-MEK, 0.125 nM WT Raf in 50 mM HEPES, pH 7.4, 60 mM NaCl, 3 mM MgCl₂, 2 mM DTT, 0.25 mM MnCl₂, 0.01% BSA, and 0.01% Tween-20 and incubated for 2 hours at room temperature. The kinase reaction was started by the addition of 50 μ M ATP to a final volume of 45 μ l and allowed to progress for 60 minutes. The reaction was stopped with 15 mM EDTA and 20 nM Streptavidin-APC, 2.5 nM Polyclonal anti p-MEK1/2 (Ser217/221), 2.5 nM Eu-labeled anti-rabbit IgG were added in Lance detection buffer and 5% Superblock in PBS for a final volume of 100 μ l. The detection reaction was incubated for 90 minutes at room temperature and then read on an Analyst plate reader using standard TR-FRET (time resolved fluorescence resonance energy transfer) settings for Eu and APC.

Mutant Raf

[0376] Inhibitors were diluted 4-fold in 100% DMSO and added to a final concentration of 10 μ M to 40 pM to a solution containing 100 nM biotin-MEK, 0.125 nM V599E Raf in 50 mM HEPES, pH 7.4, 60 mM NaCl, 3 mM MgCl₂, 2 mM DTT, 0.25 mM MnCl₂, 0.01% BSA, and 0.01% Tween-20 and incubated for 20 minutes at room temperature. The kinase reaction was started by the addition of 25 μ M ATP to a final volume of 45 μ l and allowed to progress for 60 minutes. The reaction was stopped with 15 mM EDTA and 20 nM Streptavidin-APC, 2.5 nM Polyclonal anti p-MEK1/2 (Ser217/221), 2.5 nM Eu-labeled anti-rabbit IgG were added in Lance detection buffer and 5% Superblock in PBS for a final volume of 100 μ l. The detection reaction was incubated for 90 minutes at room temperature and then read on an Analyst plate reader using standard TR-FRET (time resolved fluorescence resonance energy transfer) settings for Eu and APC.

C-Raf

[0377] Inhibitors were diluted 4-fold in 100% DMSO and added to a final concentration of 10 μ M to 40 pM to a solution containing 50 nM biotin-MEK, 0.075 nM C-Raf in 50 mM HEPES, pH 7.4, 60 mM NaCl, 3 mM MgCl₂, 2 mM DTT, 0.25 mM MnCl₂, 0.01% BSA, and 0.01% Tween-20 and incubated for 20 minutes at room temperature. The kinase reaction was started by the addition of 10 μ M ATP to a final volume of 45 μ l and allowed to progress for 60 minutes. The reaction was stopped with 15 mM EDTA and 20 nM Streptavidin-APC, 2.5 nM Polyclonal anti p-MEK1/2 (Ser217/221), 2.5 nM Eu-labeled anti-rabbit IgG were added in Lance detection buffer and 5% Superblock in PBS for a final volume of 100 μ l. The detection reaction was incubated for 90 minutes at room temperature and then read on an Analyst plate reader using standard TR-FRET (time resolved fluorescence resonance energy transfer) settings for Eu and APC.

[0378] Certain compounds of the present invention were assayed using the above Biochemical FRET assays and were found to be inhibitors of Raf kinase.

(2) Mechanistic Cellular Assay for Raf Kinase Activity

[0379] The following method was utilized for quantifying the amount of phospho-ERK in melanoma derived WM-266-4 cells (one allele each of wild type BRaf and mutant BRaf (V600D) as an indicator of Raf kinase activity in cells treated with various kinase inhibitors.

Table 9. Cellular Assay

Materials Needed	Catalog Number
WM-266-4 cells	(ATCC number: CRL-1676)
RPMI 1640 cell culture medium	
Fetal Bovine Serum (FBS)	
Phosphate Buffered Saline (PBS)	
96-well tissue culture plates	
Tissue culture 37°C incubator	
96-well V-bottom plates	
Rotary plate shaker (e.g., BELLCO GLASS Mini Orbital Shaker)	
Bio-Plex suspension array system	
Bio-Plex Cell Lysis Kit	(Bio Rad Catalog #171-304011)
Phenyl methyl sulphonyl fluoride (PMSF)	
Bio-Plex Phospho-ERK1/2 Assay Kit	(Bio Rad Catalog #171-V22238)

Day 1: Cell Seeding

(1) Detached adhered WM-266-4 cells from flask using 0.25% Trypsin. Resuspended cells in growth media (90% RPMI 1640, 10% FBS) and determine cell density.

(2) Seeded cells @ 10,000 cells/well in 96-well (flat bottom) tissue culture plates (36,000 cells/cm²). Added growth media to a final volume of 200uL/well and incubated overnight at 37°C.

Day 2: Cell Treatment

(1) Prepared compound dilutions (1000x in DMSO) as follows. Starting with a stock of 5mM compound in DMSO, diluted serially 3-fold in DMSO for a total of eight concentrations (5mM, 1.67 mM, 0.556 mM, 0.185 mM, 0.062 mM, 0.021 mM, 0.007 mM, 0.002 mM).

(2) Prepared compound-containing media by adding 1mL treatment media (100% RPMI 1640 without FBS) to 1μL of compound dilution (from step 3).

(3) Removed plates (from step 2) from incubator. Aspirated media and replace with 150 μL compound-containing media. Incubate for 1-2 hr at 37°C.

(4) Removed plates (from step 5) from incubator and treated each as follows: aspirated compound-containing media and replaced with 300 μL ice-cold 1xPBS, aspirated PBS and replaced with 45 μL lysis buffer (Biorad Bio-Plex lysis buffer containing 0.4% v/v lysis buff. Factor 1, 0.2% v/v lysis buff. Factor 2, and PMSF to 2mM final concentration), and then placed plate on ice until all plates were treated.

(5) After all plates were processed (step 6), placed plates on an orbital shaker and shook at room temperature for at least 15 min.

(6) Finally, removed plates from shaker, and transferred 40uL/well of lysate from each to new corresponding 96-well V-bottom plates. At this point, samples may be frozen and stored @ -80°C.

Day 2: Bioplex Assay

(1) Thaw (if necessary) plates (from step 8) and added 40 μL of Phospho-Protein Assay Buffer to each 40μL lysate for a 1:1 dilution.

(2) Prepared phospho-ERK1,2 Bioplex beads by diluting 1:50 with Bioplex Wash Buffer (mixing 49 μL Wash Buffer with 1μL of phospho-ERK1,2 Bioplex beads for each sample to be analyzed). Protected from light by wrapping tube in aluminum foil and kept at room temperature.

(3) Prepared Filter Plate by adding 100μL/well Bioplex Wash Buffer and removed by vacuum filtration.

(4) Add 50μL of bead solution (from step 10) to each well of a prepared Filter Plate (from step 11) and vacuum filter. Wash/filter 2x with 100uL/well Wash Buffer.

(5) Added 50μL of each lysate to appropriate well of the Filter Plate (from step 12). For this and all subsequent plate incubation steps, placed plate on an inverted plate cover (reduces background), and wrapped in aluminum foil (to protect from light). Shook overnight at room temperature. Included positive (control lysate) and negative (lysis buffer) controls.

Day 3: Bioplex Assay Continued

(1) Prepared detection antibody (phospho-ERK1,2 Ab) by diluting 1:25 with Detection Antibody Dilution Buffer Buffer (mixing 24 μ L Detection Antibody Dilution Buffer with 1 μ L of phospho-ERK1,2 Ab for each sample to be analyzed).

(2) Removed plate (from step 13) from shaker and vacuum filter. Washed/filter plate 3x with 100 μ L/well Wash Buffer. Added 25 μ L of diluted antibody to each well. Incubated on shaker at room temperature for 30-45min.

(3) Prepared streptavidin-PE by diluting 1:100 with Wash Buffer (mixing 49.5 μ L Wash Buffer with 0.5 μ L of 100x streptavidin-PE for each sample to be analyzed). Protected from light.

(4) Removed plate (from step 15) from shaker and vacuum filter. Washed/filter plate 3x with 100 μ L/well Wash Buffer. Add 50 μ L of diluted streptavidin-PE solution (from step 16) to each sample well. Incubated on shaker for 10-20min.

(5) Removed plate from shaker and vacuum filter. Wash/filter plate 3x with 100 μ L/well Bead Resuspension Buffer. After last wash resuspended beads in 125 μ L Bead Resuspension Buffer. Place plate on shaker for 2-3minutes to ensure beads are well resuspended.

(6) Quantified phospho-ERK by reading plate in the Bio-Plex plate reader (run start-up and calibration programs before this step) using bead region 38 (pERK1,2) and counting 50 beads per region.

[0380] Certain compounds of the present invention were assayed using the above Cellular Assay for Raf Kinase Activity and were found to be inhibitors of Raf kinase.

[0381] WM-266-4 cells were seeded at a density of 10,000 cells/well in RPMI 1640 cell culture media containing 10% FBS in a 96-well flat bottom and incubated overnight at 37°C. Inhibitors were diluted 3-fold in DMSO, added to serum free RPMI 1640 cell culture media to a final concentration range of 5 μ M to 2 nM, and used to treat the previously seeded WM-266-4 cells for 1-2 hours at 37°C. Cells were washed with ice-cold PBS, treated with 45 μ L of lysis buffer (Bio-Rad Bio-Plex Lysis Buffer, Cat # 171-304011, containing 0.4% v/v lysis buffer factor 1, 0.2% v/v lysis buffer Factor 2, and 2 mM PMSF) for 15 minutes on an orbital shaker at room temperature. Phosphorylated ERK was detected using a phospho-ERK Bioplex kit (Bio-Rad, Cat # 171-304011) per the manufacturer's instructions and detected on a Bio-Plex plate reader counting 50 beads per region.


[0382] Certain compounds of the present invention were assayed using the above cellular assays and were found to be inhibitors of Raf kinase.

[0383] While we have described a number of embodiments of this invention, it is apparent that our basic examples may be altered to provide other embodiments that utilize the compounds and methods of this invention. Therefore, it will be appreciated that the scope of this invention is to be defined by the appended claims rather than by the specific embodiments that have been represented by way of example.

CLAIMS

We claim:

1. A compound of formula I:



or a pharmaceutically acceptable salt thereof, wherein:

Cy^1 is an optionally substituted phenyl or an optionally substituted 5–6 membered saturated, partially unsaturated, or aromatic ring having 1–3 heteroatoms independently selected from nitrogen, oxygen, or sulfur;

Cy^2 is an optionally substituted 5–14 membered saturated, partially unsaturated, or aromatic monocyclic, bicyclic, or tricyclic ring having 0–4 heteroatoms, independently selected from nitrogen, oxygen, or sulfur;

L^1 is a direct bond or an optionally substituted, straight or branched C_{1-6} alkylene chain;

L^2 is a direct bond, or is an optionally substituted, straight or branched C_{1-6} alkylene chain wherein 1 or 2 methylene units of L^2 are optionally and independently replaced by $-O-$, $-S-$, $-N(R)-$, $-C(O)-$, $-C(O)N(R)-$, $-N(R)C(O)N(R)-$, $-N(R)C(O)-$, $-N(R)C(O)O-$, $-OC(O)N(R)-$, $-SO_2-$, $-SO_2N(R)-$, $-N(R)SO_2-$, $-OC(O)-$, $-C(O)O-$, or a 3–6 membered cycloalkylene;

each R is independently hydrogen or an optionally substituted C_{1-6} aliphatic group;

R^1 is hydrogen or an optionally substituted C_{1-6} aliphatic group;

each of R^x and R^y is independently selected from $-R^2$, $-\text{halo}$, $-\text{NO}_2$, $-\text{CN}$, $-\text{OR}^2$, $-\text{SR}^2$, $-\text{N}(\text{R}^2)_2$, $-\text{C}(\text{O})\text{R}^2$, $-\text{CO}_2\text{R}^2$, $-\text{C}(\text{O})\text{C}(\text{O})\text{R}^2$, $-\text{C}(\text{O})\text{CH}_2\text{C}(\text{O})\text{R}^2$, $-\text{S}(\text{O})\text{R}^2$, $-\text{S}(\text{O})_2\text{R}^2$, $-\text{C}(\text{O})\text{N}(\text{R}^2)_2$, $-\text{SO}_2\text{N}(\text{R}^2)_2$, $-\text{OC}(\text{O})\text{R}^2$, $-\text{N}(\text{R}^2)\text{C}(\text{O})\text{R}^2$, $-\text{N}(\text{R}^2)\text{N}(\text{R}^2)_2$, $-\text{N}(\text{R}^2)-\text{C}(=\text{NR}^2)\text{N}(\text{R}^2)_2$, $-\text{C}(=\text{NR}^2)\text{N}(\text{R}^2)_2$, $-\text{C}=\text{NOR}^2$, $-\text{N}(\text{R}^2)\text{C}(\text{O})\text{N}(\text{R}^2)_2$, $-\text{N}(\text{R}^2)\text{SO}_2\text{N}(\text{R}^2)_2$, $-\text{N}(\text{R}^2)\text{SO}_2\text{R}^2$, or $-\text{OC}(\text{O})\text{N}(\text{R}^2)_2$; and

each R^2 is independently hydrogen or an optionally substituted group selected from C_{1-6} aliphatic, a C_{6-10} monocyclic or bicyclic aryl ring, or a 5–10 membered saturated, partially

unsaturated, or aromatic monocyclic or bicyclic ring having 1–4 heteroatoms independently selected from nitrogen, oxygen, or sulfur, or

two R^2 on the same nitrogen are taken together with the nitrogen to form an optionally substituted 5–8 membered saturated, partially unsaturated, or aromatic ring having 1–4 heteroatoms independently selected from nitrogen, oxygen, or sulfur,

wherein monovalent substituents on a substitutable carbon atom of an “optionally substituted”

group are independently selected from halogen; $-(CH_2)_{0-4}R^\circ$; $-(CH_2)_{0-4}OR^\circ$; $-O-(CH_2)_{0-4}C(O)OR^\circ$; $-(CH_2)_{0-4}CH(OR^\circ)_2$; $-(CH_2)_{0-4}SR^\circ$; $-(CH_2)_{0-4}Ph$, which may be substituted with R° ; $-(CH_2)_{0-4}O(CH_2)_{0-1}Ph$ which may be substituted with R° ; $-CH=CHPh$, which may be substituted with R° ; $-NO_2$; $-CN$; $-N_3$; $-(CH_2)_{0-4}N(R^\circ)_2$; $-(CH_2)_{0-4}N(R^\circ)C(O)R^\circ$; $-N(R^\circ)C(S)R^\circ$; $-(CH_2)_{0-4}N(R^\circ)C(O)NR^\circ_2$; $-N(R^\circ)C(S)NR^\circ_2$; $-(CH_2)_{0-4}N(R^\circ)C(O)OR^\circ$; $-N(R^\circ)N(R^\circ)C(O)R^\circ$; $-N(R^\circ)N(R^\circ)C(O)NR^\circ_2$; $-N(R^\circ)N(R^\circ)C(O)OR^\circ$; $-(CH_2)_{0-4}C(O)R^\circ$; $-C(S)R^\circ$; $-(CH_2)_{0-4}C(O)OR^\circ$; $-(CH_2)_{0-4}C(O)SR^\circ$; $-(CH_2)_{0-4}C(O)OSiR^\circ_3$; $-(CH_2)_{0-4}OC(O)R^\circ$; $-OC(O)(CH_2)_{0-4}SR^\circ$; $SC(S)SR^\circ$; $-(CH_2)_{0-4}SC(O)R^\circ$; $-(CH_2)_{0-4}C(O)NR^\circ_2$; $-C(S)NR^\circ_2$; $-C(S)SR^\circ$; $-SC(S)SR^\circ$; $-(CH_2)_{0-4}OC(O)NR^\circ_2$; $-C(O)N(OR^\circ)R^\circ$; $-C(O)C(O)R^\circ$; $-C(O)CH_2C(O)R^\circ$; $-C(NOR^\circ)R^\circ$; $-(CH_2)_{0-4}SSR^\circ$; $-(CH_2)_{0-4}S(O)_2R^\circ$; $-(CH_2)_{0-4}S(O)_2OR^\circ$; $-(CH_2)_{0-4}OS(O)_2R^\circ$; $-S(O)_2NR^\circ_2$; $-(CH_2)_{0-4}S(O)R^\circ$; $-N(R^\circ)S(O)_2NR^\circ_2$; $-N(R^\circ)S(O)_2R^\circ$; $-N(OR^\circ)R^\circ$; $-C(NH)NR^\circ_2$; $-P(O)_2R^\circ$; $-P(O)R^\circ_2$; $-OP(O)R^\circ_2$; $-OP(O)(OR^\circ)_2$; SiR°_3 ; $-(C_{1-4}$ straight or branched alkylene) $O-N(R^\circ)_2$; or $-(C_{1-4}$ straight or branched alkylene) $C(O)O-N(R^\circ)_2$, wherein each R° may be substituted as defined below and is independently hydrogen, C_{1-6} aliphatic, $-CH_2Ph$, $-O(CH_2)_{0-1}Ph$, or a 4–6-membered saturated, partially unsaturated, or aryl ring having 0–4 heteroatoms independently selected from nitrogen, oxygen, or sulfur, or, notwithstanding the definition above, two independent occurrences of R° , taken together with their intervening atom(s), form a 3–12-membered saturated, partially unsaturated, or aryl mono- or bicyclic ring having 0–4 heteroatoms independently selected from nitrogen, oxygen, or sulfur which may be substituted as defined below; and

monovalent substituents on R° (or the ring formed by taking two independent occurrences of R° together with their intervening atoms), are independently selected from halogen, $-(CH_2)_{0-2}R^\bullet$, $-(haloR^\bullet)$, $-(CH_2)_{0-2}OH$, $-(CH_2)_{0-2}OR^\bullet$, $-(CH_2)_{0-2}CH(OR^\bullet)_2$; $-O(haloR^\bullet)$, $-CN$, $-N_3$,

$-(\text{CH}_2)_{0-2}\text{C}(\text{O})\text{R}^\bullet$, $-(\text{CH}_2)_{0-2}\text{C}(\text{O})\text{OH}$, $-(\text{CH}_2)_{0-2}\text{C}(\text{O})\text{OR}^\bullet$, $-(\text{CH}_2)_{0-2}\text{SR}^\bullet$, $-(\text{CH}_2)_{0-2}\text{SH}$, $-(\text{CH}_2)_{0-2}\text{NH}_2$, $-(\text{CH}_2)_{0-2}\text{NHR}^\bullet$, $-(\text{CH}_2)_{0-2}\text{NR}^{\bullet 2}$, $-\text{NO}_2$, $-\text{SiR}^{\bullet 3}$, $-\text{OSiR}^{\bullet 3}$, $-\text{C}(\text{O})\text{SR}^\bullet$, $-(\text{C}_{1-4}$ straight or branched alkylene) $\text{C}(\text{O})\text{OR}^\bullet$, and $-\text{SSR}^\bullet$ and suitable divalent substituents on a saturated carbon atom of R° are $=\text{O}$ and $=\text{S}$, wherein each $\text{R}^\bullet$ is unsubstituted or where preceded by "halo" is substituted only with one or more halogens, and is independently selected from C_{1-4} aliphatic, $-\text{CH}_2\text{Ph}$, $-\text{O}(\text{CH}_2)_{0-1}\text{Ph}$, or a 5–6-membered saturated, partially unsaturated, or aryl ring having 0–4 heteroatoms independently selected from nitrogen, oxygen, or sulfur; and

divalent substituents on a saturated carbon atom of an "optionally substituted" group are independently selected from $=\text{O}$, $=\text{S}$, $=\text{NNR}^{\bullet 2}$, $=\text{NNHC}(\text{O})\text{R}^\bullet$, $=\text{NNHC}(\text{O})\text{OR}^\bullet$, $=\text{NNHS}(\text{O})_2\text{R}^\bullet$, $=\text{NR}^\bullet$, $=\text{NOR}^\bullet$, $-\text{O}(\text{C}(\text{R}^{\bullet 2}))_{2-3}\text{O}-$, and $-\text{S}(\text{C}(\text{R}^{\bullet 2}))_{2-3}\text{S}-$, wherein each independent occurrence of R^* is selected from hydrogen, C_{1-6} aliphatic which may be substituted as defined below, or an unsubstituted 5–6-membered saturated, partially unsaturated, or aryl ring having 0–4 heteroatoms independently selected from nitrogen, oxygen, or sulfur; and

divalent substituents that are bound to vicinal substitutable carbons of an "optionally substituted" group include: $-\text{O}(\text{CR}^{\bullet 2})_{2-3}\text{O}-$, wherein each independent occurrence of R^* is selected from hydrogen, C_{1-6} aliphatic which may be substituted as defined below, or an unsubstituted 5–6-membered saturated, partially unsaturated, or aryl ring having 0–4 heteroatoms independently selected from nitrogen, oxygen, or sulfur; and

suitable substituents on the aliphatic group of R^* are independently selected from halogen, $-\text{R}^\bullet$, $-(\text{haloR}^\bullet)$, $-\text{OH}$, $-\text{OR}^\bullet$, $-\text{O}(\text{haloR}^\bullet)$, $-\text{CN}$, $-\text{C}(\text{O})\text{OH}$, $-\text{C}(\text{O})\text{OR}^\bullet$, $-\text{NH}_2$, $-\text{NHR}^\bullet$, $-\text{NR}^{\bullet 2}$, and $-\text{NO}_2$, wherein each $\text{R}^\bullet$ is unsubstituted or where preceded by "halo" is substituted only with one or more halogens, and is independently C_{1-4} aliphatic, $-\text{CH}_2\text{Ph}$, $-\text{O}(\text{CH}_2)_{0-1}\text{Ph}$, or a 5–6-membered saturated, partially unsaturated, or aryl ring having 0–4 heteroatoms independently selected from nitrogen, oxygen, or sulfur; and

substituents on a substitutable nitrogen of an "optionally substituted" group are independently selected from $-\text{R}^\dagger$, $-\text{NR}^{\dagger 2}$, $-\text{C}(\text{O})\text{R}^\dagger$, $-\text{C}(\text{O})\text{OR}^\dagger$, $-\text{C}(\text{O})\text{C}(\text{O})\text{R}^\dagger$, $-\text{C}(\text{O})\text{CH}_2\text{C}(\text{O})\text{R}^\dagger$, $-\text{S}(\text{O})_2\text{R}^\dagger$, $-\text{S}(\text{O})_2\text{NR}^{\dagger 2}$, $-\text{C}(\text{S})\text{NR}^{\dagger 2}$, $-\text{C}(\text{NH})\text{NR}^{\dagger 2}$, or $-\text{N}(\text{R}^\dagger)\text{S}(\text{O})_2\text{R}^\dagger$; wherein each $\text{R}^\dagger$ is independently hydrogen, C_{1-6} aliphatic which may be substituted as defined below, unsubstituted $-\text{OPh}$, or an unsubstituted 5–6-membered saturated, partially unsaturated, or

aryl ring having 0–4 heteroatoms independently selected from nitrogen, oxygen, or sulfur, or, notwithstanding the definition above, two independent occurrences of $R^{\dagger}$, taken together with their intervening atom(s) form an unsubstituted 3–12-membered saturated, partially unsaturated, or aryl mono- or bicyclic ring having 0–4 heteroatoms independently selected from nitrogen, oxygen, or sulfur; and substituents on the aliphatic group of $R^{\dagger}$ are independently selected from halogen, $-R^{\bullet}$, $-(\text{halo}R^{\bullet})$, $-\text{OH}$, $-\text{OR}^{\bullet}$, $-\text{O}(\text{halo}R^{\bullet})$, $-\text{CN}$, $-\text{C}(\text{O})\text{OH}$, $-\text{C}(\text{O})\text{OR}^{\bullet}$, $-\text{NH}_2$, $-\text{NHR}^{\bullet}$, $-\text{NR}^{\bullet}_2$, and $-\text{NO}_2$, wherein each $R^{\bullet}$ is unsubstituted or where preceded by “halo” is substituted only with one or more halogens, and is independently C_{1-4} aliphatic, $-\text{CH}_2\text{Ph}$, $-\text{O}(\text{CH}_2)_{0-1}\text{Ph}$, or a 5–6-membered saturated, partially unsaturated, or aryl ring having 0–4 heteroatoms independently selected from nitrogen, oxygen, or sulfur.

2. The compound according to claim 1, wherein each of R^x and R^y is independently selected from R^2 , halo, $-\text{OR}^2$, $-\text{N}(\text{R}^2)_2$, $-\text{OC}(\text{O})\text{R}^2$, $-\text{N}(\text{R}^2)\text{C}(\text{O})\text{R}^2$, $-\text{N}(\text{R}^2)\text{N}(\text{R}^2)_2$, $-\text{N}(\text{R}^2)\text{C}(\text{O})\text{N}(\text{R}^2)_2$, $-\text{N}(\text{R}^2)\text{SO}_2\text{N}(\text{R}^2)_2$, $-\text{N}(\text{R}^2)\text{SO}_2\text{R}^2$, or $-\text{OC}(\text{O})\text{N}(\text{R}^2)_2$.

3. The compound according to claim 2, wherein R^x is hydrogen, an optionally substituted C_{1-6} aliphatic group, or halo.

4. The compound according to claim 2, wherein R^y is selected from R^2 , $-\text{OR}^2$, or $-\text{N}(\text{R}^2)_2$.

5. The compound according to claim 4, wherein R^y is $-\text{NH}_2$, $-\text{NHCH}_3$, $-\text{NHCH}_2\text{CH}_3$, $-\text{NHCH}_2\text{CH}_2\text{CH}_3$, $-\text{NHCH}(\text{CH}_3)_2$, $-\text{NH}(\text{C}_3\text{H}_5)$, $-\text{NHCH}_2\text{CH}_2\text{CH}_2\text{OH}$, $-\text{N}(\text{CH}_2\text{CH}_2)_2\text{O}$, or $-\text{NHCH}_2\text{CH}_2\text{CH}_2\text{NH}(\text{CH}_3)_2$.

6. The compound according to claim 4, wherein R^y is an optionally substituted C_{1-6} aliphatic group.

7. The compound according to claim 6, wherein R^y is an optionally substituted group selected from C_{2-6} alkenyl or C_{2-6} alkynyl.

8. The compound according to claim 4, wherein R^y is an optionally substituted 5–10 membered saturated monocyclic or bicyclic ring having 0–4 heteroatoms independently selected from nitrogen, oxygen, or sulfur.

9. The compound according to claim 8, wherein R^y is an optionally substituted group selected from:

- (a) a 5-6 membered saturated monocyclic ring having 1-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur;
- (b) a 5-6 membered aromatic ring having 1-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur; or
- (c) an 8–10 membered saturated, partially unsaturated or aromatic bicyclic ring having 0–4 heteroatoms independently selected from nitrogen, oxygen, or sulfur.

10. The compound according to claim 8, wherein R^y is an optionally substituted group selected from phenyl, octahydroazocinyl, thiocyclopentanyl, thiocyclohexanyl, pyrrolidinyl, piperidinyl, piperazinyl, tetrahydrothiopyranyl, tetrahydrothiophenyl, dithiolanyl, tetrahydrofuranyl, tetrahydropyranyl, dioxanyl, thioxanyl, morpholinyl, oxathiolanyl, imidazolidinyl, pyrrolyl, pyrazolyl, imidazolyl, triazolyl, tetrazolyl, thiophenyl, furanyl, thiazolyl, isothiazolyl, thiadiazolyl, oxazolyl, isoxazolyl, oxadiazolyl, pyridinyl, pyrimidinyl, pyrazinyl, pyridazinyl, triazinyl, tetrazinyl, tetrahydropyridinyl, benzofuranyl, thianaphthenyl, pyrrolizinyl, indolyl, quinoliny, isoquinoliny, benximidazolyl, imidazopyridinyl, purinyl, indazolyl, pyrrolopyridinyl, cinnoliny, quinazoliny, phthalazinyl, naphthyridinyl, or quinoxalinyl.

11. The compound according to claim 1, wherein R^1 is hydrogen and L^1 is an optionally substituted, straight or branched C_{1-4} alkylene chain.

12. The compound according to claim 11, wherein L^1 is an optionally substituted, branched C_{1-4} alkylene chain.

13. The compound according to claim 1, wherein Cy^1 is an optionally substituted 5-membered saturated, partially unsaturated, or aromatic ring having 1–3 heteroatoms independently selected from nitrogen, oxygen, or sulfur.

14. The compound according claim 13, wherein Cy^1 is an optionally substituted pyrrolyl, pyrazolyl, imidazolyl, triazolyl, tetrazolyl, thiophenyl, furanyl, thiazolyl, isothiazolyl, thiadiazolyl, oxazolyl, isoxazolyl, or oxadiazolyl group.

15. The compound according to claim 1, wherein L^2 is a direct bond or an optionally substituted, straight or branched C_{1-4} alkylene chain wherein 1 or 2 methylene units of L^2 are replaced by $-O-$, $-S-$, $-N(R)-$, $-C(O)-$, $-C(O)N(R)-$, $-N(R)C(O)N(R)-$, $-N(R)C(O)-$, $-N(R)C(O)O-$, $-OC(O)N(R)-$, $-SO_2-$, $-SO_2N(R)-$, $-N(R)SO_2-$, $-OC(O)-$, or $-C(O)O-$.

16. The compound according to claim 15, wherein L^2 is $-C(O)N(R)-$, $-N(R)C(O)-$, $-SO_2N(R)-$, $-N(R)SO_2-$, $-OC(O)-$, or $-C(O)O-$.

17. The compound according to claim 16, wherein L^2 is $-C(O)N(H)-$ or $-N(H)C(O)-$.

18. The compound according to claim 1, wherein Cy^2 is an optionally substituted group selected from:

- (a) a 5-membered saturated, partially unsaturated, or aromatic monocyclic ring having 1–3 heteroatoms, independently selected from nitrogen, oxygen, or sulfur;
- (b) phenyl or a 6-membered saturated, partially unsaturated, or aromatic monocyclic ring having 1–4 heteroatoms, independently selected from nitrogen, oxygen, or sulfur; or
- (c) a 5–10 membered saturated, partially unsaturated, or aromatic bicyclic ring having 1–4 heteroatoms, independently selected from nitrogen, oxygen, or sulfur.


19. The compound according to claim 18, wherein Cy^2 is an optionally substituted group selected from:

- (a) a 5-membered heteroaryl ring having 1–3 heteroatoms, independently selected from nitrogen, oxygen, or sulfur;

- (b) phenyl or a 6-membered heteroaryl ring having 1–3 nitrogen atoms; or
 (c) a 5,6-fused bicyclic heteroaryl ring having 1-4 heteroatoms selected from oxygen, sulfur or nitrogen.

20. The compound according to claim 19, wherein Cy^2 is an optionally substituted group selected from phenyl, pyrrolyl, pyrazolyl, imidazolyl, triazolyl, tetrazolyl, thiophenyl, furanyl, thiazolyl, isothiazolyl, thiadiazolyl, oxazolyl, isoxazolyl, oxadiazolyl, pyridinyl, pyrimidinyl, pyrazinyl, pyridazinyl, triazinyl, tetrazinyl, pyrroliziny, indolyl, quinoliny, isoquinoliny, benzimidazolyl, imidazopyridinyl, indazolyl, purinyl, cinnoliny, quinazolinyl, phthalazinyl, naphthridinyl, quinoxalinyl, thianaphthenyl, or benzofuranyl.


21. The compound according to claim 1, wherein said compound is of formula II:



II

or a pharmaceutically acceptable salt thereof.

22. (Original) The compound according to claim 22, wherein said compound is of formula II-a or II-b:



II-a

II-b.

23. The compound according to claim 22, wherein Cy^1 is a 5-membered heteroaryl ring having 1-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur.

24. The compound according to claim 1, wherein said compound is selected from those depicted in Table 3, Table 4, or Table 5, or a pharmaceutically acceptable salt thereof.

25. A pharmaceutical composition comprising a compound according to claim 1 and a pharmaceutically acceptable carrier, adjuvant, or vehicle.

26. The composition of claim 25, in combination with a therapeutic agent selected from a chemotherapeutic or anti-proliferative agent, an anti-inflammatory agent, an immunomodulatory agent, a neurotrophic factor, an agent for treating cardiovascular disease, an agent for treating destructive bone disorders, an agent for treating liver disease, an anti-viral agent, an agent for treating blood disorders, an agent for treating diabetes, or an agent for treating immunodeficiency disorders.

27. Use of a compound according to claim 1 or a pharmaceutical composition thereof, in the manufacture of a medicament for inhibiting Raf kinase activity in a patient or a biological sample.

28. Use of a compound according to claim 1 or a pharmaceutical composition thereof, in the medicament for treatment or lessening the severity of a Raf-mediated disorder in a mammal, suffering such disorder, wherein the disorder is selected from a proliferative disorder, a cardiac disorder, a neurodegenerative disorder, an autoimmune disorder, a condition associated with organ transplant, an inflammatory disorder, an immunologically-mediated disorder, a viral disease, or a bone disorder.

29. The use according to claim 28, wherein the disorder is selected from melanoma, leukemia, colon cancer, breast cancer, gastric cancer, ovarian cancer, lung cancer, brain cancer, laryngeal cancer, cervical cancer, renal cancer, cancer of the lymphatic system, cancer of the genitourinary tract (including bladder cancer and prostate cancer), stomach cancer, bone cancer, lymphoma, glioma, papillary thyroid cancer, neuroblastoma, and pancreatic cancer.

30. Use of a compound according to claim 1 or a pharmaceutical composition thereof, and an additional therapeutic agent selected from a chemotherapeutic or anti-proliferative agent,


an anti-inflammatory agent, an immunomodulatory agent, a neutrophilic factor, an agent for treating cardiovascular disease, an agent for treating destructive bone disorders, an agent for treating liver disease, an anti-viral agent, an agent for treating blood disorders, an agent for treating diabetes, or an agent for treating immunodeficiency disorders, wherein:

said additional therapeutic agent is appropriate for disease being treated; and

said additional therapeutic agent in a single dosage form or as part of a multiple dosage form,

in the manufacture of a medicament for treatment of melanoma, leukemia, colon cancer, breast cancer, gastric cancer, ovarian cancer, lung cancer, brain cancer, laryngeal cancer, cervical cancer, renal cancer, cancer of the lymphatic system, cancer of the genitourinary tract (including bladder cancer and prostate cancer), stomach cancer, bone cancer, lymphoma, glioma, papillary thyroid cancer, neuroblastoma, and pancreatic cancer.

31. A method for preparing a compound of formula II-a':



II-a'

or a pharmaceutically acceptable salt thereof, wherein:

Cy¹ is an optionally substituted 5–6 membered saturated, partially unsaturated, or aromatic ring having 1–3 heteroatoms independently selected from nitrogen, oxygen, or sulfur;

Cy² is an optionally substituted 5–10 membered saturated, partially unsaturated, or aromatic monocyclic or bicyclic ring having 0–4 heteroatoms, independently selected from nitrogen, oxygen, or sulfur;

each of R^x and R^y is independently selected from -R², -halo, -NO₂, -CN, -OR², -SR², -N(R²)₂, -C(O)R², -CO₂R², -C(O)C(O)R², -C(O)CH₂C(O)R², -S(O)R², -S(O)₂R², -C(O)N(R²)₂, -SO₂N(R²)₂, -OC(O)R², -N(R²)C(O)R², -N(R²)N(R²)₂, -N(R²)-C(=NR²)N(R²)₂, -C(=NR²)N(R²)₂, -C=NOR², -N(R²)C(O)N(R²)₂, -N(R²)SO₂N(R²)₂, -N(R²)SO₂R², or -OC(O)N(R²)₂;

each R^2 is independently hydrogen or an optionally substituted group selected from C_{1-6} aliphatic, a C_{6-10} monocyclic or bicyclic aryl ring, or a 5–10 membered saturated, partially unsaturated, or aromatic monocyclic or bicyclic ring having 1–4 heteroatoms independently selected from nitrogen, oxygen, or sulfur, or


two R^2 on the same nitrogen are taken together with the nitrogen to form an optionally substituted 5–8 membered saturated, partially unsaturated, or aromatic ring having 1–4 heteroatoms independently selected from nitrogen, oxygen, or sulfur,

comprising the step of coupling a compound of formula **II-viii**:



II-viii


with a compound of formula **II-vii**:



II-vii

to form the compound of formula **II-a'**.

32. The method according to claim 31, wherein the compound of formula **II-vii** is prepared from a compound of formula **II-vi-b**:




II-vi-b

wherein A^- is a suitable chiral anion,


comprising the step of treating the compound of formula **II-vi-b** with a suitable base to form a compound of formula **II-vii**.

33. The method according to claim 32, wherein the compound of formula **II-vi-b** is prepared from a compound of formula **II-v**:



comprising the steps of:


(a) treating the compound of formula **II-v** with a chiral agent to form a compound of formula **II-vi-a**:




and

(b) separating the resulting diastereomers by suitable physical means to obtain a compound of formula **II-vi-b**.

34. The method according to claim 33, wherein the compound of formula **II-v**:




is prepared from a compound of formula **II-iv**:




comprising the step of converting the oxime moiety of formula **II-iv** to the amine group of formula **II-v**.

35. The method according to claim 34, wherein the compound of formula **II-iv** is prepared from a compound of formula **II-iii**:




comprising the step of treating the compound of formula **II-iii** with hydroxylamine to form the compound of formula **II-iv**.


36. The method according to claim 35, wherein the compound of formula **II-iii** is prepared by coupling a compound of formula **II-i**:



with a compound of formula **II-ii**:



37. A compound of formula **II-vi-a** or **II-vi-b**:



wherein:


A⁻ is a suitable chiral anion, wherein the anion is derived from a chiral acid selected from the group consisting of tartaric acid, a tartaric acid derivative, mandelic acid, malic acid, camphorsulfonic acid, Mosher's acid, ditoluoyl-D-tartaric acid, ditoluoyl-L-tartaric acid, camphorsulfonic acid (-); tartaric acid (+); malic acid (-); N-acetyl-L-leucine (-); di-toluloyl-L-tartaric acid (-); deoxycholic acid (+); quinic acid (-); camphoric acid (+); N-BOC-alanine

(-); tartaric acid (-); di-toluloyl-D-tartaric acid (+); camphorsulfonic acid (+); dibenzoyl-D-tartaric acid (+); L(+)-citramalic; S-acetyl mandelic acid (+); and BOC-isoleucine(+);

Cy¹ is an optionally substituted 5–6 membered saturated, partially unsaturated, or aromatic ring having 1–3 heteroatoms independently selected from nitrogen, oxygen, or sulfur; and

Cy² is an optionally substituted 5–10 membered saturated, partially unsaturated, or aromatic monocyclic or bicyclic ring having 0–4 heteroatoms, independently selected from nitrogen, oxygen, or sulfur.

38. A compound of formula **II-iv**:



wherein

Cy¹ is an optionally substituted 5–6 membered saturated, partially unsaturated, or aromatic ring having 1–3 heteroatoms independently selected from nitrogen, oxygen, or sulfur; and

Cy² is an optionally substituted 5–10 membered saturated, partially unsaturated, or aromatic monocyclic or bicyclic ring having 0–4 heteroatoms, independently selected from nitrogen, oxygen, or sulfur.