

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



FIGURE 1
PCT/GB/97/01908
10/1/98

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07B 63/04, C07C 7/20	A1	(11) International Publication Number: WO 98/02400 (43) International Publication Date: 22 January 1998 (22.01.98)
(21) International Application Number: ✓ PCT/GB97/01908 (22) International Filing Date: ✓ 15 July 1997 (15.07.97) (30) Priority Data: 9614854.9 ✓ 15 July 1996 (15.07.96) ✓ GB (71) Applicant (for all designated States except US): A.H. MARKS AND COMPANY LIMITED [GB/GB]; Wyke, Bradford, West Yorkshire BD12 9EJ (GB). ✓ (72) Inventors; and (75) Inventors/Applicants (for US only): PRYCE, Anthony [GB/GB]; 14 Alexandra Road, Pudsey, Leeds, West Yorkshire LS28 8BY (GB). CALLIN, Robert, Charles [GB/GB]; 4 Margerison Road, Ben Rhydding, Ilkley LS29 8QU (GB). ✓ (74) Agent: BROWNE, Robin, Forsythe; Urquhart-Dykes & Lord, Tower House, Merriam Way, Leeds LS2 8PA (GB).		(81) Designated States: AL, AM, AT, AU, AZ, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, GH, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published With international search report.
(54) Title: POLYMERISATION INHIBITOR (57) Abstract A monomer composition, stabilised against premature polymerisation, comprising: a) an ethylenically unsaturated monomer or mixture of monomers polymerisable by free radical initiation, and b) an effective amount, sufficient to inhibit premature polymerisation of component (a) of a mixture of: i) 1 to 99 % by weight, based on the total weight of components (i) and (ii) of a mixture of at least one aromatic amine and at least one organic acid in a molar ratio of 10:1 to 1:10, and ii) 99 to 1 % by weight based on the total weight of components (i) and (ii) of at least one stable radical compound.		