

Oil-containing washing composition

The invention relates to a transparent and moisturizing cosmetic wash compositions containing oil.

It is highly desirable for consumers to have cosmetic washes or cleansing compositions which
5 comprise natural oils since those oils can provide many different cosmetic benefits. The achieved effect directly depends on the respective natural-oil used.

It is highly desirable for consumers to have cosmetic washes or cleansing compositions which do not damage their skin. Typically, cosmetic wash preparations are aqueous formulations containing surfactants that can be delipidating or even irritating. Moreover, washes induce pH change to the skin
10 that can cause severe skin damage. Consumers are looking for a solution that would provide cleansing and foaming without drying out or damaging their skin. The second challenge is that cosmetic wash compositions are often aqueous formulations which makes the introduction of oils into these aqueous formulations especially challenging because natural oils providing cosmetic effects are immiscible with water resulting in bi-phasic products. One possible way to address that challenge is to
15 formulate oil-in-water emulsions. Another way is to formulate a bi-phasic product which has to be shaken before application wherein the shaking creates an unstable and turbid emulsion which separates over time again. Both options are well-known in the technical field of cosmetic wash and cleansing compositions.

Both options have significant drawbacks e.g. O/W and W/O emulsions are usually not transparent
20 often resulting in opaque primary packaging so that the consumer cannot see the actual formulation at the store at which she or he has to take a decision on which wash or cleansing product to buy. Bi-phasic formulations as described above always have to be shaken before application which is an additional step to carry out and may be seen as inconvenient. In addition, depending on the shaking behavior of each consumer the mixture may not be homogenous.

25 Thus, for consumers, homogeneous cosmetic wash or cleansing products which contain a cosmetically-beneficial plant-derived oil, are transparent and easily applicable with no additional preparation steps before application are highly desirable. In addition, a cosmetic wash that at the same time has moisturizing properties is even more desirable.

Researchers have surprisingly found that a cosmetic wash composition in form of a gel comprising:

- 30
- a. an anionic polysaccharide, and
 - b. a cationic polysaccharide, and
 - c. a pre-solubilized plant-derived oil, and
 - d. 1 to 8 surfactant(s), and
 - e. water

- 2 -

wherein the composition is free of inorganic and/or organic sulfates and free of polyethylene glycol and/or polyethylene glycol-derivatives and wherein the cosmetic wash composition has a viscosity of 0.5 to 5 Pa s, more preferably of 0.8 to 2.5 Pa s fulfills all the above objects.

All percentages, ratios and proportions herein are by weight, unless otherwise specified. All

- 5 temperatures are in degrees Celsius (°C) unless otherwise specified. The expression "room temperature" may be used synonymously for 20 °C. If any unit or condition is not explicitly defined in the context of the present application a person skilled in the art will always consider the most obvious unit or condition. If in the context of the application any unit or condition is missing, and this unit or condition is defined in another section of the application the same unit or condition applies where it is
- 10 missing.

The cosmetic wash composition according to the invention comprises an anionic polysaccharide. In a preferred embodiment the anionic polysaccharide is selected from xanthan gums. The anionic polysaccharide according to the invention which is preferably a xanthan gum generally has a

15 molecular weight of from 0.4 million to 25 million g mol⁻¹. Its viscosity generally ranges from 0.85 to 1,700 Pa s when measured at 25 °C using a 1% solution of the gum in 1% KCl, on a viscometer of the Brookfield LV type, at 60 rpm using Spindle No. 3.

Xanthan gum may be generally characterized as an anionic heteropolysaccharide, with a primary structure consisting of repeating penta-saccharide units formed by two glucose units, two mannose units, and a glucuronic acid unit. These repeating penta-saccharide units give xanthan gum its

20 characteristic backbone, which consists of (1→4) β-D-glucopyranosyl units substituted at C-3 on every other glucose residue with a charged trisaccharide sidechain. The trisaccharide sidechain consists of a D-glucuronic acid unit between two D-mannose units. Slightly less than half (about 40%) of the terminal D-mannose residues contain a pyruvic acid residue linked via keto groups to the four and six positions, and the D-mannose linked to the main chain mostly contains an acetyl group at position O-

25 6. Some side chains may be missing. The acetate and pyruvate contents are variable on the side chain, and depend on the bacterial strains and on the fermentation conditions used to produce the gum.

In a preferred embodiment the xanthan gum according to the invention has an average particle size of 50 to 1250 μm.

- 30 Xanthan gum is available from several commercial suppliers such as RT Vanderbilt Company, Jungbunzlauer International AG, Unionchem, KahlWax and CP Kelco. Examples of suitable xanthan gums are e.g. Keltrol®, Keltrol® CG-T, Keltrol® F, Keltrol® T, Keltrol® TF, Xantural® 180 and Vanzan® NF. For a person skilled in the art it is obvious that the above lists of suppliers and xanthan gums are non-exhaustive and not limiting the invention in any way.

- 3 -

The anionic polysaccharide which preferably is selected from xanthan gum is comprised in an amount of 0.001 to 5 % by weight of the total composition, more preferably 0.01 to 2.5 wt.-% and even more preferably 0.05 to 1 wt.-% of the total composition.

The cosmetic wash composition according to the invention also comprises a cationic polysaccharide.

5 In a preferred embodiment of the invention the cationic polysaccharide is selected from guar gums.

In general, for the purposes of the present invention, the term guar gum is defined as any guar gum containing cationic groups and/or groups which are ionizable into cationic groups in the cosmetic wash composition according to the invention and the pH range useful for a cosmetic wash composition. A pH range particularly relevant is from 4 to 10 and more preferably 4.5 to 6. The cationic groups are
10 chosen from those comprising primary amine groups, secondary, tertiary and/or quaternary.

Cationic guar gums used generally have a weight average molecular weight between 5×10^4 and 8×10^6 , and preferably between 10^3 and 3×10^6 g mol⁻¹.

According to the Invention, it is preferable to use a guar gum modified by hydroxypropyl or hydroxypropyl trimethylammonium groups. Such products are also sold under the trade names
15 JAGUAR Excel, JAGUAR HP105, JAGUAR C13 sec, JAGUAR the C 15, JAGUAR 17 and JAGUAR C162 by the companies, Solvay, BASF and Cosphatech LLC. For a person skilled in the art it is obvious that the above lists are non-exhaustive and do not limit the invention in any way. The most preferred guar gums are guar hydroxypropyl trimethylammonium chloride with the CAS number 65497-29-2 or hydroxypropyl guar with the CAS number 39421-75-5.

20 Cosmetic wash compositions according to the invention comprise the cationic polysaccharide in an amount of 0.001 to 10 % by weight of the total composition, more preferably 0.01 to 5 % by weight of the total composition, even more preferably 0.1 to 2.5 % by weight of the total composition.

The combination of the anionic and the cationic polysaccharides results in the formation of a specifically desired texture of the cosmetic wash composition which can be best described as a fluid
25 gel. In addition, only the combination of the anionic and the cationic polysaccharides together with the pre-solubilized plant-derived oil resulted in the formation of a transparent gel formulation, wherein usually the introduction of oils results in turbid formulations.

Without being bound by theory the inventors believe that the combination and the amounts of the polysaccharides as defined above are mainly responsible for the moisturizing properties of the
30 cosmetic washes of the invention. The inventors also observed that the plant-derived oil alone was not the key ingredient to explain the overall very good moisturizing properties of the cosmetic washes of the present invention, but rather the combination of all features leads to the moisturizing properties. The surfactant blend was also observed to have a supporting effect on the moisturizing properties of the cosmetic wash compositions of the invention.

- 4 -

As used herein a gel means a topical dosage form for dermatological application that is a semisolid dosage form that contains at least one gelling agent to provide stiffness to a solution or colloidal dispersion (Buhse et al., Int. J. Pharm. 2005, 295, 101–112). A gel may also comprise suspended particles. Gels according to this invention are pourable and flow at low shear stress and conform to their container at 20 °C. A conventional gel also contains more than 50 % by weight of water and other volatiles, wherein the content of water and volatiles can be measured by loss on drying test by heating at 105 °C until constant weight is achieved. Cosmetic wash compositions according to the invention preferably comprise water in an amount of at least 60 % by weight of the total composition, more preferably at least 50 % by weight of the total composition and even more preferable at least 40 % of the total composition.

In general, a gel has a specific viscosity range which is provided for the cosmetic wash composition by the anionic and the cationic polysaccharides. Thus, the cosmetic wash composition according to the invention has a viscosity from 0.5 to 5 Pa s. In a preferred embodiment the viscosity is from 0.8 to 2.5 Pa s.

A plant-derived oil according to the invention is an oil that is obtainable by extraction from plants. The plant-derived oil may be a complex mixture of multiple oil components or it may be a single oil component. Especially if the plant-derived oil is a single oil component or a mixture of a limited number of components the oil component(s) may be manufactured synthetically.

A plant-derived oil for use in the cosmetic wash compositions according to the invention are preferably selected from argan oil, castor oil, inca inchi oil, sweet almond oil, apricot oil, sunflower oil, olive oil, soy bean oil, safflower oil, cotton seed oil, corn oil, walnut oil, peanut oil, avocado oil, jojoba, palm oil, sesame oil, anise oil, lemon oil, orange oil, rosemary oil, wintergreen oil, thyme oil, lavender oil, cloves oil, hops oil, tea tree oil, citronella oil, wheat oil, barley oil, lemongrass oil, cedar leaf oil, cedarwood oil, cinnamon oil, fleagrass oil, geranium oil, sandalwood oil, violet oil, cranberry oil, eucalyptus oil, vervain oil, peppermint oil, basil fennel oil.

Plant-derived oils are highly effective substances that possess an array of different effects. The fragrance emanated from oils may affect the limbic system in the brain, e.g. by processing feeling. Therefore, some oils have a mood-enhancing effect. Individual molecules are transported into the bloodstream through the skin where they are integrated into the cell membrane, influencing bodily processes. As a result, these molecules can promote the renewal of cells or stimulate the metabolic processes. Plant-derived oils are primarily used in natural cosmetics due to their emollient, anti-inflammatory and anti-oxidant (vitamin E content) properties, and in addition they may have antibacterial and/or antiviral properties. They may also serve as natural preserving agent.

The term "pre-solubilized" in the context of the inventions and in particular in the context together with the expression "plant-derived oil" means that the plant-derived oil is solubilized in an aqueous phase so that the plant-derived oil and an aqueous phase build a mono phasic system. Plant-derived oils can for example be pre-solubilized by the process of saponification, neutralization, salification,

esterification and ionization of the plant-derived oil component(s) as it is done by the AQUASILOIL® technology marketed by Greentech GmbH. As an alternative also Floraesters® from Floratech, which also utilize chemically modified plant-derived oils may be regarded as pre-solubilized plant-derived oils. An advantage of these technologies is that the plant-derived oil itself acts as a surfactant-like compound, thus the amount of solubilizers or surfactants or the number and kind of different solubilizers and surfactants can be reduced which helps to decrease irritation risks and to decrease the environmental impact factor.

The preferred amount of pre-solubilized plant-derived oil in the cosmetic wash composition according to this invention is from 0.001 to 10 % of active by weight of the total composition, more preferred from 0.01 to 5 % of active matter by weight, even more preferred from 0.01 to 2.5 % of active matter by weight and most preferred from 0.01 to 1 % by weight of the total composition of active matter.

As mentioned above the combination of the mild surfactants, anionic and the cationic polysaccharide together with the pre-solubilized plant-derived oil results in a transparent and moisturizing aqueous gel containing a cosmetically beneficial oil.

In other cosmetic wash compositions, which are known in the prior art oils also have been used and some of them may also have been transparent. But these cosmetic wash compositions comprised additional solubilizer and especially polyethylene glycol and/or polyethylene glycol-derivatives. Polyethylene glycol and/or polyethylene glycol-derivatives like e.g. propylene glycol are deemed to have certain disadvantages especially for already damaged skin. Studies show that Polyethylene glycol and/or polyethylene glycol-derivatives have an enhanced penetration effect by altering the surface tension of the skin. Thus, PEGs may upset the natural moisture balance. In addition, PEGs are not always pure, but often come contaminated with a host of toxic impurities. Both effects taken together – the enhanced skin penetration of PEG containing composition and the impurities – may cause skin irritation or skin sensitization. As a consequence, the cosmetic wash compositions of the present invention are free of polyethylene glycol and/or polyethylene glycol-derivatives.

Since the compositions of the present invention are cosmetic wash compositions, they also comprise at least one surfactant. A wide variety of surfactants may be used, but preferred surfactants are chosen from anionic surfactants, amphiphilic surfactants and nonionic surfactants. It is even more preferred to use at least one anionic surfactant and one nonionic surfactant. In the most preferred embodiment of the invention the cosmetic wash composition comprises at least one glucoside surfactant.

Preferred anionic surfactants may be selected from disodium cocoyl glutamate, sodium lauryl sarcosinate, sodium stearte, acyl glycinates, sodium stearate or sodium gluconate.

Preferred amphiphilic surfactants may be selected from capryl/capramidopropyl betaine, sodium cocoamphoacetate or cocoamidopropylbetaine.

Preferred nonionic surfactants may be selected from caprylyl/capryl glucoside, lauryl glucoside or alkyl polyglucoside.

In a preferred embodiment the cosmetic wash composition according to the invention comprises surfactant(s) in an amount from 0.1 to 20 % by weight of the total composition, wherein the amount is based on the active material of the commercially available raw materials.

Cosmetic wash compositions according to the invention may contain at least one vitamin or vitamin derivative in an amount of 0.001 to 15 % by weight of the total composition, more preferably from 0.01 to 10 % by weight of the total composition and even more preferably from 0.1 to 7.5 % by weight of the total composition. Vitamins that are typically used in cosmetics are for example vitamin C (ascorbic acid and its derivatives), vitamin E (tocopherol and its derivatives), vitamin B3 (Niacinamide and derivatives), vitamin B5 (Panthenol and derivatives) or vitamin A (retinal, retinol, retinoic acid and retinyl palmitate and its derivatives).

Cosmetic wash compositions according to the invention comprise water in an amount of at least 60 % by weight of the total composition, more preferably at least 50 % by weight of the total composition and even more preferable at least 40 % of the total composition.

Cosmetic wash compositions according to the invention may be used to moisturize the skin. Moisturization is defined as an increase of the arbitrary units of corneometric measurements as carried out in the experimental section.

The cosmetic wash compositions of the present invention are of course not limited to the active ingredients as described above, but can comprise fillers, dyes, solvents, stabilizers, fragrancng agents, lubricants, humectants, thickeners or any other auxiliary useful for the present invention.

Examples

Formulas fit within the scope:

Ingredients (according to INCI)	1	2	3
Water	QSP		
CAPRYL/CAPRAMIDOPROPYL BETAINE	0	3% to 7%	0
COCAMIDOPROPYL BETAINE	0	0	3% to 7%
CAPRYL/CAPRYL GLUCOSIDE	1% to 3%	0	0
DISODIUM COCOYL GLUTAMATE	2% to 4%	2% to 4%	0
LAURYL GLUCOSIDE	2% to 5%	2% to 5%	2% to 5%
SODIUM COCOAMPHOACETATE	2% to 4%	0	2% to 4%
SODIUM GLUCONATE	0	0	2% to 4%
DEXPANTENOL	0% to 3%		
GLYCERINE	1% to 3%		
1,2-HEXANDIOL	1% to 2%		
GUAR HYDROXYPROPYLTRIMONIUM CHLORIDE	0.1% to 1%		
CITRIC ACID	0.1% to 6%		

- 7 -

NIACINAMIDE	0% to 1%		
LYSINE*	0.01% to 0.05%		
ARGANIA SPINOSA KERNEL OIL*	0.01% to 0.05%		
SODIUM CHLORIDE*	0.001% to 0.01%		
XANTHAN GUM	0.1% to 0.5%		
Viscosity (RS6000 cone/plan C60/1° Ti L shear rate 10 s ⁻¹)	0.8 to 2.5 Pa s	0.8 to 2.5 Pa s	0.8 to 2.5 Pa s
Appearance	Clear and homogeneous	Clear and homogeneous	Clear and homogeneous
Irritating potential	No	No	No

*These ingredients have been added in the form of a pre-solubilized oil commercially available under the brand name Aquasil Oil®

Formulas don't fit in the scope:

Ingredients (according to INCI)	4	5	6
Water	QSP		
CAPRYL/CAPRAMIDEOPROPYL BETAINE	3% to 7%	3% to 7%	3% to 7%
CAPRYL/CAPRYL GLUCOSIDE	0	0	0
DISODIUM COCOYL GLUTAMATE	2% to 4%	0	0
LAURYL GLUCOSIDE	2% to 5%	0	2% to 5%
SODIUM COCOAMPHOACETATE	0	0	2% to 5%
SODIUM LAURYL SULFATE	0	3% to 7%	0
DEXPANTENOL	1% to 3%		
GLYCERINE	1% to 3%		
1,2-HEXANDIOL	1% to 2%		
GUAR HYDROXYPROPYLTRIMONIUM CHLORIDE	0	0.1% to 1%	0
CITRIC ACID	0% to 6%		
NIACINAMIDE	0% to 2%		
LYSINE*	0	0.01% to 0.05%	0.01% to 0.05%
ARGANIA SPINOSA KERNEL OIL*	2% to 5%	0.001% to 0.01%	0.1% to 0.5%
SODIUM CHLORIDE*	0	0.001% to 0.01%	0.001% to 0.01%
HYDROXYETHYL ACRYLATE	0	0	0.1% to 0.5%
SODIUM ACRYLOYL DIMETHYL TAURATE COPOLYMER	0	0	0.1% to 0.5%
XANTHAN GUM	1%	0.1% to 0.5%	0.1% to 0.5%
Viscosity (RS6000 cone/plan C60/1° Ti L shear rate 10 s ⁻¹)	0.8 to 2.5 Pa s	0.8 to 2.5 Pa s	< 0 Pa s
Appearance	turbid	clear	turbid
Irritating potential	no	yes	no

5 *These ingredients have been added in the form of a pre-solubilized oil commercially available under the brand name Aquasil Oil®

Corneometry

10 The hydration level of the upper layers of the skin was tested via corneometry, a technique well established in the art and known by a person skilled in the art. Every commercially and calibrated

- 8 -

comeometer may be used for testing the hydration level since hydration level always is conducted against an untreated control measurement, which may be either the measurement right before applying a test composition for the first time to a skin area or parallel measurements of a treated and an untreated area of the skin over the complete test period.

5

The hydration level of the upper layers of the skin is expressed in arbitrary units (Corneometric indices) ranging from approximately 0 (no hydration) to 120 (very high hydration).

For each area and for each experimental time (T_i), the following individual data is calculated:

- means of the triplicate Corneometric indices,
- 10 • absolute variations (in comparison with T_0): $\Delta = T_i - T_0$,
- relative variations (in comparison with T_0): $\Delta\% = [(T_i - T_0) \times 100] / T_0$,
- Differences between treated area and control for each time point.

15 For the absolute variations in comparison with T_0 ($\Delta = T_i - T_0$), the following parameters are defined or calculated for treated area and negative control and for each experimental time T_i : number of subjects (n), mean, standard deviation, median, minimum, maximum.

A global comparison over time (ANOVA / Friedman test) of the instrumental measurements data for each test area will be performed. The ANOVA / Friedman test results in a p-value which is an indicator of significance. If the p-value of the ANOVA / Friedman test is $p \leq 0.05$ the result/effect is considered as being statistically relevant.

20

A total of 42 subjects (male or female) were included in the study and all subjects completed the study which means that all subjects applied the composition of the present invention twice a day for at least 14 days on the face for compositions according to example 1 and/or on a defined area on a forearm for composition according to example 2 above.

25 The 42 subjects included in the study had a mean age of 42 years (range from 42 to 70 years) and there were 95.2% women and 4.8% men and phototype II, III and IV represented 16.7%, 73.8% and 9.5%, respectively.

30 a) In this test, a comparison between a skin area that was treated with a composition according to example 2 with an untreated skin area was carried out. Given are the mean values of all 42 tested subjects and the mean difference between the treated and the untreated skin area. If a result is significant was calculated via the ANOVA / Friedman test and a p-value of < 0.05 was considered to be significant.

- 9 -

	mean value treated area	mean value untreated area	mean difference	p-value (significant at $p < 0.05$)	
D0	24.83	25.61	0.78	0.537	
D0/T1H	27.44	25.43	2.01	0.083	
D0/T2H	28.11	25.70	2.40	0.059	
D0/T5H	29.66	26.65	3.01	0.015	Significant difference
D1	27.79	25.44	2.35	0.074	
D7	27.81	24.95	2.86	0.030	Significant difference
D14	28.07	25.70	2.37	0.061	

- 5 b) In this test, the control was the corneometric measurement carried out right before applying the cosmetic wash of example 1 for the first time. Given is the mean value of the 42 subjects before the first application of the product, the mean difference and the p-value according to the ANOVA / Friedman test, wherein for this measurements results having a p-value of < 0.0125 were considered as statistically significant.

treated area vs. D0	Mean values	mean difference	p-value (significant at $p < 0.0125$)	
D0	33.31	--	--	--
D0/T1H-D0	--	14.05	<0.001	Significant increase
D0/T2H-D0	--	15.85	<0.001	Significant increase
D0/T5H-D0	--	18.13	<0.001	Significant increase
D1-D0	--	13.70	<0.001	Significant increase
D7-D0	--	16.25	<0.001	Significant increase
D14-D0	--	16.55	<0.001	Significant increase

- 10 -

Typically, cosmetic wash compositions in general tend to stress the skin e.g. because of the surfactants and other ingredients used to clean the skin which often leads to dry skin. Surprisingly, the cosmetic washes of the present invention replenish and maintain the skin's natural moisture level as can be seen in the corneometric measurements above.