

DESCRIPTION

NOVEL MICROORGANISM AND METHOD FOR
ENVIRONMENTAL PURIFICATION USING THE SAME

5 FIELD OF INVENTION

The present invention relates to new microorganisms which efficiently decompose organic halogenated compounds such as trichloroethylene, and to a method for decomposing organic halogenated compounds, specifically
10 trichloroethylene in the soil, the underground water, or waste waters.

BACKGROUND OF INVENTION

In recent years, industrial utilization of organic solvents has been producing environmental pollution
15 problems by discharge of these compounds or waste waters containing these compounds in many parts of the nation. In particular, soil pollution by organic chlorinated compounds is a major social problem, and the technology of repairing the contaminated soil has become more
20 essential. The purification method of the contaminated soil includes physical methods and biological methods.

The physical treatment methods include the air-stripping method (a method of purging air into the contaminated soil which was excavated in order to
25 volatilize organic chlorinated compounds contained therein, and of removing them by adsorption on activated charcoal), and vacuum extraction methods (a method in which pipes are driven into the contaminated soil to create the state of reduced pressure so that the organic
30 chlorinated compounds therein are volatilized and extracted from the soil). However, these methods require an enormous power for purging the air etc., and have the drawbacks that the former methods require excavation of the soil, while in the latter the extraction efficiency
35 is low and purification does not proceed smoothly under the low concentration of the contaminants. Furthermore, both methods only absorb the contaminating substances to

the activated charcoal and therefore require a separate means to detoxicate the contaminating substances.

It has been reported recently that the biological treatment method which is under development utilizes the ability of microorganisms to decompose substances and can completely decompose or detoxicate the contaminating substances, and besides less energy is needed for the treatment as compared with the physical means. Moreover, the biological means permits purification even at low concentrations of contaminants and accordingly expectations on the method are great as a low-cost method for soil purification. The known biological methods include the solid-phase treatment (the excavated soil is mixed with phosphorus, nitrogen, microorganisms etc. to promote decomposition of the contaminating substances by the microorganisms), the slurry treatment method (the excavated soil is mixed with water, phosphorus, nitrogen, microorganisms etc. to treat in the liquid form to promote purification speed of the contaminating substances by the microorganisms), and the on-site treatment method (methane, the air, phosphorus, and nitrogen are injected into the soil without excavating the soil to promote decomposition of the contaminating substances by the microorganisms).

Of the conventionally used biological treatment methods, the solid-phase treatment method and the slurry treatment method require excavation of the soil and besides have a narrow range of application, and the cost for treatment and equipment is relatively high.

On the other hand, the on-site treatment method in which indigenous microorganisms are performed as degraders is less expensive in treatment and equipment compared with the methods described above, and can be applied on a wider range. But, under the condition that the absolute number of microorganisms in the soil is small, the purification rate of the on-site treatment decreases. Especially in the case of the compounds

refractory to decomposition such as organic chlorinated compounds, purification is impossible when there are no living microorganisms which can decompose said contaminants in the soil. In such cases, it is believed that inoculation of the microorganisms having the ability of decomposing organic chlorinated compounds into the soil is essential for enhancement of the purification rate of soil.

Known microorganisms which decompose trichloroethylene include Methylosinus tricosporium OB3 (Japanese Unexamined Patent Publication (Kohyo) No. 4(1992)-501667, Japanese Unexamined Patent Publication No. 5(1993)-212371), and Methylosinus tricosporium TUKUBA (Japanese Unexamined Patent Publication No. 2(1990)-92274, Japanese Unexamined Patent Publication No. 3(1991)-292970) which are methane-degradating organisms, Pseudomonas putida F1 (Japanese Unexamined Patent Publication No. 64(1989)-34499), Pseudomonas putida BH (Fujita et al.; Chemical Engineering, 39(6):494-498, 1994), Pseudomonas putida UC-R5, UC-P2 (Japanese Unexamined Patent Publication No. 62(1987)-84780), Pseudomonas putida KWI-9 (Japanese Unexamined Patent Publication No. 6(1994)-70753), Pseudomonas mendocina KR1 (Japanese Unexamined Patent Publication No. 2(1990)-503866, 5(1993)-502593), Pseudomonas cepacia G4 (Japanese Unexamined Patent Publication No. 4(1992)-502277), and Pseudomonas cepacia KK01 (Japanese Unexamined Patent Publication No. 6(1994)-296711) which belong to the genus Pseudomonas, Alcaligenes eutropus JMP134 (A.R. Harker, Appl. Environ. Microbiol., 56(4):1179-1181, 1990), Alcaligenes eutropus KS01 (Japanese Unexamined Patent Publication No. 7(1995)-123976), Nitrosomonas europaea (D. Arciero et al., Biochem. Biophys. Res. Commun., 159(2):640-643, 1989) which is an ammonia-oxidizing bacterium, Corynebacterium J1 (Japanese Unexamined Patent Publication No. 8(1996)-66182) and the like.

The trichloroethylene-decomposing ability of these known microorganisms is not very high and most of these microorganisms can decompose 5 ppm of trichloroethylene in the liquid culture only. Furthermore, since
5 decomposing ability of trichloroethylene in a special environment as the soil is required, it is necessary that the microorganism to be used for bioremediation not only has a sufficient ability of decomposing trichloroethylene but also can remain decomposing ability of
10 trichloroethylene even in the soil. However, most of the known microorganisms are insufficient in this respect.

It is reported that Pseudomonas cepacia KK01 can decompose trichloroethylene at an initial concentration of 30 ppm to 15 ppm in the liquid culture, and
15 trichloroethylene at an initial concentration of 5 ppm to 1 ppm in the soil (Japanese Unexamined Patent Publication No. 6(1994)-296711). Furthermore, it is reported that Alcaligenes eutropus KS01 can decompose trichloroethylene at an initial concentration of 50 ppm to below the level
20 of detection in the liquid culture, and trichloroethylene at an initial concentration of 1 ppm to below the level of detection limit in the soil (Japanese Unexamined Patent Publication No. 7(1995)-123976).

It has been confirmed that these microorganisms have
25 a higher decomposing ability than the conventional microorganisms and that these abilities can be exhibited even in the soil. However, addition of at least one or more than one aromatic compound is needed to the soil environment for induction of the decomposing abilities of
30 these microorganisms. But, the aromatic compounds themselves are contaminants and therefore have a risk of causing a secondary pollution. It is a great challenge to be solved for practical application, therefore, to obtain a microorganism which enables an aromatic
35 compound, when added, to be completely decomposed and removed together with trichloroethylene, or which permits decomposition of trichloroethylene without addition of an

aromatic compound.

Accordingly, in order to put the biological purification of trichloroethylene into practical use, it has been desired to obtain a microorganism which has a high decomposing ability, and which enables an aromatic compound, when added, to be completely decomposed and removed together with trichloroethylene, or which permits decomposition of trichloroethylene without addition of an aromatic compound.

Furthermore, in many cases it is extremely difficult to increase the density of a microorganism to the level commensurate with its desired treatment capacity, because the density of the dispersed microorganism is suppressed low in the soil because of predation thereof by protozoa and competitive effects by autochthonous microorganism. In order to increase the density methods are employed such as the method of pressure pumping the air and nutrients into the soil. But despite the enormous energy required, it is difficult to increase the bacterial density by those means alone, thereby keeping the decomposing ability of microorganisms at low levels. Tremendous amounts of energy such as supply of nutrients, aeration etc. are needed to retain a high bacterial density in the closed system such as the reactor as well as in the open system.

If decomposition capability per unit amount of a bacterial mass is increased, a sufficient decomposition capability may be obtained even at low densities of the bacterial mass, thus obviating the need to put in a tremendous amount of energy for keeping the density of the bacterial mass. Though microorganisms which decompose trichloroethylene do so by expressing the enzyme capable of decomposing such substances, such expression of the enzyme requires an inducer. It has already been known that microorganisms can be allowed to exhibit their decomposition ability by adding an inducer and bringing the microorganisms into contact said inducer

during culturing, but no previous studies have focused on the length of time of the contact and thereby on the methods to enhance the decomposition ability per unit amount of bacterial mass.

5 Bio-augmentation which comprises spreading trichloroethylene-decomposing microorganisms into the soil to effect decomposition of trichloroethylene etc. is currently hard to get social acceptance, since it has a potential risk of producing far-reaching effects on the
10 ecological system by releasing a specific microorganism into the environment. But the spraying of a microorganism which has completely lost the propagating activity by sterilization treatment is equivalent to that of mere organic materials, and thus is believed to have
15 little effect on the ecological system. The invention which was disclosed in Japanese post-examined Patent Publication No. 8(1996)-3012 claims that undesirable effects on the ecological system can be minimized by crushing the decomposing bacteria and then spraying them
20 to the soil. But, it will be readily appreciated that the crushing procedure of microorganisms takes extensive equipment, a lot of time and labor, and thus the spraying of a large amount of decomposing bacterium to the contaminated soil will be in fact very difficult.

25 The above invention further lists the advantageous effects by slaiming that the crushed bacteria are easier to penetrate into the soil than the intact bacterial mass, but said invention makes no mention of duration of the decomposing ability retained by the crushed bacteria.
30 Moreover, the known trichloroethylene-oxidase requires NAD as a coenzyme. But it would be extremely difficult to supply the coenzyme in the concentration necessary for the decomposition reaction of the enzyme which is released from the bacterial mass by crushing the
35 bacterium because the coenzyme is very expensive.

DISCLOSURE OF THE INVENTION

Thus, it is an object of the present invention to

provide new microorganisms which can decompose organic halogenated compounds such as trichloroethylene in a more efficient manner than the conventionally known microorganisms, and a method for decomposing organic halogenated compounds utilizing said microorganism. It is another object of the present invention to provide an inexpensive method for minimizing the effects on the ecological system after a specific microorganism has been released to the environment.

After intensive studies to solve the above-mentioned problems the inventors have successfully isolated a new microorganism which does not belong to any of the known genera of microorganisms and which has a very high ability of decomposing trichloroethylene as compared with any of the conventionally known microorganisms, and thereby completed the present invention.

Thus, the present invention provides a bacterium which has the following taxonomical properties:

TABLE 1

20	Morphology	Coccoid, rod shaped
	Gram staining	+
	Spore forming	-
	Motility	-
	Relationship to oxygen	aerobic
25	Oxidase test	-
	Catalase test	+
	Resistance to acid	-
	Rod-coccus cycle	+
	Aerial mycelium forming	-
30	Peptide glycan type of cell wall	meso-diaminopimelic acid
	Glycolyl test	- (acetyl type)
	Mycolic acid	-
	Major isoprenoid quinones	MK-9, MK-9 (H2) (traces)
35	GC content of DNA (mole%)	73 (by HPLC)

THIS SHEET HAS BEEN
REPLACED

and, which can decompose trichloroethylene. A

This Sheet Has Been Replaced



representative strain of the bacterium is strain MO7 (FERM BP-5624).

5 The present invention further provides a method for decomposing organic halogenated compounds and/or aromatic compounds, said method comprising allowing said bacterium to act on said organic halogenated compounds and/or aromatic compounds. In this case, an organic halogenated compound is importantly trichloroethylene, and an aromatic compound is preferably phenol.

10 The present invention further provides a method for decomposing organic halogenated compounds in the soil, waste waters, or other waste products containing organic halogenated compounds, said method comprising adding a culture of said bacterium to said soil, waste waters, or
15 other waste products. An organic halogenated compound which is important in terms of practical application thereof is trichloroethylene.

The bacterial culture as used herein is preferably a cultured bacterial mass. The cultured bacterial mass may
20 be a living bacterial mass or a sterilized bacterial mass.

The present invention also provides a method for decomposing organic halogenated compounds in the soil, waste waters, or other waste products containing organic
25 halogenated compounds, said method comprising inoculating the bacterium of the present invention to said soil, waste waters, or other waste products, adding an aromatic compound or a degradable carbon source or a mixture thereof to said soil, waste waters, or other waste
30 products, and then culturing said inoculated microorganism. Said aromatic compound as used herein is preferably a phenolic compound, for example phenol, and said degradable carbon source as used herein is preferably a saccharide, for example glucose.

35 The present invention also provides a method for decomposing organic halogenated compounds and/or aromatic compounds, in which a sterilized cultured bacterial mass

of a microorganism which is capable of decomposing organic halogenated compounds and/or aromatic compounds is allowed to act on said organic halogenated compounds and/or said aromatic compounds. The present invention
5 also provides a method for decomposing organic halogenated compounds and/or aromatic compounds in the soil, waste waters, or other waste products containing organic halogenated compounds and/or aromatic compounds, in which a sterilized cultured bacterial mass of a
10 microorganism capable of decomposing said organic halogenated compounds and/or said aromatic compounds is added to said soil, waste waters, or other waste products. In the method described above, said sterilization treatment is, for example, ultraviolet
15 irradiation.

The present invention also provides a decomposition agent of organic halogenated compounds and/or aromatic compounds, said agent comprising said cultured bacterial cells of the bacterium of the present invention. Said
20 cultured bacterial mass may be a living bacterial mass or a sterilized bacterial mass. The sterilization treatment of them is, for example, ultraviolet irradiation. The bacterial mass contained in said decomposition agent is preferably in the form of a dried or freezed bacterial
25 mass from a viewpoint of storage.

BRIEF EXPLANATION OF THE DRAWINGS

Fig. 1 shows the position of strain M07 of the present invention in the phylogenetic tree constructed using the NJ (proximal conjugation) method.

30 Fig. 2 shows the position of strain M07 of the present invention in the phylogenetic tree constructed using the UPGMA (average distance) method.

Fig. 3 is a graph which shows the effect of the initial concentrations of trichloroethylene on the
35 decomposition efficiency, obtained using strain M07 of the present invention.

Fig. 4 is a graph which shows the effect of pH on

the decomposition efficiency of trichloroethylene (an initial concentration of 30 ppm), obtained using strain MO7 of the present invention.

5 Fig. 5 is a graph which shows the effect of the temperature on the decomposition efficiency of trichloroethylene (an initial concentration of 30 ppm), obtained using strain MO7 of the present invention.

10 Fig. 6 is a graph which shows the effect of the initial concentration of trichloroethylene on the decomposition efficiency of trichloroethylene (an initial concentration of 30 ppm), obtained using the dead bacterial cells (sterilized with ultra violet irradiation) of strain MO7 of the present invention.

15 Fig. 7 is a graph which shows a time course of the residual activity of decomposing trichloroethylene using the sterilized (with ultra violet irradiation) and the non-sterilized cultured bacterial cells of strain MO7 of the present invention.

20 Fig. 8 is a graph which shows growth curve (OD), time course of phenol consumption and the decomposition efficiency of trichloroethylene when strain MO7 of the present invention was cultured in a medium containing phenol and trichloroethylene.

25 Fig. 9 is a graph which shows growth curve (OD) and a time course of phenol consumption when strain MO7 of the present invention was cultured in a culture medium containing phenol.

30 Fig. 10 is a graph which shows the specific decomposition efficiency of trichloroethylene per bacterial cells (OD = 1.0) when the bacterium was cultured for various times in the time course of culture as set forth in Fig. 9.

35 Fig. 11 is a graph which shows the decomposition efficiency of trichloroethylene per total bacterial cells when the bacterium was cultured for various times in the time course of culture as set forth in Fig. 9.

Fig. 12 is a graph which shows growth curve (OD) and

a time course of phenol consumption when strain MO7 of the present invention was cultured in a medium containing phenol (500 ppm) and then further cultured by adding phenol when phenol was consumed completely.

5 Fig. 13 is a graph which shows the specific decomposition efficiency of trichloroethylene per bacterial cells (OD = 1.0) when the bacterium was cultured for various times in the time course of culture as set forth in Fig. 12.

10 Fig. 14 is a graph which shows the decomposition efficiency of trichloroethylene per total bacterial cells when the bacterium was cultured to various times in the time course of culture of Fig. 12.

15 Fig. 15 is a graph which shows the relationship between the amount of the bacterial cells and the amount of trichloroethylene decomposed when trichloroethylene was decomposed by adding the cultured bacterial cells of strain MO7 of the present invention into the soil containing trichloroethylene.

20 Fig. 16 is a graph which shows the amount of trichloroethylene decomposed when the cultured bacterial cells of strain MO7 of the present invention was added at one addition and at two additions.

25 Fig. 17 is a graph which shows the relationship between the amount of the inoculated bacterial cells and the decomposition efficiency of trichloroethylene at various initial concentration of trichloroethylene in soil when trichloroethylene was decomposed by adding the cultured bacterial mass of strain MO7 of the present invention.

30 Fig. 18 is a graph which shows a time course of the residual trichloroethylene in soil when trichloroethylene was decomposed by adding the cultured bacterial cells of strain MO7 of the present invention to the soil containing trichloroethylene.

35 Fig. 19 shows a flow chart of the analytical method of vinyl chloride, 1,1-dichloroethylene,

cis-1,2-dichloroethylene and trans-1,2-dichloroethylene.

Fig. 20 shows a flow chart of the analytical method of dichloroacetate and trichloroacetate.

Fig. 21 is shows a flow chart of the analytical
5 method of trichloroethanol.

Fig. 22 is shows a flow chart of the analytical method of hydrated chloral.

Fig. 23 is a graph which shows a result when trichloroethylene was decomposed by adding the cultured
10 bacterial cells of strain MO7 of the present invention or the sterilized product thereof to the soil containing trichloroethylene.

Fig. 24 is a graph which shows the effect of temperature (20°C, 30°C) on the decomposition efficiency
15 of trichloroethylene when trichloroethylene was decomposed by adding the cultured bacterial cells of strain MO7 of the present invention to the soil containing trichloroethylene.

Fig. 25 is a graph which shows a result when trichloroethylene was decomposed by adding a small amount
20 of the cultured bacterial cells of strain MO7 of the present invention and adding an inducer (phenol) to the soil containing trichloroethylene.

Fig. 26 is a graph which shows a result when trichloroethylene was decomposed by adding the cultured
25 bacterial mass of strain MO7 of the present invention or strain MO715 which were grown on glutamic acid to the soil containing trichloroethylene.

DETAILED DESCRIPTION

30 A representative bacterial strain of the present invention, strain MO7 may be isolated in the following manner. For example, an isolation source such as the soil or activated sludge is cultured in a culture medium containing phenol, and the microorganisms which
35 propagated therein are isolated. The isolates are subsequently incubated in a medium containing trichloroethylene to select the microorganisms which have

the ability of decomposing trichloroethylene. The method of isolation of microorganisms is described in detail in Example 1.

Strain MO7 thus isolated has the following
5 taxonomical properties:

TABLE 2

	Morphology	Coccoid, rod shaped
	Gram staining	+
	Spore forming	-
10	Motility	-
	Behavior to oxygen	aerobic
	Oxidase test	-
	Catalase test	+
	Resistance to acid	-
15	Rod-coccus cycle	+
	Aerial mycelium	-
	Peptide glycan type of cell wall	meso-diaminopimelic acid
	Glycolyl test	- (acetyl type)
20	Mycolic acid	-
	Major isoprenoid quinones	MK-9, MK-9 (H2) (traces)
	GC content of cellular DNA (mole%)	73 (by HPLC)

THIS SHEET HAS BEEN
REPLACED

25 Furthermore, strain MO7 has the following properties.

	Color of colonies	No characteristic colony color formed
	Elongation of cells surrounding colony	-
30	Arabinogalactan polymers of cell wall	- (#1)

#1: Estimated using an acid hydrolysate of the whole bacterial mass

35 The morphological and physiological characteristics of strain MO7 were investigated and the results obtained are shown in the above Table 2. Based on the results described above identification of the strain was

This Sheet Has Been Replaced



conducted with reference to the publication (N.R. Krieg and J.G. Holt, "Bergey's Manual of Systematic Bacteriology" Vol. 1 (1984), Williams and Wilkins; J.G. Holt, N.R. Krieg, P.H.A. Senath, J.T. Staley and
5 S.T. Williams, "Bergey's Manual of Systematic Bacteriology" Ninth edition (1984), Williams and Wilkins), with a result that the isolated strain MO7 did not conform to any known genus or species. Also 16S rRNA gene was
10 cloned and its sequence was compared with those of the known organisms to find that the closest relative thereto was *Terrabacter tumescens* as shown in Fig. 1 and 2. However, even the organism had a homology of 95% at most and thus it was concluded that the isolated strain does not belong to said genus. The bacterial strain of the
15 present invention, therefore, was confirmed to be a new genus which is different from any of the known bacteria.

In the method of the present invention, organic halogenated compounds and/or aromatic compounds are decomposed by allowing the bacterium of the present
20 invention to act on said organic halogenated compounds and/or said aromatic compounds. The organic halogenated compounds decomposed by the bacterium of the present invention include trichloroethylene, dichloroethylene, and vinyl chloride. From a practical point of view,
25 trichloroethylene is most important. As the aromatic compounds there are mentioned phenolic compounds, for example phenol.

In accordance with one embodiment of the present invention, there is provided a method for decomposing
30 organic halogenated compounds in the soil, waste waters, or other waste products, said method comprising adding the culture of the bacterium of the present invention to said soil, waste waters, or other waste products. The bacterial culture as used herein includes the culture
35 liquid itself obtained by culturing the bacterium of the present invention, a cultured bacterial mass isolated from said culture liquid, or a sterilized bacterium in

said culture liquid or a sterilized cultured bacterial mass which was isolated.

For culturing of the bacterium of the present invention, any culture medium in which said bacterium can propagate may be used. The culture medium may contain a carbon source such as glucose or sucrose, an organic nitrogen source such as yeast extract, peptone or meat extracts, or an inorganic nitrogen source such as an ammonium salt or nitrate. It may further contain inorganic salts comprising cations such as potassium ion, sodium ion, calcium ion, or magnesium ion, and anions such as chlorine ion, sulfate ion, or phosphate ion. The concentration of the carbon source, though varying depending upon the species, is in the range of about 0.1 to about 1%, and that of the nitrogen source, though varying depending upon the species, is in the range of about 0.01 to about 1%. And that of the inorganic salts, though varying depending upon the species, is in the range of about 0.001 to about 0.1%.

Culturing is preferably conducted by an aerobic liquid culture. The aerobic condition may be secured by a conventional means such as aeration, agitation, aeration and agitation, or shaking. In order to induce the decomposing ability of organic halogenated compounds such as trichloroethylene and that of aromatic compounds such as phenol, an aromatic compound such as phenol is preferably added to the culture medium. In this case, an aromatic compound such as phenol may be added in addition to the other carbon source, or an aromatic compound such as phenol may be added as the only carbon source. The amount of phenol etc. added to the medium is preferably in the range of about 100 ppm to about 1000 ppm.

In a case where the microorganism of the present invention is used in the form of a bacterial mass, said bacterial mass may be isolated by a conventional means of separating bacterial cells such as centrifugation. When the bacterial mass is stored prior to use, it may be

converted into the form of a freezed or dried bacterial mass. In this case, a conventional means for drying a bacterial mass may be employed such as freeze-drying, or spray-drying. In the practice of the present invention, a sterilized culture or a sterilized bacterial mass may be used in order to minimize its effect on the environment (the microbial phase). As a means of sterilization for this purpose, a conventional method such as ultraviolet irradiation to the living bacterial mass may be used. Such sterilized cells or cultures are also encompassed in the "culture" as defined in the present invention.

When the culture of the present invention is added to the subject to be treated, it is preferred to add a living bacterial mass in an amount of 10^6 to 10^9 cells/g of the subject to be treated, or a sterilized bacterial mass having the corresponding ability of decomposing organic halogenated compounds.

As a means to allow the microorganism of the present invention to act on an organic halogenated compound such as trichloroethylene, the culture of the present invention need only be added to and mixed with the subject to be treated in order to decompose the organic halogenated compounds contained in the solid such as the soil or the liquid such as waste waters (referred to herein as the subject to be treated).

In accordance with another embodiment of the present invention, organic halogenated compounds such as trichloroethylene contained in the subject to be treated can be decomposed by inoculating the microorganism of the present invention to the subject to be treated such as the soil, waste waters, or other waste products, and then allowing said organism to propagate therein. Thus, the present invention also provides a method for decomposing organic halogenated compounds in the soil, waste waters, or other waste products containing said organic

halogenated compounds, said method comprising inoculating the bacterium of the present invention to said soil, waste waters, or other waste products, adding an aromatic compound, an degradable carbon source, or a mixture thereof to the soil, waste waters, or other waste products, and then culturing said inoculated microorganism.

In this case, saccharides, for example glucose, is preferred as the degradable carbon source. The amount of such carbon source is about 0.1 to about 1% relative to the amount of the subject to be treated. Furthermore, when an aromatic compound is added to the subject to be treated, said aromatic compound need to be added. As an aromatic compound, phenol, cresol, and the like may be used. In this case, when the aromatic compound which was added remains after the decomposition of such organic halogenated compounds, it will cause another environmental pollution. Therefore, the addition of an excessive amount of the aromatic compound is not desirable and the amount of the aromatic compound to be added is preferably about 100 to about 500 ppm relative to the subject to be treated.

The inventors have found that the cultured bacterial mass of the present invention prepared as mentioned above, whether it is a living or a sterilized bacterial mass, retains the ability of decomposing organic halogenated compounds and/or aromatic compounds. Thus, the present invention provides a method for decomposing organic halogenated compounds and/or aromatic compounds in the soil, waste waters, or other waste products containing organic halogenated compounds and/or aromatic compounds, in which said sterilized cultured bacterial mass of a microorganism capable of decomposing organic halogenated compounds and/or aromatic compounds is added to said soil, waste waters, or other waste products.

The sterilization methods used includes Ultra violet irradiation, ethylene oxide treatment, radiation, and the

like. The sterilized product has an unexpected characteristics of having a higher stability during storage with regard to the ability to decompose organic halogenated compounds.

5 The present invention also provides a decomposition agent of organic halogenated compounds and/or aromatic compounds, said agent comprising a cultured fo the bacterium of the present invention. The culture is preferably a cultured bacterial mass, which may be a
10 living bacterial mass or a sterilized bacterial mass. Sterilization treatment is carried out by ultra violet irradiation, ethylene oxide treatment, radiation, or another method as described above. From a viewpoint of storage etc. the decomposition agent of the present
15 invention is preferably in the freezed or dried form, which dried product may be obtained according to a conventional method as mentioned above.

 The strain MO715, a representative mutant of the present invention (referred to hereinafter as the
20 constitutive mutant) which does not require an aromatic compound (such as phenol) for inducing the activity of trichloroethylene decomposition can be isolated as follows: the strain MO7 is mutated by the action of ultra violet, radiation, or a chemical substance such as
25 nitrosoguanidine which has a mutagenic activity and constitutive mutants are selected from the resulting mutants. The method of isolation of microorganisms is explained in detail in Example 18.

 The cultivation of the constitutive mutant of the
30 present invention may be carried out in the same manner as for the parent strain MO7 except that the cultivation of the former does not require an aromatic compound for the induction of the ability of decomposing trichloroethylene. The degradable carbon source for
35 cultivation is preferably a saccharide such as glucose, or an amino acid such as glutamic acid. The method of using it is also the same as for the parent strain MO7

except that the cultivation of the former does not require an aromatic compound for the induction of the ability of decomposing trichloroethylene. It should be noted that the living bacterial mass or the sterilized bacterial mass thereof has the same effect as the parent strain MO7 with regard to the ability of decomposing organic halogenated compounds.

The above-mentioned microorganism, the strain MO7, has been internationally deposited on August 12, 1996, with the National Institute of Bioscience and Human-Technology, Agency of Industrial Science and Technology, MITI, as FERM BP-5624 under the provisions of the Budapest Treaty. And the above-mentioned microorganism, the strain MO715, has also been internationally deposited on April 24, 1997, with the National Institute of Bioscience and Human-Technology, Agency of Industrial Science and Technology, MITI, as FERM BP-5928 under the provisions of the Budapest Treaty.

EXAMPLES

The invention will be understood more readily with reference to the following examples; however these examples are intended to illustrate the invention and are not to be construed to limit the scope of the invention.

Example 1. Isolation of the MO7 strain as well as cloning and sequencing of 16S rDNA thereof

The microorganisms for use in the present invention were isolated from the activated sludge taken at the waste water treatment plant in Aichi prefecture, Japan, in the following manner. A 0.1 ml aliquot of the harvested activated sludge collected was inoculated into 6 ml of the NMS medium containing 1% (v/v) of a vitamin solution (its composition is shown in Table 5) in a 30 ml vial, to which was added 500 ppm of phenol.

The vial was plugged with a butyl rubber septum and sealed with an aluminum cap, which was then cultured at 30°C under shaking at 160 r.p.m. for a period of from

several days to about a dozen days. The culture in which turbidity, even the slightest turbidity, was observed was passaged to the same medium and subsequently cultured under shaking. The passage was repeated for a total of
 5 four times. After the fourth passage is over, the culture medium was diluted as appropriate and plated onto the agar plate prepared by adding 1.5% agar to the NYG medium (its composition is shown in Table 5) containing
 10 500 ml phenol. The colonies which appeared were picked onto the agar plate and incubated. This operation was repeated for several times to isolate the microorganisms. In addition to the above NYG medium, another medium such as a nutrient medium can be used after selecting the
 15 optimum condition thereof for culturing of microorganisms.

TABLE 3

NMS medium

	Magnesium sulfate heptahydrate	1.0 g
	Calcium sulfate dihydrate	0.2 g
20	Potassium nitrate	0.23 g
	Ammonium sulfate	0.65 g
	Potassium dihydrogen phosphate	0.272 g
	Disodium hydrogen phosphate dodecahydrate	0.727 g
25	Trace Element Solution	0.5 ml
	Distilled water	1 liter

The Trace Element Solution

	EDTA disodium dihydrate	500 mg
	Iron (II) sulfate heptahydrate	200 mg
30	Zinc sulfate heptahydrate	10 mg
	Manganese (II) chloride tetrahydrate	3 mg
	Boric acid	30 mg
	Cobalt (II) chloride hexahydrate	20 mg
	Nickel (II) chloride hexahydrate	2 mg
35	Sodium molybdenum dihydrate	3 mg
	Distilled water	1 liter

* Trace Element Solution was separately sterilized and added after the other ingredients were sterilized.

TABLE 4

Vitamin solution		
5	Thiamine hydrochloride	3 mg
	p-amino benzoic acid	13 mg
	Adenine	1000 mg
	NAD	250 mg
	Vitamin B12	10 mg
10	Thiamine diphosphate	100 mg
	Distilled water	1 liter

TABLE 5

NYG medium		
15	Yeast Extract	0.5 g
	Glucose	0.18 g
	NMS medium	1 liter

20 The isolated microorganism was inoculated into 10 ml of the MNS medium containing 0.05% of yeast extract and 500 ppm of phenol in a 18 cm test tube by picking a colony on the agar plate for isolation using a platinum loop. After culturing at 30°C under shaking at 130 r.p.m. for 5 days, the bacterial mass was harvested by
25 centrifugation at 5000 r.p.m. for 10 minutes, and then resuspended into 4 ml of the NMS medium. The suspension of the bacterial mass was placed in a 20 ml vial and trichloroethylene was added in such an amount that it became 30 ppm after all ingredients were dissolved in the
30 liquid phase simultaneously. The vial was plugged with a Teflon-coated butyl rubber septum and sealed with an aluminum cap. After culturing overnight at 30°C, the gas phase in the vial was analyzed by a gaschromatograph equipped with a FID or an ECD detector.

35 From the results obtained, a bacterial strain having a high ability of decomposing trichloroethylene was selected and characterized for the morphological and

physiological properties to obtain the results as shown in Table 2.

Based on said results, identification was carried out with reference to the publication (N.R. Krieg and J.G. Holt, "Bergey's Manual of Systematic Bacteriology" Vol. 1 (1984) Williams and Wilkins, and J.G. Holt, N.R. Krieg, P.H.A. Senath, J.T. Stanley and S.T. Williams, "Bergey's Manual of Determinative Bacteriology" Ninth edition (1984) Williams and Wilkins) to find that this bacterial strain does not belong to any of the known genera or species. The bacterial strain was designated as the MO7 strain and was internationally deposited on August 12, 1996, with the National Institute of Bioscience and Human-Technology, Agency of Industrial Science and Technology, MITI, as FERM BP-5624 under the provisions of the Budapest Treaty.

Furthermore, the sequence of 16S rDNA of the microorganism of the present invention was determined and compared with that of the known microorganisms. The results are outlined hereinbelow.

The 16S rDNA of the microorganism of the present invention was amplified using PCR according to the method of Hiraishi (Journal of Japanese Society for Microbiological Ecology, vol. 10 (1): 31-42, 1995). The primers used for the PCR had the base sequence: 5'-GAG TTT GAT CCT GGC TCA G-3' (SEQ ID No: 1), and 5'-AGA AAG GAG GTG ATC CAG CGG CAG GTT-3' (SEQ ID No: 2). PCR was conducted using a thermal cycler (Perkin Elmer) according to the following program: i.e., preheating at 90°C for 30 seconds, 30 cycles of 96°C for 60 seconds/55°C for 120 seconds/72°C for 180 seconds, and heating at 72°C for 300 seconds.

The amplified PCR fragment was purified using QIAquick (Quiagen) and sequenced by a method which utilized a PCR directly as a template according to the method of Hiraishi (Journal of Japanese Society for Microbiological Ecology, vol. 10 (2): 81-102, 1995). The

primers used for the sequence reaction had the base sequence: 5'-GAG TTT GAT CCT GGC TCA G-3' (SEQ ID No: 1), 5'-GGC CGG ACG GGT GAG T-3' (SEQ ID No: 3), 5'-TAC GGG AGG CAG CAG-3' (SEQ ID No: 4), 5'-CTG CCA GCA GCC GCG CG-3' (SEQ ID No: 5), 5'-G ATT AGA TAC CCT GGT AG-3' (SEQ ID No: 6), 5'-ACT CAA AGG AAT TGA CGG-3' (SEQ ID No: 7), 5'-GCA ACG AGC GCA ACC C-3' (SEQ ID No: 8), or 5'-TGT ACA CAC CGC CCG T-3' (SEQ ID No: 9). PCR was conducted using the Dye terminator cycle sequencing kit (Perkin Elmer) according to the protocol instructed in the kit. Sequenced samples were subjected to electrophoresis and sequence analysis using the ABI373A Sequencer (Perkin Elmer). The base sequence determined by these procedures of the 16S rDNA of the microorganism of the present invention is shown in Table 6 (SEQ ID NO:10).

TABLE 6

Sequence	1466 BP;	352 A;	355 C;	288 T;	467 G.
	AACGCTGGCG	GCGTGCTTAA	CACATGCAAG	TGGAACGGTG	AAGCTTGGAG CTTGCTTCGA
	GTGGATCAGT	GGCGAACGGG	TGAGTAACAC	GTGAGCAACC	TGCCCCAGAC TCTGGAATAA
20	GCGCTGAAAA	CGGCGTCTAA	TACTGSATAT	GTGACGGACC	TGCATGGGTA CCGTCTGGAA
	AGTTTTTCGG	TTTGGGATGG	GCTCGCGGCC	TATCAGCTTG	TTGGTGAGGT AATGGCTCAC
	CAAGGCGACA	ACGGGTANCC	GGCCTGAGAG	GGCGACCGGC	CACACTGGCA CTGAAACACG
	GCCCCAACTC	CTACGGGAGG	CACCACTGGG	GAAATATTGC	ACAATGGGCG AAAGCCTGAT
	GCAGCGACGC	CGCGTGAGGG	ATGACGGCCT	TCGGGTTGTA	AACCTCTTTC AGCAGGGAAG
25	AAGCGAAAGT	GACGGTACCT	GCAGAATAAG	CACCGGCTAA	CTACGTGCCA GCAGCCGCGG
	TAATACGTAG	GGTGCGAGCG	TTGTCCGGAA	TTATTGGGCG	TAAAGAGCTT GTAGGCGGTT
	TGTCGCGTCT	GCTGTGAAAA	TCCGGGGCTC	AACCCCGGAC	TTGCAGTGGG TACGGGCAGA
	CTAGAGTGTG	GTAGGGGAGA	CTGGAATTCC	TGGTGTAGCG	GTGAAATGCG CAGATATCAG
	GAGGAACACC	GATGGCGAAG	GCAGGTCTCT	GGGCCACTAC	TGACGCTGAG AAGCGAAAGC
30	ATGGGGAGCG	AACAGGATTA	GATACCCCTG	TAGTCCATGC	CGTAAACGTT GGGCGCTAGG
	TGTGGGACTC	ATTCCACGAG	TTCCGTGCCG	CAGCTAACGC	ATTAAGCGCC CCGCCTGGGG
	CAGTACGGCC	GCAAGGCTAA	AACTCAAAGG	AATTGACGGG	GGCCCCCACA AGCGGCGGAG
	CATGCGGATT	AATTCGATGC	AACGCGAAGA	ACCTTACCAA	GGCTTGACAT ATACCGGAAA
	CTTCCAGAGA	TGTTTGCCCC	CTTTGGGTCG	GTATACAGGT	GGTGCATGGT TGTCGTCAGC
35	TCGTGTCGTG	AGATGTTGGG	TTAAGTCCCG	CAACGAGCGC	AACCCTCGTT CTATGTTGCC
	AGCACGTCAT	GGTGGGGACT	CATGGAAGAC	TGCCGGGGTC	AACTCGGAAG AAGGTGGGGA
	TGACGTCAAA	TCATCATGCC	CCTTATGTCT	TGGGCTTCAC	GCATGCTACA ATGGCCGGTA
	CAAGGGCTG	CGATACCGCA	AGGTGGAGCG	AATCCCCAAA	AAACCGGTCT CAGTTCGGAT
	TGGGGTCTGC	AACTCGACCC	CATGAAGTCG	GAGTCGCTAG	TAATCGCAAA TCAGCAACGC
40	TGCGGTGAAT	ACTTTCCCGG	GCCTTGATCA	CACCGCCCGT	CAAGTCACGA AAGTTCGGTA
	ACACCCGAAG	CCGGTGGCCC	AACCCTTGTG	GGGGGAGCCG	TC

The base sequence thus obtained was compared with the data in DNA database at the National Institute of Heredity (DDBJ+DDBJ NEW). Search of the homologous bacterial strains was conducted using the program "blast" to default value and at first, 50 species were selected which had the highest homology. From among them the bacterial strain (genus *Mycobacterium* which has the property of producing mycolic acid) which were believed to be definitely different from the results of physiological taxonomic tests were excluded. Then the multiple alignment analysis with the rest of the bacterial strains and the type strains of the genera which are thought to be closely related were conducted.

From the results obtained, a phylogenetic tree was constructed using the NJ (proximal conjugation) method (Fig. 1) or the UPGMA (average distance) method (Fig. 2) (alignment was conducted after the sequences at the 5'-ends were aligned). The most closely related organism turned out to be *Terrabacter tumescens* (the phylogenetic tree includes *Mycobacterium sedimentalis* for reference). The closely related bacterial strains (microorganisms belonging to the same cluster) were subjected again to homology search using Genetryx-Mac 8.0. But, the homology of the gene of even the closest *Terrabacter tumescens* was up to 95% and it was judged not to be the same genus. Therefore, it was confirmed that the microorganism of the present invention is a new microorganism different from any of the known microorganism.

The microorganism decomposes about 50% of trichloroethylene at high concentrations of about 100 ppm contaminated in the culture medium, and completely decomposes about 30 ppm of trichloroethylene in 24 hours. In order to decompose trichloroethylene in association with the propagation of the microorganism, it is necessary to add at least one aromatic compound such as phenol in the culture medium (medium, soil, water, etc.)

containing trichloroethylene.

The microorganism for use in the present invention will be explained in more detail with reference to the following examples.

5 Example 2.

Into 100 ml of the NMS medium containing 0.05% yeast extract and 500 ppm of phenol contained in a 500 ml Erlenmeyer flask was inoculated with a platinum loopful of the colony of the microorganism of the present
10 invention that had been stored by passage on the agar plate of 1.5% agar added to the NYG medium containing 500 ppm phenol, or 1.0 ml of the preculture liquid obtained by culturing overnight under shaking the microorganism of the present invention in the NYG medium containing 500
15 ppm of phenol at 30°C. After culturing under shaking at 30°C at 130 r.p.m. for 3 days, the bacterial mass was harvested by centrifugation at 5000 r.p.m. for 10 minutes and then resuspended into the NMS medium having an amount equal to the culture medium.

20 The OD₆₀₀ of the suspension was 0.5 (the cell count was 1×10^9 c.f.u./ml). 4 ml of the suspension of the bacterial mass was dispensed in a 20 ml vial and trichloroethylene was added thereto in such an amount that it became 10, 50, or 30 ppm after all the
25 ingredients were dissolved in the liquid phase. The vial was plugged with a Teflon-coated butyl rubber septum and sealed with an aluminum cap. After culturing overnight at 30°C under shaking, the gas phase in the vial was regularly analyzed by a gaschromatograph equipped with an
30 ECD detector.

The result is shown in Fig. 3. The present microorganism completely decomposed trichloroethylene in 24 hours. Trichloroethylene at such a high concentration of 50 ppm and 100 ppm was decomposed by 70% and 50%,
35 respectively in 24 hours. Even when the initial concentration of trichloroethylene as high as 50 and 100

ppm, the amount of trichloroethylene decomposed by the present microorganism was not reduced as compared to that when the initial concentration of trichloroethylene was 30 ppm, indicating that the present microorganism is resistant to decomposition inhibition even in the presence of a high concentration of trichloroethylene. Since there has been no reports which demonstrate that as high as 100 ppm of trichloroethylene was decomposed by a microorganism, the decomposition ability of the present microorganism is believed to be extremely high.

Example 3.

The bacterial mass cultured in the same method as in Example 2 was harvested by centrifugation at 5000 r.p.m. for 10 minutes and then resuspended to each of the M9 medium ($\text{Na}_2\text{HPO}_4 \cdot 7\text{H}_2\text{O}$ 12.8 g/l, KH_2PO_4 3 g/l, NaCl 0.5 g/l, NH_4Cl 1.0 g/l) prepared in varying pH's at an amount equal to that of the culture liquid. 4 ml of the suspension of the bacterial mass was dispensed in a 20 ml vial and trichloroethylene was added thereto in such an amount that it became 30 ppm after all the ingredients were dissolved in the liquid phase. The vial was plugged with a Teflon-coated butyl rubber septum and sealed with an aluminum cap. After culturing overnight at 30°C under shaking, the gas phase in the vial was regularly analyzed by a gaschromatograph equipped with an ECD detector. As shown in Fig. 4, the result indicated that at pH 5 or lower the activity is extremely decreased, whereas at pH 6 to 9 the efficiency of decomposition was as high as 100% and the activity decreased very slightly to about 90% even at pH 10.

Thus, it turned out that the optimum pH of trichloroethylene decomposition by a resting bacterial mass of the MO7 strain was between 6 and 9, indicating that a high decomposition activity is exhibited at high pH conditions. Trichloroethylene oxide produced by aerobic decomposition of trichloroethylene spontaneously

decomposes at an alkaline environment, producing
innocuous glyoxylic acid etc. On the other hand, in an
acid condition it spontaneously decomposes to produce
halo acid. Accordingly, the aerobic decomposition of
5 trichloroethylene by a microorganism is preferably
conducted at an alkaline environment. The present
microorganism is very useful for decomposition of
trichloroethylene at an alkaline environment, since it
shows a high decomposition activity at a high pH
10 condition.

Example 4.

The bacterial mass cultured in the same method as in
Example 2 was harvested by centrifugation at 5000 r.p.m.
for 10 minutes and then suspended into the same volume of
15 the NMS medium as the culture liquid. 4 ml of the
suspension of the bacterial mass was dispensed in a 20 ml
vial and trichloroethylene was added thereto in such an
amount that it became 30 ppm after all the ingredients
were dissolved in the liquid phase. The vial was plugged
20 with a Teflon-coated butyl rubber septum and sealed with
an aluminum cap. It was cultured under shaking in an
incubator each set at a different temperature. The
residual concentration of trichloroethylene was regularly
monitored by analyzing the gas phase in the vial with a
25 gaschromatograph equipped with an ECD detector.

As shown in Fig. 5, the result indicated that the
decomposition of trichloroethylene proceeded slowly over
7 days at 4°C. At 37°C, the decomposition ceased in a
day and the decomposition efficiency was low. On the
30 other hand, 80 to 90% was decomposed in a day at 20°C and
30°C. But decomposition activity was about 10% higher at
20°C than at 30°C. The temperature of the underground
soil where contamination by trichloroethylene etc. is
causing a problem is said to be stable at 15 to 20°C.
35 However, in most of the reports on the
trichloroethylene-decomposing microorganisms
decomposition activity was evaluated at 30°C which is

higher than the temperature of the soil, and very few studies have demonstrated that decomposition activity at the same temperature as the soil environment was maintained at a similar level to that at the evaluation experiments.

Since the reaction rate of the enzymatic reaction representing the basic reaction of microbial decomposition usually decreases by one-half with a temperature reduction of 10°C, it is estimated that the decomposition rate of trichloroethylene by a microorganism in the soil having a temperature lower than that at the evaluation experiment is reduced. The present microorganism was demonstrated to have a high practical characteristics without showing reduction in decomposition activity at the same temperature as the soil environment, though the mechanism thereof is unknown.

Example 5.

The bacterial mass cultured in the same method as in Example 2 was harvested by centrifugation at 5000 r.p.m. for 10 minutes and then suspended to the same volume of the NMS medium as that of the culture liquid. The suspension of the bacterial mass was placed in a petri dish and spread to a thickness of about 1 mm and then sterilized under irradiation of a 15 W ultra violet lamp with a wavelength of 260 nm for not less than 60 seconds at a distance of 40 cm from the light source. 4 ml of the suspension of bacterial mass after sterilization was dispensed in a 20 ml vial and trichloroethylene was added thereto in such an amount that it became 30 ppm after all the ingredients were dissolved in the liquid phase. The vial was plugged with a Teflon-coated butyl rubber septum and sealed with an aluminum cap. It was cultured under shaking at 30°C, and the gas phase in the vial was regularly analyzed with a gaschromatograph equipped with an ECD detector.

As shown in Fig. 6, the result shows that the

bacterial mass after sterilization retains 80% or more of the trichloroethylene-decomposition activity of the living bacterium. Furthermore, in an experiment in which the suspension of sterilized bacterial mass at an amount of 1/100 that of the NYG medium was inoculated or the suspension was plated as it was onto the NYG agar medium and cultured at 30°C, neither increase in turbidity of the culture liquid nor colony formation were observed, and thereby it was confirmed that there was complete sterilization.

The residual activity of the suspension of the sterilized bacterial mass when stored at 4°C was almost the same as that of the living bacterial mass at three days of storage and said suspension had a much higher activity than that of the living bacterial mass at seven days of storage. The results reveal that although the initial activity of the dead bacterial mass is decreased to 80% of the living bacterial mass, a higher activity of decomposing trichloroethylene is retained than the living bacterial mass during storage because of a lower reduction in activity of the dead bacterial mass during storage.

By sterilizing a microorganism without destructing the boundary structure such as the cell wall of a bacterial mass with the extracellular environment, it is possible to retain coenzymes etc. which are required for decomposition reaction of trichloroethylene at concentrations necessary for expression of enzymatic activity. Thus, supply of coenzymes for maintenance of activity of trichloroethylene decomposition is not necessary and the spread of the cultured bacterial mass is only needed. It possible therefore to effect purification of trichloroethylene etc. at low cost and with a minimum effect on the ecological system.

Example 6.

To 4 ml of the PH medium (see Table 8 for its composition) containing 0.05% yeast extract and 500 ppm

THIS SHEET HAS BEEN
REPLACED

This Sheet Has Been Replaced



of phenol in a 20 ml vial, the present microorganism inoculated by picking a platinum loopful of the colony of the present microorganism from the agar plate prepared by adding 1.5% agar to the NYG medium containing 500 ppm phenol, or by inoculating 0.04 ml (the cell count was 10^6 cells/ml) of the preculture liquid obtained by culturing the present microorganism in the NYG medium containing 500 m of phenol at 30°C overnight under shaking.

TABLE 7

10	PH medium	
	Magnesium sulfate heptahydrate	0.2 g
	Calcium sulfate	0.1 g
	Ferric chloride hexahydrate	0.02 g
	Dipotassium hydrogen phosphate	1.0 g
15	Ammonium sulfate	1.0 g
	Sodium chloride	0.1 g
	Distilled water	1 liter

Trichloroethylene in such an amount that it became 30 ppm after all ingredients were dissolved in the liquid phase simultaneously with the inoculation and 500 ppm of phenol were added. The vial was plugged with a Teflon-coated butyl rubber septum and sealed with an aluminum cap. It was cultured under shaking at 30°C, and trichloroethylene concentration was regularly measured by analyzing the gas phase in the vial with a gaschromatograph equipped with an ECD detector. The phenol concentration was determined by filtering the culture liquid with a 0.45 μ filter, and adding to 1 ml of the resulting filtrate, sequentially, 0.1 ml of an aqueous solution of $K_3Fe(CN)_6$ /0.1 M glycine and 1 ml of an aqueous solution of 4-amino antipyrine, which was then mixed, and measuring an absorbance at 505 nm.

The result as shown in Fig. 8 indicates that with the propagation of the bacterial mass phenol and trichloroethylene decreases, and they were completely decomposed at 31 hours after cultivation. It was

confirmed, therefore, that the present microorganism has a high activity of decomposing trichloroethylene when propagated in the presence of an aromatic compound such as phenol having the ability of inducing

5 trichloroethylene. When compared to known microorganisms, all reports relate to decomposition of trichloroethylene at a low concentration of up to 10 ppm except for the report that Pseudomonas cepacia KK01 decomposes 30 ppm of trichloroethylene to 15 ppm in 2

10 days and the report that Alcaligenes eutropus KS01 (Japanese Unexamined Patent Publication No. 7(1995)-123976) decomposes completely 50 and 25 ppm of trichloroethylene in 4 days.

In the decomposition of trichloroethylene by

15 Alcaligenes eutropus KS01 (Japanese Unexamined Patent Publication No. 7(1995)-123976), the amount of the microorganism inoculated to the culture medium is 10^8 cells/ml, which is about 100 times that of the present microorganism described in Example 6. The amount of the

20 bacterial mass required for trichloroethylene decomposition for the microorganism of the present invention is much smaller than that of Alcaligenes eutropus KS01. It therefore has an advantage that the cost of culturing is reduced, etc. It was also confirmed

25 that the added and mixed phenol is decomposed to below the level of detection, thereby presenting little risk of environmental pollution by phenol which is an environmental pollutant.

The inventors have focused and investigated on the

30 culturing method prior to inoculation in order to enhance the decomposition activity of a microorganism which has the ability of decomposing trichloroethylene in the soil and thereby to prepare a microorganism having the decomposition activity which can deal with pollution at

35 high concentrations. As a result we have discovered that there is an optimum value of culturing time of the

microorganism prior to inoculation, and that the activity of trichloroethylene decomposition can be enhanced by sequentially adding an inducer to the culture medium, and we have completed the present invention. The present invention will now be more fully explained with reference to the following examples.

Example 7.

Into 100 ml of the PH medium containing 0.05% yeast extract and 500 ppm of phenol contained in a 500 ml Erlenmeyer flask was inoculated a platinum loopful of the colony of the microorganism of the present invention which had been stored by passage on the agar plate of 1.5% agar added to the NYG medium containing 500 ppm phenol, or 1.0 ml of the preculture liquid obtained by culturing the microorganism of the present invention in the NYG medium containing 500 ppm of phenol at 30°C overnight under shaking. While culturing under shaking at 30°C at 130 r.p.m., turbidity of the culture liquid and phenol concentration were determined in the same methods as in Example 6. The bacterial mass in the culture was harvested by centrifugation at 5000 r.p.m. for 10 minutes and then resuspended into the NMS medium having an amount equal to the culture medium. The OD₆₀₀ of the suspension was 0.2 (the cell count was 1×10^8 c.f.u./ml).

The suspension of the bacterial mass was dispensed in a 20 ml vial and trichloroethylene was added thereto in such an amount that it became 30 ppm when all the ingredients were dissolved in the liquid phase. The vial was plugged with a Teflon-coated butyl rubber septum and sealed with an aluminum cap. After culturing at 30°C under shaking for 24 hours, the gas phase in the vial was analyzed by a gaschromatograph equipped with an ECD detector. The results are shown in Fig. 9 to Fig. 11. The line graphs in Fig. 9 represent turbidity of the culture liquid and phenol concentration. The bar graphs

in Fig. 10 represent the decomposition activity per unit amount of the bacterial mass (specific activity) of the microorganism of the present invention harvested at respective time of culturing. The bar graphs in Fig. 11
5 represent the decomposition activity per unit amount of the culture liquid (the total activity) of the microorganism of the present invention which was harvested at respective time of culturing.

From the results obtained as above on the activity
10 of trichloroethylene decomposition, the specific activity notably increases from the induction phase to the logarithmic growth phase, leveling off thereafter at a certain level. But it gradually decreased after the residual phenol became zero. However, the total activity
15 remained at the highest value at the stationary phase when phenol concentration became around zero. These observations have shown that a bacterial mass having a high activity of trichloroethylene decomposition can be obtained by culturing while monitoring the turbidity of culture liquid and phenol concentration prior to
20 inoculation into the soil, and terminating the culture at the timing that the increase in turbidity ceases to enter the stationary phase and phenol concentration becomes almost zero.

25 Example 8.

The microorganism of the present invention was cultured in a manner similar to Example 7 and turbidity of the culture liquid and phenol concentration were measured in a manner similar to Example 6. Furthermore,
30 when the phenol concentration reached zero in the middle of culturing 500 ppm of phenol was added again and culture was continued. The bacterial mass in the culture was harvested by centrifuging at 5000 r.p.m. for 10 minutes and then resuspended into the NMS medium having
35 an amount equal to that of the culture medium.

The OD₆₀₀ of the suspension was 0.2 (the cell count

was 2.5×10^8 c.f.u./ml). The suspension of the bacterial mass was dispensed in a 20 ml vial and trichloroethylene was added thereto in such an amount that it became 30 ppm when all the ingredients were dissolved in the liquid phase. The vial was plugged with a Teflon-coated butyl rubber stopper and sealed with an aluminum cap. After culturing at 30°C under shaking for 24 hours, the gas phase in the vial was analyzed by a gaschromatograph equipped with an ECD detector.

The results are shown in Fig. 12 to Fig. 14. The line graphs in Fig. 12 represent turbidity of the culture liquid and phenol concentration at the stage of culturing prior to harvesting. The bar graphs in Fig. 13 represent the specific activity of the microorganism of the present invention which was harvested at respective time of culturing. The bar graphs in Fig. 14 represent the total activity of the microorganism of the present invention which was harvested at respective time of culturing. When 500 ppm of phenol was added to the culture liquid at 45 hours of culturing when the residual phenol became almost zero, the phenol concentration decreased again and the turbidity of the culture liquid increased. Therefore, phenol was added again at 69 hours, but neither increase in turbidity of the culture liquid nor decrease in phenol concentration were observed.

With regard to the activity of trichloroethylene decomposition, high specific activities were obtained when sufficient phenol is remaining at around the stationary phase during 41 hours to 60 hours as shown in Fig. 13, but the specific activity decreased thereafter when phenol concentration decreased. When phenol was added at 93 hours, both the amount of the bacterial mass and the specific activity decreased. However, during the period from 93 hours to 114 hours when there was residual phenol, the specific activity was higher than during 45 hours to 69 hours when most of the phenol was decomposed.

On the other hand, the total activity was the highest at 60 hours when both of the amount of the bacterial mass and specific activity were high, providing the total activity 2.5 times higher than when phenol is added only at the start of culturing.

Next, trichloroethylene decomposition in the soil will be explained with reference to the following examples.

Example 9.

To a 100 ml vial was added 30 g of sandy soil (air-dried) which was artificially contaminated with trichloroethylene. The microorganism of the present invention was cultured under shaking at 30°C for 3 days in 1.5 ml, 4.5 ml, 7.5 ml, 15 ml, or 30 ml of the NMS medium which had 500 ppm of phenol and 0.05% yeast extract added thereto, and then was centrifuged to harvest the cells, which were resuspended into an appropriate amount of the NMS medium so that the water content in the soil became 25% after addition of the suspension of the bacterial mass. After adding the suspension (containing no phenol) of the bacterial mass to the soil, the vial was sealed with a Teflon-coated rubber septum and sealed with an aluminium cap, and then allowed to stand at 30°C for a certain period of time.

Thereafter, to a vial containing 30 g of the soil were added 50 ml of deionized water aerated with the air which had been passed through activated charcoal and 10 ml of n-hexane, and then the vial was sealed. The vial was sonicated in the sonicating washer for 10 minutes and then was shaken in a shaker for 10 minutes. The separated n-hexane was analyzed using a gaschromatograph equipped with an ECD detector. The result as shown in Fig. 15 indicated that trichloroethylene decomposition and the density of the bacterial mass correlated until the density of the bacterial mass reached 2.5×10^8 cfu/g wet soil (15 ml of the culture liquid), but it became

saturated at about 5×10^8 cfu/g wet soil (30 ml of the culture liquid) or higher. Therefore, it was revealed that when the microorganism of the present invention is used to decompose trichloroethylene the density of the bacterial mass of about 2.5×10^8 cfu/g wet soil is most suitable and that the microorganism of the present invention has a characteristics which enables an estimation of a minimum amount of bacterial mass required for purification of a given concentration of a contaminant in the soil.

Example 10.

To a 100 ml vial was added 30 g of sandy soil (air-dried) which was artificially contaminated with trichloroethylene. The microorganism of the present invention was cultured under shaking at 30°C for 3 days in 7.5 ml of the NMS medium which had 500 ppm of phenol and 0.05% yeast extract added thereto, and then was centrifuged to harvest the cells, which were then resuspended into an appropriate amount of the NMS medium so that the water content in the soil became 25% after addition of the suspension of the bacterial mass. After adding the suspension (containing no phenol) of the bacterial mass to the soil, the vial was sealed with a Teflon-coated rubber septum and sealed with an aluminium cap, and then allowed to stand at 30°C for a certain period of time.

Thereafter, to a vial containing 30 g of the soil were added 50 ml of deionized water aerated with the air which had been passed through activated charcoal and 10 ml of n-hexane, and the vial was sealed. The vial was sonicated in the sonicating washer for 10 minutes and then was shaken in a shaker for 10 minutes. The separated n-hexane was analyzed using a gaschromatograph equipped with an ECD detector. The result as shown in Fig. 16 indicates that when the bacterial mass corresponding to 7.5 ml of the culture liquid was added

twice to the soil, the efficiency of decomposition was equal to when the bacterial mass corresponding to 15 ml of the culture liquid was added at one addition.

5 It is difficult to supply at one addition the amount of culture liquid necessary to decompose trichloroethylene contained in the contaminated soil in terms of the actual amount needed (the actual amount of the contaminated soil is more than a few dozen m^3), and the decomposing microorganism must be sequentially
10 cultured and added to the soil. It was revealed that sequential addition of the microorganism of the present invention gives the same effect as that obtained when the required bacterial mass is added at one addition. Therefore, by adding a small amount of the culture liquid
15 repeatedly the microorganism of the present invention can deal with a vast area of the contaminated site for which one-time culture and infusion of the bacterial mass is insufficient.

Example 11.

20 To a 100 ml vial was added 30 g of sand grain soil (air-dried) which was artificially contaminated with trichloroethylene at the desired concentrations. The concentration of the contaminant in the soil was set at 15, 30, 45, 100, and 150 mg/kg. The microorganism of the
25 present invention was cultured under shaking at 30°C for 3 days in 15, 30, and 45 ml of the NMS medium which had 500 ppm of phenol and 0.05% yeast extract added thereto, and then was centrifuged to harvest the cells, which were resuspended into an appropriate amount of the NMS medium
30 so that the water content in the soil became 25% after addition of the suspension of the bacterial mass. After adding the suspension (containing no phenol) of the bacterial mass to the soil, the vial was sealed with a Teflon-coated rubber septum and sealed with an aluminium
35 cap, and then allowed to stand at 30°C for a certain period of time. The density of the bacterial mass in the

soil 9.4×10^8 , 1.9×10^9 , and 2.8×10^9 cfu/g wet soil for 15, 30, and 45 ml of the culture liquid, respectively.

Thereafter, to a vial containing 30 g of the soil were added 50 ml of deionized water aerated with the air which had been passed through activated charcoal and 10 ml of n-hexane, and then the vial was sealed. The vial was sonicated in the sonicating washer for 10 minutes and then was shaken in a shaker for 10 minutes. The separated n-hexane was analyzed using a gaschromatograph equipped with an ECD detector. The result as shown in Fig. 17 indicates that a very high concentration (about 150 mg/kg) of trichloroethylene in the soil could be decomposed. The above result indicated that purification of a high concentration (about 150 mg/kg) contaminant can be effected by increasing the amount of the bacterial mass added or by sequential addition because the amount of trichloroethylene decomposed increases in proportion to the amount of the bacterial mass added to the soil.

Example 12.

To a 100 ml vial which can be sealed with a Teflon-coated rubber septum was added 30 g of sandy soil (air-dried) which was artificially contaminated with trichloroethylene. The microorganism of the present invention was cultured under shaking at 30°C for 3 days in the NMS medium which had 500 ppm of phenol and 0.05% yeast extract added thereto, and then was centrifuged to harvest the cells, which were resuspended into an appropriate amount of the NMS medium so that the water content in the soil became 25% and the inoculated amount of bacterial mass became 10^8 to 10^9 cells/g wet soil after addition of the suspension of the bacterial mass. After adding the suspension (containing no phenol) of the bacterial mass to the soil, the vial was sealed with a Teflon-coated rubber stopper, and then allowed to stand at 30°C for a certain period of time.

Thereafter, to a vial containing 30 g of the soil were added 50 ml of deionized water aerated with the air which was passed through activated charcoal and 10 ml of n-hexane, and the vial was sealed. The vial was
5 sonicated in the sonicating washer for 10 minutes and then was shaken in a shaker for 10 minutes. The separated n-hexane was analyzed using a gaschromatograph equipped with an ECD detector. Time course of changes in trichloroethylene concentration in the soil was
10 determined by preparing a multiple of samples at the same time and extracting the entire volume of a part of samples after passage of a given time, followed by measurement. They were stored at 4°C until extraction.

The measurement of trichloroethylene concentration
15 in the soil was also conducted using a method indicated in the Environmental Standard (the Soil Environmental Standard) related to the contamination of the soil. Thus, the soil sample and the solvent (hydrochloric acid was added to purified water and pH was adjusted to 5.8 to
20 6.3) were added at a weight to volume ratio of 10% to an Erlenmeyer flask with a screw socket having a stirrer bar, and then the flask was immediately sealed. At this time, care was taken to make the volume of the mixture be not less than 500 ml and to minimize the head space in
25 the Erlenmeyer flask with a screw socket relative to the volume of the mixture. The prepared sample liquid was stirred continuously for 4 hours with the magnetic bar keeping the liquid at ordinary temperature and ordinary pressure.

30 After the sample liquid was allowed to stand for 30 minutes, it was aspirated into a glass syringe. A filter holder fitted with a membrane filter having a pore size of 0.45 μ was connected to the syringe and the plunger was pressed to exclude the air and the initial few ml of
35 the liquid, and then the filtrate was collected in fractions into stoppered test tubes, from which an amount necessary for determination was weighed out and was

subjected to analysis by gaschromatography. The result as shown in Fig. 18 indicated that 90% of the total trichloroethylene in the soil was decomposed in a day after addition of the bacterial mass.

5 It was revealed therefore that the method of purifying trichloroethylene by adding the microorganism of the present invention into the soil is a technology which has a high purification ability almost equal to decomposition capacity of about 18 mg/kg of
10 trichloroethylene in the soil in a day. When the soil sample at 7 days of standing was measured according to an analytical method designated by the nation (the Environmental Standard, the Soil Environmental Standard related to the contamination of the soil), the
15 concentration of trichloroethylene was below the base value (0.03 ppm). As a result, it was revealed that trichloroethylene can be purified to the level which satisfies the environmental standard by purifying the contaminated soil using the microorganism of the present
20 invention. It is generally accepted that the physical means of treatment such as vacuum extraction is unable to purify from low concentrations, but the use of the microorganism of the present invention enables purification to the level lower than the base value.

25 Example 13.

To a 100 ml vial which can be sealed with a Teflon-coated rubber septum was added 30 g of sandy soil (air-dried) which was artificially contaminated with trichloroethylene. The microorganism of the present
30 invention was cultured under shaking at 30°C for 3 days in the NMS medium which had 500 ppm of phenol and 0.05% yeast extract added thereto, and then was centrifuged to harvest the cells, which were resuspended into an appropriate amount of the NMS medium so that the water
35 content in the soil became 25% and the inoculated amount of bacterial mass became 10^8 to 10^9 cells/g wet soil

after addition of the suspension of the bacterial mass. After adding the suspension (containing no phenol) of the bacterial mass to the soil, the vial was sealed with a Teflon-coated rubber stopper, and then allowed to stand at 30°C for a certain period of time. Thereafter, analysis was conducted on byproducts which are said to be formed in association with trichloroethylene decomposition in the soil.

The byproducts in the soil were analyzed as follows: vinyl chloride, 1,1-dichloroethylene, cis-1,2-dichloroethylene, and trans-1,2-dichloroethylene were analyzed in the method shown in Fig. 19. Dichloro acetate and trichloro acetate, trichloroethanol, and hydrated chloral were analyzed in the method shown in Fig. 20, 21, and 22, respectively. The result as shown in Table 9 indicated that all byproducts were below the detection limit.

The dechlorination reaction of trichloroethylene under an anaerobic condition leads to accumulation of dichloroethylenes which is more toxic. It is also known that the decomposition process under an aerobic condition starting with formation of trichloroethylene oxide produces dichloro acetate etc. as intermediate products. There was no accumulation of harmful substances on decomposition of trichloroethylene by the microorganism of the present invention in the soil. Although the presence of substances other than those measured is unknown, this technology appeared to be a highly safe method.

THIS SHEET HAS BEEN
REPLACED

This Sheet Has Been Replaced



TABLE 8

	Item to be analyzed	Result	Detection limit
	Vinyl chloride	Not detected	0.01 ppm
	1,1-dichloroethylene	Not detected	0.01 ppm
5	cis-1,2-dichloroethylene	Not detected	0.01 ppm
	trans-1,2-dichloroethylene	Not detected	0.01 ppm
	Dichloroacetate	Not detected	0.05 ppm
	Trichloroethanol	Not detected	0.01 ppm
	Trichloroacetate	Not detected	0.05 ppm
10	Hydrated chloral	Not detected	0.05 ppm

Example 14.

To a 100 ml vial which can be sealed with a Teflon-coated rubber septum was added 30 g of sandy soil (air-dried) which was artificially contaminated with trichloroethylene. The microorganism of the present invention was cultured under shaking at 30°C for 3 days in the NMS medium which had 500 ppm of phenol and 0.05% yeast extract added thereto, and then was centrifuged to harvest the cells, which were resuspended into an appropriate amount of the NMS medium so that the water content in the soil became 25% and the inoculated amount of bacterial mass became 10^8 to 10^9 cells/g wet soil after addition of the suspension of the bacterial mass. After plating the suspension of the bacterial mass onto a petri dish to a thickness of about 1 mm, it was sterilized under irradiation of a 15 W Ultra violet lamp with a wavelength of 260 nm for not less than 60 seconds at a distance of 40 cm from the light source. After adding the suspension (containing no phenol) of the microorganism of the present invention into the soil, the vial was sealed with a Teflon-coated rubber stopper, and then allowed to stand at 30°C for a certain period of time.

Thereafter, to a vial containing 30 g of the soil were added 50 ml of deionized water aerated with the air which had been passed through activated charcoal and 10

ml of n-hexane, and then the vial was sealed. The vial was sonicated in the sonicating washer for 10 minutes and then was shaken in a shaker for 10 minutes. The separated n-hexane was analyzed using a gaschromatograph equipped with an ECD detector. Furthermore, in an experiment in which the suspension of sterilized bacterial mass at an amount of 1/100 of that of the NYG medium was inoculated or the suspension was plated as it was onto the NYG agar medium and cultured at 30°C, neither increase in turbidity of the culture liquid nor colony formation were observed, and thereby it was confirmed that there was complete sterilization. The result as shown in Fig. 23 indicated that the dead cells and living cells gave an equal degree of decomposition of trichloroethylene.

Although most of the purification methods using microorganisms comprise adding living organisms into the soil, addition of the bacterial mass into the soil is currently difficult from a viewpoint of public acceptance. It is also being feared that it has a potential risk of producing a far-reaching effect on the ecological system by releasing a specific microorganism into the environment. But the addition of the microorganism which has completely lost the propagating activity by sterilization treatment is equivalent to that of mere organic materials, and thus is believed to have little effect on the ecological system. Therefore an experiment was conducted in which addition of the decomposing microorganism sterilized with ultraviolet irradiation into the soil was investigated. The result indicated that the addition of the dead bacterial mass proved to be an extremely useful method.

The invention disclosed in Japanese Unexamined Patent Publication No. 8(1996)-3012 claims that the effect on the ecological system can be minimized by crushing the decomposing bacteria and then spraying it to the soil. But, it is readily anticipated that the

spraying of a large mass of decomposing bacterium to the contaminated soil is in fact difficult because crushing of microorganisms takes extensive equipment, a lot of time and labor. The known enzymes which have

5 trichloroethylene-oxidizing activity require NAD as a coenzyme. But because the coenzyme is very expensive, it would be extremely difficult to supply the coenzyme in the concentration necessary for the decomposition reaction of the enzyme which was released by crushing the

10 bacterium from the bacterial mass. On the other hand, the method using the microorganism of the present invention has no such problems, and enables purification of trichloroethylene etc. at low cost and with minimum effects on the ecological system.

15 Example 15.

To a 100 ml vial which can be sealed with a Teflon-coated rubber septum was added 30 g of sand grain soil (air-dried) which was artificially contaminated with trichloroethylene. The microorganism of the present

20 invention was cultured under shaking at 30°C for 3 days in the NMS medium which had 500 ppm of phenol and 0.05% yeast extract added thereto, and then was centrifuged to harvest the cells, which were resuspended into an appropriate amount of the NMS medium so that the water

25 content in the soil became 25% and the inoculated amount of bacterial mass became 10^8 to 10^9 cells/g wet soil after addition of the suspension of the bacterial mass. After plating the suspension of the bacterial mass onto a petri dish to a thickness of about 1 mm, and then

30 sterilized under irradiation of a 15 W ultra violet lamp with a wavelength of 260 nm for not less than 60 seconds at a distance of 40 cm from the light source.

After adding the suspension (containing no phenol) of the sterilized microorganism of the present invention

35 into the soil, the vial was sealed with a Teflon-coated rubber stopper, and then allowed to stand at 30°C for a

certain period of time. After that, trichloroethylene in the soil was allowed to be decomposed and the byproducts which is believed to be formed in association with trichloroethylene decomposition were analyzed in a similar manner to Example 13. Furthermore, in an experiment in which the suspension of sterilized bacterial mass at an amount of 1/100 of that of the NYG medium was inoculated or the suspension was plated as it was onto the NYG agar medium and cultured at 30°C, neither increase in turbidity of the culture liquid nor colony formation were observed, and thereby it was confirmed that there was complete sterilization.

The result as shown in Table 10 indicated that all byproducts were below the detection limit. The dechlorination reaction of trichloroethylene under an anaerobic condition leads to accumulation of dichloroethylenes which is more toxic. It is also known that the decomposition process under an aerobic condition which starts with formation of trichloroethylene oxide produces dichloroacetate etc. as intermediate products. There was no accumulation of harmful substances on decomposition of trichloroethylene by the microorganism of the present invention in the soil. Although the presence of substances other than those measured is unknown, it was suggested that this technology is a highly safe method.

TABLE 9

	Item to be analyzed	Result	Detection limit
	Vinyl chloride	Not detected	0.01 ppm
30	1,1-dichloroethylene	Not detected	0.01 ppm
	cis-1,2-dichloroethylene	Not detected	0.01 ppm
	trans-1,2-dichloroethylene	Not detected	0.01 ppm
	Dichloroacetate	Not detected	0.05 ppm
	Trichloroethanol	Not detected	0.01 ppm
35	Trichloroacetate	Not detected	0.05 ppm
	Hydrated chloral	Not detected	0.05 ppm

THIS SHEET HAS BEEN
REPLACED

This Sheet Has Been Replaced



Example 16.

To a 100 ml vial which can be sealed with a Teflon-coated rubber septum was added 30 g of sandy soil (air-dried) which was artificially contaminated with trichloroethylene. The microorganism of the present invention was cultured under shaking at 30°C for 3 days in the NMS medium which had 500 ppm of phenol and 0.05% yeast extract added thereto, and then was centrifuged to harvest the cells, which were resuspended into an appropriate amount of the NMS medium so that the water content in the soil became 25% and the inoculated amount of bacterial mass was 10^8 to 10^9 cells/g wet soil after addition of the suspension of the bacterial mass. After adding the suspension (containing no phenol) of the microorganism of the present invention into the soil, the vial was sealed with a Teflon-coated rubber stopper, and then allowed to stand at 20°C for a certain period of time.

Thereafter, to a vial containing 30 g of the soil were added 50 ml of deionized water aerated with the air which had been passed through activated charcoal and 10 ml of n-hexane, and the vial was sealed. The vial was sonicated in the sonicating washer for 10 minutes and then was shaken in a shaker for 10 minutes. The separated n-hexane was analyzed using a gaschromatograph equipped with an ECD detector.

The result as shown in Fig. 24 indicated that trichloroethylene was decomposed at 20°C to a degree equivalent to that at 30°C. Since it was suggested that trichloroethylene is sufficiently decomposed at the temperature of the soil, the microorganism of the present invention proved to have a high practical utility as compared to the others.

Example 17.

To a 100 ml vial was added 30 g of sand grain soil (air-dried) which was artificially contaminated with

trichloroethylene. The microorganism of the present invention was cultured under shaking at 30°C for 1 day in the NMS medium which had 500 ppm of phenol and 0.05% yeast extract added thereto. The culture liquid at a
5 volume of 1/100 of that of the NMS medium (or the PH medium) containing 500 ppm, 0.02%, and 1 mM of phenol, yeast extract, and glucose, respectively in the soil was added and then added to the soil. The density of the bacterial mass was set at 10^6 cells/g wet soil, and the
10 water content set at 25% at the time of culture liquid addition.

The vial was sealed with a Teflon-coated rubber septum, and then allowed to stand at 30°C for a certain period of time. Thereafter, to a vial containing 30 g of
15 the soil were added 50 ml of deionized water aerated with the air which had been passed through activated charcoal and 10 ml of n-hexane, and then the vial was sealed. The vial was sonicated in the sonicating washer for 10 minutes and then was shaken in a shaker for 10 minutes.
20 The separated n-hexane was analyzed using a gaschromatograph equipped with an ECD detector.

The result as shown in Fig. 25 indicated that trichloroethylene in the soil similarly decreased from about 14 mg/kg to about 1 mg/kg in the NMS medium and the
25 PH medium. Due to the addition of phenol into the soil, autochthonous microorganisms contributed to reduction of trichloroethylene, but the result indicated a greater effect came from the addition of strain MO7. Therefore, it is clear that the possibility can be ruled out that
30 the strain exerts its effect not only in the sterilized culture medium but also in the natural environment although autochthonous microorganisms are present. It was confirmed that the activity of strain MO7 is sufficiently induced by adding phenol at an amount of
35 about 10^6 cells/g wet soil even in the natural environment.

Example 18. Construction of a constitutive mutant

A platinum loopful of the colony of the microorganism of the present invention which had been stored by passage on the agar plate containing 1.5% agar was picked and inoculated to a test tube containing 2 ml of the 1/3LB medium (its composition is shown in Table 11). After cultivating overnight under shaking at 30°C at 130 r.p.m., an aliquot of the culture broth was diluted as appropriate, which was then plated to the plate of the 1/3LB medium containing 1.5% agar. After cultivating at 30°C the number of cells was counted. The remainder of the culture broth was plated on a petri dish, which was then subjected to irradiation of a 15 W ultra violet lamp with a wavelength of 260 nm under the condition of a irradiation time of 3 minutes at a distance of 30 cm from the light source.

TABLE 10

The 1/3LB Medium		
	Tryptone	3.0 g
20	Yeast extract	1.5 g
	Sodium chloride	3.0 g
	Distilled water	1.0 liter

Then, the culture broth was diluted as appropriate and was plated to the plate of the 1/3LB medium containing 1.5% agar. After culturing at 30°C the number of cells was counted to determine the death rate. When the death rate of 99% or higher was observed, it was assumed that there was enough mutation. From the culture broth at this point, the desired mutant were selected.

Catechol 2,3-dioxygenase (C230) introduces oxygen into catechol to form 2-hydroxymuconic acid semialdehyde by metafission. This product is yellow-colored. When C230 is expressed after being sprayed, it readily turns yellow. The trichloroethylene degrading enzymes of the phenol-utilizing trichloroethylene-decomposing bacteria are known to be the phenol hydroxylase (PH) which convert

THIS SHEET HAS BEEN
REPLACED

This Sheet Has Been Replaced



phenol into catechol (M. Fujita et al., J. Ferment. Bioeng., 79: 100, 1995; V. Shingler et al., J. Bacteriol., 174: 711, 1992).


5 An example has been known that C230 is expressed by one operon in which the C230 gene is adjacent to the PH gene though C230 itself was not directly involved in the decomposition of trichloroethylene (M. Fujita et al., J. Ferment. Bioeng., 79: 100, 1995; V. Shingler et al., J. Bacteriol., 174: 711, 1992). If C230 is expressed in
10 association with the PH, it is possible to select the strains which express the trichloroethylene-degrading enzyme using C230 expression as an indicator. The microorganism of the present invention, when cultivated using phenol as the only carbon source, decomposes phenol
15 and the culture broth turns yellow with the growth of the bacterium. This will make said selection method applicable to the microorganism.

Accordingly, the suspension of the bacterial mass after mutation was diluted as appropriate and was plated
20 to the plate of the 1/3LB medium containing 1.5% agar. After cultivating at 30°C for 1 to 3 days, a catechol solution (0.1% (w/v) ethereal solution) was sprayed.

Then, when the death rate was 99.99%, about 10,000 colonies which appeared after ultra violet irradiation
25 were investigated, which led to selection of 16 yellow colonies.

After the colonies were cultured in the 1/3LB liquid medium, it was plated to the plate of the 1/3LB medium containing 1.5% agar. After cultivating at 30°C, the
30 colonies which appeared were sprayed with a catechol solution, and yellow colonies were isolated again to obtain the strains which constitutively express C230.

A platinum loopful of these strains which constitutively express C230 were picked and inoculated
35 into the M medium (its composition is shown in Table 12) containing 4 ml of sodium glutamate (0.1%) in a 20 ml vial. The vial was plugged with a butyl rubber septum



This Sheet Has Been Replaced



and sealed with an aluminum cap, which was then cultivated at 30°C under shaking for 2 days. Then trichloroethylene was added in an amount so that it became 30 ppm when all were dissolved in the liquid phase. After culturing at 30°C for 5 days, trichloroethylene concentration was determined as in Example 6.

TABLE 11

M medium

10	Ammonium nitrate	3.0 g
	Disodium hydrogen phosphate	2.2 g
	Potassium dihydrogen phosphate	0.8 g
	Ferrous sulfate (II) heptahydrate	10.0 mg
	Calcium sulfate dihydrate	10.0 mg
15	Magnesium sulfate heptahydrate	10.0 mg
	Yeast extracts	50.0 mg
	Distilled water	1.0 liter, pH 7.0

The result demonstrated that the parent strain MO7 decomposed little trichloroethylene, but some of the strains which constitutively expressed C230 decomposed 52 to 34% of 30 ppm trichloroethylene. These microorganisms turned out to be a mutant which is derived from the parent strain MO7 and which can decompose trichloroethylene without requiring phenol.

Then, trichloroethylene decomposition in the soil is illustrated with reference to an example in which the strain MO715 which had the highest activity of decomposing trichloroethylene was used.

30 Example 19. Decomposition of trichloroethylene by a constitutive mutant

A soil which was artificially contaminated with trichloroethylene (11 mg/kg soil) was prepared as in Example 9. After cultivating the strain MO715 under shaking at 30°C for 2 days in 100 ml of the M medium containing sodium glutamate (0.1 w/v %), and then was centrifuged to harvest the cells, which were resuspended

into an appropriate amount of the M medium so that the water content in the soil became 25%. After the suspension of the bacterial mass (containing no phenol) was added to the soil, the vial was sealed with a Teflon-coated rubber septum and was allowed to stand at 30°C for a certain period of time. Trichloroethylene was analyzed in a method similar to that in Example 9. The culture broth cultivated on phenol and the strain MO7 were also analyzed in a similar manner.

As a result, after the addition of the suspension of the bacterial mass of the strain MO715 cultivated with sodium glutamate, about 35% of trichloroethylene was decomposed in 12 hours and about 60% in 48 hours. The strain MO7 grown on sodium glutamate decomposed little trichloroethylene. Thus, it was revealed that the microorganism of the present invention does not require phenol for induction and can decompose trichloroethylene in the soil as well.

From the foregoing it can be concluded that the present invention is an extremely safe technology since trichloroethylene can be decomposed without spreading toxic substances such as phenol into the environment.

Reference to the international deposition of microorganisms under the Budapest Treaty

International Depository Authority: National Institute of Bioscience and Human-Technology Agency of Industrial Science and Technology

Address: 1-3, Higashi 1-chome, Tsukuba-shi, Ibaraki, 305, Japan

Identification	Deposition Number	Deposition Date
MO7	FERM BP-5624	Aug. 12, 1996
MO715	FERM BP-5928	Apl. 24, 1997

SEQUENCE LISTING

SEQ ID No: 1

Sequence length: 19

Sequence type: nucleic acid

Strandedness: single

Topology: linear

Molecule type: synthetic DNA

Sequence description:

GAGTTTGATC CTGGCTCAG

SEQ ID No: 2

Sequence length: 27

Sequence type: nucleic acid

Strandedness: single

Topology: linear

Molecule type: synthetic DNA

Sequence description:

AGAAAGGAGG TGATCCAGCC GCAGGTT

SEQ ID No: 3

Sequence length: 16

Sequence type: nucleic acid

Strandedness: single

Topology: linear

Molecule type: synthetic DNA

Sequence description:

GGCCGGACGG GTGAGT

SEQ ID No: 4

Sequence length: 15

Sequence type: nucleic acid

Strandedness: single

Topology: linear

Molecule type: synthetic DNA

Sequence description:

TACGGGAGGC AGCAG

SEQ ID No: 5
Sequence length: 17
Sequence type: nucleic acid
Strandedness: single
Topology: linear
Molecule type: synthetic DNA
Sequence description:
GTGCCAGCAG CCGCGCG

SEQ ID No: 6
Sequence length: 18
Sequence type: nucleic acid
Strandedness: single
Topology: linear
Molecule type: synthetic DNA
Sequence description:
GATTAGATAC CCTGGTAG

SEQ ID No: 7
Sequence length: 18
Sequence type: nucleic acid
Strandedness: single
Topology: linear
Molecule type: synthetic DNA
Sequence description:
ACTCAAAGGA ATTGACGG

SEQ ID No: 8
Sequence length: 16
Sequence type: nucleic acid
Strandedness: single
Topology: linear
Molecule type: synthetic DNA
Sequence description:
GCAACGAGCG CAACCC

SEQ ID No: 9

Sequence length: 16
Sequence type: nucleic acid
Strandedness: single
Topology: linear
Molecule type: synthetic DNA
Sequence description:
TGTACACACC GCCCGT

SEQ ID No: 10
Sequence length: 1422
Sequence type: nucleic acid
Strandedness: double
Topology: linear
Molecule type: cDNA
Sequence description:

AACGCTGGCG	GCGTGCTTAA	CACATGCAAG	TCGAACGGTG	AAGCTTGGAG	CTTGCTTCGA	60
GTGGATCAGT	GGCGAACGGG	TGAGTAACAC	GTGAGCAACC	TGCCCCAGAC	TCTGGAATAA	120
GCGCTGGAAA	CGGGCTCTAA	TACTGGATAT	GTGACGGACC	TGCATGGGTA	CCGTCTGGAA	180
AGTTTTTTCG	TTTGGGATGG	GCTCGCGGCC	TATCAGCTTG	TTGGTGAGGT	AATGGCTCAC	240
CAAGGCGACA	ACGGGTANCC	GGCCTGAGAG	GGCGACCGGC	CACACTGGGA	CTGAAACACG	300
GCCCCAACTC	CTACGGGAGG	CACCAGTGGG	GAAATATTGC	ACAATGGGCG	AAAGCCTGAT	360
GCAGCGACGC	CGCGTGAGGG	ATGACGGCCT	TCGGGTGTA	AACCTCTTTC	AGCAGGGAAG	420
AAGCGAAAGT	GACGGTACCT	GCAGAATAAG	CACCGGCTAA	CTACGTGCCA	GCAGCCGCGG	480
TAATACGTAG	GGTCCGAGCG	TTGTCCGGAA	TTATTGGGCG	TAAAGAGCTT	GTAGGCGGTT	540
TGTCGCGTCT	GCTGTGAAAA	TCCGGGGCTC	AACCCCGGAC	TTGCAGTGGG	TACGGGCAGA	600
CTAGAGTGTG	GTAGGGGAGA	CTGGAATTCC	TGGTGTAGCG	GTGAAATGCG	CAGATATCAG	660
GAGGAACACC	GATGGCGAAG	GCAGGTCTCT	GGGCCACTAC	TGACGCTGAG	AAGCGAAAGC	720
ATGGGGAGCG	AACAGGATTA	GATACCCTGG	TAGTCCATGC	CGTAAACGTT	GGGCGCTAGG	780
TGTGGGACTC	ATTCCACGAG	TTCCGTGCCG	CAGCTAACGC	ATTAAGCGCC	CCGCCTGGGG	840
CAGTACGGCC	GCAAGGCTAA	AACTCAAAGG	AATTGACGGG	GGCCCGCACA	AGCGGCGGAG	900
CATGCGGATT	AATTCGATGC	AACGCGAAGA	ACCTTACCAA	GGCTTGACAT	ATACCGGAAA	960
CTTCCAGAGA	TGGTTGCCCC	CTTTGGGTCT	GTATACAGGT	GGTGCATGGT	TGTCGTCAGC	1020
TCGTGTCGTG	AGATGTTGGG	TTAAGTCCCC	CAACGAGCGC	AACCCTCGTT	CTATGTTGCC	1080
AGCACGTCAT	GGTGGGGACT	CATGGAAGAC	TGCCGGGGTC	AACTCGGAAG	AAGGTGGGGA	1140
TGACGTCAAA	TCATCATGCC	CCTTATGTCT	TGGGCTTCAC	GCATGCTACA	ATGGCCGGTA	1200
CAAAGGGCTG	CGATACCGCA	AGGTGGAGCG	AATCCCCAAA	AAACCGGTCT	CAGTTCGGAT	1260
TGGGGTCTGC	AACTCGACCC	CATGAAGTCG	GAGTCGCTAG	TAATCGCAAA	TCAGCAACGC	1320

TGCGGTGAAT ACTTTCCCGG GCCTTGACAC CACCGCCCGT CAAGTCACGA AAGTTCGGTA 1380
ACACCCGAAG CCGGTGGCCC AACCCCTTGTG GGGGGAGCCG TC 1422

CLAIMS

1. A bacterium which has the following properties:

	Morphology	coccoid, rod shaped
	Gram staining	+
5	Spore forming	-
	Motility	-
	Relationship to oxygen	aerobic
	Oxidase test	-
	Catalase test	+
10	Resistance to acid	-
	Rod-coccus cycle	+
	Aerial mycelium formation	-
	Peptide glycan type of cell wall	meso-diaminopimelic acid
15	Glycolyl test	- (acetyl type)
	Mycolic acid	-
	Major isoprenoid quinones	MK-9, MK-9 (H2) (traces)
	GC content of DNA (mole%)	73 (by HPLC)

THIS SHEET HAS BEEN
REPLACED

- 20 and, which can decompose trichloroethylene.

2. A bacterium according to claim 1 which is strain MO7 (FERM BP-5624).

- 25 3. A method for decomposing organic halogenated compounds and/or aromatic compounds, comprising subjecting said organic halogenated compounds and/or aromatic compounds to the action of a bacterium according to claim 1 or 2.

- 30 4. A method according to claim 3 wherein the organic halogenated compound is trichloroethylene and the aromatic compound is a phenolic compound.

- 35 5. A method for decomposing organic halogenated compounds in the soil, waste waters, or other waste products containing organic halogenated compounds, comprising adding a culture of a bacterium according to claim 1 or 2 to the soil, waste waters, or other waste products.

6. A method according to claim 5 wherein the



organic halogenated compound is trichloroethylene and the culture is a cultured bacterial cells.

7. A method according to claim 5 or 6 wherein an aromatic compound is added to a culture medium for use in culturing to obtain the bacterial culture.

8. A method according to claim 7 wherein said aromatic compound is a phenolic compound.

9. A method according to any of claims 5 to 8 wherein an degradable carbon source is added to a culture medium for use in culturing to obtain the bacterial culture.

10. A method according to claim 9 wherein the degradable carbon source is glucose.

11. A method according to any of claims 5 to 10 wherein the culture is a cultured bacterial cells.

12. A method according to claim 11 wherein the cultured bacterial cells are living bacterial cells.

13. A method according to claim 11 wherein the cultured bacterial cells are sterilized cultured bacterial cells.

14. A method according to claim 13 wherein the sterilization treatment is ultraviolet irradiation.

15. A method for decomposing organic halogenated compounds in the soil, waste waters, or other waste products containing organic halogenated compounds, comprising the steps of inoculating a bacterium according to claim 1 or 2 to the soil, waste waters, or other waste products, adding an aromatic compound, an degradable carbon source, or a mixture thereof to the soil, waste waters, or other waste products, and then culturing said inoculated microorganism.

16. A method according to claim 15 wherein the organic halogenated compound is trichloroethylene.

17. A method according to claim 15 or 16 wherein the aromatic compound is a phenolic compound and the degradable carbon source is glucose.

18. A method for decomposing organic halogenated





compounds and/or aromatic compounds, wherein sterilized cultured bacterial cells of a microorganism capable of decomposing organic halogenated compounds and/or aromatic compounds is allowed to act on the organic halogenated compounds and/or the aromatic compounds.

19. A method for decomposing organic halogenated compounds and/or aromatic compounds in the soil, waste waters, or other waste products containing organic halogenated compounds and/or aromatic compounds, in which a sterilized cultured bacterial mass of a microorganism capable of decomposing said organic halogenated compounds and/or said aromatic compounds is added to said soil, waste waters, or other waste products.

20. A method according to claim 19 wherein the sterilization treatment is ultraviolet irradiation.

21. A method according to claim 19 or 20, wherein the aromatic compound is a phenolic compound and the organic halogenated compound is trichloroethylene.

22. A decomposition agent for organic halogenated compounds and/or aromatic compounds, comprising a bacterial culture according to claim 1 or 2.

23. A decomposition agent according to claim 22 wherein the culture is cultured bacterial cells.

24. A decomposition agent according to claim 23 wherein the culture is living bacterial cells.

25. A decomposition agent according to claim 23 wherein the cultured bacterial cells are sterilized bacterial cells.

26. A decomposition agent according to claim 25 wherein the sterilization treatment is ultraviolet irradiation.

27. A decomposition agent according to any of claims 22 to 26 which is in the form of dried bacterial cells.

28. A bacterium having the decomposition activity of organic halogenated compounds and/or aromatic compounds in the presence or absence of induction of an





organic halogenated compound-decomposing enzyme by an aromatic compound, which bacterium is obtainable by subjecting a bacterium according to claim 1 to a mutation treatment and a selection procedure.

5 29. A bacterium according to claim 28 which is the strain MO715 (FERM BP-5928).

 30. A method for decomposing organic halogenated compounds and/or aromatic compounds comprising allowing a bacterium according to claim 28 or 29 to act on the
10 organic halogenated compounds and/or aromatic compounds.

 31. A method according to claim 30 wherein the organic halogenated compound is trichloroethylene and the aromatic compound is a phenolic compound.

 32. A method for decomposing organic halogenated
15 compounds in the soil, waste waters, or other waste products containing organic halogenated compounds, comprising adding a culture of a bacterium according to claim 28 or 29 to the soil, waste waters, or other waste products.

20 33. A method according to claim 32 wherein the organic halogenated compound is trichloroethylene and the culture is cultured bacterial cells.

 34. A method according to claim 32 or 33 wherein no aromatic compound is added to the culture medium for use
25 in culturing to obtain said bacterial culture.

 35. A method according to any of claims 32 to 34 wherein an degradable carbon source is added to the culture medium for use in culturing to obtain said bacterial culture.

30 36. A method according to claim 35 wherein the degradable carbon source is glutamic acid.

 37. A method according to any of claims 32 to 36 wherein the culture is cultured bacterial cells.

35 38. A method according to claim 37 wherein the cultured bacterial cells are living bacterial cells.

 39. A method according to claim 37 wherein the cultured bacterial cells are sterilized bacterial cells.

THIS SHEET HAS BEEN
REPLACED




40. A method according to claim 39 wherein the sterilization treatment is ultraviolet irradiation.

41. A method for decomposing organic halogenated compounds in the soil, waste waters, or other waste products containing organic halogenated compounds, comprising the steps of inoculating a bacterium according to claim 28 or 29 to said soil, waste waters, or other waste products, adding an degradable carbon source to said soil, waste waters, or other waste products, and then culturing the inoculated microorganism.

42. A method according to claim 41 wherein the organic halogenated compound is trichloroethylene.

43. A method according to claim 41 or 42 wherein the degradable carbon source is glutamic acid.




This Sheet Has Been Replaced



* C L M S *


$\frac{1}{23}$

Fig. 1




2/
23

Fig. 2




$\frac{3}{23}$

Fig. 3




$\frac{4}{23}$

Fig.4




5/23

Fig.5




6/23

Fig.6




$\frac{7}{23}$

Fig.7



8/23

Fig. 8



9/23

Fig.9





Fig.10



$\frac{10}{23}$

Fig.11





Fig.12



$\frac{11}{23}$

Fig.13






Fig.14




12/23

Fig.15



13/23


Fig.16



1. The MO7 culture liquid (15ml) suspended in the NMS medium was added at one addition.
2. Not added.
3. The MO7 culture liquid (7.5ml) suspended in the NMS medium was added twice.
(Total volume added is 15ml)
4. Not added.
5. The MO7 culture liquid (7.5ml) suspended in the NMS medium was added.
6. The MO7 culture liquid (7.5ml) suspended in the NMS medium was added at one addition (the final water content was adjusted to be 25%).


$^{14}/_{23}$

Fig.17



15/23


Fig.18



16/23

Fig.19

1. Test procedure



2. Operating conditions for gas chromatograph-mass spectrometer

Instrument: Shimazu Corp. GC-MS QP-1100EX

Column: 25% Silicone DC-550/Chromosorb WHP 80 to 100 mesh glass tube, $\phi 3\text{mm} \times 2.6\text{m}$ Temperature: Injection port 250°C,
column 60°C (retain 1min) →
150°C (10°C/min temperature increase)

Gas flow rate: Helium 40ml/min

Temperature of ionization source: 250°C

Separator temperature: 250°C

Ionization potential: 70eV


Ionization method: EI

Preset Mass number: 62,64 (vinyl chloride)
96,98 (1,1-dichloroethylene,
cis-1,2-dichloroethylene and
trans-1,2-dichloroethylene)

17/23

Fig.20

1. Test procedure



2. Operating conditions for gas chromatography

Instrument: Shimazu GC-14A


Detector: ECD

Column: VOCOL, $\phi 0.53\text{mm} \times 30\text{m}$ Temperature: Injection port 200°C , detector 230°C column 50°C (retain 1min) $\rightarrow$ 130°C ($3^{\circ}\text{C}/\text{min}$ temperature increase)Gas pressure: Carrier gas (helium) $0.5\text{kg}/\text{cm}^2$ Makeup gas (nitrogen): $0.9\text{kg}/\text{cm}^2$

18/23

Fig.21

1. Test procedure



2. Operating conditions for gaschromatography

Instrument: Shimazu GC-14A


Detector: ECD

Column: VOCOL, $\phi 0.53\text{mm} \times 30\text{m}$ Temperature: Injection port 200°C , detector 230°C
column 40°C (retain 6min) $\rightarrow$
 85°C ($3^{\circ}\text{C}/\text{min}$ temperature increase)Gas pressure: Carrier gas (helium) $0.5\text{kg}/\text{cm}^2$ Makeup gas (nitrogen): $0.9\text{kg}/\text{cm}^2$

19/23

Fig.22

1. Test procedure



2. Operating conditions for gas chromatograph-mass spectrometer

Instrument: Shimazu Corp. GC-MS QP-1100EX


Column: DB-1, $\Phi 0.53\text{mm} \times 30\text{m}$ Temperature: Injection port 250°C , column 50°C Gas flow rate: Helium 20ml/min Temperature of ionization source: 250°C Separator temperature: 250°C Ionization potential: 70eV

Ionization method: EI

Mass number: 82


$\frac{20}{23}$

Fig.23




$\frac{21}{23}$

Fig.24




22/23

Fig. 25



$\frac{23}{23}$

Fig.26




provide new microorganisms which can decompose organic
 'halogenated compounds such as trichloroethylene in a more
 efficient manner than the conventionally known
 microorganisms, and a method for decomposing organic
 5 halogenated compounds utilizing said microorganism. It
 is another object of the present invention to provide an
 inexpensive method for minimizing the effects on the
 ecological system after a specific microorganism has been
 released to the environment.

10 After intensive studies to solve the above-mentioned
 problems the inventors have successfully isolated a new
 microorganism which does not belong to any of the known
 genera of microorganisms and which has a very high
 ability of decomposing trichloroethylene as compared with
 15 any of the conventionally known microorganisms, and
 thereby completed the present invention.

Thus, the present invention provides a bacterium
 which has the following taxonomical properties:

TABLE 1

20	Morphology	Coccoid, rod shaped
	Gram staining	+
	Spore forming	-
	Motility	-
	Relationship to oxygen	aerobic
25	Oxidase test	-
	Catalase test	+
	Resistance to acid	-
	Rod-coccus cycle	+
	Aerial mycelium forming	-
30	Peptide glycan type of cell wall	meso-diaminopimelic acid
	Glycolyl test	- (acetyl type)
	Mycolic acid	-
	Major isoprenoid quinones	MK-8 (H4)
35	GC content of DNA (mole%)	73 (by HPLC)




and, which can decompose trichloroethylene. A

the ability of decomposing trichloroethylene. The method of isolation of microorganisms is described in detail in Example 1.

Strain MO7 thus isolated has the following
5 taxonomical properties:

TABLE 2

	Morphology	Coccoid, rod shaped
	Gram staining	+
	Spore forming	-
10	Motility	-
	Behavior to oxygen	aerobic
	Oxidase test	-
	Catalase test	+
	Resistance to acid	-
15	Rod-coccus cycle	+
	Aerial mycelium	-
	Peptide glycan type of cell wall	meso-diaminopimelic acid
	Glycolyl test	- (acetyl type)
20	Mycolic acid	-
	Major isoprenoid quinones	MK-8 (H4)
	GC content of cellular DNA (mole%)	73 (by HPLC)




25 Furthermore, strain MO7 has the following properties.

	Color of colonies	No characteristic colony color formed
	Elongation of cells surrounding colony	-
30	Arabinogalactan polymers of cell wall	- (#1)

#1: Estimated using an acid hydrolysate of the whole bacterial mass

35 The morphological and physiological characteristics of strain MO7 were investigated and the results obtained are shown in the above Table 2. Based on the results described above identification of the strain was



bacterial mass after sterilization retains 80% or more of the trichloroethylene-decomposition activity of the living bacterium. Furthermore, in an experiment in which the suspension of sterilized bacterial mass at an amount of 1/100 that of the NYG medium was inoculated or the suspension was plated as it was onto the NYG agar medium and cultured at 30°C, neither increase in turbidity of the culture liquid nor colony formation were observed, and thereby it was confirmed that there was complete sterilization.

The residual activity of the suspension of the sterilized bacterial mass when stored at 4°C was almost the same as that of the living bacterial mass at three days of storage and said suspension had a much higher activity than that of the living bacterial mass at seven days of storage. The results reveal that although the initial activity of the dead bacterial mass is decreased to 80% of the living bacterial mass, a higher activity of decomposing trichloroethylene is retained than the living bacterial mass during storage because of a lower reduction in activity of the dead bacterial mass during storage.

By sterilizing a microorganism without destructing the boundary structure such as the cell wall of a bacterial mass with the extracellular environment, it is possible to retain coenzymes etc. which are required for decomposition reaction of trichloroethylene at concentrations necessary for expression of enzymatic activity. Thus, supply of coenzymes for maintenance of activity of trichloroethylene decomposition is not necessary and the spread of the cultured bacterial mass is only needed. It possible therefore to effect purification of trichloroethylene etc. at low cost and with a minimum effect on-the ecological system.


Example 6.

To 4 ml of the PH medium (see Table 7 for its composition) containing 0.05% yeast extract and 500 ppm

after addition of the suspension of the bacterial mass. After adding the suspension (containing no phenol) of the bacterial mass to the soil, the vial was sealed with a Teflon-coated rubber stopper, and then allowed to stand at 30°C for a certain period of time. Thereafter, analysis was conducted on byproducts which are said to be formed in association with trichloroethylene decomposition in the soil.

The byproducts in the soil were analyzed as follows: vinyl chloride, 1,1-dichloroethylene, cis-1,2-dichloroethylene, and trans-1,2-dichloroethylene were analyzed in the method shown in Fig. 19. Dichloro acetate and trichloro acetate, trichloroethanol, and hydrated chloral were analyzed in the method shown in Fig. 20, 21, and 22, respectively. The result as shown in Table 8 indicated that all byproducts were below the detection limit.

The dechlorination reaction of trichloroethylene under an anaerobic condition leads to accumulation of dichloroethylenes which is more toxic. It is also known that the decomposition process under an aerobic condition starting with formation of trichloroethylene oxide produces dichloro acetate etc. as intermediate products. There was no accumulation of harmful substances on decomposition of trichloroethylene by the microorganism of the present invention in the soil. Although the presence of substances other than those measured is unknown, this technology appeared to be a highly safe method.



certain period of time. After that, trichloroethylene in the soil was allowed to be decomposed and the byproducts which is believed to be formed in association with trichloroethylene decomposition were analyzed in a similar manner to Example 13. Furthermore, in an experiment in which the suspension of sterilized bacterial mass at an amount of 1/100 of that of the NYG medium was inoculated or the suspension was plated as it was onto the NYG agar medium and cultured at 30°C, neither increase in turbidity of the culture liquid nor colony formation were observed, and thereby it was confirmed that there was complete sterilization.

The result as shown in Table 9 indicated that all byproducts were below the detection limit. The dechlorination reaction of trichloroethylene under an anaerobic condition leads to accumulation of dichloroethylenes which is more toxic. It is also known that the decomposition process under an aerobic condition which starts with formation of trichloroethylene oxide produces dichloroacetate etc. as intermediate products. There was no accumulation of harmful substances on decomposition of trichloroethylene by the microorganism of the present invention in the soil. Although the presence of substances other than those measured is unknown, it was suggested that this technology is a highly safe method.

TABLE 9

	Item to be analyzed	Result	Detection limit
	Vinyl chloride	Not detected	0.01 ppm
30	1,1-dichloroethylene	Not detected	0.01 ppm
	cis-1,2-dichloroethylene	Not detected	0.01 ppm
	trans-1,2-dichloroethylene	Not detected	0.01 ppm
	Dichloroacetate	Not detected	0.05 ppm
	Trichloroethanol	Not detected	0.01 ppm
35	Trichloroacetate	Not detected	0.05 ppm
	Hydrated chloral	Not detected	0.05 ppm

Example 18. Construction of a constitutive mutant

A platinum loopful of the colony of the microorganism of the present invention which had been stored by passage on the agar plate containing 1.5% agar was picked and inoculated to a test tube containing 2 ml of the 1/3LB medium (its composition is shown in Table 10). After cultivating overnight under shaking at 30°C at 130 r.p.m., an aliquot of the culture broth was diluted as appropriate, which was then plated to the plate of the 1/3LB medium containing 1.5% agar. After cultivating at 30°C the number of cells was counted. The remainder of the culture broth was plated on a petri dish, which was then subjected to irradiation of a 15 W ultra violet lamp with a wavelength of 260 nm under the condition of a irradiation time of 3 minutes at a distance of 30 cm from the light source.

TABLE 10

The 1/3LB Medium		
	Tryptone	3.0 g
20	Yeast extract	1.5 g
	Sodium chloride	3.0 g
	Distilled water	1.0 liter

Then, the culture broth was diluted as appropriate and was plated to the plate of the 1/3LB medium containing 1.5% agar. After culturing at 30°C the number of cells was counted to determine the death rate. When the death rate of 99% or higher was observed, it was assumed that there was enough mutation. From the culture broth at this point, the desired mutant were selected.

Catechol 2,3-dioxygenase (C230) introduces oxygen into catechol to form 2-hydroxymuconic acid semialdehyde by metafission. This product is yellow-colored. When C230 is expressed after being sprayed, it readily turns yellow. The trichloroethylene degrading enzymes of the phenol-utilizing trichloroethylene-decomposing bacteria are known to be the phenol hydroxylase (PH) which convert

phenol into catechol (M. Fujita et al., J. Ferment. Bioeng., 79: 100, 1995; V. Shingler et al., J. Bacteriol., 174: 711, 1992).

5 An example has been known that C230 is expressed by one operon in which the C230 gene is adjacent to the PH gene though C230 itself was not directly involved in the decomposition of trichloroethylene (M. Fujita et al., J. Ferment. Bioeng., 79: 100, 1995; V. Shingler et al., J. Bacteriol., 174: 711, 1992).
10 If C230 is expressed in association with the PH, it is possible to select the strains which express the trichloroethylene-degrading enzyme using C230 expression as an indicator. The microorganism of the present invention, when cultivated using phenol as the only carbon source, decomposes phenol
15 and the culture broth turns yellow with the growth of the bacterium. This will make said selection method applicable to the microorganism.

Accordingly, the suspension of the bacterial mass after mutation was diluted as appropriate and was plated
20 to the plate of the 1/3LB medium containing 1.5% agar. After cultivating at 30°C for 1 to 3 days, a catechol solution (0.1% (w/v) ethereal solution) was sprayed.

Then, when the death rate was 99.99%, about 10,000 colonies which appeared after ultra violet irradiation
25 were investigated, which led to selection of 16 yellow colonies.


After the colonies were cultured in the 1/3LB liquid medium, it was plated to the plate of the 1/3LB medium containing 1.5% agar. After cultivating at 30°C, the
30 colonies which appeared were sprayed with a catechol solution, and yellow colonies were isolated again to obtain the strains which constitutively express C230.

A platinum loopful of these strains which constitutively express C230 were picked and inoculated
35 into the M medium (its composition is shown in Table 11) containing 4 ml of sodium glutamate (0.1%) in a 20 ml vial. The vial was plugged with a butyl rubber septum

CLAIMS

1. A bacterium which has the following properties:

	Morphology	coccoid, rod shaped
	Gram staining	+
5	Spore forming	-
	Motility	-
	Relationship to oxygen	aerobic
	Oxidase test	-
	Catalase test	+
10	Resistance to acid	-
	Rod-coccus cycle	+
	Aerial mycelium formation	-
	Peptide glycan type of cell wall	meso-diaminopimelic acid
15	Glycolyl test	- (acetyl type)
	Mycolic acid	-
	Major isoprenoid quinones	MK-8 (H4)
	GC content of DNA (mole%)	73 (by HPLC)



20 and, which can decompose trichloroethylene.

2. A bacterium according to claim 1 which is strain MO7 (FERM BP-5624).

3. A method for decomposing organic halogenated compounds and/or aromatic compounds, comprising
25 subjecting said organic halogenated compounds and/or aromatic compounds to the action of a bacterium according to claim 1 or 2.

4. A method according to claim 3 wherein the
30 organic halogenated compound is trichloroethylene and the aromatic compound is a phenolic compound.

5. A method for decomposing organic halogenated compounds in the soil, waste waters, or other waste products containing organic halogenated compounds, comprising adding a culture of a bacterium according to
35 claim 1 or 2 to the soil, waste waters, or other waste products.

6. A method according to claim 5 wherein the



compounds and/or aromatic compounds, wherein sterilized cultured bacterial cells of a microorganism capable of decomposing organic halogenated compounds and/or aromatic compounds is allowed to act on the organic halogenated compounds and/or the aromatic compounds wherein the sterilization of the cultured bacterial cells is carried out by ultraviolet irradiation, ethylene oxide treatment or radiation.

19. A method for decomposing organic halogenated compounds and/or aromatic compounds in the soil, waste waters, or other waste products containing organic halogenated compounds and/or aromatic compounds, in which a sterilized cultured bacterial mass of a microorganism capable of decomposing said organic halogenated compounds and/or said aromatic compounds is added to said soil, waste waters, or other waste products.

20. A method according to claim 19 wherein the sterilization treatment is ultraviolet irradiation.

21. A method according to claim 19 or 20, wherein the aromatic compound is a phenolic compound and the organic halogenated compound is trichloroethylene.

22. A decomposition agent for organic halogenated compounds and/or aromatic compounds, comprising a bacterial culture according to claim 1 or 2.

23. A decomposition agent according to claim 22 wherein the culture is cultured bacterial cells.

24. A decomposition agent according to claim 23 wherein the culture is living bacterial cells.

25. A decomposition agent according to claim 23 wherein the cultured bacterial cells are sterilized bacterial cells.

26. A decomposition agent according to claim 25 wherein the sterilization treatment is ultraviolet irradiation.

27. A decomposition agent according to any of claims 22 to 26 which is in the form of dried bacterial cells.



01 JUL 1999

- 59 -

28. A bacterium having the decomposition activity of organic halogenated compounds and/or aromatic compounds in the presence or absence of induction of an organic halogenated compound-decomposing enzyme by an aromatic compound, which bacterium is obtainable by
5 subjecting a bacterium according to claim 1 to a mutation treatment and a selection procedure.

29. A bacterium according to claim 28 which is the strain M0715 (FERM BP-5928).

10 30. A method for decomposing organic halogenated compounds and/or aromatic compounds comprising allowing a bacterium according to claim 28 or 29 to act on the organic halogenated compounds and/or aromatic compounds.

15 31. A method according to claim 30 wherein the organic halogenated compound is trichloroethylene and the aromatic compound is a phenolic compound.

20 32. A method for decomposing organic halogenated compounds in the soil, waste waters, or other waste products containing organic halogenated compounds, comprising adding a culture of a bacterium according to claim 28 or 29 to the soil, waste waters, or other waste products.

25 33. A method according to claim 32 wherein the organic halogenated compound is trichloroethylene and the culture is cultured bacterial cells.

34. A method according to claim 32 or 33 wherein no aromatic compound is added to the culture medium for use in culturing to obtain said bacterial culture.

30 35. A method according to any of claims 32 to 34 wherein an degradable carbon source is added to the culture medium for use in culturing to obtain said bacterial culture.

36. A method according to claim 35 wherein the degradable carbon source is glutamic acid.

35 37. A method according to any of claims 32 to 36 wherein the culture is cultured bacterial cells.



* C L M S *

38. A method according to claim 37 wherein the cultured bacterial cells are living bacterial cells.


39. A method according to claim 37 wherein the cultured bacterial cells are sterilized bacterial cells.

5 40. A method according to claim 39 wherein the sterilization treatment is ultraviolet irradiation.

41. A method for decomposing organic halogenated compounds in the soil, waste waters, or other waste products containing organic halogenated compounds, comprising the steps of inoculating a bacterium according to claim 28 or 29 to said soil, waste waters, or other waste products, adding an degradable carbon source to said soil, waste waters, or other waste products, and then culturing the inoculated microorganism.

10 42. A method according to claim 41 wherein the organic halogenated compound is trichloroethylene.

15 43. A method according to claim 41 or 42 wherein the degradable carbon source is glutamic acid.



PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12N 1/20, A62D 3/00, C02F 3/34, B09C 1/10, C12S 13/00, C12P 1/04 // (C12N 1/20, C12R 1:01)		A1	(11) International Publication Number: WO 98/07831 (43) International Publication Date: 26 February 1998 (26.02.98)
(21) International Application Number: PCT/JP97/02872 (22) International Filing Date: 19 August 1997 (19.08.97) (30) Priority Data: 8/217456 19 August 1996 (19.08.96) JP 9/109553 25 April 1997 (25.04.97) JP (71) Applicant (for all designated States except US): TOYOTA JIDOSHA KABUSHIKI KAISHA [JP/JP]; 1, Toyota-cho, Toyota-shi, Aichi 471-71 (JP). (71) Applicant (for JP only): KABUSHIKI KAISHA TOYOTA CHUO KENKYUSHO [JP/JP]; 41-1, Aza Yokomichi, Oaza Nagakute, Nagakute-cho, Aichi-gun, Aichi 480-11 (JP). (72) Inventors; and (75) Inventors/Applicants (for US only): NUMATA, Koichi [JP/JP]; Toyota Jidosha Kabushiki Kaisha, 1, Toyota-cho, Toyota-shi, Aichi 471-71 (JP). ODA, Yasushi [JP/JP]; Toyota Jidosha Kabushiki Kaisha, 1, Toyota-cho, Toyota-shi, Aichi 471-71 (JP). MIYATA, Masami [JP/JP]; Toyota Jidosha Kabushiki Kaisha, 1, Toyota-cho, Toyota-shi, Aichi 471-71 (JP). OKAMURA, Yukio [JP/JP]; Toyota Jidosha Kabushiki Kaisha, 1, Toyota-cho, Toyota-shi, Aichi 471-71			(JP). KIMURA, Toshiaki [JP/JP]; Toyota Jidosha Kabushiki Kaisha, 1, Toyota-cho, Toyota-shi, Aichi 471-71 (JP). UCHIDA, Masatoshi [JP/JP]; Toyota Jidosha Kabushiki Kaisha, 1, Toyota-cho, Toyota-shi, Aichi 471-71 (JP). ASAMI, Osamu [JP/JP]; 8-1, Aza Togo, Oaza Higashino, Konan-shi, Aichi 483 (JP). (74) Agents: ISHIDA, Takashi et al.; A. Aoki & Associates, Toranomon 37 Mori Building, 5-1, Toranomon 3-chome, Minato-ku, Tokyo 105 (JP). (81) Designated States: CA, CN, JP, KR, SG, US, VN, European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: NOVEL MICROORGANISM AND METHOD FOR ENVIRONMENTAL PURIFICATION USING THE SAME			
(57) Abstract A novel microorganism which has the following characteristics: morphology (coccoid, rod shaped), gram staining (+), spore forming (-), motility (-), relationship to oxygen (aerobic), oxidase test (-), catalase test (+), resistance to acid (-), rod-coccus cycle (+), and GC content of DNA (mole%) (73 (by HPLC)), and which can decompose chloroethylene. The microorganism can decompose in 24 hours 30 ppm of trichloroethylene, and decompose 100 ppm of trichloroethylene by 50 %.			