

SPECIFIC CONJUGATION LINKERS, SPECIFIC IMMUNOCONJUGATES THEREOF, METHODS OF MAKING AND USES SUCH CONJUGATES THEREOF

FIELD OF THE INVENTION

5 The present invention relates to linkers used for the conjugation of compounds, in particular, cytotoxic agents to specific pairs of sulfur atoms of a cell-binding molecule. The present invention also relates to methods of making cell-binding agent-drug (cytotoxic agent) conjugates in a specific manner comprising either modification of drugs with these linkers first, followed by reaction with prepared cell-binding agents; or modification of cell-binding
10 agents with these linkers first, followed by reaction with drugs.

BACKGROUND OF THE INVENTION

 The targeted delivery of highly active or potent pharmaceutical drugs by antibodies or other biological ligands to specific sites of disease in human body is an exciting approach in cancer treatment that has gained significant interest over the past few years. In particular, since US
15 FDA approvals of Adcetris (brentuximab vedotin) in 2011 and Kadcyla (ado-trastuzumab emtansine) in 2013, almost every major pharmaceutical and biotech company has adopted the applications of antibody-drug conjugate (ADC) for targeted treatment of cancers (Chari, R. et al, *Angew. Chem., Int. Ed.* 2014, 53, 3796–3827; Sievers, E. L. et al. *Annu Rev Med.* 2013, 64, 15-29; Mehrling, T. *Future Oncol*, 2015, 11, 549). In conjugated form, the cytotoxic agents
20 exhibit more selective therapeutic activity, sparing non-target cells from many of the toxic effects and improving the safety profile (Van den Mooter, T. et al *Expert Opin Biol Ther.* 2015, 15, 749-60).

 The big challenge of chemotherapeutic drugs is their narrow therapeutic windows due to they normally cannot discriminate between normal and malignant cells, thus causes side effects
25 which limit the tolerated doses below the clinically effective ones. In contrast, immunotherapy, normally in the form of monoclonal antibodies (mAb) can specifically bind to certain proteins or molecules of malignant cells, leaving normal cells unharmed, and thus has less side effects and wider therapeutic windows than chemotherapy. Monoclonal antibodies (mAb) can target
30 against malignant cells by several mechanisms, such as, 1). Making the cancer cell more visible to the immune system (Villaruz, L. C. et al, 2014, *Transl Lung Cancer Res*, 3, 2-14; Camacho, L. H. 2015 *Cancer Med* 4, 661-72); 2). Blocking growth signals (Dillman, R. O. 2011, *Cancer Biother Radiopharm*, 26, 1-64; Ferris, R. L. et al 2010, *J Clin Oncol*, 28, 4390-9); 3). Stopping new blood vessels from forming (Arrillaga-Romany, I., et al, 2014, *Expert Opin Investig Drugs*, 23, 199-210); 4). Delivering radiation to cancer cells (Chapuy, B. et al, 2007, *Biotechnol J.* 2,

1435-43); 5). Delivering chemotherapy drug to cancer cells (Chari R. J. 2008 *Acc Chem Res.* 41, 98-107; Mullard A. 2013, *Nature Reviews Drug Discovery* 12, 329–32; Zhao, R. J. 2012, *J. Med. Chem.*, 55, 766-82); and 6). Delivering enzyme to cancer cells (Francis R. J. et al, 2002, *Br. J. Cancer* 87, 600-7). One of these above strategies, delivering chemotherapy to cancer cells called antibody–drug conjugates (ADCs), which enables to target and deliver drugs to cancer cells leaving normal cells largely unaffected by the exquisite targeting ability of antibodies, has undergone intense exploitation in the last two decades. Currently there are more than 50 ADC drugs in the clinic trials according to www.clinictrials.gov.

The first-generation ADCs, including Kadcyla and Adcetris, are produced through nonselective conjugation of native lysine amines or inter chain cysteine thiols on an antibody respectively to a cytotoxic drug. Since there are over 50 surface-exposed lysines and 8 hinge cysteine residues in IgG1 antibodies, this nonselective conjugation results in randomly cross-linkage of cytotoxic drugs to practically all areas of the antibody molecule, particularly having a diverse population of ADCs with a wide distribution of drugs per antibody (DAR) (Wang, L., et al. 2005 *Protein Sci.* 14, 2436; Hamblett, K. J., et al. 2004 *Clin. Cancer Res.* 10, 7063). Thus some of the undesired ADC subpopulation could lead to shorter circulation half-life, lower efficacy, potentially increased off-target toxicity and a wide range of in vivo pharmacokinetic (PK) properties (Hamblett, K. J. et al, *Clin. Cancer Res.* 2004, 10, 7063–70; Adem, Y. T. et al, *Bioconjugate Chem.* 2014, 25, 656–664; Boylan, N. J. *Bioconjugate Chem.*, 2013, 24, 1008–1016; Strop, P., et al 2013 *Chem. Biol.* 20, 161–67). In addition, with this classical conjugation, the batch-to-batch consistency in ADC production can be challenging and may require diligent manufacturing capabilities (Wakankar, A. *mAbs*, 2011, 3, 161–172).

Therefore, biotechnology companies and academic institutions are highly focusing on establishing novel reliable methods for site-specific ADC conjugation. So far, there are several approaches developed in recent years for site selective ADC preparation (Panowski, S, 2014, *mAbs* 6, 34). They include incorporation of unpaired cysteines, e.g. engineered reactive cysteine residues, called THIOMAB from Genentech (Junutula, J. R., et al 2010 *Clin. Cancer Res.* 16, 4769; Junutula, J. R., et al 2008 *Nat Biotechnol.* 26, 925-32; US Patents 8,309,300; 7,855,275; 7,521,541; 7,723,485, WO2008/141044), genetically introduced glutamine tag with *Streptococcus* transglutaminase (mTG) (Strop, P., *Bioconjugate Chem.*, 2014, 25, 855–862; Strop, P., et al., 2013, *Chem. Biol.* 20, 161–167; US Patent 8,871,908 for Rinat-Pfizer) or with Microbial transglutaminase (MTGase) (Dennler, P., et al, 2014, *Bioconjug. Chem.* 25, 569–578. US pat appl 20130189287 for Innate Pharma; US Pat 7,893,019 for Bio-Ker S.r.l. (IT)), incorporation of thiolucose (Okeley, N. M., et al 2013

Bioconjugate Chem. 24, 1650), incorporation of unnatural amino acids through mutagenesis (Axup, J.Y., et al., 2012, Proc. Natl. Acad. Sci. 109, 16101–16106; Zimmerman, E.S., et al., 2014, Bioconjug. Chem. 25, 351–361; Wu, P., et al, 2009 Proc. Natl. Acad. Sci. 106, 3000–5; Rabuka, D., et al, 2012 Nat. Protoc. 7, 1052–67; US Patent 8,778,631 and US Pat Appl. 20100184135, WO2010/081110 for Sutro Biopharma; WO2006/069246, 2007/059312, US Patents 7,332,571, 7,696,312, and 7,638,299 for Ambrx; WO2007/130453, US patents 7,632,492 and 7,829,659 for Allozyne), Incorporation of selenocysteine into antibodies (Hofer, T., et al 2009, Biochemistry 48, 12047–12057; US Patent 8,916,159 for US National Cancer Institute), Conversion of cysteines located in the CXPXR consensus sequence to formylglycine (FGly) with formylglycine generating enzyme (FGE) (Drake, P.M., et al., 2014, Bioconjug. Chem. 25, 1331–1341. Carrico, I. S. et al 7,985,783; 8,097,701; 8,349,910, and US Pat Appl 20140141025, 20100210543 for Redwood Bioscience), and through glycoengineeringly introduction of sialic acid with the use of galactosyl- and sialyltransferases (Zhou, Q., et al 2014, Bioconjug. Chem., 25, 510–520, US Pat Appl 20140294867 for Sanofi-Genzyme). These above methods have produced nearly homogeneous product profiles, but they are required antibody-engineering processes and reoptimization of cell culture conditions. Moreover, expression yields for genetic encoding of an unnatural amino acid were typically not promisingly high enough (Tian, F., et al, 2014, Proc. Natl. Acad. Sci. U. S. A. 111, 1766–71) which has a significant impact on the cost of goods of the ADC. In addition, it has been known that ADCs obtained by conjugation to cysteine side chains often display limited stability in circulation, leading to premature disconnection of the cytotoxic payload before the tumor site is reached (Junutula, J. R., et al 2008, Nat. Biotechnol. 26, 925–32).

The disulfide bond structures of the four subclasses of IgG antibodies were known in the 1960s (Milstein C. Biochem J 1966, 101: 338–51; Pink JR, Milstein C. Nature 1967, 214: 92–4; Frangione B, Milstein C. Nature 1967, 216: 939–41; Pink JR, Milstein C. Nature 1967, 216: 941–42; Frangione B, et al. Biochem J. 1968, 106, 15–21; Frangione B, Milstein C. J Mol Biol 1968, 33: 893–906; Edelman GM, et al. Proc Natl Acad Sci USA 1969; 63: 78–85; Frangione B, et al. Nature 196, 221: 145–8, Spiegelberg, H. L. et al Biochemistry, 1975, 10, 2157–63). Disulfide bond structure is critical for the structure, stability, and biological functions of IgG molecules. Among the four subclasses of IgG antibodies, IgG₁, IgG₂, IgG₃ and IgG₄, each IgG contains a total of 12 intra-chain disulfide bonds; each disulfide bond is associated with an individual IgG domain. The two heavy chains are connected in the hinge region by a variable number of disulfide bonds: 2 for IgG₁ and IgG₄, 4 for IgG₂ and 11 for IgG₃. The light chain of the IgG₁ is connected to the heavy chain by a disulfide bond between the last cysteine

residue of the light chain and the fifth cysteine residue of the heavy chain. But, for IgG₂, IgG₃ and IgG₄, the light chain is linked to the heavy chain by a disulfide bond between the last cysteine residue of the light chain and the third cysteine residue of the heavy chain (Liu, H. and May, K., 2012, mAbs 4, 17-23). On the ranks of the susceptibility of disulfide bonds in human IgG1 antibodies by experimental reduction, differential alkylation, and LC-MS analysis (Liu, H, et al, Anal. Chem., 2010, 82, 5219–26), inter chain disulfide bonds are more susceptible to reduction than intra chain disulfide bonds, and the disulfide bonds between the light chain and heavy chain were more susceptible than disulfide bonds between the two heavy chains. The upper disulfide bond of the two inter heavy chain disulfide bonds was more susceptible than the lower one. Furthermore, disulfide bonds in the CH₂ domain were the most susceptible to reduction. Disulfide bonds in VL, CL, VH, and CH₁ domains had similar and moderate susceptibility, while disulfide bonds in the CH₃ domain were the least susceptible to reduction (Liu, H, et al Anal. Chem., 2010, 82, 5219–5226).

Based on the greater susceptibility of inter chain vs. intra chain disulfide bonds in human IgG1 antibodies, several institutions and companies adopted the chemically specific conjugation strategy through rebridging reduced inter chain disulfide bonds of a native antibody, such as, using bromo or dibromo-maleimides, called next generation maleimides (NGMs) (Schumacher, F.F., et al 2014, Org. Biomol. Chem. 12, 7261–69; UCL Cancer Institute), applying bis-alkylating reagents via a three-carbon bridge (Badescu, G., et al., 2014, Bioconjug. Chem. 25, 1124–1136., WO2013/190272, WO2014/064424 for PolyTherics Ltd), with di-substituted heteroaryl bridge (US Pat Appl. 2015/0105539 for Concortis Biosystem), or through di-maleimide as a bridge (WO2014/114207). We have also used bromo maleimide and dibromomaleimide linkers to conjugate both drugs and antibodies for a quite while (WO2014/009774, PCT/IB2012/053554). However, these above bridge linkers were designed in the way to conjugate only one cytotoxic agents to a pair of disulfide bonds, and therefore at most of time they only produced ADCs at DAR less than 2 (drugs per antibody), due to limited numbers (about two pairs) of reduced disulfide bonds are more accessible for conjugation.

As one of the major issues for ADCs is the limited numbers or amount of cytotoxic compound that ultimately reaches the tumor, and thus the favorable DAR over 3 is much important factor for improvement of ADC therapeutical index (Epenetos, A. A. et al, Cancer Res., 1986, 46, 3183–91; Chari, R. V. Acc. Chem. Res., 2008, 41, 98-107, Zhao, R. Y. et al, 2011, J. Med. Chem. 54, 3606-23), we therefore disclosed novel disulfur bridge linkers (PCT/IB2015/) that not only are able to conjugate two or more drugs per linker for achieving higher DARs (≥ 4), but also can selectively rebridge pairs of reduced inter chain disulfide bonds

on surface of antibody, which are generated by overloaded TCEP or DTT reduction agents. And the over reduced pairs of thiol groups that are inaccessibly reached by the bridge linkers can be recoupled (regenerated) by an oxide, e.g. dehydroascorbic acid (DHAA) or Cu(II), to form back disulfide bonds at the end of conjugation. In principal, this rebridging back of reduced disulfide bonds results in more stable or longer half-life of ADCs in comparison with traditional thiol linked ADCs.

It has been reported that the “ring-opened” succinimide ring linker bearing mono-thioether bond has improved in vitro stability, improved PK exposure, and improved efficacy as compared to the mono-thiol-maleimide-conjugated ADCs (Tumey, L. N, et al, 2014, Bioconjug. Chem. 25, 1871–80; Lyon, R. P, et al. 2014, Nat. Biotechnol. 32, 1059–62), due to the latter is prone to payload loss via a retro-Michael type reaction of the maleimide conjugation (Shen, B. Q, et al, 2012, Nat Biotechnol. 30, 184–9; Tumey, L. N, et al, 2014 Bioconjug Chem. 25, 1871–80). In this patent application, we extend the scopes of our earlier patent application. The bridge linkers of this invention containing a 2,3-disubstituted succinic group, or 2-monosubstituted, or 2,3-disubstituted fumaric or maleic (trans (E)- or cis (Z)-butenedioic) group, or acetylenedicarboxyl group have less payload loss as compared to their nonhydrolyzed bromo or dibromo-maleimide linkers which were tested in our lab. In other words, the methods of this invention can be used for the immunoconjugates that carry a combination of drugs and functional molecules such as PEGs, which can be used to shield the hydrolysis of conjugated drugs by hydrolases in the blood circulation as well as inhibit the degradation of the antibody by proteinases. Thus the major advantages of this patent for immunoconjugates include: prolonged the half lives of the conjugates during the targeted delivery; conjugated in steps of two or more different function molecules/drugs that act in different phases of the cell cycle to increase the number of target cells exposed to the particular pharmaceutical drugs or effectors; minimized exposure to non-target cells, tissues or organs through conjugation of the function molecules; precisely controlled over drug payloads and drug ratios at the specific sites leading to homogenous final products. In short, the bridge linkers of the invention can stabilize ADCs as well as make homogeneous production of ADCs in a simple manner.


SUMMARY OF THE INVENTION

The present invention provides linkers containing a 2,3-disubstituted succinic group, or 2-monosubstituted, or 2,3-disubstituted fumaric or maleic (trans (E)- or cis (Z)-butenedioic) group, or acetylenedicarboxyl group to link a drug and/or a function molecule to a cell-binding agent

(e.g., an antibody). The preferred formula of the cell-binding molecule-linker-drug conjugates

can be represented as:
$$\text{Cb} \left[\begin{array}{c} \text{S} \diagup \text{L} \text{---} (\text{Drug}_1)_{m_1} \\ \text{S} \diagdown \text{L} \text{---} (\text{R}_2)_{m_2} \end{array} \right]_n$$
, wherein Cb is a cell-binding agent, L is a linker containing succinic, fumaric or maleic group; Drug₁ is a drug molecule; m_1 , m_2 and n are an integer from 1 to 30; and two S (sulfur) elements from Cb bridgely link to L, which covalently connects a drug, Drug₁ and a function molecule, R₂, which is for stabilizing the conjugated “Drug₁” during the circulation. The advantages in applying the linker in the cell molecule-drug conjugate are: a). Retaining the stability of the conjugates by covalently cross-linking (re-bridging) the pairs of reduced disulfur atoms of the cell-binding agents, particularly of antibodies; b). Enabling conjugation of the cytotoxic agents/drugs to specific sites of a cell-binding molecule, e.g. the inter chain disulfide bond sites of IgG antibodies, resulting in homogeneous production of ADCs.

In one aspect of the present invention, the linker is represented by Formula (I)



Wherein

— represents an optional single bond.

=== represents either a single bond, or a double bond, or a triple bond.

It provided that when === represents a single bond, both U and U' are not H; when === represents a double bond, either U or U' can be H, but they are not H at the same time; when === represents a triple bond, both U and U' can be absent.

U and U' represent the same or different leaving group that can be substituted by a thiol. Such leaving groups are, but are not limited to, a halide (e.g., fluoride, chloride, bromide, and iodide), methanesulfonyl (mesyl), toluenesulfonyl (tosyl), trifluoromethyl-sulfonyl (triflate), trifluoromethylsulfonate, nitrophenoxyl, N-succinimidyl (NHS), phenoxyl; dinitrophenoxyl; pentafluorophenoxyl, tetrafluorophenoxyl, trifluorophenoxyl, difluorophenoxyl, monofluorophenoxyl, pentachlorophenoxyl, 1H-imidazole-1-yl, chlorophenoxyl, dichlorophenoxyl, trichlorophenoxyl, tetrachlorophenoxyl, N-(benzotriazol-yl)oxyl, 2-ethyl-5-phenylisoxazolium-yl, phenyloxadiazol-yl (ODA), oxadiazol-yl, or an intermediate molecule generated with a condensation reagent for Mitsunobu reactions.

Z₁ is a function group that enables to react with a cytotoxic drug, to form a disulfide, ether, ester, thioether, thioester, peptide, hydrazone, carbamate, carbonate, amine (secondary, tertiary, or quarter), imine, cycloheteroalkane, heteroaromatic, alkyloxime or amide bond;

R₁ can be absent, or can be selected from C₁-C₈ of alkyl; C₂-C₈ of heteroalkyl, alkylcycloalkyl, heterocycloalkyl; C₃-C₈ of aryl, Ar-alkyl, heterocyclic, carbocyclic, cycloalkyl, heteroalkylcycloalkyl, alkylcarbonyl, heteroaryl; or 1-8 carbon atoms of esters, ether, or amide; or polyethyleneoxy unit of formula (OCH₂CH₂)_p or (OCH₂CH(CH₃))_p, wherein p is an integer from 0 to about 1000, or combination of above groups thereof.

Additionally R₁ is a chain of atoms selected from C, N, O, S, Si, and P, preferably having 0~500 atoms, which covalently connects to X₁ and Z₁. The atoms used in forming the R₁ may be combined in all chemically relevant ways, such as forming alkylene, alkenylene, and alkynylene, ethers, polyoxyalkylene, esters, amines, imines, polyamines, hydrazines, hydrazones, amides, ureas, semicarbazides, carbazides, alkoxyamines, alkoxyamines, urethanes, amino acids, peptides, acyloxyamines, hydroxamic acids, or combination above thereof.


X₁ and X₂ are independently selected from NH; NHNH; N(R₃); N(R₃)N(R₃′); O; S; C₁-C₈ of alkyl; C₂-C₈ of heteroalkyl, alkylcycloalkyl, heterocycloalkyl; C₃-C₈ of aryl, Ar-alkyl, heterocyclic, carbocyclic, cycloalkyl, heteroalkylcycloalkyl, alkylcarbonyl, heteroaryl; or 1~8 amino acids; Wherein R₃ and R₃′ are independently H; C₁-C₈ of alkyl; C₂-C₈ of hetero-alkyl, alkylcycloalkyl, heterocycloalkyl; C₃-C₈ of aryl, Ar-alkyl, heterocyclic, carbocyclic, cycloalkyl, heteroalkylcycloalkyl, alkylcarbonyl, heteroaryl; or 1-8 carbon atoms of esters, ether, or amide; or polyethyleneoxy unit of formula (OCH₂CH₂)_p or (OCH₂CH(CH₃))_p, wherein p is an integer from 0 to about 1000, or combination above thereof.

R₂ is independently selected from OH, H, NH₂, SH, NHNH₂, N(R₃)(R₃′), N(R₃)NH(R₃′), polyethyleneoxy unit of formula (OCH₂CH₂)_pOR₃, or (OCH₂CH(CH₃))_pOR₃, or NH(CH₂CH₂O)_pR₃, or NH(CH₂CH(CH₃)O)_pR₃, or N[(CH₂CH₂O)_pR₃][(CH₂CH₂O)_p′R₃′], or (OCH₂CH₂)_pCOOR₃, or CH₂CH₂(OCH₂CH₂)_pCOOR₃, wherein p and p′ are independently an integer selected from 0 to about 1000, or combination thereof; C₁-C₈ of alkyl; C₂-C₈ of heteroalkyl, alkylcycloalkyl, heterocycloalkyl; C₃-C₈ of aryl, Ar-alkyl, heterocyclic, carbocyclic, cycloalkyl, heteroalkylcycloalkyl, alkylcarbonyl, heteroaryl; Wherein R₃ and R₃′ are independently H; C₁-C₈ of alkyl; C₂-C₈ of heteroalkyl, alkylcycloalkyl, heterocycloalkyl; C₃-C₈ of aryl, Ar-alkyl, heterocyclic, carbocyclic, cycloalkyl, heteroalkylcycloalkyl, alkylcarbonyl, heteroaryl; or 1-8 carbon atoms of esters, ether, or amide; or 1~8 amino acids; or polyethyleneoxy unit of formula (OCH₂CH₂)_p or (OCH₂CH(CH₃))_p, wherein p is an integer from 0 to about 1000, or combination above thereof.

m₁ and m₂ are independently an integer from 1 to 30, preferably from 1 to 10.

Wherein “—”, Z₁, Z₂, n, R₁, R₂, m₁, m₂, X₁, and X₂ are defined the same as in Formula (I).
 “====” and Cb are defined the same as in Formula (II).

In an even further aspect, the present invention provides a modified drug of Formula (IV), in which the drug, Drug₁, has reacted with the linker of Formula (I), which still has the 2,3-disubstituted succinic group, or 2-monosubstituted, or 2,3-disubstituted fumaric or maleic (*trans* (E)- or *cis* (Z)-butenedioic) group, or acetylenedicarboxyl group capable of reacting with a pair of thiols of the cell-binding agent:



Wherein “—”, “====”, U, U', R₁, R₂, m₁, m₂, X₁, and X₂ are defined the same as in Formula (I). Drug₁ and Cb are defined the same as in Formula (II).

The present invention further relates to a method of making a cell-binding molecule-drug conjugate of Formula (II), wherein the drugs, “Drug₁” is linked to a cell-binding agent via the bridge linker.

The present invention also relates to a method of making a modified cell-binding molecule of Formula (III), wherein the cell-binding molecule is reacted with the bridge linker of Formula (I).

The present invention also relates to a method of making a modified drug of formula (IV), wherein the drug is reacted with the bridge linker of Formula (I).

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 shows the synthesis of a bridge linker containing a drug and a polyethylene glycol, and the application of this linker in the conjugation of the two different molecules to an antibody via amide bonds.

Figure 2 shows the synthesis of bridge linkers containing a drug and a polyethylene glycol, and the application of the linkers in the conjugation of the two different molecules to an antibody via amide bonds.

Figure 3 shows the synthesis of bridge linkers containing a drug and a polyethylene glycol, and the application of the linkers in the conjugation of the two different molecules to an antibody via amide bonds.

Figure 4 shows the synthesis of bridge linkers containing a drug and a polyethylene glycol, and the application of the linkers in the conjugation of the two different molecules to an antibody via oxime linkage.

Figure 5 shows the synthesis of bridge linkers containing a drug and a polyethylene glycol, and the application of the linkers in the conjugation of the two different molecules to an antibody via hydrazone linkage.

Figure 6 shows the synthesis of bridge linkers containing a drug and a polyethylene glycol, and the application of the linkers in the conjugation of the two different molecules to an antibody via amide bonds.

Figure 7 shows the synthesis of bridge linkers containing a drug, a polyethylene glycol or short alkyl acid, and the conjugation of the drug via these linkers to an antibody.

Figure 8 shows the synthesis of bridge linkers containing a drug, a polyethylene glycol or peptides, and the application in the conjugation of the bridge linker-drug molecules to an antibody via hinder amide linkage.

Figure 9 shows the synthesis of bridge linkers containing peptides or polyethylene glycols, and the application of these linkers in the conjugation of the bridge linker-drug molecules to an antibody via amide linkage.

Figure 10 shows the synthesis of the components of Tubulysin analogs.

Figure 11 shows the synthesis of the components of bridge linkers containing polyethylene glycol.

Figure 12 shows the the synthesis of the linkers of this patent application.

Figure 13 shows the the synthesis the conjugates of antibody-tubulysin analogs via a bridge linker of this patent application.

Figure 14 shows the the synthesis the conjugates of antibody-tubulysin analogs via a bridge-linker of this patent application.

Figure 15 shows the synthesis of the components of MMAF analogs.

Figure 16 shows the synthesis of a conjugate of antibody- PBD dimer analog via the bridge-linker.

Figure 17 shows the synthesis of conjugates of antibody- PBD dimer analogs via the bridge-linkers of this patent application.

Figure 18 shows the synthesis of conjugates of antibody- MMAF analogs via the bridge linkers of this patent application.

Figure 19 shows the synthesis of conjugates of antibody- MMAF analogs via the bridge linkers of this patent application.

Figure 20 shows the synthesis of conjugates of antibody- tubulysin analogs via the bridge linkers of this patent application.

Figure 21 shows the synthesis of the conjugates of Maytansinoid analogs via the bridge linkers of this patent application.

Figure 22 shows the synthesis of the conjugates of MMAF analogs, tubulysin analogs and amatoxin analogs via the bridge linkers of this patent application.

5 Figure 23 shows the synthesis of the conjugates of MMAF analogs, tubulysin analogs and amatoxin analogs via the bridge linkers of this patent application.

Figure 24 shows the synthesis of the conjugates of amatoxin analogs via the bridge linkers of this patent application.

10 Figure 25 shows the synthesis of the conjugates of MMAF analogs and tubulysin analogs via the bridge linkers of this patent application.

Figure 26 shows the synthesis of the conjugates containing both PBD dimer analogs and polyethylene glycols, both PBD dimer analogs and MMAF analogs, and both PBD dimer analog and Tubulysin analogs, to an antibody via the bridge linkers of this patent.

15 Figure 27 shows the synthesis of the conjugates containing both PBD dimer analogs and Tubulysin analogs, both Tubulysin analogs and polyethylene glycols, and both MMAF analogs and polyethylene glycols, to an antibody via the bridge linkers of this patent application.

Figure 28 shows the synthesis of the conjugates containing both MMAF analogs and polyethylene glycols, and Tubulysin B analogs via cyclic linkage of the bridge linkers of this patent application.

20 Figure 29 shows the synthesis of the conjugates of tubulysin analogs via the bridge linkers of this patent application.

Figure 30 shows the synthesis of the conjugates of tubulysin analogs via the bridge linkers of this patent application.

25 Figure 31 shows the synthesis of the conjugates of MMAE analogs via the bridge linkers of this patent application.

Figure 32 shows the synthesis of the conjugates of MMAE analogs via the bridge linkers of this patent application.

Figure 33 shows the synthesis of the conjugates of MMAF analogs and tubulysin analogs via the bridge linkers of this patent application.

30 Figure 34 shows the synthesis of the conjugates of tubulysin analogs via the bridge linkers of this patent application.

Figure 35 shows the SDS-PAGE gel containing reduce agent DTT in the development. Lane 1 is biomarker, Lane 2 is conjugate **201**, Lane 3 is conjugate **206**, Lane 4 is conjugate **354**, Lane 5 is Tubulysin analog conjugate through the traditional maleimidopropanoyl linker that we

disclosed previously (Huang Y. et al, Med Chem. #44, 249th ACS National Meeting, Denver, CO, Mar. 22~26, 2015; WO2014009774). The conjugates **201**, **206** and **354** via the bridge linkers of this patent application have the major band of 75KD demonstrated that the heavy chain and the light chain of the mAb were crossly linked with the bridge linkers. In comparison, the traditional maleimido linker has no 75KD band. and the 50KD band is the single heavy chain of the antibody.

Figure 36 shows the comparison of the anti-tumor effect of conjugate compounds **196a**, **201**, **206**, **354**, **358**, **377**, **385**, and **407** with T-DM1 using human gastric tumor N87 cell model at dosing of 6 mg/kg, i.v., one injection. The bottom figure indicates that all 9 conjugates did not cause the animal body weight loss. The bottom figures shows that conjugate compounds **196a**, **201**, **206**, **27**, **354**, and **407** had better antitumor activity than T-DM1. In particular, compounds **201** and **407** were able to eliminate the tumors until end of experiment.

DETAILED DESCRIPTION OF THE INVENTION

DEFINITIONS

“Alkyl” refers to an aliphatic hydrocarbon group or univalent groups derived from alkane by removal of one or two hydrogen atoms from carbon atoms. It may be straight or branched having 1 to 8 carbon atoms in the chain. “Branched” means that one or more lower C numbers of alkyl groups such as methyl, ethyl or propyl are attached to a linear alkyl chain. Exemplary alkyl groups include methyl, ethyl, *n*-propyl, *i*-propyl, *n*-butyl, *t*-butyl, *n*-pentyl, 3-pentyl, octyl, nonyl, decyl, cyclopentyl, cyclohexyl, 2,2-dimethylbutyl, 2,3-dimethylbutyl, 2,2-dimethylpentyl, 2,3-dimethylpentyl, 3,3-dimethylpentyl, 2,3,4-trimethylpentyl, 3-methyl-hexyl, 2,2-dimethylhexyl, 2,4-dimethylhexyl, 2,5-dimethylhexyl, 3,5-dimethylhexyl, 2,4-dimethylpentyl, 2-methylheptyl, 3-methylheptyl, *n*-heptyl, isoheptyl, *n*-octyl, and isooctyl. A C₁-C₈ alkyl group can be unsubstituted or substituted with one or more groups including, but not limited to, -C₁-C₈ alkyl, -O-(C₁-C₈ alkyl), -aryl, -C(O)R', -OC(O)R', -C(O)OR', -C(O)NH₂, -C(O)NHR', -C(O)N(R')₂, -NHC(O)R', -SR', -S(O)₂R', -S(O)R', -OH, -halogen, -N₃, -NH₂, -NH(R'), -N(R')₂ and -CN; where each R' is independently selected from -C₁-C₈ alkyl and aryl.

“Halogen” refers to fluorine, chlorine, bromine or iodine atom; preferably fluorine and chlorine atom.

“Heteroalkyl” refers to C₂-C₈ alkyl in which one to four carbon atoms are independently replaced with a heteroatom from the group consisting of O, S and N.

“Carbocycle” refers to a saturated or unsaturated ring having 3 to 8 carbon atoms as a monocycle or 7 to 13 carbon atoms as a bicycle. Monocyclic carbocycles have 3 to 6 ring atoms, more typically 5 or 6 ring atoms. Bicyclic carbocycles have 7 to 12 ring atoms, arranged as a

bicycle [4,5], [5,5], [5,6] or [6,6] system, or 9 or 10 ring atoms arranged as a bicycle [5,6] or [6,6] system. Representative C₃-C₈ carbocycles include, but are not limited to, -cyclopropyl, -cyclobutyl, -cyclopentyl, -cyclopentadienyl, -cyclohexyl, -cyclohexenyl, -1,3-cyclohexadienyl, -1,4-cyclohexadienyl, -cycloheptyl, -1,3-cycloheptadienyl, -1,3,5-cycloheptatrienyl, -cyclooctyl, and -cyclooctadienyl.

A “C₃-C₈ carbocycle” refers to a 3-, 4-, 5-, 6-, 7- or 8-membered saturated or unsaturated nonaromatic carbocyclic ring. A C₃-C₈ carbocycle group can be unsubstituted or substituted with one or more groups including, but not limited to, -C₁-C₈ alkyl, -O-(C₁-C₈ alkyl), -aryl, -C(O)R', -OC(O)R', -C(O)OR', -C(O)NH₂, -C(O)NHR', -C(O)N(R')₂, -NHC(O)R', -SR', -S(O)R', -S(O)₂R', -OH, -halogen, -N₃, -NH₂, -NH(R'), -N(R')₂ and -CN; where each R' is independently selected from -C₁-C₈ alkyl and aryl.

“Alkenyl” refers to an aliphatic hydrocarbon group containing a carbon-carbon double bond which may be straight or branched having 2 to 8 carbon atoms in the chain. Exemplary alkenyl groups include ethenyl, propenyl, *n*-butenyl, *i*-butenyl, 3-methylbut-2-enyl, *n*-pentenyl, hexylenyl, heptenyl, octenyl.

“Alkynyl” refers to an aliphatic hydrocarbon group containing a carbon-carbon triple bond which may be straight or branched having 2 to 8 carbon atoms in the chain. Exemplary alkynyl groups include ethynyl, propynyl, *n*-butynyl, 2-butylnyl, 3-methylbutynyl, 5-pentylnyl, *n*-pentylnyl, hexylnyl, heptynyl, and octynyl.

“Alkylene” refers to a saturated, branched or straight chain or cyclic hydrocarbon radical of 1-18 carbon atoms, and having two monovalent radical centers derived by the removal of two hydrogen atoms from the same or two different carbon atoms of a parent alkane. Typical alkylene radicals include, but are not limited to: methylene (-CH₂-), 1,2-ethyl (-CH₂CH₂-), 1,3-propyl (-CH₂CH₂CH₂-), 1,4-butyl (-CH₂CH₂CH₂CH₂-), and the like.

“Alkenylene” refers to an unsaturated, branched or straight chain or cyclic hydrocarbon radical of 2-18 carbon atoms, and having two monovalent radical centers derived by the removal of two hydrogen atoms from the same or two different carbon atoms of a parent alkene. Typical alkenylene radicals include, but are not limited to: 1,2-ethylene (-CH=CH-).

“Alkynylene” refers to an unsaturated, branched or straight chain or cyclic hydrocarbon radical of 2-18 carbon atoms, and having two monovalent radical centers derived by the removal of two hydrogen atoms from the same or two different carbon atoms of a parent alkyne. Typical alkynylene radicals include, but are not limited to: acetylene, propargyl and 4-pentylnyl.

“Aryl” or Ar refers to an aromatic or hetero aromatic group, composed of one or several rings, comprising three to fourteen carbon atoms, preferentially six to ten carbon atoms. The term

of "hetero aromatic group" refers one or several carbon on aromatic group, preferentially one, two, three or four carbon atoms are replaced by O, N, Si, Se, P or S, preferentially by O, S, and N. The term aryl or Ar also refers to an aromatic group, wherein one or several H atoms are replaced independently by -R', -halogen, -OR', or -SR', -NR'R'', -N=NR', -N=R', -NR'R'', -NO₂, -S(O)R', -S(O)₂R', -S(O)₂OR', -OS(O)₂OR', -PR'R'', -P(O)R'R'', -P(OR')(OR''), -P(O)(OR')(OR'') or -OP(O)(OR')(OR'') wherein R', R'' are independently H, alkyl, alkenyl, alkynyl, heteroalkyl, aryl, arylalkyl, carbonyl, or pharmaceutical salts.

"Heterocycle" refers to a ring system in which one to four of the ring carbon atoms are independently replaced with a heteroatom from the group of O, N, S, Se, B, Si and P. Preferable heteroatoms are O, N and S. Heterocycles are also described in The Handbook of Chemistry and Physics, 78th Edition, CRC Press, Inc., 1997-1998, p. 225 to 226, the disclosure of which is hereby incorporated by reference. Preferred nonaromatic heterocyclic include, but are not limited to epoxy, aziridinyl, thiiranyl, pyrrolidinyl, pyrazolidinyl, imidazolidinyl, oxiranyl, tetrahydrofuranyl, dioxolanyl, tetrahydropyranyl, dioxanyl, dioxolanyl, piperidyl, piperazinyl, morpholinyl, pyranal, imidazolinyl, pyrrolinyl, pyrazolinyl, thiazolidinyl, tetrahydrothiopyranal, dithianyl, thiomorpholinyl, dihydropyranyl, tetrahydropyranyl, dihydropyranyl, tetrahydropyridyl, dihydropyridyl, tetrahydropyrimidinyl, dihydrothiopyranal, azepanyl, as well as the fused systems resulting from the condensation with a phenyl group.

The term "heteroaryl" or aromatic heterocycles refers to a 3 to 14, preferably 5 to 10 membered aromatic hetero, mono-, bi- or multicyclic ring. Examples include pyrrolyl, pyridyl, pyrazolyl, thienyl, pyrimidinyl, pyrazinyl, tetrazolyl, indolyl, quinolyl, purinyl, imidazolyl, thienyl, thiazolyl, benzothiazolyl, furanyl, benzofuranyl, 1,2,4-thiadiazolyl, isothiazolyl, triazolyl, tetrazolyl, isoquinolyl, benzothienyl, isobenzofuryl, pyrazolyl, carbazolyl, benzimidazolyl, isoxazolyl, pyridyl-N-oxide, as well as the fused systems resulting from the condensation with a phenyl group.

"Alkyl", "cycloalkyl", "alkenyl", "alkynyl", "aryl", "heteroaryl", "heterocyclic" and the like refer also to the corresponding "alkylene", "cycloalkylene", "alkenylene", "alkynylene", "arylene", "heteroarylene", "heterocyclene" and the likes which are formed by the removal of two hydrogen atoms.

"Arylalkyl" refers to an acyclic alkyl radical in which one of the hydrogen atoms bonded to a carbon atom, typically a terminal or sp³ carbon atom, is replaced with an aryl radical. Typical arylalkyl groups include, but are not limited to, benzyl, 2-phenylethan-1-yl, 2-phenylethen-1-yl, naphthylmethyl, 2-naphthylethan-1-yl, 2-naphthylethen-1-yl, naphthobenzyl, 2-naphthophenylethan-1-yl and the like.

“Heteroarylalkyl” refers to an acyclic alkyl radical in which one of the hydrogen atoms bonded to a carbon atom, typically a terminal or sp^3 carbon atom, is replaced with a heteroaryl radical. Typical heteroarylalkyl groups include, but are not limited to, 2-benzimidazolymethyl, 2-furylethyl and the like.

Examples of a “hydroxyl protecting group” include, but are not limited to, methoxymethyl ether, 2-methoxyethoxymethyl ether, tetrahydropyranyl ether, benzyl ether, *p*-methoxybenzyl ether, trimethylsilyl ether, triethylsilyl ether, triisopropylsilyl ether, *t*-butyldimethylsilyl ether, triphenylmethylsilyl ether, acetate ester, substituted acetate esters, pivaloate, benzoate, methanesulfonate and *p*-toluenesulfonate.

“Leaving group” refers to a functional group that can be substituted by another functional group. Such leaving groups are well known in the art, and examples include, but are not limited to, a halide (e.g., chloride, bromide, and iodide), methanesulfonyl (mesyl), *p*-toluenesulfonyl (tosyl), trifluoromethylsulfonyl (triflate), and trifluoromethylsulfonate.

The following abbreviations may be used herein and have the indicated definitions: Boc, tert-butoxy carbonyl; BroP, bromotrispyrrolidinophosphonium hexafluorophosphate; CDI, 1,1'-carbonyldiimidazole; DCC, dicyclohexylcarbodiimide; DCE, dichloroethane; DCM, dichloromethane; DIAD, diisopropylazodicarboxylate; DIBAL-H, diisobutyl-aluminium hydride; DIPEA, diisopropylethylamine; DEPC, diethyl phosphorocyanidate; DMA, N,N-dimethyl acetamide; DMAP, 4-(N, N-dimethylamino)pyridine; DMF, N,N-dimethylformamide; DMSO, dimethylsulfoxide; DTT, dithiothreitol; EDC, 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride; ESI-MS, electrospray mass spectrometry; HATU, O-(7-azabenzotriazol-1-yl)-N, N, N', N'-tetramethyluronium hexafluorophosphate; HOBt, 1-hydroxybenzotriazole; HPLC, high pressure liquid chromatography; NHS, N-Hydroxysuccinimide; MMP, 4-methylmorpholine; PAB, *p*-aminobenzyl; PBS, phosphate-buffered saline (pH 7.0~7.5); PEG, polyethylene glycol; SEC, size-exclusion chromatography; TCEP, tris(2-carboxyethyl)phosphine; TFA, trifluoroacetic acid; THF, tetrahydrofuran; Val, valine.

The “amino acid(s)” can be natural and/or unnatural amino acids, preferably alpha-amino acids. Natural amino acids are those encoded by the genetic code, which are alanine, arginine, asparagine, aspartic acid, cysteine, glutamic acid, glutamine, glycine, histidine, isoleucine, leucine, lysine, methionine, phenylalanine, proline, serine, threonine, tyrosine, tryptophan and valine. The unnatural amino acids are derived forms of proteinogenic amino acids. Examples include hydroxyproline, lanthionine, 2-aminoisobutyric acid, dehydroalanine, gamma-aminobutyric acid (the neurotransmitter), ornithine, citrulline, beta alanine (3-aminopropanoic

acid), gamma-carboxyglutamate, selenocysteine (present in many noneukaryotes as well as most eukaryotes, but not coded directly by DNA), pyrrolysine (found only in some archaea and one bacterium), N-formylmethionine (which is often the initial amino acid of proteins in bacteria, mitochondria, and chloroplasts), 5-hydroxytryptophan, L-dihydroxyphenylalanine,

5 triiodothyronine, L-3,4-dihydroxyphenylalanine (DOPA), and O-phosphoserine. The term amino acid also includes amino acid analogs and mimetics. Analogs are compounds having the same general $\text{H}_2\text{N(R)CHCO}_2\text{H}$ structure of a natural amino acid, except that the R group is not one found among the natural amino acids. Examples of analogs include homoserine, norleucine, methionine-sulfoxide, and methionine methyl sulfonium. Preferably, an amino acid mimetic is a
10 compound that has a structure different from the general chemical structure of an alpha-amino acid but functions in a manner similar to one. The term "unnatural amino acid" is intended to represent the "D" stereochemical form, the natural amino acids being of the "L" form. When 1~8 amino acids are used in this patent application, amino acid sequence is then preferably a cleavage recognition sequence for a protease. Many cleavage recognition sequences are known in the art.

15 See, e.g., Matayoshi et al. Science 247: 954 (1990); Dunn et al. Meth. Enzymol. 241: 254 (1994); Seidah et al. Meth. Enzymol. 244: 175 (1994); Thornberry, Meth. Enzymol. 244: 615 (1994); Weber et al. Meth. Enzymol. 244: 595 (1994); Smith et al. Meth. Enzymol. 244: 412 (1994); and Bouvier et al. Meth. Enzymol. 248: 614 (1995); the disclosures of which are incorporated herein by reference. In particular, the sequence is selected from the group consisting of Val-Cit, Ala-
20 Val, Val-Ala-Val, Lys-Lys, Ala-Asn-Val, Val-Leu-Lys, Cit-Cit, Val-Lys, Ala-Ala-Asn, Lys, Cit, Ser, and Glu.

“Pharmaceutically” or “pharmaceutically acceptable” refer to molecular entities and compositions that do not produce an adverse, allergic or other untoward reaction when administered to an animal, or a human, as appropriate.

25 “Pharmaceutically acceptable solvate” or “solvate” refer to an association of one or more solvent molecules and a disclosed compound. Examples of solvents that form pharmaceutically acceptable solvates include, but are not limited to, water, isopropanol, ethanol, methanol, DMSO, ethyl acetate, acetic acid and ethanolamine.

30 “Pharmaceutically acceptable excipient” includes any carriers, diluents, adjuvants, or vehicles, such as preserving or antioxidant agents, fillers, disintegrating agents, wetting agents, emulsifying agents, suspending agents, solvents, dispersion media, coatings, antibacterial and antifungal agents, isotonic and absorption delaying agents and the like. The use of such media and agents for pharmaceutical active substances is well known in the art. Except insofar as any conventional media or agent is incompatible with the active ingredient, its use in the therapeutic

compositions is contemplated. Supplementary active ingredients can also be incorporated into the compositions as suitable therapeutic combinations.

As used herein, "pharmaceutical salts" refer to derivatives of the disclosed compounds wherein the parent compound is modified by making acid or base salts thereof. The pharmaceutically acceptable salts include the conventional non-toxic salts or the quaternary ammonium salts of the parent compound formed, for example, from non-toxic inorganic or organic acids. For example, such conventional non-toxic salts include those derived from inorganic acids such as hydrochloric, hydrobromic, sulfuric, sulfamic, phosphoric, nitric and the like; and the salts prepared from organic acids such as acetic, propionic, succinic, tartaric, citric, methanesulfonic, benzenesulfonic, glucuronic, glutamic, benzoic, salicylic, toluenesulfonic, oxalic, fumaric, maleic, lactic and the like. Further addition salts include ammonium salts such as tromethamine, meglumine, epolamine, etc., metal salts such as sodium, potassium, calcium, zinc or magnesium.

The pharmaceutical salts of the present invention can be synthesized from the parent compound which contains a basic or acidic moiety by conventional chemical methods. Generally, such salts can be prepared via reaction the free acidic or basic forms of these compounds with a stoichiometric amount of the appropriate base or acid in water or in an organic solvent, or in a mixture of the two. Generally, non-aqueous media like ether, ethyl acetate, ethanol, isopropanol, or acetonitrile are preferred. Lists of suitable salts are found in Remington's Pharmaceutical Sciences, 17th ed., Mack Publishing Company, Easton, PA, 1985, p. 1418, the disclosure of which is hereby incorporated by reference.

"Administering" or "administration" refers to any mode of transferring, delivering, introducing or transporting a pharmaceutical drug or other agent to a subject. Such modes include oral administration, topical contact, intravenous, intraperitoneal, intramuscular, intralesional, intranasal, subcutaneous or intrathecal administration. Also contemplated by the present invention is utilization of a device or instrument in administering an agent. Such device may utilize active or passive transport and may be slow-release or fast-release delivery device.


The novel conjugates disclosed herein use the bridge linkers. Examples of some suitable linkers and their synthesis are shown in Figures 1 to 34.

THE BRIDGE LINKERS

The synthetic routes to produce bridge linkers as well as the preparation of the conjugates of drugs to a cell binding molecules of the present invention are shown in Figures 1-34. The bridge linkers possess two elements: a) A Substituent that is a 2,3-disubstituted succinic group; or 2-

monosubstituted, or 2,3-disubstituted fumaric group; or 2-monosubstituted, or 2,3-disubstituted maleic group; or acetylenedicarboxyl group, which can react to a pair of thiols to form covalent thioether bonds, and b) A group, such as but not limited to, a disulfide, maleimide, haloacetyl, aldehyde, ketone, azide, amine, alkoxyamine, hydrazide, ethenesulfonyl, acyl halide (acid halide), acryl (acryloyl), and/or acid anhydride group, capable of reaction with a drug. The bridge substituents of 2,3-disubstituted succinic group; or 2-monosubstituted, or 2,3-disubstituted fumaric group; or 2-monosubstituted, or 2,3-disubstituted maleic group; or acetylenedicarboxyl group; can be introduced by direct condensation of these 2,3-disubstituted succinic acid, or 2-monosubstituted or 2,3-disubstituted fumaric or maleic, or acetylenedicarboxyl groups with an amine, an alcohol, or a thiol group to form amide, ester or thioester bonds. The synthesis of these bridge linkers and their application for antibody conjugation are exemplified in the Figures 1-34.

Preferably, the bridge linkers are compounds of the Formula (I) below:




Wherein

“—” represents an optional single bond;

“===” represents either a single bond, or a double bond, or a triple bond.

It provided that when === represents a single bond, both U and U' are not H; when === represents a double bond, either U or U' can be H, but are not H at the same time; when === represents a triple bond, both U and U' can be absent.



U and U' represent the same or different leaving group that can be substituted by a thiol. Such leaving groups are, but are not limited to, a halide (e.g., fluoride, chloride, bromide, and iodide), methanesulfonyl (mesyl), toluenesulfonyl (tosyl), trifluoromethyl-sulfonyl (triflate),

trifluoromethylsulfonate, nitrophenoxyl, N-succinimidyl (NHS), phenoxyl; dinitrophenoxyl; pentafluorophenoxyl, tetrafluorophenoxyl, trifluorophenoxyl, difluorophenoxyl, monofluorophenoxyl, pentachlorophenoxyl, 1H-imidazole-1-yl, chlorophenoxyl, dichlorophenoxyl, trichlorophenoxyl, tetrachlorophenoxyl, N-(benzotriazol-yl)oxyl, 2-ethyl-5-phenylisoxazolium-3'-sulfonyl, phenyloxadiazole-sulfonyl (-sulfone-ODA), oxadiazol-yl, or an intermediate molecule generated with a condensation reagent for Mitsunobu reactions.

Z_1 and Z_2 are the same or different a function group that enables to react with a cytotoxic drug, to form a disulfide, thioether, thioester, peptide, hydrazone, ether, ester, carbamate, carbonate, amine (secondary, tertiary, or quarter), imine, cycloheteroalkane, heteroaromatic, alkoxime or amide bond;

R_1 can be absent, or can be selected from C_1 - C_8 of alkyl; C_2 - C_8 of heteroalkyl, alkylcycloalkyl, heterocycloalkyl; C_3 - C_8 of aryl, Ar-alkyl, heterocyclic, carbocyclic, cycloalkyl, heteroalkylcycloalkyl, alkylcarbonyl, heteroaryl; or 1-8 carbon atoms of esters, ether, or amide; or polyethyleneoxy unit of formula $(OCH_2CH_2)_p$ or $(OCH_2CH(CH_3))_p$, wherein p is an integer from 0 to about 1000, or combination thereof.

Additionally R_1 is a chain of atoms selected from C, N, O, S, Si, and P, preferably having 0~500 atoms, which covalently connects to X_1 and Z_1 . The atoms used in forming the R_1 may be combined in all chemically relevant ways, such as forming alkane, alkenylene, alkynylene, ethers, polyoxyalkylene, esters, amines, imines, polyamines, hydrazines, hydrazones, amides, ureas, semicarbazides, carbazides, alkoxyamines, alkoxyamines, urethanes, amino acids, peptides, acyloxyamines, hydroxamic acids, or combination thereof.

X_1 and X_2 are independently selected from NH; $NHNH$; $N(R_3)$; $N(R_3)N(R_3')$; O; S; $B(R_3)$; $Si(R_3)N(R_3')$; $P(O)(R_3)$; C_1 - C_8 of alkyl; C_2 - C_8 of heteroalkyl, alkylcycloalkyl, heterocycloalkyl; C_3 - C_8 of aryl, Ar-alkyl, heterocyclic, carbocyclic, cycloalkyl, heteroalkylcycloalkyl, alkylcarbonyl, heteroaryl; or 1~8 amino acids; Wherein R_3 and R_3' are independently H; C_1 - C_8 of alkyl; C_2 - C_8 of heteroalkyl, alkylcycloalkyl, heterocycloalkyl; C_3 - C_8 of aryl, Ar-alkyl, heterocyclic, carbocyclic, cycloalkyl, heteroalkylcycloalkyl, alkylcarbonyl, heteroaryl; or 1-8 carbon atoms of esters, ether, or amide; or polyethyleneoxy unit of formula $(OCH_2CH_2)_p$ or $(OCH_2CH(CH_3))_p$, wherein p is an integer from 0 to about 1000, or combination thereof. In addition, X_1 and X_2 can be independently absent.

R_2 is independently selected from OH, H, NH_2 , SH, $NHNH_2$, $N(R_3)(R_3')$; $N(R_3)NH(R_3')$; polyethyleneoxy unit of formula $(OCH_2CH_2)_pOR_3$, or $(OCH_2CH(CH_3))_pOR_3$, or $NH(CH_2CH_2O)_pR_3$, or $NH(CH_2CH(CH_3)O)_pR_3$, or $N[(CH_2CH_2O)_pR_3][(CH_2CH_2O)_pR_3']$, or $(OCH_2CH_2)_pCOOR_3$, or $CH_2CH_2(OCH_2CH_2)_pCOOR_3$, wherein p and p' are independently an


integer selected from 0 to about 1000, or combination thereof; C₁-C₈ of alkyl; C₂-C₈ of heteroalkyl, alkylcycloalkyl, heterocycloalkyl; C₃-C₈ of aryl, Ar-alkyl, heterocyclic, carbocyclic, cycloalkyl, heteroalkylcycloalkyl, alkylcarbonyl, heteroaryl; Wherein R₃ and R_{3'} are independently H; C₁-C₈ of alkyl; C₂-C₈ of heteroalkyl, alkylcycloalkyl, heterocycloalkyl; C₃-C₈ of aryl, Ar-alkyl, heterocyclic, carbocyclic, cycloalkyl, heteroalkylcycloalkyl, alkylcarbonyl, heteroaryl; or 1-8 carbon atoms of esters, ether, or amide; or 1~8 amino acids; or polyethyleneoxy unit of formula (OCH₂CH₂)_p or (OCH₂CH(CH₃))_p, wherein p is an integer from 0 to about 1000, or combination thereof.

m₁ and m₂ are independently an integer from 1 to 30, preferably from 1 to 10.


In another embodiment, R₁, R₃ and R_{3'}, can be respectively a chain of atoms selected from C, N, O, S, Si, and P which covalently connects the cell-surface binding molecule and/or the conjugated drug. The atoms used in forming the bridge linker may be combined in all chemically relevant ways, such as forming alkylene, alkenylene, and alkynylene, ethers, polyoxyalkylene, esters, amines, imines, polyamines, hydrazines, hydrazones, amides, ureas, semicarbazides, carbazides, alkoxyamines, alkoxylamines, urethanes, amino acids, acyloxylamines, hydroxamic acids, and many others. In addition, it is to be understood that the atoms forming the linker (L) may be either saturated or unsaturated, or may be radicals, or may be cyclized upon each other to form divalent cyclic structures, including cycloalkanes, cyclic ethers, cyclic amines, arylenes, heteroarylenes, and the like in the linker.

Examples of the functional group, Z₁, which enables linkage of a cytotoxic drug, include groups that enable linkage via a disulfide, thioether, thioester, peptide, hydrazone, ester, carbamate, carbonate, alkoxime or an amide bond. Such functional groups include, but are not limited to, thiol, disulfide, amino, carboxyl, aldehydes, ketone, maleimido, haloacetyl, hydrazines, alkoxyamino, and/or hydroxy.


Examples of the functional group, Z₁, that enables reaction with the terminal of amine of a drug/ cytotoxic agent can be, but not limited to, *N*-hydroxysuccinimide esters, *p*-nitrophenyl esters, dinitrophenyl esters, pentafluorophenyl esters, carboxylic acid chlorides or carboxylic acid anhydride; With the terminal of thiol can be, as but not limited to, pyridyldisulfides, nitropyridyldisulfides, maleimides, haloacetates, methylsulfonephenyloxadiazole (ODA), carboxylic acid chlorides and carboxylic acid anhydride; With the terminal of ketone or aldehyde can be, as but not limited to, amines, alkoxyamines, hydrazines, acyloxylamine, or hydrazide; With the terminal of azide can be, as but not limited to, alkyne. Examples of these function groups are displayed below:




N-hydroxysuccinimide ester;




maleimide;




disulfide;




haloacetyl;




acyl halide(acid halide),




ethenesulfonyl;




acryl (acryloyl);




(tosyloxy)acetyl;




2-(mesyloxy)acetyl;




(nitrophenyloxy)acetyl;




2-(dinitrophenyloxy)acetyl;




2-(fluorophenyl)-acetyl;




(difluorophenyl)-acetyl;




2-(((trifluoromethyl)sulfonyl)oxy)acetyl;




ketone, or aldehyde,




2-(pentafluorophenyl)acetyl;



, methylsulfonylphenyloxadiazole (ODA);



10 R_1 -C(=O)-O-C(=O)-X₂'- acid anhydride, H₂N-O- alkyloxyamino; N₃- azido,



alkynyl, or





hydrazide. Wherein X₁' is F, Cl, Br, I or Lv₃; X₂' is O, NH,

N(R₁), or CH₂; R₃ and R₅ are independently H, R₁, aromatic, heteroaromatic, or aromatic group wherein one or several H atoms are replaced independently by -R₁, -halogen, -OR₁, -SR₁, -NR₁R₂, -NO₂, -S(O)R₁, -S(O)₂R₁, or -COOR₁; Lv₃ is a leaving group selected from nitrophenol; N-hydroxysuccinimide (NHS); phenol; dinitrophenol; pentafluorophenol; tetrafluorophenol; difluorophenol; monofluorophenol; pentachlorophenol; triflate; imidazole; dichlorophenol; tetrachlorophenol; 1-hydroxybenzotriazole; tosylate; mesylate; 2-ethyl-5-phenylisoxazolium-3'-sulfonate, anhydrides formed its self, or formed with the other anhydride, e.g. acetyl anhydride,

formyl anhydride; or an intermediate molecule generated with a condensation reagent for peptide coupling reactions or for Mitsunobu reactions.

In preferred embodiments, R₁, R₂, R₃ and R_{3'}, are linear alkyl having from 1-6 carbon atoms, or polyethyleneoxy unit of formula (OCH₂CH₂)_p, p = 1~100.

The key step of synthesis of the bridge linker containing 2,3-disubstituted succinic group, or 2-monosubstituted or 2,3-disubstituted fumaric group, or 2-monosubstituted or 2,3-disubstituted maleic group, or acetylenedicarboxyl group, is the condensation of the 2,3-disubstituted succinic acid, or 2-monosubstituted or 2,3-disubstituted fumaric acid, or 2-monosubstituted or 2,3-disubstituted maleic acid, or acetylenedicarboxyl group, or their acid derivatives, with the other components containing an amine (1° or 2° amines), alcohol, or thiol on their terminal, as shown in the following scheme (Ia) and (Ib):



Wherein “====”, X₁, X₂, R₁, and R₂ are described the same in Formula (I).

Lv₁ and Lv₂ are the same or independently OH; F; Cl; Br; I; nitrophenol; N-hydroxysuccinimide (NHS); phenol; dinitrophenol; pentafluorophenol; tetrafluorophenol; difluorophenol; monofluorophenol; pentachlorophenol; triflate; imidazole; dichlorophenol; tetrachlorophenol; 1-hydroxybenzotriazole; tosylate; mesylate; 2-ethyl-5-phenylisoxazolium-3'-sulfonate, anhydrides formed its self, or formed with the other anhydride, e.g. acetyl anhydride, formyl anhydride; or an intermediate molecule generated with a condensation reagent for peptide coupling reactions, or for Mitsunobu reactions, e.g. condensation reagents are: EDC (N-(3-Dimethylaminopropyl)-N'-ethylcarbodiimide), DCC (Dicyclohexyl-carbodiimide), N,N'-Diisopropylcarbodiimide (DIC), N-Cyclohexyl-N'-(2-morpholino-ethyl)carbodiimide metho-p-toluenesulfonate (CMC, or CME-CDI), 1,1'-Carbonyldiimidazole (CDI), TBTU (O-(Benzotriazol-1-yl)-N,N,N',N'-tetramethyluronium tetrafluoroborate), N,N,N',N'-Tetramethyl-O-(1H-benzotriazol-1-yl)uronium hexafluorophosphate (HBTU), (Benzotriazol-1-yloxy)tris(dimethylamino)phosphonium hexafluorophosphate (BOP), (Benzotriazol-1-yloxy)tripyrrolidinophosphonium hexafluorophosphate (PyBOP), Diethyl

cyanophosphonate (DEPC), Chloro-N,N,N',N'-tetramethylformamidiniumhexafluoro-phosphate, 1-[Bis(dimethylamino)methylene]-1H-1,2,3-triazolo[4,5-b]pyridinium 3-oxid hexafluorophosphate (HATU), 1-[(Dimethylamino)(morpholino)methylene]-1H-[1,2,3]triazolo[4,5-b]pyridine-1-ium 3-oxide hexafluorophosphate (HDMA), 2-Chloro-1,3-dimethyl-imidazolidinium hexafluorophosphate (CIP), Chlorotripyrrolidinophosphonium hexafluorophosphate (PyCloP), Fluoro-N,N,N',N'-bis(tetramethylene)formamidinium hexafluorophosphate (BTFFH), N,N,N',N'-Tetramethyl-S-(1-oxido-2-pyridyl)thiuronium hexafluorophosphate, O-(2-Oxo-1(2H)pyridyl)-N,N,N',N'-tetramethyluronium tetrafluoroborate (TPTU), S-(1-Oxido-2-pyridyl)-N,N,N',N'-tetramethylthiuronium tetrafluoroborate, O-[(Ethoxycarbonyl)-cyanomethylenamino]-N,N,N',N'-tetramethyluronium hexafluorophosphate (HOTU), (1-Cyano-2-ethoxy-2-oxoethylidenaminooxy)dimethylamino-morpholino-carbenium hexafluorophosphate (COMU), O-(Benzotriazol-1-yl)-N,N,N',N'-bis(tetramethylene)uronium hexafluorophosphate (HBPYU), N-Benzyl-N'-cyclohexylcarbodiimide (with, or without polymer-bound), Dipyrrolidino(N-succinimidyl-oxy)carbenium hexafluoro-phosphate (HSPYU), Chlorodipyrrolidinocarbenium hexafluorophosphate (PYClU), 2-Chloro-1,3-dimethylimidazolidinium tetrafluoroborate(CIB), (Benzotriazol-1-yloxy)dipiperidino-carbenium hexafluorophosphate (HBPipU), O-(6-Chlorobenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium tetrafluoroborate (TCTU), Bromotris(dimethylamino)-phosphonium hexafluorophosphate (BroP), Propylphosphonic anhydride (PPACA, T3P[®]), 2-Morpholinoethyl isocyanide (MEI), N,N,N',N'-Tetramethyl-O-(N-succinimidyl)uronium hexafluorophosphate (HSTU), 2-Bromo-1-ethyl-pyridinium tetrafluoroborate (BEP), O-[(Ethoxycarbonyl)cyano-methylenamino]-N,N,N',N'-tetra-methyluronium tetrafluoroborate (TOTU), 4-(4,6-Dimethoxy-1,3,5-triazin-2-yl)-4-methylmorpholiniumchloride (MMTM, DMTMM), N,N,N',N'-Tetramethyl-O-(N-succinimidyl)uronium tetrafluoroborate (TSTU), O-(3,4-Dihydro-4-oxo-1,2,3-benzotriazin-3-yl)-N,N,N',N'-tetramethyluronium tetrafluoro-borate (TDBTU), 1,1'-(Azodicarbonyl)-dipiperidine (ADD), Di-(4-chlorobenzyl)azodicarboxylate (DCAD), Di-tert-butyl azodicarboxylate (DBAD), Diisopropyl azodicarboxylate (DIAD), Diethyl azodicarboxylate (DEAD). In addition, Lv1 and Lv2 can be an anhydride, formed by acid themselves or formed with other C₁~C₈ acid anhydrides.

The detail examples of the synthesis of the bridge linkers are shown in the figures 1~33. Normally the bridge substituents of 2,3-disubstituted succinic group, or 2-monosubstituted or 2,3-disubstituted fumaric group, or 2-monosubstituted or 2,3-disubstituted maleic group, or acetylenedicarboxyl group, can be condensed with linker components containing function groups capable to react to drugs of desired conjugation.

CELL-BINDING AGENT-DRUG CONJUGATES


The conjugates of the present invention can be represented by the following formula,




, wherein Cb is a cell-binding agent, L is linker containing succinic, fumaric or maleic group, or acetylenedicarboxyl group, Drug₁ is a drug molecule, n is an integer from 1 to 30, and two S (sulfur) elements from Cb bridgely link to L, which covalently connects a drug and a functional molecule R₂.

The bridge linker L may be composed of one or more linker components. Exemplary linker components include 6-maleimidocaproyl ("MC"), maleimidopropanoyl ("MP"), valine-citrulline ("val-cit" or "vc"), alanine-phenylalanine ("ala-phe" or "af"), p-aminobenzyloxycarbonyl ("PAB"), 4-thiopentanoate ("SPP"), 4-(N-maleimidomethyl)-cyclohexane-1 carboxylate ("MCC"), (4-acetyl)aminobenzoate ("SIAB"), 4-thio-butyrates (SPDB), 4-thio-2-hydroxysulfonyl-butyrates (2-Sulfo-SPDB), ethyleneoxy –CH₂CH₂O– as one or more repeating units ("EO" or "PEO"). Additional linker components are known in the art and some are described herein.


Example structures of these components containing linkers are:





; (MC, 6-maleimidocaproyl containing)




(MP, maleimidopropanoyl containing)




(PAB, p-aminobenzyloxycarbonyl containing)



(ME, maleimidoethyl containing).




(valine-citrulline containing)







5 (MCC, 4-(N-maleimidomethyl)cyclohexane-1 carboxylate containing)







((4-acetyl)aminobenzoate containing)







10 4-thio-buturate (SPDB), 4-(N-maleimidomethyl)cyclo-hexane-1-







dioxybenzylthio,  diaminobenzylthio,  amino-



oxybenzylthio,  alkoxy amino (AOA),  ethyleneoxy (EO),



dithio,  4-methyl-4-dithio-pentanoic (MPDP), 




triazole,  alkylsulfonyl,  alkylsulfonamide,  sulfon-


5 bisamide,  Phosphondiamide,  alkylphosphonamide,

 phosphinic acid,  N-methylphosphonamidic acid,


 N,N'-dimethylphosphonamidic acid,  N,N'-

dimethylphosphondiamide,  hydrazine,  acetimidamide,

 oxime,  acetylaceto-hydrazide, 

10 aminoethyl-amine,  aminoethyl-aminoethyl-amine.

Preferably, the conjugates have the following Formula (II):



Wherein:

Cb represents a cell-binding agent, preferably an antibody, which conjugates to a Drug, Drug₁ and a function molecule, R₂, via a pair of sulfur atoms (thiols).

Inside the bracket (square parentheses) are the linker-drug components that are conjugated to pairs of thiols of the cell-binding agent/molecule. The thiols are preferred pairs of sulfur atoms reduced from the inter chain disulfide bonds of the cell-binding agent by a reduction agent selected from dithiothreitol (DTT), dithioerythritol (DTE), L-glutathione (GSH), tris (2-carboxyethyl) phosphine (TCEP), 2-mercaptoethylamine (β-MEA), or/and beta mercaptoethanol (β-ME, 2-ME).

Drug₁ represents a cytotoxic agent, linked to a cell-binding agent via the bridge linker of this patent through an alkyl, alkylene, alkenylene, alkynylene, ether, polyoxyalkylene, ester, amine, imine, polyamine, hydrazine, hydrazone, amide, urea, semicarbazide, carbazide, alkoxyamine, urethanes, amino acid, peptide, acyloxylamine, hydroxamic acid, disulfide, thioether, thioester, carbamate, carbonate, heterocyclic ring, heteroalkyl, heteroaromatic, or alkoxime bond, or combination above thereof.

“~~---~~” represent either single bond or double bond.

n is 1 ~ 30; R₁, R₂, m₁, m₂, X₁ and X₂ are described the same previously in Formula (I).

As described in more detail below, Drug₁ can be any of many small molecule drugs, including, but not limited to, tubulysins, calicheamicins, auristatins, maytansinoids, CC-1065 analogs, morpholinos doxorubicins, taxanes, cryptophycins, amatoxins (amanitins), epothilones, geldanamycins, duocarmycins, daunomycins, methotrexates, vindesines, vincristines, and benzodiazepine dimers (e.g., dimmers of pyrrolbenzodiazepine (PBD) or tomaymycin), indolinobenzodiazepines, imidazobenzothiadiazepines, or oxazolidinobenzodiazepines).

To synthesize the conjugate, the cell-binding agent can be first modified with the bridge linkers of the present invention through reduction of disulfide bonds of the cell-binding molecule. The yielded a pair of free thiols can react to the bridge linker of Formula (I) at pH 5~9 aqueous media with or without addition of 0~30% of water mixable (miscible) organic solvents, such as DMA, DMF, ethanol, methanol, acetone, acetonitrile, THF, isopropanol, dioxane, propylene glycol, or ethylene diol, to introduce the reactive group of Z₁ containing disulfide, maleimido, haloacetyl, azide, 1-yne, ketone, aldehyde, alkoxyamino, or hydrazide groups, as well a function molecule group R₂, which is preferably a ployethelene glycol, carboxylic acid, or carboxylic amide, or ester, or other derivatives. Then a reactive group of a cytotoxic agent reacts to the modified cell-binding molecule accordingly. For example, synthesis of the cell-binding agent-drug conjugates linked via disulfide bonds is achieved by a disulfide exchange between the disulfide bond in the modified cell-binding agent and a drug containing a free thiol group.


Synthesis of the cell-binding agent-drug conjugates linked via thioether is achieved by reaction of the maleimido or haloacetyl or ethylsulfonyl modified cell-binding agent and a drug containing a free thiol group. Synthesis of conjugates bearing an acid labile hydrazone can be achieved by reaction of a carbonyl group with the hydrazide moiety in the linker, by methods known in the art (see, for example, P. Hamann et al., *Cancer Res.* 53, 3336-34, 1993; B. Laguzza et al., *J. Med. Chem.*, 32; 548-55, 1959; P. Trail et al., *Cancer Res.*, 57; 100-5, 1997). Synthesis of conjugates bearing triazole linkage can be achieved by reaction of a 1-yne group of the drug with the azido moiety in the linker, through the click chemistry (Huisgen cycloaddition) (Lutz, J-F. et al, 2008, *Adv. Drug Del. Rev.* 60, 958–70; Sletten, E. M. et al 2011, *AccChem. Research* 44, 666–76).

Alternatively, the drug can react with the bridge linkers of the present invention that have conjugated to a cell-binding molecule to give a modified cell-binding molecule linker of Formula (III) bearing functionalities. For example, a thiol-containing drug can react with the modified cell-binding molecule bridge linker of Formula (III) bearing a maleimido, or a haloacetyl, or an ethylsulfonyl substituent at pH 5.5~9.0 in aqueous buffer to give a cell-binding molecule-drug conjugate via a thioether linkage. A thiol-containing drug can undergo disulfide exchange with a modified bridge linker of Formula (III) bearing a pyridyldithio moiety to give a conjugate a disulfide bond linkage. A drug bearing a hydroxyl group or a thiol group can be reacted with a modified bridge linker of Formula (III) bearing a halogen, particularly the alpha halide of carboxylates, in the presence of a mild base, e.g. pH 8.0~9.5, to give a modified drug bearing an ether or thiol ether link. A hydroxyl group containing drug can be condensed with a bridge cross linker of Formula (I) bearing a carboxyl group, in the presence of a dehydrating agent, such as EDC or DCC, to give ester linkage, then the subject drug modified bridge linker undergoes the conjugation with a cell-binding molecule. A drug containing an amino group can condensate with a carboxyl ester of NHS, imidazole, nitrophenol; N-hydroxysuccinimide (NHS); phenol; dinitrophenol; pentafluorophenol; tetrafluorophenol; difluorophenol; monofluorophenol; pentachlorophenol; triflate; imidazole; dichlorophenol;tetrachlorophenol;1-hydroxyben-zotriazole; tosylate; mesylate; 2-ethyl-5-phenylisoxazolium-3'-sulfonate on the cell-binding molecule-bridge linker of Formula (III) to give a conjugate via amide bond linkage.

The conjugate may be purified by standard biochemical means, such as gel filtration on a Sephadex G25 or Sephacryl S300 column, adsorption chromatography, and ion exchange or by dialysis. In some cases, a small molecule as a cell-binding agent (e.g. folic acid, melanocyte stimulating hormone, EGF etc) conjugated with a small molecular drugs can be purified by chromatography such as by HPLC, medium pressure column chromatography or ion exchange chromatography.

MODIFIED CELL-BINDING AGENTS/MOLECULES

The cell-binding agent modified by reaction with linkers of the present invention is preferably represented by the Formula (III):



Wherein --- represents either a single bond, or a double bond.

Cb, Z₁, Z₂, n, R₁, R₂, m₁, m₂, X₁, and X₂ are defined the same as in Formula (I) and (II).

In preferred embodiments, Z₁ is a disulfide substituent, maleimido, haloacetyl, alkoxyamine, azido, ketone, aldehyde, hydrazine, alkyne, an *N*-hydroxysuccinimide ester, or a carboxyl ester formed with phenol; dinitrophenol; pentafluorophenol; tetrafluoro-phenol; difluorophenol; monofluorophenol; pentachlorophenol; triflate; imidazole; dichlorophenol; tetrachlorophenol; 1-hydroxybenzotriazole; tosylate; mesylate; 2-ethyl-5-phenylisoxa-zolium-3'-sulfonate. Z₁ can then react with a cytotoxic agent through disulfide, thioether, hydrazone, amide, alkoxime, carbamate, ester, ether bond or hetero-aromatic ring. The modified cell-binding agent can be prepared via a reaction of the cell-binding agent with the bridge linkers of Formula (I) as described in Formula (II) above.

In order to achieve a higher conjugation reaction yield of 2,3-disubstituted succinic group, or 2-monosubstituted or 2,3-disubstituted fumaric group, or 2-monosubstituted or 2,3-disubstituted maleic group, or acetylenedicarboxyl group, on the bridge linkers of the Formula (I) with a pair of free thiols on the cell-binding molecule, preferably on an antibody, a small percentage of organic co-solvent may be required to add to the reaction mixture, as well in the solution after the reaction to maintain solubility of the Formula (III) in aqueous solution. To modify the cell-binding agents, the cross-linking reagent (bridge linker) of Formula (I) can be first dissolved in a polar organic solvent that is miscible with water, for example different alcohols, such as methanol, ethanol, and propanol, acetone, acetonitrile, tetrahydrofuran (THF), 1,4-dioxane, dimethyl formamide (DMF), dimethyl acetamide (DMA), or dimethylsulfoxide (DMSO) at a high concentration, for example 1-500 mM. Meanwhile, the cell-binding molecule, such as antibody dissolved in an aqueous buffer pH 5~9.5, preferably pH 6~8.5, at 1~35 mg/ml concentration was treated with 1~20 equivalent of TCEP or DTT for 20 min to 48 hour. After the reduction, DTT can be removed by SEC chromatographic purification. TCEP can be optionally removed by SEC chromatography too, or staying in the reaction mixture for the next step reaction without purification. Furthermore, the reduction of antibodies or the other cell-binding agents


with TCEP can be performed with a bridge linker of Formula (I), for which the cross-linking conjugation for the cell-binding molecules can be achieved simultaneously along with the TCEP reduction.

The aqueous solutions for the modification of cell-binding agents are buffered between pH 6 and 9, preferably between 6.5 and 7.5 and can contain any non-nucleophilic buffer salts useful for these pH ranges. Typical buffers include phosphate, triethanolamine HCl, HEPES, and MOPS buffers, which can contain additional components, such as cyclodextrins, sucrose and salts, for examples, NaCl and KCl. After the addition of the bridge linker of Formula (I) into the solution containing the reduced cell-binding molecules, the reaction mixture is incubated at a temperature of from 4 °C to 45°C, preferably at ambient temperature. The progress of the reaction can be monitored by measuring the decrease in the absorption at a certain UV wavelength, such as at 254 nm, or increase in the absorption at a certain UV wavelength, such as 280 nm, or the other appropriate wavelength. After the reaction is complete, isolation of the modified cell-binding agent can be performed in a routine way, using for example gel filtration chromatography, or adsorptive chromatography.

The extent of modification can be assessed by measuring the absorbance of the nitropyridine thione, dinitropyridine dithione, pyridine thione, carboxylamidopyridine dithione and dicarboxylamidopyridine dithione group released via UV spectra. For the conjugation without a chromophore group, the modification or conjugation reaction can be monitored by LC-MS, preferably by UPLC-QTOF mass spectrometry, or Capillary electrophoresis-mass spectrometry (CE-MS). The bridge cross-linkers described herein have diverse functional groups that can react with any drugs, preferably cytotoxic agents that possess a suitable substituent. For examples, the modified cell-binding molecules bearing an amino or hydroxyl substituent can react with drugs bearing an N-hydroxysuccinimide (NHS) ester, the modified cell-binding molecules bearing a thiol substituent can react with drugs bearing a maleimido or haloacetyl group. Additionally, the modified cell-binding molecules bearing a carbonyl (ketone or aldehyde) substituent can react with drugs bearing a hydrazide or an alkoxyamine. One skilled in the art can readily determine which linker to use based on the known reactivity of the available functional group on the linkers.

MODIFIED CYTOTOXIC DRUGS

The cytotoxic drugs modified by reaction with cross-linkers of the present invention are preferably represented by the Formula (IV):



Wherein “---” represents either a single bond, or a double bond, or a triple bond.

“—”, U, U', Drug₁, Drug₂, R₁, R₂, m₁, m₂, X₁, and X₂ are defined the same as in Formula (I) and (II).

The modified drugs can be prepared via reaction of the drug with the linkers of the Formula (I) to give a modified drug of Formula (IV) bearing functionality of an 2,3-disubstituted succinic group, or 2-monosubstituted or 2,3-disubstituted fumaric group, or 2-monosubstituted or 2,3-disubstituted maleic group, or acetylenedicarboxyl group. But for drugs containing a thiol, or the drugs undergoing to conjugation of a cell-binding molecule via the bridge linkers through thioether, thioester or disulfide bond, it is therefore preferred that the Drug₁ may be synthesized to connect to R₁ in a piece of components via the linkage of thioether, thioester or disulfide bond first. Then the synthesized R₁-Drug₁ component is assembled to 2,3-disubstituted succinic acid, or 2-monosubstituted or 2,3-disubstituted fumaric acid, or 2-monosubstituted or 2,3-disubstituted maleic acid, or acetylenedicarboxyl group, to form the bridge linker modified drugs of Formula (IV).

For examples of the synthesis, a thiol-containing drug can be reacted with the linker of components R₁ bearing a maleimido substituent at neutral pH in aqueous buffer to give a R₁-Drug₁ compartment bearing thioether linkage, and following by condensation with either 2,3-disubstituted succinic acid, or 2-monosubstituted or 2,3-disubstituted fumaric acid, or 2-monosubstituted or 2,3-disubstituted maleic acid, or acetylenedicarboxyl group, to give a modified drug of Formula (IV) bearing thioether linkage. A drug bearing a hydroxyl group can be reacted with a linker component R₁ bearing a halogen, or a tosylate, or a mesylate, in the presence of a mild base, to give a R₁-Drug₁ compartment bearing ether linkage, and following by condensation with 2,3-disubstituted succinic acid, or 2-monosubstituted or 2,3-disubstituted fumaric acid, or 2-monosubstituted or 2,3-disubstituted maleic acid, or acetylenedicarboxyl group, to give a modified drug of Formula (IV) bearing thioether linkage. A hydroxyl group containing drug can be condensed with a linker of Formula (I) bearing a carboxyl group, in the presence of a dehydrating agent, such as EDC or dicyclohexylcarbodiimide (DCC), to give a modified drug of Formula (IV) via ester linkage. A drug bearing a thiol group can also react the linker of components R₁ bearing a maleimido or a vinylsulfonyl, or a haloacetyl group, give a R₁-Drug₁ compartment bearing thioether linkage, and following by condensation with a compartment of 2,3-disubstituted succinic acid, or 2-monosubstituted or 2,3-disubstituted fumaric acid, or 2-monosubstituted or 2,3-

disubstitutedmaleic acid, or acetylenedicarboxyl group, to give a modified drug of Formula (IV) bearing thioether linkage. An amino group containing drug can similarly undergo condensation with a carboxyl group on the bridge linker of Formula (I) to give a modified drug of Formula (IV) bearing amide bonds. The modified drug can be purified by standard methods such as column chromatography over silica gel or alumina, crystallization, preparatory thin layer chromatography, ion exchange chromatography, or HPLC.

CELL-BINDING AGENTS

The cell-binding molecule, Cb, that comprises the conjugates and the modified cell-binding agents of the present invention may be of any kind presently known, or that become known, molecule that binds to, complexes with, or reacts with a moiety of a cell population sought to be therapeutically or otherwise biologically modified.

The cell binding agents include, but are not limited to, large molecular weight proteins such as, for example, full-length antibodies (polyclonal antibodies, monoclonal antibodies, dimers, multimers, multispecific antibodies (e.g., bispecific antibodies); single chain antibodies; fragments of antibodies such as Fab, Fab', F(ab')₂, F_v, [Parham, J. Immunol. 131, 2895-902 (1983)], fragments produced by a Fab expression library, anti-idiotypic (anti-Id) antibodies, CDR's, diabody, triabody, tetrabody, miniantibody, small immune proteins (SIP), and epitope-binding fragments of any of the above which immuno-specifically bind to cancer cell antigens, viral antigens, microbial antigens or a protein generated by the immune system that is capable of recognizing, binding to a specific antigen or exhibiting the desired biological activity (Miller et al (2003) J. of Immunology 170: 4854-61); interferons (such as type I, II, III); peptides; lymphokines such as IL-2, IL-3, IL-4, IL-5, IL-6, IL-10, GM-CSF, interferon-gamma (IFN- γ); hormones such as insulin, TRH (thyrotropin releasing hormones), MSH (melanocyte-stimulating hormone), steroid hormones, such as androgens and estrogens, melanocyte-stimulating hormone (MSH); growth factors and colony-stimulating factors such as epidermal growth factors (EGF), granulocyte-macrophage colony-stimulating factor (GM-CSF), transforming growth factors (TGF), such as TGF α , TGF β , insulin and insulin like growth factors (IGF-I, IGF-II) G-CSF, M-CSF and GM-CSF [Burgess, Immunology Today, 5, 155-8 (1984)]; vaccinia growth factors (VGF); fibroblast growth factors (FGFs); smaller molecular weight proteins, poly-peptide, peptides and peptide hormones, such as bombesin, gastrin, gastrin-releasing peptide; platelet-derived growth factors; interleukin and cytokines, such as interleukin-2 (IL-2), interleukin-6 (IL-6), leukemia inhibitory factors, granulocyte-macrophage colony-stimulating factor (GM-CSF); vitamins, such as folate; apoproteins and glycoproteins, such as transferrin [O'Keefe et al, 260 J.

Biol. Chem. 932-7 (1985)]; sugar-binding proteins or lipoproteins, such as lectins; cell nutrient-transport molecules; and small molecular inhibitors, such as prostate-specific membrane antigen (PSMA) inhibitors and small molecular tyrosine kinase inhibitors (TKI), non-peptides or any other cell binding molecule or substance, such as bioactive polymers (Dhar, et al, Proc. Natl. Acad. Sci. 2008, 105, 17356-61); bioactive dendrimers (Lee, et al, Nat. Biotechnol. 2005, 23, 1517-26; Almutairi, et al; Proc. Natl. Acad. Sci. 2009, 106, 685-90); nanoparticles (Liong, et al, ACS Nano, 2008, 2, 1309-12; Medarova, et al, Nat. Med. 2007, 13, 372-7; Javier, et al, Bioconjugate Chem. 2008, 19, 1309-12); liposomes (Medinai, et al, Curr. Phar. Des. 2004, 10, 2981-9); viral capsides (Flenniken, et al, Viruses Nanotechnol. 2009, 327, 71-93).

In general, a monoclonal antibody is preferred as a cell-surface binding agent if an appropriate one is available. And the antibody may be murine, human, humanized, chimeric, or derived from other species.

Production of antibodies used in the present invention involves in vivo or in vitro procedures or combinations thereof. Methods for producing polyclonal anti-receptor peptide antibodies are well-known in the art, such as in U.S. Pat. No. 4,493,795 (to Nestor et al). A monoclonal antibody is typically made by fusing myeloma cells with the spleen cells from a mouse that has been immunized with the desired antigen (Köhler, G.; Milstein, C. (1975). Nature 256: 495-7). The detailed procedures are described in “Antibodies--A Laboratory Manual”, Harlow and Lane, eds., Cold Spring Harbor Laboratory Press, New York (1988), which is incorporated herein by reference. Particularly monoclonal antibodies are produced by immunizing mice, rats, hamsters or any other mammal with the antigen of interest such as the intact target cell, antigens isolated from the target cell, whole virus, attenuated whole virus, and viral proteins. Splenocytes are typically fused with myeloma cells using polyethylene glycol (PEG) 6000. Fused hybrids are selected by their sensitivity to HAT (hypoxanthine-aminopterin-thymine). Hybridomas producing a monoclonal antibody useful in practicing this invention are identified by their ability to immunoreact specified receptors or inhibit receptor activity on target cells.

A monoclonal antibody used in the present invention can be produced by initiating a monoclonal hybridoma culture comprising a nutrient medium containing a hybridoma that secretes antibody molecules of the appropriate antigen specificity. The culture is maintained under conditions and for a time period sufficient for the hybridoma to secrete the antibody molecules into the medium. The antibody-containing medium is then collected. The antibody molecules can then be further isolated by well-known techniques, such as using protein-A affinity chromatography; anion, cation, hydrophobic, or size exclusive chromatographies (particularly by affinity for the specific antigen after protein A, and sizing column chromatography);

centrifugation, differential solubility, or by any other standard technique for the purification of proteins.

Media useful for the preparation of these compositions are both well-known in the art and commercially available and include synthetic culture media. An exemplary synthetic medium is Dulbecco's minimal essential medium (DMEM; Dulbecco et al., *Viol.* 8, 396 (1959)) supplemented with 4.5 gm/l glucose, 0~20 mM glutamine, 0~20% fetal calf serum, several ppm amount of heavy metals, such as Cu, Mn, Fe, or Zn, etc, or/and the other heavy metals added in their salt forms, and with an anti-foaming agent, such as polyoxyethylene-polyoxypropylene block copolymer.

In addition, antibody-producing cell lines can also be created by techniques other than fusion, such as direct transformation of B lymphocytes with oncogenic DNA, or transfection with an oncovirus, such as Epstein-Barr virus (EBV, also called human herpesvirus 4 (HHV-4)) or Kaposi's sarcoma-associated herpesvirus (KSHV). See, U.S. Pat. Nos. 4,341,761; 4,399,121; 4,427,783; 4,444,887; 4,451,570; 4,466,917; 4,472,500; 4,491,632; 4,493,890. A monoclonal antibody may also be produced via an anti-receptor peptide or peptides containing the carboxyl terminal as described well-known in the art. See Niman et al., *Proc. Natl. Acad. Sci. USA*, 80: 4949-53 (1983); Geysen et al., *Proc. Natl. Acad. Sci. USA*, 82: 178-82 (1985); Lei et al. *Biochemistry* 34(20): 6675-88, (1995). Typically, the anti-receptor peptide or a peptide analog is used either alone or conjugated to an immunogenic carrier, as the immunogen for producing anti-receptor peptide monoclonal antibodies.

There are also a number of other well-known techniques for making monoclonal antibodies as binding molecules in this invention. Particularly useful are methods of making fully human antibodies. One method is phage display technology which can be used to select a range of human antibodies binding specifically to the antigen using methods of affinity enrichment. Phage display has been thoroughly described in the literature and the construction and screening of phage display libraries are well known in the art, see, e.g., Dente et al, *Gene*. 148(1):7-13 (1994); Little et al, *Biotechnol Adv.* 12(3): 539-55 (1994); Clackson et al., *Nature* 352: 264-8 (1991); Huse et al., *Science* 246: 1275-81 (1989).

Monoclonal antibodies derived by hybridoma technique from another species than human, such as mouse, can be humanized to avoid human anti-mouse antibodies when infused into humans. Among the more common methods of humanization of antibodies are complementarity-determining region grafting and resurfacing. These methods have been extensively described, see e.g. U.S. Pat. Nos. 5,859,205 and 6,797,492; Liu et al, *Immunol Rev.* 222: 9-27 (2008); Almagro et al, *Front Biosci.* 13: 1619-33 (2008); Lazar et al, *Mol Immunol.* 44(8): 1986-98 (2007); Li et al,

Proc. Natl. Acad. Sci. U S A. 103(10): 3557-62 (2006) each incorporated herein by reference. Fully human antibodies can also be prepared by immunizing transgenic mice, rabbits, monkeys, or other mammals, carrying large portions of the human immunoglobulin heavy and light chains, with an immunogen. Examples of such mice are: the Xenomouse. (Abgenix/Amgen), the
5 HuMAb-Mouse (Medarex/BMS), the VelociMouse (Regeneron), see also U.S. Pat. Nos. 6,596,541, 6,207,418, 6,150,584, 6,111,166, 6,075,181, 5,922,545, 5,661,016, 5,545,806, 5,436,149 and 5,569,825. In human therapy, murine variable regions and human constant regions can also be fused to construct called "chimeric antibodies" that are considerably less immunogenic in man than murine mAbs (Kipriyanov et al, Mol Biotechnol. 26: 39-60 (2004);
10 Houdebine, Curr Opin Biotechnol. 13: 625-9 (2002) each incorporated herein by reference). In addition, site-directed mutagenesis in the variable region of an antibody can result in an antibody with higher affinity and specificity for its antigen (Brannigan et al, Nat Rev Mol Cell Biol. 3: 964-70, (2002)); Adams et al, J Immunol Methods. 231: 249-60 (1999)) and exchanging constant regions of a mAb can improve its ability to mediate effector functions of binding and cytotoxicity.

15 Antibodies immunospecific for a malignant cell antigen can also be obtained commercially or produced by any method known to one of skill in the art such as, e.g., chemical synthesis or recombinant expression techniques. The nucleotide sequence encoding antibodies immune-specific for a malignant cell antigen can be obtained commercially, e.g., from the GenBank database or a database like it, the literature publications, or by routine cloning and sequencing.

20 Apart from an antibody, a peptide or protein that bind/block/target or in some other way interact with the epitopes or corresponding receptors on a targeted cell can be used as a binding molecule. These peptides or proteins could be any random peptide or proteins that have an affinity for the epitopes or corresponding receptors and they don't necessarily have to be of the immunoglobulin family. These peptides can be isolated by similar techniques as for phage display
25 antibodies (Szardenings, J Recept Signal Transduct Res. 2003, 23(4): 307-49). The use of peptides from such random peptide libraries can be similar to antibodies and antibody fragments. The binding molecules of peptides or proteins may be conjugated on or linked to a large molecules or materials, such as, but is not limited, an albumin, a polymer, a liposome, a nano particle, a dendrimer, as long as such attachment permits the peptide or protein to retain its
30 antigen binding specificity.

Examples of antibodies used for conjugation of drugs via the bridge linkers of this prevention for treating cancer, autoimmune disease, and/or infectious disease include, but are not limited to, 3F8 (anti-GD2), Abagovomab (anti CA-125), Abciximab (anti CD41 (integrin alpha-IIb), Adalimumab (anti-TNF- α), Adecatumumab (anti-EpCAM, CD326), Afelimomab (anti-TNF- α);

Afutuzumab (anti-CD20), Alacizumab pegol (anti-VEGFR2), ALD518 (anti-IL-6), Alemtuzumab (Campath, MabCampath, anti-CD52), Altumomab (anti-CEA), Anatumomab (anti-TAG-72), Anrukinzumab (IMA-638, anti-IL-13), Apolizumab (anti-HLA-DR), Arcitumomab (anti-CEA), Aselizumab (anti-L-selectin (CD62L), Atlizumab (tocilizumab, Actemra, RoActemra, anti-IL-6 receptor), Atorolimumab (anti-Rhesus factor), Bapineuzumab (anti-beta amyloid), Basiliximab (Simulect, antiCD25 (α chain of IL-2 receptor), Bavituximab (anti-phosphatidylserine), Bectumomab (LymphoScan, anti-CD22), Belimumab (Benlysta, LymphoStat-B, anti-BAFF), Benralizumab (anti-CD125), Bertilimumab (anti-CCL11 (eotaxin-1)), Besilesomab (Scintimun, anti-CEA-related antigen), Bevacizumab (Avastin, anti-VEGF-A), Biciromab (FibriScint, anti-fibrin II beta chain), Bivatuzumab (anti-CD44 v6), Blinatumomab (BiTE, anti-CD19), Brentuximab (cAC10, anti-CD30 TNFRSF8), Briakinumab (anti-IL-12, IL-23) Canakinumab (Ilaris, anti-IL-1), Cantuzumab (C242, anti-CanAg), Capromab, Catumaxomab (Removab, anti-EpCAM, anti-CD3), CC49 (anti-TAG-72), Cedelizumab (anti-CD4), Certolizumab pegol (Cimzia anti-TNF- α), Cetuximab (Erbix, IMC-C225, anti-EGFR), Citatuzumab bogatox (anti-EpCAM), Cixutumumab (anti-IGF-1), Clenoliximab (anti-CD4), Clivatuzumab (anti-MUC1), Conatumumab (anti-TRAIL-R2), CR6261 (anti-Influenza A hemagglutinin), Dacetuzumab (anti-CD40), Daclizumab (Zenapax, anti-CD25 (α chain of IL-2 receptor)), Daratumumab (anti-CD38 (cyclic ADP ribose hydrolase), Denosumab (Prolia, anti-RANKL), Detumomab (anti-B-lymphoma cell), Dorlimomab, Dorlixizumab, Echromeximab (anti-GD3 ganglioside), Eculizumab (Soliris, anti-C5), Edobacomab (anti-endotoxin), Edrecolomab (Panorex, MAb17-1A, anti-EpCAM), Efalizumab (Raptiva, anti-LFA-1 (CD11a), Efungumab (Mycograb, anti-Hsp90), Elotuzumab (anti-SLAMF7), Elsilimomab (anti-IL-6), Enlimomab pegol (anti-ICAM-1 (CD54)), Epitumomab (anti-episialin), Epratuzumab (anti-CD22), Erlizumab (anti-ITGB2 (CD18)), Ertumaxomab (Rexomun, anti-HER2/neu, CD3), Etaracizumab (Abegrin, anti-integrin $\alpha_v\beta_3$), Exbivirumab (anti-hepatitis B surface antigen), Fanolesomab (NeutroSpec, anti-CD15), Faralimomab (anti-interferon receptor), Farletuzumab (anti-folate receptor 1), Felvizumab (anti-respiratory syncytial virus), Fezakinumab (anti-IL-22), Figitumumab (anti-IGF-1 receptor), Fontolizumab (anti-IFN- γ), Foravirumab (anti-rabies virus glycoprotein), Fresolimumab (anti-TGF- β), Galiximab (anti-CD80), Gantenerumab (anti-beta amyloid), Gavilimomab (anti-CD147 (basigin)), Gemtuzumab (anti-CD33), Girentuximab (anti-carbonic anhydrase 9), Glembatumumab (CR011, anti-GPNMB), Golimumab (Simponi, anti-TNF- α), Gomiliximab (anti-CD23 (IgE receptor)), Ibalizumab (anti-CD4), Ibritumomab (anti-CD20), Igovomab (Indimacis-125, anti-CA-125), Imciromab (Myoscint, anti-cardiac myosin), Infliximab (Remicade, anti-TNF- α), Intetumumab (anti-CD51), Inolimomab (anti-CD25 (α chain of IL-2

receptor)), Inotuzumab (anti-CD22), Ipilimumab (anti-CD152), Iratumumab (anti- CD30 (TNFRSF8)), Keliximab (anti-CD4), Labetuzumab (CEA-Cide, anti-CEA), Lebrikizumab (anti-IL-13), Lemalesomab (anti-NCA-90 (granulocyte antigen)), Lerdelimumab (anti-TGF beta 2), Lexatumumab (anti-TRAIL-R2), Libivirumab (anti-hepatitis B surface antigen), Lintuzumab (anti-CD33), Lucatumumab (anti-CD40), Lumiliximab (anti- CD23 (IgE receptor), Mapatumumab (anti-TRAIL-R1), Maslimomab (anti- T-cell receptor), Matuzumab (anti-EGFR), Mepolizumab (Bosatria, anti-IL-5), Metelimumab (anti-TGF beta 1), Milatuzumab (anti-CD74), Minretumomab (anti-TAG-72), Mitumomab (BEC-2, anti-GD3 ganglioside), Morolimumab (anti-Rhesus factor), Motavizumab (Numax, anti-respiratory syncytial virus), Muromonab-CD3 (Orthoclone OKT3, anti-CD3), Nacolomab (anti-C242), Naptumomab (anti-5T4), Natalizumab (Tysabri, anti-integrin α_4), Nebacumab (anti-endotoxin), Necitumumab (anti-EGFR), Nerelimomab (anti-TNF- α), Nimotuzumab (Theracim, Theraloc, anti-EGFR), Nofetumomab, Ocrelizumab (anti-CD20), Odulimomab (Afolimomab, anti-LFA-1 (CD11a)), Ofatumumab (Arzerra, anti-CD20), Olaratumab (anti-PDGF-R α), Omalizumab (Xolair, anti-IgE Fc region), Oportuzumab (anti-EpCAM), Oregovomab (OvaRex, anti-CA-125), Otelixizumab (anti-CD3), Pagibaximab (anti-lipoteichoic acid), Palivizumab (Synagis, Abbosynagis, anti-respiratory syncytial virus), Panitumumab (Vectibix, ABX-EGF, anti-EGFR), Panobacumab (anti-Pseudomonas aeruginosa), Pascolizumab (anti-IL-4), Pemtumomab (Theragyn, anti-MUC1), Pertuzumab (Omnitarg, 2C4, anti-HER2/neu), Pexelizumab (anti-C5), Pintumomab (anti-adenocarcinoma antigen), Priliximab (anti-CD4), Pritumumab (anti-vimentin), PRO 140 (anti-CCR5), Racotumomab (1E10, anti-(N-glycolylneuraminic acid (NeuGc, NGNA)-gangliosides GM3)), Rafivirumab (anti-rabies virus glycoprotein), Ramucirumab (anti-VEGFR2), Ranibizumab (Lucentis, anti-VEGF-A), Raxibacumab (anti-anthrax toxin, protective antigen), Regavirumab (anti-cytomegalovirus glycoprotein B), Reslizumab (anti-IL-5), Rilotumumab (anti-HGF), Rituximab (MabThera, Rituxanmab, anti-CD20), Robatumumab (anti-IGF-1 receptor), Rontalizumab (anti-IFN- α), Rovelizumab (LeukArrest, anti-CD11, CD18), Ruplizumab (Antova, anti-CD154 (CD40L)), Satumomab (anti-TAG-72), Sevirumab (anti-cytomegalovirus), Sibrotuzumab (anti-FAP), Sifalimumab (anti-IFN- α), Siltuximab (anti-IL-6), Siplizumab (anti-CD2), (Smart) MI95 (anti-CD33), Solanezumab (anti-beta amyloid), Sonepcizumab (anti-sphingosine-1-phosphate), Sontuzumab (anti-episialin), Stamulumab (anti-myostatin), Sulesomab (LeukoScan, (anti-NCA-90 (granulocyte antigen), Tacatuzumab (anti-alpha-fetoprotein), Tadocizumab (anti-integrin $\alpha_{Ib\beta_3}$), Talizumab (anti-IgE), Tanezumab (anti-NGF), Taplitumomab (anti-CD19), Tefibazumab (Aurexis, (anti-clumping factor A), Telimomab, Tenatumomab (anti-tenascin C), Teneliximab (anti-CD40), Teplizumab (anti-CD3), TGN1412 (anti-CD28),

Ticilimumab (Tremelimumab, (anti-CTLA-4), Tigatuzumab (anti-TRAIL-R2), TNX-650 (anti-IL-13), Tocilizumab (Atlizumab, Actemra, RoActemra, (anti-IL-6 receptor), Toralizumab (anti-CD154 (CD40L)), Tositumomab (anti-CD20), Trastuzumab (Herceptin, (anti-HER2/neu), Tremelimumab (anti-CTLA-4), Tucotuzumab celmoleukin (anti-EpCAM), Tuvirumab (anti-hepatitis B virus), Urtoxazumab (anti- Escherichia coli), Ustekinumab (Stelara, anti-IL-12, IL-23), Vapaliximab (anti-AOC3 (VAP-1)), Vedolizumab, (anti-integrin $\alpha_4\beta_7$), Veltuzumab (anti-CD20), Vepalimomab (anti-AOC3 (VAP-1), Visilizumab (Nuvion, anti-CD3), Vitaxin (anti-vascular integrin avb3), Volociximab (anti-integrin $\alpha_5\beta_1$), Votumumab (HumaSPECT, anti-tumor antigen CTAA16.88), Zalutumumab (HuMax-EGFr, (anti-EGFR), Zanolimumab (HuMax-CD4, anti-CD4), Ziralimumab (anti-CD147 (basigin)), Zolimomab (anti-CD5), Etanercept (Enbrel®), Alefacept (Amevive®), Abatacept (Orencia®), Rilonacept (Arcalyst), 14F7 [anti-IRP-2 (Iron Regulatory Protein 2)], 14G2a (anti-GD2 ganglioside, from Nat. Cancer Inst. for melanoma and solid tumors), J591 (anti-PSMA, Weill Cornell Medical School for prostate cancers), 225.28S [anti-HMW-MAA (High molecular weight-melanoma-associated antigen), Sorin Radiofarmaci S.R.L. (Milan, Italy) for melanoma], COL-1 (anti-CEACAM3, CGM1, from Nat. Cancer Inst. USA for colorectal and gastric cancers), CYT-356 (Oncoltad®, for prostate cancers), HNK20 (OraVax Inc. for respiratory syncytial virus), ImmuRAIT (from Immunomedics for NHL), Lym-1 (anti-HLA-DR10, Peregrine Pharm. for Cancers), MAK-195F [anti-TNF (tumor necrosis factor; TNFA, TNF-alpha; TNFSF2), from Abbott / Knoll for Sepsis toxic shock], MEDI-500 [T10B9, anti-CD3, TR $\alpha\beta$ (T cell receptor alpha/beta), complex, from MedImmune Inc for Graft-versus-host disease], RING SCAN [anti-TAG 72 (tumour associated glycoprotein 72), from Neoprobe Corp. for Breast, Colon and Rectal cancers], Avicidin (anti-EPCAM (epithelial cell adhesion molecule), anti-TACSTD1 (Tumor-associated calcium signal transducer 1), anti-GA733-2 (gastrointestinal tumor-associated protein 2), anti-EGP-2 (epithelial glycoprotein 2); anti-KSA; KS1/4 antigen; M4S; tumor antigen 17-1A; CD326, from NeoRx Corp. for Colon, Ovarian, Prostate cancers and NHL]; LymphoCide (Immunomedics, NJ), Smart ID10 (Protein Design Labs), Oncolym (Techniclone Inc, CA), Allomune (BioTransplant, CA), anti-VEGF (Genentech, CA); CEAcide (Immunomedics, NJ), IMC-1C11 (ImClone, NJ) and Cetuximab (ImClone, NJ) .

Other antibodies as cell binding molecules/ligands include, but are not limited to, are antibodies against the following antigens: Aminopeptidase N (CD13), Annexin A1, B7-H3 (CD276, various cancers), CA125 (ovarian), CA15-3 (carcinomas), CA19-9 (carcinomas), L6 (carcinomas), Lewis Y (carcinomas), Lewis X (carcinomas), alpha fetoprotein (carcinomas), CA242 (colorectal), placental alkaline phosphatase (carcinomas), prostate specific antigen (prostate), prostatic acid phosphatase (prostate), epidermal growth factor (carcinomas), CD2

(Hodgkin's disease, NHL lymphoma, multiple myeloma), CD3 epsilon (T cell lymphoma, lung, breast, gastric, ovarian cancers, autoimmune diseases, malignant ascites), CD19 (B cell malignancies), CD20 (non-Hodgkin's lymphoma), CD22 (leukemia, lymphoma, multiple myeloma, SLE), CD30 (Hodgkin's lymphoma), CD33 (leukemia, autoimmune diseases), CD38 (multiple myeloma), CD40 (lymphoma, multiple myeloma, leukemia (CLL)), CD51 (Metastatic melanoma, sarcoma), CD52 (leukemia), CD56 (small cell lung cancers, ovarian cancer, Merkel cell carcinoma, and the liquid tumor, multiple myeloma), CD66e (cancers), CD70 (metastatic renal cell carcinoma and non-Hodgkin lymphoma), CD74 (multiple myeloma), CD80 (lymphoma), CD98 (cancers), mucin (carcinomas), CD221 (solid tumors), CD227 (breast, ovarian cancers), CD262 (NSCLC and other cancers), CD309 (ovarian cancers), CD326 (solid tumors), CEACAM3 (colorectal, gastric cancers), CEACAM5 (carcinoembryonic antigen; CEA, CD66e) (breast, colorectal and lung cancers), DLL4 (delta-like-4), EGFR (Epidermal Growth Factor Receptor, various cancers), CTLA4 (melanoma), CXCR4 (CD184, Heme-oncology, solid tumors), Endoglin (CD105, solid tumors), EPCAM (epithelial cell adhesion molecule, bladder, head, neck, colon, NHL prostate, and ovarian cancers), ERBB2 (Epidermal Growth Factor Receptor 2; lung, breast, prostate cancers), FCGR1 (autoimmune diseases), FOLR (folate receptor, ovarian cancers), GD2 ganglioside (cancers), G-28 (a cell surface antigen glycolipid, melanoma), GD3 idotype (cancers), Heat shock proteins (cancers), HER1 (lung, stomach cancers), HER2 (breast, lung and ovarian cancers), HLA-DR10 (NHL), HLA-DRB (NHL, B cell leukemia), human chorionic gonadotropin (carcinoma), IGF1R (insulin-like growth factor 1 receptor, solid tumors, blood cancers), IL-2 receptor (interleukin 2 receptor, T-cell leukemia and lymphomas), IL-6R (interleukin 6 receptor, multiple myeloma, RA, Castleman's disease, IL6 dependent tumors), Integrins ($\alpha\text{v}\beta\text{3}$, $\alpha\text{5}\beta\text{1}$, $\alpha\text{6}\beta\text{4}$, $\alpha\text{11}\beta\text{3}$, $\alpha\text{5}\beta\text{5}$, $\alpha\text{v}\beta\text{5}$, for various cancers), MAGE-1 (carcinomas), MAGE-2 (carcinomas), MAGE-3 (carcinomas), MAGE 4 (carcinomas), anti-transferrin receptor (carcinomas), p97 (melanoma), MS4A1 (membrane-spanning 4-domains subfamily A member 1, Non-Hodgkin's B cell lymphoma, leukemia), MUC1 or MUC1-KLH (breast, ovarian, cervix, bronchus and gastrointestinal cancer), MUC16 (CA125) (Ovarian cancers), CEA (colorectal), gp100 (melanoma), MART1 (melanoma), MPG (melanoma), MS4A1 (membrane-spanning 4-domains subfamily A, small cell lung cancers, NHL), Nucleolin, Neu oncogene product (carcinomas), P21 (carcinomas), Paratope of anti-(N-glycolylneuraminic acid, Breast, Melanoma cancers), PLAP-like testicular alkaline phosphatase (ovarian, testicular cancers), PSMA (prostate tumors), PSA (prostate), ROBO4, TAG 72 (tumour associated glycoprotein 72, AML, gastric, colorectal, ovarian cancers), T cell transmembrane protein (cancers), Tie (CD202b), TNFRSF10B (tumor necrosis factor receptor superfamily member 10B,

cancers), TNFRSF13B (tumor necrosis factor receptor superfamily member 13B, multiple myeloma, NHL, other cancers, RA and SLE), TPBG (trophoblast glycoprotein, Renal cell carcinoma), TRAIL-R1 (Tumor necrosis apoptosis Inducing ligand Receptor 1, lymphoma, NHL, colorectal, lung cancers), VCAM-1 (CD106, Melanoma), VEGF, VEGF-A, VEGF-2 (CD309) (various cancers). Some other tumor associated antigens recognized by antibodies have been reviewed (Gerber, et al, mAbs 1:3, 247-53 (2009); Novellino et al, Cancer Immunol Immunother. 54(3), 187-207 (2005). Franke, et al, Cancer Biother Radiopharm. 2000, 15, 459-76).

The cell-binding agents, more preferred antibodies, can be any agents that are able to against tumor cells, virus infected cells, microorganism infected cells, parasite infected cells, autoimmune cells, activated cells, myeloid cells, activated T-cells, B cells, or melanocytes. More specifically the cell binding agents can be any agent/molecule that is able to against any one of the following antigens or receptors: CD3, CD4, CD5, CD6, CD7, CD8, CD9, CD10, CD11a, CD11b, CD11c, CD12w, CD14, CD15, CD16, CDw17, CD18, CD19, CD20, CD21, CD22, CD23, CD24, CD25, CD26, CD27, CD28, CD29, CD30, CD31, CD32, CD33, CD34, CD35, CD36, CD37, CD38, CD39, CD40, CD41, CD42, CD43, CD44, CD45, CD46, CD47, CD48, CD49b, CD49c, CD51, CD52, CD53, CD54, CD55, CD56, CD58, CD59, CD61, CD62E, CD62L, CD62P, CD63, CD66, CD68, CD69, CD70, CD72, CD74, CD79, CD79a, CD79b, CD80, CD81, CD82, CD83, CD86, CD87, CD88, CD89, CD90, CD91, CD95, CD96, CD98, CD100, CD103, CD105, CD106, CD109, CD117, CD120, CD125, CD126, CD127, CD133, CD134, CD135, CD137, CD138, CD141, CD142, CD143, CD144, CD147, CD151, CD147, CD152, CD154, CD156, CD158, CD163, CD166, .CD168, CD174, CD180, CD184, CDw186, CD194, CD195, CD200, CD200a, CD200b, CD209, CD221, CD227, CD235a, CD240, CD262, CD271, CD274, CD276 (B7-H3), CD303, CD304, CD309, CD326, 4-1BB, 5AC, 5T4 (Trophoblast glycoprotein, TPBG, 5T4, Wnt-Activated Inhibitory Factor 1 or WAI1), Adenocarcinoma antigen, AGS-5, AGS-22M6, Activin receptor-like kinase 1, AFP, AKAP-4, ALK, Alpha integrin, Alpha v beta6, Amino-peptidase N, Amyloid beta, Androgen receptor, Angiopoietin 2, Angiopoietin 3, Annexin A1, Anthrax toxin-protective antigen, Anti-transferrin receptor, AOC3 (VAP-1), B7-H3, Bacillus anthracis anthrax, BAFF (B-cell activating factor), B-lymphoma cell, bcr-abl, Bombesin, BORIS, C5, C242 antigen, CA125 (carbohydrate antigen 125, MUC16), CA-IX (or CAIX, carbonic anhydrase 9), CALLA, CanAg, Canis lupus familiaris IL31, Carbonic anhydrase IX, Cardiac myosin, CCL11 (C-C motif chemokine 11), CCR4 (C-C chemokine receptor type 4, CD194), CCR5, CD3E (epsilon), CEA (Carcinoembryonic antigen), CEACAM3, CEACAM5 (carcinoembryonic antigen), CFD (Factor D), Ch4D5, Cholecystokinin 2 (CCK2R), CLDN18 (Claudin-18), Clumping factor A, CRIPTO, FCSF1R (Colony stimulating factor 1 receptor, CD115), CSF2 (colony stimulating factor 2,

Granulocyte-macrophage colony-stimulating factor(GM-CSF)), CTLA4 (cytotoxic T-lymphocyte associated protein 4), CTAA16.88 tumor antigen, CXCR4 (CD184),C-X-C chemokine receptor type 4, cyclic ADP ribose hydrolase, Cyclin B1, CYP1B1, Cytomegalovirus, Cytomegalovirus glycoprotein B, Dabigatran, DLL4 (delta-like-ligand 4), DPP4 (Dipeptidyl-peptidase 4), DR5 (Death receptor 5), E. coli shiga toxintype-1, E. coli shiga toxintype-2, ED-B, EGFL7 (EGF-like domain-containing protein 7), EGFR, EGFRII, EGFRvIII, Endoglin (CD105), Endothelin B receptor, Endotoxin, EpCAM (epithelial cell adhesion molecule), EphA2, Episialin, ERBB2 (Epidermal Growth Factor Receptor 2), ERBB3, ERG (TMPRSS2 ETS fusion gene), Escherichia coli,ETV6-AML, FAP (Fibroblast activation proteinalpha), FCGR1, alpha-Fetoprotein, Fibrin II, beta chain, Fibronectin extra domain-B, FOLR (folate receptor), Folate receptor alpha, Folate hydrolase, Fos-related antigen 1, F protein of respiratory syncytial virus, Frizzled receptor, Fucosyl GM1,GD2 ganglioside, G-28 (a cell surface antigen glyvolipid), GD3 idio type, GloboH, Glypican 3, N-glycolylneuraminic acid, GM3, GMCSF receptor α -chain, Growth differentiation factor 8, GP100, GPNMB (Transmembrane glycoprotein NMB), GUCY2C (Guanylate cyclase 2C, guanylyl cyclase C(GC-C), intestinal Guanylate cyclase, Guanylate cyclase-C receptor, Heat-stable enterotoxin receptor (hSTAR)), Heat shock proteins, Hemagglutinin, Hepatitis B surface antigen, Hepatitis B virus, HER1 (human epidermal growth factor receptor 1), HER2, HER2/neu, HER3 (ERBB-3), IgG4, HGF/SF (Hepatocyte growth factor/scatter factor), HHGFR, HIV-1, Histone complex, HLA-DR (human leukocyte antigen), HLA-DR10, HLA-DRB , HMWMAA, Human chorionic gonadotropin, HNGF, Human scatter factor receptor kinase, HPV E6/E7, Hsp90, hTERT, ICAM-1 (Intercellular Adhesion Molecule 1), Idiotyp e, IGF1R (IGF-1, insulin-like growth factor 1 receptor), IGHE, IFN- γ , Influeza hemag-glutinin, IgE, IgE Fc region, IGHE, IL-1, IL-2 receptor (interleukin 2 receptor), IL-4, IL-5, IL-6, IL-6R (interleukin 6 receptor), IL-9, IL-10, IL-12, IL-13, IL-17, IL-17A, IL-20, IL-22, IL-23, IL31RA, ILGF2 (Insulin-like growth factor 2), Integrins (α 4, α IIb β 3, α v β 3, α 4 β 7, α 5 β 1, α 6 β 4, α 7 β 7, α II β 3, α 5 β 5, α v β 5), Interferon gamma-induced protein, ITGA2, ITGB2, KIR2D, LCK, Le, Legumain, Lewis-Y antigen, LFA-1(Lymphocyte function-associated antigen 1, CD11a), LHRH, LINGO-1, Lipoteichoic acid, LIV1A, LMP2, LTA, MAD-CT-1, MAD-CT-2, MAGE-1, MAGE-2, MAGE-3, MAGE A1, MAGE A3, MAGE 4, MART1, MCP-1, MIF (Macrophage migration inhibitory factor, or glycosylation-inhibiting factor (GIF)), MS4A1 (membrane-spanning 4-domains subfamily A member 1), MSLN (mesothelin), MUC1(Mucin 1, cell surfaceassociated (MUC1) orpolymorphic epithelial mucin (PEM)), MUC1-KLH, MUC16 (CA125), MCP1(monocyte chemotactic protein 1), MelanA/MART1, ML-IAP, MPG, MS4A1 (membrane-spanning 4-domains subfamily A), MYCN, Myelin-associated glycoprotein, Myostatin, NA17, NARP-1, NCA-90 (granulocyte antigen), Nectin-4 (ASG-22ME),

NGF, Neural apoptosis-regulated proteinase 1, NOGO-A, Notch receptor, Nucleolin, Neu
 oncogene product, NY-BR-1, NY-ESO-1, OX-40, OxLDL (Oxidized low-density lipoprotein),
 OY-TES1, P21, p53 nonmutant, P97, Page4, PAP, Paratope of anti-(N-glycolylneuraminic acid),
 PAX3, PAX5, PCSK9, PDCD1 (PD-1, Programmed cell death protein 1, CD279), PDGF-R α
 5 (Alpha-type platelet-derived growth factor receptor), PDGFR- β , PDL-1, PLAC1, PLAP-like
 testicular alkaline phosphatase, Platelet-derived growth factor receptor beta, Phosphate-sodium
 co-transporter, PMEL 17, Polysialic acid, Proteinase3 (PR1), Prostatic carcinoma, PS (Phos-
 phatidylserine), Prostatic carcinoma cells, Pseudomonas aeruginosa, PSMA, PSA, PSCA,
 Rabies virus glycoprotein, RHD (Rh polypeptide 1 (RhPI), CD240), Rhesus factor, RANKL,
 10 RhoC, Ras mutant, RGS5, ROBO4, Respiratory syncytial virus, RON, Sarcoma translocation
 breakpoints, SART3, Sclerostin, SLAMF7 (SLAM family member 7), Selectin P, SDC1
 (Syndecan 1), sLe(a), Somatomedin C, SIP (Sphingosine-1-phosphate), Somatostatin, Sperm
 protein 17, SSX2, STEAP1 (six-transmembrane epithelial antigen of the prostate 1), STEAP2,
 STn, TAG-72 (tumor associated glycoprotein 72), Survivin, T-cell receptor, T cell transmembrane
 15 protein, TEM1 (Tumor endothelial marker 1), TENB2, Tenascin C (TN-C), TGF- α , TGF- β
 (Transforming growth factor beta), TGF- β 1, TGF- β 2 (Transforming growth factor-beta 2), Tie
 (CD202b), Tie2, TIM-1 (CDX-014), Tn, TNF, TNF- α , TNFRSF8, TNFRSF10B (tumor necrosis
 factor receptor superfamily member 10B), TNFRSF13B (tumor necrosis factor receptor
 superfamily member 13B), TPBG (trophoblast glycoprotein), TRAIL-R1 (Tumor necrosis
 20 apoptosis Inducing ligand Receptor 1), TRAILR2 (Death receptor 5 (DR5)), tumor-associated
 calcium signal transducer 2, tumor specific glycosylation of MUC1, TWEAK receptor, TYRP1
 (glycoprotein 75), TRP-2, Tyrosinase, VCAM-1 (CD106), VEGF, VEGF-A, VEGF-2 (CD309),
 VEGFR-1, VEGFR2, or vimentin, WT1, XAGE 1, or cells expressing any insulin growth factor
 receptors, or any epidermal growth factor receptors.

25 In another specific embodiment, the cell-binding ligand-drug conjugates via the bridge
 linkers of this invention are used for the targeted treatment of cancers. The targeted cancers
 include, but are not limited, Adrenocortical Carcinoma, Anal Cancer, Bladder Cancer, Brain
 Tumor (Adult, Brain Stem Glioma, Childhood, Cerebellar Astrocytoma, Cerebral Astrocytoma,
 Ependymoma, Medulloblastoma, Supratentorial Primitive Neuroectodermal and Pineal Tumors,
 30 Visual Pathway and Hypothalamic Glioma), Breast Cancer, Carcinoid Tumor, Gastrointestinal,
 Carcinoma of Unknown Primary, Cervical Cancer, Colon Cancer, Endometrial Cancer,
 Esophageal Cancer, Extrahepatic Bile Duct Cancer, Ewings Family of Tumors (PNET),
 Extracranial Germ Cell Tumor, Eye Cancer, Intraocular Melanoma, Gallbladder Cancer, Gastric
 Cancer (Stomach), Germ Cell Tumor, Extragenital, Gestational Trophoblastic Tumor, Head and

Neck Cancer, Hypopharyngeal Cancer, Islet Cell Carcinoma, Kidney Cancer (renal cell cancer), Laryngeal Cancer, Leukemia (Acute Lymphoblastic, Acute Myeloid, Chronic Lymphocytic, Chronic Myelogenous, Hairy Cell), Lip and Oral Cavity Cancer, Liver Cancer, Lung Cancer (Non-Small Cell, Small Cell, Lymphoma (AIDS-Related, Central Nervous System, Cutaneous T-Cell, Hodgkin's Disease, Non-Hodgkin's Disease, Malignant Mesothelioma, Melanoma, Merkel Cell Carcinoma, Metastatic Squamous Neck Cancer with Occult Primary, Multiple Myeloma, and Other Plasma Cell Neoplasms, Mycosis Fungoides, Myelodysplastic Syndrome, Myeloproliferative Disorders, Nasopharyngeal Cancer, Neuroblastoma, Oral Cancer, Oropharyngeal Cancer, Osteosarcoma, Ovarian Cancer (Epithelial, Germ Cell Tumor, Low Malignant Potential Tumor), Pancreatic Cancer (Exocrine, Islet Cell Carcinoma), Paranasal Sinus and Nasal Cavity Cancer, Parathyroid Cancer, Penile Cancer, Pheochromocytoma Cancer, Pituitary Cancer, Plasma Cell Neoplasm, Prostate Cancer Rhabdomyosarcoma, Rectal Cancer, Renal Cell Cancer (kidney cancer), Renal Pelvis and Ureter (Transitional Cell), Salivary Gland Cancer, Sezary Syndrome, Skin Cancer, Skin Cancer (Cutaneous T-Cell Lymphoma, Kaposi's Sarcoma, Melanoma), Small Intestine Cancer, Soft Tissue Sarcoma, Stomach Cancer, Testicular Cancer, Thymoma (Malignant), Thyroid Cancer, Urethral Cancer, Uterine Cancer (Sarcoma), Unusual Cancer of Childhood, Vaginal Cancer, Vulvar Cancer, Wilms' Tumor.

In another specific embodiment, the cell-binding-drug conjugates via the bridge linkers of this invention are used in accordance with the compositions and methods for the treatment or prevention of an autoimmune disease. The autoimmune diseases include, but are not limited, Achlorhydra Autoimmune Active Chronic Hepatitis, Acute Disseminated Encephalomyelitis, Acute hemorrhagic leukoencephalitis, Addison's Disease, Agammaglobulinemia, Alopecia areata, Amyotrophic Lateral Sclerosis, Ankylosing Spondylitis, Anti-GBM/TBM Nephritis, Antiphospholipid syndrome, Antisynthetase syndrome, Arthritis, Atopic allergy, Atopic Dermatitis, Autoimmune Aplastic Anemia, Autoimmune cardiomyopathy, Autoimmune hemolytic anemia, Autoimmune hepatitis, Autoimmune inner ear disease, Autoimmune lymphoproliferative syndrome, Autoimmune peripheral neuropathy, Autoimmune pancreatitis, Autoimmune polyendocrine syndrome Types I, II, & III, Autoimmune progesterone dermatitis, Autoimmune thrombocytopenic purpura, Autoimmune uveitis, Balo disease/Balo concentric sclerosis, Bechets Syndrome, Berger's disease, Bickerstaff's encephalitis, Blau syndrome, Bullous Pemphigoid, Castleman's disease, Chagas disease, Chronic Fatigue Immune Dysfunction Syndrome, Chronic inflammatory demyelinating polyneuropathy, Chronic recurrent multifocal osteomyelitis, Chronic Lyme disease, Chronic obstructive pulmonary disease, Churg-Strauss syndrome, Cicatricial Pemphigoid, Coeliac Disease, Cogan syndrome, Cold agglutinin disease,

Complement component 2 deficiency, Cranial arteritis, CREST syndrome, Crohns Disease (a type of idiopathic inflammatory bowel diseases), Cushing's Syndrome, Cutaneous leukocytoclastic angiitis, Dego's disease, Dercum's disease, Dermatitis herpetiformis, Dermatomyositis, Diabetes mellitus type 1, Diffuse cutaneous systemic sclerosis, Dressler's syndrome, Discoid lupus erythematosus, Eczema, Endometriosis, Enthesitis-related arthritis, Eosinophilic fasciitis, Epidermolysis bullosa acquisita, Erythema nodosum, Essential mixed cryoglobulinemia, Evan's syndrome, Fibrodysplasia ossificans progressiva, Fibromyalgia, Fibromyositis, Fibrosing aveolitis, Gastritis, Gastrointestinal pemphigoid, Giant cell arteritis, Glomerulonephritis, Goodpasture's syndrome, Graves' disease, Guillain-Barré syndrome, Hashimoto's encephalitis, Hashimoto's thyroiditis, Haemolytic anaemia, Henoch-Schonlein purpura, Herpes gestationis, Hidradenitis suppurativa, Hughes syndrome (See Antiphospholipid syndrome), Hypogammaglobulinemia, Idiopathic Inflammatory Demyelinating Diseases, Idiopathic pulmonary fibrosis, Idiopathic thrombocytopenic purpura (See Autoimmune thrombocytopenic purpura), IgA nephropathy (Also Berger's disease), Inclusion body myositis, Inflammatory demyelinating polyneuropathy, Interstitial cystitis, Irritable Bowel Syndrome, Juvenile idiopathic arthritis, Juvenile rheumatoid arthritis, Kawasaki's Disease, Lambert-Eaton myasthenic syndrome, Leukocytoclastic vasculitis, Lichen planus, Lichen sclerosus, Linear IgA disease (LAD), Lou Gehrig's Disease (Also Amyotrophic lateral sclerosis), Lupoid hepatitis, Lupus erythematosus, Majeeed syndrome, Ménière's disease, Microscopic polyangiitis, Miller-Fisher syndrome, Mixed Connective Tissue Disease, Morphea, Mucha-Habermann disease, Muckle–Wells syndrome, Multiple Myeloma, Multiple Sclerosis, Myasthenia gravis, Myositis, Narcolepsy, Neuromyelitis optica (Devic's Disease), Neuromyotonia, Occular cicatricial pemphigoid, Opsoclonus myoclonus syndrome, Ord thyroiditis, Palindromic rheumatism, PANDAS (Pediatric Autoimmune Neuropsychiatric Disorders Associated with Streptococcus), Paraneoplastic cerebellar degeneration, Paroxysmal nocturnal hemoglobinuria, Parry Romberg syndrome, Parsonnage-Turner syndrome, Pars planitis, Pemphigus, Pemphigus vulgaris, Pernicious anaemia, Perivenous encephalomyelitis, POEMS syndrome, Polyarteritis nodosa, Polymyalgia rheumatica, Polymyositis, Primary biliary cirrhosis, Primary sclerosing cholangitis, Progressive inflammatory neuropathy, Psoriasis, Psoriatic Arthritis, Pyoderma gangrenosum, Pure red cell aplasia, Rasmussen's encephalitis, Raynaud phenomenon, Relapsing polychondritis, Reiter's syndrome, Restless leg syndrome, Retroperitoneal fibrosis, Rheumatoid arthritis, Rheumatoid fever, Sarcoidosis, Schizophrenia, Schmidt syndrome, Schnitzler syndrome, Scleritis, Scleroderma, Sjögren's syndrome, Spondyloarthropathy, Sticky blood syndrome, Still's Disease, Stiff person syndrome, Subacute bacterial endocarditis, Susac's syndrome, Sweet syndrome, Sydenham

Chorea, Sympathetic ophthalmia, Takayasu's arteritis, Temporal arteritis (giant cell arteritis), Tolosa-Hunt syndrome, Transverse Myelitis, Ulcerative Colitis (a type of idiopathic inflammatory bowel diseases), Undifferentiated connective tissue disease, Undifferentiated spondyloarthropathy, Vasculitis, Vitiligo, Wegener's granulomatosis, Wilson's syndrome, Wiskott-Aldrich syndrome

In another specific embodiment, a binding molecule used for the conjugate via the bridge linkers of this invention for the treatment or prevention of an autoimmune disease can be, but are not limited to, anti-elastin antibody; Abys against epithelial cells antibody; Anti-Basement Membrane Collagen Type IV Protein antibody; Anti-Nuclear Antibody; Anti ds DNA; Anti ss DNA, Anti Cardiolipin Antibody IgM, IgG; anti-celiac antibody; Anti Phospholipid Antibody IgK, IgG; Anti SM Antibody; Anti Mitochondrial Antibody; Thyroid Antibody; Microsomal Antibody, T-cells antibody; Thyroglobulin Antibody, Anti SCL-70; Anti-Jo; Anti-U.sub.1RNP; Anti-La/SSB; Anti SSA; Anti SSB; Anti Perital Cells Antibody; Anti Histones; Anti RNP; C-ANCA; P-ANCA; Anti centromere; Anti-Fibrillarin, and Anti GBM Antibody, Anti-ganglioside antibody; Anti-Desmoglein 3 antibody; Anti-p62 antibody; Anti-sp100 antibody; Anti-Mitochondrial(M2) antibody; Rheumatoid factor antibody; Anti-MCV antibody; Anti-topoisomerase antibody; Anti-neutrophil cytoplasmic(cANCA) antibody.

In certain preferred embodiments, the binding molecule for the conjugate in the present invention, can bind to both a receptor and a receptor complex expressed on an activated lymphocyte which is associated with an autoimmune disease. The receptor or receptor complex can comprise an immunoglobulin gene superfamily member (e.g. CD2, CD3, CD4, CD8, CD19, CD20, CD22, CD28, CD30, CD33, CD37, CD38, CD56, CD70, CD79, CD79b, CD90, CD125, CD137, CD138, CD147, CD152/CTLA-4, PD-1, or ICOS), a TNF receptor superfamily member (e.g. CD27, CD40, CD95/Fas, CD134/OX40, CD137/4-1BB, INF-R1, TNFR-2, RANK, TACI, BCMA, osteoprotegerin, Apo2/TRAIL-R1, TRAIL-R2, TRAIL-R3, TRAIL-R4, and APO-3), an integrin, a cytokine receptor, a chemokine receptor, a major histocompatibility protein, a lectin (C-type, S-type, or I-type), or a complement control protein.

In another specific embodiment, useful cell binding ligands that are immunospecific for a viral or a microbial antigen are humanized or human monoclonal antibodies. As used herein, the term "viral antigen" includes, but is not limited to, any viral peptide, polypeptide protein (e.g. HIV gp120, HIV nef, RSV F glycoprotein, influenza virus neuramini-dase, influenza virus hemagglutinin, HTLV tax, herpes simplex virus glycoprotein (e.g. gB, gC, gD, and gE) and hepatitis B surface antigen) that is capable of eliciting an immune response. As used herein, the term "microbial antigen" includes, but is not limited to, any microbial peptide, polypeptide,

protein, saccharide, polysaccharide, or lipid molecule (e.g., a bacteria, fungi, pathogenic protozoa, or yeast polypeptides including, e.g., LPS and capsular polysaccharide 5/8) that is capable of eliciting an immune response. Examples of antibodies available for the viral or microbial infection include, but are not limited to, Palivizumab which is a humanized anti-respiratory syncytial virus monoclonal antibody for the treatment of RSV infection; PRO542 which is a CD4 fusion antibody for the treatment of HIV infection; Ostavir which is a human antibody for the treatment of hepatitis B virus; PROTVIR which is a humanized IgG.sub.1 antibody for the treatment of cytomegalovirus; and anti-LPS antibodies.

The cell binding molecules–drug conjugates via the bridge linkers of this invention can be used in the treatment of infectious diseases. These infectious diseases include, but are not limited to, Acinetobacter infections, Actinomycosis, African sleeping sickness (African trypanosomiasis), AIDS (Acquired immune deficiency syndrome), Amebiasis, Anaplasmosis, Anthrax, Arcanobacterium haemolyticum infection, Argentine hemorrhagic fever, Ascariasis, Aspergillosis, Astrovirus infection, Babesiosis, Bacillus cereus infection, Bacterial pneumonia, Bacterial vaginosis, Bacteroides infection, Balantidiasis, Baylisascaris infection, BK virus infection, Black piedra, Blastocystis hominis infection, Blastomycosis, Bolivian hemorrhagic fever, Borrelia infection, Botulism (and Infant botulism), Brazilian hemorrhagic fever, Brucellosis, Burkholderia infection, Buruli ulcer, Calicivirus infection (Norovirus and Sapovirus), Campylobacteriosis, Candidiasis (Moniliasis; Thrush), Cat-scratch disease, Cellulitis, Chagas Disease (American trypanosomiasis), Chancroid, Chickenpox, Chlamydia, Chlamydophila pneumoniae infection, Cholera, Chromoblastomycosis, Clonorchiasis, Clostridium difficile infection, Coccidioidomycosis, Colorado tick fever, Common cold (Acute viral rhinopharyngitis; Acute coryza), Creutzfeldt-Jakob disease, Crimean-Congo hemorrhagic fever, Cryptococcosis, Cryptosporidiosis, Cutaneous larva migrans, Cyclosporiasis, Cysticercosis, Cytomegalovirus infection, Dengue fever, Dientamoebiasis, Diphtheria, Diphyllbothriasis, Dracunculiasis, Ebola hemorrhagic fever, Echinococcosis, Ehrlichiosis, Enterobiasis (Pinworm infection), Enterococcus infection, Enterovirus infection, Epidemic typhus, Erythema infectiosum (Fifth disease), Exanthem subitum, Fasciolopsiasis, Fasciolosis, Fatal familial insomnia, Filariasis, Food poisoning by Clostridium perfringens, Free-living amebic infection, Fusobacterium infection, Gas gangrene (Clostridial myonecrosis), Geotrichosis, Gerstmann-Sträussler-Scheinker syndrome, Giardiasis, Glanders, Gnathostomiasis, Gonorrhea, Granuloma inguinale (Donovanosis), Group A streptococcal infection, Group B streptococcal infection, Haemophilus influenzae infection, Hand, foot and mouth disease (HFMD), Hantavirus Pulmonary Syndrome, Helicobacter pylori infection, Hemolytic-uremic syndrome, Hemorrhagic fever with renal syndrome, Hepatitis A, Hepatitis B,

Hepatitis C, Hepatitis D, Hepatitis E, Herpes simplex, Histoplasmosis, Hookworm infection, Human bocavirus infection, Human ewingii ehrlichiosis, Human granulocytic anaplasmosis, Human metapneumovirus infection, Human monocytic ehrlichiosis, Human papillomavirus infection, Human parainfluenza virus infection, Hymenolepiasis, Epstein-Barr Virus Infectious

5 Mononucleosis (Mono), Influenza, Isosporiasis, Kawasaki disease, Keratitis, *Kingella kingae* infection, Kuru, Lassa fever, Legionellosis (Legionnaires' disease), Legionellosis (Pontiac fever), Leishmaniasis, Leprosy, Leptospirosis, Listeriosis, Lyme disease (Lyme borreliosis), Lymphatic filariasis (Elephantiasis), Lymphocytic choriomeningitis, Malaria, Marburg hemorrhagic fever, Measles, Melioidosis (Whitmore's disease), Meningitis, Meningococcal disease, Metagonimiasis,

10 Microsporidiosis, Molluscum contagiosum, Mumps, Murine typhus (Endemic typhus), Mycoplasma pneumonia, Mycetoma, Myiasis, Neonatal conjunctivitis (Ophthalmia neonatorum), (New) Variant Creutzfeldt-Jakob disease (vCJD, nvCJD), Nocardiosis, Onchocerciasis (River blindness), Paracoccidioidomycosis (South American blastomycosis), Paragonimiasis, Pasteurellosis, Pediculosis capitis (Head lice), Pediculosis corporis (Body lice), Pediculosis pubis

15 (Pubic lice, Crab lice), Pelvic inflammatory disease, Pertussis (Whooping cough), Plague, Pneumococcal infection, Pneumocystis pneumonia, Pneumonia, Poliomyelitis, Prevotella infection, Primary amoebic meningoencephalitis, Progressive multifocal leukoencephalopathy, Psittacosis, Q fever, Rabies, Rat-bite fever, Respiratory syncytial virus infection, Rhinosporidiosis, Rhinovirus infection, Rickettsial infection, Rickettsial-pox, Rift Valley fever, Rocky mountain

20 spotted fever, Rotavirus infection, Rubella, Salmonellosis, SARS (Severe Acute Respiratory Syndrome), Scabies, Schistosomiasis, Sepsis, Shigellosis (Bacillary dysentery), Shingles (Herpes zoster), Smallpox (Variola), Sporotrichosis, Staphylococcal food poisoning, Staphylococcal infection, Strongyloidiasis, Syphilis, Taeniasis, Tetanus (Lockjaw), Tinea barbae (Barber's itch), Tinea capitis (Ringworm of the Scalp), Tinea corporis (Ringworm of the Body), Tinea cruris

25 (Jock itch), Tinea manuum (Ringworm of the Hand), Tinea nigra, Tinea pedis (Athlete's foot), Tinea unguium (Onychomycosis), Tinea versicolor (Pityriasis versicolor), Toxocariasis (Ocular Larva Migrans), Toxocariasis (Visceral Larva Migrans), Toxoplasmosis, Trichinellosis, Trichomoniasis, Trichuriasis (Whipworm infection), Tuberculosis, Tularemia, Ureaplasma urealyticum infection, Venezuelan equine encephalitis, Venezuelan hemorrhagic fever, Viral

30 pneumonia, West Nile Fever, White piedra (Tinea blanca), *Yersinia pseudotuberculosis* infection, Yersiniosis, Yellow fever, Zygomycosis.

The cell binding molecule, which is more preferred to be an antibody described in this patent that are against pathogenic strains include, but are not limit, *Acinetobacter baumannii*, *Actinomyces israelii*, *Actinomyces gerencseriae* and *Propionibacterium propionicus*,

Trypanosoma brucei, HIV (Human immunodeficiency virus), Entamoeba histolytica, Anaplasma
 genus, Bacillus anthracis, Arcanobacterium haemolyticum, Junin virus, Ascaris lumbricoides,
 Aspergillus genus, Astroviridae family, Babesia genus, Bacillus cereus, multiple bacteria,
 Bacteroides genus, Balantidium coli, Baylisascaris genus, BK virus, Piedraia hortae, Blastocystis
 5 hominis, Blastomyces dermatitidis, Machupo virus, Borrelia genus, Clostridium botulinum,
 Sabia, Brucella genus, usually Burkholderia cepacia and other Burkholderia species,
 Mycobacterium ulcerans, Caliciviridae family, Campylobacter genus, usually Candida albicans
 and other Candida species, Bartonella henselae, Group A Streptococcus and Staphylococcus,
 Trypanosoma cruzi, Haemophilus ducreyi, Varicella zoster virus (VZV), Chlamydia trachomatis,
 10 Chlamydophila pneumoniae, Vibrio cholerae, Fonsecaea pedrosoi, Clonorchis sinensis,
 Clostridium difficile, Coccidioides immitis and Coccidioides posadasii, Colorado tick fever virus,
 rhinoviruses, coronaviruses, CJD prion, Crimean-Congo hemorrhagic fever virus, Cryptococcus
 neoformans, Cryptosporidium genus, Ancylostoma braziliense; multiple parasites, Cyclospora
 cayetanensis, Taenia solium, Cytomegalovirus, Dengue viruses (DEN-1, DEN-2, DEN-3 and
 15 DEN-4) – Flaviviruses, Dientamoeba fragilis, Corynebacterium diphtheriae, Diphyllbothrium,
 Dracunculus medinensis, Ebolavirus, Echinococcus genus, Ehrlichia genus, Enterobius
 vermicularis, Enterococcus genus, Enterovirus genus, Rickettsia prowazekii, Parvovirus B19,
 Human herpesvirus 6 and Human herpesvirus 7, Fasciolopsis buski, Fasciola hepatica and
 Fasciola gigantica, FFI prion, Filarioidea superfamily, Clostridium perfringens, Fusobacterium
 20 genus, Clostridium perfringens; other Clostridium species, Geotrichum candidum, GSS prion,
 Giardia intestinalis, Burkholderia mallei, Gnathostoma spinigerum and Gnathostoma hispidum,
 Neisseria gonorrhoeae, Klebsiella granulomatis, Streptococcus pyogenes, Streptococcus
 agalactiae, Haemophilus influenzae, Enteroviruses, mainly Coxsackie A virus and Enterovirus 71,
 Sin Nombre virus, Helicobacter pylori, Escherichia coli O157:H7, Bunyaviridae family, Hepatitis
 25 A Virus, Hepatitis B Virus, Hepatitis C Virus, Hepatitis D Virus, Hepatitis E Virus, Herpes
 simplex virus 1, Herpes simplex virus 2, Histoplasma capsulatum, Ancylostoma duodenale and
 Necator americanus, Hemophilus influenzae, Human bocavirus, Ehrlichia ewingii, Anaplasma
 phagocytophilum, Human metapneumovirus, Ehrlichia chaffeensis, Human papillomavirus,
 Human parainfluenza viruses, Hymenolepis nana and Hymenolepis diminuta, Epstein-Barr Virus,
 30 Orthomyxoviridae family, Isospora belli, Kingella kingae, Klebsiella pneumoniae, Klebsiella
 ozaenas, Klebsiella rhinoscleromatis, Kuru prion, Lassa virus, Legionella pneumophila,
 Legionella pneumophila, Leishmania genus, Mycobacterium leprae and Mycobacterium
 lepromatosis, Leptospira genus, Listeria monocytogenes, Borrelia burgdorferi and other Borrelia
 species, Wuchereria bancrofti and Brugia malayi, Lymphocytic choriomeningitis virus (LCMV),

Plasmodium genus, Marburg virus, Measles virus, Burkholderia pseudomallei, Neisseria meningitides, Metagonimus yokagawai, Microsporidia phylum, Molluscum contagiosum virus (MCV), Mumps virus, Rickettsia typhi, Mycoplasma pneumoniae, numerous species of bacteria (Actinomycetoma) and fungi (Eumycetoma), parasitic dipterous fly larvae, Chlamydia trachomatis and Neisseria gonorrhoeae, vCJD prion, Nocardia asteroides and other Nocardia species, Onchocerca volvulus, Paracoccidioides brasiliensis, Paragonimus westermani and other Paragonimus species, Pasteurella genus, Pediculus humanus capitis, Pediculus humanus corporis, Phthirus pubis, Bordetella pertussis, Yersinia pestis, Streptococcus pneumoniae, Pneumocystis jirovecii, Poliovirus, Prevotella genus, Naegleria fowleri, JC virus, Chlamydophila psittaci, Coxiella burnetii, Rabies virus, Streptobacillus moniliformis and Spirillum minus, Respiratory syncytial virus, Rhinosporidium seeberi, Rhinovirus, Rickettsia genus, Rickettsia akari, Rift Valley fever virus, Rickettsia rickettsii, Rotavirus, Rubella virus, Salmonella genus, SARS coronavirus, Sarcoptes scabiei, Schistosoma genus, Shigella genus, Varicella zoster virus, Variola major or Variola minor, Sporothrix schenckii, Staphylococcus genus, Staphylococcus aureus, Streptococcus pyogenes, Strongyloides stercoralis, Treponema pallidum, Taenia genus, Clostridium tetani, Trichophyton genus, Trichophyton tonsurans, Trichophyton genus, Epidermophyton floccosum, Trichophyton rubrum, and Trichophyton mentagrophytes, Trichophyton rubrum, Hortaea werneckii, Trichophyton genus, Malassezia genus, Toxocara canis or Toxocara cati, Toxoplasma gondii, Trichinella spiralis, Trichomonas vaginalis, Trichuris trichiura, Mycobacterium tuberculosis, Francisella tularensis, Ureaplasma urealyticum, Venezuelan equine encephalitis virus, Vibrio cholerae, Guanarito virus, West Nile virus, Trichosporon beigeli, Yersinia pseudotuberculosis, Yersinia enterocolitica, Yellow fever virus, Mucorales order (Mucormycosis) and Entomophthorales order (Entomophthora-mycosis), Pseudomonas aeruginosa, Campylobacter (Vibrio) fetus, Aeromonas hydrophila, Edwardsiella tarda, Yersinia pestis, Shigella dysenteriae, Shigella flexneri, Shigella sonnei, Salmonella typhimurium, Treponema pertense, Treponema caratense, Borrelia vincentii, Borrelia burgdorferi, Leptospira icterohemorrhagiae, Pneumocystis carinii, Brucella abortus, Brucella suis, Brucella melitensis, Mycoplasma spp., Rickettsia prowazeki, Rickettsia tsutsugumushi, Chlamydia spp.; pathogenic fungi (Aspergillus fumigatus, Candida albicans, Histoplasma capsulatum); protozoa (Entamoeba histolytica, Trichomonas tenax, Trichomonas hominis, Trypanosoma gambiense, Trypanosoma rhodesiense, Leishmania donovani, Leishmania tropica, Leishmania braziliensis, Pneumocystis pneumonia, Plasmodium vivax, Plasmodium falciparum, Plasmodium malariae); or Helminths (Schistosoma japonicum, Schistosoma mansoni, Schistosoma haematobium, and hookworms).

Other antibodies as cell binding ligands used in this invention for treatment of viral disease include, but are not limited to, antibodies against antigens of pathogenic viruses, including as examples and not by limitation: Poxviridae, Herpesviridae, Adenoviridae, Papovaviridae, Enteroviridae, Picornaviridae, Parvoviridae, Reoviridae, Retroviridae, influenza viruses, parainfluenza viruses, mumps, measles, respiratory syncytial virus, rubella, Arboviridae, Rhabdoviridae, Arenaviridae, Non-A/Non-B Hepatitis virus, Rhinoviridae, Coronaviridae, Rotoviridae, Oncovirus [such as, HBV (Hepatocellular carcinoma), HPV (Cervical cancer, Anal cancer), Kaposi's sarcoma-associated herpesvirus (Kaposi's sarcoma), Epstein-Barr virus (Nasopharyngeal carcinoma, Burkitt's lymphoma, Primary central nervous system lymphoma), MCPyV (Merkel cell cancer), SV40 (Simian virus 40), HCV (Hepatocellular carcinoma), HTLV-I (Adult T-cell leukemia/lymphoma)], Immune disorders caused virus: [such as Human Immunodeficiency Virus (AIDS)]; Central nervous system virus: [such as, JCV (Progressive multifocal leukoencephalopathy), MeV (Subacute sclerosing panencephalitis), LCV (Lymphocytic choriomeningitis), Arbovirus encephalitis, Orthomyxoviridae (probable) (Encephalitis lethargica), RV (Rabies), Chandipura virus, Herpesviral meningitis, Ramsay Hunt syndrome type II; Poliovirus (Poliomyelitis, Post-polio syndrome), HTLV-I (Tropical spastic paraparesis)]; Cytomegalovirus (Cytomegalovirus retinitis, HSV (Herpetic keratitis)); Cardiovascular virus [such as CBV (Pericarditis, Myocarditis)]; Respiratory system/acute viral nasopharyngitis/viral pneumonia: [Epstein-Barr virus (EBV infection/Infectious mononucleosis), Cytomegalovirus; SARS coronavirus (Severe acute respiratory syndrome) Orthomyxoviridae: Influenzavirus A/B/C (Influenza/Avian influenza), Paramyxovirus: Human parainfluenza viruses (Parainfluenza), RSV (Human respiratory syncytialvirus), hMPV]; Digestive system virus [MuV (Mumps), Cytomegalovirus (Cytomegalovirus esophagitis); Adenovirus (Adenovirus infection); Rotavirus, Norovirus, Astrovirus, Coronavirus; HBV (Hepatitis B virus), CBV, HAV (Hepatitis A virus), HCV (Hepatitis C virus), HDV (Hepatitis D virus), HEV (Hepatitis E virus), HGV (Hepatitis G virus)]; Urogenital virus [such as, BK virus, MuV (Mumps)].

According to a further object, the present invention also concerns pharmaceutical compositions comprising the conjugate via the bridge linkers of the invention together with a pharmaceutically acceptable carrier, diluent, or excipient for treatment of cancers, infections or autoimmune disorders. The method for treatment of cancers, infections and autoimmune disorders can be practiced in vitro, in vivo, or ex vivo. Examples of in vitro uses include treatments of cell cultures in order to kill all cells except for desired variants that do not express the target antigen; or to kill variants that express undesired antigen. Examples of ex vivo uses include treatments of hematopoietic stem cells(HSC)prior to the performance of the transplantation (HSCT) into the

same patient in order to kill diseased or malignant cells. For instance, clinical ex vivo treatment to remove tumour cells or lymphoid cells from bone marrow prior to autologous transplantation in cancer treatment or in treatment of autoimmune disease, or to remove T cells and other lymphoid cells from allogeneic bone marrow or tissue prior to transplant in order to prevent graft-versus-host disease, can be carried out as follows. Bone marrow is harvested from the patient or other individual and then incubated in medium containing serum to which is added the conjugate of the invention, concentrations range from about 1 pM to 0.1 mM, for about 30 minutes to about 48 hours at about 37 °C. The exact conditions of concentration and time of incubation (=dose) are readily determined by the skilled clinicians. After incubation, the bone marrow cells are washed with medium containing serum and returned to the patient by i.v. infusion according to known methods. In circumstances where the patient receives other treatment such as a course of ablative chemotherapy or total-body irradiation between the time of harvest of the marrow and reinfusion of the treated cells, the treated marrow cells are stored frozen in liquid nitrogen using standard medical equipment.

For clinical in vivo use, the conjugate via the linkers of the invention will be supplied as solutions or as a lyophilized solid that can be redissolved in sterile water for injection. Examples of suitable protocols of conjugate administration are as follows. Conjugates are given weekly for 8~20 weeks as an i.v. bolus. Bolus doses are given in 50 to 500 ml of normal saline to which human serum albumin (e.g. 0.5 to 1 mL of a concentrated solution of human serum albumin, 100 mg/mL) can be added. Dosages will be about 50 µg to 20 mg/kg of body weight per week, i.v. (range of 10 µg to 200 mg/kg per injection). 4~20 weeks after treatment, the patient may receive a second course of treatment. Specific clinical protocols with regard to route of administration, excipients, diluents, dosages, times, etc., can be determined by the skilled clinicians.

Examples of medical conditions that can be treated according to the in vivo or ex vivo methods of killing selected cell populations include malignancy of any types of cancer, autoimmune diseases, graft rejections, and infections (viral, bacterial or parasite).

The amount of a conjugate which is required to achieve the desired biological effect, will vary depending upon a number of factors, including the chemical characteristics, the potency, and the bioavailability of the conjugates, the type of disease, the species to which the patient belongs, the diseased state of the patient, the route of administration, all factors which dictate the required dose amounts, delivery and regimen to be administered.

In general terms, the conjugates via the linkers of this invention may be provided in an aqueous physiological buffer solution containing 0.1 to 10% w/v conjugates for parenteral administration. Typical dose ranges are from 1 µg/kg to 0.1 g/kg of body weight per day; a

preferred dose range is from 0.01 mg/kg to 20 mg/kg of body weight per day, or per week, or an equivalent dose in a human child. The preferred dosage of drug to be administered is likely to depend on such variables as the type and extent of progression of the disease or disorder, the overall health status of the particular patient, the relative biological efficacy of the compound selected, the formulation of the compound, the route of administration (intravenous, intramuscular, or other), the pharmacokinetic properties of the conjugates by the chosen delivery route, and the speed (bolus or continuous infusion) and schedule of administrations (number of repetitions in a given period of time).

The conjugates via the linkers of the present invention are also capable of being administered in unit dose forms, wherein the term "unit dose" means a single dose which is capable of being administered to a patient, and which can be readily handled and packaged, remaining as a physically and chemically stable unit dose comprising either the active conjugate itself, or as a pharmaceutically acceptable composition, as described hereinafter. As such, typical total daily/weekly/biweekly/monthly dose ranges are from 0.01 to 100 mg/kg of body weight. By way of general guidance, unit doses for humans range from 1 mg to 3000 mg per day, or per week, per two weeks (biweekly) or per month. Preferably the unit dose range is from 1 to 500 mg administered one to four times a week, and even more preferably from 1 mg to 100 mg, once a week. Conjugates provided herein can be formulated into pharmaceutical compositions by admixture with one or more pharmaceutically acceptable excipients. Such unit dose compositions may be prepared for use by oral administration, particularly in the form of tablets, simple capsules or soft gel capsules; or intranasal, particularly in the form of powders, nasal drops, or aerosols; or dermally, for example, topically in ointments, creams, lotions, gels or sprays, or via transdermal patches.

DRUGS/CYTOTOXIC AGENTS

Drugs that can be conjugated to a cell-binding molecule in the present invention are small molecule drugs including cytotoxic agents, which can be linked to or after they are modified for linkage to the cell-binding agent. A "small molecule drug" is broadly used herein to refer to an organic, inorganic, or organometallic compound that may have a molecular weight of, for example, 100 to 2500, more suitably from 120 to 1500. Small molecule drugs are well characterized in the art, such as in WO05058367A2, and in U.S. Patent No. 4,956,303, among others and are incorporated in their entirety by reference. The drugs include known drugs and those that may become known drugs.

Drugs that are known include, but not limited to,

1). Chemotherapeutic agents: a). Alkylating agents: such as Nitrogen mustards: chlorambucil, chlornaphazine, cyclophosphamide, dacarbazine, estramustine, ifosfamide, mechlorethamine, mechlorethamine oxide hydrochloride, mannomustine, mitobronitol, melphalan, mitolactol, pipobroman, novembichin, phenesterine, prednimustine, thiotepa, trofosfamide, uracil mustard; CC-1065 (including its adozelesin, carzelesin and bizelesin synthetic analogues); Duocarmycin (including the synthetic analogues, KW-2189 and CBI-TMI); Benzodiazepine dimers (e.g., dimmers of pyrrolobenzodiazepine (PBD) or tomaymycin, indolinobenzodiazepines, imidazobenzothiadiazepines, or oxazolidino-benzodiazepines); Nitrosoureas: (carmustine, lomustine, chlorozotocin, fotemustine, nimustine, ranimustine); Alkylsulphonates: (busulfan, treosulfan, improsulfan and piposulfan); Triazines: (dacarbazine); Platinum containing compounds: (carboplatin, cisplatin, oxaliplatin); aziridines, such as benzodopa, carboquone, meturedopa, and uredopa; ethylenimines and methylamelamines including altretamine, triethylenemel-amine, trietylenephosphoramidate, triethylenethiophosphoramidate and trimethylolomel-amine]; b). Plant Alkaloids: such as Vinca alkaloids: (vincristine, vinblastine, vindesine, vinorelbine, navelbin); Taxoids: (paclitaxel, docetaxol) and their analogs, Maytansinoids (DM1, DM2, DM3, DM4, maytansine and ansamitocins) and their analogs, cryptophycins (particularly cryptophycin 1 and cryptophycin 8); epothilones, eleutherobin, discodermolide, bryostatins, dolostatin, auristatins, tubulysins, cephalostatins; pancratistatin; a sarcodictyin; spongistatin; c). DNA Topoisomerase Inhibitors: such as [Epipodophyllins: (9-aminocamptothecin, camptothecin, crisnatol, daunomycin, etoposide, etoposide phosphate, irinotecan, mitoxantrone, novantrone, retinoic acids (retinols), teniposide, topotecan, 9-nitrocamptothecin (RFS 2000)); mitomycins: (mitomycin C)]; d). Anti-metabolites: such as {[Anti-folate: DHFR inhibitors: (methotrexate, trimetrexate, denopterin, pteropterin, aminopterin (4-aminopteroic acid) or the other folic acid analogues); IMP dehydrogenase Inhibitors: (mycophenolic acid, tiazofurin, ribavirin, EICAR); Ribonucleotide reductase Inhibitors: (hydroxyurea, deferoxamine)]; [Pyrimidine analogs: Uracil analogs: (ancitabine, azacitidine, 6-azauridine, capecitabine (Xeloda), carmofur, cytarabine, dideoxyuridine, doxifluridine, enocitabine, 5-Fluorouracil, floxuridine, ratitrexed (Tomudex)); Cytosine analogs: (cytarabine, cytosine arabinoside, fludarabine); Purine analogs: (azathioprine, fludarabine, mercaptopurine, thiamiprine, thioguanine)]; folic acid replenisher, such as frolinic acid}; e). Hormonal therapies: such as {Receptor antagonists: [Anti-estrogen: (megestrol, raloxifene, tamoxifen); LHRH agonists: (goserelin, leuprolide acetate); Anti-androgens: (bicalutamide, flutamide, calusterone, dromostanolone propionate, epitiofanol, goserelin, leuprolide, mepitiofanol, nilutamide, testolactone, trilostane and other androgens inhibitors)]; Retinoids/Deltoids: [Vitamin D3 analogs:

(CB 1093, EB 1089 KH 1060, cholecalciferol, ergocalciferol); Photodynamic therapies: (verteporfin, phthalocyanine, photosensitizer Pc4, demethoxy-hypocrellin A); Cytokines: (Interferon-alpha, Interferon-gamma, tumor necrosis factor (TNFs), human proteins containing a TNF domain)]; f). Kinase inhibitors, such as BIBW 2992 (anti-EGFR/Erb2), imatinib, gefitinib, 5 pegaptanib, sorafenib, dasatinib, sunitinib, erlotinib, nilotinib, lapatinib, axitinib, pazopanib, vandetanib, E7080 (anti-VEGFR2), mubritinib, ponatinib (AP24534), bafetinib (INNO-406), bosutinib (SKI-606), cabozantinib, vismodegib, iniparib, ruxolitinib, CYT387, axitinib, tivozanib, sorafenib, bevacizumab, cetuximab, Trastuzumab, Ranibizumab, Panitumumab, ispinesib; g). antibiotics, such as the enediyne antibiotics (e.g. calicheamicins, especially calicheamicin γ 1, δ 1, 10 α 1 and β 1, see, e.g., *J. Med. Chem.*, 39 (11), 2103–2117 (1996), *Angew Chem Intl. Ed. Engl.* 33:183-186 (1994); dynemicin, including dynemicin A and deoxydynemicin; esperamicin, kedarcidin, C-1027, maduropeptin, as well as neocarzinostatin chromophore and related chromoprotein enediyne antiobiotic chromomophores), aclacinomysins, actinomycin, authramycin, azaserine, bleomycins, cactinomycin, carabacin, carminomycin, carzinophilin; 15 chromomycins, dactinomycin, daunorubicin, detorubicin, 6-diazo-5-oxo-L-norleucine, doxorubicin, morpholino-doxorubicin, cyanomorpholino-doxorubicin, 2-pyrrolino-doxorubicin and deoxydoxorubicin, epirubicin, esorubicin, idarubicin, marcellomycin, nitomycins, mycophenolic acid, nogalamycin, olivomycins, peplomycin, potfiromycin, puromycin, quelamycin, rodorubicin, streptonigrin, streptozocin, tubercidin, ubenimex, zinostatin, zorubicin; 20 f). Others: such as Polyketides (acetogenins), especially bullatacin and bullatacinone; gemcitabine, epoxomicins (e. g. carfilzomib), bortezomib, thalidomide, lenalidomide, pomalidomide, tosedostat, zybrestat, PLX4032, STA-9090, Stimuvax, allovectin-7, Xegeva, Provenge, Yervoy, Isoprenylation inhibitors (such as Lovastatin), Dopaminergic neurotoxins (such as 1-methyl-4-phenylpyridinium ion), Cell cycle inhibitors (such as staurosporine), 25 Actinomycins (such as Actinomycin D, dactinomycin), Bleomycins (such as bleomycin A2, bleomycin B2, peplomycin), Anthracyclines (such as daunorubicin, doxorubicin (adriamycin), idarubicin, epirubicin, pirarubicin, zorubicin, mtoxantrone, MDR inhibitors (such as verapamil), Ca^{2+} ATPase inhibitors (such as thapsigargin), Histone deacetylase inhibitors (Vorinostat, Romidepsin, Panobinostat, Valproic acid, Mocetinostat (MGCD0103), Belinostat, PCI-24781, 30 Entinostat, SB939, Resminostat, Givinostat, AR-42, CUDC-101, sulforaphane, Trichostatin A); Thapsigargin, Celecoxib, glitazones, epigallocatechin gallate, Disulfiram, Salinosporamide A.; Anti-adrenals, such as aminoglutethimide, mitotane, trilostane; aceglatone; aldophosphamide glycoside; aminolevulinic acid; amsacrine; arabinoside, bestrabucil; bisantrene; edatraxate; defofamine; demecolcine; diaziquone; eflornithine (DFMO), elfomithine; elliptinium acetate,

etoglucid; gallium nitrate; gacytosine, hydroxyurea; ibandronate, lentinan; lonidamine; mitoguazone; mitoxantrone; mopidamol; nitracrine; pentostatin; phenamet; pirarubicin; podophyllinic acid; 2-ethylhydrazide; procarbazine; PSK[®]; razoxane; rhizoxin; sizofiran; spirogermanium; tenuazonic acid; triaziquone; 2, 2',2''-trichlorotriethylamine; trichothecenes (especially T-2 toxin, verrucarins A, roridin A and anguidine); urethane, siRNA, antisense drugs, and a nucleolytic enzyme.

2). An anti-autoimmune disease agent includes, but is not limited to, cyclosporine, cyclosporine A, aminocaproic acid, azathioprine, bromocriptine, chlorambucil, chloroquine, cyclophosphamide, corticosteroids (e.g. amcinonide, betamethasone, budesonide, hydrocortisone, flunisolide, fluticasone propionate, fluocortolone danazol, dexamethasone, Triamcinolone acetone, beclomethasone dipropionate), DHEA, enanercept, hydroxychloroquine, infliximab, meloxicam, methotrexate, mofetil, mycophenolate, prednisone, sirolimus, tacrolimus.

3). An anti-infectious disease agent includes, but is not limited to, a). Aminoglycosides: amikacin, astromycin, gentamicin (netilmicin, sisomicin, isepamicin), hygromycin B, kanamycin (amikacin, arbekacin, bekanamycin, dibekacin, tobramycin), neomycin (framycetin, paromomycin, ribostamycin), netilmicin, spectinomycin, streptomycin, tobramycin, verdamicin; b). Amphenicols: azidamphenicol, chloramphenicol, florfenicol, thiamphenicol; c). Ansamycins: geldanamycin, herbimycin; d). Carbapenems: biapenem, doripenem, ertapenem, imipenem/cilastatin, meropenem, panipenem; e). Cephems: carbacephem (loracarbef), cefacetrile, cefaclor, cefradine, cefadroxil, cefalonium, cefaloridine, cefalotin or cefalothin, cefalexin, cefaloglycin, cefamandole, cefapirin, cefatrizine, cefazaflur, cefazedone, cefazolin, cefbuperazone, cefcapene, cefdaloxime, cefepime, cefminox, cefoxitin, cefprozil, cefroxadine, ceftazidime, cefuroxime, cefixime, cefdinir, cefditoren, cefepime, cefetamet, cefmenoxime, cefodizime, cefonicid, cefoperazone, ceforanide, cefotaxime, cefotiam, cefozopran, cephalixin, cefpimizole, cefpiramide, cefpirome, cefpodoxime, cefprozil, cefquinome, cefsulodin, ceftazidime, ceftazidime, ceftibuten, ceftiole, ceftizoxime, ceftobiprole, ceftriaxone, cefuroxime, cefuzonam, cephamycin (cefoxitin, cefotetan, cefmetazole), oxacephem (flomoxef, latamoxef); f). Glycopeptides: bleomycin, vancomycin (oritavancin, telavancin), teicoplanin (dalbavancin), ramoplanin; g). Glycylcyclines: e. g. tigecycline; g). β -Lactamase inhibitors: penam (sulbactam, tazobactam), clavam (clavulanic acid); i). Lincosamides: clindamycin, lincomycin; j). Lipopeptides: daptomycin, A54145, calcium-dependent antibiotics (CDA); k). Macrolides: azithromycin, cethromycin, clarithromycin, dirithromycin, erythromycin, flurithromycin, josamycin, ketolide (telithromycin, cethromycin), midecamycin, miocamycin, oleandomycin, rifamycins (rifampicin, rifampin, rifabutin, rifapentine), rokitamycin, roxithromycin,

spectinomycin, spiramycin, tacrolimus (FK506), troleandomycin, telithromycin; l).

Monobactams: aztreonam, tigemonam; m). Oxazolidinones: linezolid; n). Penicillins: amoxicillin, ampicillin (pivampicillin, hetacillin, bacampicillin, metampicillin, talampicillin), azidocillin, azlocillin, benzylpenicillin, benzathine benzylpenicillin, benzathine phenoxymethyl-penicillin, clometocillin, procaine benzylpenicillin, carbenicillin (carindacillin), cloxacillin, dicloxacillin, epicillin, flucloxacillin, mecillinam (pivmecillinam), mezlocillin, meticillin, nafcillin, oxacillin, penamecillin, penicillin, pheneticillin, phenoxymethylpenicillin, piperacillin, propicillin, sulbenicillin, temocillin, ticarcillin; o). Polypeptides: bacitracin, colistin, polymyxin B; p).

Quinolones: alatrofloxacin, balofloxacin, ciprofloxacin, clinafloxacin, danofloxacin, difloxacin, enoxacin, enrofloxacin, floxin, garenoxacin, gatifloxacin, gemifloxacin, grepafloxacin, kano trovafloxacin, levofloxacin, lomefloxacin, marbofloxacin, moxifloxacin, nadifloxacin, norfloxacin, orbifloxacin, ofloxacin, pefloxacin, trovafloxacin, grepafloxacin, sitafloxacin, sparfloxacin, temafloxacin, tosufloxacin, trovafloxacin; q). Streptogramins: pristinamycin, quinupristin/dalfopristin; r). Sulfonamides: mafenide, prontosil, sulfacetamide, sulfamethizole, sulfanilimide, sulfasalazine, sulfisoxazole, trimethoprim, trimethoprim-sulfamethoxazole (cotrimoxazole); s). Steroid antibacterials: e.g. fusidic acid; t). Tetracyclines: doxycycline, chlortetracycline, clomocycline, demeclocycline, lymecycline, meclocycline, metacycline, minocycline, oxytetracycline, penimepicycline, rolitetracycline, tetracycline, glycylicyclines (e.g. tigecycline); u). Other types of antibiotics: annonacin, arspenamine, bactoprenol inhibitors (Bacitracin), DADAL/AR inhibitors (cycloserine), dictyostatin, discodermolide, eleutherobin, epothilone, ethambutol, etoposide, faropenem, fusidic acid, furazolidone, isoniazid, laulimalide, metronidazole, mupirocin, mycolactone, NAM synthesis inhibitors (e. g. fosfomycin), nitrofurantoin, paclitaxel, platensimycin, pyrazinamide, quinupristin/dalfopristin, rifampicin (rifampin), tazobactam tinidazole, uvaricin;

4). Anti-viral drugs: a). Entry/fusion inhibitors: aplaviroc, maraviroc, vicriviroc, gp41 (enfuvirtide), PRO 140, CD4 (ibalizumab); b). Integrase inhibitors: raltegravir, elvitegravir, globoidnan A; c). Maturation inhibitors: bevirimat, vivecon; d). Neuraminidase inhibitors: oseltamivir, zanamivir, peramivir; e). Nucleosides & nucleotides: abacavir, aciclovir, adefovir, amdoxovir, apricitabine, brivudine, cidofovir, clevudine, dexelvucitabine, didanosine (ddI), elvucitabine, emtricitabine (FTC), entecavir, famciclovir, fluorouracil (5-FU), 3'-fluoro-substituted 2', 3'-dideoxynucleoside analogues (e.g. 3'-fluoro-2',3'-dideoxythymidine (FLT) and 3'-fluoro-2',3'-dideoxyguanosine (FLG), fomivirsen, ganciclovir, idoxuridine, lamivudine (3TC), l-nucleosides (e.g. β -l-thymidine and β -l-2'-deoxycytidine), penciclovir, racivir, ribavirin, stampidine, stavudine (d4T), taribavirin (viramidine), telbivudine, tenofovir, trifluridine

valaciclovir, valganciclovir, zalcitabine (ddC), zidovudine (AZT); f). Non-nucleosides: amantadine, ateviridine, capravirine, diarylpyrimidines (etravirine, rilpivirine), delavirdine, docosanol, emivirine, efavirenz, foscarnet (phosphonoformic acid), imiquimod, interferon alfa, loviride, lodenosine, methisazone, nevirapine, NOV-205, peginterferon alfa, podophyllotoxin, rifampicin, rimantadine, resiquimod (R-848), tromantadine; g). Protease inhibitors: amprenavir, atazanavir, boceprevir, darunavir, fosamprenavir, indinavir, lopinavir, nelfinavir, pleconaril, ritonavir, saquinavir, telaprevir (VX-950), tipranavir; h). Other types of anti-virus drugs: abzyme, arbidol, calanolide A, ceragenin, cyanovirin-n, diarylpyrimidines, epigallocatechin gallate (EGCG), foscarnet, griffithsin, taribavirin (viramidine), hydroxyurea, KP-1461, miltefosine, pleconaril, portmanteau inhibitors, ribavirin, seliciclib.

5) The drugs used for conjugates via a bridge linker of the present invention also include radioisotopes. Examples of radioisotopes (radionuclides) are ^3H , ^{11}C , ^{14}C , ^{18}F , ^{32}P , ^{35}S , ^{64}Cu , ^{68}Ga , ^{86}Y , ^{99}Tc , ^{111}In , ^{123}I , ^{124}I , ^{125}I , ^{131}I , ^{133}Xe , ^{177}Lu , ^{211}At , or ^{213}Bi . Radioisotope labeled antibodies are useful in receptor targeted imaging experiments or can be for targeted treatment such as with the antibody-drug conjugates of the invention (Wu et al (2005) Nature Biotechnology 23(9): 1137-46). The cell binding molecules, e.g. an antibody can be labeled with ligand reagents through the bridge linkers of the present patent that bind, chelate or otherwise complex a radioisotope metal, using the techniques described in Current Protocols in Immunology, Volumes 1 and 2, Coligen et al, Ed. Wiley-Interscience, New York, Pubs. (1991). Chelating ligands which may complex a metal ion include DOTA, DOTP, DOTMA, DTPA and TETA (Macrocyclics, Dallas, Tex. USA).

6). The pharmaceutically acceptable salts, acids or derivatives of any of the above drugs.

In another embodiment, the drug in the Formula (II) and/or (IV) can be a chromophore molecule, for which the conjugate can be used for detection, monitoring, or study the interaction of the cell binding molecule with a target cell. Chromophore molecules are a compound that have the ability to absorb a kind of light, such as UV light, florescent light, IR light, near IR light, visual light; A chromatophore molecule includes a class or subclass of xanthophores, erythrophores, iridophores, leucophores, melanophores, and cyanophores; a class or subclass of fluorophore molecules which are fluorescent chemical compounds re-emitting light upon light; a class or subclass of visual phototransduction molecules; a class or subclass of photophore molecules; a class or subclass of luminescence molecules; and a class or subclass of luciferin compounds.

The chromophore molecule can be selected from, but not limited, non-protein organic fluorophores, such as: Xanthene derivatives (fluorescein, rhodamine, Oregon green, eosin, and Texas red); Cyanine derivatives: (cyanine, indocarbocyanine, oxacarbocyanine, thiocarbocyanine,

and merocyanine); Squaraine derivatives and ring-substituted squaraines, including Seta, SeTau, and Square dyes; Naphthalene derivatives (dansyl and prodan derivatives); Coumarin derivatives; Oxadiazole derivatives (pyridyloxazole, nitroben-zoxadiazole and benzoxadiazole); Anthracene derivatives (anthraquinones, including DRAQ5, DRAQ7 and CyTRAK Orange); Pyrene derivatives (cascade blue, etc); Oxazine derivatives (Nile red, Nile blue, cresyl violet, oxazine 170 etc). Acridine derivatives (proflavin, acridine orange, acridine yellow etc). Arylmethine derivatives (auramine, crystal violet, malachite green). Tetrapyrrole derivatives (porphin, phthalocyanine, bilirubin).



Or a chromophore molecule can be selected from any analogs and derivatives of the following fluorophore compounds: CF dye (Biotium), DRAQ and CyTRAK probes (BioStatus), BODIPY (Invitrogen), Alexa Fluor (Invitrogen), DyLight Fluor (Thermo Scientific, Pierce), Atto and Tracy (Sigma Aldrich), FluoProbes (Interchim), Abberior Dyes (Abberior), DY and MegaStokes Dyes (Dyomics), Sulfo Cy dyes (Cyandye), HiLyte Fluor (AnaSpec), Seta, SeTau and Square Dyes (SETA BioMedicals), Quasar and Cal Fluor dyes (Biosearch Technologies), SureLight Dyes (APC, RPEPerCP, Phycobilisomes)(Columbia Biosciences), APC, APCXL, RPE, BPE (Phyco-Biotech).

Examples of the widely used fluorophore compounds which are reactive or conjugatable with the linkers of the invention are: Allophycocyanin (APC), Aminocoumarin, APC-Cy7 conjugates, BODIPY-FL, Cascade Blue, Cy2, Cy3, Cy3.5, Cy3B, Cy5, Cy5.5, Cy7, Fluorescein, FluorX, Hydroxycoumarin, IR-783, Lissamine Rhodamine B, Lucifer yellow, Methoxycoumarin, NBD, Pacific Blue, Pacific Orange, PE-Cy5 conjugates, PE-Cy7 conjugates, PerCP, R-Phycoerythrin (PE), Red 613, Seta-555-Azide, Seta-555-DBCO, Seta-555-NHS, Seta-580-NHS, Seta-680-NHS, Seta-780-NHS, Seta-APC-780, Seta-PerCP-680, Seta-R-PE-670, SeTau-380-NHS, SeTau-405-Maleimide, SeTau-405-NHS, SeTau-425-NHS, SeTau-647-NHS, Texas Red, TRITC, TruRed, X-Rhodamine.

The fluorophore compounds that can be linked to the linkers of the invention for study of nucleic acids or proteins are selected from the following compounds or their derivatives: 7-AAD (7-aminoactinomycin D, CG-selective), Acridine Orange, Chromomycin A3, CyTRAK Orange (Biostatus, red excitation dark), DAPI, DRAQ5, DRAQ7, Ethidium Bromide, Hoechst33258, Hoechst33342, LDS 751, Mithramycin, Propidium Iodide (PI), SYTOX Blue, SYTOX Green, SYTOX Orange, Thiazole Orange, TO-PRO: Cyanine Monomer, TOTO-1, TO-PRO-1, TOTO-3, TO-PRO-3, YOSeta-1, YOYO-1. The fluorophore compounds that can be linked to the linkers of the invention for study cells are selected from the following compounds or their derivatives: DCFH (2',7'-Dichlorodihydro-fluorescein, oxidized form), DHR (Dihydrorhodamine 123, oxidized


form, light catalyzes oxidation), Fluo-3 (AM ester. pH > 6), Fluo-4 (AM ester. pH 7.2), Indo-1 (AM ester, low/high calcium (Ca²⁺)), and SNARF (pH 6/9). The preferred fluorophore compounds that can be linked to the linkers of the invention for study proteins/antibodies are selected from the following compounds or their derivatives: Allophycocyanin (APC), AmCyan1 (tetramer, Clontech), AsRed2 (tetramer, Clontech), Azami Green (monomer, MBL), Azurite, B-phycoerythrin (BPE), Cerulean, CyPet, DsRed monomer (Clontech), DsRed2 ("RFP", Clontech), EBFP, EBFP2, ECFP, EGFP (weak dimer, Clontech), Emerald (weak dimer, Invitrogen), EYFP (weak dimer, Clontech), GFP (S65A mutation), GFP (S65C mutation), GFP (S65L mutation), GFP (S65T mutation), GFP (Y66F mutation), GFP (Y66H mutation), GFP (Y66W mutation), GFPuv, HcRed1, J-Red, Katusha, Kusabira Orange (monomer, MBL), mCFP, mCherry, mCitrine, Midoriishi Cyan (dimer, MBL), mKate (TagFP635, monomer, Evrogen), mKeima-Red (monomer, MBL), mKO, mOrange, mPlum, mRaspberry, mRFP1 (monomer, Tsien lab), mStrawberry, mTFP1, mTurquoise2, P3 (phycobilisome complex), Peridinin Chlorophyll (PerCP), R-phycoerythrin(RPE), T-Sapphire, TagCFP (dimer, Evrogen), TagGFP (dimer, Evrogen), TagRFP (dimer, Evrogen), TagYFP (dimer, Evrogen), tdTomato (tandem dimer), Topaz, TurboFP602 (dimer, Evrogen), TurboFP635 (dimer, Evrogen), TurboGFP (dimer, Evrogen), TurboRFP (dimer, Evrogen), TurboYFP (dimer, Evrogen), Venus, Wild Type GFP, YPet, ZsGreen1 (tetramer, Clontech), ZsYellow1 (tetramer, Clontech).

The examples of the structure of the conjugates of the antibody-chromophore molecules via the bridge linker are as following Ac01, Ac02, Ac03, Ac04 and Ac05:



monophenyl glyoxals, hydroxyl PEG-monothiazolidine-2-thiones, hydroxyl PEG-monothioesters, hydroxyl PEG-monothiols, hydroxyl PEG-monotriazines and hydroxyl PEG-monovinylsulfones).

In certain such embodiments, the polyalkylene glycol has a molecular weight of from about 10 Daltons to about 200 kDa, preferably about 88 Da to about 40 kDa; two branches each with a molecular weight of about 88 Da to about 40 kDa; and more preferably two branches, each of about 88 Da to about 20 kDa. In one particular embodiment, the polyalkylene glycol is poly(ethylene) glycol and has a molecular weight of about 10 kDa; about 20 kDa, or about 40 kDa. In specific embodiments, the PEG is a PEG 10 kDa (linear or branched), a PEG 20 kDa (linear or branched), or a PEG 40 kDa (linear or branched). A number of US patents have disclosed the preparation of linear or branched "non-antigenic" PEG polymers and derivatives or conjugates thereof, e.g., U.S. Pat. Nos. 5,428,128; 5,621,039; 5,622,986; 5,643,575; 5,728,560; 5,730,990; 5,738,846; 5,811,076; 5,824,701; 5,840,900; 5,880,131; 5,900,402; 5,902,588; 5,919,455; 5,951,974; 5,965,119; 5,965,566; 5,969,040; 5,981,709; 6,011,042; 6,042,822; 6,113,906; 6,127,355; 6,132,713; 6,177,087, and 6,180,095. The structure of the conjugates of the antibody-polyalkylene glycols via the bridge linker is as following Pg01:










Wherein mAb is an antibody; R' is H or CH₃; m₃ is an integer from 1 to 5000; R₃ is OH, H, or R₁; "—" represents either single bond or double bond; m₁, m₂, n, "—", X₁, X₂, R₁, and R₂ are the same defined in Formula (I) and (II). In addition, R₁ can be absent.

In yet another embodiment, the preferred cytotoxic agents that conjugated to a cell-binding molecule via a bridge linker of this patent are tubulysins, maytansinoids, taxanoids (taxanes), CC-1065 analogs, daunorubicin and doxorubicin compounds, amatoxins, benzodiazepine dimers (e.g., dimers of pyrrolbenzodiazepine (PBD), tomaymycin, anthramycin, indolinbenzodiazepines, imidazobenzothiadiazepines, or oxazolidino-benzodiazepines), calicheamicins and the enediyne antibiotics, actinomycin, azaserines, bleomycins, epirubicin, tamoxifen, idarubicin, dolastatins, auristatins (e.g. monomethyl auristatin E, MMAE, MMAF, auristatin PYE, auristatin TP, Auristatins 2-AQ, 6-AQ, EB (AEB), and EFP (AEFP)), duocarmycins, geldanamycins, methotrexates, thiotepa, vindesines, vincristines, hemiasterlins, nazumamides, microginins, radiosumins, alterobactins, microsclerodermins, theonellamides, esperamicins, PNU-159682, and their analogues and derivatives above thereof.

Tubulysins that are preferred for conjugation in the present invention are well known in the art and can be isolated from natural sources according to known methods or prepared synthetically according to known methods (e. g. Balasubramanian, R., et al. J. Med. Chem., 2009, 52, 238–40; Wipf, P., et al. Org. Lett., 2004, 6, 4057–60; Pando, O., et al. J. Am. Chem. Soc., 2011, 133, 7692–5; Reddy, J. A., et al. Mol. Pharmaceutics, 2009, 6, 1518–25; Raghavan, B., et al. J. Med. Chem., 2008, 51, 1530–33; Patterson, A. W., et al. J. Org. Chem., 2008, 73, 4362–9; Pando, O., et al. Org. Lett., 2009, 11 (24), 5567–9; Wipf, P., et al. Org. Lett., 2007, 9 (8), 1605–7; Friestad, G. K., Org. Lett., 2004, 6, 3249–52; Peltier, H. M., et al. J. Am. Chem. Soc., 2006, 128, 16018–9; Chandrasekhar, S., et al J. Org. Chem., 2009, 74, 9531–4; Liu, Y., et al. Mol. Pharmaceutics, 2012, 9, 168–75; Friestad, G. K., et al. Org. Lett., 2009, 11, 1095–8; Kubicek, K., et al., Angew Chem Int Ed Engl, 2010.49: 4809-12; Chai, Y., et al., Chem Biol, 2010, 17: 296-309; Ullrich, A., et al., Angew Chem Int Ed Engl, 2009, 48, 4422-5; Sani, M., et al. Angew Chem Int Ed Engl, 2007, 46, 3526-9; Domling, A., et al., Angew Chem Int Ed Engl, 2006, 45, 7235-9; Patent applications: Zanda, M., et al, Can. Pat. Appl. CA 2710693 (2011); Chai, Y., et al. Eur. Pat.Appl. 2174947 (2010), WO 2010034724; Leamon, C. et al, WO2010033733, WO 2009002993; Ellman, J., et al, PCT WO2009134279; WO 2009012958, US appl. 20110263650, 20110021568; Matschiner, G., et al, WO2009095447; Vlahov, I., et al, WO2009055562, WO 2008112873; Low, P., et al, WO2009026177; Richter, W., WO2008138561; Kjems, J., et al, WO 2008125116; Davis, M.; et al, WO2008076333; Diener, J.; et al, U.S. Pat.Appl. 20070041901, WO2006096754; Matschiner, G., et al, WO2006056464; Vaghefi, F., et al, WO2006033913; Doemling, A., Ger. Offen. DE102004030227, WO2004005327, WO2004005326, WO2004005269; Stanton, M., et al, U.S. Pat. Appl. Publ. 20040249130; Hoefle, G., et al, Ger. Offen. DE10254439, DE10241152, DE10008089; Leung, D., et al, WO2002077036; Reichenbach, H., et al, Ger. Offen. DE19638870; Wolfgang, R., US20120129779; Chen, H., US appl. 20110027274. The preferred structures of tubulysins for conjugation of cell binding molecules are described in the patent application of PCT/IB2012/053554.

Examples of the structures of the conjugates of the antibody-tubulysin analogs via the bridge linker are T01, T02, T03, T04, T05, T06 and T07 as following:






Wherein mAb is an antibody; Z₃ and Z'₃ are independently H, OP(O)(OM₁)(OM₂), OCH₂OP(O)(OM₁)(OM₂), OSO₃M₁, R₁, or O-glycoside (glucoside, galactoside, mannoside, glucuronoside, alloside, fructoside, etc), NH-glycoside, S-glycoside or CH₂-glycoside; M₁ and


M₂ are independently H, Na, K, Ca, Mg, NH₄, NR₁R₂R₃; “====” represents either single bond or double bond; n, m₁, m₂, “—”, X₁, X₂, R₁, R₂ and R₃ are the same defined in Formula (I) and (II).

Calicheamicins and their related enediyne antibiotics that are preferred for cell-binding molecule-drug conjugates of this patent are described in: Nicolaou, K. C. et al, Science 1992, 256, 1172-1178; Proc. Natl. Acad. Sci USA. 1993, 90, 5881-8), U.S. Patent Nos. 4,970,198; 5,053,394; 5,108,912; 5,264,586; 5,384,412; 5,606,040; 5,712,374; 5,714,586; 5,739,116; 5,770,701; 5,770,710; 5,773,001; 5,877,296; 6,015,562; 6,124,310; 8,153,768. An Example of the structure of the conjugate of the antibody-Calicheamicin analog via the bridge linker is C01 as the following:



Wherein mAb is an antibody; “====” represents either single bond or double bond; n, m₁, m₂, “—”, X₁, X₂, R₁, and R₂ are the same defined in Formula (I) and (II).




Maytansinoids that are preferred to be used in the present invention including maytansinol and its analogues are described in U.S. Patent Nos. 4,256,746, 4,361,650, 4,307,016, 4,294,757, 4,294,757, 4,371,533, 4,424,219, 4,331,598, 4,450,254, 4,364,866, 4,313,946, 4,315,929, 4,362,663, 4,322,348, 4,371,533, 4,424,219, 5,208,020, 5,416,064, 5,208,020; 5,416,064; 6,333,410; 6,441,163; 6,716,821, 7,276,497, 7,301,019, 7,303,749, 7,368,565, 7,411,063, 7,851,432, and 8,163,888. An example of the structure of the conjugate of the antibody-Maytansinoids via the bridge linker is as the following My01:



Wherein mAb is an antibody; n, m₁, m₂, “====” represents either single bond or double bond; n, m₁, m₂, “—”, X₁, X₂, R₁, and R₂ are the same defined in Formula (I) and (II).

Taxanes, which includes Paclitaxel (Taxol), a cytotoxic natural product, and docetaxel (Taxotere), a semi-synthetic derivative, and their analogs which are preferred for conjugation via the bridge linkers of the present patent are exemplified in: K C. Nicolaou et al., J. Am. Chem. Soc. 117, 2409-20, (1995); Ojima et al, J. Med. Chem. 39:3889-3896 (1996); 40:267-78 (1997); 45, 5620-3 (2002); Ojima et al., Proc. Natl. Acad. Sci., 96:4256-61 (1999); Kim et al., Bull. Korean Chem. Soc., 20, 1389-90 (1999); Miller, et al. J. Med. Chem., 47, 4802-5(2004); U.S. Patent No. 5,475,011 5,728,849, 5,811,452; 6,340,701; 6,372,738; 6,391,913, 6,436,931; 6,589,979; 6,596,757; 6,706,708; 7,008,942; 7,186,851; 7,217,819; 7,276,499; 7,598,290; and 7,667,054.




Examples of the structures of the conjugate of the antibody-taxanes via the bridge linker are as the following Tx01, Tx02 and Tx03.



Wherein mAb is an antibody; n, m₁, m₂, “ $\equiv$ ” represents either single bond or double bond; n, m₁, m₂, “—”, X₁, X₂, R₁, and R₂ are the same defined in Formula (I) and (II).




CC-1065 analogues and doucarmycin analogs are also preferred to be used for a conjugate with the bridge linkers of the present patent. The examples of the CC-1065 analogues and doucarmycin analogs as well as their synthesis are described in: e.g. Warpehoski, et al, J. Med. Chem. 31:590-603 (1988); D. Boger et al., J. Org. Chem; 66; 6654-61, 2001; U. S. Patent Nos: 4169888, 4391904, 4671958, 4816567, 4912227, 4923990, 4952394, 4975278, 4978757,


4994578, 5037993, 5070092, 5084468, 5101038, 5117006, 5137877, 5138059, 5147786,
 5187186, 5223409, 5225539, 5288514, 5324483, 5332740, 5332837, 5334528, 5403484,
 5427908, 5475092, 5495009, 5530101, 5545806, 5547667, 5569825, 5571698, 5573922,
 5580717, 5585089, 5585499, 5587161, 5595499, 5606017, 5622929, 5625126, 5629430,
 5633425, 5641780, 5660829, 5661016, 5686237, 5693762, 5703080, 5712374, 5714586,
 5739116, 5739350, 5770429, 5773001, 5773435, 5786377, 5786486, 5789650, 5814318,
 5846545, 5874299, 5877296, 5877397, 5885793, 5939598, 5962216, 5969108, 5985908,
 6060608, 6066742, 6075181, 6103236, 6114598, 6130237, 6132722, 6143901, 6150584,
 6162963, 6172197, 6180370, 6194612, 6214345, 6262271, 6281354, 6310209, 6329497,
 6342480, 6486326, 6512101, 6521404, 6534660, 6544731, 6548530, 6555313, 6555693,
 6566336, 6,586,618, 6593081, 6630579, 6,756,397, 6759509, 6762179, 6884869, 6897034,
 6946455, 7,049,316, 7087600, 7091186, 7115573, 7129261, 7214663, 7223837, 7304032,
 7329507, 7,329,760, 7,388,026, 7,655,660, 7,655,661, 7,906,545, and 8,012,978. Examples of
 the structures of the conjugate of the antibody-CC-1065 analogs via the bridge linker are as the
 following CC01, CC02, and CC03.



Wherein mAb is an antibody; Z₄ and Z'₄ are independently H, PO(OM₁)(OM₂), SO₃M₁,
 CH₂PO(OM₁)(OM₂), CH₃N(CH₂CH₂)₂NC(O)-, O(CH₂CH₂)₂NC(O)-, R₁, or glycoside; X₃ is O,
 NH, NHC(O), OC(O), -C(O)O, R₁, or absent; “==” represents either single bond or double bond;
 n, m₁, m₂, “—”, X₁, X₂, R₁, R₂, M₁, and M₂ are the same defined in Formula (I) and (II).



Daunorubicin/Doxorubicin Analogues are also preferred for conjugation via the bridge linkers of the present patent. The preferred structures and their synthesis are exemplified in: Hurwitz, E., et al., Cancer Res. 35, 1175-81 (1975). Yang, H. M., and Reisfeld, R. A., Proc. Natl. Acad. Sci. 85, 1189-93 (1988); Pietersz, C. A., E., et al., E., et al., " Cancer Res. 48, 926-311 (1988); Trouet, et al., 79, 626-29 (1982); Z. Brich et al., J. Controlled Release, 19, 245-58 (1992); Chen et al., Syn. Comm., 33, 2377-90, 2003; King et al., Bioconj. Chem., 10, 279-88, 1999; King et al., J. Med. Chem., 45, 4336-43, 2002; Kratz et al., J Med Chem. 45, 5523-33, 2002; Kratz et al., Biol Pharm Bull. Jan. 21, 56-61, 1998; Lau et al., Bioorg. Med. Chem. 3, 1305-12, 1995; Scott et al., Bioorg. Med. Chem. Lett. 6, 1491-6, 1996; Watanabe et al., Tokai J. Experimental Clin. Med. 15, 327-34, 1990; Zhou et al., J. Am. Chem. Soc. 126, 15656-7, 2004; WO 01/38318; U.S. Patent Nos. 5,106,951; 5,122,368; 5,146,064; 5,177,016; 5,208,323; 5,824,805; 6,146,658; 6,214,345; 7,569,358; 7,803,903; 8,084,586; 8,053,205. Examples of the structures of the conjugate of the antibody-CC-1065 analogs via the bridge linker are as the following Da01, Da02, Da03 and Da04.








Wherein mAb is an antibody; X_3 is O, NH, $NHC(O)$, $NHC(O)NH$, $C(O)$, R_1 , $OC(O)$, or absent; “—” represents either single bond or double bond; n , m_1 , m_2 , “—”, X_1 , X_2 , R_1 , and R_2 are the same defined in Formula (I) and (II).








5 Auristatins and dolastatins are preferred in conjugation via the bridge linkers of this patent. The auristatins (e. g. auristatin E (AE) auristatin EB (AEB), auristatin EFP (AEFP), monomethyl auristatin E (MMAE), Monomethylauristatin (MMAF), Auristatin F phenylene diamine (AFP) and a phenylalanine variant of MMAE) which are synthetic analogs of dolastatins, are described in Int. J. Oncol. 15: 367-72 (1999); Molecular Cancer Therapeutics, vol. 3, No. 8, pp. 921-32
 10 (2004); U.S. Application Nos. 11134826, 20060074008, 2006022925. U.S. Patent Nos. 4414205, 4753894, 4764368, 4816444, 4879278, 4943628, 4978744, 5122368, 5165923, 5169774, 5286637, 5410024, 5521284, 5530097, 5554725, 5585089, 5599902, 5629197, 5635483, 5654399, 5663149, 5665860, 5708146, 5714586, 5741892, 5767236, 5767237, 5780588, 5821337, 5840699, 5965537, 6004934, 6033876, 6034065, 6048720, 6054297, 6054561, 6124431, 6143721, 6162930, 6214345, 6239104, 6323315, 6342219, 6342221, 6407213, 6569834, 6620911, 6639055, 6884869, 6913748, 7090843, 7091186, 7097840, 7098305, 7098308, 7498298, 7375078, 7462352, 7553816, 7659241, 7662387, 7745394, 7754681, 7829531, 7837980, 7837995, 7902338, 7964566, 7964567, 7851437, 7994135. Examples of the structures of the conjugate of the antibody-auristatins via the bridge linker are as the following
 15 Au01, Au02, Au03, Au04, and Au05.









Wherein mAb is an antibody; X₃ is CH₂, O, NH, NHC(O), NHC(O)NH, C(O), OC(O) R₁, or absent; X₄ is CH₂, C(O), C(O)NH, C(O)N(R₁), R₁, NHR₁, NR₁C(O)R₁ or C(O)O; Z₃ is H, R₁, OP(O)(OM₁)(OM₂), NHR₁, OCH₂OP(O)(OM₁)(OM₂), OSO₃M₁, or O-glycoside (glucoside, galactoside, mannoside, glucuronoside, alloside, fructoside), NH-glycoside, S-glycoside, or CH₂-glycoside; M₁ and M₂ are independently H, Na, K, Ca, Mg, NH₄, NR₁R₂R₃; “ $\equiv$ ” represents either single bond or double bond; n, m₁, m₂, “—”, X₁, X₂, R₁, R₂ and R₃ are the same defined in Formula (I) and (II). In addition, R₁ can be absent and R₂ can be H.

The benzodiazepine dimers (e. g. dimmers of pyrrolobenzodiazepine (PBD) or (tomaymycin), indolinobenzodiazepines, imidazobenzothiadiazepines, or oxazolidinobenzodiazepines) which are preferred cytotoxic agents according to the present invention are exemplified in the art: US Patent Nos . 8,163,736; 8,153,627; 8,034,808; 7,834,005; 7,741,319; 7,704,924; 7,691,848; 7,678,787; 7,612,062; 7,608,615; 7,557,099; 7,528,128; 7,528,126; 7,511,032; 7,429,658; 7,407,951; 7,326,700; 7,312,210; 7,265,105; 7,202,239; 7,189,710; 7,173,026; 7,109,193; 7,067,511; 7,064,120; 7,056,913; 7,049,311; 7,022,699; 7,015,215; 6,979,684; 6,951,853; 6,884,799; 6,800,622; 6,747,144; 6,660,856; 6,608,192; 6,562,806; 6,977,254; 6,951,853; 6,909,006; 6,344,451; 5,880,122; 4,935,362; 4,764,616; 4,761,412; 4,723,007; 4,723,003; 4,683,230; 4,663,453; 4,508,647; 4,464,467; 4,427,587; 4,000,304; US patent appl. 20100203007, 20100316656, 20030195196. Examples of the structures of the conjugate of the antibody- benzodiazepine dimers via the bridge linker are as the following PB01, PB02, PB03, PB04, PB05, PB06, PB07, PB08, PB09, PB10 and PB11.







5 Wherein mAb is an antibody; X₃ is CH₂, O, NH, NHC(O), NHC(O)NH, C(O), OC(O), OC(O)(NR₃), R₁NHR₁, NR₁C(O)R₁ or absent; X₄ is CH₂, C(O), C(O)NH, C(O)N(R₁), R₁NHR₁, NR₁C(O)R₁ or C(O)O; M₁ and M₂ are independently H, Na, K, Ca, Mg, NH₄, NR₁R₂R₃; “====” represents either single bond or double bond; n, m₁, m₂, “—”, X₁, X₂, R₁, R₂ and R₃ are the same defined in Formula (I) and (II). In addition, R₁ can be absent.



10 Amatoxins which are a subgroup of at least ten toxic compounds originally found in several genera of poisonous mushrooms, most notably *Amanita phalloides* and several other mushroom species, are also preferred for conjugation via the bridge linkers of the present patent. These ten amatoxins, named α-Amanitin, β-Amanitin, γ-Amanitin, ε-Amanitin, Amanullin, Amanullinic acid, Amaninamide, Amanin, Proamanullin, are rigid bicyclic peptides

15 that are synthesized as 35-amino-acid proproteins, from which the final eight amino acids are cleaved by a prolyl oligopeptidase (Litten, W. 1975 Scientific American 232 (3): 90–101; H. E. Hallen, et al 2007 Proc. Nat. Aca. Sci. USA 104, 19097–101; K. Baumann, et al, 1993

Biochemistry 32 (15): 4043–50; Karlson-Stiber C, Persson H. 2003, Toxicon 42 (4): 339–49; Horgen, P. A. et al. 1978 Arch. Microbio. 118 (3): 317–9). Amatoxins kill cells by inhibiting RNA polymerase II (Pol II), shutting down gene transcription and protein biosynthesis (Brodner, O. G. and Wieland, T. 1976 Biochemistry, 15(16): 3480–4; Fiume, L., Curr Probl Clin Biochem, 1977, 7: 23-8; Karlson-Stiber C, Persson H. 2003, Toxicon 42(4): 339–49; Chafin, D. R. , Guo, H. & Price, D. H. 1995 J. Biol. Chem. 270 (32): 19114–19; Wieland (1983) Int. J. Pept. Protein Res. 22(3): 257-76.). Amatoxins can be produced from collected Amanita phalloides mushrooms (Yocum, R. R. 1978 Biochemistry 17(18): 3786-9; Zhang, P. et al, 2005, FEMS Microbiol. Lett. 252(2), 223-8), or from fermentation using a basidiomycete (Muraoka, S. and Shinozawa T., 2000 J. Biosci. Bioeng. 89(1): 73-6) or from fermentation using A. fissa (Guo, X. W., et al, 2006 Wei Sheng Wu Xue Bao 46(3): 373-8), or from culturing Galerina fasciculata or Galerina helvoliceps, a strain belonging to the genus (WO/1990/009799, JP11137291). However the yields from these isolation and fermentation were quite low (less than 5 mg/L culture). Several preparations of amatoxins and their analogs have been reported in the past three decades (W. E. Savige, A. Fontana, Chem. Commun. 1976, 600–1; Zanotti, G., et al, Int J Pept Protein Res, 1981. 18(2): 162-8; Wieland, T., et al, Eur. J. Biochem. 1981, 117, 161–4; P. A. Bartlett, et al, Tetrahedron Lett. 1982, 23, 619–22; Zanotti, G., et al., Biochim Biophys Acta, 1986. 870(3): 454-62; Zanotti, G., et al., Int. J. Peptide Protein Res. 1987, 30, 323–9; Zanotti, G., et al., Int. J. Peptide Protein Res. 1987, 30, 450–9; Zanotti, G., et al., Int J Pept Protein Res, 1988. 32(1): 9-20; G. Zanotti, T. et al, Int. J. Peptide Protein Res. 1989, 34, 222–8; Zanotti, G., et al., Int J Pept Protein Res, 1990. 35(3): 263-70; Mullersman, J. E. and J. F. Preston, 3rd, Int J Pept Protein Res, 1991. 37(6): 544-51; Mullersman, J.E., et al, Int J Pept Protein Res, 1991. 38(5): 409-16; Zanotti, G., et al, Int J Pept Protein Res, 1992. 40(6): 551-8; Schmitt, W. et al, J. Am. Chem. Soc. 1996, 118, 4380–7; Anderson, M.O., et al, J. Org. Chem., 2005, 70(12): 4578-84; J. P. May, et al, J. Org. Chem. 2005, 70, 8424–30; F. Brueckner, P. Cramer, Nat. Struct. Mol. Biol. 2008, 15, 811–8; J. P. May, D. M. Perrin, Chem. Eur. J. 2008, 14, 3404–9; J. P. May, et al, Chem. Eur. J. 2008, 14, 3410–17; Q. Wang, et al, Eur. J. Org. Chem. 2002, 834–9; May, J. P. and D. M. Perrin, Biopolymers, 2007. 88(5): 714-24; May, J.P., et al., Chemistry, 2008. 14(11): 3410-7; S. De Lamo Marin, et al, Eur. J. Org. Chem. 2010, 3985–9; Pousse, G., et al., Org Lett, 2010. 12(16): 3582-5; Luo, H., et al., Chem Biol, 2014. 21(12): 1610-7; Zhao, L., et al., Chembiochem, 2015. 16(10): 1420-5) and most of these preparations were by partial synthesis. Because of their extreme potency and unique mechanism of cytotoxicity, amatoxins have been used as payloads for conjugations (Fiume, L., Lancet, 1969. 2 (7625): 853-4; Barbanti-Brodano, G. and L. Fiume, Nat New Biol, 1973. 243(130):






281-3; Bonetti, E., M. et al, Arch Toxicol, 1976. 35(1): p. 69-73; Davis, M. T., Preston, J. F. Science 1981, 213, 1385–1388; Preston, J.F., et al, Arch Biochem Biophys, 1981. 209(1): 63–71; H. Faulstich, et al, Biochemistry 1981, 20, 6498–504; Barak, L.S., et al., Proc Natl Acad Sci U S A, 1981. 78(5): 3034-8; Faulstich, H. and L. Fiume, Methods Enzymol, 1985. 112: 225-37; Zhelev, Z., A. et al, Toxicon, 1987. 25(9): 981-7; Khalacheva, K., et al, Eksp Med Morfol, 1990. 29(3): 26-30; U. Bernbach, H. Faulstich, Biochemistry 1990, 29, 6839–45; Mullersman, J. E. and J. F. Preston, Int. J. Peptide Protein Res. 1991, 37, 544–51; Mullersman, J.E. and J.F. Preston, Biochem Cell Biol, 1991. 69(7): 418-27; J. Anderl, H. Echner, H. Faulstich, Beilstein J. Org. Chem. 2012, 8, 2072–84; Moldenhauer, G., et al, J. Natl. Cancer Inst. 2012, 104, 622–34; A. Moshnikova, et al; Biochemistry 2013, 52, 1171–8; Zhao, L., et al., Chembiochem, 2015. 16(10): 1420-5; Zhou, B., et al., Biosens Bioelectron, 2015. 68: 189-96; WO2014/043403, US20150218220, EP 1661584). We have been working on the conjugation of amatoxins for a while. Examples of the structures of the conjugate of the antibody- amatoxins via the bridge linker are preferred as the following structures of Am01, Am02, Am03, Am04.


















Wherein mAb is an antibody; X₃ is CH₂, O, NH, NHC(O), NHC(O)NH, C(O), OC(O), OC(O)(NR₃), R₁, NHR₁, NR₁, C(O)R₁ or absent; R₇, R₈, R₉, R₁₀ and R₁₁ are independently H, OH, OR₁, NH₂, NHR₁, C₁-C₆ alkyl, or absent; Y₁ is O, O₂, or absent; X₁, X₂, R₁, R₂ and R₃ are the same defined in Formula (I).

In yet another embodiment, two or more different cytotoxic agents are preferred conjugated to a cell-binding molecule via a bridge linker of this patent. The two or more different cytotoxic agents can be selected from any combinations of tubulysins, maytansinoids, taxanoids (taxanes), CC-1065 analogs, daunorubicin and doxorubicin compounds, benzodiazepine dimers (e.g., dimers of pyrrolbenzodiazepine (PBD), tomaymycin, anthramycin, indolinbenzodiazepines, imidazobenzothiadiazepines, or oxazolidino-benzodiazepines), calicheamicins and the enediyne antibiotics, actinomycins, amanitins, azaserines, bleomycins, epirubicin, tamoxifen, idarubicin, dolastatins, auristatins (e.g. monomethyl auristatin E, MMAE, MMAF, auristatin PYE, auristatin TP, Auristatins 2-AQ, 6-AQ, EB (AEB), and EFP (AEFP)), duocarmycins, thiotepa, vincristines, hemiasterlins, nazumamides, microginins, radiosumins, alterobactins, microsclerodermins, theonellamides, esperamicins, PNU-159682, and their analogues and derivatives above thereof. Examples of the structures of the conjugates containing two or more different cytotoxic agents via the bridge linker are as the following Z01, Z02, Z02, Z04, Z05, Z06, Z07, Z08, Z09, Z10, Z12, Z13, Z14, Z15, Z16, Z17 and Z18:









Wherein mAb is an antibody; X₃ and X'₃ are independently CH₂, O, NH, NHC(O), NHC(O)NH, C(O), OC(O), OC(O)(NR₃), R₁,NHR₁, NR₁,C(O)R₁ or absent; X₄ and X'₄ are independently H, CH₂, OH, O, C(O), C(O)NH, C(O)N(R₁), R₁,NHR₁, NR₁,C(O)R₁ or C(O)O; M₁ and M₂ are independently H, Na, K, Ca, Mg, NH₄, NR₁R₂R₃; n, m₁, m₂, “—”, “—”, X₁, X₂, R₁, R₂ and R₃ are the same defined in Formula (I) and (II). In addition, R₁ and/or R₂ can be absent independently.

10 In yet another embodiment, cell-binding ligands or cell receptor agonists can be conjugated to a cell-binding molecule via a bridge linker of this patent. These conjugated cell-binding ligands or cell receptor agonists, in particular, antibody-receptor conjugates, can be not only to work as a targeting conductor/director to deliver the conjugate to malignant cells, but also be used to modulate or co-stimulate a desired immune response or altering signaling pathways.

In the immunotherapy, the cell-binding ligands or receptor agonists are preferred to conjugate to an antibody of TCR (T cell receptors) T cell, or of CARs (chimeric antigen receptors) T cells, or of B cell receptor (BCR), or the cytotoxic cells. The cell-binding ligands or receptor agonists are selected, but not limited, from: Folate derivatives (binding to the folate receptor, a protein over-expressed in ovarian cancer and in other malignancies) (Low, P. S. et al 2008, *Acc. Chem. Res.* 41, 120-9); Glutamic acid urea derivatives (binding to the prostate specific membrane antigen, a surface marker of prostate cancer cells) (Hillier, S. M. et al, 2009, *Cancer Res.* 69, 6932-40); Somato-statin (also known as growth hormone-inhibiting hormone (GHIH) or somato-tropin release-inhibiting factor (SRIF)) or somatotropin release-inhibiting hormone) and its analogues such as octreotide (Sandostatin) and lanreotide (Somatuline) (particularly for neuroendocrine tumors, GH-producing pituitary adenoma, paraganglioma, nonfunctioning pituitary adenoma, pheochromocytomas) (Ginj, M., et al, 2006, *Proc. Natl. Acad. Sci. U.S.A.* 103, 16436-41). In general, Somatostatin and its receptor subtypes (sst1, sst2, sst3, sst4, and sst5) have been found in many types of tumors, such as neuroendocrine tumors, in particular in GH-secreting pituitary adenomas (Reubi J. C., Landolt, A. M. 1984 *J. Clin. Endocrinol Metab* 59: 1148–51; Reubi J. C., Landolt A. M. 1987 *J Clin Endocrinol Metab* 65: 65–73; Moyse E, et al, *J Clin Endocrinol Metab* 61: 98–103) and gastroenteropancreatic tumors (Reubi J. C., et al, 1987 *J Clin Endocrinol Metab* 65: 1127–34; Reubi, J. C, et al, 1990 *Cancer Res* 50: 5969–77), pheochromocytomas (Epel-baum J, et al 1995 *J Clin Endocrinol Metab* 80:1837–44; Reubi J. C., et al, 1992 *J Clin Endocrinol Metab* 74: 1082–9), neuroblastomas (Prevost G, 1996 *Neuroendocrinology* 63:188–197; Moertel, C. L, et al 1994 *Am J Clin Path* 102:752–756), medullary thyroid cancers (Reubi, J. C, et al 1991 *Lab Invest* 64:567–573) small cell lung cancers (Sagman U, et al, 1990 *Cancer* 66:2129–2133), nonneuroendocrine tumors including brain tumors such as meningiomas, medulloblastomas, or gliomas (Reubi J. C., et al 1986 *J Clin Endo-crinol Metab* 63: 433–8; Reubi J. C., et al 1987 *Cancer Res* 47: 5758–64; Fruhwald, M. C, et al 1999 *Pediatr Res* 45: 697–708), breast carcinomas (Reubi J. C., et al 1990 *Int J Cancer* 46: 416–20; Srkalovic G, et al 1990 *J Clin Endocrinol Metab* 70: 661–669), lymphomas (Reubi J. C., et al 1992, *Int J Cancer* 50: 895–900), renal cell cancers (Reubi J. C., et al 1992, *Cancer Res* 52: 6074–6078), mesenchymal tumors (Reubi J. C., et al 1996 *Cancer Res* 56: 1922–31), prostatic (Reubi J. C., et al 1995, *J. Clin. Endocrinol Metab* 80: 2806–14; et al 1989, *Prostate* 14:191–208; Halmos G, et al *J. Clin. Endo-crinol Metab* 85: 2564–71), ovarian (Halmos, G, et al, 2000 *J Clin Endocrinol Metab* 85: 3509–12; Reubi J. C., et al 1991 *Am J Pathol* 138:1267–72), gastric (Reubi J. C., et al 1999, *Int J Cancer* 81: 376–86; Miller, G. V, 1992 *Br J Cancer* 66: 391–95), hepatocellular (Kouroumalis E, et al 1998 *Gut* 42: 442–7; Reubi J. C.,

et al 1999 Gut 45: 66–774) and nasopharyngeal carcinomas (Loh K. S, et al, 2002 Virchows Arch 441: 444–8); certain Aromatic sulfonamides, specific to carbonic anhydrase IX (a marker of hypoxia and of renal cell carcinoma) (Neri, D., et al, Nat. Rev. Drug Discov. 2011, 10, 767–7); Pituitary adenylate cyclase activating peptides (PACAP) (PAC1) for pheochromocytomas and paragangliomas; Vasoactive intestinal peptides (VIP) and their receptor subtypes (VPAC1, VPAC2) for cancers of lung, stomach, colon, rectum, breast, prostate, pancreatic ducts, liver, urinary bladder and epithelial tumors; α -Melanocyte-stimulating hormone (α -MSH) receptors for various tumors; Cholecystokinin (CCK)/gastrin receptors and their receptor subtypes (CCK1 (formerly CCK-A) and CCK2 for small cell lung cancers, medullary thyroid carcinomas, astrocytomas, insulinomas and ovarian cancers; Bombesin (Pyr-Gln-Arg-Leu-Gly-Asn-Gln-Trp-Ala-Val-Gly-His-Leu-Met-NH₂)/gastrin-releasing peptide (GRP) and their receptor subtypes (BB1, GRP receptor subtype (BB2), the BB3 and BB4) for renal cell, breast, lung, gastric and prostate carcinomas, and neuroblastoma (and neuroblastoma (Ohlsson, B., et al, 1999, Scand. J. Gastroenterology 34 (12): 1224–9; Weber, H. C., 2009, Cur. Opin. Endocri. Diab. Obesity 16(1): 66–71, Gonzalez N, et al, 2008, Cur. Opin. Endocri. Diab. Obesity 15(1), 58–64); Neurotensin receptors and its receptor subtypes (NTR1, NTR2, NTR3) for small cell lung cancer, neuroblastoma, pancreatic, colonic cancer and Ewing sarcoma; Substance P receptors and their receptor subtypes (such as NK1 receptor for Glial tumors, Hennig I. M., et al 1995 Int. J. Cancer 61, 786–792); Neuropeptide Y (NPY) receptors and its receptor subtypes (Y1–Y6) for breast carcinomas; Homing Peptides include RGD (Arg-Gly-Asp), NGR (Asn-Gly-Arg), the dimeric and multimeric cyclic RGD peptides (e.g. cRGDfV) that recognize receptors (integrins) on tumor surfaces (Laakkonen P, Vuorinen K. 2010, Integr Biol (Camb). 2(7–8): 326–337; Chen K, Chen X. 2011, Theranostics. 1:189–200; Garanger E, et al, Anti-Cancer Agents Med Chem. 7 (5): 552–558; Kerr, J. S. et al, Anticancer Research, 19(2A), 959–968; Thumshirn, G, et al, 2003 Chem. Eur. J. 9, 2717–2725), and TAASGVRSMD or LTLRWVGLMS (chondroitin sulfate proteoglycan NG2 receptor) and F3 peptides (31 amino acid peptide that binds to cell surface-expressed nucleolin receptor) (Zitzmann, S., 2002 Cancer Res., 62, 18, pp. 5139–5143, Temminga, K., 2005, Drug Resistance Updates, 8, 381–402; P. Laakkonen and K. Vuorinen, 2010 Integrative Biol, 2(7–8), 326–337; M. A. Burg, 1999 Cancer Res., 59(12), 2869–2874; K. Porkka, et al 2002, Proc. Nat. Acad. Sci. USA 99(11), 7444–9); Cell Penetrating Peptides (CPPs) (Nakase I, et al, 2012, J. Control Release. 159(2), 181–188); Peptide Hormones, such as luteinizing hormone-releasing hormone (LHRH) agonists and antagonists, and gonadotropin-releasing hormone (GnRH) agonist, acts by targeting follicle stimulating hormone (FSH) and luteinizing hormone (LH), as well as testosterone production, e.g. buserelin (Pyr-His-Trp-Ser-


Tyr-D-Ser(OtBu)-Leu-Arg-Pro-NHEt), Gonadorelin (Pyr-His-Trp-Ser-Tyr-Gly-Leu-Arg-Pro-Gly-NH₂), Goserelin (Pyr-His-Trp-Ser-Tyr-D-Ser(OtBu)-Leu-Arg-Pro-AzGly-NH₂), Histrelin (Pyr-His-Trp-Ser-Tyr-D-His(N-benzyl)-Leu-Arg-Pro-NHEt), leuprolide (Pyr-His-Trp-Ser-Tyr-D-Leu-Leu-Arg-Pro-NHEt), Nafarelin (Pyr-His-Trp-Ser-Tyr-2Nal-Leu-Arg-Pro-Gly-NH₂),
 5 Triptorelin (Pyr-His-Trp-Ser-Tyr-D-Trp-Leu-Arg-Pro-Gly-NH₂), Nafarelin, Deslorelin, Abarelix (Ac-D-2Nal-D-4-chloroPhe-D-3-(3-pyridyl)Ala-Ser-(N-Me)Tyr-D-Asn-Leu-isopropylLys-Pro-DAla-NH₂), Cetrorelix (Ac-D-2Nal-D-4-chloro-Phe-D-3-(3-pyridyl)Ala-Ser-Tyr-D-Cit-Leu-Arg-Pro-D-Ala-NH₂), Degarelix (Ac-D-2Nal-D-4-chloroPhe-D-3-(3-pyridyl)Ala-Ser-4-aminoPhe(L-hydroorotyl)-D-4-aminoPhe(carba-moyl)-Leu-isopropylLys-Pro-D-Ala-NH₂), and Ganirelix (Ac-
 10 D-2Nal-D-4-chloroPhe-D-3-(3-pyridyl)Ala-Ser-Tyr-D-(N⁹, N¹⁰-diethyl)-homoArg-Leu-(N⁹, N¹⁰-diethyl)-homoArg-Pro-D-Ala-NH₂) (Thundimadathil, J., J. Amino Acids, 2012, 967347, doi:10.1155/2012/967347; Boccon-Gibod, L.; et al, 2011, Therapeutic Advances in Urology 3(3): 127–140; Debruyne, F., 2006, Future Oncology, 2(6), 677–696; Schally A. V; Nagy, A. 1999 Eur J Endocrinol 141:1–14; Koppan M, et al 1999 Prostate 38:151–158); and Pattern Recognition
 15 Receptors (PRRs), such as Toll-like receptors (TLRs), C-type lectins and NodlikeReceptors (NLRs) (Fukata, M., et al, 2009, Semin. Immunol. 21, 242–253; Maisonneuve, C., et al, 2014, Proc. Natl. Acad. Sci. U. S. A. 111, 1–6; Botos, I., et al, 2011, Structure 19, 447–459; Means, T. K., et al, 2000, Life Sci. 68, 241–258) that range in size from small molecules (imiquimod, guanisine and adenosine analogs) to large and complex biomacromolecules such as
 20 lipopolysaccharide (LPS), nucleic acids (CpG DNA, polyI:C) and lipopeptides (Pam3CSK4) (Kasturi, S. P., et al, 2011, Nature 470, 543–547; Lane, T., 2001, J. R. Soc. Med. 94, 316; Hotz, C., and Bourquin, C., 2012, Oncoimmunology 1, 227–228; Dudek, A. Z., et al, 2007, Clin. Cancer Res. 13, 7119–25); Calcitonin receptors which is a 32-amino-acid neuropeptide involved in the regulation of calcium levels largely through its effects on osteoclasts and on the kidney
 25 (Zaidi M, et al, 1990 Crit Rev Clin Lab Sci 28, 109–174; Gorn, A. H., et al 1995 J Clin Invest 95:2680–91); And integrin receptors and their receptor subtypes (such as $\alpha_v\beta_1$, $\alpha_v\beta_3$, $\alpha_v\beta_5$, $\alpha_v\beta_6$, $\alpha_6\beta_4$, $\alpha_7\beta_1$, $\alpha_L\beta_2$, $\alpha_{IIb}\beta_3$, etc) which generally play important roles in angiogenesis are expressed on the surfaces of a variety of cells, in particular, of osteoclasts, endothelial cells and tumor cells (Ruoslahti, E. et al, 1994 Cell 77, 477-8; Albelda, S. M. et al, 1990 Cancer Res., 50, 6757-64).
 30 Short peptides, GRGDSPK and Cyclic RGD pentapeptides, such as cyclo(RGDfV) (L1) and its derivatives [cyclo(-N(Me)R-GDfV), cyclo(R-Sar-DfV), cyclo-(RG-N(Me)D-fV), cyclo(RGD-N(Me)f-V), cyclo(RGDf-N(Me)V)-(Cilengitide)] have shown high binding affinities of the integrin receptors (Dechantsreiter, M. A. et al, 1999 J. Med. Chem. 42, 3033-40, Goodman, S. L., et al, 2002 J. Med. Chem. 45, 1045-51).

The cell-binding ligands or cell receptor agonists can be Ig-based and non-Ig-based protein scaffold molecules. The Ig-Based scaffolds can be selected, but not limited, from Nanobody (a derivative of VHH (camelid Ig)) (Muyldermans S., 2013 *Annu Rev Biochem.* 82, 775–97); Domain antibodies (dAb, a derivative of VH or VL domain) (Holt, L. J, et al, 2003, *Trends Biotechnol.* 21, 484–90); Bispecific T cell Engager (BiTE, a bispecific diabody) (Baeuerle, P. A, et al, 2009, *Curr. Opin. Mol. Ther.* 11, 22–30); Dual Affinity ReTargeting (DART, a bispecific diabody) (Moore P. A. P, et al. 2011, *Blood* 117(17), 4542–51); Tetravalent tandem antibodies (TandAb, a dimerized bispecific diabody) (Cochlovius, B, et al. 2000, *Cancer Res.* 60(16):4336–4341). The Non-Ig scaffolds can be selected, but not limited, from Anticalin (a derivative of Lipocalins) (Skerra A. 2008, *FEBS J.*, 275(11): 2677–83; Beste G, et al, 1999 *Proc. Nat. Acad. USA.* 96(5):1898–903; Skerra, A. 2000 *Biochim Biophys Acta*, 1482(1-2): 337–50; Skerra, A. 2007, *Curr Opin Biotechnol.* 18(4): 295–304; Skerra, A. 2008, *FEBS J.* 275(11):2677–83); Adnectins (10th FN3 (Fibronectin)) (Koide, A, et al, 1998 *J. Mol. Biol.*, 284(4):1141–51; Batori V, 2002, *Protein Eng.* 15(12): 1015–20; Tolcher, A. W, 2011, *Clin. Cancer Res.* 17(2): 363–71; Hackel, B. J, 2010, *Protein Eng. Des. Sel.* 23(4): 211–19); Designed Ankyrin Repeat Proteins (DARPin) (a derivative of ankrin repeat (AR) proteins) (Boersma, Y.L, et al, 2011 *Curr Opin Biotechnol.* 22(6): 849–57), e.g. DARPin C9, DARPin Ec4 and DARPin E69_LZ3_E01 (Winkler J, et al, 2009 *Mol Cancer Ther.* 8(9), 2674–83; Patricia M-K. M., et al, *Clin Cancer Res.* 2011; 17(1):100–10; Boersma Y. L, et al, 2011 *J. Biol. Chem.* 286(48), 41273–85); Avimers (a domain A/low-density lipoprotein (LDL) receptor) (Boersma Y. L, 2011 *J. Biol. Chem.* 286(48): 41273–41285; Silverman J, et al, 2005 *Nat. Biotechnol.*, 23(12):1556–61).

Examples of the structures of the conjugate of the antibody-cell-binding ligands or cell receptor agonists via the bridge linker are as the following: LB01 (PMSA ligand conjugate), LB02 (Folate conjugate), LB03 (Somatostatin conjugate), LB04 (Octreotide, a Somatostatin analogconjugate), LB05 (Lanreotide, a Somatostatin analog conjugate), LB06 (Vapreotide (Sanvar), a Somatostatin analog conjugate), LB07 (CAIX ligandconjugate), LB08 (CAIXligand conjugate), LB09 (luteinizing hormone-releasing hormone (LH-RH) ligand and GnRH conjugate), LB10 (luteinizing hormone-releasing hormone (LH-RH) and GnRH ligand conjugate), LB11 (GnRH antagonist, Abarelix conjugate), LB12 (cobalamin, VB12 analog conjugate), LB13 (Gastrin releasing peptide receptor (GRPr), MBA conjugate), LB14 (for $\alpha_v\beta_3$ integrin receptor, cyclic RGD pentapeptide conjugate), LB15 (hetero-bivalent peptide ligand conjugate for VEGF receptor), LB16 (Neuromedin B conjugate), LB 17 (bombesin conjugate for a G-protein coupled receptor), LB18 (TLR₂ conjugate for a Toll-like receptor), LB19 (for an androgen receptor), LB20 (Cilengitide/cyclo(-RGDfV-) conjugate for an α_v intergrin receptor) and

$$\left[\left(\begin{array}{c} \text{HOOC} \\ | \\ \text{CH}_2 \\ | \\ \text{CH} \\ | \\ \text{NH} \\ | \\ \text{C=O} \\ | \\ \text{NH} \\ | \\ \text{CH} \\ | \\ \text{CH}_2 \\ | \\ \text{O} \\ | \\ \text{X}_3 \\ | \\ \text{R}_1 \end{array} \right)_{m_1} \text{X}_1 \text{C}(=\text{O}) \text{S} \left(\begin{array}{c} \text{R}_2 \\ | \\ \text{X}_2 \end{array} \right)_{m_2} \text{C}(=\text{O}) \text{S} \right]_n \text{mAb}$$





LB01




Chemical structure of a dendritic polymer with a dendritic core and a dendritic shell. The core is a dendritic polymer with a central amine group (NH₂) and a dendritic shell. The shell is composed of repeating units of a dendritic polymer, each containing a dendritic core and a dendritic shell. The dendritic core is a dendritic polymer with a central amine group (NH₂) and a dendritic shell. The dendritic shell is composed of repeating units of a dendritic polymer, each containing a dendritic core and a dendritic shell. The dendritic core is a dendritic polymer with a central amine group (NH₂) and a dendritic shell. The dendritic shell is composed of repeating units of a dendritic polymer, each containing a dendritic core and a dendritic shell.


LB04






5 LB09







LB11




LB12









LB13,




LB14




LB15






LB22 (Dexamethasone

conjugate)




LB23 (fluticasone

propionate conjugate)







LB24 (Beclomethasone

dipropionate)



LB25 (Triamcinolone

acetone conjugate)




10


Wherein mAb is an antibody; X_3 is CH_2 , O, NH, NHC(O) , NHC(O)NH , C(O) , OC(O) , $\text{OC(O)(NR}_3\text{)}$, R_1 , NHR_1 , NR_1 , C(O)R_1 or absent; X_4 is H, CH_2 , OH, O, C(O) , C(O)NH , $\text{C(O)N(R}_1\text{)}$, R_1 , NHR_1 , NR_1 , C(O)R_1 or C(O)O ; X_5 is H, CH_3 , F, or Cl; M_1 and M_2 are independently H, Na, K, Ca, Mg, NH_4 , $\text{NR}_1\text{R}_2\text{R}_3$; R_6 is 5'-deoxyadenosyl, Me, OH, or CN;

“ $\equiv$ ” represents either single bond or double bond; m_1 , m_2 , n , “—”, X_1 , X_2 , R_1 , and R_2 are the same defined in Formula (I). In addition, R_1 can be absent and R_2 can be H.


In yet another embodiment, IgG antibody conjugates conjugated with one, or two, or more differently function molecules or drugs are preferred to be conjugated specifically to a pair of thiols (through reduction of the disulfide bonds) between the light chain and heavy chain, the upper disulfide bonds between the two heavy chains, and the lower disulfide bonds between the two heavy chains as shown in the following structure, ST1, ST2, ST3, ST4, ST5, ST6, ST7 or ST8.




ST1




ST2




ST3




ST4




ST5



ST6



ST7



ST8

5 Wherein R_1 , R_1' , R_1'' , R_1''' , R_1'''' , and R_1''''' are independently defined as R_1 which is described the same in Formula (I); Wherein R_2' , R_2'' , R_2''' , R_2'''' , and R_2''''' are independently defined as R_2 which is described the same in Formula (I); X_1' , X_1'' , X_1''' , X_1'''' , and X_1''''' are

independently defined as X_1 which is described the same in claim Formula (I); X_2' , X_2'' , X_2''' , X_2'''' , and X_2''''' are independently defined as X_2 which is described the same in claim Formula (I); “====” represents either single bond or double bond; m_1 , m_2 , m_3 , m_4 , m_5 , m_6 , m_7 and m_8 are independently an integer from 1 to 30, preferably from 1 to 10; Drug₁, Drug₂, Drug₃, Drug₄, Drug₅ and Drug₆ is independently selected from tubulysins, maytansinoids, taxanoids (taxanes), CC-1065 analogs, daunorubicin and doxorubicin compounds, benzodiazepine dimers (e.g., dimers of pyrrolbenzodiazepine (PBD), tomaymycin, anthramycin, indolinbenzodiazepines, imidazobenzothiadiazepines, or oxazolidinobenzodiazepines), calicheamicins and the enediyne antibiotics, actinomycin, amanitin (amatoxins), azaserines, bleomycins, epirubicin, tamoxifen, idarubicin, dolastatins, auristatins (e.g. monomethyl auristatin E, MMAE, MMAF, auristatin PYE, auristatin TP, Auristatins 2-AQ, 6-AQ, EB (AEB), and EFP (AEFP)), duocarmycins, thiotepa, vincristines, hemiasterlins, nazumamides, microginins, radiosumins, alterobactins, microsclerodermins, theonellamides, esperamicins, PNU-159682, and their analogues and derivatives, as well as the function molecules described throughout this patent application. In addition, R_1 , R_1' , R_1'' , R_1''' , R_1'''' and/or R_1''''' can be absent.

In yet another embodiment, a pharmaceutical composition comprising a therapeutically effective amount of the conjugate of Formula (II) or any conjugates described through the present patent can be administered concurrently with the other therapeutic agents such as the chemotherapeutic agent, the radiation therapy, immunotherapy agents, autoimmune disorder agents, anti-infectious agents or the other conjugates for synergistically effective treatment or prevention of a cancer, or an autoimmune disease, or an infectious disease. The synergistic agents are preferably selected from one or several of the following drugs: Abatacept (Orencia), Abiraterone acetate (Zytiga®), Acetaminophen/hydrocodone, Adalimumab, afatinib dimaleate (Gilotrif®), alemtuzumab (Campath®), Alitretinoin (Panretin®), ado-trastuzumab emtansine (Kadcyla™), Amphetamine mixed salts (Amphetamine/dextroamphetamine, or Adderall XR), anastrozole (Arimidex®), Aripiprazole, Atazanavir, Atezolizumab (MPDL3280A), Atorvastatin, axitinib (Inlyta®), AZD9291, belinostat (Beleodaq™), Bevacizumab (Avastin®), Cabazitaxel (Jevtana®), Cabozantinib (Cometriq™), bexarotene (Targtin®), blinatumomab (Blinicyto™), Bortezomib (Velcade®), bosutinib (Bosulif®), brentuximab vedotin (Adcetris®), Budesonide, Budesonide/formoterol, Buprenorphine, Capecitabine, carfilzomib (Kyprolis®), Celecoxib, ceritinib (LDK378/Zykadia), Cetuximab (Erbix®), Ciclosporin, Cinacalcet, crizotinib (Xalkori®), Dabigatran, dabrafenib (Tafinlar®), Darbepoetin alfa, Darunavir, imatinib mesylate (Gleevec®), dasatinib (Sprycel®), denileukin diftitox (Ontak®), Denosumab (Xgeva®), Depakote, Dexlansoprazole, Dexmethylphenidate, Dinutuximab (Unituxin™), Doxycycline,

Duloxetine, Durvalumab (MEDI4736), Emtricitabine/Rilpi-virine/Tenofovir disoproxil fumarate, Emtricitabine/tenofovir/efavirenz, Enoxaparin, Enzalutamide (Xtandi®), Epoetin alfa, erlotinib (Tarceva®), Esomeprazole, Eszopiclone, Etanercept, Everolimus (Afinitor®), exemestane (Aromasin®), everolimus (Afinitor®), Ezetimibe, Ezetimibe/simvastatin, Fenofibrate, Filgrastim, fingolimod, Fluticasone propionate, Fluticasone/salmeterol, fulvestrant (Faslodex®), gefitinib (Iressa®), Glatiramer, Goserelin acetate (Zoladex), Icotinib, Imatinib (Gleevec), Ibritumomab tiuxetan (Zevalin®), ibrutinib (Imbru-vica™), idelalisib (Zydelig®), Infliximab, Insulin aspart, Insulin detemir, Insulin glargine, Insulin lispro, Interferon beta 1a, Interferon beta 1b, lapatinib (Tykerb®), Ipilimumab (Yervoy®), Ipratropium bromide/salbutamol, Lanreotide acetate (Somatuline® Depot), lenaliomide (Revlimid®), lenvatinib mesylate (Lenvima™), letrozole (Femara®), Levothyroxine, Levothyroxine, Lidocaine, Linezolid, Liraglutide, Lisdexamphetamine, MEDI4736 (AstraZeneca, Celgene), Memantine, Methylphenidate, Metoprolol, Modafinil, Mometasone, Nilotinib (Tasigna®), Nivolumab (Opdivo®), ofatumumab (Arzerra®), obinutuzumab (Gazyva™), olaparib (Lynparza™), Olmesartan, Olmesartan/hydrochlorothiazide, Omalizumab, Omega-3 fatty acid ethyl esters, Oseltamivir, Oxycodone, palbociclib (Ibrance®), Palivizumab, panitumumab (Vectibix®), panobinostat (Farydak®), pazopanib (Votrient®), pembrolizumab (Keytruda®), Pemetrexed (Alimta), pertuzumab (Perjeta™), Pneumococcal conjugate vaccine, pomalidomide (Pomalyst®), Pregabalin, Quetiapine, Rabeprazole, radium 223 chloride (Xofigo®), Raloxifene, Raltegravir, ramucirumab (Cyramza®), Ranibizumab, regorafenib (Stivarga®), Rituximab (Rituxan®), Rivaroxaban, romidepsin (Istodax®), Rosuvastatin, ruxolitinib phosphate (Jakafi™), Salbutamol, Sevelamer, Sildenafil, siltuximab (Sylvant™), Sitagliptin, Sitagliptin/ metformin, Solifenacin, Sorafenib (Nexavar®), Sunitinib (Sutent®), Tadalafil, tamoxifen, Telaprevir, temsirolimus (Torisel®), Tenofovir/emtricitabine, Testosterone gel, Thalidomide (Immunoprin, Talidex), Tiotropium bromide, toremifene (Fareston®), trametinib (Mekinist®), Trastuzumab, Tretinoin (Vesanoid®), Ustekinumab, Valsartan, vandetanib (Caprelsa®), vemurafenib (Zelboraf®), vorinostat (Zolinza®), ziv-aflibercept (Zaltrap®), Zostavax., and their analogs, derivatives, pharmaceutically acceptable salts, carriers, diluents, or excipients thereof, or a combination above thereof.

The drugs/ cytotoxic agents used for conjugation via a bridge linker of the present patent can be any analogues and/or derivatives of drugs/molecules described in the present patent. One skilled in the art of drugs/cytotoxic agents will readily understand that each of the drugs/cytotoxic agents described herein can be modified in such a manner that the resulting compound still retains the specificity and/or activity of the starting compound. The skilled artisan will also


understand that many of these compounds can be used in place of the drugs/cytotoxic agents described herein. Thus, the drugs/cytotoxic agents of the present invention include analogues and derivatives of the compounds described herein.

All references cited herein and in the examples that follow are expressly incorporated by reference in their entirety.

EXAMPLES


The invention is further described in the following examples, which are not intended to limit the scope of the invention. Cell lines described in the following examples were maintained in culture according to the conditions specified by the American Type Culture Collection (ATCC) or Deutsche Sammlung von Mikroorganismen und Zellkulturen GmbH, Braunschweig, Germany (DMSZ), or The Shanghai Cell Culture Institute of Chinese Academy of Science, unless otherwise specified. Cell culture reagents were obtained from Invitrogen Corp., unless otherwise specified. All anhydrous solvents were commercially obtained and stored in Sure-seal bottles under nitrogen. All other reagents and solvents were purchased as the highest grade available and used without further purification. The preparative HPLC separations were performed with Varian PreStar HPLC. NMR spectra were recorded on Varian Mercury 400 MHz Instrument. Chemical shifts (δ) are reported in parts per million (ppm) referenced to tetramethylsilane at 0.00 and coupling constants (J) are reported in Hz. The mass spectral data were acquired on a Waters Xevo QTOF mass spectrometer equipped with Waters Acquity UPLC separations module and Acquity TUV detector.

Example 1. Synthesis of 2,3-dibromosuccinic anhydride (**7**)




To a solution of 2,3-dibromosuccinic acid (10.00 g, 36.51 mmol) in dry CH_2Cl_2 (100 ml) at 0 °C was added phosphorus pentoxide (12.21 g, 85.84 mmol). The mixture was stirred at 0 °C for 2 h and then r.t. for 5 h, filtered through short SiO_2 column, and rinsed the column with EtOAc/ CH_2Cl_2 (1:6). The filtrate was concentrated and triturated with EtOAc/hexane to afford the title compound (6.63 g, 71% yield). ESI MS m/z $\text{C}_4\text{H}_2\text{Br}_2\text{O}_3$ $[\text{M}+\text{H}]^+$, calcd. 256.85, found 256.70.

Example 2. Synthesis of (S, E)-2-methyl-N-(3-methylbutan-2-ylidene)propane-2-sulfonamide (**122**)




To a solution of (S)-2-methylpropane-2-sulfonamide (100 g, 0.825 mol, 1.0 eq.) in 1 L THF was added Ti(OEt)₄ (345 mL, 1.82 mol, 2.2 eq.) and 3-methyl-2-butanone (81 mL, 0.825 mol, 1.0 eq.) under N₂ at r.t. The reaction mixture was refluxed for 16 h, then cooled to r.t. and poured onto iced water. The mixture was filtered and the filter cake was washed with EtOAc. The organic layer was separated, dried over anhydrous Na₂SO₄ and concentrated to give a residue which was purified by vacuum distillation (15-20 torr, 95 °C) to afford product **122** (141 g, 90% yield) as a yellow oil. ¹H NMR (500 MHz, CDCl₃) δ 2.54 – 2.44 (m, 1H), 2.25 (s, 3H), 1.17 (s, 9H), 1.06 (dd, *J* = 6.9, 5.1 Hz, 6H). MS ESI *m/z* calcd for C₉H₁₉NaNOS [M+Na]⁺ 212.12; found 212.11.

Example 3. Synthesis of (2S,3S)-2-azido-3-methylpentanoic acid (**124**)



To a solution of NaN₃ (20.0 g, 308 mmol) in a mixture of water (50 mL) and dichloromethane (80 mL), cooled at 0 °C, Tf₂O (10 mL, 59.2 mmol, 2.0 eq.) was added slowly. After addition, the reaction was stirred at 0 °C for 2 h, then the organic phase was separated and the aqueous phase was extracted with dichloromethane (2 × 40 mL). The combined organic phases were washed with saturated NaHCO₃ solution and used as is. The dichloromethane solution of triflyl azide was added to a mixture of (L)-isoleucine (4.04 g, 30.8 mmol, 1.0 eq.), K₂CO₃ (6.39 g, 46.2 mmol, 1.5 eq.), CuSO₄·5H₂O (77.4 mg, 0.31 mmol, 0.01 eq.) in water (100 mL) and methanol (200 mL). The mixture was stirred at r.t. for 16 h. The organic solvents were removed under reduced pressure and the aqueous phase was diluted with water (250 mL) and acidified to pH 6 with concentrated HCl and diluted with phosphate buffer (0.25 M, pH 6.2, 250 mL). The aqueous layer was washed with EtOAc (5 × 100 mL) to remove the sulfonamide by-product, and then acidified to pH 2 with concentrated HCl, extracted with EtOAc (3 × 150 mL). The combined organic layers were dried over anhydrous Na₂SO₄, filtered and concentrated to give product **124** (4.90 g, 99% yield) as a colorless oil. ¹H NMR (500 MHz, CDCl₃) δ 12.01 (s, 1H), 3.82 (d, *J* = 5.9 Hz, 1H), 2.00 (ddd, *J* = 10.6, 8.6, 5.5 Hz, 1H), 1.54 (dq, *J* = 14.8, 7.5, 4.4 Hz, 1H), 1.36 – 1.24 (m, 1H), 1.08 – 0.99 (m, 3H), 0.97 – 0.87 (m, 3H).


Example 4. Synthesis of D-N-methyl pipecolic acid (**126**)



To a solution of D-pipecolic acid (10.0 g, 77.4 mmol, 1.0 eq.) in methanol (100 mL) was added formaldehyde (37% aqueous solution, 30.8 mL, 154.8 mmol, 2.0 eq.), followed by Pd/C (10 wt%, 1.0 g). The reaction mixture was stirred under H₂ (1 atm) overnight, and then filtered


through Celite, with washing of the filter pad with methanol. The filtrate was concentrated under reduced pressure to afford compound **126** (10.0 g, 90% yield) as a white solid.

Example 5. Synthesis of (R)-perfluorophenyl 1-methylpiperidine-2-carboxylate (**127**)




To a solution of D-N-methyl pipercolinic acid (2.65 g, 18.5 mmol) in EtOAc (50 mL) were added pentafluorophenol (3.75 g, 20.4 mmol) and DCC (4.21 g, 20.4 mmol). The reaction mixture was stirred at r.t. for 16 h, and then filtered over Celite. The filter pad was washed with 10 mL of EtOAc. The filtrate was used immediately without further purification or concentration.

Example 6. Synthesis of 2,2-diethoxyethanethioamide (**129**)




2,2-diethoxyacetonitrile (100 g, 0.774 mol, 1.0 eq.) was mixed with (NH₄)₂S aqueous solution (48%, 143 mL, 1.05 mol, 1.36 eq.) in methanol (1.5 L) at room temperature. After stirring for 16 h, the reaction mixture was concentrated and the residue was taken up in dichloromethane, washed with saturated NaHCO₃ solution and brine, dried over anhydrous Na₂SO₄ and concentrated. The residue was triturated with a solvent mixture of petroleum ether and dichloromethane. After filtration, the desired product **2** as a white solid was collected (100 g, 79% yield). ¹H NMR (500 MHz, CDCl₃) δ 7.81 (d, *J* = 71.1 Hz, 2H), 5.03 (s, 1H), 3.73 (dq, *J* = 9.4, 7.1 Hz, 2H), 3.64 (dq, *J* = 9.4, 7.0 Hz, 2H), 1.25 (t, *J* = 7.1 Hz, 6H).

Example 7. Synthesis of ethyl 2-(diethoxymethyl)thiazole-4-carboxylate (**130**)



90 g of molecular sieves (3Å) was added to a mixture of thioamide **129** (100 g, 0.61 mol, 1.0 eq.) and ethyl bromopyruvate (142 mL, 1.1 mol, 1.8 eq.) in 1 L EtOH. The mixture was refluxed (internal temperature about 60 °C) for 1h, then ethanol was removed on rotovap and the residue was taken up in dichloromethane. The solid was filtered off and the filtrate was concentrated and purified by column chromatography (PE/EtOAc 5:1-3:1) to give thiazole carboxylate **130** (130 g, 82% yield) as a yellow oil.


Example 8. Synthesis of ethyl 2-formylthiazole-4-carboxylate (**131**)



To a solution of compound **130** (130 g, 0.50 mol) in acetone (1.3 L) was added 2 N HCl (85 mL, 0.165 mol, 0.33 eq.). The reaction mixture was refluxed (internal temperature about 60 °C), monitored by TLC analysis until starting material was completely consumed (about 1-2 h).


Acetone was removed under reduced pressure and the residue was taken up in dichloromethane (1.3 L), washed with saturated NaHCO₃ solution, water and brine, and then dried over anhydrous Na₂SO₄. The solution was filtered and concentrated under reduced pressure. The crude product was purified by recrystallization from petroleum ether and diethyl ether to afford a white solid **131** (40 g, 43% yield). ¹H NMR (500 MHz, CDCl₃) δ 10.08 – 10.06 (m, 1H), 8.53 – 8.50 (m, 1H), 4.49 (q, *J* = 7.1 Hz, 2H), 1.44 (t, *J* = 7.1 Hz, 3H). MS ESI *m/z* calcd for C₇H₈NO₃S [M+H]⁺ 186.01; found 186.01.

Example 9. Synthesis of ethyl 2-((R,E)-3-(((S)-tert-butylsulfinyl)imino)-1-hydroxy-4-methylpentyl)thiazole-4-carboxylate (**132**)




To a solution of diisopropylamine (121 mL, 0.86 mol, 4.0 eq.) in dry THF (300 mL) was added *n*-butyllithium (2.5 M, 302 mL, 0.76 mol 3.5 eq.) at -78 °C under N₂. The reaction mixture was warmed to 0 °C over 30 min and then cooled back to -78 °. Compound **122** (57 g, 0.3 mol, 1.4 eq.) in THF (200 mL) was added. The reaction mixture was stirred for 1 h before ClTi(O^{*i*}Pr)₃ (168.5 g, 0.645 mol, 3.0 eq.) in THF (350 mL) was added dropwise. After stirring for 1 h, compound **131** (40 g, 0.215 mol, 1.0 eq.) dissolved in THF (175 mL) was added dropwise and the resulting reaction mixture was stirred for 2 h. The completion of the reaction was indicated by TLC analysis. The reaction was quenched by a mixture of acetic acid and THF (v/v 1:4, 200 mL), then poured onto iced water, extracted with EtOAc (4 × 500 mL). The organic phase was washed with water and brine, dried over anhydrous Na₂SO₄, filtered and concentrated. The residue was purified by column chromatography (DCM/EtOAc/PE 2:1:2) to afford the compound **132** (60 g, 74% yield) as a colorless oil. ¹H NMR (500 MHz, CDCl₃) δ 8.13 (s, 1H), 6.63 (d, *J* = 8.2 Hz, 1H), 5.20 – 5.11 (m, 1H), 4.43 (q, *J* = 7.0 Hz, 2H), 3.42 – 3.28 (m, 2H), 2.89 (dt, *J* = 13.1, 6.5 Hz, 1H), 1.42 (t, *J* = 7.1 Hz, 3H), 1.33 (s, 9H), 1.25 – 1.22 (m, 6H). MS ESI *m/z* calcd for C₁₆H₂₆NaN₂O₄S₂ [M+Na]⁺ 397.13, found 397.11.

Example 10. Synthesis of ethyl 2-((1R,3R)-3-((S)-1,1-dimethylethylsulfinamido)-1-hydroxy-4-methylpentyl)thiazole-4-carboxylate (**133**)




A solution of compound **132** (23.5 g, 62.7 mmol) dissolved in THF (200 mL) was cooled to -45 °C. Ti(OEt)₄ (42.9 mL, 188 mmol, 3.0 eq.) was added slowly. After the completion of addition, the mixture was stirred for 1 h, before NaBH₄ (4.75 g, 126 mmol, 2.0 eq.) was added in portions. The reaction mixture was stirred at -45 °C for 3 h. TLC analysis showed some starting material still remained. The reaction was quenched with HOAc/THF (v/v 1:4, 25 mL), followed by EtOH (25 mL). The reaction mixture was poured onto ice (100 g) and warmed to r.t. After filtration over Celite, the organic phase was separated and washed with water and brine, dried over anhydrous Na₂SO₄, filtered, and concentrated. The residue was purified by column chromatography (EtOAc/PE 1:1) to deliver product **133** (16.7 g, 71% yield) as a white solid. ¹H NMR (500 MHz, CDCl₃) δ 8.10 (s, 1H), 5.51 (d, *J* = 5.8 Hz, 1H), 5.23 – 5.15 (m, 1H), 4.41 (q, *J* = 7.0 Hz, 2H), 3.48 – 3.40 (m, 1H), 3.37 (d, *J* = 8.3 Hz, 1H), 2.29 (t, *J* = 13.0 Hz, 1H), 1.95 – 1.87 (m, 1H), 1.73 – 1.67 (m, 1H), 1.40 (t, *J* = 7.1 Hz, 3H), 1.29 (s, 9H), 0.93 (d, *J* = 7.3 Hz, 3H), 0.90 (d, *J* = 7.2 Hz, 3H). MS ESI *m/z* calcd for C₁₆H₂₈NaN₂O₄S₂ [M+Na]⁺ 399.15, found 399.14.

Example 11. Synthesis of ethyl 2-((1R,3R)-3-amino-1-hydroxy-4-methylpentyl)thiazole-4-carboxylate hydrochloride (**134**)




To a solution of compound **133** (6.00 g, 16.0 mmol, 1.0 eq.) in ethanol (40 mL) was added 4 N HCl in dioxane (40 mL) slowly at 0 °C. The reaction was allowed to warm to r.t. and stirred for 2.5 h then concentrated and triturated with petroleum ether. A white solid (4.54 g, 92% yield) was collected and used in the next step.

Example 12. Synthesis of ethyl 2-((1R,3R)-3-((2S,3S)-2-azido-3-methylpentanamido)-1-hydroxy-4-methylpentyl)thiazole-4-carboxylate (**135**)




Azido-Ile-OH **124** (5.03g, 28.8 mmol, 2.0 eq.) was dissolved in THF (120 mL) and cooled to 0 °C, to which NMM (6.2 mL, 56.0 mmol, 4.0 eq.) and isobutylchloroformate (3.7 mL, 28.8 mmol, 2.0 eq.) were added in sequence. The reaction was stirred at 0 °C for 30 min and r.t. 1.0 h, and then cooled back to 0 °C. Compound **134** (4.54 g, 14.7 mmol, 1.0 eq.) was added in portions. After stirring at 0 °C for 30 min, the reaction was warmed to r.t. and stirred for 2 h. Water was added at 0 °C to quench the reaction and the resulting mixture was extracted with ethyl acetate for three times. The combined organic layers were washed with 1N HCl, saturated NaHCO₃ and brine, dried over anhydrous Na₂SO₄, filtered and concentrated. The residue was purified by column chromatography (0-30% EtOAc/PE) to give a white solid **135** (4.55 g, 74% yield).

Example 13. Synthesis of ethyl 2-((1R,3R)-3-((2S,3S)-2-azido-3-methylpentanamido)-4-methyl-1-((triethylsilyl)oxy)pentyl)thiazole-4-carboxylate (**136**)



To a solution of compound **135** (5.30 g, 12.8 mmol, 1.0 eq.) in CH₂Cl₂ (50 mL) was added imidazole (1.75 g, 25.6 mmol, 2.0 eq.), followed by chlorotriethylsilane (4.3 mL, 25.6 mmol, 2.0 eq.) at 0 °C. The reaction mixture was allowed to warm to r.t. over 1 hour and stirred for an additional hour. Brine was added to the reaction mixture, the organic layer was separated and the aqueous layer was extracted with EtOAc. The combined organic phases were dried, filtered, concentrated under reduced pressure, and purified by column chromatography with a gradient of 15-35% EtOAc in petroleum ether to afford product **136** (6.70 g, 99% yield) as a white solid. ¹H NMR (500 MHz, CDCl₃) δ 8.12 (s, 1H), 6.75 (d, *J* = 8.0 Hz, 1H), 5.20 – 5.12 (m, 1H), 4.44 (q, *J* = 7.0 Hz, 2H), 4.06 – 3.97 (m, 1H), 3.87 (d, *J* = 3.8 Hz, 1H), 2.14 (d, *J* = 3.8 Hz, 1H), 2.01 – 1.91 (m, 3H), 1.42 (t, *J* = 7.1 Hz, 3H), 1.34 – 1.25 (m, 2H), 1.06 (d, *J* = 6.8 Hz, 3H), 1.00 – 0.93 (m, 18H), 0.88 (dd, *J* = 19.1, 6.8 Hz, 6H). MS ESI *m/z* calcd for C₂₄H₄₄N₅O₄SSi [M+H]⁺ 526.28, found 526.28.


Example 14. Synthesis of ethyl 2-((1R,3R)-3-((2S,3S)-2-azido-N,3-dimethyl pentanamido)-4-methyl-1-((triethylsilyl)oxy)pentyl)thiazole-4-carboxylate (**137**)



A solution of **136** (5.20 g, 9.9 mmol, 1.0 eq.) in THF (50 mL) was cooled to -45 °C and KHMDS (1M in toluene, 23.8 mL, 23.8 mmol, 2.4 eq.) was added. The resulting mixture was stirred at -45°C for 20 min. Methyl iodide (1.85 mL, 29.7 mmol, 3.0 eq.) was then added, and the


reaction mixture was allowed to warm to r.t. over 4.5 h, at which time the reaction was quenched with EtOH (10 mL). The crude product was diluted with EtOAc (250 mL) and washed with brine (100 mL). The aqueous layer was extracted with EtOAc (3 × 50 mL). The organic layers were dried, filtered, concentrated and purified by column chromatography with a gradient of 15-35% EtOAc in petroleum ether to afford product **137** (3.33 g, 63% yield) as a light yellow oil. ¹H NMR (500 MHz, CDCl₃) δ 8.09 (s, 1H), 4.95 (d, *J* = 6.6 Hz, 1H), 4.41 (q, *J* = 7.1 Hz, 2H), 3.56 (d, *J* = 9.5 Hz, 1H), 2.98 (s, 3H), 2.27 – 2.06 (m, 4H), 1.83 – 1.70 (m, 2H), 1.41 (t, *J* = 7.2 Hz, 3H), 1.29 (ddd, *J* = 8.9, 6.8, 1.6 Hz, 3H), 1.01 (d, *J* = 6.6 Hz, 3H), 0.96 (dt, *J* = 8.0, 2.9 Hz, 15H), 0.92 (d, *J* = 6.6 Hz, 3H), 0.90 (d, *J* = 6.7 Hz, 3H). MS ESI *m/z* calcd for C₂₅H₄₆N₅O₄SSi [M+H]⁺ 540.30, found 540.30.

Example 15. Synthesis of ethyl 2-((3*S*,6*R*,8*R*)-3-((*S*)-sec-butyl)-10,10-diethyl-6-isopropyl-5-methyl-1-((*R*)-1-methylpiperidin-2-yl)-1,4-dioxo-9-oxa-2,5-diaza-10-siladodecan-8-yl)thiazole-4-carboxylate (**138**)




Dry Pd/C (10 wt%, 300 mg) and azide compound **137** (3.33 g, 6.61 mmol) were added to the above pentafluorophenyl ester **127** in EtOAc. The reaction mixture was stirred under hydrogen atmosphere for 27 h, and then filtered through a plug of Celite, with washing of the filter pad with EtOAc. The combined organic portions were concentrated and purified by column chromatography with a gradient of 0-5% methanol in EtOAc to deliver product **138** (3.90 g, 86% yield). MS ESI *m/z* calcd for C₃₂H₅₉N₄O₅SSi [M+H]⁺ 639.39, found 639.39.

Example 16. Synthesis of ethyl 2-((1*R*,3*R*)-3-((2*S*,3*S*)-N,3-dimethyl-2-((*R*)-1-methylpiperidine-2-carboxamido)pentanamido)-1-hydroxy-4-methylpentyl)thiazole-4-carboxylate (**139**)




The coupling product **138** (3.90 g, 6.1 mmol) was dissolved in deoxygenated AcOH/water/THF (v/v/v 3:1:1, 100 mL), and stirred at r.t. for 48 h. The reaction was then concentrated and purified by column chromatography (2:98 to 15:85 MeOH/EtOAc) to afford compound **139** (2.50 g, 72% yield over 2 steps). MS ESI *m/z* calcd for C₂₆H₄₅N₄O₅S [M+H]⁺ 525.30, found 525.33.

Example 17. Synthesis of 2-((1*R*,3*R*)-3-((2*S*,3*S*)-N,3-dimethyl-2-((*R*)-1-methylpiperidine-2-carboxamido)pentanamido)-1-hydroxy-4-methylpentyl)thiazole-4-carboxylic acid (**140**)




An aqueous solution of LiOH (0.4 N, 47.7 mL, 19.1 mmol, 4.0 eq.) was added to a solution of compound **139** (2.50 g, 4.76 mmol, 1.0 eq.) in dioxane (47.7 mL) at 0 °C. The reaction mixture was stirred at r.t. for 2 h and then concentrated. Column chromatography (100% CH₂Cl₂ then CH₂Cl₂/MeOH/NH₄OH 80:20:1) afforded compound **140** (2.36 g, 99% yield) as an amorphous solid. MS ESI m/z calcd for C₂₄H₄₁N₄O₅S [M+H]⁺ 497.27, found 497.28.

Example 18. Synthesis of 2-((1R,3R)-1-acetoxy-3-((2S,3S)-N,3-dimethyl-2-((R)-1-methylpiperidine-2-carboxamido)pentanamido)-4-methylpentyl)thiazole-4-carboxylic acid (**141**)




To a solution of compound **140** (2.36 g, 4.75 mmol) in pyridine (50 mL) at 0 °C, acetic anhydride (2.25 mL, 24 mmol) was added slowly. The reaction mixture was allowed to warm to r.t. over 2 h and stirred at r.t. for 24 h. The reaction was concentrated and the residue was purified by reverse phase HPLC (C₁₈ column, 10-90% acetonitrile/water) to afford compound **141** (2.25 g, 88% yield) as an amorphous white solid. MS ESI m/z calcd for C₂₆H₄₃N₄O₆S [M+H]⁺ 539.28, found 539.28.

Example 19. Synthesis of (1R,3R)-3-((2S,3S)-N,3-dimethyl-2-((R)-1-methylpiperidine-2-carboxamido)pentanamido)-4-methyl-1-(4-(perfluorobenzoyl)thiazol-2-yl)pentyl acetate (**142**)




To a solution of compound **141** (86 mg, 0.16 mmol, 1.0 eq.) in dichloromethane (2 mL) was added pentafluorophenol (44 mg, 0.24 mmol, 1.5 eq.) and *N,N'*-diisopropylcarbodiimide (22 mg, 0.175 mmol, 1.1 eq.) at 0 °C. The reaction mixture was warmed to room temperature and stirred over night. After the solvent was removed under reduced pressure, the reaction mixture was diluted with EtOAc (2 mL) then filtered over Celite. The filtrate was concentrated to afford crude pentafluorophenyl ester **142**, which was used directly without further purification.

Example 20. Synthesis of *tert*-butyl 3-(2-(2-(2-hydroxyethoxy)ethoxy)ethoxy) propanoate (**145**).




To a solution of 2,2'-(ethane-1,2-diylbis(oxy))diethanol (55.0 mL, 410.75 mmol, 3.0 eq.) in anhydrous THF (200 mL) was added sodium (0.1 g). The mixture was stirred until Na disappeared and then *tert*-butyl acrylate (20.0 mL, 137.79 mmol, 1.0 eq.) was added dropwise. The mixture was stirred overnight and then quenched by HCl solution (20.0 mL, 1N) at 0 °C. THF was removed by rotary evaporation, brine (300 mL) was added and the resulting mixture was extracted with EtOAc (3 × 100 mL). The organic layers were washed with brine (3 × 300 mL), dried over anhydrous Na₂SO₄, filtered and concentrated to afford a colorless oil (30.20 g, 79.0% yield), which was used without further purification. MS ESI m/z calcd for C₁₃H₂₇O₆ [M + H]⁺ 278.1729, found 278.1730.

Example 21. Synthesis of *tert*-butyl 3-(2-(2-(2-(tosyloxy)ethoxy)ethoxy)ethoxy) propanoate (146)




To a solution of *tert*-butyl 3-(2-(2-(2-hydroxyethoxy)ethoxy)ethoxy) propanoate (30.20 g, 108.5 mmol, 1.0 eq.) and TsCl (41.37 g, 217.0 mmol, 2.0 eq.) in anhydrous DCM (220 mL) at 0 °C was added TEA (30.0 mL, 217.0 mmol, 2.0 eq.). The mixture was stirred at room temperature overnight, and then washed with water (3 × 300 mL) and brine (300 mL), dried over anhydrous Na₂SO₄, filtered, concentrated and purified by SiO₂ column chromatography (3:1 hexanes/EtOAc) to give a colorless oil (39.4 g, 84.0% yield). MS ESI m/z calcd for C₂₀H₃₃O₈S [M + H]⁺ 433.1818, found 433.2838.

Example 22. Synthesis of *tert*-butyl 3-(2-(2-(2-azidoethoxy)ethoxy)ethoxy) propanoate (147)




To a solution of *tert*-butyl 3-(2-(2-(2-(tosyloxy)ethoxy)ethoxy)ethoxy) propanoate (39.4 g, 91.1 mmol, 1.0 eq.) in anhydrous DMF (100 mL) was added NaN₃ (20.67 g, 316.6 mmol, 3.5 eq.). The mixture was stirred at room temperature overnight. Water (500 mL) was added and extracted with EtOAc (3 × 300 mL). The combined organic layers were washed with water (3 × 900 mL) and brine (900 mL), dried over anhydrous Na₂SO₄, filtered, concentrated and purified by SiO₂ column chromatography (5:1 hexanes/EtOAc) to give a light yellow oil (23.8 g, 85.53% yield). MS ESI m/z calcd for C₁₃H₂₅O₃N₅Na [M + Na]⁺ 326.2, found 326.2.

Example 23. Synthesis of *tert*-butyl 3-(2-(2-(2-aminoethoxy)ethoxy)ethoxy) propanoate (148)




Raney-Ni (7.5 g, suspended in water) was washed with water (three times) and isopropyl alcohol (three times) and mixed with compound **147** (5.0 g, 16.5 mmol) in isopropyl alcohol. The mixture was stirred under a H₂ balloon at r.t. for 16 h and then filtered over a Celite pad, with washing of the pad with isopropyl alcohol. The filtrate was concentrated and purified by column chromatography (5-25% MeOH/DCM) to give a light yellow oil (2.60 g, 57% yield). MS ESI m/z calcd for C₁₃H₂₈NO₅ [M+H]⁺ 279.19; found 279.19.

Example 24. Synthesis of di-*tert*-butyl 14,17-dioxo-4,7,10,21,24,27-hexaoxa-13,18-diazatriacont-15-yne-1,30-dioate (**149**)




Acetylenedicarboxylic acid (0.35 g, 3.09 mmol, 1.0 eq.) was dissolved in NMP (10 mL) and cooled to 0 °C, to which compound **148** (2.06 g, 7.43 mmol, 2.4 eq.) was added, followed by DMTMM (2.39 g, 8.65 mmol, 2.8 eq.) in portions. The reaction was stirred at 0 °C for 6 h and then diluted with ethyl acetate and washed with water and brine. The organic solution was concentrated and triturated with a mixture solvent of ethyl acetate and petroleum ether. The solid was filtered off and the filtrate was concentrated and purified by column chromatography (80-90% EA/PE) to give a light yellow oil (2.26 g, >100% yield), which was used without further purification. MS ESI m/z calcd for C₃₀H₅₃N₂O₁₂ [M+H]⁺ 633.35; found 633.30.

Example 25. Synthesis of 14,17-dioxo-4,7,10,21,24,27-hexaoxa-13,18-diazatriacont-15-yne-1,30-dioic acid (**150**)



Compound **149** (2.26 g) was dissolved in dichloromethane (15 mL) and cooled to 0 °C then treated with TFA (15 mL). The reaction was warmed to r.t. and stirred for 45 min, and then the solvent and residual TFA was removed on rotovap. The crude product was purified by column chromatography (0-15% MeOH/DCM) to give a light yellow oil (1.39 g, 86% yield for two steps). MS ESI m/z calcd for C₂₂H₃₇N₂O₁₂ [M+H]⁺ 521.23; found 521.24.


Example 26. Synthesis of 2-(2-(dibenzylamino)ethoxy)ethanol (**152**).



2-(2-aminoethoxy)ethanol (21.00 g, 200 mmol, 1.0 eq.) and K₂CO₃ (83.00 g, 600 mmol, 3.0 eq.) in acetonitrile (350 mL) was added BnBr (57.0 mL, 480 mmol, 2.4 eq.). The mixture was refluxed overnight. Water (1 L) was added and extracted with EtOAc (3 × 300 mL). The combined organic layers were washed with brine (1000 mL), dried over anhydrous Na₂SO₄,


filtered, concentrated and purified by SiO₂ column chromatography (4:1 hexanes/ EtOAc) to give a colorless oil (50.97 g, 89.2% yield). MS ESI m/z calcd for C₁₈H₂₃NO₂Na [M + Na]⁺ 309.1729, found 309.1967.

Example 27. Synthesis of *tert*-butyl 3-(2-(2-(dibenzylamino)ethoxy)ethoxy) propanoate (153).




To a mixture of 2-(2-(dibenzylamino)ethoxy)ethanol (47.17 g, 165.3 mmol, 1.0 eq.) , *tert*-butyl acrylate (72.0 mL, 495.9 mmol, 3.0 eq.) and *n*-Bu₄NI (6.10 g, 16.53 mmol, 0.1 eq.) in DCM (560 mL) was added sodium hydroxide solution (300 mL, 50%). The mixture was stirred overnight. The organic layer was separated and the water layer was extracted with EtOAc (3 × 100 mL). The organic layers were washed with water(3 × 300 mL) and brine (300 mL), dried over anhydrous Na₂SO₄, filtered, concentrated and purified by SiO₂ column chromatography (7:1 hexanes/ EtOAc) to give a colorless oil (61.08 g, 89.4% yield). MS ESI m/z calcd for C₂₅H₃₆NO₄ [M + H]⁺ 414.2566, found 414.2384.

Example 28. Synthesis of *tert*-butyl 3-(2-(2-aminoethoxy)ethoxy)propanoate (154)




To a solution of *tert*-butyl 3-(2-(2-(dibenzylamino)ethoxy)ethoxy) propanoate (20.00 g, 48.36 mmol, 1.0 eq.) in THF (30 mL) and MeOH (60 mL) was added Pd/C (2.00 g, 10 wt%, 50% wet) in a hydrogenation bottle. The mixture was shaken overnight, filtered through Celite (filter aid), and the filtrate was concentrated to afford a colorless oil (10.58 g, 93.8% yield). MS ESI m/z calcd for C₁₁H₂₄NO₄ [M + H]⁺ 234.1627, found 234.1810.

Example 29. Synthesis of *tert*-butyl 3-(2-(2-hydroxyethoxy)ethoxy)propanoate (156)




To a solution of 2,2'-oxydiethanol (19.7 mL, 206.7 mmol, 3.0 eq.) in anhydrous THF (100 mL) was added sodium (0.1 g). The mixture was stirred until Na disappeared and then *tert*-butyl acrylate (10.0 mL, 68.9 mmol, 1.0 eq.) was added dropwise. The mixture was stirred overnight, and brine (200 mL) was added and extracted with EtOAc (3 × 100 mL). The organic layers were washed with brine (3 × 300 mL), dried over anhydrous Na₂SO₄, filtered, concentrated and purified by SiO₂ column chromatography (1:1 hexanes/ EtOAc) to give to a colorless oil (8.10 g, 49.4% yield). MS ESI m/z calcd for C₁₁H₂₃O₅ [M +H]⁺ 235.1467, found 235.1667.

Example 30. Synthesis of *tert*-butyl 3-(2-(2-(tosyloxy)ethoxy)ethoxy)propanoate (157)




To a solution of *tert*-butyl 3-(2-(2-hydroxyethoxy)ethoxy)propanoate (6.24 g, 26.63 mmol, 1.0 eq.) and TsCl (10.15 g, 53.27 mmol, 2.0 eq.) in anhydrous DCM (50 mL) at 0 °C was added pyridine (4.3 mL, 53.27 mmol, 2.0 eq.). The mixture was stirred at room temperature overnight, and then washed with water (100 mL) and the water layer was extracted with DCM (3 × 50 mL). The combined organic layers were washed with brine (300 mL), dried over anhydrous Na₂SO₄, filtered, concentrated and purified by SiO₂ column chromatography (5:1 hexanes/ EtOAc) to give a colorless oil (6.33 g, 61.3% yield). MS ESI m/z calcd for C₁₈H₂₇O₇S [M + H]⁺ 389.1556, found 389.2809.

Example 31. Synthesis of *tert*-butyl 3-(2-(2-azidoethoxy)ethoxy)propanoate (**158**)




To a solution of *tert*-butyl 3-(2-(2-(tosyloxy)ethoxy)ethoxy)propanoate (5.80 g, 14.93 mmol, 1.0 eq.) in anhydrous DMF (20 mL) was added NaN₃ (5.02 g, 77.22 mmol, 5.0 eq.). The mixture was stirred at room temperature overnight. Water (120 mL) was added and extracted with EtOAc (3 × 50 mL). The combined organic layers were washed with water (3 × 150 mL) and brine (150 mL), dried over anhydrous Na₂SO₄, filtered, concentrated and purified by SiO₂ column chromatography (5:1 hexanes/ EtOAc) to give a colorless oil (3.73 g, 69.6% yield). MS ESI m/z calcd for C₁₁H₂₂O₃N₄Na[M + H]⁺ 260.1532, found 260.2259.

Example 32. Synthesis of *tert*-butyl 3-(2-(2-aminoethoxy)ethoxy)propanoate (**159**)



Compound **158** (0.18 g, 0.69 mmol) was dissolved in MeOH (3.0 mL, with 60 μL concentrated HCl) and hydrogenated with Pd/C (10 wt%, 20 mg) under a H₂ balloon for 30 min. The catalyst was filtered through a celite pad, with washing of the pad with MeOH. The filtrate was concentrated to give a colorless oil (0.15 g, 93% yield). MS ESI m/z calcd for C₁₁H₂₄NO₄ [M+H]⁺ 234.16; found 234.14.


Example 33. Synthesis of di-*tert*-butyl 11,14-dioxo-4,7,18,21-tetraoxa-10,15 -diazatetracos-12-yne-1,24-dioate (**160**)



Acetylenedicarboxylic acid (26.5 mg, 0.232 mmol, 1.0 eq.) was dissolved in NMP (1.0 mL) and cooled to 0 °C, to which compound **159** (0.15 g, 0.557 mmol, 2.4 eq.) was added, followed by DMTMM (0.18 g, 0.65 mmol, 2.8 eq.) in portions. The reaction was stirred at 0 °C for 5 h and then diluted with ethyl acetate and washed with water and brine. The organic solution was concentrated and purified by column chromatography (80-90% EA/PE). The appropriate


fractions were concentrated and kept in refrigerator overnight. A solid formed and was filtered off. The filtrate was concentrated to give a light yellow oil (0.37 g, >100% yield), which was used without further purification. MS ESI m/z calcd for C₂₆H₄₅N₂O₁₀ [M+H]⁺ 545.30; found 545.30.

Example 34. Synthesis of 11,14-dioxo-4,7,18,21-tetraoxa-10,15-diaza tetracos-12-yne-1,24-dioic acid (**161**)




Compound **160** (0.21 g) was dissolved in dichloromethane (2.5 mL) and cooled to 0 °C then treated with TFA (2.5 mL). The reaction was warmed to r.t. and stirred for 45 min, and then the solvent and residual TFA was removed on rotovap. The crude product was purified by column chromatography (0-15% MeOH/DCM) to give a colorless oil (58.7 mg, 99% yield for two steps). MS ESI m/z calcd for C₁₈H₂₉N₂O₁₀ [M+H]⁺ 433.17; found 433.17.

Example 35. 3-(2-(2-azidoethoxy)ethoxy)propanoic acid (**165**)




The azide compound **158** (2.51 g, 9.68 mmol) dissolved in 1,4-dioxane (30 mL) was treated with 10 ml of HCl (conc.) at r.t. The mixture was stirred for 35 min, diluted with EtOH (30 ml) and toluene (30 ml) and concentrated under vacuum. The crude mixture was purified on silica gel using a mixture of methanol (from 5% to 10%) and 1% formic acid in methylene chloride as the eluant to give title compound **165** (1.63 g, 83% yield), ESI MS m/z C₇H₁₂N₃O₄ [M-H]⁻, calcd. 202.06, found 202.30.

Example 36. 2,5-dioxopyrrolidin-1-yl 3-(2-(2-azidoethoxy)ethoxy)propanoate (**166**)




To compound **165** (1.60 g, 7.87 mmol) in 30 mL of dichloromethane was added NHS (1.08 g, 9.39 mmol) and EDC (3.60 g, 18.75 mmol) with stirring. After 8 h TLC analysis revealed that the reaction was complete, the reaction mixture was concentrated and purified on silica gel using a mixture of ethyl acetate (from 5% to 10%) in methylene chloride as the eluant to give title compound **166** (1.93 g, 82% yield). ESI MS m/z C₁₁H₁₇N₄O₆ [M+H]⁺, calcd. 301.11, found 301.20.

Example 37. Synthesis of (E)-16-bromo-2,2-dimethyl-4,14-dioxo-3,7,10-trioxa -13-azaheptadec-15-en-17-oic acid (**167**)




To a solution of 3-bromofuran-2,5-dione (89 mg, 0.5 mmol) in THF (5 mL), *tert*-butyl 3-(2-(2-aminoethoxy)ethoxy)propanoate (117 mg, 0.5 mmol) was added. The resulting solution was stirred at r.t. for 4 h. The solvent was removed under vacuum to afford compound **167** (205 mg, theoretical yield). MS ESI *m/z* calcd for $C_{15}H_{25}BrNO_7$ $[M+H]^+$ 410.07, found 410.03.

Example 38. Synthesis of (E)-1-*tert*-butyl 18-methyl 13-bromo-11,14-dioxo-4,7-dioxo-10,15-diazaoctadec-12-ene-1,18-dioate (**169**)




Compound **167** (205 mg, 0.5 mmol) and methyl 3-aminopropanoate hydrochloride (70 mg, 0.5 mmol) were dissolved in DCM (20 mL), to which DIPEA (0.26 mL, 1.5 mmol) and EDC (144 mg, 0.75 mmol) were added. The resulting solution was stirred at r.t. overnight, and then washed with brine (50 mL), dried over anhydrous Na_2SO_4 . Concentration and purification by column chromatography (0 to 10% MeOH/DCM) yielded compound **169** (88 mg, 36% yield). MS ESI *m/z* calcd for $C_{19}H_{32}BrN_2O_8$ $[M+H]^+$ 495.13, found 495.25.

Example 39. Synthesis of (E)-8-bromo-3,7,10-trioxo-2,14,17-trioxa-6,11-diaza icos-8-en-20-oic acid (**170**)




Compound **169** (88 mg, 0.18 mmol) in DCM (3 mL) was treated with formic acid (6 ml) at 38 °C overnight. All volatiles were removed under vacuum to yield compound **170** (about 78 mg, ~90% yield).

Example 40. Synthesis of 4-(((benzyloxy)carbonyl)amino)butanoic acid (**172**)




A solution of 4-aminobutyric acid (7.5 g, 75 mmol) and NaOH (6 g, 150 mmol) in H_2O (40 mL) was cooled to 0 °C and treated with a solution of CbzCl (16.1 g, 95 mmol) in THF (32 mL) dropwise. After 1 h, the reaction was allowed to warm to r.t. and stirred for 3 h. THF was removed under vacuum, the pH of the aqueous solution was adjusted to 1.5 by addition of 6 N HCl. Extracted with ethyl acetate, and the organic layer was washed with brine, dried and concentrated to give compound **172** (16.4 g, 92% yield). MS ESI *m/z* calcd for $C_{12}H_{16}NO_5$ $[M+H]^+$ 238.10, found 238.08.

Example 41. Synthesis of *tert*-butyl 4-(((benzyloxy)carbonyl)amino)butanoate (**173**)




DMAP (0.8 g, 6.56 mmol) and DCC (17.1 g, 83 mmol) were added to a solution of 4-(((benzyloxy)carbonyl)amino)butanoic acid (16.4 g, 69.2 mmol) and *t*-BuOH (15.4 g, 208 mmol) in DCM (100 mL). After stirring at r.t. overnight, the reaction was filtered and filtrate concentrated. The residue was dissolved in ethyl acetate and the washed with 1N HCl, brine and dried over Na₂SO₄. Concentration and purification by column chromatography (10 to 50% EtOAc/hexanes) yielded compound **173** (7.5 g, 37% yield). MS ESI *m/z* calcd for C₁₆H₂₃NO₄Na [M+Na]⁺ 316.16, found 316.13.

Example 42. Synthesis of *tert*-butyl 4-aminobutanoate (**174**)




tert-Butyl 4-(((benzyloxy)carbonyl)amino)butanoate (560 mg, 1.91 mmol) was dissolved in MeOH (50 mL), and mixed with Pd/C catalyst (10 wt%, 100 mg) then hydrogenated (1 atm) at room temperature for 3 h. The catalyst was filtered off and all volatiles were removed under vacuum to afford compound **174** (272 mg, 90% yield). MS ESI *m/z* calcd for C₈H₁₈NO₂ [M+H]⁺ 160.13, found 160.13.

Example 43. Synthesis of (E)-2-bromo-4-((4-(*tert*-butoxy)-4-oxobutyl)amino)-4-oxobut-2-enoic acid (**175**)



3-Bromofuran-2,5-dione (300 mg, 1.71 mmol) was dissolved in THF (20 mL), to which *tert*-butyl 4-aminobutanoate (272 mg, 1.71 mmol) was added and the resulting solution was stirred at r.t. for 3 h. The solvent was removed under vacuum to afford compound **175** (572 mg, theoretical yield). MS ESI *m/z* calcd for C₁₂H₁₉BrNO₅ [M+H]⁺ 338.03, found 338.04.


Example 44. Synthesis of (E)-*tert*-butyl 4-(3-bromo-4-((2-methoxyethyl) amino)-4-oxobut-2-enamido)butanoate (**177**)



2-Bromo-4-((4-(*tert*-butoxy)-4-oxobutyl)amino)-4-oxobut-2-enoic acid (286mg, 0.85mmol) and 2-methoxyethanamine (128mg, 1.7mmol) were dissolved in DCM (40 mL), to which DIPEA (329 mg, 2.55 mmol) and EDC (490 mg, 2.55 mmol) were added. The resulting solution was stirred at r.t. for 24 h and then washed with brine, dried over Na₂SO₄. Concentration and


purification by column chromatography (0 to 10% MeOH/DCM) yielded compound **177** (102 mg, 31% yield). MS ESI m/z calcd for $C_{15}H_{26}BrN_2O_5$ $[M+H]^+$ 393.09, found 393.11.

Example 45. Synthesis of (E)-4-(3-bromo-4-((2-methoxyethyl)amino)-4-oxobut-2-enamido)butanoic acid (**178**)




Compound **177** (52 mg, 0.132 mmol) was dissolved in DCM (3 mL), to which formic acid (6 ml) was added. The resulting solution was stirred at 38 °C overnight then concentrated to afford compound **178** (45 mg, theoretical yield). MS ESI m/z calcd for $C_{11}H_{18}BrN_2O_5$ $[M+H]^+$ 339.03, found 339.05.

Example 46. Synthesis of (E)-2,5-dioxopyrrolidin-1-yl 4-(3-bromo-4-((2-methoxyethyl)amino)-4-oxobut-2-enamido)butanoate (**179**)




To a solution of compound **178** (45 mg, 0.132 mmol) in DCM (10 ml) NHS (23 mg, 0.199 mmol) and EDC (38 mg, 0.199 mmol) were added. After stirring at r.t. for 3 h, the reaction was concentrated and purified by column chromatography (10 to 50% EtOAc/hexanes) to yield compound **179** (57 mg, 99% yield). MS ESI m/z calcd for $C_{15}H_{21}BrN_3O_7$ $[M+H]^+$ 436.05, found 436.06.

Example 47. Synthesis of (Z)-di-*tert*-butyl 4,4'-((2-bromofumaroyl)bis(azane diyl))dibutanoate (**180**)




tert-Butyl 4-aminobutanoate (**174**) (477 mg, 3 mmol) and 2,3-dibromosuccinic acid (414 mg, 1.5 mmol) was dissolved in DCM (35 mL), to which DIPEA (1.16 g, 9 mmol) and EDC (0.86 g, 4.5 mmol) were added. The resulting solution was stirred at r.t. overnight and then washed with brine, dried over Na_2SO_4 . Filtration, concentration and purification by column chromatography (0 to 10% MeOH/DCM) yielded compound **180** (160 mg, 22% yield). MS ESI m/z calcd for $C_{20}H_{34}BrN_2O_6$ $[M+H]^+$ 477.15, found 477.16.

Example 48. Synthesis of (Z)-4,4'-((2-bromofumaroyl)bis(azanediyl))dibutanoic acid (**181**)




Compound **180** (80 mg, 0.168 mmol) was dissolved in DCM (5 mL) and treated with formic acid (8 mL) at 38 °C overnight. All volatiles were removed under vacuum to afford compound **181** (61 mg, 99% yield). MS ESI m/z calcd for $C_{12}H_{18}BrN_2O_6$ $[M+H]^+$ 365.03, found 365.05.

5 Example 49. Synthesis of (Z)-bis(2,5-dioxopyrrolidin-1-yl) 4,4'-((2-bromo fumaroyl)bis(azanediyl))dibutanoate (**182**)



10 NHS (60 mg, 0.504 mmol) and EDC (97 mg, 0.504 mmol) were added to a solution of compound **181** (61 mg, 0.168 mmol) in DCM (10 mL). After stirring at r.t. overnight, the reaction mixture was concentrated and purified by column chromatography (0 to 10% MeOH/DCM) to afford compound **182** (72 mg, 77% yield). MS ESI m/z calcd for $C_{20}H_{24}BrN_4O_{10}$ $[M+H]^+$ 559.06, found 559.78.


Example 50. Synthesis of *tert*-butyl 2-(triphenylphosphoranylidene)propanoate (**184**)



15 A mixture of *tert*-butyl-2-bromopropanoate **183** (15.5 g, 74.1 mmol, 1.0 eq.) and triphenyl phosphine (19.4 g, 74.1 mmol, 1.0 eq.) in dry acetonitrile (45 mL) was stirred at room temperature for 18 h. Acetonitrile was removed under reduced pressure and toluene was added to crash out a white precipitate. Toluene was then decanted off and the white solid was dissolved in dichloromethane (100 mL) and transferred to a separatory funnel. 10% NaOH (100 mL) was

20 added to the funnel, and the organic layer immediately turned yellow after shaking. The organic layer was separated and the aqueous layer was extracted with dichloromethane (30 mL) once. The dichloromethane layers were combined and washed with brine (50 mL) once, then dried over Na_2SO_4 , filtered and concentrated, giving the ylide as a yellow solid (16.8 g, 58%).


25 Example 51. Synthesis of (S)-methyl 3-(4-(benzyloxy)phenyl)-2-((*tert*-butoxy carbonyl)amino)propanoate (**186**)



To a mixture of Boc-L-Tyr-OMe (20.0 g, 67.7 mmol, 1.0 eq.), K_2CO_3 (14.0 g, 101.6 mmol, 1.5 eq.) and KI (1.12 g, 6.77 mmol, 0.1 eq.) in acetone (100 mL) was added BnBr (10.5 mL, 81.3


mmol, 1.2 eq.) slowly. The mixture was then refluxed overnight. Water (250 mL) was added and the reaction mixture was extracted with EtOAc (3×100 mL). The combined organic layers were washed with brine (300 mL), dried over anhydrous Na₂SO₄, filtered, concentrated and purified by SiO₂ column chromatography (4:1 hexanes/EtOAc) to give a white solid **186** (26.12 g, 99% yield). ¹H NMR (500 MHz, CDCl₃) δ 7.44 – 7.41 (m, 2H), 7.41 – 7.36 (m, 2H), 7.35 – 7.30 (m, 1H), 7.04 (d, *J* = 8.5 Hz, 2H), 6.93 – 6.89 (m, 2H), 5.04 (s, 2H), 4.97 (d, *J* = 7.7 Hz, 1H), 4.55 (d, *J* = 6.9 Hz, 1H), 3.71 (s, 3H), 3.03 (dd, *J* = 14.4, 5.7 Hz, 2H), 1.44 (d, *J* = 18.6 Hz, 10H). MS ESI *m/z* calcd for C₂₂H₂₇NO₅Na [M+Na]⁺ 408.18, found 408.11.

Example 52. Synthesis of (S)-*tert*-butyl (1-(4-(benzyloxy)phenyl)-3-oxopropan-2-yl)carbamate (**187**)




To a solution of ester **186** (26.1 g, 67.8 mmol, 1.0 eq.) in anhydrous dichloromethane (450 mL) at -78 °C was added DIBAL (1.0 M in hexanes, 163 mL, 2.2 eq.) in 1 h. The mixture was stirred at -78 °C for 3 h and then quenched with 50 mL of ethanol. 1N HCl was added dropwise until pH 4 was reached. The resulting mixture was allowed to warm to 0 °C. Layers were separated and the aqueous layer was further extracted with EtOAc (3 × 100 mL). The combined organic solution was washed with brine, dried over anhydrous Na₂SO₄, and concentrated. Trituration with PE/EtOAc and filtration gave a white solid **187** (18.3 g, 76% yield). MS ESI *m/z* calcd for C₂₂H₂₇NO₅Na [M+Na]⁺ 378.11, found 378.11.

Example 53. Synthesis of (S,Z)-*tert*-butyl 5-(4-(benzyloxy)phenyl)-4-((*tert*-butoxycarbonyl)amino)-2-methylpent-2-enoate (**188**)




Aldehyde **187** (0.84 g, 2 mmol, 1.0 eq.) was dissolved in dry dichloromethane (50 mL), to which *tert*-butyl ester ylide **184** (1.6 g, 4 mmol, 2.0 eq.) was added and the solution was stirred at r.t. for 1.5 h as determined complete by TLC. Purification by column chromatography (10-50% EtOAc/hexanes) afforded compound **188** (1.16g, 98% yield).

Example 54. Synthesis of (4R)-*tert*-butyl 4-((*tert*-butoxycarbonyl)amino)-5-(4-hydroxyphenyl)-2-methylpentanoate (**189**)




Compound **188** (467 mg, 1 mmol) was dissolved in methanol (30 mL) and hydrogenated (1 atm) with Pd/C catalyst (10 wt%, 250 mg) at r.t. overnight. The catalyst was filtered off and the filtrate were concentrated under reduced pressure to afford compound **189** (379mg, 99% yield).

Example 55. Synthesis of (4R)-tert-butyl 4-((tert-butoxycarbonyl)amino)-5-(4-hydroxy-3-nitrophenyl)-2-methylpentanoate (**190**)




Compound **189** (379mg, 1 mmol, 1.0 eq.) was dissolved in THF (20 mL), to which a solution of *tert*-butyl nitrite (315 mg, 3 mmol, 3.0 eq.) in THF (2 mL) was added. The reaction was stirred at r.t. for 3 h and then poured onto water, extracted with EtOAc (2 × 50 mL) and the combined organic phases were washed with brine (50 mL), dried over anhydrous Na₂SO₄, filtered and concentrated. Purification by column chromatography (10-50% EtOAc/hexanes) afforded compound **190** (300 mg, 71% yield).

Example 56. Synthesis of (4R)-tert-butyl 4-((tert-butoxycarbonyl)amino)-5-(4-((tert-butyldimethylsilyl)oxy)-3-nitrophenyl)-2-methylpentanoate (**191**)




To a solution of compound **190** (424 mg, 1 mmol) in DCM (20 mL), imidazole (408 mg, 6 mmol) and *tert*-butylchlorodimethylsilane (602 mg, 4 mmol) were added. The resulting solution was stirred at r.t. for 3 h. Afterwards, the reaction mixture was washed with brine (50 mL), dried over anhydrous Na₂SO₄, concentrated and purified by column chromatography (10% to 30% EtOAc/hexanes) to yield compound **191** (344 mg, 64% yield).

Example 57. Synthesis of (4R)-tert-butyl 5-(3-amino-4-((tert-butyldimethylsilyl)oxy)phenyl)-4-((tert-butoxycarbonyl)amino)-2-methylpentanoate (**192**)




Compound **191** (200 mg, 0.37 mmol) was dissolved in EtOAc (30 mL), mixed with palladium catalyst (10 wt% on carbon, 100 mg) and hydrogenated (1 atm) at r.t. for 2 h. The catalyst was filtered off and all volatiles were removed under vacuum to afford compound **192** (187 mg, 99% yield).

Example 58. Synthesis of (E)-methyl 13-bromo-1-((5-((2R)-5-(tert-butoxy)-2-((tert-butoxycarbonyl)amino)-4-methyl-5-oxopentyl)-2-((tert-butyldimethylsilyl)oxy)phenyl)-amino)-1,11,14-trioxo-4,7-dioxa-10,15-diazaoctadec-12-en-18-oate (**193**)




Compound **192** (90 mg, 0.178 mmol) and compound **170** (78 mg, 0.178 mmol) were dissolved in DCM (5 mL) and DMA (5 mL). To which TEA (36 mg, 0.356 mmol) and HATU (102 mg, 0.267 mmol) were added. The resulting mixture was stirred at r.t. for 4 h, and then diluted with EA and washed with brine, dried over anhydrous Na₂SO₄. Filtration and purification by column chromatography (0 to 10% MeOH/DCM) yielded compound **193** (87 mg, 53% yield).

Example 59. Synthesis of (E)-methyl 13-bromo-1-((5-((2R)-5-(tert-butoxy)-2-((tert-butoxycarbonyl)amino)-4-methyl-5-oxopentyl)-2-hydroxyphenyl)amino)-1,11,14-trioxo-4,7-dioxa-10,15-diazaoctadec-12-en-18-oate (**194**)




Compound **193** (87 mg, 0.0936 mmol) was dissolved in DCM (10 mL) and treated with TBAF (1M in THF, 0.3 mL). The resulting mixture was stirred at r.t. for 0.5 h and washed with brine, dried over anhydrous Na₂SO₄. Filtration and purification by column chromatography (0 to 10% MeOH/DCM) yielded compound **194** (53 mg, 69% yield). MS ESI m/z calcd for C₃₆H₅₆BrN₄O₁₂ [M+H]⁺ 817.30, found 817.51.

Example 60. Synthesis of (4R)-4-amino-5-(3-((E)-8-bromo-3,7,10-trioxo-2,14, 17-trioxa-6,11-diazaicos-8-enamido)-4-hydroxyphenyl)-2-methylpentanoic acid (**195**)




Compound **194** (18 mg, 0.022 mmol) was dissolved in DCM (2 mL) and treated with TFA (2 mL) at r.t. for 3 h. All volatiles were removed under vacuum to afford compound **195** (14.5 mg, theoretical yield). MS ESI m/z calcd for C₂₇H₄₀BrN₄O₁₀ [M+H]⁺ 659.18, found 659.18.

Example 61. Synthesis of (4R)-4-(2-((1R,3R)-1-acetoxy-3-((2S,3S)-N,3-dimethyl-2-((R)-1-methylpiperidine-2-carboxamido)pentanamido)-4-methylpentyl)thiazole-4-carboxamido)-5-(3-((E)-8-bromo-3,7,10-trioxo-2,14,17-trioxo-6,11-diazaicos-8-enamido)-4-hydroxyphenyl)-2-methylpentanoic acid (**196**)




To a solution of compound **195** (14.5 mg, 0.022 mmol) and perfluorophenyl ester **142** (18.5 mg, 0.026 mmol) in DMA (2.5 mL) was DIPEA (15 mg, 0.116 mmol) added. The resulting mixture was stirred at r.t. for 1.5 h and then the solvent was removed under vacuum. The residue was purified on reverse phase preparative HPLC (C_{18} column, 10-90% MeCN/ H_2O) to afford compound **196** (12.6 mg, 49% yield). MS ESI m/z calcd for $C_{53}H_{80}BrN_8O_{15}$ $[M+H]^+$ 1181.45, found 1181.45.

Example 62. Synthesis of (4R)-*tert*-butyl 5-(3-amino-4-hydroxyphenyl)-4-((*tert*-butoxycarbonyl)amino)-2-methylpentanoate (**197**)




Compound **190** (56 mg, 0.132 mmol) was dissolved in EtOAc (20 mL) and mixed with Pd/C catalyst (10 wt%, 50 mg) and hydrogenated (1 atm) at r.t. for 3 h. The catalyst was filtered off and all volatiles were removed under vacuum to afford compound **197** (52 mg, 99% yield). MS ESI m/z calcd for $C_{21}H_{35}N_2O_5$ $[M+H]^+$ 395.25, found 395.26.

Example 63. Synthesis of (4R)-*tert*-butyl 5-(3-(4-((E)-3-bromo-4-((2-methoxyethyl)amino)-4-oxobut-2-enamido)butanamido)-4-hydroxyphenyl)-4-((*tert*-butoxycarbonyl)amino)-2-methylpentanoate (**198**)




NaH_2PO_4 (0.1M in water, 2 ml) was added to a solution of compound **179** (57 mg, 0.1323 mmol) and compound **197** (52 mg, 0.1323 mmol) in EtOH (10 mL). The resulting solution was stirred at r.t. for 72 h, concentrated and purified by column chromatography (30 to 100% EtOAc/hexanes) to yield compound **198** (50 mg, 53% yield). MS ESI m/z calcd for $C_{32}H_{50}BrN_4O_9$ $[M+H]^+$ 715.27, found 715.27.

Example 64. Synthesis of (4R)-4-amino-5-(3-(4-((E)-3-bromo-4-((2-methoxy ethyl) amino)-4-oxobut-2-enamido)butanamido)-4-hydroxyphenyl)-2-methylpentanoic acid (**199**)




Compound **198** (25 mg, 0.035 mmol) in DCM (2 mL) was treated with TFA (2mL) at r.t. for 1 h. All volatiles were removed under vacuum to afford compound **199** (19 mg, 99% yield). MS ESI m/z calcd for C₂₃H₃₄BrN₄O₇ [M+H]⁺ 557.15, found 557.27.

Example 65. Synthesis of (4R)-4-(2-((1R,3R)-1-acetoxy-3-((2S,3S)-N,3- dimethyl-2-((R)-1-methylpiperidine-2-carboxamido)pentanamido)-4-methylpentyl)thiazole-4-carboxamido)-5-(3-(4-((E)-3-bromo-4-((2-methoxyethyl)amino)-4-oxobut-2-enamido)butanamido)-4-hydroxyphenyl)-2-methylpentanoic acid (**200**)



Compound **199** (19 mg, 0.035 mmol) and perfluorophenyl ester **142** (33 mg, 0.046 mmol) was dissolved in DMA (5 mL), to which DIPEA (10mg, 0.077 mmol) was added. The reaction mixture was stirred at r.t. for 2 h then concentrated and purified on preparative HPLC (C₁₈ column, 10-90% MeCN/H₂O) to afford the compound **200** (9 mg, 24% yield). MS ESI m/z calcd for C₄₉H₇₄BrN₈O₁₂S [M+H]⁺ 1079.43, found 1079.66.


Example 66. Synthesis of (4R)-tert-butyl 5-(3-((Z)-79-bromo-1-hydroxy-73,78, 81-trioxo-3,6,9,12,15,18,21,24,27,30,33,36,39,42,45,48,51,54,57,60,63,66,69-tricosaoxa-72,77,82-triazahexaoctacont-79-enamido)-4-hydroxyphenyl)-4-((tert-butoxycarbonyl)amino)-2-methylpentanoate (**203**)



NaH₂PO₄ (0.1M in water, 1 mL) was added to a solution of compound **182** (36 mg, 0.065 mmol) and compound **197** (25 mg, 0.063 mmol) in EtOH (5 mL). The resulting solution was stirred at r.t. overnight and then HO-(PEG)₂₄-NH₂ (95 mg) was added to the mixture and stirred at r.t. overnight. All volatiles were removed under vacuum and the residue was purified by


column chromatography (0 to 10% MeOH/DCM) to yield compound **203** (28 mg, 24% yield). MS ESI m/z 1798.93 ([M+H]⁺).

Example 67. Synthesis of (4R)-4-amino-5-(3-((Z)-79-bromo-1-hydroxy-73,78, 81-trioxo-3,6,9,12,15,18,21,24,27,30,33,36,39,42,45,48,51,54,57,60,63,66,69-tricosaoxa-72,77,82-triazahexaoctacont-79-enamido)-4-hydroxyphenyl)-2-methylpentanoic acid (**204**)




Compound **203** (28 mg, 0.0156 mmol) was dissolved in DCM (2 mL) and treated with TFA (2 mL) at r.t. for 2 h. All volatiles were removed under vacuum to afford compound **204** (25 mg, 98% yield). MS ESI m/z 1642.82 ([M+H]⁺).

Example 68. Synthesis of (4R)-4-(2-((1R,3R)-1-acetoxy-3-((2S,3S)-N,3 -dimethyl-2-((R)-1-methylpiperidine-2-carboxamido)pentanamido)-4-methylpentyl)thiazole-4-carboxamido)-5-(3-((Z)-79-bromo-1-hydroxy-73,78,81-trioxo-3,6,9,12,15,18,21,24,27,30,33,36,39,42,45,48,51,54,57,60,63,66,69-tricosaoxa-72,77,82-triazahexaoctacont-79-enamido)-4-hydroxyphenyl)-2-methylpentanoic acid (**205**)



Compound **204** (25 mg, 0.0152 mmol) and perfluorophenyl ester **142** (15 mg, 0.0213 mmol) were dissolved in DMA (5 mL). To the mixture, DIPEA (10 mg, 0.077 mmol) was added. The resulting mixture was stirred at r.t. overnight, concentrated and purified by preparative HPLC (C₁₈ column, 10-90% MeCN/H₂O) to afford compound **205** (13 mg, 40% yield). MS ESI m/z 2163.82 ([M+H]⁺).


Example 69. Synthesis of (S)-tert-butyl 2-(hydroxymethyl)pyrrolidine-1-carboxylate (**208**)



Boc-L-proline (10.0 g, 46.4 mmol) dissolved in 50 mL THF was cooled to 0 °C, to which BH₃ in THF (1.0 M, 46.4 mL) was added carefully. The mixture was stirred at 0 °C for 1.5 h then poured onto ice water and extracted with ethyl acetate. The organic layer was washed with brine (50 mL), dried over anhydrous Na₂SO₄, and concentrated under reduced pressure to give known compound **208** (8.50 g, 91% yield) as a white solid. ¹H NMR (500 MHz, CDCl₃) δ 3.94 (dd, J =


4.9, 2.7 Hz, 2H), 3.60 (ddd, $J = 18.7, 11.9, 9.3$ Hz, 2H), 3.49–3.37 (m, 1H), 3.34–3.23 (m, 1H), 2.06–1.91 (m, 1H), 1.89–1.69 (m, 2H), 1.65–1.51 (m, 1H), 1.49–1.40 (m, 9H).

Example 70. Synthesis of (S)-*tert*-butyl 2-formylpyrrolidine-1-carboxylate (**209**)




To a solution of alcohol **208** (13.0 g, 64.6 mmol) in dimethyl sulfoxide (90 mL) was added triethylamine (40 mL) and the stirring was continued for 15 min. The mixture was cooled over ice bath and sulfur trioxide-pyridine complex (35.98 g, 226 mmol) was added in portions over a 40 min period. The reaction was warmed to r.t. and stirred for 2.5 h. After addition of ice (250 g), the mixture was extracted with dichloromethane (150 mL $\times$ 3). The organic phase was washed with 50% citric acid solution (150 mL), water (150 mL), saturated sodium bicarbonate solution (150 mL), and brine (150 mL), dried over anhydrous Na_2SO_4 . Removal of solvent in vacuo yielded aldehyde **209** (10.4 g, 81% yield) as a dense oil which was used without further purification. ^1H NMR (500 MHz, CDCl_3) δ 9.45 (s, 1H), 4.04 (s, 1H), 3.53 (dd, $J = 14.4, 8.0$ Hz, 2H), 2.00 – 1.82 (m, 4H), 1.44 (d, $J = 22.6$ Hz, 9H).

Example 71. Synthesis of (4R,5S)-4-methyl-5-phenyl-3-propionyloxazolidin-2-one (**211**)




n-Butyllithium in hexane (21.6 mL, 2.2 M, 47.43 mmol) was added dropwise at -78°C to a stirred solution of 4-methyl-5-phenyloxazolidin-2-one (8.0 g, 45.17 mmol) in THF (100 mL) under N_2 . The solution was maintained at -78°C for 1 h then propionyl chloride (4.4 mL, 50.59 mmol) was added slowly. The reaction mixture was warmed to -50°C , stirred for 2 h then quenched by addition of a saturated solution of ammonium chloride (100 mL). The organic solvent was removed in vacuo and the resultant solution was extracted with ethyl acetate (3 $\times$ 100 mL). The organic layer was washed with saturated sodium bicarbonate solution (100 mL) and brine (100 mL), dried over Na_2SO_4 , filtered and concentrated in vacuo. The residue was purified by column chromatography (20% ethyl acetate/hexanes) to afford the compound **211** as a dense oil (10.5 g, 98% yield). ^1H NMR (500 MHz, CDCl_3) δ 7.45 – 7.34 (m, 3H), 7.30 (d, $J = 7.0$ Hz, 2H), 5.67 (d, $J = 7.3$ Hz, 1H), 4.82 – 4.70 (m, 1H), 2.97 (dd, $J = 19.0, 7.4$ Hz, 2H), 1.19 (t, $J = 7.4$ Hz, 3H), 0.90 (d, $J = 6.6$ Hz, 3H).

Example 72. Synthesis of (S)-*tert*-butyl 2-((1R,2R)-1-hydroxy-2-methyl-3-((4R,5S)-4-methyl-2-oxo-5-phenyloxazolidin-3-yl)-3-oxopropyl)pyrrolidine-1-carboxylate (**212**)




To a solution of **209** (9.40 g, 40.4 mmol) in dichloromethane (60 mL) was added Et₃N (6.45 mL, 46.64 mmol) at 0 °C, followed by 1M dibutylboron triflate in dichloromethane (42 mL, 42 mmol). The mixture was stirred at 0 °C for 45 min, cooled to -70 °C, aldehyde **211** (4.58 g, 22.97 mmol) in dichloromethane (40 mL) was then added slowly over a 30 min period. The reaction was stirred at -70 °C for 2 h, 0 °C 1 h, and r.t. 15 min, and then quenched with phosphate buffer solution (pH 7, 38 mL). After the addition of MeOH-30% H₂O₂ (2:1, 100 mL) at below 10 °C and stirring for 20 min, water (100 mL) was added and the mixture was concentrated in vacuo. More water (200 mL) was added to the residue and the mixture was extracted with ethyl acetate (3 × 100 mL). The organic layer was washed with 1N KHSO₄ (100 mL), sodium bicarbonate solution (100 mL) and brine (100 mL), dried over anhydrous Na₂SO₄ and concentrated in vacuo. The residue was purified by flash column chromatography (10% - 50% ethyl acetate/hexanes) to afford the compound **212** as a white solid (7.10 g, 71% yield). ¹H NMR (500 MHz, CDCl₃) δ 7.39 (dt, *J* = 23.4, 7.1 Hz, 3H), 7.30 (d, *J* = 7.5 Hz, 2H), 5.67 (d, *J* = 7.1 Hz, 1H), 4.84 – 4.67 (m, 1H), 4.08 – 3.93 (m, 3H), 3.92 – 3.84 (m, 1H), 3.50 (d, *J* = 9.0 Hz, 1H), 3.24 (d, *J* = 6.7 Hz, 1H), 2.15 (s, 1H), 1.89 (dd, *J* = 22.4, 14.8 Hz, 3H), 1.48 (d, *J* = 21.5 Hz, 9H), 1.33 (d, *J* = 6.9 Hz, 3H), 0.88 (d, *J* = 6.4 Hz, 3H).

Example 73. Synthesis of (S)-*tert*-butyl 2-((1R,2R)-1-methoxy-2-methyl-3-((4R,5S)-4-methyl-2-oxo-5-phenyloxazolidin-3-yl)-3-oxopropyl)pyrrolidine-1-carboxylate (**213**)




To a mixture of **212** (5.1 g 11.9 mmol) and molecular sieves (4 Å, 5 g) was added anhydrous dichloroethane (30 mL) under N₂. The mixture was stirred at room temperature for 20 min and cooled to 0 °C. Proton sponge (6.62 g, 30.9 mmol) was added, followed by trimethyloxonium tetrafluoroborate (4.40 g, 29.7 mmol). Stirring was continued for 2 h at 0 °C and 48 h at r.t. The reaction mixture was filtrated and the filtrate was concentrated and purified by column chromatography (20-70% ethyl acetate/hexanes) to afford compound **213** as a white solid (1.80 g, 35% yield). ¹H NMR (500 MHz, CDCl₃) δ 7.46 – 7.27 (m, 5H), 5.65 (s, 1H), 4.69 (s, 1H), 3.92 (s, 1H), 3.83 (s, 1H), 3.48 (s, 3H), 3.17 (s, 2H), 2.02 – 1.68 (m, 5H), 1.48 (d, *J* = 22.3 Hz, 9H), 1.32 (t, *J* = 6.0 Hz, 3H), 0.91 – 0.84 (m, 3H).

Example 74. Synthesis of (2R,3R)-3-((S)-1-(*tert*-butoxycarbonyl)pyrrolidin-2-yl) -3-methoxy -2-methylpropanoic acid (**214**)



To a solution of **213** (1.80 g, 4.03 mmol) in THF (30 mL) and H₂O (7.5 mL), 30% H₂O₂ (1.44 mL, 14.4 mmol) was added over a 5 min period at 0 °C, followed by a solution of LiOH (0.27 g, 6.45 mmol) in water (5 mL). After stirring at 0 °C for 3 h, 1 N sodium sulfite (15.7 mL) was added and the mixture was allowed to warm to r.t. and stirred overnight. THF was removed in vacuo and the aqueous phase was wash with dichloromethane (3 × 50 mL) to remove the oxazolidinone auxiliary. The aqueous phase was acidified to pH 3 with 1N HCl and extracted with ethyl acetate (3 × 50 mL). The organic layer was washed with brine (50 mL), dried over Na₂SO₄, filtered and concentrated in vacuo to afford the compound **214** as a colorless oil (1.15 g, 98% yield). ¹H NMR (500 MHz, CDCl₃) δ 3.99 – 3.74 (m, 2H), 3.44 (d, J = 2.6 Hz, 3H), 3.23 (s, 1H), 2.60 – 2.45 (m, 1H), 1.92 (tt, J = 56.0, 31.5 Hz, 3H), 1.79 – 1.69 (m, 1H), 1.58 – 1.39 (m, 9H), 1.30 – 1.24 (m, 3H).

Example 75. Synthesis of (4S,5S)-ethyl 4-((*tert*-butoxycarbonyl)amino)-5-methyl-3-oxo heptanoate (**217**)




To an ice-cooled solution of N-Boc-L-isoleucine (4.55 g, 19.67 mmol) in THF (20 mL) was added 1,1'-carbonyldiimidazole (3.51 g, 21.63 mmol). After evolution of gas ceased, the resultant mixture was stirred at r.t. for 3.5 h.

A solution of freshly prepared isopropylmagnesium bromide in THF (123 mmol, 30 mL) was added dropwise to a pre-cooled (0 °C) solution of ethyl hydrogen malonate (6.50 g, 49.2 mmol) at such a rate to keep the internal temperature below 5 °C. The mixture was stirred at r.t. for 1.5 h. This solution of the magnesium enolate was then cooled over an ice-water bath, followed by the gradual addition of the imidazolide solution over a 1 h period via a double-ended needle at 0 °C. The resultant mixture was stirred at 0 °C for 30 min then r.t. 64 h. The reaction mixture was quenched by addition of 10% aqueous citric acid (5 mL), and acidified to pH 3 with an additional 10% aqueous citric acid (110 mL). The mixture was extracted with ethyl acetate (3 × 150 mL). The organic extracts were washed with water (50 mL), saturated aqueous sodium hydrogen carbonate (50 mL), and saturated aqueous sodium chloride (50 mL), dried over


Na₂SO₄, and concentrated in vacuo. The residue was purified by column chromatography on silica gel using ethyl acetate/hexane (1:4) as an eluent to give compound **217** (5.50 g, 93% yield). ¹H NMR (500 MHz, CDCl₃) δ 5.04 (d, *J* = 7.8 Hz, 1H), 4.20 (p, *J* = 7.0 Hz, 3H), 3.52 (t, *J* = 10.7 Hz, 2H), 1.96 (d, *J* = 3.7 Hz, 1H), 1.69 (s, 2H), 1.44 (s, 9H), 1.28 (dd, *J* = 7.1, 2.9 Hz, 3H), 0.98 (t, *J* = 6.9 Hz, 3H), 0.92 – 0.86 (m, 3H).

Example 76. Synthesis of (3R,4S,5S)-ethyl 4-((*tert*-butoxycarbonyl)amino)-3-hydroxy-5-methylheptanoate (**218**)



To a solution of the compound **217** (5.90 g, 19.83 mmol) in ethanol (6 mL) at -60 °C was added sodium borohydride (3.77 g, 99.2 mmol) in one portion. The reaction mixture was stirred for 5.5 h below -55 °C then quenched with 10% aqueous citric acid (100 mL). The resultant solution was acidified to pH 2 with an additional 10% aqueous citric acid, followed by extraction with ethyl acetate (3 × 100 mL). The organic extracts were washed with saturated aqueous sodium chloride (100 mL), dried over Na₂SO₄, and concentrated in vacuo. The residue was purified by column chromatography (10-50% ethyl acetate/hexane) to give pure diastereomer **218** (2.20 g, 37% yield) and a mixture of two diastereomers (2.0g, 34% yield, about 9:1 ratio). ¹H NMR (500 MHz, CDCl₃) δ 4.41 (d, *J* = 9.3 Hz, 1H), 4.17 (tt, *J* = 7.1, 3.6 Hz, 2H), 4.00 (t, *J* = 6.9 Hz, 1H), 3.55 (dd, *J* = 11.7, 9.3 Hz, 1H), 2.56 – 2.51 (m, 2H), 2.44 (dd, *J* = 16.4, 9.0 Hz, 1H), 1.79 (d, *J* = 3.8 Hz, 1H), 1.60 – 1.53 (m, 1H), 1.43 (s, 9H), 1.27 (dd, *J* = 9.3, 5.0 Hz, 3H), 1.03 – 0.91 (m, 7H).


Example 77. Synthesis of (3R,4S,5S)-4-((*tert*-butoxycarbonyl)amino)-3-hydroxy-5-methylheptanoic acid (**219**)



To a solution of compound **218** (2.20 g, 7.20 mmol) in ethanol (22 mL) was added 1 N aqueous sodium hydroxide (7.57 mL, 7.57 mmol). The mixture was stirred at 0 °C for 30 min then r.t. 2 h. The resultant solution was acidified to pH 4 by addition of 1 N aqueous hydrochloric acid, which was then extracted with ethyl acetate (3 × 50 mL). The organic extracts were washed with 1 N aqueous potassium hydrogen sulfate (50 mL), and saturated aqueous sodium chloride (50 mL), dried over Na₂SO₄, and concentrated in vacuo to give compound **219** (1.90 g, 95% yield). ¹H NMR (500 MHz, CDCl₃) δ 4.50 (d, *J* = 8.7 Hz, 1H), 4.07 (d, *J* = 5.5 Hz, 1H), 3.59 (d, *J*


= 8.3 Hz, 1H), 2.56 – 2.45 (m, 2H), 1.76 – 1.65 (m, 1H), 1.56 (d, $J = 7.1$ Hz, 1H), 1.45 (s, 9H), 1.26 (t, $J = 7.1$ Hz, 3H), 0.93 (dd, $J = 14.4, 7.1$ Hz, 6H).

Example 78. Synthesis of (3R,4S,5S)-4-((*tert*-butoxycarbonyl)(methyl)amino)- 3-methoxy-5-methylheptanoic acid (**220**)




To a solution of compound **219** (1.90 g, 6.9 mmol) in THF (40 mL) was added sodium hydride (60% oil suspension, 1.93 g, 48.3 mmol) at 0 °C. After stirring for 1h, methyl iodide (6.6 mL, 103.5 mmol) was added. The stirring was continued at 0 °C for 40 h before saturated aqueous sodium hydrogen carbonate (50 mL) was added, followed by water (100 mL). The mixture was washed with diethyl ether (2 × 50 mL) and the aqueous layer was acidified to pH 3 by 1 N aqueous potassium hydrogen sulfate, then extracted with ethyl acetate (3 × 50 mL). The combined organic extracts were washed with 5% aqueous sodium thiosulfate (50 mL) and saturated aqueous sodium chloride (50 mL), dried over Na₂SO₄, and concentrated in vacuo to give compound **220** (1.00 g, 48% yield). ¹H NMR (500 MHz, CDCl₃) δ 3.95 (d, $J = 75.4$ Hz, 2H), 3.42 (d, $J = 4.4$ Hz, 3H), 2.71 (s, 3H), 2.62 (s, 1H), 2.56 – 2.47 (m, 2H), 1.79 (s, 1H), 1.47 (s, 1H), 1.45 (d, $J = 3.3$ Hz, 9H), 1.13 – 1.05 (m, 1H), 0.96 (d, $J = 6.7$ Hz, 3H), 0.89 (td, $J = 7.2, 2.5$ Hz, 3H).

Example 79. Synthesis of Boc-N-Me-L-Val-OH (**222**)



To a solution of Boc-L-Val-OH (2.00 g, 9.2 mmol) and methyl iodide (5.74 mL, 92 mmol) in anhydrous THF (40 mL) was added sodium hydride (3.68 g, 92 mmol) at 0 °C. The reaction mixture was stirred at 0 °C for 1.5 h, then warmed to r.t. and stirred for 24 h. The reaction was quenched by ice water (50 mL). After addition of water (100 mL), the reaction mixture was washed with ethyl acetate (3 × 50 mL) and the aqueous solution was acidified to pH 3 then extracted with ethyl acetate (3 × 50 mL). The combined organic phase was dried over Na₂SO₄ and concentrated to afford Boc-N-Me-Val-OH (2.00 g, 94% yield) as a white solid. ¹H NMR (500 MHz, CDCl₃) δ 4.10 (d, $J = 10.0$ Hz, 1H), 2.87 (s, 3H), 2.37 – 2.13 (m, 1H), 1.44 (d, $J = 26.7$ Hz, 9H), 1.02 (d, $J = 6.5$ Hz, 3H), 0.90 (t, $J = 8.6$ Hz, 3H).


Example 80. Synthesis of (S)-*tert*-butyl 2-((1R,2R)-1-methoxy-3-(((S)-1-methoxy-1-oxo-3-phenylpropan-2-yl)amino)-2-methyl-3-oxopropyl)pyrrolidine-1-carboxylate (**223**)



To a solution of compound **214** (100 mg, 0.347 mmol) and L-phenylalanine methyl ester hydrochloride (107.8 mg, 0.500 mmol) in DMF (5 mL) at 0 °C was added diethyl cyanophosphonate (75.6 μ L, 0.451 mmol), followed by Et₃N (131 μ L, 0.94 mmol). The reaction mixture was stirred at 0 °C for 2 h, then warmed to r.t. and stirred overnight. The reaction mixture was then diluted with ethyl acetate (80 mL), washed with 1 N aqueous potassium hydrogen sulfate (40 mL), water (40 mL), saturated aqueous sodium hydrogen carbonate (40 mL), and saturated aqueous sodium chloride (40 mL), dried over Na₂SO₄, and concentrated in vacuo. The residue was purified by column chromatography (15-75% ethyl acetate/hexanes) to afford compound **223** (130 mg, 83% yield) as a white solid. ¹H NMR (500 MHz, CDCl₃) δ 7.28 (dd, *J* = 7.9, 6.5 Hz, 2H), 7.23 (t, *J* = 7.3 Hz, 1H), 7.16 (s, 2H), 4.81 (s, 1H), 3.98 – 3.56 (m, 5H), 3.50 (s, 1H), 3.37 (d, *J* = 2.9 Hz, 3H), 3.17 (dd, *J* = 13.9, 5.4 Hz, 2H), 3.04 (dd, *J* = 14.0, 7.7 Hz, 1H), 2.34 (s, 1H), 1.81 – 1.69 (m, 2H), 1.65 (s, 3H), 1.51 – 1.40 (m, 9H), 1.16 (d, *J* = 7.0 Hz, 3H).

Example 81. General procedure for the removal of the Boc function with trifluoroacetic acid. To a solution of the *N*-Boc amino acid (1.0 mmol) in methylene chloride (2.5 mL) was added trifluoroacetic acid (1.0 mL). After being stirred at room temperature for 1-3 h, the reaction mixture was concentrated in vacuo. Co-evaporation with toluene gave the deprotected product, which was used without any further purification.


Example 82. Synthesis of (S)-methyl 2-((2R,3R)-3-((S)-1-((3R,4S,5S)-4-((tert-butoxycarbonyl)(methyl)amino)-3-methoxy-5-methylheptanoyl)pyrrolidin-2-yl)-3-methoxy-2-methylpropanamido)-3-phenylpropanoate (225**).**



To a solution of the Boc-deprotected product of compound **224** (0.29 mmol) and compound **220** (96.6 mg, 0.318 mmol) in DMF (5 mL) at 0 °C was added diethyl cyanophosphonate (58 μ L, 0.347 mmol), followed by Et₃N (109 μ L, 0.78 mmol). The reaction mixture was stirred at 0 °C for 2 h, then warmed to r.t. and stirred overnight. The reaction mixture was diluted with ethyl acetate (80 mL), washed with 1 N aqueous potassium hydrogen sulfate (40 mL), water (40 mL), saturated aqueous sodium hydrogen carbonate (40 mL), and saturated aqueous sodium chloride (40 mL), dried over Na₂SO₄ and concentrated in vacuo. The residue was purified by column


chromatography (15-75% ethyl acetate/hexanes) to afford compound **225** (150 mg, 81% yield) as a white solid. LC-MS (ESI) m/z calcd. for $C_{34}H_{55}N_3O_8$ $[M+H]^+$: 634.40, found: 634.40.

Example 83. Synthesis of (S)-methyl 2-((2R,3R)-3-((S)-1-((3R,4S,5S)-4-((S)-2-((tert-butoxycarbonyl)amino)-N,3-dimethylbutanamido)-3-methoxy-5-methylheptanoyl)pyrrolidin-2-yl)-3-methoxy-2-methylpropanamido)-3-phenylpropanoate (**227**).




To a solution of the Boc-protected product of compound **225** (0.118 mmol) and Boc-Val-OH (51.8 mg, 0.236 mmol) in DCM (5 mL) at 0 °C was added BroP (70.1 mg, 0.184 mmol), followed by diisopropylethylamine (70 μ L, 0.425 mmol). The mixture was shielded from light and stirred at 0 °C for 30 min then at r.t. for 2 days. The reaction mixture was diluted with ethyl acetate (80 mL), washed with 1 N aqueous potassium hydrogen sulfate (40 mL), water (40 mL), saturated aqueous sodium hydrogen carbonate (40 mL), and saturated aqueous sodium chloride (40 mL), dried over Na_2SO_4 and concentrated in vacuo. The residue was purified by column chromatography (20-100% ethyl acetate/hexanes) to afford compound **227** (67 mg, 77% yield) as a white solid. LC-MS (ESI) m/z calcd. for $C_{39}H_{64}N_4O_9$ $[M+H]^+$: 733.47, found: 733.46.

Example 84. Synthesis of (S)-methyl 2-((2R,3R)-3-((S)-1-((6S,9S,12S,13R)-12-((S)-sec-butyl)-6,9-diisopropyl-13-methoxy-2,2,5,11-tetramethyl-4,7,10-trioxo-3-oxa-5,8,11-triazapentadecan-15-oyl)pyrrolidin-2-yl)-3-methoxy-2-methylpropanamido)-3-phenylpropanoate (**229**).




To a solution of the Boc-protected product of compound **227** (0.091 mmol) and Boc-N-Me-Val-OH (127 mg, 0.548 mmol) in DMF (5 mL) at 0 °C was added diethyl cyanophosphonate (18.2 μ L, 0.114 mmol), followed by N-methylmorpholine (59 μ L, 0.548 mmol). The reaction mixture was stirred at 0 °C for 2 h, then warmed to r.t. and stirred overnight. The reaction mixture was diluted with ethyl acetate (80 mL), washed with 1 N aqueous potassium hydrogen sulfate (40 mL), water (40 mL), saturated aqueous sodium hydrogen carbonate (40 mL), and saturated aqueous sodium chloride (40 mL), dried over sodium sulfate, and concentrated in vacuo. The residue was purified by column chromatography (20-100% ethyl acetate/hexanes) to afford compound **229** (30 mg, 39% yield) as a white solid. LC-MS (ESI) m/z calcd. for $C_{45}H_{75}N_5O_{10}$ $[M+H]^+$: 846.55, found: 846.56.

Example 85. Synthesis of (S)-methyl 2-((2R,3R)-3-((S)-1-((3R,4S,5S)-4-((S)-N,3-dimethyl-2-((S)-3-methyl-2-(methylamino)butanamido)butanamido)-3-methoxy-5-methylheptanoyl)pyrrolidin-2-yl)-3-methoxy-2-methylpropanamido)-3-phenylpropanoate (**230**).




To a solution of compound **229** (75.0 mg, 0.0886 mmol) in methylene chloride (5 mL) was added trifluoroacetic acid (2 mL) at room temperature. After being stirred at room temperature for 1 h, the reaction mixture was concentrated in vacuo. Co-evaporation with toluene gave the deprotected product **230**, which was used without further purification.

Example 86. Synthesis of (2S,4R)-methyl 4-hydroxypyrrolidine-2-carboxylate hydrochloric (**232**).




To a solution of trans-4-hydroxy-L-proline (15.0 g, 114.3 mmol) in dry methanol (250 mL) was added thionyl chloride (17 mL, 231 mmol) dropwise at 0 to 4 °C. The resulting mixture was stirred for at r.t. overnight, concentrated, crystallized with EtOH/hexane to provide the title compound (18.0 g, 87% yield). ESI MS m/z 168.2 ($[M+Na]^+$).

Example 87. Synthesis of (2S,4R)-1-*tert*-butyl 2-methyl 4-hydroxypyrrolidine- 1,2-dicarboxyl ate (**233**).




To a solution of trans-4-hydroxy-L-proline methyl ester (18.0 g, 107.0 mmol) in the mixture of MeOH (150 ml) and sodium bicarbonate solution (2.0 M, 350 ml) was added Boc₂O (30.0 g, 137.6 mmol) in three portions in 4 h. After stirring for an additional 4 h, the reaction was concentrated to ~350 ml and extracted with EtOAc (4 × 80 mL). The combined organic layers were washed with brine (100 mL), dried (MgSO₄), filtered, concentrated and purified by SiO₂ column chromatography (1:1 hexanes/EtOAc) to give the title compound (22.54 g, 86% yield). ESI MS m/z 268.2 ($[M+Na]^+$).

Example 88. Synthesis of (S)-1-*tert*-butyl 2-methyl 4-oxopyrrolidine-1,2- dicarboxylate (**234**).




The title compound prepared through Dess-Martin oxidation was described in: Franco Manfre et al. J. Org. Chem. 1992, 57, 2060-2065. Alternatively Swern oxidation procedure is as following: To a solution of (COCl)₂ (13.0 ml, 74.38 mmol) in CH₂Cl₂ (350 ml) cooled to -78 °C was added dry DMSO (26.0 mL). The solution was stirred at -78 °C for 15 min and then compound **233** (8.0 g, 32.63 mmol) in CH₂Cl₂ (100 ml) was added. After stirring at -78 °C for 2 h, triethylamine (50 ml, 180.3 mmol) was added dropwise, and the reaction solution was warmed to room temperature. The mixture was diluted with aq. NaH₂PO₄ solution (1.0 M, 400 ml) and phases separated. The aqueous layer was extracted with CH₂Cl₂ (2 × 60 ml). The organic layers were combined, dried over MgSO₄, filtered, concentrated and purified by SiO₂ column chromatography (7:3 hexanes/EtOAc) to give the title compound (6.73 g, 85% yield). ESI MS m/z 266.2([M+Na]⁺).

Example 89. Synthesis of (S)-1-*tert*-butyl 2-methyl 4-methylenepyrrolidine-1,2-dicarboxylate (**235**).




To a suspension of methyltriphenylphosphonium bromide (19.62 g, 55.11 mmol) in THF (150 mL) at 0 °C was added potassium-*t*-butoxide (6.20 g, 55.30 mmol) in anhydrous THF (80 mL). After stirring at 0 °C for 2 h, the resulting yellow ylide was added to a solution of compound **234** (6.70 g, 27.55 mmol) in THF (40 mL). After stirring at r.t. for 1 h, the reaction mixture was concentrated, diluted with EtOAc (200 mL), washed with H₂O (150 mL), brine (150 mL), dried over MgSO₄, concentrated and purified on SiO₂ column chromatography (9:1 hexanes/EtOAc) to yield the title compound (5.77 g, 87% yield). EI MS m/z 264 ([M+Na]⁺).

Example 90. Synthesis of (S)-methyl 4-methylenepyrrolidine-2-carboxylate hydrochloride (**236**)




To a solution of compound **235** (5.70 g, 23.63 mmol) in EtOAc (40 ml) at 4 °C was added HCl (12 M, 10 ml). The mixture was stirred for 1 h, diluted with toluene (50 ml), concentrated, and crystallized with EtOH/hexane to yield the title compound as HCl salt (3.85 g, 92% yield). EI MS m/z 142.2 ([M+H]⁺).

Example 91. Synthesis of 4-(benzyloxy)-3-methoxybenzoic acid (**238**).




To a mixture of 4-hydroxy-3-methoxybenzoic acid (50.0 g, 297.5 mmol) in ethanol (350 ml) and aq. NaOH solution (2.0 M, 350 ml) was added BnBr (140.0 g, 823.5 mmol). The mixture was stirred at 65 °C for 8 h, concentrated, co-evaporated with water (2 × 400 ml) and concentrated to ~400 ml, acidified to pH 3.0 with 6 N HCl. The solid was collected by filtration, crystallized with EtOH, dried at 45 °C under vacuum to afford the title compound (63.6 g, 83% yield). ESI MS m/z 281.2 ([M+Na]⁺).

Example 92. Synthesis of 4-(benzyloxy)-5-methoxy-2-nitrobenzoic acid (**239**).




To a solution of compound **238** (63.5 g, 246.0 mmol) in CH₂Cl₂ (400 ml) and HOAc (100 ml) was added HNO₃ (fuming, 25.0 ml, 528.5 mmol). The mixture was stirred for 6 h, concentrated, crystallized with EtOH, dried at 40 °C under vacuum to afford the title compound (63.3 g, 85% yield). ESI MS m/z 326.1 ([M+Na]⁺).

Example 93. Synthesis of (S)-methyl 1-(4-(benzyloxy)-5-methoxy-2-nitro benzoyl)-4-methylenepyrrolidine-2-carboxylate (**240**).




A catalytic amount of DMF (30 µl) was added to a solution of compound **239** (2.70 g, 8.91 mmol) and oxalyl chloride (2.0 mL, 22.50 mmol) in anhydrous CH₂Cl₂ (70 mL) and the resulting mixture was stirred at room temperature for 2 h. Excess CH₂Cl₂ and oxalyl chloride was removed with rotavap. The acetyl chloride was re-suspended in fresh CH₂Cl₂ (70 mL) and was added slowly to a pre-mixed solution of **236** (1.58 g, 8.91 mmol) and Et₃N (6 mL) in CH₂Cl₂ at 0 °C under N₂ atmosphere. The reaction mixture was allowed to warm to r.t. and stirring was continued for 8 h. After removal of CH₂Cl₂ and Et₃N, the residue was partitioned between H₂O and EtOAc (70/70 mL). The aqueous layer was further extracted with EtOAc (2 × 60 mL). The combined organic layers were washed with brine (40 mL), dried (MgSO₄) and concentrated. Purification of the residue with flash chromatography (silica gel, 2:8 hexanes/EtOAc) yielded compound **240** (2.88 g, 76% yield). EI MS m/z 449.1 ([M+Na]⁺).

Example 94. Synthesis of (S)-1-(4-(benzyloxy)-5-methoxy-2-nitrobenzoyl)-4-methylenepyrrolidine-2-carbaldehyde (**241**).




To a vigorously stirred solution of compound **240** (2.80 g, 6.57 mmol) in anhydrous CH₂Cl₂ (60 mL) was added DIBAL-H (1N in CH₂Cl₂, 10 mL) dropwise at -78 °C under N₂ atmosphere. After the mixture was stirred for an additional 90 min, excess reagent was decomposed by addition of 2 ml of methanol, followed by 5% HCl (10 mL). The resulting mixture was allowed to warm to 0 °C. Layers were separated and the aqueous layer was further extracted with CH₂Cl₂ (3 × 50 mL). Combined organic layers were washed with brine, dried (MgSO₄) and concentrated. Purification of the residue with flash chromatography (silica gel, 95:5 CHCl₃/MeOH) yielded compound **241** (2.19 g, 84% yield). EIMS m/z 419.1 ([M+Na]⁺).

Example 95. Synthesis of (S)-8-(benzyloxy)-7-methoxy-2-methylene-2,3 -dihydro-1H-benzo[e]-pyrrolo[1,2-a]azepin-5(11aH)-one (**242**).




A mixture of compound **241** (2.18 g, 5.50 mmol) and Na₂S₂O₄ (8.0 g, 45.97 mmol) in THF (60 ml) and H₂O (40 ml) was stirred at r.t. for 20 h. Solvents were removed under high vacuum. The residue was re-suspended in MeOH (60 mL), and HCl (6M) was added dropwise until pH ~ 2 was reached. The resulting mixture was stirred at r.t. for 1 h. The reaction was worked-up by removing most of MeOH, then diluted with EtOAc (100 mL). The EtOAc solution was washed with sat. NaHCO₃, brine, dried (MgSO₄), and concentrated. Purification of the residue with flash chromatography (silica gel, 97:3 CHCl₃/MeOH) yielded compound **242** (1.52 g, 80%). EIMS m/z 372.1 ([M+Na]⁺).

Example 96. Synthesis of (S)-8-hydroxy-7-methoxy-2-methylene-2,3 -dihydro-1H-benzo[e]-pyrrolo[1,2-a]azepin-5(11aH)-one (**243**)




To a solution of compound **242** (1.50 g, 4.32 mmol) in 70 ml of CH₂Cl₂ was added 25 ml of CH₃SO₃H at 0 °C. The mixture was stirred at 0 °C for 10 min then r.t. for 2 h, diluted with CH₂Cl₂, pH adjusted with cold 1.0 N NaHCO₃ to 4 and filtered. The aqueous layer was extracted with CH₂Cl₂ (3 × 60 ml). The organic layers were combined, dried over Na₂SO₄, filtered, evaporated and purified on SiO₂ column chromatography (CH₃OH/CH₂Cl₂ 1:15) to afford 811 mg (73% yield) of the title product. EIMS m/z 281.1 ([M+Na]⁺).

Example 97. Synthesis of (11aS,11a'S)-8,8'-(pentane-1,5-diylbis(oxy))bis(7-methoxy-2-methylene-2,3-dihydro-1H-benzo[e]pyrrolo[1,2-a][1,4]diazepin-5(11aH)-one) (**244**).




To a stirred suspended solution of Cs_2CO_3 (0.761 g, 2.33 mmol) in butanone (8 ml) were added compound **C11** (401 mg, 1.55 mmol) and 1,5-diiodopentane (240 mg, 0.740 mmol). The mixture was stirred at r.t. overnight, concentrated, and purified on SiO_2 chromatography (EtOAc/ CH_2Cl_2 1:10) to afford 337 mg (78% yield) of the title product. EIMS m/z 607.2 ($[\text{M}+\text{Na}]^+$).

Example 98. Synthesis of (S)-7-methoxy-8-((5-(((S)-7-methoxy-2-methylene-5-oxo-2,3,5,10,11,11a-hexahydro-1H-benzo[e]pyrrolo[1,2-a][1,4]diazepin-8-yl)oxy)pentyl)oxy)-2-methylene-2,3-dihydro-1H-benzo[e]pyrrolo[1,2-a][1,4]diazepin-5(11aH)-one (**245**).




To a solution of compound **244** (150 mg, 0.256 mmol) in anhydrous dichloromethane (1 mL) and absolute ethanol (1.5 mL) was added sodium borohydride in methoxyethyl ether (85 μL , 0.5 M, 0.042 mmol) at 0 °C. The ice bath was removed after 5 minutes and the mixture was stirred at room temperature for 3 hours, then cooled to 0 °C, quenched with saturated ammonium chloride, diluted with dichloromethane, and phases separated. The organic layer was washed with brine, dried over anhydrous Na_2SO_4 , filtered through Celite and concentrated. The residue was purified by reverse phase HPLC (C_{18} column, acetonitrile/water). The corresponding fractions were extracted with dichloromethane and concentrated to afford the title compound **245** (64.7 mg, 43%), MS m/z 609.2 ($[\text{M}+\text{Na}]^+$), 625.3 ($[\text{M}+\text{K}]^+$) and 627.2 ($[\text{M}+\text{Na}+\text{H}_2\text{O}]^+$); the fully reduced compound was obtained (16.5 mg, 11%), MS m/z 611.2 ($[\text{M}+\text{Na}]^+$), 627.2 ($[\text{M}+\text{K}]^+$), 629.2 ($[\text{M}+\text{Na}+\text{H}_2\text{O}]^+$); and the unreacted starting material was also recovered (10.2 mg, 7%), MS m/z 607.2 ($[\text{M}+\text{Na}]^+$), 625.2 ($[\text{M}+\text{Na}+\text{H}_2\text{O}]^+$).

Example 99. Synthesis of (S)-8-((5-(((S)-10-(3-(2-(2-azidoethoxy)ethoxy) propanoyl)-7-methoxy-2-methylene-5-oxo-2,3,5,10,11,11a-hexahydro-1H-benzo[e]pyrrolo[1,2-a][1,4]diazepin-8-yl)oxy)pentyl)oxy)-7-methoxy-2-methylene-2,3-dihydro-1H-benzo[e]pyrrolo[1,2-a][1,4]diazepin-5(11aH)-one (**246**).




To the mixture of compound **245** (60.0 mg, 0.102 mmol) and compound **166** (40.5 mg, 0.134 mmol) in dichloromethane (5 ml) was added EDC (100.5 mg, 0.520 mmol). The mixture was stirred at r.t. overnight, concentrated and purified on SiO₂ column chromatography (EtOAc/CH₂Cl₂, 1:6) to afford 63.1 mg (81% yield) of the title product **246**. ESI MS m/z C₄₀H₅₀N₇O₉ [M+H]⁺, calcd. 772.36, found 772.30.

Example 100. Synthesis of (S)-8-(((S)-10-(3-(2-(2-aminoethoxy)ethoxy) propanoyl)-7-methoxy-2-methylene-5-oxo-2,3,5,10,11,11a-hexahydro-1H-benzo[e]pyrrolo[1,2-a][1,4]diazepin-8-yl)oxy)pentyl)oxy)-7-methoxy-2-methylene-2,3-dihydro-1H-benzo[e]pyrrolo[1,2-a][1,4]diazepin-5(11aH)-one (**247**).




To a solution of compound **246** (60 mg, 0.078 mmol) in the mixture of THF (5 ml) and NaH₂PO₄ buffer solution (pH 7.5, 1.0 M, 0.7 ml) was added PPh₃ (70 mg, 0.267 mmol). The mixture was stirred at r.t. overnight, concentrated and purified on C₁₈ preparative HPLC, eluted with water/CH₃CN (from 90% water to 35% water in 35 min) to afford 45.1 mg (79% yield) of the title product **247** after drying under high vacuum. ESI MS m/z C₄₀H₅₂N₅O₉ [M+H]⁺, calcd. 746.37, found 746.50.

Example 101. Synthesis of (2S)-2-((2R,3R)-3-((2S)-1-((11S,14S,17S)-1-azido-17-((R)-sec-butyl)-11,14-diisopropyl-18-methoxy-10,16-dimethyl-9,12,15-trioxo-3,6-dioxa-10,13,16-triazai-cosan-20-oyl)pyrrolidin-2-yl)-3-methoxy-2-methylpropanamido)-3-phenylpropanoic acid (**257**)




To the crude compound **331** (22 mg, 0.030 mmol) in a mixture of DMA (0.8 ml) and NaH₂PO₄ buffer solution (pH 7.5, 1.0 M, 0.7 ml) was added compound **166** (18.0 mg, 0.060 mmol) in four portions in 2 h. The mixture was stirred overnight, concentrated and purified on SiO₂ column chromatography (CH₃OH/CH₂Cl₂/HOAc 1:8:0.01) to afford the title compound (22.5 mg, 82% yield). LC-MS (ESI) m/z calcd. for C₄₆H₇₇N₈O₁₁ [M+H]⁺: 917.56, found: 917.60.

Example 102. Synthesis of (2S)-2-((2R,3R)-3-((2S)-1-((11S,14S,17S)-1-amino-17-((R)-sec-butyl)-11,14-diisopropyl-18-methoxy-10,16-dimethyl-9,12,15-trioxo-3,6-dioxo-10,13,16-triazaicosan-20-yl)pyrrolidin-2-yl)-3-methoxy-2-methylpropanamido)-3-phenylpropanoic acid (**258**).




To compound **257** (22.0 mg, 0.024 mmol) in methanol (5 ml) in a hydrogenation bottle was added Pd/C (5 mg, 10% Pd, 50% wet). After air was vacuumed out and 25 psi H₂ was conducted in, the mixture was shaken for 4 h, filtered through celite. The filtrate was concentrated to afford the crude title product (~20 mg, 92% yield), which was used in the next step without further purification. ESI MS m/z+ C₄₆H₇₉N₆O₁₁ (M+H), calcd.891.57, found 891.60.

Example 103. Synthesis of (S)-2-((2R,3R)-3-((S)-1-((6S,9S,12S,13R)-12-((S)-sec-butyl)-6,9-diisopropyl-13-methoxy-2,2,5,11-tetramethyl-4,7,10-trioxo-3-oxa-5,8,11-triazapenta-decan-15-yl)pyrrolidin-2-yl)-3-methoxy-2-methylpropanamido)-3-phenylpropanoic acid (**330**).




To a solution of compound **229** (30 mg, 0.035 mmol) in THF (1.0 ml) was added LiOH in water (1.0M, 0.8 ml). The mixture was stirred at r.t. for 35 min, neutralized with 0.5 M H₃PO₄ to pH 6, concentrated and purified on SiO₂ column chromatography (CH₃OH/CH₂Cl₂/HOAc 1:10:0.01) to afford the title compound **330** (25.0 mg, 85% yield). LC-MS (ESI) m/z calcd.for C₄₄H₇₄N₅O₁₀ [M+H]⁺: 832.54, found: 832.60.

Example 104. Synthesis of (S)-2-((2R,3R)-3-((S)-1-((3R,4S,5S)-4-((S)-N,3-dimethyl-2-((S)-3-methyl-2-(methylamino)butanamido)butanamido)-3-methoxy-5-methylheptanoyl)-pyrrolidin-2-yl)-3-methoxy-2-methylpropanamido)-3-phenylpropanoic acid (**331**)




Compound **330** (25 mg, 0.030 mmol) in the mixture of conc. HCl (0.3 ml) and 1,4-dioxane (0.9 ml) was stirred at r.t. for 35 min. The mixture was diluted with EtOH (1.0 ml) and toluene (1.0 ml), concentrated and co-evaporated with EtOH/toluene (2:1) to afford the title compound **331** as a white solid (22 mg, ~100% yield), which was used in the next step without further purification. LC-MS (ESI) m/z calcd.for C₃₉H₆₆N₅O₈ [M+H]⁺: 732.48, found: 732.60.

Example 105. Synthesis of (4R)-4-(2-((1R,3R)-1-acetoxy-3-((2S,3S)-N,3- dimethyl-2-((R)-1-methylpiperidine-2-carboxamido)pentanamido)-4-methylpentyl)thiazole-4-carboxamido)-5-(3-(3-(2-(2-azidoethoxy)ethoxy)propanamido)-4-hydroxyphenyl)-2-methylpentanoic acid (**335**)




To a solution of (4R)-4-(2-((1R,3R)-1-acetoxy-3-((2S,3S)-N,3-dimethyl-2-((R)-1-methylpiperidine-2-carboxamido)pentanamido)-4-methylpentyl)thiazole-4-carboxamido)-5-(3-amino-4-hydroxyphenyl)-2-methylpentanoic acid **285** (Huang Y. et al, Med Chem. #44, 249th ACS National Meeting, Denver, CO, Mar. 22~26, 2015; WO2014009774) (100 mg, 0.131 mmol) in the mixture of DMA (10 ml) and NaH₂PO₄ buffer solution (pH 7.5, 1.0 M, 0.7 ml) was added compound **166** (80.0 mg, 0.266 mmol) in four portions in 2 h. The mixture was stirred overnight, concentrated and purified on C₁₈ preparative HPLC (3.0 × 25 cm, 25 ml/min), eluted with from 80% water/methanol to 10% water/methanol in 45 min to afford the title compound (101.5 mg, 82% yield). LC-MS (ESI) m/z calcd.for C₄₅H₇₀N₉O₁₁S [M+H]⁺: 944.48, found: 944.70.

Example 106. Synthesis of (4R)-4-(2-((1R,3R)-1-acetoxy-3-((2S,3S)-N,3- dimethyl-2-((R)-1-methyl-piperidine-2-carboxamido)pentanamido)-4-methylpentyl)thiazole-4-carboxamido)-5-(3-(3-(2-(2-aminoethoxy)ethoxy)propanamido)-4-hydroxyphenyl)-2-methylpentanoic acid (**336**)




To a solution of compound **332** (100.0 mg, 0.106 mmol) in methanol (25 ml) containing 0.1% HCl in a hydrogenation bottle was added Pd/C (25 mg, 10% Pd, 50% wet). After air was vacuumed out in the vessel and 35 psi H₂ was conducted in, the mixture was shaken for 4 h, filtered through celite. The filtrate was concentrated and purified on C₁₈ preparative HPLC (3.0 × 25 cm, 25 ml/min), eluted with from 85% water/methanol to 15% water/methanol in 45 min to afford the title compound (77.5 mg, 79% yield). LC-MS (ESI) m/z calcd.for C₄₅H₇₂N₇O₁₁S [M+H]⁺: 918.49, found: 918.60.

Example 107. Synthesis of 2,3-dibromo-4-((2-(2-(3-((S)-7-methoxy-8-((5-(((S)-7-methoxy-2-methylene-5-oxo-2,3,5,11a-tetrahydro-1H-benzo[e]pyrrolo[1,2-a][1,4]diazepin-8-yl)oxy)pentyl)oxy)-2-methylene-5-oxo-2,3,11,11a-tetrahydro-1H-benzo[e]pyrrolo[1,2-a][1,4]diazepin-10(5H)-yl)-3-oxopropoxy)ethoxy)ethyl)amino)-4-oxobutanoic acid (**340**).



To a solution of compound **247** (40.0 mg, 0.068 mmol) in the mixture of DCM (4 ml) and DIPEA (12 μ L, 0.069 mmol) was added **7** (38.0 mg, 0.148 mmol) at 0 °C. The mixture was stirred at 0 °C for 2 h and then r.t. 5h. The mixture was concentrated and purified on SiO₂ column eluted with CH₃OH/CH₂Cl₂/HOAc (1:6:0.01) to afford the title compound **340** (56.5 mg, 83% yield). ESI MS *m/z* calcd. for C₄₄H₅₃Br₂N₅O₁₂ [M+H]⁺: 1002.21, found: 1002.40, 1004.40 [M+2+H]⁺.


Example 108. Synthesis of 2,5-dioxopyrrolidin-1-yl 2,3-dibromo-4-((2-(2-(3-((S)-7-methoxy-8-((5-(((S)-7-methoxy-2-methylene-5-oxo-2,3,5,11a-tetrahydro-1H-benzo[e]pyrrolo[1,2-a][1,4]diazepin-8-yl)oxy)pentyl)oxy)-2-methylene-5-oxo-2,3,11,11a-tetrahydro-1H-benzo[e]pyrrolo[1,2-a][1,4]diazepin-10(5H)-yl)-3-oxopropoxy)ethoxy)ethyl)amino)-4-oxobutanoate (**341**)



To a solution of 2,3-dibromo-4-((2-(2-(3-((S)-7-methoxy-8-((5-(((S)-7-methoxy-2-methylene-5-oxo-2,3,5,11a-tetrahydro-1H-benzo[e]pyrrolo[1,2-a][1,4]diazepin-8-yl)oxy)pentyl)oxy)-2-methylene-5-oxo-2,3,11,11a-tetrahydro-1H-benzo[e]pyrrolo[1,2-a][1,4]diazepin-10(5H)-yl)-3-oxopropoxy)ethoxy)ethyl)amino)-4-oxobutanoic acid (55.0 mg, 0.054 mmol) in CH₂Cl₂ (3 ml) was added NHS (10.0 mg, 0.086 mmol) and EDC (30.5 mg, 0.158 mmol). The mixture was stirred at r.t. overnight, concentrated and purified on SiO₂ column eluted with EtOAc/CH₂Cl₂ (1:5) to afford the title compound **341** (50.5 mg, 85% yield). ESI MS *m/z* calcd. for C₄₈H₅₆Br₂N₆O₁₄ [M+H]⁺: 1099.22, found: 1099.40, 1101.40 [M+2+H]⁺, 1119.50 [M+2+H+H₂O]⁺.


Example 109. Synthesis of (2S)-2-((2R,3R)-3-((2S)-1-((13S,26S,29S,32S)-12,13-dibromo-32-((R)-sec-butyl)-26,29-diisopropyl-33-methoxy-1-((S)-7-methoxy-8-((5-(((S)-7-methoxy-2-methylene-5-oxo-2,3,5,11a-tetrahydro-1H-benzo[e]pyrrolo[1,2-a][1,4]diazepin-8-yl)oxy)pentyl)oxy)-2-methylene-5-oxo-2,3,11,11a-tetrahydro-1H-benzo[e]pyrrolo[1,2-a][1,4]diazepin-10(5H)-yl)-25,31-dimethyl-1,11,14,24,27,30-hexaoxo-4,7,18,21-tetraoxa-

10,15,25,28,31-pentaazapentatriacontan-35-oyl)pyrrolidin-2-yl)-3-methoxy-2-methylpropanamido)-3-phenylpropanoic acid (**344**).



To a solution of compound **258** (~20 mg, 0.022 mmol) in the mixture of THF (1.5 mL) and NaH₂PO₄ buffer solution (0.9 mL, 0.15 M, pH 7.5) was added compound 2,5-dioxopyrrolidin-1-yl 2,3-dibromo-4-((2-(2-(3-((S)-7-methoxy-8-((5-(((S)-7-methoxy-2-methylene-5-oxo-2,3,5,11a-tetrahydro-1H-benzo[e]pyrrolo[1,2-a][1,4]-diazepin-8-yl)oxy)pentyl)oxy)-2-methylene-5-oxo-2,3,11,11a-tetrahydro-1H-benzo[e]pyrrolo[1,2-a][1,4]-diazepin-10(5H)-yl)-3-oxopropoxy)ethoxy)ethyl)amino)-4-oxobutanoate **341** (30.0 mg, 0.027 mmol). The mixture was stirred at r.t. for 7 h, concentrated, and purified on C₁₈ preparative HPLC (Φ 2.0 cm × 25 cm) eluted with water/CH₃CN (from 90% water to 15% water in 50 min at 10 ml/min) to afford the title product **344** (26.1 mg, 63% yield) after drying under high vacuum. ESI MS m/z C₉₀H₁₃₀Br₂N₁₁O₂₂ [M+H]⁺, calcd. 1874.77, found 1874.50.


Example 110. Conjugated compound **344** to an antibody for **345**.



To a mixture of 2.0 mL of 10 mg/ml Herceptin in pH 6.0~8.0, were added 0.70 ~ 2.0 mL PBS buffer of 100 mM NaH₂PO₄, pH 6.5~7.5 buffers, TCEP (28 μL, 20 mM in water) and the compound **344** (14 μL, 20 mM in DMA). The mixture was incubated at RT for 4~16 h, then DHAA (135 μL, 50 mM) was added. After continuous incubation at RT overnight, the mixture was purified on G-25 column eluted with 100 mM NaH₂PO₄, 50 mM NaCl pH 6.0~7.5 buffer to afford 16.5~17.7 mg of conjugate compound **345** (82%~88% yield) in 13.1~15.0 ml buffer. The drug/antibody ratio (DAR) (the combination of PBD dimer and MMAF per antibody) was 3.85, determined by UPLC-QToF mass spectrum. The product was 96~99% monomer analyzed by


SEC HPLC (Tosoh Bioscience, Tskgel G3000SW, 7.8 mm ID × 30 cm, 0.5 ml/min, 100 min) and a single band measured by SDS-PAGE gel.

Example 111. Synthesis of (4R)-4-(2-((1R,3R)-1-acetoxy-3-((2S,3S)-N,3-dimethyl-2-((R)-1-methylpiperidine-2-carboxamido)pentanamido)-4-methylpentyl)thiazole-4-carboxamido)-5-(3-(12,13-dibromo-24-((S)-7-methoxy-8-((5-(((S)-7-methoxy-2-methylene-5-oxo-2,3,5,11a-tetrahydro-1H-benzo[e]pyrrolo[1,2-a][1,4]diazepin-8-yl)oxy)pentyl)oxy)-2-methylene-5-oxo-2,3,11,11a-tetrahydro-1H-benzo[e]pyrrolo[1,2-a][1,4]diazepin-10(5H)-yl)-11,14,24-trioxo-4,7,18,21-tetraoxa-10,15-diazatetracosanamido)-4-hydroxyphenyl)-2-methylpentanoic acid (**346**)



To a solution of compound **336** (20 mg, 0.021 mmol) in the mixture of THF (1 ml) and NaH₂PO₄ buffer solution (0.6 ml, 0.15 M, pH 7.5) was added compound **341** (30.0 mg, 0.027 mmol). The mixture was stirred at r.t. for 8 h, concentrated, and purified on C₁₈ preparative HPLC (Φ 2.0 cm × 25 cm) eluted with water/CH₃CN (from 90% water to 20% water in 50 min at 10 ml/min) to afford the title product **346** (26.6 mg, 64% yield) after drying under high vacuum. ESI MS m/z C₈₉H₁₂₃Br₂N₁₂O₂₂S [M+H]⁺, calcd. 1901.69, found 1901.90.


Example 112. Conjugated compound **346** to an antibody for **347**.



To a mixture of 2.0 mL of 10 mg/ml Herceptin in pH 6.0~8.0, were added 0.70 ~ 2.0 mL PBS buffer of 100 mM NaH₂PO₄, pH 6.5~7.5 buffers, TCEP (28 μL, 20 mM in water) and the compound **346** (14 μL, 20 mM in DMA). The mixture was incubated at RT for 4~16 h, then DHAA (135 μL, 50 mM) was added. After continuous incubation at RT overnight, the mixture was purified on G-25 column eluted with 100 mM NaH₂PO₄, 50 mM NaCl pH 6.0~7.5 buffer to afford 16.4~17.6 mg of the conjugate compound **347**(82%~88% yield) in 13.2~15.1 ml buffer.

The drug/antibody ratio (DAR) (the combination of PBD dimer and tubulysin analog per antibody) was 3.9, determined by UPLC-QToF mass spectrum. The product was 96~99% monomer analyzed by SEC HPLC (Tosoh Bioscience, Tskgel G3000SW, 7.8 mm ID × 30 cm, 0.5 ml/min, 100 min) and a single band measured by SDS-PAGE gel.


Example 113. Synthesis of (4R)-*tert*-butyl 5-(4-acetoxy-3-nitrophenyl)-4-((*tert*-butoxycarbonyl)amino)-2-methylpentanoate (**349**)



To a solution of compound **190** (107.1 mg, 0.252 mmol) in dichloromethane (4.0 mL) at 0 °C was added acetic anhydride (0.11 mL, 1.17 mmol) and triethylamine (0.16 mL) in sequence.


The reaction was then warmed to r.t. and stirred for 1 h, diluted with dichloromethane and washed with water and brine, dried over anhydrous Na₂SO₄, filtered and concentrated. The residue was purified by column chromatography (0-15% EA/PE) to give a colorless oil (120.3 mg, theoretical yield). MS ESI m/z calcd for C₂₃H₃₅N₂O₈ [M+H]⁺ 467.23, found 467.23.

Example 114. Synthesis of (4R)-*tert*-butyl 5-(4-acetoxy-3-aminophenyl)-4-((*tert*-butoxycarbonyl)amino)-2-methylpentanoate (**349**)



Phenyl nitrile **348** (120.3 mg, 0.258 mmol) was dissolved in ethyl acetate (5 mL) and acetic acid (0.5 mL). To which Pd/C (10 wt%, 10 mg) was added and the mixture was stirred under H₂ balloon at r.t. for 30 min before filtration through a celite pad with washing of the pad with ethyl acetate. The filtrate was concentrated and purified by column chromatography (0-25% EA/PE) to give a yellow oil (120.9 mg, theoretical yield). MS ESI m/z calcd for C₂₃H₃₇N₂O₆ [M+H]⁺ 437.26, found 437.28.


Example 115. Synthesis of 30-((2-acetoxy-5-((2R)-5-(*tert*-butoxy)-2-((*tert*-butoxycarbonyl)amino)-4-methyl-5-oxopentyl)phenyl)amino)-14,17,30-trioxo-4,7,10,21,24,27-hexaoxa-13,18-diazatriacont-15-yn-1-oic acid (**350**)



To a solution of aniline **349** (106.9 mg, 0.245 mmol) and diacid **150** (140.2 mg, 0.269 mmol) in DMF (4.5 mL) at 0 °C was added HATU (93.2 mg, 0.245 mmol) and DIPEA (0.13 mL)


in sequence. The reaction was then warmed to r.t. and stirred for 18 h, diluted with ethyl acetate and washed with water for three times and brine, dried over anhydrous Na₂SO₄, filtered and concentrated. The residue was purified by column chromatography (0-15% MeOH/CH₂Cl₂) to give a light yellow oil (70.9 mg, 31% yield). MS ESI m/z calcd for C₄₅H₇₁N₄O₁₇ [M+H]⁺ 939.47, found 939.47.

Example 116. Synthesis of 2,5-dioxopyrrolidin-1-yl 30-((2-acetoxy-5-((2R)-5- (tert-butoxy)-2-((tert-butoxycarbonyl)amino)-4-methyl-5-oxopentyl)phenyl)amino)-14,17,30-trioxo-4,7,10,21,24,27-hexaoxa-13,18-diazatriacont-15-yn-1-oate (**351**)



A mixture of acid **350** (22.3 mg, 0.0237 mmol), N-hydroxysuccinimide (13.7 mg, 0.119 mmol) and DCC (15.0 mg, 0.0728 mmol) were stirred at r.t. for 16 h. The solid was filtered off and the filtrate concentrated and purified by column chromatography (0-10% MeOH/CH₂Cl₂) to give colorless oil (14.7 mg, 60% yield). MS ESI m/z calcd for C₄₉H₇₄N₅O₁₉ [M+H]⁺ 1036.49, found 1036.49.


Example 117. Synthesis of (4R)-butyl 5-(4-acetoxy-3-(1-hydroxy-73,86,89-trioxo-3,6,9,12,15,18,21,24,27,30,33,36,39,42,45,48,51,54,57,60,63,66,69,76,79,82,93,96,99-nonacosaoxa-72,85,90-triazadohect-87-ynamido)phenyl)-4-((tert-butoxycarbonyl)amino)-2-methylpentanoate (**352**)



To a solution of H₂N-PEG₂₄-OH (15.2 mg, 0.0142 mmol) in a mixture of EtOH/CH₂Cl₂/H₂O (v/v/v 5:3:1, 0.5 mL), triethylamine (25 μL) was added, followed by a solution of NHS ester **351** (14.7 mg, 0.0142 mmol) in EtOH/CH₂Cl₂/H₂O (v/v/v 5:3:1, 1.0 mL). After stirring at r.t. for 1 h, the reaction was concentrated and purified by column chromatography (0-15% MeOH/CH₂Cl₂) to give colorless oil. MS ESI m/z calcd for C₉₃H₁₆₈N₅O₄₀ [M+H]⁺ 1995.12, found 1995.12.


Example 118. Synthesis of (4R)-4-(2-((1R,3R)-1-acetoxy-3-((2S,3S)-N,3-di methyl-2-((R)-1-methylpiperidine-2-carboxamido)pentanamido)-4-methylpentyl)thiazole-4-carboxamido)-5-(4-acetoxy-3-(1-hydroxy-73,86,89-trioxo-

3,6,9,12,15,18,21,24,27,30,33,36,39,42,45,48,51,54,57,60,63,66,69,76,79,82,93,96,99-nonacosaoxa-72,85,90-triazadohect-87-ynamido)phenyl)-2-methylpentanoic acid (**353**)




Compound **352** (0.005 mmol) was dissolved in CH₂Cl₂ (1.0 mL) and treated with TFA (1.0 mL) at r.t. for 1.5 h then concentrated and re-dissolved in DMF (0.5 mL), to which pentafluorophenyl ester **142** (0.02 mmol) in DMF (0.5 mL) and DIPEA (3.5 μL) were added. The reaction was stirred at r.t. for 2 h and then concentrated. The residue was purified by preparative reverse phase HPLC (C₁₈ column, acetonitrile/water) to give a colorless oil (1.13 mg, 10% yield). MS ESI m/z calcd for C₁₁₀H₁₉₂N₉O₄₃S [M+H]⁺ 2359.28, found 2359.28.

Example 119. Synthesis of (3R,4S,7S,10S)-4-((S)-sec-butyl)-7,10-diisopropyl-3-(2-((S)-2-((1R,2R)-1-methoxy-3-(((S)-1-methoxy-1-oxo-3-phenylpropan-2-yl)amino)-2-methyl-3-oxopropyl)pyrrolidin-1-yl)-2-oxoethyl)-5,11-dimethyl-6,9,12,25,28-pentaoxo-2,15,18,21,32,35,38-heptaoxa-5,8,11,24,29-pentaazahentetracont-26-yn-41-oic acid (**355**)




To a solution of **230** (26.1 mg, 0.035 mmol) and diacid **150** (55 mg, 0.106mmol) in dichloromethane (2 mL) at 0 °C was added Brop (16 mg, 0.042 mmol), followed by diisopropylethylamine (22 μL, 0.12 mmol). The reaction mixture was stirred at 0 °C for 10 min, then warmed to r.t. and stirred overnight. The reaction mixture was diluted with ethyl acetate (50 mL), washed with water (10 mL), saturated aqueous sodium chloride (10 mL), dried over sodium sulfate, and concentrated in vacuo. The residue was purified by reverse phase HPLC with a gradient of MeCN/H₂O to afforded compound **355** (30 mg, 68% yield) as a white solid. MS ESI m/z calcd. for C₆₂H₁₀₁N₇O₁₉ [M+H]⁺: 1248.72, found: 1248.72.

Example 120. Synthesis of (3R,4S,7S,10S)-perfluorophenyl4-((S)-sec-butyl)-7,10-diisopropyl-3-(2-((S)-2-((1R,2R)-1-methoxy-3-(((S)-1-methoxy-1-oxo-3-phenylpropan-2-yl)amino)-2-methyl-3-oxopropyl)pyrrolidin-1-yl)-2-oxoethyl)-5,11-dimethyl-6,9,12,25,28-pentaoxo-2,15,18,21,32,35,38-heptaoxa-5,8,11,24,29-pentaazahentetracont-26-yn-41-oate (**356**).




To a solution of compound **355** (15 mg, 0.012 mmol) dissolved in ethyl acetate (1 mL) was added pentafluorophenol (3.3 mg, 0.018 mmol) and DCC (3.7 mg, 0.018 mmol). The reaction mixture was stirred at room temperature overnight and then filtered, washing the filter cake with 2 mL of EtOAc. The filtrate was used immediately without further purification or concentration.

Example 121. Synthesis of (S)-methyl 2-((2R,3R)-3-((S)-1-((104S,107S, 110S,111R)-110-((S)-sec-butyl)-1-hydroxy-104,107-diisopropyl-111-methoxy-103,109-dimethyl-73,86,89,102,105,108-hexaoxa-3,6,9,12,15,18,21,24,27,30,33,36,39,42,45,48,51,54,57,60,63,66,69,76,79,82,93,96,99-nonacosaoxa-72,85,90,103,106,109-hexaazatridecahect-87-yn-113-oyl)pyrrolidin-2-yl)-3-methoxy-2-methylpropanamido)-3-phenylpropanoate (**357**)



To a solution of compound **356** (0.012 mmol) dissolved in DMA (1 mL) was added HO-PEG₂₄-NH₂ (19 mg, 0.018 mmol) and diisopropylethylamine (4.2 μ L, 0.024 mmol). The reaction mixture was stirred overnight and concentrated in vacuo. The residue was purified by reverse phase HPLC with a gradient of MeCN/H₂O to afford compound **357** (5 mg, 18% yield) as a colorless oil. MS ESI *m/z* calcd. for C₁₁₀H₁₉₈N₈O₄₂ [M+H]⁺ 2304.36, found 2304.36.


Example 122. Synthesis of compound **361**



To a solution of compound **161** (4.7 mg, 0.011 mmol) dissolved in ethyl acetate (1 mL) was added pentafluorophenol (10.1 mg, 0.055 mmol) and DCC (11.3 mg, 0.055 mmol). The reaction mixture was stirred at room temperature overnight and then filtered through celite, washing the filter cake with 2 mL of EtOAc. The filtrate was concentrated and re-dissolved in


DMA (2 mL), to which (4R)-ethyl 4-(2-((1R,3R)-1-acetoxy-3-((2S,3S)-N,3-dimethyl-2-((R)-1-methylpiperidine-2-carboxamido)pentanamido)-4-methylpentyl)thiazole-4-carboxamido)-5-(3-amino-4-hydroxyphenyl)-2-methylpentanoate **360** (8 mg, 0.0102 mmol) and DIPEA (3.8 μ L, 0.022 mmol) were added. The reaction mixture was stirred at r.t. overnight, and then the solvent was removed under reduced pressure and the residue was purified by reverse phase HPLC with a gradient of MeCN/H₂O to give the title product **361** (4.5mg, 38% yield). ESI MS m/z: calcd for C₅₈H₈₇N₈O₁₆S [M+H]⁺ 1183.59, found 1183.58.

Example 123. Synthesis of compound **364**




To a solution of compound **161** (0.43 g, 0.997 mmol) dissolved in ethyl acetate (20 mL) was added pentafluorophenol (0.92 g, 4.98 mmol) and DCC (1.02 g, 4.98 mmol). The reaction mixture was stirred at room temperature overnight and then filtered through celite, washing the filter cake EtOAc. The filtrate was concentrated and re-dissolved in DMA (10 mL), to which (4R)-5-(3-amino-4-hydroxyphenyl)-4-((tert-butoxycarbonyl)amino)-2-methylpentanoic acid **363** (0.34 g, 0.997 mmol) and DIPEA (0.35 mL, 2 mmol) were added. The reaction mixture was stirred at r.t. overnight and concentrated under reduced pressure. The residue was diluted with H₂O and extracted with EtOAc. The combined the organic layers were dried over Na₂SO₄, concentrated and purified by SiO₂ column chromatography to give the title product **364** (0.3 g, 42% yield). ESI MS m/z: calcd for C₃₅H₅₁N₄O₁₃ [M+H]⁺:735.34, found 735.34.

Example 124. Synthesis of compound **365**




Compound **364** (0.1 g, 0.14 mmol) was dissolved in DCM (10 mL), treated with TFA (5 mL) and stirred at r.t. for 2 h, then concentrated. The residue was dissolved in DMA (8 mL), to which pentafluorophenyl ester **142** (0.15 g, 0.21 mmol) and DIPEA (37 μ L, 0.21mmol) were added and the reaction was stirred overnight. The reaction solution was concentrated and purified by reverse phase HPLC with a gradient of MeCN/H₂O to give the product **365** (47.2 mg, 30% yield). ESI MS m/z: calcd for C₅₆H₈₃N₈O₁₆S [M+H]⁺:1155.56, found 1155.56.

Example 125. Synthesis of 24-((5-((2R)-5-(*tert*-butoxy)-2-((*tert*-butoxycarbonyl) amino)-4-methyl-5-oxopentyl)-2-((*tert*-butyldimethylsilyl)oxy)phenyl)amino)-11,14,24-trioxo-4,7,18,21-tetraoxa-10,15-diazatetracos-12-yn-1-oic acid (**367**)




Compound **192** (0.085 g, 0.197 mmol), compound **161** (0.1 g, 0.197 mmol) and HATU (0.112 mg, 0.30 mmol) were mixed in DMF (6 ml), to which TEA (55 μ L, 0.40 mmol) was added. The reaction mixture was stirred at r.t. overnight. The solvent was removed under reduced pressure and the residue purified by SiO₂ column chromatography to give the title product **367** (0.065 mg, 36% yield). ESI MS *m/z*: calcd for C₄₅H₇₅N₄O₁₄Si [M+H]⁺ 923.50, found 923.50.

Example 126. Synthesis of 24-((5-((2R)-5-(*tert*-butoxy)-2-((*tert*-butoxycarbonyl) amino)-4-methyl-5-oxopentyl)-2-hydroxyphenyl)amino)-11,14,24-trioxo-4,7,18,21-tetraoxa-10,15-diazatetracos-12-yn-1-oic acid (**368**)



Compound **367** (64.8 mg, 0.070 mmol) was dissolved in DCM (3 mL), TBAF(1 M, 0.1 mL) was added, and the reaction was stirred at r.t. for 1 h, then concentrated and purified by SiO₂ column chromatography to give the title product **368** (34.5 mg, 61% yield). ESI MS *m/z*: calcd for C₃₉H₆₁N₄O₁₄ [M+H]⁺ 809.41, found 809.41.


Example 127. Synthesis of 24-((5-((2R)-2-(2-((1R,3R)-1-acetoxy-3-((2S,3S)-N,3-dimethyl-2-((R)-1-methylpiperidine-2-carboxamido)pentanamido)-4-methylpentyl)thiazole-4-carboxamido)-4-carboxypentyl)-2-hydroxyphenyl)amino)-11,14,24-trioxo-4,7,18,21-tetraoxa-10,15-diazatetracos-12-yn-1-oic acid (**369**)



Compound **368** (34.5 mg, 0.043 mmol) was dissolved in DCM (1 mL), and then TFA (0.5 mL) was added. The reaction was stirred at r.t. for 2 h, then concentrated and re-dissolved in


DMA(4 mL), to which pentafluorophenyl ester **142** (45 mg, 0.065 mmol) and DIPEA (15 μ L, 0.09mmol) were added and the reaction was stirred overnight and then concentrated and purified by reverse phase HPLC with a gradient of MeCN/H₂O to give the title product **369** (22.8 mg, 46% yield). ESI MS m/z: calcd for C₅₆H₈₅N₈O₁₇S [M+H]⁺ 1173.57, found 1173.58.

Example 128. Synthesis of (4R)-ethyl 5-(3-(4-(((benzyloxy)carbonyl)amino)butanamido)-4-((tert-butyldimethylsilyl)oxy)phenyl)-4-((tert-butoxycarbonyl)amino)-2-methylpentanoate (**372**)




2,5-dioxopyrrolidin-1-yl 4-(((benzyloxy)carbonyl)amino)butanoate (0.396 g, 1.2 mmol) and (4R)-ethyl 5-(3-amino-4-hydroxyphenyl)-4-((tert-butoxycarbonyl) amino)-2-methylpentanoate (0.44 g, 1.2 mmol) were dissolved in EtOH (10 mL), and phosphate buffer solution (pH=7.5, 0.1M, 2ml) was added. The reaction mixture was stirred at r.t. overnight and then the solvent was removed under reduced pressure and the residue purified by SiO₂ column chromatography to give the title product **372** (0.485g, 70%). ESI: m/z: calcd for C₃₁H₄₄N₃O₈ [M+H]⁺:586.31, found 586.31.

Example 129. Synthesis of (4R)-ethyl 5-(3-(4-aminobutanamido)-4-((tert-butyl dimethylsilyl)oxy)phenyl)-4-((tert-butoxycarbonyl)amino)-2-methylpentanoate (**373**)




Compound **372** (0.35 g, 0.5 mmol) was dissolved in MeOH (5 ml), and Pd/C (10 wt%, 35 mg) was then added. The reaction mixture was stirred at r.t. under H₂ balloon overnight, then filtered through celite and the filtrate was concentrated under reduced pressure to give the title product **373** (0.22 g, 79% yield). ESI MS m/z: calcd for C₂₉H₅₂N₃O₆Si [M+H]⁺:566.35, found 566.35.

Example 130. Synthesis of 4-((4-(((5-ethoxy-4-methyl-5-oxopentyl)-2-((tert-butyldimethylsilyl)oxy)phenyl)amino)-4-oxobutyl)amino)-4-oxobut-2-ynoic acid (**374**)




Compound **373** (0.22 g, 0.39 mmol) was dissolved in NMP (2 mL), and actylenedicarboxylic acid (0.089g, 0.78mmol) was added at 0 °C. After stirring for 10 min, DMTMM (0.1 g, 0.39 mmol) was added. The mixture was stirred at 0 °C overnight. The reaction was washed with H₂O (100 mL) and the aqueous layer was extracted with EtOAc. The combined organic layers were dried over Na₂SO₄, filtered, concentrated under reduced pressure and the residue was purified by SiO₂ column chromatography with a gradient of DCM/ MeOH to give the title product **374** (0.05 g, 20% yield). ESI MS m/z: calcd for C₃₃H₅₂N₃O₉Si [M+H]⁺:662.34, found 662.34.

Example 131. Synthesis of 4-((4-((5-((2R)-2-((tert-butoxycarbonyl)amino)-5-ethoxy-4-methyl-5-oxopentyl)-2-hydroxyphenyl)amino)-4-oxobutyl)amino)-4-oxobut-2-ynoic acid (**375**)




Compound **374** (50 mg, 0.076 mmol) was dissolved in DCM (2 mL), and then TBAF (1 M, 0.05 mL) was added. The reaction was stirred at r.t. for 1 h, and then concentrated and purified by SiO₂ column chromatography to give the title product **375** (10.5 mg, 25% yield). ESI MS m/z: calcd for C₂₇H₃₈N₃O₉ [M+H]⁺ 548.25, found 548.25.

Example 132. Synthesis of 4-((4-((5-((2R)-2-((1R,3R)-1-acetoxy-3-((2S,3S)-N,3-dimethyl-2-((R)-1-methylpiperidine-2-carboxamido)pentanamido)-4-methylpentyl)thiazole-4-carboxamido)-5-ethoxy-4-methyl-5-oxopentyl)-2-hydroxyphenyl)amino)-4-oxobutyl)amino)-4-oxobut-2-ynoic acid (**376**)




Compound **375** (10.5 mg, 0.02 mmol) was dissolved in DCM (1 mL), and TFA (0.5 mL) was added. The reaction was stirred at r.t. for 2 h, then concentrated and re-dissolved in DMA (2 mL), to which pentafluorophenyl ester **142** (21 mg, 0.03 mmol) and DIPEA (7 µl, 0.04 mmol) were added. The reaction was stirred overnight and then concentrated and the residue was purified by reverse phase HPLC with a gradient of MeCN/H₂O to give the title product **376** (9 mg, 47% yield). ESI MS m/z: calcd for C₄₈H₇₀N₇O₁₂S [M+H]⁺ 968.47, found 968.47.

Example 133. Synthesis of 2-((6S,9S,12R,14R)-9-((S)-sec-butyl)-14-hydroxy -6,12-diisopropyl-2,2,5,11-tetramethyl-4,7,10-trioxo-3-oxa-5,8,11-triazatetradecan-14-yl)thiazole-4-carboxylic acid (**381**)




To a solution of Boc-N-Me-L-Val-OH (33 mg, 0.14 mmol) in EtOAc was added pentafluorophenol (39 mg, 0.21 mmol) and DCC (32 mg, 0.154 mmol). The reaction mixture was stirred at r.t. for 16 h and then filtered over a celite pad, with washing of the pad with EtOAc. The filtrate was concentrated and re-dissolved in DMA (2 mL), and then 2-((1R,3R)-3-((2S,3S)-2-amino-N,3-dimethylpentanamido)-1-hydroxy-4-methylpentyl)thiazole-4-carboxylic acid **380** (52 mg, 0.14 mmol) and DIPEA (48.5 μ L, 0.28 mmol) were added. The reaction mixture was stirred at r.t. for 24 h and then concentrated and purified by reverse phase HPLC (C₁₈ column, 10-100% acetonitrile/water) to afford compound **381** (40.2 mg, 49% yield). ESI MS m/z: calcd for C₂₈H₄₉N₄O₇S [M+H]⁺: 585.32, found 585.32.

Example 134. Synthesis of 2-((6S,9S,12R,14R)-9-((S)-sec-butyl)-6,12-diisopropyl-2,2,5,11-tetramethyl-4,7,10,16-tetraoxo-3,15-dioxo-5,8,11-triazaheptadecan-14-yl)thiazole-4-carboxylic acid (**382**)



Compound **381** (40 mg, 0.069 mmol) was dissolved in pyridine (8 mL), to which acetic anhydride (20.4 mg, 0.2 mmol) was added at 0 °C and the reaction was allowed to warm to r.t. and stirred overnight. The mixture was concentrated and the residue purified by SiO₂ column chromatography with a gradient of DCM/MeOH to give the title product **382** (48.1 mg, ~100% yield). ESI MS m/z: calcd for C₃₀H₅₁N₄O₈S [M+H]⁺ 627.33, found 627.33.


Example 135. Synthesis of (4R)-4-(2-((6S,9S,12R,14R)-9-((S)-sec-butyl)-6,12-diisopropyl-2,2,5,11-tetramethyl-4,7,10,16-tetraoxo-3,15-dioxo-5,8,11-triazaheptadecan-14-yl)thiazole-4-carboxamido)-2-methyl-5-phenylpentanoic acid (**383**)



To a solution of compound **382** (48.1 mg, 0.077 mmol) in EtOAc was added pentafluorophenol (21.2 mg, 0.115 mmol) and DCC (17.4 mg, 0.085 mmol). The reaction mixture was stirred at r.t. for 16 h and then filtered over a celite pad, with washing of the pad


with EtOAc. The filtrate was concentrated and re-dissolved in DMA (4 mL), and then (4R)-4-amino-2-methyl-5-phenylpentanoic acid (20.7 mg, 0.1 mmol) and DIPEA (26.8 μ L, 0.154 mmol) were added. The reaction mixture was stirred at r.t. for 24 h and then concentrated and purified by reverse phase HPLC (C_{18} column, 10-100% acetonitrile/water) to afford compound **383** (63 mg, ~100% yield). ESI MS m/z: calcd for $C_{42}H_{66}N_5O_9S$ $[M+H]^+$ 816.45, found 816.45.

Example 136. Synthesis of 11,14,24-trioxo-24-(perfluorophenoxy)-4,7,18,21-tetraoxa-10,15-diazatetracos-12-yn-1-oic acid (**378**)



To a solution of compound **161** (45.4 mg, 0.11 mmol) in EtOAc was added pentafluorophenol (21.3 mg, 0.12 mmol) and DCC (24.7 mg, 0.12 mmol). The reaction mixture was stirred at r.t. for 16 h and then filtered over a celite pad, with washing of the pad with EtOAc. The filtrate was concentrated and used for the next step without further purification.

Example 137. Synthesis of (4R,6R,9S,12S)-9-((S)-sec-butyl)-4-(4-(((2R)-4-carboxy-1-phenylpentan-2-yl)carbamoyl)thiazol-2-yl)-6,12-diisopropyl-7,13-dimethyl-2,8,11,14,24,27-hexaoxo-3,17,20,31,34-pentaoxa-7,10,13,23,28-pentaazaheptatriacont-25-yn-37-oic acid (**384**)



Compound **383** (63 mg, 0.077 mmol) was dissolved in DCM (5 mL), and TFA (2.5 mL) was then added. The reaction was stirred at r.t. for 2 h, then concentrated and re-dissolved in DMA (4 mL), to which compound **378** (65.8mg, 0.11 mmol) and DIPEA (27 μ L, 0.154 mmol) were added. The reaction mixture was stirred at r.t. overnight and then concentrated and the residue purified by reverse phase HPLC (C_{18} column, 10-100% acetonitrile/water) to afford compound **384** (14 mg, 16% yield). ESI MS m/z: calcd for $C_{55}H_{84}N_7O_{16}S$ $[M+H]^+$: 1130.56, found 1130.57.

Example 138. General method of conjugation of compound **196, 200, 205, 259, 267, 269, 271, 273, 274, 276, 288, 292, 299, 301, 307, 310, 313, 316, 318, 320, 325, 327, 328, 333, 337, 339, 342, 344, 346, 353, 355, 357, 361, 365, 369, 376, 384, 388, 393, 394, 397, 400, 401, 404, 406, 409, 411, or 413** independently to an antibody (Herceptin).

To a mixture of 2.0 mL of 10 mg/ml Herceptin in pH 6.0~8.0, were added of 0.70 ~ 2.0 mL PBS buffer of 100 mM NaH_2PO_4 , pH 6.5~8.5 buffers, TCEP (14-35 μ L, 20 mM in water) and the

compound **196, 200, 205, 259, 260, 267, 269, 271, 273, 274, 276, 288, 292, 299, 301, 307, 310, 313, 316, 318, 320, 325, 327, 328, 333, 337, 339, 342, 344, 346, 353, 355, 357, 361, 365, 369, 376, 384, 388, 393, 394, 397, 400, 401, 404, 406, 409, 411, or 413** (14-28 μ L, 20 mM in DMA). The mixture was incubated at RT for 4~16 h, then DHAA (135 μ L, 50 mM) was added in. After continuous incubation at RT overnight, the mixture was purified on G-25 column eluted with 100 mM NaH_2PO_4 , 50 mM NaCl pH 6.0~7.5 buffer to afford 12.0~18.4 mg of the conjugate compound **196a, 201, 206, 261, 267, 270, 272, 275, 277, 289, 293, 300, 302, 308, 311, 314, 317, 319, 321, 326, 329, 334, 338, 339a, 343, 345, 347, 354, 355a, 358, 362, 366, 370, 377, 385, 386, 389, 395, 396, 398, 402, 405, 407, 410, 412, or 414** (60%~91% yield) accordingly in 13.0~15.8 ml buffer. The drug/antibody ratio (DAR) was 1.9~4.0, which was determined via UPLC-QTOF mass spectrum. It was 94~99% monomer analyzed by SEC HPLC (Tosoh Bioscience, Tskgel G3000SW, 7.8 mm ID x 30 cm, 0.5 ml/min, 100 min) and a single band measured by SDS-PAGE gel.

Example 139. General method of conjugation of compound **248, 250, 252, 254, and 255**, independently to Herceptin (an antibody).

To a mixture of 2.0 mL of 10 mg/ml Herceptin in pH 6.0~8.0, 0.70 ~ 2.0 mL of 100 mM NaH_2PO_4 , 1 mM Na_2SO_3 , pH 6.5~8.5 buffer, and the compound **248, 250, 252, 254 and 255** independently (21-28 μ L, 20 mM in DMA) incubated for 1 h, was added TCEP (14-35 μ L, 20 mM in water). After the mixture was continued to incubate at RT for 8~24 h, DHAA (135 μ L, 50 mM) was added in. And the solution was continuously incubated for another 12 h. then purified on G-25 column eluted with 100 mM NaH_2PO_4 , 50 mM NaCl pH 6.0~7.5 buffer to afford 13.8~17.6 mg of the conjugate compound **249, 251, 253, and 256** respectively (69%~88% yield) in 13.6~15.2 ml buffer. The drug/antibody ratio (DAR) was 1.9~4.0, which was determined via UPLC-QTOF mass spectrum. It was 96~99% monomer analyzed by SEC HPLC (Tosoh Bioscience, Tskgel G3000SW, 7.8 mm ID x 30 cm, 0.5 ml/min, 100 min) and a single band measured by SDS-PAGE gel or two bands when a reduce reagent DTT was in the SDS Page.

Example 140. In vitro cytotoxicity evaluation of conjugates **196a, 201, 206, 354, 358, 362, 366, 370, 377, 385 and 407** in comparison with T-DM1:

The cell line used in the cytotoxicity assays was NCI-N87, a human gastric carcinoma cell line; The cells were grown in RPMI-1640 with 10% FBS. To run the assay, the cells (180 μ L, 6000 cells) were added to each well in a 96-well plate and incubated for 24 hours at 37°C with 5% CO_2 . Next, the cells were treated with test compounds (20 μ L) at various concentrations in appropriate cell culture medium (total volume, 0.2 mL). The control wells contain cells and the medium but lack the test compounds. The plates were incubated for 120 hours at 37°C with 5%

CO₂. MTT (5mg/ml) was then added to the wells (20 µl) and the plates were incubated for 1.5hr at 37°C. The medium was carefully removed and DMSO (180 µl) was added afterward. After it was shaken for 15min, the absorbance was measured at 490nm and 570nm with a reference filter of 620nm. The inhibition% was calculated according to the following equation: inhibition% = [1- (assay-blank)/(control-blank)] × 100.

The cytotoxicity results:

	DAR	N87 cell (Ag+) IC ₅₀ (nM)
Conjugate 196a	2.8	3.9 nM
Conjugate 201	2.7	1.3 nM
Conjugate 206	2.9	1.7 nM
Conjugate 354	2.1	28.4nM
Conjugate 358	2.2	643.7 nM
Conjugate 362	2.8	3.1 nM
Conjugate 366	2.9	2.3 nM
Conjugate 370	3.2	1.3 nM
Conjugate 377	2.5	62.7 nM
Conjugate 385	2.8	561.6 nM
T-DM1	3.5	0.07 nM

The conjugates with bridge linkers had equal or less potency than T-DM1in vitro.

Example 141. Antitumor Activity In vivo.

The in vivo efficacy of conjugates **196a**, **201**, **206**, **354**, **358**, **370**, **377**, **385** and **407** along with T-DM1 were evaluated in a human gastric carcinoma N-87 cell line tumorxenograft models. Five-week-old female BALB/c Nude mice (30 animals) wereinoculated subcutaneously in the area under the right shoulder with N-87 carcinoma cells (5×10^6 cells/mouse) in 0.1mL of serum-free medium. The tumors were grown for 8 days to anaverage size of 130 mm³. The animals were then randomly divided into 12 groups (6 animals per group). The first group of mice served as the


control group and was treated with the phosphate-buffered saline vehicle. The remaining 11 groups were treated with conjugates **196a**, **201**, **206**, **354**, **358**, **362**, **365**, **370**, **377**, **385**, **407** and T-DM1 respectively at dose of 5 mg/Kg administered intravenously. Three dimensions of the tumor were measured every 4 days and the tumor volumes were calculated using the formula
5 tumor volume = $\frac{1}{2}(\text{length} \times \text{width} \times \text{height})$. The weight of the animals was also measured at the same time. A mouse was sacrificed when any one of the following criteria was met: (1) loss of body weight of more than 20% from pretreatment weight, (2) tumor volume larger than 1500 mm³, (3) too sick to reach food and water, or (4) skin necrosis. A mouse was considered to be tumor-free if no tumor was palpable.

10 The results were plotted in Figures 36. All the 9 conjugates did not cause the animal body weight loss. And the animals at control group were sacrificed at day 30 due to the tumor volume larger than 1200 mm³ and they were too sick. All 6/6 animals at the groups of compounds **201** and **407** had completely no tumor measurable at day 18 till day 68 (the end of experiment). In contrast T-DM1 at dose of 6 mg/Kg was not able to eliminate the tumors and it only inhibited the
15 tumor growth for 38 days. Conjugate compounds **196a**, **206**, **277**, and **354** did not eradicate the tumor at dose of 6 mg/Kg, but had better antitumor activity than T-DM1. The conjugates 358 and 385 had worse antitumor activity than T-DM1. More importantly, all animals treated with conjugate compounds **196a**, **201**, **206**, **277**, **354**, **358**, **385** and **407** had no or less liver toxicities than the animals treated with T-DM1 when measured levels of alanine aminotransferase (ALT) and aspartate aminotransferase (AST) in serum at the end of experiment (data not shown). This
20 demonstrates that the conjugates with the bridge linkers of this patent application would have broader therapeutical applications than the traditional conjugates.

CLAIMS

What is claimed is:

1. A bridge linker compound of the Formula (I)




Wherein:

“—” represents an optional single bond.

“---” represents either a single bond, or a double bond, or a triple bond.

It provided that when --- represents a single bond, both U and U' are not H; when --- represents a double bond, either U or U' can be H, but are not H at the same time; when --- represents a triple bond, both U and U' can be absent.



Wherein the component: $\zeta - \overset{\text{O}}{\parallel} \text{C} - \text{U} \text{---} \text{U}' - \overset{\text{O}}{\parallel} \text{C} - \zeta$, which is 2,3-disubstituted succinic group, or 2-monosubstituted or 2,3-disubstituted fumaric group, or 2-monosubstituted or 2,3-disubstituted maleic group, or acetylenedicarboxyl group, is capable of reacting with a pair of thiols of the cell-binding agent; The pair of thiols are preferred pairs of sulfur atoms reduced from the interchain disulfide bonds of the cell-binding agent by a reducing agent, such as dithiothreitol (DTT), dithioerythritol (DTE), L-glutathione (GSH), tris (2-carboxyethyl) phosphine (TCEP), 2-mercaptoethylamine (β -MEA), or/and beta mercaptoethanol (β -ME, 2-ME).

U and U' represent the same or different leaving group that can be substituted by a thiol. Such leaving groups are a halide (e.g., fluoride, chloride, bromide, and iodide), methanesulfonyl (mesyl), toluenesulfonyl (tosyl), trifluoromethyl-sulfonyl (triflate), trifluoromethylsulfonate, nitrophenoxyl, N-succinimidylloxyl (NHS), phenoxyl; dinitrophenoxyl; pentafluorophenoxyl, tetrafluorophenoxyl, trifluorophenoxyl, difluorophenoxyl, monofluorophenoxyl, pentachlorophenoxyl, 1H-imidazole-1-yl, chlorophenoxyl, dichlorophenoxyl, trichlorophenoxyl, tetrachlorophenoxyl, N-(benzotriazol-yl)oxyl, 2-ethyl-5-phenylisoxazolium-yl, phenyloxadiazol-yl (ODA), oxadiazol-yl, or an intermediate molecule generated with a condensation reagent for Mitsunobu reactions.

R₁ can be absent, or can be selected from C₁-C₈ of alkyl; C₂-C₈ of heteroalkyl, alkylcycloalkyl, heterocycloalkyl; C₃-C₈ of aryl, Ar-alkyl, heterocyclic, carbocyclic, cycloalkyl, heteroalkylcycloalkyl, alkylcarbonyl, heteroaryl; or 1-8 carbon atoms of esters, ether, or amide; or polyethyleneoxy unit of formula (OCH₂CH₂)_p or (OCH₂CH(CH₃))_p, wherein p is an integer from 0 to about 1000, or combination thereof.


Additionally R_1 is a chain of atoms selected from C, N, O, S, Si, and P, preferably having 0~500 atoms, which covalently connects to X_1 and Z_1 . The atoms used in forming the R_1 may be combined in all chemically relevant ways, such as forming alkane, alkenylene, alkynylene, ethers, polyoxyalkylene, esters, amines, imines, polyamines, hydrazines, hydrazones, amides, ureas, semicarbazides, carbazides, alkoxyamines, alkoxyamines, urethanes, amino acids, peptides, acyloxyamines, hydroxamic acids, or combination thereof.

X_1 and X_2 are independently selected from NH; $NHNH$; $N(R_3)$; $N(R_3)N(R_3')$; O; S; $B(R_3)$; $Si(R_3)N(R_3')$; $P(O)(R_3)$; C_1 - C_8 of alkyl; C_2 - C_8 of heteroalkyl, alkylcycloalkyl, heterocycloalkyl; C_3 - C_8 of aryl, Ar-alkyl, heterocyclic, carbocyclic, cycloalkyl, heteroalkylcycloalkyl, alkylcarbonyl, heteroaryl; or 1~8 amino acids; Wherein R_3 and R_3' are independently H; C_1 - C_8 of alkyl; C_2 - C_8 of heteroalkyl, alkylcycloalkyl, heterocycloalkyl; C_3 - C_8 of aryl, Ar-alkyl, heterocyclic, carbocyclic, cycloalkyl, heteroalkylcycloalkyl, alkylcarbonyl, heteroaryl; or 1-8 carbon atoms of esters, ether, or amide; or polyethyleneoxy unit of formula $(OCH_2CH_2)_p$ or $(OCH_2CH(CH_3))_p$, wherein p is an integer from 0 to about 1000, or combination thereof. In addition, X_1 and X_2 can be independently absent.

R_2 is independently selected from OH, H, NH_2 ; SH; $NHNH_2$; $N(R_3)(R_3')$; $N(R_3)NH(R_3')$; polyethyleneoxy unit of formula $(OCH_2CH_2)_pOR_3$, or $(OCH_2CH(CH_3))_pOR_3$, or $NH(CH_2CH_2O)_pR_3$, or $NH(CH_2CH(CH_3)O)_pR_3$, or $N[(CH_2CH_2O)_pR_3][(CH_2CH_2O)_pR_3']$, or $(OCH_2CH_2)_pCOOR_3$, or $CH_2CH_2(OCH_2CH_2)_pCOOR_3$, wherein p and p' are independently an integer selected from 0 to about 1000, or combination thereof; C_1 - C_8 of alkyl; C_2 - C_8 of heteroalkyl, alkylcycloalkyl, heterocycloalkyl; C_3 - C_8 of aryl, Ar-alkyl, heterocyclic, carbocyclic, cycloalkyl, heteroalkylcycloalkyl, alkylcarbonyl, heteroaryl; Wherein R_3 and R_3' are independently H; C_1 - C_8 of alkyl; C_2 - C_8 of heteroalkyl, alkylcycloalkyl, heterocycloalkyl; C_3 - C_8 of aryl, Ar-alkyl, heterocyclic, carbocyclic, cycloalkyl, heteroalkylcycloalkyl, alkylcarbonyl, heteroaryl; or 1-8 carbon atoms of esters, ether, or amide; or 1~8 amino acids; or polyethyleneoxy unit of formula $(OCH_2CH_2)_p$ or $(OCH_2CH(CH_3))_p$, wherein p is an integer from 0 to about 1000, or combination thereof.

m_1 and m_2 are independently an integer from 1 to 30, preferably from 1 to 10.


Z_1 is a function group that enables to react with a cytotoxic drug, to form a disulfide, thioether, thioester, peptide, hydrazone, ether, ester, carbamate, carbonate, amine (secondary, tertiary, or quaternary), imine, cycloheteroalkane, heteroaromatic, alkoxy or amide bond; The function groups of Z_1 is displayed below:



$\text{N(R}_1\text{), or CH}_2\text{; R}_3\text{ and R}_5\text{ are H, R}_1\text{, aromatic, heteroaromatic, or aromatic group wherein one or several H atoms are replaced independently by -R}_1\text{, -halogen, -OR}_1\text{, -SR}_1\text{, -NR}_1\text{R}_2\text{, -NO}_2\text{, -S(O)R}_1\text{, -S(O)}_2\text{R}_1\text{, or -COOR}_1\text{; Lv}_3\text{ is a leaving group selected from methanesulfonyl (mesyl), toluenesulfonyl (tosyl), trifluoromethyl-sulfonyl (triflate), trifluoromethylsulfonate, nitrophenoxyl, N-succinimidylxyl (NHS), phenoxyl; dinitrophenoxyl; pentafluorophenoxy, tetrafluorophenoxy, trifluorophenoxy, difluorophenoxy, monofluoro-phenoxyl, pentachlorophenoxy, 1H-imidazole-1-yl, chlorophenoxy, dichlorophenoxy, trichlorophenoxy, tetrachlorophenoxy, N-(benzotriazol-}$

yl)oxyl, 2-ethyl-5-phenylisoxazolium-yl, phenyloxadiazol-yl (ODA), oxadiazol-yl, or an intermediate molecule generated with a condensation reagent for Mitsunobu reactions. Wherein R₁ and R₂ are defined above.

2. A cell-binding agent-drug conjugate compound of Formula (II)



Wherein:

Cb represents a cell-binding agent, preferably an antibody.

Inside the bracket (parentheses) are the linker-drug components that are conjugated to pairs of thiols of the cell-binding agent/molecule. The thiols are preferred pairs of sulfur atoms reduced from the interchain disulfide bonds of the cell-binding agent by a reduction agent selected from dithiothreitol (DTT), dithioerythritol (DTE), L-glutathione (GSH), tris (2-carboxyethyl) phosphine (TCEP), 2-mercaptoethylamine (β-MEA), or/and beta mercaptoethanol (β-ME, 2-ME).


Drug₁ represents a cytotoxic agent or a drug, which is linked to the cell-binding agent via the bridge linker of the patent through R₁ containing an alkane, alkylene, alkenylene, alkynylene, aromatic, ether, polyoxyalkylene, ester, amine, imine, polyamine, hydrazine, hydrazone, amide, urea, semicarbazide, carbazide, alkoxyamine, urethanes, amino acid, peptide, acyloxyamine, hydroxamic acid, disulfide, thioether, thioester, carbamate, carbonate, heterocyclic ring, heteroalkyl, heteroaromatic, or alkoxime bond, or combination above thereof.

“ $\equiv$ ” represents either single bond or double bond.

Inside the square bracket are agents that are conjugated to a cell-binding molecule through a pair of sulfur atoms on the cell-binding molecule.

n is 1 ~ 30; R₁, R₂, m₁, m₂, X₁ and X₂ are the same described in Claim 1.

3. A compound of Formula (III):




Wherein:

$\equiv$ represents either a single bond, or a double bond.

Cb, Z₁, n, m₁, m₂, R₁, R₂, X₁, and X₂ are the same as defined in Claims 1 and 2.

4. A compound of Formula (IV):





Wherein:

==== represents either a single bond, or a double bond, or a triple bond.

U, U', Drug₁, n, m₁, m₂, R₁, R₂, X₁, and X₂ are the same as defined in Claims 1 and 2.

5. The bridge linker compound of Formula (I) of claims 1, wherein the 2,3-disubstituted succinic group, or 2-monosubstituted or 2,3-disubstituted fumaric group, or 2-monosubstituted or 2,3-disubstituted maleic group, or acetylenedicarboxyl group, is preferably synthesized through the condensation of the 2,3-disubstituted succinic acid, or 2-monosubstituted or 2,3-disubstituted fumaric acid, or 2-mono-substituted or 2,3-disubstituted maleic acid, or acetylenedicarboxyl group, or their acid derivatives, with the other components containing an amine (1° or 2° amines), alcohol, or thiol, on their terminal as shown in the following scheme (Ia) and (Ib):



Wherein X₁, X₂, R₁, and R₂ are described the same in Claim 1.

Lv₁ and Lv₂ are the same or independently H, OH, F, Cl, Br, I, NH₂, NHNH₂, methanesulfonyl (mesyl), toluenesulfonyl (tosyl), trifluoromethylsulfonyl (triflate), trifluoromethylsulfonate, nitrophenoxyl, N-succinimidylloxyl (NHS), phenoxyl; dinitrophenoxyl; pentafluorophenoxyl, tetrafluorophenoxyl, trifluorophenoxyl, difluorophenoxyl, monofluorophenoxyl, pentachlorophenoxyl, 1H-imidazole-1-yl, chlorophenoxyl, dichlorophenoxyl, trichlorophenoxyl, tetrachlorophenoxyl, N-(benzotriazol-yl)oxyl, 2-ethyl-5-phenylisoxazolium-3'-sulfonyl, phenyl-oxadiazole-sulfonyl (-sulfone-ODA), oxadiazol-yl, or an intermediate molecule generated with a condensation reagent for Mitsunobu reactions. The condensation reagents are: EDC (N-(3-Dimethylaminopropyl)-N'-ethylcarbodiimide), DCC (Dicyclohexyl-carbodiimide), N,N'-Diisopropylcarbodiimide (DIC), N-Cyclohexyl-N'-(2-morpholinoethyl)carbodiimide metho-p-toluenesulfonate (CMC, or CME-CDI), 1,1'-Carbonyldiimidazole (CDI), TBTU (O-(Benzotria-

zol-1-yl)-N,N,N',N'-tetramethyluronium tetrafluoroborate), N,N,N',N'-Tetramethyl-O-(1H-benzotriazol-1-yl)uronium hexafluorophosphate (HBTU), (Benzotriazol-1-yloxy)tris(dimethylamino)phosphonium hexafluorophosphate (BOP), (Benzotriazol-1-yloxy)tripyrrolidinophosphonium hexafluorophosphate (PyBOP), Diethyl cyanophosphonate (DEPC), Chloro-N,N,N',N'-tetramethylformamidinium hexafluorophosphate, 1-[Bis(dimethylamino)-methylene]-1H-1,2,3-triazolo[4,5-b]pyridinium 3-oxid hexafluorophosphate (HATU), 1-[(Dimethylamino)(morpholino) methylene]-1H-[1,2,3]triazolo[4,5-b]pyridine-1-ium 3-oxide hexafluorophosphate (HDMA), 2-Chloro-1,3-dimethylimidazolidinium hexafluorophosphate (CIP), Chlorotripyrrolidinophosphonium hexafluorophosphate (PyCloP), Fluoro-N,N,N',N'-bis(tetramethylene)formamidinium hexafluorophosphate (BTFFH), N,N,N',N'-Tetramethyl-S-(1-oxido-2-pyridyl)thiuronium hexafluorophosphate, O-(2-Oxo-1(2H)pyridyl)-N,N,N',N'-tetramethyl-uronium tetrafluoroborate (TPTU), S-(1-Oxido-2-pyridyl)-N,N,N',N'-tetramethylthiuronium tetrafluoroborate, O-[(Ethoxycarbonyl) cyano-methylenamino]-N,N,N',N'-tetramethyluronium hexafluorophosphate (HOTU), (1-Cyano-2-ethoxy-2-oxoethylidenamino-oxy)dimethylamino-morpholinocarbenium hexafluorophosphate(COMU), O-(Benzotriazol-1-yl)-N,N,N',N'-bis-(tetramethylene) uronium hexafluorophosphate (HBPYU), N-Benzyl-N'-cyclohexylcarbodi-imide (with, or without polymer-bound), Dipyrrolidino(N-succinimidyl-oxy)carbenium hexa-fluorophosphate (HSPYU), Chlorodipyrrolidinocarbenium hexafluorophosphate (PYCIU), 2-Chloro-1,3-dimethylimidazolidinium tetrafluoroborate(CIB), (Benzotriazol-1-yloxy) dipiperidinocarbenium hexafluorophosphate (HBPipU), O-(6-Chlorobenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium tetrafluoroborate (TCTU), Bromo-tris(dimethylamino)-phosphonium hexafluorophosphate (BroP), Propylphosphonic anhydride (PPACA, T3P®), 2-Morpholinoethyl isocyanide (MEI), N,N,N',N'-Tetramethyl-O-(N-succin-imidyl)uronium hexafluorophosphate (HSTU), 2-Bromo-1-ethyl-pyridinium tetrafluoroborate (BEP), O-[(Ethoxycarbonyl)cyano-methylenamino]-N,N,N',N'-tetramethyluronium tetrafluoroborate (TOTU), 4-(4,6-Dimethoxy-1,3,5-triazin-2-yl)-4-methylmorpholinium chloride (MMTM, DMTMM), N,N,N',N'-Tetramethyl-O-(N-succinimidyl)uronium tetrafluoroborate (TSTU), O-(3,4-Dihydro-4-oxo-1,2,3-benzotriazin-3-yl)-N,N,N',N'-tetramethyluronium tetrafluoroborate (TDBTU), 1,1'-(Azodicarbonyl)dipiperidine (ADD), Di-(4-chlorobenzyl) azodicarboxylate (DCAD), Di-tert-butyl azodicarboxylate (DBAD), Diisopropyl azodicarboxylate (DIAD), Diethyl azodicarboxylate(DEAD). In addition, Lv1 and Lv2 can be an anhydride, formed by acid themselves or formed with other C₁~C₈ acid anhydrides.

6. The Formula (II) of claim 2, or the Formula (IV) of claim 4, wherein the Drug₁ is independently selected from:

(1). Chemotherapeutic agents: a). Alkylating agents: such as Nitrogen mustards: chlorambucil, chlornaphazine, cyclophosphamide, dacarbazine, estramustine, ifosfamide, mechlorethamine, mechlorethamine oxide hydrochloride, mannomustine, mitobronitol, melphalan, mitolactol, pipobroman, novembichin, phenesterine, prednimustine, thiotepa, trofosfamide, uracil mustard; CC-1065 (including its adozelesin, carzelesin and bizelesin synthetic analogues); duocarmycin (including the synthetic analogues, KW-2189 and CBI-TMI); benzodiazepine dimers (e.g., dimers of pyrrolbenzodiazepine (PBD) or tomaymycin, indolinobenzodiazepines, imidazobenzothiadiazepines, or oxazolidinobenzodiazepines); Nitrosoureas: (carmustine, lomustine, chlorozotocin, fotemustine, nimustine, ranimustine); Alkylsulphonates: (busulfan, treosulfan, improsulfan and piposulfan); Triazenes: (dacarbazine); Platinum containing compounds: (carboplatin, cisplatin, oxaliplatin); aziridines, such as benzodopa, carboquone, meturedopa, and uredopa; ethylenimines and methylamelamines including altretamine, triethylenemelamine, trietylenephosphoramide, triethylenethiophosphoramide and trimethylolmelamine]; b). Plant Alkaloids: such as Vinca alkaloids: (vincristine, vinblastine, vindesine, vinorelbine, navelbin); Taxoids: (paclitaxel, docetaxol) and their analogs, Maytansinoids (DM1, DM2, DM3, DM4, DM5, DM6, DM7, maytansine and ansamitocins) and their analogs, cryptophycins (particularly cryptophycin 1 and cryptophycin 8); epothilones, eleutherobin, discodermolide, bryostatins, dolostatins, auristatins, tubulysins, cephalostatins; pancratistatin; a sarcodictyin; spongistatin; c). DNA Topoisomerase Inhibitors: such as [Epipodophyllins: (9-aminocamptothecin, camptothecin, crisnatol, daunomycin, etoposide, etoposide phosphate, irinotecan, mitoxantrone, novantrone, retinoic acids (retinols), teniposide, topotecan, 9-nitrocamptothecin (RFS 2000)); mitomycins: (mitomycin C)]; d). Antimetabolites: such as {[Anti-folate: DHFR inhibitors: (methotrexate, trimetrexate, denopterin, pteropterin, aminopterin (4-aminopteroic acid) or the other folic acid analogues); IMP dehydrogenase Inhibitors: (mycophenolic acid, tiazofurin, ribavirin, EICAR); Ribonucleotide reductase Inhibitors: (hydroxyurea, deferoxamine)]; [Pyrimidine analogs: Uracil analogs: (ancitabine, azacitidine, 6-azauridine, capecitabine (Xeloda), carmofur, cytarabine, dideoxyuridine, doxifluridine, encitabine, 5-Fluorouracil, floxuridine, ratitrexed (Tomudex)); Cytosine analogs: (cytarabine, cytosine arabinoside, fludarabine); Purine analogs: (azathioprine, fludarabine, mercaptopurine, thiamiprine, thioguanine)]; folic acid replenisher, such as frolinic acid}; e). Hormonal therapies: such as {Receptor antagonists: [Anti-estrogen: (megestrol, raloxifene, tamoxifen); LHRH agonists: (goserelin, leuprolide acetate); Anti-androgens: (bicalutamide, flutamide, calusterone, dromostanolone propionate, epitioestanol, goserelin, leuprolide, mepitioestane, nilutamide, testosterone, trilostane and other androgens inhibitors)]; Retinoids/Deltoids: [Vitamin D3 analogs:

(CB 1093, EB 1089 KH 1060, cholecalciferol, ergocalciferol); Photodynamic therapies: (verteporfin, phthalocyanine, photosensitizer Pc4, demethoxy-hypocrellin A); Cytokines: (Interferon-alpha, Interferon-gamma, tumor necrosis factor (TNFs), human proteins containing a TNF domain)]; f). Kinase inhibitors, such as BIBW 2992 (anti-EGFR/Erb2), imatinib, gefitinib, pegaptanib, sorafenib, dasatinib, sunitinib, erlotinib, nilotinib, lapatinib, axitinib, pazopanib, vandetanib, E7080 (anti-VEGFR2), mubritinib, ponatinib (AP24534), bafetinib (INNO-406), bosutinib (SKI-606), cabozantinib, vismodegib, iniparib, ruxolitinib, CYT387, axitinib, tivozanib, sorafenib, bevacizumab, cetuximab, Trastuzumab, Ranibizumab, Panitumumab, ispinesib; g). antibiotics, such as the enediyne antibiotics (e.g. calicheamicins, especially calicheamicin γ_1 , δ_1 , α_1 and β_1 ; dynemicin, including dynemicin A and deoxydynemicin; esperamicin, kedarcidin, C-1027, maduropeptin, as well as neocarzinostatin chromophore and related chromoprotein enediyne antibiotic chromomorphores), aclacinomysins, actinomycin, anthramycin, azaserine, bleomycins, cactinomycin, carabacin, carminomycin, carzinophilin; chromomycins, dactinomycin, daunorubicin, detorubicin, 6-diazo-5-oxo-L-norleucine, doxorubicin, morpholino-doxorubicin, cyanomorpholino-doxorubicin, 2-pyrrolino-doxorubicin and deoxydoxorubicin, epirubicin, esorubicin, idarubicin, marcellomycin, nitomycins, mycophe-nolic acid, nogalamycin, olivomycins, peplomycin, potfiromycin, puromycin, quelamycin, rodorubicin, streptonigrin, streptozocin, tubercidin, ubenimex, zinostatin, zorubicin; f). Others: such as Polyketides (acetogenins), especially bullatacin and bullatacinone; gemcitabine, epoxomicins (e. g. carfilzomib), bortezomib, thalidomide, lenalidomide, pomalidomide, tosedostat, zybrestat, PLX4032, STA-9090, Stimuvax, allovectin-7, Xegeva, Provenge, Yervoy, Isoprenylation inhibitors (such as Lovastatin), Dopaminergic neurotoxins (such as 1-methyl-4-phenylpyridinium ion), Cell cycle inhibitors (such as staurosporine), Actinomycins (such as Actinomycin D, dactinomycin), amanitins, Bleomycins (such as bleomycin A2, bleomycin B2, peplomycin), Anthracyclines (such as daunorubicin, doxorubicin (adriamycin), idarubicin, epirubicin, pirarubicin, zorubicin, mitoxantrone, MDR inhibitors (such as verapamil), Ca^{2+} ATPase inhibitors (such as thapsigargin), Histone deacetylase inhibitors (Vorinostat, Romidepsin, Panobinostat, Valproic acid, Mocetinostat (MGCD0103), Belinostat, PCI-24781, Entinostat, SB939, Resminostat, Givinostat, AR-42, CUDC-101, sulforaphane, Trichostatin A) ; Thapsigargin, Celecoxib, glitazones, epigallocatechin gallate, Disulfiram, Salinosporamide A.; Anti-adrenals, such as aminoglutethimide, mitotane, trilostane; aceglatone; aldophosphamide glycoside; aminolevulinic acid; amsacrine; arabinoside, bestabucil; bisantrene; edatraxate; defofamine; demecolcine; diaziquone; eflornithine (DFMO), elfomithine; elliptinium acetate, etoglucid; gallium nitrate; gacytosine, hydroxyurea; ibandronate, lentinan; lonidamine; mitoguazone; mitoxantrone; mop-

idamol; nitracrine; pentostatin; phenamet; pirarubicin; podophyllinic acid; 2-ethylhydrazide; procarbazine; PSK[®]; razoxane; rhizoxin; sizofiran; spirogermanium; tenuazonic acid; triaziquone; 2, 2',2"-trichlorotriethylamine; trichothecenes (especially T-2 toxin, verrucarins A, roridin A and anguidine); urethane, siRNA, antisense drugs;

(2). Anti-autoimmune disease agents: cyclosporine, cyclosporine A, aminocaproic acid, azathioprine, bromocriptine, chlorambucil, chloroquine, cyclophosphamide, corticosteroids (e.g. amcinonide, betamethasone, budesonide, hydrocortisone, flunisolide, fluticasone propionate, fluocortolone danazol, dexamethasone, Triamcinolone acetonide, beclometasone dipropionate), DHEA, enanercept, hydroxychloroquine, infliximab, meloxicam, methotrexate, mofetil, mycophenolate, prednisone, sirolimus, tacrolimus.

(3). Anti-infectious disease agents, a). Aminoglycosides: amikacin, astromycin, gentamicin (netilmicin, sisomicin, isepamicin), hygromycin B, kanamycin (amikacin, arbekacin, bekana-mycin, dibekacin, tobramycin), neomycin (framycetin, paromomycin, ribostamycin), netilmicin, spectinomycin, streptomycin, tobramycin, verdamicin; b). Amphenicols: azidamfenicol, chloramphenicol, florfenicol, thiamphenicol; c). Ansamycins: geldanamycin, herbimycin; d). Carbapenems: biapenem, doripenem, ertapenem, imipenem/cilastatin, meropenem, panipenem; e). Cepheems: carbacephem (loracarbef), cefacetrile, cefaclor, cefradine, cefadroxil, cefalonium, cefaloridine, cefalotin or cefalothin, cefalexin, cefaloglycin, cefamandole, cefapirin, cefatrizine, cefazaflur, cefazedone, cefazolin, cefbuperazone, cefcapene, cefdaloxime, cefepime, cefminox, cefoxitin, cefprozil, cefroxadine, ceftazidime, cefuroxime, cefixime, cefdinir, cefditoren, cefepime, cefetamet, cefmenoxime, cefodizime, cefonicid, cefoperazone, ceforanide, cefotaxime, cefotiam, ceftazidime, cefepime, cefpimizole, cefpiramide, cefpirome, cefpodoxime, cefprozil, cefquinome, cefsulodin, ceftazidime, ceftazidime, ceftibuten, ceftiole, ceftizoxime, ceftibiprole, ceftriaxone, cefuroxime, cefuzonam, *cephamycin* (cefoxitin, cefotetan, cefmetazole), oxacephem (flomoxef, latamoxef); f). Glycopeptides: bleomycin, vancomycin (oritavancin, telavancin), teicoplanin (dalbavancin), ramoplanin; g). Glycylcyclines: e. g. tigecycline; g). β -Lactamase inhibitors: penam (sulbactam, tazobactam), clavam (clavulanic acid); i). Lincosamides: clindamycin, lincomycin; j). Lipopeptides: daptomycin, A54145, calcium-dependent antibiotics (CDA); k). Macrolides: azithromycin, cethromycin, clarithromycin, dirithromycin, erythromycin, flurithromycin, josamycin, ketolide (telithromycin, cethromycin), midecamycin, miocamycin, oleandomycin, rifamycins (rifampicin, rifampin, rifabutin, rifapentine), rokitamycin, roxithromycin, spectinomycin, spiramycin, tacrolimus (FK506), troleandomycin, telithromycin; l). Monobactams: aztreonam, tigemonam; m). Oxazolidinones: linezolid; n). Penicillins: amoxicillin, ampicillin (pivampicillin, hetacillin, bacampicillin, metampicillin, talampicil-

lin), azidocillin, azlocillin, benzylpenicillin, benzathine benzylpenicillin, benzathine phenoxymethylpenicillin, clometocillin, procaine benzylpenicillin, carbenicillin (carindacillin), cloxacillin, dicloxacillin, epicillin, flucloxacillin, mecillinam (pivmecillinam), mezlocillin, meticillin, nafcillin, oxacillin, penamecillin, penicillin, pheneticillin, phenoxymethylpenicillin, piperacillin, propicillin, sulbenicillin, temocillin, ticarcillin; o). Polypeptides: bacitracin, colistin, polymyxin B; p). Quinolones: alatrofloxacin, balofloxacin, ciprofloxacin, clinafloxacin, danofloxacin, difloxacin, enoxacin, enrofloxacin, floxin, garenoxacin, gatifloxacin, gemifloxacin, grepafloxacin, kano trovafloxacin, levofloxacin, lomefloxacin, marbofloxacin, moxifloxacin, nadifloxacin, norfloxacin, orbifloxacin, ofloxacin, pefloxacin, trovafloxacin, grepafloxacin, sitafloxacin, sparfloxacin, temafloxacin, tosufloxacin, trovafloxacin; q). Streptogramins: pristnamycin, quinupristin/dalfopristin; r). Sulfonamides: mafenide, prontosil, sulfacetamide, sulfamethizole, sulfanilimide, sulfasalazine, sulfisoxazole, trimethoprim, trimethoprim-sulfamethoxazole (co-trimoxazole); s). Steroid antibacterials: e.g. fusidic acid; t). Tetracyclines: doxycycline, chlortetracycline, clomocycline, demeclocycline, lymecycline, meclocycline, metacycline, minocycline, oxytetracycline, penimepicycline, rolitetracycline, tetracycline, glycylicyclines (e.g. tigecycline); u). Other types of antibiotics: annonacin, arspenamine, bacitrenol inhibitors (Bacitracin), DADAL/AR inhibitors (cycloserine), dictyostatin, discodermolide, eleutherobin, epothilone, ethambutol, etoposide, faropenem, fusidic acid, furazolidone, isoniazid, laulimalide, metronidazole, mupirocin, mycolactone, NAM synthesis inhibitors (e.g. fosfomycin), nitrofurantoin, paclitaxel, platensimycin, pyrazinamide, quinupristin/dalfopristin, rifampicin (rifampin), tazobactam tinidazole, uvaricin;

(4). Anti-viral drugs: a). Entry/fusion inhibitors: aplaviroc, maraviroc, vicriviroc, *gp41* (enfuvirtide), PRO 140, CD4 (ibalizumab); b). Integrase inhibitors: raltegravir, elvitegravir, globoindan A; c). Maturation inhibitors: bevirimat, vivecon; d). Neuraminidase inhibitors: oseltamivir, zanamivir, peramivir; e). Nucleosides & nucleotides: abacavir, aciclovir, adefovir, amdoxovir, apricitabine, brivudine, cidofovir, clevudine, dexelvucitabine, didanosine (ddI), elvucitabine, emtricitabine (FTC), entecavir, famciclovir, fluorouracil (5-FU), 3'-fluoro-substituted 2', 3'-dideoxynucleoside analogues (e.g. 3'-fluoro-2',3'-dideoxythymidine (FLT) and 3'-fluoro-2',3'-dideoxyguanosine (FLG), fomivirsen, ganciclovir, idoxuridine, lamivudine (3TC), l-nucleosides (e.g. β -l-thymidine and β -l-2'-deoxycytidine), penciclovir, racivir, ribavirin, stampidine, stavudine (d4T), taribavirin (viramidine), telbivudine, tenofovir, trifluridine valaciclovir, valganciclovir, zalcitabine (ddC), zidovudine (AZT); f). Non-nucleosides: amantadine, ateviridine, capravirine, diarylpyrimidines (etravirine, rilpivirine), delavirdine, docosanol, emivirine, efavirenz, foscarnet (phosphonoformic acid), imiquimod, interferon alfa,

loviride, lodenosine, methisazone, nevirapine, NOV-205, peginterferon alfa, podophyllotoxin, rifampicin, rimantadine, resiquimod (R-848), tromantadine; g). Protease inhibitors: amprenavir, atazanavir, boceprevir, darunavir, fosamprenavir, indinavir, lopinavir, nelfinavir, pleconaril, ritonavir, saquinavir, telaprevir (VX-950), tipranavir; h). Other types of anti-virus drugs: abzyme, arbidol, calanolide a, ceragenin, cyanovirin-n, diarylpyrimidines, epigallocatechin gallate (EGCG), foscarnet, griffithsin, taribavirin (viramidine), hydroxyurea, KP-1461, miltefosine, pleconaril, portmanteau inhibitors, ribavirin, seliciclib.

(5). A radioisotope that can be selected from (radionuclides) ^3H , ^{11}C , ^{14}C , ^{18}F , ^{32}P , ^{35}S , ^{64}Cu , ^{68}Ga , ^{86}Y , ^{99}Tc , ^{111}In , ^{123}I , ^{124}I , ^{125}I , ^{131}I , ^{133}Xe , ^{177}Lu , ^{211}At , or ^{213}Bi .

(6). A chromophore molecule, which can be one that has the ability to absorb a kind of light, such as UV light, florescent light, IR light, near IR light, visual light; A class or subclass of xanthophores, erythrophores, iridophores, leucophores, melanophores, cyanophores, fluorophore molecules which are fluorescent chemical compounds re-emitting light upon light, visual phototransduction molecules, photophore molecules, luminescence molecules, luciferin compounds; Non-protein organic fluorophores, such as: Xanthene derivatives (fluorescein, rhodamine, Oregon green, eosin, and Texas red); Cyanine derivatives: (cyanine, indocarbocyanine, oxacarbocyanine, thiocarbocyanine, and merocyanine); Squaraine derivatives and ring-substituted squaraines, including Seta, SeTau, and Square dyes; Naphthalene derivatives (dansyl and prodan derivatives); Coumarin derivatives; Oxadiazole derivatives (pyridyloxazole, nitrobenzoxadiazole and benzoxadiazole); Anthracene derivatives (anthraquinones, including DRAQ5, DRAQ7 and CyTRAK Orange); Pyrene derivatives (cascade blue, etc); Oxazine derivatives (Nile red, Nile blue, cresyl violet, oxazine 170 etc). Acridine derivatives (proflavin, acridine orange, acridine yellow etc). Arylmethine derivatives (auramine, crystal violet, malachite green). Tetrapyrrole derivatives (porphin, phthalocyanine, bilirubin); Any analogs and derivatives of the following fluorophore compounds: CF dye (Biotium), DRAQ and CyTRAK probes (BioStatus), BODIPY (Invitrogen), Alexa Fluor (Invitrogen), DyLight Fluor (Thermo Scientific, Pierce), Atto and Tracy (Sigma Aldrich), FluoProbes (Interchim), Abberior Dyes (Abberior), DY and MegaStokes Dyes (Dyomics), Sulfo Cy dyes (Cyandye), HiLyte Fluor (AnaSpec), Seta, SeTau and Square Dyes (SETA BioMedicals), Quasar and Cal Fluor dyes (Bioscience Technologies), SureLight Dyes (APC, RPEPerCP, Phycobilisomes)(Columbia Biosciences), APC, APCXL, RPE, BPE (Phyco-Biotech), Allophycocyanin (APC), Aminocoumarin, APC-Cy7 conjugates, BODIPY-FL, Cascade Blue, Cy2, Cy3, Cy3.5, Cy3B, Cy5, Cy5.5, Cy7, Fluorescein, FluorX, Hydroxycoumarin, Lissamine Rhodamine B, Lucifer yellow, Methoxycoumarin, NBD, Pacific Blue, Pacific Orange, PE-Cy5 conjugates, PE-Cy7 conjugates, PerCP,

R-Phycoerythrin(PE), Red 613, Seta-555-Azide, Seta-555-DBCO, Seta-555-NHS, Seta-580-NHS, Seta-680-NHS, Seta-780-NHS, Seta-APC-780, Seta-PerCP-680, Seta-R-PE-670, SeTau-380-NHS, SeTau-405-Maleimide, SeTau-405-NHS, SeTau-425-NHS, SeTau-647-NHS, Texas Red, TRITC, TruRed, X-Rhodamine, 7-AAD (7-aminoactinomycin D, CG-selective), Acridine Orange, Chromomycin A3, CyTRAK Orange (Biostatus, red excitation dark), DAPI, DRAQ5, DRAQ7, Ethidium Bromide, Hoechst33258, Hoechst33342, LDS 751, Mithramycin, PropidiumIodide (PI), SYTOX Blue, SYTOX Green, SYTOX Orange, Thiazole Orange, TO-PRO: Cyanine Monomer, TOTO-1, TO-PRO-1, TOTO-3, TO-PRO-3, YOSeta-1, YOYO-1. The fluorophore compounds that can be linked to the linkers of the invention for study cells are selected from the following compounds or their derivatives: DCFH (2'7'Dichlorodihydro-fluorescein, oxidized form), DHR (Dihydrorhodamine 123, oxidized form, light catalyzes oxidation), Fluo-3 (AM ester. pH > 6), Fluo-4 (AM ester. pH 7.2), Indo-1 (AM ester, low/high calcium (Ca²⁺)), SNARF(pH 6/9), Allophycocyanin(APC), AmCyan1 (tetramer, Clontech), AsRed2 (tetramer, Clontech), Azami Green (monomer, MBL), Azurite, B-phycoerythrin(BPE), Cerulean, CyPet, DsRed monomer (Clontech), DsRed2 ("RFP", Clontech), EBFP, EBFP2, ECFP, EGFP (weak dimer, Clontech), Emerald (weak dimer, Invitrogen), EYFP (weak dimer, Clontech), GFP (S65A mutation), GFP (S65C mutation), GFP (S65L mutation), GFP (S65T mutation), GFP (Y66F mutation), GFP (Y66H mutation), GFP (Y66W mutation), GFPuv, HcRed1, J-Red, Katusha, Kusabira Orange (monomer, MBL), mCFP, mCherry, mCitrine, Midoriishi Cyan (dimer, MBL), mKate (TagFP635, monomer, Evrogen), mKeima-Red (monomer, MBL), mKO, mOrange, mPlum, mRaspberry, mRFP1 (monomer, Tsien lab), mStrawberry, mTFP1, mTurquoise2, P3 (phycobilisome complex), Peridinin Chlorophyll (PerCP), R-phycoerythrin(RPE), T-Sapphire, TagCFP (dimer, Evrogen), TagGFP (dimer, Evrogen), TagRFP (dimer, Evrogen), TagYFP (dimer, Evrogen), tdTomato (tandem dimer), Topaz, TurboFP602 (dimer, Evrogen), TurboFP635 (dimer, Evrogen), TurboGFP (dimer, Evrogen), TurboRFP (dimer, Evrogen), TurboYFP (dimer, Evrogen), Venus, Wild Type GFP, YPet, ZsGreen1 (tetramer, Clontech), ZsYellow1 (tetramer, Clontech).

(7). The cell-binding ligands or receptor agonists, which can be selected from: Folate derivatives; Glutamic acid urea derivatives; Somatostatin and its analogs (such as octreotide (Sandostatin) and lanreotide (Somatuline)); Aromatic sulfonamides; Pituitary adenylate cyclase activating peptides (PACAP) (PAC1); Vasoactive intestinal peptides (VIP/PACAP) (VPAC1, VPAC2); Melanocyte-stimulating hormones (α -MSH); Cholecystokinins (CCK) /gastrin receptor agonists; Bombesins (such as Pyr-Gln-Arg-Leu-Gly-Asn-Gln-Trp-Ala-Val-Gly-His-Leu-Met-NH₂)/gastrin-releasing peptide (GRP); Neurotensin receptor ligands (NTR1, NTR2, NTR3);

Substance P (NK1 receptor) ligands; Neuropeptide Y (Y1–Y6); Homing Peptides include RGD (Arg-Gly-Asp), NGR (Asn-Gly-Arg), the dimeric and multimeric cyclic RGD peptides (e.g. cRGDfV), TAASGVSRMH and LTLRWVGLMS (Chondroitin sulfate proteoglycan NG2 receptor ligands) and F3 peptides; Cell Penetrating Peptides (CPPs); Peptide Hormones, such as luteinizing hormone-releasing hormone (LHRH) agonists and antagonists, and gonadotropin-releasing hormone (GnRH) agonist, acts by targeting follicle stimulating hormone (FSH) and luteinising hormone (LH), as well as testosterone production, e.g. buserelin (Pyr-His-Trp-Ser-Tyr-D-Ser(OtBu)-Leu-Arg-Pro-NHEt), Gonadorelin (Pyr-His-Trp-Ser-Tyr-Gly-Leu-Arg-Pro-Gly-NH₂), Goserelin (Pyr-His-Trp-Ser-Tyr-D-Ser(OtBu)-Leu-Arg-Pro-AzGly-NH₂), Histrelin (Pyr-His-Trp-Ser-Tyr-D-His(N-benzyl)-Leu-Arg-Pro-NHEt), leuprolide (Pyr-His-Trp-Ser-Tyr-D-Leu-Leu-Arg-Pro-NHEt), Nafarelin (Pyr-His-Trp-Ser-Tyr-2Nal-Leu-Arg-Pro-Gly-NH₂), Triptorelin (Pyr-His-Trp-Ser-Tyr-D-Trp-Leu-Arg-Pro-Gly-NH₂), Nafarelin, Deslorelin, Abarelix (Ac-D-2Nal-D-4-chloroPhe-D-3-(3-pyridyl)Ala-Ser-(N-Me)Tyr-D-Asn-Leu-isopropylLys-Pro-DAla-NH₂), Cetrorelix (Ac-D-2Nal-D-4-chloroPhe-D-3-(3-pyridyl)Ala-Ser-Tyr-D-Cit-Leu-Arg-Pro-D-Ala-NH₂), Degarelix (Ac-D-2Nal-D-4-chloroPhe-D-3-(3-pyridyl)Ala-Ser-4-aminoPhe(L-hydroxyrotyl)-D-4-aminoPhe(carbamoyl)-Leu-isopropylLys-Pro-D-Ala-NH₂), and Ganirelix (Ac-D-2Nal-D-4-chloroPhe-D-3-(3-pyridyl)Ala-Ser-Tyr-D-(N₉, N₁₀-diethyl)-homoArg-Leu-(N₉, N₁₀-diethyl)-homoArg-Pro-D-Ala-NH₂); Pattern Recognition Receptor (PRRs), such as Toll-like receptors' (TLRs) ligands, C-type lectins and Nodlike Receptors' (NLRs) ligands; Calcitonin receptor agonists; integrin receptors' and their receptor subtypes' (such as $\alpha_v\beta_1$, $\alpha_v\beta_3$, $\alpha_v\beta_5$, $\alpha_v\beta_6$, $\alpha_6\beta_4$, $\alpha_7\beta_1$, $\alpha_L\beta_2$, $\alpha_{IIb}\beta_3$) agonists (such as GRGDSPK, cyclo(RGDfV) (L1) and its derivatives [cyclo(-N(Me)R-GDfV), cyclo(R-Sar-DfV), cyclo(RG-N(Me)D-fV), cyclo(RGD-N(Me)f-V), cyclo(RGDf-N(Me)V-)(Cilengitide)]; Nanobody (a derivative of VHH (camelid Ig)); Domain antibodies (dAb, a derivative of VH or VL domain); Bispecific T cell Engager (BiTE, a bispecific diabody); Dual Affinity ReTargeting (DART, a bispecific diabody); Tetravalent tandem antibodies (TandAb, a dimerized bispecific diabody); Anticalin (a derivative of Lipocalins); Adnectins (10th FN3 (Fibronectin)); Designed Ankyrin Repeat Proteins (DARPs); Avimers; EGF receptors and VEGF receptors' agonists.

(8). The pharmaceutically acceptable salts, acids or derivatives of any of the above drugs.

7. The compounds of the Formula (II) of claim 2, or the Formula (IV) of claim 4, wherein "Drug₁" is a chromophore molecule, thus the conjugate compounds of the Formula (II) of claim 2, or the conjugate compounds of the Formula (IV) of claim 4, can be used for detecting, monitoring, or studying the interactions and/or functions of the cell binding molecule, and/or the interactions of the conjugate with a target, particularly, a targeted cell.

- 8.** The compounds of the Formula (II) of claim 2, or the Formula (IV) of claim 4, wherein “Drug₁” and/or “R₂” can be independently a polyalkylene glycols or a polyalkylene glycol analogs, which are used for extending the half-life of the cell-binding molecule when administered to a mammal. Polyalkylene glycols are poly(ethylene glycols) (PEGs), poly(propylene glycol), copolymers of ethylene oxides or propylene oxides that has a molecular weight of from about 44 Daltons to about 300 kDa.
- 9.** The compounds of the Formula (II) of claim 2, or the Formula (IV) of claim 4, wherein “Drug₁” can be a cell-binding ligand or a cell receptor agonist, or a cell receptor binding molecule, so that the conjugates can work as a targeting conductor/director to deliver the conjugate to malignant cells, and also can modulate or co-stimulate a desired immune response or alter signaling pathways.
- 10.** The conjugate compound of claim 2 or 4, wherein “Drug₁” is more preferred selected from tubulysins, calicheamicins, auristatins, maytansinoids, CC-1065 analogs, daunorubicin and doxorubicin compounds, taxanoids (taxanes), cryptophycins, epothilones, benzodiazepine dimers (e.g., dimmers of pyrrolbenzodiazepine (PBD), tomaymycins, anthramycins, indolinobenzodiazepines, imidazobenzothiadiazepines, or oxazolidinobenzodiazepines), calicheamicins and the enediyne antibiotics, actinomycins, amanitins, azaserines, bleomycins, epirubicins, tamoxifen, idarubicin, dolastatins/auristatins (e.g. monomethyl auristatin E, MMAE, MMAF, auristatin PYE, auristatin TP, Auristatins 2-AQ, 6-AQ, EB (AEB), and EFP (AEFP)), duocarmycins, geldanamycins, methotrexates, thiotepa, vindesines, vincristines, hemiasterlins, nazumamides, microginins, radiosumins, alterobactins, microsclerodermins, theonellamides, esperamicins, siRNA, nucleolytic enzymes, and/or pharmaceutically acceptable salts, acids, or/and their analogues derivatives of any of the above molecules.
- 11.** The conjugate compounds of claim 2 or 3 wherein the cell binding agent/molecule, Cb, is selected from an antibody, a protein, a vitamin (e.g. folate), peptides, a polymeric micelle, a liposome, a lipoprotein-based drug carrier, a nano-particle drug carrier, a dendrimer, and a molecule said above coating with cell-binding ligands, or a combination of said above thereof.
- 12.** The cell-binding molecule/agent, Cb, according to claim 2, 3, or 11 is preferred an antibody, full-length antibodies (polyclonal antibodies, monoclonal antibodies, dimers, multimers, multispecific antibodies (e.g., bispecific or trispecific antibodies); a single chain antibody, an antibody fragment that binds to the target cell, a monoclonal antibody, a single chain monoclonal antibody, or a monoclonal antibody fragment that binds the target cell, a chimeric antibody, a chimeric antibody fragment that binds to the target cell, a domain antibody, a domain antibody fragment that binds to the target cell, a resurfaced antibody, a resurfaced single chain antibody,

or a resurfaced antibody fragment that binds to the target cell, a humanized antibody or a resurfaced antibody, a humanized single chain antibody, or a humanized antibody fragment that binds to the target cell, anti-idiotypic (anti-Id) antibodies, CDR's, diabody, triabody, tetrabody, miniantibody, small immune proteins (SIP), a lymphokine, a hormone, a vitamin, a growth factor, a colony stimulating factor, a nutrient-transport molecule, large molecular weight proteins, nanoparticles or polymers modified with antibodies or large molecular weight proteins.

13. The cell-binding molecule/agent, Cb, according to claim **2, 3, 11** or **12** can be any agent that is able to target against a tumor cell, a virus infected cell, a microorganism infected cell, a parasite infected cell, an autoimmune disease cell, an activated tumor cells, a myeloid cell, an activated T-cell, an affecting B cell, or a melanocyte.


14. The cell-binding molecule/agent, Cb, according to claim **2, 3, 11** or **12** can be any agent/molecule that is able to against any one of the following antigens or receptors: CD3, CD4, CD5, CD6, CD7, CD8, CD9, CD10, CD11a, CD11b, CD11c, CD12w, CD14, CD15, CD16, CDw17, CD18, CD19, CD20, CD21, CD22, CD23, CD24, CD25, CD26, CD27, CD28, CD29, CD30, CD31, CD32, CD33, CD34, CD35, CD36, CD37, CD38, CD39, CD40, CD41, CD42, CD43, CD44, CD45, CD46, CD47, CD48, CD49b, CD49c, CD51, CD52, CD53, CD54, CD55, CD56, CD58, CD59, CD61, CD62E, CD62L, CD62P, CD63, CD66, CD68, CD69, CD70, CD72, CD74, CD79, CD79a, CD79b, CD80, CD81, CD82, CD83, CD86, CD87, CD88, CD89, CD90, CD91, CD95, CD96, CD98, CD100, CD103, CD105, CD106, CD109, CD117, CD120, CD125, CD126, CD127, CD133, CD134, CD135, CD137, CD138, CD141, CD142, CD143, CD144, CD147, CD151, CD147, CD152, CD154, CD156, CD158, CD163, CD166, .CD168, CD174, CD180, CD184, CDw186, CD194, CD195, CD200, CD200a, CD200b, CD209, CD221, CD227, CD235a, CD240, CD262, CD271, CD274, CD276 (B7-H3), CD303, CD304, CD309, CD326, 4-1BB, 5AC, 5T4 (Trophoblast glycoprotein, TPBG, 5T4, Wnt-Activated Inhibitory Factor 1 or WAIIF1), Adenocarcinoma antigen, AGS-5, AGS-22M6, Activin receptor-like kinase 1, AFP, AKAP-4, ALK, Alpha intergrin, Alpha v beta6, Amino-peptidase N, Amyloid beta, Androgen receptor, Angiopoietin 2, Angiopoietin 3, Annexin A1, Anthrax toxin protective antigen, Anti-transferrin receptor, AOC3 (VAP-1), B7-H3, Bacillus anthracis anthrax, BAFF (B-cell activating factor), BCMA, B-lymphoma cell, bcr-abl, Bombesin, BORIS, C5, C242 antigen, CA125 (carbohydrate antigen 125, MUC16), CA-IX (or CAIX, carbonic anhydrase 9), CALLA, CanAg, Canis lupus familiaris IL31, Carbonic anhydrase IX, Cardiac myosin, CCL11(C-C motif chemokine 11), CCR4 (C-C chemokine receptor type 4, CD194), CCR5, CD3E (epsilon), CEA (Carcinoembryonic antigen), CEACAM3, CEACAM5 (carcinoembryonic antigen), CFD (Factor D), Ch4D5, Cholecystokinin 2 (CCK2R), CLDN18 (Claudin-

18), Clumping factor A, cMet, CRIPTO, FCSF1R (Colony stimulating factor 1 receptor, CD115), CSF2 (colony stimulating factor 2, Granulocyte-macrophage colony-stimulating factor (GM-CSF)), CSP4, CTLA4 (cytotoxic T-lymphocyte-associated protein 4), CTAA16.88 tumor antigen, CXCR4 (CD184), C-X-C chemokine receptor type 4, cyclic ADP ribose hydrolase, Cyclin B1, CYP1B1, Cytomegalovirus, Cytomegalovirus glycoprotein B, Dabigatran, DLL4 (delta-like-ligand 4), DPP4 (Dipeptidyl-peptidase 4), DR5 (Death receptor 5), E. coli shiga toxin type-1, E. coli shiga toxin type-2, ED-B, EGFL7 (EGF-like domain-containing protein 7), EGFR, EGFRvIII, Endoglin (CD105), Endothelin B receptor, Endotoxin, EpCAM (epithelial cell adhesion molecule), EphA2, Episialin, ERBB2 (Epidermal Growth Factor Receptor 2), ERBB3, ERG (TMPRSS2 ETS fusion gene), Escherichia coli, ETV6-AML, FAP (Fibroblast activation protein alpha), FCGR1, alpha-Fetoprotein, Fibrin II, beta chain, Fibronectin extra domain-B, FOLR (folate receptor), Folate receptor alpha, Folate hydrolase, Fc-related antigen 1F protein of respiratory syncytial virus, Frizzled receptor, Fucosyl GM1, GD2 ganglioside, G-28 (a cell surface antigen glycolipid), GD3 idio type, GloboH, Glypican 3, N-glycolylneuraminic acid, GM3, GMCSF receptor α -chain, Growth differentiation factor 8, GP100, GPNMB (Trans-membrane glycoprotein NMB), GUCY2C (Guanylate cyclase 2C, guanylyl cyclase C(GC-C), intestinal Guanylate cyclase, Guanylate cyclase-C receptor, Heat-stable enterotoxin receptor (hSTAR)), Heat shock proteins, Hemagglutinin, Hepatitis B surface antigen, Hepatitis B virus, HER1 (human epidermal growth factor receptor 1), HER2, HER2/neu, HER3 (ERBB-3), IgG4, HGF/SF (Hepatocyte growth factor/scatter factor), HHGFR, HIV-1, Histone complex, HLA-DR (human leukocyte antigen), HLA-DR10, HLA-DRB, HMWMAA, Human chorionic gonadotropin, HNGF, Human scatter factor receptor kinase, HPV E6/E7, Hsp90, hTERT, ICAM-1 (Intercellular Adhesion Molecule 1), Idiotypic, IGF1R (IGF-1, insulin-like growth factor 1 receptor), IGHE, IFN- γ , Influenza hemagglutinin, IgE, IgE Fc region, IGHE, IL-1, IL-2 receptor (interleukin 2 receptor), IL-4, IL-5, IL-6, IL-6R (interleukin 6 receptor), IL-9, IL-10, IL-12, IL-13, IL-17, IL-17A, IL-20, IL-22, IL-23, IL31RA, ILGF2 (Insulin-like growth factor 2), Integrins (α_4 , $\alpha_{1b}\beta_3$, $\alpha_v\beta_3$, $\alpha_4\beta_7$, $\alpha_5\beta_1$, $\alpha_6\beta_4$, $\alpha_7\beta_7$, $\alpha_{11}\beta_3$, $\alpha_5\beta_5$, $\alpha_v\beta_5$), Interferon gamma-induced protein, ITGA2, ITGB2, KIR2D, Kappa Ig, LCK, Le, Legumain, Lewis-Y antigen, LFA-1 (Lymphocyte function-associated antigen 1, CD11a), LHRH, LINGO-1, Lipoteichoic acid, LIV1A, LMP2, LTA, MAD-CT-1, MAD-CT-2, MAGE-1, MAGE-2, MAGE-3, MAGE A1, MAGE A3, MAGE 4, MART1, MCP-1, MIF (Macrophage migration inhibitory factor, or glycosylation-inhibiting factor (GIF)), MS4A1 (membrane-spanning 4-domains subfamily A member 1), MSLN (mesothelin), MUC1(Mucin 1, cell surface associated (MUC1) or polymorphic epithelial mucin (PEM)), MUC1-KLH, MUC16 (CA125),


MCP1(monocyte chemotactic protein 1), MelanA/MART1, ML-IAP, MPG, MS4A1 (membrane-spanning 4-domains subfamily A), MYCN, Myelin-associated glycoprotein, Myostatin, NA17, NARP-1, NCA-90 (granulocyte antigen), Nectin-4 (ASG-22ME), NGF, Neural apoptosis-regulated proteinase 1, NOGO-A, Notch receptor, Nucleolin, Neu oncogene product, NY-BR-1, NY-ESO-1, OX-40, OxLDL (Oxidized low-density lipoprotein), OY-TES1, P21, p53 nonmutant, P97, Page4, PAP, Paratope of anti-(N-glycolylneuraminic acid), PAX3, PAX5, PCSK9, PDCD1 (PD-1, Programmed cell death protein 1,CD279), PDGF-R α (Alpha-type platelet-derived growth factor receptor), PDGFR- β , PDL-1, PLAC1, PLAP-like testicular alkaline phosphatase, Platelet-derived growth factor receptor beta, Phosphate-sodium co-transporter, PMEL 17, Polysialic acid, Proteinase3 (PR1), Prostatic carcinoma, PS (Phosphatidylserine), Prostatic carcinoma cells, Pseudomonas aeruginosa, PSMA, PSA, PSCA, Rabies virus glycoprotein, RHD (Rh polypeptide 1 (RhPI), CD240), Rhesus factor, RANKL, RhoC, Ras mutant, RGS5, ROBO4, Respiratory syncytial virus, RON, ROR1, Sarcoma translocation breakpoints, SART3, Sclerostin, SLAMF7 (SLAM family member 7), Selectin P, SDC1 (Syndecan 1), sLe(a), Somatomedin C, SIP (Sphingosine-1-phosphate), Somatostatin, Sperm protein 17, SSX2, STEAP1 (six-transmembrane epithelial antigen of the prostate 1), STEAP2, STn, TAG-72 (tumor associated glycoprotein 72), Survivin, T-cell receptor, T cell transmembrane protein, TEM1 (Tumor endothelial marker 1), TENB2, Tenascin C (TN-C), TGF- α , TGF- β (Transforming growth factor beta), TGF- β 1, TGF- β 2 (Transforming growth factor-beta 2), Tie (CD202b), Tie2, TIM-1 (CDX-014), Tn, TNF, TNF- α , TNFRSF8, TNFRSF10B (tumor necrosis factor receptor superfamily member 10B), TNFRSF-13B (tumor necrosis factor receptor superfamily member 13B), TPBG (trophoblast glycoprotein), TRAIL-R1 (Tumor necrosis apoptosis Inducing ligand Receptor 1), TRAILR2 (Death receptor 5 (DR5)), tumor-associated calcium signal transducer 2, tumor specific glycosylation of MUC1, TWEAK receptor, TYRP1(glycoprotein 75), TRP-2, Tyrosinase, VCAM-1 (CD106), VEGF, VEGF-A, VEGF-2 (CD309), VEGFR-1, VEGFR2, or vimentin, WT1, XAGE 1, or cells expressing any insulin growth factor receptors, or any epidermal growth factor receptors.

15. The tumor cell according to claim **13** is selected from lymphoma cells, myeloma cells, renal cells, breast cancer cells, prostate cancer cells, ovarian cancer cells, colorectal cancer cells, gastric cancer cells, squamous cancer cells, small-cell lung cancer cells, none small-cell lung cancer cells, testicular cancer cells, malignant cells, or any cells that grow and divide at an unregulated, quickened pace to cause a cancer.


16. The linkage components R₁ or R₂ according to claim **1,2, 3, or 4**, having independently one or more, or repeating, or combining thereof, components depicted in the following:




6-maleimidocaproyl (MC),




maleimido propanoyl (MP),



valine-citrulline (val-cit),




alanine-phenylalanine (ala-phe),




lysine-phenylalanine (lys-phe),

sine-phenylalanine (lys-phe),



p-aminobenzyloxycarbonyl




(PAB),

4-thio-pentanoate (SPP),




4-thio-butyrate




(SPDB),

4-(N-maleimidomethyl)cyclo-hexane-1-carboxylate




(MCC),

maleimidoethyl (ME),




4-thio-2-sulfo-butyrate


hydroxysulfonyl-butyrate (2-Sulfo-SPDB),




aryl-thiol (PySS),



(4-acetyl)aminobenzoate (SIAB),




oxylbenzylthio,




zylthio,


aminobenzylthio,




dioxylbenzylthio,






diaminobenzylthio,





amino-oxylbenzylthio,



17. The conjugate compound of claim 2, wherein "Drug₁" is a chromophore molecule, the conjugate compound of Formula (II) is preferred structures of Ac01, Ac02, Ac03, Ac04 and Ac05 as following:








Ac03 (FNIR-774 conjugate)



Ac05 (IR800CW conjugate)

Wherein “ --- ” represents either single bond or double bond; mAb is antibody, preferably monoclonal antibody; n, m₁, m₂, X₁, X₂, R₁, R₂ and R₃ are the same defined in Claims 1.

18. The conjugate compound of claim 2, wherein “Drug₁” is a Tubulysin analog, the conjugate compound of Formula (II) is preferred structures of T01, T02, T03, T04, T05, T06 and T07 as following:





19. The conjugate compound of claim 2, wherein “Drug₁” is a Calicheamicin analog, the conjugate compound of Formula (II) is preferred structures of C01 as following:






20. The conjugate compound of claim 2, wherein “Drug₁” is a Maytansinoid analog, the conjugate compound of Formula (II) is preferred structures of My01 as following:




Wherein mAb is an antibody; n, m₁, m₂, “ $\equiv$ ” represents either single bond or double bond; n, m₁, m₂, “—”, X₁, X₂, R₁, and R₂ are the same defined in Claims 1.

21. The conjugate compound of claim 2, wherein “Drug₁” is a Taxane analog, the conjugate compound of Formula (II) is preferred structures of Tx01, Tx02 and Tx03 as following:



Wherein mAb is an antibody; n, m₁, m₂, “ $\equiv$ ” represents either single bond or double bond; n, m₁, m₂, “—”, X₁, X₂, R₁, and R₂ are the same defined in Claims 1.


22. The conjugate compound of claim 2, wherein “Drug₁” is a CC-1065 analogue and/or doxorubicin analog, the conjugate compound of Formula (II) is preferred structures of CC01, CC02, and CC03 as following:










23. The conjugate compound of claim 2, wherein “Drug₁” is a Daunorubicin or Doxorubicin analogue, the conjugate compound of Formula (II) is preferred structures of Da01, Da02, Da03 and Da04 as following:










Wherein mAb is an antibody; X_3 is O, NH, NHC(O) , NHC(O)NH , C(O) , R_1 , OC(O) , or absent; “—” represents either single bond or double bond; n , m_1 , m_2 , “—”, X_1 , X_2 , R_1 , and R_2 are the same defined in Claims 1. In addition, R_1 can be absent and R_2 can be H.







24. The conjugate compound of claim 2, wherein “Drug₁” is an Auristatin and dolastatin analogue, the conjugate compound of Formula (II) is preferred structures of Au01, Au02, Au03, Au04, and Au05 as following:



Wherein mAb is an antibody; X_3 is CH_2 , O, NH, $NHC(O)$, $NHC(O)NH$, $C(O)$, $OC(O)$ R_1 , or absent; X_4 is CH_2 , $C(O)$, $C(O)NH$, $C(O)N(R_1)$, R_1 , NHR_1 , NR_1 , $C(O)R_1$ or $C(O)O$; Z_3 is H, R_1 , $OP(O)(OM_1)(OM_2)$, NHR_1 , $OCH_2OP(O)(OM_1)(OM_2)$, OSO_3M_1 , or O-glycoside (glucoside, galactoside, mannoside, glucuronoside, alloside, fructoside), NH-glycoside, S-glycoside, or CH_2 -glycoside; M_1 and M_2 are independently H, Na, K, Ca, Mg, NH_4 , $NR_1R_2R_3$; “ $\equiv$ ” represents either single bond or double bond; n, m_1 , m_2 , “—”, X_1 , X_2 , R_1 , R_2 and R_3 are the same defined in Claims 1. In addition, R_1 can be absent and R_2 can be H.

25. The conjugate compound of claim 2, wherein “Drug₁” is a benzodiazepine dimer analogue, the conjugate compound of Formula (II) is preferred structures of PB01, PB02, PB03, PB04, PB05, PB06, PB07, PB08, PB09, PB10 and PB11.










Wherein mAb is an antibody; X₃ is CH₂, O, NH, NHC(O), NHC(O)NH, C(O), OC(O), OC(O)(NR₃), R₁, NHR₁, NR₁, C(O)R₁ or absent; X₄ is CH₂, C(O), C(O)NH, C(O)N(R₁), R₁, NHR₁, NR₁, C(O)R₁ or C(O)O; M₁ and M₂ are independently H, Na, K, Ca, Mg, NH₄, NR₁R₂R₃;

“ --- ” represents either single bond or double bond; n , m_1 , m_2 , “ --- ”, X_1 , X_2 , R_1 , R_2 and R_3 are the same defined in Claims 1. In addition, R_1 can be absent and R_2 can be H.






26. The conjugate compound of claim 2, wherein “Drug₁” is an amanitin analogue, the conjugate compound of Formula (II) is preferred structures of Am01, Am02, Am03, Am04 below.















27. The conjugate compound of claim 2, wherein “Drug₁” is a polyalkylene glycol analog, the conjugate compound of Formula (II) is preferred structures of Pg01:







[illegible]













Wherein mAb is an antibody; X_3 is CH_2 , O, NH, NHC(O) , NHC(O)NH , C(O) , OC(O) , $\text{OC(O)(NR}_3\text{)}$, R_1 , NHR_1 , NR_1 , C(O)R_1 or absent; X_4 is H, CH_2 , OH, O, C(O) , C(O)NH , $\text{C(O)N(R}_1\text{)}$, R_1 , NHR_1 , NR_1 , C(O)R_1 or C(O)O ; M_1 and M_2 are independently H, Na, K, Ca, Mg, NH_4 , $\text{NR}_1\text{R}_2\text{R}_3$; R_6 = 5'-deoxyadenosyl, Me, OH, or CN; “ $\equiv$ ” represents either single bond or double bond; m_1 , m_2 , n , “—”, X_1 , X_2 , R_1 , and R_2 are the same defined in Claims 1. In addition, R_1 can be absent and R_2 can be H.

29. The cell-binding molecule/agent, Cb, according to claim 2, 11, 12, 13 or 14 can be conjugated two or more different function molecules of claim 6 via the Formula (I) of claim 1 or via the Formula (IV) of claim 4.


30. The cell-binding molecule/agent, Cb, according to claim 2, 11, 12, 13, 14 or 29 is preferred an antibody or an antibody-like protein.

31. The IgG antibody conjugates when conjugated with one, or two, or more differently function molecules according to claim 2 or 29 are preferred to be conjugated specifically to a pair of thiols (through reduction of the disulfide bonds) between the light chain and heavy chain, the upper disulfide bonds between the two heavy chains and the lower disulfide bonds between the two heavy chains as shown in the following, ST1, ST2, ST3, ST4, ST5, ST6, ST7 or ST8.







ST3




ST4




ST5



ST6



ST7



















ST8









Wherein R_1 , R_1' , R_1'' , R_1''' , R_1'''' , and R_1''''' are independently defined as R_1 which is described the same in claim 1; Wherein R_2' , R_2'' , R_2''' , R_2'''' , and R_2''''' are independently defined as R_2 which is described the same in claim 1; X_1' , X_1'' , X_1''' , X_1'''' , and X_1''''' are independently de-









defined as X_1 which is described the same in claim 1; X_2' , X_2'' , X_2''' , X_2'''' , and X_2''''' are independently defined as X_2 which is described the same in claim 1; “ $\equiv$ ” represents either single bond or double bond; m_1 , m_2 , m_3 , m_4 , m_5 , m_6 , m_7 and m_8 are independently an integer from 1 to 30, preferably from 1 to 10; Drug₁, Drug₂, Drug₃, Drug₄, Drug₅ and Drug₆ is independently selected from tubulysins, maytansinoids, taxanoids (taxanes), CC-1065 analogs, daunorubicin and doxorubicin compounds, benzodiazepine dimers (e.g., dimers of pyrrolobenzodiazepine (PBD), tomaymycin, anthramycin, indolinobenzodiazepines, imidazobenzothiadiazepines, or oxazolidinobenzodiazepines), calicheamicins and the enediyne antibiotics, actinomycin, amanitin (amatoxins), azaserines, bleomycins, epirubicin, tamoxifen, idarubicin, dolastatins, auristatins (e.g. monomethyl auristatin E, MMAE, MMAF, auristatin PYE, auristatin TP, Auristatins 2-AQ, 6-AQ, EB (AEB), and EFP (AEFP)), duocarmycins, thiotepa, vincristines, hemi-asterlins, nazumamides, microginins, radiosumins, alterobactins, microsclerodermins, theonellamides, esperamicins, PNU-159682, and their analogues and derivatives. In addition, R_1 , R_1' , R_1'' , R_1''' , R_1'''' and/or R_1''''' can be absent.







32. The conjugates according to claim 2 or 31 are preferred having the Formula of 11, 25, 29, 33, 38, 45, 48, 57, 59, 67, 69, 78, 79a, 88, 92, 94, 99, 109, 110, 114, 120, 196a, 201, 206, 249, 251, 253, 256, 261, 268, 270, 272, 275, 277, 289, 293, 300, 302, 308, 311, 314, 317, 319, 321, 326, 329, 334, 338, 343, 345, 347, 354, 358, 362, 366, 370, 377, 385, 386, 389, 395, 396, 398, 402, 403, or 405 as shown in below:






























































































wherein R' is C₁-C₄ alkyl; m₁, m₂, drug, m, and n are defined the same as in Claim 1.









33. The compounds according to claim 4 are preferred having the Formula of 9, 10, 12, 13, 15, 16, 17, 18, 19, 20, 23, 24, 28, 30, 32, 36, 37, 40, 44, 47, 53, 54, 56, 57, 61, 62, 66, 68, 71, 72, 73, 77, 79, 81, 82, 87, 91, 93, 95, 96, 97, 98, 107, 108, 112, 113, 119, 196, 200, 205, 248, 250, 252, 254, 255, 259, 260, 267, 269, 271, 273, 274, 276, 288, 292, 299, 301, 307, 310, 313, 316, 318, 320, 325, 327, 328, 333, 337, 339, 340, 342, 344, 346, 353, 355, 357, 361, 365, 369, 376, 384, 388, 393, 394, 397, 400, 401, 406, 409, 411, 413, or 418, 420 or 423 as shown in below:










































































Wherein R' is C₁-C₄ alkyl; m₁, m₂, drug, m, and n are defined the same as in Claim 1.

- 34.** A pharmaceutical composition comprising a therapeutically effective amount of the conjugate compounds of claim 2, **18, 19, and/or 20, 21, 22, 23, 24, 25, 26, 27, 28, 31, or 32**, and a pharmaceutically acceptable salt, carrier, diluent, or excipient therefore, or a combination thereof, for the treatment or prevention of a cancer, or an autoimmune disease, or an infectious disease.
- 35.** The conjugate of claim **2, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 31, 32 or 34**, having in vitro, in vivo or ex vivo cell killing activity.
- 36.** The conjugate of claim **2, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, or 31** wherein the linkage components R_1 and/or R_2 can be cleavable by a protease.
- 37.** A pharmaceutical composition comprising a therapeutically effective amount of the conjugate of claim **2, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 31, 32 or 34**, administered concurrently with the other therapeutic agents such as the chemotherapeutic agent, the radiation therapy, immunotherapy agents, autoimmune disorder agents, anti-infectious agents or the other conjugates for synergistically effective treatment or prevention of a cancer, or an autoimmune disease, or an infectious disease.
- 38.** The synergistic agents according to claim **37** are preferably selected from one or several of the following drugs: Abatacept (Orencia), Abiraterone acetate (Zytiga®), Acetaminophen/hydrocodone, aducanumab, Adalimumab, ADXS31-142 (Advaxis'), ADXS-HER2 (Advaxis'), afatinib dimaleate (Gilotrif®), alemtuzumab (Campath®), Ali-tretinoin (Panretin®), adotrastuzumab emtansine (Kadcyla™), Amphetamine mixed salts (Amphetamine/ dextroamphetamine, or Adderall XR), anastrozole (Arimidex®), Aripiprazole, Atazanavir, Atezolizumab (MPDL3280A), Atorvastatin, axitinib (Inlyta®), Avelumab, belinostat (Beleodaq™), Bevacizumab (Avastin®), Cabazitaxel (Jevtana®), Cabozantinib (Come-triq™), bexarotene (Targretin®), blinatumomab (Blinicyto™), Bortezomib (Velcade®), bosutinib (Bosulif®), brentuximab vedotin (Adcetris®), Budesonide, Budesonide/formoterol, Buprenorphine, Capecitabine, carfilzomib (Kyprolis®), Celecoxib, ceritinib (LDK378/Zykadia), Cetuximab (Erbix®), Cyclosporin, Cinacalcet, crizotinib (Xalkori®), Cosentyx, CTL019, Dabigatran, dabrafenib (Tafinlar®), Daratumumab (Darzalex), Darbepoetin alfa, Darunavir, imatinib mesylate (Gleevec®), dasatinib (Sprycel®), denileukin diftitox (Ontak®), Denosumab (Xgeva®), Depakote, Dexlansoprazole, Dexmethylphenidate, Dexamethasone, Dignitana DigniCap Cooling System, Dinutuximab (Unituxin™), Doxycycline, Duloxetine, Duvelisib, elotuzumab, Emtricitabine/Ril-pivirine/Tenofovir disoproxil fumarate, Emtricitabine/tenofovir/ efavirenz, Enoxaparin, Enzalutamide (Xtandi®), Epoetin alfa, erlotinib (Tarceva®), Esomeprazole, Ezopiclone, Etanercept, Everolimus (Afinitor®), exemestane (Aromasin®), everolimus (Afinitor®), Ezetimibe, Ezetimibe/simvastatin, Fenofibrate, Filgrastim, fingolimod, Fluticasone propionate,

Fluticasone/salmeterol, fulvestrant (Faslodex®), gazyva, gefitinib (Iressa®), Glatiramer, Goserelin acetate (Zoladex), Icotinib, Imatinib (Gleevec), Ibritumomab tiuxetan (Zevalin®), ibrutinib (Imbruvica™), idelalisib (Zydelig®), Infliximab, Insulin aspart, Insulin detemir, Insulin glargine, Insulin lispro, Interferon beta 1a, Interferon beta 1b, lapatinib (Tykerb®), Ipilimumab (Yervoy®), Ipratropium bromide/salbutamol, Ixazomi (Ninlaro®), Kanuma, Lanreotide acetate (Somatuline® Depot), lenalidomide, lenaliomide (Revlimid®), lenvatinib mesylate (Lenvima™), letrozole (Femara®), Levothyroxine, Levothyroxine, Lidocaine, Linezolid, Liraglutide, Lisdexamfetamine, LN-144 (Lion Biotech.), MEDI4736 (AstraZeneca, Celgene), Memantine, Methylphenidate, Metoprolol, Mekinist, Modafinil, Mometasone, Nilotinib (Tasigna®), Nivolumab (Opdivo®), ofatumumab (Arzerra®), obinutuzumab (Gazyva™), olaparib (Lynparza™), Olmesartan, Olmesartan/ hydrochlorothiazide, Omalizumab, Omega-3 fatty acid ethyl esters, Oseltamivir, Oxycodone, palbociclib (Ibrance®), Palivizumab, panitumumab (Vectibix®), panobinostat (Farydak®), pazopanib (Votrient®), pembrolizumab (Keytruda®), Pemetrexed (Alimta), pertuzumab (Perjeta™), Pneumococcal conjugate vaccine, pomalidomide (Pomalyst®), Pregabalin, ProscarVax, Quetiapine, Rabeprazole, radium 223 chloride (Xofigo®), Raloxifene, Raltegravir, ramucirumab (Cyramza®), Ranibizumab, regorafenib (Stivarga®), Rituximab (Rituxan®), Rivaroxaban, romidepsin (Istodax®), Rosuvastatin, ruxolitinib phosphate (Jakafi™), Salbutamol, Sevelamer, Sildenafil, siltuximab (Sylvant™), Sitagliptin, Sitagliptin/metformin, Solifenacin, solanezumab, Sorafenib (Nexavar®), Sunitinib (Sutent®), Tadalafil, tamoxifen, Tafenlar, Telaprevir, temsirolimus (Torisel®), Tenofovir/emtricitabine, Testosterone gel, Thalidomide (Immuno-prin, Talidex), Tiotropium bromide, toremifene (Fareston®), trametinib (Mekinist®), Trastuzumab, Tretinoin (Vesanoid®), Ustekinumab, Valsartan, vandetanib (Caprelsa®), vemurafenib (Zelboraf®), venetoclax, vorinostat (Zolinza®), ziv-aflibercept (Zaltrap®), Zostavax., and their analogs, derivatives, pharmaceutically acceptable salts, carriers, diluents, or excipients thereof, or a combination above thereof.