

PYRAZOLO[1,5-a]PYRIMIDINE-BASED COMPOUNDS, COMPOSITIONS COMPRISING THEM, AND METHODS OF THEIR USE

This application claims priority to U.S. provisional patent application no. 61/608,765, filed March 9, 2012, the entirety of which is incorporated herein by reference.

5 1. FIELD OF THE INVENTION

This invention is directed to pyrazolo[1,5-a]pyrimidine-based compounds useful as inhibitors of adaptor associated kinase 1 (AAK1), compositions comprising them, and methods of their use.

2. BACKGROUND OF THE INVENTION


Adaptor associated kinase 1 (AAK1) is a member of the Ark1/Prk1 family of serine/threonine
10 kinases. AAK1 mRNA exists in two splice forms termed short and long. The long form predominates and is highly expressed in brain and heart (Henderson and Conner, Mol. Biol. Cell. 2007, 18, 2698-2706). AAK1 is enriched in synaptosomal preparations and is co-localized with endocytic structures in cultured cells. AAK1 modulates clatherin coated endocytosis, a process that is important in synaptic vesicle recycling and receptor-mediated endocytosis. AAK1 associates with the AP2
15 complex, a hetero-tetramer which links receptor cargo to the clatherin coat. The binding of clatherin to AAK1 stimulates AAK1 kinase activity (Conner et. al., Traffic 2003, 4, 885-890; Jackson et. al., J. Cell. Biol. 2003, 163, 231-236). AAK1 phosphorylates the mu-2 subunit of AP-2, which promotes the binding of mu-2 to tyrosine containing sorting motifs on cargo receptors (Ricotta et. al., J. Cell Bio. 2002, 156, 791-795; Conner and Schmid, J. Cell Bio. 2002, 156, 921-929). Mu2 phosphorylation is
20 not required for receptor uptake, but phosphorylation enhances the efficiency of internalization (Motely et. al., Mol. Biol. Cell. 2006, 17, 5298-5308).

AAK1 has been identified as an inhibitor of Neuregulin-1/ErbB4 signaling in PC12 cells. Loss of AAK1 expression through RNA interference mediated gene silencing or treatment with the kinase inhibitor K252a (which inhibits AAK1 kinase activity) results in the potentiation of Neuregulin-1
25 induced neurite outgrowth. These treatments result in increased expression of ErbB4 and accumulation of ErbB4 in or near the plasma membrane (Kuai et. al., Chemistry and Biology 2011, 18, 891-906). NRG1 and ErbB4 are putative schizophrenia susceptibility genes (Buonanno, Brain Res. Bull. 2010, 83, 122-131). SNPs in both genes have been associated with multiple schizophrenia endophenotypes (Greenwood et. al., Am. J. Psychiatry 2011, 168, 930-946).
30 Neuregulin 1 and ErbB4 KO mouse models have shown schizophrenia relevant morphological changes and behavioral phenotypes (Jaaro-Peled et. al., Schizophrenia Bulletin 2010, 36, 301-313; Wen et. al., Proc. Natl. Acad. Sci. USA. 2010, 107, 1211-1216). In addition, a single nucleotide polymorphism in an intron of the AAK1 gene has been associated with the age of onset of Parkinson's disease (Latourelle et. al., BMC Med. Genet. 2009, 10, 98). These results suggest that

inhibition of AAK1 activity may have utility in the treatment of schizophrenia, cognitive deficits in schizophrenia, Parkinson's disease, bipolar disorder, and Alzheimer's disease.

3. SUMMARY OF THE INVENTION

This invention is directed, in part, to AAK1 inhibitors of the formula:



and pharmaceutically acceptable salt thereof, wherein: R_1 is R_{1A} or optionally substituted C_{1-12} hydrocarbyl or 2-12-membered heterocarbyl, which optional substitution is with one or more R_{1A} ; each R_{1A} is independently $-OR_{1C}$, $-N(R_{1C})_2$, $-C(O)R_{1C}$, $-C(O)OR_{1C}$, $-C(O)N(R_{1C})_2$, $-N(R_{1C})C(O)OR_{1C}$, cyano, halo, or optionally substituted C_{1-12} hydrocarbyl or 2-12-membered heterocarbyl, which optional substitution is with one or more R_{1B} ; each R_{1B} is independently $-OR_{1C}$, $-N(R_{1C})_2$, $-C(O)R_{1C}$, $-C(O)OR_{1C}$, $-C(O)N(R_{1C})_2$, $-N(R_{1C})C(O)OR_{1C}$, cyano or halo; each R_{1C} is independently hydrogen or optionally substituted C_{1-12} hydrocarbyl or 2-12-membered heterocarbyl, which optional substitution is with one or more of cyano, halo or hydroxyl; R_2 is $-NR_{2A}R_{2B}$, wherein R_{2A} is hydrogen and R_{2B} is optionally substituted C_{1-12} hydrocarbyl or 2-12-membered heterocarbyl, which optional substitution is with one or more R_{2C} ; or R_{2A} and R_{2B} are taken together to form a 4-7-membered heterocycle optionally substituted with one or more R_{2C} ; each R_{2C} is independently $-OR_{2D}$, $-N(R_{2D})_2$, $-C(O)R_{2D}$, $-C(O)OR_{2D}$, $-C(O)N(R_{2D})_2$, $-N(R_{2D})C(O)OR_{2D}$, cyano, halo, or optionally substituted C_{1-12} hydrocarbyl or 2-12-membered heterocarbyl, which optional substitution is with one or more with one or more R_{2D} ; each R_{2D} is independently hydrogen or optionally substituted C_{1-12} hydrocarbyl or 2-12-membered heterocarbyl, which optional substitution is with one or more of cyano, halo or hydroxyl; and R_3 is hydrogen or C_{1-6} alkyl optionally substituted with one or more of cyano, halo or hydroxyl.

One embodiment of the invention encompasses pharmaceutical compositions and dosage forms comprising a compound disclosed herein (*i.e.*, a compound of the invention).

Another embodiment of this invention encompasses methods of inhibiting adaptor associated kinase 1 (AAK1), both *in vitro* and *in vivo*, which comprise contacting AAK1 with a compound of the invention.

Another embodiment encompasses methods of treating and managing diseases and disorders mediated by AAK1 activity. Examples of such diseases and disorders are believed to include Alzheimer's disease, bipolar disorder, pain, Parkinson's disease, and schizophrenia (including cognitive deficits in schizophrenia).

4. BRIEF DESCRIPTION OF THE FIGURES

Aspects of the invention are illustrated in Figure 1, which shows results obtained from a formalin pain model using AAK1 homozygous (-/-) knockout mice and their wild-type (+/+) littermates. The AAK1 homozygous (-/-) knockout mice show a clear reduction in both acute and tonic pain response as compared to their wild-type (+/+) littermates.

5. DETAILED DESCRIPTION OF THE INVENTION

This invention is based, in part, on the discovery that AAK1 knockout mice exhibit a high resistance to pain. That discovery prompted research that ultimately led to the discovery of AAK1 inhibitors, compositions comprising them, and methods of their use.

5.1. DEFINITIONS

Unless otherwise indicated, the phrases "compounds of the invention," "compounds of the present disclosure," and the like refer to the compounds disclosed herein.

Unless otherwise indicated, the term "hydrocarbonyl" means an aliphatic or alicyclic moiety having an all-carbon backbone and consisting of carbon and hydrogen atoms. Examples of hydrocarbonyl groups include those having 1-20, 1-12, 1-6, and 1-4 carbon atoms (referred to as C₁₋₂₀ hydrocarbonyl, C₁₋₁₂ hydrocarbonyl, C₁₋₆ hydrocarbonyl, and C₁₋₄ hydrocarbonyl, respectively). Particular examples include alkyl, alkenyl, alkynyl, aryl, benzyl, cycloalkyl, cycloalkenyl, cycloalkylalkyl, cycloalkenylalkyl, naphthyl, phenyl, and phenylethyl.

Examples of alkyl moieties include straight-chain and branched moieties having 1-20, 1-12, 1-6, 1-4 and 1-3 carbon atoms (referred to as C₁₋₂₀ alkyl, C₁₋₁₂ alkyl, C₁₋₆ alkyl, C₁₋₄ alkyl and C₁₋₃ alkyl, respectively). Particular examples include methyl, ethyl, propyl, isopropyl, n-butyl, t-butyl, isobutyl, pentyl, hexyl, isohexyl, heptyl, 4,4-dimethylpentyl, octyl, 2,2,4-trimethylpentyl, nonyl, decyl, undecyl and dodecyl.

Examples of alkenyl moieties include straight-chain and branched C₂₋₂₀, C₂₋₁₂ and C₂₋₆ alkenyl. Particular examples include vinyl, allyl, 1-butenyl, 2-butenyl, isobutenyl, 1-pentenyl, 2-pentenyl, 3-methyl-1-butenyl, 2-methyl-2-butenyl, 2,3-dimethyl-2-butenyl, 1-hexenyl, 2-hexenyl, 3-hexenyl, 1-heptenyl, 2-heptenyl, 3-heptenyl, 1-octenyl, 2-octenyl, 3-octenyl, 1-nonenyl, 2-nonenyl, 3-nonenyl, 1-decenyl, 2-decenyl and 3-decenyl.

Examples of alkynyl moieties include include straight-chain and branched C₂₋₂₀, C₂₋₁₂ and C₂₋₆ alkynyl. Particular examples include ethynyl and 2-propynyl (propargyl).

Examples of aryl moieties include anthracenyl, azulenyl, fluorenyl, indan, indenyl, naphthyl, phenyl and phenanthrenyl.

Examples of cycloalkyl moieties include C₃₋₁₂, C₃₋₇, C₄₋₆ and C₆ cycloalkyl. Particular examples include cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and adamantyl.

Unless otherwise indicated, the term "halo" encompass fluoro, chloro, bromo, and iodo.

Unless otherwise indicated, the term “heterocarbyl” refers to a moiety having a backbone made up of one or more carbon atoms and one or more heteroatoms. Particular heteroatoms are nitrogen, oxygen and sulfur. A heterocarbyl moieties can be thought of as a hydrocarbyl moiety wherein at least one carbon atom, CH, CH₂, or CH₃ group is replaced with one or more heteroatoms and the requisite number of hydrogen atoms to satisfy valencies. Examples of heterocarbyl include 2-20, 2-12, 2-8, 2-6 and 2-4 membered heterocarbyl moieties, wherein the number range refers to the sum total of carbon, nitrogen, oxygen, and/or sulfur atoms in the moiety. The term “2-12 membered heterocarbyl” thus refers to a heterocarbyl moiety having a total of 2-12 carbon, nitrogen, oxygen, and/or sulfur atoms. Particular heterocarbyl moieties include straight chain and branched heteroalkyl, heteroalkenyl, and heteroalkynyl, as well as heterocycle and heteroaryl.

Examples of heteroalkyl moieties include 2-8-membered, 2-6-membered and 2-4-membered heteroalkyl moieties. Particular examples include alkoxyl, acyl (e.g., formyl, acetyl, benzoyl), alkylamino (e.g., di-(C₁₋₃-alkyl)amino), arylamino, aryloxime, carbamates, carbamides, alkylcarbonyl, arylcarbonyl, aminocarbonyl, alkylaminocarbonyl, alkylsulfanyl, arylsulfanyl, alkylsulfinyl, arylsulfinyl, alkylsulfonyl, arylsulfonyl, alkylsulfonylamino, and arylsulfonylamino.

Unless otherwise indicated, the term “heterocycle” refers to a cyclic (monocyclic or polycyclic) heterocarbyl moiety which may be aromatic, partially aromatic or non-aromatic. Heterocycles include heteroaryls. Examples include 4-10-membered, 4-7-membered, 6-membered, and 5-membered heterocycles. Particular examples include benzo[1,3]dioxolyl, 2,3-dihydro-benzo[1,4]dioxinyl, cinnoliny, furanyl, hydantoinyl, morpholiny, oxetanyl, oxiranyl, piperazinyl, piperidinyl, pyrrolidinonyl, pyrrolidinyl, tetrahydrofuranyl, tetrahydropyranyl, tetrahydropyridinyl, tetrahydropyrimidinyl, tetrahydrothiophenyl, tetrahydrothiopyranyl and valerolactamyl. Because the term “heterocycle” refers to a ring, standing alone it does not encompass moieties such as oxazolidinone and imidazolidinone: such moieties are considered substituted heterocycles, viz. heterocycles substituted with oxo.

Examples of heteroaryl moieties include acridinyl, benzimidazolyl, benzofuranyl, benzoisothiazolyl, benzoisoxazolyl, benzoquinazolinyl, benzothiazolyl, benzoxazolyl, furyl, imidazolyl, indolyl, isothiazolyl, isoxazolyl, oxadiazolyl, oxazolyl, phthalazinyl, pyrazinyl, pyrazolyl, pyridazinyl, pyridyl, pyrimidinyl, pyrimidyl, pyrrolyl, quinazolinyl, quinoliny, tetrazolyl, thiazolyl, and triazinyl.

Unless otherwise indicated, the term “include” has the same meaning as “include, but are not limited to,” and the term “includes” has the same meaning as “includes, but is not limited to.” Similarly, the term “such as” has the same meaning as the term “such as, but not limited to.”

Unless otherwise indicated, the terms “manage,” “managing” and “management” encompass preventing the recurrence of the specified disease or disorder in a patient who has already suffered from the disease or disorder, and/or lengthening the time that a patient who has suffered from the disease or disorder remains in remission. The terms encompass modulating the

threshold, development and/or duration of the disease or disorder, or changing the way that a patient responds to the disease or disorder.


Unless otherwise indicated, a “therapeutically effective amount” of a compound is an amount sufficient to provide a therapeutic benefit in the treatment or management of a disease or condition, or to delay or minimize one or more symptoms associated with the disease or condition. A “therapeutically effective amount” of a compound means an amount of therapeutic agent, alone or in combination with other therapies, that provides a therapeutic benefit in the treatment or management of the disease or condition. The term “therapeutically effective amount” can encompass an amount that improves overall therapy, reduces or avoids symptoms or causes of a disease or condition, or enhances the therapeutic efficacy of another therapeutic agent.

Unless otherwise indicated, the terms “treat,” “treating” and “treatment” contemplate an action that occurs while a patient is suffering from the specified disease or disorder, which reduces the severity of the disease or disorder, or retards or slows the progression of the disease or disorder.

Unless otherwise indicated, one or more adjectives immediately preceding a series of nouns is to be construed as applying to each of the nouns. For example, the phrase “optionally substituted alky, aryl, or heteroaryl” has the same meaning as “optionally substituted alky, optionally substituted aryl, or optionally substituted heteroaryl.”

5.2. COMPOUNDS

This invention encompasses compounds of the formula:



and pharmaceutically acceptable salt thereof, wherein: R₁ is R_{1A} or optionally substituted C₁₋₁₂ hydrocarbyl or 2-12-membered heterocarbyl, which optional substitution is with one or more R_{1A}; each R_{1A} is independently -OR_{1C}, -N(R_{1C})₂, -C(O)R_{1C}, -C(O)OR_{1C}, -C(O)N(R_{1C})₂, -N(R_{1C})C(O)OR_{1C}, cyano, halo, or optionally substituted C₁₋₁₂ hydrocarbyl or 2-12-membered heterocarbyl, which optional substitution is with one or more R_{1B}; each R_{1B} is independently -OR_{1C}, -N(R_{1C})₂, -C(O)R_{1C}, -C(O)OR_{1C}, -C(O)N(R_{1C})₂, -N(R_{1C})C(O)OR_{1C}, cyano or halo; each R_{1C} is independently hydrogen or optionally substituted C₁₋₁₂ hydrocarbyl or 2-12-membered heterocarbyl, which optional substitution is with one or more of cyano, halo or hydroxyl; R₂ is -NR_{2A}R_{2B}, wherein R_{2A} is hydrogen and R_{2B} is optionally substituted C₁₋₁₂ hydrocarbyl or 2-12-membered heterocarbyl, which optional substitution is with one or more R_{2C}; or R_{2A} and R_{2B} are taken together to form a 4-7-membered heterocycle optionally substituted with one or more R_{2C}; each R_{2C} is independently -OR_{2D}, -N(R_{2D})₂, -C(O)R_{2D}, -C(O)OR_{2D}, -C(O)N(R_{2D})₂, -N(R_{2D})C(O)OR_{2D}, cyano, halo, or optionally substituted C₁₋₁₂ hydrocarbyl or 2-12-membered heterocarbyl, which optional substitution is with one or more with one or more R_{2D}; each

R_{2D} is independently hydrogen or optionally substituted C₁₋₁₂ hydrocarbyl or 2-12-membered heterocarbyl, which optional substitution is with one or more of cyano, halo or hydroxyl; and R₃ is hydrogen or C₁₋₆ alkyl optionally substituted with one or more of cyano, halo or hydroxyl.

In particular compounds, R₁ is R_{1A}. In some, R₁ is optionally substituted C₁₋₁₂ hydrocarbyl. In some, R₁ is optionally substituted phenyl. In some, R₁ is optionally substituted 2-12-membered heterocarbyl (e.g., 2-8 membered heterocarbyl, 2-6 membered heterocarbyl, 2-6 membered heterocarbyl). In some, R₁ is optionally substituted pyridinyl, thiophen, or imidazol.

In particular compounds, R_{1A} is halo. In some, R_{1A} is -OR_{1C}, -N(R_{1C})₂, -C(O)R_{1C}, -C(O)OR_{1C}, or -C(O)N(R_{1C})₂. In some, R_{1A} is -OR_{1C}.

In particular compounds, R_{1B} is -N(R_{1C})₂, -OR_{1C}, halo.

In particular compounds, R_{2A} and R_{2B} are taken together to form a 4-7-membered heterocycle optionally substituted with one or more R_{2C}.


In particular compounds, R_{1C} is hydrogen. In some, R_{1C} is C₁₋₁₂ hydrocarbyl (e.g., C₁₋₆ hydrocarbyl, C₁₋₄ hydrocarbyl such as methyl, ethyl, propyl). In some, R_{2C} is -C(O)OR_{2D}, -C(O)N(R_{2D})₂, or -N(R_{2D})C(O)OR_{2D}.

In particular compounds, R_{2D} is hydrogen.

In particular compounds, R_{2D} is C₁₋₁₂ hydrocarbyl (e.g., C₁₋₆ hydrocarbyl, C₁₋₄ hydrocarbyl such as methyl, ethyl, propyl).

In particular compounds, R₃ is hydrogen.

One embodiment of the invention encompasses compounds of the formula:



and pharmaceutically acceptable salt thereof, wherein: A is cyclic C₁₋₁₂ hydrocarbyl or 4-7-membered heterocycle; D is 4-7-membered heterocycle; each R_{1A} is independently -OR_{1C}, -N(R_{1C})₂, -C(O)R_{1C}, -C(O)OR_{1C}, -C(O)N(R_{1C})₂, -N(R_{1C})C(O)OR_{1C}, cyano, halo, or optionally substituted C₁₋₁₂ hydrocarbyl or 2-12-membered heterocarbyl, which optional substitution is with one or more R_{1B}; each R_{1B} is independently -OR_{1C}, -N(R_{1C})₂, -C(O)R_{1C}, -C(O)OR_{1C}, -C(O)N(R_{1C})₂, -N(R_{1C})C(O)OR_{1C}, cyano or halo; each R_{1C} is independently hydrogen or optionally substituted C₁₋₁₂ hydrocarbyl or 2-12-membered heterocarbyl, which optional substitution is with one or more of cyano, halo or hydroxyl; each R_{2C} is independently -OR_{2D}, -N(R_{2D})₂, -C(O)R_{2D}, -C(O)OR_{2D}, -C(O)N(R_{2D})₂, -N(R_{2D})C(O)OR_{2D}, cyano, halo, or optionally substituted C₁₋₁₂ hydrocarbyl or 2-12-membered heterocarbyl, which optional substitution is with one or more with one or more R_{2D}; each R_{2D} is independently hydrogen or optionally substituted C₁₋₁₂ hydrocarbyl or 2-12-membered heterocarbyl, which optional substitution is with one or more of cyano, halo or hydroxyl; n is 1-3; and m is 0-3.

In particular compounds, D is not piperidinyl.

In particular compounds, R_{2C} is not $-N(R_{2D})_2$.

In particular compounds, A is not phenyl.

In particular compounds, m is 1.

In particular compounds, m is 2.

5 In particular compounds, R_{2D} is not ethyl.


In particular compounds, when D is piperidinyl, A is phenyl, and R_{2C} is $-N(R_{2D})_2$, R_{2D} is not ethyl.

In particular compounds, D is piperazin or pyrrolidin.

In particular compounds, n is 1.


10 In particular compounds, A is pyridinyl, thiophen, or imidazol.

Particular compounds are of the formula:



wherein X is CH or N.

Another embodiment of the invention encompasses compounds of the formula:



15


and pharmaceutically acceptable salt thereof, wherein: X is CH or N; each R_{1A} is independently $-OR_{1C}$, $-N(R_{1C})_2$, $-C(O)R_{1C}$, $-C(O)OR_{1C}$, $-C(O)N(R_{1C})_2$, $-N(R_{1C})C(O)OR_{1C}$, cyano, halo, or optionally substituted C_{1-12} hydrocarbyl or 2-12-membered heterocarbyl, which optional substitution is with one or more R_{1B} ; each R_{1B} is independently $-OR_{1C}$, $-N(R_{1C})_2$, $-C(O)R_{1C}$, $-C(O)OR_{1C}$, $-C(O)N(R_{1C})_2$, $-N(R_{1C})C(O)OR_{1C}$, cyano or halo; each R_{1C} is independently hydrogen or optionally substituted C_{1-12} hydrocarbyl or 2-12-membered heterocarbyl, which optional substitution is with one or more of cyano, halo or hydroxyl; each R_{2C} is independently $-OR_{2D}$, $-N(R_{2D})_2$, $-C(O)R_{2D}$, $-C(O)OR_{2D}$, $-C(O)N(R_{2D})_2$, $-N(R_{2D})C(O)OR_{2D}$, cyano, halo, or optionally substituted C_{1-12} hydrocarbyl or 2-12-membered heterocarbyl, which optional substitution is with one or more with one or more R_{2D} ; each R_{2D} is independently hydrogen or optionally substituted C_{1-12} hydrocarbyl or 2-12-membered heterocarbyl, which optional substitution is with one or more of cyano, halo or hydroxyl; and m is 0-3.

25


In particular compounds, R_{2C} is not optionally substituted phenyl or pyridinyl.

In particular compounds, X is N and m is 1 or 2.


Particular compounds are of the formula:




Particular compounds are of the formula:




5 Particular compounds are of the formula:



Others are of the formula:




Another embodiment of the invention encompasses compounds of the formula:




10 and pharmaceutically acceptable salt thereof, wherein: X is CH or N; each R_{1A} is independently -OR_{1C}, -N(R_{1C})₂, -C(O)R_{1C}, -C(O)OR_{1C}, -C(O)N(R_{1C})₂, -N(R_{1C})C(O)OR_{1C}, cyano, halo, or optionally substituted C₁₋₁₂ hydrocarbyl or 2-12-membered heterocarbyl, which optional substitution is with one or more R_{1B}; each R_{1B} is independently -OR_{1C}, -N(R_{1C})₂, -C(O)R_{1C}, -C(O)OR_{1C}, -C(O)N(R_{1C})₂, -N(R_{1C})C(O)OR_{1C}, cyano or
 15 halo; each R_{1C} is independently hydrogen or optionally substituted C₁₋₁₂ hydrocarbyl or 2-12-membered heterocarbyl, which optional substitution is with one or more of cyano, halo or hydroxyl; each R_{2C} is independently -OR_{2D}, -N(R_{2D})₂, -C(O)R_{2D}, -C(O)OR_{2D}, -C(O)N(R_{2D})₂, -N(R_{2D})C(O)OR_{2D}, cyano, halo, or optionally substituted C₁₋₁₂ hydrocarbyl or 2-12-membered heterocarbyl, which optional substitution is with one or more with one or more R_{2D}; each R_{2D} is independently hydrogen or
 20 optionally substituted C₁₋₁₂ hydrocarbyl or 2-12-membered heterocarbyl, which optional substitution is with one or more of cyano, halo or hydroxyl; and m is 0-3.


Particular compounds are of the formula:



Particular compounds are of the formula:



5 Others are of the formula:



In particular compounds, R_{1A} is halo. In some, R_{1A} is -OR_{1C}.

In particular compounds, R_{1C} is optionally substituted C₁₋₁₂ hydrocarbyl (e.g., C₁₋₆ hydrocarbyl, C₁₋₄ hydrocarbyl).

10 In particular compounds, R_{2C} is -C(O)OR_{2D}, -C(O)N(R_{2D})₂, or -N(R_{2D})C(O)OR_{2D}. In some, R_{2C} is -C(O)R_{2D}.

In particular compounds, R_{2D} is independently hydrogen or C₁₋₁₂ hydrocarbyl (e.g., C₁₋₆ hydrocarbyl, C₁₋₄ hydrocarbyl). In some, R_{2D} is optionally substituted C₁₋₁₂ hydrocarbyl, which optional substitution is with one or more of amino, cyano, halo, hydroxyl.

15 Compounds of the invention can have one or more asymmetric centers. Unless otherwise indicated, this invention encompasses all stereoisomers of the compounds, as well as mixtures thereof. Individual stereoisomers of compounds can be prepared synthetically from commercially available starting materials which contain chiral centers or by preparation of mixtures of enantiomeric products followed by separation such as conversion to a mixture of diastereomers
20 followed by separation or recrystallization, chromatographic techniques, or direct separation of enantiomers on chiral chromatographic columns. Starting compounds of particular stereochemistry are either commercially available or can be made and resolved by techniques known in the art.

Certain compounds of the present disclosure may also exist in different stable conformational forms which may be separable. Torsional asymmetry due to restricted rotation about
25 an asymmetric single bond, for example because of steric hindrance or ring strain, may permit

separation of different conformers. The present disclosure includes each conformational isomer of these compounds and mixtures thereof.

The present disclosure is intended to include all isotopes of atoms occurring in the present compounds. Isotopes include those atoms having the same atomic number but different mass numbers. By way of general example and without limitation, isotopes of hydrogen include deuterium and tritium. Isotopes of carbon include ^{13}C and ^{14}C . Isotopically-labeled compounds of the invention can generally be prepared by conventional techniques known to those skilled in the art or by processes analogous to those described herein, using an appropriate isotopically-labeled reagent in place of the non-labeled reagent otherwise employed. Such compounds may have a variety of potential uses, for example as standards and reagents in determining biological activity. In the case of stable isotopes, such compounds may have the potential to favorably modify biological, pharmacological, or pharmacokinetic properties.

The compounds of the present disclosure can exist as pharmaceutically acceptable salts. The term "pharmaceutically acceptable salt," as used herein, represents salts or zwitterionic forms of the compounds of the present disclosure which are water or oil-soluble or dispersible, which are, within the scope of sound medical judgment, suitable for use in contact with the tissues of patients without excessive toxicity, irritation, allergic response, or other problem or complication commensurate with a reasonable benefit/risk ratio, and are effective for their intended use. The salts can be prepared during the final isolation and purification of the compounds or separately by reacting a suitable nitrogen atom with a suitable acid. Representative acid addition salts include acetate, adipate, alginate, citrate, aspartate, benzoate, benzenesulfonate, bisulfate, butyrate, camphorate, camphorsulfonate; digluconate, dihydrobromide, dihydrochloride, dihydroiodide, glycerophosphate, hemisulfate, heptanoate, hexanoate, formate, fumarate, hydrochloride, hydrobromide, hydroiodide, 2-hydroxyethanesulfonate, lactate, maleate, mesitylenesulfonate, methanesulfonate, naphthylenesulfonate, nicotinate, 2-naphthalenesulfonate, oxalate, palmoate, pectinate, persulfate, 3-phenylpropionate, picrate, pivalate, propionate, succinate, tartrate, trichloroacetate, trifluoroacetate, phosphate, glutamate, bicarbonate, para-toluenesulfonate, and undecanoate. Examples of acids which can be employed to form pharmaceutically acceptable addition salts include inorganic acids such as hydrochloric, hydrobromic, sulfuric, and phosphoric, and organic acids such as oxalic, maleic, succinic, and citric.


Basic addition salts can be prepared during the final isolation and purification of the compounds by reacting a carboxy group with a suitable base such as the hydroxide, carbonate, or bicarbonate of a metal cation or with ammonia or an organic primary, secondary, or tertiary amine. The cations of pharmaceutically acceptable salts include lithium, sodium, potassium, calcium, magnesium, and aluminum, as well as nontoxic quaternary amine cations such as ammonium, tetramethylammonium, tetraethylammonium, methylamine, dimethylamine, trimethylamine, triethylamine, diethylamine, ethylamine, tributylamine, pyridine, N,N-dimethylaniline, N-

methypiperidine, N-methylmorpholine, dicyclohexylamine, procaine, dibenzylamine, N,N-dibenzylphenethylamine, and N,N'-dibenzylethylenediamine. Other representative organic amines useful for the formation of base addition salts include ethylenediamine, ethanolamine, diethanolamine, piperidine, and piperazine.

- 5 Particular compounds of the invention inhibit AAK1 with an IC_{50} of less than 0.1, 0.01 or 0.001 μM as measured in the P81 filter plate assay described below in the Examples. Particular compounds of the invention inhibit AAK1 with an IC_{50} of less than 0.1, 0.01 or 0.001 μM as measured in the HEK281 cell-based assay described below in the Examples.

5.3. METHODS OF SYNTHESIS


- 10 Compounds of the present invention (*i.e.*, compounds disclosed herein) can be prepared using the methods described below and using methods known to those skilled in the art of organic chemistry. Particular compounds are of the general formula:



- 15 wherein R_1 , R_2 and R_3 are defined herein, and include salts thereof. These compounds can be prepared by the methods outlined below.


- Compounds of formula shown above can be prepared by the methods outlined below. In Scheme 1, the chlorine of a compound of formula 1 is displaced with an amine to produce 2. Bromination of 2 provides 3. Alternatively 1 can be brominated first to afford 4 then amine displacement to give 3. Suzuki coupling of 3 with an appropriate boronic acid [$R_1B(OH)_2$] affords compounds of formula 5.

Scheme 1



In Scheme 2, the substituted acetonitrile **6** is reacted with ethyl formate to provide **7**. Condensation of **7** with hydrazine hydrate affords **8**. Reaction of **8** with 1,3-dimethylpyrimidine-2,4(1H,3H)-dione provides **9**. Further reaction of **9** with amine gives compounds with formula **5**.


Scheme 2



5

In Scheme 3, compound with formula **10** can be prepared by reacting **2** with NIS. Reaction of **10** with methyl 2,2-difluoro-2-(fluorosulfonyl)acetate provides compounds with formula **11**.

Scheme 3



10 5.4. METHODS OF USE

One embodiment of this invention encompasses methods of inhibiting adaptor associated kinase 1 (AAK1), both *in vitro* and *in vivo*, which comprise contacting AAK1 with a compound of the invention.

Another embodiment encompasses methods of treating and managing diseases and disorders mediated by AAK1 activity. Diseases and disorders mediated by AAK1 activity are diseases and disorders that have at least one symptom, the severity or manifestation of which is affected by AAK1 activity. Examples of such diseases and disorders are believed to include Alzheimer's disease, bipolar disorder, pain, Parkinson's disease, and schizophrenia (including cognitive deficits in schizophrenia). Particular methods comprise administering to a patient (a human or other mammal) in need thereof a therapeutically or prophylactically effective amount of an AAK1 inhibitor (e.g., a compound disclosed herein).

Another embodiment of this invention encompasses a method of treating or managing a disease or disorder, which comprises administering to a patient in need thereof a therapeutically or prophylactically effective amount of an AAK1 inhibitor, wherein the disease or disorder is Alzheimer's disease, bipolar disorder, pain, Parkinson's disease, or schizophrenia (including cognitive deficits in schizophrenia). Particular types of pain include chronic pain, acute pain, and neuropathic pain. Particular types of neuropathic pain include fibromyalgia and peripheral neuropathy (e.g., diabetic neuropathy).

When used to treat or manage a disease or disorder, compounds of the invention are preferably administered as part of a pharmaceutical composition comprising one or more pharmaceutically acceptable carriers, diluents or excipients.

Pharmaceutical compositions, or formulations, may be presented in unit dose forms containing a predetermined amount of active ingredient per unit dose. Dosage levels of between about 0.01 and about 250 milligram per kilogram ("mg/kg") body weight per day, preferably between about 0.05 and about 100 mg/kg body weight per day of the compounds of the present disclosure are typical in a monotherapy for the prevention and treatment of disease. Typically, the pharmaceutical compositions of this disclosure will be administered from about 1 to about 5 times per day or alternatively, as a continuous infusion. Such administration can be used as a chronic or acute therapy. The amount of active ingredient that may be combined with the carrier materials to produce a single dosage form will vary depending on the condition being treated, the severity of the condition, the time of administration, the route of administration, the rate of excretion of the compound employed, the duration of treatment, and the age, gender, weight, and condition of the patient. Preferred unit dosage formulations are those containing a daily dose or sub-dose, as herein above recited, or an appropriate fraction thereof, of an active ingredient. Treatment may be initiated with small dosages substantially less than the optimum dose of the compound. Thereafter, the dosage is increased by small increments until the optimum effect under the circumstances is reached. In general, the compound is most desirably administered at a concentration level that will generally afford effective results without causing any harmful or deleterious side effects.

Compounds of the invention may be administered in combination with one or more additional therapeutic or prophylactic agents. For example, when used for the treatment of pain, possible additional agents include immunosuppressive and anti-inflammatory agents.

Immunosuppressants suitable for use in the methods and compositions of this invention include those known in the art. Examples include aminopterin, azathioprine, cyclosporin A, D-penicillamine, gold salts, hydroxychloroquine, leflunomide, methotrexate, minocycline, rapamycin, sulfasalazine, tacrolimus (FK506), and pharmaceutically acceptable salts thereof. A particular immunosuppressant is methotrexate.

Additional examples include anti-TNF antibodies, such as adalimumab, certolizumab pegol, etanercept, and infliximab. Others include interleukin-1 blockers, such as anakinra. Others include

anti-B cell (CD20) antibodies, such as rituximab. Others include T cell activation blockers, such as abatacept.

Additional examples include inosine monophosphate dehydrogenase inhibitors, such as mycophenolate mofetil (CellCept®) and mycophenolic acid (Myfortic®).

5 Anti-inflammatory drugs suitable for use in the methods and compositions of this invention include those known in the art. Examples include glucocorticoids and NSAIDs.

Examples of glucocorticoids include aldosterone, beclometasone, betamethasone, cortisone, deoxycorticosterone, dexamethasone, fludrocortisone, hydrocortisone, methylprednisolone, prednisolone, prednisone, triamcinolone, and pharmaceutically acceptable salts thereof.

10 Examples of NSAID include salicylates (e.g., aspirin, amoxiprin, benorilate, choline magnesium salicylate, diflunisal, faislamine, methyl salicylate, magnesium salicylate, salicyl salicylate, and pharmaceutically acceptable salts thereof), arylalkanoic acids (e.g., diclofenac, aceclofenac, acemetacin, bromfenac, etodolac, indometacin, nabumetone, sulindac, tolmetin, and pharmaceutically acceptable salts thereof), arylpropionic acids (e.g., ibuprofen, carprofen, fenbufen, 15 fenoprofen, flurbiprofen, ketoprofen, ketorolac, loxoprofen, naproxen, oxaprozin, tiaprofenic acid, suprofen, and pharmaceutically acceptable salts thereof), arylanthranilic acids (e.g., meclofenamic acid, mefenamic acid, and pharmaceutically acceptable salts thereof), pyrazolidine derivatives (e.g., azapropazone, metamizole, oxyphenbutazone, phenylbutazone, sulfinprazone, and pharmaceutically acceptable salts thereof), oxicams (e.g., lornoxicam, meloxicam, piroxicam, tenoxicam, and 20 pharmaceutically acceptable salts thereof), COX-2 inhibitors (e.g., celecoxib, etoricoxib, lumiracoxib, parecoxib, rofecoxib, valdecoxib, and pharmaceutically acceptable salts thereof), and sulphonanilides (e.g., nimesulide and pharmaceutically acceptable salts thereof).

Other agents used in the treatment of pain (including but not limited to neuropathic and inflammatory pain) include agents such as pregabalin, lidocaine, duloxetine, gabapentin, 25 carbamazepine, capsaicin, and other serotonin/norepinephrine/dopamine reuptake inhibitors, and opiates (such as oxycontin, morphine, and codeine).

In the treatment of pain caused by a known disease or condition, such as diabetes, infection (e.g., herpes zoster or HIV infection), or cancer, compounds of the invention may be administered in combination with one or more additional therapeutic or prophylactic agents directed at the 30 underlying disease or condition. For example, when used to treat diabetic neuropathy, compounds of the invention may be administered in combination with one or more anti-diabetic agents, anti-hyperglycemic agents, hypolipidemic/lipid lowering agents, anti-obesity agents, anti-hypertensive agents and appetite suppressants. Examples of anti-diabetic agents include biguanides (e.g., metformin, phenformin), glucosidase inhibitors (e.g., acarbose, miglitol), insulins (including insulin 35 secretagogues and insulin sensitizers), meglitinides (e.g., repaglinide), sulfonylureas (e.g., glimepiride, glyburide, gliclazide, chlorpropamide, and glipizide), biguanide/glyburide combinations (e.g., Glucovance), thiazolidinediones (e.g., troglitazone, rosiglitazone, and pioglitazone), PPAR-alpha

agonists, PPAR-gamma agonists, PPAR alpha/gamma dual agonists, glycogen phosphorylase inhibitors, inhibitors of fatty acid binding protein (aP2), glucagon-like peptide-1 (GLP-1) or other agonists of the GLP-1 receptor, dipeptidyl peptidase IV (DPP4) inhibitors, and sodium-glucose co-transporter 2 (SGLT2) inhibitors (e.g., dapagliflozin, canagliflozin, and LX-4211).

5 5.5. PHARMACEUTICAL COMPOSITIONS

Pharmaceutical formulations may be adapted for administration by any appropriate route, for example by the oral (including buccal or sublingual), rectal, nasal, topical (including buccal, sublingual, or transdermal), vaginal, or parenteral (including subcutaneous, intracutaneous, intramuscular, intra-articular, intrasynovial, intrasternal, intrathecal, intralesional, intravenous, or intradermal injections or infusions) route. Such formulations may be prepared by any method known in the art of pharmacy, for example by bringing into association the active ingredient with the carrier(s) or excipient(s). Oral administration or administration by injection are preferred.

Pharmaceutical formulations adapted for oral administration may be presented as discrete units such as capsules or tablets; powders or granules; solutions or suspensions in aqueous or non-
15 aqueous liquids; edible foams or whips; or oil-in-water liquid emulsions or water-in-oil emulsions.

For instance, for oral administration in the form of a tablet or capsule, the active drug component can be combined with an oral, non-toxic pharmaceutically acceptable inert carrier such as ethanol, glycerol, water, and the like. Powders are prepared by comminuting the compound to a suitable fine size and mixing with a similarly comminuted pharmaceutical carrier such as an edible
20 carbohydrate, as, for example, starch or mannitol. Flavoring, preservative, dispersing, and coloring agent can also be present.

Capsules are made by preparing a powder mixture, as described above, and filling formed gelatin sheaths. Glidants and lubricants such as colloidal silica, talc, magnesium stearate, calcium stearate, or solid polyethylene glycol can be added to the powder mixture before the filling operation.
25 A disintegrating or solubilizing agent such as agar-agar, calcium carbonate, or sodium carbonate can also be added to improve the availability of the medicament when the capsule is ingested.

Moreover, when desired or necessary, suitable binders, lubricants, disintegrating agents, and coloring agents can also be incorporated into the mixture. Suitable binders include starch, gelatin, natural sugars such as glucose or beta-lactose, corn sweeteners, natural and synthetic gums such as
30 acacia, tragacanth or sodium alginate, carboxymethylcellulose, polyethylene glycol, and the like. Lubricants used in these dosage forms include sodium oleate, sodium chloride, and the like. Disintegrators include, without limitation, starch, methyl cellulose, agar, betonite, xanthan gum, and the like. Tablets are formulated, for example, by preparing a powder mixture, granulating or slugging, adding a lubricant and disintegrant, and pressing into tablets. A powder mixture is prepared by
35 mixing the compound, suitable comminuted, with a diluent or base as described above, and optionally, with a binder such as carboxymethylcellulose, an aliginate, gelating, or polyvinyl

pyrrolidone, a solution retardant such as paraffin, a resorption accelerator such as a quaternary salt and/or and absorption agent such as betonite, kaolin, or dicalcium phosphate. The powder mixture can be granulated by wetting with a binder such as syrup, starch paste, acacia mucilage, or solutions of cellulosic or polymeric materials and forcing through a screen. As an alternative to granulating, the powder mixture can be run through the tablet machine and the result is imperfectly formed slugs broken into granules. The granules can be lubricated to prevent sticking to the tablet forming dies by means of the addition of stearic acid, a stearate salt, talc, or mineral oil. The lubricated mixture is then compressed into tablets. The compounds of the present disclosure can also be combined with a free flowing inert carrier and compressed into tablets directly without going through the granulating or slugging steps. A clear or opaque protective coating consisting of a sealing coat of shellac, a coating of sugar or polymeric material, and a polish coating of wax can be provided. Dyestuffs can be added to these coatings to distinguish different unit dosages.

Oral fluids such as solution, syrups, and elixirs can be prepared in dosage unit form so that a given quantity contains a predetermined amount of the compound. Syrups can be prepared by dissolving the compound in a suitably flavored aqueous solution, while elixirs are prepared through the use of a non-toxic vehicle. Solubilizers and emulsifiers such as ethoxylated isostearyl alcohols and polyoxyethylene sorbitol ethers, preservatives, flavor additive such as peppermint oil or natural sweeteners, or saccharin or other artificial sweeteners, and the like can also be added.

Where appropriate, dosage unit formulations for oral administration can be microencapsulated. The formulation can also be prepared to prolong or sustain the release as for example by coating or embedding particulate material in polymers, wax, or the like.

The compounds of Formula (I), and pharmaceutically acceptable salts thereof, can also be administered in the form of liposome delivery systems, such as small unilamellar vesicles, large unilamellar vesicles, and multilamellar vesicles. Liposomes can be formed from a variety of phospholipids, such as cholesterol, stearylamine, or phosphatidylcholines.

The compounds of Formula (I) and pharmaceutically acceptable salts thereof may also be delivered by the use of monoclonal antibodies as individual carriers to which the compound molecules are coupled. The compounds may also be coupled with soluble polymers as targetable drug carriers. Such polymers can include polyvinylpyrrolidone, pyran copolymer, polyhydroxypropylmethacrylamidephenol, polyhydroxyethylaspartamidephenol, or polyethyleneoxidepolylysine substituted with palitoyl residues. Furthermore, the compounds may be coupled to a class of biodegradable polymers useful in achieving controlled release of a drug, for example, polylactic acid, polyepsilon caprolactone, polyhydroxy butyric acid, polyorthoesters, polyacetals, polydihydropyrans, polycyanoacrylates, and cross-linked or amphipathic block copolymers of hydrogels.

Pharmaceutical formulations adapted for transdermal administration may be presented as discrete patches intended to remain in intimate contact with the epidermis of the recipient for a

prolonged period of time. For example, the active ingredient may be delivered from the patch by iontophoresis as generally described in *Pharmaceutical Research* 1986, 3(6), 318.

Pharmaceutical formulations adapted for topical administration may be formulated as ointments, creams, suspensions, lotions, powders, solutions, pastes, gels, sprays, aerosols, or oils.

5 Pharmaceutical formulations adapted for rectal administration may be presented as suppositories or as enemas.

Pharmaceutical formulations adapted for nasal administration wherein the carrier is a solid include a coarse powder having a particle size for example in the range 20 to 500 microns which is administered in the manner in which snuff is taken, *i.e.*, by rapid inhalation through the nasal
10 passage from a container of the powder held close up to the nose. Suitable formulations wherein the carrier is a liquid, for administration as a nasal spray or nasal drops, include aqueous or oil solutions of the active ingredient.

Pharmaceutical formulations adapted for administration by inhalation include fine particle dusts or mists, which may be generated by means of various types of metered, dose pressurized
15 aerosols, nebulizers, or insufflators.

Pharmaceutical formulations adapted for vaginal administration may be presented as pessaries, tampons, creams, gels, pastes, foams, or spray formulations.

Pharmaceutical formulations adapted for parenteral administration include aqueous and non-aqueous sterile injection solutions which may contain anti-oxidants, buffers, bacteriostats, and
20 soutes which render the formulation isotonic with the blood of the intended recipient; and aqueous and non-aqueous sterile suspensions which may include suspending agents and thickening agents. The formulations may be presented in unit-dose or multi-dose containers, for example sealed ampoules and vials, and may be stored in a freeze-dried (lyophilized) condition requiring only the addition of the sterile liquid carrier, for example water for injections, immediately prior to use.
25 Extemporaneous injection solutions and suspensions may be prepared from sterile powders, granules, and tablets.

It should be understood that in addition to the ingredients particularly mentioned above, the formulations may include other agents conventional in the art having regard to the type of
30 agents.

5.6. EXAMPLES

Certain aspects of the invention can be understood from the following examples.

5.6.1. AAK1 Knockout Mice

Mice homozygous (-/-) for the disruption of the AAK1 gene were prepared by two methods;
35 gene trapping and homologous recombination.

Gene trapping is a method of random insertional mutagenesis that uses a fragment of DNA coding for a reporter or selectable marker gene as a mutagen. Gene trap vectors have been designed to integrate into introns or genes in a manner that allows the cellular splicing machinery to splice vector encoded exons to cellular mRNAs. Commonly, gene trap vectors contain selectable
5 marker sequences that are preceded by strong splice acceptor sequences and are not preceded by a promoter. Thus, when such vectors integrate into a gene, the cellular splicing machinery splices exons from the trapped gene onto the 5' end of the selectable marker sequence. Typically, such selectable marker genes can only be expressed if the vector encoding the gene has integrated into an intron. The resulting gene trap events are subsequently identified by selecting for cells that can
10 survive selective culture.

Embryonic stem cells (Lex-1 cells from derived murine strain A129), were mutated by a process involving the insertion of at least a portion of a genetically engineered vector sequence into the gene of interest, the mutated embryonic stem cells were microinjected into blastocysts which were subsequently introduced into pseudopregnant female hosts and carried to term using
15 established methods. See, e.g., "Mouse Mutagenesis", 1998, Zambrowicz et al., eds., Lexicon Press, The Woodlands, TX. The resulting chimeric animals were subsequently bred to produce offspring capable of germline transmission of an allele containing the engineered mutation in the gene of interest.

AAK1-gene disrupted mice were also made by homologous recombination. In this case, the
20 second coding exon of the murine AAK1 gene (see GenBank Accession Number NM_177762) was removed by methods known in the art. See, e.g., U.S. Patent Nos. 5,487,992, 5,627,059, and 5,789,215.


Mice homozygous (-/-) for the disruption of the AAK1 gene were studied in conjunction with mice heterozygous (+/-) for the disruption of the AAK1 gene, and wild-type (+/+) litter mates. During
25 this analysis, the mice were subject to a medical work-up using an integrated suite of medical diagnostic procedures designed to assess the function of the major organ systems in a mammalian subject. Homozygous (-/-) "knockout" mice were studied in conjunction with their heterozygous (+/-) and wild-type (+/+) litter mates. Disruption of the AAK1 gene was confirmed by Southern analysis. Expression of the murine homolog of AAK1 was detected by RT-PCR in murine brain; spinal cord; eye;
30 thymus; spleen; lung; kidney; liver; skeletal muscle; bone; stomach, small intestine and colon; heart; adipose; asthmatic lung; LPS liver; blood; banded heart; aortic tree; prostate; and mammary gland (5 week virgin, mature virgin, 12 DPC, 3 day post-partum (lactating), 3 day post-weaning (early involution), and 7 day post-weaning (late involution)).

AAK1 homozygous (-/-) and their wild-type (+/+) littermates were tested using the formalin
35 paw test in order to assess their acute and tonic nociceptive responses. For these tests, Automatic Nociception Analyzers (purchased from the Ozaki lab at University of California, San Diego) were used. A metal band was placed around the left hind paw of each mouse 30 minutes prior to testing.


After the 30-minute acclimation period, 20 μ l of 5% formalin is subcutaneously injected in the dorsal surface of the left hind paw. Mice were individually housed in cylindrical chambers for 45 minutes. A computer recorded flinches per minute, total flinches for phase I (acute phase = first 8 minutes), and total flinches for phase II (tonic phase = time between minutes 20 - 40) through an electromagnetic field. See Yaksh TL, Ozaki G, McCumber D, Rathbun M, Svensson C, Malkmus S, Yaksh MC. *An automated flinch detecting system for use in the formalin nociceptive bioassay.* J Appl Physiol., 2001; 90:2386-402.

As shown in Figure 1, phase 1 and phase 2 data were obtained using homozygous (-/-) mice females (n = 16), wild-type females (n = 15), homozygous (-/-) mice males (n = 9), and wild-type males (n = 18). In all groups and in both phases, the AAK1 homozygous (-/-) mice exhibited significantly less recorded paw flinching than their wild-type (+/+) littermates.

5.6.2. Synthesis of [3-(4-Aminomethyl-phenyl)-pyrazolo[1,5-a]pyrimidin-5-yl]-butyl-amine




Part A. Butyl-pyrazolo[1,5-a]pyrimidin-5-yl-amine



A solution of 5-chloro-pyrazolo[1,5-a]pyrimidine [29274-24-] (955.4 mg, 6.2 mmol) in butylamine [109-73-9] (6.2 mL, 62.7 mmol), under N₂ blanket, was magnetically stirred at 65 °C for 17 h then partitioned between brine (pH adjusted to 10 with saturated aqueous sodium bicarbonate) and ethyl acetate. The phase separated extract was dried (MgSO₄) and diluted with heptane to precipitate butyl-pyrazolo[1,5-a]pyrimidin-5-yl-amine as 1.2 g of yellow powder, mp. 75-76 °C. ¹H NMR (400 MHz, DMSO-*d*₆) δ ppm 0.91 (t, *J*=7.33 Hz, 3 H) 1.37 (dq, *J*=15.00, 7.29 Hz, 2 H) 1.47 - 1.59 (m, 2 H) 3.24 - 3.34 (m, 2 H) 5.94 (d, *J*=1.52 Hz, 1 H) 6.22 (d, *J*=7.58 Hz, 1 H) 7.35 (br. s., 1 H) 7.74 (d, *J*=2.27 Hz, 1 H) 8.40 (d, *J*=7.58 Hz, 1 H). ¹³C NMR (100 MHz, DMSO-*d*₆) δ ppm 13.69, 19.71, 30.65, 91.00, 99.95, 134.70, 143.45, 148.37, 155.43. LRMS (ESI) *m/z* 191.1 [(M+H)]⁺, calc'd for C₁₀H₁₄N₄: 190.25


Part B. (3-Bromo-pyrazolo[1,5-a]pyrimidin-5-yl)-butyl-amine



Sodium acetate [127-09-3] (0.7 g, 8.9 mmol) was added to a solution of butyl-pyrazolo[1,5-a]pyrimidin-5-yl-amine (1.1 g, 5.9 mmol) in glacial acetic acid (100 mL) and allowed to stir at ambient temperature until all of the solid had dissolved. Bromine [7726-95-6] (0.3 mL, 6.4 mmol) was added drop by drop into the ambient temperature, buffered, acetic acid reaction solution over 5 minutes.

- 5 Upon completion of the bromine addition, the suspension was allowed to stir for a further 15 minutes then was slowly poured into 500 mL of stirred water. Saturated aqueous sodium bicarbonate was added to the stirred suspension until pH of the supernatant was determined to be approximately 8 by pH paper, then the suspension was extracted with ethyl acetate. The extract was dried (MgSO₄) and flash chromatographed (silica gel eluted with 60% (v/v) ethyl acetate / heptane) to provide 1.2 g of
 10 off white solid, mp. 94-95 °C. ¹H NMR (400 MHz, DMSO-*d*₆) δ ppm 0.93 (t, *J*=7.33 Hz, 3 H) 1.31 - 1.44 (m, 2 H) 1.56 (quin, *J*=7.20 Hz, 2 H) 3.31 - 3.38 (m, 2 H) 6.28 (d, *J*=7.58 Hz, 1 H) 7.61 (t, *J*=5.05 Hz, 1 H) 7.85 (s, 1 H) 8.42 (d, *J*=7.58 Hz, 1 H). ¹³C NMR (100 MHz, DMSO-*d*₆) δ ppm 13.67, 19.64, 30.53, 76.99, 100.87, 135.14, 143.03, 145.13, 156.09. LRMS (ESI) *m/z* 269.0/271.0 [(*M*+*H*)]⁺, calc'd for C₁₀H₁₃BrN₄: 269.15.


15 Part C. [3-(4-Aminomethyl-phenyl)-pyrazolo[1,5-a]pyrimidin-5-yl]-butyl-amine



- To a mixture of (3-bromo-pyrazolo[1,5-a]pyrimidin-5-yl)-butyl-amine (350.3 mg, 1.3 mmol), 4-aminomethylphenylboronic acid, hydrochloride [75705-21-4] (337.1 mg, 1.8 mmol), potassium phosphate tribasic monohydrate [27176-10-9] (601.6 mg, 2.6 mmol), and [1,1'-bis(diphenylphosphino) ferrocene] dichloropalladium(II), complex with dichloromethane [95464-05-4] (117.2 mg, 0.1 mmol) contained in a 50 mL round bottomed flask was added a solution of 30% (v/v) water in 1,2-dimethoxyethane (25 mL) and a magnetic stir bar. The reaction pot was fitted to a reflux condenser, lowered into an ambient temperature oil bath, and the system taken through 10 evacuation / N₂ blanket cycles while being rapidly stirred. The rapidly stirred, N₂ blanketed, reaction
 20 was heated to an oil bath temperature of 85 °C for 17 h then cooled and partitioned between brine and ethyl acetate. The phase separated extract was dried (MgSO₄) and evaporated to afford a brown oil which was purified by preparative RP-HPLC to yield 71.9 mg of [3-(4-Aminomethyl-phenyl)-pyrazolo[1,5-a]pyrimidin-5-yl]-butyl-amine monoacetate salt as a white solid, mp. 171-172 °C. ¹H NMR (400 MHz, DMSO-*d*₆) δ ppm 0.97 (t, *J*=7.39 Hz, 3 H) 1.43 (sxt, *J*=7.36 Hz, 2 H) 1.64 (quin, *J*=7.17 Hz, 2 H) 3.43 (q, *J*=6.47 Hz, 2 H) 3.71 - 3.84 (m, 2 H) 6.29 (d, *J*=7.50 Hz, 1 H) 7.34 (m, *J*=8.16 Hz, 2 H) 7.62 - 7.71 (m, 1 H) 8.05 (m, *J*=7.94 Hz, 2 H) 8.32 (s, 1 H) 8.46 (d, *J*=7.50 Hz, 1 H).
 30 ¹³C NMR (100 MHz, DMSO-*d*₆) δ ppm 13.03, 14.11, 19.63, 29.11, 30.33, 44.34, 45.62, 103.75,


124.02, 124.14, 126.95, 127.86, 131.85, 135.07, 137.50, 141.15, 141.36, 144.46, 155.50, 172.52. LRMS (ESI) m/z 296.1 [(M+H)]⁺, calc'd for C₁₇H₂₁N₅: 295.39.

5.6.3. Synthesis of (2-Methoxy-ethyl)-{3-[4-(1H-tetrazol-5-yl)-phenyl]-pyrazolo[1,5-a]pyrimidin-5-yl}-amine



5


Part A. N-(2-methoxyethyl)pyrazolo[1,5-a]pyrimidin-5-amine



A stirred mixture of 5-chloro-pyrazolo[1,5-a]pyrimidine [29274-24-6], (525.5 mg, 3.4 mmol) and 2-methoxy-ethylamine (3.0 mL, 34.5 mmol) was heated to 65°C for 17 h, cooled and partitioned between brine (pH adjusted to 8 with saturated aqueous sodium bicarbonate) and ethyl acetate. Extract was dried (MgSO₄) and evaporated to yield 706.4 mg of light yellow solid. LRMS (ESI) m/z 193.1[(M+H)]⁺, calc'd for C₉H₁₂N₄O: 192.22.

10


Part B. (3-Bromo-pyrazolo[1,5-a]pyrimidin-5-yl)-(2-methoxy-ethyl)-amine



Sodium acetate [127-09-3] (457.6 mg, 5.6 mmol) was added to a solution of N-(2-methoxyethyl)pyrazolo[1,5-a]pyrimidin-5-amine (706.4 mg, 3.7 mmol) in glacial acetic acid (40 mL) and allowed to stir at ambient temperature until all of the solid had dissolved. Bromine [7726-95-6] (0.2 mL, 3.7 mmol) was added into the ambient temperature, buffered acetic acid reaction solution over 2 minutes, then the reaction solution was slowly poured into stirring water (1L). The pH was increased by the addition of solid sodium bicarbonate until the pH of the supernatant was determined to be approximately 8 by pH paper. Precipitated product was isolated by filtration, the filter cake washed with water, and allowed to dry to afford 819.6 mg of white powder. LRMS (ESI) m/z 271.0/273.0 [(M+H)]⁺, calc'd for C₉H₁₁BrN₄O: 271.12.


15


20

Part C. (2-Methoxy-ethyl)-{3-[4-(1H-tetrazol-5-yl)-phenyl]-pyrazolo[1,5-a]pyrimidin-5-yl}-amine


To a mixture of (3-bromo-pyrazolo[1,5-a]pyrimidin-5-yl)-(2-methoxy-ethyl)-amine (307.7 mg, 1.1 mmol), (4-(1H-tetrazol-5-yl)phenyl)boronic acid [179942-55-3] (259.6 mg, 1.4 mmol), potassium phosphate tribasic monohydrate [27176-10-9] (522.4 mg, 2.3 mmol), and [1,1'-bis(diphenylphosphino)ferrocene] dichloro- palladium(II), complex with dichloromethane [95464-05-4] (106.3 mg, 0.1 mmol) contained in a 50 mL round bottomed flask was added a solution of 30% (v/v) water in 1,2-dimethoxyethane (12 mL) and a magnetic stir bar. The reaction pot was fitted to a reflux condenser, lowered into an ambient temperature oil bath, and the system taken through 10 evacuation / N₂ blanket cycles while being rapidly stirred. The rapidly stirred, N₂ blanketed, reaction was heated to an oil bath temperature of 85 °C for 4 h then cooled, diluted with methanol, and gravity filtered. The filtrate was preabsorbed on silica gel and flash chromatographed (silica gel eluted with 1% NH₄OH in 10% (v/v) methanol / ethyl acetate). Desired chromatography fractions were combined, evaporated, redissolved in 2-propanol and gravity filtered into stirring heptane to precipitate purified product. (2-Methoxy-ethyl)-{3-[4-(1H-tetrazol-5-yl)-phenyl]-pyrazolo[1,5-a]pyrimidin-5-yl}-amine was isolated by filtration as 96.3 mg of solid material, mp. 264-265 °C (dec.). ¹H NMR (400 MHz, DMSO-*d*₆) δ ppm 3.63 (t, *J*=4.29 Hz, 5 H) 6.41 (d, *J*=7.58 Hz, 1 H) 7.88 (br. s., 1 H) 8.04 (m, *J*=8.59 Hz, 2 H) 8.31 (m, *J*=8.34 Hz, 2 H) 8.47 (s, 1 H) 8.53 (d, *J*=7.58 Hz, 1 H). ¹³C NMR (100 MHz, DMSO-*d*₆) δ ppm 40.69, 58.48, 70.48, 100.86, 103.75, 119.86, 125.23, 127.61, 135.97, 137.08, 142.42, 145.63, 156.60, LRMS (ESI) *m/z* 337.2 [(*M*+*H*)⁺, calc'd for C₁₆H₁₆N₈O: 336.36.

5.6.4. Synthesis of Methyl-[2-(3-thiophen-2-yl-pyrazolo[1,5-a]pyrimidin-5-ylamino)-ethyl]-carbamic acid tert-butyl ester




Part A. 3-Oxo-2-thiophen-2-yl-propionitrile

Thiophen-2-yl-acetonitrile [20893-30-5] (6.7 mL, 62.6 mmol) was added to a N₂ blanketed, 0°C, stirred solution of 21wt% sodium ethoxide (25.7 mL, 68.9 mmol) in ethanol (100 mL). After 5 minutes, ethyl formate [109-94-4] (12.6 mL, 156.7 mmol) was added and the resultant solution was stirred for 20 minutes. The ice bath was then removed and replaced with a heating mantle. The reaction was heated to reflux for 2 h, cooled, and evaporated to provide a yellow solid. Crude product was dissolved in water, the solution chilled to 0°C and acidified with aqueous hydrochloric acid to precipitate an off white powder. Product was isolated by filtration, washed with water, and dried to afford 7.4 g. LRMS (ESI) m/z 152.0 [(M+H)]⁺, calc'd for C₇H₅NOS: 151.19

Part B. 4-Thiophen-2-yl-2H-pyrazol-3-ylamine


Hydrazine hydrate [7803-57-8] (4.8mL, 99.0mmol) then acetic acid [64-19-7] (5.6 mL, 97.8 mmol) were added to a solution of 3-oxo-2-thiophen-2-yl-propionitrile (7.4 g, 49.0 mmol) in ethanol (100 mL). The solution was then heated to reflux for 2 h. The cooled reaction was evaporated to dryness and the resultant solid triturated in water. Product was isolated by filtration, washed with water, and dried to provide 4-thiophen-2-yl-2H-pyrazol-3-ylamine as 7.1 g of fine off white powder, mp. 183-184°C. ¹H NMR (400 MHz, DMSO-*d*₆) δ ppm 2.51 (s, 1 H) 7.01 - 7.10 (m, 2 H) 7.27 (dd, *J*=5.05, 1.01 Hz, 1 H). ¹³C NMR (100 MHz, DMSO-*d*₆) δ ppm 121.07, 121.47, 127.58, 135.85. LRMS (ESI) m/z 166.0[(M+H)]⁺, calc'd for C₇H₇N₃S: 165.22.

Part C. 3-Thiophen-2-yl-4H-pyrazolo[1,5-a]pyrimidin-5-one

A 21wt% solution of sodium ethoxide in ethanol (32.7 mL, 87.6 mmol) was slowly added to an ambient temperature, stirred suspension of 4-thiophen-2-yl-2H-pyrazol-3-ylamine (4.1 g, 25.0 mmol) and 1,3-dimethylpyrimidine-2,4(1H,3H)-dione [874-14-6] (3.9 g, 27.5 mmol) in ethanol (217 mL). As addition progressed the solid dissolved to give a clear brown solution. The solution was heated to reflux for 2 d, cooled, then poured into 5% (w/v) aqueous ammonium chloride. Ethanol was


evaporated out and the reduced volume product solution chilled, with stirring, in an ice bath to precipitate 3-thiophen-2-yl-4H-pyrazolo[1,5-a]pyrimidin-5-one as 3.2 g of red-brown fine crystalline powder, mp. 231-232°C. ¹H NMR (400 MHz, DMSO-*d*₆) δ ppm 6.13 (br. s., 1 H) 7.10 (dd, *J*=5.05, 3.54 Hz, 1 H) 7.38 - 7.46 (m, 2 H) 8.07 (br. s., 1 H) 8.51 - 8.73 (m, 1 H) 12.06 (br. s., 1 H). ¹³C NMR (100 MHz, DMSO-*d*₆) δ ppm 123.66, 127.71, 138.70, 142.47. LRMS (ESI) *m/z* 218.0[(M+H)]⁺, calc'd for C₁₀H₇N₃OS: 217.25.


Part D. Methyl-[2-(3-thiophen-2-yl-pyrazolo[1,5-a]pyrimidin-5-ylamino)-ethyl]-carbamic acid tert-butyl ester




1,8-Diazabicycloundec-7-ene (DBU) [6674-22-2] (0.4 mL, 2.5 mmol) was added to an ambient temperature, stirred, turbid solution of 3-thiophen-2-yl-4H-pyrazolo[1,5-a]pyrimidin-5-one (354.3 mg, 1.6 mmol), (2-amino-ethyl)-methyl-carbamic acid tert-butyl ester [121492-06-6] (852.2 mg, 4.9 mmol), and (benzotriazol-1-yloxy)tris(dimethylamino)phosphonium hexafluorophosphate (BOP) [56602-33-6] (937.7 mg, 2.1 mmol) in acetonitrile (16 mL). Upon addition of the DBU the turbidity clears and the solution was permitted to stir for 2 d then partitioned between brine and ethyl acetate. The extract was dried (MgSO₄) and flash chromatographed (silica gel, eluted with 10% (v/v) methanol / ethyl acetate). Evaporation of product fractions yielded an oil which was crystallized from heptane to provide methyl-[2-(3-thiophen-2-yl-pyrazolo[1,5-a]pyrimidin-5-ylamino)-ethyl]-carbamic acid tert-butyl ester as 333.6 mg of yellow, fine crystalline powder, mp. 140-141°C. ¹H NMR (400 MHz, DMSO-*d*₆) δ ppm (rotomers present) 1.16 - 1.28 (m, 5 H) 1.37 (br. s., 3 H) 2.51 (s, 1 H) 2.85 (br. s., 3 H) 3.47 (br. s., 1 H) 3.55 (br. s., 3 H) 6.28 (d, *J*=6.82 Hz, 1 H) 7.04 (dd, *J*=5.05, 3.54 Hz, 1 H) 7.30 (dd, *J*=5.05, 1.01 Hz, 1 H) 7.41 (br. s., 1 H) 7.80 (br. s., 1 H) 8.19 (s, 1 H) 8.48 (d, *J*=7.07 Hz, 1 H). ¹³C NMR (100 MHz, DMSO-*d*₆) δ ppm 27.81, 27.96, 33.75, 38.41, 46.32, 78.13, 100.28, 100.76, 120.16, 121.75, 126.90, 135.01, 135.20, 140.78, 143.74, 154.91, 155.81. LRMS (ESI) *m/z* 374.1[(M+H)]⁺, calc'd for C₁₈H₂₃N₅O₂S: 373.48.

5.6.5. Synthesis of Isopropyl (2-((3-(3-methoxypyridin-4-yl)pyrazolo[1,5-a]pyrimidin-5-yl)amino)ethyl)(methyl)-carbamate




Part A. 3-Bromo-5-chloro-pyrazolo[1,5-a]pyrimidine

Sodium acetate [127-09-3] (24.1 g, 293.9 mmol) was added to a solution of 5-chloro-pyrazolo[1,5-a]pyrimidine [29274-24-6], (30.1 g, 195.7 mmol) in glacial acetic acid (395 mL) and
 5 allowed to stir at ambient temperature until all of the solid had dissolved. An ice bath was then put in place, and the solution chilled to 0°C. Bromine [7726-95-6] (11.0 mL, 214.7 mmol) was added drop by drop into the 0°C buffered acetic acid reaction solution over 35 minutes which gradually became a thick stirred suspension. Upon completion of the bromine addition, the suspension was allowed to stir for a further 15 minutes then was slowly poured into an aqueous (5% w/v) solution (1 L) of
 10 sodium metabisulfite [7681-57-4] stirred in an ice bath. Solid sodium hydroxide [1310-73-2] pellets were added to the stirred suspension, at a rate to maintain an internal temperature of less than 40°C, until pH of the supernatant was determined to be approximately 8 by pH paper. Precipitated product was isolated by filtration, the filter cake washed with water, and allowed to dry to afford 39.8 g of light yellow powder, mp. 173-174°C. ¹H NMR (400 MHz, DMSO-*d*₆) δ ppm 7.22 (d, *J*=7.33 Hz, 1
 15 H) 8.44 (s, 1 H) 9.21 (d, *J*=7.07 Hz, 1 H). ¹³C NMR (100 MHz, DMSO-*d*₆) δ ppm 82.84, 109.99, 138.94, 143.76, 145.61, 151.18. LRMS (ESI) *m/z* 231.8/233.8/235.8 [(*M*+*H*)]⁺, calc'd for C₆H₃BrClN₃: 232.47.

Part B. tert-butyl (2-((3-bromopyrazolo[1,5-a]pyrimidin-5-yl)amino)ethyl)(methyl)carbamate


20 To 465 mg (2.00 mmol) of 3-bromo-5-chloropyrazolo[1,5-a]pyrimidine was added tert-butyl (2-aminoethyl)(methyl)carbamate (522 mg, 3.00 mmol), 4 mL of isopropanol, and triethylamine (0.56 mL, 4.00 mmol). The resulting mixture is heated at 140°C for 0.5 hr in the microwave. It was diluted with EtOAc, and washed with water and brine. The organic layer was dried over MgSO₄, concentrated and purified on the silica gel column eluting with 20 - 100% EtOAc/Hex to obtain 607
 25 mg (82%) of the desired product.

Part C. N1-(3-bromopyrazolo[1,5-a]pyrimidin-5-yl)-N2-methylethane-1,2-diamine

The tert-butoxycarbonyl was removed by dissolving 600 mg (1.62 mmol) of tert-butyl (2-((3-bromopyrazolo[1,5-a]pyrimidin-5-yl)amino)ethyl)(methyl)carbamate in 4 mL of DCM, and then adding


4 mL TFA, stirring for 0.5 hr at rt, and concentrating the mixture to afford the TFA-salt of the desired product in 100% yield (622 mg).

Part D. Isopropyl (2-((3-bromopyrazolo[1,5-a]pyrimidin-5-yl)amino)ethyl)(methyl)carbamate



- 5 To 1.0 equivalent (384 mg, 1.000 mmol) of the amine (TFA salt) dissolved 20 mL of DCM was added 3.0 equivalents (368 mg, 3.000 mmol) of isopropylchloroformate, 4.0 equivalents triethylamine (404 mg, 4.000 mmol). The resulting mixture was stirred at rt for 0.5 hr, and then diluted with DCM (30 mL) washed with water and brine, dried over MgSO₄, and concentrated. The crude residue was purified in the silica gel (ISCO) eluting with 1-10% MeOH/DCM to obtain the
10 desired product (87% yield).


Part E. Isopropyl (2-((3-(3-methoxypyridin-4-yl)pyrazolo[1,5-a]pyrimidin-5-yl)amino)ethyl)(methyl)-carbamate



- To 80 mg (0.22 mmol) of isopropyl (2-((3-bromopyrazolo[1,5-a]pyrimidin-5-yl)amino)ethyl)(methyl)-carbamate was added (3-methoxypyridin-4-yl)boronic acid dihydrate (56 mg, 0.27 mmol), K₃PO₄ (143 mg, 0.67 mmol), PdCl₂(PPh₃)₂ (16.8 mg, 0.022 mmol), 3 mL of DME and 1 mL of water. This mixture was microwaved at 140 °C for 0.5 hr. After cooling, it was diluted with EtOAc, and washed with water and brine. The organic layer was dried over MgSO₄, concentrated, and purified on the PREP HPLC to afford the desired product in 77% yield (66.4 mg). ¹H NMR (400 MHz, Acetone) δ ppm 2.87 (br. s., 3 H) 2.99 (br. s., 3 H) 3.58 - 3.71 (m, 2 H) 3.73 - 3.82 (m, 2 H) 4.09 (s, 3 H) 4.82 (td, J=6.25, 2.91 Hz, 1 H) 6.45 (br. s., 1 H) 8.21 (br. s., 1 H) 8.35 (s, 1 H) 8.43 (d, J=6.06 Hz, 1 H) 8.58 (s, 1 H) 8.77 (br. s., 1 H); LRMS (ESI) m/e 385.0 [(M + H)⁺, calcd for C₁₉H₂₄N₆O₃ 384.0].
20


5.6.6. Synthesis of (S)-tert-butyl 2-(((3-(4-(aminomethyl)phenyl)pyrazolo[1,5-a]pyrimidin-5-yl)amino)methyl)pyrrolidine-1-carboxylate

- 25 Part A. (S)-tert-butyl 2-(((3-bromopyrazolo[1,5-a]pyrimidin-5-yl)amino)methyl)pyrrolidine-1-carboxylate




The reaction of 3-bromo-5-chloropyrazolo[1,5-a]pyrimidine and (S)-tert-butyl 2-(aminomethyl)pyrrolidine-1-carboxylate under the amine displacement conditions described in example 5.6.5 afforded 74% product.

Part B. (S)-tert-butyl 2-(((3-(4-(aminomethyl)phenyl)pyrazolo[1,5-a]pyrimidin-5-yl)amino)methyl)pyrrolidine-1-carboxylate




The Suzuki coupling of (S)-tert-butyl 2-(((3-bromopyrazolo[1,5-a]pyrimidin-5-yl)amino)methyl)pyrrolidine-1-carboxylate with (4-(aminomethyl)phenyl)boronic acid under the same conditions as described in example 5.6.5 to give 73% of the titled compound. ¹H NMR (400 MHz, METHANOL-*d*₄) δ ppm 1.44 (br. s., 9 H) 1.90 (dd, *J*=9.92, 5.51 Hz, 1 H) 1.96 - 2.11 (m, 3 H) 3.35 - 3.45 (m, 4 H) 4.07 - 4.21 (m, 3 H) 6.33 (br. s., 1 H) 7.46 (d, *J*=8.38 Hz, 2 H) 8.19 (br. s., 2 H) 8.25 - 8.36 (m, 2 H); LRMS (ESI) *m/e* 423.0 [(*M* + *H*)⁺, calcd for C₂₃H₃₀N₆O₂ 422.0].

5.6.7. Synthesis of (S)-tert-butyl 2-(((3-(2-methoxyphenyl)pyrazolo[1,5-a]pyrimidin-5-yl)amino)methyl)pyrrolidine-1-carboxylate




The Suzuki coupling of (S)-tert-butyl 2-(((3-bromopyrazolo[1,5-a]pyrimidin-5-yl)amino)methyl)pyrrolidine-1-carboxylate with (2-methoxyphenyl)boronic acid under the same conditions as described in example 5.6.5 to give 68% of the titled compound. ¹H NMR (400 MHz, CHLOROFORM-*d*) δ ppm 1.50 (s, 9 H) 1.85 - 2.12 (m, 4 H) 3.40 (dd, *J*=7.58, 4.04 Hz, 2 H) 3.46 - 3.59 (m, 1 H) 3.59 - 3.67 (m, 1 H) 3.94 (s, 3 H) 4.19 (br. s., 1 H) 6.07 (d, *J*=7.07 Hz, 1 H) 6.99 (d, *J*=7.58 Hz, 1 H) 7.06 (td, *J*=7.45, 1.01 Hz, 1 H) 7.15 - 7.22 (m, 1 H) 8.14 - 8.28 (m, 1 H) 8.45 (d, *J*=6.82 Hz, 1 H) 8.51 (s, 1 H); LRMS (ESI) *m/e* 424.0 [(*M* + *H*)⁺, calcd for C₂₃H₂₉N₅O₃ 423.0].

5.6.8. Synthesis of (S)-tert-butyl 2-(((3-(trifluoromethyl)pyrazolo[1,5-a]pyrimidin-5-yl)amino)methyl)pyrrolidine-1-carboxylate



Part A. (S)-tert-butyl 2-((pyrazolo[1,5-a]pyrimidin-5-ylamino)methyl)pyrrolidine-1-carboxylate




5

The commercially available 5-chloropyrazolo[1,5-a]pyrimidine is reacted with (S)-tert-butyl 2-(aminomethyl)pyrrolidine-1-carboxylate under the amine displacement conditions described in example 5.6.5 to give 96 % product.

Part B. (S)-tert-butyl 2-(((3-iodopyrazolo[1,5-a]pyrimidin-5-yl)amino)methyl)pyrrolidine-1-carboxylate


10



The (S)-tert-butyl 2-((pyrazolo[1,5-a]pyrimidin-5-ylamino)methyl)pyrrolidine-1-carboxylate (790 mg, 2.490 mmol) was dissolved in 10 mL of DMF, was added NIS (617 mg, 2.740 mmol). This mixture is stirred for 4 hr at rt, and then quenched with 10 mL of water. It was then diluted with 50 mL EtOAc and the organic layer was separated, washed twice with brine, and dried over MgSO₄. It was concentrated and purified on the silica gel eluting with 20-100% EtOAc/hex to obtain 1.03 g (93%) an off-white solid.

15

Part C. (S)-isopropyl 2-(((3-(trifluoromethyl)pyrazolo[1,5-a]pyrimidin-5-yl)amino)methyl)pyrrolidine-1-carboxylate




20

To 221.5 mg (0.50 mmol) of the aryl iodide dissolved in 2 mL DMF was added methyl 2,2-difluoro-2-(fluorosulfonyl)acetate (0.77g, 4.00 mmol), CuI (762 mg, 2.00 mmol), HMPA (0.41 mL, 3.50 mmol). This mixture was stirred on an oil bath preheated to 90 °C for 0.75 hr. The reaction was cooled to rt, and quenched with NH₄Cl saturated aqueous solution. Extracted with 30 mL of EtOAc, washed with brine and dry over MgSO₄. It was concentrated and purified on the PREP HPLC to obtain


25

81 mg (42%) of the desired product. ^1H NMR (400 MHz, METHANOL- d_4) δ ppm 1.44 (s, 9 H) 1.86 - 1.99 (m, 3 H) 1.99 - 2.07 (m, 1 H) 3.40 (br. s., 3 H) 3.61 (br. s., 1 H) 4.11 (br. s., 1 H) 6.37 (br. s., 1 H) 7.98 (s, 1 H) 8.33 (br. s., 1 H); LRMS (ESI) m/e 386.0 $[(M + H)^+]$, calcd for $\text{C}_{17}\text{H}_{22}\text{F}_3\text{N}_5\text{O}_2$ 385.0].

5.6.9. Synthesis of 4-[3-(3-Methoxy-pyridin-4-yl)-pyrazolo[1,5-a]pyrimidin-5-yl]-piperazine-1-carboxylic acid isopropyl ester




Part A. 3-Bromo-5-piperazin-1-yl-pyrazolo[1,5-a]pyrimidine




3-Bromo-5-chloro-pyrazolo[1,5-a]pyrimidine (15.4 g, 66.2 mmol) and piperazine [110-85-0] (56.8 g, 659.5 mmol) were ground together in a mortar to an intimate mixture and transferred to a 250 mL round bottomed flask containing a magnetic stir bar. The flask was fitted to a reflux condenser, N_2 blanketed, and the reaction pot immersed in an ambient temperature oil bath. While the neat solid mixture stirred, the bath was heated to 120°C over 0.5 h and held at nominal temperature for a total of 2 h. The bath was removed and the molten reaction allowed to cool to approximately 75°C . Ethyl acetate was cautiously added down the condenser to dissolve the reaction product to prevent its setting up into a solid mass. The reaction solution was transferred to a separatory funnel, further diluted with ethyl acetate, and washed with water. The ethyl acetate extract was dried (MgSO_4) and evaporated to obtain 16.1 g of clear yellow oil. The oil was crystallized from methanol to provide 15.8 g of light yellow crystalline powder in two crops, mp. $108-109^\circ\text{C}$. ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ ppm 2.51 (s, 1 H) 3.16 - 3.24 (m, 3 H) 3.38 (s, 1 H) 3.93 - 4.00 (m, 3 H) 6.85 (d, $J=7.83$ Hz, 1 H) 8.03 (s, 1 H) 8.78 (d, $J=8.08$ Hz, 1 H) 9.61 (br. s., 1 H). ^{13}C NMR (100 MHz, $\text{DMSO}-d_6$) δ ppm 41.25, 42.16, 77.73, 97.78, 136.77, 144.01, 144.18, 155.53. LRMS (ESI) m/z 282.0/284.0 $[(M+H)^+]$, calc'd for $\text{C}_{10}\text{H}_{12}\text{BrN}_5$: 282.14.

Part B. 4-(3-Bromo-pyrazolo[1,5-a]pyrimidin-5-yl)-piperazine-1-carboxylic acid isopropyl ester



To a rapidly stirred, 0°C, N₂ blanketed, solution of 3-bromo-5-piperazin-1-yl-pyrazolo[1,5-a]pyrimidine (15.8 g, 56.0 mmol) and Hunig's base [7087-68-5] (20.0 mL, 114.8 mmol) in ethyl acetate (280 mL) was steadily added a 1.0M solution of isopropyl chloroformate in toluene (67.0 mL) over the course of 10 minutes. The reaction was allowed to stir and warm to ambient temperature over night at which time it was washed with brine, dried (MgSO₄), preloaded onto silica gel, chromatographed (Silica gel, eluted with 100% ethyl acetate) and crystallized from ethyl acetate / heptane to afford 13.2 g of white powder, mp. 78-79°C. ¹H NMR (400 MHz, DMSO-*d*₆) δ ppm 1.22 (d, *J*=6.32 Hz, 6 H) 3.44 - 3.54 (m, 4 H) 3.67 - 3.79 (m, 4 H) 4.82 (spt, *J*=6.23 Hz, 1 H) 6.77 (d, *J*=7.83 Hz, 1 H) 7.97 (s, 1 H) 8.69 (d, *J*=7.83 Hz, 1 H). ¹³C NMR (100 MHz, DMSO-*d*₆) δ ppm 21.94, 42.80, 43.87, 68.17, 77.32, 97.66, 136.47, 143.87, 144.40, 154.29, 155.64. LRMS (ESI) *m/z* 368.0/370.0 [(M+H)]⁺, calc'd for C₁₄H₁₈BrN₅O₂: 368.24.


Part C. 4-[3-(3-Methoxy-pyridin-4-yl)-pyrazolo[1,5-a]pyrimidin-5-yl]-piperazine-1-carboxylic acid isopropyl ester



To a mixture of 4-(3-bromo-pyrazolo[1,5-a]pyrimidin-5-yl)-piperazine-1-carboxylic acid isopropyl ester (385.0 mg, 1.1 mmol), 3-methoxypyridine-4-boronic acid hydrate [1072952-50-1] (229.7 mg, 1.34 mmol), potassium phosphate tribasic monohydrate [27176-10-9] (516.3 mg, 2.2 mmol), and [1,1'-bis(diphenylphosphino)ferrocene]dichloropalladium(II), complex with dichloromethane [95464-05-4] (102.4 mg, 0.1 mmol) contained in a 50 mL round bottomed flask was added a solution of 30% (v/v) water in 1,2-dimethoxyethane (25 mL) and a magnetic stir bar. The reaction pot was fitted to a reflux condenser, lowered into an ambient temperature oil bath, and the system taken through 10 evacuation / N₂ blanket cycles while being rapidly stirred. The rapidly stirred, N₂ blanketed, reaction was heated to an oil bath temperature of 85°C for 17 h then cooled and partitioned between brine (pH adjusted to 10 with 3N aqueous sodium hydroxide) and ethyl acetate. The phase separated extract was dried (MgSO₄) and evaporated to afford an orange oil which was purified by preparative RP-HPLC. Isolated product was dissolved in methanol, diluted with concentrated hydrochloric acid, evaporated to dryness, redissolved in methanol and precipitated by the addition of diethyl ether. Filtration of the precipitate yielded 100.5 mg of the product monohydrochloride salt as a yellow powder. mp. 206-207°C (dec.) ¹H NMR (400 MHz, DMSO-*d*₆) δ ppm 1.23 (d, *J*=6.39 Hz, 6 H) 2.51 (s, 2 H) 3.53 - 3.59 (m, 4 H) 3.85 (br. s., 3 H) 4.13 (s, 3 H) 4.84 (spt, *J*=6.25 Hz, 1 H) 6.96 (d, *J*=7.94 Hz, 1 H) 8.42 (d, *J*=6.39 Hz, 1 H) 8.48 (s, 1 H) 8.76 (s, 1 H) 8.86 (d, *J*=7.94 Hz, 1 H) 9.06 (d, *J*=6.17 Hz, 1 H). ¹³C NMR (100 MHz, DMSO-*d*₆) δ ppm 21.97,

42.79, 44.23, 57.17, 68.25, 97.56, 98.39, 120.03, 123.90, 137.27, 137.05, 137.95, 146.48, 147.00, 151.80, 154.26, 156.66. LRMS (ESI) m/z 397.1 $[(M+H)]^+$, calc'd for $C_{20}H_{24}N_6O_3$: 396.45.

5.6.10. Synthesis of 4-[3-(2-Methoxy-phenyl)-pyrazolo[1,5-a]pyrimidin-5-yl]-piperazine-1-carboxylic acid isopropyl ester




5

Prepared similarly to the preparation of example 5.6.9 from 4-(3-bromo-pyrazolo[1,5-a]pyrimidin-5-yl)-piperazine-1-carboxylic acid isopropyl ester (385.0 mg, 1.1 mmol) and 2-methoxyphenylboronic acid [5720-06-9] (191.8 mg, 1.3 mmol) to afford 76.4 mg of yellow powder as a free base, mp. 57-59°C. ^1H NMR (400 MHz, $DMSO-d_6$) δ ppm 1.22 (d, $J=6.32$ Hz, 6 H) 3.48 - 3.54 (m, 4 H) 3.68 - 3.77 (m, 4H) 3.87 (s, 3 H) 4.77 - 4.85 (m, 1 H) 6.76 (d, $J=7.83$ Hz, 1 H) 6.94 - 7.07 (m, 2 H) 7.10 - 7.18 (m, 1 H) 8.38 (dd, $J=7.71$, 1.64 Hz, 1 H) 8.43 (s, 1 H) 8.71 (d, $J=8.08$ Hz, 1 H). ^{13}C NMR (100 MHz, $DMSO-d_6$) δ ppm 21.95, 44.07, 55.32, 68.14, 96.93, 101.10, 111.29, 120.48, 121.64, 125.83, 127.67, 136.28, 145.00, 154.31, 155.25, 155.51. LRMS (ESI) m/z 396.1 $[(M+H)]^+$, calc'd for $C_{21}H_{25}N_5O_3$: 395.47.

10

15


5.6.11. Synthesis of (S)-tert-butyl 2-(((3-(2-ethylphenyl)pyrazolo[1,5-a]pyrimidin-5-yl)amino)methyl)pyrrolidine-1-carboxylate




The Suzuki coupling of (S)-tert-butyl 2-(((3-bromopyrazolo[1,5-a]pyrimidin-5-yl)amino)methyl)pyrrolidine-1-carboxylate with (2-ethylphenyl)boronic acid under the same conditions as described in example 5.6.5 to give 77% of the titled compound. ^1H NMR (400 MHz, $CHLOROFORM-d$) δ ppm 1.20 (t, $J=7.58$ Hz, 3 H) 1.49 (s, 9 H) 1.79 - 1.95 (m, 3 H) 1.95 - 2.05 (m, 1 H) 2.82 (q, $J=7.58$ Hz, 2 H) 3.34 - 3.44 (m, 3 H) 3.44 - 3.53 (m, 1 H) 4.15 (br. s., 1 H) 6.08 (d, $J=6.82$ Hz, 1 H) 7.23 - 7.31 (m, 3 H) 7.32 - 7.38 (m, 1 H) 7.51 (dd, $J=7.20$, 1.39 Hz, 1 H) 7.91 (s, 1 H) 8.20 (d, $J=6.32$ Hz, 1 H); LRMS (ESI) m/e 422.0 $[(M+H)]^+$, calc'd for $C_{24}H_{31}N_5O_2$: 421.0].

20

5.6.12. Synthesis of (S)-tert-butyl 2-(((3-(2-[D3]methoxy phenyl)pyrazolo[1,5-a]pyrimidin-5-yl)amino)methyl)pyrrolidine-1-carboxylate



Part A. 2-(2-[D3]methoxyphenyl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane




5

To the 2-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenol (100 mg, 0.455 mmol) dissolved in 6 mL of DMSO was added deuterated methyl iodine (132 mg, 0.909 mmol) followed by 60% NaH (22 mg, 0.909 mmol). The resulting mixture was stirred overnight at rt. Dilute with 25 mL EtOAc, and wash with water and brine. Dry over MgSO₄ and concentrate to give 108 mg (100%) of the ether product.

10

Part B. (S)-tert-butyl 2-(((3-(2-[D3]methoxy phenyl)pyrazolo[1,5-a]pyrimidin-5-yl)amino)methyl)-pyrrolidine-1-carboxylate




The reaction of the 2-(2-[D3]methoxy phenyl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane with (S)-tert-butyl 2-(((3-bromopyrazolo[1,5-a]pyrimidin-5-yl)amino)methyl)pyrrolidine-1-carboxylate under Suzuki conditions described in example 5.6.5 afforded the titled compound in 70% yield. ¹H NMR (400 MHz, CHLOROFORM-d) δ ppm 1.51 (s, 9 H) 1.83 - 2.11 (m, 4 H) 3.40 (dd, J=7.45, 3.92 Hz, 2 H) 3.47 - 3.59 (m, 1 H) 3.59 - 3.66 (m, 1 H) 4.20 (br. s., 1 H) 6.07 (d, J=7.07 Hz, 1 H) 6.96 - 7.02 (m, 1 H) 7.06 (td, J=7.52, 1.14 Hz, 1 H) 7.15 - 7.24 (m, 1 H) 8.19 (d, J=6.32 Hz, 1 H) 8.46 (d, J=6.82 Hz, 1 H) 8.51 (s, 1 H); LRMS (ESI) m/e 427.0 [(M + H)⁺, calcd for C₂₃H₂₆D₃N₅O₃ 426.0].

15


20

5.6.13. Synthesis of 4-[3-(2-Methoxy-pyridin-3-yl)-pyrazolo[1,5-a]pyrimidin-5-yl]-piperazine-1-carboxylic acid isopropyl ester




Prepared similarly to the preparation of example 5.6.9 from 4-(3-bromo-pyrazolo[1,5-a]pyrimidin-5-yl)-piperazine-1-carboxylic acid isopropyl ester (338.7 mg, 0.9 mmol) and 2-methoxy-3-pyridineboronic acid [163105-90-6] (169.4 mg, 1.1 mmol) to afford 76.0 mg of yellow powder as a free base, mp. 131-132 °C. ¹H NMR (400 MHz, DMSO-*d*₆) δ ppm 1.22 (d, *J*=6.06 Hz, 6 H) 3.47 - 3.56 (m, 4 H) 3.71 - 3.80 (m, 4 H) 4.00 (s, 3 H) 4.82 (spt, *J*=6.23 Hz, 1 H) 6.81 (d, *J*=8.08 Hz, 1 H) 7.08 (dd, *J*=7.45, 4.93 Hz, 1 H) 7.98 (dd, *J*=4.93, 1.89 Hz, 1 H) 8.51 (s, 1 H) 8.76 (d, *J*=7.83 Hz, 1 H) 8.82 (dd, *J*=7.58, 2.02 Hz, 1 H). ¹³C NMR (100 MHz, DMSO-*d*₆) δ ppm 21.95, 42.85, 44.06, 53.10, 68.16, 97.21, 98.90, 116.40, 117.17, 134.55, 136.43, 141.99, 144.65, 144.71, 154.26, 155.46, 158.89. LRMS (ESI) *m/z* 397.1 [(*M*+*H*)]⁺, calc'd for C₂₀H₂₄N₆O₃: 396.45.

5.6.14. Synthesis of 4-[3-(1-Methyl-2-oxo-1,2-dihydro-pyridin-3-yl)-pyrazolo[1,5-a]pyrimidin-5-yl]-piperazine-1-carboxylic acid isopropyl ester




Prepared similarly to the preparation of example 5.6.9 from 4-(3-bromo-pyrazolo[1,5-a]pyrimidin-5-yl)-piperazine-1-carboxylic acid isopropyl ester and (1-methyl-2-oxo-1,2-dihydropyridin-3-yl)boronic acid to provide a yellow powder as a free base, mp. 206-207 °C. ¹H NMR (400 MHz, DMSO-*d*₆) δ ppm 1.23 (d, *J*=6.32 Hz, 6 H) 3.53 (s, 7 H) 3.70 - 3.82 (m, 4 H) 4.83 (spt, *J*=6.23 Hz, 1 H) 6.37 (t, *J*=6.82 Hz, 1 H) 6.78 (d, *J*=7.83 Hz, 1 H) 7.55 (dd, *J*=6.57, 1.26 Hz, 1 H) 8.64 (dd, *J*=7.20, 1.14 Hz, 1 H) 8.72 (d, *J*=7.83 Hz, 1 H) 8.96 (s, 1 H). ¹³C NMR (100 MHz, DMSO-*d*₆) δ ppm 21.97, 37.39, 42.84, 44.11, 68.16, 96.94, 100.56, 105.49, 122.80, 132.49, 135.00, 136.42, 144.40, 144.56, 154.29, 155.46, 160.29. LRMS (ESI) *m/z* 397.1 [(*M*+*H*)]⁺, calc'd for C₂₀H₂₄N₆O₃: 396.45.

5.6.15. Synthesis of 4-[3-(3-Fluoro-2-methoxy-phenyl)-pyrazolo[1,5-a]pyrimidin-5-yl]-piperazine-1-carboxylic acid isopropyl ester




Prepared similarly to the preparation of example 5.6.9 from 4-(3-bromo-pyrazolo[1,5-a]pyrimidin-5-yl)-piperazine-1-carboxylic acid isopropyl ester (389.9 mg, 1.1 mmol) and 3-fluoro-2-methoxyphenylboronic acid [762287-59-2] (216.2 mg, 1.3 mmol) to afford 153.8 mg of light yellow powder as a free base, mp. 52-53 °C. ¹H NMR (400 MHz, DMSO-*d*₆) δ ppm 1.22 (d, *J*=6.32 Hz, 6 H) 3.49 - 3.56 (m, 4 H) 3.75 (dd, *J*=6.44, 4.17 Hz, 4 H) 3.83 (d, *J*=1.01 Hz, 3 H) 4.82 (spt, *J*=6.23 Hz, 1 H) 6.81 (d, *J*=8.08 Hz, 1 H) 7.06 (ddd, *J*=11.37, 8.08, 1.52 Hz, 1 H) 7.16 (td, *J*=8.08, 5.81 Hz, 1 H) 8.23 (dt, *J*=7.89, 1.36 Hz, 1 H) 8.42 (s, 1 H). ¹⁹F NMR (377 MHz, DMSO-*d*₆) δ ppm -131.42. ¹³C NMR (100 MHz, DMSO-*d*₆) δ ppm 21.95, 42.83, 44.05, 60.38, 68.16, 97.24, 99.94, 99.98, 112.72, 112.90, 123.22, 124.06, 127.98, 136.39, 143.40, 143.51, 144.25, 144.49, 154.28, 154.63, 155.53, 157.04. LRMS (ESI) *m/z* 414.1 [(*M*+H)]⁺, calc'd for C₂₁H₂₄FN₅O₃: 413.46.

5.6.16. Synthesis of Isopropyl 4-(3-(2-fluoropyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate



Isopropyl 4-(3-bromopyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate (4g, 10.86 mmol), 2-fluoropyridine-3-boronic acid hydrate (3.44g, 21.7 mmol), potassium carbonate (4.5g, 32.6 mmol), Pd(OAc)₂ (24mg, 0.1086 mmol), and x-Phos (146mg, 0.217 mmol) were taken up in dioxane and water under nitrogen and stirred at 80 °C for 2 hours. Reaction then cooled to room temperature, filtered through celite plug with ethyl acetate. Reduced in vacuo and run through an isco silica column with hexane:ethyl acetate 75-100%. Product plus desbromo co elute, reduced in vacuo. Product recrystallized 3 times from IPAC, washed with heptanes dried to get 889mg product. ¹H NMR (400 MHz, DMSO-*d*₆) δ 8.95 (t, *J* = 2.15 Hz, 1H), 8.78 (d, *J* = 7.83 Hz, 1H), 8.29 (d, *J* = 4.29 Hz, 1H), 8.00 (d, *J* = 4.80 Hz, 1H), 7.43 (dt, *J* = 2.40, 4.99 Hz, 1H), 6.85 (d, *J* = 7.83 Hz, 1H), 4.78 - 4.87 (m, 1H), 3.73 - 3.83 (m, 4H), 3.48 - 3.58 (m, 4H), 3.31 (s, 1H), 1.22 (d, *J* = 6.06 Hz, 6H) LRMS (ESI) *m/z* 385 [(*M*+H)]⁺, calc'd for C₁₉H₂₁FN₆O₂: 384.42.

5.6.17. Synthesis of Isopropyl 4-(3-(5-fluoro-2-methoxypyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate




To a 200 mL round bottom flask was added 4.00 g (10.87 mmol) of isopropyl 4-(3-bromopyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate, 2.89 g (11.41 mmol) of (5-fluoro-2-methoxypyridin-3-yl)boronic acid, 0.38 g (0.33 mmol) of Pd(PPh₃)₄, and a stirring rod. After adding all the solids, 60 ml of 2% TPGS-750-M solution was added, followed by the addition of 7.56 mL (54.35 mmol) triethylamine.

The reaction flask was plunged into an oil bath pre-heated to 90°C, and stirred vigorously. It took about 0.25 hr of stirring for all the suspended solids to dissolve completely. Several color changes were observed, the last color that remained through the rest of the reaction was a very dark brown color.

After 1.5 hr, LCMS showed that reaction was complete. It was allowed to cool to RT, and diluted with 100 mL EtOAc. The organic layer was separated, and the aqueous layer was extracted twice with 40 mL EtOAc portions. The combined organic layers was washed with brine, dried over MgSO₄, and concentrated. A small volume of DCM was used to dissolve the crude mixture, and then loaded directly onto a 120 g silica gel column for separation on the ISCO. A gradient of 40% EtOAc/hex to 100% EtOAc was run for the first 25 minutes. The desired product did not elute, only some impurities eluted. However, about 10 minutes into running with 100% EtOAc, the desbromo side product started eluting together with the desired product. Even though the desbromo compound is spread through all the fractions containing the desired product, the percentage of the desbromo compound seem to drastically decrease with time. Therefore, the initial fractions containing the highest percentages of the desbromo were discarded, and the remaining fractions were concentrated to obtain 3.4 g of about 87% purity at 254 nm. (with about 13% desbromo compound).


[Base on ¹H NMR, the desbromo is over-estimated at the 254 nm on the UV, and under estimated on the UV at 220 nm]. The material was recrystallized twice using IPAC and heptanes to obtain 3.16 g (70% yield) of the desired product with purity over 99% on the UV at 254 nm. ¹H NMR (400 MHz, CHLOROFORM-*d*) δ ppm 1.31 (d, *J*=6.32 Hz, 6 H) 3.61 - 3.70 (m, 4 H) 3.77 (br. s., 4 H) 4.08 (s, 3 H) 5.00 (quin, *J*=6.19 Hz, 1 H) 6.41 (d, *J*=7.83 Hz, 1 H) 7.83 (d, *J*=3.03 Hz, 1 H) 8.37 (d, *J*=7.83 Hz, 1 H) 8.63 - 8.71 (m, 2 H). LRMS (ESI) *m/e* 415 [(M + H)⁺, calcd for C₂₀H₂₃FN₆O₃ 414].


5.6.18. Synthesis of Isopropyl 4-(3-(2-ethoxypyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate




To a 1 liter round bottom flask was added 15.00 g (40.76 mmol) of isopropyl 4-(3-
 5 bromopyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate, 8.85 g (52.99 mmol) of (2-
 ethoxypyridin-3-yl)boronic acid, 14.06 g (101.90 mmol) of K_2CO_3 , 1.41g (1.22 mmol) of $P(PPh_3)_4$,
 and a stirring rod. After adding all the solids, 200 ml of MeCN was added, followed by the addition of
 100 mL of water. The reaction flask was plunged into an oil bath pre-heated to 85°C, and stirred
 vigorously. This reaction was setup to reflux. After stirring at 85°C for about 16 hr, it was allowed to
 10 cool to RT and diluted with 200 mL of EtOAc. The organic layer was separated, and the aqueous layer
 was extracted twice with 80 mL portions of EtOAc. The combined organic layers was washed with
 brine, dried over $MgSO_4$, and concentrated. A small volume of DCM was used to dissolve the crude
 mixture, and then loaded directly onto a 750 g silica gel column for separation on the ISCO-XL. A
 gradient of 50% EtOAc/hex to 100% EtOAc was run for the first 35 minutes. The desired product did
 15 not elute, only some impurities eluted. However, about 10 minutes into running with 100% EtOAc,
 the desbromo side product starts eluting together with the desired product. Even though the
 desbromo compound is spread through all the fractions containing the desired product, the
 percentage of the desbromo compound seem to decrease with time. Therefore, the initial fractions
 containing the highest percentages of the desbromo were discarded, and the remaining fractions
 20 were concentrated to obtain 13.46 g of about 86% purity at 254 nm. (with about 14% desbromo
 compound). This material was recrystallized twice, using IPAC and heptanes, to obtain 11.4 g (68%)
 of the desired product. ^1H NMR (400 MHz, $CHLOROFORM-d$) δ ppm 1.30 (d, $J=6.32$ Hz, 6 H) 1.53 (t,
 $J=7.07$ Hz, 3 H) 3.59 - 3.70 (m, 4 H) 3.70 - 3.81 (m, 4 H) 4.53 (q, $J=7.07$ Hz, 2 H) 5.00 (dt, $J=12.44$,
 6.28 Hz, 1 H) 6.38 (d, $J=7.83$ Hz, 1 H) 6.98 (dd, $J=7.45$, 4.93 Hz, 1 H) 8.01 (dd, $J=4.93$, 1.89 Hz, 1
 25 H) 8.37 (d, $J=7.83$ Hz, 1 H) 8.70 (s, 1 H) 8.76 (dd, $J=7.45$, 1.89 Hz, 1 H). LRMS (ESI) m/e 411 [(M +
 H) $^+$, calcd for $C_{21}H_{26}N_6O_3$ 410].

5.6.19. Synthesis of 2-Methoxyethyl 4-(3-(2-methoxypyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate



Part A. 2-methoxyethyl 4-(3-bromopyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate

To 10.00 g (35.44 mmol) of 3-bromo-5-(piperazin-1-yl)pyrazolo[1,5-a]pyrimidine dissolved in 120 mL of EtOAc, at 0°C was added 5.40 g (38.99 mmol) of 2-methoxyethyl carbonochloridate, followed by 9.86 mL (70.88 mmol) of TEA. After 5 minutes, the ice bath was removed, and the reaction mixture stirred at RT. After 5 hr, LCMS showed that the reaction was complete. It was diluted with EtOAc, quenched with brine, and the organic layer dried over MgSO₄. It was concentrated and purified on the ISCO with a 330 g column eluting with 0-9% MeOH/DCM to obtain the desired product, 12.63 g (93%). ¹H NMR (400 MHz, CHLOROFORM-*d*) δ ppm 3.42 (s, 3 H) 3.62 - 3.69 (m, 6 H) 3.73 - 3.82 (m, 4 H) 4.26 - 4.35 (m, 2 H) 6.36 (d, *J*=7.83 Hz, 1 H) 7.86 (s, 1 H) 8.27 (d, *J*=7.83 Hz, 1 H). LRMS (ESI) *m/e* 384 [(M + H)⁺, (doublet), calcd for C₁₄H₁₈BrN₅O₃ 383].


Part B. 2-methoxyethyl 4-(3-(2-methoxypyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate

To a 100 mL round bottom flask was added 15.00 g (39.06 mmol) of 2-methoxyethyl 4-(3-bromopyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate, 7.17 g (46.88 mmol) of (2-methoxypyridin-3-yl)boronic acid, 38 mL (273.44 mmol) of TEA, 200 mL of MeCN was added, 100 mL of water, followed by the addition of 0.21 g (0.29 mmol) of Pd-132. The reaction flask was plunged into an oil bath pre-heated to 85°C, and stirred. After 1 hr, the reaction was complete, and it was allowed to cool to RT, and then diluted with 200 mL of EtOAc. The organic layer was separated, and the aqueous layer was extracted twice with 150 mL portions of EtOAc. The combined organic layers was washed with brine, dried over MgSO₄, and filtered through a thin pad of silica gel, and the filtrate was concentrated to obtain 14 g solid product, [90% desired product and 10% desbromo side product, based on uv absorption at 254 nm].


The resulting solid was subjected to charcoal treatment, by dissolving all 14 g in 100 mL DCE and adding 500 mg of Darco. It was heated and stirred for about 1.5 hr, cooled and then filtered through celite, and concentrated. The product obtained appeared to be clean enough to be recrystallized without subjecting it to silica gel chromatography. The recrystallization was carried out by dissolving it in a small volume of IPAC at about 75°C (a few milliliters MeOH was used to aid the

dissolution of the solids). After complete dissolution, heptane was added and precipitates begun forming, and allowed to cool at 0°C for 1 hr. It was then filtered and dried 13.1 g product with purity of about 97%. This material was recrystallized again, to obtain a pale yellow solid 12.45 g (77% yield, purity to over 99%) of the desired product. ¹H NMR (400 MHz, CHLOROFORM-*d*) δ ppm 3.42 (s, 3 H) 3.61 - 3.71 (m, 6 H) 3.72 - 3.80 (m, 4 H) 4.10 (s, 3 H) 4.29 - 4.34 (m, 2 H) 6.38 (d, *J*=7.83 Hz, 1 H) 7.00 (dd, *J*=7.45, 4.93 Hz, 1 H) 8.04 (dd, *J*=4.93, 1.89 Hz, 1 H) 8.37 (d, *J*=7.83 Hz, 1 H) 8.64 (s, 1 H) 8.74 (dd, *J*=7.45, 1.89 Hz, 1 H). ¹³C NMR (101 MHz, CHLOROFORM-*d*) δ ppm 43.49, 44.87, 53.51, 59.15, 64.96, 71.07, 76.92, 77.24, 77.56, 96.37, 100.99, 116.93, 117.05, 135.42, 136.16, 142.79, 145.37, 146.26, 155.46, 155.52, 160.08. LRMS (ESI) *m/e* 413 [(*M* + *H*)⁺, calcd for C₂₀H₂₄N₆O₄ 412].

5.6.20. Synthesis of tert-butyl 4-(3-(2-methoxypyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate




Part A. tert-butyl 4-(3-bromopyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate




3-bromo-5-chloropyrazolo[1,5-a]pyrimidine (10g, 43mmol) taken up in isopropanol. T-butylpiperazine carboxylate (9.61g, 51.6mmol), diisopropylethylamine (22.4mL, 129mmol) added and stirred at 65° C overnight. Reaction was cooled to room temperature, diluted with water and solid product filtered off. The solid was washed with water, dried to obtain 15.84g crude product for further reaction, as is. LRMS (ESI) *m/z* 382/384 [(*M*+*H*)⁺, calc'd for C₁₅H₂₀BrN₅O₂: 382.2.

Part B. tert-butyl 4-(3-(2-methoxypyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate




Tert-butyl 4-(3-bromopyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate (15.84g, 41.5mmol), 2-methoxypyridine-3-boronic acid (12.68g, 82.9mmol), potassium carbonate (17.16g, 124.5mmol), Pd(OAc)₂ (92mg, 4.15mmol), and x-Phos (394mg, 8.3mmol) were taken up in 60mL dioxane and 30mL water under nitrogen and stirred at 75° C for 2 hours. Reaction then cooled to room temperature, diluted with toluene washed with brine. The organic layer filtered through magnesium sulfate and celite plug. Reduced in vacuo and run through an ISCO silica column with hexane:ethyl acetate 0-100%. The product plus some desbromo co elute, Concentrated in vacuo. Product recrystallized 2 times from IPAC:Heptane 30:70, washed with heptanes dried to get 12g product. ¹H NMR (400 MHz, DMSO-d₆) δ 8.78 - 8.85 (m, 1H), 8.74 (d, *J* = 8.08 Hz, 1H), 8.50 (s, 1H), 7.92 - 8.02 (m, 1H), 7.04 - 7.13 (m, 1H), 6.80 (d, *J* = 8.08 Hz, 1H), 4.00 (s, 3H), 3.70 - 3.78 (m, 4H), 3.49 (d, *J* = 5.31 Hz, 4H), 1.44 (s, 9H). LRMS (ESI) *m/z* 411 [(M+H)]⁺, calc'd for C₂₁H₂₆N₆O₃: 410.48.

5.6.21. Synthesis of Isopropyl 4-(3-(5-fluoro-2-methoxyphenyl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate




Part A. isopropyl 4-(3-bromopyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate




3-bromo-5-(piperazin-1-yl)pyrazolo[1,5-a]pyrimidine (2.5g, 8.87mmol) taken up in DCM. Isopropylchloroformate 1M in toluene (13.3mL, 13.3mmol), and triethylamine (2.47mL, 17.7mmol) added and stirred at room temperature 30 minutes. Reaction washed with water, DCM layer dried over magnesium sulfate filtered reduced in vacuo. Run on an Isco silica column with DCM:MeOH 0-5%. Product fractions were combined and reduced in vacuo to obtain 2.86g crude product for further reaction as is. LRMS (ESI) *m/z* 368/370 [(M+H)]⁺, calc'd for C₁₄H₁₈BrN₅O: 368.24.

Part B. isopropyl 4-(3-(5-fluoro-2-methoxyphenyl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate



Isopropyl 4-(3-bromopyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate (2.86g, 7.8mmol),
 5 5-fluoro-2-methoxyphenyl boronic acid (2.64g, 15.5mmol), potassium carbonate (3.22g, 23.3mmol),
 Pd(OAc)₂ (35mg, 0.155mmol), and x-Phos (146mg, 0.31mmol) were taken up in 15mL dioxane and
 7mL water under nitrogen and stirred at 80° C for 2 hours. Reaction then cooled to room
 temperature, filtered through celited plug with ethyl acetate. Reduced in vacuo and run through an
 isco silica column with hexane:ethyl acetate 75-100%. Product plus desbromo co-elute, reduced in
 10 Vacuo. Product recrystallized 2 times from ethyl acetate:Hexane, washed with heptanes dried to get
 12g product. ¹H NMR (400 MHz, DMSO-d₆) δ 8.73 (d, *J* = 7.78 Hz, 1H), 8.50 (s, 1H), 8.29 (dd,
J = 3.14, 11.17 Hz, 1H), 7.04 (dd, *J* = 5.02, 9.03 Hz, 1H), 6.95(dd, *J* = 3.26, 7.78 Hz, 1H), 6.79 (d, *J*
 = 8.03 Hz, 1H), 4.77 - 4.85 (m, 1H), 3.87 (s, 3H), 3.70 - 3.78 (m, 4H), 3.48 - 3.56 (m, 4H), 1.21 (d, *J*
 = 6.27Hz, 6H). LRMS (ESI) *m/z* 414 [(M+H)]⁺, calc'd for C₂₁H₂₄N₅O₃: 413.4.

15 **5.6.22. Synthesis of Isopropyl 4-(3-(2-methoxy-6-methylpyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate**



To a 100 mL liter round bottom flask was added 2.00 g (5.43 mmol) of isopropyl 4-(3-
 bromopyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate, 1.00 g (5.98 mmol) of (2-methoxy-6-
 20 methylpyridin-3-yl)boronic acid, 0.19 g (0.16 mmol) of Pd(PPh₃)₄, and a stirring rod.

After adding all the solids, 30 ml of 2% TPGS-750-M solution was added, followed by the
 addition of 3.80 mL (27.15 mmol) triethylamine. The reaction flask was plunged into an oil bath pre-
 heated to 85°C, and stirred vigorously. It took about 0.25 hr of stirring for all the suspended solids to
 dissolve completely. After stirring at 85°C for about 2 hr, LCMS showed that the reaction was
 25 complete. It was allowed to cool to RT, dilute with 40 mL EtOAc. The organic layer was separated, and
 the aqueous layer was extracted twice with 30 mL portions EtOAc. The combined organic layers was
 washed with brine, dried over MgSO₄, and concentrated. A small volume of DCM was used to dissolve

the crude mixture, and then loaded directly onto a 80 g silica gel column for separation on the ISCO. A gradient of 50% EtOAc/hex to 100% EtOAc was run for the first 15 minutes, and then the column was run with only EtOAc. The desired product begun eluting about 8 minutes into running with only EtOAc. The desbromo side product elutes together with the desired product. Even though the


5 desbromo compound is spread through all the fractions containing the desired product, the percentage of the desbromo compound seem to decrease with time. Therefore, the first few tubes containing the highest percentages of the desbromo were discarded, and the remaining fractions were concentrated to obtain 1.73 g of about 84% purity at 254 nm (with about 16% desbromo compound). [Base on ^1H nmr, the desbromo is over-estimated at the 254 nm on the UV, and under

10 estimated on the UV at 220 nm].


The material was recrystallized twice using IPAC and heptanes, to afford 1.45 g (65% yield) of the desired product. ^1H NMR (400 MHz, CHLOROFORM-*d*) δ ppm 1.30 (d, $J=6.32$ Hz, 6 H) 2.49 (s, 3 H) 3.59 - 3.68 (m, 4 H) 3.73 (dd, $J=6.19, 3.66$ Hz, 4 H) 4.07 (s, 3 H) 4.99 (dt, $J=12.44, 6.28$ Hz, 1 H) 6.36 (d, $J=7.83$ Hz, 1 H) 6.85 (d, $J=7.58$ Hz, 1 H) 8.35 (d, $J=7.83$ Hz, 1 H) 8.55 - 8.63 (m, 2 H); ^{13}C

15 NMR (101 MHz, CHLOROFORM-*d*) δ ppm 22.48, 24.10, 43.39, 44.94, 53.38, 69.37, 96.32, 101.34, 113.40, 116.03, 135.94, 136.12, 145.14, 146.15, 152.13, 155.35, 155.45, 159.49. LRMS (ESI) m/e 411 $[(M + H)^+]$, calcd for $\text{C}_{21}\text{H}_{26}\text{N}_6\text{O}_3$ 410].

5.6.23. Synthesis of Isopropyl 4-(3-(6-methoxypyridin-2-yl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate



Part A. isopropyl 4-(3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate




To 184 mg (0.50 mmol) of isopropyl 4-(3-bromopyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-

25 carboxylate in a 50 mL round bottom flask was added 191 mg (0.75 mmol) of 4,4,4',4',5,5,5',5'-octamethyl-2,2'-bi(1,3,2-dioxaborolane), 147 mg (1.50 mmol) of KOAc, 8 mL of DMSO followed by 41 mg (0.05 mmol) of $\text{PdCl}_2(\text{dppf})_2 \cdot \text{DCM}$. The air was replaced with N_2 , and the mixture heated to 80°C with stirring. Next morning LCMS shows reaction had gone to completion. It was cooled to RT, diluted


with EtOAc, washed with brine, and the organic layer dried over MgSO₄. It was concentrated and purified in the ISCO using a 12 g column, and eluting with 25 -100% EtOAc/hex to obtain 107 mg of the desired product [Purity was only about 85%, about 15% was hydrolyzed to the corresponding boronic acid].

- 5 **Part B. Isopropyl 4-(3-(6-methoxypyridin-2-yl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate.** To 65 mg (0.157 mmol) of isopropyl 4-(3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate in a microwave vial was added, 29 mg (0.157 mmol) of 2-bromo-6-methoxypyridine, 43 mg (0.313 mmol) of K₂CO₃, 18 mg (0.016 mmol) Pd(PPh₃)₄, 3 mL of DME and then 1 mL water. The resulting mixture was microwaved at 125°C for
- 10 0.33 hr. It was diluted with EtOAc, washed with brine, and the organic layer dried over MgSO₄. It was concentrated and purified on the neutral PREP HPLC to obtain 42 mg of the desired product. ¹H NMR (400 MHz, CHLOROFORM-*d*) δ ppm 1.30 (d, *J*=6.32 Hz, 6 H) 3.60 - 3.74 (m, 4 H) 3.81 (br. s., 4 H) 4.04 (s, 3 H) 5.00 (quin, *J*=6.25 Hz, 1 H) 6.39 (d, *J*=7.83 Hz, 1 H) 6.55 (dd, *J*=8.08, 0.76 Hz, 1 H) 7.62 (dd, *J*=8.08, 7.58 Hz, 1 H) 7.83 - 7.93 (m, 1 H) 8.37 (d, *J*=7.83 Hz, 1 H) 8.66 (s, 1 H). LRMS
- 15 (ESI) *m/e* 397 [(M + H)⁺, calcd for C₂₀H₂₄N₆O₃ 396].

5.6.24. Synthesis of 5-isopropyl-3-(4-(3-(2-methoxypyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazin-1-yl)-1,2,4-oxadiazole




Part A. 4-(3-bromopyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carbonitrile



20

- 3-bromo-5-(piperazin-1-yl)pyrazolo[1,5-a]pyrimidine (500mg, 1.77mmol) taken up in 20mL DCM and stirred at 0 °C in an ice bath. Sodium bicarbonate (744mg, 8.86mmol) in 6mL water added and stirred. Cyanogen bromide (225mg, 2.12mmol) added and stirred at 0 °C for 30 minutes then allowed to reach room temperature and stirred overnight. Reaction diluted with DCM washed
- 25 with water, DCM layer dried over magnesium sulfate reduced in vacuo for 500 mg crude product for further reaction as is. LRMS (ESI) *m/z* 307/309 [(M+H)⁺, calc'd for C₁₁H₁₁BrN₆: 307.15.

Part B. 3-(4-(3-bromopyrazolo[1,5-a]pyrimidin-5-yl)piperazin-1-yl)-5-isopropyl-1,2,4-oxadiazole


4-(3-bromopyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carbonitrile (450mg, 1.46mmol) in 9mL DMF and sodium carbonate (155mg, 1.46mmol) stirred room temperature 15minutes,

- 5 hydroxylamine HCl (203mg, 2.9mmol) added and reaction stirred 30 minutes at 80°C. Then 12mL toluene added and pyridine (474μL, 5.86mmol) and isobutyric anhydride (973μL, 5.86mmol) added and reaction stirred at 80°C for 1.5 hours. Reaction cooled to room temperature reduced in vacuo then taken up in ethyl acetate and washed with water, brine, water. Organic layer dried over magnesium sulfate and reduced in vacuo. For 250 mg crude product to use as is in further reactions.
- 10 LRMS (ESI) m/z 392/394[(M+H)]⁺, calc'd for C₁₅H₁₈BrN₇O: 392.26.


Part C. 5-isopropyl-3-(4-(3-(2-methoxypyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazin-1-yl)-1,2,4-oxadiazole.

- 3-(4-(3-bromopyrazolo[1,5-a]pyrimidin-5-yl)piperazin-1-yl)-5-isopropyl-1,2,4-oxadiazole (200mg, 0.51mmol), 2-methoxypyridine-3-boronic acid (156mg, 1.0mmol), potassium carbonate (211mg, 1.53mmol), Pd(OAc)₂ (3mg, 0.01mmol), and x-Phos (10mg, 0.02mmol) were
- 15 taken up in 2mL dioxane and 1mL water in a sealed tube and heated at 85°C for 2 hours. Reaction then cooled to room temperature filtered through a celite plug with acetonitrile and DCM, reduced in vacuo. Purified on Shimadzu neutral phase prep lyophilized to get 12mg product. ¹H NMR (400 MHz, DMSO-d₆) δ 8.83 (dd, J = 1.77, 7.58 Hz, 1H), 8.77 (d, J = 7.83 Hz, 1H), 8.51 (s, 1H), 7.99 (dd, J = 1.89, 4.93 Hz, 1H), 7.10 (dd, J = 4.80, 7.58 Hz, 1H), 6.86 (d, J = 8.08 Hz, 1H), 4.00 (s, 3H), 3.84 -
- 20 3.90 (m, 4H), 3.48 - 3.53 (m, 4H), 3.15 (t, J = 7.07 Hz, 1H), 1.29 (d, J = 6.82 Hz, 6H). LRMS (ESI) m/z 421 [(M+H)]⁺, calc'd for C₂₁H₂₄N₈O₂: 420.48.

5.6.25. Synthesis of (S)-2-amino-1-(4-(3-(2-methoxypyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazin-1-yl)-4-methylpentan-1-one




Part A. (S)-2-amino-1-(4-(3-bromopyrazolo[1,5-a]pyrimidin-5-yl)piperazin-1-yl)-4-methylpentan-1-one




3-bromo-5-(piperazin-1-yl)pyrazolo[1,5-a]pyrimidine (200mg, 0.71mmol), Boc-L-Leu (196mg, 0.85mmol), HATU (245mg, 1.06mmol), and triethylamine (394uL, 2.83mmol) taken up in DMF and stirred at room temperature overnight. Reaction was diluted with DCM and washed with water. The organic layer was dried over magnesium sulfate, filtered and reduced in vacuo. The residue was taken up in 20mL of a 40%TFA in DCM solution and stirred at 35°C overnight. The mixture was concentrated in vacuo and passed through a silica plug with 10%MeOH in DCM. Solvent was removed and the residue was dried to give 600mg crude product as TFA salt for further reactions. LRMS (ESI) m/z 395/397 [(M+H)]⁺, calc'd for C₁₆H₂₃BrN₆O: 395.31.


Part B. (S)-2-amino-1-(4-(3-(2-methoxypyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazin-1-yl)-4-methylpentan-1-one. (S)-2-amino-1-(4-(3-bromopyrazolo[1,5-a]pyrimidin-5-yl)piperazin-1-yl)-4-methylpentan-1-one (300mg, 0.59mmol), 2-methoxypyridine-3-boronic acid (135mg, 0.88mmol), potassium carbonate (326mg, 2.36mmol), Pd(OAc)₂ (3mg, 0.011mmol), and x-Phos (11mg, 0.022mmol) were taken up in 2mL dioxane and 1mL water in a sealed tube and heated at 85°C for 2hours. Reaction then cooled to room temperature filtered through a celite plug with acetonitrile and DCM, reduced in vacuo. Purified on shimadzu neutral phase prep, lyophilized to get 28.1mg of formic acid salt product. ¹H NMR (400 MHz, DMSO-d₆) δ 8.82 (dd, J = 2.02, 7.58 Hz, 1H), 8.77 (d, J = 7.83 Hz, 1H), 8.51 (s, 1H), 8.27 (s, 1H), 7.99 (dd, J = 1.77, 4.80 Hz, 1H), 7.09 (dd, J = 4.80, 7.58 Hz, 1H), 6.83 (d, J = 8.08 Hz, 1H), 4.00 (s, 3H), 3.62 - 3.93 (m, 9H), 1.72 - 1.88 (m, 1H), 1.23 - 1.42 (m, 2H), 0.91 (dd, J = 6.57, 10.11 Hz, 6H). LRMS (ESI) m/z 424 [(M+H)]⁺, calc'd for C₂₂H₂₉N₇O₂: 423.52.

5.6.26. Synthesis of (S)-2-amino-N-((S)-1-(3-(2-methoxypyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)pyrrolidin-3-yl)-N,4-dimethylpentanamide




Part A. (S)-tert-butyl (1-(3-bromopyrazolo[1,5-a]pyrimidin-5-yl)pyrrolidin-3-yl)carbamate

3-bromo-5-chloropyrazolo[1,5-a]pyrimidine (2.2g, 9.5mmol) (S)-3-Boc-aminopyrrolidine (2.6g, 14.2mmol), and triethylamine (5.27mL, 37.9mmol) were taken up in isopropanol in a sealed tube and microwaved at 140°C for 30 minutes. Reaction stripped down in vacuo taken up in ethyl acetate washed with water then passed through a silica plug with ethyl acetate. Solvent was removed in vacuo to give 3.6g crude product that was carried on as is to part B. LRMS (ESI) m/z 382/384 [(M+H)⁺], calc'd for C₁₅H₁₉BrN₅O₂: 382.2.

Part B. (S)-1-(3-bromopyrazolo[1,5-a]pyrimidin-5-yl)-N-methylpyrrolidin-3-amine

(S)-tert-butyl (1-(3-bromopyrazolo[1,5-a]pyrimidin-5-yl)pyrrolidin-3-yl)carbamate (3.6g, 9.4mmol) taken up in 15mL DMF and cooled to 0°C in an ice bath. Iodomethane (1.026mL, 16.5mmol) added and stirred 5 minutes. Then sodium hydride 60% in oil (754mg, 18.8mmol) was slowly added. Reaction stirred 5 minutes at 0°C then removed from ice bath and stirred 30 minutes at room temperature. Reaction was quenched with ice then extracted with ethyl acetate 2x. Ethyl acetate fractions combined dried over magnesium sulfate filtered reduce in vacuo. This was then taken up in 26mL DCM and 4mL TFA added. Stirred 1 hour until complete by LC/MS. Reaction washed with 1N NaOH, DCM layer removed dried over magnesium sulfate filtered, reduced in vacuo to get 2.72g crude product carried on as is to part C. LRMS (ESI) m/z 296/298 [(M+H)⁺], calc'd for C₁₁H₁₄BrN₅: 296.17.

Part C. (S)-2-amino-N-((S)-1-(3-bromopyrazolo[1,5-a]pyrimidin-5-yl)pyrrolidin-3-yl)-N,4-dimethylpentanamide


(S)-1-(3-bromopyrazolo[1,5-a]pyrimidin-5-yl)-N-methylpyrrolidin-3-amine (200 mg, 0.68mmol) Boc-L-Leu (187mg, 0.81mmol), HATU (385mg, 1.01mmol), and triethylamine (375uL, 2.7mmol) Taken up in DMF and Stirred at room temp overnight. Reaction diluted with DCM, washed with water, DCM layer then dried over magnesium sulfate filtered and reduced in vacuo. This was then taken up in 26mL DCM and 4mL TFA added. Stirred 1 hour until complete by LC/MS. Reaction

washed with 1N NaOH, DCM layer removed dried over magnesium sulfate filtered, reduced in vacuo for 226mg crude product used as is in part D. LRMS (ESI) m/z 409/411 [(M+H)]⁺, calc'd for C₁₇H₂₄BrN₆O: 409.


Part D. (S)-2-amino-N-((S)-1-(3-(2-methoxypyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)pyrrolidin-

5 3-yl)-N,4-dimethylpentanamide. (S)-2-amino-N-((S)-1-(3-bromopyrazolo[1,5-a]pyrimidin-5-yl)pyrrolidin-3-yl)-N,4-dimethylpentanamide (220mg, 0.54mmol), 2-methoxypyridine-3-boronic acid (164mg, 1.1mmol), potassium carbonate (291mg, 2.15mmol), Pd(OAc)₂ (2mg, 0.011mmol) and x-phos (10mg, 0.022mmol) were taken up in 2mL acetonitrile and 1mL water and heated in a sealed tube at 85° C for 2 hours. Reaction was then filtered through celite with acetonitrile and DCM reduced in
10 vacuo, then purified on Shimadzu neutral phase prep and product fractions lyophilized to get (S)-2-amino-N-((S)-1-(3-(2-methoxypyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)pyrrolidin-3-yl)-N,4-dimethylpentanamide 41.3mg as 0.5eq formic salt. ¹H NMR (400 MHz, DMSO-d₆) δ 8.87 - 8.97 (m, 1H), 8.72 (s, 1H), 8.50 (s, 1H), 8.30 (s, 1H), 7.96 (dd, J = 2.02, 4.80 Hz, 1H), 7.00 - 7.11 (m, 1H), 6.51 (d, J = 7.83 Hz, 1H), 4.00 (s, 3H), 3.48 - 3.64 (m, 1H), 3.43 (br. s., 3H), 2.96 (s, 2H), 2.81
15 (s, 1H), 2.11 - 2.28 (m, 2H), 1.72 - 1.89 (m, 1H), 1.19 - 1.39 (m, 2H), 0.90 (dd, J = 1.89, 6.69 Hz, 6H). (ESI) m/z 438 [(M+H)]⁺, calc'd for C₂₃H₃₁N₇O₂: 437.5.

5.6.27. Synthesis of (S)-2-amino-1-(4-(3-(2-methoxypyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazin-1-yl)-3-methylbutan-1-one



20 Part A. (S)-2-amino-1-(4-(3-bromopyrazolo[1,5-a]pyrimidin-5-yl)piperazin-1-yl)-3-methylbutan-
1-one




3-bromo-5-(piperazin-1-yl)pyrazolo[1,5-a]pyrimidine (400mg, 1.42 mmol) Boc-L-Leucine (392mg, 1.7mmol), HATU (490mg, 2.13mmol), and triethylamine (788uL, 5.7mmol) taken up in DMF
25 and stirred at room temperature overnight. Reaction diluted with DCM washed with water, DCM layer then dried over magnesium sulfate filtered and reduced in Vacuo. This was then taken up in 26mL DCM and 4mL TFA added. Stirred 1 hour until complete by LC/MS. Reaction was washed with

1N NaOH, DCM layer removed dried over magnesium sulfate filtered, reduced in vacuo to obtain 349mg crude product for further reaction as is. LRMS (ESI) m/z 381/383 [(M+H)]⁺, calc'd for C₁₅H₂₁BrN₆O: 381.28.

Part B. (S)-2-amino-1-(4-(3-(2-methoxypyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazin-1-yl)-


- 5 3-methylbutan-1-one. (S)-2-amino-1-(4-(3-bromopyrazolo[1,5-a]pyrimidin-5-yl)piperazin-1-yl)-3-methylbutan-1-one (110mg, 0.29mmol), (2-methoxypyridin-3-yl)boronic acid (88mg, 0.58 mmol), potassium carbonate (120mg, 0.87mmol), Pd(OAc)₂ (1mg, 0.0058 mmol), and x-Phos (5mg, 0.012mmol) were taken up in 3mL dioxane and 1mL water under nitrogen and stirred at 85° C for 2 hours. Reaction then cooled to room temperature filtered through celited plug with acetonitrile and
- 10 DCM. Reduced in vacuo and purified on shimadzu neutral phase prep, lyophilized to get 18mg 1.5eq formic salt product. ¹H NMR (400 MHz,MSO-d₆) δ 8.83 (dd, J = 1.77, 7.58 Hz, 1H), 8.78 (d, J = 7.83 Hz, 1H), 8.52 (s, 1H), 8.35 (s, 2H), 7.99 (dd, J = 1.89, 4.93 Hz,1H), 7.09 (dd, J = 4.93, 7.45 Hz, 1H), 6.85 (d, J = 7.83 Hz, 1H), 4.01 (s, 3H), 3.81 - 3.89 (m, 1H), 3.74 (d, J = 5.31 Hz, 3H), 3.68 (d, J = 7.58 Hz,3H), 1.76 - 1.89 (m, 1H), 0.94 (d, J = 6.82 Hz, 3H), 0.86 (d, J = 6.82 Hz, 3H). LRMS
- 15 (ESI) m/z 410 [(M+H)]⁺, calc'd for C₂₁H₂₇N₇O₂: 409.5.

5.6.28. Synthesis of (S)-2-amino-3-methoxy-1-(4-(3-(2-methoxypyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazin-1-yl)propan-1-one



Part A. (S)-2-amino-1-(4-(3-bromopyrazolo[1,5-a]pyrimidin-5-yl)piperazin-1-yl)-3-

- 20 methoxypropan-1-one




- 3-Bromo-5-(piperazin-1-yl)pyrazolo[1,5-a]pyrimidine (300mg, 1.1mmol) taken up in 10mL DMF. Boc-Ser(Me)-OH (280mg, 1.3mmol), HATU (368mg, 1.6mmol), and triethylamine (590uL, 4.25mmol) was added and stirred at room temp overnight. Reaction was diluted with DCM and
- 25 washed with water. DCM layer was dried over magnesium sulfate reduced in vacuo. This was then taken up in 30% TFA in DCM solution and stirred at room temperature for 2 hours. It was then cooled to 0° C and conc Aq NaOH added slowly until aqueous layer remains basic. DCM layer

extracted dried over magnesium sulfate, reduced in vacuo for 405mg crude product for further reaction as is. LRMS (ESI) m/z 383/385 $[(M+H)]^+$, calc'd for $C_{14}H_{19}BrN_6O_2$: 383.25.


Part B. (S)-2-amino-3-methoxy-1-(4-(3-(2-methoxypyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazin-1-yl)propan-1-one.

3(S)-2-amino-1-(4-(3-bromopyrazolo[1,5-a]pyrimidin-5-yl)piperazin-1-yl)-3-methoxypropan-1-one (400mg, 1.04 mmol), 2-methoxypyridine-3-boronic acid (240mg, 1.56 mmol), potassium carbonate (435mg, 3.12 mmol), $Pd(OAc)_2$ (5mg, 0.021mmol), and x-Phos (14mg, 0.042 mmol) were taken up in 2mL dioxane and 1mL water in a sealed tube and heated at 85°C for 2 hours. Reaction then cooled to room temperature filtered through a celite plug with acetonitrile and DCM, reduced in vacuo. Purified on Shimadzu neutral phase prep lyophilized to get 39mg as the formic acid salt product. ^1H NMR (400 MHz, DMSO- d_6) δ 8.82 (dd, J = 1.89, 7.45 Hz, 1H), 8.76 (d, J = 7.83 Hz, 1H), 8.51 (s, 1H), 8.29 (s, 1H), 7.99 (dd, J = 1.89, 4.93 Hz, 1H), 7.09 (dd, J = 4.93, 7.45 Hz, 1H), 6.84 (d, J = 7.83 Hz, 1H), 3.98 - 4.02 (m, 4H), 3.61 - 3.85 (m, 8H), 3.31 - 3.45 (m, 2H) LRMS (ESI) m/z 412 $[(M+H)]^+$, calc'd for $C_{20}H_{25}N_7O_3$: 411.47.

5.6.29. Synthesis of (S)-2-amino-1-(4-(3-(2-methoxypyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazin-1-yl)-3,3-dimethylbutan-1-one




Part A. (S)-2-amino-1-(4-(3-bromopyrazolo[1,5-a]pyrimidin-5-yl)piperazin-1-yl)-3,3-dimethylbutan-1-one




3-bromo-5-(piperazin-1-yl)pyrazolo[1,5-a]pyrimidine (300mg, 1.1 mmol) N-Boc-L-tert-Leucine (295mg, 1.28mmol), HATU (368mg, 1.6mmol), and triethylamine (590uL, 4.26 mmol) taken up in DMF and stirred at room temperature overnight. Reaction diluted with DCM washed with water, DCM layer then dried over magnesium sulfate filtered and reduced in vacuo. This was then taken up in 26mL DCM and 4mL TFA added. Stirred 1 hour until complete by LC/MS. Reaction washed with 1N NaOH, DCM layer removed dried over magnesium sulfate filtered, reduced in vacuo to obtain 420mg crude product for further reaction as is. LRMS (ESI) m/z 395/397 $[(M+H)]^+$, calc'd for $C_{16}H_{23}BrN_6O$: 395.31.

Part B. (S)-2-amino-1-(4-(3-(2-methoxypyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazin-1-yl)-3,3-dimethylbutan-1-one. (S)-2-amino-1-(4-(3-bromopyrazolo[1,5-a]pyrimidin-5-yl)piperazin-1-yl)-3,3-dimethylbutan-1-one (420mg, 1.1mmol), (2-methoxypyridin-3-yl)boronic acid (240mg, 1.6 mmol), potassium carbonate (435mg, 3.2 mmol), Pd(OAc)₂ (5mg, 0.021mmol), and x-Phos (14mg, 0.043mmol) were taken up in 4mL dioxane and 1mL water under nitrogen and stirred at 85° C for 2 hours. Reaction then cooled to room temperature, filtered through celite plug with acetonitrile and DCM. Reduced in vacuo and purified on shimadzu neutral phase prep, lyophilized to get 18mg 1eq formic salt product. ¹H NMR (400 MHz, DMSO-d₆) δ 8.81 (dd, *J* = 1.77, 7.58 Hz, 1H), 8.75 (d, *J* = 7.83 Hz, 1H), 8.51 (s, 1H), 8.29 (s, 1H), 7.97 - 8.03 (m, 1H), 7.06 - 7.09 (m, 1H), 6.83 (d, *J* = 7.83 Hz, 1H), 4.00 (s, 3H), 3.69 - 3.78 (m, 3H), 3.80 (s, 4H), 3.84 (s, 2H), 0.94 (s, 9H) LRMS (ESI) *m/z* 424 [(M+H)]⁺, calc'd for C₂₂H₂₉N₇O₂: 423.52.

5.6.30. Synthesis of (R)-Tetrahydrofuran-3-yl 4-(3-(2-methoxypyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate




Part A. 4-(3-Bromo-pyrazolo[1,5-a]pyrimidin-5-yl)-piperazine-1-carboxylic acid 4-nitro-phenyl ester



To 2.00 g (7.09 mmol) of 3-bromo-5-piperazin-1-yl-pyrazolo[1,5-a]pyrimidine dissolved in 40 mL of EtOAc at 0°C, was added 1.71 g (8.51 mmol) of 4-nitrophenyl carbonochloridate, followed by 1.97 mL (14.18 mmol) of TEA. After 5 minutes stirring at 0°C, the ice bath was removed, and stirring continued at RT for about 2 hr. It was diluted with 50 mL of 50% EtOAc/hex. The desired product was filtered out, and dried to obtain 3.17 g (100% yield) of an off-white solid.


Part B. 4-(3-Bromo-pyrazolo[1,5-a]pyrimidin-5-yl)-piperazine-1-carboxylic acid (R)-(tetrahydrofuran-3-yl) ester




To 112 mg (0.25 mmol) of 4-nitrophenyl 4-(3-bromopyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate dissolved in 10 mL of THF, was added (R)-tetrahydrofuran-3-ol (44 mg, 0.5 mmol), followed by 60% NaH (40 mg, 1.0 mmol) and stirred at RT for 2 hr. After 2 hr it was diluted with 20 mL of EtOAc and slowly quench with brine. The organic layer was dried over MgSO₄ and concentrated, it was purified over 12 gram silica gel column eluting with 0-10% MeOH/DCM to obtain 71 mg (72% yield) of the desired product.

Part C. (R)-Tetrahydrofuran-3-yl 4-(3-(2-methoxypyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate. Suzuki coupling reaction was carried out under the same conditions as describe about for the synthesis of 6-[1-(2,2-Dimethyl-propyl)-1H-pyrazol-4-yl]-3-(2-methoxy-pyridin-3-yl)-imidazo-[1,2-]pyridazine yielding 72% of the desired product. ¹H NMR (400 MHz, CHLOROFORM-*d*) δ ppm 1.98 - 2.14 (m, 1 H) 2.23 (dtd, *J*=14.02, 8.17, 8.17, 6.27 Hz, 1 H) 3.66 (d, *J*=5.02 Hz, 4 H) 3.71 - 3.82 (m, 4 H) 3.83 - 4.02 (m, 4 H) 4.10 (s, 3 H) 5.27 - 5.39 (m, 1 H) 6.38 (d, *J*=7.78 Hz, 1 H) 7.00 (dd, *J*=7.53, 5.02 Hz, 1 H) 8.04 (dd, *J*=4.77, 1.76 Hz, 1 H) 8.37 (d, *J*=7.78 Hz, 1 H) 8.63 (s, 1 H) 8.73 (dd, *J*=7.53, 1.76 Hz, 1 H).

5.6.31. Synthesis of 1-Methoxypropan-2-yl 4-(3-(2-methoxypyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate



Part A. 1-methoxypropan-2-yl 4-(3-bromopyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate




This was made by using 1-methoxypropan-2-ol and following the Part B in the procedure described above for making (R)-tetrahydrofuran-3-yl 4-(3-bromopyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate.


Part B. 1-Methoxypropan-2-yl 4-(3-(2-methoxypyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-

5 yl)piperazine-1-carboxylate. The Suzuki coupling reaction was carried out by following the procedure that was used for other examples. ¹H NMR (400 MHz, CHLOROFORM-*d*) δ ppm 1.29 (d, *J*=6.57 Hz, 3 H) 3.40 (s, 3 H) 3.42 - 3.56 (m, 2 H) 3.61 - 3.70 (m, 4 H) 3.72 - 3.81 (m, 4 H) 4.10 (s, 3 H) 5.06 (td, *J*=6.25, 4.17 Hz, 1 H) 6.37 (d, *J*=7.83 Hz, 1 H) 7.00 (dd, *J*=7.58, 5.05 Hz, 1 H) 8.03 (dd, *J*=4.80, 1.77 Hz, 1 H) 8.36 (d, *J*=7.83 Hz, 1 H) 8.63 (s, 1 H) 8.73 (dd, *J*=7.45, 1.89 Hz, 1 H).

10 **5.6.32. Synthesis of Tetrahydro-2H-pyran-4-yl 4-(3-(2-methoxypyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate**



Part A. Tetrahydro-2H-pyran-4-yl 4-(3-bromopyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-
carboxylate




15

This was made by using tetrahydro-2H-pyran-4-ol and following the Part B in the procedure described above for making (R)-tetrahydrofuran-3-yl 4-(3-bromopyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate.

Part B. Tetrahydro-2H-pyran-4-yl 4-(3-(2-methoxypyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-

20 yl)piperazine-1-carboxylate. This was synthesized by following the Suzuki procedure described above for making (R)-tetrahydrofuran-3-yl 4-(3-(2-methoxypyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate. ¹H NMR (400 MHz, CHLOROFORM-*d*) δ ppm 1.66 - 1.80 (m, 2 H) 1.95 - 2.02 (m, 2 H) 3.51 - 3.62 (m, 2 H) 3.67 (d, *J*=4.55 Hz, 4 H) 3.71 - 3.82 (m, 4 H) 3.88 - 4.00 (m, 2 H) 4.09 (s, 3 H) 4.94 (tt, *J*=8.43, 3.95 Hz, 1 H) 6.36 (d, *J*=7.58 Hz, 1 H) 6.99 (dd, *J*=7.07, 5.31 Hz, 1 H)
25 8.03 (d, *J*=4.80 Hz, 1 H) 8.35 (d, *J*=7.58 Hz, 1 H) 8.62 (s, 1 H) 8.72 (d, *J*=7.33 Hz, 1 H). LRMS (ESI) *m/e* 439 [(*M* + *H*)⁺, calcd for C₂₂H₂₆FN₆O₄ 438].


5.6.33. Synthesis of (R)-tetrahydrofuran-3-yl 4-(3-(5-fluoro-2-methoxypyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate




(R)-tetrahydrofuran-3-yl 4-(3-bromopyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate

5 (100mg, 0.25mmol), 5-fluoro-2-methoxy-3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)pyridine (127mg, 0.5mmol), triethylamine (211uL, 1.5mmol), Pd(OAc)₂ (1mg, 0.0025mmol), and x-Phos (5mg, 0.005mmol) were taken up in 4mL dioxane and 1mL water under nitrogen and stirred at 85° C for 2 hours. Reaction then cooled to room temperature, filtered through celite plug with acetonitrile and DCM. Reduced in vacuo and purified on Shimadzu neutral phase prep, lyophilized to get 30mg
10 product. ¹H NMR (400 MHz, DMSO-d₆) δ 8.80 (d, *J* = 7.83 Hz, 1H), 8.72 (dd, *J* = 3.03, 10.36 Hz, 1H), 8.56 (s, 1H), 7.94 (d, *J* = 3.03 Hz, 1H), 6.86 (d, *J* = 7.83 Hz, 1H), 5.18 (d, *J* = 1.77 Hz, 1H), 4.00 (s, 3H), 3.71 - 3.83 (m, 8H), 3.32 (s, 2H), 2.08 (s, 1H), 2.13 (s, 1H), 1.89 - 2.00 (m, 1H) LRMS (ESI) *m/z* 443 [(M+H)]⁺, calc'd for C₂₁H₂₃FN₆O₄: 442.45.

5.6.34. Synthesis of (S)-tetrahydrofuran-3-yl 4-(3-(2-methoxypyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate



Part A. (S)-tetrahydrofuran-3-yl 4-(3-bromopyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate




20 To 224 mg (0.50 mmol) of 4-nitrophenyl 4-(3-bromopyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate dissolved in 18 mL of THF, was added (S)-tetrahydrofuran-3-ol (88 mg, 1.0 mmol), followed by 60% NaH (80 mg, 2.0 mmol) and stirred at RT for 2 hr. After 2 hr it was diluted with 30 mL of EtOAc and slowly quench with brine. The organic layer was dried over MgSO₄ and concentrated, it was purified using a 40 g silica gel column eluting with 0-10% MeOH/DCM to obtain

301 mg (76% yield) of the desired product. ¹H NMR (400 MHz, CHLOROFORM-*d*) δ ppm 2.01 - 2.13 (m, 1 H) 2.17 - 2.29 (m, 1 H) 3.57 - 3.70 (m, 4 H) 3.74 - 3.84 (m, 4 H) 3.86 - 4.01 (m, 4 H) 5.27 - 5.38 (m, 1 H) 6.37 (d, *J*=7.83 Hz, 1 H) 7.87 (s, 1 H) 8.28 (d, *J*=7.83 Hz, 1 H)

Part B. (S)-tetrahydrofuran-3-yl 4-(3-(2-methoxypyridin-3-yl)pyrazolo[1,5-*a*]pyrimidin-5-


5 yl)piperazine-1-carboxylate. The Suzuki coupling reaction was carried out under the same conditions as described for the synthesis of LP-943795, yielding 61% of the desired product. ¹H NMR (400 MHz, CHLOROFORM-*d*) δ ppm 2.02 - 2.14 (m, 1 H) 2.16 - 2.30 (m, 1 H) 3.66 (d, *J*=4.80 Hz, 4 H) 3.72 - 3.82 (m, 4 H) 3.84 - 4.01 (m, 4 H) 4.10 (s, 3 H) 5.33 (td, *J*=3.85, 1.89 Hz, 1 H) 6.38 (d, *J*=7.83 Hz, 1 H) 7.01 (dd, *J*=7.45, 4.93 Hz, 1 H) 8.04 (dd, *J*=5.05, 1.77 Hz, 1 H) 8.38 (d, *J*=7.83 Hz, 1 H) 8.64 (s, 1 H) 8.73 (dd, *J*=7.33, 1.77 Hz, 1 H). LRMS (ESI) *m/e* 425 [(*M* + *H*)⁺, calcd for C₂₁H₂₄N₆O₄ 424].

5.6.35. Synthesis of (S)-1-methoxypropan-2-yl 4-(3-(2-methoxypyridin-3-yl)pyrazolo[1,5-*a*]pyrimidin-5-yl)piperazine-1-carboxylate



Part A. (S)-1-methoxypropan-2-yl 4-(3-bromopyrazolo[1,5-*a*]pyrimidin-5-yl)piperazine-1-

15 carboxylate




This was carried out under same conditions and scale as used above [(S)-tetrahydrofuran-3-yl 4-(3-bromopyrazolo[1,5-*a*]pyrimidin-5-yl)piperazine-1-carboxylate] to obtain 71% of the desired product. ¹H NMR (400 MHz, CHLOROFORM-*d*) δ ppm 1.29 (d, *J*=6.27 Hz, 3 H) 3.30 - 3.41 (m, 3 H) 3.41 - 3.54 (m, 2 H) 3.58 - 3.69 (m, 4 H) 3.71 - 3.83 (m, 4 H) 4.98 - 5.12 (m, 1 H) 6.35 (d, *J*=8.03 Hz, 1 H) 7.86 (s, 1 H) 8.26 (d, *J*=7.78 Hz, 1 H).

Part B. (S)-1-methoxypropan-2-yl 4-(3-(2-methoxypyridin-3-yl)pyrazolo[1,5-*a*]pyrimidin-5-yl)piperazine-1-carboxylate. The Suzuki coupling reaction was carried out under the typical

25 conditions used for the synthesis of other examples, yielding 64% of the desired product. ¹H NMR (400 MHz, CHLOROFORM-*d*) δ ppm 1.29 (d, *J*=6.32 Hz, 3 H) 3.40 (s, 3 H) 3.42 - 3.54 (m, 2 H) 3.61 - 3.70 (m, 4 H) 3.71 - 3.81 (m, 4 H) 4.09 (s, 3 H) 5.06 (quind, *J*=6.32, 6.32, 6.32, 6.32, 4.04 Hz, 1 H) 6.38 (d, *J*=7.83 Hz, 1 H) 7.00 (dd, *J*=7.45, 4.93 Hz, 1 H) 8.03 (dd, *J*=4.93, 1.89 Hz, 1 H) 8.36 (d,


J=7.83 Hz, 1 H) 8.63 (s, 1 H) 8.74 (dd, J=7.58, 2.02 Hz, 1 H). LRMS (ESI) m/e 427 [(M + H)⁺, calcd for C₂₁H₂₆N₆O₄ 426].

5.6.36. Synthesis of 2-Methoxyethyl 4-(3-(2-methoxypyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazine(d₈)-1-carboxylate




5

Part A. 3-Bromo-5-(piperazin-1-yl)pyrazolo [1,5-a]pyrimidine(d₈)



A mixture of 3-bromo-5-chloropyrazolo[1,5-a]pyrimidine (930 mg, 4 mmol), piperazine-d₈ HCl (1 g, 6 mmol) and TEA (2.23 mL, 16 mmol) in isopropanol (8 mL) was heated in a microwave oven at 140°C for 30 min. The mixture was concentrated to give the titled compound that was used for next step without further purification.

Part B. 2-Methoxyethyl 4-(3-bromopyrazolo[1,5-a]pyrimidin-5-yl)piperazine(d₈)-1-carboxylate




To a mixture of 3-bromo-5-(piperazin-1-yl)pyrazolo [1,5-a]pyrimidine-d₈ (~ 2 mmol) and DIEA (1.74 mL, 10 mmol) in THF (20 mL) was added methoxyethyl chloroformate (279 µL, 2.4 mmol). The resulting mixture was stirred at rt for overnight. The mixture was concentrated and the residue was subjected to ISCO then prep HPLC to give the titled compound (382 mg). ¹H NMR (400 MHz, CHLOROFORM-d) δ ppm 3.39 (s, 3 H) 3.57 - 3.70 (m, 2 H) 4.24 - 4.31 (m, 2 H) 6.32 (d, J=7.83 Hz, 1 H) 7.84 (s, 1 H) 8.24 (d, J=8.08 Hz, 1 H).

Part C. 2-Methoxyethyl 4-(3-(2-methoxypyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazine(d₈)-1-carboxylate. Under typical Suzuki coupling conditions with 2-methoxyethyl 4-(3-bromopyrazolo[1,5-a]pyrimidin-5-yl)piperazine(d₈)-1-carboxylate and appropriate boronic acid to provide the titled compound. ¹H NMR (400 MHz, CHLOROFORM-d) δ ppm 3.41 (s, 3 H) 3.58 - 3.70 (t, J=3.40 Hz, 2 H) 4.09 (s, 3 H) 4.27 - 4.32 (t, J=3.40 Hz, 2 H) 6.35 (d, J=7.83 Hz, 1 H) 6.99 (dd,

$J=7.58$, 4.80 Hz, 1 H) 8.02 (dd, $J=4.80$, 1.77 Hz, 1 H) 8.35 (d, $J=7.83$ Hz, 1 H) 8.62 (s, 1 H) 8.72 (dd, $J=7.45$, 1.89 Hz, 1 H). LRMS (ESI) m/e 421.2 [(M + H)⁺, calcd for C₂₁H₂₆N₆O₄ 420.5].

5.6.37. Synthesis of Part A. Isopropyl 4-(3-bromopyrazolo[1,5-a]pyrimidin-5-yl)piperazine(d₈)-1-carboxylate



5

To a mixture of 3-bromo-5-(piperazin-1-yl)pyrazolo [1,5-a]pyrimidine-d₈ (~ 2 mmol) and DIEA (1.74 mL, 10 mmol) in THF (20 mL) was added isopropyl chloroformate (2 M solution, 1.2 mL, 2.4 mmol) dropwise. The resulting mixture was stirred at rt for overnight. The mixture was concentrated and the residue was subjected to ISCO to give the titled compound (449 mg). ¹H NMR (400 MHz, CHLOROFORM-*d*) δ ppm 1.26 (d $J = 6.32$ Hz, 6 H) $4.90 - 5.00$ (m, 1 H) 6.31 (d, $J=7.83$ Hz, 1 H) 7.83 (s, 1 H) 8.23 (d, $J=7.83$ Hz, 1 H).

Part B. Isopropyl 4-(3-(2-methoxy-6-methylpyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazine(d₈)-1-carboxylate. Under typical Suzuki coupling conditions with 2-isopropyl 4-(3-bromopyrazolo[1,5-a]pyrimidin-5-yl)piperazine(d₈)-1-carboxylate and appropriate boronic acid to provide the titled compound. ¹H NMR (400 MHz, CHLOROFORM-*d*) δ ppm 1.29 (d, $J=6.32$ Hz, 6 H) 2.48 (s, 3 H) 4.07 (s, 3 H) 4.98 (quin, $J=6.25$ Hz, 1 H) 6.32 (d, $J=7.83$ Hz, 1 H) 6.84 (d, $J=7.58$ Hz, 1 H) 8.33 (d, $J=7.83$ Hz, 1 H) $8.56 - 8.60$ (m, 2 H). LRMS (ESI) m/e 419.2 [(M + H)⁺, calcd for C₂₁H₂₆N₆O₄ 418.5].

5.6.38. P81 Filter Plate Assay

Compounds were serially diluted into a Labcyte LDV plate (Labcyte, cat# LP-0200) using a Mutiprobe (PerkinElmer) and Biomek FX (Beckman Coulter) so that the highest compound concentration was at 96 μ M. Compounds were then pinged (75 nL per well) into a Greiner 384-well reaction plate (Greiner, # 781076) using an ECHO 550 Liquid Handler (Labcyte). A total of 12 μ L reaction buffer (IMAP buffer containing Tween and DTT, from Molecular Devices) was then added to each well of columns 1 and 13 for the negative controls and 12 μ L of 2X AAK1 (0.2 nM full-length human protein, NCBI accession no. NP_055726.2) was added to the remaining wells. Enzyme was then pre-incubated with compound for 10 minutes at RT. Reactions were initiated upon Minitrak (PerkinElmer) addition of 12 μ L substrate mix containing 2X Mu2 (0.2 μ M, full length human protein), 2x cold ATP (2 μ M), and 1.3 μ Ci of hot ³³P-ATP. Reactions proceeded for one hour at RT. Meanwhile, Millipore 384-well P81 filter plates (Millipore, catalog # MZPHN0W10) were placed on a plate washer (Zoom ZW, from Titertek) and pre-wet with 50 μ L 1% phosphoric acid. Kinase reactions were then stopped upon addition of 24 μ L of 2% phosphoric acid to each well and the Minitrak was then used to

transfer 40µl from each well into the pre-wet Millipore 384-well P81 filter plates. Reaction mixtures were incubated for 10 minutes at RT in the P81 plates, followed by washing five times with 100µl/well of 1% phosphoric acid using the Zoom filter washer. The bottom of each filter plate was sealed followed by addition of 20µl Microscint 40 to each well, sealing the top of the plates with Flashplate cover, and then waiting one hour until reading on the TopCount (PerkinElmer).

5.6.39. HEK281 Cell-Based Assay

HEK293F cells were cultured in media containing DMEM (Gibco, cat. #11965), 10% FBS (SAFC Biosciences, cat. #12103C), 1X GPS (glutamine, penicillin and streptomycin). On day one, cells were plated on a 10cm dish so that they are ~80% confluent at time of transfection. Roughly 12 million cells were in a 10cm dish at time of transfection. On day two, each dish was transfected with 48 µg DNA and 144 µl Lipofectamine 2000 (Invitrogen, cat.# 11668-019). The DNA was comprised of a mixture (per 10cm dish) containing 3 µg AAK1/HA/pIRES (full length human, NCBI accession no. NP_055726.2), 45 µg Flag/AP2MI/pcDNA (full length human), and 1.5 ml OPTI-MEM. The Lipofectamine 2000 is made up of a mixture (per 10cm dish) containing 144 µl Lipofectamine 2000 and 1.5 ml OPTI-MEM. Each mixture was transferred to individual 15ml tubes and incubated at RT for 5 minutes, and then the two mixes were combined and incubated at RT for 20 minutes. Growth media was then aspirated from each 10cm plate and replaced with 10ml of DMEM+10% FBS (no GPS). Finally, 3 ml DNA/Lipofectamine mix was added to each 10cm dish and mix gently followed by incubate of plate overnight at 37 °C and 5% CO₂.

On day three, compounds were diluted in 100% DMSO at 1000X final concentration, followed by 3-fold serial dilutions for a total of 5 concentrations tested. Four compounds were tested per 10cm dish. One µl of each compound dilution was then pipetted into a deep-well, 96-well plate, followed by addition of 500 µl DMEM + 0.5% FBS into each well for a 2X final concentration of each compound. Cells were resuspended in a 10cm dish by simple pipetting (HEK293 cells come off the plate that easy at this point) and then transferred to a 50 ml conical tube and pelleted by centrifugation at 1000rpm for 5 min. Cell pellets were then resuspended in 2.75 ml DMEM + 0.5% FBS per 10cm dish and 100 µl of cell suspension transferred into each well of 96-well TC plate. Finally, 100 µl of 2X compound diluted in DMEM + 0.5% FBS was then added into wells containing cell suspension for a 1X final concentration. Plates were then incubated at 37 °C and 5% CO₂ for 3 hours followed by transferring of cell suspensions from each well into 12-tube PCR strips. The PCR strips were spun in a tip rack at 1000rpm for 5 minutes to pellet cells and media was then removed by pipetting without disturbing the cell pellet.

To prepare for Western Blot analysis, cell pellets were resuspend in 40µl 1X LDS-PAGE sample buffer (Invitrogen, cat.# NP0008) + 2X Halt phosphatase and protease inhibitor cocktail (Thermo Scientific, cat.#1861284), followed by sonicating each with microtip sonicator set at 5 for 8-10 seconds. Five µl of 10X NuPage Sample Reducing Agent (with 50 mM DTT) was to each sample followed by heat denaturing at 70C for 10 min on PCR machine. A total of 10µl per sample was

loaded into each lane of a 4-20% Tris-Glycine Criterion 26-well gel (Biorad, cat.# 345-0034) for the phospho-mu2 blot and 10µl per lane in a 4-12% Bis-Tris (+MES buffer) NuPAGE 26-well gel (Invitrogen, cat.# WG1403BX10) for the mu2 blot. For controls, 2ng of phospho-mu2 or 20ng mu2/Flag proteins were loaded in the last well of each gel. After SDS-PAGE, samples on each gel were transferred to PVDF membrane using an iBlot and membranes were blocked for one hour in TBST + 5% milk, followed by wash 3X for 5-10 min with TBST. Criterion gels were probed with rabbit anti-phospho-mu2 (1:5000; a rabbit polyclonal antibody produced by New England Peptide and affinity purified at Lexicon) in TBST + 5% BSA, whereas, NuPAGE gels were probed with mouse anti-Flag (1:500; Sigma, cat.# F1804) in TBST + 5% milk, and these primary antibodies were incubated overnight at 4°C on a rocker.

On day four, Western blots were washed 3X for 5-10 minutes with TBST, probe with anti-rabbit-HRP (1:2000; BioRad, cat.# 170-6515) or anti-mouse-HRP (1:2000; Biorad, cat.# 170-6516) in TBST + 5% milk for 1 hour at RT, washed 3X for 10 minutes with TBST, and developed with ECL reagent (GE Healthcare, cat.# RPN2132) on a Versadoc. Finally, the camera was set up to take a picture every 30 seconds for 10 minutes and the best image saved for each blot with no saturated signal (when the signal is saturated, the bands will be highlighted red). A volume analysis on each band was performed to obtain density values. Percent inhibition was calculated for each sample by first normalizing to total Mu2 expression levels and then comparing to 0% and 100% controls. IC₅₀ values were then calculated using Excel fitting software.

5.6.40. In Vitro Data

In vitro data obtained for various compounds of the invention are provided below in Table 1, wherein "MW" means molecular weight, "P81 Assay" refers to the P81 filter plate assay described above, "CBA" refers to the HEK281 cell-based assay described above, "--" means that results for the given assay were not obtained, "*" means less than or equal to 1.0 µM, "***" means a value of less than or equal to 0.1 µM, and "****" means less than or equal to 0.01 µM.

Table 1

Compound	MW	CBA IC ₅₀ µM	P81 IC ₅₀ µM
(4-(3-(2-methoxypyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazin-1-yl)(pyrrolidin-1-yl)methanone	407.5	**	***
(S)-1-(2-(((3-(2-methoxyphenyl)pyrazolo[1,5-a]pyrimidin-5-yl)amino)methyl)pyrrolidin-1-yl)butan-1-one	393.5	**	***
(S)-1-(2-(((3-(2-methoxyphenyl)pyrazolo[1,5-a]pyrimidin-5-yl)amino)methyl)pyrrolidin-1-yl)-3,3-dimethylbutan-1-one	421.5	***	***
(S)-1-(3,3-dimethylbutyl)-5-(((3-(2-methoxyphenyl)pyrazolo[1,5-a]pyrimidin-5-yl)amino)methyl)pyrrolidin-2-one	435.6	**	***

(S)-1-(3,3-dimethylbutyl)-5-(((3-(2-methoxyphenyl)pyrazolo[1,5-a]pyrimidin-5-yl)amino)methyl)pyrrolidin-2-one	421.5	***	***
(S)-1-(3,3-dimethylbutyl)-5-(((3-(2-methoxypyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)amino)methyl)pyrrolidin-2-one	422.5	**	***
(S)-1-(3,3-dimethylbutyl)-5-(((3-(3-methoxypyridin-4-yl)pyrazolo[1,5-a]pyrimidin-5-yl)amino)methyl)pyrrolidin-2-one	422.5	**	***
(S)-1-(3,3-dimethylbutyl)-5-(((3-phenylpyrazolo[1,5-a]pyrimidin-5-yl)amino)methyl)pyrrolidin-2-one	391.5	—	***
(S)-2-(((3-bromopyrazolo[1,5-a]pyrimidin-5-yl)amino)methyl)-N-(tert-butyl)pyrrolidine-1-carboxamide	395.3	***	***
(S)-2-cyclopropyl-N-methyl-N-(1-(3-(pyridin-2-yl)pyrazolo[1,5-a]pyrimidin-5-yl)pyrrolidin-3-yl)acetamide	376.5	—	**
(S)-3,3,3-trifluoro-1-(2-(((3-(2-methoxyphenyl)pyrazolo[1,5-a]pyrimidin-5-yl)amino)methyl)pyrrolidin-1-yl)propan-1-one	433.4	**	***
(S)-3,3,3-trifluoro-1-(2-(((3-(pyridin-4-yl)pyrazolo[1,5-a]pyrimidin-5-yl)amino)methyl)pyrrolidin-1-yl)propan-1-one	404.4	**	***
(S)-3,3,3-trifluoro-N-(1-(3-(2-methoxyphenyl)pyrazolo[1,5-a]pyrimidin-5-yl)pyrrolidin-3-yl)-N-methylpropanamide	433.4	—	**
(S)-3,3,3-trifluoro-N-(1-(3-(2-methoxypyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)pyrrolidin-3-yl)-N-methylpropanamide	434.4	***	***
(S)-3,3,3-trifluoro-N-methyl-N-(1-(3-(pyridin-2-yl)pyrazolo[1,5-a]pyrimidin-5-yl)pyrrolidin-3-yl)propanamide	404.4	***	***
(S)-5-(((3-bromopyrazolo[1,5-a]pyrimidin-5-yl)amino)methyl)-1-(3,3-dimethylbutyl)pyrrolidin-2-one	394.3	**	***
(S)-ethyl 2-(((3-(2-methoxyphenyl)pyrazolo[1,5-a]pyrimidin-5-yl)amino)methyl)pyrrolidine-1-carboxylate	395.5	***	***
(S)-isopropyl (1-(3-(2-methoxyphenyl)pyrazolo[1,5-a]pyrimidin-5-yl)pyrrolidin-3-yl)(methyl)carbamate	409.5	—	**
(S)-isopropyl (1-(3-(2-methoxypyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)pyrrolidin-3-yl)(methyl)carbamate	410.5	—	**
(S)-isopropyl 2-(((3-(2-methoxyphenyl)pyrazolo[1,5-a]pyrimidin-5-yl)amino)methyl)pyrrolidine-1-carboxylate	409.5	***	***
(S)-isopropyl methyl(1-(3-(pyridin-2-yl)pyrazolo[1,5-a]pyrimidin-5-yl)pyrrolidin-3-yl)carbamate	380.4	—	*
(S)-isopropyl methyl(1-(3-(pyridin-4-yl)pyrazolo[1,5-a]pyrimidin-5-yl)pyrrolidin-3-yl)carbamate	380.4	—	**
(S)-methyl 2-(((3-(2-methoxyphenyl)pyrazolo[1,5-a]pyrimidin-5-yl)amino)methyl)pyrrolidine-1-carboxylate	381.4	**	***

(S)-N-(1-(3-(2-methoxyphenyl)pyrazolo[1,5-a]pyrimidin-5-yl)pyrrolidin-3-yl)-N-methylbutyramide	393.5	*	***
(S)-N-(1-(3-(2-methoxypyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)pyrrolidin-3-yl)-N-methylbutyramide	394.5	*	***
(S)-N-(1-(3-bromopyrazolo[1,5-a]pyrimidin-5-yl)pyrrolidin-3-yl)-3,3,3-trifluoro-N-methylpropanamide	406.2	—	**
(S)-N-(1-(3-bromopyrazolo[1,5-a]pyrimidin-5-yl)pyrrolidin-3-yl)-N-methylpyrrolidine-1-carboxamide	393.3	—	**
(S)-N-(tert-butyl)-2-(((3-(2-methoxyphenyl)pyrazolo[1,5-a]pyrimidin-5-yl)amino)methyl)pyrrolidine-1-carboxamide	422.5	**	***
(S)-N-methyl-N-(1-(3-(pyridin-2-yl)pyrazolo[1,5-a]pyrimidin-5-yl)pyrrolidin-3-yl)butyramide	364.4	—	*
(S)-tert-butyl (1-(3-(2-methoxyphenyl)pyrazolo[1,5-a]pyrimidin-5-yl)pyrrolidin-3-yl)(methyl)carbamate	423.5	**	***
(S)-tert-butyl 2-(((3-(2-(methoxymethyl)phenyl)pyrazolo[1,5-a]pyrimidin-5-yl)amino)methyl)pyrrolidine-1-carboxylate	437.5	**	***
(S)-tert-butyl 2-(((3-(2-ethylphenyl)pyrazolo[1,5-a]pyrimidin-5-yl)amino)methyl)pyrrolidine-1-carboxylate	421.5	**	***
(S)-tert-butyl 2-(((3-(2-hydroxyphenyl)pyrazolo[1,5-a]pyrimidin-5-yl)amino)methyl)pyrrolidine-1-carboxylate	409.5	**	***
(S)-tert-butyl 2-(((3-(2-isopropoxyphenyl)pyrazolo[1,5-a]pyrimidin-5-yl)amino)methyl)pyrrolidine-1-carboxylate	451.6	***	***
(S)-tert-butyl 2-(((3-(2-methoxyphenyl)pyrazolo[1,5-a]pyrimidin-5-yl)amino)methyl)pyrrolidine-1-carboxylate	423.5	***	—
(S)-tert-butyl 2-(((3-(3-methoxyphenyl)pyrazolo[1,5-a]pyrimidin-5-yl)amino)methyl)pyrrolidine-1-carboxylate	423.5	***	***
(S)-tert-butyl 2-(((3-(4-(aminomethyl)phenyl)pyrazolo[1,5-a]pyrimidin-5-yl)amino)methyl)pyrrolidine-1-carboxylate	422.5	***	***
(S)-tert-butyl 2-(((3-(4-carbamoylphenyl)pyrazolo[1,5-a]pyrimidin-5-yl)amino)methyl)pyrrolidine-1-carboxylate	436.5	*	***
(S)-tert-butyl 2-(((3-(4-methoxyphenyl)pyrazolo[1,5-a]pyrimidin-5-yl)amino)methyl)pyrrolidine-1-carboxylate	423.5	***	***
(S)-tert-butyl 2-(((3-(4-methoxypyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)amino)methyl)pyrrolidine-1-carboxylate	424.5	***	***
(S)-tert-butyl 2-(((3-(pyridin-4-yl)pyrazolo[1,5-a]pyrimidin-5-yl)amino)methyl)pyrrolidine-1-carboxylate	394.5	***	—
(S)-tert-butyl 2-(((3-(pyridin-4-yl)pyrazolo[1,5-a]pyrimidin-5-yl)amino)methyl)pyrrolidine-1-carboxylate	394.5	***	—
(S)-tert-butyl 2-(((3-(trifluoromethyl)pyrazolo[1,5-a]pyrimidin-5-yl)amino)methyl)pyrrolidine-1-carboxylate	385.4	**	***
(S)-tert-butyl 2-(((3-iodopyrazolo[1,5-a]pyrimidin-5-yl)amino)methyl)pyrrolidine-1-carboxylate	443.3	**	—

(S)-tert-butyl 2-(((3-phenylpyrazolo[1,5-a]pyrimidin-5-yl)amino)methyl)pyrrolidine-1-carboxylate	393.5	**	***
(S)-tert-butyl methyl(1-(3-(2-methylpyridin-4-yl)pyrazolo[1,5-a]pyrimidin-5-yl)pyrrolidin-3-yl)carbamate	408.5	**	***
(S)-tert-butyl methyl(1-(3-(pyridin-2-yl)pyrazolo[1,5-a]pyrimidin-5-yl)pyrrolidin-3-yl)carbamate	394.5	—	*
1-(4-(3-(2-methoxypyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazin-1-yl)-3-methylbutan-1-one	394.5	**	***
2,2,2-trifluoroethyl 4-(3-(2-methoxypyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate	436.4	***	***
2-fluoroethyl 4-(3-(2-methoxypyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate	400.4	***	***
2-methoxyethyl 4-(3-(2-methoxyphenyl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate	411.5	—	***
2-methoxyethyl 4-(3-(2-methoxypyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate	412.4	**	***
3-(2-methoxypyridin-3-yl)-N-(4,4,4-trifluorobutyl)pyrazolo[1,5-a]pyrimidin-5-amine	351.3	**	***
3-(3-methoxypyridin-4-yl)-N-(2-(trifluoromethoxy)phenethyl)pyrazolo[1,5-a]pyrimidin-5-amine	429.4	***	—
3-(4-(aminomethyl)phenyl)-N-(2-(cyclopentyloxy)ethyl)pyrazolo[1,5-a]pyrimidin-5-amine	351.4	**	—
3-(4-(aminomethyl)phenyl)-N-(2-(neopentyloxy)ethyl)pyrazolo[1,5-a]pyrimidin-5-amine	353.5	**	—
3-(4-(aminomethyl)phenyl)-N-(2-(tert-butoxy)ethyl)pyrazolo[1,5-a]pyrimidin-5-amine	339.4	**	—
3-(4-(aminomethyl)phenyl)-N-(2-(trifluoromethoxy)phenethyl)pyrazolo[1,5-a]pyrimidin-5-amine	427.4	***	—
3-(4-(aminomethyl)phenyl)-N-(2-methoxyethyl)pyrazolo[1,5-a]pyrimidin-5-amine	297.4	**	—
3-(4-(aminomethyl)phenyl)-N-(3-(cyclopentyloxy)propyl)pyrazolo[1,5-a]pyrimidin-5-amine	365.5	**	—
3-(4-(aminomethyl)phenyl)-N-butylpyrazolo[1,5-a]pyrimidin-5-amine	295.4	**	—
3-(4-methoxypyridin-3-yl)-N-(2-(trifluoromethoxy)phenethyl)pyrazolo[1,5-a]pyrimidin-5-amine	429.4	**	—
3-bromo-N-((1-(2,2,2-trifluoroethyl)pyrrolidin-2-yl)methyl)pyrazolo[1,5-a]pyrimidin-5-amine	378.2	**	***
3-bromo-N-(3-(cyclopentyloxy)propyl)pyrazolo[1,5-a]pyrimidin-5-amine	339.2	*	—

4-(5-(butylamino)pyrazolo[1,5-a]pyrimidin-3-yl)-N-(2-(methylamino)ethyl)benzamide	366.5	**	--
5-(4-(isobutylsulfonyl)piperazin-1-yl)-3-(2-methoxypyridin-3-yl)pyrazolo[1,5-a]pyrimidine	430.5		**
cyclopentyl 4-(3-(2-methoxypyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate	422.5	***	***
ethyl 4-(3-(2-ethoxypyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate	396.4	***	***
ethyl 4-(3-(2-methoxypyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate	382.4	**	***
ethyl 5-(4-(isopropoxycarbonyl)piperazin-1-yl)pyrazolo[1,5-a]pyrimidine-3-carboxylate	361.4		**
ethyl methyl(2-((3-(4-(methylcarbamoyl)phenyl)pyrazolo[1,5-a]pyrimidin-5-yl)amino)ethyl)carbamate	396.4	**	--
isobutyl (2-((3-(4-(methylcarbamoyl)phenyl)pyrazolo[1,5-a]pyrimidin-5-yl)amino)ethyl)(methyl)carbamate	410.5	***	--
isobutyl methyl(2-((3-(4-(methylcarbamoyl)phenyl)pyrazolo[1,5-a]pyrimidin-5-yl)amino)ethyl)carbamate	424.5	*	--
isopropyl (1-(3-(2-methoxypyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)azetidin-3-yl)(methyl)carbamate	396.4	--	**
isopropyl (2-((3-(3-methoxypyridin-4-yl)pyrazolo[1,5-a]pyrimidin-5-yl)amino)ethyl)(methyl)carbamate	384.4	***	--
isopropyl (2-((3-(4-(aminomethyl)phenyl)pyrazolo[1,5-a]pyrimidin-5-yl)amino)ethyl)(methyl)carbamate	382.5	**	--
isopropyl (2-((3-(4-(methylcarbamoyl)phenyl)pyrazolo[1,5-a]pyrimidin-5-yl)amino)ethyl)(methyl)carbamate	396.4	***	--
isopropyl (2-((3-(4-methoxypyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)amino)ethyl)(methyl)carbamate	384.4	**	--
isopropyl 4-(3-(1,3,5-trimethyl-1H-pyrazol-4-yl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate	397.5	--	**
isopropyl 4-(3-(1-isobutyl-1H-pyrazol-4-yl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate	411.5	--	**
isopropyl 4-(3-(1-methyl-2-oxo-1,2-dihydropyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate	396.4	**	***
isopropyl 4-(3-(2-(methoxymethyl)phenyl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate	409.5	--	*
isopropyl 4-(3-(2-(methylthio)pyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate	412.5	--	**
isopropyl 4-(3-(2,6-dimethoxypyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate	426.5	***	***
isopropyl 4-(3-(2-aminopyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate	381.4	--	**

isopropyl 4-(3-(2-chloropyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate	400.9	*	**
isopropyl 4-(3-(2-ethoxypyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate	410.5	**	***
isopropyl 4-(3-(2-fluoropyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate	384.4	**	***
isopropyl 4-(3-(2-hydroxypyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate	382.4	**	***
isopropyl 4-(3-(2-isopropoxypyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate	424.5	**	***
isopropyl 4-(3-(2-methoxy-5-methylpyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate	410.5	***	***
isopropyl 4-(3-(2-methoxy-6-methylpyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate	410.5	**	***
isopropyl 4-(3-(2-methoxyphenyl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate	395.5	—	***
isopropyl 4-(3-(2-methoxypyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate	396.4	—	***
isopropyl 4-(3-(2-methoxypyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)-3-oxopiperazine-1-carboxylate	410.4	***	***
isopropyl 4-(3-(2-methoxypyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)-2,2-dimethylpiperazine-1-carboxylate	424.5	—	**
isopropyl 4-(3-(2-methylpyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate	380.4	—	**
isopropyl 4-(3-(3,6-dimethoxypyridazin-4-yl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate	427.5	***	***
isopropyl 4-(3-(3-ethoxyphenyl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate	409.5	—	***
isopropyl 4-(3-(3-fluoro-2-methoxyphenyl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate	413.4	—	**
isopropyl 4-(3-(3-methoxypyridin-2-yl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate	396.4	—	*
isopropyl 4-(3-(3-methoxypyridin-4-yl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate	396.4	**	***
isopropyl 4-(3-(4-fluoropyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate	384.4	—	**
isopropyl 4-(3-(4-methoxypyridin-2-yl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate	396.4	—	***
isopropyl 4-(3-(4-methoxypyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate	396.4	**	***
isopropyl 4-(3-(5-fluoro-2-methoxyphenyl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate	413.4	***	***
isopropyl 4-(3-(5-fluoro-2-methoxypyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate	414.4	**	***

isopropyl 4-(3-(5-fluoropyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate	384.4	—	***
isopropyl 4-(3-(5-methoxypyridin-2-yl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate	396.4	***	***
isopropyl 4-(3-(5-methoxypyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate	396.4	**	***
isopropyl 4-(3-(6-fluoropyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate	384.4	—	**
isopropyl 4-(3-(6-methoxypyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate	396.4	*	***
isopropyl 4-(3-(ethylcarbamoyl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate	360.4	—	**
isopropyl 4-(3-(isopropylcarbamoyl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate	374.4	—	*
isopropyl 4-(3-(methylcarbamoyl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate	346.4	—	**
isopropyl 4-(3-(pyrazin-2-yl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate	367.4	—	**
isopropyl 4-(3-(pyridin-2-yl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate	366.4	**	***
isopropyl 4-(3-(pyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate	366.4	—	***
isopropyl 4-(3-(pyridin-4-yl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate	366.4	**	***
isopropyl 4-(3-(pyrimidin-5-yl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate	367.4	—	**
isopropyl 4-(3-isopropylpyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate	331.4	—	**
isopropyl 4-(pyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate	289.3	—	***
isopropyl methyl(2-((3-(4-((2-(methylamino)ethyl)carbamoyl)phenyl)pyrazolo[1,5-a]pyrimidin-5-yl)amino)ethyl)carbamate	453.5	**	—
isopropyl methyl(2-((3-(4-(methylcarbamoyl)phenyl)pyrazolo[1,5-a]pyrimidin-5-yl)amino)ethyl)carbamate	410.5	***	—
isopropyl methyl(2-((3-(4-(pyrrolidin-2-yl)phenyl)pyrazolo[1,5-a]pyrimidin-5-yl)amino)ethyl)carbamate	422.5	***	—
methyl 4-(3-(2-methoxypyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate	368.4	**	**
N-(2-(cyclopentyloxy)ethyl)-3-(3-methoxypyridin-4-yl)pyrazolo[1,5-a]pyrimidin-5-amine	353.4	**	—

N-(2-(tert-butoxy)ethyl)-3-(3-methoxypyridin-4-yl)pyrazolo[1,5-a]pyrimidin-5-amine	341.4	**	--
N-(2-aminoethyl)-4-(5-((2-methoxyethyl)amino)pyrazolo[1,5-a]pyrimidin-3-yl)benzamide	354.4	*	--
N-(2-aminoethyl)-4-(5-((3,3-dimethylbutyl)amino)pyrazolo[1,5-a]pyrimidin-3-yl)benzamide	380.5	*	--
N-(2-aminoethyl)-4-(5-(butylamino)pyrazolo[1,5-a]pyrimidin-3-yl)benzamide	352.4	**	--
N-(2-methoxyethyl)-3-phenylpyrazolo[1,5-a]pyrimidin-5-amine	268.3	*	--
N-(3-(2-methoxypyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)-N-(4,4,4-trifluorobutyl)acetamide	393.4	--	*
N-(3-(cyclopentyloxy)propyl)-3-(3-methoxypyridin-4-yl)pyrazolo[1,5-a]pyrimidin-5-amine	367.4	**	--
N-(tert-butyl)-4-(3-(2-methoxypyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxamide	409.5	**	***
N-(tert-butyl)-4-(3-(2-methoxypyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)-N-methylpiperazine-1-carboxamide	423.5	**	***
N-isopropyl-4-(3-(2-methoxypyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxamide	395.5	***	***
N-isopropyl-4-(3-(2-methoxypyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)-N-methylpiperazine-1-carboxamide	409.5	**	***
tert-butyl (1-(3-(2-methoxypyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)azetidin-3-yl)(methyl)carbamate	410.5	--	*
tert-butyl (2-((3-(3-methoxypyridin-4-yl)pyrazolo[1,5-a]pyrimidin-5-yl)amino)ethyl)(methyl)carbamate	398.5	***	--
tert-butyl (2-((3-(4-carbamoylphenyl)pyrazolo[1,5-a]pyrimidin-5-yl)amino)ethyl)(methyl)carbamate	410.5	***	--
tert-butyl (2-(4-(5-((2-methoxyethyl)amino)pyrazolo[1,5-a]pyrimidin-3-yl)benzamido)ethyl)carbamate	454.5	*	--
tert-butyl 4-(3-(2-methoxypyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate	410.5	***	***
tert-butyl methyl(2-((3-(4-(methylcarbamoyl)phenyl)pyrazolo[1,5-a]pyrimidin-5-yl)amino)ethyl)carbamate	424.5	***	--
(4-(3-(2-methoxypyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazin-1-yl)(piperidin-1-yl)methanone	421.5	**	***
isopropyl 4-(3-(6-methoxypyridin-2-yl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate	396.4	**	***
isopropyl 4-(3-(5-fluoro-2-methoxy-4-methylphenyl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate	427.5	***	***

isopropyl 4-(3-(2,4-dimethoxypyrimidin-5-yl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate	427.5	**	***
isopropyl 4-(3-(3-fluoro-4-methylphenyl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate	397.4	--	**
isopropyl 4-(3-iodopyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate	415.2	--	**
(S)-isopropyl 2-((methyl(pyrazolo[1,5-a]pyrimidin-5-yl)amino)methyl)pyrrolidine-1-carboxylate	317.4	--	*
(S)-isopropyl 2-(((3-(2-methoxypyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)(methyl)amino)methyl)pyrrolidine-1-carboxylate	424.5	***	***
isopropyl 4-(3-(2-methoxypyridin-4-yl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate	396.4	--	**
(E)-isopropyl 4-(3-(3-methoxyprop-1-en-1-yl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate	359.4	--	*
(E)-isopropyl 4-(3-(2-ethoxyvinyl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate	359.4	--	**
isopropyl 4-(3-chloropyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate	323.8	--	**
isopropyl 4-(3-(2-d ₃ -methoxypyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate	399.4	***	***
(S)-isopropyl 2-(((3-(2-methoxypyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)amino)methyl)pyrrolidine-1-carboxylate	410.5	***	***
tert-butyl 4-(3-bromopyrazolo[1,5-a]pyrimidin-5-yl)-d ₈ -piperazine-1-carboxylate	390.2	--	**
isopropyl 4-(3-(2-methoxypyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)-d ₈ -piperazine-1-carboxylate	404.4	***	***
isopropyl 4-(3-(1,6-dimethyl-2-oxo-1,2-dihydropyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate	410.5	--	***
isopropyl 3-(3-(2-methoxypyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)-2-oxoimidazolidine-1-carboxylate	396.4	--	*
isopropyl 4-(3-(1H-benzo[d]imidazol-4-yl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate	405.5	--	**
2-isobutyl-7-(3-(2-methoxypyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)hexahydroimidazo[1,5-a]pyrazin-3(2H)-one	421.5	--	***
(1R,5S)-tert-butyl 3-(3-(2-methoxypyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)-3,6-diazabicyclo[3.1.1]heptane-6-carboxylate	422.5	--	**
isopropyl 4-(3-(2-methoxy-4-methylphenyl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate	409.5	**	***
(1R,5S)-isopropyl 3-(3-(2-methoxypyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)-3,6-diazabicyclo[3.1.1]heptane-6-carboxylate	408.5	--	**

isopropyl (2-((3-(2-methoxypyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)(methyl)amino)ethyl)(methyl)carbamate	398.5	--	**
isopropyl 4-(3-(2-(2-methoxyethoxy)phenyl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate	439.5	--	**
isopropyl 4-(3-(2-(2-methoxyethoxy)pyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate	440.5	--	**
ethyl 4-(3-(2-methoxy-5-methylpyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate	396.4	**	***
isopropyl (2-((3-(2-methoxypyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)amino)ethyl)carbamate	370.4	--	**
isopropyl (2-((3-(2-methoxy-6-methylpyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)(methyl)amino)ethyl)(methyl)carbamate	412.5	--	**
isopropyl (2-((3-(2-methoxy-5-methylpyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)(methyl)amino)ethyl)(methyl)carbamate	412.5	--	**
isopropyl (2-((3-(5-fluoro-2-methoxypyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)(methyl)amino)ethyl)(methyl)carbamate	416.4	--	***
ethyl 4-(3-(5-fluoro-2-methoxypyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate	400.4	**	***
ethyl 4-(3-(5-fluoro-2-methoxyphenyl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate	399.4	--	***
ethyl 4-(3-(2-methoxy-6-methylpyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate	396.4	**	***
ethyl 4-(3-(2,6-dimethoxypyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate	412.4	--	***
isopropyl (3-((3-(2-methoxypyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)(methyl)amino)propyl)(methyl)carbamate	412.5	**	***
isopropyl (3-((3-(2-methoxy-6-methylpyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)(methyl)amino)propyl)(methyl)carbamate	426.5	**	***
ethyl 4-(3-(2-d ₃ -methoxypyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate	385.4	**	***
isopropyl 4-(3-(2-(2-(dimethylamino)ethoxy)pyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate	453.5	--	**
isopropyl (3-((3-(5-fluoro-2-methoxypyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)(methyl)amino)propyl)(methyl)carbamate	430.5	--	***
N-(2-(tert-butoxy)ethyl)-3-(2-methoxypyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-amine	341.4	--	***
N-(2-(tert-butoxy)ethyl)-3-(2-methoxy-6-methylpyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-amine	355.4	--	**
tert-butyl (2-((3-(2-methoxypyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)oxy)ethyl)(methyl)carbamate	399.4	--	***

isopropyl (2-((3-(2-methoxypyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)oxy)ethyl)(methyl)carbamate	385.4	--	**
(S)-isopropyl (1-(3-(2,6-dimethoxypyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)pyrrolidin-3-yl)(methyl)carbamate	440.5	--	**
(S)-isopropyl (1-(3-(5-fluoro-2-methoxypyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)pyrrolidin-3-yl)(methyl)carbamate	428.5	--	**
(S)-isopropyl (1-(3-(2-methoxy-5-methylpyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)pyrrolidin-3-yl)(methyl)carbamate	424.5	--	***
(S)-isopropyl (1-(3-(1,6-dimethyl-2-oxo-1,2-dihydropyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)pyrrolidin-3-yl)(methyl)carbamate	424.5	--	*
(S)-isopropyl (1-(3-(2-d ₃ -methoxy-6-methylpyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)pyrrolidin-3-yl)(methyl)carbamate	424.5	**	***
(S)-isopropyl (1-(3-(2-methoxypyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)pyrrolidin-3-yl)(methyl)carbamate	413.4	**	***
isopropyl 4-(3-(imidazo[1,2-a]pyridin-5-yl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate	405.5	--	**
(S)-isopropyl (1-(3-(5-fluoro-2-methoxyphenyl)pyrazolo[1,5-a]pyrimidin-5-yl)pyrrolidin-3-yl)(methyl)carbamate	427.5	**	***
isopropyl 4-(3-(imidazo[1,2-a]pyridin-8-yl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate	405.5	--	**
propyl 4-(3-(2-methoxypyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate	396.4	**	***
propyl 4-(3-(2-methoxy-6-methylpyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate	410.5	***	***
propyl 4-(3-(5-fluoro-2-methoxypyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate	414.4	**	***
3-(2-methoxypyridin-3-yl)-5-(4-(3,3,3-trifluoropropyl)piperazin-1-yl)pyrazolo[1,5-a]pyrimidine	406.4	--	*
3-(5-fluoro-2-methoxypyridin-3-yl)-5-(4-(3,3,3-trifluoropropyl)piperazin-1-yl)pyrazolo[1,5-a]pyrimidine	424.4	--	*
3-bromo-5-(4-(2,2,2-trifluoroethyl)piperazin-1-yl)pyrazolo[1,5-a]pyrimidine	364.2	--	**
3-(2-methoxypyridin-3-yl)-5-(4-(2,2,2-trifluoroethyl)piperazin-1-yl)pyrazolo[1,5-a]pyrimidine	392.4	--	**
3-(2-methoxy-6-methylpyridin-3-yl)-5-(4-(2,2,2-trifluoroethyl)piperazin-1-yl)pyrazolo[1,5-a]pyrimidine	406.4	--	**
3-(5-fluoro-2-methoxypyridin-3-yl)-5-(4-(2,2,2-trifluoroethyl)piperazin-1-yl)pyrazolo[1,5-a]pyrimidine	410.4	--	*

tert-butyl 4-(3-(2-d ₃ -methoxypyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate	413.4	***	***
tert-butyl 4-(3-(2-methoxy-6-methylpyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate	424.5	***	***
tert-butyl 4-(3-(2-ethoxypyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate	424.5	***	***
tert-butyl 4-(3-(5-fluoro-2-methoxypyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate	428.5	**	***
tert-butyl 4-(3-(5-fluoro-2-methoxyphenyl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate	427.5	***	***
N-(3-(tert-butoxy)propyl)-3-(2-methoxypyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-amine	355.4	**	***
N-(3-(tert-butoxy)propyl)-3-(2-methoxy-6-methylpyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-amine	369.5	—	***
3-bromo-N-(3-(tert-butoxy)propyl)pyrazolo[1,5-a]pyrimidin-5-amine	327.2	—	*
isopropyl 4-(3-(trifluoromethyl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate	357.3	—	*
isopropyl 4-(3-(pyridazin-4-yl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate	367.4	—	*
N-(3-(tert-butoxy)propyl)-3-(2-methoxy-6-methylpyridin-3-yl)-N-methylpyrazolo[1,5-a]pyrimidin-5-amine	383.5	—	**
N-(3-(tert-butoxy)propyl)-3-(2-methoxypyridin-3-yl)-N-methylpyrazolo[1,5-a]pyrimidin-5-amine	369.5	—	**
5-isopropyl-3-(4-(3-(2-methoxypyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazin-1-yl)-1,2,4-oxadiazole	420.5	—	**
(S)-2-amino-1-(4-(3-(2-methoxypyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazin-1-yl)-4-methylpentan-1-one	423.5	***	***
isopropyl 4-(3-(4-(2-aminoethoxy)phenyl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate	424.5	**	***
(S)-2-amino-N-((S)-1-(3-(2-methoxypyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)pyrrolidin-3-yl)-N,4-dimethylpentanamide	437.5	***	***
(S)-2-amino-N-(2-((3-(2-methoxypyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)(methyl)amino)ethyl)-N,4-dimethylpentanamide	425.5	—	***
(S)-2-amino-1-(4-(3-bromopyrazolo[1,5-a]pyrimidin-5-yl)piperazin-1-yl)-4-methylpentan-1-one	395.3	—	**
isopropyl 4-(3-(4-(methoxycarbonyl)phenyl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate	423.5	—	**
(S)-2-amino-1-(4-(3-(5-fluoro-2-methoxyphenyl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazin-1-yl)-4-methylpentan-1-one	440.5	***	***

(S)-2-amino-4-methyl-1-(4-(3-phenylpyrazolo[1,5-a]pyrimidin-5-yl)piperazin-1-yl)pentan-1-one	392.5	***	***
isopropyl 4-(3-(4-(aminomethyl)phenyl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate	394.5	***	***
(S)-2-amino-1-(4-(3-(2-methoxypyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazin-1-yl)-3-methylbutan-1-one	409.5	***	***
(R)-2-amino-1-(4-(3-(2-methoxypyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazin-1-yl)-4-methylpentan-1-one	423.5	—	**
isopropyl 4-(3-(2-fluoro-6-methoxypyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate	414.4	—	**
(S)-2-amino-1-(4-(3-(2-fluorophenyl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazin-1-yl)-3-methylbutan-1-one	396.5	***	***
(S)-2-amino-1-(4-(3-chloropyrazolo[1,5-a]pyrimidin-5-yl)piperazin-1-yl)-4-methylpentan-1-one	350.8	—	**
(S)-2-amino-1-(4-(3-fluoropyrazolo[1,5-a]pyrimidin-5-yl)piperazin-1-yl)-4-methylpentan-1-one	334.4	—	*
2-methoxyethyl 4-(3-(5-fluoro-2-methoxyphenyl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate	429.4	***	***
2-methoxyethyl 4-(3-(5-fluoro-2-methoxypyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate	430.4	**	***
2-methoxyethyl 4-(3-(2-methoxy-6-methylpyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate	426.5	**	***
(S)-1-(4-(3-(2-methoxypyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazin-1-yl)-4-methyl-2-(methylamino)pentan-1-one	437.5	***	***
(S)-2-amino-3-methoxy-1-(4-(3-(2-methoxypyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazin-1-yl)propan-1-one	411.5	***	***
(1-aminocyclopentyl)(4-(3-(2-methoxypyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazin-1-yl)methanone	421.5	***	***
(S)-2-amino-1-(4-(3-(2-methoxypyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazin-1-yl)-3,3-dimethylbutan-1-one	423.5	***	***
2-amino-1-(4-(3-(2-methoxypyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazin-1-yl)-2-methylpropan-1-one	395.5	—	**
(S)-(4-(3-(2-methoxypyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazin-1-yl)(pyrrolidin-2-yl)methanone	407.5	**	***
2-methoxyethyl 4-(3-(2-ethoxypyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate	426.5	—	***
(S)-2-amino-1-(4-(3-(3-fluorophenyl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazin-1-yl)-3-methylbutan-1-one	396.5	***	***
(S)-2-amino-3-methyl-1-(4-(3-phenylpyrazolo[1,5-a]pyrimidin-5-yl)piperazin-1-yl)butan-1-one	378.5	***	***

(S)-2-amino-1-(4-(3-(4-fluorophenyl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazin-1-yl)-3-methylbutan-1-one	396.5	***	***
(S)-2-amino-1-(4-(3-(2,5-difluorophenyl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazin-1-yl)-3-methylbutan-1-one	414.5	***	***
(S)-2-amino-1-(4-(3-(2-fluoro-4-methylphenyl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazin-1-yl)-3-methylbutan-1-one	410.5	**	***
(S)-2-amino-1-(4-(3-(2-methoxyphenyl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazin-1-yl)-3-methylbutan-1-one	408.5	***	***
3-methoxypropyl 4-(3-(2-methoxypyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate	426.5	—	***
2-ethoxyethyl 4-(3-(2-methoxypyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate	426.5	—	**
2-(2-methoxyethoxy)ethyl 4-(3-(2-methoxypyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate	456.5	—	**
isopropyl 4-(3-(4-methoxypyrimidin-5-yl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate	397.4	—	**
(S)-2-amino-3,3-dimethyl-1-(4-(3-phenylpyrazolo[1,5-a]pyrimidin-5-yl)piperazin-1-yl)butan-1-one	392.5	**	***
(S)-2-amino-1-(4-(3-(2-methoxyphenyl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazin-1-yl)-3,3-dimethylbutan-1-one	422.5	***	***
(S)-2-amino-1-(4-(3-(5-fluoro-2-methoxyphenyl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazin-1-yl)-3,3-dimethylbutan-1-one	440.5	***	***
(S)-1-(4-(3-(2-methoxypyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazin-1-yl)-3,3-dimethyl-2-(methylamino)butan-1-one	437.5	**	***
(S)-1-(4-(3-(2-methoxypyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazin-1-yl)-4-methylpentan-2-amine	409.5	—	**
(S)-2-amino-1-(4-(3-(5-fluoro-2-methoxyphenyl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazin-1-yl)-3-methylbutan-1-one	426.5	***	***
(1-(3-(2-methoxypyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)-1H-imidazol-4-yl)methanol	322.3	—	**
(R)-tetrahydrofuran-3-yl 4-(3-(2-methoxypyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate	424.5	***	***
1-methoxypropan-2-yl 4-(3-(2-methoxypyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate	426.5	***	***
tetrahydro-2H-pyran-4-yl 4-(3-(2-methoxypyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate	438.5	**	***
1-methoxy-2-methylpropan-2-yl 4-(3-(2-methoxypyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate	440.5	*	***
5-(4-(methoxymethyl)-1H-imidazol-1-yl)-3-(2-methoxypyridin-3-yl)pyrazolo[1,5-a]pyrimidine	336.3	—	**

(2S,3R)-2-amino-3-methoxy-1-(4-(3-(2-methoxypyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazin-1-yl)butan-1-one	425.5	**	***
(S)-(4-(3-(2-methoxypyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazin-1-yl)(morpholin-3-yl)methanone	423.5	—	***
(3-aminotetrahydrofuran-3-yl)(4-(3-(2-methoxypyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazin-1-yl)methanone	423.5	**	***
2-(dimethylamino)ethyl 4-(3-(2-methoxypyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate	425.5	—	**
2-(tert-butoxy)ethyl 4-(3-(2-methoxypyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate	454.5	—	***
1,3-dimethoxypropan-2-yl 4-(3-(2-methoxypyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate	456.5	—	**
(R)-tetrahydrofuran-3-yl 4-(3-(5-fluoro-2-methoxypyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate	442.4	***	***
(R)-tetrahydrofuran-3-yl 4-(3-(5-fluoro-2-methoxyphenyl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate	441.5	***	***
(R)-tetrahydrofuran-3-yl 4-(3-(2-methoxy-6-methylpyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate	438.5	**	***
oxetan-3-yl 4-(3-(2-methoxypyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate	410.4	***	***
(S)-tetrahydrofuran-3-yl 4-(3-(2-methoxypyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate	424.5	***	***
(S)-1-methoxypropan-2-yl 4-(3-(2-methoxypyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate	426.5	***	***
(R)-1-methoxypropan-2-yl 4-(3-(2-methoxypyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate	426.5	*	***
2-methoxy-2-oxoethyl 4-(3-(2-methoxypyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate	426.4	—	**
isopropyl 4-(3-bromopyrazolo[1,5-a]pyrimidin-5-yl)-d ₈ -piperazine-1-carboxylate	376.2	—	**
2-methoxyethyl 4-(3-(2-methoxypyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)-d ₈ -piperazine-1-carboxylate	420.4	**	***
isopropyl 4-(3-(2-methoxy-6-methylpyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)-d ₈ -piperazine-1-carboxylate	418.4	***	***
N-(3-(2-methoxypyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)butyramide	311.3	—	**

5.6.41. Pharmacological Effects

Studies of AAK1 knockout mice showed that disruption of the AAK1 gene affects pain response as measured using the formalin paw test. See Example 5.6.1, above. The same test was used to confirm that the administration of an AAK1 inhibitor can also affect pain response.

5 Mice were tested for nociception with Automatic Nociception Analyzers (purchased from the Ozaki lab at University of California, San Diego). A metal band was placed around the left hind paw of each mouse with superglue 30 minutes prior to testing. After the 30-minute acclimation period, 20 μ l of 5% formalin was subcutaneously injected in the dorsal surface of the left hind paw. Mice were individually housed in cylindrical chambers for 45 minutes. Fresh 5 % formalin solution was
10 prepared by diluting formaldehyde (Formalde-fresh 20%, Fisher Scientific, Fair Lawn, NJ) with distilled water. Investigatory compounds were administered 30 minutes prior to formalin injection.

A computer software recorded flinches per minute, total flinches for Phase I (acute phase = first 8 minutes), and total flinches for Phase II (tonic phase between 20 - 40 minutes) through an electromagnetic field. See Yaksh TL, Ozaki G, McCumber D, Rathbun M, Svensson C, Malkmus S,
15 Yaksh MC. *An automated flinch detecting system for use in the formalin nociceptive bioassay.* J Appl Physiol., 2001; 90:2386-402.

Various compounds of the invention were tested at different doses. Gabapentin and pregabalin were used as positive controls. Results are shown below in Table 2, wherein “*” means an effect equal to or greater than 50 percent of that of gabapentin at 200 mpk, “**” means an
20 effect equal to or greater than 100 percent of that of gabapentin at 200 mpk, “sc” means subcutaneous administration, and “po” means oral administration.

Table 2

Compound	Dose (mpk)	Effect
Gabapentin	50 sc	*
Gabapentin	200 sc	**
Pregabalin	50 sc	*
(S)-tert-butyl 2-(((3-(2-methoxyphenyl)pyrazolo[1,5-a]pyrimidin-5-yl)amino)methyl)pyrrolidine-1-carboxylate	10 sc	*
(S)-tert-butyl 2-(((3-(2-methoxyphenyl)pyrazolo[1,5-a]pyrimidin-5-yl)amino)methyl)pyrrolidine-1-carboxylate	60 sc	*
(S)-tert-butyl 2-(((3-(pyridin-4-yl)pyrazolo[1,5-a]pyrimidin-5-yl)amino)methyl)pyrrolidine-1-carboxylate	30 sc	**
isopropyl 4-(3-(2-methoxyphenyl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate	30 sc	**
isopropyl 4-(3-(2-methoxypyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate	10 po	*
isopropyl 4-(3-(2-methoxypyridin-3-yl)pyrazolo[1,5-a]pyrimidin-5-yl)piperazine-1-carboxylate	30 po	**


These results demonstrate that AAK1 inhibitors can be used to treat pain.

All publications (e.g., patents and patent applications) cited above are incorporated herein by reference in their entireties.

CLAIMS

What is claimed is:

1. A method of inhibiting adaptor associated kinase 1 (AAK1) activity, comprising contacting AAK1 with a compound of the formula:



or a pharmaceutically acceptable salt thereof, wherein:

R₁ is R_{1A} or optionally substituted C₁₋₁₂ hydrocarbyl or 2-12-membered heterocarbyl, which optional substitution is with one or more R_{1A};

each R_{1A} is independently -OR_{1C}, -N(R_{1C})₂, -C(O)R_{1C}, -C(O)OR_{1C}, -C(O)N(R_{1C})₂, -N(R_{1C})C(O)OR_{1C}, cyano, halo, or optionally substituted C₁₋₁₂ hydrocarbyl or 2-12-membered heterocarbyl, which optional substitution is with one or more R_{1B};

each R_{1B} is independently -OR_{1C}, -N(R_{1C})₂, -C(O)R_{1C}, -C(O)OR_{1C}, -C(O)N(R_{1C})₂, -N(R_{1C})C(O)OR_{1C}, cyano or halo;

each R_{1C} is independently hydrogen or optionally substituted C₁₋₁₂ hydrocarbyl or 2-12-membered heterocarbyl, which optional substitution is with one or more of cyano, halo or hydroxyl;

R₂ is -NR_{2A}R_{2B}, wherein R_{2A} is hydrogen and R_{2B} is optionally substituted C₁₋₁₂ hydrocarbyl or 2-12-membered heterocarbyl, which optional substitution is with one or more R_{2C};


or R_{2A} and R_{2B} are taken together to form a 4-7-membered heterocycle optionally substituted with one or more R_{2C};

each R_{2C} is independently -OR_{2D}, -N(R_{2D})₂, -C(O)R_{2D}, -C(O)OR_{2D}, -C(O)N(R_{2D})₂, -N(R_{2D})C(O)OR_{2D}, cyano, halo, or optionally substituted C₁₋₁₂ hydrocarbyl or 2-12-membered heterocarbyl, which optional substitution is with one or more amino, cyano, halo, hydroxyl, or R_{2D};

each R_{2D} is independently hydrogen or optionally substituted C₁₋₁₂ hydrocarbyl or 2-12-membered heterocarbyl, which optional substitution is with one or more amino, cyano, halo, or hydroxyl; and

R₃ is hydrogen or C₁₋₆ alkyl optionally substituted with one or more cyano, halo or hydroxyl.

2. A method of treating or managing a disease or disorder mediated by AAK1 activity, which comprises administering to a patient in need thereof a therapeutically effective amount of a compound of the formula:



or a pharmaceutically acceptable salt thereof, wherein:

R₁ is R_{1A} or optionally substituted C₁₋₁₂ hydrocarbyl or 2-12-membered heterocarbyl, which optional substitution is with one or more R_{1A};

each R_{1A} is independently -OR_{1C}, -N(R_{1C})₂, -C(O)R_{1C}, -C(O)OR_{1C}, -C(O)N(R_{1C})₂, -N(R_{1C})C(O)OR_{1C},
5 cyano, halo, or optionally substituted C₁₋₁₂ hydrocarbyl or 2-12-membered heterocarbyl, which optional substitution is with one or more R_{1B};

each R_{1B} is independently -OR_{1C}, -N(R_{1C})₂, -C(O)R_{1C}, -C(O)OR_{1C}, -C(O)N(R_{1C})₂, -N(R_{1C})C(O)OR_{1C},
cyano or halo;

each R_{1C} is independently hydrogen or optionally substituted C₁₋₁₂ hydrocarbyl or 2-12-
10 membered heterocarbyl, which optional substitution is with one or more cyano, halo or hydroxyl;

R₂ is -NR_{2A}R_{2B}, wherein R_{2A} is hydrogen and R_{2B} is optionally substituted C₁₋₁₂ hydrocarbyl or 2-12-membered heterocarbyl, which optional substitution is with one or more R_{2C};

or R_{2A} and R_{2B} are taken together to form a 4-7-membered heterocycle optionally substituted
with one or more R_{2C};

each R_{2C} is independently -OR_{2D}, -N(R_{2D})₂, -C(O)R_{2D}, -C(O)OR_{2D}, -C(O)N(R_{2D})₂, -N(R_{2D})C(O)OR_{2D},
15 cyano, halo, or optionally substituted C₁₋₁₂ hydrocarbyl or 2-12-membered heterocarbyl, which optional substitution is with one or more with one or more amino, cyano, halo, hydroxyl, or R_{2D};

each R_{2D} is independently hydrogen or optionally substituted C₁₋₁₂ hydrocarbyl or 2-12-
membered heterocarbyl, which optional substitution is with one or more amino, cyano, halo, or
20 hydroxyl; and

R₃ is hydrogen or C₁₋₆ alkyl optionally substituted with one or more of cyano, halo or hydroxyl.

3. The method of claim 2, wherein the disease or disorder is Alzheimer's disease, bipolar disorder, pain, Parkinson's disease, or schizophrenia.

4. The method of claim 3, wherein the pain is neuropathic pain.

25 5. The method of claim 4, wherein the neuropathic pain is fibromyalgia or peripheral neuropathy (e.g., diabetic neuropathy).

6. The method of any of the preceding claims, wherein R₁ is R_{1A}.


7. The method of any of the preceding claims, wherein R₁ is optionally substituted C₁₋₁₂ hydrocarbyl.

30 8. The method of claim 7, wherein R₁ is optionally substituted phenyl.

9. The method of any of the preceding claims, wherein R₁ is optionally substituted 2-12-membered heterocarbyl (e.g., 2-8 membered heterocarbyl, 2-6 membered heterocarbyl, 2-6 membered heterocarbyl).

10. The method of claim 9, wherein R₁ is optionally substituted pyridinyl, thiophen, or
35 imidazol.

11. The method of any of the preceding claims, wherein R_{1A} is halo.
12. The method of any of the preceding claims, wherein R_{1A} is $-OR_{1C}$, $-N(R_{1C})_2$, $-C(O)R_{1C}$, $-C(O)OR_{1C}$, or $-C(O)N(R_{1C})_2$.
13. The method of any of the preceding claims, wherein R_{1A} is $-OR_{1C}$.
- 5 14. The method of any of the preceding claims, wherein R_{1B} is $-N(R_{1C})_2$, $-OR_{1C}$, halo.
15. The method of claim 1 or 2, wherein R_{2A} and R_{2B} are taken together to form a 4-7-membered heterocycle optionally substituted with one or more R_{2C} .
16. The method of any of the preceding claims, wherein R_{1C} is hydrogen.
17. The method of any of the preceding claims, wherein R_{1C} is C_{1-12} hydrocarbyl (e.g., C_{1-6} hydrocarbyl, C_{1-4} hydrocarbyl such as methyl, ethyl, propyl).
- 10 18. The method of any of the preceding claims, wherein R_{2C} is $-C(O)OR_{2D}$, $-C(O)N(R_{2D})_2$, or $-N(R_{2D})C(O)OR_{2D}$.
19. The method of any of the preceding claims, wherein R_{2C} is $-C(O)R_{2D}$.
20. The method of any of the preceding claims, wherein R_{2D} is hydrogen.
- 15 21. The method of any of the preceding claims, wherein R_{2D} is C_{1-12} hydrocarbyl (e.g., C_{1-6} hydrocarbyl, C_{1-4} hydrocarbyl such as methyl, ethyl, propyl).
22. The method of any of the preceding claims, wherein R_{2D} is 2-12-membered heterocarbyl comprising at least one nitrogen atom.
23. The method of any of the preceding claims, wherein at least one R_{2D} is optionally substituted C_{1-12} hydrocarbyl, which optional substitution is with one or more of amino, cyano, halo, hydroxyl.
- 20 24. The method of any of the preceding claims, wherein R_{2D} is 2-12-membered heterocarbyl comprising at least one nitrogen atom.
25. The method of any of the preceding claims, wherein R_3 is hydrogen.
- 25 26. The method of claim 1 or 2, wherein the compound is of the formula:



or a pharmaceutically acceptable salt thereof, wherein:

A is cyclic C_{1-12} hydrocarbyl or 4-7-membered heterocycle;

D is 4-7-membered heterocycle;

each R_{1A} is independently $-OR_{1C}$, $-N(R_{1C})_2$, $-C(O)R_{1C}$, $-C(O)OR_{1C}$, $-C(O)N(R_{1C})_2$, $-N(R_{1C})C(O)OR_{1C}$, cyano, halo, or optionally substituted C_{1-12} hydrocarbyl or 2-12-membered heterocarbyl, which optional substitution is with one or more R_{1B} ;

each R_{1B} is independently $-OR_{1C}$, $-N(R_{1C})_2$, $-C(O)R_{1C}$, $-C(O)OR_{1C}$, $-C(O)N(R_{1C})_2$, $-N(R_{1C})C(O)OR_{1C}$,
5 cyano or halo;

each R_{1C} is independently hydrogen or optionally substituted C_{1-12} hydrocarbyl or 2-12-membered heterocarbyl, which optional substitution is with one or more of cyano, halo or hydroxyl;

each R_{2C} is independently $-OR_{2D}$, $-N(R_{2D})_2$, $-C(O)R_{2D}$, $-C(O)OR_{2D}$, $-C(O)N(R_{2D})_2$, $-N(R_{2D})C(O)OR_{2D}$, cyano, halo, or optionally substituted C_{1-12} hydrocarbyl or 2-12-membered heterocarbyl, which
10 optional substitution is with one or more with one or more amino, cyano, halo, hydroxyl, or R_{2D} ;

each R_{2D} is independently hydrogen or optionally substituted C_{1-12} hydrocarbyl or 2-12-membered heterocarbyl, which optional substitution is with one or more of amino, cyano, halo, or hydroxyl;

n is 1-3; and

15 m is 0-3.

27. The method of claim 26, with the provision that D is not piperidinyl.

28. The method of claim 26, with the provision that R_{2C} is not $-N(R_{2D})_2$.


29. The method of claim 26, with the provision that A is not phenyl.

30. The method of claim 26, with the provision that m is 1.

20 31. The method of claim 26, with the provision that m is 2.

32. The method of claim 26, with the provision that R_{2D} is not ethyl.

33. A compound of the formula:



or a pharmaceutically acceptable salt thereof, wherein:

25 A is cyclic C_{1-12} hydrocarbyl or 4-7-membered heterocycle;

D is 4-7-membered heterocycle;

each R_{1A} is independently $-OR_{1C}$, $-N(R_{1C})_2$, $-C(O)R_{1C}$, $-C(O)OR_{1C}$, $-C(O)N(R_{1C})_2$, $-N(R_{1C})C(O)OR_{1C}$, cyano, halo, or optionally substituted C_{1-12} hydrocarbyl or 2-12-membered heterocarbyl, which optional substitution is with one or more R_{1B} ;

30 each R_{1B} is independently $-OR_{1C}$, $-N(R_{1C})_2$, $-C(O)R_{1C}$, $-C(O)OR_{1C}$, $-C(O)N(R_{1C})_2$, $-N(R_{1C})C(O)OR_{1C}$, cyano or halo;

each R_{1C} is independently hydrogen or optionally substituted C_{1-12} hydrocarbyl or 2-12-membered heterocarbyl, which optional substitution is with one or more of cyano, halo or hydroxyl;

each R_{2C} is independently $-OR_{2D}$, $-N(R_{2D})_2$, $-C(O)R_{2D}$, $-C(O)OR_{2D}$, $-C(O)N(R_{2D})_2$, $-N(R_{2D})C(O)OR_{2D}$, cyano, halo, or optionally substituted C_{1-12} hydrocarbyl or 2-12-membered heterocarbyl, which

5 optional substitution is with one or more with one or more amino, cyano, halo, hydroxyl, or R_{2D} ;

each R_{2D} is independently hydrogen or optionally substituted C_{1-12} hydrocarbyl or 2-12-membered heterocarbyl, which optional substitution is with one or more of amino, cyano, halo, hydroxyl;

n is 1-3; and

10 m is 0-3;

with the proviso that when D is piperidinyl, A is phenyl, and R_{2C} is $-N(R_{2D})_2$, R_{2D} is not ethyl.


34. The compound of claim 33, wherein D is piperazin or pyrrolidin.

35. The compound of claim 33, wherein n is 1.

36. The compound of claim 33, wherein m is 1.

15 37. The compound of claim 33, wherein A is pyridinyl, thiophen, or imidazol.


38. The compound of claim 33, which is of the formula:



wherein X is CH or N.

39. The compound of claim 38, wherein X is N and m is 1 or 2.

20 40. A compound of the formula:



or a pharmaceutically acceptable salt thereof, wherein:

X is CH or N;

25 each R_{1A} is independently $-OR_{1C}$, $-N(R_{1C})_2$, $-C(O)R_{1C}$, $-C(O)OR_{1C}$, $-C(O)N(R_{1C})_2$, $-N(R_{1C})C(O)OR_{1C}$, cyano, halo, or optionally substituted C_{1-12} hydrocarbyl or 2-12-membered heterocarbyl, which optional substitution is with one or more R_{1B} ;

each R_{1B} is independently $-OR_{1C}$, $-N(R_{1C})_2$, $-C(O)R_{1C}$, $-C(O)OR_{1C}$, $-C(O)N(R_{1C})_2$, $-N(R_{1C})C(O)OR_{1C}$, cyano or halo;

each R_{1C} is independently hydrogen or optionally substituted C_{1-12} hydrocarbyl or 2-12-membered heterocarbyl, which optional substitution is with one or more of cyano, halo or hydroxyl;

each R_{2C} is independently $-OR_{2D}$, $-N(R_{2D})_2$, $-C(O)R_{2D}$, $-C(O)OR_{2D}$, $-C(O)N(R_{2D})_2$, $-N(R_{2D})C(O)OR_{2D}$, cyano, halo, or optionally substituted C_{1-12} hydrocarbyl or 2-12-membered heterocarbyl, which


5 optional substitution is with one or more with one or more amino, cyano, halo, hydroxyl, or R_{2D} ;

each R_{2D} is independently hydrogen or optionally substituted C_{1-12} hydrocarbyl or 2-12-membered heterocarbyl, which optional substitution is with one or more of amino, cyano, halo, or hydroxyl; and


m is 0-3.

10 41. The compound of claim 40, with the provision that R_{2C} is not optionally substituted phenyl or pyridinyl.

42. The compound of claim 40, which is of the formula:




43. The compound of claim 42, which is of the formula:




15


44. The compound of claim 43, which is of the formula:



45. The compound of claim 43, which is of the formula:



46. A compound of the formula:



or a pharmaceutically acceptable salt thereof, wherein:

X is CH or N;

5 each R_{1A} is independently $-OR_{1C}$, $-N(R_{1C})_2$, $-C(O)R_{1C}$, $-C(O)OR_{1C}$, $-C(O)N(R_{1C})_2$, $-N(R_{1C})C(O)OR_{1C}$, cyano, halo, or optionally substituted C_{1-12} hydrocarbyl or 2-12-membered heterocarbyl, which optional substitution is with one or more R_{1B} ;

each R_{1B} is independently $-OR_{1C}$, $-N(R_{1C})_2$, $-C(O)R_{1C}$, $-C(O)OR_{1C}$, $-C(O)N(R_{1C})_2$, $-N(R_{1C})C(O)OR_{1C}$, cyano or halo;


10 each R_{1C} is independently hydrogen or optionally substituted C_{1-12} hydrocarbyl or 2-12-membered heterocarbyl, which optional substitution is with one or more of cyano, halo or hydroxyl;

each R_{2C} is independently $-OR_{2D}$, $-N(R_{2D})_2$, $-C(O)R_{2D}$, $-C(O)OR_{2D}$, $-C(O)N(R_{2D})_2$, $-N(R_{2D})C(O)OR_{2D}$, cyano, halo, or optionally substituted C_{1-12} hydrocarbyl or 2-12-membered heterocarbyl, which optional substitution is with one or more with one or more amino, cyano, halo, hydroxyl, or R_{2D} ;

15 each R_{2D} is independently hydrogen or optionally substituted C_{1-12} hydrocarbyl or 2-12-membered heterocarbyl, which optional substitution is with one or more of amino, cyano, halo, hydroxyl; and


m is 0-3.

47. The compound of claim 46, which is of the formula:




20

48. The compound of claim 47, which is of the formula:



49. The compound of claim 47, which is of the formula:



50. The compound of any of claims 33-49, wherein at least one R_{1A} is halo.

51. The compound of any of claims 33-49, wherein at least one R_{1A} is -OR_{1C}.

5 52. The compound of claim 51, wherein R_{1C} is optionally substituted C₁₋₁₂ hydrocarbyl (e.g., C₁₋₆ hydrocarbyl, C₁₋₄ hydrocarbyl).

53. The compound of any of claims 33-49, wherein R_{2C} is -C(O)OR_{2D}, -C(O)N(R_{2D})₂, or -N(R_{2D})C(O)OR_{2D}.

54. The compound of any of claims 33-49, wherein R_{2C} is -C(O)R_{2D}.

10 55. The compound of claim 53 or 54, wherein each R_{2D} is independently hydrogen or C₁₋₁₂ hydrocarbyl (e.g., C₁₋₆ hydrocarbyl, C₁₋₄ hydrocarbyl).

56. The compound of claim 53 or 54, wherein at least one R_{2D} is optionally substituted C₁₋₁₂ hydrocarbyl, which optional substitution is with one or more of amino, cyano, halo, hydroxyl.

15 57. The compound of claim 53 or 54, wherein R_{2D} is 2-12-membered heterocarbyl comprising at least one nitrogen atom.

58. A pharmaceutical composition comprising a compound of any of claims 33-57 and a pharmaceutically acceptable excipient or diluent.

59. A method of inhibiting AAK1 activity, which comprises contacting AAK1 with a compound of any of claims 33-57.

20 60. A method of treating or managing a disease or disorder mediated by AAK1 activity, which comprises administering to a patient in need thereof a therapeutically effective amount of a compound of any of claims 33-57, or a pharmaceutical composition of claim 58.

61. The method of claim 60, wherein the disease or disorder is Alzheimer's disease, bipolar disorder, pain, Parkinson's disease, or schizophrenia.

25 62. The method of claim 61, wherein the pain is neuropathic pain.

63. The method of claim 62, wherein the neuropathic pain is fibromyalgia or peripheral neuropathy (e.g., diabetic neuropathy).