

NOVEL ADENO-ASSOCIATED VIRUS CAPSID PROTEINS

[0001] This application claims priority benefit of United States Provisional Application No. 62/366,383, filed July 26, 2017, which is incorporated by reference herein in its entirety.

FIELD OF INVENTION

[0002] The invention relates to novel adeno-associated virus (AAV) capsid proteins, AAV particles comprising a novel capsid protein, polynucleotides encoding these capsid proteins and AAV vectors expressing these capsid proteins. The invention also relates to methods of making the herein described AAV vectors expressing the novel capsid proteins of the invention and associated therapeutic uses thereof.

BACKGROUND

[0003] AAV is a replication-deficient parvovirus, the single-stranded DNA genome of which is about 4.7 kb in length including 145 nucleotide inverted terminal repeat (ITRs). The nucleotide sequence of the AAV serotype 2 (AAV2) genome is presented in Srivastava *et al.*, *J. Virol.*, 45: 555-564 (1983) as corrected by Ruffing *et al.*, *J. Gen. Virol.*, 75: 3385-3392 (1994). Cis-acting sequences directing viral DNA replication (rep), encapsidation/packaging and host cell chromosome integration are contained within the ITRs. Three AAV promoters, p5, p19, and p40 (named for their relative map locations), drive the expression of the two AAV internal open reading frames encoding rep and cap genes. The two rep promoters (p5 and p19), coupled with the differential splicing of the single AAV intron (at nucleotides 2107 and 2227), result in the production of four rep proteins (rep 78, rep 68, rep 52, and rep 40) from the rep gene. Rep proteins possess multiple enzymatic properties which are ultimately responsible for replicating the viral genome. The cap gene is expressed from the p40 promoter and it encodes the three capsid proteins VP1, VP2, and VP3. Alternative splicing and a non-consensus translational start site are responsible for the production of the three related capsid proteins. A single consensus polyadenylation site is located at map position 95 of the AAV genome. The life cycle and genetics of AAV are reviewed in Muzyczka, *Current Topics in Microbiology and Immunology*, 158: 97-129 (1992).

[0004] When AAV infects a human cell, the viral genome can integrate into chromosome 19 resulting in latent infection of the cell. Production of infectious virus does not occur unless the

cell is infected with a helper virus (for example, adenovirus or herpesvirus). In the case of adenovirus, genes E1A, E1B, E2A, E4 and VA provide helper functions. Upon infection with a helper virus, the AAV provirus is rescued and amplified, and both AAV and adenovirus are produced.

[0005] AAV possesses unique features that make it attractive as a vaccine vector for expressing immunogenic peptides/polypeptides and as a vector for delivering foreign DNA to cells, for example, in gene therapy. AAV infection of cells in culture is non-cytopathic, and natural infection of humans and other animals is silent and asymptomatic. Moreover, AAV infects many mammalian cells allowing the possibility of targeting many different tissues *in vivo*. The AAV proviral genome is infectious as cloned DNA in plasmids which makes construction of recombinant genomes feasible. Furthermore, because the signals directing AAV replication, genome encapsidation and integration are contained within the ITRs of the AAV genome, some or all of the internal approximately 4.3 kb of the genome (encoding replication and structural capsid proteins, rep-cap) may be replaced with foreign DNA such as a gene cassette containing a promoter, a DNA of interest and a polyadenylation signal. The rep and cap proteins may be provided in trans. Another significant feature of AAV is that it is an extremely stable and hearty virus. It easily withstands the conditions used to inactivate adenovirus (56° to 65°C for several hours), making cold preservation of rAAV-vectors less critical. AAV may even be lyophilized. Finally, AAV-infected cells are not resistant to superinfection.

[0006] AAV vectors find use in numerous mammalian gene therapy applications and there is a need for new and/or modified AAV vectors and associated virus that find use in gene therapy applications. The present invention provides for novel AAV vectors expressing the novel AAV capsid proteins or the functional chimeric engineered AAV capsid proteins of the present invention, and novel, non-naturally occurring AAV virions comprising those vectors or capsid proteins.

SUMMARY OF INVENTION

[0007] The invention provides for novel AAV capsid proteins, which may be novel VP1, VP2 or VP3 capsid proteins, non-naturally occurring AAV virus comprising any of these capsid proteins, and use of such AAV virus for gene therapy applications and for use in the preparation

of medicaments for gene therapy applications. In some embodiments, the AAV capsid proteins were isolated and identified from various mammalian tissues. The amino acid sequences of certain novel mammalian-derived AAV capsid VP1 proteins are set out as SEQ ID NOS:15-89, and the associated locations of the respective VP2 and VP3 sequences are also herein described. In addition, the invention provides for novel engineered chimeric AAV capsid proteins which have a backbone amino acid sequence derived from one AAV capsid sequence and fragments of capsid protein sequence derived from at least one different AAV capsid sequence. The amino acid sequences of exemplary engineered chimeric AAV capsid VP1 proteins are set out as SEQ ID NOS:90-157. Collectively, the novel capsid proteins are referred to herein as "AAV capsid proteins of the invention." The term "non-naturally occurring" when used in regards to any composition of matter described herein means that the composition is not a product of nature, but rather is artificially synthesized by recombinant or other means.

[0008] In one embodiment the invention provides a vector and an adeno-associated virus (AAV) having a capsid protein having an amino acid sequence that is at least 95% identical to (i) any one of SEQ ID NOS:15-89, (ii) the VP2 region of any one of SEQ ID NOS:15-89, or (iii) the VP3 region of any one of SEQ ID NOS:15-89, and further having a transgene where the transgene is composed of a heterologous gene operably linked to regulatory sequences that control expression of the heterologous gene in a host cell. In another embodiment the capsid protein has the amino acid sequence of i) any one of SEQ ID NOS:15-89, (ii) the VP2 region of any one of SEQ ID NOS:15-89, or (iii) the VP3 region of any one of SEQ ID NOS:15-89. In yet another embodiment the AAV has an AAV inverted terminal repeat sequence. In further embodiments the AAV are mixed with a physiologically compatible carrier.

[0009] In another embodiment the invention provides a vector and an AAV having a chimeric capsid protein where the chimeric capsid protein has a VP1 amino acid sequence of a recipient backbone AAV capsid having variable regions I, II, III, IV, V, VI, VII, VIII, and IX, except where one or more of the variable regions I, II, III, IV, V, VI, VII, VIII, and IX is replaced by the corresponding variable region from one or more donor AAV capsids. In another embodiment, only one variable region of the recipient capsid is replaced by the corresponding variable region from the donor capsid. In a further embodiment, two or more variable regions of the recipient capsid are replaced by the corresponding variable regions from a single donor AAV capsid. In yet another embodiment, two or more variable regions of the recipient AAV capsid are replaced

by the corresponding variable regions from two or more donor AAV capsids. In another embodiment, all nine variable regions of the recipient AAV capsid are replaced by the corresponding variable regions from a single donor capsid. In yet another embodiment the recipient AAV capsid has a GBS region or a GH loop region and the GBS region or the GH loop region is replaced by the corresponding region from one or more donor AAV capsids. In a further embodiment all nine variable regions and the GBS region of the recipient AAV capsid are replaced by the corresponding variable regions and GBS region from one or more donor AAV capsids. In yet another embodiment all nine variable regions and the GBS region of the recipient AAV capsid are replaced by the corresponding regions and GBS region from two or more donor AAV capsids. In another embodiment the GH loop of the recipient AAV capsid is replaced by the corresponding GH loop region from a donor AAV capsid. In a further embodiment all nine variable regions and the GH loop region of the recipient AAV capsid are replaced by the corresponding variable regions and GH loop region from one or more donor AAV capsids. In one embodiment the recipient AAV capsid sequence is any one of SEQ ID NOS:1-14 and the donor AAV capsid sequences are selected from any one of SEQ ID NOS:1-14 and where the recipient AAV capsid and the donor AAV capsid are different. In another embodiment the recipient AAV capsid sequence is any one of SEQ ID NOS:1-89 and the donor AAV capsid sequences are selected from any one of SEQ ID NOS:1-89 and where the recipient AAV capsid and the donor AAV capsid are different. In yet another embodiment the chimeric capsid has the amino acid sequence of any one of SEQ ID NOS:90-157.

[0010] In another embodiment, the invention provides a method of delivering a transgene to a cell involving the step of contacting the cell with any AAV disclosed herein. In another embodiment, the invention provides a method of treating a subject from a disorder or disease associated with abnormal activity of an endogenous protein involving the step of administering to the subject an effective amount of an AAV disclosed herein where the AAV has a transgene that encodes a biologically active copy of the protein. In yet another embodiment, the methods involve delivering a transgene to a muscle cell. In a further embodiment, the disease is Duchenne muscular dystrophy and the transgene encodes a microdystrophin.

[0011] In an embodiment, the invention provides a composition comprising a vector or AAV disclosed herein for delivery of a transgene to a cell. In another embodiment, the invention provides a composition comprising an effective amount of a vector or AAV disclosed herein for

the treatment of a disorder or disease associated with abnormal activity of an endogenous protein, wherein the vector of AAV has a transgene that encodes a biologically active copy of the protein. In yet another embodiment, the composition delivers a transgene to a muscle cell. In a further embodiment, the disease is Duchenne muscular dystrophy and the transgene encodes a microdystrophin.

[0012] The invention also provides use of a vector or AAV disclosed herein for the preparation of a medicament effective to treat a subject suffering from a disorder or disease associated with abnormal activity of an endogenous protein, wherein the vector or AAV has a transgene that encodes a biologically active copy of the protein. In yet another embodiment, the medicament delivers a transgene to a muscle cell. In a further embodiment the disease is Duchenne muscular dystrophy and the transgene encodes a microdystrophin.

[0013]

[0014] The invention provides for fragments of any of the AAV capsid proteins of the invention that retain a biological activity of an AAV capsid protein. Exemplary fragments include VP2 and VP3 spliced variants of the capsid proteins, and fragments comprising one or more of the variable regions (VR) of the capsid protein and/or the glycan binding sequence (GBS) of a capsid protein and/or the GH loop. The invention also provides for novel, non-naturally occurring AAV particles comprising a capsid protein fragment and those comprising a capsid protein fragment having at least 90, 91, 92, 93, 94, 95, 96, 97, 98 or 99% sequence identity to a specifically defined capsid protein fragment.

[0015] In one embodiment, the invention provides for an isolated adeno-associated virus (AAV) capsid protein, wherein the capsid protein comprises (i) an amino acid sequence that is at least 95%, 96%, 97%, 98%, or 99% identical to the VP1 amino acid sequence of any one of SEQ ID NOS:15-89 or the VP2 or VP3 region of any one of SEQ ID NOS:15-89 or (ii) a VP1 amino acid sequence comprising any one of SEQ ID NOS:15-89 or the VP2 or VP3 region of any one of SEQ ID NOS:15-89. In certain embodiments, the capsid protein is linked to a heterologous amino acid sequence. The invention also provides for non-naturally occurring AAV particles having or comprising any of these capsid proteins. In certain embodiments, the non-naturally occurring AAV particle comprising any of the above described VP1, VP2 or VP3 capsid proteins comprises a nucleic acid having AAV inverted terminal repeats and a transgene comprising a

heterologous gene operably linked to regulatory sequences which direct expression of the heterologous gene in a host cell. In other embodiments, the non-naturally occurring AAV particle comprising any of the VP1, VP2 or VP3 capsid sequences described herein comprises a heterologous transgene operably linked to regulatory sequences that control transgene expression in a host cell. As used herein, the terms "heterologous gene" or "heterologous regulatory sequence" means that the referenced gene or regulatory sequence is not naturally present in the AAV vector or particle and is artificially introduced therein. The term "transgene" refers to a nucleic acid that comprises both a heterologous gene and regulatory sequences that are operably linked to the heterologous gene that control expression of that gene in a host cell.

[0016] The invention also provides for a polynucleotide comprising a nucleotide sequence encoding an adeno-associated virus (AAV) capsid protein, wherein the capsid protein comprises (i) an amino acid sequence that is at least 95%, 96%, 97%, 98% or 99% identical to the VP1 amino acid sequence of any one of SEQ ID NOS:15-89 or the VP2 or VP3 region of any one of SEQ ID NOS:15-89 or (ii) a VP1 amino acid sequence comprising any one of SEQ ID NOS:15-89 or the VP2 or VP3 region of any one of SEQ ID NOS:15-89, wherein the polynucleotide is operatively linked to a heterologous regulatory control sequence. As such, it is understood that the polynucleotides of the present invention are non-naturally occurring. The invention also provides for AAV vectors comprising any of these polynucleotide sequences operably linked to a heterologous regulatory sequence and compositions comprising these AAV vectors, including pharmaceutical compositions.

[0017] In another embodiment, the invention provides for an isolated adeno-associated virus (AAV) vector comprising a polynucleotide sequence encoding a capsid protein and a heterologous transgene sequence, wherein the capsid protein comprises (i) an amino acid sequence that is at least 95%, 96%, 97%, 98% or 99% identical to the VP1 amino acid sequence of any one of SEQ ID NOS:15-89 or the VP2 or VP3 region of any one of SEQ ID NOS:15-89 or (ii) a VP1 amino acid sequence comprising any one of SEQ ID NOS:15-89 or the VP2 or VP3 region of any one of SEQ ID NOS:15-89. The invention also provides for compositions comprising these AAV vectors, including pharmaceutical compositions.

[0018] The present invention is also based upon the novel finding described herein that AAV VP1 capsid sequences comprise nine different variable regions, a GBS region and a GH loop

region, and that replacing one of more of these regions in one AAV VP1 capsid sequence with the corresponding region(s) from an at least second, different AAV VP1 capsid sequence can generate novel chimeric AAV capsids whose associated AAVs are functional, are capable of transducing cells and delivering heterologous transgenes, and that have unique properties that may be recombinantly engineered into the chimeric AAV. In regards to the various variable regions and the GBS and GH loop regions, the term "corresponding" means the same region between two different AAV capsid sequences. For example, the region "corresponding" to VR I in a first AAV capsid sequence is the same region (i.e., the VR I region) in a second different AAV capsid sequence. The term "chimeric" in relation to an AAV capsid sequence refers to the fact that the AAV capsid sequence of interest comprises amino acid sequences derived from two or more different AAV capsid sequences.

[0019] As such, the present invention also provides for an isolated, non-naturally occurring chimeric adeno-associated virus (AAV) capsid protein, wherein the chimeric capsid protein comprises an amino acid sequence derived from a first AAV capsid sequence having at least one variable region substituted with a variable region from a second AAV capsid sequence that is different from the first AAV capsid sequence. The first AAV capsid sequence (referred to herein as the "recipient") provides the backbone amino acid sequence into which one or more variable regions are swapped or substituted by one or more variable regions from the second AAV capsid sequence (referred to herein as the "donor"). The second AAV capsid sequence is different from the first AAV capsid sequence and will provide the sequence of the variable region(s) which is/are substituted or inserted into the sequence of the backbone or recipient capsid sequence. The invention also provides for non-naturally occurring AAV virus or AAV particles that comprise any of the chimeric capsid proteins herein described. In certain embodiments, the non-naturally occurring AAV particles that comprise any of the chimeric capsid proteins herein described also comprise a heterologous transgene operably linked to regulatory sequences that control transgene expression in a host cell.

[0020] The "variable regions" that may be swapped from a donor AAV capsid sequence into a recipient backbone capsid sequence refer to the nine variable regions within the VP1 sequence of an AAV capsid protein. The variable regions (VR) are referred to herein as VR I, VR II, VR III, VR IV, VR V, VR VI, VR VII, VR VIII and VR IX and their respective locations in various VP1 sequences are herein described. The VR exhibit the highest sequence and structural variation

within the AAV VP1 capsid sequence and may also have roles in receptor attachment, transcriptional activation of transgenes, tissue transduction and antigenicity.

[0021] The “glycan binding sequence (GBS)” or “GBS domain” or “GBS region” refer to the amino acid sequence located between VR IV and VR V that governs the glycan binding specificity of the viral capsid. The locations of the GBS regions in various AAV VP1 amino acid sequences are herein described, and those from other AAV VP1 amino acid sequences are known in the art and/or may be routine identified.

[0022] The “GH loop” refers to a loop sequence that is flanked by β -strand G and β -strand H within the internal β -barrel of the capsid protein. The “GH loop” sequence comprises variable region VR IV through VR VIII, including the encompassed GBS sequence and all interspersed conserved backbone sequence from the donor. The locations of the GH loop regions in various AAV VP1 amino acid sequences are herein described and those from other AAV VP1 amino acid sequences may be routinely identified.

[0023] In regard to the herein described locations of the VR, GBS and GH loop regions, it is noted that the location of the N-terminal and/or C-terminal ends of those regions may vary by from up to 1 amino acid, 2 amino acids, 3 amino acids, 4 amino acids or 5 amino acids from the amino acid locations of those regions as they are explicitly described herein (particularly in Table 2). Novel capsid sequence comprising substituted VR, GBS and/or GH loop region(s) that vary from up to 5 amino acids on the N-terminal and/or C-terminal end as herein defined are encompassed by the present invention.

[0024] In one embodiment, the invention provides for an isolated, non-naturally occurring, chimeric adeno-associated virus (AAV) capsid protein, wherein the chimeric capsid protein comprises an amino acid sequence derived from a first AAV capsid sequence having at least one variable region substituted by a variable region derived from an at least second different AAV capsid sequence. In certain embodiments, the non-naturally occurring capsid protein is a VP1 capsid protein. In other embodiments, the chimeric capsid protein further comprises a GBS domain and/or a GH loop region from an AAV capsid sequence differing from the first recipient AAV capsid sequence. For example, the chimeric AAV capsid proteins of the invention have a backbone sequence derived from a first AAV capsid sequence (recipient) and at least one substituted variable region from a second different AAV capsid sequence (donor). In certain

embodiments, the chimeric AAV capsid proteins of the present invention have one, two, three, four, five, six, seven, eight or all nine variable regions substituted by the respective variable region(s) from one or more donor AAV capsid sequence(s) that differ from the first recipient capsid sequence. In other embodiments, the AAV capsid proteins of the invention have a GBS domain or GH loop region sequence derived from a donor capsid sequence that differs from the recipient capsid sequence. Alternatively, the chimeric AAV capsids of the invention have at least one substitute variable region and a GBS from the same AAV capsid sequence which differs from the first AAV capsid sequence. The invention also provides non-naturally occurring AAV particles comprising any of the chimeric AAV capsid proteins described herein. Such AAVs may also comprise a heterologous transgene operably linked to a regulatory sequence controlling expression of the transgene in a host cell.

[0025] In addition, the invention provides for isolated AAV capsid proteins, wherein the capsid protein comprises an amino acid sequence from a first AAV capsid sequence which has two variable regions substituted with the respective variable regions from at least one AAV capsid sequence that differs from the first AAV capsid sequence, or least three variable regions substituted with the respective variable regions from at least one AAV capsid sequence that differs from the first AAV capsid sequence, or least four variable regions substituted with the respective variable regions from at least one AAV capsid sequence that differs from the first AAV capsid sequence, or least five variable regions substituted with the respective variable regions from at least one AAV capsid sequence that differs from the first AAV capsid sequence, or least six variable regions substituted with the respective variable regions from at least one AAV capsid sequence that differs from the first AAV capsid sequence, or least seven variable regions substituted with the respective variable regions from at least one AAV capsid sequence that differs from the first AAV capsid sequence, or least eight variable regions substituted with the respective variable regions from at least one AAV capsid sequence that differs from the first AAV capsid sequence, or all nine of the variable regions substituted with the respective variable regions from at least one AAV capsid sequence that differs from the first AAV capsid sequence. For example, the substituted variable region(s) are from the same AAV capsid sequence or the substituted variable regions are from two or more different AAV capsid sequences that differ from the first AAV capsid sequence. In addition, in any of these AAV capsid proteins of the

invention, the GBS and/or the GH loop are also substituted and may be derived from any AAV donor capsid sequence that differs from the first AAV capsid sequence.

[0026] In any of the chimeric AAV capsids of the invention, the first/recipient AAV capsid sequence and the second/donor AAV capsid sequence can be any known or herein described AAV capsid sequence including, for example, capsid sequences associated with the following AAV sequences: AAV-1, AAV-2, AAV-3, AAV-3B, AAV-4, AAV-5, AAV-6, AAV-7, AAV-8, AAV-9, AAV-10, AAV-11, AAV-12, AAV-13, AAVbo, AAVmo, AAV6.2, AAVRH.8, AAV4.10, AAVanc80L65 or AAVanc110, or any of the other AAV serotypes or capsid sequences herein described. Production of pseudotyped rAAV is disclosed in, for example, WO 01/83692. Other types of rAAV variants, for example rAAV with capsid mutations, are also contemplated. See, for example, Marsic et al., *Molecular Therapy*, 22(11): 1900-1909 (2014).

[0027] In certain embodiments, the chimeric AAV capsid proteins of the invention may comprise the amino acid sequence of any one of SEQ ID NOS:90-157, each of which have at least one variable region from a donor AAV serotype swapped for the respective variable region(s) in the recipient backbone sequence.

[0028] In any of the chimeric AAV capsid proteins of the invention, the backbone sequence or the amino acid sequence from the first AAV capsid sequence derives from the amino acid sequence of any of SEQ ID NO:1-73. In addition, in any of the chimeric AAV capsid proteins of the invention, the donor sequence or the amino acid sequence from the second AAV serotype derives from the amino acid sequence of one or more variable regions, GBS domain and/or GH loop of any of SEQ ID NO:1-73.

[0029] In another embodiment, the invention provides for an isolated polynucleotide sequence comprising a nucleotide sequence encoding any of the engineered chimeric AAV capsid proteins of the invention. In addition, the invention provides for isolated AAV vectors comprising these polynucleotide sequences and AAV vectors comprising a polynucleotide sequence encoding any of the chimeric AAV capsid proteins of the invention. The invention also provides for compositions comprising these AAV vectors, including pharmaceutical compositions. The invention also provides for AAV virus comprising any of the herein described non-naturally occurring chimeric AAV capsid proteins.

[0030] The invention provides for methods of producing a recombinant adeno-associated virus (AAV) particle comprising the steps of: culturing a cell that has been transfected with any of the AAV vectors of the invention and recovering recombinant AAV particle from the supernatant of the transfected cell. In addition, the invention provides for viral particles comprising any of the viral vectors or capsid proteins of the invention and cells comprising these viral vectors.

[0031] One embodiment of the invention provides a method of producing any of the recombinant AAV described herein by culturing a viral production cell into which has been introduced a first nucleic acid vector having 5' and 3' AAV inverted terminal repeat sequences flanking a transgene having a heterologous gene operably linked to regulatory sequences that control expression of the heterologous gene in a host cell, and a second nucleic acid vector having AAV rep and cap nucleic acids sequences, wherein said cap nucleic acid sequence encodes an AAV capsid that is at least 95% identical to any of SEQ ID NOS:15-157; and recovering the AAV from the supernatant of the viral production cell culture. In another embodiment the viral production cell is an insect cell. In a preferred embodiment the insect cell is an Sf9 cell. In a further embodiment the first nucleic acid vector is introduced into the viral production cell by infection of the viral production cell by a baculovirus containing the first nucleic acid vector. In yet another embodiment the first and second nucleic acid vectors are introduced into the viral production cell by infection of the viral production cell by a first baculovirus containing the first nucleic acid vector and a second baculovirus containing the second nucleic acid vector. In further embodiments the invention AAV produced by the production methods provided herein.

[0032] In another embodiment, the invention provides for methods of treating a patient suffering from a disorder or disease comprising administering to the patient an effective amount of any of the AAV vectors or virus of the invention.

[0033] In a further embodiment, the invention provides for use of any of the AAV vectors or virus of the invention for preparation of a medicament for the treatment of a disorder or disease. The invention also provides for compositions comprising any of the AAV vectors or virus of the invention for the treatment of a disease or disorder.

[0034] In yet another embodiment, the disease or disorder in a subject is associated with abnormal activity of an endogenous protein. As used herein “endogenous protein” means a protein or gene product encoded by the genome of the subject suffering from the disease or disorder.

[0035] An “AAV virion” or “AAV viral particle” or “AAV vector particle” or “AAV virus” refers to a viral particle composed of at least one AAV capsid protein and an encapsidated polynucleotide AAV vector. If the particle comprises a heterologous polynucleotide (i.e. a polynucleotide other than a wild-type AAV genome such as a transgene to be delivered to a mammalian cell), it is typically referred to as an “AAV vector particle” or simply an “AAV vector”. Thus, production of AAV vector particle necessarily includes production of AAV vector, as such a vector is contained within an AAV vector particle.

[0036] The invention also provides for cells comprising any of the AAV vectors of the invention, and viral particles produced by these cells of the invention.

[0037] The term “inverted terminal repeat (ITR)” as used herein refers to the art-recognized regions found at the 5’ and 3’ termini of the AAV genome which function in cis as origins of DNA replication and as packaging signals for the viral genome. AAV ITRs, together with the AAV rep coding region, provide for efficient excision and rescue from a plasmid vector, and integration of a nucleotide sequence interposed between two flanking ITRs into a host cell genome. Sequences of certain AAV-associated ITRs are disclosed by Yan et al., *J. Virol.* 79(1):364-379 (2005) which is herein incorporated by reference in its entirety.

[0038] The phrase “helper functions for generating a productive AAV infection” as used herein refers to AAV-derived coding sequences that can be expressed to provide AAV gene products that, in turn, function in trans for productive AAV replication. Thus, AAV helper functions include the rep and cap regions. The rep expression products have been shown to possess many functions, including, among others: recognition, binding and nicking of the AAV origin of DNA replication; DNA helicase activity; and modulation of transcription from AAV (or other heterologous) promoters. The cap expression products supply necessary packaging functions. AAV helper functions are used herein to complement AAV functions in trans that are missing from AAV vectors. Helper functions for generating a productive AAV infection also

may include certain helper functions from baculovirus, herpes virus, adenovirus, or vaccinia virus.

[0039] In some embodiments, the viral construct comprises a nucleotide sequence encoding AAV rep and cap genes.

[0040] The term “AAV rep gene” as used herein refers to the art-recognized region of the AAV genome which encodes the replication proteins of the virus which are required to replicate the viral genome and to insert the viral genome into a host genome during latent infection. For a further description of the AAV rep coding region, see, e.g., Muzyczka et al., *Current Topics in Microbiol. and Immunol.* 158:97-129 (1992); Kotin et al., *Human Gene Therapy* 5:793-801 (1994), the disclosures of which are incorporated herein by reference in their entireties. The rep coding region, as used herein, can be derived from any viral serotype, such as the AAV serotypes described above. The region need not include all of the wild-type genes but may be altered, e.g., by the insertion, deletion or substitution of nucleotides, so long as the rep genes retain the desired functional characteristics when expressed in a suitable recipient cell.

[0041] The term “AAV cap gene” as used herein refers to the art-recognized region of the AAV genome which encodes the coat proteins of the virus which are required for packaging the viral genome. For a further description of the cap coding region, see, e.g., Muzyczka et al., *Current Topics in Microbiol. and Immunol.* 158:97-129 (1992); Kotin et al., *Human Gene Therapy* 5:793-801 (1994), the disclosures of which are incorporated herein by reference in their entireties. The AAV cap coding region, as used herein, can be derived from any AAV serotype, as described above. The region need not include all of the wild-type cap genes but may be altered, e.g., by the insertion, deletion or substitution of nucleotides, so long as the genes provide for sufficient packaging functions when present in a host cell along with an AAV vector.

[0042] The term “transfection” is used to refer to the uptake of foreign DNA by a cell. A cell has been “transfected” when exogenous DNA has been introduced inside the cell membrane. A number of transfection techniques are generally known in the art. See, e.g., Graham et al., *Virology* 52:456 (1973); Sambrook et al., Molecular Cloning: A Laboratory Manual, Cold Spring Harbor Laboratories, New York (1989); Davis et al., Basic Methods in Molecular Biology, Elsevier (1986); Chu et al., *Gene* 13:197 (1981), the disclosures of which are incorporated herein by reference in their entireties. Such techniques can be used to introduce one or more exogenous

DNA moieties, such as a nucleotide integration vector and other nucleic acid molecules, into suitable host cells. The term captures chemical, electrical, and viral-mediated transfection procedures.

[0043] The viral construct is, in some embodiments, in the form of a baculoviral vector capable of productive transformation, transfection or infection in any cell type. In some embodiments, the viral construct comprises at least one nucleotide sequence encoding a heterologous protein.

[0044] In yet another aspect, described herein is an AAV particle produced by a method described herein. In some embodiments, the AAV particle comprises in its genome at least one nucleotide encoding a heterologous protein.

[0045] The term “heterologous proteins or peptides” refer to any protein that is not expressed by wild type AAV including tags such as hexahistidine, FLAG, myc, polyhistidine, or labels or immunogens, adjuvants, selection markers, therapeutic proteins or targeting proteins or peptides, to name a few.

[0046] Exemplary heterologous protein described herein includes, but is not limited to, β -globin, hemoglobin, tissue plasminogen activator, and coagulation factors; colony stimulating factors (CSF); interleukins, such as IL-1, IL-2, IL-3, IL-4, IL-5, IL-6, IL-7, IL-8, IL-9, etc.; growth factors, such as keratinocyte growth factor (KGF), stem cell factor (SCF), fibroblast growth factor (FGF, such as basic FGF and acidic FGF), hepatocyte growth factor (HGF), insulin-like growth factors (IGFs), bone morphogenetic protein (BMP), epidermal growth factor (EGF), growth differentiation factor-9 (GDF-9), hepatoma derived growth factor (HDGF), myostatin (GDF-8), nerve growth factor (NGF), neurotrophins, platelet-derived growth factor (PDGF), thrombopoietin (TPO), transforming growth factor alpha (TGF- α), transforming growth factor beta (TGF- β), and the like; soluble receptors, such as soluble TNF- α receptors, soluble interleukin receptors (e.g., soluble IL-1 receptors and soluble type II IL-1 receptors), soluble γ/Δ T cell receptors, ligand-binding fragments of a soluble receptor, and the like; enzymes, such as α -glucosidase, imiglucrase, β -glucocerebrosidase, and alglucerase; enzyme activators, such as tissue plasminogen activator; chemokines, such as 1P-10, monokine induced by interferon-gamma (Mig), Gro α /IL-8, RANTES, MIP-1 α , MIP-1 β , MCP-1, PF-4, and the like; angiogenic agents, such as vascular endothelial growth factors (VEGFs, e.g., VEGF121, VEGF165, VEGF-

C, VEGF-2), glioma-derived growth factor, angiogenin, angiogenin-2; and the like; anti-angiogenic agents, such as a soluble VEGF receptor; protein vaccine; neuroactive peptides, such as nerve growth factor (NGF), bradykinin, cholecystokinin, gastrin, secretin, oxytocin, gonadotropin-releasing hormone, beta-endorphin, enkephalin, substance P, somatostatin, prolactin, galanin, growth hormone-releasing hormone, bombesin, dynorphin, warfarin, neurotensin, motilin, thyrotropin, neuropeptide Y, luteinizing hormone, calcitonin, insulin, glucagons, vasopressin, angiotensin II, thyrotropin-releasing hormone, vasoactive intestinal peptide, a sleep peptide, and the like; thrombolytic agents; atrial natriuretic peptide; relaxin; glial fibrillary acidic protein; follicle stimulating hormone (FSH); human alpha-1 antitrypsin; leukemia inhibitory factor (LIF); tissue factors, luteinizing hormone; macrophage activating factors; tumor necrosis factor (TNF); neutrophil chemotactic factor (NCF); tissue inhibitors of metalloproteinases; vasoactive intestinal peptide; angiogenin; angiotropin; fibrin; hirudin; IL-1 receptor antagonists; ciliary neurotrophic factor (CNTF); brain-derived neurotrophic factor (BDNF); neurotrophins 3 and 4/5 (NT-3 and 4/5); glial cell derived neurotrophic factor (GDNF); aromatic amino acid decarboxylase (AADC); Factor VIII, Factor IX, Factor X; dystrophin or mini-dystrophin; lysosomal acid lipase; phenylalanine hydroxylase (PAH); glycogen storage disease-related enzymes, such as glucose-6-phosphatase, acid maltase, glycogen debranching enzyme, muscle glycogen phosphorylase, liver glycogen phosphorylase, muscle phosphofructokinase, phosphorylase kinase, glucose transporter, aldolase A, β -enolase, glycogen synthase; and lysosomal enzymes.

DETAILED DESCRIPTION

[0047] The invention provides for novel AAV capsid proteins, nucleic acid encoding those capsid proteins and AAV virus comprising those novel capsid proteins. In some embodiments, the AAV capsid proteins were isolated and identified from various mammalian tissues. The amino acid sequences of the novel AAV capsid VP1 proteins are set out as SEQ ID NOS:15-89, and the locations of the associated VP2 and VP3 regions are disclosed herein. In addition, the invention provides for novel engineered chimeric AAV capsid proteins which have a backbone amino acid sequence derived from one AAV capsid sequence and fragments of capsid proteins from at least one other, different AAV capsid sequence, such as at least one VR, GBS and/or GH loop. The amino acid sequences of exemplary engineered chimeric AAV capsid VP1 proteins are set out as SEQ ID NOS:90-157.

AAV Vectors

[0048] As used herein, the term "AAV" is a standard abbreviation for adeno-associated virus. Adeno-associated virus is a single-stranded DNA parvovirus that grows only in cells in which certain functions are provided by a co-infecting helper virus. There are currently at least thirteen serotypes of AAV that have been characterized, as shown below in Table 1. General information and reviews of AAV can be found in, for example, Carter, Handbook of Parvoviruses, Vol. 1, pp. 169-228 (1989), and Berns, Virology, pp. 1743-1764, Raven Press, (New York, 1990). However, it is fully expected that these same principles will be applicable to additional AAV serotypes since it is well known that the various serotypes are quite closely related, both structurally and functionally, even at the genetic level. (See, for example, Blacklowe, pp. 165-174 of Parvoviruses and Human Disease, J. R. Pattison, ed. (1988); and Rose, *Comprehensive Virology* 3:1-61 (1974)). For example, all AAV serotypes apparently exhibit very similar replication properties mediated by homologous rep genes; and all bear three related capsid proteins such as those expressed in AAV6. The degree of relatedness is further suggested by heteroduplex analysis which reveals extensive cross-hybridization between serotypes along the length of the genome; and the presence of analogous self-annealing segments at the termini that correspond to "inverted terminal repeat sequences" (ITRs). The similar infectivity patterns also suggest that the replication functions in each serotype are under similar regulatory control.

[0049] An "AAV vector" as used herein refers to a vector comprising one or more polynucleotides of interest (or transgenes) that are flanked by AAV terminal repeat sequences (ITRs). Such AAV vectors can be replicated and packaged into infectious viral particles when present in a host cell that has been transfected with a vector encoding and expressing rep and cap gene products.

[0050] An "AAV virion" or "AAV viral particle" or "AAV vector particle" or "AAV virus" refers to a viral particle composed of at least one AAV capsid protein and an encapsidated polynucleotide AAV vector. If the particle comprises a heterologous polynucleotide (i.e. a polynucleotide other than a wild-type AAV genome such as a transgene to be delivered to a mammalian cell), it is typically referred to as an "AAV vector particle" or simply an "AAV vector". Thus, production of AAV vector particle necessarily includes production of AAV vector, as such a vector is contained within an AAV vector particle.

[0051] AAV "rep" and "cap" genes are genes encoding replication and encapsidation proteins, respectively. AAV rep and cap genes have been found in all AAV serotypes examined to date, and are described herein and in the references cited. In wild-type AAV, the rep and cap genes are generally found adjacent to each other in the viral genome (i.e., they are "coupled" together as adjoining or overlapping transcriptional units), and they are generally conserved among AAV serotypes. AAV rep and cap genes are also individually and collectively referred to as "AAV packaging genes." The AAV cap gene in accordance with the present invention encodes a Cap protein which is capable of packaging AAV vectors in the presence of rep and adeno helper function and is capable of binding target cellular receptors. In some embodiments, the AAV cap gene encodes a capsid protein having an amino acid sequence derived from a particular AAV serotype, for example the serotypes shown in Table 1.

Table 1. AAV serotypes

AAV Serotype	Genbank Accession No.
AAV-1	NC_002077.1
AAV-2	NC_001401.2
AAV-3	NC_001729.1
AAV-3B	AF028705.1
AAV-4	NC_001829.1
AAV-5	NC_006152.1
AAV-6	AF028704.1
AAV-7	NC_006260.1
AAV-8	NC_006261.1
AAV-9	AX753250.1
AAV-10	AY631965.1
AAV-11	AY631966.1
AAV-12	DQ813647.1
AAV-13	EU285562.1

[0052] The AAV sequences employed for the production of AAV can be derived from the genome of any AAV serotype. Generally, the AAV serotypes have genomic sequences of significant homology at the amino acid and the nucleic acid levels, provide a similar set of genetic functions, produce virions which are essentially physically and functionally equivalent, and replicate and assemble by practically identical mechanisms. For the genomic sequence of AAV serotypes and a discussion of the genomic similarities see, for example, GenBank Accession number U89790; GenBank Accession number J01901; GenBank Accession number

AF043303; GenBank Accession number AF085716; Chlorini et al., *J. Vir.* 71:6823-33(1997); Srivastava et al., *J. Vir.* 45:555-64 (1983); Chlorini et al., *J. Vir.* 73:1309-1319 (1999); Rutledge et al., *J. Vir.* 72:309-319 (1998); and Wu et al., *J. Vir.* 74: 8635-47 (2000).

[0053] The genomic organization of all known AAV serotypes is very similar. The genome of AAV is a linear, single-stranded DNA molecule that is less than about 5,000 nucleotides (nt) in length. Inverted terminal repeats (ITRs) flank the unique coding nucleotide sequences for the non-structural replication (Rep) proteins and the structural (VP) proteins. The VP proteins form the capsid. The terminal 145 nt are self-complementary and are organized so that an energetically stable intramolecular duplex forming a T-shaped hairpin may be formed. These hairpin structures function as an origin for viral DNA replication, serving as primers for the cellular DNA polymerase complex. The Rep genes encode the Rep proteins, Rep78, Rep68, Rep52, and Rep40. Rep78 and Rep68 are transcribed from the p5 promoter, and Rep 52 and Rep40 are transcribed from the p19 promoter. The cap genes encode the VP proteins, VP1, VP2, and VP3. The cap genes are transcribed from the p40 promoter.

[0054] In some embodiments, a nucleic acid sequence encoding an AAV capsid protein is operably linked to regulatory expression control sequences for expression in a specific cell type, such as Sf9 or HEK cells. Techniques known to one skilled in the art for expressing foreign genes in insect host cells or mammalian host cells can be used to practice the invention. Methodology for molecular engineering and expression of polypeptides in insect cells is described, for example, in Summers and Smith, A Manual of Methods for Baculovirus Vectors and Insect Culture Procedures, Texas Agricultural Experimental Station Bull. No. 7555, College Station, Tex. (1986); Luckow. 1991. In Prokop et al., Cloning and Expression of Heterologous Genes in Insect Cells with Baculovirus Vectors' Recombinant DNA Technology and Applications, 97-152 (1986); King, L. A. and R. D. Possee, The baculovirus expression system, Chapman and Hall, United Kingdom (1992); O'Reilly, D. R., L. K. Miller, V. A. Luckow, Baculovirus Expression Vectors: A Laboratory Manual, New York (1992); W.H. Freeman and Richardson, C. D., Baculovirus Expression Protocols, Methods in Molecular Biology, volume 39 (1995); U.S. Pat. No. 4,745,051; US2003148506; and WO 03/074714. A particularly suitable promoter for transcription of a nucleotide sequence encoding an AAV capsid protein is e.g. the polyhedron promoter. However, other promoters that are active in insect cells are known in the

art, e.g. the p10, p35 or IE-1 promoters and further promoters described in the above references are also contemplated.

[0055] Use of insect cells for expression of heterologous proteins is well documented, as are methods of introducing nucleic acids, such as vectors, e.g., insect-cell compatible vectors, into such cells and methods of maintaining such cells in culture. See, for example, METHODS IN MOLECULAR BIOLOGY, ed. Richard, Humana Press, NJ (1995); O'Reilly et al., BACULOVIRUS EXPRESSION VECTORS, A LABORATORY MANUAL, Oxford Univ. Press (1994); Samulski et al., *J. Vir.* 63:3822-8 (1989); Kajigaya et al., *Proc. Nat'l. Acad. Sci. USA* 88:4646-50 (1991); Ruffing et al., *J. Vir.* 66:6922-30 (1992); Kirnbauer et al., *Vir.* 219:37-44 (1996); Zhao et al., *Vir.* 272:382-93 (2000); and Samulski et al., U.S. Pat. No. 6,204,059. In some embodiments, the nucleic acid construct encoding AAV in insect cells is an insect cell-compatible vector. An "insect cell-compatible vector" or "vector" as used herein refers to a nucleic acid molecule capable of productive transformation or transfection of an insect or insect cell. Exemplary biological vectors include plasmids, linear nucleic acid molecules, and recombinant viruses. Any vector can be employed as long as it is insect cell-compatible. The vector may integrate into the insect cell's genome but the presence of the vector in the insect cell need not be permanent and transient episomal vectors are also included. The vectors can be introduced by any means known, for example by chemical treatment of the cells, electroporation, or infection. In some embodiments, the vector is a baculovirus, a viral vector, or a plasmid. In a more preferred embodiment, the vector is a baculovirus, i.e. the construct is a baculoviral vector. Baculoviral vectors and methods for their use are described in the above cited references on molecular engineering of insect cells.

[0056] Baculoviruses are enveloped DNA viruses of arthropods, two members of which are well known expression vectors for producing recombinant proteins in cell cultures. Baculoviruses have circular double-stranded genomes (80-200 kbp) which can be engineered to allow the delivery of large genomic content to specific cells. The viruses used as a vector are generally *Autographa californica* multicapsid nucleopolyhedrovirus (AcMNPV) or *Bombyx mori* (Bm)NPV (Kato et al., *Appl. Microbiol. Biotechnol.* 85(3):459-470 (2010)). Baculoviruses are commonly used for the infection of insect cells for the expression of recombinant proteins. In particular, expression of heterologous genes in insects can be accomplished as described in for instance U.S. Pat. No. 4,745,051; Friesen et al., Curr. Top. Microbiol. Immunol. 131:31-49.

(1986); EP 127,839; EP 155,476; Miller et al., *Ann. Rev. of Microbiol.* 42: 177-199 (1988); Carbonell et al., *Gene* 73(2):409-18 (1988); Maeda et al., *Nature* 315(6020):592-4 (1985); Lebacqz-Verheyden et al., *Mol. Cell. Biol.* 8(8):3129-35 (1988); Smith et al., *Proc. Natl. Acad. Sci. U.S.A.* 82(24):8404-8 (1985); Miyajima et al., *Gene* 58(2-3):273-81 (1987); and Martin et al., *DNA* 7(2):99-106 (1988). Numerous baculovirus strains and variants and corresponding permissive insect host cells that can be used for protein production are described in Luckow et al., *Nature Biotechnology* 6:47-55 (1988), and Maeda et al., *Nature* 315(6020):592-4 (1985).

Novel AAV Capsid Protein

[0057] In a first aspect, the invention provides for novel AAV capsid proteins that were isolated from various mammalian tissues. The novel AAV VP1 capsid proteins are provided as SEQ ID NOS:15-89 and the locations of the associated VP2 and VP3 regions are described herein. The invention also provides for polynucleotides comprising a nucleotide sequence encoding these novel AAV capsid proteins. The invention provides the amino acid sequences of the novel AAV capsid proteins including the engineered chimeric capsid proteins described herein (referred herein collectively as the “AAV capsid proteins of the invention”), and the nucleic acid sequences encoding the AAV capsid proteins of the invention. Also provided are fragments of these AAV capsid nucleic acid and amino acid sequences of the invention. Each of these sequences may be readily utilized in a variety of vector systems and host cells. Desirable fragments of the capsid VP1 proteins include VP2, VP3 and variable regions, the GBS domain and the GH loop, and polynucleotide sequences encoding these proteins. These fragments may be readily utilized in a variety of vector systems and host cells. Such fragments may be used alone, in combination with other AAV sequences or fragments, or in combination with elements from other AAV or non-AAV viral sequences. In one particularly desirable embodiment, a vector contains the AAV capsid sequences of the invention.

[0058] The AAV capsid sequences of the invention and fragments thereof are useful in production of rAAV, and are also useful as antisense delivery vectors, gene therapy vectors, or vaccine vectors. The invention further provides nucleic acid molecules, gene delivery vectors, and host cells which contain the novel AAV capsid sequences of the invention.

[0059] Suitable fragments can be determined using the information provided herein. Alignments are performed using any of a variety of publicly or commercially available Multiple

Sequence Alignment Programs, such as "Clustal W", accessible through Web Servers on the internet. Alternatively, Vector NTI utilities are also used. There are also a number of algorithms known in the art which can be used to measure nucleotide sequence identity, including those contained in the programs described above. As another example, polynucleotide sequences can be compared using FASTA, a program in GCG Version 6.1. FASTA provides alignments and percent sequence identity of the regions of the best overlap between the query and search sequences. For instance, percent sequence identity between nucleic acid sequences can be determined using FASTA with its default parameters (a word size of 6 and the NOPAM factor for the scoring matrix) as provided in GCG Version 6.1, herein incorporated by reference. Similar programs are available for amino acid sequences, e.g., the "Clustal X" program. Additional sequence alignment tools that can be used are provided by (protein sequence alignment; (http://www.ebi.ac.uk/Tools/psa/emboss_needle/)) and (nucleic acid alignment; (http://www.ebi.ac.uk/Tools/psa/emboss_needle/nucleotide.html)). Generally, any of these programs are used at default settings, although one of skill in the art can alter these settings as needed. Alternatively, one of skill in the art can utilize another algorithm or computer program which provides at least the level of identity or alignment as that provided by the referenced algorithms and programs.

[0060] The terms "substantial identity", "substantial homology" or "substantial similarity," when referring to a nucleic acid, or fragment thereof, indicates that, when optimally aligned with appropriate nucleotide insertions or deletions with another nucleic acid (or its complementary strand), there is nucleotide sequence identity in at least about 95 to 99% of the aligned sequences such as 95% identity, 96% identity, 97% identity, 98% identity and 99% identity. Preferably, the homology is over the full-length of the two sequences being compared, or an open reading frame thereof, or another suitable fragment which is at least 15 nucleotides in length. Examples of suitable fragments are described herein. Also included in the nucleic acid sequences of the invention are natural variants and engineered modifications of the nucleic acids encoding the AAV capsids of the invention and its complementary strand. Such modifications include, for example, labels which are known in the art, methylation, and substitution of one or more of the naturally occurring nucleotides with a degenerate nucleotide.

[0061] The terms "substantial identity", "substantial homology" or "substantial similarity," when referring to amino acids or fragments thereof, indicates that, when optimally aligned with

appropriate amino acid insertions or deletions with another amino acid (or its complementary strand), there is amino acid sequence identity in at least about 95 to 99% of the aligned sequences such as 95% identity, 96% identity, 97% identity, 98% identity and 99% identity. Preferably, the homology is over the full-length of the two sequences being compared, or a protein thereof, e.g., a cap protein, a rep protein, or a fragment thereof which is at least 8 amino acids, or more desirably, at least 15 amino acids in length. Examples of suitable fragments are described herein.

[0062] By the term "highly conserved" is meant at least 80% identity, preferably at least 90% identity, and more preferably, over 97% identity. Identity is readily determined by one of skill in the art by resort to algorithms and computer programs known by those of skill in the art.

[0063] The term "percent sequence identity" or "identical" in the context of nucleic acid sequences or amino acid sequences refers to the residues in the two sequences which are the same when aligned for maximum correspondence. The length of sequence identity comparison may be over the full-length of the two sequences being compared, the full-length of a gene coding sequence, or a fragment of at least about 500 to 5000 nucleotides, is desired. However, identity among smaller fragments, e.g. of at least about nine nucleotides, usually at least about 20 to 24 nucleotides, at least about 28 to 32 nucleotides, at least about 36 or more nucleotides, may also be desired. Similarly, "percent sequence identity" may be readily determined for amino acid sequences, over the full-length of a protein, or a fragment thereof. Suitably, a fragment is at least about 8 amino acids in length, and may be up to about 700 amino acids. Examples of suitable fragments are described herein.

[0064] As described herein, the vectors of the invention containing or comprising the AAV capsid proteins of the invention are particularly well-suited for use in applications in which the neutralizing antibodies diminish the effectiveness of other AAV serotype based vectors, as well as other viral vectors. The rAAV vectors of the invention are particularly advantageous in rAAV re-administration and repeat gene therapy.

[0065] Also included within the invention are fragments of the nucleic acids encoding the AAV capsid proteins of the invention, their complementary strand, cDNA and RNA complementary thereto. Suitable fragments are at least 15 nucleotides in length, and encompass functional fragments, i.e., fragments which are of biological interest. Such fragments include the

sequences encoding the three variable proteins (VP) of the capsid which are alternative splice variants: VP1, VP2 and VP3. Other suitable fragments of the nucleic acids encoding the AAV capsids of the invention include the fragment which contains the start codon for the capsid protein, and the fragments encoding the variable regions of the VP1 capsid protein, which are described herein.

[0066] The invention is not limited to the AAV capsid amino acid sequences, peptides and proteins expressed from the AAV nucleic acid sequences of the invention and encompasses amino acid sequences, peptides and proteins generated by other methods known in the art, including, e.g., by chemical synthesis, by other synthetic techniques, or by other methods. For example, the sequences of any of the capsids described herein can be readily generated using a variety of techniques.

[0067] Suitable production techniques are well known to those of skill in the art. See, e.g., Sambrook et al, Molecular Cloning: A Laboratory Manual, Cold Spring Harbor Press (Cold Spring Harbor, N.Y.). Alternatively, peptides can also be synthesized by the well-known solid phase peptide synthesis methods (Merrifield, *J. Am. Chem. Soc.*, 85:2149 (1962); Stewart and Young, Solid Phase Peptide Synthesis Freeman, (San Francisco, 1969) pp. 27-62. These and other suitable production methods are within the knowledge of those of skill in the art and are not a limitation of the present invention.

[0068] The AAV capsid is composed of three proteins, VP1, VP2 and VP3, which are alternative splice variants. The full-length capsid sequence is referred to as VP1 which encompasses the spliced variants referred to as VP2 and VP3. The invention also provides for other functional fragments of the AAV capsid proteins of the invention. Other desirable fragments of the capsid protein include the variable regions (VR), the constant regions which are located between the variable regions, the GBS domain, and the GH loop. Other desirable fragments of the capsid protein include the HPV themselves.

[0069] An algorithm has been developed to determine areas of sequence divergence in AAV2. (Chiorini et al, *J. Virol.*, 73:1309-19 (1999); Rutledge et al, *J. Virol.*, 72:309-319 (1998)). Using this algorithm and/or the alignment techniques described herein, the VR of the novel AAV capsid sequences are determined. Using the alignment provided herein performed using the Clustal X program at default settings, or using other commercially or publicly available

alignment programs at default settings, one of skill in the art can readily determine corresponding fragments of the novel AAV capsids of the invention.

[0070] Suitably, fragments of an AAV capsid protein are at least 8 amino acids in length, or at least 9 amino acids in length, or at least 10 amino acids in length, or at least 20 amino acids in length, or 30 amino acids in length or at least 50 amino acids in length, or at least 75 amino acids in length, or at least 100 amino acids in length or 200 amino acids in length or 250 amino acids in length or 300 amino acids in length or 350 amino acids in length or 400 amino acids in length. However, fragments of other desired lengths may be readily utilized. All fragments of the invention retain biological activity of a capsid AAV protein. Such fragments may be produced recombinantly or by other suitable means, e.g., chemical synthesis.

[0071] The sequences, proteins, and fragments of the invention may be produced by any suitable means, including recombinant production, chemical synthesis, or other synthetic means. Such production methods are within the knowledge of those of skill in the art and are not a limitation of the present invention.

[0072] In addition to including the nucleic acid sequences provided in the Sequence Listing, the present invention includes nucleic acid molecules and sequences which are designed to express the amino acid sequences, proteins and peptides of the AAV capsid proteins of the invention. Thus, the invention includes nucleic acid sequences which encode the following AAV capsid amino acid sequences and artificial AAV capsid proteins generated using these sequences and/or unique fragments thereof.

[0073] Artificial capsid or engineered capsid proteins may be generated by any suitable technique, using a AAV capsid protein sequence of the invention (e.g., a fragment of a VP1 capsid protein) in combination with heterologous sequences which may be obtained from another AAV serotype (known or novel), non-contiguous portions of the same AAV serotype, from a non-AAV viral source, or from a non-viral source. An artificial AAV serotype may be, without limitation, a chimeric AAV capsid, a recombinant AAV capsid, or a "humanized" AAV capsid.

Production of AAV with the Capsid Proteins of the Invention

[0074] The invention encompasses AAV capsid protein sequences and the nucleic acids encoding these proteins of which are free of DNA and/or cellular material which these viruses

are associated in nature. In another aspect, the present invention provides molecules which utilize the novel AAV sequences of the invention, including fragments thereof, for production of molecules useful in delivery of a heterologous gene or other nucleic acid sequences to a target cell.

[0075] In another aspect, the present invention provides molecules which utilize the AAV capsid protein sequences of the invention, including fragments thereof, for production of viral vectors useful in delivery of a heterologous gene or other nucleic acid sequences to a target cell.

[0076] The molecules of the invention which contain AAV capsid nucleic acid sequences include any genetic element (vector) which may be delivered to a host cell, e.g., naked DNA, a plasmid, phage, transposon, cosmid, episome, a protein in a non-viral delivery vehicle (e.g., a lipid-based carrier), virus, etc. which transfer the sequences carried thereon. The selected vector may be delivered by any suitable method, including transfection, electroporation, liposome delivery, membrane fusion techniques, high velocity DNA-coated pellets, viral infection and protoplast fusion. The methods used to construct any embodiment of this invention are known to those with skill in nucleic acid manipulation and include genetic engineering, recombinant engineering, and synthetic techniques. See, e.g., Sambrook et al, Molecular Cloning: A Laboratory Manual, Cold Spring Harbor Press, Cold Spring Harbor, N.Y.

[0077] In one embodiment, the vectors of the invention contain, at a minimum, sequences encoding the AAV capsid of the invention or a fragment thereof. In another embodiment, the vectors of the invention contain, at a minimum, sequences encoding an AAV rep protein or a fragment thereof. Optionally, such vectors may contain both AAV cap and rep proteins. In vectors in which both AAV rep and cap are provided, the AAV rep and AAV cap sequences can both be of the same AAV serotype origin. Alternatively, the present invention provides vectors in which the rep sequences are from an AAV serotype which differs from that which is providing the cap sequences. In one embodiment, the rep and cap sequences are expressed from separate sources (e.g., separate vectors, or a host cell and a vector). In another embodiment, these rep sequences are fused in frame to cap sequences of a different AAV serotype to form a chimeric AAV vector. Optionally, the vectors of the invention further contain a minigene comprising a selected transgene which is flanked by AAV 5' ITR and AAV 3' ITR.

[0078] Thus, in one embodiment, the vectors described herein contain nucleic acid sequences encoding an intact AAV capsid protein of any one of amino acid sequences SEQ ID NO: 1-73. Alternatively, these vectors contain sequences encoding artificial capsids which contain one or more fragments of the capsid fused to heterologous AAV or non-AAV capsid proteins (or fragments thereof). These artificial capsid proteins are selected from non-contiguous portions of the any of the AAV capsid proteins of the invention or from capsids of other AAV serotypes. In another example, it may be desirable to alter the start codon of the VP3 protein to GTG. Alternatively, the rAAV may contain one or more of the variable regions of one or more of the AAV capsid proteins of the invention, or other fragments. These modifications may be to increase expression, yield, and/or to improve purification in the selected expression systems, or for another desired purpose (e.g., to change tropism or alter neutralizing antibody epitopes).

[0079] The vectors described herein, e.g., a plasmid, are useful for a variety of purposes, but are particularly well suited for use in production of a rAAV containing a capsid comprising AAV sequences or a fragment thereof. These vectors, including rAAV, their elements, construction, and uses are described in detail herein.

VP1 Amino Acid Sequences of Known AAV Capsid Proteins

[0080] The invention also provides for engineered chimeric AAV capsid proteins (and AAV comprising those capsid proteins) in which one or more variable region(s), the GBS region and/or the GH loop in a backbone (or recipient) capsid protein sequence are substituted with one or more variable region(s), GBS region and/or GH loop from a different AAV capsid sequence donor. The recipient and donor sequences may derive from any previously known AAV serotype or capsid sequence, or any novel AAV capsid sequence described herein. The novel, engineered AAV capsid proteins of the invention are generated by swapping at least one variable region, GBS region or GH loop region from one capsid sequence for the respective region(s) in a recipient capsid sequence. In this regard, it is noted that one, two, three, four, five, six, seven, eight or all nine VRs in a recipient VP1 capsid sequence can be replaced by the respective region(s) from one or more different VP1 capsid sequence. Any and all of the various combinations of engineered, chimeric AAV capsid sequences that can be produced by the VR region swapping method described herein (and all associated AAV virus comprising those chimeric capsid sequences) are contemplated by this invention.

[0081] The VP1 sequence of AAVbo is set out as SEQ ID NO:1 and the locations of the associated variable regions and GBS and GH loop regions are defined in Table 2 below.

MSFVDHPPDWLESIGDGFREFLGLGAGPPKPKANQQKQDNARGLVLPGYKYLGPNGLDKGDVPNFADDEVAREHDLQSYQKQLEAGD
NPYLKYNHADADEFQEKLASDTSFGGNLGAQVFQAKKRILEPLGLVETPDKTAPAAKKRPLEQSPQEPDSSSGVGKKGKQPARKRLN
FDDEPGAGDGPPPEGPSSGAMSTETEMRAAAGGNGGDAGQGAEGVGNASGDWHCDSTWSESHVTTTSTRTWVLPITYNNHLYRLGSL
SNASDTFNGFSTPWGYFDFNRHFCHFSRDPWQRLINNNHWGLRPKSMQVRIFNIQVKEVTTSNGETTVSNNTLSTVQIFADSTYELP
YVMDAQEGSLPPFPNDVFMVPQYGYCGLVTGGSSQNQTDNRNAYCYCLEYFSPQMLRTGNNFEMVYKFENVPFHSYAHQSGLDRML
NPLLDQYLWELQSTTSGGTNLQGN SATNFAKLTKTNFSGYRKNWLP GPMKQQRFSKTASQNYKIPQGRNNSLLHYETRTTLDGRW
SNFAPGTAMATAANDATDFSAQLIFAGPNI TGN TTTDANNLMFTSEDEL RATNPRD TDLFGHLATNQQNATTVP TDDVDG VGVY
PGMVWQDRDIYYQGP I WAKIPHTDGHFHPSP LIGGFGLKSPPPQ I FIKNT PVPANPATTFS PARINSF ITQYSTGQVAVKIEWELQ
KERSKRWNP EVQFTSNYGAQDSLLWAPDNAGAYKEPRAIGSRYLTNHL

[0082] The VP1 sequence of AAVmo is set out as SEQ ID NO:2 and the locations of the associated variable regions and GBS and GH loop regions are defined in Table 2 below.

MSFFDWLGKQYAQGADEFWDLKSGPPAPKKARKDGSAGFNFGHKYLGPGNSLDRGDPVDADDAQAQKHQDSYQEQLEAGDNPYLK
YNHADREFQEALKDDTSFEGNLARGLFEAKKLVAEPLGLVEPELAPP SGRKRPVQSSQESGYSSSQDKRPNLDVDEEDREFAAAAA
ETETGSAPPTGNLPGTMAGGGSAPIDGSGYADGVGNASGDWHCDSTWLDNCVITRTTRTNWLPITYNNHIYKRLNGTTSQDQSYF
GFSTPWGYFDFNRHFCHFSRDPWQRLINNNWGLRPKSLRFKIFNIQVKEVTTQDSTKIISNNLTSTVQVFADTEYQLPYVIGSAHE
GCLPPFPADVFMPLQYGYCTRQDGNSSNPTPRSAFYCYCLEYFSPKMLRTGNSFEFTYNFEKVPFHSMAHNSQLDRMLNPLIDQYLY
YLDVTSSTGFTYQKGVHTNLPEQERNWLP GPGIRNQAWFNSATGNNPLTGTWQYSNKYVLENRASKIAPGPAMGIESTKFDNGNII
FSKEYITNVNTANPNQVNITRETEINSTNPLAGGSLGAHANNSQNTTTAP TLDHTNVMGVFP GSVWQDRDIY LQGGI WAKIPHTDG
HFHPSPLMGFGGLKNPPPI I LIKNT PVPADPPTEFNANKISSF ITQYSTGQVTVM EWELQKETSKRWNP EIQYSDDSSSTSGSIL
HFAPDDVGNYKEFRSIGTRYLTRPL

[0083] The VP1 sequence of AAV2 is set out as SEQ ID NO:3 and the locations of the associated variable regions and GBS and GH loop regions are defined in Table 2 below.

MAADGYLPDWLEDTLSEGIQWWKLKPGPPPKPAERHKDDSRGLVLPGYKYLGPFGNLDKGEVNEADAAALEHDKAYDRQLDSG
DNPYLKYNHADADEFQERLKEDTSFGGNLGRAVQAKKRILEPLGLVEEPVKTA PGKKRPVEHSPVEPDSSSGTGKAGQQPARKRLN
FGQTGDADSVDPDQPLGQPPAAPSGLGTNTMATGSGAPMADNNEGADGVGNSSGNWHDSTWMDRVIITSTRTWALPITYNNHLYK
QISSQSGASNDNHYFGYSTPWGYFDFNRHFCHFSRDPWQRLINNNWGF RPKRLNFKLFNIQVKEVTQNDGTTTTIANNLTSTVQVFT
DSEYQLPYVLGSAHQGCLPPFPADVFMVPQYGYLTNNGSQAVGRSSFYCYCLEYFSPQMLRTGNNFTFSYTFEDVPFHSSYAHQSLS
DRLMNPLIDQYLYLSRTNTPSGTTTQSR LQFSQAGASDIRDQSRNWLP GPCYRQQRVSKTSADNNNSEYSWTGATKYHLNGRDSL
VNP GPAMASHKDDEEKFFPQSGVLIFGKQGSEKTNVDIEKVMITDEEIRTNPVATEQYGSVSTNLQGRNQAAATADVNTQGVLP
GMVWQDRDVYLQGP I WAKIPHTDGHFHPSP LMGFGGLKHPPPI I LIKNT PVPANPSTTFSAAKFASFITQYSTGQVSVEIEWELQK
ENSKRWNP EIQYTSNYNKSVNVDFTVDTNGVYSEPRPIGTRYLTRNL

[0084] The VP1 sequence of AAV4 is set out as SEQ ID NO:4 and the locations of the associated variable regions and GBS and GH loop regions are defined in Table 2 below.

MTDGYLPDWLEDNLSEGVREWWALQPGAPKPKANQQHQDNARGLVLPGYKYLGPNGLDKGEVNAADAAALEHDKAYDQQLKAGD
NPYLKYNHADADEFQQRLLQGDTSFGGNLGRAVQAKKRILEPLGLVEQAGETAPGKKRPLIESPQQPDSSTGIGKKGKQPAKKKLVF
EDETGAGDGPPPEGSTSGAMSDSEMRAAAGGA AVEGGQGADGVGNASGDWHCDSTWSEGHVTTTSTRTWVLPITYNNHLYKRLGESL
QSNTYNGFSTPWGYFDFNRHFCHFSRDPWQRLINNNWGM RPKAMRVKIFNIQVKEVTTSNGETTVANNLTSTVQIFADSSYELPYV
MDAQEGESLPPFPNDVFMVPQYGYCGLVTGNTSQQTDRNAYCYCLEYFSPQMLRTGNNFEITYSFEKVPFHSYAHQSGLDRMLNP

LIDQYLWGLQSTTTGTTLNAGTATTNFTKLRPTNFSNFKKNWLPGPSIKQQGFSKTANQNYKIPATGSDSLIKYETHSTLDGRWSA
LTPGPPMATAGPADSKFSNSQLIFAGPKQNGNTATVPGTLIFTSEEELAAATNATDTDMWGNLPGGDQSNNSNLPVDRLTALGAVPG
MVWQNRDIYYQGPIWAKIPHTDGHFHPSPILIGGFGLKHPPPQIFIKNTPVPANPATTFSSSTPVNSFITQYSTGQVSVQIDWEIQKE
RSKRWNPEVQFTSNYQQNSLLWAPDAAGKYTEPRAIGTRYLTHHL

[0085] The VP1 sequence of AAV5 is set out as SEQ ID NO:5 and the locations of the associated variable regions and GBS and GH loop regions are defined in Table 2 below.

MSFVDHPPDWLEEVGEGLEFLGLEAGPPKPKPNQQHQDQARGLVLPGYNYLPGNGLDRGEPVNRADDEVAREHDI SYNEQLEAGD
NPYLKYNHADADEFQEKLADDTSFGGNLGKAVFQAKKRVLEPFLGVEEGAKTAPTGKRIDDHFPKRKKARTEEDSKPSTSSDAEAGP
SGSQQLQIPAPASSLGADTMSAGGGGPLGDNNGADGVGNASGDWHCDSTWMGDRVVTKSTRTWVLP SYNHHQYREIKSGSVDG
NANAYFGYSTPWGYFDFNRFHSHWSPRDWQRLINNYWGFRPRSLRVKIFNIQVKEVTVQDSTTTIANNLSTVQVFTDDDYQLPYV
VGNGTEGCLPAFPQVFTLPQYGYATLNRDNTENPTERSFFCLEYFPPSKMLRTGNNFEFTYNFEEVPFHSSSFAPSQNLFKLANPL
VDQYLYRFVSTNNTGGVQFNKNLAGRYANTYKNWFPGMGRITQGNLGSVGNRASVSFAATTNRMELEGASYQVPPQPNGMTNNLQ
GSNTYALENTMIFNSQPANPGTTATYLEGNMLITSESETQPVNRVAYNVGGQMATNNQSSTTAPATGTYNLQEI VPGSVWMERDVY
LQGP I WAKIPETGAHFHPSPAMGGFGLKHPPPMMLIKNTPVPGNITSFSDVPVSSFITQYSTGQVTVEMEWELKKENSKRWNPEIQ
YTNNYNDPQFVDFAPDSTGEYRTTRPIGTRYLTRPL

[0086] The VP1 sequence of AAV6 is set out as SEQ ID NO:6 and the locations of the associated variable regions and GBS and GH loop regions are defined in Table 2 below.

MAADGYLPDWLEDNLSEGIREWDLKPGAPKPKANQQKQDDGRGLVLPGYKYLGPFGNLDKGEPVNAADAAALEHDKAYDQQLKAG
DNPYLYRNHADADEFQERLQEDTSFGGNLGRAVFQAKKRVLEPFLGVEEGAKTAPGKKRPVEQSPQEPDSSSGIGKTGQQPAKKRLN
FGQTGDSESVDPDQPLGEPATPAAVGPTTASGGGAPMADNNEGADGVGNASGNWHCDSTWLGDRTVITSTRTWALPTYNNHLYK
QISSASTGASNDNHYFGYSTPWGYFDFNRFHCHFSRDPWQRLINNNWGFRPKRLNFKLFNIQVKEVTTNDGVTTIANNLTSTVQV
SDSEYQLPYVLGSAHQGCLPPFPADVFMIPQYGYLTNNNGSQAAGRSSFYCLEYFPPSMLRTGNNFTFSYTFEDVPFHSSYAHQS
LDRLMNPLIDQYLYLNRNTQNSGSAQNKDLLFSRGSAPAGMSVQPKNWLPGPCYRQQRVSKTKTDNNNSNFTWTGASKYNLNGRES
IINPGTAMASHKDDKDKFFPMGSGVMIFGKESAGASNTALDNVMTDEEEIKATNPVATERFGTVAVNLQSSSTDPATGDVHVMGAL
PGMVWQDRDVYLQGP I WAKIPHTDGHFHPSPMLGGFGLKHPPPQILIKNTPVPANPPAEFSATKFASFITQYSTGQVSVIEIEWELQ
KENSKRWNPEVQYTSNYAKSANVDFTVDNNGLYTEPRPIGTRYLTRPL

[0087] The VP1 sequence of AAV6.2 is set out as SEQ ID NO:7 and the locations of the associated variable regions and GBS and GH loop regions are defined in Table 2 below.

MAADGYLPDWLEDNLSEGIREWDLKPGAPKPKANQQKQDDGRGLVLPGYKYLGPFGNLDKGEPVNAADAAALEHDKAYDQQLKAG
DNPYLYRNHADADEFQERLQEDTSFGGNLGRAVFQAKKRVLEPFLGVEEGAKTAPGKKRPVEQSPQEPDSSSGIGKTGQQPAKKRLN
FGQTGDSESVDPDQPLGEPATPAAVGPTTASGGGAPMADNNEGADGVGNASGNWHCDSTWLGDRTVITSTRTWALPTYNNHLYK
QISSASTGASNDNHYFGYSTPWGYFDFNRFHCHFSRDPWQRLINNNWGFRPKRLNFKLFNIQVKEVTTNDGVTTIANNLTSTVQV
SDSEYQLPYVLGSAHQGCLPPFPADVFMIPQYGYLTNNNGSQAAGRSSFYCLEYFPPSMLRTGNNFTFSYTFEDVPFHSSYAHQS
LDRLMNPLIDQYLYLNRNTQNSGSAQNKDLLFSRGSAPAGMSVQPKNWLPGPCYRQQRVSKTKTDNNNSNFTWTGASKYNLNGRES
IINPGTAMASHKDDKDKFFPMGSGVMIFGKESAGASNTALDNVMTDEEEIKATNPVATERFGTVAVNLQSSSTDPATGDVHVMGAL
PGMVWQDRDVYLQGP I WAKIPHTDGHFHPSPMLGGFGLKHPPPQILIKNTPVPANPPAEFSATKFASFITQYSTGQVSVIEIEWELQ
KENSKRWNPEVQYTSNYAKSANVDFTVDNNGLYTEPRPIGTRYLTRPL

[0088] The VP1 sequence of AAV7 is set out as SEQ ID NO:8 and the locations of the associated variable regions and GBS and GH loop regions are defined in Table 2 below.

MAADGYLPDWLEDNLSEGIREWDLKPGAPKPKANQQKQDNGRGLVLPGYKYLGPFGNLDKGEPVNAADAAALEHDKAYDQQLKAG
 DNPYLRYNHADADEFQERLQEDTSFGGNLGRAVFQAKKRVLEPLGLVEEGAKTAPAKKRPVEPSPQRSPDSSTGIGKKGQQPARKRL
 NFGQTDGSESVDPDPQLGEPPAAPSSVSGSTVAAGGGAPMADNNEGADGVGNASGNWHCDSTWLGDVRVITTTSTRTWALPTYNNHLY
 KQISSETAGSTNDNTYFGYSTPWGYFDNRFHCHFSPRDWQRLINNNWGFPRPKLRFLFNIQVKEVTTNDGVTTIANNLTSTIQV
 FSDSEYQLPYVLGSAHQGCLPPFPADVFMIPQYGYLTLNNGSQSVGRSSFYCLEYFSPQMLRTGNNFEFSYSFEDVPFHSSYAHSQ
 SLDRMLNPLIDQYLYYLARTQSNPGGTAGNRELQFYQGGPSTMAEQAKNWLPGPCFRQQRVSKTLDQNNNSNFAWTGATKYHLNGR
 NSLVNPGVAMATHKDDERDFPSSGVLI FGKTGATNKTTLENVLMTNEEEIRPTNPVATEEYGVSSNLQAANTAAQTQVNNQGA
 LPGMVWQNRDVYLGQPIWAKIPHTDGNFHPSPLMGGFGLKHPPPQILIKNTVPANPPEVFTPAKFASFITQYSTGQVSVEIEWEL
 QKENS KRWNPEIQYTSNFEKQTGVDFAVDSQGVYSEPRPIGTRYLTRNL

[0089] The VP1 sequence of AAV8 is set out as SEQ ID NO:9 and the locations of the associated variable regions and GBS and GH loop regions are defined in Table 2 below.

MAADGYLPDWLEDNLSEGIREWALKPGAPKPKANQQKQDDGRGLVLPGYKYLGPFGNLDKGEPVNAADAAALEHDKAYDQQLQAG
 DNPYLRYNHADADEFQERLQEDTSFGGNLGRAVFQAKKRVLEPLGLVEEGAKTAPGKKRPVEPSPQRSPDSSTGIGKKGQQPARKRL
 NFGQTDGSESVDPDPQLGEPPAAPSGVGPNMTAAGGGAPMADNNEGADGVSSSGNWHCDSTWLGDVRVITTTSTRTWALPTYNNHLY
 KQISNGTSGGATNDNTYFGYSTPWGYFDNRFHCHFSPRDWQRLINNNWGFPRKRLSFKLFNIQVKEVTQNEGKTIANNLTSTIQ
 VFTDSEYQLPYVLGSAHQGCLPPFPADVFMIPQYGYLTLNNGSQAVGRSSFYCLEYFSPQMLRTGNNFQFTYTFEDVPFHSSYAHS
 QSLDRMLNPLIDQYLYYLSRTQTGGTANTQTGLFSQGGPNTMANQAKNWLPGPCYRQQRVSTTTGQNNNSNFAWTAGTKYHLNGR
 NSLANPGIAMATHKDDERDFPSSGILIFGKQNAARDNADYSVMLTSEEEIKTTNPVATEEYGIVADNLQQQNTAPQIGTVNSQG
 ALPGMVWQNRDVYLGQPIWAKIPHTDGNFHPSPLMGGFGLKHPPPQILIKNTVPADPPTTFNQSKLNSFITQYSTGQVSVEIEWE
 LQKENS KRWNPEIQYTSNYYKSTSVDFAVNTEGVYSEPRPIGTRYLTRNL

[0090] The VP1 sequence of AAV9 is set out as SEQ ID NO:10 and the locations of the associated variable regions and GBS and GH loop regions are defined in Table 2 below.

MAADGYLPDWLEDNLSEGIREWALKPGAPQPKANQQHQDNARGLVLPGYKYLGPFGNLDKGEPVNAADAAALEHDKAYDQQLKAG
 DNPYLKYNHADADEFQERLKEDTSFGGNLGRAVFQAKKRLLEPLGLVEEAAKTAPGKKRPVEQSPQEPDSSAGIGKSGAQPAKKRLN
 FGQTDGTEVPDPQPIGEPPAAPSGVGSGLTMAAGGGAPVADNNEGADGVSSSGNWHCDSQWLGDVRVITTTSTRTWALPTYNNHLYK
 QISNSTSGGSSNDNAYFGYSTPWGYFDNRFHCHFSPRDWQRLINNNWGFPRKRLNFKLFNIQVKEVTDNNGVKTIANNLTSTVQV
 FTDSYQLPYVLGSAHEGCLPPFPADVFMIPQYGYLTLNDGSAVGRSSFYCLEYFSPQMLRTGNNFQFSYEFENVPFHSSYAHSQ
 SLDRMLNPLIDQYLYYLSKTINGSGQNQQTLKFSVAGPSNMAVQGRNYIPGPSYRQQRVSTTVTQNNNEFAWPGASSWALNGRNS
 LMNPGPAMASHKEGEDRFPPLSGSLIFGKQGTGRDNVDADKVMITNEEEIKTTNPVATESYGVATNHQSAQAQAQTGWVQNGIL
 PGMVWQDRDVYLGQPIWAKIPHTDGNFHPSPLMGGFGMKHPPPQILIKNTVPADPPTAFNKDKLNSFITQYSTGQVSVEIEWELQ
 KENS KRWNPEIQYTSNYYKSNVVEFAVNTEGVYSEPRPIGTRYLTRNL

[0091] The VP1 sequence of AAVrh.8 is set out as SEQ ID NO:11 and the locations of the associated variable regions and GBS and GH loop regions are defined in Table 2 below.

MAADGYLPDWLEDNLSEGIREWDLKPGAPKPKANQQKQDDGRGLVLPGYKYLGPFGNLDKGEPVNAADAAALEHDKAYDQQLKAG
 DNPYLRYNHADADEFQERLQEDTSFGGNLGRAVFQAKKRVLEPLGLVEEGAKTAPGKKRPVEQSPQEPDSSSGIGKTGQQAkkRLN
 FGQTDGSESVDPDPQLGEPPAAPSGLGPNTMASGGGAPMADNNEGADGVGNSSGNWHCDSTWLGDVRVITTTSTRTWALPTYNNHLYK
 QISNGTSGGSTNDNTYFGYSTPWGYFDNRFHCHFSPRDWQRLINNNWGFPRKRLNFKLFNIQVKEVTTNEGKTIANNLTSTVQV
 FTDSEYQLPYVLGSAHQGCLPPFPADVFMVPQYGYLTLNNGSQALGRSSFYCLEYFSPQMLRTGNNFQFSYTFEDVPFHSSYAHSQ
 SLDRMLNPLIDQYLYYLVRTQTGTGGTQTlafSQAGPSSMANQARNWVPGCYRQQRVSTTTNQNNSNFAWTGAakFKLNGRDS
 LMNPGVAMASHKDDDRFPSSGVLI FGKQAGNDGVDYSQVLTDEEEIKATNPVATEEYGAVAINNQAANTQAQTGLVHNQGVI

PGMVWQNRDVYLQGPWQPIWAKIPHTDGNFHPSPMLMGGFGLKHPPPQILIKNTVPADPPLTFNQAKLNSFITQYSTGQVSVEIEWELQ
KENSKRWNPFIQYTSNYYKSTNVDFAVNTEGVYSEPRPIGTRYLTRNL

[0092] The VP1 sequence of AAVrh.10 is set out as SEQ ID NO:12 and the locations of the associated variable regions and GBS and GH loop regions are defined in Table 2 below.

MAADGYLPDWLEDNLSEGIREWDLKPGAPKPKANQKQDDGRGLVLPGYKYLGPFGNLDKGEPVNAADAAALEHDKAYDQQLKAG
DNPYLRYNHADADEFQERLQEDTSFGGNLGRAVFQAKKRVLEPLGLVEEGAKTAPGKKRPVEPSPQSPDSSTGIGKKGQQPAKKRL
NFGQTGDSESVDPDPQIGEPAPGSGLGSGTMAAGGGAPMADNNEGADGVGSSSGNWHCDSTWLGDRVITTTSTRTWALPTYNNHLY
KQISNGTSGGSTNDNTYFGYSTPWGYFDFNRHFHCHFSRDPWQRLINNNWGFPRKRLNFKLFNIQVKEVTQNEGKTIANNLSTSIQ
VFTDSEYQLPYVLGSAHQGCLPPFPADVFMIQYGYLTLNNGSQAVGRSSFYCLEYFPPSQMLRTGNNFEFSYQFEDVPFHSSYAH
QSLDRMLNPLIDQYLYLSRTQSTGGTAGTQQLLFSQAGPNNMSAQAKNWLPGPCYRQQRVSTTSLQNNNSNFAWTGATKYHLNGR
DSLVPNGVAMATHKDEERFFPSSGVLIMFGKQAGKDNVDYSSVMTSEEEIKTTNPVATEQYGVVADNLQQQNAAPVIGAVNSQG
ALPGMVWQNRDVYLQGPWQPIWAKIPHTDGNFHPSPMLMGGFGLKHPPPQILIKNTVPADPPTTFSQAKLASFITQYSTGQVSVEIEWE
LQKENSKRWNPFIQYTSNYYKSTNVDFAVNTDGTYSERPRPIGTRYLTRNL

[0093] The VP1 sequence of AAVanc80 is set out as SEQ ID NO:13 and the locations of the associated variable regions and GBS and GH loop regions are defined in Table 2 below.

MAADGYLPDWLEDNLSEGIREWDLKPGAPKPKANQKQDDGRGLVLPGYKYLGPFGNLDKGEPVNAADAAALEHDKAYDQQLKAG
DNPYLRYNHADADEFQERLQEDTSFGGNLGRAVFQAKKRVLEPLGLVEEGAKTAPGKKRPVEQSPQEPDSSSGIGKKGQQPARKRLN
FGQTGDSESVDPDPQPLGEPPAAPSGVGSNTMAAGGGAPMADNNEGADGVGNASGNWHCDSTWLGDRVITTTSTRTWALPTYNNHLYK
QISSQSGGSTNDNTYFGYSTPWGYFDFNRHFHCHFSRDPWQRLINNNWGFPRKRLNFKLFNIQVKEVTNDGTTTIANNLSTVQVF
TDSEYQLPYVLGSAHQGCLPPFPADVFMIQYGYLTLNNGSQAVGRSSFYCLEYFPPSQMLRTGNNFQFSYTFEDVPFHSSYAHSSQ
LDRMLNPLIDQYLYLSRTQTTSAGTAGNRTLQFSQAGPSSMANQAKNWLPGPCYRQQRVSKTTNQNNSNFAWTGATKYHLNGRDS
LVNPGPAMATHKDEDEDKFFPMGVLIFGKQAGNSNVDLDNVMITNEEEIKTTNPVATEEYGTATNLQSANTAPATGTVNSQAGAL
PGMVWQDRDVYLQGPWQPIWAKIPHTDGHFHPSPMLMGGFGLKHPPPQILIKNTVPANPPTTFSQAKFASFITQYSTGQVSVEIEWELQ
KENSKRWNPFIQYTSNYYKSTNVDFAVDNGVYSEPRPIGTRYLTRNL

[0094] The VP1 sequence of AAVanc110 is set out as SEQ ID NO:14 and the locations of the associated variable regions and GBS and GH loop regions are defined in Table 2 below.

MAADGYLPDWLEDNLSEGIREWDLKPGAPKPKANQKQDDGRGLVLPGYKYLGPFGNLDKGEPVNAADAAALEHDKAYDQQLKAG
DNPYLRYNHADADEFQERLQEDTSFGGNLGRAVFQAKKRVLEPLGLVEEGAKTAPGKKRPVEQSPQEPDSSSGIGKTGQQPAKKRLN
FGQTGDSESVDPDPQPLGEPPAAPSGVGSNTMASGGGAPMADNNEGADGVGNSSGNWHCDSTWLGDRVITTTSTRTWALPTYNNHLYK
QISNGTSGGSTNDNTYFGYSTPWGYFDFNRHFHCHFSRDPWQRLINNNWGFPRKRLNFKLFNIQVKEVTNEGKTIANNLSTVQVF
FTDSEYQLPYVLGSAHQGCLPPFPADVFMIQYGYLTLNNGSQAVGRSSFYCLEYFPPSQMLRTGNNFQFSYTFEDVPFHSSYAHSSQ
SLDRMLNPLIDQYLYLSRTQTGTAGTQTLQFSQAGPSSMANQARNWVPGPCYRQQRVSTTTNQNNSNFAWTGATKYHLNGRDS
LMNPGVAMASHKDEDRFFPSSGVLIFGKQAGNDNVDYSQVMITNEEEIKTTNPVATEEYGAVATNNQSANTQAQTGLVHNQGV
PGMVWQNRDVYLQGPWQPIWAKIPHTDGNFHPSPMLMGGFGLKHPPPQILIKNTVPADPPTTFNQAKLNSFITQYSTGQVSVEIEWELQ
KENSKRWNPFIQYTSNYYKSTNVDFAVNTEGVYSEPRPIGTRYLTRNL

VP1 of Novel Capsid Proteins

[0095] Novel AAV VP1 capsid proteins were isolated from tissue from the following mammals: baboon, crab-eating macaque, cynomolgus macaque, marmoset and pig.

[0096] The VP1 sequence of a novel AAV capsid isolated from baboon (denoted as Bba.21) is set out as SEQ ID NO:15 (amino acids 1-742) and the locations of the associated variable regions and GBS and GH loop regions are defined in Table 2 below. The VP2 capsid protein spans amino acids 138-742 of SEQ ID NO:15 and the VP3 capsid protein spans amino acids 206-742 of SEQ ID NO:15.

MAADGYLPDWLEDNLSEGIREWALKPGAPQPKANQQHQDNARGLVLPGYKYLPGNGLDKGEPVNEADAAALEHDKAYDQQLKSG
 DNPYLYKNHADADEFQQRLATDTSFGGNLGRAVFAKAKRILEPLGLVEEGVKTAPGKKRPLEKTPNRPTNPDSGKAPAKKKQKDGET
 ADSARRTLDFEDSGAGDGPPEGSSSGEMSHDAEMRAAPGGNAVEAGQGADGVGNASGDWHCDSTWSEGRVTTTTSTRTWVLPITYNNH
 LYLRI GTTANSNTYNGFSTPWGYFDNRFHCHFSRDPWQRLINNNWGLRPKSMRVKIFNIQVKEVTTSTNGETTANNLTSTVQIFFA
 DSTYELPYVMDAGQEGSLPPFPNDVFMVPQYGYCGVVTGENQNQTDNFAFYCLEYFPSQMLRTGNNFEVSYQFEKVPFHSYAHSQ
 SLDRMMNPLLDQYLWHPQSTTTGNSLNQGTATTTYGKITTGDFAYYRKNWLPGACIKQQKFSKNASQNYKIPASGGDALKYDTHT
 TSNGRWSNMAPGPPMATAGAGDSDFSNSQLIFAGPNQSGNTTSSNNLLFTSEEEIATTNPRDMDMFGQIADNNQNATTAPHIANL
 DAMGIVPGMVWQNRDIYYLGPIWAKVPHTDGHFHPSPLMGGFGLKHPPPQIFIKNTPVPANPNTTFSARINSFLTQYSTGQVAVQ
 IDWEIQKEHSKRWDPEVQFTSNYGTQNSMLWAPDNAGNYHEPRAIGSRFLTHHL

[0097] The VP1 sequence of a novel AAV capsid isolated from baboon (denoted as Bba.26) is set out as SEQ ID NO:16 (amino acids 1-739) and the locations of the associated variable regions and GBS and GH loop regions are defined in Table 2 below. The VP2 capsid protein spans amino acids 138-739 of SEQ ID NO:16 and the VP3 capsid protein spans amino acids 206-739 of SEQ ID NO:16.

MAADGYLPDWLEDNLSEGIREWALKPGAPQPKANQQHQGNSRGLVLPGYKYLGLFNGLDKGEPVNEADAAALEHDKAYDKQLEQG
 DDPYLYKNHADADEFQERLQEDTSFGGNLGRAVFAKAKRILEPLGLVEEAAKTAPGKKRPLEKTPNQPTDTQAAGQTPAKKRPQGEQ
 SGDSARRQLDFGPQPAAPIGQPPAAPPVGSNTMASGGGPMADDNQADGVGNASGNWHDSTWLGDRVITTTSTRTWVLPITYNNH
 IYKQISSSESATNDNHYFGYSTPWGYFDNRFHCHFSRDPWQRLINNNWGFPRKRLNFKIFNVQVKEVTQIEGGSTIANNLSTIQ
 VFADSEYQLPYVLGSAHQGLPPFPADVFMIPQYGYLTLLNNGSQSVGRSSFYCLEYFPSQMLRTGNNFQFSYTFESVPFHSSYAH
 QSLDRIMNPLVDQYLYLARTQTGTGSTTSNTRQLQFYQAGPSNMADQSRNWLPGPMYRQQRVSKTLDQNSNTNFAWTAASKYNLN
 GRKSLANPGIAMATHKDDEERFFPQHGVLVFGQTATNKTTLDNVLVTSEEEIKATNPVATEEYGTVSSNLQASNTNPTTETVNNQ
 GILPGMVWQDRDVYLQGPWAKIPHTDGHFHPSPLMGGFGLKHPPPQILIKNTPVPANPPETFTTSKFASYITQYSTGQVSVEIEW
 ELQKENS KRWNPEIQYTSNYAKSNNVEFSVDAAGVYSEPRPIGTRYLTRNL

[0098] The VP1 sequence of a novel AAV capsid isolated from baboon (denoted as Bba.27) is set out as SEQ ID NO:17 (amino acids 1-739) and the locations of the associated variable regions and GBS and GH loop regions are defined in Table 2 below. The VP2 capsid protein spans amino acids 138-739 of SEQ ID NO:17 and the VP3 capsid protein spans amino acids 206-739 of SEQ ID NO:17.

VAADGYLPDWLEDNLSEGIREWALKPGAPQPKANQQHQDNSRGLVLPGYKYLPGFNGLDKGEPVNEADAAALEHDKAYDKQLEQG
 DNPYLYKNHADAELQERLQEDTSFGGNLGRAVFAKAKRILEPLGLVEEAAKTAPGKKRPLEKTPNQPTDTQAAGQTPAKKRPQGEQ
 SGDSARRQLDFDPQPAAPIGQPPAAPPVGSNTMASGGGPMADDNQADGVGNASGNWHRDSTWLGDRVITTTSTRTWVLPITYNNH
 IYKQISSSESATNDNHYFGYSTPWGYFDNRFHCHFSRDPWQRLINNNWGFPRKRLNFKIFNIQVKEVTQIEGGSTIANNLSTIQ
 VFADSEYQLPYVLGSAHQGLPPFPADVFMIPQYGYLTLLNNGSQSVGRSSFYCLEYFPSQMLRTGNNFQFSYTFESVPFHSSYAH

QSLDRIMNPLVDQYLYLARTQTGTGTTNTRQLQFYQAGPSNMADQSRNWLPGPMYRQQRVSKTLDQNSNTNFAWTAASKYNLN
GRKSLANPGIAMATHKDDEERFFPQHGVLFVFGQTNATNKTTLDNVLITSEEEIKATNPVATEEYGTVSSNLQASNTNPTTETVNSQ
GILPGMVWQDRDVYLQGPWAKIPHTDGHFHPSPLMGGFGLKHPPPQILIKNTPVPANPPETFTTSKFASYITQYSTGQVSVEIEW
ELQKENGKRWNPEIQYTSNYAKSNNVEFSVDAAGVYSEPRPIGTRYLTRNL

[0099] The VP1 sequence of a novel AAV capsid isolated from baboon (denoted as Bba.29) is set out as SEQ ID NO:18 (amino acids 1-739) and the locations of the associated variable regions and GBS and GH loop regions are defined in Table 2 below. The VP2 capsid protein spans amino acids 138-739 of SEQ ID NO:18 and the VP3 capsid protein spans amino acids 206-739 of SEQ ID NO:18.

MAADGYLPDWLEDNLSEGIREWALKPGAPQPKANQQHQDNSRGLVLPGYKYLGPFGNLDKGEPVNEADAAALEHDKAYDKQLEQG
DNPYLKYNHADADEFQERLQEDTSFGGNLGRAVFQAKKRILEPLGLVEEAAKTAPGKKRPLEKTPNQPTDTQAAGQTPAKKRPQGEQ
SGDSARRQLDFDPQPAAPIGQPPAAPSPVGSNTMASGGGGPMADDNQADGVGNASGNWHCDSTWLGDRVITTTSTRTWVLTYYNNH
IYKQISSESGATNDNHYFGYSTPWGYFDNRFHCHFSRDPWQRLINNNWGFPRKRLNFKIFNIQVKEVTQIEGGSTIANNLSTTIQ
VFADSEYQLPYVLGSAHQGCLPPFPADVFMIPQYGYLTLLNNGSQSVGRSSFYCLEYFPPSQMLRTGNMFQSYTFESVPFHSSYAH
QSLDRIMNPLVDQCLYYLARTQTGTGTTNTRQLQFYQAGPSNMADQSRNWLPGPMYRQQRVSKTLDQNSNTNFAWTAASKYNLN
GRKSLANPGIAMATHKDDEERFFPQHGVLFVFGQTNATNKTTLDNVLITSEEEIKATNPVATEEYGTVSSNLQASNTNPTTETVNNQ
GILPGMVWQDRDVYLQGPWAKIPHTDGHFHPSPLMGGFGLKHPPPQILIKNTPVPANPPETFTTSKFASYITQYSTGQVSVEIEW
ELQKEDSKRWNPFIQYTSNYAKSNNVEFSVDAAGVYSEPRPIGTRYLTRNL

[00100] The VP1 sequence of a novel AAV capsid isolated from baboon (denoted as Bba.30) is set out as SEQ ID NO:19 (amino acids 1-739) and the locations of the associated variable regions and GBS and GH loop regions are defined in Table 2 below. The VP2 capsid protein spans amino acids 138-739 of SEQ ID NO:19 and the VP3 capsid protein spans amino acids 206-739 of SEQ ID NO:19.

MAADGYLPDWLEDNLSEGIREWALKPGAPQPKANQQHQDNSRGLVLPGYKYLGPFGNLDKGEPVNEADAAALEHDKAYDKQLEQG
DNPYLKYNHADADEFQERLQEDTSFGGNLGRAVFQAKKRILEPLGLVEEAAKTAPGKKRPLEKTPNQPTDTQAAGQTPAKKRPQGGQ
SGDSARRQLDFDPQPAAPIGQPPAAPSPVGSNTMASGGGGPMADDNQADGVGNASGNWHCDSTWLGDRVITTTSTRTWVLTYYNNH
IYKQISSESGATNDNHYFGYSTPWGYFDNRFHCHFSRDPWQRLINNNWGFPRKRLNFKIFNIQVKEVTQIEGGSTIANNLSTTIQ
VFADSEYLLPYVLGSAHQGCLPPFPADVFMIPQYGYLTLLNNGSQSVGRSSFYCLEYFPPSQMLRTGNMFQSYTFESVPFHSSYAH
QSLDRIMNPLVDQYLYLARTQTGTGTTNTRQLQFYQAGPSNMADQSRNWLPGPMYRQQRVSKTLDQNSNTNFAWTAASKYNLN
GRKSLANPGIAMATHKDDEERFFPQHGVLFVFGQTNATNKTTLDNVLITSEEEIKATNPVATEEYGTVSSNLQASNTNPTTETVNNQ
GILPGMVWQDRDVYLQGPWAKIPHTDGHFHPSPLMGGFGLKHPPPQILIKNTPVPANPPETFTTSKFASYITQYSTGQVSVEIEW
ELQKEDSKRWNPFIQYTSNYAKSNNVEFSVDAAGVYSEPRPIGTRYLTRNL

[00101] The VP1 sequence of a novel AAV capsid isolated from baboon (denoted as Bba.31) is set out as SEQ ID NO:20 (amino acids 1-742) and the locations of the associated variable regions and GBS and GH loop regions are defined in Table 2 below. The VP2 capsid protein spans amino acids 138-742 of SEQ ID NO:20 and the VP3 capsid protein spans amino acids 206-742 of SEQ ID NO:20.

MAADGYLPDWLEDNLSEGIREWALKPGAPQPKANQQHQDNARGLVLPGYKYLGPNGLDKGEPVNEADAAALEHDKAYDQQLKSG
 DNPYLKYNHADADEFQQLATDTSFGGNLGAQVFAKKRILEPLGLVEEGVKTAPGKKHPLEKTPNRPTNPDSGKAPAKKKQKDGET
 ADSARRTLDFEDSGAGDGPPEGSSSGEMSHDAEMRAAPGGNAVEAGQGADGVGNASGDWHCGSTWSEGRVTTTTSTRTWVLTPTNNH
 LYLRI GTTANSNTYNGFSTPWGYFDNRFHCHFSRDPWQRLINNNWGLRPKSMRVKIFNIQVKEVTTSNGETTVANNLTSTVQIFA
 DSTYELPYVMDAGQEGSLPPFPNDVFMVPQYGYCGVVTGENQNQTDNRNAYCYLCEFPQMLRTGNNFEISYQFEKVPFHSYAHSQ
 SLDRMMNPLLDQYLWHLQSTTTGNSLNQGTATTTYGKITTGDFAYYRKNWLPGACIKQKFSKNASQNYKIPASGGDALLKYDTH
 TLNGRWSNMAPGPPMATAGAGDSDFSNSQLIFAGPNQSGNTTTSSNNLLFTSEEEIATNPRDMDMFGQIADNNQATTAPHIANL
 DAMGIVPGMVWQNRDIYYQGPIWAKVPHTDGHFHPSPLMGGFGLKHPPQIFIKNTPVPANPNTTFSARINSFLTQYSTGQVAVQ
 IDWEIQKEHSKRWNPEVQFTSNYGTQNSMLWAPDNAGNYHEPRAIGSRFLTHHL

[00102] The VP1 sequence of a novel AAV capsid isolated from baboon (denoted as Bba.32) is set out as SEQ ID NO:21 (amino acids 1-742) and the locations of the associated variable regions and GBS and GH loop regions are defined in Table 2 below. The VP2 capsid protein spans amino acids 138-742 of SEQ ID NO:21 and the VP3 capsid protein spans amino acids 206-742 of SEQ ID NO:21.

MAADGYLPDWLEDNLSEGIREWALKPGAPQPKANQQHQDNARGLVLPGYKYLGPNGLDKGEPVNEADAAALEHDKAYDQQLKSG
 DNPYLKYNHADADEFQQLATDTSFGGNLGAQVFAKKRILEPLGLVEEGVKTAPGKKRPLEKTPNRPTNPDSGKAPAKKKQKDGET
 ADSARRTLDFEDSGAGDGPPEGSSSGEMSHDAEMRAAPGGNAVEAGQGADGVGNASGGWHDSTWSEGRVTTTTSTRTWVLTPTNNH
 LYLRI GTTANSNTYNGFSTPWGYFDNRFHCHFSRDPWQRLINNNWGLRPKSMRVKIFNIQVKEVTTSNGETTVANNLTSTVQIFA
 DSTYELPYVMDAGQEGSLPPFPNDVFMVPQYGYCGVVTGENQNQTDNRNAYCYLCEFPQMLRTGNNFEISYQFEKVPFHSYAHSQ
 SLDRMMNPLLDQYLWHLQSTTTGNSLNQGTATTTYGKITTGDFAYYRKNWLPGACIKQKFSKNASQNYKIPASGGDALLKYDTH
 TLNGRWSNMAPGPPMATAGAGDSDFSNSQLIFAGPNQSGNTTTSSNNLLFTSEEEIATNPRDMDMFGQIADNNQATTAPHIANL
 DAMGIVPGMVWQNRDIYYQGPIWAKVPHTDGHFHPSPLMGGFGLKHPPQIFIKNTPVPANPNTTFSARINSFLTQYSTGQVAVQ
 IDWEIQKEHSKRWNPEVQFTSNYGTQNSMLWAPGNAGNHHEPRAIGSRFLTHHL

[00103] The VP1 sequence of a novel AAV capsid isolated from baboon (denoted as Bba.33) is set out as SEQ ID NO:22 (amino acids 1-742) and the locations of the associated variable regions and GBS and GH loop regions are defined in Table 2 below. The VP2 capsid protein spans amino acids 138-742 of SEQ ID NO:22 and the VP3 capsid protein spans amino acids 206-742 of SEQ ID NO:22.

MAADGYLPDWLEDNLSEGIREWALKPGAPQPKANQQHQDNARGLVLPGYKYLGPNGLDKGEPVNEADAAALEHDKAYDQQLKSG
 DNPYLKYNHADADEFQQLATDTSFGGNLGAQVFAKKRILEPLGLVEEGVKTAPGKKRPLEKTPNRPTNPDSGRAPAKKKQKDGET
 ADSARRTLDFEDSGAGDGPPEGSSSGEMSHDAEMRAAPGGNAVEAGQGADGVGNASGDWHCDSTWSEGRVTTTTSTRTWVLTPTNNH
 LYLRI GTTANSNTYNGFSTPWGYFDNRFHCHFSRDPWQRLINNNWGLRPKSMRVKIFNIQVKEVTTSNGETTVANNLTSTVQIFA
 DSTYELPYVMDAGQEGSLPPFPNDVFMVPQYGYCGVVTGENQNQTDNRNAYCYLCEFPQMLRTGNNFEISYQFEKVPFHSYAHSQ
 SLDRMMNPLLDQYLWHLQSTTTGNSLNQGTATTTYGKITTGDFAYYRKNWLPGACIKQKFSKNASQNYKIPASGGDALLKYDTH
 TLNGRWSNMAPGPPMATAGAGDSDFSNSQLIFAGPNQSGNTTTSSNNLLTSEEEIATNPRDMDMFGQIADNNQATTAPHIANL
 DAMGIVPGMVWQNRDIYYQGPIWAKVPHTDGHFHPSPLMGGFGLKHPPQIFIKNTPVPANPNTTFSARINSFLTQYSTGQVAVQ
 IDWEIQKEHSKRWNPEVQFTSNYGTQNSMLWAPGNAGNYHEPRAIGSRFLTHHL

[00104] The VP1 sequence of a novel AAV capsid isolated from baboon (denoted as Bba.34) is set out as SEQ ID NO:23 (amino acids 1-739) and the locations of the associated variable

regions and GBS and GH loop regions are defined in Table 2 below. The VP2 capsid protein spans amino acids 138-739 of SEQ ID NO:23 and the VP3 capsid protein spans amino acids 206-739 of SEQ ID NO:23.

```
MAADGYLPDWLEDNLSEGIREWALKPGAPQPKANQQHQDNSRGLVLPGYKYLGPFGNLDKGEPVNEADAAALEHDKAYDKQLEQG
DNPYLKYNHADADEFQERLQEDTSFGGILGRAVFQAKKRILEPLGLVEEAAKTAPGKKRPLEKTPNQPTDTQAAGQTPAKKRPQGEQ
SGDSARRQLDFDPQPAAPIGQPPAAPSPVGSNTMASGGGGPMADDNQGADGVGNASGNWHCDSTWLGDRVITTTSTRTWVLTYYNNH
IYKQISSESGATNDNHYFGYSTPWGYFDFNRFHCHFSRDLWQRLINNNWGFPRKRLNFKIFNIQVKEVTQIEGGSTIANNLSTIQ
VFADSEYQLPYVLGSAHQGLPPFPADVFMIPQYGYLTLNNGSQSVGRSSFYCLEYFPPSQMLRTGNNFQFSYTFESVPFHSSYAHS
QSLDRIMNPLVDQYLYYLARTQTGTGTTSTNTRQLQFYQAGPSNMADQSRNWLPGPMYRQQRVSKTLDQNSNTNFAWTAASKYNLN
GRKSLVNPGLIAMATHKDDERFFPQHGVLFVFGKTNATNKTTLENVLVTDEEEVKATNPVATEEYGTVSSNLQSNNTNPTTETVNNQ
GILPGMVWQDRDVYLQGPVWAKIPHTDGHFHPSPLMGGFGLKHPPQILIKNTVPANPPETFTTSKFASYITQYSTGQVSVEIEW
ELQKENSkrwnPEIQYTSNYAKSNNVEFSADAAGVYSEPRPIGTRYLTRNL
```

[00105] The VP1 sequence of a novel AAV capsid isolated from baboon (denoted as Bba.35) is set out as SEQ ID NO:24 (amino acids 1-739) and the locations of the associated variable regions and GBS and GH loop regions are defined in Table 2 below. The VP2 capsid protein spans amino acids 138-739 of SEQ ID NO:24 and the VP3 capsid protein spans amino acids 206-739 of SEQ ID NO:24.

```
MAADGYLPDWLEDNLSEGIREWALKPGAPQPKANQQHQDNSRGLVLPGYKYLGPFGNLDKGEPVNEADAAALEHDKAYDKQLEQG
DNPYLKYNHADADEFQERLQEDTSFGGILGRAVFQAKKRILEPLGLVEEAAKTAPGKKRPLEKTPNQPTDTQAAGQTPAKKRPQGEQ
SGDSARRQLDFDPQPAAPIGQPPAAPSPVGSNTMASGGGGPMADDNQGADGVGNASGNWHCDSTWLGDRVITTTSTRTWVLTYYNNH
IYKQISSESGATNDNHYFGYSTPWGYFDFNRFHCHFSRDLWQRLINNNWGFPRKRLNFKIFNIQVKEVTQIEGGSTIANNLSTIQ
VFADSEYQLPYVLGSAHQGLPPFPADVFMIPQYGYLTLNNGSQSVGRSSFYCLEYFPPSQMLRTGNNFQFSYTFESVPFHSSYAHS
QSLDRIMNPLVDQYLYYLARTQTGTGTTSTNTRQLQFYQAGPSNMADQSRNWLPGPMYRQQRVSKTLDQNSNTNFAWTAASKYNLN
GRKSLANPGLIAMATHKDDERFFPQHRVLFVFGQTNATNKTTLDNVLITSEEEIKATNPVATEEYGTVSSNLQASNTDPTTETVNNQ
GILPGMVWQDRDVYLQGPVWIKIPHTDGHFHPSPLMGGFGLKHPPQILIKNTVPANPPETFTTSKFASYTTQYSTGQVSVEIEW
GLQKENSkrwnPEIQYTSNYAKSNNVEFSVDAAGVYSEPRPIGTRYLTRNL
```

[00106] The VP1 sequence of a novel AAV capsid isolated from baboon (denoted as Bba.36) is set out as SEQ ID NO:25 (amino acids 1-739) and the locations of the associated variable regions and GBS and GH loop regions are defined in Table 2 below. The VP2 capsid protein spans amino acids 138-739 of SEQ ID NO:25 and the VP3 capsid protein spans amino acids 206-739 of SEQ ID NO:25.

```
MAADGYLPDWLEDNLSEGIREWALKPGAPQPKANQQHQDNSRGLVLPGYKYLGPFGNLDKGEPVNEAYAAALEHDKAYDKQLEQG
DNPYLKYNHADADEFQERLQEDTSFGGILGRAVFQAKKRILEPLGLVEEAAKTAPGKKRPLEKTPNQPTDTQAAGQTPAKKRPQGEQ
SGDSARRQLDFDPQPAAPIGQPPAAPSPVGSNTMASGGGGPMADDNQGADGVGNASGNWHCDSTWLGDRVITTTSTRTWVLTYYNNH
IYKQISSESGATNDNHYFGYSTPWGYFDFNRFHCHFSRDLWQRLINNNWGFPRKRLNFKIFNIQVKEVTQIEGGSTIANNLSTIQ
VFADSEYQLPYVLGSAHQGLPPFPADVFMIPQYGYLTLNNGSQSVGRPSFYCLEYFPPSQMLRTGNNFQFSYTFESVPFHSSYAHS
QSLDRIMNPLVDQYLYYLARTQTGTGTTSTNTRQLQFYQAGPSNMADQSRNWLPGPMYRQQRVSKTLDQNSNTNFAWTAASKYNLN
GRKSLANPGLIAMATHKDDERFFPQHGVLFVFGQTNATNKTTLDNVLITSEEEIKATNPVATEEYGTVSSNLQASNTNPTTETVNNQ
```

GILPGMVWQDRDVYLQGGPIWAKIPHTDGHFHPSPLMGGFGLKHPPQILIKNTPVPANPPETFTTSKFASYITQYSTGQVSVEIEW
ELQKENS KRWNPEIQYTSNYAKSNNVEFSVDAAGVYSEPRPIGTRYLTRNL

[00107] The VP1 sequence of a novel AAV capsid isolated from baboon (denoted as Bba.37) is set out as SEQ ID NO:26 (amino acids 1-739) and the locations of the associated variable regions and GBS and GH loop regions are defined in Table 2 below. The VP2 capsid protein spans amino acids 138-739 of SEQ ID NO:26 and the VP3 capsid protein spans amino acids 206-739 of SEQ ID NO:26.

MAADGYLPDWLEDNLSEGIREWALKPGAPQPKANQQHQDNSRGLVLPGYKYLGPFGNLDKGEPVNEADAAALEHDKAYDKQLEQG
DNPYLKYNHADADEFQERLQEDTSFGGNLGRAVFQAKKRILEPLGLVEEAAETAPGKKRPLEKTPNQPTDTQAAGQTPAKKRPQGEQ
SGDSARRQLDFDPQPAAPIGQPPAAPSPVGSNTMASGGGGPMADDNQGADGVGNASGNWHCDSTWLGDRVITTTSTRTWLPTYNH
IYKQISSESATNDNHYFGYSTPWGYFDFNRHFCHFSRDPWQRLINNNWGRPKRLNFKIFNIQVKEVTQTEGGSTIANNLSTIQ
VFADSEYQLPYVLGSAHQGCLPPFPADVFMIPQYGYLTLLNNGSQSVGRSSFYCLEYFPPSQMLRTGNNFQFSYTFESVPFHSSYAH
QSLDRIINPLVDQYLYYLARTQTGTGTTSTNTRQLQFYQAGPSNMADQSRNWLPGPMYRQQRVSKTLDQNSNTNFAWTAASKYNLN
GRKSLANPGIAMATHKDDEERFFPQHGVLVFGQTATNKTTLDNVLITSEEEIKATNPVATEEYGTVSSNLQASNTNPTTETVNNQ
GILPGMVWQDRDVYLQGGPIWAKIPHTDGHFHPSPLMGGFGLKHPPQILIKNTPVPANPPETFTTSKFASYITQYSTGQVSVEIEW
ELQKENS KRWNPEIQYTSNYAKSNNVEFSVDAAGVYSEPRPIGTRYLTRNL

[00108] The VP1 sequence of a novel AAV capsid isolated from baboon (denoted as Bba.38) is set out as SEQ ID NO:27 (amino acids 1-739) and the locations of the associated variable regions and GBS and GH loop regions are defined in Table 2 below. The VP2 capsid protein spans amino acids 138-739 of SEQ ID NO:27 and the VP3 capsid protein spans amino acids 206-739 of SEQ ID NO:27.

MAADGYLPDWLEDNLSEGI RGWALKPGAPQPKANQQHQDNSRGLVLPGYKYLGPFGNLDKGEPVNEADAAALEHDKAYDKQLEQG
DNPCLKYNHADADEFQERLQEDTSFGGNLGRAVFQAKKRILEPLGLVEEAAKTAPGKKRPLEKTPNQPTDTQAAGQTPAKKRPQGEQ
SGDSARRQLDFDPQPAAPIGQPPAAPSPVGSNTMASGGGGPMADDNQGADGVGNASGNWHCDSTWLGDRVITTTSTRTWLPTYNH
IYKQISSESATNDNHYFGYSTPWGYFDFNRHFCHFSRDPWQRLINNNWGRPKRLNFKIFNIQVKEVTQIEGGSTIANNLSTIQ
VFADSEYQLPYVLGSAHQGCLPPFPADVFMIPQYGYLTLLNNGSQSVGRSSFYCLEYFPPSQMLRTGNNFQFSYTFESVPFHSSYAH
QSLDRIMNPLVDQYLYYLARTQTGTGTTSTNTRQLQFYQAGPSNMADQSRNWLPGPMYRQQRVSKTLDQNSNTNFAWTAASKYNLN
GRKSLANPGIAMATHKDDEERFFPQHGVLVFGQTATNKTTLDNVLITSEEEIKATNPVATEEYGTVSSNLQASNTNPTTETVNNQ
GILPGMVWQDRDVYLQGGPTWAKIPHTDGHFHPSPLMGGFGLKHPPQILIKNTPVPANPPETFTTSKFASYITQYSTGQVSVEIEW
ELQKENS KRWNPEIQYTSNYAKSNNVEFSVDAAGVYSEPRPIGTRYLTRNL

[00109] The VP1 sequence of a novel AAV capsid isolated from baboon (denoted as Bba.41) is set out as SEQ ID NO:28 (amino acids 1-739) and the locations of the associated variable regions and GBS and GH loop regions are defined in Table 2 below. The VP2 capsid protein spans amino acids 138-739 of SEQ ID NO:28 and the VP3 capsid protein spans amino acids 206-739 of SEQ ID NO:28.

MAADGYLPDWLEDNLSEGIREWALKPGAPQPKANQQHQDNSRGLVLPGYKYLGPFGNLDKGEPVNEADAAALEHDKAYDKQLEQG
DNPYLKYNHADADEFQERLQEDTSFGGNLGRAVFQAKKRILEPLGLVEEAAKTAPGKKRPLEKTPNQPTDTQAAGQTPAKKRPQGEQ

SGDSARRQLDFDPQPAAPIGQPPAAPSPVGSNTMASGGGPMADDNQADGVGNASGNWHCDSTWLGDRVITTTSTRTWVLTPTNNH
 IYKQISSESGATNDNHYFGYSTPWGYFDFNRHFCHFSRDLWQRLINNNWGFPRKRLNFKIFNIQVKEVTQIEGGSTIANNLSTIQ
 VFADSEYQLPYVLGSAHQGCLPPFPADVFMIPQYGYLTLNNGSQSVGRSSFYCLEYFPPSQMLRTGNSFQFSYTFESVPFHSSYAH
 QSLDRIMNPLVDQYLYYLARTQTGTGTTSTNTRQLQFYQAGPSNMADQSRNWLPGPMYRQQRVSKTLDQNSNTNFAWTAASKYNLN
 GRKSLANPGIAMATHKDDDEERFFPQHGVLVFGQTNATNKTTLDNVLITSEEEIKATNPVATEEYGTVSSNLQASNTNPTTETVNNQ
 GILPGMVWQDRDVYLQGPWAKIPHTDGHFHPSPLMGGFGLKHPPPQILIKNTPVPANPPETFTTSKFASYITQYSTGQVSVEIEW
 ELQKENSkrwnPEIQYTSNYAKSNNVEFSVDAAGVYSEPRPIGTRYLTRNL

[00110] The VP1 sequence of a novel AAV capsid isolated from baboon (denoted as Bba.42) is set out as SEQ ID NO:29 (amino acids 1-739) and the locations of the associated variable regions and GBS and GH loop regions are defined in Table 2 below. The VP2 capsid protein spans amino acids 138-739 of SEQ ID NO:29 and the VP3 capsid protein spans amino acids 206-739 of SEQ ID NO:29.

MAADGYLPDWLEDNLSEGIREWALKPGAPQPKANQQHQDNSRGLVLPGYKYLGPFGNLDKGEPVNGADAAALEHDKAYDKQLEQG
 DNPYLKYNHADADEFQERLQEDTSFGGNLGRAVFQAKKRILEPLGLVEEAAKTAPGKKRPLEKTPNQPTDTQAAGQTPAKKRPQGEQ
 SGDSARRQLDFDPQPAAPIGQPPAAPSPVGSNTMASGGGPMADDNQADGVGNASGNWHCDSTWLGDRVITTTSTRTWVLTPTNNH
 IYKQISSESGATNDNHYFGYSTPWGYFDFNRHFCHFSRDLWQRLINNNWGFPRKRLNFKIFNIQVKEVTQIEGGSTIANNLSTIQ
 VFADSEYQLPYVLGSAHQGCLPPFPADVFMIPQYGYLTLNNGSQSVGRSSFYCLEYFPPSQMLRTGNNFQFSYTFESVPFHSSYAH
 QSLDRIMNPLVDQYLYYLARTQTGTGTTSTNTRQLQFYQAGPSNMADQSRNWLPGPMYRQQRVSKTLDQNSNTNFAWTAASKYNLN
 GRKSLVNPFIAMATHKDDDEERFFPQHGVLVFGKTNATNKTTLNVLVTDEEEVKATNPVATEEYGTVSSNLQSNNTNPTTETVNNQ
 GILPGMVWQDRDVYLQGPWAKIPHTDGHFHPSPLMGGFGLKHPPPQILIKNTPVPANPPETFTTSKFASYITQYSTGQVSVEIEW
 ELQKENSkrwnPEIQYTSNYAKSNNVEFSVDAAGVYSEPRPIGTRYLTRNL

[00111] The VP1 sequence of a novel AAV capsid isolated from baboon (denoted as Bba.43) is set out as SEQ ID NO:30 (amino acids 1-739) and the locations of the associated variable regions and GBS and GH loop regions are defined in Table 2 below. The VP2 capsid protein spans amino acids 138-739 of SEQ ID NO:30 and the VP3 capsid protein spans amino acids 206-739 of SEQ ID NO:30.

MAADGYLPDWLEDNLSEGIREWALKPGAPQPKANQQHQDNSRGLVLPGYKYLGPFGNLDKGEPVNEADAAALEHDKAYDKQLEQG
 DNPYLKYNHADADEFQERLQEDTSFGGNLGRAVFQAKKRILEPLGLVEEAAKTAPGKKRPLEKTPNQPTDTQAAGQTPAKKRPQGEQ
 SGDSARRQLDFDPQPAAPIGQPPAAPSPVGSNTMASGGGPMADDNQADGVGNASGNWHCDSTWLGDRVITTTSTRTWVLTPTNNH
 IYKQISSESGATNDNHYFGYSTPWGYFDFNRHFCHFSRDLWQRLINNNWGFPRKRLNFKIFNIQVKEVTQIEGGSTIANNLSTIQ
 VFADSEYQLPYVLGSAHQGCLPPFPADVFMIPQYGYLTLNNGSQSVGRSSFYCLEYFPPSQMLRTGNNFQFSYTFESVPFHSSYAH
 QSLDRIMNPLVDQYLYYLARTQTGTGTTSTNTRQLQFYQAGPSNMADQSRNWLPGPMYRQQRVSKTLDQNSNTNFAWTAASKYNLN
 GRKSLVNPFIAMATHKDDDEERFFPQHGVLVFGKTNATNKTTLNVLVTDEEEVKATNPVATEEYGTVSSNLQSNNTNPTTETVNNQ
 GILPGMVWQDRDVYLQGPWAKIPHTDGHFHPSPLMGGFGLKHPPPQILIKNTPVPANPPETFTTSKFASYITQYSTGQVSVEIEW
 ELQKENSkrwnPEIQYTSNYAKSNNVEFSVDAAGVYSEPRPIGTRYLTRNL

[00112] The VP1 sequence of a novel AAV capsid isolated from baboon (denoted as Bba.44) is set out as SEQ ID NO:31 (amino acids 1-739) and the locations of the associated variable regions and GBS and GH loop regions are defined in Table 2 below. The VP2 capsid protein

spans amino acids 138-739 of SEQ ID NO:31 and the VP3 capsid protein spans amino acids 206-739 of SEQ ID NO:31.

MAADGYLPDWLEDNLSEGIREWALKPGAPQPKANQQHQDNRGLVLPGYKYLGPFGNLDKGEPVNEADAAALEHDKAYDKQLEQG
DNPYLYRNHADADEFQERLQEDTSFGGNLGRAVFQAKKRILEPLGLVEEAAKTAPGKKRPLEKTPNQPTDQAAGQTPAKKRPQGEQ
SGDSARRQLDFDPQPAAPIGQPPAAPSPVGSNTMASGGGPMADDNQADGVGNASGNWHCDSTWLGDRVITTTSTRTWVLTPTNNH
IYKQISSESGATNDNHYFGYSTPWGYFDNRFHCHFSPRDWQRLINNNWGFPRKRLNFKLFNIQVKEVTQIEGGSTIANNLSTIQ
VFADSEYQLPYVLGSAHQGLPPFPADVFMIPOQGYLTLNNGSQSVGRSSFYCLEYFPPSMLRTGNNFQFSYTFESVPFHSSYHS
QSLDRIMNPLVDQYLYYLARTQTGTGTTNTRQLQFYQAGPSNMADQSRNWLPGPMYRQQRVSKTLDQDSNTNFAWTAASKYNLN
GRKSLVNPGLIAMATHKDDERFFPQHGVLFVFGKTNATNKTLENVLVTDEEEVKATNPVATEEYGTVSSNLQSNNTNPTTETVNNQ
GILPGMAWQDRDVYLQGPWAKIPHTDGHFHPSPLMGGFGLKHPPPOILIKNTVPANPPETFTTSKFASYITQYSTGQVSVIEW
ELQKENSkrwnPEIQYTSNYAKSNNVEFSVDAAGVYSEPRPIGTRYLTRNL

[00113] The VP1 sequence of a novel AAV capsid isolated from crab-eating macaque (denoted as Bce.14) is set out as SEQ ID NO:32 (amino acids 1-736) and the locations of the associated variable regions and GBS and GH loop regions are defined in Table 2 below. The VP2 capsid protein spans amino acids 138-736 of SEQ ID NO:32 and the VP3 capsid protein spans amino acids 203-736 of SEQ ID NO:32.

MAADGYLPDWLEDNLSEGIREWALKPGAPQPKANQQKQDDGRGLVLPGYKYLGPFGNLDKGEPVNEADAAALEHDKAYDQQLKAG
DNPYLYRNHADADEFQERLQEDTSFGGNLGRAVFQAKKRILEPLGLVEEAAKTAPGKKRPVEQSPQEPDSSAGIGKSGAQPAKKRLN
FGQTGDTESVPDPQPIGEPPAAPSGVGSMTMASGGGAPVADNNEGADGVGSSSGNWHCDSQWLGDVITTTSTRTWALPTNNHLYK
QISNSTSGGSSNDNAYFGYSTPWGYFDNRFHCHFSPRDWQRLINNNWGFPRKRLNFKLFNIQVKEVTDNNGVKTIANNLTSTVQV
FTDSYQLPYVLGSAHEGCLPPFPADVFMIPOQGYLTLNDGSQAVGRSSFYCLEYFPPSMLRTGNNFTFSYEFENVFPFHSSYHSQ
SLDRIMNPLIDQYLYLSKTINGSGQNQQALKFSVAGPSNMAVQGRNYIPGPSYRQQRVSTTTVTQNNNEFAWPGASSWALNGRNS
LMNPGPAMASHKEGEDRFFPLSGSLIFGKQGTGRDNVDADKVMITNEEEIKTTNPVATESYGVATNHQSAQAQAQTGWVQNQGIL
PGMVWQDRDVYLQGPWAKIPHTDGNFHPSPLVGGFGMKHPPPOILIKNTVPADPPTAFNKDKLNSFITQYSTGQVRVEIEWELQ
KENSkrwnPEIQYTSNYKSNNEFAVNTGVYSEPRPIGTRYLTRNL

[00114] The VP1 sequence of a novel AAV capsid isolated from crab-eating macaque (denoted as Bce.15) is set out as SEQ ID NO:33 (amino acids 1-736) and the locations of the associated variable regions and GBS and GH loop regions are defined in Table 2 below. The VP2 capsid protein spans amino acids 138-736 of SEQ ID NO:33 and the VP3 capsid protein spans amino acids 203-736 of SEQ ID NO:33.

MAADGYLPDWLEDNLSEGIREWALKPGAPQPKANQQKQDDGRGLVLPGYKYLGPFGNLDKGEPVNEADAAALEHDKAYDQQLKAG
DNPYLYRNHADADEFQERLQEDTSFGGNLGRAVFQAKKRILEPLGLVEEAAKTAPGKKRPVEQSPQEPDSSAGIGKSGAQPAKKRLN
FGQTGDTESVPDPQPIGEPPAAPSGVGSMTMASGGGAPVADNNEGADGVGSSSGNWHCDSQWLGDVITTTSTRTWALPTNNHLYK
QISNSTSGGSSNDNAYFGYSTPWGYFDNRFHCHFSPRDWQRLINNNWGFPRKRLNFKLFNIQVKEVTDNNGVKTIANNLTSTVQV
FTDSYQLPYVLGSAHEGCLPPFPADVFMIPOQGYLTLNDGSQAVGRSSFYCLEYFPPSMPRTGNNFTFSYEFENVFPFHSSYHSQ
SLDRIMNPLIDQYLYLSKTINGSGQNQQTLKFSVAGPSNMAVQGRNYIPGPSYRQQRVSTTTVTQNNNEFAWPGASSWALNGRNS
LMNPGPAMASHKEGEDRFFPLSGSLIFGKQGTGRDNVDADKVMITNEEEIKTTNPVATESYGVATNHQSAQAQAQTGWVQNQGIL

PGMVWQDRDVYLQGPWAKIPHTDGNFHPSPMLGGFGMKHPPQILIKNTVPADPPTAFNKDKLNSFITQYSTGQVSVAEIEWELQ
KENSKRWNPETQYTSNYYKSNNVEFAVNTGEGVYSEPRPIGTRYLTRNL

[00115] The VP1 sequence of a novel AAV capsid isolated from crab-eating macaque (denoted as Bce.16) is set out as SEQ ID NO:34 (amino acids 1-736) and the locations of the associated variable regions and GBS and GH loop regions are defined in Table 2 below. The VP2 capsid protein spans amino acids 138-736 of SEQ ID NO:34 and the VP3 capsid protein spans amino acids 203-736 of SEQ ID NO:34.

MAADGYLPDWLEDNLSEGIREWALKPGAPQPKANQQKQDDGRGLVLPGYKYLGPFGNLDKGEPVNEADAAALEHDKAYDQQLKAG
DNPYLRYNHADADEFQERLQEDTSFGGNLGRAVFQAKKRVLEPLGLVEEAAKTAPGKKRPVEQSPQEPDSSAGIGKSGAQPAKKRLN
FGQTGDTESVPDPQPIGEPPAAPSGVGSMTASGGGAPVADNNEGADGVGSSGNWHCDSQLGDRVITTSRTWALPTYNHLYK
QISNSTSGSSNDNAYFGYSTPWGYFDFNRFHCHFSRDPWQRLINNNWGFPRKRLNFKLFNIQVKEVTDNNGVKTIANNLTSTVQV
FTDSYQLPYVLGSAHEGCLPPFPADVFMIQYGYLTLDNGSQAVGRSSFYCLEYFSPQMLRTGNNFTFSYEFENVVPFHSSYASQ
SLDRPMNPLIDQYLYLSKTINGSGQNQQTLKFSVAGPSNMAVQGRNYIPGPSYRQQRVSTTVTQNNNEFAWPAGASSWALNGRNS
LMNPGPAMASHKEGEDRFFPLSGSLIFGKQGTGRDNVDADKVMITNEEEIKTTNPVATESYQQVATNHQSAQAQAQTGWVQNQGIL
PGMVWQDRDVYLQGPWAKIPHTDGNFHPSPMLGGFGMKHPPQILIKNTVPADPPTAFNKDKLNSFITQYSTGQVSVAEIEWELQ
KENSKRWNPETQYTSNYYKSNNVEFAVNTGEGVYSEPRPIGTRYLTRNL

[00116] The VP1 sequence of a novel AAV capsid isolated from crab-eating macaque (denoted as Bce.17) is set out as SEQ ID NO:35 (amino acids 1-736) and the locations of the associated variable regions and GBS and GH loop regions are defined in Table 2 below. The VP2 capsid protein spans amino acids 138-736 of SEQ ID NO:35 and the VP3 capsid protein spans amino acids 203-736 of SEQ ID NO:35.

MAADGYLPDWLEDNLSEGIREWALKPGAPQPKANQQKQDDGRGLVLPGYKYLGPFGNLDKGEPVNEADAAALEHDKAYDQQLKAG
DNPYLRYNHADADEFQERLQEDTSFGGNLGRAVFQAKKRVLEPLGLVEEAAKTAPGKKRPVEQSPQEPDSSAGIGKSGAQPAIKRLN
FGQTGDTESVPDPQPIGEPPAAPSGVGSMTASGGGAPVADNNEGADGVGSSGNWHCDSQLGDRVITTSRTWALPTYNHLYK
QISNSTSGSSNDNAYFGYSTPWGYFDFSRFHCHFSRDPWQRLINNNWGFPRKRLNFKLFNIQVKEVTDNNGVKTIANNLTSTVQV
FTDSYQLPYVLGSAHEGCLPPFPADVFMIQYGYLTLDNGSQAVGRSSFYCLEYFSPQMLRTGNNFTFSYEFENVVPFHSSYASQ
SLDRMLNPLIDQYLYLSKTINGSGQNQQTLKFSVAGPSNMAVQGRNYIPGPSYRQQRVSTTVTQNNNEFAWPAGASSWALNGRNS
LMNPGPAMASHKEGEDRFFPLSGSLIFGKQGTGRDNVDADKVMITNEEEIKTTNPVATESYQQVATNHQSAQAQAQTGWVQNQGIL
PGMVWQDRDVYLQGPWAKIPHTDGNFHPSPMLGGFGMKHPPQILIKNTVPADPPTAFNKDKLNSFITQYSTGQVSVAEIEWELQ
KENSKRWNPETQYTSNYYKSNNVEFAVNTGEGVYSEPRPIGTRYLTRNL

[00117] The VP1 sequence of a novel AAV capsid isolated from crab-eating macaque (denoted as Bce.18) is set out as SEQ ID NO:36 (amino acids 1-736) and the locations of the associated variable regions and GBS and GH loop regions are defined in Table 2 below. The VP2 capsid protein spans amino acids 138-736 of SEQ ID NO:36 and the VP3 capsid protein spans amino acids 203-736 of SEQ ID NO:36.

MAADGYLPDWLEDNLSEGIREWALKPGAPQPKANQQKQDDGRGLVLPGYKYLGPFGNLDKGEPVNEADAAALEHDKAYDQQLKAG
DNPYLRYNHADADEFQERLQEDTSFGGNLGRAVFQAKKRVLEPLGLVEEAAKTAPGKKRPVEQSPQEPDSSAGIGKSGAQPAKKRLN

FGQTGDTESVPDPQPIGEPPAAPSGVGS LTMA SGGGAPVADNNEGADGVGSSSGNWHCD S QWL GDRVIT TSTR TWALPT YNNHLYK
 QISNSTSGGSSNDNAYFGYSTPWGYFDFNRFHCHFSPRDWQRLINNNWGF RPKRLNFKLFNIQVKEVTDNNGVKTIANNLTSTVQV
 FTDSYQLPYVLGSAHEGCLPPFPADVFMI PQYGYLT LNDGSQAVGRSSFYCLEYFPSQMLRTGNNFTFSYEFENVPFHSSYAHSQ
 SLDR LMNPLIDQYLYLSKTINGSGQNQQTLKFSVAGPSNMAVQGRNYIPGPSYRQQRVSTTVTQNNNSEFAWP GASSWALNGRNS
 LMNPGPAMASHKEGEDRFFPLSGSLIFGKQGTGRDNVDADKVMITNEEEIKTTNPVATESYGVATNHQSAQAQAQTGWVQNQGIL
 PGMVWQDRDVYLQGP I WAKIPHTDGNFHPSPLMGGFGMKHPPPQILIKNTVPADPPTAFNKDKLNSFITQYSTGQVSVEIEWELQ
 KENSKRWNP EIQYTSNYYKSNNVEFAVNTEGVYSEPRPIGTRYLTRNL

[00118] The VP1 sequence of a novel AAV capsid isolated from crab-eating macaque (denoted as Bce.20) is set out as SEQ ID NO:37 (amino acids 1-733) and the locations of the associated variable regions and GBS and GH loop regions are defined in Table 2 below. The VP2 capsid protein spans amino acids 138-733 of SEQ ID NO:37 and the VP3 capsid protein spans amino acids 203-733 of SEQ ID NO:37.

MAADGYLPDWLEDNLSEGI REWWALKPGAPQPKANQQKQDDGRGLVLPGYKYLGPFGNGLDKGEPVNEADAAALEHDKAYDQQLKAG
 DNPYLRYNHADA EAFQERLQEDTSFGGNLGRAVFAKAKRVLEPLGLVEEAAKTAPGKKRPVEQSPQEPDSSAGIGKSGAQPAKKRLN
 FGQTGDTESVPDPQPIGEPPAAPSGVGS LTMA SGGGAPVADNNEGADGVGSSSGNWHCD SRWL GDRVIT TSTR TWALPT YNNHLYK
 QISNSTSGGSSNDNAYFGYSTPWGYFDFNRFHCHFSPRDWQRLINNNWGF RPKRLNFKLFNIQVKEVTDNNGAKTIANNLTSTVQV
 FTDSYQLPYVLGSAHEGCLPPFPADVFMI PQYGYLT LNDGSQAVGRSSFYCLEYFPSQMLRTGNNFTFSYEFENVPFHSSYAHSQ
 SLDR LMNPLIDQYLYLSKTINGSGQNQQTLKFSVAGPSNMAVQGRNYIPGPSYRQQRVSTTVTQNNNSEFAWP GASSWALNGRNS
 LMNPGPAMASHKEGEDRFFPLSGSLIFGKQGTGRDNVDADKVMITNEEEIKTTNPVATESYGVATNHQSAQAQAQTGWVQNQGIL
 PGMVWQDRDVYLQGP I WAKIPHTDGNFHPSPLMGGFGMKHPPPQILIKNTVPADPPTAFNKDKLNSFITQYSTGQVSVEIEWELQ
 KENSKRWNP EIQYTSNYYKSNNVEFAVNTEGVYSEPRPIGTRYLTRNL

[00119] The VP1 sequence of a novel AAV capsid isolated from crab-eating macaque (denoted as Bce.35) is set out as SEQ ID NO:38 (amino acids 1-730) and the locations of the associated variable regions and GBS and GH loop regions are defined in Table 2 below. The VP2 capsid protein spans amino acids 138-730 of SEQ ID NO:38 and the VP3 capsid protein spans amino acids 199-730 of SEQ ID NO:38.

MAADGYLPDWLEDNLSEGI REWWALKPGAPQPKANQQKQDDGRGLVLPGYKYLGPFGNGLDKGEPVNEADAAALEHDKAYDQQLKAG
 DNPYLRYNHADA EAFQERLQEDTSFGGNLGRAVLQAKAKRVLEPLGLVEEAAKTAPGKKRPVDS PDSTSGIGKKGQQPARKRLNFGQT
 GDAESVPDPQPIGEPPAAPSGLSGTMAAGGGAPMADNNEGADGVGNASGNWHCDSTWLGNRVIT TSTR TWALPT YNNHLYKQISS
 SSSGATNDNHYFGYSTPWGYFDFNRFHCHFSPRDWQRLINNNWGF RPKRLRFLFKLFNIQVKEVTTNDGVTTIANNLTSTVQVFS DSE
 YQLPYVLGSAHQGCLPPFPADVFMI PQYGYLT LNNGSQSVGRSSFYCLEYFPSQMLRTGNNFEFSYEFEDVPFHSSYAHSQSLDR L
 MNPLIDQYLYLARTQSSTGTSTRELQFHQAGPATMAEQSKNWLPGPCFRQQRISKT TDNNNSNFAWTGATKYHLNGRSSLTNPG
 VPMATHKDDES VFFPINGVLVFGKTGASNKTTLENVLMTDEEEIKATNPVATEEYGVVSSNIQSQNSNPTTQT VNNQ GALPGMVWQ
 NRDVYLQGP I WAKIPHTDGNFHPSPLMGGFGLKHPPPQILIKNTVPANPPETFTPAKFASFITQYSTGQVSVEIEWELQKENSKR
 WNPEIQYTSN YDKQTGVGFVDTQGVYSEPRPIGTRYLTRNL

[00120] The VP1 sequence of a novel AAV capsid isolated from crab-eating macaque (denoted as Bce.36) is set out as SEQ ID NO:39 (amino acids 1-730) and the locations of the associated variable regions and GBS and GH loop regions are defined in Table 2 below. The

VP2 capsid protein spans amino acids 138-730 of SEQ ID NO:39 and the VP3 capsid protein spans amino acids 199-730 of SEQ ID NO:39.

MAADGYLPDWLEDNLSEGIREWALKPGAPQPKANQQKQDDGRGLVLPGYKYLGPFGNLDKGEPVNEADAAALEHDKAYDQQLKAG
 DNPYLRYNHADADEFQERLQEDTSFGGNLGRAVFQAKKRVLEPLGLVEEAAKTAPGKKRPVDSPDSTSGIGKKQQPARKRLNFGQT
 GDAESVPDPQPIGEPPAAPSGLGSGTMAAGGAPMADNNEGADGVGNASGNWHCDSTWLGNRVITTTSTRTWALPTYNNHLYKQISS
 SSSGATNDNHYFGYSTPWGYFDNRFHCHFSPRDWQRLINNNWGFPRKRLRFKLFNIQVKEVTTNDGVTTIANNLTSTVQVSDSE
 YQLPYVLGSAHQGCLPPFPADVFMIPQYGYLTLNNGSQSVGRSSFYCLEYFPSQMLRTGNNFEFSYEFEDVPFHSSYAHSQSLDRL
 MNPLIDQYLYLARQSSSTGTSTRELQFHQAGPATMAEQSKNWLPGPCFRQQRISKTTDNNNNSNFAWTGATKYHLNGRNSLTNPG
 VPMATHKDDSVFFPINGVLVFGKTGASNKTTLENVLMTDEEEIKATNPVATEEYGVVSSNIQSQNSNPTTQTVNNQGALPGMVWQ
 NRDVYLQGPWIWAKIPHTDGNFHPSPMLMGFGGLKHPPQILIKNTPVPANPPETFTPAKFASFITQYSTGQVSVEIEWELQKENSKR
 WNPEIQYTSNYDKQTGVDFAVDTQGVYSEPRPIGTRYLTRNL

[00121] The VP1 sequence of a novel AAV capsid isolated from crab-eating macaque (denoted as Bce.39) is set out as SEQ ID NO:40 (amino acids 1-736) and the locations of the associated variable regions and GBS and GH loop regions are defined in Table 2 below. The VP2 capsid protein spans amino acids 138-736 of SEQ ID NO:40 and the VP3 capsid protein spans amino acids 203-736 of SEQ ID NO:40.

MAADGYLPDWLEDNLSEGIREWDLKPGAPKPKANQQKQDDGRGLVLPGYKYPGPFGNLDKGEPVNAADAAALEHDKAYDQQLKAG
 DNPYLRYNHADADEFQERLQEDTSFGGNLGRAVFQAKKRVFEPLGLVEEGAKTAPGKKRPVEQSPQEPDSSSGIGKTGQQPAKKRLN
 FGQTDSESVDPDPQPLGEPPAAPSGLGPNTMASGGGAPMADNNEGADGVGNSSGNWHCDSTWLGDRVITTTSTRTWALPTYNNHLYK
 QISNSTSGGSTNDNTYFGYSTPWGYFDNRFHCHFSPRDWQRLINNNWGFPRKRLNFKLFNIQVKEVTTNEGTKTIANNLTSTVQV
 FTDSEYQLPYVLGSAHQGCLPPFPADVFMVPQYGYLTLNNGSQALGRSSFYCLEYFPSQMLRTGNNFQFSYTFEDVPFHSSYAHSQ
 SLDRLMNPLIDQYLYLVRTQTTGTGGTQTAFSQAGPSSMANQARNWVPGPSYRQQRVSTTKNQNNNSNFAWTGAAKFKLNGRNS
 LMNPGVAMASHKDDERDFPSSGVLI FGKQAGNDGVDYSQVLTDEEEIKATNPVATEEYGEVAINDQAANTQAQTGLVHNQGV
 PGMVWQNRDVYLQGPWIWAKIPHTDGNFHPSPMLMGFGGLMHPPQILIKNTPVPADPPLTFNQAKLNSFITQYSTGQVSVEIEWELQ
 KENSKRWNPFIQYTSNYKSTNVDFAVNSDGVYSEPRPIGTRYLTRNL

[00122] The VP1 sequence of a novel AAV capsid isolated from crab-eating macaque (denoted as Bce.40) is set out as SEQ ID NO:41 (amino acids 1-736) and the locations of the associated variable regions and GBS and GH loop regions are defined in Table 2 below. The VP2 capsid protein spans amino acids 138-736 of SEQ ID NO:41 and the VP3 capsid protein spans amino acids 203-736 of SEQ ID NO:41.

MAADGYLPDWLEDNLSEGIREWDLKPGAPKPKANQQKQDDGRGLVLPGYKYLGPFGNLDKGEPVNAADAAALEHDKAYDQQLKAG
 DNPYLRYNHADADEFQERLQEDTSFGGNLGRAVFQAKKRVLEPLGLVEEGAKTAPGKKRPVEQSPQEPDSSSGIGKTGQQPAKKRLN
 FGQTDSESVDPDPQPLGEPPAAPSGLGPNTMASGGGAPMADNNEGADGVGNSSGNWHCDSTWLGDRVITTTSTRTWALPTYNNHLYK
 QISNSTSGGSTNDNTYFGYSTPWGYFDNRFHCHFSPRDWQRLINNNWGFPRKRNPFKLFNIQVKEVTTNEGTKTIANNLTSTVQA
 FTDSEYQLPYVLGSAHQGCLPPFPADVFMVPQYGYLTLNNGSQALGRSSFYCLEYFPSQMLRTGNNFQFSYTFEDVPFHSSYAHSQ
 SLDRLMNPLIDQYLYLVRTQTTGTGGTQTAFSQAGPSSMANQARNWVPGPSYRQQRVSTTKNQNNNSNFAWTGAAKFKLNGRNS
 LMNPGVAMASHKDDERDFPSSGVLI FGKQAGNDGVDYSQVLTDEEEIKATNPVATEEYGEVAINDQAANTQAQTGLVHNQGV
 PGMVWQNRDVYLQGPWIWAKIPHTDGNFHPSPMLMGFGGLKHPPQILIKNTPVPADPPLTFNQAKLNSFITQYSTGQVSVEIEWELQ
 KENSKRWNPFIQYTSNYKSTNVDFAVNSDGVYSEPRPIGTRYLTRNL

[00123] The VP1 sequence of a novel AAV capsid isolated from crab-eating macaque (denoted as Bce.41) is set out as SEQ ID NO:42 (amino acids 1-736) and the locations of the associated variable regions and GBS and GH loop regions are defined in Table 2 below. The VP2 capsid protein spans amino acids 138-736 of SEQ ID NO:42 and the VP3 capsid protein spans amino acids 203-736 of SEQ ID NO:42.

MAADGYLPDWLEDNLSEGIREWDLKPGAPKPKANQQKQDDGRGLVLPGYKYLGPFGNLDKGEPVNAADAAALEHDKAYDQQLKAG
 DNPYLRYNHADADEFQGRLEQDTSFGGNLGRAVFQAKKRVLEPLGLVEEGAKTAPGKKRPVEQSPQEPDSSSGIGKTGQQPAKKRLN
 FGQTGDSESVDPDPQLGEPPAAPSGLGPNTMASGGGAPMADNNEGADGVGNSSGNWHCDSTWLGDRVITTTSTRTWALPTYNNHLYK
 QISNSTSGGSTNDNTYFGYSTPWGYFDNRFHCHFSPRDWQRLINNNWGFPRKRLNFKLFNIQVKEVTTNEGKTIANNLTSTVQV
 FTDSEYQLPYVLGSAHQGLPPFPADVFMVPQYGYLTLNNGSQALGRSSFYCLEYFPSQMLRTGNNFQFSYTFEDVPFHSSYAHSQ
 SLDRMLNPLIDQYLYLVRTQTTGTGGTQILAFSQAGPSSMANQARNWVPGPSYRQQRVSTTKNQNNNSNFAWTGAAKFKLNGRNS
 LMNPGVAMASHKDDERFFPSSGVLIIFGKQAGNDGMDYSQVLI TDEEEIKATNPVATEEYGEVAINDAQANTQAQTGLVHNQGV
 PGMVWQNRDVYLQGPWAKIPHTDGNFHPSPMLGGFGLKHPPPQILIKNTVPADPPLTFNQAKLNSFITQYSTGQVSVEIEWELQ
 KENSKRWNPETQYTSNYYKSTNVDFAVNSDGVYSEPRPIGTRYLTRNL

[00124] The VP1 sequence of a novel AAV capsid isolated from crab-eating macaque (denoted as Bce.42) is set out as SEQ ID NO:43 (amino acids 1-736) and the locations of the associated variable regions and GBS and GH loop regions are defined in Table 2 below. The VP2 capsid protein spans amino acids 138-736 of SEQ ID NO:43 and the VP3 capsid protein spans amino acids 203-736 of SEQ ID NO:43.

MAADGYLPDWLEDNLSEGIREWDLKPGAPKPKANQQKQDDGRGLVLPGYKYLGPFGNLDKGEPVNAADAAALEHDKAYDQQLKAG
 DNPYLRYNHADADEFQERLQEDTSFGGNLGRAVFQAKKRVLEPLGLVEEGAKTAPGKKRPVEQSPQEPDSSSGIGKTGQQPAKKRLN
 FGQTGDSESVDPDPQLGEPPAAPSGLGPNTMASGGGAPMADNNEGADGVGNSSGNWHCDSTWLGDRVITTTSTRTWALPTYNNHLYK
 QISNSTSGGSTNDNTYFGYSTPWGYFDNRFHCHFSPRDWQRLINNNWGFPRKRLNFKLFNIQVKEVTTNEGKTIANNLTSTVQV
 FTDSEYQLPYVLGSAHQGLPPFPADVFMVPQYGHLLTLNNGSQALGRSSFYCLEYFPSQMLRTGNNFQFSYTFEDVPFHSSYAHSQ
 SLDRMLNPLIDQYLYLVRTQTTGTGGTQTLAFSQAGPSSMANQARNWVPGPSYRQQRVSTTKDQNNNSNFAWTGAAKFKLNGRNS
 LMNPGVAMASHKDDERFFPSSGVLIIFGKQAGNDGVDSQVLI TDEEEIKATNPVATEEYGEVAVNDQAANTQAQTGLVHNQGV
 PGMVWQNRDVYLQGPWAKIPHTDGNFHPSPMLGGFGLKHPPPQILIKNTVPADPPLTLNQAKLNSFITQYSTGQVSVEIEWELQ
 KENSKRWNPETQYTSNYYKSTNVDFAVNSDGVYSEPRPIGTRYLTRNL

[00125] The VP1 sequence of a novel AAV capsid isolated from crab-eating macaque (denoted as Bce.43) is set out as SEQ ID NO:44 (amino acids 1-730) and the locations of the associated variable regions and GBS and GH loop regions are defined in Table 2 below. The VP2 capsid protein spans amino acids 138-730 of SEQ ID NO:44 and the VP3 capsid protein spans amino acids 199-730 of SEQ ID NO:44.

MAADGYLPDWLEDNLSEGIREWDLKPGAPQPKANQQKQDDGRGLVLPGYKYLGPFGNLDKGEPVNEADAAALEHDKAYDQQLKAG
 DNPYLRYNHADADEFQERLQEDTSFGGNLGRAVFQAKKRVLEPLGLVEEAAKTAPGKKRPVDSPDSTSGIGKKGQQPARKRLNFGQT
 GDAESVPDPQPIGEPPAAPSGLGSMTMAAGGGAPMADNNEGADGVGNAPGNWHCDSTWLGNRVITTTSTRTWALPTYNNHLYKQISS
 SSSGATNDNHYFGYSTPWGYFDNRFHCHFSPRDWPRLINNNWGFPRKRLRFKLFNIQVKEVTTNDGVTIANNLTSTVQVFSQDSE
 YQLPYVLGSAHQGLPPFPADVFMIPQYGYLTLNNGSQSVGRSSFYCLEYFPSQMLRTGNNFESYEFEDVPFHSSYAHSQSLDR

MNPLIDQYPYLLARTQSSTGTSTRELQFHQAGPATMAEQSKNWLPGPCFRQQRISKTTDNNNSNFAWTGATKYHLNGRNSLTNPG
 VPMATHKDDSVFFPINGVLVFGKTGASNKTTLENVMTDEEEIKATNPVATEEYGVVSSNIQSQNSNPATQTVNNQGALPGMVWQ
 NRDVYLQGPWAKIPHTDGNFHPSPMLGGFGLKHPPPQILIKNTVPANPPETFTPAKFASFITQYSTGQVSVEIEWELQKENS
 KRWNPEIQYTSNYDKQTGVDFAVDTQGVYSEPRPIGTRYLTRNL

[00126] The VP1 sequence of a novel AAV capsid isolated from crab-eating macaque (denoted as Bce.44) is set out as SEQ ID NO:45 (amino acids 1-730) and the locations of the associated variable regions and GBS and GH loop regions are defined in Table 2 below. The VP2 capsid protein spans amino acids 138-730 of SEQ ID NO:45 and the VP3 capsid protein spans amino acids 199-730 of SEQ ID NO:45.

MAADGYLPDWLEDNLSEGIREWALKPGAPQPKANQQKQDDGRGLMLPGYKYLGPFGNLDKGEPVNEADAAALEHDKAYDQQLKAG
 DNPYLRYNHADADEFQERLQEDTSFGGNLGRAVFQAKKRVLEPLGLVEEAAKTAPGKKRPVDSPTSGIGKKGQQPARKRLNFGQT
 GDAESVPDPQPIGEPPAAPSGLGSGTMAAGGGAPMADNNEGADGVGNASGSHWCDSTWLGNRVITTTSTRTWALPTYNNHLYKQISS
 SSSGATNDNHYFGYSTPWGYFDNRFHCHFSPRDWQRLINNNWGFPRKRLRFKLFNIQVKEVTTNDGVTTIANNLTSTVQVFSDE
 YQLPYVLGSAHQGCLPPFPADVFMIQYGYLTLNNGRQSVGRSSFYCLEYFPSQMLRTGNNFEFSYEFEDVPFHSSYAHSQSLDRL
 MNPLIDQYPYLLARTQSSTGTSTRELQFHQAGPATMAEQSKNWLPGPCFRQQRISKTTDNNNSNFAWTGATKYHLNGRNSLTNPG
 VPMATHKDDSVFFPINGVLVFGKTGASNKTTLENVMTDEEEIKATNPVATEEYGVVSSNIQSQNSNPATQTVNNQGALPGMVWQ
 NRDVYLQGPWAKIPHTDGNFHPSPMLGGFGLKHPPPQILIKNTVPANPPETFTPAKFASFITQYSTGQVSVEIEWELQKENS
 KRWNPEIQYTSNYDKQTGVDFAVDTQGVYSEPRPIGTRYLTRNL

[00127] The VP1 sequence of a novel AAV capsid isolated from crab-eating macaque (denoted as Bce.45) is set out as SEQ ID NO:46 (amino acids 1-730) and the locations of the associated variable regions and GBS and GH loop regions are defined in Table 2 below. The VP2 capsid protein spans amino acids 138-730 of SEQ ID NO:46 and the VP3 capsid protein spans amino acids 199-730 of SEQ ID NO:46.

MAADGYLPDWLEDNLSEGIREWALKPGAPQPKANQQKQDDGRGLVLPGYKYLGPFGSLDKGEPVNEADAAALEHDKAYDQQLKAG
 DNPYLRYNHADADEFQERLQEDTSFGGNLGRAVFQAKKRVLEPLGLVEEAAKTAPGKKRPVDSPTSGIGKKGQQPARKRLNFGQT
 GDAESVPDPQPIGEPPAAPSGLGSGTMAAGGGAPMADNNEGADGVGNASGNWCDSTWLGNRVITTTSTRTWALPTYNDHLYKQISS
 SSSGATNDNHYFGYSTPWGYFDNRFHCHFSPRDWQRLINNNWGFPRKRLRFKLFNIQVKEVTTNDGVTTIANNLTSTVQVFSDE
 YQLPYVLGSAHQGCLPPFPADVFMIQYGYLTLNNGSQSVGRSSFYCLEYFPSQMLRTGNNFEFSYEFEDVPFHSSYAHSQSLDRL
 MNPLIDQYLYLLARTQSSTGTSTRELQFHQAGPATMAEQSKNWLPGPCFRQQRISKTTDNNNSNFAWTGATKYHLNGRNSLTNPG
 VPMATHKDDSVFFPINGVLVFGKTGASNKTTLENVMTDEEEIKATNPVATEEYGVVSSNIQSQNSNPATQTVNNQGALPGMVWR
 NRDVYLQGPWAKIPHTDGNFHPSPMLGGFGLKHPPPQILIKNTVPANPPETFTPAKFASFITQYSTGQVSVEIEWELQKENS
 KRWNPEIQYTSNYDKQTGVDFAVDTQGVYSEPRPIGTRYLTRNL

[00128] The VP1 sequence of a novel AAV capsid isolated from crab-eating macaque (denoted as Bce.46) is set out as SEQ ID NO:47 (amino acids 1-730) and the locations of the associated variable regions and GBS and GH loop regions are defined in Table 2 below. The VP2 capsid protein spans amino acids 138-730 of SEQ ID NO:47 and the VP3 capsid protein spans amino acids 199-730 of SEQ ID NO:47.

MAADGYLPDWLEDNLSEGIREWALKPGAPQPKANQQKQDDGRGLVLPGYKYLGPFGNLDKGEPVNEADAAALEHDKAYDQQLKAG
 DNPYLRYNHADADEFQERLQEDTSFGGNLGRAVFQAKKRVLEPLGLVEEAAKTAPGKKRPVDSPDSTSGIGKKGQQPARKRLNFGQT
 GDAESVPDPQPIGEPPAAPSGLGSGTMAAGGGAPMADNNEGADGVGNASGNWHCDSTWLGNRVITTTSTRTWALPTYNNHLYKQISS
 SSSGATNDNHYFGYSTPWGYFDNRFHCHFSPRDWQRLINNNWGFPRKRLRFKLFNIQVKEVTTNDGVTTIANNLSTVQVFSDE
 YQLPYVLGSAHQGCLPPFPADVFMIQYGYLTLNNGSQSVGRSSFYCLEYFPSQMLRTGNNFEFSYEFEDVPFHSSYAHSQSLDRL
 MNPLIDQYLYLMARTQSSTGTSTREPQFHQAGPATMAEQSKNWLPGPCFRQQRISKTDDNNNNSNFAWTGATKYHLNGRNSLTNPG
 VPMATHKDDSVFFPINGVLVFGKTGASNKTTLENVLMTDEEEIKATNPVATEEYGVVSSNIQSQNSNPTTQTVNNQGALPGMVWQ
 NRDVYLQGPWAKIPHTDGNFHPSPMLMGGFGLKHPPQILIKNTPVPANPPETFTPAKFASFITQYSTGQVSVEIEWELQKENSKR
 WNPEIQYTSNYDKQTGVDFAVDTQGVYSEPRPIGTRYLTRNL

[00129] The VP1 sequence of a novel AAV capsid isolated from cynomolgus macaque (denoted as Bcy.20) is set out as SEQ ID NO:48 (amino acids 1-730) and the locations of the associated variable regions and GBS and GH loop regions are defined in Table 2 below. The VP2 capsid protein spans amino acids 138-730 of SEQ ID NO:48 and the VP3 capsid protein spans amino acids 199-730 of SEQ ID NO:48.

MAADGYLPDWLEDNLSEGIREWALKPGAPQPKANQQKQDDGRGLVLPGYKYLGPFGNLDKGEPVNEADAAALEHDKAYDQQLKAG
 DNPYLRYNHADADEFQERLQEDTSFGGNLGRAVFQAKKRVLEPLGLVEEAAKTAPGKKRPVDSPDSTSGIGKKGQQPARKRLNFGQT
 GDAESVPDPQPIGEPPAAPSGLGSGTMAAGGGAPMADNNEGADGVGNASGNWHCDSTWLGNRVITTTSTRTWALPTYNNHLYKQISS
 GSSGATNDNHYFGYSTPWGYFDNRFHCHFSPRDWQRLINNNWGFPRKRLRFKLFNIQVKEVTTNDGVTTIANNLSTVQVFSDE
 YQLPYVLGSAHQGCLPPFPADVFMIQYGYLTLNNGSQSVGRSSFYCLEYFPSQVLRRTGNNFEFSYEFEDVPFHSSYAHSQSLDRL
 MNPLIDQYLYLMARTQSSTGTSTRELQFHQAGPATMAEQSKNWLPGPCFRQQRISKTSDNNNNSNFAWTGATKYHLNGRNSLTNPG
 VPMATHKDDSVFFPINGVLVFGKTGASNKTTLENVLMTDEEEIKATNPVATEEYGVVSSNIQSQNSNPTTQTVNNQGALPGMVWQ
 NRDVYLQGPWAKIPHTDGNFHPSPMLMGGFGLKHPPQILIKNTPVPANPPETFTPAKFASFITQYSTGQVSVEIEWELQKENSKR
 WNPEIQYTSNYDKQTGVDFAVDTQGVYSEPRPIGTRYLTRNL

[00130] The VP1 sequence of a novel AAV capsid isolated from cynomolgus macaque (denoted as Bcy.22) is set out as SEQ ID NO:49 (amino acids 1-730) and the locations of the associated variable regions and GBS and GH loop regions are defined in Table 2 below. The VP2 capsid protein spans amino acids 138-730 of SEQ ID NO:49 and the VP3 capsid protein spans amino acids 199-730 of SEQ ID NO:49.

MAADGYLPDWLEDNLSEGIREWALKPGAPQPKANQQKQDDGRGLVLPGYRYLGPFGNLDKGEPVNEADAAALEHDKAYDQQLKAG
 DNPYLRYNHADADEFQERLQEDTSFGGNLGRAVFQAKKRILEPLGLVEEAAKTAPGKERPVDSPDSTSGIGKKGQQPARKRLNFGQT
 GDAESVPDPQPIGEPPAAPSGLGSGTMAAGGGAPMADNNEGADGVGNASGNWHCDSTWLGNRVITTTSTRTWALPTYDNHLYKQISS
 SSSGATNDNHYFGYSTPWGYFDNRFHCHFSPRDWQRLINNNWGFPRKRLRFKLFNIQVKEVTTNDGVTTIANNLSTVQVFSDE
 YQLPYVLGSAHQGCLPPFPADVFMIQYGYLTLNNGSQSVGRSSFYCLEYFPSQMLRTGNNFEFSYEFEDVPFHSSYAHSQSLDRL
 MNPLIDQYLYLMARTQSSTGTSTRELQFHQAGPATMAEQSKNWLPGPCFRQQRISKTDDNNNNSNFAWTGATKYHLNGRNSLTNPG
 VPMATHKDDSVFFPINGVLVFGKTGASNKTTLENVLMTDEEEIKATNPVATEEYGVVSSNIQSQNSNPTTQTVNNQGALPGMVWQ
 NRDVYLQGPWAKIPHTDGNFHPSPMLMGGFGLKHPPQILIKNTPVPANPPETFTPAKFASFITQYSTGQVSVEIEWELQKENSKR
 WNPEIQYTSNYDKQTGVDFAVDTQGVYSEPRPIGTRYLTRNL

[00131] The VP1 sequence of a novel AAV capsid isolated from cynomolgus macaque (denoted as Bcy.23) is set out as SEQ ID NO:50 (amino acids 1-730) and the locations of the

associated variable regions and GBS and GH loop regions are defined in Table 2 below. The VP2 capsid protein spans amino acids 138-730 of SEQ ID NO:50 and the VP3 capsid protein spans amino acids 199-730 of SEQ ID NO:50.

MAADGYLPDWLEDNLSEGIREWALKPGAPRPKANQQKQDDGRGLVLPGYKYLGPFGNLDKGEPVNEADAAALEHDKAYDQQLKAG
 DNPYLRYNHADADEFQERLQEDTSFGGNLGRAVFQAKKRVLEPLGLVEEAAKTAPGKKRPVDSPDSTSGIGKKGQQPARKRLNFGQT
 GDAESVPDPQPIGEPPAAPSGLGSGTMAAGGGAPMADNNEGADGVGNASGNWHCDSTWLGNRVITTTSTRTWALPTYNNHLYKQISS
 SSSGATNDNHYFGYSTPWGYFDNRFHCHFSPRDWQRLINNNWGFPRKRLRFKLFNIQVKEVTTNDGVTTIANNLTSTVQVFSDE
 YQLPYVLGSAHQGCLPPFPADVFMIPQYGYLTLNNGSQSVGRSSFYRLEYFPSQMLRTGNNFEFSYEFEDVPFHSSYAHSQSLDRL
 MNPLIDQYLYYLARTQSSTGTSTRELQFHQAGPATMAEQSKNWLPGPCFRQQRISKTDDNNNSNFAWTGATKYHLNGRNSLTNPG
 VPMATHRDDSEVFFPINGVLVFGKTGASNKTTLENVLMTDEEEIKATNPVATEEYGVVSSNIQSQNSNPTTQTVNNQGALPGMVWQ
 NRDVYLQGGPIWAKIPHTDGNFHPSPMLGGFGLKHPPPQILIKNTPVPANPPETFTPAKFASFITQYSTGQVSVEIEWELQKENSKR
 WNPEIQYTSNYDKQTGVDFAVDTQGVYSEPRPIGTRYLTRNL

[00132] The VP1 sequence of a novel AAV capsid isolated from marmoset (denoted as Bma.42) is set out as SEQ ID NO:51 (amino acids 1-736) and the locations of the associated variable regions and GBS and GH loop regions are defined in Table 2 below. The VP2 capsid protein spans amino acids 138-736 of SEQ ID NO:51 and the VP3 capsid protein spans amino acids 203-736 of SEQ ID NO:51.

MAADGYLPDWLEDNLSEGIREWDLKPGAPKPKANQQKQDDGRGLVLPGYKYLGPFGNLDKGEPVNAADAAALEHDKAYDQQLKAG
 DNPYLRYNHADADEFQERLQEDTSFGGNLGRAVFQAKKRVLEPLGLVEEGAKTAPGKKRPVEQSPQEPDSSSGIGKTGQQPAKKRLN
 FGQTGDTESEVPDPQPLREPPAAPSGLGPNTMASGGGAPMADNNEGADGVGNSSGNWHCDSTWLGDRVITTTSTRTWALPTYNNHLYK
 QISNGTSGGSTNDNTYFGYSTPWGYFDNRFHCHFSPRDWQRLINNNWGFPRKRLNFKLPFNIQVKEVTTNEGKTIANNPTSTVQV
 FTDSEYQLPYVLGSAHQGCLPPFPADVFMVPQYGYLTLNNGSQALGRSSFYCLEYFPSQMLRTGDNFQFSYTFEDVPFHSSYAHSQ
 SLDRMLNPLIDQYLYLVLTQTGTGGTQTLAFSQAGPSSMANQARNWVPGPCYRQQRVSTTTNQNNSNFAWTGAAKFKLNGRDS
 LMNPGVAMASHKDEDRFFPSSGVLI FGKQAGNDGVDYSQVLTDEEEIKATNPVATEEYGAVAINNQAAANTLAQTGLVHNQGVI
 PGMVWQNRDVYLQGGPIWAKIPHTDGNFHPSPMLGGFGLKHPPPQILIKNTPVPADPPLTFNQAKLNSFITQYSTGQVSVEIEWELQ
 KENSKRWNPFIQYTSNYKSTNVDFAVNTEGVYSEPRPIGTRYLTRNL

[00133] The VP1 sequence of a novel AAV capsid isolated from marmoset (denoted as Bma.43) is set out as SEQ ID NO:52 (amino acids 1-736) and the locations of the associated variable regions and GBS and GH loop regions are defined in Table 2 below. The VP2 capsid protein spans amino acids 138-736 of SEQ ID NO:52 and the VP3 capsid protein spans amino acids 203-736 of SEQ ID NO:52.

MAADGYLPDWLEDNLSEGIREWDLKPGAPKPKANQQKQDDGRGLVLPGYKYLGPFGNLDKGEPVNAADAAALEHDKAYDQQLKAG
 DNPYLRYNHADADEFQERLQEDTSFGGNLGRAVFQAKKRVLEPLGLVEEGANTAPGKKRPVEQSPQEPDSSSGIGKTGQQPAKKRLN
 FGQTGDSESEVPDPQPLREPPAAPSGLGPNTMASGGGAPMADNNEGADGVGNSSGNWHCDSTWLGDRVITTTSTRTWALPTYNNHLYK
 QISNGTSGGSTNDNTYFGYSTPWGYFDNRFHCHFSPRDWQRLINSNWGFPRKRLNFKLFNIQVKEVTTNEGKTIANNLTSTVQV
 FTDSEYQLPYVLGSAHQGCLPPFPADVFMVPQYGYLTLNNGSQALGRSSFYCLEYFPSQMLRTGNNFQFSYTFEDVPFHSSYAHSQ
 SLDRMLNPLIDQYLYLVLTQTGTGGTQTLAFSQAGPSSMANQARNWVPGPCYRQQRVSTTTNQNNSNFAWTGAAKFKLNGRDS
 LMNPGVAMASHKDEDRFFPSSGVLI FGKQAGNDGVDYSQVLTDEEEIKATNPVATEEYGAVAINNQAAANTQAQTGLVHNQGVI

PGMWQNRDVYLQGPWQKIPHTDGNFHPSPMLGGFGLKHPPQILIKNTVPADPPLTFNQAKLNSFITQYSTGQVSVEIEWELQ
KENSKRWNPETQYTSNYYKSTNVDFAVNTEGVYSEPRPIGTRYLTRNL

[00134] The VP1 sequence of a novel AAV capsid isolated from pig (denoted as Bpo.1) is set out as SEQ ID NO:53 (amino acids 1-716) and the locations of the associated variable regions and GBS and GH loop regions are defined in Table 2 below. The VP2 capsid protein spans amino acids 137-716 of SEQ ID NO:53 and the VP3 capsid protein spans amino acids 184-716 of SEQ ID NO:53.

MSFVDHPPDWLEEIGEGLEKFLGLEPGPPKPKPNQQKQDNARGLVLPGYNYLGPNGGLDRGEPVNRADDEVAREHDI SYNEQLQAGD
NPYLKYNHADADEFQEKLADDTSGGNGLGKAVFQAKKRVLEPFGLVEEPAKTAAGKERIDDHYPKPKKARVEETEAGTSGGQQLQIP
AQPASSLGADTMSAGGGSPLGDNQAGDGVGNASGDWHCDSTWMGDRVITKSTRTWVLP SYNHHLYKEIHSGSVDGSNANAYFGYS
TPWGYDFNRFHSHWSRQRLVNNYWGFRPRSLKVKIFNIQVKEVTVDATTIANNLSTTVQVFTDDDYQLPYVINGTEGCL
PAFPQVFTLPQYGYATLNRNNTDDPTERSFFCLEYFPSKMLRTGNSFEFTYSFEVVPFHCSFAPSQNLFKLANPLVDQYLYRFV
STDTSGNIQFQKNLKARYANTYKNWFGPMCRTOGWYTGSGTYNRSVGTNFATSNRMDLEGASYQVNPQPNGMTNTLQDSNKYALE
NTMIFNSQNAEPGTTSLYQENLLITSESETQPVNRVAYDTGGQMATNAQSTNLAPTVGTYNHQMPLPGSVWMDRDVYLQGPWQKIP
IPETGAHFHSPAMGGFGLKHPPPMMLIKNTVPVSNVTAFSEIPVKSFITQYSTGQVTVEMEWELKKEDSKRWNPETQYTNNNNP
EFVDFAPDTSGEYRTRTIGTRYLTRPL

[00135] The VP1 sequence of a novel AAV capsid isolated from pig (denoted as Bpo.2) is set out as SEQ ID NO:54 (amino acids 1-716) and the locations of the associated variable regions and GBS and GH loop regions are defined in Table 2 below. The VP2 capsid protein spans amino acids 137-716 of SEQ ID NO:54 and the VP3 capsid protein spans amino acids 184-716 of SEQ ID NO:54.

MSFVDHPPDWLEEIGEGLEKFLGLEPGPPKPKPNQQKQDNARGLVLPGYNYLGPNGGLDRGEPVNRADDEVAREHDI SYNEQLQAGD
NPYLKYNHADADEFQEKLADDTSGGNGLGKAVFQAKKRVLEPFGLVEEPAKTAAGKERIDDHYPKPKKARVEETEAGTSGGQQLQIP
AQPASSLGADTMSAGGGSPLGDNQAGDGVGNASGDWHCDSTWMGDRVITKSTRTWVLP SYNHHLYKEIHSGSVDGSNANAYFGYS
TPWGYDFNRFHSHWSRQRLVNNYWGFRPRSLKVKIFNIQVKEVTVDATTIANNLSTTVQVFTDDDYQLPYVINGTEGCL
PAFPQVFTLPQYGYATLNRNNTDDPTERSFFCLEYFPSKMLRTGNSFEFTYSFEVVPFHCSFAPSQNLFKLANPLVDQYLYRFV
GTDTSNGNIQFQKNLKARYANTYKNWFGPMCRTOGWYTGSGTYNRSVGTNFATSNRMDLEGASYQVNPQPNGMTNTLQDSNKYALE
NTMIFNSQNAEPGTTSLYQENLLITSESETQPVNRVAYDTGGQMATNAQSTNLAPTVGTYNHQMPLPGSVWMDRDVYLQGPWQKIP
IPETGAHFHSPAMGGFGLKHPPPMMLIKNTVPVSNVTAFSEIPVKSFITQYSTGQVTVEMEWELKKEDSKRWNPETQYTNNNNP
EFVDFAPDTSGEYRTRTIGTRYLTRPL

[00136] The VP1 sequence of a novel AAV capsid isolated from pig (denoted as Bpo.3) is set out as SEQ ID NO:55 (amino acids 1-716) and the locations of the associated variable regions and GBS and GH loop regions are defined in Table 2 below. The VP2 capsid protein spans amino acids 137-716 of SEQ ID NO:55 and the VP3 capsid protein spans amino acids 184-716 of SEQ ID NO:55.

MSFVDHPPDWLEEIGEGLEKFLGLEPGPPKPKPNQQKQDNARGLVLPGYNYLGPNGGLDRGEPANRADEVAREHDI SYNEQLQAGD
NPYLKYNHADADEFQEKLADDTSGGNGLGKAVFQAKKRVLEPFGLVEEPAKTAAGKERIDDHYPKPKKARVEETEAGTSGGQQLQIP

AQPASSLGLADTMSAGGGSPLGDNNGADGVGNASGDWHCDSTWMGDRVITKSTRTWVLPSSYNNHLYKEIHSGSVSDGSNANAYFGYS
 TPWGYFDNRFHSHWSRPDWQRLVNYYWGFRRPSLKVKIFNIQVKEVTVDATTTIANNLSTVQVFTDDDYQLPYVIGNGTEGCL
 PAFPPQVFTLPQYGYATLNRNNTDDPTERSFFCLEYFPSKMLRTGNSFEFTYSFGEVPLHCSFAPSQNLFKLANPLVDQHLRYFV
 STDTSGNIQFQKNLKARYANTYKNWFPGPMCRTOGWYTGSGTYNRPVGTNFATSNRMDLEGASYQVNPQPNGMTNTLQDSNKYALE
 NTMIFNSQNAEPGTTSLYQENLLITSESETQPVNRVAYDTGGQMATNAQSTNLAPTGTYNHQMELPGSVWMDRDVYLQGPWAK
 IPETGAHFHPSPAMGGFGLKHPPMMLTKNTPVPSNVTAFSEIPVKSFITQYSTGQVTVEMEWELKKENSKRWNPFIQYTNNNNP
 EFVDFAPDTSGEYRTTRAIGTRYLTRPL

[00137] The VP1 sequence of a novel AAV capsid isolated from pig (denoted as Bpo.4) is set out as SEQ ID NO:56 (amino acids 1-716) and the locations of the associated variable regions and GBS and GH loop regions are defined in Table 2 below. The VP2 capsid protein spans amino acids 137-716 of SEQ ID NO:56 and the VP3 capsid protein spans amino acids 184-716 of SEQ ID NO:56.

MSFVDHPPDWLEEIGEGLEKFLGLEPGPPKPKPNQQQDNARGLVLPGYNYLGPNGLDRGEPVNRADDEVAREHDI SYNEQLQAGD
 NPYLKYNHADADEFQEKLADDTSGGNGLGKAVFQAKKRVLEPFGLEVEPAKTAAGERIDDHYPKKKKARVEETEAGTSGGQQLQIP
 AQPASSLGLADTMSAGGGSPLGDNNGADGVGNASGDWHCDSTWMGDRVITKSTRTWVLPSSYNNHLYKEIHSGSVSDGSNANAYFGYS
 TPWGYFDNRFHSHWSRPDWQRLVNYYWGFRRPSLKVKIFNIQVKEVTVDATTTIANNLSTVQVFTDDDYQLPYVIGNGTEGCL
 PAFPPQVFTLPQYGYATLNRNNTDDPTERSFFCLEYFPSKMLRTGNSFEFTYSFEEVFPFHCSFAPSQNLFKLANPLVDQYLYRFV
 STDTSGNIQFQKNLKARYANTYKNWFPGPMCRTOGWYTGSGTYNRSVGTNFATSNRMDLEGASYQVNPQPNGMTNTLQDSNKYALE
 NTMIFNSQNAEPGTTSLYQENLLITSESETQPVNRVAYDTGGQMATNAQSTNLAPTGTYNHQMELPGSVWMDRDVYLQGPWAK
 IPETGAHFHPSPAMGGFGLKHPPMMLIKNTPVPSNVTAFSEIPVKSFITQYSTGQVTVEMEWELKKENSKRWNPFIQYTNNNNP
 EFVDFAPDTSGEYRTTRAIGTRYLTRPL

[00138] The VP1 sequence of a novel AAV capsid isolated from pig (denoted as Bpo.6) is set out as SEQ ID NO:57 (amino acids 1-716) and the locations of the associated variable regions and GBS and GH loop regions are defined in Table 2 below. The VP2 capsid protein spans amino acids 137-716 of SEQ ID NO:57 and the VP3 capsid protein spans amino acids 184-716 of SEQ ID NO:57.

MSFVDHPPDWLEEIGEGLEKFLGLEPGPPKPKPNQQQDNARGLVLPGYNYLGPNGLDRGEPVNRADDEVAREHGI SYNEQLQAGD
 DPYLKYNHADADEFQEKLADDTSGGNGLGKAVFQAKKRVLEPFGLEVEPAKTAAGERIDDHYPKKKKARVEETEAGTGGGQQLQIP
 AQPASSLGLADTMSAGGGSPLGDNNGADGVGNASGDWHCDSTWMGDRVITKSTRTWVLPSSYNNHLYKEIHSGSVSDGSNANAYFGYS
 TPWGYFDNRFHSHWSRPDWQRLVNYYWGFRRPSLKVKIFNIQVKEVTVDATTTIANNLSTVQVFTDDDYQLPYVIGNGTEGCL
 PAFPPQVFTLPQYGYATLNRNNTDDPTERSFFCLEYFPSKMLRTGNSLEFTYSFEEVFPFHCSFAPSQNLFKLANPLVDQYLYRFV
 STDTSGNIQFQKNLKARYANTYKNWFPGPMCRTOGWYTGSGTYNRSVGTNFATSNRMDLEGASYQVNPQPNGMTNTLQDSNKYALE
 NTMIFNSQNAEPGTTSLYQENLLITSESETQPVNRVAYDTGGQMATNAQSTNLAPTGTYNHQMELPGSVWMDRDVYLQGPWAK
 IPETGAHFHPSPAMGGFGLKHPPMMLIKNTPVPSNVTAFSEIPVKSFITQYSTGQVTVEMEWELKKENSKRWNPFIQYTNNNNP
 EFVDFAPDTSGEYRTTRAIGTRYLTRPL

[00139] The VP1 sequence of a novel AAV capsid isolated from pig (denoted as Bpo.8) is set out as SEQ ID NO:58 (amino acids 1-716) and the locations of the associated variable regions and GBS and GH loop regions are defined in Table 2 below. The VP2 capsid protein spans

amino acids 137-716 of SEQ ID NO:58 and the VP3 capsid protein spans amino acids 184-716 of SEQ ID NO:58.

MSFVDHPPDWLEEIGEGLKEFLGLKPGPPKPKPNQQKQDNARGLVLPGYNYLGPNGGLGRGEPVNRADDEVAREHDI SYNEQLQAGD
NPYLKYNHADADEFQEKLADDTSGGNGLGKAVFQAKKRVLEPFGLEVEEPAKTAAGERIDDHYPKPKKKARVEETEAGTSGGQQLQIP
AQPASSLGADTMSAGGGSPLGDNQAGDGVGNASGDWHCDSTWMGDRVITKSTRTWVLP SYNNHLYKEIHSGSVDSNANAYFGYS
TPWGYFDNRFHSHWSPRDWQRLVNYYWGFPRPSLKVKIFNIQVKEVTVDATTTIANNLSTVQVFTDDDYQLPYVINGTEGCL
PAFPPQVFTLPQYGYATLNRNNTDDPTERSFFCLEYFPPSKMLRTGNSFEFTYSFEEVFPFHCSFAPSQNLFKLANPLVDQYLYRFV
STDTSGNIQFQKNLKARYANTYKNWFPGPMCRTOGWYTGSGTYNRSVGTNFATSNRMDLEGASYQANPQPNGMTNTLQDSNKYALE
NTMIFNSQNAEPGTTSLYQENLLITSESETQPVNRVAYDTGGQMATNAQSTNLAPTVGTYNHQMELPGSVWMDRVDVYLQGP I WAK
IPETGAHFHPSPAMGGFGLKHPPPMMLIKNTPVPSNVTAFSEIPVKS LITQYSTGQVTVEMEWELKKENSKRWNP EIQYTNNYNNP
EFVDFAPDTSGEYRTTRAIGTRYLTRPL

[00140] The VP1 sequence of a novel AAV capsid isolated from pig (denoted as Bpo.13) is set out as SEQ ID NO:59 (amino acids 1-716) and the locations of the associated variable regions and GBS and GH loop regions are defined in Table 2 below. The VP2 capsid protein spans amino acids 137-716 of SEQ ID NO:59 and the VP3 capsid protein spans amino acids 184-716 of SEQ ID NO:59.

MSFVDHPPDWLEEIGEGLKEFLGLEP GPPKPKPNQQKQDNARGLVLPGYNYLGPNGGLDRGEPVNRADDEVAREHDI SYNEQLQAGD
NPYLKYNHADADEFQEKLADDTSGGNGLGKAVFQAKKRVLEPFGLEVEEPAKTAAGERIDDHYPKPKKKARVEETEAGTSGGQQLQIP
AQPASSLGADTMSAGGGSPLGDNQAGDGVGNASGDWHCDSTWMGDRVITKSTRTWVLP SYNNHLYKEIHSGSVDSNANAYFGYS
TPWGYFDNRFHSHWSPRDWQRLVNYYWGFPRPSLKVKIFNIQVKEVTVDATTTIANNLASTVQVFTDDDYQLPYVINGTEGCL
PAFPPQVFTLPQYGYATLNRNNTDDPTERSFFCLEYFPPSKMLRTGNSFEFTYSFEEVFPFHCSFAPSQNLFKLANPLVDQYLYRFV
STDTSGNIQFQKNLKARYANTYKNWFPGPMCRTOGWYTGSGTYNRSVGTNFATSNRMDLEGASYQVNPQPNGMTNTLQDSNKYALE
NTMIFNSQNAEPGTTSLYQENLLITSESETQPVNRVAYDTGGQMATNAQSTNLAPTVGTYNHQMELPGSVWMDRVDVYLQGP I WAK
IPETGAHFHPSPAMGGFGLKHPPPMMLIKNTPVPSNVTAFSEIPVKS FITQYSTGQVTVEMEWELKKENSKRWNP EIQYTNNYNNP
EFVDFAPDTSGEYRTTRAIGTRYLTRPL

[00141] The VP1 sequence of a novel AAV capsid isolated from pig (denoted as Bpo.18) is set out as SEQ ID NO:60 (amino acids 1-716) and the locations of the associated variable regions and GBS and GH loop regions are defined in Table 2 below. The VP2 capsid protein spans amino acids 137-716 of SEQ ID NO:60 and the VP3 capsid protein spans amino acids 184-716 of SEQ ID NO:60.

MSFVDHPPDWLEEIGEGLKEFLGLEP GPPKPKPNQQKQDNARGLVLPGYNYLGPNGGLDRGEPVNRADDEVAREHDI SYNEQLQAGD
DPYLKYNHADADEFQEKLADDTSGGNGLGKAVFQAKKRVLEPFGLEVEEPAKTAAGERIDDHYPKPKKKARVEETEAGTSGGQQLQIP
AQPASSLGADTMSAGGGSPLGDNQAGDGVGNASGDWHCDSTWMGDRVITKSTRTWVLP SYNNHLYKEIHSGSVDSNANAYFGYS
TPWGYFDNRFHSHWSPRDWQRLVNYYWGFPRPSLKVKIFNIQVKEVTVDATTTIANNLSTVQVFTDDDYQLPYVINGTEGCL
PAFPPQVFTLPQYGYATLNRNNTDDPTERSFFCLEYFPPSKMLRTGNSFEFTYSFEEVFPFHCSFAPSQNLFKLANPLVDQYLYRFV
STDTSGNIQFQKNLKARYANTYKNWFPGPMRRTQGWYTGSGTYNRSVGTNFATSNRMDLEGASYQVNPQPNGMTNTLQDSNKYALE
NTMIFNSQNAEPGTTSLYQENLLITSESETQPVNRVAYDTGGQMATNAQSTNLAPTVGTYNHQMELPGSVWMDRVDVYLQGP I WAK
IPETGAHFHPSPAMGGFGLKHPPPMMLIKNTPVPSNVTAFSEIPVKS FITQYSTGQVTVEMEWELKKENSKRWNP EIQYTNNYNNP
EFVDFAPDTSGEYRTTRAIGTRYLTRPL

[00142] The VP1 sequence of a novel AAV capsid isolated from pig (denoted as Bpo.20) is set out as SEQ ID NO:61 (amino acids 1-716) and the locations of the associated variable regions and GBS and GH loop regions are defined in Table 2 below. The VP2 capsid protein spans amino acids 137-716 of SEQ ID NO:61 and the VP3 capsid protein spans amino acids 184-716 of SEQ ID NO:61.

```
MSFVDHPPDWLEEIGEGLEKFLGLEPGPPKPKPNQQKDNRGLVLPGYNYLGPNGGLDRGEPVNRADDEVAREHDI SYNEQLQAGD
NPYLKYNHADA EFQEK LADDT SFGGNLGKAVFQAKKRVLEPFGLVEEPAKTA AKGERID DHYPK KKKARVEETEAGTSGGQQLQIP
AQPASSLGADTMSAGGGSPLGDNNGADGVGNASGDWHCDSTWMGDRVITKSTRTWVLP SYNNHLYKEI HSGSVDGSNANAYFGYS
TPWGYFDNRFHSHWGPRDWQRLVN NYWGF RPRSLKVKIFNIQVKEVTVDATTTIANNLTSTVQVFTDDDDYQLPYVINGTEGCL
PAFPQVFTLPQYGYATLNRNNTDDP TERS SFFCLEYF P SKMLRTGNSFEFTYSFEGVPFHCSFAPSQNLFKLANPLVDQYLYRFV
STDTSGNIQFQKNLKARYANTYKNWFP GPMCR TQGWYT GSGTYNRSGVTNFATSNRMDLEGASYQVNPQPNGMTNTLQDSNKYALE
NTMIFNSQNAEPGTTSLYQENLLITSESETQPVNRVAYDTGGQMATNAQSTNLAPTVGTYNH QEMLPGSVWMDRVDVYLQGP I WAK
IPETGAHFHPSPAMGGFGLKHPPPMMLIKNTPVPSNVTA FSEIPVKSFITQYSTGQVT VEMEWELKKENSKRWNP EIQYTNNYNNP
EFVDFAPDTSGEYRTTRAIGTRYLTRPL
```

[00143] The VP1 sequence of a novel AAV capsid isolated from pig (denoted as Bpo.23) is set out as SEQ ID NO:62 (amino acids 1-716) and the locations of the associated variable regions and GBS and GH loop regions are defined in Table 2 below. The VP2 capsid protein spans amino acids 137-716 of SEQ ID NO:62 and the VP3 capsid protein spans amino acids 184-716 of SEQ ID NO:62.

```
MSFVDHPPDWLEEIGEGLEKFLGLEPGPPKPKPNQQKRDNRGLVLPGYNYLGPNGGLDRGEPVNRVDEVARERDI SYNEQLQAGD
NPYLKYNHADA EFQEK LADDT SFGGNLGKAVFQAKKRVLEPFGLVEEPAKTA AKGERID DHYPK KKKARVEETEAGTSGGQQLQIP
AQPASSLGADTMSAGGGSPLGDNNGADGVGNASGDWHCDSTWMGDRVITKSTRTWVLP SYNNHLYKEI HSGSVDGSNANAYFGYS
TPWGYFDNRFHSHWSPRDWQRLVN NYWGF RPRSLKVKIFNIQVKEVTVDATTTIANNLTSTVQVFTDDDDYQLPYVINGTEGCL
PAFPQLQVFTLPQYGYATLNRNNTDDP TERS SFFCLEYF P SKMLRTGNSFEFTYSFEVVPFHCSFAPSQNLFKLANPLVDQYLYRFV
STDTSGNIQFQKNLKARYANTYKNWFP GPMCR TQGWYT GSGTYNRSGVTNFATSNRMDLEGASYQVNPQPNGMTNTLQDSNKYALE
NTMIFNSQNAEPGTTSLYQENLLITSESETQPVNRVAYDTGGQMATNAQSTNLAPTVGTYNH QEMLPGSVWMDRVDVYLQGP I WAK
IPETGAHFHPSPAMGGFGLKHPPPMMLIKNTPVPSNVTA FSEIPVKSFITQYSTGQVT VEMEWELKKENSKRWNP EIQYTNNYNNP
EFVDFAPDTSGEYRTTRAIGTRYLTRPL
```

[00144] The VP1 sequence of a novel AAV capsid isolated from pig (denoted as Bpo.24) is set out as SEQ ID NO:63 (amino acids 1-716) and the locations of the associated variable regions and GBS and GH loop regions are defined in Table 2 below. The VP2 capsid protein spans amino acids 137-716 of SEQ ID NO:63 and the VP3 capsid protein spans amino acids 184-716 of SEQ ID NO:63.

```
MSFVDHPPDWLEEIGEGLEKFLGLEPGPPKPKPNQQKDNRGLVLPGYNYLGPNGGLDRGEPVNRADDEVAREHDI SYNEQLQAGD
NPYLKYNHADA EFQEK LADDT SFGGNLGKAVFQAKKRVLEPFGLVEEPAKTA AKGERID DHYPK KKKARVEETEAGTSGGQQLQIP
AQPASSLGADTMSAGGGSPLGDNNGADGVGNASGDWHCDSTWMGDRVITKSTRTWVLP SYNNHLYKEI HSGSVDGSNANAYFGYS
TPWGYFDNRFHSHWSPRDWQRLVN NYWGF RPRSLKVKIFNIQVKEVTVDATTTIANNLTSTVQVFTDDDDYQLPYVINGTEGCL
PAFPQVFTLPQYGYATLNRNNTDDP TERS SFFCLEYF P SKMLRTGNSFEFTYSFEVVPFHCSFAPSQNLFKLANPLVDQYLYRFV
```

STDTSIGNIQFQKNLKARYANTYKNWFPGPMCRTOGWYTGSGTYNRSVGNFATSNRMDLEGASYQVNPQPNGMTNTLQDSNKYALE
 NTMIFNSQNAEPGTSTLYQENLLITSESETQPVNRVAYDTGGQMATNAQSTNLAPTGTYNHQMELPGSVWMDRDVYLQGGPIWAK
 IPETGAHFHPSAMGGFGLKHPPPMMLIKNTPVPSNVTAFSEIPVKSFITQYSTGQVTVEMEWELKKENSKRWNPFIQYTNNNYNNP
 EFVDFAPDTSGEYRTRTTRIGTRYLTRPL

[00145] The VP1 sequence of a novel AAV capsid isolated from pig (denoted as Bpo.27) is set out as SEQ ID NO:64 (amino acids 1-716) and the locations of the associated variable regions and GBS and GH loop regions are defined in Table 2 below. The VP2 capsid protein spans amino acids 137-716 of SEQ ID NO:64 and the VP3 capsid protein spans amino acids 184-716 of SEQ ID NO:64.

MSFVDHPPDWLEEIGEGLKEFLGLEPGPPKPKPNQQKQDNARGLVLPGYNYLGPNGLDGRGEPVNRADDEVAREHDI SYNEQLQAGD
 NPYLKYNHADAEEFQEKLADDTSGGNGLGKAVFQAKKRVLEPFGLVEEPAKTAAGERIDDHYPKKKKARVEETEAGTSGGQQLQIP
 AQPASSLGADTMSAGGGSPLGDNNGADGVGNASGDWHCDSTWMGDRVITKSTRTWVLPSSYNNHLYKEIHSGSVDGSNANAYFGYS
 TPWGYFDNRFHSHWSRWDQRLVDNYWGRPRSLKVKIFNIQVKEVTVDATTTIANNLSTTVQVFTDDDYQLPYVIGNGTEGCL
 PAFPPQVFTLPQYGYATLNRNNTDDPTERSFFCLEYFPSKMLRTGNSFEFTYSFEEVFPFHCSFAPSQNLFKLANPLVDQYLYRFV
 STDTSIGNIQFQKNLKARYANTYKNWFPGPMCRTOGWYTGSGTYNRSVGNFATSNRMDLEGASYQVNPQPNGMTNTLQDSNKYALE
 NTMIFNSQNAEPGTSTLYQENLLITSESETQPVNRVAYNTGGQMATNAQSTNLAPTGTYNHQMELPGSVWMDRDVYLQGGPIWAK
 IPETGAHFHPSAMGGFGLKHPPPMMLIKNTPVPSNVTAFSEIPVKSFITQYSTGQVTVEMEWELKKENSKRWNPFIQYTNNNYNNP
 EFVDFAPDTSGEYRTRTTRIGTRYLTRPL

[00146] The VP1 sequence of a novel AAV capsid isolated from pig (denoted as Bpo.28) is set out as SEQ ID NO:65 (amino acids 1-716) and the locations of the associated variable regions and GBS and GH loop regions are defined in Table 2 below. The VP2 capsid protein spans amino acids 137-716 of SEQ ID NO:65 and the VP3 capsid protein spans amino acids 184-716 of SEQ ID NO:65.

MSFVDHPPDWLEEIGEGLKEFLGLEPGPPKPKPNQQKRDNARGLVLPGYNYLGPNGLDGRGEPVNRADDEVAREHDI SYNEQLQAGD
 NPYLKYNHADAEEFQEKLADDTSGGNGLGKAVFQAKKRVLEPFGLVEEPAKTAAGERIDDHYPKKKKARVEETEAGTSGGQQLQIP
 AQPASSLGADTMSAGGGSPLGDNNGADGVGNASGDWHCDSTWMGDRVITKSTRTWVLPSSYNNHLYKEIHSGSVDGSNANAYFGYS
 TPWGYFDNRFHSHWSRWDQRLVNNYWGRLRPSLKVIFNIQVKEVTVDATTTIANNLSTTVQVFTDDDYQLPYVIGNGTEGCL
 PAFPPQVFTLPQYGYATLNRNNTDDPTERSFFCLEYFPSKMLRTGNSFEFTYSFEEVFPFHCSFAPSQNLFKLANPLADQYLYRFV
 STDTSIGNIQFQKNLKARYANTYKNWFPGPMCRTOGWYTGSGTYNRSVGNFATSNRMDLEGASYQVNPQPNGMTNTLQDSNKYALE
 NTMISNSQNAEPGTSTLYRENLLITSESETQPVNRVAYDTGGQMATNAQSTNLAPTGTYNHQMELPGSVWMDRDVYLQGGPIWAK
 IPETGAHFHPSAMGGFGLKHPPPMMLIKNTPVPSNVTAFSEIPVKSFITQYSTGQVTVEMEWELKKENSKRWNPFIQYTNNNYNNP
 EFVDFAPDTSGEYRTRTTRIGTRYLTRPL

[00147] The VP1 sequence of a novel AAV capsid isolated from pig (denoted as Bpo.29) is set out as SEQ ID NO:66 (amino acids 1-716) and the locations of the associated variable regions and GBS and GH loop regions are defined in Table 2 below. The VP2 capsid protein spans amino acids 137-716 of SEQ ID NO:66 and the VP3 capsid protein spans amino acids 184-716 of SEQ ID NO:66.

MSFVDHPPDWLEEIGEGLEKFLGLEPGPPKPKPNQQKDNRGLVLPGYNYLGPNGGLDRGEPVNRADDEVAREHDI SYNEQLQAGD
 NPYLKYNHADADEFQEKLADDTSFGGNLGKAVFQAKKRVLEPFGLVEEPAKTAAGERIDDHYPKKKKARVEETEAGTSGGQQLQIP
 AQPASSLGADTMSAGGGSPLGDNQAGDGVGNASGDWHCDSTWMGDRVITKSTRTWVLP SYN NHLYKEIHSGSVDGSNANAYFGYS
 TPWGYFDNRFHSHWSPRDWQRLVN NYWGF RPRSLKVKIFNIQVKEVTVDATTTIANNLSTVQVFTDDDYQLPYVIGNGTEGCL
 PAFPPQVFTLPQYGYATLNRNNTDDPTERSFFCLEYFPSKMLRTGNSFEFTYSFEVVPFHCSFAPSQNLFKLANPLVDQYLYRFV
 STDTSGNIQFQKNLKARYANTYKNWFGPMCR TQGWYTGSGTYNRSGVTNFATSNRMDLEGASYQVNPQPNGMTNTLQDSNKYALE
 NTMIFNSQNAEPGTTSLYQENLLITSESGTQPVNRVAYDTGGQMATNAQSTNLAPTVGTYNHQEMLPGSVWMDRGVYLQGP IWAK
 IPETGAHFHPSPAMGGFGLKHPPPMMLIKNTPVPSNVTA FSEIPVKSFITQYSTGQVTVEMEWELKKENSKRWNP EIQYTNNYNNP
 EFVDFAPDTSGEYRTTRAIGTRYLTRPL

[00148] The VP1 sequence of a novel AAV capsid isolated from pig (denoted as Bpo.33) is set out as SEQ ID NO:67 (amino acids 1-716) and the locations of the associated variable regions and GBS and GH loop regions are defined in Table 2 below. The VP2 capsid protein spans amino acids 137-716 of SEQ ID NO:67 and the VP3 capsid protein spans amino acids 184-716 of SEQ ID NO:67.

MSFVDHPPDWLEEIGEGLEKFLGLEPGPPKPKPNQQKDNRGLVLPGYNYLGPNGGLDRGEPVNRADDEVAREHDI SYNEQLQAGD
 NPYLKYNHADADEFQEKLADDTSFGGNLGKAVFQAKKRVLEPFGLVEEPAKTAAGERIDDHYPKKKKARVEETEAGTSGGQQLQIP
 AQPASSLGADTMSAGGGSPLGDNQAGDGVGNASGDWHCDSTWMGDRVITKSTRTWVLP SYN NHLYKEIHSGSVDGSNANAYFGYS
 TPWGYFDNRFHSHWSPRDWQRLVN NYWGF RPRSLKVKIFNIQVKEVTVDATTTIANNLSTVQVFTDDDYQLPYVIGNGTEGCL
 PAFPPQVFTLPQYGYATLNRNNTDDPTERSFFCLEYFPSKMLRTGNSFEFTYSFEVVPFHCSFAPSQNLFKLANPLVDQYLYRFV
 STDTSGNIQFQKNLKARYANTYKNWFGPMCR TQGWYTGSGTYNRSGVTNFATSNRMDLEGASYQVNPQPNGMTNTLQDSNKYALE
 NTMIFNSQNAEPGTTSLYQENLLITSESETQPVNRVAYDTGGQMATNAQSTNLAPTVGTYNHQEMLPGSVWMDRDVYLQGP IWAK
 IPETGAHFHPSPAMGGFGLKHPPPMMLIKNTPVPSNVTA FSEIPVKSFITQYSTGQVTVEMEWELKKENSKRWNP EIQYTNNYNNP
 EFVDFAPDTSGEYRTTRAIGTRYLTRPL

[00149] The VP1 sequence of a novel AAV capsid isolated from pig (denoted as Bpo.35) is set out as SEQ ID NO:68 (amino acids 1-716) and the locations of the associated variable regions and GBS and GH loop regions are defined in Table 2 below. The VP2 capsid protein spans amino acids 137-716 of SEQ ID NO:68 and the VP3 capsid protein spans amino acids 184-716 of SEQ ID NO:68.

MSFVDHPPDWLEEIGEGLEKFLGLEPGPPKPKPNQQKRDNRGLVLPGYNYLGPNGGLDRGEPVNRADDEVAREHDI SYNEQLQAGD
 NPYLKYNHADADEFQEKLADDTSFGGNLGKAVFQAKKRVLEPFGLVEEPAKTAAGERIDDHYPKKKKARVEETEAGTSGGQQLQIP
 AQPASSLGADTMSAGGGSPLGDNQAGDGVGNASGDWHCDSTWMGDRVITKSTRTWVLP SYN NHLYKEIHSGSVDGSNANAYFGYS
 TPWGYFDNRFHSHWSPRDWQRLVN NYWGF RPRSLKVKIFNIQVKEVTVDATTTIANNLSTVQVFTDDDYQLPYVIGNGTEGCL
 PAFPPQVFTLPQYGYATLNRNNTDDPTERSFFCLEYFPSKMLRTGNSFEFTYSFEVVPFHCSFAPSQNLFKLANPLVDQYLYRFV
 STDTSGNIQFQKNLKARYANTYKNWFGPMCR TQGWYTGSGTYNRSGVTNFATSNRMDLEGASYQVNPQPNGMTNTLQDSNKYALE
 NTMIFNSQNAEPGTTSLYQENLLITSESETQPVNRVAYDTGGQMATNAQSTNLAPTVGTYNHQEMLPGSVWMDRDVYLQGP IWAK
 IPETGAHFHPSPAMGGFGLKHPPPMMLIKNTPVPSNVTA FSEIPVKSFITQYSTGQVTVEMEWELKKENSKRWNP EIQYTNNYNNP
 EFVGFAPDTSGEYRTTRAIGTRYLTRPL

[00150] The VP1 sequence of a novel AAV capsid isolated from pig (denoted as Bpo.36) is set out as SEQ ID NO:69 (amino acids 1-716) and the locations of the associated variable

regions and GBS and GH loop regions are defined in Table 2 below. The VP2 capsid protein spans amino acids 137-716 of SEQ ID NO:69 and the VP3 capsid protein spans amino acids 184-716 of SEQ ID NO:69.

MSFVDHPPDWLEEIGEGLKEFLGLEPGPPKPKPNQQKQDNARGLVLPGYNYLGPNGGLDRGEPVNRADDEVAREHDI SYNEQLQAGD
NPYLKYNHADADEFQEKLADDTSGGNLKGAVFQAKKRVLEPFGLVVEEPAKTAAGGERIGDHYPKPKKARVEETEAGTSGGQQPQIP
AQPASSLGADTMSAGGGSPLGDNNQGADGVGNASGDWHCDSTWMDRVTITKSTRTWVLP SYNNHLYKEIHSGSVGDSNANAYFGYS
TPWGYFDNRFHSHWSPRDQRLVNYYWGFRRPSLKVKIFNIQVKEVTVDATTTIANNLSTTVQVFTDDDYQLPYVINGNTEGCL
PAFPPQVFTLPQYGYATLNRNNTDDPTERSFFCLEYLP SKMLRTGNSFEFTYSFEEVFPFHCSFAPSQNLFKLANPLVDQYLYRFV
STDTSGNIQFQKNLKARYANTYKNWFP GPMCRTOGWYTGSGTYNRSVGNFATSNRMDLEGASYQVNPQPNGMTNTLQDSNKYALE
NTMIFNSQNAEPGTTSLYQENNLITSESETQPVNRVAYNTGGQMATNAQSTNLAPT VGTYNHQMELPGSVWMDRDVYLQGP I WAK
IPETGAHFHPSPAMGGFGLKHPPPMMLIKNTVPVSNVTAFSEIPVKSFTITQYSTGQVTVEMEWELKKENSKRWNP EIQYTNNYNNP
EFVDFAPDTSGEYRTRTTRIGTRYLTRPL

[00151] The VP1 sequence of a novel AAV capsid isolated from pig (denoted as Bpo.37) is set out as SEQ ID NO:70 and (amino acids 1-716) and the locations of the associated variable regions and GBS and GH loop regions are defined in Table 2 below. The VP2 capsid protein spans amino acids 137-716 of SEQ ID NO:70 and the VP3 capsid protein spans amino acids 184-716 of SEQ ID NO:70.

MSFVDHPPDWLEEIGEGLKEFLGLKPGPLKPKPNQQKQDNARGLVLPGYNYLGPNGGLDRGEPVNRADDEVAREHDI SYNEQLQAGD
NPYLKYNHADADEFQEKLADDTSGGNLKGAVFQAKKRVLEPFGLVVEEPAKTAAGGERIDDHYPKPKKARVEETEAGTSGGQQPQIP
AQPASSLGADTMSAGGGSPLGDNNQGADGVGNASGDWHCDSTWMDRVTITKSTRTWVLP SYNNHLYKEIHSGSVGDSNANAYFGYS
TPWGYFDNRFHSHWSPRDQRLVNYYWGFRRPSLKVKIFNIQVKEVTVDATTTIANNLSTTVQVFTDDDYQLPYVINGNTEGCL
PAFPPQVFTLPQYGYATLNRNNTDDPTERSFFCLEYFP SKMLRTGNSFEFTYSFEEVFPFHCSFAPSQNLFKLANPLVDQYLYRFV
STDTPGNIQFQKNLKARYANTYKNWFP GPMCRTOGWYTGSGTYNRSVGNFATSNRMDLEGASYQANPQPNGMTNTLQDSNKYALE
NTMIFNSQNAEPGTTSLYQENNLITSESETQPVNRVAYDTGGQMATNAQSTNLAPT VGTYNHQMELPGSVWMDRDVYLQGP I WAK
IPETGAHFHPSPAMGGFGLKHPPPMMLIKNTVPVSNVTAFSEIPVKSFTITQYSTGQVTVEMEWELKKENSKRWNP EIQYTNNYNNP
EFVDFAPDTSGEYRTRTTRIGTRYLTRPL

[00152] The VP1 sequence of a novel AAV capsid isolated from rhesus macaque (denoted as Brh.26) is set out as SEQ ID NO:71 and (amino acids 1-736) and the locations of the associated variable regions and GBS and GH loop regions are defined in Table 2 below. The VP2 capsid protein spans amino acids 138-736 of SEQ ID NO:71 and the VP3 capsid protein spans amino acids 203-736 of SEQ ID NO:71.

MAADGYLPDWLEDNLSEGIREWDLKPGAPKPKANQQKQDDGRGLVLPGYKYLGPFGNLDKGEPANAADAAALEHDKAYDQQLKAG
DNPYLRYNHADADEFQERLQEDTSFGGNLGRAIFQAKKRVLEPLGLVEEGAKTAPGKKGPVEQSPQEPDSSSGIGKTGQQPAKKRLN
FGQTGDSESVDPDQPLGEPPAAPSGLGPNTMASGGGAPMADNNEGADGVGNSSGNWHCDSTWLGDRVITSTRTWALPTYNNHLYK
QISNGTSGGSTNDNTYFGYSTPWGYFDNRFHCHFSRWDQRLINNNWGFRRPKRLNFKLFNIQVKEVTNEGTKTIANNLSTTVQV
FTDSEYQLPYVLGSAHQGCLPPFPADVFMPQYGYLT LNNGSQALGRSSFYCLEYFPSQMLRTGNNFQFSYTFEDVPFHSYAHSQ
SLDRMLNPLIDQYLYLVLTQTGTGGTQT LAFSQAGPSSMANQARNWVGP CYRQQRVSTTTNQNNNSNFAWTGAAKFKLNGRDS
LMNPGVAMASHKDDDRFFPSSGVLFQKGTGNDGVDYSQVLTDEEEIKATNPVATEEYGAVAINNQAANTQAQTGLVHNQGVI

PGMVWQNRDVYLQGP I WAKIPHTDGNFHPSPMLGGFGLKHPPPQILIKNTFPVPADPPLTFNQAKLNSFITQYSTGQVSVEIEWELQ
KENSKRWNP EIQYTSNYYKSTNVDFAVNTEGVYSEPRPIGTRYLTRNL

[00153] The VP1 sequence of a novel AAV capsid isolated from rhesus macaque (denoted as Brh.27) is set out as SEQ ID NO:72 and (amino acids 1-736) and the locations of the associated variable regions and GBS and GH loop regions are defined in Table 2 below. The VP2 capsid protein spans amino acids 138-736 of SEQ ID NO:72 and the VP3 capsid protein spans amino acids 203-736 of SEQ ID NO:72.

MAADGYLPDWLEDNLSEGIREWDLKPGAPKPKANQQKQDDGRGLVLPGYKYLGPFGNGLDKGEPVNAADAAALEHDKAYDQQLKAG
DNPYLRYNHADA E FQERLQEDTSFGGNLGRAIFQAKKRVLEPLGLVEEGAKTAPGKKRPVEQSPQEPDSSSGIGKTGQQPAKKRLN
FGQTGDSESVDPDPQLGEPPAAPSGLGPNTMASGGGAPMADNNEGADGVGNPSGNWHCDSTRLGDRVITTTSTRTWALPTYNNHLYK
QISNGTSGGSTNDNTYFGYSTPWGYFDNRFHCHFSPRDWQRLINNNWGFPRKRLNFKLFNIQVKEVTTNEGTKTIANNLTSTVQV
FTDSEYQLPYVLGSAHQGCLPPFPADVFMPVQYGYLTLNNGSQALGRSSFYCLEYFPSQMLRTSNNFQFSYTFEDVPPFHSSYAHSQ
SLDRMLNPLIDQYLYLVRTQTTGTGGTQTLAFSQAGPSSMANQARNWVGP CYRQQRVSTTTNQNNSNFAWTGA AKFKLNGRDS
LMNPGVAMASHKDDDDRFFPSSGVLIFGKQGTGNDGVDYSQVLI TDEEEIKATNPVATEEYGAVAINNQAANTQAQTGLVHNQGVI
PGMVWQNRDVYLQGP I WAKIPHTDGNFHPSPMLGGFGLKLPPPQILIKNTFPVPADPPLTFNQAKLNSFITQYSTGQVSVEIEWELQ
KENSKRWNP EIQYTSNYYKSTNVDFAVNTEGVYSEPRPIGTRYLTRNL

[00154] The VP1 sequence of a novel AAV capsid isolated from rhesus macaque (denoted as Brh.28) is set out as SEQ ID NO:73 and (amino acids 1-736) and the locations of the associated variable regions and GBS and GH loop regions are defined in Table 2 below. The VP2 capsid protein spans amino acids 138-736 of SEQ ID NO:73 and the VP3 capsid protein spans amino acids 203-736 of SEQ ID NO:73.

MAADGYLPDWLEDNLSEGIREWDLKPGAPKPKANQQKQDDGRGLVLPGYKYLGPFGNGLDKGEPVNAADAAALEHDKAYDQQLKAG
DNPYLRYNHADA E FQERLQEDTSFGGNLGRAIFQAKKRVLEPLGLVEEGAKTAPGKKRPVEQSPQEPDSSSGIGKTGQQPAKKRLN
FGQTGDSESVDPDPQLGEPPAAPSGLGPNTMASGGGAPMADNNEGADGVGNSSGNWHCDSTWLGDVRVITTTSTRTWALPTYNNHLYK
QISNGTSGGSTNDNTYFGYSTPWGYFDNRFHCHFSPRDWQRLINNNWGFPRKRLNFKLFNIQVKEVTTNEGTKTIANNLTSTVQV
FTDSEYQLPYVLGSAHQGCLPPFPADVFMPVQYGYLTLNNGSQALGRSSFYCLEYFPSQMLRTGNNFQFSYTFEDVPPFHSSYAHSQ
SLDRMLNPLIDQYLYLVRTQTTGTGGTQTLAFSQAGPSSMANQARNWVGP CYRQQRVSTTTNQNNSNFAWTGA AKFKLNGRDS
LMNPGVAMASHKDDDDRFFPSSGVLIFGKQGTGNDGVDYSQVLI TDEEEIKATNPVATEEYGAVAINNQAANTQAQTGLVHNQGVI
PGMVWQNRDVYLQGP I WAKIPHTDGNFHPSPMLGGFGLKHPPPQILIKNTFPVPADPPLTFNQAKLNSFITQYSTGQVSVEIEWELQ
KENSKRWNP EIQYTSNYYKSTNVDFAVNTEGVYSEPRPIGTRYLTRNL

[00155] The VP1 sequence of a novel AAV capsid isolated from rhesus macaque (denoted as Brh.29) is set out as SEQ ID NO:74 and (amino acids 1-736) and the locations of the associated variable regions and GBS and GH loop regions are defined in Table 2 below. The VP2 capsid

protein spans amino acids 138-736 of SEQ ID NO:74 and the VP3 capsid protein spans amino acids 203-736 of SEQ ID NO:74.

MAADGYLPDWLEDNLSEGIREWDLKPGAPKPKANQQKQDDGRGLVLPGYKYLGPFGNLDKGEPVNAADAAALEHDKAYDQQLKAG
 DNPYLRYNHADADEFQERLQEDTSFGGNLGRAIFQAKRRVLEPLGLVEEGAKTAPGKKRPVEQSPQEPDSSSGIGKTGQQPAKKRLN
 FGQTDGSESVDPDPQLGEPPAAPSGLGPNTMASGGGAPMADNNEGADGVGNSSGNWHCDSTWLGDRVITTTSTRTWALPTYNNHLYK
 QISNGTSGGSTNDNTYFGYSTPWGYFDNRFHCHFSPRDWQRLINNNWGFRPKRLNFKLFNIQVKEVTTNEGTKTIANNLTSTVQV
 FTDSEYQLPYVLGSAHQGCLPPFPADVFMVPPQYGYLTLNNGSQALGRSSFYCLEYFPSQMLRTSNNFQFSYTFEDVPFHSSCAHSQ
 SLDRMLNPLIDQYLYLVRTQTTGTGGTQTLAFSQAGPGSMANQARNWVPGPCYRQQRVSTTTNQNNSNFAWTGAAKEFKLNGRDS
 LMNPGVAMASHKDDDDRRFPSSGVLI FGKQGTGNDGVDYSQVLI TDEEEIKATNPVATEEYGAVAINNQAANTQAQTGLVHNQGV
 PGMVWQNRDYYLQGPWAKIPHTDGNFHPPLMGGFGLKHPPPILIKNTVPVADPPLTFNQAKLNSFITQYSTGQVSVEIEWELQ
 KENSKRWNPETQYTSNYYKSTNVDFAVNTEGVYSEPRPIGTRYLTRNL

[00156] The VP1 sequence of a novel AAV capsid isolated from rhesus macaque (denoted as Brh.30) is set out as SEQ ID NO:75 and (amino acids 1-736) and the locations of the associated variable regions and GBS and GH loop regions are defined in Table 2 below. The VP2 capsid protein spans amino acids 138-736 of SEQ ID NO:75 and the VP3 capsid protein spans amino acids 203-736 of SEQ ID NO:75.

MAADGYLPDWLEDNLSEGIREWDLKPGAPKPKANQQKQDDGRGLVFPGYKYLGPFGNLDKGEPVNAADAAALEHDKAYDQQLKAG
 DNPYLRYNHADADEFQERLQEDTSFGGNLGRAIFQAKRRVLEPLGLVEEGAKTAPGKKRPVEQSPQEPDSSSGIGKTGRQPAKKRLN
 FGQTDGSESVDPDPQLGEPPAAPSGLGPNTMASGGGAPMADNNEGADGVGNSSGNWHCDSTWLGDRVITTTSTRTWALPTYNNHLYK
 QISNGTPGGSTNDNTYFGYSAPWGYFDNRFHCHFSPRDWQRLINNNWGFRPKRLNFKLFNIQVKEVTTNEGTKTIANNLTSTVQV
 FTDSEYQLPYVLGSAHQGCLPPFPADVFMVPPQYGYLTLNNGSQALGRSSFYCLEYFPSQMLRTGNFQFSYTFEDVPFHSSYAHSQ
 SLDRMLNPLIDQYLYLVRTQTTGTGGTQTLAFSQAGPSSMANQARNWVPGPCYRQQRVSTTTNQNNSNFAWTGAAKEFKLNGRDS
 LMNPGVAMASHKDDDDRRFPSSGVLI FGKQGTGNDGVDYSQVLI TDEEEIKATNPVATEEYGAVAINNQAANTQAQTGLVHNQGV
 PGMVWQNRDYYQGPWAKIPHTDGNFHPSPMLGGFGLKHPPPILIKNTVPVADPPLTFNQAKLNSFITQYSTGQVSVEIEWELQ
 KENSKRWNPETQYTSNYYKSTNVDFAVNTEGVYSEPRPIGTRYLTRNL

[00157] The VP1 sequence of a novel AAV capsid isolated from rhesus macaque (denoted as Brh.31) is set out as SEQ ID NO:76 and (amino acids 1-736) and the locations of the associated variable regions and GBS and GH loop regions are defined in Table 2 below. The VP2 capsid protein spans amino acids 138-736 of SEQ ID NO:76 and the VP3 capsid protein spans amino acids 203-736 of SEQ ID NO:76.

MAADGYLPDWLEDNLSEGIREWDLKPGAPKPKANQQKQDDGRGLVLPGYKYLGPFGNLDKGEPVNAADAAALEHDKAYDQQLKAG
 DNPYLRYNHADADEFQERLQEDTSFGGNLGRAIFQAKRRVLEPLGLVEEGAKTAPGKKRPVEQSPQEPDSSSGIGKTGQQPAKKRLN
 SGQTDGSESVDPDPQLGEPPAAPSGLGPNTMASGGGAPMADNNEGADGVGNSSGNWHCDSTWLGDRVITTTSTRTWALPTYNNHLYK

QISNGTSGGSTNDNTYFGYSTPWGYFDNRFHCHFSRWDQRLINNNWGFPRKRLNFKLFNIQVKEVTTNEGKTIANNLTSTVQV
 FTDSEYQLPYVLGSAYQGCLPPFPADVFMVPQYGYLTLNNGSQALGRSSFYCLEYFPSQMLRTGNNFQSSYTFEDVPPFHSSYAHSQ
 SLDRMLNPLIDQYLYLVRTQTTGTGGTQTLAFSQAGPSSMANQARNWVPGPCYRQQRVSTTTNQNNNSNFAWTGAAKFKLNGRDS
 LMNPGVAMASHKDDDDRFPPSSGVLI FGKQGTGNDGVDYSQVLI TDEEEIKATNPVATEEYGAVAINNQAANTQAQTGLVHNQGVI
 PGMVWQNRDYYLQGP I WAKIPHTDGNFHPSPLMGGFGLKHPPPQILIKNTFPVPADPPLTFNQAKLNSFITQYSTGQVSVEIEWELQ
 KENSKRWNP EIQYTSNYYKSTNVDFAVNTEGVYSEPRPIGTRYLTRNL

[00158] The VP1 sequence of a novel AAV capsid isolated from rhesus macaque (denoted as Brh.32) is set out as SEQ ID NO:77 and (amino acids 1-736) and the locations of the associated variable regions and GBS and GH loop regions are defined in Table 2 below. The VP2 capsid protein spans amino acids 138-736 of SEQ ID NO:77 and the VP3 capsid protein spans amino acids 203-736 of SEQ ID NO:77.

MAADGYLPDWLEDNLSEGIREWDLKPGASKPKANQQKQDDGRGLVLPGYKYLGPFGNLDKGEPVNAADAAALEHDKAYDQQLKAG
 DNPYLRYNHADA E FQERLQEDTSFGGNLGRAVFQAKKRVLEPLGLVEEGAKTVP GK KRPVEQSPQEPDSSSGIGKTGQQPAKKRLN
 FGQTGDSESVDPDQPLGEPPAAPSGLGPNTMASGGGAPMADNNEGADGVGNSSGNWHCDSTWLGDRVITSTRTWALPTYNNHLYK
 QISNGTSGGSTNDNTYFGYSTPWGYFDNRFHCHFSRWDQRLINNNWGFPRKRLNFKLFNIQVKEVTTNEGKTIANNLTSTVQV
 FTDSEYQLPYVLGSAYQGCLPPFPADVFMVPQYGYLTLNNGSQALGRSSFYCLEYFPSQMLRTGNNFQSSYTFEDVPPFHSSYAHSQ
 SLDRMLNPLIDQYLYLVRTQTTGTGGTQTLAFSQAGPSSMANQARNWVPGPCYRQQRVSTTTNQNNNSNFAWTGAAKFKLNGRDS
 LMNPGVAMASHKDDDDRFPPSSGVLI FGKQGTGNDGVDYSQVLI TDEEEIKATNPVATEEYGAVAINNQAANTQAQTGLVHNQGVI
 PGMVWQNRDYYLQGP I WAKIPHTDGNFHPSPLMGGFGLKHPPPQILIKNTFPVPADPPLTFNQAKLNSFITQYSTGQVSVEIEWELQ
 KENSKRWNP EIQYTSNYYKSTNVDFAVNTEGVYSEPRPIGTRYLTRNL

[00159] The VP1 sequence of a novel AAV capsid isolated from rhesus macaque (denoted as Brh.33) is set out as SEQ ID NO:78 and (amino acids 1-736) and the locations of the associated variable regions and GBS and GH loop regions are defined in Table 2 below. The VP2 capsid protein spans amino acids 138-736 of SEQ ID NO:78 and the VP3 capsid protein spans amino acids 203-736 of SEQ ID NO:78.

MAADGYLPDWLEDNLSEGIREWDLKPGASKPKANQQKQDDGRGLVLPGYKYLGPFGNLDKGEPVNAADAAALEHDKAYDQQLKAG
 DNPYLRYNHADA E FQERLQEDTSFGGNLGRAVFQAKKRVLEPLGLVEEGAKTVP GK KRPVEQSPQEPDSSSGIGKTGQQPAKKRLN
 FGQTGDSESVDPDQPLGEPPAAPSGLGPNTMASGGGAPMADNNEGADGVGNSSGNWHCDSTWLGDRVITSTRTWALPTYNNHLYK
 QISNGTSGGSTNDNTYFGYSTPWGYFDNRFHCHFSRWDQRLINNNWGFPRKRLNFKLFNIQVKEVTTNEGKTIANNLTSTVQV
 FTDSEYQLPYVLGSAYQGCLPPFPADVFMVPQYGYLTLNNGSQALGRSSFYCLEYFPSQMLRTGNNFQSSYTFEDVPPFHSSYAHSQ
 SLDRMLNPLIDQYLYLVRTQTTGTGGTQTLAFSQAGPSSMANQARNWVPGPCYRQQRVSTTTNQNNNSNFAWTGAAKFKLNGRDS
 LMNPGVAMASHKDDDDRFPPSSGVLI FGKQGTGNDGVDYSQVLI TDEEEIKATNPVATEEYGAVAINNQAANTQAQTGLVHNQGVI

PGMVWQNRDVYLQGP I WAKIPHTDGNFHPSPLMGGFGLKHPPPQILIKNTVPADPPLTFNQAKLNSFITQYSTGQVSVEIEWELQ
KENSKRWNP EIQYTSNYYKSTNVDFAVNTEGVYSEPRPIGTRYLTRNL

[00160] The VP1 sequence of a novel AAV capsid isolated from formosan macaque (denoted as Bfm.17) is set out as SEQ ID NO:79 and (amino acids 1-737) and the locations of the associated variable regions and GBS and GH loop regions are defined in Table 2 below. The VP2 capsid protein spans amino acids 138-736 of SEQ ID NO:79 and the VP3 capsid protein spans amino acids 203-736 of SEQ ID NO:79.

MAADGYLPDWLEDNLSEGIREWALKPGAPKPKANQQKQDDGRGLVLPGYKYLGPFGNLDKGEPVNAADAAALEHDKAYDQQLQAG
DNPYLRYNHADA E FQERLQEDTSFGGNLGRAVFQAKKRVLEPLGLVEEGAKTAPGKKRPVEQSPQEPDSSSGIGKKGQQPAKKRLN
FGQTGDSESVDPDQPLGEPPAAPSSLGLNTMAAGGGAPMADNNEGADGVGSSSGNWHCDSTRLGDRVITSTRTWALPTYNNHLYK
QISNGTSGGSTNDNTYFGYSTPWGYFDNRFHCHFSRDPWQRLINNNWGFRPKKLSFKLFNIQVKEVTQNEGTKTIANNLTSTIQV
FTDSEYQLSYVLGSAHQGLPPFPADVFMIQYGYLTLNNGSQAVGRSSFYCLEYFPSQMLRTGNNFSFSYTFEDVPPFHSSYAHSQ
SLDRLMNPLIDQYLYLSRTQSTGGTAGTQQLLFSQAGPNNMSAQAKNWLPGPMYRQQRVSTTSLQNNNSNFAWTGGTKYHLNGRD
SLVNPGVAMATNKDDEDRFFPSSGVL MFGKQGAGKDNVDYSSVMLTSEEEIKTTNPVATEQYGVVADNLQQQNTAPIVGAVNSQEA
LPGMVWQNRDVYLQGP I WAKIPHTDGNFHPSPLMGGFGLKHPPPQILIKNTVPADPPTAFNQAKLNSFITQYSTGQVSVEIEWEL
QKENS KRWNP EIQYTSNYYKSTNVDFAVDTEGVYSEHRPIGTRYLTRNL

[00161] The VP1 sequence of a novel AAV capsid isolated from formosan macaque (denoted as Bfm.18) is set out as SEQ ID NO:80 and (amino acids 1-737) and the locations of the associated variable regions and GBS and GH loop regions are defined in Table 2 below. The VP2 capsid protein spans amino acids 138-736 of SEQ ID NO:80 and the VP3 capsid protein spans amino acids 203-736 of SEQ ID NO:80.

MAADGYLPDWLEDNLSEGIREWALKPGAPKPKANQQKQDDGRGLVLPGYKYLGPFGNLDKGEPVNAADAAALEHDKAYDQQLQAG
DNPYLRYNHADA E FQERLQEDTSFGGNLGRAVFQAKKRVLEPLGLVEEGAKTAPGKKRPVEQSPQEPDSSSGIGKKGQQPAKKRLN
FGQTGDSESVDPDQPLGEPPAAPSSLGLNTMAAGGGAPMADNNEGADGVGSSSGNWHCDSTWLGDRVITSTRTWALPTYNNHLYK
QISNGTSGGSTNDNTYFGYSTPWGYFDNRFHCHFSRDPWQRLINNNWGFRPKKLSFKLFNIQVKEVTQNEGTKTIANNLTSTIQV
FTDSEYQLPYVPGSAHQGLPPFPADVFMIQYGYLTLNNGSQAVGRSSFYCLEYFPSQMLRTGNNFSFSYTFEDVPPFHSSYAHSQ
SLDRLMNPLIDQYLYLSRTQSTGGTAGTQQLLFSQAGPNNMSAQAKNWLPGPMYRQQRVSTTSLQNNNSNFAWTGGTKYHLNGRD
SLVNPGVAMATNKDDEDRFFPSSGVL MFGKQGAGKDNVDYSSVMLTSEEEIKTTNPVATEQYGVVADNLQQQNTAPIVGAVNSQGA
LPGMVWQNRDVYPQGP I WAKIPHTDGNFHPSPLMGGFGLKHPPPQILIKNTVPADPPTAFNQAKLNSFITQYSTGQVSVEIEWEL
QKENS KRWNP EIQYTSNYYKSTNVDFAVNTEGVYSEPRPIGTRYLTRNL

[00162] The VP1 sequence of a novel AAV capsid isolated from formosan macaque (denoted as Bfm.20) is set out as SEQ ID NO:81 and (amino acids 1-737) and the locations of the

associated variable regions and GBS and GH loop regions are defined in Table 2 below. The VP2 capsid protein spans amino acids 138-736 of SEQ ID NO:81 and the VP3 capsid protein spans amino acids 203-736 of SEQ ID NO:81.

```
MAADGYLPDWLEDNLSEGIREWALKPGAPKPKANQQKQDDGRGLVLPGYKYLGPFGNLDKGEPVNAADAAALEHDKAYDQQLQAG
DNPYLRYNHADADEFQERLQEDTSFGGNLGRAVFQAKKRVLEPPGLVEEGAKTAPGKKRPVEQSPQEPDSSSGIGKKGQQPAKKRLN
FGQTGDSESVDPDPQLGEPPAAPSSSLGLNTMAAGGGAPMADNNEGADGVGSSSGNWHCDSTWLGDVRVITTTSTRTWALPTYNNHLYK
QISNGTSGGSTNDNTYFGYSTPWGYFDNRFHCHFSPRDWQRLINNNWGFPRPKLSFKLFNIQVKEVTQNEGKTIANNLTSTIQV
FTDSEYQLPYVLGSAHQGCLPPFPADVFMIPQYGYLTLNNGSQAVGRSFFCYCLEYFPSQMLRTGNNFSFSYTFEDVPPHSSYAHSQ
SLDRMLNPLIDQYLYLSRTQSTGGTAGTQQLLFSQAGPNNMSAQAKNWLPGPMYRQQRVSTTSLQNNNSNFAWTGGTKYHLNGRD
SLVNPGVAMATNKDDEDRFPSSGVLMFGKQGAGKDNVDYSSVMLTSEEEIKTTNPVATEQYGVVADNLQQQNTAPIVGAVNSQGA
LPGMVWQNRDVYLQGPiWAKIPHTDGNFHPSPLMGGFGLKHPPPIILIKNTVPADPPTAFNQAKLNSFITQYSTGQVSVEIEREL
QKENSKRWNPEIQYTSNYYKSTNVDFAVNTEGVYSEPRPIGTRYLTRNL
```

[00163] The VP1 sequence of a novel AAV capsid isolated from formosan macaque (denoted as Bfm.21) is set out as SEQ ID NO:82 and (amino acids 1-737) and the locations of the associated variable regions and GBS and GH loop regions are defined in Table 2 below. The VP2 capsid protein spans amino acids 138-736 of SEQ ID NO:82 and the VP3 capsid protein spans amino acids 203-736 of SEQ ID NO:82.

```
MAADGYLPDWLEDNLSEGIREWALKPGAPKPKANQQKQDDGRGLVLPGYKYLGPFGNLDKGEPVNAADAAALEHDKAYDQQLQAG
DNPYLRYNHADADEFQERLQEDTSFGGNLGRAVFQAKKRVLEPLGLVEEGAKTAPGKERPVEQSPQEPDSSSGIGKKGQQPAKKRLN
FGQTGDSESVDPDPQLGEPPAAPSSSLGLNTMAAGGGAPMADNNEGADGVGSSSGNWHCDSTWLGDVRVITTTSTRTWALPTYNNHLYK
QISNGTSGGSTNDNTYFGYSTPWGYFDNRFHCHFSPRDWQRLINNNWGFPRPKLSFKLFNIQVKEVTQNEGKTIANNLTSTIQV
FTDSEYQLPYVLGSAHQGCLPPFPADVFMIPQYGYLTLNNGSQAVGRSFFCYCLEYFPSQMLRTGNNFSFSYTFEDVPPHSSYAHSQ
SLDRMLNPLIDQYLYLSRTQSTGGTAGTQQLLFSQAGPNNMSAQAKNWLPGPMYRQQRVSTTSLQNNNSNFAWTGGTKYHLNGRD
SLVNPGVAMATNKDDEDRFPSSAVLMFGKQGAGKDNVDYSSVMLTSEEEIKTTNPVTTEQYGVVADNLQQQNTAPIVGAVNSQGA
LPGMVWQNRDVYLQGPiWAKIPHTDGNFHPSPLMGGFGLKHPPPIILIKNTVPADPPTAFNQAKLNSFITQYSTGQVSVEIEWEL
QKENSKRWNPEIQYTSNYYKSTNVDFAVNTESVYSEPRPIGTRYLTRNL
```

[00164] The VP1 sequence of a novel AAV capsid isolated from formosan macaque (denoted as Bfm.24) is set out as SEQ ID NO:83 and (amino acids 1-737) and the locations of the associated variable regions and GBS and GH loop regions are defined in Table 2 below. The VP2 capsid protein spans amino acids 138-736 of SEQ ID NO:83 and the VP3 capsid protein spans amino acids 203-736 of SEQ ID NO:83.

```
MAADGYLPDWLEDNLSEGIREWALKPGAPKPKANQQKQDDGRGLVLPGYKYLGPFGNPDKGEPVNAADAAALEHDKAYDQQLQAG
DNPYLRYNHADADEFQERLQEDTSFGGNLGRAVFQAKKRVLEPLGLVEEGAKTAPGKKRPVEQSPQEPDSSSGIGKKGQQPAKKRLN
FGQTGDSESVDPDPQLGEPPAAPSSSLGLNTMAAGGGAPMADNNEGADGVGSSSGNWHCDSTWLGDVRVITTTSTRTWALPTYNNHLYK
```

QISNGTSGGSTNDNTYFGYSTPWGYFDNFRSHCHFSPRDWQRLINNNWGFPRPKLSFKLFNIQVKEVTQNEGKTIANNLTSTIQV
 FTDSEYQLPYVLGSAHQGCLPPFPADVFMIPQYGYLTLNNGSQAVGRSSFYCLEYFPSQMLRTGNNFSFSYTFEDVPPFHSSYAHSQ
 SLDRMLNPLIDQYLYLSRTQSTGGTAGTQQLLFSQAGPNNMSAQAKNWLPMPYRQQRVSTTSLQNNNSNFAWTGGTKYHLNLRD
 SLVNPVAMATNKDDEDRFPSSGVLMTFGKQGAGKDNVDYSSVMTLSEEEIIRTNPVATEQYGVVADNLQQQNTAPIVGAVNSQGA
 LPGMVWQNRDVYLGQPIWAKIPHTDGNFHPSPLMGGFGLKHPPQILIKNTPVPADPPTAFNQAKLNSFITQYSTGQVSVEIEWEL
 QKENSQRWNPEIQYTSNYYKSTNVDFAVNTEGVYSEPRPIGTRYLTRNL

[00165] The VP1 sequence of a novel AAV capsid isolated from formosan macaque (denoted as Bfm.25) is set out as SEQ ID NO:84 and (amino acids 1-737) and the locations of the associated variable regions and GBS and GH loop regions are defined in Table 2 below. The VP2 capsid protein spans amino acids 138-736 of SEQ ID NO:84 and the VP3 capsid protein spans amino acids 203-736 of SEQ ID NO:84.

MAADGYLPDWLEDNLSEGIREWWALKPGAPKPKANQQKQDDGRGLVLPGYKYLGPFGNLDKGEPVNAADAAALEHDKAYDQQLQAG
 DNPYLRYNHADADEFQERLQEDTSFGGNLGRAVFQAKKRVLEPLGLVEEGAKTAPGKKRPVEQSPQEPDSSSGIGKKGQQAQKRLN
 FGQTGDSSEVPDPQPLGEPPAAPSSLGLNTMAAGGAPMADNNEGADGVGSSSGNWHCDSTWLGDRTVITSTRTWALPTYNNHLYK
 QISNGTSGGSTNDNTYFGYSTPWGYFDNFRHCHFSPRDWQRLINNNWGFPRPKLSFKLFNIQVKEVTQNEGKTIANNLTSTIQV
 FTDSEYQLPYVLGSAHQGCLPPFPADVFMIPQYGYLTLNNGSQAVGRSSFYCLEYFPSQMLRTGNNFSFSYTFEDVPPFHSSYAHSQ
 SLDRMLNPLIDQYLYLSRTQSTGGTAGTQQLLFSQAGPNNMSAQAKNWLPMPYRQQRVSTTSLQNNNSNFAWTGGTKYHLNLRD
 SLVNPVAMATNKDDEDRFPSSGVLMTFGKQGAGKDNVDYSSVMTLSEEEVKTNPVATEQYGVVADNLQQQNTAPIVGAVNSQGA
 LPGMVWQNRDVYLGQPIWAKIPHTDGNFHPSPLMGGFGLKHPPQILIKNTPVPADPPTAFNQAKLNSFITQYSTGQVSVEIEWEL
 QKENSQRWNPEIQYTSNYYKSTNVDFAVNTEGVYSEPRPIGTRYLTRNL

[00166] The VP1 sequence of a novel AAV capsid isolated from formosan macaque (denoted as Bfm.27) is set out as SEQ ID NO:85 and (amino acids 1-737) and the locations of the associated variable regions and GBS and GH loop regions are defined in Table 2 below. The VP2 capsid protein spans amino acids 138-736 of SEQ ID NO:85 and the VP3 capsid protein spans amino acids 203-736 of SEQ ID NO:85.

MAADGYLPDWLEDNLSEGIREWWALKPGAPKPKANQQKQDDGRGLVLPGYKYLGPFGNLDKGEPVNAADAAALEHDKAYDQQLQAG
 DNPYLRYNHADADEFQERLQEDTSFGGNLGRAVFQAKKRVLEPLGLVEEGAKTAPGKKRPVEQSPQEPDSSSGIGKKGQQAQKRLN
 FGQTGDSSEVPDPQPLGEPPAAPSSLGLNTMAAGGAPMADNNEGADGVGSSSGNWHCDSTWLGDRTVITSTRTWALPTYNNHLYK
 QISNGTSGGSTNDNTYFGYSTPWGYFDNFRHCHFSPRDWQRLINNNWGFPRPKLSFKLFNIQVKEVTQNEGKTIANNLTSTIQV
 FTDSEYQLPYVLGSAHQGCLPPFPADVFMIPQYGYLTLNNGSQAVGRSSFYCLEYFPSQMLRTGNNFSFSYTFEDVPPFHSSYAHSQ
 SLDRMLNPLIDQYLYLSRTQSTGGTAGTQQLLFSQAGPNNMSAQAKNWLPMPYRQQRVSTTSLQNNNSNFAWTGGTKYHLNLRD
 SLVNPVAMATNKDDEDRFPSSGVLMTFGKQGAGKDNVDYSSVMTLSEEEIKTNPVATEQYGVVADNLQQQNTAPIVGAVNSQGA
 LPGMVWQNRDVYLGQPIWAKIPHTDGNFHPSPLMGGFGLKHPPQILIKNTPVPADPPTAFNQAKLNSFITQYSTGQVSVEIEWEL
 QKENSQRWNPEIQYTSNYYKSTNVDFAVDTEGVYSEPRPIGTRYLTRNL

[00167] The VP1 sequence of a novel AAV capsid isolated from formosan macaque (denoted as Bfm.32) is set out as SEQ ID NO:86 and (amino acids 1-737) and the locations of the associated variable regions and GBS and GH loop regions are defined in Table 2 below. The VP2 capsid protein spans amino acids 138-736 of SEQ ID NO:86 and the VP3 capsid protein spans amino acids 203-736 of SEQ ID NO:86.

```
MAADGYLPDWLEDNLSEGIREWALKPGAPKPKANQQKQDDGRGLVLPGYKYLGPFGNLDKGEPVNAADAAALEHDKAYDQQLQAG
DNPYLRYNHADAEEFQERLQEDTSFGGNLGRAVFQAKKRVLEPLGLVEEGAKTAPGKKRPVEQSPQEPDSSSGIGKKGQPAKKRLN
FGQTGDSESVDPDPQLGEPAPAPSSSLGLNTMAAGGGAPMADNNEGADGVGSSSGNWHCDSTWLGDVRVITTTSTRTWALPTYNHLYK
QISNGTSGGSTNDNTYFGYSTPWGYFDFNRHFHCHFSRDPWQRLINNNWGFPRPKLSFKLFDIQVKEVTQNEGKTIANNLTSTIQV
FTDSEYQLPYVLGSAHQGLPPFPADVFMIQYGYLTLNNGSQAVGRSSFYCLEYFSPQLRGTGNNFSFSYTFEDVPFHSSYAHSQ
SLDRLMNPLIDQYLYLSRTQSTGGTAGTQQLLFSQAGPNNMSAQAKNWLPGPMYRQQRVSTTSLQNNNSNFAWTGGTKYHLNLRD
SLVNPGVAMATNKDDEDRFFPSSAVLMFGKQGAGKDNVDYSSVMLTSEEEIKTTNPVATEQYGVVADNLQQQNTAPIVGAVNSQGA
LPGMVWQNRDVYLGPIWAKIPHTDGNFHPSPLMGGFGLKHPPPQILIKNTPVPADPPTAFNQAKLNSFITQYSTGQVSVEIEWEL
QKENSkrwnPEIQYTSNYYKSTNVDFAVNTEGVYSEPRPIGTRYLTRNL
```

[00168] The VP1 sequence of a novel AAV capsid isolated from formosan macaque (denoted as Bfm.33) is set out as SEQ ID NO:87 and (amino acids 1-737) and the locations of the associated variable regions and GBS and GH loop regions are defined in Table 2 below. The VP2 capsid protein spans amino acids 138-736 of SEQ ID NO:87 and the VP3 capsid protein spans amino acids 203-736 of SEQ ID NO:87.

```
MAADGYLPDWLEDNLSEGIREWALKPGAPKPKANQQKQDDGRGLVLPGYKYLGPFGNLDKGEPVNAADAAALEHDKAYDQQLQAG
DNPYLRYNHADAEEFQERLQEDTSFGGNLGRAVFQAKKRVLEPLGLVEEGAKTAPGKKRPVEQSPQEPDSSSGIGKKGQPAKKRLN
FGQTGDSESVDPDPQLGEPAPAPSSSLGLNTMAAGGGAPMADNNEGADGVGSSSGNWHCDSTWLGDVRVITTTSTRTWALPTYNHLYK
QISNGTSGGSTNDNTYFGYSTPWGYFDFNRHFHCHFSRDPWQRLINNNWGFPRPKLSFKLFNIQVKEVTQNEGKTIANNLTSTIQV
FTDSEYQLPYVLGSAHQGLPPFPADVFMIQYGYLTLNNGSQAVGRSSFYCLEYFSPQLRGTGNNFSFSYTFEDVPFHSSYAHSQ
SLDRLMNPLIDQYLYLSRTQSTGGTAGTQQLLFSQAGPNNMSAQAKNWLPGPMYRQQRVSTTSLQNNNSNFAWTGGTKYHLNLRD
SLVNPGVAMATNKDDEDRFFPSSAVLMFGKQGAGKDNVDYSSVMLTSEEEIKTTNPVATEQYGVVADNLQQQNTAPIVGAVNSQGA
LPGMVWQNRDVYLGPIWAKIPHTDGNFHPSPLMGGFGLKHPPPQILIKNTPVPADPPTAFNQAKLNSFITQYSTGQVSVEIEWEL
QKENSkrwnPEIQYTSNYYKSTNVDFAVNTESVYSEPRPIGTRYLTRNL
```

[00169] The VP1 sequence of a novel AAV capsid isolated from formosan macaque (denoted as Bfm.34) is set out as SEQ ID NO:88 and (amino acids 1-737) and the locations of the associated variable regions and GBS and GH loop regions are defined in Table 2 below. The

VP2 capsid protein spans amino acids 138-736 of SEQ ID NO:88 and the VP3 capsid protein spans amino acids 203-736 of SEQ ID NO:88.

MAADGYLPDWLEDNLSEGIREWALKPGAPKPKANQQKQDDGRGLVLPGYKYLGPFGNLDKGEPVNAADAAALEHDKAYDQQLQAG
 DNPYLRYNHADAEEFQERLQEDTSFGGNLGRAVFQAKKRVLEPLGLVEEGAKTAPGKKRPVEQSPQEPDSSSGIGKKGQPAKKRLN
 FGQTGDSESVDPDPQLGEPAPAPSSSLGLNTMAAGGGAPMADNNEGADGVGSSSGNWHCDSTWLGDRVITTTSTRTWALPTYNNHLYK
 QISNGTSGGSTNDNTYFGYSTPWGYFDFNRHFCHFSRWDQRLNNNWGFRPKKLSFKLFNIQVKEVTQNEGKTIANNLTSTIQV
 FTDSEYQLPYVLGSAHQGCLPPFPADVFMIPQYGYLTLLNNGSQAVGRSSFYCLEYFPSQMLRTGNNFSFSYTFEDVPFHSSYAHSQ
 SLDRMLNPLIDQYLYLSRTQSTGGTAGTQQLLFSQAGPNMMSAQAKNWLPGPMPYRQQRVSTTSLQNNNSNFAWTGGTKYHLNGRD
 SLVYPGVAMATNKDDEDRFPSSGVLMFGKQGAGKDNVDYSSVMLTSEEEIKTTNPVATEQYGVVADNLQQQNTAPIVGAVNSQGA
 LPGMVWQNRDVLQGPWAKIPHTDGNFHPSPLMGGFGLKHPPIQILIKNTVPADPPTAFNQAKLNSFITQYSTGQVSVEIEWEL
 QKENSKRWNPEIQYTSNYYKSTNVDFAVNTEGVYSEPRPIGTRYLTRNL

[00170] The VP1 sequence of a novel AAV capsid isolated from formosan macaque (denoted as Bfm.35) is set out as SEQ ID NO:89 and (amino acids 1-737) and the locations of the associated variable regions and GBS and GH loop regions are defined in Table 2 below. The VP2 capsid protein spans amino acids 138-736 of SEQ ID NO:89 and the VP3 capsid protein spans amino acids 203-736 of SEQ ID NO:89.

MAADGYLPDWLEDNLSEGIREWALKPGAPKPKANQQKQDDGRGLVLPGYKYLGPFGNLDKGGPVNAADAAALEHDKAYDQQLQAG
 DNPYLRYNHADAEEFRERLQEDTSFGGNLGRAVFQAKKRVLEPLGLVEEGAKTAPGKKRPVEQSPQEPDSSSGIGKKGQPAKKRLN
 FGQTGDSESVDPDPQLGEPAPAPSSSLGLNTMAAGGGAPMADNNEGADGVGSSSGNWHCDSTWLGDRVITTTSTRTWALPTYNNHLYK
 QISNGTSGGSTNDSTYFGYSTPWGYFDFNRSHCHFSRWDQRLNNNWGFRPKKLSFKLFNIQVKEVTQNEGKTIANNLTSTIQV
 FTDSEYQLPYVLGSAHQGCLPPFPADVFMIPQYGYLTLLNNGQAVGRSSFYCLEYFPSQMLRTGNNFSFSYIFEDVPFHSSYAHSQ
 SLDRMLNPLIDQYLYLSRTQSTGGTAGTQQLLFSQAGPNMMSAQAKNWLPGPMPYRQQRVSTTSLQNNNSNFAWTGGTKYHLNGRD
 SLVNPVGVAMATNKDDEDRFPSSGVLMFGKQGAGKDNVDYSSVMLTSEEEIKTTNPVATEQYGVVADNLQQQNTAPIVGAVNSQGA
 LPGMVWQNRDVLQGPWAKIPHTDGNFHPSPLMGGFGLKHPPIQILIKNTVPADPPTAFNQAKLNSFITQYSTGQVSVEIEWEL
 LKESKRWNPEIQYTSNYYKSTNVDFAVNTEGVYSEPRPIGTRYLTRNL

[00171] In Table 2 immediately below, “VR” refers to the variable region and the numbers refer to the amino acid residues each variable region or the GBS and GH loop regions span in the amino acid sequence.

Table 2

AAV	ARI	ARII	ARIII	ARIV	GBS	VRV	VRVI	VRVII	VRVIII	VRIX	GHloop
AAV2 (SEQ ID NO:3)	262- 268	325- 330	380- 384	450- 459	464- 475	487- 504	526- 538	544- 557	580- 592	703- 711	446- 600
AAV4 (SEQ ID NO:4)	256- 260	316- 321	372- 378	444- 454	459- 470	481- 504	526- 536	542- 556	579- 591	702- 710	440- 599

AAV	VRI	VRII	VRIII	VRIV	GBS	VRV	VRVI	VRVII	VRVIII	VRIX	GHloop
AAV5 (SEQ ID NO:5)	252- 258	316- 321	372- 377	443- 446	452- 462	474- 491	513- 525	531- 547	570- 582	692- 700	439- 590
AAV6 (SEQ ID NO:6)	262- 269	326- 331	381- 385	451- 460	465- 476	488- 505	527- 539	545- 558	581- 593	704- 712	447- 601
AAV6.2 (SEQ ID NO:7)	262- 269	326- 331	381- 385	451- 460	465- 476	488- 505	527- 539	545- 558	581- 593	704- 712	447- 601
AAV7 (SEQ ID NO:8)	263- 270	327- 332	382- 386	452- 462	467- 478	490- 507	529- 541	547- 559	582- 594	705- 713	448- 602
AAV8 (SEQ ID NO:9)	263- 271	328- 333	383- 387	453- 462	467- 478	490- 507	529- 541	548- 560	583- 595	706- 714	449- 603
AAV9 (SEQ ID NO:10)	262- 270	327- 332	382- 386	452- 460	465- 476	488- 505	527- 539	545- 558	581- 593	704- 712	448- 602
AAVrh8 (SEQ ID NO:11)	262- 270	327- 332	382- 386	452- 460	465- 476	488- 505	527- 539	545- 558	581- 593	704- 712	448- 602
AAVrh10 (SEQ ID NO:12)	263- 271	328- 333	383- 387	453- 462	467- 478	490- 507	529- 541	548- 560	583- 595	706- 714	449- 603
AAVbo (SEQ ID NO:1)	258- 262	318- 323	374- 380	446- 456	462- 472	484- 506	528- 538	544- 558	581- 593	704- 712	442- 601
AAVBba.21 (SEQ ID NO:15)	265- 269	325- 330	381- 386	452- 462	468- 478	490- 512	534- 544	550- 564	587- 599	710- 718	448- 607
AAVBba.31 (SEQ ID NO:20)	265- 269	325- 330	381- 386	452- 462	468- 478	490- 512	534- 544	550- 564	587- 599	710- 718	448- 607
AAVBba.32 (SEQ ID NO:21)	265- 269	325- 330	381- 386	452- 462	468- 478	490- 512	534- 544	550- 564	587- 599	710- 718	448- 607
AAVBba.33 (SEQ ID NO:22)	265- 269	325- 330	381- 386	452- 462	468- 478	490- 512	534- 544	550- 564	587- 599	710- 718	448- 607
AAVBpo.1 (SEQ ID NO:53)	243- 249	307- 312	363- 368	434- 437	443- 453	465- 483	505- 517	523- 539	562- 574	684- 692	430- 582
AAVBpo.2 (SEQ ID NO:54)	243- 249	307- 312	363- 368	434- 437	443- 453	465- 483	505- 517	523- 539	562- 574	684- 692	430- 582
AAVBpo.3 (SEQ ID NO:55)	243- 249	307- 312	363- 368	434- 437	443- 453	465- 483	505- 517	523- 539	562- 574	684- 692	430- 582
AAVBpo.4 (SEQ ID NO:56)	243- 249	307- 312	363- 368	434- 437	443- 453	465- 483	505- 517	523- 539	562- 574	684- 692	430- 582
AAVBpo.6 (SEQ ID NO:57)	243- 249	307- 312	363- 368	434- 437	443- 453	465- 483	505- 517	523- 539	562- 574	684- 692	430- 582
AAVBpo.8 (SEQ ID NO:58)	243- 249	307- 312	363- 368	434- 437	443- 453	465- 483	505- 517	523- 539	562- 574	684- 692	430- 582
AAVBpo.13 (SEQ ID NO:59)	243- 249	307- 312	363- 368	434- 437	443- 453	465- 483	505- 517	523- 539	562- 574	684- 692	430- 582
AAVBpo.18 (SEQ ID NO:60)	243- 249	307- 312	363- 368	434- 437	443- 453	465- 483	505- 517	523- 539	562- 574	684- 692	430- 582
AAVBpo.20 (SEQ ID NO:61)	243- 249	307- 312	363- 368	434- 437	443- 453	465- 483	505- 517	523- 539	562- 574	684- 692	430- 582
AAVBpo.23 (SEQ ID NO:62)	243- 249	307- 312	363- 368	434- 437	443- 453	465- 483	505- 517	523- 539	562- 574	684- 692	430- 582
AAVBpo.24 (SEQ ID NO:63)	243- 249	307- 312	363- 368	434- 437	443- 453	465- 483	505- 517	523- 539	562- 574	684- 692	430- 582
AAVBpo.27 (SEQ ID NO:64)	243- 249	307- 312	363- 368	434- 437	443- 453	465- 483	505- 517	523- 539	562- 574	684- 692	430- 582

AAV	VRI	VRII	VRIII	VRIV	GBS	VRV	VRVI	VRVII	VRVIII	VRIX	GHloop
AAVBpo.28 (SEQ ID NO:65)	243- 249	307- 312	363- 368	434- 437	443- 453	465- 483	505- 517	523- 539	562- 574	684- 692	430- 582
AAVBpo.29 (SEQ ID NO:66)	243- 249	307- 312	363- 368	434- 437	443- 453	465- 483	505- 517	523- 539	562- 574	684- 692	430- 582
AAVBpo.33 (SEQ ID NO:67)	243- 249	307- 312	363- 368	434- 437	443- 453	465- 483	505- 517	523- 539	562- 574	684- 692	430- 582
AAVBpo.35 (SEQ ID NO:68)	243- 249	307- 312	363- 368	434- 437	443- 453	465- 483	505- 517	523- 539	562- 574	684- 692	430- 582
AAVBpo.36 (SEQ ID NO:69)	243- 249	307- 312	363- 368	434- 437	443- 453	465- 483	505- 517	523- 539	562- 574	684- 692	430- 582
AAVBpo.37 (SEQ ID NO:70)	243- 249	307- 312	363- 368	434- 437	443- 453	465- 483	505- 517	523- 539	562- 574	684- 692	430- 582
AAVmo (SEQ ID NO:2)	249- 252	310- 315	366- 371	437- 439	443- 455	467- 484	507- 519	525- 533	556- 568	679- 689	433- 576
AAVBce.14 (SEQ ID NO:32)	262- 270	327- 332	382- 386	452- 460	465- 476	488- 505	527- 539	545- 558	581- 593	704- 712	448- 602
AAVBce.15 (SEQ ID NO:33)	262- 270	327- 332	382- 386	452- 460	465- 476	488- 505	527- 539	545- 558	581- 593	704- 712	448- 602
AAVBce.16 (SEQ ID NO:34)	262- 270	327- 332	382- 386	452- 460	465- 476	488- 505	527- 539	545- 558	581- 593	704- 712	448- 602
AAVBce.17 (SEQ ID NO:35)	262- 270	327- 332	382- 386	452- 460	465- 476	488- 505	527- 539	545- 558	581- 593	704- 712	448- 602
AAVBce.18 (SEQ ID NO:36)	262- 270	327- 332	382- 386	452- 460	465- 476	488- 505	527- 539	545- 558	581- 593	704- 712	448- 602
AAVBce.20 (SEQ ID NO:37)	262- 270	327- 332	382- 386	452- 460	465- 476	488- 505	527- 539	545- 558	581- 593	704- 712	448- 602
AAVBma.42 (SEQ ID NO:51)	262- 270	327- 332	382- 386	452- 460	465- 476	488- 505	527- 539	545- 558	581- 593	704- 712	448- 602
AAVBma.43 (SEQ ID NO:52)	262- 270	327- 332	382- 386	452- 460	465- 476	488- 505	527- 539	545- 558	581- 593	704- 712	448- 602
AAVBce.39 (SEQ ID NO:40)	262- 270	327- 332	382- 386	452- 460	465- 476	488- 505	527- 539	545- 558	581- 593	704- 712	448- 602
AAVBce.40 (SEQ ID NO:41)	262- 270	327- 332	382- 386	452- 460	465- 476	488- 505	527- 539	545- 558	581- 593	704- 712	448- 602
AAVBce.41 (SEQ ID NO:42)	262- 270	327- 332	382- 386	452- 460	465- 476	488- 505	527- 539	545- 558	581- 593	704- 712	448- 602
AAVBce.42 (SEQ ID NO:43)	262- 270	327- 332	382- 386	452- 460	465- 476	488- 505	527- 539	545- 558	581- 593	704- 712	448- 602
AAVanc110 (SEQ ID NO:14)	262- 270	327- 332	382- 386	452- 460	465- 476	488- 505	527- 539	545- 558	581- 593	704- 712	448- 602
AAVanc80 (SEQ ID NO:13)	262- 269	326- 331	381- 385	451- 460	465- 476	488- 505	527- 539	545- 558	581- 593	704- 712	447- 601
AAVBce.35 (SEQ ID NO:38)	258- 265	322- 327	377- 381	447- 455	460- 471	483- 500	522- 534	540- 552	575- 587	698- 706	443- 595
AAVBce.36 (SEQ ID NO:39)	258- 265	322- 327	377- 381	447- 455	460- 471	483- 500	522- 534	540- 552	575- 587	698- 706	443- 595
AAVBce.43 (SEQ ID NO:44)	258- 265	322- 327	377- 381	447- 455	460- 471	483- 500	522- 534	540- 552	575- 587	698- 706	443- 595
AAVBce.44 (SEQ ID NO:45)	258- 265	322- 327	377- 381	447- 455	460- 471	483- 500	522- 534	540- 552	575- 587	698- 706	443- 595

AAV	VRI	VRII	VRIII	VRIV	GBS	VRV	VRVI	VRVII	VRVIII	VRIX	GHloop
AAVBce.45 (SEQ ID NO:46)	258- 265	322- 327	377- 381	447- 455	460- 471	483- 500	522- 534	540- 552	575- 587	698- 706	443- 595
AAVBce.46 (SEQ ID NO:47)	258- 265	322- 327	377- 381	447- 455	460- 471	483- 500	522- 534	540- 552	575- 587	698- 706	443- 595
AAVBcy.20 (SEQ ID NO:48)	258- 265	322- 327	377- 381	447- 455	460- 471	483- 500	522- 534	540- 552	575- 587	698- 706	443- 595
AAVBcy.22 (SEQ ID NO:49)	258- 265	322- 327	377- 381	447- 455	460- 471	483- 500	522- 534	540- 552	575- 587	698- 706	443- 595
AAVBcy.23 (SEQ ID NO:50)	258- 265	322- 327	377- 381	447- 455	460- 471	483- 500	522- 534	540- 552	575- 587	698- 706	443- 595
AAVBba.26 (SEQ ID NO:16)	265- 271	328- 333	383- 387	453- 464	469- 480	492- 508	531- 543	549- 561	584- 596	707- 715	449- 604
AAVBba.27 (SEQ ID NO:17)	265- 271	328- 333	383- 387	453- 464	469- 480	492- 508	531- 543	549- 561	584- 596	707- 715	449- 604
AAVBba.29 (SEQ ID NO:18)	265- 271	328- 333	383- 387	453- 464	469- 480	492- 508	531- 543	549- 561	584- 596	707- 715	449- 604
AAVBba.30 (SEQ ID NO:19)	265- 271	328- 333	383- 387	453- 464	469- 480	492- 508	531- 543	549- 561	584- 596	707- 715	449- 604
AAVBba.34 (SEQ ID NO:23)	265- 271	328- 333	383- 387	453- 464	469- 480	492- 508	531- 543	549- 561	584- 596	707- 715	449- 604
AAVBba.35 (SEQ ID NO:24)	265- 271	328- 333	383- 387	453- 464	469- 480	492- 508	531- 543	549- 561	584- 596	707- 715	449- 604
AAVBba.36 (SEQ ID NO:25)	265- 271	328- 333	383- 387	453- 464	469- 480	492- 508	531- 543	549- 561	584- 596	707- 715	449- 604
AAVBba.37 (SEQ ID NO:26)	265- 271	328- 333	383- 387	453- 464	469- 480	492- 508	531- 543	549- 561	584- 596	707- 715	449- 604
AAVBba.38 (SEQ ID NO:27)	265- 271	328- 333	383- 387	453- 464	469- 480	492- 508	531- 543	549- 561	584- 596	707- 715	449- 604
AAVBba.41 (SEQ ID NO:28)	265- 271	328- 333	383- 387	453- 464	469- 480	492- 508	531- 543	549- 561	584- 596	707- 715	449- 604
AAVBba.42 (SEQ ID NO:29)	265- 271	328- 333	383- 387	453- 464	469- 480	492- 508	531- 543	549- 561	584- 596	707- 715	449- 604
AAVBba.43 (SEQ ID NO:30)	265- 271	328- 333	383- 387	453- 464	469- 480	492- 508	531- 543	549- 561	584- 596	707- 715	449- 604
AAVBba.44 (SEQ ID NO:31)	265- 271	328- 333	383- 387	453- 464	469- 480	492- 508	531- 543	549- 561	584- 596	707- 715	449- 604
AAVBrh.26 (SEQ ID NO:71)	262- 270	327- 332	382- 386	452- 460	465- 476	488- 505	527- 539	545- 558	581- 593	704- 712	448- 601
AAVBrh.27 (SEQ ID NO:72)	262- 270	327- 332	382- 386	452- 460	465- 476	488- 505	527- 539	545- 558	581- 593	704- 712	448- 601
AAVBrh.28 (SEQ ID NO:73)	262- 270	327- 332	382- 386	452- 460	465- 476	488- 505	527- 539	545- 558	581- 593	704- 712	448- 601
AAVBrh.29 (SEQ ID NO:74)	262- 270	327- 332	382- 386	452- 460	465- 476	488- 505	527- 539	545- 558	581- 593	704- 712	448- 601
AAVBrh.30 (SEQ ID NO:75)	262- 270	327- 332	382- 386	452- 460	465- 476	488- 505	527- 539	545- 558	581- 593	704- 712	448- 601
AAVBrh.31 (SEQ ID NO:76)	262- 270	327- 332	382- 386	452- 460	465- 476	488- 505	527- 539	545- 558	581- 593	704- 712	448- 601
AAVBrh.32 (SEQ ID NO:77)	262- 270	327- 332	382- 386	452- 460	465- 476	488- 505	527- 539	545- 558	581- 593	704- 712	448- 601

AAV	VRI	VRII	VRIII	VRIV	GBS	VRV	VRVI	VRVII	VRVIII	VRIX	GHloop
AAVBrh.33 (SEQ ID NO:78)	262- 270	327- 332	382- 386	452- 460	465- 476	488- 505	527- 539	545- 558	581- 593	704- 712	448- 601
AAVBfm.17 (SEQ ID NO:79)	262- 270	327- 332	382- 386	452- 461	466- 477	489- 506	528- 540	546- 559	582- 594	705- 713	448- 602
AAVBfm.18 (SEQ ID NO:80)	262- 270	327- 332	382- 386	452- 461	466- 477	489- 506	528- 540	546- 559	582- 594	705- 713	448- 602
AAVBfm.20 (SEQ ID NO:81)	262- 270	327- 332	382- 386	452- 461	466- 477	489- 506	528- 540	546- 559	582- 594	705- 713	448- 602
AAVBfm.21 (SEQ ID NO:82)	262- 270	327- 332	382- 386	452- 461	466- 477	489- 506	528- 540	546- 559	582- 594	705- 713	448- 602
AAVBfm.24 (SEQ ID NO:83)	262- 270	327- 332	382- 386	452- 461	466- 477	489- 506	528- 540	546- 559	582- 594	705- 713	448- 602
AAVBfm.25 (SEQ ID NO:84)	262- 270	327- 332	382- 386	452- 461	466- 477	489- 506	528- 540	546- 559	582- 594	705- 713	448- 602
AAVBfm.27 (SEQ ID NO:85)	262- 270	327- 332	382- 386	452- 461	466- 477	489- 506	528- 540	546- 559	582- 594	705- 713	448- 602
AAVBfm.32 (SEQ ID NO:86)	262- 270	327- 332	382- 386	452- 461	466- 477	489- 506	528- 540	546- 559	582- 594	705- 713	448- 602
AAVBfm.33 (SEQ ID NO:87)	262- 270	327- 332	382- 386	452- 461	466- 477	489- 506	528- 540	546- 559	582- 594	705- 713	448- 602
AAVBfm.34 (SEQ ID NO:88)	262- 270	327- 332	382- 386	452- 461	466- 477	489- 506	528- 540	546- 559	582- 594	705- 713	448- 602
AAVBfm.35 (SEQ ID NO:89)	262- 270	327- 332	382- 386	452- 461	466- 477	489- 506	528- 540	546- 559	582- 594	705- 713	448- 602

Transgene

[00172] The transgene is a nucleic acid sequence, heterologous to the vector sequences flanking the transgene, which encodes a polypeptide, protein, or other product, of interest. The nucleic acid coding sequence is operatively linked to regulatory components in a manner which permits transgene transcription, translation, and/or expression in a host cell.

[00173] The composition of the transgene sequence will depend upon the use to which the resulting vector will be put. For example, one type of transgene sequence includes a reporter sequence, which upon expression produces a detectable signal. Such reporter sequences include, without limitation, DNA sequences encoding β -lactamase, β -galactosidase (LacZ), alkaline phosphatase, thymidine kinase, green fluorescent protein (GFP), chloramphenicol acetyltransferase (CAT), luciferase, membrane bound proteins including, for example, CD2, CD4, CD8, the influenza hemagglutinin protein, and others well known in the art, to which high affinity antibodies directed thereto exist or can be produced by conventional means, and fusion proteins comprising a membrane bound protein appropriately fused to an antigen tag domain from, among others, hemagglutinin or Myc.

[00174] These coding sequences, when associated with regulatory elements which drive their expression, provide signals detectable by conventional means, including enzymatic, radiographic, colorimetric, fluorescence or other spectrographic assays, fluorescent activating cell sorting assays and immunological assays, including enzyme linked immunosorbent assay (ELISA), radioimmunoassay (RIA) and immunohistochemistry. For example, where the marker sequence is the LacZ gene, the presence of the vector carrying the signal is detected by assays for beta-galactosidase activity. Where the transgene is green fluorescent protein or luciferase, the vector carrying the signal may be measured visually by color or light production in a luminometer.

[00175] However, desirably, the transgene is a non-marker sequence encoding a product which is useful in biology and medicine, such as proteins, peptides, RNA, enzymes, dominant negative mutants, or catalytic RNAs. Desirable RNA molecules include tRNA, dsRNA, ribosomal RNA, catalytic RNAs, siRNA, small hairpin RNA, trans-splicing RNA, and antisense RNAs. One example of a useful RNA sequence is a sequence which inhibits or extinguishes expression of a targeted nucleic acid sequence in the treated animal. Typically, suitable target sequences include oncologic targets and viral diseases. See, for examples of such targets the oncologic targets and viruses identified below in the section relating to immunogens.

[00176] The transgene may be used to correct or ameliorate gene deficiencies, which may include deficiencies in which normal genes are expressed at less than normal levels or deficiencies in which the functional gene product is not expressed. A preferred type of transgene sequence encodes a therapeutic protein or polypeptide which is expressed in a host cell. The invention further includes using multiple transgenes, e.g., to correct or ameliorate a gene defect caused by a multi-subunit protein. In certain situations, a different transgene may be used to encode each subunit of a protein, or to encode different peptides or proteins. This is desirable when the size of the DNA encoding the protein subunit is large, e.g., for an immunoglobulin, the platelet-derived growth factor, or a dystrophin protein. In order for the cell to produce the multi-subunit protein, a cell is infected with the recombinant virus containing each of the different subunits. Alternatively, different subunits of a protein may be encoded by the same transgene. In this case, a single transgene includes the DNA encoding each of the subunits, with the DNA for each subunit separated by an internal ribozyme entry site (IRES). This is desirable when the size of the DNA encoding each of the subunits is small, e.g., the total size of the DNA encoding the

subunits and the IRES is less than five kilobases. As an alternative to an IRES, the DNA may be separated by sequences encoding a 2A peptide, which self-cleaves in a post-translational event. See, e.g., Donnelly *et al*, *J. Gen. Virol.*, 78(Pt 1):13-21 (January 1997); Furler, *et al*, *Gene Ther.*, 8(11):864-873 (June 2001); Klump *et al.*, *Gene Ther.*, 8(10):811-817 (May 2001). This 2A peptide is significantly smaller than an IRES, making it well suited for use when space is a limiting factor. More often, when the transgene is large, consists of multi-subunits, or two transgenes are co-delivered, rAAV carrying the desired transgene(s) or subunits are co-administered to allow them to concatamerize *in vivo* to form a single vector genome. In such an embodiment, a first AAV may carry an expression cassette which expresses a single transgene and a second AAV may carry an expression cassette which expresses a different transgene for co-expression in the host cell. However, the selected transgene may encode any biologically active product or other product, e.g., a product desirable for study.

[00177] Suitable transgenes may be readily selected by one of skill in the art. The selection of the transgene is not considered to be a limitation of this invention.

[00178] In some embodiments, the transgene is a heterologous protein, and this heterologous protein is a therapeutic protein. Exemplary therapeutic proteins include, but are not limited to, blood factors, such as β -globin, hemoglobin, tissue plasminogen activator, and coagulation factors; colony stimulating factors (CSF); interleukins, such as IL-1, IL-2, IL-3, IL-4, IL-5, IL-6, IL-7, IL-8, IL-9, etc.; growth factors, such as keratinocyte growth factor (KGF), stem cell factor (SCF), fibroblast growth factor (FGF, such as basic FGF and acidic FGF), hepatocyte growth factor (HGF), insulin-like growth factors (IGFs), bone morphogenetic protein (BMP), epidermal growth factor (EGF), growth differentiation factor-9 (GDF-9), hepatoma derived growth factor (HDGF), myostatin (GDF-8), nerve growth factor (NGF), neurotrophins, platelet-derived growth factor (PDGF), thrombopoietin (TPO), transforming growth factor alpha (TGF- α), transforming growth factor beta (TGF- β), and the like; soluble receptors, such as soluble TNF- α receptors, soluble VEGF receptors, soluble interleukin receptors (e.g., soluble IL-1 receptors and soluble type II IL-1 receptors), soluble γ/δ T cell receptors, ligand-binding fragments of a soluble receptor, and the like; enzymes, such as α -glucosidase, imiglucrase, β -glucocerebrosidase, and alglucerase; enzyme activators, such as tissue plasminogen activator; chemokines, such as IP-10, monokine induced by interferon-gamma (Mig), Gro α /IL-8, RANTES, MIP-1 α , MIP-1 β , MCP-1, PF-4, and the like; angiogenic agents, such as vascular endothelial growth factors (VEGFs, e.g.,

VEGF121, VEGF165, VEGF-C, VEGF-2), glioma-derived growth factor, angiogenin, angiogenin-2; and the like; anti-angiogenic agents, such as a soluble VEGF receptor; protein vaccine; neuroactive peptides, such as nerve growth factor (NGF), bradykinin, cholecystokinin, gastrin, secretin, oxytocin, gonadotropin-releasing hormone, beta-endorphin, enkephalin, substance P, somatostatin, prolactin, galanin, growth hormone-releasing hormone, bombesin, dynorphin, warfarin, neurotensin, motilin, thyrotropin, neuropeptide Y, luteinizing hormone, calcitonin, insulin, glucagons, vasopressin, angiotensin II, thyrotropin-releasing hormone, vasoactive intestinal peptide, a sleep peptide, and the like; thrombolytic agents; atrial natriuretic peptide; relaxin; glial fibrillary acidic protein; follicle stimulating hormone (FSH); human alpha-1 antitrypsin; leukemia inhibitory factor (LIF); tissue factors, luteinizing hormone; macrophage activating factors; tumor necrosis factor (TNF); neutrophil chemotactic factor (NCF); tissue inhibitors of metalloproteinases; vasoactive intestinal peptide; angiogenin; angiotropin; fibrin; hirudin; IL-1 receptor antagonists; and the like. Some other non-limiting examples of protein of interest include ciliary neurotrophic factor (CNTF); brain-derived neurotrophic factor (BDNF); neurotrophins 3 and 4/5 (NT-3 and 4/5); glial cell derived neurotrophic factor (GDNF); aromatic amino acid decarboxylase (AADC); hemophilia related clotting proteins, such as Factor VIII, Factor IX, Factor X; dystrophin, mini-dystrophin, or microdystrophin; lysosomal acid lipase; phenylalanine hydroxylase (PAH); glycogen storage disease-related enzymes, such as glucose-6-phosphatase, acid maltase, glycogen debranching enzyme, muscle glycogen phosphorylase, liver glycogen phosphorylase, muscle phosphofructokinase, phosphorylase kinase (e.g., PHKA2), glucose transporter (e.g., GLUT2), aldolase A, β -enolase, and glycogen synthase; lysosomal enzymes (e.g., beta-N-acetylhexosaminidase A); and any variants thereof.

[00179] In one preferred embodiment, the transgene encodes a protein that restores dystrophin function and is useful to treat Duchenne muscular dystrophy. Duchenne muscular dystrophy is a degenerative muscle disease caused by deletions or mutations in the X-linked gene that encodes the protein dystrophin. The absence of functional dystrophin causes muscle fiber degeneration, inflammation, and necrosis. The dystrophin gene is the largest known gene in the human genome, covering over 2.5Mb of the human X chromosome. The dystrophin gene has 79 exons, transcription of which results in an 11kb RNA transcript that encodes a protein consisting of 3,685 amino acids. A transgene encoding wild-type dystrophin exceeds the packing limit of known gene therapy vector systems. To overcome the packaging limitation, engineered

synthetic versions of dystrophin have been generated. As used herein “microdystrophin” or “minidystrophin” refer to transgenes that encode truncation but functional dystrophins and have been described in WO2016177911, WO2015197869, and U.S. Patent Application Publication No. US2008249052 each of which is incorporated by reference in their entirety herein.

Illustrative examples of microdystrophins include MD1 (a microdystrophin lacking spectrin like repeats 4 through 23 and lacking a C-terminal domain), MD3 (a codon optimized microdystrophin lacking spectrin-like repeats 4-23 while retaining spectrin-like repeats 1, 2, 3, and 24, and further retaining exons 70-75 of the C-terminal domain encoding the coiled-coil region helix 1 and 2 of the C-terminal domain), and MD4 (identical to human MD3, except that it retains the entire C-terminal domain (all of exons 70-79)).

Regulatory Control Elements

[00180] The AAV vector also includes conventional control elements or sequences which are operably linked to the transgene in a manner which permits its transcription, translation and/or expression in a cell transfected with the plasmid vector or infected with the virus produced by the invention. As used herein, "operably linked" sequences include both expression control sequences that are contiguous with the gene of interest and expression control sequences that act in trans or at a distance to control the gene of interest.

[00181] Expression control sequences include appropriate transcription initiation, termination, promoter and enhancer sequences; efficient RNA processing signals such as splicing and polyadenylation (polyA) signals; sequences that stabilize cytoplasmic mRNA; sequences that enhance translation efficiency (*i.e.*, Kozak consensus sequence); sequences that enhance protein stability; and when desired, sequences that enhance secretion of the encoded product. A great number of expression control sequences, including promoters which are native, constitutive, inducible and/or tissue-specific, are known in the art and may be utilized.

[00182] Examples of constitutive promoters include, without limitation, the retroviral Rous sarcoma virus (RSV) LTR promoter (optionally with the RSV enhancer), the cytomegalovirus (CMV) promoter (optionally with the CMV enhancer) (*see, e.g., Boshart et al, Cell, 41:521-530 (1985)*), the SV40 promoter, the dihydrofolate reductase promoter, the β -actin promoter, the phosphoglycerol kinase (PGK) promoter, and the EF1 promoter [Invitrogen]. Inducible promoters allow regulation of gene expression and can be regulated by exogenously supplied

compounds, environmental factors such as temperature, or the presence of a specific physiological state, e.g., acute phase, a particular differentiation state of the cell, or in replicating cells only. Inducible promoters and inducible systems are available from a variety of commercial sources, including, without limitation, Invitrogen, Clontech and Ariad. Many other systems have been described and can be readily selected by one of skill in the art. Examples of inducible promoters regulated by exogenously supplied compounds, include, the zinc-inducible sheep metallothionein (MT) promoter, the dexamethasone (Dex)-inducible mouse mammary tumor virus (MMTV) promoter, the T7 polymerase promoter system [WO 98/10088]; the ecdysone insect promoter [No *et al*, *Proc. Natl. Acad. Sci. USA*, 93:3346-3351 (1996)], the tetracycline-repressible system [Gossen *et al*, *Proc. Natl. Acad. Sci. USA*, 89:5547-5551 (1992)], the tetracycline-inducible system [Gossen *et al*, *Science*, 268:1766-1769 (1995), see also Harvey *et al*, *Curr. Opin. Chem. Biol.*, 2:512-518 (1998)], the RU486-inducible system [Wang *et al*, *Nat. Biotech.*, 15:239-243 (1997) and Wang *et al*, *Gene Ther.*, 4:432-441 (1997)] and the rapamycin-inducible system [Magari *et al*, *J. Clin. Invest.*, 100:2865-2872 (1997)]. Other types of inducible promoters which may be useful in this context are those which are regulated by a specific physiological state, e.g., temperature, acute phase, a particular differentiation state of the cell, or in replicating cells only.

[00183] In another embodiment, the native promoter for the transgene will be used. The native promoter may be preferred when it is desired that expression of the transgene should mimic the native expression. The native promoter may be used when expression of the transgene must be regulated temporally or developmentally, or in a tissue-specific manner, or in response to specific transcriptional stimuli. In a further embodiment, other native expression control elements, such as enhancer elements, polyadenylation sites or Kozak consensus sequences may also be used to mimic the native expression.

[00184] Another embodiment of the transgene includes a gene operably linked to a tissue-specific promoter. For instance, if expression in skeletal muscle is desired, a promoter active in muscle should be used. These include the promoters from genes encoding skeletal β -actin, myosin light chain 2A, dystrophin, muscle creatine kinase, as well as synthetic muscle promoters with activities higher than naturally-occurring promoters (see Li *et al*, *Nat. Biotech.*, 17:241-245 (1999)). Examples of promoters that are tissue-specific are known for liver (albumin, Miyatake *et al*, *J. Virol.*, 71:5124-32 (1997); hepatitis B virus core promoter, Sandig *et al*, *Gene Ther.*,

3:1002-9 (1996); alpha-fetoprotein (AFP), Arbutnot *et al.*, *Hum. Gene Ther.*, 7:1503-14 (1996)), bone osteocalcin (Stein *et al.*, *Mol. Biol. Rep.*, 24:185-96 (1997)); bone sialoprotein (Chen *et al.*, *J. Bone Miner. Res.*, 11:654-64 (1996)), lymphocytes (CD2, Hansal *et al.*, *J. Immunol.*, 161:1063-8 (1998); immunoglobulin heavy chain; T cell receptor chain), neuronal such as neuron-specific enolase (NSE) promoter (Andersen *et al.*, *Cell. Mol. Neurobiol.*, 13:503-15 (1993)), neurofilament light-chain gene (Piccioli *et al.*, *Proc. Natl. Acad. Sci. USA*, 88:5611-5 (1991)), and the neuron-specific vgf gene (Piccioli *et al.*, *Neuron*, 15:373-84 (1995)), among others.

[00185] Optionally, plasmids carrying therapeutically useful transgenes may also include selectable markers or reporter genes may include sequences encoding geneticin, hygromycin or purimycin resistance, among others. Such selectable reporters or marker genes (preferably located outside the viral genome to be rescued by the method of the invention) can be used to signal the presence of the plasmids in bacterial cells, such as ampicillin resistance. Other components of the plasmid may include an origin of replication. Selection of these and other promoters and vector elements are conventional and many such sequences are available [see, e.g., Sambrook *et al.*, and references cited therein].

Methods for Producing Recombinant AAVs

[00186] The present disclosure provides materials and methods for producing recombinant AAVs in insect or mammalian cells. In some embodiments, the viral construct further comprises a promoter and a restriction site downstream of the promoter to allow insertion of a polynucleotide encoding one or more proteins of interest, wherein the promoter and the restriction site are located downstream of the 5' AAV ITR and upstream of the 3' AAV ITR. In some embodiments, the viral construct further comprises a posttranscriptional regulatory element downstream of the restriction site and upstream of the 3' AAV ITR. In some embodiments, the viral construct further comprises a polynucleotide inserted at the restriction site and operably linked with the promoter, where the polynucleotide comprises the coding region of a protein of interest. As a skilled artisan will appreciate, any one of the AAV vector disclosed in the present application can be used in the method as the viral construct to produce the recombinant AAV.

[00187] In some embodiments, the helper functions are provided by one or more helper plasmids or helper viruses comprising adenoviral or baculoviral helper genes. Non-limiting

examples of the adenoviral or baculoviral helper genes include, but are not limited to, E1A, E1B, E2A, E4 and VA, which can provide helper functions to AAV packaging.

[00188] Helper viruses of AAV are known in the art and include, for example, viruses from the family *Adenoviridae* and the family *Herpesviridae*. Examples of helper viruses of AAV include, but are not limited to, SAdV-13 helper virus and SAdV-13-like helper virus described in US Publication No. 20110201088 (the disclosure of which is incorporated herein by reference), helper vectors pHELP (Applied Viromics). A skilled artisan will appreciate that any helper virus or helper plasmid of AAV that can provide adequate helper function to AAV can be used herein.

[00189] In some embodiments, the AAV cap genes are present in a plasmid. The plasmid can further comprise an AAV rep gene. The cap genes and/or rep gene from any AAV serotype (including, but not limited to, AAV1, AAV2, AAV4, AAV5, AAV6, AAV7, AAV8, AAV9, AAV10, AAV11, AAV12, AAV13 and any variants thereof) can be used herein to produce the recombinant AAV. In some embodiments, the AAV cap genes encode a capsid from serotype 1, serotype 2, serotype 4, serotype 5, serotype 6, serotype 7, serotype 8, serotype 9, serotype 10, serotype 11, serotype 12, serotype 13 or a variant thereof.

[00190] In some embodiments, the insect or mammalian cell can be transfected with the helper plasmid or helper virus, the viral construct and the plasmid encoding the AAV cap genes; and the recombinant AAV virus can be collected at various time points after co-transfection. For example, the recombinant AAV virus can be collected at about 12 hours, about 24 hours, about 36 hours, about 48 hours, about 72 hours, about 96 hours, about 120 hours, or a time between any of these two time points after the co-transfection.

[00191] Recombinant AAV can also be produced using any conventional methods known in the art suitable for producing infectious recombinant AAV. In some instances, a recombinant AAV can be produced by using an insect or mammalian cell that stably expresses some of the necessary components for AAV particle production. For example, a plasmid (or multiple plasmids) comprising AAV rep and cap genes, and a selectable marker, such as a neomycin resistance gene, can be integrated into the genome of the cell. The insect or mammalian cell can then be co-infected with a helper virus (e.g., adenovirus or baculovirus providing the helper functions) and the viral vector comprising the 5' and 3' AAV ITR (and the nucleotide sequence encoding the heterologous protein, if desired). The advantages of this method are that the cells

are selectable and are suitable for large-scale production of the recombinant AAV. As another non-limiting example, adenovirus or baculovirus rather than plasmids can be used to introduce rep and cap genes into packaging cells. As yet another non-limiting example, both the viral vector containing the 5' and 3' AAV LTRs and the rep-cap genes can be stably integrated into the DNA of producer cells, and the helper functions can be provided by a wild-type adenovirus to produce the recombinant AAV.

Cell Types Used in AAV Production

[00192] The viral particles comprising the AAV vectors of the invention may be produced using any invertebrate cell type which allows for production of AAV or biologic products and which can be maintained in culture. For example, the insect cell line used can be from *Spodoptera frugiperda*, such as Sf9, SF21, SF900+, drosophila cell lines, mosquito cell lines, e.g., *Aedes albopictus* derived cell lines, domestic silkworm cell lines, e.g. Bombyxmori cell lines, *Trichoplusia ni* cell lines such as High Five cells or Lepidoptera cell lines such as *Ascalapha odorata* cell lines. Preferred insect cells are cells from the insect species which are susceptible to baculovirus infection, including High Five, Sf9, Se301, SeIZD2109, SeUCR1, Sf900+, Sf21, BTI-TN-5B1-4, MG-1, Tn368, HzAm1, BM-N, Ha2302, Hz2E5 and Ao38.

[00193] Baculoviruses are enveloped DNA viruses of arthropods, two members of which are well known expression vectors for producing recombinant proteins in cell cultures. Baculoviruses have circular double-stranded genomes (80-200 kbp) which can be engineered to allow the delivery of large genomic content to specific cells. The viruses used as a vector are generally *Autographa californica* multicapsid nucleopolyhedrovirus (AcMNPV) or *Bombyx mori* (Bm-NPV) (Kato et al., 2010).

[00194] Baculoviruses are commonly used for the infection of insect cells for the expression of recombinant proteins. In particular, expression of heterologous genes in insects can be accomplished as described in for instance U.S. Pat. No. 4,745,051; Friesen et al (1986); EP 127,839; EP 155,476; Vlak et al (1988); Miller et al (1988); Carbonell et al (1988); Maeda et al (1985); Lebacqz-Verheyden et al (1988); Smith et al (1985); Miyajima et al (1987); and Martin et al (1988). Numerous baculovirus strains and variants and corresponding permissive insect host cells that can be used for protein production are described in Luckow et al (1988), Miller et al (1986); Maeda et al (1985) and McKenna (1989).

[00195] In another aspect of the invention, the methods of the invention are also carried out with any mammalian cell type which allows for replication of AAV or production of biologic products, and which can be maintained in culture. Preferred mammalian cells used can be HEK293, HeLa, CHO, NS0, SP2/0, PER.C6, Vero, RD, BHK, HT 1080, A549, Cos-7, ARPE-19 and MRC-5 cells.

Production of Heterologous Proteins In Vitro

[00196] As a non-limiting example, the recombinant AAV disclosed herein can be used to produce a protein of interest *in vitro*, for example, in a cell culture. As one non-limiting example, in some embodiments, a method for producing a protein of interest *in vitro*, where the method includes providing a recombinant AAV comprising a nucleotide sequence encoding the heterologous protein; and contacting the recombinant AAV with a cell in a cell culture, whereby the recombinant AAV expresses the protein of interest in the cell. The size of the nucleotide sequence encoding the protein of interest can vary. For example, the nucleotide sequence can be at least about 1.4 kb, at least about 1.5 kb, at least about 1.6 kb, at least about 1.7 kb, at least about 1.8 kb, at least about 2.0 kb, at least about 2.2 kb, at least about 2.4 kb, at least about 2.6 kb, at least about 2.8 kb, at least about 3.0 kb, at least about 3.2 kb, at least about 3.4 kb, at least about 3.5 kb in length, at least about 4.0 kb in length, at least about 5.0 kb in length, at least about 6.0 kb in length, at least about 7.0 kb in length, at least about 8.0 kb in length, at least about 9.0 kb in length, or at least about 10.0 kb in length. In some embodiments, the nucleotide is at least about 1.4 kb in length.

Production of Heterologous Proteins In Vivo

[00197] The recombinant AAV disclosed herein can be used to produce a protein of interest *in vivo*, for example in an animal such as a mammal. Some embodiments provide a method for producing a protein of interest *in vivo*, where the method includes providing a recombinant AAV comprising a nucleotide sequence encoding the protein of interest; and administering the recombinant AAV to the subject, whereby the recombinant AAV expresses the protein of interest in the subject. The subject can be, in some embodiments, a non-human mammal, for example, a monkey, a dog, a cat, a mouse, or a cow. The size of the nucleotide sequence encoding the protein of interest can vary. For example, the nucleotide sequence can be at least about 1.4 kb, at least about 1.5 kb, at least about 1.6 kb, at least about 1.7 kb, at least about 1.8 kb, at least about

2.0 kb, at least about 2.2 kb, at least about 2.4 kb, at least about 2.6 kb, at least about 2.8 kb, at least about 3.0 kb, at least about 3.2 kb, at least about 3.4 kb, at least about 3.5 kb in length, at least about 4.0 kb in length, at least about 5.0 kb in length, at least about 6.0 kb in length, at least about 7.0 kb in length, at least about 8.0 kb in length, at least about 9.0 kb in length, or at least about 10.0 kb in length. In some embodiments, the nucleotide is at least about 1.4 kb in length.

Therapeutic Uses

[00198] The recombinant AAV produced by the methods described can be used to express one or more therapeutic proteins to treat various diseases or disorders. Non-limiting examples of the diseases include cancer such as carcinoma, sarcoma, leukemia, lymphoma; and autoimmune diseases such as multiple sclerosis. Non-limiting examples of carcinomas include esophageal carcinoma; hepatocellular carcinoma; basal cell carcinoma, squamous cell carcinoma (various tissues); bladder carcinoma, including transitional cell carcinoma; bronchogenic carcinoma; colon carcinoma; colorectal carcinoma; gastric carcinoma; lung carcinoma, including small cell carcinoma and non-small cell carcinoma of the lung; adrenocortical carcinoma; thyroid carcinoma; pancreatic carcinoma; breast carcinoma; ovarian carcinoma; prostate carcinoma; adenocarcinoma; sweat gland carcinoma; sebaceous gland carcinoma; papillary carcinoma; papillary adenocarcinoma; cystadenocarcinoma; medullary carcinoma; renal cell carcinoma; ductal carcinoma in situ or bile duct carcinoma; choriocarcinoma; seminoma; embryonal carcinoma; Wilm's tumor; cervical carcinoma; uterine carcinoma; testicular carcinoma; osteogenic carcinoma; epithelial carcinoma; and nasopharyngeal carcinoma. Non-limiting examples of sarcomas include fibrosarcoma, myxosarcoma, liposarcoma, chondrosarcoma, chordoma, osteogenic sarcoma, osteosarcoma, angiosarcoma, endotheliosarcoma, lymphangiosarcoma, lymphangioendotheliosarcoma, synovioma, mesothelioma, Ewing's sarcoma, leiomyosarcoma, rhabdomyosarcoma, and other soft tissue sarcomas. Non-limiting examples of solid tumors include glioma, astrocytoma, medulloblastoma, craniopharyngioma, ependymoma, pinealoma, hemangioblastoma, acoustic neuroma, oligodendroglioma, meningioma, melanoma, neuroblastoma, and retinoblastoma. Non-limiting examples of leukemias include chronic myeloproliferative syndromes; acute myelogenous leukemias; chronic lymphocytic leukemias, including B-cell CLL, T-cell CLL prolymphocytic leukemia, and hairy cell leukemia; and acute lymphoblastic leukemias. Examples of lymphomas include, but are not limited to, B-cell lymphomas, such as Burkitt's lymphoma; Hodgkin's lymphoma; and the like.

Other non-limiting examples of the diseases that can be treated using the AAV vectors, recombinant viruses and methods disclosed herein include genetic disorders including sickle cell anemia, cystic fibrosis, lysosomal acid lipase (LAL) deficiency 1, Tay-Sachs disease, Phenylketonuria, Mucopolysaccharidoses, Glycogen storage diseases (GSD, e.g., GSD types I, II, III, IV, V, VI, VII, VIII, IX, X, XI, XII, XIII, and XIV), Galactosemia, muscular dystrophy (e.g., Duchenne muscular dystrophy), and hemophilia such as hemophilia A (classic hemophilia) and hemophilia B (Christmas Disease).

[00199] The amount of the heterologous protein expressed in the subject (e.g., the serum of the subject) can vary. For example, in some embodiments the protein can be expressed in the serum of the subject in the amount of at least about 9 $\mu\text{g/ml}$, at least about 10 $\mu\text{g/ml}$, at least about 50 $\mu\text{g/ml}$, at least about 100 $\mu\text{g/ml}$, at least about 200 $\mu\text{g/ml}$, at least about 300 $\mu\text{g/ml}$, at least about 400 $\mu\text{g/ml}$, at least about 500 $\mu\text{g/ml}$, at least about 600 $\mu\text{g/ml}$, at least about 700 $\mu\text{g/ml}$, at least about 800 $\mu\text{g/ml}$, at least about 900 $\mu\text{g/ml}$, or at least about 1000 $\mu\text{g/ml}$. In some embodiments, the protein of interest is expressed in the serum of the subject in the amount of about 9 $\mu\text{g/ml}$, about 10 $\mu\text{g/ml}$, about 50 $\mu\text{g/ml}$, about 100 $\mu\text{g/ml}$, about 200 $\mu\text{g/ml}$, about 300 $\mu\text{g/ml}$, about 400 $\mu\text{g/ml}$, about 500 $\mu\text{g/ml}$, about 600 $\mu\text{g/ml}$, about 700 $\mu\text{g/ml}$, about 800 $\mu\text{g/ml}$, about 900 $\mu\text{g/ml}$, about 1000 $\mu\text{g/ml}$, about 1500 $\mu\text{g/ml}$, about 2000 $\mu\text{g/ml}$, about 2500 $\mu\text{g/ml}$, or a range between any two of these values. A skilled artisan will understand that the expression level in which a protein of interest is needed for the method to be effective can vary depending on non-limiting factors such as the particular protein of interest and the subject receiving the treatment, and an effective amount of the protein can be readily determined by a skilled artisan using conventional methods known in the art without undue experimentation.

EXAMPLES

Example 1

Isolation of Novel Naturally-Occurring Capsid Proteins

[00200] Novel naturally-occurring capsid proteins were isolated from the liver tissue from various mammals. Fresh liver tissue (porcine) was obtained from local farmers. Frozen liver tissue (baboon, cynomolgous macaque, marmoset, and crab-eating macaque) was obtained from

Texas Biomedical or the New England Primate Research Center. Genomic DNA was prepared from liver tissue using the DNeasy Blood & Tissue kit (Qiagen catalog #69504).

[00201] Polymerase chain reaction (PCR) was carried out on the genomic DNA using the following primers: primer rep-1397-F (5'-GTGCCCTTTTACGGCTGCGTGAAGTGGACCAATGAAAACCTTTCC- 3' SEQ ID NO:158) and primer cap-2872-R (5'-CCGACGGAGTGGGCAATGCCTCAGGAAATTGGCATTGCGATTCC- 3' SEQ ID NO:159) under the following conditions: initial incubation: 97°C, 120 sec, denaturation step: 97°C, 15 sec, annealing step: 58°C, 60°C, or 62°C, 15 sec, extension step: 72°C, 240 sec. The denaturation, annealing, and extension steps were performed for 35 cycles. Then the reaction was incubated at 72°C, 7 min and stored at 4°C until analyzed. The PCR products were separated by electrophoresis on 1% agarose gels, isolated using the Gel Extraction Kit (Qiagen catalog #28704), and cloned into pCR4-TOPO-TA (Invitrogen catalog #450030) according to the manufacturer's instructions. After transformation of *E. coli*, NEB5 α cells, DNA was prepared from ampicillin resistant colonies and sequenced from both ends to determine if the insert encoded an AAV-related sequence.

[00202] If the inserts in pCR4-TOPO TA were related to AAV sequences, sequence-specific primers were designed to the rep portion of the sequence to perform "around the episome PCR" (hereinafter "ATE PCR") to obtain a complete capsid gene. ATE PCR is based on the notion that persistent AAV genomes forms circular episomes in animal tissues. Accordingly, one can use a "divergent" set of primers corresponding to a sequence in the rep gene to perform polymerase chain reactions to isolate most or all of any AAV sequence that may exist in that episome but in particular one could isolate a complete contiguous capsid gene. Multimers of episomes can form, for example by homologous recombination, and in that case it is possible to isolate more than one capsid gene (which usually are not the same) from a single ATE PCR reaction.

[00203] An ATE PCR was carried-out in a standard polymerase chain reaction instrument using a 2-step program as follows: initial incubation: 95°C, 240 sec, denaturation step: 95°C, 30 sec annealing / extension step: 72°C, 300 sec. The denaturation and combined annealing / extension steps were performed for 40 cycles. The reaction was then incubated at 72°C, 7 min and stored at 4°C until analyzed. The PCR products were electrophoresed on 1% agarose gels.

PCR products that were the length of multimers of an AAV genome (~4.5 kilobases) were excised from the gel, purified using the QIAquick Gel Extraction Kit (Qiagen catalog #28704), and cloned into pCR4-TOPO-TA (Invitrogen catalog # 450030) according to the manufacturer's instructions. After transformation of *E. coli*, NEB5 α cells, DNA was prepared from ampicillin resistant colonies and the entire sequence of the insert was determined.

[00204] If the 2-step program described above did not produce PCR products of the correct size the following 3-step program was used: Initial incubation: 95°C, 240 sec, Denaturation step: 95°C, 30 sec Annealing step: 62°C, 64°C, 66°C, or 68°C, 30 sec, Extension step: 72°C, 300 sec. The denaturation, annealing, and extension steps were performed for 40 cycles. Then the reaction was incubated at 72°C, 7 min and stored at 4°C until analyzed as above.

[00205] Once complete insert sequences in pCR4-TOPO TA were determined they were identified as being AAV capsid genes using the BLAST algorithm (available at the NCBI website). Their relationship to known AAVs was determined using various nucleotide or amino acid sequence alignment programs such as Clustal Omega (available at the EBI web site) or Vector NTI (Invitrogen, Inc.).

[00206] To produce AAV, the unique AAV capsid genes were subcloned into an expression plasmid (pAAV-RC; Agilent, Inc.), then transfected into 293 cells along with a vector (pAAV luciferase) and adenovirus helper plasmid (pHELPER; Agilent, Inc.). AAV production was allowed to occur for 3 days and then crude lysates were made by freeze-thawing the cells three times. Debris was pelleted and the supernatant (crude AAV) was titered by Q-PCR to determine a genomic titer (which confirms the capsid is capable of assembly and DNA packaging) and then used to assess transduction by the AAVs on various cells.

[00207] The VP1 amino acid sequences of the novel mammalian tissue-derived AAV capsid proteins identified are herein described as SEQ ID NOS:15-89. The locations of the associated VP2 and VP3 regions are also herein described. The present invention is directed to (i) isolated AAV capsid proteins having at least 95%, 96%, 97%, 98% or 99% sequence identity to any of the VP1 capsid sequences of SEQ ID NOS:15-89, or the VP2 or VP3 regions of any of the capsid sequences of SEQ ID NOS:15-89, or (ii) isolated AAV capsid proteins comprising or consisting of any of the VP1 capsid sequences of SEQ ID NOS:15-89, or the VP2 or VP3 regions of any of the capsid sequences of SEQ ID NOS:15-89. The invention is also directed to

an AAV particle that comprises any of the above described AAV capsid proteins, wherein the AAV particle further comprises either (i) a nucleic acid having AAV inverted terminal repeats and a transgene comprising a heterologous gene operably linked to regulatory sequences that direct expression of the heterologous gene in a host cell, or (ii) a nucleic acid comprising a heterologous gene operably linked to regulatory sequences that control expression of the heterologous gene in a host cell.

Example 2

Generation of Engineered Chimeric Capsid Proteins

[00208] Alignment of the primary VP1 amino acid sequence of various AAV serotypes identified that conserved and variable regions exist in AAV capsid proteins. This analysis identified nine variable regions (denoted herein as VR I - VR IX) in the capsid protein amino acid sequence. In the present invention, one or more of the nine variable regions of a backbone ("recipient") capsid protein and/or the glycan binding sequence (GBS) region or GH loop region were substituted with the corresponding variable region(s), GBS or GH loop regions from a donor capsid protein having a different amino acid sequence than the recipient. The pHLP19-AAV-BMRNX expression plasmid vector was used to generate the chimeric capsid proteins. Rep proteins from AAV2 were used consistently for all different capsid proteins (VP1, VP2 and VP3). Engineered capsid genes were subcloned into pHLP19-AAV-BMRNX using unique *Swa*I and *Age*I restriction enzyme sites in the plasmid and that also flank each capsid gene.

[00209] Table 3 provides exemplary engineered chimeric capsid proteins that were generated as described here and tested as described in Example 3 below. "Backbone Sequence" refers to the backbone VP1 capsid sequence into which VR, GBS and/or GH loop region(s) are substituted or swapped, i.e., the "recipient." "Donor Sequence" refers to the VP1 capsid sequence from which the variable, GBS and/or GH loop region(s) are obtained and swapped into the recipient backbone sequence, i.e., the "donor." The "VR" substitution refers to a substitution where all nine variable regions (VR I – VR IX) from the donor are swapped into the recipient to produce the resulting engineered chimeric capsid. The "VRGBS" substitution refers to a substitution where all nine variable regions (VR I-VR IX) and the GBS sequence are swapped from the donor into the recipient to produce the resulting engineered chimeric capsid. The "GH"

substitution refers to a substitution where only the GH loop region is swapped from the donor into the recipient to produce the resulting engineered chimeric capsid. The “GH loop” sequence comprises variable regions VR IV through VR VIII, including the encompassed GBS sequence and all conserved sequence that is interspersed between those regions from the donor. The “VRGH” substitution refers to a substitution in which all nine variable regions (VR I - VR IX) and the GH loop sequence (including the encompassed GBS sequence and the conserved sequence that is interspersed between those regions from the donor sequence) are swapped from the donor to the recipient. The “PHP” modification refers to an insertion of the seven amino acid sequence TLAVPFK (SEQ ID NO: 160) into the chimeric capsid sequence to enhance targeting to brain tissue as described in Deverman et al., *Nature Biotechnology* 34:204-209 (2016).

Table 3:

Name/ SEQ ID NO:	Backbone Sequence	Donor Sequence	Sub- stitution	VP1 Amino Acid Sequence
AAVrh.10_9GH SEQ ID NO: 90	AAVrh.10	AAV9	GH	MAADGYLPDWLEDNLSEGIREWDLKPGAPKPKANQQKQD DGRGLVLPGYKYLGPFGNLDKGEFVNAAADAAALEHDKAYD QQLKAGDNPYLRYNHADADEFQERLQEDTSFGGNLGRAVFQ AKKRVLEPLGLVEEGAKTAPGKKRPVEPSPQSPDSSSTGI GKKGQQPAKKRLNFGQTGDSESVDPDPQPIGEPPAGPSGLG SGTMAAGGGAPMADNNEGADGVGSSSGNWHCDSTWLGDREV ITTSTRTWALPTYNNHLYKQISNGTSGGSTNDNTYFGYST PWGYFDFNRFHCHFSPRDWQRLINNNWGFPRKRLNFKLFN IQVKEVTQNEGKTIANNLTSTIQVFTDSEYQLPYVLGSA HQGCLPPFPADVFMIPQYGYLTLDNGSQAAGRSSFYCLEY FPSQMLRTGNNFEFSYQFEDVFPFHSSYAHSSQLDRLMNPL IDQYLYLSKTINGSGQNQQTTLKFSVAGPSNMAVQGRNYI PGPSYRQQRVSTTVTQNNSEFAWPGASSWALNGRNSLMN PGPAMASHKEGEDRFFPLSGSLIFGKQGTGRDNDVADKVM ITNEEEIKTTNPVATESYQGVATNHQSAQAQAQTGWVQNO GILPGMVWQNRDVYLQGPWAKIPHTDGNFHPSPLMGGFG LKHPPPQILIKNTPVPADPPTTFSQAKLASFITQYSTGQV SVEIEWELQKENSKRWNPEIQYTSNYYKSTNVDFAVNTDG TYSEPRPIGTRYLTRNL
AAVrh.10_9VRGBS SEQ ID NO: 91	AAVrh.10	AAV9	VRGBS	MAADGYLPDWLEDNLSEGIREWDLKPGAPKPKANQQKQD DGRGLVLPGYKYLGPFGNLDKGEFVNAAADAAALEHDKAYD QQLKAGDNPYLRYNHADADEFQERLQEDTSFGGNLGRAVFQ AKKRVLEPLGLVEEGAKTAPGKKRPVEPSPQSPDSSSTGI GKKGQQPAKKRLNFGQTGDSESVDPDPQPIGEPPAGPSGLG SGTMAAGGGAPMADNNEGADGVGSSSGNWHCDSTWLGDREV ITTSTRTWALPTYNNHLYKQISNSTSGGSTNDNTYFGYST PWGYFDFNRFHCHFSPRDWQRLINNNWGFPRKRLNFKLFN IQVKEVTQNEGKTIANNLTSTIQVFTDSEYQLPYVLGSA HQGCLPPFPADVFMIPQYGYLTLDNGSQAAGRSSFYCLEY FPSQMLRTGNNFEFSYQFEDVFPFHSSYAHSSQLDRLMNPL IDQYLYLSRTINGSGQNQQTTLKFSVAGPSNMAVQAKNWL PGPCYRQQRVSTTVTQNNSEFAWPGATKYHLNGRDSLNV PGVAMASHKEGEDRFFPLSGSLMFGKQGTGRDNDVADKVM LTSEEEIKTTNPVATEQYGVVATNHQSAQAQAQTGWVQNO

				GILPGMVWQNRDVYLQGP I WAKIPHTDGNFHPSPLMGGFG LKHPPPPQILIKNTPVPADPPTTFSQAKLASFITQYSTGQV SVEIEWELQKENS KRWNPEIQYTSNYYKSNNVEFAVNTEG TYSEPRPIGTRYLTRNL
AAVBce.36_9GH SEQ ID NO: 92	AAVBce.36	AAV9	GH	MAADGYLPDWLEDNLSEGIREWALKPGAPQPKANQQKQD DGRGLVLPGYKYLGPFGNLDKGEPVNEADAAALEHDKAYD QQLKAGDNPYLRYNHADAEEFQERLQEDTSFGGNLGRAVFQ AKKRVLEPLGLVEEAAKTAPGKKRPVDSPDSTSGIGKKGQ QPARKRLNFGQTGDAESVPDPQPIGEPPAAPSGLGSGTMA AGGGAPMADNNEGADGVGNASGNWHCDSTWLGNRVITTTST RTWALPTYNNHLYKQISSSSSGATNDNHYFGYSTPWGYFD VNRHFCHFSRDPWQRLINNNWGFPRKRLRFKLFNIQVKEV TTNDGVTTIANNLSTTVQVFSDEYQLPYVLGSAHQGCLP PPPADVFMIQYGYLTLNNGSQSVGRSSFYCLEYFPPSQML RTGNNFEFSYEFEDVPPHSSYAHSSQSLDRLMNP LIDQYLY YLSKTINGSGQNQQTLKF SVAGPSNMVQGRNYPGPYSYR QQRVSTTVTQNNSEFAWP GASSWALNGRNSLMNPGPAMA SHKEGEDRFFPLSGSLIFGKQGTGRDNVDADKVMITNEEE IKTNPVATESYQVATNHQSAQAQAQTGWVQNGGILPGM VWQNRDVYLQGP I WAKIPHTDGNFHPSPLMGGFGLKHPP QILIKNTPVPANPPETFTPAKFASFITQYSTGQVSVIEW ELQKENS KRWNPEIQYTSNYYDKQTGVDFAVDTQGVYSEPR PIGTRYLTRNL
AAVBce.36_9VRGBS SEQ ID NO: 93	AAVBce.36	AAV9	VRGBS	MAADGYLPDWLEDNLSEGIREWALKPGAPQPKANQQKQD DGRGLVLPGYKYLGPFGNLDKGEPVNEADAAALEHDKAYD QQLKAGDNPYLRYNHADAEEFQERLQEDTSFGGNLGRAVFQ AKKRVLEPLGLVEEAAKTAPGKKRPVDSPDSTSGIGKKGQ QPARKRLNFGQTGDAESVPDPQPIGEPPAAPSGLGSGTMA AGGGAPMADNNEGADGVGNASGNWHCDSTWLGNRVITTTST RTWALPTYNNHLYKQISNSTSGSSNDNHYFGYSTPWGYF DFNRHFCHFSRDPWQRLINNNWGFPRKRLRFKLFNIQVKEV VTDNNGVKTIANNLSTTVQVFSDEYQLPYVLGSAHQGCL PPPADVFMIQYGYLTLNDGSQAVGRSSFYCLEYFPPSQM LRTGNNFEFSYEFEDVPPHSSYAHSSQSLDRLMNP LIDQYLY YYLARTINGSGQNQQTLKF SVAGPSNMVQSKNWLPGPCF RQQRVSTTVTQNNSEFAWP GATKYHLNGRNSLTNPGVPM ASHKEGEDRFFPLSGSLVFGKQGTGRDNVDADKVLMTDEE EIKATNPVATEEYGVVATNHQSAQAQAQTGWVQNGGILPG MVWQNRDVYLQGP I WAKIPHTDGNFHPSPLMGGFGLKHPP PQILIKNTPVPANPPETFTPAKFASFITQYSTGQVSVIE WELQKENS KRWNPEIQYTSNYYKSNNVEFAVNTEGVYSEPR PIGTRYLTRNL
AAV7_9VRGBS SEQ ID NO: 94	AAV7	AAV9	VRGBS	MAADGYLPDWLEDNLSEGIREWDLKPGAPKPKANQQKQD NGRGLVLPGYKYLGPFGNLDKGEPVNAADAAALEHDKAYD QQLKAGDNPYLRYNHADAEEFQERLQEDTSFGGNLGRAVFQ AKKRVLEPLGLVEEAAKTAPAKKRPVESPQRSPDSSSTGI GKKGQQPARKRLNFGQTGDSSESVPDPQLGEPPAAPSSVG SGTVAAGGGAPMADNNEGADGVGNASGNWHCDSTWLGDRV ITTSTRTWALPTYNNHLYKQISNSTSGSSNDNTYFGYST PWGYFDNRHFCHFSRDPWQRLINNNWGFPRKRLRFKLFN IQVKEVTDNNGVKTIANNLSTTVQVFSDEYQLPYVLGSA HQGCLPPPADVFMIQYGYLTLNDGSQAVGRSSFYCLEY FPSQMLRTGNNFEFSYEFEDVPPHSSYAHSSQSLDRLMNP L IDQYLYLARTINGSGQNQQTLKF SVAGPSNMVQAKNWL PGPCFRQQRVSTTVTQNNSEFAWP GATKYHLNGRNSLVN PGVAMASHKEGEDRFFPLSGSLIFGKQGTGRDNVDADKLM TNEEEIRPTNPVATEEYGIVATNHQSAQAQAQTGWVQNGG ILPGMVWQNRDVYLQGP I WAKIPHTDGNFHPSPLMGGFGL KHPPPPQILIKNTPVPANPPEVFTPAKFASFITQYSTGQV VEIEWELQKENS KRWNPEIQYTSNYYKSNNVEFAVNTEGV YSEPRPIGTRYLTRNL
AAV7_9GH SEQ ID NO: 95	AAV7	AAV9	GH	MAADGYLPDWLEDNLSEGIREWDLKPGAPKPKANQQKQD NGRGLVLPGYKYLGPFGNLDKGEPVNAADAAALEHDKAYD

				<p>QQLKAGDNPYLRYNHADAEEFQERLQEDTSFGGNLGRAVFQ AKKRVLEPLGLVEEGAKTAPAKKRPVEQSPQSPDSSSTGI GKKQQPARKRLNFGQTGDSESVDPDPQLGEPPAAPSSVSG SGTVAAGGGAPMADNNEGADGVGNASGNWHCDSTWLGDRV ITTSTRTWALPTYNNHLYKQISSETAGSTNDNTYFGYSTP WGYFDNRFHCHFSRQRLINNNWGFPRKRLFKLFNI QVKEVTTNDGVTTIANNLSTIQVFSSEYQLPYVLGSAH QGCLPPFPADVFMIPQYGYLTLNNGSQSVGRSSFYCLEYF PSQMLRTGNNFEFSYSFEDVPPHSSYAHSQSLDRLMNPLI DQYLYLSKTINGSGQNQQLKFSVAGPSNMAVQGRNYIP GPSYRQQRVSTTVTQNNNEFAWPAGASSWALNGRNSLMNP GPAMASHKEGEDRFFPLSGSLIFGKQGTGRDNVDADKVM TNEEEIKTTPNVATESYQVATNHQSAQAQAQTGWVQNG ILPGMVWQNRDVYLQGPWAKIPHTDGNFHPSPLMGGFGL KHPPQILIKNTVPANPPEVFTPAKFASFITQYSTGQVS VEIEWELQKENSKRWNPEIQYTSNFEKQGTGVDFAVDSQGV YSEPRPIGTRYLTRNL</p>
AAV6.2_9VRGBS SEQ ID NO: 96	AAV6.2	AAV9	VRGBS	<p>MAADGYLPDWLEDNLSEGIREWDLKPGAPKPKANQQKQD DGRGLVLPYGYLGPFGNLDKGEFVNAADAAAEHDKAYD QQLKAGDNPYLRYNHADAEEFQERLQEDTSFGGNLGRAVFQ AKKRVLEPLGLVEEGAKTAPGKKRPVEQSPQEPDSSSGIG KTGQQPAKKRLNFGQTGDSESVDPDPQLGEPPATPAAVGP TTMASGGGAPMADNNEGADGVGNASGNWHCDSTWLGDRVI TTSTRTWALPTYNNHLYKQISNSTSGGSSNDNHYFGYSTP WGYFDNRFHCHFSRQRLINNNWGFPRKRLNFKLFNI QVKEVTDNNGVKTIANNLSTVQVFSSEYQLPYVLGSAH QGCLPPFPADVFMIPQYGYLTLNNGSQAVGRSSFYCLEYF PSQMLRTGNNFTFSYTFEDVPPHSSYAHSQSLDRLMNPLI DQYLYLNRTINGSGQNQQLKFSVAGPSNMAVQPKNWLP GPCYRQQRVSTTVTQNNNEFAWPAGASKYNLNGRESIINP GTAMASHKEGEDRFFPLSGSMIFGKQGTGRDNVDADKVM TDEEEIKATNPVATERFGTVATNHQSAQAQAQTGWVQNG ILPGMVWQDRDVYLQGPWAKIPHTDGHFHPSPLMGGFGL KHPPQILIKNTVPANPPAEFSATKFASFITQYSTGQVS VEIEWELQKENSKRWNPEVQYTSNYKSNNEVEAVNTEGL YTEPRPIGTRYLTRPL</p>
AAV6.2_9GH SEQ ID NO: 97	AAV6.2	AAV9	GH	<p>MAADGYLPDWLEDNLSEGIREWDLKPGAPKPKANQQKQD DGRGLVLPYGYLGPFGNLDKGEFVNAADAAAEHDKAYD QQLKAGDNPYLRYNHADAEEFQERLQEDTSFGGNLGRAVFQ AKKRVLEPLGLVEEGAKTAPGKKRPVEQSPQEPDSSSGIG KTGQQPAKKRLNFGQTGDSESVDPDPQLGEPPATPAAVGP TTMASGGGAPMADNNEGADGVGNASGNWHCDSTWLGDRVI TTSTRTWALPTYNNHLYKQISSASTGASNDNHYFGYSTPW GYFDNRFHCHFSRQRLINNNWGFPRKRLNFKLFNIQ VKEVTTNDGVTTIANNLSTVQVFSSEYQLPYVLGSAHQ GCLPPFPADVFMIPQYGYLTLNNGSQAVGRSSFYCLEYFP SQMLRTGNNFTFSYTFEDVPPHSSYAHSQSLDRLMNPLID QYLYLSKTINGSGQNQQLKFSVAGPSNMAVQGRNYIPG PSYRQQRVSTTVTQNNNEFAWPAGASSWALNGRNSLMNPG PAMASHKEGEDRFFPLSGSLIFGKQGTGRDNVDADKVMIT NEEEIKTTPNVATESYQVATNHQSAQAQAQTGWVQNGGI LPGMVWQDRDVYLQGPWAKIPHTDGHFHPSPLMGGFGLK HPPQILIKNTVPANPPAEFSATKFASFITQYSTGQVSV EIEWELQKENSKRWNPEVQYTSNYAKSANVDFTVDNNGLY TEPRPIGTRYLTRPL</p>
AAVmo_2VR SEQ ID NO: 98	AAVmo	AAV2	VR	<p>MSFFDWLGKQYAQGAEEFWDLKGPPAPKKARKDGSAGFN FPGHKYLGPGNSLDRGDPVDADAAAQKHDQSYQEQLLEAG DNPYLKYNHADREFQEALKDDTSFEGNLARGLEFAKKLVA EPLGLVEPELAPPSGRKRPVQSSQESGYSSSQDKRPNLDV DEEDREFAAAAAETETGSAPPTGNLGPMTMAGGGSAPIDD GSYGADGVGNASGDWHCDSTWLDNCVITRTTRTNLPTYN NHIYKRLSSQSGASNDNHSYFGFSTPWGYFDNRFHCHFS PRDWQRLINNNWGLRPKSLRFKIFNIQVKEVTQNDGTTTI</p>

				SNNLTSTVQVFADTEYQLPYVIGSAHEGCLPPFPADVFM LPQYGYCTLNNGSQPTPRSAFYCLEYFPPSKMLRTGNSFEFT YNEFEKVPFHSMAHNSQLDRLMNPLIDQYLYYLDVTNTPS GTTTQSRFTYQKGVHTNLPQERNWLPGPGRNQVRVSKTS ADNNNSEYSWQYSNKYVLENGRASKIAPGPAMASHKDDEE KFFPQSGVLIFGKQGSEKTNVDIEKVMITRETEINSTNPL AGGSLGAVSTNLQRGNRQAALDHTNVMGVFPGSVWQDRDI YLQGGIWAQIPHTDGHFHPSPLMGGFGLKNPPQILIKNT PVPADPPTEFNANKISSFITQYSTGQVTVEMEWELQKETS KRWNPFIQYTSNYNKSVMVHFAPDDVGNYKEFRSIGTRYL TRPL
AAVmo_2VRGH SEQ ID NO: 99	AAVmo	AAV2	VRGH	MSFFDWLGKQYAQGAAEFWDLKSGPPAPKKARKDGSAGFN FPGHKYLGPGNSLDRGDPVDADAAAQKHDQSYQEQLLEAG DNPYLKYNHADREFQEALKDDTSFEGNLARGLFEAKKLVA EPLGLVEPELAPPSGRKRVPQSSQESGYSSSQDKRPNLDV DEEDREFAAAAAETETGSAPPTGNLPGMTMAGGGSAPIDD GSYGADGVGNASGDWHCDSTWLDNCVITRTTRTNWNLPTYN NHIYKRLSSQSGASNDNHSYFGFSTPWGYFDFNRFHCHF PRDWQRLINNNWGLRPKSLRFKIFNIQVKEVTQNDGTTTI SNNLTSTVQVFADTEYQLPYVIGSAHEGCLPPFPADVFM LPQYGYCTLNNGSQPTPRSAFYCLEYFPPSKMLRTGNSFEFT YNEFEKVPFHSMAHNSQLDRLMNPLIDQYLYYLSRTNTPS GTTTQSRQFSQAGASDIRDQSRNWLPGPCYRQQRVSKTS ADNNNSEYSWTGATKYHLNGRDSLVPNGPAMASHKDDEEK FFPQSGVLIFGKQGSEKTNVDIEKVMITDEEERTTNPVA TEQYGSVSTNLQRGNRQAATADVNTQGVFPGSVWQDRDIY LQGGIWAQIPHTDGHFHPSPLMGGFGLKNPPQILIKNT VPADPPTEFNANKISSFITQYSTGQVTVEMEWELQKETS KRWNPFIQYTSNYNKSVMVHFAPDDVGNYKEFRSIGTRYLT RPL
AAVmo_9VR SEQ ID NO: 100	AAVmo	AAV9	VR	MSFFDWLGKQYAQGAAEFWDLKSGPPAPKKARKDGSAGFN FPGHKYLGPGNSLDRGDPVDADAAAQKHDQSYQEQLLEAG DNPYLKYNHADREFQEALKDDTSFEGNLARGLFEAKKLVA EPLGLVEPELAPPSGRKRVPQSSQESGYSSSQDKRPNLDV DEEDREFAAAAAETETGSAPPTGNLPGMTMAGGGSAPIDD GSYGADGVGNASGDWHCDSTWLDNCVITRTTRTNWNLPTYN NHIYKQISNSTSGSSNDNAYFGFSTPWGYFDFNRFHCHF SPRDWQRLINNNWGLRPKSLRFKIFNIQVKEVTDNNGVKI ISNNLTSTVQVFADTEYQLPYVIGSAHEGCLPPFPADVFM LPQYGYLTLNDGSQAVGRSSFYCLEYFPPSKMLRTGNSFEF TYNEFEKVPFHSMAHNSQLDRLMNPLIDQYLYYLSKTING SGQNQQTLLKFSVAGPSNMAVQGRNWLPGPGRQQRVSTTV TQNNSEFAWPAGSSWALNGRASKIAPGPAMASHKEGEDR FFPLSGSLIFGKQGTGRDNVDADKVNI TRETEINSTNPLA GGSLGQVATNHQSAQAQQTGWVQNGQVFPGSVWQDRDIY LQGGIWAQIPHTDGHFHPSPLMGGFGLKNPPQILIKNT VPADPPTEFNANKISSFITQYSTGQVTVEMEWELQKETS KRWNPFIQYTSNYKSNNEFAVNTGEGVYKEFRSIGTRYLT RPL
AAVmo_9GH SEQ ID NO: 101	AAVmo	AAV9	GH	MSFFDWLGKQYAQGAAEFWDLKSGPPAPKKARKDGSAGFN FPGHKYLGPGNSLDRGDPVDADAAAQKHDQSYQEQLLEAG DNPYLKYNHADREFQEALKDDTSFEGNLARGLFEAKKLVA EPLGLVEPELAPPSGRKRVPQSSQESGYSSSQDKRPNLDV DEEDREFAAAAAETETGSAPPTGNLPGMTMAGGGSAPIDD GSYGADGVGNASGDWHCDSTWLDNCVITRTTRTNWNLPTYN NHIYKRLNGTTSQSYFGFSTPWGYFDFNRFHCHFSPRD WQRLINNNWGLRPKSLRFKIFNIQVKEVTQDSTKIIISNN LTSTVQVFADTEYQLPYVIGSAHEGCLPPFPADVFM LPQYGYCTLNNGSQPTPRSAFYCLEYFPPSKMLRTGNSFEFT GYCTRQDGNSSNPPTPRSAFYCLEYFPPSKMLRTGNSFEFT YNEFEKVPFHSMAHNSQLDRLMNPLIDQYLYYLSKTINGSG QNQQTLLKFSVAGPSNMAVQGRNYPGPYRQQRVSTTVTQ NNNSEFAWPAGSSWALNGRNSLMNPGPAMASHKEGEDRFF PLSGSLIFGKQGTGRDNVDADKVMI TNEEEIKTTNPVATE

				SYGQVATNHQSAQAQAQTGWVQNQGVFPGSVWQDRDIYLO GQIWAKIPHTDGHFHPSPLMGGFGLKNPPPQILIKNTPVP ADPPTEFNANKISSFITQYSTGQVTVEMEWELQKETS NPEIQYSDSSSTSGSILHFAPDVGVNYKEFRSIGTRYLT RPL
AAVmo_9VRGH SEQ ID NO: 102	AAVmo	AAV9	VRGH	MSFFDWLGKQYAQAAGAEFWDLKS GPPAPKKARKDGSAGFN FPGHKYLGPGNSLDRGDPVDADAAAQKHDQSYQEQL EAG DNPYLKYNHADREFQEALKDDTSFEGNLARGLFEAKKLVA EPLGLVEPELAPP SGRKRPVQSSQESGYSSSQDKRP NLDV DEEDREFAAAAAETETGSAPPTGNLPGTMAGGGSAP ID GSYGADGVGNASGDWHCDSTWLDNCVITRTTRTWNLP TYN NHIYKQISNSTSGSSNDNAYFGFSTPWGYFDFNRHCH F SPRDWQRLINNNWGLRPKSLRFKIFNIQVKEVTDNNGVKI ISNNLTSTVQVFADTEYQLPYVIGSAHEGCLPPFPADVFM LPQYGYLTLDNGSQAVGRSSFYCLEYFPSKMLRTGNSFEF TYNFEKVPFHSMAHNNQSLDRLMNPLIDQYLYLSKT ING SGQNQQTLKF SVAGPSNMAVQGRNYPGP SYRQQRVSTTV TQNNNSEFAWPGASSWALNGRNSLMNPGPAMASHKEGEDR FFPLSGSLIFGKQGTGRDNVDADKVMITNEEEIKTTNPVA TESYGQVATNHQSAQAQAQTGWVQNQGVFPGSVWQDRDIY LQGGIWAKIPHTDGHFHPSPLMGGFGLKNPPPQILIKNT P VPADPPTEFNANKISSFITQYSTGQVTVEMEWELQKETS RWNPEIQYTSNYYKSNNVEFAVNTGEGVYKEFRSIGTRYLT RPL
AAVmo_8VR SEQ ID NO: 103	AAVmo	AAV8	VR	MSFFDWLGKQYAQAAGAEFWDLKS GPPAPKKARKDGSAGFN FPGHKYLGPGNSLDRGDPVDADAAAQKHDQSYQEQL EAG DNPYLKYNHADREFQEALKDDTSFEGNLARGLFEAKKLVA EPLGLVEPELAPP SGRKRPVQSSQESGYSSSQDKRP NLDV DEEDREFAAAAAETETGSAPPTGNLPGTMAGGGSAP ID GSYGADGVGNASGDWHCDSTWLDNCVITRTTRTWNLP TYN NHIYKQISNGTSGGATNDNTYFGFSTPWGYFDFNRHCH F SPRDWQRLINNNWGLRPKSLRFKIFNIQVKEVTQNEGTKI ISNNLTSTVQVFADTEYQLPYVIGSAHEGCLPPFPADVFM LPQYGYLTLDNGSQAVGRSSFYCLEYFPSKMLRTGNSFEF TYNFEKVPFHSMAHNNQSLDRLMNPLIDQYLYLSRTQT T GGTANTQTLGFSQGGPNTMANQAKNWLP GPGIRQQRVSTT TGQNNNSNFAWTAGTKYHLNGRASKIAPGPAMATHKDD E RFFPSNGILIFGKQNAARDNADYSVNITRETEINSTNPL AGGSLGIVADNLQQNTAPQIGTVNSQGVFPGSVWQDRDIY LQGGIWAKIPHTDGHFHPSPLMGGFGLKNPPPQILIKNT P VPADPPTEFNANKISSFITQYSTGQVTVEMEWELQKETS KRWNP EIQYTSNYYKSTSVDFAVNTGEGVYKEFRSIGTRYL TRPL
AAVmo_8GH SEQ ID NO: 104	AAVmo	AAV8	GH	MSFFDWLGKQYAQAAGAEFWDLKS GPPAPKKARKDGSAGFN FPGHKYLGPGNSLDRGDPVDADAAAQKHDQSYQEQL EAG DNPYLKYNHADREFQEALKDDTSFEGNLARGLFEAKKLVA EPLGLVEPELAPP SGRKRPVQSSQESGYSSSQDKRP NLDV DEEDREFAAAAAETETGSAPPTGNLPGTMAGGGSAP ID GSYGADGVGNASGDWHCDSTWLDNCVITRTTRTWNLP TYN NHIYKRLNGTTSGDQSYFGFSTPWGYFDFNRHCH FSPRD WQRLINNNWGLRPKSLRFKIFNIQVKEVTQDSTKII SNN LTSTVQVFADTEYQLPYVIGSAHEGCLPPFPADVFM LPOY GYCTRQDGNSSNNTPRSFYCLEYFPSKMLRTGNSFEFTY NFEKVPFHSMAHNNQSLDRLMNPLIDQYLYLSRTQT TGG TANTQTLGFSQGGPNTMANQAKNWLP GPCYRQQRVSTTGG QNNNSNFAWTAGTKYHLNGRNSLANPGIAMATHKDD EERF FPSNGILIFGKQNAARDNADYSVMLTSEEEIKTTNPVAT EEYGIVADNLQQNTAPQIGTVNSQGVFPGSVWQDRDIY L QGGIWAKIPHTDGHFHPSPLMGGFGLKNPPPQILIKNT P PADPPTEFNANKISSFITQYSTGQVTVEMEWELQKETS KRWNP EIQYSDSSSTSGSILHFAPDVGVNYKEFRSIGTRYL TRPL
AAVmo_8VRGH	AAVmo	AAV8	VRGH	MSFFDWLGKQYAQAAGAEFWDLKS GPPAPKKARKDGSAGFN

SEQ ID NO: 105				FPGHKYLGPGNSLDRGDPVDADDAQAQKHDQSYQEQLQLEAG DNPYLKYNHADREFQEALKDDTSFEGNLARGLFEAKKLVA EPLGLVEPELAPP SGRKRPVQSSQESGYSSSQDKRPNLDV DEEDREFAAAAAETETGSAPPTGNLPGPTMAGGGSAPIDD GSYGADGVGNASGDWHCDSTWLDNCVITRTTRTWNLP TYN NHIYKQISNGTSGGATNDNTYFGFSTPWGYFDFNRFHCHF SPRDWQRLINNNWGLRPKSLRFKIFNIQVKEVTQNEGTKI ISNNLTSTVQVFADTEYQLPYVIGSAHEGCLPPFPADVFM LPQYGYLTLNNGSQAVGRSSFYCLEYFPSKMLRTGNSFEF TYNFEKVPFHSMAHNSQLDRLMNLIDQYLYLSRTQTT GGTANTQTLGFSQGGPNTMANQAKNWLPGPCYRQQRVSTT TGQNNNSNFAWTAGTKYHLNGRNSLANPGIAMATHKDDDEE RFFPSNGILIFGKQNAARDNADYSDVMLTSEEEIKTTNPV ATEEYGI VADNLQQONTAPQIGTVNSQGVFPGSVWQDRDI YLQQGIWAKIPHTDGHFHP SPLMGFGFLKNPPQILIKNT PVPADPPTEFNANKISSFITQYSTGQVTMEMEWELQKETS KRWNPEIQYTSNYKSTSVDFAVNTEGVYKEFRSIGTRYL TRPL
AAV4_2VR SEQ ID NO: 106	AAV4	AAV2	VR	MTDGYLPDWLEDNLSEGVREWWALQPGAPKPKANQQHQDN ARGVLVPGYKYLPGNGLDKGEPVNAADAAALEHDKAYDQ QLKAGDNPYLKYNHADAEEFQQLQGDTSFGGNLGRAVFQA KKRVLLEPLGLVEQAGETAPGKKRPLIESPQQPDSSTGIGK KGKQPAKKKLVFEDETGAGDGPPEGSTSGAMSDDSEMRAA AGGAAVEGGQAGADGVGNASGDWHCDSTWSEGHVTTTSTR WVLPTYNNHLYKQISSQSGASNDNHYFGFSTPWGYFDFNR FHCHFSRDPWQRLINNNWGMRPKAMRVKIFNIQVKEVTQ DGTTVANNLTSTVQIFADSSYELPYVMDAGQEGSLPPFP NDVFMVPPQYGYLTLNNGSQAVGRSSFYCLEYFPSQMLRTG NNFEITYSFEKVPFHSMAHNSQLDRLMNLIDQYLYLS RTNTPSGTTTQSRQLQFSQAGASDIRDQSRNWLPGPSIRQQ RVSKTSADNNNSEYSWTGATKYHLNGRWSALTGPMPATH KDDEEKFFPQSGVLIFGKQGSEKTNVDIEKLIFTSEEEELA ATNATDTDMWGSVSTNLQRGNRQAATADVNTQGA VPGMVW QNRDIYYQGP I WAKIPHTDGHFHP SPLIGGFLKHPPPPQI FIKNTPVPANPATTFSSTPVNSFITQYSTGQVSVQIDWEI QKERSKRWNPEVQFTSNYKSVNVDFTVDTNGVYTEPRAI GTRYLTHHL
AAV4_2GH SEQ ID NO: 107	AAV4	AAV2	GH	MTDGYLPDWLEDNLSEGVREWWALQPGAPKPKANQQHQDN ARGVLVPGYKYLPGNGLDKGEPVNAADAAALEHDKAYDQ QLKAGDNPYLKYNHADAEEFQQLQGDTSFGGNLGRAVFQA KKRVLLEPLGLVEQAGETAPGKKRPLIESPQQPDSSTGIGK KGKQPAKKKLVFEDETGAGDGPPEGSTSGAMSDDSEMRAA AGGAAVEGGQAGADGVGNASGDWHCDSTWSEGHVTTTSTR WVLPTYNNHLYKRLGESLQSN TYNGFSTPWGYFDFNRFHC HFSPRDWQRLINNNWGMRPKAMRVKIFNIQVKEVTTSNGE TTVANNLTSTVQIFADSSYELPYVMDAGQEGSLPPFPNDV FMVPQYGYCGLVTGNTSQQOTDRNAFYCLEYFPSQMLRTG NNFEITYSFEKVPFHSMAHNSQLDRLMNLIDQYLYLS RTNTPSGTTTQSRQLQFSQAGASDIRDQSRNWLPGPSIRQQ RVSKTSADNNNSEYSWTGATKYHLNGRSLVNP GPAMASH KDDEEKFFPQSGVLIFGKQGSEKTNVDIEKVMITDEEEIR TTNPVATEQYGSVSTNLQRGNRQAATADVNTQGA VPGMVW QNRDIYYQGP I WAKIPHTDGHFHP SPLIGGFLKHPPPPQI FIKNTPVPANPATTFSSTPVNSFITQYSTGQVSVQIDWEI QKERSKRWNPEVQFTSNYQQNSLLWAPDAAGKYTEPRAI GTRYLTHHL
AAV4_2VRGH SEQ ID NO: 108	AAV4	AAV2	VRGH	MTDGYLPDWLEDNLSEGVREWWALQPGAPKPKANQQHQDN ARGVLVPGYKYLPGNGLDKGEPVNAADAAALEHDKAYDQ QLKAGDNPYLKYNHADAEEFQQLQGDTSFGGNLGRAVFQA KKRVLLEPLGLVEQAGETAPGKKRPLIESPQQPDSSTGIGK KGKQPAKKKLVFEDETGAGDGPPEGSTSGAMSDDSEMRAA AGGAAVEGGQAGADGVGNASGDWHCDSTWSEGHVTTTSTR WVLPTYNNHLYKQISSQSGASNDNHYFGFSTPWGYFDFNR

				FHCHFSPRDWQRLINNNWGMRPKAMRVKIFNIQVKEVTQN DGTTVANNLTSTVQIFADSSYELPYVMDAGQEGSLPPFP NDVFMVPQYGYLTLNNGSQAVGRSSFYCLEYFPSQMLRTG NNFEITYSFEKVPFHSYAHSQSLDRLMNPLIDQYLYYLS RTNTPSGTTTQSRLOFSQAGASDIRDQSRNWLPGPCYRQQ RVSKTSADNNNSEYSWTGATKYHLNGRDSLVPNGPAMASH KDDEEKFFPQSGVLIFGKQGSEKTNVDIEKVMITDEEEIR TTNPVATEQYGSVSTNLQRGNRQAATADVNTQGAVPGMVW QNRDIYYQGPIWAKIPHDTGHHFSPPLIGGFGLKHPPPQI FIKNTVPANPATTFSSTPVNSFITQYSTGQVSVQIDWEI QKERSKRWNPEVQFTSNYKSVNVDFTVDTNGVYTEPRAI GTRYLTHHL
AAV4_8VR SEQ ID NO: 109	AAV4	AAV8	VR	MTDGYLPDWLEDNLSEGVREWWALQPGAPKPKANQQHQDN ARGLVLPGYKYLPGPNGLDKGEPVNAADAAALEHDKAYDQ QLKAGDNPYLYKNHADAEEFQQRLQGDTSFGGNGRAVFA KKRVLEPLGLVEQAGETAPGKKRPLIESPQQPDSSTGIGK KGKQPAKKKLVEFEDETGAGDGPPEGSTSGAMSDSEMRAA AGGAAVEGGQGADGVGNASGDWHCDSTWSEGHVTTTSTR WVLPYNNHLYKQISNGTSGGATNDNTYFGFSTPWGYFDF NRFHCHFSPRDWQRLINNNWGMRPKAMRVKIFNIQVKEVT QNEGKTVANNLTSTVQIFADSSYELPYVMDAGQEGSLPP FPNDVFMVPQYGYLTLNNGSQAVGRSSFYCLEYFPSQMLR TGNNEITYSFEKVPFHSYAHSQSLDRLMNPLIDQYLYY LSRTQTTGGTANTQTLGFSQGGPNTMANQAKNWLPGPSIR QQRVSTTTGQNNNSNFAWTAGTKYHLNGRWSALTPGPMA THKDDEERFFPSNGILIFGKQNAARDNADYSDLIFTSEEE LAATNATDTDMWGIADNLQQQNTAPQIGTVNSQGAVPGM VWQNRDIYYQGPIWAKIPHDTGHHFSPPLIGGFGLKHPPP QIFIKNTVPANPATTFSSTPVNSFITQYSTGQVSVQIDW EIQKERSKRWNPEVQFTSNYKSTSVDFAVNTEGVYTEPR AIGTRYLTHHL
AAV4_8GH SEQ ID NO: 110	AAV4	AAV8	GH	MTDGYLPDWLEDNLSEGVREWWALQPGAPKPKANQQHQDN ARGLVLPGYKYLPGPNGLDKGEPVNAADAAALEHDKAYDQ QLKAGDNPYLYKNHADAEEFQQRLQGDTSFGGNGRAVFA KKRVLEPLGLVEQAGETAPGKKRPLIESPQQPDSSTGIGK KGKQPAKKKLVEFEDETGAGDGPPEGSTSGAMSDSEMRAA AGGAAVEGGQGADGVGNASGDWHCDSTWSEGHVTTTSTR WVLPYNNHLYKRLGESLQSNYNGFSTPWGYFDFNRFHC HFSRPDWQRLINNNWGMRPKAMRVKIFNIQVKEVTTSNGE TTVANNLTSTVQIFADSSYELPYVMDAGQEGSLPPFPNDV FMVPQYGYCGLVTGNTSQQTDRNAFYCLEYFPSQMLRTG NNFEITYSFEKVPFHSYAHSQSLDRLMNPLIDQYLYYLS RTQTTGGTANTQTLGFSQGGPNTMANQAKNWLPGPCYRQQ RVSTTTGQNNNSNFAWTAGTKYHLNGRNSLANPGIAMATH KDDEERFFPSNGILIFGKQNAARDNADYSVMLTSEEEIK TTNPVATEEYGIADNLQQQNTAPQIGTVNSQGAVPGMVW QNRDIYYQGPIWAKIPHDTGHHFSPPLIGGFGLKHPPPQI FIKNTVPANPATTFSSTPVNSFITQYSTGQVSVQIDWEI QKERSKRWNPEVQFTSNYQQNSLLWAPDAAGKYTEPRAI GTRYLTHHL
AAV4_8VRGH SEQ ID NO: 111	AAV4	AAV8	VRGH	MTDGYLPDWLEDNLSEGVREWWALQPGAPKPKANQQHQDN ARGLVLPGYKYLPGPNGLDKGEPVNAADAAALEHDKAYDQ QLKAGDNPYLYKNHADAEEFQQRLQGDTSFGGNGRAVFA KKRVLEPLGLVEQAGETAPGKKRPLIESPQQPDSSTGIGK KGKQPAKKKLVEFEDETGAGDGPPEGSTSGAMSDSEMRAA AGGAAVEGGQGADGVGNASGDWHCDSTWSEGHVTTTSTR WVLPYNNHLYKQISNGTSGGATNDNTYFGFSTPWGYFDF NRFHCHFSPRDWQRLINNNWGMRPKAMRVKIFNIQVKEVT QNEGKTVANNLTSTVQIFADSSYELPYVMDAGQEGSLPP FPNDVFMVPQYGYLTLNNGSQAVGRSSFYCLEYFPSQMLR TGNNEITYSFEKVPFHSYAHSQSLDRLMNPLIDQYLYY LSRTQTTGGTANTQTLGFSQGGPNTMANQAKNWLPGPCYR QQRVSTTTGQNNNSNFAWTAGTKYHLNGRNSLANPGIAMA

				THKDDEERFFPSNGILIFGKQNAARDNADYSDVMLTSEEE IKTTNPVATEEYGIVADNLQQQNTAPQIGTVNSQGAVPGM VWQNRDIYYQGPIWAKIPHTDGHFHPSPPLIGGFGLKHPPP QIFIKNTPVPANPATTFSSSTPVNSFITQYSTGQVSVQIDW EIQKERSKRWNPEVQFTSNYYKSTSVDFAVNTGCVYTEPR AIGTRYLTHHL
AAV4_9VR SEQ ID NO: 112	AAV4	AAV9	VR	MTDGYLPDWLEDNLSEGVREWWALQPGAPKPKANQQHQDN ARGLVLPGYKYLGPNGLDKGEPVNAADAAALEHDKAYDQ QLKAGDNPYLKYNHADAEEFQRLQGDTSFGGNLGRAVFQA KKRVLEPLGLVEQAGETAPGKKRPLIESPQQPDSSTGIGK KGKQPAKKKLVFEDETGAGDGPPEGSTSGAMSDDSEMR AGGAAVEGGQGADGVGNASGDWHCDSTWSEGHVTTTSTR WVLPITYNNHLYKQISNSTSGGSSNDNAYFGFSTPWGYFDF NRFHCHFSRDPWQRLINNNWGMRPKAMRVKIFNIQVKEVT DNNGVKTVANNLTSTVQIFADSSYELPYVMDAGQEGSLPP FPNDVFMVPQYGYLTLNDGSQAVGRSSFYCLEYFPSQMLR TGNNFEITYSFEKVPFHSYAHSSQLDRLMNPIDQYLYY LSKTINGSGQNQQTLKFSVAGPSNMAVQGRNWLPGPSIRQ QRVSTTVTQNNNEFAWPGASSWALNGRWSALTPGPPMAT HKEGEDRFFPLSGSLIFGKQGTGRDNVDADKLIFTSEEL AATNATDMDWGQVATNHQSAQAQAQTGWVQNGAVPGMV WQNRDIYYQGPIWAKIPHTDGHFHPSPPLIGGFGLKHPPPQ IFIKNTPVPANPATTFSSSTPVNSFITQYSTGQVSVQIDWE IQKERSKRWNPEVQFTSNYYKSTSVDFAVNTGCVYTEPRA IGTRYLTHHL
AAV4_9GH SEQ ID NO: 113	AAV4	AAV9	GH	MTDGYLPDWLEDNLSEGVREWWALQPGAPKPKANQQHQDN ARGLVLPGYKYLGPNGLDKGEPVNAADAAALEHDKAYDQ QLKAGDNPYLKYNHADAEEFQRLQGDTSFGGNLGRAVFQA KKRVLEPLGLVEQAGETAPGKKRPLIESPQQPDSSTGIGK KGKQPAKKKLVFEDETGAGDGPPEGSTSGAMSDDSEMR AGGAAVEGGQGADGVGNASGDWHCDSTWSEGHVTTTSTR WVLPITYNNHLYKRLGESLQSNITYNGFSTPWGYFDFNRFH HFSRDPWQRLINNNWGMRPKAMRVKIFNIQVKEVTTSNGE TTVANNLTSTVQIFADSSYELPYVMDAGQEGSLPPFPNDV FMVPQYGYCGLVTGNTSQQOTDRNAFYCLEYFPSQMLRTG NNFEITYSFEKVPFHSYAHSSQLDRLMNPIDQYLYYLS KTINGSGQNQQTLKFSVAGPSNMAVQGRNYIPGPSYRQQR VSTTVTQNNNEFAWPGASSWALNGRNSLMNPGPAMASHK EGEDRFFPLSGSLIFGKQGTGRDNVDADKVMITNEEEIKT TNPVATESYQVATNHQSAQAQAQTGWVQNGAVPGMVWQ NRDIYYQGPIWAKIPHTDGHFHPSPPLIGGFGLKHPPPQIF IKNTPVPANPATTFSSSTPVNSFITQYSTGQVSVQIDWEIQ KERSKRWNPEVQFTSNYQQNSLLWAPDAAGKYTEPRAIG TRYLTHHL
AAV4_9VRGH SEQ ID NO: 114	AAV4	AAV9	VRGH	MTDGYLPDWLEDNLSEGVREWWALQPGAPKPKANQQHQDN ARGLVLPGYKYLGPNGLDKGEPVNAADAAALEHDKAYDQ QLKAGDNPYLKYNHADAEEFQRLQGDTSFGGNLGRAVFQA KKRVLEPLGLVEQAGETAPGKKRPLIESPQQPDSSTGIGK KGKQPAKKKLVFEDETGAGDGPPEGSTSGAMSDDSEMR AGGAAVEGGQGADGVGNASGDWHCDSTWSEGHVTTTSTR WVLPITYNNHLYKQISNSTSGGSSNDNAYFGFSTPWGYFDF NRFHCHFSRDPWQRLINNNWGMRPKAMRVKIFNIQVKEVT DNNGVKTVANNLTSTVQIFADSSYELPYVMDAGQEGSLPP FPNDVFMVPQYGYLTLNDGSQAVGRSSFYCLEYFPSQMLR TGNNFEITYSFEKVPFHSYAHSSQLDRLMNPIDQYLYY LSKTINGSGQNQQTLKFSVAGPSNMAVQGRNYIPGPSYRQ QRVSTTVTQNNNEFAWPGASSWALNGRNSLMNPGPAMAS HKEGEDRFFPLSGSLIFGKQGTGRDNVDADKVMITNEEEI KTTNPVATESYQVATNHQSAQAQAQTGWVQNGAVPGMV WQNRDIYYQGPIWAKIPHTDGHFHPSPPLIGGFGLKHPPPQ IFIKNTPVPANPATTFSSSTPVNSFITQYSTGQVSVQIDWE IQKERSKRWNPEVQFTSNYYKSSNNVEFAVNTGCVYTEPRA IGTRYLTHHL

AAVbo_9VR SEQ ID NO: 115	AAVbo	AAV9	VR	MSFVDHPPDWLESIGDGFREFLGLGAGPPKPKANQQKQDN ARGLVLPGYKYLGPNGLDKGDVPNFADDEVAREHDLSTYQK QLEAGDNPYLKYNHADAEEFQEKLASDTSF'GGNLGKAVFQA KKRILEPLGLVETPDKTAPAAKKRPLEQSPQEPDSSSGVG KKGKQPARKRLNFDEPGAGDGPPPEGPSSGAMSTETEMR AAAGGNGGDAGQGAEGVGNASGDWHCDSTWSESHVTTTST RTWVLPTYNNHLYLQISNSTSGGSSNDNAYFGFSTPWGYF DFNRFHCHFSPRDWQRLINNHWGLRPKSMQVRIFNIQVKE VTDNNGVKSNNLTSTVQIFADSTYELPYVMDAGQEGSLPP FPNDVFMVPQYGYLTLNDGSQAVGRSSFYCLEYF'PSQMLR TGNNFEMVYKFENVPFHSMYAHSSQLDRMLNPLLDQYLYY LSKTINGSGQNQQTLKFSVAGPSNMAVQGRNWLP'GPMRQ QRVSTTVTQNNNEFAWPGASSWALNGRWSNFAPGTAMAT HKEGEDRFFPLSGSLIFGKQGTGRDNVDADKLMFTSEDEL RATNPRD'TDLFGQVATNHQSAQAQAQTGWVQNGVYPGMV WQDRDIYYQGPIWAKIPHTDGHFHPSP'LI'GGFGLKSPPPQ IFIKNTPVPANPATTFS'PARINSF'ITQYSTGQVAVKIEWE IQKERSKRWNPEVQFTSNYYKSNNVEFAVNTGEGVYKEPRA IGSRYLTNHL
AAVbo_9GH SEQ ID NO: 116	AAVbo	AAV9	GH	MSFVDHPPDWLESIGDGFREFLGLGAGPPKPKANQQKQDN ARGLVLPGYKYLGPNGLDKGDVPNFADDEVAREHDLSTYQK QLEAGDNPYLKYNHADAEEFQEKLASDTSF'GGNLGKAVFQA KKRILEPLGLVETPDKTAPAAKKRPLEQSPQEPDSSSGVG KKGKQPARKRLNFDEPGAGDGPPPEGPSSGAMSTETEMR AAAGGNGGDAGQGAEGVGNASGDWHCDSTWSESHVTTTST RTWVLPTYNNHLYLRLGSSNASDTFNGFSTPWGYFDFNRF HCHFSPRDWQRLINNHWGLRPKSMQVRIFNIQVKEVTTSN GETTVSNNLTSTVQIFADSTYELPYVMDAGQEGSLPPFPN DVFMVPQYGYCGLVTGGSSQNQTD'RNAFYCLEYF'PSQMLR TGNNFEMVYKFENVPFHSMYAHSSQLDRMLNPLLDQYLYY LSKTINGSGQNQQTLKFSVAGPSNMAVQGRNYP'G'PSYRQ QRVSTTVTQNNNEFAWPGASSWALNGRNSLMNPGPAMAS HKEGEDRFFPLSGSLIFGKQGTGRDNVDADKVMITNEEEI KTTNPVATESY'GQVATNHQSAQAQAQTGWVQNGVYPGMV WQDRDIYYQGPIWAKIPHTDGHFHPSP'LI'GGFGLKSPPPQ IFIKNTPVPANPATTFS'PARINSF'ITQYSTGQVAVKIEWE IQKERSKRWNPEVQFTSNYGAQD'SLLWAPDNAGAYKEPRA IGSRYLTNHL
AAVbo_9VRGH SEQ ID NO: 117	AAVbo	AAV9	VRGH	MSFVDHPPDWLESIGDGFREFLGLGAGPPKPKANQQKQDN ARGLVLPGYKYLGPNGLDKGDVPNFADDEVAREHDLSTYQK QLEAGDNPYLKYNHADAEEFQEKLASDTSF'GGNLGKAVFQA KKRILEPLGLVETPDKTAPAAKKRPLEQSPQEPDSSSGVG KKGKQPARKRLNFDEPGAGDGPPPEGPSSGAMSTETEMR AAAGGNGGDAGQGAEGVGNASGDWHCDSTWSESHVTTTST RTWVLPTYNNHLYLQISNSTSGGSSNDNAYFGFSTPWGYF DFNRFHCHFSPRDWQRLINNHWGLRPKSMQVRIFNIQVKE VTDNNGVKSNNLTSTVQIFADSTYELPYVMDAGQEGSLPP FPNDVFMVPQYGYLTLNDGSQAVGRSSFYCLEYF'PSQMLR TGNNFEMVYKFENVPFHSMYAHSSQLDRMLNPLLDQYLYY LSKTINGSGQNQQTLKFSVAGPSNMAVQGRNYP'G'PSYRQ QRVSTTVTQNNNEFAWPGASSWALNGRNSLMNPGPAMAS HKEGEDRFFPLSGSLIFGKQGTGRDNVDADKVMITNEEEI KTTNPVATESY'GQVATNHQSAQAQAQTGWVQNGVYPGMV WQDRDIYYQGPIWAKIPHTDGHFHPSP'LI'GGFGLKSPPPQ IFIKNTPVPANPATTFS'PARINSF'ITQYSTGQVAVKIEWE IQKERSKRWNPEVQFTSNYYKSNNVEFAVNTGEGVYKEPRA IGSRYLTNHL
AAVbo_8VR SEQ ID NO: 118	AAVbo	AAV8	VR	MSFVDHPPDWLESIGDGFREFLGLGAGPPKPKANQQKQDN ARGLVLPGYKYLGPNGLDKGDVPNFADDEVAREHDLSTYQK QLEAGDNPYLKYNHADAEEFQEKLASDTSF'GGNLGKAVFQA KKRILEPLGLVETPDKTAPAAKKRPLEQSPQEPDSSSGVG KKGKQPARKRLNFDEPGAGDGPPPEGPSSGAMSTETEMR AAAGGNGGDAGQGAEGVGNASGDWHCDSTWSESHVTTTST

				RTWVLPPTYNNHLYLQISNGTSGGATNDNTYFGFSTPWGYF DFNRFHCHFSPRDWQRLINNHWGLRPKSMQVRIFNIQVKE VTQNEGTKSNNLTSTVQIFADSTYELPYVMDAQEGSLPP FPNDVFMVPQYGYLTLNNGSQAVGRSSFYCLEYFPSQMLR TGNNFEMVYKFENVPFHSMYAHSQSLDRLMNPLLDQYLYY LSRTQTTGGTANTQTLGFSQGGPNTMANQAKNWLPGPMMR QQRVSTTTGQNNNSNFAWTAGTKYHLNGRWSNFAPGTAMA THKDDEERFFPSNGILIFGKQNAARDNADYSDLMFTSEDE LRATNPRDLDLFGIVADNLQQQNTAPQIGTVNSQGVYPGM VWQDRDIYYQGPWAKIPHTDGHFHPSPPLIGGFGLKSPPP QIFIKNTPVPANPATTFSPARINSFITQYSTGQVAVKIEW EIQKERSKRWNPEVQFTSNYYKSTSVDFAVNTEGVYKEPR AIGSRYLTNHL
AAVbo_8GH SEQ ID NO: 119	AAVbo	AAV8	GH	MSFVDHPPDWLESIGDGFREFLGLAAGPPKPKANQQKQDN ARGLVLPGYKYLGPNGLDKGDVPNFADEVAREHDLSTYQK QLEAGDNPYLKYNHADADEFQEKLASDTSFGGNLKGAVFQA KKRILEPLGLVETPDKTAPAAKKRPLEQSPQEPDSSSGVG KKGKQPARKRLNFDDEPGAGDGPPEGPSSGAMSTETEMR AAAGNGGDAGQGAEGVGNASGDWHCDSTWSESHVTTTST RTWVLPPTYNNHLYLRLGSSNASDTFNGFSTPWGYDFNRF HCHFSPRDWQRLINNHWGLRPKSMQVRIFNIQVKEVTTSN GETTVSNNLTSTVQIFADSTYELPYVMDAQEGSLPPFPN DVMVPQYGYGLVTGGSSQNQTDNRFYCLEYFPSQMLR TGNNFEMVYKFENVPFHSMYAHSQSLDRLMNPLLDQYLYY LSRTQTTGGTANTQTLGFSQGGPNTMANQAKNWLPGPCYR QQRVSTTTGQNNNSNFAWTAGTKYHLNGRNSLANPGIAMA THKDDEERFFPSNGILIFGKQNAARDNADYSDVMLTSEEE IKTTNPVATEEYGIVADNLQQQNTAPQIGTVNSQGVYPGM VWQDRDIYYQGPWAKIPHTDGHFHPSPPLIGGFGLKSPPP QIFIKNTPVPANPATTFSPARINSFITQYSTGQVAVKIEW EIQKERSKRWNPEVQFTSNYGAQDSLLWAPDNAGAYKEPR AIGSRYLTNHL
AAVbo_8VRGH SEQ ID NO: 120	AAVbo	AAV8	VRGH	MSFVDHPPDWLESIGDGFREFLGLAAGPPKPKANQQKQDN ARGLVLPGYKYLGPNGLDKGDVPNFADEVAREHDLSTYQK QLEAGDNPYLKYNHADADEFQEKLASDTSFGGNLKGAVFQA KKRILEPLGLVETPDKTAPAAKKRPLEQSPQEPDSSSGVG KKGKQPARKRLNFDDEPGAGDGPPEGPSSGAMSTETEMR AAAGNGGDAGQGAEGVGNASGDWHCDSTWSESHVTTTST RTWVLPPTYNNHLYLQISNGTSGGATNDNTYFGFSTPWGYF DFNRFHCHFSPRDWQRLINNHWGLRPKSMQVRIFNIQVKE VTQNEGTKSNNLTSTVQIFADSTYELPYVMDAQEGSLPP FPNDVFMVPQYGYLTLNNGSQAVGRSSFYCLEYFPSQMLR TGNNFEMVYKFENVPFHSMYAHSQSLDRLMNPLLDQYLYY LSRTQTTGGTANTQTLGFSQGGPNTMANQAKNWLPGPCYR QQRVSTTTGQNNNSNFAWTAGTKYHLNGRNSLANPGIAMA THKDDEERFFPSNGILIFGKQNAARDNADYSDVMLTSEEE IKTTNPVATEEYGIVADNLQQQNTAPQIGTVNSQGVYPGM VWQDRDIYYQGPWAKIPHTDGHFHPSPPLIGGFGLKSPPP QIFIKNTPVPANPATTFSPARINSFITQYSTGQVAVKIEW EIQKERSKRWNPEVQFTSNYYKSTSVDFAVNTEGVYKEPR AIGSRYLTNHL
AAV8_9VRGBS SEQ ID NO: 121	AAV8	AAV9	VRGBS	MAADGYLPDWLEDNLSEGIREWALKPGAPKPKANQQKQD DGRGLVLPGYKYLGPFGNLDKGEFVNAAADAAALEHDKAYD QQLQAGDNPYLRYNHADADEFQERLQEDTSFGGNLGRAVFQ AKKRVLEPLGLVEEGAKTAPGKKRPVEPSPQRSPPDSSSTGI GKKGQQPARKRLNFGQTGDSESVDPQPPLGEPPAAPSGVG PNTMAAGGAPMADNNEGADGVGSSSGNWHCDSTWLGDRV ITTSTRTWALPTYNNHLYLQISNSTSGGSSNDNTYFGYST PWGYFDFNRFHCHFSPRDWQRLINNNWGF'RPKRLSF'KLFN IQVKEVTDNNGVKTIANNLSTIQVFTDSEYQLPYVLGSA HQGCLPPFPADVFMIPQYGYLTLNDGSQAVGRSSFYCLEY FPSQMLRTGNNFQFTYTFEDVPFHSSYAHSQSLDRLMNPL IDQYLYLSRTINGSGQNQQTTLGFSVAGPSNMAVQGRNWL

				PGPCYRQQRVSTTQNNSEFAWPGGTYHLNGRNSIAN PGIAMATHKEGEDRFFPLSGILIFGKQGTGRDNVDADKVM LTSEEEIKTTNPVATEEYGIVATNHQSAQAQAQTGTVNSQ GALPGMVWQNRDVYLQGPWAKIPHTDGNFHPSPLMGGFG LKHPPQILIKNTVPADPPTTFNQSKLNSFITQYSTGQV SVEIEWELQKENS KRWNPEIQYTSNYYKSNNVEFAVNTEG VYSEPRPIGTRYLTRNL
AAV8_9VRGH SEQ ID NO: 122	AAV8	AAV9	VRGH	MAADGYLPDWLEDNLSEGIREWALKPGAPKPKANQQKQD DGRGLVLPGYKYLGPFGNLDKGEFVNADAAALEHDKAYD QQLQAGDNPYLRYNHADADEFQERLQEDTSFGGNLGRAVFQ AKKRVLEPLGLVEEGAKTAPGKKRPVEPSPQSPDSSSTGI GKKGQQPARKRLNFGQTGDSESVDPDQPLGEPPAAPSGVGS PNTMAAGGGAPMADNNEGADGVGSSSGNWHCDSTWLGDRV ITSTRTWALPTYNNHLYKQISNSTSGGSSNDNTYFGYST PWGYFDNRFHCHFSPRDWQRLINNNWGFPRKRLSFKLFN IQVKEVTDNNGVKTIANNLTSTVQVFTDSEYQLPYVLGSA HQGLPPFPADVFMIQYGYLTLNDGSQAVGRSSFYCLEY FPSQMLRTGNNFQFSYTFEDVPPHSSYAHQSGLDRLMNPLI IDQYLYLSKTINGSGQNQQTLKFVSVAGPSNMAVQGRNYI PGPSYRQQRVSTTQNNSEFAWPGASSWALNGRNSLMN PGPAMASHKEGEDRFFPLSGSLIFGKQGTGRDNVDADKVM ITNEEEIKTTNPVATESYQVATNHQSAQAQAQTGWVQNG GILPGMVWQNRDVYLQGPWAKIPHTDGNFHPSPLMGGFG LKHPPQILIKNTVPADPPTTFNQSKLNSFITQYSTGQV SVEIEWELQKENS KRWNPEIQYTSNYYKSNNVEFAVNTEG VYSEPRPIGTRYLTRNL
AAVanc.110_9VR.PHP SEQ ID NO: 123	AAVanc.110	AAV9	VR.PHP	MAADGYLPDWLEDNLSEGIREWDLKPGAPKPKANQQKQD DGRGLVLPGYKYLGPFGNLDKGEFVNADAAALEHDKAYD QQLKAGDNPYLRYNHADADEFQERLQEDTSFGGNLGRAVFQ AKKRVLEPLGLVEEGAKTAPGKKRPVEQSPQEPDSSSGIG KTGQQPAKKRLNFGQTGDSESVDPDQPLGEPPAAPSGVGS NTMASGGGAPMADNNEGADGVGNSSGNWHCDSTWLGDRVI TTSTRTWALPTYNNHLYKQISNSTSGGSSNDNTYFGYSTP WGYFDNRFHCHFSPRDWQRLINNNWGFPRKRLNFKLFNI IQVKEVTDNNGVKTIANNLTSTVQVFTDSEYQLPYVLGSAH QGCLPPFPADVFMIQYGYLTLNDGSQAVGRSSFYCLEYF PSQMLRTGNNFQFSYTFEDVPPHSSYAHQSGLDRLMNPLI DQYLYFLSRTINGSGQNQQTLQFSQAGPSSMANQARNWVP GPCYRQQRVSTTQNNSEFAWPGATKYHLNGRDSLMP GVAMASHKEGEDRFFPLSGVLIFGKQGTGRDNVDADKVM TNEEEIKTTNPVATEEYGAVATNHQSAQTLAVPFKAQAQT GLVHNQGVLPGMVWQNRDVYLQGPWAKIPHTDGNFHPSP LMGGFGLKHPPQILIKNTVPADPPTTFNQAKLNSFITQ YSTGQVSVEIEWELQKENS KRWNPEIQYTSNYYKSNNVEF AVNTEGVYSEPRPIGTRFLTRNL
AAVanc.110_9VR SEQ ID NO: 124	AAVanc.110	AAV9	VR	MAADGYLPDWLEDNLSEGIREWDLKPGAPKPKANQQKQD DGRGLVLPGYKYLGPFGNLDKGEFVNADAAALEHDKAYD QQLKAGDNPYLRYNHADADEFQERLQEDTSFGGNLGRAVFQ AKKRVLEPLGLVEEGAKTAPGKKRPVEQSPQEPDSSSGIG KTGQQPAKKRLNFGQTGDSESVDPDQPLGEPPAAPSGVGS NTMASGGGAPMADNNEGADGVGNSSGNWHCDSTWLGDRVI TTSTRTWALPTYNNHLYKQISNSTSGGSSNDNTYFGYSTP WGYFDNRFHCHFSPRDWQRLINNNWGFPRKRLNFKLFNI QVKEVTDNNGVKTIANNLTSTVQVFTDSEYQLPYVLGSAH QGCLPPFPADVFMIQYGYLTLNDGSQAVGRSSFYCLEYF PSQMLRTGNNFQFSYTFEDVPPHSSYAHQSGLDRLMNPLI DQYLYFLSRTINGSGQNQQTLQFSQAGPSSMANQARNWVP GPCYRQQRVSTTQNNSEFAWPGATKYHLNGRDSLMP GVAMASHKEGEDRFFPLSGVLIFGKQGTGRDNVDADKVM TNEEEIKTTNPVATEEYGAVATNHQSAQAQAQTGLVHNQ VLPGMVWQNRDVYLQGPWAKIPHTDGNFHPSPLMGGFGL KHPPQILIKNTVPADPPTTFNQAKLNSFITQYSTGQVS VEIEWELQKENS KRWNPEIQYTSNYYKSNNVEFAVNTEGV

				YSEPRPIGTRFLTRNL
AAVanc.110_9GH.PHP SEQ ID NO: 125	AAVanc.110	AAV9	GH.PHP	MAADGYLPDWLEDNLSEGIREWDLKPGAPKPKANQQKQD DGRGLVLPGYKYLGPFNGLDKGEPVNAADAAALEHDKAYD QQLKAGDNPYLRYNHADADEFQERLQEDTSFGGNLGRAVFQ AKKRVLEPLGLVEEGAKTAPGKKRPVEQSPQEPDSSSGIG KTGQQPAKKRLNFGQTGDSESVDPQPPLGEPPAAPSGVGS NTMASGGGAPMADNNEGADGVGNSSGNWHCDSTWLGDRVI TTSTRTWALPTYNNHLYKQISNGTSGGSTNDNTYFGYSTP WGYDFNRFHCHFSPRDWQRLINNNWGFRPKRLNFKLFNI QVKEVTTNEGKTIANNLTSTVQVFTDSEYQLPYVLGSAH QGCLPPFPADVFMIPQYGYLTLNNGSQAVGRSSFYCLEYF PSQMLRTGNMFQFSYTFEDVPPHSSYAHSQSLDRLMNPLI DQYLYFLSRTINGSGQNQQTLKFSVAGPSNMAVQGRNYIP GPSYRQQRVSTTVTQNNNSEFAWPGASSWALNGRNSLMNP GPAMASHKEGEDRFFPLSGSLIFGKQGTGRDNVDADKVM TNEEEIKTTNPVATESYQQVATNHQSAQTAVPFKAQAQT GWVQNGQVLPGMVWQNRDVYLQGPWAKIPHTDGNFHPSP LMGGFGLKHPPPIILIKNTVPADPPTTFNQAKLNSFITQ YSTGQVSVEIEWELQKENSQRWNPEIQYTSNYYKSTNVDF AVNTEGVYSEPRPIGTRFLTRNL
AAVanc.110_9GH SEQ ID NO: 126	AAVanc.110	AAV9	GH	MAADGYLPDWLEDNLSEGIREWDLKPGAPKPKANQQKQD DGRGLVLPGYKYLGPFNGLDKGEPVNAADAAALEHDKAYD QQLKAGDNPYLRYNHADADEFQERLQEDTSFGGNLGRAVFQ AKKRVLEPLGLVEEGAKTAPGKKRPVEQSPQEPDSSSGIG KTGQQPAKKRLNFGQTGDSESVDPQPPLGEPPAAPSGVGS NTMASGGGAPMADNNEGADGVGNSSGNWHCDSTWLGDRVI TTSTRTWALPTYNNHLYKQISNGTSGGSTNDNTYFGYSTP WGYDFNRFHCHFSPRDWQRLINNNWGFRPKRLNFKLFNI QVKEVTTNEGKTIANNLTSTVQVFTDSEYQLPYVLGSAH QGCLPPFPADVFMIPQYGYLTLNNGSQAVGRSSFYCLEYF PSQMLRTGNMFQFSYTFEDVPPHSSYAHSQSLDRLMNPLI DQYLYFLSRTINGSGQNQQTLKFSVAGPSNMAVQGRNYIP GPSYRQQRVSTTVTQNNNSEFAWPGASSWALNGRNSLMNP GPAMASHKEGEDRFFPLSGSLIFGKQGTGRDNVDADKVM TNEEEIKTTNPVATESYQQVATNHQSAQAQAQTGWVQNG VLPGMVWQNRDVYLQGPWAKIPHTDGNFHPSPLMGGFGL KHPPPIILIKNTVPADPPTTFNQAKLNSFITQYSTGQVS VEIEWELQKENSQRWNPEIQYTSNYYKSTNVDFAVNTEGV YSEPRPIGTRFLTRNL
AAVanc.110_9VRGBS SEQ ID NO: 127	AAVanc.110	AAV9	VRGBS	MAADGYLPDWLEDNLSEGIREWDLKPGAPKPKANQQKQD DGRGLVLPGYKYLGPFNGLDKGEPVNAADAAALEHDKAYD QQLKAGDNPYLRYNHADADEFQERLQEDTSFGGNLGRAVFQ AKKRVLEPLGLVEEGAKTAPGKKRPVEQSPQEPDSSSGIG KTGQQPAKKRLNFGQTGDSESVDPQPPLGEPPAAPSGVGS NTMASGGGAPMADNNEGADGVGNSSGNWHCDSTWLGDRVI TTSTRTWALPTYNNHLYKQISNSTSGGSSNDNTYFGYSTP WGYDFNRFHCHFSPRDWQRLINNNWGFRPKRLNFKLFNI QVKEVTDNNGVKTIANNLTSTVQVFTDSEYQLPYVLGSAH QGCLPPFPADVFMIPQYGYLTLNDGSQAVGRSSFYCLEYF PSQMLRTGNMFQFSYTFEDVPPHSSYAHSQSLDRLMNPLI DQYLYFLSRTINGSGQNQQTLQFSVAGPSNMAVQARNWVP GPCYRQQRVSTTVTQNNNSEFAWPGATKYHLNGRDSLMNP GVMASHKEGEDRFFPLSGVLIFGKQGTGRDNVDADKVM TNEEEIKTTNPVATEEYGAVATNHQSAQAQAQTGLVHNQ VLPGMVWQNRDVYLQGPWAKIPHTDGNFHPSPLMGGFGL KHPPPIILIKNTVPADPPTTFNQAKLNSFITQYSTGQVS VEIEWELQKENSQRWNPEIQYTSNYYKSNVVEAVNTEGV YSEPRPIGTRFLTRNL
AAVanc.110_8GH SEQ ID NO: 128	AAVanc.110	AAV8	GH	MAADGYLPDWLEDNLSEGIREWDLKPGAPKPKANQQKQD DGRGLVLPGYKYLGPFNGLDKGEPVNAADAAALEHDKAYD QQLKAGDNPYLRYNHADADEFQERLQEDTSFGGNLGRAVFQ AKKRVLEPLGLVEEGAKTAPGKKRPVEQSPQEPDSSSGIG KTGQQPAKKRLNFGQTGDSESVDPQPPLGEPPAAPSGVGS

				NTMASGGGAPMADNNEGADGVGNSSGNWHCDSTWLGDRVI TTSTRTWALPTYNNHLYKQISNGTSGGSTNDNTYFGYSTP WGYDFNRFHCHFSPRDWQRLINNNWGFRPKRLNFKLFNI QVKEVTTNEGKTIANNLTSTVQVFTDSEYQLPYVLGSAH QGCLPPFPADVFMIPQYGYLTLNNGSQAVGRSSFYCLEYF PSQMLRTGNNFQFSYTFEDVPPHSSYAHSSQSLDRLMNPLI DQYLYFLSRTQTTGGTANTQTLGFSQGGPNTMANQAKNWL PGPCYRQQRVSTTTGQNNNSNFAWTAGTKYHLNGRNSLAN PGIAMATHKDDDEERFFPSNGILIFGKQNAARDNADYSDVM LTSEEEIKTTNPVATEEYGVADNLQQQNTAPQIGTVNSQ GVLPGMVWQNRDVYLQGPWAKIPHTDGNFHPSPLMGGFG LKHPPQILIKNTPVPADPPTTFNQAKLNSFITQYSTGQV SVEIEWELQKENSkrwnPEIQYTSNYYKSTNVDFAVNTEG VYSEPRPIGTRFLTRNL
AAV9_anc.110VR SEQ ID NO: 129	AAV9	AAVanc.110	VR	MAADGYLPDWLEDNLSEGIREWALKPGAPQPKANQQHQD NARGVLVPGYKYLGPNGLDKGEFVNAAADAAALEHDKAYD QQLKAGDNPYLKYNHADADEFQERLKEDTSFGGNLGRAVFQ AKKRLLEPLGLVEEAAKTAPGKKRPVEQSPQEPDSSAGIG KSGAQPAKKRLNFGQTGDTEVPDPQPIGEPPAAPSGVGS LTMASGGGAPVADNNEGADGVGSSGNWHCDSQWLGDRVI TTSTRTWALPTYNNHLYKQISNGTSGGSTNDNAYFGYSTP WGYDFNRFHCHFSPRDWQRLINNNWGFRPKRLNFKLFNI QVKEVTTNEGKTIANNLTSTVQVFTDSDYQLPYVLGSAH EGCLPPFPADVFMIPQYGYLTLNNGSQAVGRSSFYCLEYF PSQMLRTGNNFQFSYEFENVPPHSSYAHSSQSLDRLMNPLI DQYLYFLSRTQTTGTAGTQTLKFSVAGPSNMAVQGRNYP GPSYRQQRVSTTTNQNNSNFAWTGASSWALNGRNSLMNP GPAMASHKDDDEDRFFPSSGSLIFGKQGAGNDNVDSQVMI TNEEEIKTTNPVATESYQQVATNNQSANTQAQTGWVQNGG ILPGMVWQDRDVYLQGPWAKIPHTDGNFHPSPLMGGFGM KHPPQILIKNTPVPADPPTAFNKDKLNSFITQYSTGQVS VEIEWELQKENSkrwnPEIQYTSNYYKSTNVDFAVNTEGV YSEPRPIGTRFLTRNL
AAV9_anc.110GH SEQ ID NO: 130	AAV9	AAVanc.110	GH	MAADGYLPDWLEDNLSEGIREWALKPGAPQPKANQQHQD NARGVLVPGYKYLGPNGLDKGEFVNAAADAAALEHDKAYD QQLKAGDNPYLKYNHADADEFQERLKEDTSFGGNLGRAVFQ AKKRLLEPLGLVEEAAKTAPGKKRPVEQSPQEPDSSAGIG KSGAQPAKKRLNFGQTGDTEVPDPQPIGEPPAAPSGVGS LTMASGGGAPVADNNEGADGVGSSGNWHCDSQWLGDRVI TTSTRTWALPTYNNHLYKQISNSTSGGSSNDNAYFGYSTP WGYDFNRFHCHFSPRDWQRLINNNWGFRPKRLNFKLFNI QVKEVTDNNGVKTIANNLTSTVQVFTDSDYQLPYVLGSAH EGCLPPFPADVFMIPQYGYLTLNDGSQAVGRSSFYCLEYF PSQMLRTGNNFQFSYEFENVPPHSSYAHSSQSLDRLMNPLI DQYLYFLSRTQTTGTAGTQTLQFSQAGPSSMANQARNWVP GPCYRQQRVSTTTNQNNSNFAWTGATKYHLNGRDSLMP GVAMASHKDDDEDRFFPSSGVLIFGKQGAGNDNVDSQVMI TNEEEIKTTNPVATEEYGAVATNNQSANTQAQTGLVHNQG ILPGMVWQDRDVYLQGPWAKIPHTDGNFHPSPLMGGFGM KHPPQILIKNTPVPADPPTAFNKDKLNSFITQYSTGQVS VEIEWELQKENSkrwnPEIQYTSNYYKSNVVEFAVNTEGV YSEPRPIGTRFLTRNL
AAV8_anc.110VR SEQ ID NO: 131	AAV8	AAVanc.110	VR	MAADGYLPDWLEDNLSEGIREWALKPGAPKPKANQQKQD DGRGLVLPGYKYLGFNGLDKGEFVNAAADAAALEHDKAYD QQLQAGDNPYLRYNHADADEFQERLQEDTSFGGNLGRAVFQ AKKRVLEPLGLVEEGAKTAPGKKRPVEPSPQRSPTSSTGI GKKGQQPARKRLNFGQTGDSESVDPDPQLGEPPAAPSGVG PNTMAAGGGAPMADNNEGADGVGSSGNWHCDSTWLGDRV ITTSTRTWALPTYNNHLYKQISNGTSGGSTNDNTYFGYST PWGYDFNRFHCHFSPRDWQRLINNNWGFRPKRLSFKLFN IQVKEVTTNEGKTIANNLTSTIQVFTDSEYQLPYVLGSA HQGCLPPFPADVFMIPQYGYLTLNNGSQAVGRSSFYCLEY FPSQMLRTGNNFQFTYTFEDVPPHSSYAHSSQSLDRLMNPL

				IDQYLYFLSRTQTTGTAGTQTLGFSQGGPNTMANQAKNWL PGPCYRQQRVSTTTNQNNSNFAWTGGTKYHLNGRNSLAN PGIAMATHKDDDRFFPSSGILIFGKQGAGNDNDVYSQVM LTSEEEIKTTNPVATEEYGVATNNQSANTQAQTGTVNSQ GALPGMVWQNRDVYLQGPWAKIPHTDGNFHPSPMLMGFG LKHPPPQILIKNTPVPADPPTTFNQSKLNSFITQYSTGQV SVEIEWELQKENSkrwnPEIQYTSNYYKSTNVDFAVNTEG VYSEPRPIGTRFLTRNL
AAV8_anc.110GH SEQ ID NO: 132	AAV8	AAVanc.110	GH	MAADGYLPDWLEDNLSEGIREWALKPGAPKPKANQQKQD DGRGLVLPGYKYLGPFGNLDKGEPVNAADAAALEHDKAYD QQLQAGDNPYLRYNHADADEFQERLQEDTSFGGNLGRAVFQ AKKRVLEPLGLVEEGAKTAPGKKRPVEPSQSPDSSSTGI GKKGQQPARKRLNFGQTGDSSESVPDPQPLGEPPAAPSGVG PNTMAAGGGAPMADNNEGADGVGSSSGNWHCDSTWLGD RV ITTSTRTWALPTYNNHLYKQISNGTSGGATNDNTYFGYST PWGYDFNRFHCHFSPRDWQRLINNNWGF'RPKRLSFKLFN IQVKEVTQNEGKTIANNLTSTIQVFTDSEYQLPYVLGSA HQGCLPPFPADVFMIPQYGYLTLNNGSQAVGRSSFYCLEY FPSQMLRTGNNFQFTYTFEDVPPHSSYAHSSQSLDRLMNPL IDQYLYFLSRTQTTGTAGTQTLQFSQAGPSSMANQARNWV PGPCYRQQRVSTTTNQNNSNFAWTGATKYHLNGRDLSMN PGVAMASHKDDDRFFPSSGVLI'FGKQGAGNDNDVYSQVM ITNEEEIKTTNPVATEEYGAVATNNQSANTQAQTGLVHNQ GALPGMVWQNRDVYLQGPWAKIPHTDGNFHPSPMLMGFG LKHPPPQILIKNTPVPADPPTTFNQSKLNSFITQYSTGQV SVEIEWELQKENSkrwnPEIQYTSNYYKSTSVDFAVNTEG VYSEPRPIGTRFLTRNL
Anc.110_8VR SEQ ID NO: 133	AAVanc.110	AAV8	VR	MAADGYLPDWLEDNLSEGIREWDLKPGAPKPKANQQKQD DGRGLVLPGYKYLGPFGNLDKGEPVNAADAAALEHDKAYD QQLKAGDNPYLRYNHADADEFQERLQEDTSFGGNLGRAVFQ AKKRVLEPLGLVEEGAKTAPGKKRPVEQSPQEPDSSSGIG KTGQQPAKKRLNFGQTGDSSESVPDPQPLGEPPAAPSGVGS NTMASGGGAPMADNNEGADGVGSSSGNWHCDSTWLGD RV TTSTRTWALPTYNNHLYKQISNGTSGGATNDNTYFGYSTP WGYYDFNRFHCHFSPRDWQRLINNNWGF'RPKRLNFKLFN QVKEVTQNEGKTIANNLTSTVQVFTDSEYQLPYVLGSAH QGCLPPFPADVFMIPQYGYLTLNNGSQAVGRSSFYCLEYF PSQMLRTGNNFQFSYTFEDVPPHSSYAHSSQSLDRLMNPLI DQYLYFLSRTQTTGGTANTQTLQFSQAGPSSMANQARNWV PGPCYRQQRVSTTTGQNNNSNFAWTAATKYHLNGRDLSMN PGVAMASHKDDDEERFFP'NGVLI'FGKQNAARDNADYSVM ITNEEEIKTTNPVATEEYGAVADNLQQNTAPQIGLVHNQ GVLPGMVWQNRDVYLQGPWAKIPHTDGNFHPSPMLMGFG LKHPPPQILIKNTPVPADPPTTFNQAKLNSFITQYSTGQV SVEIEWELQKENSkrwnPEIQYTSNYYKSTSVDFAVNTEG VYSEPRPIGTRFLTRNL
AAV8_6VR SEQ ID NO: 134	AAV8	AAV6	VR	MAADGYLPDWLEDNLSEGIREWALKPGAPKPKANQQKQD DGRGLVLPGYKYLGPFGNLDKGEPVNAADAAALEHDKAYD QQLQAGDNPYLRYNHADADEFQERLQEDTSFGGNLGRAVFQ AKKRVLEPLGLVEEGAKTAPGKKRPVEPSQSPDSSSTGI GKKGQQPARKRLNFGQTGDSSESVPDPQPLGEPPAAPSGVG PNTMAAGGGAPMADNNEGADGVGSSSGNWHCDSTWLGD RV ITTSTRTWALPTYNNHLYKQISSASTGASNDNTYFGYSTP WGYYDFNRFHCHFSPRDWQRLINNNWGF'RPKRLSFKLFNI QVKEVTTNDGVTTIANNLTSTIQVFTDSEYQLPYVLGSAH QGCLPPFPADVFMIPQYGYLTLNNGSQAVGRSSFYCLEYF PSQMLRTGNNFQFTYTFEDVPPHSSYAHSSQSLDRLMNPLI DQYLYFLSRTQNSGSAQNKDLGFSQGGPNTMANQAKNWL PGPCYRQQRVSKTKTDNNNSNFTWTGATKYHLNGRNSLAN PGIAMASHKDDKDKFFPMSGILIFGKESAGASNTALDNVM LTSEEEIKTTNPVATEEYGVAVNLQSSSTDPATGTVNSQ GALPGMVWQNRDVYLQGPWAKIPHTDGNFHPSPMLMGFG LKHPPPQILIKNTPVPADPPTTFNQSKLNSFITQYSTGQV

				SVEIEWELQKENS KRWNPEIQYTSNYAKSANVDFAVNTEG VYSEPRPIGTRYLTRNL
AAV8_6VRGBS SEQ ID NO: 135	AAV8	AAV6	VRGBS	MAADGYLPDWLEDNLSEGIREW WALKPGAPKPKANQQKQD DGRGLVLPGYKYLGPFGNLDKGEPVNAADAAALEHDKAYD QQLQAGDNPYLRYNHADA EFQERLQEDTSFGGNLGRAVFQ AKKRVLEPLGLVEEGAKTAPGKKRPVEPSPQRSPTSSTGI GKKGQQPARKRLNFGQTGDSESVDPDPQLGEPAPAAPSGVG PNTMAAGGGAPMADNNEGADGVGSSSGNWHCDSTWLGDRV ITTSTRTWALPTYNNHLYKQISSASTGASNDNTYFGYSTP WGYFDNRFHCHFSRQWLINNNWGF RPKRLSFKLFNI QVKEVTTNDGVTTIANNLTSTIQVFTDSEYQLPYVLGSAH QGCLPPFPADVFMIPQYGYLTLNNGSQAVGRSSFYCLEYF PSQMLRTGNNFQFTYTFEDVPPHSSYAHSQSLDRLMNPLI DQYLYYLSRTQNGSGSAQNKDLGF'SRGSPAGMSVQAKNWL PGPCYRQQRVSKTKTDNNNSNFTWTGATKYHLNGRNSLAN PGIAMASHKDDKDFPMSGILIF'GKESAGASNTALDNVM LTSEEEIKTNPVATEEY GIVAVNLQSSSTDPATGTVNSQ GALPGMVWQNRDVYLQGP I WAKIPHTDGNFHPSPMLMGFG LKHPPPQILIKNTPVPADPPTTFNQSKLNSFITQYSTGQV SVEIEWELQKENS KRWNPEIQYTSNYAKSANVDFAVNTEG VYSEPRPIGTRYLTRNL
AAV8_6VRGH SEQ ID NO: 136	AAV8	AAV6	VRGH	MAADGYLPDWLEDNLSEGIREW WALKPGAPKPKANQQKQD DGRGLVLPGYKYLGPFGNLDKGEPVNAADAAALEHDKAYD QQLQAGDNPYLRYNHADA EFQERLQEDTSFGGNLGRAVFQ AKKRVLEPLGLVEEGAKTAPGKKRPVEPSPQRSPTSSTGI GKKGQQPARKRLNFGQTGDSESVDPDPQLGEPAPAAPSGVG PNTMAAGGGAPMADNNEGADGVGSSSGNWHCDSTWLGDRV ITTSTRTWALPTYNNHLYKQISSASTGASNDNTYFGYSTP WGYFDNRFHCHFSRQWLINNNWGF RPKRLSFKLFNI QVKEVTTNDGVTTIANNLTSTIQVFTDSEYQLPYVLGSAH QGCLPPFPADVFMIPQYGYLTLNNGSQAVGRSSFYCLEYF PSQMLRTGNNFQFTYTFEDVPPHSSYAHSQSLDRLMNPLI DQYLYYLSRTQNGSGSAQNKDLGF'SRGSPAGMSVQPKNWL PGPCYRQQRVSKTKTDNNNSNFTWTGASKYNLNGRESIIN PGTAMASHKDDKDFPMSGVMI'GKESAGASNTALDNVM ITDEEEIKATNPVATERFGTVAVNLQSSSTDPATGDVHVM GALPGMVWQNRDVYLQGP I WAKIPHTDGNFHPSPMLMGFG LKHPPPQILIKNTPVPADPPTTFNQSKLNSFITQYSTGQV SVEIEWELQKENS KRWNPEIQYTSNYAKSANVDFAVNTEG VYSEPRPIGTRYLTRNL
AAV8_9VRGBS.SA SEQ ID NO: 137	AAV8	AAV9	VRGBS. SA	MAADGYLPDWLEDNLSEGIREW WALKPGAPKPKANQQKQD DGRGLVLPGYKYLGPFGNLDKGEPVNAADAAALEHDKAYD QQLQAGDNPYLRYNHADA EFQERLQEDTSFGGNLGRAVFQ AKKRVLEPLGLVEEGAKTAPGKKRPVEPSPQRSPTSSTGI GKKGQQPARKRLNFGQTGDSESVDPDPQLGEPAPAAPSGVG PNTMAAGGGAPMADNNEGADGVGSSSGNWHCDSTWLGDRV ITTSTRTWALPTYNNHLYKQISNSTSGSSNDNTYFGYST PWGYFDNRFHCHFSRQWLINNNWGF RPKRLSFKLFNI IQVKEVTDNNGVKTIANNLSTIQVFTDSEYQLPYVLGSA HQGCLPPFPADVFMIPQYGYLTLNDGSAVGRSSFYCLEY FPSQMLRTGNNFQFTYTFEDVPPHSSYAHSQSLDRLMNPLI IDQYLYYLSRTINGSGQNQQT LGSVAGPANMAVQGRNWL PGPCYRQQRVSTTVTQNNSEFAWPGGTYHLNGRNSLAN PGIAMATHKEGEDRFPLSGILIF'GKQGTGRDNVDADKVM LTSEEEIKTNPVATEEY GIVATNHQSAQAQAQTGTVNSQ GALPGMVWQNRDVYLQGP I WAKIPHTDGNFHPSPMLMGFG LKHPPPQILIKNTPVPADPPTTFNQSKLNSFITQYSTGQV SVEIEWELQKENS KRWNPEIQYTSNYKSNNVFAVNTEG VYSEPRPIGTRYLTRNL
AAV8_9VRGBS.ST SEQ ID NO: 138	AAV8	AAV9	VRGBS. ST	MAADGYLPDWLEDNLSEGIREW WALKPGAPKPKANQQKQD DGRGLVLPGYKYLGPFGNLDKGEPVNAADAAALEHDKAYD QQLQAGDNPYLRYNHADA EFQERLQEDTSFGGNLGRAVFQ AKKRVLEPLGLVEEGAKTAPGKKRPVEPSPQRSPTSSTGI GKKGQQPARKRLNFGQTGDSESVDPDPQLGEPAPAAPSGVG

				PNTMAAGGGAPMADNNEGADGVGSSSGNWHCDSTWLGDRV ITTSTRTWALPTYNNHLYKQISNSTSGGSSNDNTYFGYST PWGYFDFNRFHCHFSPRDWQRLINNNWGF'RPKRLSFKLFN IQVKEVTDNNGVKTIANNLTSTIQVFTDSEYQLPYVLGSA HQGCLPPFPADVFMI PQYGYLTLDNGSQAVGRSSFYCLEY FPSQMLRTGNNFQFTYTFEDVPFHSSYAHSSQLDRLMNPL IDQYLYLSRTINGSGQNQOTLGF'SVAGP'TNMAVQGRNWL PGPCYRQQRVSTTQNNSEFAWPGGTYHLNGRNSLAN PGIAMATHKEGEDRFFPLSGILIFGKQGTGRDNVDADKVM LTSEEEIKTTNPVATEEYGIVATNHQSAQAQAQTGTVNSQ GALPGMVWQNRDVLQGPWAKIPHTDGNFHPSPMLMGFG LKHPPPPQILIKNTPVPADPPTTFNQSKLNSFITQYSTGQV SVEIEWELQKENS KRWNPEIQYTSNYYKSNNVEFAVNTEG VYSEPRPIGTRYLTRNL
AAV8_9VRGBS.MA SEQ ID NO: 139	AAV8	AAV9	VRGBS. MA	MAADGYLPDWLEDNLSEGIREWALKPGAPKPKANQQKQD DGRGLVLPGYKYLGPFGNLDKGEFVNAAADAAAEHDKAYD QQLQAGDNPYLRYNHADAEEFQERLQEDTSFGGNLGRAVFQ AKKRVLEPLGLVEEGAKTAPGKKRPVEPSPQRSPPDSSGTGI GKKGQQPARKRLNFGQTGDSESVDPDQPLGEPPAAPSGVG PNTMAAGGGAPMADNNEGADGVGSSSGNWHCDSTWLGDRV ITTSTRTWALPTYNNHLYKQISNSTSGGSSNDNTYFGYST PWGYFDFNRFHCHFSPRDWQRLINNNWGF'RPKRLSFKLFN IQVKEVTDNNGVKTIANNLTSTIQVFTDSEYQLPYVLGSA HQGCLPPFPADVFMI PQYGYLTLDNGSQAVGRSSFYCLEY FPSQMLRTGNNFQFTYTFEDVPFHSSYAHSSQLDRLMNPL IDQYLYLSRTINGSGQNQOTLGF'SVAGP'SNAAVQGRNWL PGPCYRQQRVSTTQNNSEFAWPGGTYHLNGRNSLAN PGIAMATHKEGEDRFFPLSGILIFGKQGTGRDNVDADKVM LTSEEEIKTTNPVATEEYGIVATNHQSAQAQAQTGTVNSQ GALPGMVWQNRDVLQGPWAKIPHTDGNFHPSPMLMGFG LKHPPPPQILIKNTPVPADPPTTFNQSKLNSFITQYSTGQV SVEIEWELQKENS KRWNPEIQYTSNYYKSNNVEFAVNTEG VYSEPRPIGTRYLTRNL
AAV8_9VRGBS.MI SEQ ID NO: 140	AAV8	AAV9	VRGBS. MI	MAADGYLPDWLEDNLSEGIREWALKPGAPKPKANQQKQD DGRGLVLPGYKYLGPFGNLDKGEFVNAAADAAAEHDKAYD QQLQAGDNPYLRYNHADAEEFQERLQEDTSFGGNLGRAVFQ AKKRVLEPLGLVEEGAKTAPGKKRPVEPSPQRSPPDSSGTGI GKKGQQPARKRLNFGQTGDSESVDPDQPLGEPPAAPSGVG PNTMAAGGGAPMADNNEGADGVGSSSGNWHCDSTWLGDRV ITTSTRTWALPTYNNHLYKQISNSTSGGSSNDNTYFGYST PWGYFDFNRFHCHFSPRDWQRLINNNWGF'RPKRLSFKLFN IQVKEVTDNNGVKTIANNLTSTIQVFTDSEYQLPYVLGSA HQGCLPPFPADVFMI PQYGYLTLDNGSQAVGRSSFYCLEY FPSQMLRTGNNFQFTYTFEDVPFHSSYAHSSQLDRLMNPL IDQYLYLSRTINGSGQNQOTLGF'SVAGP'SNAAVQGRNWL PGPCYRQQRVSTTQNNSEFAWPGGTYHLNGRNSLAN PGIAMATHKEGEDRFFPLSGILIFGKQGTGRDNVDADKVM LTSEEEIKTTNPVATEEYGIVATNHQSAQAQAQTGTVNSQ GALPGMVWQNRDVLQGPWAKIPHTDGNFHPSPMLMGFG LKHPPPPQILIKNTPVPADPPTTFNQSKLNSFITQYSTGQV SVEIEWELQKENS KRWNPEIQYTSNYYKSNNVEFAVNTEG VYSEPRPIGTRYLTRNL
AAV8_9VRGBS.MK SEQ ID NO: 141	AAV8	AAV9	VRGBS. MK	MAADGYLPDWLEDNLSEGIREWALKPGAPKPKANQQKQD DGRGLVLPGYKYLGPFGNLDKGEFVNAAADAAAEHDKAYD QQLQAGDNPYLRYNHADAEEFQERLQEDTSFGGNLGRAVFQ AKKRVLEPLGLVEEGAKTAPGKKRPVEPSPQRSPPDSSGTGI GKKGQQPARKRLNFGQTGDSESVDPDQPLGEPPAAPSGVG PNTMAAGGGAPMADNNEGADGVGSSSGNWHCDSTWLGDRV ITTSTRTWALPTYNNHLYKQISNSTSGGSSNDNTYFGYST PWGYFDFNRFHCHFSPRDWQRLINNNWGF'RPKRLSFKLFN IQVKEVTDNNGVKTIANNLTSTIQVFTDSEYQLPYVLGSA HQGCLPPFPADVFMI PQYGYLTLDNGSQAVGRSSFYCLEY FPSQMLRTGNNFQFTYTFEDVPFHSSYAHSSQLDRLMNPL IDQYLYLSRTINGSGQNQOTLGF'SVAGP'SNAAVQGRNWL PGPCYRQQRVSTTQNNSEFAWPGGTYHLNGRNSLAN PGIAMATHKEGEDRFFPLSGILIFGKQGTGRDNVDADKVM LTSEEEIKTTNPVATEEYGIVATNHQSAQAQAQTGTVNSQ

				GALPGMVWQNRDVYLQGP IWAKIPHTDGNFHPSPMLMGFG LKHPPQILIKNTPVPADPPTTFNQSKLNSFITQYSTGQV SVEIEWELQKENS KRWNPEIQYTSNYYKSNVFEAVNTEG VYSEPRPIGTRYLTRNL
AAV8_9VRGBS.PA SEQ ID NO: 142	AAV8	AAV9	VRGBS. PA	MAADGYLPDWLEDNLSEGIREWALKPGAPKPKANQQKQD DGRGLVLPGYKYLGPFGNLDKGEFVNAAADAAAEHDKAYD QQLQAGDNPYLRYNHADADEFQERLQEDTSFGGNLGRAVFQ AKKRVLEPLGLVEEGAKTAPGKKRPVEPSPQRSPPDSSSTGI GKKGQQPARKRLNFGQTGDSESVDPDPQLGEPPAAPSGVG PNTMAAGGAPMADNNEGADGVGSSSGNWHCDSTWLGDRV ITTSTRTWALPTYNNHLYKQISNSTSGGSSNDNTYFGYST PWGYDFNRFHCHFSRDPWQRLINNNWGFPRKRLSFKLFNI IQKEVTDNNGVKTIANNLTSTIQVFTDSEYQLPYVLGSA HQGCLPPFPADVFMIPQYGYLTLDGSAQAVGRSSFYCLEY FPSQMLRTGNNFQFTYTFEDVPFHSSYAHSSQSLDRLMNP LIDQYLYLSRTINGSGQNQQTLLGF'SVAGASNMVQGRNWL PGPCYRQQRVSTTVTQNNNSEFAWPGGTYHLNGRNSLAN PGIAMATHKEGEDRFFPLSGILIFGKQGTGRDNVDADKVM LTSEEEIKTTNPVATEEYGI VATNHQSAQAQAGTGTVNSQ GALPGMVWQNRDVYLQGP IWAKIPHTDGNFHPSPMLMGFG LKHPPQILIKNTPVPADPPTTFNQSKLNSFITQYSTGQV SVEIEWELQKENS KRWNPEIQYTSNYYKSNVFEAVNTEG VYSEPRPIGTRYLTRNL
AAVanc110_9VRHS SEQ ID NO: 143	AAVanc.110	AAV9	VRHS	MAADGYLPDWLEDNLSEGIREWDLKPGAPKPKANQQKQD DGRGLVLPGYKYLGPFGNLDKGEFVNAAADAAAEHDKAYD QQLKAGDNPYLRYNHADADEFQERLQEDTSFGGNLGRAVFQ AKKRVLEPLGLVEEGAKTAPGKKRPVEQSPQEPDSSSGIG KTGQQPAKKRLNFGQTGDSESVDPDPQLGEPPAAPSGVGS NTMASGGAPMADNNEGADGVGSSSGNWHCDSTWLGDRVI TTSTRTWALPTYNNHLYKQISNSTSGGSSNDNTYFGYSTP WGYDFNRFHCHFSRDPWQRLINNNWGFPRKRLNFKLFNI QVKEVTDNNGVKTIANNLTSTVQVFTDSEYQLPYVLGSAH QGCLPPFPADVFMIPQYGYLTLDGSAQAVGRSSFYCLEYF PSQMLRTGNNFQFTYTFEDVPFHSSYAHSSQSLDRLMNP LIDQYLYLSRTINGSGQNQQTLLGF'SQAGPSSMANQARNWVP GPCYRQQRVSTTVTQNNNSEFAWPGATKYHLNGRDSLMP GVAMASHKEGEDRFFPLSGVLIIFGKQGTGRDNVDADKVM TNEEEIKTTNPVATEEYGA VATNHQSAQAQAGTGLVHNQ VLPGMVWQNRDVYLQGP IWAKIPHTDGNFHPSPMLMGFG LKHPPQILIKNTPVPADPPTTFNQAKLNSFITQYSTGQVS VEIEWELQKENS KRWNPEIQYTSNYYKSNVFEAVNTEGV YSEPRPIGTRFLTRNL
AAV8_5VR SEQ ID NO: 144	AAV8	AAV5	VR	MAADGYLPDWLEDNLSEGIREWALKPGAPKPKANQQKQD DGRGLVLPGYKYLGPFGNLDKGEFVNAAADAAAEHDKAYD QQLQAGDNPYLRYNHADADEFQERLQEDTSFGGNLGRAVFQ AKKRVLEPLGLVEEGAKTAPGKKRPVEPSPQRSPPDSSSTGI GKKGQQPARKRLNFGQTGDSESVDPDPQLGEPPAAPSGVG PNTMAAGGAPMADNNEGADGVGSSSGNWHCDSTWLGDRV ITTSTRTWALPTYNNHLYKQIKSGSVDGSSNDNTYFGYSTP WGYDFNRFHCHFSRDPWQRLINNNWGFPRKRLSFKLFNI QVKEVTVQDSTTTIANNLTSTIQVFTDSEYQLPYVLGSAH QGCLPPFPADVFMIPQYGYLTLDNRDNTENPTERSFYCLE YFPSQMLRTGNNFQFTYTFEDVPFHSSYAHSSQSLDRLMNP LIDQYLYLSRTNTGGVQFSQGGPNTMANQAKNWLPGPCY RQGWNLGSGVNRASVSFAFATGTYHLNGRNSLANPGIAM ATNNLQGSNTYALENTLIFNSQPANPGTTATYLEGNMLTS EEEEKTTNPVATEEYGI VATNNQSSSTPATGTVNSQ GALPGMVWQNRDVYLQGP IWAKIPHTDGNFHPSPMLMGFG LKHPPQILIKNTPVPADPPTTFNQSKLNSFITQYSTGQVS VEIEWELQKENS KRWNPEIQYTSNYYKSNVFEAVNTEGV YSEPRPIGTRYLTRNL
Anc110_5VR SEQ ID NO: 145	Anc110	AAV5	VR	MAADGYLPDWLEDNLSEGIREWDLKPGAPKPKANQQKQD DGRGLVLPGYKYLGPFGNLDKGEFVNAAADAAAEHDKAYD QQLKAGDNPYLRYNHADADEFQERLQEDTSFGGNLGRAVFQ AKKRVLEPLGLVEEGAKTAPGKKRPVEQSPQEPDSSSGIG

				KTGQQPAKKRLNFGQTDGSESVDPQPPLGEPAPPSGVGS NTMASGGGAPMADNNEGADGVGNSSGNWHCDSTWLGDRVI TTSTRTWALPTYNNHLYKQIKSGSVGSDSNDNTYFGYSTPW GYFDNRFHCHFSPRDWQRLINNNWGFPRKRLNFKLFNIQ VKEVTVQDSTTTIANNLSTVQVFTDSEYQLPYVLGSAHQ GCLPPFPADVFMIPQYGYLTLNRDNTENPTERSSFYCLEY FPSQMLRTGNNFQFSYTFEDVFPFHSSYAHSQSLDRLMNP IDQYLYLRSRTNTGGVQFSQAGPSSMANQARNWVPGPCYR QQGWNLGSGVNRASVSFAFATATKYHLNGRDSLMNPGVAMA SNNLQGSNTYALENTLIFNSQPANPGTTATYLEGNMITNE EEIKTNPVATEEYGAVATNNQSSTTAPATGLVHNQGVLP GMVWQNRDVYLQGPWAKIPHTDGNFHPSPLMGGFGLKHP PPQILIKNTPVPADPPTTFNQAKLNSFITQYSTGQVSVEI EWELQKENS KRWNPEIQYTSNYNDPQFVDFAVNTEGVYSE PRPIGTRFLTRNL
AAV8_6VR SEQ ID NO: 146	AAV8	AAV6	VR	MAADGYLPDWLEDNLSEGIREWALKPGAPKPKANQQKQD DGRGLVLPYGYKYLGPFNGLDKGEPVNAADAAALEHDKAYD QQLQAGDNPYLRYNHADAEEFQERLQEDTSFGGNLGRAVFQ AKKRVLEPLGLVEEGAKTAPGKKRPVEPSPQRSPTSSTGI GKKGQQPARKRLNFGQTDGSESVDPQPPLGEPAPPSGVG PNTMAAGGAPMADNNEGADGVGSSSGNWHCDSTWLGDRV ITTSTRTWALPTYNNHLYKQISSASTGASNDNTYFGYSTP WGYFDNRFHCHFSPRDWQRLINNNWGFPRKRLSFKLFNI QVKEVTTNDGVTTIANNLSTIQVFTDSEYQLPYVLGSAH QGCLPPFPADVFMIPQYGYLTLNNGSQAVGRSSFYCLEYF PSQMLRTGNNFQFTYTFEDVFPFHSSYAHSQSLDRLMNP DQYLYLRSRTQNSGSAQNKDLGFSQGGPNTMANQAKNWL PGPCYRQQRVSKTKTDNNNSNFTWTGATKYHLNGRNSLAN PGIAMASHKDDKDKFFPMGILIFGKESAGASNTALDNVM LTSEEEIKTNPVATEEYGIVAVNLQSSSTD PATGTVNSQ GALPGMVWQNRDVYLQGPWAKIPHTDGNFHPSPLMGGFGL KHPPPPQILIKNTPVPADPPTTFNQSKLNSFITQYSTGQV SVEIEWELQKENS KRWNPEIQYTSNYAKSANVDFAVNTEG VYSEPRPIGTRYLTRNL
AAV8_6VRGBS SEQ ID NO: 147	AAV8	AAV6	VRGBS	MAADGYLPDWLEDNLSEGIREWALKPGAPKPKANQQKQD DGRGLVLPYGYKYLGPFNGLDKGEPVNAADAAALEHDKAYD QQLQAGDNPYLRYNHADAEEFQERLQEDTSFGGNLGRAVFQ AKKRVLEPLGLVEEGAKTAPGKKRPVEPSPQRSPTSSTGI GKKGQQPARKRLNFGQTDGSESVDPQPPLGEPAPPSGVG PNTMAAGGAPMADNNEGADGVGSSSGNWHCDSTWLGDRV ITTSTRTWALPTYNNHLYKQISSASTGASNDNTYFGYSTP WGYFDNRFHCHFSPRDWQRLINNNWGFPRKRLSFKLFNI QVKEVTTNDGVTTIANNLSTIQVFTDSEYQLPYVLGSAH QGCLPPFPADVFMIPQYGYLTLNNGSQAVGRSSFYCLEYF PSQMLRTGNNFQFTYTFEDVFPFHSSYAHSQSLDRLMNP DQYLYLRSRTQNSGSAQNKDLGFSRGSPAGMSVQAKNWL PGPCYRQQRVSKTKTDNNNSNFTWTGATKYHLNGRNSLAN PGIAMASHKDDKDKFFPMGILIFGKESAGASNTALDNVM LTSEEEIKTNPVATEEYGIVAVNLQSSSTD PATGTVNSQ GALPGMVWQNRDVYLQGPWAKIPHTDGNFHPSPLMGGFGL KHPPPPQILIKNTPVPADPPTTFNQSKLNSFITQYSTGQV SVEIEWELQKENS KRWNPEIQYTSNYAKSANVDFAVNTEG VYSEPRPIGTRYLTRNL
AAV8_6VRGH SEQ ID NO: 148	AAV8	AAV6	VRGH	MAADGYLPDWLEDNLSEGIREWALKPGAPKPKANQQKQD DGRGLVLPYGYKYLGPFNGLDKGEPVNAADAAALEHDKAYD QQLQAGDNPYLRYNHADAEEFQERLQEDTSFGGNLGRAVFQ AKKRVLEPLGLVEEGAKTAPGKKRPVEPSPQRSPTSSTGI GKKGQQPARKRLNFGQTDGSESVDPQPPLGEPAPPSGVG PNTMAAGGAPMADNNEGADGVGSSSGNWHCDSTWLGDRV ITTSTRTWALPTYNNHLYKQISSASTGASNDNTYFGYSTP WGYFDNRFHCHFSPRDWQRLINNNWGFPRKRLSFKLFNI QVKEVTTNDGVTTIANNLSTIQVFTDSEYQLPYVLGSAH QGCLPPFPADVFMIPQYGYLTLNNGSQAVGRSSFYCLEYF

				PSQMLRTGNNFQFTYTFEDVPFHSSYAHQSQSLDRLMNPLI DQYLYLNRQTQNSGSAQNKDLLF'SRGSPPAGMSVQPKNWL PGPCYRQQRVSKTKTDNNNSNFTWTGASKYNLNGRESIIN PGTAMASHKDDKDKFPMMSGVMIFFGKESAGASNTALDNVM ITDEEEIKATNPVATERFGTVAVNLQSSSTDPATGDDVHVM GALPGMVWQNRDVYLQGPWAKIPHTDGNFHPSPMLMGFG LKHPPQILIKNTPVPADPPTTFNQSKLNSFITQYSTGQV SVEIEWELQKENS KRWNPEIQYTSNYAKSANVDFAVNTEG VYSEPRPIGTRYLTRNL
AAV8_7VRGBS SEQ ID NO: 149	AAV8	AAV7	VRGBS	MAADGYLPDWLEDNLSEGIREWALKPGAPKPKANQQKQD DGRGLVLPGYKYLGPFNGLDKGEPVNAADAAALEHDKAYD QQLQAGDNPYLRYNHADADEFQERLQEDTSFGGNLGRAVFQ AKKRVLEPLGLVEEGAKTAPGKKRPVEPSPQSPDSSSTGI GKKGQQPARKRLNFGQTGDSESVDPDQPLGEPPAAPSGVG PNTMAAGGGAPMADNNEGADGVGSSSGNWHCDSTWLGDRV ITTSTRTWALPTYNNHLYKQISSSETAGSTNDNTYFGYSTP WGYFDFNRFHCHFSPRDWQRLINNNWGFPRKLSFKLFNI QVKEVTTNDGVTTIANNLSTIQVFTDSEYQLPYVLGSAH QGCLPPFPADVFMIPQYGYLTLNNGSQAVGRSSFYCLEYF PSQMLRTGNNFQFTYTFEDVPFHSSYAHQSQSLDRLMNPLI DQYLYLNRQTQNSGSAQNKDLLF'SRGSPPAGMSVQPKNWL LPGPCYRQQRVSKTKTDQNNNSNFAWTGATKYHLNGRNSLA NPGIAMATHKDDERDFPSSGILIFGKTGATNKTTLNENVM LTSEEEIKTNPVATEEYGVSSNLQAANTAAQTGTVNSQ GALPGMVWQNRDVYLQGPWAKIPHTDGNFHPSPMLMGFG LKHPPQILIKNTPVPADPPTTFNQSKLNSFITQYSTGQV SVEIEWELQKENS KRWNPEIQYTSNFEKQGTGVDFAVNTEG VYSEPRPIGTRYLTRNL
AAV8_7VRGH SEQ ID NO: 150	AAV8	AAV7	VRGH	MAADGYLPDWLEDNLSEGIREWALKPGAPKPKANQQKQD DGRGLVLPGYKYLGPFNGLDKGEPVNAADAAALEHDKAYD QQLQAGDNPYLRYNHADADEFQERLQEDTSFGGNLGRAVFQ AKKRVLEPLGLVEEGAKTAPGKKRPVEPSPQSPDSSSTGI GKKGQQPARKRLNFGQTGDSESVDPDQPLGEPPAAPSGVG PNTMAAGGGAPMADNNEGADGVGSSSGNWHCDSTWLGDRV ITTSTRTWALPTYNNHLYKQISSSETAGSTNDNTYFGYSTP WGYFDFNRFHCHFSPRDWQRLINNNWGFPRKLSFKLFNI QVKEVTTNDGVTTIANNLSTIQVFTDSEYQLPYVLGSAH QGCLPPFPADVFMIPQYGYLTLNNGSQAVGRSSFYCLEYF PSQMLRTGNNFQFTYTFEDVPFHSSYAHQSQSLDRLMNPLI DQYLYLNRQTQNSGSAQNKDLLF'SRGSPPAGMSVQPKNWL LPGPCYRQQRVSKTKTDQNNNSNFAWTGATKYHLNGRNSLV NPGVAMATHKDDERDFPSSGVLIFGKTGATNKTTLNENVM MTNEEEIRPTNPVATEEYGVSSNLQAANTAAQTQVNNQ GALPGMVWQNRDVYLQGPWAKIPHTDGNFHPSPMLMGFG LKHPPQILIKNTPVPADPPTTFNQSKLNSFITQYSTGQV SVEIEWELQKENS KRWNPEIQYTSNFEKQGTGVDFAVNTEG VYSEPRPIGTRYLTRNL
Anc110_8VR SEQ ID NO: 151	Anc110	AAV8	VR	MAADGYLPDWLEDNLSEGIREWDLKPGAPKPKANQQKQD DGRGLVLPGYKYLGPFNGLDKGEPVNAADAAALEHDKAYD QQLKAGDNPYLRYNHADADEFQERLQEDTSFGGNLGRAVFQ AKKRVLEPLGLVEEGAKTAPGKKRPVEQSPQEPDSSSGIG KTGQQPAKKRLNFGQTGDSESVDPDQPLGEPPAAPSGVGS NTMASGGGAPMADNNEGADGVGNSSGNWHCDSTWLGDRV ITTSTRTWALPTYNNHLYKQISSSETAGSTNDNTYFGYSTP WGYFDFNRFHCHFSPRDWQRLINNNWGFPRKLSFKLFNI QVKEVTQNEGKTIANNLTSTVQVFTDSEYQLPYVLGSAH QGCLPPFPADVFMIPQYGYLTLNNGSQAVGRSSFYCLEYF PSQMLRTGNNFQFTYTFEDVPFHSSYAHQSQSLDRLMNPLI DQYLYLNRQTQNSGSAQNKDLLF'SRGSPPAGMSVQPKNWL PGPCYRQQRVSTTGQNNNSNFAWTAAATKYHLNGRDLSLMN PGVAMASHKDDERDFPSSGVLIFGKQNAARDNADYSDVM ITNEEEIKTNPVATEEYGAADNLQQNTAPQIGLVHNQ GVLPGMVWQNRDVYLQGPWAKIPHTDGNFHPSPMLMGFG

				LKHPPPQILIKNTPVPADPPTTFNQAKLNSFITQYSTGQV SVEIEWELQKENS KRWNPEIQYTSNYYKSTSVDFAVNTEG VYSEPRPIGTRFLTRNL
Anc110_8VRGBS SEQ ID NO: 152	Anc110	AAV8	VRGBS	MAADGYLPDWLEDNLSEGIREWDLKPGAPKPKANQQKQD DGRGLVLPGYKYLGPFGNLDKGEFVNAAADAAAEHDKAYD QQLKAGDNPYLRYNHADADEFQERLQEDTSFGGNLGRAVFQ AKKRVLEPLGLVEEGAKTAPGKKRPVEQSPQEPDSSSGIG KTGQQPAKKRLNFGQTGDSESVDPQPPLGEPPAAPSGVGS NTMASGGGAPMADNNEGADGVGNSSGNWHCDSTWLGDRVI TTSTRTWALPTYNNHLYKQISNGTSGGATNDNTYFGYSTP WGYFDFNRFHCHFSPRDWQRLINNNWGFPRKRLNFKLFNI QVKEVTQNEGKTIANNLTSTVQVFTDSEYQLPYVLGSAH QGCLPPFPADVFMIQYGYLTLNNGSQAVGRSSFYCLEYF PSQMLRTGNNFQFSYTFEDVPFHSSYAHSQSLDRLMNP LI DQYLYFLSRTQTGGTANTQTLOFSQGGPNTMANQARNWV PGPCYRQQRVSTTGQNNNSNFAWTAATKYHLNGRDSLMN PGVAMASHKDDEERFFPSNGVLIFGKQNAARDNADYSDVM ITNEEEIKTNPVATEEYGAVDNLQQQNTAPQIGLVHNQ GVLPGMVWQNRDVYLQGP I WAKIPHTDGNFHPSPLMGGFG LKHPPPQILIKNTPVPADPPTTFNQAKLNSFITQYSTGQV SVEIEWELQKENS KRWNPEIQYTSNYYKSTSVDFAVNTEG VYSEPRPIGTRFLTRNL
AAV7_rh10VR SEQ ID NO: 153	AAV7	rh10	VR	MAADGYLPDWLEDNLSEGIREWDLKPGAPKPKANQQKQD NGRGLVLPGYKYLGPFGNLDKGEFVNAAADAAAEHDKAYD QQLKAGDNPYLRYNHADADEFQERLQEDTSFGGNLGRAVFQ AKKRVLEPLGLVEEGAKTAPAKKRPVEPSQSPDSSSTGI GKKGQQPARKRLNFGQTGDSESVDPQPPLGEPPAAPSSVG SGTVAAGGGAPMADNNEGADGVGNASGNWHCDSTWLGDRV ITTSTRTWALPTYNNHLYKQISNGTSGGSTNDNTYFGYST PWGYFDFNRFHCHFSPRDWQRLINNNWGFPRKRLNFKLFNI IQVKEVTQNEGKTIANNLTSTIQVFSSEYQLPYVLGSA HQGCLPPFPADVFMIQYGYLTLNNGSQSVGRSSFYCLEY FPSQMLRTGNNFEFSYSFEDVPFHSSYAHSQSLDRLMNP LI IDQYLYLARTQSTGGTAGTQQLQFYQGGPSTMAEQAKNW LPGPCFRQQRVSTTSLQNNNSNFAWTGATKYHLNGRNSLV NPGVAMATHKDDEERFFPSNGVLIFGKQAGKDNVDYSSV LMTNEEEIRTPNPVATEEYGIVADNLQQQNAAPIVQVVNN QGALPGMVWQNRDVYLQGP I WAKIPHTDGNFHPSPLMGGFG GLKHPPPQILIKNTPVPANPPEVFTPAKFASFITQYSTGQ VSVEIEWELQKENS KRWNPEIQYTSNYYKSTNVDFAVDSQ GVYSEPRPIGTRYLTRNL
AAV7_rh10VRGBS SEQ ID NO: 154	AAV7	rh10	VRGBS	MAADGYLPDWLEDNLSEGIREWDLKPGAPKPKANQQKQD NGRGLVLPGYKYLGPFGNLDKGEFVNAAADAAAEHDKAYD QQLKAGDNPYLRYNHADADEFQERLQEDTSFGGNLGRAVFQ AKKRVLEPLGLVEEGAKTAPAKKRPVEPSQSPDSSSTGI GKKGQQPARKRLNFGQTGDSESVDPQPPLGEPPAAPSSVG SGTVAAGGGAPMADNNEGADGVGNASGNWHCDSTWLGDRV ITTSTRTWALPTYNNHLYKQISNGTSGGSTNDNTYFGYST PWGYFDFNRFHCHFSPRDWQRLINNNWGFPRKRLNFKLFNI IQVKEVTQNEGKTIANNLTSTIQVFSSEYQLPYVLGSA HQGCLPPFPADVFMIQYGYLTLNNGSQSVGRSSFYCLEY FPSQMLRTGNNFEFSYSFEDVPFHSSYAHSQSLDRLMNP LI IDQYLYLARTQSTGGTAGTQQLQFYQAGPNNMSAQAKNW LPGPCFRQQRVSTTSLQNNNSNFAWTGATKYHLNGRNSLV NPGVAMATHKDDEERFFPSNGVLIFGKQAGKDNVDYSSV LMTNEEEIRTPNPVATEEYGIVADNLQQQNAAPIVQVVNN QGALPGMVWQNRDVYLQGP I WAKIPHTDGNFHPSPLMGGFG GLKHPPPQILIKNTPVPANPPEVFTPAKFASFITQYSTGQ

				VSVEIEWELQKENS KRWNPEIQYTSNYYKSTNVDFAVDSQ GVYSEPRPIGTRYLTRNL
AAV7_rh10VRGH SEQ ID NO: 155	AAV7	rh10	VRGH	MAADGYLPDWLEDNLSEGIREWWDLKPGAPKPKANQQKQD NGRGLVLPGYKYLGPFGNLDKGEPVNAADAAALEHDKAYD QQLKAGDNPYLRYNHADADEFQERLQEDTSFGGNLGRAVFQ AKKRVLEPLGLVEEGAKTAPAKKRPVEQSPQSPDSSSTGI GKKGQQPARKRLNFGQTGDSSESVPDPQPLGEPPAAPSSVG SGTVAAGGGAPMADNNEGADGVGNASGNWHCDSTWLGDRV ITTSTRTWALPTYNNHLYKQISNGTSGGSTNDNTYFGYST PWGYFDNRFHCHFSPRDWQRLINNNWGFPRPKLRFKLFNI IQKEVTQNEGKTIANNLSTIQVFSDEYQLPYVLGSAH HQGLPPFPADVFMIPQYGYLTLNNGSQSVGRSSFYCLEY FPSQMLRTGNNFESYSFEDVPFHSSYAHSSQLDRLMNLPL IDQYLYLSRTQSTGGTAGTQQLLFSQAGPNNMSAQAKNW LPGPCYRQQRVSTTSLQNNNSNFAWTGATKYHLNGRDSL NPGVAMATHKDDEERFFPSSGVLMEFGKQAGKDNVDYSSV MLTSEEEIKTTNPVATEQYGVVADNLQQQNAAPIVGAVNS QGALPGMVWQNRDVYLQGP I WAKIPHTDGNFHP S PLMGGF GLKHPPPQILIKNTPVPANPPEVFTPAKFASFITQYSTGQ VSVEIEWELQKENS KRWNPEIQYTSNYYKSTNVDFAVDSQ GVYSEPRPIGTRYLTRNL
AAV9_rh10VRGBS SEQ ID NO: 156	AAV9	rh10	VRGBS	MAADGYLPDWLEDNLSEGIREWWDLKPGAPQPKANQQHQD NARGLVLPGYKYLGPFGNLDKGEPVNAADAAALEHDKAYD QQLKAGDNPYLRYNHADADEFQERLKEDTSFGGNLGRAVFQ AKKRLLEPLGLVEEAAKTAPGKKRPVEQSPQEPDSSAGIG KSGAQPAKKRLNFGQTGDTESVPDPQPIGEPPAAPSGVGS LTMASGGGAPVADNNEGADGVGSSSGNWHCDSQWLGDRVI TTSTRTWALPTYNNHLYKQISNGTSGGSTNDNAYFGYSTP WGYFDNRFHCHFSPRDWQRLINNNWGFPRPKLRFKLFNI QVKEVTQNEGKTIANNLSTVQVFTDSYQLPYVLGSAH EGCLPPFPADVFMIPQYGYLTLNNGSQAVGRSSFYCLEYF PSQMLRTGNNFQFSYEFENVFPFHSSYAHSSQLDRLMNLPL IDQYLYLSKTIISTGGTAGTQQLKFSVAGPSNMAVQGRNYI PGPSYRQQRVSTTSLQNNNSNFAWTGASSWALNCRNSLMN PGPAMATHKDDEERFFPSSGSLIFGKQAGKDNVDYSSVM ITNEEEIKTTNPVATESYGVVADNLQQQNAAPIVGWVQNG GILPGMVWQDRDVYLQGP I WAKIPHTDGNFHP S PLMGGF MKHPPPQILIKNTPVPADPPTAFNKDKLNSFITQYSTGQV SVEIEWELQKENS KRWNPEIQYTSNYYKSTNVDFAVNTEG VYSEPRPIGTRYLTRNL
AAV9_rh10VRGH SEQ ID NO: 157	AAV9	rh10	VRGH	MAADGYLPDWLEDNLSEGIREWWDLKPGAPQPKANQQHQD NARGLVLPGYKYLGPFGNLDKGEPVNAADAAALEHDKAYD QQLKAGDNPYLRYNHADADEFQERLKEDTSFGGNLGRAVFQ AKKRLLEPLGLVEEAAKTAPGKKRPVEQSPQEPDSSAGIG KSGAQPAKKRLNFGQTGDTESVPDPQPIGEPPAAPSGVGS LTMASGGGAPVADNNEGADGVGSSSGNWHCDSQWLGDRVI TTSTRTWALPTYNNHLYKQISNGTSGGSTNDNAYFGYSTP WGYFDNRFHCHFSPRDWQRLINNNWGFPRPKLRFKLFNI QVKEVTQNEGKTIANNLSTVQVFTDSYQLPYVLGSAH EGCLPPFPADVFMIPQYGYLTLNNGSQAVGRSSFYCLEYF PSQMLRTGNNFQFSYEFENVFPFHSSYAHSSQLDRLMNLPL IDQYLYLSRTQSTGGTAGTQQLLFSQAGPNNMSAQAKNW LPGPCYRQQRVSTTSLQNNNSNFAWTGATKYHLNGRDSL NPGVAMATHKDDEERFFPSSGVLMEFGKQAGKDNVDYSSVM LTSEEEIKTTNPVATEQYGVVADNLQQQNAAPIVGAVNSQ GALPGMVWQDRDVYLQGP I WAKIPHTDGNFHP S PLMGGF MKHPPPQILIKNTPVPADPPTAFNKDKLNSFITQYSTGQV SVEIEWELQKENS KRWNPEIQYTSNYYKSTNVDFAVNTEG VYSEPRPIGTRYLTRNL

Example 3**Analysis of Engineered Chimeric Capsid Proteins**

[00210] DNaseI resistant virus titer analysis was used to characterize the virus packaging efficiency and production of engineered AAV variants. Five million HEK293 cells were seeded on a 10 cm cell culture dish and incubated overnight at 37°C. AAV plasmid transfection was performed using a conventional calcium phosphate transfection protocol using ~20µg total DNA. Cells were harvested at 72 hours post-transfection and centrifuged at 1,000xg for 15 minutes. Lysis buffer was added to the cell pellet and samples were subjected to three freeze-thaw cycles with the addition of DNaseI in the final thaw, followed by an incubation for 30 minutes. Crude expressed AAV virus was collected in the supernatant after centrifugation to remove cellular debris at 10,000xg for 30 minutes. EDTA was added to stop the DNaseI activity and then the expressed AAV was incubated at 95°C for 15 minutes to further inactivate the DNase. Quantitative PCR was performed using a LightCyclerII according to the Fast Expression Taqman PCR mix protocol provided from the manufacturer (Roche). Virus titers were calculated and compared in vg/µL amounts. Each virus titer measurement was performed 3 times. The results of this DNaseI resistant titer analysis are provided below in Table 4 and are ordered from highest to lowest titer.

[00211] Table 4

Crude AAV	AVERAGE	STDEV	Crude AAV	AVERAGE	STDEV
AAV9_rh10VRGBS	6.60E+08	5.32E+07	AAV4_2VRGH	2.17E+07	5.87E+06
AAV7_rh10VRGBS	5.58E+08	7.27E+07	AAVanc110_9GH	2.13E+07	1.16E+07
AAV7_rh10VR	4.72E+08	1.79E+07	AAVmo_8GH	2.13E+07	6.34E+06
AAV7_rh10VRGH	4.19E+08	3.49E+07	AAVbo_8VR	2.12E+07	2.84E+06
Anc110_8VRGBS	3.68E+08	3.60E+07	AAV4	2.04E+07	5.02E+06
AAV9_rh10VRGH	3.31E+08	4.11E+07	AAV8_9VRGBS	2.02E+07	6.82E+06
AAV8_5VR	2.47E+08	2.72E+07	AAV6.2_9GH	1.93E+07	4.25E+06
AAV8	2.10E+08	8.83E+07	AAVmo_2VRGH	1.69E+07	3.95E+06

AAV2	1.71E+08	1.01E+08	AAV6	1.67E+07	3.40E+06
AAV8_7VRGBS	1.58E+08	8.57E+06	AAV5	1.64E+07	2.35E+06
AAV7	1.50E+08	5.71E+07	AAV8_anc110GH	1.54E+07	2.82E+06
AAV8_6VRGH	1.20E+08	9.65E+06	AAV6.2_9VRGBS	1.50E+07	3.92E+06
AAV8_6VR	1.14E+08	1.21E+07	AAVbo_WT	1.38E+07	1.49E+07
AAV8_6VRGBS	1.12E+08	9.44E+06	AAV8_9VRGBS.MK	1.25E+07	6.73E+06
AAV8_7VRGH	1.03E+08	1.76E+07	AAV8_9VRGBS.PA	1.12E+07	3.51E+06
Anc110_5VR	9.13E+07	1.27E+07	AAV8_6VRGBS	9.42E+06	5.27E+05
AAV9	7.63E+07	2.64E+07	AAVbo_9GH	9.35E+06	6.75E+06
AAV12	6.76E+07	3.18E+07	AAV7_9VRGBS	9.14E+06	2.24E+06
AAVanc110_9VRHS	6.64E+07	5.38E+07	AAVmo_2VR	8.83E+06	3.08E+06
AAVmo_9VRGH	6.18E+07	1.12E+07	AAV8_9RGRS.SA	8.65E+06	2.21E+06
AAVanc110_9VR.PHP	6.06E+07	1.37E+07	AAV8_9RGRS.ST	8.20E+06	5.97E+06
AAVmo_8VR	5.04E+07	1.09E+07	AAV8_6VR	7.76E+06	3.65E+06
AAVmo_9VR	4.11E+07	8.80E+06	AAV8_9VRGBS.MI	7.72E+06	1.59E+05
AAV8_9VRGBS.SmT	4.07E+07	2.11E+07	AAV8_9VRGBS.PHP	7.55E+06	2.52E+06
AAV4_9GH	3.97E+07	6.45E+06	AAVanc110	7.05E+06	2.26E+06
AAVmo_9GH	3.67E+07	6.29E+06	AAV8_9VRGBS.MA	5.36E+06	5.02E+05
AAVbo_8VRGH	3.49E+07	1.12E+07	AAVanc110_9VRGBS	5.36E+06	1.12E+06
AAVmo1_WT	3.34E+07	7.17E+06	AAVrh10_9GH	5.13E+06	7.73E+05
AAV4_2GH	3.21E+07	1.06E+07	AAVanc110_8VR	4.77E+06	1.60E+06
AAV4_8VR	3.16E+07	4.68E+06	AAVanc110_8GH	4.64E+06	1.32E+06
AAVanc110_9GH.PHP	3.12E+07	4.10E+06	AAVanc80L65	4.07E+06	3.85E+05
AAVmo_8VRGH	3.11E+07	1.10E+07	AAV8_9VRGH.PHP	3.66E+06	2.33E+06
AAVbo_9VR	2.97E+07	1.20E+07	AAVbo_8GH	2.80E+06	8.36E+05

AAV4_8GH	2.90E+07	3.48E+06	AAV4_9VR	2.14E+06	N/A
AAVrh10	2.88E+07	5.51E+06	AAV9_anc110VR	1.65E+06	3.58E+05
AAVanc110_9VR	2.76E+07	4.72E+06	AAV8_9VRGH	1.48E+06	2.37E+05
AAV4_9VRGH	2.68E+07	3.75E+06	AAV8_anc110VR	1.21E+06	4.89E+05
AAV4_2VR	2.32E+07	6.96E+06	AAV9_anc110GH	1.19E+06	3.07E+05
AAV8_6VRGH	2.32E+07	4.88E+06	Bce36_9GH	8.44E+05	1.96E+05
AAV4_8VRGH	2.26E+07	2.97E+06	No Cap Control	5.03E+05	3.81E+04
			AAVbo_9VRGH	1.13E+05	7.52E+03

[00212] The results in Table 4 demonstrate that AAV virus comprising the engineered chimeric AAV capsids of the present invention may be successfully produced, although with varying titers.

[00213] Next, the ability of AAV particles comprising engineered chimeric capsid proteins of the present invention to transduce cultured cells *in vitro* was determined. HEK293 and HepG2 cells were seeded at 5×10^4 cells/well on a 96 well plate and incubated overnight. Etoposide was added on the day of infection to a final concentration of 4 μ M and 20 μ M for HEK293 and HepG2 cells, respectively. Different crude AAV particles were added at an MOI of 2000 and transduction data was measured in relative luciferase units (RLU) 72 hours post-infection. The efficiency of transduction data is provided in Table 5 below and is ordered from the highest to the lowest transduction ability. As expected, recombinant AAV2 (positive control) exhibited the best *in vitro* transduction for both cell lines and most of the engineered chimeric AAVs tested were shown to have positive ($RLU > 10^4$) *in vitro* transduction properties. These results demonstrate that not only may AAV particles comprising the chimeric engineered capsid proteins of the present invention be successfully produced, but those novel recombinant AAV particles are functional in that they are capable of transducing either HEK293 or HepG2 cells with varying efficiencies.

Table 5

HEK293

HepG2

<u>AAV</u>	<u>AVRG</u>	<u>ERROR</u>
AAV2	9.07E+06	8.54E+05
AAVBce36	2.11E+05	7.57E+04
AAV7	1.60E+05	6.77E+04
AAV8_9VRGBS	1.06E+05	6.65E+03
AAV8_anc110GH	9.45E+04	8.35E+03
AAVanc110	7.89E+04	6.14E+04
AAVanc110_8GH	6.96E+04	1.01E+04
AAV5	6.33E+04	4.82E+03
AAVboWT	5.90E+04	4.36E+03
AAV9	3.85E+04	3.93E+03
AAV8_9VRGBS.ST	2.96E+04	4.45E+03
AAV8	2.93E+04	1.80E+03
AAV6.2_9GH	2.61E+04	4.71E+03
AAV8_9VRGH.PHP	2.45E+04	1.43E+04
AAV8_9VRGBS.PA	1.64E+04	5.58E+03
AAV6	1.64E+04	2.29E+03
AAVanc110_9VR	1.49E+04	3.33E+03
AAV8_9VRGBS.SA	1.37E+04	1.65E+03
rh10	1.09E+04	1.79E+03
AAVanc110_9VRHS	1.02E+04	2.26E+03
AAVanc110_8VR	9.67E+03	1.76E+03
AAV8_9VRGBS.MI	7.13E+03	3.07E+03
AAV12	5.93E+03	1.97E+03
AAV8_9VRGBS.PHP	5.70E+03	3.59E+03
AAV9_anc110VR	5.42E+03	4.20E+03
AAV8_9VRGH	5.24E+03	4.99E+02
AAVanc110_9VRGBS	3.77E+03	2.09E+03
AAV8_anc110VR	3.22E+03	4.17E+03
AAV9_anc110GH	2.57E+03	1.81E+03
AAV7_rh10VRGH	1.70E+03	3.44E+02
AAVanc110_9GH	1.51E+03	9.65E+01
AAVbo_9GH	1.01E+03	9.05E+01
AAV_Anc110	9.61E+02	1.78E+02
AAVmo_9VRGH	7.29E+02	3.87E+02
AAVanc110_9VR.PHP	6.95E+02	8.71E+02
no Cap control	6.85E+02	7.13E+01
AAVrh10_9VRGBS	6.55E+02	6.87E+02
AAV8_7VRGH	4.93E+02	4.72E+02
AAV7_9GH	4.93E+02	1.35E+02
AAV8_9VRGBS.MA	4.91E+02	4.35E+01
AAV8_6VR	4.52E+02	1.22E+02
AAV4_2VR	4.06E+02	2.40E+02

<u>AAV</u>	<u>AVRG</u>	<u>ERROR</u>
AAV2	2.12E+07	1.59E+06
AAV9_anc110VR	2.41E+06	2.01E+05
AAV8_anc110GH	6.50E+05	3.64E+03
AAV8_9VRGBS	6.14E+05	8.37E+03
AAVanc110_8GH	6.07E+05	5.75E+04
AAV9	4.14E+05	3.22E+03
AAV8_anc110VR	2.76E+05	5.89E+04
AAV5	2.33E+05	3.16E+03
AAVBce36	1.28E+05	4.46E+04
AAV7	9.96E+04	4.20E+04
AAVanc110_8VR	7.26E+04	3.83E+03
AAV9_anc110GH	6.36E+04	1.45E+04
rh10	5.83E+04	1.63E+03
AAV6	4.13E+04	4.44E+03
AAV8_9VRGH.PHP	3.89E+04	2.22E+04
AAV8	3.54E+04	4.86E+03
AAVanc110_9VR	3.35E+04	3.91E+03
AAVanc110	3.18E+04	2.32E+04
AAVanc110_9GH	2.42E+04	5.33E+02
AAV7_rh10VRGH	8.44E+03	7.72E+02
AAV_Anc110	8.14E+03	5.22E+01
AAV8_9VRGBS.PHP	4.19E+03	2.47E+03
AAV8_6VR	3.87E+03	8.42E+02
AAVanc110_9VR.PHP	2.66E+03	3.94E+02
AAVboWT	2.58E+03	8.17E+02
no Cap control	2.18E+03	3.70E+02
AAV4_2GH	1.61E+03	6.60E+02
AAVbo_9GH	1.58E+03	6.33E+02
Anc110_8VRGBS	1.45E+03	4.08E+02
AAV9_rh10VRGH	1.30E+03	2.03E+02
AAV9_rh10VRGBS	1.26E+03	1.10E+02
AAVrh10_9VRGBS	7.92E+02	6.42E+02
AAVanc110_9GH.PHP	6.78E+02	2.18E+02
AAV8_7VRGH	5.35E+02	1.13E+02
AAV4_2VRGH	4.85E+02	3.04E+02
AAVmo_8GH	4.71E+02	1.01E+02
AAVmo_9GH	3.92E+02	9.89E+01
AAV4_9VRGH	3.76E+02	1.07E+02
AAV4_8VRGH	3.10E+02	1.40E+01
AAV7_9GH	3.01E+02	9.52E+01
AAV4_2VR	2.83E+02	8.36E+00
AAV8_6VRGBS	2.63E+02	5.38E+01

AAV9_rh10VRGH	3.94E+02	1.44E+02	AAVmo_8VRGH	2.52E+02	1.48E+01
Neg	3.19E+02	7.49E+01	AAVBce36_9VRGBS	2.44E+02	2.03E+02
AAV4_2GH	3.04E+02	5.38E+01	AAVBce41_8VRGBS	2.33E+02	5.34E+01
Anc110_8VRGBS	3.02E+02	5.09E+01	AAV4_9GH	2.10E+02	3.50E+01
AAV4_8VRGH	2.69E+02	2.62E+01	AAV4_8VR	1.97E+02	1.38E+01
AAV8_9VRGBS.MK	2.53E+02	1.10E+01	AAV4_8GH	1.89E+02	2.26E+01
AAVanc110_9GH.PHP	2.49E+02	9.90E+01	AAV8_7VRGBS	1.86E+02	5.92E+01
AAV4_2VRGH	2.49E+02	1.03E+02	AAVbo_8VR	1.57E+02	6.67E-01
AAVmo_8GH	2.49E+02	2.25E+01	AAVmo_8VR	1.53E+02	5.21E+00
AAV4_8GH	2.34E+02	6.31E+01	AAVbo_9VR	1.43E+02	1.07E+01
AAVmo_9GH	2.28E+02	5.11E+01	AAVmo_9VRGH	1.39E+02	6.77E+00
AAVBce36_9VRGBS	2.06E+02	1.55E+02	AAVbo_8VRGH	1.32E+02	1.04E+01
AAVmo_8VRGH	1.92E+02	1.36E+01	AAVmo_9VR	1.26E+02	1.62E+01
AAV9_rh10VRGBS	1.62E+02	9.99E+00	Anc110_5VR	1.05E+02	1.51E+01
AAVmo_8VR	1.60E+02	3.38E+01	AAV8_5VR	8.55E+01	1.64E+01
AAVBce41_8VRGBS	1.58E+02	5.01E+01	AAV7_rh10VR	6.67E+01	2.38E+01
AAV4_9VRGH	1.52E+02	1.21E+01	AAV7_rh10VRGBS	5.13E+01	8.74E+00
Anc110_5VR	1.49E+02	5.87E+00	AAV8_6VRGH	4.75E+01	1.11E+01
AAVbo_9VR	1.37E+02	1.21E+01			
AAV4_9GH	1.33E+02	9.27E+00			
AAV8_7VRGBS	1.29E+02	1.04E+01			
AAV8_7VRGH	1.29E+02	2.00E+01			
AAV4_8VR	1.25E+02	8.36E+00			
AAVbo_8VR	1.22E+02	2.00E+00			
AAVmo_9VR	1.21E+02	1.09E+01			
AAVbo_8VRGH	1.21E+02	1.31E+01			
AAV8_5VR	1.11E+02	1.94E+01			
AAV7_rh10VR	1.10E+02	1.28E+01			
AAV8_6VRGBS	9.60E+01	7.39E+00			
AAV6.2_9VRGBS	9.53E+01	1.22E+01			
AAV7_rh10VRGBS	8.40E+01	3.46E+00			
AAV8_6VRGH	7.84E+01	5.62E+00			

[00214] Next, the ability of the certain engineered AAV particles of the present invention to transduce glial cells was investigated. Human glioblastoma U87MG cells were seeded at 5×10^4 cells/well on a 96 well plate and incubated overnight. Etoposide was added on the day of infection to a final concentration of $4 \mu\text{M}$. Different crude AAV particles were added at an MOI of 2000 and transduction data was measured in RLU units 72 hours post-infection. As shown in

Table 6, engineered AAV variants retained the ability to transduce glial cells with various efficiencies.

Table 6: U87MG glial cell transduction data:

AAV	AVERAGE	ERROR
AAV2	3.89E+06	5.32E+05
AAV8_9VRGBS	1.34E+05	1.74E+04
AAV5	9.51E+04	1.26E+04
AAV8_anc110GH	6.05E+04	8.85E+03
AAV9	2.52E+04	1.68E+03
anc110_9VR	1.05E+04	1.47E+03
anc110_9VR.PHP	2.53E+03	5.81E+02
AAVanc110_8VR	1.38E+03	5.67E+01
anc110_9GH	1.19E+03	3.05E+02
anc110_9GH.PHP	9.22E+02	2.39E+01
anc110_8GH	4.55E+02	1.57E+01

[00215] Neutralization of the engineered AAV particles of the present invention by antibodies in human serum was also investigated. HEK293T cells were seeded in density 5×10^4 cells/well and incubated overnight. Purified rAAVs were diluted to final titer of 2×10^6 vg/uL and mixed with serial dilutions (0 – 10mg/mL) of IVIG for 1 hour. Recombinant AAVs were added onto HEK293T cells using MOI of 1000 and incubated in 37°C. Seventy-two hours post-infection, IVIG neutralization was analyzed based on relative luciferase unit (RLU) reading. No etoposide was used in this study. The results are provided in Table 7. As expected, AAV2 transduction (positive control) was abolished by the addition of human IVIG. In contrast, certain of the engineered AAVs tested exhibited IVIG resistant properties.

Table 7: IVIG Neutralization Data:

IVIG (mg/ml)	10	5	2.5	1.25	0.625	0.312	0.156	0.0781	0.039	0.01
AAV2	44	55	46	65	73	76	1351	11462	71835	681882
AAV5	60	56	46	39	725	4872	9210	12821	14065	14154
AAV8	29	45	49	39	139	2029	13742	32098	49373	66066

AAV9	27	40	53	59	45	444	1462	6820	13934	23233
AAVrh10	45	37	59	39	31	537	3429	13190	25726	34653
Bce36	32	41	18	78	64	340	3398	21732	39388	64691
AAV8_9VRGBS	57	46	36	42	60	236	1966	7223	12080	20501
AAVanc110_9VR	44	55	57	40	45	126	363	1473	2161	3462
Bce41	36	33	47	44	58	643	3644	10161	15275	25008
Bba41	40	38	55	47	45	57	1096	4459	6661	8308
Bba42	27	17	32	31	47	42	260	2125	2964	4203
Bba43	44	35	42	36	28	288	2130	6457	12626	18284
AAV6.2-9GH	26	35	45	37	39	82	159	2691	12087	38922
AAV8-9VRGH	39	31	37	39	38	110	1408	6235	10488	16625
AAV7	38	44	26	44	58	276	500	2891	5432	10224
AAV6	47	39	103	206	81	101	577	2319	5734	9976
AAVBba30	32	34	34	57	48	45	349	1688	3335	6203
AAVanc110	42	39	14	43	36	105	462	1354	2822	4005
AAVanc110-9VRGBS	29	37	35	39	48	39	172	819	2172	2922

[00216] Some of the engineered AAV particles have their GBS region substituted with a GBS region donated from a different AAV capsid sequence. The effect of this substitution on glycan binding properties was investigated. For example, galactose binding was investigated in an engineered AAV variant having the AAV8 backbone sequence with a substitution of all of the variable regions (VR I through VR IX) plus the GBS region from AAV9 (AAV8_9VRGBS). Pro5, Lec1 and Lec2 cells were seeded at 5×10^4 cells/well on a 96 well plate and incubated overnight, then chilled at 4°C for 30 minutes prior to infection. Recombinant AAV particles were added at an MOI of 10,000 and incubated at 4°C for another 2 hours. After two washes with DMEM media, cells were incubated at 37°C in DMEM, 10% fetal calf sera for 48 hours and RLU were measured. As shown in Table 8, this particular engineered AAV variant exhibited the ability to bind to Lec2, similar to wild type AAV9, demonstrating that the glycan binding properties of an AAV virus comprising an engineered chimeric AAV capsid protein of the present invention are determined by the donor source of the GBS region swapped into the backbone sequence.

Table 8: Glycan Binding Data

	Pro5 (sialic acid)		Lec1 (mannose)		Lec2 (galactose)	
	AVERAGE	STDEV	AVERAGE	STDEV	AVERAGE	STDEV
AAV9	4.21E+03	1.76E+03	3.56E+03	6.82E+02	1.22E+05	2.90E+04
AAV8_9VRGH	3.11E+03	9.29E+02	1.79E+03	4.04E+02	1.19E+05	2.34E+04

AAV8_9VRGBS	3.72E+03	7.71E+02	3.36E+03	1.14E+03	1.09E+05	4.21E+04
AAV6.2_9GH	5.58E+02	1.06E+02	1.90E+02	7.55E+01	7.70E+04	2.11E+04
AAVanc110_9VRGBS	4.26E+02	1.05E+02	3.71E+02	6.52E+01	2.67E+04	7.26E+03
AAV8	4.74E+03	1.46E+03	6.37E+03	4.70E+03	1.34E+04	7.21E+03
AAVBce41	8.79E+03	2.50E+03	4.74E+03	2.17E+03	8.20E+03	2.20E+03
AAVanc110	1.48E+03	7.36E+02	6.61E+02	2.88E+02	2.25E+03	2.32E+02
AAVBce36	1.24E+03	3.10E+02	2.15E+03	1.60E+03	1.52E+03	1.92E+02
AAV7	2.57E+02	8.02E+01	3.66E+02	2.08E+02	6.10E+02	2.67E+02
AAVanc110_9VR	3.37E+02	9.74E+01	2.85E+02	6.08E+01	5.77E+02	2.20E+02
AAV6	3.91E+03	1.18E+03	4.12E+02	1.75E+02	5.62E+02	7.22E+01
AAVanc110_9GH	4.25E+01	7.55E+00	4.05E+01	1.34E+01	1.64E+02	1.26E+02
AAVBba43	5.20E+01	1.34E+01	5.45E+01	8.70E+00	1.31E+02	5.94E+01
AAV5	3.66E+04	3.45E+03	1.84E+02	5.09E+01	1.28E+02	3.07E+01
AAVBba42	7.30E+01	1.37E+01	6.00E+01	2.16E+01	1.16E+02	2.03E+01
AAVBba41	7.25E+01	1.33E+01	5.55E+01	8.54E+00	9.60E+01	3.07E+01
AAVBba30	9.35E+01	7.24E+01	8.50E+01	4.56E+01	4.75E+01	1.70E+01

[00217] In order to determine the tissue specific infectivity of the AAV capsids disclosed herein, AAV comprising each of the capsids and expressing the luciferase transgene were generated (AAV-RSV-egfp-T2A-Fluc2). Male Balb/C mice were purchased from Charles River Breeding Laboratories. A dose of 2×10^{13} vg/kg of AAV-RSV-egfp-T2A-Fluc2 vector was injected into the tail vein of 8 week old mice. At 3 and 5 weeks post injection, *in vivo* bioluminescent imaging was performed using an *in vivo* imaging device (IVIS Lumina LT obtained from PerkinElmer Inc., Waltham, MA). In brief, the mice were anesthetized with 2% isoflurane and oxygen. 150 μ l of 30 mg/ml of RediJect D-Luciferin Bioluminescent Substrate was injected intraperitoneally. Ten minutes after substrate injection, the animals were imaged with the *in vivo* imaging device using its cooled charge-coupled device (CCD) camera. Images were taken in the dorsal positions of the animals. Anesthesia was maintained throughout the entire imaging session by isoflurane-oxygen delivery in the light-tight imaging chamber.

[00218] The mice were sacrificed after the imaging sessions at 5 weeks post AAV injection. Various organs were harvested and imaged using the imaging device. The measurement conditions were the same as those used for *in vivo* imaging.

[00219] For imaging, a gray-scale photograph of the animals was acquired, followed by bioluminescence image acquisition. Image data was processed and analyzed using living image software version 4.5.2 (PerkinsElmer Waltham, MA). Regions of interest (ROIs) were traced surrounding each animal as well as individual organs to quantify the total flux (TF) (photons/second) being released by luciferase activity. Total flux activity is a proxy for AAV infectivity of each organ system and is shown for each AAV capsid in Table 9. To further characterize the tissue specific infectivity imparted by various capsids, tissue from infected mice was harvested and sectioned. The percent of cells expressing GFP was quantitated for each (see Table 10). These data show that AAV harboring specific capsids and chimeric capsids demonstrate different tissue specificities, for example some AAV are more active in liver while others are more active in muscle tissue. In particular, the data demonstrate that capsids AAVanc110_9VR and Bba41 produce recombinant AAV that have a high degree of specificity for muscle cells. Accordingly, recombinant AAV comprising capsid AAVanc110_9VR or capsid Bba41 would be useful for targeted transgene delivery to muscle cells.

Table 9: Total Flux in Tissue (photons/sec/cm²/radian): presented as AVE/(STD)

Capsid	Brain	Lung	Heart	Liver	Spleen	Kidney	Gast.*	Quad.**	Front Limb
AAV5	2.14E+04 (5.75E+03)	9.41E+05 (4.50E+05)	9.57E+03 (2.80E+03)	2.75E+06 (6.58E+05)	2.63E+03 (1.94E+03)	1.11E+04 (1.75E+03)	2.38E+04 (1.44E+04)	1.12E+04 (7.74E+03)	1.45E+03 (6.40E+02)
AAV8	2.31E+06 (7.88E+05)	4.83E+06 (2.34E+06)	6.57E+06 (1.71E+06)	4.58E+08 (1.82E+08)	7.34E+05 (2.84E+05)	2.63E+06 (5.58E+05)	6.87E+06 (4.71E+06)	9.25E+06 (4.83E+06)	1.91E+05 (1.55E+04)
AAV9	2.35E+06 (4.35E+05)	2.21E+06 (7.75E+05)	1.71E+07 (4.45E+06)	1.75E+08 (4.65E+07)	5.57E+05 (2.64E+05)	1.54E+06 (8.70E+05)	4.59E+07 (4.75E+07)	4.63E+07 (2.21E+07)	2.58E+06 (2.10E+06)
AAV8_9VRGBS	8.18E+05 (2.02E+05)	2.02E+06 (1.58E+06)	2.75E+06 (6.56E+05)	1.80E+08 (3.48E+07)	1.79E+05 (1.56E+04)	8.84E+05 (1.70E+05)	1.49E+06 (5.65E+05)	1.46E+06 (1.07E+05)	1.00E+05 (5.20E+04)
Anc110_9VR	1.87E+06 (1.68E+06)	7.79E+05 (2.73E+05)	1.28E+07 (4.54E+06)	1.80E+07 (4.25E+06)	5.83E+05 (7.37E+05)	9.34E+05 (5.53E+05)	1.72E+08 (1.84E+08)	8.88E+07 (6.55E+07)	2.69E+06 (1.74E+06)
Anc110_9GH	6.22E+03 (2.26E+03)	1.07E+04 (3.30E+03)	5.56E+03 (2.25E+03)	3.62E+05 (1.87E+05)	5.75E+03 (2.01E+03)	1.07E+04 (2.49E+03)	1.57E+04 (1.86E+04)	8.06E+03 (2.92E+03)	3.96E+03 (1.25E+03)
Anc110_9VRGBS	9.71E+04 (2.92E+04)	2.64E+05 (8.05E+04)	2.75E+06 (1.55E+06)	1.30E+07 (3.93E+06)	2.91E+04 (2.52E+03)	4.50E+05 (1.12E+05)	1.95E+07 (1.77E+07)	2.17E+07 (1.23E+07)	5.64E+05 (3.67E+05)
AAV6	5.72E+03 (3.77E+03)	6.60E+04 (4.18E+04)	7.61E+04 (3.87E+04)	9.23E+06 (5.37E+06)	4.98E+03 (4.23E+03)	1.87E+04 (1.14E+04)	1.68E+04 (2.82E+04)	3.19E+04 (4.26E+04)	1.72E+03 (2.34E+03)
AAVanc110	1.12E+05 (3.58E+04)	9.13E+05 (3.58E+05)	5.18E+05 (3.01E+05)	1.50E+08 (5.98E+07)	1.14E+05 (2.32E+04)	5.25E+05 (1.86E+05)	2.31E+05 (1.02E+05)	3.34E+05 (1.80E+05)	1.65E+04 (1.16E+04)
AAV6.2_9GH	3.20E+05 (1.59E+05)	9.94E+05 (3.22E+05)	1.41E+07 (4.99E+06)	5.63E+07 (1.66E+07)	9.32E+04 (3.59E+04)	1.21E+06 (3.78E+05)	9.56E+07 (7.74E+07)	6.96E+07 (2.16E+07)	1.46E+06 (6.74E+05)
AAV8_9VRGH	9.10E+04 (3.29E+04)	1.32E+06 (5.46E+05)	7.74E+05 (6.25E+05)	2.02E+08 (6.00E+07)	1.36E+05 (5.76E+04)	5.88E+05 (9.00E+04)	5.33E+05 (4.37E+05)	1.02E+06 (8.28E+05)	4.86E+04 (3.59E+04)
AAV7	1.47E+05 (1.16E+05)	1.03E+06 (7.65E+05)	8.04E+05 (5.40E+05)	1.47E+08 (6.17E+07)	1.47E+05 (3.72E+04)	1.13E+06 (1.10E+06)	7.04E+05 (4.48E+05)	6.52E+05 (4.34E+05)	4.42E+04 (2.04E+04)
AAVBce36	1.58E+05 (1.31E+04)	3.55E+05 (5.40E+04)	1.35E+06 (6.18E+05)	1.53E+08 (5.03E+06)	4.15E+05 (1.13E+05)	3.77E+05 (8.99E+04)	4.06E+05 (2.02E+05)	7.02E+05 (6.58E+05)	1.76E+04 (1.91E+04)
AAVBce41	1.07E+05 (4.06E+04)	3.92E+05 (4.13E+04)	7.54E+05 (6.44E+04)	1.49E+08 (5.00E+06)	5.79E+05 (2.54E+05)	5.35E+05 (4.75E+05)	3.32E+06 (4.62E+06)	1.64E+06 (5.94E+05)	5.18E+04 (3.47E+04)
AAVBba41	2.43E+05 (1.67E+05)	3.75E+05 (1.73E+05)	9.63E+06 (1.65E+06)	1.36E+07 (5.13E+06)	6.70E+04 (3.89E+04)	9.90E+05 (3.93E+05)	1.79E+08 (8.15E+07)	2.32E+08 (2.25E+07)	3.62E+06 (1.34E+06)
AAVBba42	2.17E+05 (2.10E+04)	4.35E+05 (4.16E+05)	9.54E+06 (8.14E+05)	1.65E+07 (7.75E+06)	5.23E+04 (1.76E+04)	1.12E+06 (4.41E+05)	2.68E+08 (2.35E+08)	1.62E+08 (2.70E+07)	4.94E+06 (1.99E+06)

AAVBba43	1.50E+05 (7.00E+04)	4.11E+05 (1.98E+05)	7.41E+06 (3.12E+06)	2.00E+07 (3.64E+06)	5.17E+04 (3.20E+04)	7.45E+05 (1.86E+05)	3.90E+07 (1.04E+07)	6.60E+07 (3.12E+07)	8.51E+05 (4.08E+05)
AAVBba30	1.10E+05 (6.96E+04)	4.74E+05 (2.55E+05)	3.10E+06 (1.43E+06)	1.24E+07 (3.66E+06)	1.99E+04 (1.00E+04)	5.18E+05 (9.89E+04)	2.00E+07 (8.52E+06)	2.38E+07 (7.17E+06)	5.77E+05 (2.65E+05)
AAV12	5.08E+03 (3.05E+03)	1.72E+05 (8.26E+04)	1.74E+04 (1.86E+03)	4.17E+06 (1.29E+06)	1.44E+03 (9.26E+02)	2.22E+04 (1.56E+04)	2.48E+05 (1.45E+05)	6.63E+05 (1.90E+05)	2.02E+04 (6.39E+04)
AAVBba33	1.86E+04 (1.11E+04)	2.20E+06 (1.20E+06)	2.37E+04 (1.35E+04)	1.18E+07 (5.28E+06)	2.65E+04 (1.36E+04)	3.56E+04 (1.80E+04)	2.79E+04 (3.18E+04)	1.22E+04 (6.27E+03)	7.75E+03 (8.64E+03)
AAVBba26	8.00E+02 (2.33E+03)	7.41E+03 (6.31E+03)	2.17E+04 (1.18E+04)	2.51E+04 (7.43E+03)	8.10E+02 (1.82E+03)	2.28E+03 (2.27E+03)	7.38E+04 (4.11E+04)	8.65E+04 (6.90E+04)	9.35E+03 (4.67E+03)
AAVBba44	1.41E+04 (1.05E+04)	8.76E+04 (4.40E+04)	5.29E+05 (2.87E+05)	8.05E+05 (4.43E+05)	6.90E+03 (8.53E+03)	1.05E+05 (1.11E+05)	2.17E+06 (1.47E+06)	1.94E+06 (1.63E+06)	6.40E+04 (3.60E+04)
AAVrh10	4.47E+04 (1.29E+04)	1.29E+05 (4.52E+04)	1.90E+05 (1.05E+05)	6.84E+07 (1.84E+07)	4.88E+04 (1.34E+04)	2.03E+05 (1.13E+05)	6.30E+05 (5.22E+05)	4.59E+05 (2.48E+05)	4.38E+04 (6.63E+04)
AAVBfm25	2.84E+04 (1.71E+04)	2.15E+05 (1.60E+05)	1.65E+05 (1.14E+05)	4.72E+07 (3.31E+07)	4.37E+04 (2.95E+04)	2.46E+05 (1.99E+05)	1.29E+07 (2.52E+07)	1.77E+05 (8.78E+04)	2.63E+04 (4.27E+04)
AAVBfm34	4.08E+04 (3.95E+03)	1.29E+05 (5.91E+04)	1.12E+06 (8.04E+05)	4.58E+07 (1.40E+07)	5.93E+04 (2.20E+04)	2.90E+05 (1.26E+05)	4.86E+06 (4.35E+06)	4.56E+06 (3.15E+06)	5.36E+04 (2.67E+04)
AAVBce18	5.22E+04 (5.51E+04)	2.10E+05 (1.89E+05)	1.85E+05 (1.23E+05)	1.03E+08 (1.30E+08)	8.57E+04 (8.51E+04)	2.65E+05 (3.36E+05)	1.19E+05 (7.19E+04)	1.72E+05 (1.30E+05)	2.61E+03 (1.64E+03)
AAVBrh28	7.72E+04 (3.27E+04)	2.62E+05 (8.93E+04)	1.21E+06 (4.38E+05)	7.27E+07 (2.83E+07)	6.86E+04 (3.00E+04)	3.76E+05 (1.31E+05)	1.93E+06 (7.29E+05)	2.29E+06 (1.09E+06)	8.59E+04 (4.89E+04)
AAVBce20	4.35E+04 (1.80E+04)	2.18E+05 (8.33E+04)	6.48E+05 (2.59E+05)	4.48E+07 (1.14E+07)	5.82E+05 (2.58E+04)	4.85E+05 (1.74E+05)	3.59E+07 (7.32E+07)	2.77E+06 (1.06E+05)	1.92E+05 (6.47E+04)

*Gast. = Gastrocnemius

**Quad. = Quadriceps

Table 10

	Diaphragm	Quadriceps	Gast.	Heart	Liver
AAV5	0%	5%	2%	NA	8%
AAV8	15%	2%	22%	NA	18%
AAVanc110_9VR	49%	29%	48%	18%	0.6%
Bba41	70%	24%	34%	32%	0.2%

What is Claimed:

1. An adeno-associated virus (AAV) comprising a capsid protein, wherein the capsid protein comprises an amino acid sequence that is at least 95% identical to (i) any one of SEQ ID NOS:15-89, (ii) the VP2 region of any one of SEQ ID NOS:15-89, or (iii) the VP3 region of any one of SEQ ID NOS:15-89, and further comprising a transgene comprising a heterologous gene operably linked to regulatory sequences that control expression of the heterologous gene in a host cell.
2. The AAV of claim 1, wherein the capsid protein comprises the amino acid sequence of (i) any one of SEQ ID NOS:15-89, (ii) the VP2 region of any one of SEQ ID NOS:15-89, or (iii) the VP3 region of any one of SEQ ID NOS:15-89.
3. The AAV of claim 1 or 2 further comprising AAV inverted terminal repeats.
4. A composition comprising the AAV of any one of claims 1 to 3 and a physiologically compatible carrier.
5. A method of delivering a transgene to a cell comprising contacting the cell with the AAV of any one of claims 1-3.
6. The method of claim 5, wherein the cell is a muscle cell.
7. The method of claim 5 or 6, wherein the transgene encodes a microdystrophin.
8. A method of treating a subject suffering from a disorder or disease associated with abnormal activity of an endogenous protein comprising administering to the subject an effective amount of an AAV comprising a capsid protein, wherein the capsid protein comprises an amino acid sequence that is at least 95% identical to (i) any one of SEQ ID NOS:15-89, (ii) the VP2 region of any one of SEQ ID NOS:15-89, or (iii) the VP3 region of any one of SEQ ID NOS:15-89, and wherein the AAV comprises a transgene encoding a biologically active copy of the protein.
9. The method of claim 8, wherein the disorder or disease is Duchenne muscular dystrophy and wherein the transgene encodes a microdystrophin.

10. A vector comprising a nucleic acid sequence encoding an adeno-associated virus (AAV) capsid protein having an amino acid sequence that is at least 95% identical to (i) any one of SEQ ID NOS:15-89, (ii) the VP2 region of any one of SEQ ID NOS:15-89, or (iii) the VP3 region of any one of SEQ ID NOS:15-89.
11. The vector of claim 10, wherein the nucleic acid sequence is operably linked to a heterologous regulatory element that controls expression of the capsid protein in a host cell.
12. A cell comprising the vector of claim 10 or 11.
13. Use of an AAV according to any one of claims 1-3 in the preparation of a medicament for the treatment of a disorder or disease.
14. The use of claim 13, wherein the disorder or disease is associated with abnormal activity of an endogenous protein and further wherein the AAV comprises a transgene that encodes a biologically active copy of the protein.
15. The use of claim 14, wherein the disorder or disease is Duchenne muscular dystrophy and the transgene encodes a microdystrophin.
16. A composition comprising an AAV of any one of claims 1-3 for the treatment of a disease or disorder.
17. The composition of claim 16, wherein the disorder or disease is Duchenne muscular dystrophy and the transgene encodes a microdystrophin.
18. An adeno-associated virus (AAV) comprising a chimeric capsid protein having a VP1 amino acid sequence of a recipient backbone AAV capsid comprising variable regions I, II, III, IV, V, VI, VII, VIII and IX, except wherein one or more of variable regions I, II, III, IV, V, VI, VII, VIII, or IX is replaced by the corresponding variable region from one or more donor AAV capsids.
19. The AAV of claim 18, wherein only one variable region from the recipient AAV capsid is replaced by the corresponding variable region from the donor AAV capsid.

20. The AAV of claim 18, wherein two or more variable regions from the recipient AAV capsid are replaced by the corresponding variable regions from a single donor AAV capsid.
21. The AAV of claim 18, wherein two or more variable regions from the recipient AAV capsid are replaced by the corresponding variable regions from two or more donor AAV capsids.
22. The AAV of claim 18, wherein all nine variable regions of the recipient AAV capsid are replaced by the corresponding variable regions from a single donor AAV capsid.
23. The AAV of any one of claims 18-22, wherein the recipient AAV capsid further comprises a GBS region or a GH loop region and the GBS region or the GH loop region is replaced by the corresponding region from one or more donor AAV capsids.
24. The AAV of claim 23, wherein all nine variable regions and the GBS region of the recipient AAV capsid are replaced by the corresponding variable regions and GBS region from one or more donor AAV capsids.
25. The AAV of claim 23, wherein all nine variable regions and the GBS region of the recipient AAV capsid are replaced by the corresponding variable regions and GBS region from two or more donor AAV capsids.
26. The AAV of claim 23, wherein the GH loop of the recipient AAV capsid is replaced by the corresponding GH loop region from a donor AAV capsid.
27. The AAV of claim 23, wherein all nine variable regions and the GH loop region of the recipient AAV capsid are replaced by the corresponding variable regions and GH loop region from one or more donor AAV capsids.
28. The AAV of any one of claims 18-27, wherein the recipient AAV capsid sequence is any one of SEQ ID NOS:1-14 and the donor AAV capsid sequences are selected from the group of sequences consisting of SEQ ID NOS:1-14, and the recipient AAV capsid sequence and the donor AAV capsid sequences are different.
29. The AAV of any one of claims 18-27, wherein the recipient AAV capsid sequence is any one of SEQ ID NOS:1-89 and the donor AAV capsid sequences are selected from the group of

sequences consisting of SEQ ID NOS:1-89, and the recipient AAV capsid sequence and the donor AAV capsid sequences are different.

30. The AAV of any one of claims 18-27, wherein the chimeric capsid protein comprises the amino acid sequence of any one of SEQ ID NOS:90-157.

31. The AAV of any one of claims 18-27 further comprising AAV inverted terminal repeats and a transgene comprising a heterologous gene operably linked to regulatory sequences that control expression of the heterologous gene in a host cell.

32. A composition comprising the AAV according to any one of claims 18 to 31 and a physiologically compatible carrier.

33. A method of delivering a transgene to a cell, the method comprising the step of contacting the cell with the AAV of claim 31 or the composition of claim 32.

34. The method of claim 33, wherein the cell is a muscle cell and the transgene encodes microdystrophin.

35. A method of treating a subject suffering from a disorder or disease associated with abnormal activity of an endogenous protein comprising administering to the subject an effective amount of the AAV of claim 31, wherein the heterologous gene encodes a biologically active copy of the protein.

36. The method of claim 35, wherein the disorder or disease is duchenne muscular dystrophy and the transgene encodes microdystrophin.

37. A vector comprising a nucleic acid sequence encoding a chimeric adeno-associated virus (AAV) capsid protein, wherein the chimeric AAV capsid protein comprises a VP1 amino acid sequence of a recipient AAV capsid sequence having variable regions I, II, III, IV, V, VI, VII, VIII and IX, and wherein one or more of variable regions I, II, III, IV, V, VI, VII, VIII, or IX is replaced by the corresponding variable region from a donor AAV capsid sequence, wherein the amino acid sequences of the chimeric AAV capsid protein and the recipient capsid sequence are different.

38. The vector of claim 37, wherein the nucleic acid sequence is operably linked to a heterologous regulatory element that controls expression of the chimeric AAV capsid protein in a host cell.
39. A cDNA comprising a nucleic acid sequence encoding a a chimeric capsid protein having a VP1 amino acid sequence of a recipient backbone AAV capsid comprising variable regions I, II, III, IV, V, VI, VII, VIII and IX, and wherein one or more of variable regions I, II, III, IV, V, VI, VII, VIII, or IX is replaced by the corresponding variable region from one or more donor AAV capsids.
40. A cell comprising the vector of claim 37 or 38 or the cDNA of claim 39.
41. Use of an AAV according to any of claims 18-31 in the preparation of a medicament for the treatment of a disorder or disease.
42. The use of claim 41, wherein the disorder or disease is associated with abnormal activity of a gene expressed by the AAV.
43. The use of claim 42, wherein the disorder or disease is Duchenne muscular dystrophy and the transgene encodes microdystrophin.
44. A composition comprising an AAV of any one of claims 18-31 for the treatment of a disease or disorder.
45. The composition of claim 44, wherein the disease or disorder is Duchenne muscular dystrophy and the transgene encodes microdystrophin.
46. A method of targeting a transgene to a muscle cell, comprising contacting the muscle cell with an AAV of any one of claims 18-31.
47. A method of producing an adeno-associated virus (AAV) of any one of claims 1-3 or 18-31 comprising:
- (a) culturing a viral production cell into which has been introduced a first nucleic acid vector comprising 5' and 3' AAV inverted terminal repeat sequences flanking a transgene comprising a

heterologous gene operably linked to regulatory sequences that control expression of the heterologous gene in a host cell, and a second nucleic acid vector comprising AAV rep and cap nucleic acids sequences, wherein said cap nucleic acid sequence encodes an AAV capsid that is at least 95% identical to any of SEQ ID NOS:15-157; and

(b) recovering the AAV from the supernatant of the viral production cell culture.

48. The method of claim 47, wherein the viral production cell is an insect cell.

49. The method of claim 48, wherein the insect cell is an Sf9 cell.

50. The method of claim 47, wherein the first nucleic acid vector is introduced into the viral production cell by infection of the viral production cell by a baculovirus comprising the first nucleic acid vector.

51. The method of claim 47, wherein the first and second nucleic acid vectors are introduced into the viral production cell by infection of the viral production cell by a first baculovirus comprising the first nucleic acid vector and a second baculovirus comprising the second nucleic acid vector.

52. An adeno-associated virus (AAV) produced by the method of any one of claims 47-51.