MANUFACTURING OF TAILORED OILS IN RECOMBINANT HETEROTROPHIC MICROORGANISMS

CROSS-REFERENCE TO RELATED APPLICATIONS

[0001] This application claims the benefit under 35 U.S.C. 119(e) of US Provisional Patent Application No. 61/118,590, filed November 28, 2008, US Provisional Patent Application No. 61/118,994, filed December 1, 2008, US Provisional Patent Application No. 61/174,357, filed April 30, 2009, and US Provisional Patent Application No. 61/219,525, filed June 23, 2009. Each of these applications is incorporated herein by reference in its entirety for all purposes.

REFERENCE TO A SEQUENCE LISTING

[0002] This application includes a sequence listing as shown in pages 1-72, appended hereto.

FIELD OF THE INVENTION

[0003] The present inveniton relates to the production of oils, fuels, and oleochemicals made from microorganisms. In particular, the disclosure relates to oil-bearing microalgae, methods of cultivating them for the production of useful compounds, including lipids, fatty acid esters, fatty acids, aldehydes, alcohols, and alkanes, and methods and reagents for genetically altering them to improve production efficiency and alter the type and composition of the oils produced by them.

BACKGROUND OF THE INVENTION

[0004] Fossil fuel is a general term for buried combustible geologic deposits of organic materials, formed from decayed plants and animals that have been converted to crude oil, coal, natural gas, or heavy oils by exposure to heat and pressure in the earth's crust over hundreds of millions of years. Fossil fuels are a finite, non-renewable resource.

[0005] Increased demand for energy by the global economy has also placed increasing pressure on the cost of hydrocarbons. Aside from energy, many industries, including plastics and chemical manufacturers, rely heavily on the availability of hydrocarbons as a feedstock for their manufacturing processes. Cost-effective alternatives to current sources of supply could help mitigate the upward pressure on energy and these raw material costs.

[0006] PCT Pub. No. 2008/151149 describes methods and materials for cultivating microalgae for the production of oil and particularly exemplifies the production of diesel fuel from oil produced by the microalgae *Chlorella protothecoides*. There remains a need for improved methods for producing oil in microalgae, particularly for methods that produce oils

with shorter chain length and a higher degree of saturation and without pigments, with greater yield and efficiency. The present invention meets this need.

SUMMARY OF THE INVENTION

[0007] The invention provides cells of the genus *Prototheca* comprising an exogenous gene, and in some embodiments the cell is a strain of the species *Prototheca moriformis, Prototheca krugani, Prototheca stagnora* or *Prototheca zopfii* and in other embodiment the cell has a 23S rRNA sequence with at least 70, 75, 80, 85 or 95% nucleotide identity to one or more of SEQ ID NOs: 11-19. In some cells the exogenous gene is coding sequence and is in operable linkage with a promoter, and in some embodiments the promoter is from a gene endogenous to a species of the genus *Prototheca*. In further embodiments the coding sequence encodes a protein selected from the group consisting of a sucrose invertase, a fatty acyl-ACP thioesterase, a fatty acyl-CoA/aldehyde reductase, a fatty acyl-CoA reductase, a fatty aldehyde reductase, a fatty aldehyde decarbonylase, an acyl carrier protein and a protein that imparts resistance to an antibiotic. Some embodiments of a fatty acyl-ACP thioesterase that has hydrolysis activity towards one or more fatty acyl-ACP substrates of chain length C8, C10, C12 or C14, including acyl-ACP thioesterases with at least 50, 60, 70, 80, or 90% amino acid identity with one or more sequences selected from the group consisting of SEQ ID NOs: 59, 61, 63 and 138-140. In further embodiments the coding sequence comprises a plastid targeting sequence from microalgae, and in some embodiments the microalgae is a species of the genus *Prototheca* or *Chlorella* as well as other genera from the family *Chlorellaceae*. In some embodiments the plastid targeting sequence has at least 20, 25, 35, 45, or 55% amino acid sequence identity to one or more of SEQ ID NOs: 127-133 and is capable of targeting a protein encoded by an exogenous gene not located in the plastid genome to the plastid. In other embodiments the promoter is upregulated in response to reduction or elimination of nitrogen in the culture media of the cell, such as at least a 3-fold upregulation as determined by transcript abundance in a cell of the genus *Prototheca* when the extracellular environment changes from containing at least 10mM or 5mM nitrogen to containing no nitrogen. In further embodiments the promoter comprises a segment of 50 or more nucleotides of one of SEQ ID NOs: 91-102. In other embodiments the cell has a 23S rRNA sequence with at least 70, 75, 80, 85 or 95% nucleotide identity to one or more of SEQ ID NOs: 11-19. In other embodiments the exogenous gene is integrated into a chromosome of the cell.

[0008] In additional embodiments of cells of the invention, the cell is of the genus *Prototheca* and comprises an exogenous fatty acyl-ACP thioesterase gene and a lipid profile

of at least 4% C8-C14 of total lipids of the cell, an amount of C8 that is at least 0.3% of total lipids of the cell, an amount of C10 that is at least 2% of total lipids of the cell, an amount of C12 that is at least 2% of total lipids of the cell, an amount of C14 that is at least 4% of total lipids of the cell, and an amount of C8-C14 that is 10-30%, 20-30%, or at least 10, 20, or 30% of total lipids of the cell. In some embodiments the cell further comprises an exogenous sucrose invertase gene. In some embodiments the cell is a strain of the species *Prototheca moriformis, Prototheca krugani, Prototheca stagnora* or *Prototheca zopfii*, and in other embodiment the cell has a 23S rRNA sequence with at least 70, 75, 80, 85 or 95% nucleotide identity to one or more of SEQ ID NOs: 11-19. In other embodiments the exogenous fatty acyl-ACP thioesterase gene is integrated into a chromosome of the cell. Other embodiments of the invention comprise methods of making triglyceride compositions of a lipid profile of at least 4% C8-C14 w/w or area percent of the triglyceride composition, an amount of C8 that is at least 0.3% w/w or area percent, an amount of C10 that is at least 2% w/w or area percent, an amount of C12 that is at least 2% w/w or area percent, an amount of C14 that is at least 4% w/w or area percent, and an amount of C8-C14 that is 10-30%, 20-30%, or at least 10, 20, or 30% w/w or area percent. The invention also comprises methods of making triglyceride compositions comprising cultivating the foregoing cells, wherein the cells also comprise an exogenous gene encoding a sucrose invertase and sucrose is provided as a carbon source. In some embodiments the sucrose invertase has at least 50, 60, 70, 80, or 90% amino acid identity to one or more of SEQ ID NOs: 3, 20-29 and 90.

[0009] Embodiments of the invention include triglyceride oil compositions as well as cells containing triglyceride oil compositions comprising a lipid profile of at least 4% C8-C14 and one or more of the following attributes: 0.1-0.4 micrograms/ml total carotenoids, less than 0.4 micrograms/ml total carotenoids, less than 0.001 micrograms/ml lycopene; less than 0.02 micrograms/ml beta carotene, less than 0.02 milligrams of chlorophyll per kilogram of oil; 0.40-0.60 milligrams of gamma tocopherol per 100 grams of oil; 0.2-0.5 milligrams of total tocotrienols per gram of oil, less than 0.4 milligrams of total tocotrienols per gram of oil, 4-8 mg per 100 grams of oil of campesterol, and 40-60 mg per 100 grams of oil of stigmasterol. In some embodiments of the invention the triglyceride oil compositions have a lipid profile of at least 4% C8-C14 w/w or area percent of the triglyceride composition, an amount of C8 that is at least 0.3% w/w or area percent, an amount of C10 that is at least 2% w/w or area percent, an amount of C12 that is at least 2% w/w or area percent, an amount of C14 that is at least 4% w/w or area percent, and an amount of C8-C14 that is 10-30%, 20-30%, or at least 10, 20, or 30% w/w or area percent. In other embodiments the triglyceride oil composition is

blended with at least one other composition selected from the group consisting of soy, rapeseed, canola, palm, palm kernel, coconut, corn, waste vegetable, Chinese tallow, olive, sunflower, cotton seed. chicken fat, beef tallow, porcine tallow, microalgae, macroalgae, *Cuphea*, flax, peanut, choice white grease, lard, *Camelina* sativa, mustard seed cashew nut, oats, lupine, kenaf, calendula, hemp, coffee, linseed (flax), hazelnut, euphorbia, pumpkin seed, coriander, camellia, sesame, safflower, rice, tung tree, cocoa, copra, pium poppy, castor beans, pecan, jojoba, jatropha, macadamia, Brazil nuts, avocado, petroleum, or a distillate fraction of any of the preceding oils.

[0010] Methods of the invention also include processing the aforementioned oils of by performing one or more chemical reactions from the list consisting of transesterification, hydrogenation, hydrocracking, deoxygenation, isomerization, interesterification, hydroxylation, hydrolysis to yield free fatty acids, and saponification. The invention also includes hydrocarbon fuels made from hydrogenation and isomerization of the aforementioned oils and fatty acid alkyl esters made from transesterification of the aforementioned oils. In some embodiments the hydrocarbon fuel is made from triglyceride isolated from cells of the genus *Prototheca* wherein the ASTM D86 T10-T90 distillation range is at least 25°C. In other embodiments the fatty acid alkyl ester fuel is made from triglyceride isolated from cells of the genus *Prototheca*, wherein the composition has an ASTM D6751 A1 cold soak time of less than 120 seconds.

[0011] The invention also includes composition comprising (a) polysaccharide comprising one or more monosaccharides from the list consisting of 20-30 mole percent galactose; 55-65 mole percent glucose; and 5-15 mole percent mannose; (b) protein; and (c) DNA comprising a 23S rRNA sequence with at least 70, 75, 80, 85 or 95% nucleotide identity to one or more of SEQ ID NOs: 11-19; and (d) an exogenous gene. In some embodiments the exogenous gene is selected from a sucrose invertase and a fatty acyl-ACP thioesterase, and in further embodiments the composition further comprises lipid with a lipid profile of at least 4% C8-C14. In other embodiments the composition is formulated for consumption as an animal feed.

[0012] The invention includes recombinant nucleic acids encoding promoters that are upregulated in response to reduction or elimination of nitrogen in the culture media of a cell of the genus *Prototheca*, such as at least a 3-fold upregulation as determined by transcript abundance when the extracellular environment changes from containing at least 10mM or 5mM nitrogen to containing no nitrogen. In some embodiments the recombinant nucleic acid comprises a segment of 50 or more nucleotides of one of SEQ ID NOs: 91-102. The

invention also includes nucleic acid vectors comprising an expression cassette comprising (a) a promoter that is active in a cell of the genus *Prototheca*; and (b) a coding sequence in operable linkage with the promoter wherein the coding sequence contains the most or second most preferred codons of Table 1 for at least 20, 30, 40, 50, 60, or 80% of the codons of the coding sequence. In some vectors the coding sequence comprises a plastid targeting sequence in-frame with a fatty acyl-ACP thioesterase, including thioesterase that have hydrolysis activity towards one or more fatty acyl-ACP substrates of chain length C8, C10, C12 or C14. Some vectors include plastid targeting sequences that encode peptides that are capable of targeting a protein to the plastid of a cell of the genus *Prototheca*, including those from microalgae and those wherein the plastid targeting sequence has at least 20, 25, 35, 45, or 55% amino acid sequence identity to one or more of SEQ ID NOs. 127-133 and is capable of targeting a protein to the plastid of a cell of the genus *Prototheca*. Additional vectors of the invention comprise nucleic acid sequences endogenous to the nuclear genome of a cell of the genus *Prototheca*, wherein the sequence is at least 200 nucleotides long, and some vectors comprise first and second nucleic acid sequences endogenous to the nuclear genome of a cell of the genus *Prototheca*, wherein the first and second sequences (a) are each at least 200 nucleotides long; (b) flank the expression cassette; and (c) are located on the same *Prototheca* chromosome no more than 5, 10, 15, 20, and 50kB apart.

[0013] The invention also includes a recombinant nucleic acid with at least 80, 90, 95 or 98% nucleotide identity to one or both of SEQ ID NOs: 134-135 and a recombinant nucleic acid encoding a protein with at least 80, 90, 95 or 98% amino acid identity to one or both of SEQ ID NOs: 136-137.

[0014] The invention also comprises methods of producing triglyceride compositions, comprising (a) culturing a population of cells of the genus *Prototheca* in the presence of a fixed carbon source, wherein: (i) the cells contain an exogenous gene; (ii) the cells accumulate at least 10, 20, 30, 40, 60, or 70% of their dry cell weight as lipid; and (iii) the fixed carbon source is selected from the group consisting of sorghum and depolymerized cellulosic material; and (b) isolating lipid components from the cultured microorganisms. In some embodiments the the fixed carbon source is depolymerized cellulosic material selected from the group consisting of corn stover, *Miscanthus*, forage sorghum, sugar beet pulp and sugar cane bagasse, optionally that has been subjected to washing with water prior to the culturing step. In some methods the fixed carbon source is depolymerized cellulosic material and the glucose level of the depolymerized cellulosic material is concentrated to a level of at least 300 g/liter, at least 400 g/liter, at least 500 g/liter, or at least 600 g/liter of prior to the

culturing step and is fed to the culture over time as the cells grow and accumulate lipid. In some methods the exogenous gene encodes a fatty acyl-ACP thioesterase that has hydrolysis activity towards one or more fatty acyl-ACP substrates of chain length C8, C10, C12 or C14, and in some methods the triglyceride has a lipid profile of at least 4% C8-C14 and one or more of the following attributes: 0.1-0.4 micrograms/ml total carotenoids; less than 0.02 milligrams of chlorophyll per kilogram of oil; 0.40-0.60 milligrams of gamma tocopherol per 100 grams of oil; 0.2-0.5 milligrams of total tocotrienols per gram of oil, 4-8 mg per 100 grams of oil of campesterol, and 40-60 mg per 100 grams of oil of stigmasterol.

[0015] Further methods of the invention include producing a triglyceride composition, comprising: (a) culturing a population of microorganisms in the presence of depolymerized cellulosic material, wherein: (i) the depolymerized cellulosic material is subjected to washing with water prior to the culturing step; (ii) the cells accumulate at least 10, 20, 30, 40, 60, or 70% of their dry cell weight as lipid; and (iii) the depolymerized cellulosic material is concentrated to at least 300, 400, 500, or 600 g/liter of glucose prior to the cultivation step; (iv) the microorganisms are cultured in a fed-batch reaction in which depolymerized cellulosic material of at least 300, 400, 500, or 600 g/liter of glucose is fed to the microorganisms; and (b) isolating lipid components from the cultured microorganisms. In some embodiments the the fixed carbon source is depolymerized cellulosic material selected from the group consisting of corn stover, *Miscanthus*, forage sorghum, sugar beet pulp and sugar cane bagasse. In further embodiments the microorganisms are a species of the genus *Prototheca* and contain an exogenous gene, including a fatty acyl-ACP thioesterase that has hydrolysis activity towards one or more fatty acyl-ACP substrates of chain length C8, C10, C12 or C14. A further method of the invention comprises manufacturing triglyceride oil comprising cultivating a cell that has a 23S rRNA sequence with at least 90 or 96% nucleotide identity to SEQ ID NO: 30 in the presence of sucrose as a carbon source.

[0016] The invention also includes methods of manufacturing a chemical comprising performing one or more chemical reactions from the list consisting of transesterification, hydrogenation, hydrocracking, deoxygenation, isomerization, interesterification, hydroxylation, hydrolysis, and saponification on a triglyceride oil, wherein the oil has a lipid profile of at least 4% C8-C14 and one or more of the following attributes: 0.1-0.4 micrograms/ml total carotenoids; less than 0.02 milligrams of chlorophyll per kilogram of oil; 0.10-0.60 milligrams of gamma tocopherol per 100 grams of oil; 0.1-0.5 milligrams of total tocotrienols per gram of oil, 1-8 mg per 100 grams of oil of campesterol, and 10-60 mg per 100 grams of oil of stigmasterol. Some methods are performed by manufacturing the oil by

cultivating a cell of the genus *Prototheca* that comprises an exogenous fatty acyl-ACP thioesterase gene that encodes a fatty acyl-ACP thioesterase having hydrolysis activity towards one or more fatty acyl-ACP substrates of chain length C8, C10, C12 or C14. In some methods the hydrolysis reaction is selected from the group consisting of saponification, acid hydrolysis, alkaline hydrolysis, enzymatic hydrolysis, catalytic hydrolysis, and hot-compressed water hydrolysis, including a catalytic hydrolysis reaction wherein the oil is split into glycerol and fatty acids. In further methods the fatty acids undergo an amination reaction to produce fatty nitrogen compounds or an ozonolysis reaction to produce mono- and dibasic-acids. In some embodiments the oil undergoes a triglyceride splitting method selected from the group consisting of enzymatic splitting and pressure splitting. In some methods a condensation reaction follows the hydrolysis reaction. Other methods include performing a hydroprocessing reaction on the oil, optionally wherein the product of the hydroprocessing reaction undergoes a deoxygenation reaction or a condensation reaction prior to or simultaneous with the hydroprocessing reaction. Some methods additionally include a gas removal reaction. Additioanl methods include processing the aforementioned oils by performing a deoxygenation reaction selected from the group consisting of: a hydrogenolysis reaction, hydrogenation, a consecutive hydrogenation-hydrogenolysis reaction, a consecutive hydrogenolysis-hydrogenation reaction, and a combined hydrogenation-hydrogenolysis reaction. In some methods a condensation reaction follows the deoxygenation reaction. Other methods include performing an esterification reaction on the aforementioned oils, optionally an interestification reaction or a transesterification reaction. Other methods include performing a hydroxylation reaction on the aforementioned oils, optionally wherein a condensation reaction follows the hydroxylation reaction.

BRIEF DESCRIPTION OF THE DRAWINGS

[0017] Figures 1 and 2 illustrate the growth curves of *Prototheca* species and *Chlorella luteoviridis* strain SAG 2214 grown on sorghum as the carbon source.

[0018] Figure 3 shows time course growth of SAG 2214 on glucose and sucrose.

[0019] Figure 4 shows maps of the cassettes using in *Prototheca* transformations, as described in Example 3.

[0020] Figure 5 shows the results of Southern blot analysis on three transformants of UTEX strain 1435, as described in Example 3.

[0021] Figure 6 shows a schematic of the codon optimized and non-codon optimized *suc2* (yeast sucrose invertase (yInv)) transgene construct. The relevant restriction cloning sites are indicated and arrows indicate the direction of transcription.

[0022] Figure 7a shows the results of *Prototheca moriformis* grown on cellulosic-derived sugars (corn stover, beet pulp, sorghum cane, *Miscanthus* and glucose control). Growth is expressed in optical density measurements (A750 readings).

[0023] Figure 7b shows the results of growth experiments using *Prototheca moriformis* using different levels of corn stover-derived cellulosic sugar as compared to glucose/xylose control.

[0024] Figure 7c shows the impact that xylose has on the lipid production in *Prototheca* cultures.

[0025] Figure 7d shows the impact of salt concentration (Na_2SO_4) and antifoam on the growth (in dry cell weight (DCW)) of *Prototheca*.

[0026] Figure 8 shows the impact of hydrothermal treatment of various cellulosic materials (sugar cane bagasse, sorghum cane, *Miscanthus* and beet pulp) and the resulting sugar stream on the growth of *Prototheca*.

[0027] Figure 9 shows decreasing levels of hydroxymethyl furfurals (HMF) and furfurals in cellulosic biomass (sugar cane bagasse, sorghum cane, *Miscanthus* and beet pulp) after repeated cycles of hydrothermal treatment.

[0028] Figure 10 shows a schematic of a saccharification process of cellulosic materials to generate sugar streams suitable for use in heterotrophic oil production in a fermentor.

[0029] Figure 11 shows decreasing levels of HMF and furfurals in exploded sugar cane bagasse after repeated cycles of hydrothermal treatment.

[0030] Figure 12 shows a schematic of thioesterase constructs used in *Prototheca* transformations. The heterologous beta-tubulin (driving Neo^R) and glutamate dehydrogenase promoters are derived from *Chlamydomonas reinhardtii* and *Chlorella sorokiniana*, respectively. The nitrate reductase 3'UTR was derived from *Chlorella vulgaris*. The relevant restriction cloning sites are indicated and arrows indicate the direction of transcription.

[0031] Figure 13 shows a chromatogram of renewable diesel produced from *Prototheca* triglyceride oil.

DETAILED DESCRIPTION OF THE INVENTION

[0032] The present invention arises from the discovery that *Prototheca* and certain related microorganisms have unexpectedly advantageous properties for the production of oils, fuels, and other hydrocarbon or lipid compositions economically and in large quantities, as well as from the discovery of methods and reagents for genetically altering these microorganisms to improve these properties. The oils produced by these microorganisms can be used in the transportation fuel, petrochemical, and/or food and cosmetic industries, among other

applications. Transesterification of lipids yields long-chain fatty acid esters useful as biodiesel. Other enzymatic and chemical processes can be tailored to yield fatty acids, aldehydes, alcohols, alkanes, and alkenes. In some applications, renewable diesel, jet fuel, or other hydrocarbon compounds are produced. The present invention also provides methods of cultivating microalgae for increased productivity and increased lipid yield, and/or for more cost-effective production of the compositions described herein.

[0033] This detailed description of the invention is divided into sections for the convenience of the reader. Section I provides definitions of terms used herein. Section 2 provides a description of culture conditions useful in the methods of the invention. Section 3 provides a description of genetic engineering methods and materials. Section 4 provides a description of genetic engineering of *Prototheca* to enable sucrose utilization. Section 5 provides a description of genetic engineering of *Prototheca* to modify lipid biosynthesis. Section 6 describes methods for making fuels and chemicals. Section 7 discloses examples and embodiments of the invention. The detailed description of the invention is followed by examples that illustrate the various aspects and embodiments of the invention.

I. DEFINITIONS

[0034] Unless defined otherwise, all technical and scientific terms used herein have the meaning commonly understood by a person skilled in the art to which this invention belongs. The following references provide one of skill with a general definition of many of the terms used in this invention: Singleton *et al.*, *Dictionary of Microbiology and Molecular Biology* (2nd ed. 1994); *The Cambridge Dictionary of Science and Technology* (Walker ed., 1988); *The Glossary of Genetics*, 5th Ed., R. Rieger *et al.* (eds.), Springer Verlag (1991); and Hale & Marham, *The Harper Collins Dictionary of Biology* (1991). As used herein, the following terms have the meanings ascribed to them unless specified otherwise.

[0035] "Active in microalgae" refers to a nucleic acid that is functional in microalgae. For example, a promoter that has been used to drive an antibiotic resistance gene to impart antibiotic resistance to a transgenic microalgae is active in microalgae.

[0036] "Acyl carrier protein" or "ACP" is a protein that binds a growing acyl chain during fatty acid synthesis as a thiol ester at the distal thiol of the 4'-phosphopantetheine moiety and comprises a component of the fatty acid synthase complex.

[0037] "Acyl-CoA molecule" or "acyl-CoA" is a molecule comprising an acyl moiety covalently attached to coenzyme A through a thiol ester linkage at the distal thiol of the 4'-phosphopantetheine moiety of coenzyme A.

[0038] "Area Percent" refers to the area of peaks observed using FAME GC/FID detection methods in which every fatty acid in the sample is converted into a fatty acid methyl ester (FAME) prior to detection. For example, a separate peak is observed for a fatty acid of 14 carbon atoms with no unsaturation (C14:0) compared to any other fatty acid such as C14:1. The peak area for each class of FAME is directly proportional to its percent composition in the mixture and is calculated based on the sum of all peaks present in the sample (i.e. [area under specific peak/ total area of all measured peaks] X 100). When referring to lipid profiles of oils and cells of the invention, "at least 4% C8-C14" means that at least 4% of the total fatty acids in the cell or in the extracted glycerolipid composition have a chain length that includes 8, 10, 12 or 14 carbon atoms.

[0039] "Axenic" is a culture of an organism free from contamination by other living organisms.

[0040] "Biodiesel" is a biologically produced fatty acid alkyl ester suitable for use as a fuel in a diesel engine.

[0041] "Biomass" is material produced by growth and/or propagation of cells. Biomass may contain cells and/or intracellular contents as well as extracellular material, includes, but is not limited to, compounds secreted by a cell.

[0042] "Bioreactor" is an enclosure or partial enclosure in which cells are cultured, optionally in suspension.

[0043] "Catalyst" is an agent, such as a molecule or macromolecular complex, capable of facilitating or promoting a chemical reaction of a reactant to a product without becoming a part of the product. A catalyst increases the rate of a reaction, after which, the catalyst may act on another reactant to form the product. A catalyst generally lowers the overall activation energy required for the reaction such that it proceeds more quickly or at a lower temperature. Thus, a reaction equilibrium may be more quickly attained. Examples of catalysts include enzymes, which are biological catalysts; heat, which is a non-biological catalyst; and metals used in fossil oil refining processes.

[0044] "Cellulosic material" is the product of digestion of cellulose, including glucose and xylose, and optionally additional compounds such as disaccharides, oligosaccharides, lignin, furfurals and other compounds. Nonlimiting examples of sources of cellulosic material include sugar cane bagasses, sugar beet pulp, corn stover, wood chips, sawdust and switchgrass.

[0045] "Co-culture", and variants thereof such as "co-cultivate" and "co-ferment", refer to the presence of two or more types of cells in the same bioreactor. The two or more types of

cells may both be microorganisms, such as microalgae, or may be a microalgal cell cultured with a different cell type. The culture conditions may be those that foster growth and/or propagation of the two or more cell types or those that facilitate growth and/or proliferation of one, or a subset, of the two or more cells while maintaining cellular growth for the remainder.

[0046] "Cofactor" is any molecule, other than the substrate, required for an enzyme to carry out its enzymatic activity.

[0047] "Complementary DNA" or "cDNA" is a DNA copy of mRNA, usually obtained by reverse transcription of messenger RNA (mRNA) or amplification (e.g., via polymerase chain reaction ("PCR")).

[0048] "Cultivated", and variants thereof such as "cultured" and "fermented", refer to the intentional fostering of growth (increases in cell size, cellular contents, and/or cellular activity) and/or propagation (increases in cell numbers via mitosis) of one or more cells by use of selected and/or controlled conditions. The combination of both growth and propagation may be termed proliferation. Examples of selected and/or controlled conditions include the use of a defined medium (with known characteristics such as pH, ionic strength, and carbon source), specified temperature, oxygen tension, carbon dioxide levels, and growth in a bioreactor. Cultivate does not refer to the growth or propagation of microorganisms in nature or otherwise without human intervention; for example, natural growth of an organism that ultimately becomes fossilized to produce geological crude oil is not cultivation.

[0049] "Cytolysis" is the lysis of cells in a hypotonic environment. Cytolysis is caused by excessive osmosis, or movement of water, towards the inside of a cell (hyperhydration). The cell cannot withstand the osmotic pressure of the water inside, and so it explodes.

[0050] "Delipidated meal" and "delipidated microbial biomass" is microbial biomass after oil (including lipids) has been extracted or isolated from it, either through the use of mechanical (i.e., exerted by an expeller press) or solvent extraction or both. Delipidated meal has a reduced amount of oil/lipids as compared to before the extraction or isolation of oil/lipids from the microbial biomass but does contain some residual oil/lipid.

[0051] "Expression vector" or "expression construct" or "plasmid" or "recombinant DNA construct" refer to a nucleic acid that has been generated via human intervention, including by recombinant means or direct chemical synthesis, with a series of specified nucleic acid elements that permit transcription and/or translation of a particular nucleic acid in a host cell. The expression vector can be part of a plasmid, virus, or nucleic acid fragment. Typically, the expression vector includes a nucleic acid to be transcribed operably linked to a promoter.

[0052] "Exogenous gene" is a nucleic acid that codes for the expression of an RNA and/or protein that has been introduced ("transformed") into a cell. A transformed cell may be referred to as a recombinant cell, into which additional exogenous gene(s) may be introduced. The exogenous gene may be from a different species (and so heterologous), or from the same species (and so homologous), relative to the cell being transformed. Thus, an exogenous gene can include a homologous gene that occupies a different location in the genome of the cell or is under different control, relative to the endogenous copy of the gene. An exogenous gene may be present in more than one copy in the cell. An exogenous gene may be maintained in a cell as an insertion into the genome or as an episomal molecule.

[0053] "Exogenously provided" refers to a molecule provided to the culture media of a cell culture.

[0054] "Expeller pressing" is a mechanical method for extracting oil from raw materials such as soybeans and rapeseed. An expeller press is a screw type machine, which presses material through a caged barrel-like cavity. Raw materials enter one side of the press and spent cake exits the other side while oil seeps out between the bars in the cage and is collected. The machine uses friction and continuous pressure from the screw drives to move and compress the raw material. The oil seeps through small openings that do not allow solids to pass through. As the raw material is pressed, friction typically causes it to heat up.

[0055] "Fatty acyl-ACP thioesterase" is an enzyme that catalyzes the cleavage of a fatty acid from an acyl carrier protein (ACP) during lipid synthesis.

[0056] "Fatty acyl-CoA/aldehyde reductase" is an enzyme that catalyzes the reduction of an acyl-CoA molecule to a primary alcohol.

[0057] "Fatty acyl-CoA reductase" is an enzyme that catalyzes the reduction of an acyl-CoA molecule to an aldehyde.

[0058] "Fatty aldehyde decarbonylase" is an enzyme that catalyzes the conversion of a fatty aldehyde to an alkane.

[0059] "Fatty aldehyde reductase" is an enzyme that catalyzes the reduction of an aldehyde to a primary alcohol.

[0060] "Fixed carbon source" is a molecule(s) containing carbon, typically an organic molecule, that is present at ambient temperature and pressure in solid or liquid form in a culture media that can be utilized by a microorganism cultured therein.

[0061] "Homogenate" is biomass that has been physically disrupted.

[0062] "Hydrocarbon" is (a) a molecule containing only hydrogen and carbon atoms wherein the carbon atoms are covalently linked to form a linear, branched, cyclic, or partially

cyclic backbone to which the hydrogen atoms are attached. The molecular structure of hydrocarbon compounds varies from the simplest, in the form of methane (CH_4), which is a constituent of natural gas, to the very heavy and very complex, such as some molecules such as asphaltenes found in crude oil, petroleum, and bitumens. Hydrocarbons may be in gaseous, liquid, or solid form, or any combination of these forms, and may have one or more double or triple bonds between adjacent carbon atoms in the backbone. Accordingly, the term includes linear, branched, cyclic, or partially cyclic alkanes, alkenes, lipids, and paraffin. Examples include propane, butane, pentane, hexane, octane, and squalene.

[0063] "Hydrogen:carbon ratio" is the ratio of hydrogen atoms to carbon atoms in a molecule on an atom-to-atom basis. The ratio may be used to refer to the number of carbon and hydrogen atoms in a hydrocarbon molecule. For example, the hydrocarbon with the highest ratio is methane CH_4 (4:1).

[0064] "Hydrophobic fraction" is the portion, or fraction, of a material that is more soluble in a hydrophobic phase in comparison to an aqueous phase. A hydrophobic fraction is substantially insoluble in water and usually non-polar.

[0065] "Increase lipid yield" refers to an increase in the productivity of a microbial culture by, for example, increasing dry weight of cells per liter of culture, increasing the percentage of cells that constitute lipid, or increasing the overall amount of lipid per liter of culture volume per unit time.

[0066] "Inducible promoter" is a promoter that mediates transcription of an operably linked gene in response to a particular stimulus.

[0067] "In operable linkage" is a functional linkage between two nucleic acid sequences, such a control sequence (typically a promoter) and the linked sequence (typically a sequence that encodes a protein, also called a coding sequence). A promoter is in operable linkage with an exogenous gene if it can mediate transcription of the gene.

[0068] "In situ" means "in place" or "in its original position".

[0069] "Limiting concentration of a nutrient" is a concentration of a compound in a culture that limits the propagation of a cultured organism. A "non-limiting concentration of a nutrient" is a concentration that supports maximal propagation during a given culture period. Thus, the number of cells produced during a given culture period is lower in the presence of a limiting concentration of a nutrient than when the nutrient is non-limiting. A nutrient is said to be "in excess" in a culture, when the nutrient is present at a concentration greater than that which supports maximal propagation.

[0070] "Lipase" is a water-soluble enzyme that catalyzes the hydrolysis of ester bonds in water-insoluble, lipid substrates. Lipases catalyze the hydrolysis of lipids into glycerols and fatty acids.

[0071] "Lipid modification enzyme" refers to an enayme that alters the covalent structure of a lipid. Examples of lipid modification enzymes include a lipase, a fatty acyl-ACP thioesterase, a fatty acyl-CoA/aldehyde reductase, a fatty acyl-CoA reductase, a fatty aldehyde reductase, and a fatty aldehyde decarbonylase.

[0072] "Lipid pathway enzyme" is any enzyme that plays a role in lipid metabolism, i.e., either lipid synthesis, modification, or degradation, and any proteins that chemically modify lipids, as well as carrier proteins.

[0073] "Lipids" are a class of molecules that are soluble in nonpolar solvents (such as ether and chloroform) and are relatively or completely insoluble in water. Lipid molecules have these properties, because they consist largely of long hydrocarbon tails which are hydrophobic in nature. Examples of lipids include fatty acids (saturated and unsaturated); glycerides or glycerolipids (such as monoglycerides, diglycerides, triglycerides or neutral fats, and phosphoglycerides or glycerophospholipids); nonglycerides (sphingolipids, sterol lipids including cholesterol and steroid hormones, prenol lipids including terpenoids, fatty alcohols, waxes, and polyketides); and complex lipid derivatives (sugar-linked lipids, or glycolipids, and protein-linked lipids). "Fats" are a subgroup of lipids called "triacylglycerides."

[0074] "Lysate" is a solution containing the contents of lysed cells.

[0075] "Lysis" is the breakage of the plasma membrane and optionally the cell wall of a biological organism sufficient to release at least some intracellular content, often by mechanical, viral or osmotic mechanisms that compromise its integrity.

[0076] "Lysing" is disrupting the cellular membrane and optionally the cell wall of a biological organism or cell sufficient to release at least some intracellular content.

[0077] "Microalgae" is a eukaryotic microbial organism that contains a chloroplast or plastid, and optionally that is capable of performing photosynthesis, or a prokaryotic microbial organism capable of performing photosynthesis. Microalgae include obligate photoautotrophs, which cannot metabolize a fixed carbon source as energy, as well as heterotrophs, which can live solely off of a fixed carbon source. Microalgae include unicellular organisms that separate from sister cells shortly after cell division, such as *Chlamydomonas*, as well as microbes such as, for example, *Volvox*, which is a simple multicellular photosynthetic microbe of two distinct cell types. Microalgae include cells such

as *Chlorella*, *Dunaliella*, and *Prototheca*. Microalgae also include other microbial photosynthetic organisms that exhibit cell-cell adhesion, such as *Agmenellum*, *Anabaena*, and *Pyrobotrys*. Microalgae also include obligate heterotrophic microorganisms that have lost the ability to perform photosynthesis, such as certain dinoflagellate algae species and species of the genus *Prototheca*.

[0078] "Microorganism" and "microbe" are microscopic unicellular organisms.

[0079] "Naturally co-expressed" with reference to two proteins or genes means that the proteins or their genes are co-expressed naturally in a tissue or organism from which they are derived, e.g., because the genes encoding the two proteins are under the control of a common regulatory sequence or because they are expressed in response to the same stimulus.

[0080] "Osmotic shock" is the rupture of cells in a solution following a sudden reduction in osmotic pressure. Osmotic shock is sometimes induced to release cellular components of such cells into a solution.

[0081] "Polysaccharide-degrading enzyme" is any enzyme capable of catalyzing the hydrolysis, or saccharification, of any polysaccharide. For example, cellulases catalyze the hydrolysis of cellulose.

[0082] "Polysaccharides" or "glycans" are carbohydrates made up of monosaccharides joined together by glycosidic linkages. Cellulose is a polysaccharide that makes up certain plant cell walls. Cellulose can be depolymerized by enzymes to yield monosaccharides such as xylose and glucose, as well as larger disaccharides and oligosaccharides.

[0083] "Promoter" is a nucleic acid control sequence that directs transcription of a nucleic acid. As used herein, a promoter includes necessary nucleic acid sequences near the start site of transcription, such as, in the case of a polymerase II type promoter, a TATA element. A promoter also optionally includes distal enhancer or repressor elements, which can be located as much as several thousand base pairs from the start site of transcription.

[0084] "Recombinant" is a cell, nucleic acid, protein or vector, that has been modified due to the introduction of an exogenous nucleic acid or the alteration of a native nucleic acid. Thus, e.g., recombinant cells express genes that are not found within the native (non-recombinant) form of the cell or express native genes differently than those genes are expressed by a non-recombinant cell. A "recombinant nucleic acid" is a nucleic acid originally formed in vitro, in general, by the manipulation of nucleic acid, e.g., using polymerases and endonucleases, or otherwise is in a form not normally found in nature. Recombinant nucleic acids may be produced, for example, to place two or more nucleic acids in operable linkage. Thus, an isolated nucleic acid or an expression vector formed in vitro by

ligating DNA molecules that are not normally joined in nature, are both considered recombinant for the purposes of this invention. Once a recombinant nucleic acid is made and introduced into a host cell or organism, it may replicate using the in vivo cellular machinery of the host cell; however, such nucleic acids, once produced recombinantly, although subsequently replicated intracellularly, are still considered recombinant for purposes of this invention. Similarly, a "recombinant protein" is a protein made using recombinant techniques, i.e., through the expression of a recombinant nucleic acid.

[0085] "Renewable diesel" is a mixture of alkanes (such as C10:0, C12:0, C14:0, C16:0 and C18:0) produced through hydrogenation and deoxygenation of lipids.

[0086] "Saccharification" is a process of converting biomass, usually cellulosic or lignocellulosic biomass, into monomeric sugars, such as glucose and xylose. "Saccharified" or "depolymerized" cellulosic material or biomass refers to cellulosic material or biomass that has been converted into monomeric sugars through saccharification.

[0087] "Sonication" is a process of disrupting biological materials, such as a cell, by use of sound wave energy.

[0088] "Species of furfural" is 2-furancarboxaldehyde or a derivative that retains the same basic structural characteristics.

[0089] "Stover" is the dried stalks and leaves of a crop remaining after a grain has been harvested.

[0090] "Sucrose utilization gene" is a gene that, when expressed, aids the ability of a cell to utilize sucrose as an energy source. Proteins encoded by a sucrose utilization gene are referred to herein as "sucrose utilization enzymes" and include sucrose transporters, sucrose invertases, and hexokinases such as glucokinases and fructokinases.

II. CULTIVATION

[0091] The present invention generally relates to cultivation of *Prototheca* strains, particularly recombinant *Prototheca* strains, for the production of lipid. For the convenience of the reader, this section is subdivided into subsections. Subsection 1 describes *Prototheca* species and strains and how to identify new *Prototheca* species and strains and related microalgae by genomic DNA comparison. Subsection 2 describes bioreactors useful for cultivation. Subsection 3 describes media for cultivation. Subsection 4 describes oil production in accordance with illustrative cultivation methods of the invention.

1. Prototheca species and strains

[0092] *Prototheca* is a remarkable microorganism for use in the production of lipid, because it can produce high levels of lipid, particularly lipid suitable for fuel production. The

lipid produced by *Prototheca* has hydrocarbon chains of shorter chain length and a higher degree of saturation than that produced by other microalgae. Moreover, *Prototheca* lipid is generally free of pigment (low to undetectable levels of chlorophyll and certain carotenoids) and in any event contains much less pigment than lipid from other microalgae. Moreover, recombinant *Prototheca* cells provided by the invention can be used to produce lipid in greater yield and efficiency, and with reduced cost, relative to the production of lipid from other microorganisms. Illustrative Prototheca strains for use in the methods of the invention include In addition, this microalgae grows heterotrophically and can be genetically engineered as *Prototheca wickerhamii, Prototheca stagnora (including UTEX 327), Prototheca portoricensis, Prototheca moriformis (including UTEX strains 1441, 1435),* and *Prototheca zopfii.* Species of the genus *Prototheca* are obligate heterotrophs.

[0093] Species of *Prototheca* for use in the invention can be identified by amplification of certain target regions of the genome. For example, identification of a specific *Prototheca* species or strain can be achieved through amplification and sequencing of nuclear and/or chloroplast DNA using primers and methodology using any region of the genome, for example using the methods described in Wu *et al.*, *Bot. Bull. Acad. Sin.* (2001) 42:115-121 Identification of *Chlorella spp.* isolates using ribosomal DNA sequences. Well established methods of phylogenetic analysis, such as amplification and sequencing of ribosomal internal transcribed spacer (ITS1 and ITS2 rDNA), 23S rRNA, 18S rRNA, and other conserved genomic regions can be used by those skilled in the art to identify species of not only *Prototheca*, but other hydrocarbon and lipid producing organisms with similar lipid profiles and production capability. For examples of methods of identification and classification of algae also see for example *Genetics*, 2005 Aug;170(4):1601-10 and *RNA*, 2005 Apr;11(4):361-4.

[0094] Thus, genomic DNA comparison can be used to identify suitable species of microalgae to be used in the present invention. Regions of conserved genomic DNA, such as but not limited to DNA encoding for 23S rRNA, can be amplified from microalgal species and compared to consensus sequences in order to screen for microalgal species that are taxonomically related to the preferred microalgae used in the present invention. Examples of such DNA sequence comparison for species within the *Prototheca* genus are shown below. Genomic DNA comparison can also be useful to identify microalgal species that have been misidentified in a strain collection. Often a strain collection will identify species of microalgae based on phenotypic and morphological characteristics. The use of these characteristics may lead to miscategorization of the species or the genus of a microalgae. The

use of genomic DNA comparison can be a better method of categorizing microalgae species based on their phylogenetic relationship.

[0095] Microalgae for use in the present invention typically have genomic DNA sequences encoding for 23S rRNA that have at least 99%, least 95%, at least 90%, or at least 85% nucleotide identity to at least one of the sequences listed in SEQ ID NOs: 11-19.

[0096] For sequence comparison to determine percent nucleotide or amino acid identity, typically one sequence acts as a reference sequence, to which test sequences are compared. When using a sequence comparison algorithm, test and reference sequences are input into a computer, subsequence coordinates are designated, if necessary, and sequence algorithm program parameters are designated. The sequence comparison algorithm then calculates the percent sequence identity for the test sequence(s) relative to the reference sequence, based on the designated program parameters.

[0097] Optimal alignment of sequences for comparison can be conducted, *e.g.*, by the local homology algorithm of Smith & Waterman, *Adv. Appl. Math.* 2:482 (1981), by the homology alignment algorithm of Needleman & Wunsch, *J. Mol. Biol.* 48:443 (1970), by the search for similarity method of Pearson & Lipman, *Proc. Nat'l. Acad. Sci. USA* 85:2444 (1988), by computerized implementations of these algorithms (GAP, BESTFIT, FASTA, and TFASTA in the Wisconsin Genetics Software Package, Genetics Computer Group, 575 Science Dr., Madison, WI), or by visual inspection (see generally Ausubel *et al., supra*).

[0098] Another example algorithm that is suitable for determining percent sequence identity and sequence similarity is the BLAST algorithm, which is described in Altschul *et al., J. Mol. Biol.* 215:403-410 (1990). Software for performing BLAST analyses is publicly available through the National Center for Biotechnology Information (at the web address www.ncbi.nlm.nih.gov). This algorithm involves first identifying high scoring sequence pairs (HSPs) by identifying short words of length W in the query sequence, which either match or satisfy some positive-valued threshold score T when aligned with a word of the same length in a database sequence. T is referred to as the neighborhood word score threshold (Altschul *et al., supra.*). These initial neighborhood word hits act as seeds for initiating searches to find longer HSPs containing them. The word hits are then extended in both directions along each sequence for as far as the cumulative alignment score can be increased. Cumulative scores are calculated using, for nucleotide sequences, the parameters M (reward score for a pair of matching residues; always > 0) and N (penalty score for mismatching residues; always < 0). For amino acid sequences, a scoring matrix is used to calculate the cumulative score. Extension of the word hits in each direction are halted when: the cumulative alignment score

falls off by the quantity X from its maximum achieved value; the cumulative score goes to zero or below due to the accumulation of one or more negative-scoring residue alignments; or the end of either sequence is reached. For identifying whether a nucleic acid or polypeptide is within the scope of the invention, the default parameters of the BLAST programs are suitable. The BLASTN program (for nucleotide sequences) uses as defaults a word length (W) of 11, an expectation (E) of 10, M=5, N=-4, and a comparison of both strands. For amino acid sequences, the BLASTP program uses as defaults a word length (W) of 3, an expectation (E) of 10, and the BLOSUM62 scoring matrix. The TBLATN program (using protein sequence for nucleotide sequence) uses as defaults a word length (W) of 3, an expectation (E) of 10, and a BLOSUM 62 scoring matrix. (see Henikoff & Henikoff, *Proc. Natl. Acad. Sci. USA* 89:10915 (1989)).

[0099] In addition to calculating percent sequence identity, the BLAST algorithm also performs a statistical analysis of the similarity between two sequences (see, *e.g.*, Karlin & Altschul, *Proc. Nat'l. Acad. Sci. USA* 90:5873-5787 (1993)). One measure of similarity provided by the BLAST algorithm is the smallest sum probability (P(N)), which provides an indication of the probability by which a match between two nucleotide or amino acid sequences would occur by chance. For example, a nucleic acid is considered similar to a reference sequence if the smallest sum probability in a comparison of the test nucleic acid to the reference nucleic acid is less than about 0.1, more preferably less than about 0.01, and most preferably less than about 0.001.

[0100] Other considerations affecting the selection of microorganisms for use in the invention include, in addition to production of suitable lipids or hydrocarbons for production of oils, fuels, and oleochemicals: (1) high lipid content as a percentage of cell weight; (2) ease of growth; (3) ease of genetic engineering; and (4) ease of biomass processing. In particular embodiments, the wild-type or genetically engineered microorganism yields cells that are at least 40%, at least 45%, at least 50%, at least 55%, at least 60%, at least 65%, or at least 70% or more lipid. Preferred organisms grow heterotrophically (on sugars in the absence of light).

2. Bioreactor

[0101] Microrganisms are cultured both for purposes of conducting genetic manipulations and for production of hydrocarbons (e.g., lipids, fatty acids, aldehydes, alcohols, and alkanes). The former type of culture is conducted on a small scale and initially, at least, under conditions in which the starting microorganism can grow. Culture for purposes of hydrocarbon production is usually conducted on a large scale (e.g., 10,000 L, 40,000 L,

100,000 L or larger bioreactors) in a bioreactor. *Prototheca* are typically cultured in the methods of the invention in liquid media within a bioreactor. Typically, the bioreactor does not allow light to enter.

[0102] The bioreactor or fermentor is used to culture microalgal cells through the various phases of their physiological cycle. Bioreactors offer many advantages for use in heterotrophic growth and propagation methods. To produce biomass for use in food, microalgae are preferably fermented in large quantities in liquid, such as in suspension cultures as an example. Bioreactors such as steel fermentors can accommodate very large culture volumes (40,000 liter and greater capacity bioreactors are used in various embodiments of the invention). Bioreactors also typically allow for the control of culture conditions such as temperature, pH, oxygen tension, and carbon dioxide levels. For example, bioreactors are typically configurable, for example, using ports attached to tubing, to allow gaseous components, like oxygen or nitrogen, to be bubbled through a liquid culture. Other culture parameters, such as the pH of the culture media, the identity and concentration of trace elements, and other media constituents can also be more readily manipulated using a bioreactor.

[0103] Bioreactors can be configured to flow culture media though the bioreactor throughout the time period during which the microalgae reproduce and increase in number. In some embodiments, for example, media can be infused into the bioreactor after inoculation but before the cells reach a desired density. In other instances, a bioreactor is filled with culture media at the beginning of a culture, and no more culture media is infused after the culture is inoculated. In other words, the microalgal biomass is cultured in an aqueous medium for a period of time during which the microalgae reproduce and increase in number; however, quantities of aqueous culture medium are not flowed through the bioreactor throughout the time period. Thus in some embodiments, aqueous culture medium is not flowed through the bioreactor after inoculation.

[0104] Bioreactors equipped with devices such as spinning blades and impellers, rocking mechanisms, stir bars, means for pressurized gas infusion can be used to subject microalgal cultures to mixing. Mixing may be continuous or intermittent. For example, in some embodiments, a turbulent flow regime of gas entry and media entry is not maintained for reproduction of microalgae until a desired increase in number of said microalgae has been achieved.

[0105] Bioreactor ports can be used to introduce, or extract, gases, solids, semisolids, and liquids, into the bioreactor chamber containing the microalgae. While many bioreactors have

more than one port (for example, one for media entry, and another for sampling), it is not necessary that only one substance enter or leave a port. For example, a port can be used to flow culture media into the bioreactor and later used for sampling, gas entry, gas exit, or other purposes. Preferably, a sampling port can be used repeatedly without altering compromising the axenic nature of the culture. A sampling port can be configured with a valve or other device that allows the flow of sample to be stopped and started or to provide a means of continuous sampling. Bioreactors typically have at least one port that allows inoculation of a culture, and such a port can also be used for other purposes such as media or gas entry.

[0106] Bioreactors ports allow the gas content of the culture of microalgae to be manipulated. To illustrate, part of the volume of a bioreactor can be gas rather than liquid, and the gas inlets of the bioreactor to allow pumping of gases into the bioreactor. Gases that can be beneficially pumped into a bioreactor include air, air/CO_2 mixtures, noble gases, such as argon, and other gases. Bioreactors are typically equipped to enable the user to control the rate of entry of a gas into the bioreactor. As noted above, increasing gas flow into a bioreactor can be used to increase mixing of the culture.

[0107] Increased gas flow affects the turbidity of the culture as well. Turbulence can be achieved by placing a gas entry port below the level of the aqueous culture media so that gas entering the bioreactor bubbles to the surface of the culture. One or more gas exit ports allow gas to escape, thereby preventing pressure buildup in the bioreactor. Preferably a gas exit port leads to a "one-way" valve that prevents contaminating microorganisms from entering the bioreactor.

3. Media

[0108] Microalgal culture media typically contains components such as a fixed nitrogen source, a fixed carbon source, trace elements, optionally a buffer for pH maintenance, and phosphate (typically provided as a phosphate salt). Other components can include salts such as sodium chloride, particularly for seawater microalgae. Nitrogen sources include organic and inorganic nitrogen sources, including, for example, without limitation, molecular nitrogen, nitrate, nitrate salts, ammonia (pure or in salt form, such as, $(NH_4)_2SO_4$ and NH_4OH), protein, soybean meal, cornsteep liquor, and yeast extract. Examples of trace elements include zinc, boron, cobalt, copper, manganese, and molybdenum in, for example, the respective forms of $ZnCl_2$, H_3BO_3, $CoCl_2 \cdot 6H_2O$, $CuCl_2 \cdot 2H_2O$, $MnCl_2 \cdot 4H_2O$ and $(NH_4)_6Mo_7O_{24} \cdot 4H_2O$.

[0109] Microorganisms useful in accordance with the methods of the present invention are found in various locations and environments throughout the world. As a consequence of their isolation from other species and their resulting evolutionary divergence, the particular growth medium for optimal growth and generation of lipid and/or hydrocarbon constituents can be difficult to predict. In some cases, certain strains of microorganisms may be unable to grow on a particular growth medium because of the presence of some inhibitory component or the absence of some essential nutritional requirement required by the particular strain of microorganism.

[0110] Solid and liquid growth media are generally available from a wide variety of sources, and instructions for the preparation of particular media that is suitable for a wide variety of strains of microorganisms can be found, for example, online at http://www.utex.org/, a site maintained by the University of Texas at Austin, 1 University Station A6700, Austin, Texas, 78712-0183, for its culture collection of algae (UTEX). For example, various fresh water and salt water media include those described in PCT Pub. No. 2008/151149, incorporated herein by reference.

[0111] In a particular example, Proteose Medium is suitable for axenic cultures, and a 1L volume of the medium (pH ~6.8) can be prepared by addition of 1g of proteose peptone to 1 liter of Bristol Medium. Bristol medium comprises 2.94 mM $NaNO_3$, 0.17 mM $CaCl_2 \cdot 2H_2O$, 0.3 mM $MgSO_4 \cdot 7H_2O$, 0.43 mM, 1.29 mM KH_2PO_4, and 1.43 mM NaCl in an aqueous solution. For 1.5% agar medium, 15 g of agar can be added to 1 L of the solution. The solution is covered and autoclaved, and then stored at a refrigerated temperature prior to use. Another example is the *Prototheca* isolation medium (PIM), which comprises 10g/L postassium hydrogen phthalate (KHP), 0.9g/L sodium hydroxide, 0.1g/L magnesium sulfate, 0.2g/L potassium hydrogen phosphate, 0.3g/L ammonium chloride, 10g/L glucose 0.001g/L thiamine hydrochloride, 20g/L agar, 0.25g/L 5-fluorocytosine, at a pH in the range of 5.0 to 5.2 (see Pore, 1973, App. Microbiology, 26: 648-649). Other suitable media for use with the methods of the invention can be readily identified by consulting the URL identified above, or by consulting other organizations that maintain cultures of microorganisms, such as SAG, CCAP, or CCALA. SAG refers to the Culture Collection of Algae at the University of Göttingen (Göttingen, Germany), CCAP refers to the culture collection of algae and protozoa managed by the Scottish Association for Marine Science (Scotland, United Kingdom), and CCALA refers to the culture collection of algal laboratory at the Institute of Botany (Třeboň, Czech Republic). Additionally, US Patent No. 5,900,370 describes media formulations and conditions suitable for heterotrophic fermentation of *Prototheca* species.

[0112] For oil production, selection of a fixed carbon source is important, as the cost of the fixed carbon source must be sufficiently low to make oil production economical. Thus, while suitable carbon sources include, for example, acetate, floridoside, fructose, galactose, glucuronic acid, glucose, glycerol, lactose, mannose, N-acetylglucosamine, rhamnose, sucrose, and/or xylose, selection of feedstocks containing those compounds is an important aspect of the methods of the invention. Suitable feedstocks useful in accordance with the methods of the invention include, for example, black liquor, corn starch, depolymerized cellulosic material, milk whey, molasses, potato, sorghum, sucrose, sugar beet, sugar cane, rice, and wheat. Carbon sources can also be provided as a mixture, such as a mixture of sucrose and depolymerized sugar beet pulp. The one or more carbon source(s) can be supplied at a concentration of at least about 50 μM, at least about 100 μM, at least about 500 μM, at least about 5 mM, at least about 50 mM, and at least about 500 mM, of one or more exogenously provided fixed carbon source(s). Carbon sources of particular interest for purposes of the present invention include cellulose (in a depolymerized form), glycerol, sucrose, and sorghum, each of which is discussed in more detal below.

[0113] In accordance with the present invention, microorganisms can be cultured using depolymerized cellulosic biomass as a feedstock. Cellulosic biomass (e.g., stover, such as corn stover) is inexpensive and readily available; however, attempts to use this material as a feedstock for yeast have failed. In particular, such feedstocks have been found to be inhibitory to yeast growth, and yeast cannot use the 5-carbon sugars produced from cellulosic materials (e.g., xylose from hemi-cellulose). By contrast, microalgae can grow on processed cellulosic material. Cellulosic materials generally include about 40-60% cellulose; about 20-40% hemicellulose; and 10-30% lignin.

[0114] Suitable cellulosic materials include residues from herbaceous and woody energy crops, as well as agricultural crops, i.e., the plant parts, primarily stalks and leaves, not removed from the fields with the primary food or fiber product. Examples include agricultural wastes such as sugarcane bagasse, rice hulls, corn fiber (including stalks, leaves, husks, and cobs), wheat straw, rice straw, sugar beet pulp, citrus pulp, citrus peels; forestry wastes such as hardwood and softwood thinnings, and hardwood and softwood residues from timber operations; wood wastes such as saw mill wastes (wood chips, sawdust) and pulp mill waste; urban wastes such as paper fractions of municipal solid waste, urban wood waste and urban green waste such as municipal grass clippings; and wood construction waste. Additional cellulosics include dedicated cellulosic crops such as switchgrass, hybrid poplar

wood, and miscanthus, fiber cane, and fiber sorghum. Five-carbon sugars that are produced from such materials include xylose.

[0115] Cellulosic materials are treated to increase the efficiency with which the microbe can utilize the sugar(s) contained within the materials. The invention provides novel methods for the treatment of cellulosic materials after acid explosion so that the materials are suitable for use in a heterotrophic culture of microbes (e.g., microalgae and oleaginous yeast). As discussed above, lignocellulosic biomass is comprised of various fractions, including cellulose, a crystalline polymer of beta 1,4 linked glucose (a six-carbon sugar), hemicellulose, a more loosely associated polymer predominantly comprised of xylose (a five-carbon sugar) and to a lesser extent mannose, galactose, arabinose, lignin, a complex aromatic polymer comprised of sinapyl alcohol and its derivatives, and pectins, which are linear chains of an alpha 1,4 linked polygalacturonic acid. Because of the polymeric structure of cellulose and hemicellulose, the sugars (e.g., monomeric glucose and xylose) in them are not in a form that can be efficiently used (metabolized) by many microbes. For such microbes, further processing of the cellulosic biomass to generate the monomeric sugars that make up the polymers can be very helpful to ensuring that the cellulosic materials are efficiently utilized as a feedstock (carbon source).

[0116] Celluose or cellulosic biomass is subjected to a process, termed "explosion", in which the biomass is treated with dilute sulfuric (or other) acid at elevated temperature and pressure. This process conditions the biomass such that it can be efficiently subjected to enzymatic hydrolysis of the cellulosic and hemicellulosic fractions into glucose and xylose monomers. The resulting monomeric sugars are termed cellulosic sugars. Cellulosic sugars can subsequently be utilized by microorganisms to produce a variety of metabolites (e.g., lipid). The acid explosion step results in a partial hydrolysis of the hemicellulose fraction to constitutent monosaccharides. These sugars can be completely liberated from the biomass with further treatment. In some embodiments, the further treatment is a hydrothermal treatment that includes washing the exploded material with hot water, which removes contaminants such as salts. This step is not necessary for cellulosic ethanol fermentations due to the more dilute sugar concentrations used in such processes. In other embodiments, the further treatment is additional acid treatment. In still other embodiments, the further treatment is enzymatic hydrolysis of the exploded material. These treatments can also be used in any combination. The type of treatment can affect the type of sugars liberated (e.g., five carbon sugars versus six carbon sugars) and the stage at which they are liberated in the process. As a consequence, different streams of sugars, whether they are predominantly five-carbon or six-

carbon, can be created. These enriched five-carbon or six-carbon streams can thus be directed to specific microorganisms with different carbon utilization cabilities.

[0117] The methods of the present invention typically involve fermentation to higher cell densities than what is achieved in ethanol fermentation. , Because of the higher densities of the cultures for heterotrophic cellulosic oil production, the fixed carbon source (e.g., the cellulosic derived sugar stream(s)) is preferably in a concentrated form. The glucose level of the depolymerized cellulosic material is preferably at least 300 g/liter, at least 400 g/liter, at least 500 g/liter or at least 600 g/liter prior to the cultivation step, which is optionally a fed batch cultivation in which the material is fed to the cells over time as the cells grow and accumulate lipid. Cellulosic sugar streams are not used at or near this concentration range in the production of cellulosic ethanol. Thus, in order to generate and sustain the very high cell densities during the production of lignocellulosic oil, the carbon feedstock(s) must be delivered into the heterotrophic cultures in a highly concentrated form. However, any component in the feedstream that is not a substrate for, and is not metabolized by, the oleaginous microorganism will accumulate in the bioreactor, which can lead to problems if the component is toxic or inhibitory to production of the desired end product. While ligin and lignin-derived by-products, carbohydrate-derived byproducts such as furfurals and hydroxymethyl furfurals and salts derived from the generation of the cellulosic materials (both in the explosion process and the subsequent neutralization process), and even non-metabolized pentose/hexose sugars can present problems in ethanolic fermentations, these effects are amplified significantly in a process in which their concentration in the initial feedstock is high. To achieve sugar concentrations in the 300g/L range (or higher) for six-carbon sugars that may be used in large scale production of lignocellulosic oil described in the present invention, the concentration of these toxic materials can be 20 times higher than the concentrations typically present in ethanolic fermentations of cellulosic biomass.

[0118] The explosion process treatment of the cellulosic material utilizes significant amounts of sulfuric acid, heat and pressure, thereby liberating by-products of carbohydrates, namely furfurals and hydroxymethyl furfurals. Furfurals and hydroxymethyl furfurals are produced during hydrolysis of hemicellulose through dehydration of xylose into furfural and water. In some embodiments of the present invention, these by-products (e.g., furfurals and hydroxymethyl furfurals) are removed from the saccharified lignocellulosic material prior to introduction into the bioreactor. In certain embodiments of the present invention, the process for removal of the by-products of carbohydrates is hydrothermal treatment of the exploded cellulosic materials. In addition, the present invention provides methods in which strains

capable of tolerating compounds such as furfurals or hydroxymethyl furfurals are used for lignocellulosic oil production. In another embodiment, the present invention also provides methods and microorganisms that are not only capable of tolerating furfurals in the fermentation media, but are actually able to metabolize these by-products during the production of lignocellulosic oil.

[0119] The explosion process also generates significant levels of salts. For example, typical conditions for explosion can result in conductivites in excess of 5 mS/cm when the exploded cellulosic biomass is resuspended at a ratio of 10:1 water:solids (dry weight). In certain embodiments of the present invention, the diluted exploded biomass is subjected to enzymatic saccharification, and the resulting supernatant is concentrated up to 25 fold for use in the bioreactor. The salt level (as measured by conductivity) in the concentrated sugar stream(s) can be unacceptably high (up to 1.5 M Na^+ equivalents). Additional salts are generated upon neutralization of the exploded materials for the subsequent enzymatic saccharification process as well. The present invention provides methods for removing these salts so that the resulting concentrated cellulosic sugar stream(s) can be used in heterotrophic processes for producing lignocellulosic oil. In some embodiments, the method of removing these salts is deionization with resins, such as, but not limited to, DOWEX Marathon MR3. In certain embodiments, the deionization with resin step occurs before sugar concentration or pH adjustment and hydrothermal treatment of biomass prior to saccharification, or any combination of the preceding; in other embodiments, the step is conducted after one or more of these processes. In other embodiments, the explosion process itself is changed so as to avoid the generation of salts at unacceptably high levels. For example, a suitable alternative to sulfuric acid (or other acid) explosion of the cellulosic biomass is mechanical pulping to render the cellulosic biomass receptive to enzymatic hydrolysis (saccharification). In still other embodiments, native strains of microorganisms resistant to high levels of salts or genetically engineered strains with resistance to high levels of salts are used.

[0120] A preferred embodiment for the process of preparing of exploded cellulosic biomass for use in heterotrophic lignocellulosic oil production using oleaginous microbes is diagramed in Figure 10. Step I. comprises adjusting the pH of the resuspended exploded cellulosic biomass to the range of 5.0-5.3 followed by washing the cellulosic biomass three times. This washing step can be accomplished by a variety of means including the use of desalting and ion exchange resins, reverse omosis, hydrothermal treatment (as described above), or just repeated re-suspension and centrifugation in deionized water. This wash step results in a cellulosic stream whose conductivity is between 100-300 μS/cm and the removal

of significant amounts of furfurals and hydroxymethyl furfurals. Decants from this wash step can be saved to concentrate five-carbon sugars liberated from the hemicellulose fraction. Step II comprises enzymatic saccharification of the washed cellulosic biomass. In a preferred embodiment, Accellerase (Genencor) is used. Step III comprises the recovery of sugars via centrifugation or decanting and rinsing of the saccharified biomass. The resulting biomass (solids) is an energy dense, lignin rich component that can be used as fuel or sent to waste. The recovered sugar stream in the centrifugation/decanting and rinse process is collected. Step IV comprises microfiltration to remove contaminating solids with recovery of the permeate. Step V comprises a concentration step which can be accomplished using a vacuum evaporator. This step can optionally include the addition of antifoam agents such as P'2000 (Sigma/Fluka), which is sometimes necessary due to the protein content of the resulting sugar feedstock.

[0121] In another embodiment of the methods of the invention, the carbon source is glycerol, including acidulated and non-acidulated glycerol byproduct from biodiesel transesterification. In one embodiment, the carbon source includes glycerol and at least one other carbone source. In some cases, all of the glycerol and the at least one other fixed carbon source are provided to the microorganism at the beginning of the fermentation. In some cases, the glycerol and the at least one other fixed carbon source are provided to the microorganism simultaneously at a predetermined ratio. In some cases, the glycerol and the at least one other fixed carbon source are fed to the microbes at a predetermined rate over the course of fermentation.

[0122] Some microalgae undergo cell division faster in the presence of glycerol than in the presence of glucose (see PCT Pub. No. 2008/151149). In these instances, two-stage growth processes in which cells are first fed glycerol to rapidly increase cell density, and are then fed glucose to accumulate lipids can improve the efficiency with which lipids are produced. The use of the glycerol byproduct of the transesterification process provides significant economic advantages when put back into the production process. Other feeding methods are provided as well, such as mixtures of glycerol and glucose. Feeding such mixtures also captures the same economic benefits. In addition, the invention provides methods of feeding alternative sugars to microalgae such as sucrose in various combinations with glycerol.

[0123] In another embodiment of the methods of the invention, the carbon source is sucrose, including a complex feedstock containing sucrose, such as thick cane juice from sugar cane processing. In one embodiment, the culture medium further includes at least one sucrose utilization enzyme. In some cases, the culture medium includes a sucrose invertase.

In one embodiment, the sucrose invertase enzyme is a secrectable sucrose invertase enzyme encoded by an exogenous sucrose invertase gene expressed by the population of microorganisms. Thus, in some cases, as described in more detail in Section IV, below, the microalgae has been genetically engineered to express a sucrose utilization enzyme, such as a sucrose transporter, a sucrose invertase, a hexokinase, a glucokinase, or a fructokinase.

[0124] Complex feedstocks containing sucrose include waste molasses from sugar cane processing; the use of this low-value waste product of sugar cane processing can provide significant cost savings in the production of hydrocarbons and other oils. Another complex feedstock containing sucrose that is useful in the methods of the invention is sorghum, including sorghum syrup and pure sorghum. Sorghum syrup is produced from the juice of sweet sorghum cane. Its sugar profile consists of mainly glucose (dextrose), fructose and sucrose.

4. Oil production

[0125] For the production of oil in accordance with the methods of the invention, it is preferable to culture cells in the dark, as is the case, for example, when using extremely large (40,000 liter and higher) fermentors that do not allow light to strike the culture. *Prototheca* species are grown and propagated for the production of oil in a medium containing a fixed carbon source and in the absence of light; such growth is known as heterotrophic growth.

[0126] As an example, an inoculum of lipid-producing microalgal cells are introduced into the medium; there is a lag period (lag phase) before the cells begin to propagate. Following the lag period, the propagation rate increases steadily and enters the log, or exponential, phase. The exponential phase is in turn followed by a slowing of propagation due to decreases in nutrients such as nitrogen, increases in toxic substances, and quorum sensing mechanisms. After this slowing, propagation stops, and the cells enter a stationary phase or steady growth state, depending on the particular environment provided to the cells. For obtaining lipid rich biomass, the culture is typically harvested well after then end of the exponential phase, which may be terminated early by allowing nitrogen or another key nutrient (other than carbon) to become depleted, forcing the cells to convert the carbon sources, present in excess, to lipid. Culture condition parameters can be manipulated to optimize total oil production, the combination of lipid species produced, and/or production of a specific oil.

[0127] As discussed above, a bioreactor or fermentor is used to allow cells to undergo the various phases of their growth cycle. As an example, an inoculum of lipid-producing cells can be introduced into a medium followed by a lag period (lag phase) before the cells begin

growth. Following the lag period, the growth rate increases steadily and enters the log, or exponential, phase. The exponential phase is in turn followed by a slowing of growth due to decreases in nutrients and/or increases in toxic substances. After this slowing, growth stops, and the cells enter a stationary phase or steady state, depending on the particular environment provided to the cells. Lipid production by cells disclosed herein can occur during the log phase or thereafter, including the stationary phase wherein nutrients are supplied, or still available, to allow the continuation of lipid production in the absence of cell division.

[0128] Preferably, microorganisms grown using conditions described herein and known in the art comprise at least about 20% by weight of lipid, preferably at least about 40% by weight, more preferably at least about 50% by weight, and most preferably at least about 60% by weight. Process conditions can be adjusted to increase the yield of lipids suitable for a particular use and/or to reduce production cost. For example, in certain embodiments, a microalgae is cultured in the presence of a limiting concentration of one or more nutrients, such as, for example, nitrogen, phosphorous, or sulfur, while providing an excess of fixed carbon energy such as glucose. Nitrogen limitation tends to increase microbial lipid yield over microbial lipid yield in a culture in which nitrogen is provided in excess. In particular embodiments, the increase in lipid yield is at least about: 10%, 50%, 100%, 200%, or 500%. The microbe can be cultured in the presence of a limiting amount of a nutrient for a portion of the total culture period or for the entire period. In particular embodiments, the nutrient concentration is cycled between a limiting concentration and a non-limiting concentration at least twice during the total culture period. Lipid content of cells can be increased by continuing the culture for increased periods of time while providing an excess of carbon, but limiting or no nitrogen.

[0129] In another embodiment, lipid yield is increased by culturing a lipid-producing microbe (e.g., microalgae) in the presence of one or more cofactor(s) for a lipid pathway enzyme (e.g., a fatty acid synthetic enzyme). Generally, the concentration of the cofactor(s) is sufficient to increase microbial lipid (e.g., fatty acid) yield over microbial lipid yield in the absence of the cofactor(s). In a particular embodiment, the cofactor(s) are provided to the culture by including in the culture a microbe (e.g., microalgae) containing an exogenous gene encoding the cofactor(s). Alternatively, cofactor(s) may be provided to a culture by including a microbe (e.g., microalgae) containing an exogenous gene that encodes a protein that participates in the synthesis of the cofactor. In certain embodiments, suitable cofactors include any vitamin required by a lipid pathway enzyme, such as, for example: biotin, pantothenate. Genes encoding cofactors suitable for use in the invention or that participate in

the synthesis of such cofactors are well known and can be introduced into microbes (e.g., microalgae), using contructs and techniques such as those described above.

[0130] The specific examples of bioreactors, culture conditions, and heterotrophic growth and propagation methods described herein can be combined in any suitable manner to improve efficiencies of microbial growth and lipid and/or protein production.

[0131] Microalgal biomass with a high percentage of oil/lipid accumulation by dry weight has been generated using different methods of culture, which are known in the art (see PCT Pub. No. 2008/151149). Microalgal biomass generated by the culture methods described herein and useful in accordance with the present invention comprises at least 10% microalgal oil by dry weight. In some embodiments, the microalgal biomass comprises at least 25%, at least 50%, at least 55%, or at least 60% microalgal oil by dry weight. In some embodiments, the microalgal biomass contains from 10-90% microalgal oil, from 25-75% microalgal oil, from 40-75% microalgal oil, or from 50-70% microalgal oil by dry weight.

[0132] The microalgal oil of the biomass described herein, or extracted from the biomass for use in the methods and compositions of the present invention can comprise glycerolipids with one or more distinct fatty acid ester side chains. Glycerolipids are comprised of a glycerol molecule esterified to one, two or three fatty acid molecules, which can be of varying lengths and have varying degrees of saturation. The length and saturation characteristics of the fatty acid molecules (and the microalgal oils) can be manipulated to modify the properties or proportions of the fatty acid molecules in the microalgal oils of the present invention via culture conditions or via lipid pathway engineering, as described in more detail in Section IV, below. Thus, specific blends of algal oil can be prepared either within a single species of algae by mixing together the biomass or algal oil from two or more species of microalgae, or by blending algal oil of the invention with oils from other sources such as soy, rapeseed, canola, palm, palm kernel, coconut, corn, waste vegetable, Chinese tallow, olive, sunflower, cottonseed, chicken fat, beef tallow, porcine tallow, microalgae, macroalgae, microbes, Cuphea, flax, peanut, choice white grease, lard, Camelina sativa, mustard seed, cashew nut, oats, lupine, kenaf, calendula, help, coffee, linseed (flax), hazelnut, euphorbia, pumpkin seed, coriander, camellia, sesame, safflower, rice, tung tree, cocoa, copra, pium poppy, castor beans, pecan, jojoba, macadamia, Brazil nuts, avocado, petroleum, or a distillate fraction of any of the preceding oils.

[0133] The oil composition, *i.e.*, the properties and proportions of the fatty acid consitutents of the glycerolipids, can also be manipulated by combining biomass or oil from at least two distinct species of microalgae. In some embodiments, at least two of the distinct

species of microalgae have different glycerolipid profiles. The distinct species of microalgae can be cultured together or separately as described herein, preferably under heterotrophic conditions, to generate the respective oils. Different species of microalgae can contain different percentages of distinct fatty acid consituents in the cell's glycerolipids.

[0134] Generally, *Prototheca* strains have very little or no fatty acids with the chain length C8-C14. For example, *Prototheca moriformis* (UTEX 1435), *Prototheca krugani* (UTEX 329), *Prototheca stagnora* (UTEX 1442) and *Prototheca zopfii* (UTEX 1438) contains no (or undectable amounts) C8 fatty acids, between 0-0.01% C10 fatty acids, between 0.03-2.1% C12 fatty acids and between 1.0-1.7% C14 fatty acids.

[0135] In some cases, the *Protheca* strains containing a transgene encoding a fatty acyl-ACP thioesterase that has activity towards fatty acyl-ACP substrate of chain lengths C8-10 has at least 0.3%, at least 0.8%, at least 1.5% or more fatty acids of chain length C8 and at least 0.3%, at least 1.0%, at least 3.0%, at least 5% or more fatty acids of chain length C10. In other instances, the *Prototheca* strains containing a transgene encoding a fatty acyl-ACP thioesterase that has activity towards fatty acyl-ACP substrate of chain length C12 has at least 3.0%, at least 5%, at least 7%, at least 10%, at least 13% or more fatty acids of the chain length C12 and at least 1.5%, at least 2%, or at least 3% or more fatty acids of the chain length C14. In other cases, the *Prototheca* strains containing a transgene encoding a fatty acyl-ACP thioesterase that has activity towards fatty acyl-ACP substrate of chain length C14 has at least 4.0%, at least 7%, at least 10%, at least 15%, at least 20%, at least 25% or more fatty acids of the chain length C14, and at least 0.4%, at least 1%, at least 1.5%, or more fatty acids of the chain length C12.

[0136] In non-limiting examples, the *Prototheca* strains containing a transgene encoding a fatty acyl-ACP thioesterase that has activity towards fatty acyl-ACP substrate of chain length C8 and C10 has between 0.3-1.58 % fatty acids of chain length C8 and between 0.35-6.76 % fatty acids of the chain length C10. In other non-limiting examples, *Prototheca* strains containing a transgene encoding a fatty acyl-ACP thioesterase that has activity towards fatty acyl-ACP substrate of chain length C12 has between 3.9-14.11% fatty acids of the chain length C12 and between 1.95-3.05% fatty acids of the chain length C14. In other non-limiting examples, *Prototheca* strains containing a transgene encoding a fatty acyl-ACP thioesterase that has activity towards fatty acyl-ACP substrate of chain length C14 has between 4.40-17.35 % fatty acids of the chain length C14 and between 0.4-1.83 Area % fatty acids of the chain length C12. In some cases, the *Prototheca* strains containing a transgene encoding a fatty acyl-ACP thioesterase that has activity towards fatty acyl-ACP substrate of chain

lengths between C8 and C14 have between 3.5- 20% medium chain (C8-C14) fatty acids. In some instances, keeping the transgenic *Prototheca* strains under constant and high selective pressure to retain exogenous genes is advantageous due to the increase in the desired fatty acid of a specific chain length. In a non-limiting example, Example 5 demonstrates a two fold increase in C14 chain length fatty acids (more than 30% C8-C14 chain length fatty acids) when the culture of *Prototheca moriformis* containing a C14 preferring thioesterase exogenous gene is retained. High levels of exogenous gene retention can also be achieved by inserting exogenous genes into the nuclear chromosomes of the cells using homologous recombination vectors and methods disclosed herein. Recombinant cells containing exogenous genes integrated into nuclear chromosomes are an object of the invention.

[0137] Microalgal oil can also include other constituents produced by the microalgae, or incorporated into the microalgal oil from the culture medium. These other constituents can be present in varying amount depending on the culture conditions used to culture the microalgae, the species of microalgae, the extraction method used to recover microalgal oil from the biomass and other factors that may affect microalgal oil composition. Non-limiting examples of such constituents include carotenoids, present from 0.1-0.4 micrograms/ml, chlorophyll present from 0-0.02 milligrams/kilogram of oil, gamma tocopherol present from 0.4-0.6 milligrams/100 grams of oil, and total tocotrienols present from 0.2-0.5 milligrams/gram of oil.

[0138] The other constituents can include, without limitation, phospholipids, tocopherols, tocotrienols, carotenoids (*e.g.*, alpha-carotene, beta-carotene, lycopene, etc.), xanthophylls (*e.g.*, lutein, zeaxanthin, alpha-cryptoxanthin and beta-crytoxanthin), and various organic or inorganic compounds.

[0139] In some cases, the oil extracted from *Prototheca* species comprises no more than 0.02mg/kg chlorophyll. In some cases, the oil extracted from *Prototheca* species comprises no more than 0.4 mcg/ml total carotenoids. In some cases the *Prototheca* oil comprises between 0.40-0.60 milligrams of gamma tocopherol per 100 grams of oil. In other cases, the *Prototheca* oil comprises between 0.2-0.5 milligrams of total tocotrienols per gram of oil.

III. GENETIC ENGINEERING METHODS AND MATERIALS

[0140] The present invention provides methods and materials for genentically modifying Prototheca cells and recombinant host cells useful in the methods of the present invention, including but not limited to recombinant *Prototheca moriformis, Prototheca zopfii, Prototheca krugani,* and *Prototheca stagnora* host cells. The description of these methods and materials is divided into subsections for the convenience of the reader. In subsection 1,

transformation methods are described. In subsection 2, genetic engineering methods using homologous recombination are described. In subsection 3, expression vectors and components are described.

1. Engineering Methods - Transformation

[0141] Cells can be transformed by any suitable technique including, e.g., biolistics, electroporation (see Maruyama et al. (2004), Biotechnology Techniques 8:821-826), glass bead transformation and silicon carbide whisker transformation. Another method that can be used involves forming protoplasts and using $CaCl_2$ and polyethylene glycol (PEG) to introduce recombinant DNA into microalgal cells (see Kim et al. (2002), *Mar. Biotechnol.* 4:63-73, which reports the use of this method for the transformation of *Chorella ellipsoidea*). Co-transformation of microalgae can be used to introduce two distinct vector molecules into a cell simultaneously (see for example Protist 2004 Dec;155(4):381-93).

[0142] Biolistic methods (see, for example, Sanford, Trends In Biotech. (1988) 6:299 302, U.S. Patent No. 4,945,050; electroporation (Fromm et al., Proc. Nat'l. Acad. Sci. (USA) (1985) 82:5824 5828); use of a laser beam, microinjection or any other method capable of introducing DNA into a microalgae can also be used for transformation of a *Prototheca* cell.

2. Engineering Methods - Homologous Recombination

[0143] Homologous recombination is the ability of complementary DNA sequences to align and exchange regions of homology. Transgenic DNA ("donor") containing sequences homologous to the genomic sequences being targeted ("template") is introduced into the organism and then undergoes recombination into the genome at the site of the corresponding genomic homologous sequences. The mechanistic steps of this process, in most casees, include: (1) pairing of homologous DNA segments; (2) introduction of double-stranded breaks into the donor DNA molecule; (3) invasion of the template DNA molecule by the free donor DNA ends followed by DNA synthesis; and (4) resolution of double-strand break repair events that result in final recombination products.

[0144] The ability to carry out homologous recombination in a host organism has many practical implications for what can be carried out at the molecular genetic level and is useful in the generation of an oleaginous microbe that can produced tailored oils. By its very nature homologous recombination is a precise gene targeting event, hence, most transgenic lines generated with the same targeting sequence will be essentially identical in terms of phenotype, necessitating the screening of far fewer transformation events. Homologous recombination also targets gene insertion events into the host chromosome, resulting in excellent genetic stability, even in the absence of genetic selection. Because different

chromosomal loci will likey impact gene expression, even from heterologous promoters/UTRs, homologous recombination can be a method of querying loci in an unfamiliar genome environment and to assess the impact of these environments on gene expression.

[0145] Particularly useful genetic engineering applications using homologous recombination is to co-opt specific host regulatory elements such as promoters/UTRs to drive heterologous gene expression in a highly specific fashion. For example, precise ablation of the endogenous stearoyl ACP desaturase gene with a heterologous C12:0 specific FATB (thioesterase) gene cassette and suitable selective marker, might be expected to dramatically decrease endogenous levels of C18:1 fatty acids concomitant with increased levels of the C12:0 fatty acids. Example 13 describes the homologous recombination targeting construct that is suitable for the eblation of an endogenous *Prototheca moriformis* stearoyl ACP destaurase gene.

[0146] Because homologous recombination is a precise gene targeting event, it can be used to precisely modify any nucleotide(s) within a gene or region of interest, so long as sufficient flanking regions have been identified. Therefore, homologous recombination can be used as a means to modify regulatory sequences impacting gene expression of RNA and/or proteins. It can also be used to modify protein coding regions in an effort to modify enzyme activites such as substrate specificity, affinities and Km, and thus affecting the desired change in metabolism of the host cell. Homologous recombination provides a powerful means to manipulate the gost genome resulting in gene targeting, gene conversion, gene deletion, gene duplication, gene inversion and exchanging gene expression regulatory elements such as promoters, enhancers and 3'UTRs.

[0147] Homologous recombination can be achieve by using targeting constructs containing pieces of endogenous sequences to "target" the gene or region of interest within the endogenous host cell genome. Such targeting sequences can either be located 5' of the gene or region of interest, 3' of the gene/region of interest or even flank the gene/region of interest. Such targeting constructs can be transformed into the host cell either as a supercoiled plasmid DNA with additional vector backbone, a PCR product with no vector backbone, or as a linearized molecule. In some cases, it may be advantageous to first expose the homologous sequences within the transgenic DNA (donor DNA) with a restriction enzyme. This step can increase the recombination efficiency and decrease the occurance of undesired events. Other methods of increasing recombination efficiency include using PCR to generate transforming transgenic DNA containing linear ends homologous to the genomic sequences being targeted.

3. Vectors and Vector Components

[0148] Vectors for transformation of microorganisms in accordance with the present invention can be prepared by known techniques familiar to those skilled in the art in view of the disclosure herein. A vector typically contains one or more genes, in which each gene codes for the expression of a desired product (the gene product) and is operably linked to one or more control sequences that regulate gene expression or target the gene product to a particular location in the recombinant cell. To aid the reader, this subsection is divided into subsections. Subsection A describes control sequences typically contained on vectors as well as novel control sequences provided by the present invention. Subsection B describes genes typically contained in vectors as well as novel codon optimization methods and genes prepared using them provided by the invention.

A. Control Sequences

[0149] Control sequences are nucleic acids that regulate the expression of a coding sequence or direct a gene product to a particular location in or outside a cell. Control sequences that regulate expression include, for example, promoters that regulate transcription of a coding sequence and terminators that terminate transcription of a coding sequence. Another control sequence is a 3' untranslated sequence located at the end of a coding sequence that encodes a polyadenylation signal. Control sequences that direct gene products to particular locations include those that encode signal peptides, which direct the protein to which they are attached to a particular location in or outside the cell.

[0150] Thus, an exemplary vector design for expression of an exogenous gene in a microalgae contains a coding sequence for a desired gene product (for example, a selectable marker, a lipid pathway modification enzyme, or a sucrose utilization enzyme) in operable linkage with a promoter active in microalgae. Alternatively, if the vector does not contain a promoter in operable linkage with the coding sequence of interest, the coding sequence can be transformed into the cells such that it becomes operably linked to an endogenous promoter at the point of vector integration. The promoterless method of transformation has been proven to work in microalgae (see for example Plant Journal 14:4, (1998), pp.441-447).

[0151] Many promoters are active in microalgae, including promoters that are endogenous to the algae being transformed, as well as promoters that are not endogenous to the algae being transformed (i.e., promoters from other algae, promoters from higher plants, and promoters from plant viruses or algae viruses). Illustrative exogenous and/or endogenous promoters that are active in microalgae (as well as antibiotic resistance genes functional in microalgae) are described in PCT Pub. No. 2008/151149 and references cited therein).

[0152] The promoter used to express an exogenous gene can be the promoter naturally linked to that gene or can be a heterologous gene. Some promoters are active in more than one species of microalgae. Other promoters are species-specific. Illustrative promoters include promoters such as β-tubulin from *Chlamydomonas reinhardtii*, used in the Examples below,and viral promoters, such as cauliflower mosaic virus (CMV) and chlorella virus, which have been shown to be active in multiple species of microalgae (see for example Plant Cell Rep. 2005 Mar;23(10-11):727-35; J Microbiol. 2005 Aug;43(4):361-5; Mar Biotechnol (NY). 2002 Jan;4(1):63-73). Another promoter that is suitable for use for expression of exogenous genes in *Prototheca* is the *Chlorella sorokiniana* glutamate dehydrogenase promoter/5'UTR (SEQ ID NO: 69). Optionally, at least 10, 20, 30, 40, 50, or 60 nucleotides or more of these sequences containing a promoter are used. Illustrative promoters useful for expression of exogenous genes in *Prototheca* are listed in the sequence listing of this application, such as the promoter of the *Chlorella* HUP1 gene (SEQ ID NO:1) and the *Chlorella ellipsoidea* nitrate reductase promoter (SEQ ID NO:2). Chlorella virus promoters can also be used to express genes in *Prototheca*, such as SEQ ID NOs: 1-7 of U.S. Patent 6,395,965. Additional promoters active in *Prototheca* can be found, for example, in Biochem Biophys Res Commun. 1994 Oct 14;204(1):187-94; Plant Mol Biol. 1994 Oct;26(1):85-93; Virology. 2004 Aug 15;326(1):150-9; and Virology. 2004 Jan 5;318(1):214-23.

[0153] A promoter can generally be characterized as either constitutive or inducible. Constitutive promoters are generally active or function to drive expression at all times (or at certain times in the cell life cycle) at the same level. Inducible promoters, conversely, are active (or rendered inactive) or are significantly up- or down-regulated only in response to a stimulus. Both types of promoters find application in the methods of the invention. Inducible promoters useful in the invention include those that mediate transcription of an operably linked gene in response to a stimulus, such as an exogenously provided small molecule (e.g, glucose, as in SEQ ID NO:1), temperature (heat or cold), lack of nitrogen in culture media, etc. Suitable promoters can activate transcription of an essentially silent gene or upregulate, preferably substantially, transcription of an operably linked gene that is transcribed at a low level.

[0154] Inclusion of termination region control sequence is optional, and if employed, then the choice is be primarily one of convenience, as the termination region is relatively interchangeable. The termination region may be native to the transcriptional initiation region (the promoter), may be native to the DNA sequence of interest, or may be obtainable from another source. See, for example, Chen and Orozco, Nucleic Acids Res. (1988) 16:8411.

[0155] The present invention also provides control sequences and recombinant genes and vectors containing them that provide for the compartmentalized expression of a gene of interest. Organelles for targeting are chloroplasts, plastids, mitochondria, and endoplasmic reticulum. In addition, the present invention provides control sequences and recombinant genes and vectors containing them that provide for the secretion of a protein outside the cell.

[0156] Proteins expressed in the nuclear genome of *Prototheca* can be targeted to the plastid using plastid targeting signals. Plastid targeting sequences endogenous to *Chlorella* are known, such as genes in the *Chlorella* nuclear genome that encode proteins that are targeted to the plastid; see for example GenBank Accession numbers AY646197 and AF499684, and in one embodiment, such control sequences are used in the vectors of the present invention to target expression of a protein to a *Prototheca* plastid.

[0157] The Examples below describe the use of algal plastid targeting sequences to target heterologous proteins to the correct compartment in the host cell. cDNA libraries were made using *Prototheca moriformis* and *Chlorella protothecodies* cells and are described in Examples 12 and Example 11 below. Sequences were BLASTed and analyzed for homology to known proteins that traffic to the plastid/chloroplast. The cDNAs encoding these proteins were cloned and plastid targeting sequences were isolated from these cDNAs. The amino acid sequences of the algal plastid targeting sequences identified from the cDNA libraries and the amino acid sequences of plant fatty acyl-ACP thioesterases that are used in the heterologous expression Examples below are listed in SEQ ID NOs: 127-133.

[0158] In another embodiment of the present invention, the expression of a polypeptide in *Prototheca* is targeted to the endoplasmic reticulum. The inclusion of an appropriate retention or sorting signal in an expression vector ensure that proteins are retained in the endoplasmic reticulum (ER) and do not go downstream into Golgi. For example, the IMPACTVECTOR1.3 vector, from Wageningen UR- Plant Research International, includes the well known KDEL retention or sorting signal. With this vector, ER retention has a practical advantage in that it has been reported to improve expression levels 5-fold or more. The main reason for this appears to be that the ER contains lower concentrations and/or different proteases responsible for post-translational degradation of expressed proteins than are present in the cytoplasm. ER retention signals functional in green microalgae are known. For example, see Proc Natl Acad Sci U S A. 2005 Apr 26;102(17):6225-30.

[0159] In another embodiment of the present invention, a polypeptide is targeted for secretion outside the cell into the culture media. See Hawkins et al., Current Microbiology

Vol. 38 (1999), pp. 335–341 for examples of secretion signals active in *Chlorella* that can be used, in accordance with the methods of the invention, in *Prototheca*.

B. Genes and Codon Optimization

[0160] Typically, a gene includes a promoter, coding sequence, and termination control sequences. When assembled by recombinant DNA technology, a gene may be termed an expression cassette and may be flanked by restriction sites for convenient insertion into a vector that is used to introduce the recombinant gene into a host cell. The expression cassette can be flanked by DNA sequences from the genome or other nucleic acid target to facilitate stable integration of the expression cassette into the genome by homologous recombination. Alternatively, the vector and its expression cassette may remain unintegrated, in which case, the vector typically includes an origin of replication, which is capable of providing for replication of the heterologous vector DNA.

[0161] A common gene present on a vector is a gene that codes for a protein, the expression of which allows the recombinant cell containing the protein to be differentiated from cells that do not express the protein. Such a gene, and its corresponding gene product, is called a selectable marker. Any of a wide variety of selectable markers can be employed in a transgene construct useful for transforming *Prototheca*. Examples of suitable selectable markers include the G418 resistance gene, the nitrate reductase gene (see Dawson et al. (1997), Current Microbiology 35:356-362), the hygromycin phosphotransferase gene (HPT; see Kim et al. (2002), Mar. Biotechnol. 4:63-73), the neomycin phosphotransferase gene, and the ble gene, which confers resistance to phleomycin (Huang et al. (2007), Appl. Microbiol. Biotechnol. 72:197-205). Methods of determining sensitivity of microalgae to antibiotics are well known. For example, Mol Gen Genet. 1996 Oct 16;252(5):572-9.

[0162] For purposes of the present invention, the expression vector used to prepare a recombinant host cell of the invention will include at least two, and often three, genes, if one of the genes is a selectable marker. For example, a genetically engineered *Prototheca* of the invention can be made by transformation with vectors of the invention that comprise, in addition to a selectable marker, one or more exogenous genes, such as, for example, sucrose invertase gene or acyl ACP-thioesterase gene. One or both genes can be expressed using an inducible promoter, which allows the relative timing of expression of these genes to be controlled to enhance the lipid yield and conversion to fatty acid esters. Expression of the two or more exogenous genes may be under control of the same inducible promoter or under control of different inducible (or constitutive) promoters. In the latter situation, expression of a first exogenous gene can be induced for a first period of time (during which expression of a

second exogenous gene may or may not be induced) and expression of a second exogenous gene can be induced for a second period of time (during which expression of a first exogenous gene may or may not be induced).

[0163] In other embodiments, the two or more exogenous genes (in addition to any selectable marker) are: a fatty acyl-ACP thioesterase and a fatty acyl-CoA/aldehyde reductase, the combined action of which yields an alcohol product. Further provided are other combinations of exogenous genes, including without limitation, a fatty acyl-ACP thioesterase and a fatty acyl-CoA reductase to generate aldehydes. In one embodiment, the vector provides for the combination of a fatty acyl-ACP thioesterase, a fatty acyl-CoA reductase, and a fatty aldehyde decarbonylase to generate alkanes. In each of these embodiments, one or more of the exogenous genes can be expressed using an inducible promoter.

[0164] Other illustrative vectors of the invention that express two or more exogenous genes include those encoding both a sucrose transporter and a sucrose invertase enzyme and those encoding both a selectable marker and a secreted sucrose invertase. The recombinant *Prototheca* transformed with either type of vector produce lipids at lower manufacturing cost due to the engineered ability to use sugar cane (and sugar cane-derived sugars) as a carbon source. Insertion of the two exogenous genes described above can be combined with the disruption of polysaccharide biosynthesis through directed and/or random mutagenesis, which steers ever greater carbon flux into lipid production. Individually and in combination, trophic conversion, engineering to alter lipid production and treatment with exogenous enzymes alter the lipid composition produced by a microorganism. The alteration can be a change in the amount of lipids produced, the amount of one or more hydrocarbon species produced relative to other lipids, and/or the types of lipid species produced in the microorganism. For example, microalgae can be engineered to produce a higher amount and/or percentage of TAGs.

[0165] For optimal expression of a recombinant protein, it is beneficial to employ coding sequences that produce mRNA with codons preferentially used by the host cell to be transformed. Thus, proper expression of transgenes can require that the codon usage of the transgene matches the specific codon bias of the organism in which the transgene is being expressed. The precise mechanisms underlying this effect are many, but include the proper balancing of available aminoacylated tRNA pools with proteins being synthesized in the cell, coupled with more efficient translation of the transgenic messenger RNA (mRNA) when this need is met. When codon usage in the transgene is not optimized, available tRNA pools are

not sufficient to allow for efficient translation of the heterologous mRNA resulting in ribosomal stalling and termination and possible instability of the transgenic mRNA.

[0166] The present invention provides codon-optimized nucleic acids useful for the successful expression of recombinant proteins in *Prototheca*. Codon usage in *Prototheca* species was analyzed by studying cDNA sequences isolated from *Prototheca moriformis*. This analysis represents the interrogation over 24, 000 codons and resulted in Table 1 below.

[0167] Table 1. Preferred codon usage in *Prototheca* strains.

Ala	GCG	345 (0.36)	Asn	AAT	8 (0.04)
	GCA	66 (0.07)		AAC	201 (0.96)
	GCT	101 (0.11)			
	GCC	442 (0.46)	Pro	CCG	161 (0.29)
				CCA	49 (0.09)
Cys	TGT	12 (0.10)		CCT	71 (0.13)
	TGC	105 (0.90)		CCC	267 (0.49)
Asp	GAT	43 (0.12)	Gln	CAG	226 (0.82)
	GAC	316 (0.88)		CAA	48 (0.18)
Glu	GAG	377 (0.96)	Arg	AGG	33 (0.06)
	GAA	14 (0.04)		AGA	14 (0.02)
				CGG	102 (0.18)
Phe	TTT	89 (0.29)		CGA	49 (0.08)
	TTC	216 (0.71)		CGT	51 (0.09)
				CGC	331 (0.57)
Gly	GGG	92 (0.12)			
	GGA	56 (0.07)	Ser	AGT	16 (0.03)
	GGT	76 (0.10)		AGC	123 (0.22)
	GGC	559 (0.71)		TCG	152 (0.28)
				TCA	31 (0.06)
His	CAT	42 (0.21)		TCT	55 (0.10)
	CAC	154 (0.79)		TCC	173 (0.31)
Ile	ATA	4 (0.01)	Thr	ACG	184 (0.38)
	ATT	30 (0.08)		ACA	24 (0.05)
	ATC	338 (0.91)		ACT	21 (0.05)
				ACC	249 (0.52)
Lys	AAG	284 (0.98)			
	AAA	7 (0.02)	Val	GTG	308 (0.50)
				GTA	9 (0.01)
Leu	TTG	26 (0.04)		GTT	35 (0.06)
	TTA	3 (0.00)		GTC	262 (0.43)
	CTG	447 (0.61)			
	CTA	20 (0.03)	Trp	TGG	107 (1.00)
	CTT	45 (0.06)			
	CTC	190 (0.26)	Tyr	TAT	10 (0.05)
				TAC	180 (0.95)

Met	ATG	191 (1.00)	
		Stop	TGA/TAG/TAA

[0168] In other embodiments, the gene in the recombinant vector has been codon-optimized with reference to a microalgal strain other than a *Prototheca* strain. For example, methods of recoding genes for expression in microalgae are described in U.S. Patent 7,135,290. Additional information for codon optimization is available, e.g., at the codon usage database of GenBank.

[0169] While the methods and materials of the invention allow for the introduction of any exogenous gene into *Prototheca*, genes relating to sucrose utilization and lipid pathway modification are of particular interest, as discussed in the following sections.

IV. SUCROSE UTILIZATION

[0170] In embodiment, the recombinant *Prototheca* cell of the invention further contains one or more exogenous sucrose utilization genes. In various embodiments, the one or more genes encode one or more proteins selected from the group consisting of a fructokinase, a glucokinase, a hexokinase, a sucrose invertase, a sucrose transporter. For example, expression of a sucrose transporter and a sucrose invertase allows *Prototheca* to transport sucrose into the cell from the culture media and hydrolyze sucrose to yield glucose and fructose. Optionally, a fructokinase can be expressed as well in instances where endogenous hexokinase activity is insufficient for maximum phosphorylation of fructose. Examples of suitable sucrose transporters are Genbank accession numbers CAD91334, CAB92307, and CAA53390. Examples of suitable fructokinases are Genbank accession numbers P26984, P26420 and CAA43322.

[0171] In one embodiment, the present invention provides a *Prototheca* host cell that secretes a sucrose invertase. Secretion of a sucrose invertase obviates the need for expression of a transporter that can transport sucrose into the cell. This is because a secreted invertase catalyzes the conversion of a molecule of sucrose into a molecule of glucose and a molecule of fructose, both of which can be transported and utilized by microbes provided by the invention. For example, expression of a sucrose invertase (such as SEQ ID NO:3) with a secretion signal (such as that of SEQ ID NO: 4 (from yeast), SEQ ID NO: 5 (from higher plants), SEQ ID NO: 6 (eukaryotic consensus secretion signal), and SEQ ID NO: 7 (combination of signal sequence from higher plants and eukaryotic consensus) generates invertase activity outside the cell. Expression of such a protein, as enabled by the genetic

engineering methodology disclosed herein, allows cells already capable of utilizing extracellular glucose as an energy source to utilize sucrose as an extracellular energy source.

[0172] *Prototheca* species expressing an invertase in media containing sucrose are a preferred microalgal species for the production of oil. Example 3 illustrates how the methods and reagents of the invention can be used to express a recombinant yeast invertase and secrete it from a recombinant *Prototheca* cell. The expression and extracellular targeting of this fully active protein allows the resulting host cells to grow on sucrose, whereas their non-transformed counterparts cannot. Thus, the present invention provides *Prototheca* recombinant cells with a codon-optimized invertase gene, including but not limited to the yeast invertase gene, integrated into their genome such that the invertase gene is expressed as assessed by invertase activity and sucrose hydrolysis. The present invention also provides invertase genes useful as selectable markers in *Prototheca* recombinant cells, as such cells are able to grow on sucrose, while their non-transformed counterparts cannot; and methods for selecting recombinant host cells using an invertase as a powerful, selectable marker for algal molecular genetics.

[0173] The successful expression of a sucrose invertase in *Prototheca* also illustrates another aspect of the present invention in that it demonstrates that heterologous (recombinant) proteins can be expressed in the algal cell and successfully transit outside of the cell and into the culture medium in a fully active and functional form. Thus, the present invention provides methods and reagents for expressing a wide and diverse array of heterologous proteins in microalgae and secreting them outside of the host cell. Such proteins include, for example, industrial enzymes such as, for example, lipases, proteases, cellulases, pectinases, amylases, esterases, oxidoreductases, transferases, lactases, isomerases, and invertases, as well as therapeutic proteins such as, for example, growth factors, cytokines, full length antibodies comprising two light and two heavy chains, Fabs, scFvs (single chain variable fragment), camellid-type antibodies, antibody fragments, antibody fragment-fusions, antibody-receptor fusions, insulin, interferons, and insulin-like growth factors.

[0174] The successful expression of a sucrose invertase in *Prototheca* also illustrates another aspect of the present invention in that it provides methods and reagents for the use of fungal transit peptides in algae to direct secretion of proteins in *Prototheca*; and methods and reagents for determining if a peptide can function, and the ability of it to function, as a transit peptide in *Prototheca* cells. The methods and reagents of the invention can be used as a tool and platform to identify other transit peptides that can successfully traffic proteins outside of a cell, and that the yeast invertase has great utility in these methods. As demonstrated in this

example, removal of the endogenous yeast invertase transit peptide and its replacement by other transit peptides, either endogenous to the host algae or from other sources (eukaryotic, prokaryotic and viral), can identify whether any peptide of interest can function as a transit peptide in guiding protein egress from the cell.

[0175] Examples of suitable sucrose invertases include those identified by Genbank accession numbers CAB95010, NP_012104 and CAA06839. Non-limiting examples of suitable invertases are listed below in Table 2. Amino acid sequences for each listed invertase are included in the Sequence Listing below. In some cases, the exogenous sucrose utilization gene suitable for use in the methods and vectors of the invention encodes a sucrose invertase that has at least 40, 50, 60, 75, or 90% or higher amino acid identity with a sucrose invertase selected from Table 2.

[0176] Table 2. Sucrose invertases.

Description	Organism	GenBank Accession No.	SEQ ID NO:
Invertase	*Chicorium intybus*	Y11124	SEQ ID NO:20
Invertase	*Schizosaccharomyces pombe*	AB011433	SEQ ID NO:21
beta-fructofuranosidase (invertase)	*Pichia anomala*	X80640	SEQ ID NO:22
Invertase	*Debaryomyces occidentalis*	X17604	SEQ ID NO:23
Invertase	*Oryza sativa*	AF019113	SEQ ID NO:24
Invertase	*Allium cepa*	AJ006067	SEQ ID NO:25
Invertase	*Beta vulgaris subsp. Vulgaris*	AJ278531	SEQ ID NO:26
beta-fructofuranosidase (invertase)	*Bifidobacterium breve UCC2003*	AAT28190	SEQ ID NO:27
Invertase	*Saccharomyces cerevisiae*	NP_012104	SEQ ID NO:8 (nucleotide) SEQ ID NO:28 (amino acid)
Invertase A	*Zymomonas mobilis*	AAO38865	SEQ ID NO:29

[0177] The secretion of an invertase to the culture medium by *Prototheca* enable the cells to grow as well on waste molasses from sugar cane processing as they do on pure reagent-grade glucose; the use of this low-value waste product of sugar cane processing can provide significant cost savings in the production of lipids and other oils. Thus, the present invention provides a microbial culture containing a population of *Prototheca* microorganisms, and a culture medium comprising (i) sucrose and (ii) a sucrose invertase enzyme. In various embodiments the sucrose in the culture comes from sorghum, sugar beet, sugar cane, molasses, or depolymerized cellulosic material (which may optionally contain lignin). In another aspect, the methods and reagents of the invention significantly increase the number and type of feedstocks that can be utilized by recombinant *Prototheca*. While the microbes exemplified here are altered such that they can utilize sucrose, the methods and reagents of the invention can be applied so that feedstocks such as cellulosics are utilizable by an engineered host microbe of the invention with the ability to secrete cellulases, pectinases, isomerases, or the like, such that the breakdown products of the enzymatic reactions are no longer just simply tolerated but rather utilized as a carbon source by the host.

V. LIPID PATHWAY ENGINEERING

[0178] In addition to altering the ability of *Prototheca* to utilize feedstocks such as sucrose-containing feedstocks, the present invention also provides recombinant *Prototheca* that have been modified to alter the properties and/or proportions of lipids produced. The pathway can further, or alternatively, be modified to alter the properties and/or proportions of various lipid molecules produced through enzymatic processing of lipids and intermediates in the fatty acid pathway. In various embodiments, the recombinant *Prototheca* cells of the invention have, relative to their untransformed counterparts, optimized lipid yield per unit volume and/or per unit time, carbon chain length (e.g., for renewable diesel production or for industrial chemicals applications requiring lipid feedstock), reduced number of double or triple bonds, optionally to zero, and increasing the hydrogen:carbon ratio of a particular species of lipid or of a population of distinct lipid.

[0179] In particular embodiments, one or more key enzymes that control branch points in metabolism to fatty acid synthesis have been up-regulated or down-regulated to improve lipid production. Up-regulation can be achieved, for example, by transforming cells with expression constructs in which a gene encoding the enzyme of interest is expressed, e.g., using a strong promoter and/or enhancer elements that increase transcription. Such constructs can include a selectable marker such that the transformants can be subjected to selection, which can result in amplification of the construct and an increase in the expression level of

the encoded enzyme. Examples of enzymes suitable for up-regulation according to the methods of the invention include pyruvate dehydrogenase, which plays a role in converting pyruvate to acetyl-CoA (examples, some from microalgae, include Genbank accession numbers NP_415392; AAA53047; Q1XDM1; and CAF05587). Up-regulation of pyruvate dehydrogenase can increase production of acetyl-CoA, and thereby increase fatty acid synthesis. Acetyl-CoA carboxylase catalyzes the initial step in fatty acid synthesis. Accordingly, this enzyme can be up-regulated to increase production of fatty acids (examples, some from microalgae, include Genbank accession numbers BAA94752; AAA75528; AAA81471; YP_537052; YP_536879; NP_045833; and BAA57908). Fatty acid production can also be increased by up-regulation of acyl carrier protein (ACP), which carries the growing acyl chains during fatty acid synthesis (examples, some from microalgae, include Genbank accession numbers A0T0F8; P51280; NP_849041; YP_874433). Glycerol-3-phosphate acyltransferase catalyzes the rate-limiting step of fatty acid synthesis. Up-regulation of this enzyme can increase fatty acid production (examples, some from microalgae, include Genbank accession numbers AAA74319; AAA33122; AAA37647; P44857; and ABO94442).

[0180] Up- and/or down-regulation of genes can be applied to global regulators controlling the expression of the genes of the fatty acid biosynthetic pathways. Accordingly, one or more global regulators of fatty acid synthesis can be up- or down-regulated, as appropriate, to inhibit or enhance, respectively, the expression of a plurality of fatty acid synthetic genes and, ultimately, to increase lipid production. Examples include sterol regulatory element binding proteins (SREBPs), such as SREBP-1a and SREBP-1c (for examples see Genbank accession numbers NP_035610 and Q9WTN3).

[0181] The present invention also provides recombinant *Prototheca* cells that have been modified to contain one or more exogenous genes encoding lipid modification enzymes such as, for example, fatty acyl-ACP thioesterases (see Table 3), fatty acyl-CoA/aldehyde reductases (see Table 4), fatty acyl-CoA reductases (see Table 5), fatty aldehyde decarbonylase (see Table 6), fatty aldehyde reductases, and squalene synthases (see GenBank Accession number AF205791). In some embodiments, genes encoding a fatty acyl-ACP thioesterase and a naturally co-expressed acyl carrier protein are transformed into a *Prototheca* cell, optionally with one or more genes encoding other lipid modification enzymes. In other embodiments, the ACP and the fatty acyl-ACP thioesterase may have an affinity for one another that imparts an advantage when the two are used together in the microbes and methods of the present invention, irrespective of whether they are or are not

naturally co-expressed in a particular tissue or organism. Thus, the present invention contemplates both naturally co-expressed pairs of these enzymes as well as those that share an affinity for interacting with one another to facilitate cleavage of a length-specific carbon chain from the ACP.

[0182] In still other embodiments, an exogenous gene encoding a desaturase is transformed into the *Prototheca* cell in conjunction with one or more genes encoding other lipid modification enzymes to provide modifications with respect to lipid saturation. Stearoyl-ACP desaturase (see, e.g., GenBank Accession numbers AAF15308; ABM45911; and AAY86086), for example, catalyzes the conversion of stearoyl-ACP to oleoyl-ACP. Up-regulation of this gene can increase the proportion of monounsaturated fatty acids produced by a cell; whereas down-regulation can reduce the proportion of monounsaturates. Similarly, the expression of one or more glycerolipid desaturases can be controlled to alter the ratio of unsaturated to saturated fatty acids such as ω-6 fatty acid desaturase, ω-3 fatty acid desaturase, or ω-6-oleate desaturase. In some embodiments, the desaturase can be selected with reference to a desired carbon chain length, such that the desaturase is capable of making location specific modifications within a specified carbon-length substrate, or substrates having a carbon-length within a specified range.

[0183] Thus, in particular embodiments, microbes of the present invention are genetically engineered to express one or more exogenous genes selected from an acyl-ACP thioesterase, an acyl-CoA/aldehyde reductase, a fatty acyl-CoA reductase, a fatty aldehyde reductase, a fatty aldehyde decarbonylase, or a naturally co-expressed acyl carrier protein. Suitable expression methods are described above with respect to the expression of a lipase gene, including, among other methods, inducible expression and compartmentalized expression. A fatty acyl-ACP thioesterase cleaves a fatty acid from an acyl carrier protein (ACP) during lipid synthesis. Through further enzymatic processing, the cleaved fatty acid is then combined with a coenzyme to yield an acyl-CoA molecule. This acyl-CoA is the substrate for the enzymatic activity of a fatty acyl-CoA reductase to yield an aldehyde, as well as for a fatty acyl-CoA/aldehyde reductase to yield an alcohol. The aldehyde produced by the action of the fatty acyl-CoA reductase identified above is the substrate for further enzymatic activity by either a fatty aldehyde reductase to yield an alcohol, or a fatty aldehyde decarbonylase to yield an alkane or alkene.

[0184] In some embodiments, fatty acids, glycerolipids, or the corresponding primary alcohols, aldehydes, alkanes or alkenes, generated by the methods described herein, contain 8, 10, 12,or 14 carbon atoms. Preferred fatty acids for the production of diesel, biodiesel,

renewable diesel, or jet fuel, or the corresponding primary alcohols, aldehydes, alkanes and alkenes, for industrial applications contain 8 to 14 carbon atoms. In certain embodiments, the above fatty acids, as well as the other corresponding hydrocarbon molecules, are saturated (with no carbon-carbon double or triple bonds); mono unsaturated (single double bond); poly unsturated (two or more double bonds); are linear (not cyclic) or branched. For fuel production, greater saturation is preferred.

[0185] The enzymes described directly above have a preferential specificity for hydrolysis of a substrate containing a specific number of carbon atoms. For example, a fatty acyl-ACP thioesterase may have a preference for cleaving a fatty acid having 12 carbon atoms from the ACP. In some embodiments, the ACP and the length-specific thioesterase may have an affinity for one another that makes them particularly useful as a combination (e.g., the exogenous ACP and thioesterase genes may be naturally co-expressed in a particular tissue or organism from which they are derived). Therefore, in various embodiments, the recombinant *Prototheca* cell of the invention can contain an exogenous gene that encodes a protein with specificity for catalyzing an enzymatic activity (e.g., cleavage of a fatty acid from an ACP, reduction of an acyl-CoA to an aldehyde or an alcohol, or conversion of an aldehyde to an alkane) with regard to the number of carbon atoms contained in the substrate. The enzymatic specificity can, in various embodiments, be for a substrate having from 8 to 34 carbon atoms, preferably from 8 to 18 carbon atoms, and more preferably from 8 to 14 carbon atoms. A preferred specificity is for a substrate having fewer, i.e., 12, rather than more, i.e., 18, carbon atoms.

[0186] In non-limiting but illustrative examples, the present invention provides vectors and Prototheca host cells that express an exogenous thioesterase and accordingly produce lipid enriched, relative to the lipid profile of untransformed *Prototheca* cells, in the chain length for which the thioesterase is specific. The thioesterases illustrated are (i) *Cinnamomum camphorum* FatB1 (GenBank Accension No. Q39473, amino acid sequence is in SEQ ID NO: 59, amino acid sequence without plastid targeting sequence (PTS) is in SEQ ID NO: 139, and codon optimized cDNA sequence based on Table 1 is in SEQ ID NO: 60), which has a preference for fatty acyl-ACP substrate with a carbon chain length of 14; (ii) *Cuphea hookeriana* FatB2 (GenBank Accension No. AAC49269, amino acid sequence is in SEQ ID NO: 61, amino acid sequence without PTS is in SEQ ID NO: 138, and codon optimized cDNA sequence based on Table 1 is in SEQ ID NO: 62), which has a preference for a fatty acyl-ACP substrate with a carbon chain length of 8-10; and (iii) *Umbellularia* Fat B1 (GenBank Accesion No. Q41635, amino acid sequence is included in SEQ ID NO: 63, amino

acid sequence without PTS is in SEQ ID NO: 139, and codon optimized cDNA sequence based on Table 1 is included in SEQ ID NO: 64), which has a preference for a fatty acyl-ACP substrate with a carbon chain length of 12.

[0187] Other fatty acyl-ACP thioesterases suitable for use with the microbes and methods of the invention include, without limitation, those listed in Table 3.

[0188] Table 3. Fatty acyl-ACP thioesterases and GenBank accession numbers.

Umbellularia californica fatty acyl-ACP thioesterase (GenBank #AAC49001)
Cinnamomum camphora fatty acyl-ACP thioesterase (GenBank #Q39473)
Umbellularia californica fatty acyl-ACP thioesterase (GenBank #Q41635)
Myristica fragrans fatty acyl-ACP thioesterase (GenBank #AAB71729)
Myristica fragrans fatty acyl-ACP thioesterase (GenBank #AAB71730)
Elaeis guineensis fatty acyl-ACP thioesterase (GenBank #ABD83939)
Elaeis guineensis fatty acyl-ACP thioesterase (GenBank #AAD42220)
Populus tomentosa fatty acyl-ACP thioesterase (GenBank #ABC47311)
Arabidopsis thaliana fatty acyl-ACP thioesterase (GenBank #NP_172327)
Arabidopsis thaliana fatty acyl-ACP thioesterase (GenBank #CAA85387)
Arabidopsis thaliana fatty acyl-ACP thioesterase (GenBank #CAA85388)
Gossypium hirsutum fatty acyl-ACP thioesterase (GenBank #Q9SQI3)
Cuphea lanceolata fatty acyl-ACP thioesterase (GenBank #CAA54060)
Cuphea hookeriana fatty acyl-ACP thioesterase (GenBank #AAC72882)
Cuphea calophylla subsp. mesostemon fatty acyl-ACP thioesterase (GenBank #ABB71581)
Cuphea lanceolata fatty acyl-ACP thioesterase (GenBank #CAC19933)
Elaeis guineensis fatty acyl-ACP thioesterase (GenBank #AAL15645)
Cuphea hookeriana fatty acyl-ACP thioesterase (GenBank #Q39513)
Gossypium hirsutum fatty acyl-ACP thioesterase (GenBank #AAD01982)
Vitis vinifera fatty acyl-ACP thioesterase (GenBank #CAN81819)
Garcinia mangostana fatty acyl-ACP thioesterase (GenBank #AAB51525)
Brassica juncea fatty acyl-ACP thioesterase (GenBank #ABI18986)
Madhuca longifolia fatty acyl-ACP thioesterase (GenBank #AAX51637)
Brassica napus fatty acyl-ACP thioesterase (GenBank #ABH11710)
Oryza sativa (indica cultivar-group) fatty acyl-ACP thioesterase (GenBank #EAY86877)
Oryza sativa (japonica cultivar-group) fatty acyl-ACP thioesterase (GenBank #NP_001068400)
Oryza sativa (indica cultivar-group) fatty acyl-ACP thioesterase (GenBank #EAY99617)
Cuphea hookeriana fatty acyl-ACP thioesterase (GenBank #AAC49269)
Ulmus Americana fatty acyl-ACP thioesterase (GenBank #AAB71731)
Cuphea lanceolata fatty acyl-ACP thioesterase (GenBank #CAB60830)
Cuphea palustris fatty acyl-ACP thioesterase (GenBank #AAC49180)
Iris germanica fatty acyl-ACP thioesterase (GenBank #AAG43858)
Cuphea palustris fatty acyl-ACP thioesterase (GenBank #AAC49179)
Myristica fragrans fatty acyl-ACP thioesterase (GenBank# AAB71729)
Cuphea hookeriana fatty acyl-ACP thioesterase (GenBank #U39834)
Umbelluaria californica fatty acyl-ACP thioesterase (GenBank # M94159)
Cinnamomum camphora fatty acyl-ACP thioesterase (GenBank #U31813)

[0189] The Examples below describe the successful targeting and expression of heterologous fatty acyl-ACP thioesterases from *Cuphea hookeriana, Umbellularia californica, Cinnamomun camphora* in *Prototheca* species. Additionally, alterations in fatty acid profiles were confirmed in the host cells expression these heterologous fatty acyl-ACP thioesterases. These results were quite unexpected given the lack of sequence identity between algal and higher plant thioesterases in general, and between *Prototheca moriformis* fatty acyl-ACP thioesterase and the above listed heterologous fatty acyl-ACP thioesterases. Two *Prototheca moriformis* acyl-ACP thioesterases were isolated and sequenced. The sequences of the two cDNAs showed a high degree of identity between each other, differing in only 12 positions at the nucleotide level and five positions at the amino acid level, four of these in the plastid transit peptide. Further analysis of genomic sequence from *Prototheca moriformis* confirmed that these two cDNAs were indeed encoded on separate contigs, and although highly homolous, are encoded by two distinct genes. The cDNA and amino acid sequence of the two *Prototheca moriformis* fatty acyl-ACP thioesterase, *P.moriformis* fatty acyl-ACP thioesterase-1 and *P.moriformis* fatty acyl-ACP thioesterase-2, are listed as SEQ ID NOs: 134-137.

[0190] When the amino acid sequences of these two cDNAs were BLASTed against the NCBI database, the two most homologous sequences were fatty acyl-ACP thioesterases from *Chlamydomonas reinhardtii* and *Arabidopsis thaliana*. Surprisingly, the level of amino acid identity between the *Prototheca moriformis* fatty acyl-ACP thioesterases and higher plant thioesterases was fairly low, at only 49 and 37% identity. In addition, there also is a subtle difference in the sequences surrounding the amino terminal portion of the catalytic triad ($NXHX_{36}C$) among these fatty acyl-ACP thioesterases. Thirty nine of forty higher plant fatty acyl-ACP thioesterases surveyed showed the sequence LDMNQH surrounding the N and H residues at the amino terminus of the triad, while all of the algal sequences identified had the sequence MDMNGH. Given the low amino acid sequence identity and the differences surrounding the catalytic triad of the thioesterases, the successful results of expression of exogenous fatty acyl-ACP thioesterases obtained and described in the Examples were unexpected, particularly given the fact that activity of the exogenous fatty acyl-ACP thioesterases was dependent on a functional protein-protein interaction with the endogenous *Prototheca* acyl carrier protein.

[0191] Fatty acyl-CoA/aldehyde reductases suitable for use with the microbes and methods of the invention include, without limitation, those listed in Table 4.

[0192] Table4. Fatty acyl-CoA/aldehyde reductases listed by GenBank accession numbers.

AAC45217, YP_047869, BAB85476, YP_001086217, YP_580344, YP_001280274, YP_264583, YP_436109, YP_959769, ZP_01736962, ZP_01900335, ZP_01892096, ZP_01103974, ZP_01915077, YP_924106, YP_130411, ZP_01222731, YP_550815, YP_983712, YP_001019688, YP_524762, YP_856798, ZP_01115500, YP_001141848, NP_336047, NP_216059, YP_882409, YP_706156, YP_001136150, YP_952365, ZP_01221833, YP_130076, NP_567936, AAR88762, ABK28586, NP_197634, CAD30694, NP_001063962, BAD46254, NP_001030809, EAZ10132, EAZ43639, EAZ07989, NP_001062488, CAB88537, NP_001052541, CAH66597, CAE02214, CAH66590, CAB88538, EAZ39844, AAZ06658, CAA68190, CAA52019, and BAC84377

[0193] Fatty acyl-CoA reductases suitable for use with the microbes and methods of the invention include, without limitation, those listed in Table 5.

[0194] Table 5. Fatty acyl-CoA reductases listed by GenBank accession numbers.

NP_187805, ABO14927, NP_001049083, CAN83375, NP_191229, EAZ42242, EAZ06453, CAD30696, BAD31814, NP_190040, AAD38039, CAD30692, CAN81280, NP_197642, NP_190041, AAL15288, and NP_190042

[0195] Fatty aldehyde decarbonylases suitable for use with the microbes and methods of the invention include, without limitation, those listed in Table 6.

[0196] Table 6. Fatty aldehyde decarbonylases listed by GenBank accession numbers.

NP_850932, ABN07985, CAN60676, AAC23640, CAA65199, AAC24373, CAE03390, ABD28319, NP_181306, EAZ31322, CAN63491, EAY94825, EAY86731, CAL55686, XP_001420263, EAZ23849, NP_200588, NP_001063227, CAN83072, AAR90847, and AAR97643

[0197] Combinations of naturally co-expressed fatty acyl-ACP thioesterases and acyl carrier proteins are suitable for use with the microbes and methods of the invention.

[0198] Additional examples of hydrocarbon or lipid modification enzymes include amino acid sequences contained in, referenced in, or encoded by nucleic acid sequences contained or referenced in, any of the following US patents: 6,610,527; 6,451,576; 6,429,014; 6,342,380; 6,265,639; 6,194,185; 6,114,160; 6,083,731; 6,043,072 ; 5,994,114; 5,891,697; 5,871,988; 6,265,639, and further described in GenBank Accession numbers: AAO18435;

ZP_00513891; Q38710; AAK60613; AAK60610; AAK60611; NP_113747; CAB75874; AAK60612; AAF20201; BAA11024; AF205791; and CAA03710.

[0199] Other suitable enzymes for use with the microbes and the methods of the invention include those that have at least 70% amino acid identity with one of the proteins listed in Tables 3-6, and that exhibit the corresponding desired enzymatic activity (e.g., cleavage of a fatty acid from an acyl carrier protein, reduction of an acyl-CoA to an aldehyde or an alcohol, or conversion of an aldehyde to an alkane). In additional embodiments, the enzymatic activity is present in a sequence that has at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 95%, or at least about 99% identity with one of the above described sequences, all of which are hereby incorporated by reference as if fully set forth.

[0200] By selecting the desired combination of exogenous genes to be expressed, one can tailor the product generated by the microbe, which may then be extracted from the aqueous biomass. For example, the microbe can contain: (i) an exogenous gene encoding a fatty acyl-ACP thioesterase; and, optionally, (ii) a naturally co-expressed acyl carrier protein or an acyl carrier protein otherwise having affinity for the fatty acyl-ACP thioesterase (or conversely); and, optionally, (iii) an exogenous gene encoding a fatty acyl-CoA/aldehyde reductase or a fatty acyl-CoA reductase; and, optionally, (iv) an exogenous gene encoding a fatty aldehyde reductase or a fatty aldehyde decarbonylase. The microbe, under culture conditions described herein, synthesizes a fatty acid linked to an ACP and the fatty acyl-ACP thioesterase catalyzes the cleavage of the fatty acid from the ACP to yield, through further enzymatic processing, a fatty acyl-CoA molecule. When present, the fatty acyl-CoA/aldehyde reducatase catalyzes the reduction of the acyl-CoA to an alcohol. Similarly, the fatty acyl-CoA reductase, when present, catalyzes the reduction of the acyl-CoA to an aldehyde. In those embodiments in which an exogenous gene encoding a fatty acyl-CoA reductase is present and expressed to yield an aldehyde product, a fatty aldehyde reductase, encoded by the third exogenous gene, catalyzes the reduction of the aldehyde to an alcohol. Similarly, a fatty aldehyde decarbonylase catalyzes the conversion of the aldehyde to an alkane or an alkene, when present.

[0201] Genes encoding such enzymes can be obtained from cells already known to exhibit significant lipid production such as *Chlorella protothecoides*. Genes already known to have a role in lipid production, e.g., a gene encoding an enzyme that saturates double bonds, can be transformed individually into recipient cells. However, to practice the invention it is not necessary to make *a priori* assumptions as to which genes are required. Methods for

identifiying genes that can alter (improve) lipid production in microalgae are described in PCT Pub. No. 2008/151149.

[0202] Thus, the present invention provides a *Prototheca* cell that has been genetically engineered to express a lipid pathway enzyme at an altered level compared to a wild-type cell of the same species. In some cases, the cell produces more lipid compared to the wild-type cell when both cells are grown under the same conditions. In some cases, the cell has been genetically engineered and/or selected to express a lipid pathway enzyme at a higher level than the wild-type cell. In some cases, the lipid pathway enzyme is selected from the group consisting of pyruvate dehydrogenase, acetyl-CoA carboxylase, acyl carrier protein, and glycerol-3 phosphate acyltransferase. In some cases, the cell has been genetically engineered and/or selected to express a lipid pathway enzyme at a lower level than the wild-type cell. In at least one embodiment in which the cell expresses the lipid pathway enzyme at a lower level, the lipid pathway enzyme comprises citrate synthase.

[0203] In some embodiments, the cell has been genetically engineered and/or selected to express a global regulator of fatty acid synthesis at an altered level compared to the wild-type cell, whereby the expression levels of a plurality of fatty acid synthetic genes are altered compared to the wild-type cell. In some cases, the lipid pathway enzyme comprises an enzyme that modifies a fatty acid. In some cases, the lipid pathway enzyme is selected from a stearoyl-ACP desaturase and a glycerolipid desaturase.

[0204] In other embodiments, the present invention is directed to an oil-producing microbe containing one or more exogenous genes, wherein the exogenous genes encode protein(s) selected from the group consisting of a fatty acyl-ACP thioesterase, a fatty acyl-CoA reductase, a fatty aldehyde reductase, a fatty acyl-CoA/aldehyde reductase, a fatty aldehyde decarbonylase, and an acyl carrier protein. In one embodiment, the exogenous gene is in operable linkage with a promoter, which is inducible or repressible in response to a stimulus. In some cases, the stimulus is selected from the group consisting of an exogenously provided small molecule, heat, cold, and limited or no nitrogen in the culture media. In some cases, the exogenous gene is expressed in a cellular compartment. In some embodiments, the cellular compartment is selected from the group consisting of a chloroplast, a plastid and a mitochondrion. In some embodiments the microbe is *Prototheca moriformis, Prototheca krugani, Prototheca stagnora* or *Prototheca zopfii.*

[0205] In one embodiment, the exogenous gene encodes a fatty acid acyl-ACP thioesterase. In some cases, the thioesterase encoded by the exogenous gene catalyzes the cleavage of an 8 to 18-carbon fatty acid from an acyl carrier protein (ACP). In some cases, the thioesterase

encoded by the exogenous gene catalyzes the cleavage of a 10 to 14-carbon fatty acid from an ACP. In one embodiment, the thioesterase encoded by the exogenous gene catalyzes the cleavage of a 12-carbon fatty acid from an ACP.

[0206] In one embodiment, the exogenous gene encodes a fatty acyl-CoA/aldehyde reductase. In some cases, the reductase encoded by the exogenous gene catalyzes the reduction of an 8 to 18-carbon fatty acyl-CoA to a corresponding primary alcohol. In some cases, the reductase encoded by the exogenous gene catalyzes the reduction of a 10 to 14-carbon fatty acyl-CoA to a corresponding primary alcohol. In one embodiment, the reductase encoded by the exogenous gene catalyzes the reduction of a 12-carbon fatty acyl-CoA to dodecanol.

[0207] The present invention also provides a recombinant *Prototheca* cell containing two exogenous genes, wherein a first exogenous gene encodes a fatty acyl-ACP thioesterase and a second exogenous gene encodes a protein selected from the group consisting of a fatty acyl-CoA reductase, a fatty acyl-CoA/aldehyde reductase, and an acyl carrier protein. In some cases, the two exogenous genes are each in operable linkage with a promoter, which is inducible in response to a stimulus. In some cases, each promoter is inducible in response to an identical stimulus, such as limited or no nitrogen in the culture media. Limitation or complete lack of nitrogen in the culture media stimulates oil production in some microorganisms such as Prototheca species, and can be used as a trigger to inducec oil production to high levels. When used in combination with the genetic engineering methods disclosed herein, the lipid as a percentage of dry cell weight can be pushed to high levels such as at least 30%, at least 40%, at least 50%, at least 60%, at least 70% and at least 75%; methods disclosed herein provide for cells with these levels of lipid, wherein the lipid is at least 4% C8-C14, at least 0.3% C8, at least 2% C10, at least 2% C12, and at least 2% C14. In some embodiments the cells are over 25% lipid by dry cell weight and contain lipid that is at least 10% C8-C14, at least 20% C8-C14, at least 30% C8-C14, 10-30% C8-C14 and 20-30% C8-C14.

[0208] The novel oils disclosed herein are distinct from other naturally occurring oils that are high in mic-chain fatty acids, such as palm oil, palm kernel oil, and coconut oil. For example, levels of contaminants such as carotenoids are far higher in palm oil and palm kernel oil than in the oils of the invention. Palm and palm kernel oils in particular contain alpha and beta carotenes and lycopene in much higher amounts than is in the oils of the invention. In addition, over 20 different carotenoids are found in palm and palm kernel oil, whereas the Examples demonstrate that the oils of the invention contain very few carotenoids

species and very low levels. In addition, the levels of vitamin E compounds such as tocotrienols are far higher in palm, palm kernel, and coconut oil than in the oils of the invention.

[0209] In one embodiment, the thioesterase encoded by the first exogenous gene catalyzes the cleavage of an 8 to 18-carbon fatty acid from an ACP. In some embodiments, the second exogenous gene encodes a fatty acyl-CoA/aldehyde reductase which catalyzes the reduction of an 8 to 18-carbon fatty acyl-CoA to a corresponding primary alcohol. In some cases, the thioesterase encoded by the first exogenous gene catalyzes the cleavage of a 10 to 14-carbon fatty acid from an ACP, and the reductase encoded by the second exogenous gene catalyzes the reduction of a 10 to 14-carbon fatty acyl-CoA to the corresponding primary alcohol, wherein the thioesterase and the reductase act on the same carbon chain length. In one embodiment, the thioesterase encoded by the first exogenous gene catalyzes the cleavage of a 12-carbon fatty acid from an ACP, and the reductase encoded by the second exogenous gene catalyzes the reduction of a 12-carbon fatty acyl-CoA to dodecanol. In some embodiments, the second exogenous gene encodes a fatty acyl-CoA reductase which catalyzes the reduction of an 8 to 18-carbon fatty acyl-CoA to a corresponding aldehyde. In some embodiments, the second exogenous gene encodes an acyl carrier protein that is naturally co-expressed with the fatty acyl-ACP thioesterase.

[0210] In some embodiments, the second exogenous gene encodes a fatty acyl-CoA reductase, and the microbe further contains a third exogenous gene encoding a fatty aldehyde decarbonylase. In some cases, the thioesterase encoded by the first exogenous gene catalyzes the cleavage of an 8 to 18-carbon fatty acid from an ACP, the reductase encoded by the second exogenous gene catalyzes the reduction of an 8 to 18-carbon fatty acyl-CoA to a corresponding fatty aldehyde, and the decarbonylase encoded by the third exogenous gene catalyzes the conversion of an 8 to 18-carbon fatty aldehyde to a corresponding alkane, wherein the thioesterase, the reductase, and the decarbonylase act on the same carbon chain length.

[0211] In some embodiments, the second exogenous gene encodes an acyl carrier protein, and the microbe further contains a third exogenous gene encoding a protein selected from the group consisting of a fatty acyl-CoA reductase and a fatty acyl-CoA/aldehyde reductase. In some cases, the third exogenous gene encodes a fatty acyl-CoA reductase, and the microbe further contains a fourth exogenous gene encoding a fatty aldehyde decarbonylase.

[0212] The present invention also provides methods for producing an alcohol comprising culturing a population of recombinant *Prototheca* cells in a culture medium, wherein the cells

contain (i) a first exogenous gene encoding a fatty acyl-ACP thioesterase, and (ii) a second exogenous gene encoding a fatty acyl-CoA/aldehyde reductase, and the cells synthesize a fatty acid linked to an acyl carrier protein (ACP), the fatty acyl-ACP thioesterase catalyzes the cleavage of the fatty acid from the ACP to yield, through further processing, a fatty acyl-CoA, and the fatty acyl-CoA/aldehyde reductase catalyzes the reduction of the acyl-CoA to an alcohol.

[0213] The present invention also provides methods of producing a lipid molecule in a *Prototheca* cell. In one embodiment, the method comprises culturing a population of *Prototheca* cells in a culture medium, wherein the cells contain (i) a first exogenous gene encoding a fatty acyl-ACP thioesterase, and (ii) a second exogenous gene encoding a fatty acyl-CoA reductase, and wherein the microbes synthesize a fatty acid linked to an acyl carrier protein (ACP), the fatty acyl-ACP thioesterase catalyzes the cleavage of the fatty acid from the ACP to yield, through further processing, a fatty acyl-CoA, and the fatty acyl-CoA reductase catalyzes the reduction of the acyl-CoA to an aldehyde.

[0214] The present invention also provides methods of producing a fatty acid molecule having a specified carbon chain length in a *Prototheca* cell. In one embodiment, the method comprises culturing a population of lipid-producing *Prototheca* cells in a culture medium, wherein the microbes contain an exogenous gene encoding a fatty acyl-ACP thioesterase having an activity specific or preferential to a certain carbon chain length, such as 8, 10, 12 or 14 carbon atoms, and wherein the microbes synthesize a fatty acid linked to an acyl carrier protein (ACP) and the thioesterase catalyzes the cleavage of the fatty acid from the ACP when the fatty acid has been synthesized to the specific carbon chain length.

[0215] In the various embodiments described above, the *Prototheca* cell can contain at least one exogenous gene encoding a lipid pathway enzyme. In some cases, the lipid pathway enzyme is selected from the group consisting of a stearoyl-ACP desaturase, a glycerolipid desaturase, a pyruvate dehydrogenase, an acetyl-CoA carboxylase, an acyl carrier protein, and a glycerol-3 phosphate acyltransferase. In other cases, the *Prototheca* cell contains a lipid modification enzyme selected from the group consisting of a fatty acyl-ACP thioesterase, a fatty acyl-CoA/aldehyde reductase, a fatty acyl-CoA reductase, a fatty aldehyde reductase, a fatty aldehyde decarbonylase, and/or an acyl carrier protein.

VI. FUELS AND CHEMICALS PRODUCTION

[0216] For the production of fuel in accordance with the methods of the invention lipids produced by cells of the invention are harvested, or otherwise collected, by any convenient means. Lipids can be isolated by whole cell extraction. The cells are first disrupted, and then

intracellular and cell membrane/cell wall-associated lipids as well as extracellular hydrocarbons can be separated from the cell mass, such as by use of centrifugation as described above. Intracellular lipids produced in microorganisms are, in some embodiments, extracted after lysing the cells of the microorganism. Once extracted, the lipids are further refined to produce oils, fuels, or oleochemicals.

[0217] After completion of culturing, the microorganisms can be separated from the fermentation broth. Optionally, the separation is effected by centrifugation to generate a concentrated paste. Centrifugation does not remove significant amounts of intracellular water from the microorganisms and is not a drying step. The biomass can then optionally be washed with a washing solution (e.g., DI water) to get rid of the fermentation broth and debris. Optionally, the washed microbial biomass may also be dried (oven dried, lyophilized, etc.) prior to cell disruption. Alternatively, cells can be lysed without separation from some or all of the fermentation broth when the fermentation is complete. For example, the cells can be at a ratio of less than 1:1 v:v cells to extracellular liquid when the cells are lysed.

[0218] Microorganisms containing a lipid can be lysed to produce a lysate. As detailed herein, the step of lysing a microorganism (also referred to as cell lysis) can be achieved by any convenient means, including heat-induced lysis, adding a base, adding an acid, using enzymes such as proteases and polysaccharide degradation enzymes such as amylases, using ultrasound, mechanical lysis, using osmotic shock, infection with a lytic virus, and/or expression of one or more lytic genes. Lysis is performed to release intracellular molecules which have been produced by the microorganism. Each of these methods for lysing a microorganism can be used as a single method or in combination simultaneously or sequentially. The extent of cell disruption can be observed by microscopic analysis. Using one or more of the methods described herein, typically more than 70% cell breakage is observed. Preferably, cell breakage is more than 80%, more preferably more than 90% and most preferred about 100%.

[0219] In particular embodiments, the microorganism is lysed after growth, for example to increase the exposure of cellular lipid and/or hydrocarbon for extraction or further processing. The timing of lipase expression (e.g., via an inducible promoter) or cell lysis can be adjusted to optimize the yield of lipids and/or hydrocarbons. Below are described a number of lysis techniques. These techniques can be used individually or in combination.

[0220] In one embodiment of the present invention, the step of lysing a microorganism comprises heating of a cellular suspension containing the microorganism. In this embodiment, the fermentation broth containing the microorganisms (or a suspension of

microorganisms isolated from the fermentation broth) is heated until the microorganisms, i.e., the cell walls and membranes of microorganisms degrade or breakdown. Typically, temperatures applied are at least 50°C. Higher temperatures, such as, at least 30°C at least 60°C, at least 70°C, at least 80°C, at least 90°C, at least 100°C, at least 110°C, at least 120°C, at least 130°C or higher are used for more efficient cell lysis. Lysing cells by heat treatment can be performed by boiling the microorganism. Alternatively, heat treatment (without boiling) can be performed in an autoclave. The heat treated lysate may be cooled for further treatment. Cell disruption can also be performed by steam treatment, i.e., through addition of pressurized steam. Steam treatment of microalgae for cell disruption is described, for example, in U.S. Patent No. 6,750,048. In some embodiments, steam treatment may be achieved by sparging steam into the fermentor and maintaining the broth at a desired temperature for less than about 90 minutes, preferably less than about 60 minutes, and more preferably less than about 30 minutes.

[0221] In another embodiment of the present invention, the step of lysing a microorganism comprises adding a base to a cellular suspension containing the microorganism. The base should be strong enough to hydrolyze at least a portion of the proteinaceous compounds of the microorganisms used. Bases which are useful for solubilizing proteins are known in the art of chemistry. Exemplary bases which are useful in the methods of the present invention include, but are not limited to, hydroxides, carbonates and bicarbonates of lithium, sodium, potassium, calcium, and mixtures thereof. A preferred base is KOH. Base treatment of microalgae for cell disruption is described, for example, in U.S. Patent No. 6,750,048.

[0222] In another embodiment of the present invention, the step of lysing a microorganism comprises adding an acid to a cellular suspension containing the microorganism. Acid lysis can be effected using an acid at a concentration of 10-500 mN or preferably 40-160 nM. Acid lysis is preferably performed at above room temperature (e.g., at 40-160°, and preferably a temperature of 50-130°. For moderate temperatures (e.g., room temperature to 100°C and particularly room temperature to 65°, acid treatment can usefully be combined with sonication or other cell disruption methods.

[0223] In another embodiment of the present invention, the step of lysing a microorganism comprises lysing the microorganism by using an enzyme. Preferred enzymes for lysing a microorganism are proteases and polysaccharide-degrading enzymes such as hemicellulase (e.g., hemicellulase from *Aspergillus niger*; Sigma Aldrich, St. Louis, MO; #H2125), pectinase (e.g., pectinase from *Rhizopus sp.*; Sigma Aldrich, St. Louis, MO; #P2401), Mannaway 4.0 L (Novozymes), cellulase (e.g., cellulose from Trichoderma viride; Sigma

Aldrich, St. Louis, MO; #C9422), and driselase (e.g., driselase from *Basidiomycetes sp.*; Sigma Aldrich, St. Louis, MO; #D9515.

[0224] In other embodiments of the present invention, lysis is accomplished using an enzyme such as, for example, a cellulase such as a polysaccharide-degrading enzyme, optionally from *Chlorella* or a *Chlorella* virus, or a proteases, such as *Streptomyces griseus* protease, chymotrypsin, proteinase K, proteases listed in Degradation of Polylactide by Commercial Proteases, Oda Yet al., Journal of Polymers and the Environment, Volume 8, Number 1, January 2000 , pp. 29-32(4), Alcalase 2.4 FG (Novozymes), and Flavourzyme 100 L (Novozymes). Any combination of a protease and a polysaccharide-degrading enzyme can also be used, including any combination of the preceding proteases and polysaccharide-degrading enzymes.

[0225] In another embodiment, lysis can be performed using an expeller press. In this process, biomass is forced through a screw-type device at high pressure, lysing the cells and causing the intracellular lipid to be released and separated from the protein and fiber (and other components) in the cell.

[0226] In another embodiment of the present invention, the step of lysing a microorganism is performed by using ultrasound, i.e., sonication. Thus, cells can also by lysed with high frequency sound. The sound can be produced electronically and transported through a metallic tip to an appropriately concentrated cellular suspension. This sonication (or ultrasonication) disrupts cellular integrity based on the creation of cavities in cell suspension.

[0227] In another embodiment of the present invention, the step of lysing a microorganism is performed by mechanical lysis. Cells can be lysed mechanically and optionally homogenized to facilitate hydrocarbon (e.g., lipid) collection. For example, a pressure disrupter can be used to pump a cell containing slurry through a restricted orifice valve. High pressure (up to 1500 bar) is applied, followed by an instant expansion through an exiting nozzle. Cell disruption is accomplished by three different mechanisms: impingement on the valve, high liquid shear in the orifice, and sudden pressure drop upon discharge, causing an explosion of the cell. The method releases intracellular molecules. Alternatively, a ball mill can be used. In a ball mill, cells are agitated in suspension with small abrasive particles, such as beads. Cells break because of shear forces, grinding between beads, and collisions with beads. The beads disrupt the cells to release cellular contents. Cells can also be disrupted by shear forces, such as with the use of blending (such as with a high speed or Waring blender as examples), the french press, or even centrifugation in case of weak cell walls, to disrupt cells.

[0228] In another embodiment of the present invention, the step of lysing a microorganism is performed by applying an osmotic shock.

[0229] In another embodiment of the present invention, the step of lysing a microorganism comprises infection of the microorganism with a lytic virus. A wide variety of viruses are known to lyse microorganisms suitable for use in the present invention, and the selection and use of a particular lytic virus for a particular microorganism is within the level of skill in the art. For example, *paramecium bursaria chlorella* virus (PBCV-1) is the prototype of a group (family Phycodnaviridae, genus Chlorovirus) of large, icosahedral, plaque-forming, double-stranded DNA viruses that replicate in, and lyse, certain unicellular, eukaryotic chlorella-like green algae. Accordingly, any susceptible microalgae can be lysed by infecting the culture with a suitable chlorella virus. Methods of infecting species of *Chlorella* with a chlorella virus are known. See for example *Adv. Virus Res.* 2006;66:293-336; *Virology*, 1999 Apr 25;257(1):15-23; *Virology*, 2004 Jan 5;318(1):214-23; *Nucleic Acids Symp. Ser.* 2000;(44):161-2; *J. Virol.* 2006 Mar;80(5):2437-44; and *Annu. Rev. Microbiol.* 1999;53:447-94.

[0230] In another embodiment of the present invention, the step of lysing a microorganism comprises autolysis. In this embodiment, a microorganism according to the invention is genetically engineered to produce a lytic protein that will lyse the microorganism. This lytic gene can be expressed using an inducible promoter so that the cells can first be grown to a desirable density in a fermentor, followed by induction of the promoter to express the lytic gene to lyse the cells. In one embodiment, the lytic gene encodes a polysaccharide-degrading enzyme. In certain other embodiments, the lytic gene is a gene from a lytic virus. Thus, for example, a lytic gene from a Chlorella virus can be expressed in an algal cell; see *Virology* 260, 308-315 (1999); *FEMS Microbiology Letters* 180 (1999) 45-53; *Virology* 263, 376-387 (1999); and *Virology* 230, 361–368 (1997). Expression of lytic genes is preferably done using an inducible promoter, such as a promoter active in microalgae that is induced by a stimulus such as the presence of a small molecule, light, heat, and other stimuli.

[0231] Various methods are available for separating lipids from cellular lysates produced by the above methods. For example, lipids and lipid derivatives such as fatty aldehydes, fatty alcohols, and hydrocarbons such as alkanes can be extracted with a hydrophobic solvent such as hexane (see Frenz et al. 1989, Enzyme Microb. Technol., 11:717). Lipids and lipid derivatives can also be extracted using liquefaction (see for example Sawayama et al. 1999, Biomass and Bioenergy 17:33-39 and Inoue et al. 1993, Biomass Bioenergy 6(4):269-274); oil liquefaction (see for example Minowa et al. 1995, Fuel 74(12):1735-1738); and

supercritical CO_2 extraction (see for example Mendes et al. 2003, Inorganica Chimica Acta 356:328-334). Miao and Wu describe a protocol of the recovery of microalgal lipid from a culture of *Chlorella prototheocoides* in which the cells were harvested by centrifugation, washed with distilled water and dried by freeze drying. The resulting cell powder was pulverized in a mortar and then extracted with *n*-hexane. Miao and Wu, Biosource Technology (2006) 97:841-846.

[0232] Thus, lipids, lipid derivatives and hydrocarbons generated by the microorganisms of the present invention can be recovered by extraction with an organic solvent. In some cases, the preferred organic solvent is hexane. Typically, the organic solvent is added directly to the lysate without prior separation of the lysate components. In one embodiment, the lysate generated by one or more of the methods described above is contacted with an organic solvent for a period of time sufficient to allow the lipid and/or hydrocarbon components to form a solution with the organic solvent. In some cases, the solution can then be further refined to recover specific desired lipid or hydrocarbon components. Hexane extraction methods are well known in the art.

[0233] Lipids and lipid derivatives such as fatty aldehydes, fatty alcohols, and hydrocarbons such as alkanes produced by cells as described herein can be modified by the use of one or more enzymes, including a lipase, as described above. When the hydrocarbons are in the extracellular environment of the cells, the one or more enzymes can be added to that environment under conditions in which the enzyme modifies the hydrocarbon or completes its synthesis from a hydrocarbon precursor. Alternatively, the hydrocarbons can be partially, or completely, isolated from the cellular material before addition of one or more catalysts such as enzymes. Such catalysts are exogenously added, and their activity occurs outside the cell or *in vitro*.

[0234] Thus, lipids and hydrocarbons produced by cells *in vivo*, or enzymatically modified *in vitro*, as described herein can be optionally further processed by conventional means. The processing can include "cracking" to reduce the size, and thus increase the hydrogen:carbon ratio, of hydrocarbon molecules. Catalytic and thermal cracking methods are routinely used in hydrocarbon and triglyceride oil processing. Catalytic methods involve the use of a catalyst, such as a solid acid catalyst. The catalyst can be silica-alumina or a zeolite, which result in the heterolytic, or asymmetric, breakage of a carbon-carbon bond to result in a carbocation and a hydride anion. These reactive intermediates then undergo either rearrangement or hydride transfer with another hydrocarbon. The reactions can thus regenerate the intermediates to result in a self-propagating chain mechanism. Hydrocarbons

can also be processed to reduce, optionally to zero, the number of carbon-carbon double, or triple, bonds therein. Hydrocarbons can also be processed to remove or eliminate a ring or cyclic structure therein. Hydrocarbons can also be processed to increase the hydrogen:carbon ratio. This can include the addition of hydrogen ("hydrogenation") and/or the "cracking" of hydrocarbons into smaller hydrocarbons.

[0235] Thermal methods involve the use of elevated temperature and pressure to reduce hydrocarbon size. An elevated temperature of about 800°C and pressure of about 700kPa can be used. These conditions generate "light," a term that is sometimes used to refer to hydrogen-rich hydrocarbon molecules (as distinguished from photon flux), while also generating, by condensation, heavier hydrocarbon molecules which are relatively depleted of hydrogen. The methodology provides homolytic, or symmetrical, breakage and produces alkenes, which may be optionally enzymatically saturated as described above.

[0236] Catalytic and thermal methods are standard in plants for hydrocarbon processing and oil refining. Thus hydrocarbons produced by cells as described herein can be collected and processed or refined via conventional means. See Hillen et al. (Biotechnology and Bioengineering, Vol. XXIV:193-205 (1982)) for a report on hydrocracking of microalgae-produced hydrocarbons. In alternative embodiments, the fraction is treated with another catalyst, such as an organic compound, heat, and/or an inorganic compound. For processing of lipids into biodiesel, a transesterification process is used as described in Section IV herein.

[0237] Hydrocarbons produced via methods of the present invention are useful in a variety of industrial applications. For example, the production of linear alkylbenzene sulfonate (LAS), an anionic surfactant used in nearly all types of detergents and cleaning preparations, utilizes hydrocarbons generally comprising a chain of 10-14 carbon atoms. See, for example, US Patent Nos.: 6,946,430; 5,506,201; 6,692,730; 6,268,517; 6,020,509; 6,140,302; 5,080,848; and 5,567,359. Surfactants, such as LAS, can be used in the manfacture of personal care compositions and detergents, such as those described in US Patent Nos.: 5,942,479; 6,086,903; 5,833,999; 6,468,955; and 6,407,044.

[0238] Increasing interest is directed to the use of hydrocarbon components of biological origin in fuels, such as biodiesel, renewable diesel, and jet fuel, since renewable biological starting materials that may replace starting materials derived from fossil fuels are available, and the use thereof is desirable. There is an urgent need for methods for producing hydrocarbon components from biological materials. The present invention fulfills this need by providing methods for production of biodiesel, renewable diesel, and jet fuel using the

lipids generated by the methods described herein as a biological material to produce biodiesel, renewable diesel, and jet fuel.

[0239] Traditional diesel fuels are petroleum distillates rich in paraffinic hydrocarbons. They have boiling ranges as broad as 370° to 780°F, which are suitable for combustion in a compression ignition engine, such as a diesel engine vehicle. The American Society of Testing and Materials (ASTM) establishes the grade of diesel according to the boiling range, along with allowable ranges of other fuel properties, such as cetane number, cloud point, flash point, viscosity, aniline point, sulfur content, water content, ash content, copper strip corrosion, and carbon residue. Technically, any hydrocarbon distillate material derived from biomass or otherwise that meets the appropriate ASTM specification can be defined as diesel fuel (ASTM D975), jet fuel (ASTM D1655), or as biodiesel if it is a fatty acid methyl ester (ASTM D6751).

[0240] After extraction, lipid and/or hydrocarbon components recovered from the microbial biomass described herein can be subjected to chemical treatment to manufacture a fuel for use in diesel vehicles and jet engines.

[0241] Biodiesel is a liquid which varies in color - between golden and dark brown - depending on the production feedstock. It is practically immiscible with water, has a high boiling point and low vapor pressure. Biodiesel refers to a diesel-equivalent processed fuel for use in diesel-engine vehicles. Biodiesel is biodegradable and non-toxic. An additional benefit of biodiesel over conventional diesel fuel is lower engine wear. Typically, biodiesel comprises C14-C18 alkyl esters. Various processes convert biomass or a lipid produced and isolated as described herein to diesel fuels. A preferred method to produce biodiesel is by transesterification of a lipid as described herein. A preferred alkyl ester for use as biodiesel is a methyl ester or ethyl ester.

[0242] Biodiesel produced by a method described herein can be used alone or blended with conventional diesel fuel at any concentration in most modern diesel-engine vehicles. When blended with conventional diesel fuel (petroleum diesel), biodiesel may be present from about 0.1% to about 99.9%. Much of the world uses a system known as the "B" factor to state the amount of biodiesel in any fuel mix. For example, fuel containing 20% biodiesel is labeled B20. Pure biodiesel is referred to as B100.

[0243] Biodiesel can also be used as a heating fuel in domestic and commercial boilers. Existing oil boilers may contain rubber parts and may require conversion to run on biodiesel. The conversion process is usually relatively simple, involving the exchange of rubber parts for synthetic parts due to biodiesel being a strong solvent. Due to its strong solvent power,


burning biodiesel will increase the efficiency of boilers. Biodiesel can be used as an additive in formulations of diesel to increase the lubricity of pure Ultra-Low Sulfur Diesel (ULSD) fuel, which is advantageous because it has virtually no sulfur content. Biodiesel is a better solvent than petrodiesel and can be used to break down deposits of residues in the fuel lines of vehicles that have previously been run on petrodiesel.

[0244] Biodiesel can be produced by transesterification of triglycerides contained in oil-rich biomass. Thus, in another aspect of the present invention a method for producing biodiesel is provided. In a preferred embodiment, the method for producing biodiesel comprises the steps of (a) cultivating a lipid-containing microorganism using methods disclosed herein (b) lysing a lipid-containing microorganism to produce a lysate, (c) isolating lipid from the lysed microorganism, and (d) transesterifying the lipid composition, whereby biodiesel is produced. Methods for growth of a microorganism, lysing a microorganism to produce a lysate, treating the lysate in a medium comprising an organic solvent to form a heterogeneous mixture and separating the treated lysate into a lipid composition have been described above and can also be used in the method of producing biodiesel.

[0245] The lipid profile of the biodiesel is usually highly similar to the lipid profile of the feedstock oil. Other oils provided by the methods and compositions of the invention can be subjected to transesterification to yield biodiesel with lipid profiles including (a) at least 4% C8-C14; (b) at least 0.3% C8; (c) at least 2% C10; (d) at least 2% C12; and (3) at least 30% C8-C14.

[0246] Lipid compositions can be subjected to transesterification to yield long-chain fatty acid esters useful as biodiesel. Preferred transesterification reactions are outlined below and include base catalyzed transesterification and transesterification using recombinant lipases. In a base-catalyzed transesterification process, the triacylglycerides are reacted with an alcohol, such as methanol or ethanol, in the presence of an alkaline catalyst, typically potassium hydroxide. This reaction forms methyl or ethyl esters and glycerin (glycerol) as a byproduct.

[0247] Animal and plant oils are typically made of triglycerides which are esters of free fatty acids with the trihydric alcohol, glycerol. In transesterification, the glycerol in a triacylglyceride (TAG) is replaced with a short-chain alcohol such as methanol or ethanol. A typical reaction scheme is as follows:



In this reaction, the alcohol is deprotonated with a base to make it a stronger nucleophile. Commonly, ethanol or methanol is used in vast excess (up to 50-fold). Normally, this reaction will proceed either exceedingly slowly or not at all. Heat, as well as an acid or base can be used to help the reaction proceed more quickly. The acid or base are not consumed by the transesterification reaction, thus they are not reactants but catalysts. Almost all biodiesel has been produced using the base-catalyzed technique as it requires only low temperatures and pressures and produces over 98% conversion yield (provided the starting oil is low in moisture and free fatty acids).

[0248] Transesterification has also been carried out, as discussed above, using an enzyme, such as a lipase instead of a base. Lipase-catalyzed transesterification can be carried out, for example, at a temperature between the room temperature and 80° C, and a mole ratio of the TAG to the lower alcohol of greater than 1:1, preferably about 3:1. Lipases suitable for use in transesterification include, but are not limited to, those listed in Table7. Other examples of lipases useful for transesterification are found in, e.g. U.S. Patent Nos. 4,798,793; 4,940,845 5,156,963; 5,342,768; 5,776,741 and WO89/01032. Such lipases include, but are not limited to, lipases produced by microorganisms of Rhizopus, Aspergillus, Candida, Mucor, Pseudomonas, Rhizomucor, Candida, and Humicola and pancreas lipase.

[0249] Table 7. Lipases suitable for use in transesterification.

Aspergillus niger lipase ABG73614, *Candida antarctica* lipase B (novozym-435) CAA83122, *Candida cylindracea* lipase AAR24090, *Candida lipolytica* lipase (Lipase L; Amano Pharmaceutical Co., Ltd.), *Candida rugosa* lipase (e.g., Lipase-OF; Meito Sangyo Co., Ltd.), *Mucor miehei* lipase (Lipozyme IM 20), *Pseudomonas fluorescens* lipase AAA25882, *Rhizopus japonicas* lipase (Lilipase A-10FG) Q7M4U7_1, *Rhizomucor miehei* lipase B34959, *Rhizopus oryzae lipase* (Lipase F) AAF32408, *Serratia marcescens* lipase (SM Enzyme) ABI13521, *Thermomyces lanuginosa* lipase CAB58509, Lipase P (Nagase ChemteX Corporation), and Lipase QLM (Meito Sangyo Co., Ltd., Nagoya, Japan)

[0250] One challenge to using a lipase for the production of fatty acid esters suitable for biodiesel is that the price of lipase is much higher than the price of sodium hydroxide (NaOH) used by the strong base process. This challenge has been addressed by using an immobilized lipase, which can be recycled. However, the activity of the immobilized lipase must be maintained after being recycled for a minimum number of cycles to allow a lipase-

based process to compete with the strong base process in terms of the production cost. Immobilized lipases are subject to poisoning by the lower alcohols typically used in transesterification. U.S. Patent No. 6,398,707 (issued June 4, 2002 to Wu et al.) describes methods for enhancing the activity of immobilized lipases and regenerating immobilized lipases having reduced activity. Some suitable methods include immersing an immobilized lipase in an alcohol having a carbon atom number not less than 3 for a period of time, preferably from 0.5-48 hours, and more preferably from 0.5-1.5 hours. Some suitable methods also include washing a deactivated immobilized lipase with an alcohol having a carbon atom number not less than 3 and then immersing the deactivated immobilized lipase in a vegetable oil for 0.5-48 hours.

[0251] In particular embodiments, a recombinant lipase is expressed in the same microorganisms that produce the lipid on which the lipase acts. Suitable recombinant lipases include those listed above in Table 7 and/or having GenBank Accession numbers listed above in Table 7, or a polypeptide that has at least 70% amino acid identity with one of the lipases listed above in Table 7 and that exhibits lipase activity. In additional embodiments, the enzymatic activity is present in a sequence that has at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 95%, or at least about 99% identity with one of the above described sequences, all of which are hereby incorporated by reference as if fully set forth. DNA encoding the lipase and selectable marker is preferably codon-optimized cDNA. Methods of recoding genes for expression in microalgae are described in US Patent 7,135,290.

[0252] The common international standard for biodiesel is EN 14214. ASTM D6751 is the most common biodiesel standard referenced in the United States and Canada. Germany uses DIN EN 14214 and the UK requires compliance with BS EN 14214. Basic industrial tests to determine whether the products conform to these standards typically include gas chromatography, HPLC, and others. Biodiesel meeting the quality standards is very non-toxic, with a toxicity rating (LD_{50}) of greater than 50 mL/kg.

[0253] Although biodiesel that meets the ASTM standards has to be non-toxic, there can be contaminants which tend to crystallize and/or precipitate and fall out of solution as sediment. Sediment formation is particularly a problem when biodiesel is used at lower temperatures. The sediment or precipitates may cause problems such as decreasing fuel flow, clogging fuel lines, clogging filters, etc. Processes are well-known in the art that specifically deal with the removal of these contaminants and sediments in biodiesel in order to produce a higher quality product. Examples for such processes include, but are not limited to, pretreatment of the oil to

remove contaiminants such as phospholipids and free fatty acids (*e.g.*, degumming, caustic refining and silica adsorbant filtration) and cold filtration. Cold filtration is a process that was developed specifically to remove any particulates and sediments that are present in the biodiesel after production. This process cools the biodiesel and filters out any sediments or precipitates that might form when the fuel is used at a lower temperature. Such a process is well known in the art and is described in US Patent Application Publication No. 2007-0175091. Suitable methods may include cooling the biodiesel to a temperature of less than about 38°C so that the impurities and contaminants precipitate out as particulates in the biodiesel liquid. Diatomaceous earth or other filtering material may then added to the cooled biodiesel to form a slurry, which may then filtered through a pressure leaf or other type of filter to remove the particulates. The filtered biodiesel may then be run through a polish filter to remove any remaining sediments and diatomaceous earth, so as to produce the final biodiesel product.

[0254] Example 14 described the production of biodiesel using triglyceride oil from *Prototheca moriformis*. The Cold Soak Filterability by the ASTM D6751 A1 method of the biodiesel produced in Example 14 was 120 seconds for a volume of 300ml. This test involves filtration of 300 ml of B100, chilled to 40°F for 16 hours, allowed to warm to room temp, and filtered under vacuum using 0.7 micron glass fiber filter with stainless steel support. Oils of the invention can be transesterified to generate biodiesel with a cold soak time of less than 120 seconds, less than 100 seconds, and less than 90 seconds.

[0255] Subsequent processes may also be used if the biodiesel will be used in particularly cold temperatures. Such processes include winterization and fractionation. Both processes are designed to improve the cold flow and winter performance of the fuel by lowering the cloud point (the temperature at which the biodiesel starts to crystallize). There are several approaches to winterizing biodiesel. One approach is to blend the biodiesel with petroleum diesel. Another approach is to use additives that can lower the cloud point of biodiesel. Another approach is to remove saturated methyl esters indiscriminately by mixing in additives and allowing for the crystallization of saturates and then filtering out the crystals. Fractionation selectively separates methyl esters into individual components or fractions, allowing for the removal or inclusion of specific methyl esters. Fractionation methods include urea fractionation, solvent fractionation and thermal distillation.

[0256] Another valuable fuel provided by the methods of the present invention is renewable diesel, which comprises alkanes, such as C10:0, C12:0, C14:0, C16:0 and C18:0 and thus, are distinguishable from biodiesel. High quality renewable diesel conforms to the

ASTM D975 standard. The lipids produced by the methods of the present invention can serve as feedstock to produce renewable diesel. Thus, in another aspect of the present invention, a method for producing renewable diesel is provided. Renewable diesel can be produced by at least three processes: hydrothermal processing (hydrotreating); hydroprocessing; and indirect liquefaction. These processes yield non-ester distillates. During these processes, triacylglycerides produced and isolated as described herein, are converted to alkanes.

[0257] In one embodiment, the method for producing renewable diesel comprises (a) cultivating a lipid-containing microorganism using methods disclosed herein (b) lysing the microorganism to produce a lysate, (c) isolating lipid from the lysed microorganism, and (d) deoxygenating and hydrotreating the lipid to produce an alkane, whereby renewable diesel is produced. Lipids suitable for manufacturing renewable diesel can be obtained via extraction from microbial biomass using an organic solvent such as hexane, or via other methods, such as those described in US Patent 5,928,696. Some suitable methods may include mechanical pressing and centrifuging.

[0258] In some methods, the microbial lipid is first cracked in conjunction with hydrotreating to reduce carbon chain length and saturate double bonds, respectively. The material is then isomerized, also in conjunction with hydrotreating. The naptha fraction can then be removed through distillation, followed by additional distillation to vaporize and distill components desired in the diesel fuel to meet an ASTM D975 standard while leaving components that are heavier than desired for meeting the D975 standard. Hydrotreating, hydrocracking, deoxygenation and isomerization methods of chemically modifying oils, including triglyceride oils, are well known in the art. See for example European patent applications EP1741768 (A1); EP1741767 (A1); EP1682466 (A1); EP1640437 (A1); EP1681337 (A1); EP1795576 (A1); and U.S. Patents 7,238,277; 6,630,066; 6,596,155; 6,977,322; 7,041,866; 6,217,746; 5,885,440; 6,881,873.

[0259] In one embodiment of the method for producing renewable diesel, treating the lipid to produce an alkane is performed by hydrotreating of the lipid composition. In hydrothermal processing, typically, biomass is reacted in water at an elevated temperature and pressure to form oils and residual solids. Conversion temperatures are typically 300° to 660°F, with pressure sufficient to keep the water primarily as a liquid, 100 to 170 standard atmosphere (atm). Reaction times are on the order of 15 to 30 minutes. After the reaction is completed, the organics are separated from the water. Thereby a distillate suitable for diesel is produced.

[0260] In some methods of making renewable diesel, the first step of treating a triglyceride is hydroprocessing to saturate double bonds, followed by deoxygenation at elevated

temperature in the presence of hydrogen and a catalyst. In some methods, hydrogenation and deoxygenation occur in the same reaction. In other methods deoxygenation occurs before hydrogenation. Isomerization is then optionally performed, also in the presence of hydrogen and a catalyst. Naphtha components are preferably removed through distillation. For examples, see U.S. Patents 5,475,160 (hydrogenation of triglycerides); 5,091,116 (deoxygenation, hydrogenation and gas removal); 6,391,815 (hydrogenation); and 5,888,947 (isomerization).

[0261] One suitable method for the hydrogenation of triglycerides includes preparing an aqueous solution of copper, zinc, magnesium and lanthanum salts and another solution of alkali metal or preferably, ammonium carbonate. The two solutions may be heated to a temperature of about 20°C to about 85°C and metered together into a precipitation container at rates such that the pH in the precipitation container is maintained between 5.5 and 7.5 in order to form a catalyst. Additional water may be used either initially in the precipitation container or added concurrently with the salt solution and precipitation solution. The resulting precipitate may then be thoroughly washed, dried, calcined at about 300°C and activated in hydrogen at temperatures ranging from about 100°C to about 400°C. One or more triglycerides may then be contacted and reacted with hydrogen in the presence of the above-described catalyst in a reactor. The reactor may be a trickle bed reactor, fixed bed gas-solid reactor, packed bubble column reactor, continuously stirred tank reactor, a slurry phase reactor, or any other suitable reactor type known in the art. The process may be carried out either batchwise or in continuous fashion. Reaction temperatures are typically in the range of from about 170°C to about 250°C while reaction pressures are typically in the range of from about 300 psig to about 2000 psig. Moreover, the molar ratio of hydrogen to triglyceride in the process of the present invention is typically in the range of from about 20:1 to about 700:1. The process is typically carried out at a weight hourly space velocity (WHSV) in the range of from about 0.1 hr^{-1} to about 5 hr^{-1}. One skilled in the art willrecognize that the time period required for reaction will vary according to the temperature used, the molar ratio of hydrogen to triglyceride, and the partial pressure of hydrogen. The products produced by the such hydrogenation processes include fatty alcohols, glycerol, traces of paraffins and unreacted triglycerides. These products are typically separated by conventional means such as, for example, distillation, extraction, filtration, crystallization, and the like.

[0262] Petroleum refiners use hydroprocessing to remove impurities by treating feeds with hydrogen. Hydroprocessing conversion temperatures are typically 300° to 700°F. Pressures are typically 40 to 100 atm. The reaction times are typically on the order of 10 to 60 minutes.

Solid catalysts are employed to increase certain reaction rates, improve selectivity for certain products, and optimize hydrogen consumption.

[0263] Suitable methods for the deoxygenation of an oil includes heating an oil to a temperature in the range of from about 350°F to about 550°F and continuously contacting the heated oil with nitrogen under at least pressure ranging from about atmospeheric to above for at least about 5 minutes.

[0264] Suitable methods for isomerization includes using alkali isomerization and other oil isomerization known in the art.

[0265] Hydrotreating and hydroprocessing ultimately lead to a reduction in the molecular weight of the triglyceride feed. The triglyceride molecule is reduced to four hydrocarbon molecules under hydroprocessing conditions: a propane molecule and three heavier hydrocarbon molecules, typically in the C8 to C18 range.

[0266] Thus, in one embodiment, the product of one or more chemical reaction(s) performed on lipid compositions of the invention is an alkane mixture that comprises ASTM D975 renewable diesel. Production of hydrocarbons by microorganisms is reviewed by Metzger et al. Appl Microbiol Biotechnol (2005) 66: 486–496 and A Look Back at the U.S. Department of Energy's Aquatic Species Program: Biodiesel from Algae, NREL/TP-580-24190, John Sheehan, Terri Dunahay, John Benemann and Paul Roessler (1998).

[0267] The distillation properties of a diesel fuel is described in terms of T10-T90 (temperature at 10% and 90%, respectively, volume distilled). Renewable diesel was produced from *Prototheca moriformis* triglyceride oil and is described in Example 14. The T10-T90 of the material produced in Example 14 was 57.9°C. Methods of hydrotreating, isomerization, and other covalent modification of oils disclosed herein, as well as methods of distillation and fractionation (such as cold filtration) disclosed herein, can be employed to generate renewable diesel compositions with other T10-T90 ranges, such as 20, 25, 30, 35, 40, 45, 50, 60 and 65°C using triglyceride oils produced according to the methods disclosed herein.

[0268] The T10 of the material produced in Example 14 was 242.1°C. Methods of hydrotreating, isomerization, and other covalent modification of oils disclosed herein, as well as methods of distillation and fractionation (such as cold filtration) disclosed herein, can be employed to generate renewable diesel compositions with other T10 values, such as T10 between 180 and 295, between 190 and 270, between 210 and 250, between 225 and 245, and at least 290.

[0269] The T90 of the material produced in Example 14 was 300°C. Methods of hydrotreating, isomerization, and other covalent modification of oils disclosed herein, as well as methods of distillation and fractionation (such as cold filtration) disclosed herein can be employed to generate renewable diesel compositions with other T90 values, such as T90 between 280 and 380, between 290 and 360, between 300 and 350, between 310 and 340, and at least 290.

[0270] The FBP of the material produced in Example 14 was 300°C. Methods of hydrotreating, isomerization, and other covalent modification of oils disclosed herein, as well as methods of distillation and fractionation (such as cold filtration) disclosed herein, can be employed to generate renewable diesel compositions with other FBP values, such as FBP between 290 and 400, between 300 and 385, between 310 and 370, between 315 and 360, and at least 300.

[0271] Other oils provided by the methods and compositions of the invention can be subjected to combinations of hydrotreating, isomerization, and other covalent modification including oils with lipid profiles including (a) at least 4% C8-C14; (b) at least 0.3% C8; (c) at least 2% C10; (d) at least 2% C12; and (3) at least 30% C8-C14.

[0272] A traditional ultra-low sulfur diesel can be produced from any form of biomass by a two-step process. First, the biomass is converted to a syngas, a gaseous mixture rich in hydrogen and carbon monoxide. Then, the syngas is catalytically converted to liquids. Typically, the production of liquids is accomplished using Fischer-Tropsch (FT) synthesis. This technology applies to coal, natural gas, and heavy oils. Thus, in yet another preferred embodiment of the method for producing renewable diesel, treating the lipid composition to produce an alkane is performed by indirect liquefaction of the lipid composition.

[0273] The present invention also provides methods to produce jet fuel. Jet fuel is clear to straw colored. The most common fuel is an unleaded/paraffin oil-based fuel classified as Aeroplane A-1, which is produced to an internationally standardized set of specifications. Jet fuel is a mixture of a large number of different hydrocarbons, possibly as many as a thousand or more. The range of their sizes (molecular weights or carbon numbers) is restricted by the requirements for the product, for example, freezing point or smoke point. Kerosone-type Aeroplane fuel (including Jet A and Jet A-1) has a carbon number distribution between about 8 and 16 carbon numbers. Wide-cut or naphta-type Aeroplane fuel (including Jet B) typically has a carbon number distribution between about 5 and 15 carbons.

[0274] Both Aeroplanes (Jet A and Jet B) may contain a number of additives. Useful additives include, but are not limited to, antioxidants, antistatic agents, corrosion inhibitors,

and fuel system icing inhibitor (FSII) agents. Antioxidants prevent gumming and usually, are based on alkylated phenols, for example, AO-30, AO-31, or AO-37. Antistatic agents dissipate static electricity and prevent sparking. Stadis 450 with dinonylnaphthylsulfonic acid (DINNSA) as the active ingredient, is an example. Corrosion inhibitors, e.g., DCI-4A is used for civilian and military fuels and DCI-6A is used for military fuels. FSII agents, include, e.g., Di-EGME.

[0275] In one embodiment of the invention, a jet fuel is produced by blending algal fuels with existing jet fuel. The lipids produced by the methods of the present invention can serve as feedstock to produce jet fuel. Thus, in another aspect of the present invention, a method for producing jet fuel is provided. Herewith two methods for producing jet fuel from the lipids produced by the methods of the present invention are provided: fluid catalytic cracking (FCC); and hydrodeoxygenation (HDO).

[0276] Fluid Catalytic Cracking (FCC) is one method which is used to produce olefins, especially propylene from heavy crude fractions. The lipids produced by the method of the present invention can be converted to olefins. The process involves flowing the lipids produced through an FCC zone and collecting a product stream comprised of olefins, which is useful as a jet fuel. The lipids produced are contacted with a cracking catalyst at cracking conditions to provide a product stream comprising olefins and hydrocarbons useful as jet fuel.

[0277] In one embodiment, the method for producing jet fuel comprises (a) cultivating a lipid-containing microorganism using methods disclosed herein, (b) lysing the lipid-containing microorganism to produce a lysate, (c) isolating lipid from the lysate, and (d) treating the lipid composition, whereby jet fuel is produced. In one embodiment of the method for producing a jet fuel, the lipid composition can be flowed through a fluid catalytic cracking zone, which, in one embodiment, may comprise contacting the lipid composition with a cracking catalyst at cracking conditions to provide a product stream comprising C_2-C_5 olefins.

[0278] In certain embodiments of this method, it may be desirable to remove any contaminants that may be present in the lipid composition. Thus, prior to flowing the lipid composition through a fluid catalytic cracking zone, the lipid composition is pretreated. Pretreatment may involve contacting the lipid composition with an ion-exchange resin. The ion exchange resin is an acidic ion exchange resin, such as Amberlyst™-15 and can be used as a bed in a reactor through which the lipid composition is flowed, either upflow or downflow. Other pretreatments may include mild acid washes by contacting the lipid

composition with an acid, such as sulfuric, acetic, nitric, or hydrochloric acid. Contacting is done with a dilute acid solution usually at ambient temperature and atmospheric pressure.

[0279] The lipid composition, optionally pretreated, is flowed to an FCC zone where the hydrocarbonaceous components are cracked to olefins. Catalytic cracking is accomplished by contacting the lipid composition in a reaction zone with a catalyst composed of finely divided particulate material. The reaction is catalytic cracking, as opposed to hydrocracking, and is carried out in the absence of added hydrogen or the consumption of hydrogen. As the cracking reaction proceeds, substantial amounts of coke are deposited on the catalyst. The catalyst is regenerated at high temperatures by burning coke from the catalyst in a regeneration zone. Coke-containing catalyst, referred to herein as "coked catalyst", is continually transported from the reaction zone to the regeneration zone to be regenerated and replaced by essentially coke-free regenerated catalyst from the regeneration zone. Fluidization of the catalyst particles by various gaseous streams allows the transport of catalyst between the reaction zone and regeneration zone. Methods for cracking hydrocarbons, such as those of the lipid composition described herein, in a fluidized stream of catalyst, transporting catalyst between reaction and regeneration zones, and combusting coke in the regenerator are well known by those skilled in the art of FCC processes. Exemplary FCC applications and catalysts useful for cracking the lipid composition to produce C_2-C_5 olefins are described in U.S. Pat. Nos. 6,538,169, 7,288,685, which are incorporated in their entirety by reference.

[0280] Suitable FCC catalysts generally comprise at least two components that may or may not be on the same matrix. In some embodiments, both two components may be circulated throughout the entire reaction vessel. The first component generally includes any of the well-known catalysts that are used in the art of fluidized catalytic cracking, such as an active amorphous clay-type catalyst and/or a high activity, crystalline molecular sieve. Molecular sieve catalysts may be preferred over amorphous catalysts because of their much-improved selectivity to desired products. IN some preferred embodiments, zeolites may be used as the molecular sieve in the FCC processes. Preferably, the first catalyst component comprises a large pore zeolite, such as an Y-type zeolite, an active alumina material, a binder material, comprising either silica or alumina and an inert filler such as kaolin.

[0281] In one embodiment, cracking the lipid composition of the present invention, takes place in the riser section or, alternatively, the lift section, of the FCC zone. The lipid composition is introduced into the riser by a nozzle resulting in the rapid vaporization of the lipid composition. Before contacting the catalyst, the lipid composition will ordinarily have a

temperature of about 149°C to about 316°C (300°F to 600°F). The catalyst is flowed from a blending vessel to the riser where it contacts the lipid composition for a time of abort 2 seconds or less.

[0282] The blended catalyst and reacted lipid composition vapors are then discharged from the top of the riser through an outlet and separated into a cracked product vapor stream including olefins and a collection of catalyst particles covered with substantial quantities of coke and generally referred to as "coked catalyst." In an effort to minimize the contact time of the lipid composition and the catalyst which may promote further conversion of desired products to undesirable other products, any arrangement of separators such as a swirl arm arrangement can be used to remove coked catalyst from the product stream quickly. The separator, e.g. swirl arm separator, is located in an upper portion of a chamber with a stripping zone situated in the lower portion of the chamber. Catalyst separated by the swirl arm arrangement drops down into the stripping zone. The cracked product vapor stream comprising cracked hydrocarbons including light olefins and some catalyst exit the chamber via a conduit which is in communication with cyclones. The cyclones remove remaining catalyst particles from the product vapor stream to reduce particle concentrations to very low levels. The product vapor stream then exits the top of the separating vessel. Catalyst separated by the cyclones is returned to the separating vessel and then to the stripping zone. The stripping zone removes adsorbed hydrocarbons from the surface of the catalyst by counter-current contact with steam.

[0283] Low hydrocarbon partial pressure operates to favor the production of light olefins. Accordingly, the riser pressure is set at about 172 to 241 kPa (25 to 35 psia) with a hydrocarbon partial pressure of about 35 to 172 kPa (5 to 25 psia), with a preferred hydrocarbon partial pressure of about 69 to 138 kPa (10 to 20 psia). This relatively low partial pressure for hydrocarbon is achieved by using steam as a diluent to the extent that the diluent is 10 to 55 wt-% of lipid composition and preferably about 15 wt-% of lipid composition. Other diluents such as dry gas can be used to reach equivalent hydrocarbon partial pressures.

[0284] The temperature of the cracked stream at the riser outlet will be about 510°C to 621°C (950°F to 1150°F). However, riser outlet temperatures above 566°C (1050°F) make more dry gas and more olefins. Whereas, riser outlet temperatures below 566°C (1050°F) make less ethylene and propylene. Accordingly, it is preferred to run the FCC process at a preferred temperature of about 566°C to about 630°C, preferred pressure of about 138 kPa to about 240 kPa (20 to 35 psia). Another condition for the process is the catalyst to lipid

composition ratio which can vary from about 5 to about 20 and preferably from about 10 to about 15.

[0285] In one embodiment of the method for producing a jet fuel, the lipid composition is introduced into the lift section of an FCC reactor. The temperature in the lift section will be very hot and range from about 700°C (1292°F) to about 760°C (1400°F) with a catalyst to lipid composition ratio of about 100 to about 150. It is anticipated that introducing the lipid composition into the lift section will produce considerable amounts of propylene and ethylene.

[0286] In another embodiment of the method for producing a jet fuel using the lipid composition or the lipids produced as described herein, the structure of the lipid composition or the lipids is broken by a process referred to as hydrodeoxygenation (HDO). HDO means removal of oxygen by means of hydrogen, that is, oxygen is removed while breaking the structure of the material. Olefinic double bonds are hydrogenated and any sulphur and nitrogen compounds are removed. Sulphur removal is called hydrodesulphurization (HDS). Pretreatment and purity of the raw materials (lipid composition or the lipids) contribute to the service life of the catalyst.

[0287] Generally in the HDO/HDS step, hydrogen is mixed with the feed stock (lipid composition or the lipids) and then the mixture is passed through a catalyst bed as a co-current flow, either as a single phase or a two phase feed stock. After the HDO/MDS step, the product fraction is separated and passed to a separate isomerzation reactor. An isomerization reactor for biological starting material is described in the literature (FI 100 248) as a co-current reactor.

[0288] The process for producing a fuel by hydrogenating a hydrocarbon feed, e.g., the lipid composition or the lipids herein, can also be performed by passing the lipid composition or the lipids as a co-current flow with hydrogen gas through a first hydrogenation zone, and thereafter the hydrocarbon effluent is further hydrogenated in a second hydrogenation zone by passing hydrogen gas to the second hydrogenation zone as a counter-current flow relative to the hydrocarbon effluent. Exemplary HDO applications and catalysts useful for cracking the lipid composition to produce C_2-C_5 olefins are described in U.S. Pat. No. 7,232,935, which is incorporated in its entirety by reference.

[0289] Typically, in the hydrodeoxygenation step, the structure of the biological component, such as the lipid composition or lipids herein, is decomposed, oxygen, nitrogen, phosphorus and sulphur compounds, and light hydrocarbons as gas are removed, and the olefinic bonds are hydrogenated. In the second step of the process, i.e. in the so-called

isomerization step, isomerzation is carried out for branching the hydrocarbon chain and improving the performance of the paraffin at low temperatures.

[0290] In the first step, i.e. HDO step, of the cracking process, hydrogen gas and the lipid composition or lipids herein which are to be hydrogenated are passed to a HDO catalyst bed system either as co-current or counter-current flows, said catalyst bed system comprising one or more catalyst bed(s), preferably 1-3 catalyst beds. The HDO step is typically operated in a co-current manner. In case of a HDO catalyst bed system comprising two or more catalyst beds, one or more of the beds may be operated using the counter-current flow principle. In the HDO step, the pressure varies between 20 and 150 bar, preferably between 50 and 100 bar, and the temperature varies between 200 and 500°C, preferably in the range of 300-400°C. In the HDO step, known hydrogenation catalysts containing metals from Group VII and/or VIB of the Periodic System may be used. Preferably, the hydrogenation catalysts are supported Pd, Pt, Ni, NiMo or a CoMo catalysts, the support being alumina and/or silica. Typically, $NiMo/Al_2O_3$ and $CoMo/Al_2O_3$ catalysts are used.

[0291] Prior to the HDO step, the lipid composition or lipids herein may optionally be treated by prehydrogenation under milder conditions thus avoiding side reactions of the double bonds. Such prehydrogenation is carried out in the presence of a prehydrogenation catalyst at temperatures of 50 400°C and at hydrogen pressures of 1 200 bar, preferably at a temperature between 150 and 250°C and at a hydrogen pressure between 10 and 100 bar. The catalyst may contain metals from Group VIII and/or VIB of the Periodic System. Preferably, the prehydrogenation catalyst is a supported Pd, Pt, Ni, NiMo or a CoMo catalyst, the support being alumina and/or silica.

[0292] A gaseous stream from the HDO step containing hydrogen is cooled and then carbon monoxide, carbon dioxide, nitrogen, phosphorus and sulphur compounds, gaseous light hydrocarbons and other impurities are removed therefrom. After compressing, the purified hydrogen or recycled hydrogen is returned back to the first catalyst bed and/or between the catalyst beds to make up for the withdrawn gas stream. Water is removed from the condensed liquid. The liquid is passed to the first catalyst bed or between the catalyst beds.

[0293] After the HDO step, the product is subjected to an isomerization step. It is substantial for the process that the impurities are removed as completely as possible before the hydrocarbons are contacted with the isomerization catalyst. The isomerization step comprises an optional stripping step, wherein the reaction product from the HDO step may be purified by stripping with water vapour or a suitable gas such as light hydrocarbon, nitrogen

or hydrogen. The optional stripping step is carried out in counter-current manner in a unit upstream of the isomerization catalyst, wherein the gas and liquid are contacted with each other, or before the actual isomerization reactor in a separate stripping unit utilizing counter-current principle.

[0294] After the stripping step the hydrogen gas and the hydrogenated lipid composition or lipids herein, and optionally an n-paraffin mixture, are passed to a reactive isomerization unit comprising one or several catalyst bed(s). The catalyst beds of the isomerization step may operate either in co-current or counter-current manner.

[0295] It is important for the process that the counter-current flow principle is applied in the isomerization step. In the isomerization step this is done by carrying out either the optional stripping step or the isomerization reaction step or both in counter-current manner. In the isomerzation step, the pressure varies in the range of 20 150 bar, preferably in the range of 20 100 bar, the temperature being between 200 and 500°C, preferably between 300 and 400°C. In the isomerization step, isomerization catalysts known in the art may be used. Suitable isomerization catalysts contain molecular sieve and/or a metal from Group VII and/or a carrier. Preferably, the isomerization catalyst contains SAPO-11 or SAPO41 or ZSM-22 or ZSM-23 or ferrierite and Pt, Pd or Ni and Al_2O_3 or SiO_2. Typical isomerization catalysts are, for example, Pt/SAPO-11/Al_2O_3, Pt/ZSM-22/Al_2O_3, Pt/ZSM-23/Al_2O_3 and Pt/SAPO-11/SiO_2. The isomerization step and the HDO step may be carried out in the same pressure vessel or in separate pressure vessels. Optional prehydrogenation may be carried out in a separate pressure vessel or in the same pressure vessel as the HDO and isomerization steps.

[0296] Thus, in one embodiment, the product of the one or more chemical reactions is an alkane mixture that comprises ASTM D1655 jet fuel. In some embodiments, the composition comforming to the specification of ASTM 1655 jet fuel has a sulfur content that is less than 10 ppm. In other embodiments, the composition conforming to the specification of ASTM 1655 jet fuel has a T10 value of the distillation curve of less than 205° C. In another embodiment, the composition conforming to the specification of ASTM 1655 jet fuel has a final boiling point (FBP) of less than 300° C. In another embodiment, the composition conforming to the specification of ASTM 1655 jet fuel has a flash point of at least 38° C. In another embodiment, the composition conforming to the specification of ASTM 1655 jet fuel has a density between 775K/M^3 and 840K/M^3. In yet another embodiment, the composition conforming to the specification of ASTM 1655 jet fuel has a freezing point that is below -47° C. In another embodiment, the composition conforming to the specification of ASTM 1655

jet fuel has a net Heat of Combustion that is at least 42.8 MJ/K. In another embodiment, the composition conforming to the specification of ASTM 1655 jet fuel has a hydrogen content that is at least 13.4 mass %. In another embodiment, the composition conforming to the specification of ASTM 1655 jet fuel has a thermal stability, as tested by quantitative gravimetric JFTOT at 260° C, that is below 3mm of Hg. In another embodiment, the composition conforming to the specification of ASTM 1655 jet fuel has an existent gum that is below 7 mg/dl.

[0297] Thus, the present invention discloses a variety of methods in which chemical modification of microalgal lipid is undertaken to yield products useful in a variety of industrial and other applications. Examples of processes for modifying oil produced by the methods disclosed herein include, but are not limited to, hydrolysis of the oil, hydroprocessing of the oil, and esterification of the oil. The modification of the microalgal oil produces basic oleochemicals that can be further modified into selected derivative oleochemicals for a desired function. In a manner similar to that described above with reference to fuel producing processes, these chemical modifications can also be performed on oils generated from the microbial cultures described herein. Examples of basic oleochemicals include, but are not limited to, soaps, fatty acids, fatty acid methyl esters, and glycerol. Examples of derivative oleochemicals include, but are not limited to, fatty nitriles, esters, dimer acids, quats, surfactants, fatty alkanolamides, fatty alcohol sulfates, resins, emulsifiers, fatty alcohols, olefins, and higher alkanes.

[0298] Hydrolysis of the fatty acid constituents from the glycerolipids produced by the methods of the invention yields free fatty acids that can be derivatized to produce other useful chemicals. Hydrolysis occurs in the presence of water and a catalyst which may be either an acid or a base. The liberated free fatty acids can be derivatized to yield a variety of products, as reported in the following: US Patent Nos. 5,304,664 (Highly sulfated fatty acids); 7,262,158 (Cleansing compositions); 7,115,173 (Fabric softener compositions); 6,342,208 (Emulsions for treating skin); 7,264,886 (Water repellant compositions); 6,924,333 (Paint additives); 6,596,768 (Lipid-enriched ruminant feedstock); and 6,380,410 (Surfactants for detergents and cleaners).

[0299] With regard to hydrolysis, in one embodiment of the invention, a triglyceride oil is optionally first hydrolyzed in a liquid medium such as water or sodium hydroxide so as to obtain glycerol and soaps. There are various suitable triglyceride hydrolysis methods, including, but not limited to, saponification, acid hydrolysis, alkaline hydrolysis, enzymatic hydrolysis (referred herein as splitting), and hydrolysis using hot-compressed water. One

skilled in the art will recognize that a triglyceride oil need not be hydrolyzed in order to produce an oleochemical; rather, the oil may be converted directly to the desired oleochemical by other known process. For example, the triglyceride oil may be directly converted to a methyl ester fatty acid through esterification.

[0300] In some embodiments, catalytic hydrolysis of the oil produced by methods disclosed herein occurs by splitting the oil into glycerol and fatty acids. As discussed above, the fatty acids may then be further processed through several other modifications to obtained derivative oleochemicals. For example, in one embodiment the fatty acids may undergo an amination reaction to produce fatty nitrogen compounds. In another embodiment, the fatty acids may undergo ozonolysis to produce mono- and dibasic-acids.

[0301] In other embodiments hydrolysis may occur via the, splitting of oils produced herein to create oleochemicals. In some preferred embodiments of the invention, a triglyceride oil may be split before other processes is performed. One skilled in the art will recognize that there are many suitable triglyceride splitting methods, including, but not limited to, enzymatic splitting and pressure splitting.

[0302] Generally, enzymatic oil splitting methods use enzymes, lipases, as biocatalysts acting on a water/oil mixture. Enzymatic splitting then slpits the oil or fat, respectively, is into glycerol and free fatty acids. The glycerol may then migrates into the water phase whereas the organic phase enriches with free fatty acids.

[0303] The enzymatic splitting reactions generally take place at the phase boundary between organic and aqueous phase, where the enzyme is present only at the phase boundary. Triglycerides that meet the phase boundary then contribute to or participate in the splitting reaction. As the reaction proceeds, the occupation density or concentration of fatty acids still chemically bonded as glycerides, in comparison to free fatty acids, decreases at the phase boundary so that the reaction is slowed down. In certain embodiments, enzymatic splitting may occur at room temperature. One of ordinary skill in the art would know the suitable conditions for splitting oil into the desired fatty acids.

[0304] By way of example, the reaction speed can be accelerated by increasing the interface boundary surface. Once the reaction is complete, free fatty acids are then separated from the organic phase freed from enzyme, and the residue which still contains fatty acids chemically bonded as glycerides is fed back or recycled and mixed with fresh oil or fat to be subjected to splitting. In this manner, recycled glycerides are then subjected to a further enzymatic splitting process. In some embodiments, the free fatty acids are extracted from an oil or fat partially split in such a manner. In that way, if the chemically bound fatty acids

(triglycerides) are returned or fed back into the splitting process, the enzyme consumption can be drastically reduced.

[0305] The splitting degree is determined as the ratio of the measured acid value divided by the theoretically possible acid value which can be computed for a given oil or fat. Preferably, the acid value is measured by means of titration according to standard common methods. Alternatively, the density of the aqueous glycerol phase can be taken as a measure for the splitting degree.

[0306] In one embodiment, the slitting process as described herein is also suitable for splitting the mono-, di- and triglyceride that are contained in the so-called soap-stock from the alkali refining processes of the produced oils. In this manner, the soap-stock can be quantitatively converted without prior saponification of the neutral oils into the fatty acids. For this purpose, the fatty acids being chemically bonded in the soaps are released, preferably before splitting, through an addition of acid. In certain embodiments, a buffer solution is used in addition to water and enzyme for the splitting process.

[0307] In one embodiment, oils produced in accordance with the methods of the invention can also be subjected to saponification as a method of hydrolysis. Animal and plant oils are typically made of triacylglycerols (TAGs), which are esters of fatty acids with the trihydric alcohol, glycerol. In an alkaline hydrolysis reaction, the glycerol in a TAG is removed, leaving three carboxylic acid anions that can associate with alkali metal cations such as sodium or potassium to produce fatty acid salts. In this scheme, the carboxylic acid constituents are cleaved from the glycerol moiety and replaced with hydroxyl groups. The quantity of base (e.g., KOH) that is used in the reaction is determined by the desired degree of saponification. If the objective is, for example, to produce a soap product that comprises some of the oils originally present in the TAG composition, an amount of base insufficient to convert all of the TAGs to fatty acid salts is introduced into the reaction mixture. Normally, this reaction is performed in an aqueous solution and proceeds slowly, but may be expedited by the addition of heat. Precipitation of the fatty acid salts can be facilitated by addition of salts, such as water-soluble alkali metal halides (e.g., NaCl or KCl), to the reaction mixture. Preferably, the base is an alkali metal hydroxide, such as NaOH or KOH. Alternatively, other bases, such as alkanolamines, including for example triethanolamine and aminomethylpropanol, can be used in the reaction scheme. In some cases, these alternatives may be preferred to produce a clear soap product.

[0308] In some methods, the first step of chemical modification may be hydroprocessing to saturate double bonds, followed by deoxygenation at elevated temperature in the presence of

hydrogen and a catalyst. In other methods, hydrogenation and deoxygenation may occur in the same reaction. In still other methods deoxygenation occurs before hydrogenation. Isomerization may then be optionally performed, also in the presence of hydrogen and a catalyst. Finally, gases and naphtha components can be removed if desired. For example, see U.S. Patents 5,475,160 (hydrogenation of triglycerides); 5,091,116 (deoxygenation, hydrogenation and gas removal); 6,391,815 (hydrogenation); and 5,888,947 (isomerization).

[0309] In some embodiments of the invention, the triglyceride oils are partially or completely deoxygenated. The deoxygenation reactions form desired products, including, but not limited to, fatty acids, fatty alcohols, polyols, ketones, and aldehydes. In general, without being limited by any particular theory, the deoxygenation reactions involve a combination of various different reaction pathways, including without limitation: hydrogenolysis, hydrogenation, consecutive hydrogenation-hydrogenolysis, consecutive hydrogenolysis-hydrogenation, and combined hydrogenation-hydrogenolysis reactions, resulting in at least the partial removal of oxygen from the fatty acid or fatty acid ester to produce reaction products, such as fatty alcohols, that can be easily converted to the desired chemicals by further processing. For example, in one embodiment, a fatty alcohol may be converted to olefins through FCC reaction or to higher alkanes through a condensation reaction.

[0310] One such chemical modification is hydrogenation, which is the addition of hydrogen to double bonds in the fatty acid constituents of glycerolipids or of free fatty acids. The hydrogenation process permits the transformation of liquid oils into semi-solid or solid fats, which may be more suitable for specific applications.

[0311] Hydrogenation of oil produced by the methods described herein can be performed in conjunction with one or more of the methods and/or materials provided herein, as reported in the following: US Patent Nos. 7,288,278 (Food additives or medicaments); 5,346,724 (Lubrication products); 5,475,160 (Fatty alcohols); 5,091,116 (Edible oils); 6,808,737 (Structural fats for margarine and spreads); 5,298,637 (Reduced-calorie fat substitutes); 6,391,815 (Hydrogenation catalyst and sulfur adsorbent); 5,233,099 and 5,233,100 (Fatty alcohols); 4,584,139 (Hydrogenation catalysts); 6,057,375 (Foam suppressing agents); and 7,118,773 (Edible emulsion spreads).

[0312] One skilled in the art will recognize that various processes may be used to hydrogenate carbohydrates. One suitable method includes contacting the carbohydrate with hydrogen or hydrogen mixed with a suitable gas and a catalyst under conditions sufficient in a hydrogenation reactor to form a hydrogenated product. The hydrogenation catalyst

generally can include Cu, Re, Ni, Fe, Co, Ru, Pd, Rh, Pt, Os, Ir, and alloys or any combination thereof, either alone or with promoters such as W, Mo, Au, Ag, Cr, Zn, Mn, Sn, B, P, Bi, and alloys or any combination thereof. Other effective hydrogenation catalyst materials include either supported nickel or ruthenium modified with rhenium. In an embodiment, the hydrogenation catalyst also includes any one of the supports, depending on the desired functionality of the catalyst. The hydrogenation catalysts may be prepared by methods known to those of ordinary skill in the art.

[0313] In some embodiments the hydrogenation catalyst includes a supported Group VIII metal catalyst and a metal sponge material (e.g., a sponge nickel catalyst). Raney nickel provides an example of an activated sponge nickel catalyst suitable for use in this invention. In other embodiment, the hydrogenation reaction in the invention is performed using a catalyst comprising a nickel-rhenium catalyst or a tungsten-modified nickel catalyst. One example of a suitable catalyst for the hydrogenation reaction of the invention is a carbon-supported nickel-rhenium catalyst.

[0314] In an embodiment, a suitable Raney nickel catalyst may be prepared by treating an alloy of approximately equal amounts by weight of nickel and aluminum with an aqueous alkali solution, e.g., containing about 25 weight % of sodium hydroxide. The aluminum is selectively dissolved by the aqueous alkali solution resulting in a sponge shaped material comprising mostly nickel with minor amounts of aluminum. The initial alloy includes promoter metals (i.e., molybdenum or chromium) in the amount such that about 1 to 2 weight % remains in the formed sponge nickel catalyst. In another embodiment, the hydrogenation catalyst is prepared using a solution of ruthenium(III) nitrosylnitrate, ruthenium (III) chloride in water to impregnate a suitable support material. The solution is then dried to form a solid having a water content of less than about 1% by weight. The solid may then be reduced at atmospheric pressure in a hydrogen stream at 300°C (uncalcined) or 400°C (calcined) in a rotary ball furnace for 4 hours. After cooling and rendering the catalyst inert with nitrogen, 5% by volume of oxygen in nitrogen is passed over the catalyst for 2 hours.

[0315] In certain embodiments, the catalyst described includes a catalyst support. The catalyst support stabilizes and supports the catalyst. The type of catalyst support used depends on the chosen catalyst and the reaction conditions. Suitable supports for the invention include, but are not limited to, carbon, silica, silica-alumina, zirconia, titania, ceria, vanadia, nitride, boron nitride, heteropolyacids, hydroxyapatite, zinc oxide, chromia, zeolites, carbon nanotubes, carbon fullerene and any combination thereof.

[0316] The catalysts used in this invention can be prepared using conventional methods known to those in the art. Suitable methods may include, but are not limited to, incipient wetting, evaporative impregnation, chemical vapor deposition, wash-coating, magnetron sputtering techniques, and the like.

[0317] The conditions for which to carry out the hydrogenation reaction will vary based on the type of starting material and the desired products. One of ordinary skill in the art, with the benefit of this disclosure, will recognize the appropriate reaction conditions. In general, the hydrogenation reaction is conducted at temperatures of 80°C to 250°C, and preferably at 90°C to 200°C, and most preferably at 100°C to 150°C. In some embodiments, the hydrogenation reaction is conducted at pressures from 500 KPa to 14000 KPa.

[0318] The hydrogen used in the hydrogenolysis reaction of the current invention may include external hydrogen, recycled hydrogen, in situ generated hydrogen, and any combination thereof. As used herein, the term "external hydrogen" refers to hydrogen that does not originate from the biomass reaction itself, but rather is added to the system from another source.

[0319] In some embodiments of the invention, it is desirable to convert the starting carbohydrate to a smaller molecule that will be more readily converted to desired higher hydrocarbons. One suitable method for this conversion is through a hydrogenolysis reaction. Various processes are known for performing hydrogenolysis of carbohydrates. One suitable method includes contacting a carbohydrate with hydrogen or hydrogen mixed with a suitable gas and a hydrogenolysis catalyst in a hydrogenolysis reactor under conditions sufficient to form a reaction product comprising smaller molecules or polyols. As used herein, the term "smaller molecules or polyols" includes any molecule that has a smaller molecular weight, which can include a smaller number of carbon atoms or oxygen atoms than the starting carbohydrate. In an embodiment, the reaction products include smaller molecules that include polyols and alcohols. Someone of ordinary skill in the art would be able to choose the appropriate method by which to carry out the hydrogenolysis reaction.

[0320] In some embodiments, a 5 and/or 6 carbon sugar or sugar alcohol may be converted to propylene glycol, ethylene glycol, and glycerol using a hydrogenolysis catalyst. The hydrogenolysis catalyst may include Cr, Mo, W, Re, Mn, Cu, Cd, Fe, Co, Ni, Pt, Pd, Rh, Ru, Ir, Os, and alloys or any combination thereof, either alone or with promoters such as Au, Ag, Cr, Zn, Mn, Sn, Bi, B, O, and alloys or any combination thereof. The hydrogenolysis catalyst may also include a carbonaceous pyropolymer catalyst containing transition metals (e.g., chromium, molybdemum, tungsten, rhenium, manganese, copper, cadmium) or Group VIII

metals (e.g., iron, cobalt, nickel, platinum, palladium, rhodium, ruthenium, iridium, and osmium). In certain embodiments, the hydrogenolysis catalyst may include any of the above metals combined with an alkaline earth metal oxide or adhered to a catalytically active support. In certain embodiments, the catalyst described in the hydrogenolysis reaction may include a catalyst support as described above for the hydrogenation reaction.

[0321] The conditions for which to carry out the hydrogenolysis reaction will vary based on the type of starting material and the desired products. One of ordinary skill in the art, with the benefit of this disclosure, will recognize the appropriate conditions to use to carry out the reaction. In general, they hydrogenolysis reaction is conducted at temperatures of 110°C to 300°C, and preferably at 170°C to 220°C, and most preferably at 200°C to 225°C. In some embodiments, the hydrogenolysis reaction is conducted under basic conditions, preferably at a pH of 8 to 13, and even more preferably at a pH of 10 to 12. In some embodiments, the hydrogenolysis reaction is conducted at pressures in a range between 60 KPa and 16500 KPa, and preferably in a range between 1700 KPa and 14000 KPa, and even more preferably between 4800 KPa and 11000 KPa.

[0322] The hydrogen used in the hydrogenolysis reaction of the current invention can include external hydrogen, recycled hydrogen, in situ generated hydrogen, and any combination thereof.

[0323] In some embodiments, the reaction products discussed above may be converted into higher hydrocarbons through a condensation reaction in a condensation reactor (shown schematically as condensation reactor 110 in Figure 1). In such embodiments, condensation of the reaction products occurs in the presence of a catalyst capable of forming higher hydrocarbons. While not intending to be limited by theory, it is believed that the production of higher hydrocarbons proceeds through a stepwise addition reaction including the formation of carbon-carbon, or carbon-oxygen bond. The resulting reaction products include any number of compounds containing these moieties, as described in more detail below.

[0324] In certain embodiments, suitable condensation catalysts include an acid catalyst, a base catalyst, or an acid/base catalyst. As used herein, the term "acid/base catalyst" refers to a catalyst that has both an acid and a base functionality. In some embodiments the condensation catalyst can include, without limitation, zeolites, carbides, nitrides, zirconia, alumina, silica, aluminosilicates, phosphates, titanium oxides, zinc oxides, vanadium oxides, lanthanum oxides, yttrium oxides, scandium oxides, magnesium oxides, cerium oxides, barium oxides, calcium oxides, hydroxides, heteropolyacids, inorganic acids, acid modified resins, base modified resins, and any combination thereof. In some embodiments, the

condensation catalyst can also include a modifier. Suitable modifiers include La, Y, Sc, P, B, Bi, Li, Na, K, Rb, Cs, Mg, Ca, Sr, Ba, and any combination thereof. In some embodiments, the condensation catalyst can also include a metal. Suitable metals include Cu, Ag, Au, Pt, Ni, Fe, Co, Ru, Zn, Cd, Ga, In, Rh, Pd, Ir, Re, Mn, Cr, Mo, W, Sn, Os, alloys, and any combination thereof.

[0325] In certain embodiments, the catalyst described in the condensation reaction may include a catalyst support as described above for the hydrogenation reaction. In certain embodiments, the condensation catalyst is self-supporting. As used herein, the term "self-supporting" means that the catalyst does not need another material to serve as support. In other embodiments, the condensation catalyst in used in conjunction with a separate support suitable for suspending the catalyst. In an embodiment, the condensation catalyst support is silica.

[0326] The conditions under which the condensation reaction occurs will vary based on the type of starting material and the desired products. One of ordinary skill in the art, with the benefit of this disclosure, will recognize the appropriate conditions to use to carry out the reaction. In some embodiments, the condensation reaction is carried out at a temperature at which the thermodynamics for the proposed reaction are favorable. The temperature for the condensation reaction will vary depending on the specific starting polyol or alcohol. In some embodiments, the temperature for the condensation reaction is in a range from 80°C to 500°C, and preferably from 125°C to 450°C, and most preferably from 125°C to 250°C. In some embodiments, the condensation reaction is conducted at pressures in a range between 0 Kpa to 9000 KPa, and preferably in a range between 0 KPa and 7000 KPa, and even more preferably between 0 KPa and 5000 KPa.

[0327] The higher alkanes formed by the invention include, but are not limited to, branched or straight chain alkanes that have from 4 to 30 carbon atoms, branched or straight chain alkenes that have from 4 to 30 carbon atoms, cycloalkanes that have from 5 to 30 carbon atoms, cycloalkenes that have from 5 to 30 carbon atoms, aryls, fused aryls, alcohols, and ketones. Suitable alkanes include, but are not limited to, butane, pentane, pentene, 2-methylbutane, hexane, hexene, 2-methylpentane, 3-methylpentane, 2,2,-dimethylbutane, 2,3-dimethylbutane, heptane, heptene, octane, octene, 2,2,4-trimethylpentane, 2,3-dimethyl hexane, 2,3,4-trimethylpentane, 2,3-dimethylpentane, nonane, nonene, decane, decene, undecane, undecene, dodecane, dodecene, tridecane, tridecene, tetradecane, tetradecene, pentadecane, pentadecene, nonyldecane, nonyldecene, eicosane, eicosene, uneicosane,

uneicosene, doeicosane, doeicosene, trieicosane, trieicosene, tetraeicosane, tetraeicosene, and isomers thereof. Some of these products may be suotable for use as fuels.

[0328] In some embodiments, the cycloalkanes and the cycloalkenes are unsubstituted. In other embodiments, the cycloalkanes and cycloalkenes are mono-substituted. In still other embodiments, the cycloalkanes and cycloalkenes are multi-substituted. In the embodiments comprising the substituted cycloalkanes and cycloalkenes, the substituted group includes, without limitation, a branched or straight chain alkyl having 1 to 12 carbon atoms, a branched or straight chain alkylene having 1 to 12 carbon atoms, a phenyl, and any combination thereof. Suitable cycloalkanes and cycloalkenes include, but are not limited to, cyclopentane, cyclopentene, cyclohexane, cyclohexene, methyl-cyclopentane, methyl-cyclopentene, ethyl-cyclopentane, ethyl-cyclopentene, ethyl-cyclohexane, ethyl-cyclohexene, isomers and any combination thereof.

[0329] In some embodiments, the aryls formed are unsubstituted. In another embodiment, the aryls formed are mono-substituted. In the embodiments comprising the substituted aryls, the substituted group includes, without limitation, a branched or straight chain alkyl having 1 to 12 carbon atoms, a branched or straight chain alkylene having 1 to 12 carbon atoms, a phenyl, and any combination thereof. Suitable aryls for the invention include, but are not limited to, benzene, toluene, xylene, ethyl benzene, para xylene, meta xylene, and any combination thereof.

[0330] The alcohols produced in the invention have from 4 to 30 carbon atoms. In some embodiments, the alcohols are cyclic. In other embodiments, the alcohols are branched. In another embodiment, the alcohols are straight chained. Suitable alcohols for the invention include, but are not limited to, butanol, pentanol, hexanol, heptanol, octanol, nonanol, decanol, undecanol, dodecanol, tridecanol, tetradecanol, pentadecanol, hexadecanol, heptyldecanol, octyldecanol, nonyldecanol, eicosanol, uneicosanol, doeicosanol, trieicosanol, tetraeicosanol, and isomers thereof.

[0331] The ketones produced in the invention have from 4 to 30 carbon atoms. In an embodiment, the ketones are cyclic. In another embodiment, the ketones are branched. In another embodiment, the ketones are straight chained. Suitable ketones for the invention include, but are not limited to, butanone, pentanone, hexanone, heptanone, octanone, nonanone, decanone, undecanone, dodecanone, tridecanone, tetradecanone, pentadecanone, hexadecanone, heptyldecanone, octyldecanone, nonyldecanone, eicosanone, uneicosanone, doeicosanone, trieicosanone, tetraeicosanone, and isomers thereof.

[0332] Another such chemical modification is interesterification. Naturally produced glycerolipids do not have a uniform distribution of fatty acid constituents. In the context of oils, interesterification refers to the exchange of acyl radicals between two esters of different glycerolipids. The interesterification process provides a mechanism by which the fatty acid constituents of a mixture of glycerolipids can be rearranged to modify the distribution pattern. Interesterification is a well-known chemical process, and generally comprises heating (to about 200°C) a mixture of oils for a period (e.g, 30 minutes) in the presence of a catalyst, such as an alkali metal or alkali metal alkylate (e.g., sodium methoxide). This process can be used to randomize the distribution pattern of the fatty acid constituents of an oil mixture, or can be directed to produce a desired distribution pattern. This method of chemical modification of lipids can be performed on materials provided herein, such as microbial biomass with a percentage of dry cell weight as lipid at least 20%.

[0333] Directed interesterification, in which a specific distribution pattern of fatty acids is sought, can be performed by maintaining the oil mixture at a temperature below the melting point of some TAGs which might occur. This results in selective crystallization of these TAGs, which effectively removes them from the reaction mixture as they crystallize. The process can be continued until most of the fatty acids in the oil have precipitated, for example. A directed interesterification process can be used, for example, to produce a product with a lower calorie content via the substitution of longer-chain fatty acids with shorter-chain counterparts. Directed interesterification can also be used to produce a product with a mixture of fats that can provide desired melting characteristics and structural features sought in food additives or products (e.g., margarine) without resorting to hydrogenation, which can produce unwanted trans isomers.

[0334] Interesterification of oils produced by the methods described herein can be performed in conjuction with one or more of the methods and/or materials, or to produce products, as reported in the following: US Patent Nos. 6,080,853 (Nondigestible fat substitutes); 4,288,378 (Peanut butter stabilizer); 5,391,383 (Edible spray oil); 6,022,577 (Edible fats for food products); 5,434,278 (Edible fats for food products); 5,268,192 (Low calorie nut products); 5,258,197 (Reduce calorie edible compositions); 4,335,156 (Edible fat product); 7,288,278 (Food additives or medicaments); 7,115,760 (Fractionation process); 6,808,737 (Structural fats); 5,888,947 (Engine lubricants); 5,686,131 (Edible oil mixtures); and 4,603,188 (Curable urethane compositions).

[0335] In one embodiment in accordance with the invention, transesterification of the oil, as described above, is followed by reaction of the transesterified product with polyol, as

reported in US Patent No. 6,465,642, to produce polyol fatty acid polyesters. Such an esterification and separation process may comprise the steps as follows: reacting a lower alkyl ester with polyol in the presence of soap; removing residual soap from the product mixture; water-washing and drying the product mixture to remove impurities; bleaching the product mixture for refinement; separating at least a portion of the unreacted lower alkyl ester from the polyol fatty acid polyester in the product mixture; and recycling the separated unreacted lower alkyl ester.

[0336] Transesterification can also be performed on microbial biomass with short chain fatty acid esters, as reported in U.S. Patent 6,278,006. In general, transesterification may be performed by adding a short chain fatty acid ester to an oil in the presence of a suitable catalyst and heating the mixture. In some embodiments, the oil comprises about 5% to about 90% of the reaction mixture by weight. In some embodiments, the short chain fatty acid esters can be about 10% to about 50% of the reaction mixture by weight. Non-limiting examples of catalysts include base catalysts, sodium methoxide, acid catalysts including inorganic acids such as sulfuric acid and acidified clays, organic acids such as methane sulfonic acid, benzenesulfonic acid, and toluenesulfonic acid, and acidic resins such as Amberlyst 15. Metals such as sodium and magnesium, and metal hydrides also are useful catalysts.

[0337] Another such chemical modification is hydroxylation, which involves the addition of water to a double bond resulting in saturation and the incorporation of a hydroxyl moiety. The hydroxylation process provides a mechanism for converting one or more fatty acid constituents of a glycerolipid to a hydroxy fatty acid. Hydroxylation can be performed, for example, via the method reported in US Patent No. 5,576,027. Hydroxylated fatty acids, including castor oil and its derivatives, are useful as components in several industrial applications, including food additives, surfactants, pigment wetting agents, defoaming agents, water proofing additives, plasticizing agents, cosmetic emulsifying and/or deodorant agents, as well as in electronics, pharmaceuticals, paints, inks, adhesives, and lubricants. One example of how the hydroxylation of a glyceride may be performed is as follows: fat may be heated, preferably to about 30-50°C combined with heptane and maintained at temperature for thirty minutes or more; acetic acid may then be added to the mixture followed by an aqueous solution of sulfuric acid followed by an aqueous hydrogen peroxide solution which is added in small increments to the mixture over one hour; after the aqueous hydrogen peroxide, the temperature may then be increased to at least about 60°C and stirred for at least six hours; after the stirring, the mixture is allowed to settle and a lower aqueous layer formed

by the reaction may be removed while the upper heptane layer formed by the reaction may be washed with hot water having a temperature of about 60°C; the washed heptane layer may then be neutralized with an aqueous potassium hydroxide solution to a pH of about 5 to 7 and then removed by distillation under vacuum; the reaction product may then be dried under vacuum at 100°C and the dried product steam-deodorized under vacuum conditions and filtered at about 50° to 60°C using diatomaceous earth.

[0338] Hydroxylation of microbial oils produced by the methods described herein can be performed in conjuction with one or more of the methods and/or materials, or to produce products, as reported in the following: US Patent Nos. 6,590,113 (Oil-based coatings and ink); 4,049,724 (Hydroxylation process); 6,113,971 (Olive oil butter); 4,992,189 (Lubricants and lube additives); 5,576,027 (Hydroxylated milk); and 6,869,597 (Cosmetics).

[0339] Hydroxylated glycerolipids can be converted to estolides. Estolides consist of a glycerolipid in which a hydroxylated fatty acid constituent has been esterified to another fatty acid molecule. Conversion of hydroxylated glycerolipids to estolides can be carried out by warming a mixture of glycerolipids and fatty acids and contacting the mixture with a mineral acid, as described by Isbell et al., *JAOCS* 71(2):169-174 (1994). Estolides are useful in a variety of applications, including without limitation those reported in the following: US Patent Nos. 7,196,124 (Elastomeric materials and floor coverings); 5,458,795 (Thickened oils for high-temperature applications); 5,451,332 (Fluids for industrial applications); 5,427,704 (Fuel additives); and 5,380,894 (Lubricants, greases, plasticizers, and printing inks).

[0340] Other chemical reactions that can be performed on microbial oils include reacting triacylglycerols with a cyclopropanating agent to enhance fluidity and/or oxidative stability, as reported in U.S. Patent 6,051,539; manufacturing of waxes from triacylglycerols, as reported in U.S. Patent 6,770,104; and epoxidation of triacylglycerols, as reported in "The effect of fatty acid composition on the acrylation kinetics of epoxidized triacylglycerols", Journal of the American Oil Chemists' Society, 79:1, 59-63, (2001) and Free Radical Biology and Medicine, 37:1, 104-114 (2004).

[0341] The generation of oil-bearing microbial biomass for fuel and chemical products as described above results in the production of delipidated biomass meal. Delipidated meal is a byproduct of preparing algal oil and is useful as animal feed for farm animals, e.g., ruminants, poultry, swine and aquaculture. The resulting meal, although of reduced oil content, still contains high quality proteins, carbohydrates, fiber, ash, residual oil and other nutrients appropriate for an animal feed. Because the cells are predominantly lysed by the oil separation process, the delipidated meal is easily digestible by such animals. Delipidated

meal can optionally be combined with other ingredients, such as grain, in an animal feed. Because delipidated meal has a powdery consistency, it can be pressed into pellets using an extruder or expander or another type of machine, which are commercially available.

[0342] The invention, having been described in detail above, is exemplified in the following examples, which are offered to illustrate, but not to limit, the claimed invention.

VII. EXAMPLES

[0343] EXAMPLE 1: Methods for Culturing *Prototheca*

[0344] *Prototheca* strains were cultivated to achieve a high percentage of oil by dry cell weight. Cryopreserved cells were thawed at room temperature and 500 ul of cells were added to 4.5 ml of medium (4.2 g/L K_2HPO_4, 3.1 g/L NaH_2PO_4, 0.24 g/L $MgSO_4 \cdot 7H_2O$, 0.25 g/L Citric Acid monohydrate, 0.025 g/L $CaCl_2$ $2H_2O$, 2g/L yeast extract) plus 2% glucose and grown for 7 days at 28°C with agitation (200 rpm) in a 6-well plate. Dry cell weights were determined by centrifuging 1 ml of culture at 14,000 rpm for 5 min in a pre-weighed Eppendorf tube. The culture supernatant was discarded and the resulting cell pellet washed with 1 ml of deionized water. The culture was again centrifuged, the supernatant discarded, and the cell pellets placed at -80°C until frozen. Samples were then lyophilized for 24 hrs and dry cell weights calculated. For determination of total lipid in cultures, 3 ml of culture was removed and subjected to analysis using an Ankom system (Ankom Inc., Macedon, NY) according to the manufacturer's protocol. Samples were subjected to solvent extraction with an Amkom XT10 extractor according to the manufacturer's protocol. Total lipid was determined as the difference in mass between acid hydrolyzed dried samples and solvent extracted, dried samples. Percent oil dry cell weight measurements are shown in Table 8.

[0345] Table 8. Percent oil by dry cell weight

Species	Strain	% Oil
Prototheca stagnora	UTEX 327	13.14
Prototheca moriformis	UTEX 1441	18.02
Prototheca moriformis	UTEX 1435	27.17

[0346] Microalgae samples from the strains listed in Table 22 above were genotyped. Genomic DNA was isolated from algal biomass as follows. Cells (approximately 200 mg) were centifuged from liquid cultures 5 minutes at 14,000 x g. Cells were then resuspended in sterile distilled water, centrifuged 5 minutes at 14,000 x g and the supernatant discarded. A single glass bead ~2mm in diameter was added to the biomass and tubes were placed at -80°C for at least 15 minutes. Samples were removed and 150 µl of grinding buffer (1% Sarkosyl, 0.25 M Sucrose, 50 mM NaCl, 20 mM EDTA, 100 mM Tris-HCl, pH 8.0, RNase A 0.5

ug/ul) was added. Pellets were resuspended by vortexing briefly, followed by the addition of 40 ul of 5M NaCl. Samples were vortexed briefly, followed by the addition of 66 µl of 5% CTAB (Cetyl trimethylammonium bromide) and a final brief vortex. Samples were next incubated at 65°C for 10 minutes after which they were centrifuged at 14,000 x g for 10 minutes. The supernatant was transferred to a fresh tube and extracted once with 300 µl of Phenol:Chloroform:Isoamyl alcohol 12:12:1, followed by centrifugation for 5 minutes at 14,000 x g. The resulting aqueous phase was transferred to a fresh tube containing 0.7 vol of isopropanol (~190 µl), mixed by inversion and incubated at room temperature for 30 minutes or overnight at 4°C. DNA was recovered via centrifugation at 14,000 x g for 10 minutes. The resulting pellet was then washed twice with 70% ethanol, followed by a final wash with 100% ethanol. Pellets were air dried for 20-30 minutes at room temperature followed by resuspension in 50 µl of 10mM TrisCl, 1mM EDTA (pH 8.0).

[0347] Five µl of total algal DNA, prepared as described above, was diluted 1:50 in 10 mM Tris, pH 8.0. PCR reactions, final volume 20 µl, were set up as follows. Ten µl of 2 x iProof HF master mix (BIO-RAD) was added to 0.4 µl primer SZ02613 (5'-TGTTGAAGAATGAGCCGGCGAC-3' (SEQ ID NO:9) at 10mM stock concentration). This primer sequence runs from position 567-588 in Gen Bank accession no. L43357 and is highly conserved in higher plants and algal plastid genomes. This was followed by the addition of 0.4 µl primer SZ02615 (5'-CAGTGAGCTATTACGCACTC-3' (SEQ ID NO:10) at 10 mM stock concentration). This primer sequence is complementary to position 1112-1093 in Gen Bank accession no. L43357 and is highly conserved in higher plants and algal plastid genomes. Next, 5 µl of diluted total DNA and 3.2 µl dH_2O were added. PCR reactions were run as follows: 98°C, 45''; 98°C, 8''; 53°C, 12''; 72°C, 20'' for 35 cycles followed by 72°C for 1 min and holding at 25°C. For purification of PCR products, 20 µl of 10 mM Tris, pH 8.0, was added to each reaction, followed by extraction with 40 µl of Phenol:Chloroform:isoamyl alcohol 12:12:1, vortexing and centrifuging at 14,000 x g for 5 minutes. PCR reactions were applied to S-400 columns (GE Healthcare) and centrifuged for 2 minutes at 3,000 x g. Purified PCR products were subsequently TOPO cloned into PCR8/GW/TOPO and positive clones selected for on LB/Spec plates. Purified plasmid DNA was sequenced in both directions using M13 forward and reverse primers. In total, twelve *Prototheca* strains were selected to have their 23S rRNA DNA sequenced and the sequences are listed in the Sequence Listing. A summary of the strains and Sequence Listing Numbers is included below. The sequences were analyzed for overall divergence from the UTEX 1435 (SEQ ID NO: 15) sequence. Two pairs emerged (UTEX 329/UTEX 1533 and UTEX

329/UTEX 1440) as the most divergent. In both cases, pairwise alignment resulted in 75.0% pairwise sequence identity. The percent sequence identity to UTEX 1435 is also included below.

Species	Strain	% nt identity	SEQ ID NO.
Prototheca kruegani	UTEX 329	75.2	SEQ ID NO: 11
Prototheca wickerhamii	UTEX 1440	99	SEQ ID NO: 12
Prototheca stagnora	UTEX 1442	75.7	SEQ ID NO: 13
Prototheca moriformis	UTEX 288	75.4	SEQ ID NO: 14
Prototheca moriformis	UTEX 1439; 1441; 1435; 1437	100	SEQ ID NO: 15
Prototheca wikerhamii	UTEX 1533	99.8	SEQ ID NO: 16
Prototheca moriformis	UTEX 1434	75.9	SEQ ID NO: 17
Prototheca zopfii	UTEX 1438	75.7	SEQ ID NO: 18
Prototheca moriformis	UTEX 1436	88.9	SEQ ID NO: 19

[0348] Lipid samples from a subset of the above-listed strains were analyzed for lipid profile using HPLC. Results are shown below in Table 9.

[0349] Table 9. Diversity of lipid chains in microalgal species

Strain	C14:0	C16:0	C16:1	C18:0	C18:1	C18:2	C18:3	C20:0	C20:1
UTEX 327	0	12.01	0	0	50.33	17.14	0	0	0
UTEX 1441	1.41	29.44	0.70	3.05	57.72	12.37	0.97	0.33	0
UTEX 1435	1.09	25.77	0	2.75	54.01	11.90	2.44	0	0

[0350] Algal plastid transit peptides were identified through the analysis of UTEX 1435 (*Prototheca moriformis*) or UTEX 250 (*Chlorella protothecoides*) cDNA libraries as described in Examples 12 and Example 11 below. cDNAs encoding potentially plastid targeted proteins based upon BLAST hit homology to other known plastid targeted proteins were subjected to further analysis by the sowftawre programs PSORT (http://psort.ims.u-tokyo.ac.jp/form.html), ChloroP (http://www.cbs.dtu.dk/services/ChloroP/) are TargetP (http://www.cbs.dtu.dk/services/TargetP/) . Candidate plastid transit peptides identified through at least one of these three programs were then PCR amplified from the appropriate genomic DNA. Below is a summary of the amino acid sequences algal plastid targeting sequences (PTS) that were identified from this screen. Also included are the amino acid sequences of plant fatty acyl-ACP thioesterases that are used in the heterologous expression Examples below.

cDNA	SEQ ID NO.
P.moriformis isopentenyl diphosphate synthase PTS	SEQ ID NO: 127
P.moriformis delta 12 fatty acid desaturase PTS	SEQ ID NO: 128

P.moriformis stearoyl ACP desaturase PTS	SEQ ID NO: 129
C.protothecoides stearoyl ACP desaturase PTS	SEQ ID NO: 130
Cuphea hookeriana fatty acyl-ACP thioesterase (C8-10)	SEQ ID NO: 131
Umbellularia californica fatty acyl-ACP thioesterase (C12)	SEQ ID NO: 132
Cinnamomum camphora fatty acyl-ACP thioesterase (C14)	SEQ ID NO: 133

[0351] EXAMPLE 2: Culturing *Prototheca* on Various Feedstocks

A. Sorghum

[0352] The following strains were shown to be capable of utilizing sorghum as a sole carbon source: *Prototheca moriformis* strains UTEX 1435, UTEX 1437, UTEX 288, UTEX 1439, UTEX 1441 and UTEX 1434, and *Prototheca stagnora* strain UTEX 1442. The "UTEX" designation indicates the strain number from the algal culture collection of the University of Texas, 1 University State A6700, Austin, Texas 78712-0183.

[0353] Pure sorghum was purchased from Maasdam Sorghum Mills (Lynnville, Iowa) with a sugar profile of fructose 21.0% w/w, dextrose 28.0% w/w, sucrose 16.0% w/w and maltose <0.5%w/w. The cultures were grown in liquid medium containing 2%, 5%, or 7% (v/v) pure sorghum (diluted from the pure stock) as the sole carbon source and the cultures were grown heterotrophically in the dark, agitating at ~ 350 rpm. Samples from the cultures were pulled at 24, 40, 48, 67 and 89 hours and growth was measured using A750 readings on a spectrophotometer. Growth was observed for each of the strains tested as shown in Figures 1-2.

B. Cellulose

[0354] Wet, exploded corn stover, *Miscanthus*, forage sorghum, beet pulp and sugar cane bagasse were prepared by The National Renewable Energy Laboratory (Golden, CO) by cooking in a 1.4% sulfuric acid solution and dewatering the resultant slurry. Percent solids were determined gravimetrically by drying and were as follows: corn stover, 25% solids; *Miscanthus*, 28.7% solids; forage sorghum, 26.7% solids; and sugar cane bagasse, 26% solids.

[0355] 100 gram wet samples of exploded cellulosic materials (corn stover or switch grass) were resuspended in deionized water to a final volume of 420 mL and the pH was adjusted to 4.8 using 10N NaOH. For beet pulp, 9.8 grams dry solids were brought to 350 mL with deionized water and pH was adjusted to 4.8 with 10 N NaOH. For all of the above feedstocks, Accellerase 1000 (Genencor, New York) was used at a ratio of 0.25 ml enzyme per gram of dry biomass for saccharification of the cellulosic materials. Samples were incubated with

agitation (110 rpm) at 50°C for 72 hours. The pH of each of the samples was adjusted to 7.0 with NaOH (with negligible volume change), filter sterilized through a 0.22 μm filter and used in the processes detailed below. For larger scale processes, the same procedure for saccharification was followed except an additional step of tangential flow filtration (TFF) or microfiltration step was performed to aid in filter sterilization of feedstocks. A sample from each of the feedstocks prepared was reserved for determination of glucose and xylose concentration using an HPLC/ELSD-based system or a hexokinase-based kit (Sigma). Additionally, for beet pulp, the material was initially brought to volume as with the other feedstocks, the pH was then adjusted to 4.0 and a pectinase treatment was carried out at 50°C for 24 hours. The pH was then adjusted to 4.8 if no washing steps were conducted or 5.3 if washing steps were conducted. Enzymatic saccharification was then performed with the same procedure used for the other feedstocks as described above.

[0356] Microalgae *Prototheca moriformis* strain UTEX 1435 was assessed for its ability to grow on a series of cellulosic feedstocks prepared as described above (corn stover, beet pulp, sorghum cane, *Miscanthus* and glucose control). The microalgae culture was grown in conditions described in Example 1 above with the exception of the carbon source. The carbon source was either 4% glucose (for control conditions) or 4% glucose as measured by available glucose in the cellulosic materials. Growth was assessed by A750 readings and the culturing time was 168 hours, with A750 readings at 48, 72, 96, 120, 144 and 168 hours after initiation of the culture. As can be seen in Figure 7a, the *Prototheca moriformis* culture grew best in corn stover. The other cellulosic feedstocks used, *Miscanthus*, sorghum cane and beet pulp, all exhibited inhibition of growth.

[0357] Based on the above results with corn stover derived cellulosic sugars, lipid accumulation was also assessed in *Prototheca moriformis* using different levels of corn stover derived cellulosic sugars and reagent glucose as a control. Cultures were grown in 18g/L glucose that was completely from corn stover derived cellulosic sugars (100% corn stover condition in Figure 7b), 9g/L glucose from corn stover derived cellulosic sugars supplemented with 9g/L reagent glucose (50% corn stover supplemented with glucose to 18g/L condition in Figure 7b), 9g/L glucose from corn stover derived cellulosic sugars (50% corn stover, not supplemented; glucose at 9g/L condition in Figure 7b) and a control culture of 42g/L reagent glucose and 13 g/L reagent xylose for osmolarity control. All cultures were fed with cellulosic sugars to maintain the glucose concentration at 20 g/L, except for the control culture, which was fed with reagent glucose to maintain the glucose concentration at 20 g/L. Growth was measured based on the dry cell weight of the culture and lipid

productivity was determined as a percent dry cell weight. Total lipids were determined gravimetrically using an Ankom acid hydrolysis/solvent extraction system as described in Example 1 above.

[0358] As can be seen in Figure 7b, based on biomass accumulation (as measured by DCW), all concentrations of the corn stover derived cellulosics out-performed (higher DCW) the control media that was fed glucose alone. Lipid production as a percentage of DCW was also calculated for all of the conditions. In addition to the higher biomass accumulation seen for growth on corn stover, lipid accumulation was also higher in the corn stover derived cellulosics conditions as compared to the glucose control condition. These data demonstrate that, in addition to providing cellulosic derived sugars, corn stover provides additional nutrients/components that contribute to an increased biomass accumulation (growth) and increased product yield.

[0359] Because the cellulosic feedstocks contain components in addition to glucose, some of these additional components can accumulate to undesirable levels during culture as more cellulosic derived sugars are fed into the culture as the main carbon source (usually, but not limited to, glucose) is consumed. For example, the xylose present in the cellulosic derived sugar feedstock may build up during the high density cultivation of microalgae to levels inhibitory to growth and end product production. To test the effects of xylose build up during *Prototheca* cultivation, cultures were grown with 4% glucose in the media and supplemented with 0, 10 g/L, 25 g/L, 50 g/L and 100 g/L xylose. After 6 days of culture, growth and lipid accumulation were assessed using the methods described above. As seen in Figure 7c, surprisingly, the highest concentrations of xylose tested were not inhibitory to *Prototheca moriformis*' ability to grow and accumulate lipid, and the culture actually grew better and accumulated more lipids at the highest xylose concentrations. To explore this phenomenon, a similar experiment was carried out with sucrose, a carbon source which wild type *Prototheca moriformis* is unable to metabolize. No positive impact was observed with sucrose, suggesting that the increased growth and lipid accumulation seen with xylose is attributable to a mechanism other than the osmotic stress from high concentrations of unmetabolized components in the media and is xylose-specific.

[0360] In addition to non-metabolized sugars, salts may accumulate to inhibitory levels as a result of concentrating lignocellulosic derived sugars. Due to the acid hydrolysis step with H_2SO_4 during the typical preparation of cellulosic materials followed by neutralization of the acid with NaOH, Na_2SO_4 is formed during the generation of lignocellulosic sugars. To assess the impact of salt concentration on growth and lipid production, *Prototheca moriformis*

cultures were grown at Na_2SO_4 concentrations ranging from 0-700 mM in media supplemented with 4% glucose. As shown in Figure 7d, a significant inhibition of growth was observed, as measured by DCW accumulation, where Na_2SO_4 concentrations exceeded 25 mM, specifically at the 80 mM, 240 mM and 700 mM concentrations. In addition, the impact of antifoam P2000 was assessed in the same test. The antifoam compound had a significant, positive impact on biomass productivity. Lipid productivity was also assessed for each condition, and Na2SO4 concentrations above 80 mM, specifically 240 mM and 700 mM, were inhibitory while the addition of antifoam P2000 significantly increased lipid productivity. Thus, in one embodiment, the culturing steps of the methods of the present invention include culturing in media containing an antifoaming agent.

[0361] Based on the results discussed above and summarized in Figure 7a, inhibitors were likely present in the cellulosic feedstocks exhibiting poor growth. The present invention provides means of removing such compounds by washing the materials with hot water (hydrothermal treatment). Figure 8 summarizes the growth results, as measured by A750, using sugar derived from cellulosic feedstock with a single hot water wash. The culture conditions were identical to those used in the processes summarized in Figure 7a. Compared to the results shown in Figure 7a, after just one hot water wash, *Prototheca moriformis* cultures grew better in all cellulosic feedstocks tested, specifically sugar cane bagasse, sorghum cane, *Miscanthus* and beet pulp, as compared to glucose control. Lipid productivity was also assessed in each of the conditions. Except for the beet pulp condition, which was comparable to the glucose control, cultures grown in sugars derived from cellulosic materials subjected to one hot water wash exhibited better lipid productivity than the glucose control.

[0362] One potential impact of hydrothermal treatment (hot water washing) of cellulosic biomass is the removal of furfurals and hydroxymethyl furfurals released by acid explosion of the material. The presence of furfurals and hydroxymethyl furfurals may have contributed to limited growth observed in some of the processes summarized in Figure 7a. To assess how hydrothermal treatment affected the levels of furfurals (FA) and hydroxymethyl furfurals (HMF), supernatants resulting from one to three washes of cellulosic biomass derived from sugarcane bagasse (B), sorghum cane (S), *Miscanthus* (M) or beet pulp (BP) were assayed for FA and HMF by HPLC. As shown in Figure 8, FA and HMF levels decrease significantly with each washing step. This result is consistent with the observation that FA and HMF can be inhibitory to microalgal growth (as seen in Figure 7a) and that hydrothermal treatment removes these compounds and results in improved microalgal growth, even better than the growth in the control glucose conditions (as seen in Figure 8).

[0363] The impact on the lipid profile of *Prototheca moriformis* cultures grown on the various hydrothermally treated lignocellulosic derived sugars was assessed. *Prototheca moriformis* cultures were grown on the following 4x-washed cellulosic feedstocks: *Miscanthus*, sugar cane bagasse and sorghum cane, with glucose levels maintained at 20 g/L through feeding of the cellulosic sugars. At the conclusion of the culturing, microalgae biomass from each condition was analyzed for lipid profile using the methods described in Example 1. The results of the lipid profile analysis (expressed in Area %) are summarized in Table 10 below. Each condition was tested in duplicates, and the results from each of the duplicate test conditions are included. Growth on cellulosic feedstocks resulted in a significant re-distribution in the lipid profile as compared to the glucose control. For example, there was a significant increase in C18:0 Area % in all of the cellulosic feedstock conditions as compared to the glucose control condition.

[0364] Table 10. Lipid profile of *Prototheca moriformis* grown on glucose and cellulosics derived sugars.

	glucose 1 (ctrl)	glucose 2 (ctrl)	bagasse 1	bagasse 2	sorgh 1	sorgh 2	*Miscan* 1	*Miscan* 2
C10:0	n.d.	n.d.	0.03	0.02	n.d.	n.d.	n.d.	n.d.
C12:0	0.04	0.05	0.04	0.04	0.05	0.04	0.04	0.04
C14:0	1.64	1.64	1.07	1.10	1.17	1.14	1.08	1.12
C14:1	0.03	0.04	0.04	0.04	0.06	0.06	0.03	0.03
C15:0	0.04	0.05	0.07	0.05	0.08	0.08	0.06	0.06
C16:0	26.80	26.81	22.32	22.81	22.09	22.19	23.45	23.62
C16:1	0.75	0.82	1.68	1.70	1.92	2.12	1.38	1.23
C17:0	0.14	0.16	0.28	0.17	0.29	0.27	0.21	0.19
C17:1	0.07	0.06	0.10	0.10	0.13	0.12	0.10	0.09
C18:0	3.56	3.64	15.88	10.40	15.30	12.37	10.15	8.69
C18:1	54.22	54.01	49.87	53.87	49.35	50.80	54.05	55.26
C18:2	11.23	11.11	6.54	7.91	7.47	8.80	7.71	7.88
C18:3 alpha	0.84	0.85	0.39	0.56	0.47	0.53	0.56	0.60
C20:0	0.31	0.30	0.85	0.63	0.76	0.69	0.63	0.56
C20:1	0.15	0.15	0.33	0.28	0.32	0.32	0.27	0.25
C20:3	0.06	0.06	0.13	0.12	0.14	0.12	0.11	0.11
C24:0	0.12	0.12	0.22	0.19	0.22	0.20	0.18	0.15

n.d. denotes none detected

[0365] Cellulosic sugar stream was generated from exploded corn stover, saccharified using Accellerase enzyme and concentrated using vacuum evaportation. This sugar stream was tested in *Prototheca moriformis* growth assays at a 4% glucose concentration. The results of the growth assays showed very poor growth and the cellulosic sugar stream was tested for conductivity (salt content). The conductivity was very high, far greater than 700 mM sodium equivalents, a level that was shown to be inhibitory to growth as described above and shown in Figure 7d. Methods of the invention include methods in which salt is reduced or removed

from lignocellulosic derived sugars prior to utilizing these feedstocks in the production of of lignocellulosic derived microalgal oil. Surprisingly, however, one cannot use resins to desalt concentrated sugar streams, one must first dilute the concentrated sugar stream. To demonstrate this embodiment of the invention, cellulosic sugars derived from corn stover material were diluted eight-fold prior to removing contaminating salts with the resin. The initial conductivity of the concentrated starting material was 87 mS/cm while that of the eight-fold diluted stream was 10990 µS/cm at a pH of 5.61. Previous studies had indicated that failure to dilute the concentrated sugar streamprior to de-ionization resulted in an inability to remove salts quantitatively as well as a significant loss of glucose from the sugar stream. Three different bed volumes of IEX resin (DOWEX Marathon MR3) were used (1:2, 1:4 and 1:10). Table 11 summarize results demonstrating the ability of a mixed bed ion exchange (IEX) resin to reduce salts (as measured by conductivity) significantly in a previously concentrated corn stover derived cellulosic sugar stream in diluted feedstocks.

[0366] Table 11. Ability of IEX resin to reduce salts.

Bed volume resin: cellulosics	pH post-deionization	Conductivity post-deionization (µS/cm)	Calculated conductivity post deionization and 8x re-concentration (µS/cm)	Na^+ equivalents (based on std curve) in mM
1:2	3.1	74	592	7.42
1:4	3.1	97	776	9.7
1:10	5.25	6320	50560	634

[0367] A process employing a 1:4 bed volume:cellulosic feedstock and re-concentration of the material eight-fold would result in a sodium concentration is well within the range for normal biomass and lipid accumulation. Alternatively, deionization or salt removal can be performed prior to saccharification or after saccharification, but before concentration of the sugar stream. If salt removal is performed before the concentration of the sugar stream, a dilution step of the sugar stream before salt removal would likely not be necessary.

[0368] This example demonstrates the efficacy of washing of exploded cellulosic material for the use in cellulosic oil production. As described above, concentration of cellulosically derived sugars without the removal of salts (inherent to the production of exploded cellulosic material and subsequent treatment) results in less than optimal fermentations. The materials treated in the process described below were of the appropriate pH for subsequent saccharifaication. In addition, the conductivity of this material was significantly reduced (over 100 fold) from the starting feedstock. Therefore, the subsequenct concentrated sugars to be used in fermentations were not inhibitory due to the presence of excessive salts. An

additional advantage is seen by the removal of furfurals from the cellulosic material. Any xylose or glucose removed in the hemicellulosic fraction can either be discarded or prefereably re-concentrated to be used in fermentations.

[0369] Wet, exploded sugar cane bagasse (NREL, Colorado) with an initial starting mass of 65 kg wet weight and conductivity of 15,000 μS/cm, pH 2.4 was brought to 128 kg with deionized water and the pH adjusted to 4.6 with 10 N NaOH, making the resulting conductivity 6,800 μS/cm). The percent solids were assessed by removal of an aliquot of the suspended materials to a tared (weight = t) aluminum pan, recording the wet weight (weight = w) followed by drying for three hours at 110°C. After drying samples were removed to a desiccator and allowed to come to room temperature (25 °C) at which point, they were weighed again (weight = d). Percent solids were calculated as: % solids = [(d-t/w-t)]x 100. Conductivities were measured on a Thermo Electron Orion 3 Star Conductivity meter.

[0370] The sugar cane bagasse was washed in a semi-continuous fashion by continuously mixing the cellulosic slurry (initial percent solids of 8.2%) at a temperature of 50°C in a stainless steel reactor (150L capacity). Cellulosics were discharged from the reactor vessel via a rotary load pump at a flow rate of 1.9-3.8 kg/min to a Sharples Model 660 decanter centrifuge. Liquid permeate was retained batch wise (ca. 35-175 kg aliquots, see Table 12 below) and homogenous aliquots removed for assessment of total sugars (glucose and xylose) and percent solids as described in Table 12. Conductivity and pH of the cellulosic material were controlled via the addition of de-ionized water and 10 N NaOH, respectively. Samples 1-10 in Table 12 respresent decanted centrifuge permeate, and as such, solids and sugars present in these fractions are removed from the final, washed cellulosic materials. A mass balance calculation of total solids compared to solids removed minus solids lost plus final solids for saccharification, resulted in a 99% recovery in the above process. Figure 8 summarizes the furfural and hydroxymethyl furfurals concentration (mg/L) in each of the 11 centrifuge permeates collected and described in Table 12. These data demonstrate a clear removal of furfurals and hydroxymethyl furfurals from the sugar cane bagasse.

[0371] Table 12. Mass balance for semi-continuous hydrothermal treatment of sugar cane bagasse.

Sample	kg (wet)	kg (dry)	pH	Conductivity μS/cm	total xylose removed (g)	total glucose removed (g)
1 (initial material)	128	10.50	4.60	6,880	0	0
2	81.8	2.03		3,280	1030.68	286.3
3	76.5	0.49		2,500	298.35	76.50
4	106	0.41			254.40	63.60
5	173.9	0.30	3.74	1,260	226.07	69.56

6	101.8	0.08	4.40	791	71.26	20.36
7	110.6	0.04	4.86	327	44.24	0
8	77.2	0			0	0
9	108.6	0.02	4.7	221	0	0
10	101.5	0			0	0
11	34.8	0	4.7	146	0	0
Solids removed (samples 1-10) lost in process		3.37				
Total xylose removed					1925.00	
Total glucose removed						516.32
Final solids for saccharification	7.03					

[0372] In another demonstration of the ability of Prototheca to utilize cellulosic-derived feedstock, *Prototheca moriformis* (UTEX 1435) was cultivated in three-liter bioreactors using cellulosic derived sugar as a fixed carbon feedstock. The inoculum was prepared from cryopreserved cells, which were thawed at room temperature and 1 mL of cells were added to 300 mL of inoculum medium based on the basal microalgae medium described in Example 1 with 1 g/L $(NH_4)_2SO_4$, 4 g/L yeast extract and a trace element solution, plus 4% glucose and grown for 1 day at 28°C with agitation (200 rpm). This culture was used to inoculate a three-liter bioreactor containing 1L medium plus 0.26mL of Antifoam 204 (Sigma, USA). The fermentor was controlled at 28°C and pH was maintained at 6.8 by addition of KOH. Dissolved oxygen was maintained at 30% saturation by cascading agitation and airflow. Cellulosic sugar feedstock from corn stover was fed to the culture to maintain 0-10g/L glucose. Desalination of cellulosic sugar feedstocks to less than 300 mM salt was essential to assure similar dry cell weight and lipid accumulation performance as compared to purified sugar feedstock controls. Desalination of the cellulosic sugar feedstock was performed using the methods described above. Fermentor samples were removed to monitor fermentation performance. Cell mass accumulation was monitored by optical density and dry cell weight. Glucose, xylose, ammonia, potassium, sodium and furfural concentrations were also determined and monitored throughout the fermentation time course. Lipid concentration was determined by gravimetric methods discussed above.

[0373] EXAMPLE 3: Methods for Transforming *Prototheca*

A. <u>General Method for Biolistic transformation of *Prototheca*</u>

[0374] S550d gold carriers from Seashell Technology were prepared according to the protocol from manufacturer. Linearized plasmid (20 μg) was mixed with 50 μl of binding buffer and 60 μl (30 mg) of S550d gold carriers and incubated in ice for 1 min. Precipitation

buffer (100 µl) was added, and the mixture was incubated in ice for another 1 min. After vortexing, DNA-coated particles were pelleted by spinning at 10,000 rpm in an Eppendorf 5415C microfuge for 10 seconds. The gold pellet was washed once with 500 µl of cold 100% ethanol, pelleted by brief spinning in the microfuge, and resuspended with 50 µl of ice-cold ethanol. After a brief (1-2 sec) sonication, 10 µl of DNA-coated particles were immediately transferred to the carrier membrane.

[0375] *Prototheca* strains were grown in proteose medium (2g/L yeast extract, 2.94mM NaNO3, 0.17mM CaCl2•2H2O, 0.3mM MgSO4•7H2O, 0.4mM K2HPO4, 1.28mM KH2PO4, 0.43mM NaCl) on a gyratory shaker until it reaches a cell density of $2x10^6$cells/ml. The cells were harvested, washed once with sterile distilled water, and resuspended in 50 µl of medium. 1 x 10^7 cells were spread in the center third of a non-selective proteose media plate. The cells were bombarded with the PDS-1000/He Biolistic Particle Delivery system (Bio-Rad). Rupture disks (1100 and 1350 psi) were used, and the plates are placed 9 and 12 cm below the screen/macrocarrier assembly. The cells were allowed to recover at 25°C for 12–24 h. Upon recovery, the cells were scraped from the plates with a rubber spatula, mixed with 100 µl of medium and spread on plates containing the appropriate antibiotic selection. After 7-10 days of incubation at 25°C, colonies representing transformed cells were visible on the plates from 1100 and 1350 psi rupture discs and from 9 and 12 cm distances. Colonies were picked and spotted on selective agar plates for a second round of selection.

B. Transformation of *Prototheca* with G418 Resistance Gene

[0376] *Prototheca moriformis* and other *Prototheca* strains sensitive to G418 can be transformed using the methods described below. G418 is an aminoglycoside antibiotic that inhibits the function of 80S ribosomes and thereby inhibits protein synthesis. The corresponding resistance gene fuctions through phosphorylation, resulting in inactivation of G418. *Prototheca* strains UTEX 1435, UTEX 1439 and UTEX 1437 were selected for transformation. All three *Prototheca* strains were genotyped using the methods described above. All three *Prototheca* strains had identical 23s rRNA genomic sequences (SEQ ID NO:15).

[0377] All transformation cassettes were cloned as EcoRI-SacI fragments into pUC19. Standard molecular biology techniques were used in the construction of all vectors according to Sambrook and Russell, 2001. The *C. reinhardtii* beta-tubulin promoter/5'UTR was obtained from plasmid pHyg3 (Berthold *et al.*, (2002) Protist: 153(4), pp 401-412) by PCR as an EcoRI-AscI fragment. The *Chlorella vulgaris* nitrate reductase 3'UTR was obtained

from genomic DNA isolated from UTEX strain 1803 via PCR using the following primer pairs:

Forward:

5' TGACCTAGGTGATTAATTAACTCGAGGCAGCAGCAGCTCGGATAGTATCG 3' (SEQ ID NO:35)

Reverse:

5' CTACGAGCTCAAGCTTTCCATTTGTGTTC CCATCCCACTACTTCC 3' (SEQ ID NO:36)

[0378] The *Chlorella sorokiniana* glutamate dehydrogenase promoter/UTR was otained via PCR of genomic DNA isolated from UTEX strain 1230 via PCR using the following primer pairs:

Forward: 5' GATCAGAATTCCGCCTGCAACGCAAGG GCAGC 3' (SEQ ID NO:37)

Reverse: 5' GCATACTAGTGGCGGGACGGAGAGA GGGCG 3' (SEQ ID NO:38)

[0379] Codon optimization was based on the codons in Table 1 for*Prototheca moriformis*. The sequence of the non-codon optimized neomycin phosphotransferase (nptII) cassette was synthesized as an AscI-XhoI fragment and was based on upon the sequence of Genbank Accession No. YP_788126. The codon optimized nptII cassette was also based on this Genbank Accession number.

[0380] The three *Prototheca* strains were transformed using biolistic methods described above. Briefly, the *Prototheca* strains were grown heterophically in liquid medium containing 2% glucose until they reached the desired cell density ($1x10^7$ cells/mL to $5x10^7$ cells/mL). The cells were harvested, washed once with sterile distilled water and resuspended at $1x10^8$ cells/mL. 0.5mL of cells were then spread out on a non-selective solid media plate and allowed to dry in a sterile hood. The cells were bombarded with the PDS-1000/He Biolistic Particle Delivery System (BioRad). The cells were allowed to recover at 25° C for 24 hours. Upon recovery, the cells were removed by washing plates with 1 mL of sterile media and transferring to fresh plates containing 100 µg/mL G418. Cells were allowed to dry in a sterile hood and colonies were allowed to form on the plate at room temperature for up to three weeks. Colonies of UTEX 1435, UTEX 1439 and UTEX 1437 were picked and spotted on selective agar plates for a second round of selection.

[0381] A subset of colonies that survived a second round of selection described above, were cultured in small volume and genomic DNA and RNA were extracted using standard molecular biology methods. Southern blots were done on genomic DNA extracted from untransformed (WT), the transformants and plasmid DNA. DNA from each sample was run

on 0.8% agarose gels after the following treatments: undigested (U), digested with AvrII (A), digested with NcoI (N), digested with SacI (S). DNA from these gels was blotted on Nylon+ membranes (Amersham). These membranes were probed with a fragment corresponding to the entire coding region of the nptII gene (NeoR probe). Figure 4 shows maps of the cassettes used in the transformations. Figure 5 shows the results of Southern blot analysis on three transformants (all generated in UTEX strain 1435) (1, 2, and 3) transformed with either the beta-tubulin::neo::nit (SEQ ID NO: 39) (transformants 1 and 2) or glutamate dehydrogenase:neo:nit (SEQ ID NO: 40) (transformant 3). The glutamate dehydrogenase:neo:nit transforming plasmid was run as a control and cut with both NcoI and SacI. AvrII does not cut in this plasmid. Genomic DNA isolated from untransformed UTEX strain 1435 shows no hybridization to the NeoR probe.

[0382] Additional transformants containing the codon-optimized glutamate dehydrogenase:neo:nit (SEQ ID NO: 41) and codon-optimized β-tubulin::neo::nit (SEQ ID NO:42) constructs were picked and analyzed by Southern blot analysis. As expected, only digests with SacI show linearization of the transforming DNA. These transformation events are consistent with integration events that occur in the form of oligomers of the transforming plasmid. Only upon digestion with restriction enzymes that cut within the transforming plasmid DNA do these molecules collapse down the size of the transforming plasmid.

[0383] Southern blot analysis was also performed on transformants generated upon transformation of *Prototheca* strains UTEX 1437 and UTEX 1439 with the glutamate dehydrogenase::neo::nit cassette. The blot was probed with the NeoR probe probe and the results are similar to the UTEX 1435 transformants. The results are indicative of integration events characterized by oligomerization and integration of the transforming plasmid. This type of integration event is known to occur quite commonly in *Dictyostelium discoideum* (see, for example, Kuspa, A. and Loomis, W. (1992) *PNAS*, 89:8803-8807 and Morio *et al.*, (1995) *J. Plant Res.* 108:111-114).

[0384] To further confirm expression of the transforming plasmid, Northern blot analysis and RT-PCR analysis were performed on selected transformants. RNA extraction was performed using Trizol Reagent according to manufacturer's instructions. Northern blot analysis were run according to methods published in Sambrook and Russel, 2001. Total RNA (15 μg) isolated from five UTEX 1435 transformants and untransformed UTEX 1435 (control lanes) was separated on 1% agarose-formaldehyde gel and blotted on nylon membrane. The blot was hybridized to the neo-non-optimized probe specific for transgene sequences in transformants 1 and 3. The two other transformants RNAs express the codon-optimized

version of the neo-transgene and, as expected, based on the sequence homology between the optimized and non-optimized neo genes, showed significantly lower hybridization signal.

[0385] RNA (1 μg) was extracted from untransformed *Prototheca* strain UTEX 1435 and two representative UTEX 1435 transformants and reverse transcribed using an oligio dT primer or a gene specific primer. Subsequently these cDNAs (in duplicate) were subjected to qPCR analysis on ABI Veriti Thermocycler using SYBR-Green qPCR chemistry using the following primers (nptII):

Forward: 5' GCCGCGACTGGCTGCTGCTGG 3' (SEQ ID NO:43)

Reverse: 5' AGGTCCTCGCCGTCGGGCATG 3' (SEQ ID NO:44)

[0386] Possible genomic DNA contamination was ruled out by a no reverse transcriptase negative control sample. The results indicated that the NeoR genes used to transform these strains is actively transcribed in the transformants.

C. Transformation of *Prototheca* with Secreted Heterologous Sucrose Invertase

[0387] All of the following experiments were performed using liquid medium/agar plates based on the basal medium described in Ueno et al., (2002) *J Bioscience and Bioengineering* 94(2):160-65, with the addition of trace minerals described in US Patent No. 5,900,370, and 1x DAS Vitamin Cocktail (1000x solution): tricine: 9g, thiamine HCL: 0.67g, biotin: 0.01g, cyannocobalamin (vitamin B12): 0.008g, calcium pantothenate: 0.02g and p-aminobenzoic acid: 0.04g).

[0388] Two plasmid constructs were assembled using standard recombinant DNA techniques. The yeast sucrose invertase genes (one codon optimized and one non-codon optimized), suc2, were under the control of the *Chlorella reinhardtii* beta-tubulin promoter/5'UTR and had the *Chlorella vulgaris* nitrate reductase 3'UTR. The sequences (including the 5'UTR and 3'UTR sequences) for the non-codon optimized (Crβ-tub::NCO-suc2::CvNitRed) construct, SEQ ID NO: 57, and codon optimized (Crβ-tub::CO-suc2::CvNitRed) construct, SEQ ID NO: 58, are listed in the Sequence Listing. Codon optimization was based on Table 1 for *Prototheca sp.* Figure 6 shows a schematic of the two constructs with the relevant restriction cloning sites and arrows indicating the direction of transcription. Selection was provided by Neo R (codon optimized using Table 1).

[0389] Preparation of the DNA/gold microcarrier: DNA/gold microcarriers were prepared immediately before use and stored on ice until applied to macrocarriers. The plasmid DNA (in TE buffer) was added to 50 μl of binding buffer. Saturation of the gold beads was achieved at 15 μg plasmid DNA for 3 mg gold carrier. The binding buffer and DNA were

mixed well via vortexing. The DNA and binding buffer should be pre-mix prior to gold addition to ensure uniformed plasmid binding to gold carrier particles. 60 µl of S550d (Seashell Technologies, San Diego, CA) gold carrier was added to the DNA/binding buffer mixture. For a gold stock at 50 mg/ml, addition of 60 µl results in an optimal ratio of 15 µg DNA/3 mg gold carrier. The gold carrier/DNA mixture was allowed to incubate on ice for 1 minute and then 100 µl of precipitation buffer was added. The mixture was allowed to incubate again on ice for 1 minute and then briefly vortexed and centrifuged at 10,000 rpm at room temperature for 10 seconds to pellet the gold carrier. The supernatant was carefully removed with a pipette and the pellet was washed with 500 µl of ice cold 100% ethanol. The gold particles were re-pelleted by centrifuging again at 10,000 rpm for 10 seconds. The ethanol was removed and 50 µl of ice cold ethanol was added to the gold mixture. Immediately prior to applying the gold to macrocarriers, the gold/ethanol was resuspended with a brief 1-2 second pulse at level 2 on a MISONIX sonicator using the micro tip. Immediately after resuspension, 10 µl of the dispersed gold particles was transferred to the macrocarrier and allowed to dry in a sterile hood.

[0390] The two *Prototheca moriformis* strains (UTEX 1435 and 1441) were grown heterotrophically in liquid medium containing 2% glucose from cryopreserved vials. Each strain was grown to a density of 10^7 cells/ml. This seed culture was then diluted with fresh media to a density of 10^5 cells/ml and allowed to grow for 12-15 hours to achive a final cell density of approximately 10^6 cells/ml. The microalgae were aliquoted into 50 ml conical tubes and centrifuged for 10 minutes at 3500 rpm. The cells were washed with fresh medium and centrifuged again for 10 minutes at 3500 rpm. The cells were then resuspended at a density of 1.25 x 10^8 cells/ml in fresh medium.

[0391] In a sterile hood, 0.4ml of the above-prepared cells were removed and placed directly in the center of an agar plate (without selection agent). The plate was gently swirled with a level circular motion to evenly distribute the cells to a diameter of no more than 3cm. The cells were allowed to dry onto the plates in the sterile hood for approximately 30-40 minutes and then were bombarded at a rupture disk pressure of 1350 psi and a plate to macrocarrier distance of 6 cm. The plates were then covered and wrapped with parafilm and allowed to incubate under low light for 24 hours.

[0392] After the 24 hour recovery, 1ml of sterile medium (with no glucose) was added to the lawn of cells. The cells were resuspended using a sterile loop, applied in a circular motion to the lawn of cells and the resuspended cells were collected using a sterile pipette. The cells were then plated onto a fresh agar plate with 2% glucose and 100 µg/ml G418. The

appearance of colonies occurred 7-12 days after plating. Individual colonies were picked and grown in selective medium with 2% glucose and 100 μg/ml G418. The wildtype (untransformed) and transgenic cells were then analyzed for successful introduction, integration and expression of the transgene.

[0393] Genomic DNA from transformed *Prototheca moriformis* UTEX 1435 and 1441 and their wildtype (untransformed) counterparts were isolated using standard methods. Briefly, the cells were centrifuged for 5 minutes at 14,000 rpm in a standard table top Eppendorf centrifuge (model 5418) and flash frozen prior to DNA extraction. Cell pellets were lysed by adding 200 uL of Lysis buffer (100 mM Tris HCl, pH 8.0, 1% Lauryl Sarcosine, 50 mM NaCl, 20 mM EDTA, 0.25 M sucrose, 0.5 mg/ml RNase A) for every 100-200 mg of cells (wet weight) and vortexing for 30-60 seconds. Cetyl trimethyammonium bromide (CTAB) and NaCl were brought to 1% and 1 M, respectively, and cell extracts were incubated at 60-65°C for 10 minutes. Subsequently, extracts were clarified via centrifugation at 14,000 rpm for 10 minutes and the resulting supernatant was extracted with an equal volume of phenol/chloroform/isoamyl alcohol (25:24:1). Samples were then centrifuged for 5 minutes at 14,000 rpm and the aqueous phase removed. DNA was precipitated with 0.7 volumes of isopropanol. DNA was pelleted via centrifugation at 14,000 rpm for 10 minutes and washed twice with 80% ethanol, and once with ethanol. After drying, DNA was resuspended in 10 mM Tris HCl, pH 8.0 and DNA concentrations were determined by using PicoGreen fluorescence quantification assay (Molecular Probes).

[0394] RNA from transformed *Prototheca moriformis* UTEX 1435 and 1441 and their wildtype (untransformed) counterparts were isolated using standard methods. Briefly, the cells were centrifuged for 5 minutes at 14,000 rpm in a standard table top Eppendorf centrifuge (model 5418) and flash frozen before RNA extraction. Cell pellets were lysed by addition of 1 mL of Trizol reagent (Sigma) for every 100 mg of cells (wet weight) and by vortexing for 1-2 minutes. Samples were incubated at room temperature for 5 minutes and subsequently adjusted with 200 uL of chloroform per 1 mL of Trizol reagent. After extensive shaking, cells were incubated at room temperature for 15 minutes and then subjected to centrifugation at 14000 rpm for 15 minutes in a refrigerated table top microcentrifuge. RNA partitioning to the upper aqueous phase was removed and precipitated by addition of isopropanol (500 uL per 1ml of Trizol reagent). RNA was collected by centrifugation for 10 minutes and the resulting pellet washed twice with 1 mL of 80% ethanol, dried, and resuspended in RNAse free water. RNA concentration was estimated by RiboGreen fluorescence quantification assay (Molecular Probes).

[0395] Expression of neomycin phophotransferase gene conferring G418 antibotic resistance and yeast invertase was assayed in non-transformed *Prototheca moriformis* UTEX 1435 and 1441 and transformants T98 (UTEX 1435 transformant) and T97 (UTEX 1441 transformant) using reverse transcription quantitative PCR analysis (RT-qPCR) . 20 ng total RNA (isolated as described above) was subjected to one step RT-qPCR analysis using iScript SYBR Green RT-PCR kit (BioRad Laboratories) and primer pairs targeting the neomycin resistance gene (forward primer 5'CCGCCGTGCTGGACGTGGTG 3' and reverse primer 5' GGTGGCGGGGTCCAGGGTGT 3'; SEQ ID NOs: 65 and 66, respectively) and *suc2* invertase transcripts (forward primer 5' CGGCCGGCGGCTCCTTCAAC 3' and reverse primer 5' GGCGCTCCCGTAGGTCGGGT 3'; SEQ ID NO: 67 and 68, respectively). Endogenous beta-tubulin transcripts served as an internal positive control for PCR amplification and as a normalization reference to estimate relative transcript levels.

[0396] Both codon optimized and non-codon optimized constructs were transformed into UTEX 1435 and 1441 *Prototheca moriformis* cells as described above. Initially, transformants were obtained with both constructs and the presence of the transgene was verified by Southern blot analysis followed by RTPCR to confirm the presence of the DNA and mRNA from the transgene. For the Southern blot analysis, genomic DNA isolated as described above was electrophoresed on 0.7% agarose gels in 1x TAE buffer. Gells were processed as described in Sambrook et al. (Molecular Cloning; A Laboratory Manual, 2nd Edition. Cold Spring Harbor Laboratory Press, 1989). Probes were prepared by random priming and hybridizations carried out as described in Sambrook et al. Transformants from both the codon optimized and the non-codon optimized constructs showed the presence of the invertase cassette, while the non-transformed control was negative. Invertase mRNA was also detected in transformants with both the codon optimized and non-codon optimized constructs.

[0397] To confirm that the transformants were expressing an active invertase protein, the transformants were plated on sucrose plates. The transformants containing the non-codon optimized cassette failed to grow on the sucrose containing plates, indicating that, while the gene and the mRNA encoding the SUC2 protein were present, the protein was either (1) not being translated, or (2) being translated, but not accumulating to levels sufficient to allow for growth on sucrose as the sole carbon source. The transformants with the codon optimized cassette grew on the sucrose containing plates. To assess the levels of invertase being expressed by these transformants, two clones (T98 and T97) were subjected to an invertase assay of whole cells scraped from solid medium and direct sampling and quantitation of

sugars in the culture supernatants after 48 hours of growth in liquid medium containing 2% sucrose as the sole carbon source.

[0398] For the invertase assay, the cells (T98 and T97) were grown on plates containing 2% sucrose, scraped off and assyed for invertase activity. 10 µl of the scraped cells was mixed with 40 µl of 50mM NaOAc pH 5.1. 12.5 µl of 0.5M sucrose was added to the cell mixture and incubated at 37°C for 10-30 minutes. To stop the reaction, 75 µl of 0.2M K_2HPO_4 was added. To assay for glucose liberated, 500 µl of reconstituted reagent (glucose oxidase/peroxidase + o-Dianisidine) from Sigma (GAGO-20 assay kit) was added to each tube and incubated at 37°C for 30 minutes. A glucose standard curve was also created at this time (range: 25 µg to 0.3 µg glucose). After incubation, 500 µl of 6N HCl was added to stop the reaction and to develop the color. The samples were read at 540nm. The amount of glucose liberated was calculated from the glucose standard curve using the formula y=mx+c, where y is the 540nm reading, and x is µg of glucose. Weight of glucose was converted to moles of glucose, and given the equimolar relationship between moles of sucrose hydrolyzed to moles of glucose generated, the data was expressed as nmoles of sucrose hydrolyzed per unit time. The assay showed that both T98 and T97 clones were able to hydrolyze sucrose, indicating that a functional sucrose invertase was being produced and secreted by the cells.

[0399] For the sugar analysis on liquid culture media after 48 hours of algal growth, T97 and T98 cells were grown in 2% sucrose containing medium for 48 hours and the culture media were processed for sugar analysis. Culture broths from each transformant (and negative non-transformed cell control) were centrifuged at 14,000 rpm for 5 minutes. The resulting supernatant was removed and subjected to HPLC/ELSD (evaporative light scattering detection). The amount of sugar in each sample was determined using external standards and liner regression analysis. The sucrose levels in the culture media of the transformants were very low (less than 1.2g/L, and in most cases 0g/L). In the negative controls, the sucrose levels remained high, at approximately 19g/L after 48 hours of growth.

[0400] These results were consistant with the invertase activity results, and taken together, indicated that the codon optimized transformants, T97 and T98, secreted an active sucrose invertase that allowed the microalgae to utilize sucrose as the sole carbon source in contrast to (1) the non-codon optimized transformants and (2) the non-transformed wildtype microalgae, both of which could not utilize sucrose as the sole carbon source in the culture medium.

[0401] *Prototheca moriformis* strains, T98 and T97, expressing a functional, secreted sucrose invertase (SUC2) transgene were assayed for growth and lipid production using sucrose as the sole carbon source.

[0402] Wild type (untransformed), T98 and T97 strains were grown in growth media (as described above) containing either 4% glucose or 4% sucrose as the sole carbon source under heterotrophic conditions for approximately 6 days. Growth, as determined by A750 optical density readings were taken of all four samples every 24 hours and the dry cell weight of the cultures and lipid profiles were determined after the 6 days of growth. The optical density readings of the transgenic strains grown in both the glucose and sucrose conditions were comparable to the wildtype strains grown in the glucose conditions. These results indicate that the transgenic strains were able to grow on either glucose or sucrose as the sole carbon source at a rate equal to wildtype strains in glucose conditions. The non-transformed, wildtype strains did not grow in the sucrose-only condition.

[0403] The biomass for the wildtype strain grown on glucose and T98 strain grown on sucrose was analyzed for lipid profile. Lipid samples were prepared from dried biomass (lyophilized) using an Acid Hydrolysis System (Ankom Technology, NY) according to manufacturer's instructions. Lipid profile determinations were carried as described in Example 4. The lipid profile for the non-transformed *Prototheca moriformis* UTEX 1435 strain, grown on glucose as the sole carbon source and two colonal T98 strains (UTEX 1435 transformed with a sucrose invertase transgene), grown on sucrose as the sole carbon source, are disclosed in Table 13 (wildtype UTEX 1435 and T98 clone 8 and clone 11 below. C:19:0 lipid was used as an internal calibration control.

[0404] Table 13. Lipid profile of wildtype UTEX 1435 and UTEX 1435 clones with *suc2* transgene.

Name	wildtype (Area % - ISTD)	T98 clone 11 (Area % - ISTD)	T98 clone 8 (Area % - ISTD)
C 12:0	0.05	0.05	0.05
C 14:0	1.66	1.51	1.48
C 14:1	0.04	nd	nd
C 15:0	0.05	0.05	0.04
C 16:0	27.27	26.39	26.50
C 16:1	0.86	0.80	0.84
C 17:0	0.15	0.18	0.14
C 17:1	0.05	0.07	0.05
C 18:0	3.35	4.37	4.50
C 18:1	53.05	54.48	54.50
C 18:2	11.79	10.33	10.24
C 19:0 (ISTD)	--	--	--
C 18:3 alpha	0.90	0.84	0.81
C 20:0	0.32	0.40	0.38

C 20:1	0.10	0.13	0.12
C 20:1	0.04	0.05	0.04
C 22:0	0.12	0.16	0.12
C 20:3	0.07	0.08	0.07
C 24:0	0.12	0.11	0.10
nd - denotes none detected			

[0405] Oil extracted from wildtype *Prototheca moriformis* UTEX 1435 (via solvent extraction or using an expeller press (see methods in Example 44 above) was analyzed for carotenoids, chlorophyll, tocopherols, other sterols and tocotrienols. The results are summarized below in Table 14.

[0406] Table 14. Carotenoid, chlorophyll, tocopherol/sterols and tocotrienol analysis in oil extracted from *Prototheca moriformis* (UTEX 1435).

	Pressed oil (mcg/ml)	Solvent extracted oil (mcg/ml)
cis-Lutein	0.041	0.042
trans-Lutein	0.140	0.112
trans-Zeaxanthin	0.045	0.039
cis-Zeaxanthin	0.007	0.013
t-alpha-Crytoxanthin	0.007	0.010
t-beta-Crytoxanthin	0.009	0.010
t-alpha-Carotene	0.003	0.001
c-alpha-Carotene	none detected	none detected
t-beta-Carotene	0.010	0.009
9-cis-beta-Carotene	0.004	0.002
Lycopene	none detected	none detected
Total Carotenoids	**0.267**	**0.238**
Chlorophyll	<0.01 mg/kg	<0.01 mg/kg
Tocopherols and Sterols		
	Pressed oil (mg/100g)	Solvent extracted oil (mg/100g)
gamma Tocopherol	0.49	0.49
Campesterol	6.09	6.05
Stigmasterol	47.6	47.8
Beta-sitosterol	11.6	11.5
Other sterols	445	446
Tocotrienols		
	Pressed oil (mg/g)	Solvent extracted oil (mg/g)
alpha Tocotrienol	0.26	0.26
beta Tocotrienol	<0.01	<0.01
gamma Tocotrienol	0.10	0.10
detal Tocotrienol	<0.01	<0.01
Total Tocotrienols	**0.36**	**0.36**

[0407] The ability of using sucrose as the sole carbon source as the selection factor for clones containing the *suc2* transgene construct instead of G418 (or another antibiotic) was assessed using the positive *suc2* gene transformants. A subset of the positive transformants was grown on plates containing sucrose as the sole carbon source and without antibiotic

selection for 24 doublings. The clones were then challenged with plates containing glucose as the sole carbon source and G418. There was a subset of clones that did not grow on the glucose + G418 condition, indicating a loss of expression of the transgene. An additional experiment was performed using a plate containing sucrose as the sole carbon source and no G418 and streaking out a *suc2* transgene expressing clone on one half of the plate and wild-type *Prototheca moriformis* on the other half of the plate. Growth was seen with both the wild-type and transgene-containing *Prototheca moriformis* cells. Wild-type *Prototheca moriformis* has not demonstrated the ability to grow on sucrose, therefore, this result shows that unlike antibiotic resistance, the use of sucrose/invertase selection is not cell-autonomous. It is very likely that the transformants were secreting enough sucrose invertase into the plate/media to support wildtype growth as the sucrose was hydrolyzed into fructose and glucose.

[0408] EXAMPLE 4: Recombinant *Prototheca* with Exogenous TE Gene

[0409] As described above, *Prototheca* strains can be transformed with exogenous genes. *Prototheca moriformis* (UTEX 1435) was transformed, using methods described above, with either *Umbellularia californica* C12 thioesterase gene or *Cinnamomum camphora* C14 thiotesterase gene (both codon optimized according to Table 1). Each of the transformation constructs contained a *Chlorella sorokiniana* glutamate dehydrogenase promoter/5'UTR region (SEQ ID NO: 69) to drive expression of the thioesterase transgene. The thioesterase transgenes coding regions of *Umbellularia californica* C12 thioesterase (SEQ ID NO: 70) or *Cinnamomum camphora* C14 thioesterase (SEQ ID NO: 71), each with the native putative plastid targeting sequence. Immediately following the thioesterase coding sequence is the coding sequence for a c-terminal 3x-FLAG tag (SEQ ID NO: 72), followed by the *Chlorella vulgaris* nitrate reductase 3'UTR (SEQ ID NO: 73). A diagram of the thioesterase constructs that were used in the *Prototheca moriformis* transformations is shown in Figure 9.

[0410] Preparation of the DNA, gold microcarrier and *Prototheca moriformis* (UTEX 1435) cells were perfomed using the methods described above in Example 3. The microalgae were bombarded using the gold microcarrier—DNA mixture and plated on selection plates containing 2% glucose and 100μg/ml G418. The colonies were allowed to develop for 7 to 12 days and colonies were picked from each transformation plate and screened for DNA construct incorporation using Southern blots assays and expression of the thioesterase constructs were screened using RT-PCR.

[0411] Positive clones were picked from both the C12 and C14 thioesterase transformation plates and screened for construct incorporation using Southern blot assays. Southern blot

assays were carried out using standard methods (and described above in Example 3) using an optimized c probes, based on the sequence in SEQ ID NO: 70 and SEQ ID NO: 71. Transforming plasmid DNA was run as a positive control. Out of the clones that were positive for construct incorporation, a subset was selected for reverse transcription quantitative PCR (RT-qPCR) analysis for C12 thioesterase and C14 thioesterase expression.

[0412] RNA isolation was performed using methods described in Example 3 above and RT-qPCR of the positive clones were performed using 20 ng of total RNA from each clone using the below-described primer pair and iScript SYBR Green RT-PCR kit (Bio-Rad Laboratories) according to manufacturer's protocol. Wildtype (non-transformed) *Prototheca moriformis* total RNA was included as a negative control. mRNA expression was expreesed as relative fold expression (RFE) as compared to negative control. . The primers that were used in the C12 thioesterase transformation RT-qPCR screening were:

[0413] *U. californica* C12 thioesterase PCR primers:

Forward: 5' CTGGGCGACGGCTTCGGCAC 3' (SEQ ID NO: 74)

Reverse: 5' AAGTCGCGGCGCATGCCGTT 3' (SEQ ID NO: 75)

[0414] The primers that were used in the C14 thioesterase transformation RT-qPCR screening were:

[0415] *Cinnamomum camphora* C14 thioesterase PCR primers:

Forward: 5' TACCCCGCCTGGGGCGACAC 3' (SEQ ID NO: 76)

Reverse: 5' CTTGCTCAGGCGGCGGGTGC 3' (SEQ ID NO: 77)

[0416] RT-qPCR results for C12 thioesterase expression in the positive clones showed an increased RFE of about 40 fold to over 2000 fold increased expression as compared to negative control. Similar results were seen with C14 thioesterase expression in the positive clones with an increase RFE of about 60-fold to over 1200 fold increased expression as compared to negative control.

[0417] A subset of the positive clones from each transformation (as screened by Southern blotting and RT-qPCR assays) were selected and grown under nitrogen-replete conditions and analyzed for total lipid production and profile. Lipid samples were prepared from dried biomass from each clone. 20-40 mg of dried biomass from each transgenic clone was resuspended in 2 mL of 3% H_2SO_4 in MeOH, and 200ul of toluene containing an appropriate amount of a suitable internal standard (C19:0) was added. The mixture was sonicated briefly to disperse the biomass, then heated at 65-70°C for two hours. 2 mL of heptane was added to extract the fatty acid methyl esters, followed by addition of 2 mL of 6% K_2CO_3 (aq) to neutralize the acid. The mixture was agitated vigorously, and a portion of the upper layer

was transferred to a vial containing Na_2SO_4 (anhydrous) for gas chromatography analysis using standard FAME GC/FID (fatty acid methyl ester gas chromatography flame ionization detection) methods. Lipid profile (expressed as Area %) of the positive clones as compared to wildtype negative control are summarized in Tables 15 and 16 below. As shown in Table 15, the fold increase of C12 production in the C12 transformants ranged from about a 5-fold increase (clone C12-5) to over 11-fold increase (clone C12-1). Fold increase of C14 production in the C14 transformants ranged from about a 1.5 fold increase to about a 2.5 fold increase.

[0418] Table 15. Summary of total lipid profile of the *Prototheca moriformis* C12 thioesterase transformants.

	Wildtype	C12-1	C12-2	C12-3	C12-4	C12-5	C12-6	C12-7	C12-8
C6:0	0.03	nd	nd	nd	nd	nd	nd	nd	nd
C8:0	0.11	0.09	nd	0.11	nd	nd	nd	nd	nd
C10:0	nd	nd	nd	0.01	0.01	nd	nd	0.01	nd
C12:0	0.09	1.04	0.27	0.72	0.71	0.50	0.67	0.61	0.92
C14:0	2.77	2.68	2.84	2.68	2.65	2.79	2.73	2.56	2.69
C14:1	0.01	nd	nd	0.02	nd	nd	nd	0.01	nd
C15:0	0.30	0.09	0.10	0.54	0.19	0.09	0.13	0.97	0.09
C15:1	0.05	nd	nd	0.02	nd	nd	nd	nd	nd
C16:0	24.13	23.12	24.06	22.91	22.85	23.61	23.14	21.90	23.18
C16:1	0.57	0.62	0.10	0.52	0.69	0.63	0.69	0.49	0.63
C17:0	0.47	0.24	0.27	1.02	0.36	0.17	0.26	2.21	0.19
C17:1	0.08	nd	0.09	0.27	0.10	0.05	0.09	0.80	0.05
C18:0	nd	nd	2.14	1.75	2.23	2.16	2.38	1.62	2.47
C18:1	22.10	23.15	24.61	21.90	23.52	19.30	22.95	20.22	22.85
C18:1	nd	0.33	0.24	nd	nd	0.09	0.09	nd	0.11
C18:2	37.16	34.71	35.29	35.44	35.24	36.29	35.54	36.01	35.31
C18:3 alpha	11.68	11.29	9.26	11.62	10.76	13.61	10.64	11.97	10.81
C20:0	0.15	0.16	0.19	0.16	0.16	0.14	0.18	0.14	0.18
C20:1	0.22	0.17	0.19	0.20	0.21	0.19	0.21	0.20	0.21
C20:2	0.05	nd	0.04	0.05	0.05	0.05	0.04	0.05	0.04
C22:0	nd	nd	nd	0.01	nd	nd	nd	0.02	nd
C22:1	nd	nd	nd	nd	nd	0.01	nd	0.01	nd
C20:3	0.05	nd	0.07	0.06	0.06	0.10	0.07	0.05	0.06
C20:4	nd	nd	nd	nd	nd	0.02	nd	nd	nd
C24:0	nd	nd	0.24	0.01	0.20	0.19	0.19	0.14	0.20

[0419] Table 16. Summary of total lipid profile of the *Prototheca moriformis* C14 thioesterase transformants.

	Wildtype	C14-1	C14-2	C14-3	C14-4	C14-5	C14-6	C14-7
C6:0	0.03	nd	nd	nd	nd	nd	nd	nd
C8:0	0.11	nd	nd	nd	nd	nd	nd	nd
C10:0	nd	0.01	nd	0.01	nd	0.01	nd	nd
C12:0	0.09	0.20	0.16	0.25	0.21	0.19	0.40	0.17
C14:0	2.77	4.31	4.76	4.94	4.66	4.30	6.75	4.02
C14:1	0.01	nd	0.01	nd	nd	0.01	nd	nd
C15:0	0.30	0.43	0.45	0.12	0.09	0.67	0.10	0.33
C15:1	0.05	nd	nd	nd	nd	nd	nd	nd

C16:0	24.13	22.85	23.20	23.83	23.84	23.48	24.04	23.34
C16:1	0.57	0.65	0.61	0.60	0.60	0.47	0.56	0.67
C17:0	0.47	0.77	0.76	0.21	0.19	1.11	0.18	0.54
C17:1	0.08	0.23	0.15	0.06	0.05	0.24	0.05	0.12
C18:0	nd	1.96	1.46	2.48	2.34	1.84	2.50	2.06
C18:1	22.10	22.25	19.92	22.36	20.57	19.50	20.63	22.03
C18:1	nd	nd	nd	nd	nd	nd	0.10	nd
C18:2	37.16	34.97	36.11	34.35	35.70	35.49	34.03	35.60
C18:3 alpha	11.68	10.71	12.00	10.15	11.03	12.08	9.98	10.47
C20:0	0.15	0.16	0.19	0.17	0.17	0.14	0.18	0.16
C20:1	0.22	0.20	0.12	.019	0.19	0.19	0.17	0.20
C20:2	0.05	0.04	0.02	0.03	0.04	0.05	0.03	0.04
C22:0	nd	nd	nd	nd	0.02	0.01	nd	nd
C22:1	nd	0.01	nd	nd	nd	nd	nd	0.01
C20:3	0.05	0.08	0.03	0.06	0.09	0.05	0.05	0.07
C20:4	nd	0.01	nd	nd	nd	nd	0.02	nd
C24:0	nd	0.17	0.14	0.19	0.20	0.16	0.22	0.17

[0420] The above-described experiments indicate the successful transformation of *Prototheca moriformis* (UTEX 1435) with transgene constructs of two different thioesterases (C12 and C14), which involved not only the successful expression of the transgene, but also the correct targeting of the expressed protein to the plastid and a functional effect (the expected change in lipid profile) as a result of the transformation. The same transformation experiment was performed using an expression construct containing a codon-optimized (according to Table 1) *Cuphea hookeriana* C8-10 thioesterase coding region with the native plastid targeting sequence (SEQ ID NO: 78) yielded no change in lipid profile. While the introduction of the *Cuphea hookeriana* C8-10 transgene into *Prototheca moriformis* (UTEX 1435) was successful and confirmed by Southern blot analysis, no change in C8 or C10 fatty acid production was detected in the transformants compared to the wildtype strain.

[0421] EXAMPLE 5: Generation of *Prototheca moriformis* strain with exogenous plant TE with algal plastid targeting sequence

[0422] In order to investigate whether the use of algal chloroplast/plastid targeting sequences would improve medium chain (C8-C14) thioesterase expression and subsequent medium chain lipid production in *Prototheca moriformis* (UTEX 1435), several putative algal plastid targeting sequences were cloned from *Chlorella protothecoides* and *Prototheca moriformis*. Thioesterase constructs based on *Cuphea hookeriana* C8-10 thioesterase, *Umbellularia californica* C12 thioesterase, and *Cinnamomum camphora* C14 thioesterase were made using made with a *Chlorella sorokiniana* glutamate dehydrogenase promoter/5'UTR and a *Chlorella vulgaris* nitrate reductase 3'UTR. The thioesterase coding sequences were modified by removing the native plastid targeting sequences and replacing

them with plastid targeting sequences from the *Chlorella protothecoides* and the *Prototheca moriformis* genomes. The thioesterase expression constructs and their corresponding sequence identification numbers are listed below. Each transformation plasmid also contained a Neo resistance construct that was identical to the ones described in Example 3 above. Additionally, another algal-derived promoter, the *Chlamydomonas reinhardtii* β-tubulin promoter, was also tested in conjunction with the thioesterase constructs. "Native" plastid targeting sequence refers to the higher plant thioesterase plastid targeting sequence. A summary of the constructs used in these experiments is provided below:

Construct Name	Promoter/ 5'UTR	Plastid targeting seq	Gene	3'UTR	SEQ ID NO.
Construct 1	*C. sorokiniana* glutamate dehydrogenase	*C.protothecoides* stearoyl ACP desaturase	*Cuphea hookeriana* C8-10 TE	*C. vulgaris* nitrate reductase	SEQ ID NO: 79
Construct 2	*C. sorokiniana* glutamate dehydrogenase	*P. moriformis* delta 12 fatty acid desaturase	*Cuphea hookeriana* C8-10 TE	*C. vulgaris* nitrate reductase	SEQ ID NO: 80
Construct 3	*C. sorokiniana* glutamate dehydrogenase	*P. moriformis* isopentenyl diphosphate synthase	*Cuphea hookeriana* C8-10 TE	*C. vulgaris* nitrate reductase	SEQ ID NO: 81
Construct 4	*C. sorokiniana* glutamate dehydrogenase	*P. moriformis* isopentenyl diphosphate synthase	*Umbellularia californica* C12 TE	*C. vulgaris* nitrate reductase	SEQ ID NO: 82
Construct 5	*C. sorokiniana* glutamate dehydrogenase	*P. moriformis* stearoyl ACP desaturase	*Umbellularia californica* C12 TE	*C. vulgaris* nitrate reductase	SEQ ID NO: 83
Construct 6	*C. sorokiniana* glutamate dehydrogenase	*C.protothecoides* stearoyl ACP desaturase	*Umbellularia californica* C12 TE	*C. vulgaris* nitrate reductase	SEQ ID NO: 84
Construct 7	*C. sorokiniana* glutamate dehydrogenase	*P. moriformis* delta 12 fatty acid desaturase	*Umbellularia californica* C12 TE	*C. vulgaris* nitrate reductase	SEQ ID NO: 85
Construct 8	*C. sorokiniana* glutamate dehydrogenase	*C.protothecoides* stearoyl ACP desaturase	*Cinnamomum camphora* C14 TE	*C. vulgaris* nitrate reductase	SEQ ID NO: 86
Construct 9	*Chlamydomonas reinhardtii* β-tubulin	Native	*Cuphea hookeriana* C8-10 TE	*C. vulgaris* nitrate reductase	SEQ ID NO:113
Construct 10	*Chlamydomonas reinhardtii* β-tubulin	*P. moriformis* isopentenyl diphosphate synthase	*Cuphea hookeriana* C8-10 TE	*C. vulgaris* nitrate reductase	SEQ ID NO: 114
Construct 11	*Chlamydomonas*	*P. moriformis*	*Cuphea*	*C. vulgaris*	SEQ ID NO: 115

	reinhardtii β-tubulin	delta 12 fatty acid desaturase	*hookeriana* C8-10 TE	nitrate reductase	
Construct 12	*Chlamydomonas reinhardtii* β-tubulin	*C.protothecoides* stearoyl ACP desaturase	*Cuphea hookeriana* C8-10 TE	*C. vulgaris* nitrate reductase	SEQ ID NO: 116
Construct 13	*Chlamydomonas reinhardtii* β-tubulin	*P. moriformis* stearoyl ACP desaturase	*Cuphea hookeriana* C8-10 TE	*C. vulgaris* nitrate reductase	SEQ ID NO: 117
Construct 14	*Chlamydomonas reinhardtii* β-tubulin	Native	*Umbellularia californica* C12 TE	*C. vulgaris* nitrate reductase	SEQ ID NO: 118
Construct 15	*Chlamydomonas reinhardtii* β-tubulin	*P. moriformis* isopentenyl diphosphate	*Umbellularia californica* C12 TE	*C. vulgaris* nitrate reductase	SEQ ID NO: 119
Construct 16	*Chlamydomonas reinhardtii* β-tubulin	*P. moriformis* delta 12 fatty acid desaturase	*Umbellularia californica* C12 TE	*C. vulgaris* nitrate reductase	SEQ ID NO: 120
Construct 17	*Chlamydomonas reinhardtii* β-tubulin	*C.protothecoides* stearoyl ACP desaturase	*Umbellularia californica* C12 TE	*C. vulgaris* nitrate reductase	SEQ ID NO: 121
Construct 18	*Chlamydomonas reinhardtii* β-tubulin	*P. moriformis* stearoyl ACP desaturase	*Umbellularia californica* C12 TE	*C. vulgaris* nitrate reductase	SEQ ID NO: 122
Construct 19	*Chlamydomonas reinhardtii* β-tubulin	Native	*Cinnamomum camphora* C14 TE	*C. vulgaris* nitrate reductase	SEQ ID NO: 123
Construct 20	*Chlamydomonas reinhardtii* β-tubulin	*P. moriformis* isopentenyl diphosphate synthase	*Cinnamomum camphora* C14 TE	*C. vulgaris* nitrate reductase	SEQ ID NO: 124
Construct 21	*Chlamydomonas reinhardtii* β-tubulin	*P. moriformis* delta 12 fatty acid desaturase	*Cinnamomum camphora* C14 TE	*C. vulgaris* nitrate reductase	SEQ ID NO:
Construct 22	*Chlamydomonas reinhardtii* β-tubulin	*C.protothecoides* stearoyl ACP desaturase	*Cinnamomum camphora* C14 TE	*C. vulgaris* nitrate reductase	SEQ ID NO: 87
Construct 23	*Chlamydomonas reinhardtii* β-tubulin	*P. moriformis* stearoyl ACP desaturase	*Cinnamomum camphora* C14 TE	*C. vulgaris* nitrate reductase	SEQ ID NO: 88

[0423] Each construct was transformed into *Prototheca moriformis* (UTEX 1435) and selection was performed using G418 using the methods described in Example 4 above. Several positive clones from each transformation were picked and screened for the presence thioesterase transgene using Southern blotting analysis. Expression of the thioesterase

transgene was confirmed using RT-PCR. A subset of the positive clones (as confirmed by Southern blotting analysis and RT-PCR) from each transformation was selected and grown for lipid profile analysis. Lipid samples were prepared from dried biomass samples of each clone and lipid profile analysis was performed using acid hydrolysis methods described in Example 4. Changes in area percent of the fatty acid corresponding to the thioesterase transgene were compared to wildtype levels, and clones transformed with a thioesterase with the native plastid targeting sequence.

[0424] As mentioned in Example 4, the clones transformed with *Cuphea hookeriana* C8-10 thioesterase constructs with the native plastid targeting sequence had the same level of C8 and C10 fatty acids as wildtype. The clones transformed with *Cuphea hookeriana* C8-10 thioesterase constructs (Constructs 1-3) with algal plastid targeting sequences had over a 10-fold increase in C10 fatty acids for Construct 3 and over 40-fold increase in C10 fatty acids for Constructs 1 and 2 (as compared to wildtype). The clones transformed with *Umbellularia californica* C12 thioesterase constructs with the native plastid targeting sequence had a modest 6-8 fold increase in C12 fatty acid levels as compared to wildtype. The clones transformed with the *Umbellularia californica* C12 thioesterase constructs with the algal plasid targeting constructs (Constructs 4-7) had over an 80-fold increase in C12 fatty acid level for Construct 4, about an 20-fold increase in C12 fatty acid level for Construct 6, about a 10-fold increase in C12 fatty acid level for Construct 7 and about a 3-fold increase in C12 fatty acid level for Construct 5 (all compared to wildtype). The clones transformed with *Cinnamomum camphora* C14 thioesterase with either the native plastid targeting sequence or the construct 8 (with the *Chlorella protothecoides* stearoyl ACP desaturase plastid targeting sequence) had about a 2-3 fold increase in C14 fatty acid levels as compared to wildtype. In general clones transformed with an algal plastid targeting sequence thioesterase constructs had a higher fold increase in the corresponding chain-length fatty acid levels than when using the native higher plant targeting sequence.

A. *Clamydomonas reinhartii* β-tubulin promoter

[0425] Additional heterologous thioesterase expression constructs were prepared using the *Chlamydomonas reinhardtii* β- tubulin promoter instead of the *C. sorokinana* glutamate dehydrogenase promoter. The construct elements and sequence of the expression constructs are listed above. Each construct was transformed into *Prototheca moriformis* UTEX 1435 host cells using the methods described above. Lipid profiles were generated from a subset of positive clones for each construct in order to assess the success and productivity of each construct. The lipid profiles compare the fatty acid levels (expressed in area %) to wildtype

host cells. The "Mean" column represents the numerical average of the subset of positive clones. The "Sample" column represents the best positive clone that was screened (best being defined as the sample that produced the greatest change in area % of the corresponding chain-length fatty acid production). The "low-high" column represents the lowest area % and the highest area % of the fatty acid from the clones that were screened. The lipid profiles results of Constructs 9-23 are summarized below.

Construct 9. ***Cuphea hookeriana*** **C8-10 TE**				
Fatty Acid	wildtype	Mean	Sample	low/high
C 8:0	0	0.05	0.30	0-0.29
C 10:0	0.01	0.63	2.19	0-2.19
C 12:0	0.03	0.06	0.10	0-0.10
C 14:0	1.40	1.50	1.41	1.36-3.59
C 16:0	24.01	24.96	24.20	
C 16:1	0.67	0.80	0.85	
C 17:0	0	0.16	0.16	
C 17:1	0	0.91	0	
C 18:0	4.15	17.52	3.19	
C 18:1	55.83	44.81	57.54	
C 18:2	10.14	7.58	8.83	
C 18:3α	0.93	0.68	0.76	
C 20:0	0.33	0.21	0.29	
C 24:0	0	0.05	0.11	

Construct 10. ***Cuphea hookeriana*** **C8-10 TE**				
Fatty Acid	wildtype	Mean	Sample	low/high
C 8:0	0	0.01	0.02	0-0.03
C 10:0	0	0.16	0.35	0-0.35
C 12:0	0.04	0.05	0.07	0-0.07
C 14:0	1.13	1.62	1.81	0-0.05
C 14:1	0	0.04	0.04	
C 15:0	0.06	0.05	0.05	
C 16:0	19.94	26.42	28.08	
C 16:1	0.84	0.96	0.96	
C 17:0	0.19	0.14	0.13	
C 17:1	0.10	0.06	0.05	
C 18:0	2.68	3.62	3.43	
C 18:1	63.96	54.90	53.91	
C 18:2	9.62	9.83	9.11	
C18:3 γ	0	0.01	0	
C 18:3α	0.63	0.79	0.73	
C 20:0	0.26	0.35	0.33	
C 20:1	0.06	0.08	0.09	
C 20:1	0.08	0.06	0.07	
C 22:0	0	0.08	0.09	
C 24:0	0.13	0.13	0.11	

Construct 11. ***Cuphea hookeriana*** **C8-10 TE**				
Fatty Acid	wildtype	Mean	Sample	low/high
C 8:0	0	0.82	1.57	0-1.87
C 10:0	0	3.86	6.76	0-6.76

C 12:0	0.04	0.13	0.20	0.03-0.20
C 14:0	1.13	1.80	1.98	1.64-2.05
C 14:1	0	0.04	0.04	
C 15:0	0.06	0.06	0.06	
C 16:0	19.94	25.60	25.44	
C 16:1	0.84	1.01	1.02	
C 17:0	0.19	0.13	0.11	
C 17:1	0.10	0.06	0.05	
C 18:0	2.68	2.98	2.38	
C 18:1	63.96	51.59	48.85	
C 18:2	9.62	9.85	9.62	
C18:3 γ	0	0.01	0	
C 18:3α	0.63	0.91	0.92	
C 20:0	0.26	0.29	0.26	
C 20:1	0.06	0.06	0	
C 20:1	0.08	0.06	0.03	
C 22:0	0	0.08	0.08	
C 24:0	0.13	0.06	0	

Construct 12. *Cuphea hookeriana* C8-10 TE				
Fatty Acid	wildtype	Mean	Sample	low/high
C 8:0	0	0.31	0.85	0-0.85
C 10:0	0	2.16	4.35	0.20-4.35
C 12:0	0.04	0.10	0.15	0-0.18
C 14:0	1.13	1.96	1.82	1.66-2.97
C 14:1	0	0.03	0.04	
C 15:0	0.06	0.07	0.07	
C 16:0	19.94	26.08	25.00	
C 16:1	0.84	1.04	0.88	
C 17:0	0.19	0.16	0.16	
C 17:1	0.10	0.05	0.07	
C 18:0	2.68	3.02	3.19	
C 18:1	63.96	51.08	52.15	
C 18:2	9.62	11.44	9.47	
C18:3 γ	0	0.01	0	
C 18:3α	0.63	0.98	0.90	
C 20:0	0.26	0.30	0.28	
C 20:1	0.06	0.06	0.05	
C 20:1	0.08	0.04	0	
C 22:0	0	0.07	0	
C 24:0	0.13	0.05	0	

Construct 14. *Umbellularia californica* C12 TE				
Fatty Acid	wildtype	Mean	Sample	low/high
C 10:0	0.01	0.02	0.03	0.02-0.03
C 12:0	0.03	2.62	3.91	0.04-3.91
C 14:0	1.40	1.99	2.11	1.83-2.19
C 16:0	24.01	27.64	27.01	
C 16:1	0.67	0.92	0.92	

C 18:0	4.15	2.99	2.87	
C 18:1	55.83	53.22	52.89	
C 18:2	10.14	8.68	8.41	
C 18:3α	0.93	0.78	0.74	
C 20:0	0.33	0.29	0.27	

Construct 15. ***Umbellularia californica*** **C12 TE**				
Fatty Acid	wildtype	Mean	Sample	low/high
C 10:0	0	0.05	0.08	0-0.08
C 12:0	0.04	8.12	12.80	4.35-12.80
C 13:0	0	0.02	0.03	0-0.03
C 14:0	1.13	2.67	3.02	2.18-3.37
C 14:1	0	0.04	0.03	0.03-0.10
C 15:0	0.06	0.07	0.06	
C 16:0	19.94	25.26	23.15	
C 16:1	0.84	0.99	0.86	
C 17:0	0.19	0.14	0.14	
C 17:1	0.10	0.05	0.05	
C 18:0	2.68	2.59	2.84	
C 18:1	63.96	46.91	44.93	
C 18:2	9.62	10.59	10.01	
C 18:3α	0.63	0.92	0.83	
C 20:0	0.26	0.27	0.24	
C 20:1	0.06	0.06	0.06	
C 20:1	0.08	0.05	0.04	
C 22:0	0	0.07	0.09	
C 24:0	0.13	0.13	0.12	

Construct 16. ***Umbellularia californica*** **C12 TE**				
Fatty Acid	wildtype	Mean	Sample	low/high
C 10:0	0	0.03	0.04	0.02-0.04
C 12:0	0.04	2.43	5.32	0.98-5.32
C 13:0	0	0.01	0.02	0-0.02
C 14:0	1.13	1.77	1.93	1.62-1.93
C 14:1	0	0.03	0.02	0.02-0.04
C 15:0	0.06	0.06	0.05	
C 16:0	19.94	24.89	22.29	
C 16:1	0.84	0.91	0.82	
C 17:0	0.19	0.16	0.15	
C 17:1	0.10	0.06	0.06	
C 18:0	2.68	3.81	3.67	
C 18:1	63.96	53.19	52.82	
C 18:2	9.62	10.38	10.57	
C 18:3α	0.63	0.80	0.77	
C 20:0	0.26	0.35	0.32	
C 20:1	0.06	0.06	0.07	
C 20:1	0.08	0.07	0.08	
C 22:0	0	0.08	0.07	
C 24:0	0.13	0.15	0.14	

Construct 17. ***Umbellularia californica*** **C12 TE**				
Fatty Acid	wildtype	Mean	Sample	low/high
C 10:0	0	0.04	0.07	0.03-0.08
C 12:0	0.04	7.02	14.11	4.32-14.11
C 13:0	0	0.03	0.04	0.01-0.04
C 14:0	1.13	2.25	3.01	1.95-3.01
C 14:1	0	0.03	0.03	0.02-0.03
C 15:0	0.06	0.06	0.06	
C 16:0	19.94	23.20	21.46	
C 16:1	0.84	0.82	0.77	
C 17:0	0.19	0.15	0.14	
C 17:1	0.10	0.06	0.06	
C 18:0	2.68	3.47	2.93	
C 18:1	63.96	50.30	45.17	
C 18:2	9.62	10.33	9.98	
C18:3 γ	0	0.01	0	
C 18:3α	0.63	0.84	0.86	
C 20:0	0.26	0.32	0.27	
C 20:1	0.06	0.07	0.06	
C 20:1	0.08	0.06	0.06	
C 22:0	0	0.08	0.09	
C 24:0	0.13	0.14	0.13	

Construct 18. ***Umbellularia californica*** **C12 TE**				
Fatty Acid	wildtype	Mean	Sample	low/high
C 10:0	0	0.03	0.05	0.01-0.05
C 12:0	0.04	5.06	7.77	0.37-7.77
C 13:0	0	0.02	0	0-0.03
C 14:0	1.13	2.11	2.39	1.82-2.39
C 14:1	0	0.03	0.03	0.02-0.05
C 15:0	0.06	0.06	0.06	
C 16:0	19.94	24.60	23.95	
C 16:1	0.84	0.86	0.83	
C 17:0	0.19	0.15	0.14	
C 17:1	0.10	0.06	0.05	
C 18:0	2.68	3.31	2.96	
C 18:1	63.96	51.26	49.70	
C 18:2	9.62	10.18	10.02	
C18:3 γ	0	0.01	0.02	
C 18:3α	0.63	0.86	0.86	
C 20:0	0.26	0.32	0.29	
C 20:1	0.06	0.05	0.05	
C 20:1	0.08	0.07	0.04	
C 22:0	0	0.08	0.08	
C 24:0	0.13	0.13	0.13	

Construct 19. ***Cinnamomum camphora*** **C14 TE**				
Fatty Acid	wildtype	Mean	Sample	low/high
C 10:0	0.02	0.01	0.01	0.01-0.02
C 12:0	0.05	0.27	0.40	0.08-0.41
C 14:0	1.52	4.47	5.81	2.10-5.81
C 16:0	25.16	28.14	28.55	
C 16:1	0.72	0.84	0.82	
C 18:0	3.70	3.17	2.87	
C 18:1	54.28	51.89	51.01	
C 18:2	12.24	9.36	8.62	
C 18:3α	0.87	0.74	0.75	
C 20:0	0.33	0.33	0.31	

Construct 20. ***Cinnamomum camphora*** **C14 TE**				
Fatty Acid	wildtype	Mean	Sample	low/high
C 10:0	0.01	0.01	0.02	0.01-0.02
C 12:0	0.03	0.39	0.65	0.08-0.65
C 13:0	0	0.01	0.01	0.01-0.02
C 14:0	1.40	5.61	8.4	2.1-8.4
C 14:1	0	0.03	0.03	0.02-0.03
C 15:0	0	0.06	0.07	
C 16:0	24.01	25.93	25.57	
C 16:1	0.67	0.75	0.71	
C 17:0	0	0.13	0.12	
C 17:1	0	0.05	0.05	
C 18:0	4.15	3.30	3.23	
C 18:1	55.83	51.00	48.48	
C 18:2	10.14	10.38	10.35	
C 18:3α	0.93	0.91	0.88	
C 20:0	0.33	0.35	0.32	
C 20:1	0	0.08	0.08	
C 20:1	0	0.07	0.07	
C 22:0	0	0.08	0.08	
C 24:0	0	0.14	0.13	

Construct 21. ***Cinnamomum camphora*** **C14 TE**				
Fatty Acid	wildtype	Mean	Sample	low/high
C 10:0	0.01	0.01	0.01	0-0.01
C 12:0	0.03	0.10	0.27	0.04-0.27
C 14:0	1.40	2.28	4.40	1.47-4.40
C 16:0	24.01	26.10	26.38	
C 16:1	0.67	0.79	0.73	
C 17:0	0	0.15	0.16	
C 17:1	0	0.06	0.06	
C 18:0	4.15	3.59	3.51	
C 18:1	55.83	53.53	50.86	
C 18:2	10.14	10.83	11.11	

C 18:3α	0.93	0.97	0.87	
C 20:0	0.33	0.36	0.37	
C 20:1	0	0.09	0.08	
C 20:1	0	0.07	0.07	
C 22:0	0	0.09	0.09	

Construct 22. *Cinnamomum camphora* C14 TE				
Fatty Acid	wildtype	Mean	Sample	low/high
C 10:0	0.01	0.02	0.02	0.02-0.02
C 12:0	0.03	1.22	1.83	0.59-1.83
C 13:0	0	0.02	0.03	0.01-0.03
C 14:0	1.40	12.77	17.33	7.97-17.33
C 14:1	0	0.02	0.02	0.02-0.04
C 15:0	0	0.07	0.08	
C 16:0	24.01	24.79	24.22	
C 16:1	0.67	0.64	0.58	
C 17:0	0	0.11	0.10	
C 17:1	0	0.04	0.04	
C 18:0	4.15	2.85	2.75	
C 18:1	55.83	45.16	41.23	
C 18:2	10.14	9.96	9.65	
C 18:3α	0.93	0.91	0.85	
C 20:0	0.33	0.30	0.30	
C 20:1	0	0.07	0.06	
C 20:1	0	0.06	0.05	
C 22:0	0	0.08	0.08	

Construct 23. *Cinnamomum camphora* C14 TE				
Fatty Acid	wildtype	Mean	Sample	low/high
C 10:0	0.01	0.01	0.02	0-0.02
C 12:0	0.05	0.57	1.08	0.16-1.08
C 13:0	0	0.02	0.02	0-0.02
C 14:0	1.45	7.18	11.24	2.96-11.24
C 14:1	0.02	0.03	0.03	0.02-0.03
C 15:0	0.06	0.07	0.07	
C 16:0	24.13	25.78	25.21	
C 16:1	0.77	0.72	0.66	
C 17:0	0.19	0.13	0.11	
C 17:1	0.08	0.05	0.04	
C 18:0	3.53	3.35	3.12	
C 18:1	56.15	49.65	46.35	
C 18:2	11.26	10.17	9.72	
C 18:3α	0.84	0.95	0.83	
C 20:0	0.32	0.34	0.32	
C 20:1	0.09	0.08	0.09	
C 20:1	0.07	0.05	0.06	
C 22:0	0.07	0.08	0.08	
C 24:0	0.13	0.13	0.12	

[0426] Constructs 9-13 were expression vectors containing the *Cuphea hookeriana* C8-10 thioesterase construct. As can be seen in the data summaries above, the best results were seen with Construct 11, with the Sample C8 fatty acid being 1.57 Area % (as compared to 0 in wildtype) and C10 fatty acid being 6.76 Area % (as compared to 0 in wildtype). There was also a modest increase in C12 fatty acids (approximately 2-5 fold increase). While the native plastid targeting sequence produced no change when under the control of the *C.sorokinana* glutamate dehydrogenase promoter, the same expression construct driven by the *C.reinhardtii* β-tubulin promoter produced significant changes in C8-10 fatty acids in the host cell. This is further evidence of the idiosyncrasies of heterologous expression of thioesterases in *Prototheca* species. All of the clones containing the *C.reinhardtii* β-tubulin promoter C8-10 thioesterase construct had greater increases in C8-10 fatty acids than the clones containing the *C.sorokinana* glutamate dehydrogenase promoter C8-10 thioesterase construct. Lipid profile data for Construct 13 was not obtained and therefore, not included above.

[0427] Constructs 14-18 were expression vectors containing the *Umbellularia californica* C12 thioesterase construct. As can be seen in the data summaries above, the best results were seen with Constructs 15 (*P.moriformis* isopentenyl diphosphate synthase plastid targeting sequence) and 17 (*C.protothecoides* stearoyl ACP desaturase plastid targeting sequence). The greatest change in C12 fatty acid production was seen with Construct 17, with C12 fatty acids levels of up to 14.11 area %, as compared to 0.04 area % in wildtype. Modest changes (about 2-fold) were also seen with C14 fatty acid levels. When compared to the same constructs with the *C.sorokinana* glutamate dehydrogenase promoter, the same trends were true with the *C.reinhardtii* β-tubulin promoter—the *C.protothecoides* stearoyl ACP desaturase and *P.moriformis* isopentenyl diphosphate synthase plastid targeting sequences produced the greatest change in C12 fatty acid levels with both promoters.

[0428] Constructs 19-23 were expression vectors containing the *Cinnamomum camphora* C14 thioesterase construct. As can be seen in the data summaries above, the best results were seen with Constructs 22 and Construct 23. The greatest change in C14 fatty acid production was seen with Construct 22, with C14 fatty acid levels of up to 17.35 area % (when the values for C140 and C141 are combined), as compared to 1.40% in wildtype. Changes in C12 fatty acids were also seen (5-60 fold). When compared to the same constructs with the *C.sorokinana* glutamate dehydrogenase promoter, the same trends were true with the *C.reinhardtii* β-tubulin promoter—the *C.protothecoides* stearoyl ACP desaturase and *P.moriformis* stearoyl ACP desaturase plastid targeting sequences produced the greatest

change in C14 fatty acid levels with both promoters. Consistently with all thioesterase expression constructs, the *C.reinhardtii* β-tubulin promoter constructs produced greater changes in C8-14 fatty acid levels than the *C.sorokiniana* glutamate dehydrogenase

[0429] Two positive clones from the Construct 22 were selected and grown under high selective pressure (50mg/L G418). After 6 days in culture, the clones were harvested and their lipid profile was determined using the methods described above. The lipid profile data is summarized below and is expressed in area %.

Construct 22 clones + 50mg/L G418		
Fatty Acid	Construct 22 A	Construct 22 B
C 12:0	3.21	3.37
C 14:0	27.55	26.99
C 16:0	25.68	24.37
C 16:1	0.99	0.92
C 18:0	1.37	1.23
C 18:1	28.35	31.07
C 18:2	11.73	11.05
C 18:3α	0.92	0.81
C 20:0	0.16	0.17

[0430] Both clones, when grown under constant, high selective pressure, produced an increased amount of C14 and C12 fatty acids, about double the levels seen with Construct 22 above. These clones yielded over 30 area % of C12-14 fatty acids, as compared to 1.5 area % of C12-14 fatty acids seen in wildtype cells.

[0431] EXAMPLE 6: Heterologous Expression of *Cuphea palustris* and *Ulmus americanca* Thioesterase in *Prototheca*

[0432] Given the success of the above-described heterologous expression thioesterases in *Prototheca* species, expression cassettes containing codon-optimized (according to Table 1) sequences encoding fatty acyl-ACP thioesterases from *Cuphea palustris* and *Ulmus americana* were constructed and described below.

Construct Name	Promoter/ 5'UTR	Plastid targeting seq	Gene	3'UTR	SEQ ID NO.
Construct 27	*C.reinhardtii* β-tubulin	*C.protothecoides* stearoyl ACP desaturase	*Cuphea palustris* thioesterase	*C. vulgaris* nitrate reductase	SEQ ID NO: 107

[0433] The *Ulmus americana* (codon-optimized coding sequence) can be inserted into the expression cassette. The codon-optimized coding sequence without the native plastid targeting sequence for the *Ulmus americana* thioesterase is listed as SEQ ID NO: 108 and can be fused any desired plastid targeting sequence and expression element (i.e., promoter/5'UTR and 3'UTR).

[0434] These expression cassettes can be transformed in to *Prototheca* species using the methods described above. Positive clones can be screened with the inclusion of an antibiotic resistance gene (e.g, neoR) on the expression construct and screened on G418-containing plates/media. Positive clones can be confirmed using Southern blot assays with probes specific to the heterologous thioesterase coding region and expression of the construct can also be confirmed using RT-PCR and primers specific to the coding region of the heterologous thioesterase. Secondary confirmation of positive clones can be achieved by looking for changes in levels of fatty acids in the host cell's lipid profile. As seen in the above Examples, heterologous expression in *Prototheca* species of thioesterase can be idiosyncratic to the particular thioesterase. Promoter elements and plastid targeting sequences (and other expression regulatory elements) can be interchanged until the expression of the thioesterase (and the subsequent increase in the corresponding fatty acid) reaches a desired level.

[0435] EXAMPLE 7: Dual Transformants—Simultaneous Expression of Two Heterologous Proteins

[0436] Microalgae strain *Prototheca moriformis* (UTEX 1435) was transformed using the above disclosed methods with a expression construct containing the yeast sucrose invertase *suc2* gene encoding the secreted form of the *S. cerevisiae* invertase. Successful expression of this gene and targeting to the periplasm results in the host cell's ability to grow on (and utilize) sucrose as a sole carbon source in heterotrophic conditions (as demonstrated in Example 3 above). The second set of genes expressed are thioesterases which are responsible for the cleavage of the acyl moiety from the acyl carrier protein. Specifically, thioesterases from *Cuphea hookeriana* (a C8-10 preferring thioesterase), *Umbellularia californica* (a C12 preferring thioesterase), and *Cinnamomum camphora* (a C14 preferring thioesterase). These thioesterase expression cassettes were cloned as fusions with N-termial microalgal plastid targeting sequences from either *Prototheca moriformis* or *Chlorella protothecoides*, which have been shown (in the above Examples) to be more optimal than the native higher plant plastid targeting sequences. The successful expression of the thioesterase genes and the targeting to the plastid resulted in measurable changes in the fatty acid profiles within the host cell. These changes in profiles are consistent with the enzymatic specificity or preference of each thioesterase. Below is a summary of dual expression contructs that were assembled and transformed into *Prototheca moriformis* (UTEX 1435). Each construct contained the yeast *suc2* gene under the control of the *C .reinhardtii* β-tubulin 5'UTR/promoter and contained the *C. vulgaris* nitrate reductase 3'UTR and a higher plant thioesterase with a

microalgal plastid targeting sequence replacing the native sequence under the control of *C. sorokinana* glutamate dehydrogenase 5'UTR and contained the *C. vulgaris* nitrate reductase 3'UTR. Below is a summary of the thioesterase portion of the constructs that were assembled and transformed into *Prototheca moriformis* (UTEX 1435). The entire dual expression cassette with the *suc2* gene and the thioesterase gene and the is listed in the Sequence Identification Listing.

Construct Name	Promoter/ 5'UTR	Plastid targeting seq	Gene	3'UTR	SEQ ID NO.
Construct 24	*C. sorokiniana* glutamate dehydrogenase	*C.protothecoides* stearoyl ACP desaturase	*Cuphea hookeriana* C8-10 TE	*C. vulgaris* nitrate reductase	SEQ ID NO: 109
Construct 25	*C. sorokinana* glutamate dehydrogenase	*P. moriformis* isopentenyl diphosphate synthase	*Umbellularia californica* C12 TE	*C. vulgaris* nitrate reductase	SEQ ID NO: 110
Construct 26	*C. sorokinana* glutamate dehydrogenase	*C.protothecoides* stearoyl ACP desaturase	*Cinnamomum camphora* C14 TE	*C. vulgaris* nitrate reductase	SEQ ID NO: 111

Similar dual expression constructs with the thioesterase cassettes described in Example 5 (e.g., under the control of a different promoter such as *C. reinhardtii* β-tubulin promoter/5'UTR) can also be generated using standard molecular biology methods and methods described herein.

[0437] Positive clones containing each of expression constructs were screened using their ability to grow on sucrose-containing plates, where sucrose is the sole-carbon source, as the selection factor. A subset of these positive clones from each construct transformation was selected and the presence of the expression construct was confirmed using Southern blot assays. The function of the yeast sucrose invertase was also confirmed using a sucrose hydrolysis assay. Positive clones were selected and grown in media containing sucrose as the sole carbon source at a starting concentration of 40g/L. A negative control of wildtype *Prototheca moriformis* (UTEX 1435) grown in media containing glucose as the sole carbon source at the same 40 g/L starting concentration was also included. Utilization of sucrose was measured throughout the course of the experiment by measuring the level of sucrose in the media using a YSI 2700 Biochemistry Analyzer with a sucrose-specific membrane. After six days in culture, the cultures were harvested and processed for lipid profile using the same methods as described above. The lipid profile results are summarized below in Table 17 and are show in area %.

[0438] Table 17. Lipid profiles of dual transformants with *suc2* sucrose invertase and thioesterase.

Fatty Acid	Wt	C24 A	C24 B	C24 C	C25 A	C25 B	C25 C	C26 A	C26 B	C26 C
C 10:0	0.01	**0.03**	**0.04**	**0.08**	0.01	0.01	0.01	0.01	0.01	0.0
C 12:0	0.04	0.04	0.04	0.04	**0.28**	**0.40**	**0.10**	0.04	0.04	0.13
C 14:0	1.6	1.55	1.53	1.56	1.59	1.59	1.60	**1.65**	**1.56**	**2.69**
C 14:1	0.03	0.03	0.03	0.02	0.03	0.03	0.03	0.03	0.03	0.03
C 15:0	0.04	0.03	0.03	0.04	0.04	0.03	0.03	0.03	0.03	0.04
C 16:0	29.2	29.1	29.0	28.6	28.9	28.6	29.0	28.8	29.5	27.5
C 16:1	0.86	0.80	0.79	0.82	0.77	0.81	0.82	0.79	0.79	0.86
C 17:0	0.1	0.08	0.08	0.09	0.09	0.08	0.09	0.08	0.08	0.09
C 17:1	0.04	0.03	0.03	0.04	0.03	0.03	0.03	0.03	0.03	0.04
C 18:0	3.26	3.33	3.37	3.27	3.36	3.28	3.18	3.33	3.36	3.03
C 18:1	54.5	53.9	54.1	53.9	53.5	53.7	53.5	54.2	53.9	52.7
C 18:2	8.72	9.35	9.22	9.45	9.68	9.65	9.87	9.31	9.06	10.8
C 18:3 alpha	0.63	0.71	0.69	0.73	0.74	0.73	0.75	0.71	0.66	0.83
C 20:0	0.29	0.31	0.31	0.31	0.32	0.32	0.31	0.32	0.31	0.29

[0439] All of the positive clones selected for the sucrose utilization assay were able to hydrolyze the sucrose in the media and at the end of the 6 day culture period, there were no measurable levels of sucrose in the media. This data, in addition to the successful use of sucrose as a selection tool for positive clones, indicates that the exogenous yeast *suc2* sucrose invertase gene was targeted correctly and expressed in the transformants. As show in Table 17 above, the clones expressing Construct 24 (C8-10 thioesterase) had a measurable increase in C10 fatty acids (as high as an eight-fold increase). Likewise there were measurable increases in clones expressing Construct 25 (C12 thioesterase) and Construct 26 (C14 thioesterase) in the corresponding medium chain fatty acids. Taken together, the data shows the successful simultaneous expression in *Prototheca moriformis* two recombinant proteins (e.g., sucrose invertase and a fatty acid acyl-ACP thioesterase), both of which confer useful and quantifiable phenotypic changes on the host organism.

[0440] EXAMPLE 8: Effects of glycerol on C10-C14 fatty acid production in C14 thioesterase transformants

[0441] Clones from all the thioesterase transformations were selected and further evaluated. One clone expressing Construct 8 (*Cinnamomum camphora* C14 TE) was grown heterotrophically using different carbon sources: glucose only, fructose only and glycerol only. The glucose only condition resulted in higher cell growth and total lipid production when compared to the fructose only and glycerol only conditions. However, the proportion of C12-14 fatty acids produced in the glycerol only condition was two-fold higher than that attained in the glucose only condition.

[0442] EXAMPLE 9: Expression of *Arabidopsis thaliana* invertase in *Protoheca moriformis*

[0443] Microalgae strain *Prototheca moriformis* (UTEX 1435) was transformed using methods described above, with an expression construct containing a codon-optimized (according to Table 1) cell wall associated invertase from *Arabidopsis thaliana*. The *Arabidoposis* invertase sequence was modified to include the the N-terminal 39 amino acids from yeast invertase (SUC2 protein) to ensure efficient targeting to the ER and ultimately the periplasm. To aid detection, a Flag epitope was added to the C-terminus of the recombinant protein. The transgene was cloned into an expression vector with a *Chlorella sorokinianna* glutamate dehydrogenase promoter/5'UTR region and a *Chlorella vulgaris* nitrate reductase 3'UTR region. The DNA sequence of this transgene cassette is listed as SEQ ID NO: 89 and the translated amino acid sequence is listed as SEQ ID NO: 90. Positive clones were screened and selected using sucrose-containing media/plates. A subset of the positive clones were confirmed for the presence of the transgene and expression of invertase using Southern blot analysis and Western blot analysis for the Flag-tagged invertase. From these screens, 10 positive clones were chosen for lipid productivity and sucrose utilization assays. All 10 clones were grown on media containing sucrose as the sole carbon source and a positive control suc2 invertase transformant was also included. The negative control, wildtype *Prototheca moriformis*, was also grown but on glucose containing media. After six days, the cells were harvested and dried and the total percent lipid by dry cell weight was determined. The media was also analyzed for total sucrose consumption.

[0444] All ten positive clones were able to hydrolyze sucrose, however, most clones grew about half as well as either wildtype or the positive control suc2 yeast invertase transformant as determined by dry cell weight at the end of the experiment. Similarly, all ten positive clones produced about half as much total lipid when compared to wildtype or the positive control transformant. This data demonstrate the successful heterologous expression of diverse sucrose invertases in *Prototheca*.

[0445] EXAMPLE 10: Heterologous expression of yeast invertase (suc2) in *Prototheca krugani, Prototheca stagnora* and *Prototheca zopfii*

[0446] To test the general applicability of the transformation methods for use in species of the genus *Prototheca*, three other *Prototheca* species were selected: *Prototheca krugani* (UTEX 329), *Prototheca stagnora* (UTEX 1442) and *Prototheca zopfii* (UTEX 1438). These three strains were grown in the media and conditions described in Example 1 and their lipid

profiles were determined using the above described methods. A summary of the lipid profiles from the three *Prototheca* strains are summarized below in Area %.

Fatty Acid	*P. krugani* (UTEX 329)	*P. stagnora* (UTEX 1442)	*P. zopfii* (UTEX 1438)
C 10:0	0.0	0.0	0.0
C 10:1	0.0	0.0	0.0
C 12:0	1.5	0.8	2.1
C 14:0	1.2	0.9	1.7
C 16	15.1	17.1	19.7
C 18:0	3.3	4.1	5.4
C 18:1	66.0	61.5	53.8
C 18:2	12.9	15.6	17.3

[0447] These three strains were transformed with a yeast invertase (suc2) expression cassette (SEQ ID NO: 58) using the methods described in Example 3 above. This yeast invertase (suc2) expression cassette has been demonstrated to work in *Prototheca moriformis* (UTEX 1435) above in Example 3. The transformants were screened using sucrose containing plates/media. A subset of the positive clones for each *Prototheca* species was selected and the presence of the transgene was confirmed by Southern blot analysis. Ten of confirmed positive clones from each species were selected for sucrose hydrolysis analysis and lipid productivity. The clones were grown in media containing sucrose as the sole carbon source and compared to its wildtype counterpart grown on glucose. After 6 days, the cultures were harvested and dried and total percent lipid and dry cell weight was assessed. The media from each culture was also analyzed for sucrose hydrolysis using a YSI2700 Biochemistry Analyzer for sucrose content over the course of the experiment. Clones from all three species were able to hydrolyze sucrose, with *Prototheca stagnora* and *Prototheca zopfii* transformants being able to hydrolyze sucrose more efficiently than *Prototheca krugani*. Total lipid production and dry cell weight of the three species of transformants were comparable to their wildtype counterpart grown on glucose. This data demonstrates the successful transformation and expression exogenous genes in multiple species of the genus *Prototheca*.

[0448] EXAMPLE 11: Algal-derived Promoters and Genes for Use in Microalgae

A. 5'UTR and Promoter Sequences from *Chlorella protothecoides*

[0449] A cDNA library was generated from mixotrophically grown *Chlorella protothecoides* (UTEX 250) using standard techniques. Based upon the cDNA sequences, primers were designed in certain known housekeeping genes to "walk" upstream of the coding regions using Seegene's DNA Walking kit (Rockville, MD). Sequences isolated include an actin (SEQ ID NO:31) and elongation factor-1a (EF1a) (SEQ ID NO:32)

promoter/UTR, both of which contain introns (as shown in the lower case) and exons (upper case italicized) and the predicted start site (in bold) and two beta-tubulin promoter/UTR elements: Isoform A (SEQ ID NO:33) and Isoform B (SEQ ID NO:34).

B. Lipid Biosynthesis Enzyme and Plastid Targeting Sequences from *C. protothecoides*

[0450] From the cDNA library described above, three cDNAs encoding proteins functional in lipid metabolism in *Chlorella protothecoides* (UTEX 250) were cloned using the same methods as described above. The nucleotide and amino acid sequences for an acyl ACP desaturase (SEQ ID NOs: 45 and 46) and two geranyl geranyl diphosphate synthases (SEQ ID NOs:47-50) are included in the Sequence Listing below. Additionally, three cDNAs with putative signal sequences targeting to the plastid were also cloned. The nucleotide and amino acid sequences for a glyceraldehyde-3-phosphate dehydrogenase (SEQ ID NOs:51 and 52), an oxygen evolving complex protein OEE33 (SEQ ID NOs:53 and 54) and a Clp protease (SEQ ID NOs:55 and 56) are included in the Sequence Listing below. The putative plastid targeting sequence has been underlined in both the nucleotide and amino acid sequence. The plastid targeting sequences can be used to target the producs of transgenes to the plastid of microbes, such as lipid modification enzymes.

[0451] EXAMPLE 12: 5'UTR/promoters that are nitrogen responsive from *Prototheca moriformis*

[0452] A cDNA library was generated from *Prototheca moriformis* (UTEX 1435) using standard techniques. The *Prototheca moriformis* cells were grown for 48 hours under nitrogen replete conditions. Then a 5% innoculum (v/v) was then transferred to low nitrogen and the cells were harvested every 24 hours for seven days. After about 24 hours in culture, the nitrogen supply in the media was completely depleted. The collected samples were immediately frozen using dry ice and isopropanol. Total RNA was subsequently isolated from the frozen cell pellet samples and a portion from each sample was held in reserve for RT-PCR studies. The rest of the total RNA harvested from the samples was subjected to polyA selection. Equimolar amounts of polyA selected RNA from each condition was then pooled and used to generate a cDNA library in vector pcDNA 3.0 (Invitrogen). Roughly 1200 clones were randomly picked from the resulting pooled cDNA libray and subjected to sequencing on both strands. Approximately 68 different cDNAs were selected from among these 1200 sequences and used to design cDNA-specific primers for use in real-time RT-PCR studies.

[0453] RNA isolated from the cell pellet samples that were held in reserve was used as substrate in the real time RT-PCR studies using the cDNA-specific primer sets generated above. This reserved RNA was converted into cDNA and used as substrate for RT-PCR for each of the 68 gne specific primer sets. Threshold cylcle or C_T numbers were used to indicate relative transcript abundance for each of the 68 cDNAs within each RNA sample collected throughout the time course. cDNAs showing significant increase (greater than three fold) between nitrogen replete and nitrogen-depleted conditions were flagged as potential genes whose expression was up-regulated by nitrogen depletion. As discussed in the specification, nitrogen depletion/limitation is a known inducer of lipogenesis in oleaginous microorganisms.

[0454] In order to identify putative promoters/5'UTR sequences from the cDNAs whose expression was upregulated during nitrogen depletion/limitation, total DNA was isolated from *Prototheca moriformis* (UTEX 1435) grown under nitrogen replete conditions and were then subjected to sequencing using 454 sequencing technology (Roche). cDNAs flagged as being up-regulated by the RT-PCR results above were compared using BLAST against assembled contigs arising from the 454 genomic sequencing reads. The 5' ends of cDNAs were mapped to specific contigs, and where possible, greater than 500bp of 5' flanking DNA was used to putatively identify promoters/UTRs. The presence of promoters/5'UTR were subsequently confirmed and cloned using PCR amplification of genomic DNA. Individual cDNA 5' ends were used to design 3' primers and 5' end of the 454 contig assemblies were used to design 5' gene-specific primers.

[0455] As a first screen, one of the putative promoter, the 5'UTR/promoter isolated from Aat2 (Ammonium transporter, SEQ ID NO: 99), was cloned into the *Cinnamomum camphora* C14 thioesterase construct with the *Chlorella protothecoides* stearoyl ACP desaturase transit peptide described in Example 5 above, replacing the *C.sorokinana* glutamate dehydrogenase promoter. This construct is listed as SEQ ID NO: 112 . To test the putative promoter, the thioesterase construct is transformed into *Prototheca moriformis* cells to confirm actual promoter activity by screening for an increase in C14/C12 fatty acids under low/no nitrogen conditions, using the methods described above. Similar testing of the putative nitrogen-regulated promoters isolated from the cDNA/genomic screen can be done using the same methods.

[0456] Other putative nitrogen-regulated promoters/5'UTRs that were isolated from the cDNA/genomic screen were:

Promoter/5'UTR	SEQ ID NO.	Fold increased

FatB/A promoter/5'UTR	SEQ ID NO: 91	n/a
NRAMP metal transporter promoter/5'UTR	SEQ ID NO: 92	9.65
Flap Flagellar-associated protein promoter/5'UTR	SEQ ID NO: 93	4.92
SulfRed Sulfite reductase promoter/5'UTR	SEQ ID NO: 94	10.91
SugT Sugar transporter promoter/5'UTR	SEQ ID NO: 95	17.35
Amt03-Ammonium transporter 03 promoter/5'UTR	SEQ ID NO: 96	10.1
Amt02-Ammonium transporter 02 promoter/5'UTR	SEQ ID NO: 97	10.76
Aat01-Amino acid transporter 01 promoter/5'UTR	SEQ ID NO: 98	6.21
Aat02-Amino acid transporter 02 promoter/5'UTR	SEQ ID NO: 99	6.5
Aat03-Amino acid transporter 03 promoter/5'UTR	SEQ ID NO: 100	7.87
Aat04-Amino acid transporter 04 promoter/5'UTR	SEQ ID NO: 101	10.95
Aat05-Amino acid transporter 05 promoter/5'UTR	SEQ ID NO: 102	6.71

Fold increase refers to the fold increase in cDNA abundance after 24 hours of culture in low nitrogen medium.

[0457] EXAMPLE 13: Homologous Recombination in *Prototheca* species

[0458] Homologous recombination of transgenes has several advantages over the transformation methods described in the above Examples. First, the introduction of transgenes without homologous recombination can be unpredictable because there is no control over the number of copies of the plasmid that gets introduced into the cell. Also, the introduction of transgenes without homologous recombination can be unstable because the plasmid may remain episomal and is lost over subsequent cell divisions. Another advantage of homologous recombination is the ability to "knock-out" gene targets, introduce epitope tags, switch promoters of endogenous genes and otherwise alter gene targets (e.g., the introduction of point mutations.

[0459] Two vectors were constructed using a specific region of the *Prototheca moriformis* (UTEX 1435) genome, designated KE858. KE858 is a 1.3 kb, genomic fragment that encompasses part of the coding region for a protein that shares homology with the transfer RNA (tRNA) family of proteins. Southern blots have shown that the KE858 sequence is present in a single copy in the *Prototheca moriformis* (UTEX 1435) genome. The first type of vector that was constructed, designated SZ725 (SEQ ID NO: 103), consisted of the entire 1.3 kb KE858 fragment cloned into a pUC19 vector backbone that also contains the optimized yeast invertase (*suc2*) gene used in Example 3 above. The KE858 fragment contains an unique SnaB1 site that does not occur anywhere else in the targeting construct. The second type of vector that was constructed, designated SZ726 (SEQ ID NO: 126),

consisted of the KE858 sequence that had been disrupted by the insertion of the yeast invertase gene *(suc2)* at the SnaB1 site within the KE858 genomic sequence. The entire DNA fragment containing the KE858 sequences flanking the yeast invertase gene can be excised from the vector backbone by digestion with EcoRI, which cuts at either end of the KE858 region.

[0460] Both vectors were used to direct homologous recombination of the yeast invertase gene (*suc2*) into the corresponding KE858 region of the *Prototheca moriformis* (UTEX 1435) genome. The linear DNA ends homologous to the genomic region that was being targeted for homologous recombination were exposed by digesting the vector construct SZ725 with SnaB1 and vector construct SZ726 with EcoRI. The digested vector constructs were then introduced into *Prototheca moriformis* cultures using methods described above in Example 3. Transformants from each vector construct were then selected using sucrose plates. Ten independent, clonally pure transformants from each vector transformation were analyzed for successful recombination of the yeast invertase gene into the desired genomic location (using Southern blots) and for transgene stability.

[0461] Southern blot analysis of the SZ725 transformants showed that 4 out of the 10 transformants picked for analysis contained the predicted recombinant bands, indicating that a single crossover event had occurred between the KE858 sequences on the vector and the KE858 sequences in the genome. In contrast, all ten of the SZ726 transformants contained the predicted recombinat bands, indicating that double crossover events had occurred between the EcoRI fragment of pSZ726 carrying KE858 sequence flanking the yeast invertase transgene and the corresponding KE858 region of the genome.

[0462] Sucrose invertase expression and transgene stability were assessed by growing the transformants for over 15 generations in the absence of selection. The four SZ725 transformants and the ten SZ276 transformants that were positive for the transgene by Southern blotting were selected and 48 single colonies from each of the transformants were grown serially: first without selection in glucose containing media and then with selection in media containing sucrose as the sole carbon source. All ten SZ276 transformants (100%) retained their ability to grow on sucrose after 15 generations, whereas about 97% of the SZ725 transformants retained their ability to grow on sucrose after 15 generations. Transgenes introduced by a double crossover event (SZ726 vector) have extremely high stability over generation doublings. In contrast, transgenes introduced by a single cross over event (SZ725 vector) can result in some instability over generation doublings because is

tandem copies of the transgenes were introduced, the repeated homologous regions flanking the transgenes may recombine and excise the transgenic DNA located between them.

[0463] These experiments demonstrate the successful use of homologous recombination to generate *Prototheca* transformants containing a heterologous sucrose invertase gene that is stably integrated into the nuclear chromosomes of the organism. The success of the homologous recombination enables other genomic alterations in *Prototheca*, including gene deletions, point mutations and epitope tagging a desired gene product. These experiments also demonstrate the first documented system for homologous recombination in the nuclear genome of an eukaryotic microalgae.

A. Use of Homologous Recombination to Knock-out an Endogenous *Prototheca moriformis* gene

[0464] In the *Prototheca moriformis* cDNA/genomic screen described in Example 11 above, an endogenous stearoyl ACP desaturase (SAPD) cDNA was identified. Stearoyl ACP desaturase enzymes are part of the lipid synthesis pathway and they function to introduce double bonds into the fatty acyl chains. In some cases, it may be advantages to knock-out or reduce the expression of lipid pathway enzymes in order to alter a fatty acid profile. A homologous recombination construct was created to assess whether the expression of an endogenous stearoyl ACP desaturase enzyme can be reduced (or knocked out) and if a corresponding reduction in unsaturated fatty acids can be observed in the lipid profile of the host cell. An approximately 1.5kb coding sequence of a stearoyl ACP desaturase gene from *Prototheca moriformis* (UTEX 1435) was identified and cloned (SEQ ID NO: 104). The homologous recombination construct was constructed using 0.5kb of the SAPD coding sequence at the 5'end (5' targeting site), followed by the *Chlamydomonas reinhardtii* β-tublin promoter driving a codon-optimized yeast sucrose invertase *suc2* gene with the *Chlorella vulgaris* 3'UTR. The rest (~1kb) of the *Prototheca moriformis* SAPD coding sequence was then inserted after the *C.vulgaris* 3'UTR to make up the 3' targeting site. The sequence for this homologous recombination cassette is listed in SEQ ID NO: 105. As shown above, the success-rate for integration of the homologous recombination cassette into the nuclear genome can be increased by linearizing the cassette before transforming the microalgae, leaving exposed ends. The homologous recombination cassette targeting an endogenous SAPD enzyme in *Prototheca moriformis* is linearized and then transformed into the host cell (*Prototheca moriformis*, UTEX 1435). A successful integration will eliminate the endogenous SAPD enzyme coding region from the host genome via a double reciprocal recombination event, while expression of the newly inserted *suc2* gene will be regulated by

the *C.reinhardtii* β-tubulin promoter. The resulting clones can be screened using plates/media containing sucrose as the sole carbon source. Clones containing a successful integration of the homologous recombination cassette will have the ability to grow on sucrose as the sole carbon source and changes in overall saturation of the fatty acids in the lipid profile will serve as a secondary confirmation factor. Additionally, Southern blotting assays using a probe specific for the yeast sucrose invertase *suc2* gene and RT-PCR can also confirm the presence and expression of the invertase gene in positive clones. As an alternative, the same construct without the β-tubulin promoter can be used to excise the endogenous SAPD enzyme coding region. In this case, the newly inserted yeast sucrose invertase *suc2* gene will be regulated by the endogenous SAPD promoter/5'UTR.

[0465] EXAMPLE 14: Fuel Production

A. Extraction of oil from microalgae using an expeller press and a press aid

[0466] Microalgal biomass containing 38% oil by DCW was dried using a drum dryer resulting in resulting moisture content of 5-5.5%. The biomass was fed into a French L250 press. 30.4 kg (67 lbs.) of biomass was fed through the press and no oil was recovered. The same dried microbial biomass combined with varying percentage of switchgrass as a press aid was fed through the press. The combination of dried microbial biomass and 20% w/w switchgrass yielded the best overall percentage oil recovery. The pressed cakes were then subjected to hexane extraction and the final yield for the 20% switchgrass condition was 61.6% of the total available oil (calculated by weight). Biomass with above 50% oil dry cell weight did not require the use of a pressing aid such as switchgrass in order to liberate oil.

B. Monosaccharide Composition of Delipidated *Prototheca moriformis* biomass

[0467] *Prototheca moriformis* (UTEX 1435) was grown in conditions and nutrient media (with 4% glucose) as described in Example 45 above. The microalgal biomass was then harvested and dried using a drum dryer. The dried algal biomass was lysed and the oil extracted using an expeller press as described in Example 44 above. The residual oil in the pressed biomass was then solvent extracted using petroleum ether. Residual petroleum ether was evaporated from the delipidated meal using a Rotovapor (Buchi Labortechnik AG, Switzerland). Glycosyl (monosaccharide) composition analysis was then performed on the delipidated meal using combined gas chromatography/mass spectrometry (GC/MS) of the per-O-trimethylsily (TMS) derivatives of the monosaccharide methyl glycosides produced from the sample by acidic methanolysis. A sample of delipidated meal was subjected to methanolysis in 1M HCl in methanol at 80°C for approximately 20 hours, followed by re-N-

acetylation with pyridine and acetic anhydride in methanol (for detection of amino sugars). The samples were then per-O-trimethylsiylated by treatment with Tri-Sil (Pierce) at 80°C for 30 minutes (see methods in Merkle and Poppe (1994) *Methods Enzymol.* 230: 1-15 and York et al., (1985) *Methods Enzymol.* 118:3-40). GC/MS analysis of the TMS methyl glycosides was performed on an HP 6890 GC interfaced to a 5975b MSD, using a All Tech EC-1 fused silica capillary column (30m x 0.25 mm ID). The monosaccharides were identified by their retention times in comparison to standards, and the carbohydrate character of these are authenticated by their mass spectra. 20 micrograms per sample of inositol was added to the sample before derivatization as an internal standard. The monosaccharide profile of the delipidated *Prototheca moriformis* (UTEX 1435) biomass is summarized in Table 18 below. The total percent carbohydrate from the sample was calculated to be 28.7%.

[0468] Table 18. Monosaccharide (glycosyl) composition analysis of *Prototheca moriformis* (UTEX 1435) delipidated biomass.

	Mass (µg)	**Mole % (of total carbohydrate)**
Arabinose	0.6	1.2
Xylose	n.d.	n.d.
Galacturonic acid (GalUA)	n.d.	n.d.
Mannose	6.9	11.9
Galactose	14.5	25.2
Glucose	35.5	61.7
N Acetyl Galactosamine (GalNAc)	n.d.	n.d.
N Acetyl Glucosamine (GlcNAc)	n.d.	n.d.
Heptose	n.d.	n.d.
3 Deoxy-2-manno-2 Octulsonic acid (KDO)	n.d.	n.d.
Sum	**57**	**100**

n.d. = none detected

[0469] The carbohydrate content and monosaccharide composition of the delipidated meal makes it suitable for use as an animal feed or as part of an animal feed formulation. Thus, in one aspect, the present invention provides delipidated meal having the product content set forth in the table above.

C. Production of biodiesel from *Prototheca* oil

[0470] Degummed oil from *Prototheca moriformis* UTEX 1435, produced according to the methods described above, was subjected to transesterification to produce fatty acid methyl esters. Results are shown below:

The lipid profile of the oil was:
C10:0 0.02
C12:0 0.06
C14:0 1.81
C14.1 0.07

C16:0 24.53
C16:1 1.22
C18:0 2.34
C18:1 59.21
C18:2 8.91
C18:3 0.28
C20:0 0.23
C20:1 0.10
C20:1 0.08
C21:0 0.02
C22:0 0.06
C24:0 0.10

[0471] Table 19. Biodiesel profile from *Prototheca moriformis* triglyceride oil.

Method	Test		Result	Units
ASTM D6751 A1	Cold Soak Filterability of Biodiesel Blend Fuels	Filtration Time	120	sec
		Volume Filtered	300	ml
ASTM D93	Pensky-Martens Closed Cup Flash Point	Procedure Used	A	
		Corrected Flash Point	165.0	°C
ASTM D2709	Water and Sediment in Middle Distillate Fuels (Centrifuge Method)	Sediment and Water	0.000	Vol %
EN 14538	Determination of Ca and Mg Content by ICP OES	Sum of (Ca and Mg)	<1	mg/kg
EN 14538	Determination of Ca and Mg Content by ICP OES	Sum of (Na and K)	<1	mg/kg
ASTM D445	Kinematic / Dynamic Viscosity	Kinematic Viscosity @ 104 °F/ 40 °C	4.873	mm^2/s
ASTM D874	Sulfated Ash from Lubricating Oils and Additives	Sulfated Ash	< 0.005	Wt %
ASTM D5453	Determination of Total Sulfur in Light Hydrocarbons, Spark Ignition Engine Fuel, Diesel Engine Fuel, and Engine Oil by Ultraviolet Fluorescence.	Sulfur, mg /kg	1.7	mg/kg
ASTM D130	Corrosion - Copper Strip	Biodiesel-Cu Corrosion 50°C (122°F)/3 hr	1a	
ASTM D2500	Cloud Point	Cloud Point	6	°C
ASTM D4530	Micro Carbon Residue	Average Micro Method Carbon Residue	< 0.10	Wt %
ASTM D664	Acid Number of Petroleum Products by Potentiometric Titration	Procedure Used	A	
		Acid Number	0.20	mg KOH/g
ASTM	Determination of Free and	Free Glycerin	< 0.005	Wt %

Method	Test		Result	Units
D6584	Total Glycerin in B-100 Biodiesel Methyl Esters By Gas Chromatography	Total Glycerin	0.123	Wt %
ASTM D4951	Additive Elements in Lubricating Oils by ICP-AES	Phosphorus	0.000200	Wt %
ASTM D1160	Distillation of Petroleum Products at Reduced Pressure	IBP	248	°C
		AET @ 5% Recovery	336	°C
		AET @ 10% Recovery	338	°C
		AET @ 20% Recovery	339	°C
		AET @ 30% Recovery	340	°C
		AET @ 40% Recovery	342	°C
		AET @ 50% Recovery	344	°C
		AET @ 60% Recovery	345	°C
		AET @ 70% Recovery	347	°C
		AET @ 80% Recovery	349	°C
		AET @ 90% Recovery	351	°C
		AET @ 95% Recovery	353	°C
		FBP	362	°C
		% Recovered	98.5	%
		% Loss	1.5	%
		% Residue	0.0	%
		Cold Trap Volume	0.0	ml
		IBP	248	°C
EN 14112	Determination of Oxidation Stability (Accelerated Oxidation Test)	Oxidation Stability	> 12	hr
		Operating Temp (usually 110 deg C)	110	°C
ASTM D4052	Density of Liquids by Digital Density Meter	API Gravity @ 60°F	29.5	°API
ASTM D 6890	Determination of Ignition Delay (ID) and Derived Cetane Number (DCN)	Derived Cetane Number (DCN)	> 61.0	

[0472] The lipid profile of the biodiesel was highly similar to the lipid profile of the feedstock oil. Other oils provided by the methods and compositions of the invention can be subjected to transesterification to yield biodiesel with lipid profiles including (a) at least 4%

C8-C14; (b) at least 0.3% C8; (c) at least 2% C10; (d) at least 2% C12; and (3) at least 30% C8-C14.

[0473] The Cold Soak Filterability by the ASTM D6751 A1 method of the biodiesel produced was 120 seconds for a volume of 300ml. This test involves filtration of 300 ml of B100, chilled to 40°F for 16 hours, allowed to warm to room temp, and filtered under vacuum using 0.7 micron glass fiber filter with stainless steel support. Oils of the invention can be transesterified to generate biodiesel with a cold soak time of less than 120 seconds, less than 100 seconds, and less than 90 seconds.

D. Production of Renewable Diesel

[0474] Degummed oil from *Prototheca moriformis* UTEX 1435, produced according to the methods described above and having the same lipid profile as the oil used to make biodiesel in Example X above, was subjected to transesterification to produce renewable diesel.

[0475] The oil was first hydrotreated to remove oxygen and the glycerol backbone, yielding n-paraffins. The n-parrafins were then subjected to cracking and isomerization. A chromatogram of the material is shown in Figure 13. The material was then subjected to cold filtration, which removed about 5% of the C18 material. Following the cold filtration the total volume material was cut to flash point and evaluated for flash point, ASTM D-86 distillation distribution, cloud point and viscosity. Flash point was 63°C; viscosity was 2.86 cSt (centistokes); cloud point was 4°C. ASTM D86 distillation values are shown in Table 20:

[0476] Table 20. Readings in °C:

Volume	Temperature
IBP	173
5	217.4
10	242.1
15	255.8
20	265.6
30	277.3
40	283.5
50	286.6
60	289.4
70	290.9
80	294.3
90	300
95	307.7
FBP	331.5

[0477] The T10-T90 of the material produced was 57.9°C. Methods of hydrotreating, isomerization, and other covalent modification of oils disclosed herein, as well as methods of distillation and fractionation (such as cold filtration) disclosed herein, can be employed to

generate renewable diesel compositions with other T10-T90 ranges, such as 20, 25, 30, 35, 40, 45, 50, 60 and 65°C using triglyceride oils produced according to the methods disclosed herein.

[0478] The T10 of the material produced was 242.1°C. Methods of hydrotreating, isomerization, and other covalent modification of oils disclosed herein, as well as methods of distillation and fractionation (such as cold filtration) disclosed herein, can be employed to generate renewable diesel compositions with other T10 values, such as T10 between 180 and 295, between 190 and 270, between 210 and 250, between 225 and 245, and at least 290.

[0479] The T90 of the material produced was 300°C. Methods of hydrotreating, isomerization, and other covalent modification of oils disclosed herein, as well as methods of distillation and fractionation (such as cold filtration) disclosed herein can be employed to generate renewable diesel compositions with other T90 values, such as T90 between 280 and 380, between 290 and 360, between 300 and 350, between 310 and 340, and at least 290.

[0480] The FBP of the material produced was 300°C. Methods of hydrotreating, isomerization, and other covalent modification of oils disclosed herein, as well as methods of distillation and fractionation (such as cold filtration) disclosed herein, can be employed to generate renewable diesel compositions with other FBP values, such as FBP between 290 and 400, between 300 and 385, between 310 and 370, between 315 and 360, and at least 300.

[0481] Other oils provided by the methods and compositions of the invention can be subjected to combinations of hydrotreating, isomerization, and other covalent modification including oils with lipid profiles including (a) at least 4% C8-C14; (b) at least 0.3% C8; (c) at least 2% C10; (d) at least 2% C12; and (3) at least 30% C8-C14.

[0482] EXAMPLE 15: Utilization of Sucrose by *Chlorella luteoviridis*

A. SAG 2214 Growth on Glucose and Sucrose

[0483] SAG 2214 (designated as *Chlorella luteoviridis*) was tested for growth in the dark on media containing either glucose or sucrose. Heterotrophic liquid cultures were initiated using inoculum from a frozen vial in either media containing 4% glucose or 4% sucrose as the sole carbon source. Cultures were grown in the dark, shaking at 200 rpm. Samples from the cultures were taken at 0, 24, 48 and 72 hour timepoints and growth was measured by relative absorbance at 750nm (UV Mini1240, Shimadzu). SAG 2214 grew equally well on glucose as on sucrose, showing that this microalgae can utilize sucrose as effectively as glucose as a sole carbon source. The result of this experiment is represented graphically in Figure 3.

B. Lipid Productivity and Fatty Acid Profile for SAG 2214

[0484] Microalgal strain SAG 2214 was cultivated in liquid medium containing either glucose or sucrose as the sole carbon source in similar conditions as described in Example 32 above. After 7 days, cells were harvested for dry cell weight calculation. Cells were centrifuged and lyophilized for 24 hours. The dried cell pellets were weighed and the dry cell weight per liter was calculated. Cells for lipid analysis were also harvested and centrifuged at 4000 x g for 10 minutes at room temperature. The supernatant was discarded and the samples were processed for lipid analysis and fatty acid profile using standard gas chromatography (GC/FID) procedures. The results are summarized below in Tables 21 and 22.

[0485] Table 21. Lipid productivity and DCW for SAG 2214.

Sample	Lipid (g/L)	DCW (g/L)	% Lipid DCW
SAG 2214 glucose	2.43	5.73	42.44%
SAG 2214 sucrose	0.91	2.00	45.56%

[0486] Table 22. Fatty acid profile for SAG 2214.

Fatty Acid	Percent (w/w)
C:16:0	21
C:18:1	38
C:18:2	41

C. Genomic Comparison of SAG 2214 to other *Chlorella luteoviridis* strains

[0487] Microalgal strain SAG 2214 proved to be of general interest due to its ability to grow on sucrose as a carbon source (illustrated above). In addition to the growth characteristics of this strain, its taxonomic relationship to other microalgal species was also of interest. Designated by the SAG collection as a *Chlorella luteoviridis* strain, the 23s rRNA gene of SAG 2214 was sequenced and compared to the 23s rRNA genomic sequence of nine other strains also identified by the SAG and UTEX collections as *Chlorella luteoviridis*. These strains were UTEX 21, 22, 28, 257 and 258, and SAG strains 2133, 2196, 2198 and 2203. The DNA genotyping methods used were the same as the methods described above in Example 1. Sequence alignments and unrooted trees were generated using Geneious DNA analysis software. Out of the nine other strains that were genotypes, UTEX 21, 22, 28 and 257 had identical 23s rRNA DNA sequence (SEQ ID NO: 106). The other five *Chlorella luteoviridis* strains had 23s rRNA sequences that were highly homologous to UTEX 21, 22, 28, and 257.

[0488] The 23s rRNA gene sequence from SAG 2214 (SEQ ID NO: 30)is decidedly different from that of the the other nine *C. luteoviridis* strains, having a large insertion that was not found in the other strains. Further analysis of this 23s rRNA gene sequence using BLAST indicated that it shared the greatest homology with members of the genus *Leptosira* and *Trebouxia* (members of phycobiont portion of lichens). These results indicate that SAG 2214 may not be *Chlorella luteoviridis* strain as categorized by the strain collection, but instead shares significant 23S rRNA nucleotide identity to algal symbionts found in lichen. The genomic analysis along with the growth characteristics indicate that SAG 2214 may be a source for genes and proteins involved in the metabolism of sucrose, as well as signaling and transit peptides responsible for the correct localization of such enzymes. SAG 2214 and other strains with a high degree of genomic similarity may also be strains useful for oil production using sucrose as a source of fixed carbon.

[0489] Although this invention has been described in connection with specific embodiments thereof, it will be understood that it is capable of further modifications. This application is intended to cover any variations, uses, or adaptations of the invention following, in general, the principles of the invention and including such departures from the present disclosure as come within known or customary practice within the art to which the invention pertains and as may be applied to the essential features hereinbefore set forth.

[0490] All references cited herein, including patents, patent applications, and publications, are hereby incorporated by reference in their entireties, whether previously specifically incorporated or not. The publications mentioned herein are cited for the purpose of describing and disclosing reagents, methodologies and concepts that may be used in connection with the present invention. Nothing herein is to be construed as an admission that these references are prior art in relation to the inventions described herein. In particular, the following patent applications are hereby incorporated by reference in their entireties for all purposes: US Provisional Application No. 60/941,581, filed June 1, 2007, entitled "Production of Hydrocarbons in Microorganisms"; US Provisional Application No. 60/959,174, filed July 10, 2007, entitled "Production of Hydrocarbons in Microorganisms"; US Provisional Application No. 60/968,291, filed August 27, 2007, entitled "Production of Hydrocarbons in Microorganisms"; US Provisional Application No. 61/024,069, filed January 28, 2008, entitled "Production of Hydrocarbons in Microorganims"; PCT Application No. PCT/US08/65563, filed June 2, 2008, entitled "Production of Oil in Microorganisms"; US Patent Application No. 12/131,783, filed June 2, 2008, entitled "Use of Cellulosic Material for Cultivation of Microorganisms"; US Patent Application No. 12/131,773, filed June 2,

2008, entitled "Renewable Diesel and Jet Fuel from Microbial Sources"; US Patent Application No. 12/131,793, filed June 2, 2008, entitled "Sucrose Feedstock Utilization for Oil-Based Fuel Manufacturing"; US Patent Application No. 12/131,766, filed June 2, 2008, entitled "Glycerol Feedstock Utilization for Oil-Based Fuel Manufacturing"; US Patent Application No. 12/131,804, filed June 2, 2008, entitled "Lipid Pathway Modification in Oil-Bearing Microorganisms"; US Patent Application No. 61/118,590, filed November 28, 2008, entitled "Production of Oil in Microorganisms"; US Provisional Patent Application No. 61/118,994, filed December 1, 2008, entitled "Production of Oil in Microorganisms"; US Provisional Patent Application No. 61/174,357, filed April 3, 2009, entitled "Production of Oil in Microorganisms"; US Provisional Patent Application No. 61/219,525, filed June 23, 2009, entitled "Production of Oil in Microorganisms"; and PCT Application No. _____ [[Attorney Docket No. 026172-005010PC]], filed November 30, 2009, entitled "Production of Tailored Oils in Heterotrophic Microorganisms".

WHAT IS CLAIMED IS:

1. A recombinant nucleic acid encoding a promoter that is upregulated at least 3-fold in a cell of the genus *Prototheca* when the extracellular environment changes from containing at least 5mM nitrogen to containing no nitrogen.

2. The recombinant nucleic acid of claim 1, wherein the nucleic acid comprises a segment of 50 or more nucleotides of one of SEQ ID NOs: 91-102.

3. A nucleic acid vector comprising an expression cassette comprising (a) a promoter that is active in a cell of the genus *Prototheca*; and (b) a coding sequence in operable linkage with the promoter wherein the coding sequence contains the most or second most preferred codons of Table 1 for at least 20% of the codons of the coding sequence.

4. The vector of claim 3, wherein the coding sequence comprises a plastid targeting sequence in-frame with a fatty acyl-ACP thioesterase.

5. The vector of claim 4, wherein the fatty acyl-ACP thioesterase has hydrolysis activity towards one or more fatty acyl-ACP substrates of chain length C8, C10, C12 or C14.

6. The vector of claim 3, wherein the promoter is upregulated at least 3-fold in a cell of the genus *Prototheca* when the extracellular environment changes from containing at least 5mM nitrogen to containing no nitrogen.

7. The vector of claim 4, wherein the plastid targeting sequence encodes a peptide that is capable of targeting a protein to the plastid of a cell of the genus *Prototheca*.

8. The vector of claim 4, wherein the plastid targeting sequence is from microalgae.

9. The vector of claim 8, wherein the plastid targeting sequence has at least 25% amino acid sequence identity to one or more of SEQ ID NOs: 127-133 and is capable of targeting a protein to the plastid of a cell of the genus *Prototheca*.

10. The vector of claim 3, further comprising a nucleic acid sequence endogenous to the nuclear genome of a cell of the genus *Prototheca*, wherein the sequence is at least 200 nucleotides long.

11. The vector of claim 10, further comprising first and second nucleic acid sequences endogenous to the nuclear genome of a cell of the genus *Prototheca*, wherein the first and second sequences (a) are each at least 200 nucleotides long; (b) flank the expression cassette; and (c) are located on the same *Prototheca* chromosome no more than 20kB apart.

12. A method of producing a triglyceride composition, comprising:
(a) culturing a population of cells of the genus *Prototheca* in the presence of a fixed carbon source, wherein:

(i) the cells contain an exogenous gene;

(ii) the cells accumulate at least 10% of their dry cell weight as lipid; and

(iii) the fixed carbon source is selected from the group consisting of sorghum and depolymerized cellulosic material; and
(b) isolating lipid components from the cultured microorganisms.

13. The method of claim 12, wherein the fixed carbon source is depolymerized cellulosic material selected from the group consisting of corn stover, *Miscanthus*, forage sorghum, sugar beet pulp and sugar cane bagasse.

14. The method of claim 12, wherein the fixed carbon source is depolymerized cellulosic material that has been subjected to washing with water prior to the culturing step.

15. The method of claim 12, wherein the fixed carbon source is depolymerized cellulosic material and the glucose level of the depolymerized cellulosic material is concentrated to a level of at least 300 g/liter, at least 400 g/liter, at least 500 g/liter, or at least 600 g/liter of prior to the culturing step and is fed to the culture over time as the cells grow and accumulate lipid.

16. The method of claim 12, wherein the exogenous gene encodes a fatty acyl-ACP thioesterase that has hydrolysis activity towards one or more fatty acyl-ACP substrates of chain length C8, C10, C12 or C14.

17. The method of claim 12, wherein the triglyceride composition has a profile of:

a. at least 4% C8-C14; and

b. one or more of the following attributes:

i. less than 0.4 micrograms/ml total carotenoids;

ii. less than 0.001 micrograms/ml lycopene;

iii. less than 0.02 micrograms/ml beta carotene;

iv. less than 0.02 milligrams of chlorophyll per kilogram of oil;

v. 0.40-0.60 milligrams of gamma tocopherol per 100 grams of oil;

vi. 3-9 mg campesterol per 100 grams of oil; and

vii. less than 0.5 milligrams of total tocotrienols per gram of oil.

18. A method of producing a triglyceride composition, comprising:
(a) culturing a population of microorganisms in the presence of depolymerized cellulosic material, wherein:

(i) the depolymerized cellulosic material is subjected to washing with water prior to the culturing step;

(ii) the cells accumulate at least 10% of their dry cell weight as lipid; and

(iii) the depolymerized cellulosic material is concentrated to at least 300 g/liter of glucose prior to the cultivation step;

(iv) the microorganisms are cultured in a fed-batch reaction in which depolymerized cellulosic material of at least 300 g/liter of glucose is fed to the microorganisms; and
(b) isolating lipid components from the cultured microorganisms.

19. The method of claim 18, wherein the depolymerized cellulosic material is selected from the group consisting of corn stover, *Miscanthus*, forage sorghum, sugar beet pulp and sugar cane bagasse.

20. The method of claim 18, wherein the microorganisms are a species of the genus *Prototheca* and contain an exogenous gene.

21. The method of claim 20, wherein the exogenous gene encodes a fatty acyl-ACP thioesterase that has hydrolysis activity towards one or more fatty acyl-ACP substrates of chain length C8, C10, C12 or C14.

22. A method of manufacturing triglyceride oil comprising cultivating a cell that has a 23S rRNA sequence with at least 96% nucleotide identity to SEQ ID NO: 30 in the presence of sucrose as a carbon source.

23. A method of manufacturing a chemical, comprising performing one or more chemical reactions selected from the group consisting of transesterification, hydrogenation, hydrocracking, deoxygenation, isomerization, interesterification, hydroxylation, hydrolysis, and saponification on a triglyceride oil, wherein the oil:

(a) has a lipid profile of at least 4% C8-C14; and

(b) at least one of the following attributes:

i. less than 0.4 micrograms/ml total carotenoids;

ii. less than 0.001 micrograms/ml lycopene;

iii. less than 0.02 micrograms/ml beta carotene;

iv. less than 0.02 milligrams of chlorophyll per kilogram of oil;

v. 0.40-0.60 milligrams of gamma tocopherol per 100 grams of oil;

vi. 3-9 mg campesterol per 100 grams of oil; and

vii. less than 0.5 milligrams of total tocotrienols per gram of oil.

24. The method of claim 23, further comprising a step of manufacturing the oil by cultivating a cell of the genus *Prototheca* that comprises an exogenous fatty acyl-ACP thioesterase gene that encodes a fatty acyl-ACP thioesterase having hydrolysis activity towards one or more fatty acyl-ACP substrates of chain length C8, C10, C12 or C14.

25. The method of claim 23, comprising performing a hydrolysis reaction.

26. The method of claim 25, wherein the hydrolysis reaction comprises a hydrolysis reaction selected from the group consisting of saponification, acid hydrolysis, alkaline hydrolysis, enzymatic hydrolysis, catalytic hydrolysis, and hot-compressed water hydrolysis.

27. The method of claim 25, wherein the hydrolysis reaction comprises a catalytic hydrolysis reaction and wherein the oil is split into glycerol and fatty acids.

28. The method of claim 27, wherein the fatty acids formed undergo an amination reaction to produce fatty nitrogen compounds.

29. The method of claim 27, wherein the fatty acids formed undergo an ozonolysis reaction to produce mono- and dibasic-acids.

30. The method of claim 25, wherein the oil undergoes a triglyceride splitting method selected from the group consisting of enzymatic splitting and pressure splitting.

31. The method of claim 25, wherein the hydrolysis reaction comprises a saponification reaction.

32. The method of claim 25, wherein a condensation reaction follows the hydrolysis reaction.

33. A method of manufacturing an oil, comprising performing a hydroprocessing reaction, wherein the oil:

(a) has a lipid profile of at least 4% C8-C14; and

(b) at least one of the following attributes:

i. less than 0.4 micrograms/ml total carotenoids;

ii. less than 0.001 micrograms/ml lycopene;

iii. less than 0.02 micrograms/ml beta carotene;

iv. less than 0.02 milligrams of chlorophyll per kilogram of oil;

v. 0.40-0.60 milligrams of gamma tocopherol per 100 grams of oil;

vi. 3-9 mg campesterol per 100 grams of oil; and

vii. less than 0.5 milligrams of total tocotrienols per gram of oil.

34. The method of claim 33, wherein the product of the hydroprocessing reaction undergoes a deoxygenation reaction.

35. The method of claim 33, wherein the oil undergoes a deoxygenation reaction prior to the performance of a hydroprocessing reaction.

36. The method of claim 33, wherein the oil undergoes a hydroprocessing reaction and a deoxygenation reaction simultaneously.

37. The method of claim 33, further comprising an isomerization reaction.

38. The method of claim 33, further comprising a gas removal reaction.

39. The method of claim 33, wherein a condensation reaction follows the hydroprocessing reaction.

40. A method of manufacturing an oil, comprising performing a deoxygenation reaction selected from the group consisting of a hydrogenolysis reaction, hydrogenation, a consecutive hydrogenation-hydrogenolysis reaction, a consecutive hydrogenolysis-hydrogenation reaction, and a combined hydrogenation-hydrogenolysis reaction, wherein the oil:

(a) has a lipid profile of at least 4% C8-C14; and

(b) at least one of the following attributes:

i. less than 0.4 micrograms/ml total carotenoids;

ii. less than 0.001 micrograms/ml lycopene;

iii. less than 0.02 micrograms/ml beta carotene;

iv. less than 0.02 milligrams of chlorophyll per kilogram of oil;

v. 0.40-0.60 milligrams of gamma tocopherol per 100 grams of oil;

vi. 3-9 mg campesterol per 100 grams of oil; and

vii. less than 0.5 milligrams of total tocotrienols per gram of oil.

41. The method of claim 40, wherein a condensation reaction follows the deoxygenation reaction.

42. A method of manufacturing an oil, comprising performing an esterification reaction, wherein the oil:

(a) has a lipid profile of at least 4% C8-C14; and

(b) at least one of the following attributes:

i. less than 0.4 micrograms/ml total carotenoids;

ii. less than 0.001 micrograms/ml lycopene;

iii. less than 0.02 micrograms/ml beta carotene;

iv. less than 0.02 milligrams of chlorophyll per kilogram of oil;
v. 0.40-0.60 milligrams of gamma tocopherol per 100 grams of oil;
vi. 3-9 mg campesterol per 100 grams of oil; and
vii. less than 0.5 milligrams of total tocotrienols per gram of oil.

43. The method of claim 42, wherein the esterification reaction comprises an interestification reaction.

44. The method of claim 42, wherein the esterification reaction comprises a transesterification reaction.

45. The method of claim 42, wherein a condensation reaction follows the esterification reaction.

46. A method of manufacturing an oil, comprising performing a hydroxylation reaction, wherein the oil:

(a) has a lipid profile of at least 4% C8-C14; and
(b) at least one of the following attributes:
i. less than 0.4 micrograms/ml total carotenoids;
ii. less than 0.001 micrograms/ml lycopene;
iii. less than 0.02 micrograms/ml beta carotene;
iv. less than 0.02 milligrams of chlorophyll per kilogram of oil;
v. 0.40-0.60 milligrams of gamma tocopherol per 100 grams of oil;
vi. 3-9 mg campesterol per 100 grams of oil; and
vii. less than 0.5 milligrams of total tocotrienols per gram of oil.

47. The method of claim 46, wherein a condensation reaction follows the hydroxylation reaction.

48. A recombinant nucleic acid with at least 95% nucleotide identity to one or both of SEQ ID NOs: 134-135.

49. A recombinant nucleic acid encoding a protein with at least 90% amino acid identity to one or both of SEQ ID NOs: 136-137.

SEQUENCE LISTING

SEQ ID NO: 1
HUP promoter from Chlorella (subsequence of GenBank accession number X55349)

gatcagacgggcctgacctgcgagataatcaagtgctcgtaggcaaccaactcagcagctgcttggtgttgggtctgcaggatagtgtt
gcagggccccaaggacagcaggggaacttacaccttgtccccgacccagttttatggagtgcattgcctcaagagcctagccggagc
gctaggctacatacttgccgcaccggtatgaggggatatagtactcgcactgcgctgtctagtgagatgggcagtgctgcccataaac
aactggctgctcagccatttgttggcggaccattctgggggggccagcaatgcctgactttcgggtagggtgaaaactgaacaaagac
taccaaaacagaatttcttcctccttggaggtaagcgcaggccggcccgcctgcgcccacatggcgctccgaacacctccatagctgt
aagggcgcaaacatggccggactgttgtcagcactctttcatggccatacaaggtcatgtcgagattagtgctgagtaagacactatca
ccccatgttcgattgaagccgtgacttcatgccaacctgcccctgggcgtagcagacgtatgccatcatgaccactagccgacatgcg
ctgtcttttgccaccaaaacaactggtacaccgctcgaagtcgtgccgcacacctccgggagtgagtccggcgactcctccccggcg
ggccgcggccctacctgggtagggtcgccatacgcccacgaccaaacgacgcaggaggggattggggtagggaatcccaaccag
cctaaccaagacggcacctataataataggtgggggggactaacagccctatatcgcaagctttgggtgcctatcttgagaagcacgag
ttggagtggctgtgtacggtcgaccctaaggtgggtgtgccgcagcctgaaacaaagcgtctagcagctgcttctataatgtgtcagcc
gttgtgtttcagttatattgtatgctattgtttgttcgtgctagggtggcgcaggcccacctactgtggcgggccattggttggtgcttgaatt
gcctcaccatctaaggtctgaacgctcactcaaacgcctttgtacaactgcagaactttccttggcgctgcaactacagtgtgcaaacca
gcacatagcactcccttacatcacccagcagtacaaca

SEQ ID NO: 2
Chlorella ellipsoidea nitrate reductase promoter from AY307383

cgctgcgcaccagggccgccagctcgctgatgtcgctccaaatgcggtcccccgatttttgttcttcatcttctccaccttggtggccttc
ttggccagggccttcagctgcatgcgcacagaccgttgagctcctgatcagcatcctcaggaggccctttgacaagcaagcccctgtg
caagcccattcacggggtaccagtggtgctgaggtagatgggtttgaaaaggattgctcggtcgattgctgctcatggaattggcatgt
gcatgcatgttcacaatatgccaccaggctttggagcaagagagcatgaatgccttcaggcaggttgaaagttcctgggggtgaagag
gcagggccgaggattggaggaggaaagcatcaagtcgtcgctcatgctcatgttttcagtcagagtttgccaagctcacaggagcag
agacaagactggctgctcaggtgttgcatcgtgtgtgtggtgggggggggggggttaatacggtacgaaatgcacttggaattcccac
ctcatgccagcggacccacatgcttgaattcgaggcctgtggggtgagaaatgctcactctgccctcgttgctgaggtacttcaggccg
ctgagctcaaagtcgatgccctgctcgtctatcagggcctgcacctctgggctgaccggctcagcctccttcgcgggcatggagtagg
cgccggcagcgttcatgtccgggcccagggcagcggtggtgccataaatgtcggtgatggtggggagggggccgtcgccacacc
attgccgttgctggctgacgcatgcacatgtggcctggctggcaccggcagcactggtctccagccagccagcaagtggctgttcag
gaaagcggccatgttgttggtccctgcgcatgtaattccccagatcaaaggagggaacagcttggatttgatgtagtgcccaaccgga
ctgaatgtgcgatggcaggtccctttgagtctcccgaattactagcagggcactgtgacctaacgcagcatgccaaccgcaaaaaaat
gattgacagaaaatgaagcggtgtgtcaatatttgctgtatttattcgttttaatcagcaaccaagttcgaaacgcaactatcgtggtgatca
agtgaacctcatcagacttacctcgttcggcaaggaaacggaggcaccaaattccaatttgatattatcgcttgccaagctagagctgat
ctttgggaaaccaactgccagacagtggactgtgatggagtgccccgagtggtggagcctcttcgattcggttagtcattactaacgtg
aaccctcagtgaagggaccatcagaccagaaagaccagatctcctcctcgacaccgagagagtgttgcggcagtaggacgacaag

SEQ ID NO: 3
Yeast sucrose invertase

MTNETSDRPLVHFTPNKGWMNDPNGLWYDEKDAKWHLYFQYNPNDTVWGTPLF
WGHATSDDLTNWEDQPIAIAPKRNDSGAFSGSMVVDYNNTSGFFNDTIDPRQRCVAI
WTYNTPESEEQYISYSLDGGYTFTEYQKNPVLAANSTQFRDPKVFWYEPSQKWIMT
AAKSQDYKIEIYSSDDLKSWKLESAFANEGFLGYQYECPGLIEVPTEQDPSKSYWVM
FISINPGAPAGGSFNQYFVGSFNGTHFEAFDNQSRVVDFGKDYYALQTFFNTDPTYGS
ALGIAWASNWEYSAFVPTNPWRSSMSLVRKFSLNTEYQANPETELINLKAEPILNISN
AGPWSRFATNTTLTKANSYNVDLSNSTGTLEFELVYAVNTTQTISKSVFADLSLWFK

GLEDPEEYLRMGFEVSASSFFLDRGNSKVKFVKENPYFTNRMSVNNQPFKSENDLSY
YKVYGLLDQNILELYFNDGDVVSTNTYFMTTGNALGSVNMTTGVDNLFYIDKFQVR
EVK

SEQ ID NO: 4
Yeast secretion signal

MLLQAFLFLLAGFAAKISAS

SEQ ID NO: 5
Higher plants secretion signal

MANKSLLLLLLLGSLASG

SEQ ID NO: 6
Consensus eukaryotic secretion signal

MARLPLAALG

SEQ ID NO:7
Combination higher plant/eukaryotic secretion signal

MANKLLLLLLLLLPLAASG

SEQ ID NO: 8
S. cerevisiae sucrose invertase NP_012104

gaattccccaacatggtggagcacgacactctcgtctactccaagaatatcaaagatacagtctcagaagaccaaagggctattgaga
cttttcaacaaagggtaatatcgggaaacctcctcggattccattgcccagctatctgtcacttcatcaaaaggacagtagaaaaggaag
gtggcacctacaaatgccatcattgcgataaaggaaaggctatcgttcaagatgcctctgccgacagtggtcccaaagatggacccccc
acccacgaggagcatcgtggaaaaagaagacgttccaaccacgtcttcaaagcaagtggattgatgtgaacatggtggagcacgac
actctcgtctactccaagaatatcaaagatacagtctcagaagaccaaagggctattgagacttttcaacaaagggtaatatcgggaaac
ctcctcggattccattgcccagctatctgtcacttcatcaaaaggacagtagaaaaggaaggtggcacctacaaatgccatcattgcgat
aaaggaaaggctatcgttcaagatgcctctgccgacagtggtcccaaagatggacccccacccacgaggagcatcgtggaaaaaga
agacgttccaaccacgtcttcaaagcaagtggattgatgtgatatctccactgacgtaagggatgacgcacaatcccactatccttcgca
agacccttcctctatataaggaagttcatttcatttggagaggacacgctgaaatcaccagtctctctctacaaatctatctctggcgcgcc
atatcaATGCTTCTTCAGGCCTTTCTTTTTCTTCTTGCTGGTTTTGCTGCCAAGATCAG
CGCCTCTATGACGAACGAAACCTCGGATAGACCACTTGTGCACTTTACACCAAAC
AAGGGCTGGATGAATGACCCCAATGGACTGTGGTACGACGAAAAAGATGCCAAG
TGGCATCTGTACTTTCAATACAACCCGAACGATACTGTCTGGGGGACGCCATTGT
TTTGGGGCCACGCCACGTCCGACGACCTGACCAATTGGGAGGACCAACCAATAG
CTATCGCTCCGAAGAGGAACGACTCCGGAGCATTCTCGGGTTCCATGGTGGTTGA
CTACAACAATACTTCCGGCTTTTTCAACGATACCATTGACCCGAGACAACGCTGC
GTGGCCATATGGACTTACAACACACCGGAGTCCGAGGAGCAGTACATCTCGTAT
AGCCTGGACGGTGGATACACTTTTACAGAGTATCAGAAGAACCCTGTGCTTGCTG
CAAATTCGACTCAGTTCCGAGATCCGAAGGTCTTTTGGTACGAGCCCTCGCAGAA
GTGGATCATGACAGCGGCAAAGTCACAGGACTACAAGATCGAAATTTACTCGTC
TGACGACCTTAAATCCTGGAAGCTCGAATCCGCGTTCGCAAACGAGGGCTTTCTC
GGCTACCAATACGAATGCCCAGGCCTGATAGAGGTCCCAACAGAGCAAGATCCC
AGCAAGTCCTACTGGGTGATGTTTATTTCCATTAATCCAGGAGCACCGGCAGGAG
GTTCTTTTAATCAGTACTTCGTCGGAAGCTTTAACGGAACTCATTTCGAGGCATTT
GATAACCAATCAAGAGTAGTTGATTTTGGAAAGGACTACTATGCCCTGCAGACTT

TCTTCAATACTGACCCGACCTATGGGAGCGCTCTTGGCATTGCGTGGGCTTCTAA
CTGGGAGTATTCCGCATTCGTTCCTACAAACCCTTGGAGGTCCTCCATGTCGCTC
GTGAGGAAATTCTCTCTCAACACTGAGTACCAGGCCAACCCGGAAACCGAACTC
ATAAACCTGAAAGCCGAACCGATCCTGAACATTAGCAACGCTGGCCCCTGGAGC
CGGTTTGCAACCAACACCACGTTGACGAAAGCCAACAGCTACAACGTCGATCTTT
CGAATAGCACCGGTACACTTGAATTTGAACTGGTGTATGCCGTCAATACCACCCA
AACGATCTCGAAGTCGGTGTTCGCGGACCTCTCCCTCTGGTTTAAAGGCCTGGAA
GACCCCGAGGAGTACCTCAGAATGGGTTTCGAGGTTTCTGCGTCCTCCTTCTTCCT
TGATCGCGGGAACAGCAAAGTAAAATTTGTTAAGGAGAACCCATATTTTACCAA
CAGGATGAGCGTTAACAACCAACCATTCAAGAGCGAAAACGACCTGTCGTACTA
CAAAGTGTATGGTTTGCTTGATCAAAATATCCTGGAACTCTACTTCAACGATGGT
GATGTCGTGTCCACCAACACATACTTCATGACAACCGGGAACGCACTGGGCTCCG
TGAACATGACGACGGGTGTGGATAACCTGTTCTACATCGACAAATTCCAGGTGA
GGGAAGTCAAGTGAgatctgtcgatcgacaagctcgagtttctccataataatgtgtgagtagttcccagataagggaatta
gggttcctatagggtttcgctcatgtgttgagcatataagaaacccttagtatgtatttgtatttgtaaaatacttctatcaataaaatttctaatt
cctaaaaccaaaatccagtactaaaatccagatcccccgaattaa

SEQ ID NO: 9
TGTTGAAGAATGAGCCGGCGAC

SEQ ID NO: 10
CAGTGAGCTATTACGCACTC

SEQ ID NO: 11
UTEX 329 *Prototheca kruegani*

TGTTGAAGAATGAGCCGGCGAGTTAAAAAGAGTGGCATGGTTAAAGAAAATACT
CTGGAGCCATAGCGAAAGCAAGTTTAGTAAGCTTAGGTCATTCTTTTTAGACCCG
AAACCGAGTGATCTACCCATGATCAGGGTGAAGTGTTAGTAAAATAACATGGAG
GCCCGAACCGACTAATGTTGAAAAATTAGCGGATGAATTGTGGGTAGGGGCGAA
AAACCAATCGAACTCGGAGTTAGCTGGTTCTCCCCGAAATGCGTTTAGGCGCAGC
AGTAGCAGTACAAATAGAGGGGTAAAGCACTGTTTCTTTTGTGGGCTTCGAAAGT
TGTACCTCAAAGTGGCAAACTCTGAATACTCTATTTAGATATCTACTAGTGAGAC
CTTGGGGGATAAGCTCCTTGGTCAAAAGGGAAACAGCCCAGATCACCAGTTAAG
GCCCCAAAATGAAAATGATAGTGACTAAGGATGTGGGTATGTCAAAACCTCCAG
CAGGTTAGCTTAGAAGCAGCAATCCTTTCAAGAGTGCGTAATAGCTCACTG

SEQ ID NO: 12
UTEX 1440 *Prototheca wickerhamii*
TGTTGAAGAATGAGCCGGCGACTTAAAATAAATGGCAGGCTAAGAGATTTAATA
ACTCGAAACCTAAGCGAAAGCAAGTCTTAATAGGGCGTCAATTTAACAAAACTT
TAAATAAATTATAAAGTCATTTATTTTAGACCCGAACCTGAGTGATCTAACCATG
GTCAGGATGAAACTTGGGTGACACCAAGTGGAAGTCCGAACCGACCGATGTTGA
AAAATCGGCGGATGAACTGTGGTTAGTGGTGAAATACCAGTCGAACTCAGAGCT
AGCTGGTTCTCCCCGAAATGCGTTGAGGCGCAGCAATATATCTCGTCTATCTAGG
GGTAAAGCACTGTTTCGGTGCGGGCTATGAAAATGGTACCAAATCGTGGCAAAC
TCTGAATACTAGAAATGACGATATATTAGTGAGACTATGGGGGATAAGCTCCAT
AGTCGAGAGGGAAACAGCCCAGACCACCAGTTAAGGCCCCAAAATGATAATGAA
GTGGTAAAGGAGGTGAAAATGCAAATACAACCAGGAGGTTGGCTTAGAAGCAGC
CATCCTTTAAAGAGTGCGTAATAGCTCACTG

SEQ ID NO: 13
UTEX 1442 *Prototheca stagnora*
TGTTGAAGAATGAGCCGGCGAGTTAAAAAAAATGGCATGGTTAAAGATATTTCT
CTGAAGCCATAGCGAAAGCAAGTTTTACAAGCTATAGTCATTTTTTTTAGACCCG
AAACCGAGTGATCTACCCATGATCAGGGTGAAGTGTTGGTCAAATAACATGGAG
GCCCGAACCGACTAATGGTGAAAAATTAGCGGATGAATTGTGGGTAGGGGCGAA
AAACCAATCGAACTCGGAGTTAGCTGGTTCTCCCCGAAATGCGTTTAGGCGCAGC
AGTAGCAACACAAATAGAGGGGTAAAGCACTGTTTCTTTTGTGGGCTTCGAAAGT
TGTACCTCAAAGTGGCAAACTCTGAATACTCTATTTAGATATCTACTAGTGAGAC
CTTGGGGGATAAGCTCCTTGGTCAAAAGGGAAACAGCCCAGATCACCAGTTAAG
GCCCCAAAATGAAAATGATAGTGACTAAGGACGTGAGTATGTCAAAACCTCCAG
CAGGTTAGCTTAGAAGCAGCAATCCTTTCAAGAGTGCGTAATAGCTCACTG

SEQ ID NO: 14
UTEX 288 *Prototheca moriformis*
TGTTGAAGAATGAGCCGGCGAGTTAAAAAGAGTGGCATGGTTAAAGATAATTCT
CTGGAGCCATAGCGAAAGCAAGTTTAACAAGCTAAAGTCACCCTTTTTAGACCCG
AAACCGAGTGATCTACCCATGATCAGGGTGAAGTGTTGGTAAAATAACATGGAG
GCCCGAACCGACTAATGGTGAAAAATTAGCGGATGAATTGTGGGTAGGGGCGAA
AAACCAATCGAACTCGGAGTTAGCTGGTTCTCCCCGAAATGCGTTTAGGCGCAGC
AGTAGCAACACAAATAGAGGGGTAAAGCACTGTTTCTTTTGTGGGCTTCGAAAGT
TGTACCTCAAAGTGGCAAACTCTGAATACTCTATTTAGATATCTACTAGTGAGAC
CTTGGGGGATAAGCTCCTTGGTCAAAAGGGAAACAGCCCAGATCACCAGTTAAG
GCCCCAAAATGAAAATGATAGTGACTAAGGATGTGGGTATGTTAAAACCTCCAG
CAGGTTAGCTTAGAAGCAGCAATCCTTTCAAGAGTGCGTAATAGCTCACTG

SEQ ID NO: 15
UTEX 1439, UTEX 1441, UTEX 1435, UTEX 1437 *Prototheca moriformis*
TGTTGAAGAATGAGCCGGCGACTTAAAATAAATGGCAGGCTAAGAGAATTAATA
ACTCGAAACCTAAGCGAAAGCAAGTCTTAATAGGGCGCTAATTTAACAAAACAT
TAAATAAAATCTAAAGTCATTTATTTTAGACCCGAACCTGAGTGATCTAACCATG
GTCAGGATGAAACTTGGGTGACACCAAGTGGAAGTCCGAACCGACCGATGTTGA
AAAATCGGCGGATGAACTGTGGTTAGTGGTGAAATACCAGTCGAACTCAGAGCT
AGCTGGTTCTCCCCGAAATGCGTTGAGGCGCAGCAATATATCTCGTCTATCTAGG
GGTAAAGCACTGTTTCGGTGCGGGCTATGAAAATGGTACCAAATCGTGGCAAAC
TCTGAATACTAGAAATGACGATATATTAGTGAGACTATGGGGGATAAGCTCCAT
AGTCGAGAGGGAAACAGCCCAGACCACCAGTTAAGGCCCCAAAATGATAATGAA
GTGGTAAAGGAGGTGAAAATGCAAATACAACCAGGAGGTTGGCTTAGAAGCAGC
CATCCTTTAAAGAGTGCGTAATAGCTCACTG

SEQ ID NO: 16
UTEX 1533 *Prototheca wickerhamii*
TGTTGAAGAATGAGCCGTCGACTTAAAATAAATGGCAGGCTAAGAGAATTAATA
ACTCGAAACCTAAGCGAAAGCAAGTCTTAATAGGGCGCTAATTTAACAAAACAT
TAAATAAAATCTAAAGTCATTTATTTTAGACCCGAACCTGAGTGATCTAACCATG
GTCAGGATGAAACTTGGGTGACACCAAGTGGAAGTCCGAACCGACCGATGTTGA
AAAATCGGCGGATGAACTGTGGTTAGTGGTGAAATACCAGTCGAACTCAGAGCT
AGCTGGTTCTCCCCGAAATGCGTTGAGGCGCAGCAATATATCTCGTCTATCTAGG
GGTAAAGCACTGTTTCGGTGCGGGCTATGAAAATGGTACCAAATCGTGGCAAAC

TCTGAATACTAGAAATGACGATATATTAGTGAGACTATGGGGGATAAGCTCCAT
AGTCGAGAGGGAAACAGCCCAGACCACCAGTTAAGGCCCCAAAATGATAATGAA
GTGGTAAAGGAGGTGAAAATGCAAATACAACCAGGAGGTTGGCTTAGAAGCAGC
CATCCTTTAAAGAGTGCGTAATAGCTCACTG

SEQ ID NO: 17
UTEX 1434 *Prototheca moriformis*
TGTTGAAGAATGAGCCGGCGAGTTAAAAAGAGTGGCGTGGTTAAAGAAAATTCT
CTGGAACCATAGCGAAAGCAAGTTTAACAAGCTTAAGTCACTTTTTTTAGACCCG
AAACCGAGTGATCTACCCATGATCAGGGTGAAGTGTTGGTAAAATAACATGGAG
GCCCGAACCGACTAATGGTGAAAAATTAGCGGATGAATTGTGGGTAGGGGCGAA
AAACCAATCGAACTCGGAGTTAGCTGGTTCTCCCCGAAATGCGTTTAGGCGCAGC
AGTAGCAACACAAATAGAGGGGTAAAGCACTGTTTCTTTTGTGGGCTCCGAAAG
TTGTACCTCAAAGTGGCAAACTCTGAATACTCTATTTAGATATCTACTAGTGAGA
CCTTGGGGGGATAAGCTCCTTGGTCGAAAGGGAAACAGCCCAGATCACCAGTTAA
GGCCCCAAAATGAAAATGATAGTGACTAAGGATGTGAGTATGTCAAAACCTCCA
GCAGGTTAGCTTAGAAGCAGCAATCCTTTCAAGAGTGCGTAATAGCTCACTG

SEQ ID NO: 18
UTEX 1438 *Prototheca zopfii*
TGTTGAAGAATGAGCCGGCGAGTTAAAAAGAGTGGCATGGTTAAAGAAAATTCT
CTGGAGCCATAGCGAAAGCAAGTTTAACAAGCTTAAGTCACTTTTTTTAGACCCG
AAACCGAGTGATCTACCCATGATCAGGGTGAAGTGTTGGTAAAATAACATGGAG
GCCCGAACCGACTAATGGTGAAAAATTAGCGGATGAATTGTGGGTAGGGGCGAA
AAACCAATCGAACTCGGAGTTAGCTGGTTCTCCCCGAAATGCGTTTAGGCGCAGC
AGTAGCAACACAAATAGAGGGGTAAAGCACTGTTTCTTTCGTGGGCTTCGAAAG
TTGTACCTCAAAGTGGCAAACTCTGAATACTCTATTTAGATATCTACTAGTGAGA
CCTTGGGGGATAAGCTCCTTGGTCAAAAGGGAAACAGCCCAGATCACCAGTTAA
GGCCCCAAAATGAAAATGATAGTGACTAAGGATGTGAGTATGTCAAAACCTCCA
GCAGGTTAGCTTAGAAGCAGCAATCCTTTCAAGAGTGCGTAATAGCTCACTG

SEQ ID NO: 19
UTEX 1436 *Prototheca moriformis*
TGTTGAAGAATGAGCCGGCGACTTAGAAAAGGTGGCATGGTTAAGGAAATATTC
CGAAGCCGTAGCAAAAGCGAGTCTGAATAGGGCGATAAAATATATTAATATTTA
GAATCTAGTCATTTTTTCTAGACCCGAACCCGGGTGATCTAACCATGACCAGGAT
GAAGCTTGGGTGATACCAAGTGAAGGTCCGAACCGACCGATGTTGAAAAATCGG
CGGATGAGTTGTGGTTAGCGGTGAAATACCAGTCGAACCCGGAGCTAGCTGGTT
CTCCCCGAAATGCGTTGAGGCGCAGCAGTACATCTAGTCTATCTAGGGGTAAAGC
ACTGTTTCGGTGCGGGCTGTGAGAACGGTACCAAATCGTGGCAAACTCTGAATAC
TAGAAATGACGATGTAGTAGTGAGACTGTGGGGGATAAGCTCCATTGTCAAGAG
GGAAACAGCCCAGACCACCAGCTAAGGCCCCAAAATGGTAATGTAGTGACAAAG
GAGGTGAAAATGCAAATACAACCAGGAGGTTGGCTTAGAAGCAGCCATCCTTTA
AAGAGTGCGTAATAGCTCACTG

SEQ ID NO: 20
Chicorium intybus invertase: Genbank Accession No. Y11124

MSNSSNASESLFPATSEQPYRTAFHFQPPQNWMNDPNGPMCYNGVYHLFYQYNPFG
PLWNLRMYWAHSVSHDLINWIHLDLAFAPTEPFDINGCLSGSATVLPGNKPIMLYTG

IDTENRQVQNLAVPKDLSDPYLREWVKHTGNPIISLPEEIQPDDFRDPTTTWLEEDGT
WRLLVGSQKDKTGIAFLYHSGDFVNWTKSDSPLHKVSGTGMWECVDFFPVWVDST
NGVDTSIINPSNRVKHVLKLGIQDHGKDCYLIGKYSADKENYVPEDELTLSTLRLDY
GMYYASKSFFDPVKNRRIMTAWVNESDSEADVIARGWSGVQSFPRSLWLDKNQKQ
LLQWPIEEIEMLHQNEVSFHNKKLDGGSSLEVLGITASQADVKISFKLANLEEAEELD
PSWVDPQLICSENDASKKGKFGPFGLLALASSDLREQTAIFFRVFRKNGRYVVLMCS
DQSRSSMKNGIEKRTYGAFVDIDPQQDEISLRTLIDHSIVESFGGRGKTCITTRVYPTL
AIGEQARLFAFNHGTESVEISELSAWSMKKAQMKVEEP

SEQ ID NO: 21
Schizosaccharomyces pombe Invertase: Genbank Accession No. AB011433

MFLKYILASGICLVSLLSSTNAAPRHLYVKRYPVIYNASNITEVSNSTTVPPPPFVNTT
APNGTCLGNYNEYLPSGYYNATDRPKIHFTPSSGFMNDPNGLVYTGGVYHMFFQYS
PKTLTAGEVHWGHTVSKDLIHWENYPIAIYPDEHENGVLSLPFSGSAVVDVHNSSGL
FSNDTIPEERIVLIYTDHWTGVAERQAIAYTTDGGYTFKKYSGNPVLDINSLQFRDPK
VIWDFDANRWVMIVAMSQNYGIAFYSSYDLIHWTELSVFSTSGYLGLQYECPGMAR
VPVEGTDEYKWVLFISINPGAPLGGSVVQYFVGDWNGTNFVPDDGQTRFVDLGKDF
YASALYHSSSANADVIGVGWASNWQYTNQAPTQVFRSAMTVARKFTLRDVPQNPM
TNLTSLIQTPLNVSLLRDETLFTAPVINSSSSLSGSPITLPSNTAFEFNVTLSINYTEGCT
TGYCLGRIIIDSDDPYRLQSISVDVDFAASTLVINRAKAQMGWFNSLFTPSFANDIYIY
GNVTLYGIVDNGLLELYVNNGEKTYTNDFFFLQGATPGQISFAAFQGVSFNNVTVTP
LKTIWNC

SEQ ID NO: 22
Picha anomala beta-fructofuranosidase (invertase): Genbank Accession No. X80640

MIQLSPLLLLPLFSVFNSIADASTEYLRPQIHLTPDQGWMNDPNGMFYDRKDKLWHV
YFQHNPDKKSIWATPVTWGHSTSKDLLTWDYHGNALEPENDDEGIFSGSVVVDRNN
TSGFFNDSTDPEQRIVAIYTNNAQLQTQEIAYSLDKGYSFIKYDQNPVINVNSSQQRD
PKVLWHDESNQWIMVVAKTQEFKVQIYGSPDLKKWDLKSNFTSNGYLGFQYECPG
LFKLPIENPLNDTVTSKWVLLLAINPGSPLGGSINEYFIGDFDGTTFHPDDGATRFMDI
GKDFYAFQSFDNTEPEDGALGLAWASNWQYANTVPTENWRSSMSLVRNYTLKYVD
VNPENYGLTLIQKPVYDTKETRLNETLKTLETINEYEVNDLKLDKSSFVATDFNTERN
ATGVFEFDLKFTQTDLKMGYSNMTTQFGLYIHSQTVKGSQETLQLVFDTLSTTWYID
RTTQHSFQRNSPVFTERISTYVEKIDTTDQGNVYTLYGVVDRNILELYFNDGSIAMTN
TFFFREGKIPTSFEVVCDSEKSFITIDELSVRELARK

SEQ ID NO: 23
Debaryomyces occidentalis Invertase: Genbank Accession No. X17604

MVQVLSVLVIPLLTLFFGYVASSSIDLSVDTSEYNRPLIHFTPEKGWMNDPNGLFYDK
TAKLWHLYFQYNPNATAWGQPLYWGHATSNDLVHWDEHEIAIGPEHDNEGIFSGSI
VVDHNNTSGFFNSSIDPNQRIVAIYTNNIPDLQTQDIAFSLDGGYTFTKYENNPVIDVS
SNQFRDPKVFWHERFKSMDHGCSEIARVKIQIFGSANLKNWVLNSNFSSGYYGNQY
GMSRLIEVPIENSDKSKWVMFLAINPGSPLGGSINQYFVGDFDGFQFVPDDSQTRFVD
IGKDFYAFQTFSEVEHGVLGLAWASNWQYADQVPTNPWRSSTSLARNYTLRYVIQM
LKLTANIDKSVLPDSINVVDKLKKKNVKLTNKKPIKTNFKGSTGLFDFNITFKVLNLN
VSPGKTHFDILINSQELNSSVDSIKIGFDSSQSLFYIDRHIPNVEFPRKQFFTDKLAAYL

EPLDYDQDLRVFSLYGIVDKNIIELYFNDGTVAMTNTFFMGEGKYPHDIQIVTDTEEP
LFELESVIIRELNK

SEQ ID NO: 24
Oryza sativa Invertase: Genbank Accession No. AF019113

MATSRLTPAYDLKNAAAAVYTPLPEQPHSAEVEIRDRKPFKIISAIILSSLLLLALILVA
VNYQAPPSHSSGDNSQPAAVMPPSRGVSQGVSEKAFRGASGAGNGVSFAWSNLMLS
WQRTSYHFQPVKNWMNDPNGPLYYKGWYHLFYQYNPDSAVWGNITWGHAVSTD
LINWLHLPFAMVPDQWYDVNGVWTGSATILPDGRIVMLYTGDTDDYVQDQNLAFP
ANLSDPLLVDWVKYPNNPVIYPPPGIGVKDFRDPTTAGTAGMQNGQRLVTIGSKVG
KTGISLVYETTNFTTFKLLYGVLHAVPGTGMWECVDLYPVSTTGENGLDTSVNGLG
VKHVLKTSLDDDKHDYYALGTYDPVKNKWTPDNPDLDVGIGLRLDYGKYYAARTF
YDQNKQRRILWGWIGETDLEAVDLMKGWASLQAIPRTIVFDKKTGTNVLQRPEEEV
ESWSSGDPITQRRIFEPGSVVPIHVSGATQLDITASFEVDETLLETTSESHDAGYDCSN
SGGAGTRGSLGPFGLLVVADEKLSELTPVYLYVAKGGDGKAKAHLCAYQTRSSMAS
GVEKEVYGSAVPVLDGENYSARILIDHSIVESFAQAGRTCVRSRDYPTKDIYGAARCF
FFNNATEASVRASLKAWQMKSFIRPYPFIPDQKS

SEQ ID NO: 25
Allium cepa Invertase: Genbank Accession No. AJ006067

MSSDDLESPPSSYLPIPPSDEFHDQPPPLRSWLRLLSIPLALMFLLFLATFLSNLESPPSD
SGLVSDPVTFDVNPAVVRRGKDAGVSDKTSGVDSGFVLDPVAVDANSVVVHRGKD
AGVSDKTSGVDSGLLKDSPLGPYPWTNQMLSWQRTGFHFQPVKNWMNDPNGPLYY
KGWYHFFYQYNPEGAVWGNIAWGHAVSRDLVHWTHLPLAMVPDQWYDINGVWT
GSATILPDGQIVMLYTGATNESVQVQNLAVPADQSDTLLLRWKKSEANPILVPPPGIG
DKDFRDPTTAWYEPSDDTWRIVIGSKDSSHSGIAIVYSTKDFINYKLIPGILHAVERVG
MWECVDFYPVATADSSHANHGLDPSARPSPAVKHVLKASMDDDRHDYYAIGTYDP
AQNTWVPDDASVDVGIGLRYDWGKFYASKTFYDHAKKRRILWSWIGETDSETADIA
KGWASLQGVPRTVLLDVKTGSNLITWPVVEIESLRTRPRDFSGITVDAGSTFKLDVG
GAAQLDIEAEFKISSEELEAVKEADVSYNCSSSGGAAERGVLGPFGLLVLANQDLTE
QTATYFYVSRGMDGGLNTHFCQDEKRSSKASDIVKRIVGHSVPVLDGESFALRILVD
HSIVESFAQGGRASATSRVYPTEAIYNNARVFVFNNATGAKVTAQSLKVWHMSTAI
NEIYDPATSVM

SEQ ID NO: 26
Beta vulgaris subsp. vulgaris Invertase: Genbank Accession No. AJ278531

LFYQYNPNGVIWGPPVWGHSTSKDLVNWVPQPLTMEPEMAANINGSWSGSATILPG
NKPAILFTGLDPKYEQVQVLAYPKDTSDPNLKEWFLAPQNPVMFPTPQNQINATSFR
DPTTAWRLPDGVWRLLIGSKRGQRGLSLLFRSRDFVHWVQAKHPLYSDKLSGMWE
CPDFFPVYANGDQMGVDTSIIGSHVKHVLKNSLDITKHDIYTIGDYNIKKDAYTPDIG
YMNDSSLRYDYGKYYASKTFFDDAKKERILLGWANESSSVEDDIKKGWSGIHTIPRK
IWLDKLGKQLIQWPIANIEKLRQKPVNIYRKVLKGGSQIEVSGITAAQADVEISFKIKD
LKNVEKFDASWTSPQLLCSKKGASVKGGLGPFGLLTLASXGLEEYTAVFFRIFKAYD
NKFVVLMCSDQSRSSLNPTNDKTTYGTFVDVNPIREGLSLRVLIDHSVVESFGAKGK
NVITARVYPTLAINEKAHLYVFNRGTSNVEITGLTAWSMKKANIA

SEQ ID NO: 27

Bifidobacterium breve UCC2003 beta-fructofuranosidase (invertase): Genbank Accession No. AAT28190

MTDFTPETPVLTPIRDHAAELAKAEAGVAEMAAKRNNRWYPKYHIASNGGWINDPN
GLCFYKGRWHVFYQLHPYGTQWGPMHWGHVSSTDMLNWKREPIMFAPSLEQEKD
GVFSGSAVIDDNGDLRFYYTGHRWANGHDNTGGDWQVQMTALPDNDELTSATKQ
GMIIDCPTDKVDHHYRDPKVWKTGDTWYMTFGVSSEDKRGQMWLFSSKDMVRWE
YERVLFQHPDPDVFMLECPDFFPIKDKDGNEKWVIGFSAMGSKPSGFMNRNVNNAG
YMIGTWEPGGEFKPETEFRLWDCGHNYYAPQSFNVDGRQIVYGWMSPFVQPIPMED
DGWCGQLTLPREITLDDDGDVVTAPVAEMEGLREDTLDHGSITLDMDGEQVIADDA
EAVEIEMTIDLAASTADRAGLKIHATEDGAYTYVAYDDQIGRVVVDRQAMANGDH
GYRAAPLTDAELASGKLDLRVFVDRGSVEVYVNGGHQVLSSYSYASEGPRAIKLVA
EFGNLKVESLKLHHMKSIGLE

SEQ ID NO: 28
Saccharomyces cerevisiae Invertase: Genbank Accession No. NP_012104
MLLQAFLFLLAGFAAKISASMTNETSDRPLVHFTPNKGWMNDPNGLWYDEKDAKW
HLYFQYNPNDTVWGTPLFWGHATSDDLTNWEDQPIAIAPKRNDSGAFSGSMVVDY
NNTSGFFNDTIDPRQRCVAIWTYNTPESEEQYISYSLDGGYTFTEYQKNPVLAANSTQ
FRDPKVFWYEPSQKWIMTAAKSQDYKIEIYSSDDLKSWKLESAFANEGFLGYQYECP
GLIEVPTEQDPSKSYWVMFISINPGAPAGGSFNQYFVGSFNGTHFEAFDNQSRVVDFG
KDYYALQTFFNTDPTYGSALGIAWASNWEYSAFVPTNPWRSSMSLVRKFSLNTEYQ
ANPETELINLKAEPILNISNAGPWSRFATNTTLTKANSYNVDLSNSTGTLEFELVYAV
NTTQTISKSVFADLSLWFKGLEDPEEYLRMGFEVSASSFFLDRGNSKVKFVKENPYFT
NRMSVNNQPFKSENDLSYYKVYGLLDQNILELYFNDGDVVSTNTYFMTTGNALGSV
NMTTGVDNLFYIDKFQVREVK

SEQ ID NO: 29
Zymomonas mobilis Invertase A: Genbank Accession No. AY171597

MESPSYKNLIKAEDAQKKAGKRLLSSEWYPGFHVTPLTGWMNDPNGLIFFKGEYHL
FYQYYPFAPVWGPMHWGHAKSRDLVHWETLPVALAPGDLFDRDGCFSGCAVDNN
GVLTLIYTGHIVLSNDSPDAIREVQCMATSIDGIHFQKEGIVLEKAPMPQVAHFRDPR
VWKENDHWFMVVGYRTDDEKHQGIGHVALYRSENLKDWIFVKTLLGDNSQLPLGK
RAFMWECPDFFSLGNRSVLMFSPQGLKASGYKNRNLFQNGYILGKWQAPQFTPETS
FQELDYGHDFYAAQRFEAKDGRQILIAWFDMWENQKPSQRDGWAGCMTLPRKLDL
IDNKIVMTPVREMEILRQSEKIESVVTLSDAEHPFTMDSPLQEIELIFDLEKSSAYQAG
LALRCNGKGQETLLYIDRSQNRIILDRNRSGQNVKGIRSCPLPNTSKVRLHIFLDRSSIE
IFVGDDQTQGLYSISSRIFPDKDSLKGRLFAIEGYAVFDSFKRWTLQDANLAAFSSDA
C

SEQ ID NO:30
SAG 2214 *Chlorella luteoviridis*

TGTTGAAGAATGAGCCGGCGACTTATAGGAAGTGGCTTGGTTAAGGATACTTTCC
GAAGCCTAAGCGAAAGCAAGTTGTAACAATAGCGATATACCTCTTTGTAGGTCA
GTCACTTCTTATGGACCCGAACCCGGGTGATCTAACCATGACCAGGATGAAGCTT
GGGTAACACCAAGTGAAGGTCCGAACTCTTCGATCTTTAAAAATCGTGAGATGA
GTTATGGTTAGGGGTAAATCTGGCAGTTTTGCCCCGCAAAAGGGTAACCTTTTGT
AATTACTGACTCATAACGGTGAAGCCTAAGGCGTTAGCTATGGTAATACCGTGGG

AAGTTTCAATACCTTCTTGCATATTTTTTATTTGCACCTTTAGTGCAAACAGTGTA
AAGAAAGCGTTTTGAAACCCCTTAACGACTAATTTTTTGCTTTTGCAAGAACGTC
AGCACTCACCAATACACTTTCCGTTTTTTTCTTTTATTAATTAAAGCAACATAAAA
ATATATTTTATAGCTTTAATCATAAAACTATGTTAGCACTTCGTGCTAATGTGCTA
ATGTGCTAATCAAATGAAAAGTGTTCTTAAAAGTGAGTTGAAGGTAGAGTCTAAT
CTTGCCTGAAAGGGCAAGCTGCACATTTTTTTTTGAATGTGCAACAATGGAAATG
CCAATCGAACTCGGAGCTAGCTGGTTCTCCCCGAAATGTGTTGAGGCGCAGCGAT
TCATGATTAGTACGGTGTAGGGGTAAAGCACTGTTTCGGTGCGGGCTGTGAAAAC
GGTACCAAATCGTGGCAAACTAAGAATACTACGCTTGTATACCATGGATCAGTG
AGACTATGGGGGATAAGCTCCATAGTCAAGAGGGAAACAGCCCAGATCACCAGT
TAAGGCCCCAAAATGACAGCTAAGTGGCAAAGGAGGTGAAAGTGCAGAAACAA
CCAGGAGGTTTGCCCAGAAGCAGCCATCCTTTAAAGAGTGCGTAATAGCTCACTG

SEQ ID NO: 31
Endogenous *Chlorella protothecoides* actin promoter and 5'UTR

GAATTCGAGTTTAGGTCCAGCGTCCGTGGGGGGGGACGGGCTGGGAGCTTGGGC
CGGGAAGGGCAAGACGATGCAGTCCCTCTGGGGAGTCACAGCCGACTGTGTGTG
TTGCACTGTGCGGCCCGCAGCACTCACACGCAAAATGCCTGGCCGACAGGCAGG
CCCTGTCCAGTGCAACATCCACGGTCCCTCTCATCAGGCTCACCTTGCTCATTGAC
ATAACGGAATGCGTACCGCTCTTTCAGATCTGTCCATCCAGAGAGGGGAGCAGG
CTCCCCACCGACGCTGTCAAACTTGCTTCCTGCCCAACCGAAAACATTATTGTTT
GAGGGGGGGGGGGGGGGGGCAGATTGCATGGCGGGATATCTCGTGAGGAACAT
CACTGGGACACTGTGGAACACAGTGAGTGCAGTATGCAGAGCATGTATGCTAGG
GGTCAGCGCAGGAAGGGGGCCTTTCCCAGTCTCCCATGCCACTGCACCGTATCCA
CGACTCACCAGGACCAGCTTCTTGATCGGCTTCCGCTCCCGTGGACACCAGTGTG
TAGCCTCTGGACTCCAGGTATGCGTGCACCGCAAAGGCCAGCCGATCGTGCCGAT
TCCTGGGGTGGAGGATATGAGTCAGCCAACTTGGGGCTCAGAGTGCACACTGGG
GCACGATACGAAACAACATCTACACCGTGTCCTCCATGCTGACACACCACAGCTT
CGCTCCACCTGAATGTGGGCGCATGGGCCCGAATCACAGCCAATGTCGCTGCTGC
CATAATGTGATCCAGACCCTCTCCGCCCAGATGCCGAGCGGATCGTGGGCGCTGA
ATAGATTCCTGTTTCGATCACTGTTTGGGTCCTTTCCTTTTCGTCTCGGATGCGCG
TCTCGAAACAGGCTGCGTCGGGCTTTCGGATCCCTTTTGCTCCCTCCGTCACCATC
CTGCGCGCGGGCAAGTTGCTTGACCCTGGGCTGGTACCAGGGTTGGAGGGTATTA
CCGCGTCAGGCCATTCCCAGCCCGGATTCAATTCAAAGTCTGGGCCACCACCCTC
CGCCGCTCTGTCTGATCACTCCACATTCGTGCATACACTACGTTCAAGTCCTGATC
CAGGCGTGTCTCGGGACAAGGTGTGCTTGAGTTTGAATCTCAAGGACCCACTCCA
GCACAGCTGCTGGTTGACCCCGCCCTCGCAACTCCCTACC***ATGTCTGCTG***gtaggtcca
gggatctttgccatgcacacaggaccccgtttgtgggggtccccggtgcatgctgtcgctgtgcaggcgccggtgtggggcctgggc
cccgcgggagctcaactcctccccatatgcctgccgtccctcccacccaccgcgacctggccccctttgcag*AGGAAGGCGA*
AGTCAGCGCCATCGTGTGCGATAATGGATCCGG

SEQ ID NO: 32
Endogenous *Chlorella protothecoides* EF1a promoter and 5'UTR

GAATTCGCCCTTGAGTTTAGGTCCAGCGTCCGTGGGGGGGGCGTGAGACTCCCCC
CTGACCTTCGTATGGCAGGGACTCCTACTTGCCAAGTAATCAGTTGACAATGCCA
CTTCAATGCTCGTTGTGGTACACTGACGCGGGTCTAACATACTGGGAAGCATGAA
TTGCCGACATGGACTCAGTTGGAGACAGTAACAGCTCTTTGTGTTCTATCTTCAG
GAACACATTTGGCAGCGCACCCATACAGTGGCGCACACGCAGCTGTACCTGATG

TGGCTCTATTCCCACATGTTTCAACTTGATCCAAAAGTCACTCAGACTCTCAGCA
GCTAGACTTGATCGCATCTTTGGCCATGAAGATGCTTGCGCAACTCTAGGAATGG
GACGAGAAAAGAGCCTGCTCTGATCGGATATTTCCATTCTCTGGATGGGACTGAG
ATGATTCTGAAGAAATGCTGCTCGACTTATTTGGAAGAACAGCACCTGACGCATG
CTTTGAGGCTGCTGTGGCTGGGATGTGCTGTATTTGTCAGCATTGAGCATCTACG
GGTAGATGGCCATAACCACGCGCTGCCTATCATGCGGTGGGTTGTGTGGAAAAC
GTACAATGGACAGAAATCAATCCCATTGCGAGCCTAGCGTGCAGCCATGCGCTC
CCTCTGTAGCCCCGCTCCAAGACAAAGCCAGCCAATGCCAGAACCCACATAGAG
AGGGTATCTTCCTAATGACCTCGCCCATCATTTCCTCCAAATTAACTATAATGCCT
TGATTGTGGAGTTGGCTTTGGCTTGCAGCTGCTCGCGCTGGCACTTTTGTAGGCA
GCACAGGGTATGCCAGCGCCGAACTTTGTGCCCTTGAGCAGGCCACAAGGGCAC
AAGACTACACCATGCAGCTGGTATACTTGGAACTGATACCATTCTTACCAAGCAA
GGCACAGCACAGCCTGCACCGACTCACTTTGCTTGAGCGGGGCACAGCGCCGCG
ACTGATCCTGCGAGCTGTGGGGAGTTCCGACTGTTCTGGACCTCGGTCTCTGAAA
GATGTGTACGATGGGATCAAGTCATTCAAGTATGCTCTTCACATGAGCAATCGGG
GGAGACACGGTGGCCCTAAAGGTGTTCATCTGATTCAAGTGTAGTGGGGGGGTG
CTGTTTGTCCCGGGGCGCCCCCCGCTCCCCGACCCCGGAGAAGGGCCCCAGAGG
ACTCGGCCGCCCACAGAGGAATAACCGGGCGTGGCTCGGCCCTGCGCCTCCCTCT
TTCAATATTTCACCTGGTGTTCAGTGCACGGACACGTAAAGAACTAGATACA***ATG***
*GCCGAGGGAAAGACG*gtgagagcttggcgttggtggaccgggcagcatcagaaactcctcttccccgcccgccttgaaa
ctcactgtaactccctcctcttcccccctcgcag*CATCTGTCTATCGTTATC*gtgagtgaaagggactgccatgtgtcgggt
cgttgaccacggtcggctcgggcgctgctgcccgcgtcgcgaacgttccctgcaaacgccgcgcagccgtccctttttctgccgccg
ccccacccccctcgctcccccccttcaatcacaccgcag*TGCGGACATGTCGATTCCGGCAAGTCCACC*

SEQ ID NO: 33
Endogenous *Chlorella protothecoides* beta-tubulin promoter (isoform A)
GAATTCCCTGCAGGAAGAAGGCCGGCAGCAGCTGGTACTTGTCCTTCACCTCCTT
GATCGGCTGGGTGAGCTTGGCCGGGTCGCAGTCGTCGATGCCGGCATCGCCCAG
CACGCTGTGCGGGGAGCCGGCATCGACAACCTTGGCACTGCTCACCTTGGTCACC
GGCATGGGGTCATGGCGCTGCAGACCAGCGGCCTGTCAGCATGCTGCAGGCATC
TGTGTTTTGTAGTAGATACTTTCTGATGCATCACCACACGTTTGGAAGGTCCCCA
AGCCCCTTCAACAGTCTCGACATATGACACTCGCGCCCTCTTCCTCGTCCCGTGG
CCTGATGAGGGTACGCAGGTACCGCAGCTGCGCCCCGTCCCGCCAGTTGCCCTGG
CCCCGCCGGGCCCAATCTGTTCATTGCCGCTCCCTGGCAGCCGTGAACTTCACAC
TACCGCTCTCTGTGACCTTCAGCACAGCAGGAATCGCCATTTCACCGGCGGTCGT
TGCTGCGGAGCCTCAGCTGATCTCGCCTGCGAGACCCCACAGTTTGAATTTGCGG
TCCCCACACAACCTCTGACGCC

SEQ ID NO: 34
Endogenous *Chlorella protothecoides* beta-tubulin promoter (isoform B)

GAATTCCCTCAGGAAGAAGGCCGGCAGCAGCTGGTACTTGTCCTTCACCTCCTTG
ATCGGCTGGGTGAGCTTCGCAGGATCGCAGTCGTCGATGCCGGCATCGCCCAGC
ACGCTGTGCGGGGAGCCGGCATCNACAACCTTGGCACTGCTCCCCTTGGTCACCG
GCATGGGGTCATGGCGCTGCAGCCCAGCGGCCTGTCAGCATGCTGCAGGCATCT
GTGTATTGTAGTAGGTACTTCCTGATGCATCAACACACGTTTGGAAGCTCCCCAA
GCCCCTTCAACAGTCTCGACGTATGACACTCGCGCCCTCTTCCTCGCCCCGTGGC
CTGATGAGGGTACGCAGGTACCACAGCTGCGCCCCGTCCCGCCAGTTGCCCTGGC
CCGGCCGGGCCCAATCTGTTCATTGCCGCTCCCTGGTAGCCGTGAACTCACATTA
CCGCTCTCTGTGACCTTCAGCACAGCAGGAATCGCCATTTCACCGGCGGTCGTTG

CTGCGGAGCCTCAGCTGATCTCGCCTGCGAGACCCCACAGTTTGAATTTGCGGTC
CCCACACAACCTCTGACGCC

SEQ ID NO: 35
TGACCTAGGTGATTAATTAACTCGAGGCAGCAGCAGCTCGGATAGTATCG

SEQ ID NO: 36
CTACGAGCTCAAGCTTTCCATTTGTGTTCCCATCCCACTACTTCC

SEQ ID NO: 37
GATCAGAATTCCGCCTGCAACGCAAGGGCAGC

SEQ ID NO: 38
GCATACTAGTGGCGGGACGGAGAGAGGGCG

SEQ ID NO: 39
Beta-tubulin::neo::nit

gaattcctttcttgcgctatgacacttccagcaaaaggtagggcgggctgcgagacggcttcccggcgctgcatgcaacaccgatgat
gcttcgaccccccgaagctccttcggggctgcatgggcgctccgatgccgctccagggcgagcgctgtttaaatagccaggccccc
gattgcaaagacattatagcgagctaccaaagccatattcaaacacctagatcactaccacttctacacaggccactcgagcttgtgatc
gcactccgctaagggggcgcctcttcctcttcgtttcagtcacaacccgcaaac**ggcgcgcc**atatca***atg***attgaacaagatggattg
cacgcaggttctccggccgcttgggtggagaggctattcggctatgactgggcacaacagacaatcggctgctctgatgccgccgtgt
tccggctgtcagcgcaggggcgcccggttcttttttgtcaagaccgacctgtccggtgccctgaatgaactgcaggacgaggcagcgc
ggctatcgtggctggccacgacgggcgttccttgcgcagctgtgctcgacgttgtcactgaagcgggaagggactggctgctattgg
gcgaagtgccggggcaggatctcctgtcatctcaccttgctcctgccgagaaagtatccatcatggctgatgcaatgcggcggctgca
tacgcttgatccggctacctgcccattcgaccaccaagcgaaacatcgcatcgagcgagcacgtactcggatggaagccggtcttgtc
gatcaggatgatctggacgaagagcatcaggggctcgcgccagccgaactgttcgccaggctcaaggcgcgcatgcccgacggcg
aggatctcgtcgtgacccatggcgatgcctgcttgccgaatatcatggtggaaaatggccgcttttctggattcatcgactgtggccggc
tgggtgtggcggaccgctatcaggacatagcgttggctacccgtgatattgctgaagagcttggcggcgaatgggctgaccgcttcct
cgtgctttacggtatcgccgctcccgattcgcagcgcatcgccttctatcgccttcttgacgagttcttctaagatctgtcgatcgacaagt
gactcgaggcagcagcagctcggatagtatcgacacactctggacgctggtcgtgtgatggactgttgccgccacacttgctgccttg
acctgtgaatatccctgccgcttttatcaaacagcctcagtgtgtttgatcttgtgtgtacgcgcttttgcgagttgctagctgcttgtgctatt
tgcgaataccacccccagcatccccttccctcgtttcatatcgcttgcatcccaaccgcaacttatctacgctgtcctgctatccctcagcg
ctgctcctgctcctgctcactgcccctcgcacagccttggtttgggctccgcctgtattctcctggtactgcaacctgtaaaccagcactg
caatgctgatgcacgggaagtagtgggatgggaacacaaatggaaagctt**gagctc**

SEQ ID NO: 40
Glutamate dehydrogenase::neo::nit

gaattccgcctgcaacgcaagggcagccacagccgctcccacccgccgctgaaccgacacgtgcttgggcgcctgccgcctgcct
gccgcatgcttgtgctggtgaggctgggcagtgctgccatgctgattgaggcttggttcatcgggtggaagcttatgtgtgtgctgggct
tgcatgccgggcaatgcgcatggtggcaagagggcggcagcacttgctggacgtgccgcggtgcctccaggtggttcaatcgcggc
agccagagggatttcagatgatcgcgcgtacaggttgagcagcagtgtcagcaaaggtagcagtttgccagaatgatcggttcagctg
ttaatcaatgccagcaagagaaggggtcaagtgcaaacacgggcatgccacagcacgggcaccggggagtggaatggcaccacc
aagtgtgtgcgagccagcatcgccgcctggctgtttcagctacaacggcaggagtcatccaactaaccatagctgatcaacactgcaa
tcatcggcggctgatgcaagcatcctgcaagacacatgctgtgcgatgctgcgctgctgcctgctgcgcacgccgttgagttggcagc
agctcagccatgcactggatcaggctgggctgccactgcaatgtggtggataggatgcaagtggagcgaataccaaaccctctggct
gcttgctgggttgcatggcatcgcaccatcagcaggagcgcatgcgaagggactggccccatgcacgccatgccaaaccggagcg
caccgagtgtccacactgtcaccaggcccgcaagctttgcagaaccatgctcatggacgcatgtagcgctgacgtcccttgacggcg

ctcctctcgggtgtgggaaacgcaatgcagcacaggcagcagaggcggcggcagcagagcggcggcagcagcggcgggggcc
acccttcttgcggggtcgcgccccagccagcggtgatgcgctgatcnnnccaaacgagttcacattcatttgcagcctggagaagcg
aggctggggcctttgggctggtgcagcccgcaatggaatgcgggaccgccaggctagcagcaaaggcgcctccctactccgcat
cgatgttccatagtgcattggactgcatttgggtggggcggccggctgtttctttcgtgttgcaaaacgcgccacgtcagcaacctgtcc
cgtgggtcccccgtgccgatgaaatcgtgtgcacgccgatcagctgattgcccggctcgcgaagtaggcgccctctttctgctcgccc
tctctccgtcccgccactagt**ggcgcgcc**atatca***atg***attgaacaagatggattgcacgcaggttctccggccgcttgggtggagag
gctattcggctatgactgggcacaacagacaatcggctgctctgatgccgccgtgttccggctgtcagcgcaggggcgcccggttcttt
ttgtcaagaccgacctgtccggtgccctgaatgaactgcaggacgaggcagcgcggctatcgtggctggccacgacgggcgttcctt
gcgcagctgtgctcgacgttgtcactgaagcgggaagggactggctgctattgggcgaagtgccggggcaggatctcctgtcatctc
accttgctcctgccgagaaagtatccatcatggctgatgcaatgcggcggctgcatacgcttgatccggctacctgcccattcgaccac
caagcgaaacatcgcatcgagcgagcacgtactcggatggaagccggtcttgtcgatcaggatgatctggacgaagagcatcaggg
gctcgcgccagccgaactgttcgccaggctcaaggcgcgcatgcccgacggcgaggatctcgtcgtgacccatggcgatgcctgct
tgccgaatatcatggtggaaaatggccgcttttctggattcatcgactgtggccggctgggtgtggcggaccgctatcaggacatagcg
ttggctacccgtgatattgctgaagagcttggcggcgaatgggctgaccgcttcctcgtgctttacggtatcgccgctcccgattcgcag
cgcatcgccttctatcgccttcttgacgagttcttctaagatctgtcgatcgacaag**tgactcgag**gcagcagcagctcggatagtatcg
acacactctggacgctggtcgtgtgatggactgttgccgccacacttgctgccttgacctgtgaatatccctgccgcttttatcaaacagc
ctcagtgtgtttgatcttgtgtgtacgcgcttttgcgagttgctagctgcttgtgctatttgcgaataccacccccagcatcccccttccctcgt
ttcatatcgcttgcatcccaaccgcaacttatctacgctgtcctgctatccctcagcgctgctcctgctcctgctcactgcccctcgcacag
ccttggtttgggctccgcctgtattctcctggtactgcaacctgtaaaccagcactgcaatgctgatgcacgggaagtagtgggatggg
aacacaaatggaaagctt**gagctc**

SEQ ID NO: 41
Glutamate dehydrogenase::neo-opt::nit

gaattccgcctgcaacgcaagggcagccacagccgctcccacccgccgctgaaccgacacgtgcttgggcgcctgccgcctgcctgccgcat
gcttgtgctggtgaggctgggcagtgctgccatgctgattgaggcttggttcatcgggtggaagcttatgtgtgtgctgggcttgcatgccgggcaa
tgcgcatggtggcaagagggcggcagcacttgctggacgtgccgcggtgcctccaggtggttcaatcgcggcagccagagggatttcagatga
tcgcgcgtacaggttgagcagcagtgtcagcaaaggtagcagtttgccagaatgatcggttcagctgttaatcaatgccagcaagagaaggggtc
aagtgcaaacacgggcatgccacagcacgggcaccggggagtggaatggcaccaccaagtgtgtgcgagccagcatcgccgcctggctgttt
cagctacaacggcaggagtcatccaactaaccatagctgatcaacactgcaatcatcggcggctgatgcaagcatcctgcaagacacatgctgtg
cgatgctgcgctgctgcctgctgcgcacgccgttgagttggcagcagctcagccatgcactggatcaggctgggctgccactgcaatgtggtgg
ataggatgcaagtggagcgaataccaaaccctctggctgcttgctgggttgcatggcatcgcaccatcagcaggagcgcatgcgaagggactg
gccccatgcacgccatgccaaaccggagcgcaccgagtgtccacactgtcaccaggcccgcaagctttgcagaaccatgctcatggacgcatg
tagcgctgacgtcccttgacggcgctcctctcgggtgtgggaaacgcaatgcagcacaggcagcagaggcggcggcagcagagcggcggca
gcagcggcgggggccacccttcttgcggggtcgcgccccagccagcggtgatgcgctgatcnnnccaaacgagttcacattcatttgcagcct
ggagaagcgaggctggggccttgggctggtgcagcccgcaatggaatgcgggaccgccaggctagcagcaaaggcgcctccctactccg
catcgatgttccatagtgcattggactgcatttgggtggggcggccggctgtttctttcgtgttgcaaaacgcgccacgtcagcaacctgtcccgtg
ggtcccccgtgccgatgaaatcgtgtgcacgccgatcagctgattgcccggctcgcgaagtaggcgccctctttctgctcgccctctctccgtccc
gccactagt**ggcgcgcc**atatca***atg***atcgagcaggacggcctccacgccggctcccccgccgcctgggtggagcgcctgttcggctacgact
gggcccagcagaccatcggctgctccgacgccgccgtgttccgcctgtccgcccagggccgccccgtgctgttcgtgaagaccgacctgtccg
gcgccctgaacgagctgcaggacgaggccgcccgcctgtcctggctggccaccaccggcgtgccctgcgccgccgtgctggacgtggtgac
cgaggccggccgcgactggctgctgtgggcgaggtgcccggccaggacctgctgtcctcccacctggcccccgccgagaaggtgtccatca
tggccgacgccatgcgccgcctgcacaccctggacccccgccacctgcccttcgaccaccaggccaagcaccgcatcgagcgcgcccgcac
ccgcatggaggccggcctggtggaccaggacgacctggacgaggagcaccagggcctggcccccgccgagctgttcgcccgcctgaaggc
ccgcatgcccgacggcgaggacctggtggtgacccacggcgacgcctgcctgcccaacatcatggtggagaacggccgcttctccggcttcat
cgactgcggccgcctgggcgtggccgaccgctaccaggacatcgccctggccacccgcgacatcgccgaggagctgggcggcgagtgggc
cgaccgcttcctggtgctgtacggcatcgccgcccccgactcccagcgcatcgccttctaccgcctgctggacgagttcttc**tgactcgag**gcag
cagcagctcggatagtatcgacacactctggacgctggtcgtgtgatggactgttgccgccacacttgctgccttgacctgtgaatatccctgccgc
ttttatcaaacagcctcagtgtgtttgatcttgtgtgtacgcgcttttgcgagttgctagctgcttgtgctatttgcgaataccacccccagcatcccccttc
cctcgtttcatatcgcttgcatcccaaccgcaacttatctacgctgtcctgctatccctcagcgctgctcctgctcctgctcactgcccctcgcacagc

cttggtttgggctccgcctgtattctcctggtactgcaacctgtaaaccagcactgcaatgctgatgcacgggaagtagtgggatgggaacacaaat
ggaaagctt**gagctc**

SEQ ID NO: 42
Beta-tubulin::neo-opt::nit

gaattcctttcttgcgctatgacacttccagcaaaaggtagggcgggctgcgagacggcttcccggcgctgcatgcaacaccgatgat
gcttcgaccccccgaagctccttcggggctgcatgggcgctccgatgccgctccagggcgagcgctgtttaaatagccaggccccc
gattgcaaagacattatagcgagctaccaaagccatattcaaacacctagatcactaccacttctacacaggccactcgagcttgtgatc
gcactccgctaaggggggcgcctcttcctcttcgtttcagtcacaacccgcaaac**ggcgcgcc**atatca***atg***atcgagcaggacggcct
ccacgccggctccccccgccgcctgggtggagcgcctgttcggctacgactgggcccagcagaccatcggctgctccgacgccgcc
gtgttccgcctgtccgcccagggccgccccgtgctgttcgtgaagaccgacctgtccggcgccctgaacgagctgcaggacgaggc
cgcccgcctgtcctggctggccaccaccggcgtgccctgcgccgccgtgctggacgtggtgaccgaggccggccgcgactggctg
ctgctgggcgaggtgcccggccaggacctgctgtcctcccacctggcccccgccgagaaggtgtccatcatggccgacgccatgcg
ccgcctgcacaccctggaccccgccacctgcccccttcgaccaccaggccaagcaccgcatcgagcgcgcccgcaccccgcatgga
ggccggcctggtggaccaggacgacctggacgaggagcaccagggcctggcccccgccgagctgttcgcccgcctgaaggcccc
gcatgcccgacggcgaggacctggtggtgacccacggcgacgcctgcctgcccaacatcatggtggagaacggccgcttctccgg
cttcatcgactgcggccgcctgggcgtggccgaccgctaccaggacatcgccctggccacccgcgacatcgccgaggagctgggc
ggcgagtgggccgaccgcttcctggtgctgtacggcatcgccgcccccgactcccagcgcatcgccttctaccgcctgctggacga
gttcttctga**ctcgag**gcagcagcagctcggatagtatcgacacactctggacgctggtcgtgtgatggactgttgccgccacacttgc
tgccttgacctgtgaatatccctgccgcttttatcaaacagcctcagtgtgtttgatcttgtgtgtacgcgcttttgcgagttgctagctgctt
gtgctatttgcgaataccacccccagcatcccccttccctcgtttcatatcgcttgcatcccaaccgcaacttatctacgctgtcctgctatcc
ctcagcgctgctcctgctcctgctcactgcccctcgcacagccttggtttgggctccgcctgtattctcctggtactgcaacctgtaaacc
agcactgcaatgctgatgcacgggaagtagtgggatgggaacacaaatggaaagctt**gagctc**

SEQ ID NO: 43
GCCGCGACTGGCTGCTGCTGG

SEQ ID NO: 44
AGGTCCTCGCCGTCGGGCATGC

SEQ ID NO: 45
Acyl ACP desaturase

atcaaaggcatagattcacatttgttggcattgcagagcaatcatcgcgcaggacgaacatcgctcaccaagcacgtactgggcatcc
ggaggcctccgcaaattcctgcaacaggactcgctgatcagttcgcccaaggtctacgacgctccctatcggcgctagacttcaacac
atatttcactgtcacagcctcggcATGCATCAGGCCTCAGTCTCCACCATGAAGACCATCCAGTC
TCGGCACGCCGGTCCCATCGGACATGTGCAGTCGGGTCGCCGATCGGCGGGGCG
CGCGGGATCCCGCATGGCGACCCCCGTGGCCGCAGCTACCGTCGCAGCCCCTCGC
TCGGCCCTCAACCTCTCCCCCACCATCATTCGACAGGAGGTGCTCCACTCCGCCA
GCGCCCAGCAACTAGACTGCGTGGCCTCCCTGGCGCCCGTCTTCGAGTCCCAGAT
CCTCCCCCTCCTGACGCCCGTGGACGAGATGTGGCAGCCCACCGACTTCCTCCCC
GCCTCGAACTCGGAGGCATTCTTCGACCAGATCGGCGACCTGCGGGCGCGATCG
GCGGCCATCCCCGACGACCTGCTGGTCTGCCTGGTGGGGGACATGATCACGGAG
GAGGCCCTGCCCACCTACATGGCCATGCTGAACACCCTGGACGTCGTGCGCGATG
AGACAGGGCACAGCCAGCACCCCTACGCCAAGTGGACCAGGGCTTGGATCGCGG
AGGAGAACCGCCATGGCGACCTGCTGAACAAGTACATGTGGCTGACGGGGCGGG
TGGGACATGCTGGCGGTGGAGCGCACCATCCAGCCATGCTGGCGGTGGAGCGCA
CCATCCAGCGCCTCATCTCATCGGGCATGGACCCGGGCACGGAGAACCACCCCT
ACCACGCCTTTGTGTTCACCAGCTTCCAGGAGCGCGCCACCAAGCTGAGCCACGG

CTCCACCGCCCGCCTGGCGGTCGCCGCCGGGGACGAGGCCCTGGCCAAGATCTG
CGGGACCATTGCGCGGGACGAGTCGCGCCACGAGGCGGCGTACACGCGGACCAT
GGATGCCATCTTCCAGCGCGACCCCAGCGGGGCCATGGTGGCGTTTGCGCACATG
ATGATGCGCAAGATCACCATGCCCGCCCACCTCATGGACGACGGCCAGCACGGC
GCGCGCAACGGGGGGGCGCAACTTGTTCGACGACTTTGCGGCAGTGGCGGAGCG
GGCAGGGGTGTACACCGCCGGCGACTACATCGGCATCCTGCGCCACCTCATCCG
GCGCTGGGACGTGGAGGG

SEQ ID NO: 46
Acyl ACP desaturase
MHQASVSTMKTIQSRHAGPIGHVQSGRRSAGRAGSRMATPVAAATVAAPRSALNLS
PTIIRQEVLHSASAQQLDCVASLAPVFESQILPLLTPVDEMWQPTDFLPASNSEAFFDQ
IGDLRARSAAIPDDLLVCLVGDMITEEALPTYMAMLNTLDVVRDETGHSQHPYAKW
TRAWIAEENRHGDLLNKYMWLTGRVGHAGGGAHHPAMLAVERTIQRLISSGMDPG
TENHPYHAFVFTSFQERATKLSHGSTARLAVAAGDEALAKICGTIARDESRHEAAYT
RTMDAIFQRDPSGAMVAFAHMMMMRKITMPAHLMDDGQHGARNGGAQLVRRLCGS
GGAGRGVHRRRLHRHPAPPHPALGRGG

SEQ ID NO: 47
Geranyl geranyl diphosphate synthase

attatacatcggcatcgtctcaggtttcacgatctgcatgctatctatgggactgtgactccgccggccaggttgtggtgcgcgagaatc
ctccccgctcctgccttctcatttccctgacgggagtcgccgctgagcaccgggcggatcATGGGCGTCGGCACACT
CCAAACCCCATATACATGTGGTCGTGCATTCACGCATAGCGCACGGTATGTCCCG
CGACGCGCGGCTCGAAGCCGTGGCCATCCGACGCGCTGCACGGCCGAGGTGAGG
GCACGCCCCTCCGCCAATGGCGCGCAGCCCATGACCGCCTTCGACTTCCGGCAGT
ACATGCAGCAGCGCGCCGCGCTGGTGGACGCAGCGCTGGACCTGGCAGTGCCGC
TGCAGTACCCCGAGAAGATCAACGAGGCCATGCGGTACAGCCTGCTGGCCGGGG
GCAAGCGCGTGCGCCCCGCGCTCTGCCTCGCTGCCTGCGAGCTCGTGGGCGGCCC
CCTGGAGGCGGCCATGCCCGCCGCCTGCGCCATGGAGATGATCCACACCATGAG
CCTCATCCACGACGACCTCCCCGCCATGGACAACGACGACTTCCGGCGCGGCCA
GCCCGCCAACCACAAGGCCTATGGCGAGGAGATTGCGATCCTGGCGGGCGACGC
GCTGCTGTCGCTGAGCTTTGAGCACATCGCGCGCGAGACGCGAGGCGTGGACCC
GGTGCGCGTCCTGGCCGCCATCTCGGAGTGGCGCGCGGTGGGCAGCCGCGGGCT
GGTGGCGGGGCAGGTGGTGGACCTGGGTTTCGAGGGCGGCGGCGTGGGGCTGGC
CCCGCTGCGCTACATCCACGAGCACAAAACCGCGGCGCTGCTGGAGGCGGCGGT
GGTGTCCGGCGCGCTGCTGGGCGGCGCGGAGGAGGCGGACCTGGAGCGCCTGCG
CACCTACAACCGCGCCATCGGCCTCGCTTTCCAGGTGGTGGGGGACATCCTGGAC
ATCCCGGGGACCAGCGAGGAGCTGGGCAAGACCGCGGGCAAGGACCTGAGCTCC
CCCAAAACCCCCTACCCGTCCCTGGTGGGGCTGGCCAGGTCCAAAAAAATTGCG
GACGAACTGATTGAGGACGCGAAAACCCAACTCACCCAGTACGAGCCGGCCCGA
GCGGCGCCCCTCGTAACCCTGGCCGAAAACATTTGAaaccggaagaactgactgggggccccccc
ctgcccccagatacggcggggctcctccatccagttttgggatgggaggagcgacaaccgaccccgtaaccctgtgacgcgtttgcc
ttgcatacgtacgcatgccttgaaacccatccatgaccctcaacaatacctggttgtgtgtagcttggtcctgaaaaaaaaaaaaaaaaa
aaaaaaaaaaa

SEQ ID NO: 48
Geranyl geranyl diphosphate synthase

MGVGTLQTPYTCGRAFTHSARYVPRRAARSRGHPTRCTAEVRARPSANGAQPMTAF
DFRQYMQQRAALVDAALDLAVPLQYPEKINEAMRYSLLAGGKRVRPALCLAACEL
VGGPLEAAMPAACAMEMIHTMSLIHDDLPAMDNDDFRRGQPANHKAYGEEIAILAG
DALLSLSFEHIARETRGVDPVRVLAAISEWRAVGSRGLVAGQVVDLGFEGGGVGLA
PLRYIHEHKTAALLEAAVVSGALLGGAEEADLERLRTYNRAIGLAFQVVGDILDIPGT
SEELGKTAGKDLSSPKTPYPSLVGLARSKKIADELIEDAKTQLTQYEPARAAPLVTLA
ENI

SEQ ID NO: 49
Geranyl geranyl diphosphate synthase
cagatgccATGCGCCCTCGGGCCGCGGGCCTGAGGGTCCACGCAGCGTCCTCGGTGG
CCCAGACGCACCAGGCCGCCCCCCCGGCGGACAGGAGGTTCGACGACTACCAGC
CCCGCACCGCCATCCTCTTCCCCGGCCAGGGCGCGCAGAGCGTGGGCATGGCGG
GAGAGCTGGCGAAGGCCGTCCCCGCCGCCGCGGCGCTGTTCGACGCCGCCTCCG
ACCAGCTCGGCTATGACCTGCTCCGCGTGTGCGTTGAGGGCCCCAAGGCGCGCCT
GGACAGCACCGCCGTCAGCCAGCCCGCCATCTACGTGGCCAGCCTGGCGGCGGT
GGAGAAGCTGCGCGCGGAGGGCGGGGAGGAGGCACTGGCCGCCATCGACGTCG
CTGCCGGTCTGTCCTTGGGCGAGTACACCGCGCTGGCCTTTGCCGGCGCCTTCTC
CTTCGCCGACGGGCTGCGCCTGGTGGCCCTGCGCGGCGCCAGCATGCAGGCCGC
CGCCGACGCCGCACCCTCGGGCATGGTCTCCGTCATCGGTCTGCCCTCCGACGCG
GTGGCCGCGCTGTGCGAGGCCGCCAACGCGCAGGTGGCCCCCGACCAGGCCGTG
CGCATCGCCAACTACCTCTGCGACGGCAACTACGCCGTCAGCGGTGGGCTGGAG
GGCTGCGCGGCGGTGGAGGGCCTGGCCAAGGCCCACAAGGCGCGCATGACGGTG
CGCCTGGCGGTGGCGGGCGCCTTCCACACCCCCTTCATGCAGCCGGCGGTGGAG
GCGCTGAGCGCGGGCGCTGGCGGACACGCCGCTGGTCGCGCCGCGCATCCCCGT
GGTCAGCAACGGGACGCC

SEQ ID NO: 50
Geranyl geranyl diphosphate synthase

MRPRAAGLRVHAASSVAQTHQAAPPADRRFDDYQPRTAILFPGQGAQSVGMAGEL
AKAVPAAAALFDAASDQLGYDLLRVCVEGPKARLDSTAVSQPAIYVASLAAVEKLR
AEGGEEALAAIDVAAGLSLGEYTALAFAGAFSFADGLRLVALRGASMQAAADAAPS
GMVSVIGLPSDAVAALCEAANAQVAPDQAVRIANYLCDGNYAVSGGLEGCAAVEG
LAKAHKARMTVRLAVAGAFHTPFMQPAVEALSAGAGGHAAGRAAHPRGQQRDA

SEQ ID NO: 51
Gylceraldehyde 3-phosphate dehydrogenase cDNA sequence

TGTCCATCTCCCCCCACCCTCCATCCAACCATCGTCGACGGCATGCAGGCGCTGT
GTTCTCACCCCGCGTCCCTCACGGCGCGTGCGGTACCCCATGGGCGGGCCAGCCC
AGCACAGCGGGTGTCCAGCGCCGGCCCGGCCTACACCGGCCTGTCCCGGCACAC
CCTGGGCTGCCCCAGCACCCCCACCCTCCAGTCCCGCGCCGCGGTCCAGACCCGC
GGCTCCTCCTCCGGCTCCACCACGCGCATGACCACCACCGCCCAGCGCAAGATCA
AGGTGGCCATCAACGGGTTCGGCCGCATCGGCCGCCAGTTCCTGCGCTGCGTGGA
GGGGCGCGAGGACTCGCTGCTGGAGATCGTGGCCGTGAACGACTCCGGCGGCGT
GAAGCAGGCCAGCCACCTGCTCAAGTACGACTCCACCATGGGCACCTTCAACGC
CGACATCAAGATCTCGGGCGAGGGCACCTTCTCCGTCAACGGCCGCGACATCCG
CGTCGTCTCCTCCCGCGACCCCCTGGCCCTGCCCTGGGGCGAGCTGGGCGTGGAC
CTGGTGATCGAGGGGACGGGAGTGTTTGTGGACCGCAAGGGTGCCAGCAAGCAC

CTGCAGGCGGGGGCCAAGAAGGTCATCATCACCGCGCCGGCCAAGGGCTCCGAC
GTGCCCACCTACGTCATGGGCGTGAACGCGGACCAGTACTCCAACTCCGACGAC
ATCATCTCCAACGCCTCCTGCACCACCAACTGCCTGGCGCCCTTTGTCAAGGTGC
TCAACGACCGCTTCGGCATCGTGA

SEQ ID NO: 52
Glyceraldehyde 3-phosphate dehydrogenase

MQALCSHPASLTARAVPHGRASPAQRVSSAGPAYTGLSRHTLGCPSTPTLQSRAAVQ
TRGSSSGSTTRMTTTAQRKIKVAINGFGRIGRQFLRCVEGREDSLLEIVAVNDSGGVK
QASHLLKYDSTMGTFNADIKISGEGTFSVNGRDIRVVSSRDPLALPWGELGVDLVIEG
TGVFVDRKGASKHLQAGAKKVIITAPAKGSDVPTYVMGVNADQYSNSDDIISNASCT
TNCLAPFVKVLNDRFGIV

SEQ ID NO: 53
Oxygen Evolving Complex protein (OEE33) cDNA sequence

gatgttgagaatagtagcttgctgccttgtcgccatgcagagcgtgtgcgcgcagtcggtttcatgcaaggggccttcacccagtccc
tgcggacccccccgatgcagcaggagccagctcgtctgccgggctgatggcaaggccggagccttcatcaagaccgtaaagagcgg
tgctgccgctctggctgcctccctcctcctgtctgggggtgcgggcgcactgacctttgatgagctgcagggcctgacctacctgcag
gtgaagggctctggcatcgccaacacctgccccaccctgtctggcggctcctccaacatcaaggacctgaagagcgggacctactcc
gtcaacaagatgtgcctggagcccacgtccttcaaggtcaaggaggaggcacagttcaagaacggcgaggccgactttgtgcccac
caagctcgtcacgcgtctgacctacaccctggacgagatctctggccagatgaagatcgacggcagcggcggcgtggagttcaagg
aggaggatggcatcgactatgctgcagtcaccgtgcagcttccgggcggggagcgcgtgccctcctcttcaccatcaaggagcttg
acgccaaggggactgccgacggcttcaagggcgagttcaccgtgccctcctaccgtgggtcctccttcctggaccccaagggccgc
ggcgcctccaccggctacgacaacgccgtggccctgcccgccgccggcgattccgaggagttggagaaggagaacaacaagtcc
accaaggctctgaaggggaggccatcttctccatcgccaaggtggacgccgggacaggggaggtggcgggcatctttgagt

SEQ ID NO: 54
Oxygen Evolving Complex protein (OEE33)

MQSVCAQSVSCKGAFTQSLRTPRCSRSQLVCRADGKAGAFIKTVKSGAAALAASLL
LSGGAGALTFDELQGLTYLQVKGSGIANTCPTLSGGSSNIKDLKSGTYSVNKMCLEP
TSFKVKEEAQFKNGEADFVPTKLVTRLTYTLDEISGQMKIDGSGGVEFKEEDGIDYA
AVTVQLPGGERVPFLFTIKELDAKGTADGFKGEFTVPSYRGSSFLDPKGRGASTGYD
NAVALPAAGDSEELEKENNKSTKALKGEAIFSIAKVDAGTGEVAGIFE

SEQ ID NO: 55
Clp protease cDNA sequence
ataatcggaacccagctgcacgcaccatcagtgcggcagcatgcagaccgtcgcagccagctatggcgtattggcgccctccggct
ccagcgtgacccggggctcgaccagcagcaagcagcacttcaccaccctcactccctttccggcttcaggcgcctgaatcatgtgga
tcgggcggggcaggcggggtctgggagcccccagaccctgcagcaggccgtgggcaaggccgtgcgccggtcgcggggccgc
accaccagcgccgtgcgcgtgacccgcatgatgtttgagcggttcaccgagaaggccatcaaggtggtcatgctcgcgcaggagga
ggctcgccgtctgggccacaacttcgtggggacggagcaaatcctgctggggttgattggggagtccacaggcatcgccgccaagg
tcctcaagtcgatgggcgtcacgctgaaagatgcgcgtgtggaggtcgagaagatcatcggccgggggagcggctttgtggccgtg
gagatccccttcaccccccgcgccaagcgtgtgctggagctgtccctggaggaggctcgccagctcggccacaactacattggcac
ggagcacatcctgctgggcctgctgcgcgagggtgagggcgtggcctcccgcgtgctggagaccctgggcgccgaccccagaa
gatccgcactcaggtggtacgcatggtgggtgagtcgcaggagcccgtgggcaccacggtgggcggagggtccaccggctccaa
caagatgcccaccctggaggagtacggcaccaacctgaccgcccaggccg

SEQ ID NO: 56
Clp protease

MQTVAASYGVLAPSGSSVTRGSTSSKQHFTTLTPFSGFRRLNHVDRAGQAGSGSPQT
LQQAVGKAVRRSRGRTTSAVRVTRMMFERFTEKAIKVVMLAQEEARRLGHNFVGT
EQILLGLIGESTGIAAKVLKSMGVTLKDARVEVEKIIGRGSGFVAVEIPFTPRAKRVLE
LSLEEARQLGHNYIGTEHILLGLLREGEGVASRVLETLGADPQKIRTQVVRMVGESQ
EPVGTTVGGGSTGSNKMPTLEEYGTNLTAQA

SEQ ID NO: 57
Crβ-tub::NCO-suc2::CvNitRed
CTTTCTTGCGCTATGACACTTCCAGCAAAAGGTAGGGCGGGCTGCGAGACGGCTT
CCCGGCGCTGCATGCAACACCGATGATGCTTCGACCCCCCGAAGCTCCTTCGGGG
CTGCATGGGCGCTCCGATGCCGCTCCAGGGCGAGCGCTGTTTAAATAGCCAGGCC
CCCGATTGCAAAGACATTATAGCGAGCTACCAAAGCCATATTCAAACACCTAGA
TCACTACCACTTCTACACAGGCCACTCGAGCTTGTGATCGCACTCCGCTAAGGGG
GCGCCTCTTCCTCTTCGTTTCAGTCACAACCCGCAAACGGCGCGCCATATCAATG
CTTCTTCAGGCCTTTCTTTTTCTTCTTGCTGGTTTTGCTGCCAAGATCAGCGCCTCT
ATGACGAACGAAACCTCGGATAGACCACTTGTGCACTTTACACCAAACAAGGGC
TGGATGAATGACCCCAATGGACTGTGGTACGACGAAAAAGATGCCAAGTGGCAT
CTGTACTTTCAATACAACCCGAACGATACTGTCTGGGGGACGCCATTGTTTTGGG
GCCACGCCACGTCCGACGACCTGACCAATTGGGAGGACCAACCAATAGCTATCG
CTCCGAAGAGGAACGACTCCGGAGCATTCTCGGGTTCCATGGTGGTTGACTACAA
CAATACTTCCGGCTTTTTCAACGATACCATTGACCCGAGACAACGCTGCGTGGCC
ATATGGACTTACAACACACCGGAGTCCGAGGAGCAGTACATCTCGTATAGCCTG
GACGGTGGATACACTTTTACAGAGTATCAGAAGAACCCTGTGCTTGCTGCAAATT
CGACTCAGTTCCGAGATCCGAAGGTCTTTTGGTACGAGCCCTCGCAGAAGTGGAT
CATGACAGCGGCAAAGTCACAGGACTACAAGATCGAAATTTACTCGTCTGACGA
CCTTAAATCCTGGAAGCTCGAATCCGCGTTCGCAAACGAGGGCTTTCTCGGCTAC
CAATACGAATGCCCAGGCCTGATAGAGGTCCCAACAGAGCAAGATCCCAGCAAG
TCCTACTGGGTGATGTTTATTTCCATTAATCCAGGAGCACCGGCAGGAGGTTCTT
TTAATCAGTACTTCGTCGGAAGCTTTAACGGAACTCATTTCGAGGCATTTGATAA
CCAATCAAGAGTAGTTGATTTTGGAAAGGACTACTATGCCCTGCAGACTTTCTTC
AATACTGACCCGACCTATGGGAGCGCTCTTGGCATTGCGTGGGCTTCTAACTGGG
AGTATTCCGCATTCGTTCCTACAAACCCTTGGAGGTCCTCCATGTCGCTCGTGAG
GAAATTCTCTCTCAACACTGAGTACCAGGCCAACCCGGAAACCGAACTCATAAA
CCTGAAAGCCGAACCGATCCTGAACATTAGCAACGCTGGCCCCTGGAGCCGGTTT
GCAACCAACACCACGTTGACGAAAGCCAACAGCTACAACGTCGATCTTTCGAAT
AGCACCGGTACACTTGAATTTGAACTGGTGTATGCCGTCAATACCACCCAAACGA
TCTCGAAGTCGGTGTTCGCGGACCTCTCCCTCTGGTTTAAAGGCCTGGAAGACCC
CGAGGAGTACCTCAGAATGGGTTTCGAGGTTTCTGCGTCCTCCTTCTTCCTTGATC
GCGGGAACAGCAAAGTAAAATTTGTTAAGGAGAACCCATATTTTACCAACAGGA
TGAGCGTTAACAACCAACCATTCAAGAGCGAAAACGACCTGTCGTACTACAAAG
TGTATGGTTTGCTTGATCAAAATATCCTGGAACTCTACTTCAACGATGGTGATGT
CGTGTCCACCAACACATACTTCATGACAACCGGGAACGCACTGGGCTCCGTGAA
CATGACGACGGGTGTGGATAACCTGTTCTACATCGACAAATTCCAGGTGAGGGA
AGTCAAGTGAGATCTGTCGATCGACAAGCTCGAGGCAGCAGCAGCTCGGATAGT
ATCGACACACTCTGGACGCTGGTCGTGTGATGGACTGTTGCCGCCACACTTGCTG
CCTTGACCTGTGAATATCCCTGCCGCTTTTATCAAACAGCCTCAGTGTGTTTGATC
TTGTGTGTACGCGCTTTTGCGAGTTGCTAGCTGCTTGTGCTATTTGCGAATACCAC

CCCCAGCATCCCCTTCCCTCGTTTCATATCGCTTGCATCCCAACCGCAACTTATCT
ACGCTGTCCTGCTATCCCTCAGCGCTGCTCCTGCTCCTGCTCACTGCCCCTCGCAC
AGCCTTGGTTTGGGCTCCGCCTGTATTCTCCTGGTACTGCAACCTGTAAACCAGC
ACTGCAATGCTGATGCACGGGAAGTAGTGGGATGGGAACACAAATGGAAAGCTT

SEQ ID NO: 58 Crβ-tub::CO-suc2::CvNitRed

CTTTCTTGCGCTATGACACTTCCAGCAAAAGGTAGGGCGGGCTGCGAGACGGCTT
CCCGGCGCTGCATGCAACACCGATGATGCTTCGACCCCCCGAAGCTCCTTCGGGG
CTGCATGGGCGCTCCGATGCCGCTCCAGGGCGAGCGCTGTTTAAATAGCCAGGCC
CCCGATTGCAAAGACATTATAGCGAGCTACCAAAGCCATATTCAAACACCTAGA
TCACTACCACTTCTACACAGGCCACTCGAGCTTGTGATCGCACTCCGCTAAGGGG
GCGCCTCTTCCTCTTCGTTTCAGTCACAACCCGCAAACGGCGCGCCATGCTGCTG
CAGGCCTTCCTGTTCCTGCTGGCCGGCTTCGCCGCCAAGATCAGCGCCTCCATGA
CGAACGAGACGTCCGACCGCCCCCTGGTGCACTTCACCCCCAACAAGGGCTGGA
TGAACGACCCCAACGGCCTGTGGTACGACGAGAAGGACGCCAAGTGGCACCTGT
ACTTCCAGTACAACCCGAACGACACCGTCTGGGGGACGCCCTTGTTCTGGGGCCA
CGCCACGTCCGACGACCTGACCAACTGGGAGGACCAGCCCATCGCCATCGCCCC
GAAGCGCAACGACTCCGGCGCCTTCTCCGGCTCCATGGTGGTGGACTACAACAA
CACCTCCGGCTTCTTCAACGACACCATCGACCCGCGCCAGCGCTGCGTGGCCATC
TGGACCTACAACACCCCGGAGTCCGAGGAGCAGTACATCTCCTACAGCCTGGAC
GGCGGCTACACCTTCACCGAGTACCAGAAGAACCCCGTGCTGGCCGCCAACTCC
ACCCAGTTCCGCGACCCGAAGGTCTTCTGGTACGAGCCCTCCCAGAAGTGGATCA
TGACCGCGGCCAAGTCCCAGGACTACAAGATCGAGATCTACTCCTCCGACGACCT
GAAGTCCTGGAAGCTGGAGTCCGCGTTCGCCAACGAGGGCTTCCTCGGCTACCA
GTACGAGTGCCCCGGCCTGATCGAGGTCCCCACCGAGCAGGACCCCAGCAAGTC
CTACTGGGTGATGTTCATCTCCATCAACCCCGGCGCCCCGGCCGGCGGCTCCTTC
AACCAGTACTTCGTCGGCAGCTTCAACGGCACCCACTTCGAGGCCTTCGACAACC
AGTCCCGCGTGGTGGACTTCGGCAAGGACTACTACGCCCTGCAGACCTTCTTCAA
CACCGACCCGACCTACGGGAGCGCCCTGGGCATCGCGTGGGCCTCCAACTGGGA
GTACTCCGCCTTCGTGCCCACCAACCCCTGGCGCTCCTCCATGTCCCTCGTGCGCA
AGTTCTCCCTCAACACCGAGTACCAGGCCAACCCGGAGACGGAGCTGATCAACC
TGAAGGCCGAGCCGATCCTGAACATCAGCAACGCCGGCCCCTGGAGCCGGTTCG
CCACCAACACCACGTTGACGAAGGCCAACAGCTACAACGTCGACCTGTCCAACA
GCACCGGCACCCTGGAGTTCGAGCTGGTGTACGCCGTCAACACCACCCAGACGA
TCTCCAAGTCCGTGTTCGCGGACCTCTCCCTCTGGTTCAAGGGCCTGGAGGACCC
CGAGGAGTACCTCCGCATGGGCTTCGAGGTGTCCGCGTCCTCCTTCTTCCTGGAC
CGCGGGAACAGCAAGGTGAAGTTCGTGAAGGAGAACCCCTACTTCACCAACCGC
ATGAGCGTGAACAACCAGCCCTTCAAGAGCGAGAACGACCTGTCCTACTACAAG
GTGTACGGCTTGCTGGACCAGAACATCCTGGAGCTGTACTTCAACGACGGCGAC
GTCGTGTCCACCAACACCTACTTCATGACCACCGGGAACGCCCTGGGCTCCGTGA
ACATGACGACGGGGGTGGACAACCTGTTCTACATCGACAAGTTCCAGGTGCGCG
AGGTCAAGTGATTAATTAACTCGAGGCAGCAGCAGCTCGGATAGTATCGACACA
CTCTGGACGCTGGTCGTGTGATGGACTGTTGCCGCCACACTTGCTGCCTTGACCT
GTGAATATCCCTGCCGCTTTTATCAAACAGCCTCAGTGTGTTTGATCTTGTGTGTA
CGCGCTTTTGCGAGTTGCTAGCTGCTTGTGCTATTTGCGAATACCACCCCCAGCAT
CCCCTTCCCTCGTTTCATATCGCTTGCATCCCAACCGCAACTTATCTACGCTGTCC
TGCTATCCCTCAGCGCTGCTCCTGCTCCTGCTCACTGCCCCTCGCACAGCCTTGGT
TTGGGCTCCGCCTGTATTCTCCTGGTACTGCAACCTGTAAACCAGCACTGCAATG
CTGATGCACGGGAAGTAGTGGGATGGGAACACAAATGGA

SEQ ID NO: 59
Cinnamomum camphorum FatB1 (Q39473)

MATTSLASAFCSMKAVMLARDGRGMKPRSSDLQLRAGNAQTSLKMINGTKFSYTES
LKKLPDWSMLFAVITTIFSAAEKQWTNLEWKPKPNPPQLLDDHFGPHGLVFRRTFAI
RSYEVGPDRSTSIVAVMNHLQEAALNHAKSVGILGDGFGTTLEMSKRDLIWVVKRT
HVAVERYPAWGDTVEVECWVGASGNNGRRHDFLVRDCKTGEILTRCTSLSVMMNT
RTRRLSKIPEEVRGEIGPAFIDNVAVKDEEIKKPQKLNDSTADYIQGGLTPRWNDLDI
NQHVNNIKYVDWILETVPDSIFESHHISSFTIEYRRECTMDSVLQSLTTVSGGSSEAGL
VCEHLLQLEGGSEVLRAKTEWRPKLTDSFRGISVIPAESSV

SEQ ID NO: 60
Cinnamomum camphorum FatB1 codon optimized for Prototheca

ggcgcgccatggccaccacctccctggcctccgccttctgcagcatgaaggccgtgatgctggcccgcgacggcc
gcggcatgaagccccgctccagcgacctgcagctgcgcgccggcaacgcccagacctccctgaagatgatcaacg
gcaccaagttctcctacaccgagagcctgaagaagctgcccgactggtccatgctgttcgccgtgatcaccacca
tcttctccgccgccgagaagcagtggaccaacctggagtggaagcccaagcccaaccccccccagctgctggacg
accacttcggccccccacggcctggtgttccgccgcaccttcgccatccgcagctacgaggtgggccccgaccgct
ccaccagcatcgtggccgtgatgaaccacctgcaggaggccgccctgaaccacgccaagtccgtgggcatcctgg
gcgacggcttcggcaccaccctggagatgtccaagcgcgacctgatctgggtggtgaagcgcacccacgtggccg
tggagcgctaccccgcctggggcgacaccgtggaggtggagtgctgggtgggcgcctccggcaacaacggccgcc
gccacgacttcctggtgcgcgactgcaagaccggcgagatcctgacccgctgcacctccctgagcgtgatgatga
acacccgcacccgccgcctgagcaagatccccgaggaggtgcgcggcgagatcggccccgccttcatcgacaacg
tggccgtgaaggacgaggagatcaagaagccccagaagctgaacgactccaccgccgactacatccagggcggcc
tgaccccccgctggaacgacctggacatcaaccagcacgtgaacaacatcaagtacgtggactggatcctggaga
ccgtgcccgacagcatcttcgagagccaccacatctcctccttcaccatcgagtaccgccgcgagtgcaccatgg
acagcgtgctgcagtccctgaccaccgtgagcggcggctcctccgaggccggcctggtgtgcgagcacctgctgc
agctggagggcggcagcgaggtgctgcgcgccaagaccgagtggcgccccaagctgaccgactccttccgcggca
tcagcgtgatccccgccgagtccagcgtgatggactacaaggaccacgacggcgactacaaggaccacgacatcg
actacaaggacgacgacgacaagtgactcgagttaattaa

SEQ ID NO: 61
Cuphea hookeriana FatB2 (GenBank AAC49269)

MVAAAASSAFFPVPAPGASPKPGKFGNWPSSLSPSFKPKSIPNGGFQVKANDSAHPK
ANGSAVSLKSGSLNTQEDTSSSPPPRTFLHQLPDWSRLLTAITTVFVKSKRPDMHDRK
SKRPDMLVDSFGLESTVQDGLVFRQSFSIRSYEIGTDRTASIETLMNHLQETSLNHCK
STGILLDGFGRTLEMCKRDLIWVVIKMQIKVNRYPAWGDTVEINTRFSRLGKIGMGR
DWLISDCNTGEILVRATSAYAMMNQKTRRLSKLPYEVHQEIVPLFVDSPVIEDSDLK
VHKFKVKTGDSIQKGLTPGWNDLDVNQHVSNVKYIGWILESMPTEVLETQELCSLA
LEYRRECGRDSVLESVTAMDPSKVGVRSQYQHLLRLEDGTAIVNGATEWRPKNAGA
NGAISTGKTSNGNSVS

SEQ ID NO: 62
Cuphea hookeriana FatB2 codon optimized for Prototheca

GGCGCGCCATGGTGGCCGCCGCCGCCTCCAGCGCCTTCTTCCCCGTGCCCGCCCC
CGGCGCCTCCCCCAAGCCCGGCAAGTTCGGCAACTGGCCCTCCAGCCTGAGCCCC
TCCTTCAAGCCCAAGTCCATCCCCAACGGCGGCTTCCAGGTGAAGGCCAACGAC
AGCGCCCACCCCAAGGCCAACGGCTCCGCCGTGAGCCTGAAGAGCGGCAGCCTG
AACACCCAGGAGGACACCTCCTCCAGCCCCCCCCCCCGCACCTTCCTGCACCAGC
TGCCCGACTGGAGCCGCCTGCTGACCGCCATCACCACCGTGTTCGTGAAGTCCAA
GCGCCCCGACATGCACGACCGCAAGTCCAAGCGCCCCGACATGCTGGTGGACAG

CTTCGGCCTGGAGTCCACCGTGCAGGACGGCCTGGTGTTCCGCCAGTCCTTCTCC
ATCCGCTCCTACGAGATCGGCACCGACCGCACCGCCAGCATCGAGACCCTGATG
AACCACCTGCAGGAGACCTCCCTGAACCACTGCAAGAGCACCGGCATCCTGCTG
GACGGCTTCGGCCGCACCCTGGAGATGTGCAAGCGCGACCTGATCTGGGTGGTG
ATCAAGATGCAGATCAAGGTGAACCGCTACCCCGCCTGGGGCGACACCGTGGAG
ATCAACACCCGCTTCAGCCGCCTGGGCAAGATCGGCATGGGCCGCGACTGGCTG
ATCTCCGACTGCAACACCGGCGAGATCCTGGTGCGCGCCACCAGCGCCTACGCC
ATGATGAACCAGAAGACCCGCCGCCTGTCCAAGCTGCCCTACGAGGTGCACCAG
GAGATCGTGCCCCTGTTCGTGGACAGCCCCGTGATCGAGGACTCCGACCTGAAG
GTGCACAAGTTCAAGGTGAAGACCGGCGACAGCATCCAGAAGGGCCTGACCCCC
GGCTGGAACGACCTGGACGTGAACCAGCACGTGTCCAACGTGAAGTACATCGGC
TGGATCCTGGAGAGCATGCCCACCGAGGTGCTGGAGACCCAGGAGCTGTGCTCC
CTGGCCCTGGAGTACCGCCGCGAGTGCGGCCGCGACTCCGTGCTGGAGAGCGTG
ACCGCCATGGACCCCAGCAAGGTGGGCGTGCGCTCCCAGTACCAGCACCTGCTG
CGCCTGGAGGACGGCACCGCCATCGTGAACGGCGCCACCGAGTGGCGCCCCAAG
AACGCCGGCGCCAACGGCGCCATCTCCACCGGCAAGACCAGCAACGGCAACTCC
GTGTCCATGGACTACAAGGACCACGACGGCGACTACAAGGACCACGACATCGAC
TACAAGGACGACGACGACAAGTGACTCGAGTTAATTAA

SEQ ID NO: 63
Umbellularia FatB1 (GenBank Q41635)

MATTSLASAFCSMKAVMLARDGRGMKPRSSDLQLRAGNAPTSLKMINGTKFSYTES
LKRLPDWSMLFAVITTIFSAAEKQWTNLEWKPKPKLPQLLDDHFGLHGLVFRRTFAI
RSYEVGPDRSTSILAVMNHMQEATLNHAKSVGILGDGFGTTLEMSKRDLMWVVRR
THVAVERYPTWGDTVEVECWIGASGNNGMRRDFLVRDCKTGEILTRCTSLSVLMNT
RTRRLSTIPDEVRGEIGPAFIDNVAVKDDEIKKLQKLNDSTADYIQGGLTPRWNDLDV
NQHVNNLKYVAWVFETVPDSIFESHHISSFTLEYRRECTRDSVLRSLTTVSGGSSEAG
LVCDHLLQLEGGSEVLRARTEWRPKLTDSFRGISVIPAEPRV

SEQ ID NO: 64
Umbellularia FatB1 codon optimized for Prototheca

GGCGCGCCATGGCCACCACCAGCCTGGCCTCCGCCTTCTGCTCCATGAAGGCCGT
GATGCTGGCCCGCGACGGCCGCGGCATGAAGCCCCGCAGCTCCGACCTGCAGCT
GCGCGCCGGCAACGCCCCCACCTCCCTGAAGATGATCAACGGCACCAAGTTCAG
CTACACCGAGAGCCTGAAGCGCCTGCCCGACTGGTCCATGCTGTTCGCCGTGATC
ACCACCATCTTCAGCGCCGCCGAGAAGCAGTGGACCAACCTGGAGTGGAAGCCC
AAGCCCAAGCTGCCCCAGCTGCTGGACGACCACTTCGGCCTGCACGGCCTGGTGT
TCCGCCGCACCTTCGCCATCCGCTCCTACGAGGTGGGCCCCGACCGCAGCACCTC
CATCCTGGCCGTGATGAACCACATGCAGGAGGCCACCCTGAACCACGCCAAGAG
CGTGGGCATCCTGGGCGACGGCTTCGGCACCACCCTGGAGATGTCCAAGCGCGA
CCTGATGTGGGTGGTGCGCCGCACCCACGTGGCCGTGGAGCGCTACCCCACCTGG
GGCGACACCGTGGAGGTGGAGTGCTGGATCGGCGCCAGCGGCAACAACGGCATG
CGCCGCGACTTCCTGGTGCGCGACTGCAAGACCGGCGAGATCCTGACCCGCTGC
ACCTCCCTGAGCGTGCTGATGAACACCCGCACCCGCCGCCTGAGCACCATCCCCG
ACGAGGTGCGCGGCGAGATCGGCCCCGCCTTCATCGACAACGTGGCCGTGAAGG
ACGACGAGATCAAGAAGCTGCAGAAGCTGAACGACTCCACCGCCGACTACATCC
AGGGCGGCCTGACCCCCCGCTGGAACGACCTGGACGTGAACCAGCACGTGAACA
ACCTGAAGTACGTGGCCTGGGTGTTCGAGACCGTGCCCGACAGCATCTTCGAGTC

CCACCACATCAGCTCCTTCACCCTGGAGTACCGCCGCGAGTGCACCCGCGACTCC
GTGCTGCGCAGCCTGACCACCGTGAGCGGCGGCAGCTCCGAGGCCGGCCTGGTG
TGCGACCACCTGCTGCAGCTGGAGGGCGGCAGCGAGGTGCTGCGCGCCCGCACC
GAGTGGCGCCCCAAGCTGACCGACTCCTTCCGCGGCATCAGCGTGATCCCCGCCG
AGCCCCGCGTGATGGACTACAAGGACCACGACGGCGACTACAAGGACCACGACA
TCGACTACAAGGACGACGACGACAAGTGACTCGAGTTAATTAA

SEQ ID NO: 65
NeoR forward primer

CCGCCGTGCTGGACGTGGTG

SEQ ID NO: 66
NeoR reverse primer

GGTGGCGGGGTCCAGGGTGT

SEQ ID NO: 67
Yeast invertase *suc2* forward primer

CGGCCGGCGGCTCCTTCAAC

SEQ ID NO: 68
Yeast invertase *suc2* reverse primer

GGCGCTCCCGTAGGTCGGGT

SEQ ID NO: 69
5'UTR/promoter *Chlorella sorokiniana* glutamate dehydrogenase

CGCCTGCAACGCAAGGGCAGCCACAGCCGCTCCCACCCGCCGCTGAACCGACAC
GTGCTTGGGCGCCTGCCGCCTGCCTGCCGCATGCTTGTGCTGGTGAGGCTGGGCA
GTGCTGCCATGCTGATTGAGGCTTGGTTCATCGGGTGGAAGCTTATGTGTGTGCT
GGGCTTGCATGCCGGGCAATGCGCATGGTGGCAAGAGGGCGGCAGCACTTGCTG
GAGCTGCCGCGGTGCCTCCAGGTGGTTCAATCGCGGCAGCCAGAGGGATTTCAG
ATGATCGCGCGTACAGGTTGAGCAGCAGTGTCAGCAAAGGTAGCAGTTTGCCAG
AATGATCGGTTCAGCTGTTAATCAATGCCAGCAAGAGAAGGGGTCAAGTGCAAA
CACGGGCATGCCACAGCACGGGCACCGGGGAGTGGAATGGCACCACCAAGTGTG
TGCGAGCCAGCATCGCCGCCTGGCTGTTTCAGCTACAACGGCAGGAGTCATCCAA
CGTAACCATGAGCTGATCAACACTGCAATCATCGGGCGGGCGTGATGCAAGCAT
GCCTGGCGAAGACACATGGTGTGCGGATGCTGCCGGCTGCTGCCTGCTGCGCAC
GCCGTTGAGTTGGCAGCAGGCTCAGCCATGCACTGGATGGCAGCTGGGCTGCCA
CTGCAATGTGGTGGATAGGATGCAAGTGGAGCGAATACCAAACCCTCTGGCTGC
TTGCTGGGTTGCATGGCATCGCACCATCAGCAGGAGCGCATGCGAAGGGACTGG
CCCCATGCACGCCATGCCAAACCGGAGCGCACCGAGTGTCCACACTGTCACCAG
GCCCGCAAGCTTTGCAGAACCATGCTCATGGACGCATGTAGCGCTGACGTCCCTT
GACGGCGCTCCTCTCGGGTGTGGGAAACGCAATGCAGCACAGGCAGCAGAGGCG
GCGGCAGCAGAGCGGCGGCAGCAGCGGCGGGGGCCACCCTTCTTGCGGGGTCGC
GCCCCAGCCAGCGGTGATGCGCTGATCCCAAACGAGTTCACATTCATTTGCATGC
CTGGAGAAGCGAGGCTGGGGCCTTTGGGCTGGTGCAGCCCGCAATGGAATGCGG

GACCGCCAGGCTAGCAGCAAAGGCGCCTCCCCTACTCCGCATCGATGTTCCATAG
TGCATTGGACTGCATTTGGGTGGGGCGGCCGGCTGTTTCTTTCGTGTTGCAAAAC
GCGCCAGCTCAGCAACCTGTCCCGTGGGTCCCCCGTGCCGATGAAATCGTGTGCA
CGCCGATCAGCTGATTGCCCGGCTCGCGAAGTAGGCGCCCTCCTTTCTGCTCGCC
CTCTCTCCGTCCCGCCACTAGTGGCGCGCC

SEQ ID NO: 70
Umbellularia californica C12 thioesterase coding region

ATGGCCACCACCAGCCTGGCCTCCGCCTTCTGCTCCATGAAGGCCGTGATGCTGG
CCCGCGACGGCCGCGGCATGAAGCCCCGCAGCTCCGACCTGCAGCTGCGCGCCG
GCAACGCCCCCACCTCCCTGAAGATGATCAACGGCACCAAGTTCAGCTACACCG
AGAGCCTGAAGCGCCTGCCCGACTGGTCCATGCTGTTCGCCGTGATCACCACCAT
CTTCAGCGCCGCCGAGAAGCAGTGGACCAACCTGGAGTGGAAGCCCAAGCCCAA
GCTGCCCCAGCTGCTGGACGACCACTTCGGCCTGCACGGCCTGGTGTTCCGCCGC
ACCTTCGCCATCCGCTCCTACGAGGTGGGCCCCGACCGCAGCACCTCCATCCTGG
CCGTGATGAACCACATGCAGGAGGCCACCCTGAACCACGCCAAGAGCGTGGGCA
TCCTGGGCGACGGCTTCGGCACCACCCTGGAGATGTCCAAGCGCGACCTGATGTG
GGTGGTGCGCCGCACCCACGTGGCCGTGGAGCGCTACCCCACCTGGGGCGACAC
CGTGGAGGTGGAGTGCTGGATCGGCGCCAGCGGCAACAACGGCATGCGCCGCGA
CTTCCTGGTGCGCGACTGCAAGACCGGCGAGATCCTGACCCGCTGCACCTCCCTG
AGCGTGCTGATGAACACCCGCACCCGCCGCCTGAGCACCATCCCCGACGAGGTG
CGCGGCGAGATCGGCCCCGCCTTCATCGACAACGTGGCCGTGAAGGACGACGAG
ATCAAGAAGCTGCAGAAGCTGAACGACTCCACCGCCGACTACATCCAGGGCGGC
CTGACCCCCCGCTGGAACGACCTGGACGTGAACCAGCACGTGAACAACCTGAAG
TACGTGGCCTGGGTGTTCGAGACCGTGCCCGACAGCATCTTCGAGTCCCACCACA
TCAGCTCCTTCACCCTGGAGTACCGCCGCGAGTGCACCCGCGACTCCGTGCTGCG
CAGCCTGACCACCGTGAGCGGCGGCAGCTCCGAGGCCGGCCTGGTGTGCGACCA
CCTGCTGCAGCTGGAGGGCGGCAGCGAGGTGCTGCGCGCCCGCACCGAGTGGCG
CCCCAAGCTGACCGACTCCTTCCGCGGCATCAGCGTGATCCCCGCCGAGCCCCGC
GTG

SEQ ID NO: 71
Cinnamomum camphora C14 thioesterase coding region

ATGGCCACCACCTCCCTGGCCTCCGCCTTCTGCAGCATGAAGGCCGTGATGCTGG
CCCGCGACGGCCGCGGCATGAAGCCCCGCTCCAGCGACCTGCAGCTGCGCGCCG
GCAACGCCCAGACCTCCCTGAAGATGATCAACGGCACCAAGTTCTCCTACACCG
AGAGCCTGAAGAAGCTGCCCGACTGGTCCATGCTGTTCGCCGTGATCACCACCAT
CTTCTCCGCCGCCGAGAAGCAGTGGACCAACCTGGAGTGGAAGCCCAAGCCCAA
CCCCCCCCAGCTGCTGGACGACCACTTCGGCCCCCACGGCCTGGTGTTCCGCCGC
ACCTTCGCCATCCGCAGCTACGAGGTGGGCCCCGACCGCTCCACCAGCATCGTGG
CCGTGATGAACCACCTGCAGGAGGCCGCCCTGAACCACGCCAAGTCCGTGGGCA
TCCTGGGCGACGGCTTCGGCACCACCCTGGAGATGTCCAAGCGCGACCTGATCTG
GGTGGTGAAGCGCACCCACGTGGCCGTGGAGCGCTACCCCGCCTGGGGCGACAC
CGTGGAGGTGGAGTGCTGGGTGGGCGCCTCCGGCAACAACGGCCGCCGCCACGA
CTTCCTGGTGCGCGACTGCAAGACCGGCGAGATCCTGACCCGCTGCACCTCCCTG
AGCGTGATGATGAACACCCGCACCCGCCGCCTGAGCAAGATCCCCGAGGAGGTG
CGCGGCGAGATCGGCCCCGCCTTCATCGACAACGTGGCCGTGAAGGACGAGGAG
ATCAAGAAGCCCCAGAAGCTGAACGACTCCACCGCCGACTACATCCAGGGCGGC

CTGACCCCCCGCTGGAACGACCTGGACATCAACCAGCACGTGAACAACATCAAG
TACGTGGACTGGATCCTGGAGACCGTGCCCGACAGCATCTTCGAGAGCCACCAC
ATCTCCTCCTTCACCATCGAGTACCGCCGCGAGTGCACCATGGACAGCGTGCTGC
AGTCCCTGACCACCGTGAGCGGCGGCTCCTCCGAGGCCGGCCTGGTGTGCGAGC
ACCTGCTGCAGCTGGAGGGCGGCAGCGAGGTGCTGCGCGCCAAGACCGAGTGGC
GCCCCAAGCTGACCGACTCCTTCCGCGGCATCAGCGTGATCCCCGCCGAGTCCAG
CGTG

SEQ ID NO: 72
FLAG-tag sequence

ATGGACTACAAGGACCACGACGGCGACTACAAGGACCACGACATCGACTACAAG
GACGACGACGACAAGTGA

SEQ ID NO: 73
3' UTR *Chlorella vulgaris* nitrate reductase

CTCGAGGCAGCAGCAGCTCGGATAGTATCGACACACTCTGGACGCTGGTCGTGT
GATGGACTGTTGCCGCCACACTTGCTGCCTTGACCTGTGAATATCCCTGCCGCTTT
TATCAAACAGCCTCAGTGTGTTTGATCTTGTGTGTACGCGCTTTTGCGAGTTGCTA
GCTGCTTGTGCTATTTGCGAATACCACCCCCAGCATCCCCTTCCCTCGTTTCATAT
CGCTTGCATCCCAACCGCAACTTATCTACGCTGTCCTGCTATCCCTCAGCGCTGCT
CCTGCTCCTGCTCACTGCCCCTCGCACAGCCTTGGTTTGGGCTCCGCCTGTATTCT
CCTGGTACTGCAACCTGTAAACCAGCACTGCAATGCTGATGCACGGGAAGTAGT
GGGATGGGAACACAAATGGA

SEQ ID NO: 74
C12 thioesterase forward primer

5' CTGGGCGACGGCTTCGGCAC 3'

SEQ ID NO: 75
C12 thioesterase reverse primer

5' AAGTCGCGGCGCATGCCGTT 3'

SEQ ID NO: 76
C14 thioesterase forward primer

5' TACCCCGCCTGGGGCGACAC 3'

SEQ ID NO: 77
C14 thioesterase reverse primer

5' CTTGCTCAGGCGGCGGGTGC 3'

SEQ ID NO: 78
Codon optimized *Cuphea hookeriana* C8-10 thioesterase coding region with native plastid targeting sequence with 3x FLAG tag

ATGGTGGCCGCCGCCGCCTCCAGCGCCTTCTTCCCCGTGCCCGCCCCCGGCGCCT
CCCCCAAGCCCGGCAAGTTCGGCAACTGGCCCTCCAGCCTGAGCCCCTCCTTCAA
GCCCAAGTCCATCCCCAACGGCGGCTTCCAGGTGAAGGCCAACGACAGCGCCCA
CCCCAAGGCCAACGGCTCCGCCGTGAGCCTGAAGAGCGGCAGCCTGAACACCCA
GGAGGACACCTCCTCCAGCCCCCCCCCCCGCACCTTCCTGCACCAGCTGCCCGAC
TGGAGCCGCCTGCTGACCGCCATCACCACCGTGTTCGTGAAGTCCAAGCGCCCCG
ACATGCACGACCGCAAGTCCAAGCGCCCCGACATGCTGGTGGACAGCTTCGGCC
TGGAGTCCACCGTGCAGGACGGCCTGGTGTTCCGCCAGTCCTTCTCCATCCGCTC
CTACGAGATCGGCACCGACCGCACCGCCAGCATCGAGACCCTGATGAACCACCT
GCAGGAGACCTCCCTGAACCACTGCAAGAGCACCGGCATCCTGCTGGACGGCTT
CGGCCGCACCCTGGAGATGTGCAAGCGCGACCTGATCTGGGTGGTGATCAAGAT
GCAGATCAAGGTGAACCGCTACCCCGCCTGGGGCGACACCGTGGAGATCAACAC
CCGCTTCAGCCGCCTGGGCAAGATCGGCATGGGCCGCGACTGGCTGATCTCCGAC
TGCAACACCGGCGAGATCCTGGTGCGCGCCACCAGCGCCTACGCCATGATGAAC
CAGAAGACCCGCCGCCTGTCCAAGCTGCCCTACGAGGTGCACCAGGAGATCGTG
CCCCTGTTCGTGGACAGCCCCGTGATCGAGGACTCCGACCTGAAGGTGCACAAGT
TCAAGGTGAAGACCGGCGACAGCATCCAGAAGGGCCTGACCCCCGGCTGGAACG
ACCTGGACGTGAACCAGCACGTGTCCAACGTGAAGTACATCGGCTGGATCCTGG
AGAGCATGCCCACCGAGGTGCTGGAGACCCAGGAGCTGTGCTCCCTGGCCCTGG
AGTACCGCCGCGAGTGCGGCCGCGACTCCGTGCTGGAGAGCGTGACCGCCATGG
ACCCCAGCAAGGTGGGCGTGCGCTCCCAGTACCAGCACCTGCTGCGCCTGGAGG
ACGGCACCGCCATCGTGAACGGCGCCACCGAGTGGCGCCCCAAGAACGCCGGCG
CCAACGGCGCCATCTCCACCGGCAAGACCAGCAACGGCAACTCCGTGTCCATGG
ACTACAAGGACCACGACGGCGACTACAAGGACCACGACATCGACTACAAGGACG
ACGACGACAAGTGA

SEQ ID NO: 79

Cuphea hookeriana C8-10 thioesterase coding region and 3x FLAG tag, with *Chlorella protothecoides* stearoyl ACP desaturase plastid targeting sequence

ATGGCCACCGCATCCACTTTCTCGGCGTTCAATGCCCGCTGCGGCGACCTGCGTC
GCTCGGCGGGCTCCGGGCCCCGGCGCCCAGCGAGGCCCCTCCCCGTGCGCGGGC
GCGCCCAGCTGCCCGACTGGAGCCGCCTGCTGACCGCCATCACCACCGTGTTCGT
GAAGTCCAAGCGCCCCGACATGCACGACCGCAAGTCCAAGCGCCCCGACATGCT
GGTGGACAGCTTCGGCCTGGAGTCCACCGTGCAGGACGGCCTGGTGTTCCGCCA
GTCCTTCTCCATCCGCTCCTACGAGATCGGCACCGACCGCACCGCCAGCATCGAG
ACCCTGATGAACCACCTGCAGGAGACCTCCCTGAACCACTGCAAGAGCACCGGC
ATCCTGCTGGACGGCTTCGGCCGCACCCTGGAGATGTGCAAGCGCGACCTGATCT
GGGTGGTGATCAAGATGCAGATCAAGGTGAACCGCTACCCCGCCTGGGGCGACA
CCGTGGAGATCAACACCCGCTTCAGCCGCCTGGGCAAGATCGGCATGGGCCGCG
ACTGGCTGATCTCCGACTGCAACACCGGCGAGATCCTGGTGCGCGCCACCAGCG
CCTACGCCATGATGAACCAGAAGACCCGCCGCCTGTCCAAGCTGCCCTACGAGG
TGCACCAGGAGATCGTGCCCCTGTTCGTGGACAGCCCCGTGATCGAGGACTCCGA
CCTGAAGGTGCACAAGTTCAAGGTGAAGACCGGCGACAGCATCCAGAAGGGCCT
GACCCCCGGCTGGAACGACCTGGACGTGAACCAGCACGTGTCCAACGTGAAGTA
CATCGGCTGGATCCTGGAGAGCATGCCCACCGAGGTGCTGGAGACCCAGGAGCT
GTGCTCCCTGGCCCTGGAGTACCGCCGCGAGTGCGGCCGCGACTCCGTGCTGGAG
AGCGTGACCGCCATGGACCCCAGCAAGGTGGGCGTGCGCTCCCAGTACCAGCAC
CTGCTGCGCCTGGAGGACGGCACCGCCATCGTGAACGGCGCCACCGAGTGGCGC
CCCAAGAACGCCGGCGCCAACGGCGCCATCTCCACCGGCAAGACCAGCAACGGC

AACTCCGTGTCCATGGACTACAAGGACCACGACGGCGACTACAAGGACCACGAC
ATCGACTACAAGGACGACGACGACAAGTGA

SEQ ID NO: 80
Cuphea hookeriana C8-10 thioesterase coding region and 3x FLAG tag, with *Prototheca moriformis* delta 12 fatty acid desaturase plastid targeting sequence

ATGGCTATCAAGACGAACAGGCAGCCTGTGGAGAAGCCTCCGTTCACGATCGGG
ACGCTGCGCAAGGCCATCCCCGCGCACTGTTTCGAGCGCTCGGCGCTTCGTGGGC
GCGCCCAGCTGCCCGACTGGAGCCGCCTGCTGACCGCCATCACCACCGTGTTCGT
GAAGTCCAAGCGCCCCGACATGCACGACCGCAAGTCCAAGCGCCCCGACATGCT
GGTGGACAGCTTCGGCCTGGAGTCCACCGTGCAGGACGGCCTGGTGTTCCGCCA
GTCCTTCTCCATCCGCTCCTACGAGATCGGCACCGACCGCACCGCCAGCATCGAG
ACCCTGATGAACCACCTGCAGGAGACCTCCCTGAACCACTGCAAGAGCACCGGC
ATCCTGCTGGACGGCTTCGGCCGCACCCTGGAGATGTGCAAGCGCGACCTGATCT
GGGTGGTGATCAAGATGCAGATCAAGGTGAACCGCTACCCCGCCTGGGGCGACA
CCGTGGAGATCAACACCCGCTTCAGCCGCCTGGGCAAGATCGGCATGGGCCGCG
ACTGGCTGATCTCCGACTGCAACACCGGCGAGATCCTGGTGCGCGCCACCAGCG
CCTACGCCATGATGAACCAGAAGACCCGCCGCCTGTCCAAGCTGCCCTACGAGG
TGCACCAGGAGATCGTGCCCCTGTTCGTGGACAGCCCCGTGATCGAGGACTCCGA
CCTGAAGGTGCACAAGTTCAAGGTGAAGACCGGCGACAGCATCCAGAAGGGCCT
GACCCCCGGCTGGAACGACCTGGACGTGAACCAGCACGTGTCCAACGTGAAGTA
CATCGGCTGGATCCTGGAGAGCATGCCCACCGAGGTGCTGGAGACCCAGGAGCT
GTGCTCCCTGGCCCTGGAGTACCGCCGCGAGTGCGGCCGCGACTCCGTGCTGGAG
AGCGTGACCGCCATGGACCCCAGCAAGGTGGGCGTGCGCTCCCAGTACCAGCAC
CTGCTGCGCCTGGAGGACGGCACCGCCATCGTGAACGGCGCCACCGAGTGGCGC
CCCAAGAACGCCGGCGCCAACGGCGCCATCTCCACCGGCAAGACCAGCAACGGC
AACTCCGTGTCCATGGACTACAAGGACCACGACGGCGACTACAAGGACCACGAC
ATCGACTACAAGGACGACGACGACAAGTGA

SEQ ID NO: 81
Cuphea hookeriana C8-10 thioesterase coding region and 3x FLAG tag, with *Prototheca moriformis* isopentenyl diphosphate plastid targeting sequence

ATGACGTTCGGGGTCGCCCTCCCGGCCATGGGCCGCGGTGTCTCCCTTCCCCGGC
CCAGGGTCGCGGTGCGCGCCCAGTCGGCGAGTCAGGTTTTGGAGAGCGGGCGCG
CCCAGCTGCCCGACTGGAGCCGCCTGCTGACCGCCATCACCACCGTGTTCGTGAA
GTCCAAGCGCCCCGACATGCACGACCGCAAGTCCAAGCGCCCCGACATGCTGGT
GGACAGCTTCGGCCTGGAGTCCACCGTGCAGGACGGCCTGGTGTTCCGCCAGTCC
TTCTCCATCCGCTCCTACGAGATCGGCACCGACCGCACCGCCAGCATCGAGACCC
TGATGAACCACCTGCAGGAGACCTCCCTGAACCACTGCAAGAGCACCGGCATCC
TGCTGGACGGCTTCGGCCGCACCCTGGAGATGTGCAAGCGCGACCTGATCTGGGT
GGTGATCAAGATGCAGATCAAGGTGAACCGCTACCCCGCCTGGGGCGACACCGT
GGAGATCAACACCCGCTTCAGCCGCCTGGGCAAGATCGGCATGGGCCGCGACTG
GCTGATCTCCGACTGCAACACCGGCGAGATCCTGGTGCGCGCCACCAGCGCCTAC
GCCATGATGAACCAGAAGACCCGCCGCCTGTCCAAGCTGCCCTACGAGGTGCAC
CAGGAGATCGTGCCCCTGTTCGTGGACAGCCCCGTGATCGAGGACTCCGACCTGA
AGGTGCACAAGTTCAAGGTGAAGACCGGCGACAGCATCCAGAAGGGCCTGACCC
CCGGCTGGAACGACCTGGACGTGAACCAGCACGTGTCCAACGTGAAGTACATCG
GCTGGATCCTGGAGAGCATGCCCACCGAGGTGCTGGAGACCCAGGAGCTGTGCT

CCCTGGCCCTGGAGTACCGCCGCGAGTGCGGCCGCGACTCCGTGCTGGAGAGCG
TGACCGCCATGGACCCCAGCAAGGTGGGCGTGCGCTCCCAGTACCAGCACCTGC
TGCGCCTGGAGGACGGCACCGCCATCGTGAACGGCGCCACCGAGTGGCGCCCCA
AGAACGCCGGCGCCAACGGCGCCATCTCCACCGGCAAGACCAGCAACGGCAACT
CCGTGTCCATGGACTACAAGGACCACGACGGCGACTACAAGGACCACGACATCG
ACTACAAGGACGACGACGACAAGTGA

SEQ ID NO: 82
Umbellularia californica C12 thioesterase coding region and 3x FLAG tag, with *Prototheca moriformis* isopentenyl diphosphate plastid targeting sequence

ATGACGTTCGGGGTCGCCCTCCCGGCCATGGGCCGCGGTGTCTCCCTTCCCCGGC
CCAGGGTCGCGGTGCGCGCCCAGTCGGCGAGTCAGGTTTTGGAGAGCGGGCGCG
CCCCCGACTGGTCCATGCTGTTCGCCGTGATCACCACCATCTTCAGCGCCGCCGA
GAAGCAGTGGACCAACCTGGAGTGGAAGCCCAAGCCCAAGCTGCCCCAGCTGCT
GGACGACCACTTCGGCCTGCACGGCCTGGTGTTCCGCCGCACCTTCGCCATCCGC
TCCTACGAGGTGGGCCCCGACCGCAGCACCTCCATCCTGGCCGTGATGAACCACA
TGCAGGAGGCCACCCTGAACCACGCCAAGAGCGTGGGCATCCTGGGCGACGGCT
TCGGCACCACCCTGGAGATGTCCAAGCGCGACCTGATGTGGGTGGTGCGCCGCA
CCCACGTGGCCGTGGAGCGCTACCCCACCTGGGGCGACACCGTGGAGGTGGAGT
GCTGGATCGGCGCCAGCGGCAACAACGGCATGCGCCGCGACTTCCTGGTGCGCG
ACTGCAAGACCGGCGAGATCCTGACCCGCTGCACCTCCCTGAGCGTGCTGATGA
ACACCCGCACCCGCCGCCTGAGCACCATCCCCGACGAGGTGCGCGGCGAGATCG
GCCCCGCCTTCATCGACAACGTGGCCGTGAAGGACGACGAGATCAAGAAGCTGC
AGAAGCTGAACGACTCCACCGCCGACTACATCCAGGGCGGCCTGACCCCCCGCT
GGAACGACCTGGACGTGAACCAGCACGTGAACAACCTGAAGTACGTGGCCTGGG
TGTTCGAGACCGTGCCCGACAGCATCTTCGAGTCCCACCACATCAGCTCCTTCAC
CCTGGAGTACCGCCGCGAGTGCACCCGCGACTCCGTGCTGCGCAGCCTGACCACC
GTGAGCGGCGGCAGCTCCGAGGCCGGCCTGGTGTGCGACCACCTGCTGCAGCTG
GAGGGCGGCAGCGAGGTGCTGCGCGCCCGCACCGAGTGGCGCCCCAAGCTGACC
GACTCCTTCCGCGGCATCAGCGTGATCCCCGCCGAGCCCCGCGTGATGGACTACA
AGGACCACGACGGCGACTACAAGGACCACGACATCGACTACAAGGACGACGAC
GACAAGTGA

SEQ ID NO: 83
Umbellularia californica C12 thioesterase coding region and 3x FLAG tag, with *Prototheca moriformis* stearoyl ACP desaturase plastid targeting sequence

ATGGCTTCCGCGGCATTCACCATGTCGGCGTGCCCCGCGATGACTGGCAGGGCCC
CTGGGGCACGTCGCTCCGGACGGCCAGTCGCCACCCGCCTGAGGGGGCGCGCCC
CCGACTGGTCCATGCTGTTCGCCGTGATCACCACCATCTTCAGCGCCGCCGAGAA
GCAGTGGACCAACCTGGAGTGGAAGCCCAAGCCCAAGCTGCCCCAGCTGCTGGA
CGACCACTTCGGCCTGCACGGCCTGGTGTTCCGCCGCACCTTCGCCATCCGCTCC
TACGAGGTGGGCCCCGACCGCAGCACCTCCATCCTGGCCGTGATGAACCACATG
CAGGAGGCCACCCTGAACCACGCCAAGAGCGTGGGCATCCTGGGCGACGGCTTC
GGCACCACCCTGGAGATGTCCAAGCGCGACCTGATGTGGGTGGTGCGCCGCACC
CACGTGGCCGTGGAGCGCTACCCCACCTGGGGCGACACCGTGGAGGTGGAGTGC
TGGATCGGCGCCAGCGGCAACAACGGCATGCGCCGCGACTTCCTGGTGCGCGAC
TGCAAGACCGGCGAGATCCTGACCCGCTGCACCTCCCTGAGCGTGCTGATGAAC
ACCCGCACCCGCCGCCTGAGCACCATCCCCGACGAGGTGCGCGGCGAGATCGGC

CCCGCCTTCATCGACAACGTGGCCGTGAAGGACGACGAGATCAAGAAGCTGCAG
AAGCTGAACGACTCCACCGCCGACTACATCCAGGGCGGCCTGACCCCCCGCTGG
AACGACCTGGACGTGAACCAGCACGTGAACAACCTGAAGTACGTGGCCTGGGTG
TTCGAGACCGTGCCCGACAGCATCTTCGAGTCCCACCACATCAGCTCCTTCACCC
TGGAGTACCGCCGCGAGTGCACCCGCGACTCCGTGCTGCGCAGCCTGACCACCGT
GAGCGGCGGCAGCTCCGAGGCCGGCCTGGTGTGCGACCACCTGCTGCAGCTGGA
GGGCGGCAGCGAGGTGCTGCGCGCCCGCACCGAGTGGCGCCCCAAGCTGACCGA
CTCCTTCCGCGGCATCAGCGTGATCCCCGCCGAGCCCCGCGTGATGGACTACAAG
GACCACGACGGCGACTACAAGGACCACGACATCGACTACAAGGACGACGACGA
CAAGTGA

SEQ ID NO: 84
Umbellularia californica C12 thioesterase coding region and 3x FLAG tag, with *Chlorella protothecoides* stearoyl ACP desaturase plastid targeting sequence

ATGGCCACCGCATCCACTTTCTCGGCGTTCAATGCCCGCTGCGGCGACCTGCGTC
GCTCGGCGGGCTCCGGGCCCCGGCGCCCAGCGAGGCCCCTCCCCGTGCGCGGGC
GCGCCCCCGACTGGTCCATGCTGTTCGCCGTGATCACCACCATCTTCAGCGCCGC
CGAGAAGCAGTGGACCAACCTGGAGTGGAAGCCCAAGCCCAAGCTGCCCCAGCT
GCTGGACGACCACTTCGGCCTGCACGGCCTGGTGTTCCGCCGCACCTTCGCCATC
CGCTCCTACGAGGTGGGCCCCGACCGCAGCACCTCCATCCTGGCCGTGATGAACC
ACATGCAGGAGGCCACCCTGAACCACGCCAAGAGCGTGGGCATCCTGGGCGACG
GCTTCGGCACCACCCTGGAGATGTCCAAGCGCGACCTGATGTGGGTGGTGCGCC
GCACCCACGTGGCCGTGGAGCGCTACCCCACCTGGGGCGACACCGTGGAGGTGG
AGTGCTGGATCGGCGCCAGCGGCAACAACGGCATGCGCCGCGACTTCCTGGTGC
GCGACTGCAAGACCGGCGAGATCCTGACCCGCTGCACCTCCCTGAGCGTGCTGAT
GAACACCCGCACCCGCCGCCTGAGCACCATCCCCGACGAGGTGCGCGGCGAGAT
CGGCCCCCGCCTTCATCGACAACGTGGCCGTGAAGGACGACGAGATCAAGAAGCT
GCAGAAGCTGAACGACTCCACCGCCGACTACATCCAGGGCGGCCTGACCCCCCG
CTGGAACGACCTGGACGTGAACCAGCACGTGAACAACCTGAAGTACGTGGCCTG
GGTGTTCGAGACCGTGCCCGACAGCATCTTCGAGTCCCACCACATCAGCTCCTTC
ACCCTGGAGTACCGCCGCGAGTGCACCCGCGACTCCGTGCTGCGCAGCCTGACC
ACCGTGAGCGGCGGCAGCTCCGAGGCCGGCCTGGTGTGCGACCACCTGCTGCAG
CTGGAGGGCGGCAGCGAGGTGCTGCGCGCCCGCACCGAGTGGCGCCCCAAGCTG
ACCGACTCCTTCCGCGGCATCAGCGTGATCCCCGCCGAGCCCCGCGTGATGGACT
ACAAGGACCACGACGGCGACTACAAGGACCACGACATCGACTACAAGGACGAC
GACGACAAGTGATGA

SEQ ID NO: 85
Umbellularia californica C12 thioesterase coding region and 3x FLAG tag, with *Prototheca moriformis* delta 12 fatty acid desaturase plastid targeting sequence

ATGGCTATCAAGACGAACAGGCAGCCTGTGGAGAAGCCTCCGTTCACGATCGGG
ACGCTGCGCAAGGCCATCCCCGCGCACTGTTTCGAGCGCTCGGCGCTTCGTGGGC
GCGCCCCCGACTGGTCCATGCTGTTCGCCGTGATCACCACCATCTTCAGCGCCGC
CGAGAAGCAGTGGACCAACCTGGAGTGGAAGCCCAAGCCCAAGCTGCCCCAGCT
GCTGGACGACCACTTCGGCCTGCACGGCCTGGTGTTCCGCCGCACCTTCGCCATC
CGCTCCTACGAGGTGGGCCCCGACCGCAGCACCTCCATCCTGGCCGTGATGAACC
ACATGCAGGAGGCCACCCTGAACCACGCCAAGAGCGTGGGCATCCTGGGCGACG
GCTTCGGCACCACCCTGGAGATGTCCAAGCGCGACCTGATGTGGGTGGTGCGCC

GCACCCACGTGGCCGTGGAGCGCTACCCCACCTGGGGCGACACCGTGGAGGTGG
AGTGCTGGATCGGCGCCAGCGGCAACAACGGCATGCGCCGCGACTTCCTGGTGC
GCGACTGCAAGACCGGCGAGATCCTGACCCGCTGCACCTCCCTGAGCGTGCTGAT
GAACACCCGCACCCGCCGCCTGAGCACCATCCCCGACGAGGTGCGCGGCGAGAT
CGGCCCCGCCTTCATCGACAACGTGGCCGTGAAGGACGACGAGATCAAGAAGCT
GCAGAAGCTGAACGACTCCACCGCCGACTACATCCAGGGCGGCCTGACCCCCCG
CTGGAACGACCTGGACGTGAACCAGCACGTGAACAACCTGAAGTACGTGGCCTG
GGTGTTCGAGACCGTGCCCGACAGCATCTTCGAGTCCCACCACATCAGCTCCTTC
ACCCTGGAGTACCGCCGCGAGTGCACCCGCGACTCCGTGCTGCGCAGCCTGACC
ACCGTGAGCGGCGGCAGCTCCGAGGCCGGCCTGGTGTGCGACCACCTGCTGCAG
CTGGAGGGCGGCAGCGAGGTGCTGCGCGCCCGCACCGAGTGGCGCCCCAAGCTG
ACCGACTCCTTCCGCGGCATCAGCGTGATCCCCGCCGAGCCCCGCGTGATGGACT
ACAAGGACCACGACGGCGACTACAAGGACCACGACATCGACTACAAGGACGAC
GACGACAAGTGA

SEQ ID NO: 86
Cinnamomum camphorum C14 thioesterase coding region and 3x FLAG tag, with *Chlorella protothecoides* stearoyl ACP desaturase plastid targeting sequence

ATGGCCACCGCATCCACTTTCTCGGCGTTCAATGCCCGCTGCGGCGACCTGCGTC
GCTCGGCGGGCTCCGGGCCCCGGCGCCCAGCGAGGCCCCTCCCCGTGCGCGGGC
GCGCCCCCGACTGGTCCATGCTGTTCGCCGTGATCACCACCATCTTCTCCGCCGC
CGAGAAGCAGTGGACCAACCTGGAGTGGAAGCCCAAGCCCAACCCCCCCCAGCT
GCTGGACGACCACTTCGGCCCCCACGGCCTGGTGTTCCGCCGCACCTTCGCCATC
CGCAGCTACGAGGTGGGCCCCGACCGCTCCACCAGCATCGTGGCCGTGATGAAC
CACCTGCAGGAGGCCGCCCTGAACCACGCCAAGTCCGTGGGCATCCTGGGCGAC
GGCTTCGGCACCACCCTGGAGATGTCCAAGCGCGACCTGATCTGGGTGGTGAAG
CGCACCCACGTGGCCGTGGAGCGCTACCCCGCCTGGGGCGACACCGTGGAGGTG
GAGTGCTGGGTGGGCGCCTCCGGCAACAACGGCCGCCGCCACGACTTCCTGGTG
CGCGACTGCAAGACCGGCGAGATCCTGACCCGCTGCACCTCCCTGAGCGTGATG
ATGAACACCCGCACCCGCCGCCTGAGCAAGATCCCCGAGGAGGTGCGCGGCGAG
ATCGGCCCCGCCTTCATCGACAACGTGGCCGTGAAGGACGAGGAGATCAAGAAG
CCCCAGAAGCTGAACGACTCCACCGCCGACTACATCCAGGGCGGCCTGACCCCC
CGCTGGAACGACCTGGACATCAACCAGCACGTGAACAACATCAAGTACGTGGAC
TGGATCCTGGAGACCGTGCCCGACAGCATCTTCGAGAGCCACCACATCTCCTCCT
TCACCATCGAGTACCGCCGCGAGTGCACCATGGACAGCGTGCTGCAGTCCCTGAC
CACCGTGAGCGGCGGCTCCTCCGAGGCCGGCCTGGTGTGCGAGCACCTGCTGCA
GCTGGAGGGCGGCAGCGAGGTGCTGCGCGCCAAGACCGAGTGGCGCCCCAAGCT
GACCGACTCCTTCCGCGGCATCAGCGTGATCCCCGCCGAGTCCAGCGTGATGGAC
TACAAGGACCACGACGGCGACTACAAGGACCACGACATCGACTACAAGGACGAC
GACGACAAGTGATGA

SEQ ID NO: 87
C. reinhardtii β-tublulin::*C. camphora* C14 thioesterase with *C.protothecoides* stearoyl ACP desaturase plastid targeting sequence::3xFLAG::*C.vulgaris* NitRed expression construct element

gaattcctttcttgcgctatgacacttccagcaaaaggtagggcgggctgcgagacggcttcccggcgctgcatgcaacaccgatgat
gcttcgaccccccgaagctccttcggggctgcatgggcgctccgatgccgctccagggcgagcgctgtttaaatagccaggccccc
gattgcaaagacattatagcgagctaccaaagccatattcaaacacctagatcactaccacttctacacaggccactcgagcttgtgatc

gcactccgctaaggggcgcctcttcctcttcgtttcagtcacaacccgcaaacactagtATGGCCACCGCATCCACT
TTCTCGGCGTTCAATGCCCGCTGCGGCGACCTGCGTCGCTCGGCGGGCTCCGGGC
CCCGGCGCCCAGCGAGGCCCCTCCCCGTGCGCGGGCGCGCCCCCGACTGGTCCAT
GCTGTTCGCCGTGATCACCACCATCTTCTCCGCCGCCGAGAAGCAGTGGACCAAC
CTGGAGTGGAAGCCCAAGCCCAACCCCCCCCCAGCTGCTGGACGACCACTTCGGC
CCCCACGGCCTGGTGTTCCGCCGCACCTTCGCCATCCGCAGCTACGAGGTGGGCC
CCGACCGCTCCACCAGCATCGTGGCCGTGATGAACCACCTGCAGGAGGCCGCCC
TGAACCACGCCAAGTCCGTGGGCATCCTGGGCGACGGCTTCGGCACCACCCTGG
AGATGTCCAAGCGCGACCTGATCTGGGTGGTGAAGCGCACCCACGTGGCCGTGG
AGCGCTACCCCCGCCTGGGGCGACACCGTGGAGGTGGAGTGCTGGGTGGGCGCCT
CCGGCAACAACGGCCGCCGCCACGACTTCCTGGTGCGCGACTGCAAGACCGGCG
AGATCCTGACCCGCTGCACCTCCCTGAGCGTGATGATGAACACCCGCACCCGCCG
CCTGAGCAAGATCCCCGAGGAGGTGCGCGGCGAGATCGGCCCCGCCTTCATCGA
CAACGTGGCCGTGAAGGACGAGGAGATCAAGAAGCCCCAGAAGCTGAACGACT
CCACCGCCGACTACATCCAGGGCGGCCTGACCCCCCGCTGGAACGACCTGGACA
TCAACCAGCACGTGAACAACATCAAGTACGTGGACTGGATCCTGGAGACCGTGC
CCGACAGCATCTTCGAGAGCCACCACATCTCCTCCTTCACCATCGAGTACCGCCG
CGAGTGCACCATGGACAGCGTGCTGCAGTCCCTGACCACCGTGAGCGGCGGCTC
CTCCGAGGCCGGCCTGGTGTGCGAGCACCTGCTGCAGCTGGAGGGCGGCAGCGA
GGTGCTGCGCGCCAAGACCGAGTGGCGCCCCAAGCTGACCGACTCCTTCCGCGG
CATCAGCGTGATCCCCGCCGAGTCCAGCGTGATGGACTACAAGGACCACGACGG
CGACTACAAGGACCACGACATCGACTACAAGGACGACGACGACAAGTGATGActc
gaggcagcagcagctcggatagtatcgacacactctggacgctggtcgtgtgatggactgttgccgccacacttgctgccttgacctgt
gaatatccctgccgcttttatcaaacagcctcagtgtgtttgatcttgtgtgtacgcgcttttgcgagttgctagctgcttgtgctatttgcga
ataccacccccagcatccccttccctcgtttcatatcgcttgcatcccaaccgcaacttatctacgctgtcctgctatccctcagcgctgct
cctgctcctgctcactgcccctcgcacagccttggtttgggctccgcctgtattctcctggtactgcaacctgtaaaccagcactgcaatg
ctgatgcacgggaagtagtgggatgggaacacaaatggaaagctt

SEQ ID NO: 88

C. reinhardtii β-tublulin::*C. camphora* C14 thioesterase with *P.moriformis* stearoyl ACP desaturase plastid targeting sequence::3xFLAG::*C.vulgaris* NitRed expression construct element

gaattcctttcttgcgctatgacacttccagcaaaaggtagggcgggctgcgagacggcttcccggcgctgcatgcaacaccgatgat
gcttcgaccccccgaagctccttcggggctgcatgggcgctccgatgccgctccagggcgagcgctgtttaaatagccaggccccc
gattgcaaagacattatagcgagctaccaaagccatattcaaacacctagatcactaccacttctacacaggccactcgagcttgtgatc
gcactccgctaaggggcgcctcttcctcttcgtttcagtcacaacccgcaaacactagtATGGCTTCCGCGGCATTC
ACCATGTCGGCGTGCCCCGCGATGACTGGCAGGGCCCCTGGGGCACGTCGCTCC
GGACGGCCAGTCGCCACCCGCCTGAGGGGGCGCGCCCCCGACTGGTCCATGCTG
TTCGCCGTGATCACCACCATCTTCTCCGCCGCCGAGAAGCAGTGGACCAACCTGG
AGTGGAAGCCCAAGCCCAACCCCCCCCCAGCTGCTGGACGACCACTTCGGCCCCC
ACGGCCTGGTGTTCCGCCGCACCTTCGCCATCCGCAGCTACGAGGTGGGCCCCGA
CCGCTCCACCAGCATCGTGGCCGTGATGAACCACCTGCAGGAGGCCGCCCTGAA
CCACGCCAAGTCCGTGGGCATCCTGGGCGACGGCTTCGGCACCACCCTGGAGAT
GTCCAAGCGCGACCTGATCTGGGTGGTGAAGCGCACCCACGTGGCCGTGGAGCG
CTACCCCGCCTGGGGCGACACCGTGGAGGTGGAGTGCTGGGTGGGCGCCTCCGG
CAACAACGGCCGCCGCCACGACTTCCTGGTGCGCGACTGCAAGACCGGCGAGAT
CCTGACCCGCTGCACCTCCCTGAGCGTGATGATGAACACCCGCACCCGCCGCCTG
AGCAAGATCCCCGAGGAGGTGCGCGGCGAGATCGGCCCCGCCTTCATCGACAAC
GTGGCCGTGAAGGACGAGGAGATCAAGAAGCCCCAGAAGCTGAACGACTCCACC

GCCGACTACATCCAGGGCGGCCTGACCCCCCGCTGGAACGACCTGGACATCAAC
CAGCACGTGAACAACATCAAGTACGTGGACTGGATCCTGGAGACCGTGCCCGAC
AGCATCTTCGAGAGCCACCACATCTCCTCCTTCACCATCGAGTACCGCCGCGAGT
GCACCATGGACAGCGTGCTGCAGTCCCTGACCACCGTGAGCGGCGGCTCCTCCG
AGGCCGGCCTGGTGTGCGAGCACCTGCTGCAGCTGGAGGGCGGCAGCGAGGTGC
TGCGCGCCAAGACCGAGTGGCGCCCCAAGCTGACCGACTCCTTCCGCGGCATCA
GCGTGATCCCCCGCCGAGTCCAGCGTGATGGACTACAAGGACCACGACGGCGACT
ACAAGGACCACGACATCGACTACAAGGACGACGACGACAAGTGATGActcgaggcag
cagcagctcggatagtatcgacacactctggacgctggtcgtgtgatggactgttgccgccacacttgctgccttgacctgtgaatatcc
ctgccgcttttatcaaacagcctcagtgtgtttgatcttgtgtgtacgcgcttttgcgagttgctagctgcttgtgctatttgcgaataccacc
cccagcatcccctttccctcgtttcatatcgcttgcatcccaaccgcaacttatctacgctgtcctgctatccctcagcgctgctcctgctcct
gctcactgcccctcgcacagccttggtttgggctccgcctgtattctcctggtactgcaacctgtaaaccagcactgcaatgctgatgca
cgggaagtagtgggatgggaacacaaatggaaagctt

SEQ ID NO: 89

Arabidopsis thaliana invertase sequence (with targeting signal from yeast suc2: *C. sorokiniana* glutamate dehydrogenase promoter and *C. vulgaris* nitrate reductase 3'UTR:3xFLAG)

GAATTCCGCCTGCAACGCAAGGGCAGCCACAGCCGCTCCCACCCGCCGCTGAAC
CGACACGTGCTTGGGCGCCTGCCGCCTGCCTGCCGCATGCTTGTGCTGGTGAGGC
TGGGCAGTGCTGCCATGCTGATTGAGGCTTGGTTCATCGGGTGGAAGCTTATGTG
TGTGCTGGGCTTGCATGCCGGGCAATGCGCATGGTGGCAAGAGGGCGGCAGCAC
TTGCTGGAGCTGCCGCGGTGCCTCCAGGTGGTTCAATCGCGGCAGCCAGAGGGA
TTTCAGATGATCGCGCGTACAGGTTGAGCAGCAGTGTCAGCAAAGGTAGCAGTTT
GCCAGAATGATCGGTTCAGCTGTTAATCAATGCCAGCAAGAGAAGGGGTCAAGT
GCAAACACGGGCATGCCACAGCACGGGCACCGGGGAGTGGAATGGCACCACCA
AGTGTGTGCGAGCCAGCATCGCCGCCTGGCTGTTTCAGCTACAACGGCAGGAGTC
ATCCAACGTAACCATGAGCTGATCAACACTGCAATCATCGGGCGGGCGTGATGC
AAGCATGCCTGGCGAAGACACATGGTGTGCGGATGCTGCCGGCTGCTGCCTGCT
GCGCACGCCGTTGAGTTGGCAGCAGGCTCAGCCATGCACTGGATGGCAGCTGGG
CTGCCACTGCAATGTGGTGGATAGGATGCAAGTGGAGCGAATACCAAACCCTCT
GGCTGCTTGCTGGGTTGCATGGCATCGCACCATCAGCAGGAGCGCATGCGAAGG
GACTGGCCCCATGCACGCCATGCCAAACCGGAGCGCACCGAGTGTCCACACTGT
CACCAGGCCCGCAAGCTTTGCAGAACCATGCTCATGGACGCATGTAGCGCTGAC
GTCCCTTGACGGCGCTCCTCTCGGGTGTGGGAAACGCAATGCAGCACAGGCAGC
AGAGGCGGCGGCAGCAGAGCGGCGGCAGCAGCGGCGGGGGCCACCCTTCTTGCG
GGGTCGCGCCCCAGCCAGCGGTGATGCGCTGATCCCAAACGAGTTCACATTCATT
TGCATGCCTGGAGAAGCGAGGCTGGGGCCTTTGGGCTGGTGCAGCCCGCAATGG
AATGCGGGACCGCCAGGCTAGCAGCAAAGGCGCCTCCCCTACTCCGCATCGATG
TTCCATAGTGCATTGGACTGCATTTGGGTGGGGCGGCCGGCTGTTTCTTTCGTGTT
GCAAAACGCGCCAGCTCAGCAACCTGTCCCGTGGGTCCCCCGTGCCGATGAAAT
CGTGTGCACGCCGATCAGCTGATTGCCCGGCTCGCGAAGTAGGCGCCCTCCTTTC
TGCTCGCCCTCTCTCCGTCCCGCCACTAGTATGCTGCTGCAGGCCTTCCTGTTCCT
GCTGGCCGGCTTCGCCGCCAAGATCAGCGCCTCCATGACGAACGAGACGTCCGA
CCGCCCCCTGGTGCACTTCACCCCCAACAAGGGCTGGGGGCGCGCCAGCCACCA
CGTGTACAAGCGCCTGACCCAGAGCACCAACACCAAGTCCCCCAGCGTGAACCA
GCCCTACCGCACCGGCTTCCACTTCCAGCCCCCCAAGAACTGGATGAACGACCCC
AACGGCCCCATGATCTACAAGGGCATCTACCACCTGTTCTACCAGTGGAACCCCA
AGGGCGCCGTGTGGGGCAACATCGTGTGGGCCCACTCCACCAGCACCGACCTGA

TCAACTGGGACCCCCACCCCCCCGCCATCTTCCCCAGCGCCCCCTTCGACATCAA
CGGCTGCTGGTCCGGCAGCGCCACCATCCTGCCCAACGGCAAGCCCGTGATCCTG
TACACCGGCATCGACCCCAAGAACCAGCAGGTGCAGAACATCGCCGAGCCCAAG
AACCTGTCCGACCCCTACCTGCGCGAGTGGAAGAAGAGCCCCCTGAACCCCCTG
ATGGCCCCCGACGCCGTGAACGGCATCAACGCCTCCAGCTTCCGCGACCCCACCA
CCGCCTGGCTGGGCCAGGACAAGAAGTGGCGCGTGATCATCGGCTCCAAGATCC
ACCGCCGCGGCCTGGCCATCACCTACACCAGCAAGGACTTCCTGAAGTGGGAGA
AGTCCCCCGAGCCCCTGCACTACGACGACGGCAGCGGCATGTGGGAGTGCCCCG
ACTTCTTCCCCGTGACCCGCTTCGGCAGCAACGGCGTGGAGACCTCCAGCTTCGG
CGAGCCCAACGAGATCCTGAAGCACGTGCTGAAGATCTCCCTGGACGACACCAA
GCACGACTACTACACCATCGGCACCTACGACCGCGTGAAGGACAAGTTCGTGCC
CGACAACGGCTTCAAGATGGACGGCACCGCCCCCCGCTACGACTACGGCAAGTA
CTACGCCAGCAAGACCTTCTTCGACTCCGCCAAGAACCGCCGCATCCTGTGGGGC
TGGACCAACGAGTCCTCCAGCGTGGAGGACGACGTGGAGAAGGGCTGGTCCGGC
ATCCAGACCATCCCCCGCAAGATCTGGCTGGACCGCAGCGGCAAGCAGCTGATC
CAGTGGCCCGTGCGCGAGGTGGAGCGCCTGCGCACCAAGCAGGTGAAGAACCTG
CGCAACAAGGTGCTGAAGTCCGGCAGCCGCCTGGAGGTGTACGGCGTGACCGCC
GCCCAGGCCGACGTGGAGGTGCTGTTCAAGGTGCGCGACCTGGAGAAGGCCGAC
GTGATCGAGCCCTCCTGGACCGACCCCCAGCTGATCTGCAGCAAGATGAACGTGT
CCGTGAAGTCCGGCCTGGGCCCCTTCGGCCTGATGGTGCTGGCCAGCAAGAACCT
GGAGGAGTACACCTCCGTGTACTTCCGCATCTTCAAGGCCCGCCAGAACAGCAA
CAAGTACGTGGTGCTGATGTGCTCCGACCAGTCCCGCAGCTCCCTGAAGGAGGA
CAACGACAAGACCACCTACGGCGCCTTCGTGGACATCAACCCCCACCAGCCCCT
GAGCCTGCGCGCCCTGATCGACCACTCCGTGGTGGAGAGCTTCGGCGGCAAGGG
CCGCGCCTGCATCACCTCCCGCGTGTACCCCAAGCTGGCCATCGGCAAGTCCAGC
CACCTGTTCGCCTTCAACTACGGCTACCAGTCCGTGGACGTGCTGAACCTGAACG
CCTGGAGCATGAACTCCGCCCAGATCAGCATGGACTACAAGGACCACGACGGCG
ACTACAAGGACCACGACATCGACTACAAGGACGACGACGACAAGTGATTAATTA
ACCGG<u>CTCGAG</u>*GCAGCAGCAGCTCGGATAGTATCGA*CACACTCTGGACGCTGGTCG
TGTGATGGACTGTTGCCGCCACACTTGCTGCCTTGACCTGTGAATATCCCTGCCG
CTTTTATCAAACAGCCTCAGTGTGTTTGATCTTGTGTGTACGCGCTTTTGCGAGTT
GCTAGCTGCTTGTGCTATTTGCGAATACCACCCCCAGCATCCCCTTCCCTCGTTTC
ATATCGCTTGCATCCCAACCGCAACTTATCTACGCTGTCCTGCTATCCCTCAGCGC
TGCTCCTGCTCCTGCTCACTGCCCCTCGCACAGCCTTGGTTTGGGCTCCGCCTGTA
TTCTCCTGGTACTGCAACCTGTAAACCAGCACTGCAATGCTGATGCACGGGAAGT
AGTGGGATGGGAACACAAATGGAAAGCTT<u>GAGCTC</u>

SEQ ID NO: 90

Translated amino acid sequence of the *Arabidopsis thaliana* invertase expression construct

MLLQAFLFLLAGFAAKISASMTNETSDRPLVHFTPNKGWGRASHHVYKRLTQSTNT
KSPSVNQPYRTGFHFQPPKNWMNDPNGPMIYKGIYHLFYQWNPKGAVWGNIVWAH
STSTDLINWDPHPPAIFPSAPFDINGCWSGSATILPNGKPVILYTGIDPKNQQVQNIAEP
KNLSDPYLREWKKSPLNPLMAPDAVNGINASSFRDPTTAWLGQDKKWRVIIGSKIHR
RGLAITYTSKDFLKWEKSPEPLHYDDGSGMWECPDFFPVTRFGSNGVETSSFGEPNEI
LKHVLKISLDDTKHDYYTIGTYDRVKDKFVPDNGFKMDGTAPRYDYGKYYASKTFF
DSAKNRRILWGWTNESSSVEDDVEKGWSGIQTIPRKIWLDRSGKQLIQWPVREVERL
RTKQVKNLRNKVLKSGSRLEVYGVTAAQADVEVLFKVRDLEKADVIEPSWTDPQLI
CSKMNVSVKSGLGPFGLMVLASKNLEEYTSVYFRIFKARQNSNKYVVLMCSDQSRS

SLKEDNDKTTYGAFVDINPHQPLSLRALIDHSVVESFGGKGRACITSRVYPKLAIGKS
SHLFAFNYGYQSVDVLNLNAWSMNSAQISMDYKDHDGDYKDHDIDYKDDDDK

SEQ ID NO: 91
Prototheca moriformis FatB/A promoter/5'UTR
CCTGTCGATCGAAGAGAAGGAGACATGTGTACATTATTGGTGTGAGGGCGCTGA
ATCGGCCATTTTTTAAAATGATCACGCTCATGCCAATAGACGCGGCACATAACGA
CGTTCAAACCCCCGCCAAAGCCGCGGACAACCCCATCCCTCCACACCCCCCACAC
AAAGAACCCGCCACCGCTTACCTTGCCCACGAGGTAGGCCTTTCGTTGCGCAAAA
CCGGCCTCGGTGATGAATGCATGCCCGTTCCTGACGAGCGCTGCCCGGGCCAACA
CGCTCTTTTGCTGCGTCTCCTCAGGCTTGGGGGCCTCCTTGGGCTTGGGTGCCGCC
ATGATCTGCGCGCATCAGAGAAACGTTGCTGGTAAAAAGGAGCGCCCGGCTGCG
CAATATATATATAGGCATGCCAACACAGCCCAACCTCACTCGGGAGCCCGTCCCA
CCACCCCCAAGTCGCGTGCCTTGACGGCATACTGCTGCAGAAGCTTCATGAGAAT
GATGCCGAACAAGAGGGGCACGAGGACCCAATCCCGGACATCCTTGTCGATAAT
GATCTCGTGAGTCCCCATCGTCCGCCCGACGCTCCGGGGAGCCCGCCGATGCTCA
AGACGAGAGGGCCCTCGACCAGGAGGGGCTGGCCCGGGCGGGCACTGGCGTCG
AAGGTGCGCCCGTCGTTCGCCTGCAGTCCTATGCCACAAAACAAGTCTTCTGACG
GGGTGCGTTTGCTCCCGTGCGGGCAGGCAACAGAGGTATTCACCCTGGTCATGGG
GAGATCGGCGATCGAGCTGGGATAAGAGATACTTCTGGCAAGCAATGACAACTT
GTCAGGACCGGACCGTGCCATATATTTCTCACCTAGCGCCGCAAAACCTAACAAT
TTGGGAGTCACTGTGCCACTGAGTTCGACTGGTAGCTGAATGGAGTCGCTGCTCC
ACTAAACGAATTGTCAGCACCGCCAGCCGGCCGAGGACCCGAGTCATAGCGAGG
GTAGTAGCGCGCC

SEQ ID NO: 92
Prototheca moriformis NRAMP metal transporter promoter/5'UTR
ACTAATTGCAATCGTGCAGTAATCATCGATATGGTCACAAGTAGATCCCCTACTG
ACACCCTCTCGTACATGTAGGCAATGTCATCGGCGCCGTCCTGCTGACCGATGCC
GACGTAGCAGAGCAGACCCGGGCCGATCTGGGATACGAGCCGGCCCTCCACCTG
CGCTCGAGGTGGAATCAAGTAAATAACCAATACACTTTTCGACACCACACAGAG
TTGCACGGACGGTGGCGTACCTCTACGCTCGCGCTCTTCACGCGCTGGACGACCG
CACGCATGAGCCCGGGTGGCTTGGTCTGGGCTGCAAAAATGCACAACAAACAAG
TATCAGACGCTCATGGATGCACACGCGCTCCCAAGCACGCTCAGACTAAATATTA
CAGTAGCTCGTATCTGATAAGATATCGAGACATACCGCTCAACTCACCCGCAAAC
TGCGCCCCGCCAGGTGATGCGCACAGGGCCCCACCATGCGATCCATCGCATCGCT
CCTCGAGGGCGCTATCACGTGGCCGGAGAGCGTTCACAGCGTACGCCACTGTATC
TGGGCGGTATGCGGTCCGTCAACATGGAGACAGATACCCGCACCACCACCTTGC
AAGCTCTTCCATATTGGAAGTAGAAAATTGTAATTGTATCATCGCACGAGGGGCC
AACTTGCCGTCGGCGAGCTGGGCGACGAACACCACCTGGACGTTGTCGAGACTC
GCTCGTGCCGTGCGCCGGGCCGCTGGGTATCCAGACCGTCGCC

SEQ ID NO: 93
Prototheca moriformis FLAP Flagellar-associated protein promoter/5'UTR
CAACGACAACCAGCAGGCAACTCGGTCAGCGACCCAACACGCGAGTCAAATTGT
TGCGTGTTCTTGCCTTGTCTATTTACTGTGATAGCAAGACTGTCGGTCAGTCAATA
CCGCGGTGCGCACGTCGGGGTGCCAAGCCTAGCAGAGCACGGGACGGCTGGTGC
TGTGCGCCAGCTCAGCTCGCTTCGCGACCAATTGTAGGACCGGCAAAGTCACCAA
AACATGCCAGCGGTGCGATTCAATTGGTCATGAGCTCTACAAAATTGTTTTGTGC
GTCGCGCAGGTATCCAACGGCGCGGCAGAGAAAGTTTGACAGCTCTCGATTTCAT

CTCGGAAAAATGGGGAGAATTTATGACACACAAGTGCGCAGGCGGCCCAGGCGG
CCAGCATATTCTGGCGTGACCTGGGCCGCCCACAAAATGCTTGGATGCACTCTAA
AATAATTATATTTGCCATGAACAAGGGAAGAGTTACCGCACCCAGCCCTAGACTT
GGGCGCCCGAGCAAGGTTACGTCAAGCCACCTTCGCCCATCGCCCAACTCCGTAT
TCCCCGACAGCCGCACGTGGCCCTCGCCGGAATGAACCCTGAATCGGCATCACG
CCACGCGTTCGCCAATCGTTCCGCTCTCTGGCTTCATCGGCCTGCGCCTTCACGTC
GTGGTCACGACAGTGCATTCATACTTCCATTTGCACCTCGGCACACACTTTTACG
CATCGCCTACCCTTGCTGCGGCAGTCTAGGGTCACTTTGCAGCCATGGGACAGTG
CTACACCACCGTCGGTGCGCAAAGCTATTTCAAGTGAACCGTGGGCGGAAAAAA
GGAATGTACACTGTCTCAACCGACTCCTACAATTGTTTACCATGCAGATCAGAGC
TCGACGGCCATCATCGAGCAGGTGTGGGGCCTTGGTGGCGCGGCGCGGGGCCCC
AGGGCGTCGCAGGCATTGATGGCACTCTGAGACTTTCGCACGCGCATGAGGGAC
CCCATCAAGAGAAGAGTGTGTCTTTATGTCCCCATTCATGATGATGTATCTTGTG
ATTGTCGCAGTTTGGCAAGTTTAACCGGATCGCCGCTCCAGGTGTGGCGTGGCGG
ATTTTTCTAGGGGTGCTTGAGCAGTCG

SEQ ID NO: 94

Prototheca moriformis SulfRed Sulfite reductase promoter/5'UTR

GGCCCAGGGCCCTGCGGATGGCCCACACCAGATCTAGCCTCTCTTATGCCATGCC
CGCCTCGCTGCCCGTCGTATCCCCCCGCCGATCCGCGCGTAGGGGACCGCGGCCT
GACCCACGCCACGAAAGAGCTTTGCTCCTCAATTTCTCGCCAACAGAACCGTATC
AAACGCTCAACGCCTATCCCGAACAATCCGTATTCACACCAAATCGAGTATACCG
GACTGGTTTGCCTAGTCTTGAAGGAAATGATCCCGTCCATGCTCGGAAGGGGGA
GCGGGCGGAGGATCCTACTCATCTCTGAAATGGGATTGGTCCGAAGATGGGTTG
GGCAAGCACGTGCCAAACCCCAGCGAGTTGCTGACGAGCAGGCTCATCCAATCC
CCCGGCGAATCCTCCCTCACGCCCCGCATGCATACAAGTCCCTCCCACACGCCCC
CTCCCATCCATTTTCGCCTGGTCCGAACGCGAGCGGCGTCGAGGCGGACCACTTG
CTCCGCAGCGCCGTCTGGGTCTCCACCCCACAGCGGCTTTGCTGCCAGAGGCACC
CCCCTTGCCCCACCTCCTCTTGCAGCC

SEQ ID NO: 95

Prototheca moriformis SugT Sugar tranporter promoter/5"UTR

CCAGGCAGGCGGTAGGGTTGCCGATTGCTTGAGCGAATTGGAAGATATAATTTTT
TGTGGTGTCCCTGGACGCTGTTTGTGGCGCTCCTTTTTGGAGAAGATTGCGTGGG
GGAGCTTTCCATGTACCACGCTTCCTTCTGAAAGGATTCTGGCCGAGTCCTGATG
AGCCCAAAGAAAACACCTGCCTTTCAGTGCTGGCACTCTGAAAACGTCAACAGA
TGATTATACATGTCACAAAAGGCAGCCGATTAGGAACGGGAGCTCTGGCCGTTC
GTTTGGCTGCCTGGGCTGATTGAAGTGATCCACCCTGTTCGAATGAAGGCGGTCG
AGTCGAATTATCGACCGGAGCTGTCGGGAAGGCGTCCGGGGCAGAGTGAGGTGC
TGCGGCCTGGTTGTCGTTCAAAAAGACCCCGGTAGCCCAACAATCACGAACGAA
AGGAATATAATTGCTTGCATACTATACATTCAGTTTCTATGTGGCGGGTAGACAA
GTCTCATGGGCTTCTAAAGGCTGTCCCTTGAAGGCTACTTATAAAAACTTGCTGC
GCCATGGCACGGATCGCGCTTGCGCAGGCTGCAACCCTGCGCGCAAGGTCAAAT
ACACAGCAAAAGATACTAACAGAATTTCTAAAAACATTTAAATATTTGTTTCGAC
CAGCCAATTGTGGTCGTAGGCACGCAAAAGACTTTGTTTTGCGCCCACCGAGCAT
CCACGCTGGCAGTCAAGCCAGTCCGATGTGCATTGCGTGGCAGCATCGAGGAGC
ATCAAAAACCTCGTGCACGCTTTTCTGTCAATCATCATCAACCACTCCACCATGT
ATACCCGATGCATCGCGGTGCGCAGCGCGCCACGCGTCCCAGACCCGCCCAAAA
ACCCAGCAGCGGCGAAAGCAAATCTTCACTTGCCCGAAACCCCGAGCAGCGGCA
TTCACACGTGGGCGAAAACCCCACTTGCCCTAACAGGCGTATGTCTGCTGTCACG

ATGCCTGACAACGGTATTATAGATATACACTGATTAATGTTTGAGTGTGTGCGAG
TCGCGAATCAGGAATGAATTGCTAGTAGGCACTCCGACCGGGCGGGGGCCGAGG
GACCA

SEQ ID NO: 96
Prototheca moriformis Amt03-Ammonium transporter promoter/5'UTR
GGCCGACAGGACGCGCGTCAAAGGTGCTGGGCGTGTATGCCCTGGTCGGCAGGT
CGTTGCTGTTGCTGCGCTCGTGGTTCCGCAACCCTGATTTTGGCGTCTTATTCTGG
CGTGGCAAGCGCTGACGCCCGCGAGCCGGGCCGGCGGCGATGCGGTGTCTCACG
GCTGCCGAGCTCCAAGGGAGGCAAGAGCGCCCGGATCAGCTGAAGGGCTTTACA
CGCAAGGTACAGCCGCTCCTGCAAGGCTGCGTGGTGGACTTGAACCTGTAGGTCC
TCTGCTGAAGTTCCTCCACTACCTCACCAGGCCCAGCAGACCAAAGCACAGGCTT
TTCAGGTCCGTGTCATCCACTCTAAAACACTCGACTACGACCTACTGATGGCCCT
AGATTCTTCATCAACAATGCCTGAGACACTTGCTCAGAATTGAAACTCCCTGAAG
GGACCACCAGAGGCCCTGAGTTGTTCCTTCCCCCCGTGGCGAGCTGCCAGCCAGG
CTGTACCTGTGATCGAGGCTGGCGGGAAAATAGGCTTCGTGTGCTCAGGTCATGG
GAGGTGCAGGACAGCTCATGAAACGCCAACAATCGCACAATTCATGTCAAGCTA
ATCAGCTATTTCCTCTTCACGAGCTGTAATTGTCCCAAAATTCTGGTCTACCGGGG
GTGATCCTTCGTGTACGGGCCCTTCCCTCAACCCTAGGTATGCGCGCATGCGGTC
GCCGCGCAACTCGCGCGAGGGCCGAGGGTTTGGGACGGGCCGTCCCGAAATGCA
GTTGCACCCGGATGCGCGGCGCCTTTCTTGCGATAATTTATGCAATGGACTGCTC
TGCAAATTTCTGGGTCTGTCGCCAACCCTAGGATCAGCGGCGTAGGATTTCGTAA
TCATTCGTCCTGATGGGGAGCTACCGACTACCCTAATATCAGCCCGGCTGCCTGA
CGCCAGCGTCCACTTTTGCGTACACATTCCATTCGTGCCCAAGACATTTCATTGTG
GTGCGAAGCGTCCCCAGTTACGCTCACCTGTTTCCCGACCTCCTTACTGTTCTGTC
GACAGAGCGGGCCCACAGGCCGGTCGCAGCC

SEQ ID NO: 97
Prototheca moriformis Amt02-Ammonium transporter promoter/5'UTR
TCACCAGCGGACAAAGCACCGGTGTATCAGGTCCGTGTCATCCACTCTAAAGAG
CTCGACTACGACCTACTGATGGCCCTAGATTCTTCATCAAAAACGCCTGAGACAC
TTGCCCAGGATTGAAACTCCCTGAAGGGACCACCAGGGGCCCTGAGTTGTTCCTT
CCCCCCGTGGCGAGCTGCCAGCCAGGCTGTACCTGTGATCGGGGCTGGCGGGAA
AACAGGCTTCGTGTGCTCAGGTTATGGGAGGTGCAGGACAGCTCATTAAACGCC
AACAATCGCACAATTCATGGCAAGCTAATCAGTTATTTCCCATTAACGAGCTATA
ATTGTCCCAAAATTCTGGTCTACCGGGGGTGATCCTTCGTGTACGGGCCCTTCCCT
CAACCCTAGGTATGCGCACATGCGGTCGCCGCGCAACGCGCGCGAGGGCCGAGG
GTTTGGGACGGGCCGTCCCGAAATGCAGTTGCACCCGGATGCGTGGCACCTTTTT
TGCGATAATTTATGCAATGGACTGCTCTGCAAAATTCTGGCTCTGTCGCCAACCC
TAGGATCAGCGGTGTAGGATTTCGTAATCATTCGTCCTGATGGGGAGCTACCGAC
TGCCCTAGTATCAGCCCGACTGCCTGACGCCAGCGTCCACTTTTGTGCACACATT
CCATTCGTGCCCAAGACATTTCATTGTGGTGCGAAGCGTCCCCAGTTACGCTCAC
CTGATCCCCAACCTCCTTATTGTTCTGTCGACAGAGTGGGCCCAGAGGCCGGTCG
CAGCC

SEQ ID NO: 98
Protoheca moriformis Aat-01Amino Acid Transporter promoter/5'UTR
CGAAGGGGTCTGCATCGATTCGCGCGGTCTGGAGGCCAGCGTGACTGCTCGCGA
AAATGCTCTGCCGTGTCGGGCTCTGGCTGGGGCGGCCAGAGATCTCACCGTGCCA
CACGCAACTGCCGCACTCTGTGCCCGCCACCTGGCGCGCACATGCGACCTCTTCC

CCGTCATACCCTCTCCTCATGTGATCTTTCCACACGAGTGACGCAGGTGCGCGGA
GTGGAGGGAATCAGGACGTTTTCAAGGTACCTGCTCGAGCCGTACCAACAGCTG
CCGCCCGGCAAGGAAGAGATCGAGGCAGAGATTGCCCGGCTGGAGGCCCGGATA
ACGGAGCTCAAGAGCAAGCTGTCCGAGTGAGACCGCCCAGGTGCACGTGTCGAC
TCGCTATGACATGTACTCGACACAACATGAGGAATTCATCGAATTTGTAGGAAGC
GGGCATTGGTACGGGAGTGGGAAAGCGAAAAAACCTCCCTCCGGCAGTGCCATC
TGCCGGAGTCGAACGTTGATAGGGTTCTCGTGACAGGGTGTGACCTCTCAGCCTT
GCATCAATTAAACGCTATAGACATTATCAGTAACCGTGAATCCCGCATTGGATGC
CACCCGCGCGACCATTGGGGACCTGCATTACAGATCTAGGTGAGATGACAGCGA
GGCAACTTCGGCCCGCGGCCCAGCTTGCGGCGCACCAATATTGGTCACGGGAAG
CCACACACCGACCATAAATGAATACTTGTAAGCTATGTCAACCGATCAATGGCGT
CGAAAGTGTGCCACGAGGATCCATCTGGCGGGGCGGCGTGGCGCACAAGCGCAG
TCGCAATTTCTCGGACCCATCTGACCTAGGCCCAGCGCCGCGGGAGAAATCCCCG
GCGGGTCCTCCACGCAGTAACCCTAATGAGTATCGAGCGCCGACCATTTACACCA
TCGCCCCCGAAATCCTTCCGACATTATTATTATCTTTTAGATCTTGGAACAGACTC
TGCCAACC

SEQ ID NO: 99

Prototheca moriformis Aat02-Amino acid transporter promoter/5'UTR

AGAGAGCGGAGGTGGGGTTGTGAGGTGGGGTTGCTGACCAGGAGCTCGCGTCGC
CGAGCGCGACTCGCACACGGTCCAGTTACCCCCCCCTCCGCCCAAACGCAAGCCT
CCCATCTTGATGCCTTTCCGGCCACCTATACTATTTCTTAGTTCGCTGTAACATCC
AGACCGTCCTGAATAATAACAATGCCCTGTGTCAAGTGCATTCCTAAAAAAATTC
TGTCCCAACCAACAATCCCACCTGAAATACCACCAGCCCTGCCCAGTACACTCTT
CCAATACCATCTCCCTACCTCCACGCGCAAGCGACCCCCATGCGCGACCAGGCTC
GAAAGTGATTTATGACTTGAGACGAGCGAGTGGCGGCGCGGTCGACTGCCTTTTC
ATCACGTGCCGTACGTCGGCGACCGCTAGGGCTTTGCACGGCAACGCACGGCTTC
GCCAACCCGACCAGCCAGGACCTCGACTACTCTACCGCGAATTCGCCTCAAGAA
GTCGCCAAATGTGCCATACACCATTCCTTACAGCACTGTTCAAACTTGATGCCAA
TTTTGACATTCGGGTTGCTCGTTGGCTGCGCCCACATCGGCCGTGAGTGCAGCAG
GCGGGATCGGACACGGAGGACGCGGCGTCACGCCCCGAACGCAGCCCGTAACTC
TACATCAACACGACGTGTTGCGTAATCCCGCCCGGCTGCGCATCGTGCCAACCCA
TTCGCGATGGATGGTCGGAAAATGGTGTGCCAACTGCCCTGAGGGAGGCTCTCG
CGAAACGGGCACGTCCCTGAAACCGAAACTGTGGCCTTGTCGTCGGCCACGCAA
GCACGTGGACCCTAAACACCAAGAAAATCAGTAAACAAGGTTGACATCCTCTAC
GGGCGAATTGTTTGCCCAACCCTTCATCGCACACTGCCATTATAATGCATCTAGC
TCGGCGACAAGTTTAGAAAAGGCAGGCTGCATTGTTCCATTTCGCCGTGGCGGCG
TGGGTGCCCATTTTACGAGGTTTGGGCTCCCGGGCAGCGACCGAGCCAGGTCGA
GTCCCTCTCGCCCGTCGACAATGTTGCGAACCCCACAAGCGGCTAACAACAACTT
GATGGTACCTGTACACTGCCAATTCCTTCTTCCCCGGCCGAGGTTTACACGTGAT
GGCCATGGCTTCGCATTCAGGCCGACTTCCCATTCCGACTTTCCAGAGGGTCCGC
GGACGCTGGGGGTTGGCTGCCTGAGGCCCACCCTTTGTTCCCCGCGTCCCGACAA
ACACAATTGCGTTACATAAGGGGGAGCCGCCCCCGTTCAGAGTGCAGAAATCTTT
CACTATATTTTCCAGTCGTCAGCGAAATCAAGT

SEQ ID NO: 100

Prototheca moriformis Aat03-Amino acid transporter promoter/5"UTR

GATGGTGGGGTGTCTGCCTTGGGCTGGGTGATGGAGGCTGGTGGTGCGCGGGTTT
CCTGATGCATTCTATCTACGCAGTGTCATGGTGTCCATTCCACACACCAGTACAC
CCTTACACTAAGGATCCATCCCTCCTTCCCTCTTCAGGACTACATGGACCCCACG

AGCTACCGACCGGGCTTTCTCAAAAACGTCAAGGTCATGTTTGACATGCGGGACG
TGGTGGACGACGTGCAAGGTGCGTCCGGAGTGCGCGCAAATGAGCAAGTCGGGC
AATGTGTCGGGGTGGGCACCGGGGCTGGAGATCCGCGATCCCCGAGAAAACGCC
GTACCACCCCCCCGCGCTATTCCCTCGATTGCGCGCAGATGTGGTGACCGACACGG
GGGACAACCTGGCGGACATGGGGCGCCGGACCTGGAAGCACGCCAAGTCGCACA
CGGGGAGGCTCGTGCAGTCCCCCCCCATCGTACCTCAAGGGTCTCTTTGGTCGCGA
TCCAAAGTACGCTGGTGGCATGGCATGCCCGAAATGAACATCATGTGTGATCTCC
GATTGCCAATGGCCACCTCCACGGACCACCTTGCAGGCGGAAGCGCAATCCAGG
GCCCGAGCCTGACGAGGACGGAGACTCCTCGTCCAGCGCGGGGTCCCCGACCCG
ACGCAGCAGCCGACCCCTGCTAACCCGGCAACGATCGGACCAGCAACCTTGCTG
TAGTTCCGATCCGTGATGACGGGCATTGCCGCCGCTCGATCCGCTTTGATGACTG
TCTATTATTTGCGCGGAGCCCCCTCGGAACCCTACCCCGCTCTTGCAAGCCCCTTG
CATCGGAGATCCTCGTGCGCCCGCCATGACCCCACTGGATTGCCCAACATCCTTC
TTTATCGTGTAAAATGTGATTCCTCGGCTGCAATCGACTGGCCTTCGCTTCTGGCC
CCAAGAGGGCTCGAACGTGCGGCAGCGAGGGCGCTGACACACCCAAGCCCTAGG
GCTTTCAACGTCGGCTGCCAGGCCGGATAGGGGGATCGCCTCCTTTCCACCACCC
ACCTACGAGGGATTCGAGTCGGCTTCCAGCTCAGCTATTCGGCCGCGCCCCCGGC
CCTGCAGACGTCCTCCAGTTTCCGAACAGGTCGCTCTCAGAACACCTGCCGCGGC
TGCGATACGGCAGGCTCTCAAAGCGTCGAC

SEQ ID NO: 101
Prototheca moriformis Aat04-Amino acid transporter promoter/5'UTR
CGCGTGGAGCGGTGCGTGCGGATGCCGCGCGCCTGCCAAGGCCTTTTGTATGCCT
GGCCTGGGAAGTTTCCTGACTGAAGCATCTTCAAGATGCTCTCTCACGACCAGCG
ACACCAACACCGTCACTTTTTGCCCCTCCTGCCGCAGGTGCCACTTTCTACTTTGA
CGTCTTCTCCAGGCGGTACATTGCGGGACTGAGCGCCAATTCGGCCAAGAACAG
CGCTGTCGACTTGAGGAGGCAGGGGTCCGTCGACTCTGCCGAGTGACACGCCTTC
GACCCGACTGTACTACGGCCTGCTGAAGAGTGGGTCTCGCCGGCCGGCGTGACC
GGCCCTGTGCCCACAATCGACCATCTATTCGCTCCTTGTCATCTGGCGCCGTCAAT
TGCCCGCGACTTGACGGCAACTGGCTCGATCGAGTCGTATTGAAAAAGCACGTTT
TGTCCTACAGGGCCGCGGTCCGTTACCAACGTGGTTCTCGTTAGGTTTTCGTCGG
GCGGTGGTGCGCGAACTGTCCGATGCCATCCCGGCAAACCCCAGCAAGGTCGCC
AGTCTGGTTCTGACGCAATAGAGTGCGTTTTGGGCCAGTCTAAAAATTCGTCTGG
CATGACGTGGCTCCACATCGTACCCGGAGCCTGCCTTGGTAATGTGAGGCACCGG
TGCCAACTCCATTATGGCAGGCATCGAGCGCGCAGGTGAGTACATGACCTTCCGT
GAATTGGGAAGGCGAGCTTGTGTAACGCCTGCGATCGTGCCAGTGAGGCATCGT
AAACTCAAAATATTTTGTAGAAAGTGTCTGATGCCTGGTGAGGCTGCGTAGGGCA
AGGGCAAGCCCTTGGCAGATGGGTAATGGGTCCGGACCTCACAACAGCAACCCC
GCGTCCCCCTTAGGGCCCCTGAGGCTCGATGGCAGGGCCAGCGAGCCCGCGGCC
AAAGGGCGCCATCCCACGGTCGCCCAACGACTCCACGGGTCCTATACCTCATCTT
GAATGGCACTAAAAACTATAGAATATCGGGCACTGGTGGGCGTCTGGGGTACAG
CTGGCCGAGCGCAGTGGCAAACCCTAGGTCCCGCCTCAAGGGCGATTCCCGGGT
CAATGACACGCAAGCAAGATCACATGGCGCGGTCCGCCTCGCGGCTCCACACCC
AGGCCCTAGTTTCGCAACCCATAAATATCGCCCCGATACCATCATAAGCCAGCAA
ATAATTTTTTATCAGAGTTCCAAACCTCCTCAGCTGTGGGAAACCAGCCCACTCT
GAACG

SEQ ID NO: 102
Prototheca moriformis Aat05-Amino acid transporter promoter/5'UTR

CCGAGCAGTTCATGGCCAAGTACAAGGACTAGAGACCGGAGGTCGGTAGGCTGA
ATGGAGCTGGCGTCGTCGTGCGCGACGTGCACGCGATGCGATACTACGACCCCA
CAAACGCATGCCTCCCATCTTGATGCCTTTCCGGCCATTTATACTATTTCTCATTT
CGCTGTAACATCTTGAATAATAGAATTGCCCTGTGTCAAGTGGATTCCAAGAAAT
ATTCTGTCCCAACAAAACAACCCAACCTGAAAACAACCTCAAATACCACCAGCC
CTGCCCACCTGCCCAGTACACTTTTCCAATACCATCTCCCTACCTTCACGCGCAAG
CGGCACCCATGCGCGACCAGGCTCGAAAGGATTTCACGACTCAGGACGAGCGAG
TGGCGGCGCGACCGCCTGCCTGTTCGTCACGTGCCGTACGTCGGCGACCGCTAGA
GCTTTGCCTGGCAACCCCCGGCTTCGTCAACCCGGCCAGCCAGGATCTCGACCAC
TCTACCGCGAAATCGCCTCAAGAAGTCGCCAAAAGTGCCGTACACCATGCTTCGC
AGCGCTGTTCAAACTTGATGCCAATCTTGACAATCAGGTTGCTCGTTGGCTGCGT
CCACATCGGCCGTGATTGCAGCAGGCGGGGATCGGACACGGAGGACGCGGCGTC
ACGCCGCGAACGCAGCCCGTAACTCTACATCAACGCGATATGTTGCGTAATCCCG
CCCGGCTGCGCATTGTGACAACCCATTCGCGATGGATGGTCGGAAAATGGTGTGC
CAACTGCCCTGAGGGACTCTCTCGCGAAACGGGCACGTCCCTGTATCCGAAACTG
TGGCATGGCCTTGTCGACCACGCAAGCACGTGGACCCTAACACCACGAAAATAA
GTAAAAAAGGTTGACATCCTCTACGAGCGAATTGTTTGCTCGACCCTTCATCGCA
CACTGTCATTATAATGCATCTAGCTCGGCGACAAGTTTAAAAAAGGCAGGCTGCA
TTATTCCATTTTGCCGTGGCGGCATGGGTGCCCATTTTATGAGGTTTGGGCTCTTG
GGCAGCGACCGAGCCAGGTTGAGTCCCTCTCGCCCGTCGACAACGTTCCAAAGC
CCATAAGTGGCTAATAAACAACTTGATGGTACCTGTACACTGCCAGTTCCTTCTT
CCCCGGCCGAGGTTTACACGTGATGGCCATGGCTTCGCGTTTCAGGCTGACTTCC
CATTCCGACTTTCCAGAGGGTCCGCGGACGCCGGGGGTTGGCTGCGTGAGGCCC
ACCCCTTGTTCCCCGCGTCCCGACAAACACAATTGCGTTACATAAGGGGGAAGCC
GCCCCCCCGTTCAGAGTGCAAACATCTTTCATTATATTTTTCAGTCGTCAGCGAAAT
CAAGTATGTCGCTGACAGGCATGAAGGCC

SEQ ID NO: 103
KE858 Homologous recombination construct SZ725

GCCCTTTGTCATCGTTGGCATGCTTTTTGCGTATGTACCATATGTTGAATGTATAA
TACGAACGGTTGACCGTCTGAGATGCGAGCTTTGGGTCTTGTCAAATGCGTGGCC
GCACGGCTCCCTCGCACCCAGCCCCGAGGCGTCGCGCACCTGGCGAGGAGCAGA
CCCACGCCAAGAAAGTCTAGTCCAGCATGTAACAACATCAGGCAATGTGACGTT
TTCGGTTCCCGATTTCTCTGCCGCTCTTTGACGGCAGGCACGGGCGAGCAACCGG
CGGCGCTCGCGTCAGGCACGATGGATGCGGCGCTGCCCACCTGTCAATGTACCCC
ACCAGTCTGTCGATCGCTACAAGCAACCTTGTGCTCCACATTCCCACTTGCAGAC
AGTCTAGTCGATTTTGCCAAGCTGGATGTGAGGATTGGCCATATCTTGGAGGCCA
AGATTCACCCGGATGCTGATGGGTACGTACGCGAGCCAGGCAGGCAGCTGCGTT
GACTTTCTGATTGGCACAAAGCTTTGGCTACTCTCAATACCAACCACGTGCCCCT
TCTGCACACCTGCTTCCTTCTGATGACCACTCGCCACGCATGTCGCAGTCTGTACG
TCGAGCAGATCGACCTCGGCGAGGAGGGGGGCCCTCGCACCATCGTGAGTGGCC
TGGTCCGGCACGTGACCCTGGAGGACCTTGTCGGCCGGCGGGTGGTGGTGCTGG
CCAACCTCAAGCCTCGGAGCATGCGCGGGGTCAAATCGGCTGGGATGCTGCTCT
GCGCCGCCAACGCGGATCACACCGCGGTGGAGCCGCTGCGGGTCCCGGACGCCG
CCGTGACGGGGGAGCGGGTCTGGGCGGGGGACGAGGCACTCCTGTCCACGGAGC
CTGCCACACCCAATCAGGTAAGGACACGTTATTGGTGCGCATGGTGCGAATGCGT
GGTCTGACCTGCTGTGGGTATGTGTTGTGGGATTGGAAACCGAATGAGGGCCGTT
CAGGATTGAGCCCTTGGCCCCACCCTGCTCATCCTCTCACGCCCGCAGGTCCAGA
AGAAGAAAATCTGGGAGGCAGTACAGCCGCTGCTGAGAGTGAACGCCCAGGGG

ATCGCTACTGTGGCAGGAGAGGCTATGGTGACCAGTGCGGGGCCACTGACCGCG
CCCACGCTGGTTGACGCCGCGATTTCCTGACGCGAGCGACTGATTCTTGACCTTT
GAGAAGCCACCACAGCACCATTTTCATTGTTCATCCTTGATTTCAGTACGACTTCT
CACCATTTCAGTACTGTAGGACCCCCAAAATAGTGTGATCACGCTCGCAAGGCAC
CTGTGTGATCACGGGGAAGGGCGAATTCCTTTCTTGCGCTATGACACTTCCAGCA
AAAGGTAGGGCGGGCTGCGAGACGGCTTCCCGGCGCTGCATGCAACACCGATGA
TGCTTCGACCCCCCGAAGCTCCTTCGGGGCTGCATGGGCGCTCCGATGCCGCTCC
AGGGCGAGCGCTGTTTAAATAGCCAGGCCCCCGATTGCAAAGACATTATAGCGA
GCTACCAAAGCCATATTCAAACACCTAGATCACTACCACTTCTACACAGGCCACT
CGAGCTTGTGATCGCACTCCGCTAAGGGGGCGCCTCTTCCTCTTCGTTTCAGTCAC
AACCCGCAAACGGCGCGCCATGCTGCTGCAGGCCTTCCTGTTCCTGCTGGCCGGC
TTCGCCGCCAAGATCAGCGCCTCCATGACGAACGAGACGTCCGACCGCCCCCTG
GTGCACTTCACCCCCAACAAGGGCTGGATGAACGACCCCAACGGCCTGTGGTAC
GACGAGAAGGACGCCAAGTGGCACCTGTACTTCCAGTACAACCCGAACGACACC
GTCTGGGGGACGCCCTTGTTCTGGGGCCACGCCACGTCCGACGACCTGACCAACT
GGGAGGACCAGCCCATCGCCATCGCCCCGAAGCGCAACGACTCCGGCGCCTTCT
CCGGCTCCATGGTGGTGGACTACAACAACACCTCCGGCTTCTTCAACGACACCAT
CGACCCGCGCCAGCGCTGCGTGGCCATCTGGACCTACAACACCCCGGAGTCCGA
GGAGCAGTACATCTCCTACAGCCTGGACGGCGGCTACACCTTCACCGAGTACCA
GAAGAACCCCGTGCTGGCCGCCAACTCCACCCAGTTCCGCGACCCGAAGGTCTTC
TGGTACGAGCCCTCCCAGAAGTGGATCATGACCGCGGCCAAGTCCCAGGACTAC
AAGATCGAGATCTACTCCTCCGACGACCTGAAGTCCTGGAAGCTGGAGTCCGCGT
TCGCCAACGAGGGCTTCCTCGGCTACCAGTACGAGTGCCCCGGCCTGATCGAGGT
CCCCACCGAGCAGGACCCCAGCAAGTCCTACTGGGTGATGTTCATCTCCATCAAC
CCCGGCGCCCCGGCCGGCGGCTCCTTCAACCAGTACTTCGTCGGCAGCTTCAACG
GCACCCACTTCGAGGCCTTCGACAACCAGTCCCGCGTGGTGGACTTCGGCAAGG
ACTACTACGCCCTGCAGACCTTCTTCAACACCGACCCGACCTACGGGAGCGCCCT
GGGCATCGCGTGGGCCTCCAACTGGGAGTACTCCGCCTTCGTGCCCACCAACCCC
TGGCGCTCCTCCATGTCCCTCGTGCGCAAGTTCTCCCTCAACACCGAGTACCAGG
CCAACCCGGAGACGGAGCTGATCAACCTGAAGGCCGAGCCGATCCTGAACATCA
GCAACGCCGGCCCCTGGAGCCGGTTCGCCACCAACACCACGTTGACGAAGGCCA
ACAGCTACAACGTCGACCTGTCCAACAGCACCGGCACCCTGGAGTTCGAGCTGG
TGTACGCCGTCAACACCACCCAGACGATCTCCAAGTCCGTGTTCGCGGACCTCTC
CCTCTGGTTCAAGGGCCTGGAGGACCCCGAGGAGTACCTCCGCATGGGCTTCGA
GGTGTCCGCGTCCTCCTTCTTCCTGGACCGCGGGAACAGCAAGGTGAAGTTCGTG
AAGGAGAACCCCTACTTCACCAACCGCATGAGCGTGAACAACCAGCCCTTCAAG
AGCGAGAACGACCTGTCCTACTACAAGGTGTACGGCTTGCTGGACCAGAACATC
CTGGAGCTGTACTTCAACGACGGCGACGTCGTGTCCACCAACACCTACTTCATGA
CCACCGGGAACGCCCTGGGCTCCGTGAACATGACGACGGGGGTGGACAACCTGT
TCTACATCGACAAGTTCCAGGTGCGCGAGGTCAAGTGATTAATTAACTCGAGGCA
GCAGCAGCTCGGATAGTATCGACACACTCTGGACGCTGGTCGTGTGATGGACTGT
TGCCGCCACACTTGCTGCCTTGACCTGTGAATATCCCTGCCGCTTTTATCAAACAG
CCTCAGTGTGTTTGATCTTGTGTGTACGCGCTTTTGCGAGTTGCTAGCTGCTTGTG
CTATTTGCGAATACCACCCCCAGCATCCCCTTCCCTCGTTTCATATCGCTTGCATC
CCAACCGCAACTTATCTACGCTGTCCTGCTATCCCTCAGCGCTGCTCCTGCTCCTG
CTCACTGCCCCTCGCACAGCCTTGGTTTGGGCTCCGCCTGTATTCTCCTGGTACTG
CAACCTGTAAACCAGCACTGCAATGCTGATGCACGGGAAGTAGTGGGATGGGAA
CACAAATGGAAAGCTT

SEQ ID NO: 104

Homologous recombination: Targeting cassette for disruption of *Prototheca moriformis* stearoyl ACP desaturase coding region (β-tubulin driven *suc2* cassette)

TTTGGCCCCGCTTTCCAGCTCCGGATCTGCTGGCGTCCGCCGCGAGACGTGACAT
CGCACGTCGCCGGGAGCGCCAGCTTGATCACTTGGCAGGGGGCCGTGCTCTACA
AATACCAGGCCCCGCGGCGGTCAGTTCGCACATCCAATACCTGCCGAGCCATCTT
GCCTACACTTTTTATCGACTCCTCTACTCTGTTCGCGAGAGCGCTCGGTCCAGGCT
TGGAATTCGCCGAATTCAGCTCGATCAGTCGCTTCTGCAACTGATCTCGGCCGTT
CGCAGACTGCCTTTTCTCAGCTTGTCAGGTAGCGAGTTGTTGTTTTATATTTATTC
GATTTCATCTGTGTTGCATGTCTTGTTCGTGCTGTGCGTTCTTTCTGGGCCGCGCT
GTCGGGTCGCATGGGCTAGCTGTACTCATGTTAGTCATGCCGGTCCGACCTTGTT
CGAGGAAGGCCCCACACTGAGCGTGCCCTCTTTCTACACCCCTTGTGCAGAAATT
AGATAGAAAGCAGAATTCCTTTCTTGCGCTATGACACTTCCAGCAAAAGGTAGG
GCGGGCTGCGAGACGGCTTCCCGGCGCTGCATGCAACACCGATGATGCTTCGAC
CCCCCGAAGCTCCTTCGGGGCTGCATGGGCGCTCCGATGCCGCTCCAGGGCGAGC
GCTGTTTAAATAGCCAGGCCCCCGATTGCAAAGACATTATAGCGAGCTACCAAA
GCCATATTCAAACACCTAGATCACTACCACTTCTACACAGGCCACTCGAGCTTGT
GATCGCACTCCGCTAAGGGGGCGCCTCTTCCTCTTCGTTTCAGTCACAACCCGCA
AACGGCGCGCCATGCTGCTGCAGGCCTTCCTGTTCCTGCTGGCCGGCTTCGCCGC
CAAGATCAGCGCCTCCATGACGAACGAGACGTCCGACCGCCCCCTGGTGCACTTC
ACCCCCAACAAGGGCTGGATGAACGACCCCAACGGCCTGTGGTACGACGAGAAG
GACGCCAAGTGGCACCTGTACTTCCAGTACAACCCGAACGACACCGTCTGGGGG
ACGCCCTTGTTCTGGGGCCACGCCACGTCCGACGACCTGACCAACTGGGAGGAC
CAGCCCATCGCCATCGCCCCGAAGCGCAACGACTCCGGCGCCTTCTCCGGCTCCA
TGGTGGTGGACTACAACAACACCTCCGGCTTCTTCAACGACACCATCGACCCGCG
CCAGCGCTGCGTGGCCATCTGGACCTACAACACCCCGGAGTCCGAGGAGCAGTA
CATCTCCTACAGCCTGGACGGCGGCTACACCTTCACCGAGTACCAGAAGAACCCC
GTGCTGGCCGCCAACTCCACCCAGTTCCGCGACCCGAAGGTCTTCTGGTACGAGC
CCTCCCAGAAGTGGATCATGACCGCGGCCAAGTCCCAGGACTACAAGATCGAGA
TCTACTCCTCCGACGACCTGAAGTCCTGGAAGCTGGAGTCCGCGTTCGCCAACGA
GGGCTTCCTCGGCTACCAGTACGAGTGCCCCGGCCTGATCGAGGTCCCCACCGAG
CAGGACCCCAGCAAGTCCTACTGGGTGATGTTCATCTCCATCAACCCCGGCGCCC
CGGCCGGCGGCTCCTTCAACCAGTACTTCGTCGGCAGCTTCAACGGCACCCACTT
CGAGGCCTTCGACAACCAGTCCCGCGTGGTGGACTTCGGCAAGGACTACTACGC
CCTGCAGACCTTCTTCAACACCGACCCGACCTACGGGAGCGCCCTGGGCATCGCG
TGGGCCTCCAACTGGGAGTACTCCGCCTTCGTGCCCACCAACCCCTGGCGCTCCT
CCATGTCCCTCGTGCGCAAGTTCTCCCTCAACACCGAGTACCAGGCCAACCCGGA
GACGGAGCTGATCAACCTGAAGGCCGAGCCGATCCTGAACATCAGCAACGCCGG
CCCCTGGAGCCGGTTCGCCACCAACACCACGTTGACGAAGGCCAACAGCTACAA
CGTCGACCTGTCCAACAGCACCGGCACCCTGGAGTTCGAGCTGGTGTACGCCGTC
AACACCACCCAGACGATCTCCAAGTCCGTGTTCGCGGACCTCTCCCTCTGGTTCA
AGGGCCTGGAGGACCCCGAGGAGTACCTCCGCATGGGCTTCGAGGTGTCCGCGT
CCTCCTTCTTCCTGGACCGCGGGAACAGCAAGGTGAAGTTCGTGAAGGAGAACC
CCTACTTCACCAACCGCATGAGCGTGAACAACCAGCCCTTCAAGAGCGAGAACG
ACCTGTCCTACTACAAGGTGTACGGCTTGCTGGACCAGAACATCCTGGAGCTGTA
CTTCAACGACGGCGACGTCGTGTCCACCAACACCTACTTCATGACCACCGGGAAC
GCCCTGGGCTCCGTGAACATGACGACGGGGGTGGACAACCTGTTCTACATCGAC
AAGTTCCAGGTGCGCGAGGTCAAGTGATTAATTAACTCGAGGCAGCAGCAGCTC
GGATAGTATCGACACACTCTGGACGCTGGTCGTGTGATGGACTGTTGCCGCCACA
CTTGCTGCCTTGACCTGTGAATATCCCTGCCGCTTTTATCAAACAGCCTCAGTGTG

TTTGATCTTGTGTGTACGCGCTTTTGCGAGTTGCTAGCTGCTTGTGCTATTTGCGA
ATACCACCCCCAGCATCCCCTTCCCTCGTTTCATATCGCTTGCATCCCAACCGCAA
CTTATCTACGCTGTCCTGCTATCCCTCAGCGCTGCTCCTGCTCCTGCTCACTGCCC
CTCGCACAGCCTTGGTTTGGGCTCCGCCTGTATTCTCCTGGTACTGCAACCTGTAA
ACCAGCACTGCAATGCTGATGCACGGGAAGTAGTGGGATGGGAACACAAATGGA
CCGACACGCCCCCGGCCCAGGTCCAGTTCTCCTGGGTCTTCCAGAGGCCCGTCGC
CATGTAAAGTGGCAGAGATTGGCGCCTGATTCGATTTGGATCCAAGGATCTCCAA
TCGGTGATGGGGACTGAGTGCCCAACTACCACCCTTGCACTATCGTCCTCGCACT
ATTTATTCCCACCTTCTGCTCGCCCTGCCGGGCGATTGCGGGCGTTTCTGCCCTTG
ACGTATCAATTTCGCCCCTGCTGGCGCGAGGATTCTTCATTCTAATAAGAACTCA
CTCCCGCCAGCTCTGTACTTTTCCTGCGGGGCCCCTGCATGGCTTGTTCCCAATGC
TTGCTCGATCGACGGCGCCCATTGCCCACGGCGCTGCCGCATCCATGTGAAGAAA
CACGGAAGAGTGCGAAGACTGGAAGTGAATTAAGAGTATAAGAAGAGGTACCA
AGGGATTCTCAGGTGCTCTTAGGAACGGCTTTTCCTTCGCCAAGAGAAACTGCTA
CTGCTCGTGTCGCCACGGTGGTCAAGCCGCCCCATCTGCGATCCACCAGGCCCAT
CCGCGGACTCGCGATCAGCCTGCTGGATCCGGACTGCCGACCTGACCGCTCGCAT
CCACCATTACAACCCTCCAATTGGACACCACTCCCACGTCCTAAAGTTCACCATG
CAAGCTGATCGATCGCATTCGCCGATGCACTCGCCTGCCACAGAGGTGTGCGCTT
CGGACTAGCGTGCAGGCGCCCCGAGGCCACCAGCATGCACCGATGGAAGCGGGC
ACGGCCGCTGCTCCAGGTCGCTGGCTCGCTCAGACCCATAGCAACCTCCGCTGCG
TCCCTAAATGTCACACAGAGCGTCTTTGATGGGTACGGATGGGAGAGAATCTGAT
TGGGCATTGCTGGTGCAGTGCAGGAAGATGGCAAGTGCACAGTCAGTCATGCTG
TACAAACTGGTGCCTCGTAGTATTGACTCGTATAGTGCATAGTATCATGCATGGT
CGTTACTTGCAA

SEQ ID NO: 105
Homologous recombination: Targeting cassette for disruption of *Prototheca moriformis* stearoyl ACP desaturase coding region (*suc2* cassette alone)

TTTGGCCCCGCTTTCCAGCTCCGGATCTGCTGGCGTCCGCCGCGAGACGTGA
CATCGCACGTCGCCGGGAGCGCCAGCTTGATCACTTGGCAGGGGGCCGTGCT
CTACAAATACCAGGCCCCGCGGCGGTCAGTTCGCACATCCAATACCTGCCGA
GCCATCTTGCCTACACTTTTTATCGACTCCTCTACTCTGTTCGCGAGAGCGCTC
GGTCCAGGCTTGGAATTCGCCGAATTCAGCTCGATCAGTCGCTTCTGCAACT
GATCTCGGCCGTTCGCAGACTGCCTTTTCTCAGCTTGTCAGGTAGCGAGTTGT
TGTTTTATATTTATTCGATTTCATCTGTGTTGCATGTCTTGTTCGTGCTGTGCGT
TCTTTCTGGGCCGCGCTGTCGGGTCGCATGGGCTAGCTGTACTCATGTTAGTC
ATGCCGGTCCGACCTTGTTCGAGGAAGGCCCCACACTGAGCGTGCCCTCTTT
CTACACCCCTTGTGCAGAAATTAGATAGAAAGCAATGCTGCTGCAGGCCTTC
CTGTTCCTGCTGGCCGGCTTCGCCGCCAAGATCAGCGCCTCCATGACGAACG
AGACGTCCGACCGCCCCCTGGTGCACTTCACCCCCAACAAGGGCTGGATGA
ACGACCCCAACGGCCTGTGGTACGACGAGAAGGACGCCAAGTGGCACCTGT
ACTTCCAGTACAACCCGAACGACACCGTCTGGGGGACGCCCTTGTTCTGGGG
CCACGCCACGTCCGACGACCTGACCAACTGGGAGGACCAGCCCATCGCCAT
CGCCCCGAAGCGCAACGACTCCGGCGCCTTCTCCGGCTCCATGGTGGTGGAC
TACAACAACACCTCCGGCTTCTTCAACGACACCATCGACCCGCGCCAGCGCT
GCGTGGCCATCTGGACCTACAACACCCCGGAGTCCGAGGAGCAGTACATCT

CCTACAGCCTGGACGGCGGCTACACCTTCACCGAGTACCAGAAGAACCCCG
TGCTGGCCGCCAACTCCACCCAGTTCCGCGACCCGAAGGTCTTCTGGTACGA
GCCCTCCCAGAAGTGGATCATGACCGCGGCCAAGTCCCAGGACTACAAGAT
CGAGATCTACTCCTCCGACGACCTGAAGTCCTGGAAGCTGGAGTCCGCGTTC
GCCAACGAGGGCTTCCTCGGCTACCAGTACGAGTGCCCCGGCCTGATCGAG
GTCCCCACCGAGCAGGACCCCAGCAAGTCCTACTGGGTGATGTTCATCTCCA
TCAACCCCGGCGCCCCGGCCGGCGGCTCCTTCAACCAGTACTTCGTCGGCAG
CTTCAACGGCACCCACTTCGAGGCCTTCGACAACCAGTCCCGCGTGGTGGAC
TTCGGCAAGGACTACTACGCCCTGCAGACCTTCTTCAACACCGACCCGACCT
ACGGGAGCGCCCTGGGCATCGCGTGGGCCTCCAACTGGGAGTACTCCGCCTT
CGTGCCCACCAACCCCTGGCGCTCCTCCATGTCCCTCGTGCGCAAGTTCTCCC
TCAACACCGAGTACCAGGCCAACCCGGAGACGGAGCTGATCAACCTGAAGG
CCGAGCCGATCCTGAACATCAGCAACGCCGGCCCCTGGAGCCGGTTCGCCA
CCAACACCACGTTGACGAAGGCCAACAGCTACAACGTCGACCTGTCCAACA
GCACCGGCACCCTGGAGTTCGAGCTGGTGTACGCCGTCAACACCACCCAGA
CGATCTCCAAGTCCGTGTTCGCGGACCTCTCCCTCTGGTTCAAGGGCCTGGAG
GACCCCGAGGAGTACCTCCGCATGGGCTTCGAGGTGTCCGCGTCCTCCTTCTT
CCTGGACCGCGGGAACAGCAAGGTGAAGTTCGTGAAGGAGAACCCCTACTT
CACCAACCGCATGAGCGTGAACAACCAGCCCTTCAAGAGCGAGAACGACCT
GTCCTACTACAAGGTGTACGGCTTGCTGGACCAGAACATCCTGGAGCTGTAC
TTCAACGACGGCGACGTCGTGTCCACCAACACCTACTTCATGACCACCGGGA
ACGCCCTGGGCTCCGTGAACATGACGACGGGGGTGGACAACCTGTTCTACAT
CGACAAGTTCCAGGTGCGCGAGGTCAAGTGACCGACACGCCCCCGGCCCAG
GTCCAGTTCTCCTGGGTCTTCCAGAGGCCCGTCGCCATGTAAAGTGGCAGAG
ATTGGCGCCTGATTCGATTTGGATCCAAGGATCTCCAATCGGTGATGGGGAC
TGAGTGCCCAACTACCACCCTTGCACTATCGTCCTCGCACTATTTATTCCCAC
CTTCTGCTCGCCCTGCCGGGCGATTGCGGGCGTTTCTGCCCTTGACGTATCAA
TTTCGCCCCTGCTGGCGCGAGGATTCTTCATTCTAATAAGAACTCACTCCCGC
CAGCTCTGTACTTTTCCTGCGGGGCCCCTGCATGGCTTGTTCCCAATGCTTGCT
CGATCGACGGCGCCCATTGCCCACGGCGCTGCCGCATCCATGTGAAGAAAC
ACGGAAGAGTGCGAAGACTGGAAGTGAATTAAGAGTATAAGAAGAGGTAC
CAAGGGATTCTCAGGTGCTCTTAGGAACGGCTTTTCCTTCGCCAAGAGAAAC
TGCTACTGCTCGTGTCGCCACGGTGGTCAAGCCGCCCCATCTGCGATCCACC
AGGCCCATCCGCGGACTCGCGATCAGCCTGCTGGATCCGGACTGCCGACCTG
ACCGCTCGCATCCACCATTACAACCCTCCAATTGGACACCACTCCCACGTCC
TAAAGTTCACCATGCAAGCTGATCGATCGCATTCGCCGATGCACTCGCCTGC
CACAGAGGTGTGCGCTTCGGACTAGCGTGCAGGCGCCCCGAGGCCACCAGC
ATGCACCGATGGAAGCGGGCACGGCCGCTGCTCCAGGTCGCTGGCTCGCTC
AGACCCATAGCAACCTCCGCTGCGTCCCTAAATGTCACACAGAGCGTCTTTG
ATGGGTACGGATGGGAGAGAATCTGATTGGGCATTGCTGGTGCAGTGCAGG
AAGATGGCAAGTGCACAGTCAGTCATGCTGTACAAACTGGTGCCTCGTAGTA
TTGACTCGTATAGTGCATAGTATCATGCATGGTCGTTACTTGCAA

SEQ ID NO: 106
UTEX 21, 22, 28, 257

TGTTGAAGAATGAGCCGGCGACTTATAGGAAGTGGCGTGGTTAAGGAATTTTCC
GAAGCCCAAGCGAAAGCAAGTTTTAAAAATAGCGATATTTGTCACTTTTTATGGA
CCCGAACCCGGGTGATCTAACCGTGACCAGGATGAAGCTTGGGTAACACCAAGT
GAAGGTCCGAACTCTTCGATCTTTAAAAATCGTGAGATGAGTTGCGGTTAGTAGG
TGAAATGCCAATCGAACTCGGAGCTAGCTGGTTCTCCCCGAAATGTGTTGAGGCG
CAGCGATGAATGACAAAACAAATAGTACGGTGTAGGGGTAAAGCACTGTTTCGG
TGCGGGCTGCGAAAGCGGTACCAAATCGTGGCAAACTCAGAATACTACGCTTGT
ATACCATTCATCAGTGAGACTGTGGGGGATAAGCTCCATAGTCAAGAGGGAAAC
AGCCCAGATCACCAGTTAAGGCCCCAAAATGACAGCTAAGTGGCAAAGGAGGTG
AAAGTGCAGAAACAACCAGGAGGTTTGCCCAGAAGCAGCCATCCTTTAAAGAGT
GCGTAATAGCTCACTG

SEQ ID NO: 107
Cuphea palustris thioesterase expression cassette

GAATTCCTTTCTTGCGCTATGACACTTCCAGCAAAAGGTAGGGCGGGCTGCGAGA
CGGCTTCCCGGCGCTGCATGCAACACCGATGATGCTTCGACCCCCCGAAGCTCCT
TCGGGGCTGCATGGGCGCTCCGATGCCGCTCCAGGGCGAGCGCTGTTTAAATAGC
CAGGCCCCCGATTGCAAAGACATTATAGCGAGCTACCAAAGCCATATTCAAACA
CCTAGATCACTACCACTTCTACACAGGCCACTCGAGCTTGTGATCGCACTCCGCT
AAGGGGGCGCCTCTTCCTCTTCGTTTCAGTCACAACCCGCAAACACTAGTATGGC
CACCGCATCCACTTTCTCGGCGTTCAATGCCCGCTGCGGCGACCTGCGTCGCTCG
GCGGGCTCCGGGCCCCGGCGCCCAGCGAGGCCCCTCCCCGTGCGCGGGCGCGCC
AGCATGCTGCTGTCGGCGGTGACCACGGTCTTCGGCGTGGCCGAGAAGCAGTGG
CCCATGCTGGACCGCAAGTCCAAGCGCCCCGACATGCTGGTCGAGCCCCTGGGC
GTGGACCGCATCGTCTACGACGGCGTGAGCTTCCGCCAGTCGTTCTCCATCCGCA
GCTACGAGATCGGCGCCGACCGCACCGCCTCGATCGAGACGCTGATGAACATGT
TCCAGGAGACCTCCCTGAACCACTGCAAGATCATCGGCCTGCTGAACGACGGCTT
CGGCCGCACGCCCGAGATGTGCAAGCGCGACCTGATCTGGGTCGTGACCAAGAT
GCAGATCGAGGTGAACCGCTACCCCACGTGGGGCGACACCATCGAGGTCAACAC
GTGGGTGAGCGCCTCGGGCAAGCACGGCATGGGCCGCGACTGGCTGATCTCCGA
CTGCCACACCGGCGAGATCCTGATCCGCGCGACGAGCGTCTGGGCGATGATGAA
CCAGAAGACCCGCCGCCTGTCGAAGATCCCCTACGAGGTGCGCCAGGAGATCGA
GCCCCAGTTCGTCGACTCCGCCCCCGTGATCGTGGACGACCGCAAGTTCCACAAG
CTGGACCTGAAGACGGGCGACAGCATCTGCAACGGCCTGACCCCCCGCTGGACG
GACCTGGACGTGAACCAGCACGTCAACAACGTGAAGTACATCGGCTGGATCCTG
CAGTCGGTCCCCACCGAGGTGTTCGAGACGCAGGAGCTGTGCGGCCTGACCCTG
GAGTACCGCCGCGAGTGCGGCCGCGACTCCGTGCTGGAGAGCGTCACGGCCATG
GACCCCTCGAAGGAGGGCGACCGCTCCCTGTACCAGCACCTGCTGCGCCTGGAG
GACGGCGCGGACATCGTGAAGGGCCGCACCGAGTGGCGCCCCAAGAACGCCGGC
GCCAAGGGCGCCATCCTGACGGGCAAGACCAGCAACGGCAACTCGATCTCCTGA
CTCGAGTTAATTAACTCGAGGCAGCAGCAGCTCGGATAGTATCGACACACTCTGG
ACGCTGGTCGTGTGATGGACTGTTGCCGCCACACTTGCTGCCTTGACCTGTGAAT
ATCCCTGCCGCTTTTATCAAACAGCCTCAGTGTGTTTGATCTTGTGTGTACGCGCT
TTTGCGAGTTGCTAGCTGCTTGTGCTATTTGCGAATACCACCCCCAGCATCCCCTT
CCCTCGTTTCATATCGCTTGCATCCCAACCGCAACTTATCTACGCTGTCCTGCTAT
CCCTCAGCGCTGCTCCTGCTCCTGCTCACTGCCCCTCGCACAGCCTTGGTTTGGGC

TCCGCCTGTATTCTCCTGGTACTGCAACCTGTAAACCAGCACTGCAATGCTGATG
CACGGGAAGTAGTGGGATGGGAACACAAATGGAAAGCTT

SEQ ID NO: 108
Codon-optimized *Ulmus americana* thioesterase

GGCGCGCCCAGCTGCCCGACTGGAGCATGCTGCTGGCCGCGATCACCACCCTGTT
CCTGGCGGCCGAGAAGCAGTGGATGATGCTGGACTGGAAGCCCAAGCGCCCCGA
CATGCTGGTGGACCCCTTCGGCCTGGGCCGCTTCGTGCAGGACGGCCTGGTGTTC
CGCAACAACTTCAGCATCCGCAGCTACGAGATCGGCGCGGACCGCACCGCCAGC
ATCGAGACCCTGATGAACCACCTGCAGGAGACCGCCCTGAACCACGTGAAGAGC
GTGGGCCTGCTGGAGGACGGCCTGGGCAGCACCCGCGAGATGAGCCTGCGCAAC
CTGATCTGGGTGGTGACCAAGATGCAGGTGGCGGTGGACCGCTACCCCACCTGG
GGCGACGAGGTGCAGGTGAGCAGCTGGGCGACCGCCATCGGCAAGAACGGCAT
GCGCCGCGAGTGGATCGTGACCGACTTCCGCACCGGCGAGACCCTGCTGCGCGC
CACCAGCGTGTGGGTGATGATGAACAAGCTGACCCGCCGCATCAGCAAGATCCC
CGAGGAGGTGTGGCACGAGATCGGCCCCAGCTTCATCGACGCGCCCCCCCTGCC
CACCGTGGAGGACGACGGCCGCAAGCTGACCCGCTTCGACGAGAGCAGCGCCGA
CTTCATCCGCAAGGGCCTGACCCCCCGCTGGAGCGACCTGGACATCAACCAGCA
CGTGAACAACGTGAAGTACATCGGCTGGCTGCTGGAGAGCGCGCCCCCCGAGAT
CCACGAGAGCCACGAGATCGCCAGCCTGACCCTGGAGTACCGCCGCGAGTGCGG
CCGCGACAGCGTGCTGAACAGCGCCACCAAGGTGAGCGACAGCAGCCAGCTGGG
CAAGAGCGCCGTGGAGTGCAACCACCTGGTGCGCCTGCAGAACGGCGGCGAGAT
CGTGAAGGGCCGCACCGTGTGGCGCCCCAAGCGCCCCCTGTACAACGACGGCGC
CGTGGTGGACGTGCCCGCCAAGACCAGCTGACTCGAG

SEQ ID NO: 109
Dual transformation construct: *suc2* sucrose invertase:: *Cuphea hookeriana* C8-10 thioesterase

ggtacc**ctttcttgcgctatgacacttccagcaaaaggtagggcgggctgcgagacggcttcccggcgctgcatgcaacaccgatg**
atgcttcgaccccccgaagctccttcggggctgcatgggcgctccgatgccgctccagggcgagcgctgtttaaatagccaggccc
ccgattgcaaagacattatagcgagctaccaaagccatattcaaacacctagatcactaccacttctacacaggccactcgagctt
gtgatcgcactccgctaaggggcgcctcttcctcttcgtttcagtcacaacccgcaaactctagaatatca**ATG**CTGCTGCAG
GCCTTCCTGTTCCTGCTGGCCGGCTTCGCCGCCAAGATCAGCGCCTCCATGACGAACGAGACGTCCG
ACCGCCCCCTGGTGCACTTCACCCCCAACAAGGGCTGGATGAACGACCCCAACGGCCTGTGGTACG
ACGAGAAGGACGCCAAGTGGCACCTGTACTTCCAGTACAACCCGAACGACACCGTCTGGGGGACGC
CCTTGTTCTGGGGCCACGCCACGTCCGACGACCTGACCAACTGGGAGGACCAGCCCATCGCCATCGC
CCCGAAGCGCAACGACTCCGGCGCCTTCTCCGGCTCCATGGTGGTGGACTACAACAACACCTCCGGC
TTCTTCAACGACACCATCGACCCGCGCCAGCGCTGCGTGGCCATCTGGACCTACAACACCCCGGAGT
CCGAGGAGCAGTACATCTCCTACAGCCTGGACGGCGGCTACACCTTCACCGAGTACCAGAAGAACC
CCGTGCTGGCCGCCAACTCCACCCAGTTCCGCGACCCGAAGGTCTTCTGGTACGAGCCCTCCCAGAA
GTGGATCATGACCGCGGCCAAGTCCCAGGACTACAAGATCGAGATCTACTCCTCCGACGACCTGAA
GTCCTGGAAGCTGGAGTCCGCGTTCGCCAACGAGGGCTTCCTCGGCTACCAGTACGAGTGCCCCGG
CCTGATCGAGGTCCCCACCGAGCAGGACCCCAGCAAGTCCTACTGGGTGATGTTCATCTCCATCAAC
CCCGGCGCCCCGGCCGGCGGCTCCTTCAACCAGTACTTCGTCGGCAGCTTCAACGGCACCCACTTCG
AGGCCTTCGACAACCAGTCCCGCGTGGTGGACTTCGGCAAGGACTACTACGCCCTGCAGACCTTCTT
CAACACCGACCCGACCTACGGGAGCGCCCTGGGCATCGCGTGGGCCTCCAACTGGGAGTACTCCGC
CTTCGTGCCCACCAACCCCTGGCGCTCCTCCATGTCCCTCGTGCGCAAGTTCTCCCTCAACACCGAGT

ACCAGGCCAACCCGGAGACGGAGCTGATCAACCTGAAGGCCGAGCCGATCCTGAACATCAGCAACG
CCGGCCCCTGGAGCCGGTTCGCCACCAACACCACGTTGACGAAGGCCAACAGCTACAACGTCGACC
TGTCCAACAGCACCGGCACCCTGGAGTTCGAGCTGGTGTACGCCGTCAACACCACCCAGACGATCTC
CAAGTCCGTGTTCGCGGACCTCTCCCTCTGGTTCAAGGGCCTGGAGGACCCCGAGGAGTACCTCCGC
ATGGGCTTCGAGGTGTCCGCGTCCTCCTTCTTCCTGGACCGCGGGAACAGCAAGGTGAAGTTCGTG
AAGGAGAACCCCTACTTCACCAACCGCATGAGCGTGAACAACCAGCCCTTCAAGAGCGAGAACGAC
CTGTCCTACTACAAGGTGTACGGCTTGCTGGACCAGAACATCCTGGAGCTGTACTTCAACGACGGCG
ACGTCGTGTCCACCAACACCTACTTCATGACCACCGGGAACGCCCTGGGCTCCGTGAACATGACGAC
GGGGGTGGACAACCTGTTCTACATCGACAAGTTCCAGGTGCGCGAGGTCAAG**TGA**caattg*gcagcagc*
agctcggatagtatcgacacactctggacgctggtcgtgtgatggactgttgccgccacacttgctgccttgacctgtgaatatc
cctgccgcttttatcaaacagcctcagtgtgtttgatcttgtgtgtacgcgcttttgcgagttgctagctgcttgtgctatttgcgaa
taccacccccagcatcccccttccctcgtttcatatcgcttgcatcccaaccgcaacttatctacgctgtcctgctatccctcagcgct
gctcctgctcctgctcactgcccctcgcacagccttggtttgggctccgcctgtattctcctggtactgcaacctgtaaaccagcact
*gcaatgctgatgcacgggaagtagtgggatgggaacacaaatggaggatcc*cgcgtctcgaacagagcgcgcagaggaa
cgctgaaggtctcgcctctgtcgcacctcagcgcggcatacaccacaataaccacctgacgaatgcgcttggttcttcgtccatt
agcgaagcgtccggttcacacacgtgccacgttggcgaggtggcaggtgacaatgatcggtggagctgatggtcgaaacgtt
*cacagcctagggatatc*gaattc**cgcctgcaacgcaagggcagccacagccgctcccacccgccgctgaaccgacacgtgcttgg**
gcgcctgccgcctgcctgccgcatgcttgtgctggtgaggctgggcagtgctgccatgctgattgaggcttggttcatcgggtggaa
gcttatgtgtgtgctgggcttgcatgccgggcaatgcgcatggtggcaagagggcggcagcacttgctggagctgccgcggtgcct
ccaggtggttcaatcgcggcagccagagggatttcagatgatcgcgcgtacaggttgagcagcagtgtcagcaaaggtagcagtt
tgccagaatgatcggttcagctgttaatcaatgccagcaagagaaggggtcaagtgcaaacacgggcatgccacagcacgggca
ccggggagtggaatggcaccaccaagtgtgtgcgagccagcatcgccgcctggctgtttcagctacaacggcaggagtcatccaa
cgtaaccatgagctgatcaacactgcaatcatcgggcgggcgtgatgcaagcatgcctggcgaagacacatggtgtgcggatgct
gccggctgctgcctgctgcgcacgccgttgagttggcagcaggctcagccatgcactggatggcagctgggctgccactgcaatgt
ggtggataggatgcaagtggagcgaataccaaaccctctggctgcttgctgggttgcatggcatcgcaccatcagcaggagcgca
tgcgaagggactggccccatgcacgccatgccaaaccggagcgcaccgagtgtccacactgtcaccaggcccgcaagctttgcag
aaccatgctcatggacgcatgtagcgctgacgtcccttgacggcgctcctctcgggtgtgggaaacgcaatgcagcacaggcagc
agaggcggcggcagcagagcggcggcagcagcggcgggggccacccttcttgcggggtcgcgccccagccagcggtgatgcgct
gatcccaaacgagttcacattcatttgcatgcctggagaagcgaggctggggcctttgggctggtgcagcccgcaatggaatgcg
ggaccgccaggctagcagcaaaggcgcctccctactccgcatcgatgttccatagtgcattggactgcatttgggtggggcggcc
ggctgtttctttcgtgttgcaaaacgcgccagctcagcaacctgtcccgtgggtccccgtgccgatgaaatcgtgtgcacgccgat
cagctgattgcccggctcgcgaagtaggcgccctcctttctgctcgccctctctccgtcccgccactagt**ATG**GCCACCGCATC
CACTTTCTCGGCGTTCAATGCCCGCTGCGGCGACCTGCGTCGCTCGGCGGGCTCCGGGCCCCGGCGC
CCAGCGAGGCCCCTCCCCGTGCGCGGGCGCGCCCAGCTGCCCGACTGGAGCCGCCTGCTGACCGCC
ATCACCACCGTGTTCGTGAAGTCCAAGCGCCCCGACATGCACGACCGCAAGTCCAAGCGCCCCGACA
TGCTGGTGGACAGCTTCGGCCTGGAGTCCACCGTGCAGGACGGCCTGGTGTTCCGCCAGTCCTTCTC
CATCCGCTCCTACGAGATCGGCACCGACCGCACCGCCAGCATCGAGACCCTGATGAACCACCTGCAG
GAGACCTCCCTGAACCACTGCAAGAGCACCGGCATCCTGCTGGACGGCTTCGGCCGCACCCTGGAG
ATGTGCAAGCGCGACCTGATCTGGGTGGTGATCAAGATGCAGATCAAGGTGAACCGCTACCCCGCC
TGGGGCGACACCGTGGAGATCAACACCCGCTTCAGCCGCCTGGGCAAGATCGGCATGGGCCGCGA
CTGGCTGATCTCCGACTGCAACACCGGCGAGATCCTGGTGCGCGCCACCAGCGCCTACGCCATGATG
AACCAGAAGACCCGCCGCCTGTCCAAGCTGCCCTACGAGGTGCACCAGGAGATCGTGCCCCTGTTC
GTGGACAGCCCCGTGATCGAGGACTCCGACCTGAAGGTGCACAAGTTCAAGGTGAAGACCGGCGA
CAGCATCCAGAAGGGCCTGACCCCCGGCTGGAACGACCTGGACGTGAACCAGCACGTGTCCAACGT
GAAGTACATCGGCTGGATCCTGGAGAGCATGCCCACCGAGGTGCTGGAGACCCAGGAGCTGTGCTC
CCTGGCCCTGGAGTACCGCCGCGAGTGCGGCCGCGACTCCGTGCTGGAGAGCGTGACCGCCATGG
ACCCCAGCAAGGTGGGCGTGCGCTCCCAGTACCAGCACCTGCTGCGCCTGGAGGACGGCACCGCCA

TCGTGAACGGCGCCACCGAGTGGCGCCCCAAGAACGCCGGCGCCAACGGCGCCATCTCCACCGGCA
AGACCAGCAACGGCAACTCCGTGTCC*ATGGACTACAAGGACCACGACGGCGACTACAAGGACCACG*
*ACATCGACTACAAGGACGACGACGACAAGTGA*ctcgagg*cagcagcagctcggatagtatcgacacactctgga*
cgctggtcgtgtgatggactgttgccgccacacttgctgccttgacctgtgaatatccctgccgcttttatcaaacagcctcagtgt
gtttgatcttgtgtgtacgcgcttttgcgagttgctagctgcttgtgctatttgcgaataccacccccagcatcccctttccctcgtttc
atatcgcttgcatcccaaccgcaacttatctacgctgtcctgctatccctcagcgctgctcctgctcctgctcactgcccctcgcaca
gccttggtttgggctccgcctgtattctcctggtactgcaacctgtaaaccagcactgcaatgctgatgcacgggaagtagtggg
*atgggaacacaaatgga*aagctt

SEQ ID NO: 110
Dual transformation construct: *suc2* sucrose invertase:: *Umbellularia californica* C12 thioesterase

ggtacc**ctttcttgcgctatgacacttccagcaaaaggtagggcgggctgcgagacggcttcccggcgctgcatgcaacaccgatg**
atgcttcgaccccccgaagctccttcggggctgcatgggcgctccgatgccgctccagggcgagcgctgtttaaatagccaggccc
ccgattgcaaagacattatagcgagctaccaaagccatattcaaacacctagatcactaccacttctacacaggccactcgagctt
gtgatcgcactccgctaaggggcgcctcttcctcttcgtttcagtcacaacccgcaaactctaga**ATG**CTGCTGCAG
GCCTTCCTGTTCCTGCTGGCCGGCTTCGCCGCCAAGATCAGCGCCTCCATGACGAACGAGACGTCCG
ACCGCCCCCTGGTGCACTTCACCCCCAACAAGGGCTGGATGAACGACCCCAACGGCCTGTGGTACG
ACGAGAAGGACGCCAAGTGGCACCTGTACTTCCAGTACAACCCGAACGACACCGTCTGGGGGACGC
CCTTGTTCTGGGGCCACGCCACGTCCGACGACCTGACCAACTGGGAGGACCAGCCCATCGCCATCGC
CCCGAAGCGCAACGACTCCGGCGCCTTCTCCGGCTCCATGGTGGTGGACTACAACAACACCTCCGGC
TTCTTCAACGACACCATCGACCCGCGCCAGCGCTGCGTGGCCATCTGGACCTACAACACCCCGGAGT
CCGAGGAGCAGTACATCTCCTACAGCCTGGACGGCGGCTACACCTTCACCGAGTACCAGAAGAACC
CCGTGCTGGCCGCCAACTCCACCCAGTTCCGCGACCCGAAGGTCTTCTGGTACGAGCCCTCCCAGAA
GTGGATCATGACCGCGGCCAAGTCCCAGGACTACAAGATCGAGATCTACTCCTCCGACGACCTGAA
GTCCTGGAAGCTGGAGTCCGCGTTCGCCAACGAGGGCTTCCTCGGCTACCAGTACGAGTGCCCCGG
CCTGATCGAGGTCCCCACCGAGCAGGACCCCAGCAAGTCCTACTGGGTGATGTTCATCTCCATCAAC
CCCGGCGCCCCGGCCGGCGGCTCCTTCAACCAGTACTTCGTCGGCAGCTTCAACGGCACCCACTTCG
AGGCCTTCGACAACCAGTCCCGCGTGGTGGACTTCGGCAAGGACTACTACGCCCTGCAGACCTTCTT
CAACACCGACCCGACCTACGGGAGCGCCCTGGGCATCGCGTGGGCCTCCAACTGGGAGTACTCCGC
CTTCGTGCCCACCAACCCCTGGCGCTCCTCCATGTCCCTCGTGCGCAAGTTCTCCCTCAACACCGAGT
ACCAGGCCAACCCGGAGACGGAGCTGATCAACCTGAAGGCCGAGCCGATCCTGAACATCAGCAACG
CCGGCCCCTGGAGCCGGTTCGCCACCAACACCACGTTGACGAAGGCCAACAGCTACAACGTCGACC
TGTCCAACAGCACCGGCACCCTGGAGTTCGAGCTGGTGTACGCCGTCAACACCACCCAGACGATCTC
CAAGTCCGTGTTCGCGGACCTCTCCCTCTGGTTCAAGGGCCTGGAGGACCCCGAGGAGTACCTCCGC
ATGGGCTTCGAGGTGTCCGCGTCCTCCTTCTTCCTGGACCGCGGGAACAGCAAGGTGAAGTTCGTG
AAGGAGAACCCCTACTTCACCAACCGCATGAGCGTGAACAACCAGCCCTTCAAGAGCGAGAACGAC
CTGTCCTACTACAAGGTGTACGGCTTGCTGGACCAGAACATCCTGGAGCTGTACTTCAACGACGGCG
ACGTCGTGTCCACCAACACCTACTTCATGACCACCGGGAACGCCCTGGGCTCCGTGAACATGACGAC
GGGGGTGGACAACCTGTTCTACATCGACAAGTTCCAGGTGCGCGAGGTCAAG**TGA**caattg*gcagcag*
cagctcggatagtatcgacacactctggacgctggtcgtgtgatggactgttgccgccacacttgctgccttgacctgtgaatat
ccctgccgcttttatcaaacagcctcagtgtgtttgatcttgtgtgtacgcgcttttgcgagttgctagctgcttgtgctatttgcga
ataccacccccagcatcccctttccctcgtttcatatcgcttgcatcccaaccgcaacttatctacgctgtcctgctatccctcagcgc
tgctcctgctcctgctcactgcccctcgcacagccttggtttgggctccgcctgtattctcctggtactgcaacctgtaaaccagcac
*tgcaatgctgatgcacgggaagtagtgggatgggaacacaaatggaggatcc*cgcgtctcgaacagagcgcgcagaggaac
gctgaaggtctcgcctctgtcgcacctcagcgcggcatacaccacaataaccacctgacgaatgcgcttggttcttcgtccattagcg

aagcgtccggttcacacacgtgccacgttggcgaggtggcaggtgacaatgatcggtggagctgatggtcgaaacgttcacagcct
agggatatcgaattc**cgcctgcaacgcaagggcagccacagccgctcccacccgccgctgaaccgacacgtgcttgggcgcctgc**
cgcctgcctgccgcatgcttgtgctggtgaggctgggcagtgctgccatgctgattgaggcttggttcatcgggtggaagcttatgtg
tgtgctgggcttgcatgccgggcaatgcgcatggtggcaagagggcggcagcacttgctggagctgccgcggtgcctccaggtggt
tcaatcgcggcagccagagggatttcagatgatcgcgcgtacaggttgagcagcagtgtcagcaaaggtagcagtttgccagaat
gatcggttcagctgttaatcaatgccagcaagagaaggggtcaagtgcaaacacgggcatgccacagcacgggcaccggggag
tggaatggcaccaccaagtgtgtgcgagccagcatcgccgcctggctgtttcagctacaacggcaggagtcatccaacgtaacca
tgagctgatcaacactgcaatcatcgggcgggcgtgatgcaagcatgcctggcgaagacacatggtgtgcggatgctgccggctg
ctgcctgctgcgcacgccgttgagttggcagcaggctcagccatgcactggatggcagctgggctgccactgcaatgtggtggata
ggatgcaagtggagcgaataccaaaccctctggctgcttgctgggttgcatggcatcgcaccatcagcaggagcgcatgcgaagg
gactggccccatgcacgccatgccaaaccggagcgcaccgagtgtccacactgtcaccaggcccgcaagctttgcagaaccatgc
tcatggacgcatgtagcgctgacgtcccttgacggcgctcctctcgggtgtgggaaacgcaatgcagcacaggcagcagaggcgg
cggcagcagagcggcggcagcagcggcgggggccaccctttcttgcggggtcgcgccccagccagcggtgatgcgctgatcccaa
acgagttcacattcatttgcatgcctggagaagcgaggctgggggcctttgggctggtgcagcccgcaatggaatgcgggaccgcc
aggctagcagcaaaggcgcctcccctactccgcatcgatgttccatagtgcattggactgcatttgggtggggcggccggctgtttc
tttcgtgttgcaaaacgcgccagctcagcaacctgtcccgtgggtcccccgtgccgatgaaatcgtgtgcacgccgatcagctgatt
gcccggctcgcgaagtaggcgccctcctttctgctcgccctctctccgtcccgccactagt**ATG**ACGTTCGGGGTCGCCCTC
CCGGCCATGGGCCGCGGTGTCTCCCTTCCCCGGCCCAGGGTCGCGGTGCGCGCCCAGTCGGCGAGT
CAGGTTTTGGAGAGCGGGCGCGCCCCCGACTGGTCCATGCTGTTCGCCGTGATCACCACCATCTTCA
GCGCCGCCGAGAAGCAGTGGACCAACCTGGAGTGGAAGCCCAAGCCCAAGCTGCCCCAGCTGCTG
GACGACCACTTCGGCCTGCACGGCCTGGTGTTCCGCCGCACCTTCGCCATCCGCTCCTACGAGGTGG
GCCCCGACCGCAGCACCTCCATCCTGGCCGTGATGAACCACATGCAGGAGGCCACCCTGAACCACG
CCAAGAGCGTGGGCATCCTGGGCGACGGCTTCGGCACCACCCTGGAGATGTCCAAGCGCGACCTGA
TGTGGGTGGTGCGCCGCACCCACGTGGCCGTGGAGCGCTACCCCACCTGGGGCGACACCGTGGAG
GTGGAGTGCTGGATCGGCGCCAGCGGCAACAACGGCATGCGCCGCGACTTCCTGGTGCGCGACTG
CAAGACCGGCGAGATCCTGACCCGCTGCACCTCCCTGAGCGTGCTGATGAACACCCGCACCCGCCGC
CTGAGCACCATCCCCGACGAGGTGCGCGGCGAGATCGGCCCCGCCTTCATCGACAACGTGGCCGTG
AAGGACGACGAGATCAAGAAGCTGCAGAAGCTGAACGACTCCACCGCCGACTACATCCAGGGCGG
CCTGACCCCCCGCTGGAACGACCTGGACGTGAACCAGCACGTGAACAACCTGAAGTACGTGGCCTG
GGTGTTCGAGACCGTGCCCGACAGCATCTTCGAGTCCCACCACATCAGCTCCTTCACCCTGGAGTAC
CGCCGCGAGTGCACCCGCGACTCCGTGCTGCGCAGCCTGACCACCGTGAGCGGCGGCAGCTCCGAG
GCCGGCCTGGTGTGCGACCACCTGCTGCAGCTGGAGGGCGGCAGCGAGGTGCTGCGCGCCCGCAC
CGAGTGGCGCCCCAAGCTGACCGACTCCTTCCGCGGCATCAGCGTGATCCCCGCCGAGCCCCGCGT
GATGGACTACAAGGACCACGACGGCGACTACAAGGACCACGACATCGACTACAAGGACGACGACGA
*CAAGTGA*ctcgag*gcagcagcagctcggatagtatcgacacactctggacgctggtcgtgtgatggactgttgccgccacac*
ttgctgccttgacctgtgaatatccctgccgcttttatcaaacagcctcagtgtgtttgatcttgtgtgtacgcgcttttgcgagttg
ctagctgcttgtgctatttgcgaataccacccccagcatcccctttccctcgtttcatatcgcttgcatcccaaccgcaacttatctac
gctgtcctgctatccctcagcgctgctcctgctcctgctcactgcccctcgcacagccttggtttgggctccgcctgtattctcctggt
*actgcaacctgtaaaccagcactgcaatgctgatgcacgggaagtagtgggatgggaacacaaatgga*aagctt

SEQ ID NO: 111

Dual transformation construct: *suc2* sucrose invertase:: *Cinnamomum camphora* C14 thioesterase

ggtacc**ctttcttgcgctatgacacttccagcaaaaggtagggcgggctgcgagacggcttcccggcgctgcatgcaacaccgatg**
atgcttcgaccccccgaagctccttcggggctgcatgggcgctccgatgccgctccagggcgagcgctgtttaaatagccaggccc
**ccgattgcaaagacattatagcgagctaccaaagccatattcaaacacctagatcactaccacttctacacaggccactcgagctt

gtgatcgcactccgctaaggggcgcctcttcctcttcgtttcagtcacaacccgcaaactctaga**atatca**ATG**CTGCTGCAG**
GCCTTCCTGTTCCTGCTGGCCGGCTTCGCCGCCAAGATCAGCGCCTCCATGACGAACGAGACGTCCG
ACCGCCCCCTGGTGCACTTCACCCCCAACAAGGGCTGGATGAACGACCCCAACGGCCTGTGGTACG
ACGAGAAGGACGCCAAGTGGCACCTGTACTTCCAGTACAACCCGAACGACACCGTCTGGGGGACGC
CCTTGTTCTGGGGCCACGCCACGTCCGACGACCTGACCAACTGGGAGGACCAGCCCATCGCCATCGC
CCCGAAGCGCAACGACTCCGGCGCCTTCTCCGGCTCCATGGTGGTGGACTACAACAACACCTCCGGC
TTCTTCAACGACACCATCGACCCGCGCCAGCGCTGCGTGGCCATCTGGACCTACAACACCCCGGAGT
CCGAGGAGCAGTACATCTCCTACAGCCTGGACGGCGGCTACACCTTCACCGAGTACCAGAAGAACC
CCGTGCTGGCCGCCAACTCCACCCAGTTCCGCGACCCGAAGGTCTTCTGGTACGAGCCCTCCCAGAA
GTGGATCATGACCGCGGCCAAGTCCCAGGACTACAAGATCGAGATCTACTCCTCCGACGACCTGAA
GTCCTGGAAGCTGGAGTCCGCGTTCGCCAACGAGGGCTTCCTCGGCTACCAGTACGAGTGCCCCGG
CCTGATCGAGGTCCCCACCGAGCAGGACCCCAGCAAGTCCTACTGGGTGATGTTCATCTCCATCAAC
CCCGGCGCCCCGGCCGGCGGCTCCTTCAACCAGTACTTCGTCGGCAGCTTCAACGGCACCCACTTCG
AGGCCTTCGACAACCAGTCCCGCGTGGTGGACTTCGGCAAGGACTACTACGCCCTGCAGACCTTCTT
CAACACCGACCCGACCTACGGGAGCGCCCTGGGCATCGCGTGGGCCTCCAACTGGGAGTACTCCGC
CTTCGTGCCCACCAACCCCTGGCGCTCCTCCATGTCCCTCGTGCGCAAGTTCTCCCTCAACACCGAGT
ACCAGGCCAACCCGGAGACGGAGCTGATCAACCTGAAGGCCGAGCCGATCCTGAACATCAGCAACG
CCGGCCCCTGGAGCCGGTTCGCCACCAACACCACGTTGACGAAGGCCAACAGCTACAACGTCGACC
TGTCCAACAGCACCGGCACCCTGGAGTTCGAGCTGGTGTACGCCGTCAACACCACCCAGACGATCTC
CAAGTCCGTGTTCGCGGACCTCTCCCTCTGGTTCAAGGGCCTGGAGGACCCCGAGGAGTACCTCCGC
ATGGGCTTCGAGGTGTCCGCGTCCTCCTTCTTCCTGGACCGCGGGAACAGCAAGGTGAAGTTCGTG
AAGGAGAACCCCTACTTCACCAACCGCATGAGCGTGAACAACCAGCCCTTCAAGAGCGAGAACGAC
CTGTCCTACTACAAGGTGTACGGCTTGCTGGACCAGAACATCCTGGAGCTGTACTTCAACGACGGCG
ACGTCGTGTCCACCAACACCTACTTCATGACCACCGGGAACGCCCTGGGCTCCGTGAACATGACGAC
GGGGGTGGACAACCTGTTCTACATCGACAAGTTCCAGGTGCGCGAGGTCAAG**TGA**caattg*gcagcagc*
agctcggatagtatcgacacactctggacgctggtcgtgtgatggactgttgccgccacacttgctgccttgacctgtgaatatc
cctgccgcttttatcaaacagcctcagtgtgtttgatcttgtgtgtacgcgcttttgcgagttgctagctgcttgtgctatttgcgaa
taccacccccagcatcccctttccctcgtttcatatcgcttgcatcccaaccgcaacttatctacgctgtcctgctatccctcagcgct
gctcctgctcctgctcactgcccctcgcacagccttggtttgggctccgcctgtattctcctggtactgcaacctgtaaaccagcact
gcaatgctgatgcacgggaagtagtgggatgggaacacaaatggaggatcccgcgtctcgaacagagcgcgcagaggaa
cgctgaaggtctcgcctctgtcgcacctcagcgcggcatacaccacaataaccacctgacgaatgcgcttggttcttcgtccatt
agcgaagcgtccggttcacacacgtgccacgttggcgaggtggcaggtgacaatgatcggtggagctgatggtcgaaacgtt
*cacagcctagggatatcg*aattc**cgcctgcaacgcaagggcagccacagccgctcccacccgccgctgaaccgacacgtgcttgg**
gcgcctgccgcctgcctgccgcatgcttgtgctggtgaggctgggcagtgctgccatgctgattgaggcttggttcatcgggtggaa
gcttatgtgtgtgctgggcttgcatgccgggcaatgcgcatggtggcaagagggcggcagcacttgctggagctgccgcggtgcct
ccaggtggttcaatcgcggcagccagagggatttcagatgatcgcgcgtacaggttgagcagcagtgtcagcaaaggtagcagtt
tgccagaatgatcggttcagctgttaatcaatgccagcaagagaaggggtcaagtgcaaacacgggcatgccacagcacgggca
ccggggagtggaatggcaccaccaagtgtgtgcgagccagcatcgccgcctggctgtttcagctacaacggcaggagtcatccaa
cgtaaccatgagctgatcaacactgcaatcatcgggcgggcgtgatgcaagcatgcctggcgaagacacatggtgtgcggatgct
gccggctgctgcctgctgcgcacgccgttgagttggcagcaggctcagccatgcactggatggcagctgggctgccactgcaatgt
ggtggataggatgcaagtggagcgaataccaaaccctctggctgcttgctgggttgcatggcatcgcaccatcagcaggagcgca
tgcgaagggactggccccatgcacgccatgccaaaccggagcgcaccgagtgtccacactgtcaccaggcccgcaagctttgcag
aaccatgctcatggacgcatgtagcgctgacgtcccttgacggcgctcctctcgggtgtgggaaacgcaatgcagcacaggcagc
agaggcggcggcagcagagcggcggcagcagcggcgggggccacccttcttgcggggtcgcgccccagccagcggtgatgcgct
gatcccaaacgagttcacattcatttgcatgcctggagaagcgaggctggggcctttgggctggtgcagcccgcaatggaatgcg
ggaccgccaggctagcagcaaaggcgcctcccctactccgcatcgatgttccatagtgcattggactgcatttgggtggggcggcc
ggctgtttctttcgtgttgcaaaacgcgccagctcagcaacctgtcccgtgggtcccccgtgccgatgaaatcgtgtgcacgccgat

cagctgattgcccggctcgcgaagtaggcgccctcctttctgctcgccctctctccgtcccgccactagtATGGCCACCGCATC
CACTTTCTCGGCGTTCAATGCCCGCTGCGGCGACCTGCGTCGCTCGGCGGGCTCCGGGCCCCGGCGC
CCAGCGAGGCCCCTCCCCGTGCGCGGGCGCGCCCCCGACTGGTCCATGCTGTTCGCCGTGATCACCA
CCATCTTCTCCGCCGCCGAGAAGCAGTGGACCAACCTGGAGTGGAAGCCCAAGCCCAACCCCCCCCA
GCTGCTGGACGACCACTTCGGCCCCCACGGCCTGGTGTTCCGCCGCACCTTCGCCATCCGCAGCTAC
GAGGTGGGCCCCGACCGCTCCACCAGCATCGTGGCCGTGATGAACCACCTGCAGGAGGCCGCCCTG
AACCACGCCAAGTCCGTGGGCATCCTGGGCGACGGCTTCGGCACCACCCTGGAGATGTCCAAGCGC
GACCTGATCTGGGTGGTGAAGCGCACCCACGTGGCCGTGGAGCGCTACCCCGCCTGGGGCGACACC
GTGGAGGTGGAGTGCTGGGTGGGCGCCTCCGGCAACAACGGCCGCCGCCACGACTTCCTGGTGCG
CGACTGCAAGACCGGCGAGATCCTGACCCGCTGCACCTCCCTGAGCGTGATGATGAACACCCGCAC
CCGCCGCCTGAGCAAGATCCCCGAGGAGGTGCGCGGCGAGATCGGCCCCGCCTTCATCGACAACGT
GGCCGTGAAGGACGAGGAGATCAAGAAGCCCCAGAAGCTGAACGACTCCACCGCCGACTACATCC
AGGGCGGCCTGACCCCCCGCTGGAACGACCTGGACATCAACCAGCACGTGAACAACATCAAGTACG
TGGACTGGATCCTGGAGACCGTGCCCGACAGCATCTTCGAGAGCCACCACATCTCCTCCTTCACCATC
GAGTACCGCCGCGAGTGCACCATGGACAGCGTGCTGCAGTCCCTGACCACCGTGAGCGGCGGCTCC
TCCGAGGCCGGCCTGGTGTGCGAGCACCTGCTGCAGCTGGAGGGCGGCAGCGAGGTGCTGCGCGC
CAAGACCGAGTGGCGCCCCAAGCTGACCGACTCCTTCCGCGGCATCAGCGTGATCCCCGCCGAGTCC
AGCGTG*ATGGACTACAAGGACCACGACGGCGACTACAAGGACCACGACATCGACTACAAGGACGAC*
*GACGACAAGTGA*ctcgag*gcagcagcagctcggatagtatcgacacactctggacgctggtcgtgtgatggactgttgcc*
gccacacttgctgccttgacctgtgaatatccctgccgcttttatcaaacagcctcagtgtgtttgatcttgtgtgtacgcgcttttg
cgagttgctagctgcttgtgctatttgcgaataccacccccagcatcccctccctcgtttcatatcgcttgcatcccaaccgcaact
tatctacgctgtcctgctatccctcagcgctgctcctgctcctgctcactgcccctcgcacagccttggtttgggctccgcctgtattc
*tcctggtactgcaacctgtaaaccagcactgcaatgctgatgcacgggaagtagtgggatgggaacacaaatgga*aagctt

SEQ ID NO: 112
P.moriformis Aat02 promoter/5'UTR::*C.protothecoides* stearoyl ACP desaturase PTS::*Cinnamomum camphora* C14 thioesterase::*C.vulgaris* nitrate reductase 3"UTR

AGAGAGCGGAGGTGGGGTTGTGAGGTGGGGTTGCTGACCAGGAGCTCGCGTCGC
CGAGCGCGACTCGCACACGGTCCAGTTACCCCCCCCTCCGCCCAAACGCAAGCCT
CCCATCTTGATGCCTTTCCGGCCACCTATACTATTTCTTAGTTCGCTGTAACATCC
AGACCGTCCTGAATAATAACAATGCCCTGTGTCAAGTGCATTCCTAAAAAAATTC
TGTCCCAACCAACAATCCCACCTGAAATACCACCAGCCCTGCCCAGTACACTCTT
CCAATACCATCTCCCTACCTCCACGCGCAAGCGACCCCCATGCGCGACCAGGCTC
GAAAGTGATTTATGACTTGAGACGAGCGAGTGGCGGCGCGGTCGACTGCCTTTTC
ATCACGTGCCGTACGTCGGCGACCGCTAGGGCTTTGCACGGCAACGCACGGCTTC
GCCAACCCGACCAGCCAGGACCTCGACTACTCTACCGCGAATTCGCCTCAAGAA
GTCGCCAAATGTGCCATACACCATTCCTTACAGCACTGTTCAAACTTGATGCCAA
TTTTGACATTCGGGTTGCTCGTTGGCTGCGCCCACATCGGCCGTGAGTGCAGCAG
GCGGGATCGGACACGGAGGACGCGGCGTCACGCCCCGAACGCAGCCCGTAACTC
TACATCAACACGACGTGTTGCGTAATCCCGCCCGGCTGCGCATCGTGCCAACCCA
TTCGCGATGGATGGTCGGAAAATGGTGTGCCAACTGCCCTGAGGGAGGCTCTCG
CGAAACGGGCACGTCCCTGAAACCGAAACTGTGGCCTTGTCGTCGGCCACGCAA
GCACGTGGACCCTAAACACCAAGAAAATCAGTAAACAAGGTTGACATCCTCTAC
GGGCGAATTGTTTGCCCAACCCTTCATCGCACACTGCCATTATAATGCATCTAGC
TCGGCGACAAGTTTAGAAAAGGCAGGCTGCATTGTTCCATTTCGCCGTGGCGGCG
TGGGTGCCCATTTTACGAGGTTTGGGCTCCCGGGCAGCGACCGAGCCAGGTCGA
GTCCCTCTCGCCCGTCGACAATGTTGCGAACCCCACAAGCGGCTAACAACAACTT
GATGGTACCTGTACACTGCCAATTCCTTCTTCCCCGGCCGAGGTTTACACGTGAT

GGCCATGGCTTCGCATTCAGGCCGACTTCCCATTCCGACTTTCCAGAGGGTCCGC
GGACGCTGGGGGGTTGGCTGCCTGAGGCCCACCCTTTGTTCCCCGCGTCCCGACAA
ACACAATTGCGTTACATAAGGGGGAGCCGCCCCCGTTCAGAGTGCAGAAATCTTT
CACTATATTTTCCAGTCGTCAGCGAAATCAAGTACTAGTATGGCCACCGCATCCA
CTTTCTCGGCGTTCAATGCCCGCTGCGGCGACCTGCGTCGCTCGGCGGGCTCCGG
GCCCCGGCGCCCAGCGAGGCCCCTCCCCGTGCGCGGGCGCGCCCCCGACTGGTC
CATGCTGTTCGCCGTGATCACCACCATCTTCTCCGCCGCCGAGAAGCAGTGGACC
AACCTGGAGTGGAAGCCCAAGCCCAACCCCCCCCAGCTGCTGGACGACCACTTC
GGCCCCCACGGCCTGGTGTTCCGCCGCACCTTCGCCATCCGCAGCTACGAGGTGG
GCCCCGACCGCTCCACCAGCATCGTGGCCGTGATGAACCACCTGCAGGAGGCCG
CCCTGAACCACGCCAAGTCCGTGGGCATCCTGGGCGACGGCTTCGGCACCACCCT
GGAGATGTCCAAGCGCGACCTGATCTGGGTGGTGAAGCGCACCCACGTGGCCGT
GGAGCGCTACCCCGCCTGGGGCGACACCGTGGAGGTGGAGTGCTGGGTGGGCGC
CTCCGGCAACAACGGCCGCCGCCACGACTTCCTGGTGCGCGACTGCAAGACCGG
CGAGATCCTGACCCGCTGCACCTCCCTGAGCGTGATGATGAACACCCGCACCCGC
CGCCTGAGCAAGATCCCCGAGGAGGTGCGCGGCGAGATCGGCCCCGCCTTCATC
GACAACGTGGCCGTGAAGGACGAGGAGATCAAGAAGCCCCAGAAGCTGAACGA
CTCCACCGCCGACTACATCCAGGGCGGCCTGACCCCCCGCTGGAACGACCTGGA
CATCAACCAGCACGTGAACAACATCAAGTACGTGGACTGGATCCTGGAGACCGT
GCCCGACAGCATCTTCGAGAGCCACCACATCTCCTCCTTCACCATCGAGTACCGC
CGCGAGTGCACCATGGACAGCGTGCTGCAGTCCCTGACCACCGTGAGCGGCGGC
TCCTCCGAGGCCGGCCTGGTGTGCGAGCACCTGCTGCAGCTGGAGGGCGGCAGC
GAGGTGCTGCGCGCCAAGACCGAGTGGCGCCCCAAGCTGACCGACTCCTTCCGC
GGCATCAGCGTGATCCCCGCCGAGTCCAGCGTGATGGACTACAAGGACCACGAC
GGCGACTACAAGGACCACGACATCGACTACAAGGACGACGACGACAAGTGACTC
GAGTTAATTAACTCGAGGCAGCAGCAGCTCGGATAGTATCGACACACTCTGGAC
GCTGGTCGTGTGATGGACTGTTGCCGCCACACTTGCTGCCTTGACCTGTGAATAT
CCCTGCCGCTTTTATCAAACAGCCTCAGTGTGTTTGATCTTGTGTGTACGCGCTTT
TGCGAGTTGCTAGCTGCTTGTGCTATTTGCGAATACCACCCCCAGCATCCCCTTCC
CTCGTTTCATATCGCTTGCATCCCAACCGCAACTTATCTACGCTGTCCTGCTATCC
CTCAGCGCTGCTCCTGCTCCTGCTCACTGCCCCTCGCACAGCCTTGGTTTGGGCTC
CGCCTGTATTCTCCTGGTACTGCAACCTGTAAACCAGCACTGCAATGCTGATGCA
CGGGAAGTAGTGGGATGGGAACACAAATGGAAAGCTT

SEQ ID NO: 113
C.reinhartii β-tubulin promoter::native PTS::*Cuphea hookeriana* C8-10 thioesterase::*C.vulgaris* nitrate reductase 3'UTR

ggtacc**cgcctgcaacgcaagggcagccacagccgctcccacccgccgctgaaccgacacgtgcttgggcgcctgccgcctgcctg**
ccgcatgcttgtgctggtgaggctgggcagtgctgccatgctgattgaggcttggttcatcgggtggaagcttatgtgtgtgctggg
cttgcatgccgggcaatgcgcatggtggcaagagggcggcagcacttgctggagctgccgcggtgcctccaggtggttcaatcgc
ggcagccagagggatttcagatgatcgcgcgtacaggttgagcagcagtgtcagcaaaggtagcagtttgccagaatgatcggtt
cagctgttaatcaatgccagcaagagaaggggtcaagtgcaaacacgggcatgccacagcacgggcaccggggagtggaatgg
caccaccaagtgtgtgcgagccagcatcgccgcctggctgtttcagctacaacggcaggagtcatccaacgtaaccatgagctga
tcaacactgcaatcatcgggcgggcgtgatgcaagcatgcctggcgaagacacatggtgtgcggatgctgccggctgctgcctgct
gcgcacgccgttgagttggcagcaggctcagccatgcactggatggcagctgggctgccactgcaatgtggtggataggatgcaa
gtggagcgaataccaaaccctctggctgcttgctgggttgcatggcatcgcaccatcagcaggagcgcatgcgaagggactggcc
ccatgcacgccatgccaaaccggagcgcaccgagtgtccacactgtcaccaggcccgcaagctttgcagaaccatgctcatggac
gcatgtagcgctgacgtcccttgacggcgctcctctcgggtgtgggaaacgcaatgcagcacaggcagcagaggcggcggcagc

agagcggcggcagcagcggcgggggccacccttcttgcggggtcgcgccccagccagcggtgatgcgctgatcccaaacgagttc
acattcatttgcatgcctggagaagcgaggctggggcctttgggctggtgcagcccgcaatggaatgcgggaccgccaggctagc
agcaaaggcgcctcccctactccgcatcgatgttccatagtgcattggactgcatttgggtggggcggccggctgtttctttcgtgtt
gcaaaacgcgccagctcagcaacctgtcccgtgggtcccccgtgccgatgaaatcgtgtgcacgccgatcagctgattgcccggct
cgcgaagtaggcgccctcctttctgctcgccctctctccgtcccgcctctaga*atatcaATGATCGAGCAGGACGGCCTCC*
ACGCCGGCTCCCCCGCCGCCTGGGTGGAGCGCCTGTTCGGCTACGACTGGGCCCAGCAGACCATCG
GCTGCTCCGACGCCGCCGTGTTCCGCCTGTCCGCCCAGGGCCGCCCCGTGCTGTTCGTGAAGACCGA
CCTGTCCGGCGCCCTGAACGAGCTGCAGGACGAGGCCGCCCGCCTGTCCTGGCTGGCCACCACCGG
CGTGCCCTGCGCCGCCGTGCTGGACGTGGTGACCGAGGCCGGCCGCGACTGGCTGCTGCTGGGCG
AGGTGCCCGGCCAGGACCTGCTGTCCTCCCACCTGGCCCCCGCCGAGAAGGTGTCCATCATGGCCG
ACGCCATGCGCCGCCTGCACACCCTGGACCCCGCCACCTGCCCCTTCGACCACCAGGCCAAGCACCG
CATCGAGCGCGCCCGCACCCGCATGGAGGCCGGCCTGGTGGACCAGGACGACCTGGACGAGGAGC
ACCAGGGCCTGGCCCCCGCCGAGCTGTTCGCCCGCCTGAAGGCCCGCATGCCCGACGGCGAGGACC
TGGTGGTGACCCACGGCGACGCCTGCCTGCCCAACATCATGGTGGAGAACGGCCGCTTCTCCGGCT
TCATCGACTGCGGCCGCCTGGGCGTGGCCGACCGCTACCAGGACATCGCCCTGGCCACCCGCGACA
TCGCCGAGGAGCTGGGCGGCGAGTGGGCCGACCGCTTCCTGGTGCTGTACGGCATCGCCGCCCCCG
ACTCCCAGCGCATCGCCTTCTACCGCCTGCTGGACGAGTTCTTCTGA*caattggcagcagcagctcggatagt*
atcgacacactctggacgctggtcgtgtgatggactgttgccgccacacttgctgccttgacctgtgaatatccctgccgcttttat
caaacagcctcagtgtgtttgatcttgtgtgtacgcgcttttgcgagttgctagctgcttgtgctatttgcgaataccacccccagc
atccccttccctcgtttcatatcgcttgcatcccaaccgcaacttatctacgctgtcctgctatccctcagcgctgctcctgctcctgct
cactgcccctcgcacagccttggtttgggctccgcctgtattctcctggtactgcaacctgtaaaccagcactgcaatgctgatgc
*acgggaagtagtgggatgggaacacaaatgg*aggatcc*cgcgtctcgaacagagcgcgcagaggaacgctgaaggtctcgcc*
tctgtcgcacctcagcgcggcatacaccacaataaccacctgacgaatgcgcttggttcttcgtccattagcgaagcgtccggttcac
acacgtgccacgttggcgaggtggcaggtgacaatgatcggtggagctgatggtcgaaacgttcacagcctagggatatcgaattc
ctttcttgcgctatgacacttccagcaaaaggtagggcgggctgcgagacggcttcccggcgctgcatgcaacaccgatgatgctt
cgacccccgaagctccttcggggctgcatgggcgctccgatgccgctccagggcgagcgctgtttaaatagccaggccccgatt
gcaaagacattatagcgagctaccaaagccatattcaaacacctagatcactaccacttctacacaggccactcgagcttgtgatc
gcactccgctaaggggcgcctcttcctcttcgtttcagtcacaacccgcaaacg*ggcgcgcc***ATG**GTGGCCGCCGCCGCCT
CCAGCGCCTTCTTCCCCGTGCCCGCCCCCGGCGCCTCCCCCAAGCCCGGCAAGTTCGGCAACTGGCC
CTCCAGCCTGAGCCCCTCCTTCAAGCCCAAGTCCATCCCCAACGGCGGCTTCCAGGTGAAGGCCAAC
GACAGCGCCCACCCCAAGGCCAACGGCTCCGCCGTGAGCCTGAAGAGCGGCAGCCTGAACACCCAG
GAGGACACCTCCTCCAGCCCCCCCCCCCGCACCTTCCTGCACCAGCTGCCCGACTGGAGCCGCCTGCT
GACCGCCATCACCACCGTGTTCGTGAAGTCCAAGCGCCCCGACATGCACGACCGCAAGTCCAAGCG
CCCCGACATGCTGGTGGACAGCTTCGGCCTGGAGTCCACCGTGCAGGACGGCCTGGTGTTCCGCCA
GTCCTTCTCCATCCGCTCCTACGAGATCGGCACCGACCGCACCGCCAGCATCGAGACCCTGATGAAC
CACCTGCAGGAGACCTCCCTGAACCACTGCAAGAGCACCGGCATCCTGCTGGACGGCTTCGGCCGC
ACCCTGGAGATGTGCAAGCGCGACCTGATCTGGGTGGTGATCAAGATGCAGATCAAGGTGAACCGC
TACCCCGCCTGGGGCGACACCGTGGAGATCAACACCCGCTTCAGCCGCCTGGGCAAGATCGGCATG
GGCCGCGACTGGCTGATCTCCGACTGCAACACCGGCGAGATCCTGGTGCGCGCCACCAGCGCCTAC
GCCATGATGAACCAGAAGACCCGCCGCCTGTCCAAGCTGCCCTACGAGGTGCACCAGGAGATCGTG
CCCCTGTTCGTGGACAGCCCCGTGATCGAGGACTCCGACCTGAAGGTGCACAAGTTCAAGGTGAAG
ACCGGCGACAGCATCCAGAAGGGCCTGACCCCCGGCTGGAACGACCTGGACGTGAACCAGCACGT
GTCCAACGTGAAGTACATCGGCTGGATCCTGGAGAGCATGCCCACCGAGGTGCTGGAGACCCAGGA
GCTGTGCTCCCTGGCCCTGGAGTACCGCCGCGAGTGCGGCCGCGACTCCGTGCTGGAGAGCGTGAC
CGCCATGGACCCCAGCAAGGTGGGCGTGCGCTCCCAGTACCAGCACCTGCTGCGCCTGGAGGACGG
CACCGCCATCGTGAACGGCGCCACCGAGTGGCGCCCCAAGAACGCCGGCGCCAACGGCGCCATCTC
CACCGGCAAGACCAGCAACGGCAACTCCGTGTCC*ATGGACTACAAGGACCACGACGGCGACTACAA*

*GGACCACGACATCGACTACAAGGACGACGACGACAAG**TGA**ctcgaggcagcagcagctcggatagtatcgac*
acactctggacgctggtcgtgtgatggactgttgccgccacacttgctgccttgacctgtgaatatccctgccgcttttatcaaac
agcctcagtgtgtttgatcttgtgtgtacgcgcttttgcgagttgctagctgcttgtgctatttgcgaataccacccccagcatcccc
ttccctcgtttcatatcgcttgcatcccaaccgcaacttatctacgctgtcctgctatccctcagcgctgctcctgctcctgctcactg
cccctcgcacagccttggtttgggctccgcctgtattctcctggtactgcaacctgtaaaccagcactgcaatgctgatgcacggg
*aagtagtgggatgggaacacaaatgga*aagctt

SEQ ID NO: 114
C.reinhartii β-tubulin promoter::*P.moriformis* isopentenyl diphosphate synthase PTS::*Cuphea hookeriana* C8-10 thioesterase::*C.vulgaris* nitrate reductase 3'UTR

ggtacc**cgcctgcaacgcaagggcagccacagccgctcccacccgccgctgaaccgacacgtgcttgggcgcctgccgcctgcctg**
ccgcatgcttgtgctggtgaggctgggcagtgctgccatgctgattgaggcttggttcatcgggtggaagcttatgtgtgtgctggg
cttgcatgccgggcaatgcgcatggtggcaagagggcggcagcacttgctggagctgccgcggtgcctccaggtggttcaatcgc
ggcagccagagggatttcagatgatcgcgcgtacaggttgagcagcagtgtcagcaaaggtagcagtttgccagaatgatcggtt
cagctgttaatcaatgccagcaagagaaggggtcaagtgcaaacacgggcatgccacagcacgggcaccggggagtggaatgg
caccaccaagtgtgtgcgagccagcatcgccgcctggctgtttcagctacaacggcaggagtcatccaacgtaaccatgagctga
tcaacactgcaatcatcgggcgggcgtgatgcaagcatgcctggcgaagacacatggtgtgcggatgctgccggctgctgcctgct
gcgcacgccgttgagttggcagcaggctcagccatgcactggatggcagctgggctgccactgcaatgtggtggataggatgcaa
gtggagcgaataccaaaccctctggctgcttgctgggttgcatggcatcgcaccatcagcaggagcgcatgcgaagggactggcc
ccatgcacgccatgccaaaccggagcgcaccgagtgtccacactgtcaccaggcccgcaagctttgcagaaccatgctcatggac
gcatgtagcgctgacgtcccttgacggcgctcctctcgggtgtgggaaacgcaatgcagcacaggcagcagaggcggcggcagc
agagcggcggcagcagcggcgggggccacccttcttgcggggtcgcgccccagccagcggtgatgcgctgatcccaaacgagttc
acattcatttgcatgcctggagaagcgaggctggggcctttgggctggtgcagcccgcaatggaatgcgggaccgccaggctagc
agcaaaggcgcctcccctactccgcatcgatgttccatagtgcattggactgcatttgggtggggcggccggctgtttctttcgtgtt
gcaaaacgcgccagctcagcaacctgtcccgtgggtcccccgtgccgatgaaatcgtgtgcacgccgatcagctgattgcccggct
cgcgaagtaggcgccctcctttctgctcgccctctctccgtcccgcctctaga*atatcaATGATCGAGCAGGACGGCCTCC*
ACGCCGGCTCCCCCGCCGCCTGGGTGGAGCGCCTGTTCGGCTACGACTGGGCCCAGCAGACCATCG
GCTGCTCCGACGCCGCCGTGTTCCGCCTGTCCGCCCAGGGCCGCCCCGTGCTGTTCGTGAAGACCGA
CCTGTCCGGCGCCCTGAACGAGCTGCAGGACGAGGCCGCCCGCCTGTCCTGGCTGGCCACCACCGG
CGTGCCCTGCGCCGCCGTGCTGGACGTGGTGACCGAGGCCGGCCGCGACTGGCTGCTGCTGGGCG
AGGTGCCCGGCCAGGACCTGCTGTCCTCCCACCTGGCCCCCGCCGAGAAGGTGTCCATCATGGCCG
ACGCCATGCGCCGCCTGCACACCCTGGACCCCGCCACCTGCCCCTTCGACCACCAGGCCAAGCACCG
CATCGAGCGCGCCCGCACCCGCATGGAGGCCGGCCTGGTGGACCAGGACGACCTGGACGAGGAGC
ACCAGGGCCTGGCCCCCGCCGAGCTGTTCGCCCGCCTGAAGGCCCGCATGCCCGACGGCGAGGACC
TGGTGGTGACCCACGGCGACGCCTGCCTGCCCAACATCATGGTGGAGAACGGCCGCTTCTCCGGCT
TCATCGACTGCGGCCGCCTGGGCGTGGCCGACCGCTACCAGGACATCGCCCTGGCCACCCGCGACA
TCGCCGAGGAGCTGGGCGGCGAGTGGGCCGACCGCTTCCTGGTGCTGTACGGCATCGCCGCCCCCG
ACTCCCAGCGCATCGCCTTCTACCGCCTGCTGGACGAGTTCTTCTGA*caattggcagcagcagctcggatagt*
atcgacacactctggacgctggtcgtgtgatggactgttgccgccacacttgctgccttgacctgtgaatatccctgccgcttttat
caaacagcctcagtgtgtttgatcttgtgtgtacgcgcttttgcgagttgctagctgcttgtgctatttgcgaataccacccccagc
atcccttccctcgtttcatatcgcttgcatcccaaccgcaacttatctacgctgtcctgctatccctcagcgctgctcctgctcctgct
cactgcccctcgcacagccttggtttgggctccgcctgtattctcctggtactgcaacctgtaaaccagcactgcaatgctgatgc
*acgggaagtagtgggatgggaacacaaatgga*ggatccgcgtctcgaacagagcgcgcagaggaacgctgaaggtctcgcc
tctgtcgcacctcagcgcggcatacaccacaataaccacctgacgaatgcgcttggttcttcgtccattagcgaagcgtccggttcac
acacgtgccacgttggcgaggtggcaggtgacaatgatcggtggagctgatggtcgaaacgttcacagcctagggatatcgaattc
ctttcttgcgctatgacacttccagcaaaaggtagggcgggctgcgagacggcttcccggcgctgcatgcaacaccgatgatgctt

cgacccccgaagctccttcggggctgcatgggcgctccgatgccgctccagggcgagcgctgtttaaatagccaggcccccgatt
gcaaagacattatagcgagctaccaaagccatattcaaacacctagatcactaccacttctacacaggccactcgagcttgtgatc
gcactccgctaagggggcgcctcttcctcttcgtttcagtcacaacccgcaaacactagt**ATG**ACGTTCGGGGTCGCCCTC
CCGGCCATGGGCCGCGGTGTCTCCCTTCCCCGGCCCAGGGTCGCGGTGCGCGCCCAGTCGGCGAGT
CAGGTTTTGGAGAGCGGGCGCGCCCAGCTGCCCGACTGGAGCCGCCTGCTGACCGCCATCACCACC
GTGTTCGTGAAGTCCAAGCGCCCCGACATGCACGACCGCAAGTCCAAGCGCCCCGACATGCTGGTG
GACAGCTTCGGCCTGGAGTCCACCGTGCAGGACGGCCTGGTGTTCCGCCAGTCCTTCTCCATCCGCT
CCTACGAGATCGGCACCGACCGCACCGCCAGCATCGAGACCCTGATGAACCACCTGCAGGAGACCT
CCCTGAACCACTGCAAGAGCACCGGCATCCTGCTGGACGGCTTCGGCCGCACCCTGGAGATGTGCA
AGCGCGACCTGATCTGGGTGGTGATCAAGATGCAGATCAAGGTGAACCGCTACCCCGCCTGGGGCG
ACACCGTGGAGATCAACACCCGCTTCAGCCGCCTGGGCAAGATCGGCATGGGCCGCGACTGGCTGA
TCTCCGACTGCAACACCGGCGAGATCCTGGTGCGCGCCACCAGCGCCTACGCCATGATGAACCAGA
AGACCCGCCGCCTGTCCAAGCTGCCCTACGAGGTGCACCAGGAGATCGTGCCCCTGTTCGTGGACA
GCCCCGTGATCGAGGACTCCGACCTGAAGGTGCACAAGTTCAAGGTGAAGACCGGCGACAGCATCC
AGAAGGGCCTGACCCCCGGCTGGAACGACCTGGACGTGAACCAGCACGTGTCCAACGTGAAGTACA
TCGGCTGGATCCTGGAGAGCATGCCCACCGAGGTGCTGGAGACCCAGGAGCTGTGCTCCCTGGCCC
TGGAGTACCGCCGCGAGTGCGGCCGCGACTCCGTGCTGGAGAGCGTGACCGCCATGGACCCCAGC
AAGGTGGGCGTGCGCTCCCAGTACCAGCACCTGCTGCGCCTGGAGGACGGCACCGCCATCGTGAAC
GGCGCCACCGAGTGGCGCCCCAAGAACGCCGGCGCCAACGGCGCCATCTCCACCGGCAAGACCAG
CAACGGCAACTCCGTGTCC*ATGGACTACAAGGACCACGACGGCGACTACAAGGACCACGACATCGA*
*CTACAAGGACGACGACGACAAG**TGA**ctcgaggcagcagcagctcggatagtatcgacacactctggacgctggtcg*
tgtgatggactgttgccgccacacttgctgccttgacctgtgaatatccctgccgcttttatcaaacagcctcagtgtgtttgatctt
gtgtgtacgcgcttttgcgagttgctagctgcttgtgctatttgcgaataccacccccagcatcccctttccctcgtttcatatcgctt
gcatcccaaccgcaacttatctacgctgtcctgctatccctcagcgctgctcctgctcctgctcactgcccctcgcacagccttggtt
tgggctccgcctgtattctcctggtactgcaacctgtaaaccagcactgcaatgctgatgcacgggaagtagtgggatgggaa
*cacaaatgga*aagctt

SEQ ID NO: 115

C.reinhartii β-tubulin promoter::*P.moriformis* delta 12 fatty acid desaturase PTS::*Cuphea hookeriana* C8-10 thioesterase::*C.vulgaris* nitrate reductase 3'UTR

ggtacc**cgcctgcaacgcaagggcagccacagccgctcccacccgccgctgaaccgacacgtgcttgggcgcctgccgcctgcctg**
ccgcatgcttgtgctggtgaggctgggcagtgctgccatgctgattgaggcttggttcatcgggtggaagcttatgtgtgtgctggg
cttgcatgccgggcaatgcgcatggtggcaagagggcggcagcacttgctggagctgccgcggtgcctccaggtggttcaatcgc
ggcagccagagggatttcagatgatcgcgcgtacaggttgagcagcagtgtcagcaaaggtagcagtttgccagaatgatcggtt
cagctgttaatcaatgccagcaagagaaggggtcaagtgcaaacacgggcatgccacagcacgggcaccggggagtggaatgg
caccaccaagtgtgtgcgagccagcatcgccgcctggctgtttcagctacaacggcaggagtcatccaacgtaaccatgagctga
tcaacactgcaatcatcgggcgggcgtgatgcaagcatgcctggcgaagacacatggtgtgcggatgctgccggctgctgcctgct
gcgcacgccgttgagttggcagcaggctcagccatgcactggatggcagctgggctgccactgcaatgtggtggataggatgcaa
gtggagcgaataccaaaccctctggctgcttgctgggttgcatggcatcgcaccatcagcaggagcgcatgcgaagggactggcc
ccatgcacgccatgccaaaccggagcgcaccgagtgtccacactgtcaccaggcccgcaagctttgcagaaccatgctcatggac
gcatgtagcgctgacgtcccttgacggcgctcctctcgggtgtgggaaacgcaatgcagcacaggcagcagaggcggcggcagc
agagcggcggcagcagcggcgggggccacccttcttgcggggtcgcgccccagccagcggtgatgcgctgatcccaaacgagttc
acattcatttgcatgcctggagaagcgaggctggggccttgggctggtgcagcccgcaatggaatgcgggaccgccaggctagc
agcaaaggcgcctcccctactccgcatcgatgttccatagtgcattggactgcatttgggtggggcggccggctgtttctttcgtgtt
gcaaaacgcgccagctcagcaacctgtcccgtgggtcccccgtgccgatgaaatcgtgtgcacgccgatcagctgattgcccggct
cgcgaagtaggcgccctcctttctgctcgccctctctccgtcccgcctctaga**atatcaATGATCGAGCAGGACGGCCTCC**

ACGCCGGCTCCCCCGCCGCCTGGGTGGAGCGCCTGTTCGGCTACGACTGGGCCCAGCAGACCATCG
GCTGCTCCGACGCCGCCGTGTTCCGCCTGTCCGCCCAGGGCCGCCCCGTGCTGTTCGTGAAGACCGA
CCTGTCCGGCGCCCTGAACGAGCTGCAGGACGAGGCCGCCCGCCTGTCCTGGCTGGCCACCACCGG
CGTGCCCTGCGCCGCCGTGCTGGACGTGGTGACCGAGGCCGGCCGCGACTGGCTGCTGCTGGGCG
AGGTGCCCGGCCAGGACCTGCTGTCCTCCCACCTGGCCCCCGCCGAGAAGGTGTCCATCATGGCCG
ACGCCATGCGCCGCCTGCACACCCTGGACCCCGCCACCTGCCCCTTCGACCACCAGGCCAAGCACCG
CATCGAGCGCGCCCGCACCCGCATGGAGGCCGGCCTGGTGGACCAGGACGACCTGGACGAGGAGC
ACCAGGGCCTGGCCCCCGCCGAGCTGTTCGCCCGCCTGAAGGCCCGCATGCCCGACGGCGAGGACC
TGGTGGTGACCCACGGCGACGCCTGCCTGCCCAACATCATGGTGGAGAACGGCCGCTTCTCCGGCT
TCATCGACTGCGGCCGCCTGGGCGTGGCCGACCGCTACCAGGACATCGCCCTGGCCACCCGCGACA
TCGCCGAGGAGCTGGGCGGCGAGTGGGCCGACCGCTTCCTGGTGCTGTACGGCATCGCCGCCCCCG
ACTCCCAGCGCATCGCCTTCTACCGCCTGCTGGACGAGTTCTTCTGA*caattggcagcagcagctcggatagt*
atcgacacactctggacgctggtcgtgtgatggactgttgccgccacacttgctgccttgacctgtgaatatccctgccgcttttat
caaacagcctcagtgtgtttgatcttgtgtgtacgcgcttttgcgagttgctagctgcttgtgctatttgcgaataccacccccagc
atccccttccctcgtttcatatcgcttgcatcccaaccgcaacttatctacgctgtcctgctatccctcagcgctgctcctgctcctgct
cactgcccctcgcacagccttggtttgggctccgcctgtattctcctggtactgcaacctgtaaaccagcactgcaatgctgatgc
*acgggaagtagtgggatgggaacacaaatgga*ggatccgcgtctcgaacagagcgcgcagaggaacgctgaaggtctcgcc
tctgtcgcacctcagcgcggcatacaccacaataaccacctgacgaatgcgcttggttcttcgtccattagcgaagcgtccggttcac
acacgtgccacgttggcgaggtggcaggtgacaatgatcggtggagctgatggtcgaaacgttcacagcctagggatatcgaattc
ctttcttgcgctatgacacttccagcaaaaggtagggcgggctgcgagacggcttcccggcgctgcatgcaacaccgatgatgctt
cgaccccccgaagctccttcggggctgcatgggcgctccgatgccgctccagggcgagcgctgtttaaatagccaggcccccgatt
gcaaagacattatagcgagctaccaaagccatattcaaacacctagatcactaccacttctacacaggccactcgagcttgtgatc
gcactccgctaaggggcgcctcttcctcttcgtttcagtcacaacccgcaaacactagt**ATG**GCTATCAAGACGAACAG
GCAGCCTGTGGAGAAGCCTCCGTTCACGATCGGGACGCTGCGCAAGGCCATCCCCGCGCACTGTTT
CGAGCGCTCGGCGCTTCGTGGGCGCGCCCAGCTGCCCGACTGGAGCCGCCTGCTGACCGCCATCAC
CACCGTGTTCGTGAAGTCCAAGCGCCCCGACATGCACGACCGCAAGTCCAAGCGCCCCGACATGCT
GGTGGACAGCTTCGGCCTGGAGTCCACCGTGCAGGACGGCCTGGTGTTCCGCCAGTCCTTCTCCATC
CGCTCCTACGAGATCGGCACCGACCGCACCGCCAGCATCGAGACCCTGATGAACCACCTGCAGGAG
ACCTCCCTGAACCACTGCAAGAGCACCGGCATCCTGCTGGACGGCTTCGGCCGCACCCTGGAGATGT
GCAAGCGCGACCTGATCTGGGTGGTGATCAAGATGCAGATCAAGGTGAACCGCTACCCCGCCTGGG
GCGACACCGTGGAGATCAACACCCGCTTCAGCCGCCTGGGCAAGATCGGCATGGGCCGCGACTGGC
TGATCTCCGACTGCAACACCGGCGAGATCCTGGTGCGCGCCACCAGCGCCTACGCCATGATGAACCA
GAAGACCCGCCGCCTGTCCAAGCTGCCCTACGAGGTGCACCAGGAGATCGTGCCCCTGTTCGTGGA
CAGCCCCGTGATCGAGGACTCCGACCTGAAGGTGCACAAGTTCAAGGTGAAGACCGGCGACAGCAT
CCAGAAGGGCCTGACCCCCGGCTGGAACGACCTGGACGTGAACCAGCACGTGTCCAACGTGAAGTA
CATCGGCTGGATCCTGGAGAGCATGCCCACCGAGGTGCTGGAGACCCAGGAGCTGTGCTCCCTGGC
CCTGGAGTACCGCCGCGAGTGCGGCCGCGACTCCGTGCTGGAGAGCGTGACCGCCATGGACCCCA
GCAAGGTGGGCGTGCGCTCCCAGTACCAGCACCTGCTGCGCCTGGAGGACGGCACCGCCATCGTGA
ACGGCGCCACCGAGTGGCGCCCCAAGAACGCCGGCGCCAACGGCGCCATCTCCACCGGCAAGACCA
GCAACGGCAACTCCGTGTC*CATGGACTACAAGGACCACGACGGCGACTACAAGGACCACGACATCG*
*ACTACAAGGACGACGACGACAAG**TGA**ctcgaggcagcagcagctcggatagtatcgacacactctggacgctggtc*
gtgtgatggactgttgccgccacacttgctgccttgacctgtgaatatccctgccgcttttatcaaacagcctcagtgtgtttgatc
ttgtgtgtacgcgcttttgcgagttgctagctgcttgtgctatttgcgaataccacccccagcatcccctttccctcgtttcatatcgct
tgcatcccaaccgcaacttatctacgctgtcctgctatccctcagcgctgctcctgctcctgctcactgcccctcgcacagccttggt
ttgggctccgcctgtattctcctggtactgcaacctgtaaaccagcactgcaatgctgatgcacgggaagtagtgggatggga
*acacaaatgga*aagctt

SEQ ID NO: 116

C.reinhartii β-tubulin promoter::*C.protothecoides* stearoyl ACP desaturase PTS::*Cuphea hookeriana* C8-10 thioesterase::*C.vulgaris* nitrate reductase 3'UTR

ggtaccgcctgcaacgcaagggcagccacagccgctcccacccgccgctgaaccgacacgtgcttgggcgcctgccgcctgcctg
ccgcatgcttgtgctggtgaggctgggcagtgctgccatgctgattgaggcttggttcatcgggtggaagcttatgtgtgtgctggg
cttgcatgccgggcaatgcgcatggtggcaagagggcggcagcacttgctggagctgccgcggtgcctccaggtggttcaatcgc
ggcagccagagggatttcagatgatcgcgcgtacaggttgagcagcagtgtcagcaaaggtagcagtttgccagaatgatcggtt
cagctgttaatcaatgccagcaagagaaggggtcaagtgcaaacacgggcatgccacagcacgggcaccggggagtggaatgg
caccaccaagtgtgtgcgagccagcatcgccgcctggctgtttcagctacaacggcaggagtcatccaacgtaaccatgagctga
tcaacactgcaatcatcgggcgggcgtgatgcaagcatgcctggcgaagacacatggtgtgcggatgctgccggctgctgcctgct
gcgcacgccgttgagttggcagcaggctcagccatgcactggatggcagctgggctgccactgcaatgtggtggataggatgcaa
gtggagcgaataccaaaccctctggctgcttgctgggttgcatggcatcgcaccatcagcaggagcgcatgcgaagggactggcc
ccatgcacgccatgccaaaccggagcgcaccgagtgtccacactgtcaccaggcccgcaagctttgcagaaccatgctcatggac
gcatgtagcgctgacgtcccttgacggcgctcctctcgggtgtgggaaacgcaatgcagcacaggcagcagaggcggcggcagc
agagcggcggcagcagcggcgggggccacccttcttgcggggtcgcgccccagccagcggtgatgcgctgatcccaaacgagttc
acattcatttgcatgcctggagaagcgaggctggggcctttgggctggtgcagcccgcaatggaatgcgggaccgccaggctagc
agcaaaggcgcctcccctactccgcatcgatgttccatagtgcattggactgcatttgggtggggcggccggctgtttctttcgtgtt
gcaaaacgcgccagctcagcaacctgtcccgtgggtcccccgtgccgatgaaatcgtgtgcacgccgatcagctgattgcccggct
cgcgaagtaggcgccctcctttctgctcgccctctctccgtcccgcctctagaatatcaATGATCGAGCAGGACGGCCTCC
ACGCCGGCTCCCCCGCCGCCTGGGTGGAGCGCCTGTTCGGCTACGACTGGGCCCAGCAGACCATCG
GCTGCTCCGACGCCGCCGTGTTCCGCCTGTCCGCCCAGGGCCGCCCCGTGCTGTTCGTGAAGACCGA
CCTGTCCGGCGCCCTGAACGAGCTGCAGGACGAGGCCGCCCGCCTGTCCTGGCTGGCCACCACCGG
CGTGCCCTGCGCCGCCGTGCTGGACGTGGTGACCGAGGCCGGCCGCGACTGGCTGCTGCTGGGCG
AGGTGCCCGGCCAGGACCTGCTGTCCTCCCACCTGGCCCCCGCCGAGAAGGTGTCCATCATGGCCG
ACGCCATGCGCCGCCTGCACACCCTGGACCCCGCCACCTGCCCCTTCGACCACCAGGCCAAGCACCG
CATCGAGCGCGCCCGCACCCGCATGGAGGCCGGCCTGGTGGACCAGGACGACCTGGACGAGGAGC
ACCAGGGCCTGGCCCCCGCCGAGCTGTTCGCCCGCCTGAAGGCCCGCATGCCCGACGGCGAGGACC
TGGTGGTGACCCACGGCGACGCCTGCCTGCCCAACATCATGGTGGAGAACGGCCGCTTCTCCGGCT
TCATCGACTGCGGCCGCCTGGGCGTGGCCGACCGCTACCAGGACATCGCCCTGGCCACCCGCGACA
TCGCCGAGGAGCTGGGCGGCGAGTGGGCCGACCGCTTCCTGGTGCTGTACGGCATCGCCGCCCCCG
ACTCCCAGCGCATCGCCTTCTACCGCCTGCTGGACGAGTTCTTCTGAcaattg*gcagcagcagctcggatagt*
atcgacacactctggacgctggtcgtgtgatggactgttgccgccacacttgctgccttgacctgtgaatatccctgccgcttttat
caaacagcctcagtgtgtttgatcttgtgtgtacgcgcttttgcgagttgctagctgcttgtgctatttgcgaataccacccccagc
atccccttccctcgtttcatatcgcttgcatcccaaccgcaacttatctacgctgtcctgctatccctcagcgctgctcctgctcctgct
cactgcccctcgcacagccttggtttgggctccgcctgtattctcctggtactgcaacctgtaaaccagcactgcaatgctgatgc
*acgggaagtagtgggatgggaacacaaatgga*ggatccgcgtctcgaacagagcgcgcagaggaacgctgaaggtctcgcc
tctgtcgcacctcagcgcggcatacaccacaataaccacctgacgaatgcgcttggttcttcgtccattagcgaagcgtccggttcac
acacgtgccacgttggcgaggtggcaggtgacaatgatcggtggagctgatggtcgaaacgttcacagcctagggatatcgaattc
ctttcttgcgctatgacacttccagcaaaaggtagggcgggctgcgagacggcttcccggcgctgcatgcaacaccgatgatgctt
cgaccccccgaagctccttcggggctgcatgggcgctccgatgccgctccagggcgagcgctgtttaaatagccaggccccgatt
gcaaagacattatagcgagctaccaaagccatattcaaacacctagatcactaccacttctacacaggccactcgagcttgtgatc
gcactccgctaaggggggcgcctcttcctcttcgtttcagtcacaacccgcaaacactagt**ATG**GCCACCGCATCCACTTTCT
CGGCGTTCAATGCCCGCTGCGGCGACCTGCGTCGCTCGGCGGGCTCCGGGCCCCGGCGCCCAGCGA
GGCCCCTCCCCGTGCGCGGGCGCGCCCAGCTGCCCGACTGGAGCCGCCTGCTGACCGCCATCACCA
CCGTGTTCGTGAAGTCCAAGCGCCCCGACATGCACGACCGCAAGTCCAAGCGCCCCGACATGCTGG
TGGACAGCTTCGGCCTGGAGTCCACCGTGCAGGACGGCCTGGTGTTCCGCCAGTCCTTCTCCATCCG

CTCCTACGAGATCGGCACCGACCGCACCGCCAGCATCGAGACCCTGATGAACCACCTGCAGGAGAC
CTCCCTGAACCACTGCAAGAGCACCGGCATCCTGCTGGACGGCTTCGGCCGCACCCTGGAGATGTGC
AAGCGCGACCTGATCTGGGTGGTGATCAAGATGCAGATCAAGGTGAACCGCTACCCCGCCTGGGGC
GACACCGTGGAGATCAACACCCGCTTCAGCCGCCTGGGCAAGATCGGCATGGGCCGCGACTGGCTG
ATCTCCGACTGCAACACCGGCGAGATCCTGGTGCGCGCCACCAGCGCCTACGCCATGATGAACCAG
AAGACCCGCCGCCTGTCCAAGCTGCCCTACGAGGTGCACCAGGAGATCGTGCCCCTGTTCGTGGAC
AGCCCCGTGATCGAGGACTCCGACCTGAAGGTGCACAAGTTCAAGGTGAAGACCGGCGACAGCATC
CAGAAGGGCCTGACCCCCGGCTGGAACGACCTGGACGTGAACCAGCACGTGTCCAACGTGAAGTAC
ATCGGCTGGATCCTGGAGAGCATGCCCACCGAGGTGCTGGAGACCCAGGAGCTGTGCTCCCTGGCC
CTGGAGTACCGCCGCGAGTGCGGCCGCGACTCCGTGCTGGAGAGCGTGACCGCCATGGACCCCAG
CAAGGTGGGCGTGCGCTCCCAGTACCAGCACCTGCTGCGCCTGGAGGACGGCACCGCCATCGTGAA
CGGCGCCACCGAGTGGCGCCCCAAGAACGCCGGCGCCAACGGCGCCATCTCCACCGGCAAGACCA
GCAACGGCAACTCCGTGTCC*ATGGACTACAAGGACCACGACGGCGACTACAAGGACCACGACATCG*
*ACTACAAGGACGACGACGACAAG**TGA***ctcgagg*cagcagcagctcggatagtatcgacacactctggacgctggtc*
gtgtgatggactgttgccgccacacttgctgccttgacctgtgaatatccctgccgcttttatcaaacagcctcagtgtgtttgatc
ttgtgtgtacgcgcttttgcgagttgctagctgcttgtgctatttgcgaataccacccccagcatcccctttccctcgtttcatatcgct
tgcatcccaaccgcaacttatctacgctgtcctgctatccctcagcgctgctcctgctcctgctcactgcccctcgcacagccttggt
ttgggctccgcctgtattctcctggtactgcaacctgtaaaccagcactgcaatgctgatgcacgggaagtagtgggatggga
*acacaaatgga*aagctt

SEQ ID NO: 117

C.reinhartii β-tubulin promoter::*P.moriformis* stearoyl ACP desaturase PTS::*Cuphea hookeriana* C8-10 thioesterase::*C.vulgaris* nitrate reductase 3'UTR

ggtacc**cgcctgcaacgcaagggcagccacagccgctcccacccgccgctgaaccgacacgtgcttgggcgcctgccgcctgcctg**
ccgcatgcttgtgctggtgaggctgggcagtgctgccatgctgattgaggcttggttcatcgggtggaagcttatgtgtgtgctggg
cttgcatgccgggcaatgcgcatggtggcaagagggcggcagcacttgctggagctgccgcggtgcctccaggtggttcaatcgc
ggcagccagagggatttcagatgatcgcgcgtacaggttgagcagcagtgtcagcaaaggtagcagtttgccagaatgatcggtt
cagctgttaatcaatgccagcaagagaaggggtcaagtgcaaacacgggcatgccacagcacgggcaccggggagtggaatgg
caccaccaagtgtgtgcgagccagcatcgccgcctggctgtttcagctacaacggcaggagtcatccaacgtaaccatgagctga
tcaacactgcaatcatcgggcgggcgtgatgcaagcatgcctggcgaagacacatggtgtgcggatgctgccggctgctgcctgct
gcgcacgccgttgagttggcagcaggctcagccatgcactggatggcagctgggctgccactgcaatgtggtggataggatgcaa
gtggagcgaataccaaaccctctggctgcttgctgggttgcatggcatcgcaccatcagcaggagcgcatgcgaagggactggcc
ccatgcacgccatgccaaaccggagcgcaccgagtgtccacactgtcaccaggcccgcaagctttgcagaaccatgctcatggac
gcatgtagcgctgacgtcccttgacggcgctcctctcgggtgtgggaaacgcaatgcagcacaggcagcagaggcggcggcagc
agagcggcggcagcagcggcgggggccacccttcttgcggggtcgcgccccagccagcggtgatgcgctgatcccaaacgagttc
acattcatttgcatgcctggagaagcgaggctggggcctttgggctggtgcagcccgcaatggaatgcgggaccgccaggctagc
agcaaaggcgcctcccctactccgcatcgatgttccatagtgcattggactgcatttgggtggggcggccggctgtttctttcgtgtt
gcaaaacgcgccagctcagcaacctgtcccgtgggtcccccgtgccgatgaaatcgtgtgcacgccgatcagctgattgcccggct
cgcgaagtaggcgccctcctttctgctcgccctctctccgtcccgcctctaga*atatca*ATGATCGAGCAGGACGGCCTCC
ACGCCGGCTCCCCCGCCGCCTGGGTGGAGCGCCTGTTCGGCTACGACTGGGCCCAGCAGACCATCG
GCTGCTCCGACGCCGCCGTGTTCCGCCTGTCCGCCCAGGGCCGCCCCGTGCTGTTCGTGAAGACCGA
CCTGTCCGGCGCCCTGAACGAGCTGCAGGACGAGGCCGCCCGCCTGTCCTGGCTGGCCACCACCGG
CGTGCCCTGCGCCGCCGTGCTGGACGTGGTGACCGAGGCCGGCCGCGACTGGCTGCTGCTGGGCG
AGGTGCCCGGCCAGGACCTGCTGTCCTCCCACCTGGCCCCCGCCGAGAAGGTGTCCATCATGGCCG
ACGCCATGCGCCGCCTGCACACCCTGGACCCCGCCACCTGCCCCTTCGACCACCAGGCCAAGCACCG
CATCGAGCGCGCCCGCACCCGCATGGAGGCCGGCCTGGTGGACCAGGACGACCTGGACGAGGAGC

ACCAGGGCCTGGCCCCCGCCGAGCTGTTCGCCCGCCTGAAGGCCCGCATGCCCGACGGCGAGGACC
TGGTGGTGACCCACGGCGACGCCTGCCTGCCCAACATCATGGTGGAGAACGGCCGCTTCTCCGGCT
TCATCGACTGCGGCCGCCTGGGCGTGGCCGACCGCTACCAGGACATCGCCCTGGCCACCCGCGACA
TCGCCGAGGAGCTGGGCGGCGAGTGGGCCGACCGCTTCCTGGTGCTGTACGGCATCGCCGCCCCCG
ACTCCCAGCGCATCGCCTTCTACCGCCTGCTGGACGAGTTCTTCTGAcaattg*gcagcagcagctcggatagt*
atcgacacactctggacgctggtcgtgtgatggactgttgccgccacacttgctgccttgacctgtgaatatccctgccgcttttat
caaacagcctcagtgtgtttgatcttgtgtgtacgcgcttttgcgagttgctagctgcttgtgctatttgcgaataccacccccagc
atccccttccctcgtttcatatcgcttgcatcccaaccgcaacttatctacgctgtcctgctatccctcagcgctgctcctgctcctgct
cactgcccctcgcacagccttggtttgggctccgcctgtattctcctggtactgcaacctgtaaaccagcactgcaatgctgatgc
*acgggaagtagtgggatgggaacacaaatgga*ggatccgcgtctcgaacagagcgcgcagaggaacgctgaaggtctcgcc
tctgtcgcacctcagcgcggcatacaccacaataaccacctgacgaatgcgcttggttcttcgtccattagcgaagcgtccggttcac
acacgtgccacgttggcgaggtggcaggtgacaatgatcggtggagctgatggtcgaaacgttcacagcctagggatatcgaattc
ctttcttgcgctatgacacttccagcaaaaggtagggcgggctgcgagacggcttcccggcgctgcatgcaacaccgatgatgctt
cgacccccgaagctccttcggggctgcatgggcgctccgatgccgctccagggcgagcgctgtttaaatagccaggcccccgatt
gcaaagacattatagcgagctaccaaagccatattcaaacacctagatcactaccacttctacacaggccactcgagcttgtgatc
gcactccgctaaggggcgcctcttcctcttcgtttcagtcacaacccgcaaacactagt**ATG**GCTTCCGCGGCATTCACC
ATGTCGGCGTGCCCCGCGATGACTGGCAGGGCCCCTGGGGCACGTCGCTCCGGACGGCCAGTCGCC
ACCCGCCTGAGGGGGCGCGCCCCCGACTGGAGCCGCCTGCTGACCGCCATCACCACCGTGTTCGTG
AAGTCCAAGCGCCCCGACATGCACGACCGCAAGTCCAAGCGCCCCGACATGCTGGTGGACAGCTTC
GGCCTGGAGTCCACCGTGCAGGACGGCCTGGTGTTCCGCCAGTCCTTCTCCATCCGCTCCTACGAGA
TCGGCACCGACCGCACCGCCAGCATCGAGACCCTGATGAACCACCTGCAGGAGACCTCCCTGAACC
ACTGCAAGAGCACCGGCATCCTGCTGGACGGCTTCGGCCGCACCCTGGAGATGTGCAAGCGCGACC
TGATCTGGGTGGTGATCAAGATGCAGATCAAGGTGAACCGCTACCCCGCCTGGGGCGACACCGTGG
AGATCAACACCCGCTTCAGCCGCCTGGGCAAGATCGGCATGGGCCGCGACTGGCTGATCTCCGACT
GCAACACCGGCGAGATCCTGGTGCGCGCCACCAGCGCCTACGCCATGATGAACCAGAAGACCCGCC
GCCTGTCCAAGCTGCCCTACGAGGTGCACCAGGAGATCGTGCCCCTGTTCGTGGACAGCCCCGTGAT
CGAGGACTCCGACCTGAAGGTGCACAAGTTCAAGGTGAAGACCGGCGACAGCATCCAGAAGGGCC
TGACCCCCGGCTGGAACGACCTGGACGTGAACCAGCACGTGTCCAACGTGAAGTACATCGGCTGGA
TCCTGGAGAGCATGCCCACCGAGGTGCTGGAGACCCAGGAGCTGTGCTCCCTGGCCCTGGAGTACC
GCCGCGAGTGCGGCCGCGACTCCGTGCTGGAGAGCGTGACCGCCATGGACCCCAGCAAGGTGGGC
GTGCGCTCCCAGTACCAGCACCTGCTGCGCCTGGAGGACGGCACCGCCATCGTGAACGGCGCCACC
GAGTGGCGCCCCAAGAACGCCGGCGCCAACGGCGCCATCTCCACCGGCAAGACCAGCAACGGCAA
CTCCGTGTCC*ATGGACTACAAGGACCACGACGGCGACTACAAGGACCACGACATCGACTACAAGGA*
*CGACGACGACAAG***TGA**ctcgag*gcagcagcagctcggatagtatcgacacactctggacgctggtcgtgtgatggactg*
ttgccgccacacttgctgccttgacctgtgaatatccctgccgcttttatcaaacagcctcagtgtgtttgatcttgtgtgtacgcgc
ttttgcgagttgctagctgcttgtgctatttgcgaataccacccccagcatccccttccctcgtttcatatcgcttgcatcccaaccgc
aacttatctacgctgtcctgctatccctcagcgctgctcctgctcctgctcactgcccctcgcacagccttggtttgggctccgcctgt
*attctcctggtactgcaacctgtaaaccagcactgcaatgctgatgcacgggaagtagtgggatgggaacacaaatgga*aaag
ctt

SEQ ID NO: 118

C.reinhartii β-tubulin promoter::native PTS::*Umbellularia californica* C12 thioesterase::*C.vulgaris* nitrate reductase 3'UTR

ggtacc**cgcctgcaacgcaagggcagccacagccgctcccacccgccgctgaaccgacacgtgcttgggcgcctgccgcctgcctg**
ccgcatgcttgtgctggtgaggctgggcagtgctgccatgctgattgaggcttggttcatcgggtggaagcttatgtgtgtgctggg
cttgcatgccgggcaatgcgcatggtggcaagagggcggcagcacttgctggagctgccgcggtgcctccaggtggttcaatcgc
ggcagccagagggatttcagatgatcgcgcgtacaggttgagcagcagtgtcagcaaaggtagcagtttgccagaatgatcggtt

cagctgttaatcaatgccagcaagagaaggggtcaagtgcaaacacgggcatgccacagcacgggcaccggggagtggaatgg
caccaccaagtgtgtgcgagccagcatcgccgcctggctgtttcagctacaacggcaggagtcatccaacgtaaccatgagctga
tcaacactgcaatcatcgggcgggcgtgatgcaagcatgcctggcgaagacacatggtgtgcggatgctgccggctgctgcctgct
gcgcacgccgttgagttggcagcaggctcagccatgcactggatggcagctgggctgccactgcaatgtggtggataggatgcaa
gtggagcgaataccaaaccctctggctgcttgctgggttgcatggcatcgcaccatcagcaggagcgcatgcgaagggactggcc
ccatgcacgccatgccaaaccggagcgcaccgagtgtccacactgtcaccaggcccgcaagctttgcagaaccatgctcatggac
gcatgtagcgctgacgtcccttgacggcgctcctctcgggtgtgggaaacgcaatgcagcacaggcagcagaggcggcggcagc
agagcggcggcagcagcggcgggggccacccttcttgcggggtcgcgccccagccagcggtgatgcgctgatcccaaacgagttc
acattcatttgcatgcctggagaagcgaggctggggcctttgggctggtgcagcccgcaatggaatgcgggaccgccaggctagc
agcaaaggcgcctcccctactccgcatcgatgttccatagtgcattggactgcatttgggtggggcggccggctgtttctttcgtgtt
gcaaaacgcgccagctcagcaacctgtcccgtgggtcccccgtgccgatgaaatcgtgtgcacgccgatcagctgattgcccggct
cgcgaagtaggcgccctcctttctgctcgccctctctccgtcccgcctctaga**atatca**ATGATCGAGCAGGACGGCCTCC
ACGCCGGCTCCCCCGCCGCCTGGGTGGAGCGCCTGTTCGGCTACGACTGGGCCCAGCAGACCATCG
GCTGCTCCGACGCCGCCGTGTTCCGCCTGTCCGCCCAGGGCCGCCCCGTGCTGTTCGTGAAGACCGA
CCTGTCCGGCGCCCTGAACGAGCTGCAGGACGAGGCCGCCCGCCTGTCCTGGCTGGCCACCACCGG
CGTGCCCTGCGCCGCCGTGCTGGACGTGGTGACCGAGGCCGGCCGCGACTGGCTGCTGCTGGGCG
AGGTGCCCGGCCAGGACCTGCTGTCCTCCCACCTGGCCCCCGCCGAGAAGGTGTCCATCATGGCCG
ACGCCATGCGCCGCCTGCACACCCTGGACCCCGCCACCTGCCCCTTCGACCACCAGGCCAAGCACCG
CATCGAGCGCGCCCGCACCCGCATGGAGGCCGGCCTGGTGGACCAGGACGACCTGGACGAGGAGC
ACCAGGGCCTGGCCCCCGCCGAGCTGTTCGCCCGCCTGAAGGCCCGCATGCCCGACGGCGAGGACC
TGGTGGTGACCCACGGCGACGCCTGCCTGCCCAACATCATGGTGGAGAACGGCCGCTTCTCCGGCT
TCATCGACTGCGGCCGCCTGGGCGTGGCCGACCGCTACCAGGACATCGCCCTGGCCACCCGCGACA
TCGCCGAGGAGCTGGGCGGCGAGTGGGCCGACCGCTTCCTGGTGCTGTACGGCATCGCCGCCCCCG
ACTCCCAGCGCATCGCCTTCTACCGCCTGCTGGACGAGTTCTTCTGA**caattg***gcagcagcagctcggatagt*
atcgacacactctggacgctggtcgtgtgatggactgttgccgccacacttgctgccttgacctgtgaatatccctgccgcttttat
caaacagcctcagtgtgtttgatcttgtgtgtacgcgcttttgcgagttgctagctgcttgtgctatttgcgaataccacccccagc
atccccttccctcgtttcatatcgcttgcatcccaaccgcaacttatctacgctgtcctgctatccctcagcgctgctcctgctcctgct
cactgcccctcgcacagccttggtttgggctccgcctgtattctcctggtactgcaacctgtaaaccagcactgcaatgctgatgc
*acgggaagtagtgggatgggaacacaaatgga*ggatcc**cgcgtctcgaacagagcgcgcagaggaacgctgaaggtctcgcc**
tctgtcgcacctcagcgcggcatacaccacaataaccacctgacgaatgcgcttggttcttcgtccattagcgaagcgtccggttcac
acacgtgccacgttggcgaggtggcaggtgacaatgatcggtggagctgatggtcgaaacgttcacagcctagggatatcgaattc
ctttcttgcgctatgacacttccagcaaaaggtagggcgggctgcgagacggcttcccggcgctgcatgcaacaccgatgatgctt
cgacccccgaagctccttcggggctgcatgggcgctccgatgccgctccagggcgagcgctgtttaaatagccaggccccgatt
gcaaagacattatagcgagctaccaaagccatattcaaacacctagatcactaccacttctacacaggccactcgagcttgtgatc
gcactccgctaaggggcgcctcttcctcttcgtttcagtcacaacccgcaaacggcgcgcc**ATG**GCCACCACCAGCCTGG
CCTCCGCCTTCTGCTCCATGAAGGCCGTGATGCTGGCCCGCGACGGCCGCGGCATGAAGCCCCGCA
GCTCCGACCTGCAGCTGCGCGCCGGCAACGCCCCCACCTCCCTGAAGATGATCAACGGCACCAAGTT
CAGCTACACCGAGAGCCTGAAGCGCCTGCCCGACTGGTCCATGCTGTTCGCCGTGATCACCACCATC
TTCAGCGCCGCCGAGAAGCAGTGGACCAACCTGGAGTGGAAGCCCAAGCCCAAGCTGCCCCAGCTG
CTGGACGACCACTTCGGCCTGCACGGCCTGGTGTTCCGCCGCACCTTCGCCATCCGCTCCTACGAGG
TGGGCCCCGACCGCAGCACCTCCATCCTGGCCGTGATGAACCACATGCAGGAGGCCACCCTGAACC
ACGCCAAGAGCGTGGGCATCCTGGGCGACGGCTTCGGCACCACCCTGGAGATGTCCAAGCGCGACC
TGATGTGGGTGGTGCGCCGCACCCACGTGGCCGTGGAGCGCTACCCCACCTGGGGCGACACCGTG
GAGGTGGAGTGCTGGATCGGCGCCAGCGGCAACAACGGCATGCGCCGCGACTTCCTGGTGCGCGA
CTGCAAGACCGGCGAGATCCTGACCCGCTGCACCTCCCTGAGCGTGCTGATGAACACCCGCACCCGC
CGCCTGAGCACCATCCCCGACGAGGTGCGCGGCGAGATCGGCCCCGCCTTCATCGACAACGTGGCC
GTGAAGGACGACGAGATCAAGAAGCTGCAGAAGCTGAACGACTCCACCGCCGACTACATCCAGGG

CGGCCTGACCCCCCGCTGGAACGACCTGGACGTGAACCAGCACGTGAACAACCTGAAGTACGTGGC
CTGGGTGTTCGAGACCGTGCCCGACAGCATCTTCGAGTCCCACCACATCAGCTCCTTCACCCTGGAG
TACCGCCGCGAGTGCACCCGCGACTCCGTGCTGCGCAGCCTGACCACCGTGAGCGGCGGCAGCTCC
GAGGCCGGCCTGGTGTGCGACCACCTGCTGCAGCTGGAGGGCGGCAGCGAGGTGCTGCGCGCCCG
CACCGAGTGGCGCCCCAAGCTGACCGACTCCTTCCGCGGCATCAGCGTGATCCCCGCCGAGCCCCGC
GTG*ATGGACTACAAGGACCACGACGGCGACTACAAGGACCACGACATCGACTACAAGGACGACGAC*
*GACAAGTGA**TGA***<u>ctcgag</u>*gcagcagcagctcggatagtatcgacacactctggacgctggtcgtgtgatggactgttgcc*
gccacacttgctgccttgacctgtgaatatccctgccgcttttatcaaacagcctcagtgtgtttgatcttgtgtgtacgcgcttttg
cgagttgctagctgcttgtgctatttgcgaataccacccccagcatccccttccctcgtttcatatcgcttgcatcccaaccgcaact
tatctacgctgtcctgctatccctcagcgctgctcctgctcctgctcactgcccctcgcacagccttggtttgggctccgcctgtattc
tcctggtactgcaacctgtaaaccagcactgcaatgctgatgcacgggaagtagtgggatgggaacacaaatgga<u>aagctt</u>

SEQ ID NO: 119
C.reinhartii β-tubulin promoter::*P.moriformis* isopentenyl diphosphate synthase PTS::*Umbellularia californica* C12 thioesterase::*C.vulgaris* nitrate reductase 3'UTR

<u>ggtacc</u>**cgcctgcaacgcaagggcagccacagccgctcccacccgccgctgaaccgacacgtgcttgggcgcctgccgcctgcctg**
ccgcatgcttgtgctggtgaggctgggcagtgctgccatgctgattgaggcttggttcatcgggtggaagcttatgtgtgtgctggg
cttgcatgccgggcaatgcgcatggtggcaagagggcggcagcacttgctggagctgccgcggtgcctccaggtggttcaatcgc
ggcagccagagggatttcagatgatcgcgcgtacaggttgagcagcagtgtcagcaaaggtagcagtttgccagaatgatcggtt
cagctgttaatcaatgccagcaagagaaggggtcaagtgcaaacacgggcatgccacagcacgggcaccggggagtggaatgg
caccaccaagtgtgtgcgagccagcatcgccgcctggctgtttcagctacaacggcaggagtcatccaacgtaaccatgagctga
tcaacactgcaatcatcgggcgggcgtgatgcaagcatgcctggcgaagacacatggtgtgcggatgctgccggctgctgcctgct
gcgcacgccgttgagttggcagcaggctcagccatgcactggatggcagctgggctgccactgcaatgtggtggataggatgcaa
gtggagcgaataccaaaccctctggctgcttgctgggttgcatggcatcgcaccatcagcaggagcgcatgcgaagggactggcc
ccatgcacgccatgccaaaccggagcgcaccgagtgtccacactgtcaccaggcccgcaagctttgcagaaccatgctcatggac
gcatgtagcgctgacgtcccttgacggcgctcctctcgggtgtgggaaacgcaatgcagcacaggcagcagaggcggcggcagc
agagcggcggcagcagcggcgggggccacccttcttgcggggtcgcgccccagccagcggtgatgcgctgatcccaaacgagttc
acattcatttgcatgcctggagaagcgaggctggggcctttgggctggtgcagcccgcaatggaatgcgggaccgccaggctagc
agcaaaggcgcctccctactccgcatcgatgttccatagtgcattggactgcatttgggtggggcggccggctgtttctttcgtgtt
gcaaaacgcgccagctcagcaacctgtcccgtgggtcccccgtgccgatgaaatcgtgtgcacgccgatcagctgattgcccggct
cgcgaagtaggcgccctcctttctgctcgccctctctccgtcccgcc<u>tctaga</u>atatcaATGATCGAGCAGGACGGCCTCC
ACGCCGGCTCCCCCGCCGCCTGGGTGGAGCGCCTGTTCGGCTACGACTGGGCCCAGCAGACCATCG
GCTGCTCCGACGCCGCCGTGTTCCGCCTGTCCGCCCAGGGCCGCCCCGTGCTGTTCGTGAAGACCGA
CCTGTCCGGCGCCCTGAACGAGCTGCAGGACGAGGCCGCCCGCCTGTCCTGGCTGGCCACCACCGG
CGTGCCCTGCGCCGCCGTGCTGGACGTGGTGACCGAGGCCGGCCGCGACTGGCTGCTGCTGGGCG
AGGTGCCCGGCCAGGACCTGCTGTCCTCCCACCTGGCCCCCGCCGAGAAGGTGTCCATCATGGCCG
ACGCCATGCGCCGCCTGCACACCCTGGACCCCGCCACCTGCCCCTTCGACCACCAGGCCAAGCACCG
CATCGAGCGCGCCCGCACCCGCATGGAGGCCGGCCTGGTGGACCAGGACGACCTGGACGAGGAGC
ACCAGGGCCTGGCCCCCGCCGAGCTGTTCGCCCGCCTGAAGGCCCGCATGCCCGACGGCGAGGACC
TGGTGGTGACCCACGGCGACGCCTGCCTGCCCAACATCATGGTGGAGAACGGCCGCTTCTCCGGCT
TCATCGACTGCGGCCGCCTGGGCGTGGCCGACCGCTACCAGGACATCGCCCTGGCCACCCGCGACA
TCGCCGAGGAGCTGGGCGGCGAGTGGGCCGACCGCTTCCTGGTGCTGTACGGCATCGCCGCCCCCG
ACTCCCAGCGCATCGCCTTCTACCGCCTGCTGGACGAGTTCTTCTGA<u>caattgg</u>*cagcagcagctcggatagt*
atcgacacactctggacgctggtcgtgtgatggactgttgccgccacacttgctgccttgacctgtgaatatccctgccgcttttat
caaacagcctcagtgtgtttgatcttgtgtgtacgcgcttttgcgagttgctagctgcttgtgctatttgcgaataccacccccagc
atcccccttccctcgtttcatatcgcttgcatcccaaccgcaacttatctacgctgtcctgctatccctcagcgctgctcctgctcctgct

cactgcccctcgcacagccttggtttgggctccgcctgtattctcctggtactgcaacctgtaaaccagcactgcaatgctgatgc
*acgggaagtagtgggatgggaacacaaatgga*ggatccgcgtctcgaacagagcgcgcagaggaacgctgaaggtctcgcc
tctgtcgcacctcagcgcggcatacaccacaataaccacctgacgaatgcgcttggttcttcgtccattagcgaagcgtccggttcac
acacgtgccacgttggcgaggtggcaggtgacaatgatcggtggagctgatggtcgaaacgttcacagcctagggatatcgaattc
ctttcttgcgctatgacacttccagcaaaaggtagggcgggctgcgagacggcttcccggcgctgcatgcaacaccgatgatgctt
cgacccccgaagctccttcggggctgcatgggcgctccgatgccgctccagggcgagcgctgtttaaatagccaggcccccgatt
gcaaagacattatagcgagctaccaaagccatattcaaacacctagatcactaccacttctacacaggccactcgagcttgtgatc
gcactccgctaaggggcgcctcttcctcttcgtttcagtcacaacccgcaaacactagt**ATG**ACGTTCGGGGTCGCCCTC
CCGGCCATGGGCCGCGGTGTCTCCCTTCCCCGGCCCAGGGTCGCGGTGCGCGCCCAGTCGGCGAGT
CAGGTTTTGGAGAGCGGGCGCGCCCCCGACTGGTCCATGCTGTTCGCCGTGATCACCACCATCTTCA
GCGCCGCCGAGAAGCAGTGGACCAACCTGGAGTGGAAGCCCAAGCCCAAGCTGCCCCAGCTGCTG
GACGACCACTTCGGCCTGCACGGCCTGGTGTTCCGCCGCACCTTCGCCATCCGCTCCTACGAGGTGG
GCCCCGACCGCAGCACCTCCATCCTGGCCGTGATGAACCACATGCAGGAGGCCACCCTGAACCACG
CCAAGAGCGTGGGCATCCTGGGCGACGGCTTCGGCACCACCCTGGAGATGTCCAAGCGCGACCTGA
TGTGGGTGGTGCGCCGCACCCACGTGGCCGTGGAGCGCTACCCCACCTGGGGCGACACCGTGGAG
GTGGAGTGCTGGATCGGCGCCAGCGGCAACAACGGCATGCGCCGCGACTTCCTGGTGCGCGACTG
CAAGACCGGCGAGATCCTGACCCGCTGCACCTCCCTGAGCGTGCTGATGAACACCCGCACCCGCCGC
CTGAGCACCATCCCCGACGAGGTGCGCGGCGAGATCGGCCCCGCCTTCATCGACAACGTGGCCGTG
AAGGACGACGAGATCAAGAAGCTGCAGAAGCTGAACGACTCCACCGCCGACTACATCCAGGGCGG
CCTGACCCCCCGCTGGAACGACCTGGACGTGAACCAGCACGTGAACAACCTGAAGTACGTGGCCTG
GGTGTTCGAGACCGTGCCCGACAGCATCTTCGAGTCCCACCACATCAGCTCCTTCACCCTGGAGTAC
CGCCGCGAGTGCACCCGCGACTCCGTGCTGCGCAGCCTGACCACCGTGAGCGGCGGCAGCTCCGAG
GCCGGCCTGGTGTGCGACCACCTGCTGCAGCTGGAGGGCGGCAGCGAGGTGCTGCGCGCCCGCAC
CGAGTGGCGCCCCAAGCTGACCGACTCCTTCCGCGGCATCAGCGTGATCCCCGCCGAGCCCCGCGT
GATGGACTACAAGGACCACGACGGCGACTACAAGGACCACGACATCGACTACAAGGACGACGACGA
*CAAGTGA***TGA**ctcgag*gcagcagcagctcggatagtatcgacacactctggacgctggtcgtgtgatggactgttgccgcc*
acacttgctgccttgacctgtgaatatccctgccgcttttatcaaacagcctcagtgtgtttgatcttgtgtgtacgcgcttttgcga
gttgctagctgcttgtgctatttgcgaataccacccccagcatcccctttccctcgtttcatatcgcttgcatcccaaccgcaacttat
ctacgctgtcctgctatccctcagcgctgctcctgctcctgctcactgcccctcgcacagccttggtttgggctccgcctgtattctcc
*tggtactgcaacctgtaaaccagcactgcaatgctgatgcacgggaagtagtgggatgggaacacaaatgga*aagctt

SEQ ID NO: 120
C.reinhartii β-tubulin promoter::*P.moriformis* delta 12 fatty acid desaturase
PTS::*Umbellularia californica* C12 thioesterase::*C.vulgaris* nitrate reductase 3'UTR

ggtacc**cgcctgcaacgcaagggcagccacagccgctcccacccgccgctgaaccgacacgtgcttgggcgcctgccgcctgcctg**
ccgcatgcttgtgctggtgaggctgggcagtgctgccatgctgattgaggcttggttcatcgggtggaagcttatgtgtgtgctggg
cttgcatgccgggcaatgcgcatggtggcaagagggcggcagcacttgctggagctgccgcggtgcctccaggtggttcaatcgc
ggcagccagagggatttcagatgatcgcgcgtacaggttgagcagcagtgtcagcaaaggtagcagtttgccagaatgatcggtt
cagctgttaatcaatgccagcaagagaaggggtcaagtgcaaacacgggcatgccacagcacgggcaccggggagtggaatgg
caccaccaagtgtgtgcgagccagcatcgccgcctggctgtttcagctacaacggcaggagtcatccaacgtaaccatgagctga
tcaacactgcaatcatcgggcgggcgtgatgcaagcatgcctggcgaagacacatggtgtgcggatgctgccggctgctgcctgct
gcgcacgccgttgagttggcagcaggctcagccatgcactggatggcagctgggctgccactgcaatgtggtggataggatgcaa
gtggagcgaataccaaaccctctggctgcttgctgggttgcatggcatcgcaccatcagcaggagcgcatgcgaagggactggcc
ccatgcacgccatgccaaaccggagcgcaccgagtgtccacactgtcaccaggcccgcaagctttgcagaaccatgctcatggac
gcatgtagcgctgacgtcccttgacggcgctcctctcgggtgtgggaaacgcaatgcagcacaggcagcagaggcggcggcagc
agagcggcggcagcagcggcgggggccacccttcttgcggggtcgcgccccagccagcggtgatgcgctgatcccaaacgagttc

acattcatttgcatgcctggagaagcgaggctggggcctttgggctggtgcagcccgcaatggaatgcgggaccgccaggctagc
agcaaaggcgcctcccctactccgcatcgatgttccatagtgcattggactgcatttgggtggggcggccggctgtttctttcgtgtt
gcaaaacgcgccagctcagcaacctgtcccgtgggtcccccgtgccgatgaaatcgtgtgcacgccgatcagctgattgcccggct
cgcgaagtaggcgccctcctttctgctcgccctctctccgtcccgcctctagaatatcaATGATCGAGCAGGACGGCCTCC
ACGCCGGCTCCCCCGCCGCCTGGGTGGAGCGCCTGTTCGGCTACGACTGGGCCCAGCAGACCATCG
GCTGCTCCGACGCCGCCGTGTTCCGCCTGTCCGCCCAGGGCCGCCCCGTGCTGTTCGTGAAGACCGA
CCTGTCCGGCGCCCTGAACGAGCTGCAGGACGAGGCCGCCCGCCTGTCCTGGCTGGCCACCACCGG
CGTGCCCTGCGCCGCCGTGCTGGACGTGGTGACCGAGGCCGGCCGCGACTGGCTGCTGCTGGGCG
AGGTGCCCGGCCAGGACCTGCTGTCCTCCCACCTGGCCCCCGCCGAGAAGGTGTCCATCATGGCCG
ACGCCATGCGCCGCCTGCACACCCTGGACCCCGCCACCTGCCCCTTCGACCACCAGGCCAAGCACCG
CATCGAGCGCGCCCGCACCCGCATGGAGGCCGGCCTGGTGGACCAGGACGACCTGGACGAGGAGC
ACCAGGGCCTGGCCCCCGCCGAGCTGTTCGCCCGCCTGAAGGCCCGCATGCCCGACGGCGAGGACC
TGGTGGTGACCCACGGCGACGCCTGCCTGCCCAACATCATGGTGGAGAACGGCCGCTTCTCCGGCT
TCATCGACTGCGGCCGCCTGGGCGTGGCCGACCGCTACCAGGACATCGCCCTGGCCACCCGCGACA
TCGCCGAGGAGCTGGGCGGCGAGTGGGCCGACCGCTTCCTGGTGCTGTACGGCATCGCCGCCCCCG
ACTCCCAGCGCATCGCCTTCTACCGCCTGCTGGACGAGTTCTTCTGA<u>caattg</u>*gcagcagcagctcggatagt*
atcgacacactctggacgctggtcgtgtgatggactgttgccgccacacttgctgccttgacctgtgaatatccctgccgcttttat
caaacagcctcagtgtgtttgatcttgtgtgtacgcgcttttgcgagttgctagctgcttgtgctatttgcgaataccacccccagc
atccccttccctcgtttcatatcgcttgcatcccaaccgcaacttatctacgctgtcctgctatccctcagcgctgctcctgctcctgct
cactgcccctcgcacagccttggtttgggctccgcctgtattctcctggtactgcaacctgtaaaccagcactgcaatgctgatgc
acgggaagtagtgggatgggaacacaaatgga<u>ggatcc</u>*c*gcgtctcgaacagagcgcgcagaggaacgctgaaggtctcgcc
tctgtcgcacctcagcgcggcatacaccacaataaccacctgacgaatgcgcttggttcttcgtccattagcgaagcgtccggttcac
acacgtgccacgttggcgaggtggcaggtgacaatgatcggtggagctgatggtcgaaacgttcacagcctagggatatc<u>gaattc</u>
ctttcttgcgctatgacacttccagcaaaaggtagggcgggctgcgagacggcttcccggcgctgcatgcaacaccgatgatgctt
cgacccccgaagctccttcggggctgcatgggcgctccgatgccgctccagggcgagcgctgtttaaatagccaggccccgatt
gcaaagacattatagcgagctaccaaagccatattcaaacacctagatcactaccacttctacacaggccactcgagcttgtgatc
gcactccgctaaggggcgcctcttcctcttcgtttcagtcacaacccgcaaac<u>actagt</u>**ATG**GCTATCAAGACGAACAG
GCAGCCTGTGGAGAAGCCTCCGTTCACGATCGGGACGCTGCGCAAGGCCATCCCCGCGCACTGTTT
CGAGCGCTCGGCGCTTCGTG<u>GGCGCGCC</u>CCCGACTGGTCCATGCTGTTCGCCGTGATCACCACCATC
TTCAGCGCCGCCGAGAAGCAGTGGACCAACCTGGAGTGGAAGCCCAAGCCCAAGCTGCCCCAGCTG
CTGGACGACCACTTCGGCCTGCACGGCCTGGTGTTCCGCCGCACCTTCGCCATCCGCTCCTACGAGG
TGGGCCCCGACCGCAGCACCTCCATCCTGGCCGTGATGAACCACATGCAGGAGGCCACCCTGAACC
ACGCCAAGAGCGTGGGCATCCTGGGCGACGGCTTCGGCACCACCCTGGAGATGTCCAAGCGCGACC
TGATGTGGGTGGTGCGCCGCACCCACGTGGCCGTGGAGCGCTACCCCACCTGGGGCGACACCGTG
GAGGTGGAGTGCTGGATCGGCGCCAGCGGCAACAACGGCATGCGCCGCGACTTCCTGGTGCGCGA
CTGCAAGACCGGCGAGATCCTGACCCGCTGCACCTCCCTGAGCGTGCTGATGAACACCCGCACCCGC
CGCCTGAGCACCATCCCCGACGAGGTGCGCGGCGAGATCGGCCCCGCCTTCATCGACAACGTGGCC
GTGAAGGACGACGAGATCAAGAAGCTGCAGAAGCTGAACGACTCCACCGCCGACTACATCCAGGG
CGGCCTGACCCCCCGCTGGAACGACCTGGACGTGAACCAGCACGTGAACAACCTGAAGTACGTGGC
CTGGGTGTTCGAGACCGTGCCCGACAGCATCTTCGAGTCCCACCACATCAGCTCCTTCACCCTGGAG
TACCGCCGCGAGTGCACCCGCGACTCCGTGCTGCGCAGCCTGACCACCGTGAGCGGCGGCAGCTCC
GAGGCCGGCCTGGTGTGCGACCACCTGCTGCAGCTGGAGGGCGGCAGCGAGGTGCTGCGCGCCCG
CACCGAGTGGCGCCCCAAGCTGACCGACTCCTTCCGCGGCATCAGCGTGATCCCCGCCGAGCCCCGC
GTG*ATGGACTACAAGGACCACGACGGCGACTACAAGGACCACGACATCGACTACAAGGACGACGAC*
*GACAAGTGA***TGA**<u>ctcgag</u>*gcagcagcagctcggatagtatcgacacactctggacgctggtcgtgtgatggactgttgcc*
gccacacttgctgccttgacctgtgaatatccctgccgcttttatcaaacagcctcagtgtgtttgatcttgtgtgtacgcgcttttg
cgagttgctagctgcttgtgctatttgcgaataccacccccagcatcccttccctcgtttcatatcgcttgcatcccaaccgcaact

tatctacgctgtcctgctatccctcagcgctgctcctgctcctgctcactgcccctcgcacagccttggtttgggctccgcctgtattc
tcctggtactgcaacctgtaaaccagcactgcaatgctgatgcacgggaagtagtgggatgggaacacaaatggaaagctt

SEQ ID NO: 121
C.reinhartii β-tubulin promoter::*C.protothecoides* stearoyl ACP desaturase
PTS::*Umbellularia californica* C12 thioesterase::*C.vulgaris* nitrate reductase 3'UTR

ggtacc**cgcctgcaacgcaagggcagccacagccgctcccacccgccgctgaaccgacacgtgcttgggcgcctgccgcctgcctg**
ccgcatgcttgtgctggtgaggctgggcagtgctgccatgctgattgaggcttggttcatcgggtggaagcttatgtgtgtgctggg
cttgcatgccgggcaatgcgcatggtggcaagagggcggcagcacttgctggagctgccgcggtgcctccaggtggttcaatcgc
ggcagccagagggatttcagatgatcgcgcgtacaggttgagcagcagtgtcagcaaaggtagcagtttgccagaatgatcggtt
cagctgttaatcaatgccagcaagagaaggggtcaagtgcaaacacgggcatgccacagcacgggcaccggggagtggaatgg
caccaccaagtgtgtgcgagccagcatcgccgcctggctgtttcagctacaacggcaggagtcatccaacgtaaccatgagctga
tcaacactgcaatcatcgggcgggcgtgatgcaagcatgcctggcgaagacacatggtgtgcggatgctgccggctgctgcctgct
gcgcacgccgttgagttggcagcaggctcagccatgcactggatggcagctgggctgccactgcaatgtggtggataggatgcaa
gtggagcgaataccaaaccctctggctgcttgctgggttgcatggcatcgcaccatcagcaggagcgcatgcgaagggactggcc
ccatgcacgccatgccaaaccggagcgcaccgagtgtccacactgtcaccaggcccgcaagctttgcagaaccatgctcatggac
gcatgtagcgctgacgtcccttgacggcgctcctctcgggtgtgggaaacgcaatgcagcacaggcagcagaggcggcggcagc
agagcggcggcagcagcggcgggggccacccttcttgcggggtcgcgccccagccagcggtgatgcgctgatcccaaacgagttc
acattcatttgcatgcctggagaagcgaggctggggcctttgggctggtgcagcccgcaatggaatgcgggaccgccaggctagc
agcaaaggcgcctccctactccgcatcgatgttccatagtgcattggactgcatttgggtggggcggccggctgtttctttcgtgtt
gcaaaacgcgccagctcagcaacctgtcccgtgggtcccccgtgccgatgaaatcgtgtgcacgccgatcagctgattgcccggct
cgcgaagtaggcgccctcctttctgctcgccctctctccgtcccgcctctaga**atatcaATGATCGAGCAGGACGGCCTCC**
ACGCCGGCTCCCCCGCCGCCTGGGTGGAGCGCCTGTTCGGCTACGACTGGGCCCAGCAGACCATCG
GCTGCTCCGACGCCGCCGTGTTCCGCCTGTCCGCCCAGGGCCGCCCCGTGCTGTTCGTGAAGACCGA
CCTGTCCGGCGCCCTGAACGAGCTGCAGGACGAGGCCGCCCGCCTGTCCTGGCTGGCCACCACCGG
CGTGCCCTGCGCCGCCGTGCTGGACGTGGTGACCGAGGCCGGCCGCGACTGGCTGCTGCTGGGCG
AGGTGCCCGGCCAGGACCTGCTGTCCTCCCACCTGGCCCCCGCCGAGAAGGTGTCCATCATGGCCG
ACGCCATGCGCCGCCTGCACACCCTGGACCCCGCCACCTGCCCCTTCGACCACCAGGCCAAGCACCG
CATCGAGCGCGCCCGCACCCGCATGGAGGCCGGCCTGGTGGACCAGGACGACCTGGACGAGGAGC
ACCAGGGCCTGGCCCCCGCCGAGCTGTTCGCCCGCCTGAAGGCCCGCATGCCCGACGGCGAGGACC
TGGTGGTGACCCACGGCGACGCCTGCCTGCCCAACATCATGGTGGAGAACGGCCGCTTCTCCGGCT
TCATCGACTGCGGCCGCCTGGGCGTGGCCGACCGCTACCAGGACATCGCCCTGGCCACCCGCGACA
TCGCCGAGGAGCTGGGCGGCGAGTGGGCCGACCGCTTCCTGGTGCTGTACGGCATCGCCGCCCCCG
ACTCCCAGCGCATCGCCTTCTACCGCCTGCTGGACGAGTTCTTCTGA*caattggcagcagcagctcggatagt*
atcgacacactctggacgctggtcgtgtgatggactgttgccgccacacttgctgccttgacctgtgaatatccctgccgcttttat
caaacagcctcagtgtgtttgatcttgtgtgtacgcgcttttgcgagttgctagctgcttgtgctatttgcgaataccacccccagc
atccccttccctcgtttcatatcgcttgcatcccaaccgcaacttatctacgctgtcctgctatccctcagcgctgctcctgctcctgct
cactgcccctcgcacagccttggtttgggctccgcctgtattctcctggtactgcaacctgtaaaccagcactgcaatgctgatgc
*acgggaagtagtgggatgggaacacaaatgga*ggatcc**cgcgtctcgaacagagcgcgcagaggaacgctgaaggtctcgcc**
tctgtcgcacctcagcgcggcatacaccacaataaccacctgacgaatgcgcttggttcttcgtccattagcgaagcgtccggttcac
acacgtgccacgttggcgaggtggcaggtgacaatgatcggtggagctgatggtcgaaacgttcacagcctagggatatcgaattc
ctttcttgcgctatgacacttccagcaaaaggtagggcgggctgcgagacggcttcccggcgctgcatgcaacaccgatgatgctt
cgacccccgaagctccttcggggctgcatgggcgctccgatgccgctccagggcgagcgctgtttaaatagccaggcccccgatt
gcaaagacattatagcgagctaccaaagccatattcaaacacctagatcactaccacttctacacaggccactcgagcttgtgatc
gcactccgctaaggggcgcctcttcctcttcgtttcagtcacaacccgcaaacactagt**ATG**GCCACCGCATCCACTTTCT
CGGCGTTCAATGCCCGCTGCGGCGACCTGCGTCGCTCGGCGGGCTCCGGGCCCCGGCGCCCAGCGA

GGCCCCTCCCCGTGCGCGGGCGCGCCCCCGACTGGTCCATGCTGTTCGCCGTGATCACCACCATCTT
CAGCGCCGCCGAGAAGCAGTGGACCAACCTGGAGTGGAAGCCCAAGCCCAAGCTGCCCCAGCTGCT
GGACGACCACTTCGGCCTGCACGGCCTGGTGTTCCGCCGCACCTTCGCCATCCGCTCCTACGAGGTG
GGCCCCGACCGCAGCACCTCCATCCTGGCCGTGATGAACCACATGCAGGAGGCCACCCTGAACCAC
GCCAAGAGCGTGGGCATCCTGGGCGACGGCTTCGGCACCACCCTGGAGATGTCCAAGCGCGACCTG
ATGTGGGTGGTGCGCCGCACCCACGTGGCCGTGGAGCGCTACCCCACCTGGGGCGACACCGTGGA
GGTGGAGTGCTGGATCGGCGCCAGCGGCAACAACGGCATGCGCCGCGACTTCCTGGTGCGCGACT
GCAAGACCGGCGAGATCCTGACCCGCTGCACCTCCCTGAGCGTGCTGATGAACACCCGCACCCGCC
GCCTGAGCACCATCCCCGACGAGGTGCGCGGCGAGATCGGCCCCGCCTTCATCGACAACGTGGCCG
TGAAGGACGACGAGATCAAGAAGCTGCAGAAGCTGAACGACTCCACCGCCGACTACATCCAGGGC
GGCCTGACCCCCCGCTGGAACGACCTGGACGTGAACCAGCACGTGAACAACCTGAAGTACGTGGCC
TGGGTGTTCGAGACCGTGCCCGACAGCATCTTCGAGTCCCACCACATCAGCTCCTTCACCCTGGAGT
ACCGCCGCGAGTGCACCCGCGACTCCGTGCTGCGCAGCCTGACCACCGTGAGCGGCGGCAGCTCCG
AGGCCGGCCTGGTGTGCGACCACCTGCTGCAGCTGGAGGGCGGCAGCGAGGTGCTGCGCGCCCGC
ACCGAGTGGCGCCCCAAGCTGACCGACTCCTTCCGCGGCATCAGCGTGATCCCCGCCGAGCCCCGC
GTG*ATGGACTACAAGGACCACGACGGCGACTACAAGGACCACGACATCGACTACAAGGACGACGAC*
*GACAAGTGA**TGA***ctcgagg*cagcagcagctcggatagtatcgacacactctggacgctggtcgtgtgatggactgttgcc*
gccacacttgctgccttgacctgtgaatatccctgccgcttttatcaaacagcctcagtgtgtttgatcttgtgtgtacgcgcttttg
cgagttgctagctgcttgtgctatttgcgaataccacccccagcatccccttccctcgtttcatatcgcttgcatcccaaccgcaact
tatctacgctgtcctgctatccctcagcgctgctcctgctcctgctcactgcccctcgcacagccttggtttgggctccgcctgtattc
*tcctggtactgcaacctgtaaaccagcactgcaatgctgatgcacgggaagtagtgggatgggaacacaaatgga*aagctt

SEQ ID NO: 122
C.reinhartii β-tubulin promoter::*P.moriformis* stearoyl ACP desaturase PTS::*Umbellularia californica* C12 thioesterase::*C.vulgaris* nitrate reductase 3'UTR

ggtacc**cgcctgcaacgcaagggcagccacagccgctcccacccgccgctgaaccgacacgtgcttgggcgcctgccgcctgcctg**
ccgcatgcttgtgctggtgaggctgggcagtgctgccatgctgattgaggcttggttcatcgggtggaagcttatgtgtgtgctggg
cttgcatgccgggcaatgcgcatggtggcaagagggcggcagcacttgctggagctgccgcggtgcctccaggtggttcaatcgc
ggcagccagagggatttcagatgatcgcgcgtacaggttgagcagcagtgtcagcaaaggtagcagtttgccagaatgatcggtt
cagctgttaatcaatgccagcaagagaaggggtcaagtgcaaacacgggcatgccacagcacgggcaccggggagtggaatgg
caccaccaagtgtgtgcgagccagcatcgccgcctggctgtttcagctacaacggcaggagtcatccaacgtaaccatgagctga
tcaacactgcaatcatcgggcgggcgtgatgcaagcatgcctggcgaagacacatggtgtgcggatgctgccggctgctgcctgct
gcgcacgccgttgagttggcagcaggctcagccatgcactggatggcagctgggctgccactgcaatgtggtggataggatgcaa
gtggagcgaataccaaaccctctggctgcttgctgggttgcatggcatcgcaccatcagcaggagcgcatgcgaagggactggcc
ccatgcacgccatgccaaaccggagcgcaccgagtgtccacactgtcaccaggcccgcaagctttgcagaaccatgctcatggac
gcatgtagcgctgacgtcccttgacggcgctcctctcgggtgtgggaaacgcaatgcagcacaggcagcagaggcggcggcagc
agagcggcggcagcagcggcgggggccaccctttcttgcggggtcgcgccccagccagcggtgatgcgctgatcccaaacgagttc
acattcatttgcatgcctggagaagcgaggctggggcctttgggctggtgcagcccgcaatggaatgcgggaccgccaggctagc
agcaaaggcgcctcccctactccgcatcgatgttccatagtgcattggactgcatttgggtggggcggccggctgtttctttcgtgtt
gcaaaacgcgccagctcagcaacctgtcccgtgggtcccccgtgccgatgaaatcgtgtgcacgccgatcagctgattgcccggct
cgcgaagtaggcgccctcctttctgctcgccctctctccgtcccgcctctaga*atatcaATGATCGAGCAGGACGGCCTCC*
ACGCCGGCTCCCCCGCCGCCTGGGTGGAGCGCCTGTTCGGCTACGACTGGGCCCAGCAGACCATCG
GCTGCTCCGACGCCGCCGTGTTCCGCCTGTCCGCCCAGGGCCGCCCCGTGCTGTTCGTGAAGACCGA
CCTGTCCGGCGCCCTGAACGAGCTGCAGGACGAGGCCGCCCGCCTGTCCTGGCTGGCCACCACCGG
CGTGCCCTGCGCCGCCGTGCTGGACGTGGTGACCGAGGCCGGCCGCGACTGGCTGCTGCTGGGCG
AGGTGCCCGGCCAGGACCTGCTGTCCTCCCACCTGGCCCCCGCCGAGAAGGTGTCCATCATGGCCG

ACGCCATGCGCCGCCTGCACACCCTGGACCCCGCCACCTGCCCCTTCGACCACCAGGCCAAGCACCG
CATCGAGCGCGCCCGCACCCGCATGGAGGCCGGCCTGGTGGACCAGGACGACCTGGACGAGGAGC
ACCAGGGCCTGGCCCCCGCCGAGCTGTTCGCCCGCCTGAAGGCCCGCATGCCCGACGGCGAGGACC
TGGTGGTGACCCACGGCGACGCCTGCCTGCCCAACATCATGGTGGAGAACGGCCGCTTCTCCGGCT
TCATCGACTGCGGCCGCCTGGGCGTGGCCGACCGCTACCAGGACATCGCCCTGGCCACCCGCGACA
TCGCCGAGGAGCTGGGCGGCGAGTGGGCCGACCGCTTCCTGGTGCTGTACGGCATCGCCGCCCCCG
ACTCCCAGCGCATCGCCTTCTACCGCCTGCTGGACGAGTTCTTCTGA*caattggcagcagcagctcggatagt*
atcgacacactctggacgctggtcgtgtgatggactgttgccgccacacttgctgccttgacctgtgaatatccctgccgcttttat
caaacagcctcagtgtgtttgatcttgtgtgtacgcgcttttgcgagttgctagctgcttgtgctatttgcgaataccacccccagc
atccccttccctcgtttcatatcgcttgcatcccaaccgcaacttatctacgctgtcctgctatccctcagcgctgctcctgctcctgct
cactgcccctcgcacagccttggtttgggctccgcctgtattctcctggtactgcaacctgtaaaccagcactgcaatgctgatgc
*acgggaagtagtgggatgggaacacaaatgga*ggatccgcgtctcgaacagagcgcgcagaggaacgctgaaggtctcgcc
tctgtcgcacctcagcgcggcatacaccacaataaccacctgacgaatgcgcttggttcttcgtccattagcgaagcgtccggttcac
acacgtgccacgttggcgaggtggcaggtgacaatgatcggtggagctgatggtcgaaacgttcacagcctagggatatcgaattc
ctttcttgcgctatgacacttccagcaaaaggtagggcgggctgcgagacggcttcccggcgctgcatgcaacaccgatgatgctt
cgacccccgaagctccttcggggctgcatgggcgctccgatgccgctccagggcgagcgctgtttaaatagccaggccccgatt
gcaaagacattatagcgagctaccaaagccatattcaaacacctagatcactaccacttctacacaggccactcgagcttgtgatc
gcactccgctaaggggcgcctcttcctcttcgtttcagtcacaacccgcaaacactagt**ATG**GCTTCCGCGGCATTCACC
ATGTCGGCGTGCCCCGCGATGACTGGCAGGGCCCCTGGGGCACGTCGCTCCGGACGGCCAGTCGCC
ACCCGCCTGAGGGGGCGCGCCCCCGACTGGTCCATGCTGTTCGCCGTGATCACCACCATCTTCAGCG
CCGCCGAGAAGCAGTGGACCAACCTGGAGTGGAAGCCCAAGCCCAAGCTGCCCCAGCTGCTGGAC
GACCACTTCGGCCTGCACGGCCTGGTGTTCCGCCGCACCTTCGCCATCCGCTCCTACGAGGTGGGCC
CCGACCGCAGCACCTCCATCCTGGCCGTGATGAACCACATGCAGGAGGCCACCCTGAACCACGCCA
AGAGCGTGGGCATCCTGGGCGACGGCTTCGGCACCACCCTGGAGATGTCCAAGCGCGACCTGATGT
GGGTGGTGCGCCGCACCCACGTGGCCGTGGAGCGCTACCCCACCTGGGGCGACACCGTGGAGGTG
GAGTGCTGGATCGGCGCCAGCGGCAACAACGGCATGCGCCGCGACTTCCTGGTGCGCGACTGCAA
GACCGGCGAGATCCTGACCCGCTGCACCTCCCTGAGCGTGCTGATGAACACCCGCACCCGCCGCCTG
AGCACCATCCCCGACGAGGTGCGCGGCGAGATCGGCCCCGCCTTCATCGACAACGTGGCCGTGAAG
GACGACGAGATCAAGAAGCTGCAGAAGCTGAACGACTCCACCGCCGACTACATCCAGGGCGGCCTG
ACCCCCCGCTGGAACGACCTGGACGTGAACCAGCACGTGAACAACCTGAAGTACGTGGCCTGGGTG
TTCGAGACCGTGCCCGACAGCATCTTCGAGTCCCACCACATCAGCTCCTTCACCCTGGAGTACCGCCG
CGAGTGCACCCGCGACTCCGTGCTGCGCAGCCTGACCACCGTGAGCGGCGGCAGCTCCGAGGCCG
GCCTGGTGTGCGACCACCTGCTGCAGCTGGAGGGCGGCAGCGAGGTGCTGCGCGCCCGCACCGAG
TGGCGCCCCAAGCTGACCGACTCCTTCCGCGGCATCAGCGTGATCCCCGCCGAGCCCCGCGTG*ATG*
GACTACAAGGACCACGACGGCGACTACAAGGACCACGACATCGACTACAAGGACGACGACGACAAG
*TGA***TGA**ctcgag*gcagcagcagctcggatagtatcgacacactctggacgctggtcgtgtgatggactgttgccgccacact*
tgctgccttgacctgtgaatatccctgccgcttttatcaaacagcctcagtgtgtttgatcttgtgtgtacgcgcttttgcgagttgc
tagctgcttgtgctatttgcgaataccacccccagcatccccttccctcgtttcatatcgcttgcatcccaaccgcaacttatctacg
ctgtcctgctatccctcagcgctgctcctgctcctgctcactgcccctcgcacagccttggtttgggctccgcctgtattctcctggta
*ctgcaacctgtaaaccagcactgcaatgctgatgcacgggaagtagtgggatgggaacacaaatgga*aagctt

SEQ ID NO: 123
C.reinhartii β-tubulin promoter::native PTS::*Cinnamomum camphora* C14 thioesterase::*C.vulgaris* nitrate reductase 3'UTR

ggtaccgcctgcaacgcaagggcagccacagccgctcccacccgccgctgaaccgacacgtgcttgggcgcctgccgcctgcctg
ccgcatgcttgtgctggtgaggctgggcagtgctgccatgctgattgaggcttggttcatcgggtggaagcttatgtgtgtgctgggct

tgcatgccgggcaatgcgcatggtggcaagagggcggcagcacttgctggagctgccgcggtgcctccaggtggttcaatcgcggc
agccagagggatttcagatgatcgcgcgtacaggttgagcagcagtgtcagcaaaggtagcagtttgccagaatgatcggttcagc
tgttaatcaatgccagcaagagaaggggtcaagtgcaaacacgggcatgccacagcacgggcaccggggagtggaatggcacca
ccaagtgtgtgcgagccagcatcgccgcctggctgtttcagctacaacggcaggagtcatccaacgtaaccatgagctgatcaaca
ctgcaatcatcgggcgggcgtgatgcaagcatgcctggcgaagacacatggtgtgcggatgctgccggctgctgcctgctgcgcac
gccgttgagttggcagcaggctcagccatgcactggatggcagctgggctgccactgcaatgtggtggataggatgcaagtggagc
gaataccaaaccctctggctgcttgctgggttgcatggcatcgcaccatcagcaggagcgcatgcgaagggactggccccatgcac
gccatgccaaaccggagcgcaccgagtgtccacactgtcaccaggcccgcaagctttgcagaaccatgctcatggacgcatgtagc
gctgacgtcccttgacggcgctcctctcgggtgtgggaaacgcaatgcagcacaggcagcagaggcggcggcagcagagcggcgg
cagcagcggcgggggccacccttcttgcggggtcgcgccccagccagcggtgatgcgctgatcccaaacgagttcacattcatttgc
atgcctggagaagcgaggctggggcctttgggctggtgcagcccgcaatggaatgcgggaccgccaggctagcagcaaaggcgcc
tcccctactccgcatcgatgttccatagtgcattggactgcatttgggtggggcggccggctgtttctttcgtgttgcaaaacgcgcca
gctcagcaacctgtcccgtgggtcccccgtgccgatgaaatcgtgtgcacgccgatcagctgattgcccggctcgcgaagtaggcgc
cctcctttctgctcgccctctctccgtcccgcctctagaatatcaATGATCGAGCAGGACGGCCTCCACGCCGGCTCCC
CCGCCGCCTGGGTGGAGCGCCTGTTCGGCTACGACTGGGCCCAGCAGACCATCGGCTGCTCCGACG
CCGCCGTGTTCCGCCTGTCCGCCCAGGGCCGCCCCGTGCTGTTCGTGAAGACCGACCTGTCCGGCGC
CCTGAACGAGCTGCAGGACGAGGCCGCCCGCCTGTCCTGGCTGGCCACCACCGGCGTGCCCTGCGC
CGCCGTGCTGGACGTGGTGACCGAGGCCGGCCGCGACTGGCTGCTGCTGGGCGAGGTGCCCGGCC
AGGACCTGCTGTCCTCCCACCTGGCCCCCGCCGAGAAGGTGTCCATCATGGCCGACGCCATGCGCCG
CCTGCACACCCTGGACCCCGCCACCTGCCCCTTCGACCACCAGGCCAAGCACCGCATCGAGCGCGCC
CGCACCCGCATGGAGGCCGGCCTGGTGGACCAGGACGACCTGGACGAGGAGCACCAGGGCCTGGC
CCCCGCCGAGCTGTTCGCCCGCCTGAAGGCCCGCATGCCCGACGGCGAGGACCTGGTGGTGACCCA
CGGCGACGCCTGCCTGCCCAACATCATGGTGGAGAACGGCCGCTTCTCCGGCTTCATCGACTGCGGC
CGCCTGGGCGTGGCCGACCGCTACCAGGACATCGCCCTGGCCACCCGCGACATCGCCGAGGAGCTG
GGCGGCGAGTGGGCCGACCGCTTCCTGGTGCTGTACGGCATCGCCGCCCCCGACTCCCAGCGCATC
GCCTTCTACCGCCTGCTGGACGAGTTCTTCTGAcaattg*gcagcagcagctcggatagtatcgacacactctggac*
gctggtcgtgtgatggactgttgccgccacacttgctgccttgacctgtgaatatccctgccgcttttatcaaacagcctcagtgtg
tttgatcttgtgtgtacgcgcttttgcgagttgctagctgcttgtgctatttgcgaataccacccccagcatcccctttccctcgtttca
tatcgcttgcatcccaaccgcaacttatctacgctgtcctgctatccctcagcgctgctcctgctcctgctcactgcccctcgcacag
ccttggtttgggctccgcctgtattctcctggtactgcaacctgtaaaccagcactgcaatgctgatgcacgggaagtagtggga
*tgggaacacaaatgga*ggatccgcgtctcgaacagagcgcgcagaggaacgctgaaggtctcgcctctgtcgcacctcagcgc
ggcatacaccacaataaccacctgacgaatgcgcttggttcttcgtccattagcgaagcgtccggttcacacacgtgccacgttggc
gaggtggcaggtgacaatgatcggtggagctgatggtcgaaacgttcacagcctagggatatcgaattcctttcttgcgctatgaca
cttccagcaaaaggtagggcgggctgcgagacggcttcccggcgctgcatgcaacaccgatgatgcttcgacccccgaagctcctt
cggggctgcatgggcgctccgatgccgctccagggcgagcgctgtttaaatagccaggcccccgattgcaaagacattatagcgag
ctaccaaagccatattcaaacacctagatcactaccacttctacacaggccactcgagcttgtgatcgcactccgctaaggggcgc
ctcttcctcttcgtttcagtcacaacccgcaaacggcgcgccc**ATG**GCCACCACCTCCCTGGCCTCCGCCTTCTGCAGC
ATGAAGGCCGTGATGCTGGCCCGCGACGGCCGCGGCATGAAGCCCCGCTCCAGCGACCTGCAGCTG
CGCGCCGGCAACGCCCAGACCTCCCTGAAGATGATCAACGGCACCAAGTTCTCCTACACCGAGAGCC
TGAAGAAGCTGCCCGACTGGTCCATGCTGTTCGCCGTGATCACCACCATCTTCTCCGCCGCCGAGAA
GCAGTGGACCAACCTGGAGTGGAAGCCCAAGCCCAACCCCCCCCAGCTGCTGGACGACCACTTCGG
CCCCCACGGCCTGGTGTTCCGCCGCACCTTCGCCATCCGCAGCTACGAGGTGGGCCCCGACCGCTCC
ACCAGCATCGTGGCCGTGATGAACCACCTGCAGGAGGCCGCCCTGAACCACGCCAAGTCCGTGGGC
ATCCTGGGCGACGGCTTCGGCACCACCCTGGAGATGTCCAAGCGCGACCTGATCTGGGTGGTGAAG
CGCACCCACGTGGCCGTGGAGCGCTACCCCGCCTGGGGCGACACCGTGGAGGTGGAGTGCTGGGT
GGGCGCCTCCGGCAACAACGGCCGCCGCCACGACTTCCTGGTGCGCGACTGCAAGACCGGCGAGAT
CCTGACCCGCTGCACCTCCCTGAGCGTGATGATGAACACCCGCACCCGCCGCCTGAGCAAGATCCCC

GAGGAGGTGCGCGGCGAGATCGGCCCCGCCTTCATCGACAACGTGGCCGTGAAGGACGAGGAGAT
CAAGAAGCCCCAGAAGCTGAACGACTCCACCGCCGACTACATCCAGGGCGGCCTGACCCCCCGCTG
GAACGACCTGGACATCAACCAGCACGTGAACAACATCAAGTACGTGGACTGGATCCTGGAGACCGT
GCCCGACAGCATCTTCGAGAGCCACCACATCTCCTCCTTCACCATCGAGTACCGCCGCGAGTGCACC
ATGGACAGCGTGCTGCAGTCCCTGACCACCGTGAGCGGCGGCTCCTCCGAGGCCGGCCTGGTGTGC
GAGCACCTGCTGCAGCTGGAGGGCGGCAGCGAGGTGCTGCGCGCCAAGACCGAGTGGCGCCCCAA
GCTGACCGACTCCTTCCGCGGCATCAGCGTGATCCCCGCCGAGTCCAGCGTG*ATGGACTACAAGGA*
*CCACGACGGCGACTACAAGGACCACGACATCGACTACAAGGACGACGACGACAAGTGA***TGA**ctcgag
gcagcagcagctcggatagtatcgacacactctggacgctggtcgtgtgatggactgttgccgccacacttgctgccttgacct
gtgaatatccctgccgcttttatcaaacagcctcagtgtgtttgatcttgtgtgtacgcgcttttgcgagttgctagctgcttgtgct
atttgcgaataccacccccagcatcccccttccctcgtttcatatcgcttgcatcccaaccgcaacttatctacgctgtcctgctatcc
ctcagcgctgctcctgctcctgctcactgcccctcgcacagccttggtttgggctccgcctgtattctcctggtactgcaacctgtaa
*accagcactgcaatgctgatgcacgggaagtagtgggatgggaacacaaatgga*aagctt

SEQ ID NO: 124
C.reinhartii β-tubulin promoter::*P.moriformis* isopentenyl diphosphate synthase PTS::*Cinnamomum camphora* C14 thioesterase::*C.vulgaris* nitrate reductase 3'UTR

ggtacc**cgcctgcaacgcaagggcagccacagccgctcccacccgccgctgaaccgacacgtgcttgggcgcctgccgcctgcctg**
ccgcatgcttgtgctggtgaggctgggcagtgctgccatgctgattgaggcttggttcatcgggtggaagcttatgtgtgtgctggg
cttgcatgccgggcaatgcgcatggtggcaagagggcggcagcacttgctggagctgccgcggtgcctccaggtggttcaatcgc
ggcagccagagggatttcagatgatcgcgcgtacaggttgagcagcagtgtcagcaaaggtagcagtttgccagaatgatcggtt
cagctgttaatcaatgccagcaagagaaggggtcaagtgcaaacacgggcatgccacagcacgggcaccggggagtggaatgg
caccaccaagtgtgtgcgagccagcatcgccgcctggctgtttcagctacaacggcaggagtcatccaacgtaaccatgagctga
tcaacactgcaatcatcgggcgggcgtgatgcaagcatgcctggcgaagacacatggtgtgcggatgctgccggctgctgcctgct
gcgcacgccgttgagttggcagcaggctcagccatgcactggatggcagctgggctgccactgcaatgtggtggataggatgcaa
gtggagcgaataccaaaccctctggctgcttgctgggttgcatggcatcgcaccatcagcaggagcgcatgcgaagggactggcc
ccatgcacgccatgccaaaccggagcgcaccgagtgtccacactgtcaccaggcccgcaagctttgcagaaccatgctcatggac
gcatgtagcgctgacgtcccttgacggcgctcctctcgggtgtgggaaacgcaatgcagcacaggcagcagaggcggcggcagc
agagcggcggcagcagcggcgggggccacccttcttgcggggtcgcgccccagccagcggtgatgcgctgatcccaaacgagttc
acattcatttgcatgcctggagaagcgaggctggggcctttgggctggtgcagcccgcaatggaatgcgggaccgccaggctagc
agcaaaggcgcctcccctactccgcatcgatgttccatagtgcattggactgcatttgggtggggcggccggctgtttctttcgtgtt
gcaaaacgcgccagctcagcaacctgtcccgtgggtcccccgtgccgatgaaatcgtgtgcacgccgatcagctgattgcccggct
cgcgaagtaggcgccctcctttctgctcgccctctctccgtcccgcctctaga**atatcaATGATCGAGCAGGACGGCCTCC**
ACGCCGGCTCCCCCGCCGCCTGGGTGGAGCGCCTGTTCGGCTACGACTGGGCCCAGCAGACCATCG
GCTGCTCCGACGCCGCCGTGTTCCGCCTGTCCGCCCAGGGCCGCCCCGTGCTGTTCGTGAAGACCGA
CCTGTCCGGCGCCCTGAACGAGCTGCAGGACGAGGCCGCCCGCCTGTCCTGGCTGGCCACCACCGG
CGTGCCCTGCGCCGCCGTGCTGGACGTGGTGACCGAGGCCGGCCGCGACTGGCTGCTGCTGGGCG
AGGTGCCCGGCCAGGACCTGCTGTCCTCCCACCTGGCCCCCGCCGAGAAGGTGTCCATCATGGCCG
ACGCCATGCGCCGCCTGCACACCCTGGACCCCGCCACCTGCCCCTTCGACCACCAGGCCAAGCACCG
CATCGAGCGCGCCCGCACCCGCATGGAGGCCGGCCTGGTGGACCAGGACGACCTGGACGAGGAGC
ACCAGGGCCTGGCCCCCGCCGAGCTGTTCGCCCGCCTGAAGGCCCGCATGCCCGACGGCGAGGACC
TGGTGGTGACCCACGGCGACGCCTGCCTGCCCAACATCATGGTGGAGAACGGCCGCTTCTCCGGCT
TCATCGACTGCGGCCGCCTGGGCGTGGCCGACCGCTACCAGGACATCGCCCTGGCCACCCGCGACA
TCGCCGAGGAGCTGGGCGGCGAGTGGGCCGACCGCTTCCTGGTGCTGTACGGCATCGCCGCCCCCG
ACTCCCAGCGCATCGCCTTCTACCGCCTGCTGGACGAGTTCTTCTGA**caattg***gcagcagcagctcggatagt*
atcgacacactctggacgctggtcgtgtgatggactgttgccgccacacttgctgccttgacctgtgaatatccctgccgcttttat

caaacagcctcagtgtgtttgatcttgtgtgtacgcgcttttgcgagttgctagctgcttgtgctatttgcgaataccacccccagc
atccccttccctcgtttcatatcgcttgcatcccaaccgcaacttatctacgctgtcctgctatccctcagcgctgctcctgctcctgct
cactgcccctcgcacagccttggtttgggctccgcctgtattctcctggtactgcaacctgtaaaccagcactgcaatgctgatgc
*acgggaagtagtgggatgggaacacaaatgga*ggatcccgcgtctcgaacagagcgcgcagaggaacgctgaaggtctcgcc
tctgtcgcacctcagcgcggcatacaccacaataaccacctgacgaatgcgcttggttcttcgtccattagcgaagcgtccggttcac
acacgtgccacgttggcgaggtggcaggtgacaatgatcggtggagctgatggtcgaaacgttcacagcctagggatatcgaattc
ctttcttgcgctatgacacttccagcaaaaggtagggcgggctgcgagacggcttcccggcgctgcatgcaacaccgatgatgctt
cgacccccgaagctccttcggggctgcatgggcgctccgatgccgctccagggcgagcgctgtttaaatagccaggcccccgatt
gcaaagacattatagcgagctaccaaagccatattcaaacacctagatcactaccacttctacacaggccactcgagcttgtgatc
gcactccgctaaggggcgcctcttcctcttcgtttcagtcacaacccgcaaacactagt**ATG**ACGTTCGGGGTCGCCCTC
CCGGCCATGGGCCGCGGTGTCTCCCTTCCCCGGCCCAGGGTCGCGGTGCGCGCCCAGTCGGCGAGT
CAGGTTTTGGAGAGCGGGCGCGCCCCCGACTGGTCCATGCTGTTCGCCGTGATCACCACCATCTTCT
CCGCCGCCGAGAAGCAGTGGACCAACCTGGAGTGGAAGCCCAAGCCCAACCCCCCCCAGCTGCTGG
ACGACCACTTCGGCCCCCACGGCCTGGTGTTCCGCCGCACCTTCGCCATCCGCAGCTACGAGGTGGG
CCCCGACCGCTCCACCAGCATCGTGGCCGTGATGAACCACCTGCAGGAGGCCGCCCTGAACCACGC
CAAGTCCGTGGGCATCCTGGGCGACGGCTTCGGCACCACCCTGGAGATGTCCAAGCGCGACCTGAT
CTGGGTGGTGAAGCGCACCCACGTGGCCGTGGAGCGCTACCCCGCCTGGGGCGACACCGTGGAGG
TGGAGTGCTGGGTGGGCGCCTCCGGCAACAACGGCCGCCGCCACGACTTCCTGGTGCGCGACTGCA
AGACCGGCGAGATCCTGACCCGCTGCACCTCCCTGAGCGTGATGATGAACACCCGCACCCGCCGCCT
GAGCAAGATCCCCGAGGAGGTGCGCGGCGAGATCGGCCCCGCCTTCATCGACAACGTGGCCGTGA
AGGACGAGGAGATCAAGAAGCCCCAGAAGCTGAACGACTCCACCGCCGACTACATCCAGGGCGGC
CTGACCCCCCGCTGGAACGACCTGGACATCAACCAGCACGTGAACAACATCAAGTACGTGGACTGG
ATCCTGGAGACCGTGCCCGACAGCATCTTCGAGAGCCACCACATCTCCTCCTTCACCATCGAGTACCG
CCGCGAGTGCACCATGGACAGCGTGCTGCAGTCCCTGACCACCGTGAGCGGCGGCTCCTCCGAGGC
CGGCCTGGTGTGCGAGCACCTGCTGCAGCTGGAGGGCGGCAGCGAGGTGCTGCGCGCCAAGACCG
AGTGGCGCCCCAAGCTGACCGACTCCTTCCGCGGCATCAGCGTGATCCCCGCCGAGTCCAGCGTG*AT*
GGACTACAAGGACCACGACGGCGACTACAAGGACCACGACATCGACTACAAGGACGACGACGACAA
*GTGA***TGA**ctcgag*gcagcagcagctcggatagtatcgacacactctggacgctggtcgtgtgatggactgttgccgccaca*
cttgctgccttgacctgtgaatatccctgccgcttttatcaaacagcctcagtgtgtttgatcttgtgtgtacgcgcttttgcgagtt
gctagctgcttgtgctatttgcgaataccacccccagcatcccttccctcgtttcatatcgcttgcatcccaaccgcaacttatcta
cgctgtcctgctatccctcagcgctgctcctgctcctgctcactgcccctcgcacagccttggtttgggctccgcctgtattctcctgg
*tactgcaacctgtaaaccagcactgcaatgctgatgcacgggaagtagtgggatgggaacacaaatgga*aagctt

SEQ ID NO: 125

C.reinhartii β-tubulin promoter::*P.moriformis* delta 12 fatty acid desaturase
PTS::*Cinnamomum camphora* C14 thioesterase::*C.vulgaris* nitrate reductase 3'UTR

ggtacc**cgcctgcaacgcaagggcagccacagccgctcccacccgccgctgaaccgacacgtgcttgggcgcctgccgcctgcctg**
ccgcatgcttgtgctggtgaggctgggcagtgctgccatgctgattgaggcttggttcatcgggtggaagcttatgtgtgtgctggg
cttgcatgccgggcaatgcgcatggtggcaagagggcggcagcacttgctggagctgccgcggtgcctccaggtggttcaatcgc
ggcagccagagggatttcagatgatcgcgcgtacaggttgagcagcagtgtcagcaaaggtagcagtttgccagaatgatcggtt
cagctgttaatcaatgccagcaagagaaggggtcaagtgcaaacacgggcatgccacagcacgggcaccggggagtggaatgg
caccaccaagtgtgtgcgagccagcatcgccgcctggctgtttcagctacaacggcaggagtcatccaacgtaaccatgagctga
tcaacactgcaatcatcgggcgggcgtgatgcaagcatgcctggcgaagacacatggtgtgcggatgctgccggctgctgcctgct
gcgcacgccgttgagttggcagcaggctcagccatgcactggatggcagctgggctgccactgcaatgtggtggataggatgcaa
gtggagcgaataccaaaccctctggctgcttgctgggttgcatggcatcgcaccatcagcaggagcgcatgcgaagggactggcc
ccatgcacgccatgccaaaccggagcgcaccgagtgtccacactgtcaccaggcccgcaagctttgcagaaccatgctcatggac

gcatgtagcgctgacgtcccttgacggcgctcctctcgggtgtgggaaacgcaatgcagcacaggcagcagaggcggcggcagc
agagcggcggcagcagcggcgggggccacccttcttgcggggtcgcgccccagccagcggtgatgcgctgatcccaaacgagttc
acattcatttgcatgcctggagaagcgaggctggggcctttgggctggtgcagcccgcaatggaatgcgggaccgccaggctagc
agcaaaggcgcctcccctactccgcatcgatgttccatagtgcattggactgcatttgggtggggcggccggctgtttctttcgtgtt
gcaaaacgcgccagctcagcaacctgtcccgtgggtcccccgtgccgatgaaatcgtgtgcacgccgatcagctgattgcccggct
cgcgaagtaggcgccctcctttctgctcgcccctctctccgtcccgcctctaga**atatcaATGATCGAGCAGGACGGCCTCC**
ACGCCGGCTCCCCCGCCGCCTGGGTGGAGCGCCTGTTCGGCTACGACTGGGCCCAGCAGACCATCG
GCTGCTCCGACGCCGCCGTGTTCCGCCTGTCCGCCCAGGGCCGCCCCGTGCTGTTCGTGAAGACCGA
CCTGTCCGGCGCCCTGAACGAGCTGCAGGACGAGGCCGCCCGCCTGTCCTGGCTGGCCACCACCGG
CGTGCCCTGCGCCGCCGTGCTGGACGTGGTGACCGAGGCCGGCCGCGACTGGCTGCTGCTGGGCG
AGGTGCCCGGCCAGGACCTGCTGTCCTCCCACCTGGCCCCCGCCGAGAAGGTGTCCATCATGGCCG
ACGCCATGCGCCGCCTGCACACCCTGGACCCCGCCACCTGCCCCTTCGACCACCAGGCCAAGCACCG
CATCGAGCGCGCCCGCACCCGCATGGAGGCCGGCCTGGTGGACCAGGACGACCTGGACGAGGAGC
ACCAGGGCCTGGCCCCCGCCGAGCTGTTCGCCCGCCTGAAGGCCCGCATGCCCGACGGCGAGGACC
TGGTGGTGACCCACGGCGACGCCTGCCTGCCCAACATCATGGTGGAGAACGGCCGCTTCTCCGGCT
TCATCGACTGCGGCCGCCTGGGCGTGGCCGACCGCTACCAGGACATCGCCCTGGCCACCCGCGACA
TCGCCGAGGAGCTGGGCGGCGAGTGGGCCGACCGCTTCCTGGTGCTGTACGGCATCGCCGCCCCCG
ACTCCCAGCGCATCGCCTTCTACCGCCTGCTGGACGAGTTCTTCTGA**caattg***gcagcagcagctcggatagt*
atcgacacactctggacgctggtcgtgtgatggactgttgccgccacacttgctgccttgacctgtgaatatccctgccgcttttat
caaacagcctcagtgtgtttgatcttgtgtgtacgcgcttttgcgagttgctagctgcttgtgctatttgcgaataccacccccagc
atccccttccctcgtttcatatcgcttgcatcccaaccgcaacttatctacgctgtcctgctatccctcagcgctgctcctgctcctgct
cactgcccctcgcacagccttggtttgggctccgcctgtattctcctggtactgcaacctgtaaaccagcactgcaatgctgatgc
*acgggaagtagtgggatgggaacacaaatgga*ggatcc**cgcgtctcgaacagagcgcgcagaggaacgctgaaggtctcgcc**
tctgtcgcacctcagcgcggcatacaccacaataaccacctgacgaatgcgcttggttcttcgtccattagcgaagcgtccggttcac
acacgtgccacgttggcgaggtggcaggtgacaatgatcggtggagctgatggtcgaaacgttcacagcctagggatatcgaattc
ctttcttgcgctatgacacttccagcaaaaggtagggcgggctgcgagacggcttcccggcgctgcatgcaacaccgatgatgctt
cgaccccccgaagctccttcggggctgcatgggcgctccgatgccgctccagggcgagcgctgtttaaatagccaggccccgatt
gcaaagacattatagcgagctaccaaagccatattcaaacacctagatcactaccacttctacacaggccactcgagcttgtgatc
gcactccgctaaggggcgcctcttcctcttcgtttcagtcacaacccgcaaacactagt**ATG**GCTATCAAGACGAACAG
GCAGCCTGTGGAGAAGCCTCCGTTCACGATCGGGACGCTGCGCAAGGCCATCCCCGCGCACTGTTT
CGAGCGCTCGGCGCTTCGTGGGCGCGCCCCCGACTGGTCCATGCTGTTCGCCGTGATCACCACCATC
TTCTCCGCCGCCGAGAAGCAGTGGACCAACCTGGAGTGGAAGCCCAAGCCCAACCCCCCCCAGCTG
CTGGACGACCACTTCGGCCCCCACGGCCTGGTGTTCCGCCGCACCTTCGCCATCCGCAGCTACGAGG
TGGGCCCCGACCGCTCCACCAGCATCGTGGCCGTGATGAACCACCTGCAGGAGGCCGCCCTGAACC
ACGCCAAGTCCGTGGGCATCCTGGGCGACGGCTTCGGCACCACCCTGGAGATGTCCAAGCGCGACC
TGATCTGGGTGGTGAAGCGCACCCACGTGGCCGTGGAGCGCTACCCCGCCTGGGGCGACACCGTG
GAGGTGGAGTGCTGGGTGGGCGCCTCCGGCAACAACGGCCGCCGCCACGACTTCCTGGTGCGCGA
CTGCAAGACCGGCGAGATCCTGACCCGCTGCACCTCCCTGAGCGTGATGATGAACACCCGCACCCGC
CGCCTGAGCAAGATCCCCGAGGAGGTGCGCGGCGAGATCGGCCCCGCCTTCATCGACAACGTGGCC
GTGAAGGACGAGGAGATCAAGAAGCCCCAGAAGCTGAACGACTCCACCGCCGACTACATCCAGGG
CGGCCTGACCCCCCGCTGGAACGACCTGGACATCAACCAGCACGTGAACAACATCAAGTACGTGGA
CTGGATCCTGGAGACCGTGCCCGACAGCATCTTCGAGAGCCACCACATCTCCTCCTTCACCATCGAGT
ACCGCCGCGAGTGCACCATGGACAGCGTGCTGCAGTCCCTGACCACCGTGAGCGGCGGCTCCTCCG
AGGCCGGCCTGGTGTGCGAGCACCTGCTGCAGCTGGAGGGCGGCAGCGAGGTGCTGCGCGCCAAG
ACCGAGTGGCGCCCCAAGCTGACCGACTCCTTCCGCGGCATCAGCGTGATCCCCGCCGAGTCCAGC
GTG*ATGGACTACAAGGACCACGACGGCGACTACAAGGACCACGACATCGACTACAAGGACGACGAC*
*GACAAGTGA**TGA**ctcgag**gcagcagcagctcggatagtatcgacacactctggacgctggtcgtgtgatggactgttgcc*

gccacacttgctgccttgacctgtgaatatccctgccgcttttatcaaacagcctcagtgtgtttgatcttgtgtgtacgcgcttttg
cgagttgctagctgcttgtgctatttgcgaataccacccccagcatcccctttccctcgtttcatatcgcttgcatcccaaccgcaact
tatctacgctgtcctgctatccctcagcgctgctcctgctcctgctcactgcccctcgcacagccttggtttgggctccgcctgtattc
*tcctggtactgcaacctgtaaaccagcactgcaatgctgatgcacgggaagtagtgggatgggaacacaaatgga*aagctt

SEQ ID NO: 126
Homologous recombination construct SZ726

CCCGTGATCACACAGGTGCCTTGCGAGCGTGATCACACTATTTTGGGGGTCCTAC
AGTACTGAAATGGTGAGAAGTCGTACTGAAATCAAGGATGAACAATGAAAATGG
TGCTGTGGTGGCTTCTCAAAGGTCAAGAATCAGTCGCTCGCGTCAGGAAATCGCG
GCGTCAACCAGCGTGGGCGCGGTCAGTGGCCCCGCACTGGTCACCATAGCCTCTC
CTGCCACAGTAGCGATCCCCTGGGCGTTCACTCTCAGCAGCGGCTGTACTGCCTC
CCAGATTTTCTTCTTCTGGACCTGCGGGCGTGAGAGGATGAGCAGGGTGGGGCCA
AGGGCTCAATCCTGAACGGCCCTCATTCGGTTTCCAATCCCACAACACATACCCA
CAGCAGGTCAGACCACGCATTCGCACCATGCGCACCAAATAACGTGTCCTTACCT
GATTGGGTGTGGCAGGCTCCGTGGACAGGAGTGCCTCGTCCCCCGCCCAGACCC
GCTCCCCCCGTCACGGCGGCGTCCGGGACCCGCAGCGGCTCCACCGCGGTGTGATC
CGCGTTGGCGGCGCAGAGCAGCATCCCAGCCGATTTGACCCCGCGCATGCTCCG
AGGCTTGAGGTTGGCCAGCACCACCACCCGCCGGCCGACAAGGTCCTCCAGGGT
CACGTGCCGGACCAGGCCACTCACGATGGTGCGAGGGCCCCCCTCCTCGCCGAG
GTCGATCTGCTCGACGTACAGACTGCGACATGCGTGGCGAGTGGTCATCAGAAG
GAAGCAGGTGTGCAGAAGGGGCACGTGGTTGGTATTGAGAGTAGCCAAAGCTTT
GTGCCAATCAGAAAGTCAACGCAGCTGCCTGCCTGGCTCGCGTACAATTCCTTTC
TTGCGCTATGACACTTCCAGCAAAAGGTAGGGCGGGCTGCGAGACGGCTTCCCG
GCGCTGCATGCAACACCGATGATGCTTCGACCCCCCGAAGCTCCTTCGGGGCTGC
ATGGGCGCTCCGATGCCGCTCCAGGGCGAGCGCTGTTTAAATAGCCAGGCCCCC
GATTGCAAAGACATTATAGCGAGCTACCAAAGCCATATTCAAACACCTAGATCA
CTACCACTTCTACACAGGCCACTCGAGCTTGTGATCGCACTCCGCTAAGGGGGCG
CCTCTTCCTCTTCGTTTCAGTCACAACCCGCAAACGGCGCGCCATGCTGCTGCAG
GCCTTCCTGTTCCTGCTGGCCGGCTTCGCCGCCAAGATCAGCGCCTCCATGACGA
ACGAGACGTCCGACCGCCCCCTGGTGCACTTCACCCCCAACAAGGGCTGGATGA
ACGACCCCAACGGCCTGTGGTACGACGAGAAGGACGCCAAGTGGCACCTGTACT
TCCAGTACAACCCGAACGACACCGTCTGGGGGACGCCCTTGTTCTGGGGCCACGC
CACGTCCGACGACCTGACCAACTGGGAGGACCAGCCCATCGCCATCGCCCCGAA
GCGCAACGACTCCGGCGCCTTCTCCGGCTCCATGGTGGTGGACTACAACAACACC
TCCGGCTTCTTCAACGACACCATCGACCCGCGCCAGCGCTGCGTGGCCATCTGGA
CCTACAACACCCCGGAGTCCGAGGAGCAGTACATCTCCTACAGCCTGGACGGCG
GCTACACCTTCACCGAGTACCAGAAGAACCCCGTGCTGGCCGCCAACTCCACCCA
GTTCCGCGACCCGAAGGTCTTCTGGTACGAGCCCTCCCAGAAGTGGATCATGACC
GCGGCCAAGTCCCAGGACTACAAGATCGAGATCTACTCCTCCGACGACCTGAAG
TCCTGGAAGCTGGAGTCCGCGTTCGCCAACGAGGGCTTCCTCGGCTACCAGTACG
AGTGCCCCGGCCTGATCGAGGTCCCCACCGAGCAGGACCCCAGCAAGTCCTACT
GGGTGATGTTCATCTCCATCAACCCCGGCGCCCCGGCCGGCGGCTCCTTCAACCA
GTACTTCGTCGGCAGCTTCAACGGCACCCACTTCGAGGCCTTCGACAACCAGTCC
CGCGTGGTGGACTTCGGCAAGGACTACTACGCCCTGCAGACCTTCTTCAACACCG
ACCCGACCTACGGGAGCGCCCTGGGCATCGCGTGGGCCTCCAACTGGGAGTACT
CCGCCTTCGTGCCCACCAACCCCTGGCGCTCCTCCATGTCCCTCGTGCGCAAGTTC
TCCCTCAACACCGAGTACCAGGCCAACCCGGAGACGGAGCTGATCAACCTGAAG
GCCGAGCCGATCCTGAACATCAGCAACGCCGGCCCCTGGAGCCGGTTCGCCACC

AACACCACGTTGACGAAGGCCAACAGCTACAACGTCGACCTGTCCAACAGCACC
GGCACCCTGGAGTTCGAGCTGGTGTACGCCGTCAACACCACCCAGACGATCTCCA
AGTCCGTGTTCGCGGACCTCTCCCTCTGGTTCAAGGGCCTGGAGGACCCCGAGGA
GTACCTCCGCATGGGCTTCGAGGTGTCCGCGTCCTCCTTCTTCCTGGACCGCGGG
AACAGCAAGGTGAAGTTCGTGAAGGAGAACCCCTACTTCACCAACCGCATGAGC
GTGAACAACCAGCCCTTCAAGAGCGAGAACGACCTGTCCTACTACAAGGTGTAC
GGCTTGCTGGACCAGAACATCCTGGAGCTGTACTTCAACGACGGCGACGTCGTGT
CCACCAACACCTACTTCATGACCACCGGGAACGCCCTGGGCTCCGTGAACATGAC
GACGGGGGTGGACAACCTGTTCTACATCGACAAGTTCCAGGTGCGCGAGGTCAA
GTGATTAATTAACTCGAGGCAGCAGCAGCTCGGATAGTATCGACACACTCTGGA
CGCTGGTCGTGTGATGGACTGTTGCCGCCACACTTGCTGCCTTGACCTGTGAATA
TCCCTGCCGCTTTTATCAAACAGCCTCAGTGTGTTTGATCTTGTGTGTACGCGCTT
TTGCGAGTTGCTAGCTGCTTGTGCTATTTGCGAATACCACCCCCAGCATCCCCTTC
CCTCGTTTCATATCGCTTGCATCCCAACCGCAACTTATCTACGCTGTCCTGCTATC
CCTCAGCGCTGCTCCTGCTCCTGCTCACTGCCCCTCGCACAGCCTTGGTTTGGGCT
CCGCCTGTATTCTCCTGGTACTGCAACCTGTAAACCAGCACTGCAATGCTGATGC
ACGGGAAGTAGTGGGATGGGAACACAAATGGAAAGCTTGAGCTCGGTACCCGTA
CCCATCAGCATCCGGGTGAATCTTGGCCTCCAAGATATGGCCAATCCTCACATCC
AGCTTGGCAAAATCGACTAGACTGTCTGCAAGTGGGAATGTGGAGCACAAGGTT
GCTTGTAGCGATCGACAGACTGGTGGGGTACATTGACAGGTGGGCAGCGCCGCA
TCCATCGTGCCTGACGCGAGCGCCGCCGGTTGCTCGCCCGTGCCTGCCGTCAAAG
AGCGGCAGAGAAATCGGGAACCGAAAACGTCACATTGCCTGATGTTGTTACATG
CTGGACTAGACTTTCTTGGCGTGGGTCTGCTCCTCGCCAGGTGCGCGACGCCTCG
GGGCTGGGTGCGAGGGAGCCGTGCGGCCACGCATTTGACAAGACCCAAAGCTCG
CATCTCAGACGGTCAACCGTTCGTATTATACATTCAACATATGGTACATACGCAA
AAAGCATG

SEQ ID NO: 127
P.moriformis isopentenyl diphosphate synthase plastid targeting sequence

MTFGVALPAMGRGVSLPRPRVAVRAQSASQVLESGRAQL

SEQ ID NO: 128
P.moriformis delta 12 fatty acid desaturase plastid targeting sequence

MAIKTNRQPVEKPPFTIGTLRKAIPAHCFERSALRGRAQL

SEQ ID NO: 129
P.moriformis stearoyl ACP desaturase plastid targeting sequence

MASAAFTMSACPAMTGRAPGARRSGRPVATRLRGRA

SEQ ID NO: 130
C.protothecoides stearoyl ACP desaturase plastid targeting sequence

MATASTFSAFNARCGDLRRSAGSGPRRPARPLPVRGRAQL

SEQ ID NO: 131
Cuphea hookeriana fatty acyl-ACP thioesterase (C8-10) plastid targeting sequence

MVAAAASSAFFPVPAPGASPKPGKFGNWPSSLSPSFKPKSIPNGGFQVKANDSAHPK
ANGSAVSLKSGSLNTQEDTSSSPPPRTFLH

SEQ ID NO: 132
Umbellularia californica fatty acyl-ACP thioesterase (C12) plastid targeting sequence

MATTSLASAFCSMKAVMLARDGRGMKPRSSDLQLRAGNAPTSLKMINGTKFSYTES
LKRL

SEQ ID NO: 133
Cinnamomum camphora fatty acyl-ACP thioesterase (C14) plastid targeting sequence

MATTSLASAFCSMKAVMLARDGRGMKPRSSDLQLRAGNAQTSLKMINGTKFSYTES
LKKL

SEQ ID NO: 134
P.moriformis fatty acyl-ACP thioesterase-1 cDNA sequence

ATGGCACCGACCAGCCTGCTTGCCAGTACTGGCGTCTCTTCCGCTTCTCTGTGGTC
CTCTGCGCGCTCCAGCGCGTGCGCTTTTCCGGTGGATCATGCGGTCCGTGGCGCA
CCGCAGCGGCCGCTGCCCATGCAGCGCCGCTGCTTCCGAACAGTGGCGGTCAGG
GCCGCACCCGCGGTAGCCGTCCGTCCGGAACCCGCCCAAGAGTTTTGGGAGCAG
CTTGAGCCCTGCAAGATGGCGGAGGACAAGCGCATCTTCCTGGAGGAGCACCGC
ATTCGGGGCAACGAGGTGGGCCCCTCGCAGCGGCTGACGATCACGGCGGTGGCC
AACATCCTGCAGGAGGCGGCGGGCAACCACGCGGTGGCCATGTGGGGCCGGAGC
TCGGAGGGTTTCGCGACGGACCCGGAGCTGCAGGAGGCGGGTCTCATCTTTGTG
ATGACGCGCATGCAGATCCAAATGTACCGCTACCCGCGCTGGGGCGACCTGATG
CAGGTGGAGACCTGGTTCCAGACGGCGGGCAAGCTAGGCGCGCAGCGCGAGTGG
GTGCTGCGCGACAAGCTGACCGGCGAGGCGCTGGGCGCGGCCACCTCCAGCTGG
GTCATGATCAACATCCGCACGCGCCGGCCGTGCCGCATGCCCGAGCTCGTCCGCG
TCAAGTCGGCCTTCTTCGCGCGCGAGCCGCCGCGCCTGGCGCTGCCGCCCACGGT
CACGCGCGCCAAGCTGCCCAACATCGCGACGCCGGCGCCGCTGCGCGGGCACCG
CCAGGTCGCGCGCCGCACCGACATGGACATGAACGGGCACGTGAACAACGTGGC
CTACCTGGCCTGGTGCCTGGAGGCCGTGCCCGAGCACGTCTTCAGCGACTACCAC
CTCTACCAGATGGAGATCGACTTCAAGGCCGAGTGCCACGCGGGCGACGTCATC
TCCTCCCAGGCCGAGCAGATCCCGCCCCAGGAGGCGCTCACGCACAACGGCGCC
GGCCGCAACCCCTCCTGCTTCGTCCATAGCATTCTGCGCGCCGAGACCGAGCTCG
TCCGCGCGCGAACCACATGGTCGGCCCCCATCGACGCGCCCGCCGCCAAGCCGC
CCAAGGCGAGCCACTGA

SEQ ID NO: 135
P.moriformis fatty acyl-ACP thioesterase-2 cDNA sequence

ATGGCACCGACCAGCCTGCTTGCCCGTACTGGCGTCTCTTCCGCTTCTCTGTGCTC
CTCTACGCGCTCCGGCGCGTGCGCTTTTCCGGTGGATCATGCGGTCCGTGGCGCA
CCGCAGCGGCCGCTGCCCATGCAGCGCCGCTGCTTCCGAACAGTGGCTGTCAGG
GCCGCACCCGCAGTAGCCGTCCGTCCGGAACCCGCCCAAGAGTTTTGGGAGCAG
CTTGAGCCCTGCAAGATGGCGGAGGACAAGCGCATCTTCCTGGAGGAGCACCGC
ATTCGTGGCAACGAGGTGGGCCCCTCGCAGCGGCTGACGATCACGGCGGTGGCC
AACATCCTGCAGGAGGCGGCGGGCAACCACGCGGTGGCCATGTGGGGTCGGAGC

TCGGAGGGTTTCGCGACGGACCCGGAGCTGCAGGAGGCGGGCCTCATCTTTGTG
ATGACGCGCATGCAGATCCAAATGTACCGCTACCCGCGCTGGGGCGACCTGATG
CAGGTGGAGACCTGGTTCCAGACGGCGGGCAAGCTAGGCGCGCAGCGCGAGTGG
GTGCTGCGCGACAAGCTGACCGGCGAGGCGCTGGGCGCGGCCACCTCCAGCTGG
GTCATGATCAACATCCGCACGCGCCGGCCGTGCCGCATGCCCGAGCTCGTCCGCG
TCAAGTCGGCCTTCTTCGCGCGCGAGCCGCCGCGCCTGGCGCTGCCGCCCGCGGT
CACGCGTGCCAAGCTGCCCAACATCGCGACGCCGGCGCCGCTGCGCGGGCACCG
CCAGGTCGCGCGCCGCACCGACATGGACATGAACGGCCACGTGAACAACGTTGC
CTACCTGGCCTGGTGCCTGGAGGCCGTGCCCGAGCACGTCTTCAGCGACTACCAC
CTCTACCAGATGGAGATCGACTTCAAGGCCGAGTGCCACGCGGGCGACGTCATC
TCCTCCCAGGCCGAGCAGATCCCGCCCCAGGAGGCGCTCACGCACAACGGCGCC
GGCCGCAACCCCTCCTGCTTCGTCCATAGCATTCTGCGCGCCGAGACCGAGCTCG
TCCGCGCGCGAACCACATGGTCGGCCCCCATCGACGCGCCCGCCGCCAAGCCGC
CCAAGGCGAGCCACTGA

SEQ ID NO: 136
P.moriformis fatty acyl-ACP thioesterase-1 amino acid sequence

MAPTSLLASTGVSSASLWSSARSSACAFPVDHAVRGAPQRPLPMQRRCFRTVAVRA
APAVAVRPEPAQEFWEQLEPCKMAEDKRIFLEEHRIRGNEVGPSQRLTITAVANILQE
AAGNHAVAMWGRSSEGFATDPELQEAGLIFVMTRMQIQMYRYPRWGDLMQVETW
FQTAGKLGAQREWVLRDKLTGEALGAATSSWVMINIRTRRPCRMPELVRVKSAFFA
REPPRLALPPTVTRAKLPNIATPAPLRGHRQVARRTDMDMNGHVNNVAYLAWCLE
AVPEHVFSDYHLYQMEIDFKAECHAGDVISSQAEQIPPQEALTHNGAGRNPSCFVHSI
LRAETELVRARTTWSAPIDAPAAKPPKASH

SEQ ID NO: 137
P.moriformis fatty acyl-ACP thioesterase-2 amino acid sequence

MAPTSLLARTGVSSASLCSSTRSGACAFPVDHAVRGAPQRPLPMQRRCFRTVAVRA
APAVAVRPEPAQEFWEQLEPCKMAEDKRIFLEEHRIRGNEVGPSQRLTITAVANILQE
AAGNHAVAMWGRSSEGFATDPELQEAGLIFVMTRMQIQMYRYPRWGDLMQVETW
FQTAGKLGAQREWVLRDKLTGEALGAATSSWVMINIRTRRPCRMPELVRVKSAFFA
REPPRLALPPAVTRAKLPNIATPAPLRGHRQVARRTDMDMNGHVNNVAYLAWCLE
AVPEHVFSDYHLYQMEIDFKAECHAGDVISSQAEQIPPQEALTHNGAGRNPSCFVHSI
LRAETELVRARTTWSAPIDAPAAKPPKASH

SEQ ID NO: 138
C.hookeriana fatty acyl-ACP thioesterase amino acid sequence without PTS

QLPDWSRLLTAITTVFVKSKRPDMHDRKSKRPDMLVDSFGLESTVQDGLVFRQSFSI
RSYEIGTDRTASIETLMNHLQETSLNHCKSTGILLDGFGRTLEMCKRDLIWVVIKMQI
KVNRYPAWGDTVEINTRFSRLGKIGMGRDWLISDCNTGEILVRATSAYAMMNQKTR
RLSKLPYEVHQEIVPLFVDSPVIEDSDLKVHKFKVKTGDSIQKGLTPGWNDLDVNQH
VSNVKYIGWILESMPTEVLETQELCSLALEYRRECGRDSVLESVTAMDPSKVGVRSQ
YQHLLRLEDGTAIVNGATEWRPKNAGANGAISTGKTSNGNSVS

SEQ ID NO: 139
U.californica fatty acyl-ACP thioesterase amino acid sequence without PTS

PDWSMLFAVITTIFSAAEKQWTNLEWKPKPKLPQLLDDHFGLHGLVFRRTFAIRSYE
VGPDRSTSILAVMNHMQEATLNHAKSVGILGDGFGTTLEMSKRDLMWVVRRTHVA
VERYPTWGDTVEVECWIGASGNNGMRRDFLVRDCKTGEILTRCTSLSVLMNTRTRR
LSTIPDEVRGEIGPAFIDNVAVKDDEIKKLQKLNDSTADYIQGGLTPRWNDLDVNQH
VNNLKYVAWVFETVPDSIFESHHISSFTLEYRRECTRDSVLRSLTTVSGGSSEAGLVC
DHLLQLEGGSEVLRARTEWRPKLTDSFRGISVIPAEPRV

SEQ ID NO: 140
Cinnamomum camphora fatty acyl-ACP thioesterase amino acid sequence without PTS

PDWSMLFAVITTIFSAAEKQWTNLEWKPKPNPPQLLDDHFGPHGLVFRRTFAIRSYE
VGPDRSTSIVAVMNHLQEAALNHAKSVGILGDGFGTTLEMSKRDLIWVVKRTHVAV
ERYPAWGDTVEVECWVGASGNNGRRHDFLVRDCKTGEILTRCTSLSVMMNTRTRR
LSKIPEEVRGEIGPAFIDNVAVKDEEIKKPQKLNDSTADYIQGGLTPRWNDLDINQHV
NNIKYVDWILETVPDSIFESHHISSFTIEYRRECTMDSVLQSLTTVSGGSSEAGLVCEH
LLQLEGGSEVLRAKTEWRPKLTDSFRGISVIPAESSVMDYKDHDGDYKDHDIDYKD
DDDK

SEQUENCE LISTING

<110> SOLAZYME, INC.

<120> MANUFACTURING OF TAILORED OILS IN RECOMBINANT HETEROTROPHIC MICROORGANISMS

<130> 026172-005020PC

<140> PCT/US2009/066141
<141> 2009-11-30

<150> 61/219,525
<151> 2009-06-23

<150> 61/174,357
<151> 2009-04-30

<150> 61/118,994
<151> 2008-12-01

<150> 61/118,590
<151> 2008-11-28

<160> 143

<170> PatentIn version 3.5

<210> 1
<211> 1187
<212> DNA
<213> Chlorella sp.

<400> 1
gatcagacgg gcctgacctg cgagataatc aagtgctcgt aggcaaccaa ctcagcagct 60

gcttggtgtt gggtctgcag gatagtgttg cagggcccca aggacagcag gggaacttac 120

accttgtccc cgacccagtt ttatggagtg cattgcctca agagcctagc cggagcgcta 180

ggctacatac ttgccgcacc ggtatgaggg gatatagtac tcgcactgcg ctgtctagtg 240

agatgggcag tgctgcccat aaacaactgg ctgctcagcc atttgttggc ggaccattct 300

gggggggcca gcaatgcctg actttcgggt agggtgaaaa ctgaacaaag actaccaaaa 360

cagaatttct tcctccttgg aggtaagcgc aggccggccc gcctgcgccc acatggcgct 420

ccgaacacct ccatagctgt aagggcgcaa acatggccgg actgttgtca gcactctttc 480

atggccatac aaggtcatgt cgagattagt gctgagtaag acactatcac cccatgttcg 540

attgaagccg tgacttcatg ccaacctgcc cctgggcgta gcagacgtat gccatcatga 600

ccactagccg acatgcgctg tcttttgcca ccaaaacaac tggtacaccg ctcgaagtcg 660

tgccgcacac ctccgggagt gagtccggcg actcctcccc ggcgggccgc ggccctacct 720

gggtagggtc gccatacgcc cacgaccaaa cgacgcagga ggggattggg gtagggaatc 780

ccaaccagcc taaccaagac ggcacctata ataataggtg gggggactaa cagccctata 840

tcgcaagctt tgggtgccta tcttgagaag cacgagttgg agtggctgtg tacggtcgac 900

cctaaggtgg gtgtgccgca gcctgaaaca aagcgtctag cagctgcttc tataatgtgt 960

cagccgttgt gtttcagtta tattgtatgc tattgtttgt tcgtgctagg gtggcgcagg 1020

cccacctact gtggcgggcc attggttggt gcttgaattg cctcaccatc taaggtctga 1080

acgctcactc aaacgccttt gtacaactgc agaactttcc ttggcgctgc aactacagtg 1140

tgcaaaccag cacatagcac tcccttacat cacccagcag tacaaca 1187

<210> 2
<211> 1414
<212> DNA
<213> Chlorella ellipsoidea

<400> 2
cgctgcgcac cagggccgcc agctcgctga tgtcgctcca aatgcggtcc cccgattttt 60

tgttcttcat cttctccacc ttggtggcct tcttggccag ggccttcagc tgcatgcgca 120

cagaccgttg agctcctgat cagcatcctc aggaggccct ttgacaagca agcccctgtg 180

caagcccatt cacggggtac cagtggtgct gaggtagatg ggtttgaaaa ggattgctcg 240

gtcgattgct gctcatggaa ttggcatgtg catgcatgtt cacaatatgc caccaggctt 300

tggagcaaga gagcatgaat gccttcaggc aggttgaaag ttcctggggg tgaagaggca 360

gggccgagga ttggaggagg aaagcatcaa gtcgtcgctc atgctcatgt tttcagtcag 420

agtttgccaa gctcacagga gcagagacaa gactggctgc tcaggtgttg catcgtgtgt 480

gtggtggggg ggggggggtt aatacggtac gaaatgcact tggaattccc acctcatgcc 540

agcggaccca catgcttgaa ttcgaggcct gtggggtgag aaatgctcac tctgccctcg 600

ttgctgaggt acttcaggcc gctgagctca aagtcgatgc cctgctcgtc tatcagggcc 660

tgcacctctg ggctgaccgg ctcagcctcc ttcgcgggca tggagtaggc gccggcagcg 720

ttcatgtccg ggcccagggc agcggtggtg ccataaatgt cggtgatggt ggggaggggg 780

gccgtcgcca caccattgcc gttgctggct gacgcatgca catgtggcct ggctggcacc 840

ggcagcactg gtctccagcc agccagcaag tggctgttca ggaaagcggc catgttgttg 900

gtccctgcgc atgtaattcc ccagatcaaa ggagggaaca gcttggattt gatgtagtgc 960

ccaaccggac tgaatgtgcg atggcaggtc cctttgagtc tcccgaatta ctagcagggc 1020

actgtgacct aacgcagcat gccaaccgca aaaaaatgat tgacagaaaa tgaagcggtg 1080

tgtcaatatt tgctgtattt attcgtttta atcagcaacc aagttcgaaa cgcaactatc 1140

gtggtgatca agtgaacctc atcagactta cctcgttcgg caaggaaacg gaggcaccaa 1200

attccaattt gatattatcg cttgccaagc tagagctgat ctttgggaaa ccaactgcca 1260

gacagtggac tgtgatggag tgccccgagt ggtggagcct cttcgattcg gttagtcatt 1320

actaacgtga accctcagtg aagggaccat cagaccagaa agaccagatc tcctcctcga 1380

caccgagaga gtgttgcggc agtaggacga caag 1414

<210> 3
<211> 512
<212> PRT
<213> Unknown

<220>

<223> Description of Unknown: Yeast sequence

<400> 3

```
Met Thr Asn Glu Thr Ser Asp Arg Pro Leu Val His Phe Thr Pro Asn
1               5                  10                  15

Lys Gly Trp Met Asn Asp Pro Asn Gly Leu Trp Tyr Asp Glu Lys Asp
            20                  25                  30

Ala Lys Trp His Leu Tyr Phe Gln Tyr Asn Pro Asn Asp Thr Val Trp
        35                  40                  45

Gly Thr Pro Leu Phe Trp Gly His Ala Thr Ser Asp Asp Leu Thr Asn
    50                  55                  60

Trp Glu Asp Gln Pro Ile Ala Ile Ala Pro Lys Arg Asn Asp Ser Gly
65                  70                  75                  80

Ala Phe Ser Gly Ser Met Val Val Asp Tyr Asn Asn Thr Ser Gly Phe
                85                  90                  95

Phe Asn Asp Thr Ile Asp Pro Arg Gln Arg Cys Val Ala Ile Trp Thr
            100                 105                 110

Tyr Asn Thr Pro Glu Ser Glu Glu Gln Tyr Ile Ser Tyr Ser Leu Asp
        115                 120                 125

Gly Gly Tyr Thr Phe Thr Glu Tyr Gln Lys Asn Pro Val Leu Ala Ala
    130                 135                 140

Asn Ser Thr Gln Phe Arg Asp Pro Lys Val Phe Trp Tyr Glu Pro Ser
145                 150                 155                 160

Gln Lys Trp Ile Met Thr Ala Ala Lys Ser Gln Asp Tyr Lys Ile Glu
                165                 170                 175

Ile Tyr Ser Ser Asp Asp Leu Lys Ser Trp Lys Leu Glu Ser Ala Phe
            180                 185                 190

Ala Asn Glu Gly Phe Leu Gly Tyr Gln Tyr Glu Cys Pro Gly Leu Ile
        195                 200                 205

Glu Val Pro Thr Glu Gln Asp Pro Ser Lys Ser Tyr Trp Val Met Phe
    210                 215                 220

Ile Ser Ile Asn Pro Gly Ala Pro Ala Gly Gly Ser Phe Asn Gln Tyr
225                 230                 235                 240

Phe Val Gly Ser Phe Asn Gly Thr His Phe Glu Ala Phe Asp Asn Gln
                245                 250                 255

Ser Arg Val Val Asp Phe Gly Lys Asp Tyr Tyr Ala Leu Gln Thr Phe
```

260 265 270

Phe Asn Thr Asp Pro Thr Tyr Gly Ser Ala Leu Gly Ile Ala Trp Ala
275 280 285

Ser Asn Trp Glu Tyr Ser Ala Phe Val Pro Thr Asn Pro Trp Arg Ser
290 295 300

Ser Met Ser Leu Val Arg Lys Phe Ser Leu Asn Thr Glu Tyr Gln Ala
305 310 315 320

Asn Pro Glu Thr Glu Leu Ile Asn Leu Lys Ala Glu Pro Ile Leu Asn
325 330 335

Ile Ser Asn Ala Gly Pro Trp Ser Arg Phe Ala Thr Asn Thr Thr Leu
340 345 350

Thr Lys Ala Asn Ser Tyr Asn Val Asp Leu Ser Asn Ser Thr Gly Thr
355 360 365

Leu Glu Phe Glu Leu Val Tyr Ala Val Asn Thr Thr Gln Thr Ile Ser
370 375 380

Lys Ser Val Phe Ala Asp Leu Ser Leu Trp Phe Lys Gly Leu Glu Asp
385 390 395 400

Pro Glu Glu Tyr Leu Arg Met Gly Phe Glu Val Ser Ala Ser Ser Phe
405 410 415

Phe Leu Asp Arg Gly Asn Ser Lys Val Lys Phe Val Lys Glu Asn Pro
420 425 430

Tyr Phe Thr Asn Arg Met Ser Val Asn Asn Gln Pro Phe Lys Ser Glu
435 440 445

Asn Asp Leu Ser Tyr Tyr Lys Val Tyr Gly Leu Leu Asp Gln Asn Ile
450 455 460

Leu Glu Leu Tyr Phe Asn Asp Gly Asp Val Val Ser Thr Asn Thr Tyr
465 470 475 480

Phe Met Thr Thr Gly Asn Ala Leu Gly Ser Val Asn Met Thr Thr Gly
485 490 495

Val Asp Asn Leu Phe Tyr Ile Asp Lys Phe Gln Val Arg Glu Val Lys
500 505 510

<210> 4
<211> 20
<212> PRT
<213> Unknown

<220>
<223> Description of Unknown: Yeast sequence

<400> 4
Met Leu Leu Gln Ala Phe Leu Phe Leu Leu Ala Gly Phe Ala Ala Lys
1 5 10 15

Ile Ser Ala Ser
20

<210> 5
<211> 18
<212> PRT
<213> Unknown

<220>
<223> Description of Unknown: Higher plant secretion signal

<400> 5
Met Ala Asn Lys Ser Leu Leu Leu Leu Leu Leu Leu Gly Ser Leu Ala
1 5 10 15

Ser Gly

<210> 6
<211> 10
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic consensus sequence

<400> 6
Met Ala Arg Leu Pro Leu Ala Ala Leu Gly
1 5 10

<210> 7
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic peptide

<400> 7
Met Ala Asn Lys Leu Leu Leu Leu Leu Leu Leu Leu Leu Leu Pro Leu
1 5 10 15

Ala Ala Ser Gly
20

<210> 8
<211> 2615
<212> DNA
<213> Saccharomyces cerevisiae

<400> 8
gaattcccca acatggtgga gcacgacact ctcgtctact ccaagaatat caaagataca 60

gtctcagaag accaaagggc tattgagact tttcaacaaa gggtaatatc gggaaacctc 120
ctcggattcc attgcccagc tatctgtcac ttcatcaaaa ggacagtaga aaaggaaggt 180
ggcacctaca aatgccatca ttgcgataaa ggaaaggcta tcgttcaaga tgcctctgcc 240
gacagtggtc ccaaagatgg acccccaccc acgaggagca tcgtggaaaa agaagacgtt 300
ccaaccacgt cttcaaagca agtggattga tgtgaacatg gtggagcacg acactctcgt 360
ctactccaag aatatcaaag atacagtctc agaagaccaa agggctattg agacttttca 420
acaaagggta atatcgggaa acctcctcgg attccattgc ccagctatct gtcacttcat 480
caaaaggaca gtagaaaagg aaggtggcac ctacaaatgc catcattgcg ataaaggaaa 540
ggctatcgtt caagatgcct ctgccgacag tggtcccaaa gatggacccc cacccacgag 600
gagcatcgtg gaaaaagaag acgttccaac cacgtcttca aagcaagtgg attgatgtga 660
tatctccact gacgtaaggg atgacgcaca atcccactat ccttcgcaag acccttcctc 720
tatataagga agttcatttc atttggagag gacacgctga aatcaccagt ctctctctac 780
aaatctatct ctggcgcgcc atatcaatgc ttcttcaggc ctttcttttt cttcttgctg 840
gttttgctgc caagatcagc gcctctatga cgaacgaaac ctcggataga ccacttgtgc 900
actttacacc aaacaagggc tggatgaatg accccaatgg actgtggtac gacgaaaaag 960
atgccaagtg gcatctgtac tttcaataca acccgaacga tactgtctgg gggacgccat 1020
tgttttgggg ccacgccacg tccgacgacc tgaccaattg ggaggaccaa ccaatagcta 1080
tcgctccgaa gaggaacgac tccggagcat tctcgggttc catggtggtt gactacaaca 1140
atacttccgg ctttttcaac gataccattg acccgagaca acgctgcgtg gccatatgga 1200
cttacaacac accggagtcc gaggagcagt acatctcgta tagcctggac ggtggataca 1260
cttttacaga gtatcagaag aaccctgtgc ttgctgcaaa ttcgactcag ttccgagatc 1320
cgaaggtctt ttggtacgag ccctcgcaga agtggatcat gacagcggca aagtcacagg 1380
actacaagat cgaaatttac tcgtctgacg accttaaatc ctggaagctc gaatccgcgt 1440
tcgcaaacga gggctttctc ggctaccaat acgaatgccc aggcctgata gaggtcccaa 1500
cagagcaaga tcccagcaag tcctactggg tgatgtttat ttccattaat ccaggagcac 1560
cggcaggagg ttcttttaat cagtacttcg tcggaagctt taacggaact catttcgagg 1620
catttgataa ccaatcaaga gtagttgatt ttggaaagga ctactatgcc ctgcagactt 1680
tcttcaatac tgacccgacc tatgggagcg ctcttggcat tgcgtgggct tctaactggg 1740
agtattccgc attcgttcct acaaaccctt ggaggtcctc catgtcgctc gtgaggaaat 1800
tctctctcaa cactgagtac caggccaacc cggaaaccga actcataaac ctgaaagccg 1860
aaccgatcct gaacattagc aacgctggcc cctggagccg gtttgcaacc aacaccacgt 1920
tgacgaaagc caacagctac aacgtcgatc tttcgaatag caccggtaca cttgaatttg 1980
aactggtgta tgccgtcaat accacccaaa cgatctcgaa gtcggtgttc gcggacctct 2040
ccctctggtt taaaggcctg gaagaccccg aggagtacct cagaatgggt ttcgaggttt 2100

ctgcgtcctc cttcttcctt gatcgcggga acagcaaagt aaaatttgtt aaggagaacc 2160

catattttac caacaggatg agcgttaaca accaaccatt caagagcgaa aacgacctgt 2220

cgtactacaa agtgtatggt ttgcttgatc aaaatatcct ggaactctac ttcaacgatg 2280

gtgatgtcgt gtccaccaac acatacttca tgacaaccgg gaacgcactg ggctccgtga 2340

acatgacgac gggtgtggat aacctgttct acatcgacaa attccaggtg agggaagtca 2400

agtgagatct gtcgatcgac aagctcgagt ttctccataa taatgtgtga gtagttccca 2460

gataagggaa ttagggttcc tatagggttt cgctcatgtg ttgagcatat aagaaaccct 2520

tagtatgtat ttgtatttgt aaaatacttc tatcaataaa atttctaatt cctaaaacca 2580

aaatccagta ctaaaatcca gatcccccga attaa 2615

<210> 9
<211> 22
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic primer

<400> 9
tgttgaagaa tgagccggcg ac 22

<210> 10
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic primer

<400> 10
cagtgagcta ttacgcactc 20

<210> 11
<211> 541
<212> DNA
<213> Prototheca kruegani

<400> 11
tgttgaagaa tgagccggcg agttaaaaag agtggcatgg ttaaagaaaa tactctggag 60

ccatagcgaa agcaagttta gtaagcttag gtcattcttt ttagacccga aaccgagtga 120

tctacccatg atcagggtga agtgttagta aaataacatg gaggcccgaa ccgactaatg 180

ttgaaaaatt agcggatgaa ttgtgggtag gggcgaaaaa ccaatcgaac tcggagttag 240

ctggttctcc ccgaaatgcg tttaggcgca gcagtagcag tacaaataga ggggtaaagc 300

actgtttctt ttgtgggctt cgaaagttgt acctcaaagt ggcaaactct gaatactcta 360

tttagatatc tactagtgag accttggggg ataagctcct tggtcaaaag ggaaacagcc 420

cagatcacca gttaaggccc caaaatgaaa atgatagtga ctaaggatgt gggtatgtca 480

aaacctccag caggttagct tagaagcagc aatcctttca agagtgcgta atagctcact 540

g 541

<210> 12
<211> 573
<212> DNA
<213> Prototheca wickerhamii

<400> 12
tgttgaagaa tgagccggcg acttaaaata aatggcaggc taagagattt aataactcga 60

aacctaagcg aaagcaagtc ttaatagggc gtcaatttaa caaaacttta aataaattat 120

aaagtcattt attttagacc cgaacctgag tgatctaacc atggtcagga tgaaacttgg 180

gtgacaccaa gtggaagtcc gaaccgaccg atgttgaaaa atcggcggat gaactgtggt 240

tagtggtgaa ataccagtcg aactcagagc tagctggttc tccccgaaat gcgttgaggc 300

gcagcaatat atctcgtcta tctaggggta aagcactgtt tcggtgcggg ctatgaaaat 360

ggtaccaaat cgtggcaaac tctgaatact agaaatgacg atatattagt gagactatgg 420

gggataagct ccatagtcga gagggaaaca gcccagacca ccagttaagg ccccaaaatg 480

ataatgaagt ggtaaaggag gtgaaaatgc aaatacaacc aggaggttgg cttagaagca 540

gccatccttt aaagagtgcg taatagctca ctg 573

<210> 13
<211> 541
<212> DNA
<213> Prototheca stagnora

<400> 13
tgttgaagaa tgagccggcg agttaaaaaa aatggcatgg ttaaagatat ttctctgaag 60

ccatagcgaa agcaagtttt acaagctata gtcatttttt ttagacccga aaccgagtga 120

tctacccatg atcagggtga agtgttggtc aaataacatg gaggcccgaa ccgactaatg 180

gtgaaaaatt agcggatgaa ttgtgggtag gggcgaaaaa ccaatcgaac tcggagttag 240

ctggttctcc ccgaaatgcg tttaggcgca gcagtagcaa cacaaataga ggggtaaagc 300

actgtttctt ttgtgggctt cgaaagttgt acctcaaagt ggcaaactct gaatactcta 360

tttagatatc tactagtgag accttggggg ataagctcct tggtcaaaag ggaaacagcc 420

cagatcacca gttaaggccc caaaatgaaa atgatagtga ctaaggacgt gagtatgtca 480

aaacctccag caggttagct tagaagcagc aatcctttca agagtgcgta atagctcact 540

g 541

<210> 14
<211> 541
<212> DNA
<213> Prototheca moriformis

<400> 14
tgttgaagaa tgagccggcg agttaaaaag agtggcatgg ttaaagataa ttctctggag 60

ccatagcgaa agcaagttta acaagctaaa gtcacccttt ttagacccga aaccgagtga 120

tctacccatg atcagggtga agtgttggta aaataacatg gaggcccgaa ccgactaatg 180

gtgaaaaatt agcggatgaa ttgtgggtag gggcgaaaaa ccaatcgaac tcggagttag 240

ctggttctcc ccgaaatgcg tttaggcgca gcagtagcaa cacaaataga ggggtaaagc 300

actgtttctt ttgtgggctt cgaaagttgt acctcaaagt ggcaaactct gaatactcta 360

tttagatatc tactagtgag accttggggg ataagctcct tggtcaaaag ggaaacagcc 420

cagatcacca gttaaggccc caaaatgaaa atgatagtga ctaaggatgt gggtatgtta 480

aaacctccag caggttagct tagaagcagc aatcctttca agagtgcgta atagctcact 540

g 541

<210> 15
<211> 573
<212> DNA
<213> Prototheca moriformis

<400> 15
tgttgaagaa tgagccggcg acttaaaata aatggcaggc taagagaatt aataactcga 60

aacctaagcg aaagcaagtc ttaatagggc gctaatttaa caaaacatta aataaaatct 120

aaagtcattt attttagacc cgaacctgag tgatctaacc atggtcagga tgaaacttgg 180

gtgacaccaa gtggaagtcc gaaccgaccg atgttgaaaa atcggcggat gaactgtggt 240

tagtggtgaa ataccagtcg aactcagagc tagctggttc tccccgaaat gcgttgaggc 300

gcagcaatat atctcgtcta tctaggggta aagcactgtt tcggtgcggg ctatgaaaat 360

ggtaccaaat cgtggcaaac tctgaatact agaaatgacg atatattagt gagactatgg 420

gggataagct ccatagtcga gagggaaaca gcccagacca ccagttaagg ccccaaaatg 480

ataatgaagt ggtaaaggag gtgaaaatgc aaatacaacc aggaggttgg cttagaagca 540

gccatccttt aaagagtgcg taatagctca ctg 573

<210> 16
<211> 573
<212> DNA
<213> Prototheca wickerhamii

<400> 16
tgttgaagaa tgagccgtcg acttaaaata aatggcaggc taagagaatt aataactcga 60

aacctaagcg aaagcaagtc ttaatagggc gctaatttaa caaaacatta aataaaatct 120

aaagtcattt attttagacc cgaacctgag tgatctaacc atggtcagga tgaaacttgg 180

gtgacaccaa gtggaagtcc gaaccgaccg atgttgaaaa atcggcggat gaactgtggt 240

tagtggtgaa ataccagtcg aactcagagc tagctggttc tccccgaaat gcgttgaggc 300

gcagcaatat atctcgtcta tctaggggta aagcactgtt tcggtgcggg ctatgaaaat 360

ggtaccaaat cgtggcaaac tctgaatact agaaatgacg atatattagt gagactatgg 420

gggataagct ccatagtcga gagggaaaca gcccagacca ccagttaagg ccccaaaatg 480

ataatgaagt ggtaaaggag gtgaaaatgc aaatacaacc aggaggttgg cttagaagca 540

gccatccttt aaagagtgcg taatagctca ctg 573

<210> 17
<211> 541
<212> DNA
<213> Prototheca moriformis

<400> 17
tgttgaagaa tgagccggcg agttaaaaag agtggcgtgg ttaaagaaaa ttctctggaa 60

ccatagcgaa agcaagttta acaagcttaa gtcacttttt ttagacccga aaccgagtga 120

tctacccatg atcagggtga agtgttggta aaataacatg gaggcccgaa ccgactaatg 180

gtgaaaaatt agcggatgaa ttgtgggtag gggcgaaaaa ccaatcgaac tcggagttag 240

ctggttctcc ccgaaatgcg tttaggcgca gcagtagcaa cacaaataga ggggtaaagc 300

actgtttctt ttgtgggctc cgaaagttgt acctcaaagt ggcaaactct gaatactcta 360

tttagatatc tactagtgag accttggggg ataagctcct tggtcgaaag ggaaacagcc 420

cagatcacca gttaaggccc caaaatgaaa atgatagtga ctaaggatgt gagtatgtca 480

aaacctccag caggttagct tagaagcagc aatcctttca agagtgcgta atagctcact 540

g 541

<210> 18
<211> 541
<212> DNA
<213> Prototheca zopfii

<400> 18
tgttgaagaa tgagccggcg agttaaaaag agtggcatgg ttaaagaaaa ttctctggag 60

ccatagcgaa agcaagttta acaagcttaa gtcacttttt ttagacccga aaccgagtga 120

tctacccatg atcagggtga agtgttggta aaataacatg gaggcccgaa ccgactaatg 180

gtgaaaaatt agcggatgaa ttgtgggtag gggcgaaaaa ccaatcgaac tcggagttag 240

ctggttctcc ccgaaatgcg tttaggcgca gcagtagcaa cacaaataga ggggtaaagc 300

actgtttctt tcgtgggctt cgaaagttgt acctcaaagt ggcaaactct gaatactcta 360

tttagatatc tactagtgag accttggggg ataagctcct tggtcaaaag ggaaacagcc 420

cagatcacca gttaaggccc caaaatgaaa atgatagtga ctaaggatgt gagtatgtca 480

aaacctccag caggttagct tagaagcagc aatcctttca agagtgcgta atagctcact 540

g 541

<210> 19
<211> 565
<212> DNA
<213> Prototheca moriformis

<400> 19
tgttgaagaa tgagccggcg acttagaaaa ggtggcatgg ttaaggaaat attccgaagc 60

cgtagcaaaa gcgagtctga atagggcgat aaaatatatt aatatttaga atctagtcat 120

```
tttttctaga cccgaacccg ggtgatctaa ccatgaccag gatgaagctt gggtgatacc      180

aagtgaaggt ccgaaccgac cgatgttgaa aaatcggcgg atgagttgtg gttagcggtg      240

aaataccagt cgaacccgga gctagctggt tctccccgaa atgcgttgag gcgcagcagt      300

acatctagtc tatctagggg taaagcactg tttcggtgcg ggctgtgaga acggtaccaa      360

atcgtggcaa actctgaata ctagaaatga cgatgtagta gtgagactgt gggggataag      420

ctccattgtc aagagggaaa cagcccagac caccagctaa ggccccaaaa tggtaatgta      480

gtgacaaagg aggtgaaaat gcaaatacaa ccaggaggtt ggcttagaag cagccatcct      540

ttaaagagtg cgtaatagct cactg                                            565
```

<210> 20
<211> 550
<212> PRT
<213> Cichorium intybus

<400> 20

```
Met Ser Asn Ser Ser Asn Ala Ser Glu Ser Leu Phe Pro Ala Thr Ser
1               5                  10                  15

Glu Gln Pro Tyr Arg Thr Ala Phe His Phe Gln Pro Pro Gln Asn Trp
            20                  25                  30

Met Asn Asp Pro Asn Gly Pro Met Cys Tyr Asn Gly Val Tyr His Leu
        35                  40                  45

Phe Tyr Gln Tyr Asn Pro Phe Gly Pro Leu Trp Asn Leu Arg Met Tyr
    50                  55                  60

Trp Ala His Ser Val Ser His Asp Leu Ile Asn Trp Ile His Leu Asp
65                  70                  75                  80

Leu Ala Phe Ala Pro Thr Glu Pro Phe Asp Ile Asn Gly Cys Leu Ser
                85                  90                  95

Gly Ser Ala Thr Val Leu Pro Gly Asn Lys Pro Ile Met Leu Tyr Thr
            100                 105                 110

Gly Ile Asp Thr Glu Asn Arg Gln Val Gln Asn Leu Ala Val Pro Lys
        115                 120                 125

Asp Leu Ser Asp Pro Tyr Leu Arg Glu Trp Val Lys His Thr Gly Asn
    130                 135                 140

Pro Ile Ile Ser Leu Pro Glu Glu Ile Gln Pro Asp Asp Phe Arg Asp
145                 150                 155                 160

Pro Thr Thr Thr Trp Leu Glu Glu Asp Gly Thr Trp Arg Leu Leu Val
                165                 170                 175

Gly Ser Gln Lys Asp Lys Thr Gly Ile Ala Phe Leu Tyr His Ser Gly
```

180 185 190

Asp Phe Val Asn Trp Thr Lys Ser Asp Ser Pro Leu His Lys Val Ser
195 200 205

Gly Thr Gly Met Trp Glu Cys Val Asp Phe Phe Pro Val Trp Val Asp
210 215 220

Ser Thr Asn Gly Val Asp Thr Ser Ile Ile Asn Pro Ser Asn Arg Val
225 230 235 240

Lys His Val Leu Lys Leu Gly Ile Gln Asp His Gly Lys Asp Cys Tyr
245 250 255

Leu Ile Gly Lys Tyr Ser Ala Asp Lys Glu Asn Tyr Val Pro Glu Asp
260 265 270

Glu Leu Thr Leu Ser Thr Leu Arg Leu Asp Tyr Gly Met Tyr Tyr Ala
275 280 285

Ser Lys Ser Phe Phe Asp Pro Val Lys Asn Arg Arg Ile Met Thr Ala
290 295 300

Trp Val Asn Glu Ser Asp Ser Glu Ala Asp Val Ile Ala Arg Gly Trp
305 310 315 320

Ser Gly Val Gln Ser Phe Pro Arg Ser Leu Trp Leu Asp Lys Asn Gln
325 330 335

Lys Gln Leu Leu Gln Trp Pro Ile Glu Glu Ile Glu Met Leu His Gln
340 345 350

Asn Glu Val Ser Phe His Asn Lys Lys Leu Asp Gly Gly Ser Ser Leu
355 360 365

Glu Val Leu Gly Ile Thr Ala Ser Gln Ala Asp Val Lys Ile Ser Phe
370 375 380

Lys Leu Ala Asn Leu Glu Glu Ala Glu Glu Leu Asp Pro Ser Trp Val
385 390 395 400

Asp Pro Gln Leu Ile Cys Ser Glu Asn Asp Ala Ser Lys Lys Gly Lys
405 410 415

Phe Gly Pro Phe Gly Leu Leu Ala Leu Ala Ser Ser Asp Leu Arg Glu
420 425 430

Gln Thr Ala Ile Phe Phe Arg Val Phe Arg Lys Asn Gly Arg Tyr Val
435 440 445

Val Leu Met Cys Ser Asp Gln Ser Arg Ser Ser Met Lys Asn Gly Ile

450 455 460

Glu Lys Arg Thr Tyr Gly Ala Phe Val Asp Ile Asp Pro Gln Gln Asp
465 470 475 480

Glu Ile Ser Leu Arg Thr Leu Ile Asp His Ser Ile Val Glu Ser Phe
485 490 495

Gly Gly Arg Gly Lys Thr Cys Ile Thr Thr Arg Val Tyr Pro Thr Leu
500 505 510

Ala Ile Gly Glu Gln Ala Arg Leu Phe Ala Phe Asn His Gly Thr Glu
515 520 525

Ser Val Glu Ile Ser Glu Leu Ser Ala Trp Ser Met Lys Lys Ala Gln
530 535 540

Met Lys Val Glu Glu Pro
545 550

<210> 21
<211> 581
<212> PRT
<213> Schizosaccharomyces pombe

<400> 21
Met Phe Leu Lys Tyr Ile Leu Ala Ser Gly Ile Cys Leu Val Ser Leu
1 5 10 15

Leu Ser Ser Thr Asn Ala Ala Pro Arg His Leu Tyr Val Lys Arg Tyr
20 25 30

Pro Val Ile Tyr Asn Ala Ser Asn Ile Thr Glu Val Ser Asn Ser Thr
35 40 45

Thr Val Pro Pro Pro Pro Phe Val Asn Thr Thr Ala Pro Asn Gly Thr
50 55 60

Cys Leu Gly Asn Tyr Asn Glu Tyr Leu Pro Ser Gly Tyr Tyr Asn Ala
65 70 75 80

Thr Asp Arg Pro Lys Ile His Phe Thr Pro Ser Ser Gly Phe Met Asn
85 90 95

Asp Pro Asn Gly Leu Val Tyr Thr Gly Gly Val Tyr His Met Phe Phe
100 105 110

Gln Tyr Ser Pro Lys Thr Leu Thr Ala Gly Glu Val His Trp Gly His
115 120 125

Thr Val Ser Lys Asp Leu Ile His Trp Glu Asn Tyr Pro Ile Ala Ile
130 135 140

```
Tyr Pro Asp Glu His Glu Asn Gly Val Leu Ser Leu Pro Phe Ser Gly
145                 150                 155                 160

Ser Ala Val Val Asp Val His Asn Ser Ser Gly Leu Phe Ser Asn Asp
                165                 170                 175

Thr Ile Pro Glu Glu Arg Ile Val Leu Ile Tyr Thr Asp His Trp Thr
            180                 185                 190

Gly Val Ala Glu Arg Gln Ala Ile Ala Tyr Thr Thr Asp Gly Gly Tyr
        195                 200                 205

Thr Phe Lys Lys Tyr Ser Gly Asn Pro Val Leu Asp Ile Asn Ser Leu
    210                 215                 220

Gln Phe Arg Asp Pro Lys Val Ile Trp Asp Phe Asp Ala Asn Arg Trp
225                 230                 235                 240

Val Met Ile Val Ala Met Ser Gln Asn Tyr Gly Ile Ala Phe Tyr Ser
                245                 250                 255

Ser Tyr Asp Leu Ile His Trp Thr Glu Leu Ser Val Phe Ser Thr Ser
            260                 265                 270

Gly Tyr Leu Gly Leu Gln Tyr Glu Cys Pro Gly Met Ala Arg Val Pro
        275                 280                 285

Val Glu Gly Thr Asp Glu Tyr Lys Trp Val Leu Phe Ile Ser Ile Asn
    290                 295                 300

Pro Gly Ala Pro Leu Gly Gly Ser Val Val Gln Tyr Phe Val Gly Asp
305                 310                 315                 320

Trp Asn Gly Thr Asn Phe Val Pro Asp Asp Gly Gln Thr Arg Phe Val
                325                 330                 335

Asp Leu Gly Lys Asp Phe Tyr Ala Ser Ala Leu Tyr His Ser Ser Ser
            340                 345                 350

Ala Asn Ala Asp Val Ile Gly Val Gly Trp Ala Ser Asn Trp Gln Tyr
        355                 360                 365

Thr Asn Gln Ala Pro Thr Gln Val Phe Arg Ser Ala Met Thr Val Ala
    370                 375                 380

Arg Lys Phe Thr Leu Arg Asp Val Pro Gln Asn Pro Met Thr Asn Leu
385                 390                 395                 400

Thr Ser Leu Ile Gln Thr Pro Leu Asn Val Ser Leu Leu Arg Asp Glu
                405                 410                 415
```

```
Thr Leu Phe Thr Ala Pro Val Ile Asn Ser Ser Ser Ser Leu Ser Gly
            420                 425                 430

Ser Pro Ile Thr Leu Pro Ser Asn Thr Ala Phe Glu Phe Asn Val Thr
        435                 440                 445

Leu Ser Ile Asn Tyr Thr Glu Gly Cys Thr Thr Gly Tyr Cys Leu Gly
    450                 455                 460

Arg Ile Ile Ile Asp Ser Asp Asp Pro Tyr Arg Leu Gln Ser Ile Ser
465                 470                 475                 480

Val Asp Val Asp Phe Ala Ala Ser Thr Leu Val Ile Asn Arg Ala Lys
                485                 490                 495

Ala Gln Met Gly Trp Phe Asn Ser Leu Phe Thr Pro Ser Phe Ala Asn
            500                 505                 510

Asp Ile Tyr Ile Tyr Gly Asn Val Thr Leu Tyr Gly Ile Val Asp Asn
        515                 520                 525

Gly Leu Leu Glu Leu Tyr Val Asn Asn Gly Glu Lys Thr Tyr Thr Asn
    530                 535                 540

Asp Phe Phe Phe Leu Gln Gly Ala Thr Pro Gly Gln Ile Ser Phe Ala
545                 550                 555                 560

Ala Phe Gln Gly Val Ser Phe Asn Asn Val Thr Val Thr Pro Leu Lys
                565                 570                 575

Thr Ile Trp Asn Cys
            580
```

<210> 22
<211> 550
<212> PRT
<213> Pichia anomala

<400> 22

```
Met Ile Gln Leu Ser Pro Leu Leu Leu Leu Pro Leu Phe Ser Val Phe
1               5                   10                  15

Asn Ser Ile Ala Asp Ala Ser Thr Glu Tyr Leu Arg Pro Gln Ile His
            20                  25                  30

Leu Thr Pro Asp Gln Gly Trp Met Asn Asp Pro Asn Gly Met Phe Tyr
        35                  40                  45

Asp Arg Lys Asp Lys Leu Trp His Val Tyr Phe Gln His Asn Pro Asp
    50                  55                  60

Lys Lys Ser Ile Trp Ala Thr Pro Val Thr Trp Gly His Ser Thr Ser
```

65 70 75 80

Lys Asp Leu Leu Thr Trp Asp Tyr His Gly Asn Ala Leu Glu Pro Glu
85 90 95

Asn Asp Asp Glu Gly Ile Phe Ser Gly Ser Val Val Val Asp Arg Asn
100 105 110

Asn Thr Ser Gly Phe Phe Asn Asp Ser Thr Asp Pro Glu Gln Arg Ile
115 120 125

Val Ala Ile Tyr Thr Asn Asn Ala Gln Leu Gln Thr Gln Glu Ile Ala
130 135 140

Tyr Ser Leu Asp Lys Gly Tyr Ser Phe Ile Lys Tyr Asp Gln Asn Pro
145 150 155 160

Val Ile Asn Val Asn Ser Ser Gln Gln Arg Asp Pro Lys Val Leu Trp
165 170 175

His Asp Glu Ser Asn Gln Trp Ile Met Val Val Ala Lys Thr Gln Glu
180 185 190

Phe Lys Val Gln Ile Tyr Gly Ser Pro Asp Leu Lys Lys Trp Asp Leu
195 200 205

Lys Ser Asn Phe Thr Ser Asn Gly Tyr Leu Gly Phe Gln Tyr Glu Cys
210 215 220

Pro Gly Leu Phe Lys Leu Pro Ile Glu Asn Pro Leu Asn Asp Thr Val
225 230 235 240

Thr Ser Lys Trp Val Leu Leu Leu Ala Ile Asn Pro Gly Ser Pro Leu
245 250 255

Gly Gly Ser Ile Asn Glu Tyr Phe Ile Gly Asp Phe Asp Gly Thr Thr
260 265 270

Phe His Pro Asp Asp Gly Ala Thr Arg Phe Met Asp Ile Gly Lys Asp
275 280 285

Phe Tyr Ala Phe Gln Ser Phe Asp Asn Thr Glu Pro Glu Asp Gly Ala
290 295 300

Leu Gly Leu Ala Trp Ala Ser Asn Trp Gln Tyr Ala Asn Thr Val Pro
305 310 315 320

Thr Glu Asn Trp Arg Ser Ser Met Ser Leu Val Arg Asn Tyr Thr Leu
325 330 335

Lys Tyr Val Asp Val Asn Pro Glu Asn Tyr Gly Leu Thr Leu Ile Gln

340 345 350

Lys Pro Val Tyr Asp Thr Lys Glu Thr Arg Leu Asn Glu Thr Leu Lys
355 360 365

Thr Leu Glu Thr Ile Asn Glu Tyr Glu Val Asn Asp Leu Lys Leu Asp
370 375 380

Lys Ser Ser Phe Val Ala Thr Asp Phe Asn Thr Glu Arg Asn Ala Thr
385 390 395 400

Gly Val Phe Glu Phe Asp Leu Lys Phe Thr Gln Thr Asp Leu Lys Met
405 410 415

Gly Tyr Ser Asn Met Thr Thr Gln Phe Gly Leu Tyr Ile His Ser Gln
420 425 430

Thr Val Lys Gly Ser Gln Glu Thr Leu Gln Leu Val Phe Asp Thr Leu
435 440 445

Ser Thr Thr Trp Tyr Ile Asp Arg Thr Thr Gln His Ser Phe Gln Arg
450 455 460

Asn Ser Pro Val Phe Thr Glu Arg Ile Ser Thr Tyr Val Glu Lys Ile
465 470 475 480

Asp Thr Thr Asp Gln Gly Asn Val Tyr Thr Leu Tyr Gly Val Val Asp
485 490 495

Arg Asn Ile Leu Glu Leu Tyr Phe Asn Asp Gly Ser Ile Ala Met Thr
500 505 510

Asn Thr Phe Phe Phe Arg Glu Gly Lys Ile Pro Thr Ser Phe Glu Val
515 520 525

Val Cys Asp Ser Glu Lys Ser Phe Ile Thr Ile Asp Glu Leu Ser Val
530 535 540

Arg Glu Leu Ala Arg Lys
545 550

<210> 23
<211> 533
<212> PRT
<213> Debaryomyces occidentalis

<400> 23
Met Val Gln Val Leu Ser Val Leu Val Ile Pro Leu Leu Thr Leu Phe
1 5 10 15

Phe Gly Tyr Val Ala Ser Ser Ser Ile Asp Leu Ser Val Asp Thr Ser
20 25 30

```
Glu Tyr Asn Arg Pro Leu Ile His Phe Thr Pro Glu Lys Gly Trp Met
        35                  40                  45

Asn Asp Pro Asn Gly Leu Phe Tyr Asp Lys Thr Ala Lys Leu Trp His
    50                  55                  60

Leu Tyr Phe Gln Tyr Asn Pro Asn Ala Thr Ala Trp Gly Gln Pro Leu
65                  70                  75                  80

Tyr Trp Gly His Ala Thr Ser Asn Asp Leu Val His Trp Asp Glu His
                85                  90                  95

Glu Ile Ala Ile Gly Pro Glu His Asp Asn Glu Gly Ile Phe Ser Gly
            100                 105                 110

Ser Ile Val Val Asp His Asn Asn Thr Ser Gly Phe Phe Asn Ser Ser
        115                 120                 125

Ile Asp Pro Asn Gln Arg Ile Val Ala Ile Tyr Thr Asn Asn Ile Pro
    130                 135                 140

Asp Leu Gln Thr Gln Asp Ile Ala Phe Ser Leu Asp Gly Gly Tyr Thr
145                 150                 155                 160

Phe Thr Lys Tyr Glu Asn Asn Pro Val Ile Asp Val Ser Ser Asn Gln
                165                 170                 175

Phe Arg Asp Pro Lys Val Phe Trp His Glu Arg Phe Lys Ser Met Asp
            180                 185                 190

His Gly Cys Ser Glu Ile Ala Arg Val Lys Ile Gln Ile Phe Gly Ser
        195                 200                 205

Ala Asn Leu Lys Asn Trp Val Leu Asn Ser Asn Phe Ser Ser Gly Tyr
    210                 215                 220

Tyr Gly Asn Gln Tyr Gly Met Ser Arg Leu Ile Glu Val Pro Ile Glu
225                 230                 235                 240

Asn Ser Asp Lys Ser Lys Trp Val Met Phe Leu Ala Ile Asn Pro Gly
                245                 250                 255

Ser Pro Leu Gly Gly Ser Ile Asn Gln Tyr Phe Val Gly Asp Phe Asp
            260                 265                 270

Gly Phe Gln Phe Val Pro Asp Asp Ser Gln Thr Arg Phe Val Asp Ile
        275                 280                 285

Gly Lys Asp Phe Tyr Ala Phe Gln Thr Phe Ser Glu Val Glu His Gly
    290                 295                 300
```

```
Val Leu Gly Leu Ala Trp Ala Ser Asn Trp Gln Tyr Ala Asp Gln Val
305                 310                 315                 320

Pro Thr Asn Pro Trp Arg Ser Ser Thr Ser Leu Ala Arg Asn Tyr Thr
                325                 330                 335

Leu Arg Tyr Val Ile Gln Met Leu Lys Leu Thr Ala Asn Ile Asp Lys
            340                 345                 350

Ser Val Leu Pro Asp Ser Ile Asn Val Val Asp Lys Leu Lys Lys Lys
        355                 360                 365

Asn Val Lys Leu Thr Asn Lys Lys Pro Ile Lys Thr Asn Phe Lys Gly
    370                 375                 380

Ser Thr Gly Leu Phe Asp Phe Asn Ile Thr Phe Lys Val Leu Asn Leu
385                 390                 395                 400

Asn Val Ser Pro Gly Lys Thr His Phe Asp Ile Leu Ile Asn Ser Gln
                405                 410                 415

Glu Leu Asn Ser Ser Val Asp Ser Ile Lys Ile Gly Phe Asp Ser Ser
            420                 425                 430

Gln Ser Leu Phe Tyr Ile Asp Arg His Ile Pro Asn Val Glu Phe Pro
        435                 440                 445

Arg Lys Gln Phe Phe Thr Asp Lys Leu Ala Ala Tyr Leu Glu Pro Leu
    450                 455                 460

Asp Tyr Asp Gln Asp Leu Arg Val Phe Ser Leu Tyr Gly Ile Val Asp
465                 470                 475                 480

Lys Asn Ile Ile Glu Leu Tyr Phe Asn Asp Gly Thr Val Ala Met Thr
                485                 490                 495

Asn Thr Phe Phe Met Gly Glu Gly Lys Tyr Pro His Asp Ile Gln Ile
            500                 505                 510

Val Thr Asp Thr Glu Glu Pro Leu Phe Glu Leu Glu Ser Val Ile Ile
        515                 520                 525

Arg Glu Leu Asn Lys
    530
```

<210> 24
<211> 654
<212> PRT
<213> Oryza sativa

<400> 24

```
Met Ala Thr Ser Arg Leu Thr Pro Ala Tyr Asp Leu Lys Asn Ala Ala
```

1 5 10 15

Ala Ala Val Tyr Thr Pro Leu Pro Glu Gln Pro His Ser Ala Glu Val
20 25 30

Glu Ile Arg Asp Arg Lys Pro Phe Lys Ile Ile Ser Ala Ile Ile Leu
35 40 45

Ser Ser Leu Leu Leu Leu Ala Leu Ile Leu Val Ala Val Asn Tyr Gln
50 55 60

Ala Pro Pro Ser His Ser Ser Gly Asp Asn Ser Gln Pro Ala Ala Val
65 70 75 80

Met Pro Pro Ser Arg Gly Val Ser Gln Gly Val Ser Glu Lys Ala Phe
85 90 95

Arg Gly Ala Ser Gly Ala Gly Asn Gly Val Ser Phe Ala Trp Ser Asn
100 105 110

Leu Met Leu Ser Trp Gln Arg Thr Ser Tyr His Phe Gln Pro Val Lys
115 120 125

Asn Trp Met Asn Asp Pro Asn Gly Pro Leu Tyr Tyr Lys Gly Trp Tyr
130 135 140

His Leu Phe Tyr Gln Tyr Asn Pro Asp Ser Ala Val Trp Gly Asn Ile
145 150 155 160

Thr Trp Gly His Ala Val Ser Thr Asp Leu Ile Asn Trp Leu His Leu
165 170 175

Pro Phe Ala Met Val Pro Asp Gln Trp Tyr Asp Val Asn Gly Val Trp
180 185 190

Thr Gly Ser Ala Thr Ile Leu Pro Asp Gly Arg Ile Val Met Leu Tyr
195 200 205

Thr Gly Asp Thr Asp Asp Tyr Val Gln Asp Gln Asn Leu Ala Phe Pro
210 215 220

Ala Asn Leu Ser Asp Pro Leu Leu Val Asp Trp Val Lys Tyr Pro Asn
225 230 235 240

Asn Pro Val Ile Tyr Pro Pro Pro Gly Ile Gly Val Lys Asp Phe Arg
245 250 255

Asp Pro Thr Thr Ala Gly Thr Ala Gly Met Gln Asn Gly Gln Arg Leu
260 265 270

Val Thr Ile Gly Ser Lys Val Gly Lys Thr Gly Ile Ser Leu Val Tyr

275 280 285

Glu Thr Thr Asn Phe Thr Thr Phe Lys Leu Leu Tyr Gly Val Leu His
290 295 300

Ala Val Pro Gly Thr Gly Met Trp Glu Cys Val Asp Leu Tyr Pro Val
305 310 315 320

Ser Thr Thr Gly Glu Asn Gly Leu Asp Thr Ser Val Asn Gly Leu Gly
325 330 335

Val Lys His Val Leu Lys Thr Ser Leu Asp Asp Asp Lys His Asp Tyr
340 345 350

Tyr Ala Leu Gly Thr Tyr Asp Pro Val Lys Asn Lys Trp Thr Pro Asp
355 360 365

Asn Pro Asp Leu Asp Val Gly Ile Gly Leu Arg Leu Asp Tyr Gly Lys
370 375 380

Tyr Tyr Ala Ala Arg Thr Phe Tyr Asp Gln Asn Lys Gln Arg Arg Ile
385 390 395 400

Leu Trp Gly Trp Ile Gly Glu Thr Asp Leu Glu Ala Val Asp Leu Met
405 410 415

Lys Gly Trp Ala Ser Leu Gln Ala Ile Pro Arg Thr Ile Val Phe Asp
420 425 430

Lys Lys Thr Gly Thr Asn Val Leu Gln Arg Pro Glu Glu Glu Val Glu
435 440 445

Ser Trp Ser Ser Gly Asp Pro Ile Thr Gln Arg Arg Ile Phe Glu Pro
450 455 460

Gly Ser Val Val Pro Ile His Val Ser Gly Ala Thr Gln Leu Asp Ile
465 470 475 480

Thr Ala Ser Phe Glu Val Asp Glu Thr Leu Leu Glu Thr Thr Ser Glu
485 490 495

Ser His Asp Ala Gly Tyr Asp Cys Ser Asn Ser Gly Gly Ala Gly Thr
500 505 510

Arg Gly Ser Leu Gly Pro Phe Gly Leu Leu Val Val Ala Asp Glu Lys
515 520 525

Leu Ser Glu Leu Thr Pro Val Tyr Leu Tyr Val Ala Lys Gly Gly Asp
530 535 540

Gly Lys Ala Lys Ala His Leu Cys Ala Tyr Gln Thr Arg Ser Ser Met

545 550 555 560

Ala Ser Gly Val Glu Lys Glu Val Tyr Gly Ser Ala Val Pro Val Leu
565 570 575

Asp Gly Glu Asn Tyr Ser Ala Arg Ile Leu Ile Asp His Ser Ile Val
580 585 590

Glu Ser Phe Ala Gln Ala Gly Arg Thr Cys Val Arg Ser Arg Asp Tyr
595 600 605

Pro Thr Lys Asp Ile Tyr Gly Ala Ala Arg Cys Phe Phe Phe Asn Asn
610 615 620

Ala Thr Glu Ala Ser Val Arg Ala Ser Leu Lys Ala Trp Gln Met Lys
625 630 635 640

Ser Phe Ile Arg Pro Tyr Pro Phe Ile Pro Asp Gln Lys Ser
645 650

<210> 25
<211> 690
<212> PRT
<213> Allium cepa

<400> 25
Met Ser Ser Asp Asp Leu Glu Ser Pro Pro Ser Ser Tyr Leu Pro Ile
1 5 10 15

Pro Pro Ser Asp Glu Phe His Asp Gln Pro Pro Pro Leu Arg Ser Trp
20 25 30

Leu Arg Leu Leu Ser Ile Pro Leu Ala Leu Met Phe Leu Leu Phe Leu
35 40 45

Ala Thr Phe Leu Ser Asn Leu Glu Ser Pro Pro Ser Asp Ser Gly Leu
50 55 60

Val Ser Asp Pro Val Thr Phe Asp Val Asn Pro Ala Val Val Arg Arg
65 70 75 80

Gly Lys Asp Ala Gly Val Ser Asp Lys Thr Ser Gly Val Asp Ser Gly
85 90 95

Phe Val Leu Asp Pro Val Ala Val Asp Ala Asn Ser Val Val Val His
100 105 110

Arg Gly Lys Asp Ala Gly Val Ser Asp Lys Thr Ser Gly Val Asp Ser
115 120 125

Gly Leu Leu Lys Asp Ser Pro Leu Gly Pro Tyr Pro Trp Thr Asn Gln
130 135 140

```
Met Leu Ser Trp Gln Arg Thr Gly Phe His Phe Gln Pro Val Lys Asn
145                 150                 155                 160

Trp Met Asn Asp Pro Asn Gly Pro Leu Tyr Tyr Lys Gly Trp Tyr His
                165                 170                 175

Phe Phe Tyr Gln Tyr Asn Pro Glu Gly Ala Val Trp Gly Asn Ile Ala
            180                 185                 190

Trp Gly His Ala Val Ser Arg Asp Leu Val His Trp Thr His Leu Pro
        195                 200                 205

Leu Ala Met Val Pro Asp Gln Trp Tyr Asp Ile Asn Gly Val Trp Thr
    210                 215                 220

Gly Ser Ala Thr Ile Leu Pro Asp Gly Gln Ile Val Met Leu Tyr Thr
225                 230                 235                 240

Gly Ala Thr Asn Glu Ser Val Gln Val Gln Asn Leu Ala Val Pro Ala
                245                 250                 255

Asp Gln Ser Asp Thr Leu Leu Leu Arg Trp Lys Lys Ser Glu Ala Asn
            260                 265                 270

Pro Ile Leu Val Pro Pro Pro Gly Ile Gly Asp Lys Asp Phe Arg Asp
        275                 280                 285

Pro Thr Thr Ala Trp Tyr Glu Pro Ser Asp Asp Thr Trp Arg Ile Val
    290                 295                 300

Ile Gly Ser Lys Asp Ser Ser His Ser Gly Ile Ala Ile Val Tyr Ser
305                 310                 315                 320

Thr Lys Asp Phe Ile Asn Tyr Lys Leu Ile Pro Gly Ile Leu His Ala
                325                 330                 335

Val Glu Arg Val Gly Met Trp Glu Cys Val Asp Phe Tyr Pro Val Ala
            340                 345                 350

Thr Ala Asp Ser Ser His Ala Asn His Gly Leu Asp Pro Ser Ala Arg
        355                 360                 365

Pro Ser Pro Ala Val Lys His Val Leu Lys Ala Ser Met Asp Asp Asp
    370                 375                 380

Arg His Asp Tyr Tyr Ala Ile Gly Thr Tyr Asp Pro Ala Gln Asn Thr
385                 390                 395                 400

Trp Val Pro Asp Asp Ala Ser Val Asp Val Gly Ile Gly Leu Arg Tyr
                405                 410                 415
```

Asp Trp Gly Lys Phe Tyr Ala Ser Lys Thr Phe Tyr Asp His Ala Lys
420 425 430

Lys Arg Arg Ile Leu Trp Ser Trp Ile Gly Glu Thr Asp Ser Glu Thr
435 440 445

Ala Asp Ile Ala Lys Gly Trp Ala Ser Leu Gln Gly Val Pro Arg Thr
450 455 460

Val Leu Leu Asp Val Lys Thr Gly Ser Asn Leu Ile Thr Trp Pro Val
465 470 475 480

Val Glu Ile Glu Ser Leu Arg Thr Arg Pro Arg Asp Phe Ser Gly Ile
485 490 495

Thr Val Asp Ala Gly Ser Thr Phe Lys Leu Asp Val Gly Gly Ala Ala
500 505 510

Gln Leu Asp Ile Glu Ala Glu Phe Lys Ile Ser Ser Glu Glu Leu Glu
515 520 525

Ala Val Lys Glu Ala Asp Val Ser Tyr Asn Cys Ser Ser Ser Gly Gly
530 535 540

Ala Ala Glu Arg Gly Val Leu Gly Pro Phe Gly Leu Leu Val Leu Ala
545 550 555 560

Asn Gln Asp Leu Thr Glu Gln Thr Ala Thr Tyr Phe Tyr Val Ser Arg
565 570 575

Gly Met Asp Gly Gly Leu Asn Thr His Phe Cys Gln Asp Glu Lys Arg
580 585 590

Ser Ser Lys Ala Ser Asp Ile Val Lys Arg Ile Val Gly His Ser Val
595 600 605

Pro Val Leu Asp Gly Glu Ser Phe Ala Leu Arg Ile Leu Val Asp His
610 615 620

Ser Ile Val Glu Ser Phe Ala Gln Gly Gly Arg Ala Ser Ala Thr Ser
625 630 635 640

Arg Val Tyr Pro Thr Glu Ala Ile Tyr Asn Asn Ala Arg Val Phe Val
645 650 655

Phe Asn Asn Ala Thr Gly Ala Lys Val Thr Ala Gln Ser Leu Lys Val
660 665 670

Trp His Met Ser Thr Ala Ile Asn Glu Ile Tyr Asp Pro Ala Thr Ser
675 680 685

Val Met
690

<210> 26
<211> 501
<212> PRT
<213> Beta vulgaris

<220>
<221> MOD_RES
<222> (382)..(382)
<223> Any amino acid

<400> 26
Leu Phe Tyr Gln Tyr Asn Pro Asn Gly Val Ile Trp Gly Pro Pro Val
1 5 10 15

Trp Gly His Ser Thr Ser Lys Asp Leu Val Asn Trp Val Pro Gln Pro
20 25 30

Leu Thr Met Glu Pro Glu Met Ala Ala Asn Ile Asn Gly Ser Trp Ser
35 40 45

Gly Ser Ala Thr Ile Leu Pro Gly Asn Lys Pro Ala Ile Leu Phe Thr
50 55 60

Gly Leu Asp Pro Lys Tyr Glu Gln Val Gln Val Leu Ala Tyr Pro Lys
65 70 75 80

Asp Thr Ser Asp Pro Asn Leu Lys Glu Trp Phe Leu Ala Pro Gln Asn
85 90 95

Pro Val Met Phe Pro Thr Pro Gln Asn Gln Ile Asn Ala Thr Ser Phe
100 105 110

Arg Asp Pro Thr Thr Ala Trp Arg Leu Pro Asp Gly Val Trp Arg Leu
115 120 125

Leu Ile Gly Ser Lys Arg Gly Gln Arg Gly Leu Ser Leu Leu Phe Arg
130 135 140

Ser Arg Asp Phe Val His Trp Val Gln Ala Lys His Pro Leu Tyr Ser
145 150 155 160

Asp Lys Leu Ser Gly Met Trp Glu Cys Pro Asp Phe Phe Pro Val Tyr
165 170 175

Ala Asn Gly Asp Gln Met Gly Val Asp Thr Ser Ile Ile Gly Ser His
180 185 190

Val Lys His Val Leu Lys Asn Ser Leu Asp Ile Thr Lys His Asp Ile
195 200 205

```
Tyr Thr Ile Gly Asp Tyr Asn Ile Lys Lys Asp Ala Tyr Thr Pro Asp
    210                 215                 220

Ile Gly Tyr Met Asn Asp Ser Ser Leu Arg Tyr Asp Tyr Gly Lys Tyr
225                 230                 235                 240

Tyr Ala Ser Lys Thr Phe Phe Asp Asp Ala Lys Lys Glu Arg Ile Leu
                245                 250                 255

Leu Gly Trp Ala Asn Glu Ser Ser Ser Val Glu Asp Asp Ile Lys Lys
            260                 265                 270

Gly Trp Ser Gly Ile His Thr Ile Pro Arg Lys Ile Trp Leu Asp Lys
        275                 280                 285

Leu Gly Lys Gln Leu Ile Gln Trp Pro Ile Ala Asn Ile Glu Lys Leu
    290                 295                 300

Arg Gln Lys Pro Val Asn Ile Tyr Arg Lys Val Leu Lys Gly Gly Ser
305                 310                 315                 320

Gln Ile Glu Val Ser Gly Ile Thr Ala Ala Gln Ala Asp Val Glu Ile
                325                 330                 335

Ser Phe Lys Ile Lys Asp Leu Lys Asn Val Glu Lys Phe Asp Ala Ser
            340                 345                 350

Trp Thr Ser Pro Gln Leu Leu Cys Ser Lys Lys Gly Ala Ser Val Lys
        355                 360                 365

Gly Gly Leu Gly Pro Phe Gly Leu Leu Thr Leu Ala Ser Xaa Gly Leu
    370                 375                 380

Glu Glu Tyr Thr Ala Val Phe Phe Arg Ile Phe Lys Ala Tyr Asp Asn
385                 390                 395                 400

Lys Phe Val Val Leu Met Cys Ser Asp Gln Ser Arg Ser Ser Leu Asn
                405                 410                 415

Pro Thr Asn Asp Lys Thr Thr Tyr Gly Thr Phe Val Asp Val Asn Pro
            420                 425                 430

Ile Arg Glu Gly Leu Ser Leu Arg Val Leu Ile Asp His Ser Val Val
        435                 440                 445

Glu Ser Phe Gly Ala Lys Gly Lys Asn Val Ile Thr Ala Arg Val Tyr
    450                 455                 460

Pro Thr Leu Ala Ile Asn Glu Lys Ala His Leu Tyr Val Phe Asn Arg
465                 470                 475                 480
```

```
Gly Thr Ser Asn Val Glu Ile Thr Gly Leu Thr Ala Trp Ser Met Lys
                485                 490                 495

Lys Ala Asn Ile Ala
            500
```

<210> 27
<211> 518
<212> PRT
<213> Bifidobacterium breve

<400> 27

```
Met Thr Asp Phe Thr Pro Glu Thr Pro Val Leu Thr Pro Ile Arg Asp
1               5                   10                  15

His Ala Ala Glu Leu Ala Lys Ala Glu Ala Gly Val Ala Glu Met Ala
            20                  25                  30

Ala Lys Arg Asn Asn Arg Trp Tyr Pro Lys Tyr His Ile Ala Ser Asn
        35                  40                  45

Gly Gly Trp Ile Asn Asp Pro Asn Gly Leu Cys Phe Tyr Lys Gly Arg
    50                  55                  60

Trp His Val Phe Tyr Gln Leu His Pro Tyr Gly Thr Gln Trp Gly Pro
65                  70                  75                  80

Met His Trp Gly His Val Ser Ser Thr Asp Met Leu Asn Trp Lys Arg
                85                  90                  95

Glu Pro Ile Met Phe Ala Pro Ser Leu Glu Gln Glu Lys Asp Gly Val
            100                 105                 110

Phe Ser Gly Ser Ala Val Ile Asp Asp Asn Gly Asp Leu Arg Phe Tyr
        115                 120                 125

Tyr Thr Gly His Arg Trp Ala Asn Gly His Asp Asn Thr Gly Gly Asp
    130                 135                 140

Trp Gln Val Gln Met Thr Ala Leu Pro Asp Asn Asp Glu Leu Thr Ser
145                 150                 155                 160

Ala Thr Lys Gln Gly Met Ile Ile Asp Cys Pro Thr Asp Lys Val Asp
                165                 170                 175

His His Tyr Arg Asp Pro Lys Val Trp Lys Thr Gly Asp Thr Trp Tyr
            180                 185                 190

Met Thr Phe Gly Val Ser Ser Glu Asp Lys Arg Gly Gln Met Trp Leu
        195                 200                 205

Phe Ser Ser Lys Asp Met Val Arg Trp Glu Tyr Glu Arg Val Leu Phe
```

210 215 220

Gln His Pro Asp Pro Asp Val Phe Met Leu Glu Cys Pro Asp Phe Phe
225 230 235 240

Pro Ile Lys Asp Lys Asp Gly Asn Glu Lys Trp Val Ile Gly Phe Ser
245 250 255

Ala Met Gly Ser Lys Pro Ser Gly Phe Met Asn Arg Asn Val Asn Asn
260 265 270

Ala Gly Tyr Met Ile Gly Thr Trp Glu Pro Gly Gly Glu Phe Lys Pro
275 280 285

Glu Thr Glu Phe Arg Leu Trp Asp Cys Gly His Asn Tyr Tyr Ala Pro
290 295 300

Gln Ser Phe Asn Val Asp Gly Arg Gln Ile Val Tyr Gly Trp Met Ser
305 310 315 320

Pro Phe Val Gln Pro Ile Pro Met Glu Asp Asp Gly Trp Cys Gly Gln
325 330 335

Leu Thr Leu Pro Arg Glu Ile Thr Leu Asp Asp Asp Gly Asp Val Val
340 345 350

Thr Ala Pro Val Ala Glu Met Glu Gly Leu Arg Glu Asp Thr Leu Asp
355 360 365

His Gly Ser Ile Thr Leu Asp Met Asp Gly Glu Gln Val Ile Ala Asp
370 375 380

Asp Ala Glu Ala Val Glu Ile Glu Met Thr Ile Asp Leu Ala Ala Ser
385 390 395 400

Thr Ala Asp Arg Ala Gly Leu Lys Ile His Ala Thr Glu Asp Gly Ala
405 410 415

Tyr Thr Tyr Val Ala Tyr Asp Asp Gln Ile Gly Arg Val Val Val Asp
420 425 430

Arg Gln Ala Met Ala Asn Gly Asp His Gly Tyr Arg Ala Ala Pro Leu
435 440 445

Thr Asp Ala Glu Leu Ala Ser Gly Lys Leu Asp Leu Arg Val Phe Val
450 455 460

Asp Arg Gly Ser Val Glu Val Tyr Val Asn Gly Gly His Gln Val Leu
465 470 475 480

Ser Ser Tyr Ser Tyr Ala Ser Glu Gly Pro Arg Ala Ile Lys Leu Val

485 490 495

Ala Glu Phe Gly Asn Leu Lys Val Glu Ser Leu Lys Leu His His Met
500 505 510

Lys Ser Ile Gly Leu Glu
515

<210> 28
<211> 532
<212> PRT
<213> Saccharomyces cerevisiae

<400> 28
Met Leu Leu Gln Ala Phe Leu Phe Leu Leu Ala Gly Phe Ala Ala Lys
1 5 10 15

Ile Ser Ala Ser Met Thr Asn Glu Thr Ser Asp Arg Pro Leu Val His
20 25 30

Phe Thr Pro Asn Lys Gly Trp Met Asn Asp Pro Asn Gly Leu Trp Tyr
35 40 45

Asp Glu Lys Asp Ala Lys Trp His Leu Tyr Phe Gln Tyr Asn Pro Asn
50 55 60

Asp Thr Val Trp Gly Thr Pro Leu Phe Trp Gly His Ala Thr Ser Asp
65 70 75 80

Asp Leu Thr Asn Trp Glu Asp Gln Pro Ile Ala Ile Ala Pro Lys Arg
85 90 95

Asn Asp Ser Gly Ala Phe Ser Gly Ser Met Val Val Asp Tyr Asn Asn
100 105 110

Thr Ser Gly Phe Phe Asn Asp Thr Ile Asp Pro Arg Gln Arg Cys Val
115 120 125

Ala Ile Trp Thr Tyr Asn Thr Pro Glu Ser Glu Glu Gln Tyr Ile Ser
130 135 140

Tyr Ser Leu Asp Gly Gly Tyr Thr Phe Thr Glu Tyr Gln Lys Asn Pro
145 150 155 160

Val Leu Ala Ala Asn Ser Thr Gln Phe Arg Asp Pro Lys Val Phe Trp
165 170 175

Tyr Glu Pro Ser Gln Lys Trp Ile Met Thr Ala Ala Lys Ser Gln Asp
180 185 190

Tyr Lys Ile Glu Ile Tyr Ser Ser Asp Asp Leu Lys Ser Trp Lys Leu
195 200 205

```
Glu Ser Ala Phe Ala Asn Glu Gly Phe Leu Gly Tyr Gln Tyr Glu Cys
    210                 215                 220

Pro Gly Leu Ile Glu Val Pro Thr Glu Gln Asp Pro Ser Lys Ser Tyr
225                 230                 235                 240

Trp Val Met Phe Ile Ser Ile Asn Pro Gly Ala Pro Ala Gly Gly Ser
                245                 250                 255

Phe Asn Gln Tyr Phe Val Gly Ser Phe Asn Gly Thr His Phe Glu Ala
            260                 265                 270

Phe Asp Asn Gln Ser Arg Val Val Asp Phe Gly Lys Asp Tyr Tyr Ala
        275                 280                 285

Leu Gln Thr Phe Phe Asn Thr Asp Pro Thr Tyr Gly Ser Ala Leu Gly
    290                 295                 300

Ile Ala Trp Ala Ser Asn Trp Glu Tyr Ser Ala Phe Val Pro Thr Asn
305                 310                 315                 320

Pro Trp Arg Ser Ser Met Ser Leu Val Arg Lys Phe Ser Leu Asn Thr
                325                 330                 335

Glu Tyr Gln Ala Asn Pro Glu Thr Glu Leu Ile Asn Leu Lys Ala Glu
            340                 345                 350

Pro Ile Leu Asn Ile Ser Asn Ala Gly Pro Trp Ser Arg Phe Ala Thr
        355                 360                 365

Asn Thr Thr Leu Thr Lys Ala Asn Ser Tyr Asn Val Asp Leu Ser Asn
    370                 375                 380

Ser Thr Gly Thr Leu Glu Phe Glu Leu Val Tyr Ala Val Asn Thr Thr
385                 390                 395                 400

Gln Thr Ile Ser Lys Ser Val Phe Ala Asp Leu Ser Leu Trp Phe Lys
                405                 410                 415

Gly Leu Glu Asp Pro Glu Glu Tyr Leu Arg Met Gly Phe Glu Val Ser
            420                 425                 430

Ala Ser Ser Phe Phe Leu Asp Arg Gly Asn Ser Lys Val Lys Phe Val
        435                 440                 445

Lys Glu Asn Pro Tyr Phe Thr Asn Arg Met Ser Val Asn Asn Gln Pro
    450                 455                 460

Phe Lys Ser Glu Asn Asp Leu Ser Tyr Tyr Lys Val Tyr Gly Leu Leu
465                 470                 475                 480
```

Asp Gln Asn Ile Leu Glu Leu Tyr Phe Asn Asp Gly Asp Val Val Ser
485 490 495

Thr Asn Thr Tyr Phe Met Thr Thr Gly Asn Ala Leu Gly Ser Val Asn
500 505 510

Met Thr Thr Gly Val Asp Asn Leu Phe Tyr Ile Asp Lys Phe Gln Val
515 520 525

Arg Glu Val Lys
530

<210> 29
<211> 512
<212> PRT
<213> Zymomonas mobilis

<400> 29
Met Glu Ser Pro Ser Tyr Lys Asn Leu Ile Lys Ala Glu Asp Ala Gln
1 5 10 15

Lys Lys Ala Gly Lys Arg Leu Leu Ser Ser Glu Trp Tyr Pro Gly Phe
20 25 30

His Val Thr Pro Leu Thr Gly Trp Met Asn Asp Pro Asn Gly Leu Ile
35 40 45

Phe Phe Lys Gly Glu Tyr His Leu Phe Tyr Gln Tyr Tyr Pro Phe Ala
50 55 60

Pro Val Trp Gly Pro Met His Trp Gly His Ala Lys Ser Arg Asp Leu
65 70 75 80

Val His Trp Glu Thr Leu Pro Val Ala Leu Ala Pro Gly Asp Leu Phe
85 90 95

Asp Arg Asp Gly Cys Phe Ser Gly Cys Ala Val Asp Asn Asn Gly Val
100 105 110

Leu Thr Leu Ile Tyr Thr Gly His Ile Val Leu Ser Asn Asp Ser Pro
115 120 125

Asp Ala Ile Arg Glu Val Gln Cys Met Ala Thr Ser Ile Asp Gly Ile
130 135 140

His Phe Gln Lys Glu Gly Ile Val Leu Glu Lys Ala Pro Met Pro Gln
145 150 155 160

Val Ala His Phe Arg Asp Pro Arg Val Trp Lys Glu Asn Asp His Trp
165 170 175

Phe Met Val Val Gly Tyr Arg Thr Asp Asp Glu Lys His Gln Gly Ile

180 185 190

Gly His Val Ala Leu Tyr Arg Ser Glu Asn Leu Lys Asp Trp Ile Phe
195 200 205

Val Lys Thr Leu Leu Gly Asp Asn Ser Gln Leu Pro Leu Gly Lys Arg
210 215 220

Ala Phe Met Trp Glu Cys Pro Asp Phe Phe Ser Leu Gly Asn Arg Ser
225 230 235 240

Val Leu Met Phe Ser Pro Gln Gly Leu Lys Ala Ser Gly Tyr Lys Asn
245 250 255

Arg Asn Leu Phe Gln Asn Gly Tyr Ile Leu Gly Lys Trp Gln Ala Pro
260 265 270

Gln Phe Thr Pro Glu Thr Ser Phe Gln Glu Leu Asp Tyr Gly His Asp
275 280 285

Phe Tyr Ala Ala Gln Arg Phe Glu Ala Lys Asp Gly Arg Gln Ile Leu
290 295 300

Ile Ala Trp Phe Asp Met Trp Glu Asn Gln Lys Pro Ser Gln Arg Asp
305 310 315 320

Gly Trp Ala Gly Cys Met Thr Leu Pro Arg Lys Leu Asp Leu Ile Asp
325 330 335

Asn Lys Ile Val Met Thr Pro Val Arg Glu Met Glu Ile Leu Arg Gln
340 345 350

Ser Glu Lys Ile Glu Ser Val Val Thr Leu Ser Asp Ala Glu His Pro
355 360 365

Phe Thr Met Asp Ser Pro Leu Gln Glu Ile Glu Leu Ile Phe Asp Leu
370 375 380

Glu Lys Ser Ser Ala Tyr Gln Ala Gly Leu Ala Leu Arg Cys Asn Gly
385 390 395 400

Lys Gly Gln Glu Thr Leu Leu Tyr Ile Asp Arg Ser Gln Asn Arg Ile
405 410 415

Ile Leu Asp Arg Asn Arg Ser Gly Gln Asn Val Lys Gly Ile Arg Ser
420 425 430

Cys Pro Leu Pro Asn Thr Ser Lys Val Arg Leu His Ile Phe Leu Asp
435 440 445

Arg Ser Ser Ile Glu Ile Phe Val Gly Asp Asp Gln Thr Gln Gly Leu

450 455 460

Tyr Ser Ile Ser Ser Arg Ile Phe Pro Asp Lys Asp Ser Leu Lys Gly
465 470 475 480

Arg Leu Phe Ala Ile Glu Gly Tyr Ala Val Phe Asp Ser Phe Lys Arg
485 490 495

Trp Thr Leu Gln Asp Ala Asn Leu Ala Ala Phe Ser Ser Asp Ala Cys
500 505 510

<210> 30
<211> 987
<212> DNA
<213> Chlorella luteoviridis

<400> 30
tgttgaagaa tgagccggcg acttatagga agtggcttgg ttaaggatac tttccgaagc 60

ctaagcgaaa gcaagttgta acaatagcga tatacctctt tgtaggtcag tcacttctta 120

tggacccgaa cccgggtgat ctaaccatga ccaggatgaa gcttgggtaa caccaagtga 180

aggtccgaac tcttcgatct ttaaaaatcg tgagatgagt tatggttagg ggtaaatctg 240

gcagttttgc cccgcaaaag ggtaaccttt tgtaattact gactcataac ggtgaagcct 300

aaggcgttag ctatggtaat accgtgggaa gtttcaatac cttcttgcat atttttttatt 360

tgcaccttta gtgcaaacag tgtaaagaaa gcgttttgaa acccccttaac gactaatttt 420

ttgcttttgc aagaacgtca gcactcacca atacactttc cgtttttttc ttttattaat 480

taaagcaaca taaaaatata ttttatagct ttaatcataa aactatgtta gcacttcgtg 540

ctaatgtgct aatgtgctaa tcaaatgaaa agtgttctta aaagtgagtt gaaggtagag 600

tctaatcttg cctgaaaggg caagctgcac attttttttt gaatgtgcaa caatggaaat 660

gccaatcgaa ctcggagcta gctggttctc cccgaaatgt gttgaggcgc agcgattcat 720

gattagtacg gtgtaggggt aaagcactgt ttcggtgcgg gctgtgaaaa cggtaccaaa 780

tcgtggcaaa ctaagaatac tacgcttgta taccatggat cagtgagact atgggggata 840

agctccatag tcaagaggga aacagcccag atcaccagtt aaggccccaa aatgacagct 900

aagtggcaaa ggaggtgaaa gtgcagaaac aaccaggagg tttgcccaga agcagccatc 960

ctttaaagag tgcgtaatag ctcactg 987

<210> 31
<211> 1412
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic polynucleotide

<400> 31
gaattcgagt ttaggtccag cgtccgtggg gggggacggg ctgggagctt gggccgggaa 60

gggcaagacg atgcagtccc tctggggagt cacagccgac tgtgtgtgtt gcactgtgcg 120

gcccgcagca ctcacacgca aaatgcctgg ccgacaggca ggccctgtcc agtgcaacat 180

ccacggtccc tctcatcagg ctcaccttgc tcattgacat aacggaatgc gtaccgctct 240

ttcagatctg tccatccaga gaggggagca ggctccccac cgacgctgtc aaacttgctt 300

cctgcccaac cgaaaacatt attgtttgag gggggggggg gggggcaga ttgcatggcg 360

ggatatctcg tgaggaacat cactgggaca ctgtggaaca cagtgagtgc agtatgcaga 420

gcatgtatgc taggggtcag cgcaggaagg gggcctttcc cagtctccca tgccactgca 480

ccgtatccac gactcaccag gaccagcttc ttgatcggct tccgctcccg tggacaccag 540

tgtgtagcct ctggactcca ggtatgcgtg caccgcaaag gccagccgat cgtgccgatt 600

cctggggtgg aggatatgag tcagccaact tggggctcag agtgcacact ggggcacgat 660

acgaaacaac atctacaccg tgtcctccat gctgacacac cacagcttcg ctccacctga 720

atgtgggcgc atgggcccga atcacagcca atgtcgctgc tgccataatg tgatccagac 780

cctctccgcc cagatgccga gcggatcgtg ggcgctgaat agattcctgt ttcgatcact 840

gtttgggtcc tttccttttc gtctcggatg cgcgtctcga aacaggctgc gtcgggcttt 900

cggatccctt ttgctccctc cgtcaccatc ctgcgcgcgg gcaagttgct tgaccctggg 960

ctggtaccag ggttggaggg tattaccgcg tcaggccatt cccagcccgg attcaattca 1020

aagtctgggc caccaccctc cgccgctctg tctgatcact ccacattcgt gcatacacta 1080

cgttcaagtc ctgatccagg cgtgtctcgg gacaaggtgt gcttgagttt gaatctcaag 1140

gacccactcc agcacagctg ctggttgacc ccgccctcgc aactccctac catgtctgct 1200

ggtaggtcca gggatctttg ccatgcacac aggaccccgt ttgtgggggt ccccggtgca 1260

tgctgtcgct gtgcaggcgc cggtgtgggg cctgggcccc gcgggagctc aactcctccc 1320

catatgcctg ccgtccctcc cacccaccgc gacctggccc cctttgcaga ggaaggcgaa 1380

gtcagcgcca tcgtgtgcga taatggatcc gg 1412

<210> 32
<211> 1627
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic polynucleotide

<400> 32
gaattcgccc ttgagtttag gtccagcgtc cgtgggggggg gcgtgagact ccccccctgac 60

cttcgtatgg cagggactcc tacttgccaa gtaatcagtt gacaatgcca cttcaatgct 120

cgttgtggta cactgacgcg ggtctaacat actgggaagc atgaattgcc gacatggact 180

cagttggaga cagtaacagc tctttgtgtt ctatcttcag gaacacattt ggcagcgcac 240

ccatacagtg gcgcacacgc agctgtacct gatgtggctc tattcccaca tgtttcaact 300

tgatccaaaa gtcactcaga ctctcagcag ctagacttga tcgcatcttt ggccatgaag 360

atgcttgcgc aactctagga atgggacgag aaaagagcct gctctgatcg gatatttcca 420
ttctctggat gggactgaga tgattctgaa gaaatgctgc tcgacttatt tggaagaaca 480
gcacctgacg catgctttga ggctgctgtg gctgggatgt gctgtatttg tcagcattga 540
gcatctacgg gtagatggcc ataaccacgc gctgcctatc atgcggtggg ttgtgtggaa 600
aacgtacaat ggacagaaat caatcccatt gcgagcctag cgtgcagcca tgcgctccct 660
ctgtagcccc gctccaagac aaagccagcc aatgccagaa cccacataga gagggtatct 720
tcctaatgac ctcgcccatc atttcctcca aattaactat aatgccttga ttgtggagtt 780
ggctttggct tgcagctgct cgcgctggca cttttgtagg cagcacaggg tatgccagcg 840
ccgaactttg tgcccttgag caggccacaa gggcacaaga ctacaccatg cagctggtat 900
acttggaact gataccattc ttaccaagca aggcacagca cagcctgcac cgactcactt 960
tgcttgagcg gggcacagcg ccgcgactga tcctgcgagc tgtggggagt tccgactgtt 1020
ctggacctcg gtctctgaaa gatgtgtacg atgggatcaa gtcattcaag tatgctcttc 1080
acatgagcaa tcgggggaga cacggtggcc ctaaaggtgt tcatctgatt caagtgtagt 1140
ggggggtgtgc tgtttgtccc ggggcgcccc ccgctccccg accccggaga agggccccag 1200
aggactcggc cgcccacaga ggaataaccg ggcgtggctc ggccctgcgc ctccctcttt 1260
caatatttca cctggtgttc agtgcacgga cacgtaaaga actagataca atggccgagg 1320
gaaagacggt gagagcttgg cgttggtgga ccgggcagca tcagaaactc ctcttccccg 1380
cccgccttga aactcactgt aactccctcc tcttcccct cgcagcatct gtctatcgtt 1440
atcgtgagtg aaagggactg ccatgtgtcg ggtcgttgac cacggtcggc tcgggcgctg 1500
ctgcccgcgt cgcgaacgtt ccctgcaaac gccgcgcagc cgtccctttt tctgccgccg 1560
ccccacccc tcgctccccc cttcaatcac accgcagtgc ggacatgtcg attccggcaa 1620
gtccacc 1627

<210> 33
<211> 570
<212> DNA
<213> Chlorella protothecoides

<400> 33
gaattccctg caggaagaag gccggcagca gctggtactt gtccttcacc tccttgatcg 60
gctgggtgag cttggccggg tcgcagtcgt cgatgccggc atcgcccagc acgctgtgcg 120
gggagccggc atcgacaacc ttggcactgc tcaccttggt caccggcatg gggtcatggc 180
gctgcagacc agcggcctgt cagcatgctg caggcatctg tgttttgtag tagatacttt 240
ctgatgcatc accacacgtt tggaaggtcc ccaagcccct tcaacagtct cgacatatga 300
cactcgcgcc ctcttcctcg tcccgtggcc tgatgagggt acgcaggtac cgcagctgcg 360
ccccgtcccg ccagttgccc tggccccgcc gggcccaatc tgttcattgc cgctccctgg 420
cagccgtgaa cttcacacta ccgctctctg tgaccttcag cacagcagga atcgccattt 480

caccggcggt cgttgctgcg gagcctcagc tgatctcgcc tgcgagaccc cacagtttga 540

atttgcggtc cccacacaac ctctgacgcc 570

<210> 34
<211> 568
<212> DNA
<213> Chlorella protothecoides

<220>
<221> modified_base
<222> (133)..(133)
<223> a, c, t, g, unknown or other

<400> 34
gaattccctc aggaagaagg ccggcagcag ctggtacttg tccttcacct ccttgatcgg 60

ctgggtgagc ttcgcaggat cgcagtcgtc gatgccggca tcgcccagca cgctgtgcgg 120

ggagccggca tcnacaacct tggcactgct ccccttggtc accggcatgg ggtcatggcg 180

ctgcagccca gcggcctgtc agcatgctgc aggcatctgt gtattgtagt aggtacttcc 240

tgatgcatca acacacgttt ggaagctccc caagcccctt caacagtctc gacgtatgac 300

actcgcgccc tcttcctcgc cccgtggcct gatgagggta cgcaggtacc acagctgcgc 360

cccgtcccgc cagttgccct ggcccggccg ggcccaatct gttcattgcc gctccctggt 420

agccgtgaac tcacattacc gctctctgtg accttcagca cagcaggaat cgccatttca 480

ccggcggtcg ttgctgcgga gcctcagctg atctcgcctg cgagacccca cagtttgaat 540

ttgcggtccc cacacaacct ctgacgcc 568

<210> 35
<211> 50
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
primer

<400> 35
tgacctaggt gattaattaa ctcgaggcag cagcagctcg gatagtatcg 50

<210> 36
<211> 45
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
primer

<400> 36
ctacgagctc aagctttcca tttgtgttcc catcccacta cttcc 45

<210> 37
<211> 32
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic primer

<400> 37
gatcagaatt ccgcctgcaa cgcaagggca gc 32

<210> 38
<211> 30
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic primer

<400> 38
gcatactagt ggcgggacgg agagagggcg 30

<210> 39
<211> 1568
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic polynucleotide

<400> 39
gaattccttt cttgcgctat gacacttcca gcaaaaggta gggcgggctg cgagacggct 60

tcccggcgct gcatgcaaca ccgatgatgc ttcgaccccc cgaagctcct tcggggctgc 120

atgggcgctc cgatgccgct ccagggcgag cgctgtttaa atagccaggc ccccgattgc 180

aaagacatta tagcgagcta ccaaagccat attcaaacac ctagatcact accacttcta 240

cacaggccac tcgagcttgt gatcgcactc cgctaagggg gcgcctcttc ctcttcgttt 300

cagtcacaac ccgcaaacgg cgcgccatat caatgattga acaagatgga ttgcacgcag 360

gttctccggc cgcttgggtg gagaggctat tcggctatga ctgggcacaa cagacaatcg 420

gctgctctga tgccgccgtg ttccggctgt cagcgcaggg gcgcccggtt ctttttgtca 480

agaccgacct gtccggtgcc ctgaatgaac tgcaggacga ggcagcgcgg ctatcgtggc 540

tggccacgac gggcgttcct tgcgcagctg tgctcgacgt tgtcactgaa gcgggaaggg 600

actggctgct attgggcgaa gtgccggggc aggatctcct gtcatctcac cttgctcctg 660

ccgagaaagt atccatcatg gctgatgcaa tgcggcggct gcatacgctt gatccggcta 720

cctgcccatt cgaccaccaa gcgaaacatc gcatcgagcg agcacgtact cggatggaag 780

ccggtcttgt cgatcaggat gatctggacg aagagcatca ggggctcgcg ccagccgaac 840

tgttcgccag gctcaaggcg cgcatgcccg acggcgagga tctcgtcgtg acccatggcg 900

atgcctgctt gccgaatatc atggtggaaa atggccgctt ttctggattc atcgactgtg 960

gccggctggg tgtggcggac cgctatcagg acatagcgtt ggctacccgt gatattgctg 1020

aagagcttgg cggcgaatgg gctgaccgct tcctcgtgct ttacggtatc gccgctcccg 1080

attcgcagcg catcgccttc tatcgccttc ttgacgagtt cttctaagat ctgtcgatcg 1140

acaagtgact cgaggcagca gcagctcgga tagtatcgac acactctgga cgctggtcgt 1200

gtgatggact gttgccgcca cacttgctgc cttgacctgt gaatatccct gccgctttta 1260

tcaaacagcc tcagtgtgtt tgatcttgtg tgtacgcgct tttgcgagtt gctagctgct 1320

tgtgctattt gcgaatacca cccccagcat ccccttccct cgtttcatat cgcttgcatc 1380

ccaaccgcaa cttatctacg ctgtcctgct atccctcagc gctgctcctg ctcctgctca 1440

ctgcccctcg cacagccttg gtttgggctc cgcctgtatt ctcctggtac tgcaacctgt 1500

aaaccagcac tgcaatgctg atgcacggga agtagtggga tgggaacaca aatggaaagc 1560

ttgagctc 1568

<210> 40
<211> 2571
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic polynucleotide

<220>
<221> modified_base
<222> (997)..(999)
<223> a, c, t, g, unknown or other

<400> 40
gaattccgcc tgcaacgcaa gggcagccac agccgctccc acccgccgct gaaccgacac 60

gtgcttgggc gcctgccgcc tgcctgccgc atgcttgtgc tggtgaggct gggcagtgct 120

gccatgctga ttgaggcttg gttcatcggg tggaagctta tgtgtgtgct gggcttgcat 180

gccgggcaat gcgcatggtg gcaagagggc ggcagcactt gctggacgtg ccgcggtgcc 240

tccaggtggt tcaatcgcgg cagccagagg gatttcagat gatcgcgcgt acaggttgag 300

cagcagtgtc agcaaaggta gcagtttgcc agaatgatcg gttcagctgt taatcaatgc 360

cagcaagaga aggggtcaag tgcaaacacg ggcatgccac agcacgggca ccggggagtg 420

gaatggcacc accaagtgtg tgcgagccag catcgccgcc tggctgtttc agctacaacg 480

gcaggagtca tccaactaac catagctgat caacactgca atcatcggcg gctgatgcaa 540

gcatcctgca agacacatgc tgtgcgatgc tgcgctgctg cctgctgcgc acgccgttga 600

gttggcagca gctcagccat gcactggatc aggctgggct gccactgcaa tgtggtggat 660

aggatgcaag tggagcgaat accaaaccct ctggctgctt gctgggttgc atggcatcgc 720

accatcagca ggagcgcatg cgaagggact ggccccatgc acgccatgcc aaaccggagc 780

gcaccgagtg tccacactgt caccaggccc gcaagctttg cagaaccatg ctcatggacg 840

catgtagcgc tgacgtccct tgacggcgct cctctcgggt gtgggaaacg caatgcagca 900

caggcagcag aggcggcggc agcagagcgg cggcagcagc ggcgggggcc acccttcttg 960

cggggtcgcg ccccagccag cggtgatgcg ctgatcnnnc caaacgagtt cacattcatt 1020

tgcagcctgg agaagcgagg ctggggcctt tgggctggtg cagcccgcaa tggaatgcgg 1080

gaccgccagg ctagcagcaa aggcgcctcc cctactccgc atcgatgttc catagtgcat 1140

tggactgcat ttgggtgggg cggccggctg tttctttcgt gttgcaaaac gcgccacgtc 1200

agcaacctgt cccgtgggtc ccccgtgccg atgaaatcgt gtgcacgccg atcagctgat 1260

tgcccggctc gcgaagtagg cgccctcttt ctgctcgccc tctctccgtc ccgccactag 1320

tggcgcgcca tatcaatgat tgaacaagat ggattgcacg caggttctcc ggccgcttgg 1380

gtggagaggc tattcggcta tgactgggca caacagacaa tcggctgctc tgatgccgcc 1440

gtgttccggc tgtcagcgca ggggcgcccg gttctttttg tcaagaccga cctgtccggt 1500

gccctgaatg aactgcagga cgaggcagcg cggctatcgt ggctggccac gacgggcgtt 1560

ccttgcgcag ctgtgctcga cgttgtcact gaagcgggaa gggactggct gctattgggc 1620

gaagtgccgg ggcaggatct cctgtcatct caccttgctc ctgccgagaa agtatccatc 1680

atggctgatg caatgcggcg gctgcatacg cttgatccgg ctacctgccc attcgaccac 1740

caagcgaaac atcgcatcga gcgagcacgt actcggatgg aagccggtct tgtcgatcag 1800

gatgatctgg acgaagagca tcaggggctc gcgccagccg aactgttcgc caggctcaag 1860

gcgcgcatgc ccgacggcga ggatctcgtc gtgacccatg gcgatgcctg cttgccgaat 1920

atcatggtgg aaaatggccg cttttctgga ttcatcgact gtggccggct gggtgtggcg 1980

gaccgctatc aggacatagc gttggctacc cgtgatattg ctgaagagct tggcggcgaa 2040

tgggctgacc gcttcctcgt gctttacggt atcgccgctc ccgattcgca gcgcatcgcc 2100

ttctatcgcc ttcttgacga gttcttctaa gatctgtcga tcgacaagtg actcgaggca 2160

gcagcagctc ggatagtatc gacacactct ggacgctggt cgtgtgatgg actgttgccg 2220

ccacacttgc tgccttgacc tgtgaatatc cctgccgctt ttatcaaaca gcctcagtgt 2280

gtttgatctt gtgtgtacgc gcttttgcga gttgctagct gcttgtgcta tttgcgaata 2340

ccacccccag catccccttc cctcgtttca tatcgcttgc atcccaaccg caacttatct 2400

acgctgtcct gctatccctc agcgctgctc ctgctcctgc tcactgcccc tcgcacagcc 2460

ttggtttggg ctccgcctgt attctcctgg tactgcaacc tgtaaaccag cactgcaatg 2520

ctgatgcacg ggaagtagtg ggatgggaac acaaatggaa agcttgagct c 2571

<210> 41
<211> 2550
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic polynucleotide

<220>
<221> modified_base
<222> (997)..(999)
<223> a, c, t, g, unknown or other

<400> 41
gaattccgcc tgcaacgcaa gggcagccac agccgctccc acccgccgct gaaccgacac 60
gtgcttgggc gcctgccgcc tgcctgccgc atgcttgtgc tggtgaggct gggcagtgct 120
gccatgctga ttgaggcttg gttcatcggg tggaagctta tgtgtgtgct gggcttgcat 180
gccgggcaat gcgcatggtg gcaagagggc ggcagcactt gctggacgtg ccgcggtgcc 240
tccaggtggt tcaatcgcgg cagccagagg gatttcagat gatcgcgcgt acaggttgag 300
cagcagtgtc agcaaaggta gcagtttgcc agaatgatcg gttcagctgt taatcaatgc 360
cagcaagaga aggggtcaag tgcaaacacg ggcatgccac agcacgggca ccggggagtg 420
gaatggcacc accaagtgtg tgcgagccag catcgccgcc tggctgtttc agctacaacg 480
gcaggagtca tccaactaac catagctgat caacactgca atcatcggcg gctgatgcaa 540
gcatcctgca agacacatgc tgtgcgatgc tgcgctgctg cctgctgcgc acgccgttga 600
gttggcagca gctcagccat gcactggatc aggctgggct gccactgcaa tgtggtggat 660
aggatgcaag tggagcgaat accaaaccct ctggctgctt gctgggttgc atggcatcgc 720
accatcagca ggagcgcatg cgaagggact ggccccatgc acgccatgcc aaaccggagc 780
gcaccgagtg tccacactgt caccaggccc gcaagctttg cagaaccatg ctcatggacg 840
catgtagcgc tgacgtccct tgacggcgct cctctcgggt gtgggaaacg caatgcagca 900
caggcagcag aggcggcggc agcagagcgg cggcagcagc ggcgggggcc acccttcttg 960
cggggtcgcg ccccagccag cggtgatgcg ctgatcnnnc caaacgagtt cacattcatt 1020
tgcagcctgg agaagcgagg ctggggcctt tgggctggtg cagcccgcaa tggaatgcgg 1080
gaccgccagg ctagcagcaa aggcgcctcc cctactccgc atcgatgttc catagtgcat 1140
tggactgcat ttgggtgggg cggccggctg tttctttcgt gttgcaaaac gcgccacgtc 1200
agcaacctgt cccgtgggtc ccccgtgccg atgaaatcgt gtgcacgccg atcagctgat 1260
tgcccggctc gcgaagtagg cgccctcttt ctgctcgccc tctctccgtc ccgccactag 1320
tggcgcgcca tatcaatgat cgagcaggac ggcctccacg ccggctcccc cgccgcctgg 1380
gtggagcgcc tgttcggcta cgactgggcc cagcagacca tcggctgctc cgacgccgcc 1440
gtgttccgcc tgtccgccca gggccgcccc gtgctgttcg tgaagaccga cctgtccggc 1500
gccctgaacg agctgcagga cgaggccgcc cgcctgtcct ggctggccac caccggcgtg 1560
ccctgcgccg ccgtgctgga cgtggtgacc gaggccggcc gcgactggct gctgctgggc 1620
gaggtgcccg gccaggacct gctgtcctcc cacctggccc ccgccgagaa ggtgtccatc 1680
atggccgacg ccatgcgccg cctgcacacc ctggaccccg ccacctgccc cttcgaccac 1740
caggccaagc accgcatcga gcgcgcccgc acccgcatgg aggccggcct ggtggaccag 1800
gacgacctgg acgaggagca ccagggcctg gcccccgccg agctgttcgc ccgcctgaag 1860
gcccgcatgc ccgacggcga ggacctggtg gtgacccacg gcgacgcctg cctgcccaac 1920
atcatggtgg agaacggccg cttctccggc ttcatcgact gcggccgcct gggcgtggcc 1980

gaccgctacc aggacatcgc cctggccacc cgcgacatcg ccgaggagct gggcggcgag 2040

tgggccgacc gcttcctggt gctgtacggc atcgccgccc ccgactccca gcgcatcgcc 2100

ttctaccgcc tgctggacga gttcttctga ctcgaggcag cagcagctcg gatagtatcg 2160

acacactctg gacgctggtc gtgtgatgga ctgttgccgc cacacttgct gccttgacct 2220

gtgaatatcc ctgccgcttt tatcaaacag cctcagtgtg tttgatcttg tgtgtacgcg 2280

cttttgcgag ttgctagctg cttgtgctat ttgcgaatac cacccccagc atccccttcc 2340

ctcgtttcat atcgcttgca tcccaaccgc aacttatcta cgctgtcctg ctatccctca 2400

gcgctgctcc tgctcctgct cactgccccc cgcacagcct tggtttgggc tccgcctgta 2460

ttctcctggt actgcaacct gtaaaccagc actgcaatgc tgatgcacgg gaagtagtgg 2520

gatgggaaca caaatggaaa gcttgagctc 2550

<210> 42
<211> 1547
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic polynucleotide

<400> 42
gaattccttt cttgcgctat gacacttcca gcaaaaggta gggcgggctg cgagacggct 60

tcccggcgct gcatgcaaca ccgatgatgc ttcgaccccc cgaagctcct tcggggctgc 120

atgggcgctc cgatgccgct ccagggcgag cgctgtttaa atagccaggc ccccgattgc 180

aaagacatta tagcgagcta ccaaagccat attcaaacac ctagatcact accacttcta 240

cacaggccac tcgagcttgt gatcgcactc cgctaagggg gcgcctcttc ctcttcgttt 300

cagtcacaac ccgcaaacgg cgcgccatat caatgatcga gcaggacggc ctccacgccg 360

gctcccccgc cgcctgggtg gagcgcctgt tcggctacga ctgggcccag cagaccatcg 420

gctgctccga cgccgccgtg ttccgcctgt ccgcccaggg ccgccccgtg ctgttcgtga 480

agaccgacct gtccggcgcc ctgaacgagc tgcaggacga ggccgcccgc ctgtcctggc 540

tggccaccac cggcgtgccc tgcgccgccg tgctggacgt ggtgaccgag gccggccgcg 600

actggctgct gctgggcgag gtgcccggcc aggacctgct gtcctcccac ctggcccccg 660

ccgagaaggt gtccatcatg gccgacgcca tgcgccgcct gcacaccctg gaccccgcca 720

cctgcccctt cgaccaccag gccaagcacc gcatcgagcg cgcccgcacc cgcatggagg 780

ccggcctggt ggaccaggac gacctggacg aggagcacca gggcctggcc cccgccgagc 840

tgttcgcccg cctgaaggcc cgcatgcccg acggcgagga cctggtggtg acccacggcg 900

acgcctgcct gcccaacatc atggtggaga acggccgctt ctccggcttc atcgactgcg 960

gccgcctggg cgtggccgac cgctaccagg acatcgccct ggccacccgc gacatcgccg 1020

aggagctggg cggcgagtgg gccgaccgct tcctggtgct gtacggcatc gccgcccccg 1080

actcccagcg catcgccttc taccgcctgc tggacgagtt cttctgactc gaggcagcag 1140

cagctcggat agtatcgaca cactctggac gctggtcgtg tgatggactg ttgccgccac 1200

acttgctgcc ttgacctgtg aatatccctg ccgcttttat caaacagcct cagtgtgttt 1260

gatcttgtgt gtacgcgctt ttgcgagttg ctagctgctt gtgctatttg cgaataccac 1320

ccccagcatc cccttccctc gtttcatatc gcttgcatcc caaccgcaac ttatctacgc 1380

tgtcctgcta tccctcagcg ctgctcctgc tcctgctcac tgcccctcgc acagccttgg 1440

tttgggctcc gcctgtattc tcctggtact gcaacctgta aaccagcact gcaatgctga 1500

tgcacgggaa gtagtgggat gggaacacaa atggaaagct tgagctc 1547

<210> 43
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic primer

<400> 43
gccgcgactg gctgctgctg g 21

<210> 44
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic primer

<400> 44
aggtcctcgc cgtcgggcat g 21

<210> 45
<211> 1292
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic polynucleotide

<400> 45
atcaaaggca tagattcaca tttgttggca ttgcagagca atcatcgcgc aggacgaaca 60

tcgctcacca agcacgtact gggcatccgg aggcctccgc aaattcctgc aacaggactc 120

gctgatcagt tcgcccaagg tctacgacgc tccctatcgg cgctagactt caacacatat 180

ttcactgtca cagcctcggc atgcatcagg cctcagtctc caccatgaag accatccagt 240

ctcggcacgc cggtcccatc ggacatgtgc agtcgggtcg ccgatcggcg gggcgcgcgg 300

gatcccgcat ggcgaccccc gtggccgcag ctaccgtcgc agcccctcgc tcggccctca 360

acctctcccc caccatcatt cgacaggagg tgctccactc cgccagcgcc cagcaactag 420

actgcgtggc ctccctggcg cccgtcttcg agtcccagat cctccccctc ctgacgcccg 480

tggacgagat gtggcagccc accgacttcc tccccgcctc gaactcggag gcattcttcg 540

accagatcgg cgacctgcgg gcgcgatcgg cggccatccc cgacgacctg ctggtctgcc 600

tggtggggga catgatcacg gaggaggccc tgcccaccta catggccatg ctgaacaccc 660

tggacgtcgt gcgcgatgag acagggcaca gccagcaccc ctacgccaag tggaccaggg 720

cttggatcgc ggaggagaac cgccatggcg acctgctgaa caagtacatg tggctgacgg 780

ggcgggtggg acatgctggc ggtggagcgc accatccagc catgctggcg gtggagcgca 840

ccatccagcg cctcatctca tcgggcatgg acccgggcac ggagaaccac ccctaccacg 900

cctttgtgtt caccagcttc caggagcgcg ccaccaagct gagccacggc tccaccgccc 960

gcctggcggt cgccgccggg gacgaggccc tggccaagat ctgcgggacc attgcgcggg 1020

acgagtcgcg ccacgaggcg gcgtacacgc ggaccatgga tgccatcttc cagcgcgacc 1080

ccagcggggc catggtggcg tttgcgcaca tgatgatgcg caagatcacc atgcccgccc 1140

acctcatgga cgacggccag cacggcgcgc gcaacggggg ggcgcaactt gttcgacgac 1200

tttgcggcag tggcggagcg ggcaggggtg tacaccgccg gcgactacat cggcatcctg 1260

cgccacctca tccggcgctg ggacgtggag gg 1292

<210> 46
<211> 364
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic polypeptide

<400> 46

Met His Gln Ala Ser Val Ser Thr Met Lys Thr Ile Gln Ser Arg His
1 5 10 15

Ala Gly Pro Ile Gly His Val Gln Ser Gly Arg Arg Ser Ala Gly Arg
20 25 30

Ala Gly Ser Arg Met Ala Thr Pro Val Ala Ala Ala Thr Val Ala Ala
35 40 45

Pro Arg Ser Ala Leu Asn Leu Ser Pro Thr Ile Ile Arg Gln Glu Val
50 55 60

Leu His Ser Ala Ser Ala Gln Gln Leu Asp Cys Val Ala Ser Leu Ala
65 70 75 80

Pro Val Phe Glu Ser Gln Ile Leu Pro Leu Leu Thr Pro Val Asp Glu
85 90 95

Met Trp Gln Pro Thr Asp Phe Leu Pro Ala Ser Asn Ser Glu Ala Phe
100 105 110

Phe Asp Gln Ile Gly Asp Leu Arg Ala Arg Ser Ala Ala Ile Pro Asp

115 120 125

Asp Leu Leu Val Cys Leu Val Gly Asp Met Ile Thr Glu Glu Ala Leu
130 135 140

Pro Thr Tyr Met Ala Met Leu Asn Thr Leu Asp Val Val Arg Asp Glu
145 150 155 160

Thr Gly His Ser Gln His Pro Tyr Ala Lys Trp Thr Arg Ala Trp Ile
165 170 175

Ala Glu Glu Asn Arg His Gly Asp Leu Leu Asn Lys Tyr Met Trp Leu
180 185 190

Thr Gly Arg Val Gly His Ala Gly Gly Gly Ala His His Pro Ala Met
195 200 205

Leu Ala Val Glu Arg Thr Ile Gln Arg Leu Ile Ser Ser Gly Met Asp
210 215 220

Pro Gly Thr Glu Asn His Pro Tyr His Ala Phe Val Phe Thr Ser Phe
225 230 235 240

Gln Glu Arg Ala Thr Lys Leu Ser His Gly Ser Thr Ala Arg Leu Ala
245 250 255

Val Ala Ala Gly Asp Glu Ala Leu Ala Lys Ile Cys Gly Thr Ile Ala
260 265 270

Arg Asp Glu Ser Arg His Glu Ala Ala Tyr Thr Arg Thr Met Asp Ala
275 280 285

Ile Phe Gln Arg Asp Pro Ser Gly Ala Met Val Ala Phe Ala His Met
290 295 300

Met Met Arg Lys Ile Thr Met Pro Ala His Leu Met Asp Asp Gly Gln
305 310 315 320

His Gly Ala Arg Asn Gly Gly Ala Gln Leu Val Arg Arg Leu Cys Gly
325 330 335

Ser Gly Gly Ala Gly Arg Gly Val His Arg Arg Arg Leu His Arg His
340 345 350

Pro Ala Pro Pro His Pro Ala Leu Gly Arg Gly Gly
355 360

<210> 47
<211> 1395
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic polynucleotide

<400> 47
attatacatc ggcatcgtct caggtttcac gatctgcatg ctatctatgg gactgtgact 60

ccgccggcca ggttgtggtg cgcgagaatc ctccccgctc ctgccttctc atttccctga 120

cgggagtcgc cgctgagcac cgggcggatc atgggcgtcg gcacactcca aaccccatat 180

acatgtggtc gtgcattcac gcatagcgca cggtatgtcc cgcgacgcgc ggctcgaagc 240

cgtggccatc cgacgcgctg cacggccgag gtgagggcac gcccctccgc caatggcgcg 300

cagcccatga ccgccttcga cttccggcag tacatgcagc agcgcgccgc gctggtggac 360

gcagcgctgg acctggcagt gccgctgcag taccccgaga agatcaacga ggccatgcgg 420

tacagcctgc tggccggggg caagcgcgtg cgccccgcgc tctgcctcgc tgcctgcgag 480

ctcgtgggcg gcccccctgga ggcggccatg cccgccgcct gcgccatgga gatgatccac 540

accatgagcc tcatccacga cgacctcccc gccatggaca acgacgactt ccggcgcggc 600

cagcccgcca accacaaggc ctatggcgag gagattgcga tcctggcggg cgacgcgctg 660

ctgtcgctga gctttgagca catcgcgcgc gagacgcgag gcgtggaccc ggtgcgcgtc 720

ctggccgcca tctcggagtg gcgcgcggtg ggcagccgcg ggctggtggc ggggcaggtg 780

gtggacctgg gtttcgaggg cggcggcgtg gggctggccc cgctgcgcta catccacgag 840

cacaaaaccg cggcgctgct ggaggcggcg gtggtgtccg gcgcgctgct gggcggcgcg 900

gaggaggcgg acctggagcg cctgcgcacc tacaaccgcg ccatcggcct cgctttccag 960

gtggtggggg acatcctgga catcccgggg accagcgagg agctgggcaa gaccgcgggc 1020

aaggacctga gctcccccaa aacccccctac ccgtccctgg tggggctggc caggtccaaa 1080

aaaattgcgg acgaactgat tgaggacgcg aaaacccaac tcacccagta cgagccggcc 1140

cgagcggcgc ccctcgtaac cctggccgaa aacatttgaa accggaagaa ctgactgggg 1200

gccccccctg cccccagata cggcggggct cctccatcca gttttgggat gggaggagcg 1260

acaaccgacc ccgtaaccct gtgacgcgtt tgccttgcat acgtacgcat gccttgaaac 1320

ccatccatga ccctcaacaa tacctggttg tgtgtagctt ggtcctgaaa aaaaaaaaaa 1380

aaaaaaaaaa aaaaa 1395

<210> 48
<211> 342
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic polypeptide

<400> 48
Met Gly Val Gly Thr Leu Gln Thr Pro Tyr Thr Cys Gly Arg Ala Phe
1 5 10 15

```
Thr His Ser Ala Arg Tyr Val Pro Arg Arg Ala Ala Arg Ser Arg Gly
            20                  25                  30

His Pro Thr Arg Cys Thr Ala Glu Val Arg Ala Arg Pro Ser Ala Asn
        35                  40                  45

Gly Ala Gln Pro Met Thr Ala Phe Asp Phe Arg Gln Tyr Met Gln Gln
    50                  55                  60

Arg Ala Ala Leu Val Asp Ala Ala Leu Asp Leu Ala Val Pro Leu Gln
65                  70                  75                  80

Tyr Pro Glu Lys Ile Asn Glu Ala Met Arg Tyr Ser Leu Leu Ala Gly
                85                  90                  95

Gly Lys Arg Val Arg Pro Ala Leu Cys Leu Ala Ala Cys Glu Leu Val
            100                 105                 110

Gly Gly Pro Leu Glu Ala Ala Met Pro Ala Ala Cys Ala Met Glu Met
        115                 120                 125

Ile His Thr Met Ser Leu Ile His Asp Asp Leu Pro Ala Met Asp Asn
    130                 135                 140

Asp Asp Phe Arg Arg Gly Gln Pro Ala Asn His Lys Ala Tyr Gly Glu
145                 150                 155                 160

Glu Ile Ala Ile Leu Ala Gly Asp Ala Leu Leu Ser Leu Ser Phe Glu
                165                 170                 175

His Ile Ala Arg Glu Thr Arg Gly Val Asp Pro Val Arg Val Leu Ala
            180                 185                 190

Ala Ile Ser Glu Trp Arg Ala Val Gly Ser Arg Gly Leu Val Ala Gly
        195                 200                 205

Gln Val Val Asp Leu Gly Phe Glu Gly Gly Gly Val Gly Leu Ala Pro
    210                 215                 220

Leu Arg Tyr Ile His Glu His Lys Thr Ala Ala Leu Leu Glu Ala Ala
225                 230                 235                 240

Val Val Ser Gly Ala Leu Leu Gly Gly Ala Glu Glu Ala Asp Leu Glu
                245                 250                 255

Arg Leu Arg Thr Tyr Asn Arg Ala Ile Gly Leu Ala Phe Gln Val Val
            260                 265                 270

Gly Asp Ile Leu Asp Ile Pro Gly Thr Ser Glu Glu Leu Gly Lys Thr
        275                 280                 285
```

Ala Gly Lys Asp Leu Ser Ser Pro Lys Thr Pro Tyr Pro Ser Leu Val
290 295 300

Gly Leu Ala Arg Ser Lys Lys Ile Ala Asp Glu Leu Ile Glu Asp Ala
305 310 315 320

Lys Thr Gln Leu Thr Gln Tyr Glu Pro Ala Arg Ala Ala Pro Leu Val
325 330 335

Thr Leu Ala Glu Asn Ile
340

<210> 49
<211> 833
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic polynucleotide

<400> 49
cagatgccat gcgccctcgg gccgcgggcc tgagggtcca cgcagcgtcc tcggtggccc 60

agacgcacca ggccgccccc ccggcggaca ggaggttcga cgactaccag ccccgcaccg 120

ccatcctctt ccccggccag ggcgcgcaga gcgtgggcat ggcgggagag ctggcgaagg 180

ccgtccccgc cgccgcggcg ctgttcgacg ccgcctccga ccagctcggc tatgacctgc 240

tccgcgtgtg cgttgagggc cccaaggcgc gcctggacag caccgccgtc agccagcccg 300

ccatctacgt ggccagcctg gcggcggtgg agaagctgcg cgcggagggc ggggaggagg 360

cactggccgc catcgacgtc gctgccggtc tgtccttggg cgagtacacc gcgctggcct 420

ttgccggcgc cttctccttc gccgacgggc tgcgcctggt ggccctgcgc ggcgccagca 480

tgcaggccgc cgccgacgcc gcaccctcgg gcatggtctc cgtcatcggt ctgccctccg 540

acgcggtggc cgcgctgtgc gaggccgcca acgcgcaggt ggcccccgac caggccgtgc 600

gcatcgccaa ctacctctgc gacggcaact acgccgtcag cggtgggctg gagggctgcg 660

cggcggtgga gggcctggcc aaggcccaca aggcgcgcat gacggtgcgc ctggcggtgg 720

cgggcgcctt ccacaccccc ttcatgcagc cggcggtgga ggcgctgagc gcgggcgctg 780

gcggacacgc cgctggtcgc gccgcgcatc cccgtggtca gcaacgggac gcc 833

<210> 50
<211> 275
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic polypeptide

<400> 50
Met Arg Pro Arg Ala Ala Gly Leu Arg Val His Ala Ala Ser Ser Val
1 5 10 15

```
Ala Gln Thr His Gln Ala Ala Pro Pro Ala Asp Arg Arg Phe Asp Asp
            20                  25                  30

Tyr Gln Pro Arg Thr Ala Ile Leu Phe Pro Gly Gln Gly Ala Gln Ser
        35                  40                  45

Val Gly Met Ala Gly Glu Leu Ala Lys Ala Val Pro Ala Ala Ala Ala
    50                  55                  60

Leu Phe Asp Ala Ala Ser Asp Gln Leu Gly Tyr Asp Leu Leu Arg Val
65                  70                  75                  80

Cys Val Glu Gly Pro Lys Ala Arg Leu Asp Ser Thr Ala Val Ser Gln
                85                  90                  95

Pro Ala Ile Tyr Val Ala Ser Leu Ala Ala Val Glu Lys Leu Arg Ala
            100                 105                 110

Glu Gly Gly Glu Glu Ala Leu Ala Ala Ile Asp Val Ala Ala Gly Leu
        115                 120                 125

Ser Leu Gly Glu Tyr Thr Ala Leu Ala Phe Ala Gly Ala Phe Ser Phe
    130                 135                 140

Ala Asp Gly Leu Arg Leu Val Ala Leu Arg Gly Ala Ser Met Gln Ala
145                 150                 155                 160

Ala Ala Asp Ala Ala Pro Ser Gly Met Val Ser Val Ile Gly Leu Pro
                165                 170                 175

Ser Asp Ala Val Ala Ala Leu Cys Glu Ala Ala Asn Ala Gln Val Ala
            180                 185                 190

Pro Asp Gln Ala Val Arg Ile Ala Asn Tyr Leu Cys Asp Gly Asn Tyr
        195                 200                 205

Ala Val Ser Gly Gly Leu Glu Gly Cys Ala Ala Val Glu Gly Leu Ala
    210                 215                 220

Lys Ala His Lys Ala Arg Met Thr Val Arg Leu Ala Val Ala Gly Ala
225                 230                 235                 240

Phe His Thr Pro Phe Met Gln Pro Ala Val Glu Ala Leu Ser Ala Gly
                245                 250                 255

Ala Gly Gly His Ala Ala Gly Arg Ala Ala His Pro Arg Gly Gln Gln
            260                 265                 270

Arg Asp Ala
        275
```

<210> 51
<211> 787
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic polynucleotide

<400> 51
tgtccatctc cccccaccct ccatccaacc atcgtcgacg gcatgcaggc gctgtgttct 60

caccccgcgt ccctcacggc gcgtgcggta ccccatgggc gggccagccc agcacagcgg 120

gtgtccagcg ccggcccggc ctacaccggc ctgtcccggc acaccctggg ctgccccagc 180

acccccaccc tccagtcccg cgccgcggtc cagacccgcg gctcctcctc cggctccacc 240

acgcgcatga ccaccaccgc ccagcgcaag atcaaggtgg ccatcaacgg gttcggccgc 300

atcggccgcc agttcctgcg ctgcgtggag gggcgcgagg actcgctgct ggagatcgtg 360

gccgtgaacg actccggcgg cgtgaagcag gccagccacc tgctcaagta cgactccacc 420

atgggcacct tcaacgccga catcaagatc tcgggcgagg gcaccttctc cgtcaacggc 480

cgcgacatcc gcgtcgtctc ctcccgcgac cccctggccc tgccctgggg cgagctgggc 540

gtggacctgg tgatcgaggg gacgggagtg tttgtggacc gcaagggtgc cagcaagcac 600

ctgcaggcgg gggccaagaa ggtcatcatc accgcgccgg ccaagggctc cgacgtgccc 660

acctacgtca tgggcgtgaa cgcggaccag tactccaact ccgacgacat catctccaac 720

gcctcctgca ccaccaactg cctggcgccc tttgtcaagg tgctcaacga ccgcttcggc 780

atcgtga 787

<210> 52
<211> 248
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic polypeptide

<400> 52
Met Gln Ala Leu Cys Ser His Pro Ala Ser Leu Thr Ala Arg Ala Val
1 5 10 15

Pro His Gly Arg Ala Ser Pro Ala Gln Arg Val Ser Ser Ala Gly Pro
20 25 30

Ala Tyr Thr Gly Leu Ser Arg His Thr Leu Gly Cys Pro Ser Thr Pro
35 40 45

Thr Leu Gln Ser Arg Ala Ala Val Gln Thr Arg Gly Ser Ser Ser Gly
50 55 60

Ser Thr Thr Arg Met Thr Thr Thr Ala Gln Arg Lys Ile Lys Val Ala
65 70 75 80

Ile Asn Gly Phe Gly Arg Ile Gly Arg Gln Phe Leu Arg Cys Val Glu
85 90 95

Gly Arg Glu Asp Ser Leu Leu Glu Ile Val Ala Val Asn Asp Ser Gly
100 105 110

Gly Val Lys Gln Ala Ser His Leu Leu Lys Tyr Asp Ser Thr Met Gly
115 120 125

Thr Phe Asn Ala Asp Ile Lys Ile Ser Gly Glu Gly Thr Phe Ser Val
130 135 140

Asn Gly Arg Asp Ile Arg Val Val Ser Ser Arg Asp Pro Leu Ala Leu
145 150 155 160

Pro Trp Gly Glu Leu Gly Val Asp Leu Val Ile Glu Gly Thr Gly Val
165 170 175

Phe Val Asp Arg Lys Gly Ala Ser Lys His Leu Gln Ala Gly Ala Lys
180 185 190

Lys Val Ile Ile Thr Ala Pro Ala Lys Gly Ser Asp Val Pro Thr Tyr
195 200 205

Val Met Gly Val Asn Ala Asp Gln Tyr Ser Asn Ser Asp Asp Ile Ile
210 215 220

Ser Asn Ala Ser Cys Thr Thr Asn Cys Leu Ala Pro Phe Val Lys Val
225 230 235 240

Leu Asn Asp Arg Phe Gly Ile Val
245

<210> 53
<211> 860
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic polynucleotide

<400> 53
gatgttgaga atagtagctt gctgccttgt cgccatgcag agcgtgtgcg cgcagtcggt 60

ttcatgcaag ggggccttca cccagtccct gcggaccccc cgatgcagca ggagccagct 120

cgtctgccgg gctgatggca aggccggagc cttcatcaag accgtaaaga gcggtgctgc 180

cgctctggct gcctccctcc tcctgtctgg gggtgcgggc gcactgacct ttgatgagct 240

gcagggcctg acctacctgc aggtgaaggg ctctggcatc gccaacacct gccccaccct 300

gtctggcggc tcctccaaca tcaaggacct gaagagcggg acctactccg tcaacaagat 360

gtgcctggag cccacgtcct tcaaggtcaa ggaggaggca cagttcaaga acggcgaggc 420

cgactttgtg cccaccaagc tcgtcacgcg tctgacctac accctggacg agatctctgg 480

ccagatgaag atcgacggca gcggcggcgt ggagttcaag gaggaggatg gcatcgacta 540

tgctgcagtc accgtgcagc ttccgggcgg ggagcgcgtg cccttcctct tcaccatcaa 600

ggagcttgac gccaagggga ctgccgacgg cttcaagggc gagttcaccg tgccctccta 660

ccgtgggtcc tccttcctgg accccaaggg ccgcggcgcc tccaccggct acgacaacgc 720

cgtggccctg cccgccgccg gcgattccga ggagttggag aaggagaaca acaagtccac 780

caaggctctg aaggggagg ccatcttctc catcgccaag gtggacgccg ggacaggggá 840

ggtggcgggc atctttgagt 860

<210> 54
<211> 275
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic polypeptide

<400> 54

Met Gln Ser Val Cys Ala Gln Ser Val Ser Cys Lys Gly Ala Phe Thr
1 5 10 15

Gln Ser Leu Arg Thr Pro Arg Cys Ser Arg Ser Gln Leu Val Cys Arg
20 25 30

Ala Asp Gly Lys Ala Gly Ala Phe Ile Lys Thr Val Lys Ser Gly Ala
35 40 45

Ala Ala Leu Ala Ala Ser Leu Leu Leu Ser Gly Gly Ala Gly Ala Leu
50 55 60

Thr Phe Asp Glu Leu Gln Gly Leu Thr Tyr Leu Gln Val Lys Gly Ser
65 70 75 80

Gly Ile Ala Asn Thr Cys Pro Thr Leu Ser Gly Gly Ser Ser Asn Ile
85 90 95

Lys Asp Leu Lys Ser Gly Thr Tyr Ser Val Asn Lys Met Cys Leu Glu
100 105 110

Pro Thr Ser Phe Lys Val Lys Glu Glu Ala Gln Phe Lys Asn Gly Glu
115 120 125

Ala Asp Phe Val Pro Thr Lys Leu Val Thr Arg Leu Thr Tyr Thr Leu
130 135 140

Asp Glu Ile Ser Gly Gln Met Lys Ile Asp Gly Ser Gly Gly Val Glu
145 150 155 160

Phe Lys Glu Glu Asp Gly Ile Asp Tyr Ala Ala Val Thr Val Gln Leu
165 170 175

Pro Gly Gly Glu Arg Val Pro Phe Leu Phe Thr Ile Lys Glu Leu Asp
180 185 190

Ala Lys Gly Thr Ala Asp Gly Phe Lys Gly Glu Phe Thr Val Pro Ser
195 200 205

Tyr Arg Gly Ser Ser Phe Leu Asp Pro Lys Gly Arg Gly Ala Ser Thr
210 215 220

Gly Tyr Asp Asn Ala Val Ala Leu Pro Ala Ala Gly Asp Ser Glu Glu
225 230 235 240

Leu Glu Lys Glu Asn Asn Lys Ser Thr Lys Ala Leu Lys Gly Glu Ala
245 250 255

Ile Phe Ser Ile Ala Lys Val Asp Ala Gly Thr Gly Glu Val Ala Gly
260 265 270

Ile Phe Glu
275

<210> 55
<211> 818
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic polynucleotide

<400> 55
ataatcggaa cccagctgca cgcaccatca gtgcggcagc atgcagaccg tcgcagccag 60

ctatggcgta ttggcgccct ccggctccag cgtgacccgg ggctcgacca gcagcaagca 120

gcacttcacc accctcactc ccttttccgg cttcaggcgc ctgaatcatg tggatcgggc 180

ggggcaggcg gggtctggga gcccccagac cctgcagcag gccgtgggca aggccgtgcg 240

ccggtcgcgg ggccgcacca ccagcgccgt gcgcgtgacc cgcatgatgt ttgagcggtt 300

caccgagaag gccatcaagg tggtcatgct cgcgcaggag gaggctcgcc gtctgggcca 360

caacttcgtg gggacggagc aaatcctgct ggggttgatt ggggagtcca caggcatcgc 420

cgccaaggtc ctcaagtcga tgggcgtcac gctgaaagat gcgcgtgtgg aggtcgagaa 480

gatcatcggc cgggggagcg gctttgtggc cgtggagatc cccttcaccc cccgcgccaa 540

gcgtgtgctg gagctgtccc tggaggaggc tcgccagctc ggccacaact acattggcac 600

ggagcacatc ctgctgggcc tgctgcgcga gggtgagggc gtggcctccc gcgtgctgga 660

gaccctgggc gccgaccccc agaagatccg cactcaggtg gtacgcatgg tgggtgagtc 720

gcaggagccc gtgggcacca cggtgggcgg agggtccacc ggctccaaca agatgcccac 780

cctggaggag tacggcacca acctgaccgc ccaggccg 818

<210> 56
<211> 259
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic polypeptide

<400> 56

```
Met Gln Thr Val Ala Ala Ser Tyr Gly Val Leu Ala Pro Ser Gly Ser
1               5                  10                  15

Ser Val Thr Arg Gly Ser Thr Ser Ser Lys Gln His Phe Thr Thr Leu
            20                  25                  30

Thr Pro Phe Ser Gly Phe Arg Arg Leu Asn His Val Asp Arg Ala Gly
        35                  40                  45

Gln Ala Gly Ser Gly Ser Pro Gln Thr Leu Gln Gln Ala Val Gly Lys
    50                  55                  60

Ala Val Arg Arg Ser Arg Gly Arg Thr Thr Ser Ala Val Arg Val Thr
65                  70                  75                  80

Arg Met Met Phe Glu Arg Phe Thr Glu Lys Ala Ile Lys Val Val Met
                85                  90                  95

Leu Ala Gln Glu Glu Ala Arg Arg Leu Gly His Asn Phe Val Gly Thr
            100                 105                 110

Glu Gln Ile Leu Leu Gly Leu Ile Gly Glu Ser Thr Gly Ile Ala Ala
        115                 120                 125

Lys Val Leu Lys Ser Met Gly Val Thr Leu Lys Asp Ala Arg Val Glu
    130                 135                 140

Val Glu Lys Ile Ile Gly Arg Gly Ser Gly Phe Val Ala Val Glu Ile
145                 150                 155                 160

Pro Phe Thr Pro Arg Ala Lys Arg Val Leu Glu Leu Ser Leu Glu Glu
                165                 170                 175

Ala Arg Gln Leu Gly His Asn Tyr Ile Gly Thr Glu His Ile Leu Leu
            180                 185                 190

Gly Leu Leu Arg Glu Gly Glu Gly Val Ala Ser Arg Val Leu Glu Thr
        195                 200                 205

Leu Gly Ala Asp Pro Gln Lys Ile Arg Thr Gln Val Val Arg Met Val
    210                 215                 220
```

Gly Glu Ser Gln Glu Pro Val Gly Thr Thr Val Gly Gly Gly Ser Thr
225 230 235 240

Gly Ser Asn Lys Met Pro Thr Leu Glu Glu Tyr Gly Thr Asn Leu Thr
245 250 255

Ala Gln Ala

<210> 57
<211> 2357
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic polynucleotide

<400> 57
ctttcttgcg ctatgacact tccagcaaaa ggtagggcgg gctgcgagac ggcttcccgg 60

cgctgcatgc aacaccgatg atgcttcgac cccccgaagc tccttcgggg ctgcatgggc 120

gctccgatgc cgctccaggg cgagcgctgt ttaaatagcc aggcccccga ttgcaaagac 180

attatagcga gctaccaaag ccatattcaa acacctagat cactaccact tctacacagg 240

ccactcgagc ttgtgatcgc actccgctaa gggggcgcct cttcctcttc gtttcagtca 300

caacccgcaa acggcgcgcc atatcaatgc ttcttcaggc ctttcttttt cttcttgctg 360

gttttgctgc caagatcagc gcctctatga cgaacgaaac ctcggataga ccacttgtgc 420

actttacacc aaacaagggc tggatgaatg accccaatgg actgtggtac gacgaaaaag 480

atgccaagtg gcatctgtac tttcaataca acccgaacga tactgtctgg gggacgccat 540

tgttttgggg ccacgccacg tccgacgacc tgaccaattg ggaggaccaa ccaatagcta 600

tcgctccgaa gaggaacgac tccggagcat tctcgggttc catggtggtt gactacaaca 660

atacttccgg ctttttcaac gataccattg acccgagaca acgctgcgtg gccatatgga 720

cttacaacac accggagtcc gaggagcagt acatctcgta tagcctggac ggtggataca 780

cttttacaga gtatcagaag aaccctgtgc ttgctgcaaa ttcgactcag ttccgagatc 840

cgaaggtctt ttggtacgag ccctcgcaga agtggatcat gacagcggca aagtcacagg 900

actacaagat cgaaatttac tcgtctgacg accttaaatc ctggaagctc gaatccgcgt 960

tcgcaaacga gggctttctc ggctaccaat acgaatgccc aggcctgata gaggtcccaa 1020

cagagcaaga tcccagcaag tcctactggg tgatgtttat ttccattaat ccaggagcac 1080

cggcaggagg ttcttttaat cagtacttcg tcggaagctt taacggaact catttcgagg 1140

catttgataa ccaatcaaga gtagttgatt ttggaaagga ctactatgcc ctgcagactt 1200

tcttcaatac tgacccgacc tatgggagcg ctcttggcat tgcgtgggct tctaactggg 1260

agtattccgc attcgttcct acaaaccctt ggaggtcctc catgtcgctc gtgaggaaat 1320

tctctctcaa cactgagtac caggccaacc cggaaaccga actcataaac ctgaaagccg 1380

aaccgatcct gaacattagc aacgctggcc cctggagccg gtttgcaacc aacaccacgt 1440

tgacgaaagc caacagctac aacgtcgatc tttcgaatag caccggtaca cttgaatttg 1500

aactggtgta tgccgtcaat accacccaaa cgatctcgaa gtcggtgttc gcggacctct 1560

ccctctggtt taaaggcctg gaagaccccg aggagtacct cagaatgggt ttcgaggttt 1620

ctgcgtcctc cttcttcctt gatcgcggga acagcaaagt aaaatttgtt aaggagaacc 1680

catattttac caacaggatg agcgttaaca accaaccatt caagagcgaa aacgacctgt 1740

cgtactacaa agtgtatggt ttgcttgatc aaaatatcct ggaactctac ttcaacgatg 1800

gtgatgtcgt gtccaccaac acatacttca tgacaaccgg gaacgcactg ggctccgtga 1860

acatgacgac gggtgtggat aacctgttct acatcgacaa attccaggtg agggaagtca 1920

agtgagatct gtcgatcgac aagctcgagg cagcagcagc tcggatagta tcgacacact 1980

ctggacgctg gtcgtgtgat ggactgttgc cgccacactt gctgccttga cctgtgaata 2040

tccctgccgc ttttatcaaa cagcctcagt gtgtttgatc ttgtgtgtac gcgcttttgc 2100

gagttgctag ctgcttgtgc tatttgcgaa taccaccccc agcatccccc tccctcgttt 2160

catatcgctt gcatcccaac cgcaacttat ctacgctgtc ctgctatccc tcagcgctgc 2220

tcctgctcct gctcactgcc cctcgcacag ccttggtttg ggctccgcct gtattctcct 2280

ggtactgcaa cctgtaaacc agcactgcaa tgctgatgca cgggaagtag tgggatggga 2340

acacaaatgg aaagctt 2357

<210> 58
<211> 2335
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic polynucleotide

<400> 58
ctttcttgcg ctatgacact tccagcaaaa ggtagggcgg gctgcgagac ggcttcccgg 60

cgctgcatgc aacaccgatg atgcttcgac cccccgaagc tccttcgggg ctgcatgggc 120

gctccgatgc cgctccaggg cgagcgctgt ttaaatagcc aggcccccga ttgcaaagac 180

attatagcga gctaccaaag ccatattcaa acacctagat cactaccact tctacacagg 240

ccactcgagc ttgtgatcgc actccgctaa gggggcgcct cttcctcttc gtttcagtca 300

caacccgcaa acggcgcgcc atgctgctgc aggccttcct gttcctgctg gccggcttcg 360

ccgccaagat cagcgcctcc atgacgaacg agacgtccga ccgccccctg gtgcacttca 420

cccccaacaa gggctggatg aacgacccca acggcctgtg gtacgacgag aaggacgcca 480

agtggcacct gtacttccag tacaacccga acgacaccgt ctgggggacg cccttgttct 540

ggggccacgc cacgtccgac gacctgacca actgggagga ccagcccatc gccatcgccc 600

cgaagcgcaa cgactccggc gccttctccg gctccatggt ggtggactac aacaacacct 660

ccggcttctt caacgacacc atcgacccgc gccagcgctg cgtggccatc tggacctaca 720

acaccccgga gtccgaggag cagtacatct cctacagcct ggacggcggc tacaccttca 780

ccgagtacca gaagaacccc gtgctggccg ccaactccac ccagttccgc gacccgaagg 840

tcttctggta cgagccctcc cagaagtgga tcatgaccgc ggccaagtcc caggactaca 900

agatcgagat ctactcctcc gacgacctga agtcctggaa gctggagtcc gcgttcgcca 960

acgagggctt cctcggctac cagtacgagt gccccggcct gatcgaggtc cccaccgagc 1020

aggaccccag caagtcctac tgggtgatgt tcatctccat caaccccggc gccccggccg 1080

gcggctcctt caaccagtac ttcgtcggca gcttcaacgg cacccacttc gaggccttcg 1140

acaaccagtc ccgcgtggtg gacttcggca aggactacta cgccctgcag accttcttca 1200

acaccgaccc gacctacggg agcgccctgg gcatcgcgtg ggcctccaac tgggagtact 1260

ccgccttcgt gcccaccaac ccctggcgct cctccatgtc cctcgtgcgc aagttctccc 1320

tcaacaccga gtaccaggcc aacccggaga cggagctgat caacctgaag gccgagccga 1380

tcctgaacat cagcaacgcc ggcccctgga gccggttcgc caccaacacc acgttgacga 1440

aggccaacag ctacaacgtc gacctgtcca acagcaccgg caccctggag ttcgagctgg 1500

tgtacgccgt caacaccacc cagacgatct ccaagtccgt gttcgcggac ctctccctct 1560

ggttcaaggg cctggaggac cccgaggagt acctccgcat gggcttcgag gtgtccgcgt 1620

cctccttctt cctggaccgc gggaacagca aggtgaagtt cgtgaaggag aacccctact 1680

tcaccaaccg catgagcgtg aacaaccagc ccttcaagag cgagaacgac ctgtcctact 1740

acaaggtgta cggcttgctg gaccagaaca tcctggagct gtacttcaac gacggcgacg 1800

tcgtgtccac caacacctac ttcatgacca ccgggaacgc cctgggctcc gtgaacatga 1860

cgacgggggt ggacaacctg ttctacatcg acaagttcca ggtgcgcgag gtcaagtgat 1920

taattaactc gaggcagcag cagctcggat agtatcgaca cactctggac gctggtcgtg 1980

tgatggactg ttgccgccac acttgctgcc ttgacctgtg aatatccctg ccgcttttat 2040

caaacagcct cagtgtgttt gatcttgtgt gtacgcgctt ttgcgagttg ctagctgctt 2100

gtgctatttg cgaataccac ccccagcatc cccttccctc gtttcatatc gcttgcatcc 2160

caaccgcaac ttatctacgc tgtcctgcta tccctcagcg ctgctcctgc tcctgctcac 2220

tgcccctcgc acagccttgg tttgggctcc gcctgtattc tcctggtact gcaacctgta 2280

aaccagcact gcaatgctga tgcacgggaa gtagtgggat gggaacacaa atgga 2335

<210> 59
<211> 382
<212> PRT
<213> Cinnamomum camphorum

<400> 59
Met Ala Thr Thr Ser Leu Ala Ser Ala Phe Cys Ser Met Lys Ala Val
1 5 10 15

Met Leu Ala Arg Asp Gly Arg Gly Met Lys Pro Arg Ser Ser Asp Leu
20 25 30

Gln Leu Arg Ala Gly Asn Ala Gln Thr Ser Leu Lys Met Ile Asn Gly
35 40 45

Thr Lys Phe Ser Tyr Thr Glu Ser Leu Lys Lys Leu Pro Asp Trp Ser
50 55 60

Met Leu Phe Ala Val Ile Thr Thr Ile Phe Ser Ala Ala Glu Lys Gln
65 70 75 80

Trp Thr Asn Leu Glu Trp Lys Pro Lys Pro Asn Pro Pro Gln Leu Leu
85 90 95

Asp Asp His Phe Gly Pro His Gly Leu Val Phe Arg Arg Thr Phe Ala
100 105 110

Ile Arg Ser Tyr Glu Val Gly Pro Asp Arg Ser Thr Ser Ile Val Ala
115 120 125

Val Met Asn His Leu Gln Glu Ala Ala Leu Asn His Ala Lys Ser Val
130 135 140

Gly Ile Leu Gly Asp Gly Phe Gly Thr Thr Leu Glu Met Ser Lys Arg
145 150 155 160

Asp Leu Ile Trp Val Val Lys Arg Thr His Val Ala Val Glu Arg Tyr
165 170 175

Pro Ala Trp Gly Asp Thr Val Glu Val Glu Cys Trp Val Gly Ala Ser
180 185 190

Gly Asn Asn Gly Arg Arg His Asp Phe Leu Val Arg Asp Cys Lys Thr
195 200 205

Gly Glu Ile Leu Thr Arg Cys Thr Ser Leu Ser Val Met Met Asn Thr
210 215 220

Arg Thr Arg Arg Leu Ser Lys Ile Pro Glu Glu Val Arg Gly Glu Ile
225 230 235 240

Gly Pro Ala Phe Ile Asp Asn Val Ala Val Lys Asp Glu Glu Ile Lys
245 250 255

Lys Pro Gln Lys Leu Asn Asp Ser Thr Ala Asp Tyr Ile Gln Gly Gly
260 265 270

Leu Thr Pro Arg Trp Asn Asp Leu Asp Ile Asn Gln His Val Asn Asn
275 280 285

Ile Lys Tyr Val Asp Trp Ile Leu Glu Thr Val Pro Asp Ser Ile Phe
290 295 300

Glu Ser His His Ile Ser Ser Phe Thr Ile Glu Tyr Arg Arg Glu Cys
305 310 315 320

Thr Met Asp Ser Val Leu Gln Ser Leu Thr Thr Val Ser Gly Gly Ser
325 330 335

Ser Glu Ala Gly Leu Val Cys Glu His Leu Leu Gln Leu Glu Gly Gly
340 345 350

Ser Glu Val Leu Arg Ala Lys Thr Glu Trp Arg Pro Lys Leu Thr Asp
355 360 365

Ser Phe Arg Gly Ile Ser Val Ile Pro Ala Glu Ser Ser Val
370 375 380

<210> 60
<211> 1240
<212> DNA
<213> Cinnamomum camphorum

<400> 60
ggcgcgccat ggccaccacc tccctggcct ccgccttctg cagcatgaag gccgtgatgc 60

tggcccgcga cggccgcggc atgaagcccc gctccagcga cctgcagctg cgcgccggca 120

acgcccagac ctccctgaag atgatcaacg gcaccaagtt ctcctacacc gagagcctga 180

agaagctgcc cgactggtcc atgctgttcg ccgtgatcac caccatcttc tccgccgccg 240

agaagcagtg gaccaacctg gagtggaagc ccaagcccaa ccccccccag ctgctggacg 300

accacttcgg cccccacggc ctggtgttcc gccgcacctt cgccatccgc agctacgagg 360

tgggccccga ccgctccacc agcatcgtgg ccgtgatgaa ccacctgcag gaggccgccc 420

tgaaccacgc caagtccgtg ggcatcctgg gcgacggctt cggcaccacc ctggagatgt 480

ccaagcgcga cctgatctgg gtggtgaagc gcacccacgt ggccgtggag cgctaccccg 540

cctggggcga caccgtggag gtggagtgct gggtgggcgc ctccggcaac aacggccgcc 600

gccacgactt cctggtgcgc gactgcaaga ccggcgagat cctgacccgc tgcacctccc 660

tgagcgtgat gatgaacacc cgcacccgcc gcctgagcaa gatccccgag gaggtgcgcg 720

gcgagatcgg ccccgccttc atcgacaacg tggccgtgaa ggacgaggag atcaagaagc 780

cccagaagct gaacgactcc accgccgact acatccaggg cggcctgacc ccccgctgga 840

acgacctgga catcaaccag cacgtgaaca acatcaagta cgtggactgg atcctggaga 900

ccgtgcccga cagcatcttc gagagccacc acatctcctc cttcaccatc gagtaccgcc 960

gcgagtgcac catggacagc gtgctgcagt ccctgaccac cgtgagcggc ggctcctccg 1020

aggccggcct ggtgtgcgag cacctgctgc agctggaggg cggcagcgag gtgctgcgcg 1080

ccaagaccga gtggcgcccc aagctgaccg actccttccg cggcatcagc gtgatccccg 1140

ccgagtccag cgtgatggac tacaaggacc acgacggcga ctacaaggac cacgacatcg 1200

actacaagga cgacgacgac aagtgactcg agttaattaa 1240

<210> 61
<211> 415
<212> PRT
<213> Cuphea hookeriana

<400> 61

```
Met Val Ala Ala Ala Ala Ser Ser Ala Phe Phe Pro Val Pro Ala Pro
1               5                  10                  15

Gly Ala Ser Pro Lys Pro Gly Lys Phe Gly Asn Trp Pro Ser Ser Leu
            20                  25                  30

Ser Pro Ser Phe Lys Pro Lys Ser Ile Pro Asn Gly Gly Phe Gln Val
        35                  40                  45

Lys Ala Asn Asp Ser Ala His Pro Lys Ala Asn Gly Ser Ala Val Ser
    50                  55                  60

Leu Lys Ser Gly Ser Leu Asn Thr Gln Glu Asp Thr Ser Ser Ser Pro
65                  70                  75                  80

Pro Pro Arg Thr Phe Leu His Gln Leu Pro Asp Trp Ser Arg Leu Leu
                85                  90                  95

Thr Ala Ile Thr Thr Val Phe Val Lys Ser Lys Arg Pro Asp Met His
            100                 105                 110

Asp Arg Lys Ser Lys Arg Pro Asp Met Leu Val Asp Ser Phe Gly Leu
        115                 120                 125

Glu Ser Thr Val Gln Asp Gly Leu Val Phe Arg Gln Ser Phe Ser Ile
    130                 135                 140

Arg Ser Tyr Glu Ile Gly Thr Asp Arg Thr Ala Ser Ile Glu Thr Leu
145                 150                 155                 160

Met Asn His Leu Gln Glu Thr Ser Leu Asn His Cys Lys Ser Thr Gly
                165                 170                 175

Ile Leu Leu Asp Gly Phe Gly Arg Thr Leu Glu Met Cys Lys Arg Asp
            180                 185                 190

Leu Ile Trp Val Val Ile Lys Met Gln Ile Lys Val Asn Arg Tyr Pro
        195                 200                 205

Ala Trp Gly Asp Thr Val Glu Ile Asn Thr Arg Phe Ser Arg Leu Gly
    210                 215                 220

Lys Ile Gly Met Gly Arg Asp Trp Leu Ile Ser Asp Cys Asn Thr Gly
```

225 230 235 240

Glu Ile Leu Val Arg Ala Thr Ser Ala Tyr Ala Met Met Asn Gln Lys
245 250 255

Thr Arg Arg Leu Ser Lys Leu Pro Tyr Glu Val His Gln Glu Ile Val
260 265 270

Pro Leu Phe Val Asp Ser Pro Val Ile Glu Asp Ser Asp Leu Lys Val
275 280 285

His Lys Phe Lys Val Lys Thr Gly Asp Ser Ile Gln Lys Gly Leu Thr
290 295 300

Pro Gly Trp Asn Asp Leu Asp Val Asn Gln His Val Ser Asn Val Lys
305 310 315 320

Tyr Ile Gly Trp Ile Leu Glu Ser Met Pro Thr Glu Val Leu Glu Thr
325 330 335

Gln Glu Leu Cys Ser Leu Ala Leu Glu Tyr Arg Arg Glu Cys Gly Arg
340 345 350

Asp Ser Val Leu Glu Ser Val Thr Ala Met Asp Pro Ser Lys Val Gly
355 360 365

Val Arg Ser Gln Tyr Gln His Leu Leu Arg Leu Glu Asp Gly Thr Ala
370 375 380

Ile Val Asn Gly Ala Thr Glu Trp Arg Pro Lys Asn Ala Gly Ala Asn
385 390 395 400

Gly Ala Ile Ser Thr Gly Lys Thr Ser Asn Gly Asn Ser Val Ser
405 410 415

<210> 62
<211> 1339
<212> DNA
<213> Cuphea hookeriana

<400> 62
ggcgcgccat ggtggccgcc gccgcctcca gcgccttctt ccccgtgccc gcccccggcg 60

cctcccccaa gcccggcaag ttcggcaact ggccctccag cctgagcccc tccttcaagc 120

ccaagtccat ccccaacggc ggcttccagg tgaaggccaa cgacagcgcc caccccaagg 180

ccaacggctc cgccgtgagc ctgaagagcg gcagcctgaa cacccaggag gacacctcct 240

ccagcccccc cccccgcacc ttcctgcacc agctgcccga ctggagccgc ctgctgaccg 300

ccatcaccac cgtgttcgtg aagtccaagc gccccgacat gcacgaccgc aagtccaagc 360

gccccgacat gctggtggac agcttcggcc tggagtccac cgtgcaggac ggcctggtgt 420

tccgccagtc cttctccatc cgctcctacg agatcggcac cgaccgcacc gccagcatcg 480

agaccctgat gaaccacctg caggagacct ccctgaacca ctgcaagagc accggcatcc 540

tgctggacgg cttcggccgc accctggaga tgtgcaagcg cgacctgatc tgggtggtga 600

tcaagatgca gatcaaggtg aaccgctacc ccgcctgggg cgacaccgtg gagatcaaca 660

cccgcttcag ccgcctgggc aagatcggca tgggccgcga ctggctgatc tccgactgca 720

acaccggcga gatcctggtg cgcgccacca gcgcctacgc catgatgaac cagaagaccc 780

gccgcctgtc caagctgccc tacgaggtgc accaggagat cgtgcccctg ttcgtggaca 840

gccccgtgat cgaggactcc gacctgaagg tgcacaagtt caaggtgaag accggcgaca 900

gcatccagaa gggcctgacc cccggctgga acgacctgga cgtgaaccag cacgtgtcca 960

acgtgaagta catcggctgg atcctggaga gcatgcccac cgaggtgctg gagacccagg 1020

agctgtgctc cctggccctg gagtaccgcc gcgagtgcgg ccgcgactcc gtgctggaga 1080

gcgtgaccgc catggacccc agcaaggtgg gcgtgcgctc ccagtaccag cacctgctgc 1140

gcctggagga cggcaccgcc atcgtgaacg gcgccaccga gtggcgcccc aagaacgccg 1200

gcgccaacgg cgccatctcc accggcaaga ccagcaacgg caactccgtg tccatggact 1260

acaaggacca cgacggcgac tacaaggacc acgacatcga ctacaaggac gacgacgaca 1320

agtgactcga gttaattaa 1339

<210> 63
<211> 382
<212> PRT
<213> Umbellularia sp.

<400> 63

Met Ala Thr Thr Ser Leu Ala Ser Ala Phe Cys Ser Met Lys Ala Val
1 5 10 15

Met Leu Ala Arg Asp Gly Arg Gly Met Lys Pro Arg Ser Ser Asp Leu
20 25 30

Gln Leu Arg Ala Gly Asn Ala Pro Thr Ser Leu Lys Met Ile Asn Gly
35 40 45

Thr Lys Phe Ser Tyr Thr Glu Ser Leu Lys Arg Leu Pro Asp Trp Ser
50 55 60

Met Leu Phe Ala Val Ile Thr Thr Ile Phe Ser Ala Ala Glu Lys Gln
65 70 75 80

Trp Thr Asn Leu Glu Trp Lys Pro Lys Pro Lys Leu Pro Gln Leu Leu
85 90 95

Asp Asp His Phe Gly Leu His Gly Leu Val Phe Arg Arg Thr Phe Ala
100 105 110

Ile Arg Ser Tyr Glu Val Gly Pro Asp Arg Ser Thr Ser Ile Leu Ala
115 120 125

```
Val Met Asn His Met Gln Glu Ala Thr Leu Asn His Ala Lys Ser Val
    130                 135                 140

Gly Ile Leu Gly Asp Gly Phe Gly Thr Thr Leu Glu Met Ser Lys Arg
145                 150                 155                 160

Asp Leu Met Trp Val Val Arg Arg Thr His Val Ala Val Glu Arg Tyr
                165                 170                 175

Pro Thr Trp Gly Asp Thr Val Glu Val Glu Cys Trp Ile Gly Ala Ser
            180                 185                 190

Gly Asn Asn Gly Met Arg Arg Asp Phe Leu Val Arg Asp Cys Lys Thr
        195                 200                 205

Gly Glu Ile Leu Thr Arg Cys Thr Ser Leu Ser Val Leu Met Asn Thr
    210                 215                 220

Arg Thr Arg Arg Leu Ser Thr Ile Pro Asp Glu Val Arg Gly Glu Ile
225                 230                 235                 240

Gly Pro Ala Phe Ile Asp Asn Val Ala Val Lys Asp Asp Glu Ile Lys
                245                 250                 255

Lys Leu Gln Lys Leu Asn Asp Ser Thr Ala Asp Tyr Ile Gln Gly Gly
            260                 265                 270

Leu Thr Pro Arg Trp Asn Asp Leu Asp Val Asn Gln His Val Asn Asn
        275                 280                 285

Leu Lys Tyr Val Ala Trp Val Phe Glu Thr Val Pro Asp Ser Ile Phe
    290                 295                 300

Glu Ser His His Ile Ser Ser Phe Thr Leu Glu Tyr Arg Arg Glu Cys
305                 310                 315                 320

Thr Arg Asp Ser Val Leu Arg Ser Leu Thr Thr Val Ser Gly Gly Ser
                325                 330                 335

Ser Glu Ala Gly Leu Val Cys Asp His Leu Leu Gln Leu Glu Gly Gly
            340                 345                 350

Ser Glu Val Leu Arg Ala Arg Thr Glu Trp Arg Pro Lys Leu Thr Asp
        355                 360                 365

Ser Phe Arg Gly Ile Ser Val Ile Pro Ala Glu Pro Arg Val
    370                 375                 380
```

<210> 64
<211> 1240

<212> DNA
<213> Umbellularia sp.

<400> 64
ggcgcgccat ggccaccacc agcctggcct ccgccttctg ctccatgaag gccgtgatgc 60

tggcccgcga cggccgcggc atgaagcccc gcagctccga cctgcagctg cgcgccggca 120

acgcccccac ctccctgaag atgatcaacg gcaccaagtt cagctacacc gagagcctga 180

agcgcctgcc cgactggtcc atgctgttcg ccgtgatcac caccatcttc agcgccgccg 240

agaagcagtg gaccaacctg gagtggaagc ccaagcccaa gctgccccag ctgctggacg 300

accacttcgg cctgcacggc ctggtgttcc gccgcacctt cgccatccgc tcctacgagg 360

tgggccccga ccgcagcacc tccatcctgg ccgtgatgaa ccacatgcag gaggccaccc 420

tgaaccacgc caagagcgtg ggcatcctgg gcgacggctt cggcaccacc ctggagatgt 480

ccaagcgcga cctgatgtgg gtggtgcgcc gcacccacgt ggccgtggag cgctacccca 540

cctggggcga caccgtggag gtggagtgct ggatcggcgc cagcggcaac aacggcatgc 600

gccgcgactt cctggtgcgc gactgcaaga ccggcgagat cctgacccgc tgcacctccc 660

tgagcgtgct gatgaacacc cgcacccgcc gcctgagcac catccccgac gaggtgcgcg 720

gcgagatcgg ccccgccttc atcgacaacg tggccgtgaa ggacgacgag atcaagaagc 780

tgcagaagct gaacgactcc accgccgact acatccaggg cggcctgacc ccccgctgga 840

acgacctgga cgtgaaccag cacgtgaaca acctgaagta cgtggcctgg gtgttcgaga 900

ccgtgcccga cagcatcttc gagtcccacc acatcagctc cttcaccctg gagtaccgcc 960

gcgagtgcac ccgcgactcc gtgctgcgca gcctgaccac cgtgagcggc ggcagctccg 1020

aggccggcct ggtgtgcgac cacctgctgc agctggaggg cggcagcgag gtgctgcgcg 1080

cccgcaccga gtggcgcccc aagctgaccg actccttccg cggcatcagc gtgatccccg 1140

ccgagccccg cgtgatggac tacaaggacc acgacggcga ctacaaggac cacgacatcg 1200

actacaagga cgacgacgac aagtgactcg agttaattaa 1240

<210> 65
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic primer

<400> 65
ccgccgtgct ggacgtggtg 20

<210> 66
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic primer

<400> 66
ggtggcgggg tccagggtgt 20

<210> 67
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic primer

<400> 67
cggccggcgg ctccttcaac 20

<210> 68
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic primer

<400> 68
ggcgctcccg taggtcgggt 20

<210> 69
<211> 1335
<212> DNA
<213> Chlorella sorokiniana

<400> 69
cgcctgcaac gcaagggcag ccacagccgc tcccacccgc cgctgaaccg acacgtgctt 60

gggcgcctgc cgcctgcctg ccgcatgctt gtgctggtga ggctgggcag tgctgccatg 120

ctgattgagg cttggttcat cgggtggaag cttatgtgtg tgctgggctt gcatgccggg 180

caatgcgcat ggtggcaaga gggcggcagc acttgctgga gctgccgcgg tgcctccagg 240

tggttcaatc gcggcagcca gagggatttc agatgatcgc gcgtacaggt tgagcagcag 300

tgtcagcaaa ggtagcagtt tgccagaatg atcggttcag ctgttaatca atgccagcaa 360

gagaaggggt caagtgcaaa cacgggcatg ccacagcacg ggcaccgggg agtggaatgg 420

caccaccaag tgtgtgcgag ccagcatcgc cgcctggctg tttcagctac aacggcagga 480

gtcatccaac gtaaccatga gctgatcaac actgcaatca tcgggcgggc gtgatgcaag 540

catgcctggc gaagacacat ggtgtgcgga tgctgccggc tgctgcctgc tgcgcacgcc 600

gttgagttgg cagcaggctc agccatgcac tggatggcag ctgggctgcc actgcaatgt 660

ggtggatagg atgcaagtgg agcgaatacc aaaccctctg gctgcttgct gggttgcatg 720

gcatcgcacc atcagcagga gcgcatgcga agggactggc cccatgcacg ccatgccaaa 780

ccggagcgca ccgagtgtcc acactgtcac caggcccgca agctttgcag aaccatgctc 840

atggacgcat gtagcgctga cgtcccttga cggcgctcct ctcgggtgtg ggaaacgcaa 900

tgcagcacag gcagcagagg cggcggcagc agagcggcgg cagcagcggc gggggccacc 960

cttcttgcgg ggtcgcgccc cagccagcgg tgatgcgctg atcccaaacg agttcacatt 1020

catttgcatg cctggagaag cgaggctggg gcctttgggc tggtgcagcc cgcaatggaa 1080

tgcgggaccg ccaggctagc agcaaaggcg cctcccctac tccgcatcga tgttccatag 1140

tgcattggac tgcatttggg tggggcggcc ggctgtttct ttcgtgttgc aaaacgcgcc 1200

agctcagcaa cctgtcccgt gggtcccccg tgccgatgaa atcgtgtgca cgccgatcag 1260

ctgattgccc ggctcgcgaa gtaggcgccc tcctttctgc tcgccctctc tccgtcccgc 1320

cactagtggc gcgcc 1335

<210> 70
<211> 1146
<212> DNA
<213> Umbellularia californica

<400> 70
atggccacca ccagcctggc ctccgccttc tgctccatga aggccgtgat gctggcccgc 60

gacggccgcg gcatgaagcc ccgcagctcc gacctgcagc tgcgcgccgg caacgccccc 120

acctccctga agatgatcaa cggcaccaag ttcagctaca ccgagagcct gaagcgcctg 180

cccgactggt ccatgctgtt cgccgtgatc accaccatct tcagcgccgc cgagaagcag 240

tggaccaacc tggagtggaa gcccaagccc aagctgcccc agctgctgga cgaccacttc 300

ggcctgcacg gcctggtgtt ccgccgcacc ttcgccatcc gctcctacga ggtgggcccc 360

gaccgcagca cctccatcct ggccgtgatg aaccacatgc aggaggccac cctgaaccac 420

gccaagagcg tgggcatcct gggcgacggc ttcggcacca ccctggagat gtccaagcgc 480

gacctgatgt gggtggtgcg ccgcacccac gtggccgtgg agcgctaccc cacctggggc 540

gacaccgtgg aggtggagtg ctggatcggc gccagcggca acaacggcat gcgccgcgac 600

ttcctggtgc gcgactgcaa gaccggcgag atcctgaccc gctgcacctc cctgagcgtg 660

ctgatgaaca cccgcacccg ccgcctgagc accatccccg acgaggtgcg cggcgagatc 720

ggccccgcct tcatcgacaa cgtggccgtg aaggacgacg agatcaagaa gctgcagaag 780

ctgaacgact ccaccgccga ctacatccag ggcggcctga ccccccgctg gaacgacctg 840

gacgtgaacc agcacgtgaa caacctgaag tacgtggcct gggtgttcga gaccgtgccc 900

gacagcatct tcgagtccca ccacatcagc tccttcaccc tggagtaccg ccgcgagtgc 960

acccgcgact ccgtgctgcg cagcctgacc accgtgagcg gcggcagctc cgaggccggc 1020

ctggtgtgcg accacctgct gcagctggag ggcggcagcg aggtgctgcg cgcccgcacc 1080

gagtggcgcc ccaagctgac cgactccttc cgcggcatca gcgtgatccc cgccgagccc 1140

cgcgtg 1146

<210> 71
<211> 1146
<212> DNA
<213> Cinnamomum camphora

<400> 71
atggccacca cctccctggc ctccgccttc tgcagcatga aggccgtgat gctggcccgc 60

gacggccgcg gcatgaagcc ccgctccagc gacctgcagc tgcgcgccgg caacgcccag 120

acctccctga agatgatcaa cggcaccaag ttctcctaca ccgagagcct gaagaagctg 180

cccgactggt ccatgctgtt cgccgtgatc accaccatct tctccgccgc cgagaagcag 240

tggaccaacc tggagtggaa gcccaagccc aacccccccc agctgctgga cgaccacttc 300

ggcccccacg gcctggtgtt ccgccgcacc ttcgccatcc gcagctacga ggtgggcccc 360

gaccgctcca ccagcatcgt ggccgtgatg aaccacctgc aggaggccgc cctgaaccac 420

gccaagtccg tgggcatcct gggcgacggc ttcggcacca ccctggagat gtccaagcgc 480

gacctgatct gggtggtgaa gcgcacccac gtggccgtgg agcgctaccc cgcctggggc 540

gacaccgtgg aggtggagtg ctgggtgggc gcctccggca acaacggccg ccgccacgac 600

ttcctggtgc gcgactgcaa gaccggcgag atcctgaccc gctgcacctc cctgagcgtg 660

atgatgaaca cccgcacccg ccgcctgagc aagatccccg aggaggtgcg cggcgagatc 720

ggccccgcct tcatcgacaa cgtggccgtg aaggacgagg agatcaagaa gccccagaag 780

ctgaacgact ccaccgccga ctacatccag ggcggcctga ccccccgctg gaacgacctg 840

gacatcaacc agcacgtgaa caacatcaag tacgtggact ggatcctgga gaccgtgccc 900

gacagcatct tcgagagcca ccacatctcc tccttcacca tcgagtaccg ccgcgagtgc 960

accatggaca gcgtgctgca gtccctgacc accgtgagcg gcggctcctc cgaggccggc 1020

ctggtgtgcg agcacctgct gcagctggag ggcggcagcg aggtgctgcg cgccaagacc 1080

gagtggcgcc ccaagctgac cgactccttc cgcggcatca gcgtgatccc cgccgagtcc 1140

agcgtg 1146

<210> 72
<211> 72
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
oligonucleotide

<400> 72
atggactaca aggaccacga cggcgactac aaggaccacg acatcgacta caaggacgac 60

gacgacaagt ga 72

<210> 73
<211> 408
<212> DNA
<213> Chlorella vulgaris

<400> 73
ctcgaggcag cagcagctcg gatagtatcg acacactctg gacgctggtc gtgtgatgga 60

ctgttgccgc cacacttgct gccttgacct gtgaatatcc ctgccgcttt tatcaaacag 120

cctcagtgtg tttgatcttg tgtgtacgcg cttttgcgag ttgctagctg cttgtgctat 180

ttgcgaatac cacccccagc atcccccttcc ctcgtttcat atcgcttgca tcccaaccgc 240

aacttatcta cgctgtcctg ctatccctca gcgctgctcc tgctcctgct cactgccccct 300

cgcacagcct tggtttgggc tccgcctgta ttctcctggt actgcaacct gtaaaccagc 360

actgcaatgc tgatgcacgg gaagtagtgg gatgggaaca caaatgga 408

<210> 74
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic primer

<400> 74
ctgggcgacg gcttcggcac 20

<210> 75
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic primer

<400> 75
aagtcgcggc gcatgccgtt 20

<210> 76
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic primer

<400> 76
taccccgcct ggggcgacac 20

<210> 77
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic primer

<400> 77
cttgctcagg cggcgggtgc 20

<210> 78
<211> 1317
<212> DNA
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic polynucleotide

<400> 78
atggtggccg ccgccgcctc cagcgccttc ttccccgtgc ccgccccgg cgcctccccc 60

aagcccggca agttcggcaa ctggccctcc agcctgagcc cctccttcaa gcccaagtcc 120

atccccaacg gcggcttcca ggtgaaggcc aacgacagcg cccaccccaa ggccaacggc 180

tccgccgtga gcctgaagag cggcagcctg aacacccagg aggacacctc ctccagcccc 240

ccccccgca ccttcctgca ccagctgccc gactggagcc gcctgctgac cgccatcacc 300

accgtgttcg tgaagtccaa gcgccccgac atgcacgacc gcaagtccaa gcgccccgac 360

atgctggtgg acagcttcgg cctggagtcc accgtgcagg acggcctggt gttccgccag 420

tccttctcca tccgctccta cgagatcggc accgaccgca ccgccagcat cgagaccctg 480

atgaaccacc tgcaggagac ctccctgaac cactgcaaga gcaccggcat cctgctggac 540

ggcttcggcc gcaccctgga gatgtgcaag cgcgacctga tctgggtggt gatcaagatg 600

cagatcaagg tgaaccgcta ccccgcctgg ggcgacaccg tggagatcaa cacccgcttc 660

agccgcctgg gcaagatcgg catgggccgc gactggctga tctccgactg caacaccggc 720

gagatcctgg tgcgcgccac cagcgcctac gccatgatga accagaagac ccgccgcctg 780

tccaagctgc cctacgaggt gcaccaggag atcgtgcccc tgttcgtgga cagccccgtg 840

atcgaggact ccgacctgaa ggtgcacaag ttcaaggtga agaccggcga cagcatccag 900

aagggcctga cccccggctg gaacgacctg gacgtgaacc agcacgtgtc caacgtgaag 960

tacatcggct ggatcctgga gagcatgccc accgaggtgc tggagaccca ggagctgtgc 1020

tccctggccc tggagtaccg ccgcgagtgc ggccgcgact ccgtgctgga gagcgtgacc 1080

gccatggacc ccagcaaggt gggcgtgcgc tcccagtacc agcacctgct gcgcctggag 1140

gacggcaccg ccatcgtgaa cggcgccacc gagtggcgcc ccaagaacgc cggcgccaac 1200

ggcgccatct ccaccggcaa gaccagcaac ggcaactccg tgtccatgga ctacaaggac 1260

cacgacggcg actacaagga ccacgacatc gactacaagg acgacgacga caagtga 1317

<210> 79
<211> 1170
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic polynucleotide

<400> 79
atggccaccg catccacttt ctcggcgttc aatgcccgct gcggcgacct gcgtcgctcg 60

gcgggctccg ggccccggcg cccagcgagg cccctccccg tgcgcgggcg cgcccagctg 120

cccgactgga gccgcctgct gaccgccatc accaccgtgt tcgtgaagtc caagcgcccc 180

gacatgcacg accgcaagtc caagcgcccc gacatgctgg tggacagctt cggcctggag 240

tccaccgtgc aggacggcct ggtgttccgc cagtccttct ccatccgctc ctacgagatc 300

ggcaccgacc gcaccgccag catcgagacc ctgatgaacc acctgcagga gacctccctg 360

aaccactgca agagcaccgg catcctgctg gacggcttcg gccgcaccct ggagatgtgc 420

aagcgcgacc tgatctgggt ggtgatcaag atgcagatca aggtgaaccg ctaccccgcc 480

tggggcgaca ccgtggagat caacacccgc ttcagccgcc tgggcaagat cggcatgggc 540

cgcgactggc tgatctccga ctgcaacacc ggcgagatcc tggtgcgcgc caccagcgcc 600

tacgccatga tgaaccagaa gacccgccgc ctgtccaagc tgccctacga ggtgcaccag 660

gagatcgtgc ccctgttcgt ggacagcccc gtgatcgagg actccgacct gaaggtgcac 720

aagttcaagg tgaagaccgg cgacagcatc cagaagggcc tgacccccgg ctggaacgac 780

ctggacgtga accagcacgt gtccaacgtg aagtacatcg gctggatcct ggagagcatg 840

cccaccgagg tgctggagac ccaggagctg tgctccctgg ccctggagta ccgccgcgag 900

tgcggccgcg actccgtgct ggagagcgtg accgccatgg accccagcaa ggtgggcgtg 960

cgctcccagt accagcacct gctgcgcctg gaggacggca ccgccatcgt gaacggcgcc 1020

accgagtggc gccccaagaa cgccggcgcc aacggcgcca tctccaccgg caagaccagc 1080

aacggcaact ccgtgtccat ggactacaag gaccacgacg gcgactacaa ggaccacgac 1140

atcgactaca aggacgacga cgacaagtga 1170

<210> 80
<211> 1170
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic polynucleotide

<400> 80
atggctatca agacgaacag gcagcctgtg gagaagcctc cgttcacgat cgggacgctg 60

cgcaaggcca tccccgcgca ctgtttcgag cgctcggcgc ttcgtgggcg cgcccagctg 120

cccgactgga gccgcctgct gaccgccatc accaccgtgt tcgtgaagtc caagcgcccc 180

gacatgcacg accgcaagtc caagcgcccc gacatgctgg tggacagctt cggcctggag 240

tccaccgtgc aggacggcct ggtgttccgc cagtccttct ccatccgctc ctacgagatc 300

ggcaccgacc gcaccgccag catcgagacc ctgatgaacc acctgcagga gacctccctg 360

aaccactgca agagcaccgg catcctgctg gacggcttcg gccgcaccct ggagatgtgc 420

aagcgcgacc tgatctgggt ggtgatcaag atgcagatca aggtgaaccg ctaccccgcc 480

tggggcgaca ccgtggagat caacacccgc ttcagccgcc tgggcaagat cggcatgggc 540

cgcgactggc tgatctccga ctgcaacacc ggcgagatcc tggtgcgcgc caccagcgcc 600

tacgccatga tgaaccagaa gacccgccgc ctgtccaagc tgccctacga ggtgcaccag 660

gagatcgtgc ccctgttcgt ggacagcccc gtgatcgagg actccgacct gaaggtgcac 720

aagttcaagg tgaagaccgg cgacagcatc cagaagggcc tgacccccgg ctggaacgac 780

ctggacgtga accagcacgt gtccaacgtg aagtacatcg gctggatcct ggagagcatg 840

cccaccgagg tgctggagac ccaggagctg tgctccctgg ccctggagta ccgccgcgag 900

tgcggccgcg actccgtgct ggagagcgtg accgccatgg accccagcaa ggtgggcgtg 960

cgctcccagt accagcacct gctgcgcctg gaggacggca ccgccatcgt gaacggcgcc 1020

accgagtggc gccccaagaa cgccggcgcc aacggcgcca tctccaccgg caagaccagc 1080

aacggcaact ccgtgtccat ggactacaag gaccacgacg gcgactacaa ggaccacgac 1140

atcgactaca aggacgacga cgacaagtga 1170

<210> 81
<211> 1167
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic polynucleotide

<400> 81
atgacgttcg gggtcgccct cccggccatg ggccgcggtg tctcccttcc ccggcccagg 60

gtcgcggtgc gcgcccagtc ggcgagtcag gttttggaga gcgggcgcgc ccagctgccc 120

gactggagcc gcctgctgac cgccatcacc accgtgttcg tgaagtccaa gcgccccgac 180

atgcacgacc gcaagtccaa gcgccccgac atgctggtgg acagcttcgg cctggagtcc 240

accgtgcagg acggcctggt gttccgccag tccttctcca tccgctccta cgagatcggc 300

accgaccgca ccgccagcat cgagaccctg atgaaccacc tgcaggagac ctccctgaac 360

cactgcaaga gcaccggcat cctgctggac ggcttcggcc gcaccctgga gatgtgcaag 420

cgcgacctga tctgggtggt gatcaagatg cagatcaagg tgaaccgcta ccccgcctgg 480

ggcgacaccg tggagatcaa cacccgcttc agccgcctgg gcaagatcgg catgggccgc 540

gactggctga tctccgactg caacaccggc gagatcctgg tgcgcgccac cagcgcctac 600

gccatgatga accagaagac ccgccgcctg tccaagctgc cctacgaggt gcaccaggag 660

atcgtgcccc tgttcgtgga cagccccgtg atcgaggact ccgacctgaa ggtgcacaag 720

ttcaaggtga agaccggcga cagcatccag aagggcctga ccccggctg gaacgacctg 780

gacgtgaacc agcacgtgtc caacgtgaag tacatcggct ggatcctgga gagcatgccc 840

accgaggtgc tggagaccca ggagctgtgc tccctggccc tggagtaccg ccgcgagtgc 900

ggccgcgact ccgtgctgga gagcgtgacc gccatggacc ccagcaaggt gggcgtgcgc 960

tcccagtacc agcacctgct gcgcctggag gacggcaccg ccatcgtgaa cggcgccacc 1020

gagtggcgcc ccaagaacgc cggcgccaac ggcgccatct ccaccggcaa gaccagcaac 1080

ggcaactccg tgtccatgga ctacaaggac cacgacggcg actacaagga ccacgacatc 1140

gactacaagg acgacgacga caagtga 1167

<210> 82
<211> 1149

<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic polynucleotide

<400> 82
atgacgttcg gggtcgccct cccggccatg ggccgcggtg tctcccttcc ccggcccagg 60

gtcgcggtgc gcgcccagtc ggcgagtcag gttttggaga gcgggcgcgc ccccgactgg 120

tccatgctgt tcgccgtgat caccaccatc ttcagcgccg ccgagaagca gtggaccaac 180

ctggagtgga agcccaagcc caagctgccc cagctgctgg acgaccactt cggcctgcac 240

ggcctggtgt tccgccgcac cttcgccatc cgctcctacg aggtgggccc cgaccgcagc 300

acctccatcc tggccgtgat gaaccacatg caggaggcca ccctgaacca cgccaagagc 360

gtgggcatcc tgggcgacgg cttcggcacc accctggaga tgtccaagcg cgacctgatg 420

tgggtggtgc gccgcaccca cgtggccgtg gagcgctacc ccacctgggg cgacaccgtg 480

gaggtggagt gctggatcgg cgccagcggc aacaacggca tgcgccgcga cttcctggtg 540

cgcgactgca agaccggcga gatcctgacc cgctgcacct ccctgagcgt gctgatgaac 600

acccgcaccc gccgcctgag caccatcccc gacgaggtgc gcggcgagat cggccccgcc 660

ttcatcgaca acgtggccgt gaaggacgac gagatcaaga agctgcagaa gctgaacgac 720

tccaccgccg actacatcca gggcggcctg acccccgct ggaacgacct ggacgtgaac 780

cagcacgtga acaacctgaa gtacgtggcc tgggtgttcg agaccgtgcc cgacagcatc 840

ttcgagtccc accacatcag ctccttcacc ctggagtacc gccgcgagtg cacccgcgac 900

tccgtgctgc gcagcctgac caccgtgagc ggcggcagct ccgaggccgg cctggtgtgc 960

gaccacctgc tgcagctgga gggcggcagc gaggtgctgc gcgcccgcac cgagtggcgc 1020

cccaagctga ccgactcctt ccgcggcatc agcgtgatcc ccgccgagcc ccgcgtgatg 1080

gactacaagg accacgacgg cgactacaag gaccacgaca tcgactacaa ggacgacgac 1140

gacaagtga 1149

<210> 83
<211> 1146
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic polynucleotide

<400> 83
atggcttccg cggcattcac catgtcggcg tgccccgcga tgactggcag ggcccctggg 60

gcacgtcgct ccggacggcc agtcgccacc cgcctgaggg ggcgcgcccc cgactggtcc 120

atgctgttcg ccgtgatcac caccatcttc agcgccgccg agaagcagtg gaccaacctg 180

gagtggaagc ccaagcccaa gctgccccag ctgctggacg accacttcgg cctgcacggc 240

ctggtgttcc gccgcacctt cgccatccgc tcctacgagg tgggccccga ccgcagcacc 300

tccatcctgg ccgtgatgaa ccacatgcag gaggccaccc tgaaccacgc caagagcgtg 360

ggcatcctgg gcgacggctt cggcaccacc ctggagatgt ccaagcgcga cctgatgtgg 420

gtggtgcgcc gcacccacgt ggccgtggag cgctacccca cctggggcga caccgtggag 480

gtggagtgct ggatcggcgc cagcggcaac aacggcatgc gccgcgactt cctggtgcgc 540

gactgcaaga ccggcgagat cctgacccgc tgcacctccc tgagcgtgct gatgaacacc 600

cgcacccgcc gcctgagcac catccccgac gaggtgcgcg gcgagatcgg ccccgccttc 660

atcgacaacg tggccgtgaa ggacgacgag atcaagaagc tgcagaagct gaacgactcc 720

accgccgact acatccaggg cggcctgacc ccccgctgga acgacctgga cgtgaaccag 780

cacgtgaaca acctgaagta cgtggcctgg gtgttcgaga ccgtgcccga cagcatcttc 840

gagtcccacc acatcagctc cttcaccctg gagtaccgcc gcgagtgcac ccgcgactcc 900

gtgctgcgca gcctgaccac cgtgagcggc ggcagctccg aggccggcct ggtgtgcgac 960

cacctgctgc agctggaggg cggcagcgag gtgctgcgcg cccgcaccga gtggcgcccc 1020

aagctgaccg actccttccg cggcatcagc gtgatccccg ccgagccccg cgtgatggac 1080

tacaaggacc acgacggcga ctacaaggac cacgacatcg actacaagga cgacgacgac 1140

aagtga 1146

<210> 84
<211> 1155
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic polynucleotide

<400> 84
atggccaccg catccacttt ctcggcgttc aatgcccgct gcggcgacct gcgtcgctcg 60

gcgggctccg ggccccggcg cccagcgagg cccctccccg tgcgcgggcg cgccccgac 120

tggtccatgc tgttcgccgt gatcaccacc atcttcagcg ccgccgagaa gcagtggacc 180

aacctggagt ggaagcccaa gcccaagctg ccccagctgc tggacgacca cttcggcctg 240

cacggcctgg tgttccgccg caccttcgcc atccgctcct acgaggtggg ccccgaccgc 300

agcacctcca tcctggccgt gatgaaccac atgcaggagg ccaccctgaa ccacgccaag 360

agcgtgggca tcctgggcga cggcttcggc accaccctgg agatgtccaa gcgcgacctg 420

atgtgggtgg tgcgccgcac ccacgtggcc gtggagcgct accccacctg gggcgacacc 480

gtggaggtgg agtgctggat cggcgccagc ggcaacaacg gcatgcgccg cgacttcctg 540

gtgcgcgact gcaagaccgg cgagatcctg acccgctgca cctccctgag cgtgctgatg 600

aacacccgca cccgccgcct gagcaccatc cccgacgagg tgcgcggcga gatcggcccc 660

gccttcatcg acaacgtggc cgtgaaggac gacgagatca agaagctgca gaagctgaac 720

gactccaccg ccgactacat ccagggcggc ctgacccccc gctggaacga cctggacgtg 780

aaccagcacg tgaacaacct gaagtacgtg gcctgggtgt tcgagaccgt gcccgacagc 840

atcttcgagt cccaccacat cagctccttc accctggagt accgccgcga gtgcacccgc 900

gactccgtgc tgcgcagcct gaccaccgtg agcggcggca gctccgaggc cggcctggtg 960

tgcgaccacc tgctgcagct ggagggcggc agcgaggtgc tgcgcgcccg caccgagtgg 1020

cgccccaagc tgaccgactc cttccgcggc atcagcgtga tccccgccga gccccgcgtg 1080

atggactaca aggaccacga cggcgactac aaggaccacg acatcgacta caaggacgac 1140

gacgacaagt gatga 1155

<210> 85
<211> 1152
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic polynucleotide

<400> 85
atggctatca agacgaacag gcagcctgtg gagaagcctc cgttcacgat cgggacgctg 60

cgcaaggcca tccccgcgca ctgtttcgag cgctcggcgc ttcgtgggcg cgccccgac 120

tggtccatgc tgttcgccgt gatcaccacc atcttcagcg ccgccgagaa gcagtggacc 180

aacctggagt ggaagcccaa gcccaagctg ccccagctgc tggacgacca cttcggcctg 240

cacggcctgg tgttccgccg caccttcgcc atccgctcct acgaggtggg ccccgaccgc 300

agcacctcca tcctggccgt gatgaaccac atgcaggagg ccaccctgaa ccacgccaag 360

agcgtgggca tcctgggcga cggcttcggc accaccctgg agatgtccaa gcgcgacctg 420

atgtgggtgg tgcgccgcac ccacgtggcc gtggagcgct accccacctg gggcgacacc 480

gtggaggtgg agtgctggat cggcgccagc ggcaacaacg gcatgcgccg cgacttcctg 540

gtgcgcgact gcaagaccgg cgagatcctg acccgctgca cctccctgag cgtgctgatg 600

aacacccgca cccgccgcct gagcaccatc cccgacgagg tgcgcggcga gatcggcccc 660

gccttcatcg acaacgtggc cgtgaaggac gacgagatca agaagctgca gaagctgaac 720

gactccaccg ccgactacat ccagggcggc ctgaccccc gctggaacga cctggacgtg 780

aaccagcacg tgaacaacct gaagtacgtg gcctgggtgt tcgagaccgt gcccgacagc 840

atcttcgagt cccaccacat cagctccttc accctggagt accgccgcga gtgcacccgc 900

gactccgtgc tgcgcagcct gaccaccgtg agcggcggca gctccgaggc cggcctggtg 960

tgcgaccacc tgctgcagct ggagggcggc agcgaggtgc tgcgcgcccg caccgagtgg 1020

cgccccaagc tgaccgactc cttccgcggc atcagcgtga tccccgccga gccccgcgtg 1080

atggactaca aggaccacga cggcgactac aaggaccacg acatcgacta caaggacgac 1140

gacgacaagt ga 1152

<210> 86
<211> 1155

<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic polynucleotide

<400> 86
atggccaccg catccacttt ctcggcgttc aatgcccgct gcggcgacct gcgtcgctcg 60

gcgggctccg ggccccggcg cccagcgagg cccctccccg tgcgcgggcg cgccccgac 120

tggtccatgc tgttcgccgt gatcaccacc atcttctccg ccgccgagaa gcagtggacc 180

aacctggagt ggaagcccaa gcccaacccc ccccagctgc tggacgacca cttcggcccc 240

cacggcctgg tgttccgccg caccttcgcc atccgcagct acgaggtggg ccccgaccgc 300

tccaccagca tcgtggccgt gatgaaccac ctgcaggagg ccgccctgaa ccacgccaag 360

tccgtgggca tcctgggcga cggcttcggc accaccctgg agatgtccaa gcgcgacctg 420

atctgggtgg tgaagcgcac ccacgtggcc gtggagcgct accccgcctg gggcgacacc 480

gtggaggtgg agtgctgggt gggcgcctcc ggcaacaacg gccgccgcca cgacttcctg 540

gtgcgcgact gcaagaccgg cgagatcctg acccgctgca cctccctgag cgtgatgatg 600

aacacccgca cccgccgcct gagcaagatc cccgaggagg tgcgcggcga gatcggcccc 660

gccttcatcg acaacgtggc cgtgaaggac gaggagatca agaagcccca gaagctgaac 720

gactccaccg ccgactacat ccagggcggc ctgaccccc gctggaacga cctggacatc 780

aaccagcacg tgaacaacat caagtacgtg gactggatcc tggagaccgt gcccgacagc 840

atcttcgaga gccaccacat ctcctccttc accatcgagt accgccgcga gtgcaccatg 900

gacagcgtgc tgcagtccct gaccaccgtg agcggcggct cctccgaggc cggcctggtg 960

tgcgagcacc tgctgcagct ggagggcggc agcgaggtgc tgcgcgccaa gaccgagtgg 1020

cgccccaagc tgaccgactc cttccgcggc atcagcgtga tccccgccga gtccagcgtg 1080

atggactaca aggaccacga cggcgactac aaggaccacg acatcgacta caaggacgac 1140

gacgacaagt gatga 1155

<210> 87
<211> 1893
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic polynucleotide

<400> 87
gaattccttt cttgcgctat gacacttcca gcaaaaggta gggcgggctg cgagacggct 60

tcccggcgct gcatgcaaca ccgatgatgc ttcgaccccc cgaagctcct tcggggctgc 120

atgggcgctc cgatgccgct ccagggcgag cgctgtttaa atagccaggc ccccgattgc 180

aaagacatta tagcgagcta ccaaagccat attcaaacac ctagatcact accacttcta 240

cacaggccac tcgagcttgt gatcgcactc cgctaagggg gcgcctcttc ctcttcgttt 300

cagtcacaac ccgcaaacac tagtatggcc accgcatcca ctttctcggc gttcaatgcc 360

cgctgcggcg acctgcgtcg ctcggcgggc tccgggcccc ggcgcccagc gaggcccctc 420

cccgtgcgcg ggcgcgcccc cgactggtcc atgctgttcg ccgtgatcac caccatcttc 480

tccgccgccg agaagcagtg gaccaacctg gagtggaagc ccaagcccaa cccccccccag 540

ctgctggacg accacttcgg cccccacggc ctggtgttcc gccgcacctt cgccatccgc 600

agctacgagg tgggccccga ccgctccacc agcatcgtgg ccgtgatgaa ccacctgcag 660

gaggccgccc tgaaccacgc caagtccgtg ggcatcctgg gcgacggctt cggcaccacc 720

ctggagatgt ccaagcgcga cctgatctgg gtggtgaagc gcacccacgt ggccgtggag 780

cgctaccccg cctggggcga caccgtggag gtggagtgct gggtgggcgc ctccggcaac 840

aacggccgcc gccacgactt cctggtgcgc gactgcaaga ccggcgagat cctgacccgc 900

tgcacctccc tgagcgtgat gatgaacacc cgcacccgcc gcctgagcaa gatccccgag 960

gaggtgcgcg gcgagatcgg ccccgccttc atcgacaacg tggccgtgaa ggacgaggag 1020

atcaagaagc cccagaagct gaacgactcc accgccgact acatccaggg cggcctgacc 1080

ccccgctgga acgacctgga catcaaccag cacgtgaaca acatcaagta cgtggactgg 1140

atcctggaga ccgtgcccga cagcatcttc gagagccacc acatctcctc cttcaccatc 1200

gagtaccgcc gcgagtgcac catggacagc gtgctgcagt ccctgaccac cgtgagcggc 1260

ggctcctccg aggccggcct ggtgtgcgag cacctgctgc agctggaggg cggcagcgag 1320

gtgctgcgcg ccaagaccga gtggcgcccc aagctgaccg actccttccg cggcatcagc 1380

gtgatccccg ccgagtccag cgtgatggac tacaaggacc acgacggcga ctacaaggac 1440

cacgacatcg actacaagga cgacgacgac aagtgatgac tcgaggcagc agcagctcgg 1500

atagtatcga cacactctgg acgctggtcg tgtgatggac tgttgccgcc acacttgctg 1560

ccttgacctg tgaatatccc tgccgctttt atcaaacagc ctcagtgtgt ttgatcttgt 1620

gtgtacgcgc ttttgcgagt tgctagctgc ttgtgctatt tgcgaatacc acccccagca 1680

tcccccttccc tcgtttcata tcgcttgcat cccaaccgca acttatctac gctgtcctgc 1740

tatccctcag cgctgctcct gctcctgctc actgcccctc gcacagcctt ggtttgggct 1800

ccgcctgtat tctcctggta ctgcaacctg taaaccagca ctgcaatgct gatgcacggg 1860

aagtagtggg atgggaacac aaatggaaag ctt 1893

<210> 88
<211> 1887
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic polynucleotide

<400> 88
gaattccttt cttgcgctat gacacttcca gcaaaaggta gggcgggctg cgagacggct 60

tcccggcgct gcatgcaaca ccgatgatgc ttcgaccccc cgaagctcct tcggggctgc 120

atgggcgctc cgatgccgct ccagggcgag cgctgtttaa atagccaggc ccccgattgc 180

aaagacatta tagcgagcta ccaaagccat attcaaacac ctagatcact accacttcta 240

cacaggccac tcgagcttgt gatcgcactc cgctaagggg gcgcctcttc ctcttcgttt 300

cagtcacaac ccgcaaacac tagtatggct tccgcggcat tcaccatgtc ggcgtgcccc 360

gcgatgactg gcagggcccc tggggcacgt cgctccggac ggccagtcgc cacccgcctg 420

agggggcgcg cccccgactg gtccatgctg ttcgccgtga tcaccaccat cttctccgcc 480

gccgagaagc agtggaccaa cctggagtgg aagcccaagc ccaacccccc ccagctgctg 540

gacgaccact tcggccccca cggcctggtg ttccgccgca ccttcgccat ccgcagctac 600

gaggtgggcc ccgaccgctc caccagcatc gtggccgtga tgaaccacct gcaggaggcc 660

gccctgaacc acgccaagtc cgtgggcatc ctgggcgacg gcttcggcac caccctggag 720

atgtccaagc gcgacctgat ctgggtggtg aagcgcaccc acgtggccgt ggagcgctac 780

cccgcctggg gcgacaccgt ggaggtggag tgctgggtgg gcgcctccgg caacaacggc 840

cgccgccacg acttcctggt gcgcgactgc aagaccggcg agatcctgac ccgctgcacc 900

tccctgagcg tgatgatgaa cacccgcacc cgccgcctga gcaagatccc cgaggaggtg 960

cgcggcgaga tcggccccgc cttcatcgac aacgtggccg tgaaggacga ggagatcaag 1020

aagccccaga agctgaacga ctccaccgcc gactacatcc agggcggcct gacccccccgc 1080

tggaacgacc tggacatcaa ccagcacgtg aacaacatca agtacgtgga ctggatcctg 1140

gagaccgtgc ccgacagcat cttcgagagc caccacatct cctccttcac catcgagtac 1200

cgccgcgagt gcaccatgga cagcgtgctg cagtccctga ccaccgtgag cggcggctcc 1260

tccgaggccg gcctggtgtg cgagcacctg ctgcagctgg agggcggcag cgaggtgctg 1320

cgcgccaaga ccgagtggcg ccccaagctg accgactcct tccgcggcat cagcgtgatc 1380

cccgccgagt ccagcgtgat ggactacaag gaccacgacg gcgactacaa ggaccacgac 1440

atcgactaca aggacgacga cgacaagtga tgactcgagg cagcagcagc tcggatagta 1500

tcgacacact ctggacgctg gtcgtgtgat ggactgttgc cgccacactt gctgccttga 1560

cctgtgaata tccctgccgc ttttatcaaa cagcctcagt gtgtttgatc ttgtgtgtac 1620

gcgcttttgc gagttgctag ctgcttgtgc tatttgcgaa taccaccccc agcatccccct 1680

tccctcgttt catatcgctt gcatcccaac cgcaacttat ctacgctgtc ctgctatccc 1740

tcagcgctgc tcctgctcct gctcactgcc cctcgcacag ccttggtttg ggctccgcct 1800

gtattctcct ggtactgcaa cctgtaaacc agcactgcaa tgctgatgca cgggaagtag 1860

tgggatggga acacaaatgg aaagctt 1887

<210> 89
<211> 3631
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic polynucleotide

<400> 89

gaattccgcc tgcaacgcaa gggcagccac agccgctccc acccgccgct gaaccgacac 60

gtgcttgggc gcctgccgcc tgcctgccgc atgcttgtgc tggtgaggct gggcagtgct 120

gccatgctga ttgaggcttg gttcatcggg tggaagctta tgtgtgtgct gggcttgcat 180

gccgggcaat gcgcatggtg gcaagagggc ggcagcactt gctggagctg ccgcggtgcc 240

tccaggtggt tcaatcgcgg cagccagagg gatttcagat gatcgcgcgt acaggttgag 300

cagcagtgtc agcaaaggta gcagtttgcc agaatgatcg gttcagctgt taatcaatgc 360

cagcaagaga aggggtcaag tgcaaacacg ggcatgccac agcacgggca ccggggagtg 420

gaatggcacc accaagtgtg tgcgagccag catcgccgcc tggctgtttc agctacaacg 480

gcaggagtca tccaacgtaa ccatgagctg atcaacactg caatcatcgg gcgggcgtga 540

tgcaagcatg cctggcgaag acacatggtg tgcggatgct gccggctgct gcctgctgcg 600

cacgccgttg agttggcagc aggctcagcc atgcactgga tggcagctgg gctgccactg 660

caatgtggtg gataggatgc aagtggagcg aataccaaac cctctggctg cttgctgggt 720

tgcatggcat cgcaccatca gcaggagcgc atgcgaaggg actggcccca tgcacgccat 780

gccaaaccgg agcgcaccga gtgtccacac tgtcaccagg cccgcaagct ttgcagaacc 840

atgctcatgg acgcatgtag cgctgacgtc ccttgacggc gctcctctcg ggtgtgggaa 900

acgcaatgca gcacaggcag cagaggcggc ggcagcagag cggcggcagc agcggcgggg 960

gccacccttc ttgcggggtc gcgccccagc cagcggtgat gcgctgatcc caaacgagtt 1020

cacattcatt tgcatgcctg gagaagcgag gctggggcct ttgggctggt gcagcccgca 1080

atggaatgcg ggaccgccag gctagcagca aaggcgcctc ccctactccg catcgatgtt 1140

ccatagtgca ttggactgca tttgggtggg gcggccggct gtttctttcg tgttgcaaaa 1200

cgcgccagct cagcaacctg tcccgtgggt cccccgtgcc gatgaaatcg tgtgcacgcc 1260

gatcagctga ttgcccggct cgcgaagtag gcgccctcct ttctgctcgc cctctctccg 1320

tcccgccact agtatgctgc tgcaggcctt cctgttcctg ctggccggct tcgccgccaa 1380

gatcagcgcc tccatgacga acgagacgtc cgaccgcccc ctggtgcact tcacccccaa 1440

caagggctgg gggcgcgcca gccaccacgt gtacaagcgc ctgacccaga gcaccaacac 1500

caagtccccc agcgtgaacc agccctaccg caccggcttc cacttccagc cccccaagaa 1560

ctggatgaac gaccccaacg gccccatgat ctacaagggc atctaccacc tgttctacca 1620

gtggaacccc aagggcgccg tgtggggcaa catcgtgtgg gcccactcca ccagcaccga 1680

cctgatcaac tgggaccccc acccccccgc catcttcccc agcgcccccc tcgacatcaa 1740

cggctgctgg tccggcagcg ccaccatcct gcccaacggc aagcccgtga tcctgtacac 1800

cggcatcgac cccaagaacc agcaggtgca gaacatcgcc gagcccaaga acctgtccga 1860

cccctacctg cgcgagtgga agaagagccc cctgaacccc ctgatggccc ccgacgccgt 1920

gaacggcatc aacgcctcca gcttccgcga ccccaccacc gcctggctgg gccaggacaa 1980

gaagtggcgc gtgatcatcg gctccaagat ccaccgccgc ggcctggcca tcacctacac 2040

cagcaaggac ttcctgaagt gggagaagtc ccccgagccc ctgcactacg acgacggcag 2100

cggcatgtgg gagtgccccg acttcttccc cgtgacccgc ttcggcagca acggcgtgga 2160

gacctccagc ttcggcgagc ccaacgagat cctgaagcac gtgctgaaga tctccctgga 2220

cgacaccaag cacgactact acaccatcgg cacctacgac cgcgtgaagg acaagttcgt 2280

gcccgacaac ggcttcaaga tggacggcac cgcccccccgc tacgactacg gcaagtacta 2340

cgccagcaag accttcttcg actccgccaa gaaccgccgc atcctgtggg gctggaccaa 2400

cgagtcctcc agcgtggagg acgacgtgga gaagggctgg tccggcatcc agaccatccc 2460

ccgcaagatc tggctggacc gcagcggcaa gcagctgatc cagtggcccg tgcgcgaggt 2520

ggagcgcctg cgcaccaagc aggtgaagaa cctgcgcaac aaggtgctga agtccggcag 2580

ccgcctggag gtgtacggcg tgaccgccgc ccaggccgac gtggaggtgc tgttcaaggt 2640

gcgcgacctg gagaaggccg acgtgatcga gccctcctgg accgaccccc agctgatctg 2700

cagcaagatg aacgtgtccg tgaagtccgg cctgggcccc ttcggcctga tggtgctggc 2760

cagcaagaac ctggaggagt acacctccgt gtacttccgc atcttcaagg cccgccagaa 2820

cagcaacaag tacgtggtgc tgatgtgctc cgaccagtcc cgcagctccc tgaaggagga 2880

caacgacaag accacctacg gcgccttcgt ggacatcaac ccccaccagc ccctgagcct 2940

gcgcgccctg atcgaccact ccgtggtgga gagcttcggc ggcaagggcc gcgcctgcat 3000

cacctcccgc gtgtacccca agctggccat cggcaagtcc agccacctgt tcgccttcaa 3060

ctacggctac cagtccgtgg acgtgctgaa cctgaacgcc tggagcatga actccgccca 3120

gatcagcatg gactacaagg accacgacgg cgactacaag gaccacgaca tcgactacaa 3180

ggacgacgac gacaagtgat taattaaccg gctcgaggca gcagcagctc ggatagtatc 3240

gacacactct ggacgctggt cgtgtgatgg actgttgccg ccacacttgc tgccttgacc 3300

tgtgaatatc cctgccgctt ttatcaaaca gcctcagtgt gtttgatctt gtgtgtacgc 3360

gcttttgcga gttgctagct gcttgtgcta tttgcgaata ccacccccag catcccccttc 3420

cctcgtttca tatcgcttgc atcccaaccg caacttatct acgctgtcct gctatccctc 3480

agcgctgctc ctgctcctgc tcactgcccc tcgcacagcc ttggtttggg ctccgcctgt 3540

attctcctgg tactgcaacc tgtaaaccag cactgcaatg ctgatgcacg ggaagtagtg 3600

ggatgggaac acaaatggaa agcttgagct c 3631

<210> 90
<211> 621
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic polypeptide

<400> 90

```
Met Leu Leu Gln Ala Phe Leu Phe Leu Leu Ala Gly Phe Ala Ala Lys
1               5                  10                  15

Ile Ser Ala Ser Met Thr Asn Glu Thr Ser Asp Arg Pro Leu Val His
            20                  25                  30

Phe Thr Pro Asn Lys Gly Trp Gly Arg Ala Ser His His Val Tyr Lys
        35                  40                  45

Arg Leu Thr Gln Ser Thr Asn Thr Lys Ser Pro Ser Val Asn Gln Pro
    50                  55                  60

Tyr Arg Thr Gly Phe His Phe Gln Pro Pro Lys Asn Trp Met Asn Asp
65                  70                  75                  80

Pro Asn Gly Pro Met Ile Tyr Lys Gly Ile Tyr His Leu Phe Tyr Gln
                85                  90                  95

Trp Asn Pro Lys Gly Ala Val Trp Gly Asn Ile Val Trp Ala His Ser
            100                 105                 110

Thr Ser Thr Asp Leu Ile Asn Trp Asp Pro His Pro Pro Ala Ile Phe
        115                 120                 125

Pro Ser Ala Pro Phe Asp Ile Asn Gly Cys Trp Ser Gly Ser Ala Thr
    130                 135                 140

Ile Leu Pro Asn Gly Lys Pro Val Ile Leu Tyr Thr Gly Ile Asp Pro
145                 150                 155                 160

Lys Asn Gln Gln Val Gln Asn Ile Ala Glu Pro Lys Asn Leu Ser Asp
                165                 170                 175

Pro Tyr Leu Arg Glu Trp Lys Lys Ser Pro Leu Asn Pro Leu Met Ala
            180                 185                 190

Pro Asp Ala Val Asn Gly Ile Asn Ala Ser Ser Phe Arg Asp Pro Thr
        195                 200                 205

Thr Ala Trp Leu Gly Gln Asp Lys Lys Trp Arg Val Ile Ile Gly Ser
    210                 215                 220

Lys Ile His Arg Arg Gly Leu Ala Ile Thr Tyr Thr Ser Lys Asp Phe
225                 230                 235                 240

Leu Lys Trp Glu Lys Ser Pro Glu Pro Leu His Tyr Asp Asp Gly Ser
                245                 250                 255

Gly Met Trp Glu Cys Pro Asp Phe Phe Pro Val Thr Arg Phe Gly Ser
            260                 265                 270
```

```
Asn Gly Val Glu Thr Ser Ser Phe Gly Glu Pro Asn Glu Ile Leu Lys
        275                 280                 285

His Val Leu Lys Ile Ser Leu Asp Asp Thr Lys His Asp Tyr Tyr Thr
    290                 295                 300

Ile Gly Thr Tyr Asp Arg Val Lys Asp Lys Phe Val Pro Asp Asn Gly
305                 310                 315                 320

Phe Lys Met Asp Gly Thr Ala Pro Arg Tyr Asp Tyr Gly Lys Tyr Tyr
                325                 330                 335

Ala Ser Lys Thr Phe Phe Asp Ser Ala Lys Asn Arg Arg Ile Leu Trp
            340                 345                 350

Gly Trp Thr Asn Glu Ser Ser Ser Val Glu Asp Asp Val Glu Lys Gly
        355                 360                 365

Trp Ser Gly Ile Gln Thr Ile Pro Arg Lys Ile Trp Leu Asp Arg Ser
    370                 375                 380

Gly Lys Gln Leu Ile Gln Trp Pro Val Arg Glu Val Glu Arg Leu Arg
385                 390                 395                 400

Thr Lys Gln Val Lys Asn Leu Arg Asn Lys Val Leu Lys Ser Gly Ser
                405                 410                 415

Arg Leu Glu Val Tyr Gly Val Thr Ala Ala Gln Ala Asp Val Glu Val
            420                 425                 430

Leu Phe Lys Val Arg Asp Leu Glu Lys Ala Asp Val Ile Glu Pro Ser
        435                 440                 445

Trp Thr Asp Pro Gln Leu Ile Cys Ser Lys Met Asn Val Ser Val Lys
    450                 455                 460

Ser Gly Leu Gly Pro Phe Gly Leu Met Val Leu Ala Ser Lys Asn Leu
465                 470                 475                 480

Glu Glu Tyr Thr Ser Val Tyr Phe Arg Ile Phe Lys Ala Arg Gln Asn
                485                 490                 495

Ser Asn Lys Tyr Val Val Leu Met Cys Ser Asp Gln Ser Arg Ser Ser
            500                 505                 510

Leu Lys Glu Asp Asn Asp Lys Thr Thr Tyr Gly Ala Phe Val Asp Ile
        515                 520                 525

Asn Pro His Gln Pro Leu Ser Leu Arg Ala Leu Ile Asp His Ser Val
    530                 535                 540
```

Val Glu Ser Phe Gly Gly Lys Gly Arg Ala Cys Ile Thr Ser Arg Val
545 550 555 560

Tyr Pro Lys Leu Ala Ile Gly Lys Ser Ser His Leu Phe Ala Phe Asn
565 570 575

Tyr Gly Tyr Gln Ser Val Asp Val Leu Asn Leu Asn Ala Trp Ser Met
580 585 590

Asn Ser Ala Gln Ile Ser Met Asp Tyr Lys Asp His Asp Gly Asp Tyr
595 600 605

Lys Asp His Asp Ile Asp Tyr Lys Asp Asp Asp Asp Lys
610 615 620

<210> 91
<211> 997
<212> DNA
<213> Prototheca moriformis

<400> 91
cctgtcgatc gaagagaagg agacatgtgt acattattgg tgtgagggcg ctgaatcggc 60

catttttaa aatgatcacg ctcatgccaa tagacgcggc acataacgac gttcaaaccc 120

ccgccaaagc cgcggacaac cccatccctc cacacccccc acacaaagaa cccgccaccg 180

cttaccttgc ccacgaggta ggcctttcgt tgcgcaaaac cggcctcggt gatgaatgca 240

tgcccgttcc tgacgagcgc tgcccgggcc aacacgctct tttgctgcgt ctcctcaggc 300

ttgggggcct ccttgggctt gggtgccgcc atgatctgcg cgcatcagag aaacgttgct 360

ggtaaaaagg agcgcccggc tgcgcaatat atatataggc atgccaacac agcccaacct 420

cactcgggag cccgtcccac cacccccaag tcgcgtgcct tgacggcata ctgctgcaga 480

agcttcatga gaatgatgcc gaacaagagg ggcacgagga cccaatcccg gacatccttg 540

tcgataatga tctcgtgagt ccccatcgtc cgcccgacgc tccggggagc ccgccgatgc 600

tcaagacgag agggccctcg accaggaggg gctggcccgg gcgggcactg gcgtcgaagg 660

tgcgcccgtc gttcgcctgc agtcctatgc cacaaaacaa gtcttctgac ggggtgcgtt 720

tgctcccgtg cgggcaggca acagaggtat tcaccctggt catggggaga tcggcgatcg 780

agctgggata agagatactt ctggcaagca atgacaactt gtcaggaccg gaccgtgcca 840

tatatttctc acctagcgcc gcaaaaccta acaatttggg agtcactgtg ccactgagtt 900

cgactggtag ctgaatggag tcgctgctcc actaaacgaa ttgtcagcac cgccagccgg 960

ccgaggaccc gagtcatagc gagggtagta gcgcgcc 997

<210> 92
<211> 753
<212> DNA
<213> Prototheca moriformis

<400> 92
actaattgca atcgtgcagt aatcatcgat atggtcacaa gtagatcccc tactgacacc 60

ctctcgtaca tgtaggcaat gtcatcggcg ccgtcctgct gaccgatgcc gacgtagcag 120

agcagacccg ggccgatctg ggatacgagc cggccctcca cctgcgctcg aggtggaatc 180

aagtaaataa ccaatacact tttcgacacc acacagagtt gcacggacgg tggcgtacct 240

ctacgctcgc gctcttcacg cgctggacga ccgcacgcat gagcccgggt ggcttggtct 300

gggctgcaaa aatgcacaac aaacaagtat cagacgctca tggatgcaca cgcgctccca 360

agcacgctca gactaaatat tacagtagct cgtatctgat aagatatcga gacataccgc 420

tcaactcacc cgcaaactgc gccccgccag gtgatgcgca cagggcccca ccatgcgatc 480

catcgcatcg ctcctcgagg gcgctatcac gtggccggag agcgttcaca gcgtacgcca 540

ctgtatctgg gcggtatgcg gtccgtcaac atggagacag atacccgcac caccaccttg 600

caagctcttc catattggaa gtagaaaatt gtaattgtat catcgcacga ggggccaact 660

tgccgtcggc gagctgggcg acgaacacca cctggacgtt gtcgagactc gctcgtgccg 720

tgcgccgggc cgctgggtat ccagaccgtc gcc 753

<210> 93
<211> 1122
<212> DNA
<213> Prototheca moriformis

<400> 93
caacgacaac cagcaggcaa ctcggtcagc gacccaacac gcgagtcaaa ttgttgcgtg 60

ttcttgcctt gtctatttac tgtgatagca agactgtcgg tcagtcaata ccgcggtgcg 120

cacgtcgggg tgccaagcct agcagagcac gggacggctg gtgctgtgcg ccagctcagc 180

tcgcttcgcg accaattgta ggaccggcaa agtcaccaaa acatgccagc ggtgcgattc 240

aattggtcat gagctctaca aaattgtttt gtgcgtcgcg caggtatcca acggcgcggc 300

agagaaagtt tgacagctct cgatttcatc tcggaaaaat ggggagaatt tatgacacac 360

aagtgcgcag gcggcccagg cggccagcat attctggcgt gacctgggcc gcccacaaaa 420

tgcttggatg cactctaaaa taattatatt tgccatgaac aagggaagag ttaccgcacc 480

cagccctaga cttgggcgcc cgagcaaggt tacgtcaagc caccttcgcc catcgcccaa 540

ctccgtattc cccgacagcc gcacgtggcc ctcgccggaa tgaaccctga atcggcatca 600

cgccacgcgt tcgccaatcg ttccgctctc tggcttcatc ggcctgcgcc ttcacgtcgt 660

ggtcacgaca gtgcattcat acttccattt gcacctcggc acacactttt acgcatcgcc 720

taccccttgct gcggcagtct agggtcactt tgcagccatg ggacagtgct acaccaccgt 780

cggtgcgcaa agctatttca agtgaaccgt gggcggaaaa aaggaatgta cactgtctca 840

accgactcct acaattgttt accatgcaga tcagagctcg acggccatca tcgagcaggt 900

gtggggcctt ggtggcgcgg cgcggggccc cagggcgtcg caggcattga tggcactctg 960

agactttcgc acgcgcatga gggaccccat caagagaaga gtgtgtcttt atgtccccat 1020

tcatgatgat gtatcttgtg attgtcgcag tttggcaagt ttaaccggat cgccgctcca 1080

ggtgtggcgt ggcggatttt tctaggggtg cttgagcagt cg 1122

<210> 94
<211> 574
<212> DNA
<213> Prototheca moriformis

<400> 94
ggcccagggc cctgcggatg gcccacacca gatctagcct ctcttatgcc atgcccgcct 60

cgctgcccgt cgtatccccc cgccgatccg cgcgtagggg accgcggcct gacccacgcc 120

acgaaagagc tttgctcctc aatttctcgc caacagaacc gtatcaaacg ctcaacgcct 180

atcccgaaca atccgtattc acaccaaatc gagtataccg gactggtttg cctagtcttg 240

aaggaaatga tcccgtccat gctcggaagg gggagcgggc ggaggatcct actcatctct 300

gaaatgggat tggtccgaag atgggttggg caagcacgtg ccaaacccca gcgagttgct 360

gacgagcagg ctcatccaat cccccggcga atcctccctc acgccccgca tgcatacaag 420

tccctcccac acgccccctc ccatccattt tcgcctggtc cgaacgcgag cggcgtcgag 480

gcggaccact tgctccgcag cgccgtctgg gtctccaccc cacagcggct ttgctgccag 540

aggcaccccc cttgccccac ctcctcttgc agcc 574

<210> 95
<211> 1096
<212> DNA
<213> Prototheca moriformis

<400> 95
ccaggcaggc ggtagggttg ccgattgctt gagcgaattg gaagatataa ttttttgtgg 60

tgtccctgga cgctgtttgt ggcgctcctt tttggagaag attgcgtggg ggagctttcc 120

atgtaccacg cttccttctg aaaggattct ggccgagtcc tgatgagccc aaagaaaaca 180

cctgcctttc agtgctggca ctctgaaaac gtcaacagat gattatacat gtcacaaaag 240

gcagccgatt aggaacggga gctctggccg ttcgtttggc tgcctgggct gattgaagtg 300

atccaccctg ttcgaatgaa ggcggtcgag tcgaattatc gaccggagct gtcgggaagg 360

cgtccggggc agagtgaggt gctgcggcct ggttgtcgtt caaaaagacc ccggtagccc 420

aacaatcacg aacgaaagga atataattgc ttgcatacta tacattcagt ttctatgtgg 480

cgggtagaca agtctcatgg gcttctaaag gctgtccctt gaaggctact tataaaaact 540

tgctgcgcca tggcacggat cgcgcttgcg caggctgcaa ccctgcgcgc aaggtcaaat 600

acacagcaaa agatactaac agaatttcta aaaacattta aatatttgtt tcgaccagcc 660

aattgtggtc gtaggcacgc aaaagacttt gttttgcgcc caccgagcat ccacgctggc 720

agtcaagcca gtccgatgtg cattgcgtgg cagcatcgag gagcatcaaa aacctcgtgc 780

acgcttttct gtcaatcatc atcaaccact ccaccatgta tacccgatgc atcgcggtgc 840

gcagcgcgcc acgcgtccca gacccgccca aaaacccagc agcggcgaaa gcaaatcttc 900

acttgcccga aaccccgagc agcggcattc acacgtgggc gaaaacccca cttgccctaa 960

caggcgtatg tctgctgtca cgatgcctga caacggtatt atagatatac actgattaat 1020

gtttgagtgt gtgcgagtcg cgaatcagga atgaattgct agtaggcact ccgaccgggc 1080

gggggccgag ggacca 1096

<210> 96
<211> 1075
<212> DNA
<213> Prototheca moriformis

<400> 96
ggccgacagg acgcgcgtca aaggtgctgg gcgtgtatgc cctggtcggc aggtcgttgc 60

tgttgctgcg ctcgtggttc cgcaaccctg attttggcgt cttattctgg cgtggcaagc 120

gctgacgccc gcgagccggg ccggcggcga tgcggtgtct cacggctgcc gagctccaag 180

ggaggcaaga gcgcccggat cagctgaagg gctttacacg caaggtacag ccgctcctgc 240

aaggctgcgt ggtggacttg aacctgtagg tcctctgctg aagttcctcc actacctcac 300

caggcccagc agaccaaagc acaggctttt caggtccgtg tcatccactc taaaacactc 360

gactacgacc tactgatggc cctagattct tcatcaacaa tgcctgagac acttgctcag 420

aattgaaact ccctgaaggg accaccagag gccctgagtt gttccttccc cccgtggcga 480

gctgccagcc aggctgtacc tgtgatcgag gctggcggga aaataggctt cgtgtgctca 540

ggtcatggga ggtgcaggac agctcatgaa acgccaacaa tcgcacaatt catgtcaagc 600

taatcagcta tttcctcttc acgagctgta attgtcccaa aattctggtc taccgggggt 660

gatccttcgt gtacgggccc ttccctcaac cctaggtatg cgcgcatgcg gtcgccgcgc 720

aactcgcgcg agggccgagg gtttgggacg ggccgtcccg aaatgcagtt gcacccggat 780

gcgcggcgcc tttcttgcga taatttatgc aatggactgc tctgcaaatt tctgggtctg 840

tcgccaaccc taggatcagc ggcgtaggat ttcgtaatca ttcgtcctga tggggagcta 900

ccgactaccc taatatcagc ccggctgcct gacgccagcg tccacttttg cgtacacatt 960

ccattcgtgc ccaagacatt tcattgtggt gcgaagcgtc cccagttacg ctcacctgtt 1020

tcccgacctc cttactgttc tgtcgacaga gcgggcccac aggccggtcg cagcc 1075

<210> 97
<211> 772
<212> DNA
<213> Prototheca moriformis

<400> 97
tcaccagcgg acaaagcacc ggtgtatcag gtccgtgtca tccactctaa agagctcgac 60

tacgacctac tgatggccct agattcttca tcaaaaacgc ctgagacact tgcccaggat 120

tgaaactccc tgaagggacc accaggggcc ctgagttgtt ccttcccccc gtggcgagct 180

gccagccagg ctgtacctgt gatcggggct ggcgggaaaa caggcttcgt gtgctcaggt 240

tatgggaggt gcaggacagc tcattaaacg ccaacaatcg cacaattcat ggcaagctaa 300

tcagttattt cccattaacg agctataatt gtcccaaaat tctggtctac cgggggtgat 360

ccttcgtgta cgggcccttc cctcaaccct aggtatgcgc acatgcggtc gccgcgcaac 420

gcgcgcgagg gccgagggtt tgggacgggc cgtcccgaaa tgcagttgca cccggatgcg 480

tggcaccttt tttgcgataa tttatgcaat ggactgctct gcaaaattct ggctctgtcg 540

ccaaccctag gatcagcggt gtaggatttc gtaatcattc gtcctgatgg ggagctaccg 600

actgccctag tatcagcccg actgcctgac gccagcgtcc acttttgtgc acacattcca 660

ttcgtgccca agacatttca ttgtggtgcg aagcgtcccc agttacgctc acctgatccc 720

caacctcctt attgttctgt cgacagagtg ggcccagagg ccggtcgcag cc 772

<210> 98
<211> 991
<212> DNA
<213> Protoheca moriformis

<400> 98
cgaaggggtc tgcatcgatt cgcgcggtct ggaggccagc gtgactgctc gcgaaaatgc 60

tctgccgtgt cgggctctgg ctggggcggc cagagatctc accgtgccac acgcaactgc 120

cgcactctgt gcccgccacc tggcgcgcac atgcgacctc ttccccgtca taccctctcc 180

tcatgtgatc tttccacacg agtgacgcag gtgcgcggag tggagggaat caggacgttt 240

tcaaggtacc tgctcgagcc gtaccaacag ctgccgcccg gcaaggaaga gatcgaggca 300

gagattgccc ggctggaggc ccggataacg gagctcaaga gcaagctgtc cgagtgagac 360

cgcccaggtg cacgtgtcga ctcgctatga catgtactcg acacaacatg aggaattcat 420

cgaatttgta ggaagcgggc attggtacgg gagtgggaaa gcgaaaaaac ctccctccgg 480

cagtgccatc tgccggagtc gaacgttgat agggttctcg tgacagggtg tgacctctca 540

gccttgcatc aattaaacgc tatagacatt atcagtaacc gtgaatcccg cattggatgc 600

cacccgcgcg accattgggg acctgcatta cagatctagg tgagatgaca gcgaggcaac 660

ttcggcccgc ggcccagctt gcggcgcacc aatattggtc acgggaagcc acacaccgac 720

cataaatgaa tacttgtaag ctatgtcaac cgatcaatgg cgtcgaaagt gtgccacgag 780

gatccatctg gcggggcggc gtggcgcaca agcgcagtcg caatttctcg gacccatctg 840

acctaggccc agcgccgcgg gagaaatccc cggcgggtcc tccacgcagt aaccctaatg 900

agtatcgagc gccgaccatt tacaccatcg cccccgaaat ccttccgaca ttattattat 960

cttttagatc ttggaacaga ctctgccaac c 991

<210> 99
<211> 1347
<212> DNA
<213> Prototheca moriformis

<400> 99
agagagcgga ggtggggttg tgaggtgggg ttgctgacca ggagctcgcg tcgccgagcg 60

cgactcgcac acggtccagt tacccccccc tccgcccaaa cgcaagcctc ccatcttgat 120

gcctttccgg ccacctatac tatttcttag ttcgctgtaa catccagacc gtcctgaata 180

ataacaatgc cctgtgtcaa gtgcattcct aaaaaaattc tgtcccaacc aacaatccca 240

cctgaaatac caccagccct gcccagtaca ctcttccaat accatctccc tacctccacg 300

cgcaagcgac ccccatgcgc gaccaggctc gaaagtgatt tatgacttga gacgagcgag 360

tggcggcgcg gtcgactgcc ttttcatcac gtgccgtacg tcggcgaccg ctagggcttt 420

gcacggcaac gcacggcttc gccaacccga ccagccagga cctcgactac tctaccgcga 480

attcgcctca agaagtcgcc aaatgtgcca tacaccattc cttacagcac tgttcaaact 540

tgatgccaat tttgacattc gggttgctcg ttggctgcgc ccacatcggc cgtgagtgca 600

gcaggcggga tcggacacgg aggacgcggc gtcacgcccc gaacgcagcc cgtaactcta 660

catcaacacg acgtgttgcg taatcccgcc cggctgcgca tcgtgccaac ccattcgcga 720

tggatggtcg gaaaatggtg tgccaactgc cctgagggag gctctcgcga aacgggcacg 780

tccctgaaac cgaaactgtg gccttgtcgt cggccacgca agcacgtgga ccctaaacac 840

caagaaaatc agtaaacaag gttgacatcc tctacgggcg aattgtttgc ccaacccttc 900

atcgcacact gccattataa tgcatctagc tcggcgacaa gtttagaaaa ggcaggctgc 960

attgttccat ttcgccgtgg cggcgtgggt gcccatttta cgaggtttgg gctcccgggc 1020

agcgaccgag ccaggtcgag tccctctcgc ccgtcgacaa tgttgcgaac cccacaagcg 1080

gctaacaaca acttgatggt acctgtacac tgccaattcc ttcttccccg gccgaggttt 1140

acacgtgatg gccatggctt cgcattcagg ccgacttccc attccgactt tccagagggt 1200

ccgcggacgc tgggggttgg ctgcctgagg cccacccttt gttccccgcg tcccgacaaa 1260

cacaattgcg ttacataagg gggagccgcc cccgttcaga gtgcagaaat ctttcactat 1320

attttccagt cgtcagcgaa atcaagt 1347

<210> 100
<211> 1180
<212> DNA
<213> Prototheca moriformis

<400> 100
gatggtgggg tgtctgcctt gggctgggtg atggaggctg gtggtgcgcg ggtttcctga 60

tgcattctat ctacgcagtg tcatggtgtc cattccacac accagtacac ccttacacta 120

aggatccatc cctccttccc tcttcaggac tacatggacc ccacgagcta ccgaccgggc 180

tttctcaaaa acgtcaaggt catgtttgac atgcgggacg tggtggacga cgtgcaaggt 240

gcgtccggag tgcgcgcaaa tgagcaagtc gggcaatgtg tcggggtggg caccggggct 300

ggagatccgc gatccccgag aaaacgccgt accacccccc gcgctattcc ctcgattgcg 360

cgcagatgtg gtgaccgaca cgggggacaa cctggcggac atggggcgcc ggacctggaa 420

gcacgccaag tcgcacacgg ggaggctcgt gcagtccccc ccatcgtacc tcaagggtct 480

ctttggtcgc gatccaaagt acgctggtgg catggcatgc ccgaaatgaa catcatgtgt 540

gatctccgat tgccaatggc cacctccacg gaccaccttg caggcggaag cgcaatccag 600

ggcccgagcc tgacgaggac ggagactcct cgtccagcgc ggggtccccg acccgacgca 660

gcagccgacc cctgctaacc cggcaacgat cggaccagca accttgctgt agttccgatc 720

cgtgatgacg ggcattgccg ccgctcgatc cgctttgatg actgtctatt atttgcgcgg 780

agccccctcg gaaccctacc ccgctcttgc aagccccttg catcggagat cctcgtgcgc 840

ccgccatgac cccactggat tgcccaacat ccttctttat cgtgtaaaat gtgattcctc 900

ggctgcaatc gactggcctt cgcttctggc cccaagaggg ctcgaacgtg cggcagcgag 960

ggcgctgaca cacccaagcc ctagggcttt caacgtcggc tgccaggccg gatagggga 1020

tcgcctcctt tccaccaccc acctacgagg gattcgagtc ggcttccagc tcagctattc 1080

ggccgcgccc ccggccctgc agacgtcctc cagtttccga acaggtcgct ctcagaacac 1140

ctgccgcggc tgcgatacgg caggctctca aagcgtcgac 1180

<210> 101
<211> 1263
<212> DNA
<213> Prototheca moriformis

<400> 101
cgcgtggagc ggtgcgtgcg gatgccgcgc gcctgccaag gccttttgta tgcctggcct 60

gggaagtttc ctgactgaag catcttcaag atgctctctc acgaccagcg acaccaacac 120

cgtcactttt tgcccctcct gccgcaggtg ccactttcta ctttgacgtc ttctccaggc 180

ggtacattgc gggactgagc gccaattcgg ccaagaacag cgctgtcgac ttgaggaggc 240

aggggtccgt cgactctgcc gagtgacacg ccttcgaccc gactgtacta cggcctgctg 300

aagagtgggt ctcgccggcc ggcgtgaccg gccctgtgcc cacaatcgac catctattcg 360

ctccttgtca tctggcgccg tcaattgccc gcgacttgac ggcaactggc tcgatcgagt 420

cgtattgaaa aagcacgttt tgtcctacag ggccgcggtc cgttaccaac gtggttctcg 480

ttaggttttc gtcgggcggt ggtgcgcgaa ctgtccgatg ccatcccggc aaaccccagc 540

aaggtcgcca gtctggttct gacgcaatag agtgcgtttt gggccagtct aaaaattcgt 600

ctggcatgac gtggctccac atcgtacccg gagcctgcct tggtaatgtg aggcaccggt 660

gccaactcca ttatggcagg catcgagcgc gcaggtgagt acatgacctt ccgtgaattg 720

ggaaggcgag cttgtgtaac gcctgcgatc gtgccagtga ggcatcgtaa actcaaaata 780

ttttgtagaa agtgtctgat gcctggtgag gctgcgtagg gcaagggcaa gcccttggca 840

gatgggtaat gggtccggac ctcacaacag caaccccgcg tccccttag ggcccctgag 900

gctcgatggc agggccagcg agcccgcggc caaagggcgc catcccacgg tcgcccaacg 960

actccacggg tcctatacct catcttgaat ggcactaaaa actatagaat atcgggcact 1020

ggtgggcgtc tggggtacag ctggccgagc gcagtggcaa accctaggtc ccgcctcaag 1080

ggcgattccc gggtcaatga cacgcaagca agatcacatg gcgcggtccg cctcgcggct 1140

ccacacccag gccctagttt cgcaacccat aaatatcgcc ccgataccat cataagccag 1200

caaataattt tttatcagag ttccaaacct cctcagctgt gggaaaccag cccactctga 1260

acg 1263

<210> 102
<211> 1400
<212> DNA
<213> Prototheca moriformis

<400> 102
ccgagcagtt catggccaag tacaaggact agagaccgga ggtcggtagg ctgaatggag 60

ctggcgtcgt cgtgcgcgac gtgcacgcga tgcgatacta cgaccccaca aacgcatgcc 120

tcccatcttg atgcctttcc ggccatttat actatttctc atttcgctgt aacatcttga 180

ataatagaat tgccctgtgt caagtggatt ccaagaaata ttctgtccca acaaaacaac 240

ccaacctgaa aacaacctca aataccacca gccctgccca cctgcccagt acacttttcc 300

aataccatct ccctaccttc acgcgcaagc ggcacccatg cgcgaccagg ctcgaaagga 360

tttcacgact caggacgagc gagtggcggc gcgaccgcct gcctgttcgt cacgtgccgt 420

acgtcggcga ccgctagagc tttgcctggc aacccccggc ttcgtcaacc cggccagcca 480

ggatctcgac cactctaccg cgaaatcgcc tcaagaagtc gccaaaagtg ccgtacacca 540

tgcttcgcag cgctgttcaa acttgatgcc aatcttgaca atcaggttgc tcgttggctg 600

cgtccacatc ggccgtgatt gcagcaggcg gggatcggac acggaggacg cggcgtcacg 660

ccgcgaacgc agcccgtaac tctacatcaa cgcgatatgt tgcgtaatcc cgcccggctg 720

cgcattgtga caacccattc gcgatggatg gtcggaaaat ggtgtgccaa ctgccctgag 780

ggactctctc gcgaaacggg cacgtccctg tatccgaaac tgtggcatgg ccttgtcgac 840

cacgcaagca cgtggaccct aacaccacga aaataagtaa aaaaggttga catcctctac 900

gagcgaattg tttgctcgac ccttcatcgc acactgtcat tataatgcat ctagctcggc 960

gacaagttta aaaaaggcag gctgcattat tccattttgc cgtggcggca tgggtgccca 1020

ttttatgagg tttgggctct tgggcagcga ccgagccagg ttgagtccct ctcgcccgtc 1080

gacaacgttc caaagcccat aagtggctaa taaacaactt gatggtacct gtacactgcc 1140

agttccttct tccccggccg aggtttacac gtgatggcca tggcttcgcg tttcaggctg 1200

acttcccatt ccgactttcc agagggtccg cggacgccgg gggttggctg cgtgaggccc 1260

acccccttgtt ccccgcgtcc cgacaaacac aattgcgtta cataaggggg aagccgcccc 1320

ccgttcagag tgcaaacatc tttcattata tttttcagtc gtcagcgaaa tcaagtatgt 1380

cgctgacagg catgaaggcc 1400

<210> 103
<211> 3681
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic polynucleotide

<400> 103
gccctttgtc atcgttggca tgctttttgc gtatgtacca tatgttgaat gtataatacg 60
aacggttgac cgtctgagat gcgagctttg ggtcttgtca aatgcgtggc cgcacggctc 120
cctcgcaccc agccccgagg cgtcgcgcac ctggcgagga gcagacccac gccaagaaag 180
tctagtccag catgtaacaa catcaggcaa tgtgacgttt tcggttcccg atttctctgc 240
cgctctttga cggcaggcac gggcgagcaa ccggcggcgc tcgcgtcagg cacgatggat 300
gcggcgctgc ccacctgtca atgtacccca ccagtctgtc gatcgctaca agcaaccttg 360
tgctccacat tcccacttgc agacagtcta gtcgattttg ccaagctgga tgtgaggatt 420
ggccatatct tggaggccaa gattcacccg gatgctgatg ggtacgtacg cgagccaggc 480
aggcagctgc gttgactttc tgattggcac aaagctttgg ctactctcaa taccaaccac 540
gtgccccttc tgcacacctg cttccttctg atgaccactc gccacgcatg tcgcagtctg 600
tacgtcgagc agatcgacct cggcgaggag gggggccctc gcaccatcgt gagtggcctg 660
gtccggcacg tgaccctgga ggaccttgtc ggccggcggg tggtggtgct ggccaacctc 720
aagcctcgga gcatgcgcgg ggtcaaatcg gctgggatgc tgctctgcgc cgccaacgcg 780
gatcacaccg cggtggagcc gctgcgggtc ccggacgccg ccgtgacggg ggagcgggtc 840
tgggcggggg acgaggcact cctgtccacg gagcctgcca cacccaatca ggtaaggaca 900
cgttattggt gcgcatggtg cgaatgcgtg gtctgacctg ctgtgggtat gtgttgtggg 960
attggaaacc gaatgagggc cgttcaggat tgagcccttg gccccaccct gctcatcctc 1020
tcacgcccgc aggtccagaa gaagaaaatc tgggaggcag tacagccgct gctgagagtg 1080
aacgcccagg ggatcgctac tgtggcagga gaggctatgg tgaccagtgc ggggccactg 1140
accgcgccca cgctggttga cgccgcgatt tcctgacgcg agcgactgat tcttgacctt 1200
tgagaagcca ccacagcacc attttcattg ttcatccttg atttcagtac gacttctcac 1260
catttcagta ctgtaggacc cccaaaatag tgtgatcacg ctcgcaaggc acctgtgtga 1320
tcacggggaa gggcgaattc ctttcttgcg ctatgacact tccagcaaaa ggtagggcgg 1380
gctgcgagac ggcttcccgg cgctgcatgc aacaccgatg atgcttcgac cccccgaagc 1440
tccttcgggg ctgcatgggc gctccgatgc cgctccaggg cgagcgctgt ttaaatagcc 1500
aggcccccga ttgcaaagac attatagcga gctaccaaag ccatattcaa acacctagat 1560
cactaccact tctacacagg ccactcgagc ttgtgatcgc actccgctaa gggggcgcct 1620
cttcctcttc gtttcagtca caacccgcaa acggcgcgcc atgctgctgc aggccttcct 1680
gttcctgctg gccggcttcg ccgccaagat cagcgcctcc atgacgaacg agacgtccga 1740
ccgccccctg gtgcacttca cccccaacaa gggctggatg aacgacccca acggcctgtg 1800
gtacgacgag aaggacgcca agtggcacct gtacttccag tacaacccga acgacaccgt 1860
ctgggggacg cccttgttct ggggccacgc cacgtccgac gacctgacca actgggagga 1920
ccagcccatc gccatcgccc cgaagcgcaa cgactccggc gccttctccg gctccatggt 1980

ggtggactac aacaacacct ccggcttctt caacgacacc atcgacccgc gccagcgctg 2040

cgtggccatc tggacctaca acaccccgga gtccgaggag cagtacatct cctacagcct 2100

ggacggcggc tacaccttca ccgagtacca gaagaacccc gtgctggccg ccaactccac 2160

ccagttccgc gacccgaagg tcttctggta cgagccctcc cagaagtgga tcatgaccgc 2220

ggccaagtcc caggactaca agatcgagat ctactcctcc gacgacctga agtcctggaa 2280

gctggagtcc gcgttcgcca acgagggctt cctcggctac cagtacgagt gccccggcct 2340

gatcgaggtc cccaccgagc aggaccccag caagtcctac tgggtgatgt tcatctccat 2400

caaccccggc gccccggccg gcggctcctt caaccagtac ttcgtcggca gcttcaacgg 2460

cacccacttc gaggccttcg acaaccagtc ccgcgtggtg gacttcggca aggactacta 2520

cgccctgcag accttcttca acaccgaccc gacctacggg agcgccctgg gcatcgcgtg 2580

ggcctccaac tgggagtact ccgccttcgt gcccaccaac ccctggcgct cctccatgtc 2640

cctcgtgcgc aagttctccc tcaacaccga gtaccaggcc aacccggaga cggagctgat 2700

caacctgaag gccgagccga tcctgaacat cagcaacgcc ggcccctgga gccggttcgc 2760

caccaacacc acgttgacga aggccaacag ctacaacgtc gacctgtcca acagcaccgg 2820

caccctggag ttcgagctgg tgtacgccgt caacaccacc cagacgatct ccaagtccgt 2880

gttcgcggac ctctccctct ggttcaaggg cctggaggac cccgaggagt acctccgcat 2940

gggcttcgag gtgtccgcgt cctccttctt cctggaccgc gggaacagca aggtgaagtt 3000

cgtgaaggag aacccctact tcaccaaccg catgagcgtg aacaaccagc ccttcaagag 3060

cgagaacgac ctgtcctact acaaggtgta cggcttgctg gaccagaaca tcctggagct 3120

gtacttcaac gacggcgacg tcgtgtccac caacacctac ttcatgacca ccgggaacgc 3180

cctgggctcc gtgaacatga cgacgggggt ggacaacctg ttctacatcg acaagttcca 3240

ggtgcgcgag gtcaagtgat taattaactc gaggcagcag cagctcggat agtatcgaca 3300

cactctggac gctggtcgtg tgatggactg ttgccgccac acttgctgcc ttgacctgtg 3360

aatatccctg ccgcttttat caaacagcct cagtgtgttt gatcttgtgt gtacgcgctt 3420

ttgcgagttg ctagctgctt gtgctatttg cgaataccac ccccagcatc cccttccctc 3480

gtttcatatc gcttgcatcc caaccgcaac ttatctacgc tgtcctgcta tccctcagcg 3540

ctgctcctgc tcctgctcac tgcccctcgc acagccttgg tttgggctcc gcctgtattc 3600

tcctggtact gcaacctgta aaccagcact gcaatgctga tgcacgggaa gtagtgggat 3660

gggaacacaa atggaaagct t 3681

<210> 104
<211> 3850
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic polynucleotide

<400> 104

tttggccccg ctttccagct ccggatctgc tggcgtccgc cgcgagacgt gacatcgcac 60

gtcgccggga gcgccagctt gatcacttgg caggggccg tgctctacaa ataccaggcc 120

ccgcggcggt cagttcgcac atccaatacc tgccgagcca tcttgcctac acttttatc 180

gactcctcta ctctgttcgc gagagcgctc ggtccaggct tggaattcgc cgaattcagc 240

tcgatcagtc gcttctgcaa ctgatctcgg ccgttcgcag actgcctttt ctcagcttgt 300

caggtagcga gttgttgttt tatatttatt cgatttcatc tgtgttgcat gtcttgttcg 360

tgctgtgcgt tctttctggg ccgcgctgtc gggtcgcatg ggctagctgt actcatgtta 420

gtcatgccgg tccgaccttg ttcgaggaag gccccacact gagcgtgccc tctttctaca 480

ccccttgtgc agaaattaga tagaaagcag aattcctttc ttgcgctatg acacttccag 540

caaaaggtag ggcgggctgc gagacggctt cccggcgctg catgcaacac cgatgatgct 600

tcgacccccc gaagctcctt cggggctgca tgggcgctcc gatgccgctc cagggcgagc 660

gctgtttaaa tagccaggcc cccgattgca aagacattat agcgagctac caaagccata 720

ttcaaacacc tagatcacta ccacttctac acaggccact cgagcttgtg atcgcactcc 780

gctaaggggg cgcctcttcc tcttcgtttc agtcacaacc cgcaaacggc gcgccatgct 840

gctgcaggcc ttcctgttcc tgctggccgg cttcgccgcc aagatcagcg cctccatgac 900

gaacgagacg tccgaccgcc ccctggtgca cttcaccccc aacaagggct ggatgaacga 960

ccccaacggc ctgtggtacg acgagaagga cgccaagtgg cacctgtact tccagtacaa 1020

cccgaacgac accgtctggg ggacgccctt gttctggggc cacgccacgt ccgacgacct 1080

gaccaactgg gaggaccagc ccatcgccat cgccccgaag cgcaacgact ccggcgcctt 1140

ctccggctcc atggtggtgg actacaacaa cacctccggc ttcttcaacg acaccatcga 1200

cccgcgccag cgctgcgtgg ccatctggac ctacaacacc ccggagtccg aggagcagta 1260

catctcctac agcctggacg gcggctacac cttcaccgag taccagaaga accccgtgct 1320

ggccgccaac tccacccagt tccgcgaccc gaaggtcttc tggtacgagc cctcccagaa 1380

gtggatcatg accgcggcca agtcccagga ctacaagatc gagatctact cctccgacga 1440

cctgaagtcc tggaagctgg agtccgcgtt cgccaacgag ggcttcctcg gctaccagta 1500

cgagtgcccc ggcctgatcg aggtccccac cgagcaggac cccagcaagt cctactgggt 1560

gatgttcatc tccatcaacc ccggcgcccc ggccggcggc tccttcaacc agtacttcgt 1620

cggcagcttc aacggcaccc acttcgaggc cttcgacaac cagtcccgcg tggtggactt 1680

cggcaaggac tactacgccc tgcagacctt cttcaacacc gacccgacct acgggagcgc 1740

cctgggcatc gcgtgggcct ccaactggga gtactccgcc ttcgtgccca ccaacccctg 1800

gcgctcctcc atgtccctcg tgcgcaagtt ctccctcaac accgagtacc aggccaaccc 1860

ggagacggag ctgatcaacc tgaaggccga gccgatcctg aacatcagca acgccggccc 1920

ctggagccgg ttcgccacca acaccacgtt gacgaaggcc aacagctaca acgtcgacct 1980

gtccaacagc accggcaccc tggagttcga gctggtgtac gccgtcaaca ccacccagac 2040

gatctccaag tccgtgttcg cggacctctc cctctggttc aagggcctgg aggaccccga 2100

ggagtacctc cgcatgggct tcgaggtgtc cgcgtcctcc ttcttcctgg accgcgggaa 2160

cagcaaggtg aagttcgtga aggagaaccc ctacttcacc aaccgcatga gcgtgaacaa 2220

ccagcccttc aagagcgaga acgacctgtc ctactacaag gtgtacggct tgctggacca 2280

gaacatcctg gagctgtact tcaacgacgg cgacgtcgtg tccaccaaca cctacttcat 2340

gaccaccggg aacgccctgg gctccgtgaa catgacgacg ggggtggaca acctgttcta 2400

catcgacaag ttccaggtgc gcgaggtcaa gtgattaatt aactcgaggc agcagcagct 2460

cggatagtat cgacacactc tggacgctgg tcgtgtgatg gactgttgcc gccacacttg 2520

ctgccttgac ctgtgaatat ccctgccgct tttatcaaac agcctcagtg tgtttgatct 2580

tgtgtgtacg cgcttttgcg agttgctagc tgcttgtgct atttgcgaat accaccccca 2640

gcatcccctt ccctcgtttc atatcgcttg catcccaacc gcaacttatc tacgctgtcc 2700

tgctatccct cagcgctgct cctgctcctg ctcactgccc ctcgcacagc cttggtttgg 2760

gctccgcctg tattctcctg gtactgcaac ctgtaaacca gcactgcaat gctgatgcac 2820

gggaagtagt gggatgggaa cacaaatgga ccgacacgcc cccggcccag gtccagttct 2880

cctgggtctt ccagaggccc gtcgccatgt aaagtggcag agattggcgc ctgattcgat 2940

ttggatccaa ggatctccaa tcggtgatgg ggactgagtg cccaactacc acccttgcac 3000

tatcgtcctc gcactattta ttcccacctt ctgctcgccc tgccgggcga ttgcgggcgt 3060

ttctgccctt gacgtatcaa tttcgccccc gctggcgcga ggattcttca ttctaataag 3120

aactcactcc cgccagctct gtacttttcc tgcggggccc ctgcatggct tgttcccaat 3180

gcttgctcga tcgacggcgc ccattgccca cggcgctgcc gcatccatgt gaagaaacac 3240

ggaagagtgc gaagactgga agtgaattaa gagtataaga agaggtacca agggattctc 3300

aggtgctctt aggaacggct tttccttcgc caagagaaac tgctactgct cgtgtcgcca 3360

cggtggtcaa gccgccccat ctgcgatcca ccaggcccat ccgcggactc gcgatcagcc 3420

tgctggatcc ggactgccga cctgaccgct cgcatccacc attacaaccc tccaattgga 3480

caccactccc acgtcctaaa gttcaccatg caagctgatc gatcgcattc gccgatgcac 3540

tcgcctgcca cagaggtgtg cgcttcggac tagcgtgcag gcgccccgag gccaccagca 3600

tgcaccgatg gaagcgggca cggccgctgc tccaggtcgc tggctcgctc agacccatag 3660

caacctccgc tgcgtcccta aatgtcacac agagcgtctt tgatgggtac ggatgggaga 3720

gaatctgatt gggcattgct ggtgcagtgc aggaagatgg caagtgcaca gtcagtcatg 3780

ctgtacaaac tggtgcctcg tagtattgac tcgtatagtg catagtatca tgcatggtcg 3840

ttacttgcaa 3850

<210> 105
<211> 3108
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic polynucleotide

<400> 105
tttggccccg ctttccagct ccggatctgc tggcgtccgc cgcgagacgt gacatcgcac 60

gtcgccggga gcgccagctt gatcacttgg caggggccg tgctctacaa ataccaggcc 120

ccgcggcggt cagttcgcac atccaatacc tgccgagcca tcttgcctac actttttatc 180

gactcctcta ctctgttcgc gagagcgctc ggtccaggct tggaattcgc cgaattcagc 240

tcgatcagtc gcttctgcaa ctgatctcgg ccgttcgcag actgcctttt ctcagcttgt 300

caggtagcga gttgttgttt tatatttatt cgatttcatc tgtgttgcat gtcttgttcg 360

tgctgtgcgt tctttctggg ccgcgctgtc gggtcgcatg ggctagctgt actcatgtta 420

gtcatgccgg tccgaccttg ttcgaggaag gccccacact gagcgtgccc tctttctaca 480

ccccttgtgc agaaattaga tagaaagcaa tgctgctgca ggccttcctg ttcctgctgg 540

ccggcttcgc cgccaagatc agcgcctcca tgacgaacga gacgtccgac cgccccctgg 600

tgcacttcac ccccaacaag ggctggatga acgaccccaa cggcctgtgg tacgacgaga 660

aggacgccaa gtggcacctg tacttccagt acaacccgaa cgacaccgtc tgggggacgc 720

ccttgttctg gggccacgcc acgtccgacg acctgaccaa ctgggaggac cagcccatcg 780

ccatcgcccc gaagcgcaac gactccggcg ccttctccgg ctccatggtg gtggactaca 840

acaacacctc cggcttcttc aacgacacca tcgacccgcg ccagcgctgc gtggccatct 900

ggacctacaa caccccggag tccgaggagc agtacatctc ctacagcctg gacggcggct 960

acaccttcac cgagtaccag aagaaccccg tgctggccgc caactccacc cagttccgcg 1020

acccgaaggt cttctggtac gagccctccc agaagtggat catgaccgcg gccaagtccc 1080

aggactacaa gatcgagatc tactcctccg acgacctgaa gtcctggaag ctggagtccg 1140

cgttcgccaa cgagggcttc ctcggctacc agtacgagtg ccccggcctg atcgaggtcc 1200

ccaccgagca ggaccccagc aagtcctact gggtgatgtt catctccatc aaccccggcg 1260

ccccggccgg cggctccttc aaccagtact tcgtcggcag cttcaacggc acccacttcg 1320

aggccttcga caaccagtcc cgcgtggtgg acttcggcaa ggactactac gccctgcaga 1380

ccttcttcaa caccgacccg acctacggga gcgccctggg catcgcgtgg gcctccaact 1440

gggagtactc cgccttcgtg cccaccaacc cctggcgctc ctccatgtcc ctcgtgcgca 1500

agttctccct caacaccgag taccaggcca acccggagac ggagctgatc aacctgaagg 1560

ccgagccgat cctgaacatc agcaacgccg gcccctggag ccggttcgcc accaacacca 1620

cgttgacgaa ggccaacagc tacaacgtcg acctgtccaa cagcaccggc accctggagt 1680

tcgagctggt gtacgccgtc aacaccaccc agacgatctc caagtccgtg ttcgcggacc 1740

tctccctctg gttcaagggc ctggaggacc ccgaggagta cctccgcatg ggcttcgagg 1800

tgtccgcgtc ctccttcttc ctggaccgcg ggaacagcaa ggtgaagttc gtgaaggaga 1860

acccctactt caccaaccgc atgagcgtga acaaccagcc cttcaagagc gagaacgacc 1920

tgtcctacta caaggtgtac ggcttgctgg accagaacat cctggagctg tacttcaacg 1980

acggcgacgt cgtgtccacc aacacctact tcatgaccac cgggaacgcc ctgggctccg 2040

tgaacatgac gacgggggtg gacaacctgt tctacatcga caagttccag gtgcgcgagg 2100

tcaagtgacc gacacgcccc cggcccaggt ccagttctcc tgggtcttcc agaggcccgt 2160

cgccatgtaa agtggcagag attggcgcct gattcgattt ggatccaagg atctccaatc 2220

ggtgatgggg actgagtgcc caactaccac ccttgcacta tcgtcctcgc actatttatt 2280

cccaccttct gctcgccctg ccgggcgatt gcgggcgttt ctgcccttga cgtatcaatt 2340

tcgcccctgc tggcgcgagg attcttcatt ctaataagaa ctcactcccg ccagctctgt 2400

acttttcctg cggggccccc gcatggcttg ttcccaatgc ttgctcgatc gacggcgccc 2460

attgcccacg gcgctgccgc atccatgtga agaaacacgg aagagtgcga agactggaag 2520

tgaattaaga gtataagaag aggtaccaag ggattctcag gtgctcttag gaacggcttt 2580

tccttcgcca agagaaactg ctactgctcg tgtcgccacg gtggtcaagc cgccccatct 2640

gcgatccacc aggcccatcc gcggactcgc gatcagcctg ctggatccgg actgccgacc 2700

tgaccgctcg catccaccat tacaaccctc caattggaca ccactcccac gtcctaaagt 2760

tcaccatgca agctgatcga tcgcattcgc cgatgcactc gcctgccaca gaggtgtgcg 2820

cttcggacta gcgtgcaggc gccccgaggc caccagcatg caccgatgga agcgggcacg 2880

gccgctgctc caggtcgctg gctcgctcag acccatagca acctccgctg cgtccctaaa 2940

tgtcacacag agcgtctttg atgggtacgg atgggagaga atctgattgg gcattgctgg 3000

tgcagtgcag gaagatggca agtgcacagt cagtcatgct gtacaaactg gtgcctcgta 3060

gtattgactc gtatagtgca tagtatcatg catggtcgtt acttgcaa 3108

<210> 106
<211> 559
<212> DNA
<213> Chlorella luteoviridis

<400> 106
tgttgaagaa tgagccggcg acttatagga agtggcgtgg ttaaggaatt ttccgaagcc 60

caagcgaaag caagttttaa aaatagcgat atttgtcact ttttatggac ccgaacccgg 120

gtgatctaac cgtgaccagg atgaagcttg ggtaacacca agtgaaggtc cgaactcttc 180

gatctttaaa aatcgtgaga tgagttgcgg ttagtaggtg aaatgccaat cgaactcgga 240

gctagctggt tctccccgaa atgtgttgag gcgcagcgat gaatgacaaa acaaatagta 300

cggtgtaggg gtaaagcact gtttcggtgc gggctgcgaa agcggtacca aatcgtggca 360

aactcagaat actacgcttg tataccattc atcagtgaga ctgtggggga taagctccat 420

agtcaagagg gaaacagccc agatcaccag ttaaggcccc aaaatgacag ctaagtggca 480

aaggaggtga aagtgcagaa acaaccagga ggtttgccca gaagcagcca tcctttaaag 540

agtgcgtaat agctcactg 559

<210> 107
<211> 1841
<212> DNA
<213> Cuphea palustris

<400> 107
gaattccttt cttgcgctat gacacttcca gcaaaaggta gggcgggctg cgagacggct 60

tcccggcgct gcatgcaaca ccgatgatgc ttcgaccccc cgaagctcct tcggggctgc 120

atgggcgctc cgatgccgct ccagggcgag cgctgtttaa atagccaggc ccccgattgc 180

aaagacatta tagcgagcta ccaaagccat attcaaacac ctagatcact accacttcta 240

cacaggccac tcgagcttgt gatcgcactc cgctaagggg gcgcctcttc ctcttcgttt 300

cagtcacaac ccgcaaacac tagtatggcc accgcatcca ctttctcggc gttcaatgcc 360

cgctgcggcg acctgcgtcg ctcggcgggc tccgggcccc ggcgcccagc gaggcccctc 420

cccgtgcgcg ggcgcgccag catgctgctg tcggcggtga ccacggtctt cggcgtggcc 480

gagaagcagt ggcccatgct ggaccgcaag tccaagcgcc ccgacatgct ggtcgagccc 540

ctgggcgtgg accgcatcgt ctacgacggc gtgagcttcc gccagtcgtt ctccatccgc 600

agctacgaga tcggcgccga ccgcaccgcc tcgatcgaga cgctgatgaa catgttccag 660

gagacctccc tgaaccactg caagatcatc ggcctgctga acgacggctt cggccgcacg 720

cccgagatgt gcaagcgcga cctgatctgg gtcgtgacca agatgcagat cgaggtgaac 780

cgctacccca cgtggggcga caccatcgag gtcaacacgt gggtgagcgc ctcgggcaag 840

cacggcatgg gccgcgactg gctgatctcc gactgccaca ccggcgagat cctgatccgc 900

gcgacgagcg tctgggcgat gatgaaccag aagacccgcc gcctgtcgaa gatcccctac 960

gaggtgcgcc aggagatcga gccccagttc gtcgactccg cccccgtgat cgtggacgac 1020

cgcaagttcc acaagctgga cctgaagacg ggcgacagca tctgcaacgg cctgacccca 1080

cgctggacgg acctggacgt gaaccagcac gtcaacaacg tgaagtacat cggctggatc 1140

ctgcagtcgg tccccaccga ggtgttcgag acgcaggagc tgtgcggcct gaccctggag 1200

taccgccgcg agtgcggccg cgactccgtg ctggagagcg tcacggccat ggacccctcg 1260

aaggagggcg accgctccct gtaccagcac ctgctgcgcc tggaggacgg cgcggacatc 1320

gtgaagggcc gcaccgagtg gcgccccaag aacgccggcg ccaagggcgc catcctgacg 1380

ggcaagacca gcaacggcaa ctcgatctcc tgactcgagt taattaactc gaggcagcag 1440

cagctcggat agtatcgaca cactctggac gctggtcgtg tgatggactg ttgccgccac 1500

acttgctgcc ttgacctgtg aatatccctg ccgcttttat caaacagcct cagtgtgttt 1560

gatcttgtgt gtacgcgctt ttgcgagttg ctagctgctt gtgctatttg cgaataccac 1620

ccccagcatc cccttccctc gtttcatatc gcttgcatcc caaccgcaac ttatctacgc 1680

tgtcctgcta tccctcagcg ctgctcctgc tcctgctcac tgcccctcgc acagccttgg 1740

tttgggctcc gcctgtattc tcctggtact gcaacctgta aaccagcact gcaatgctga 1800

tgcacgggaa gtagtgggat gggaacacaa atggaaagct t 1841

<210> 108
<211> 1010
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic polynucleotide

<400> 108
ggcgcgccca gctgcccgac tggagcatgc tgctggccgc gatcaccacc ctgttcctgg 60

cggccgagaa gcagtggatg atgctggact ggaagcccaa gcgccccgac atgctggtgg 120

acccccttcgg cctgggccgc ttcgtgcagg acggcctggt gttccgcaac aacttcagca 180

tccgcagcta cgagatcggc gcggaccgca ccgccagcat cgagaccctg atgaaccacc 240

tgcaggagac cgccctgaac cacgtgaaga gcgtgggcct gctggaggac ggcctgggca 300

gcacccgcga gatgagcctg cgcaacctga tctgggtggt gaccaagatg caggtggcgg 360

tggaccgcta ccccacctgg ggcgacgagg tgcaggtgag cagctgggcg accgccatcg 420

gcaagaacgg catgcgccgc gagtggatcg tgaccgactt ccgcaccggc gagaccctgc 480

tgcgcgccac cagcgtgtgg gtgatgatga acaagctgac ccgccgcatc agcaagatcc 540

ccgaggaggt gtggcacgag atcggcccca gcttcatcga cgcgcccccc ctgcccaccg 600

tggaggacga cggccgcaag ctgacccgct tcgacgagag cagcgccgac ttcatccgca 660

agggcctgac cccccgctgg agcgacctgg acatcaacca gcacgtgaac aacgtgaagt 720

acatcggctg gctgctggag agcgcgcccc ccgagatcca cgagagccac gagatcgcca 780

gcctgaccct ggagtaccgc cgcgagtgcg gccgcgacag cgtgctgaac agcgccacca 840

aggtgagcga cagcagccag ctgggcaaga gcgccgtgga gtgcaaccac ctggtgcgcc 900

tgcagaacgg cggcgagatc gtgaagggcc gcaccgtgtg gcgccccaag cgccccctgt 960

acaacgacgg cgccgtggtg gacgtgcccg ccaagaccag ctgactcgag 1010

<210> 109
<211> 5472
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic polynucleotide

<400> 109
ggtacccttt cttgcgctat gacacttcca gcaaaaggta gggcgggctg cgagacggct 60

tcccggcgct gcatgcaaca ccgatgatgc ttcgaccccc cgaagctcct tcggggctgc 120

atgggcgctc cgatgccgct ccagggcgag cgctgtttaa atagccaggc ccccgattgc 180

aaagacatta tagcgagcta ccaaagccat attcaaacac ctagatcact accacttcta 240

cacaggccac tcgagcttgt gatcgcactc cgctaagggg gcgcctcttc ctcttcgttt 300

cagtcacaac ccgcaaactc tagaatatca atgctgctgc aggccttcct gttcctgctg 360

gccggcttcg ccgccaagat cagcgcctcc atgacgaacg agacgtccga ccgccccctg 420

gtgcacttca cccccaacaa gggctggatg aacgacccca acggcctgtg gtacgacgag 480

aaggacgcca agtggcacct gtacttccag tacaacccga acgacaccgt ctgggggacg 540

cccttgttct ggggccacgc cacgtccgac gacctgacca actgggagga ccagcccatc 600

gccatcgccc cgaagcgcaa cgactccggc gccttctccg gctccatggt ggtggactac 660

aacaacacct ccggcttctt caacgacacc atcgacccgc gccagcgctg cgtggccatc 720

tggacctaca acaccccgga gtccgaggag cagtacatct cctacagcct ggacggcggc 780

tacaccttca ccgagtacca gaagaacccc gtgctggccg ccaactccac ccagttccgc 840

gacccgaagg tcttctggta cgagccctcc cagaagtgga tcatgaccgc ggccaagtcc 900

caggactaca agatcgagat ctactcctcc gacgacctga agtcctggaa gctggagtcc 960

gcgttcgcca acgagggctt cctcggctac cagtacgagt gccccggcct gatcgaggtc 1020

cccaccgagc aggaccccag caagtcctac tgggtgatgt tcatctccat caaccccggc 1080

gccccggccg gcggctcctt caaccagtac ttcgtcggca gcttcaacgg cacccacttc 1140

gaggccttcg acaaccagtc ccgcgtggtg gacttcggca aggactacta cgccctgcag 1200

accttcttca acaccgaccc gacctacggg agcgccctgg gcatcgcgtg ggcctccaac 1260

tgggagtact ccgccttcgt gcccaccaac ccctggcgct cctccatgtc cctcgtgcgc 1320

aagttctccc tcaacaccga gtaccaggcc aacccggaga cggagctgat caacctgaag 1380

gccgagccga tcctgaacat cagcaacgcc ggcccctgga gccggttcgc caccaacacc 1440

acgttgacga aggccaacag ctacaacgtc gacctgtcca acagcaccgg caccctggag 1500

ttcgagctgg tgtacgccgt caacaccacc cagacgatct ccaagtccgt gttcgcggac 1560

ctctccctct ggttcaaggg cctggaggac cccgaggagt acctccgcat gggcttcgag 1620

gtgtccgcgt cctccttctt cctggaccgc gggaacagca aggtgaagtt cgtgaaggag 1680

aacccctact tcaccaaccg catgagcgtg aacaaccagc ccttcaagag cgagaacgac 1740

ctgtcctact acaaggtgta cggcttgctg gaccagaaca tcctggagct gtacttcaac 1800

gacggcgacg tcgtgtccac caacacctac ttcatgacca ccgggaacgc cctgggctcc 1860

gtgaacatga cgacgggggt ggacaacctg ttctacatcg acaagttcca ggtgcgcgag 1920

gtcaagtgac aattggcagc agcagctcgg atagtatcga cacactctgg acgctggtcg 1980

tgtgatggac tgttgccgcc acacttgctg ccttgacctg tgaatatccc tgccgctttt 2040

atcaaacagc ctcagtgtgt ttgatcttgt gtgtacgcgc ttttgcgagt tgctagctgc 2100

ttgtgctatt tgcgaatacc acccccagca tcccccttccc tcgtttcata tcgcttgcat 2160

cccaaccgca acttatctac gctgtcctgc tatccctcag cgctgctcct gctcctgctc 2220

actgcccctc gcacagcctt ggtttgggct ccgcctgtat tctcctggta ctgcaacctg 2280

taaaccagca ctgcaatgct gatgcacggg aagtagtggg atgggaacac aaatggagga 2340

tcccgcgtct cgaacagagc gcgcagagga acgctgaagg tctcgcctct gtcgcacctc 2400

agcgcggcat acaccacaat aaccacctga cgaatgcgct tggttcttcg tccattagcg 2460

aagcgtccgg ttcacacacg tgccacgttg gcgaggtggc aggtgacaat gatcggtgga 2520

gctgatggtc gaaacgttca cagcctaggg atatcgaatt ccgcctgcaa cgcaagggca 2580

gccacagccg ctcccacccg ccgctgaacc gacacgtgct tgggcgcctg ccgcctgcct 2640

gccgcatgct tgtgctggtg aggctgggca gtgctgccat gctgattgag gcttggttca 2700

tcgggtggaa gcttatgtgt gtgctgggct tgcatgccgg gcaatgcgca tggtggcaag 2760

agggcggcag cacttgctgg agctgccgcg gtgcctccag gtggttcaat cgcggcagcc 2820

agagggattt cagatgatcg cgcgtacagg ttgagcagca gtgtcagcaa aggtagcagt 2880

ttgccagaat gatcggttca gctgttaatc aatgccagca agagaagggg tcaagtgcaa 2940

acacgggcat gccacagcac gggcaccggg gagtggaatg gcaccaccaa gtgtgtgcga 3000

gccagcatcg ccgcctggct gtttcagcta caacggcagg agtcatccaa cgtaaccatg 3060

agctgatcaa cactgcaatc atcgggcggg cgtgatgcaa gcatgcctgg cgaagacaca 3120

tggtgtgcgg atgctgccgg ctgctgcctg ctgcgcacgc cgttgagttg gcagcaggct 3180

cagccatgca ctggatggca gctgggctgc cactgcaatg tggtggatag gatgcaagtg 3240

gagcgaatac caaaccctct ggctgcttgc tgggttgcat ggcatcgcac catcagcagg 3300

agcgcatgcg aagggactgg ccccatgcac gccatgccaa accggagcgc accgagtgtc 3360

cacactgtca ccaggcccgc aagctttgca gaaccatgct catggacgca tgtagcgctg 3420

acgtcccttg acggcgctcc tctcgggtgt gggaaacgca atgcagcaca ggcagcagag 3480

gcggcggcag cagagcggcg gcagcagcgg cgggggccac ccttcttgcg gggtcgcgcc 3540

ccagccagcg gtgatgcgct gatcccaaac gagttcacat tcatttgcat gcctggagaa 3600

gcgaggctgg ggcctttggg ctggtgcagc ccgcaatgga atgcgggacc gccaggctag 3660

cagcaaaggc gcctccccta ctccgcatcg atgttccata gtgcattgga ctgcatttgg 3720

gtggggcggc cggctgtttc tttcgtgttg caaaacgcgc cagctcagca acctgtcccg 3780

tgggtccccc gtgccgatga aatcgtgtgc acgccgatca gctgattgcc cggctcgcga 3840

agtaggcgcc ctcctttctg ctcgccctct ctccgtcccg ccactagtat ggccaccgca 3900

tccactttct cggcgttcaa tgcccgctgc ggcgacctgc gtcgctcggc gggctccggg 3960

ccccggcgcc cagcgaggcc cctccccgtg cgcgggcgcg cccagctgcc cgactggagc 4020

cgcctgctga ccgccatcac caccgtgttc gtgaagtcca agcgccccga catgcacgac 4080

cgcaagtcca agcgccccga catgctggtg gacagcttcg gcctggagtc caccgtgcag 4140

gacggcctgg tgttccgcca gtccttctcc atccgctcct acgagatcgg caccgaccgc 4200

accgccagca tcgagaccct gatgaaccac ctgcaggaga cctccctgaa ccactgcaag 4260

agcaccggca tcctgctgga cggcttcggc cgcaccctgg agatgtgcaa gcgcgacctg 4320

atctgggtgg tgatcaagat gcagatcaag gtgaaccgct accccgcctg gggcgacacc 4380

gtggagatca acacccgctt cagccgcctg ggcaagatcg gcatgggccg cgactggctg 4440

atctccgact gcaacaccgg cgagatcctg gtgcgcgcca ccagcgccta cgccatgatg 4500

aaccagaaga cccgccgcct gtccaagctg ccctacgagg tgcaccagga gatcgtgccc 4560

ctgttcgtgg acagccccgt gatcgaggac tccgacctga aggtgcacaa gttcaaggtg 4620

aagaccggcg acagcatcca gaagggcctg acccccggct ggaacgacct ggacgtgaac 4680

cagcacgtgt ccaacgtgaa gtacatcggc tggatcctgg agagcatgcc caccgaggtg 4740

ctggagaccc aggagctgtg ctccctggcc ctggagtacc gccgcgagtg cggccgcgac 4800

tccgtgctgg agagcgtgac cgccatggac cccagcaagg tgggcgtgcg ctcccagtac 4860

cagcacctgc tgcgcctgga ggacggcacc gccatcgtga acggcgccac cgagtggcgc 4920

cccaagaacg ccggcgccaa cggcgccatc tccaccggca agaccagcaa cggcaactcc 4980

gtgtccatgg actacaagga ccacgacggc gactacaagg accacgacat cgactacaag 5040

gacgacgacg acaagtgact cgaggcagca gcagctcgga tagtatcgac acactctgga 5100

cgctggtcgt gtgatggact gttgccgcca cacttgctgc cttgacctgt gaatatccct 5160

gccgctttta tcaaacagcc tcagtgtgtt tgatcttgtg tgtacgcgct tttgcgagtt 5220

gctagctgct tgtgctattt gcgaatacca cccccagcat ccccttccct cgtttcatat 5280

cgcttgcatc ccaaccgcaa cttatctacg ctgtcctgct atccctcagc gctgctcctg 5340

ctcctgctca ctgcccctcg cacagccttg gtttgggctc cgcctgtatt ctcctggtac 5400

tgcaacctgt aaaccagcac tgcaatgctg atgcacggga agtagtggga tgggaacaca 5460

aatggaaagc tt 5472

<210> 110
<211> 5451
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic polynucleotide

<400> 110
ggtacccttt cttgcgctat gacacttcca gcaaaaggta gggcgggctg cgagacggct 60

tcccggcgct gcatgcaaca ccgatgatgc ttcgaccccc cgaagctcct tcggggctgc 120

atgggcgctc cgatgccgct ccagggcgag cgctgtttaa atagccaggc ccccgattgc 180

aaagacatta tagcgagcta ccaaagccat attcaaacac ctagatcact accacttcta 240

cacaggccac tcgagcttgt gatcgcactc cgctaagggg gcgcctcttc ctcttcgttt 300

cagtcacaac ccgcaaactc tagaatatca atgctgctgc aggccttcct gttcctgctg 360

gccggcttcg ccgccaagat cagcgcctcc atgacgaacg agacgtccga ccgccccctg 420

gtgcacttca cccccaacaa gggctggatg aacgacccca acggcctgtg gtacgacgag 480

aaggacgcca agtggcacct gtacttccag tacaacccga acgacaccgt ctgggggacg 540

cccttgttct ggggccacgc cacgtccgac gacctgacca actgggagga ccagcccatc 600

gccatcgccc cgaagcgcaa cgactccggc gccttctccg gctccatggt ggtggactac 660
aacaacacct ccggcttctt caacgacacc atcgacccgc gccagcgctg cgtggccatc 720
tggacctaca acaccccgga gtccgaggag cagtacatct cctacagcct ggacggcggc 780
tacaccttca ccgagtacca gaagaacccc gtgctggccg ccaactccac ccagttccgc 840
gacccgaagg tcttctggta cgagccctcc cagaagtgga tcatgaccgc ggccaagtcc 900
caggactaca agatcgagat ctactcctcc gacgacctga agtcctggaa gctggagtcc 960
gcgttcgcca acgagggctt cctcggctac cagtacgagt gccccggcct gatcgaggtc 1020
cccaccgagc aggaccccag caagtcctac tgggtgatgt tcatctccat caaccccggc 1080
gccccggccg gcggctcctt caaccagtac ttcgtcggca gcttcaacgg cacccacttc 1140
gaggccttcg acaaccagtc ccgcgtggtg gacttcggca aggactacta cgccctgcag 1200
accttcttca acaccgaccc gacctacggg agcgccctgg gcatcgcgtg ggcctccaac 1260
tgggagtact ccgccttcgt gcccaccaac ccctggcgct cctccatgtc cctcgtgcgc 1320
aagttctccc tcaacaccga gtaccaggcc aacccggaga cggagctgat caacctgaag 1380
gccgagccga tcctgaacat cagcaacgcc ggcccctgga gccggttcgc caccaacacc 1440
acgttgacga aggccaacag ctacaacgtc gacctgtcca acagcaccgg caccctggag 1500
ttcgagctgg tgtacgccgt caacaccacc cagacgatct ccaagtccgt gttcgcggac 1560
ctctccctct ggttcaaggg cctggaggac cccgaggagt acctccgcat gggcttcgag 1620
gtgtccgcgt cctccttctt cctggaccgc gggaacagca aggtgaagtt cgtgaaggag 1680
aaccccctact tcaccaaccg catgagcgtg aacaaccagc ccttcaagag cgagaacgac 1740
ctgtcctact acaaggtgta cggcttgctg gaccagaaca tcctggagct gtacttcaac 1800
gacggcgacg tcgtgtccac caacacctac ttcatgacca ccgggaacgc cctgggctcc 1860
gtgaacatga cgacgggggt ggacaacctg ttctacatcg acaagttcca ggtgcgcgag 1920
gtcaagtgac aattggcagc agcagctcgg atagtatcga cacactctgg acgctggtcg 1980
tgtgatggac tgttgccgcc acacttgctg ccttgacctg tgaatatccc tgccgctttt 2040
atcaaacagc ctcagtgtgt ttgatcttgt gtgtacgcgc ttttgcgagt tgctagctgc 2100
ttgtgctatt tgcgaatacc acccccagca tccccttccc tcgtttcata tcgcttgcat 2160
cccaaccgca acttatctac gctgtcctgc tatccctcag cgctgctcct gctcctgctc 2220
actgcccctc gcacagcctt ggtttgggct ccgcctgtat tctcctggta ctgcaacctg 2280
taaaccagca ctgcaatgct gatgcacggg aagtagtggg atgggaacac aaatggagga 2340
tcccgcgtct cgaacagagc gcgcagagga acgctgaagg tctcgcctct gtcgcacctc 2400
agcgcggcat acaccacaat aaccacctga cgaatgcgct tggttcttcg tccattagcg 2460
aagcgtccgg ttcacacacg tgccacgttg gcgaggtggc aggtgacaat gatcggtgga 2520
gctgatggtc gaaacgttca cagcctaggg atatcgaatt ccgcctgcaa cgcaagggca 2580
gccacagccg ctcccacccg ccgctgaacc gacacgtgct tgggcgcctg ccgcctgcct 2640

gccgcatgct tgtgctggtg aggctgggca gtgctgccat gctgattgag gcttggttca 2700
tcgggtggaa gcttatgtgt gtgctgggct tgcatgccgg gcaatgcgca tggtggcaag 2760
agggcggcag cacttgctgg agctgccgcg gtgcctccag gtggttcaat cgcggcagcc 2820
agagggattt cagatgatcg cgcgtacagg ttgagcagca gtgtcagcaa aggtagcagt 2880
ttgccagaat gatcggttca gctgttaatc aatgccagca agagaagggg tcaagtgcaa 2940
acacgggcat gccacagcac gggcaccggg gagtggaatg gcaccaccaa gtgtgtgcga 3000
gccagcatcg ccgcctggct gtttcagcta caacggcagg agtcatccaa cgtaaccatg 3060
agctgatcaa cactgcaatc atcgggcggg cgtgatgcaa gcatgcctgg cgaagacaca 3120
tggtgtgcgg atgctgccgg ctgctgcctg ctgcgcacgc cgttgagttg gcagcaggct 3180
cagccatgca ctggatggca gctgggctgc cactgcaatg tggtggatag gatgcaagtg 3240
gagcgaatac caaaccctct ggctgcttgc tgggttgcat ggcatcgcac catcagcagg 3300
agcgcatgcg aagggactgg ccccatgcac gccatgccaa accggagcgc accgagtgtc 3360
cacactgtca ccaggcccgc aagctttgca gaaccatgct catggacgca tgtagcgctg 3420
acgtcccttg acggcgctcc tctcgggtgt gggaaacgca atgcagcaca ggcagcagag 3480
gcggcggcag cagagcggcg gcagcagcgg cgggggccac ccttcttgcg gggtcgcgcc 3540
ccagccagcg gtgatgcgct gatcccaaac gagttcacat tcatttgcat gcctggagaa 3600
gcgaggctgg ggcctttggg ctggtgcagc ccgcaatgga atgcgggacc gccaggctag 3660
cagcaaaggc gcctccccta ctccgcatcg atgttccata gtgcattgga ctgcatttgg 3720
gtggggcggc cggctgtttc tttcgtgttg caaaacgcgc cagctcagca acctgtcccg 3780
tgggtccccc gtgccgatga aatcgtgtgc acgccgatca gctgattgcc cggctcgcga 3840
agtaggcgcc ctcctttctg ctcgccctct ctccgtcccg ccactagtat gacgttcggg 3900
gtcgccctcc cggccatggg ccgcggtgtc tcccttcccc ggcccagggt cgcggtgcgc 3960
gcccagtcgg cgagtcaggt tttggagagc gggcgcgccc ccgactggtc catgctgttc 4020
gccgtgatca ccaccatctt cagcgccgcc gagaagcagt ggaccaacct ggagtggaag 4080
cccaagccca agctgcccca gctgctggac gaccacttcg gcctgcacgg cctggtgttc 4140
cgccgcacct tcgccatccg ctcctacgag gtgggccccg accgcagcac ctccatcctg 4200
gccgtgatga accacatgca ggaggccacc ctgaaccacg ccaagagcgt gggcatcctg 4260
ggcgacggct tcggcaccac cctggagatg tccaagcgcg acctgatgtg ggtggtgcgc 4320
cgcacccacg tggccgtgga gcgctacccc acctggggcg acaccgtgga ggtggagtgc 4380
tggatcggcg ccagcggcaa caacggcatg cgccgcgact tcctggtgcg cgactgcaag 4440
accggcgaga tcctgacccg ctgcacctcc ctgagcgtgc tgatgaacac ccgcacccgc 4500
cgcctgagca ccatccccga cgaggtgcgc ggcgagatcg gccccgcctt catcgacaac 4560
gtggccgtga aggacgacga gatcaagaag ctgcagaagc tgaacgactc caccgccgac 4620
tacatccagg gcggcctgac cccccgctgg aacgacctgg acgtgaacca gcacgtgaac 4680

aacctgaagt acgtggcctg ggtgttcgag accgtgcccg acagcatctt cgagtcccac 4740

cacatcagct ccttcaccct ggagtaccgc cgcgagtgca cccgcgactc cgtgctgcgc 4800

agcctgacca ccgtgagcgg cggcagctcc gaggccggcc tggtgtgcga ccacctgctg 4860

cagctggagg gcggcagcga ggtgctgcgc gcccgcaccg agtggcgccc caagctgacc 4920

gactccttcc gcggcatcag cgtgatcccc gccgagcccc gcgtgatgga ctacaaggac 4980

cacgacggcg actacaagga ccacgacatc gactacaagg acgacgacga caagtgactc 5040

gaggcagcag cagctcggat agtatcgaca cactctggac gctggtcgtg tgatggactg 5100

ttgccgccac acttgctgcc ttgacctgtg aatatccctg ccgcttttat caaacagcct 5160

cagtgtgttt gatcttgtgt gtacgcgctt ttgcgagttg ctagctgctt gtgctatttg 5220

cgaataccac ccccagcatc cccttccctc gtttcatatc gcttgcatcc caaccgcaac 5280

ttatctacgc tgtcctgcta tccctcagcg ctgctcctgc tcctgctcac tgcccctcgc 5340

acagccttgg tttgggctcc gcctgtattc tcctggtact gcaacctgta aaccagcact 5400

gcaatgctga tgcacgggaa gtagtgggat gggaacacaa atggaaagct t 5451

<210> 111
<211> 5454
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic polynucleotide

<400> 111
ggtacccttt cttgcgctat gacacttcca gcaaaaggta gggcgggctg cgagacggct 60

tcccggcgct gcatgcaaca ccgatgatgc ttcgaccccc cgaagctcct tcggggctgc 120

atgggcgctc cgatgccgct ccagggcgag cgctgtttaa atagccaggc ccccgattgc 180

aaagacatta tagcgagcta ccaaagccat attcaaacac ctagatcact accacttcta 240

cacaggccac tcgagcttgt gatcgcactc cgctaagggg gcgcctcttc ctcttcgttt 300

cagtcacaac ccgcaaactc tagaatatca atgctgctgc aggccttcct gttcctgctg 360

gccggcttcg ccgccaagat cagcgcctcc atgacgaacg agacgtccga ccgccccctg 420

gtgcacttca cccccaacaa gggctggatg aacgacccca acggcctgtg gtacgacgag 480

aaggacgcca agtggcacct gtacttccag tacaacccga acgacaccgt ctgggggacg 540

cccttgttct ggggccacgc cacgtccgac gacctgacca actgggagga ccagcccatc 600

gccatcgccc cgaagcgcaa cgactccggc gccttctccg gctccatggt ggtggactac 660

aacaacacct ccggcttctt caacgacacc atcgacccgc gccagcgctg cgtggccatc 720

tggacctaca acaccccgga gtccgaggag cagtacatct cctacagcct ggacggcggc 780

tacaccttca ccgagtacca gaagaacccc gtgctggccg ccaactccac ccagttccgc 840

gacccgaagg tcttctggta cgagccctcc cagaagtgga tcatgaccgc ggccaagtcc 900

caggactaca agatcgagat ctactcctcc gacgacctga agtcctggaa gctggagtcc 960

gcgttcgcca acgagggctt cctcggctac cagtacgagt gccccggcct gatcgaggtc 1020

cccaccgagc aggaccccag caagtcctac tgggtgatgt tcatctccat caaccccggc 1080

gccccggccg gcggctcctt caaccagtac ttcgtcggca gcttcaacgg cacccacttc 1140

gaggccttcg acaaccagtc ccgcgtggtg gacttcggca aggactacta cgccctgcag 1200

accttcttca acaccgaccc gacctacggg agcgccctgg gcatcgcgtg ggcctccaac 1260

tgggagtact ccgccttcgt gcccaccaac ccctggcgct cctccatgtc cctcgtgcgc 1320

aagttctccc tcaacaccga gtaccaggcc aacccggaga cggagctgat caacctgaag 1380

gccgagccga tcctgaacat cagcaacgcc ggcccctgga gccggttcgc caccaacacc 1440

acgttgacga aggccaacag ctacaacgtc gacctgtcca acagcaccgg caccctggag 1500

ttcgagctgg tgtacgccgt caacaccacc cagacgatct ccaagtccgt gttcgcggac 1560

ctctccctct ggttcaaggg cctggaggac cccgaggagt acctccgcat gggcttcgag 1620

gtgtccgcgt cctccttctt cctggaccgc gggaacagca aggtgaagtt cgtgaaggag 1680

aacccctact tcaccaaccg catgagcgtg aacaaccagc ccttcaagag cgagaacgac 1740

ctgtcctact acaaggtgta cggcttgctg gaccagaaca tcctggagct gtacttcaac 1800

gacggcgacg tcgtgtccac caacacctac ttcatgacca ccgggaacgc cctgggctcc 1860

gtgaacatga cgacgggggt ggacaacctg ttctacatcg acaagttcca ggtgcgcgag 1920

gtcaagtgac aattggcagc agcagctcgg atagtatcga cacactctgg acgctggtcg 1980

tgtgatggac tgttgccgcc acacttgctg ccttgacctg tgaatatccc tgccgctttt 2040

atcaaacagc ctcagtgtgt ttgatcttgt gtgtacgcgc ttttgcgagt tgctagctgc 2100

ttgtgctatt tgcgaatacc acccccagca tcccccttccc tcgtttcata tcgcttgcat 2160

cccaaccgca acttatctac gctgtcctgc tatccctcag cgctgctcct gctcctgctc 2220

actgcccctc gcacagcctt ggtttgggct ccgcctgtat tctcctggta ctgcaacctg 2280

taaaccagca ctgcaatgct gatgcacggg aagtagtggg atgggaacac aaatggagga 2340

tcccgcgtct cgaacagagc gcgcagagga acgctgaagg tctcgcctct gtcgcacctc 2400

agcgcggcat acaccacaat aaccacctga cgaatgcgct tggttcttcg tccattagcg 2460

aagcgtccgg ttcacacacg tgccacgttg gcgaggtggc aggtgacaat gatcggtgga 2520

gctgatggtc gaaacgttca cagcctaggg atatcgaatt ccgcctgcaa cgcaagggca 2580

gccacagccg ctcccacccg ccgctgaacc gacacgtgct tgggcgcctg ccgcctgcct 2640

gccgcatgct tgtgctggtg aggctgggca gtgctgccat gctgattgag gcttggttca 2700

tcgggtggaa gcttatgtgt gtgctgggct tgcatgccgg gcaatgcgca tggtggcaag 2760

agggcggcag cacttgctgg agctgccgcg gtgcctccag gtggttcaat cgcggcagcc 2820

agagggattt cagatgatcg cgcgtacagg ttgagcagca gtgtcagcaa aggtagcagt 2880

ttgccagaat gatcggttca gctgttaatc aatgccagca agagaagggg tcaagtgcaa 2940

acacgggcat gccacagcac gggcaccggg gagtggaatg gcaccaccaa gtgtgtgcga 3000

gccagcatcg ccgcctggct gtttcagcta caacggcagg agtcatccaa cgtaaccatg 3060

agctgatcaa cactgcaatc atcgggcggg cgtgatgcaa gcatgcctgg cgaagacaca 3120

tggtgtgcgg atgctgccgg ctgctgcctg ctgcgcacgc cgttgagttg gcagcaggct 3180

cagccatgca ctggatggca gctgggctgc cactgcaatg tggtggatag gatgcaagtg 3240

gagcgaatac caaaccctct ggctgcttgc tgggttgcat ggcatcgcac catcagcagg 3300

agcgcatgcg aagggactgg ccccatgcac gccatgccaa accggagcgc accgagtgtc 3360

cacactgtca ccaggcccgc aagctttgca gaaccatgct catggacgca tgtagcgctg 3420

acgtcccttg acggcgctcc tctcgggtgt gggaaacgca atgcagcaca ggcagcagag 3480

gcggcggcag cagagcggcg gcagcagcgg cgggggccac ccttcttgcg gggtcgcgcc 3540

ccagccagcg gtgatgcgct gatcccaaac gagttcacat tcatttgcat gcctggagaa 3600

gcgaggctgg ggcctttggg ctggtgcagc ccgcaatgga atgcgggacc gccaggctag 3660

cagcaaaggc gcctccccta ctccgcatcg atgttccata gtgcattgga ctgcatttgg 3720

gtggggcggc cggctgtttc tttcgtgttg caaaacgcgc cagctcagca acctgtcccg 3780

tgggtccccc gtgccgatga aatcgtgtgc acgccgatca gctgattgcc cggctcgcga 3840

agtaggcgcc ctcctttctg ctcgccctct ctccgtcccg ccactagtat ggccaccgca 3900

tccactttct cggcgttcaa tgcccgctgc ggcgacctgc gtcgctcggc gggctccggg 3960

ccccggcgcc cagcgaggcc cctccccgtg cgcgggcgcg cccccgactg gtccatgctg 4020

ttcgccgtga tcaccaccat cttctccgcc gccgagaagc agtggaccaa cctggagtgg 4080

aagcccaagc ccaacccccc ccagctgctg gacgaccact tcggccccca cggcctggtg 4140

ttccgccgca ccttcgccat ccgcagctac gaggtgggcc ccgaccgctc caccagcatc 4200

gtggccgtga tgaaccacct gcaggaggcc gccctgaacc acgccaagtc cgtgggcatc 4260

ctgggcgacg gcttcggcac caccctggag atgtccaagc gcgacctgat ctgggtggtg 4320

aagcgcaccc acgtggccgt ggagcgctac cccgcctggg gcgacaccgt ggaggtggag 4380

tgctgggtgg gcgcctccgg caacaacggc cgccgccacg acttcctggt gcgcgactgc 4440

aagaccggcg agatcctgac ccgctgcacc tccctgagcg tgatgatgaa cacccgcacc 4500

cgccgcctga gcaagatccc cgaggaggtg cgcggcgaga tcggccccgc cttcatcgac 4560

aacgtggccg tgaaggacga ggagatcaag aagccccaga agctgaacga ctccaccgcc 4620

gactacatcc agggcggcct gacccccgc tggaacgacc tggacatcaa ccagcacgtg 4680

aacaacatca agtacgtgga ctggatcctg gagaccgtgc ccgacagcat cttcgagagc 4740

caccacatct cctccttcac catcgagtac cgccgcgagt gcaccatgga cagcgtgctg 4800

cagtccctga ccaccgtgag cggcggctcc tccgaggccg gcctggtgtg cgagcacctg 4860

ctgcagctgg agggcggcag cgaggtgctg cgcgccaaga ccgagtggcg ccccaagctg 4920

accgactcct tccgcggcat cagcgtgatc cccgccgagt ccagcgtgat ggactacaag 4980

gaccacgacg gcgactacaa ggaccacgac atcgactaca aggacgacga cgacaagtga 5040

ctcgaggcag cagcagctcg gatagtatcg acacactctg gacgctggtc gtgtgatgga 5100

ctgttgccgc cacacttgct gccttgacct gtgaatatcc ctgccgcttt tatcaaacag 5160

cctcagtgtg tttgatcttg tgtgtacgcg cttttgcgag ttgctagctg cttgtgctat 5220

ttgcgaatac caccccccagc atccccttcc ctcgtttcat atcgcttgca tcccaaccgc 5280

aacttatcta cgctgtcctg ctatccctca gcgctgctcc tgctcctgct cactgccccct 5340

cgcacagcct tggtttgggc tccgcctgta ttctcctggt actgcaacct gtaaaccagc 5400

actgcaatgc tgatgcacgg gaagtagtgg gatgggaaca caaatggaaa gctt 5454

<210> 112
<211> 2933
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic polynucleotide

<400> 112
agagagcgga ggtggggttg tgaggtgggg ttgctgacca ggagctcgcg tcgccgagcg 60

cgactcgcac acggtccagt tacccccccc tccgcccaaa cgcaagcctc ccatcttgat 120

gcctttccgg ccacctatac tatttcttag ttcgctgtaa catccagacc gtcctgaata 180

ataacaatgc cctgtgtcaa gtgcattcct aaaaaaattc tgtcccaacc aacaatccca 240

cctgaaatac caccagccct gcccagtaca ctcttccaat accatctccc tacctccacg 300

cgcaagcgac ccccatgcgc gaccaggctc gaaagtgatt tatgacttga gacgagcgag 360

tggcggcgcg gtcgactgcc ttttcatcac gtgccgtacg tcggcgaccg ctagggcttt 420

gcacggcaac gcacggcttc gccaacccga ccagccagga cctcgactac tctaccgcga 480

attcgcctca agaagtcgcc aaatgtgcca tacaccattc cttacagcac tgttcaaact 540

tgatgccaat tttgacattc gggttgctcg ttggctgcgc ccacatcggc cgtgagtgca 600

gcaggcggga tcggacacgg aggacgcggc gtcacgcccc gaacgcagcc cgtaactcta 660

catcaacacg acgtgttgcg taatcccgcc cggctgcgca tcgtgccaac ccattcgcga 720

tggatggtcg gaaaatggtg tgccaactgc cctgagggag gctctcgcga aacgggcacg 780

tccctgaaac cgaaactgtg gccttgtcgt cggccacgca agcacgtgga ccctaaacac 840

caagaaaatc agtaaacaag gttgacatcc tctacgggcg aattgtttgc ccaacccttc 900

atcgcacact gccattataa tgcatctagc tcggcgacaa gtttagaaaa ggcaggctgc 960

attgttccat ttcgccgtgg cggcgtgggt gcccatttta cgaggtttgg gctcccgggc 1020

agcgaccgag ccaggtcgag tccctctcgc ccgtcgacaa tgttgcgaac cccacaagcg 1080

gctaacaaca acttgatggt acctgtacac tgccaattcc ttcttccccg gccgaggttt 1140

acacgtgatg gccatggctt cgcattcagg ccgacttccc attccgactt tccagagggt 1200

ccgcggacgc tgggggttgg ctgcctgagg cccacccttt gttccccgcg tcccgacaaa 1260

cacaattgcg ttacataagg gggagccgcc cccgttcaga gtgcagaaat ctttcactat 1320

attttccagt cgtcagcgaa atcaagtact agtatggcca ccgcatccac tttctcggcg 1380

ttcaatgccc gctgcggcga cctgcgtcgc tcggcgggct ccgggccccg gcgcccagcg 1440

aggcccctcc ccgtgcgcgg gcgcgccccc gactggtcca tgctgttcgc cgtgatcacc 1500

accatcttct ccgccgccga gaagcagtgg accaacctgg agtggaagcc caagcccaac 1560

cccccccagc tgctggacga ccacttcggc ccccacggcc tggtgttccg ccgcaccttc 1620

gccatccgca gctacgaggt gggccccgac cgctccacca gcatcgtggc cgtgatgaac 1680

cacctgcagg aggccgccct gaaccacgcc aagtccgtgg gcatcctggg cgacggcttc 1740

ggcaccaccc tggagatgtc caagcgcgac ctgatctggg tggtgaagcg cacccacgtg 1800

gccgtggagc gctaccccgc ctggggcgac accgtggagg tggagtgctg ggtgggcgcc 1860

tccggcaaca acggccgccg ccacgacttc ctggtgcgcg actgcaagac cggcgagatc 1920

ctgacccgct gcacctccct gagcgtgatg atgaacaccc gcacccgccg cctgagcaag 1980

atccccgagg aggtgcgcgg cgagatcggc cccgccttca tcgacaacgt ggccgtgaag 2040

gacgaggaga tcaagaagcc ccagaagctg aacgactcca ccgccgacta catccagggc 2100

ggcctgaccc cccgctggaa cgacctggac atcaaccagc acgtgaacaa catcaagtac 2160

gtggactgga tcctggagac cgtgcccgac agcatcttcg agagccacca catctcctcc 2220

ttcaccatcg agtaccgccg cgagtgcacc atggacagcg tgctgcagtc cctgaccacc 2280

gtgagcggcg gctcctccga ggccggcctg gtgtgcgagc acctgctgca gctggagggc 2340

ggcagcgagg tgctgcgcgc caagaccgag tggcgcccca agctgaccga ctccttccgc 2400

ggcatcagcg tgatccccgc cgagtccagc gtgatggact acaaggacca cgacggcgac 2460

tacaaggacc acgacatcga ctacaaggac gacgacgaca agtgactcga gttaattaac 2520

tcgaggcagc agcagctcgg atagtatcga cacactctgg acgctggtcg tgtgatggac 2580

tgttgccgcc acacttgctg ccttgacctg tgaatatccc tgccgctttt atcaaacagc 2640

ctcagtgtgt ttgatcttgt gtgtacgcgc ttttgcgagt tgctagctgc ttgtgctatt 2700

tgcgaatacc acccccagca tccccttccc tcgtttcata tcgcttgcat cccaaccgca 2760

acttatctac gctgtcctgc tatccctcag cgctgctcct gctcctgctc actgcccctc 2820

gcacagcctt ggtttgggct ccgcctgtat tctcctggta ctgcaacctg taaaccagca 2880

ctgcaatgct gatgcacggg aagtagtggg atgggaacac aaatggaaag ctt 2933

<210> 113
<211> 4817
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic polynucleotide

<400> 113

ggtacccgcc tgcaacgcaa gggcagccac agccgctccc acccgccgct gaaccgacac 60

gtgcttgggc gcctgccgcc tgcctgccgc atgcttgtgc tggtgaggct gggcagtgct 120

gccatgctga ttgaggcttg gttcatcggg tggaagctta tgtgtgtgct gggcttgcat 180

gccgggcaat gcgcatggtg gcaagagggc ggcagcactt gctggagctg ccgcggtgcc 240

tccaggtggt tcaatcgcgg cagccagagg gatttcagat gatcgcgcgt acaggttgag 300

cagcagtgtc agcaaaggta gcagtttgcc agaatgatcg gttcagctgt taatcaatgc 360

cagcaagaga aggggtcaag tgcaaacacg ggcatgccac agcacgggca ccggggagtg 420

gaatggcacc accaagtgtg tgcgagccag catcgccgcc tggctgtttc agctacaacg 480

gcaggagtca tccaacgtaa ccatgagctg atcaacactg caatcatcgg gcgggcgtga 540

tgcaagcatg cctggcgaag acacatggtg tgcggatgct gccggctgct gcctgctgcg 600

cacgccgttg agttggcagc aggctcagcc atgcactgga tggcagctgg gctgccactg 660

caatgtggtg gataggatgc aagtggagcg aataccaaac cctctggctg cttgctgggt 720

tgcatggcat cgcaccatca gcaggagcgc atgcgaaggg actggcccca tgcacgccat 780

gccaaaccgg agcgcaccga gtgtccacac tgtcaccagg cccgcaagct ttgcagaacc 840

atgctcatgg acgcatgtag cgctgacgtc ccttgacggc gctcctctcg ggtgtgggaa 900

acgcaatgca gcacaggcag cagaggcggc ggcagcagag cggcggcagc agcggcgggg 960

gccacccttc ttgcggggtc gcgccccagc cagcggtgat gcgctgatcc caaacgagtt 1020

cacattcatt tgcatgcctg gagaagcgag gctggggcct ttgggctggt gcagcccgca 1080

atggaatgcg ggaccgccag gctagcagca aaggcgcctc ccctactccg catcgatgtt 1140

ccatagtgca ttggactgca tttgggtggg gcggccggct gtttctttcg tgttgcaaaa 1200

cgcgccagct cagcaacctg tcccgtgggt cccccgtgcc gatgaaatcg tgtgcacgcc 1260

gatcagctga ttgcccggct cgcgaagtag gcgccctcct ttctgctcgc cctctctccg 1320

tcccgcctct agaatatcaa tgatcgagca ggacggcctc cacgccggct cccccgccgc 1380

ctgggtggag cgcctgttcg gctacgactg ggcccagcag accatcggct gctccgacgc 1440

cgccgtgttc cgcctgtccg cccagggccg ccccgtgctg ttcgtgaaga ccgacctgtc 1500

cggcgccctg aacgagctgc aggacgaggc cgcccgcctg tcctggctgg ccaccaccgg 1560

cgtgccctgc gccgccgtgc tggacgtggt gaccgaggcc ggccgcgact ggctgctgct 1620

gggcgaggtg cccggccagg acctgctgtc ctcccacctg gcccccgccg agaaggtgtc 1680

catcatggcc gacgccatgc gccgcctgca caccctggac cccgccacct gccccttcga 1740

ccaccaggcc aagcaccgca tcgagcgcgc ccgcacccgc atggaggccg gcctggtgga 1800

ccaggacgac ctggacgagg agcaccaggg cctggccccc gccgagctgt tcgcccgcct 1860

gaaggcccgc atgcccgacg gcgaggacct ggtggtgacc cacggcgacg cctgcctgcc 1920

caacatcatg gtggagaacg gccgcttctc cggcttcatc gactgcggcc gcctgggcgt 1980

ggccgaccgc taccaggaca tcgccctggc cacccgcgac atcgccgagg agctgggcgg 2040

cgagtgggcc gaccgcttcc tggtgctgta cggcatcgcc gcccccgact cccagcgcat 2100

cgccttctac cgcctgctgg acgagttctt ctgacaattg gcagcagcag ctcggatagt 2160

atcgacacac tctggacgct ggtcgtgtga tggactgttg ccgccacact tgctgccttg 2220

acctgtgaat atccctgccg cttttatcaa acagcctcag tgtgtttgat cttgtgtgta 2280

cgcgcttttg cgagttgcta gctgcttgtg ctatttgcga ataccacccc cagcatcccc 2340

ttccctcgtt tcatatcgct tgcatcccaa ccgcaactta tctacgctgt cctgctatcc 2400

ctcagcgctg ctcctgctcc tgctcactgc ccctcgcaca gccttggttt gggctccgcc 2460

tgtattctcc tggtactgca acctgtaaac cagcactgca atgctgatgc acgggaagta 2520

gtgggatggg aacacaaatg gaggatcccg cgtctcgaac agagcgcgca gaggaacgct 2580

gaaggtctcg cctctgtcgc acctcagcgc ggcatacacc acaataacca cctgacgaat 2640

gcgcttggtt cttcgtccat tagcgaagcg tccggttcac acacgtgcca cgttggcgag 2700

gtggcaggtg acaatgatcg gtggagctga tggtcgaaac gttcacagcc tagggatatc 2760

gaattccttt cttgcgctat gacacttcca gcaaaaggta gggcgggctg cgagacggct 2820

tcccggcgct gcatgcaaca ccgatgatgc ttcgaccccc cgaagctcct tcggggctgc 2880

atgggcgctc cgatgccgct ccagggcgag cgctgtttaa atagccaggc ccccgattgc 2940

aaagacatta tagcgagcta ccaaagccat attcaaacac ctagatcact accacttcta 3000

cacaggccac tcgagcttgt gatcgcactc cgctaagggg gcgcctcttc ctcttcgttt 3060

cagtcacaac ccgcaaacgg cgcgccatgg tggccgccgc cgcctccagc gccttcttcc 3120

ccgtgcccgc ccccggcgcc tcccccaagc ccggcaagtt cggcaactgg ccctccagcc 3180

tgagcccctc cttcaagccc aagtccatcc ccaacggcgg cttccaggtg aaggccaacg 3240

acagcgccca ccccaaggcc aacggctccg ccgtgagcct gaagagcggc agcctgaaca 3300

cccaggagga cacctcctcc agccccccccc cccgcacctt cctgcaccag ctgcccgact 3360

ggagccgcct gctgaccgcc atcaccaccg tgttcgtgaa gtccaagcgc cccgacatgc 3420

acgaccgcaa gtccaagcgc cccgacatgc tggtggacag cttcggcctg gagtccaccg 3480

tgcaggacgg cctggtgttc cgccagtcct tctccatccg ctcctacgag atcggcaccg 3540

accgcaccgc cagcatcgag accctgatga accacctgca ggagacctcc ctgaaccact 3600

gcaagagcac cggcatcctg ctggacggct tcggccgcac cctggagatg tgcaagcgcg 3660

acctgatctg ggtggtgatc aagatgcaga tcaaggtgaa ccgctacccc gcctggggcg 3720

acaccgtgga gatcaacacc cgcttcagcc gcctgggcaa gatcggcatg ggccgcgact 3780

ggctgatctc cgactgcaac accggcgaga tcctggtgcg cgccaccagc gcctacgcca 3840

tgatgaacca gaagacccgc cgcctgtcca agctgcccta cgaggtgcac caggagatcg 3900

tgcccctgtt cgtggacagc cccgtgatcg aggactccga cctgaaggtg cacaagttca 3960

aggtgaagac cggcgacagc atccagaagg gcctgacccc cggctggaac gacctggacg 4020

tgaaccagca cgtgtccaac gtgaagtaca tcggctggat cctggagagc atgcccaccg 4080

aggtgctgga gacccaggag ctgtgctccc tggccctgga gtaccgccgc gagtgcggcc 4140

gcgactccgt gctggagagc gtgaccgcca tggaccccag caaggtgggc gtgcgctccc 4200

agtaccagca cctgctgcgc ctggaggacg gcaccgccat cgtgaacggc gccaccgagt 4260

ggcgccccaa gaacgccggc gccaacggcg ccatctccac cggcaagacc agcaacggca 4320

actccgtgtc catggactac aaggaccacg acggcgacta caaggaccac gacatcgact 4380

acaaggacga cgacgacaag tgactcgagg cagcagcagc tcggatagta tcgacacact 4440

ctggacgctg gtcgtgtgat ggactgttgc cgccacactt gctgccttga cctgtgaata 4500

tccctgccgc ttttatcaaa cagcctcagt gtgtttgatc ttgtgtgtac gcgcttttgc 4560

gagttgctag ctgcttgtgc tatttgcgaa taccaccccc agcatccccc tccctcgttt 4620

catatcgctt gcatcccaac cgcaacttat ctacgctgtc ctgctatccc tcagcgctgc 4680

tcctgctcct gctcactgcc cctcgcacag ccttggtttg ggctccgcct gtattctcct 4740

ggtactgcaa cctgtaaacc agcactgcaa tgctgatgca cgggaagtag tgggatggga 4800

acacaaatgg aaagctt 4817

<210> 114
<211> 4665
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic polynucleotide

<400> 114
ggtacccgcc tgcaacgcaa gggcagccac agccgctccc acccgccgct gaaccgacac 60

gtgcttgggc gcctgccgcc tgcctgccgc atgcttgtgc tggtgaggct gggcagtgct 120

gccatgctga ttgaggcttg gttcatcggg tggaagctta tgtgtgtgct gggcttgcat 180

gccgggcaat gcgcatggtg gcaagagggc ggcagcactt gctggagctg ccgcggtgcc 240

tccaggtggt tcaatcgcgg cagccagagg gatttcagat gatcgcgcgt acaggttgag 300

cagcagtgtc agcaaaggta gcagtttgcc agaatgatcg gttcagctgt taatcaatgc 360

cagcaagaga aggggtcaag tgcaaacacg ggcatgccac agcacgggca ccggggagtg 420

gaatggcacc accaagtgtg tgcgagccag catcgccgcc tggctgtttc agctacaacg 480

gcaggagtca tccaacgtaa ccatgagctg atcaacactg caatcatcgg gcgggcgtga 540

tgcaagcatg cctggcgaag acacatggtg tgcggatgct gccggctgct gcctgctgcg 600

cacgccgttg agttggcagc aggctcagcc atgcactgga tggcagctgg gctgccactg 660

caatgtggtg gataggatgc aagtggagcg aataccaaac cctctggctg cttgctgggt 720

tgcatggcat cgcaccatca gcaggagcgc atgcgaaggg actggcccca tgcacgccat 780

gccaaaccgg agcgcaccga gtgtccacac tgtcaccagg cccgcaagct ttgcagaacc 840

atgctcatgg acgcatgtag cgctgacgtc ccttgacggc gctcctctcg ggtgtgggaa 900

acgcaatgca gcacaggcag cagaggcggc ggcagcagag cggcggcagc agcggcgggg 960

gccacccttc ttgcggggtc gcgccccagc cagcggtgat gcgctgatcc caaacgagtt 1020
cacattcatt tgcatgcctg gagaagcgag gctggggcct ttgggctggt gcagcccgca 1080
atggaatgcg ggaccgccag gctagcagca aaggcgcctc ccctactccg catcgatgtt 1140
ccatagtgca ttggactgca tttgggtggg gcggccggct gtttctttcg tgttgcaaaa 1200
cgcgccagct cagcaacctg tcccgtgggt cccccgtgcc gatgaaatcg tgtgcacgcc 1260
gatcagctga ttgcccggct cgcgaagtag gcgccctcct ttctgctcgc cctctctccg 1320
tcccgcctct agaatatcaa tgatcgagca ggacggcctc cacgccggct cccccgccgc 1380
ctgggtggag cgcctgttcg gctacgactg ggcccagcag accatcggct gctccgacgc 1440
cgccgtgttc cgcctgtccg cccagggccg ccccgtgctg ttcgtgaaga ccgacctgtc 1500
cggcgccctg aacgagctgc aggacgaggc cgcccgcctg tcctggctgg ccaccaccgg 1560
cgtgccctgc gccgccgtgc tggacgtggt gaccgaggcc ggccgcgact ggctgctgct 1620
gggcgaggtg cccggccagg acctgctgtc ctcccacctg gcccccgccg agaaggtgtc 1680
catcatggcc gacgccatgc gccgcctgca caccctggac cccgccacct gccccttcga 1740
ccaccaggcc aagcaccgca tcgagcgcgc ccgcacccgc atggaggccg gcctggtgga 1800
ccaggacgac ctggacgagg agcaccaggg cctggccccc gccgagctgt tcgcccgcct 1860
gaaggcccgc atgcccgacg gcgaggacct ggtggtgacc cacggcgacg cctgcctgcc 1920
caacatcatg gtggagaacg gccgcttctc cggcttcatc gactgcggcc gcctgggcgt 1980
ggccgaccgc taccaggaca tcgccctggc cacccgcgac atcgccgagg agctgggcgg 2040
cgagtgggcc gaccgcttcc tggtgctgta cggcatcgcc gcccccgact cccagcgcat 2100
cgccttctac cgcctgctgg acgagttctt ctgacaattg gcagcagcag ctcggatagt 2160
atcgacacac tctggacgct ggtcgtgtga tggactgttg ccgccacact tgctgccttg 2220
acctgtgaat atccctgccg cttttatcaa acagcctcag tgtgtttgat cttgtgtgta 2280
cgcgcttttg cgagttgcta gctgcttgtg ctatttgcga ataccacccc cagcatcccc 2340
ttccctcgtt tcatatcgct tgcatcccaa ccgcaactta tctacgctgt cctgctatcc 2400
ctcagcgctg ctcctgctcc tgctcactgc ccctcgcaca gccttggttt gggctccgcc 2460
tgtattctcc tggtactgca acctgtaaac cagcactgca atgctgatgc acgggaagta 2520
gtgggatggg aacacaaatg gaggatcccg cgtctcgaac agagcgcgca gaggaacgct 2580
gaaggtctcg cctctgtcgc acctcagcgc ggcatacacc acaataacca cctgacgaat 2640
gcgcttggtt cttcgtccat tagcgaagcg tccggttcac acacgtgcca cgttggcgag 2700
gtggcaggtg acaatgatcg gtggagctga tggtcgaaac gttcacagcc tagggatatc 2760
gaattccttt cttgcgctat gacacttcca gcaaaaggta gggcgggctg cgagacggct 2820
tcccggcgct gcatgcaaca ccgatgatgc ttcgaccccc cgaagctcct tcggggctgc 2880
atgggcgctc cgatgccgct ccagggcgag cgctgtttaa atagccaggc ccccgattgc 2940
aaagacatta tagcgagcta ccaaagccat attcaaacac ctagatcact accacttcta 3000

cacaggccac tcgagcttgt gatcgcactc cgctaagggg gcgcctcttc ctcttcgttt 3060

cagtcacaac ccgcaaacac tagtatgacg ttcggggtcg ccctcccggc catgggccgc 3120

ggtgtctccc ttccccggcc cagggtcgcg gtgcgcgccc agtcggcgag tcaggttttg 3180

gagagcgggc gcgcccagct gcccgactgg agccgcctgc tgaccgccat caccaccgtg 3240

ttcgtgaagt ccaagcgccc cgacatgcac gaccgcaagt ccaagcgccc cgacatgctg 3300

gtggacagct tcggcctgga gtccaccgtg caggacggcc tggtgttccg ccagtccttc 3360

tccatccgct cctacgagat cggcaccgac cgcaccgcca gcatcgagac cctgatgaac 3420

cacctgcagg agacctccct gaaccactgc aagagcaccg gcatcctgct ggacggcttc 3480

ggccgcaccc tggagatgtg caagcgcgac ctgatctggg tggtgatcaa gatgcagatc 3540

aaggtgaacc gctaccccgc ctggggcgac accgtggaga tcaacacccg cttcagccgc 3600

ctgggcaaga tcggcatggg ccgcgactgg ctgatctccg actgcaacac cggcgagatc 3660

ctggtgcgcg ccaccagcgc ctacgccatg atgaaccaga agacccgccg cctgtccaag 3720

ctgccctacg aggtgcacca ggagatcgtg cccctgttcg tggacagccc cgtgatcgag 3780

gactccgacc tgaaggtgca caagttcaag gtgaagaccg gcgacagcat ccagaagggc 3840

ctgacccccg gctggaacga cctggacgtg aaccagcacg tgtccaacgt gaagtacatc 3900

ggctggatcc tggagagcat gcccaccgag gtgctggaga cccaggagct gtgctccctg 3960

gccctggagt accgccgcga gtgcggccgc gactccgtgc tggagagcgt gaccgccatg 4020

gaccccagca aggtgggcgt gcgctcccag taccagcacc tgctgcgcct ggaggacggc 4080

accgccatcg tgaacggcgc caccgagtgg cgccccaaga acgccggcgc caacggcgcc 4140

atctccaccg gcaagaccag caacggcaac tccgtgtcca tggactacaa ggaccacgac 4200

ggcgactaca aggaccacga catcgactac aaggacgacg acgacaagtg actcgaggca 4260

gcagcagctc ggatagtatc gacacactct ggacgctggt cgtgtgatgg actgttgccg 4320

ccacacttgc tgccttgacc tgtgaatatc cctgccgctt ttatcaaaca gcctcagtgt 4380

gtttgatctt gtgtgtacgc gcttttgcga gttgctagct gcttgtgcta tttgcgaata 4440

ccacccccag catccccttc cctcgtttca tatcgcttgc atcccaaccg caacttatct 4500

acgctgtcct gctatccctc agcgctgctc ctgctcctgc tcactgcccc tcgcacagcc 4560

ttggtttggg ctccgcctgt attctcctgg tactgcaacc tgtaaaccag cactgcaatg 4620

ctgatgcacg ggaagtagtg ggatgggaac acaaatggaa agctt 4665

<210> 115
<211> 4668
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic polynucleotide

<400> 115

ggtacccgcc tgcaacgcaa gggcagccac agccgctccc acccgccgct gaaccgacac 60

gtgcttgggc gcctgccgcc tgcctgccgc atgcttgtgc tggtgaggct gggcagtgct 120

gccatgctga ttgaggcttg gttcatcggg tggaagctta tgtgtgtgct gggcttgcat 180

gccgggcaat gcgcatggtg gcaagagggc ggcagcactt gctggagctg ccgcggtgcc 240

tccaggtggt tcaatcgcgg cagccagagg gatttcagat gatcgcgcgt acaggttgag 300

cagcagtgtc agcaaaggta gcagtttgcc agaatgatcg gttcagctgt taatcaatgc 360

cagcaagaga aggggtcaag tgcaaacacg ggcatgccac agcacgggca ccggggagtg 420

gaatggcacc accaagtgtg tgcgagccag catcgccgcc tggctgtttc agctacaacg 480

gcaggagtca tccaacgtaa ccatgagctg atcaacactg caatcatcgg gcgggcgtga 540

tgcaagcatg cctggcgaag acacatggtg tgcggatgct gccggctgct gcctgctgcg 600

cacgccgttg agttggcagc aggctcagcc atgcactgga tggcagctgg gctgccactg 660

caatgtggtg gataggatgc aagtggagcg aataccaaac cctctggctg cttgctgggt 720

tgcatggcat cgcaccatca gcaggagcgc atgcgaaggg actggcccca tgcacgccat 780

gccaaaccgg agcgcaccga gtgtccacac tgtcaccagg cccgcaagct ttgcagaacc 840

atgctcatgg acgcatgtag cgctgacgtc ccttgacggc gctcctctcg ggtgtgggaa 900

acgcaatgca gcacaggcag cagaggcggc ggcagcagag cggcggcagc agcggcgggg 960

gccacccttc ttgcggggtc gcgccccagc cagcggtgat gcgctgatcc caaacgagtt 1020

cacattcatt tgcatgcctg gagaagcgag gctggggcct ttgggctggt gcagcccgca 1080

atggaatgcg ggaccgccag gctagcagca aaggcgcctc ccctactccg catcgatgtt 1140

ccatagtgca ttggactgca tttgggtggg gcggccggct gtttctttcg tgttgcaaaa 1200

cgcgccagct cagcaacctg tcccgtgggt cccccgtgcc gatgaaatcg tgtgcacgcc 1260

gatcagctga ttgcccggct cgcgaagtag gcgccctcct ttctgctcgc cctctctccg 1320

tcccgcctct agaatatcaa tgatcgagca ggacggcctc cacgccggct cccccgccgc 1380

ctgggtggag cgcctgttcg gctacgactg ggcccagcag accatcggct gctccgacgc 1440

cgccgtgttc cgcctgtccg cccagggccg ccccgtgctg ttcgtgaaga ccgacctgtc 1500

cggcgccctg aacgagctgc aggacgaggc cgcccgcctg tcctggctgg ccaccaccgg 1560

cgtgccctgc gccgccgtgc tggacgtggt gaccgaggcc ggccgcgact ggctgctgct 1620

gggcgaggtg cccggccagg acctgctgtc ctcccacctg gcccccgccg agaaggtgtc 1680

catcatggcc gacgccatgc gccgcctgca caccctggac cccgccacct gcccccttcga 1740

ccaccaggcc aagcaccgca tcgagcgcgc ccgcacccgc atggaggccg gcctggtgga 1800

ccaggacgac ctggacgagg agcaccaggg cctggccccc gccgagctgt tcgcccgcct 1860

gaaggcccgc atgcccgacg gcgaggacct ggtggtgacc cacggcgacg cctgcctgcc 1920

caacatcatg gtggagaacg gccgcttctc cggcttcatc gactgcggcc gcctgggcgt 1980

ggccgaccgc taccaggaca tcgccctggc cacccgcgac atcgccgagg agctgggcgg 2040

cgagtgggcc gaccgcttcc tggtgctgta cggcatcgcc gcccccgact cccagcgcat 2100

cgccttctac cgcctgctgg acgagttctt ctgacaattg gcagcagcag ctcggatagt 2160

atcgacacac tctggacgct ggtcgtgtga tggactgttg ccgccacact tgctgccttg 2220

acctgtgaat atccctgccg cttttatcaa acagcctcag tgtgtttgat cttgtgtgta 2280

cgcgcttttg cgagttgcta gctgcttgtg ctatttgcga ataccacccc cagcatcccc 2340

ttccctcgtt tcatatcgct tgcatcccaa ccgcaactta tctacgctgt cctgctatcc 2400

ctcagcgctg ctcctgctcc tgctcactgc ccctcgcaca gccttggttt gggctccgcc 2460

tgtattctcc tggtactgca acctgtaaac cagcactgca atgctgatgc acgggaagta 2520

gtgggatggg aacacaaatg gaggatcccg cgtctcgaac agagcgcgca gaggaacgct 2580

gaaggtctcg cctctgtcgc acctcagcgc ggcatacacc acaataacca cctgacgaat 2640

gcgcttggtt cttcgtccat tagcgaagcg tccggttcac acacgtgcca cgttggcgag 2700

gtggcaggtg acaatgatcg gtggagctga tggtcgaaac gttcacagcc tagggatatc 2760

gaattccttt cttgcgctat gacacttcca gcaaaaggta gggcgggctg cgagacggct 2820

tcccggcgct gcatgcaaca ccgatgatgc ttcgaccccc cgaagctcct tcggggctgc 2880

atgggcgctc cgatgccgct ccagggcgag cgctgtttaa atagccaggc ccccgattgc 2940

aaagacatta tagcgagcta ccaaagccat attcaaacac ctagatcact accacttcta 3000

cacaggccac tcgagcttgt gatcgcactc cgctaagggg gcgcctcttc ctcttcgttt 3060

cagtcacaac ccgcaaacac tagtatggct atcaagacga acaggcagcc tgtggagaag 3120

cctccgttca cgatcgggac gctgcgcaag gccatccccg cgcactgttt cgagcgctcg 3180

gcgcttcgtg ggcgcgccca gctgcccgac tggagccgcc tgctgaccgc catcaccacc 3240

gtgttcgtga agtccaagcg ccccgacatg cacgaccgca agtccaagcg ccccgacatg 3300

ctggtggaca gcttcggcct ggagtccacc gtgcaggacg gcctggtgtt ccgccagtcc 3360

ttctccatcc gctcctacga gatcggcacc gaccgcaccg ccagcatcga gaccctgatg 3420

aaccacctgc aggagacctc cctgaaccac tgcaagagca ccggcatcct gctggacggc 3480

ttcggccgca ccctggagat gtgcaagcgc gacctgatct gggtggtgat caagatgcag 3540

atcaaggtga accgctaccc cgcctggggc gacaccgtgg agatcaacac ccgcttcagc 3600

cgcctgggca agatcggcat gggccgcgac tggctgatct ccgactgcaa caccggcgag 3660

atcctggtgc gcgccaccag cgcctacgcc atgatgaacc agaagacccg ccgcctgtcc 3720

aagctgccct acgaggtgca ccaggagatc gtgcccctgt tcgtggacag ccccgtgatc 3780

gaggactccg acctgaaggt gcacaagttc aaggtgaaga ccggcgacag catccagaag 3840

ggcctgaccc ccggctggaa cgacctggac gtgaaccagc acgtgtccaa cgtgaagtac 3900

atcggctgga tcctggagag catgcccacc gaggtgctgg agacccagga gctgtgctcc 3960

ctggccctgg agtaccgccg cgagtgcggc cgcgactccg tgctggagag cgtgaccgcc 4020

atggacccca gcaaggtggg cgtgcgctcc cagtaccagc acctgctgcg cctggaggac 4080

ggcaccgcca tcgtgaacgg cgccaccgag tggcgcccca agaacgccgg cgccaacggc 4140

gccatctcca ccggcaagac cagcaacggc aactccgtgt ccatggacta caaggaccac 4200

gacggcgact acaaggacca cgacatcgac tacaaggacg acgacgacaa gtgactcgag 4260

gcagcagcag ctcggatagt atcgacacac tctggacgct ggtcgtgtga tggactgttg 4320

ccgccacact tgctgccttg acctgtgaat atccctgccg cttttatcaa acagcctcag 4380

tgtgtttgat cttgtgtgta cgcgcttttg cgagttgcta gctgcttgtg ctatttgcga 4440

ataccacccc cagcatcccc ttccctcgtt tcatatcgct tgcatcccaa ccgcaactta 4500

tctacgctgt cctgctatcc ctcagcgctg ctcctgctcc tgctcactgc ccctcgcaca 4560

gccttggttt gggctccgcc tgtattctcc tggtactgca acctgtaaac cagcactgca 4620

atgctgatgc acgggaagta gtgggatggg aacacaaatg gaaagctt 4668

<210> 116
<211> 4668
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic polynucleotide

<400> 116
ggtacccgcc tgcaacgcaa gggcagccac agccgctccc acccgccgct gaaccgacac 60

gtgcttgggc gcctgccgcc tgcctgccgc atgcttgtgc tggtgaggct gggcagtgct 120

gccatgctga ttgaggcttg gttcatcggg tggaagctta tgtgtgtgct gggcttgcat 180

gccgggcaat gcgcatggtg gcaagagggc ggcagcactt gctggagctg ccgcggtgcc 240

tccaggtggt tcaatcgcgg cagccagagg gatttcagat gatcgcgcgt acaggttgag 300

cagcagtgtc agcaaaggta gcagtttgcc agaatgatcg gttcagctgt taatcaatgc 360

cagcaagaga aggggtcaag tgcaaacacg ggcatgccac agcacgggca ccggggagtg 420

gaatggcacc accaagtgtg tgcgagccag catcgccgcc tggctgtttc agctacaacg 480

gcaggagtca tccaacgtaa ccatgagctg atcaacactg caatcatcgg gcgggcgtga 540

tgcaagcatg cctggcgaag acacatggtg tgcggatgct gccggctgct gcctgctgcg 600

cacgccgttg agttggcagc aggctcagcc atgcactgga tggcagctgg gctgccactg 660

caatgtggtg gataggatgc aagtggagcg aataccaaac cctctggctg cttgctgggt 720

tgcatggcat cgcaccatca gcaggagcgc atgcgaaggg actggcccca tgcacgccat 780

gccaaaccgg agcgcaccga gtgtccacac tgtcaccagg cccgcaagct ttgcagaacc 840

atgctcatgg acgcatgtag cgctgacgtc ccttgacggc gctcctctcg ggtgtgggaa 900

acgcaatgca gcacaggcag cagaggcggc ggcagcagag cggcggcagc agcggcgggg 960

gccacccttc ttgcggggtc gcgccccagc cagcggtgat gcgctgatcc caaacgagtt 1020

cacattcatt tgcatgcctg gagaagcgag gctggggcct ttgggctggt gcagcccgca 1080

atggaatgcg ggaccgccag gctagcagca aaggcgcctc ccctactccg catcgatgtt 1140

ccatagtgca ttggactgca tttgggtggg gcggccggct gtttctttcg tgttgcaaaa 1200

cgcgccagct cagcaacctg tcccgtgggt cccccgtgcc gatgaaatcg tgtgcacgcc 1260

gatcagctga ttgcccggct cgcgaagtag gcgccctcct ttctgctcgc cctctctccg 1320

tcccgcctct agaatatcaa tgatcgagca ggacggcctc cacgccggct cccccgccgc 1380

ctgggtggag cgcctgttcg gctacgactg ggcccagcag accatcggct gctccgacgc 1440

cgccgtgttc cgcctgtccg cccagggccg ccccgtgctg ttcgtgaaga ccgacctgtc 1500

cggcgccctg aacgagctgc aggacgaggc cgcccgcctg tcctggctgg ccaccaccgg 1560

cgtgccctgc gccgccgtgc tggacgtggt gaccgaggcc ggccgcgact ggctgctgct 1620

gggcgaggtg cccggccagg acctgctgtc ctcccacctg gcccccgccg agaaggtgtc 1680

catcatggcc gacgccatgc gccgcctgca caccctggac cccgccacct gcccccttcga 1740

ccaccaggcc aagcaccgca tcgagcgcgc ccgcacccgc atggaggccg gcctggtgga 1800

ccaggacgac ctggacgagg agcaccaggg cctggccccc gccgagctgt tcgcccgcct 1860

gaaggcccgc atgcccgacg gcgaggacct ggtggtgacc cacggcgacg cctgcctgcc 1920

caacatcatg gtggagaacg gccgcttctc cggcttcatc gactgcggcc gcctgggcgt 1980

ggccgaccgc taccaggaca tcgccctggc cacccgcgac atcgccgagg agctgggcgg 2040

cgagtgggcc gaccgcttcc tggtgctgta cggcatcgcc gcccccgact cccagcgcat 2100

cgccttctac cgcctgctgg acgagttctt ctgacaattg gcagcagcag ctcggatagt 2160

atcgacacac tctggacgct ggtcgtgtga tggactgttg ccgccacact tgctgccttg 2220

acctgtgaat atccctgccg cttttatcaa acagcctcag tgtgtttgat cttgtgtgta 2280

cgcgcttttg cgagttgcta gctgcttgtg ctatttgcga ataccacccc cagcatcccc 2340

ttccctcgtt tcatatcgct tgcatcccaa ccgcaactta tctacgctgt cctgctatcc 2400

ctcagcgctg ctcctgctcc tgctcactgc ccctcgcaca gccttggttt gggctccgcc 2460

tgtattctcc tggtactgca acctgtaaac cagcactgca atgctgatgc acgggaagta 2520

gtgggatggg aacacaaatg gaggatcccg cgtctcgaac agagcgcgca gaggaacgct 2580

gaaggtctcg cctctgtcgc acctcagcgc ggcatacacc acaataacca cctgacgaat 2640

gcgcttggtt cttcgtccat tagcgaagcg tccggttcac acacgtgcca cgttggcgag 2700

gtggcaggtg acaatgatcg gtggagctga tggtcgaaac gttcacagcc tagggatatc 2760

gaattccttt cttgcgctat gacacttcca gcaaaaggta gggcgggctg cgagacggct 2820

tcccggcgct gcatgcaaca ccgatgatgc ttcgaccccc cgaagctcct tcggggctgc 2880

atgggcgctc cgatgccgct ccagggcgag cgctgtttaa atagccaggc ccccgattgc 2940

aaagacatta tagcgagcta ccaaagccat attcaaacac ctagatcact accacttcta 3000

cacaggccac tcgagcttgt gatcgcactc cgctaagggg gcgcctcttc ctcttcgttt 3060

cagtcacaac ccgcaaacac tagtatggcc accgcatcca ctttctcggc gttcaatgcc 3120

cgctgcggcg acctgcgtcg ctcggcgggc tccgggcccc ggcgcccagc gaggcccctc 3180

cccgtgcgcg ggcgcgccca gctgcccgac tggagccgcc tgctgaccgc catcaccacc 3240

gtgttcgtga agtccaagcg ccccgacatg cacgaccgca agtccaagcg ccccgacatg 3300

ctggtggaca gcttcggcct ggagtccacc gtgcaggacg gcctggtgtt ccgccagtcc 3360

ttctccatcc gctcctacga gatcggcacc gaccgcaccg ccagcatcga gaccctgatg 3420

aaccacctgc aggagacctc cctgaaccac tgcaagagca ccggcatcct gctggacggc 3480

ttcggccgca ccctggagat gtgcaagcgc gacctgatct gggtggtgat caagatgcag 3540

atcaaggtga accgctaccc cgcctggggc gacaccgtgg agatcaacac ccgcttcagc 3600

cgcctgggca agatcggcat gggccgcgac tggctgatct ccgactgcaa caccggcgag 3660

atcctggtgc gcgccaccag cgcctacgcc atgatgaacc agaagacccg ccgcctgtcc 3720

aagctgccct acgaggtgca ccaggagatc gtgcccctgt tcgtggacag ccccgtgatc 3780

gaggactccg acctgaaggt gcacaagttc aaggtgaaga ccggcgacag catccagaag 3840

ggcctgaccc ccggctggaa cgacctggac gtgaaccagc acgtgtccaa cgtgaagtac 3900

atcggctgga tcctggagag catgcccacc gaggtgctgg agacccagga gctgtgctcc 3960

ctggccctgg agtaccgccg cgagtgcggc cgcgactccg tgctggagag cgtgaccgcc 4020

atggacccca gcaaggtggg cgtgcgctcc cagtaccagc acctgctgcg cctggaggac 4080

ggcaccgcca tcgtgaacgg cgccaccgag tggcgcccca agaacgccgg cgccaacggc 4140

gccatctcca ccggcaagac cagcaacggc aactccgtgt ccatggacta caaggaccac 4200

gacggcgact acaaggacca cgacatcgac tacaaggacg acgacgacaa gtgactcgag 4260

gcagcagcag ctcggatagt atcgacacac tctggacgct ggtcgtgtga tggactgttg 4320

ccgccacact tgctgccttg acctgtgaat atccctgccg cttttatcaa acagcctcag 4380

tgtgtttgat cttgtgtgta cgcgcttttg cgagttgcta gctgcttgtg ctatttgcga 4440

ataccacccc cagcatcccc ttccctcgtt tcatatcgct tgcatcccaa ccgcaactta 4500

tctacgctgt cctgctatcc ctcagcgctg ctcctgctcc tgctcactgc ccctcgcaca 4560

gccttggttt gggctccgcc tgtattctcc tggtactgca acctgtaaac cagcactgca 4620

atgctgatgc acgggaagta gtgggatggg aacacaaatg gaaagctt 4668

<210> 117
<211> 4656
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic polynucleotide

<400> 117
ggtacccgcc tgcaacgcaa gggcagccac agccgctccc acccgccgct gaaccgacac 60

gtgcttgggc gcctgccgcc tgcctgccgc atgcttgtgc tggtgaggct gggcagtgct 120

gccatgctga ttgaggcttg gttcatcggg tggaagctta tgtgtgtgct gggcttgcat 180

gccgggcaat gcgcatggtg gcaagagggc ggcagcactt gctggagctg ccgcggtgcc 240

tccaggtggt tcaatcgcgg cagccagagg gatttcagat gatcgcgcgt acaggttgag 300

cagcagtgtc agcaaaggta gcagtttgcc agaatgatcg gttcagctgt taatcaatgc 360

cagcaagaga aggggtcaag tgcaaacacg ggcatgccac agcacgggca ccggggagtg 420

gaatggcacc accaagtgtg tgcgagccag catcgccgcc tggctgtttc agctacaacg 480

gcaggagtca tccaacgtaa ccatgagctg atcaacactg caatcatcgg gcgggcgtga 540

tgcaagcatg cctggcgaag acacatggtg tgcggatgct gccggctgct gcctgctgcg 600

cacgccgttg agttggcagc aggctcagcc atgcactgga tggcagctgg gctgccactg 660

caatgtggtg gataggatgc aagtggagcg aataccaaac cctctggctg cttgctgggt 720

tgcatggcat cgcaccatca gcaggagcgc atgcgaaggg actggcccca tgcacgccat 780

gccaaaccgg agcgcaccga gtgtccacac tgtcaccagg cccgcaagct ttgcagaacc 840

atgctcatgg acgcatgtag cgctgacgtc ccttgacggc gctcctctcg ggtgtgggaa 900

acgcaatgca gcacaggcag cagaggcggc ggcagcagag cggcggcagc agcggcgggg 960

gccacccttc ttgcggggtc gcgccccagc cagcggtgat gcgctgatcc caaacgagtt 1020

cacattcatt tgcatgcctg gagaagcgag gctggggcct ttgggctggt gcagcccgca 1080

atggaatgcg ggaccgccag gctagcagca aaggcgcctc ccctactccg catcgatgtt 1140

ccatagtgca ttggactgca tttgggtggg gcggccggct gtttctttcg tgttgcaaaa 1200

cgcgccagct cagcaacctg tcccgtgggt cccccgtgcc gatgaaatcg tgtgcacgcc 1260

gatcagctga ttgcccggct cgcgaagtag gcgccctcct ttctgctcgc cctctctccg 1320

tcccgcctct agaatatcaa tgatcgagca ggacggcctc cacgccggct cccccgccgc 1380

ctgggtggag cgcctgttcg gctacgactg ggcccagcag accatcggct gctccgacgc 1440

cgccgtgttc cgcctgtccg cccagggccg ccccgtgctg ttcgtgaaga ccgacctgtc 1500

cggcgccctg aacgagctgc aggacgaggc cgcccgcctg tcctggctgg ccaccaccgg 1560

cgtgccctgc gccgccgtgc tggacgtggt gaccgaggcc ggccgcgact ggctgctgct 1620

gggcgaggtg cccggccagg acctgctgtc ctcccacctg gcccccgccg agaaggtgtc 1680

catcatggcc gacgccatgc gccgcctgca caccctggac cccgccacct gccccttcga 1740

ccaccaggcc aagcaccgca tcgagcgcgc ccgcacccgc atggaggccg gcctggtgga 1800

ccaggacgac ctggacgagg agcaccaggg cctggccccc gccgagctgt tcgcccgcct 1860

gaaggcccgc atgcccgacg gcgaggacct ggtggtgacc cacggcgacg cctgcctgcc 1920

caacatcatg gtggagaacg gccgcttctc cggcttcatc gactgcggcc gcctgggcgt 1980

ggccgaccgc taccaggaca tcgccctggc cacccgcgac atcgccgagg agctgggcgg 2040

cgagtgggcc gaccgcttcc tggtgctgta cggcatcgcc gcccccgact cccagcgcat 2100

cgccttctac cgcctgctgg acgagttctt ctgacaattg gcagcagcag ctcggatagt 2160

atcgacacac tctggacgct ggtcgtgtga tggactgttg ccgccacact tgctgccttg 2220

acctgtgaat atccctgccg cttttatcaa acagcctcag tgtgtttgat cttgtgtgta 2280

cgcgcttttg cgagttgcta gctgcttgtg ctatttgcga ataccacccc cagcatcccc 2340

ttccctcgtt tcatatcgct tgcatcccaa ccgcaactta tctacgctgt cctgctatcc 2400

ctcagcgctg ctcctgctcc tgctcactgc ccctcgcaca gccttggttt gggctccgcc 2460

tgtattctcc tggtactgca acctgtaaac cagcactgca atgctgatgc acgggaagta 2520

gtgggatggg aacacaaatg gaggatcccg cgtctcgaac agagcgcgca gaggaacgct 2580

gaaggtctcg cctctgtcgc acctcagcgc ggcatacacc acaataacca cctgacgaat 2640

gcgcttggtt cttcgtccat tagcgaagcg tccggttcac acacgtgcca cgttggcgag 2700

gtggcaggtg acaatgatcg gtggagctga tggtcgaaac gttcacagcc tagggatatc 2760

gaattccttt cttgcgctat gacacttcca gcaaaaggta gggcgggctg cgagacggct 2820

tcccggcgct gcatgcaaca ccgatgatgc ttcgacccccc cgaagctcct tcggggctgc 2880

atgggcgctc cgatgccgct ccagggcgag cgctgtttaa atagccaggc ccccgattgc 2940

aaagacatta tagcgagcta ccaaagccat attcaaacac ctagatcact accacttcta 3000

cacaggccac tcgagcttgt gatcgcactc cgctaagggg gcgcctcttc ctcttcgttt 3060

cagtcacaac ccgcaaacac tagtatggct tccgcggcat tcaccatgtc ggcgtgcccc 3120

gcgatgactg gcagggcccc tggggcacgt cgctccggac ggccagtcgc cacccgcctg 3180

agggggcgcg cccccgactg gagccgcctg ctgaccgcca tcaccaccgt gttcgtgaag 3240

tccaagcgcc ccgacatgca cgaccgcaag tccaagcgcc ccgacatgct ggtggacagc 3300

ttcggcctgg agtccaccgt gcaggacggc ctggtgttcc gccagtcctt ctccatccgc 3360

tcctacgaga tcggcaccga ccgcaccgcc agcatcgaga ccctgatgaa ccacctgcag 3420

gagacctccc tgaaccactg caagagcacc ggcatcctgc tggacggctt cggccgcacc 3480

ctggagatgt gcaagcgcga cctgatctgg gtggtgatca agatgcagat caaggtgaac 3540

cgctaccccg cctggggcga caccgtggag atcaacaccc gcttcagccg cctgggcaag 3600

atcggcatgg gccgcgactg gctgatctcc gactgcaaca ccggcgagat cctggtgcgc 3660

gccaccagcg cctacgccat gatgaaccag aagacccgcc gcctgtccaa gctgccctac 3720

gaggtgcacc aggagatcgt gcccctgttc gtggacagcc ccgtgatcga ggactccgac 3780

ctgaaggtgc acaagttcaa ggtgaagacc ggcgacagca tccagaaggg cctgaccccc 3840

ggctggaacg acctggacgt gaaccagcac gtgtccaacg tgaagtacat cggctggatc 3900

ctggagagca tgcccaccga ggtgctggag acccaggagc tgtgctccct ggccctggag 3960

taccgccgcg agtgcggccg cgactccgtg ctggagagcg tgaccgccat ggaccccagc 4020

aaggtgggcg tgcgctccca gtaccagcac ctgctgcgcc tggaggacgg caccgccatc 4080

gtgaacggcg ccaccgagtg gcgccccaag aacgccggcg ccaacggcgc catctccacc 4140

ggcaagacca gcaacggcaa ctccgtgtcc atggactaca aggaccacga cggcgactac 4200

aaggaccacg acatcgacta caaggacgac gacgacaagt gactcgaggc agcagcagct 4260

cggatagtat cgacacactc tggacgctgg tcgtgtgatg gactgttgcc gccacacttg 4320

ctgccttgac ctgtgaatat ccctgccgct tttatcaaac agcctcagtg tgtttgatct 4380

tgtgtgtacg cgcttttgcg agttgctagc tgcttgtgct atttgcgaat accaccccca 4440

gcatcccctt ccctcgtttc atatcgcttg catcccaacc gcaacttatc tacgctgtcc 4500

tgctatccct cagcgctgct cctgctcctg ctcactgccc ctcgcacagc cttggtttgg 4560

gctccgcctg tattctcctg gtactgcaac ctgtaaacca gcactgcaat gctgatgcac 4620

gggaagtagt gggatgggaa cacaaatgga aagctt 4656

<210> 118
<211> 4721
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic polynucleotide

<400> 118
ggtacccgcc tgcaacgcaa gggcagccac agccgctccc acccgccgct gaaccgacac 60

gtgcttgggc gcctgccgcc tgcctgccgc atgcttgtgc tggtgaggct gggcagtgct 120

gccatgctga ttgaggcttg gttcatcggg tggaagctta tgtgtgtgct gggcttgcat 180

gccgggcaat gcgcatggtg gcaagagggc ggcagcactt gctggagctg ccgcggtgcc 240

tccaggtggt tcaatcgcgg cagccagagg gatttcagat gatcgcgcgt acaggttgag 300

cagcagtgtc agcaaaggta gcagtttgcc agaatgatcg gttcagctgt taatcaatgc 360

cagcaagaga aggggtcaag tgcaaacacg ggcatgccac agcacgggca ccggggagtg 420

gaatggcacc accaagtgtg tgcgagccag catcgccgcc tggctgtttc agctacaacg 480

gcaggagtca tccaacgtaa ccatgagctg atcaacactg caatcatcgg gcgggcgtga 540

tgcaagcatg cctggcgaag acacatggtg tgcggatgct gccggctgct gcctgctgcg 600

cacgccgttg agttggcagc aggctcagcc atgcactgga tggcagctgg gctgccactg 660

caatgtggtg gataggatgc aagtggagcg aataccaaac cctctggctg cttgctgggt 720

tgcatggcat cgcaccatca gcaggagcgc atgcgaaggg actggcccca tgcacgccat 780

gccaaaccgg agcgcaccga gtgtccacac tgtcaccagg cccgcaagct ttgcagaacc 840

atgctcatgg acgcatgtag cgctgacgtc ccttgacggc gctcctctcg ggtgtgggaa 900

acgcaatgca gcacaggcag cagaggcggc ggcagcagag cggcggcagc agcggcgggg 960

gccacccttc ttgcggggtc gcgccccagc cagcggtgat gcgctgatcc caaacgagtt 1020

cacattcatt tgcatgcctg gagaagcgag gctggggcct ttgggctggt gcagcccgca 1080

atggaatgcg ggaccgccag gctagcagca aaggcgcctc ccctactccg catcgatgtt 1140

ccatagtgca ttggactgca tttgggtggg gcggccggct gtttctttcg tgttgcaaaa 1200

cgcgccagct cagcaacctg tcccgtgggt cccccgtgcc gatgaaatcg tgtgcacgcc 1260

gatcagctga ttgcccggct cgcgaagtag gcgccctcct ttctgctcgc cctctctccg 1320

tcccgcctct agaatatcaa tgatcgagca ggacggcctc cacgccggct cccccgccgc 1380
ctgggtggag cgcctgttcg gctacgactg ggcccagcag accatcggct gctccgacgc 1440
cgccgtgttc cgcctgtccg cccagggccg ccccgtgctg ttcgtgaaga ccgacctgtc 1500
cggcgccctg aacgagctgc aggacgaggc cgcccgcctg tcctggctgg ccaccaccgg 1560
cgtgccctgc gccgccgtgc tggacgtggt gaccgaggcc ggccgcgact ggctgctgct 1620
gggcgaggtg cccggccagg acctgctgtc ctcccacctg gcccccgccg agaaggtgtc 1680
catcatggcc gacgccatgc gccgcctgca caccctggac cccgccacct gccccttcga 1740
ccaccaggcc aagcaccgca tcgagcgcgc ccgcacccgc atggaggccg gcctggtgga 1800
ccaggacgac ctggacgagg agcaccaggg cctggccccc gccgagctgt tcgcccgcct 1860
gaaggcccgc atgcccgacg gcgaggacct ggtggtgacc cacggcgacg cctgcctgcc 1920
caacatcatg gtggagaacg gccgcttctc cggcttcatc gactgcggcc gcctgggcgt 1980
ggccgaccgc taccaggaca tcgccctggc cacccgcgac atcgccgagg agctgggcgg 2040
cgagtgggcc gaccgcttcc tggtgctgta cggcatcgcc gcccccgact cccagcgcat 2100
cgccttctac cgcctgctgg acgagttctt ctgacaattg gcagcagcag ctcggatagt 2160
atcgacacac tctggacgct ggtcgtgtga tggactgttg ccgccacact tgctgccttg 2220
acctgtgaat atccctgccg cttttatcaa acagcctcag tgtgtttgat cttgtgtgta 2280
cgcgcttttg cgagttgcta gctgcttgtg ctatttgcga ataccacccc cagcatcccc 2340
ttccctcgtt tcatatcgct tgcatcccaa ccgcaactta tctacgctgt cctgctatcc 2400
ctcagcgctg ctcctgctcc tgctcactgc ccctcgcaca gccttggttt gggctccgcc 2460
tgtattctcc tggtactgca acctgtaaac cagcactgca atgctgatgc acgggaagta 2520
gtgggatggg aacacaaatg gaggatcccg cgtctcgaac agagcgcgca gaggaacgct 2580
gaaggtctcg cctctgtcgc acctcagcgc ggcatacacc acaataacca cctgacgaat 2640
gcgcttggtt cttcgtccat tagcgaagcg tccggttcac acacgtgcca cgttggcgag 2700
gtggcaggtg acaatgatcg gtggagctga tggtcgaaac gttcacagcc tagggatatc 2760
gaattccttt cttgcgctat gacacttcca gcaaaaggta gggcgggctg cgagacggct 2820
tcccggcgct gcatgcaaca ccgatgatgc ttcgaccccc cgaagctcct tcggggctgc 2880
atgggcgctc cgatgccgct ccagggcgag cgctgtttaa atagccaggc ccccgattgc 2940
aaagacatta tagcgagcta ccaaagccat attcaaacac ctagatcact accacttcta 3000
cacaggccac tcgagcttgt gatcgcactc cgctaagggg gcgcctcttc ctcttcgttt 3060
cagtcacaac ccgcaaacgg cgcgccatgg ccaccaccag cctggcctcc gccttctgct 3120
ccatgaaggc cgtgatgctg gcccgcgacg gccgcggcat gaagccccgc agctccgacc 3180
tgcagctgcg cgccggcaac gcccccacct ccctgaagat gatcaacggc accaagttca 3240
gctacaccga gagcctgaag cgcctgcccg actggtccat gctgttcgcc gtgatcacca 3300
ccatcttcag cgccgccgag aagcagtgga ccaacctgga gtggaagccc aagcccaagc 3360

tgccccagct gctggacgac cacttcggcc tgcacggcct ggtgttccgc cgcaccttcg 3420

ccatccgctc ctacgaggtg ggccccgacc gcagcacctc catcctggcc gtgatgaacc 3480

acatgcagga ggccaccctg aaccacgcca agagcgtggg catcctgggc gacggcttcg 3540

gcaccaccct ggagatgtcc aagcgcgacc tgatgtgggt ggtgcgccgc acccacgtgg 3600

ccgtggagcg ctaccccacc tggggcgaca ccgtggaggt ggagtgctgg atcggcgcca 3660

gcggcaacaa cggcatgcgc cgcgacttcc tggtgcgcga ctgcaagacc ggcgagatcc 3720

tgacccgctg cacctccctg agcgtgctga tgaacacccg cacccgccgc ctgagcacca 3780

tccccgacga ggtgcgcggc gagatcggcc ccgccttcat cgacaacgtg gccgtgaagg 3840

acgacgagat caagaagctg cagaagctga acgactccac cgccgactac atccagggcg 3900

gcctgacccc ccgctggaac gacctggacg tgaaccagca cgtgaacaac ctgaagtacg 3960

tggcctgggt gttcgagacc gtgcccgaca gcatcttcga gtcccaccac atcagctcct 4020

tcaccctgga gtaccgccgc gagtgcaccc gcgactccgt gctgcgcagc ctgaccaccg 4080

tgagcggcgg cagctccgag gccggcctgg tgtgcgacca cctgctgcag ctggagggcg 4140

gcagcgaggt gctgcgcgcc cgcaccgagt ggcgccccaa gctgaccgac tccttccgcg 4200

gcatcagcgt gatccccgcc gagccccgcg tgatggacta caaggaccac gacggcgact 4260

acaaggacca cgacatcgac tacaaggacg acgacgacaa gtgatgactc gaggcagcag 4320

cagctcggat agtatcgaca cactctggac gctggtcgtg tgatggactg ttgccgccac 4380

acttgctgcc ttgacctgtg aatatccctg ccgcttttat caaacagcct cagtgtgttt 4440

gatcttgtgt gtacgcgctt ttgcgagttg ctagctgctt gtgctatttg cgaataccac 4500

ccccagcatc cccttccctc gtttcatatc gcttgcatcc caaccgcaac ttatctacgc 4560

tgtcctgcta tccctcagcg ctgctcctgc tcctgctcac tgcccctcgc acagccttgg 4620

tttgggctcc gcctgtattc tcctggtact gcaacctgta aaccagcact gcaatgctga 4680

tgcacgggaa gtagtgggat gggaacacaa atggaaagct t 4721

<210> 119
<211> 4650
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic polynucleotide

<400> 119
ggtacccgcc tgcaacgcaa gggcagccac agccgctccc acccgccgct gaaccgacac 60

gtgcttgggc gcctgccgcc tgcctgccgc atgcttgtgc tggtgaggct gggcagtgct 120

gccatgctga ttgaggcttg gttcatcggg tggaagctta tgtgtgtgct gggcttgcat 180

gccgggcaat gcgcatggtg gcaagagggc ggcagcactt gctggagctg ccgcggtgcc 240

tccaggtggt tcaatcgcgg cagccagagg gatttcagat gatcgcgcgt acaggttgag 300

cagcagtgtc agcaaaggta gcagtttgcc agaatgatcg gttcagctgt taatcaatgc 360

cagcaagaga aggggtcaag tgcaaacacg ggcatgccac agcacgggca ccggggagtg 420

gaatggcacc accaagtgtg tgcgagccag catcgccgcc tggctgtttc agctacaacg 480

gcaggagtca tccaacgtaa ccatgagctg atcaacactg caatcatcgg gcgggcgtga 540

tgcaagcatg cctggcgaag acacatggtg tgcggatgct gccggctgct gcctgctgcg 600

cacgccgttg agttggcagc aggctcagcc atgcactgga tggcagctgg gctgccactg 660

caatgtggtg gataggatgc aagtggagcg aataccaaac cctctggctg cttgctgggt 720

tgcatggcat cgcaccatca gcaggagcgc atgcgaaggg actggcccca tgcacgccat 780

gccaaaccgg agcgcaccga gtgtccacac tgtcaccagg cccgcaagct ttgcagaacc 840

atgctcatgg acgcatgtag cgctgacgtc ccttgacggc gctcctctcg ggtgtgggaa 900

acgcaatgca gcacaggcag cagaggcggc ggcagcagag cggcggcagc agcggcgggg 960

gccacccttc ttgcggggtc gcgccccagc cagcggtgat gcgctgatcc caaacgagtt 1020

cacattcatt tgcatgcctg gagaagcgag gctggggcct ttgggctggt gcagcccgca 1080

atggaatgcg ggaccgccag gctagcagca aaggcgcctc ccctactccg catcgatgtt 1140

ccatagtgca ttggactgca tttgggtggg gcggccggct gtttctttcg tgttgcaaaa 1200

cgcgccagct cagcaacctg tcccgtgggt cccccgtgcc gatgaaatcg tgtgcacgcc 1260

gatcagctga ttgcccggct cgcgaagtag gcgccctcct ttctgctcgc cctctctccg 1320

tcccgcctct agaatatcaa tgatcgagca ggacggcctc cacgccggct cccccgccgc 1380

ctgggtggag cgcctgttcg gctacgactg ggcccagcag accatcggct gctccgacgc 1440

cgccgtgttc cgcctgtccg cccagggccg ccccgtgctg ttcgtgaaga ccgacctgtc 1500

cggcgccctg aacgagctgc aggacgaggc cgcccgcctg tcctggctgg ccaccaccgg 1560

cgtgccctgc gccgccgtgc tggacgtggt gaccgaggcc ggccgcgact ggctgctgct 1620

gggcgaggtg cccggccagg acctgctgtc ctcccacctg gcccccgccg agaaggtgtc 1680

catcatggcc gacgccatgc gccgcctgca caccctggac cccgccacct gccccttcga 1740

ccaccaggcc aagcaccgca tcgagcgcgc ccgcacccgc atggaggccg gcctggtgga 1800

ccaggacgac ctggacgagg agcaccaggg cctggccccc gccgagctgt tcgcccgcct 1860

gaaggcccgc atgcccgacg gcgaggacct ggtggtgacc cacggcgacg cctgcctgcc 1920

caacatcatg gtggagaacg gccgcttctc cggcttcatc gactgcggcc gcctgggcgt 1980

ggccgaccgc taccaggaca tcgccctggc cacccgcgac atcgccgagg agctgggcgg 2040

cgagtgggcc gaccgcttcc tggtgctgta cggcatcgcc gcccccgact cccagcgcat 2100

cgccttctac cgcctgctgg acgagttctt ctgacaattg gcagcagcag ctcggatagt 2160

atcgacacac tctggacgct ggtcgtgtga tggactgttg ccgccacact tgctgccttg 2220

acctgtgaat atccctgccg cttttatcaa acagcctcag tgtgtttgat cttgtgtgta 2280

cgcgcttttg cgagttgcta gctgcttgtg ctatttgcga ataccacccc cagcatcccc 2340

ttccctcgtt tcatatcgct tgcatcccaa ccgcaactta tctacgctgt cctgctatcc 2400

ctcagcgctg ctcctgctcc tgctcactgc ccctcgcaca gccttggttt gggctccgcc 2460

tgtattctcc tggtactgca acctgtaaac cagcactgca atgctgatgc acgggaagta 2520

gtgggatggg aacacaaatg gaggatcccg cgtctcgaac agagcgcgca gaggaacgct 2580

gaaggtctcg cctctgtcgc acctcagcgc ggcatacacc acaataacca cctgacgaat 2640

gcgcttggtt cttcgtccat tagcgaagcg tccggttcac acacgtgcca cgttggcgag 2700

gtggcaggtg acaatgatcg gtggagctga tggtcgaaac gttcacagcc tagggatatc 2760

gaattccttt cttgcgctat gacacttcca gcaaaaggta gggcgggctg cgagacggct 2820

tcccggcgct gcatgcaaca ccgatgatgc ttcgaccccc cgaagctcct tcggggctgc 2880

atgggcgctc cgatgccgct ccagggcgag cgctgtttaa atagccaggc ccccgattgc 2940

aaagacatta tagcgagcta ccaaagccat attcaaacac ctagatcact accacttcta 3000

cacaggccac tcgagcttgt gatcgcactc cgctaagggg gcgcctcttc ctcttcgttt 3060

cagtcacaac ccgcaaacac tagtatgacg ttcggggtcg ccctcccggc catgggccgc 3120

ggtgtctccc ttccccggcc cagggtcgcg gtgcgcgccc agtcggcgag tcaggttttg 3180

gagagcgggc gcgcccccga ctggtccatg ctgttcgccg tgatcaccac catcttcagc 3240

gccgccgaga agcagtggac caacctggag tggaagccca agcccaagct gccccagctg 3300

ctggacgacc acttcggcct gcacggcctg gtgttccgcc gcaccttcgc catccgctcc 3360

tacgaggtgg gccccgaccg cagcacctcc atcctggccg tgatgaacca catgcaggag 3420

gccaccctga accacgccaa gagcgtgggc atcctgggcg acggcttcgg caccaccctg 3480

gagatgtcca agcgcgacct gatgtgggtg gtgcgccgca cccacgtggc cgtggagcgc 3540

taccccacct ggggcgacac cgtggaggtg gagtgctgga tcggcgccag cggcaacaac 3600

ggcatgcgcc gcgacttcct ggtgcgcgac tgcaagaccg gcgagatcct gacccgctgc 3660

acctccctga gcgtgctgat gaacacccgc acccgccgcc tgagcaccat ccccgacgag 3720

gtgcgcggcg agatcggccc cgccttcatc gacaacgtgg ccgtgaagga cgacgagatc 3780

aagaagctgc agaagctgaa cgactccacc gccgactaca tccagggcgg cctgaccccc 3840

cgctggaacg acctggacgt gaaccagcac gtgaacaacc tgaagtacgt ggcctgggtg 3900

ttcgagaccg tgcccgacag catcttcgag tcccaccaca tcagctcctt caccctggag 3960

taccgccgcg agtgcacccg cgactccgtg ctgcgcagcc tgaccaccgt gagcggcggc 4020

agctccgagg ccggcctggt gtgcgaccac ctgctgcagc tggagggcgg cagcgaggtg 4080

ctgcgcgccc gcaccgagtg gcgccccaag ctgaccgact ccttccgcgg catcagcgtg 4140

atccccgccg agccccgcgt gatggactac aaggaccacg acggcgacta caaggaccac 4200

gacatcgact acaaggacga cgacgacaag tgatgactcg aggcagcagc agctcggata 4260

gtatcgacac actctggacg ctggtcgtgt gatggactgt tgccgccaca cttgctgcct 4320

tgacctgtga atatccctgc cgcttttatc aaacagcctc agtgtgtttg atcttgtgtg 4380

tacgcgcttt tgcgagttgc tagctgcttg tgctatttgc gaataccacc cccagcatcc 4440

ccttccctcg tttcatatcg cttgcatccc aaccgcaact tatctacgct gtcctgctat 4500

ccctcagcgc tgctcctgct cctgctcact gcccctcgca cagccttggt ttgggctccg 4560

cctgtattct cctggtactg caacctgtaa accagcactg caatgctgat gcacgggaag 4620

tagtgggatg ggaacacaaa tggaaagctt 4650

<210> 120
<211> 4653
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic polynucleotide

<400> 120
ggtacccgcc tgcaacgcaa gggcagccac agccgctccc acccgccgct gaaccgacac 60

gtgcttgggc gcctgccgcc tgcctgccgc atgcttgtgc tggtgaggct gggcagtgct 120

gccatgctga ttgaggcttg gttcatcggg tggaagctta tgtgtgtgct gggcttgcat 180

gccgggcaat gcgcatggtg gcaagagggc ggcagcactt gctggagctg ccgcggtgcc 240

tccaggtggt tcaatcgcgg cagccagagg gatttcagat gatcgcgcgt acaggttgag 300

cagcagtgtc agcaaaggta gcagtttgcc agaatgatcg gttcagctgt taatcaatgc 360

cagcaagaga aggggtcaag tgcaaacacg ggcatgccac agcacgggca ccggggagtg 420

gaatggcacc accaagtgtg tgcgagccag catcgccgcc tggctgtttc agctacaacg 480

gcaggagtca tccaacgtaa ccatgagctg atcaacactg caatcatcgg gcgggcgtga 540

tgcaagcatg cctggcgaag acacatggtg tgcggatgct gccggctgct gcctgctgcg 600

cacgccgttg agttggcagc aggctcagcc atgcactgga tggcagctgg gctgccactg 660

caatgtggtg gataggatgc aagtggagcg aataccaaac cctctggctg cttgctgggt 720

tgcatggcat cgcaccatca gcaggagcgc atgcgaaggg actggcccca tgcacgccat 780

gccaaaccgg agcgcaccga gtgtccacac tgtcaccagg cccgcaagct ttgcagaacc 840

atgctcatgg acgcatgtag cgctgacgtc ccttgacggc gctcctctcg ggtgtgggaa 900

acgcaatgca gcacaggcag cagaggcggc ggcagcagag cggcggcagc agcggcgggg 960

gccacccttc ttgcggggtc gcgccccagc cagcggtgat gcgctgatcc caaacgagtt 1020

cacattcatt tgcatgcctg gagaagcgag gctggggcct ttgggctggt gcagcccgca 1080

atggaatgcg ggaccgccag gctagcagca aaggcgcctc ccctactccg catcgatgtt 1140

ccatagtgca ttggactgca tttgggtggg gcggccggct gtttctttcg tgttgcaaaa 1200

cgcgccagct cagcaacctg tcccgtgggt cccccgtgcc gatgaaatcg tgtgcacgcc 1260

gatcagctga ttgcccggct cgcgaagtag gcgccctcct ttctgctcgc cctctctccg 1320

tcccgcctct agaatatcaa tgatcgagca ggacggcctc cacgccggct cccccgccgc 1380

ctgggtggag cgcctgttcg gctacgactg ggcccagcag accatcggct gctccgacgc 1440

cgccgtgttc cgcctgtccg cccagggccg ccccgtgctg ttcgtgaaga ccgacctgtc 1500

cggcgccctg aacgagctgc aggacgaggc cgcccgcctg tcctggctgg ccaccaccgg 1560

cgtgccctgc gccgccgtgc tggacgtggt gaccgaggcc ggccgcgact ggctgctgct 1620

gggcgaggtg cccggccagg acctgctgtc ctcccacctg gcccccgccg agaaggtgtc 1680

catcatggcc gacgccatgc gccgcctgca caccctggac cccgccacct gcccccttcga 1740

ccaccaggcc aagcaccgca tcgagcgcgc ccgcacccgc atggaggccg gcctggtgga 1800

ccaggacgac ctggacgagg agcaccaggg cctggccccc gccgagctgt tcgcccgcct 1860

gaaggcccgc atgcccgacg gcgaggacct ggtggtgacc cacggcgacg cctgcctgcc 1920

caacatcatg gtggagaacg gccgcttctc cggcttcatc gactgcggcc gcctgggcgt 1980

ggccgaccgc taccaggaca tcgccctggc cacccgcgac atcgccgagg agctgggcgg 2040

cgagtgggcc gaccgcttcc tggtgctgta cggcatcgcc gcccccgact cccagcgcat 2100

cgccttctac cgcctgctgg acgagttctt ctgacaattg gcagcagcag ctcggatagt 2160

atcgacacac tctggacgct ggtcgtgtga tggactgttg ccgccacact tgctgccttg 2220

acctgtgaat atccctgccg cttttatcaa acagcctcag tgtgtttgat cttgtgtgta 2280

cgcgcttttg cgagttgcta gctgcttgtg ctatttgcga ataccacccc cagcatcccc 2340

ttccctcgtt tcatatcgct tgcatcccaa ccgcaactta tctacgctgt cctgctatcc 2400

ctcagcgctg ctcctgctcc tgctcactgc ccctcgcaca gccttggttt gggctccgcc 2460

tgtattctcc tggtactgca acctgtaaac cagcactgca atgctgatgc acgggaagta 2520

gtgggatggg aacacaaatg gaggatcccg cgtctcgaac agagcgcgca gaggaacgct 2580

gaaggtctcg cctctgtcgc acctcagcgc ggcatacacc acaataacca cctgacgaat 2640

gcgcttggtt cttcgtccat tagcgaagcg tccggttcac acacgtgcca cgttggcgag 2700

gtggcaggtg acaatgatcg gtggagctga tggtcgaaac gttcacagcc tagggatatc 2760

gaattccttt cttgcgctat gacacttcca gcaaaaggta gggcgggctg cgagacggct 2820

tcccggcgct gcatgcaaca ccgatgatgc ttcgaccccc cgaagctcct tcggggctgc 2880

atgggcgctc cgatgccgct ccagggcgag cgctgtttaa atagccaggc ccccgattgc 2940

aaagacatta tagcgagcta ccaaagccat attcaaacac ctagatcact accacttcta 3000

cacaggccac tcgagcttgt gatcgcactc cgctaagggg gcgcctcttc ctcttcgttt 3060

cagtcacaac ccgcaaacac tagtatggct atcaagacga acaggcagcc tgtggagaag 3120

cctccgttca cgatcgggac gctgcgcaag gccatccccg cgcactgttt cgagcgctcg 3180

gcgcttcgtg ggcgcgcccc cgactggtcc atgctgttcg ccgtgatcac caccatcttc 3240

agcgccgccg agaagcagtg gaccaacctg gagtggaagc ccaagcccaa gctgccccag 3300

ctgctggacg accacttcgg cctgcacggc ctggtgttcc gccgcacctt cgccatccgc 3360

tcctacgagg tgggccccga ccgcagcacc tccatcctgg ccgtgatgaa ccacatgcag 3420

gaggccaccc tgaaccacgc caagagcgtg ggcatcctgg gcgacggctt cggcaccacc 3480

ctggagatgt ccaagcgcga cctgatgtgg gtggtgcgcc gcacccacgt ggccgtggag 3540

cgctacccca cctggggcga caccgtggag gtggagtgct ggatcggcgc cagcggcaac 3600

aacggcatgc gccgcgactt cctggtgcgc gactgcaaga ccggcgagat cctgacccgc 3660

tgcacctccc tgagcgtgct gatgaacacc cgcacccgcc gcctgagcac catccccgac 3720

gaggtgcgcg gcgagatcgg ccccgccttc atcgacaacg tggccgtgaa ggacgacgag 3780

atcaagaagc tgcagaagct gaacgactcc accgccgact acatccaggg cggcctgacc 3840

ccccgctgga acgacctgga cgtgaaccag cacgtgaaca acctgaagta cgtggcctgg 3900

gtgttcgaga ccgtgcccga cagcatcttc gagtcccacc acatcagctc cttcaccctg 3960

gagtaccgcc gcgagtgcac ccgcgactcc gtgctgcgca gcctgaccac cgtgagcggc 4020

ggcagctccg aggccggcct ggtgtgcgac cacctgctgc agctggaggg cggcagcgag 4080

gtgctgcgcg cccgcaccga gtggcgcccc aagctgaccg actccttccg cggcatcagc 4140

gtgatccccg ccgagccccg cgtgatggac tacaaggacc acgacggcga ctacaaggac 4200

cacgacatcg actacaagga cgacgacgac aagtgatgac tcgaggcagc agcagctcgg 4260

atagtatcga cacactctgg acgctggtcg tgtgatggac tgttgccgcc acacttgctg 4320

ccttgacctg tgaatatccc tgccgctttt atcaaacagc ctcagtgtgt ttgatcttgt 4380

gtgtacgcgc ttttgcgagt tgctagctgc ttgtgctatt tgcgaatacc acccccagca 4440

tccccttccc tcgtttcata tcgcttgcat cccaaccgca acttatctac gctgtcctgc 4500

tatccctcag cgctgctcct gctcctgctc actgcccctc gcacagcctt ggtttgggct 4560

ccgcctgtat tctcctggta ctgcaacctg taaaccagca ctgcaatgct gatgcacggg 4620

aagtagtggg atgggaacac aaatggaaag ctt 4653

<210> 121
<211> 4653
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic polynucleotide

<400> 121
ggtacccgcc tgcaacgcaa gggcagccac agccgctccc acccgccgct gaaccgacac 60

gtgcttgggc gcctgccgcc tgcctgccgc atgcttgtgc tggtgaggct gggcagtgct 120

gccatgctga ttgaggcttg gttcatcggg tggaagctta tgtgtgtgct gggcttgcat 180

gccgggcaat gcgcatggtg gcaagagggc ggcagcactt gctggagctg ccgcggtgcc 240

tccaggtggt tcaatcgcgg cagccagagg gatttcagat gatcgcgcgt acaggttgag 300

cagcagtgtc agcaaaggta gcagtttgcc agaatgatcg gttcagctgt taatcaatgc 360

cagcaagaga aggggtcaag tgcaaacacg ggcatgccac agcacgggca ccggggagtg 420

gaatggcacc accaagtgtg tgcgagccag catcgccgcc tggctgtttc agctacaacg 480

gcaggagtca tccaacgtaa ccatgagctg atcaacactg caatcatcgg gcgggcgtga 540

tgcaagcatg cctggcgaag acacatggtg tgcggatgct gccggctgct gcctgctgcg 600

cacgccgttg agttggcagc aggctcagcc atgcactgga tggcagctgg gctgccactg 660

caatgtggtg gataggatgc aagtggagcg aataccaaac cctctggctg cttgctgggt 720

tgcatggcat cgcaccatca gcaggagcgc atgcgaaggg actggcccca tgcacgccat 780

gccaaaccgg agcgcaccga gtgtccacac tgtcaccagg cccgcaagct ttgcagaacc 840

atgctcatgg acgcatgtag cgctgacgtc ccttgacggc gctcctctcg ggtgtgggaa 900

acgcaatgca gcacaggcag cagaggcggc ggcagcagag cggcggcagc agcggcgggg 960

gccacccttc ttgcggggtc gcgccccagc cagcggtgat gcgctgatcc caaacgagtt 1020

cacattcatt tgcatgcctg gagaagcgag gctggggcct ttgggctggt gcagcccgca 1080

atggaatgcg ggaccgccag gctagcagca aaggcgcctc ccctactccg catcgatgtt 1140

ccatagtgca ttggactgca tttgggtggg gcggccggct gtttctttcg tgttgcaaaa 1200

cgcgccagct cagcaacctg tcccgtgggt cccccgtgcc gatgaaatcg tgtgcacgcc 1260

gatcagctga ttgcccggct cgcgaagtag gcgccctcct ttctgctcgc cctctctccg 1320

tcccgcctct agaatatcaa tgatcgagca ggacggcctc cacgccggct cccccgccgc 1380

ctgggtggag cgcctgttcg gctacgactg ggcccagcag accatcggct gctccgacgc 1440

cgccgtgttc cgcctgtccg cccagggccg ccccgtgctg ttcgtgaaga ccgacctgtc 1500

cggcgccctg aacgagctgc aggacgaggc cgcccgcctg tcctggctgg ccaccaccgg 1560

cgtgccctgc gccgccgtgc tggacgtggt gaccgaggcc ggccgcgact ggctgctgct 1620

gggcgaggtg cccggccagg acctgctgtc ctcccacctg gcccccgccg agaaggtgtc 1680

catcatggcc gacgccatgc gccgcctgca caccctggac cccgccacct gccccttcga 1740

ccaccaggcc aagcaccgca tcgagcgcgc ccgcacccgc atggaggccg gcctggtgga 1800

ccaggacgac ctggacgagg agcaccaggg cctggccccc gccgagctgt tcgcccgcct 1860

gaaggcccgc atgcccgacg gcgaggacct ggtggtgacc cacggcgacg cctgcctgcc 1920

caacatcatg gtggagaacg gccgcttctc cggcttcatc gactgcggcc gcctgggcgt 1980

ggccgaccgc taccaggaca tcgccctggc cacccgcgac atcgccgagg agctgggcgg 2040

cgagtgggcc gaccgcttcc tggtgctgta cggcatcgcc gcccccgact cccagcgcat 2100

cgccttctac cgcctgctgg acgagttctt ctgacaattg gcagcagcag ctcggatagt 2160

atcgacacac tctggacgct ggtcgtgtga tggactgttg ccgccacact tgctgccttg 2220

acctgtgaat atccctgccg cttttatcaa acagcctcag tgtgtttgat cttgtgtgta 2280

cgcgcttttg cgagttgcta gctgcttgtg ctatttgcga ataccacccc cagcatcccc 2340

ttccctcgtt tcatatcgct tgcatcccaa ccgcaactta tctacgctgt cctgctatcc 2400

ctcagcgctg ctcctgctcc tgctcactgc ccctcgcaca gccttggttt gggctccgcc 2460

tgtattctcc tggtactgca acctgtaaac cagcactgca atgctgatgc acgggaagta 2520

gtgggatggg aacacaaatg gaggatcccg cgtctcgaac agagcgcgca gaggaacgct 2580

gaaggtctcg cctctgtcgc acctcagcgc ggcatacacc acaataacca cctgacgaat 2640

gcgcttggtt cttcgtccat tagcgaagcg tccggttcac acacgtgcca cgttggcgag 2700

gtggcaggtg acaatgatcg gtggagctga tggtcgaaac gttcacagcc tagggatatc 2760

gaattccttt cttgcgctat gacacttcca gcaaaaggta gggcgggctg cgagacggct 2820

tcccggcgct gcatgcaaca ccgatgatgc ttcgaccccc cgaagctcct tcggggctgc 2880

atgggcgctc cgatgccgct ccagggcgag cgctgtttaa atagccaggc ccccgattgc 2940

aaagacatta tagcgagcta ccaaagccat attcaaacac ctagatcact accacttcta 3000

cacaggccac tcgagcttgt gatcgcactc cgctaagggg gcgcctcttc ctcttcgttt 3060

cagtcacaac ccgcaaacac tagtatggcc accgcatcca ctttctcggc gttcaatgcc 3120

cgctgcggcg acctgcgtcg ctcggcgggc tccgggcccc ggcgcccagc gaggcccctc 3180

cccgtgcgcg ggcgcgcccc cgactggtcc atgctgttcg ccgtgatcac caccatcttc 3240

agcgccgccg agaagcagtg gaccaacctg gagtggaagc ccaagcccaa gctgccccag 3300

ctgctggacg accacttcgg cctgcacggc ctggtgttcc gccgcacctt cgccatccgc 3360

tcctacgagg tgggccccga ccgcagcacc tccatcctgg ccgtgatgaa ccacatgcag 3420

gaggccaccc tgaaccacgc caagagcgtg ggcatcctgg gcgacggctt cggcaccacc 3480

ctggagatgt ccaagcgcga cctgatgtgg gtggtgcgcc gcacccacgt ggccgtggag 3540

cgctacccca cctggggcga caccgtggag gtggagtgct ggatcggcgc cagcggcaac 3600

aacggcatgc gccgcgactt cctggtgcgc gactgcaaga ccggcgagat cctgacccgc 3660

tgcacctccc tgagcgtgct gatgaacacc cgcacccgcc gcctgagcac catccccgac 3720

gaggtgcgcg gcgagatcgg ccccgccttc atcgacaacg tggccgtgaa ggacgacgag 3780

atcaagaagc tgcagaagct gaacgactcc accgccgact acatccaggg cggcctgacc 3840

ccccgctgga acgacctgga cgtgaaccag cacgtgaaca acctgaagta cgtggcctgg 3900

gtgttcgaga ccgtgcccga cagcatcttc gagtcccacc acatcagctc cttcaccctg 3960

gagtaccgcc gcgagtgcac ccgcgactcc gtgctgcgca gcctgaccac cgtgagcggc 4020

ggcagctccg aggccggcct ggtgtgcgac cacctgctgc agctggaggg cggcagcgag 4080

gtgctgcgcg cccgcaccga gtggcgcccc aagctgaccg actccttccg cggcatcagc 4140

gtgatccccg ccgagccccg cgtgatggac tacaaggacc acgacggcga ctacaaggac 4200

cacgacatcg actacaagga cgacgacgac aagtgatgac tcgaggcagc agcagctcgg 4260

atagtatcga cacactctgg acgctggtcg tgtgatggac tgttgccgcc acacttgctg 4320

ccttgacctg tgaatatccc tgccgctttt atcaaacagc ctcagtgtgt ttgatcttgt 4380

gtgtacgcgc ttttgcgagt tgctagctgc ttgtgctatt tgcgaatacc acccccagca 4440

tcccccttccc tcgtttcata tcgcttgcat cccaaccgca acttatctac gctgtcctgc 4500

tatccctcag cgctgctcct gctcctgctc actgcccctc gcacagcctt ggtttgggct 4560

ccgcctgtat tctcctggta ctgcaacctg taaaccagca ctgcaatgct gatgcacggg 4620

aagtagtggg atgggaacac aaatggaaag ctt 4653

<210> 122
<211> 4647
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic polynucleotide

<400> 122
ggtacccgcc tgcaacgcaa gggcagccac agccgctccc acccgccgct gaaccgacac 60

gtgcttgggc gcctgccgcc tgcctgccgc atgcttgtgc tggtgaggct gggcagtgct 120

gccatgctga ttgaggcttg gttcatcggg tggaagctta tgtgtgtgct gggcttgcat 180

gccgggcaat gcgcatggtg gcaagagggc ggcagcactt gctggagctg ccgcggtgcc 240

tccaggtggt tcaatcgcgg cagccagagg gatttcagat gatcgcgcgt acaggttgag 300

cagcagtgtc agcaaaggta gcagtttgcc agaatgatcg gttcagctgt taatcaatgc 360

cagcaagaga aggggtcaag tgcaaacacg ggcatgccac agcacgggca ccggggagtg 420

gaatggcacc accaagtgtg tgcgagccag catcgccgcc tggctgtttc agctacaacg 480

gcaggagtca tccaacgtaa ccatgagctg atcaacactg caatcatcgg gcgggcgtga 540

tgcaagcatg cctggcgaag acacatggtg tgcggatgct gccggctgct gcctgctgcg 600

cacgccgttg agttggcagc aggctcagcc atgcactgga tggcagctgg gctgccactg 660

caatgtggtg gataggatgc aagtggagcg aataccaaac cctctggctg cttgctgggt 720

tgcatggcat cgcaccatca gcaggagcgc atgcgaaggg actggcccca tgcacgccat 780

gccaaaccgg agcgcaccga gtgtccacac tgtcaccagg cccgcaagct ttgcagaacc 840

atgctcatgg acgcatgtag cgctgacgtc ccttgacggc gctcctctcg ggtgtgggaa 900

acgcaatgca gcacaggcag cagaggcggc ggcagcagag cggcggcagc agcggcgggg 960

gccacccttc ttgcggggtc gcgccccagc cagcggtgat gcgctgatcc caaacgagtt 1020

cacattcatt tgcatgcctg gagaagcgag gctggggcct ttgggctggt gcagcccgca 1080

atggaatgcg ggaccgccag gctagcagca aaggcgcctc ccctactccg catcgatgtt 1140

ccatagtgca ttggactgca tttgggtggg gcggccggct gtttctttcg tgttgcaaaa 1200

cgcgccagct cagcaacctg tcccgtgggt cccccgtgcc gatgaaatcg tgtgcacgcc 1260

gatcagctga ttgcccggct cgcgaagtag gcgccctcct ttctgctcgc cctctctccg 1320

tcccgcctct agaatatcaa tgatcgagca ggacggcctc cacgccggct cccccgccgc 1380

ctgggtggag cgcctgttcg gctacgactg ggcccagcag accatcggct gctccgacgc 1440

cgccgtgttc cgcctgtccg cccagggccg ccccgtgctg ttcgtgaaga ccgacctgtc 1500

cggcgccctg aacgagctgc aggacgaggc cgcccgcctg tcctggctgg ccaccaccgg 1560

cgtgccctgc gccgccgtgc tggacgtggt gaccgaggcc ggccgcgact ggctgctgct 1620

gggcgaggtg cccggccagg acctgctgtc ctcccacctg gcccccgccg agaaggtgtc 1680
catcatggcc gacgccatgc gccgcctgca caccctggac cccgccacct gcccccttcga 1740
ccaccaggcc aagcaccgca tcgagcgcgc ccgcacccgc atggaggccg gcctggtgga 1800
ccaggacgac ctggacgagg agcaccaggg cctggccccc gccgagctgt tcgcccgcct 1860
gaaggcccgc atgcccgacg gcgaggacct ggtggtgacc cacggcgacg cctgcctgcc 1920
caacatcatg gtggagaacg gccgcttctc cggcttcatc gactgcggcc gcctgggcgt 1980
ggccgaccgc taccaggaca tcgccctggc cacccgcgac atcgccgagg agctgggcgg 2040
cgagtgggcc gaccgcttcc tggtgctgta cggcatcgcc gcccccgact cccagcgcat 2100
cgccttctac cgcctgctgg acgagttctt ctgacaattg gcagcagcag ctcggatagt 2160
atcgacacac tctggacgct ggtcgtgtga tggactgttg ccgccacact tgctgccttg 2220
acctgtgaat atccctgccg cttttatcaa acagcctcag tgtgtttgat cttgtgtgta 2280
cgcgcttttg cgagttgcta gctgcttgtg ctatttgcga ataccacccc cagcatcccc 2340
ttccctcgtt tcatatcgct tgcatcccaa ccgcaactta tctacgctgt cctgctatcc 2400
ctcagcgctg ctcctgctcc tgctcactgc ccctcgcaca gccttggttt gggctccgcc 2460
tgtattctcc tggtactgca acctgtaaac cagcactgca atgctgatgc acgggaagta 2520
gtgggatggg aacacaaatg gaggatcccg cgtctcgaac agagcgcgca gaggaacgct 2580
gaaggtctcg cctctgtcgc acctcagcgc ggcatacacc acaataacca cctgacgaat 2640
gcgcttggtt cttcgtccat tagcgaagcg tccggttcac acacgtgcca cgttggcgag 2700
gtggcaggtg acaatgatcg gtggagctga tggtcgaaac gttcacagcc tagggatatc 2760
gaattccttt cttgcgctat gacacttcca gcaaaaggta gggcgggctg cgagacggct 2820
tcccggcgct gcatgcaaca ccgatgatgc ttcgaccccc cgaagctcct tcggggctgc 2880
atgggcgctc cgatgccgct ccagggcgag cgctgtttaa atagccaggc ccccgattgc 2940
aaagacatta tagcgagcta ccaaagccat attcaaacac ctagatcact accacttcta 3000
cacaggccac tcgagcttgt gatcgcactc cgctaagggg gcgcctcttc ctcttcgttt 3060
cagtcacaac ccgcaaacac tagtatggct tccgcggcat tcaccatgtc ggcgtgcccc 3120
gcgatgactg gcagggcccc tggggcacgt cgctccggac ggccagtcgc cacccgcctg 3180
agggggcgcg cccccgactg gtccatgctg ttcgccgtga tcaccaccat cttcagcgcc 3240
gccgagaagc agtggaccaa cctggagtgg aagcccaagc ccaagctgcc ccagctgctg 3300
gacgaccact tcggcctgca cggcctggtg ttccgccgca ccttcgccat ccgctcctac 3360
gaggtgggcc ccgaccgcag cacctccatc ctggccgtga tgaaccacat gcaggaggcc 3420
accctgaacc acgccaagag cgtgggcatc ctgggcgacg gcttcggcac caccctggag 3480
atgtccaagc gcgacctgat gtgggtggtg cgccgcaccc acgtggccgt ggagcgctac 3540
cccacctggg gcgacaccgt ggaggtggag tgctggatcg gcgccagcgg caacaacggc 3600
atgcgccgcg acttcctggt gcgcgactgc aagaccggcg agatcctgac ccgctgcacc 3660

tccctgagcg tgctgatgaa cacccgcacc cgccgcctga gcaccatccc cgacgaggtg 3720

cgcggcgaga tcggccccgc cttcatcgac aacgtggccg tgaaggacga cgagatcaag 3780

aagctgcaga agctgaacga ctccaccgcc gactacatcc agggcggcct gacccccccgc 3840

tggaacgacc tggacgtgaa ccagcacgtg aacaacctga agtacgtggc ctgggtgttc 3900

gagaccgtgc ccgacagcat cttcgagtcc caccacatca gctccttcac cctggagtac 3960

cgccgcgagt gcacccgcga ctccgtgctg cgcagcctga ccaccgtgag cggcggcagc 4020

tccgaggccg gcctggtgtg cgaccacctg ctgcagctgg agggcggcag cgaggtgctg 4080

cgcgcccgca ccgagtggcg ccccaagctg accgactcct tccgcggcat cagcgtgatc 4140

cccgccgagc cccgcgtgat ggactacaag gaccacgacg gcgactacaa ggaccacgac 4200

atcgactaca aggacgacga cgacaagtga tgactcgagg cagcagcagc tcggatagta 4260

tcgacacact ctggacgctg gtcgtgtgat ggactgttgc cgccacactt gctgccttga 4320

cctgtgaata tccctgccgc ttttatcaaa cagcctcagt gtgtttgatc ttgtgtgtac 4380

gcgcttttgc gagttgctag ctgcttgtgc tatttgcgaa taccaccccc agcatccccct 4440

tccctcgttt catatcgctt gcatcccaac cgcaacttat ctacgctgtc ctgctatccc 4500

tcagcgctgc tcctgctcct gctcactgcc cctcgcacag ccttggtttg ggctccgcct 4560

gtattctcct ggtactgcaa cctgtaaacc agcactgcaa tgctgatgca cgggaagtag 4620

tgggatggga acacaaatgg aaagctt 4647

<210> 123
<211> 4721
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic polynucleotide

<400> 123
ggtacccgcc tgcaacgcaa gggcagccac agccgctccc acccgccgct gaaccgacac 60

gtgcttgggc gcctgccgcc tgcctgccgc atgcttgtgc tggtgaggct gggcagtgct 120

gccatgctga ttgaggcttg gttcatcggg tggaagctta tgtgtgtgct gggcttgcat 180

gccgggcaat gcgcatggtg gcaagagggc ggcagcactt gctggagctg ccgcggtgcc 240

tccaggtggt tcaatcgcgg cagccagagg gatttcagat gatcgcgcgt acaggttgag 300

cagcagtgtc agcaaaggta gcagtttgcc agaatgatcg gttcagctgt taatcaatgc 360

cagcaagaga aggggtcaag tgcaaacacg ggcatgccac agcacgggca ccggggagtg 420

gaatggcacc accaagtgtg tgcgagccag catcgccgcc tggctgtttc agctacaacg 480

gcaggagtca tccaacgtaa ccatgagctg atcaacactg caatcatcgg gcgggcgtga 540

tgcaagcatg cctggcgaag acacatggtg tgcggatgct gccggctgct gcctgctgcg 600

cacgccgttg agttggcagc aggctcagcc atgcactgga tggcagctgg gctgccactg 660

caatgtggtg gataggatgc aagtggagcg aataccaaac cctctggctg cttgctgggt 720

tgcatggcat cgcaccatca gcaggagcgc atgcgaaggg actggcccca tgcacgccat 780

gccaaaccgg agcgcaccga gtgtccacac tgtcaccagg cccgcaagct ttgcagaacc 840

atgctcatgg acgcatgtag cgctgacgtc ccttgacggc gctcctctcg ggtgtgggaa 900

acgcaatgca gcacaggcag cagaggcggc ggcagcagag cggcggcagc agcggcgggg 960

gccacccttc ttgcggggtc gcgccccagc cagcggtgat gcgctgatcc caaacgagtt 1020

cacattcatt tgcatgcctg gagaagcgag gctggggcct ttgggctggt gcagcccgca 1080

atggaatgcg ggaccgccag gctagcagca aaggcgcctc ccctactccg catcgatgtt 1140

ccatagtgca ttggactgca tttgggtggg gcggccggct gtttctttcg tgttgcaaaa 1200

cgcgccagct cagcaacctg tcccgtgggt cccccgtgcc gatgaaatcg tgtgcacgcc 1260

gatcagctga ttgcccggct cgcgaagtag gcgccctcct ttctgctcgc cctctctccg 1320

tcccgcctct agaatatcaa tgatcgagca ggacggcctc cacgccggct cccccgccgc 1380

ctgggtggag cgcctgttcg gctacgactg ggcccagcag accatcggct gctccgacgc 1440

cgccgtgttc cgcctgtccg cccagggccg ccccgtgctg ttcgtgaaga ccgacctgtc 1500

cggcgccctg aacgagctgc aggacgaggc cgcccgcctg tcctggctgg ccaccaccgg 1560

cgtgccctgc gccgccgtgc tggacgtggt gaccgaggcc ggccgcgact ggctgctgct 1620

gggcgaggtg cccggccagg acctgctgtc ctcccacctg gcccccgccg agaaggtgtc 1680

catcatggcc gacgccatgc gccgcctgca caccctggac cccgccacct gcccccttcga 1740

ccaccaggcc aagcaccgca tcgagcgcgc ccgcacccgc atggaggccg gcctggtgga 1800

ccaggacgac ctggacgagg agcaccaggg cctggccccc gccgagctgt tcgcccgcct 1860

gaaggcccgc atgcccgacg gcgaggacct ggtggtgacc cacggcgacg cctgcctgcc 1920

caacatcatg gtggagaacg gccgcttctc cggcttcatc gactgcggcc gcctgggcgt 1980

ggccgaccgc taccaggaca tcgccctggc cacccgcgac atcgccgagg agctgggcgg 2040

cgagtgggcc gaccgcttcc tggtgctgta cggcatcgcc gcccccgact cccagcgcat 2100

cgccttctac cgcctgctgg acgagttctt ctgacaattg gcagcagcag ctcggatagt 2160

atcgacacac tctggacgct ggtcgtgtga tggactgttg ccgccacact tgctgccttg 2220

acctgtgaat atccctgccg cttttatcaa acagcctcag tgtgtttgat cttgtgtgta 2280

cgcgcttttg cgagttgcta gctgcttgtg ctatttgcga ataccacccc cagcatcccc 2340

ttccctcgtt tcatatcgct tgcatcccaa ccgcaactta tctacgctgt cctgctatcc 2400

ctcagcgctg ctcctgctcc tgctcactgc ccctcgcaca gccttggttt gggctccgcc 2460

tgtattctcc tggtactgca acctgtaaac cagcactgca atgctgatgc acgggaagta 2520

gtgggatggg aacacaaatg gaggatcccg cgtctcgaac agagcgcgca gaggaacgct 2580

gaaggtctcg cctctgtcgc acctcagcgc ggcatacacc acaataacca cctgacgaat 2640

gcgcttggtt cttcgtccat tagcgaagcg tccggttcac acacgtgcca cgttggcgag 2700

gtggcaggtg acaatgatcg gtggagctga tggtcgaaac gttcacagcc tagggatatc 2760

gaattccttt cttgcgctat gacacttcca gcaaaaggta gggcgggctg cgagacggct 2820

tcccggcgct gcatgcaaca ccgatgatgc ttcgaccccc cgaagctcct tcggggctgc 2880

atgggcgctc cgatgccgct ccagggcgag cgctgtttaa atagccaggc ccccgattgc 2940

aaagacatta tagcgagcta ccaaagccat attcaaacac ctagatcact accacttcta 3000

cacaggccac tcgagcttgt gatcgcactc cgctaagggg gcgcctcttc ctcttcgttt 3060

cagtcacaac ccgcaaacgg cgcgccatgg ccaccacctc cctggcctcc gccttctgca 3120

gcatgaaggc cgtgatgctg gcccgcgacg gccgcggcat gaagccccgc tccagcgacc 3180

tgcagctgcg cgccggcaac gcccagacct ccctgaagat gatcaacggc accaagttct 3240

cctacaccga gagcctgaag aagctgcccg actggtccat gctgttcgcc gtgatcacca 3300

ccatcttctc cgccgccgag aagcagtgga ccaacctgga gtggaagccc aagcccaacc 3360

ccccccagct gctggacgac cacttcggcc cccacggcct ggtgttccgc cgcaccttcg 3420

ccatccgcag ctacgaggtg ggccccgacc gctccaccag catcgtggcc gtgatgaacc 3480

acctgcagga ggccgccctg aaccacgcca agtccgtggg catcctgggc gacggcttcg 3540

gcaccaccct ggagatgtcc aagcgcgacc tgatctgggt ggtgaagcgc acccacgtgg 3600

ccgtggagcg ctaccccgcc tggggcgaca ccgtggaggt ggagtgctgg gtgggcgcct 3660

ccggcaacaa cggccgccgc cacgacttcc tggtgcgcga ctgcaagacc ggcgagatcc 3720

tgacccgctg cacctccctg agcgtgatga tgaacacccg cacccgccgc ctgagcaaga 3780

tccccgagga ggtgcgcggc gagatcggcc ccgccttcat cgacaacgtg gccgtgaagg 3840

acgaggagat caagaagccc cagaagctga acgactccac cgccgactac atccagggcg 3900

gcctgacccc ccgctggaac gacctggaca tcaaccagca cgtgaacaac atcaagtacg 3960

tggactggat cctggagacc gtgcccgaca gcatcttcga gagccaccac atctcctcct 4020

tcaccatcga gtaccgccgc gagtgcacca tggacagcgt gctgcagtcc ctgaccaccg 4080

tgagcggcgg ctcctccgag gccggcctgg tgtgcgagca cctgctgcag ctggagggcg 4140

gcagcgaggt gctgcgcgcc aagaccgagt ggcgccccaa gctgaccgac tccttccgcg 4200

gcatcagcgt gatccccgcc gagtccagcg tgatggacta caaggaccac gacggcgact 4260

acaaggacca cgacatcgac tacaaggacg acgacgacaa gtgatgactc gaggcagcag 4320

cagctcggat agtatcgaca cactctggac gctggtcgtg tgatggactg ttgccgccac 4380

acttgctgcc ttgacctgtg aatatccctg ccgcttttat caaacagcct cagtgtgttt 4440

gatcttgtgt gtacgcgctt ttgcgagttg ctagctgctt gtgctatttg cgaataccac 4500

ccccagcatc cccttccctc gtttcatatc gcttgcatcc caaccgcaac ttatctacgc 4560

tgtcctgcta tccctcagcg ctgctcctgc tcctgctcac tgcccctcgc acagccttgg 4620

tttgggctcc gcctgtattc tcctggtact gcaacctgta aaccagcact gcaatgctga 4680

tgcacgggaa gtagtgggat gggaacacaa atggaaagct t 4721

<210> 124
<211> 4650
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic polynucleotide

<400> 124
ggtacccgcc tgcaacgcaa gggcagccac agccgctccc acccgccgct gaaccgacac 60

gtgcttgggc gcctgccgcc tgcctgccgc atgcttgtgc tggtgaggct gggcagtgct 120

gccatgctga ttgaggcttg gttcatcggg tggaagctta tgtgtgtgct gggcttgcat 180

gccgggcaat gcgcatggtg gcaagagggc ggcagcactt gctggagctg ccgcggtgcc 240

tccaggtggt tcaatcgcgg cagccagagg gatttcagat gatcgcgcgt acaggttgag 300

cagcagtgtc agcaaaggta gcagtttgcc agaatgatcg gttcagctgt taatcaatgc 360

cagcaagaga aggggtcaag tgcaaacacg ggcatgccac agcacgggca ccggggagtg 420

gaatggcacc accaagtgtg tgcgagccag catcgccgcc tggctgtttc agctacaacg 480

gcaggagtca tccaacgtaa ccatgagctg atcaacactg caatcatcgg gcgggcgtga 540

tgcaagcatg cctggcgaag acacatggtg tgcggatgct gccggctgct gcctgctgcg 600

cacgccgttg agttggcagc aggctcagcc atgcactgga tggcagctgg gctgccactg 660

caatgtggtg gataggatgc aagtggagcg aataccaaac cctctggctg cttgctgggt 720

tgcatggcat cgcaccatca gcaggagcgc atgcgaaggg actggcccca tgcacgccat 780

gccaaaccgg agcgcaccga gtgtccacac tgtcaccagg cccgcaagct ttgcagaacc 840

atgctcatgg acgcatgtag cgctgacgtc ccttgacggc gctcctctcg ggtgtgggaa 900

acgcaatgca gcacaggcag cagaggcggc ggcagcagag cggcggcagc agcggcgggg 960

gccacccttc ttgcggggtc gcgccccagc cagcggtgat gcgctgatcc caaacgagtt 1020

cacattcatt tgcatgcctg gagaagcgag gctggggcct ttgggctggt gcagcccgca 1080

atggaatgcg ggaccgccag gctagcagca aaggcgcctc ccctactccg catcgatgtt 1140

ccatagtgca ttggactgca tttgggtggg gcggccggct gtttctttcg tgttgcaaaa 1200

cgcgccagct cagcaacctg tcccgtgggt cccccgtgcc gatgaaatcg tgtgcacgcc 1260

gatcagctga ttgcccggct cgcgaagtag gcgccctcct ttctgctcgc cctctctccg 1320

tcccgcctct agaatatcaa tgatcgagca ggacggcctc cacgccggct cccccgccgc 1380

ctgggtggag cgcctgttcg gctacgactg ggcccagcag accatcggct gctccgacgc 1440

cgccgtgttc cgcctgtccg cccagggccg ccccgtgctg ttcgtgaaga ccgacctgtc 1500

cggcgccctg aacgagctgc aggacgaggc cgcccgcctg tcctggctgg ccaccaccgg 1560

cgtgccctgc gccgccgtgc tggacgtggt gaccgaggcc ggccgcgact ggctgctgct 1620

gggcgaggtg cccggccagg acctgctgtc ctcccacctg gcccccgccg agaaggtgtc 1680

catcatggcc gacgccatgc gccgcctgca caccctggac cccgccacct gccccttcga 1740

ccaccaggcc aagcaccgca tcgagcgcgc ccgcacccgc atggaggccg gcctggtgga 1800
ccaggacgac ctggacgagg agcaccaggg cctggccccc gccgagctgt tcgcccgcct 1860
gaaggcccgc atgcccgacg gcgaggacct ggtggtgacc cacggcgacg cctgcctgcc 1920
caacatcatg gtggagaacg gccgcttctc cggcttcatc gactgcggcc gcctgggcgt 1980
ggccgaccgc taccaggaca tcgccctggc cacccgcgac atcgccgagg agctgggcgg 2040
cgagtgggcc gaccgcttcc tggtgctgta cggcatcgcc gcccccgact cccagcgcat 2100
cgccttctac cgcctgctgg acgagttctt ctgacaattg gcagcagcag ctcggatagt 2160
atcgacacac tctggacgct ggtcgtgtga tggactgttg ccgccacact tgctgccttg 2220
acctgtgaat atccctgccg cttttatcaa acagcctcag tgtgtttgat cttgtgtgta 2280
cgcgcttttg cgagttgcta gctgcttgtg ctatttgcga ataccacccc cagcatcccc 2340
ttccctcgtt tcatatcgct tgcatcccaa ccgcaactta tctacgctgt cctgctatcc 2400
ctcagcgctg ctcctgctcc tgctcactgc ccctcgcaca gccttggttt gggctccgcc 2460
tgtattctcc tggtactgca acctgtaaac cagcactgca atgctgatgc acgggaagta 2520
gtgggatggg aacacaaatg gaggatcccg cgtctcgaac agagcgcgca gaggaacgct 2580
gaaggtctcg cctctgtcgc acctcagcgc ggcatacacc acaataacca cctgacgaat 2640
gcgcttggtt cttcgtccat tagcgaagcg tccggttcac acacgtgcca cgttggcgag 2700
gtggcaggtg acaatgatcg gtggagctga tggtcgaaac gttcacagcc tagggatatc 2760
gaattccttt cttgcgctat gacacttcca gcaaaaggta gggcgggctg cgagacggct 2820
tcccggcgct gcatgcaaca ccgatgatgc ttcgaccccc cgaagctcct tcggggctgc 2880
atgggcgctc cgatgccgct ccagggcgag cgctgtttaa atagccaggc ccccgattgc 2940
aaagacatta tagcgagcta ccaaagccat attcaaacac ctagatcact accacttcta 3000
cacaggccac tcgagcttgt gatcgcactc cgctaagggg gcgcctcttc ctcttcgttt 3060
cagtcacaac ccgcaaacac tagtatgacg ttcggggtcg ccctcccggc catgggccgc 3120
ggtgtctccc ttccccggcc cagggtcgcg gtgcgcgccc agtcggcgag tcaggttttg 3180
gagagcgggc gcgcccccga ctggtccatg ctgttcgccg tgatcaccac catcttctcc 3240
gccgccgaga agcagtggac caacctggag tggaagccca agcccaaccc cccccagctg 3300
ctggacgacc acttcggccc ccacggcctg gtgttccgcc gcaccttcgc catccgcagc 3360
tacgaggtgg gccccgaccg ctccaccagc atcgtggccg tgatgaacca cctgcaggag 3420
gccgccctga accacgccaa gtccgtgggc atcctgggcg acggcttcgg caccaccctg 3480
gagatgtcca agcgcgacct gatctgggtg gtgaagcgca cccacgtggc cgtggagcgc 3540
taccccgcct ggggcgacac cgtggaggtg gagtgctggg tgggcgcctc cggcaacaac 3600
ggccgccgcc acgacttcct ggtgcgcgac tgcaagaccg gcgagatcct gacccgctgc 3660
acctccctga gcgtgatgat gaacacccgc acccgccgcc tgagcaagat ccccgaggag 3720
gtgcgcggcg agatcggccc cgccttcatc gacaacgtgg ccgtgaagga cgaggagatc 3780

aagaagcccc agaagctgaa cgactccacc gccgactaca tccagggcgg cctgaccccc 3840

cgctggaacg acctggacat caaccagcac gtgaacaaca tcaagtacgt ggactggatc 3900

ctggagaccg tgcccgacag catcttcgag agccaccaca tctcctcctt caccatcgag 3960

taccgccgcg agtgcaccat ggacagcgtg ctgcagtccc tgaccaccgt gagcggcggc 4020

tcctccgagg ccggcctggt gtgcgagcac ctgctgcagc tggagggcgg cagcgaggtg 4080

ctgcgcgcca agaccgagtg gcgccccaag ctgaccgact ccttccgcgg catcagcgtg 4140

atccccgccg agtccagcgt gatggactac aaggaccacg acggcgacta caaggaccac 4200

gacatcgact acaaggacga cgacgacaag tgatgactcg aggcagcagc agctcggata 4260

gtatcgacac actctggacg ctggtcgtgt gatggactgt tgccgccaca cttgctgcct 4320

tgacctgtga atatccctgc cgcttttatc aaacagcctc agtgtgtttg atcttgtgtg 4380

tacgcgcttt tgcgagttgc tagctgcttg tgctatttgc gaataccacc cccagcatcc 4440

ccttccctcg tttcatatcg cttgcatccc aaccgcaact tatctacgct gtcctgctat 4500

ccctcagcgc tgctcctgct cctgctcact gcccctcgca cagccttggt ttgggctccg 4560

cctgtattct cctggtactg caacctgtaa accagcactg caatgctgat gcacgggaag 4620

tagtgggatg ggaacacaaa tggaaagctt 4650

<210> 125
<211> 4653
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic polynucleotide

<400> 125
ggtacccgcc tgcaacgcaa gggcagccac agccgctccc acccgccgct gaaccgacac 60

gtgcttgggc gcctgccgcc tgcctgccgc atgcttgtgc tggtgaggct gggcagtgct 120

gccatgctga ttgaggcttg gttcatcggg tggaagctta tgtgtgtgct gggcttgcat 180

gccgggcaat gcgcatggtg gcaagagggc ggcagcactt gctggagctg ccgcggtgcc 240

tccaggtggt tcaatcgcgg cagccagagg gatttcagat gatcgcgcgt acaggttgag 300

cagcagtgtc agcaaaggta gcagtttgcc agaatgatcg gttcagctgt taatcaatgc 360

cagcaagaga aggggtcaag tgcaaacacg ggcatgccac agcacgggca ccggggagtg 420

gaatggcacc accaagtgtg tgcgagccag catcgccgcc tggctgtttc agctacaacg 480

gcaggagtca tccaacgtaa ccatgagctg atcaacactg caatcatcgg gcgggcgtga 540

tgcaagcatg cctggcgaag acacatggtg tgcggatgct gccggctgct gcctgctgcg 600

cacgccgttg agttggcagc aggctcagcc atgcactgga tggcagctgg gctgccactg 660

caatgtggtg gataggatgc aagtggagcg aataccaaac cctctggctg cttgctgggt 720

tgcatggcat cgcaccatca gcaggagcgc atgcgaaggg actggcccca tgcacgccat 780

gccaaaccgg agcgcaccga gtgtccacac tgtcaccagg cccgcaagct ttgcagaacc 840

atgctcatgg acgcatgtag cgctgacgtc ccttgacggc gctcctctcg ggtgtgggaa 900

acgcaatgca gcacaggcag cagaggcggc ggcagcagag cggcggcagc agcggcgggg 960

gccacccttc ttgcggggtc gcgccccagc cagcggtgat gcgctgatcc caaacgagtt 1020

cacattcatt tgcatgcctg gagaagcgag gctggggcct ttgggctggt gcagcccgca 1080

atggaatgcg ggaccgccag gctagcagca aaggcgcctc ccctactccg catcgatgtt 1140

ccatagtgca ttggactgca tttgggtggg gcggccggct gtttctttcg tgttgcaaaa 1200

cgcgccagct cagcaacctg tcccgtgggt cccccgtgcc gatgaaatcg tgtgcacgcc 1260

gatcagctga ttgcccggct cgcgaagtag gcgccctcct ttctgctcgc cctctctccg 1320

tcccgcctct agaatatcaa tgatcgagca ggacggcctc cacgccggct cccccgccgc 1380

ctgggtggag cgcctgttcg gctacgactg ggcccagcag accatcggct gctccgacgc 1440

cgccgtgttc cgcctgtccg cccagggccg ccccgtgctg ttcgtgaaga ccgacctgtc 1500

cggcgccctg aacgagctgc aggacgaggc cgcccgcctg tcctggctgg ccaccaccgg 1560

cgtgccctgc gccgccgtgc tggacgtggt gaccgaggcc ggccgcgact ggctgctgct 1620

gggcgaggtg cccggccagg acctgctgtc ctcccacctg gcccccgccg agaaggtgtc 1680

catcatggcc gacgccatgc gccgcctgca caccctggac cccgccacct gccccttcga 1740

ccaccaggcc aagcaccgca tcgagcgcgc ccgcacccgc atggaggccg gcctggtgga 1800

ccaggacgac ctggacgagg agcaccaggg cctggccccc gccgagctgt tcgcccgcct 1860

gaaggcccgc atgcccgacg gcgaggacct ggtggtgacc cacggcgacg cctgcctgcc 1920

caacatcatg gtggagaacg gccgcttctc cggcttcatc gactgcggcc gcctgggcgt 1980

ggccgaccgc taccaggaca tcgccctggc cacccgcgac atcgccgagg agctgggcgg 2040

cgagtgggcc gaccgcttcc tggtgctgta cggcatcgcc gcccccgact cccagcgcat 2100

cgccttctac cgcctgctgg acgagttctt ctgacaattg gcagcagcag ctcggatagt 2160

atcgacacac tctggacgct ggtcgtgtga tggactgttg ccgccacact tgctgccttg 2220

acctgtgaat atccctgccg cttttatcaa acagcctcag tgtgtttgat cttgtgtgta 2280

cgcgcttttg cgagttgcta gctgcttgtg ctatttgcga ataccacccc cagcatcccc 2340

ttccctcgtt tcatatcgct tgcatcccaa ccgcaactta tctacgctgt cctgctatcc 2400

ctcagcgctg ctcctgctcc tgctcactgc ccctcgcaca gccttggttt gggctccgcc 2460

tgtattctcc tggtactgca acctgtaaac cagcactgca atgctgatgc acgggaagta 2520

gtgggatggg aacacaaatg gaggatcccg cgtctcgaac agagcgcgca gaggaacgct 2580

gaaggtctcg cctctgtcgc acctcagcgc ggcatacacc acaataacca cctgacgaat 2640

gcgcttggtt cttcgtccat tagcgaagcg tccggttcac acacgtgcca cgttggcgag 2700

gtggcaggtg acaatgatcg gtggagctga tggtcgaaac gttcacagcc tagggatatc 2760

gaattccttt cttgcgctat gacacttcca gcaaaaggta gggcgggctg cgagacggct 2820

tcccggcgct gcatgcaaca ccgatgatgc ttcgaccccc cgaagctcct tcggggctgc 2880

atgggcgctc cgatgccgct ccagggcgag cgctgtttaa atagccaggc ccccgattgc 2940

aaagacatta tagcgagcta ccaaagccat attcaaacac ctagatcact accacttcta 3000

cacaggccac tcgagcttgt gatcgcactc cgctaagggg gcgcctcttc ctcttcgttt 3060

cagtcacaac ccgcaaacac tagtatggct atcaagacga acaggcagcc tgtggagaag 3120

cctccgttca cgatcgggac gctgcgcaag gccatccccg cgcactgttt cgagcgctcg 3180

gcgcttcgtg ggcgcgcccc cgactggtcc atgctgttcg ccgtgatcac caccatcttc 3240

tccgccgccg agaagcagtg gaccaacctg gagtggaagc ccaagcccaa ccccccccag 3300

ctgctggacg accacttcgg cccccacggc ctggtgttcc gccgcacctt cgccatccgc 3360

agctacgagg tgggccccga ccgctccacc agcatcgtgg ccgtgatgaa ccacctgcag 3420

gaggccgccc tgaaccacgc caagtccgtg ggcatcctgg gcgacggctt cggcaccacc 3480

ctggagatgt ccaagcgcga cctgatctgg gtggtgaagc gcacccacgt ggccgtggag 3540

cgctaccccg cctggggcga caccgtggag gtggagtgct gggtgggcgc ctccggcaac 3600

aacggccgcc gccacgactt cctggtgcgc gactgcaaga ccggcgagat cctgacccgc 3660

tgcacctccc tgagcgtgat gatgaacacc cgcacccgcc gcctgagcaa gatccccgag 3720

gaggtgcgcg gcgagatcgg ccccgccttc atcgacaacg tggccgtgaa ggacgaggag 3780

atcaagaagc cccagaagct gaacgactcc accgccgact acatccaggg cggcctgacc 3840

ccccgctgga acgacctgga catcaaccag cacgtgaaca acatcaagta cgtggactgg 3900

atcctggaga ccgtgcccga cagcatcttc gagagccacc acatctcctc cttcaccatc 3960

gagtaccgcc gcgagtgcac catggacagc gtgctgcagt ccctgaccac cgtgagcggc 4020

ggctcctccg aggccggcct ggtgtgcgag cacctgctgc agctggaggg cggcagcgag 4080

gtgctgcgcg ccaagaccga gtggcgcccc aagctgaccg actccttccg cggcatcagc 4140

gtgatccccg ccgagtccag cgtgatggac tacaaggacc acgacggcga ctacaaggac 4200

cacgacatcg actacaagga cgacgacgac aagtgatgac tcgaggcagc agcagctcgg 4260

atagtatcga cacactctgg acgctggtcg tgtgatggac tgttgccgcc acacttgctg 4320

ccttgacctg tgaatatccc tgccgctttt atcaaacagc ctcagtgtgt ttgatcttgt 4380

gtgtacgcgc ttttgcgagt tgctagctgc ttgtgctatt tgcgaatacc acccccagca 4440

tccccttccc tcgtttcata tcgcttgcat cccaaccgca acttatctac gctgtcctgc 4500

tatccctcag cgctgctcct gctcctgctc actgcccctc gcacagcctt ggtttgggct 4560

ccgcctgtat tctcctggta ctgcaacctg taaaccagca ctgcaatgct gatgcacggg 4620

aagtagtggg atgggaacac aaatggaaag ctt 4653

<210> 126
<211> 3669
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic polynucleotide

<400> 126

cccgtgatca cacaggtgcc ttgcgagcgt gatcacacta ttttgggggt cctacagtac 60

tgaaatggtg agaagtcgta ctgaaatcaa ggatgaacaa tgaaaatggt gctgtggtgg 120

cttctcaaag gtcaagaatc agtcgctcgc gtcaggaaat cgcggcgtca accagcgtgg 180

gcgcggtcag tggccccgca ctggtcacca tagcctctcc tgccacagta gcgatcccct 240

gggcgttcac tctcagcagc ggctgtactg cctcccagat tttcttcttc tggacctgcg 300

ggcgtgagag gatgagcagg gtggggccaa gggctcaatc ctgaacggcc ctcattcggt 360

ttccaatccc acaacacata cccacagcag gtcagaccac gcattcgcac catgcgcacc 420

aaataacgtg tccttacctg attgggtgtg gcaggctccg tggacaggag tgcctcgtcc 480

cccgcccaga cccgctcccc cgtcacggcg gcgtccggga cccgcagcgg ctccaccgcg 540

gtgtgatccg cgttggcggc gcagagcagc atcccagccg atttgacccc gcgcatgctc 600

cgaggcttga ggttggccag caccaccacc cgccggccga caaggtcctc cagggtcacg 660

tgccggacca ggccactcac gatggtgcga gggcccccct cctcgccgag gtcgatctgc 720

tcgacgtaca gactgcgaca tgcgtggcga gtggtcatca gaaggaagca ggtgtgcaga 780

aggggcacgt ggttggtatt gagagtagcc aaagctttgt gccaatcaga aagtcaacgc 840

agctgcctgc ctggctcgcg tacaattcct ttcttgcgct atgacacttc cagcaaaagg 900

tagggcgggc tgcgagacgg cttcccggcg ctgcatgcaa caccgatgat gcttcgaccc 960

cccgaagctc cttcggggct gcatgggcgc tccgatgccg ctccagggcg agcgctgttt 1020

aaatagccag gcccccgatt gcaaagacat tatagcgagc taccaaagcc atattcaaac 1080

acctagatca ctaccacttc tacacaggcc actcgagctt gtgatcgcac tccgctaagg 1140

gggcgcctct tcctcttcgt ttcagtcaca acccgcaaac ggcgcgccat gctgctgcag 1200

gccttcctgt tcctgctggc cggcttcgcc gccaagatca gcgcctccat gacgaacgag 1260

acgtccgacc gccccctggt gcacttcacc cccaacaagg gctggatgaa cgaccccaac 1320

ggcctgtggt acgacgagaa ggacgccaag tggcacctgt acttccagta caacccgaac 1380

gacaccgtct gggggacgcc cttgttctgg ggccacgcca cgtccgacga cctgaccaac 1440

tgggaggacc agcccatcgc catcgccccg aagcgcaacg actccggcgc cttctccggc 1500

tccatggtgg tggactacaa caacacctcc ggcttcttca acgacaccat cgacccgcgc 1560

cagcgctgcg tggccatctg gacctacaac accccggagt ccgaggagca gtacatctcc 1620

tacagcctgg acggcggcta caccttcacc gagtaccaga agaaccccgt gctggccgcc 1680

aactccaccc agttccgcga cccgaaggtc ttctggtacg agccctccca gaagtggatc 1740

atgaccgcgg ccaagtccca ggactacaag atcgagatct actcctccga cgacctgaag 1800

tcctggaagc tggagtccgc gttcgccaac gagggcttcc tcggctacca gtacgagtgc 1860

cccggcctga tcgaggtccc caccgagcag gaccccagca agtcctactg ggtgatgttc 1920

atctccatca accccggcgc cccggccggc ggctccttca accagtactt cgtcggcagc 1980

ttcaacggca cccacttcga ggccttcgac aaccagtccc gcgtggtgga cttcggcaag 2040

gactactacg ccctgcagac cttcttcaac accgacccga cctacgggag cgccctgggc 2100

atcgcgtggg cctccaactg ggagtactcc gccttcgtgc ccaccaaccc ctggcgctcc 2160

tccatgtccc tcgtgcgcaa gttctccctc aacaccgagt accaggccaa cccggagacg 2220

gagctgatca acctgaaggc cgagccgatc ctgaacatca gcaacgccgg cccctggagc 2280

cggttcgcca ccaacaccac gttgacgaag gccaacagct acaacgtcga cctgtccaac 2340

agcaccggca ccctggagtt cgagctggtg tacgccgtca acaccaccca gacgatctcc 2400

aagtccgtgt tcgcggacct ctccctctgg ttcaagggcc tggaggaccc cgaggagtac 2460

ctccgcatgg gcttcgaggt gtccgcgtcc tccttcttcc tggaccgcgg gaacagcaag 2520

gtgaagttcg tgaaggagaa cccctacttc accaaccgca tgagcgtgaa caaccagccc 2580

ttcaagagcg agaacgacct gtcctactac aaggtgtacg gcttgctgga ccagaacatc 2640

ctggagctgt acttcaacga cggcgacgtc gtgtccacca acacctactt catgaccacc 2700

gggaacgccc tgggctccgt gaacatgacg acgggggtgg acaacctgtt ctacatcgac 2760

aagttccagg tgcgcgaggt caagtgatta attaactcga ggcagcagca gctcggatag 2820

tatcgacaca ctctggacgc tggtcgtgtg atggactgtt gccgccacac ttgctgcctt 2880

gacctgtgaa tatccctgcc gcttttatca aacagcctca gtgtgtttga tcttgtgtgt 2940

acgcgctttt gcgagttgct agctgcttgt gctatttgcg aataccaccc ccagcatccc 3000

cttccctcgt ttcatatcgc ttgcatccca accgcaactt atctacgctg tcctgctatc 3060

cctcagcgct gctcctgctc ctgctcactg cccctcgcac agccttggtt tgggctccgc 3120

ctgtattctc ctggtactgc aacctgtaaa ccagcactgc aatgctgatg cacgggaagt 3180

agtgggatgg gaacacaaat ggaaagcttg agctcggtac ccgtacccat cagcatccgg 3240

gtgaatcttg gcctccaaga tatggccaat cctcacatcc agcttggcaa aatcgactag 3300

actgtctgca agtgggaatg tggagcacaa ggttgcttgt agcgatcgac agactggtgg 3360

ggtacattga caggtgggca gcgccgcatc catcgtgcct gacgcgagcg ccgccggttg 3420

ctcgcccgtg cctgccgtca aagagcggca gagaaatcgg gaaccgaaaa cgtcacattg 3480

cctgatgttg ttacatgctg gactagactt tcttggcgtg ggtctgctcc tcgccaggtg 3540

cgcgacgcct cggggctggg tgcgagggag ccgtgcggcc acgcatttga caagacccaa 3600

agctcgcatc tcagacggtc aaccgttcgt attatacatt caacatatgg tacatacgca 3660

aaaagcatg 3669

<210> 127
<211> 39
<212> PRT
<213> Prototheca moriformis

<400> 127

Met Thr Phe Gly Val Ala Leu Pro Ala Met Gly Arg Gly Val Ser Leu
1 5 10 15

Pro Arg Pro Arg Val Ala Val Arg Ala Gln Ser Ala Ser Gln Val Leu
20 25 30

Glu Ser Gly Arg Ala Gln Leu
35

<210> 128
<211> 40
<212> PRT
<213> Prototheca moriformis

<400> 128
Met Ala Ile Lys Thr Asn Arg Gln Pro Val Glu Lys Pro Pro Phe Thr
1 5 10 15

Ile Gly Thr Leu Arg Lys Ala Ile Pro Ala His Cys Phe Glu Arg Ser
20 25 30

Ala Leu Arg Gly Arg Ala Gln Leu
35 40

<210> 129
<211> 36
<212> PRT
<213> Prototheca moriformis

<400> 129
Met Ala Ser Ala Ala Phe Thr Met Ser Ala Cys Pro Ala Met Thr Gly
1 5 10 15

Arg Ala Pro Gly Ala Arg Arg Ser Gly Arg Pro Val Ala Thr Arg Leu
20 25 30

Arg Gly Arg Ala
35

<210> 130
<211> 40
<212> PRT
<213> Chlorella prototheccoides

<400> 130
Met Ala Thr Ala Ser Thr Phe Ser Ala Phe Asn Ala Arg Cys Gly Asp
1 5 10 15

Leu Arg Arg Ser Ala Gly Ser Gly Pro Arg Arg Pro Ala Arg Pro Leu
20 25 30

Pro Val Arg Gly Arg Ala Gln Leu
35 40

<210> 131
<211> 87

<212> PRT
<213> Cuphea hookeriana

<400> 131
Met Val Ala Ala Ala Ala Ser Ser Ala Phe Phe Pro Val Pro Ala Pro
1 5 10 15

Gly Ala Ser Pro Lys Pro Gly Lys Phe Gly Asn Trp Pro Ser Ser Leu
20 25 30

Ser Pro Ser Phe Lys Pro Lys Ser Ile Pro Asn Gly Gly Phe Gln Val
35 40 45

Lys Ala Asn Asp Ser Ala His Pro Lys Ala Asn Gly Ser Ala Val Ser
50 55 60

Leu Lys Ser Gly Ser Leu Asn Thr Gln Glu Asp Thr Ser Ser Ser Pro
65 70 75 80

Pro Pro Arg Thr Phe Leu His
85

<210> 132
<211> 60
<212> PRT
<213> Umbellularia californica

<400> 132
Met Ala Thr Thr Ser Leu Ala Ser Ala Phe Cys Ser Met Lys Ala Val
1 5 10 15

Met Leu Ala Arg Asp Gly Arg Gly Met Lys Pro Arg Ser Ser Asp Leu
20 25 30

Gln Leu Arg Ala Gly Asn Ala Pro Thr Ser Leu Lys Met Ile Asn Gly
35 40 45

Thr Lys Phe Ser Tyr Thr Glu Ser Leu Lys Arg Leu
50 55 60

<210> 133
<211> 60
<212> PRT
<213> Cinnamomum camphora

<400> 133
Met Ala Thr Thr Ser Leu Ala Ser Ala Phe Cys Ser Met Lys Ala Val
1 5 10 15

Met Leu Ala Arg Asp Gly Arg Gly Met Lys Pro Arg Ser Ser Asp Leu
20 25 30

Gln Leu Arg Ala Gly Asn Ala Gln Thr Ser Leu Lys Met Ile Asn Gly
35 40 45

Thr Lys Phe Ser Tyr Thr Glu Ser Leu Lys Lys Leu
50 55 60

<210> 134
<211> 1104
<212> DNA
<213> Prototheca moriformis

<400> 134
atggcaccga ccagcctgct tgccagtact ggcgtctctt ccgcttctct gtggtcctct 60

gcgcgctcca gcgcgtgcgc ttttccggtg gatcatgcgg tccgtggcgc accgcagcgg 120

ccgctgccca tgcagcgccg ctgcttccga acagtggcgg tcagggccgc acccgcggta 180

gccgtccgtc cggaacccgc ccaagagttt tgggagcagc ttgagccctg caagatggcg 240

gaggacaagc gcatcttcct ggaggagcac cgcattcggg gcaacgaggt gggcccctcg 300

cagcggctga cgatcacggc ggtggccaac atcctgcagg aggcggcggg caaccacgcg 360

gtggccatgt ggggccggag ctcggagggt ttcgcgacgg acccggagct gcaggaggcg 420

ggtctcatct ttgtgatgac gcgcatgcag atccaaatgt accgctaccc gcgctggggc 480

gacctgatgc aggtggagac ctggttccag acggcgggca agctaggcgc gcagcgcgag 540

tgggtgctgc gcgacaagct gaccggcgag gcgctgggcg cggccacctc cagctgggtc 600

atgatcaaca tccgcacgcg ccggccgtgc cgcatgcccg agctcgtccg cgtcaagtcg 660

gccttcttcg cgcgcgagcc gccgcgcctg gcgctgccgc ccacggtcac gcgcgccaag 720

ctgcccaaca tcgcgacgcc ggcgccgctg cgcgggcacc gccaggtcgc gcgccgcacc 780

gacatggaca tgaacgggca cgtgaacaac gtggcctacc tggcctggtg cctggaggcc 840

gtgcccgagc acgtcttcag cgactaccac ctctaccaga tggagatcga cttcaaggcc 900

gagtgccacg cgggcgacgt catctcctcc caggccgagc agatcccgcc ccaggaggcg 960

ctcacgcaca acggcgccgg ccgcaacccc tcctgcttcg tccatagcat tctgcgcgcc 1020

gagaccgagc tcgtccgcgc gcgaaccaca tggtcggccc ccatcgacgc gcccgccgcc 1080

aagccgccca aggcgagcca ctga 1104

<210> 135
<211> 1104
<212> DNA
<213> Prototheca moriformis

<400> 135
atggcaccga ccagcctgct tgcccgtact ggcgtctctt ccgcttctct gtgctcctct 60

acgcgctccg gcgcgtgcgc ttttccggtg gatcatgcgg tccgtggcgc accgcagcgg 120

ccgctgccca tgcagcgccg ctgcttccga acagtggctg tcagggccgc acccgcagta 180

gccgtccgtc cggaacccgc ccaagagttt tgggagcagc ttgagccctg caagatggcg 240

gaggacaagc gcatcttcct ggaggagcac cgcattcgtg gcaacgaggt gggcccctcg 300

cagcggctga cgatcacggc ggtggccaac atcctgcagg aggcggcggg caaccacgcg 360

gtggccatgt ggggtcggag ctcggagggt ttcgcgacgg acccggagct gcaggaggcg 420

ggcctcatct ttgtgatgac gcgcatgcag atccaaatgt accgctaccc gcgctggggc 480

gacctgatgc aggtggagac ctggttccag acggcgggca agctaggcgc gcagcgcgag 540

tgggtgctgc gcgacaagct gaccggcgag gcgctgggcg cggccacctc cagctgggtc 600

atgatcaaca tccgcacgcg ccggccgtgc cgcatgcccg agctcgtccg cgtcaagtcg 660

gccttcttcg cgcgcgagcc gccgcgcctg gcgctgccgc ccgcggtcac gcgtgccaag 720

ctgcccaaca tcgcgacgcc ggcgccgctg cgcgggcacc gccaggtcgc gcgccgcacc 780

gacatggaca tgaacggcca cgtgaacaac gttgcctacc tggcctggtg cctggaggcc 840

gtgcccgagc acgtcttcag cgactaccac ctctaccaga tggagatcga cttcaaggcc 900

gagtgccacg cgggcgacgt catctcctcc caggccgagc agatcccgcc ccaggaggcg 960

ctcacgcaca acggcgccgg ccgcaacccc tcctgcttcg tccatagcat tctgcgcgcc 1020

gagaccgagc tcgtccgcgc gcgaaccaca tggtcggccc ccatcgacgc gcccgccgcc 1080

aagccgccca aggcgagcca ctga 1104

<210> 136
<211> 367
<212> PRT
<213> Prototheca moriformis

<400> 136

Met Ala Pro Thr Ser Leu Leu Ala Ser Thr Gly Val Ser Ser Ala Ser
1 5 10 15

Leu Trp Ser Ser Ala Arg Ser Ser Ala Cys Ala Phe Pro Val Asp His
20 25 30

Ala Val Arg Gly Ala Pro Gln Arg Pro Leu Pro Met Gln Arg Arg Cys
35 40 45

Phe Arg Thr Val Ala Val Arg Ala Ala Pro Ala Val Ala Val Arg Pro
50 55 60

Glu Pro Ala Gln Glu Phe Trp Glu Gln Leu Glu Pro Cys Lys Met Ala
65 70 75 80

Glu Asp Lys Arg Ile Phe Leu Glu Glu His Arg Ile Arg Gly Asn Glu
85 90 95

Val Gly Pro Ser Gln Arg Leu Thr Ile Thr Ala Val Ala Asn Ile Leu
100 105 110

Gln Glu Ala Ala Gly Asn His Ala Val Ala Met Trp Gly Arg Ser Ser
115 120 125

Glu Gly Phe Ala Thr Asp Pro Glu Leu Gln Glu Ala Gly Leu Ile Phe
130 135 140

```
Val Met Thr Arg Met Gln Ile Gln Met Tyr Arg Tyr Pro Arg Trp Gly
145                 150                 155                 160

Asp Leu Met Gln Val Glu Thr Trp Phe Gln Thr Ala Gly Lys Leu Gly
                165                 170                 175

Ala Gln Arg Glu Trp Val Leu Arg Asp Lys Leu Thr Gly Glu Ala Leu
            180                 185                 190

Gly Ala Ala Thr Ser Ser Trp Val Met Ile Asn Ile Arg Thr Arg Arg
        195                 200                 205

Pro Cys Arg Met Pro Glu Leu Val Arg Val Lys Ser Ala Phe Phe Ala
    210                 215                 220

Arg Glu Pro Pro Arg Leu Ala Leu Pro Pro Thr Val Thr Arg Ala Lys
225                 230                 235                 240

Leu Pro Asn Ile Ala Thr Pro Ala Pro Leu Arg Gly His Arg Gln Val
                245                 250                 255

Ala Arg Arg Thr Asp Met Asp Met Asn Gly His Val Asn Asn Val Ala
            260                 265                 270

Tyr Leu Ala Trp Cys Leu Glu Ala Val Pro Glu His Val Phe Ser Asp
        275                 280                 285

Tyr His Leu Tyr Gln Met Glu Ile Asp Phe Lys Ala Glu Cys His Ala
    290                 295                 300

Gly Asp Val Ile Ser Ser Gln Ala Glu Gln Ile Pro Pro Gln Glu Ala
305                 310                 315                 320

Leu Thr His Asn Gly Ala Gly Arg Asn Pro Ser Cys Phe Val His Ser
                325                 330                 335

Ile Leu Arg Ala Glu Thr Glu Leu Val Arg Ala Arg Thr Thr Trp Ser
            340                 345                 350

Ala Pro Ile Asp Ala Pro Ala Ala Lys Pro Pro Lys Ala Ser His
        355                 360                 365


<210> 137
<211> 367
<212> PRT
<213> Prototheca moriformis

<400> 137
Met Ala Pro Thr Ser Leu Leu Ala Arg Thr Gly Val Ser Ser Ala Ser
1               5                   10                  15

Leu Cys Ser Ser Thr Arg Ser Gly Ala Cys Ala Phe Pro Val Asp His
            20                  25                  30
```

```
Ala Val Arg Gly Ala Pro Gln Arg Pro Leu Pro Met Gln Arg Arg Cys
        35                  40                  45

Phe Arg Thr Val Ala Val Arg Ala Ala Pro Ala Val Ala Val Arg Pro
    50                  55                  60

Glu Pro Ala Gln Glu Phe Trp Glu Gln Leu Glu Pro Cys Lys Met Ala
65                  70                  75                  80

Glu Asp Lys Arg Ile Phe Leu Glu Glu His Arg Ile Arg Gly Asn Glu
                85                  90                  95

Val Gly Pro Ser Gln Arg Leu Thr Ile Thr Ala Val Ala Asn Ile Leu
            100                 105                 110

Gln Glu Ala Ala Gly Asn His Ala Val Ala Met Trp Gly Arg Ser Ser
        115                 120                 125

Glu Gly Phe Ala Thr Asp Pro Glu Leu Gln Glu Ala Gly Leu Ile Phe
    130                 135                 140

Val Met Thr Arg Met Gln Ile Gln Met Tyr Arg Tyr Pro Arg Trp Gly
145                 150                 155                 160

Asp Leu Met Gln Val Glu Thr Trp Phe Gln Thr Ala Gly Lys Leu Gly
                165                 170                 175

Ala Gln Arg Glu Trp Val Leu Arg Asp Lys Leu Thr Gly Glu Ala Leu
            180                 185                 190

Gly Ala Ala Thr Ser Ser Trp Val Met Ile Asn Ile Arg Thr Arg Arg
        195                 200                 205

Pro Cys Arg Met Pro Glu Leu Val Arg Val Lys Ser Ala Phe Phe Ala
    210                 215                 220

Arg Glu Pro Pro Arg Leu Ala Leu Pro Pro Ala Val Thr Arg Ala Lys
225                 230                 235                 240

Leu Pro Asn Ile Ala Thr Pro Ala Pro Leu Arg Gly His Arg Gln Val
                245                 250                 255

Ala Arg Arg Thr Asp Met Asp Met Asn Gly His Val Asn Asn Val Ala
            260                 265                 270

Tyr Leu Ala Trp Cys Leu Glu Ala Val Pro Glu His Val Phe Ser Asp
        275                 280                 285

Tyr His Leu Tyr Gln Met Glu Ile Asp Phe Lys Ala Glu Cys His Ala
    290                 295                 300
```

```
Gly Asp Val Ile Ser Ser Gln Ala Glu Gln Ile Pro Pro Gln Glu Ala
305                 310                 315                 320

Leu Thr His Asn Gly Ala Gly Arg Asn Pro Ser Cys Phe Val His Ser
                325                 330                 335

Ile Leu Arg Ala Glu Thr Glu Leu Val Arg Ala Arg Thr Thr Trp Ser
            340                 345                 350

Ala Pro Ile Asp Ala Pro Ala Ala Lys Pro Pro Lys Ala Ser His
        355                 360                 365


<210> 138
<211> 328
<212> PRT
<213> Cuphea hookeriana

<400> 138
Gln Leu Pro Asp Trp Ser Arg Leu Leu Thr Ala Ile Thr Thr Val Phe
1               5                   10                  15

Val Lys Ser Lys Arg Pro Asp Met His Asp Arg Lys Ser Lys Arg Pro
            20                  25                  30

Asp Met Leu Val Asp Ser Phe Gly Leu Glu Ser Thr Val Gln Asp Gly
        35                  40                  45

Leu Val Phe Arg Gln Ser Phe Ser Ile Arg Ser Tyr Glu Ile Gly Thr
    50                  55                  60

Asp Arg Thr Ala Ser Ile Glu Thr Leu Met Asn His Leu Gln Glu Thr
65                  70                  75                  80

Ser Leu Asn His Cys Lys Ser Thr Gly Ile Leu Leu Asp Gly Phe Gly
                85                  90                  95

Arg Thr Leu Glu Met Cys Lys Arg Asp Leu Ile Trp Val Val Ile Lys
            100                 105                 110

Met Gln Ile Lys Val Asn Arg Tyr Pro Ala Trp Gly Asp Thr Val Glu
        115                 120                 125

Ile Asn Thr Arg Phe Ser Arg Leu Gly Lys Ile Gly Met Gly Arg Asp
    130                 135                 140

Trp Leu Ile Ser Asp Cys Asn Thr Gly Glu Ile Leu Val Arg Ala Thr
145                 150                 155                 160

Ser Ala Tyr Ala Met Met Asn Gln Lys Thr Arg Arg Leu Ser Lys Leu
                165                 170                 175
```

Pro Tyr Glu Val His Gln Glu Ile Val Pro Leu Phe Val Asp Ser Pro
180 185 190

Val Ile Glu Asp Ser Asp Leu Lys Val His Lys Phe Lys Val Lys Thr
195 200 205

Gly Asp Ser Ile Gln Lys Gly Leu Thr Pro Gly Trp Asn Asp Leu Asp
210 215 220

Val Asn Gln His Val Ser Asn Val Lys Tyr Ile Gly Trp Ile Leu Glu
225 230 235 240

Ser Met Pro Thr Glu Val Leu Glu Thr Gln Glu Leu Cys Ser Leu Ala
245 250 255

Leu Glu Tyr Arg Arg Glu Cys Gly Arg Asp Ser Val Leu Glu Ser Val
260 265 270

Thr Ala Met Asp Pro Ser Lys Val Gly Val Arg Ser Gln Tyr Gln His
275 280 285

Leu Leu Arg Leu Glu Asp Gly Thr Ala Ile Val Asn Gly Ala Thr Glu
290 295 300

Trp Arg Pro Lys Asn Ala Gly Ala Asn Gly Ala Ile Ser Thr Gly Lys
305 310 315 320

Thr Ser Asn Gly Asn Ser Val Ser
325

<210> 139
<211> 322
<212> PRT
<213> Umbellularia californica

<400> 139
Pro Asp Trp Ser Met Leu Phe Ala Val Ile Thr Thr Ile Phe Ser Ala
1 5 10 15

Ala Glu Lys Gln Trp Thr Asn Leu Glu Trp Lys Pro Lys Pro Lys Leu
20 25 30

Pro Gln Leu Leu Asp Asp His Phe Gly Leu His Gly Leu Val Phe Arg
35 40 45

Arg Thr Phe Ala Ile Arg Ser Tyr Glu Val Gly Pro Asp Arg Ser Thr
50 55 60

Ser Ile Leu Ala Val Met Asn His Met Gln Glu Ala Thr Leu Asn His
65 70 75 80

Ala Lys Ser Val Gly Ile Leu Gly Asp Gly Phe Gly Thr Thr Leu Glu
85 90 95

```
Met Ser Lys Arg Asp Leu Met Trp Val Val Arg Arg Thr His Val Ala
            100                 105                 110

Val Glu Arg Tyr Pro Thr Trp Gly Asp Thr Val Glu Val Glu Cys Trp
        115                 120                 125

Ile Gly Ala Ser Gly Asn Asn Gly Met Arg Arg Asp Phe Leu Val Arg
    130                 135                 140

Asp Cys Lys Thr Gly Glu Ile Leu Thr Arg Cys Thr Ser Leu Ser Val
145                 150                 155                 160

Leu Met Asn Thr Arg Thr Arg Arg Leu Ser Thr Ile Pro Asp Glu Val
                165                 170                 175

Arg Gly Glu Ile Gly Pro Ala Phe Ile Asp Asn Val Ala Val Lys Asp
            180                 185                 190

Asp Glu Ile Lys Lys Leu Gln Lys Leu Asn Asp Ser Thr Ala Asp Tyr
        195                 200                 205

Ile Gln Gly Gly Leu Thr Pro Arg Trp Asn Asp Leu Asp Val Asn Gln
    210                 215                 220

His Val Asn Asn Leu Lys Tyr Val Ala Trp Val Phe Glu Thr Val Pro
225                 230                 235                 240

Asp Ser Ile Phe Glu Ser His His Ile Ser Ser Phe Thr Leu Glu Tyr
                245                 250                 255

Arg Arg Glu Cys Thr Arg Asp Ser Val Leu Arg Ser Leu Thr Thr Val
            260                 265                 270

Ser Gly Gly Ser Ser Glu Ala Gly Leu Val Cys Asp His Leu Leu Gln
        275                 280                 285

Leu Glu Gly Gly Ser Glu Val Leu Arg Ala Arg Thr Glu Trp Arg Pro
    290                 295                 300

Lys Leu Thr Asp Ser Phe Arg Gly Ile Ser Val Ile Pro Ala Glu Pro
305                 310                 315                 320

Arg Val
```

<210> 140
<211> 345
<212> PRT
<213> Cinnamomum camphora

<400> 140

```
Pro Asp Trp Ser Met Leu Phe Ala Val Ile Thr Thr Ile Phe Ser Ala
1               5                  10                  15

Ala Glu Lys Gln Trp Thr Asn Leu Glu Trp Lys Pro Lys Pro Asn Pro
            20                  25                  30

Pro Gln Leu Leu Asp Asp His Phe Gly Pro His Gly Leu Val Phe Arg
        35                  40                  45

Arg Thr Phe Ala Ile Arg Ser Tyr Glu Val Gly Pro Asp Arg Ser Thr
    50                  55                  60

Ser Ile Val Ala Val Met Asn His Leu Gln Glu Ala Ala Leu Asn His
65                  70                  75                  80

Ala Lys Ser Val Gly Ile Leu Gly Asp Gly Phe Gly Thr Thr Leu Glu
                85                  90                  95

Met Ser Lys Arg Asp Leu Ile Trp Val Val Lys Arg Thr His Val Ala
            100                 105                 110

Val Glu Arg Tyr Pro Ala Trp Gly Asp Thr Val Glu Val Glu Cys Trp
        115                 120                 125

Val Gly Ala Ser Gly Asn Asn Gly Arg Arg His Asp Phe Leu Val Arg
    130                 135                 140

Asp Cys Lys Thr Gly Glu Ile Leu Thr Arg Cys Thr Ser Leu Ser Val
145                 150                 155                 160

Met Met Asn Thr Arg Thr Arg Arg Leu Ser Lys Ile Pro Glu Glu Val
                165                 170                 175

Arg Gly Glu Ile Gly Pro Ala Phe Ile Asp Asn Val Ala Val Lys Asp
            180                 185                 190

Glu Glu Ile Lys Lys Pro Gln Lys Leu Asn Asp Ser Thr Ala Asp Tyr
        195                 200                 205

Ile Gln Gly Gly Leu Thr Pro Arg Trp Asn Asp Leu Asp Ile Asn Gln
    210                 215                 220

His Val Asn Asn Ile Lys Tyr Val Asp Trp Ile Leu Glu Thr Val Pro
225                 230                 235                 240

Asp Ser Ile Phe Glu Ser His His Ile Ser Ser Phe Thr Ile Glu Tyr
                245                 250                 255

Arg Arg Glu Cys Thr Met Asp Ser Val Leu Gln Ser Leu Thr Thr Val
            260                 265                 270
```

```
Ser Gly Gly Ser Ser Glu Ala Gly Leu Val Cys Glu His Leu Leu Gln
        275                 280                 285

Leu Glu Gly Gly Ser Glu Val Leu Arg Ala Lys Thr Glu Trp Arg Pro
    290                 295                 300

Lys Leu Thr Asp Ser Phe Arg Gly Ile Ser Val Ile Pro Ala Glu Ser
305                 310                 315                 320

Ser Val Met Asp Tyr Lys Asp His Asp Gly Asp Tyr Lys Asp His Asp
                325                 330                 335

Ile Asp Tyr Lys Asp Asp Asp Asp Lys
            340                 345
```

<210> 141
<211> 4
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic peptide

<400> 141
Lys Asp Glu Leu
1

<210> 142
<211> 6
<212> PRT
<213> Unknown

<220>
<223> Description of Unknown: Higher plant fatty acyl-ACP thioesterase sequence

<400> 142
Leu Asp Met Asn Gln His
1   5

<210> 143
<211> 6
<212> PRT
<213> Unknown

<220>
<223> Description of Unknown: Algal fatty acyl-ACP thioesterase sequence

<400> 143
Met Asp Met Asn Gly His
1   5