



NOVEL ORGANOMETALLIC COMPLEXES COMPRISING CATIONIC
HETEROCYCLIC CARBENES AND THEIR PROCESS OF PREPARATION

The present invention relates to novel organometallic complexes comprising cationic
5 heterocyclic carbenes.

It also relates, as means for the preparation of such organometallic complexes, to dicationic heterocyclic precursor compounds.

A further subject-matter of the invention is
10 a process for the preparation of the above organometallic complexes.

Finally, it also comprises the use, as catalysts, of the said organometallic complexes in a number of chemical reactions.

15 Organometallic complexes comprising heterocyclic carbenes have already been disclosed in the prior art. Thus, Patent EP-A-0,721,953 discloses complexes comprising noncationic heterocyclic carbenes derived from imidazoline or from pyrazoline and a metal
20 from Groups 8, 9 and 10 of the Periodic Classification of the Elements. Patent EP-A-0,798,041 discloses complexes comprising noncationic heterocyclic carbenes derived from various compounds having 5-, 6- or 7-membered rings and comprising one or more nitrogen
25 and/or sulphur atoms and a metal chosen from palladium, rhodium, nickel, ruthenium and cobalt.

Such complexes are very stable compounds which additionally have the advantage of not forming



toxic byproducts during their decomposition. However, they exhibit the disadvantage of not being able to be used in an excessively acidic medium and complexes of metals with the 0 degree of oxidation are not soluble
5 in water, which limits the field of their use.

The present invention therefore first of all relates to organometallic complexes comprising heterocyclic carbenes, characterized in that they correspond to the general formula (I)



in which:


- Z^+ represents a 1,2,4-triazolium-5-ylidene ion, at least a portion of the atoms of the ring of which are substituted by hydrocarbon-comprising
15 radicals,
- L represents a ligand, which can be ionic or neutral,
- M represents a metal chosen from the transition elements from Groups 1b, 2b, 3b, 4b, 5b, 6b,
20 7b and 8 of the Periodic Classification of the Elements as published in "Handbook of Chemistry and Physics, 51st Edition" (1970-1971) of The Chemical Rubber Company,
- X^- represents an organic or inorganic
25 anion,
- m represents an integer from 1 to 6,
- n represents an integer from 0 to 5,
- the sum of m and of n is equal to or less

than 6,

- Y represents an anion or a cation such that the metal complex is electrically neutral.

For convenience, compounds of formula (I) also including the Y counterion are known as organometallic complexes or metal complexes in the present text.

The 1,2,4-triazolium-5-ylidene ions Z^+ correspond to the general formula (II):



10 in which R_1 , R_2 , R_3 and R_4 , which are identical or different, represent:

- a linear or branched alkyl radical or a cycloalkyl radical,

- an aryl radical,

15 - an alkyl radical which comprises one or more substituents, such as an aryl radical, an alkoxy radical, a halogen atom or a hydrophilic group, for example:

-COOM₁, -SO₃M₁, or -PO₃M₁, M₁ representing an
20 inorganic or organic cationic residue chosen from a proton, cations derived from alkali metals or alkaline

earth metals, or ammonium cations $-N(R)_4$, in the formula of which cations the R symbols, which are identical or different, represent a hydrogen atom or an alkyl radical having from 1 to 4 carbon atoms,

5 $-N(R)_3Y_a$, in the formula of which the R symbols, which are identical or different, represent a hydrogen atom or an alkyl radical having from 1 to 12 carbon atoms and Y_a represents an inorganic or organic anion,

10 $-OH$,

- an aryl or aralkyl or cycloalkyl radical which comprises, on the ring, one or more substituents, such as an alkyl radical, an alkoxy radical, a halogen atom or a hydrophilic group, for example:

15 $-COOM_1$, $-SO_3M_1$, or $-PO_3M_1$, M_1 representing an inorganic or organic cationic residue chosen from a proton, cations derived from alkali metals or alkaline earth metals, or ammonium cations $-N(R)_4$, in the formula of which cations the R symbols, which are
20 identical or different, represent a hydrogen atom or an alkyl radical having from 1 to 12 carbon atoms,

$-N(R)_3Y_a$, in the formula of which the R symbols, which are identical or different, represent a hydrogen atom or an alkyl radical having from 1 to 12
25 carbon atoms and Y_a represents an inorganic or organic anion,

$-OH$,

R_3 also represents a hydrogen atom,

R_1 and/or R_4 can also each represent an organic ligand residue bonded to the nitrogen atom of the 1,2,4-triazolium-5-ylidene ion via an alkylene or arylene radical, it being possible for the said ligand residue
5 to be a phosphite residue, a phosphonite residue, a phosphinite residue, a phosphine residue or an amine residue which is tertiary and aliphatic, cycloaliphatic, aromatic or heterocyclic and which acts as ligand L to the metal M.

10 The preceding organic ligand residues can derive, for example, from phenyl or alkyl phosphites, substituted or unsubstituted by substituents as defined above, or from phenyl or alkyl phosphinites, substituted or unsubstituted by substituents as defined
15 above, or from phenyl or alkyl phosphonites, substituted or unsubstituted by substituents as defined above, or from phenylphosphines or alkylphosphines, substituted or unsubstituted by substituents as defined above.

20 In such a case, the 1,2,4-triazolium-5-ylidene ion substituted by a R_1 and/or R_4 radical as defined above is a bidentate or tridentate ligand.

 L represents an ionic ligand, such as a halide or a cyanide, or a neutral ligand, such as
25 carbon monoxide, an isonitrile, a phosphine, an organic phosphite, a phosphonate or a phosphonite.

 M preferably represents a metal chosen from nickel, cobalt, iron, ruthenium, rhodium, palladium,


osmium, iridium, platinum, copper, silver, gold, zinc, cadmium or mercury.

These metals can be in their various degrees of oxidation, including the 0 degree of oxidation.

5 The X^- anions are organic or inorganic anions, such as, for example, halides, but are preferably chosen from weakly coordinating anions which are defined with respect to the metal M under consideration. They predominantly constitute the
10 counterions of the 1,2,4-triazolium-5-ylidene ions.

Mention may be made, as non-limiting examples of such weakly coordinating anions, of trifluoromethanesulphonate, tetrafluoroborate, hexafluorophosphate, aluminium tetrachloride, aluminium
15 tetrabromide, aluminium tetrafluoride, aluminium tetraiodide, gallium tetrachloride, gallium tetrabromide, gallium tetrafluoride or gallium tetraiodide.

Another subject-matter of the present
20 invention is the use of dicationic heterocyclic compounds as precursors in the preparation of the organometallic complexes of general formula (I). These dicationic heterocyclic compounds correspond to the general formula (III)



in which

- the R_1 , R_2 , R_3 and R_4 symbols have the meanings shown above for the 1,2,4-triazolium-5-ylidene ions of formula (II),

5 $-\text{X}^-$ represents an anion such as those shown for the general formula (I).

The dicationic heterocyclic compounds of general formula (III) can be prepared by reacting a compound of formula XR_2 , in which X and R_2 have the meanings shown above, with a triazole derivative

10 carrying the R_1 , R_3 and R_4 substituents. The preparation can also be carried out in several stages, each stage consisting in reacting a triazole derivative with a compound of formula XR_1 or XR_3 or XR_4 , depending on the

15 substituent which it is desired to introduce into the formula of the triazole employed, in order to obtain the dicationic heterocyclic compound of formula (III). The reaction conditions can be adapted from the procedure described in the article by T.J. Curphey and

20 K.S. Prasad, Journal of Org. Chem., 1972, 37, 2259.

The organometallic complexes of formula (I) can be prepared by reacting a dicationic heterocyclic compound of general formula (III) with a compound of

the metal M in the presence of an inorganic or organic base. Such a base can be the associated anion of the compound of the metal M or can be added independently to the reaction mixture. This reaction can be carried
5 out in liquid medium, generally by adding the compound of the metal M and, if appropriate, the inorganic or organic base to the solution or suspension of the compound (III). Very clearly, the order of introduction of the various reactants can be modified. The reaction
10 can be carried out at room temperature or, preferably, at a temperature of 25°C to 150°C, for example. This temperature can conveniently be that of the reflux of the liquid in which the synthesis is carried out. The separation of the complex of formula (I) can be carried
15 out according to the methods commonly used in chemistry, for example by filtration, by centrifuging or by extraction. The complexes of formula (I) and the dicationic heterocyclic compounds of formula (III) are generally characterized by Nuclear Magnetic Resonance
20 (NMR) or by Infrared or by X-ray diffraction.

The organometallic complexes of formula (I) can be used as catalysts in numerous chemical reactions.


The majority of the organometallic complexes
25 of formula (I) as well as the dicationic heterocyclic compounds of formula (III) have little solubility in or are insoluble in the majority of nonpolar or only slightly polar organic solvents, such as alkanes,

cycloalkanes, haloalkanes, aromatic hydrocarbons, alkyларomatic hydrocarbons or ethers. On the other hand, they are soluble in polar solvents. Thus, the dicationic compounds are soluble in particular in
5 tetrahydrofuran, dimethyl sulphoxide (DMSO), nitriles (for example, acetonitrile) or water. Thus, the organometallic complexes of the invention are soluble in particular in nitriles (for example, acetonitrile), DMSO or water. These properties make possible easier
10 separation of these various compounds, in particular by liquid/liquid extraction. Furthermore, they make it possible to carry out catalysis in two-phase medium.

Mention may be made, as non-limiting-examples of such reactions, of the hydrosilylation of alkenes or
15 of alkynes, in particular in the presence of ruthenium complexes, the hydrosilylation of ketones in the presence of ruthenium or rhodium complexes, the Heck reaction in the presence of palladium complexes, the hydrogenation of olefins, aldehydes, acids, enamides
20 and nitroaromatic compounds in the presence of ruthenium, rhodium, platinum or palladium complexes, the hydroformylation and the hydrocarbonylation of olefins in the presence of rhodium complexes, the hydrocyanation of olefins in the presence of nickel
25 complexes, the synthesis of furan in the presence of ruthenium complexes, the metathesis of olefins in the presence of ruthenium complexes, or the polymerization of acrylates in the presence of nickel complexes.

EXAMPLES OF THE SYNTHESIS OF PRECURSORS OF FORMULA
(III)

EXAMPLE 1. Synthesis of the compound (IIIa):



By following the procedure described in J.


5 Org. Chem., 1972, 37, page 2259, 1H-1,2,4-methyltriazole is methylated twice (at the 2 and 4 positions of the ring) with methyl trifluoromethanesulphonate at reflux in 1,2-dichloroethane with stirring.

10 The brown precipitate formed is filtered off and then washed with 1,2-dichloroethane until a white powder is obtained. This solid (IIIa) exhibits a melting point of 160-162°C.


It is characterized by:

15 - ¹H NMR (d₆-DMSO): δ 4.22 (s, 3H, CH₃N), 4.50 (s, 6H, CH₃N), 10.79 (s, 2H, H_{cycle}).
 - ¹³C NMR (H) (CD₃CN): δ 38.67 (s, CH₃N), 38.87 (s, 2CH₃N), 120.35 (q, ¹J_{CF} = 320.1 Hz, CF₃), 147.72 (s, 2C_{cycle}).

EXAMPLE 2. Synthesis of the compound (IIIb):



The synthesis is carried out in two stages. First of all, an isopropyl group is attached at the 4 position (R₄ radical) by stoichiometric addition of 1H-1,2,4-
 5 methyltriazole to a solution of isopropyl trifluoromethanesulphonate in 1,2-dichloroethane (at reflux with stirring for 3 h). The compound of following formula:



is obtained, with a yield of 91%, in the form of a
 10 yellow oil.

2.5 ml of methyl trifluoromethanesulphonate (22 mmol) are added to a solution of 5 g (18 mmol) of this compound in 20 ml of 1,2-dichloroethane and the mixture is brought to reflux with stirring for 24 h. A
 15 brown precipitate is formed, which precipitate is filtered off and then washed with 1,2-dichloroethane


until a white powder is obtained.

This solid (IIIb) exhibits a melting point of 163-165°C and corresponds to a yield of 73% with respect to the intermediate compound charged.


5 It is characterized by:

- ^1H NMR (CD_3CN): δ 1.65 (d, $^3J_{\text{HH}} = 6.58$ Hz, 6H, CH_3CH), 4.34 (s, 3H, CH_3N), 5.03 (sept, $^3J_{\text{HH}} = 6.58$ Hz, 1H, $[\text{CH}_3]_2\text{CHN}$), 10.22 (s, 2H, H_{cycle}).
 - ^{13}C NMR (CD_3CN): δ 21.51 (s, CH_3CH), 38.71 (s, CH_3N), 58.88 (s, $[\text{CH}_3]_2\text{CHN}$), 120.35 (q, $^1J_{\text{CF}} = 320.1$ Hz, CF_3), 145.22 (s, 2C_{cycle}).
- 10

EXAMPLE 3. Synthesis of the compound (IIIc):



The synthesis is carried out in two stages.
First of all, the compound of following formula:



15 is prepared according to the procedure described in J. Prakt. Chem., 1988, 330, 3, 325.


This compound is heated at reflux in methyl

trifluoromethanesulphonate with stirring for 4 days. A brown precipitate is formed, which precipitate is filtered off and then washed with 1,2-dichloroethane until a white powder is obtained. This solid (IIIc) is obtained with a yield of 50% with respect to the compound charged in the second stage.


It is characterized by:

- ^1H NMR (CD_3CN): δ 4.22 (s, 3H, CH_3N), 7.4-8.4 (unresolved peak, phenyl H), 10.34 (s, 1H, H_{cycle}).

10 EXAMPLE 4. Synthesis of the compound (IIIId):



The synthesis is carried out in two stages. First of all, the compound of following formula:



is prepared according to the procedure described in J. Prakt. Chem., 1988, 330, 3, 325.

15 This compound is heated at reflux with 3 equivalents of methyl trifluoromethanesulphonate in


1,2-dichloroethane with stirring for 2 days. A brown precipitate is formed, which precipitate is filtered off and then washed with 1,2-dichloroethane until a white powder is obtained. This solid (IIIId) is obtained
 5 with a yield of 69% with respect to the compound charged in the second stage and it exhibits a melting point of 190.5-191.3°C.

It is characterized by:

- ^1H NMR (CD_3CN): δ 3.04 (s, 3H, CH_3), 4.19 (s, 3H, CH_3N), 7.8-8.1 (unresolved peak, 10H, 1 phenyl H),
 10 10.51 (s, 1H, H_{cycle}).
- ^{13}C NMR (CD_3CN): δ 12.31 (s, CH_3C), 38.04 (s, CH_3N), 126.70 (s, 2 phenyl CH), 128.02 (s, phenyl C), 128.49 (s, 2 phenyl CH), 130.07 (s, phenyl C), 131.56 (s, 2
 15 phenyl CH), 131.83 (s, 2 phenyl CH), 134.04 (s, 1 phenyl CH), 135.88 (s, 1 phenyl CH), 120.35 (q, $^1J_{\text{CF}} = 320.1$ Hz, CF_3), 145.58 (s, CH_{cycle}), 157.66 (s, $\text{C}_{\text{cycle}}\text{CH}_3$).


The proton coupled spectrum shows a coupling constant
 20 $^1J_{\text{CH}} = 240$ Hz for the CH of the triazole ring.

EXAMPLE 5. Synthesis of the compound (IIIe):



The synthesis is carried out in two stages.

First of all, the compound of following formula:




is prepared according to the procedure described in J. Prakt. Chem., 1988, 330, 3, 325.

This compound is heated at reflux with 3
 5 equivalents of methyl trifluoromethanesulphonate in
 1,2-dichloroethane with stirring for 2 days. A brown
 precipitate is formed, which precipitate is filtered
 off and then washed with 1,2-dichloroethane until a
 white powder is obtained. This solid (IIIe) is obtained
 10 with a yield of 71% with respect to the compound
 charged in the second stage and it exhibits a melting
 point of 180.0°C.


It is characterized by:

- ^1H NMR (CD_3CN): δ 4.08 (s, 3H, CH_3N), 4.20 (s, 3H,
 15 CH_3N), 4.46 (s, 3H, CH_3N), 7.8-8.1 (unresolved peak,
 10H, 5 phenyl H), 10.16 (s, 1H, H_{cycle}).
- ^{13}C NMR (CD_3CN): δ 37.88 (s, CH_3N), 38.36 (s, CH_3N),
 39.42 (s, CH_3N), 130.78 (s, phenyl C), 131.97 (s, 2
 phenyl CH), 132.15 (s, 2 phenyl CH), 137.35 (s, phenyl
 20 CH), 131.83 (s, 2 phenyl CH), 120.35 (q, $^1J_{\text{CF}} = 320.1$
 Hz, CF_3), 147.35 (s, CH_{cycle}), 156.48 (s, C_{cycle}).

EXAMPLE 6. Synthesis of the compound (IIIIf):



The synthesis is carried out in two stages.
First of all, the compound of following formula:



is prepared according to the procedure described in
5 J. Prakt. Chem., 1988, 330, 3, 325.

This compound is heated at reflux with 3
equivalents of methyl trifluoromethanesulphonate in
1,2-dichloroethane with stirring for 2 days. A brown
precipitate is formed, which precipitate is filtered
10 off and then washed with 1,2-dichloroethane until a
white powder is obtained. This solid (IIIIf) is obtained
with a yield of 95% with respect to the compound
charged in the second stage and it exhibits a melting
point of 194.4°C.

15 It is characterized by:


- ¹H NMR (d₆-acetone): δ 3.35 (s, 3H, CH₃C), 4.46 (s,
3H, CH₃N), 4.69 (s, 3H, CH₃N), 4.77 (s, 3H, CH₃N), 10.77
(s, 1H, H_{cycle}).

17

- ^{13}C NMR (d_6 -acetone): δ 10.53 (s, $\underline{\text{C}}\text{H}_3\text{C}$), 35.91 (s, CH_3N), 36.26 (s, CH_3N), 37.99 (s, CH_3N), 120.35 (q, $^1J_{\text{CF}} = 320.1$ Hz, CF_3), 145.35 (s, CH_{cycle}), 157.31 (s, $\underline{\text{C}}_{\text{cycle}}\text{CH}_3$).

5 EXAMPLES OF THE SYNTHESIS OF THE ORGANOMETALLIC COMPLEXES OF FORMULA (I)

EXAMPLE 7. Synthesis of the Ag(I) compound (Ia):




0.4 g (2.4 mmol) of Ag acetate is added to a suspension of 1 g of dicationic compound (IIIa), prepared in Example 1, in 20 ml of tetrahydrofuran (THF). The mixture is stirred for 2 h at reflux. A THF-immiscible brown oil is formed, which oil is washed several times with 10 ml of THF. After evaporating the solvent, a light-sensitive brown solid is obtained (a yield of 82% with respect to the dicationic compound (IIIa)).

This complex exhibits the following characteristics:

- ^1H NMR (CD_3CN): δ 4.01 (s, 3H, NCH_3), 4.19 (s, 3H, $\text{CH}_3\text{N-NCH}_3$), 4.22 (s, 3H, $\text{CH}_3\text{N-NCH}_3$), 9.72 (s, 1H, CH).
 - ^{13}C NMR (CD_3CN): δ 37.2 (NCH_3), 38.5 (NCH_3), 38.6 (NCH_3), 121.1 (q, $^1J_{\text{CF}} = 320.1$ Hz, CF_3), 144.8 (CH).

189.2 (CAG).

EXAMPLE 8. Synthesis of the Hg(II) compound (Ib):



0.16 g of Hg diacetate (which was dried
beforehand under vacuum for 12 h at 50°C) and 1 ml of
5 dimethyl sulphoxide (DMSO) are added to a suspension of
0.43 g of dicationic compound (IIIe), prepared in
Example 5, in 20 ml of tetrahydrofuran (THF). The
mixture is stirred for 24 h at ambient temperature. A
brown precipitate is formed which is soluble solely in
10 highly polar solvents. This solid is washed several
times with THF in order to remove any traces of acetic
acid. The complex thus obtained is recrystallized by
slow diffusion of ether into a methanolic solution of
the said complex. The recrystallized complex (Ib)
15 exhibits a melting point of 85.4-87°C. A yield of 86%
is obtained with respect to the dicationic compound
(IIIe).


This complex exhibits the following
characteristics:

20 - ¹H NMR (CD₃OD): δ 4.70 (s, 12H, NCH₃, ⁴J_{H-Hg} = 12.8),
4.29 (s, 6H, CH₃N, ⁴J_{H-Hg} = 8.5 Hz), 7.9-8.2 (unresolved
peak, 10H, phenyl H).

- ^{13}C NMR (CD_3OD): δ 38.79 ($\text{CH}_3\text{N}-\text{NCH}_3$), 41.75 (NCH_3),
 121.1 (q, $^1J_{\text{CF}} = 320.1$ Hz, CF_3), 156.88 ($\text{C}-\text{Ph}$,
 ($^3J+^2J$) $_{\text{C}-\text{Hg}} = 86$ Hz), 183.78 (CHg , $^1J_{\text{CHg}} = 3275$ Hz).

This structure was also established by X-ray
 5 diffraction.

EXAMPLE 9. Synthesis of the Ni(II) compound (Ic):



2.4 g of dicationic compound (IIIe), prepared
 in Example 5, are added to a suspension of 0.442 g of
 Ni diacetate and 0.75 g of Na iodide in 50 ml of THF.
 10 The mixture is stirred at reflux for 2 h. A coloured
 precipitate is formed, which precipitate is filtered
 off and washed several times with THF. The complex thus
 obtained is recrystallized from an acetone/ether
 mixture at -30°C . The recrystallized complex (Ic) is
 15 red and it exhibits a melting point of 310.7°C
 (decomposition: the crystals become black). A yield of
 85% is obtained with respect to the dicationic compound
 (IIIe).

This complex exhibits the following
 20 characteristics:


- ^1H NMR (d_6 -acetone): δ 7.8-8.0 (broad, 10H, phenyl H),
 4.88-4.84 (2s, 6H, NCH_3), 4.54-4.49 (2s, 6H, CH_3N),

4.42-4.41 (2s, 6H, CH₃N).

- ¹³C NMR (CD₃OD): δ 37.0 (CH₃N), 37.9 (CH₃N), 38.4 (CH₃N), 121.1 (q, ¹J_{CF} = 320.1 Hz, CF₃), 130.5, 130.8 and 134.8 (phenyl C), 153.4 (C-Ph), 188.2 (C-Ni).

5 Its structure was also established by X-ray diffraction.

EXAMPLE 10. Synthesis of the Ni(II) compound (Id):



0.425 g of dicationic compound (IIIIf), prepared in Example 6, is added to a suspension of
 10 0.090 g of Ni diacetate and 0.15 g of Na iodide in 25 ml of THF. The mixture is stirred at reflux for 2 h. A coloured precipitate is formed, which precipitate is filtered off and washed several times with THF. The complex thus obtained is recrystallized from an
 15 acetone/ether mixture at -30°C. The recrystallized complex (Id) is red and it exhibits a melting point of 300-305°C (decomposition: the crystals become black). A yield of 79% is obtained with respect to the dicationic compound (IIIIf).

20 This complex exhibits the following characteristics:

- ¹H NMR (d₆-acetone): δ 4.88-4.84 (2s, 6H, NCH₃), 4.54-


4.49 (2s, 6H, CH₃N), 4.42-4.41 (2s, 6H, CH₃N), 2.63 (s, 6H, C-CH₃).

- ¹³C NMR (CD₃OD): δ 13.86 (C-CH₃), 36.1 (CH₃N), 37.6 (CH₃N), 38.5 (CH₃N), 121.1 (q, ¹J_{CF} = 320.1 Hz, CF₃),

5 154.8 (C-Me), 187.6 (C-Ni).

Its structure was also established by X-ray diffraction.

EXAMPLE 11. Synthesis of the Ni(II) compound (Ie):




0.935 g of dicationic compound (IIIa),
 10 prepared in Example 1, is added to a suspension of
 0.200 g of Ni diacetate and 0.34 g of Na iodide in 30
 ml of THF. The mixture is stirred at reflux for 2 h. A
 coloured precipitate is formed, which precipitate is
 filtered off and washed several times with THF. The
 15 complex thus obtained is recrystallized by slow
 diffusion of ether into a solution of the said complex
 in acetonitrile. Recrystallization is carried out from
 an acetone/ether mixture at -30°C. The recrystallized
 complex (Ie) is red. A yield of 70% is obtained with
 20 respect to the dicationic compound (IIIa).

This complex exhibits the following characteristics:

- ^1H NMR (d_6 -acetone): δ 4.38-4.35 (2s, 6H, CH_3N), 4.68-4.65 (2s, 6H, CH_3N), 4.99-4.96 (2s, 6H, NCH_3), 9.32 (s, 2H, CH).

EXAMPLE 12. Synthesis of the Pd(II) compound (If):



0.472 g of dicationic compound (IIIe), prepared in Example 5, is added to a suspension of 0.112 g of Pd diacetate and 0.30 g of Na iodide in 60 ml of THF. The mixture is stirred at reflux for 2 h. A coloured precipitate is formed, which precipitate is filtered off and washed several times with ether. The complex thus obtained is recrystallized at -30°C from an acetone/ether mixture. It exists in the form of yellow crystals. The recrystallized complex (If) exhibits a melting point of $211-213^\circ\text{C}$ (decomposition: the crystals become brown). During the reaction, all the trifluoromethanesulphonate anions were substituted by iodide anions. A yield of 51% is obtained with respect to the dicationic compound (IIIe).

This complex exhibits the following characteristics:


- ^1H NMR (d_6 -DMSO): δ 7.8-8.1 (broad, 10H, phenyl H), 4.49-4.53 (2s, 6H, NCH_3), 4.21 (s, 6H, CH_3N), 4.10-4.13

(2s, 6H, CH₃N).

- ¹³C NMR (d₆-DMSO): δ 37.2 (CH₃N), 38.3 (CH₃N), 39.9 (CH₃N), 130.1, 130.9 and 134.5 (phenyl C), 152.4 (C-Ph), 174.5 (C-Pd).

5 Its structure was also established by X-ray diffraction.

EXAMPLE 13. Synthesis of the Pd(II) compound (Ig):




0.425 g of dicationic compound (IIIIf), prepared in Example 6, is added to a suspension of
 10 0.112 g of Pd diacetate and 0.30 g of Na iodide in 60 ml of THF. The mixture is stirred at reflux for 2 h. A coloured precipitate is formed, which precipitate is filtered off and washed several times with ether. The complex thus obtained is recrystallized at -30°C from
 15 an acetone/ether mixture. It exists in the form of red crystals. The recrystallized complex (Ig) exhibits a melting point of 206-212°C (decomposition: the crystals become brown). During the reaction, all the trifluoromethanesulphonate anions were substituted by
 20 iodide anions. A yield of 60% is obtained with respect to the dicationic compound (IIIIf).

This complex exhibits the following

characteristics:

- ^1H NMR (d_6 -DMSO): δ 4.41-4.40 (2s, 6H, NCH_3), 4.29 (s, 6H, CH_3N), 4.16-4.15 (2s, 6H, CH_3N), 2.95 (6H, C- CH_3).

EXAMPLE 14. Synthesis of the Ni(0) compound (Ih):



5 Synthetic Method A

1 ml of nickel tetracarbonyl is diluted in 40 ml of anhydrous and degassed acetonitrile. 380 μl of pyridine and 660 μl of triethylamine are added to the solution obtained.

10 A solution of 2.0 g of dicationic compound (IIIIf), prepared in Example 6, in 60 ml of anhydrous and degassed acetonitrile is added dropwise using a dropping funnel. Gas is observed to be given off and the solution turns yellow in colour. When all the

15 solution has been introduced, stirring is maintained for a further 1 h. The volume of the reaction solution is subsequently reduced by two thirds by evaporation of the solvent under vacuum and then 200 ml of degassed ether are added. A precipitate of the complex (Ih) is

20 then formed, which precipitate is filtered off and washed several times with ether. The complex thus obtained is recrystallized from an acetone/ether

mixture at -30°C . The complex (Ih) exists in the form of yellow crystals. Its melting point could not be determined exactly (decomposition: the crystals become brown). A yield of 26% is obtained with respect to the
5 dicationic compound (IIIIf).

This complex exhibits the following characteristics:

- ^1H NMR (d_6 -DMSO): δ 4.00 (s, 6H, NCH_3), 3.99 (s, 6H, CH_3N), 3.78 (s, 6H, CH_3N), 2.72 (6H, $\text{C}-\text{CH}_3$).

10 Bands at 1982 cm^{-1} and 1906 cm^{-1} (CO) are observed by Infrared spectrometry.

Its structure was also established by X-ray diffraction.

The following were found by elemental
15 analysis:


- 28.99% of C (theory 28.89%)
- 3.36% of H (theory 3.63%)
- 12.39% of N (theory 12.62%).

Synthetic Method B

20 150 mg of activated metallic Zn powder (10 equivalents) are added to a solution of 0.2 g of the organometallic complex (Id), prepared in Example 10, in 10 ml of anhydrous and degassed acetonitrile. The solution is placed under a carbon monoxide atmosphere.
25 The reaction begins at the same time as the stirring is begun. The solution becomes yellow. The solution is filtered using a tube and then it is treated in the way

described for Method A (from the addition of ether in order to precipitate the organometallic complex). The yield obtained is 48% with respect to the organometallic complex (Id) charged. The same analyses and characterizations are carried out as in Method A
 5 and the results are the same.

EXAMPLE 15. Synthesis of the Rh(I) compound (Ij):



A solution of 0.277 g of di(carbonyl)rhodium acetylacetonate in 10 ml of THF is added to a
 10 suspension of 0.456 g of dicationic compound (IIIIf), prepared in Example 6, and of 0.16 g of NaI in 40 ml of THF. The mixture is stirred at 50°C for 2 days. A brown solid is recovered after evaporation of the solvent and washing several times with ether. The yield is 70% with
 15 respect to the dicationic compound (IIIIf).


This complex exhibits the following characteristics:

- ^1H NMR (d_6 -DMSO): δ 4.47 (s, 3H, NCH_3), 4.28 (s, 3H, CH_3N), 4.20 (s, 3H, CH_3N), 2.96 (s, 3H, C-CH_3).
- 20 - ^{13}C NMR (d_6 -DMSO): δ 10.5 (s, C-CH_3), 35.2 (s, NCH_3), 36.8 (s, CH_3N), 37.9 (s, CH_3N), 154.2 (s, C-CH_3), 183.0 (d, $^1\text{J}(\text{Rh-C}) = 82.9$ Hz, CO), 188.1 (d, $^1\text{J}(\text{Rh-C}) = 42.2$

Hz, carbene C).

A band at 1975 cm^{-1} (CO) is observed by Infrared.

EXAMPLE 16. Synthesis of the Rh(I) compound (Ik):



5 A solution of 0.200 g of di(carbonyl)rhodium acetylacetonate in 10 ml of THF is added to a suspension of 0.16 g of dicationic compound (IIIa), prepared in Example 1, in 10 ml of THF. The mixture is stirred at 50°C for 15 h. A brown solid precipitate is formed, which precipitate is washed several times with
10 ether. The yield of complex (Ik) is 67% with respect to the dicationic compound (IIIa).

This complex exhibits the following characteristics:

- 15 - ^1H NMR (d_6 -DMSO): δ 4.20 (s, 3H, NCH₃), 4.22 (s, 3H, CH₃N), 4.25 (s, 3H, CH₃N), 9.66 (s, 1H, CH).
- ^{13}C NMR (d_6 -DMSO): δ 38.6 (s, NCH₃), 38.7 (s, NCH₃), 39.1 (s, CH₃N), 37.9 (s, CH₃N), 121.1 (q, $^1J_{\text{CF}} = 320.1$ Hz, CF₃), 145.5 (s, CH), 186.1 (d, $^1J(\text{Rh}-\text{C}) = 76$ Hz, CO), 187.9 (d, $^1J(\text{Rh}-\text{C}) = 43.3$ Hz, carbene C).
- 20

Bands at 2029 (i) cm^{-1} and 1985 (m) cm^{-1} (CO) are observed by Infrared.

EXAMPLE 17. Catalysis of the hydroformylation of styrene by the Rh(I) compound (Ij)

The hydroformylation reaction of styrene is carried out in a 100 ml stainless steel reactor comprising a jacket allowing heating by circulation of oil at a temperature controlled using a thermostat. The lid of the reactor is equipped with a valve for introducing or discharging gases under pressure, with an accurate manometer, with a safety valve and with a ball valve allowing the transfer of solutions with a syringe. A Teflon[®] pot is fitted inside the autoclave and comprises a magnetic bar for stirring the reactants.

The Rh(I) complex (Ij) prepared in Example 15 is charged (40 mg) to the Teflon[®] pot, as well as 100 mg of triphenylphosphine (5 equivalents with respect to the catalyst). The autoclave is assembled and purged three times with argon. 15 ml of tetrahydrofuran, 5 ml of styrene and 60 mg of triethylamine (4 equivalents with respect to the catalyst) are subsequently charged. A carbon monoxide pressure (15 bar) and a hydrogen pressure (15 bar) are then established. The autoclave is heated to 60°C with magnetic stirring. After reacting for 4 h at this temperature, the autoclave is quickly cooled in an acetone/liquid nitrogen mixture and depressurization is carried out over one hour. A sample of the reaction mixture is analysed by vapour phase chromatography

(VPC) .

The following results are obtained:

- degree of conversion of the styrene: 95%
- selectivity for aldehydes: 97%
- 5 - branched aldehyde/linear aldehyde ratio: 95/5
- frequency of rotation of the catalyst: 75 h^{-1}
(calculated by average with regard to the reaction time, taking an identical VPC response factor for all the products) .

10 EXAMPLE 18 Catalysis of the hydroformylation of styrene by the Rh(I) compound (Ik)

Example 17 is repeated with 40 mg of complex (Ik) prepared in Example 16 and without employing triphenylphosphine.


15 After reacting for 3 h at 80°C , the autoclave is quickly cooled in an acetone/liquid nitrogen mixture and depressurization is carried out over one hour. A sample of the reaction mixture is analysed by vapour phase chromatography (VPC) :

20 The following results are obtained:

- degree of conversion of the styrene: 96%
- selectivity for aldehydes: 100%
- branched aldehyde/linear aldehyde ratio: 25/75
- frequency of rotation of the catalyst: 79 h^{-1}
25 (calculated by average with regard to the reaction time, taking an identical VPC response factor for all the products) .

CLAIMS


1. Organometallic complexes comprising
heterocyclic carbenes, characterized in that they
5 correspond to the general formula (I)



in which:

- Z^+ represents a 1,2,4-triazolium-5-ylidene
ion, at least a portion of the atoms of the ring of
10 which are substituted by hydrocarbon-comprising
radicals,
- L represents a ligand, which can be ionic
or neutral,
- M represents a metal chosen from the
15 transition elements from Groups 1b, 2b, 3b, 4b, 5b, 6b,
7b and 8 of the Periodic Classification of the
Elements,
- X^- represents an organic or inorganic anion,
- m represents an integer from 1 to 6,
- 20 - n represents an integer from 0 to 5,
- the sum of m and of n is equal to or less
than 6,
- Y represents an anion or a cation such that
the metal complex is electrically neutral.

2. Organometallic complexes according to claim 1, characterized in that the 1,2,4-triazolium-5-ylidene ions Z^+ correspond to the general formula (II):



5 in which R_1 , R_2 , R_3 and R_4 , which are identical or different, represent:

- a linear or branched alkyl radical or a cycloalkyl radical,
- an aryl radical,
- 10 - an alkyl radical which comprises one or more substituents, such as an aryl radical, an alkoxy radical, a halogen atom or a hydrophilic group, for example:

-COOM₁, -SO₃M₁, or -PO₃M₁, M₁ representing an

15 inorganic or organic cationic residue chosen from a proton, cations derived from alkali metals or alkaline earth metals, or ammonium cations -N(R)₄, in the formula of which cations the R symbols, which are identical or different, represent a hydrogen atom or an alkyl

20 radical having from 1 to 4 carbon atoms,

-N(R)₃Y_a, in the formula of which the R symbols, which are identical or different, represent a

hydrogen atom or an alkyl radical having from 1 to 12 carbon atoms and Y_a represents an inorganic or organic anion,

-OH,

- 5 - an aryl or aralkyl or cycloalkyl radical which comprises, on the ring, one or more substituents, such as an alkyl radical, an alkoxy radical, a halogen atom or a hydrophilic group, for example:

-COOM₁, -SO₃M₁, or -PO₃M₁, M₁ representing an
10 inorganic or organic cationic residue chosen from a proton, cations derived from alkali metals or alkaline earth metals, or ammonium cations -N(R)₄, in the formula of which cations the R symbols, which are identical or different, represent a hydrogen atom or an alkyl
15 radical having from 1 to 12 carbon atoms,

-N(R)₃Y_a, in the formula of which the R symbols, which are identical or different, represent a hydrogen atom or an alkyl radical having from 1 to 12 carbon atoms and Y_a represents an inorganic or organic
20 anion,

-OH,

R₃ also represents a hydrogen atom,

R₁ and/or R₄ can also each represent an organic ligand residue bonded to the nitrogen atom of the 1,2,4-
25 triazolium-5-ylidene ion via an alkylene or arylene radical, it being possible for the said ligand residue

to be a phosphite residue, a phosphonite residue, a phosphinite residue, a phosphine residue or an amine residue which is tertiary and aliphatic, cycloaliphatic, aromatic or heterocyclic and which acts
5 as ligand L to the metal M.

3. Organometallic complexes according to either of claims 1 and 2, characterized in that R_1 and/or R_4 represent an organic ligand residue bonded to the nitrogen atom of the 1,2,4-triazolium-5-ylidene ion
10 via an alkylene or phenylene radical, the said ligand residue deriving from phenyl or alkyl phosphites, substituted or unsubstituted by substituents as defined above, or from phenyl or alkyl phosphonites, substituted or unsubstituted by substituents as defined
15 above, or from phenyl or alkyl phosphinites, substituted or unsubstituted by substituents as defined above, or from phenylphosphines or alkylphosphines, substituted or unsubstituted by substituents as defined above.

20 4. Organometallic complexes according to one of claims 1 to 3, characterized in that L represents an ionic ligand chosen from halides or cyanides or a neutral ligand chosen from carbon monoxide, an isonitrile, a phosphine, an organic
25 phosphite, a phosphonate or a phosphonite.


5. Organometallic complexes according to one of claims 1 to 4, characterized in that M represents a metal chosen from nickel, cobalt, iron, ruthenium, rhodium, palladium, osmium, iridium, platinum, copper, silver, gold, zinc, cadmium or mercury.

6. Organometallic complexes according to one of claims 1 to 5, characterized in that the X^- anions are weakly coordinating anions and predominantly constitute the counterions of the 1,2,4-triazolium-5-ylidene ions.

7. Organometallic complexes according to claim 6, characterized in that the weakly coordinating anions X^- are chosen from trifluoromethanesulphonate, tetrafluoroborate, hexafluorophosphate, aluminium tetrachloride, aluminium tetrabromide, aluminium tetrafluoride, aluminium tetraiodide, gallium tetrachloride, gallium tetrabromide, gallium tetrafluoride or gallium tetraiodide.

8. Process for the preparation of the organometallic complexes of general formula (I), characterized in that a dicationic heterocyclic compound of general formula (III):

35



in which

- the R_1 , R_2 , R_3 and R_4 symbols have the meanings shown above for the 1,2,4-triazolium-5-ylidene ions of formula (II),

- X^- represents an anion such as those shown for the general formula (I),

is reacted with a compound of the metal M in the presence of an inorganic or organic base.


9. Preparation process according to claim 8, characterized in that the reaction is in liquid medium, the compound of the metal M and, if appropriate, the inorganic or organic base being added to the solution or suspension of the compound (III) or the preparation being carried out with another order of introduction of the reactants, at room temperature or, preferably, at a temperature of 25°C to 150°C.

10. Use of the organometallic complexes of formula (I) according to one of claims 1 to 7, as catalysts of chemical reactions.

11. Use according to claim 10 in the hydrosilylation of alkenes or of alkynes, in particular

in the presence of ruthenium complexes, the hydrosilylation of ketones in the presence of ruthenium or rhodium complexes, the Heck reaction in the presence of palladium complexes, the hydrogenation of olefins, aldehydes, acids, enamides and nitroaromatic compounds in the presence of ruthenium, rhodium, platinum or palladium complexes, the hydroformylation and the hydrocarbonylation of olefins in the presence of rhodium complexes, the hydrocyanation of olefins in the presence of nickel complexes, the synthesis of furan in the presence of ruthenium complexes, the metathesis of olefins in the presence of ruthenium complexes, or the polymerization of acrylates in the presence of nickel complexes.

12. Dicationic heterocyclic compounds, characterized in that they correspond to the following general formula (III):



in which R_1 , R_2 , R_3 and R_4 , which are identical or different, represent:

- a linear or branched alkyl radical or a cycloalkyl radical,

- an aryl radical,

- an alkyl radical which comprises one or more substituents, such as an aryl radical, an alkoxy radical, a halogen atom or a hydrophilic group, for

5 example:

-COOM₁, -SO₃M₁, or -PO₃M₁, M₁ representing an inorganic or organic cationic residue chosen from a proton, cations derived from alkali metals or alkaline earth metals, or ammonium cations -N(R)₄, in the formula
10 of which cations the R symbols, which are identical or different, represent a hydrogen atom or an alkyl radical having from 1 to 4 carbon atoms,

-N(R)₃Y_a, in the formula of which the R symbols, which are identical or different, represent a
15 hydrogen atom or an alkyl radical having from 1 to 12 carbon atoms and Y_a represents an inorganic or organic anion,

-OH,

- an aryl or aralkyl or cycloalkyl radical which
20 comprises, on the ring, one or more substituents, such as an alkyl radical, an alkoxy radical, a halogen atom or a hydrophilic group, for example:

-COOM₁, -SO₃M₁, or -PO₃M₁, M₁ representing an inorganic or organic cationic residue chosen from a
25 proton, cations derived from alkali metals or alkaline earth metals, or ammonium cations -N(R)₄, in the formula

of which cations the R symbols, which are identical or different, represent a hydrogen atom or an alkyl radical having from 1 to 12 carbon atoms,

-N(R)₃Y_a, in the formula of which the R symbols, which are identical or different, represent a hydrogen atom or an alkyl radical having from 1 to 12 carbon atoms and Y_a represents an inorganic or organic anion,

-OH,

R₃ also represents a hydrogen atom,

R₁ and/or R₄ can also each represent an organic ligand residue bonded to the nitrogen atom of the 1,2,4-triazolium-5-ylidene ion via an alkylene or arylene radical, it being possible for the said ligand residue to be a phosphite residue, a phosphonite residue, a phosphinite residue, a phosphine residue or an amine residue which is tertiary and aliphatic, cycloaliphatic, aromatic or heterocyclic,


X⁻ represent organic or inorganic anions.

13. Compounds according to claim 12, characterized in that the X⁻ anions are weakly coordinating anions and predominantly constitute the counterions of the 1,2,4-triazolium-5-ylidene ions.

14. Compounds according to claim 13, characterized in that the weakly coordinating anions X⁻ are chosen from trifluoromethanesulphonate,

tetrafluoroborate, hexafluorophosphate, aluminium
 tetrachloride, aluminium tetrabromide, aluminium
 tetrafluoride, aluminium tetraiodide, gallium
 tetrachloride, gallium tetrabromide, gallium
 5 tetrafluoride or gallium tetraiodide.

15. Process for the manufacture of a
 dicationic heterocyclic compound according to one of
 claims 12 to 14, characterized in that it consists in
 reacting a compound of formula XR_2 , in which X and R_2
 10 have the meanings indicated above, with a monocationic
 triazoline derivative of following general formula



in which:

R_1 , R_3 , R_4 and X have the meanings indicated
 15 above.