

STYRYLPYRIDINE DERIVATIVES AND THEIR USE FOR BINDING AND IMAGING AMYLOID PLAQUES

BACKGROUND OF THE INVENTION

Field of the Invention

[0001] This invention relates to novel styrylpyridine compounds, the uses thereof in diagnostic imaging and inhibiting amyloid- β aggregation, and methods of making these compounds.

Background Art

[0002] Alzheimer's disease (AD) is a progressive neurodegenerative disorder characterized by cognitive decline, irreversible memory loss, disorientation, and language impairment. Postmortem examination of AD brain sections reveals abundant senile plaques (SPs) composed of amyloid- β (A β) peptides and numerous neurofibrillary tangles (NFTs) formed by filaments of highly phosphorylated tau proteins (for recent reviews and additional citations see Ginsberg, S. D., *et al.*, "Molecular Pathology of Alzheimer's Disease and Related Disorders," in *Cerebral Cortex: Neurodegenerative and Age-related Changes in Structure and Function of Cerebral Cortex*, Kluwer Academic/Plenum, NY (1999), pp. 603-654; Vogelsberg-Ragaglia, V., *et al.*, "Cell Biology of Tau and Cytoskeletal Pathology in Alzheimer's Disease," *Alzheimer's Disease*, Lippincot, Williams & Wilkins, Philadelphia, PA (1999), pp. 359-372).

[0003] Amyloidosis is a condition characterized by the accumulation of various insoluble, fibrillar proteins in the tissues of a patient. An amyloid deposit is formed by the aggregation of amyloid proteins, followed by the further combination of aggregates and/or amyloid proteins. Formation and accumulation of aggregates of β -amyloid (A β) peptides in the brain are critical factors in the development and progression of AD.

[0004] In addition to the role of amyloid deposits in Alzheimer's disease, the presence of amyloid deposits has been shown in diseases such as Mediterranean fever, Muckle-Wells syndrome, idiopathic myeloma, amyloid

polyneuropathy, amyloid cardiomyopathy, systemic senile amyloidosis, amyloid polyneuropathy, hereditary cerebral hemorrhage with amyloidosis, Down's syndrome, Scrapie, Creutzfeldt-Jacob disease, Kuru, Gerstmann-Straussler-Scheinker syndrome, medullary carcinoma of the thyroid, Isolated atrial amyloid, β_2 -microglobulin amyloid in dialysis patients, inclusion body myositis, β_2 -amyloid deposits in muscle wasting disease, and Islets of Langerhans diabetes Type II insulinoma.

- [0005] The fibrillar aggregates of amyloid peptides, $A\beta_{1-40}$ and $A\beta_{1-42}$, are major metabolic peptides derived from amyloid precursor protein found in senile plaques and cerebrovascular amyloid deposits in AD patients (Xia, W., *et al.*, *J. Proc. Natl. Acad. Sci. U.S.A.* 97:9299-9304 (2000)). Prevention and reversal of $A\beta$ plaque formation are being targeted as a treatment for this disease (Selkoe, D., *J. JAMA* 283:1615-1617 (2000); Wolfe, M.S., *et al.*, *J. Med. Chem.* 41:6-9 (1998); Skovronsky, D.M., and Lee, V.M., *Trends Pharmacol. Sci.* 21:161-163 (2000)).
- [0006] Familial AD (FAD) is caused by multiple mutations in the A precursor protein (APP), presenilin 1 (PS1) and presenilin 2 (PS2) genes (Ginsberg, S. D., *et al.*, "Molecular Pathology of Alzheimer's Disease and Related Disorders," in *Cerebral Cortex: Neurodegenerative and Age-related Changes in Structure and Function of Cerebral Cortex*, Kluwer Academic/Plenum, NY (1999), pp. 603-654; Vogelsberg-Ragaglia, V., *et al.*, "Cell Biology of Tau and Cytoskeletal Pathology in Alzheimer's Disease," *Alzheimer's Disease*, Lippincot, Williams & Wilkins, Philadelphia, PA (1999), pp. 359-372).
- [0007] While the exact mechanisms underlying AD are not fully understood, all pathogenic FAD mutations studied thus far increase production of the more amyloidogenic 42-43 amino-acid long form of the $A\beta$ peptide. Thus, at least in FAD, dysregulation of $A\beta$ production appears to be sufficient to induce a cascade of events leading to neurodegeneration. Indeed, the amyloid cascade hypothesis suggests that formation of extracellular fibrillar $A\beta$ aggregates in the brain may be a pivotal event in AD pathogenesis (Selkoe, D. J., "Biology of β -amyloid Precursor Protein and the Mechanism of Alzheimer's Disease,"

Alzheimer's Disease, Lippincot Williams & Wilkins, Philadelphia, PA (1999), pp. 293-310; Selkoe, D. J., *J. Am. Med. Assoc.* 283:1615-1617 (2000); Naslund, J., *et al.*, *J. Am. Med. Assoc.* 283:1571-1577 (2000); Golde, T. E., *et al.*, *Biochimica et Biophysica Acta* 1502:172-187 (2000)).

[0008] Various approaches in trying to inhibit the production and reduce the accumulation of fibrillar A β in the brain are currently being evaluated as potential therapies for AD (Skovronsky, D. M. and Lee, V. M., *Trends Pharmacol. Sci.* 21:161-163 (2000); Vassar, R., *et al.*, *Science* 286:735-741 (1999); Wolfe, M. S., *et al.*, *J. Med. Chem.* 41:6-9 (1998); Moore, C. L., *et al.*, *J. Med. Chem.* 43:3434-3442 (2000); Findeis, M. A., *Biochimica et Biophysica Acta* 1502:76-84 (2000); Kuner, P., Bohrmann, *et al.*, *J. Biol. Chem.* 275:1673-1678 (2000)). It is therefore of interest to develop ligands that specifically bind fibrillar A β aggregates. Since extracellular SPs are accessible targets, these new ligands could be used as *in vivo* diagnostic tools and as probes to visualize the progressive deposition of A β in studies of AD amyloidogenesis in living patients.

[0009] To this end, several interesting approaches for developing fibrillar A β aggregate-specific ligands have been reported (Ashburn, T. T., *et al.*, *Chem. Biol.* 3:351-358 (1996); Han, G., *et al.*, *J. Am. Chem. Soc.* 118:4506-4507 (1996); Klunk, W. E., *et al.*, *Biol. Psychiatry* 35:627 (1994); Klunk, W. E., *et al.*, *Neurobiol. Aging* 16:541-548 (1995); Klunk, W. E., *et al.*, *Society for Neuroscience Abstract* 23:1638 (1997); Mathis, C. A., *et al.*, *Proc. XIIth Intl. Symp. Radiopharm. Chem., Uppsala, Sweden*:94-95 (1997); Lorenzo, A. and Yankner, B. A., *Proc. Natl. Acad. Sci. U.S.A.* 91:12243-12247 (1994); Zhen, W., *et al.*, *J. Med. Chem.* 42:2805-2815 (1999)). The most attractive approach is based on highly conjugated chrysamine-G (CG) and Congo red (CR), and the latter has been used for fluorescent staining of SPs and NFTs in postmortem AD brain sections (Ashburn, T. T., *et al.*, *Chem. Biol.* 3:351-358 (1996); Klunk, W. E., *et al.*, *J. Histochem. Cytochem.* 37:1273-1281 (1989)). The inhibition constants (K_i) for binding to fibrillar A β aggregates of CR, CG, and 3'-bromo- and 3'-iodo derivatives of CG are 2,800, 370, 300 and 250 nM, respectively (Mathis, C. A., *et al.*, *Proc. XIIth Intl. Symp. Radiopharm. Chem.*,

Uppsala, Sweden:94-95 (1997)). These compounds have been shown to bind selectively to A β (1-40) peptide aggregates *in vitro* as well as to fibrillar A β deposits in AD brain sections (Mathis, C. A., *et al.*, *Proc. XIIIth Intl. Symp. Radiopharm. Chem., Uppsala, Sweden:94-95 (1997)*).

[0010] There are several potential benefits of imaging A β aggregates in the brain. The imaging technique will improve diagnosis by identifying potential patients with excess A β plaques in the brain; therefore, they may be likely to develop Alzheimer's disease. It will also be useful to monitor the progression of the disease. When anti-plaque drug treatments become available, imaging A β plaques in the brain may provide an essential tool for monitoring treatment. Thus, a simple, noninvasive method for detecting and quantitating amyloid deposits in a patient has been eagerly sought. Presently, detection of amyloid deposits involves histological analysis of biopsy or autopsy materials. Both methods have drawbacks. For example, an autopsy can only be used for a postmortem diagnosis.

[0011] The direct imaging of amyloid deposits *in vivo* is difficult, as the deposits have many of the same physical properties (e.g., density and water content) as normal tissues. Attempts to image amyloid deposits using magnetic resonance imaging (MRI) and computer-assisted tomography (CAT) have been disappointing and have detected amyloid deposits only under certain favorable conditions. In addition, efforts to label amyloid deposits with antibodies, serum amyloid P protein, or other probe molecules have provided some selectivity on the periphery of tissues, but have provided for poor imaging of tissue interiors.

[0012] Potential ligands for detecting A β aggregates in the living brain must cross the intact blood-brain barrier. Thus brain uptake can be improved by using ligands with relatively smaller molecular size (compared to Congo Red) and increased lipophilicity. Highly conjugated thioflavins (S and T) are commonly used as dyes for staining the A β aggregates in the AD brain (Elhaddaoui, A., *et al.*, *Biospectroscopy 1:351-356 (1995)*).

[0013] A highly lipophilic tracer, [^{18}F]FDDNP, for binding both tangles (mainly composed of hyperphosphorylated tau protein) and plaques

(containing A β protein aggregates) has been reported. (Shoghi-Jadid K, *et al.*, *Am J Geriatr Psychiatry*. 2002;10:24-35). Using positron-emission tomography (PET), it was reported that this tracer specifically labeled deposits of plaques and tangles in nine AD patients and seven comparison subjects. (Nordberg A. *Lancet Neurol*. 2004;3:519-27). Using a novel pharmacokinetic analysis procedure called the relative residence time of the brain region of interest versus the pons, differences between AD patients and comparison subjects were demonstrated. The relative residence time was significantly higher in AD patients. This is further complicated by an intriguing finding that FDDNP competes with some NSAIDs for binding to A β fibrils in vitro and to A β plaques *ex vivo* (Agdeppa ED, *et al.* 2001; Agdeppa ED, *et al.*, *Neuroscience*. 2003;117:723-30).

[0014] Imaging β -amyloid in the brain of AD patients by using a benzothiazole aniline derivative, [^{11}C]6-OH-BTA-1 (also referred to as [^{11}C]PIB), was recently reported. (Mathis CA, *et al.*, *Curr Pharm Des*. 2004;10:1469-92; Mathis CA, *et al.*, *Arch. Neurol*. 2005, 62:196-200.). Contrary to that observed for [^{18}F]FDDNP, [^{11}C]6-OH-BTA-1 binds specifically to fibrillar A β in vivo. Patients with diagnosed mild AD showed marked retention of [^{11}C]6-OH-BTA-1 in the cortex, known to contain large amounts of amyloid deposits in AD. In the AD patient group, [^{11}C]6-OH-BTA-1 retention was increased most prominently in the frontal cortex. Large increases also were observed in parietal, temporal, and occipital cortices and in the striatum. [^{11}C]6-OH-BTA-1 retention was equivalent in AD patients and comparison subjects in areas known to be relatively unaffected by amyloid deposition (such as subcortical white matter, pons, and cerebellum). Recently, another ^{11}C labeled A β plaque-targeting probe, a stilbene derivative- [^{11}C]SB-13, has been studied. In vitro binding using the [^3H]SB-13 suggests that the compound showed excellent binding affinity and binding can be clearly measured in the cortical gray matter, but not in the white matter of AD cases. (Kung M-P, *et al.*, *Brain Res*. 2004;1025:89-105. There was a very low specific binding in cortical tissue homogenates of control brains. The K_d values of [^3H]SB-13 in AD cortical homogenates were 2.4 $\pm$ 0.2 nM. High

binding capacity and comparable values were observed (14-45 pmol/mg protein) (*Id.*). As expected, in AD patients [¹¹C]SB-13 displayed a high accumulation in the frontal cortex (presumably an area containing a high density of A β plaques) in mild to moderate AD patients, but not in age-matched control subjects. (Verhoeff NP, *et al.*, *Am J Geriatr Psychiatry*. 2004;12:584-95).

- [0015] It would be useful to have a noninvasive technique for imaging and quantitating amyloid deposits in a patient. In addition, it would be useful to have compounds that inhibit the aggregation of amyloid proteins to form amyloid deposits and a method for determining a compound's ability to inhibit amyloid protein aggregation.

SUMMARY OF THE INVENTION

- [0016] The present invention provides novel compounds of Formulae I, Ia, II and III.
- [0017] The present invention also provides diagnostic compositions comprising a radiolabeled compound of Formulae I, Ia, II and III a pharmaceutically acceptable carrier or diluent.
- [0018] The invention further provides a method of imaging amyloid deposits, the method comprising introducing into a patient a detectable quantity of a labeled compound of Formulae I, Ia, II and III or a pharmaceutically acceptable salt, ester, amide or prodrug thereof.
- [0019] The present invention also provides a method for inhibiting the aggregation of amyloid proteins, the method comprising administering to a mammal an amyloid inhibiting amount of a compound Formulae I, Ia, II and III or a pharmaceutically acceptable salt, ester, amide, or prodrug thereof.
- [0020] A further aspect of this invention is directed to methods and intermediates useful for synthesizing the amyloid inhibiting and imaging compounds of Formulae I, Ia, II and III described herein.


DESCRIPTION OF THE FIGURES

- [0021] Fig. 1 depicts a film resulting from the imaging of a compound of the present invention.
- [0022] Fig. 2 depicts brain and bone uptake of styrylpyridine **2** in comparison to a stilbene analog.
- [0023] Fig. 3 depicts film autoradiography comparing styrylpyridine **2** in comparison to a stilbene analog.
- [0024] Fig. 4 depicts a saturation curve of styrylpyridine **2** in AD brain homogenates.
- [0025] Fig. 5 depicts several compounds of the present invention and their respective binding data.
- [0026] Fig. 6 depicts *in vitro* autoradiography of macroarray brain sections.
- [0027] Fig. 7 depicts the *in vitro* stability of F-18 labeled tracers toward pooled human liver microsomal fractions. Tracers in PBS without microsomal fractions served as the control. Values (% of unchanged parent compound) were average of duplicates.
- [0028] Fig. 8 depicts specific binding of [^{18}F]**2** to pooled AD and control brain tissue homogenates. Gray and white matters were dissected from the cortical regions. High specific binding was detected mainly in gray matter. The values presented are the mean $\pm$ SEM of six measurements. Relatively low binding was observed in white matter homogenates. In contrast, homogenates of control brain, either grey or white matters, showed significantly lower specific binding of [^{18}F]**2**.
- [0029] Fig. 9 depicts: (top) HPLC profile of compound [^{18}F]**2**; (bottom) UV trace of non-radioactive reference compound **2**, (350 nm). HPLC condition: Agilent 1100 series; Phenomenex Gemini C-18 column 5 μ 250 x 4.6 mm, CH₃CN/Ammonium formate buffer (1 mM) 8/2 v/v, 1 mL/min. Rt. 6.34 min (radioactive), 6.05 min (UV). Retention time gap was due to the detector configuration.

DETAILED DESCRIPTION OF THE INVENTION

[0030]


A compound of Formula I,




or a pharmaceutically acceptable salt thereof; wherein, n is an integer from one to six; at least one, no more than three, of A_1 , A_2 , A_3 , A_4 and A_5 is N, the others are $-CH$ or $-CR^2$ as permitted; R^1 is selected from the group consisting of:

- a. $-(CH_2)_pNR^aR^b$, wherein R^a and R^b are independently hydrogen, C_{1-4} alkyl, hydroxy(C_{1-4})alkyl or halo(C_{1-4})alkyl, and p is an integer from 0 to 5;
- b. hydroxy,
- c. C_{1-4} alkoxy,
- d. hydroxy(C_{1-4})alkyl,
- e. halogen,
- f. cyano,
- g. hydrogen,
- h. nitro,
- i. (C_1-C_4) alkyl,
- j. halo(C_1-C_4)alkyl,
- k. formyl,
- l. $-NHCO(C_{1-4}$ alkyl), and
- m. $-OCO(C_{1-4}$ alkyl);


R^2 is selected from the group consisting of:




wherein q is an integer from 1 to 10; Z is selected from the group consisting of halogen, halogen substituted benzoyloxy, halogen substituted benzyloxy, halogen substituted phenyl(C₁₋₄)alkyl, halogen substituted aryloxy, and a halogen substituted C₆₋₁₀ aryl, or Z can also be hydroxy; and R³⁰, R³¹, R³² and R³³ are in each instance independently selected from the group consisting of hydrogen, hydroxy, C₁₋₄ alkoxy, C₁₋₄ alkyl, and hydroxy(C₁₋₄)alkyl; or Z is hydroxy;



wherein q is an integer from 1 to 10, R^x and R^y are hydrogen, hydroxy or C₁₋₄ alkyl; t is 0, 1, 2 or 3; and Z, R³⁰, R³¹, R³² and R³³ are as described above;



wherein Y is selected from the group consisting of halogen, halogen substituted benzoyloxy, halogen substituted phenyl(C₁₋₄)alkyl, halogen substituted aryloxy, and halogen substituted C₆₋₁₀ aryl; U is selected from the group consisting of hydrogen, hydroxy, halogen, halogen substituted benzoyloxy, halogen substituted phenyl(C₁₋₄)alkyl, halogen substituted aryloxy, and halogen substituted C₆₋₁₀ aryl; and R³⁴, R³⁵, R³⁶, R³⁷, R³⁸, R³⁹ and R⁴⁰ are in each instance independently selected from the group consisting of hydrogen, halogen, hydroxy, C₁₋₄ alkoxy, C₁₋₄ alkyl, and hydroxy(C₁₋₄)alkyl;



wherein R^x and R^y are hydrogen, hydroxy or C_{1-4} alkyl; t is 0, 1, 2 or 3; and Y , U , R^{34} , R^{35} , R^{36} , R^{37} , R^{38} , R^{39} and R^{40} are as described above;

iii. $NR'R''$, wherein at least one of R' and R'' is $(CH_2)_dX$, where X is halogen, preferably F or ^{18}F , and d is an integer from 1 to 4; the other of R' and R'' is selected from the group consisting of hydrogen, C_{1-4} alkyl, halo(C_{1-4})alkyl, and hydroxy(C_{1-4})alkyl;

iv. $NR'R''-(C_{1-4})$ alkyl, wherein at least one of R' and R'' is $(CH_2)_dX$, where X is halogen, preferably F or ^{18}F , and d is an integer from 1 to 4; the other of R' and R'' is selected from the group consisting of hydrogen, C_{1-4} alkyl, halo(C_{1-4})alkyl, and hydroxy(C_{1-4})alkyl;

v. halo(C_{1-4})alkyl; and

vi. an ether ($R-O-R$) having the following structure:
[halo(C_{1-4})alkyl-O-(C_{1-4})alkyl]-; and

R^7 and R^8 are in each instance independently selected from the group consisting of hydrogen, hydroxy, amino, methylamino, dimethylamino, C_{1-4} alkoxy, C_{1-4} alkyl, and hydroxy(C_{1-4})alkyl.

[0031] Preferred compounds include those where the halogen, in one or more occurrence on the structure, is a radiolabeled halogen. Also preferred are compounds wherein the halogen is selected from the group consisting of I , ^{123}I , ^{125}I , ^{131}I , Br , ^{76}Br , ^{77}Br , F or ^{18}F . Especially preferred compounds are those that contain ^{18}F . Compounds containing ^{123}I are also especially preferred.

[0032] Useful values of R^1 are listed above. Preferred values are hydroxy or $NR^aR^b(CH_2)_p-$, wherein p is an integer from 0 to 5, and R^a and R^b are independently hydrogen, C_{1-4} alkyl or $(CH_2)_dX$, where X is halogen, and d is an integer from 1 to 4. Useful values of p include integers from 0 to 5. Preferably, p is 0, 1 or 2. Most preferably, p is 0 such that R^1 represents

NR^aR^b . In preferred embodiments, R^1 is either in the meta or para position relative to the respective bridge. A preferred value of R^1 is NR^aR^b , wherein R^a and R^b are independently hydrogen or C_{1-4} alkyl. In this embodiment, it is preferable that the C_{1-4} alkyl is methyl. Preferably one of R^a and R^b is hydrogen, the other is C_{1-4} alkyl, such as methyl. Most preferably, both R^a and R^b are methyl. Another preferred value of R^1 is hydroxy. Also preferred are any prodrug groups that after administration yield a preferred value of R^1 . Such prodrug groups are well-known in the art.

[0033] Useful values of n include integers from 1 to 6. Preferably, the value of n is from 1 to 4. Most preferably, the value of n is from 1 to 3. It is especially preferred that n is one.


[0034] Useful values of R^7 and R^8 are in each instance independently selected from the group consisting of hydrogen, hydroxy, amino, methylamino, dimethylamino, C_{1-4} alkoxy, C_{1-4} alkyl, and hydroxy(C_{1-4})alkyl. The value of n determines the number of R^7 and R^8 group(s) present in the compound. If present more than once in a particular compound, in each instance of R^7 and R^8 the value can be different from any other value of R^7 and R^8 . In preferred embodiments, R^7 and R^8 are each hydrogen in every instance.

[0035] Useful values of R^2 include substructures i, i', ii, ii', iii, iv, v, and vi, as depicted above. In preferred embodiments of Formula I, R^2 is either in the meta or para position relative to the respective bridge. Preferably, R^2 is substructure i or ii. Also preferred are substructures i' and ii'. In these embodiments, useful values of q include integers from one to ten. Preferably, in a compound where R^2 is i or i', q is an integer from 1 to 5. Most preferably, q is 1 to 4, especially 3 or 4. In substructure i or i', useful values of R^{30} , R^{31} , R^{32} and R^{33} independently include hydrogen, hydroxy, C_{1-4} alkoxy, C_{1-4} alkyl, and hydroxy(C_{1-4})alkyl. Preferred compounds include those where one or more of R^{30} , R^{31} , R^{32} and R^{33} are hydrogen. More preferred compounds include those where each of R^{30} , R^{31} , R^{32} and R^{33} is hydrogen.



[0036] In substructure ii or ii', useful values of Y, U and R^{34} , R^{35} , R^{36} , R^{37} , R^{38} , R^{39} and R^{40} are described above. Preferred compounds include those where U is hydroxy.

[0037] Useful compounds include those compounds where at least one, no more than three, of A_1 , A_2 , A_3 , A_4 and A_5 is N, and the others are -CH or -CR² as permitted. It is more preferred that N be in position A_4 .


[0038] Preferred compounds of Formula I include those compounds wherein A_4 is N, having the following formula:




wherein R^a and R^b are independently selected from hydrogen or C₁-C₄ alkyl, Z is as described above and q is an integer from 1 to 5. Examples of preferred compounds include:




wherein q is an integer from 1 to 4; such as,




[0039] Other preferred compounds of Formula I, when R^2 is ii, include:




and




[0040] In another aspect, the present invention is directed to compounds of Formula I, having the following structure:




or a pharmaceutically acceptable salt thereof, wherein: at least one, no more than three, of A₁, A₂, A₃, A₄ and A₅ is N, the others are -CH, or -CR² as permitted; n is an integer from 1 to 6; R¹ includes all useful values described above, preferably hydroxy or NR^aR^b(CH₂)_p-, wherein p is an integer from 0 to 5, and R^a and R^b are independently hydrogen, C₁₋₄ alkyl or (CH₂)_dX, where X is halogen, and d is an integer from 1 to 6; R² is selected from the group consisting of:




wherein q is an integer from 2 to 10; Z is -Ch;




wherein q is an integer from 1 to 10, R^x and R^y are hydrogen, hydroxy or C₁₋₄ alkyl; t is 0, 1, 2 or 3; and Z, R³⁰, R³¹, R³² and R³³ are as described; and Z is -Ch;



wherein Z is -Ch, R³⁰, R³¹, R³² and R³³ are as described above, and




wherein Y is -Ch; U is selected from the group consisting of hydrogen, hydroxy, halogen, halogen substituted benzoyloxy, halogen substituted phenyl(C₁₋₄)alkyl, halogen substituted aryloxy, and halogen substituted C₆₋₁₀ aryl; and R³⁴, R³⁵, R³⁶, R³⁷, R³⁸, R³⁹ and R⁴⁰ are in each instance independently selected from the group consisting of hydrogen, halogen, hydroxy, C₁₋₄ alkoxy, C₁₋₄ alkyl, and hydroxy(C₁₋₄)alkyl;




wherein, R^x and R^y are hydrogen, hydroxy or C₁₋₄ alkyl; t is 0, 1, 2 or 3; and Y, U, R³⁴, R³⁵, R³⁶, R³⁷, R³⁸, R³⁹ and R⁴⁰ are as described above;

- iv. -(CH₂)_w-O-Ch, wherein w is an integer from 1 to 10;
- v. -Ch; and
- vi. -(CH₂)_w-Ch, wherein w is an integer from 1 to 10;

wherein, the moiety "-Ch" is a chelating ligand capable of complexing with a metal to form a metal chelate. Many ligands are known in the art and are suitable for use as a labeling moiety for the compounds of the present invention. Those of skill in the art will understand that such ligands provide a way to label compounds and the invention is not limited to particular ligands, many of which are interchangeable. Preferably, this ligand is a tri- or tetradentate ligand, such as N₃, N₂S, NS₂, N₄ and those of the N₂S₂ type, represented by, but not limited to, the following structure:



[0041] wherein R^P is hydrogen or a sulfhydryl protecting group, and R⁹, R¹⁰, R¹¹, R¹², R¹³, R¹⁴, R¹⁵, R¹⁶, R⁴³ and R⁴⁴ are in each instance independently selected from the group consisting of hydrogen, hydroxy, amino, methylamino, dimethylamino, C₁₋₄ alkoxy, C₁₋₄ alkyl, and hydroxy(C₁₋₄)alkyl. When complexed with a metal such as 99m-Tc, -Ch has the following structure:



[0042] Additionally, a rhenium radioisotope can be complexed with the tetradentate ligand, rather than technetium. When the chelating moiety is not complexed with a metal, R^P are both hydrogen, or can be any of the variety of protecting groups available for sulfur, including methoxymethyl, methoxyethoxymethyl, *p*-methoxybenzyl or benzyl. Sulfur protecting groups are described in detail in Greene, T.W. and Wuts, P.G.M., *Protective Groups in Organic Synthesis*, 2nd Edition, John Wiley and Sons, Inc., New York (1991). Protecting group R^P can be removed by appropriate methods well known in the art of organic synthesis, such as trifluoroacetic acid, mercuric chloride or sodium in liquid ammonia. In the case of Lewis acid labile groups, including acetamidomethyl and benzamidomethyl, R^P can be left intact. Labeling of the ligand with technetium in this case will cleave the protecting group, rendering the protected diaminedithiol equivalent to the unprotected form. Further, several ligands of the general N₂S₂ type are known, and can be used interchangeably without changing the scope of the invention; and

R^7 and R^8 are in each instance independently selected from the group consisting of hydrogen, hydroxy, amino, methylamino, dimethylamino, C_{1-4} alkoxy, C_{1-4} alkyl, and hydroxy(C_{1-4})alkyl.

[0043] Preferred values of R^1 are hydroxy or $NR^aR^b(CH_2)_p$, wherein p is an integer from 0 to 5, and R^a and R^b are independently hydrogen, C_{1-4} alkyl or $(CH_2)_dX$, where X is halogen, and d is an integer from 1 to 4. Useful values of p include integers from 0 to 5. Preferably, p is 0, 1 or 2. Most preferably, p is 0 such that R^1 represents NR^aR^b . In preferred embodiments, R^1 is either in the meta or para position relative to the respective bridge. A preferred value of R^1 is NR^aR^b , wherein R^a and R^b are independently hydrogen or C_{1-4} alkyl. In this embodiment, it is preferable that the C_{1-4} alkyl is methyl. Preferably one of R^a and R^b is hydrogen, the other is C_{1-4} alkyl, such as methyl or both R^a and R^b are methyl. Another preferred value of R^1 is hydroxy. Also preferred for R^1 are any groups that after administration into the body metabolize or degrade to the preferred values of R^1 listed above. Such groups are known in the art to constitute a prodrug and the groups capable of forming prodrugs are well-known to one of ordinary skill in the art.

[0044] Useful values of n include integers from 1 to 6. Preferably, the value of n is from 1 to 4. Most preferably, the value of n is from 1 to 3. It is especially preferred that n is one.

[0045] Useful values of R^7 and R^8 are in each instance independently selected from the group consisting of hydrogen, hydroxy, amino, methylamino, dimethylamino, C_{1-4} alkoxy, C_{1-4} alkyl, and hydroxy(C_{1-4})alkyl. The value of n determines the number of R^7 and R^8 group(s) present in the compound. If present more than once in a particular compound, in each instance of R^7 and R^8 the value can be different from any other value of R^7 and R^8 . In preferred embodiments, R^7 and R^8 are each hydrogen in every instance.


[0046] Useful values of R^2 include substructures i, i', ii, iii and iii' as depicted above. In preferred embodiments of Formula I, R^2 is either in the meta or para position relative to the respective bridge. Preferably, in a compound where R^2 is i or i', q is an integer from 2 to 5. Most preferably, q is 3 or 4. In substructure i or i', useful values of R^{30} , R^{31} , R^{32} and R^{33} independently

include hydrogen, hydroxy, C₁₋₄ alkoxy, C₁₋₄ alkyl, and hydroxy(C₁₋₄)alkyl. Preferred compounds include those where one or more of R³⁰, R³¹, R³² and R³³ are hydrogen. More preferred compounds include those where each of R³⁰, R³¹, R³² and R³³ is hydrogen.

[0047] In substructure iii or iii', useful values of U and R³⁴, R³⁵, R³⁶, R³⁷, R³⁸, R³⁹ and R⁴⁰ are described above. Preferred compounds include those where U is hydroxy.

[0048] Useful compounds include those compounds where one, no more than three, of A₁, A₂, A₃, A₄ and A₅ is N, and the others are -CH or -CR² as permitted. It is preferred that if only one of A₁, A₂, A₃, A₄ and A₅ is N, that it is A₄.

[0049] In another aspect, the present invention is directed to a compound of Formula II, having the following structure:



or a pharmaceutically acceptable salt thereof, wherein: at least one, no more than three, of A₁, A₂, A₃, A₄ and A₅ is N, the others are -CH, -CR³ or -CR⁴ as permitted; R⁵ and R^{5'} are independently hydrogen or C₁₋₄ alkyl; R¹ and R², in each instance, is independently selected from the group consisting of hydrogen, halogen, C₁₋₄ alkyl, cyano, carboxy(C₁₋₅)alkyl, trifluoromethyl, nitro, halo(C₁₋₄)alkyl, formyl and NR⁶R⁷(CH₂)_p-, wherein p is an integer from 0 to 5, and R⁶ and R⁷ are independently hydrogen, C₁₋₄ alkyl or (CH₂)_dX, where X is halogen, and d is an integer from 1 to 4; in addition to the values listed above for R¹ and R², R¹ and/or R² can independently also be hydroxy; R³ is selected from the group consisting of hydrogen, halogen, C₁₋₄ alkyl, cyano, carboxy(C₁₋₅)alkyl, trifluoromethyl, nitro, halo(C₁₋₄)alkyl, formyl, NR⁶R⁷(CH₂)_p-, wherein p is an integer from 0 to 5, and R⁶ and R⁷ are independently hydrogen, C₁₋₄ alkyl or (CH₂)_dX, where X is halogen, and d is

an integer from 1 to 4, ^{18}F fluoromethyl, ^{18}F fluoroethyl, ^{18}F fluoropropyl and $\text{Sn}(\text{alkyl})_3$;

R^4 is selected from the group consisting of:

- a. C_{1-4} alkylthio,
- b. C_{1-4} alkylsulfonyl,
- c. hydroxy,
- d. C_{1-4} alkoxy,
- e. $\text{NR}^6\text{R}^7(\text{CH}_2)_p$, wherein p is an integer from 0 to 5, and R^6 and R^7 are independently hydrogen, C_{1-4} alkyl or $(\text{CH}_2)_d\text{X}$, where X is halogen, and d is an integer from 1 to 4,
- f. phenyl(C_{1-4})alkyl,
- g. C_{6-10} aryl,
- h. heteroaryl,
- i. heterocycle,
- j. heterocycle(C_{1-4})alkyl, and
- k. C_{3-6} cycloalkyl,

wherein said phenyl(C_{1-4})alkyl, C_{6-10} aryl, heteroaryl, heterocycle, heterocycle(C_{1-4})alkyl or C_{3-6} cycloalkyl is substituted with one of the following: C_{1-4} alkylthio, C_{1-4} alkyl sulfonyl, methoxy, hydroxy, dimethylamino or methylamino; and, X' is hydrogen, ^{18}F fluoromethyl, ^{18}F fluoroethyl, ^{18}F fluoropropyl, ^{125}I , ^{123}I , ^{131}I , ^{18}F , ^{76}Br , ^{77}Br or $\text{Sn}(\text{alkyl})_3$; provided that one of R^1 , R^2 , R^3 or R^4 is $\text{NR}^6\text{R}^7(\text{CH}_2)_p$. In another embodiment, it is also provided that one of R^1 , R^2 or R^4 is hydroxy.

[0050] Useful values of R^5 and $\text{R}^{5'}$ include all the values listed above. Preferably, R^5 and $\text{R}^{5'}$ are independently hydrogen or a C_{1-4} alkyl such as methyl. Also preferred, R^1 and R^2 are independently hydroxy, monomethylamine or dimethylamine.

[0051] Useful values of R^3 include all those values listed above. More preferably, R^3 is hydrogen, ^{18}F fluoromethyl, ^{18}F fluoroethyl, ^{18}F fluoropropyl, ^{125}I , ^{123}I , ^{131}I or ^{18}F .





[0052] Useful values of R^1 and R^2 include all the values listed above. Preferably, R^1 and R^2 are independently hydrogen or a C_{1-4} alkyl such as methyl.

[0053] Useful values of R^4 include all those values listed above. Preferably, R^4 is methylthio, methylsulfonyl, hydroxy, methoxy or $NR^6R^7(CH_2)_p$.

[0054] Useful values of X' include all those listed above. Preferred values include hydrogen, ^{18}F fluoromethyl, ^{18}F fluoroethyl or ^{18}F fluoropropyl, ^{125}I , ^{123}I , ^{131}I and ^{18}F .


[0055] In all compounds of the present invention wherein only one of A_1 , A_2 , A_3 , A_4 and A_5 is N, it is more preferred that A_4 is N.

[0056] Representative compounds of the present invention include:




wherein -Ch is an N_2S_2 type chelating moiety, X, q, R^a , R^b , R^7 , R^8 , R^{30} , R^{31} , R^{32} and R^{33} are as described above. Most preferably, R^a and R^b are both methyl.

[0057] In another embodiment are compounds of Formula I'a having the following general structure:




wherein at least one, no more than three, of A_1 , A_2 , A_3 , A_4 and A_5 is N, the others are -CH; q is an integer from 1 to 10; R' and R'' are each independently hydrogen or C_{1-4} alkyl and X is a radiolabeled halogen or -Ch moiety. An example of these compounds include compounds of Formula Ia having the following structure:




wherein at least one, no more than three, of A_1 , A_2 , A_3 , A_4 and A_5 is N, the others are -CH; q is an integer from 1 to 10; and X is a radiolabeled halogen or -Ch moiety. Preferably, a mono or di C_{1-4} alkylamino, more preferably monomethylamino or dimethylamino and PEG substituents are in the para position relative to the ethylene bridge. Also, it is preferred that A_4 is N, and A_1 , A_2 , A_3 and A_5 are each -CH. Preferred values of q are integers from 2 to 5; and especially preferred values are 3 and 4. Preferred values of X include ^{123}I and ^{18}F . The most preferred value of X is ^{18}F .

[0058] In another embodiment of the present invention are compounds of Formula III having the following structure:




or a pharmaceutically acceptable salt thereof; wherein, n is an integer from one to six; at least one, no more than three, of A_1 , A_2 , A_3 , A_4 and A_5 is N, the others are -CH, $-\text{CR}^2$ or $-\text{CR}^3$ as permitted; R^1 includes all the useful values listed above under Formula I, preferably hydroxy or $\text{NR}^a\text{R}^b(\text{CH}_2)_p-$, wherein p is an integer from 0 to 5, and R^a and R^b are independently hydrogen, C_{1-4} alkyl


or $(\text{CH}_2)_d\text{X}$, where X is halogen, and d is an integer from 1 to 4, R^3 is selected from the group of ^{125}I , ^{123}I , ^{131}I , ^{18}F , $^{18}\text{F}(\text{C}_1\text{-C}_4)$ alkyl, ^{76}Br , ^{77}Br or $\text{Sn}(\text{alkyl})_3$; R^2 is selected from the group consisting of:




wherein q is an integer from 1 to 10; Z is selected from the group consisting of hydrogen, hydroxy, halogen, C_{1-4} alkoxy, C_{1-4} alkyl, and hydroxy(C_{1-4})alkyl; and R^{30} , R^{31} , R^{32} and R^{33} are in each instance independently selected from the group consisting of hydrogen, hydroxy, C_{1-4} alkoxy, C_{1-4} alkyl, and hydroxy(C_{1-4})alkyl;



Wherein q is an integer from 1 to 10, R^x and R^y are hydrogen, hydroxy or C_{1-4} alkyl; t is 0, 1, 2 or 3; and Z, R^{30} , R^{31} , R^{32} and R^{33} are as described;



wherein Y and U are independently selected from the group consisting of hydrogen, hydroxyl, halogen, C_{1-4} alkoxy, C_{1-4} alkyl, and hydroxy(C_{1-4})alkyl; U is selected from the group consisting of hydrogen, hydroxy, halogen, halogen substituted benzoyloxy, halogen substituted phenyl(C_{1-4})alkyl, halogen substituted aryloxy, and halogen substituted C_{6-10} aryl; and R^{34} , R^{35} , R^{36} , R^{37} , R^{38} , R^{39} and R^{40} are in each instance independently selected from the group consisting of hydrogen, halogen, hydroxy, C_{1-4} alkoxy, C_{1-4} alkyl, and hydroxy(C_{1-4})alkyl;



wherein R^x and R^y are hydrogen, hydroxy or C_{1-4} alkyl; t is 0, 1, 2 or 3; and Y , U , R^{34} , R^{35} , R^{36} , R^{37} , R^{38} , R^{39} and R^{40} are as described above;

iii. $NR'R''$, wherein at least one of R' and R'' is $(CH_2)_dX$, where X is halogen, preferably F or ^{18}F , and d is an integer from 1 to 4; the other of R' and R'' is selected from the group consisting of hydrogen, C_{1-4} alkyl, halo(C_{1-4})alkyl, and hydroxy(C_{1-4})alkyl;

iv. $NR'R''-(C_{1-4})$ alkyl, wherein at least one of R' and R'' is $(CH_2)_dX$, where X is halogen, preferably F or ^{18}F , and d is an integer from 1 to 4; the other of R' and R'' is selected from the group consisting of hydrogen, C_{1-4} alkyl, halo(C_{1-4})alkyl, and hydroxy(C_{1-4})alkyl;

v. halo(C_{1-4})alkyl; and

vi. an ether ($R-O-R$) having the following structure:

[halo(C_{1-4})alkyl-O-(C_{1-4})alkyl]-; and

[0059] R^7 and R^8 are in each instance independently selected from the group consisting of hydrogen, hydroxy, amino, methylamino, dimethylamino, C_{1-4} alkoxy, C_{1-4} alkyl, and hydroxy(C_{1-4})alkyl.

[0060] Preferred compounds include those where the halogen, in one or more occurrence on the structure, is a radiolabeled halogen. Also preferred are compounds wherein the halogen is selected from the group consisting of I , ^{123}I , ^{125}I , ^{131}I , Br , ^{76}Br , ^{77}Br , F or ^{18}F . Especially preferred compounds are those that contain ^{18}F . Compounds containing ^{123}I are also especially preferred.

[0061] Useful values of R^1 are listed above. Preferred values are hydroxy or $NR^aR^b(CH_2)_p-$, wherein p is an integer from 0 to 5, and R^a and R^b are independently hydrogen, C_{1-4} alkyl or $(CH_2)_dX$, where X is halogen, and d is an integer from 1 to 4. Useful values of p include integers from 0 to 5. Preferably, p is 0, 1 or 2. Most preferably, p is 0 such that R^1 represents

NR^aR^b . In preferred embodiments, R^1 is either in the meta or para position relative to the respective bridge. A preferred value of R^1 is NR^aR^b , wherein R^a and R^b are independently hydrogen or C_{1-4} alkyl. In this embodiment, it is preferable that the C_{1-4} alkyl is methyl. Preferably one of R^a and R^b is hydrogen, the other is C_{1-4} alkyl, such as methyl. Most preferably, both R^a and R^b are methyl. Another preferred value of R^1 is hydroxy. Also preferred for R^1 are any groups that after administration into the body metabolize or degrade to the preferred values of R^1 listed above. Such groups are known in the art to constitute a prodrug and the groups capable of forming prodrugs are well-known to one of ordinary skill in the art.

[0062] Useful values of n include integers from 1 to 6. Preferably, the value of n is from 1 to 4. Most preferably, the value of n is from 1 to 3. It is especially preferred that n is one.

[0063] Useful values of R^7 and R^8 are in each instance independently selected from the group consisting of hydrogen, hydroxy, amino, methylamino, dimethylamino, C_{1-4} alkoxy, C_{1-4} alkyl, and hydroxy(C_{1-4})alkyl. The value of n determines the number of R^7 and R^8 group(s) present in the compound. If present more than once in a particular compound, in each instance of R^7 and R^8 the value can be different from any other value of R^7 and R^8 . In preferred embodiments, R^7 and R^8 are each hydrogen in every instance.


[0064] Useful values of R^2 include substructures i, i', ii, ii', iii, iv, v, and vi, as depicted above. In preferred embodiments of Formula I, R^2 is either in the meta or para position relative to the respective bridge. Preferably, R^2 is substructure i or ii. Also preferred are substructures i' and ii'. In these embodiments, useful values of q include integers from one to ten. Preferably, in a compound where R^2 is i or i', q is an integer from 1 to 5. Most preferably, q is 1 to 4, especially 3 or 4. In substructure i or i', useful values of R^{30} , R^{31} , R^{32} and R^{33} independently include hydrogen, hydroxy, C_{1-4} alkoxy, C_{1-4} alkyl, and hydroxy(C_{1-4})alkyl. Preferred compounds include those where one or more of R^{30} , R^{31} , R^{32} and R^{33} are hydrogen. More preferred compounds include those where each of R^{30} , R^{31} , R^{32} and R^{33} is hydrogen.

[0065] In substructure ii or ii', useful values of Y, U and R^{34} , R^{35} , R^{36} , R^{37} , R^{38} , R^{39} and R^{40} are described above. Preferred compounds include those where U is hydroxy.

[0066] Useful compounds include those compounds where at least one, no more than three, of A_1 , A_2 , A_3 , A_4 and A_5 is N, and the others are -CH or -CR² as permitted. It is more preferred that N be in position A_4 .




Particularly useful compounds of Formula III include those compounds where A_4 is N, and the others are -CH, -CR² or -CR³ as permitted.

[0067] Especially preferred compounds of Formula III wherein A_4 is N, include the following:




wherein R^a and R^b are independently selected from hydrogen or C₁-C₄ alkyl and q is an integer from 1 to 4 and R^3 is preferably ^{123}I or ^{18}F ;

[0068] Examples of most preferred compounds of Formula III include the following:




and



16b


[0069]

Other preferred compounds of Formula III, when R^2 is ii, include:



17a

and




18a

wherein Y is selected from the group consisting of hydrogen and F.

[0070]


Compounds of Formula III when R^2 is i, or i' when t is 0, include hydroxy ethers such as:



wherein R^1 and R^3 are as described above under Formula III.

[0071]

In all embodiments of Formulae I and III containing $-(CR^xR^y)_t$ where t is other than zero, the compounds have the following general structure wherein there is at least one carbon-carbon bond between a substituent and the nitrogen-containing ring:



[0072]

The compounds of the present invention can also contain a radioactive isotope of carbon as the radiolabel. This refers to a compound that comprises one or more radioactive carbon atoms, preferably ^{11}C , with a specific activity above that of the background level for that atom. It is well known, in this respect, that naturally occurring elements are present in the form of varying

isotopes, some of which are radioactive isotopes. The radioactivity of the naturally occurring elements is a result of the natural distribution or abundance of these isotopes, and is commonly referred to as a background level. The carbon labeled compounds of the present invention have a specific activity that is higher than the natural abundance, and therefore above the background level. The composition claimed herein comprising a carbon-labeled compound(s) of the present invention will have an amount of the compound such that the composition can be used for tracing, imaging, radiotherapy, and the like.

[0073] In certain embodiments of the compounds disclosed herein, a halogen, preferably ^{18}F , or a chelating agent is linked to the styrylpyridine backbone through a PEG chain, having a variable number of ethoxy groups. The fluorinated styrylpyridine, 2, displayed high binding affinity ($K_i = 2.5 \pm 0.4$ nM). The dimethylamino analog showed the greatest affinity. This is in contrast to stilbene analogs, which tend to possess higher affinity when monomethylamino substituted. As shown in Schemes 1-3 herein, radiolabeling was successfully performed giving the target compounds. The synthesis of compound 2 in Scheme 5 resulted in a preparation time of about 60 mins; radiochemical yield of ~35% (decay corrected); radiochemical purity of >98%; and specific activity of from about 1,000 to about 1,500 Ci/mmol. In vivo biodistribution of a ^{18}F pegylated styrylpyridine in normal mice exhibited excellent brain penetrations and rapid washouts after an iv injection. Autoradiography of postmortem AD brain sections of 2 confirmed the specific binding related to the presence of A β plaques.

[0074] Preferable values under the scope of C_{6-10} aryl include phenyl, naphthyl or tetrahydronaphthyl. Preferable values of under the scope of heteroaryl include thienyl, furyl, pyranyl, pyrrolyl, pyridinyl, indolyl, and imidazolyl. Preferable values under the scope of heterocycle include piperidinyl, pyrrolidinyl, and morpholinyl. A preferred embodiment of a C_{6-10} aryl, heteroaryl, heterocycle, heterocycle(C_{1-4})alkyl or C_{3-6} cycloalkyl, contains a ring substituted with one of the following: C_{1-4} alkylthio, C_{1-4} alkyl sulfonyl, methoxy, hydroxy, dimethylamino or methylamino.

[0075] The compounds of Formulae I, Ia, II, and III may also be solvated, especially hydrated. Hydration may occur during manufacturing of the compounds or compositions comprising the compounds, or the hydration may occur over time due to the hygroscopic nature of the compounds. In addition, the compounds of the present invention can exist in unsolvated as well as solvated forms with pharmaceutically acceptable solvents such as water, ethanol, and the like. In general, the solvated forms are considered equivalent to the unsolvated forms for the purposes of the present invention.

[0076] When any variable occurs more than one time in any constituent or in Formula I, Ia, II or III its definition on each occurrence is independent of its definition at every other occurrence. Also combinations of substituents and/or variables are permissible only if such combinations result in stable compounds.

[0077] The term "alkyl" as employed herein by itself or as part of another group refers to both straight and branched chain radicals of up to 8 carbons, preferably 6 carbons, more preferably 4 carbons, such as methyl, ethyl, propyl, isopropyl, butyl, *t*-butyl, and isobutyl.

[0078] The term "alkoxy" is used herein to mean a straight or branched chain alkyl radical, as defined above, unless the chain length is limited thereto, bonded to an oxygen atom, including, but not limited to, methoxy, ethoxy, *n*-propoxy, isopropoxy, and the like. Preferably the alkoxy chain is 1 to 6 carbon atoms in length, more preferably 1-4 carbon atoms in length.

[0079] The term "monoalkylamine" as employed herein by itself or as part of another group refers to an amino group which is substituted with one alkyl group as defined above.

[0080] The term "dialkylamine" as employed herein by itself or as part of another group refers to an amino group which is substituted with two alkyl groups as defined above.

[0081] The term "halo" or "halogen" employed herein by itself or as part of another group refers to chlorine, bromine, fluorine or iodine and their isotopes. The term "radiohalogen" refers specifically to radioactive halogen isotopes.

- [0082] The term "haloalkyl" as employed herein refers to any of the above alkyl groups substituted by one or more chlorine, bromine, fluorine or iodine with fluorine and chlorine being preferred, such as chloromethyl, iodomethyl, trifluoromethyl, 2,2,2-trifluoroethyl, and 2-chloroethyl.
- [0083] The term "alkylthio" as employed herein by itself or as part of another group refers to a thioether of the structure: R-S, wherein R is a C₁₋₄ alkyl as defined above.
- [0084] The term "alkylsulfonyl" as employed herein by itself or as part of another group refers to a sulfone of the structure: R-SO₂, wherein R is a C₁₋₄ alkyl as defined above.
- [0085] The term "aryl" as employed herein by itself or as part of another group refers to monocyclic or bicyclic aromatic groups containing from 6 to 12 carbons in the ring portion, preferably 6-10 carbons in the ring portion, such as phenyl, naphthyl or tetrahydronaphthyl.
- [0086] The term "heterocycle" or "heterocyclic ring", as used herein except where noted, represents a stable 5- to 7- membered mono-heterocyclic ring system which may be saturated or unsaturated, and which consists of carbon atoms and from one to three heteroatoms selected from the group consisting of N, O, and S, and wherein the nitrogen and sulfur heteroatom may optionally be oxidized. Especially useful are rings contain one nitrogen combined with one oxygen or sulfur, or two nitrogen heteroatoms. Examples of such heterocyclic groups include piperidinyl, pyrrolyl, pyrrolidinyl, imidazolyl, imidaziny, imidazolidinyl, pyridyl, pyrazinyl, pyrimidinyl, oxazolyl, oxazolidinyl, isoxazolyl, isoxazolidinyl, thiazolyl, thiazolidinyl, isothiazolyl, homopiperidinyl, homopiperazinyl, pyridazinyl, pyrazolyl, and pyrazolidinyl, most preferably thiamorpholinyl, piperazinyl, and morpholinyl.
- [0087] The term "heteroatom" is used herein to mean an oxygen atom ("O"), a sulfur atom ("S") or a nitrogen atom ("N"). It will be recognized that when the heteroatom is nitrogen, it may form an NRR moiety, wherein the R groups independently from one another may be hydrogen or C₁₋₄ alkyl, C₂₋₄ aminoalkyl, C₁₋₄ halo alkyl, halo benzyl, or R¹ and R² are taken together to

form a 5- to 7-member heterocyclic ring optionally having O, S or NR^c in said ring, where R^c is hydrogen or C₁₋₄ alkyl.

[0088] The term "heteroaryl" as employed herein refers to groups having 5 to 14 ring atoms; 6, 10 or 14 Π electrons shared in a cyclic array; and containing carbon atoms and 1, 2, 3 or 4 oxygen, nitrogen or sulfur heteroatoms (where examples of heteroaryl groups are: thienyl, benzo[b]thienyl, naphtho[2,3-b]thienyl, thianthrenyl, furyl, pyranyl, isobenzofuranyl, benzoxazolyl, chromenyl, xanthenyl, phenoxathiinyl, 2*H*-pyrrolyl, pyrrolyl, imidazolyl, pyrazolyl, pyridyl, pyrazinyl, pyrimidinyl, pyridazinyl, indoliziny, isoindolyl, 3*H*-indolyl, indolyl, indazolyl, purinyl, 4*H*-quinoliziny, isoquinolyl, quinolyl, phthalazinyl, naphthyridinyl, quinazolinyl, cinnoliny, pteridinyl, 4*aH*-carbazolyl, carbazolyl, α , β , or γ -carbolinyl, phenanthridinyl, acridinyl, perimidinyl, phenanthrolinyl, phenazinyl, isothiazolyl, phenothiazinyl, isoxazolyl, furazanyl and phenoxazinyl groups).

[0089] The term "aralkyl" or "arylalkyl" as employed herein by itself or as part of another group refers to C₁₋₆ alkyl groups as discussed above having an aryl substituent, such as benzyl, phenylethyl or 2-naphthylmethyl.


[0090] Another aspect of this invention is related to methods of preparing compounds of Formulae I, Ia, II and III.

[0091] The synthesis of dimethylamino substituted styrylpyridine derivative 1 and its fluoropegylated compound 2 is shown in scheme 1. Compound 1 was obtained by a Wittig reaction between diethyl 4-(dimethylamino)benzylphosphonate and 6-chloronicotinaldehyde at the present of potassium *tert*-butoxide in DMF (yield 62 %). A direct alkylation of compound 1 with 2-(2-(2-fluoroethoxy)ethoxy)ethanol² using sodium hydride in THF obtained the fluoropegylated compound 2 (yield 33 %), which can be used as the cold standard for the radio-labeling. The preparation of monomethyl amino substituted derivative 6 was accomplished through a route showing in Scheme 2. A Wittig reaction between 4-nitro-benzylphosphonate and 6-chloronicotinaldehyde at the present of sodium methoxide in methanol under refluxing condition obtained compound 3 in a high yield (88 %). Compound 3 can be easily filtered out after the reaction and used directly for


next step; no further purification is required. The alkylation of **3** with 2-(2-(2-fluoroethoxy)ethoxy)ethanol using sodium hydride in THF obtained compound **4** (yield 30 %). The nitro group of compound **4** was reduced using stannous chloride in ethanol to obtain compound **5** (yield 58 %). Monomethylation of **5** was achieved using paraformaldehyde, sodium methoxide and sodium borohydride to obtain compound **6** in a relatively high yield (73 %).

[0092] To make desired F-18 labeled dimethylamino substituted styrylpyridine derivative [^{18}F]**2**, the tosylate **10** (Scheme 3) was used as the precursor. The preparation of **10** started with a microwave-assisted alkylation of **3** with triethylene glycol in DMF to obtain compound **7** (yield 77 %). The nitro group of **7** was then reduced to amine using stannous chloride to give compound **8** (yield 76 %) then followed by a dimethylation using paraformaldehyde, sodium cyanoborohydride in acetic acid to obtain compound **9** in a high yield (95 %). Mesylation of **9** was tried first, however, the mesylate of **9** was very unstable and decomposed during the preparation. Tosylation of **9** was accomplished successfully using tosylchloride in pyridine to give desired tosylate **10** (yield 41 %) as the precursor for making radio-labeled [^{18}F]**2**.


SCHEME 1




SCHEME 2



SCHEME 3




SCHEME 4




[0093] Schemes 5-7 depict a synthetic route for compounds of Formula III. Scheme 5 depicts a synthesis of several intermediates useful for preparing the compounds of the invention. Scheme 6 and 7 depict the synthesis of radiolabeled and non-radiolabeled compounds of the invention. In compounds I7-I10, "I" in the compound name means "intermediate."

SCHEME 5




Reagents and conditions: (a) NIS, CH₃CN, reflux, 1 h; (b) F(CH₂CH₂O)₃H, Ph₃P, DIAD, THF, -5 °C to rt, 2 h; (c) (1) HOCH₂CH₂OTBDMS, Ph₃P, DIAD, THF, -5 °C to rt, 2 h; (2) 1% HCl in 95% EtOH, rt, 1 h.

SCHEME 6




- 11: R₂ = (CH₂CH₂O)₂CH₂CH₂F, R₃ = Br
 12: R₂ = (CH₂CH₂O)₂CH₂CH₂F, R₃ = SnBu₃
 13: R₂ = (CH₂CH₂O)₂CH₂CH₂F, R₃ = I
 14: R₂ = CH₂CH₂OH, R₃ = Br
 15: R₂ = CH₂CH₂OH, R₃ = SnBu₃
 16: R₂ = CH₂CH₂OH, R₃ = I

- a: R₁ = N(CH₃)₂
 b: R₁ = NH(CH₃)
 c: R₁ = N(Boc)(CH₃)
 d: R₁ = OCOCH₃
 e: R₁ = OH



Reagents and conditions: (a) 4-substituted styrenes, K₂CO₃, Bu₄NBr, Pd(OAc)₂, DMF, 55-65 °C; (b) (Et₃Sn)₂, Pd(PPh₃)₄, toluene, 110 °C; (c) K₂CO₃, EtOH/THF, rt, 2 h; (d) I₂, THF, 0 °C to rt; (e) TMSOTf, 2,6-lutidine, DCM, -78 °C to rt.

SCHEME 7



[0094] Tc-99m complexes can be prepared as follows. A small amount of non-radiolabeled compound (1-2 mg) is dissolved in 100 μL EtOH and mixed with 200 μL HCl (1 N) and 1 mL Sn-glucoheptonate solution (containing 8-32 μg SnCl₂ and 80-320 μg Na-glucoheptonate, pH 6.67) and 50 μL EDTA solution (0.1 N). [^{99m}Tc]Pertechnetate (100-200 μL ; ranging from 2-20 mCi) saline solution are then added. The reaction is heated for 30 min at 100°C, then cooled to room temperature. The reaction mixture is analyzed on TLC (EtOH:conc. NH₃ 9:1) for product formation and purity check. The mixture can be neutralized with phosphate buffer to pH 5.0.

[0095] The present invention further relates to a method of preparing a technetium-99m complex according to the present invention by reacting technetium-99m in the form of a pertechnetate in the presence of a reducing agent and optionally a suitable chelator with an appropriate Ch-containing compound.

[0096] The reducing agent serves to reduce the Tc-99m pertechnetate which is eluted from a molybdenum-technetium generator in a physiological saline solution. Suitable reducing agents are, for example, dithionite, formamidine sulphinic acid, diaminoethane disulphinate or suitable metallic reducing agents

such as Sn(II), Fe(II), Cu(I), Ti(III) or Sb(III). Sn(II) has proven to be particularly suitable.

[0097] For the above-mentioned complex-forming reaction, technetium-99m is reacted with an appropriate compound of the invention as a salt or in the form of technetium bound to comparatively weak chelators. In the latter case the desired technetium-99m complex is formed by ligand exchange. Examples of suitable chelators for the radionuclide are dicarboxylic acids, such as oxalic acid, malonic acid, succinic acid, maleic acid, orthophthalic acid, malic acid, lactic acid, tartaric acid, citric acid, ascorbic acid, salicylic acid or derivatives of these acids; phosphorus compounds such as pyrophosphates; or enolates. Citric acid, tartaric acid, ascorbic acid, glucoheptonic acid or a derivative thereof are particularly suitable chelators for this purpose, because a chelate of technetium-99m with one of these chelators undergoes the desired ligand exchange particularly easily.

[0098] The most commonly used procedure for preparing $[\text{Tc}^{\text{V}}\text{O}]^{+3}\text{N}_2\text{S}_2$ complexes is based on stannous (II) chloride reduction of $[\text{}^{99\text{m}}\text{Tc}]$ pertechnetate, the common starting material. The labeling procedure normally relies on a Tc-99m ligand exchange reaction between Tc-99m (Sn)-glucoheptonate and the N_2S_2 ligand. Preparation of stannous (II) chloride and preserving it in a consistent stannous (II) form is critically important for the success of the labeling reaction. To stabilize the air-sensitive stannous ion it is a common practice in nuclear medicine to use a lyophilized kit, in which the stannous ion is in a lyophilized powder form mixed with an excess amount of glucoheptonate under an inert gas like nitrogen or argon. The preparation of the lyophilized stannous chloride/sodium glucoheptonate kits ensures that the labeling reaction is reproducible and predictable. The N_2S_2 ligands are usually air-sensitive (thiols are easily oxidized by air) and there are subsequent reactions which lead to decomposition of the ligands. The most convenient and predictable method to preserve the ligands is to produce lyophilized kits containing 100-500 μg of the ligands under argon or nitrogen.

[0099] When the compounds of this invention are to be used as imaging agents, they must be labeled with suitable radioactive halogen isotopes.

Although ^{125}I -isotopes are useful for laboratory testing, they will generally not be useful for actual diagnostic purposes because of the relatively long half-life (60 days) and low gamma-emission (30-65 Kev) of ^{125}I . The isotope ^{123}I has a half life of thirteen hours and gamma energy of 159 KeV, and it is therefore expected that labeling of ligands to be used for diagnostic purposes would be with this isotope. Other isotopes which may be used include ^{131}I (half life of 2 hours). Suitable bromine isotopes include ^{77}Br and ^{76}Br .

[0100] The radiohalogenated compounds of this invention lend themselves easily to formation from materials which could be provided to users in kits. Kits for forming the imaging agents can contain, for example, a vial containing a physiologically suitable solution of an intermediate of Formula I, Ia, II or III in a concentration and at a pH suitable for optimal complexing conditions. The user would add to the vial an appropriate quantity of the radioisotope, e.g., Na^{123}I , and an oxidant, such as hydrogen peroxide. The resulting labeled ligand may then be administered intravenously to a patient, and receptors in the brain imaged by means of measuring the gamma ray or photo emissions therefrom.

[0101] Since the radiopharmaceutical composition according to the present invention can be prepared easily and simply, the preparation can be carried out readily by the user. Therefore, the present invention also relates to a kit, comprising:

(1) A non-radiolabeled compound of the invention, the compound optionally being in a dry condition; and also optionally having an inert, pharmaceutically acceptable carrier and/or auxiliary substances added thereto; and

(2) a reducing agent and optionally a chelator;

wherein ingredients (1) and (2) may optionally be combined; and

further wherein instructions for use with a prescription for carrying out the above-described method by reacting ingredients (1) and (2) with technetium-99m in the form of a pertechnetate solution may be optionally included.

- [0102] Examples of suitable reducing agents and chelators for the above kit have been listed above. The pertechnetate solution can be obtained by the user from a molybdenum-technetium generator. Such generators are available in a number of institutions that perform radiodiagnostic procedures. As noted above the ingredients (1) and (2) may be combined, provided they are compatible. Such a monocomponent kit, in which the combined ingredients are preferably lyophilized, is excellently suitable to be reacted by the user with the pertechnetate solution in a simple manner.
- [0103] When desired, the radioactive diagnostic agent may contain any additive such as pH controlling agents (e.g., acids, bases, buffers), stabilizers (e.g., ascorbic acid) or isotonicizing agents (e.g., sodium chloride).
- [0104] The term "pharmaceutically acceptable salt" as used herein refers to those carboxylate salts or acid addition salts of the compounds of the present invention which are, within the scope of sound medical judgment, suitable for use in contact with the tissues of patients without undue toxicity, irritation, allergic response, and the like, commensurate with a reasonable benefit/risk ratio, and effective for their intended use, as well as the zwitterionic forms, where possible, of the compounds of the invention. The term "salts" refers to the relatively nontoxic, inorganic and organic acid addition salts of compounds of the present invention. Also included are those salts derived from non-toxic organic acids such as aliphatic mono and dicarboxylic acids, for example acetic acid, phenyl-substituted alkanic acids, hydroxy alkanic acids and alkanedioic acids, aromatic acids, and aliphatic and aromatic sulfonic acids. These salts can be prepared *in situ* during the final isolation and purification of the compounds or by separately reacting the purified compound in its free base form with a suitable organic or inorganic acid and isolating the salt thus formed. Further representative salts include the hydrobromide, hydrochloride, sulfate, bisulfate, nitrate, acetate, oxalate, valerate, oleate, palmitate, stearate, laurate, borate, benzoate, lactate, phosphate, tosylate, citrate, maleate, fumarate, succinate, tartrate, naphthylate mesylate, glucoheptonate, lactobionate and laurylsulphonate salts, propionate, pivalate, cyclamate, isethionate, and the like. These may include cations based on the

alkali and alkaline earth metals, such as sodium, lithium, potassium, calcium, magnesium, and the like, as well as, nontoxic ammonium, quaternary ammonium and amine cations including, but not limited to ammonium, tetramethylammonium, tetraethylammonium, methylamine, dimethylamine, trimethylamine, triethylamine, ethylamine, and the like. (See, for example, Berge S. M., *et al.*, *Pharmaceutical Salts*, *J. Pharm. Sci.* 66:1-19 (1977) which is incorporated herein by reference.)

[0105] In the first step of the present method of imaging, a labeled compound of Formula I, Ia, II or III is introduced into a tissue or a patient in a detectable quantity. The compound is typically part of a pharmaceutical composition and is administered to the tissue or the patient by methods well known to those skilled in the art.

[0106] The administration of the labeled compound to a patient can be by a general or local administration route. For example, the compound can be administered either orally, rectally, parenterally (intravenous, by intramuscularly or subcutaneously), intracisternally, intravaginally, intraperitoneally, intravesically, locally (powders, ointments or drops), or as a buccal or nasal spray. The labeled compound may be administered to the patient such that it is delivered throughout the body. Alternatively, the labeled compound can be administered to a specific organ or tissue of interest. For example, it is desirable to locate and quantitate amyloid deposits in the brain in order to diagnose or track the progress of Alzheimer's disease in a patient. One of the most desirable characteristics of an *in vivo* imaging agent of the brain is the ability to cross the intact blood-brain barrier after a bolus *iv* injection.

[0107] In a preferred embodiment of the invention, the labeled compound is introduced into a patient in a detectable quantity and after sufficient time has passed for the compound to become associated with amyloid deposits, the labeled compound is detected noninvasively inside the patient. In another embodiment of the invention, a radiolabeled compound of Formula I, Ia, II or III is introduced into a patient, sufficient time is allowed for the compound to become associated with amyloid deposits, and then a sample of tissue from the

patient is removed and the labeled compound in the tissue is detected apart from the patient. In a third embodiment of the invention, a tissue sample is removed from a patient and a labeled compound of Formula I, Ia, II or III is introduced into the tissue sample. After a sufficient amount of time for the compound to become bound to amyloid deposits, the compound is detected.

[0108] The term "tissue" means a part of a patient's body. Examples of tissues include the brain, heart, liver, blood vessels, and arteries. A detectable quantity is a quantity of labeled compound necessary to be detected by the detection method chosen. The amount of a labeled compound to be introduced into a patient in order to provide for detection can readily be determined by those skilled in the art. For example, increasing amounts of the labeled compound can be given to a patient until the compound is detected by the detection method of choice. A label is introduced into the compounds to provide for detection of the compounds.

[0109] The term "patient" means humans and other animals. Those skilled in the art are also familiar with determining the amount of time sufficient for a compound to become associated with amyloid deposits. The amount of time necessary can easily be determined by introducing a detectable amount of a labeled compound of Formula I, Ia, II or III into a patient and then detecting the labeled compound at various times after administration.

[0110] The term "associated" means a chemical interaction between the labeled compound and the amyloid deposit. Examples of associations include covalent bonds, ionic bonds, hydrophilic-hydrophilic interactions, hydrophobic-hydrophobic interactions, and complexes.

[0111] Those skilled in the art are familiar with the various ways to detect labeled compounds. For example, magnetic resonance imaging (MRI), positron emission tomography (PET), or single photon emission computed tomography (SPECT) can be used to detect radiolabeled compounds. The label that is introduced into the compound will depend on the detection method desired. For example, if PET is selected as a detection method, the compound must possess a positron-emitting atom, such as ^{11}C or ^{18}F .

- [0112] The radioactive diagnostic agent should have sufficient radioactivity and radioactivity concentration which can assure reliable diagnosis. For instance, in case of the radioactive metal being technetium-99m, it may be included usually in an amount of 0.1 to 50 mCi in about 0.5 to 5.0 ml at the time of administration. The amount of a compound of Formula I, Ia, II or III may be such as sufficient to form a stable chelate compound with the radioactive metal.
- [0113] The thus formed chelate compound as a radioactive diagnostic agent is sufficiently stable, and therefore it may be immediately administered as such or stored until its use. When desired, the radioactive diagnostic agent may contain any additive such as pH controlling agents (e.g., acids, bases, buffers), stabilizers (e.g., ascorbic acid) or isotonicizing agents (e.g., sodium chloride).
- [0114] The imaging of amyloid deposits can also be carried out quantitatively so that the amount of amyloid deposits can be determined.
- [0115] Preferred compounds for imaging include a radioisotope such as ^{11}C , ^{123}I , ^{125}I , ^{131}I , ^{18}F , ^{76}Br or ^{77}Br .
- [0116] Another aspect of the invention is a method of inhibiting amyloid plaque aggregation. The present invention also provides a method of inhibiting the aggregation of amyloid proteins to form amyloid deposits, by administering to a patient an amyloid inhibiting amount of a compound of the above Formula I, Ia, II or III.
- [0117] Those skilled in the art are readily able to determine an amyloid inhibiting amount by simply administering a compound of Formula I, Ia, II or III to a patient in increasing amounts until the growth of amyloid deposits is decreased or stopped. The rate of growth can be assessed using imaging as described above or by taking a tissue sample from a patient and observing the amyloid deposits therein. The compounds of the present invention can be administered to a patient at dosage levels in the range of about 0.1 to about 1,000 mg per day. For a normal human adult having a body weight of about 70 kg, a dosage in the range of about 0.01 to about 100 mg per kilogram of body weight per day is sufficient. The specific dosage used, however, can vary. For example, the dosage can depend on a number of factors including the

requirements of the patient, the severity of the condition being treated, and the pharmacological activity of the compound being used. The determination of optimum dosages for a particular patient is well known to those skilled in the art.

[0118] The following examples are illustrative, but not limiting, of the method and compositions of the present invention. Other suitable modifications and adaptations of the variety of conditions and parameters normally encountered and obvious to those skilled in the art are within the spirit and scope of the invention.

[0119] All reagents used in synthesis were commercial products and were used without further purification unless otherwise indicated. ¹H NMR spectra were obtained on a Bruker DPX spectrometer (200 MHz) in CDCl₃. Chemical shifts are reported as δ values (parts per million) relative to internal TMS. Coupling constants are reported in hertz. The multiplicity is defined by s (singlet), d (doublet), t (triplet), br (broad), m (multiplet). Elemental analyses were performed by Atlantic Microlab INC. For each procedure, "standard workup" refers to the following steps: addition of indicated organic solvent, washing the organic layer with water then brine, separation of the organic layer from the aqueous layer, drying off the combined the organic layers with anhydrous sodium sulfate, filtering off the sodium sulfate and removing the organic solvent under reduced pressure.

EXAMPLES

EXAMPLE 1

Synthesis of Compound 2

(E)-2-chloro-5-(4-dimethylaminostyryl)pyridine (1)

[0120] Potassium *tert*-butoxide (99 mg, 0.89 mmol) was added to a solution of diethyl-(4-dimethylamino-benzyl)-phosphonate (80 mg, 0.30 mmol) in anhydrous DMF (5.0 ml) at 0 °C. 2-Chloro-5-pyridyl aldehyde (42 mg, 0.30 mmol) was then added. The reaction mixture was warmed to room

temperature and stirred for 4 h. Water was added and mixture was extracted with MeOH/DCM (1: 9, v/v). Organic layer was separated, washed with brine, dried over sodium sulfate and evaporated. The residue was purified by PTLC (20 % Hexanes in DCM as developing solvent) to give product 1 (48 mg, Yield: 62 %). ^1H NMR (200 MHz, CDCl_3): δ 8.42 (1H, d, $J = 2.2$ Hz), 7.77 (1H, d, d, $J_1 = 8.4$ Hz, $J_2 = 2.4$ Hz), 7.41 (2H, d, $J = 8.6$ Hz), 7.27 (1H, d, $J = 8.2$ Hz), 7.08 (1H, d, $J = 16.4$ Hz), 6.77 (3H, m), HRMS (EI) m/z calcd. for $[\text{C}_{13}\text{H}_9\text{ClN}_2\text{O}_2]^+$ 260.0353.

(E)-2-(2-(2-(2-fluoroethoxy)ethoxy)ethoxy)-5-(4-dimethylaminostyryl)pyridine (2)

- [0121] Sodium hydride (95%, 10 mg, 0.39 mmol) was added to a solution of 2-(2-(2-(2-fluoroethoxy)ethoxy)ethanol (39 mg, 0.26 mmol) in anhydrous DMF (5.0 ml). After stirring at room temperature for 20 min, compound 5 (35 mg, 0.13 mmol) was added and reaction mixture was heated to 100 °C for 2 h. After cooling down to room temperature, water was added and the reaction mixture was extracted with ethyl acetate. Organic layer was separated, washed with brine, dried over anhydrous sodium sulfate, and evaporated. The residue was purified by PTLC. (4% MeOH in DCM as developing solvent) to give product 2 (16 mg, Yield; 32.9%): ^1H NMR (200 MHz, CDCl_3): δ 8.14 (1H, d, $J = 2.4$ Hz), 7.76 (1H, d, d, $J_1 = 8.6$ Hz, $J_2 = 2.4$ Hz), 7.39 (2H, d, $J = 8.8$ Hz), 6.87 (2H, m), 6.76 (3H, m), 4.53 (2H, d, t, $J_1 = 47.6$ Hz, $J_2 = 4.2$ Hz), 4.50 (2H, t, $J = 4.8$ Hz), 3.85 (3H, m), 3.70 (5H, m), 2.99 (6H, s). HRMS (EI) m/z calcd. for $[\text{C}_{21}\text{H}_{28}\text{N}_2\text{O}_4]^+$ 372.2049.

EXAMPLE 2

Synthesis of Compound 6

(E)-2-chloro-5-(4-nitrostyryl)pyridine (3):

- [0122] Sodium methoxide (1 M in methanol, 5.0 ml) was added slowly into a solution of diethyl-(4-nitro-benzyl)-phosphonate (546 mg, 2.0 mmol) and 2-chloro-5-pyridyl aldehyde (283 mg, 2.0 mmol) in methanol (5.0 ml). The

reaction mixture was then refluxed for 1 h. After cooled down to 0 °C, yellow precipitate was filtered and washed with cold methanol to obtain product 3 (458 mg, Yield: 88%), which was used directly for next step without further purification. 3: ¹H NMR (200 MHz, CDCl₃): δ 8.53 (1H, d, *J* = 2.4 Hz), 8.25 (2H, d, *J* = 8.8 Hz), 7.85 (1H, d, d, *J*₁ = 8.4 Hz, *J*₂ = 2.4 Hz), 7.65 (2H, d, *J* = 8.8 Hz), 7.36 (1H, d, *J* = 8.4 Hz), 7.19 (2H, s), HRMS (EI) *m/z* calcd. for [C₁₃H₉ClN₂O₂]⁺ 260.0353.

(E)-2-(2-(2-(2-fluoroethoxy)ethoxy)ethoxy)-5-(4-nitrostyryl)pyridine (4)

[0123] Under the protection of nitrogen atmosphere, 2-(2-(2-fluoroethoxy)-ethoxy)-ethanol^a (60 mg, 0.39 mmol) was added into a mixture of sodium hydride (26.4 mg, 60 % dispersion in mineral oil, 0.66 mmol) in anhydrous DMF (5 ml) at 0 °C. The mixture was stirred at room temperature for half an hour and compound 3 (85.7 mg, 0.33 mmol) was added. The reaction mixture was then heated to 100 °C for 2 hours and cooled down. Ethyl acetate and water was added, organic layer was separated, washed with brine, dried over anhydrous sodium sulfate and evaporated. The residue was purified by PTLC (2% MeOH in DCM as developing solvent) to give product 4 (37 mg, Yield: 30%): ¹H NMR (200 MHz, CDCl₃): δ 8.822 (3H, d, *J* = 8.8 Hz), 7.84 (1H, d, d, *J*₁ = 8.6 Hz, *J*₂ = 2.4 Hz), 7.61 (2H, d, *J* = 8.8 Hz), 7.20 (1H, d, *J* = 16.4 Hz), 7.02 (1H, d, *J* = 16.4 Hz), 6.84 (1H, d, *J* = 8.6 Hz), 4.53 (2H, d, t, *J*₁ = 47.6 Hz, *J*₂ = 4.2 Hz), 4.52 (2H, t, *J* = 4.8 Hz), 3.85 (3H, m), 3.70 (5H, m); HRMS (EI) *m/z* calcd. for [C₁₉H₂₁FN₂O₅]⁺ 376.1435.

(E)-2-(2-(2-(2-fluoroethoxy)ethoxy)ethoxy)-5-(4-aminostyryl)pyridine (5)

[0124] Compound 4 (34 mg, 0.09 mmol) was dissolved in ethanol (5 ml) followed by the addition of stannous chloride (51.4 mg, 0.27 mmol) and concentrated HCl (0.25 ml). The reaction mixture was refluxed for 2 hours and cooled down. 2N NaOH was used to adjust pH to 10. Dichloromethane was added and the organic layer was separated, washed with brine, dried over anhydrous sodium sulfate and evaporated. The residue was purified by PTLC (3% MeOH in DCM as developing solvent) to give product 5 (18 mg, Yield:

58%): ^1H NMR (200 MHz, CDCl_3): δ 8.14 (1H, d, $J = 2.2$ Hz), 7.76 (1H, d, d, $J_1 = 8.6$ Hz, $J_2 = 2.4$ Hz), 7.32 (2H, d, $J = 8.4$ Hz), 6.80 (5H, m), 4.53 (2H, d, t, $J_1 = 47.6$ Hz, $J_2 = 4.2$ Hz), 4.49 (2H, t, $J = 4.8$ Hz), 3.85 (3H, m), 3.70 (5H, m), 1.8 -3.0 (2H, br); HRMS (EI) m/z calcd. for $[\text{C}_{19}\text{H}_{23}\text{FN}_2\text{O}_3]^+$ 376.1693.

(E)-2-(2-(2-(2-fluoroethoxy)ethoxy)ethoxy)-5-(4-methylaminostyryl)pyridine (6)

[0125] Sodium methoxide (1M in methanol, 0.23 ml) was added to a solution of compound 5 (15.8 mg, 0.046 mmol) in methanol (5 ml) followed by the addition of paraformaldehyde (6.6 mg, 0.23 mmol). The reaction mixture was refluxed for 1.5 hour then cooled to 0 °C with an ice bath. Sodium borohydride (10.4 mg, 0.27 mmol) was added with caution. The mixture was refluxed again for 1 hour and cooled down. Dichloromethane and water was added. Organic layer was separated, washed with brine, dried over anhydrous sodium sulfate and evaporated. The residue was purified by PTLC (3% MeOH in DCM as developing solvent) to give product 6 (12 mg, Yield: 73%): ^1H NMR (200 MHz, CDCl_3): δ 8.14 (1H, d, $J = 2.2$ Hz), 7.76 (1H, d, d, $J_1 = 8.6$ Hz, $J_2 = 2.4$ Hz), 7.35 (2H, d, $J = 8.6$ Hz), 6.92 (1H, d, $J = 16.4$ Hz), 6.80 (1H, d, $J = 16.4$ Hz), 6.76 (2H, d, $J = 8.6$ Hz), 4.53 (2H, d, t, $J_1 = 47.6$ Hz, $J_2 = 4.2$ Hz), 4.49 (2H, t, $J = 4.8$ Hz), 3.85 (3H, m), 3.70 (5H, m), 2.88 (3H, s). HRMS (EI) m/z calcd. for $[\text{C}_{20}\text{H}_{25}\text{FN}_2\text{O}_3]^+$ 360.1849.

EXAMPLE 3

Synthesis of Compound 10

(E)-2-(2-(2-(2-hydroxyethoxy)ethoxy)ethoxy)-5-(4-nitrostyryl)pyridine (7)

[0126] The mixture of potassium carbonate (158.7 mg, 1.15 mmol), compound 3 (100 mg, 0.38 mmol) and triethylene glycol (576 mg, 3.8 mmol) in anhydrous DMF (5.0 ml) was sealed in a microwavable vial (from Biotage) and subjected to microwave irradiation (Biotage Initiator system) at 180 °C for 25 min. After cooling to room temperature, water was added and reaction

mixture was extracted with ethyl acetate. Organic layer was separated, washed with brine, dried over anhydrous sodium sulfate and evaporated. The residue was purified with PTLC (4% MeOH in DCM as developing solvent) gave product **7** (110 mg, Yield: 77%): ^1H NMR (200 MHz, CDCl_3): δ 8.20 (3H, m), 7.83 (1H, d, d, $J_1 = 8.6$ Hz, $J_2 = 2.4$ Hz), 7.61 (2H, d, $J = 8.8$ Hz), 7.10 (2H, m), 6.84 (1H, d, $J = 8.6$ Hz), 4.53 (2H, t, $J = 4.8$ Hz), 3.88 (2H, t, $J = 4.8$ Hz), 3.71 (6H, m), 3.61 (2H, m), 2.10 (1H, b), HRMS (EI) m/z calcd. for $[\text{C}_{19}\text{H}_{22}\text{N}_2\text{O}_6]^+$ 374.1478.

(E)-2-(2-(2-(2-hydroxyethoxy)ethoxy)ethoxy)-5-(4-aminostyryl)pyridine (**8**)

[0127] Stannous chloride (202.8 mg, 1.07 mmol) was added to a solution of compound **7** (100 mg 0.27 mmol) in ethanol (10 ml) followed by the addition of concentrated HCl (0.5 ml). The reaction mixture was refluxed for 1.5 h and then cooled to 0 °C. Yellow precipitate was collected by filtration and then suspended in ethyl acetate. Saturated NaHCO_3 was added to adjust pH to 9. Organic layer was separated, dried over anhydrous sodium sulfate and evaporated. The residue was purified by PTLC (5% MeOH in DCM as developing solvent) to give product **8** (70 mg, Yield: 76%): ^1H NMR (200 MHz, CDCl_3): δ 8.12 (1H, d, $J = 2.4$ Hz), 7.73 (1H, d, d, $J_1 = 8.6$ Hz, $J_2 = 2.4$ Hz), 7.29 (2H, d, $J = 8.5$ Hz), 6.84 (2H, m), 6.75 (1H, d, $J = 8.6$ Hz), 6.69 (2H, d, $J = 8.5$ Hz), 4.48 (2H, t, $J = 4.8$ Hz), 3.86 (2H, t, $J = 4.8$ Hz), 3.71 (6H, m), 3.60 (2H, m), 3.32 (3H, b), HRMS (EI) m/z calcd. for $[\text{C}_{19}\text{H}_{24}\text{N}_2\text{O}_4]^+$ 344.1736.

(E)-2-(2-(2-(2-Hydroxyethoxy)ethoxy)ethoxy)-5-(4-dimethylaminostyryl)pyridine (**9**)

[0128] Sodium cyanoborohydride (36 mg, 0.57 mmol) was added to a solution of compound **8** (65 mg, 0.19 mmol) and paraformaldehyde (57 mg, 1.9 mmol) in acetic acid (10 ml). The reaction mixture was stirred at room temperature overnight and poured onto ice. Sodium bicarbonate was used to adjust pH to 9. Reaction mixture was extracted with ethyl acetate. Organic layer was separated, washed with brine, dried over anhydrous sodium sulfate and

evaporated. The residue was purified with PTLC (5% MeOH in DCM as the developing solvent) to give product **9** (67 mg, Yield: 95%): ^1H NMR (200 MHz, CDCl_3): δ 8.14 (1H, d, $J = 2.4$ Hz), 7.76 (1H, d, d, $J_1 = 8.6$ Hz, $J_2 = 2.4$ Hz), 7.39 (2H, d, $J = 8.8$ Hz), 6.87 (2H, m), 6.76 (3H, m), 4.50 (2H, t, $J = 4.8$ Hz), 3.87 (2H, t, $J = 4.8$ Hz), 3.70 (6H, m), 3.61 (2H, m), 2.98 (6H, s), 2.49 (1H, b), HRMS (EI) m/z calcd. for $[\text{C}_{21}\text{H}_{28}\text{N}_2\text{O}_4]^+$ 372.2049.

(E)-2-(2-(2-(2-tosyloxyethoxy)ethoxy)ethoxy)-5-(4-dimethylaminostyryl)pyridine (**10**)

[0129] Tosyl chloride (52 mg, 0.27 mmol) was added to a solution of compound **9** (43 mg, 0.116 mmol) in pyridine (5.0 ml) at 0 °C. The reaction mixture was stirred at 0 °C for 1 h and then warmed up to room temperature and stirred for 3 h. Water was added and reaction mixture was extracted with ethyl acetate. Organic layer was separated, washed with brine, dried over sodium sulfate and evaporated. The residue was purified by PTLC (4% MeOH in DCM as developing solvent) to give product **10** (25 mg, Yield: 41%): ^1H NMR (200 MHz, CDCl_3): δ 8.14 (1H, d, $J = 2.0$ Hz), 7.76 (3H, m), 7.39 (2H, d, $J = 8.8$ Hz), 7.32 (2H, d, $J = 8.0$ Hz), 6.87 (2H, m), 6.75 (3H, m), 4.46 (2H, t, $J = 4.6$ Hz), 4.16 (2H, t, $J = 4.8$ Hz), 3.81 (2H, t, $J = 4.8$ Hz), 3.66 (6H, m), 2.99 (6H, s), 2.43 (3H, s), HRMS (EI) m/z calcd. for $[\text{C}_{28}\text{H}_{34}\text{N}_2\text{O}_6\text{S}]^+$ 526.2138.

EXAMPLE 4

Synthesis of Compound 11a

a. Synthesis of intermediates **I8** and **I9**

2-Hydroxy-3-bromo-5-iodopyridine (**I8**)

[0130] Following a previously reported method (Meana A, et al, Synlett 2003, 1678-1682) compound **I8** was prepared from N-iodosuccinimide (2.48 g, 11.0 mmol) and 3-bromo-2-hydroxypyridine **I7** (1.74 g, 10.0 mmol) as a pale

brown solid (2.55 g, 85%). ^1H NMR (DMSO- d_6) δ 12.27 (br s, 1H), 8.08 (d, 1H, $J = 2.3$ Hz), 7.71 (d, 1H, $J = 2.3$ Hz).

{2-[2-(2-Fluoroethoxy)ethoxy]ethoxy}-3-bromo-5-iodopyridine (I9)

[0131] To a stirring suspension of I8 (0.393 g, 1.3 mmol), 2-(2-(2-fluoroethoxy)ethoxy)ethanol (0.200 g, 1.3 mmol) and PPh_3 (0.511 g, 1.95 mmol) in 10 mL of THF at -10°C was added dropwise of diisopropyl azodicarboxylate (DIAD, 0.394 g, 1.95 mmol) in 5 mL of THF. The ice-salt bath was removed and the reaction was kept at room temperature (r.t.) 2 h. The reaction solution was concentrated and purified by FC (MeOH/ CHCl_3 , 1/99) to yield I9, a colorless viscous liquid (0.423 g, 75%). ^1H NMR δ 8.21 (d, 1H, $J = 2.0$ Hz), 8.02 (d, 1H, $J = 2.0$ Hz), 4.66 (t, 1H, $J = 4.1$ Hz), 4.50-4.39 (m, 3H), 3.89-3.64 (m, 8H). ^{13}C NMR δ 159.4, 151.2, 148.5, 108.5, 84.9, 81.6, 81.5, 71.1, 71.0, 70.8, 70.4, 69.3, 66.9. HRMS calcd for $\text{C}_{11}\text{H}_{14}\text{BrFINO}_3$ (M^+), 432.9186; found, 432.9173.

b. Synthesis of compound 11a

(*E*)-(5-Bromo-6-{2-[2-(2-fluoroethoxy)ethoxy]ethoxy}pyridin-3-yl)-2-(4-dimethylaminophenyl)-ethylene (11a)

[0132] A mixture of 4-dimethylaminostyrene (0.110 g, 0.75 mmol), I9 (0.217 g, 0.5 mmol), K_2CO_3 (0.173 g, 1.25 mmol), tetrabutylammonium bromide (TBAB, 0.322 g, 1.0 mmol) and palladium acetate ($\text{Pd}(\text{OAc})_2$, 0.006 g, 0.025 mmol) in 2 mL DMF was deoxygenated by purging into nitrogen for 15 min and then heated to 65°C for 2 h. The reaction mixture was cooled to r.t. and submitted to standard workup with ethyl acetate (EtOAc). The crude product was purified by FC (EtOAc/Hexanes, 30/70) and resulted in 11a as a light yellow solid (0.178 g, 79%). ^1H NMR δ 8.08 (d, 1H, $J = 2.1$ Hz), 8.00 (d, 1H, $J = 2.1$ Hz), 7.39 (d, 2H, $J = 8.8$ Hz), 6.92 (d, 1H, $J = 16.3$ Hz), 6.74 (d, 1H, $J = 16.3$ Hz), 6.72 (d, 2H, $J = 8.1$ Hz), 4.69 (t, 1H, $J = 4.2$ Hz), 4.55 (t, 2H, $J = 4.8$ Hz), 4.45 (t, 1H, $J = 4.2$ Hz), 3.94-3.68 (m, 8H), 3.00 (s, 6H). ^{13}C NMR δ

158.3, 150.4, 143.5, 138.0, 129.6, 129.5, 127.7, 125.2, 118.8, 112.5, 107.5, 85.0, 81.6, 71.2, 71.0, 70.8, 70.4, 69.6, 66.7, 40.5. HRMS calcd for $C_{21}H_{26}BrFN_2O_3$ (M^+), 452.1111; found, 452.1099.

EXAMPLE 5

Synthesis of Compound 11b

(*E*)-(5-Bromo-6-{2-[2-(2-fluoroethoxy)ethoxy]ethoxy}pyridin-3-yl)-2-(4-methylaminophenyl)-ethylene (11b)

[0133] Compound 11b was prepared from 4-methylaminostyrene (0.073 g, 0.55 mmol) and **I9** (0.217 g, 0.50 mmol) as a light yellow viscous liquid (0.113g, 52% yield). 1H NMR δ 8.07 (d, 1H, $J = 2.1$ Hz), 8.00 (d, 1H, $J = 2.1$ Hz), 7.35 (d, 2H, $J = 8.6$ Hz), 6.91 (d, 1H, $J = 16.3$ Hz), 6.74 (d, 1H, $J = 16.3$ Hz), 6.60 (d, 2H, $J = 8.6$ Hz), 4.69 (t, 1H, $J = 4.2$ Hz), 4.55 (t, 2H, $J = 4.8$ Hz), 4.45 (t, 1H, $J = 4.2$ Hz), 3.94-3.68 (m, 8H), 2.88 (s, 3H). ^{13}C NMR δ 158.4, 149.5, 143.6, 138.0, 129.8, 129.5, 127.9, 126.1, 118.9, 112.6, 107.5, 85.0, 81.7, 71.2, 71.1, 70.8, 70.4, 69.6, 66.8, 30.7. HRMS calcd for $C_{20}H_{24}BrFN_2O_3$ (M^+), 438.0954; found, 438.0967.

EXAMPLE 6

Synthesis of Compound 11e

(*E*)-(5-Bromo-6-{2-[2-(2-fluoroethoxy)ethoxy]ethoxy}pyridin-3-yl)-2-[4-*N*-methyl-4-*N*-(tert-butyloxycarbonyl)aminophenyl]-ethylene (11c)

[0134] Compound 11c was prepared from 4-*N*-methyl-4-*N*-(tert-butyloxycarbonyl)aminostyrene (0.219 g, 0.94 mmol) and **I9** (0.273 g, 0.63 mmol) as a white viscous liquid (0.319 g, 94% yield). 1H NMR δ 8.12 (d, 1H, $J = 2.1$ Hz), 8.03 (d, 1H, $J = 2.1$ Hz), 7.44 (d, 2H, $J = 8.6$ Hz), 7.25 (d, 2H, $J = 9.0$ Hz), 6.94 (d, 2H, $J = 2.1$ Hz), 4.69 (t, 1H, $J = 4.2$ Hz), 4.56 (t, 2H, $J = 4.9$ Hz), 4.45 (t, 1H, $J = 4.2$ Hz), 3.94-3.68 (m, 8H), 3.28 (s, 3H), 1.48 (s, 9H). ^{13}C NMR δ 158.8, 154.5, 144.0, 143.5, 138.2, 133.6, 128.5, 128.4, 126.8,

126.6, 125.4, 122.9, 107.4, 84.8, 81.4, 80.4, 71.0, 70.9, 70.6, 70.2, 69.4, 66.7, 53.5, 37.1, 28.4. HRMS calcd for $C_{25}H_{32}BrFN_2O_5$ (M^+), 538.1479; found, 538.1476.

(*E*)-(5-Bromo-6-{2-[2-(2-fluoroethoxy)ethoxy]ethoxy}pyridin-3-yl)-2-(4-acetoxyphenyl)-ethylene (11d)

[0135] Compound 11d was prepared from 4-acetoxystyrene (0.122 g, 0.75 mmol) and **I9** (0.217 g, 0.5 mmol) as a white viscous liquid (0.181 g, 77% yield). 1H NMR δ 8.12 (d, 1H, $J = 2.1$ Hz), 8.03 (d, 1H, $J = 2.1$ Hz), 7.50 (d, 2H, $J = 8.6$ Hz), 7.10 (d, 2H, $J = 8.6$ Hz), 6.94 (d, 2H, $J = 3.3$ Hz), 4.69 (t, 1H, $J = 4.2$ Hz), 4.56 (t, 2H, $J = 4.9$ Hz), 4.45 (t, 1H, $J = 4.2$ Hz), 3.94-3.68 (m, 8H), 2.32 (s, 3H), 1.48 (s, 9H). ^{13}C NMR δ 169.3, 158.9, 150.3, 144.1, 138.2, 134.5, 128.24, 128.16, 127.4, 123.4, 121.9, 107.5, 84.8, 81.5, 71.0, 70.9, 70.6, 70.3, 69.4, 66.7, 21.1. HRMS calcd for $C_{21}H_{23}BrFNO_5$ (M^+), 467.0744; found, 467.0731.

(*E*)-(5-Bromo-6-{2-[2-(2-fluoroethoxy)ethoxy]ethoxy}pyridin-3-yl)-2-(4-hydroxyphenyl)-ethylene (11e)

[0136] Acetate 11d (0.145 g, 0.31 mmol) and K_2CO_3 (0.064 g, 0.465 mmol) were placed in EtOH/THF (5 mL/5 mL) and the reaction mixture was stirred at r.t. 2 h. After standard workup with EtOAc, the crude product was purified by PTLC to give 11e as a white solid (0.128 g, 97%). 1H NMR δ 8.07 (d, 1H, $J = 2.1$ Hz), 7.99 (d, 1H, $J = 2.1$ Hz), 7.35 (d, 2H, $J = 8.6$ Hz), 6.96-6.74 (m, 4H), 5.22 (br s, 1H), 4.69 (t, 1H, $J = 4.2$ Hz), 4.54 (t, 2H, $J = 4.8$ Hz), 4.45 (t, 1H, $J = 4.2$ Hz), 3.94-3.68 (m, 8H). ^{13}C NMR δ 158.5, 156.4, 143.6, 138.2, 129.2, 129.0, 127.9, 120.7, 116.0, 107.6, 84.9, 81.6, 71.1, 71.0, 70.8, 70.4, 69.6, 66.8. HRMS calcd for $C_{19}H_{21}BrFNO_4$ (M^+), 425.0638; found, 425.0651.

EXAMPLE 7

Synthesis of Compound 12b

(*E*)-(5-tri-Butylstannyl-6-{2-[2-(2-fluoroethoxy)ethoxy]ethoxy}pyridin-3-yl)-2-(4-methylaminophenyl)-ethylene (12b)

[0137] Compound 12b was prepared from 11b (0.069 g, 0.156 mmol) as a light yellow oil (0.068 g, 68% yield). ^1H NMR δ 8.10 (d, 1H, $J = 2.5$ Hz), 7.80 (d, 1H, $J = 2.5$ Hz), 7.36 (d, 2H, $J = 8.6$ Hz), 6.92 (d, 1H, $J = 16.3$ Hz), 6.80 (d, 1H, $J = 16.3$ Hz), 6.61 (d, 2H, $J = 8.6$ Hz), 4.69 (t, 1H, $J = 4.2$ Hz), 4.45 (t, 3H, $J = 5.1$ Hz), 3.83 (t, 3H, $J = 4.4$ Hz), 3.71-3.66 (m, 5H), 2.88 (s, 3H), 1.68-1.48 (m, 6H), 1.43-1.25 (m, 6H), 1.15-1.02 (m, 6H), 0.91 (t, 9H, $J = 7.1$ Hz). ^{13}C NMR δ 166.8, 149.1, 145.4, 143.6, 127.8, 127.7, 127.0, 123.8, 121.2, 112.6, 85.0, 81.6, 71.1, 70.9, 70.8, 70.5, 70.1, 65.0, 30.8, 29.5, 29.3, 29.1, 28.1, 27.5, 26.9, 13.9, 13.4, 13.3, 9.9, 6.6, 6.4. HRMS calcd for $\text{C}_{32}\text{H}_{51}\text{FN}_2\text{O}_3\text{Sn}$ (M^+), 650.2906; found, 650.2894.

EXAMPLE 8

Synthesis of Compound 12e

(*E*)-(5-tri-Butylstannyl-6-{2-[2-(2-fluoroethoxy)ethoxy]ethoxy}pyridin-3-yl)-2-(4-hydroxyphenyl)-ethylene (12e)

[0138] Compound 12e was prepared from 11e (0.032 g, 0.075 mmol) as a white viscous liquid (0.040 g, 84% yield). ^1H NMR δ 8.11 (d, 1H, $J = 2.5$ Hz), 7.82 (d, 1H, $J = 2.5$ Hz), 7.39 (d, 2H, $J = 8.6$ Hz), 6.98-6.74 (m, 4H), 5.19 (br s, 1H), 4.71-4.66 (m, 1H), 4.48-4.43 (m, 3H), 3.90-3.62 (m, 8H), 1.70-1.02 (m, 18H), 0.91 (t, 9H, $J = 7.1$ Hz). ^{13}C NMR δ 166.9, 156.0, 145.4, 144.0, 130.1, 127.9, 127.6, 127.4, 124.3, 123.0, 115.9, 85.0, 81.6, 71.0, 70.9, 70.7, 70.5, 70.0, 65.2, 29.5, 29.3, 29.1, 28.0, 27.5, 26.9, 13.9, 13.4, 13.3, 9.9, 6.6, 6.4. HRMS calcd for $\text{C}_{31}\text{H}_{48}\text{FNO}_4\text{Sn}$ (M^+), 637.2589; found, 637.2573.

EXAMPLE 9

Synthesis of Compound 13a

(*E*)-(5-tri-Butylstannyl-6-{2-[2-(2-fluoroethoxy)ethoxy]ethoxy}pyridin-3-yl)-2-(4-dimethylaminophenyl)-ethylene (**12a**)

[0139] A mixture of **11a** (0.052 g, 0.115 mmol), bis(tributyltin) ((Bu₃Sn)₂, 0.333 g, 0.57 mmol), and palladium tetrakis(triphenylphosphine) (Pd(PPh₃)₄, 0.013 g, 10 mol%) in toluene was heated at 110 °C for 18 h. The reaction solution was cooled to r.t. and treated with 5 mL 10% KF. After vigorously stirring for additional 0.5 h, the standard workup with EtOAc and following FC (EtOAc/Hexanes, 25/75) afforded **12a** as a light yellow oil (0.052 g, 68%). ¹H NMR δ 8.11 (d, 1H, *J* = 2.5 Hz), 7.81 (d, 1H, *J* = 2.5 Hz), 7.41 (d, 2H, *J* = 8.8 Hz), 6.93 (d, 1H, *J* = 16.5 Hz), 6.81 (d, 1H, *J* = 16.5 Hz), 6.72 (d, 2H, *J* = 8.7 Hz), 4.69 (t, 1H, *J* = 4.2 Hz), 4.46 (t, 3H, *J* = 4.9 Hz), 3.83 (t, 3H, *J* = 4.8 Hz), 3.71-3.66 (m, 5H), 3.00 (s, 6H), 1.68-1.48 (m, 6H), 1.43-1.21 (m, 6H), 1.15-1.02 (m, 6H), 0.91 (t, 9H, *J* = 7.1 Hz). ¹³C NMR δ 166.7, 150.2, 145.4, 143.6, 127.8, 127.7, 127.5, 126.0, 123.7, 121.2, 112.6, 85.0, 81.6, 71.0, 70.8, 70.7, 70.4, 70.0, 65.0, 40.6, 29.5, 29.3, 29.1, 28.1, 27.5, 26.9, 13.9, 13.4, 13.3, 9.9, 6.6, 6.4. HRMS calcd for C₃₃H₅₃FN₂O₃Sn (M⁺), 664.3062; found, 664.3037.

(*E*)-(5-Iodo-6-{2-[2-(2-fluoroethoxy)ethoxy]ethoxy}pyridin-3-yl)-2-(4-dimethylaminophenyl)-ethylene (**13a**)

[0140] A solution of iodine (I₂, 0.063 g, 0.24 mmol) in THF (2 mL) was added dropwise to an ice bath cooled solution of **12a** (0.114 g, 0.172 mmol) in THF (3 mL). After the addition, the reaction was stirred at 0 °C for 1 h. Following standard workup with CH₂Cl₂, the crude product was purified by FC (EtOAc/Hexanes, 25/75) to give a light yellow solid **13a** (0.037 g, 48%). ¹H NMR δ 8.22 (d, 1H, *J* = 2.1 Hz), 8.10 (d, 1H, *J* = 2.1 Hz), 7.38 (d, 2H, *J* = 8.8 Hz), 6.92 (d, 1H, *J* = 16.3 Hz), 6.72 (d, 1H, *J* = 16.3 Hz), 6.71 (d, 2H, *J* =

8.8 Hz), 4.72-4.67 (m, 1H), 4.54-4.44 (m, 3H), 3.93-3.69 (m, 8H), 3.00 (s, 6H). ^{13}C NMR δ 160.4, 150.5, 144.6, 144.55, 129.8, 129.5, 127.8, 125.3, 118.8, 112.6, 85.1, 81.7, 80.6, 71.3, 71.1, 70.8, 70.5, 69.6, 67.1, 40.6. HRMS calcd for $\text{C}_{21}\text{H}_{26}\text{FIN}_2\text{O}_3$ (M^+), 500.0972; found, 500.0959.

EXAMPLE 10

Synthesis of Compound 13b

(*E*)-(5-tri-Butylstannyl-6-{2-[2-(2-fluoroethoxy)ethoxy]ethoxy}pyridin-3-yl)-2-[4-*N*-methyl-4-*N*-(tert-butyloxycarbonyl)aminophenyl]-ethylene (**12c**)

[0141] Compound **12c** was prepared from **11c** (0.072 g, 0.133 mmol) as a white viscous liquid (0.077 g, 77% yield). ^1H NMR δ 8.14 (d, 1H, $J = 2.5$ Hz), 7.83 (d, 1H, $J = 2.5$ Hz), 7.46 (d, 2H, $J = 8.6$ Hz), 7.23 (d, 2H, $J = 8.5$ Hz), 6.96 (s, 2H), 4.70-4.66 (m, 1H), 4.49-4.42 (m, 3H), 3.86-3.66 (m, 8H), 3.28 (s, 3H), 1.80-1.02 (m, 27H), 0.90 (t, 9H, $J = 7.1$ Hz). ^{13}C NMR δ 167.3, 146.1, 143.8, 143.2, 134.6, 127.0, 126.8, 126.6, 125.7, 125.4, 124.1, 85.0, 81.6, 80.6, 71.1, 70.9, 70.8, 70.5, 70.0, 65.1, 37.4, 29.5, 29.3, 29.1, 28.1, 27.5, 26.9, 13.9, 13.4, 9.9, 6.4. HRMS calcd for $\text{C}_{37}\text{H}_{59}\text{FN}_2\text{O}_5\text{Sn}$ (M^+), 750.343; found, 750.3425.

(*E*)-(5-Iodo-6-{2-[2-(2-fluoroethoxy)ethoxy]ethoxy}pyridin-3-yl)-2-[4-*N*-methyl-4-*N*-(tert-butyloxycarbonyl)aminophenyl]-ethylene (**13c**)

[0142] Compound **13c** was prepared from **12c** (0.024 g, 0.032 mmol) as a white viscous liquid (0.018 g, 98%). ^1H NMR δ 8.25 (d, 1H, $J = 1.6$ Hz), 8.13 (d, 1H, $J = 1.6$ Hz), 7.44 (d, 2H, $J = 8.4$ Hz), 7.24 (d, 2H, $J = 8.4$ Hz), 6.97 (d, 1H, $J = 16.4$ Hz), 6.86 (d, 1H, $J = 16.4$ Hz), 4.69 (t, 1H, $J = 4.1$ Hz), 4.53 (t, 2H, $J = 4.8$ Hz), 4.45 (t, 1H, $J = 4.1$ Hz), 3.94-3.69 (m, 8H), 3.28 (s, 3H), 1.47 (s, 9H). ^{13}C NMR δ 161.0, 154.8, 145.3, 144.9, 143.7, 133.9, 128.9, 128.6, 126.8, 125.7, 123.1, 85.1, 81.7, 80.7, 77.4, 71.3, 71.1, 70.9, 70.5, 69.6, 67.2, 37.4, 28.6. HRMS calcd for $\text{C}_{21}\text{H}_{26}\text{FIN}_2\text{O}_3$ (M^+), 500.0972; found, 500.0959.

(*E*)-(5-Iodo-6-{2-[2-(2-fluoroethoxy)ethoxy]ethoxy}pyridin-3-yl)-2-(4-methylaminophenyl)-ethylene (**13b**)

- [0143] To a stirred solution of **13c** (0.014 g, 0.024 mmol) and 2,6-lutidine (28 μ L, 0.24 mmol) in 2 mL CH_2Cl_2 at 0 $^\circ\text{C}$ was added trimethylsilyl triflate (34 μ L, 0.19 mmol). After 15 min, the reaction solution was submitted to the standard workup with CH_2Cl_2 . The crude product was purified by PTLC to give a light yellow viscous liquid **13b** (0.010 g, 88%). ^1H NMR δ 8.22 (d, 1H, $J = 2.1$ Hz), 8.10 (d, 1H, $J = 2.1$ Hz), 7.34 (d, 2H, $J = 8.6$ Hz), 6.91 (d, 1H, $J = 16.3$ Hz), 6.70 (d, 1H, $J = 16.3$ Hz), 6.60 (d, 2H, $J = 8.6$ Hz), 4.71-4.67 (m, 1H), 4.54-4.43 (m, 3H), 3.94-3.69 (m, 9H), 2.88 (s, 3H). ^{13}C NMR δ 160.5, 149.5, 144.6, 129.8, 129.7, 128.0, 126.3, 118.9, 112.6, 85.1, 81.7, 80.6, 77.4, 71.3, 71.2, 70.9, 70.5, 69.7, 67.2, 30.8. HRMS calcd for $\text{C}_{20}\text{H}_{24}\text{FIN}_2\text{O}_3$ (M^+), 486.0816; found, 486.0818.

EXAMPLE 11

Synthesis of Compound 13e

(*E*)-(5-tri-Butylstannyl-6-{2-[2-(2-fluoroethoxy)ethoxy]ethoxy}pyridin-3-yl)-2-(4-hydroxyphenyl)-ethylene (**12e**)

- [0144] Compound **12e** was prepared from **11e** (0.032 g, 0.075 mmol) as a white viscous liquid (0.040 g, 84% yield). ^1H NMR δ 8.11 (d, 1H, $J = 2.5$ Hz), 7.82 (d, 1H, $J = 2.5$ Hz), 7.39 (d, 2H, $J = 8.6$ Hz), 6.98-6.74 (m, 4H), 5.19 (br s, 1H), 4.71-4.66 (m, 1H), 4.48-4.43 (m, 3H), 3.90-3.62 (m, 8H), 1.70-1.02 (m, 18H), 0.91 (t, 9H, $J = 7.1$ Hz). ^{13}C NMR δ 166.9, 156.0, 145.4, 144.0, 130.1, 127.9, 127.6, 127.4, 124.3, 123.0, 115.9, 85.0, 81.6, 71.0, 70.9, 70.7, 70.5, 70.0, 65.2, 29.5, 29.3, 29.1, 28.0, 27.5, 26.9, 13.9, 13.4, 13.3, 9.9, 6.6, 6.4. HRMS calcd for $\text{C}_{31}\text{H}_{48}\text{FNO}_4\text{Sn}$ (M^+), 637.2589; found, 637.2573.

(*E*)-(5-Iodo-6-{2-[2-(2-fluoroethoxy)ethoxy]ethoxy}pyridin-3-yl)-2-(4-hydroxyphenyl)-ethylene (**13e**)

- [0145] Compound **13e** was prepared from **12e** (0.012 g, 0.019 mmol) as a white solid (0.008 g, 90%). ¹H NMR δ 8.21 (d, 1H, *J* = 2.1 Hz), 8.08 (d, 1H, *J* = 2.1 Hz), 7.33 (d, 2H, *J* = 8.6 Hz), 6.94-6.69 (m, 4H), 4.71-4.67 (m, 1H), 4.53-4.43 (m, 3H), 3.94-3.69 (m, 8H). HRMS calcd for C₁₉H₂₁FINO₄ (M⁺), 473.0499; found, 473.0498.

EXAMPLE 12

Synthesis of Compound 14a

2-Hydroxyethoxy-3-bromo-5-iodopyridine (9b)

- [0146] To a stirring suspension of **I8** (see Example 4 above) (0.906 g, 3.0 mmol), 2-(tert-butyl-dimethyl-silanyloxy)ethanol (0.554 g, 3.15 mmol) and PPh₃ (0.944 g, 3.6 mmol) in 20 mL of THF at -10 °C was added dropwise of diisopropylazodicarboxylate (DIAD) (0.728 g, 3.6 mmol) in 10 mL of THF. The ice-salt bath was removed and the reaction was kept at r.t. 2 h. The reaction solution was concentrated and purified by FC (EtOAc/Hexanes, 5/95) to afford 2-(tert-butyl-dimethyl-silanyloxy)ethoxy-3-bromo-5-iodopyridine, a colorless viscous liquid (0.995 g, 72%). ¹H NMR δ 8.23 (d, 1H, *J* = 2.0 Hz), 8.05 (d, 1H, *J* = 2.0 Hz), 4.42 (t, 2H, *J* = 4.9 Hz), 3.98 (t, 2H, *J* = 4.9 Hz), 0.90 (s, 9H), 0.10 (s, 6H). HRMS calcd for C₁₂H₁₈BrINO₂Si (M-CH₃⁺), 441.9335; found, 441.9312.

(*E*)-[5-Bromo-6-(2-hydroxyethoxy)pyridin-3-yl]-2-(4-dimethylaminophenyl)-ethylene (**14a**)

- [0147] Compound **14a** was prepared from 4-dimethylaminostyrene (0.031 g, 0.212 mmol) and **I10** (0.073 g, 0.212 mmol) as a light yellow solid (0.022 g, 29% yield). ¹H NMR δ 8.07 (d, 1H, *J* = 2.1 Hz), 8.03 (d, 1H, *J* = 2.1 Hz), 7.39 (d, 2H, *J* = 8.8 Hz), 6.94 (d, 1H, *J* = 16.3 Hz), 6.78-6.69 (m, 3H), 4.57-4.52 (m, 2H), 3.99 (t, 2H, *J* = 4.3 Hz), 3.21 (br s, 1H), 3.00 (s, 6H). ¹³C NMR

δ 158.3, 150.4, 143.0, 138.2, 129.9, 129.8, 127.6, 124.9, 118.3, 112.3, 107.5, 69.6, 62.1, 40.3. HRMS calcd for $C_{17}H_{19}BrN_2O_2$ (M^+), 362.063; found, 362.0629.

EXAMPLE 13

Synthesis of Compound 14b

(*E*)-[5-Bromo-6-(2-hydroxyethoxy)pyridin-3-yl]-2-(4-methylaminophenyl)-ethylene (**14b**)

[0148] Compound **14b** was prepared from 4-methylaminostyrene (0.140 g, 1.05 mmol) and **I10** (0.241 g, 0.7 mmol) as a light yellow viscous liquid (0.149 g, 61% yield). 1H NMR δ 8.07 (d, 1H, $J = 2.1$ Hz), 8.03 (d, 1H, $J = 2.1$ Hz), 7.35 (d, 2H, $J = 8.6$ Hz), 6.93 (d, 1H, $J = 16.3$ Hz), 6.74 (d, 1H, $J = 16.3$ Hz), 6.61 (d, 2H, $J = 8.6$ Hz), 4.57-4.52 (m, 2H), 3.99 (br s, 2H), 3.18 (br s, 1H), 2.88 (s, 3H). ^{13}C NMR δ 149.6, 143.3, 138.5, 130.1, 130.0, 128.0, 126.0, 118.6, 112.6, 107.7, 69.8, 62.2, 30.7. HRMS calcd for $C_{17}H_{19}BrN_2O_2$ (M^+), 348.0473; found, 348.0468.

EXAMPLE 14

Synthesis of Compound 14d

(*E*)-[5-Bromo-6-(2-hydroxyethoxy)pyridin-3-yl]-2-(4-acetoxyphenyl)-ethylene (**14d**)

[0149] Compound **14d** was prepared from 4-acetoxystyrene (0.130 g, 0.80 mmol) and **10** (0.244 g, 0.7 mmol) as a white viscous liquid (0.031 g, 12% yield). 1H NMR δ 8.12 (d, 1H, $J = 2.1$ Hz), 8.08 (d, 1H, $J = 2.1$ Hz), 7.50 (d, 2H, $J = 6.8$ Hz), 7.11 (d, 2H, $J = 6.8$ Hz), 6.95 (d, 2H, $J = 5.2$ Hz), 4.58-4.54 (m, 2H), 4.01 (br s, 2H), 3.08 (br s, 1H), 2.32 (s, 3H).

EXAMPLE 15

Synthesis of Compound 14e

(*E*)-[5-Bromo-6-(2-hydroxyethoxy)pyridin-3-yl]-2-(4-hydroxyphenyl)-ethylene (14e)

[0150] In a similar procedure as described in the preparation of 11e, compound 14e was prepared from Acetate 14d (0.031 g, 0.082 mmol) as a white solid (0.020 g, 73%). ¹H NMR (DMSO-d₆) δ 9.60 (br s, 1H), 8.31 (s, 1H), 8.23 (s, 1H), 7.39 (d, 2H, *J* = 8.3 Hz), 7.19 (d, 1H, *J* = 16.8 Hz), 6.94 (d, 1H, *J* = 16.6 Hz), 6.77 (d, 2H, *J* = 8.3 Hz), 4.35 (t, 2H, *J* = 5.1 Hz), 3.73 (t, 2H, *J* = 5.1 Hz). ¹³C NMR (DMSO-d₆) δ 157.9, 157.4, 143.7, 138.1, 129.2, 129.0, 127.8, 119.8, 115.6, 106.7, 68.4, 59.2. HRMS calcd for C₁₅H₁₄BrNO₃ (M⁺), 335.0157; found, 335.0165.

EXAMPLE 16

Synthesis of Compound 15e

(*E*)-[5-tri-Butylstannyl-6-(2-hydroxyethoxy)pyridin-3-yl]-2-(4-hydroxyphenyl)-ethylene (15e)

[0151] Compound 15e was prepared from 14e (0.031 g, 0.092 mmol) as a white viscous liquid (0.012 g, 24% yield). ¹H NMR δ 8.07 (d, 1H, *J* = 2.5 Hz), 7.85 (d, 1H, *J* = 2.5 Hz), 7.39 (d, 2H, *J* = 8.6 Hz), 6.99-6.80 (m, 4H), 5.97 (br s, 1H), 5.01 (br s, 1H), 4.50-4.46 (m, 2H), 3.98-3.94 (m, 2H), 1.69-1.01 (m, 18H), 0.91 (t, 9H, *J* = 7.1 Hz). ¹³C NMR δ 167.2, 156.0, 144.9, 144.7, 144.5, 130.1, 128.0, 127.96, 124.7, 122.8, 116.0, 69.9, 63.4, 29.9, 29.5, 29.3, 29.1, 28.1, 27.5, 26.9, 13.9, 13.6, 13.5, 10.1, 6.7, 6.6. HRMS calcd for C₂₇H₄₁NO₃Sn (M⁺), 547.2108; found, 547.2112.

EXAMPLE 17

Synthesis of Compound 16a

(*E*)-[5-tri-Butylstannyl-6-(2-hydroxyethoxy)pyridin-3-yl]-2-(4-dimethylaminophenyl)-ethylene (**15a**)

[0152] Compound **15a** was prepared from **14a** (0.100 g, 0.275 mmol) as a light yellow oil (0.105 g, 66% yield). ^1H NMR δ 8.10 (d, 1H, $J = 2.5$ Hz), 7.85 (d, 1H, $J = 2.4$ Hz), 7.41 (d, 2H, $J = 8.7$ Hz), 6.95 (d, 1H, $J = 16.3$ Hz), 6.81 (d, 1H, $J = 16.6$ Hz), 6.73 (d, 2H, $J = 8.8$ Hz), 4.48-4.44 (m, 2H), 3.96-3.92 (m, 2H), 2.99 (s, 6H), 1.68-1.01 (m, 18H), 0.92 (t, 9H, $J = 7.2$ Hz). ^{13}C NMR δ 166.6, 150.1, 144.5, 144.1, 128.2, 128.1, 127.4, 125.6, 124.0, 120.5, 112.4, 69.4, 63.0, 40.4, 29.0, 27.2, 13.6, 9.8. HRMS calcd for $\text{C}_{29}\text{H}_{46}\text{N}_2\text{O}_2\text{Sn}$ (M^+), 574.2581; found, 574.2584.

(*E*)-[5-Iodo-6-(2-hydroxyethoxy)pyridin-3-yl]-2-(4-dimethylaminophenyl)-ethylene (**16a**)

[0153] Compound **16a** was prepared from **15a** (0.011 g, 0.019 mmol) as a light yellow solid (0.004 g, 50%). ^1H NMR δ 8.25 (s, 1H), 8.10 (s, 1H), 7.39 (d, 2H, $J = 8.6$ Hz), 6.94 (d, 1H, $J = 16.3$ Hz), 6.76-6.70 (m, 3H), 4.51 (t, 2H, $J = 4.2$ Hz), 4.02-3.95 (m, 2H), 3.19 (s, 1H), 3.00 (s, 6H). HRMS calcd for $\text{C}_{17}\text{H}_{19}\text{IN}_2\text{O}_2$ (M^+), 410.0491; found, 410.0489.

EXAMPLE 18

Synthesis of Compound 16b

(*E*)-[5-tri-Butylstannyl-6-(2-hydroxyethoxy)pyridin-3-yl]-2-(4-methylaminophenyl)-ethylene (**15b**)

[0154] Compound **15b** was prepared from **14b** (0.052 g, 0.15 mmol) as a light yellow oil (0.059 g, 64% yield). ^1H NMR δ 8.08 (d, 1H, $J = 2.5$ Hz), 7.84 (d, 1H, $J = 2.4$ Hz), 7.37 (d, 2H, $J = 8.6$ Hz), 6.93 (d, 1H, $J = 16.3$ Hz), 6.80 (d, 1H, $J = 16.4$ Hz), 6.61 (d, 2H, $J = 8.6$ Hz), 4.48-4.43 (m, 2H), 3.95-3.91 (m, 2H), 2.88 (s, 3H), 1.69-1.01 (m, 18H), 0.91 (t, 9H, $J = 7.1$ Hz). ^{13}C NMR δ

166.9, 149.2, 144.7, 144.3, 128.4, 128.3, 127.8, 126.7, 124.2, 120.7, 112.6, 69.6, 63.2, 30.8, 29.5, 29.3, 29.1, 28.0, 27.5, 26.9, 13.9, 13.5, 13.4, 10.0, 6.6, 6.5. HRMS calcd for $C_{28}H_{44}N_2O_2Sn$ (M^+), 560.2425; found, 560.2419.

(*E*)-[5-Iodo-6-(2-hydroxyethoxy)pyridin-3-yl]-2-(4-methylaminophenyl)-ethylene (**16b**)

[0155] Compound **16b** was prepared from **15b** (0.032 g, 0.057 mmol) as a light yellow solid (0.005 g, 21%). 1H NMR δ 8.24 (d, 1H, $J = 2.1$ Hz), 8.09 (d, 1H, $J = 2.0$ Hz), 7.36 (d, 2H, $J = 8.5$ Hz), 6.92 (d, 1H, $J = 16.3$ Hz), 6.76-6.64 (m, 3H), 4.53-4.49 (m, 2H), 4.01-3.96 (m, 2H), 2.96 (s, 1H), 2.89 (s, 3H). HRMS calcd for $C_{16}H_{17}IN_2O_2$ (M^+), 396.0335; found, 396.0335.

EXAMPLE 19

Radioiodination

[0156] Radioiodinated compounds, [^{125}I]**13a**, **13b**, **16a**, **16b** and **16e**, were prepared via iododestannylation reactions from the corresponding tributyltin precursors according to the method described previously (ref). Hydrogen peroxide (50 μ L, 3% w/v) was added to a mixture of 50 μ L of the tributyltin precursor (4 μ g/ μ L EtOH), 50 μ L of 1N HCl and [^{125}I]NaI (1-5 mCi purchased from Perkin Elmer) in a sealed vial. The reaction was allowed to proceed for 5-10 min at room temperature and terminated by addition of 100 μ L of sat. NaHSO₃. The reaction mixture was extracted with ethyl acetate (3 x 1 mL) after neutralization with 1.5 mL of saturated sodium bicarbonate solution. The combined extracts were evaporated to dryness. The residues were dissolved in 100 μ L of EtOH and purified by HPLC using a reversed-phase column (Phenomenex Gemini C18 analytical column, 4.6 x 250 mm, 5 μ m, CH₃CN/Ammonium formate buffer (1 mM) 8/2 or 7/3; flow rate 0.5-1.0 mL/min). The no-carrier-added products were evaporated to dryness and redissolved in 100% EtOH (1 μ Ci/ μ L) to be stored at -20 °C up to 6 weeks for animal studies and autoradiography studies.

EXAMPLE 20

Binding Studies

[0157] $[^{125}\text{I}]\text{IMPY}$ with 2,200 Ci/mmol specific activity and greater than 95% radiochemical purity was prepared using the standard iododestannylation reaction, and purified by a simplified C-4 mini-column as described previously in Kung, M.-P.; Hou, C.; Zhuang, Z.-P.; Cross, A. J.; Maier, D. L.; Kung, H. F., "Characterization of IMPY as a potential imaging agent for β -amyloid plaques in double transgenic PSAPP mice." *Eur. J. Nucl. Med. Mol. Imaging* 2004, 31, 1136-1145. Competition binding assays were carried out in 12 $\times$ 75 mm borosilicate glass tubes. The reaction mixture contained 50 μl of pooled AD brain homogenates (20-50 μg), 50 μl of $[^{125}\text{I}]\text{IMPY}$ (0.04-0.06 nM diluted in PBS) and 50 μl of inhibitors (10^{-5} - 10^{-10} M diluted serially in PBS containing 0.1 % bovine serum albumin) in a final volume of 1 ml. Nonspecific binding was defined in the presence of 600 nM IMPY in the same assay tubes. The mixture was incubated at 37°C for 2 h and the bound and the free radioactivity were separated by vacuum filtration through Whatman GF/B filters using a Brandel M-24R cell harvester followed by 2 $\times$ 3 ml washes of PBS at room temperature. Filters containing the bound I-125 ligand were counted in a gamma counter (Packard 5000) with 70% counting efficiency. Under the assay conditions, the specifically bound fraction was less than 15% of the total radioactivity. The results of inhibition experiments were subjected to nonlinear regression analysis using equilibrium binding data analysis from which K_i values were calculated. Figures 1 and 6 show K_i values for selected compounds of the present invention.

EXAMPLE 21

Film Autoradiography

[0158] $[^{18}\text{F}]\text{tracers}$: Brain sections from AD subjects were obtained by freezing the brain in powdered dry ice and cut into 20 micrometer-thick

sections. The sections were incubated with [^{18}F]tracers (200,000-250,000 cpm/200 μl) for 1 hr at room temperature. The sections were then dipped in saturated Li_2CO_3 in 40% EtOH (two two-minute washes) and washed with 40% EtOH (one two-minute wash) followed by rinsing with water for 30 sec. After drying, the ^{18}F -labeled sections were exposed to Kodak MR film overnight. The results are depicted in the film in Fig. 2.

[0159] [^{125}I]tracers : To compare different probes using similar sections of human brain tissue, human macro-array brain sections from 6 confirmed AD cases and one control subject were assembled. The presence and localization of plaques on the sections was confirmed with immunohistochemical staining with monoclonal $\text{A}\beta$ antibody 4G8 (Sigma). The sections were incubated with [^{125}I]tracers (200,000-250,000 cpm/200 μL) for 1 h at room temperature. The sections were then dipped in saturated Li_2CO_3 in 40% EtOH (two two-minute washes) and washed with 40% EtOH (one two-minute wash) followed by rinsing with water for 30 sec. After drying, the ^{125}I -labeled sections were exposed to Kodak Biomax MR film overnight.

EXAMPLE 22

Organ distribution in mice

[0160] While under isoflurane anesthesia, 0.15 mL of a 0.1% bovine serum albumin solution containing [^{125}I]tracers (5-10 μCi) was injected directly into the tail vein of ICR mice (22-25 g, male). The mice ($n = 3$ for each time point) were sacrificed by cervical dislocation at designated time-points post injection. The organs of interest were removed and weighed, and the radioactivity was counted with an automatic gamma counter. The percentage dose per organ was calculated by a comparison of the tissue counts to suitably diluted aliquots of the injected material. The total activity of the blood was calculated under the assumption that it is 7% of the total body weight. The % dose/g of samples was calculated by comparing the sample counts with the count of the diluted initial dose.

Table 1. Biodistribution in ICR mice after iv injection of [^{18}F]10 in 5% EtOH in saline (%dose/g, avg of 3 mice $\pm$ SD)

Organ	2 min	30 min	1 hr	2 hr
Blood	6.05 $\pm$ 0.33	2.65 $\pm$ 0.22	3.48 $\pm$ 0.47	2.15 $\pm$ 0.25
Heart	0.75 $\pm$ 0.14	0.17 $\pm$ 0.03	0.22 $\pm$ 0.03	0.18 $\pm$ 0.08
Muscle	7.03 $\pm$ 1.30	8.58 $\pm$ 0.26	10.62 $\pm$ 2.59	5.96 $\pm$ 0.06
Lung	1.07 $\pm$ 0.20	0.30 $\pm$ 0.01	0.35 $\pm$ 0.07	0.20 $\pm$ 0.36
Kidney	6.38 $\pm$ 0.95	1.68 $\pm$ 0.11	1.96 $\pm$ 0.21	0.96 $\pm$ 1.58
Spleen	0.43 $\pm$ 0.11	0.15 $\pm$ 0.05	0.13 $\pm$ 0.03	0.10 $\pm$ 0.17
Liver	24.90 $\pm$ 1.49	9.26 $\pm$ 0.83	10.52 $\pm$ 2.18	6.86 $\pm$ 0.59
Skin	2.52 $\pm$ 0.24	3.99 $\pm$ 0.34	4.42 $\pm$ 0.65	2.91 $\pm$ 0.16
Brain	3.49 $\pm$ 0.58	0.48 $\pm$ 0.07	0.55 $\pm$ 0.10	0.37 $\pm$ 0.08
Bone	5.97 $\pm$ 0.56	2.52 $\pm$ 0.34	4.39 $\pm$ 0.40	6.49 $\pm$ 0.08

Organ	2 min	30 min	1 hr	2 hr
Blood	3.04 $\pm$ 0.29	1.33 $\pm$ 0.16	1.80 $\pm$ 0.16	1.08 $\pm$ 0.06
Heart	6.00 $\pm$ 1056	1.28 $\pm$ 0.16	1.66 $\pm$ 0.24	1.32 $\pm$ 0.33
Muscle	0.62 $\pm$ 0.10	0.75 $\pm$ 0.04	0.95 $\pm$ 0.18	0.52 $\pm$ 0.08
Lung	5.65 $\pm$ 0.89	1.73 $\pm$ 0.17	1.82 $\pm$ 0.31	0.98 $\pm$ 0.08
Kidney	14.19 $\pm$ 2.34	3.77 $\pm$ 0.36	4.29 $\pm$ 0.52	2.19 $\pm$ 0.36
Spleen	4.65 $\pm$ 0.76	1.57 $\pm$ 0.51	1.56 $\pm$ 0.17	1.14 $\pm$ 0.18
Liver	17.00 $\pm$ 0.69	7.21 $\pm$ 0.69	8.13 $\pm$ 1.42	4.96 $\pm$ 0.90
Skin	0.59 $\pm$ 0.03	0.93 $\pm$ 0.13	1.06 $\pm$ 0.09	0.68 $\pm$ 0.16
Brain	7.77 $\pm$ 1.34	1.03 $\pm$ 0.11	1.28 $\pm$ 0.20	0.84 $\pm$ 0.08
Bone	1.49 $\pm$ 0.08	0.63 $\pm$ 0.12	1.13 $\pm$ 0.01	1.64 $\pm$ 0.50

Table 2. Biodistribution in ICR mice after iv injections of [125 I]-labeled tracers
(%dose/g, avg of 3 mice $\pm$ SD)

[125 I]13a (logP = 2.59)								
Organ	2 min		30 min		1 hr		2 hr	
Blood	2.70	$\pm$ 0.58	2.05	$\pm$ 0.18	1.65	$\pm$ 0.45	1.45	$\pm$ 0.41
Heart	12.76	$\pm$ 1.24	1.63	$\pm$ 0.03	0.97	$\pm$ 0.16	0.73	$\pm$ 0.17
Muscle	0.90	$\pm$ 0.20	1.00	$\pm$ 0.08	0.59	$\pm$ 0.13	0.53	$\pm$ 0.08
Lung	10.08	$\pm$ 2.15	2.50	$\pm$ 0.14	1.62	$\pm$ 0.46	1.33	$\pm$ 0.39
Kidney	16.62	$\pm$ 1.96	3.32	$\pm$ 0.11	2.30	$\pm$ 0.54	1.71	$\pm$ 0.24
Spleen	4.47	$\pm$ 1.28	1.42	$\pm$ 0.05	0.99	$\pm$ 0.47	0.79	$\pm$ 0.27
Liver	22.15	$\pm$ 4.34	9.54	$\pm$ 1.30	5.34	$\pm$ 2.22	5.62	$\pm$ 1.31
Skin	0.54	$\pm$ 0.05	1.47	$\pm$ 0.26	1.59	$\pm$ 0.68	1.23	$\pm$ 0.41
Brain	4.03	$\pm$ 0.43	1.93	$\pm$ 0.18	0.68	$\pm$ 0.17	0.26	$\pm$ 0.04
Thyroid	3.89	$\pm$ 0.67	16.23	$\pm$ 11.75	24.19	$\pm$ 8.26	60.76	$\pm$ 6.09

[125 I]13b (log P = 2.54)								
Organ	2 min		30 min ⁺		1 hr		2 hr	
Blood	4.37	$\pm$ 1.07	3.83	$\pm$ 1.11	2.88	$\pm$ 0.28	2.21	$\pm$ 0.73
Heart	9.85	$\pm$ 1.78	2.54	$\pm$ 0.37	1.75	$\pm$ 0.26	1.22	$\pm$ 0.28
Muscle	1.04	$\pm$ 0.25	1.11	$\pm$ 0.34	0.85	$\pm$ 0.06	0.44	$\pm$ 0.19
Lung	6.85	$\pm$ 0.27	3.01	$\pm$ 0.96	2.37	$\pm$ 0.29	1.85	$\pm$ 0.74
Kidney	9.03	$\pm$ 6.81	3.40	$\pm$ 0.76	2.81	$\pm$ 0.70	1.86	$\pm$ 0.36
Spleen	4.41	$\pm$ 1.05	2.49	$\pm$ 0.75	1.75	$\pm$ 0.33	1.27	$\pm$ 0.24
Liver	26.24	$\pm$ 4.47	11.47	$\pm$ 2.10	7.70	$\pm$ 1.22	6.25	$\pm$ 1.79
Skin	1.48	$\pm$ 0.07	2.95	$\pm$ 0.81	2.46	$\pm$ 0.16	1.32	$\pm$ 0.41
Brain	6.22	$\pm$ 1.01	1.23	$\pm$ 0.13	0.62	$\pm$ 0.17	0.26	$\pm$ 0.01
Thyroid	5.74	$\pm$ 0.42	24.09	$\pm$ 27.44	38.09	$\pm$ 6.37	215.05	$\pm$ 74.59

[125 I]16a (log P = 2.64)								
Organ	2 min		30 min		1 hr ⁺		2 hr	
Blood	2.71	$\pm$ 0.07	2.24	$\pm$ 0.38	2.18	$\pm$ 0.66	1.01	$\pm$ 0.02
Heart	10.24	$\pm$ 0.45	1.93	$\pm$ 0.27	1.12	$\pm$ 0.02	0.62	$\pm$ 0.12
Muscle	0.71	$\pm$ 0.46	1.05	$\pm$ 0.20	0.55	$\pm$ 0.03	0.22	$\pm$ 0.04
Lung	9.41	$\pm$ 0.56	3.02	$\pm$ 0.38	1.98	$\pm$ 0.21	1.00	$\pm$ 0.15
Kidney	14.25	$\pm$ 1.98	4.19	$\pm$ 0.45	2.49	$\pm$ 0.33	1.48	$\pm$ 0.20
Spleen	4.40	$\pm$ 1.89	1.94	$\pm$ 0.19	1.32	$\pm$ 0.10	0.80	$\pm$ 0.11
Liver	19.12	$\pm$ 2.68	12.38	$\pm$ 1.29	6.22	$\pm$ 0.96	4.87	$\pm$ 0.46
Skin	0.46	$\pm$ 0.13	1.18	$\pm$ 0.26	1.16	$\pm$ 0.00	0.40	$\pm$ 0.05
Brain	5.43	$\pm$ 0.85	3.56	$\pm$ 0.32	1.32	$\pm$ 0.00	0.46	$\pm$ 0.05
Thyroid	4.15	$\pm$ 0.43	11.21	$\pm$ 7.88	59.13	$\pm$ 6.26	24.81	$\pm$ 0.62

[¹²⁵I]16b (log P = 2.20)

Organ	2 min		30 min		1 hr		2 hr	
Blood	4.14	± 0.41	3.08	± 0.35	1.81	± 0.56	1.96	± 0.14
Heart	7.16	± 1.16	1.50	± 0.18	0.88	± 0.30	0.76	± 0.03
Muscle	1.15	± 0.38	0.91	± 0.06	0.42	± 0.08	0.38	± 0.02
Lung	7.43	± 1.21	2.67	± 0.46	1.76	± 0.32	1.58	± 0.10
Kidney	11.51	± 1.48	3.73	± 0.75	2.16	± 0.08	1.53	± 0.20
Spleen	4.08	± 0.68	1.34	± 0.29	0.87	± 0.37	1.08	± 0.15
Liver	20.84	± 2.38	12.57	± 3.03	5.62	± 0.68	3.41	± 0.20
Skin	0.95	± 0.09	1.86	± 0.50	1.29	± 0.51	1.43	± 0.10
Brain	8.04	± 0.82	0.88	± 0.30	0.26	± 0.03	0.15	± 0.02
Thyroid	6.31	± 1.59	17.23	± 14.23	36.69	± 37.17	99.88	± 69.45


[¹²⁵I]16e (log P = 1.98)

Organ	2 min		30 min		1 hr		2 hr	
Blood	10.09	± 1.12	3.92	± 0.07	1.29	± 0.05	1.56	± 0.04
Heart	6.66	± 0.31	1.35	± 0.16	0.65	± 0.21	0.51	± 0.09
Muscle	1.01	± 0.34	0.59	± 0.05	0.21	± 0.02	0.12	± 0.01
Lung	14.22	± 0.92	3.10	± 0.05	1.34	± 0.11	1.02	± 0.01
Kidney	20.40	± 2.20	10.03	± 2.12	2.94	± 0.17	2.50	± 1.32
Spleen	4.20	± 0.31	1.28	± 0.44	0.50	± 0.03	0.50	± 0.06
Liver	18.27	± 1.29	5.15	± 0.61	2.38	± 0.58	2.63	± 1.30
Skin	0.64	± 0.20	1.36	± 0.07	0.62	± 0.01	0.37	± 0.08
Brain	0.99	± 0.24	0.26	± 0.03	0.09	± 0.01	0.06	± 0.01
Thyroid	4.38	± 0.46	3.99	± 3.56	13.02	± 8.11	16.02	± 11.52

[0161] It will be understood to those of ordinary skill in the art that the same can be performed within a wide and equivalent range of conditions, formulations, and other parameters without affecting the scope of the invention or any embodiment thereof. All patents, patent applications, and publications cited herein are fully incorporated by reference herein in their entirety.

What Is Claimed Is:

1. A compound of Formula I,



or a pharmaceutically acceptable salt thereof; wherein,

n is an integer from one to six;


at least one, no more than three, of A_1 , A_2 , A_3 , A_4 and A_5 is N, the others are $-CH$, or $-CR^2$ as permitted;

R^1 is selected from the group consisting of:


- a. $-(CH_2)_pNR'R''$, wherein R' and R'' are independently hydrogen, C_{1-4} alkyl, hydroxy(C_{1-4})alkyl or halo(C_{1-4})alkyl, and p is an integer from 0 to 5;
- b. hydroxy,
- c. C_{1-4} alkoxy,
- d. hydroxy(C_{1-4})alkyl,
- e. halogen,
- f. cyano,
- g. hydrogen,
- h. nitro,
- i. (C_1-C_4) alkyl,
- j. halo(C_1-C_4)alkyl,
- k. formyl,
- l. $-NHCO(C_{1-4}$ alkyl), and

m. $-\text{OCO}(\text{C}_{1-4} \text{ alkyl})$;

R^2 is selected from the group consisting of:



wherein, q is an integer from 1 to 10, R^x and R^y are hydrogen, hydroxy or C_{1-4} alkyl; t is 0, 1, 2 or 3; Z is selected from the group consisting of hydroxy, halogen, halogen substituted benzoyloxy, halogen substituted benzyloxy, halogen substituted phenyl(C_{1-4})alkyl, halogen substituted aryloxy, and a halogen substituted C_{6-10} aryl; and R^{30} , R^{31} , R^{32} and R^{33} are in each instance independently selected from the group consisting of hydrogen, hydroxy, C_{1-4} alkoxy, C_{1-4} alkyl, and hydroxy(C_{1-4})alkyl;



wherein R^x and R^y are hydrogen, hydroxy or C_{1-4} alkyl; t is 0, 1, 2 or 3;

Y is selected from the group consisting of halogen, halogen substituted benzoyloxy, halogen substituted phenyl(C_{1-4})alkyl, halogen substituted aryloxy, and halogen substituted C_{6-10} aryl;

U is selected from the group consisting of hydrogen, hydroxy, halogen, halogen substituted benzoyloxy, halogen

substituted phenyl(C₁₋₄)alkyl, halogen substituted aryloxy, and halogen substituted C₆₋₁₀ aryl; and

R³⁴, R³⁵, R³⁶, R³⁷, R³⁸, R³⁹ and R⁴⁰ are in each instance independently selected from the group consisting of hydrogen, halogen, hydroxy, C₁₋₄ alkoxy, C₁₋₄ alkyl, and hydroxy(C₁₋₄)alkyl;

iii. NR'R'', wherein at least one of R' and R'' is (CH₂)_dX, where X is halogen, preferably F or ¹⁸F, and d is an integer from 1 to 4; the other of R' and R'' is selected from the group consisting of hydrogen, C₁₋₄ alkyl, halo(C₁₋₄)alkyl, and hydroxy(C₁₋₄)alkyl;

iv. NR'R''-(C₁₋₄)alkyl, wherein at least one of R' and R'' is (CH₂)_dX, where X is halogen, preferably F or ¹⁸F, and d is an integer from 1 to 4; the other of R' and R'' is selected from the group consisting of hydrogen, C₁₋₄ alkyl, halo(C₁₋₄)alkyl, and hydroxy(C₁₋₄)alkyl;


v. halo(C₁₋₄)alkyl; and

vi. an ether (R-O-R) having the following structure:
[halo(C₁₋₄)alkyl-O-(C₁₋₄)alkyl]-;

and

R⁷ and R⁸ are in each instance independently selected from the group consisting of hydrogen, hydroxy, amino, methylamino, dimethylamino, C₁₋₄ alkoxy, C₁₋₄ alkyl, and hydroxy(C₁₋₄)alkyl.

2. The compound of claim 1 having the following structure:



I


or a pharmaceutically acceptable salt thereof; wherein,

n is an integer from one to six;


at least one, no more than three, of A_1 , A_2 , A_3 , A_4 and A_5 is N, the others are $-CH_2$ or $-CR^2$ as permitted;

R^1 is hydroxy or $NR^aR^b(CH_2)_p-$, wherein p is an integer from 0 to 5, and R^a and R^b are independently hydrogen, C_{1-4} alkyl or $(CH_2)_dX$, where X is halogen, and d is an integer from 1 to 4,

R^2 is selected from the group consisting of:



wherein q is an integer from 1 to 10; Z is selected from the group consisting of halogen, halogen substituted benzoyloxy, halogen substituted benzyloxy, halogen substituted phenyl(C_{1-4})alkyl, halogen substituted aryloxy, and a halogen substituted C_{6-10} aryl; and R^{30} , R^{31} , R^{32} and R^{33} are in each instance independently selected from the group consisting of hydrogen, hydroxy, C_{1-4} alkoxy, C_{1-4} alkyl, and hydroxy(C_{1-4})alkyl;



wherein Y is selected from the group consisting of halogen, halogen substituted benzoyloxy, halogen substituted phenyl(C₁₋₄)alkyl, halogen substituted aryloxy, and halogen substituted C₆₋₁₀ aryl;

U is selected from the group consisting of hydrogen, hydroxy, halogen, halogen substituted benzoyloxy, halogen substituted phenyl(C₁₋₄)alkyl, halogen substituted aryloxy, and halogen substituted C₆₋₁₀ aryl; and

R³⁴, R³⁵, R³⁶, R³⁷, R³⁸, R³⁹ and R⁴⁰ are in each instance independently selected from the group consisting of hydrogen, halogen, hydroxy, C₁₋₄ alkoxy, C₁₋₄ alkyl, and hydroxy(C₁₋₄)alkyl;

iii. NR'R'', wherein at least one of R' and R'' is (CH₂)_dX, where X is halogen, preferably F or ¹⁸F, and d is an integer from 1 to 4; the other of R' and R'' is selected from the group consisting of hydrogen, C₁₋₄ alkyl, halo(C₁₋₄)alkyl, and hydroxy(C₁₋₄)alkyl;

iv. NR'R''-(C₁₋₄)alkyl, wherein at least one of R' and R'' is (CH₂)_dX, where X is halogen, preferably F or ¹⁸F, and d is an integer from 1 to 4; the other of R' and R'' is selected from the group consisting of hydrogen, C₁₋₄ alkyl, halo(C₁₋₄)alkyl, and hydroxy(C₁₋₄)alkyl;


v. halo(C₁₋₄)alkyl; and

vi. an ether (R-O-R) having the following structure:
[halo(C₁₋₄)alkyl-O-(C₁₋₄)alkyl]-;


and

R^7 and R^8 are in each instance independently selected from the group consisting of hydrogen, hydroxy, amino, methylamino, dimethylamino, C_{1-4} alkoxy, C_{1-4} alkyl, and hydroxy(C_{1-4})alkyl.

3. The compound of claim 1 wherein n is one.
4. The compound of claim 1, wherein said halogen, in each instance, is a radiolabeled halogen.
5. The compound of claim 1, wherein said halogen, in each instance, is selected from the group consisting of I, ^{123}I , ^{125}I , ^{131}I , Br, ^{76}Br , ^{77}Br , F or ^{18}F .
6. The compound of claim 5, wherein R^1 is $\text{NR}^a\text{R}^b(\text{CH}_2)_p$ -, wherein R^a and R^b are independently hydrogen or C_{1-4} alkyl and p is 0.
7. The compound of claim 3, wherein R^2 is




8. The compound of claim 7, wherein q is an integer from 1 to 5.
9. The compound of claim 7, wherein R^7 and R^8 are each hydrogen.
10. The compound of claim 8, wherein R^{30} , R^{31} , R^{32} and R^{33} are, in each instance, hydrogen.
11. The compound of claim 10, wherein A_4 is N, having the following formula:




wherein, R^a and R^b are independently hydrogen or C_{1-4} alkyl, Z is I, ^{123}I , ^{125}I , ^{131}I , Br, ^{76}Br , ^{77}Br , F or ^{18}F .


12. The compound of claim 10, having the following formula:



13. The compound of claim 10, having the formula:




14. The compound of claim 10, having the following formula:




wherein Z is I, ^{123}I , ^{125}I , ^{131}I , Br, ^{76}Br , ^{77}Br , F or ^{18}F .

15. The compound of claim 10, having the following formula:




16. The compound of claim 3, wherein R^2 is



wherein Z is hydroxy.


17. The compound of claim 16, wherein q is an integer from 1 to 5.

18. The compound of claim 17, wherein R^7 and R^8 are each hydrogen.
19. The compound of claim 18, wherein R^{30} , R^{31} , R^{32} and R^{33} are, in each instance, hydrogen, and q is 1, 2 or 3.
20. The compound of claim 1, wherein R^2 is




wherein U is hydroxy.

21. The compound of claim 20, wherein R^{34} , R^{35} , R^{36} and R^{37} , R^{38} , R^{39} and R^{40} are, in each instance, hydrogen.
22. The compound of claim 20, having the following structure:




wherein R^a and R^b are independently hydrogen or C_{1-4} alkyl.

23. The compound of claim 20, having the following structure:



24. A compound having the following structure:



or a pharmaceutically acceptable salt thereof; wherein,


at least one, no more than three, of A₁, A₂, A₃, A₄ and A₅ is N, the others are -CH, or -CR² as permitted;

n is an integer from 1 to 6;


R¹ is selected from the group consisting of:

- a. $-(CH_2)_pNR'R''$, wherein R' and R'' are independently hydrogen, C₁₋₄ alkyl, hydroxy(C₁₋₄)alkyl or halo(C₁₋₄)alkyl, and p is an integer from 0 to 5;
- b. hydroxy,
- c. C₁₋₄ alkoxy,
- d. hydroxy(C₁₋₄)alkyl,
- e. halogen,
- f. cyano,
- g. hydrogen,
- h. nitro,
- i. (C₁₋₄)alkyl,
- j. halo(C₁₋₄)alkyl,
- k. formyl,
- l. -NHCO(C₁₋₄ alkyl), and
- m. -OCO(C₁₋₄ alkyl);

R² is selected from the group consisting of:



wherein, q is an integer from 1 to 10, R^x and R^y are hydrogen, hydroxy or C_{1-4} alkyl; t is 0, 1, 2 or 3; Z is selected from the group consisting of hydroxy, halogen, halogen substituted benzoyloxy, halogen substituted benzyloxy, halogen substituted phenyl(C_{1-4})alkyl, halogen substituted aryloxy, and a halogen substituted C_{6-10} aryl; and R^{30} , R^{31} , R^{32} and R^{33} are in each instance independently selected from the group consisting of hydrogen, hydroxy, C_{1-4} alkoxy, C_{1-4} alkyl, and hydroxy(C_{1-4})alkyl;



wherein R^x and R^y are hydrogen, hydroxy or C_{1-4} alkyl; t is 0, 1, 2 or 3;

Y is selected from the group consisting of halogen, halogen substituted benzoyloxy, halogen substituted phenyl(C_{1-4})alkyl, halogen substituted aryloxy, and halogen substituted C_{6-10} aryl;

U is selected from the group consisting of hydrogen, hydroxy, halogen, halogen substituted benzoyloxy, halogen substituted phenyl(C_{1-4})alkyl, halogen substituted aryloxy, and halogen substituted C_{6-10} aryl; and

R^{34} , R^{35} , R^{36} , R^{37} , R^{38} , R^{39} and R^{40} are in each instance independently selected from the group consisting of hydrogen, halogen, hydroxy, C_{1-4} alkoxy, C_{1-4} alkyl, and hydroxy(C_{1-4})alkyl;

iii. $NR'R''$, wherein at least one of R' and R'' is $(CH_2)_dX$, where X is halogen, preferably F or ^{18}F , and d is an integer from 1 to 4; the other of R' and R'' is selected from the group consisting of hydrogen, C_{1-4} alkyl, halo(C_{1-4})alkyl, and hydroxy(C_{1-4})alkyl;

iv. $NR'R''-(C_{1-4})alkyl$, wherein at least one of R' and R'' is $(CH_2)_dX$, where X is halogen, preferably F or ^{18}F , and d is an integer from 1 to 4; the other of R' and R'' is selected from the group consisting of hydrogen, C_{1-4} alkyl, halo(C_{1-4})alkyl, and hydroxy(C_{1-4})alkyl;


v. halo(C_{1-4})alkyl; and

vi. an ether ($R-O-R$) having the following structure:
[halo(C_{1-4})alkyl- O -(C_{1-4})alkyl]-;

and

R^7 and R^8 are in each instance independently selected from the group consisting of hydrogen, hydroxy, amino, methylamino, dimethylamino, C_{1-4} alkoxy, C_{1-4} alkyl, and hydroxy(C_{1-4})alkyl.

25. A compound of claim 24, having the following structure:




or a pharmaceutically acceptable salt thereof; wherein,

at least one, no more than three, of A_1 , A_2 , A_3 , A_4 and A_5 is N, the others are -CH, or -CR² as permitted;


n is an integer from 1 to 6;

R^1 is hydroxy or $NR^aR^b(CH_2)_p-$, wherein p is an integer from 0 to 5, and R^a and R^b are independently hydrogen, C₁₋₄ alkyl or $(CH_2)_dX$, where X is halogen, and d is an integer from 1 to 6,

R^2 is selected from the group consisting of:




wherein q is an integer from 2 to 10; Z is -Ch;



wherein Z is -Ch, R^{30} , R^{31} , R^{32} and R^{33} are as described

above, and



wherein Y is -Ch;

U is selected from the group consisting of hydrogen, hydroxy, halogen, halogen substituted benzoyloxy, halogen substituted phenyl(C₁₋₄)alkyl, halogen substituted aryloxy, and halogen substituted C₆₋₁₀ aryl; and

R³⁴, R³⁵, R³⁶, R³⁷, R³⁸, R³⁹ and R⁴⁰ are in each instance independently selected from the group consisting of hydrogen, halogen, hydroxy, C₁₋₄ alkoxy, C₁₋₄ alkyl, and hydroxy(C₁₋₄)alkyl;

iv. -(CH₂)_w-O-Ch, wherein w is an integer from 1 to 10;


v. -Ch; and

vi. -(CH₂)_w-Ch, wherein w is an integer from 1 to 10;

wherein, -Ch is an N₂S₂-type chelating moiety, and

R⁷ and R⁸ are in each instance independently selected from the group consisting of hydrogen, hydroxy, amino, methylamino, dimethylamino, C₁₋₄ alkoxy, C₁₋₄ alkyl, and hydroxy(C₁₋₄)alkyl.


26. The compound of claim 24 wherein, said N₂S₂ moiety has the following structure:




wherein R^P is hydrogen or a sulfhydryl protecting group, and R⁹ R¹⁰, R¹¹, R¹², R¹³, R¹⁴, R¹⁵, R¹⁶, R⁴³ and R⁴⁴ are in each instance independently

selected from the group consisting of hydrogen, hydroxy, amino, methylamino, dimethylamino, C₁₋₄ alkoxy, C₁₋₄ alkyl, and hydroxy(C₁₋₄)alkyl.

27. A radiometal complex of a compound of claim 26, wherein said -Ch has the following structure:



28. A compound of Formula II, having the following structure:



or a pharmaceutically acceptable salt thereof, wherein:

at least one, no more than three, of A₁, A₂, A₃, A₄ and A₅ is N,
the others are -CH, -CR³ or -CR⁴ as permitted;

R^5 and $R^{5'}$ are independently hydrogen or C_{1-4} alkyl;

R^1 and R^2 , in each instance, is independently selected from the group consisting of hydrogen, hydroxy, halogen, C_{1-4} alkyl, cyano, carboxy(C_{1-5})alkyl, trifluoromethyl, nitro, halo(C_{1-4})alkyl, formyl and $NR^6R^7(CH_2)_p$, wherein p is an integer from 0 to 5, and R^6 and R^7 are independently hydrogen, C_{1-4} alkyl or $(CH_2)_dX$, where X is halogen, and d is an integer from 1 to 4;

R^3 is selected from the group consisting of hydrogen, halogen, C_{1-4} alkyl, cyano, carboxy(C_{1-5})alkyl, trifluoromethyl, nitro, halo(C_{1-4})alkyl, formyl, $NR^6R^7(CH_2)_p$ -, wherein p is an integer from 0 to 5, and R^6 and R^7 are independently hydrogen, C_{1-4} alkyl or $(CH_2)_dX$, where X is halogen, and d is an integer from 1 to 4, ^{18}F fluoromethyl, ^{18}F fluoroethyl, ^{18}F fluoropropyl and $Sn(alkyl)_3$;

R^4 is selected from the group consisting of:

- a. C_{1-4} alkylthio,
- b. C_{1-4} alkylsulfonyl,
- c. hydroxy,
- d. C_{1-4} alkoxy,
- e. $NR^6R^7(CH_2)_p$ -, wherein p is an integer from 0 to 5, and R^6 and R^7 are independently hydrogen, C_{1-4} alkyl or $(CH_2)_dX$, where X is halogen, and d is an integer from 1 to 4,
- f. phenyl(C_{1-4})alkyl,
- g. C_{6-10} aryl,
- h. heteroaryl,
- i. heterocycle,
- j. heterocycle(C_{1-4})alkyl, and
- k. C_{3-6} cycloalkyl,

wherein said phenyl(C_{1-4})alkyl, C_{6-10} aryl, heteroaryl, heterocycle, heterocycle(C_{1-4})alkyl or C_{3-6} cycloalkyl is substituted with one of the following: C_{1-4} alkylthio, C_{1-4} alkyl sulfonyl, methoxy, hydroxy, dimethylamino or methylamino;

and,

X' is hydrogen, ¹⁸Fluoromethyl, ¹⁸Fluoroethyl or ¹⁸Fluoropropyl, ¹²⁵I, ¹²³I, ¹³¹I, ¹⁸F, ⁷⁶Br, ⁷⁷Br or Sn(alkyl)₃,

provided that one of R¹, R², R³ and R⁴ is NR⁶R⁷(CH₂)_p- or one of R¹, R² and R⁴ is hydroxy.

29. A compound of claim 28 wherein,

at least one, no more than three, of A₁, A₂, A₃, A₄ and A₅ is N, the others are -CH, -CR³ or -CR⁴ as permitted;

R⁵ and R^{5'} are independently hydrogen or C₁₋₄ alkyl;

R¹ and R², in each instance, is independently selected from the group consisting of hydrogen, halogen, C₁₋₄ alkyl, cyano, carboxy(C₁₋₅)alkyl, trifluoromethyl, nitro, halo(C₁₋₄)alkyl, formyl and NR⁶R⁷(CH₂)_p-, wherein p is an integer from 0 to 5, and R⁶ and R⁷ are independently hydrogen, C₁₋₄ alkyl or (CH₂)_dX, where X is halogen, and d is an integer from 1 to 4;

R³ is selected from the group consisting of hydrogen, halogen, C₁₋₄ alkyl, cyano, carboxy(C₁₋₅)alkyl, trifluoromethyl, nitro, halo(C₁₋₄)alkyl, formyl, NR⁶R⁷(CH₂)_p-, wherein p is an integer from 0 to 5, and R⁶ and R⁷ are independently hydrogen, C₁₋₄ alkyl or (CH₂)_dX, where X is halogen, and d is an integer from 1 to 4, ¹⁸Fluoromethyl, ¹⁸Fluoroethyl, ¹⁸Fluoropropyl and Sn(alkyl)₃;

R⁴ is selected from the group consisting of:

- a. C₁₋₄ alkylthio,
- b. C₁₋₄ alkylsulfonyl,
- c. hydroxy,

- d. C₁₋₄ alkoxy,
- e. NR⁶R⁷(CH₂)_p-, wherein p is an integer from 0 to 5, and R⁶ and R⁷ are independently hydrogen, C₁₋₄ alkyl or (CH₂)_dX, where X is halogen, and d is an integer from 1 to 4,
- f. phenyl(C₁₋₄)alkyl,
- g. C₆₋₁₀ aryl,
- h. heteroaryl,
- i. heterocycle,
- j. heterocycle(C₁₋₄)alkyl, and
- k. C₃₋₆ cycloalkyl,

wherein said phenyl(C₁₋₄)alkyl, C₆₋₁₀ aryl, heteroaryl, heterocycle, heterocycle(C₁₋₄)alkyl or C₃₋₆ cycloalkyl is substituted with one of the following: C₁₋₄ alkylthio, C₁₋₄ alkyl sulfonyl, methoxy, hydroxy, dimethylamino or methylamino;

and,

X' is hydrogen, ¹⁸F fluoromethyl, ¹⁸F fluoroethyl or ¹⁸F fluoropropyl, ¹²⁵I, ¹²³I, ¹³¹I, ¹⁸F, ⁷⁶Br, ⁷⁷Br or Sn(alkyl)₃,

provided that one of R¹, R², R³ and R⁴ is NR⁶R⁷(CH₂)_p-.

30. A compound of claim 28 wherein,

R⁵ is hydrogen or methyl;

R³ is hydrogen; and


R¹ and R² are hydrogen or C₁₋₄ alkyl.

31. A compound of claim 28, wherein

R^1 is hydrogen, and

R^4 is NR^6R^7 , wherein R^6 and R^7 are hydrogen or C_{1-4} alkyl.


32. A compound of claim 30, wherein X' is ^{123}I .
33. A compound of claim 30, wherein X' is ^{18}F fluoromethyl, ^{18}F fluoroethyl or ^{18}F fluoropropyl.
34. A compound of claim 30, wherein R^3 is ^{18}F fluoromethyl, ^{18}F fluoroethyl, ^{18}F fluoropropyl, ^{125}I , ^{123}I , ^{131}I , ^{18}F , ^{76}Br or ^{77}Br .
35. A composition comprising a compound of Formula I, Ia, II or III, wherein said compound contains a ^{11}C isotope, wherein said isotope is present in said composition in an amount above natural abundance, and said compound containing said ^{11}C isotope has a specific activity above background level.
36. A compound of Formula Ia, having the following general structure:



wherein at least one, no more than three, of A_1 , A_2 , A_3 , A_4 and A_5 is N, the others are $-CH$; q is an integer from 1 to 10; R' and R'' are each independently hydrogen or C_{1-4} alkyl and X is a radiolabeled halogen or $-Ch$ moiety.

37. The compound of claim 36, wherein one of R' and R'' is hydrogen, the other is C_{1-4} alkyl.

38. A compound of Formula III, having the following general structure:



III

or a pharmaceutically acceptable salt thereof, wherein,

at least one, no more than three, of A₁, A₂, A₃, A₄ and A₅ is N, the others are -CH-, -CR² or -CR³ as permitted;


n is an integer from 1 to 6;

R¹ is selected from the group consisting of:


- a. -(CH₂)_pNR'R'', wherein R' and R'' are independently hydrogen, C₁₋₄ alkyl, hydroxy(C₁₋₄)alkyl or halo(C₁₋₄)alkyl, and p is an integer from 0 to 5;
- b. hydroxy,
- c. C₁₋₄ alkoxy,
- d. hydroxy(C₁₋₄)alkyl,
- e. halogen,
- f. cyano,
- g. hydrogen,
- h. nitro,
- i. (C₁₋₄)alkyl,
- j. halo(C₁₋₄)alkyl,
- k. formyl,
- l. -NHCO(C₁₋₄ alkyl), and
- m. -OCO(C₁₋₄ alkyl);

R³ is radiohalogen, radiohalo(C₁₋₄)alkyl or -Sn(C₁₋₄ alkyl)₃; and

R² is selected from the group consisting of:




wherein q is an integer from 1 to 10, R^x and R^y are hydrogen, hydroxy or C_{1-4} alkyl; t is 0, 1, 2 or 3; Z is selected from the group consisting of hydrogen, hydroxy, halogen, C_{1-4} alkoxy, C_{1-4} alkyl, and hydroxy(C_{1-4})alkyl, and R^{30} , R^{31} , R^{32} and R^{33} are in each instance independently selected from the group consisting of hydrogen, hydroxy, C_{1-4} alkoxy, C_{1-4} alkyl, and hydroxy(C_{1-4})alkyl;



wherein Y is selected from the group consisting of halogen, halogen substituted benzoyloxy, halogen substituted phenyl(C_{1-4})alkyl, halogen substituted aryloxy, and halogen substituted C_{6-10} aryl;

U is selected from the group consisting of hydrogen, hydroxy, halogen, halogen substituted benzoyloxy, halogen substituted phenyl(C_{1-4})alkyl, halogen substituted aryloxy, and halogen substituted C_{6-10} aryl; and

R^{34} , R^{35} , R^{36} , R^{37} , R^{38} , R^{39} and R^{40} are in each instance independently selected from the group consisting of hydrogen, halogen, hydroxy, C_{1-4} alkoxy, C_{1-4} alkyl, and hydroxy(C_{1-4})alkyl;



wherein R^x and R^y are hydrogen, hydroxy or C_{1-4} alkyl; t is 0, 1, 2 or 3; and Y , U , R^{34} , R^{35} , R^{36} , R^{37} , R^{38} , R^{39} and R^{40} are as described above;

iii. $NR'R''$, wherein at least one of R' and R'' is $(CH_2)_dX$, where X is halogen, preferably F or ^{18}F , and d is an integer from 1 to 4; the other of R' and R'' is selected from the group consisting of hydrogen, C_{1-4} alkyl, halo(C_{1-4})alkyl, and hydroxy(C_{1-4})alkyl;

iv. $NR'R''-(C_{1-4})$ alkyl, wherein at least one of R' and R'' is $(CH_2)_dX$, where X is halogen, preferably F or ^{18}F , and d is an integer from 1 to 4; the other of R' and R'' is selected from the group consisting of hydrogen, C_{1-4} alkyl, halo(C_{1-4})alkyl, and hydroxy(C_{1-4})alkyl;

v. halo(C_{1-4})alkyl; and


vi. an ether ($R-O-R$) having the following structure:
[halo(C_{1-4})alkyl-O-(C_{1-4})alkyl]-;

and

R^7 and R^8 are in each instance independently selected from the group consisting of hydrogen, hydroxy, amino, methylamino, dimethylamino, C_{1-4} alkoxy, C_{1-4} alkyl, and hydroxy(C_{1-4})alkyl.


39. The compound of claim 38 wherein, R^3 is selected from the group consisting of ^{125}I , ^{123}I , ^{131}I , ^{18}F , $^{18}F(C_{1-4}$ alkyl), ^{76}Br , ^{77}Br and $-Sn(C_{1-4}$ alkyl) $_3$.

40. The compound of claim 38 wherein, R^1 is selected from the group consisting of hydroxy, $-\text{OCO}(\text{C}_{1-4}\text{alkyl})$ and $-(\text{CH}_2)_p\text{NR}'\text{R}''$, wherein R' and R'' are independently hydrogen, C_{1-4} alkyl, hydroxy(C_{1-4})alkyl or halo(C_{1-4})alkyl, and p is an integer from 0 to 5.
41. The compound of claim 38 wherein, R^1 is hydroxy or $-(\text{CH}_2)_p\text{NR}'\text{R}''$, wherein R' and R'' are independently hydrogen or C_{1-4} alkyl, and p is 0.
42. The compound of claim 38 wherein, A^4 is N and A^1 , A^2 , A^3 and A^5 are each C.
43. The compound of claim 38 wherein, R^2 is



wherein q is an integer from 1 to 4; Z is halogen or hydroxy, and R^{30} , R^{31} , R^{32} and R^{33} are in each instance independently selected from the group consisting of hydrogen, hydroxy, C_{1-4} alkoxy, C_{1-4} alkyl, and hydroxy(C_{1-4})alkyl.

44. The compound of claim 38 wherein, Z is fluoro and R^{30} , R^{31} , R^{32} and R^{33} are in each instance hydrogen.
45. The compound of claim 38 wherein, R^2 is






wherein Y is selected from the group consisting of halogen, halogen substituted benzoyloxy, halogen substituted


phenyl(C₁₋₄)alkyl, halogen substituted aryloxy, and halogen substituted C₆₋₁₀ aryl;

U is selected from the group consisting of hydrogen, hydroxy, halogen, halogen substituted benzoyloxy, halogen substituted phenyl(C₁₋₄)alkyl, halogen substituted aryloxy, and halogen substituted C₆₋₁₀ aryl; and

R³⁴, R³⁵, R³⁶, R³⁷, R³⁸, R³⁹ and R⁴⁰ are in each instance independently selected from the group consisting of hydrogen, halogen, hydroxy, C₁₋₄ alkoxy, C₁₋₄ alkyl, and hydroxy(C₁₋₄)alkyl.


46. The compound of claim 45 wherein, U is hydroxy.
47. The compound of claim 45 wherein, Y is halogen or hydrogen.
48. The compound of claim 45 wherein, R³⁴, R³⁵, R³⁶, R³⁷, R³⁸, R³⁹ and R⁴⁰ are in each instance hydrogen.
49. The compound of claim 38 having one of the following structures:





16b


;



17a

,

and



18a

,

wherein Y is selected from the group consisting of hydrogen and F.

50. A pharmaceutical composition comprising a compound of any one of claims 1, 3, 28 and 38.
51. A diagnostic composition for imaging amyloid deposits, comprising a radiolabeled compound of any one of claims 1, 3, 28 and 38.
52. A method of imaging amyloid deposits, comprising:
 - a. introducing into a mammal a detectable quantity of a diagnostic composition of claim 51;
 - b. allowing sufficient time for the labeled compound to be associated with amyloid deposits; and
 - c. detecting the labeled compound associated with one or more amyloid deposits.
53. A method of inhibiting amyloid plaque aggregation in a mammal, comprising administering a composition of claim 50 in an amount effective to inhibit amyloid plaque aggregation.