

Description

The present invention relates to biocidal compositions and their use in controlling the growth of slime forming microbes in an aqueous system.

5 The formation of slimes by microorganisms is a problem that is encountered in many aqueous systems. For example, the problem is not only found in natural waters such as, for example, lagoons, lakes or ponds, and confined waters (as, for example, in pools), but also in such industrial systems as, for example, cooling water systems, air washer systems and pulp and paper mill systems. Such locations all possess conditions which are conducive to the growth and reproduction of slime-forming microorganisms. 10 In both once-through and recirculating cooling systems, for example, which employ large quantities of water as a cooling medium, the formation of slime by microorganisms is an extensive and constant problem.

Airborne organisms are readily entrained in the water from cooling towers and find this warm medium an ideal environment for growth and multiplication. Aerobic and heliotropic organisms flourish on the tower proper while other organisms colonize and grow in areas such as, for example, the tower sump and the piping and passages of the cooling system. The slime formation not only aids in the deterioration of the tower structure in the case of wooden towers, but also, by its deposition on metal surfaces, promotes corrosion. In addition, slime carried through the cooling system plugs and fouls items such as, for example, lines, valves and strainers, and also deposits on heat exchange surfaces. In the latter case, the impedance 20 of heat transfer can greatly reduce the efficiency of the cooling system.

In pulp and paper mill systems, slime formed by microorganisms is commonly encountered and causes fouling, plugging, or corrosion thereof. The slime also becomes entrained in the paper produced to cause breakouts on the paper machines, with consequent work stoppages and the loss of production time, and/or is responsible for unsightly blemishes in the final product, which result in rejects and wasted output.

25 The previously discussed problems have resulted in the extensive utilization of biocides in cooling water and pulp and paper mill systems. Materials which have enjoyed widespread use in such applications include chlorine, chlorinated phenols, organo-bromines, and various organo-sulphur compounds. All of these compounds are generally useful for this purpose but each is attended by a variety of impediments. For example, chlorination is limited both by its specific toxicity for slime-forming organisms at economic 30 levels and by the tendency of chlorine to react, which results in the expenditure of the chlorine before its full biocidal function is achieved. Other biocides are attended by odour problems and hazards in respect to storage, use or handling which limit their utility. To date, no one compound or type of compound has achieved a clearly established predominance in respect to the applications discussed. Likewise, lagoons, ponds, lakes, and even pools, either used for pleasure purposes or used for industrial purposes for the disposal and storage of industrial wastes, become, during the warm weather, besieged by slime due to 35 microorganism growth and reproduction. In the case of the recreational areas the problem of infection is obvious. In the case of industrial storage or disposal of industrial materials, the microorganisms cause additional problems which must be eliminated prior to use of the material or disposal of the waste.

Naturally, economy is a major consideration in respect to all of these biocides. Such economic 40 considerations attach to both the cost of the biocide and the expense of its application. The cost performance index of any biocide is derived from the basic cost of the material, its effectiveness per unit of weight, the duration of its biocidal or biostatic effect in the system treated, and the ease and frequency of its addition to the system treated. To date, none of the commercially available biocides has exhibited a prolonged biocidal effect. Instead, their effectiveness is rapidly reduced as the result of exposure to 45 physical conditions such as, for example, temperature, association with ingredients contained by the system toward which they exhibit an affinity of substantivity, etc., with a resultant restriction or elimination of their biocidal effectiveness, or by dilution.

As a consequence, the use of such biocides involves their continuous or frequent addition to systems to be treated and their addition to a plurality of points or zones in the systems to be treated. Accordingly, 50 the cost of the biocide and the labour cost of such means of applying it are considerable. In other instances, the difficulty of access to the zone in which slime formation is experienced precludes the effective use of a biocide. For example, if in a particular system there is no access to an area at which slime formation occurs the biocide can only be applied at a point which is upstream in the flow system. However, the physical or chemical conditions, e.g., chemical reactivity or thermal degradation, which exist between the point at 55 which the biocide may be added to the system and the point at which its biocidal effect is desired render the effective use of a biocide impossible.

Similarly, in a system experiencing relatively slow flow, such as, for example, a paper mill, if a biocide is added at the beginning of the system, its biocidal effect may be completely dissipated before it has reached all of the points at which this effect is desired or required. As a consequence, the biocide must be 60 added at a plurality of points, and even then a diminishing biocidal effect will be experienced between one point of addition to the system and the next point downstream at which the biocides may be added. In addition to the increased cost of utilizing and maintaining plural feed points, a gross uneconomic situation with respect to the cost of the biocide is experienced. Specifically, at each point of addition, an excess of the biocide must be added to the system in order to compensate for that portion of the biocide which will be 65

expended in reacting with other constituents present in the system or experience physical changes which impair its biocidal activity.

Use of specific n-alkyl dimethyl benzyl ammonium chlorides and n-dodecylguanidine salts, singly, as a biocide is known in the art. For instance n-dodecylguanidine hydrochloride and n-dodecylguanidine acetate are both listed at 21 CFR §176.170 as being F.D.A. acceptable antimicrobial agents for paper and paperboard use under certain conditions.

Similarly, n-alkyl (C_{12} — C_{18}) dimethyl benzyl ammonium chloride is listed at 21 CFR 176.300 as being a slimicide that may be safely used in the manufacture of paper and paperboard products that contact food. Use of specific n-alkyl dimethyl benzyl ammonium chlorides in various biocidal compositions is also disclosed in US—A—3 934 025, US—A—3 827 873, and US—A—3 881 008.

Other patent specifications which may be noted of possible interest include US—A—3 264 172; US—A—3 142 615; US—A—2 906 595; and US—A—2 409 883.

It has now been surprisingly found that mixtures of n-alkyl dimethyl benzyl ammonium chlorides and n-dodecylguanidine hydrochloride or water soluble salts therefore are especially efficacious in controlling the growth of slime-forming microbes.

According to the present invention there is provided a biocidal composition which comprises a synergistic mixture of n-alkyl dimethyl benzyl ammonium chloride having an alkyl distribution of about 50% C_{14} , 40% C_{12} , 10% C_{16} and n-dodecylguanidine or water soluble salt thereof, the weight ratio of the n-dodecylguanidine or water soluble salt thereof to the n-alkyl dimethyl benzyl ammonium chloride being from 1:2 to 1:8.

The present invention also provides a method for controlling the growth of slime forming microbes in an aqueous system which comprises adding to the system from 0.1 to 200 parts per weight of a synergistic composition per million parts by weight of the aqueous system, the composition comprising a mixture of n-alkyl dimethyl benzyl ammonium chloride having an alkyl distribution of about 50% C_{14} , 40% C_{12} , 10% C_{16} and n-dodecylguanidine or water soluble salt thereof, the weight ratio of the n-dodecylguanidine or water soluble salt thereof to the n-alkyl dimethyl benzyl ammonium chloride being from 1:2 to 1:8.

The n-alkyl dimethyl benzyl ammonium chloride is sometimes hereinafter referred to as "quat" and the n-dodecylguanidine or water soluble salt thereof is sometimes hereinafter referred to as "DGH".

The composition of the present invention has been found to be especially efficacious in controlling the growth of bacterial microbes, particularly the *Klebsiella pneumoniae* species. This particular species is a member of the capsulated, facultative class of bacteria and is generally present in air, water and soil. These bacteria continually contaminate open cooling systems and pulping and papermaking systems and are among the most common slime formers. This slime may be viewed as being a mass of agglomerated cells stuck together by the cementing action of the gelatinous polysaccharide or proteinaceous secretions around each cell. The slimy mass entraps other debris, restricts water flow and heat transfer, and may serve as a site for corrosion.

The fact that the *Klebsiella* species used in the tests is a facultative species is important as, by definition, such bacteria may thrive under either aerobic or anaerobic conditions. Accordingly, by reason of demonstrated efficacy in the growth inhibition of this particular species, one can expect similar growth inhibition attributes when other aerobic and anaerobic bacterial species are encountered.

The n-alkyl (50% C_{14} , 40% C_{12} , 10% C_{16}) dimethyl benzyl ammonium chloride is commercially available from various sources. For instance, it is sold under the trademark "Maquat 1412" from Mason Chemical Co., "Onyx 8358" from Onyx Chemical Co., or "Hyamine 3500" by Lonza Inc.

As to the water soluble n-dodecylguanidine salt that may be used, the hydrochloride, acetate, carbonate and nitrate salts may be mentioned as being exemplary. The preferred hydrochloride salt is commercially available under the trademark "Cytox 2013" and "Cytox 2014" from American Cyanamid.

In accordance with the present invention, the combined quat:DGH treatment may be added to the desired aqueous system in need of biocidal treatment, in an amount of from about 0.001 to about 200 parts of the combined treatment to one million parts (by weight) of the aqueous medium. Preferably, about 5 to about 50 parts of the combined treatment per one million parts (by weight) of the aqueous medium is added.

The combined treatment is added, for example, to cooling water systems, paper and pulp mill systems, pools, ponds, lagoons, or lakes, to control the formation of bacterial microorganisms, which may be contained by, or which may become entrained in, the system to be treated. It has been found that the quat/DGH compositions and methods of utilization of the treatment are efficacious in controlling the facultative bacterium, *Klebsiella pneumoniae*, which may populate these systems. It is thought that the combined treatment composition and method of the present invention will also be efficacious in inhibiting and controlling all types of aerobic and anaerobic bacteria.

Surprisingly, it has been found that when the quat/DGH ingredients are mixed, in certain instances, the resulting mixtures possess a higher degree of bactericidal activity than that of the individual ingredients comprising the mixture. Accordingly, it is possible to produce a highly efficacious bactericide. Because of the enhance activity of the mixture, the total quantity of the bacterial treatment may be reduced. In addition, the high degree of bactericidal effectiveness which is provided by each of the ingredients may be exploited without use of higher concentrations of each.

The following experimental data were developed. It is to be remembered that the following examples

are to be regarded solely as being illustrative, and not as restricting the scope of the invention.

Quat and DGH were added in varying ratios and over a wide range of concentrations to a liquid nutrient medium which was subsequently inoculated with a standard volume of a suspension of the facultative bacterium *Klebsiella pneumoniae*. Growth was measured by determining the amount of radioactivity accumulated by the cells when ^{14}C -glucose was added as the sole source of carbon in the nutrient medium. The effect of the biocide chemicals, alone and in combination, is to reduce the rate and amount of ^{14}C incorporation into the cells during incubation, as compared to controls not treated with the chemicals. Additions of the biocides, alone and in varying combinations and concentrations, were made according to the accepted "checkerboard" technique described by M. T. Kelley and J. M. Matsen, *Antimicrobial Agents and Chemotherapy*, 9: 440, (1976). Following a four hour incubation, the amount of radioactivity incorporated in the cells was determined by counting (^{14}C liquid scintillation procedures) for all treated and untreated samples. The percent reduction of each treated sample was calculated from the relationship.

$$\frac{\text{Control } ^{14}\text{C(cpm)} - \text{Treated } ^{14}\text{C(cpm)}}{\text{Control } ^{14}\text{C(cpm)}} \times 100 = \% \text{ reduction}$$

Plotting the % reduction of ^{14}C level against the concentration of each biocide acting alone results in a dose-response curve, from which the biocide dose necessary to achieve any given % reduction can be interpolated.

Synergism was determined by the method of calculation described by F. C. Kull, P. C. Eisman, H. D. Sylwestrowicz and R. L. Mayer, *Applied Microbiology* 9, 538, (1961) using the relationship:

$$\frac{Q_A}{Q_a} + \frac{Q_B}{Q_b} = \text{synergism index (SI)}$$

where:

Q_A = quantity of compound A, acting alone, producing an end point

Q_b = quantity of compound B, acting alone, producing an end point

Q_A = quantity of compound A in mixture, producing an end point

Q_b = quantity of compound B in mixture, producing an end point

The end point used in the calculations is the % reduction caused by each mixture of A and B. Q_A and Q_b are the individual concentrations in the A/B mixture causing a given % reduction. Q_a and Q_b are determined by interpolation from the respective dose-response curves of A and B as those concentrations of A and B acting alone which produce the same % reduction as each specific mixture produced.

Dose-response curves for each active acting alone were determined by linear regression analysis of the dose-response data. Data were fitted to a curve represented by the equation shown with each data set. After linearizing the data, the contributions of each biocide component in the biocide mixtures to the inhibition of radioisotope uptake were determined by interpolation with the dose response curve of the respective biocide. If, for example, quantities of Q_A plus Q_b are sufficient to give a 50% reduction in ^{14}C content, Q_a and Q_b are those quantities of A or B acting alone, respectively, found to give 50% reduction in ^{14}C content. A synergism index (SI) is calculated for each combination of A and B.

Where the SI is <1 , synergism exists. Where the SI = 1, additivity exists. Where SI >1 , antagonism exists.

The data in the following Tables come from treating *Klebsiella pneumoniae*, a common nuisance bacterial type found in industrial cooling waters and in pulping and paper making systems, with varying ratios and concentrations of quat and DGH. Shown for each combination is the % reduction of ^{14}C content, the calculated SI, and the weight ratio of quat to DGH.

Table I

	DGH	vs	Quat		
	ppm DGH(1)	ppm Quat(2)	ratio DGH:Quat	%I	SI
	40	0	100:0	99	
	20	0	100:0	97	
	10	0	100:0	80	
	5	0	100:0	0	
	2.5	0	100:0	0	
	1.25	0	100:0	0	
	0	20	0:100	99	
	0	10	0:100	97	

EP 0 255 239 B1

Table I (Continued)

ppm DGH(1)	ppm Quat(2)	ratio DGH:Quat	%I	SI
0	5	0:100	78	
0	4	0:100	62	
0	2.5	0:100	0	
0	1.25	0:100	0	
40	20	2:1	99	4.26
20	20	1:1	99	3.05
10	20	1:2	99	2.44
5	20	1:4	99	2.14
2.5	20	1:8	99	1.99
1.25	20	1:16	99	1.91
40	10	4:1	99	3.34
20	10	2:1	99	2.13
10	10	1:1	98	1.55
5	10	1:2	97	1.26
2.5	10	1:4	97	1.11
1.25	10	1:8	97	1.03
40	5	8:1	99	2.88
20	5	4:1	99	1.67
10	5	2:1	99	1.06
5	5	1:1	92	0.85*
2.5	5	1:2	89	0.72*
1.25	5	1:4	85	0.67*
40	4	10:1	99	2.79
20	4	5:1	99	1.58
10	4	2.5:1	96	1.02
5	4	1.25:1	92	0.75*
2.5	4	1:1.6	85	0.65*
1.25	4	1:3.2	82	0.59*
40	2.5	16:1	99	2.65
20	2.5	8:1	99	1.44
10	2.5	4:1	91	0.93*
5	2.5	2:1	76	0.75*
2.5	2.5	1:1	0	
1.25	2.5	1:2	0	
40	1.25	1:32	99	2.50
20	1.25	1:16	99	1.33
10	1.25	1:8	93	0.78*
5	1.25	1:4	12	1.39
2.5	1.25	1:2	0	
1.25	1.25	1:1	0	

(1) product containing 35% actives

(2) product containing 80% actives

Asterisks (*) in the SI column indicate synergistic combinations in accordance with the Kull method supra.

EP 0 255 239 B1

Table II
DGH vs Quat

ppm DGH(1)	ppm Quat(2)	ratio DGH:Quat	%I	SI
15	0	100:0	54	
10	0	100:0	28	
7.5	0	100:0	29	
6	0	100:0	20	
5	0	100:0	10	
4.5	0	100:0	15	
0	7.5	0:100	51	
0	6	0:100	38	
0	4.5	0:100	25	
0	3.75	0:100	18	
0	3.5	0:100	17	
0	3	0:100	11	
15	7.5	2:1	72	1.54
10	7.5	1.3:1	58	1.54
7.5	7.5	1:1	59	1.36
6	7.5	1:1.25	57	1.31
5	7.5	1:1.5	53	1.31
4.5	7.5	1:1.6	54	1.26
15	6	2.5:1	65	1.51
10	6	1.6:1	56	1.40
7.5	6	1.25:1	57	1.21
6	6	1:1	51	1.23
5	6	1:1.2	47	1.23
4.5	6	1:1.3	49	1.16
15	4.5	3.3:1	62	1.41
10	4.5	2.2:1	51	1.31
7.5	4.5	1.7:1	50	1.15
6	4.5	1.3:1	46	1.12
5	4.5	1.1:1	41	1.14
4.5	4.5	1:1	42	1.07
15	3.75	4:1	59	2.91
10	3.75	2.7:1	46	1.32
7.5	3.75	2:1	48	1.08
6	3.75	1.6:1	43	1.06
5	3.75	1.3:1	38	1.08
4.5	3.75	1.2:1	38	1.03
15	3.5	4.3:1	58	1.37
10	3.5	2.9:1	48	1.24
7.5	3.5	2.1:1	45	1.11
6	3.5	1.7:1	43	1.03
5	3.5	1.4:1	38	1.04
4.5	3.5	1.3:1	37	1.01
15	3	5:1	58	1.31
10	3	3.3:1	46	1.21
7.5	3	2.5:1	43	1.53
6	3	2:1	43	0.95*
5	3	1.7:1	36	1.00
4.5	3	1.5:1	39	0.91*

(1) product containing 35% actives (2) product containing 80% actives
Asterisks (*) in the SI column indicate synergistic combinations
in accordance with the Kull method supra.

EP 0 255 239 B1

Table III
DGH vs Quat

ppm DGH(1)	ppm Quat(2)	ratio DGH:Quat	%I	SI
40	0	100:0	95	
20	0	100:0	80	
10	0	100:0	46	
5	0	100:0	18	
2.5	0	100:0	5	
1.25	0	100:0	2	
0	20	0:100	95	
0	10	0:100	78	
0	5	0:100	40	
0	2.5	0:100	6	
0	1.25	0:100	0	
0	0.625	0:100	0	
40	20	2:1	99	1.88
20	20	1:1	98	1.47
10	20	1:2	97	1.27
5	20	1:4	96	1.19
2.5	20	1:8	96	1.13
1.25	20	1:16	97	1.07
40	10	4:1	98	1.43
20	10	2:1	94	1.09
10	10	1:1	87	0.98
5	10	1:2	82	0.95*
2.5	10	1:4	80	0.90*
1.25	10	1:8	80	0.86*
40	5	8:1	98	1.18
20	5	4:1	89	0.93*
10	5	2:1	71	1.04
5	5	1:1	58	1.08
2.5	5	1:2	49	1.15
1.25	5	1:4	49	1.01
40	2.5	16:1	97	1.08
20	2.5	8:1	84	0.90*
10	2.5	4:1	64	0.66*
5	2.5	2:1	41	1.24
2.5	2.5	1:1	34	1.08
1.25	2.5	1:2	29	1.00
40	1.25	32:1	97	1.02
20	1.25	16:1	82	0.86*
10	1.25	8:1	53	1.16
5	1.25	4:1	32	1.28
2.5	1.25	2:1	21	1.10
1.25	1.25	1:1	19	0.82*
40	0.625	64:1	97	0.99
20	0.625	32:1	81	0.84*
10	0.625	16:1	49	1.21
5	0.625	8:1	31	1.14
2.5	0.625	4:1	20	0.93*
1.25	0.625	2:1	20	0.58*

(1) product containing 35% actives (2) product containing 80% actives

Asterisks (*) in the SI column indicate synergistic combinations in accordance with the Kull method supra.

In accordance with Tables I—III supra., unexpected results occurred more frequently within the product ratios of DGH product: Quat product of from about 1:1 to 1:8. Since the tested DGH product was about 35% active and the Quat product about 80% active, this range translates to a range of DGH:Quat (100% actives basis) of from about 1:2 to about 1:8. At present, it is preferred that the commercial product embodying the invention comprise a weight ratio of about 1:2 DGH:Quat.

Claims

1. A biocidal composition which comprises a synergistic mixture of n-alkyl dimethyl benzyl ammonium chloride having an alkyl distribution of about 50% C₁₄, 40% C₁₂, 10% C₁₆ and n-dodecylguanidine or water soluble salt thereof, the weight ratio of the n-dodecylguanidine or water soluble salt thereof to the n-alkyl dimethyl benzyl ammonium chloride being from 1:2 to 1:8.
2. A method for controlling the growth of slime forming microbes in an aqueous system which comprises adding to the system from 0.1 to 200 parts per weight of a synergistic composition per million parts by weight of the aqueous system, the composition comprising a mixture of n-alkyl dimethyl benzyl ammonium chloride having an alkyl distribution of about 50% C₁₄, 40% C₁₂, 10% C₁₆ and n-dodecylguanidine or water soluble salt thereof, the weight ratio of the n-dodecylguanidine or water soluble salt thereof to the n-alkyl dimethyl benzyl ammonium chloride being from 1:2 to 1:8.
3. A method according to claim 2, wherein the composition is added to the system in an amount of from 5 to 50 parts per million of the aqueous system.
4. A method according to claim 2 or 3, wherein the microbes comprise bacteria.
5. A method according to any of claims 2 to 4, wherein the microbes comprise *Klebsiella pneumoniae* bacteria.
6. A method according to any of claims 2 to 5, wherein the aqueous system comprises a cooling water system.
7. A method according to any of claims 2 to 5, wherein the aqueous system comprises a pulping and papermaking system.

Patentansprüche

1. Biozide Zusammensetzung, enthaltend ein synergistisches Gemisch aus n-Alkyldimethylbenzylammoniumchlorid mit einer Alkyl-Verteilung von etwa 50% C₁₄, 40% C₁₂, 10% C₁₆ und n-Dodecylguanidin oder ein wasserlösliches Salz davon, wobei das Gewichtsverhältnis von n-Dodecylguanidin oder eines wasserlöslichen Salzes davon zu dem n-Alkyldimethylbenzylammoniumchlorid 1:2 bis 1:8 beträgt.
2. Verfahren zum Einschränken des Wachstums von schleimbildenden Mikroben in einem wässrigen System, bei dem man zu dem System 0,1 bis 200 Gewichtsteile einer synergistischen Zusammensetzung pro Million Gewichtsteilen des wässrigen Systems gibt, und die Zusammensetzung ein Gemisch aus n-Alkyldimethylbenzylammoniumchlorid mit einer Alkylverteilung von etwa 50% C₁₄, 40% C₁₂, 10% C₁₆ und n-Dodecylguanidin oder ein wasserlösliches Salz davon enthält, wobei das Gewichtsverhältnis von n-Dodecylguanidin oder des wasserlöslichen Salzes davon zu n-Alkyldimethylbenzylammoniumchlorid 1:2 bis 1:8 beträgt.
3. Verfahren gemäß Anspruch 2, bei dem man die Zusammensetzung zu dem System in einer Menge von 5 bis 20 Teilen pro Million Teilen des wässrigen Systems gibt.
4. Verfahren gemäß Anspruch 2 oder 3, bei dem die Mikroben Bakterien umfassen.
5. Verfahren gemäß einem der Ansprüche 2 bis 4, bei dem die Mikroben *Klebsiella pneumoniae*-Bakterien umfassen.
6. Verfahren gemäß einem der Ansprüche 2 bis 5, bei dem das wässrige System ein Kühlwassersystem umfaßt.
7. Verfahren gemäß einem der Ansprüche 2 bis 5, bei dem das wässrige System ein Papieraufschluß- und Papierherstellungssystem umfaßt.

Revendications

1. Composition biocide qui comprend un mélange synergique de chlorure de n-alkyl-diméthyl-benzylammonium ayant une distribution de groupes alkyle d'environ 50% en C₁₄, 40% C₁₂ et 10% en C₁₆, et de n-dodécylguanidine ou d'un sel hydrosoluble de celle-ci, le rapport en poids de la n-dodécylguanidine ou de son sel hydrosoluble au chlorure de n-alkyl-diméthyl-benzylammonium étant de 1:2 à 1:8.
2. Procédé pour combattre la croissance de microbes formateurs de vase dans un système aqueux, qui consiste à ajouter au système 0,1 à 200 parties en poids d'une composition synergique par million de parties en poids du système aqueux, la composition comprenant un mélange de chlorure de n-alkyl-diméthyl-benzylammonium ayant une distribution de groupes alkyle d'environ 50% en C₁₄, 40% en C₁₂ et 10% en C₁₆, et de n-dodécylguanidine ou d'un sel hydrosoluble de celle-ci, le rapport en poids de la n-dodécylguanidine ou de son sel hydrosoluble au chlorure de n-alkyl-diméthyl-benzylammonium étant de 1:2 à 1:8.

In accordance with Tables I—III supra., unexpected results occurred more frequently within the product ratios of DGH product: Quat product of from about 1:1 to 1:8. Since the tested DGH product was about 35% active and the Quat product about 80% active, this range translates to a range of DGH:Quat (100% actives basis) of from about 1:2 to about 1:8. At present, it is preferred that the commercial product embodying the invention comprise a weight ratio of about 1:2 DGH:Quat.

Claims

1. A biocidal composition which comprises a synergistic mixture of n-alkyl dimethyl benzyl ammonium chloride having an alkyl distribution of about 50% C₁₄, 40% C₁₂, 10% C₁₆ and n-dodecylguanidine or water soluble salt thereof, the weight ratio of the n-dodecylguanidine or water soluble salt thereof to the n-alkyl dimethyl benzyl ammonium chloride being from 1:2 to 1:8.
2. A method for controlling the growth of slime forming microbes in an aqueous system which comprises adding to the system from 0.1 to 200 parts per weight of a synergistic composition per million parts by weight of the aqueous system, the composition comprising a mixture of n-alkyl dimethyl benzyl ammonium chloride having an alkyl distribution of about 50% C₁₄, 40% C₁₂, 10% C₁₆ and n-dodecylguanidine or water soluble salt thereof, the weight ratio of the n-dodecylguanidine or water soluble salt thereof to the n-alkyl dimethyl benzyl ammonium chloride being from 1:2 to 1:8.
3. A method according to claim 2, wherein the composition is added to the system in an amount of from 5 to 50 parts per million of the aqueous system.
4. A method according to claim 2 or 3, wherein the microbes comprise bacteria.
5. A method according to any of claims 2 to 4, wherein the microbes comprise *Klebsiella pneumoniae* bacteria.
6. A method according to any of claims 2 to 5, wherein the aqueous system comprises a cooling water system.
7. A method according to any of claims 2 to 5, wherein the aqueous system comprises a pulping and papermaking system.

Patentansprüche

1. Biozide Zusammensetzung, enthaltend ein synergistisches Gemisch aus n-Alkyldimethylbenzylammoniumchlorid mit einer Alkyl-Verteilung von etwa 50% C₁₄, 40% C₁₂, 10% C₁₆ und n-Dodecylguanidin oder ein wasserlösliches Salz davon, wobei das Gewichtsverhältnis von n-Dodecylguanidin oder eines wasserlöslichen Salzes davon zu dem n-Alkyldimethylbenzylammoniumchlorid 1:2 bis 1:8 beträgt.
2. Verfahren zum Einschränken des Wachstums von schleimbildenden Mikroben in einem wässrigen System, bei dem man zu dem System 0,1 bis 200 Gewichtsteile einer synergistischen Zusammensetzung pro Million Gewichtsteilen des wässrigen Systems gibt, und die Zusammensetzung ein Gemisch aus n-Alkyldimethylbenzylammoniumchlorid mit einer Alkylverteilung von etwa 50% C₁₄, 40% C₁₂, 10% C₁₆ und n-Dodecylguanidin oder ein wasserlösliches Salz davon enthält, wobei das Gewichtsverhältnis von n-Dodecylguanidin oder des wasserlöslichen Salzes davon zu n-Alkyldimethylbenzylammoniumchlorid 1:2 bis 1:8 beträgt.
3. Verfahren gemäß Anspruch 2, bei dem man die Zusammensetzung zu dem System in einer Menge von 5 bis 20 Teilen pro Million Teilen des wässrigen Systems gibt.
4. Verfahren gemäß Anspruch 2 oder 3, bei dem die Mikroben Bakterien umfassen.
5. Verfahren gemäß einem der Ansprüche 2 bis 4, bei dem die Mikroben *Klebsiella pneumoniae*-Bakterien umfassen.
6. Verfahren gemäß einem der Ansprüche 2 bis 5, bei dem das wässrige System ein Kühlwassersystem umfaßt.
7. Verfahren gemäß einem der Ansprüche 2 bis 5, bei dem das wässrige System ein Papieraufschluß- und Papierherstellungssystem umfaßt.

Revendications

1. Composition biocide qui comprend un mélange synergique de chlorure de n-alkyl-diméthyl-benzyl-ammonium ayant une distribution de groupes alkyle d'environ 50% en C₁₄, 40% C₁₂ et 10% en C₁₆, et de n-dodécylguanidine ou d'un sel hydrosoluble de celle-ci, le rapport en poids de la n-dodécylguanidine ou de son sel hydrosoluble au chlorure de n-alkyl-diméthyl-benzylammonium étant de 1:2 à 1:8.
2. Procédé pour combattre la croissance de microbes formateurs de vase dans un système aqueux, qui consiste à ajouter au système 0,1 à 200 parties en poids d'une composition synergique par million de parties en poids du système aqueux, la composition comprenant un mélange de chlorure de n-alkyl-diméthyl-benzylammonium ayant une distribution de groupes alkyle d'environ 50% en C₁₄, 40% en C₁₂ et 10% en C₁₆, et de n-dodécylguanidine ou d'un sel hydrosoluble de celle-ci, le rapport en poids de la n-dodécylguanidine ou de son sel hydrosoluble au chlorure de n-alkyl-diméthyl-benzylammonium étant de 1:2 à 1:8.



EP 0 255 239 B1

3. Procédé selon la revendication 2, dans lequel la composition est ajoutée au système en une quantité de 5 à 50 parties par million de parties du système aqueux.

4. Procédé selon la revendication 2 ou 3, dans lequel les microbes comprennent des bactéries.

5. Procédé selon l'une quelconque des revendications 2 à 4, dans lequel les microbes sont des bactéries de l'espèce *Klebsiella pneumoniae*.

6. Procédé selon l'une quelconque des revendications 2 à 5, dans lequel le système aqueux est un système de refroidissement à eau.

7. Procédé selon l'une quelconque des revendications 2 à 5, dans lequel le système aqueux est un système de fabrication de pâte et de papier.

10

15

20

25

30

35

40

45

50

55

60

65

