What is Claimed Is:

1. A topical formulation, comprising:

(i) an active agent comprising salvigenin in an amount of about 0.0001% to 0.5% (w/w) and asiaticoside in an amount of about 0.05% to 5% (w/w);

(ii) a viscosity increasing agent in an amount of about 1.0-10% (w/w);

(iii) an ointment base in an amount of about 5-30% (w/w);

(iv) an antimicrobial preservative in an amount of about 0.005-0.2% (w/w); and

(v) an emulsifying agent in an amount of about 0.5-10% (w/w);

wherein the viscosity increasing agent comprises cetostearyl alcohol in an amount of about 3-7% (w/w);

wherein the ointment base comprises liquid petrolatum and white petrolatum in a total amount of about 15-25% (w/w); and

wherein the emulsifying agent comprises sorbitan monostearate and polysorbate 60 in a total amount of about 2-4% (w/w).

2. The topical formulation of claim 1, wherein the active agent is a *Plectranthus amboinicus* extract, a *Centella asiatica* extract, or a combination thereof, and wherein the *Plectranthus amboinicus* extract, the *Centella asiatica* extract, or the combination thereof comprises the salvigenin and asiaticoside.

3. The topical formulation of claim 2, wherein the *Plectranthus amboinicus* extract, the *Centella asiatica* extract, or the combination thereof is in an amount of about 0.1-30% (w/w) in the topical formulation.

4. The topical formulation of claim 1, wherein the topical formulation comprises salvigenin ranging from 0.0001% to 0.1% (w/w), and/or wherein the topical formulation comprises asiaticoside ranging from 0.05% to 1% (w/w).

5. The topical formulation of claim 1, wherein the active agent further comprises cirsimaritin, madecassoside, or a combination thereof.

6. The topical formulation of claim 2, wherein the *Plectranthus amboinicus* extract is prepared by contacting a whole *Plectranthus amboinicus* or a part thereof with a solvent having a polarity index lower than 7 to produce a solution and drying the solution to produce the *Plectranthus amboinicus* extract.

7. The topical formulation of claim 2, wherein the topical formulation comprises both the *Plectranthus amboinicus* extract and the *Centella asiatica* extract at a weight ratio of about 1:10 to 10:1.

8. The topical formulation of claim 1, wherein the viscosity increasing agent further comprises cholesterol, stearyl alcohol, chlorocresol, white wax, stearic acid, cetyl alcohol, or a combination thereof; and wherein the antimicrobial preservative comprises one or more paraben compounds.

9. The topical formulation of claim 1, wherein the ointment base is a cream base.

10. The topical formulation of claim 1, wherein the formulation further comprises one or more solvents.

11. The topical formulation of claim 10, wherein the one or more solvents comprise propylene glycol.

12. The topical formulation of claim 2, wherein the topical formulation comprises a combination of the *Plectranthus amboinicus* extract and the *Centella asiatica* extract in an amount of about 0.1-30% (w/w), cetostearyl alcohol in an amount of about 5-7% (w/w), a combination of white petrolatum and liquid petrolatum in a total amount of

about 20% (w/w), a combination of methyl paraben and propyl paraben in a total amount of about 0.005-0.2% (w/w), and a combination of sorbitan monostearate and polysorbate 60 in a total amount of about 3% (w/w).

13. The topical formulation of claim 12, which further comprises propylene glycol in an amount of about 2-20% (w/w).