
METHODS OF IMMUNOTHERAPY

RELATED APPLICATIONS

5 This application claims the benefit of priority to U.S. Provisional Patent Application serial number 62/341,360, filed May 25, 2016, and U.S. Provisional Patent Application serial number 62/487,814, filed April 20, 2017, each of which is hereby incorporated by reference in its entirety.

BACKGROUND

10 Adoptive immunotherapy involves implanting or infusing disease-specific T cells, such as cytotoxic T cells (CTLs), into individuals with the aim of recognizing, targeting, and destroying disease-associated cells. Adoptive immunotherapy has become a promising route for the treatment of many diseases and disorders, including cancer, infectious diseases and autoimmune diseases. However, the efficiency of adoptive immunotherapy can vary from cell sample to cell sample, thus creating a need for methods of predicting the likely
15 therapeutic efficacy of an immunotherapeutic sample.

SUMMARY

In certain aspects, provided herein are methods related to selecting samples comprising T cells (*e.g.*, CD4 T cells or CD8 T cells, such as CTLs) for use in adoptive immunotherapy and/or for inclusion into a cell bank based on the expression of CD107a,
20 interferon gamma (IFNg), interleukin 2 (IL-2) tumor necrosis factor (TNF), granzyme B (GzmB), granzyme K (GzmK) and/or perforin (Prf) by the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells in the samples. In some embodiments, the samples are selected based on the expression of CD8, CD4, PD-1, TIM-3, LAG-3 and/or CTLA-4 by the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells in the samples. In some embodiments,
25 the samples are selected based on the expression of antigen-specific T cell receptors (TCR) by the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells in the samples. In some embodiments, the samples are selected if the expression of CD107a, interferon gamma IFNg, IL-2, TNF, GzmB, GzmK, Prf, CD8, CD4, PD-1, TIM-3, LAG-3, CTLA-4 and/or antigen-specific TCR is above a threshold level provided herein. In some embodiments, the samples
30 are selected if the expression of IFNg, IL-2, TNF and CD-107a is above a threshold level

provided herein. In some embodiments, the samples are not selected if the expression of CD107a, interferon gamma (IFN γ), interleukin 2 (IL-2), tumor necrosis factor (TNF), granzyme B (GzmB), granzyme K (GzmK), perforin (Prf), CD8, CD4, PD-1, TIM-3, LAG-3, CTLA-4 and/or antigen-specific TCR is below a threshold level provided herein. In some
5 embodiments, the samples are not selected if the expression of IFN γ , IL-2, TNF and CD-107a is below a threshold level provided herein. In certain aspects, provided herein are methods of treating a subject in need thereof (*e.g.*, a subject with cancer, a viral infection and/or an autoimmune disease) by administering to the subject a sample comprising T cells (*e.g.*, CD4 T cells or CD8 T cells, such as CTLs) selected according to a method described
10 herein. In certain embodiments, provided herein is a method of generating a cell bank of samples comprising T cells (*e.g.*, CD4 T cells or CD8 T cells, such as CTLs) for adoptive immunotherapy, the method comprising adding one or more samples selected according to a method provided herein to a cell bank. In some aspects, provided herein is a cell bank generated according to a method described herein.

15 In certain aspects, provided herein are methods related to selecting samples comprising T cells (*e.g.*, CD8 and/or CD4 T cells) for use in adoptive immunotherapy and/or for inclusion into a cell bank based on the expression of IFN γ , IL-2, TNF, GzmB, GzmK, Prf, CD8, CD4, PD-1, TIM-3, LAG-3, CTLA-4 and/or antigen-specific TCR by the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells in the samples. In some embodiments,
20 the samples are selected if the expression of IFN γ , IL-2, TNF, GzmB, GzmK, Prf, CD8, CD4, PD-1, TIM-3, LAG-3, CTLA-4 and/or antigen-specific TCR is above a threshold level provided herein. In some embodiments, the samples are selected if the expression of IFN γ , IL-2, TNF and CD-107a is above a threshold level provided herein. In some embodiments, the samples are not selected if the expression of IFN γ , IL-2, TNF, GzmB, GzmK, Prf, CD8,
25 CD4, PD-1, TIM-3, LAG-3, CTLA-4 and/or antigen-specific TCR is below a threshold level provided herein. In some embodiments, the samples are not selected if the expression of IFN γ , IL-2, TNF and CD-107a is below a threshold level provided herein. In certain aspects, provided herein are methods of treating a subject in need thereof (*e.g.*, a subject with cancer, a viral infection and/or an autoimmune disease) by administering to the subject a sample
30 comprising T cells (*e.g.*, CD8 and/or CD4 T cells) selected according to a method described herein. In certain embodiments, provided herein is a method of generating a cell bank of samples comprising T cells (*e.g.*, CD8 and/or CD4 T cells) for adoptive immunotherapy, the

method comprising adding one or more samples selected according to a method provided herein to a cell bank. In some aspects, provided herein is a cell bank generated according to a method described herein.

In some aspects, provided herein are methods of selecting a sample comprising T
5 cells (*e.g.*, CD8 and/or CD4 T cells) for *ex vivo* expansion (*e.g.*, a large scale expansion for inclusion in a cell bank or for adoptive transfer) based on the expression of IFN γ , IL-2, TNF, GzmB, GzmK, Prf, CD8, CD4, PD-1, TIM-3, LAG-3, CTLA-4 and/or antigen-specific TCR by a portion of the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells in the sample. For example, in some embodiments the samples are selected by first performing a small
10 scale expansion and/or inducing proliferation in T cells from a portion of the sample, measuring the expression of one or more of IFN γ , IL-2, TNF, GzmB, GzmK, Prf, CD8, CD4, PD-1, TIM-3, LAG-3, CTLA-4 and/or antigen-specific TCR by the portion of the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells. In some embodiments, the samples are selected if the expression of IFN γ , IL-2, TNF, GzmB, GzmK, Prf, CD8, CD4, PD-1,
15 TIM-3, LAG-3, CTLA-4 and/or antigen-specific TCR is above a threshold level provided herein. In some embodiments, the samples are selected if the expression of IFN γ , IL-2, TNF and CD-107a is above a threshold level provided herein. In some embodiments, the samples are not selected if the expression of IFN γ , IL-2, TNF, GzmB, GzmK, Prf, CD8, CD4, PD-1, TIM-3, LAG-3, CTLA-4 and/or antigen-specific TCR is below a threshold level provided
20 herein. In some embodiments, the samples are not selected if the expression of IFN γ , IL-2, TNF and CD-107a is below a threshold level provided herein.

In some aspects, provided herein are methods of selecting a subject for autologous adoptive T cell immunotherapy (*e.g.*, using CD8 or CD4 T cells) and/or as a T cell donor based on the expression of IFN γ , IL-2, TNF, GzmB, GzmK, Prf, CD8, CD4, PD-1, TIM-3,
25 LAG-3, CTLA-4 and/or antigen-specific TCR by the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells in a sample obtained from the subject. For example, in some embodiments the subject is selected by first performing an expansion and/or inducing proliferation in T cells (*e.g.*, CD8 and/or CD4 T cells) in a sample obtained from the subject, and measuring the expression of one or more of IFN γ , IL-2, TNF, GzmB, GzmK, Prf, CD8,
30 CD4, PD-1, TIM-3, LAG-3, CTLA-4 and/or antigen-specific TCR by the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells. In some embodiments, the subject is selected if the expression of IFN γ , IL-2, TNF, GzmB, GzmK, Prf, CD8, CD4, PD-1, TIM-3, LAG-3,

CTLA-4 and/or antigen-specific TCR is above a threshold level provided herein. In some embodiments, the subjects are selected if the expression of IFN γ , IL-2, TNF and CD-107a is above a threshold level provided herein. In some embodiments, the subject is not selected if the expression of IFN γ , IL-2, TNF, GzmB, GzmK, Prf, CD8, CD4, PD-1, TIM-3, LAG-3,

- 5 CTLA-4 and/or antigen-specific TCR is below a threshold level provided herein. In some embodiments, the subjects are not selected if the expression of IFN γ , IL-2, TNF and CD-107a is below a threshold level provided herein.

In some embodiments, the threshold level is met if at least about 0.5%, 0.6%, 0.7%, 0.8%, 0.9%, 1%, 1.1%, 1.2%, 1.3%, 1.4%, 1.5%, 1.6%, 1.7%, 1.8%, 1.9%, 2%, 2.5%, 3%,
10 3.5%, 4%, 4.5%, 5%, 6%, 7%, 8%, 9%, 10%, 11%, 12%, 13%, 14%, 15%, 16%, 17%, 18%, 19%, 20%, 21%, 22%, 23%, 24% or 25% of the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells express CD107a.

In some embodiments, the threshold level is met if at least about 0.5%, 0.6%, 0.7%, 0.8%, 0.9%, 1%, 1.1%, 1.2%, 1.3%, 1.4%, 1.5%, 1.6%, 1.7%, 1.8%, 1.9%, 2%, 2.5%, 3%,
15 3.5%, 4%, 4.5%, 5%, 6%, 7%, 8%, 9%, 10%, 11%, 12%, 13%, 14%, 15%, 16%, 17%, 18%, 19%, 20%, 21%, 22%, 23%, 24% or 25% of the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells express IFN γ .

In some embodiments, the threshold level is met if at least about 0.5%, 0.6%, 0.7%, 0.8%, 0.9%, 1%, 1.1%, 1.2%, 1.3%, 1.4%, 1.5%, 1.6%, 1.7%, 1.8%, 1.9%, 2%, 2.5%, 3%,
20 3.5%, 4%, 4.5%, 5%, 6%, 7%, 8%, 9%, 10%, 11%, 12%, 13%, 14%, 15%, 16%, 17%, 18%, 19%, 20%, 21%, 22%, 23%, 24% or 25% of the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells express IL-2.

In some embodiments, the threshold level is met if at least about 0.5%, 0.6%, 0.7%, 0.8%, 0.9%, 1%, 1.1%, 1.2%, 1.3%, 1.4%, 1.5%, 1.6%, 1.7%, 1.8%, 1.9%, 2%, 2.5%, 3%,
25 3.5%, 4%, 4.5%, 5%, 6%, 7%, 8%, 9%, 10%, 11%, 12%, 13%, 14%, 15%, 16%, 17%, 18%, 19%, 20%, 21%, 22%, 23%, 24% or 25% of the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells express TNF.

In some embodiments, the threshold level is met if at least about 5%, 6%, 7%, 8%, 9%, 10%, 11%, 12%, 13%, 14%, 15%, 16%, 17%, 18%, 19%, 20%, 21%, 22%, 23%, 24%
30 or 25% of the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells express granzyme B.

In some embodiments, the threshold level is met if at least about 3%, 4%, 5%, 6%, 7%, 8%, 9%, 10%, 11%, 12%, 13%, 14%, 15%, 16%, 17%, 18%, 19% or 20% of the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells express granzyme K.

In some embodiments, the threshold level is met if at least about 2%, 3%, 4%, 5%, 6%, 7%, 8%, 9%, 10%, 11%, 12%, 13%, 14% or 15% of the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells express perforin.

In some embodiments, the threshold level is met if at least about 15%, 16%, 17%, 18%, 19%, 20%, 21%, 22%, 23%, 24%, 25%, 26%, 27%, 28%, 29%, 30%, 40%, 50%, 60%, 70% or 80% of the total lymphocytes in the sample express CD8.

In some embodiments, the threshold level is met if at least about 1%, 2%, 3%, 4%, 5%, 6%, 7%, 8%, 9%, 10%, 11%, 12%, 13%, 14%, 15%, 16%, 17%, 18%, 19%, 20%, 21%, 22%, 23%, 24% or 25% of the total lymphocytes in the sample express CD4.

In some embodiments, the threshold level is met if at least about 3%, 4%, 5%, 6%, 7%, 8%, 9%, 10%, 11%, 12%, 13%, 14%, 15%, 16%, 17%, 18%, 19% or 20% of the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells express PD-1.

In some embodiments, the threshold level is met if at least about 3%, 4%, 5%, 6%, 7%, 8%, 9%, 10%, 11%, 12%, 13%, 14%, 15%, 16%, 17%, 18%, 19% or 20% of the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells express TIM-3.

In some embodiments, the threshold level is met if at least about 3%, 4%, 5%, 6%, 7%, 8%, 9%, 10%, 11%, 12%, 13%, 14%, 15%, 16%, 17%, 18%, 19% or 20% of the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells express LAG-3.

In some embodiments, the threshold level is met if at least about 3%, 4%, 5%, 6%, 7%, 8%, 9%, 10%, 11%, 12%, 13%, 14%, 15%, 16%, 17%, 18%, 19% or 20% of the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells express CTLA-4.

In some embodiments, the threshold level is met if at least about 3%, 4%, 5%, 6%, 7%, 8%, 9%, 10%, 11%, 12%, 13%, 14%, 15%, 16%, 17%, 18%, 19% or 20% of the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells express CTLA-4.

In some embodiments, the threshold level is met if at least about 15%, 16%, 17%, 18%, 19%, 20%, 21%, 22%, 23%, 24%, 25%, 26%, 27%, 28%, 29%, 30%, 40%, 50%, 60%, 70% or 80% of the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells in the sample express an antigen-specific TCR.

In certain aspects, provided herein is a method of treating a disease or disorder in a subject in need thereof comprising administering to the subject in need thereof a sample obtained from a cell bank provided herein and/or the T cells (*e.g.*, CD4 T cells or CD8 T cells, such as CTLs) from a cell bank provided herein. In some embodiments, the disease or disorder is cancer (*e.g.*, EBV-associated cancer, such as nasopharyngeal carcinoma, NK/T cell lymphoma, EBV-associated gastric carcinoma, or EBV-associated leiomyosarcoma). In some embodiments, the subject has post-transplant lymphoproliferative disorder (PTLD). In some embodiments, the subject has post-transplant lymphoproliferative disorder (PTLD) and an immunodeficiency disorder (*e.g.*, HIV/AIDS or X-linked inhibitor Apoptosis (XIAP)). In some embodiments, the subject has an autoimmune disorder, such as multiple sclerosis, rheumatoid arthritis, systemic lupus erythematosus, celiac disease, other systemic autoimmune diseases (SAD), or inflammatory bowel disease (IBD). In some embodiments, the T cells (*e.g.*, CD4 T cells or CD8 T cells, such as CTLs) are allogeneic to the subject.

In some embodiments, the expression of IFN γ , IL-2, TNF, GzmB, GzmK, Prf, CD8, CD4, PD-1, TIM-3, LAG-3, CTLA-4 and/or antigen-specific TCR is determined by Fluorescence Activated Cell Sorting (FACS). In some embodiments, the expression of IFN γ , IL-2, TNF, GzmB, GzmK, Prf, CD8, CD4, PD-1, TIM-3, LAG-3, CTLA-4 and/or antigen-specific TCR is determined by ELISpot. In some embodiments, the T cells (*e.g.*, CD8 and/or CD4 T cells) in the sample express a TCR specific for an Epstein-Barr virus (EBV) peptide (*e.g.*, a LMP1 peptide, a LMP2A peptide or an EBNA1 peptide) presented on a MHC (*e.g.*, a class I or a class II MHC).

In some embodiments, the selected sample is administered to a subject in need thereof. In some embodiments, the T cells (*e.g.*, CD8 and/or CD4 T cells) are allogeneic to the subject (*e.g.*, the sample comprising the T cells is obtained from a cell bank). In some embodiments, the T cells (*e.g.*, CD8 and/or CD4 T cells) are autologous to the subject.

In certain aspects, provided herein is a method of generating a sample comprising T cells (*e.g.*, CD8 and/or CD4 T cells) for adoptive immunotherapy, the method comprising incubating a sample comprising T cells (*e.g.*, CD8 and/or CD4 T cells) with antigen-presenting cells (APCs) presenting a peptide antigen on a MHC (*e.g.*, class I and/or class II MHC) such that T cells expressing a TCR specific for the peptide antigen presented on the MHC proliferate and then determining the expression of IFN γ , IL-2, TNF, GzmB, GzmK, Prf, CD8, CD4, PD-1, TIM-3, LAG-3, CTLA-4 and/or antigen-specific TCR by T cells in

the sample. In some embodiments, the peptide is an EBV peptide (*e.g.*, a LMP1 peptide, a LMP2A peptide or an EBNA1 peptide).

In certain aspects, provided herein is a method of treating a disease or disorder in a subject in need thereof comprising administering to the subject in need thereof T cells from a sample selected according to a method provided herein and/or obtained from a cell bank provided herein. In some embodiments, the disease or disorder is cancer (*e.g.*, EBV-associated cancer, such as nasopharyngeal carcinoma, NK/T cell lymphoma, EBV-associated gastric carcinoma, or EBV-associated leiomyosarcoma). In some embodiments, the subject has post-transplant lymphoproliferative disorder (PTLD). In some embodiments, the subject has post-transplant lymphoproliferative disorder (PTLD) and an immunodeficiency disorder (*e.g.*, HIV/AIDS or X-linked inhibitor Apoptosis (XIAP)). In some embodiments, the subject has an autoimmune disorder, such as multiple sclerosis, rheumatoid arthritis, systemic lupus erythematosus, celiac disease, other systemic autoimmune diseases (SAD), or inflammatory bowel disease (IBD). In some embodiments, the T cells are allogeneic to the subject.

In certain aspects, provided herein is a cell bank of samples comprising T cells (*e.g.*, CD8 and/or CD4 T cells) that is enriched for samples in which at least a threshold level of lymphocytes in the sample express IFN γ , IL-2, TNF, GzmB, GzmK, Prf, CD8, CD4, PD-1, TIM-3, LAG-3, CTLA-4 and/or antigen-specific TCR. In certain embodiments, at least 1%, 2%, 3%, 4%, 5%, 6%, 7%, 8%, 9%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or 100% of the samples in the cell bank are samples comprising T cells (*e.g.*, CD8 and/or CD4 T cells) in which at least a threshold level of lymphocytes in the sample express IFN γ , IL-2, TNF, GzmB, GzmK, Prf, CD8, CD4, PD-1, TIM-3, LAG-3, CTLA-4 and/or antigen-specific TCR.

Provided herein are methods of selecting subject for adoptive immunotherapy by performing an expansion and/or inducing proliferation in T cells (*e.g.*, CD8 and/or CD4 T cells) in a sample obtained from the subject, and measuring the expression of one or more of IFN γ , IL-2, TNF, GzmB, GzmK, Prf, CD8, CD4, PD-1, TIM-3, LAG-3, CTLA-4 and/or antigen-specific TCR by the T cells. In some embodiments, the subject is selected if the expression of IFN γ , IL-2, TNF, GzmB, GzmK, Prf, CD8, CD4, PD-1, TIM-3, LAG-3, CTLA-4 and/or antigen-specific TCR is above a threshold level provided herein. In some embodiments, the subject is not selected if the expression of IFN γ , IL-2, TNF, GzmB,

GzmK, Prf, CD8, CD4, PD-1, TIM-3, LAG-3, CTLA-4 and/or antigen-specific TCR is below a threshold level provided herein. In some embodiments, the subject has a disease or disorder. In some embodiments, the disease or disorder is cancer (*e.g.*, EBV-associated cancer, such as nasopharyngeal carcinoma, NK/T cell lymphoma, EBV-associated gastric carcinoma, or EBV-associated leiomyosarcoma). In some embodiments, the subject has post-transplant lymphoproliferative disorder (PTLD). In some embodiments, the subject has post-transplant lymphoproliferative disorder (PTLD) and an immunodeficiency disorder (*e.g.*, HIV/AIDS or X-linked inhibitor Apoptosis (XIAP)). In some embodiments, the subject has an autoimmune disorder, such as multiple sclerosis, rheumatoid arthritis, systemic lupus erythematosus, celiac disease, other systemic autoimmune diseases (SAD), or inflammatory bowel disease (IBD).

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 has three panels showing that E1-LMPpoly expanded T cells are functionally competent and express immune checkpoint molecules. Panel A depicts percentage of HLA-multimer positive CD8-positive lymphocytes that have express CD107a, IFN- γ , IL-2 and/or TNF. Panel B depicts percentage of HLA-multimer positive CD8-positive lymphocytes that express granzyme B (GzmB), granzyme K (GzmK) and/or perforin (Prf). Panel C depicts percentage of HLA-multimer positive CD8-positive lymphocytes that express PD-1, TIM-3, LAG-3 and CTLA-4.

Figure 2 has two panels and shows the T cell phenotype of CTL compositions administered to patients who had minimal residual disease (N/MRD) and patients who had active-recurrent/metastatic disease (ARMD). ARMD patients either achieved stable disease (SD) or continued to show progressive disease (PD) after CTL immunotherapy. Panel A shows the percentage of CD8-positive T cells in the CTL immunotherapy administered, while panel B shows the total number of LMP/EBNA-1-specific T cells in the CTL immunotherapy administered.

Figure 3 has two panels and shows the T cell phenotype of CTL compositions administered to patients who had minimal residual disease (N/MRD) and patients who had active-recurrent/metastatic disease (ARMD). ARMD patients either achieved stable disease (SD) or continued to show progressive disease (PD) CTL immunotherapy. Panel A shows the percentage of granzyme B positive (GzmB+) granzyme K positive (GzmK+) and

perforin positive (PRF+) lymphocytes in the CTL immunotherapy administered. Panel B shows the percentage of PD-1 positive, TIM-3 positive, LAG-3 positive and CTLA-4 positive lymphocytes in the CTL immunotherapy administered.

5 **Figure 4** shows the percentage of CD8 T cells expressing granzyme B (GzmB), granzyme K (GzmK) and/or perforin (Prf).

Figure 5 shows the percentage of LMP/EBNA1-specific T cells in different categories of responders versus non-responders.

Figure 6 shows that responses to adoptive immunotherapy correlate to EBV reactivity as measured based on CD107A, IFN γ , IL-2 and TNF expression.

10 **Figure 7** shows percentage of total lymphocytes expressing CD107a, IFN- γ , IL-2 and/or TNF in responders versus non-responders.

Figure 8 shows percentage of CD8 T cells expressing CD107a, IFN- γ , IL-2 and/or TNF in responders versus non-responders.

15 **Figure 9** shows percentage of CD8 T cells expressing CD107a, IFN- γ , IL-2 and TNF in responders versus non-responders.

DETAILED DESCRIPTION

Definitions

For convenience, certain terms employed in the specification, examples, and appended claims are collected here.

20 The articles “a” and “an” are used herein to refer to one or to more than one (*i.e.*, to at least one) of the grammatical object of the article. By way of example, “an element” means one element or more than one element.

 In some embodiments, the CTL composition further comprises an adjuvant. As used herein, the term “*adjuvant*” broadly refers to an agent that affects an immunological or
25 physiological response in a patient or subject. For example, an adjuvant might increase the presence of an antigen over time or to an area of interest like a tumor, help absorb an antigen-presenting cell antigen, activate macrophages and lymphocytes and support the production of cytokines. By changing an immune response, an adjuvant might permit a
30 smaller dose of an immune interacting agent to increase the effectiveness or safety of a particular dose of the immune interacting agent. For example, an adjuvant might prevent T cell exhaustion and thus increase the effectiveness or safety of a particular immune

interacting agent. Examples of adjuvants include, but are not limited to, an immune modulatory protein, Adjuvant 65, α -GalCer, aluminum phosphate, aluminum hydroxide, calcium phosphate, β -Glucan Peptide, CpG DNA, GPI-0100, lipid A, lipopolysaccharide, Lipovant, Montanide, N-acetyl-muramyl-L-alanyl-D-isoglutamine, Pam3CSK4, quil A and
5 trehalose dimycolate.

As used herein, the term "*administering*" means providing a pharmaceutical agent or composition to a subject, and includes, but is not limited to, administering by a medical professional and self-administering. Such an agent can contain, for example, peptide described herein, an antigen-presenting cell provided herein and/or a CTL provided herein.

10 The term "*biological sample*," "*tissue sample*," or simply "*sample*" each refers to a collection of cells obtained from a tissue of a subject. The source of the tissue sample may be solid tissue, as from a fresh, frozen and/or preserved organ, tissue sample, biopsy, or aspirate; blood or any blood constituents, serum, blood; bodily fluids such as cerebral spinal fluid, amniotic fluid, peritoneal fluid or interstitial fluid, urine, saliva, stool, tears; or cells
15 from any time in gestation or development of the subject.

The term "*binding*" or "*interacting*" refers to an association, which may be a stable association, between two molecules, *e.g.*, between a T cell receptor (TCR) and a peptide/MHC, due to, for example, electrostatic, hydrophobic, ionic and/or hydrogen-bond interactions under physiological conditions.

20 As used herein, the term "*cancer*" includes, but is not limited to, solid tumors and blood borne tumors. The term cancer includes diseases of the skin, tissues, organs, bone, cartilage, blood and vessels. The term "*cancer*" further encompasses primary and metastatic cancers.

The term "*epitope*" means a protein determinant capable of specific binding to an
25 antibody. Epitopes usually consist of chemically active surface groupings of molecules such as amino acids or sugar side chains. Certain epitopes can be defined by a particular sequence of amino acids to which a T cell receptor or antibody is capable of binding.

As used herein, the phrase "*pharmaceutically acceptable*" refers to those agents, compounds, materials, compositions, and/or dosage forms which are, within the scope of sound medical
30 judgment, suitable for use in contact with the tissues of human beings and animals without excessive toxicity, irritation, allergic response, or other problem or complication, commensurate with a reasonable benefit/risk ratio.

As used herein, the phrase "*pharmaceutically-acceptable carrier*" means a pharmaceutically-acceptable material, composition or vehicle, such as a liquid or solid filler, diluent, excipient, or solvent encapsulating material, involved in carrying or transporting an agent from one organ, or portion of the body, to another organ, or portion of the body. Each carrier must be "acceptable" in the sense of being compatible with the other ingredients of the formulation and not injurious to the patient. Some examples of materials which can serve as pharmaceutically-acceptable carriers include: (1) sugars, such as lactose, glucose and sucrose; (2) starches, such as corn starch and potato starch; (3) cellulose, and its derivatives, such as sodium carboxymethyl cellulose, ethyl cellulose and cellulose acetate; (4) powdered tragacanth; (5) malt; (6) gelatin; (7) talc; (8) excipients, such as cocoa butter and suppository waxes; (9) oils, such as peanut oil, cottonseed oil, safflower oil, sesame oil, olive oil, corn oil and soybean oil; (10) glycols, such as propylene glycol; (11) polyols, such as glycerin, sorbitol, mannitol and polyethylene glycol; (12) esters, such as ethyl oleate and ethyl laurate; (13) agar; (14) buffering agents, such as magnesium hydroxide and aluminum hydroxide; (15) alginic acid; (16) pyrogen-free water; (17) isotonic saline; (18) Ringer's solution; (19) ethyl alcohol; (20) pH buffered solutions; (21) polyesters, polycarbonates and/or polyanhydrides; and (22) other non-toxic compatible substances employed in pharmaceutical formulations.

As used herein, a therapeutic that "*prevents*" a condition refers to a compound that, when administered to a statistical sample prior to the onset of the disorder or condition, reduces the occurrence of the disorder or condition in the treated sample relative to an untreated control sample, or delays the onset or reduces the severity of one or more symptoms of the disorder or condition relative to the untreated control sample.

As used herein, the term "*subject*" means a human or non-human animal selected for treatment or therapy.

The phrases "*therapeutically-effective amount*" and "*effective amount*" as used herein means the amount of an agent which is effective for producing the desired therapeutic effect in at least a sub-population of cells in a subject at a reasonable benefit/risk ratio applicable to any medical treatment.

"*Treating*" a disease in a subject or "*treating*" a subject having a disease refers to subjecting the subject to a pharmaceutical treatment, *e.g.*, the administration of a drug, such that at least one symptom of the disease is decreased or prevented from worsening.

T cells

In certain aspects, provided herein are methods related to selecting T cells (*e.g.*, CD8 T cells, such as CTLs and/or CD4 T cells) or a sample comprising T cells (*e.g.*, CD8 T cells, such as CTLs and/or CD4 T cells) for adoptive immunotherapy, for expansion and/or for inclusion in a cell bank, *e.g.*, by determining the expression of a biomarker (*e.g.*, IFN γ , IL-2, TNF, GzmB, GzmK, Prf, CD8, CD4, PD-1, TIM-3, LAG-3, CTLA-4 and/or antigen-specific TCR) in the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells and selecting the sample for adoptive immunotherapy, expansion or inclusion into a cell bank if at least a threshold portion of the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells in the sample express one or more of the biomarkers.

In certain aspects, also provided herein are methods related to selecting a subject for adoptive immunotherapy or to provide a sample comprising T cells (*e.g.*, for inclusion in a cell bank or use in adoptive therapy) by obtaining a sample comprising T cells from the subject and determining the expression of a biomarker (*e.g.*, IFN γ , IL-2, TNF, GzmB, GzmK, Prf, CD8, CD4, PD-1, TIM-3, LAG-3, CTLA-4 and/or antigen-specific TCR) in the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells in the sample and selecting the subject for adoptive immunotherapy or to provide a sample if at least a threshold portion of the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells in the sample express the biomarker. In some embodiments, the T cell is a cytotoxic T lymphocyte (CTL).

In certain embodiments, the method includes determining the expression of CD107a in the sample and selecting the sample or subject if at least a threshold portion of the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells in the sample express CD107a. In certain embodiments, the method includes determining the expression of IFN γ in the sample and selecting the sample or subject if at least a threshold portion of the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells in the sample express IFN γ . In certain embodiments, the method includes determining the expression of IL-2 in the sample and selecting the sample or subject if at least a threshold portion of the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells in the sample express IL-2. In certain embodiments, the method includes determining the expression of TNF in the sample and selecting the sample or subject if at least a threshold portion of the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells in the sample express TNF. In certain embodiments, the method includes determining the expression of granzyme B in the sample and selecting the sample or subject if at least a

threshold portion of the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells in the sample express granzyme B. In certain embodiments, the method includes determining the expression of granzyme K in the sample and selecting the sample or subject if at least a threshold portion of the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells in the sample express granzyme K. In certain embodiments, the method includes determining the expression of perforin in the sample and selecting the sample or subject if at least a threshold portion of the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells in the sample express perforin.

In certain embodiments, the method includes determining the expression of CD107a and IFN γ in the sample and selecting the sample or subject if at least a threshold portion of the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells in the sample express CD107a and IFN γ . In certain embodiments, the method includes determining the expression of CD107a and IL-2 in the sample and selecting the sample or subject if at least a threshold portion of the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells in the sample express CD107a and IL-2. In certain embodiments, the method includes determining the expression of CD107a and TNF in the sample and selecting the sample or subject if at least a threshold portion of the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells in the sample express CD107a and TNF. In certain embodiments, the method includes determining the expression of IL-2 and TNF in the sample and selecting the sample or subject if at least a threshold portion of the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells in the sample express IL-2 and TNF. In certain embodiments, the method includes determining the expression of IFN γ and TNF in the sample and selecting the sample or subject if at least a threshold portion of the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells in the sample express IFN γ and TNF. In certain embodiments, the method includes determining the expression of IL-2 and IFN γ in the sample and selecting the sample or subject if at least a threshold portion of the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells in the sample express IL-2 and IFN γ .

In certain embodiments, the method includes determining the expression of CD107a and granzyme B in the sample and selecting the sample or subject if at least a threshold portion of the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells in the sample express CD107a and granzyme B. In certain embodiments, the method includes determining the expression of CD107a and granzyme K in the sample and selecting the sample or subject

if at least a threshold portion of the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells in the sample express CD107a and granzyme K. In certain embodiments, the method includes determining the expression of CD107a and perforin in the sample and selecting the sample or subject if at least a threshold portion of the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells in the sample express CD107a and perforin. In certain embodiments, the method includes determining the expression of IL-2 and granzyme K in the sample and selecting the sample or subject if at least a threshold portion of the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells in the sample express IL-2 and granzyme K. In certain embodiments, the method includes determining the expression of IL-2 and granzyme B in the sample and selecting the sample or subject if at least a threshold portion of the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells in the sample express IL-2 and granzyme B. In certain embodiments, the method includes determining the expression of IL-2 and perforin in the sample and selecting the sample or subject if at least a threshold portion of the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells in the sample express IL-2 and perforin. In certain embodiments, the method includes determining the expression of IFN γ and granzyme B in the sample and selecting the sample or subject if at least a threshold portion of the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells in the sample express IFN γ and granzyme B. In certain embodiments, the method includes determining the expression of IFN γ and granzyme K in the sample and selecting the sample or subject if at least a threshold portion of the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells in the sample express IFN γ and granzyme K. In certain embodiments, the method includes determining the expression of IFN γ and perforin in the sample and selecting the sample or subject if at least a threshold portion of the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells in the sample express IFN γ and perforin. In certain embodiments, the method includes determining the expression of TNF and granzyme B in the sample and selecting the sample or subject if at least a threshold portion of the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells in the sample express TNF and granzyme B. In certain embodiments, the method includes determining the expression of TNF and granzyme K in the sample and selecting the sample or subject if at least a threshold portion of the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells in the sample express TNF and granzyme K. In certain embodiments, the method includes determining the expression of TNF and perforin in the sample and selecting the sample or subject if at least a threshold portion of the total

lymphocytes, T cells, CD8 T cells and/or CD4 T cells in the sample express TNF and perforin.

In certain embodiments, the method includes determining the expression of granzyme B and granzyme K in the sample and selecting the sample or subject if at least a threshold
5 portion of the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells in the sample express granzyme B and granzyme K. In certain embodiments, the method includes determining the expression of perforin and granzyme K in the sample and selecting the sample or subject if at least a threshold portion of the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells in the sample express perforin and granzyme K. In certain embodiments,
10 the method includes determining the expression of granzyme B and perforin in the sample and selecting the sample or subject if at least a threshold portion of the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells in the sample express granzyme B and perforin.

In certain embodiments, the method includes determining the expression of CD107a, IL-2, and IFN γ in the sample and selecting the sample or subject if at least a threshold
15 portion of the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells in the sample express CD107a, IL-2, and IFN γ .

In certain embodiments, the method includes determining the expression of CD107a, TNF, and IFN γ in the sample and selecting the sample or subject if at least a threshold portion of the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells in the sample
20 express CD107a, TNF, and IFN γ .

In certain embodiments, the method includes determining the expression of IL-2, TNF, and IFN γ in the sample and selecting the sample or subject if at least a threshold portion of the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells in the sample express IL-2, TNF, and IFN γ .

25 In certain embodiments, the method includes determining the expression of CD107a, IL-2, and granzyme B in the sample and selecting the sample or subject if at least a threshold portion of the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells in the sample express CD107a, IL-2, and granzyme B.

In certain embodiments, the method includes determining the expression of CD107a,
30 IL-2, and granzyme K in the sample and selecting the sample or subject if at least a threshold portion of the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells in the sample express CD107a, IL-2, and granzyme K.

In certain embodiments, the method includes determining the expression of CD107a, IL-2, and perforin in the sample and selecting the sample or subject if at least a threshold portion of the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells in the sample express CD107a, IL-2, and perforin.

- 5 In certain embodiments, the method includes determining the expression of CD107a, TNF, and granzyme B in the sample and selecting the sample or subject if at least a threshold portion of the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells in the sample express CD107a, TNF, and granzyme B.

- In certain embodiments, the method includes determining the expression of CD107a,
10 TNF, and granzyme K in the sample and selecting the sample or subject if at least a threshold portion of the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells in the sample express CD107a, TNF, and granzyme K.

- In certain embodiments, the method includes determining the expression of CD107a, TNF, and perforin in the sample and selecting the sample or subject if at least a threshold
15 portion of the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells in the sample express CD107a, TNF, and perforin.

- In certain embodiments, the method includes determining the expression of CD107a, IFNg, and granzyme B in the sample and selecting the sample or subject if at least a threshold portion of the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells in the
20 sample express IFNg, IL-2, and granzyme B.

 In certain embodiments, the method includes determining the expression of CD107a, IFNg, and granzyme K in the sample and selecting the sample or subject if at least a threshold portion of the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells in the sample express IFNg, IL-2, and granzyme K.

- 25 In certain embodiments, the method includes determining the expression of CD107a, IFNg, and perforin in the sample and selecting the sample or subject if at least a threshold portion of the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells in the sample express IFNg, IL-2, and perforin.

- In certain embodiments, the method includes determining the expression of IL-2,
30 TNF, and granzyme B in the sample and selecting the sample or subject if at least a threshold portion of the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells in the sample express IL-2, TNF, and granzyme B.

In certain embodiments, the method includes determining the expression of IL-2, TNF, and granzyme K in the sample and selecting the sample or subject if at least a threshold portion of the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells in the sample express IL-2, TNF, and granzyme K.

- 5 In certain embodiments, the method includes determining the expression of IL-2, TNF, and perforin in the sample and selecting the sample or subject if at least a threshold portion of the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells in the sample express IFNg, TNF, and perforin.

- 10 In certain embodiments, the method includes determining the expression of IFNg, TNF, and granzyme B in the sample and selecting the sample or subject if at least a threshold portion of the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells in the sample express IFNg, TNF, and granzyme B.

- 15 In certain embodiments, the method includes determining the expression of IFNg, TNF, and granzyme K in the sample and selecting the sample or subject if at least a threshold portion of the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells in the sample express IFNg, TNF, and granzyme K.

- 20 In certain embodiments, the method includes determining the expression of IFNg, TNF, and perforin in the sample and selecting the sample or subject if at least a threshold portion of the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells in the sample express IFNg, TNF, and perforin.

In certain embodiments, the method includes determining the expression of IL-2, TNF, and IFNg in the sample and selecting the sample or subject if at least a threshold portion of the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells in the sample express IL-2, TNF, and IFNg.

- 25 In certain embodiments, the method includes determining the expression of granzyme K, granzyme B, and/or perforin in the sample and selecting the sample or subject if at least a threshold portion of the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells in the sample express granzyme K, granzyme B, and/or perforin.

- 30 In certain embodiments, the method includes determining the expression of CD107a, IL-2, TNF, and IFNg in the sample and selecting the sample or subject if at least a threshold portion of the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells in the sample express CD107a, IL-2, TNF, and IFNg.

In certain embodiments, the method includes determining the expression of CD107a, IL-2, TNF, and granzyme B in the sample and selecting the sample or subject if at least a threshold portion of the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells in the sample express CD107a, IL-2, TNF, and granzyme B.

- 5 In certain embodiments, the method includes determining the expression of CD107a, IL-2, TNF, and granzyme K in the sample and selecting the sample or subject if at least a threshold portion of the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells in the sample express CD107a, IL-2, TNF, and granzyme K.

- 10 In certain embodiments, the method includes determining the expression of CD107a, IL-2, TNF, and perforin in the sample and selecting the sample or subject if at least a threshold portion of the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells in the sample express CD107a, IL-2, TNF, and perforin.

- 15 In certain embodiments, the method includes determining the expression of CD107a, IL-2, IFNg, and granzyme B in the sample and selecting the sample or subject if at least a threshold portion of the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells in the sample express CD107a, IL-2, IFNg, and granzyme B.

- 20 In certain embodiments, the method includes determining the expression of CD107a, IL-2, IFNg, and granzyme K in the sample and selecting the sample or subject if at least a threshold portion of the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells in the sample express CD107a, IL-2, IFNg, and granzyme K.

In certain embodiments, the method includes determining the expression of CD107a, IL-2, IFNg, and perforin in the sample and selecting the sample or subject if at least a threshold portion of the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells in the sample express CD107a, IL-2, IFNg, and perforin.

- 25 In certain embodiments, the method includes determining the expression of TNF, IL-2, IFNg, and granzyme B in the sample and selecting the sample or subject if at least a threshold portion of the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells in the sample express TNF, IL-2, IFNg, and granzyme B.

- 30 In certain embodiments, the method includes determining the expression of TNF, IL-2, IFNg, and granzyme K in the sample and selecting the sample or subject if at least a threshold portion of the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells in the sample express TNF, IL-2, IFNg, and granzyme K.

In certain embodiments, the method includes determining the expression of TNF, IL-2, IFN γ , and perforin in the sample and selecting the sample or subject if at least a threshold portion of the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells in the sample express TNF, IL-2, IFN γ , and perforin.

5 In certain embodiments, the method includes determining the expression of TNF, granzyme K, granzyme B, and perforin in the sample and selecting the sample or subject if at least a threshold portion of the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells in the sample express TNF, granzyme K, granzyme B, and perforin.

In certain embodiments, the method includes determining the expression of CD107a, granzyme K, granzyme B, and perforin in the sample and selecting the sample or subject if at
10 least a threshold portion of the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells in the sample express CD107a, granzyme K, granzyme B, and perforin.

In certain embodiments, the method includes determining the expression of IFN γ , granzyme K, granzyme B, and perforin in the sample and selecting the sample or subject if at
15 least a threshold portion of the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells in the sample express IFN γ , granzyme K, granzyme B, and perforin.

In certain embodiments, the method includes determining the expression of IL-2, granzyme K, granzyme B, and perforin in the sample and selecting the sample or subject if at
20 least a threshold portion of the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells in the sample express IL-2, granzyme K, granzyme B, and perforin.

In certain embodiments, the method includes determining the expression of TNF, CD107a, granzyme K, granzyme B, and perforin in the sample and selecting the sample or subject if at least a threshold portion of the total lymphocytes, T cells, CD8 T cells and/or
CD4 T cells in the sample express TNF, CD107a, granzyme K, granzyme B, and perforin.

25 In certain embodiments, the method includes determining the expression of TNF, IFN γ , granzyme K, granzyme B, and perforin in the sample and selecting the sample or subject if at least a threshold portion of the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells in the sample express TNF, IFN γ , granzyme K, granzyme B, and perforin.

In certain embodiments, the method includes determining the expression of TNF, IL-2, granzyme K, granzyme B, and perforin in the sample and selecting the sample or subject if
30 at least a threshold portion of the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells in the sample express TNF, IL-2, granzyme K, granzyme B, and perforin.

In certain embodiments, the method includes determining the expression of IL-2, CD107a, granzyme K, granzyme B, and perforin in the sample and selecting the sample or subject if at least a threshold portion of the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells in the sample express IL-2, CD107a, granzyme K, granzyme B, and perforin.

5 In certain embodiments, the method includes determining the expression of IFNg, CD107a, granzyme K, granzyme B, and perforin in the sample and selecting the sample or subject if at least a threshold portion of the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells in the sample express IFNg, CD107a, granzyme K, granzyme B, and perforin.

In certain embodiments, the method includes determining the expression of IFNg, IL-2, granzyme K, granzyme B, and perforin in the sample and selecting the sample or subject if
10 at least a threshold portion of the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells in the sample express IFNg, IL-2, granzyme K, granzyme B, and perforin.

In certain embodiments, the method includes determining the expression of IL-2, CD107a, TNF, granzyme B, and perforin in the sample and selecting the sample or subject if
15 at least a threshold portion of the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells in the sample express IL-2, CD107a, TNF, granzyme B, and perforin.

In certain embodiments, the method includes determining the expression of IFNg, CD107a, TNF, granzyme B, and perforin in the sample and selecting the sample or subject if
20 at least a threshold portion of the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells in the sample express IFNg, CD107a, TNF, granzyme B, and perforin.

In certain embodiments, the method includes determining the expression of IL-2, CD107a, TNF, granzyme K, and perforin in the sample and selecting the sample or subject if
at least a threshold portion of the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells in the sample express IL-2, CD107a, TNF, granzyme K, and perforin.

25 In certain embodiments, the method includes determining the expression of IFNg, CD107a, TNF, granzyme K, and perforin in the sample and selecting the sample or subject if
at least a threshold portion of the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells in the sample express IFNg, CD107a, TNF, granzyme K, and perforin.

In certain embodiments, the method includes determining the expression of IL-2, CD107a, TNF, granzyme K, and perforin in the sample and selecting the sample or subject if
30 at least a threshold portion of the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells in the sample express IL-2, CD107a, TNF, granzyme K, and granzyme B.

In certain embodiments, the method includes determining the expression of IFN γ , CD107a, TNF, granzyme K, and perforin in the sample and selecting the sample or subject if at least a threshold portion of the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells in the sample express IFN γ , CD107a, TNF, granzyme K, and granzyme B.

- 5 In certain embodiments, the method includes determining the expression of IL-2, IFN γ , TNF, granzyme K, and granzyme B in the sample and selecting the sample or subject if at least a threshold portion of the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells in the sample express IL-2, IFN γ , TNF, granzyme K, and granzyme B.

- 10 In certain embodiments, the method includes determining the expression of IFN γ , CD107a, TNF, granzyme K, granzyme B, and perforin in the sample and selecting the sample or subject if at least a threshold portion of the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells in the sample express IFN γ , CD107a, TNF, granzyme K, granzyme B, and perforin.

- 15 In certain embodiments, the method includes determining the expression of IL-2, CD107a, TNF, granzyme K, granzyme B, and perforin in the sample and selecting the sample or subject if at least a threshold portion of the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells in the sample express IL-2, CD107a, TNF, granzyme K, granzyme B, and perforin.

- 20 In certain embodiments, the method includes determining the expression of IL-2, IFN γ , TNF, granzyme K, granzyme B, and perforin in the sample and selecting the sample or subject if at least a threshold portion of the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells in the sample express IL-2, IFN γ , TNF, granzyme K, granzyme B, and perforin.

- 25 In certain embodiments, the method includes determining the expression of IL-2, CD107a, TNF, IFN γ , granzyme B, and perforin in the sample and selecting the sample or subject if at least a threshold portion of the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells in the sample express IL-2, CD107a, TNF, IFN γ , granzyme B, and perforin.

- 30 In certain embodiments, the method includes determining the expression of IL-2, CD107a, TNF, IFN γ , granzyme K, and perforin in the sample and selecting the sample or subject if at least a threshold portion of the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells in the sample express IL-2, CD107a, TNF, IFN γ , granzyme K, and perforin.

In certain embodiments, the method includes determining the expression of IL-2, CD107a, TNF, IFN γ , granzyme K, and granzyme B in the sample and selecting the sample

or subject if at least a threshold portion of the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells in the sample express IL-2, CD107a, TNF, IFN γ , granzyme K, and granzyme B.

In certain embodiments, the method includes determining the expression of IL-2, CD107a, TNF, IFN γ , granzyme K, perforin, and granzyme B in the sample and selecting the sample or subject if at least a threshold portion of the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells in the sample express IL-2, CD107a, TNF, IFN γ , perforin, granzyme K, and granzyme B.

In certain embodiments, the threshold level of the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells expressing CD107a in a sample is met if at least 0.5%, 0.6%, 0.7%, 0.8%, 0.9%, 1.0%, 2%, 3%, 4%, 5%, 6%, 7%, 8%, 9%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or 100% of the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells in the sample express CD107a.

In certain embodiments, the threshold level of the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells expressing IFN γ in a sample is met if at least 0.5%, 0.6%, 0.7%, 0.8%, 0.9%, 1.0%, 2%, 3%, 4%, 5%, 6%, 7%, 8%, 9%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or 100% of the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells in the sample express IFN γ .

In certain embodiments, the threshold level of the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells expressing IL-2 in a sample is met if at least 0.5%, 0.6%, 0.7%, 0.8%, 0.9%, 1.0%, 2%, 3%, 4%, 5%, 6%, 7%, 8%, 9%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or 100% of the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells in the sample express IL-2.

In certain embodiments, the threshold level of the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells expressing TNF in a sample is met if at least 0.5%, 0.6%, 0.7%, 0.8%, 0.9%, 1.0%, 2%, 3%, 4%, 5%, 6%, 7%, 8%, 9%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or 100% of the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells in the sample express TNF.

In some embodiments, the threshold level of the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells expressing CD107a in a sample is not met if less than 15%, 14%, 13%, 12%, 11%, 10%, 9%, 8%, 7%, 6%, 5%, 4%, 3%, 2%, 1%, 0.9%, 0.8%, 0.7%, 0.6%, 0.5%, 0.4%, 0.3%, 0.2%, or 0.1% of the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells in the sample express CD107a.

In some embodiments, the threshold level of the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells expressing IFN γ in a sample is not met if less than 15%, 14%, 13%, 12%, 11%, 10%, 9%, 8%, 7%, 6%, 5%, 4%, 3%, 2%, 1%, 0.9%, 0.8%, 0.7%, 0.6%, 0.5%, 0.4%, 0.3%, 0.2%, or 0.1% of the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells in the sample express IFN γ .

In some embodiments, the threshold level of the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells expressing IL-2 in a sample is not met if less than 15%, 14%, 13%, 12%, 11%, 10%, 9%, 8%, 7%, 6%, 5%, 4%, 3%, 2%, 1%, 0.9%, 0.8%, 0.7%, 0.6%, 0.5%, 0.4%, 0.3%, 0.2%, or 0.1% of the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells in the sample express IL-2.

In some embodiments, the threshold level of the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells expressing TNF in a sample is not met if less than 15%, 14%, 13%, 12%, 11%, 10%, 9%, 8%, 7%, 6%, 5%, 4%, 3%, 2%, 1%, 0.9%, 0.8%, 0.7%, 0.6%, 0.5%, 0.4%, 0.3%, 0.2%, or 0.1% of the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells in the sample express TNF. In some embodiments, the threshold level of the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells expressing granzyme B in a sample is met if at least 5%, 6%, 7%, 8%, 9%, 10%, 11%, 12%, 13%, 14%, 15%, 16%, 17%, 18%, 19%, 20%, 21%, 22%, 23%, 24% or 25% of the lymphocytes in the sample express granzyme B.

In some embodiments, the threshold level of the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells expressing granzyme K in a sample is met if at least 3%, 4%, 5%, 6%, 7%, 8%, 9%, 10%, 11%, 12%, 13%, 14%, 15%, 16%, 17%, 18%, 19% or 20% of the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells in the sample express granzyme K.

In some embodiments, the threshold level of the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells expressing perforin in a sample is met if at least 2%, 3%, 4%, 5%,

6%, 7%, 8%, 9%, 10%, 11%, 12%, 13%, 14% or 15% of the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells in the sample express perforin.

In some embodiments, the threshold level of the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells expressing granzyme B in a sample is not met if less than 5%, 6%,
5 7%, 8%, 9%, 10%, 11%, 12%, 13%, 14%, 15%, 16%, 17%, 18%, 19%, 20%, 21%, 22%, 23%, 24% or 25% of the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells in the sample express granzyme B.

In some embodiments, the threshold level of the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells expressing granzyme K in a sample is not met if less than 3%, 4%,
10 5%, 6%, 7%, 8%, 9%, 10%, 11%, 12%, 13%, 14%, 15%, 16%, 17%, 18%, 19% or 20% of the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells in the sample express granzyme K.

In some embodiments, the threshold level of the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells expressing perforin in a sample is not met if less than 2%, 3%, 4%,
15 5%, 6%, 7%, 8%, 9%, 10%, 11%, 12%, 13%, 14% or 15% of the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells in the sample express perforin.

In some embodiments, the method further comprises determining the expression of CD8, CD4, PD-1, TIM-3, LAG-3 and/or CTLA-4 in the sample and selecting the sample or subject if at least a threshold portion of the total lymphocytes, T cells, CD8 T cells and/or
20 CD4 T cells in the sample express granzyme B, granzyme K, perforin, CD8, CD4, PD-1, TIM-3, LAG-3 and/or CTLA-4.

In some embodiments, the threshold level of the total lymphocytes expressing CD8 in a sample is met if at least 15%, 16%, 17%, 18%, 19%, 20%, 21%, 22%, 23%, 24%, 25%,
26%, 27%, 28%, 29% or 30% of the lymphocytes in the sample express CD8.

25 In certain embodiments, the threshold level of total lymphocytes expressing CD4 in a sample is met if at least 0.5%, 0.6%, 0.7%, 0.8%, 0.9%, 1.0%, 2%, 3%, 4%, 5%, 6%, 7%, 8%, 9%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or 100% of the lymphocytes in the sample express CD4.

30 In some embodiments, the threshold level of the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells expressing PD-1 in a sample is met if at least 3%, 4%, 5%, 6%, 7%,

8%, 9%, 10%, 11%, 12%, 13%, 14%, 15%, 16%, 17%, 18%, 19% or 20% of the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells in the sample express PD-1.

In some embodiments, the threshold level of the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells expressing TIM-3 in a sample is met if at least 3%, 4%, 5%, 6%,
5 7%, 8%, 9%, 10%, 11%, 12%, 13%, 14%, 15%, 16%, 17%, 18%, 19% or 20% of the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells in the sample express TIM-3.

In some embodiments, the threshold level of the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells expressing LAG-3 in a sample is met if at least 3%, 4%, 5%, 6%,
7%, 8%, 9%, 10%, 11%, 12%, 13%, 14%, 15%, 16%, 17%, 18%, 19% or 20% of the total
10 lymphocytes, T cells, CD8 T cells and/or CD4 T cells in the sample express LAG-3.

In some embodiments, the threshold level of the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells expressing CTLA4 in a sample is met if at least 3%, 4%, 5%, 6%,
7%, 8%, 9%, 10%, 11%, 12%, 13%, 14%, 15%, 16%, 17%, 18%, 19% or 20% of the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells in the sample express CTLA-4.

15 In some embodiments, the threshold level of the total lymphocytes expressing CD8 in a sample is not met if less than 15%, 16%, 17%, 18%, 19%, 20%, 21%, 22%, 23%, 24%,
25%, 26%, 27%, 28%, 29% or 30% of the total lymphocytes in the sample express CD8.

In certain embodiments, the threshold level of the total lymphocytes expressing CD4 in a sample is not met if less than 1%, 2%, 3%, 4%, 5%, 6%, 7%, 8%, 9%, 10%, 11%, 12%,
20 13%, 14%, 15%, 16%, 17%, 18%, 19%, 20%, 21%, 22%, 23%, 24%, 25%, 26%, 27%, 28%,
29% or 30% of the total lymphocytes in the sample express CD4.

In some embodiments, the threshold level of the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells expressing PD-1 in a sample is not met if less than 3%, 4%, 5%,
6%, 7%, 8%, 9%, 10%, 11%, 12%, 13%, 14%, 15%, 16%, 17%, 18%, 19% or 20% of the
25 total lymphocytes, T cells, CD8 T cells and/or CD4 T cells in the sample express PD-1.

In some embodiments, the threshold level of the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells expressing TIM-3 in a sample is not met if less than 3%, 4%, 5%,
6%, 7%, 8%, 9%, 10%, 11%, 12%, 13%, 14%, 15%, 16%, 17%, 18%, 19% or 20% of the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells in the sample express TIM-3.

30 In some embodiments, the threshold level of the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells expressing LAG-3 in a sample is not met if less than 3%, 4%, 5%,

6%, 7%, 8%, 9%, 10%, 11%, 12%, 13%, 14%, 15%, 16%, 17%, 18%, 19% or 20% of the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells in the sample express LAG-3.

In some embodiments, the threshold level of the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells expressing CTLA-4 in a sample is not met if less than 3%, 4%, 5%,
5 6%, 7%, 8%, 9%, 10%, 11%, 12%, 13%, 14%, 15%, 16%, 17%, 18%, 19% or 20% of the total lymphocytes, T cells, CD8 T cells and/or CD4 T cells in the sample express CTLA-4.

In some embodiments, the T cells (*e.g.*, CD8 and/or CD4 T cells) provided herein express a T cell receptor that specifically binds to a peptide presented on an MHC (*e.g.*, a class I MHC and/or a class II MHC). In some embodiments, the MHC is a class I MHC. In
10 some embodiments, the class I MHC has an α chain polypeptide that is HLA-A, HLA-B, HLA-C, HLA-E, HLA-F, HLA-g, HLA-K or HLA-L. In some embodiments, the peptide is a peptide described herein. In some embodiments, the T cells in the sample express a TCR specific for an Epstein-Barr Virus (EBV) peptide (*e.g.*, a LMP1 peptide, a LMP2A peptide or an EBNA1 peptide) presented on a class I MHC. In some embodiments, the T cells
15 express one or more EBV peptides or fragments thereof.

In some embodiments, any assay capable of detecting expression of the relevant biomarker can be used in the methods provided herein. In some embodiments, the biomarkers are detected using fluorescence activated cell sorting (FACS). In some
embodiments, the biomarkers are detected using ELISpot. In some embodiments, the
20 biomarkers are detected using microscopy (*e.g.*, fluorescence microscopy).

T cells described herein may be generated, activated (*e.g.*, prior to selection) by inducing peptide-specific T cell proliferation and/or activation by incubating a sample comprising T cells with the antigen-presenting cells (APCs), thereby inducing the T cells to proliferate. In some embodiments, the APCs that present a peptide described herein (*e.g.*, a
25 peptide comprising one or more LMP1, LMP2A, or EBNA1 epitope sequences). In some embodiments the APCs are B cells, antigen-presenting T-cells, dendritic cells, or artificial antigen-presenting cells (*e.g.*, aK562 cells).

Dendritic cells for use in the process may be prepared by taking peripheral blood mononuclear cells (PBMCs) from a patient sample and adhering them to plastic. Generally
30 the monocyte population sticks and all other cells can be washed off. The adherent population is then differentiated with IL-4 and GM-CSF to produce monocyte derived dendritic cells. These cells may be matured by the addition of IL-1 β , IL-6, PGE-1 and TNF- α

(which upregulates the important co-stimulatory molecules on the surface of the dendritic cell) and are then transduced with one or more of the peptides provided herein.

APCs that present the one or more of peptides described herein may be generated by contacting an APC with a peptide comprising a T cell epitope and/or with a nucleic acid encoding a peptide comprising a T cell epitope. In some embodiments, the APCs are irradiated. In some embodiments, the APCs that present a peptide described herein (*e.g.*, a peptide comprising one or more LMP1, LMP2A, or EBNA1 epitope sequences). A cell presenting a peptide described herein can be produced by standard techniques known in the art. For example, a cell may be pulsed to encourage peptide uptake. In some embodiments, the cells are transfected with a nucleic acid encoding a peptide provided herein. Provided herein are methods of producing antigen-presenting cells (APCs), comprising pulsing a cell with the peptides described herein. Exemplary examples of producing antigen-presenting cells can be found in WO2013088114, hereby incorporated in its entirety.

In some embodiments, the methods provided herein include steps of generating, activating and/or inducing proliferation of T cells (*e.g.*, CTLs, CD8 T cells, and/or CD4 T cells) that recognize one or more of the T cell (*e.g.*, CTL) epitopes described herein prior to selection. In some embodiments, a sample comprising T cells (*i.e.*, a PBMC sample) is incubated in culture with an APC provided herein (*e.g.*, an APC that presents a peptide comprising a T cell epitope on a class I MHC complex). In some embodiments, the APCs are autologous to the subject from whom the T cells were obtained. In some embodiments, the APCs are not autologous (*i.e.*, allogeneic) to the subject from whom the T cells were obtained. In some embodiments, the sample containing T cells are incubated 2 or more times with APCs provided herein. In some embodiments, the T cells are incubated with the APCs in the presence of at least one cytokine. In some embodiments, the cytokine is IL-4, IL-7 and/or IL-15. Exemplary methods for inducing proliferation of T cells using APCs are provided, for example, in U.S. Pat. No. 2015/0017723, which is hereby incorporated by reference.

In some aspects, provided herein are methods comprising the administration of samples and/or T cells (*e.g.*, CTLs, CD8 T cells, and/or CD4 T cells) from selected according a method provided herein to a subject in order to treat and/or prevent a disease or disorder (*e.g.*, cancer, an infectious disease and/or an autoimmune disorder). In some embodiments, the method includes administering to the subject an effective amount of the T

cells provided herein. In some embodiments, the composition includes a combination of multiple (e.g., two or more) T cells (e.g., CD4 T cells or CD8 T cells, such as CTLs) provided herein. In some embodiments, the T cells are autologous to the subject. In some embodiments, the T cells are stored in a cell bank before they are administered to the subject.

5 Peptides

In some embodiments, the methods and compositions provided herein relate to peptide-specific T cells (e.g., CTLs, CD8 T cells, and/or CD4 T cells). In some embodiments, the methods include the generation of such T cells, for example, by incubating a sample comprising T cells (i.e., a PBMC sample) with antigen-presenting cells (APCs) that
10 present one or more of the T cell epitopes described herein (e.g., APCs that present a peptide described herein comprising a CTL epitope on a class I MHC complex).

In some embodiments, the peptides provided herein comprise a sequence of any EBV viral protein (e.g., a sequence of at least 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19 or
15 20 contiguous amino acids of any EBV protein). In some embodiments, the peptides provided herein comprise no more than 25, 20, 19, 18, 17, 16, 15, 14, 13, 12, 11 or 10 contiguous amino acids of the EBV viral protein.

In some embodiments, the peptides provided herein comprise a sequence of LMP1 (e.g., a sequence of at least 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19 or 20
20 contiguous amino acids of LMP1). In some embodiments, the peptides provided herein comprise no more than 25, 20, 19, 18, 17, 16, 15, 14, 13, 12, 11 or 10 contiguous amino acids of LMP1. An exemplary LMP1 amino acid sequence is provided below (SEQ ID NO:

1):

```
1 mdldlergpp gprpprgpp lssyalall lllallfwl yiimsnwtgg allvlyafal
61 mlviiliif ifrdllcpl galclllmi tllialwnl hggalylgiv lfifgcllvf
25 121 giwvyfleil wrigatiwql lafflaflld illlialy1 qqnwwtllvd llwllflai
181 liwmyyhgqr hsdchhdds lphqqatdd ssnhdsnsn egrhhlivsg agdapplcsq
241 nlgapgggpd ngpqpdpntd dngpdpnt ddngphdplp qdpntddng pqpdpntddn
301 gphdplphnp sdsagndggp pnltcevenk ggdrpppsmt dggggdphlp tllltsgsg
361 gddddphgpv qlsyd
```

30 In some embodiments, the peptides provided herein comprise a sequence of LMP2A (e.g., a sequence of at least 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19 or 20 contiguous amino acids of LMP2A). In some embodiments, the peptides provided herein

comprise no more than 25, 20, 19, 18, 17, 16, 15, 14, 13, 12, 11 or 10 contiguous amino acids of LMP2A. An exemplary LMP2A amino acid sequence is provided below (SEQ ID NO: 2):

1 mgslemvpmg agppspgdp dgddggmsq ypsasgsdgn tptppndeer esneeppppy
 5 61 edldwngdr hsdypqlgnq dpslylgqh dgndglpppp ysprddssqh iycagrgsm
 121 npvcfpviva pylfwlaaia ascftasvst vvtatglals lllaavass yaaqrklit
 181 pvtvltavvt ffaicltwri edppfnsllf allaaagglq giyvltmvlv lilayrrwr
 241 rltvcggimf lacvvlivd avlqlspilg avtvvsmll llafvwlss pggltglgaa
 301 lllaalal laililgtln ltmfllmll wtlvvllic scsscpltki llarfilyal
 10 361 allilasali aggsilqtnf kslstefip nlfcmlliv agilfilail tewgsgnrty
 421 gpvfmcglgl ltmvagavwl tvmtntlsa wltagflif ligfalfgvi rccryccyc
 481 ltleseerpp tpyrntv

In some embodiments, the peptides provided herein comprise a sequence of EBNA1 (e.g., a sequence of at least 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19 or 20
 15 contiguous amino acids of EBNA1). In some embodiments, the peptides provided herein comprise no more than 25, 20, 19, 18, 17, 16, 15, 14, 13, 12, 11 or 10 contiguous amino acids of EBNA1. An exemplary EBNA1 amino acid sequence is provided below (SEQ ID NO: 3):

1 pffhpvgead yfeylqeggp dgepdvppga icqgpaddpg egpstgprgq gdggrtkkkg
 20 61 wfgkhrqgg snpkfeniae glrvllarsh vertteegt wvagvfvyggs ktslynlrrg
 121 talaipqcr ltpslrpfgm apgpgpqpqp lresivcyfm vflqthifae vlkdaikdlv
 181 mtkpaptcni kvvcfsddg vdlppwfppm vegaaegdd gddgdegdgd degeegqe

In some embodiments, the peptide comprises the sequence of an epitope listed in Table 1.

25 **Table 1. Exemplary EBV viral protein epitopes**

Peptide Sequence	HLA Restriction	SEQ ID No:
PYLFWLAAL	A*2301/A*2402/03	4
SSCSSCPLSKI	A*1101	5
TYGPVFMCL	A*2402	6
RRRWRLTV	B*27/02/04/05/06/09	7
LLSAWILTA	A*0203	8
LTAGFLIFL	A*0206	9

CLGGLTMTV	A*0201	10
VMSNTLLSAW	A*25/A*26	11
MSNTLLSAW	B*58	12
IEDPPFNSL	B*4001	13
YLLEMLWRL	A*02	14
YLQQNWWTL	A*02	15
ALLVLYSFA	A*02	16
IALYLQQNW	B*57/B*58	17
FLYALALLL	A*0201	18
WTLVVLLI	A*24	19
CPLSKILL	B*0801	20
HPVGEADYFEY	B*35	21
RPQKRPSCI	B*0702	22
IPQCRLTPL	B*0702	23
LSRLPFGMA	B*5701	24
YNLRRGTAL	B*0801	25
VLKDAIKDL	A*0203	26
FVYGGSKTSL	C*0303/C*0304	27
FVYGGSKTSLY	A*26	28
HPVGEADYF	B*53	29
LQTHFAEV	A*0206	30
FMVFLQTHI	A*0201	31

In some embodiments, the peptides provided herein comprise two or more of the T cell epitopes (*e.g.*, viral epitopes). In some embodiments, the peptides provided herein comprise at least 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19 or 20 T cell epitopes. For example, in some embodiments, the peptides provided herein comprise two or more of the T cell epitopes connected by linkers (*e.g.*, polypeptide linkers).

In some embodiments, the sequence of the peptides comprises a viral protein sequence except for 1 or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) conservative sequence modifications. As used herein, the term “conservative sequence modifications” is

intended to refer to amino acid modifications that do not significantly affect or alter the interaction between a T cell receptor (TCR) and a peptide containing the amino acid sequence presented on an MHC. Such conservative modifications include amino acid substitutions, additions (*e.g.*, additions of amino acids to the N or C terminus of the peptide) and deletions (*e.g.*, deletions of amino acids from the N or C terminus of the peptide).

Conservative amino acid substitutions are ones in which the amino acid residue is replaced with an amino acid residue having a similar side chain. Families of amino acid residues having similar side chains have been defined in the art. These families include amino acids with basic side chains (*e.g.*, lysine, arginine, histidine), acidic side chains (*e.g.*, aspartic acid, glutamic acid), uncharged polar side chains (*e.g.*, glycine, asparagine, glutamine, serine, threonine, tyrosine, cysteine, tryptophan), nonpolar side chains (*e.g.*, alanine, valine, leucine, isoleucine, proline, phenylalanine, methionine), beta-branched side chains (*e.g.*, threonine, valine, isoleucine) and aromatic side chains (*e.g.*, tyrosine, phenylalanine, tryptophan, histidine). Thus, one or more amino acid residues of the peptides described herein can be replaced with other amino acid residues from the same side chain family and the altered peptide can be tested for retention of TCR binding using methods known in the art. Modifications can be introduced into an antibody by standard techniques known in the art, such as site-directed mutagenesis and PCR-mediated mutagenesis.

In some embodiments, the peptides provided herein comprise a sequence that is at least 80%, 85%, 90%, 95% or 100% identical to a protein sequence (*e.g.*, the sequence of a fragment of a viral protein). To determine the percent identity of two amino acid sequences, the sequences are aligned for optimal comparison purposes (*e.g.*, gaps can be introduced in one or both of a first and a second amino acid sequence for optimal alignment and non-identical sequences can be disregarded for comparison purposes). The amino acid residues at corresponding amino acid positions are then compared. When a position in the first sequence is occupied by the same amino acid residue as the corresponding position in the second sequence, then the molecules are identical at that position. The percent identity between the two sequences is a function of the number of identical positions shared by the sequences, taking into account the number of gaps, and the length of each gap, which need to be introduced for optimal alignment of the two sequences.

In some embodiments, the peptide is chimeric or fusion peptide. As used herein, a "chimeric peptide" or "fusion peptide" comprises a peptide having a sequence provided

herein linked to a distinct peptide having sequence to which it is not linked in nature. For example, the distinct peptide can be fused to the N-terminus or C-terminus of the peptide provided herein either directly, through a peptide bond, or indirectly through a chemical linker. In some embodiments, the peptide of the provided herein is linked to another peptide
5 comprising a distinct epitopes. In some embodiments, the peptide provided herein is linked to peptides comprising epitopes from other viral and/or infectious diseases.

A chimeric or fusion peptide provided herein can be produced by standard recombinant DNA techniques. For example, DNA fragments coding for the different peptide sequences are ligated together in-frame in accordance with conventional techniques, for
10 example by employing blunt-ended or stagger-ended termini for ligation, restriction enzyme digestion to provide for appropriate termini, filling-in of cohesive ends as appropriate, alkaline phosphatase treatment to avoid undesirable joining, and enzymatic ligation. In another embodiment, the fusion gene can be synthesized by conventional techniques including automated DNA synthesizers. Alternatively, PCR amplification of gene fragments
15 can be carried out using anchor primers which give rise to complementary overhangs between two consecutive gene fragments which can subsequently be annealed and re-amplified to generate a chimeric gene sequence (see, for example, Current Protocols in Molecular Biology, Ausubel et al., eds., John Wiley & Sons: 1992). Moreover, many expression vectors are commercially available that already encode a fusion moiety.

20 The peptides provided herein can be isolated from cells or tissue sources by an appropriate purification scheme using standard protein purification techniques, and can be produced by recombinant DNA techniques, and/or can be chemically synthesized using standard peptide synthesis techniques. The peptides described herein can be produced in prokaryotic or eukaryotic host cells by expression of nucleotides encoding a peptide(s) of the
25 present invention. Alternatively, such peptides can be synthesized by chemical methods. Methods for expression of heterologous peptides in recombinant hosts, chemical synthesis of peptides, and *in vitro* translation are well known in the art and are described further in Maniatis et al., Molecular Cloning: A Laboratory Manual (1989), 2nd Ed., Cold Spring Harbor, N. Y.; Berger and Kimmel, Methods in Enzymology, Volume 152, Guide to
30 Molecular Cloning Techniques (1987), Academic Press, Inc., San Diego, Calif.; Merrifield, J. (1969) J. Am. Chem. Soc. 91:501; Chaiken I. M. (1981) CRC Crit. Rev. Biochem. 11:255; Kaiser et al. (1989) Science 243:187; Merrifield, B. (1986) Science 232:342; Kent, S. B. H.

(1988) *Annu. Rev. Biochem.* 57:957; and Offord, R. E. (1980) *Semisynthetic Proteins*, Wiley Publishing, which are incorporated herein by reference.

In certain aspects, provided herein are nucleic acid molecules encoding the peptides described herein. In some embodiments, the nucleic acid molecule is a vector. In some
5 embodiments, the nucleic acid molecule is a viral vector, such as an adenovirus based expression vector, that comprises the nucleic acid molecules described herein. In some embodiments, the vector provided herein encodes a plurality of epitopes provided herein (*e.g.*, as a polyepitope). In some embodiments, the vector provided herein encodes at least 2,
3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19 or 20 epitopes provided herein (*e.g.*,
10 epitopes provided in Table 1).

In some embodiments, the vector is AdE1-LMPpoly. The AdE1-LMPpoly vector encodes a polyepitope of defined CTL epitopes from LMP1 and LMP2 fused to a Gly-Ala repeat-depleted EBNA1 sequence. The AdE1-LMPpoly vector is described, for example, in Smith *et al.*, *Cancer Research* 72:1116 (2012); Duraiswamy *et al.*, *Cancer Research*
15 64:1483-9 (2004); and Smith *et al.*, *J. Immunol* 117:4897-906, each of which is hereby incorporated by reference.

As used herein, the term “vector,” refers to a nucleic acid molecule capable of transporting another nucleic acid to which it has been linked. One type of vector is a “plasmid”, which refers to a circular double-stranded DNA loop into which additional DNA
20 segments may be ligated. Another type of vector is a viral vector, wherein additional DNA segments may be ligated into the viral genome. Certain vectors are capable of autonomous replication in a host cell into which they are introduced (*e.g.*, bacterial vectors having a bacterial origin of replication, episomal mammalian vectors). Other vectors (*e.g.*, non-episomal mammalian vectors) can be integrated into the genome of a host cell upon
25 introduction into the host cell, and thereby be replicated along with the host genome. Moreover, certain vectors are capable of directing the expression of genes. Such vectors are referred to herein as “recombinant expression vectors” (or simply, “expression vectors”). In some embodiments, provided herein are nucleic acids operably linked to one or more regulatory sequences (*e.g.*, a promotor) in an expression vector. In some embodiments, the
30 cell transcribes the nucleic acid provided herein and thereby expresses a peptide described herein. The nucleic acid molecule can be integrated into the genome of the cell or it can be extrachromosomal.

In some embodiments, provided herein are cells that contain a nucleic acid described herein (*e.g.*, a nucleic acid encoding a peptide described herein). The cell can be, for example, prokaryotic, eukaryotic, mammalian, avian, murine and/or human. In some embodiments, the cell is a mammalian cell. In some embodiments the cell is an APC (*e.g.*, an antigen-presenting T cell, a dendritic cell, a B cell, or an aK562 cell). In the present methods, a nucleic acid described herein can be administered to the cell, for example, as nucleic acid without delivery vehicle, in combination with a delivery reagent. In some embodiments, any nucleic acid delivery method known in the art can be used in the methods described herein. Suitable delivery reagents include, but are not limited to, *e.g.*, the Mirus Transit TKO lipophilic reagent; lipofectin; lipofectamine; cellfectin; polycations (*e.g.*, polylysine), atelocollagen, nanoplexes and liposomes. In some embodiments of the methods described herein, liposomes are used to deliver a nucleic acid to a cell or subject. Liposomes suitable for use in the methods described herein can be formed from standard vesicle-forming lipids, which generally include neutral or negatively charged phospholipids and a sterol, such as cholesterol. The selection of lipids is generally guided by consideration of factors such as the desired liposome size and half-life of the liposomes in the blood stream. A variety of methods are known for preparing liposomes, for example, as described in Szoka *et al.* (1980), *Ann. Rev. Biophys. Bioeng.* 9:467; and U.S. Pat. Nos. 4,235,871, 4,501,728, 4,837,028, and 5,019,369, the entire disclosures of which are herein incorporated by reference.

Therapeutic Methods

In some embodiments, the provided herein are methods of treating a cancer, an infection or an autoimmune disorder in a subject by administering to the subject a sample selected according to a method described herein and/or peptide-specific T cells from a sample selected according to a method provided herein.

In some embodiments, the methods provided herein can be used to treat any disease or disorder (*e.g.*, cancer). Examples of cancers include some embodiments, the methods and T cells (*e.g.*, CTLs, CD8 T cells, and/or CD4 T cells) described herein may be used to treat any cancerous or pre-cancerous tumor. In some embodiments, the cancer includes a solid tumor. Cancers that may be treated by methods and compositions provided herein include, but are not limited to, cancer cells from the bladder, blood, bone, bone marrow, brain, breast,

- colon, esophagus, gastrointestinal, gum, head, kidney, liver, lung, nasopharynx, neck, ovary, prostate, skin, stomach, testis, tongue, or uterus. In addition, the cancer may specifically be of the following histological type, though it is not limited to these: neoplasm, malignant; carcinoma; carcinoma, undifferentiated; giant and spindle cell carcinoma; small cell
- 5 carcinoma; papillary carcinoma; squamous cell carcinoma; lymphoepithelial carcinoma; basal cell carcinoma; pilomatrix carcinoma; transitional cell carcinoma; papillary transitional cell carcinoma; adenocarcinoma; gastrinoma, malignant; cholangiocarcinoma; hepatocellular carcinoma; combined hepatocellular carcinoma and cholangiocarcinoma; trabecular adenocarcinoma; adenoid cystic carcinoma; adenocarcinoma in adenomatous polyp;
- 10 adenocarcinoma, familial polyposis coli; solid carcinoma; carcinoid tumor, malignant; branchiolo-alveolar adenocarcinoma; papillary adenocarcinoma; chromophobe carcinoma; acidophil carcinoma; oxyphilic adenocarcinoma; basophil carcinoma; clear cell adenocarcinoma; granular cell carcinoma; follicular adenocarcinoma; papillary and follicular adenocarcinoma; nonencapsulating sclerosing carcinoma; adrenal cortical carcinoma;
- 15 endometrioid carcinoma; skin appendage carcinoma; apocrine adenocarcinoma; sebaceous adenocarcinoma; ceruminous adenocarcinoma; mucoepidermoid carcinoma; cystadenocarcinoma; papillary cystadenocarcinoma; papillary serous cystadenocarcinoma; mucinous cystadenocarcinoma; mucinous adenocarcinoma; signet ring cell carcinoma; infiltrating duct carcinoma; medullary carcinoma; lobular carcinoma; inflammatory
- 20 carcinoma; mammary paget's disease; acinar cell carcinoma; adenosquamous carcinoma; adenocarcinoma w/squamous metaplasia; malignant thymoma; malignant ovarian stromal tumor; malignant thecoma; malignant granulosa cell tumor; and malignant robblastoma; sertoli cell carcinoma; malignant leydig cell tumor; malignant lipid cell tumor; malignant paraganglioma; malignant extra-mammary paraganglioma; pheochromocytoma;
- 25 glomangiosarcoma; malignant melanoma; amelanotic melanoma; superficial spreading melanoma; malignant melanoma in giant pigmented nevus; epithelioid cell melanoma; malignant blue nevus; sarcoma; fibrosarcoma; malignant fibrous histiocytoma; myxosarcoma; liposarcoma; leiomyosarcoma; rhabdomyosarcoma; embryonal rhabdomyosarcoma; alveolar rhabdomyosarcoma; stromal sarcoma; malignant mixed tumor;
- 30 mullerian mixed tumor; nephroblastoma; hepatoblastoma; carcinosarcoma; malignant mesenchymoma; malignant brenner tumor; malignant phyllodes tumor; synovial sarcoma; malignant mesothelioma; dysgerminoma; embryonal carcinoma; malignant teratoma;

malignant struma ovarii; choriocarcinoma; malignant mesonephroma; hemangiosarcoma;
 malignant hemangioendothelioma; kaposi's sarcoma; malignant hemangiopericytoma;
 lymphangiosarcoma; osteosarcoma; juxtacortical osteosarcoma; chondrosarcoma; malignant
 chondroblastoma; mesenchymal chondrosarcoma; giant cell tumor of bone; ewing's sarcoma;
 5 malignant odontogenic tumor; ameloblastic odontosarcoma; malignant ameloblastoma;
 ameloblastic fibrosarcoma; malignant pinealoma; chordoma; malignant glioma;
 ependymoma; astrocytoma; protoplasmic astrocytoma; fibrillary astrocytoma; astroblastoma;
 glioblastoma; oligodendroglioma; oligodendroblastoma; primitive neuroectodermal;
 cerebellar sarcoma; ganglioneuroblastoma; neuroblastoma; retinoblastoma; olfactory
 10 neurogenic tumor; malignant meningioma; neurofibrosarcoma; malignant neurilemmoma;
 malignant granular cell tumor; malignant lymphoma; Hodgkin's disease; Hodgkin's
 lymphoma; paraganuloma; small lymphocytic malignant lymphoma; diffuse large cell
 malignant lymphoma; follicular malignant lymphoma; mycosis fungoides; other specified
 non-Hodgkin's lymphomas; malignant histiocytosis; multiple myeloma; mast cell sarcoma;
 15 immunoproliferative small intestinal disease; leukemia; lymphoid leukemia; plasma cell
 leukemia; erythroleukemia; lymphosarcoma cell leukemia; myeloid leukemia; basophilic
 leukemia; eosinophilic leukemia; monocytic leukemia; mast cell leukemia; megakaryoblastic
 leukemia; myeloid sarcoma; and hairy cell leukemia.

In some embodiments, the methods provided herein are used to treat EBV-associated
 20 cancer. In some embodiments, the EBV-associated cancer is EBV-associated nasopharyngeal
 carcinoma, post-transplant lymphoproliferative disorder (PTLD), NK/T cell lymphoma,
 EBV+ gastric cancer, or EBV+ leiomyosarcoma. In some embodiments, the subject has
 PTLD and immunodeficiency disorder (*e.g.*, HIV/AIDS or X-linked inhibitor Apoptosis
 (XIAP)). In some embodiments, the subject is in remission. In some embodiments, the
 25 subject has no radiographically or molecularly detectable disease but where the subject
 remains at high risk of relapse.

In some embodiments, provided herein are methods to treat autoimmune disease
 using the T cells (*e.g.*, CD4 T cells or CD8 T cells, such as CTLs) disclosed herein.
 Examples of autoimmune diseases include, for example, glomerular nephritis, arthritis,
 30 dilated cardiomyopathy-like disease, ulcerous colitis, Sjogren syndrome, Crohn disease,
 systemic erythematodes, chronic rheumatoid arthritis, multiple sclerosis, psoriasis, allergic
 contact dermatitis, polymyositis, pachyderma, periarteritis nodosa, rheumatic fever, vitiligo

vulgaris, Behcet disease, Hashimoto disease, Addison disease, dermatomyositis, myasthenia gravis, Reiter syndrome, Graves' disease, anaemia perniciosa, sterility disease, pemphigus, autoimmune thrombopenic purpura, autoimmune hemolytic anemia, active chronic hepatitis, Addison's disease, anti-phospholipid syndrome, atopic allergy, autoimmune atrophic

5 gastritis, achlorhydra autoimmune, celiac disease, Cushing's syndrome, dermatomyositis, discoid lupus erythematosus, Goodpasture's syndrome, Hashimoto's thyroiditis, idiopathic adrenal atrophy, idiopathic thrombocytopenia, insulin-dependent diabetes, Lambert-Eaton syndrome, lupoid hepatitis, lymphopenia, mixed connective tissue disease, pemphigoid, pemphigus vulgaris, pernicious anemia, phacogenic uveitis, polyarteritis nodosa,

10 polyglandular autosyndromes, primary biliary cirrhosis, primary sclerosing cholangitis, Raynaud's syndrome, relapsing polychondritis, Schmidt's syndrome, limited scleroderma (or crest syndrome), sympathetic ophthalmia, systemic lupus erythematosus, Takayasu's arteritis, temporal arteritis, thyrotoxicosis, type b insulin resistance, ulcerative colitis and Wegener's granulomatosis.

15 In some embodiments, the methods provided herein are used to treat MS. In some embodiments, the MS is relapsing-remitting MS, secondary progressive MS, primary progressive MS or progressively relapsing MS.

In some embodiments, the methods provided herein are used to treat a SAD. For example, in certain embodiments, the methods provided herein are used to treat rheumatoid

20 arthritis, systemic lupus erythematosus and/or Sjögren's syndrome.

In some embodiments, the methods provided herein are used to treat IBD. For example, in certain embodiments the methods provided herein are used to treat Crohn's disease (regional bowel disease, *e.g.*, inactive and active forms) and/or ulcerative colitis (*e.g.*, inactive and active forms). In some embodiments, the methods provided herein are

25 used to treat irritable bowel syndrome, microscopic colitis, lymphocytic-plasmocytic enteritis, coeliac disease, collagenous colitis, lymphocytic colitis, eosinophilic enterocolitis, indeterminate colitis, infectious colitis (viral, bacterial or protozoan, *e.g.*, amoebic colitis) (*e.g.*, clostridium difficile colitis), pseudomembranous colitis (necrotizing colitis), ischemic inflammatory bowel disease, Behcet's disease, sarcoidosis, scleroderma, IBD-associated

30 dysplasia, dysplasia-associated masses or lesions, and/or primary sclerosing cholangitis.

Actual dosage levels of the active ingredients in the pharmaceutical compositions provided herein may be varied so as to obtain an amount of the active ingredient which is

effective to achieve the desired therapeutic response for a particular patient, composition, and mode of administration, without being toxic to the patient.

The selected dosage level will depend upon a variety of factors including the activity of the particular agent employed, the route of administration, the time of administration, the
5 rate of excretion or metabolism of the particular compound being employed, the duration of the treatment, other drugs, compounds and/or materials used in combination with the particular compound employed, the age, sex, weight, condition, general health and prior medical history of the patient being treated, and like factors well known in the medical arts.

In some embodiments, the subject has been exposed to a virus (*e.g.*, EBV or CMV)
10 such that virus particles are detectable in the subject's blood. In some embodiments, the method further comprises measuring viral load in the subject (*e.g.*, before or after administering the peptide-specific T cells (*e.g.*, CD4 T cells or CD8 T cells, such as CTLs) to the subject). Determining viral load in a subject may be a good prognostic marker for immunotherapy effectiveness. In some embodiments, selecting T cells (*e.g.*, CD4 T cells or
15 CD8 T cells, such as CTLs) further comprises determining the number of viral DNA copies in the subject (*e.g.*, in a tissue or blood sample). In some embodiments, viral load is measured two or more times.

In some embodiments, the method includes selecting allogeneic T cells (*e.g.*, CD4 T cells or CD8 T cells, such as CTLs) from a cell bank (*e.g.*, a pre-generated third party donor
20 derived bank of epitope-specific CTLs) for adoptive immunotherapy by determining the level expression of a biomarker within the CTL population. In some embodiments, the level of expression of two or more biomarkers is determined. In some embodiments, the method further includes selecting allogeneic T cells (*e.g.*, CD4 T cells or CD8 T cells, such as CTLs) because they express a TCR restricted to a class I MHC that is encoded by an HLA allele
25 that is present in the subject. In some embodiments, the T cells (*e.g.*, CD4 T cells or CD8 T cells, such as CTLs) are selected if the T cells (*e.g.*, CD4 T cells or CD8 T cells, such as CTLs) and subject share at least 2 (*e.g.*, at least 3, at least 4, at least 5, at least 6) HLA alleles and the CTLs are restricted through a shared HLA allele. In some embodiments, the method comprises testing the TCR repertoire of the pre-generated third-party-donor-derived epitope-
30 specific T cells (*i.e.*, allogeneic T cells) with flow cytometry. In some embodiments epitope-specific T cells are detected using a tetramer assay, an ELISA assay, a western blot assay, a fluorescent microscopy assay, an Edman degradation assay and/or a mass spectrometry assay

(e.g., protein sequencing). In some embodiments, the TCR repertoire is analyzed using a nucleic acid probe, a nucleic acid amplification assay and/or a sequencing assay.

EXEMPLIFICATION

Example 1

5 Fifty-two nasopharyngeal carcinoma (NPC) patients were enrolled in a study applying LMP1&2 and EBNA1-specific CTL immunotherapy for the treatment of EBV-associated NPC, including 41 with active progressive disease following palliative chemotherapy (active recurrent/metastatic disease, or ARMD) and 11 patients with minimal or no residual disease (N/MRD) following standard radio/chemotherapeutic treatment.

10 Twenty active disease patients and 9 N/MRD patients received the minimum 2 doses (range 2-8 doses) and a median total of 1.1×10^8 cells (range: 5.7×10^7 to 2.4×10^8). The clinical characteristics of the patients who received adoptive T cell therapy are provided in Table 1 and 2. Of the remaining 23 patients, 1 patient died after the administration of a single dose, T cell therapy was manufactured for 5 patients but not administered due to illness,

15 12 failed to meet release criteria due to low specificity or cell yield and 5 were withdrawn prior to the commencement of T cell manufacture.

 To generate LMP/EBNA1-specific T cells 100-300mL of peripheral blood was harvested and used to generate peripheral blood mononuclear cells (PBMC). The AdE1-LMPpoly vector was then used to infect 30% of the PBMC (MOI of 10:1) that were then

20 irradiated and co-cultured with the remaining PBMC for two weeks. Cultures were supplemented with fresh growth medium and 120IU/mL of recombinant IL-2 every 3-4 days (Komtur Pharmaceuticals, Friburg, Germany). Cultured T cells were tested for antigen specificity using intracellular cytokine analysis and microbial contamination prior to release for infusion.

25 Polychromatic profiling was performed to characterize the T cells administered to the subjects. MHC tetramers were generated in house. T cells were incubated for 20 minutes at 4°C with APC-labelled MHC class I tetramers specific for the HLA A11-restricted epitope SSCSSCPLSKI (LMP2A), the HLA A24 restricted epitope TYGPVFMCL (LMP2A) and the HLA Cw03 restricted epitope FVYGGSKTSL (EBNA1). For the

30 assessment of surface phenotype cells were then incubated for a further 30 minutes with the following antibodies: V500-conjugated anti-CD8, PECy7-conjugated anti-CD4 and AF700-

conjugated anti-CD3; or with V500- conjugated anti-CD8, AF700-conjugated anti-CD4, PECy7-conjugated anti-CD56, eFluor 450- conjugated anti-CD19 and -conjugated anti-CD14; or with V500-conjugated anti-CD8, PECy7- conjugated anti-CD4, Biotin-conjugated anti-CD57 followed by Streptavidin Cascade Yellow, PE-conjugated anti-CD27, perCPCy5.5-conjugated anti-CD28 , FITC-conjugated anti-CD45RA and AF700-conjugated anti-CCR7; or with V500-conjugated anti-CD8, PECy7-conjugated anti- CD4, PE-conjugated anti-TIM3, FITC conjugated anti-LAG3 and BV786-conjugated anti-PD-1. For intracellular analysis, cells were treated with BD TF Fixation/Permeabilisation buffer and then stained in the presence Perm/Wash with the following antibodies: BV421-conjugated anti-Perforin, AF700-conjugated anti-Granzyme B and FITC-conjugated anti-Granzyme K, or BV421-conjugated anti-CTLA-4. Cells were acquired using a BD LSR Fortessa with FACSDiva software (BD Biosciences) and post-acquisition analysis was performed using FlowJo software (TreeStar). As seen in Figure 1, the E1-LMPpoly expanded T cells were functionally competent.

As seen in Figure 1, the level of expression of GzmB, GzmK, Prf, PD-1, TIM-3, LAG-3 and/or CTLA4 can be determined in T cells prior to use in adoptive immunotherapy.

ARMD patients were divided into two groups: patients who achieved stable disease (referred to as SD) and patients who continued to show progressive disease (referred to as PD) after CTL immunotherapy. Circles represent individual patients. As seen in Figure 2, ARMD patients who received CTL compositions with a greater percentage of CD8⁺ T cells and/or a greater number of LMP/EBNA-1-specific T cells were more likely to have disease stabilization.. Similarly, as seen in Figure 3, ARMD patients who received CTL compositions with greater GzmB, GzmK, Prf, PD-1, TIM-3, LAG-3 and/or CTLA4 expression were also more likely to have disease stabilization.

Example 2

Autologous EBV-specific T cell therapy was used to treat patients with progressive MS. Each patient received their own T cells stimulated *ex vivo* to enhance reactivity to EBNA1, LMP1 and LMP2A and their clinical response was monitored. Following *ex vivo* stimulation, and prior to administration, the T cell samples were assessed for expression of CD107a, interferon gamma (IFN γ), interleukin 2 (IL-2), tumor necrosis factor (TNF), granzyme B (GzmB), granzyme K (GzmK) and perforin (Prf) by FACs. Four escalating dose

infusions were administered biweekly. Outcome measures were as follows: vital signs; changes in neurological symptoms; neurological examination, including EDSS; cognitive assessment; fatigue assessment; screening for depression; quality of life (QOL) assessment; blood testing; MRI of the brain and spinal cord with gadolinium contrast; and CSF analysis of intrathecal IgG production.

As seen in Figure 4, responders tended to have higher percentages of CD8 T cells expressing GzmB, GzmK and Prf compared to non-responders. Additional data reflecting the phenotypic characterization of the adoptively transferred T cells is provided in Table 2.

Table 2: Percent of total lymphocytes that express CD107a, IFNg, IL-2 and TNF (ND = no data)

Patient	Responder?	CD107a	IFNg	IL-2	TNF
1	Yes	0.0%	0.0%	0.0%	0.0%
2	No	0.1%	0.1%	0.0%	0.0%
3	Yes	8.9%	8.5%	5.1%	9.5%
4	Yes	2.5%	2.3%	1.4%	2.4%
5	Yes	26.4%	23.1%	13.7%	24.3%
6	No	0.1%	0.1%	0.0%	0.1%
7	ND	0.0%	0.0%	ND	0.0%
8	ND	0.0%	0.0%	ND	0.0%
9	ND	14.3%	13.2%	ND	12.7%

The percentage of LMP/EBNA1-specific T cells in samples based on patient response and expression of combinations of CD107(a), IFNg, IL-2 and TNF is provided in Figure 5 and Figure 9. As can be seen, in these figures, a positive patient response was highly correlated with expression of all four genes by LMP/EBNA 1-specific T cells. Notably, patient response correlated with EBV reactivity of the administered T cells, as measured by CD107a, IFNg, IL-2 and TNF expression (Figures 6 – 8). Additional data reflecting the phenotypic characterization of the adoptively transferred T cells is provided in Table 3.

Table 3: Percent of CD8 T cells that express CD107a, IFN γ , IL-2 and TNF (ND = no data)

Patient	Responder?	CD107a	IFN γ	IL-2	TNF
1	Yes	0.0%	0.0%	0.0%	0.0%
2	No	0.8%	0.7%	0.2%	0.4%
3	Yes	23.2%	22.3%	12.0%	23.8%
4	Yes	13.3%	14.2%	8.6%	14.6%
5	Yes	51.7%	45.5%	27.0%	47.8%
6	No	0.4%	0.3%	0.2%	0.4%
7	ND	0.0%	0.0%	ND	0.1%
8	ND	0.1%	0.1%	ND	0.0%
9	ND	32.4%	29.8%	ND	28.8%

5 All publications, patents, patent applications and sequence accession numbers mentioned herein are hereby incorporated by reference in their entirety as if each individual publication, patent or patent application was specifically and individually indicated to be incorporated by reference. In case of conflict, the present application, including any definitions herein, will control.

10 Those skilled in the art will recognize, or be able to ascertain using no more than routine experimentation, many equivalents to the specific embodiments of the invention described herein. Such equivalents are intended to be encompassed by the following claims.

What is claimed is:

1. A method of selecting a sample comprising T cells for adoptive immunotherapy, the method comprising determining the expression of granzyme B by lymphocytes in the sample and selecting the sample for adoptive immunotherapy if at least 5% of the lymphocytes in
5 the sample express granzyme B.
2. A method of selecting a sample comprising T cells for inclusion in a cell bank, the method comprising determining the expression of granzyme B by lymphocytes in the sample and selecting the sample for inclusion in a cell bank if at least 5% of the lymphocytes in the sample express granzyme B.
- 10 3. A method of selecting for adoptive immunotherapy a sample comprising T cells from a library of samples, the method comprising screening the library of samples for the expression of granzyme B by lymphocytes in the samples and selecting a sample from the library of samples for adoptive immunotherapy in which at least 5% of the lymphocytes in the sample express granzyme B.
- 15 4. A method of selecting for inclusion in a cell bank samples comprising T cells from a library of samples, the method comprising screening the library of samples for the expression of granzyme B by lymphocytes in the samples and selecting samples for inclusion in a cell bank in which at least 5% of the lymphocytes in the sample express granzyme B.
5. The method of any one of claims 1 to 4, wherein the sample is selected if at least
20 10% of the lymphocytes in the sample express granzyme B.
6. The method of any one of claims 1 to 4, wherein the sample is selected if at least 15% of the lymphocytes in the sample express granzyme B.
7. The method of any one of claims 1 to 4, wherein the sample is selected if at least 20% of the lymphocytes in the sample express granzyme B.
- 25 8. The method of any one of claims 1 to 7, wherein the expression of granzyme B is determined by Fluorescence Activated Cell Sorting (FACS).
9. The method of any one of claims 1 to 8, further comprising determining the expression of granzyme K by lymphocytes in the sample and selecting the sample if at least 3% of the lymphocytes in the sample express granzyme K.

10. The method of claim 9, wherein the sample is selected if at least 5% of the lymphocytes in the sample express granzyme K.
11. The method of claim 9, wherein the sample is selected if at least 8% of the lymphocytes in the sample express granzyme K.
- 5 12. The method of claim 9, wherein the sample is selected if at least 10% of the lymphocytes in the sample express granzyme K.
13. The method of claim 9, wherein the sample is selected if at least 15% of the lymphocytes in the sample express granzyme K.
14. The method of any one of claims 9 to 13, wherein the expression of granzyme K is
10 determined by Fluorescence Activated Cell Sorting (FACS).
15. The method of any one of claims 1 to 14, further comprising determining the expression of perforin by lymphocytes in the sample and selecting the sample if at least 2% of the lymphocytes in the sample express perforin.
16. The method of claim 15, wherein the sample is selected if at least 5% of the
15 lymphocytes in the sample express perforin.
17. The method of claim 15, wherein the sample is selected if at least 8% of the lymphocytes in the sample express perforin.
18. The method of claim 15, wherein the sample is selected if at least 10% of the lymphocytes in the sample express perforin.
- 20 19. The method of any one of claims 15 to 18, wherein the expression of perforin is determined by Fluorescence Activated Cell Sorting (FACS).
20. The method of any one of claims 1 to 19, wherein the T cells are incubated with antigen-presenting cells to induct proliferation of the T cells before the expression of granzyme B by the lymphocytes is determined.
- 25 21. The method of any one of claims 9 to 20, wherein the T cells are incubated with antigen-presenting cells to induct proliferation of the T cells before the expression of granzyme K by the lymphocytes is determined.

22. The method of any one of claims 15 to 21, wherein the T cells are incubated with antigen-presenting cells to induce proliferation of the T cells before the expression of perforin by the lymphocytes is determined.
23. A method of selecting a sample comprising T cells for adoptive immunotherapy, the
5 method comprising determining the expression of granzyme K by lymphocytes in the sample and selecting the sample for adoptive immunotherapy if at least 3% of the lymphocytes in the sample express granzyme K.
24. A method of selecting a sample comprising T cells for inclusion in a cell bank, the method comprising determining the expression of granzyme K by lymphocytes in the sample
10 and selecting the sample for inclusion in a cell bank if at least 3% of the lymphocytes in the sample express granzyme K.
25. A method of selecting for adoptive immunotherapy a sample comprising T cells from a library of samples, the method comprising screening the library of samples for the expression of granzyme K by lymphocytes in the samples and selecting a sample from the
15 library of samples for adoptive immunotherapy in which at least 3% of the lymphocytes in the sample express granzyme K.
26. A method of selecting for inclusion in a cell bank samples comprising T cells from a library of samples, the method comprising screening the library of samples for the expression of granzyme K in the lymphocytes in the sample and selecting samples for
20 inclusion in a cell bank in which at least 3% of the lymphocytes in the sample express granzyme K.
27. The method of any one of claims 23 to 26, wherein the sample is selected if at least 5% of the lymphocytes in the sample express granzyme K.
28. The method of any one of claims 23 to 26, wherein the sample is selected if at least
25 8% of the lymphocytes in the sample express granzyme K.
29. The method of any one of claims 23 to 26, wherein the sample is selected if at least 10% of the lymphocytes in the sample express granzyme K.
30. The method of any one of claims 23 to 26, wherein the sample is selected if at least 15% of the lymphocytes in the sample express granzyme K.

31. The method of any one of claims 23 to 30, wherein the expression of granzyme K is determined by Fluorescence Activated Cell Sorting (FACS).
32. The method of any one of claims 23 to 31, further comprising determining the expression of perforin by lymphocytes in the sample and selecting the sample if at least 2%
5 of the lymphocytes in the sample express perforin.
33. The method of claim 32, wherein the sample is selected if at least 5% of the lymphocytes in the sample express perforin.
34. The method of claim 32, wherein the sample is selected if at least 8% of the lymphocytes in the sample express perforin.
- 10 35. The method of claim 32, wherein the sample is selected if at least 10% of the lymphocytes in the sample express perforin.
36. The method of any one of claims 23 to 35, wherein the expression of perforin is determined by Fluorescence Activated Cell Sorting (FACS).
37. The method of any one of claims 23 to 36, wherein the T cells are incubated with
15 antigen-presenting cells to induct proliferation of the T cells before the expression of granzyme K by the lymphocytes is determined.
38. The method of any one of claims 32 to 37, wherein the T cells are incubated with antigen-presenting cells to induct proliferation of the T cells before the expression of perforin by the lymphocytes is determined.
- 20 39. A method of selecting a sample comprising T cells for adoptive immunotherapy, the method comprising determining the expression of perforin by lymphocytes in the sample and selecting the sample for adoptive immunotherapy if at least 2% of the lymphocytes in the sample express perforin.
40. A method of selecting a sample comprising T cells for inclusion in a cell bank, the
25 method comprising determining the expression of perforin by lymphocytes in the sample and selecting the sample for inclusion in a cell bank if at least 2% of the lymphocytes in the sample express perforin.
41. A method of selecting for adoptive immunotherapy a sample comprising T cells from a library of samples, the method comprising screening the library of samples for the

expression of perforin by lymphocytes in the samples and selecting a sample from the library for adoptive immunotherapy in which at least 2% of the lymphocytes in the sample express perforin.

42. A method of selecting for inclusion in a cell bank samples comprising T cells from a library of samples, the method comprising screening the library of samples for the expression of perforin by lymphocytes in the samples and selecting samples for inclusion in a cell bank in which at least 2% of the lymphocytes in the sample express perforin.

43. The method of any one of claims 39 to 42, wherein the sample is selected if at least 5% of the lymphocytes in the sample express perforin.

44. The method of any one of claims 39 to 42, wherein the sample is selected if at least 8% of the lymphocytes in the sample express perforin.

45. The method of any one of claims 39 to 42, wherein the sample is selected if at least 10% of the lymphocytes in the sample express perforin.

46. The method of any one of claims 39 to 45, wherein the expression of perforin is determined by Fluorescence Activated Cell Sorting (FACS).

47. The method of any one of claims 39 to 46, wherein the T cells are incubated with antigen-presenting cells to induce proliferation of the T cells before the expression of perforin by the lymphocytes is determined.

48. The method of any one of claims 1 to 47, further comprising determining the expression of CD8 by the lymphocytes in the sample and selecting the sample if at least 20% of the lymphocytes express CD8.

49. The method of claim 48, wherein the sample is selected if at least 25% of the lymphocytes in the sample express CD8.

50. The method of claim 48, wherein the sample is selected if at least 30% of the lymphocytes in the sample express CD8.

51. The method of any one of claims 1 to 50, further comprising determining the expression of PD-1 by the lymphocytes in the sample and selecting the sample if at least 5% of the lymphocytes express PD-1.

52. The method of claim 51, wherein the sample is selected if at least 7% of the lymphocytes in the sample express PD-1.
53. The method of claim 51, wherein the sample is selected if at least 10% of the lymphocytes in the sample express PD-1.
- 5 54. The method of any one of claims 1 to 53, further comprising determining the expression of TIM-3 by the lymphocytes in the sample and selecting the sample if at least 8% of the lymphocytes express TIM-3.
55. The method of claim 54, wherein the sample is selected if at least 10% of the lymphocytes in the sample express TIM-3.
- 10 56. The method of claim 54, wherein the sample is selected if at least 12% of the lymphocytes in the sample express TIM-3.
57. The method of any one of claims 1 to 56, further comprising determining the expression of LAG-3 by the lymphocytes in the sample and selecting the sample if at least 3% of the lymphocytes express LAG-3.
- 15 58. The method of claim 57, wherein the sample is selected if at least 5% of the lymphocytes in the sample express LAG-3.
59. The method of claim 57, wherein the sample is selected if at least 8% of the lymphocytes in the sample express LAG-3.
60. The method of any one of claims 1 to 59, further comprising determining the expression of CTLA-4 by the lymphocytes in the sample and selecting the sample if at least 8% of the lymphocytes express CTLA-4.
- 20 61. The method of claim 60, wherein the sample is selected if at least 10% of the lymphocytes in the sample express CTLA-4.
62. The method of claim 60, wherein the sample is selected if at least 12% of the lymphocytes in the sample express CTLA-4.
- 25 63. The method of any one of claims 1 to 62, wherein the selected T cells are stored in a cell bank after selection.

64. The method of any one of claims 1 to 63, wherein the T cells express a TCR specific for an Epstein-Barr Virus peptide presented on a class I MHC.
65. The method of claim 64, wherein the EBV peptide comprises a LMP1 peptide.
66. The method of claim 65, wherein the EBV peptide comprises a LMP2A peptide.
- 5 67. The method of claim 64 wherein the EBV peptide comprises an EBNA1 peptide.
68. The method of any one of claims 1 to 67, further comprising administration of the selected T cells are administered to a subject in need thereof.
69. The method of claim 68, wherein the subject in need thereof has cancer.
70. The method of claim 69, wherein the cancer is an Epstein-Barr Virus-associated
10 cancer.
71. The method of claim 69 or claim 70, wherein the subject has nasopharyngeal carcinoma, NK/T cell lymphoma, Epstein-Barr Virus-associated gastric carcinoma, or Epstein-Barr Virus-associated leiomyosarcoma.
72. The method of claim 68, wherein the subject in need thereof has post-transplant
15 lymphoproliferative disorder.
73. The method of claim 68, wherein the subject has an autoimmune disorder.
74. The method of claim 73, wherein the autoimmune disorder is multiple sclerosis.
75. The method of any one of claims 68 to 74, wherein the T cells are allogeneic to the subject.
- 20 76. The method of any one of claims 68 to 75, wherein the T cells are obtained from a cell bank.
77. The method of any one of claims 68 to 74, wherein the T cells are autologous to the subject.
78. A cell bank of samples comprising T cells, wherein the cell bank is enriched for
25 samples in which at least a threshold level of lymphocytes in the sample express a biomarker, wherein the threshold level of lymphocytes expressing the biomarker is:
- (a) at least 5% of lymphocytes in the sample express granzyme B;

- (b) at least 3% of the lymphocytes in the sample express granzyme K;
 - (c) at least 2% of the lymphocytes in the sample express perforin;
 - (d) at least 20% of the lymphocytes in the sample express CD8;
 - (e) at least 5% of the lymphocytes in the sample express PD-1;
 - 5 (f) at least 8% of the lymphocytes in the sample express TIM-3;
 - (g) at least 3% of the lymphocytes in the sample express LAG-3; or
 - (h) at least 8% of the lymphocytes in the sample express CTLA-4.
79. The cell bank of claim 78, wherein the cell bank is enriched for samples in which at least a threshold level of lymphocytes in the sample express two or more biomarkers.
- 10 80. The cell bank of claim 78, wherein the cell bank is enriched for samples in which at least a threshold level of lymphocytes in the sample express three or more biomarkers.
81. The cell bank of claim 78, wherein the cell bank is enriched for samples in which at least a threshold level of lymphocytes in the sample express four or more biomarkers.
82. The cell bank of any one of claims 78 to 81, wherein at least 50% of the samples in
- 15 the cell bank have at least the threshold level of lymphocytes in the sample expressing the biomarker.
83. The cell bank of any one of claims 78 to 81, wherein at least 75% of the samples in the cell bank have at least the threshold level of lymphocytes in the sample expressing the biomarker.
- 20 84. The cell bank of any one of claims 78 to 81, wherein at least 90% of the samples in the cell bank have at least the threshold level of lymphocytes in the sample expressing the biomarker.
85. The cell bank of any one of claims 78 to 81, wherein 100% of the samples in the cell bank have at least the threshold level of lymphocytes in the sample expressing the
- 25 biomarker.
86. A method of selecting a sample comprising T cells for adoptive immunotherapy, the method comprising determining the expression of CD107a by lymphocytes in the sample

and selecting the sample for adoptive immunotherapy if at least a threshold level of the lymphocytes in the sample express CD107a.

87. A method of selecting a sample comprising T cells for inclusion in a cell bank, the method comprising determining the expression of CD107a by lymphocytes in the sample
5 and selecting the sample for inclusion in a cell bank if at least a threshold level of the lymphocytes in the sample express CD107a.

88. A method of selecting for adoptive immunotherapy a sample comprising T cells from a library of samples, the method comprising screening the library of samples for the expression of CD107a by lymphocytes in the samples and selecting a sample from the
10 library of samples for adoptive immunotherapy in which at least a threshold level of the lymphocytes in the sample express CD107a.

89. A method of selecting for inclusion in a cell bank samples comprising T cells from a library of samples, the method comprising screening the library of samples for the expression of CD107a by lymphocytes in the samples and selecting samples for inclusion in
15 a cell bank in which at least a threshold level of the lymphocytes in the sample express CD107a.

90. A method of selecting a sample comprising T cells for *ex vivo* expansion, comprising:

(a) inducing proliferation in T cells from a portion of the sample,
20 (b) determining the expression of CD107a by the lymphocytes in the portion of the sample; and
(c) selecting the sample for expansion if at least a threshold level of the lymphocytes express CD107a.

91. The method of claim 90, further comprising selecting the sample for inclusion into a
25 cell bank.

92. The method of claim 90, further comprising selecting the sample for adoptive immunotherapy.

93. The method of any one of claims 86 to 92, wherein the threshold level is 0.5%.

94. The method of any one of claims 86 to 92, wherein the threshold level is 1%.

95. The method of any one of claims 86 to 92, wherein the threshold level is 2%.
96. The method of any one of claims 86 to 95, wherein the expression of CD107a is determined by Fluorescence Activated Cell Sorting (FACS) or ELISpot.
97. The method of any one of claims 86 to 96, further comprising determining the
5 expression of IFN γ by the lymphocytes and selecting the sample if at least 0.5% of the lymphocytes express IFN γ .
98. The method of claim 97, wherein the sample is selected if at least 1% of the lymphocytes express IFN γ .
99. The method of claim 97, wherein the sample is selected if at least 2% of the
10 lymphocytes express IFN γ .
100. The method of any one of claims 97 to 99, wherein the expression of IFN γ is determined by Fluorescence Activated Cell Sorting (FACS) or ELISpot.
101. The method of any one of claims 86 to 100, further comprising determining the expression of IL-2 by the lymphocytes and selecting the sample if at least 0.5% of the
15 lymphocytes express IL-2.
102. The method of claim 101, wherein the sample is selected if at least 1% of the lymphocytes express IL-2.
103. The method of claim 101, wherein the sample is selected if at least 2% of the lymphocytes express IL-2.
- 20 104. The method of claim 101, wherein the sample is selected if at least 10% of the lymphocytes in the sample express IL-2.
105. The method of any one of claims 101 to 104, wherein the expression of IL-2 is determined by Fluorescence Activated Cell Sorting (FACS) or ELISpot.
106. The method of any one of claims 86 to 105, further comprising determining the
25 expression of TNF by the lymphocytes and selecting the sample if at least 0.5% of the lymphocytes express TNF.
107. The method of claim 106, wherein the sample is selected if at least 1% of the lymphocytes express TNF.

108. The method of claim 107, wherein the sample is selected if at least 2% of the lymphocytes express TNF.
109. The method of any one of claims 106 to 108, wherein the expression of TNF is determined by Fluorescence Activated Cell Sorting (FACS) or ELISpot.
- 5 110. The method of any one of claims 86 to 109, further comprising determining the expression of granzyme B by the lymphocytes and selecting the sample if at least 5% of the lymphocytes express granzyme B.
111. The method of claim 110, wherein the sample is selected if at least 10% of the lymphocytes express granzyme B.
- 10 112. The method of claim 110, wherein the sample is selected if at least 15% of the lymphocytes express granzyme B.
113. The method of claim 110, wherein the sample is selected if at least 20% of the lymphocytes express granzyme B.
114. The method of any one of claims 110 to 113, wherein the expression of granzyme B
15 is determined by Fluorescence Activated Cell Sorting (FACS) or ELISpot.
115. The method of any one of claims 86 to 114, further comprising determining the expression of granzyme K by the lymphocytes and selecting the sample if at least 3% of the lymphocytes express granzyme K.
116. The method of claim 115, wherein the sample is selected if at least 5% of the
20 lymphocytes express granzyme K.
117. The method of claim 115, wherein the sample is selected if at least 8% of the lymphocytes express granzyme K.
118. The method of claim 115, wherein the sample is selected if at least 10% of the lymphocytes express granzyme K.
- 25 119. The method of claim 115, wherein the sample is selected if at least 15% of the lymphocytes express granzyme K.
120. The method of any one of claims 115 to 119, wherein the expression of granzyme K is determined by Fluorescence Activated Cell Sorting (FACS) or ELISpot.

121. The method of any one of claims 86 to 120, further comprising determining the expression of perforin by the lymphocytes and selecting the sample if at least 2.0% of the lymphocytes express perforin.
122. The method of claim 121, wherein the sample is selected if at least 5% of the
5 lymphocytes express perforin.
123. The method of claim 121, wherein the sample is selected if at least 8% of the lymphocytes express perforin.
124. The method of claim 121, wherein the sample is selected if at least 10% of the lymphocytes in the sample express perforin.
- 10 125. The method of any one of claims 121 to 124, wherein the expression of perforin is determined by Fluorescence Activated Cell Sorting (FACS) or ELISpot.
126. The method of any one of claims 86 to 125, wherein the T cells are incubated with antigen-presenting cells to induce proliferation of the T cells before the expression of CD107a by the lymphocytes is determined.
- 15 127. The method of any one of claims 97 to 126, wherein the T cells are incubated with antigen-presenting cells to induce proliferation of the T cells before the expression of IFN γ by the lymphocytes is determined.
128. The method of any one of claims 101 to 127, wherein the T cells are incubated with antigen-presenting cells to induce proliferation of the T cells before the expression of IL-2 by
20 the lymphocytes is determined.
129. The method of any one of claims 106 to 128, wherein the T cells are incubated with antigen-presenting cells to induce proliferation of the T cells before the expression of TNF by the lymphocytes is determined.
130. The method of any one of claims 110 to 129, wherein the T cells are incubated with
25 antigen-presenting cells to induce proliferation of the T cells before the expression of granzyme B by the lymphocytes is determined.
131. The method of any one of claims 115 to 130, wherein the T cells are incubated with antigen-presenting cells to induce proliferation of the T cells before the expression of granzyme K by the lymphocytes is determined.

132. The method of any one of claims 121 to 131, wherein the T cells are incubated with antigen-presenting cells to induce proliferation of the T cells before the expression of perforin by the lymphocytes is determined.

5 133. A method of selecting a sample comprising T cells for adoptive immunotherapy, the method comprising determining the expression of IFN γ by T cells in the sample and selecting the sample for adoptive immunotherapy if at least a threshold level of the lymphocytes in the sample express IFN γ .

10 134. A method of selecting a sample comprising T cells for inclusion in a cell bank, the method comprising determining the expression of IFN γ by lymphocytes in the sample and selecting the sample for inclusion in a cell bank if at least a threshold level of the lymphocytes in the sample express IFN γ .

15 135. A method of selecting for adoptive immunotherapy a sample comprising T cells from a library of samples, the method comprising screening the library of samples for the expression of IFN γ by lymphocytes in the samples and selecting a sample from the library of samples for adoptive immunotherapy in which at least a threshold level of the lymphocytes in the sample express IFN γ .

20 136. A method of selecting for inclusion in a cell bank samples comprising T cells from a library of samples, the method comprising screening the library of samples for the expression of IFN γ in the lymphocytes in the sample and selecting samples for inclusion in a cell bank in which at least a threshold level of the lymphocytes in the sample express IFN γ .

137. A method of selecting a sample comprising T cells for *ex vivo* expansion, comprising:

- 25 (a) inducing proliferation in T cells from a portion of the sample,
(b) determining the expression of IFN γ by the lymphocytes in the portion of the sample; and
(c) selecting the sample for further expansion if at least a threshold level of the lymphocytes express IFN γ .

138. The method of claim 137, further comprising selecting the sample for inclusion into a cell bank.

139. The method of claim 137, further comprising selecting the sample for adoptive immunotherapy.
140. The method of any one of claims 133 to 139, wherein the threshold level is 0.5%.
141. The method of any one of claims 133 to 139, wherein the threshold level is 1%.
- 5 142. The method of any one of claims 133 to 139, wherein the threshold level is 2%.
143. The method of any one of claims 133 to 139, wherein the expression of IFN γ is determined by Fluorescence Activated Cell Sorting (FACS) or ELISpot.
144. The method of any one of claims 133 to 143, further comprising determining the expression of IL-2 by the lymphocytes and selecting the sample if at least 0.5% of the
10 lymphocytes express IL-2.
145. The method of claim 144, wherein the sample is selected if at least 1% of the lymphocytes express IL-2.
146. The method of claim 144, wherein the sample is selected if at least 2% of the lymphocytes express IL-2.
- 15 147. The method of any one of claims 144 to 146, wherein the expression of IL-2 is determined by Fluorescence Activated Cell Sorting (FACS) or ELISpot.
148. The method of any one of claims 133 to 147, further comprising determining the expression of TNF by the lymphocytes and selecting the sample if at least 0.5% of the lymphocytes express TNF.
- 20 149. The method of claim 148, wherein the sample is selected if at least 1% of the lymphocytes express TNF.
150. The method of claim 148, wherein the sample is selected if at least 2% of the lymphocytes express TNF.
151. The method of any one of claims 148 to 150, wherein the expression of TNF is
25 determined by Fluorescence Activated Cell Sorting (FACS) or ELISpot.
152. The method of any one of claims 133 to 151, further comprising determining the expression of granzyme B by the lymphocytes and selecting the sample if at least 5% of the lymphocytes express granzyme B.

153. The method of claim 152, wherein the sample is selected if at least 10% of the lymphocytes express granzyme B.
154. The method of claim 152, wherein the sample is selected if at least 15% of the lymphocytes express granzyme B.
- 5 155. The method of claim 152, wherein the sample is selected if at least 20% of the lymphocytes express granzyme B.
156. The method of any one of claims 152 to 155, wherein the expression of granzyme B is determined by Fluorescence Activated Cell Sorting (FACS) or ELISpot.
157. The method of any one of claims 133 to 156, further comprising determining the
10 expression of granzyme K by the lymphocytes and selecting the sample if at least 3% of the lymphocytes express granzyme K.
158. The method of claim 157, wherein the sample is selected if at least 5% of the lymphocytes express granzyme K.
159. The method of claim 157, wherein the sample is selected if at least 8% of the
15 lymphocytes express granzyme K.
160. The method of claim 157, wherein the sample is selected if at least 10% of the lymphocytes express granzyme K.
161. The method of claim 157, wherein the sample is selected if at least 15% of the lymphocytes express granzyme K.
- 20 162. The method of any one of claims 157 to 161, wherein the expression of granzyme K is determined by Fluorescence Activated Cell Sorting (FACS) or ELISpot.
163. The method of any one of claims 133 to 162, further comprising determining the expression of perforin by the lymphocytes and selecting the sample if at least 2.0% of the lymphocytes express perforin.
- 25 164. The method of claim 163, wherein the sample is selected if at least 5% of the lymphocytes express perforin.
165. The method of claim 163, wherein the sample is selected if at least 8% of the lymphocytes express perforin.

166. The method of claim 163, wherein the sample is selected if at least 10% of the lymphocytes in the sample express perforin.
167. The method of any one of claims 163 to 166, wherein the expression of perforin is determined by Fluorescence Activated Cell Sorting (FACS) or ELISpot.
- 5 168. The method of any one of claims 133 to 167, wherein the T cells are incubated with antigen-presenting cells to induce proliferation of the T cells before the expression of IFN γ by the lymphocytes is determined.
169. The method of any one of claims 144 to 168, wherein the T cells are incubated with antigen-presenting cells to induce proliferation of the T cells before the expression of IL-2 by
10 the lymphocytes is determined.
170. The method of any one of claims 148 to 169, wherein the T cells are incubated with antigen-presenting cells to induce proliferation of the T cells before the expression of TNF by the lymphocytes is determined.
171. The method of any one of claims 152 to 170, wherein the T cells are incubated with
15 antigen-presenting cells to induce proliferation of the T cells before the expression of granzyme B by the lymphocytes is determined.
172. The method of any one of claims 157 to 171, wherein the T cells are incubated with antigen-presenting cells to induce proliferation of the T cells before the expression of granzyme K by the lymphocytes is determined.
- 20 173. The method of any one of claims 163 to 172, wherein the T cells are incubated with antigen-presenting cells to induce proliferation of the T cells before the expression of perforin by the lymphocytes is determined.
174. A method of selecting a sample comprising T cells for adoptive immunotherapy, the method comprising determining the expression of IL-2 by lymphocytes in the sample and
25 selecting the sample for adoptive immunotherapy if at least a threshold level of the lymphocytes in the sample express IL-2.
175. A method of selecting a sample comprising T cells for inclusion in a cell bank, the method comprising determining the expression of IL-2 by lymphocytes in the sample and

selecting the sample for inclusion in a cell bank if at least a threshold level of the lymphocytes in the sample express IL-2.

176. A method of selecting for adoptive immunotherapy a sample comprising T cells from a library of samples, the method comprising screening the library of samples for the expression of IL-2 by lymphocytes in the samples and selecting a sample from the library for adoptive immunotherapy in which at least a threshold level of the lymphocytes in the sample express IL-2.

177. A method of selecting for inclusion in a cell bank samples comprising T cells from a library of samples, the method comprising screening the library of samples for the expression of IL-2 by lymphocytes in the samples and selecting samples for inclusion in a cell bank in which at least a threshold level of the lymphocytes in the sample express IL-2.

178. A method of selecting a sample comprising T cells for *ex vivo* expansion, comprising:

- (a) inducing proliferation by the T cells in a portion of the sample,
- (b) determining the expression of IL-2 by the lymphocytes in the portion of the sample; and
- (c) selecting the sample for further expansion if at least a threshold level of the lymphocytes express IL-2.

179. The method of claim 178, further comprising selecting the sample for inclusion into a cell bank.

180. The method of claim 178, further comprising selecting the sample for adoptive immunotherapy.

181. The method of any one of claims 174 to 180, wherein the threshold level is 0.5%.

182. The method of any one of claims 174 to 180, wherein the threshold level is 1%.

183. The method of any one of claims 174 to 180, wherein the threshold level is 2%.

184. The method of any one of claims 174 to 183, wherein the expression of IL-2 is determined by Fluorescence Activated Cell Sorting (FACS) or ELISpot.

185. The method of any one of claims 174 to 184, further comprising determining the expression of TNF by the lymphocytes and selecting the sample if at least 0.5% of the lymphocytes express TNF.
186. The method of claim 185, wherein the sample is selected if at least 1% of the lymphocytes express TNF.
- 5 187. The method of claim 185, wherein the sample is selected if at least 2% of the lymphocytes express TNF.
188. The method of any one of claims 185 to 187, wherein the expression of TNF is determined by Fluorescence Activated Cell Sorting (FACS) or ELISpot.
- 10 189. The method of any one of claims 174 to 188, further comprising determining the expression of granzyme B by the lymphocytes and selecting the sample if at least 5% of the lymphocytes express granzyme B.
190. The method of claim 189, wherein the sample is selected if at least 10% of the lymphocytes express granzyme B.
- 15 191. The method of claim 189, wherein the sample is selected if at least 15% of the lymphocytes express granzyme B.
192. The method of claim 189, wherein the sample is selected if at least 20% of the lymphocytes express granzyme B.
193. The method of any one of claims 189 to 192, wherein the expression of granzyme B is determined by Fluorescence Activated Cell Sorting (FACS) or ELISpot.
- 20 194. The method of any one of claims 174 to 193, further comprising determining the expression of granzyme K by the lymphocytes and selecting the sample if at least 3% of the lymphocytes express granzyme K.
195. The method of claim 194, wherein the sample is selected if at least 5% of the lymphocytes express granzyme K.
- 25 196. The method of claim 194, wherein the sample is selected if at least 8% of the lymphocytes express granzyme K.

197. The method of claim 194, wherein the sample is selected if at least 10% of the lymphocytes express granzyme K.
198. The method of claim 194, wherein the sample is selected if at least 15% of the lymphocytes express granzyme K.
- 5 199. The method of any one of claims 194 to 198, wherein the expression of granzyme K is determined by Fluorescence Activated Cell Sorting (FACS) or ELISpot.
200. The method of any one of claims 174 to 199, further comprising determining the expression of perforin by the lymphocytes and selecting the sample if at least 2.0% of the lymphocytes express perforin.
- 10 201. The method of claim 200, wherein the sample is selected if at least 5% of the lymphocytes express perforin.
202. The method of claim 200, wherein the sample is selected if at least 8% of the lymphocytes express perforin.
203. The method of claim 200, wherein the sample is selected if at least 10% of the
15 lymphocytes in the sample express perforin.
204. The method of any one of claims 200 to 203, wherein the expression of perforin is determined by Fluorescence Activated Cell Sorting (FACS) or ELISpot.
205. The method of any one of claims 174 to 204, wherein the T cells are incubated with antigen-presenting cells to induct proliferation of the T cells before the expression of IL-2 by
20 the lymphocytes is determined.
206. The method of any one of claims 185 to 205, wherein the T cells are incubated with antigen-presenting cells to induct proliferation of the T cells before the expression of TNF by the lymphocytes is determined.
207. The method of any one of claims 189 to 206, wherein the T cells are incubated with
25 antigen-presenting cells to induct proliferation of the T cells before the expression of granzyme B by the lymphocytes is determined.
208. The method of any one of claims 194 to 207, wherein the T cells are incubated with antigen-presenting cells to induct proliferation of the T cells before the expression of granzyme K by the lymphocytes is determined.

209. The method of any one of claims 200 to 208, wherein the T cells are incubated with antigen-presenting cells to induct proliferation of the T cells before the expression of perforin by the lymphocytes is determined.

210. A method of selecting a sample comprising T cells for adoptive immunotherapy, the method comprising determining the expression of TNF by lymphocytes in the sample and selecting the sample for adoptive immunotherapy if at least a threshold level of the lymphocytes in the sample express TNF.

211. A method of selecting a sample comprising T cells for inclusion in a cell bank, the method comprising determining the expression of TNF by lymphocytes in the sample and selecting the sample for inclusion in a cell bank if at least a threshold level of the lymphocytes in the sample express TNF.

212. A method of selecting for adoptive immunotherapy a sample comprising T cells from a library of samples, the method comprising screening the library of samples for the expression of TNF by lymphocytes in the samples and selecting a sample from the library for adoptive immunotherapy in which at least a threshold level of the lymphocytes in the sample express TNF.

213. A method of selecting for inclusion in a cell bank samples comprising T cells from a library of samples, the method comprising screening the library of samples for the expression of TNF by lymphocytes in the samples and selecting samples for inclusion in a cell bank in which at least a threshold level of the lymphocytes in the sample express TNF.

214. A method of selecting a sample comprising T cells for *ex vivo* expansion, comprising:

- (a) inducing proliferation in T cells from a portion of the sample,
- (b) determining the expression of TNF by the lymphocytes in the portion of the sample; and
- (c) selecting the sample for expansion if at least a threshold level of the lymphocytes express TNF.

215. The method of claim 214, further comprising selecting the sample for inclusion into a cell bank.

216. The method of claim 214, further comprising selecting the sample for adoptive immunotherapy.
217. The method of any one of claims 210 to 216, wherein the threshold level is 0.5%.
218. The method of any one of claims 210 to 216, wherein the threshold level is 1%.
- 5 219. The method of any one of claims 210 to 216, wherein the threshold level is 2%.
220. The method of any one of claims 210 to 219, wherein the expression of TNF is determined by Fluorescence Activated Cell Sorting (FACS) or ELISpot.
221. The method of any one of claims 210 to 220, further comprising determining the expression of granzyme B by the lymphocytes and selecting the sample if at least 5% of the
10 lymphocytes express granzyme B.
222. The method of claim 221, wherein the sample is selected if at least 10% of the lymphocytes express granzyme B.
223. The method of claim 221, wherein the sample is selected if at least 15% of the lymphocytes express granzyme B.
- 15 224. The method of claim 221, wherein the sample is selected if at least 20% of the lymphocytes express granzyme B.
225. The method of any one of claims 221 to 224, wherein the expression of granzyme B is determined by Fluorescence Activated Cell Sorting (FACS) or ELISpot.
226. The method of any one of claims 210 to 225, further comprising determining the
20 expression of granzyme K by the lymphocytes and selecting the sample if at least 3% of the lymphocytes express granzyme K.
227. The method of claim 226, wherein the sample is selected if at least 5% of the lymphocytes express granzyme K.
228. The method of claim 226, wherein the sample is selected if at least 8% of the
25 lymphocytes express granzyme K.
229. The method of claim 226, wherein the sample is selected if at least 10% of the lymphocytes express granzyme K.

230. The method of claim 226, wherein the sample is selected if at least 15% of the lymphocytes express granzyme K.
231. The method of any one of claims 226 to 230, wherein the expression of granzyme K is determined by Fluorescence Activated Cell Sorting (FACS) or ELISpot.
- 5 232. The method of any one of claims 210 to 231, further comprising determining the expression of perforin by the lymphocytes and selecting the sample if at least 2.0% of the lymphocytes express perforin.
233. The method of claim 232, wherein the sample is selected if at least 5% of the lymphocytes express perforin.
- 10 234. The method of claim 232, wherein the sample is selected if at least 8% of the lymphocytes express perforin.
235. The method of claim 232, wherein the sample is selected if at least 10% of the lymphocytes in the sample express perforin.
236. The method of any one of claims 232 to 235, wherein the expression of perforin is
15 determined by Fluorescence Activated Cell Sorting (FACS) or ELISpot.
237. The method of any one of claims 210 to 236, wherein the T cells are incubated with antigen-presenting cells to induct proliferation of the T cells before the expression of TNF by the lymphocytes is determined.
238. The method of any one of claims 221 to 237, wherein the T cells are incubated with
20 antigen-presenting cells to induct proliferation of the T cells before the expression of granzyme B by the lymphocytes is determined.
239. The method of any one of claims 226 to 238, wherein the T cells are incubated with antigen-presenting cells to induct proliferation of the T cells before the expression of granzyme K by the lymphocytes is determined.
- 25 240. The method of any one of claims 232 to 239, wherein the T cells are incubated with antigen-presenting cells to induct proliferation of the T cells before the expression of perforin by the lymphocytes is determined.
241. A method of selecting a sample comprising T cells for adoptive immunotherapy, the method comprising determining the expression of granzyme B by lymphocytes in the sample

and selecting the sample for adoptive immunotherapy if at least a threshold level of the lymphocytes in the sample express granzyme B.

242. A method of selecting a sample comprising T cells for inclusion in a cell bank, the method comprising determining the expression of granzyme B by lymphocytes in the sample
5 and selecting the sample for inclusion in a cell bank if at least a threshold level of the lymphocytes in the sample express granzyme B.

243. A method of selecting for adoptive immunotherapy a sample comprising T cells from a library of samples, the method comprising screening the library of samples for the expression of granzyme B by lymphocytes in the samples and selecting a sample from the
10 library of samples for adoptive immunotherapy in which at least a threshold level of the lymphocytes in the sample express granzyme B.

244. A method of selecting for inclusion in a cell bank samples comprising T cells from a library of samples, the method comprising screening the library of samples for the expression of granzyme B by lymphocytes in the samples and selecting samples for inclusion
15 in a cell bank in which at least a threshold level of the lymphocytes in the sample express granzyme B.

245. A method of selecting a sample comprising T cells for *ex vivo* expansion, comprising:

- (a) inducing proliferation in T cells from a portion of the sample,
- 20 (b) determining the expression of granzyme B by the lymphocytes in the portion of the sample; and
- (c) selecting the sample for expansion if at least a threshold level of the lymphocytes express granzyme B.

246. The method of claim 245, further comprising selecting the sample for inclusion into a
25 cell bank.

247. The method of claim 245, further comprising selecting the sample for adoptive immunotherapy.

248. The method of any one of claims 241 to 247, wherein the threshold level is 5%.

249. The method of any one of claims 241 to 247, wherein the threshold level is 10%.

250. The method of any one of claims 241 to 247, wherein the threshold level is 15%.
251. The method of any one of claims 241 to 247, wherein the threshold level is 20%.
252. The method of any one of claims 241 to 251, wherein the expression of granzyme B is determined by Fluorescence Activated Cell Sorting (FACS) or ELISpot.
- 5 253. The method of any one of claims 241 to 252, further comprising determining the expression of granzyme K by the lymphocytes and selecting the sample if at least 3% of the lymphocytes express granzyme K.
254. The method of claim 253, wherein the sample is selected if at least 5% of the lymphocytes express granzyme K.
- 10 255. The method of claim 253, wherein the sample is selected if at least 8% of the lymphocytes express granzyme K.
256. The method of claim 253, wherein the sample is selected if at least 10% of the lymphocytes express granzyme K.
257. The method of claim 253, wherein the sample is selected if at least 15% of the
- 15 lymphocytes express granzyme K.
258. The method of any one of claims 241 to 257, wherein the expression of granzyme K is determined by Fluorescence Activated Cell Sorting (FACS) or ELISpot.
259. The method of any one of claims 241 to 258, further comprising determining the expression of perforin by the lymphocytes and selecting the sample if at least 2.0% of the
- 20 lymphocytes express perforin.
260. The method of claim 259, wherein the sample is selected if at least 5% of the lymphocytes express perforin.
261. The method of claim 259, wherein the sample is selected if at least 8% of the lymphocytes express perforin.
- 25 262. The method of claim 259, wherein the sample is selected if at least 10% of the lymphocytes in the sample express perforin.
263. The method of any one of claims 241 to 262, wherein the expression of perforin is determined by Fluorescence Activated Cell Sorting (FACS) or ELISpot.

264. The method of any one of claims 241 to 263, wherein the T cells are incubated with antigen-presenting cells to induct proliferation of the T cells before the expression of granzyme B by the lymphocytes is determined.
265. The method of any one of claims 253 to 264, wherein the T cells are incubated with antigen-presenting cells to induct proliferation of the T cells before the expression of granzyme K by the lymphocytes is determined.
266. The method of any one of claims 259 to 265, wherein the T cells are incubated with antigen-presenting cells to induct proliferation of the T cells before the expression of perforin by the lymphocytes is determined.
- 10 267. A method of selecting a sample comprising T cells for adoptive immunotherapy, the method comprising determining the expression of granzyme K by lymphocytes in the sample and selecting the sample for adoptive immunotherapy if at least a threshold level of the lymphocytes in the sample express granzyme K.
268. A method of selecting a sample comprising T cells for inclusion in a cell bank, the method comprising determining the expression of granzyme K by lymphocytes in the sample and selecting the sample for inclusion in a cell bank if at least a threshold level of the lymphocytes in the sample express granzyme K.
- 15 269. A method of selecting for adoptive immunotherapy a sample comprising T cells from a library of samples, the method comprising screening the library of samples for the expression of granzyme K by lymphocytes in the samples and selecting a sample from the library of samples for adoptive immunotherapy in which at least a threshold level of the lymphocytes in the sample express granzyme K.
- 20 270. A method of selecting for inclusion in a cell bank samples comprising T cells from a library of samples, the method comprising screening the library of samples for the expression of granzyme K in the lymphocytes in the sample and selecting samples for inclusion in a cell bank in which at least a threshold level of the lymphocytes in the sample express granzyme K.
- 25 271. A method of selecting a sample comprising T cells for *ex vivo* expansion, comprising:

- (a) inducing proliferation in T cells from a portion of the sample,
- (b) determining the expression of granzyme K by the lymphocytes in the portion of the sample; and
- (c) selecting the sample for further expansion if at least a threshold level of the lymphocytes express granzyme K.
- 5 272. The method of claim 271, further comprising selecting the sample for inclusion into a cell bank.
273. The method of claim 271, further comprising selecting the sample for adoptive immunotherapy.
- 10 274. The method of any one of claims 267 to 273, wherein the threshold level is 3%.
275. The method of any one of claims 267 to 273, wherein the threshold level is 5%.
276. The method of any one of claims 267 to 273, wherein the threshold level is 8%.
277. The method of any one of claims 267 to 273, wherein the threshold level is 10%.
278. The method of any one of claims 267 to 273, wherein the threshold level is 15%.
- 15 279. The method of any one of claims 267 to 278, wherein the expression of granzyme K is determined by Fluorescence Activated Cell Sorting (FACS) or ELISpot.
280. The method of any one of claims 267 to 279, further comprising determining the expression of perforin by the lymphocytes and selecting the sample if at least 2% of the lymphocytes express perforin.
- 20 281. The method of claim 280, wherein the sample is selected if at least 5% of the lymphocytes express perforin.
282. The method of claim 280, wherein the sample is selected if at least 8% of the lymphocytes express perforin.
283. The method of claim 280, wherein the sample is selected if at least 10% of the lymphocytes express perforin.
- 25 284. The method of any one of claims 280 to 283, wherein the expression of perforin is determined by Fluorescence Activated Cell Sorting (FACS) or ELISpot.

285. The method of any one of claims 267 to 284, wherein the T cells are incubated with antigen-presenting cells to induct proliferation of the T cells before the expression of granzyme K by the lymphocytes is determined.

5 286. The method of any one of claims 280 to 285, wherein the T cells are incubated with antigen-presenting cells to induct proliferation of the T cells before the expression of perforin by the lymphocytes is determined.

287. A method of selecting a sample comprising T cells for adoptive immunotherapy, the method comprising determining the expression of perforin by lymphocytes in the sample and selecting the sample for adoptive immunotherapy if at least a threshold level of the
10 lymphocytes in the sample express perforin.

288. A method of selecting a sample comprising T cells for inclusion in a cell bank, the method comprising determining the expression of perforin by lymphocytes in the sample and selecting the sample for inclusion in a cell bank if at least a threshold level of the lymphocytes in the sample express perforin.

15 289. A method of selecting for adoptive immunotherapy a sample comprising T cells from a library of samples, the method comprising screening the library of samples for the expression of perforin by lymphocytes in the samples and selecting a sample from the library for adoptive immunotherapy in which at least a threshold level of the lymphocytes in the sample express perforin.

20 290. A method of selecting for inclusion in a cell bank samples comprising T cells from a library of samples, the method comprising screening the library of samples for the expression of perforin by lymphocytes in the samples and selecting samples for inclusion in a cell bank in which at least a threshold level of the lymphocytes in the sample express perforin.

25 291. A method of selecting a sample comprising T cells for *ex vivo* expansion, comprising:

(a) inducing proliferation by T cells from a portion of the sample,

(b) determining the expression of perforin by the lymphocytes in the portion of the sample; and

(c) selecting the sample for further expansion if at least a threshold level of the lymphocytes express perforin.

292. The method of claim 291, further comprising selecting the sample for inclusion into a cell bank.

5 293. The method of claim 291, further comprising selecting the sample for adoptive immunotherapy.

294. The method of any one of claims 287 to 293, wherein the threshold level is 2%.

295. The method of any one of claims 287 to 293, wherein the threshold level is 5%.

296. The method of any one of claims 287 to 293, wherein the threshold level is 8%.

10 297. The method of any one of claims 287 to 293, wherein the threshold level is 10%.

298. The method of any one of claims 287 to 297, wherein the expression of perforin is determined by Fluorescence Activated Cell Sorting (FACS) or ELISpot.

15 299. The method of any one of claims 287 to 298, wherein the T cells are incubated with antigen-presenting cells to induct proliferation of the T cells before the expression of perforin by the lymphocytes is determined.

300. The method of any one of claims 86 to 299, further comprising determining the expression of CD8 by cells in the sample and selecting the sample if at least 20% of the cells express CD8.

20 301. The method of claim 300, wherein the sample is selected if at least 25% of the cells express CD8.

302. The method of claim 300, wherein the sample is selected if at least 30% of the cells express CD8.

25 303. The method of any one of claims 86 to 302, further comprising determining the expression of CD8 by cells in the sample and selecting the sample if at least 10% of the cells express CD4.

304. The method of claim 303, wherein the sample is selected if at least 15% of the cells in the sample express CD4.

305. The method of claim 303, wherein the sample is selected if at least 25% of the cells in the sample express CD4.

306. The method of any one of claims 86 to 305, further comprising determining the expression of PD-1 by the lymphocytes and selecting the sample if at least 5% of the lymphocytes express PD-1.

307. The method of claim 306, wherein the sample is selected if at least 7% of the lymphocytes express PD-1.

308. The method of claim 306, wherein the sample is selected if at least 10% of the lymphocytes express PD-1.

309. The method of any one of claims 86 to 308, further comprising determining the expression of TIM-3 by the lymphocytes and selecting the sample if at least 8% of the lymphocytes express TIM-3.

310. The method of claim 309, wherein the sample is selected if at least 10% of the lymphocytes express TIM-3.

311. The method of claim 309, wherein the sample is selected if at least 12% of the lymphocytes express TIM-3.

312. The method of any one of claims 86 to 311, further comprising determining the expression of LAG-3 by the lymphocytes and selecting the sample if at least 3% of the lymphocytes express LAG-3.

313. The method of claim 312, wherein the sample is selected if at least 5% of the lymphocytes express LAG-3.

314. The method of claim 312, wherein the sample is selected if at least 8% of the lymphocytes express LAG-3.

315. The method of any one of claims 86 to 314, further comprising determining the expression of CTLA-4 by the lymphocytes and selecting the sample if at least 8% of the lymphocytes express CTLA-4.

316. The method of claim 315, wherein the sample is selected if at least 10% of the lymphocytes express CTLA-4.

317. The method of claim 315, wherein the sample is selected if at least 12% of the lymphocytes express CTLA-4.
318. The method of any one of claims 86 to 317, wherein the selected T cells are stored in a cell bank after selection.
- 5 319. The method of any one of claims 86 to 318, wherein the T cells express a TCR specific for an Epstein-Barr Virus peptide presented on a class I MHC.
320. The method of claim 319, wherein the EBV peptide comprises a LMP1 peptide.
321. The method of claim 319, wherein the EBV peptide comprises a LMP2A peptide.
322. The method of claim 319, wherein the EBV peptide comprises an EBNA1 peptide.
- 10 323. The method of any one of claims 86 to 322, further comprising administration of the selected T cells are administered to a subject in need thereof.
324. The method of claim 323, wherein the subject in need thereof has cancer.
325. The method of claim 324, wherein the cancer is an Epstein-Barr Virus-associated cancer.
- 15 326. The method of claim 324 or 325, wherein the subject has nasopharyngeal carcinoma, NK/T cell lymphoma, Epstein-Barr Virus-associated gastric carcinoma, or Epstein-Barr Virus-associated leiomyosarcoma.
327. The method of claim 326, wherein the subject in need thereof has post-transplant lymphoproliferative disorder.
- 20 328. The method of claim 327, wherein the subject has an autoimmune disorder.
329. The method of claim 328, wherein the autoimmune disorder is multiple sclerosis.
330. The method of any one of claims 318 to 329, wherein the T cells are allogeneic to the subject.
331. The method of any one of claims 318 to 330, wherein the T cells are obtained from a
25 cell bank.
332. The method of any one of claims 318 to 329, wherein the T cells are autologous to the subject.

333. A cell bank of samples comprising T cells, wherein the cell bank is enriched for samples in which at least a threshold level of lymphocytes in the sample express a biomarker, wherein the threshold level of lymphocytes expressing the biomarker is:

- (a) at least 1% of lymphocytes in the sample express CD107a;
- 5 (b) at least 1% of the lymphocytes in the sample express IFN γ ;
- (c) at least 1% of the lymphocytes in the sample express IL-2;
- (d) at least 1% of the lymphocytes in the sample express TNF;
- (e) at least 5% of lymphocytes in the sample express granzyme B;
- (f) at least 3% of the lymphocytes in the sample express granzyme K;
- 10 (h) at least 2% of the lymphocytes in the sample express perforin;
- (i) at least 20% of the lymphocytes in the sample express CD8;
- (j) at least 10% of the lymphocytes in the sample express CD4;
- (k) at least 5% of the lymphocytes in the sample express PD-1;
- (l) at least 8% of the lymphocytes in the sample express TIM-3;
- 15 (m) at least 3% of the lymphocytes in the sample express LAG-3; or
- (n) at least 8% of the lymphocytes in the sample express CTLA-4.

334. The cell bank of claim 333, wherein the cell bank is enriched for samples in which at least a threshold level of lymphocytes in the sample express two or more biomarkers.

20 335. The cell bank of claim 333, wherein the cell bank is enriched for samples in which at least a threshold level of lymphocytes in the sample express three or more biomarkers.

336. The cell bank of claim 333, wherein the cell bank is enriched for samples in which at least a threshold level of lymphocytes in the sample express four or more biomarkers.

337. The cell bank of claim 333, wherein the cell bank is enriched for samples in which at least a threshold level of lymphocytes in the sample express five or more biomarkers.

25 338. The cell bank of claim 333, wherein the cell bank is enriched for samples in which at least a threshold level of lymphocytes in the sample express six or more biomarkers.

339. The cell bank of claim 333, wherein the cell bank is enriched for samples in which at least a threshold level of lymphocytes in the sample express seven or more biomarkers.
340. The cell bank of claim 333, wherein the cell bank is enriched for samples in which at least a threshold level of lymphocytes in the sample express eight or more biomarkers.
- 5 341. The cell bank of claim 333, wherein the cell bank is enriched for samples in which at least a threshold level of lymphocytes in the sample express nine or more biomarkers.
342. The cell bank of claim 333, wherein the cell bank is enriched for samples in which at least a threshold level of lymphocytes in the sample express ten or more biomarkers.
343. The cell bank of claim 333, wherein the cell bank is enriched for samples in which at
10 least a threshold level of lymphocytes in the sample express eleven or more biomarkers.
344. The cell bank of claim 333, wherein the cell bank is enriched for samples in which at least a threshold level of lymphocytes in the sample express twelve or more biomarkers.
345. The cell bank of any one of claims 333 to 344, wherein at least 25% of the samples in the cell bank have at least the threshold level of lymphocytes in the sample expressing the
15 biomarker.
346. The cell bank of any one of claims 333 to 344, wherein at least 50% of the samples in the cell bank have at least the threshold level of lymphocytes in the sample expressing the biomarker.
347. The cell bank of any one of claims 333 to 344, wherein at least 75% of the samples in
20 the cell bank have at least the threshold level of lymphocytes in the sample expressing the biomarker.
348. The cell bank of any one of claims 333 to 344, wherein 100% of the samples in the cell bank have at least the threshold level of lymphocytes in the sample expressing the biomarker.
- 25 349. A method of selecting a subject for autologous adoptive immunotherapy, comprising:
a) inducing proliferation of the T cells obtained from the subject,
b) determining the expression of CD107a by the T cells, and

c) selecting the subject for autologous adoptive immunotherapy if at least a threshold level of the lymphocytes express CD107a.

350. The method of claim 349, wherein the T cells are from a blood sample obtained from the subject.

5 351. The method of claim 349 or claim 350, wherein the subject has cancer.

352. The method of claim 351, wherein the cancer is an Epstein-Barr Virus-associated cancer.

353. The method of claim 351 or claim 352, wherein the subject has nasopharyngeal carcinoma, NK/T cell lymphoma, Epstein-Barr Virus-associated gastric carcinoma, or
10 Epstein-Barr Virus-associated leiomyosarcoma.

354. The method of claim 349 or claim 350, wherein the subject has post-transplant lymphoproliferative disorder.

355. The method of claim 349, wherein the subject has an autoimmune disorder.

356. The method of claim 355, wherein the autoimmune disorder is multiple sclerosis.

15 357. The method of any one of claims 349 to 356, further comprising administering a sample comprising autologous T cells to the subject.

358. A method of selecting a subject as a T cell donor, comprising:

a) inducing proliferation of the T cells obtained from the subject, and

b) determining the expression of CD107a by the T cells, and

20 c) selecting the subject as a T cell donor if at least a threshold level of the T cells express CD107a.

359. The method of any one of claims 349 to 358, wherein the threshold level is 0.5%.

360. The method of any one of claims 349 to 358, wherein the threshold level is 1%.

361. The method of any one of claims 349 to 358, wherein the threshold level is 2%.

25 362. The method of any one of claims 349 to 361, wherein the subject is selected if at least 0.5% of the T cells express IFN γ .

363. The method of claim any one of claims 349 to 362, wherein the subject is selected if at least 1% of the T cells express IFNg.
364. The method of claim any one of claims 349 to 363, wherein the subject is selected if at least 2% of the T cells express IFNg.
- 5 365. The method of claim any one of claims 349 to 364, wherein the subject is selected if at least 0.5% of the T cells express IL-2.
366. The method of claim any one of claims 349 to 365, wherein the subject is selected if at least 1% of the T cells express IL-2.
367. The method of claim any one of claims 349 to 366, wherein the subject is selected if
10 at least 2% of the T cells express IL-2.
368. The method of claim any one of claims 349 to 367, wherein the subject is selected if at least 0.5% of the T cells express TNF.
369. The method of claim any one of claims 349 to 368, wherein the subject is selected if at least 1% of the T cells express TNF.
- 15 370. The method of claim any one of claims 349 to 365, wherein the subject is selected if at least 2% of the T cells express TNF.
371. The method of claim any one of claims 349 to 370, wherein the subject is selected if at least 5% of the T cells express granzyme B.
372. The method of claim any one of claims 349 to 371, wherein the subject is selected if
20 at least 10% of the T cells express granzyme B.
373. The method of claim any one of claims 349 to 372, wherein the subject is selected if at least 15% of the T cells express granzyme B.
374. The method of claim any one of claims 349 to 373, wherein the subject is selected if at least 20% of the T cells express granzyme B.
- 25 375. The method of claim any one of claims 349 to 374, wherein the subject is selected if at least 3% of the T cells express granzyme K.
376. The method of claim any one of claims 349 to 375, wherein the subject is selected if at least 5% of the T cells express granzyme K.

377. The method of claim any one of claims 349 to 376, wherein the subject is selected if at least 8% of the T cells express granzyme K.
378. The method of claim any one of claims 349 to 377, wherein the subject is selected if at least 10% of the T cells express granzyme K.
- 5 379. The method of claim any one of claims 349 to 378, wherein the subject is selected if at least 15% of the T cells express granzyme K.
380. The method of claim any one of claims 349 to 379, wherein the subject is selected if at least 2% of the T cells express perforin.
381. The method of claim any one of claims 349 to 380, wherein the subject is selected if
10 at least 5% of the T cells express perforin.
382. The method of claim any one of claims 349 to 381, wherein the subject is selected if at least 8% of the T cells express perforin.
383. The method of claim any one of claims 349 to 382, wherein the subject is selected if at least 10% of the T cells express perforin.
- 15 384. The method of any one of claims 349 to 383, wherein the expression of CD107a is determined by Fluorescence Activated Cell Sorting (FACS) or ELISpot.
385. The method of any one of claims 362 to 384, wherein the expression of IFN γ is determined by Fluorescence Activated Cell Sorting (FACS) or ELISpot.
386. The method of any one of claims 365 to 385, wherein the expression of IL-2 is
20 determined by Fluorescence Activated Cell Sorting (FACS) or ELISpot.
387. The method of any one of claims 368 to 386, wherein the expression of TNF is determined by Fluorescence Activated Cell Sorting (FACS) or ELISpot.
388. The method of any one of claims 371 to 387, wherein the expression of granzyme B is determined by Fluorescence Activated Cell Sorting (FACS) or ELISpot.
- 25 389. The method of any one of claims 375 to 388, wherein the expression of granzyme K is determined by Fluorescence Activated Cell Sorting (FACS) or ELISpot.
390. The method of any one of claims 380 to 389, wherein the expression of perforin is determined by Fluorescence Activated Cell Sorting (FACS) or ELISpot.

391. The method of any one of claims 349 to 390, further comprising determining the expression of PD-1 by the T cells and selecting the subject if at least 5% of the T cells express PD-1.
392. The method of claim 391, wherein the subject is selected if at least 7% of the T cells
5 express PD-1.
393. The method of claim 391, wherein the subject is selected if at least 10% of the T cells express PD-1.
394. The method of any one of claims 349 to 393, further comprising determining the expression of TIM-3 by the T cells and selecting the subject if at least 8% of the T cells
10 express TIM-3.
395. The method of claim 394, wherein the subject is selected if at least 10% of the T cells express TIM-3.
396. The method of claim 394, wherein the subject is selected if at least 12% of the T cells express TIM-3.
- 15 397. The method of any one of claims 349 to 396, further comprising determining the expression of LAG-3 by the T cells and selecting the subject if at least 3% of the T cells express LAG-3.
398. The method of claim 397, wherein the subject is selected if at least 5% of the T cells express LAG-3.
- 20 399. The method of claim 398, wherein the subject is selected if at least 8% of the T cells express LAG-3.
400. The method of any one of claims 349 to 399, further comprising determining the expression of CTLA-4 by the T cells and selecting the subject if at least 8% of the T cells express CTLA-4.
- 25 401. The method of claim 400, wherein the subject is selected if at least 10% of the T cells express CTLA-4.
402. The method of claim 400, wherein the subject is selected if at least 12% of the T cells express CTLA-4.

403. The method of any one of claims 1 to 77, 86-332 and 349 to 402, wherein the T cells are CD8 T cells.
404. The method of claim 403, wherein the CD8 T cells are cytotoxic T lymphocytes (CTLs).
- 5 405. The method of claim 404, wherein the CTLs express IFN γ , TNF, IL-2 and CD107.
406. The method of any one of claims 1 to 77, 86-332 and 349 to 402, wherein the T cells are CD4 T cells.
407. The cell bank of any one of claims 78-85 and 333-348, wherein the T cells are CD8 T cells.
- 10 408. The cell bank of claim 407, wherein the CD8 T cells are cytotoxic T lymphocytes (CTLs).
409. The cell bank of claim 408, wherein the CTLs express IFN γ , TNF, IL-2 and CD107.
410. The cell bank of any one of claims 78-85 and 333-348, wherein the T cells are CD4 T cells.