

- 1 -

CELL-FREE SYSTEM FOR CONVERTING METHANE INTO FUEL, PYRUVATE OR ISOBUTANOL

RELATED APPLICATIONS

5 This application claims priority under 35 U.S.C. § 119(e) to U.S. provisional application, U.S.S.N. 61/740,972, filed December 21, 2012, which is incorporated by reference herein in its entirety.

BACKGROUND OF INVENTION

10 Isobutanol is produced naturally during the fermentation of carbohydrates and may also be a byproduct of the decay process of organic matter (Atsumi *et al.* 2008 *Nature* 451:86–89). The biosynthetic pathway used to produce isobutanol was first discovered in species of bacteria from the genus *Clostridium*. This pathway has been genetically engineered into several species of microorganisms which are more easily manipulated by current scientific
15 methods than microorganisms of the genus *Clostridium* (Peralta-Yahya *et al.* 2012 *Nature* 488: 320–328).

SUMMARY OF INVENTION

 The present invention provides a cell-free system for the conversion of natural gas, in
20 particular, methane, to target biofuels (*e.g.*, isobutanol) and other chemical compounds. Natural gas is a hydrocarbon gas mixture of primarily methane, but also includes varying amounts of other higher alkanes and a lesser percentage of carbon dioxide, nitrogen, and hydrogen sulfide. Aspects of the invention uniquely combine biosynthetic and chemical engineering processes to provide a cell-free biosynthetic system capable of producing from
25 natural gas greater than, for example, 10 g to 25 g biofuel per liter-hr. For example, an industrial or commercial plant sized to use a 155 m³ reactor can produce greater than 30,000 metric tons of biofuel (*e.g.*, isobutanol) per year (*e.g.*, >500 BPD scale), in some instances with raw material costs of less than 20% of the product's value.

 The cell-free process provided herein, in some embodiments, combines (i) a cell-free
30 lysate from methane-metabolizing methanotrophs (*e.g.*, *Methylococcus capsulatus* Bath) that express enzymes (*e.g.*, methane monooxygenase, MMO) for the conversion of methane to methanol or formaldehyde with (ii) a cell-free lysate(s) from recombinant bacterial cells (*e.g.*,

- 2 -

Escherichia coli) that express enzymes for the conversion of methanol or formaldehyde to biofuel or other chemical compound. The cell lysates are combined with methane and oxygen, both required for MMO activation, in a pressurized bioreactor (*e.g.*, at least 1 bar) to drive the transformation of methane to biofuel or other chemical compound. The cell-free process of the present disclosure typically uses a multiphase reaction system that is particularly useful for delivering methane and oxygen to exposed lipid bilayer membranes (*e.g.*, of intracellular membrane vesicles of the cell lysate(s)) by first diffusing the gases into an aqueous phase. In some embodiments, enzymes for the conversion of methane to formaldehyde and/or for the conversion of formaldehyde to biofuel or other chemical compound are added exogenously to the cell-free system. Such exogenously added enzymes may be purified or partially purified.

It should be understood that while the present disclosure describes the use of methanotrophs for the expression/production of enzymes for the conversion of methane to formaldehyde, other organisms for use in producing enzymes for the conversion of methane to formaldehyde are contemplated herein.

In some aspects, provided herein are cell-free methods for large-scale conversion of methane to a biofuel or other chemical compound comprising: combining, in a bioreactor at elevated pressure, one or more cell lysates containing enzyme(s) that catalyze the conversion of methane to formaldehyde, and enzymes that catalyze the conversion of formaldehyde to a biofuel or another chemical compound, methane, and oxygen to form a cell-free reaction mixture; and incubating under suitable conditions (*e.g.*, high pressure, 37 °C) the cell-free reaction to convert methane to a biofuel or other chemical compound. In some embodiments, the bioreactor may further comprise an organic solvent (*e.g.*, decane). In some embodiments, the organic phase is saturated or supersaturated with methane. "Suitable conditions" include, without limitation, a pressure of at least 1 bar (*e.g.*, 1 bar to 10 bar, or 5 bar to 10 bar) and/or a temperature of 35 °C to 40 °C (*e.g.*, 37 °C).

In some embodiments, provided herein are cell-free methods for large-scale conversion of methane to isobutanol comprising: combining, in a bioreactor at elevated pressure, one or more cell lysates containing methane monooxygenase and methanol dehydrogenase (*e.g.*, a recombinant NAD-linked methanol dehydrogenase), which catalyze the conversion of methane to formaldehyde, and one or more of hexulose-6-phosphate synthase, 6-phospho-3-hexuloisomerase, 6-phosphofructokinase, fructose biphosphate aldolase, triose phosphate isomerase, transketolase, ribose-5-phosphate isomerase, ribose-5-phosphate 3-epimerase,

- 3 -

glyceraldehyde 3-phosphate dehydrogenase, phosphoglycerate kinase, phosphoglycerate mutase, enolase, pyruvate kinase, acetolactate synthase, acetohydroxy acid isomeroreductase, dihydroxy acid dehydratase, alpha-ketoisovalerate decarboxylase, and isobutanol dehydrogenase, which catalyze the conversion of formaldehyde to isobutanol, methane, and oxygen to form a cell-free reaction mixture; and incubating under suitable conditions the cell-free reaction to convert methane to isobutanol. In some embodiments, the bioreactor may further comprise an organic solvent (*e.g.*, decane).

One of the intermediate products of the conversion formaldehyde to biofuel or other chemical compound is pyruvate. Thus, in some aspects, provided herein are cell-free methods for large-scale conversion of methane to pyruvate comprising: combining, in a bioreactor at elevated pressure, one or more cell lysates containing enzyme(s) that catalyze the conversion of methane to formaldehyde, and enzymes that catalyze the conversion of formaldehyde to pyruvate, methane, and oxygen to form a cell-free reaction mixture; and incubating under suitable conditions the cell-free reaction to convert methane to pyruvate. In some embodiments, the bioreactor may further comprise an organic solvent (*e.g.*, decane).

In some embodiments, provided herein are cell-free methods for large-scale conversion of methane to pyruvate comprising: combining, in a bioreactor at elevated pressure, one or more cell lysates containing methane monooxygenase and methanol dehydrogenase (*e.g.*, a recombinant NAD-linked methanol dehydrogenase), which catalyze the conversion of methane to formaldehyde, and one or more of hexulose-6-phosphate synthase, 6-phospho-3-hexuloisomerase, 6-phosphofructokinase, fructose biphosphate aldolase, triose phosphate isomerase, transketolase, ribose-5-phosphate isomerase, ribose-5-phosphate 3-epimerase, glyceraldehyde 3-phosphate dehydrogenase, phosphoglycerate kinase, phosphoglycerate mutase, enolase, and pyruvate kinase, which catalyze the conversion of formaldehyde to pyruvate, methane, and oxygen to form a cell-free reaction mixture; and incubating under suitable conditions the cell-free reaction to convert methane to pyruvate. In some embodiments, the bioreactor may further comprise an organic solvent (*e.g.*, decane).

In other aspects, provided herein are cell-free systems and compositions for large-scale conversion of methane to a biofuel or other chemical compound comprising a bioreactor that comprises: a gas phase comprising methane and oxygen; and an aqueous phase comprising a cell lysate containing enzyme(s) that catalyze the conversion of methane to formaldehyde and enzymes that catalyze the conversion of formaldehyde to a biofuel or chemical compound. In

- 4 -

some embodiments, the bioreactor further comprises an organic phase comprising an organic solvent (*e.g.*, decane).

In some embodiments, provided herein are cell-free systems and compositions for large-scale conversion of methane to isobutanol comprising a bioreactor that comprises: a gas phase comprising methane and oxygen; and an aqueous phase comprising one or more cell lysates containing methane monooxygenase, and methanol dehydrogenase (*e.g.*, a recombinant NAD-linked methanol dehydrogenase), which catalyze the conversion of methane to formaldehyde and hexulose-6-phosphate synthase, 6-phospho-3-hexuloisomerase, 6-phosphofructokinase, fructose biphosphate aldolase, triose phosphate isomerase, transketolase, ribose-5-phosphate isomerase, ribose-5-phosphate 3-epimerase, glyceraldehyde 3-phosphate dehydrogenase, phosphoglycerate kinase, phosphoglycerate mutase, enolase, pyruvate kinase, acetolactate synthase, acetohydroxy acid isomeroreductase, dihydroxy acid dehydratase, alpha-ketoisovalerate decarboxylase, and isobutanol dehydrogenase, which catalyze the conversion of formaldehyde to isobutanol. In some embodiments, the bioreactor further comprises an organic phase comprising an organic solvent (*e.g.*, decane).

In some embodiments, provided herein are cell-free systems and compositions for large-scale conversion of methane to pyruvate comprising a bioreactor that comprises: a gas phase comprising methane and oxygen; and an aqueous phase comprising one or more cell lysates containing methane monooxygenase and methanol dehydrogenase (*e.g.*, a recombinant NAD-linked methanol dehydrogenase), which catalyze the conversion of methane to formaldehyde, and hexulose-6-phosphate synthase, 6-phospho-3-hexuloisomerase, 6-phosphofructokinase, fructose biphosphate aldolase, triose phosphate isomerase, transketolase, ribose-5-phosphate isomerase, ribose-5-phosphate 3-epimerase, glyceraldehyde 3-phosphate dehydrogenase, phosphoglycerate kinase, phosphoglycerate mutase, enolase, and pyruvate kinase, which catalyze the conversion of formaldehyde to pyruvate. In some embodiments, the bioreactor further comprises an organic phase comprising an organic solvent (*e.g.*, decane).

In some aspects, provided herein are cell-free methods for large-scale conversion of methane to pyruvate, comprising: (a) expressing in methanotrophs one or more enzymes for the conversion of methane to formaldehyde, (b) expressing in recombinant bacteria one or more enzymes for the conversion of formaldehyde to pyruvate, (c) combining cell lysate from the methanotrophs and the recombinant bacteria with methane, an organic solvent (*e.g.*, decane), and oxygen to form a cell-free reaction mixture, and (d) incubating under suitable

- 5 -

conditions the cell-free reaction to convert methane to pyruvate. In some embodiments, the one or more enzymes for the conversion of methane to formaldehyde are selected from methane monooxygenase and methanol dehydrogenase. In some embodiments, the one or more enzymes for the conversion of formaldehyde to pyruvate are selected from hexulose-6-phosphate synthase, 6-phospho-3-hexuloisomerase, 6-phosphofructokinase, fructose biphosphate aldolase, triose phosphate isomerase, transketolase, ribose-5-phosphate isomerase, ribose-5-phosphate 3-epimerase, glyceraldehyde 3-phosphate dehydrogenase, phosphoglycerate kinase, phosphoglycerate mutase, enolase, and pyruvate kinase.

In some embodiments, the method further comprises lysing the methanotroph and/or recombinant bacterial cells (*e.g.*, prior to the combining step).

In some embodiments, the method further comprises expressing in the recombinant bacterial cells a protease that cleaves and inactivates an enzyme. For example, the protease may cleave and inactivate pyruvate dehydrogenase and/or formaldehyde dehydrogenase. Alternatively, the protease may be added exogenously.

Also provided herein are cell lysates that comprise one or more enzymes for the conversion of methane to formaldehyde, as well as cell lysates comprising one or more enzymes for the conversion of formaldehyde to a biofuel or other chemical compound. In some embodiments, the cell lysates comprise one or more enzymes for the conversion of formaldehyde to pyruvate, or one or more enzymes for the conversion of formaldehyde to isobutanol. Cell lysates may further comprise small molecules and cofactors (*e.g.*, adenosine triphosphate, NAD(PH)), and salts, such as magnesium, for enzyme function.

Further provided herein are biofuels and other chemical compound or intermediate produced by any one or more of the cell-free processes described herein.

BRIEF DESCRIPTION OF DRAWINGS

Figure 1 shows a diagram of an exemplary cell-free multiphase reaction system of the invention designed to deliver methane and oxygen into the aqueous phase at a high rate.

Figure 2 shows a diagram of an exemplary cell-free transformation platform of the invention.

Figure 3 shows a diagram of an exemplary cell-free transformation platform of the invention.

Figure 4 shows a diagram of the isobutanol pathway.

- 6 -

Figure 5 shows a diagram of the methane monooxygenase (MMO) catalytic system assuming that adequate NADH is supplied by a nicotinamide adenine dinucleotide (NAD⁺)-coupled methanol dehydrogenase.

5 *Figure 6* shows a diagram of an apparatus for use in some embodiments to assess gas/liquid interface formation and stability.

Figure 7 shows a diagram of an apparatus for use in some embodiments to assess system performance and safety.

Figure 8 shows a diagram of multiplexed instrumented reactors for use in assessing MMO activity and methanotroph cell growth.

10 *Figure 9* shows a diagram demonstrating feedstock flexibility.

Figure 10 shows a graph of data demonstrating that a single extract can accept glucose or ribose-5-P.

Figure 11 shows a diagram demonstrating use of 5-carbon (C5) and 6-carbon (C6) sugars in cell-free reactions.

15 *Figure 12* shows a diagram of methane metabolism.

Figure 13 shows a diagram of the isoprene pathway, requiring energy and reducing equivalents.

Figure 14 shows a diagram of methane as the sole carbon source for isoprene.

20 DETAILED DESCRIPTION OF THE INVENTION

Provided herein, in some embodiments, is a cell-free system for the conversion of natural gas, in particular methane, to a biofuel and other chemical compounds. The process combines cell-free lysates from methanotrophs with those from other recombinant bacteria in a reactor typically operated at high pressure (*e.g.*, at least 1 bar, or at least 5 bar). The cell-free processes, systems, reactions mixtures, compositions, recombinant organisms, and nucleic acid constructs of the invention are considered parts of a platform technology enabling the efficient production of many different biofuels or other chemical compounds from methane or natural gas. The use of a cell-free biosynthetic system does not require the maintenance of cell viability, enables direct monitoring and adjustment of reaction parameters (*e.g.*, substrate feed rates, temperature, pH, pressure, dissolved oxygen), provides a flexible reaction environment, and enables control of carbon and energy flux (*see, e.g.*, International Publication Numbers WO 2010/077806 and WO 2012/030980, incorporated by reference herein). In some

25
30

- 7 -

embodiments, the cell-free process of the invention addresses mass transfer challenges associated with methane and oxygen solubility by operating under high pressure, providing an organic phase in which methane has greater solubility, and increasing methane bubble and air bubble surface area by injecting gases at high pressure.

5 Generally, a cell-free process of the invention includes (i) use of methanotrophs to express enzymes (*e.g.*, MMO) for the conversion of methane to formaldehyde, (ii) use of recombinant bacteria (*e.g.*, *E. coli*) to express enzymes in biosynthetic pathways that produce biofuels and other chemical compounds and (iii) use of control systems and a reaction environment with flexibility that previously was only offered by chemical processes using
10 inorganic catalysts.

 Some aspects of the invention provide cell-free processes that use a multiphase reaction system, as shown in Figure 1. The reaction mixture of the cell-free bioreactor includes, in some embodiments: (i) a gaseous phase to deliver methane and oxygen and to remove CO₂, and (ii) an aqueous, or continuous, phase that includes the enzymes and membrane vesicles. In
15 some embodiments, the cell-free bioreactor also includes an organic (apolar) phase to deliver the natural gas (including methane) and to preferentially absorb the desired product (Figure 1). In some embodiments, for example when an organic phase is included, the aqueous phase, includes an emulsion of the cell lysate (*e.g.*, including enzymes and membrane vesicles of the cell contents), the gaseous phase (*e.g.*, as air bubbles to deliver oxygen and remove CO₂ and as
20 methane bubbles to deliver methane), and the organic phase (*e.g.*, droplets of organic solvent to deliver methane and capture the product). In some embodiments, the emulsion is formed outside of the reactor. In other embodiments, the emulsion is formed via high pressure injection of an organic phase into the bioreactor. "High pressure," as used herein, refers to pressure that is at least 1 bar. For example, high pressure may refer to pressure that is at least 1
25 bar, at least 2 bar, at least 5 bar, or at least 10 bar. In some embodiments, high pressure may refer to a pressure that is 1 to 5 bar, 1 to 10 bar, or 5 to 10 bar.

 The organic solvent of the organic phase may be an alkane. Examples of alkanes for use as provided herein include, without limitation, pentane, hexane, heptane, octane, nonane, and decane. Other organic solvents may be used. For example, organic solvents with the
30 following properties are contemplated herein: (i) use of which results in less than 10% reduction in pyruvate production rate in cell-free reaction, and (ii) methane solubility greater than 25 mg/L to 30 mg/L (*e.g.*, 23 mg/L) at atmospheric pressure.

- 8 -

Figure 2 shows a diagram of a non-limiting exemplary cell-free process of the present invention. Each step of the process is indicated numerically, 1-8. In step 1, recombinant methanotrophs express enzymes for the conversion of methane in natural gas to formaldehyde. In step 2, recombinant *Escherichia coli* (*E. coli*) express enzymes for the conversion of formaldehyde to biofuel, in this case, isobutanol, and also provide a compartmentalized protease (*see, e.g.*, International Publication Numbers WO 2010/077806 and WO 2012/030980, incorporated by reference herein). The protease, for example, cleaves a targeted formaldehyde dehydrogenase (expressed by the methanotroph) that has been engineered to contain a cognate protease recognition site. Formaldehyde dehydrogenase converts formaldehyde to formate, and therefore, it interferes with the conversion of formaldehyde to a biofuel or other chemical compound in a cell-free reaction system. Thus, when the organisms from step 1 and step 2 are lysed and combined in step 3, the protease is released from its cellular compartment (*e.g.*, periplasm) and cleaves the formaldehyde dehydrogenase, thereby inactivating it. This inactivation of formaldehyde dehydrogenase increases the conversion yield of biofuel. The conversion of methane into the biofuel occurs in step 4, in an engineered and computer controlled bioreactor pressurized at, for example, 10 bar to drive mass transfer rates for a volumetric productivity of, for example, at least 1, at least 5, at least 10, at least 15, at least 20, or at least 25 g/liter-hr. The bioreactor of step 4 includes a multiphase fluid containing an aqueous phase to support the enzyme catalysts of the cell lysates and, optionally, an organic phase to help deliver the natural gas or methane and absorb the product biofuel. These phases, in some embodiments, are separated in step 5 to deliver the organic phase for fuel removal in step 6. In some embodiments, the organic phase is then recharged with natural gas (or methane) and injected back into the bioreactor at multiple points. Air and natural gas may be separately injected, in some embodiments. Further, in some instances, external heat exchangers as well as a cooling jacket remove process heat and control reactor temperature. To ensure safe operation, typically multiple temperature sensors throughout the reactor detect any "hot spots" possibly caused by spontaneous combustion. This triggers an immediate reduction in methane and/or air injection and a rebalancing of the distributed inlets.

It should be understood that some of the steps of the cell-free process may be carried out simultaneously or sequentially. For example, expression of enzymes for the conversion of methane to formaldehyde and expression of enzymes for the conversion of formaldehyde to

- 9 -

biofuel may occur simultaneously (*e.g.*, in separate compartments of a reactor) before cell lysis.

Figure 3 shows a diagram of another non-limiting example of a cell-free process of the present invention. For the sake of clarity, elements such as the inoculum preparation fermentors, compressors, coolers, heat exchangers, and pumps are not shown. In this example, the methanotrophs are concentrated by centrifugation before being combined with the uncentrifuged recombinant *E. coli* culture. The combined cells are held in a chilled tank while waiting for lysis with a high pressure homogenizer. The cell lysate is then centrifuged and passed into a second holding tank. Based on an estimated biofuel adsorption capacity in alkane (*e.g.*, decane) of 16%, approximately half of the cell-free reaction fluid is pumped from the reactor to a decanting centrifuge each hour, in some embodiments. The gas phase is removed and the liquid phases separated. The organic phase, in some embodiments, is transferred to the distillation device for biofuel removal. The re-purified alkane is then cooled and injected with methane to produce a chilled methane-in-alkane foam for distributed entry back into the cell-free reactor. About 10% of the aqueous phase leaving the decanting centrifuge is removed to maintain a constant volume in the cell-free reactor and to remove any possible water soluble toxins. The aqueous catalyst stream is then cooled before single point entry into the bioreactor.

Cell-free Systems

The cell-free processes provided herein permit multiple enzymes to be expressed (*e.g.*, from a bacterial artificial chromosome) during bacterial (*e.g.*, *E. coli*) fermentation. In some embodiments, enzymes for the conversion of substrates (*e.g.*, methane) to the desired chemical product (*e.g.*, pyruvate) are expressed in a recombinant, typically rapidly-growing, bacterial strain (*e.g.*, *E. coli*). Prior to cell harvest and lysis, the enzymes accumulate to optimal levels. Cell lysis enables periplasmically-expressed proteases (discussed below) to eliminate side pathways, shunting carbon to the product of interest. An optimized chemical environment activates respiration catalyzed by inner membrane vesicles formed during cell lysis, thereby providing a plentiful supply of ATP and removing excess reducing equivalents to recycle NAD⁺ and/or nicotinamide adenine dinucleotide phosphate (NADP⁺), if needed (*see* International Publication Numbers WO 2010/077806 and WO 2012/030980, incorporated by reference herein).

- 10 -

A “cell-free” composition, as used herein, refers to a composition substantially free of intact cells. One of skill in the art would understand that a certain percentage of the cells after lysing may be intact, *e.g.*, less than 10%, less than 5%, less than 2%, less than 1%, or less than 0.5%. A “cell-free system,” as used herein, is an isolated cell-free system containing a cell lysate or extract expressly engineered to include an enzyme or cascade of enzymes that, when acting in a given sequence (*e.g.*, in an enzymatic pathway) and proportion over a determined substrate, results in the generation of a desired product (*e.g.*, a biofuel or other chemical compound, or an intermediate thereto).

A “cell lysate,” as used herein, refers to a fluid containing the contents of lysed cells. Cell lysates may be whole cell lysates and/or crude (unpurified) cell lysates. In some embodiments, cell lysates may be partially purified (*e.g.*, to remove cellular debris/particulate such as damaged outer cell membranes). In some embodiments, cell lysates are prepared by high pressure lysis, thereby forming intracellular membrane vesicles that enable oxidative phosphorylation in the cell lysates. Other methods of preparing cell lysate are well-known in the art and include, without limitation, sonication, homogenization, enzymatic lysis using lysozyme, and freezing and grinding.

Sonication includes lysing cells by liquid shear and cavitation. DNA is also sheared during sonication, so it may not be necessary to add DNase to the cell suspension. Typically, the cell suspension is kept on ice to control temperature, and short pulses (5-10 sec) with pauses (10-30 sec) may be used to re-establish a low temperature.

Homogenizers may be used to press lyse cells by pressurizing the cell suspension and suddenly releasing the pressure. This creates a liquid shear capable of lysing cells. Multiple (2-3) passes are generally required to achieve adequate lysis. The high operating pressures, however, result in a rise in operating temperatures. Therefore, pressure cells may be cooled (4 °C) prior to use. In addition to temperature control, care should be taken to avoid inactivating proteins by foaming.

Enzymatic lysis is based on the digestion of the peptidoglycan layer of the bacterial cell wall by lysozyme. Gram-negative bacteria, however, have an outer membrane that is external to the cell wall and may need to be permeabilized to expose the peptidoglycan layer. Tris-buffered saline, often used as a buffer in lysis methods, effectively permeabilizes outer membranes. This effect can be enhanced by the addition of a chelating agent such as ethylenediaminetetraacetic acid (EDTA, *e.g.*, 1 mM). EDTA chelates the magnesium ions that

- 11 -

stabilize membranes. During cell lysis, DNA is liberated and it may be necessary to add DNase (1 mg/ml) to reduce the viscosity of the preparation. Enzymatic cell lysis can be carried out on any scale. To increase the level of cell lysis the solution may also be sonicated.

Another lysis method is to freeze the cells directly in liquid nitrogen and grind the frozen cells to a powder using a mortar and pestle that are chilled with liquid nitrogen. The powder can be stored indefinitely at -80 °C, and the cell lysate can be prepared by adding the powder to buffer.

Cell lysates of methanotrophs and recombinant bacteria are combined in a cell-free system with at least one substrate, cofactor, or a combination thereof required for producing the desired biofuel or other chemical compound. “Substrate,” as used herein, is a compound or mixture of compounds capable of providing the required elements needed to synthesize a compound of interest. In certain embodiments, a substrate may refer to a carbon source. In some embodiments, methane is the substrate.

“Cofactor,” as used herein, is a non-protein chemical compound required for a protein’s (*e.g.*, enzyme’s) biological activity. In some embodiments, cofactors are required for the cell-free biotransformation of methane to a biofuel or other chemical compound. Examples of cofactors include, without limitation, NADH, NAD⁺, ATP, ADP, Pi, and NADPH. In some embodiments, cofactors are provided by (*e.g.*, present in) the cell extract.

Protease targeting

Some aspects of the present disclosure use methods for silencing (*e.g.*, eliminating or reducing) unwanted side reactions by inactivating one or more enzymes that is detrimental to a particular stage of the biotransformation process. In some embodiments, a detrimental enzyme is one that catalyzes a rate-limiting step in a biosynthetic pathway. This is achieved, in some embodiments, by engineering an enzyme to include a protease recognition site. Such recombinant enzymes typically have a protease recognition sequence selectively located in their primary amino acid sequence such that, despite the presence of the recognition sequence, the activity of the recombinant protein is sufficient to enable wild-type growth of the cell. The recombinant enzymes can be selectively inactivated by the introduction, expression, and/or activation of a cognate protease, which cleaves the recombinant target protein specifically at the protease recognition sequence, thereby rendering the recombinant enzyme inactive (or with reduced activity). The cognate protease is sequestered into a cellular compartment (*e.g.*,

- 12 -

periplasm) until inactivation of the recombinant enzyme is needed, at which time the protease is brought into contact with the recombinant enzyme to cleave and inactivate the enzyme.

The recombinant enzymes provided herein may be inactivated by any one of a variety of proteases that cleave at specific recognition sequences. As used herein, “protease

5 recognition sequence,” in the context of an enzyme, refers to an amino acid sequence that is recognized and cleaved by a cognate protease. In the context of a nucleic acid that encodes a protein, a “protease recognition sequence” refers to a sequence that encodes the amino acid sequence recognized and cleaved by a cognate protease. As used herein, “cognate protease” refers to a protease that cleaves and thereby inactivates a recombinant enzyme. Cognate
10 proteases that may be used herein include those with a single, specific recognition sequence, meaning the proteases cleave within or adjacent to a specific sequence of one or more amino acids. In some embodiments, the proteins of the invention are prepared with an engineered human rhinovirus 3C protease recognition sequence.

Other examples of proteases that may be used in accordance with the invention include,
15 without limitation, alanine carboxypeptidase, *Armillaria mellea* astacin, bacterial leucyl aminopeptidase, cancer procoagulant, cathepsin B, clostripain, cytosol alanyl aminopeptidase, elastase, endoproteinase Arg-C, enterokinase, gastricsin, gelatinase, Gly-X carboxypeptidase, glycyl endopeptidase, human rhinovirus 3C protease, hypodermin C, Iga-specific serine endopeptidase, leucyl aminopeptidase, leucyl endopeptidase, lysC, lysosomal pro-X
20 carboxypeptidase, lysyl aminopeptidase, methionyl aminopeptidase, myxobacter, nardilysin, pancreatic endopeptidase E, picornain 2A, picornain 3C, proendopeptidase, prolyl aminopeptidase, proprotein convertase I, proprotein convertase II, russellysin, saccharopepsin, semenogelase, T-plasminogen activator, thrombin, tissue kallikrein, tobacco etch virus (TEV), togavirin, tryptophanyl aminopeptidase, U-plasminogen activator, V8, venombin A, venombin
25 AB and Xaa-pro aminopeptidase (see Rawlings, S. D., *et al.*, Handbook of Proteolytic Enzymes, Academic Press, 2013, Science, Elsevier Ltd., 4094 pages, incorporated herein by reference for its teachings relating to the structural chemistry and biological aspects of the proteases described therein). Other proteases may be used in accordance with the invention.

Compartmentalization of enzymes in either the cytoplasm or periplasm of bacteria (*e.g.*,
30 *E. coli*) provides an opportunity to overexpress components of biosynthetic pathways and to accumulate a specific protease without affecting cell health and growth. When the cells are lysed by, for example, scalable high pressure homogenizers, the intracellular compartments

- 13 -

mix to activate the pathways and inactivate the deleterious enzymes; essentially to remodel the catalytic system. This permits the enzyme catalysts to be produced inexpensively, in some instances. Additionally, the absence of the cell wall and dispersion of the macromolecular catalysts throughout the entire reaction volume permits, in some embodiments, precise
5 sampling for on-line monitoring as well as immediate dispersion of added substrates and reaction control reagents (*e.g.*, effectors of pH, substrates). This enables, in some embodiments, the ability to approach complex biological conversions (*e.g.*, converting methane to a biofuel) using technologies employed by traditional heterogeneous catalysis processes as described herein.

10 As discussed herein, some aspects of the invention relate to the inactivation of enzymes detrimental to biofuel (*e.g.*, isobutanol) production. For example, in some embodiments, pyruvate dehydrogenase may be targeted by a protease in the manner described herein. Pyruvate dehydrogenase (E1) is the first component enzyme of pyruvate dehydrogenase complex (PDC). The pyruvate dehydrogenase complex contributes to transforming pyruvate
15 into acetyl-CoA by a process called pyruvate decarboxylation. Acetyl-CoA may then be used in the citric acid cycle to carry out cellular respiration, thus pyruvate dehydrogenase contributes to linking the glycolysis metabolic pathway to the citric acid cycle and releasing energy via NADH.

20 In some embodiments, formaldehyde dehydrogenase may be targeted by a protease in the manner described herein. Formaldehyde dehydrogenase is an enzyme that catalyzes the chemical conversion of formaldehyde to formate.

Methanotrophs

Some aspects of the invention relate to enzymes that catalyze the conversion of
25 methane to formaldehyde and are derived from a methanotroph. Enzymes are considered to be “derived from” an organism if they are expressed in that organism or obtained from that organism. Thus, in some embodiments, cell-free processes of the invention use cell extracts from methanotrophs (*e.g.*, *Methylococcus capsulatus* Bath) to provide methane monooxygenase, a biological catalyst for methane activation. “Methanotrophs,” as used
30 herein, refers to prokaryotes that are able to metabolize methane as their only source of carbon and energy. Methanotrophs can grow aerobically or anaerobically and require single-carbon compounds to survive.

- 14 -

In some embodiments, factors to consider when using methanotrophs for the conversion of methane to a biofuel or other chemical compound, include without limitation: methanotrophic culture densities on the order of 5-15 grams cell dry weight per liter (*e.g.*, 10 grams cell dry weight per liter) or 25-100 grams dry weight per liter (*e.g.*, 50 grams dry weight per liter grams dry weight per liter within 2-5 years (*e.g.*, 2.5 years); a turnover frequency of methane oxidation by the methane monooxygenase on the order of 25 to 75 50 sec⁻¹ (*e.g.*, 50 sec⁻¹) and an effective catalyst lifetime of 10 to 30 hours; net zero consumption of reducing equivalents in the oxidation of methane to precursor compounds for biofuel production and protease targeting of side processes to ensure maximum carbon flux from methane to a biofuel or other chemical compound.

In some embodiments, batch culture systems and/or continuous culture systems are used to systematically manipulate growth conditions, *e.g.*, varying nitrogen, phosphorous, iron, copper, and temperature, along with the addition of paraffin (Han B, *et al.* 2009 *Appl Microbiol Biotechnol* 83:669-77) to stimulate both high growth rates and cell densities in methanotrophs (*e.g.*, *M. capsulatus* Bath). Thus, in some embodiments, any one or more of nitrogen, phosphorous, iron, copper, and paraffin may be used in the cell-free process of the invention to improve methanotroph growth rates and to improve the accumulation of MMO.

In some embodiments, the genome of the methanotrophs is modified to have enhanced pMMO expression and activity.

Recombinant bacteria

Some aspects of the invention relate to enzymes that catalyze the conversion of formaldehyde to a biofuel or other chemical compound and which are derived from recombinant bacteria. As used herein, “recombinant bacteria,” or a “recombinant cell,” refers to a cell into which a foreign, or exogenous, nucleic acid (*i.e.*, nucleic acid not native to the cell) is introduced. A recombinant cell may include a recombinant nucleic acid (*e.g.*, DNA or RNA), which is formed artificially by combining nucleic acids from different sources. In some embodiments, the foreign nucleic acid is integrated into the genome of the cell, whereas in other embodiments, the foreign nucleic acid is not integrated into the genome of the cell. The terms “recombinant cell” and “genetically engineered cell” may be used interchangeably herein.

- 15 -

While many of the embodiments provided herein describe the use of recombinant *Escherichia coli* (*E. coli*), it should be understood that the invention is not so limited, but rather contemplates the use of any bacterial cells that grow rapidly and can be easily genetically manipulated. Thus, methods associated with the present disclosure encompass lysates from any type of cell, *e.g.*, prokaryotic and eukaryotic cells. In some embodiments, the cell is a bacterial cell, such as *Escherichia* spp., *Streptomyces* spp., *Zymonas* spp., *Acetobacter* spp., *Citrobacter* spp., *Synechocystis* spp., *Rhizobium* spp., *Clostridium* spp., *Corynebacterium* spp., *Streptococcus* spp., *Xanthomonas* spp., *Lactobacillus* spp., *Lactococcus* spp., *Bacillus* spp., *Alcaligenes* spp., *Pseudomonas* spp., *Aeromonas* spp., *Azotobacter* spp., *Comamonas* spp., *Mycobacterium* spp., *Rhodococcus* spp., *Gluconobacter* spp., *Ralstonia* spp., *Acidithiobacillus* spp., *Microtholunatus* spp., *Geobacter* spp., *Geobacillus* spp., *Arthrobacter* spp., *Flavobacterium* spp., *Serratia* spp., *Saccharopolyspora* spp., *Thermus* spp., *Stenotrophomonas* spp., *Chromobacterium* spp., *Sinorhizobium* spp., *Saccharopolyspora* spp., *Agrobacterium* spp., and *Pantoea* spp. The bacterial cell can be a Gram-negative cell such as an *Escherichia coli* (*E. coli*) cell, or a Gram-positive cell such as a species of *Bacillus*. In other embodiments the cell is a fungal cell such as a yeast cell, *e.g.*, *Saccharomyces* spp., *Schizosaccharomyces* spp., *Pichia* spp., *Paffia* spp., *Kluyveromyces* spp., *Candida* spp., *Talaromyces* spp., *Brettanomyces* spp., *Pachysolen* spp., *Debaryomyces* spp., *Yarrowia* spp. and industrial polyploid yeast strains. Other non-limiting examples of fungi include *Aspergillus* spp., *Penicillium* spp., *Fusarium* spp., *Rhizopus* spp., *Acremonium* spp., *Neurospora* spp., *Sordaria* spp., *Magnaporthe* spp., *Allomyces* spp., *Ustilago* spp., *Botrytis* spp., *Pectobacterium* spp., and *Trichoderma* spp. In some embodiments, the cell is an algal cell, a plant cell, an insect cell, or a mammalian cell. It should be appreciated that some cells compatible with the invention may express an endogenous copy of one or more of the genes associated with the invention as well as a recombinant copy. It should also be appreciated that some cells for use in accordance with the invention do not contain the wild-type chromosomal copy of a gene encoding the wild-type enzyme (*e.g.*, the wild-type enzyme corresponding to the genetically-engineered enzyme introduced into the cell).

In some embodiments, *E. coli* cells are used as they are economical, well studied, and amenable to genetic manipulation. In some embodiments, yeast cells are used.

In some embodiments, the cell lysate that contains enzymes that catalyze the conversion of formaldehyde to a biofuel or other chemical compound is a lysate of a

- 16 -

recombinant *E. coli* cell engineered to overexpress one or more enzymes described herein. In some embodiments, the cell lysate is a lysate of an *E. coli* cell engineered to overexpress a group of enzymes, *e.g.*, two, three, four, five, six, seven, eight, nine, ten, eleven, twelve, thirteen, fourteen, fifteen, sixteen, seventeen, eighteen, nineteen, twenty, or more than twenty enzymes. In some embodiments, the cell lysate is a combination of different cell lysates, *e.g.*, a combination of two, three, four, five, six, seven, eight, nine, ten, or more than ten different cell lysates, obtained from two, three, four, five, six, seven, eight, nine, ten, or more than ten different cells, *e.g.*, from different organisms, each engineered to overexpress one or more enzymes that the conversion of formaldehyde to a biofuel or chemical compound. In some
5 embodiments, lysates from different organisms (*e.g.*, different strains of bacteria) are combined. In some embodiments, different engineered *E. coli* strains (*e.g.*, overexpressing different production pathway proteins) are combined to optimize enzyme levels prior to creating a single engineered strain with all overexpressed proteins.

The cells provided herein, in some embodiments, are prokaryotic cells that may be
15 transformed with nucleic acids encoding as enzymes as provided herein. Transformation and transfection are processes by which exogenous genetic material is introduced into a prokaryotic cell and into a eukaryotic cell, respectively. Transformation can be achieved by electroporation or by chemical means. The cells to be transformed are typically in a state of competence. Thus, in some embodiments, the cells provided herein are electrocompetent or
20 chemically competent cells (*see, e.g.*, Donahue *et al. Focus* 20, 1998, (2):54-56; Donahue *et al. Focus* 20, 1998, (2):77-78; Inoue *et al. Gene* 96, 1990, (1): 23-28, each of which is incorporated by reference herein). A variety of electrocompetent and chemically competent cells are known in the art and may be used in accordance with the invention.

Cells, in some embodiments, may comprise selectable markers. Selectable markers
25 include, without limitation, genes encoding proteins that increase or decrease either resistance or sensitivity to antibiotics (*e.g.*, ampicillin resistance genes, kanamycin resistance genes, neomycin resistance genes, tetracyclin resistance genes and chloramphenicol resistance genes) or other compounds, genes encoding enzymes with activities detectable by standard assays known in the art (*e.g.*, β -galactosidase, luciferase or alkaline phosphatase), and genes that
30 visibly affect the phenotype of transformed or transfected cells, hosts, colonies, or plaques (*e.g.*, green fluorescent protein). Other selectable markers known in the art may also be used in accordance with the invention.

- 17 -

A “nucleic acid,” as used herein, refers to a polymer of at least three nucleotides (*e.g.*, adenine, thymine, cytosine, guanine, uracil) covalently linked together. The polymer may include natural nucleosides (*i.e.*, adenosine, thymidine, guanosine, cytidine, uridine, deoxyadenosine, deoxythymidine, deoxyguanosine, and deoxycytidine), nucleoside analogs (*e.g.*, 2-aminoadenosine, 2-thiothymidine, inosine, pyrrolo-pyrimidine, 3-methyl adenosine, 5-methylcytidine, C5-bromouridine, C5-fluorouridine, C5-iodouridine, C5-propynyl-uridine, C5-propynyl-cytidine, C5-methylcytidine, 7-deazaadenosine, 7-deazaguanosine, 8-oxoadenosine, 8-oxoguanosine, O(6)-methylguanine, 4-acetylcytidine, 5-(carboxyhydroxymethyl)uridine, dihydrouridine, methylpseudouridine, 1-methyl adenosine, 1-methyl guanosine, N6-methyl adenosine, and 2-thiocytidine), chemically modified bases, biologically modified bases (*e.g.*, methylated bases), intercalated bases, modified sugars (*e.g.*, 2'-fluororibose, ribose, 2'-deoxyribose, 2'-O-methylcytidine, arabinose, and hexose), or modified phosphate groups (*e.g.*, phosphorothioates and 5'-N-phosphoramidite linkages). A nucleic acid of the present disclosure will generally contain phosphodiester bonds. A nucleic acid may be single-stranded (ss) or double-stranded (ds), DNA or RNA. In some embodiments, a nucleic acid is in the form of cDNA. In some embodiments, a nucleic acid is in the form of genomic DNA. Nucleic acids of the present disclosure may be in the form of vectors including, without limitation, plasmids, cosmids, and artificial chromosomes (*e.g.*, bacterial artificial chromosomes (BACs) and yeast artificial chromosomes (YACs)). Other nucleic acid-based vectors are described below.

The nucleic acids used herein, whether RNA, iRNA, antisense nucleic acid, cDNA, genomic DNA, vectors, viruses or hybrids thereof, may be isolated from a variety of sources, genetically engineered, amplified, and/or expressed/generated recombinantly. Recombinant polypeptides generated from these nucleic acids can be individually isolated or cloned and tested for a desired activity. Any recombinant expression system can be used, including bacterial, mammalian, yeast, insect or plant cell expression systems.

Techniques for the manipulation of nucleic acids, such as, *e.g.*, subcloning, labeling probes (*e.g.*, random-primer labeling using Klenow polymerase, nick translation, amplification), sequencing, hybridization and the like are well described in the scientific and patent literature, see, *e.g.*, Sambrook, ed., *Molecular Cloning: A Laboratory Manual* (2nd ed.), Vols. 1-3, Cold Spring Harbor Laboratory, (1989); *Current Protocols In Molecular Biology*, Ausubel, ed. John Wiley & Sons, Inc., New York (1997); *Laboratory Techniques In*

- 18 -

Biochemistry And Molecular Biology: Hybridization With Nucleic Acid Probes, Part I. Theory and Nucleic Acid Preparation, Tijssen, ed. Elsevier, N.Y. (1993), incorporated by reference herein.

Alternatively, these nucleic acids can be synthesized *in vitro* by well-known chemical synthesis techniques, as described in, *e.g.*, Adams (1983) *J. Am. Chem. Soc.* 105:661; Belousov (1997) *Nucleic Acids Res.* 25:3440-3444; Frenkel (1995) *Free Radic. Biol. Med.* 19:373-380; Blommers (1994) *Biochemistry* 33:7886-7896; Narang (1979) *Meth. Enzymol.* 68:90; Brown (1979) *Meth. Enzymol.* 68:109; Beaucage (1981) *Tetra. Lett.* 22:1859; U.S. Pat. No. 4,458,066, incorporated by reference herein.

Nucleic acids encoding enzymes of the present disclosure may comprise regulatory sequences operably linked to coding sequences. As used herein, a regulatory sequence (*e.g.*, promoter sequence) and a coding sequence are said to be “operably” linked when they are covalently linked in such a way as to place the expression or transcription of the coding sequence under the influence or control of the regulatory sequences (*e.g.*, such that the regulatory sequence “drives” transcriptional initiation and/or expression of the coding sequence). For example, for a coding sequences that is to be translated into a functional enzyme, two DNA sequences are considered operably linked if induction of a promoter in the 5′ regulatory sequences results in the transcription of the coding sequence and if the nature of the linkage between the two DNA sequences does not (1) result in the introduction of a frame-shift mutation, (2) interfere with the ability of the promoter region to direct the transcription of the coding sequences, or (3) interfere with the ability of the corresponding RNA transcript to be translated into the enzyme. Thus, a promoter region would be operably linked to a coding sequence if the promoter region can effect transcription of that DNA sequence such that the resulting transcript can be translated into the desired enzyme.

When a nucleic acid that encodes any of the enzymes provided herein is expressed in a cell, a variety of transcription control sequences may be used to direct its expression. For example, a nucleic acid may contain a promoter, an enhancer, and/or a terminator. Alternatively, the vector into which the nucleic acid is inserted may contain such regulatory sequences. A “promoter,” as used herein, refers to a control region of a nucleic acid sequence at which initiation and rate of transcription of the remainder of a nucleic acid sequence are controlled. A promoter may also contain subregions at which regulatory proteins and molecules may bind, such as RNA polymerase and other transcription factors. Promoters may

- 19 -

be constitutive (*e.g.*, unregulated), inducible, activatable, repressible, tissue-specific or any combination thereof. A promoter drives expression or drives transcription of the nucleic acid sequence that it regulates. A promoter may be one naturally associated with a gene or sequence, as may be obtained by isolating the 5'-non-coding sequences located upstream of the coding segment and/or exon of a given gene or sequence. Such a promoter can be referred to as "endogenous" or "native." A coding nucleic acid segment, in some embodiments may be positioned under the control of a recombinant or heterologous promoter, which refers to a promoter that is not normally associated with the encoded nucleic acid sequence in its natural environment. Such promoters may include promoters of other genes, promoters isolated from any other prokaryotic, viral or eukaryotic cell, and synthetic promoters that are not "naturally occurring" such as, for example, those that contain different elements of different transcriptional regulatory regions and/or mutations that alter expression through methods of genetic engineering that are known in the art. In addition to producing nucleic acid sequences of promoters synthetically, sequences may be produced using recombinant cloning and/or nucleic acid amplification technology, including polymerase chain reaction (PCR).

Promoters suitable for use with prokaryotic hosts include, without limitation, the β -lactamase and lactose promoter systems, alkaline phosphatase, a tryptophan (*trp*) promoter system, and numerous hybrid promoters such as the *tac* promoter. However, other known bacterial promoters are also suitable, *e.g.*, the *lacI* promoter, the T3 promoter, the T7 promoter, the arabinose promoter, the *gpt* promoter, the lambda PR promoter, the lambda PL promoter, promoters from operons encoding glycolytic enzymes such as 3-phosphoglycerate kinase (PGK), and the acid phosphatase promoter. Their nucleotide sequences have been published, thereby enabling a skilled worker operably to ligate them to a sequence of interest using linkers or adaptors. Promoters for use in bacterial systems also will contain a Shine-Dalgarno (S.D.) sequence operably linked to the coding sequence. In some embodiments, the host cell may be modified genetically to adjust concentrations of metabolite or inducer transporter proteins so that all cells in a culture will be induced equivalently.

Promoters suitable for eukaryotic cells, *e.g.*, yeast cells, are also known in the art. Virtually all eukaryotic genes have an AT-rich region located approximately 25 to 30 bases upstream from the site where transcription is initiated. Another sequence found 70 to 80 bases upstream from the start of transcription of many genes is a CXCAAT region where X may be any nucleotide. At the 3' end of most eukaryotic genes is an AATAAA sequence that may be

- 20 -

the signal for addition of the poly A tail to the 3' end of the coding sequence. All of these sequences are suitably inserted into eukaryotic expression vectors. Examples of suitable promoting sequences for use with yeast hosts include, without limitation, promoters for 3-phosphoglyceratekinase or other glycolytic enzymes, such as enolase, glyceraldehyde-3-phosphate dehydrogenase, hexokinase, pyruvate decarboxylase, phosphofructokinase, glucose-6-phosphate isomerase, 3-phosphoglycerate mutase, pyruvate kinase, triose phosphate isomerase, phosphoglucose isomerase, and glucokinase.

An “inducible promoter,” as used herein, is one that is characterized by initiating or enhancing transcriptional activity when in the presence of, influenced by or contacted by an inducer or inducing agent. An “inducer” or “inducing agent” may be endogenous or a normally exogenous compound or protein that is administered in such a way as to be active in inducing transcriptional activity from the inducible promoter. Inducible promoters for use in accordance with the invention include any inducible promoter described herein or known to one of ordinary skill in the art. Examples of inducible promoters include, without limitation, chemically/biochemically-regulated and physically-regulated promoters such as isopropyl β -D-1-thiogalactopyranoside (IPTG)-regulated promoters, alcohol-regulated promoters, tetracycline-regulated promoters (*e.g.*, anhydrotetracycline (aTc)-responsive promoters and other tetracycline-responsive promoter systems, which include a tetracycline repressor protein (tetR), a tetracycline operator sequence (tetO) and a tetracycline transactivator fusion protein (tTA)), steroid-regulated promoters (*e.g.*, promoters based on the rat glucocorticoid receptor, human estrogen receptor, moth ecdysone receptors, and promoters from the steroid/retinoid/thyroid receptor superfamily), metal-regulated promoters (*e.g.*, promoters derived from metallothionein (proteins that bind and sequester metal ions) genes from yeast, mouse and human), pathogenesis-regulated promoters (*e.g.*, induced by salicylic acid, ethylene or benzothiadiazole (BTH)), temperature/heat-inducible promoters (*e.g.*, heat shock promoters), and light-regulated promoters (*e.g.*, light responsive promoters from plant cells). Other inducible promoters may be used in accordance with the invention.

A “codon,” as used herein, refers to a set of three adjacent nucleotides that encode an amino acid. In some embodiments, nucleic acids are codon-optimized for improved expression of the genetically engineered enzymes useful in the present invention. Codon optimization, also referred to as biased codon usage, refers to differences in the frequency of occurrence of synonymous codons in coding DNA.

- 21 -

“Isozymes,” as used herein, refer to enzymes that differs in amino acid sequence but catalyzes the same chemical reaction or produces the same reaction product from starting material.

Enzymes of the present disclosure may, in some embodiments, be expressed
5 episomally. Thus, the present disclosure contemplates the use of vectors comprising a nucleic acid that express enzymes as described herein. A “vector,” as used herein, may be any of a number of nucleic acids into which a desired sequence or sequences may be inserted by restriction and ligation for transport between different genetic environments or for expression in a cell. Vectors are typically composed of DNA although RNA vectors are also available.
10 Examples of vectors in accordance with the present disclosure include, without limitation, plasmids, cosmids, fosmids, phagemids, virus genomes, and artificial chromosomes (*e.g.*, BACs, YACs). In some embodiments, a nucleic acid of the present disclosure is provided in a recombinant cloning vector. In some embodiments, a nucleic acid variant is expressed in a recombinant expression vector.

15 A cloning vector of the present disclosure is able to replicate autonomously or integrated in the genome of a cell. A cloning vector has an endonuclease restriction sequence at which the vector may be cut in a determinable fashion and into which a desired DNA sequence may be ligated such that the new recombinant vector retains its ability to replicate in a cell. In the case of plasmids, replication of the desired sequence may occur many times as
20 the plasmid increases in copy number within a cell such as a bacterium or just a single time per cell before the cell reproduces by mitosis.

An expression vector of the present disclosure is one into which a desired DNA coding sequence may be inserted by restriction and ligation such that it is operably linked to regulatory sequences and may be expressed as an RNA transcript.

25 Vectors of the invention may further comprise a marker sequence for use in the identification of cells that have or have not been transformed or transfected with the vector. Markers include, for example, genes encoding proteins that increase or decrease either resistance or sensitivity to antibiotics (*e.g.*, ampicillin resistance genes, kanamycin resistance genes, neomycin resistance genes, tetracycline resistance genes and chloramphenicol resistance
30 genes) or other compounds, genes encoding enzymes with activities detectable by standard assays known in the art (*e.g.*, β -galactosidase, luciferase or alkaline phosphatase), and genes that visibly affect the phenotype of transformed or transfected cells, hosts, colonies or plaques

- 22 -

(e.g., green fluorescent protein). In some embodiments, the vectors used herein are capable of autonomous replication and expression of the structural gene products present in the DNA segments to which they are operably linked.

The nucleic acids encoding enzymes associated with the present disclosure can be
5 obtained from a variety of sources. As one of ordinary skill in the art would be aware, homologous genes for these enzymes exist in many species and can be identified by homology searches, for example, through using the internet-based Basic Local Alignment Search Tool (BLAST) made available by the National Center for Biotechnology Information (NCBI). Nucleic acids encoding these enzymes can be amplified by polymerase chain reaction (PCR)
10 from DNA from any source which contains the given enzyme, for example, using degenerate primers, as would be understood by one of ordinary skill in the art. In some embodiments, nucleic acids encoding a given enzyme can be synthetic. Any means of obtaining nucleic acids encoding the enzymes discussed herein are compatible with aspects of the present disclosure.

Expression of the enzymes of the present disclosure may be determined using routine
15 methods known to those of ordinary skill in the art. These methods include, but are not limited to, direct RNA amplification, reverse transcription of RNA to cDNA, real-time RT-PCR, amplification of cDNA, hybridization, and immunologically based assay methods, which include, but are not limited to Western blotting, immunohistochemistry, antibody sandwich capture assay, ELISA, and enzyme-linked immunospot assay (EliSpot assay). For example,
20 the determination of the presence of levels of nucleic acid molecules of the invention in a sample such as a tissue or cell lysate can be carried out via any standard nucleic acid determination assay, including PCR, or assaying with labeled hybridization probes. Such hybridization methods include, but are not limited to, microarray techniques.

The present disclosure thus involves, in one aspect, methods involving enzymes,
25 nucleic acids encoding those enzymes, functional modifications and variants of the foregoing, as well as uses relating thereto. Homologs and alleles of the nucleic acids described herein can be identified by conventional techniques (*see, e.g., Molecular Cloning: A Laboratory Manual*, J. Sambrook, et al., eds., Second Edition, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York, 1989, or *Current Protocols in Molecular Biology*, F.M. Ausubel, et al.,
30 eds., John Wiley & Sons, Inc., New York.

Aspects of the present disclosure relate to the expression of one or more enzymes in a genetically engineered cell followed by cell lysis. In some embodiments, one or more enzymes

- 23 -

are expressed and optionally sequestered in the cytoplasm of a cell prior to lysis. In some embodiments, one or more enzymes are expressed and optionally sequestered in the periplasmic space. Sequestration of enzymes in the periplasmic space is known in the art, see, *e.g.*, PCT Application No. PCT/US2011/035639, incorporated herein by reference. Upon lysis
5 of a cell to provide a cell lysate, the sequestered enzymes are free to react with one or more substrates also present in the lysate or different lysate.

Cell-free biotransformation of methane to biofuel and other chemical compounds

The cell-free biosynthetic system provided herein is useful for producing a biofuel or
10 other chemical compound (*e.g.*, pyruvate) from methane. For example, isobutanol of relatively high purity (99%), suitable for use as a second-generation biofuel, can be produced from methane in accordance with the invention. Isobutanol has several properties which make it ideal as a replacement for gasoline: its energy density is 98% that of gasoline, it does not readily absorb water (unlike ethanol), thereby avoiding corrosion of engine parts, and it can be
15 combined with gasoline in any proportion, permitting it to be “dropped-in” to the current combustible infrastructure. Isobutanol derived from natural gas can be combined with conventional gasoline for rapid commercialization, or it can be used as a full replacement of oil-derived fuel, depending on economic performance of the process and global oil prices.

Enzymes that catalyze the conversion of methane to formaldehyde include, without
20 limitation, methane monooxygenase (catalyzes methane to methanol) and methanol dehydrogenase (catalyzes methanol to formaldehyde). In some embodiments, methane monooxygenase and methanol dehydrogenase are provided in one or more cell lysate(s) obtained from methanotrophs. In other embodiments, methane monooxygenase is provided in a cell lysate obtained from methanotrophs, while methanol dehydrogenase is provided in a cell
25 lysate from recombinant bacteria, such as, for example, *Escherichia coli*. The invention also contemplates providing methane monooxygenase and/or methanol dehydrogenase exogenously in, for example, purified or partially purified form.

Enzymes that catalyze the conversion of formaldehyde to pyruvate include, without limitation, hexulose-6-phosphate synthase, 6-phospho-3-hexuloisomerase, 6-
30 phosphofructokinase, fructose biphosphate aldolase, triose phosphate isomerase, transketolase, ribose-5-phosphate isomerase, ribose-5-phosphate 3-epimerase, glyceraldehyde 3-phosphate

- 24 -

dehydrogenase, phosphoglycerate kinase, phosphoglycerate mutase, enolase, and pyruvate kinase.

Enzymes that catalyze the conversion of pyruvate to isobutanol include, without limitation, acetolactate synthase, acetohydroxy acid isomeroreductase, dihydroxy acid

5 dehydratase, alpha-ketoisovalerate decarboxylase, and isobutanol dehydrogenase.

Enzymes for use in accordance with the invention are shown in Table I.

Table I

Substrate(s)	Product(s)	Enzyme	EC number(s)
methane	methanol	methane monooxygenase	1.14.13.25, 1.14.18.3
methanol	formaldehyde	methanol dehydrogenase	1.1.1.244, 1.1.2.7, 1.1.99.37
formaldehyde, D-ribulose 5-phosphate	hexulose 6-phosphate	hexulose-6-phosphate synthase	4.1.2.43
hexulose 6-phosphate	beta-D-fructofuranose 6-phosphate	6-phospho-3-hexuloisomerase	5.3.1.27
beta-D-fructofuranose 6-phosphate	fructose-1,6-bisphosphate	6-phosphofructokinase	2.7.1.11
fructose-1,6-bisphosphate	D-glyceraldehyde 3-phosphate, dihydroxyacetone phosphate	fructose bisphosphate aldolase	4.1.2.13
dihydroxyacetone phosphate	D-glyceraldehyde 3-phosphate	triose phosphate isomerase	5.3.1.1
D-glyceraldehyde 3-phosphate, beta-D-fructofuranose 6-phosphate	D-erythrose 4-phosphate, D-xylulose 5-phosphate	transketolase	2.2.1.1
D-erythrose 4-phosphate, beta-D-fructofuranose 6-phosphate	D-glyceraldehyde 3-phosphate, D-sedoheptulose 7-phosphate	transaldolase	2.2.1.2
D-glyceraldehyde 3-phosphate, D-sedoheptulose 7-phosphate	D-ribose 5-phosphate, D-xylulose 5-phosphate	transketolase	2.2.1.1
D-ribose 5-phosphate	D-ribulose 5-phosphate	ribose-5-phosphate isomerase	5.3.1.6
D-xylulose 5-phosphate	D-ribulose 5-phosphate	ribulose-5-phosphate 3-epimerase	5.1.3.1
D-glyceraldehyde 3-phosphate	1,3-bisphospho-D-glycerate	glyceraldehyde 3-phosphate dehydrogenase	1.2.1.12
1,3-bisphospho-D-glycerate	3-phospho-D-glycerate	phosphoglycerate kinase	2.7.2.3
3-phospho-D-glycerate	2-phospho-D-glycerate	phosphoglycerate mutase	5.4.2.11/5.4.2.12
2-phospho-D-glycerate	phosphoenolpyruvate	enolase	4.2.1.11
phosphoenolpyruvate	pyruvate	pyruvate kinase	2.7.1.40
2 pyruvate	(S)-2-acetolactate	acetolactate synthase	2.2.1.6
(S)-2-acetolactate	2,3-dihydroxy-3-methylbutanoate	acetohydroxy acid isomeroreductase	1.1.1.86

- 25 -

Substrate(s)	Product(s)	Enzyme	EC number(s)
2,3-dihydroxy-3-methylbutanoate	3-methyl-2-oxobutanoate	dihydroxy acid dehydratase	4.2.1.9
3-methyl-2-oxobutanoate	isobutanal	alpha-ketoisovalerate decarboxylase	4.1.1.72
isobutanal	isobutanol	isobutanol dehydrogenase	1.1.1.1

It should be understood that any one or more of the enzymes provided herein may be used in any one or more cell lysates of a cell-free biosynthesis process of the invention. The invention also contemplates the use of one or more cell lysates, each containing, for example, a different combination of enzymes that convert a substrate (*e.g.*, methane) to a product (*e.g.*, isobutanol) or product intermediate (*e.g.*, formaldehyde or pyruvate).

An example of a biosynthetic pathway for cell-free conversion of methane to isobutanol in accordance with the invention is depicted in Figure 5. The cell-free approach provided herein, which combines cell extracts from at least two different bacterial cells, including a methanotroph (*e.g.*, *Methylococcus capsulatus* Bath), minimizes the issues associated with engineering a single organism with all of the enzymes necessary for conversion of methane to biofuel. Nonetheless, in some embodiments, a single organism may be used to express all enzymes necessary for the conversion of methane to a biofuel or other chemical compound. In the example depicted in Figure 5, cell lysates from *M. capsulatus* Bath cells and from *E. coli* cells are used. The first step, conversion of methane to methanol, is catalyzed by the methane monooxygenase (MMO) provided in the lysate of the methanotroph. The second step, conversion of methanol to formaldehyde, is catalyzed by a heterologous (*e.g.*, from different species) NAD-linked methanol dehydrogenase (*e.g.*, from *Bacillus methanolicus*, de Vries GE, *et al.* 1992 *J Bacteriol* 174:5346-53, incorporated by reference herein), which replaces methanol dehydrogenase native to the methanotroph. This NAD-linked methanol dehydrogenase provides the necessary NADH reductant for the methane monooxygenase, resulting in a net zero consumption of electrons for conversion of methane to formaldehyde. Thus, in some embodiments, methanotrophs of a cell-free system of the invention may express methane monooxygenase and an exogenous, or recombinant, NAD-linked methanol dehydrogenase.

In native methanotroph metabolism, formaldehyde is further oxidized by formaldehyde dehydrogenase to formate, then to CO₂ to generate energy for methanotroph cell function. This is a limitation in the use of intact methanotrophs for bioproduction. Because only MMO

- 26 -

and methanol dehydrogenase from the methanotroph are typically used in the cell-free process of the invention, this carbon route, in some embodiments, can be eliminated without affecting methanotroph function by targeting the formaldehyde dehydrogenase for cleavage by a protease expressed in recombinant bacterial (*e.g.*, *E. coli*) cells.

5 In some embodiments, the NAD-linked methanol dehydrogenase is not expressed in the methanotroph but rather in another bacterial cell, such as *E. coli*, as discussed herein.

Additional steps depicted in Figure 5 for the conversion of formaldehyde to isobutanol are catalyzed by enzymes provided in an extract of another recombinant cell(s) (*e.g.*, *E. coli*). Two ribulose monophosphate cycle enzymes, hexulose-6-phosphate synthase and 6-phospho-
10 3-hexuloisomerase, are overexpressed in the cell(s) along with a ketoisovalerate decarboxylase and an isobutanol oxidoreductase, which are required for the conversion of pyruvate to isobutanol. In addition to eliminating further oxidation of formaldehyde to CO₂, protease targeting methods may be used in certain embodiments to eliminate flux from pyruvate to other molecules by targeting pyruvate dehydrogenase for cleavage by a protease. Thus, in some
15 embodiments, recombinant bacteria of the cell-free system of the invention may express any one or more of hexulose-6-phosphate synthase, 6-phospho-3-hexuloisomerase, ketoisovalerate decarboxylase, and isobutanol oxidoreductase. In some embodiments, the recombinant bacteria may also express a protease that targets an engineered formaldehyde dehydrogenase and/or a pyruvate dehydrogenase. In some embodiments, a protease may be added
20 exogenously.

Delivering methane and oxygen to the methane monooxygenase may be accomplished by first diffusing the gases into the water phase, in some embodiments. By removing the cell wall barrier that has to be traversed in a living cell, the cell-free process provided herein simplifies the mass transfer process. In one embodiment, the gases are fed separately into a
25 reactor in order to avoid gas mixing until they have entered the aqueous phase, thereby avoiding problems associated with spontaneous combustion. An example of a stable multiphase system is depicted in Figure 1. This diagram includes the decane phase, which is used to help deliver the methane and to absorb the isobutanol as it is being produced. Small bubbles of methane are introduced into the decane phase before it enters the cell-free reactor.
30 Methane has much higher solubility in decane than in water, thus the decane surface area in contact with the aqueous phase helps to diffuse the methane into the aqueous phase.

- 27 -

Nonetheless, in some embodiments, some of the methane gas is injected directly into the aqueous phase.

To assess the formation and stability of the gas/liquid and liquid/liquid interfaces, for example, the formation and stability of methane gas bubbles, the apparatus depicted in Figure 7 may be used in some embodiments. Methane may be injected into the cell extract solution as it flows past the injection nozzles. An initial high speed camera permits accurate measurement of the size and uniformity of the bubbles formed. This fluid then passes into a holding loop and, after a set interval, is monitored by a second high speed camera to measure the stability of the gas bubbles. In some embodiments, the incubation time after bubble formation is adjusted by changing the size of the holding loop and/or the flow rate. The apparatus depicted in Figure 7 may also be used to examine the effects of nozzle design and injection pressure as well as the effect of multiple nozzles on bubble size and stability.

In some embodiments, fatty acids and/or surfactants are used to help form and stabilize the gas/liquid interface. Without being bound by theory, the hydrophobic tails and hydrophilic carboxylic acid heads of fatty acids cause them to collect at the gas/liquid interface to stabilize it.

In some embodiments, methane gas bubbles produced are smaller than 3 μm to 7 μm in diameter. For example, the methane gas bubbles produced may be smaller than 3 μm , 4 μm , 5 μm , 6 μm , or 7 μm in diameter. In some embodiments, the methane gas bubbles remain smaller than 8 μm to 12 μm in diameter for a certain period of time after initial bubble formation. For example, the methane gas bubbles may remain smaller than 8 μm , 9 μm , 10 μm , 11 μm , or 12 μm for at least one, two, three, four or five minutes after initial bubble formation.

Further, the apparatus depicted in Figure 7 may be used to study the formation and stability of methane in decane foam as well as the formation of the emulsion diagrammed in Figure 1. As discussed above, surface active agents may be used to help form and stabilize the desired interfaces that define the methane bubbles in the decane phase as well as the decane droplets in the cell-free reactor emulsion.

To assess the safety and productivity of the cell-free biotransformation processes of the invention, the apparatus shown in Figure 8 may be used in some embodiments. This reactor permits separate adjustment of methane and air injection rates and pressures. These vessels are typically small with working volumes of 10 to 100 ml, which helps to conserve extract and to

- 28 -

minimize safety risks. The back pressure is controlled, and temperature and dissolved oxygen monitored. The cooling/heating loop is not shown in the reactor depicted in Figure 8. Such reactors may be insulated to permit calculation of heat balance by monitoring the flow rate and temperature rise (or decrease) of the cooling/heating fluid. Further, the entering gases flow rates may be recorded and the off-gas composition may be monitored by an exit gas mass spec such that material balances indicate methane and oxygen utilization rates. Liquid samples may be analyzed by HPLC or a separate mass spec to indicate accumulation of intermediates and products. These values may be compared with measured heat evolution rates to provide confidence for performance assessments.

In some embodiments, the apparatus shown in Figure 8 may be used without the decane phase to provide an initial assessment of any safety limits related to possible gas mixing. As confidence is gained and safety demonstrated, decane alone may be introduced to form an emulsion. In some embodiments, decane/methane foams may be injected to fully replicate the final system.

In other embodiments, the apparatus shown in Figure 8 may be used to assess MMO activities and methanotroph growth. In yet other embodiments, a multiplexed reactor system, as shown in Figure 9, may be used to measure MMO activity and cell extract performance under working conditions to improve relevant performance characteristics and longevities.

In addition, in some embodiments, it may be economically advantageous to grow the methanotrophs without the support of paraffin or other additives. Thus, in some embodiments, oxygen and methane may be delivered to the organism to support acceptable growth rates and cell accumulation levels without using paraffin.

The cell-free biotransformation processes provided herein may be used to produce biofuel or other chemical compounds at a production rate of at least 1 g/L-h, at least 5 g/L-h, at least 10 g/L-h, at least 15 g/L-h, or at least 25 g/L-h. For example, the biofuel or other chemical compound may be produced at a rate of 0.5 g/L-h, 1 g/L-h, 2 g/L-h, 3 g/L-h, 4 g/L-h, 5 g/L-h, 6 g/L-h, 7 g/L-h, 8 g/L-h, 9 g/L-h, 10 g/L-h, 11 g/L-h, 12 g/L-h, 13 g/L-h, 14 g/L-h, 15 g/L-h, 16 g/L-h, 17 g/L-h, 18 g/L-h, 19 g/L-h, 20 g/L-h, 21 g/L-h, 22 g/L-h, 23 g/L-h, 24 g/L-h, 25 g/L-h, or more. In some embodiments, the biofuel or other chemical compound may be produced at a rate of 1 to 5 g/L-h, 1 to 10 g/L-h, 1 to 25 g/L-h, 5 to 10 g/L-h, 5 to 25 g/L-h, or 10 to 25 g/L-h.

- 29 -

Large-scale production

Some aspects of the invention relate to large-scale (*e.g.*, using reaction volumes of greater than 10 L) production of biofuels (*e.g.*, isobutanol) or other chemical compounds. Prior to the invention, several factors prevented the application of cell-free biotransformation to, for example, industrial and/or commercial processes. Such factors included the inability to control the redox balance in cell-free systems and the lack of molecular biology tools that would enable the rapid and efficient control of carbon flux. The present disclosure enables the use of enzymes from various microorganisms in a fully controlled and tailored environment capable of processing natural gas, in particular methane, into high-value chemicals, such as, for example, isobutanol.

In some embodiments, cell-free biotransformation reactions are conducted using reaction volumes (*e.g.*, in a bioreactor) of 10 L to 1,000 L. For example, reaction volumes may be 10 L to 100 L, 10 L to 500 L, 100 L to 500 L, 100 L to 1000 L, or 500 L to 1000 L.

In some embodiments, cell-free biotransformation processes provided herein are used to produce at least 1 L of isobutanol per week. In some embodiments, cell-free biotransformation processes provided herein are used to produce 1 to 100 metric tons of isobutanol per week. For example, a cell-free system may be used to produce 10, 20, 30, 40, 50, 60, 70, 80, 90, or 100 metric tons of isobutanol per week.

Flexible feedstock in cell free systems

Cell-free systems enable complete decoupling of cell growth from target production. Typical challenges encountered when using mixed carbon sources (*e.g.*, glucose and xylose as mixed C6 and C5 sugars from lignocellulosic biomass) in bioproduction include catabolite repression, membrane transport, and cytotoxicity. Cell-free systems avoid these issues, enabling more efficient use of these inexpensive sugar sources. Use of other technologies such as protease targeting enables efficient simultaneous utilization of C5 and C6 sugars for target molecule carbon skeletons as well as for sources of energy and reducing equivalents.

The following examples of isoprene production are included: (i) mixed C6 (glucose) and C1 (methane) feedstocks, and (ii) C1 (methane) feedstock.

- 30 -

CLAIMS

What is claimed is:

1. A cell-free method for large-scale conversion of methane to isobutanol, comprising:

5 combining, in a bioreactor at elevated pressure, methane, oxygen, and cell lysates containing methane monooxygenase, methanol dehydrogenase, and enzymes that catalyze the conversion of formaldehyde to isobutanol, to form a cell-free reaction mixture; and

incubating under suitable conditions the cell-free reaction to convert methane to isobutanol.

10 2. The cell-free method of claim 1, wherein the methane monooxygenase is obtained from a methanotroph.

3. The cell-free method of claim 1 or 2, wherein the methanol dehydrogenase is a
15 heterologous NAD-linked methanol dehydrogenase.

4. The cell-free method of any one of claims 1-3, wherein the enzymes that catalyze the conversion of formaldehyde to isobutanol are derived from recombinant bacteria.

20 5. The cell-free method of claim 4, wherein the recombinant bacteria are recombinant *Escherichia coli*.

6. The cell-free method of any one of claims 1-5, wherein the enzymes that catalyze the conversion of formaldehyde to isobutanol comprise one or more of hexulose-6-phosphate
25 synthase, 6-phospho-3-hexuloisomerase, 6-phosphofructokinase, fructose biphosphate aldolase, triose phosphate isomerase, transketolase, ribose-5-phosphate isomerase, ribose-5-phosphate 3-epimerase, glyceraldehyde 3-phosphate dehydrogenase, phosphoglycerate kinase, phosphoglycerate mutase, enolase, pyruvate kinase, acetolactate synthase, acetohydroxy acid

- 31 -

isomeroreductase, dihydroxy acid dehydratase, alpha-ketoisovalerate decarboxylase, and isobutanol dehydrogenase.

7. The cell-free method of any one of claims 1-6, wherein the pressure is at least 1 bar.

5

8. The cell-free method of claim 7, wherein the pressure is at least 2 bar.

9. The cell-free method of claim 8, wherein the pressure is at least 5 bar.

10 10. The cell-free method of any one of claims 1-9, wherein the isobutanol is produced at a production rate of at least 1 g/L-h.

11. The cell-free method of claim 10, wherein the isobutanol is produced at a production rate of at least 10 g/L-h.

15

12. The cell-free method of claim 11, wherein the isobutanol is produced at a production rate of at least 25 g/L-h.

13. The cell-free method of any one of claims 1-12, wherein the bioreactor comprises a gas
20 phase and an aqueous phase.

14. The cell-free method of claim 13, wherein the bioreactor further comprises an organic solvent.

25 15. The cell-free method of claim 14, wherein the organic solvent is an alkane.

- 32 -

16. The cell-free method of claim 15, wherein the alkane is selected from the group consisting of pentane, hexane, heptane, octane, nonane, and decane.

17. A cell-free method for large-scale conversion of methane to pyruvate, comprising:

5 combining, in a bioreactor at elevated pressure, methane, oxygen, and cell lysates containing methane monooxygenase, methanol dehydrogenase, and enzymes that catalyze the conversion of formaldehyde to pyruvate, to form a cell-free reaction mixture; and

incubating under suitable conditions the cell-free reaction to convert methane to pyruvate.

10 18. The cell-free method of claim 17, wherein the methane monooxygenase is obtained from a methanotroph.

19. The cell-free method of claim 17 or 18, wherein the methanol dehydrogenase is a
15 heterologous NAD-linked methanol dehydrogenase.

20. The cell-free method of any one of claims 17-19, wherein the enzymes that catalyze the conversion of formaldehyde to isobutanol are derived from recombinant bacteria.

20 21. The cell-free method of claim 20, wherein the recombinant bacteria are recombinant *Escherichia coli*.

22. The cell-free method of any one of claims 17-21, wherein the enzymes that catalyze the conversion of formaldehyde to pyruvate comprise one or more of hexulose-6-phosphate
25 synthase, 6-phospho-3-hexuloisomerase, 6-phosphofructokinase, fructose biphosphate aldolase, triose phosphate isomerase, transketolase, ribose-5-phosphate isomerase, ribose-5-phosphate 3-epimerase, glyceraldehyde 3-phosphate dehydrogenase, phosphoglycerate kinase, phosphoglycerate mutase, enolase, and pyruvate kinase.

- 33 -

23. The cell-free method of any one of claims 17-22, wherein the pressure is at least 1 bar.

24. The cell-free method of claim 23, wherein the pressure is at least 2 bar.

5

25. The cell-free method of claim 24, wherein the pressure is at least 5 bar.

26. The cell-free method of any one of claims 17-25, wherein the isobutanol is produced at a production rate of at least 1 g/L-h.

10

27. The cell-free method of claim 26, wherein the isobutanol is produced at a production rate of at least 10 g/L-h.

28. The cell-free method of claim 27, wherein the isobutanol is produced at a production rate of at least 25 g/L-h.

15

29. The cell-free method of any one of claims 17-28, wherein the bioreactor comprises a gas phase and an aqueous phase.

30. The cell-free method of claim 29, wherein the bioreactor further comprises an organic solvent.

20

31. The cell-free method of claim 30, wherein the organic solvent is an alkane.

32. The cell-free method of claim 31, wherein the alkane is selected from the group consisting of pentane, hexane, heptane, octane, nonane, and decane.

25

- 34 -

33. A cell-free method for large-scale conversion of methane to a biofuel or other chemical compound, comprising:

combining, in a bioreactor at elevated pressure, methane, oxygen, and cell lysates
5 containing methane monooxygenase, methanol dehydrogenase, enzymes that catalyze the conversion of formaldehyde to pyruvate, and enzymes that catalyze the conversion of pyruvate to a biofuel or other chemical compound, to form a cell-free reaction mixture; and

incubating under suitable conditions the cell-free reaction to convert methane to the biofuel or other chemical compound.

10

34. The cell-free method of claim 33, wherein the methane monooxygenase is obtained from a methanotroph.

15

35. The method of claim 33 or 34, wherein the methanol dehydrogenase is a heterologous NAD-linked methanol dehydrogenase.

20

36. The cell-free method of any one of claims 33-35, wherein the enzymes that catalyze the conversion of formaldehyde to pyruvate and/or pyruvate to the biofuel or other compound are derived from recombinant bacteria.

37. The cell-free method of claim 36, wherein the recombinant bacteria are recombinant *Escherichia coli*.

25

38. The cell-free method of any one of claims 33-37, wherein the enzymes that catalyze the conversion of formaldehyde to pyruvate comprise one or more of hexulose-6-phosphate synthase, 6-phospho-3-hexuloisomerase, 6-phosphofructokinase, fructose biphosphate aldolase, triose phosphate isomerase, transketolase, ribose-5-phosphate isomerase, ribose-5-

- 35 -

phosphate 3-epimerase, glyceraldehyde 3-phosphate dehydrogenase, phosphoglycerate kinase, phosphoglycerate mutase, enolase, and pyruvate kinase.

39. The cell-free method of any one of claims 33-38, wherein the pressure is at least 1 bar.

5

40. The cell-free method of claim 39, wherein the pressure is at least 2 bar.

41. The cell-free method of claim 40, wherein the pressure is at least 5 bar.

10 42. The cell-free method of any one of claims 33-41, wherein the isobutanol is produced at a production rate of at least 1 g/L-h.

43. The cell-free method of claim 42, wherein the isobutanol is produced at a production rate of at least 10 g/L-h.

15

44. The cell-free method of claim 43, wherein the isobutanol is produced at a production rate of at least 25 g/L-h.

20 45. The cell-free method of any one of claims 33-44, wherein the bioreactor comprises a gas phase and an aqueous phase.

46. The cell-free method of claim 45, wherein the bioreactor further comprises an organic solvent.

25 47. The cell-free method of claim 46, wherein the organic solvent is an alkane.

- 36 -

48. The cell-free method of claim 47, wherein the alkane is selected from the group consisting of pentane, hexane, heptane, octane, nonane, and decane.

49. A cell-free system for large-scale conversion of methane to isobutanol, the system comprising a bioreactor that comprises:

a gas phase comprising methane and oxygen; and

an aqueous phase comprising a cell lysate containing methane monooxygenase, methanol dehydrogenase, and enzymes that catalyze the conversion of formaldehyde to isobutanol.

50. The cell-free system of claim 49, wherein the methane monooxygenase is obtained from a methanotroph.

51. The cell-free system of claim 49 or 50, wherein the methanol dehydrogenase is a heterologous NAD-linked methanol dehydrogenase.

52. The cell-free system of any one of claims 49-51, wherein the enzymes that catalyze the conversion of formaldehyde to isobutanol are derived from recombinant bacteria.

53. The cell-free system of claim 52, wherein the recombinant bacteria are recombinant *Escherichia coli*.

54. The cell-free system of any one of claims 49-53, wherein the enzymes that catalyze the conversion of formaldehyde to isobutanol comprise one or more of hexulose-6-phosphate synthase, 6-phospho-3-hexuloisomerase, 6-phosphofructokinase, fructose biphosphate aldolase, triose phosphate isomerase, transketolase, ribose-5-phosphate isomerase, ribose-5-phosphate 3-epimerase, glyceraldehyde 3-phosphate dehydrogenase, phosphoglycerate kinase, phosphoglycerate mutase, enolase, pyruvate kinase, acetolactate synthase, acetohydroxy acid

- 37 -

isomeroreductase, dihydroxy acid dehydratase, alpha-ketoisovalerate decarboxylase, and isobutanol dehydrogenase.

55. The cell-free system of any one of claims 49-54, wherein the bioreactor further
5 comprises an organic solvent.

56. The cell-free system of claim 55, wherein the organic solvent is an alkane.

57. The cell-free system of claim 56, wherein the alkane is selected from the group
10 consisting of pentane, hexane, heptane, octane, nonane, and decane.

58. A cell-free system for large-scale conversion of methane to pyruvate, the system comprising a bioreactor that comprises:
a gas phase comprising methane and oxygen; and
15 an aqueous phase comprising a cell lysate containing methane monooxygenase, methanol dehydrogenase, and enzymes that catalyze the conversion of formaldehyde to pyruvate.

59. The cell-free system of claim 58, wherein the methane monooxygenase is obtained
20 from a methanotroph.

60. The cell-free system of claim 58 or 59, wherein the methanol dehydrogenase is a heterologous NAD-linked methanol dehydrogenase.

25 61. The cell-free system of any one of claims 58-60, wherein the enzymes that catalyze the conversion of formaldehyde to isobutanol are derived from recombinant bacteria.

- 38 -

62. The cell-free system of claim 61, wherein the recombinant bacteria are recombinant *Escherichia coli*.

63. The cell-free system of any one of claims 58-62, wherein the enzymes that catalyze the conversion of formaldehyde to pyruvate comprise one or more of hexulose-6-phosphate synthase, 6-phospho-3-hexuloisomerase, 6-phosphofructokinase, fructose biphosphate aldolase, triose phosphate isomerase, transketolase, ribose-5-phosphate isomerase, ribose-5-phosphate 3-epimerase, glyceraldehyde 3-phosphate dehydrogenase, phosphoglycerate kinase, phosphoglycerate mutase, enolase, and pyruvate kinase.

64. The cell-free system of any one of claims 58-63, wherein the bioreactor further comprises an organic solvent.

65. The cell-free system of claim 64, wherein the organic solvent is an alkane.

66. The cell-free system of claim 65, wherein the alkane is selected from the group consisting of pentane, hexane, heptane, octane, nonane, and decane.

67. A cell-free system for large-scale conversion of methane to pyruvate, the system comprising a bioreactor that comprises:

a gas phase comprising methane and oxygen; and

an aqueous phase comprising a cell lysate containing methane monooxygenase, methanol dehydrogenase, enzymes that catalyze the conversion of formaldehyde to pyruvate, and enzymes that catalyze the conversion of pyruvate to a biofuel or other chemical compound, to form a cell-free reaction mixture.

68. The cell-free system of claim 67, wherein the methane monooxygenase is obtained from a methanotroph.

- 39 -

69. The cell-free system of claim 67 or 68, wherein the methanol dehydrogenase is a heterologous NAD-linked methanol dehydrogenase.

5 70. The cell-free system of any one of claims 67-69, wherein the enzymes that catalyze the conversion of formaldehyde to isobutanol are derived from recombinant bacteria.

71. The cell-free system of claim 70, wherein the recombinant bacteria are recombinant *Escherichia coli*.

10

72. The cell-free system of any one of claims 67-71, wherein the enzymes that catalyze the conversion of formaldehyde to pyruvate comprise one or more of hexulose-6-phosphate synthase, 6-phospho-3-hexuloisomerase, 6-phosphofructokinase, fructose biphosphate aldolase, triose phosphate isomerase, transketolase, ribose-5-phosphate isomerase, ribose-5-phosphate 3-epimerase, glyceraldehyde 3-phosphate dehydrogenase, phosphoglycerate kinase, 15 phosphoglycerate mutase, enolase, and pyruvate kinase.

73. The cell-free system of any one of claims 67-71, wherein the bioreactor further comprises an organic solvent.

20

74. The cell-free system of claim 73, wherein the organic solvent is an alkane.

75. The cell-free system of claim 74, wherein the alkane is selected from the group consisting of pentane, hexane, heptane, octane, nonane, and decane.

25

76. A cell lysate of any one of claims 1-75.