

Description

This invention relates to novel 2-(1-piperazinyl)-4-phenylcycloalkanopyridine derivatives having psychotropic activity, processes for the production thereof, and a pharmaceutical composition containing the said compound as an active ingredient.

Prior Art

There have hitherto been known some 2-(1-piperazinyl)-4-arylpyridine derivatives and related compounds which have pharmacological activities. For example, U.S. Patent No. 4,469,696 [Japanese First Publication (Kokai) No. 58-963] discloses 2-(1-piperazinyl)-4-arylpyridine derivatives, however, the compounds disclosed therein are merely the compounds having an aryl group or a 2-furyl group at 5-position of pyridine ring, of which structure is completely different from that of the compounds of the present invention. Besides, the pharmacological activity of these compounds disclosed in the above reference is lipid absorption-inhibitory action which is also completely different from that of the compounds of the present invention. It is clear from the experiments by the present inventors, whereby the 4,5-bis-(4-fluorophenyl)-6-methyl-2-(4-phenyl-1-piperazinyl)pyridine disclosed in the above mentioned U.S. patent did not show any psychotropic activity, that is, it shows no inhibitory effect on an apomorphine-induced vomiting, which can be an index for anti-psychotic drug and no binding property to dopamine (D₂) and serotonin (S₂) receptors.

Moreover, a pyridine derivative having a piperazinyl group and a phenyl group at 2-position and 4-position of pyridine ring thereof respectively, 2-[4-(4-methylbenzyl)-1-piperazinyl]-4-phenylpyridine has been reported to have a weak activity as an anti-psychotic drug or as a neuroleptic drug, [cf: U.S. Patent No. 4,831,034, Japanese First Publication (Kokai) No. 63-48267]. However, the structure of this compound is obviously different from that of the compounds of the present invention in respect to that this compound does not form the condensed ring between 5- and 6-positions of pyridine ring.

EP-A-68 259 discloses compounds which are structurally similar to the compounds of the present invention, but does not define exactly the carbocyclic ring condensed to the pyridine, nor does it define exactly the ring attached in the 2-position of the pyridine. Thus the subject-matter of this invention is novel. In addition, EP-A-68 259 does not exemplify any compound having a carbocyclic ring condensed to the pyridine. Therefore the skilled man has no hint that a carbocyclic ring attached to the pyridine would lead to compounds having a psychotropic activity, which is different from the activity reported for the compounds of

GB-A-2 017 698 describes a compound differing from the compounds from the present application in that the ring condensed to the pyridine ring is unsaturated. GB-A-2 017 698 does not suggest that a modification of this unsaturated ring would result in compounds having psychotropic activity.


Brief Description of the Invention

The present inventors have extensively searched for compounds having an activity on the central nervous system and have found that novel 2-(1-piperazinyl)-4-phenylcycloalkanopyridine derivatives of the formula (I) disclosed hereinafter have excellent psychotropic activity and are useful as an anti-psychotic drug or an anti-anxiety drug and further as a drug for the treatment of cerebral insufficiency diseases.

An object of the invention is to provide novel 2-(1-piperazinyl)-4-phenylcycloalkanopyridine derivatives having excellent psychotropic activity. Another object of the invention is to provide processes for the production of these compounds. A further object of the invention is to provide a pharmaceutical composition being useful as an anti-psychotic or anti-anxiety drug containing said compound as an active ingredient. These and other objects and advantages of the invention will be apparent to those skilled in the art from the following description.

Detailed Description of the Invention

The novel 2-(1-piperazinyl)-4-phenylcycloalkanopyridine derivatives of the present invention have the following formula (I):



15 wherein n is 3, 4, 5, 6 or 7,

R¹ is hydrogen atom, C₁-C₁₀ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₃-C₆ cycloalkyl-(C₁-C₄)alkyl, hydroxy-(C₂-C₆) alkyl, C₁-C₃ alkoxy-(C₂-C₆) alkyl, acyloxy-(C₂-C₆) alkyl, unsubstituted or substituted aroyl-(C₁-C₆) alkyl, unsubstituted or substituted aryl, heteroaryl, or acyl,

20 R² and R³ are the same or different and are each hydrogen atom, halogen atom, C₁-C₆ alkyl, C₁-C₆ alkoxy, trifluoromethyl, or hydroxy,

R⁴, R⁵ and R⁶ are the same or different and are each hydrogen atom, C₁-C₆ alkyl, or phenyl, or two of R⁴, R⁵ and R⁶ combine to form a single bond or C₁-C₃ alkylene,

R⁷ and R⁸ are the same or different and are each hydrogen atom or C₁-C₃ alkyl,

m is 2 or 3,

25 or an acid addition salt thereof.

The salt of the compounds of the formula (I) includes salts of inorganic acids (e.g. hydrochloride, hydrobromide, hydroiodide, sulfate, phosphate), and salts of organic acids (e.g. maleate, fumarate, citrate, oxalate, tartrate, lactate, benzoate, methanesulfonate). Besides, these salts may optionally be present in the form of a hydrate, and hence, the compounds of the present invention include also these hydrate

30 compounds. Besides, when the compounds of the formula (I) contain asymmetric carbons, these compounds include stereoisomers, a mixture thereof, and a racemic mixture, which are also included as the active compound in the present invention.

In the present specification and claims, the groups in the formulae denote the following groups.

35 The "alkyl" and "alkyl moiety" and "alkylene" include straight chain or branched chain alkyl groups or alkylene groups.

The "alkyl" includes methyl, ethyl, propyl, isopropyl, butyl, isobutyl, sec-butyl, pentyl, isopentyl, neopentyl, hexyl, isohexyl, heptyl, octyl, nonyl, decyl.

The "alkylene" includes methylene, ethylene, propylene.

40 The "halogen atom" includes fluorine, chlorine, bromine, and iodine.

The "alkoxy" includes methoxy, ethoxy, propoxy, isopropoxy, butoxy, isobutoxy, pentyloxy, hexyloxy.

The "cycloalkyl" includes, for example, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl.

The "cycloalkyl-alkyl" includes cyclopropylmethyl, cyclobutylmethyl, cyclopentylmethyl.

45 The "hydroxyalkyl" includes 2-hydroxyethyl, 2-hydroxypropyl, 3-hydroxypropyl, 2,3-dihydroxypropyl, 4-hydroxybutyl, 2-hydroxybutyl.

The "alkoxyalkyl" includes methoxymethyl, methoxyethyl, ethoxyethyl.

The "alkenyl" includes vinyl, allyl, 1-propenyl, isopropenyl, 1-butenyl, 2-butenyl.

The "alkynyl" includes ethynyl, propargyl.

50 The "aryl" includes phenyl and naphthyl. The "unsubstituted or substituted aryl" includes aryl groups having no substituent or one or two substituents selected from halogen atom, C₁-C₃ alkyl, C₁-C₃ alkoxy, and trifluoromethyl, and the examples are phenyl, 4-fluorophenyl.

The "unsubstituted or substituted aroylalkyl" includes aroylalkyl groups in which the aryl moiety is the above-mentioned unsubstituted or substituted aryl, and the examples are benzoylmethyl, p-fluorobenzoyl-propyl.

55 The "heteroaryl" means a monocyclic or bicyclic heterocyclic group containing at least one hetero atom selected from nitrogen, oxygen and sulfur, for example, furyl, thienyl, pyridyl, pyrimidyl, isoquinolyl.

The "acyl" includes C₁-C₄ alkanoyl, C₅-C₆ cycloalkylcarbonyl, a benzoyl group which may optionally be substituted by halogen, C₁-C₃ alkyl or C₁-C₃ alkoxy, a heteroarylcarbonyl group in which the heteroaryl

moiety is the above-mentioned heteroaryl, and the examples are formyl, acetyl, propionyl, butyryl, cyclohexanecarbonyl, benzoyl, nicotinoyl, isonicotinoyl, 4-fluorobenzoyl, furoyl, thenoyl.


The "acyloxyalkyl" includes acyloxyalkyl groups in which the acyl moiety is the above-mentioned acyl, and the examples are 2-acetyloxyethyl, 3-acetoxypropyl, benzoyloxyethyl.

5 Among the compounds of the formula (I) of the present invention, preferred compounds are those of the formula (I) wherein R^1 is hydrogen atom, C_1 - C_{10} alkyl, C_3 - C_8 cycloalkyl, hydroxy- (C_2-C_6) alkyl, C_1 - C_3 alkoxy- (C_2-C_4) alkyl, C_2 - C_4 alkanoyloxy- (C_2-C_6) alkyl, a benzoyl- (C_2-C_5) alkyl group in which the phenyl moiety may optionally be substituted by halogen, C_1 - C_3 alkyl or C_1 - C_3 alkoxy, C_2 - C_4 alkenyl, C_2 - C_4 alkynyl, pyridyl, pyrimidyl, C_2 - C_5 alkanoyl or furoyl, R^2 and R^3 are the same or different and are each
10 hydrogen atom, halogen atom, methyl or methoxy, R^4 , R^5 and R^6 are the same or different and are each hydrogen atom or C_1 - C_4 alkyl, or two of them combine to form C_1 - C_2 alkylene, R^7 and R^8 are the same or different, and are each hydrogen atom or C_1 - C_3 alkyl, and m is 2, and an acid addition salt thereof.

Further preferred compounds of the present invention are the compounds of the formula (I) wherein R^1 is hydrogen atom, C_1 - C_8 alkyl, C_3 - C_8 cycloalkyl, hydroxy- (C_2-C_6) alkyl, C_1 - C_2 alkoxy- (C_2-C_3) alkyl,
15 acetyloxy- (C_2-C_4) alkyl, C_3 - C_4 alkenyl, or C_2 - C_3 alkanoyl, R^2 and R^3 are the same and both are hydrogen atom or halogen atom, or one of them is hydrogen atom and the other is halogen atom, methyl, or methoxy, and an acid addition salt thereof.

More preferred compounds of the present invention are the compounds of the formula (I) wherein R^2 and R^3 are the same or different and are each hydrogen atom or fluorine atom, and an acid addition salt
20 thereof.

Particularly preferred compounds are the compounds of the following formula (I-1):




wherein n_1 is 3, 6 or 7,

R^{11} is hydrogen atom, C_1 - C_8 alkyl, C_3 - C_6 cycloalkyl, hydroxy- (C_2-C_4) alkyl, C_1 - C_2 alkoxy- (C_2-C_3) alkyl, or C_3 alkenyl,

40 R^{21} and R^{31} are the same or different and are each hydrogen atom or fluorine atom,

R^{71} and R^{81} are the same or different and are each hydrogen atom or C_1 - C_3 alkyl, or an acid addition salt thereof, and of the following formula (I-2):



wherein n_2 is 4 or 5,


R^{12} is hydrogen atom, C_1 - C_8 alkyl, C_3 - C_8 cycloalkyl, hydroxy-(C_2 - C_4) alkyl, C_1 - C_2 alkoxy-(C_2 - C_3) alkyl or C_3 alkenyl,

R^{22} and R^{32} are the same or different and are each hydrogen atom or fluorine atom,

5 R^{42} , R^{52} and R^{62} are the same or different and are each hydrogen atom or C_1 - C_4 alkyl, or two of them combine to form C_1 - C_2 alkylene, or an acid addition salt thereof.

Especially most preferred compounds in the present invention are the compounds of the formula (I-1) wherein n_1 is 6, R^{11} is C_1 - C_6 alkyl, C_3 - C_6 cycloalkyl, or hydroxy-(C_2 - C_4) alkyl, R^{21} and R^{31} are the same or
10 different and are each hydrogen atom or fluorine atom substituted at 2- or 4-position, R^{71} and R^{81} are each hydrogen atom, and a salt thereof, and the compounds of the formula (I-2) wherein n_2 is 5, R^{12} is C_1 - C_6 alkyl, C_3 - C_6 cycloalkyl or hydroxy-(C_2 - C_4) alkyl, R^{22} and R^{32} are the same or different and are each hydrogen atom or fluorine atom substituted at 2- or 4-position, R^{42} , R^{52} and R^{62} are all hydrogen atom, or two of them combine to form C_1 - C_2 alkylene, and another one is hydrogen atom, and an acid addition salt thereof.

15 The most preferred compounds of the present invention are the compounds of the following formula (I-3):



wherein R^{13} is methyl, ethyl, propyl, butyl, pentyl or hydroxyethyl, and R^{33} is hydrogen atom or fluorine atom, and a salt thereof.


Specifically preferred compounds of the present invention are as follows.

35 2-(4-Methyl-1-piperazinyl)-4-(4-fluorophenyl)-5,6,7,8,9,10-hexahydrocycloocta[b]pyridine
2-(4-Methyl-1-piperazinyl)-4-(2,4-difluorophenyl)-5,6,7,8,9,10-hexahydrocycloocta[b]pyridine
2-(4-Ethyl-1-piperazinyl)-4-(4-fluorophenyl)-5,6,7,8,9,10-hexahydrocycloocta[b]pyridine
2-(4-Ethyl-1-piperazinyl)-4-(2,4-difluorophenyl)-5,6,7,8,9,10-hexahydrocycloocta[b]pyridine
40 2-(4-n-Propyl-1-piperazinyl)-4-(4-fluorophenyl)-5,6,7,8,9,10-hexahydrocycloocta[b]pyridine
2-(4-n-Propyl-1-piperazinyl)-4-(2,4-difluorophenyl)-5,6,7,8,9,10-hexahydrocycloocta[b]pyridine
2-(4-n-Butyl-1-piperazinyl)-4-(4-fluorophenyl)-5,6,7,8,9,10-hexahydrocycloocta[b]pyridine
2-(4-n-Pentyl-1-piperazinyl)-4-(4-fluorophenyl)-5,6,7,8,9,10-hexahydrocycloocta[b]pyridine
2-[4-(2-Hydroxyethyl)-1-piperazinyl]-4-(4-fluorophenyl)-5,6,7,8,9,10-hexahydrocycloocta[b]pyridine

The compounds of the present invention can be prepared, for example, by the following processes.

(1) Process A

The compounds of the formula (I) are prepared by reacting a compound of the formula (II):



wherein X is a leaving atom or group, and R^2 , R^3 , R^4 , R^5 , R^6 and n are as defined above, with a compound

20 of the formula (III):



30 wherein R^1 , R^7 , R^8 and m are as defined above.

The leaving atom or group X in the formula (II) denotes any atom or group which can leave off in the form of HX under the reaction conditions together with hydrogen atom bonded to the nitrogen atom at 4-position of 1-substituted piperazines or homopiperazines. Examples of the leaving atom or group are halogen atoms, lower alkythio groups (e.g. methylthio, ethylthio, propylthio, butylthio), arylsulfonyloxy groups (e.g. benzenesulfonyloxy, p-toluenesulfonyloxy), and alkylsulfonyloxy groups (e.g. methanesulfonyloxy).

35

The reaction of the compound of the formula (II) and the compound of the formula (III) is carried out in an appropriate solvent or without using any solvent under atmospheric pressure or under pressure. Suitable examples of the solvent are aromatic hydrocarbons (e.g. toluene, xylene), ketones (e.g. methyl ethyl ketone), ethers (e.g. dioxane, diglyme), alcohols (e.g. ethanol, isopropyl alcohol, butanol), N,N-dimethylformamide, dimethylsulfoxide. The reaction is preferably carried out in the presence of a basic substance. Suitable examples of the basic substance are alkali metal carbonates (e.g. sodium carbonate, potassium carbonate), alkali metal hydrogen carbonates (e.g. sodium hydrogen carbonate, potassium hydrogen carbonate), tertiary amines (e.g. triethylamine), but an excess amount of the compound of the formula (III) may be used instead of using the basic substance. When the compound of the formula (III) is in the form of a hydrate, the hydrate may be used. The reaction temperature is usually in the range of 40 to 200 °C. The starting compound (II) can be prepared in the procedure as described in Reference Examples 1 to 94 hereinafter or in a similar process.


40

45

50 (2) Process B

The compounds of the formula (I) wherein R^1 is hydrogen atom can be prepared by hydrogenolysis of a compound of the formula (I-4):

55




wherein R^{14} is unsubstituted or substituted benzyl or benzyloxycarbonyl (the substituent on the phenyl ring of said benzyl and benzyloxycarbonyl is a member selected from lower alkyl, lower alkoxy and halogen atom), and R^2 , R^3 , R^4 , R^5 , R^6 , R^7 , R^8 , m and n are as defined above.


The hydrogenolysis of the compounds of the formula (I-4) is usually carried out by a conventional catalytic reduction in a solvent such as alcohols (e.g. ethanol) at room temperature under atmospheric pressure. The compounds of the formula (I-4) can be prepared by the same process as the above-mentioned Process A. The compounds of the formula (I-4) wherein R^{14} is benzyloxycarbonyl can alternatively be prepared by a conventional process from a compound of the formula (I) wherein R^{14} is methyl or benzyl having optionally a substituent.

(3) Process C


The compounds of the formula (I) wherein R^1 is hydrogen atom can also be prepared by reacting a compound of the formula (I-5):



wherein R^{15} is methyl or unsubstituted or substituted benzyl (the substituent on the phenyl ring of the benzyl is a member selected from lower alkyl, lower alkoxy and halogen atom), and R^2 , R^3 , R^4 , R^5 , R^6 , R^7 , R^8 , m and n are as defined above, with ethyl chlorocarbonate or 1-chloroethyl chlorocarbonate to give a compound of the formula (I-6):



15 wherein R^{16} is ethoxycarbonyl, 1-chloroethoxycarbonyl, and R^2 , R^3 , R^4 , R^5 , R^6 , R^7 , R^8 , m and n are as defined above, or a compound of the formula (I-7):



35 wherein R^{17} is acyl, and R^2 , R^3 , R^4 , R^5 , R^6 , R^7 , R^8 , m and n are as defined above, followed by hydrolysis of the compound of the formula (I-6) or (I-7).


40 The hydrolysis of the compound of the formula (I-6) or (I-7) is usually carried out by a conventional method, for example, by heating the compound in an appropriate solvent such as ethanol which is miscible with water in the presence of a basic substance (e.g. sodium hydroxide, potassium hydroxide) or an acid (e.g. hydrochloric acid, sulfuric acid). The hydrolysis of the compound of the formula (I-6) wherein R^{16} is 1-chloroethoxycarbonyl is usually carried out by heating the compound in methanol. The compounds of the formulae (I-5) and (I-7) can be prepared by the same process as the above-mentioned Process A.

(4) Process D

45 The compound of the formula (I) wherein R^1 is a group other than hydrogen atom can be prepared by reacting a compound of the formula (I-8):

50

55



15 wherein R^2 , R^3 , R^4 , R^5 , R^6 , R^7 , R^8 , m and n are as defined above, with a compound of the formula (IV):

R^{18} -Z (IV)

wherein Z is a reactive residue of an alcohol, and R^{18} is the same as R^1 except hydrogen atom.

20 The reactive residue of an alcohol represented by the group Z includes, for example, halogen atom (e.g. chlorine, bromine, iodine), lower alkylsulfonyloxy (e.g. methanesulfonyloxy), arylsulfonyloxy (e.g. benzenesulfonyloxy, p-toluenesulfonyloxy).

The reaction of the compound of the formula (I-8) and the compound of the formula (IV) is usually carried out in an appropriate solvent. Suitable examples of the solvent are aromatic hydrocarbons (e.g. benzene, xylene), ketones (e.g. methyl ethyl ketone), ethers (e.g. dioxane), N,N-dimethylformamide. The reaction is preferably carried out in the presence of a basic substance. The basic substance includes the same substances as used in the above Process A. The reaction temperature is usually in the range of 30 to 150°C. The compounds of the formula (I-8) can be prepared by the same processes as the above-mentioned Processes A to C.

30 When the compounds of the formula (I) prepared by the Processes A and D have a hydroxyl group in the structure thereof, they can be converted into corresponding ester derivatives or ether derivatives by reacting the compound with an appropriate acylating agent or a lower alkylating agent. These reactions are usually carried out by a conventional method.

35 The compounds of the formula (I) prepared by the above mentioned Processes can be isolated and purified from the reaction mixture by a conventional method.

40 The compounds of the formula (I) of the present invention are obtained in the form of a free base or a salt or a hydrate depending on the kinds of the starting compound, the kinds of reaction, the reaction conditions, and the like. When the compounds are obtained in the form of a salt, they can be converted into the corresponding free base by a conventional method, for example, by treating them with a basic substance such as an alkali metal hydroxide. Besides, when the compounds are obtained in the form of a free base, they can be converted into the corresponding salt by a conventional method, for example, by treating them with various acids.

45 The compounds of the formula (I) of the present invention show inhibitory effect on exploratory activity, antagonistic effect on apomorphine-induced vomiting, binding property to dopamine (D_2) and serotonin (5_2) receptors, increasing effect on concentration of brain monoamine metabolites, and little toxicity. Accordingly, the compounds of the present invention are useful as an anti-psychotic (neuroleptic) drug or an anti-anxiety (anxiolytic) drug.

50 Furthermore, the compounds of the formula (I) of the present invention show excellent improving effect in some animal models of memory impairment. Accordingly, the compounds of the present invention are also useful as a medicament for the treatment of various symptoms of cerebral insufficiency.

The compounds of the present invention which show potent antagonistic effect on apomorphine-induced vomiting, binding property to serotonin (5_2) and dopamine (D_2) receptors, and increasing effect on concentration of brain monoamine metabolites are, for example, the following compounds and a pharmaceutically acceptable salt thereof.

- 55
- (1) 2-(4-Ethyl-1-piperazinyl)-4-(4-fluorophenyl)-5,6,7,8,9,10-hexahydrocycloocta[b]pyridine
 - (2) 2-(4-Ethyl-1-piperazinyl)-4-(2,4-difluorophenyl)-5,6,7,8,9,10-hexahydrocycloocta[b]pyridine
 - (3) 2-(4-Methyl-1-piperazinyl)-4-(4-fluorophenyl)-5,6,7,8,9,10-hexahydrocycloocta[b]pyridine
 - (4) 2-(4-Methyl-1-piperazinyl)-4-(2,4-difluorophenyl)-5,6,7,8,9,10-hexahydrocycloocta[b]pyridine

- (5) 2-(4-n-Propyl-1-piperazinyl)-4-(4-fluorophenyl)-5,6,7,8,9,10-hexahydrocycloocta[b]pyridine
- (6) 2-(4-n-Butyl-1-piperazinyl)-4-(4-fluorophenyl)-5,6,7,8,9,10-hexahydrocycloocta[b]pyridine
- (7) 2-(4-n-Pentyl-1-piperazinyl)-4-(4-fluorophenyl)-5,6,7,8,9,10-hexahydrocycloocta[b]pyridine
- (8) 2-[4-(2-Hydroxyethyl)-1-piperazinyl]-4-(4-fluorophenyl)-5,6,7,8,9,10-hexahydrocycloocta[b]pyridine
- 5 (9) 2-(1-Piperazinyl)-4-(4-fluorophenyl)-6,7-dihydro-5H-1-pyridine
- (10) 2-(4-Ethyl-1-piperazinyl)-4-(4-fluorophenyl)-6,7-dihydro-5H-1-pyridine
- (11) 2-(4-Ethyl-1-piperazinyl)-4-phenyl-6,7,8,9-tetrahydro-5H-cyclohepta[b]pyridine
- (12) 2-(4-Ethyl-1-piperazinyl)-4-(4-fluorophenyl)-6,7,8,9-tetrahydro-5H-cyclohepta[b]pyridine
- (13) 2-(4-Ethyl-1-piperazinyl)-4-(4-fluorophenyl)-5,6,7,8-tetrahydro-5,8-methanoquinoline
- 10 (14) 2-(4-Ethyl-1-piperazinyl)-4-(2,4-difluorophenyl)-5,6,7,8-tetrahydro-5,8-methanoquinoline
- (15) 2-(4-Ethyl-1-piperazinyl)-4-(4-fluorophenyl)-6,7,8,9-tetrahydro-5H-6,9-methanocyclohepta[b]pyridine

The compounds of the present invention which have excellent improving effects on behavioral and/or memory deficits induced by scopolamine or cycloheximide are, for example, the following compounds and a pharmaceutically acceptable salt thereof.

- 15 (1) 2-(1-Piperazinyl)-4-phenyl-6,7-dihydro-5H-1-pyridine
- (2) 2-(1-Piperazinyl)-4-(4-fluorophenyl)-6,7-dihydro-5H-1-pyridine
- (3) 2-(1-Piperazinyl)-4-phenyl-5,6,7,8-tetrahydroquinoline
- (4) 2-(1-Piperazinyl)-4-(4-fluorophenyl)-5,6,7,8-tetrahydroquinoline
- (5) 2-(1-Piperazinyl)-4-(4-fluorophenyl)-6,7,8,9-tetrahydro-5H-cyclohepta[b]pyridine

20 The pharmacological activities of the representative compounds of the present invention are illustrated by the following Experiments.

In the Experiments, the following compounds, which are disclosed in the above mentioned U.S. Patent No. 4,469,696, were used as reference.

- Test Compound A (disclosed in Example 4):
- 25 4,5-Bis-(4-fluorophenyl)-6-methyl-2-(4-phenylpiperazino)pyridine
- Test Compound B (disclosed in Example 1):
- 4,5-Bis-(4-methoxyphenyl)-6-methyl-2-(4-phenylpiperazino)pyridine
- Test Compound C (disclosed in Example 2):
- 4,5-Bis-(4-hydroxyphenyl)-6-methyl-2-(4-phenylpiperazino)pyridine

30 Experiment 1

Inhibitory effect on exploratory activity:

35 A group of 5 male mice (Std-ddy strain, 20 - 25 g) was used. Two hours after oral administration of the test compound, mice were placed individually in a test box (23 x 35 x 30 cm) on Animex activity meter (Farad Co.). Immediately thereafter, activity counting was started and lasted for three minutes. The mean counts of the compound-treated group were compared with those of the corresponding control (non-dosed) group, and the percent inhibition was calculated. The results are shown in Table 1.

40

45

50

55

Table 1 Inhibitory effect on exploratory activity

Test compound	Inhibitory ratio (%)	Test compound	Inhibitory ratio (%)
Ex. 1b*	83.7	83	53.5
" 2a	77.5	106	74.9
" 31	62.5	107	53.2
" 36	68.4	111	74.0
" 38	58.9	112	73.2
" 53	68.1	114	81.9
" 57	77.2	122	81.3
" 71	61.9	123	72.2
" 75	78.9	125	57.9
" 76	87.6	127	80.3

*) The compound of Example 1b (hereinafter, the same)

Experiment 2

Antagonistic effect on apomorphine-induced vomiting:

A group of 3 - 4 dogs (Beagle, 8 - 15 kg) was used for examining effects on the test compounds on apomorphine-induced vomiting, a known test for evaluating neuroleptic drugs.

Each dog was given a subcutaneous injection of apomorphine hydrochloride (0.3 mg/kg) two hours after the treatment of the test compounds. Then, the frequency of oral vomiting was counted for one hour. The mean counts of the compound-treated group were compared with those of the corresponding control (non-dosed) group, and the percent inhibition was calculated. The results are shown in Table 2.

Table 2 Antagonistic effect on apomorphine-induced vomiting

Test comp.	Dose (mg/kg)	Inhibitory ratio (%)	Test comp.	Dose (mg/kg)	Inhibitory ratio (%)
Ex. 1a* ¹	0.2	100	Ex. 73	1.0	100
" 1b	0.3	88	" 76	0.3	71
" 2b	0.2	100	" 77	0.5	61
" 6	1.0	89	" 82	1.0	100
" 7	1.0	80	" 88	3.0	94
" 15	3.0	100	" 112	0.2	94
" 25	3.0	100	" 114	0.2	89
" 27	3.0	47	" 116	1.0	100
" 29	3.0	63	" 118	0.3	94
" 36	0.5	81	" 121	0.5	93
" 37	0.3	80	" 122	0.5	100
" 43	1.0	96	" 135	1.0	100
" 56	0.5	85	A* ²	3.0	11
" 66	1.0	95	B* ²	3.0	13
" 71	3.0	84	C* ²	3.0	16

*¹) The compound of Example 1a (hereinafter, the same)

*²) The compounds disclosed in the U.S. Patent No. 4,469,696

Experiment 3

In vitro binding property to dopamine (D₂), serotonin (S₁, S₂) and adrenaline (α₁) receptors:

Dopamine (D₂), serotonin (S₁, S₂) and adrenaline (α₁) receptor binding assays were carried out according to the methods of I. Creese et al. [Eur. J. Pharmacol., 46, 377 (1977), S.J. Peroutka et al. [Mol. Pharmacol. 16, 687 (1979)], J.E. Leysen et al. [Mol. Pharmacol., 21, 301 (1982)] and D.C. U'Prichard et al. [Mol. Pharmacol., 13, 454 (1977)], respectively.

Crude synaptosome fractions were prepared from some brain regions in rats for receptor sources. Radioactive ligands used were [³H] spiperone (D₂), [³H] serotonin (S₁), [³H] ketanserin (S₂) and [³H] WB-4101 (α₁). The binding assay was performed by incubating aliquots of synaptosome fraction in buffer solution (final volume: 1 ml) containing a [³H] labelled ligand and a test compound. The assay was terminated by rapid filtration through Whatman GF/B glass fiber filters attached to a cell-harvester (Brandel) and radioactivity on the filters was counted in a Packard Triscarb scintillation counter. Specific binding was calculated as a difference between amounts of radioactivity in the presence and absence of an unlabelled ligand [spiperone (D₂), serotonin (S₁), methysergide (S₂) and prazosin (α₁)]. The IC₅₀ value of the test compounds (concentration causing 50 % inhibition of [³H] ligand specific binding) was determined by probit analysis. The results are shown in Table 3.

Table 3 Binding property to dopamine (D_2),
serotonin (S_1 , S_2) and adrenaline (α_1)
receptor

5

10

15

20

25

30

35

40

Test compound	IC ₅₀ (nM)			
	D_2	S_1	S_2	α_1
Ex. 1a ^{*1}	24	-	9.9	83
" 1b	46	-	7.0	92
" 2a	24	-	28	88
" 6	19	1500	7.6	6.9
" 7	30	-	9.5	-
" 15	58	1700	19	35
" 23	96	2900	7.7	320
" 25	77	2600	2.3	120
" 71	25	-	14	13
" 73	18	-	2.4	4.6
" 86	730	1400	81	530
" 87	270	820	28	94
" 88	70	670	9.1	65
" 92	300	1900	27	380
" 96	660	4900	62	1100
" 97	260	4100	5.1	890
" 103	140	-	41	-
A ^{*2}	>10000	-	>10000	>10000
B ^{*2}	>10000	-	-	-
C ^{*2}	>10000	-	-	-

*1) The compound of Example 1a (hereinafter, the same)

*2) The compounds disclosed in the U.S. Patent No.
4,469,696

45

Experiment 4

50

Increasing effect on concentration of brain monoamine metabolites:

55

A group of 5 male mice (Std-ddy strain, 25 - 30 g) was used for examining effect of the test compounds on concentration of brain monoamine metabolites. It is generally accepted that an increase in each monoamine is mainly caused by each monoamine receptor blockade.

Mice were killed by decapitation 2 hours after treatment with the test compounds. Brains were quickly taken out, homogenized in 1N formic acid-acetone solution, and centrifuged in a refrigerated ultracentrifuge. The supernatant was evaporated by blowing with N_2 gas. Then, the residue was again resolved in 0.01 N

acetic acid, and served for determining concentration of dopamine metabolites, 3,4-dihydroxyphenylacetic acid (DOPAC) and homovanillic acid (HVA), a norepinephrine metabolite, 3-methyl-4-hydroxyphenylethylene glycol (MOPEG), and a serotonin metabolite, 5-hydroxyindole-3-acetic acid (5-HIAA) concentrations by HPLC with electrochemical detection. The effect of the test compounds on concentration of each monoamine metabolite is shown as % of control (level of non-dosed animals = 100) in Table 4.

Table 4 Increasing effect on the concentration of monoamine metabolites

10

Test compound	Dose (mg/kg)	DOPAC	(%) HVA	MOPEG	5-HIAA
EX. 1b*	1	235	212	118	102
" 2b	2	364	271	-	-
" 6	10	321	242	115	103
" 15	10	277	235	118	108
" 23	10	344	320	115	145
" 24	10	290	317	139	150
" 25	1	272	207	106	113
" 26	10	344	263	-	-
" 27	10	208	193	-	-
" 29	10	296	274	-	-
" 35	10	165	170	-	-
" 37	3	268	203	143	95
" 39	10	307	233	117	97
" 42	10	258	211	-	-
" 44	10	272	210	-	-
" 46	10	296	235	-	-
" 49	10	197	195	-	-
" 52	10	233	216	-	-
" 54	2	251	250	-	-
" 55	2	379	240	-	-
" 56	10	302	249	106	110
" 59	2	246	236	126	106
" 66	2	311	308	111	116
" 68	2	206	181	-	-
" 84	10	237	216	-	-
" 87	10	180	194	117	136
" 88	10	329	209	136	120

55

(to be continued)

Table 4 (Continue)

Test compound	Dose (mg/kg)	DOPAC	(%) HVA	MOPEG	5-HIAA
EX. 92	10	208	192	121	139
" 96	10	201	194	115	127
" 97	10	232	176	125	128
" 104	10	344	247	160	97
" 117	10	242	210	133	100
" 118	3	302	250	129	110
" 121	10	312	215	-	111
" 122	2	306	229	-	100
" 124	10	292	238	102	107
" 131	1	319	224	-	-
" 132	10	333	209	-	-
" 137	10	300	270	-	-
" 138	2	236	173	-	-
" 139	10	267	206	-	-

*) The compound of Example 1b (hereinafter, the same)

Experiment 5

Improving effect on scopolamine-induced deficit of spontaneous alternation behavior:

A group of 15 - 25 male mice (Std-ddY strain, 22 - 28 g) was used for evaluating effect of the test compounds on scopolamine-induced deficit of spontaneous alternation in a T-maze, which is a known animal model of memory impairment due to hypofunction of the cholinergic nervous system. The T-maze used consists of a stem and two arms which are 25 cm long, 5 cm wide and 10 cm high. The first 10 cm of the stem and last 10 cm of each arm are divided by sliding doors into start and goal boxes.

A test compound and scopolamine hydrobromide (1 mg/kg) were intraperitoneally administered to each mouse, and after 30 minutes a test of spontaneous alternation task in the T-maze was continuously repeated for 8 trials. Commonly, naive mice alternate each (right and left) goal box in turn, but scopolamine-treated animals tend to enter the same goal box repeatedly. The effect of the test compounds was expressed as % improvement (complete improvement to the alternation level of non-dosed mice = 100 %). The results are shown in Table 5.

Table 5 Improving effect on scopolamine-induced deficit of spontaneous alternation behavior

Test compound	Dose (mg/kg)	Improvement (%)
Ex. 86*	10.0	47.6
" 87	0.5	48.0
	2.0	60.0
" 88	10.0	37.0
" 97	10.0	32.0

*) The compound of Example 86 (hereinafter the same)

Experiment 6

Improving effect on cycloheximide-induced amnesia of passive avoidance response:

Anti-amnesic effect of the test compounds was examined using mice given cycloheximide, which is a known amnesia-inducing agent.

A group of 15 - 20 male mice (Std-ddY strain, 27 - 33 g) was subjected to training and retention trials for a passive avoidance task in a step-down apparatus (30 x 30 x 50 cm) with a grid floor and a wooden platform (4 x 4 x 4 cm) in a center of the floor. In the training trial, each mouse was first placed on the platform. When the mouse stepped down on the grid floor, an electric shock (1 Hz, 0.5 sec, 60 VDC) was delivered to the feet for 15 seconds. Immediately after the training trial, cycloheximide (60 mg/kg, s.c.) and a test compound (i.p.) were administered. The retention trial was carried out 24 hours thereafter, the time from placing again each mouse placed on the platform until stepping down on the grid floor (step-down latency) was measured. The step-down latency in the retention trial was markedly shortened by treatment of cycloheximide (amnesia). The effect of test compounds was assessed by % improvement (complete improvement to the latency level of non-dosed animals = 100 %). The results are shown in Table 6.

Table 6 Improving effect on cycloheximide-induced amnesia of passive avoidance response

Test compound	Dose (mg/kg)	Improvement (%)
Ex. 87*	0.5	48
	2.0	77
" 88	2.0	60
" 92	2.0	44
	10.0	78
" 96	0.5	81
	2.0	64
" 97	0.5	38
	2.0	36

*) The compound of Example 87 (hereinafter the same)

Experiment 7

Acute toxicity:

A group of 5 male mice (Std-ddy strain, 25 - 30 g) was used. The test compound was orally administered to the test animal in the form of a 0.5 % tragacanth solution or suspension, and for 7 days after the administration of the test compound, the lethality of animals was observed. The results are shown in Table 7.

Table 7 Acute toxicity

Test compound	Dose (mg/kg)	Number of dead animals
		Number of test animals
Ex. 1a*	500	0/5
" 1b	500	0/5
" 2b	500	0/5
" 6	500	0/5
" 25	500	0/5
" 76	500	0/5

*) The compound of Example 1a (hereinafter, the same)

The compounds of the present invention can be administered either in oral route, parenteral route or intrarectal route, but preferably in oral route. The dose of the compounds may vary depending on the kinds of the compounds, administration routes, severity of the disease and age of patients, but is usually in the

range of 0.01 to 50 mg/kg/day, preferably 0.01 to 5 mg/kg/day.

The compounds of the present invention are usually administered in the form of a conventional pharmaceutical preparation in admixture with a conventional pharmaceutically acceptable carrier or diluent. The pharmaceutically acceptable carrier or diluent includes the conventional pharmaceutically acceptable carriers or diluents which do not react with the compounds of the present invention. Suitable examples of the carrier or diluent are lactose, glucose, mannitol, sorbitol, dextrin, cyclodextrin, starch, sucrose, magnesium metasilicate aluminate, synthetic aluminum silicate, crystalline cellulose, sodium carboxymethyl cellulose, hydroxypropyl starch, calcium carboxymethyl cellulose, ion exchange resin, methyl cellulose, gelatin, acacia, pullulan, hydroxypropyl cellulose, low substituted hydroxypropyl cellulose, hydroxypropyl methyl cellulose, polyvinylpyrrolidone, polyvinyl alcohol, light silicic anhydride, magnesium stearate, talc, tragacanth, bentonite, veegum, carboxyvinyl polymer, titanium oxide, sorbitan fatty acid ester, sodium laurylsulfate, glycerin, glycerin fatty acid ester, anhydrous lanolin, glycerogelatin, polysorbate, macrogol, vegetable oil, wax, propylene glycol and water. The pharmaceutical preparation includes tablets, capsules, granules, fine granules, powders, syrups, suspensions, injections, suppositories. These preparations can be prepared by a conventional method. The liquid preparations may be in the form that they are dissolved or suspended in water or any other conventional medium when used. The tablets, granules and fine granules may be coated with a conventional coating agent. The injections are usually prepared by dissolving the compound of the present invention in water, but occasionally in a physiological saline solution or glucose solution, which is optionally incorporated with a buffer or a preservative. The pharmaceutical preparations may also contain other pharmaceutically active compounds.

The present invention is illustrated by the following Reference Examples, Examples and Preparations but should not be construed to be limited thereto. The compounds are identified by elementary analysis, mass spectrum, IR spectrum, UV spectrum, NMR spectrum, and the like.

In the Reference Examples and Examples, the following abbreviations may occasionally be used.

25	Me:	methyl
	Et:	ethyl
	t-Bu:	tertiary butyl
	Ph:	phenyl
	A:	ethanol
30	AC:	acetonitrile
	AT:	acetone
	CF:	chloroform
	D:	N,N-dimethylformamide
	E:	diethyl ether
35	EA:	ethyl acetate
	HX:	hexane
	IP:	isopropyl alcohol
	M:	methanol
	MC:	methylene chloride
40	PE:	petroleum ether
	T:	toluene
	W:	water

Besides, the solvent shown in brackets as to the melting point in the following Reference Examples and Examples means a solvent for recrystallization.

Reference Example 1


Preparation of 4-phenyl-5,6,7,8-tetrahydro-2(1H)-quinolinone:

A mixture of benzoylacetone (25 g), cyclohexanone (25 g) and 75 % polyphosphoric acid (250 g) is stirred at 50°C for 30 minutes and further at 110°C for 1.5 hour. After cooling, the reaction mixture is poured into ice-water and thereto is added diethyl ether (300 ml). The mixture is stirred and the precipitated crystals are collected by filtration. The crystals are recrystallized from N,N-dimethylformamide-ethanol to give the desired compound (27 g), m.p. 285 - 288°C.

Reference Examples 2 to 46

In the same manner as described in Reference Example 1 except that the corresponding starting materials are used, there are obtained the compounds as shown in the following Tables 8 and 9.

Table 8



Ref. Ex.	n	R ²	R ³	Melting point (°C)	Solvent for recrystallization
2	3	H	H	221 - 223	M
3	3	4-F	H	258 - 265	M
4	3	3-F	H	234 - 235	A
5	3	2-F	H	210 - 212	A-W
6	3	4-OMe	H	243 - 245	M
7	3	4-F	2-F	235 - 245	A
8	4	4-F	H	287 - 288	D-A
9	4	3-F	H	253 - 255	A
10	4	2-F	H	174 - 176	M
11	4	4-OMe	H	239 - 242	M
12	5	H	H	280 - 281	M
13	5	4-F	H	245 - 246	M
14	5	3-F	H	273 - 275	M
15	5	2-F	H	251 - 252	A-W
16	5	4-OMe	H	258 - 260	M
17	5	2-OMe	H	245 - 247	A
18	5	4-Me	H	253 - 258	M

(to be continued)

Table 8 (Continue)

5	Ref. Ex.	n	R ²	R ³	Melting point (°C)	Solvent for recrystallization
	19	5	4-Cl	H	251 - 255	D-W
	20	5	2-Cl	H	262 - 263	D-W
10	21	5	4-F	2-F	233 - 235	A
	22	6	H	H	265 - 266	M
	23	6	4-F	H	235 - 238	IP
15	24	6	3-F	H	260 - 270	A
	25	6	2-F	H	245 - 247	IP
	26	6	4-F	3-F	255 - 261	M
	27	6	4-F	2-F	240 - 243	M
20	28	6	6-F	2-F	254 - 256	A
	29	6	4-OMe	H	267 - 271	M
	30	6	4-Me	H	265 - 275	A
25	31	6	4-Cl	H	284 - 287	M
	32	6	3-Cl	H	248 - 250	A
	33	6	4-Br	H	286 - 292	M
30	34	7	4-F	H	241 - 243	EA

35


40

45

50

55







Table 9



Ref. Ex.	W Y	R ²	R ³	Melting point (°C)	Solvent for recrystallization
35	Me	H	H	275 - 279	M
36	Me	F	H	270 - 275	M
37	Me	F	H	255 - 257	M
38	Me	F	H	278 - 284	M
39	Me	F	H	266 - 267	M
40	Me	F	H	-	-

(to be continued)

Table 9 (Continue)

Ref. Ex.	W Y	R ²	R ³	Melting point (°C)	Solvent for recrystallization
41	t-Bu 	H	H	272 - 276	M
42	Ph 	F	H	273 - 277	M
43		H	H	220 - 225	IP
44		F	H	263 - 274	A
45		F	F	-	-
46		F	H	>300	M

Reference Example 47

Preparation of 4-(4-fluorophenyl)-1,5,6,7,8,9-hexahydro-2H-6,9-methanocyclohepta[b]pyridin-2-one:

Bicyclo[3.2.1]octan-2-one (2 g) and 4-fluorobenzoylacetone (2.6 g) are dissolved in 1,1,2,2-tetrachloroethane (5 ml) and thereto is added 75 % polyphosphoric acid (25 g). The mixture is stirred at 80 °C for 30 minutes, at 100 °C for 1 hour, and further at 130 °C for 30 minutes. After cooling, the reaction mixture is poured into ice-water and neutralized with potassium carbonate. The precipitated crystals are collected by filtration and washed successively with water and ethyl acetate. The resultant is recrystallized from methanol to give the desired product (2.7 g), m.p. >300 °C.

Reference Example 48

Preparation of 4-(4-fluorophenyl)-1,5,6,7,8,9-hexahydro-2H-5,8-methanocyclohepta[b]pyridin-2-one:

In the same manner as described in Reference Example 47 except that the corresponding starting materials are used, there is obtained the desired compound.

Reference Example 49

Preparation of 2-chloro-4-(4-fluorophenyl)-5,6,7,8-tetrahydroquinoline:


A mixture of 4-(4-fluorophenyl)-5,6,7,8-tetrahydro-2(1H)-quinolinone (18.7 g) and phenylphosphonic dichloride (29 ml) is stirred at 170 °C for 1 hour. After cooling, the reaction mixture is dissolved in

chloroform (200 ml) and the mixture is added dropwise into ice-water with stirring over a period of about 30 minutes. The mixture is made alkaline by dropwise addition of conc. aqueous ammonia thereto. The organic layer is separated, washed with water, dried over anhydrous sodium sulfate, and concentrated under reduced pressure. The residue is recrystallized from isopropanol-petroleum ether to give the desired product (15.1 g), m.p. 111 - 112°C.

Reference Examples 50 to 94

In the same manner as described in Reference Example 49 except that the corresponding starting materials are used, there are obtained the compounds as shown in Tables 10 and 11.

Table 10



Ref. Ex.	n	R ²	R ³	Melting point (°C)	Solvent for recrystallization
50	3	H	H	86 - 87	IP-PE
51	3	4-F	H	144 - 146	A
52	3	3-F	H	55 - 57	IP
53	3	2-F	H	Oil	-
54	3	4-OMe	H	Oil	-
55	3	4-F	2-F	Oil	-
56	4	H	H	86 - 88	A
57	4	3-F	H	84 - 86	A
58	4	2-F	H	84 - 85	IP-PE
59	4	4-OMe	H	104 - 106	A
60	5	H	H	71 - 72	IP
61	5	4-F	H	84 - 85	PE
62	5	3-F	H	Oil	-
63	5	2-F	H	89 - 90	A
64	5	4-OMe	H	75 - 76	A
65	5	4-Me	H	66 - 67	A
66	5	4-Cl	H	117 - 118	MC-HX

(to be continued)

Table 10 (Continue)

5	Ref. Ex.	n	R ²	R ³	Melting point (°C)	Solvent for recrystallization
	67	5	4-F	2-F	55 - 56	IP
	68	6	H	H	96 - 97	IP-PE
10	69	6	4-F	H	136 - 137	A
	70	6	3-F	H	110 - 111	A
	71	6	2-F	H	81 - 82	IP
15	72	6	4-F	3-F	137 - 138	A
	73	6	4-F	2-F	74 - 75	IP
	74	6	6-F	2-F	83 - 84	IP
	75	6	4-OMe	H	147 - 150	MC-HX
20	76	6	4-Me	H	142 - 143	M
	77	6	4-Cl	H	183 - 185	CF-T
	78	6	3-Cl	H	100 - 101	A
25	79	6	4-Br	H	Oil	-
	80	7	4-F	H	60 - 68	E

30

35


40

45

50

55

Table 11



Ref. Ex.	W Y	R ²	R ³	Melting point (°C)	Solvent for recrystallization
81	Me 	H	H	94 - 95	IP
82	Me 	F	H	94 - 95	A
83	 Me	F	H	74 - 75	IP
84	Me Me 	F	H	102 - 104	MC-HX
85	t-Bu 	H	H	109 - 110	A
86	Ph 	F	H	159 - 162	CF-A
87	Me Me	F	H	83 - 84	IP

(to be continued)

Table 11 (Continue)

Ref. Ex.	W Y	R ²	R ³	Melting point (°C)	Solvent for recrystallization
88		F	H	-	-
89		H	H	63 - 65	IP-PE
90		F	H	74 - 76	IP
91		F	F	Oil	-
92		F	H	132 - 135	MC-HX
93		H	H	Oil	-
94		F	H	92 - 94	MC-HX

Example 1

Preparation of 2-(4-ethyl-1-piperazinyl)-4-(4-fluorophenyl)-5,6,7,8,9,10-hexahydrocycloocta[b]pyridine:

A mixture of 2-chloro-4-(4-fluorophenyl)-5,6,7,8,9,10-hexahydrocycloocta[b]pyridine (2.0 g), N-ethyl-piperazine (2.4 g), and potassium iodide (1.1 g) is stirred at 170°C for 5 hours. After cooling, the reaction mixture is dissolved in ethyl acetate and water. The organic layer is washed with water and extracted with 5 % hydrochloric acid. The extract is made alkaline with potassium carbonate, and extracted with ethyl acetate. The extract is washed with water, dried over anhydrous sodium sulfate, and concentrated under reduced pressure.

(a) The residue is recrystallized from acetonitrile to give the desired product (1.2 g), m.p. 123 - 124°C.

This product obtained in the above (a) is converted to the following salt thereof by treating the product with various acids.

(b) Dimaleate, m.p. 165 - 167°C (ethanol)

(c) Dihydrochloride·½ hydrate, m.p. 215 - 222°C (acetone)

(d) Fumarate, m.p. 228 - 230°C (ethanol)

(e) Citrate, m.p. 184 - 187°C (ethanol)

Example 2

Preparation of 2-(4-ethyl-1-piperazinyl)-4-(2,4-difluorophenyl)-5,6,7,8,9,10-hexahydrocycloocta[b]pyridine:

- 5 A mixture of 2-chloro-4-(2,4-difluorophenyl)-5,6,7,8,9,10-hexahydrocycloocta[b]pyridine (10 g), N-ethylpiperazine (11 g), and potassium iodide (5.4 g) is stirred at 170 °C for 5 hours. After cooling, the reaction mixture is dissolved in chloroform and 5 % aqueous potassium carbonate solution. The organic layer is washed with water, dried over anhydrous sodium sulfate, and concentrated under reduced pressure.
- 10 (a) The residue is dissolved in toluene, and subjected to silica gel column chromatography. The fractions eluted with toluene and a mixture of toluene-ethyl acetate (1:1) are collected, and recrystallized from isopropyl alcohol to give the desired product (5.5 g), m.p. 124 - 125 °C.
- (b) The product obtained in the above (a) is treated with a solution of maleic acid in ethanol to give the dimaleate of the desired product, m.p. 133 - 135 °C (ethanol).

15 Example 3

Preparation of 2-(4-ethyl-1-piperazinyl)-4-phenyl-5,6,7,8-tetrahydroquinoline:

- 20 A mixture of 2-chloro-4-phenyl-5,6,7,8-tetrahydroquinoline (1.0 g), N-ethylpiperazine (1.2 g), and potassium iodide (0.66 g) is stirred at 170 °C for 15 hours. After cooling, the reaction mixture is dissolved in chloroform, and 5 % aqueous potassium carbonate solution. The organic layer is washed with water, dried over anhydrous sodium sulfate, and concentrated under reduced pressure. The resulting oily residue is subjected to basic alumina column chromatography. The fractions eluted with toluene and a mixture of toluene and ethyl acetate (9:1) are collected, and treated with a solution of maleic acid in ethanol. The
- 25 resulting maleate product is recrystallized from ethanol-ethyl acetate to give the dimaleate of the desired product (0.45 g), m.p. 139 - 142 °C.

Examples 4 - 77

- 30 In the same manner as described in Example 3 except that the corresponding starting materials are used, there are obtained the compounds as shown in Tables 12 and 13.

35


40

45

50

55

Table 12




15	Ex. n	R ²	R ³	R ¹	Q	M.p. (°C)	Sol. recry.	
	4	3	H	H	Et	2/3 Maleate	132 - 136	A
	5	3	H	H	CH ₂ CH ₂ OH	Maleate	178 - 183	A
20	6	3	4-F	H	Et	Dimaleate	164 - 166	A
	7	3	4-F	H	CH ₂ CH ₂ OH	½ Fumarate	207 - 210	M
	8	3	3-F	H	Et	Dimaleate	124 - 126	A
25	9	3	3-F	H	CH ₂ CH ₂ OH	Maleate·¼ H ₂ O	182 - 185	A
	10	3	2-F	H	Et	Maleate·1/5 H ₂ O	189 - 191	A
	11	3	2-F	H	CH ₂ CH ₂ OH	Maleate·½ H ₂ O	186 - 188	A
	12	3	4-OMe	H	Et	Dimaleate	174 - 176	A
30	13	3	4-OMe	H	CH ₂ CH ₂ OH	Dioxalate	183 - 187	M
	14	4	H	H	CH ₂ CH ₂ OH	Dimaleate	129 - 131	A
	15	4	4-F	H	Et	Dimaleate	146 - 148	A
	16	4	4-F	H	CH ₂ CH ₂ OH	Dimaleate	149 - 151	A
35	17	4	3-F	H	Et	Dimaleate	133 - 136	M
	18	4	3-F	H	CH ₂ CH ₂ OH	Dioxalate	180 - 182	M
	19	4	2-F	H	Et	Dimaleate	148 - 150	A
40	20	4	2-F	H	CH ₂ CH ₂ OH	Dioxalate·¼ H ₂ O	169 - 172	A
	21	4	4-OMe	H	Et	Dimaleate	179 - 181	A
	22	4	4-OMe	H	CH ₂ CH ₂ OH	Dioxalate·½ H ₂ O	160 - 165	A
45	23	5	H	H	Et	Dimaleate	150 - 152	IP
	24	5	H	H	CH ₂ CH ₂ OH	½ Fumarate	142 - 144	A
	25	5	4-F	H	Et	Dimaleate	150 - 152	A
	26	5	4-F	H	CH ₂ CH ₂ OH	1.7 Oxalate	180 - 182	A
50	27	5	3-F	H	Et	Dimaleate	138 - 140	A

(to be continued)

Table 12 (Continue)

5	Ex. n	R ²	R ³	R ¹	Q	M.p. (°C)	Sol. recry.
	28	5	3-F	H	CH ₂ CH ₂ OH	Maleate·½ H ₂ O	155 - 158 IP
	29	5	2-F	H	Et	Oxalate	231 - 234 A-M
10	30	5	2-F	H	CH ₂ CH ₂ OH	½ Fumarate·½ H ₂ O	163 - 164 M
	31	5	4-Cl	H	Et	-	150 - 151 MC-A
	32	5	4-OMe	H	Et	Dimalate	162 - 164 A
15	33	5	4-OMe	H	CH ₂ CH ₂ OH	Fumarate·½ H ₂ O	190 - 195 A-M
	34	5	4-Me	H	Et	Dimalate	180 - 183 A
	35	5	4-Me	H	CH ₂ CH ₂ OH	Dimalate·½ H ₂ O	151 - 153 IP
	36	6	H	H	Et	Fumarate	236 - 238 M
20	37	6	4-F	H	CH ₂ CH ₂ OH	Dimalate	135 - 137 A
	38	6	4-F	H		-	167 - 168 A-AT
	39	6	4-F	H		-	165 - 167 M
25	40	6	4-F	H		-	156 - 157 A-AT
	41	6	4-F	H		-	206 - 208 M-AT
30	42	6	3-F	H	Et	Dimalate	153 - 155 A
	43	6	2-F	H	Et	Fumarate	201 - 204 A
	44	6	4-Cl	H	Et	-	133 - 135 CF-IP
35	45	6	4-Cl	H	CH ₂ CH ₂ OH	-	156 - 158 CF-A
	46	6	3-Cl	H	Et	-	114 - 115 IP
	47	6	3-Cl	H	CH ₂ CH ₂ OH	Oxalate	210 - 212 M
40	48	6	4-Br	H	Et	-	134 - 136 A
	49	6	4-OMe	H	Et	-	113 - 114 MC-HX
	50	6	4-OMe	H	CH ₂ CH ₂ OH	HCl·3/4 H ₂ O	153 - 156 A-E
	51	6	4-Me	H	Et	Dimalate	163 - 164 A
45	52	6	4-Me	H	CH ₂ CH ₂ OH	Oxalate	207 - 209 A
	53	6	4-F	2-F	CH ₂ CH ₂ OH	Dioxalate·1/10 H ₂ O	169 - 172 A
	54	6	4-F	3-F	Et	-	105 - 106 AC
	55	6	6-F	2-F	Et	-	104 - 105 AC
50	56	7	4-F	H	Et	Fumarate	184 - 189 A-E

Table 13



Ex.	W Y	R ²	R ¹	Q	M.p. (°C)	Sol. recry.
57	Me	H	Et	Dimaleate	138 - 140	A
58	Me	H	CH ₂ CH ₂ OH	Dimaleate	140 - 141	A
59	Me	F	Et	Dioxalate · ½ H ₂ O	126 - 127	A
60	Me	F	CH ₂ CH ₂ OH	Fumarate	160 - 161	A
61	t-Bu	H	Et	-	130 - 131	A
62	t-Bu	H	CH ₂ CH ₂ OH	-	162 - 163	A
63	Ph	F	Et	Fumarate	212 - 215	M-A
64	Me	F	Et	Dimaleate	141 - 142	IP




(to be continued)

Table 13 (Continue)

Ex.	W Y	R ²	R ¹	Q	M.p. (°C)	Sol. recry.
65		F	CH ₂ CH ₂ OH	Dimaleate · ½ H ₂ O	120 - 122	A
66		F	Et	Fumarate	209 - 211	M-A
67		F	CH ₂ CH ₂ OH	HCl	200 - 203	A-E
68		F	Et	-	107 - 108	IP
69		F	CH ₂ CH ₂ OH	Dioxalate · ½ H ₂ O	125 - 126	A
70		F	Et	Fumarate	203 - 204	A
71		H	Et	Dimaleate · ½ H ₂ O	173 - 176	A
72		H	CH ₂ CH ₂ OH	Dimaleate	140 - 142	A
73		F	Et	Dimaleate	163 - 169	IP

(to be continued)

Table 13 (Continue)

Ex.	W Y	R ²	R ¹	Q	M.p. (°C)	Sol. recry.
74		F	CH ₂ CH ₂ OH	Dimaleate	159 - 161	A
75		F	Et	Fumarate	234 - 236	M-A
76		F	Et	Fumarate	216 - 219	M-A
77		F	Et	Fumarate · ½ H ₂ O	226 - 230	M-A

Examples 78 - 85

In the same manner as described in Example 3 except that the corresponding starting materials are used, there are obtained the following compounds.

(Example 78)

2-(4-Methyl-1-homopiperaziny)-4-(4-fluorophenyl)-6,7,8,9-tetrahydro-5H-cyclohepta[b]pyridine maleate, m.p. 187 - 189 °C (ethanol)

(Example 79)

2-(4-Butanoyl-1-homopiperaziny)-4-(4-fluorophenyl)-6,7,8,9-tetrahydro-5H-cyclohepta[b]pyridine, oily product

(Example 80)

2-(4-Methyl-1-homopiperaziny)-4-(4-fluorophenyl)-5,6,7,8,9,10-hexahydrocycloocta[b]pyridine dimaleate, m.p. 141 - 143 °C (ethanol)

(Example 81)

2-(4-Benzoyl-1-homopiperaziny)-4-(4-fluorophenyl)-5,6,7,8,9,10-hexahydrocycloocta[b]pyridine, oily product

(Example 82)

2-[4-(2-Methoxyethyl)-1-piperaziny]-4-(4-fluorophenyl)-5,6,7,8,9,10-hexahydrocycloocta[b]pyridine dimaleate, m.p. 119 - 120 °C (acetonitrile)

(Example 83)

2-(3-Methyl-1-piperaziny)-4-(4-fluorophenyl)-5,6,7,8,9,10-hexahydrocycloocta[b]pyridine, m.p. 138 - 141 °C (isopropyl alcohol-hexane)

(Example 84)

2-(3,5-Dimethyl-1-piperaziny)-4-(4-fluorophenyl)-5,6,7,8,9,10-hexahydrocycloocta[b]pyridine $\frac{1}{2}$ fumarate, m.p. 262 - 266 °C (decomposed) (methanol)

(Example 85)

2-(1-Homopiperaziny)-4-(4-fluorophenyl)-5,6,7,8,9,10-hexahydrocycloocta[b]pyridine maleate, m.p. 198 - 200 °C (ethanol)

Example 86


Preparation of 2-(1-piperaziny)-4-phenyl-5,6,7,8-tetrahydroquinoline:

A mixture of 2-(4-benzyl-1-piperaziny)-4-phenyl-5,6,7,8-tetrahydroquinoline (m.p. 105 - 107 °C, 2.1 g) obtained in the same manner as in Example 3, chloroethyl chloroformate (0.86 g) and methylene chloride (40 ml) is refluxed for 1 hour. The reaction mixture is concentrated under reduced pressure, and methanol (40 ml) is added to the residue. The mixture is refluxed for 30 minutes and concentrated under reduced pressure. The resulting residue is dissolved in water, washed with diethyl ether, neutralized with potassium carbonate, and extracted with chloroform. The extract is washed with water, dried over anhydrous sodium sulfate, and the solvent is distilled off under reduced pressure. To the residue is added a solution of maleic acid in ethanol. The resulting maleate product is recrystallized from ethanol to give the dimaleate of the desired product (1.2 g), m.p. 150 - 151 °C.

Examples 87 - 110






In the same manner as described in Example 86 except that the corresponding starting materials are used, there are obtained the compounds as shown in Tables 14 and 15.

Table 14



Ex.	n	R^2	R^3	Q	Melting point (°C)	Solvent for re-crystalln.
87	3	H	H	Maleate	175 - 179	A
88	3	4-F	H	Dimaleate	152 - 155	A
89	3	3-F	H	Maleate	176 - 179	A
90	3	2-F	H	Maleate	186 - 188	A
91	3	4-OMe	H	Dimaleate	148 - 149	A
92	4	4-F	H	Dimaleate · $\frac{1}{2}$ H ₂ O	143 - 145	A
93	4	3-F	H	Dimaleate · $\frac{1}{2}$ H ₂ O	148 - 151	A
94	4	2-F	H	Dimaleate	136 - 137	A
95	4	4-OMe	H	Dioxalate · $\frac{1}{2}$ H ₂ O	210 - 213	M
96	5	H	H	Maleate	173 - 176	A
97	5	4-F	H	Dimaleate	144 - 147	A
98	5	3-F	H	Maleate	173 - 175	A
99	5	2-F	H	Maleate	173 - 180	A
100	5	4-Cl	H	Maleate · $\frac{1}{2}$ H ₂ O	176 - 178	M-A
101	5	4-OMe	H	Dimaleate	133 - 137	IP
102	5	4-Me	H	Dimaleate · $\frac{3}{4}$ H ₂ O	95 - 97	IP
103	6	H	H	Dimaleate · $\frac{1}{2}$ H ₂ O	124 - 127	A
104	6	4-F	H	Maleate	186 - 190	A
105	6	4-Me	H	Fumarate	218 - 221	A
106	6	4-F	2-F	$\frac{1}{2}$ Fumarate · $\frac{1}{2}$ H ₂ O	177 - 179	A

Table 15

					
Ex.	W Y	R ²	Q	Melting point (°C)	Solvent for re- crystalln.
107		H	Dimaleate · 1/2 H ₂ O	147 - 149	A
108	Me 	H	3/2 Oxalate · 1/2 H ₂ O	178 - 181	M
109	Me 	F	1/2 Fumarate · 1/10 H ₂ O	181 - 186	A
110	t-Bu 	H	Maleate	207 - 208	A

Example 111

Preparation of 2-(4-methyl-1-piperazinyl)-4-(4-fluorophenyl)-6,7-dihydro-5H-1-pyridine:

A mixture of 2-(1-piperazinyl)-4-(4-fluorophenyl)-6,7-dihydro-5H-1-pyridine (2 g), 37 % aqueous formaldehyde solution (0.66 g), formic acid (0.68 g) and water (15 ml) is refluxed for 20 minutes. After cooling, the reaction mixture is made alkaline with diluted aqueous sodium hydroxide solution, and extracted with ethyl acetate. The extract is washed with water, dried over anhydrous sodium sulfate, and concentrated under reduced pressure. To the residue is added a solution of maleic acid in ethanol, and the resulting maleate product is recrystallized from ethanol to give the maleate of the desired compound, m.p. 135 - 137 °C.

Examples 112 - 117

In the same manner as described in Example 111 except that the corresponding starting materials are used, there are obtained the following compounds.

(Example 112)

2-(4-Methyl-1-piperazinyl)-4-(4-fluorophenyl)-5,6,7,8,9,10-hexahydrocycloocta[b]pyridine dimaleate, m.p. 136 - 138 °C (ethanol)

(Example 113)

2-(4-Methyl-1-piperazinyl)-4-(4-methylphenyl)-5,6,7,8,9,10-hexahydrocycloocta[b]pyridine dimaleate, m.p. 152 - 154 °C (ethanol)

(Example 114)

2-(4-Methyl-1-piperazinyl)-4-(2,4-difluorophenyl)-5,6,7,8,9,10-hexahydrocycloocta[b]pyridine, m.p. 132 - 133 °C (ethanol)

(Example 115)

2-(4-Methyl-1-piperazinyl)-4-(4-fluorophenyl)-6-methyl-5,6,7,8-tetrahydroquinoline dimaleate, m.p. 161 - 164 °C (ethanol)

(Example 116)

2-(3,4-Dimethyl-1-piperazinyl)-4-(4-fluorophenyl)-5,6,7,8,9,10-hexahydrocycloocta[b]pyridine fumarate ½ hydrate, m.p. 173 - 175 °C (ethanol-diethyl ether)

(Example 117)

2-(cis-3,5-Dimethyl-4-methyl-1-piperazinyl)-4-(4-fluorophenyl)-5,6,7,8,9,10-hexahydrocycloocta[b]pyridine fumarate, m.p. 208 - 210 °C (methanol-ethanol)

Example 118


Preparation of 2-(4-n-propyl-1-piperazinyl)-4-(4-fluorophenyl)-5,6,7,8,9,10-hexahydrocycloocta[b]pyridine:

A mixture of 2-(1-piperazinyl)-4-(4-fluorophenyl)-5,6,7,8,9,10-hexahydrocycloocta[b]pyridine (1.4 g), n-propyl bromide (0.56 g), potassium carbonate (0.68 g), potassium iodide (0.1 g) and methanol (50 ml) is refluxed for 15 hours. The reaction mixture is concentrated under reduced pressure and thereto is added water. The mixture is extracted with ethyl acetate, washed with water, dried over anhydrous sodium sulfate, and the solvent is distilled off under reduced pressure. To the residue is added a solution of maleic acid in ethanol, and the resulting maleate product is recrystallized from ethanol to give the dimaleate of the desired compound (0.6 g), m.p. 149 - 152 °C.

Examples 119 - 126

In the same manner as described in Example 118 except that the corresponding starting materials are used, there are obtained the compounds as shown in the following Table 16.

Table 16



Ex.	n	R¹	Q	M.p. (°C)	Solvent. for re- recrystal.
119	3	-(CH₂)₃OH	Maleate	174 - 178	A
120	3	$\begin{array}{c} \text{Me} \\ \diagup \\ -\text{CH} \\ \diagdown \\ \text{Me} \end{array}$	Maleate	215 - 218	A
121	6	$\begin{array}{c} \text{Me} \\ \diagup \\ -\text{CH} \\ \diagdown \\ \text{Me} \end{array}$	Dimalate	154 - 155	A
122	6	-(CH₂)₃CH₃	-	89 - 90	A
123	6	-(CH₂)₄CH₃	Maleate	213 - 217	A-M
124	6	-(CH₂)₅CH₃	Maleate	196 - 197	A
125	6	-CH₂CH=CH₂	$\frac{1}{2}$ H₂O	109 - 110	AC
126	6	-CH₂C≡CH	-	75 - 77	HX

Example 127

Preparation of 2-(4-n-propyl-1-piperazinyl)-4-(2,4-difluorophenyl)-5,6,7,8,9,10-hexahydrocycloocta[b]pyridine:

In the same manner as described in Example 118 except that the corresponding starting materials are used, the desired product is obtained, m.p. 108 - 109 °C (ethanol).

Example 128

Preparation of 2-[4-(2-furoyl)-1-piperazinyl]-4-phenyl-5,6,7,8-tetrahydroquinoline:

A mixture of 2-(1-piperazinyl)-4-phenyl-5,6,7,8-tetrahydroquinoline (1.2 g), 2-furancarboxylic acid (0.46 g), chloroform (40 ml) and 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (0.79 g) is stirred at 25 °C for 2 hours. The reaction mixture is washed with water, dried over anhydrous sodium sulfate, and the solvent is distilled off under reduced pressure. The residue is dissolved in toluene and subjected to silica gel column chromatography. The fractions eluted with toluene and toluene-ethyl acetate (9:1) are collected and recrystallized from ethanol to give the desired product (0.4 g), m.p. 128 - 130 °C.

Example 129

Preparation of 2-[4-(2-furoyl)-1-piperazinyl]-4-(4-fluorophenyl)-6,7-dihydro-5H-1-pyridine:

- 5 In the same manner as described in Example 128 except that the corresponding starting materials are used, the desired product is obtained, m.p. 165 - 166 °C (methanol).

Example 130

- 10 Preparation of 2-(4-ethyl-1-piperazinyl)-4-(4-hydroxyphenyl)-5,6,7,8,9,10-hexahydrocycloocta[b]pyridine:

- 2-(4-Ethyl-1-piperazinyl)-4-(4-methoxyphenyl)-5,6,7,8,9,10-hexahydrocycloocta[b]pyridine (1.3 g) is dissolved in 48 % hydrobromic acid (10 ml) and the mixture is stirred at 120 °C for 2 hours. After cooling, water is added thereto and the mixture is neutralized with 1N aqueous sodium hydroxide solution and
15 sodium hydrogen carbonate. The resulting precipitate is collected by filtration, washed with water, and recrystallized from methanol to give the desired product (0.6 g), m.p. 250 - 253 °C.

Example 131

- 20 Preparation of 2-(4-ethyl-1-piperazinyl)-4-(2,4-difluorophenyl)-6,7,8,9-tetrahydro-5H-cyclohepta[b]pyridine:

In the same manner as described in Example 3 except that the corresponding starting materials are used, there is obtained the oxalate • $\frac{1}{2}$ hydrate of the desired product, m.p. 225 - 227 °C (methanol).

- 25 Example 132

Preparation of 2-(4-ethyl-1-piperazinyl)-4-(2,4-difluorophenyl)-6,7-dihydro-5H-1-pyridine:

- In the same manner as described in Example 3 except that the corresponding starting materials are
30 used, there is obtained the maleate of the desired product, m.p. 195 - 196 °C (ethanol).

Example 133

Preparation of 2-(4-cyclopropyl-1-piperazinyl)-4-(4-fluorophenyl)-5,6,7,8,9,10-hexahydrocycloocta[b]pyridine:

- 35 In the same manner as described in Example 3 except that the corresponding starting materials are used, there is obtained the desired product, m.p. 125 - 128 °C (ethanol).

Example 134

- 40 Preparation of 2-[4-(4-fluorophenyl)-1-piperazinyl]-4-(4-fluorophenyl)-5,6,7,8,9,10-hexahydrocycloocta[b]pyridine:

- In the same manner as described in Example 3 except that the corresponding starting materials are
45 used, there is obtained the desired product, m.p. 130 - 131 °C (methanol).

Example 135

Preparation of 2-(4-ethyl-1-piperazinyl)-4-(2,4-difluorophenyl)-5,6,7,8-tetrahydro-5,8-methanoquinoline:

- 50 In the same manner as described in Example 3 except that the corresponding starting materials are used, there is obtained the oxalate • $\frac{1}{2}$ hydrate of the desired product, m.p. 234 - 238 °C (ethanol).

55

Example 136

Preparation of 2-(cis-3,5-dimethyl-4-methyl-1-piperazinyl)-4-(4-fluorophenyl)-6,7-dihydro-5H-1-pyridine:

- 5 In the same manner as described in Example 111 except that the corresponding starting materials are used, there is obtained the fumarate $\cdot \frac{1}{2}$ hydrate of the desired product, m.p. 204 - 208 °C (methanol-ethanol).

Example 137

10

Preparation of 2-[4-[3-(4-fluorobenzoyl)propyl]-1-piperazinyl]-4-(4-fluorophenyl)-6,7-dihydro-5H-1-pyridine:

In the same manner as described in Example 118 except that the corresponding starting materials are used, there is obtained the desired product, m.p. 115 - 116 °C (ethanol).

15

Example 138

Preparation of 2-[4-(3-hydroxypropyl)-1-piperazinyl]-4-(4-fluorophenyl)-5,6,7,8,9,10-hexahydrocycloocta[b]-pyridine:

20

In the same manner as described in Example 118 except that the corresponding starting materials are used, there is obtained the dioxalate $\cdot \frac{1}{2}$ hydrate of the desired product, m.p. 136 - 138 °C (methanol).

Example 139

25

Preparation of 2-[4-(2-acetoxyethyl)-1-piperazinyl]-4-(4-fluorophenyl)-5,6,7,8,9,10-hexahydrocycloocta[b]-pyridine:

30

A mixture of 2-[4-(2-hydroxyethyl)-1-piperazinyl]-4-(4-fluorophenyl)-5,6,7,8,9,10-hexahydrocycloocta[b]-pyridine (1.5 g), acetic anhydride (0.6 g), triethylamine (0.7 g) and ethyl acetate (30 ml) is refluxed for 2 hours. After cooling, the reaction mixture is washed with water, dried over anhydrous sodium sulfate, and concentrated under reduced pressure. The residue is dissolved in toluene and the mixture is subjected to silica gel column chromatography. The fractions eluted with toluene and toluene-ethyl acetate (1:1) are collected and thereto is added a solution of maleic acid in ethanol. The resulting maleate product is recrystallized from ethanol to give the maleate of the desired product (0.65 g), m.p. 187 - 191 °C.

35

Example 140

40

Preparation of 2-(4-cyclopropylmethyl-1-piperazinyl)-4-(4-fluorophenyl)-5,6,7,8,9,10-hexahydrocycloocta[b]-pyridine:

In the same manner as described in Example 118 except that the corresponding starting materials are used, there is obtained the desired product, m.p. 109 - 110 °C (methylene chloride-hexane).

- 45 The preparation of the pharmaceutical composition of the present invention is illustrated by the following Preparations.

50

55

Preparation 1

Preparation of capsules:

5

Components	Amount
2-(4-Ethyl-1-piperaziny)-4-(4-fluorophenyl)-6,7,8,9-tetrahydro-5H-cyclohepta[b]pyridine dimaleate	5 g
Corn starch	57 g
Lactose	10 g
Crystalline cellulose	25 g
Hydroxypropyl cellulose	2 g
Light silicic anhydride	0.5 g
Magnesium stearate	0.5 g

15

According to a conventional method, the above components are mixed and granulated, and the granules thus obtained are packed in capsules (1000 capsules) to give capsules containing the granules of 100 mg per one capsule.

20

Preparation 2

Preparation of tablets:

25

Components	Amount
2-(4-Ethyl-1-piperaziny)-4-(4-fluorophenyl)-5,6,7,8,9,10-hexahydrocycloocta[b]pyridine dimaleate	5 g
Corn starch	20 g
Lactose	30 g
Crystalline cellulose	30 g
Hydroxypropyl cellulose	5 g
Low substituted hydroxypropyl cellulose	10 g

35

According to a conventional method, the above components are mixed and granulated, and the granules thus obtained are mixed with light silicic anhydride and magnesium stearate, and the mixture is tabletted to give tablets containing the active ingredient of 5 mg per one tablet.

Preparation 3

40

Preparation of powders:

Components	Amount
2-(4-Ethyl-1-piperaziny)-4-(4-fluorophenyl)-5,6,7,8,9,10-hexahydrocycloocta[b]pyridine dimaleate	5 g
Corn starch	173 g
Lactose	300 g
Hydroxypropyl cellulose	20 g

50

According to a conventional method, the above components are mixed, granulated and screened, and the granules thus obtained are mixed with an appropriate amount of light silicic anhydride to give powders (100 triturations).

55

Preparation 4

Preparation of injections:

Components	Amount
2-(4-Ethyl-1-piperazinyl)-4-(4-fluorophenyl)-5,6,7,8,9,10-h-exahydrocycloocta[b]pyridine dimaleate	5 g
D-Sorbitol	45 g
1N Aqueous solution of maleic acid or sodium hydroxide	q.s.
Distilled water for injection	q.s.
	Totally 1000 ml

The above active ingredient and D-sorbitol are mixed with distilled water for injection, and thereto is added 1N aqueous solution of maleic acid or sodium hydroxide to adjust the solution to pH 4.0. The solution is filtered with a membrane filter (pore size, 0.22 μ m) and packed in ampoule (content 10 ml). The ampoule is sealed by melting and sterilized with high pressure steam at 121 °C for 20 minutes to give injection solutions.

Preparation 5

Preparation of lyophilized preparation:

Components	Amount
2-(4-Ethyl-1-piperazinyl)-4-(4-fluorophenyl)-5,6,7,8,9,10-h-exahydrocycloocta[b]pyridine dimaleate	5 g
D-Mannitol	45 g
1N Aqueous solution of maleic acid of sodium hydroxide	q.s.
Distilled water for injection	q.s.
	Totally 1000 ml


The above active ingredient and D-mannitol are mixed with distilled water for injection, and thereto is added 1N aqueous solution of maleic acid or sodium hydroxide to adjust the solution to pH 4.0. The solution is filtered with a membrane filter (pore size, 0.22 μ m) and packed in a vial (content 10 ml). The vial is sealed with a rubber stopper in halfway and subjected to lyophilization, that is, pre-freezing, primary drying at -50 °C, secondary drying at -20 °C, and then final drying at 20 °C. After completely sealed with a rubber stopper within a chamber, the vial is covered with a flip-off cap to give lyophilized preparation.



Claims

Claims for the following Contracting States : AT, BE, CH, DE, DK, FR, GB, IT, LI, LU, NL, SE

1. A compound of the formula (I):



wherein n is 3, 4, 5, 6 or 7,

R¹ is hydrogen atom, C₁-C₁₀ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₃-C₆ cycloalkyl-(C₁-C₄) alkyl, hydroxy-(C₂-C₆) alkyl, C₁-C₃ alkoxy-(C₂-C₆) alkyl, acyloxy-(C₂-C₆) alkyl, unsubstituted or substituted aroyl-(C₁-C₆) alkyl, unsubstituted or substituted aryl, heteroaryl, or acyl,

R² and R³ are the same or different and are each hydrogen atom, halogen atom, C₁-C₆ alkyl, C₁-C₆ alkoxy, trifluoromethyl, or hydroxy,

R⁴, R⁵ and R⁶ are the same or different and are each hydrogen atom, C₁-C₆ alkyl, or phenyl, or two of R⁴, R⁵ and R⁶ combine to form a single bond or C₁-C₃ alkylene,

R⁷ and R⁸ are the same or different and are each hydrogen atom or C₁-C₃ alkyl,

m is 2 or 3,

or an acid addition salt thereof.

2. A compound according to claim 1, wherein R¹ is hydrogen atom, C₁-C₁₀ alkyl, C₃-C₈ cycloalkyl, hydroxy-(C₂-C₆) alkyl, C₁-C₃ alkoxy-(C₂-C₄) alkyl, C₂-C₄ alkanoyloxy-(C₂-C₆) alkyl, a benzoyl-(C₂-C₅) alkyl group in which the phenyl moiety may optionally be substituted by halogen, C₁-C₃ alkyl or C₁-C₃ alkoxy, C₂-C₄ alkenyl, C₂-C₄ alkynyl, pyridyl, pyrimidyl, C₂-C₅ alkanoyl or furoyl,

R² and R³ are the same or different and are each hydrogen atom, halogen atom, methyl or methoxy,

R⁴, R⁵ and R⁶ are the same or different and are each hydrogen atom or C₁-C₄ alkyl, or two of them combine to form C₁-C₂ alkylene,

R⁷ and R⁸ are the same or different and are each hydrogen atom or C₁-C₃ alkyl,

m is 2,

or an acid addition salt thereof.


3. A compound according to claim 2, wherein R¹ is hydrogen atom, C₁-C₈ alkyl, C₃-C₈ cycloalkyl, hydroxy-(C₂-C₆) alkyl, C₁-C₂ alkoxy-(C₂-C₃) alkyl, acetyloxy-(C₂-C₄) alkyl, C₃-C₄ alkenyl or C₂-C₃ alkanoyl,

R² and R³ are the same or different and are each hydrogen atom or halogen atom, or one of them is hydrogen atom and another one is halogen atom, methyl or methoxy, or an acid addition salt thereof.

4. A compound according to claim 3, wherein R² and R³ are the same or different and are each hydrogen atom or fluorine atom, or an acid addition salt thereof.



5. A compound of the formula:




wherein n_1 is 3, 6 or 7,

R^{11} is hydrogen atom, C_1 - C_8 alkyl, C_3 - C_6 cycloalkyl, hydroxy- (C_2-C_4) alkyl, C_1 - C_2 alkoxy- (C_2-C_3) alkyl or C_3 alkenyl,

R^{21} and R^{31} are the same or different and are each hydrogen atom or fluorine atom,

R^{71} and R^{81} are the same or different and are each hydrogen atom or C_1 - C_3 alkyl, or an acid addition salt thereof.

6. A compound according to claim 1, which is a compound of the formula:



wherein n_2 is 4 or 5,

R^{12} is hydrogen atom, C_1 - C_8 alkyl, C_3 - C_8 cycloalkyl, hydroxy- (C_2-C_4) alkyl, C_1 - C_2 alkoxy- (C_2-C_3) alkyl or C_3 alkenyl,

R^{22} and R^{32} are the same or different and are each hydrogen atom or fluorine atom,

R^{42} , R^{52} and R^{62} are the same or different and are each hydrogen atom or C_1 - C_4 alkyl, or two of them combine to form C_1 - C_2 alkylene, or an acid addition salt thereof.

7. A compound according to claim 5, wherein n_1 is 6,

R^{11} is C_1 - C_6 alkyl, C_3 - C_6 cycloalkyl or hydroxy- (C_2-C_4) alkyl,

R^{21} and R^{31} are the same or different and are each hydrogen atom or fluorine atom substituted at 2- or 4-position of the phenyl ring,

R^{71} and R^{81} are each hydrogen atom,

or an acid addition salt thereof.

8. A compound according to claim 6, wherein n_2 is 5,


R^{12} is C_1 - C_6 alkyl, C_3 - C_6 cycloalkyl or hydroxy- (C_2-C_4) alkyl,



R^{22} and R^{32} are the same or different and are each hydrogen atom or fluorine atom substituted at 2- or 4-position of the phenyl ring,

R^{42} , R^{52} and R^{62} are all hydrogen atom, or two of them combine to form C_1 - C_2 alkylene and the other is hydrogen atom,
or an acid addition salt thereof.

9. A compound according to claim 5, which is a compound of the formula:



wherein R^{13} is methyl, ethyl, propyl, butyl, pentyl or hydroxyethyl,

R^{33} is hydrogen atom or fluorine atom, or an acid addition salt thereof.

10. 2-(4-Ethyl-1-piperazinyl)-4-(4-fluorophenyl)-5,6,7,8,9,10-hexahydrocycloocta[b]pyridine, or an acid addition salt thereof.


11. 2-(4-Ethyl-1-piperazinyl)-4-(2,4-difluorophenyl)-5,6,7,8,9,10-hexahydrocycloocta[b]pyridine, or an acid addition salt thereof.

12. 2-(4-n-Propyl-1-piperazinyl)-4-(4-fluorophenyl)-5,6,7,8,9,10-hexahydrocycloocta[b]pyridine, or an acid addition salt thereof.

13. 2-(4-Methyl-1-piperazinyl)-4-(4-fluorophenyl)-5,6,7,8,9,10-hexahydrocycloocta[b]pyridine, or an acid addition salt thereof.

14. 2-(4-Methyl-1-piperazinyl)-4-(2,4-difluorophenyl)-5,6,7,8,9,10-hexahydrocycloocta[b]pyridine, or an acid addition salt thereof.

15. A process for preparing a compound of the formula (I):



(I)

wherein n is 3, 4, 5, 6 or 7,

R^1 is hydrogen atom, C_1 - C_{10} alkyl, C_2 - C_6 alkenyl, C_2 - C_6 -alkynyl, C_3 - C_8 cycloalkyl, C_3 - C_6 cycloalkyl-(C_1 - C_4) alkyl, hydroxy-(C_2 - C_6) alkyl, C_1 - C_3 alkoxy-(C_2 - C_6) alkyl, acyloxy-(C_2 - C_6) alkyl, unsubstituted



or substituted aroyl-(C₁-C₆) alkyl, unsubstituted or substituted aryl, heteroaryl, or acyl,


R² and R³ are the same or different and are each hydrogen atom, halogen atom, C₁-C₆ alkyl, C₁-C₆ alkoxy, trifluoromethyl, or hydroxy,

R⁴, R⁵ and R⁶ are the same or different and are each hydrogen atom, C₁-C₆ alkyl, or phenyl, or two of R⁴, R⁵ and R⁶ combine to form a single bond or C₁-C₃ alkylene,


R⁷ and R⁸ are the same or different and are each hydrogen atom or C₁-C₃ alkyl, m is 2 or 3,

or an acid addition salt thereof, which comprises

(a) reacting a compound of the formula (II):




wherein X is a leaving atom or group and R², R³, R⁴, R⁵, R⁶ and n are as defined above, with a compound of the formula (III):



wherein R¹, R⁷, R⁸ and m are as defined above,


(b) in case of R¹ being hydrogen atom, subjecting to hydrogenolysis of a compound of the formula (I-4):



wherein R¹⁴ is unsubstituted or substituted benzyl or benzloxycarbonyl (the substituent on the phenyl ring of said benzyl and benzyloxycarbonyl is a member selected from lower alkyl, lower alkoxy and halogen atom), and R², R³, R⁴, R⁵, R⁶, R⁷, m and n are as defined above,




(c) in case of R¹ being hydrogen atom, reacting a compound of the formula (I-5):




20

wherein R¹⁵ is methyl or unsubstituted or substituted benzyl (the substituent on the phenyl ring of the benzyl is a member selected from lower alkyl, lower alkoxy and halogen atom), and R², R³, R⁴, R⁵, R⁶, R⁷, R⁸, m and n are as defined above, with ethyl chlorocarbonate or 1-chloroethyl chlorocarbonate to give a compound of the formula (I-6):



wherein R¹⁶ is ethoxycarbonyl or 1-chloroethoxycarbonyl, and R², R³, R⁴, R⁵, R⁶, R⁷, R⁸, m and n are as defined above, or a compound of the formula (I-7):




55


wherein R¹⁷ is acyl, R², R³, R⁴, R⁵, R⁶, R⁷, R⁸, m and n are as defined above, followed by hydrolysis of the compound of the formula (I-6) or (I-7), or



(d) in case of R¹ being other than hydrogen atom, reacting a compound of the formula (I-8):



wherein R², R³, R⁴, R⁵, R⁶, R⁷, R⁸, m and n are as defined above, with a compound of the formula (IV):




wherein Z is a reactive residue of an alcohol, and R¹⁸ is the same as R¹ except hydrogen atom, and optionally converting the form of the compound of the formula (I) into an acid addition salt thereof.

16. A psychotropic composition comprising as an active ingredient an efficient amount of the compound of the formula (I) as set forth in claim 1, or a pharmaceutically acceptable salt thereof in admixture with a pharmaceutically acceptable carrier or diluent.
17. A composition according to claim 16, wherein the active ingredient is the compound as set forth in claim 5, or a pharmaceutically acceptable salt thereof.
18. A composition according to claim 16, wherein the active ingredient is the compound as set forth in claim 6, or a pharmaceutically acceptable salt thereof.
19. A composition according to claim 16, wherein the active ingredient is the compound as set forth in claim 9, or a pharmaceutically acceptable salt thereof.
20. A compound of the formula (I) as set forth in claim 16 for use as a pharmaceutical preparation for psychotic diseases.
21. A use of a compound of the formula (I) as set forth in claim 16 for preparing a pharmaceutical preparation useful for treatment of psychotic diseases.



Claims for the following Contracting State : ES

1. A process for preparing a compound of the formula (I):



wherein n is 3, 4, 5, 6 or 7,

R¹ is hydrogen atom, C₁-C₁₀ alkyl, C₂-C₆ alkenyl, C₂-C₆-alkynyl, C₃-C₈ cycloalkyl, C₃-C₆ cycloalkyl-(C₁-C₄) alkyl, hydroxy-(C₂-C₆) alkyl, C₁-C₃ alkoxy-(C₂-C₆) alkyl, acyloxy-(C₂-C₆) alkyl, unsubstituted or substituted aroyl-(C₁-C₆) alkyl, unsubstituted or substituted aryl, heteroaryl, or acyl,

R² and R³ are the same or different and are each hydrogen atom, halogen atom, C₁-C₆ alkyl, C₁-C₆ alkoxy, trifluoromethyl, or hydroxy,


R⁴, R⁵ and R⁶ are the same or different and are each hydrogen atom, C₁-C₆ alkyl, or phenyl, or two of R⁴, R⁵ and R⁶ combine to form a single bond or C₁-C₃ alkylene,

R⁷ and R⁸ are the same or different and are each hydrogen atom or C₁-C₃ alkyl,


m is 2 or 3,

or an acid addition salt thereof, which comprises

(a) reacting a compound of the formula (II):




wherein X is a leaving atom or group and R², R³, R⁴, R⁵, R⁶ and n are as defined above, with a compound of the formula (III):






wherein R^1 , R^7 , R^8 and m are as defined above,

(b) in case of R^1 being hydrogen atom, subjecting to hydrogenolysis of a compound of the formula (I-4):




wherein R^{14} is unsubstituted or substituted benzyl or benzyloxycarbonyl (the substituent on the phenyl ring of said benzyl and benzyloxycarbonyl is a member selected from lower alkyl, lower alkoxy and halogen atom), and R^2 , R^3 , R^4 , R^5 , R^6 , R^7 , m and n are as defined above,

(c) in case of R^1 being hydrogen atom, reacting a compound of the formula (I-5):




wherein R^{15} is methyl or unsubstituted or substituted benzyl (the substituent on the phenyl ring of the benzyl is a member selected from lower alkyl, lower alkoxy and halogen atom), and R^2 , R^3 , R^4 , R^5 , R^6 , R^7 , R^8 , m and n are as defined above, with ethyl chlorocarbonate or 1-chloroethyl chlorocarbonate to give a compound of the formula (I-6):



wherein R^{16} is ethoxycarbonyl or 1-chloroethoxycarbonyl, and R^2 , R^3 , R^4 , R^5 , R^6 , R^7 , R^8 , m and n




are as defined above, or a compound of the formula (I-7):



wherein R^{17} is acyl, R^2 , R^3 , R^4 , R^5 , R^6 , R^7 , R^8 , m and n are as defined above, followed by hydrolysis of the compound of the formula (I-6) or (I-7), or

20 (d) in case of R^1 being other than hydrogen atom, reacting a compound of the formula (I-8):



wherein R^2 , R^3 , R^4 , R^5 , R^6 , R^7 , R^8 , m and n are as defined above, with a compound of the formula (IV):

40 R^{18} -Z (IV)

wherein Z is a reactive residue of an alcohol, and R^{18} is the same as R^1 except hydrogen atom, and optionally converting the a form of the compound of the formula (I) into an acid addition salt thereof.

- 45
2. A process according to claim 1, characterized in that a compound is prepared wherein R^1 is hydrogen atom, C_1 - C_{10} alkyl, C_3 - C_8 cycloalkyl, hydroxy-(C_2 - C_6) alkyl, C_1 - C_3 alkoxy-(C_2 - C_4) alkyl, C_2 - C_4 alkanoyloxy-(C_2 - C_6) alkyl, a benzoyl-(C_2 - C_5) alkyl group in which the phenyl moiety may optionally be substituted by halogen, C_1 - C_3 alkyl or C_1 - C_3 alkoxy, C_2 - C_4 alkenyl, C_2 - C_4 alkynyl, pyridyl, pyrimidyl, C_2 - C_5 alkanol or furoyl,

R^2 and R^3 are the same or different and are each hydrogen atom, halogen atom, methyl or methoxy,

R^4 , R^5 and R^6 are the same or different and are each hydrogen atom or C_1 - C_4 alkyl, or two of them combine to form C_1 - C_2 alkylene,

55 R^7 and R^8 are the same or different and are each hydrogen atom or C_1 - C_3 alkyl,

m is 2,

or an acid addition salt thereof.




3. A process according to claim 2, characterized in that a compound is prepared wherein R^1 is hydrogen atom, C_1-C_8 alkyl, C_3-C_8 cycloalkyl, hydroxy-(C_2-C_6) alkyl, C_1-C_2 alkoxy-(C_2-C_3) alkyl, acetyloxy-(C_2-C_4) alkyl, C_3-C_4 alkenyl or C_2-C_3 alkanoyl,

R^2 and R^3 are the same or different and are each hydrogen atom or halogen atom, or one of them is hydrogen atom and another one is halogen atom, methyl or methoxy, or an acid addition salt thereof.

4. A process according to claim 3, characterized in that a compound is prepared wherein R^2 and R^3 are the same or different and are each hydrogen atom or fluorine atom, or an acid addition salt thereof.

5. A process for preparing a compound of the formula:




characterized in that a compound is prepared wherein n_1 is 3, 6 or 7,

R^{11} is hydrogen atom, C_1-C_8 alkyl, C_3-C_6 cycloalkyl, hydroxy-(C_2-C_4) alkyl, C_1-C_2 alkoxy-(C_2-C_3) alkyl or C_3 alkenyl,

R^{21} and R^{31} are the same or different and are each hydrogen atom or fluorine atom,

R^{71} and R^{81} are the same or different and are each hydrogen atom or C_1-C_3 alkyl, or an acid addition salt thereof.

6. A process according to claim 1, characterized in that a compound is prepared which is a compound of the formula:



wherein n_2 is 4 or 5,


R^{12} is hydrogen atom, C_1-C_8 alkyl, C_3-C_8 cycloalkyl, hydroxy-(C_2-C_4) alkyl, C_1-C_2 alkoxy-(C_2-C_3) alkyl or C_3 alkenyl,

R^{22} and R^{32} are the same or different and are each hydrogen atom or fluorine atom,

R^{42} , R^{52} and R^{62} are the same or different and are each hydrogen atom or C_1-C_4 alkyl, or two of them combine to form C_1-C_2 alkylene, or an acid addition salt thereof.



7. A process according to claim 5 characterized in that a compound is prepared wherein n_1 is 6,
 R^{11} is C_1 - C_6 alkyl, C_3 - C_6 cycloalkyl or hydroxy- (C_2-C_4) alkyl,
 R^{21} and R^{31} are the same or different and are each hydrogen atom or fluorine atom substituted at
 2- or 4-position of the phenyl ring,
 R^{71} and R^{81} are each hydrogen atom,
 or an acid addition salt thereof.
8. A process according to claim 6, characterized in that a compound is prepared wherein n_2 is 5,
 R^{12} is C_1 - C_6 alkyl, C_3 - C_6 cycloalkyl or hydroxy- (C_2-C_4) alkyl,
 R^{22} and R^{32} are the same or different and are each hydrogen atom or fluorine atom substituted at
 2- or 4-position of the phenyl ring,
 R^{42} , R^{52} and R^{62} are all hydrogen atom, or two of them combine to form C_1 - C_2 alkylene and the
 other is hydrogen atom,
 or an acid addition salt thereof.
9. A process according to claim 5, characterized in that a compound is prepared which is a compound of
 the formula:




- wherein R^{13} is methyl, ethyl, propyl, butyl, pentyl or hydroxyethyl,
 R^{33} is hydrogen atom or fluorine atom,
 or an acid addition salt thereof.
10. The process of claim 1 characterized in that 2-(4-Ethyl-1-piperazinyl)-4-(4-fluorophenyl)-5,6,7,8,9,10-hexahydrocycloocta[b]pyridine, or an acid addition salt thereof is prepared.
11. The process of claim 1 characterized in that 2-(4-Ethyl-1-piperazinyl)-4-(2,4-difluorophenyl)-5,6,7,8,9,10-hexahydrocycloocta[b]pyridine, or an acid addition salt thereof is prepared.
12. The process of claim 1 characterized in that 2-(4-n-Propyl-1-piperazinyl)-4-(4-fluorophenyl)-5,6,7,8,9,10-hexahydrocycloocta[b]pyridine, or an acid addition salt thereof is prepared.
13. The process of claim 1 characterized in that 2-(4-Methyl-1-piperazinyl)-4-(4-fluorophenyl)-5,6,7,8,9,10-hexahydrocycloocta[b]pyridine, or an acid addition salt thereof is prepared.
14. The process of claim 1 characterized in that 2-(4-Methyl-1-piperazinyl)-4-(2,4-difluorophenyl)-5,6,7,8,9,10-hexahydrocycloocta[b]pyridine, or an acid addition salt thereof is prepared.
15. A process for preparing a psychotropic composition comprising mixing as an active ingredient an efficient amount of the compound of the formula (I) as set forth in claim 1, or a pharmaceutically acceptable salt thereof and a pharmaceutically acceptable carrier or diluent.
16. The process according to claim 15, wherein the active ingredient is the compound as prepared in claim 5, or a pharmaceutically acceptable salt thereof.
17. The process according to claim 15, wherein the active ingredient is the compound as prepared in claim 6, or a pharmaceutically acceptable salt thereof.



18. The process according to claim 15, wherein the active ingredient is the compound as prepared in claim 9, or a pharmaceutically acceptable salt thereof.

Claims for the following Contracting State : GR

1. A compound of the formula (I):



wherein n is 3, 4, 5, 6 or 7,

R¹ is hydrogen atom, C₁-C₁₀ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₃-C₆ cycloalkyl-(C₁-C₄) alkyl, hydroxy-(C₂-C₆) alkyl, C₁-C₃ alkoxy-(C₂-C₆) alkyl, acyloxy-(C₂-C₆) alkyl, unsubstituted or substituted aroyl-(C₁-C₆) alkyl, unsubstituted or substituted aryl, heteroaryl, or acyl,

R² and R³ are the same or different and are each hydrogen atom, halogen atom, C₁-C₆ alkyl, C₁-C₆ alkoxy, trifluoromethyl, or hydroxy,

R⁴, R⁵ and R⁶ are the same or different and are each hydrogen atom, C₁-C₆ alkyl, or phenyl, or two of R⁴, R⁵ and R⁶ combine to form a single bond or C₁-C₃ alkylene,

R⁷ and R⁸ are the same or different and are each hydrogen atom or C₁-C₃ alkyl,

m is 2 or 3,

or an acid addition salt thereof.

2. A compound according to claim 1, wherein R¹ is hydrogen atom, C₁-C₁₀ alkyl, C₃-C₈ cycloalkyl, hydroxy-(C₂-C₆) alkyl, C₁-C₃ alkoxy-(C₂-C₄) alkyl, C₂-C₄ alkanoyloxy-(C₂-C₆) alkyl, a benzoyl-(C₂-C₅) alkyl group in which the phenyl moiety may optionally be substituted by halogen, C₁-C₃ alkyl or C₁-C₃ alkoxy, C₂-C₄ alkenyl, C₂-C₄ alkynyl, pyridyl, pyrimidyl, C₂-C₅ alkanolyl or furoyl,

R² and R³ are the same or different and are each hydrogen atom, halogen atom, methyl or methoxy,

R⁴, R⁵ and R⁶ are the same or different and are each hydrogen atom or C₁-C₄ alkyl, or two of them combine to form C₁-C₂ alkylene,

R⁷ and R⁸ are the same or different and are each hydrogen atom or C₁-C₃ alkyl,

m is 2,

or an acid addition salt thereof.


3. A compound according to claim 2, wherein R¹ is hydrogen atom, C₁-C₈ alkyl, C₃-C₈ cycloalkyl, hydroxy-(C₂-C₆) alkyl, C₁-C₂ alkoxy-(C₂-C₃) alkyl, acetyloxy-(C₂-C₄) alkyl, C₃-C₄ alkenyl or C₂-C₃ alkanoyl,

R² and R³ are the same or different and are each hydrogen atom or halogen atom, or one of them is hydrogen atom and another one is halogen atom, methyl or methoxy, or an acid addition salt thereof.

4. A compound according to claim 3, wherein R² and R³ are the same or different and are each hydrogen atom or fluorine atom, or an acid addition salt thereof.



5. A compound of the formula:




wherein n_1 is 3, 6 or 7,

R^{11} is hydrogen atom, C_1 - C_8 alkyl, C_3 - C_6 cycloalkyl, hydroxy- $(C_2$ - $C_4)$ alkyl, C_1 - C_2 alkoxy- $(C_2$ - $C_3)$ alkyl or C_3 alkenyl,

R^{21} and R^{31} are the same or different and are each hydrogen atom or fluorine atom,

R^{71} and R^{81} are the same or different and are each hydrogen atom or C_1 - C_3 alkyl, or an acid addition salt thereof.

6. A compound according to claim 1, which is a compound of the formula:



wherein n_2 is 4 or 5,

R^{12} is hydrogen atom, C_1 - C_8 alkyl, C_3 - C_8 cycloalkyl, hydroxy- $(C_2$ - $C_4)$ alkyl, C_1 - C_2 alkoxy- $(C_2$ - $C_3)$ alkyl or C_3 alkenyl,

R^{22} and R^{32} are the same or different and are each hydrogen atom or fluorine atom,

R^{42} , R^{52} and R^{62} are the same or different and are each hydrogen atom or C_1 - C_4 alkyl, or two of them combine to form C_1 - C_2 alkylene, or an acid addition salt thereof.

7. A compound according to claim 5, wherein n_1 is 6,

R^{11} is C_1 - C_6 alkyl, C_3 - C_6 cycloalkyl or hydroxy- $(C_2$ - $C_4)$ alkyl,

R^{21} and R^{31} are the same or different and are each hydrogen atom or fluorine atom substituted at 2- or 4-position of the phenyl ring,

R^{71} and R^{81} are each hydrogen atom, or an acid addition salt thereof.

8. A compound according to claim 6, wherein n_2 is 5,

R^{12} is C_1 - C_6 alkyl, C_3 - C_6 cycloalkyl or hydroxy- $(C_2$ - $C_4)$ alkyl,


R^{22} and R^{32} are the same or different and are each hydrogen atom or fluorine atom substituted at



2- or 4-position of the phenyl ring,

R^{42} , R^{52} and R^{62} are all hydrogen atom, or two of them combine to form C_1 - C_2 alkylene and the other is hydrogen atom, or an acid addition salt thereof.

9. A compound according to claim 5, which is a compound of the formula:



wherein R^{13} is methyl, ethyl, propyl, butyl, pentyl or hydroxyethyl,

R^{33} is hydrogen atom or fluorine atom, or an acid addition salt thereof.

10. 2-(4-Ethyl-1-piperazinyl)-4-(4-fluorophenyl)-5,6,7,8,9,10-hexahydrocycloocta[b]pyridine, or an acid addition salt thereof.


11. 2-(4-Ethyl-1-piperazinyl)-4-(2,4-difluorophenyl)-5,6,7,8,9,10-hexahydrocycloocta[b]pyridine, or an acid addition salt thereof.

12. 2-(4-n-Propyl-1-piperazinyl)-4-(4-fluorophenyl)-5,6,7,8,9,10-hexahydrocycloocta[b]pyridine, or an acid addition salt thereof.

13. 2-(4-Methyl-1-piperazinyl)-4-(4-fluorophenyl)-5,6,7,8,9,10-hexahydrocycloocta[b]pyridine, or an acid addition salt thereof.

14. 2-(4-Methyl-1-piperazinyl)-4-(2,4-difluorophenyl)-5,6,7,8,9,10-hexahydrocycloocta[b]pyridine, or an acid addition salt thereof.

15. A process for preparing a compound of the formula (I):



(I)

wherein n is 3, 4, 5, 6 or 7,

R^1 is hydrogen atom, C_1 - C_{10} alkyl, C_2 - C_6 alkenyl, C_2 - C_6 -alkynyl, C_3 - C_8 cycloalkyl, C_3 - C_6 cycloalkyl- $(C_1$ - $C_4)$ alkyl, hydroxy- $(C_2$ - $C_6)$ alkyl, C_1 - C_3 alkoxy- $(C_2$ - $C_6)$ alkyl, acyloxy- $(C_2$ - $C_6)$ alkyl, unsubstituted



or substituted aroyl-(C₁-C₆) alkyl, unsubstituted or substituted aryl, heteroaryl, or acyl,


R² and R³ are the same or different and are each hydrogen atom, halogen atom, C₁-C₆ alkyl, C₁-C₆ alkoxy, trifluoromethyl, or hydroxy,

R⁴, R⁵ and R⁶ are the same or different and are each hydrogen atom, C₁-C₆ alkyl, or phenyl, or two of R⁴, R⁵ and R⁶ combine to form a single bond or C₁-C₃ alkylene,


R⁷ and R⁸ are the same or different and are each hydrogen atom or C₁-C₃ alkyl, m is 2 or 3,

or an acid addition salt thereof, which comprises

(a) reacting a compound of the formula (II):




wherein X is a leaving atom or group and R², R³, R⁴, R⁵, R⁶ and n are as defined above, with a compound of the formula (III):



wherein R¹, R⁷, R⁸ and m are as defined above,


(b) in case of R¹ being hydrogen atom, subjecting to hydrogenolysis of a compound of the formula (I-4):



wherein R¹⁴ is unsubstituted or substituted benzyl or benzyloxycarbonyl (the substituent on the phenyl ring of said benzyl and benzyloxycarbonyl is a member selected from lower alkyl, lower alkoxy and halogen atom), and R², R³, R⁴, R⁵, R⁶, R⁷, m and n are as defined above,




(c) in case of R^1 being hydrogen atom, reacting a compound of the formula (I-5):




20

wherein R^{15} is methyl or unsubstituted or substituted benzyl (the substituent on the phenyl ring of the benzyl is a member selected from lower alkyl, lower alkoxy and halogen atom), and R^2 , R^3 , R^4 , R^5 , R^6 , R^7 , R^8 , m and n are as defined above, with ethyl chlorocarbonate or 1-chloroethyl chlorocarbonate to give a compound of the formula (I-6):



wherein R^{16} is ethoxycarbonyl or 1-chloroethoxycarbonyl, and R^2 , R^3 , R^4 , R^5 , R^6 , R^7 , R^8 , m and n are as defined above, or a compound of the formula (I-7):




55

wherein R^{17} is acyl, R^2 , R^3 , R^4 , R^5 , R^6 , R^7 , R^8 , m and n are as defined above, followed by hydrolysis of the compound of the formula (I-6) or (I-7), or



(d) in case of R^1 being other than hydrogen atom, reacting a compound of the formula (I-8):



wherein R^2 , R^3 , R^4 , R^5 , R^6 , R^7 , R^8 , m and n are as defined above, with a compound of the formula (IV):



wherein Z is a reactive residue of an alcohol, and R^{18} is the same as R^1 except hydrogen atom, and optionally converting the a form of the compound of the formula (I) into an acid addition salt thereof.

25

16. A composition comprising as an active ingredient an efficient amount of the compound of the formula (I) as set forth in claim 1, or a salt thereof in admixture with a carrier or diluent.

30

17. A composition according to claim 16, wherein the active ingredient is the compound as set forth in claim 5, or a salt thereof.

18. A composition according to claim 16, wherein the active ingredient is the compound as set forth in claim 6, or a salt thereof.


35

19. A composition according to claim 16, wherein the active ingredient is the compound as set forth in claim 9, or a salt thereof.

Patentansprüche

40 Patentansprüche für folgende Vertragsstaaten : AT, BE, CH, DE, DK, FR, GB, IT, LI, LU, NL, SE

1. Verbindung der Formel (I):





worin

n 3, 4, 5, 6 oder 7 bedeutet,

R^1 ein Wasserstoffatom, C_1 - C_{10} -Alkyl, C_2 - C_6 -Alkenyl, C_2 - C_6 -Alkynyl, C_3 - C_8 -Cycloalkyl, C_3 - C_6 -Cycloalkyl(C_1 - C_4)-alkyl, Hydroxy(C_2 - C_6)alkyl, C_1 - C_3 -Alkoxy(C_2 - C_6)alkyl, Acyloxy(C_2 - C_6)alkyl, unsubstituiertes oder substituiertes Aryl-(C_1 - C_5)alkyl, unsubstituiertes oder substituiertes Aryl, Heteroaryl oder Acyl bedeutet,

R^2 und R^3 gleich oder unterschiedlich sind und je ein Wasserstoffatom, ein Halogenatom, C_1 - C_6 -Alkyl, C_1 - C_6 -Alkoxy, Trifluormethyl oder Hydroxy bedeuten,

R^4 , R^5 und R^6 gleich oder unterschiedlich sind und je ein Wasserstoffatom, C_1 - C_6 -Alkyl oder Phenyl bedeuten, oder worin zwei von R^4 , R^5 und R^6 unter Bildung einer Einfachbindung oder C_1 - C_3 -Alkylen gebunden sind,

R^7 und R^8 gleich oder unterschiedlich sind und je ein Wasserstoffatom oder C_1 - C_3 -Alkyl bedeuten, m 2 oder 3 bedeutet,

oder ein Säureadditionssalz davon.

2. Verbindung nach Anspruch 1, dadurch gekennzeichnet, daß R^1 ein Wasserstoffatom, C_1 - C_{10} -Alkyl, C_3 - C_8 -Cycloalkyl, Hydroxy(C_2 - C_6)alkyl, C_1 - C_3 -Alkoxy(C_2 - C_4)-alkyl, C_2 - C_4 -Alkanoyloxy(C_2 - C_6)alkyl, eine Benzoyl(C_2 - C_5)-alkylgruppe, worin die Phenylgruppierung gegebenenfalls durch Halogen, C_1 - C_3 -Alkyl oder C_1 - C_3 -Alkoxy substituiert sein kann, C_2 - C_4 -Alkenyl, C_2 - C_4 -Alkynyl, Pyridyl, Pyrimidyl, C_2 - C_5 -Alkanoyl oder Furoyl bedeutet,

R^2 und R^3 gleich oder unterschiedlich sind und je ein Wasserstoffatom, ein Halogenatom, Methyl oder Methoxy bedeuten,

R^4 , R^5 und R^6 gleich oder unterschiedlich sind und je ein Wasserstoffatom oder C_1 - C_4 -Alkyl bedeuten oder worin zwei oder mehrere von diesen unter Bildung von C_1 - C_2 -Alkylen gebunden sind,

R^7 und R^8 gleich oder unterschiedlich sind und je ein Wasserstoffatom oder C_1 - C_3 -Alkyl bedeuten, m 2 bedeutet,

oder ein Säureadditionssalz davon.


3. Verbindung nach Anspruch 2, dadurch gekennzeichnet, daß R^1 ein Wasserstoffatom, C_1 - C_8 -Alkyl, C_3 - C_8 -Cycloalkyl, Hydroxy(C_2 - C_6)alkyl, C_1 - C_2 -Alkoxy(C_2 - C_3)-alkyl, Acetyloxy(C_2 - C_4)alkyl, C_3 - C_4 -Alkenyl oder C_2 - C_3 -Alkanoyl bedeutet,

R^2 und R^3 gleich oder unterschiedlich sind und je ein Wasserstoffatom oder Halogenatom bedeuten oder worin einer von ihnen ein Wasserstoffatom und der andere ein Halogenatom, Methyl oder Methoxy bedeutet,

oder ein Säureadditionssalz davon.

4. Verbindung nach Anspruch 3, dadurch gekennzeichnet, daß R^2 und R^3 gleich oder unterschiedlich sind und je ein Wasserstoffatom oder ein Fluoratom bedeuten, oder ein Säureadditionssalz davon.

5. Verbindung der Formel:



worin


n_1 3, 6 oder 7 bedeutet,

R^{11} ein Wasserstoffatom, C_1 - C_8 -Alkyl, C_3 - C_6 -Cycloalkyl, Hydroxy(C_2 - C_4)alkyl, C_1 - C_2 -Alkoxy(C_2 - C_3)alkyl oder C_3 -Alkenyl bedeutet,

R^{21} und R^{31} gleich oder unterschiedlich sind und je ein Wasserstoffatom oder ein Fluoratom bedeuten,

R^{71} und R^{81} gleich oder unterschiedlich sind und je ein Wasserstoffatom oder C_1 - C_3 -Alkyl bedeuten, oder ein Säureadditionssalz davon.

6. Verbindung nach Anspruch 1, dadurch gekennzeichnet, daß sie eine Verbindung der Formel:



ist, worin

n_2 4 oder 5 bedeutet,

R^{12} ein Wasserstoffatom, C_1 - C_8 -Alkyl, C_3 - C_6 -Cycloalkyl, Hydroxy(C_2 - C_4)alkyl, C_1 - C_2 -Alkoxy(C_2 - C_3)alkyl oder C_3 -Alkenyl bedeutet,

R^{22} und R^{32} gleich oder unterschiedlich sind und je ein Wasserstoffatom oder ein Fluoratom bedeuten,

R^{42} , R^{52} und R^{62} gleich oder unterschiedlich sind und je ein Wasserstoffatom oder C_1 - C_4 -Alkyl bedeuten oder worin zwei davon unter Bildung von C_1 - C_2 -Alkylen gebunden sind, oder ein Säureadditionssalz davon.

7. Verbindung nach Anspruch 5, dadurch gekennzeichnet, daß n_1 6 bedeutet,

R^{11} C_1 - C_6 -Alkyl, C_3 - C_6 -Cycloalkyl oder Hydroxy(C_2 - C_4)-alkyl bedeutet,

R^{21} und R^{31} gleich oder unterschiedlich sind und je ein Wasserstoffatom oder Fluoratom, substituiert in der 2- oder 4-Stellung des Phenylrings, bedeuten,

R^{71} und R^{81} je ein Wasserstoffatom bedeuten, oder ein Säureadditionssalz davon.


8. Verbindung nach Anspruch 6, dadurch gekennzeichnet, daß n_2 5 bedeutet,

R^{12} C_1 - C_6 -Alkyl, C_3 - C_6 -Cycloalkyl oder Hydroxy(C_2 - C_4)-alkyl bedeutet,

R^{22} und R^{32} gleich oder unterschiedlich sind und je ein Wasserstoffatom oder Fluoratom, substituiert in der 2- oder 4-Stellung des Phenylrings, bedeuten,

R^{42} , R^{52} und R^{62} alle ein Wasserstoffatom bedeuten oder worin zwei davon unter Bildung von C_1 - C_2 -Alkylen gebunden sind und der andere ein Wasserstoffatom bedeutet, oder ein Säureadditionssalz davon.

9. Verbindung nach Anspruch 5, dadurch **gekennzeichnet**, daß sie eine Verbindung der Formel:




ist, worin

R^{13} Methyl, Ethyl, Propyl, Butyl, Pentyl oder Hydroxyethyl bedeutet,

R^{33} ein Wasserstoffatom oder Fluoratom bedeutet, oder ein Säureadditionssalz davon.

10. 2-(4-Ethyl-1-piperazinyl)-4-(4-fluorphenyl)-5,6,7,8,9,10-hexahydrocycloocta[b]pyridin oder ein Säureadditionssalz davon.
11. 2-(4-Ethyl-1-piperazinyl)-4-(2,4-difluorphenyl)-5,6,7,8,9,10-hexahydrocycloocta[b]pyridin oder ein Säureadditionssalz davon.
12. 2-(4-n-Propyl-1-piperazinyl)-4-(4-fluorphenyl)-5,6,7,8,9,10-hexahydrocycloocta[b]pyridin oder ein Säureadditionssalz davon.
13. 2-(4-Methyl-1-piperazinyl)-4-(4-fluorphenyl)-5,6,7,8,9,10-hexahydrocycloocta[b]pyridin oder ein Säureadditionssalz davon.
14. 2-(4-Methyl-1-piperazinyl)-4-(2,4-difluorphenyl)-5,6,7,8,9,10-hexahydrocycloocta[b]pyridin oder ein Säureadditionssalz davon.
15. Verfahren zur Herstellung einer Verbindung der Formel (I):



worin

n 3, 4, 5, 6 oder 7 bedeutet,

R^1 ein Wasserstoffatom, C_1 - C_{10} -Alkyl, C_2 - C_6 -Alkenyl, C_2 - C_6 -Alkynyl, C_3 - C_8 -Cycloalkyl, C_3 - C_6 -Cycloalkyl(C_1 - C_4)-alkyl, Hydroxy(C_2 - C_6)alkyl, C_1 - C_3 -Alkoxy(C_2 - C_6)alkyl, Acyloxy(C_2 - C_6)alkyl, unsubstituiertes oder substituiertes Aroyl(C_1 - C_6)alkyl, unsubstituiertes oder substituiertes Aryl, Heteroaryl oder Acyl bedeutet,

R^2 und R^3 gleich oder unterschiedlich sind und je ein Wasserstoffatom, ein Halogenatom, C_1 - C_6 -




Alkyl, C₁-C₆-Alkoxy, Trifluormethyl oder Hydroxy bedeuten,

R⁴, R⁵ und R⁶ gleich oder unterschiedlich sind und je ein Wasserstoffatom, C₁-C₆-Alkyl oder Phenyl bedeuten oder worin zwei von R⁴, R⁵ und R⁶ unter Bildung einer Einfachbindung oder C₁-C₃-Alkylen gebunden sind,


R⁷ und R⁸ gleich oder unterschiedlich sind und je ein Wasserstoffatom oder C₁-C₃-Alkyl bedeuten, m 2 oder 3 bedeutet,

oder eines Säureadditionssalzes davon, dadurch gekennzeichnet, daß

(a) eine Verbindung der Formel (II):




worin X ein Austrittsatom oder eine Austrittsgruppe bedeutet und R², R³, R⁴, R⁵, R⁶ und n die oben gegebenen Definitionen besitzen, mit einer Verbindung der Formel (III):



worin R¹, R⁷, R⁸ und m die oben gegebenen Definitionen besitzen, umgesetzt wird,


(b) im Falle, wenn R¹ ein Wasserstoffatom bedeutet, eine Verbindung der Formel (I-4):




worin R¹⁴ unsubstituiertes oder substituiertes Benzyl oder Benzyloxycarbonyl bedeutet (wobei der Substituent an dem Phenylring von Benzyl und Benzyloxycarbonyl ein Glied, ausgewählt aus Niedrigalkyl, Niedrigalkoxy und einem Halogenatom, ist) und R², R³, R⁴, R⁵, R⁶, R⁷, m und n die oben gegebenen Definitionen besitzen, der Hydrogenolyse unterworfen wird,




(c) im Falle, wenn R^1 ein Wasserstoffatom bedeutet, eine Verbindung der Formel (I-5):



worin R^{15} Methyl oder unsubstituiertes oder substituiertes Benzyl bedeutet (wobei der Substituent an dem Phenylring des Benzyls ein Glied, ausgewählt aus Niedrigalkyl, Niedrigalkoxy und einem Halogenatom, ist) und R^2 , R^3 , R^4 , R^5 , R^6 , R^7 , R^8 , m und n die oben gegebenen Definitionen besitzen, mit Ethylchlorcarbonat oder 1-Chlorethylchlorcarbonat unter Bildung einer Verbindung der Formel (I-6):




worin R^{16} Ethoxycarbonyl oder 1-Chlorethoxycarbonyl bedeutet und R^2 , R^3 , R^4 , R^5 , R^6 , R^7 , R^8 , m und n die oben gegebenen Definitionen besitzen, oder einer Verbindung der Formel (I-7):



worin R^{17} Acyl bedeutet, R^2 , R^3 , R^4 , R^5 , R^6 , R^7 , R^8 , m und n die oben gegebenen Definitionen besitzen, umgesetzt wird, gefolgt von einer Hydrolyse der Verbindung der Formel (I-6) oder (I-7), oder



(d) in Falle, wenn R^1 eine andere Bedeutung als ein Wasserstoffatom besitzt, eine Verbindung der Formel (I-8):



worin R^2 , R^3 , R^4 , R^5 , R^6 , R^7 , R^8 , m und n die oben gegebenen Definitionen besitzen, mit einer Verbindung der Formel (IV):

20 $R^{18}-Z$ (IV)


25 worin Z den reaktiven Rest eines Alkohols bedeutet und R^{18} die gleiche Bedeutung wie R^1 , ausgenommen ein Wasserstoffatom, besitzt, umgesetzt wird, und gegebenenfalls die Form der Verbindung der Formel (I) in ein Säureadditionssalz davon umgewandelt wird.

- 30 16. Psychotropische Zusammensetzung, dadurch gekennzeichnet, daß sie als aktiven Bestandteil eine wirksame Menge der Verbindung der Formel (I), wie in Anspruch 1 angegeben, oder ein pharmazeutisch annehmbares Salz davon im Gemisch mit einem pharmazeutisch annehmbaren Träger oder Verdünnungsmittel enthält.
- 35 17. Zusammensetzung nach Anspruch 16, dadurch gekennzeichnet, daß der aktive Bestandteil die Verbindung nach Anspruch 5 oder ein pharmazeutisch annehmbares Salz davon ist.
18. Zusammensetzung nach Anspruch 16, dadurch gekennzeichnet, daß der aktive Bestandteil die Verbindung nach Anspruch 6 oder ein pharmazeutisch annehmbares Salz davon ist.
- 40 19. Zusammensetzung nach Anspruch 16, dadurch gekennzeichnet, daß der aktive Bestandteil die Verbindung nach Anspruch 9 oder ein pharmazeutisch annehmbares Salz davon ist.
20. Verbindung der Formel (I) nach Anspruch 16 für die Verwendung als pharmazeutisches Präparat für psychotische Krankheiten.
- 45 21. Verwendung einer Verbindung der Formel (I) nach Anspruch 16 für die Herstellung eines pharmazeutischen Präparats, das für die Behandlung psychotischer Krankheiten nützlich ist.



Patentansprüche für folgenden Vertragsstaat : ES

1. Verfahren zur Herstellung einer Verbindung der Formel (I):



worin

n 3, 4, 5, 6 oder 7 bedeutet,

R^1 ein Wasserstoffatom, C_1 - C_{10} -Alkyl, C_2 - C_6 -Alkenyl, C_2 - C_6 -Alkynyl, C_3 - C_8 -Cycloalkyl, C_3 - C_6 -Cycloalkyl(C_1 - C_4)alkyl, Hydroxy(C_2 - C_6)alkyl, C_1 - C_3 -Alkoxy(C_2 - C_6)alkyl, Acyloxy(C_2 - C_6)alkyl, unsubstituiertes oder substituiertes Aroyl-(C_1 - C_6)alkyl, unsubstituiertes oder substituiertes Aryl, Heteroaryl oder Acyl bedeutet,

R^2 und R^3 gleich oder unterschiedlich sind und je ein Wasserstoffatom, ein Halogenatom, C_1 - C_6 -Alkyl, C_1 - C_6 -Alkoxy, Trifluormethyl oder Hydroxy bedeuten,


R^4 , R^5 und R^6 gleich oder unterschiedlich sind und je ein Wasserstoffatom, C_1 - C_6 -Alkyl oder Phenyl bedeuten oder worin zwei von R^4 , R^5 und R^6 unter Bildung einer Einfachbindung oder C_1 - C_3 -Alkylen gebunden sind,

R^7 und R^8 gleich oder unterschiedlich sind und je ein Wasserstoffatom oder C_1 - C_3 -Alkyl bedeuten,


m 2 oder 3 bedeutet,

oder eines Säureadditionssalzes davon, dadurch gekennzeichnet, daß

(a) eine Verbindung der Formel (II):



worin X ein Austrittsatom oder eine Austrittsgruppe bedeutet und R^2 , R^3 , R^4 , R^5 , R^6 und n die oben gegebenen Definitionen besitzen, mit einer Verbindung der Formel (III):




15

20

25

30


worin R^1 , R^7 , R^8 und m die oben gegebenen Definitionen besitzen, umgesetzt wird,
 (b) im Falle, wenn R^1 ein Wasserstoffatom bedeutet, eine Verbindung der Formel (I-4):



50

55

worin R^{14} unsubstituiertes oder substituiertes Benzyl oder Benzyloxycarbonyl bedeutet (wobei der Substituent an dem Phenylring von Benzyl und Benzyloxycarbonyl ein Glied, ausgewählt aus Niedrigalkyl, Niedrigalkoxy und einem Halogenatom, ist) und R^2 , R^3 , R^4 , R^5 , R^6 , R^7 , m und n die oben gegebenen Definitionen besitzen, der Hydrogenolyse unterworfen wird,
 (c) im Falle, wenn R^1 ein Wasserstoffatom bedeutet, eine Verbindung der Formel (I-5):




75


80

85

worin R^{15} Methyl oder unsubstituiertes oder substituiertes Benzyl bedeutet (wobei der Substituent an dem Phenylring des Benzyls ein Glied, ausgewählt aus Niedrigalkyl, Niedrigalkoxy und einem Halogenatom, ist) und R^2 , R^3 , R^4 , R^5 , R^6 , R^7 , R^8 , m und n die oben gegebenen Definitionen besitzen, mit Ethylchlorcarbonat oder 1-Chlorethylchlorcarbonat unter Bildung einer Verbindung der Formel (I-6):




15 worin R^{16} Ethoxycarbonyl oder 1-Chlorethoxycarbonyl bedeutet und $R^2, R^3, R^4, R^5, R^6, R^7, R^8, m$ und n die oben gegebenen Definitionen besitzen, oder einer Verbindung der Formel (I-7):



35 worin R^{17} Acyl bedeutet, $R^2, R^3, R^4, R^5, R^6, R^7, R^8, m$ und n die oben gegebenen Definitionen besitzen, umgesetzt wird, gefolgt von einer Hydrolyse der Verbindung der Formel (I-6) oder (I-7), oder

(d) in Falle, wenn R^1 eine andere Bedeutung als ein Wasserstoffatom besitzt, eine Verbindung der Formel (I-8):



55 worin $R^2, R^3, R^4, R^5, R^6, R^7, R^8, m$ und n die oben gegebenen Definitionen besitzen, mit einer Verbindung der Formel (IV):

$R^{18}-Z$ (IV)

worin Z den reaktiven Rest eines Alkohols bedeutet und R^{18} die gleiche Bedeutung wie R^1 ,



ausgenommen ein Wasserstoffatom, besitzt, umgesetzt wird,
und gegebenenfalls die Form der Verbindung der Formel (I) in ein Säureadditionssalz davon
umgewandelt wird.

- 5 2. Verfahren nach Anspruch 1, dadurch **gekennzeichnet**, daß eine Verbindung, worin R¹ ein Wasserstoffatom, C₁-C₁₀-Alkyl, C₃-C₈-Cycloalkyl, Hydroxy(C₂-C₆)alkyl, C₁-C₃-Alkoxy(C₂-C₄)alkyl, C₂-C₄-Alkanoyloxy(C₂-C₅)alkyl, eine Benzoyl(C₂-C₅)alkylgruppe, worin die Phenylgruppierung gegebenenfalls durch Halogen, C₁-C₃-Alkyl oder C₁-C₃-Alkoxy substituiert sein kann, C₂-C₄-Alkenyl, C₂-C₄-Alkynyl, Pyridyl, Pyrimidyl, C₂-C₅-Alkanoyl oder Furoyl bedeutet,

10 R² und R³ gleich oder unterschiedlich sind und je ein Wasserstoffatom, ein Halogenatom, Methyl oder Methoxy bedeuten,

R⁴, R⁵ und R⁶ gleich oder unterschiedlich sind und je ein Wasserstoffatom oder C₁-C₄-Alkyl bedeuten oder worin zwei von diesen unter Bildung von C₁-C₂-Alkylen gebunden sind,

15 R⁷ und R⁸ gleich oder unterschiedlich sind und je ein Wasserstoffatom oder C₁-C₃-Alkyl bedeuten, m 2 bedeutet,

oder ein Säureadditionssalz davon hergestellt wird.


3. Verfahren nach Anspruch 2, dadurch **gekennzeichnet**, daß eine Verbindung, worin R¹ ein Wasserstoffatom, C₁-C₈-Alkyl, C₃-C₈-Cycloalkyl, Hydroxy(C₂-C₆)alkyl, C₁-C₂-Alkoxy(C₂-C₃)alkyl, Acetyloxy(C₂-C₄)alkyl, C₃-C₄-Alkenyl oder C₂-C₃-Alkanoyl bedeutet,

20 R² und R³ gleich oder unterschiedlich sind und je ein Wasserstoffatom oder Halogenatom bedeuten oder worin einer von ihnen ein Wasserstoffatom und der andere ein Halogenatom, Methyl oder Methoxy bedeutet,

25 oder ein Säureadditionssalz davon hergestellt wird.

4. Verfahren nach Anspruch 3, dadurch **gekennzeichnet**, daß eine Verbindung, worin R² und R³ gleich oder unterschiedlich sind und je ein Wasserstoffatom oder ein Fluoratom bedeuten, oder ein Säureadditionssalz davon hergestellt wird.

- 30 5. Verfahren zur Herstellung einer Verbindung der Formel:



45 dadurch **gekennzeichnet**, daß eine Verbindung, worin n₁ 3, 6 oder 7 bedeutet,


R¹¹ ein Wasserstoffatom, C₁-C₈-Alkyl, C₃-C₆-Cycloalkyl, Hydroxy(C₂-C₄)alkyl, C₁-C₂-Alkoxy(C₂-C₃)alkyl oder C₃-Alkenyl bedeutet,

50 R²¹ und R³¹ gleich oder unterschiedlich sind und je ein Wasserstoffatom oder ein Fluoratom bedeuten,

R⁷¹ und R⁸¹ gleich oder unterschiedlich sind und je ein Wasserstoffatom oder C₁-C₃-Alkyl bedeuten,

oder ein Säureadditionssalz davon hergestellt wird.

- 55 6. Verfahren nach Anspruch 1, dadurch **gekennzeichnet**, daß eine Verbindung, die eine Verbindung der Formel:



ist, worin

n_2 4 oder 5 bedeutet,

R^{12} ein Wasserstoffatom, C_1 - C_8 -Alkyl, C_3 - C_8 -Cycloalkyl, Hydroxy(C_2 - C_4)alkyl, C_1 - C_2 -Alkoxy(C_2 - C_3)alkyl oder C_3 -Alkenyl bedeutet,

R^{22} und R^{32} gleich oder unterschiedlich sind und je ein Wasserstoffatom oder ein Fluoratom bedeuten,

R^{42} , R^{52} und R^{62} gleich oder unterschiedlich sind und je ein Wasserstoffatom oder C_1 - C_4 -Alkyl bedeuten oder worin zwei davon unter Bildung von C_1 - C_2 -Alkylen gebunden sind, oder ein Säureadditionssalz davon hergestellt wird.

7. Verfahren nach Anspruch 5, dadurch **gekennzeichnet**, daß eine Verbindung, worin n_1 6 bedeutet,

R^{11} C_1 - C_6 -Alkyl, C_3 - C_6 -Cycloalkyl oder Hydroxy(C_2 - C_4)alkyl bedeutet,

R^{21} und R^{31} gleich oder unterschiedlich sind und je ein Wasserstoffatom oder Fluoratom, substituiert in der 2- oder 4-Stellung des Phenylrings, bedeuten,

R^{71} und R^{81} je ein Wasserstoffatom bedeuten, oder ein Säureadditionssalz davon hergestellt wird.


8. Verfahren nach Anspruch 6, dadurch **gekennzeichnet**, daß eine Verbindung, worin n_2 5 bedeutet,

R^{12} C_1 - C_6 -Alkyl, C_3 - C_6 -Cycloalkyl oder Hydroxy(C_2 - C_4)alkyl bedeutet,

R^{22} und R^{32} gleich oder unterschiedlich sind und je ein Wasserstoffatom oder Fluoratom, substituiert in der 2- oder 4-Stellung des Phenylrings, bedeuten,

R^{42} , R^{52} und R^{62} alle ein Wasserstoffatom bedeuten oder worin zwei davon unter Bildung von C_1 - C_2 -Alkylen gebunden sind und der andere ein Wasserstoffatom bedeutet, oder ein Säureadditionssalz davon hergestellt wird.

9. Verfahren nach Anspruch 5, dadurch **gekennzeichnet**, daß eine Verbindung, die eine Verbindung der Formel:



ist, worin

R^{13} Methyl, Ethyl, Propyl, Butyl, Pentyl oder Hydroxyethyl bedeutet,

R^{33} ein Wasserstoffatom oder Fluoratom bedeutet,




oder ein Säureadditionssalz davon hergestellt wird.

10. Verfahren nach Anspruch 1, dadurch **gekennzeichnet**, daß 2-(4-Ethyl-1-piperazinyl)-4-(4-fluorphenyl)-5,6,7,8,9,10-hexahydrocycloocta[b]pyridin oder ein Säureadditionssalz davon hergestellt wird.
11. Verfahren nach Anspruch 1, dadurch **gekennzeichnet**, daß 2-(4-Ethyl-1-piperazinyl)-4-(2,4-difluorphenyl)-5,6,7,8,9,10-hexahydrocycloocta[b]pyridin oder ein Säureadditionssalz davon hergestellt wird.
12. Verfahren nach Anspruch 1, dadurch **gekennzeichnet**, daß 2-(4-n-Propyl-1-piperazinyl)-4-(4-fluorphenyl)-5,6,7,8,9,10-hexahydrocycloocta[b]pyridin oder ein Säureadditionssalz davon hergestellt wird.
13. Verfahren nach Anspruch 1, dadurch **gekennzeichnet**, daß 2-(4-Methyl-1-piperazinyl)-4-(4-fluorphenyl)-5,6,7,8,9,10-hexahydrocycloocta[b]pyridin oder ein Säureadditionssalz davon hergestellt wird.
14. Verfahren nach Anspruch 1, dadurch **gekennzeichnet**, daß 2-(4-Methyl-1-piperazinyl)-4-(2,4-difluorphenyl)-5,6,7,8,9,10-hexahydrocycloocta[b]pyridin oder ein Säureadditionssalz davon hergestellt wird.
15. Verfahren zur Herstellung einer psychotropischen Zusammensetzung, dadurch **gekennzeichnet**, daß als aktiver Bestandteil eine wirksame Menge der Verbindung der Formel (I), wie in Anspruch 1 angegeben, oder ein pharmazeutisch annehmbares Salz davon und ein pharmazeutisch annehmbarer Träger oder ein Verdünnungsmittel vermischt werden.
16. Zusammensetzung nach Anspruch 15, dadurch **gekennzeichnet**, daß der aktive Bestandteil die Verbindung, die gemäß Anspruch 5 hergestellt worden ist, oder ein pharmazeutisch annehmbares Salz davon ist.
17. Zusammensetzung nach Anspruch 15, dadurch **gekennzeichnet**, daß der aktive Bestandteil die Verbindung, die gemäß Anspruch 6 hergestellt worden ist, oder ein pharmazeutisch annehmbares Salz davon ist.
18. Zusammensetzung nach Anspruch 15, dadurch **gekennzeichnet**, daß der aktive Bestandteil die Verbindung, die gemäß Anspruch 9 hergestellt worden ist, oder ein pharmazeutisch annehmbares Salz davon ist.

Patentansprüche für folgenden Vertragsstaat : GR

1. Verbindung der Formel (I):



worin

n 3, 4, 5, 6 oder 7 bedeutet,

R¹ ein Wasserstoffatom, C₁-C₁₀-Alkyl, C₂-C₆-Alkenyl, C₂-C₆-Alkynyl, C₃-C₈-Cycloalkyl, C₃-C₆-Cycloalkyl(C₁-C₄)alkyl, Hydroxy(C₂-C₆)alkyl, C₁-C₃-Alkoxy(C₂-C₆)alkyl, Acyloxy(C₂-C₆)alkyl, unsubstituiertes oder substituiertes Aroyl(C₁-C₆)alkyl, unsubstituiertes oder substituiertes Aryl, Heteroaryl oder



Acyl bedeutet,

R^2 und R^3 gleich oder unterschiedlich sind und je ein Wasserstoffatom, ein Halogenatom, C_1 - C_6 -Alkyl, C_1 - C_6 -Alkoxy, Trifluormethyl oder Hydroxy bedeuten,

5 R^4 , R^5 und R^6 gleich oder unterschiedlich sind und je ein Wasserstoffatom, C_1 - C_6 -Alkyl oder Phenyl bedeuten, oder worin zwei von R^4 , R^5 und R^6 unter Bildung einer Einfachbindung oder C_1 - C_3 -Alkylen gebunden sind,

R^7 und R^8 gleich oder unterschiedlich sind und je ein Wasserstoffatom oder C_1 - C_3 -Alkyl bedeuten, m 2 oder 3 bedeutet,

oder ein Säureadditionssalz davon.

10

2. Verbindung nach Anspruch 1, dadurch **gekennzeichnet**, daß R^1 ein Wasserstoffatom, C_1 - C_{10} -Alkyl, C_3 - C_8 -Cycloalkyl, Hydroxy(C_2 - C_6)alkyl, C_1 - C_3 -Alkoxy(C_2 - C_4)alkyl, C_2 - C_4 -Alkanoyloxy(C_2 - C_6)alkyl, eine Benzoyl(C_2 - C_5)alkylgruppe, worin die Phenylgruppierung gegebenenfalls durch Halogen, C_1 - C_3 -Alkyl oder C_1 - C_3 -Alkoxy substituiert sein kann, C_2 - C_4 -Alkenyl, C_2 - C_4 -Alkynyl, Pyridyl, Pyrimidyl, C_2 - C_5 -Alkanoyl oder Furoyl bedeutet,

15

R^2 und R^3 gleich oder unterschiedlich sind und je ein Wasserstoffatom, ein Halogenatom, Methyl oder Methoxy bedeuten,

R^4 , R^5 und R^6 gleich oder unterschiedlich sind und je ein Wasserstoffatom oder C_1 - C_4 -Alkyl bedeuten oder worin zwei oder mehrere von diesen unter Bildung von C_1 - C_2 -Alkylen gebunden sind,

20

R^7 und R^8 gleich oder unterschiedlich sind und je ein Wasserstoffatom oder C_1 - C_3 -Alkyl bedeuten, m 2 bedeutet,

oder ein Säureadditionssalz davon.

25

3. Verbindung nach Anspruch 2, dadurch **gekennzeichnet**, daß R^1 ein Wasserstoffatom, C_1 - C_8 -Alkyl, C_3 - C_8 -Cycloalkyl, Hydroxy(C_2 - C_6)alkyl, C_1 - C_2 -Alkoxy(C_2 - C_3)alkyl, Acetyloxy(C_2 - C_4)alkyl, C_3 - C_4 -Alkenyl oder C_2 - C_3 -Alkanoyl bedeutet,

R^2 und R^3 gleich oder unterschiedlich sind und je ein Wasserstoffatom oder Halogenatom bedeuten oder worin einer von ihnen ein Wasserstoffatom und der andere ein Halogenatom, Methyl oder Methoxy bedeutet,

30

oder ein Säureadditionssalz davon.

4. Verbindung nach Anspruch 3, dadurch **gekennzeichnet**, daß R^2 und R^3 gleich oder unterschiedlich sind und je ein Wasserstoffatom oder ein Fluoratom bedeuten, oder ein Säureadditionssalz davon.


35

5. Verbindung der Formel:

40

45

50



worin

n_1 3, 6 oder 7 bedeutet,

55 R^{11} ein Wasserstoffatom, C_1 - C_8 -Alkyl, C_3 - C_6 -Cycloalkyl, Hydroxy(C_2 - C_4)alkyl, C_1 - C_2 -Alkoxy(C_2 - C_3)alkyl oder C_3 -Alkenyl bedeutet,


R^{21} und R^{31} gleich oder unterschiedlich sind und je ein Wasserstoffatom oder ein Fluoratom bedeuten,

R^{71} und R^{81} gleich oder unterschiedlich sind und je ein Wasserstoffatom oder C_1 - C_3 -Alkyl



bedeuten,
oder ein Säureadditionssalz davon.

6. Verbindung nach Anspruch 1, dadurch **gekennzeichnet**, daß sie eine Verbindung der Formel:



ist, worin

n_2 4 oder 5 bedeutet,

R^{12} ein Wasserstoffatom, C_1 - C_8 -Alkyl, C_3 - C_8 -Cycloalkyl, Hydroxy(C_2 - C_4)alkyl, C_1 - C_2 -Alkoxy(C_2 - C_3)alkyl oder C_3 -Alkenyl bedeutet,

R^{22} und R^{32} gleich oder unterschiedlich sind und je ein Wasserstoffatom oder ein Fluoratom bedeuten,

R^{42} , R^{52} und R^{62} gleich oder unterschiedlich sind und je ein Wasserstoffatom oder C_1 - C_4 -Alkyl bedeuten oder worin zwei davon unter Bildung von C_1 - C_2 -Alkylen gebunden sind, oder ein Säureadditionssalz davon.

7. Verbindung nach Anspruch 5, dadurch **gekennzeichnet**, daß n_1 6 bedeutet,

R^{11} C_1 - C_6 -Alkyl, C_3 - C_6 -Cycloalkyl oder Hydroxy(C_2 - C_4)alkyl bedeutet,

R^{21} und R^{31} gleich oder unterschiedlich sind und je ein Wasserstoffatom oder Fluoratom, substituiert in der 2- oder 4-Stellung des Phenylrings, bedeuten,

R^{71} und R^{81} je ein Wasserstoffatom bedeuten, oder ein Säureadditionssalz davon.

8. Verbindung nach Anspruch 6, dadurch **gekennzeichnet**, daß n_2 5 bedeutet,


R^{12} C_1 - C_6 -Alkyl, C_3 - C_6 -Cycloalkyl oder Hydroxy(C_2 - C_4)alkyl bedeutet,

R^{22} und R^{32} gleich oder unterschiedlich sind und je ein Wasserstoffatom oder Fluoratom, substituiert in der 2- oder 4-Stellung des Phenylrings, bedeuten,

R^{42} , R^{52} und R^{62} alle ein Wasserstoffatom bedeuten oder worin zwei davon unter Bildung von C_1 - C_2 -Alkylen gebunden sind und der andere ein Wasserstoffatom bedeutet, oder ein Säureadditionssalz davon.



9. Verbindung nach Anspruch 5, dadurch gekennzeichnet, daß sie eine Verbindung der Formel:



ist, worin

R^{13} Methyl, Ethyl, Propyl, Butyl, Pentyl oder Hydroxyethyl bedeutet,

R^{33} ein Wasserstoffatom oder Fluoratom bedeutet,
oder ein Säureadditionssalz davon.

10. 2-(4-Ethyl-1-piperazinyl)-4-(4-fluorphenyl)-5,6,7,8,9,10-hexahydrocycloocta[b]pyridin oder ein Säureadditionssalz davon.


11. 2-(4-Ethyl-1-piperazinyl)-4-(2,4-difluorphenyl)-5,6,7,8,9,10-hexahydrocycloocta[b]pyridin oder ein Säureadditionssalz davon.

12. 2-(4-n-Propyl-1-piperazinyl)-4-(4-fluorphenyl)-5,6,7,8,9,10-hexahydrocycloocta[b]pyridin oder ein Säureadditionssalz davon.

13. 2-(4-Methyl-1-piperazinyl)-4-(4-fluorphenyl)-5,6,7,8,9,10-hexahydrocycloocta[b]pyridin oder ein Säureadditionssalz davon.

14. 2-(4-Methyl-1-piperazinyl)-4-(2,4-difluorphenyl)-5,6,7,8,9,10-hexahydrocycloocta[b]pyridin oder ein Säureadditionssalz davon.

15. Verfahren zur Herstellung einer Verbindung der Formel (I):



(I)

worin

n 3, 4, 5, 6 oder 7 bedeutet,

R^1 ein Wasserstoffatom, C_1 - C_{10} -Alkyl, C_2 - C_6 -Alkenyl, C_2 - C_6 -Alkynyl, C_3 - C_8 -Cycloalkyl, C_3 - C_6 -Cycloalkyl(C_1 - C_4)alkyl, Hydroxy(C_2 - C_6)alkyl, C_1 - C_3 -Alkoxy(C_2 - C_6)alkyl, Acyloxy(C_2 - C_6)alkyl, unsubstituiertes oder substituiertes Aroyl(C_1 - C_6)alkyl, unsubstituiertes oder substituiertes Aryl, Heteroaryl oder



Acyl bedeutet,


R^2 und R^3 gleich oder unterschiedlich sind und je ein Wasserstoffatom, ein Halogenatom, C_1 - C_6 -Alkyl, C_1 - C_6 -Alkoxy, Trifluormethyl oder Hydroxy bedeuten,

R^4 , R^5 und R^6 gleich oder unterschiedlich sind und je ein Wasserstoffatom, C_1 - C_6 -Alkyl oder Phenyl bedeuten oder worin zwei von R^4 , R^5 und R^6 unter Bildung einer Einfachbindung oder C_1 - C_3 -Alkylen gebunden sind,


R^7 und R^8 gleich oder unterschiedlich sind und je ein Wasserstoffatom oder C_1 - C_3 -Alkyl bedeuten, m 2 oder 3 bedeutet,

oder eines Säureadditionssalzes davon, dadurch gekennzeichnet, daß

(a) eine Verbindung der Formel (II):




worin X ein Austrittsatom oder eine Austrittsgruppe bedeutet und R^2 , R^3 , R^4 , R^5 , R^6 und n die oben gegebenen Definitionen besitzen, mit einer Verbindung der Formel (III):



worin R^1 , R^7 , R^8 und m die oben gegebenen Definitionen besitzen, umgesetzt wird,


(b) im Falle, wenn R^1 ein Wasserstoffatom bedeutet, eine Verbindung der Formel (I-4):




worin R^{14} unsubstituiertes oder substituiertes Benzyl oder Benzyloxycarbonyl bedeutet (wobei der Substituent an dem Phenylring von Benzyl und Benzyloxycarbonyl ein Glied, ausgewählt aus Niedrigalkyl, Niedrigalkoxy und einem Halogenatom, ist) und R^2 , R^3 , R^4 , R^5 , R^6 , R^7 , m und n die oben gegebenen Definitionen besitzen, der Hydrogenolyse unterworfen wird,




(c) im Falle, wenn R^1 ein Wasserstoffatom bedeutet, eine Verbindung der Formel (I-5):



worin R^{15} Methyl oder unsubstituiertes oder substituiertes Benzyl bedeutet (wobei der Substituent an dem Phenylring des Benzyls ein Glied, ausgewählt aus Niedrigalkyl, Niedrigalkoxy und einem Halogenatom, ist) und R^2 , R^3 , R^4 , R^5 , R^6 , R^7 , R^8 , m und n die oben gegebenen Definitionen besitzen, mit Ethylchlorcarbonat oder 1-Chlorethylchlorcarbonat unter Bildung einer Verbindung der Formel (I-6):




worin R^{16} Ethoxycarbonyl oder 1-Chlorethoxycarbonyl bedeutet und R^2 , R^3 , R^4 , R^5 , R^6 , R^7 , R^8 , m und n die oben gegebenen Definitionen besitzen, oder einer Verbindung der Formel (I-7):



worin R^{17} Acyl bedeutet, R^2 , R^3 , R^4 , R^5 , R^6 , R^7 , R^8 , m und n die oben gegebenen Definitionen besitzen, umgesetzt wird, gefolgt von einer Hydrolyse der Verbindung der Formel (I-6) oder (I-7), oder



(d) in Falle, wenn R^1 eine andere Bedeutung als ein Wasserstoffatom besitzt, eine Verbindung der Formel (I-8):



worin R^2 , R^3 , R^4 , R^5 , R^6 , R^7 , R^8 , m und n die oben gegebenen Definitionen besitzen, mit einer Verbindung der Formel (IV):

$R^{18}-Z$ (IV)

worin Z den reaktiven Rest eines Alkohols bedeutet und R^{18} die gleiche Bedeutung wie R^1 , ausgenommen ein Wasserstoffatom, besitzt, umgesetzt wird, und gegebenenfalls die Form der Verbindung der Formel (I) in ein Säureadditionssalz davon umgewandelt wird.


16. Zusammensetzung, dadurch gekennzeichnet, daß sie als aktiven Bestandteil eine wirksame Menge der Verbindung der Formel (I), wie in Anspruch 1 angegeben, oder ein Salz davon im Gemisch mit einem Träger oder Verdünnungsmittel enthält.
17. Zusammensetzung nach Anspruch 16, dadurch gekennzeichnet, daß der aktive Bestandteil die Verbindung, wie sie in Anspruch 5 angegeben ist, oder ein Salz davon ist.
18. Zusammensetzung nach Anspruch 16, dadurch gekennzeichnet, daß der aktive Bestandteil die Verbindung, wie sie in Anspruch 6 angegeben ist, oder ein Salz davon ist.
19. Zusammensetzung nach Anspruch 16, dadurch gekennzeichnet, daß der aktive Bestandteil die Verbindung, wie sie in Anspruch 9 angegeben ist, oder ein Salz davon ist.



Revendications

Revendications pour les Etats contractants suivants : AT, BE, CH, DE, DK, FR, GB, IT, LI, LU, NL, SE

1. Un composé de formule (I) :



dans laquelle n est égal à 3, 4, 5, 6 ou 7,

R¹ est un atome d'hydrogène, un groupe C₁-C₁₀ alkyle, C₂-C₆ alcényle, C₂-C₆ alcynyle, C₃-C₈ cycloalkyle, C₃-C₆ cycloalkyl-(C₁-C₄)alkyle, hydroxy-(C₂-C₆)alkyle, C₁-C₃ alcoxy-(C₂-C₆)alkyle, acyloxy-(C₂-C₆)alkyle, aroyl-(C₁-C₆)alkyle substitué ou non substitué, aryle substitué ou non substitué, hétéroaryle ou acyle,

R² et R³ sont identiques ou différents et désignent chacun un atome d'hydrogène, un atome d'halogène, un groupe C₁-C₆ alkyle, C₁-C₆ alcoxy, trifluorométhyle ou hydroxy,

R⁴, R⁵ et R⁶ sont identiques ou différents et désignent chacun un atome d'hydrogène, un groupe C₁-C₆ alkyle ou phényle, ou deux des R⁴, R⁵ et R⁶ sont combinés pour former une liaison simple ou un groupe C₁-C₃ alkylène,

R⁷ et R⁸ sont identiques ou différents et désignent chacun un atome d'hydrogène ou un groupe C₁-C₃ alkyle,

m est égal à 2 ou 3,

ou un sel d'addition d'acide dudit composé.

2. Un composé selon la revendication 1, selon laquelle R¹ est un atome d'hydrogène, un groupe C₁-C₁₀ alkyle, C₃-C₈ cycloalkyle, hydroxy-(C₂-C₆)alkyle, C₁-C₃ alcoxy-(C₂-C₄)alkyle, C₂-C₄ alcanoyloxy-(C₂-C₆)alkyle, un groupe benzoyl-(C₂-C₅)alkyle dans lequel la portion phényle peut éventuellement être substituée par un halogène, un groupe C₁-C₃ alkyle ou C₁-C₃ alcoxy, C₂-C₄ alcényle, C₂-C₄ alcynyle, pyridyle, pyrimidyle, C₂-C₅ alcanoyloxy ou furoyle,

R² et R³ sont identiques ou différents et désignent chacun un atome d'hydrogène, un atome d'halogène, un groupe méthyle ou méthoxy,

R⁴, R⁵ et R⁶ sont identiques ou différents et désignent chacun un atome d'hydrogène, un groupe C₁-C₄ alkyle, ou bien deux d'entre eux sont combinés pour former un groupe C₁-C₂ alkylène,

R⁷ et R⁸ sont identiques ou différents et désignent chacun un atome d'hydrogène ou un groupe C₁-C₃ alkyle,

m est égal à 2,

ou bien un sel d'addition d'acide dudit composé.

3. Un composé selon la revendication 2, selon laquelle R¹ est un atome d'hydrogène, un groupe C₁-C₈ alkyle, C₃-C₈ cycloalkyle, hydroxy-(C₂-C₆)alkyle, C₁-C₂ alcoxy-(C₂-C₃)alkyle, acétyloxy-(C₂-C₄)alkyle, C₃-C₄ alcényle ou C₂-C₃ alcanoyloxy,


R² et R³ sont identiques ou différents et désignent chacun un atome d'hydrogène ou un atome d'halogène, ou l'un des deux est un atome d'hydrogène et l'autre est un atome d'halogène, un groupe méthyle ou méthoxy, ou un sel d'addition d'acide dudit composé.

4. Un composé selon la revendication 3, selon laquelle R² et R³ sont identiques ou différents et désignent chacun un atome d'hydrogène ou un atome de fluor,



ou un sel d'addition d'acide dudit composé.

5. Un composé de formule :




dans laquelle n_1 est égal à 3, 6 ou 7,

R^{11} est un atome d'hydrogène, un groupe C_1-C_8 alkyle, C_3-C_6 cycloalkyle, hydroxy- (C_2-C_4) alkyle, C_1-C_2 alcoxy- (C_2-C_3) alkyle ou C_3 alcényle,

R^{21} et R^{31} sont identiques ou différents et désignent chacun un atome d'hydrogène ou un atome de fluor,

R^{71} et R^{81} sont identiques ou différents et désignent chacun un atome d'hydrogène ou un groupe C_1-C_3 alkyle, ou un sel d'addition d'acide dudit composé.

6. Un composé selon la revendication 1 qui est un composé de formule :



dans laquelle n_2 est égal à 4 ou 5,

R^{12} est un atome d'hydrogène, un groupe C_1-C_8 alkyle, C_3-C_8 cycloalkyle, hydroxy- (C_2-C_4) alkyle, C_1-C_2 -alcoxy- (C_2-C_3) alkyle ou C_3 alcényle,

R^{22} et R^{32} sont identiques ou différents et désignent chacun un atome d'hydrogène ou un atome de fluor,

R^{42} , R^{52} et R^{62} sont identiques ou différents et désignent chacun un atome d'hydrogène ou un groupe C_1-C_4 alkyle, ou deux d'entre eux sont combinés pour former un groupe C_1-C_2 alkylène, ou un sel d'addition d'acide dudit composé.

7. Un composé selon la revendication 5, selon laquelle n_1 est égal à 6,

R^{11} est un groupe C_1-C_6 alkyle, C_3-C_6 cycloalkyle ou hydroxy- (C_2-C_4) alkyle,

R^{21} et R^{31} sont identiques ou différents et désignent chacun un atome d'hydrogène ou un atome de fluor substitué dans la position 2- ou 4- du noyau phényle,


R^{71} et R^{81} désignent chacun un atome d'hydrogène,



ou un sel d'addition d'acide dudit composé.

8. Un composé selon la revendication 6, selon laquelle n_2 est égal à 5,
 R^{12} est un groupe C_1-C_6 alkyle, C_3-C_6 cycloalkyle ou hydroxy- (C_2-C_4) alkyle,
 R^{22} et R^{32} sont identiques ou différents et désignent chacun un atome d'hydrogène ou un atome de fluor substitué dans la position 2- ou 4- du noyau phényle,
 R^{42} , R^{52} et R^{62} sont tous des atomes d'hydrogène, ou deux d'entre eux sont combinés pour former un groupe C_1-C_2 alkylène et l'autre est un atome d'hydrogène,
 ou un sel d'addition d'acide dudit composé.

9. Un composé selon la revendication 5 qui est un composé de formule :




dans laquelle R^{13} est un groupe méthyle, éthyle, propyle, butyle, pentyle ou hydroxyéthyle,
 R^{33} est un atome d'hydrogène ou un atome de fluor,
 ou un sel d'addition d'acide dudit composé.

10. 2-(4-éthyl-1-pipérazinyl)-4-(4-fluorophényl)-5,6,7,8,9,10-hexahydrocycloocta[b]pyridine ou un sel d'addition d'acide de celle-ci.
11. 2-(4-éthyl-1-pipérazinyl)-4-(2,4-difluorophényl)-5,6,7,8,9,10-hexahydrocycloocta[b]pyridine, ou un sel d'addition d'acide de celle-ci.
12. 2-(4-n-propyl-1-pipérazinyl)-4-(4-fluorophényl)-5,6,7,8,9,10-hexahydrocycloocta[b]pyridine, ou un sel d'addition d'acide de celle-ci.
13. 2-(4-méthyl-1-pipérazinyl)-4-(4-fluorophényl)-5,6,7,8,9,10-hexahydrocycloocta[b]pyridine, ou un sel d'addition d'acide de celle-ci.
14. 2-(4-méthyl-1-pipérazinyl)-4-(2,4-difluorophényl)-5,6,7,8,9,10-hexahydrocycloocta[b]pyridine, ou un sel d'addition d'acide de celle-ci.



15. Un procédé de préparation d'un composé de formule (I) :



dans laquelle n est égal à 3, 4, 5, 6 ou 7,

R¹ est un atome d'hydrogène, un groupe C₁-C₁₀ alkyle, C₂-C₆ alcényle, C₂-C₆ alcynyle, C₃-C₈ cycloalkyle, C₃-C₆ cycloalkyl-(C₁-C₄)alkyle, hydroxy-(C₂-C₆)alkyle, C₁-C₃ alcoxy-(C₂-C₆)alkyle, acyloxy-(C₂-C₆)alkyle, aroyl-(C₁-C₆)alkyle substitué ou non substitué, aryle substitué ou non substitué, hétéroaryle ou acyle,

R² et R³ sont identiques ou différents et désignent chacun un atome d'hydrogène, un atome d'halogène, un groupe C₁-C₆ alkyle, C₁-C₆ alcoxy, trifluorométhyle ou hydroxy,


R⁴, R⁵ et R⁶ sont identiques ou différents et désignent chacun un atome d'hydrogène, un groupe C₁-C₆ alkyle, ou phényle ou bien deux des R⁴, R⁵ et R⁶ sont combinés pour former une liaison simple ou un groupe C₁-C₆ alkylène,

R⁷ et R⁸ sont identiques ou différents et désignent chacun un atome d'hydrogène ou un groupe C₁-C₃ alkyle,


m est égal à 2 ou 3,

ou un sel d'addition d'acide dudit composé qui comprend

(a) la réaction d'un composé de formule (II) :




dans laquelle X est un atome ou groupe séparable et R², R³, R⁴, R⁵, R⁶ et n sont comme définis précédemment, avec un composé de formule (III) :



dans laquelle R¹, R⁷, R⁸ et m sont comme définis précédemment,




(b) dans le cas où R^1 est un atome d'hydrogène, le traitement par hydrogénolyse d'un composé de formule (I-4) :




dans laquelle R^{14} est un groupe benzyle ou benzyloxycarbonyle substitué ou non substitué (le substituant sur le noyau phényle dudit groupe benzyle ou benzyloxycarbonyle est un élément choisi parmi les groupes alkyles inférieurs, alcoxy inférieurs et un atome d'halogène), et R^2 , R^3 , R^4 , R^5 , R^6 , R^7 , m et n sont comme définis précédemment,


(c) dans le cas où R^1 est un atome d'hydrogène, la réaction d'un composé de formule (I-5) :



dans laquelle R^{15} est un groupe méthyle ou un groupe benzyle substitué ou non substitué (le substituant sur le noyau phényle du groupe benzyle est un élément choisi parmi les groupes alkyles inférieurs, alcoxy inférieurs et un atome d'halogène), et R^2 , R^3 , R^4 , R^5 , R^6 , R^7 , R^8 , m et n sont comme définis précédemment, avec le chlorocarbonate d'éthyle ou le chlorocarbonate de 1-chloroéthyle pour donner un composé de formule (I-6) :




dans laquelle R^{16} est un groupe éthoxycarbonyl ou 1-chloroéthoxycarbonyl, et R^2 , R^3 , R^4 , R^5 , R^6 , R^7 , R^8 , m et n sont comme définis ci-dessus, ou un composé de formule (I-7) :



dans laquelle R^{17} est un groupe acyle, R^2 , R^3 , R^4 , R^5 , R^6 , R^7 , R^8 , m et n sont comme définis ci-dessus, suivie de l'hydrolyse du composé de formule (I-6) ou (I-7), ou bien

(d) dans le cas où R^1 est autre qu'un atome d'hydrogène, la réaction d'un composé de formule (I-8) :



dans laquelle R^2 , R^3 , R^4 , R^5 , R^6 , R^7 , R^8 , m et n sont comme définis ci-dessus, avec un composé de formule (IV) :

R^{18} -Z (IV)




dans laquelle Z est un résidu réactif d'un alcool, et R¹⁸ est identique à R¹, sauf pour l'atome d'hydrogène,
et la conversion éventuelle de la forme du composé de formule (I) en un sel d'addition d'acide.

16. Une composition psychotrope comprenant comme ingrédient actif une quantité efficace du composé de formule (I) mentionné dans la revendication 1, ou d'un sel pharmaceutiquement acceptable de ce composé, en mélange avec un support ou diluant pharmaceutiquement acceptable.
17. Une composition selon la revendication 16, selon laquelle l'ingrédient actif est le composé mentionné dans la revendication 5, ou un sel pharmaceutiquement acceptable de ce composé.
18. Une composition selon la revendication 16, selon laquelle l'ingrédient actif est le composé mentionné dans la revendication 6, ou un sel pharmaceutiquement acceptable de ce composé.
19. Une composition selon la revendication 16, selon laquelle l'ingrédient actif est le composé mentionné dans la revendication 9, ou un sel pharmaceutiquement acceptable de ce composé.
20. Un composé de formule (I) mentionné dans la revendication 16 pour l'utilisation comme préparation pharmaceutique pour les maladies psychotiques.
21. Utilisation d'un composé de formule (I) mentionné dans la revendication 16 pour la préparation d'une préparation pharmaceutique utile pour le traitement des maladies psychotiques.

Revendications pour l'Etat contractant suivant : ES

1. Un procédé de préparation d'un composé de formule (I) :



dans laquelle n est égal à 3, 4, 5, 6 ou 7,

R¹ est un atome d'hydrogène, un groupe C₁-C₁₀ alkyle, C₂-C₆ alcényle, C₂-C₆ alcynyle, C₃-C₈ cycloalkyle, C₃-C₆ cycloalkyl-(C₁-C₄)alkyle, hydroxy-(C₂-C₆)alkyle, C₁-C₃ alcoxy-(C₂-C₆)alkyle, acyloxy-(C₂-C₆)alkyle, aroyl-(C₁-C₆)alkyle substitué ou non substitué, aryle substitué ou non substitué, hétéroaryle ou acyle,

R² et R³ sont identiques ou différents et désignent chacun un atome d'hydrogène, un atome d'halogène, un groupe C₁-C₆ alkyle, C₁-C₆ alcoxy, trifluorométhyle ou hydroxy,

R⁴, R⁵ et R⁶ sont identiques ou différents et désignent chacun un atome d'hydrogène, un groupe C₁-C₆ alkyle, ou phényle ou bien deux des R⁴, R⁵ et R⁶ sont combinés pour former une liaison simple ou un groupe C₁-C₆ alkylène,


R⁷ et R⁸ sont identiques ou différents et désignent chacun un atome d'hydrogène ou un groupe C₁-C₃ alkyle,

m est égal à 2 ou 3,


ou un sel d'addition d'acide dudit composé qui comprend



(a) la réaction d'un composé de formule (II) :




dans laquelle X est un atome ou groupe séparable et R^2 , R^3 , R^4 , R^5 , R^6 et n sont comme définis précédemment, avec un composé de formule (III) :



dans laquelle R^1 , R^7 , R^8 et m sont comme définis précédemment,


(b) dans le cas où R^1 est un atome d'hydrogène, le traitement par hydrogénolyse d'un composé de formule (I-4) :




dans laquelle R^{14} est un groupe benzyle ou benzyloxycarbonyle substitué ou non substitué (le substituant sur le noyau phényle dudit groupe benzyle ou benzyloxycarbonyle est un élément choisi parmi les groupes alkyles inférieurs, alcoxy inférieurs et un atome d'halogène), et R^2 , R^3 , R^4 , R^5 , R^6 , R^7 , m et n sont comme définis précédemment,




(c) dans le cas où R^1 est un atome d'hydrogène, la réaction d'un composé de formule (I-5) :



dans laquelle R^{15} est un groupe méthyle ou un groupe benzyle substitué ou non substitué (le substituant sur le noyau phényle du groupe benzyle est un élément choisi parmi les groupes alkyles inférieurs, alcoxy inférieurs et un atome d'halogène), et R^2 , R^3 , R^4 , R^5 , R^6 , R^7 , R^8 , m et n sont comme définis précédemment, avec le chlorocarbonate d'éthyle ou le chlorocarbonate de 1-chloroéthyle pour donner un composé de formule (I-6) :




dans laquelle R^{16} est un groupe éthoxycarbonyl ou 1-chloroéthoxycarbonyl, et R^2 , R^3 , R^4 , R^5 , R^6 , R^7 , R^8 , m et n sont comme définis ci-dessus, ou un composé de formule (I-7) :



dans laquelle R^{17} est un groupe acyle, R^2 , R^3 , R^4 , R^5 , R^6 , R^7 , R^8 , m et n sont comme définis ci-dessus, suivie de l'hydrolyse du composé de formule (I-6) ou (I-7), ou bien



(d) dans le cas où R^1 est autre qu'un atome d'hydrogène, la réaction d'un composé de formule (I-8)



dans laquelle R^2 , R^3 , R^4 , R^5 , R^6 , R^7 , R^8 , m et n sont comme définis ci-dessus, avec un composé de formule (IV) :

R^{18} -Z (IV)

dans laquelle Z est un résidu réactif d'un alcool, et R^{18} est identique à R^1 , sauf pour l'atome d'hydrogène, et la conversion éventuelle de la forme du composé de formule (I) en un sel d'addition d'acide.

2. Un procédé selon la revendication 1, caractérisé en ce que l'on prépare un composé dans lequel R^1 est un atome d'hydrogène, un groupe C_1 - C_{10} alkyle, C_3 - C_8 cycloalkyle, hydroxy- $(C_2$ - $C_6)$ alkyle, C_1 - C_3 alcoxy- $(C_2$ - $C_4)$ alkyle, C_2 - C_4 alcanoyloxy- $(C_2$ - $C_6)$ alkyle, un groupe benzoyl- $(C_2$ - $C_5)$ alkyle dans lequel la portion phényle peut éventuellement être substituée par un halogène, un groupe C_1 - C_3 alkyle ou C_1 - C_3 alcoxy, C_2 - C_4 alcényle, C_2 - C_4 alcynyle, pyridyle, pyrimidyle, C_2 - C_5 alcanoyloxy ou furoyle,

R^2 et R^3 sont identiques ou différents et désignent chacun un atome d'hydrogène, un atome d'halogène, un groupe méthyle ou méthoxy,

R^4 , R^5 et R^6 sont identiques ou différents et désignent chacun un atome d'hydrogène, un groupe C_1 - C_4 alkyle, ou bien deux d'entre eux sont combinés pour former un groupe C_1 - C_2 alkylène,

R^7 et R^8 sont identiques ou différents et désignent chacun un atome d'hydrogène ou un groupe C_1 - C_3 alkyle,

m est égal à 2,

ou bien un sel d'addition d'acide dudit composé.


3. Un procédé selon la revendication 2, caractérisé en ce que l'on prépare un composé dans lequel R^1 est un atome d'hydrogène, un groupe C_1 - C_8 alkyle, C_3 - C_8 cycloalkyle, hydroxy- $(C_2$ - $C_6)$ alkyle, C_1 - C_2 alcoxy- $(C_2$ - $C_3)$ alkyle, acétyloxy- $(C_2$ - $C_4)$ alkyle, C_3 - C_4 alcényle ou C_2 - C_3 alcanoyloxy,

R^2 et R^3 sont identiques ou différents et désignent chacun un atome d'hydrogène ou un atome d'halogène, ou l'un des deux est un atome d'hydrogène et l'autre est un atome d'halogène, un groupe méthyle ou méthoxy, ou un sel d'addition d'acide dudit composé.

4. Un procédé selon la revendication 3, caractérisé en ce qu'on prépare un composé dans lequel R^2 et R^3 sont identiques ou différents et désignent chacun un atome d'hydrogène ou un atome de fluor, ou un sel d'addition d'acide dudit composé.



5. Un procédé de préparation d'un composé de formule :




dans laquelle n_1 est égal à 3, 6 ou 7,

R^{11} est un atome d'hydrogène, un groupe C_1-C_8 alkyle, C_3-C_6 cycloalkyle, hydroxy- (C_2-C_4) alkyle, C_1-C_2 alcoxy- (C_2-C_3) alkyle ou C_3 alcényle,

R^{21} et R^{31} sont identiques ou différents et désignent chacun un atome d'hydrogène ou un atome de fluor,

R^{71} et R^{81} sont identiques ou différents et désignent chacun un atome d'hydrogène ou un groupe C_1-C_3 alkyle, ou un sel d'addition d'acide dudit composé.

6. Un procédé selon la revendication 1, caractérisé en ce que l'on prépare un composé qui est un composé de formule :



dans laquelle n_2 est égal à 4 ou 5,

R^{12} est un atome d'hydrogène, un groupe C_1-C_8 alkyle, C_3-C_8 cycloalkyle, hydroxy- (C_2-C_4) alkyle, C_1-C_2 alcoxy- (C_2-C_3) alkyle ou C_3 alcényle,

R^{22} et R^{32} sont identiques ou différents et désignent chacun un atome d'hydrogène ou un atome de fluor,

R^{42} , R^{52} et R^{62} sont identiques ou différents et désignent chacun un atome d'hydrogène ou un groupe C_1-C_4 alkyle, ou deux d'entre eux sont combinés pour former un groupe C_1-C_2 alkylène, ou un sel d'addition d'acide dudit composé.

7. Un procédé selon la revendication 5, caractérisé en ce que l'on prépare un composé dans lequel n_1 est égal à 6,

R^{11} est un groupe C_1-C_6 alkyle, C_3-C_6 cycloalkyle ou hydroxy- (C_2-C_4) alkyle,

R^{21} et R^{31} sont identiques ou différents et désignent chacun un atome d'hydrogène ou un atome de fluor substitué dans la position 2- ou 4- du noyau phényle,

R^{71} et R^{81} désignent chacun un atome d'hydrogène, ou un sel d'addition d'acide dudit composé.




8. Un procédé selon la revendication 6, caractérisé en ce que l'on prépare un composé dans lequel n_2 est égal à 5,

R^{12} est un groupe C_1-C_6 alkyle, C_3-C_6 cycloalkyle ou hydroxy- (C_2-C_4) alkyle,

- 5 R^{22} et R^{32} sont identiques ou différents et désignent chacun un atome d'hydrogène ou un atome de fluor substitué dans la position 2- ou 4- du noyau phényle,

R^{42} , R^{52} et R^{62} sont tous des atomes d'hydrogène, ou deux d'entre eux sont combinés pour former un groupe C_1-C_2 alkylène et l'autre est un atome d'hydrogène, ou un sel d'addition d'acide dudit composé.

- 10 9. Un procédé selon la revendication 5, caractérisé en ce que l'on prépare un composé qui est un composé de formule :



dans laquelle R^{13} est un groupe méthyle, éthyle, propyle, butyle, pentyle ou hydroxyéthyle,

R^{33} est un atome d'hydrogène ou un atome de fluor, ou un sel d'addition d'acide dudit composé.

- 30 10. Le procédé selon la revendication 1, caractérisé en ce que la 2-(4-éthyl-1-pipérazinyl)-4-(4-fluorophényl)-5,6,7,8,9,10-hexahydrocycloocta[b]pyridine ou un sel d'addition d'acide est préparé.

11. Le procédé selon la revendication 1, caractérisé en ce que la 2-(4-éthyl-1-pipérazinyl)-4-(2,4-difluorophényl)-5,6,7,8,9,10-hexahydrocycloocta[b]pyridine ou un sel d'addition d'acide est préparé.

- 35 12. Le procédé selon la revendication 1, caractérisé en ce que la 2-(4-n-propyl-1-pipérazinyl)-4-(4-fluorophényl)-5,6,7,8,9,10-hexahydrocycloocta[b]pyridine ou un sel d'addition d'acide est préparé.

- 40 13. Le procédé selon la revendication 1, caractérisé en ce que la 2-(4-méthyl-1-pipérazinyl)-4-(4-fluorophényl)-5,6,7,8,9,10-hexahydrocycloocta[b]pyridine ou un sel d'addition d'acide est préparé.

14. Le procédé selon la revendication 1, caractérisé en ce que la 2-(4-méthyl-1-pipérazinyl)-4-(2,4-difluorophényl)-5,6,7,8,9,10-hexahydrocycloocta[b]pyridine ou un sel d'addition d'acide est préparé.

- 45 15. Un procédé de préparation d'une composition psychotrope comprenant le mélangeage comme ingrédient actif d'une quantité efficace du composé de formule (I) comme mentionné dans la revendication 1, ou d'un sel pharmaceutiquement acceptable dudit composé et d'un support ou diluant pharmaceutiquement acceptable.

- 50 16. Le procédé selon la revendication 15, selon laquelle l'ingrédient actif est le composé préparé dans la revendication 5, ou un sel pharmaceutiquement acceptable dudit composé.


17. Le procédé selon la revendication 15, selon laquelle l'ingrédient actif est le composé préparé dans la revendication 6, ou un sel pharmaceutiquement acceptable dudit composé.

- 55 18. Le procédé selon la revendication 15, selon laquelle l'ingrédient actif est le composé préparé dans la revendication 9, ou un sel pharmaceutiquement acceptable dudit composé.



Revendications pour l'Etat contractant suivant : GR

1. Un composé de formule (I) :



dans laquelle n est égal à 3, 4, 5, 6 ou 7,

R¹ est un atome d'hydrogène, un groupe C₁-C₁₀ alkyle, C₂-C₆ alcényle, C₂-C₆ alcynyle, C₃-C₈ cycloalkyle, C₃-C₆ cycloalkyl-(C₁-C₄)alkyle, hydroxy-(C₂-C₆)alkyle, C₁-C₃ alcoxy-(C₂-C₆)alkyle, acyloxy-(C₂-C₆)alkyle, aroyl-(C₁-C₆)alkyle substitué ou non substitué, aryle substitué ou non substitué, hétéroaryle ou acyle,

R² et R³ sont identiques ou différents et désignent chacun un atome d'hydrogène, un atome d'halogène, un groupe C₁-C₆ alkyle, C₁-C₆ alcoxy, trifluorométhyle ou hydroxy,

R⁴, R⁵ et R⁶ sont identiques ou différents et désignent chacun un atome d'hydrogène, un groupe C₁-C₆ alkyle ou phényle, ou deux des R⁴, R⁵ et R⁶ sont combinés pour former une liaison simple ou un groupe C₁-C₃ alkylène,

R⁷ et R⁸ sont identiques ou différents et désignent chacun un atome d'hydrogène ou un groupe C₁-C₃ alkyle,

m est égal à 2 ou 3,

ou un sel d'addition d'acide dudit composé.

2. Un composé selon la revendication 1, selon laquelle R¹ est un atome d'hydrogène, un groupe C₁-C₁₀ alkyle, C₃-C₈ cycloalkyle, hydroxy-(C₂-C₆)alkyle, C₁-C₃ alcoxy-(C₂-C₄)alkyle, C₂-C₄ alcanoyloxy-(C₂-C₆)alkyle, un groupe benzoyl-(C₂-C₅)alkyle dans lequel la portion phényle peut éventuellement être substituée par un halogène, un groupe C₁-C₃ alkyle ou C₁-C₃ alcoxy, C₂-C₄ alcényle, C₂-C₄ alcynyle, pyridyle, pyrimidyle, C₂-C₅ alcanoyloxy ou furoyle,

R² et R³ sont identiques ou différents et désignent chacun un atome d'hydrogène, un atome d'halogène, un groupe méthyle ou méthoxy,

R⁴, R⁵ et R⁶ sont identiques ou différents et désignent chacun un atome d'hydrogène, un groupe C₁-C₄ alkyle, ou bien deux d'entre eux sont combinés pour former un groupe C₁-C₂ alkylène,

R⁷ et R⁸ sont identiques ou différents et désignent chacun un atome d'hydrogène ou un groupe C₁-C₃ alkyle,

m est égal à 2,

ou bien un sel d'addition d'acide dudit composé.


3. Un composé selon la revendication 2, selon laquelle R¹ est un atome d'hydrogène, un groupe C₁-C₈ alkyle, C₃-C₈ cycloalkyle, hydroxy-(C₂-C₆)alkyle, C₁-C₂ alcoxy-(C₂-C₃)alkyle, acétyloxy-(C₂-C₄)alkyle, C₃-C₄ alcényle ou C₂-C₃ alcanoyloxy,

R² et R³ sont identiques ou différents et désignent chacun un atome d'hydrogène ou un atome d'halogène, ou l'un des deux est un atome d'hydrogène et l'autre est un atome d'halogène, un groupe méthyle ou méthoxy, ou un sel d'addition d'acide dudit composé.

4. Un composé selon la revendication 3, selon laquelle R² et R³ sont identiques ou différents et désignent chacun un atome d'hydrogène ou un atome de fluor, ou un sel d'addition d'acide dudit composé.



5. Un composé de formule :




dans laquelle n_1 est égal à 3, 6 ou 7,

R^{11} est un atome d'hydrogène, un groupe C_1-C_3 alkyle, C_3-C_6 cycloalkyle, hydroxy- (C_2-C_4) alkyle, C_1-C_2 alcoxy- (C_2-C_3) alkyle ou C_3 alcényle,

R^{21} et R^{31} sont identiques ou différents et désignent chacun un atome d'hydrogène ou un atome de fluor,

R^{71} et R^{81} sont identiques ou différents et désignent chacun un atome d'hydrogène ou un groupe C_1-C_3 alkyle, ou un sel d'addition d'acide dudit composé.

6. Un composé selon la revendication 1 qui est un composé de formule :



dans laquelle n_2 est égal à 4 ou 5,

R^{12} est un atome d'hydrogène, un groupe C_1-C_3 alkyle, C_3-C_6 cycloalkyle, hydroxy- (C_2-C_4) alkyle, C_1-C_2 alcoxy- (C_2-C_3) alkyle ou C_3 alcényle,

R^{22} et R^{32} sont identiques ou différents et désignent chacun un atome d'hydrogène ou un atome de fluor,

R^{42} , R^{52} et R^{62} sont identiques ou différents et désignent chacun un atome d'hydrogène ou un groupe C_1-C_4 alkyle, ou deux d'entre eux sont combinés pour former un groupe C_1-C_2 alkylène, ou un sel d'addition d'acide dudit composé.

7. Un composé selon la revendication 5, selon laquelle n_1 est égal à 6,

R^{11} est un groupe C_1-C_6 alkyle, C_3-C_6 cycloalkyle ou hydroxy- (C_2-C_4) alkyle,


R^{21} et R^{31} sont identiques ou différents et désignent chacun un atome d'hydrogène ou un atome de fluor substitué dans la position 2- ou 4- du noyau phényle,

R^{71} et R^{81} désignent chacun un atome d'hydrogène, ou un sel d'addition d'acide dudit composé.



8. Un composé selon la revendication 6, selon laquelle n_2 est égal à 5,
 R^{12} est un groupe C_1-C_6 alkyle, C_3-C_6 cycloalkyle ou hydroxy- (C_2-C_4) alkyle,
 R^{22} et R^{32} sont identiques ou différents et désignent chacun un atome d'hydrogène ou un atome de fluor substitué dans la position 2- ou 4- du noyau phényle,
 R^{42} , R^{52} et R^{62} sont tous des atomes d'hydrogène, ou deux d'entre eux sont combinés pour former un groupe C_1-C_2 alkylène et l'autre est un atome d'hydrogène, ou un sel d'addition d'acide dudit composé.

9. Un composé selon la revendication 5 qui est un composé de formule :



dans laquelle R^{13} est un groupe méthyle, éthyle, propyle, butyle, pentyle ou hydroxyéthyle,
 R^{33} est un atome d'hydrogène ou un atome de fluor, ou un sel d'addition d'acide dudit composé.

10. 2-(4-éthyl-1-pipérazinyl)-4-(4-fluorophényl)-5,6,7,8,9,10-hexahydrocycloocta[b]pyridine, ou un sel d'addition d'acide de celle-ci.


11. 2-(4-éthyl-1-pipérazinyl)-4-(2,4-difluorophényl)-5,6,7,8,9,10-hexahydrocycloocta[b]pyridine, ou un sel d'addition d'acide de celle-ci.

12. 2-(4-n-propyl-1-pipérazinyl)-4-(4-fluorophényl)-5,6,7,8,9,10-hexahydrocycloocta[b]pyridine, ou un sel d'addition d'acide de celle-ci.

13. 2-(4-méthyl-1-pipérazinyl)-4-(4-fluorophényl)-5,6,7,8,9,10-hexahydrocycloocta[b]pyridine, ou un sel d'addition d'acide de celle-ci.

14. 2-(4-méthyl-1-pipérazinyl)-4-(2,4-difluorophényl)-5,6,7,8,9,10-hexahydrocycloocta[b]pyridine, ou un sel d'addition de celle-ci.

15. Un procédé de préparation d'un composé de formule (I) :



(I)



dans laquelle n est égal à 3, 4, 5, 6 ou 7,

R¹ est un atome d'hydrogène, un groupe C₁-C₁₀ alkyle, C₂-C₆ alcényle, C₂-C₆ alcynyle, C₃-C₈ cycloalkyle, C₃-C₆ cycloalkyl-(C₁-C₄)alkyle, hydroxy-(C₂-C₆)alkyle, C₁-C₃ alcoxy-(C₂-C₆)alkyle, acyloxy-(C₂-C₆)alkyle, aroyl-(C₁-C₆)alkyle substitué ou non substitué, aryle substitué ou non substitué, hétéroaryle ou acyle,

R² et R³ sont identiques ou différents et désignent chacun un atome d'hydrogène, un atome d'halogène, un groupe C₁-C₆ alkyle, C₁-C₆ alcoxy, trifluorométhyle ou hydroxy,


R⁴, R⁵ et R⁶ sont identiques ou différents et désignent chacun un atome d'hydrogène, un groupe C₁-C₆ alkyle, ou phényle ou bien deux des R⁴, R⁵ et R⁶ sont combinés pour former une liaison simple ou un groupe C₁-C₆ alkylène,

R⁷ et R⁸ sont identiques ou différents et désignent chacun un atome d'hydrogène ou un groupe C₁-C₃ alkyle,

m est égal à 2 ou 3,


ou un sel d'addition d'acide dudit composé qui comprend

(a) la réaction d'un composé de formule (II) :



(II)


dans laquelle X est un atome ou groupe séparable et R², R³, R⁴, R⁵, R⁶ et n sont comme définis précédemment, avec un composé de formule (III) :



(III)

dans laquelle R¹, R⁷, R⁸ et m sont comme définis précédemment,


(b) dans le cas où R¹ est un atome d'hydrogène, le traitement par hydrogénolyse d'un composé de formule (I-4) :



(I-4)


dans laquelle R^{14} est un groupe benzyle ou benzyloxycarbonyl substitué ou non substitué (le substituant sur le noyau phényle dudit groupe benzyle ou benzyloxycarbonyl est un élément choisi parmi les groupes alkyles inférieurs, alcoxy inférieurs et un atome d'halogène), et R^2 , R^3 , R^4 , R^5 , R^6 , R^7 , m et n sont comme définis précédemment,

(c) dans le cas où R^1 est un atome d'hydrogène, la réaction d'un composé de formule (I-5) :




(I-5)

dans laquelle R^{15} est un groupe méthyle ou un groupe benzyle substitué ou non substitué (le substituant sur le noyau phényle du groupe benzyle est un élément choisi parmi les groupes alkyles inférieurs, alcoxy inférieurs et un atome d'halogène), et R^2 , R^3 , R^4 , R^5 , R^6 , R^7 , R^8 , m et n sont comme définis précédemment, avec le chlorocarbonate d'éthyle ou le chlorocarbonate de 1-chloroéthyle pour donner un composé de formule (I-6) :



(I-6)


dans laquelle R^{16} est un groupe éthoxycarbonyl ou 1-chloroéthoxycarbonyl, et R^2 , R^3 , R^4 , R^5 , R^6 , R^7 , R^8 , m et n sont comme définis ci-dessus, ou un composé de formule (I-7) :



20

25

dans laquelle R^{17} est un groupe acyle, R^2 , R^3 , R^4 , R^5 , R^6 , R^7 , R^8 , m et n sont comme définis ci-dessus, suivie de l'hydrolyse du composé de formule (I-6) ou (I-7), ou bien
(d) dans le cas où R^1 est autre qu'un atome d'hydrogène, la réaction d'un composé de formule (I-8)



dans laquelle R^2 , R^3 , R^4 , R^5 , R^6 , R^7 , R^8 , m et n sont comme définis ci-dessus, avec un composé de formule (IV) :

40

$R^{18}-Z$ (IV)

dans laquelle Z est un résidu réactif d'un alcool, et R^{18} est identique à R^1 , sauf pour l'atome d'hydrogène,
et la conversion éventuelle de la forme du composé de formule (I) en un sel d'addition d'acide.

- 45
16. Une composition comprenant comme ingrédient actif une quantité efficace du composé de formule (I) mentionné dans la revendication 1, ou d'un sel dudit composé en mélange avec un support ou diluant.
17. Une composition selon la revendication 16, selon laquelle l'ingrédient actif est le composé mentionné dans la revendication 5 ou un sel de ce composé.
- 50
18. Une composition selon la revendication 16, selon laquelle l'ingrédient actif est le composé mentionné dans la revendication 6 ou un sel de ce composé.
- 55
19. Une composition selon la revendication 16, selon laquelle l'ingrédient actif est le composé mentionné dans la revendication 9, ou un sel dudit composé.