

Aminochromane, aminothiochromane and amino-1,2,3,4-tetrahydroquinoline derivatives, pharmaceutical compositions containing them, and their use in therapy

Background of the Invention

5

The present invention relates to aminochromane, aminothiochromane and amino-1,2,3,4-tetrahydroquinoline derivatives, pharmaceutical compositions comprising such aminochromane, aminothiochromane and amino-1,2,3,4-tetrahydroquinoline derivatives, and the use of such aminochromane, aminothiochromane and amino-1,2,3,4-tetrahydroquinoline derivatives for therapeutic
10 purposes. The aminochromane, aminothiochromane and amino-1,2,3,4-tetrahydroquinoline derivatives are GlyT1 inhibitors.

Dysfunction of glutamatergic pathways has been implicated in a number of disease states in the human central nervous system (CNS) including but not limited to schizophrenia, cognitive deficits,
15 dementia, Parkinson disease, Alzheimer disease and bipolar disorder. A large number of studies in animal models lend support to the NMDA hypofunction hypothesis of schizophrenia.

NMDA receptor function can be modulated by altering the availability of the co-agonist glycine. This approach has the critical advantage of maintaining activity-dependent activation of the NMDA
20 receptor because an increase in the synaptic concentration of glycine will not produce an activation of NMDA receptors in the absence of glutamate. Since synaptic glutamate levels are tightly maintained by high affinity transport mechanisms, an increased activation of the glycine site will only enhance the NMDA component of activated synapses.


25 Two specific glycine transporters, GlyT1 and GlyT2 have been identified and shown to belong to the Na/Cl-dependent family of neurotransmitter transporters which includes taurine, gamma-aminobutyric acid (GABA), proline, monoamines and orphan transporters. GlyT1 and GlyT2 have been isolated from different species and shown to have only 50% identity at the amino acid level. They also have a different pattern of expression in mammalian central nervous system, with GlyT2
30 being expressed in spinal cord, brainstem and cerebellum and GlyT1 present in these regions as well as forebrain areas such as cortex, hippocampus, septum and thalamus. At the cellular level, GlyT2 has been reported to be expressed by glycinergic nerve endings in rat spinal cord whereas GlyT1 appears to be preferentially expressed by glial cells. These expression studies have led to the suggestion that GlyT2 is predominantly responsible for glycine uptake at glycinergic synapses
35 whereas GlyT1 is involved in monitoring glycine concentration in the vicinity of NMDA receptor

expressing synapses. Recent functional studies in rat have shown that blockade of GlyT1 with the potent inhibitor (N-[3-(4'-fluorophenyl)-3-(4'-phenylphenoxy)propyl])-sarcosine (NFPS) potentiates NMDA receptor activity and NMDA receptor-dependent long-term potentiation in rat.


- 5 Molecular cloning has further revealed the existence of three variants of GlyT1, termed GlyT-1a, GlyT-1b and GlyT-1c, each of which displays a unique distribution in the brain and peripheral tissues. The variants arise by differential splicing and exon usage, and differ in their N-terminal regions.
- 10 The physiological effects of GlyT1 in forebrain regions together with clinical reports showing the beneficial effects of GlyT1 inhibitor sarcosine in improving symptoms in schizophrenia patients suggest that selective GlyT1 inhibitors represent a new class of antipsychotic drugs.

Glycine transporter inhibitors are already known in the art, for example:


15




US 200626364




US2002169197




EP 1 284 257



WO 2003053942




WO 2004096761




WO 2003031435


DE 10315570




WO 2003055478




WO 2004113280




WO 2004112787




WO 2004113301




WO 2005049023




WO 2003089411




WO 2004013100




WO 2004013101




WO 2005037783




WO 2005037792




WO 2005037781




WO 2005037782




WO 2005037785




WO 2005037785




WO 2004072034







WO 2005014563




WO 2005023260




WO 2005023261




WO 2005040166




WO 2005058882




WO 2005058885




WO 2005058317



WO 2005046601




WO 2003087086



WO2003076420


WO2004022528



- (see also Hashimoto K., Recent Patents on CNS Drug Discovery, 2006, 1, 43–53; Harsing L.G. et al., Current Medicinal Chemistry, 2006, 13, 1017–1044; Javitt D.C., Molecular Psychiatry (2004) 9, 984–997; Lindsley, C.W. et al., Current Topics in Medicinal Chemistry, 2006, 6, 771–785;
- 5 Lindsley C.W. et al., Current Topics in Medicinal Chemistry, 2006, 6, 1883–1896).


Further glycine transporter inhibitors are known from the following documents.

WO 2009024611 describes 4-benzylaminoquinolines of formula:




10


WO 2009121872 describes tetrahydroisoquinoline of formula:



- 15 WO 2010092180 describes aminotetraline derivatives of formula:




WO 2010092181 describes heterocyclic compounds of formula:




5

WO 2012020131 describes aminoindane derivatives of formula:




WO 2012020130 describes phenalkylamine derivatives of formula:




10

WO 2012020133 describes tetraline and indane derivatives of formula:





WO 2012152915 describes benzazepine derivatives of formula:




5





WO 2012020134 describes phenalkylamine derivatives of formulae:




10 WO 2013020930 describes aminochromane, aminothiochromane and amino-1,2,3,4-tetrahydroquinoline derivatives of formula:



WO 2013072520 describes N-substituted aminobenzocycloheptene, aminotetraline, aminoindane and phenalkylamine derivatives of formula:




WO 2013120835 describes isoindoline derivatives of formula



It was one object of the present invention to provide further glycine transporter inhibitors. It was a further object of the present invention to provide glycine transporter inhibitors which combine high stability with high affinity. It was a further object of the present invention to provide glycine transporter inhibitors which show favorable efflux properties. It was a further object of the present invention to provide glycine transporter inhibitors which combine high stability and high affinity with favorable efflux properties.

Summary of the Invention

The present invention relates to aminochromane, aminothiochromane and amino-1,2,3,4-tetrahydroquinoline derivatives of the formula (I)



15

wherein

A is a 5- or 6-membered ring;

20 R¹ is hydrogen, alkyl, cycloalkylalkyl, halogenated alkyl, trialkylsilylalkyl, hydroxyalkyl, alkoxyalkyl, aminoalkyl, alkylaminoalkyl, dialkylaminoalkyl, alkylcarbonylaminoalkyl, alkyloxycarbonylaminoalkyl, alkylaminocarbonylaminoalkyl, dialkylaminocarbonylaminoalkyl, alkylsulfonylaminoalkyl, (optionally substituted arylalkyl)aminoalkyl, optionally substituted arylalkyl, optionally substituted heterocyclalkyl, cycloalkyl, alkylcarbonyl, alkoxy carbonyl, halogenated alkoxy carbonyl, aryloxycarbonyl, aminocarbonyl, alkyl-

25

- minocarbonyl, (halogenated alkyl)aminocarbonyl, arylaminocarbonyl, alkenyl, alkynyl, optionally substituted aryl, hydroxy, alkoxy, halogenated alkoxy, hydroxyalkoxy, alkoxyalkoxy, aminoalkoxy, alkylaminoalkoxy, dialkylaminoalkoxy, alkylcarbonylaminoalkoxy, arylcarbonylaminoalkoxy, alkoxycarbonylaminoalkoxy, arylalkoxy, alkylsulfonylaminoalkoxy, (halogenated alkyl)sulfonylaminoalkoxy, arylsulfonylaminoalkoxy, (arylalkyl)sulfonylaminoalkoxy, heterocyclylsulfonylaminoalkoxy, heterocyclylalkoxy, aryloxy, heterocyclyloxy, alkylthio, halogenated alkylthio, alkylamino, (halogenated alkyl)amino, dialkylamino, di-(halogenated alkyl)amino, alkylcarbonylamino, (halogenated alkyl)carbonylamino, arylcarbonylamino, alkylsulfonylamino, (halogenated alkyl)sulfonylamino, arylsulfonylamino or optionally substituted heterocyclyl;
- 5
- 10
- W is $-NR^7-$ or a bond;
- 15
- A¹ is optionally substituted alkylene or a bond;
- Q is $-S(O)_2-$ or $-C(O)-$;
- Y is $-NR^8-$ or a bond;
- 20
- n1 is 0, 1, 2, or 3;
- n2 is 0, 1, 2, or 3;
- 25
- X¹ is $>N-$ or $>CH-$;
- R⁶ is hydrogen, halogen, alkyl, halogenated alkyl, $-CN$, hydroxy, alkoxy or halogenated alkoxy, or two radicals R⁶ together with the carbon atom to which they are attached form a carbonyl group;
- 30
- R² is hydrogen, halogen, alkyl, halogenated alkyl, $-CN$, alkenyl, alkynyl, optionally substituted aryl, hydroxy, alkoxy, halogenated alkoxy, alkoxy carbonyl, alkenyloxy, arylalkoxy, alkylcarbonyloxy, alkylthio, alkylsulfinyl, alkylsulfonyl, aminosulfonyl, amino, alkylamino, alkenylamino, nitro or optionally substituted heterocyclyl, or two radicals R² together with the ring atoms of A to which they are bound form a 5- or 6 membered ring;
- 35
- A² is $-O-$, $-S-$ or $-NR^9-$;
- R³ is hydrogen, halogen, alkyl or alkoxy, or two radicals R³ together with the carbon atom to which they are attached form a carbonyl group;
- 40
- Y¹ is a bond or optionally substituted alkylene;

- 5 R^{4a} is hydrogen, alkyl, cycloalkylalkyl, halogenated alkyl, hydroxyalkyl, alkoxyalkyl, aminoalkyl, $-\text{CH}_2\text{CN}$, arylalkyl, optionally substituted cycloalkyl, $-\text{CHO}$, alkylcarbonyl, (halogenated alkyl)carbonyl, arylcarbonyl, alkoxycarbonyl, aryloxy carbonyl, alkylaminocarbonyl, alkenyl, $-\text{C}(=\text{NH})\text{NH}_2$, $-\text{C}(=\text{NH})\text{NHCN}$, alkylsulfonyl, arylsulfonyl, amino, $-\text{NO}$ or optionally substituted heterocyclyl; or
- R^{4a} is optionally substituted alkylene that is bound to a carbon atom in Y^1 ;
- 10 R^{4b} is hydrogen, alkyl, halogenated alkyl, hydroxyalkyl, alkoxyalkyl, aminoalkyl, $-\text{CH}_2\text{CN}$, $-\text{CHO}$, alkylcarbonyl, (halogenated alkyl)carbonyl, arylcarbonyl, alkoxycarbonyl, aryloxy carbonyl, alkylaminocarbonyl, alkenyl, $-\text{C}(=\text{NH})\text{NH}_2$, $-\text{C}(=\text{NH})\text{NHCN}$, alkylsulfonyl, arylsulfonyl, amino, $-\text{NO}$ or heterocyclyl; or
- 15 R^{4a} , R^{4b} together are optionally substituted alkylene, wherein one $-\text{CH}_2-$ of alkylene may be replaced by an oxygen atom or $-\text{NR}^{10}$;
- X^2 is $-\text{O}-$, $-\text{NR}^{11a}-$, $-\text{S}-$, $>\text{CR}^{12a}\text{R}^{12b}$ or a bond;
- 20 X^3 is $-\text{O}-$, $-\text{NR}^{11b}-$, $-\text{S}-$, $>\text{CR}^{13a}\text{R}^{13b}$ or a bond;
- R^5 is optionally substituted aryl, optionally substituted cycloalkyl or optionally substituted heterocyclyl;
- 25 R^7 is hydrogen or alkyl;
- R^8 is hydrogen, alkyl, cycloalkyl, aminoalkyl, optionally substituted arylalkyl or heterocyclyl; or
- 30 R^8 , R^1 together are alkylene;
- R^9 is hydrogen, alkyl, cycloalkylalkyl, halogenated alkyl, hydroxyalkyl, alkoxyalkyl, aminoalkyl, CH_2CN , arylalkyl, cycloalkyl, $-\text{CHO}$, alkylcarbonyl, (halogenated alkyl)carbonyl, arylcarbonyl, alkoxycarbonyl, aryloxy carbonyl, alkylaminocarbonyl, alkenyl, $-\text{C}(=\text{NH})\text{NH}_2$, $-\text{C}(=\text{NH})\text{NHCN}$, alkylsulfonyl, arylsulfonyl, amino, $-\text{NO}$ or heterocyclyl;
- 35 R^{10} is hydrogen or alkyl;
- 40 R^{11a} is hydrogen or alkyl;
- R^{11b} is hydrogen or alkyl;

- R^{12a} is hydrogen, optionally substituted alkyl, alkylaminoalkyl, dialkylaminoalkyl, heterocycl-
alkyl, optionally substituted aryl or hydroxy;
- 5 R^{12b} is hydrogen or alkyl, or
- R^{12a} , R^{12b}
together with the carbon atom to which they are attached form a carbonyl or are optionally
substituted alkylene, wherein one $-CH_2-$ of alkylene may be replaced by an oxygen atom or -
10 $NR^{14}-$;
- R^{13a} is hydrogen, optionally substituted alkyl, alkylaminoalkyl, dialkylaminoalkyl, heterocycl-
alkyl, optionally substituted aryl or hydroxy;
- 15 R^{13b} is hydrogen or alkyl,
- R^{13a} , R^{13b}
together with the carbon atom to which they are attached form a carbonyl or are optionally
substituted alkylene, wherein one $-CH_2-$ of alkylene may be replaced by an oxygen atom or -
20 $NR^{15}-$;
- R^{14} is hydrogen or alkyl; and
- R^{15} is hydrogen or alkyl,
- 25 or a physiologically tolerated salt thereof.

Thus, the terms aminochromane, aminothiochromane and amino-1,2,3,4-tetrahydroquinoline deriv-
atives are used herein to denote in particular aminochromanes (A^2 is $-O-$), thiochromanes (A^2 is -
30 $S-$) and 1,2,3,4-tetrahydroquinolines (A^2 is $-NR^9-$) as well as fused tetrahydropyranes, tetrahy-
drothiopyranes and tetrahydropyridines wherein the benzene ring of the chromanes, thiochromanes
and 1,2,3,4-tetrahydroquinolines is replaced by a 5- or 6-membered heterocyclic ring.

Said compounds of formula (I), i.e., the aminochromane, aminothiochromane and amino-1,2,3,4-
35 tetrahydroquinoline derivatives of formula (I) and their physiologically tolerated salts, are glycine
transporter inhibitors and thus useful as pharmaceuticals. Compounds of formula (I) combine high
metabolic stability with high affinity. Compounds of formula (I) show favorable efflux properties
which may lead to enhanced oral bioavailability and/or increased brain availability. Compounds of
formula (I) combine high metabolic stability and high affinity with favorable efflux properties.

40

The present invention thus further relates to the compounds of formula (I) for use in therapy.

The present invention also relates to pharmaceutical compositions which comprise a carrier and a compound of formula (I).

5 In particular, said compounds, i.e., the aminochromane, aminothiochromane and amino-1,2,3,4-tetrahydroquinoline derivatives and their physiologically tolerated salts, are inhibitors of the glycine transporter GlyT1.

10 The present invention thus further relates to the compounds of formula (I) for use in inhibiting the glycine transporter GlyT1.

The present invention also relates to the use of the compounds of formula (I) in the manufacture of a medicament for inhibiting the glycine transporter GlyT1 and corresponding methods of inhibiting the glycine transporter GlyT1.

15 Glycine transport inhibitors and in particular inhibitors of the glycine transporter GlyT1 are known to be useful in treating a variety of neurologic and psychiatric disorders.

20 The present invention thus further relates to the compounds of formula (I) for use in treating a neurologic or psychiatric disorder.


The present invention further relates to the compounds of formula (I) for use in treating pain.

25 The present invention also relates to the use of the compounds of formula (I) in the manufacture of a medicament for treating a neurologic or psychiatric disorder and corresponding methods of treating said disorders. The present invention also relates to the use of the compounds of formula (I) in the manufacture of a medicament for treating pain and corresponding methods of treating pain.

Detailed Description Of The Invention

30 Provided that the aminochromane, aminothiochromane and amino-1,2,3,4-tetrahydroquinoline derivatives of the formula (I) of a given constitution may exist in different spatial arrangements, for example if they possess one or more centers of asymmetry, polysubstituted rings or double bonds, or as different tautomers, it is also possible to use enantiomeric mixtures, in particular racemates, diastereomeric mixtures and tautomeric mixtures, preferably, however, the respective essentially
35 pure enantiomers, diastereomers and tautomers of the compounds of formula (I) and/or of their salts.


According to one embodiment, an enantiomer of the compounds of the present invention has the following formula:



wherein R^1 , W , A^1 , Q , Y , R^6 , $n1$, $n2$, X^1 , A , R^2 , A^2 , R^3 , Y^1 , R^{4a} , R^{4b} , X^2 , X^3 , R^5 are as defined herein.


5

According to another embodiment, an enantiomer of the compounds of the present invention has the following formula:



10 wherein R^1 , W , A^1 , Q , Y , R^6 , $n1$, $n2$, X^1 , A , R^2 , A^2 , R^3 , Y^1 , R^{4a} , R^{4b} , X^2 , X^3 , R^5 are as defined herein.


According to one embodiment, an enantiomer of the compounds of the present invention has the following formula:



15

wherein R^1 , W, A^1 , Q, Y, R^6 , n1, n2, X^1 , A, R^2 , A^2 , R^3 , Y^1 , R^{4a} , R^{4b} , X^2 , X^3 , R^5 are as defined herein.

- 5 According to another embodiment, an enantiomer of the compounds of the present invention has the following formula:



- 10 wherein R^1 , W, A^1 , Q, Y, R^6 , n1, n2, X^1 , A, R^2 , A^2 , R^3 , Y^1 , R^{4a} , R^{4b} , X^2 , X^3 , R^5 are as defined herein.

- The physiologically tolerated salts of the aminochromane, aminothiochromane and amino-1,2,3,4-tetrahydroquinoline derivatives of the formula (I) are especially acid addition salts with physiologically tolerated acids. Examples of suitable physiologically tolerated organic and inorganic acids are
- 15 hydrochloric acid, hydrobromic acid, phosphoric acid, sulfuric acid, C_1 - C_4 -alkylsulfonic acids, such as methanesulfonic acid, cycloaliphatic sulfonic acids, such as S-(+)-10-camphor sulfonic acid, aromatic sulfonic acids, such as benzenesulfonic acid and toluenesulfonic acid, di- and tricarboxylic acids and hydroxycarboxylic acids having 2 to 10 carbon atoms, such as oxalic acid, malonic acid, maleic acid, fumaric acid, lactic acid, tartaric acid, citric acid, glycolic acid, adipic acid and
- 20 benzoic acid. Other utilizable acids are described, e.g., in Fortschritte der Arzneimittelforschung [Advances in drug research], Volume 10, pages 224 ff., Birkhäuser Verlag, Basel and Stuttgart, 1966. The physiologically tolerated salts of the aminochromane, aminothiochromane and amino-1,2,3,4-tetrahydroquinoline derivatives also include salts of a physiologically tolerated anion with aminochromane, aminothiochromane and amino-1,2,3,4-tetrahydroquinoline derivatives wherein
- 25 one or more than one nitrogen atom is quaternized, e.g. with an alkyl residue (e.g. methyl or ethyl).

- The present invention moreover relates to compounds of formula (I) as defined herein, wherein at least one of the atoms has been replaced by its stable, non-radioactive isotope (e.g., hydrogen by deuterium, ^{12}C by ^{13}C , ^{14}N by ^{15}N , ^{16}O by ^{18}O) and preferably wherein at least one hydrogen atom
- 30 has been replaced by a deuterium atom.

Of course, such compounds contain more of the respective isotope than this naturally occurs and thus is anyway present in the compounds (I).

Stable isotopes (e.g., deuterium, ^{13}C , ^{15}N , ^{18}O) are nonradioactive isotopes which contain one or more additional neutron than the normally abundant isotope of the respective atom. Deuterated compounds have been used in pharmaceutical research to investigate the in vivo metabolic fate of the compounds by evaluation of the mechanism of action and metabolic pathway of the non-deuterated parent compound (Blake et al. *J. Pharm. Sci.* 64, 3, 367-391 (1975)). Such metabolic studies are important in the design of safe, effective therapeutic drugs, either because the in vivo active compound administered to the patient or because the metabolites produced from the parent compound prove to be toxic or carcinogenic (Foster et al., *Advances in Drug Research* Vol. 14, pp. 2-36, Academic Press, London, 1985; Kato et al., *J. Labelled Comp. Radiopharmaceut.*, 36(10):927-932 (1995); Kushner et al., *Can. J. Physiol. Pharmacol.*, 77, 79-88 (1999)).

Incorporation of a heavy atom particularly substitution of deuterium for hydrogen, can give rise to an isotope effect that could alter the pharmacokinetics of the drug. This effect is usually insignificant if the label is placed at a metabolically inert position of the molecule.

Stable isotope labeling of a drug can alter its physico-chemical properties such as pKa and lipid solubility. These changes may influence the fate of the drug at different steps along its passage through the body. Absorption, distribution, metabolism or excretion can be changed. Absorption and distribution are processes that depend primarily on the molecular size and the lipophilicity of the substance. These effects and alterations can affect the pharmacodynamic response of the drug molecule if the isotopic substitution affects a region involved in a ligand-receptor interaction.

Drug metabolism can give rise to large isotopic effect if the breaking of a chemical bond to a deuterium atom is the rate limiting step in the process. While some of the physical properties of a stable isotope-labeled molecule are different from those of the unlabeled one, the chemical and biological properties are the same, with one important exception: because of the increased mass of the heavy isotope, any bond involving the heavy isotope and another atom will be stronger than the same bond between the light isotope and that atom. In any reaction in which the breaking of this bond is the rate limiting step, the reaction will proceed slower for the molecule with the heavy isotope due to "kinetic isotope effect". A reaction involving breaking a C--D bond can be up to 700 percent slower than a similar reaction involving breaking a C--H bond. If the C--D bond is not involved in any of the steps leading to the metabolite, there may not be any effect to alter the behavior of the drug. If a deuterium is placed at a site involved in the metabolism of a drug, an isotope effect will be observed only if breaking of the C--D bond is the rate limiting step. There is evidence to suggest that whenever cleavage of an aliphatic C--H bond occurs, usually by oxidation catalyzed by a mixed-function oxidase, replacement of the hydrogen by deuterium will lead to observable isotope effect. It is also important to understand that the incorporation of deuterium at the site of metabolism slows its rate to the point where another metabolite produced by attack at a carbon atom not substituted by deuterium becomes the major pathway a process called "metabolic switching".

Deuterium tracers, such as deuterium-labeled drugs and doses, in some cases repeatedly, of thousands of milligrams of deuterated water, are also used in healthy humans of all ages, including neonates and pregnant women, without reported incident (e.g. Pons G and Rey E, *Pediatrics* 1999 104: 633; Coward W A et al., *Lancet* 1979 7: 13; Schwarcz H P, *Control. Clin. Trials* 1984 5(4 Suppl): 573; Rodewald L E et al., *J. Pediatr.* 1989 114: 885; Butte N F et al. *Br. J. Nutr.* 1991 65: 3; MacLennan A H et al. *Am. J. Obstet Gynecol.* 1981 139: 948). Thus, it is clear that any deuterium released, for instance, during the metabolism of compounds of this invention poses no health risk.

The weight percentage of hydrogen in a mammal (approximately 9%) and natural abundance of deuterium (approximately 0.015%) indicates that a 70 kg human normally contains nearly a gram of deuterium. Furthermore, replacement of up to about 15% of normal hydrogen with deuterium has been effected and maintained for a period of days to weeks in mammals, including rodents and dogs, with minimal observed adverse effects (Czajka D M and Finkel A J, *Ann. N.Y. Acad. Sci.* 1960 84: 770; Thomson J F, *Ann. New York Acad. Sci* 1960 84: 736; Czajka D M et al., *Am. J. Physiol.* 1961 201: 357). Higher deuterium concentrations, usually in excess of 20%, can be toxic in animals. However, acute replacement of as high as 15%-23% of the hydrogen in humans' fluids with deuterium was found not to cause toxicity (Blagojevic N et al. in "Dosimetry & Treatment Planning for Neutron Capture Therapy", Zamenhof R, Solares G and Harling O Eds. 1994. Advanced Medical Publishing, Madison Wis. pp.125-134; *Diabetes Metab.* 23: 251 (1997)).

Increasing the amount of deuterium present in a compound above its natural abundance is called enrichment or deuterium-enrichment. Examples of the amount of enrichment include from about 0.5, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 12, 16, 21, 25, 29, 33, 37, 42, 46, 50, 54, 58, 63, 67, 71, 75, 79, 84, 88, 92, 96, to about 100 mol %.

The hydrogens present on a particular organic compound have different capacities for exchange with deuterium. Certain hydrogen atoms are easily exchangeable under physiological conditions and, if replaced by deuterium atoms, it is expected that they will readily exchange for protons after administration to a patient. Certain hydrogen atoms may be exchanged for deuterium atoms by the action of a deuterium acid such as D₂SO₄/D₂O. Alternatively, deuterium atoms may be incorporated in various combinations during the synthesis of compounds of the invention. Certain hydrogen atoms are not easily exchangeable for deuterium atoms. However, deuterium atoms at the remaining positions may be incorporated by the use of deuterated starting materials or intermediates during the construction of compounds of the invention.

Deuterated and deuterium-enriched compounds of the invention can be prepared by using known methods described in the literature. Such methods can be carried out utilizing corresponding deuterated and optionally, other isotope-containing reagents and/or intermediates to synthesize the compounds delineated herein, or invoking standard synthetic protocols known in the art for introducing isotopic atoms to a chemical structure. Relevant procedures and intermediates are disclosed, for instance in Lizondo, J et al., *Drugs Fut*, 21(11), 1116 (1996); Brickner, S J et al., *J Med Chem*, 39(3), 673 (1996); Mallesham, B et al., *Org Lett*, 5(7), 963 (2003); PCT publications

WO1997010223, WO2005099353, WO1995007271, WO2006008754; US Patent Nos. 7538189; 7534814; 7531685; 7528131; 7521421; 7514068; 7511013; and US Patent Application Publication Nos. 20090137457; 20090131485; 20090131363; 20090118238; 20090111840; 20090105338; 20090105307; 20090105147; 20090093422; 20090088416; 20090082471, the methods are hereby
 5 incorporated by reference.

The organic moieties mentioned in the above definitions of the variables are - like the term halogen - collective terms for individual listings of the individual group members. The prefix C_n-C_m indicates in each case the possible number of carbon atoms in the group. The prefix M_n-M_m indicates in
 10 each case the possible number of ring forming atoms (ring members) in the group.

Unless indicated otherwise, the term "substituted" means that a radical is substituted with 1, 2 or 3, especially 1, substituent which, according to a particular embodiment of the invention, are independently selected from the group consisting of halogen, C₁-C₄-alkyl, C₃-C₆-aryl-C₁-C₄-alkyl, halogenated-C₁-C₄-alkyl, hydroxy-C₁-C₄-alkyl, hydroxy-(halogenated C₁-C₄-alkyl), C₁-C₄-alkoxy-C₁-C₄-alkyl, amino-C₁-C₄-alkyl, M₃-M₁₂-heterocyclyl-C₁-C₄-alkyl, C₃-C₇-cycloalkyl, C₂-C₄-alkenyl, -CN, -CO₂H, C₁-C₄-alkoxycarbonyl, aminocarbonyl, C₁-C₄-alkylaminocarbonyl, (di-C₁-C₄-alkylamino)carbonyl, C₆-C₁₂-arylaminocarbonyl, M₃-M₁₂-heterocyclylaminocarbonyl, C₆-C₁₂-aryl, oxo (=O), OH, C₁-C₄-alkoxy, halogenated-C₁-C₄-alkoxy, C₃-C₇-cycloalkoxy, carboxy-C₁-C₄-alkoxy, C₆-C₁₂-aryl-C₁-C₄-alkoxy, C₆-C₁₂-aryloxy, M₃-M₁₂-heterocyclyl-C₁-C₄-alkoxy, SH, C₁-C₄-alkylthio, C₁-C₄-alkylsulfonyl, C₁-C₄-alkylaminosulfonyl, di-C₁-C₄-alkylaminosulfonyl, C₃-C₆-arylsulfonyl, aminosulfonyl, C₃-C₆-arylaminosulfonyl, M₃-M₁₂-heterocyclylaminosulfonyl, NH₂, C₁-C₄-alkylamino, di-C₁-C₄-alkylamino, C₆-C₁₂-aryl-C₁-C₄-alkylamino, C₁-C₄-alkylcarbonylamino, C₆-C₁₂-arylcarbonylamino, M₃-M₁₂-heterocyclylcarbonylamino, C₁-C₆-alkylsulfonylamino, C₆-C₁₂-arylsulfonylamino, M₃-M₁₂-heterocyclylsulfonylamino and M₃-M₁₂-heterocyclyl, wherein aryl and heterocyclyl may be unsubstituted or substituted with 1, 2 or 3 substituents selected from the group consisting of halogen, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy and C₁-C₄-haloalkoxy.
 15
 20
 25

30 The term halogen denotes in each case fluorine, bromine, chlorine or iodine, in particular fluorine or chlorine.

C₁-C₄-Alkyl is a straight-chain or branched alkyl group having from 1 to 4 carbon atoms. Examples of an alkyl group are methyl, C₂-C₄-alkyl such as ethyl, n-propyl, iso-propyl, n-butyl, 2-butyl, iso-butyl or tert-butyl. C₁-C₂-Alkyl is methyl or ethyl, C₁-C₃-alkyl is additionally n-propyl or isopropyl.
 35

C₁-C₆-Alkyl is a straight-chain or branched alkyl group having from 1 to 6 carbon atoms. Examples include methyl, C₂-C₄-alkyl as mentioned herein and also pentyl, 1-methylbutyl, 2-methylbutyl, 3-methylbutyl, 2,2-dimethylpropyl, 1-ethylpropyl, hexyl, 1,1-dimethylpropyl, 1,2-dimethylpropyl, 1-methylpentyl, 2-methylpentyl, 3-methylpentyl, 4-methylpentyl, 1,1-dimethylbutyl, 1,2-dimethylbutyl, 1,3-dimethylbutyl, 2,2-dimethylbutyl, 2,3-dimethylbutyl, 3,3-dimethylbutyl, 1-
 40

ethylbutyl, 2-ethylbutyl, 1,1,2-trimethylpropyl, 1,2,2-trimethylpropyl, 1-ethyl-1-methylpropyl and 1-ethyl-2-methylpropyl.

5 Halogenated C₁-C₆-alkyl is a straight-chain or branched alkyl group having 1 to 6 carbon atoms, preferably 1 to 3 carbon atoms, more preferably 1 or 2 carbon atoms, wherein at least one, e.g. 1, 2, 3, 4 or all of the hydrogen atoms are replaced by 1, 2, 3, 4 or a corresponding number of identical or different halogen atoms, such as in halogenomethyl, dihalogenomethyl, trihalogenomethyl, (R)-1-halogenoethyl, (S)-1-halogenoethyl, 2-halogenoethyl, 1,1-dihalogenoethyl, 2,2-dihalogenoethyl, 2,2,2-trihalogenoethyl, (R)-1-halogenopropyl, (S)-1-halogenopropyl, 2-halogenopropyl, 3-
10 halogenopropyl, 1,1-dihalogenopropyl, 2,2-dihalogenopropyl, 3,3-dihalogenopropyl, 3,3,3-trihalogenopropyl, (R)-2-halogeno-1-methylethyl, (S)-2-halogeno-1-methylethyl, (R)-2,2-dihalogeno-1-methylethyl, (S)-2,2-dihalogeno-1-methylethyl, (R)-1,2-dihalogeno-1-methylethyl, (S)-1,2-dihalogeno-1-methylethyl, (R)-2,2,2-trihalogeno-1-methylethyl, (S)-2,2,2-trihalogeno-1-methylethyl, 2-halogeno-1-(halogenomethyl)ethyl, 1-(dihalogenomethyl)-2,2-dihalogenoethyl, (R)-
15 1-halogenobutyl, (S)-1-halogenobutyl, 2-halogenobutyl, 3-halogenobutyl, 4-halogenobutyl, 1,1-dihalogenobutyl, 2,2-dihalogenobutyl, 3,3-dihalogenobutyl, 4,4-dihalogenobutyl, 4,4,4-trihalogenobutyl, etc. Particular examples include the fluorinated C₁-C₆ alkyl groups as defined, such as trifluoromethyl.

20 C₃-C₁₂-Cycloalkyl-C₁-C₄-alkyl is a straight-chain or branched alkyl group having 1 to 4 carbon atoms, preferably 1 to 3 carbon atoms, more preferably 1 or 2 carbon atoms, in particular 1 or two carbon atoms, wherein one hydrogen atom is replaced by a cycloaliphatic radical having from 3 to 12 carbon atoms such as in cyclopropylmethyl, cyclobutylmethyl, cyclopentylmethyl and cyclohexylmethyl.

25 C₆-C₁₂-Aryl-C₁-C₄-alkyl is a straight-chain or branched alkyl group having 1 to 4 carbon atoms, preferably 1 to 3 carbon atoms, more preferably 1 or 2 carbon atoms, in particular 1 or two carbon atoms, wherein one hydrogen atom is replaced by C₆-C₁₂-aryl, such as in benzyl.

30 Hydroxy-C₁-C₄-alkyl is a straight-chain or branched alkyl group having 1 to 4 carbon atoms, preferably 1 to 3 carbon atoms, more preferably 1 or 2 carbon atoms, wherein one or two hydrogen atoms are replaced by one or two hydroxyl groups, such as in hydroxymethyl, (R)-1-hydroxyethyl, (S)-1-hydroxyethyl, 2-hydroxyethyl, (R)-1-hydroxypropyl, (S)-1-hydroxypropyl, 2-hydroxypropyl, 3-hydroxypropyl, (R)-2-hydroxy-1-methylethyl, (S)-2-hydroxy-1-methylethyl, 2-hydroxy-1-
35 (hydroxymethyl)ethyl, (R)-1-hydroxybutyl, (S)-1-hydroxybutyl, 2-hydroxybutyl, 3-hydroxybutyl, 4-hydroxybutyl.

C₁-C₆-Alkoxy-C₁-C₄-alkyl is a straight-chain or branched alkyl group having 1 to 4 carbon atoms, preferably 1 to 3 carbon atoms, more preferably 1 or 2 carbon atoms, wherein one or two hydrogen
40 atoms are replaced by one or two alkoxy groups having 1 to 6, preferably 1 to 4, in particular 1 or 2 carbon atoms, such as in methoxymethyl, (R)-1-methoxyethyl, (S)-1-methoxyethyl, 2-methoxyethyl, (R)-1-methoxypropyl, (S)-1-methoxypropyl, 2-methoxypropyl, 3-methoxypropyl,

(R)-2-methoxy-1-methylethyl, (S)-2-methoxy-1-methylethyl, 2-methoxy-1-(methoxymethyl)ethyl, (R)-1-methoxybutyl, (S)-1-methoxybutyl, 2-methoxybutyl, 3-methoxybutyl, 4-methoxybutyl, ethoxymethyl, (R)-1-ethoxyethyl, (S)-1-ethoxyethyl, 2-ethoxyethyl, (R)-1-ethoxypropyl, (S)-1-ethoxypropyl, 2-ethoxypropyl, 3-ethoxypropyl, (R)-2-ethoxy-1-methylethyl, (S)-2-ethoxy-1-methylethyl, 2-ethoxy-1-(ethoxymethyl)ethyl, (R)-1-ethoxybutyl, (S)-1-ethoxybutyl, 2-ethoxybutyl, 3-ethoxybutyl, 4-ethoxybutyl.

Amino-C₁-C₄-alkyl is a straight-chain or branched alkyl group having 1 to 4 carbon atoms, preferably 1 to 3 carbon atoms, more preferably 1 or 2 carbon atoms, in particular 1 or two carbon atoms, wherein one hydrogen atom is replaced by an amino group, such as in aminomethyl, 2-aminoethyl.

C₁-C₆-Alkylamino-C₁-C₄-alkyl is a straight-chain or branched alkyl group having 1 to 4 carbon atoms, preferably 1 to 3 carbon atoms, more preferably 1 or 2 carbon atoms, in particular 1 or two carbon atoms, wherein one hydrogen atom is replaced by a C₁-C₆-alkylamino group, in particular by a C₁-C₄-alkylamino group, such as in methylaminomethyl, ethylaminomethyl, n-propylaminomethyl, iso-propylaminomethyl, n-butylaminomethyl, 2-butylaminomethyl, iso-butylaminomethyl or tert-butylaminomethyl.

Di-C₁-C₆-Alkylamino-C₁-C₄-alkyl is a straight-chain or branched alkyl group having 1 to 4 carbon atoms, preferably 1 to 3 carbon atoms, more preferably 1 or 2 carbon atoms, in particular 1 or two carbon atoms, wherein one hydrogen atom is replaced by a di-C₁-C₆-Alkylamino group, in particular by a di-C₁-C₄-alkylamino group, such as in dimethylaminomethyl.

C₁-C₆-Alkylcarbonylamino-C₁-C₄-alkyl is a straight-chain or branched alkyl group having 1 to 4 carbon atoms, preferably 1 to 3 carbon atoms, more preferably 1 or 2 carbon atoms, in particular 1 or two carbon atoms, wherein one hydrogen atom is replaced by a C₁-C₆-alkylcarbonylamino group, in particular by a C₁-C₄-alkylcarbonylamino group, such as in methylcarbonylaminomethyl, ethylcarbonylaminomethyl, n-propylcarbonylaminomethyl, iso-propylcarbonylaminomethyl, n-butylcarbonylaminomethyl, 2-butylcarbonylaminomethyl, iso-butylcarbonylaminomethyl or tert-butylcarbonylaminomethyl.

C₁-C₆-Alkylaminocarbonylamino-C₁-C₄-alkyl is a straight-chain or branched alkyl group having 1 to 4 carbon atoms, preferably 1 to 3 carbon atoms, more preferably 1 or 2 carbon atoms, in particular 1 or two carbon atoms, wherein one hydrogen atom is replaced by a C₁-C₆-alkylaminocarbonylamino group, in particular by a C₁-C₄-alkylaminocarbonylamino group, such as in methylaminocarbonylaminomethyl, ethylaminocarbonylaminomethyl, n-propylaminocarbonylaminomethyl, iso-propylaminocarbonylaminomethyl, n-butylaminocarbonylaminomethyl, 2-butylaminocarbonylaminomethyl, iso-butylaminocarbonylaminomethyl or tert-butylaminocarbonylaminomethyl.

Di-C₁-C₆-alkylaminocarbonylamino-C₁-C₄-alkyl is a straight-chain or branched alkyl group having 1 to 4 carbon atoms, preferably 1 to 3 carbon atoms, more preferably 1 or 2 carbon atoms, in par-

ticular 1 or two carbon atoms, wherein one hydrogen atom is replaced by a di-C₁-C₆-alkylaminocarbonylamino group, in particular by a di-C₁-C₄-alkylaminocarbonylamino group, such as in dimethylaminocarbonylaminomethyl, dimethylaminocarbonylaminoethyl, dimethylaminocarbonylaminon-propyl.

5

C₁-C₆-Alkylsulfonylamino-C₁-C₄-alkyl is a straight-chain or branched alkyl group having 1 to 4 carbon atoms, preferably 1 to 3 carbon atoms, more preferably 1 or 2 carbon atoms, in particular 1 or two carbon atoms, wherein one hydrogen atom is replaced by a C₁-C₆-alkylsulfonylamino group, in particular by a C₁-C₄-alkylsulfonylamino group, such as in methylsulfonylaminomethyl, ethylsulfonylaminomethyl, n-propylsulfonylaminomethyl, iso-propylsulfonylaminomethyl, n-butylsulfonylaminomethyl, 2-butylsulfonylaminomethyl, iso-butylsulfonylaminomethyl or tert-butylsulfonylaminomethyl.

10

(C₆-C₁₂-Aryl-C₁-C₆-alkyl)amino-C₁-C₄ alkyl is a straight-chain or branched alkyl group having 1 to 4 carbon atoms, preferably 1 to 3 carbon atoms, more preferably 1 or 2 carbon atoms, in particular 1 or two carbon atoms, wherein one hydrogen atom is replaced by a (C₆-C₁₂-aryl-C₁-C₆-alkyl)amino group, in particular a (C₆-C₁₂-aryl-C₁-C₂-alkyl)amino group, such as in benzylaminomethyl.

15

M₃-M₁₂-Heterocyclyl-C₁-C₄-alkyl is a straight-chain or branched alkyl group having 1 to 4 carbon atoms, preferably 1 to 3 carbon atoms, more preferably 1 or 2 carbon atoms, in particular 1 or two carbon atoms, wherein one hydrogen atom is replaced by M₃-M₁₂-heterocyclyl, such as in N-pyrrolidinylmethyl, N-piperidinylmethyl, N-morpholinylmethyl.

20

C₃-C₁₂-Cycloalkyl is a cycloaliphatic radical having from 3 to 12 carbon atoms. In particular, 3 to 6 carbon atoms form the cyclic structure, such as cyclopropyl, cyclobutyl, cyclopentyl and cyclohexyl. The cyclic structure may be unsubstituted or may carry 1, 2, 3 or 4 C₁-C₄ alkyl radicals, preferably one or more methyl radicals.

25

30 Carbonyl is >C=O.

C₁-C₆-Alkylcarbonyl is a radical of the formula R-C(O)-, wherein R is an alkyl radical having from 1 to 6, preferably from 1 to 4, in particular 1 or 2 carbon atoms as defined herein. Examples include acetyl, propionyl, n-butyryl, 2-methylpropionyl, pivaloyl.

35

Halogenated C₁-C₆-alkylcarbonyl is C₁-C₆-alkylcarbonyl as defined herein, wherein at least one, e.g. 1, 2, 3, 4 or all of the hydrogen atoms are replaced by 1, 2, 3, 4 or a corresponding number of identical or different halogen atoms. Examples include fluoromethylcarbonyl, difluoromethylcarbonyl, trifluoromethylcarbonyl. Further examples are 1,1,1-trifluoroeth-2-ylcarbonyl, 1,1,1-trifluoroprop-3-ylcarbonyl.

40

C₆-C₁₂-Arylcarbonyl is a radical of the formula R-C(O)-, wherein R is an aryl radical having from 6 to 12 carbon atoms as defined herein. Examples include benzoyl.

5 C₁-C₆-Alkoxy carbonyl is a radical of the formula R-O-C(O)-, wherein R is an alkyl radical having from 1 to 6, preferably from 1 to 4, in particular 1 or 2 carbon atoms as defined herein. Examples include methoxycarbonyl and tert-butyloxycarbonyl.

10 Halogenated C₁-C₆-alkoxy carbonyl is a C₁-C₆-alkoxy carbonyl as defined herein, wherein at least one, e.g. 1, 2, 3, 4 or all of the hydrogen atoms are replaced by 1, 2, 3, 4 or a corresponding number of identical or different halogen atoms.

C₆-C₁₂-Aryloxy carbonyl is a radical of the formula R-O-C(O)-, wherein R is an aryl radical having from 6 to 12 carbon atoms as defined herein. Examples include phenoxycarbonyl.

15 Cyano is -C≡N.

Aminocarbonyl is NH₂C(O)-.

20 C₁-C₆-Alkylaminocarbonyl is a radical of the formula R-NH-C(O)-, wherein R is an alkyl radical having from 1 to 6, preferably from 1 to 4, in particular 1 or 2 carbon atoms as defined herein. Examples include methylaminocarbonyl.

25 (Halogenated C₁-C₄-alkyl)aminocarbonyl is a C₁-C₄-alkylaminocarbonyl as defined herein, wherein at least one, e.g. 1, 2, 3, 4 or all of the hydrogen atoms are replaced by 1, 2, 3, 4 or a corresponding number of identical or different hydrogen atoms.

C₆-C₁₂-Arylaminocarbonyl is a radical of the formula R-NH-C(O)-, wherein R is an aryl radical having from 6 to 12 carbon atoms as defined herein. Examples include phenylaminocarbonyl.

30 C₂-C₆-Alkenyl is a singly unsaturated hydrocarbon radical having 2, 3, 4, 5 or 6 carbon atoms, e.g. vinyl, allyl (2-propen-1-yl), 1-propen-1-yl, 2-propen-2-yl, methallyl (2-methylprop-2-en-1-yl) and the like. C₃-C₅-Alkenyl is, in particular, allyl, 1-methylprop-2-en-1-yl, 2-buten-1-yl, 3-buten-1-yl, methallyl, 2-penten-1-yl, 3-penten-1-yl, 4-penten-1-yl, 1-methylbut-2-en-1-yl or 2-ethylprop-2-en-1-yl.

35 C₂-C₆-Alkynyl is a singly unsaturated hydrocarbon radical having 2, 3, 4, 5 or 6 carbon atoms, e.g. ethynyl, 2-propyn-1-yl, 1-propyn-1-yl, 2-propyn-2-yl and the like. C₃-C₅-Alkynyl is, in particular, 2-propyn-1-yl, 2-butyne-1-yl, 3-butyne-1-yl, 2-pentyne-1-yl, 3-pentyne-1-yl, 4-pentyne-1-yl.

40 C₁-C₄-Alkylene is straight-chain or branched alkylene group having from 1 to 4 carbon atoms. Examples include methylene and ethylene. A further example is propylene.

C₂-C₆-Alkylene is straight-chain or branched alkylene group having from 2 to 6 carbon atoms. Examples include ethylene. A further example is propylene.

C₂-C₄-Alkenylene is straight-chain or branched alkenylene group having from 2 to 4 carbon atoms.

5

C₂-C₄-Alkynylene is straight-chain or branched alkynylene group having from 2 to 4 carbon atoms. Examples include propynylene.

10 C₆-C₁₂-Aryl is a 6- to 12-membered, in particular 6- to 10-membered, aromatic cyclic radical. Examples include phenyl and naphthyl.

C₃-C₁₂-Arylene is an aryl diradical. Examples include phen-1,4-ylene and phen-1,3-ylene.

Hydroxy is -OH.

15

C₁-C₆-Alkoxy is a radical of the formula R-O-, wherein R is a straight-chain or branched alkyl group having from 1 to 6, in particular 1 to 4 carbon atoms. Examples include methoxy, ethoxy, n-propoxy, isopropoxy, n-butoxy, 2-butoxy, iso-butoxy (2-methylpropoxy), tert.-butoxy pentyloxy, 1-methylbutoxy, 2-methylbutoxy, 3-methylbutoxy, 2,2-dimethylpropoxy, 1-ethylpropoxy, hex-
20 yloxy, 1,1-dimethylpropoxy, 1,2-dimethylpropoxy, 1-methylpentyloxy, 2-methylpentyloxy, 3-methylpentyloxy, 4-methylpentyloxy, 1,1-dimethylbutoxy, 1,2-dimethylbutoxy, 1,3-dimethylbutoxy, 2,2-dimethylbutoxy, 2,3-dimethylbutoxy, 3,3-dimethylbutoxy, 1-ethylbutoxy, 2-ethylbutoxy, 1,1,2-trimethylpropoxy, 1,2,2-trimethylpropoxy, 1-ethyl-1-methylpropoxy and 1-ethyl-2-methylpropoxy.

25

Halogenated C₁-C₆-alkoxy is a straight-chain or branched alkoxy group having from 1 to 6, preferably from 1 to 4, in particular 1 or 2 carbon atoms, wherein at least one, e.g. 1, 2, 3, 4 or all of the hydrogen atoms are replaced by 1, 2, 3, 4 or a corresponding number of identical or different halogen atoms, such as in halogenomethoxy, dihalogenomethoxy, trihalogenomethoxy, (R)-1-halogenoethoxy, (S)-1-halogenoethoxy, 2-halogenoethoxy, 1,1-dihalogenoethoxy, 2,2-dihalogenoethoxy, 2,2,2-trihalogenoethoxy, (R)-1-halogenopropoxy, (S)-1-halogenopropoxy, 2-halogenopropoxy, 3-halogenopropoxy, 1,1-dihalogenopropoxy, 2,2-dihalogenopropoxy, 3,3-dihalogenopropoxy, 3,3,3-trihalogenopropoxy, (R)-2-halogeno-1-methylethoxy, (S)-2-halogeno-1-methylethoxy, (R)-2,2-dihalogeno-1-methylethoxy, (S)-2,2-dihalogeno-1-methylethoxy, (R)-1,2-dihalogeno-1-methylethoxy, (S)-1,2-dihalogeno-1-methylethoxy, (R)-2,2,2-trihalogeno-1-methylethoxy, (S)-2,2,2-trihalogeno-1-methylethoxy, 2-halogeno-1-(halogenomethyl)ethoxy, 1-(dihalogenomethyl)-2,2-dihalogenoethoxy, (R)-1-halogenobutoxy, (S)-1-halogenobutoxy, 2-halogenobutoxy, 3-halogenobutoxy, 4-halogenobutoxy, 1,1-dihalogenobutoxy, 2,2-dihalogenobutoxy, 3,3-dihalogenobutoxy, 4,4-dihalogenobutoxy, 4,4,4-trihalogenobutoxy, etc.
40 Particular examples include the fluorinated C₁-C₄ alkoxy groups as defined, such as trifluoromethoxy.

C₁-C₆-Hydroxyalkoxy is an alkoxy radical having from 1 to 6, preferably from 1 to 4 carbon atoms as defined herein, wherein one or two hydrogen atoms are replaced by hydroxy. Examples include 2-hydroxyethoxy, 3-hydroxypropoxy, 2-hydroxypropoxy, 1-methyl-2-hydroxyethoxy and the like.

- 5 C₁-C₆-Alkoxy-C₁-C₄-alkoxy is an alkoxy radical having from 1 to 4 carbon atoms, preferably 1 or 2 carbon atoms as defined herein, wherein one or two hydrogen atoms are replaced by one or two alkoxy radicals having from 1 to 6, preferably from 1 to 4 carbon atoms as defined herein. Examples include methoxymethoxy, 2-methoxyethoxy, 1-methoxyethoxy, 3-methoxypropoxy, 2-methoxypropoxy, 1-methyl-1-methoxyethoxy, ethoxymethoxy, 2-ethoxyethoxy, 1-ethoxyethoxy, 3-ethoxypropoxy, 2-ethoxypropoxy, 1-methyl-1-ethoxyethoxy and the like.
- 10

Amino-C₁-C₄-alkoxy is an alkoxy radical having from 1 to 4, preferably 1 or 2 carbon atoms as defined herein, wherein one hydrogen atom is replaced by an amino group. Examples include 2-aminoethoxy.

- 15 C₁-C₆-Alkylamino-C₁-C₄-alkoxy is an alkoxy radical having from 1 to 4, preferably 1 or 2 carbon atoms as defined herein, wherein one hydrogen atom is replaced by an alkylamino group having from 1 to 6, preferably from 1 to 4 carbon atoms as defined herein. Examples include methylaminomethoxy, ethylaminomethoxy, n-propylaminomethoxy, iso-propylaminomethoxy, n-butylaminomethoxy, 2-butylaminomethoxy, iso-butylaminomethoxy, tert-butylaminomethoxy, 2-(methylamino)ethoxy, 2-(ethylamino)ethoxy, 2-(n-propylamino)ethoxy, 2-(iso-propylamino)ethoxy, 2-(n-butylamino)ethoxy, 2-(2-butylamino)ethoxy, 2-(iso-butylamino)ethoxy, 2-(tert-butylamino)ethoxy.
- 20

- 25 Di-C₁-C₆-alkylamino-C₁-C₄-alkoxy is an alkoxy radical having from 1 to 4, preferably 1 or 2 carbon atoms as defined herein, wherein one hydrogen atom is replaced by a di-alkylamino group having from 1 to 6, preferably from 1 to 4 carbon atoms as defined herein. Examples include dimethylaminomethoxy, diethylaminomethoxy, N-methyl-N-ethylamino)ethoxy, 2-(dimethylamino)ethoxy, 2-(diethylamino)ethoxy, 2-(N-methyl-N-ethylamino)ethoxy.
- 30

- C₁-C₆-Alkylcarbonylamino-C₁-C₄-alkoxy is an alkoxy radical having from 1 to 4, preferably 1 or 2 carbon atoms as defined herein, wherein one hydrogen atom is replaced by an alkylcarbonylamino group wherein the alkyl group has from 1 to 6, preferably from 1 to 4 carbon atoms as defined herein. Examples include methylcarbonylaminomethoxy, ethylcarbonylaminomethoxy, n-propylcarbonylaminomethoxy, iso-propylcarbonylaminomethoxy, n-butylcarbonylaminomethoxy, 2-butylcarbonylaminomethoxy, iso-butylcarbonylaminomethoxy, tert-butylcarbonylaminomethoxy, 2-(methylcarbonylamino)ethoxy, 2-(ethylcarbonylamino)ethoxy, 2-(n-propylcarbonylamino)ethoxy, 2-(iso-propylcarbonylamino)ethoxy, 2-(n-butylcarbonylamino)ethoxy, 2-(2-butylcarbonylamino)ethoxy, 2-(iso-butylcarbonylamino)ethoxy, 2-(tert-butylcarbonylamino)ethoxy.
- 35
- 40

C₆-C₁₂-Arylcarbonylamino-C₁-C₄-alkoxy is an alkoxy radical having from 1 to 4, preferably 1 or 2 carbon atoms as defined herein, wherein one hydrogen atom is replaced by a C₆-C₁₂-arylcarbonylamino group as defined herein. Examples include 2-(benzoylamino)ethoxy.

- 5 C₁-C₆-Alkoxy carbonylamino-C₁-C₄-alkoxy is an alkoxy radical having from 1 to 4, preferably 1 or 2 carbon atoms as defined herein, wherein one hydrogen atom is replaced by an alkoxy carbonylamino group wherein the alkoxy group has from 1 to 6, preferably from 1 to 4 carbon atoms as defined herein. Examples include methoxycarbonylaminomethoxy, ethoxycarbonylaminomethoxy, n-propoxycarbonylaminomethoxy, iso-propoxycarbonylaminomethoxy, n-butoxycarbonylaminomethoxy, 2-butoxycarbonylaminomethoxy, iso-butoxycarbonylaminomethoxy, tert-butoxycarbonylaminomethoxy, 2-(methoxycarbonylamino)ethoxy, 2-(ethoxycarbonylamino)ethoxy, 2-(n-propoxycarbonylamino)ethoxy, 2-(iso-propoxycarbonylamino)ethoxy, 2-(n-butoxycarbonylamino)ethoxy, 2-(2-butoxycarbonylamino)ethoxy, 2-(iso-butoxycarbonylamino)ethoxy, 2-(tert-butoxycarbonylamino)ethoxy.

- 15 C₂-C₆-Alkenyloxy is a radical of the formula R-O-, wherein R is a straight-chain or branched alkenyl group having from 2 to 6, in particular 2 to 4 carbon atoms. Examples include vinyloxy, allyloxy (2-propen-1-yloxy), 1-propen-1-yloxy, 2-propen-2-yloxy, methallyloxy (2-methylprop-2-en-1-yloxy) and the like. C₃-C₅-Alkenyloxy is, in particular, allyloxy, 1-methylprop-2-en-1-yloxy, 2-buten-1-yloxy, 3-buten-1-yloxy, methallyloxy, 2-penten-1-yloxy, 3-penten-1-yloxy, 4-penten-1-yloxy, 1-methylbut-2-en-1-yloxy or 2-ethylprop-2-en-1-yloxy.

- 20 C₆-C₁₂-Aryl-C₁-C₄-alkoxy is an alkoxy radical having from 1 to 4, preferably 1 or 2 carbon atoms as defined herein, wherein one hydrogen atom is replaced by a C₆-C₁₂-aryl group as defined herein. Examples include benzyloxy.

- 25 C₁-C₆-Alkylsulfonylamino-C₁-C₄-alkoxy is an alkoxy radical having from 1 to 4, preferably 1 or 2 carbon atoms as defined herein, wherein one hydrogen atom is replaced by an alkylsulfonylamino group having from 1 to 6, preferably from 1 to 4 carbon atoms as defined herein. Examples include 2-(methylsulfonylamino)ethoxy, 2-(ethylsulfonylamino)ethoxy, 2-[(2-methylpropyl)sulfonylamino]ethoxy.

- 30 (Halogenated C₁-C₆-alkyl)sulfonylamino-C₁-C₄-alkoxy is an alkoxy radical having from 1 to 4, preferably 1 or 2 carbon atoms as defined herein, wherein one hydrogen atom is replaced by an alkylsulfonylamino group having from 1 to 6, preferably from 1 to 4 carbon atoms as defined herein, wherein the alkyl group is halogenated. Examples include 2-(trifluoromethylsulfonylamino)ethoxy.

- 35 C₆-C₁₂-Arylsulfonylamino-C₁-C₄-alkoxy is an alkoxy radical having from 1 to 4, preferably 1 or 2 carbon atoms as defined herein, wherein one hydrogen atom is replaced by a C₆-C₁₂-arylsulfonylamino group as defined herein. Examples include 2-(phenylsulfonylamino)ethoxy, 2-(naphthylsulfonylamino)ethoxy.

(C₆-C₁₂-Aryl-C₁-C₆-alkyl)sulfonylamino-C₁-C₄-alkoxy is an alkoxy radical having from 1 to 4, preferably 1 or 2 carbon atoms as defined herein, wherein one hydrogen atom is replaced by a (C₆-C₁₂-aryl-C₁-C₆-alkyl)sulfonylamino group, preferably by a (C₆-C₁₂-aryl-C₁-C₂-alkyl)sulfonylamino group. Examples include 2-(benzylsulfonylamino)ethoxy.

M₃-M₁₂-Heterocyclylsulfonylamino-C₁-C₄-alkoxy is an alkoxy radical having from 1 to 4, preferably 1 or 2 carbon atoms as defined herein, wherein one hydrogen atom is replaced by a M₃-M₁₂-heterocyclylsulfonylamino group as defined herein. Examples include 2-(pyridin-3-yl-sulfonylamino)ethoxy.

M₃-M₁₂-Heterocyclyl-C₁-C₄-alkoxy is an alkoxy radical having from 1 to 4, preferably 1 or 2 carbon atoms as defined herein, wherein one hydrogen atom is replaced by a M₃-M₁₂-heterocyclyl group as defined herein. Examples include 2-(N-pyrrolidinyl)ethoxy, 2-(N-morpholinyl)ethoxy and 2-(N-imidazolyl)ethoxy.

C₁-C₂-Alkylenedioxo is a radical of the formula -O-R-O-, wherein R is a straight-chain or branched alkylene group having from 1 or 2 carbon atoms as defined herein. Examples include methylenedioxo.

C₆-C₁₂-Aryloxy is a radical of the formula R-O-, wherein R is an aryl group having from 6 to 12, in particular 6 carbon atoms as defined herein. Examples include phenoxy.

M₃-M₁₂-Heterocyclyloxy is a radical of the formula R-O-, wherein R is a M₃-M₁₂-heterocyclyl group having from 3 to 12, in particular from 3 to 7 ring forming atoms (ring members) as defined herein. Examples include pyridin-2-yloxy.

C₁-C₆-Alkylthio is a radical of the formula R-S-, wherein R is an alkyl radical having from 1 to 6, preferably from 1 to 4 carbon atoms as defined herein. Examples include methylthio, ethylthio, propylthio, butylthio, pentylthio, 1-methylbutylthio, 2-methylbutylthio, 3-methylbutylthio, 2,2-dimethylpropylthio, 1-ethylpropylthio, hexylthio, 1,1-dimethylpropylthio, 1,2-dimethylpropylthio, 1-methylpentylthio, 2-methylpentylthio, 3-methylpentylthio, 4-methylpentylthio, 1,1-dimethylbutylthio, 1,2-dimethylbutylthio, 1,3-dimethylbutylthio, 2,2-dimethylbutylthio, 2,3-dimethylbutylthio, 3,3-dimethylbutylthio, 1-ethylbutylthio, 2-ethylbutylthio, 1,1,2-trimethylpropylthio, 1,2,2-trimethylpropylthio, 1-ethyl-1-methylpropyl and 1-ethyl-2-methylpropyl.

Halogenated C₁-C₆-alkylthio is a radical of the formula R-S-, wherein R is a halogenated alkyl radical having from 1 to 6, preferably from 1 to 4 carbon atoms as defined herein. Examples include halogenomethylthio, dihalogenomethylthio, trihalogenomethylthio, (R)-1-halogenoethylthio, (S)-1-halogenoethylthio, 2-halogenoethylthio, 1,1-dihalogenoethylthio, 2,2-dihalogenoethylthio, 2,2,2-trihalogenoethylthio, (R)-1-halogenopropylthio, (S)-1-halogenopropylthio, 2-halogenopro-

pylthio, 3-halogenopropylthio, 1,1-dihalogenopropylthio, 2,2-dihalogenopropylthio, 3,3-dihalogenopropylthio, 3,3,3-trihalogenopropylthio, (R)-2-halogeno-1-methylethylthio, (S)-2-halogeno-1-methylethylthio, (R)-2,2-dihalogeno-1-methylethylthio, (S)-2,2-dihalogeno-1-methylethylthio, (R)-1,2-dihalogeno-1-methylethylthio, (S)-1,2-dihalogeno-1-methylethylthio, (R)-2,2,2-trihalogeno-1-methylethylthio, (S)-2,2,2-trihalogeno-1-methylethylthio, 2-halogeno-1-(halogenomethyl)ethylthio, 1-(dihalogenomethyl)-2,2-dihalogenoethylthio, (R)-1-halogenobutylthio, (S)-1-halogenobutylthio, 2-halogenobutylthio, 3-halogenobutylthio, 4-halogenobutylthio, 1,1-dihalogenobutylthio, 2,2-dihalogenobutylthio, 3,3-dihalogenobutylthio, 4,4-dihalogenobutylthio, 4,4,4-trihalogenobutylthio, etc. Particular examples include the fluorinated C₁-C₄ alkylthio groups as defined, such as trifluoromethylthio.

C₁-C₆-Alkylsulfinyl is a radical of the formula R-S(O)-, wherein R is an alkyl radical having from 1 to 6, preferably from 1 to 4 carbon atoms as defined herein. Examples include methylsulfinyl, ethylsulfinyl, propylsulfinyl, butylsulfinyl, pentylsulfinyl, 1-methylbutylsulfinyl, 2-methylbutylsulfinyl, 3-methylbutylsulfinyl, 2,2-dimethylpropylsulfinyl, 1-ethylpropylsulfinyl, hexylsulfinyl, 1,1-dimethylpropylsulfinyl, 1,2-dimethylpropylsulfinyl, 1-methylpentylsulfinyl, 2-methylpentylsulfinyl, 3-methylpentylsulfinyl, 4-methylpentylsulfinyl, 1,1-dimethylbutylsulfinyl, 1,2-dimethylbutylsulfinyl, 1,3-dimethylbutylsulfinyl, 2,2-dimethylbutylsulfinyl, 2,3-dimethylbutylsulfinyl, 3,3-dimethylbutylsulfinyl, 1-ethylbutylsulfinyl, 2-ethylbutylsulfinyl, 1,1,2-trimethylpropylsulfinyl, 1,2,2-trimethylpropylsulfinyl, 1-ethyl-1-methylpropyl and 1-ethyl-2-methylpropyl.

C₁-C₆-Alkylsulfonyl is a radical of the formula R-S(O)₂-, wherein R is an alkyl radical having from 1 to 6, preferably from 1 to 4 carbon atoms as defined herein. Examples include methylsulfonyl, ethylsulfonyl, propylsulfonyl, butylsulfonyl, pentylsulfonyl, 1-methylbutylsulfonyl, 2-methylbutylsulfonyl, 3-methylbutylsulfonyl, 2,2-dimethylpropylsulfonyl, 1-ethylpropylsulfonyl, hexylsulfonyl, 1,1-dimethylpropylsulfonyl, 1,2-dimethylpropylsulfonyl, 1-methylpentylsulfonyl, 2-methylpentylsulfonyl, 3-methylpentylsulfonyl, 4-methylpentylsulfonyl, 1,1-dimethylbutylsulfonyl, 1,2-dimethylbutylsulfonyl, 1,3-dimethylbutylsulfonyl, 2,2-dimethylbutylsulfonyl, 2,3-dimethylbutylsulfonyl, 3,3-dimethylbutylsulfonyl, 1-ethylbutylsulfonyl, 2-ethylbutylsulfonyl, 1,1,2-trimethylpropylsulfonyl, 1,2,2-trimethylpropylsulfonyl, 1-ethyl-1-methylpropyl and 1-ethyl-2-methylpropyl.

(Halogenated C₁-C₆-alkyl)sulfonyl is a C₁-C₆-alkylsulfonyl as defined herein, wherein at least one, e.g. 1, 2, 3, 4 or all of the hydrogen atoms are replaced by 1, 2, 3, 4 or a corresponding number of identical or different halogen atoms.

C₆-C₁₂-Arylsulfonyl is a radical of the formula R-S(O)₂-, wherein R is an aryl radical having from 6 to 12 carbon atoms as defined herein. Examples include phenylsulfonyl.

(C₆-C₁₂-Aryl-C₁-C₄-alkyl)sulfonyl is a radical of the formula R-S(O)₂·, wherein R is a C₆-C₁₂-aryl-C₁-C₄-alkyl radical, in particular a C₆-C₁₂-aryl-C₁-C₂-alkyl radical as defined herein. Examples include benzylsulfonyl.

- 5 M₃-M₁₂-Heterocyclylsulfonyl is a radical of the formula R-S(O)₂·, wherein R is M₃-M₁₂-heterocyclyl as defined herein.

Aminosulfonyl is NH₂-S(O)₂·.

- 10 C₁-C₆-Alkylaminosulfonyl is a radical of the formula R-NH-S(O)₂· wherein R is an alkyl radical having from 1 to 6, preferably from 1 to 4 carbon atoms as defined herein. Examples include methylaminosulfonyl, ethylaminosulfonyl, n-propylaminosulfonyl, iso-propylaminosulfonyl, n-butylaminosulfonyl, 2-butylaminosulfonyl, iso-butylaminosulfonyl, tert-butylaminosulfonyl.
- 15 Di-C₁-C₆-alkylaminosulfonyl is a radical of the formula RR'N-S(O)₂· wherein R and R' are independently of each other an alkyl radical having from 1 to 6, preferably from 1 to 4 carbon atoms as defined herein. Examples include dimethylaminosulfonyl, diethylaminosulfonyl, N-methyl-N-ethylaminosulfonyl.
- 20 C₆-C₁₂-Arylaminosulfonyl is a radical of the formula R-NH-S(O)₂· wherein R is an aryl radical having from 6 to 12, preferably 6 carbon atoms as defined herein.

Amino is NH₂.

- 25 C₁-C₆-Alkylamino is a radical of the formula R-NH· wherein R is an alkyl radical having from 1 to 6, in particular from 1 to 4 carbon atoms as defined herein. Examples include methylamino, ethylamino, n-propylamino, iso-propylamino, n-butylamino, 2-butylamino, iso-butylamino, tert-butylamino.
- 30 (Halogenated C₁-C₆-alkyl)amino is a C₁-C₆-alkylamino as defined herein, wherein at least one, e.g. 1, 2, 3, 4 or all of the hydrogen atoms are replaced by 1, 2, 3, 4 or a corresponding number of identical or different halogen atoms.

- Di-C₁-C₆-alkylamino is a radical of the formula RR'N· wherein R and R' are independently of each other an alkyl radical having from 1 to 6, in particular from 1 to 4 carbon atoms as defined herein. Examples include dimethylamino, diethylamino, N-methyl-N-ethylamino.
- 35

- Di-(halogenated C₁-C₆-alkyl)amino is a di-C₁-C₆-alkylamino as defined herein, wherein at least one, e.g. 1, 2, 3, 4 or all of the hydrogen atoms are replaced by 1, 2, 3, 4 or a corresponding number of identical or different halogen atoms.
- 40

C₁-C₆-Alkylcarbonylamino is a radical of the formula R-C(O)-NH-, wherein R is an alkyl radical having from 1 to 6, in particular from 1 to 4 carbon atoms as defined herein. Examples include acetamido (methylcarbonylamino), propionamido, n-butyramido, 2-methylpropionamido (isopropylcarbonylamino), 2,2-dimethylpropionamido and the like.

5

(Halogenated C₁-C₆-alkyl)carbonylamino is a C₁-C₆-alkylcarbonylamino as defined herein, wherein at least one, e.g. 1, 2, 3, 4 or all of the hydrogen atoms are replaced by 1, 2, 3, 4 or a corresponding number of identical or different halogen atoms.

- 10 C₆-C₁₂-Arylcarbonylamino is a radical of the formula R-C(O)-NH-, wherein R is an aryl radical having from 6 to 12 carbon atoms as defined herein. Examples include phenylcarbonylamino.

- 15 C₂-C₆-Alkenylamino is a radical of the formula R-NH-, wherein R is a straight-chain or branched alkenyl group having from 2 to 6, in particular 2 to 4 carbon atoms. Examples include vinylamino, allylamino (2-propen-1-ylamino), 1-propen-1-ylamino, 2-propen-2-ylamino, methallylamino (2-methylprop-2-en-1-ylamino) and the like. C₃-C₅-Alkenylamino is, in particular, allylamino, 1-methylprop-2-en-1-ylamino, 2-buten-1-ylamino, 3-buten-1-ylamino, methallylamino, 2-penten-1-ylamino, 3-penten-1-ylamino, 4-penten-1-ylamino, 1-methylbut-2-en-1-ylamino or 2-ethylprop-2-en-1-ylamino.

20

C₁-C₆-Alkylsulfonylamino is a radical of the formula R-S(O)₂-NH-, wherein R is an alkyl radical having from 1 to 6, in particular from 1 to 4 carbon atoms as defined herein. Examples include methylsulfonylamino, ethylsulfonylamino, n-propylsulfonylamino, iso-propylsulfonylamino, n-butylsulfonylamino, 2-butylsulfonylamino, iso-butylsulfonylamino, tert-butylsulfonylamino.

25

(Halogenated C₁-C₆ alkyl)sulfonylamino is a C₁-C₆-alkylsulfonylamino as defined herein, wherein at least one, e.g. 1, 2, 3, 4 or all of the hydrogen atoms are replaced by 1, 2, 3, 4 or a corresponding number of identical or different halogen atoms.

- 30 C₆-C₁₂-Arylsulfonylamino is a radical of the formula R-S(O)₂-NH-, wherein R is an aryl radical having from 6 to 12 carbon atoms as defined herein. Examples include phenylsulfonylamino.

Nitro is -NO₂.

- 35 M₃-M₁₂-Heterocyclyl is a 3- to 12-membered heterocyclic radical including a saturated heterocyclic radical, which generally has 3, 4, 5, 6, or 7 ring forming atoms (ring members), an unsaturated non-aromatic heterocyclic radical, which generally has 5, 6 or 7 ring forming atoms, and a heteroaromatic radical (hetaryl), which generally has 5, 6 or 7 ring forming atoms. The heterocyclic radicals may be bound via a carbon atom (C-bound) or a nitrogen atom (N-bound). Preferred heterocyclic radicals comprise 1 nitrogen atom as ring member atom and optionally 1, 2 or 3 further heteroatoms as ring members, which are selected, independently of each other from O, S and N. Likewise
- 40

preferred heterocyclic radicals comprise 1 heteroatom as ring member, which is selected from O, S and N, and optionally 1, 2 or 3 further nitrogen atoms as ring members.

Examples of M₃-M₁₂-heterocyclyl include:

- 5 C- or N-bound 3-4-membered, saturated rings, such as
2-oxiranyl, 2-oxetanyl, 3-oxetanyl, 2-aziridinyl, 3-thiethanyl, 1-azetidiny, 2-azetidiny, 3-azetidiny;
- 10 C-bound, 5-membered, saturated rings, such as
tetrahydrofuran-2-yl, tetrahydrofuran-3-yl, tetrahydrothien-2-yl, tetrahydrothien-3-yl, tetrahydro-
pyrrol-2-yl, tetrahydropyrrol-3-yl, tetrahydropyrazol-3-yl, tetrahydro-pyrazol-4-yl, tetrahydroisox-
azol-3-yl, tetrahydroisoxazol-4-yl, tetrahydroisoxazol-5-yl, 1,2-oxathiolan-3-yl, 1,2-oxathiolan-4-
yl, 1,2-oxathiolan-5-yl, tetrahydroisothiazol-3-yl, tetrahydroisothiazol-4-yl, tetrahydroisothiazol-5-
15 yl, 1,2-dithiolan-3-yl, 1,2-dithiolan-4-yl, tetrahydroimidazol-2-yl, tetrahydroimidazol-4-yl, tetrahy-
drooxazol-2-yl, tetrahydrooxazol-4-yl, tetrahydrooxazol-5-yl, tetrahydrothiazol-2-yl, tetrahydrothi-
azol-4-yl, tetrahydrothiazol-5-yl, 1,3-dioxolan-2-yl, 1,3-dioxolan-4-yl, 1,3-oxathiolan-2-yl, 1,3-
oxathiolan-4-yl, 1,3-oxathiolan-5-yl, 1,3-dithiolan-2-yl, 1,3-dithiolan-4-yl, 1,3,2-dioxathiolan-4-yl;
- 20 C-bound, 6-membered, saturated rings, such as
tetrahydropyran-2-yl, tetrahydropyran-3-yl, tetrahydropyran-4-yl, piperidin-2-yl, piperidin-3-yl,
piperidin-4-yl, tetrahydrothiopyran-2-yl, tetrahydrothiopyran-3-yl, tetrahydrothiopyran-4-yl, 1,3-
dioxan-2-yl, 1,3-dioxan-4-yl, 1,3-dioxan-5-yl, 1,4-dioxan-2-yl, 1,3-dithian-2-yl, 1,3-dithian-4-yl,
1,3-dithian-5-yl, 1,4-dithian-2-yl, 1,3-oxathian-2-yl, 1,3-oxathian-4-yl, 1,3-oxathian-5-yl, 1,3-
25 oxathian-6-yl, 1,4-oxathian-2-yl, 1,4-oxathian-3-yl, 1,2-dithian-3-yl, 1,2-dithian-4-yl, hexahydro-
pyrimidin-2-yl, hexahydropyrimidin-4-yl, hexahydropyrimidin-5-yl, hexahydropyrazin-2-yl, hexa-
hydropyridazin-3-yl, hexahydropyridazin-4-yl, tetrahydro-1,3-oxazin-2-yl, tetrahydro-1,3-oxazin-
4-yl, tetrahydro-1,3-oxazin-5-yl, tetrahydro-1,3-oxazin-6-yl, tetrahydro-1,3-thiazin-2-yl, tetrahy-
dro-1,3-thiazin-4-yl, tetrahydro-1,3-thiazin-5-yl, tetrahydro-1,3-thiazin-6-yl, tetrahydro-1,4-thiazin-
30 2-yl, tetrahydro-1,4-thiazin-3-yl, tetrahydro-1,4-oxazin-2-yl, tetrahydro-1,4-oxazin-3-yl, tetrahy-
dro-1,2-oxazin-3-yl, tetrahydro-1,2-oxazin-4-yl, tetrahydro-1,2-oxazin-5-yl, tetrahydro-1,2-oxazin-
6-yl;
- N-bound, 5-membered, saturated rings, such as
35 tetrahydropyrrol-1-yl (pyrrolidin-1-yl), tetrahydropyrazol-1-yl, tetrahydroisoxazol-2-yl, tetrahy-
droisothiazol-2-yl, tetrahydroimidazol-1-yl, tetrahydrooxazol-3-yl, tetrahydrothiazol-3-yl;
- N-bound, 6-membered, saturated rings, such as
piperidin-1-yl, hexahydropyrimidin-1-yl, hexahydropyrazin-1-yl (piperazin-1-yl), hexahydro-
pyridazin-1-yl, tetrahydro-1,3-oxazin-3-yl, tetrahydro-1,3-thiazin-3-yl, tetrahydro-1,4-thiazin-4-yl,
40 tetrahydro-1,4-oxazin-4-yl (morpholin-1-yl), tetrahydro-1,2-oxazin-2-yl;

- C-bound, 5-membered, partially unsaturated rings, such as
- 2,3-dihydrofuran-2-yl, 2,3-dihydrofuran-3-yl, 2,5-dihydrofuran-2-yl, 2,5-dihydrofuran-3-yl, 4,5-dihydrofuran-2-yl, 4,5-dihydrofuran-3-yl, 2,3-dihydrothien-2-yl, 2,3-dihydrothien-3-yl, 2,5-dihydrothien-2-yl, 2,5-dihydrothien-3-yl, 4,5-dihydrothien-2-yl, 4,5-dihydrothien-3-yl, 2,3-dihydro-1H-pyrrol-2-yl, 2,3-dihydro-1H-pyrrol-3-yl, 2,5-dihydro-1H-pyrrol-2-yl, 2,5-dihydro-1H-pyrrol-3-yl, 4,5-dihydro-1H-pyrrol-2-yl, 4,5-dihydro-1H-pyrrol-3-yl, 3,4-dihydro-2H-pyrrol-2-yl, 3,4-dihydro-2H-pyrrol-3-yl, 3,4-dihydro-5H-pyrrol-2-yl, 3,4-dihydro-5H-pyrrol-3-yl, 4,5-dihydro-1H-pyrazol-3-yl, 4,5-dihydro-1H-pyrazol-4-yl, 4,5-dihydro-1H-pyrazol-5-yl, 2,5-dihydro-1H-pyrazol-3-yl, 2,5-dihydro-1H-pyrazol-4-yl, 2,5-dihydro-1H-pyrazol-5-yl, 4,5-dihydroisoxazol-3-yl, 4,5-dihydroisoxazol-4-yl, 4,5-dihydroisoxazol-5-yl, 2,5-dihydroisoxazol-3-yl, 2,5-dihydroisoxazol-4-yl, 2,5-dihydroisoxazol-5-yl, 2,3-dihydroisoxazol-3-yl, 2,3-dihydroisoxazol-4-yl, 2,3-dihydroisoxazol-5-yl, 4,5-dihydroisothiazol-3-yl, 4,5-dihydroisothiazol-4-yl, 4,5-dihydroisothiazol-5-yl, 2,5-dihydroisothiazol-3-yl, 2,5-dihydroisothiazol-4-yl, 2,5-dihydroisothiazol-5-yl, 2,3-dihydroisothiazol-3-yl, 2,3-dihydroisothiazol-4-yl, 2,3-dihydroisothiazol-5-yl, 4,5-dihydro-1H-imidazol-2-yl, 4,5-dihydro-1H-imidazol-4-yl, 4,5-dihydro-1H-imidazol-5-yl, 2,5-dihydro-1H-imidazol-2-yl, 2,5-dihydro-1H-imidazol-4-yl, 2,5-dihydro-1H-imidazol-5-yl, 2,3-dihydro-1H-imidazol-2-yl, 2,3-dihydro-1H-imidazol-4-yl, 4,5-dihydro-oxazol-2-yl, 4,5-dihydrooxazol-4-yl, 4,5-dihydrooxazol-5-yl, 2,5-dihydrooxazol-2-yl, 2,5-dihydrooxazol-4-yl, 2,5-dihydrooxazol-5-yl, 2,3-dihydrooxazol-2-yl, 2,3-dihydrooxazol-4-yl, 2,3-dihydrooxazol-5-yl, 4,5-dihydrothiazol-2-yl, 4,5-dihydrothiazol-4-yl, 4,5-dihydrothiazol-5-yl, 2,5-dihydrothiazol-2-yl, 2,5-dihydrothiazol-4-yl, 2,5-dihydrothiazol-5-yl, 2,3-dihydrothiazol-2-yl, 2,3-dihydrothiazol-4-yl, 2,3-dihydrothiazol-5-yl, 1,3-dioxol-2-yl, 1,3-dioxol-4-yl, 1,3-dithiol-2-yl, 1,3-dithiol-4-yl, 1,3-oxathiol-2-yl, 1,3-oxathiol-4-yl, 1,3-oxathiol-5-yl;
- 25 C-bound, 6-membered, partially unsaturated rings, such as
- 2H-3,4-dihydropyran-6-yl, 2H-3,4-dihydropyran-5-yl, 2H-3,4-dihydropyran-4-yl, 2H-3,4-dihydropyran-3-yl, 2H-3,4-dihydropyran-2-yl, 2H-3,4-dihydrothiopyran-6-yl, 2H-3,4-dihydrothiopyran-5-yl, 2H-3,4-dihydrothiopyran-4-yl, 2H-3,4-dihydrothiopyran-3-yl, 2H-3,4-dihydrothiopyran-2-yl, 1,2,3,4-tetrahydropyridin-6-yl, 1,2,3,4-tetrahydropyridin-5-yl, 1,2,3,4-tetrahydropyridin-4-yl, 1,2,3,4-tetrahydropyridin-3-yl, 1,2,3,4-tetrahydropyridin-2-yl, 2H-5,6-dihydropyran-2-yl, 2H-5,6-dihydropyran-3-yl, 2H-5,6-dihydropyran-4-yl, 2H-5,6-dihydropyran-5-yl, 2H-5,6-dihydropyran-6-yl, 2H-5,6-dihydrothiopyran-2-yl, 2H-5,6-dihydrothiopyran-3-yl, 2H-5,6-dihydrothiopyran-4-yl, 2H-5,6-dihydrothiopyran-5-yl, 2H-5,6-dihydrothiopyran-6-yl, 1,2,5,6-tetrahydropyridin-2-yl, 1,2,5,6-tetrahydropyridin-3-yl, 1,2,5,6-tetrahydropyridin-4-yl, 1,2,5,6-tetrahydropyridin-5-yl, 1,2,5,6-tetrahydropyridin-6-yl, 2,3,4,5-tetrahydropyridin-2-yl, 2,3,4,5-tetrahydropyridin-3-yl, 2,3,4,5-tetrahydropyridin-4-yl, 2,3,4,5-tetrahydropyridin-5-yl, 2,3,4,5-tetrahydropyridin-6-yl, 4H-pyran-2-yl, 4H-pyran-3-yl, 4H-pyran-4-yl, 4H-thiopyran-2-yl, 4H-thiopyran-3-yl, 4H-thiopyran-4-yl, 1,4-dihydropyridin-2-yl, 1,4-dihydropyridin-3-yl, 1,4-dihydropyridin-4-yl, 2H-pyran-2-yl, 2H-pyran-3-yl, 2H-pyran-4-yl, 2H-pyran-5-yl, 2H-pyran-6-yl, 2H-thiopyran-2-yl, 2H-thiopyran-3-yl, 2H-thiopyran-4-yl, 2H-thiopyran-5-yl, 2H-thiopyran-6-yl, 1,2-dihydropyridin-2-yl, 1,2-dihydro-pyridin-3-yl, 1,2-dihydropyridin-4-yl, 1,2-dihydropyridin-5-yl, 1,2-dihydro-pyridin-6-yl, 3,4-dihydropyridin-2-yl, 3,4-dihydropyridin-3-yl, 3,4-dihydro-pyridin-

4-yl, 3,4-dihydropyridin-5-yl, 3,4-dihydropyridin-6-yl, 2,5-dihydropyridin-2-yl, 2,5-dihydropyridin-3-yl, 2,5-dihydropyridin-4-yl, 2,5-dihydropyridin-5-yl, 2,5-dihydropyridin-6-yl, 2,3-dihydropyridin-2-yl, 2,3-dihydropyridin-3-yl, 2,3-dihydropyridin-4-yl, 2,3-dihydropyridin-5-yl, 2,3-dihydropyridin-6-yl, 2H-5,6-dihydro-1,2-oxazin-3-yl, 2H-5,6-dihydro-1,2-oxazin-4-yl, 2H-5,6-dihydro-1,2-oxazin-5-yl, 2H-5,6-dihydro-1,2-oxazin-6-yl, 2H-5,6-dihydro-1,2-thiazin-3-yl, 2H-5,6-dihydro-1,2-thiazin-4-yl, 2H-5,6-dihydro-1,2-thiazin-5-yl, 2H-5,6-dihydro-1,2-thiazin-6-yl, 4H-5,6-dihydro-1,2-oxazin-3-yl, 4H-5,6-dihydro-1,2-oxazin-4-yl, 4H-5,6-dihydro-1,2-oxazin-5-yl, 4H-5,6-dihydro-1,2-oxazin-6-yl, 4H-5,6-dihydro-1,2-thiazin-3-yl, 4H-5,6-dihydro-1,2-thiazin-4-yl, 4H-5,6-dihydro-1,2-thiazin-5-yl, 4H-5,6-dihydro-1,2-thiazin-6-yl, 2H-3,6-dihydro-1,2-oxazin-3-yl, 2H-3,6-dihydro-1,2-oxazin-4-yl, 2H-3,6-dihydro-1,2-oxazin-5-yl, 2H-3,6-dihydro-1,2-oxazin-6-yl, 2H-3,6-dihydro-1,2-thiazin-3-yl, 2H-3,6-dihydro-1,2-thiazin-4-yl, 2H-3,6-dihydro-1,2-thiazin-5-yl, 2H-3,6-dihydro-1,2-thiazin-6-yl, 2H-3,4-dihydro-1,2-oxazin-3-yl, 2H-3,4-dihydro-1,2-oxazin-4-yl, 2H-3,4-dihydro-1,2-oxazin-5-yl, 2H-3,4-dihydro-1,2-oxazin-6-yl, 2H-3,4-dihydro-1,2-thiazin-3-yl, 2H-3,4-dihydro-1,2-thiazin-4-yl, 2H-3,4-dihydro-1,2-thiazin-5-yl, 2H-3,4-dihydro-1,2-thiazin-6-yl, 2,3,4,5-tetrahydropyridazin-3-yl, 2,3,4,5-tetrahydropyridazin-4-yl, 2,3,4,5-tetrahydropyridazin-5-yl, 2,3,4,5-tetrahydropyridazin-6-yl, 3,4,5,6-tetrahydropyridazin-3-yl, 3,4,5,6-tetrahydropyridazin-4-yl, 1,2,5,6-tetrahydropyridazin-3-yl, 1,2,5,6-tetrahydropyridazin-4-yl, 1,2,5,6-tetrahydropyridazin-5-yl, 1,2,5,6-tetrahydropyridazin-6-yl, 1,2,3,6-tetrahydropyridazin-3-yl, 1,2,3,6-tetrahydropyridazin-4-yl, 4H-5,6-dihydro-1,3-oxazin-2-yl, 4H-5,6-dihydro-1,3-oxazin-4-yl, 4H-5,6-dihydro-1,3-oxazin-5-yl, 4H-5,6-dihydro-1,3-oxazin-6-yl, 4H-5,6-dihydro-1,3-thiazin-2-yl, 4H-5,6-dihydro-1,3-thiazin-4-yl, 4H-5,6-dihydro-1,3-thiazin-5-yl, 4H-5,6-dihydro-1,3-thiazin-6-yl, 3,4,5,6-tetrahydropyrimidin-2-yl, 3,4,5,6-tetrahydropyrimidin-4-yl, 3,4,5,6-tetrahydropyrimidin-5-yl, 3,4,5,6-tetrahydropyrimidin-6-yl, 1,2,3,4-tetrahydropyrazin-2-yl, 1,2,3,4-tetrahydropyrazin-5-yl, 1,2,3,4-tetrahydropyrimidin-2-yl, 1,2,3,4-tetrahydropyrimidin-4-yl, 1,2,3,4-tetrahydropyrimidin-5-yl, 1,2,3,4-tetrahydropyrimidin-6-yl, 2,3-dihydro-1,4-thiazin-2-yl, 2,3-dihydro-1,4-thiazin-3-yl, 2,3-dihydro-1,4-thiazin-5-yl, 2,3-dihydro-1,4-thiazin-6-yl, 2H-1,3-oxazin-2-yl, 2H-1,3-oxazin-4-yl, 2H-1,3-oxazin-5-yl, 2H-1,3-oxazin-6-yl, 2H-1,3-thiazin-2-yl, 2H-1,3-thiazin-4-yl, 2H-1,3-thiazin-5-yl, 2H-1,3-thiazin-6-yl, 4H-1,3-oxazin-2-yl, 4H-1,3-oxazin-4-yl, 4H-1,3-oxazin-5-yl, 4H-1,3-oxazin-6-yl, 4H-1,3-thiazin-2-yl, 4H-1,3-thiazin-4-yl, 4H-1,3-thiazin-5-yl, 4H-1,3-thiazin-6-yl, 6H-1,3-oxazin-2-yl, 6H-1,3-oxazin-4-yl, 6H-1,3-oxazin-5-yl, 6H-1,3-oxazin-6-yl, 6H-1,3-thiazin-2-yl, 6H-1,3-oxazin-4-yl, 6H-1,3-oxazin-5-yl, 6H-1,3-thiazin-6-yl, 2H-1,4-oxazin-2-yl, 2H-1,4-oxazin-3-yl, 2H-1,4-oxazin-5-yl, 2H-1,4-oxazin-6-yl, 2H-1,4-thiazin-2-yl, 2H-1,4-thiazin-3-yl, 2H-1,4-thiazin-5-yl, 2H-1,4-thiazin-6-yl, 4H-1,4-oxazin-2-yl, 4H-1,4-oxazin-3-yl, 4H-1,4-thiazin-2-yl, 4H-1,4-thiazin-3-yl, 1,4-dihydropyridazin-3-yl, 1,4-dihydropyridazin-4-yl, 1,4-dihydropyridazin-5-yl, 1,4-dihydropyridazin-6-yl, 1,4-dihydropyrazin-2-yl, 1,2-dihydropyrazin-2-yl, 1,2-dihydropyrazin-3-yl, 1,2-dihydropyrazin-5-yl, 1,2-dihydropyrazin-6-yl, 1,4-dihydropyrimidin-2-yl, 1,4-dihydropyrimidin-4-yl, 1,4-dihydropyrimidin-5-yl, 1,4-dihydropyrimidin-6-yl, 3,4-dihydropyrimidin-2-yl, 3,4-dihydropyrimidin-4-yl, 3,4-dihydropyrimidin-5-yl or 3,4-dihydropyrimidin-6-yl;

40

N-bound, 5-membered, partially unsaturated rings, such as

2,3-dihydro-1H-pyrrol-1-yl, 2,5-dihydro-1H-pyrrol-1-yl, 4,5-dihydro-1H-pyrazol-1-yl, 2,5-dihydro-1H-pyrazol-1-yl, 2,3-dihydro-1H-pyrazol-1-yl, 2,5-dihydroisoxazol-2-yl, 2,3-dihydroisoxazol-2-yl, 2,5-dihydroisothiazol-2-yl, 2,3-dihydroisoxazol-2-yl, 4,5-dihydro-1H-imidazol-1-yl, 2,5-dihydro-1H-imidazol-1-yl, 2,3-dihydro-1H-imidazol-1-yl, 2,3-dihydrooxazol-3-yl, 2,3-dihydrothiazol-3-yl;

N-bound, 6-membered, partially unsaturated rings, such as
1,2,3,4-tetrahydropyridin-1-yl, 1,2,5,6-tetrahydropyridin-1-yl, 1,4-dihydro-pyridin-1-yl, 1,2-dihydropyridin-1-yl, 2H-5,6-dihydro-1,2-oxazin-2-yl, 2H-5,6-dihydro-1,2-thiazin-2-yl, 2H-3,6-dihydro-1,2-oxazin-2-yl, 2H-3,6-dihydro-1,2-thiazin-2-yl, 2H-3,4-dihydro-1,2-oxazin-2-yl, 2H-3,4-dihydro-1,2-thiazin-2-yl, 2,3,4,5-tetrahydropyridazin-2-yl, 1,2,5,6-tetrahydropyridazin-1-yl, 1,2,5,6-tetrahydropyridazin-2-yl, 1,2,3,6-tetrahydropyridazin-1-yl, 3,4,5,6-tetrahydropyrimidin-3-yl, 1,2,3,4-tetrahydropyrazin-1-yl, 1,2,3,4-tetrahydropyrimidin-1-yl, 1,2,3,4-tetrahydropyrimidin-3-yl, 2,3-dihydro-1,4-thiazin-4-yl, 2H-1,2-oxazin-2-yl, 2H-1,2-thiazin-2-yl, 4H-1,4-oxazin-4-yl, 4H-1,4-thiazin-4-yl, 1,4-dihydropyridazin-1-yl, 1,4-dihydropyrazin-1-yl, 1,2-dihydropyrazin-1-yl, 1,4-dihydropyrimidin-1-yl or 3,4-dihydropyrimidin-3-yl;

C-bound, 5-membered, heteroaromatic rings, such as
2-furyl, 3-furyl, 2-thienyl, 3-thienyl, pyrrol-2-yl, pyrrol-3-yl, pyrazol-3-yl, pyrazol-4-yl, isoxazol-3-yl, isoxazol-4-yl, isoxazol-5-yl, isothiazol-3-yl, isothiazol-4-yl, isothiazol-5-yl, imidazol-2-yl, imidazol-4-yl, oxazol-2-yl, oxazol-4-yl, oxazol-5-yl, thiazol-2-yl, thiazol-4-yl, thiazol-5-yl, 1,2,3-oxadiazol-4-yl, 1,2,3-oxadiazol-5-yl, 1,2,4-oxadiazol-3-yl, 1,2,4-oxadiazol-5-yl, 1,3,4-oxadiazol-2-yl, 1,2,3-thiadiazol-4-yl, 1,2,3-thiadiazol-5-yl, 1,2,4-thiadiazol-3-yl, 1,2,4-thiadiazol-5-yl, 1,3,4-thiadiazolyl-2-yl, 1,2,3-triazol-4-yl, 1,2,4-triazol-3-yl, tetrazol-5-yl;


C-bound, 6-membered, heteroaromatic rings, such as
pyridin-2-yl, pyridin-3-yl, pyridin-4-yl (4-pyridyl), pyridazin-3-yl, pyridazin-4-yl, pyrimidin-2-yl, pyrimidin-4-yl, pyrimidin-5-yl, pyrazin-2-yl, 1,3,5-triazin-2-yl, 1,2,4-triazin-3-yl, 1,2,4-triazin-5-yl, 1,2,4-triazin-6-yl, 1,2,4,5-tetrazin-3-yl;

N-bound, 5-membered, heteroaromatic rings, such as
pyrrol-1-yl, pyrazol-1-yl, imidazol-1-yl, 1,2,3-triazol-1-yl, 1,2,4-triazol-1-yl, tetrazol-1-yl.


Heterocyclyl also includes bicyclic heterocycles, which comprise one of the described 5- or 6-membered heterocyclic rings and a further anellated, saturated or unsaturated or aromatic carbocycle, such as a benzene, cyclohexane, cyclohexene or cyclohexadiene ring, or a further anellated 5- or 6-membered heterocyclic ring, this heterocyclic ring being saturated or unsaturated or aromatic. These include quinolinyl, isoquinolinyl, indolyl, indoliziny, isoindolyl, indazolyl, benzofuryl, benzthienyl, benzo[b]thiazolyl, benzoxazolyl, benzodioxol, benzthiazolyl and benzimidazolyl. Examples of 5- or 6-membered heteroaromatic compounds comprising an anellated cycloalkenyl ring include dihydroindolyl, dihydroindoliziny, dihydroisoindolyl, dihydroquinolinyl, dihydroisoquinolinyl, chromenyl and chromanyl.

M_3 - M_{12} -Heteroarylene is a heteroaryl diradical. Examples include pyrid-2,5-ylene and pyrid-2,4-ylene.


- 5 With respect to the compounds' capability of inhibiting glycine transporter 1, the variables R^1 , W, A^1 , Q, Y, R^6 , n1, n2, X^1 , A, R^2 , A^2 , R^3 , Y^1 , R^{4a} , R^{4b} , X^2 , X^3 , R^5 have in particular the following meanings which, when taken alone or in combination, represent particular embodiments of the compounds of the formula (I) or any other formula disclosed herein.
- 10 In said formula (I), there may be one or more than one substituent R^2 , R^3 , R^6 and one or more than one substituent



- 15 More particularly, there may be up to 3 substituents R^2 , up to 4 substituents R^3 , and up to 6 substituents R^6 . Preferably, there is one substituent



and 1, 2 or 3 substituents R^2 . Formula (I) may thus be depicted as follows:




20

wherein A, R^1 , W, A^1 , Q, Y, R^6 , n1, n2, X^1 , R^2 , A^2 , R^3 , Y^1 , R^{4a} , R^{4b} , X^2 , X^3 and R^5 are as defined herein, a is 1, 2 or 3, b is 1, 2, 3 or 4, c is 1 and d is 1, 2, 3, 4, 5 or 6. If there is more than one radical R^2 , these may be the same or different radicals. If there is more than one radical R^3 , these may be the same or different radicals. If there is more than one radical R^6 , these may be the same or


25 different radicals. According to one embodiment, a is 1, b is 1 or 2, c is 1, and d is 1 or 2.


In the following the radical



is also referred to as R.


- 5 A is a 5- or 6-membered ring which includes two carbon atoms from the tetrahydropyran, tetrahydrothiopyrane and tetrahydropyridine moiety to which A is fused. A may be a homocyclic or heterocyclic ring. The ring may be saturated, unsaturated non-aromatic or aromatic. According to a particular embodiment, A is a benzene ring. As a heterocyclic ring, A may include 1, 2 or 3 heteroatoms as ring member atoms, which are selected, independently of each other from N, S and O. Preferred heterocyclic rings comprise 1 nitrogen atom as ring member atom and optionally 1 or 2 further heteroatoms as ring members, which are selected, independently of each other from O, S and N. Likewise preferred heterocyclic rings comprise 1 heteroatom as ring member atom, which is selected from O, S and N, and optionally 1 or 2 further nitrogen atoms as ring member atoms. According to a particular embodiment, A is a heterocyclic ring selected from the group consisting of
- 10 the following 5- or 6-membered heterocyclic rings:
- 15






In said formulae, hydrogen atoms are not depicted. This is meant to illustrate that the free valency
 5 of a carbon or nitrogen atom may be either bound to a hydrogen atom, to R or to R^2 . Accordingly, R and R^2 may be C- or N-bound at any position of ring A.

The skilled person will appreciate that some of the rings depicted above may be represented with a
 different structure, e.g. with hydrogen atoms having other positions than those shown above, for
 10 instance as given in the following structures:




Preferably, A is a heterocyclic ring selected from the group consisting of the following 5- or 6-
 15 membered heterocyclic rings:



If ring A is a 5-membered heterocyclic ring it is preferred that R is bound to G^1 or G^2 , in particular G^2 :


5



In said formula, G^1 , G^2 and G^3 independently are $-\text{CH}=\text{}$, $-\text{CH}_2-$, $-\text{N}=\text{}$, $-\text{NH}-$, S or O, at least one of G^1 , G^2 and G^3 is $-\text{CH}=\text{}$ or $-\text{CH}_2-$, the dotted line represents a single or a double bond and A^2 , R^3 , Y^1 , R^{4a} , R^{4b} , X^2 , X^3 , R^5 are as defined herein.

10


If ring A is 6-membered heterocyclic ring it is preferred that R is bound to G^1 or G^2 , in particular G^2 :




15

In said formula, G^1 , G^2 , G^3 and G^4 independently are $-\text{CH}=\text{}$, $-\text{CH}_2-$, $-\text{N}=\text{}$, $-\text{NH}-$, S or O, at least one of G^1 , G^2 , G^3 and G^4 is $-\text{CH}=\text{}$ or $-\text{CH}_2-$, the dotted line represents a single or a double bond and A^2 , R^3 , Y^1 , R^{4a} , R^{4b} , X^2 , X^3 , R^5 are as defined herein.

Heterocyclic compounds having the following partial structures are preferred:



Heterocyclic compounds having the following partial structures are particularly preferred:



In said formulae, R and R² are as defined herein. If there is more than one radical R², these may be the same or different radicals.

5

- R¹ is hydrogen, C₁-C₆-alkyl (e.g. methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl or n-pentyl), C₃-C₁₂-cycloalkyl-C₁-C₄-alkyl (e.g. cyclopropylmethyl, cyclopentylmethyl or cyclohexylmethyl), halogenated C₁-C₆-alkyl (e.g. 3-fluoroprop-1-yl, 3-chloroprop-1-yl or 3,3,3-trifluoroprop-1-yl), tri-(C₁-C₄-alkyl)-silyl-C₁-C₄-alkyl (e.g. trimethylsilylethyl), hydroxy-C₁-C₄-alkyl, C₁-C₆-alkoxy-C₁-C₄-alkyl (e.g. ethoxyethyl), amino-C₁-C₄-alkyl, C₁-C₆-alkylamino-C₁-C₄-alkyl, di-C₁-C₆-alkylamino-C₁-C₄-alkyl, C₁-C₆-alkylcarbonylamino-C₁-C₄-alkyl, C₁-C₆-alkyloxycarbonylamino-C₁-C₄-alkyl, C₁-C₆-alkylaminocarbonylamino-C₁-C₄-alkyl, di-C₁-C₆-alkylaminocarbonylamino-C₁-C₄-alkyl, C₁-C₆-alkylsulfonylamino-C₁-C₄-alkyl, (optionally substituted C₆-C₁₂-aryl-C₁-C₆-alkyl)amino-C₁-C₄-alkyl, optionally substituted C₆-C₁₂-aryl-C₁-C₄-alkyl, optionally substituted M₃-M₁₂-heterocyclyl-C₁-C₄-alkyl, C₃-C₁₂-cycloalkyl (e.g. cyclopropyl or cyclobutyl), C₁-C₆-alkylcarbonyl, C₁-C₆-alkoxycarbonyl, halogenated C₁-C₆-alkoxycarbonyl, C₆-C₁₂-aryloxycarbonyl, aminocarbonyl, C₁-C₆-alkylaminocarbonyl, (halogenated C₁-C₄-alkyl)aminocarbonyl, C₆-C₁₂-arylaminocarbonyl, C₂-C₆-alkenyl (e.g. prop-1,2-en-1-yl), C₂-C₆-alkynyl, optionally substituted C₆-C₁₂-aryl (e.g. phenyl, 2-methylphenyl), hydroxy, C₁-C₆-alkoxy (e.g. tert-butyloxy), halogenated C₁-C₆-alkoxy, C₁-C₆-hydroxyalkoxy, C₁-C₆-alkoxy-C₁-C₄-alkoxy, amino-C₁-C₄-alkoxy, C₁-C₆-alkylamino-C₁-C₄-alkoxy, di-C₁-C₆-alkylamino-C₁-C₄-alkoxy, C₁-C₆-alkylcarbonylamino-C₁-C₄-alkoxy, C₆-C₁₂-arylcarbonylamino-C₁-C₄-alkoxy, C₁-C₆-alkoxycarbonylamino-C₁-C₄-alkoxy, C₆-C₁₂-aryl-C₁-C₄-alkoxy, C₁-C₆-alkylsulfonylamino-C₁-C₄-alkoxy, (halogenated C₁-C₆-alkyl)sulfonylamino-C₁-C₄-alkoxy, C₆-C₁₂-arylsulfonylamino-C₁-C₄-alkoxy, (C₆-C₁₂-aryl-C₁-C₆-alkyl)sulfonylamino-C₁-C₄-alkoxy, M₃-M₁₂-heterocyclylsulfonylamino-C₁-C₄-alkoxy, M₃-M₁₂-heterocyclyl-C₁-C₄-alkoxy, C₆-C₁₂-aryloxy, M₃-M₁₂-heterocyclyloxy, C₁-C₆-alkylthio, halogenated C₁-C₆-alkylthio, C₁-C₆-alkylamino, (halogenated C₁-C₆-alkyl)amino, di-C₁-C₆-alkylamino (e.g. dimethylamino), di-(halogenated C₁-C₆-alkyl)amino, C₁-C₆-alkylcarbonylamino, (halogenated C₁-C₆-alkyl)carbonylamino, C₆-C₁₂-arylcarbonylamino, C₁-C₆-alkylsulfonylamino, (halogenated C₁-C₆-alkyl)sulfonylamino, C₆-C₁₂-arylsulfonylamino or optionally substituted M₃-M₁₂-heterocyclyl (e.g. 3-pyridyl, 2-pyridyl, 2-thienyl, 4-methyl-2-thienyl, 5-methyl-2-thienyl, 5-chloro-2-thienyl, 2,5-dimethyl-3-thienyl, 1,2-diazol-4-yl, 1-methyl-1,2-diazol-4-yl, 1,3-dimethyl-1,2-diazol-4-yl, 1,1-ethyl-1,2-diazol-4-yl, 1-difluormethyl-1,2-diazol-4-yl, 2-methyl-1,3-diazol-4-yl, 1-methyl-1,3-diazol-4-yl, 2-methyl-1,3-thiazol-5-yl, 2,4-dimethyl-1,3-thiazol-5-yl, 3-pyrrolidinyl, 1-methyl-pyrrol-3-yl, 2-pyridyl, 1-methyl-1,2-diazol-3-yl, 1-methyl-3-trifluoromethyl-1,2-diazol-4-yl, 1,2-dimethyl-1,3-diazol-4-yl, 5-methylisoxazol-3-yl, 1-methyl-1,2,4-triazol-3-yl, 1-methyl-1,2,3-triazol-4-yl, 1-ethyl-1,2,3-triazol-4-yl, furan-3-yl, 5-methyl-furan-2-yl, 2,5-dimethyl-furan-3-yl, 3-

methyl-piperidinyl, thiophen-2-yl, 4-methyl-thiophen-2-yl, 5-methyl-thiophen-2-yl, thiophen-3-yl, or morpholin-4-yl).

Preferably, R¹ is C₁-C₆-alkyl (e.g. methyl, ethyl, n-propyl, isopropyl, sec-butyl, n-butyl or n-pentyl), C₃-C₁₂-cycloalkyl-C₁-C₄-alkyl (e.g. cyclopropylmethyl, cyclopentylmethyl or cyclohexylmethyl), halogenated C₁-C₆-alkyl (e.g. 3-fluoroprop-1-yl, 3-chloroprop-1-yl or 3,3,3-trifluoroprop-1-yl), tri-(C₁-C₄-alkyl)-silyl-C₁-C₄-alkyl (e.g. trimethylsilylethyl), C₁-C₆-alkoxy-C₁-C₄-alkyl (e.g. ethoxyethyl), amino-C₁-C₄-alkyl, C₁-C₆-alkylamino-C₁-C₄-alkyl, di-C₁-C₆-alkylamino-C₁-C₄-alkyl, C₁-C₆-alkyloxycarbonylamino-C₁-C₄-alkyl, C₁-C₆-alkylaminocarbonylamino-C₁-C₄-alkyl, C₆-C₁₂-aryl-C₁-C₄-alkyl, C₃-C₁₂-cycloalkyl (e.g. cyclopropyl or cyclobutyl), C₂-C₆-alkenyl (e.g. prop-1,2-en-1-yl), optionally substituted C₆-C₁₂-aryl (e.g. phenyl), hydroxy, C₁-C₆-alkylamino, (halogenated C₁-C₆-alkyl)amino, di-C₁-C₆-alkylamino or optionally substituted M₃-M₁₂-heterocyclyl (e.g. 3-pyridyl, 2-pyridyl, 2-thienyl, 4-methyl-2-thienyl, 5-methyl-2-thienyl, 5-chloro-2-thienyl, 2,5-dimethyl-3-thienyl, 1,2-diazol-4-yl, 1-methyl-1,2-diazol-4-yl, 1,3-dimethyl-1,2-diazol-4-yl, 1-ethyl-1,2-diazol-4-yl, 1-difluormethyl-1,2-diazol-4-yl, 2-methyl-1,3-diazol-4-yl, 1-methyl-1,3-diazol-4-yl, 2-methyl-1,3-thiazol-5-yl, 2,4-dimethyl-1,3-thiazol-5-yl, 1-methyl-1,2,3-triazol-4-yl, 1-ethyl-1,2,3-triazol-4-yl, 3-pyrrolidinyl, furan-3-yl, 5-methyl-furan-2-yl, 2,5-dimethyl-furan-3-yl, 3-methyl-piperidinyl, thiophen-2-yl, 4-methyl-thiophen-2-yl, 5-methyl-thiophen-2-yl, thiophen-3-yl, or morpholin-4-yl).

More preferably, R¹ is C₁-C₆-alkyl (e.g. ethyl, n-propyl, isopropyl, 2-butyl), C₃-C₁₂-cycloalkyl-C₁-C₄-alkyl (e.g. cyclopropylmethyl), C₃-C₁₂-cycloalkyl (e.g. cyclobutyl), or optionally substituted M₃-M₁₂-heterocyclyl (e.g. 3-pyridyl, 2-pyridyl, 1-methyl-1,2-diazol-4-yl, 1,3-dimethyl-1,2-diazol-4-yl, 1-ethyl-1,2-diazol-4-yl, 1-methyl-1,3-diazol-4-yl, 1-methyl-1,2,3-triazol-4-yl, 1-ethyl-1,2,3-triazol-4-yl, 3-oxetanyl, 1-methyl-pyrrol-3-yl, furan-3-yl, 5-methyl-furan-2-yl, 2,5-dimethyl-furan-3-yl, 3-methyl-piperidinyl, thiophen-2-yl, 4-methyl-thiophen-2-yl, 5-methyl-thiophen-2-yl, thiophen-3-yl, or morpholin-4-yl).

According to a particular embodiment, R¹ is C₁-C₆-alkyl (e.g. ethyl, n-propyl, isopropyl, 2-butyl), C₃-C₁₂-cycloalkyl-C₁-C₄-alkyl (e.g. cyclopropylmethyl), or C₃-C₁₂-cycloalkyl (e.g. cyclobutyl).

In connection with R¹, substituted C₆-C₁₂-aryl in particular includes C₆-C₁₂-aryl, such as phenyl or naphthyl, substituted with 1, 2 or 3 substituents selected from the group consisting of halogen, C₁-C₄-alkyl, C₁-C₄-haloalkyl, cyano, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy, amino, C₁-C₄-alkylamino, C₁-C₄-dialkylamino, morpholinyl and piperidinyl. The same applies to substituted C₆-C₁₂-aryl in substituted C₆-C₁₂-aryl-C₁-C₄-alkyl.

In connection with R¹, substituted M₃-M₁₂-heterocyclyl in particular includes M₃-M₁₂-heterocyclyl, such as pyridyl, thienyl, diazolyl, quinolinyl, furanyl, thiophenyl, piperidinyl, piperazinyl or morpholinyl, pyrrolyl, isoxazolyl and triazolyl being further examples of such M₃-M₁₂-heterocyclyl, substituted with 1, 2 or 3 substituents selected from the group consisting of halogen, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxycarbonyl, cyano, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy, C₁-C₄-

alkylsulfonyl, amino, C₁-C₄-alkylamino, C₁-C₄-dialkylamino, C₆-C₁₂-aryl amino and M₃-M₁₂-heterocyclyl (e.g., morpholinyl or piperidinyl). The same applies to substituted M₃-M₁₂-heterocyclyl in substituted M₃-M₁₂-heterocyclyl-C₁-C₄-alkyl.

- 5 W is -NR⁷- or a bond. Y is -NR⁸- or a bond. According to one embodiment, W is -NR⁷- and Y is a bond. According to an alternative embodiment, W is a bond and Y is -NR⁸-. According to a further alternative embodiment, W is a bond and Y is a bond, especially if R¹ is a nitrogen-bound radical, e.g. nitrogen-bound heterocyclyl such as piperazinyl or morpholinyl.

- 10 Q is -S(O)₂- or -C(O)-. According to one embodiment, Q is -S(O)₂-, especially if Y is -NR⁸-. According to a preferred embodiment, -Q-Y- is -S(O)₂-NR⁸-.

- According to a particular embodiment, -W-A¹-Q-Y- is -W-A¹-S(O)₂-NR⁸-, -NR⁷-S(O)₂-, -A¹-S(O)₂- or -S(O)₂-. According to a further particular embodiment, -W-A¹-Q-Y- is -W-A¹-CO-NR⁸- or -NR⁷-CO-.
- 15

- A¹ is optionally substituted C₁-C₄-alkylene or a bond. In connection with A¹, substituted C₁-C₄-alkylene in particular includes C₁-C₄-alkylene substituted with 1, 2 or 3 substituents selected from the group consisting of halogen, C₁-C₄-alkyl and cyano. Preferably, A¹ is a bond. If A¹ is C₁-C₄-alkylene, W is preferably NR⁷-.
- 20

According to a particular embodiment, R¹-W-A¹-Q-Y- is R¹-S(O)₂-NH-, R¹-NH-S(O)₂-, R¹-C(O)-NH- or R¹-NH-C(O)-.


- 25 According to a further particular embodiment, W is a bond and A¹ is a bond.

The index n1 is 0, 1, 2, or 3. Preferably, n1 is 1, 2 or 3. In particular, n1 is 1 or 2.

- The index n2 is 0, 1, 2, or 3. Preferably, n2 is 1, 2, or 3. In particular, n2 is 1 or 2.
- 30

According to a particular embodiment, at least one of n1 and n2 is 1, 2, or 3.

The following examples of cyclic moieties illustrate combinations of n_1 and n_2 :



wherein X^1 and R^6 are as defined herein.

5


According to a further particular embodiment, the sum of n_1 and n_2 is 2, 3, or 4.

According to a further particular embodiment, combinations of n_1 and n_2 include $n_1=1, n_2=1$; $n_1=1, n_2=2$; $n_1=2, n_2=1$; $n_1=2, n_2=2$; $n_1=1, n_2=3$; or $n_1=3, n_2=1$.

10

According to one embodiment, X^1 is $>N-$. According to an alternative embodiment, X^1 is $>CH-$.

The following examples of cyclic moieties illustrate preferred combinations of n_1 , n_2 and X^1 :




wherein R^6 is as defined herein.

More preferred combinations of n_1 , n_2 and X^1 include cyclic moieties where

- 5 n_1 is 1, n_2 is 1 and X^1 is $\rightarrow N-$; n_1 is 1, n_2 is 1 and X^1 is $>CH-$; n_1 is 1, n_2 is 2 and X^1 is $\rightarrow N-$; n_1 is 1, n_2 is 2 and X^1 is $>CH-$; n_1 is 2, n_2 is 1 and X^1 is $\rightarrow N-$; n_1 is 2, n_2 is 1 and X^1 is $>CH-$; n_1 is 2, n_2 is 2 and X^1 is $\rightarrow N-$; n_1 is 2, n_2 is 2 and X^1 is $>CH-$; n_1 is 1, n_2 is 3 and X^1 is $\rightarrow N-$; n_1 is 1, n_2 is 3 and X^1 is $>CH-$; n_1 is 3, n_2 is 1 and X^1 is $\rightarrow N-$; or n_1 is 3, n_2 is 1 and X^1 is $>CH-$.





- 10 The cyclic moieties may thus be depicted by the following formulae:



wherein R^6 is as defined herein.

Particularly preferred combinations of n_1 , n_2 and X^1 include moieties where n_1 is 1, n_2 is 1 and X^1 is $\rightarrow N$ - (azetidiny); or n_1 is 1, n_2 is 1 and X^1 is $>CH$ - (cyclobutyl).


- 5 The substituents R^1 -W- A^1 -Q-Y- and -A on the cyclic moiety can be cis- or trans-configuration as depicted by the following formula:



wherein R^1 , W, A^1 , Q, Y, R^6 , n_1 , n_2 , X^1 and A are as defined herein.

- 10 According to a particular embodiment, the substituents R^1 -W- A^1 -Q-Y- and -A are in trans-configuration.


- 15 In formula (I), there may be one or more than one radical R^6 . More particularly, there may be up to 6 radicals R^6 . Preferably, there may be up to 4 radical R^6 . In particular, there may be one or 2 radicals R^6 . If there is more than one radical R^6 , these may be the same or different radicals. The compounds of the invention may therefore be represented by the following formula:



wherein R^{6a} , R^{6b} , R^{6c} , R^{6d} , R^{6e} independently have one of the meanings given for R^6 , and A , R^1 , W , A^1 , Q , Y , R^6 , X^1 , $n1$, $n2$, R^2 , A^2 , R^3 , Y^1 , R^{4a} , R^{4b} , X^2 , X^3 , R^5 are as defined herein (with X^1 being $>N-$ or $>CR^{6f}-$ and R^{6f} having one of the meanings given for R^6).

5

According to a particular embodiment, the compounds of the invention have the following formula:



wherein R^{6e} has one of the meanings given for R^6 , and A , R^1 , W , A^1 , Q , Y , R^6 , X^1 , $n1$, $n2$, R^2 , A^2 , R^3 , Y^1 , R^{4a} , R^{4b} , X^2 , X^3 , R^5 are as defined herein.

10


R^6 is hydrogen, halogen (e.g. fluorine), C_1 - C_4 -alkyl (e.g. methyl or ethyl), halogenated C_1 - C_4 -alkyl (e.g. 1,1,1-trifluoromethyl-yl), $-CN$, hydroxy, C_1 - C_6 -alkoxy (e.g. methoxy), or halogenated C_1 - C_6 -alkoxy.

15

According to an alternative embodiment, two R^6 together with the carbon atom to which they are bound may form a carbonyl.


20

The following examples of cyclic moieties illustrate combinations of $n1$, $n2$ and R^6 , wherein two R^6 together with the carbon atom to which they are bound form a carbonyl:




The following examples of cyclic moieties illustrate particular combinations of n_1 , n_2 and R^6 , wherein two R^6 together with the carbon atom to which they are bound form a carbonyl:

5




The following examples of cyclic moieties illustrate preferred combinations of n_1 , n_2 , X^1 and R^6 , wherein two R^6 together with the carbon atom to which they are bound form a carbonyl.




10

Preferably, R^6 is hydrogen or C_1 - C_4 -alkyl (e.g. methyl, ethyl). In particular, R^6 is hydrogen.

If A is a benzene ring, the radical




may, in principle, be bound to the 5-, 6-, 7- or 8-position of the skeleton of the compounds of the invention:



In said formulae, R^1 , W , A^1 , Q , Y , R^6 , X^1 , n_1 , n_2 , R^2 , A^2 , R^3 , Y^1 , R^{4a} , R^{4b} , X^2 , X^3 , R^5 are as defined herein.


5

Compounds of the invention having the radical



in the 5-, 6-, 7-position are preferred.


Particularly preferred are compounds of the invention having the radical




5

in the 6-position.

In addition to the radical



- 10 the compounds of the invention may have one or more than one further substituent bound to the ring A. In these positions, the skeleton of the compounds of the invention may thus be substituted with one or more than one radical R^2 . If there is more than one radical R^2 , these may be the same or different radicals. In particular, the skeleton of the compounds of the invention may be substituted with one or more than one radical R^2 in 5-, 6-, 7- and/or 8-position if A is a benzene ring. The
- 15 compounds of the invention may therefore be represented by one of the following formulae:



wherein R^{2a} , R^{2b} , R^{2c} , R^{2d} independently have one of the meanings given for R^2 , and R^1 , W , A^1 , Q , Y , R^6 , X^1 , $n1$, $n2$, R^2 , A^2 , R^3 , Y^1 , R^{4a} , R^{4b} , X^2 , X^3 , R^5 are as defined herein.

R^2 is hydrogen, halogen (e.g. fluorine), C_1 - C_6 -alkyl, halogenated C_1 - C_4 -alkyl, -CN, C_2 - C_6 -alkenyl, C_2 - C_6 -alkynyl, optionally substituted C_6 - C_{12} -aryl, hydroxy, C_1 - C_6 -alkoxy, halogenated C_1 - C_6 -alkoxy, C_1 - C_6 -alkoxycarbonyl, C_2 - C_6 -alkenyloxy, C_6 - C_{12} -aryl- C_1 - C_4 -alkoxy, C_1 - C_6 -alkylcarbonyloxy, C_1 - C_6 -alkylthio, C_1 - C_6 -alkylsulfinyl, C_1 - C_6 -alkylsulfonyl, aminosulfonyl, amino, C_1 - C_6 -alkylamino, C_2 - C_6 -alkenylamino, nitro or optionally substituted M_3 - M_{12} -heterocyclyl, or two radicals R^2 together with the ring atoms of A to which they are bound form a 5- or 6 membered ring.




An optionally substituted 5- or 6-membered ring that is formed by two radicals R^2 together with the ring atoms of A to which they are bound is, for instance, a benzene ring.

In connection with R^2 , substituted C_6 - C_{12} -aryl in particular includes C_6 - C_{12} -aryl, such as phenyl, substituted with 1, 2 or 3 substituents selected from the group consisting of halogen, C_1 - C_4 -alkyl, C_1 - C_4 -haloalkyl, cyano, C_1 - C_4 -alkoxy and C_1 - C_4 -haloalkoxy.

In connection with R^2 , substituted M_3 - M_{12} -heterocyclyl in particular includes M_3 - M_{12} -heterocyclyl, such as morpholinyl, pyrrolidinyl and piperidinyl, substituted with 1, 2 or 3 substituents selected from the group consisting of halogen, C_1 - C_4 -alkyl, C_1 - C_4 -haloalkyl, cyano, C_1 - C_4 -alkoxy and C_1 - C_4 -haloalkoxy.

Preferably, R^2 is hydrogen, halogen (e.g. fluorine), CN or C_1 - C_6 -alkoxy. In particular, R^2 is hydrogen or halogen (e.g. fluorine).

According to a particular embodiment, the compounds of the invention have one of the following formulae:




wherein R^1 , W , A^1 , Q , Y , R^6 , $n1$, $n2$, X^1 , R^2 , A^2 , R^3 , Y^1 , R^{4a} , R^{4b} , X^2 , X^3 and R^5 are as defined herein.

5

A^2 is -O-, -S- or -NR⁹-. According to a preferred embodiment, A^2 is -O-.




In 2-, 3- and/or 4-position, the compounds of the invention may be substituted with one or more than one radical R^3 . If there is more than one radical R^3 , these may be the same or different radicals. The compounds of the invention may therefore be represented by the following formula:

10



wherein R^{3a} , R^{3b} , R^{3c} , R^{3d} independently have one of the meanings given for R^3 , and A , R^1 , W , A^1 , Q , Y , R^6 , $n1$, $n2$, X^1 , R^2 , A^2 , R^3 , Y^1 , R^{4a} , R^{4b} , X^2 , X^3 , R^5 are as defined herein.

- 5 According to a particular embodiment, the compounds of the invention have one of the following formulae:



wherein R^{3a} , R^{3b} , R^{3d} independently have the meaning of R^3 and A , R^1 , W , A^1 , Q , Y , R^6 , $n1$, $n2$, X^1 , R^2 , A^2 , R^3 , Y^1 , R^{4a} , R^{4b} , X^2 , X^3 , R^5 are as defined herein.

R^3 is hydrogen, halogen, C_1 - C_6 -alkyl, C_1 - C_6 -alkoxy, or two radicals R^3 together with the carbon atom to which they are attached form a carbonyl group.

Preferably, R^3 is hydrogen or C_1 - C_6 -alkyl (e.g. methyl). In particular, R^3 is hydrogen.

5

Y^1 is a bond or optionally substituted C_1 - C_4 -alkylene (e.g. methylene or 1,2-ethylene). In connection with Y^1 , substituted C_1 - C_4 -alkylene in particular includes C_1 - C_4 -alkylene substituted with 1, 2 or 3 substituents selected from the group consisting of halogen, C_1 - C_4 -alkyl, C_1 - C_4 -haloalkyl, C_3 - C_{12} -cycloalkyl and cyano. In particular, Y^1 is a bond.


10

R^{4a} is hydrogen, C_1 - C_6 -alkyl (e.g. methyl, ethyl, n-propyl or isopropyl), C_3 - C_{12} -cycloalkyl- C_1 - C_4 -alkyl (e.g. cyclopropylmethyl), halogenated C_1 - C_4 -alkyl (e.g. 2-fluoroethyl or 2,2,2-trifluoroethyl), hydroxy- C_1 - C_4 -alkyl, C_1 - C_6 -alkoxy- C_1 - C_4 -alkyl, amino- C_1 - C_4 -alkyl, $-CH_2CN$, C_6 - C_{12} -aryl- C_1 - C_4 -alkyl (e.g. benzyl), optionally substituted C_3 - C_{12} -cycloalkyl (e.g. cyclopropyl or cyclobutyl), $-CHO$, C_1 - C_4 -alkylcarbonyl (e.g. methylcarbonyl, ethylcarbonyl or isopropylcarbonyl), (halogenated C_1 - C_4 -alkyl)carbonyl (e.g. fluoromethylcarbonyl, difluoromethylcarbonyl, trifluoromethylcarbonyl, 1,1,1-trifluoroeth-2-ylcarbonyl or 1,1,1-trifluoroprop-3-ylcarbonyl), C_6 - C_{12} -arylcarbonyl (e.g. phenylcarbonyl), C_1 - C_4 -alkoxycarbonyl (e.g. ethoxycarbonyl or tert-butyloxycarbonyl), C_6 - C_{12} -aryloxycarbonyl (e.g. phenoxycarbonyl), C_1 - C_6 -alkylaminocarbonyl, C_2 - C_6 -alkenyl, $-C(=NH)NH_2$, $-C(=NH)NHCN$, C_1 - C_6 -alkylsulfonyl, C_6 - C_{12} -arylsulfonyl, amino, $-NO$ or optionally substituted M_3 - M_{12} -heterocyclyl (e.g. 3-oxetanyl).

In connection with R^{4a} , substituted C_3 - C_{12} -cycloalkyl in particular includes C_3 - C_{12} -cycloalkyl such as cyclopropyl, cyclobutyl or cyclohexyl, substituted with 1, 2 or 3 substituents selected from the group consisting of halogen, optionally substituted C_1 - C_6 -alkyl, halogenated C_1 - C_6 -alkyl, CN, hydroxy, C_1 - C_6 -alkoxy, halogenated C_1 - C_6 -alkoxy, amino, C_1 - C_6 -alkylamino, di- C_1 - C_6 -alkylamino and M_3 - M_{12} -heterocyclyl.

In connection with R^{4a} , substituted M_3 - M_{12} -heterocyclyl in particular includes M_3 - M_{12} -heterocyclyl substituted with 1 or more substituents R^{4g} and/or $R^{4b'}$. The compounds of the invention may therefore be represented by the following formula:

30



wherein A, R^1 , W, A^1 , Q, Y, R^6 , $n1$, $n2$, X^1 , R^2 , A^2 , R^3 , Y^1 , R^{4b} , X^2 , X^3 and R^5 are as defined herein,

35

R^{4b} is hydrogen, halogen, C₁-C₆-alkyl, C₃-C₁₂-cycloalkyl-C₁-C₄-alkyl, halogenated C₁-C₆-alkyl, tri-(C₁-C₄-alkyl)-silyl-C₁-C₄-alkyl, hydroxy-C₁-C₄-alkyl, C₁-C₆-alkoxy-C₁-C₄-alkyl, amino-C₁-C₄-alkyl, C₁-C₆-alkylamino-C₁-C₄-alkyl, di-C₁-C₆-alkylamino-C₁-C₄-alkyl, C₁-C₆-alkylcarbonylamino-C₁-C₄-alkyl, C₁-C₆-alkyloxycarbonylamino-C₁-C₄-alkyl, C₁-C₆-alkylaminocarbonylamino-C₁-C₄-alkyl, di-C₁-C₆-alkylaminocarbonylamino-C₁-C₄-alkyl, C₁-C₆-alkylsulfonylamino-C₁-C₄-alkyl, (optionally substituted C₆-C₁₂-aryl-C₁-C₆-alkyl)amino-C₁-C₄-alkyl, optionally substituted C₆-C₁₂-aryl-C₁-C₄-alkyl, optionally substituted C₃-C₁₂-heterocyclyl-C₁-C₄-alkyl, C₃-C₁₂-cycloalkyl, C₁-C₆-alkylcarbonyl, C₁-C₆-alkoxycarbonyl, halogenated C₁-C₆-alkoxycarbonyl, C₆-C₁₂-aryloxycarbonyl, aminocarbonyl, C₁-C₆-alkylaminocarbonyl, (halogenated C₁-C₄-alkyl)aminocarbonyl, C₆-C₁₂-arylamino-C₁-C₄-alkyl, C₂-C₆-alkenyl, C₂-C₆-alkynyl, optionally substituted C₆-C₁₂-aryl, cyano; hydroxy, C₁-C₆-alkoxy, halogenated C₁-C₆-alkoxy, C₁-C₆-hydroxyalkoxy, C₁-C₆-alkoxy-C₁-C₄-alkoxy, amino-C₁-C₄-alkoxy, C₁-C₆-alkylamino-C₁-C₄-alkoxy, di-C₁-C₆-alkylamino-C₁-C₄-alkoxy, C₁-C₆-alkylcarbonylamino-C₁-C₄-alkoxy, C₆-C₁₂-arylcarbonylamino-C₁-C₄-alkoxy, C₁-C₆-alkoxycarbonylamino-C₁-C₄-alkoxy, C₆-C₁₂-aryl-C₁-C₄-alkoxy, C₁-C₆-alkylsulfonylamino-C₁-C₄-alkoxy, (halogenated C₁-C₆-alkyl)sulfonylamino-C₁-C₄-alkoxy, C₆-C₁₂-arylsulfonylamino-C₁-C₄-alkoxy, (C₆-C₁₂-aryl-C₁-C₆-alkyl)sulfonylamino-C₁-C₄-alkoxy, C₃-C₁₂-heterocyclylsulfonylamino-C₁-C₄-alkoxy, C₃-C₁₂-heterocyclyl-C₁-C₄-alkoxy, C₆-C₁₂-aryloxy, C₃-C₁₂-heterocyclyloxy, C₁-C₆-alkylthio, halogenated C₁-C₆-alkylthio, C₁-C₆-alkylamino, (halogenated C₁-C₆-alkyl)amino, di-C₁-C₆-alkylamino, di-(halogenated C₁-C₆-alkyl)amino, C₁-C₆-alkylcarbonylamino, (halogenated C₁-C₆-alkyl)carbonylamino, C₆-C₁₂-arylcarbonylamino, C₁-C₆-alkylsulfonylamino, (halogenated C₁-C₆-alkyl)sulfonylamino, C₆-C₁₂-arylsulfonylamino or optionally substituted C₃-C₁₂-heterocyclyl,

R^{4g} is hydrogen, halogen, C₁-C₆-alkyl, C₃-C₁₂-cycloalkyl-C₁-C₄-alkyl, halogenated C₁-C₆-alkyl, tri-(C₁-C₄-alkyl)-silyl-C₁-C₄-alkyl, hydroxy-C₁-C₄-alkyl, C₁-C₆-alkoxy-C₁-C₄-alkyl, amino-C₁-C₄-alkyl, C₁-C₆-alkylamino-C₁-C₄-alkyl, di-C₁-C₆-alkylamino-C₁-C₄-alkyl, C₁-C₆-alkylcarbonylamino-C₁-C₄-alkyl, C₁-C₆-alkyloxycarbonylamino-C₁-C₄-alkyl, C₁-C₆-alkylaminocarbonylamino-C₁-C₄-alkyl, di-C₁-C₆-alkylaminocarbonylamino-C₁-C₄-alkyl, C₁-C₆-alkylsulfonylamino-C₁-C₄-alkyl, (optionally substituted C₆-C₁₂-aryl-C₁-C₆-alkyl)amino-C₁-C₄-alkyl, optionally substituted C₆-C₁₂-aryl-C₁-C₄-alkyl, optionally substituted C₃-C₁₂-heterocyclyl-C₁-C₄-alkyl, C₃-C₁₂-cycloalkyl, C₁-C₆-alkylcarbonyl, C₁-C₆-alkoxycarbonyl, halogenated C₁-C₆-alkoxycarbonyl, C₆-C₁₂-aryloxycarbonyl, aminocarbonyl, C₁-C₆-alkylaminocarbonyl, (halogenated C₁-C₄-alkyl)aminocarbonyl, C₆-C₁₂-arylamino-C₁-C₄-alkyl, C₂-C₆-alkenyl, C₂-C₆-alkynyl, optionally substituted C₆-C₁₂-aryl, cyano, hydroxy, C₁-C₆-alkoxy, halogenated C₁-C₆-alkoxy, C₁-C₆-hydroxyalkoxy, C₁-C₆-alkoxy-C₁-C₄-alkoxy, amino-C₁-C₄-alkoxy, C₁-C₆-alkylamino-C₁-C₄-alkoxy, di-C₁-C₆-alkylamino-C₁-C₄-alkoxy, C₁-C₆-alkylcarbonylamino-C₁-C₄-alkoxy, C₆-C₁₂-arylcarbonylamino-C₁-C₄-alkoxy, C₁-C₆-alkoxycarbonylamino-C₁-C₄-alkoxy, C₆-C₁₂-aryl-C₁-C₄-alkoxy, C₁-C₆-alkylsulfonylamino-C₁-C₄-alkoxy, (halogenated C₁-C₆-alkyl)sulfonylamino-C₁-C₄-alkoxy, C₆-C₁₂-arylsulfonylamino-C₁-C₄-alkoxy, (C₆-C₁₂-aryl-C₁-C₆-alkyl)sulfonylamino-C₁-C₄-alkoxy, C₃-C₁₂-heterocyclylsulfonylamino-C₁-C₄-alkoxy, C₃-C₁₂-heterocyclyl-C₁-C₄-alkoxy, C₆-C₁₂-aryloxy, C₃-C₁₂-heterocyclyloxy, C₁-C₆-alkylthio, halogenated C₁-C₆-alkylthio, C₁-C₆-alkylamino, (halogenated C₁-C₆-alkyl)amino, di-C₁-C₆-alkylamino, di-(halogenated C₁-C₆-alkyl)amino, C₁-C₆-alkylcarbonylamino, (halogenated C₁-

C₆-alkyl)carbonylamino, C₆-C₁₂-arylcarbonylamino, C₁-C₆-alkylsulfonylamino, (halogenated C₁-C₆-alkyl)sulfonylamino, C₆-C₁₂-arylsulfonylamino or optionally substituted C₃-C₁₂-heterocyclyl,

the index q is 1, 2 or 3; and in particular, q is 1 or 2,

5

the index r is 1, 2 or 3; and in particular, r is 1 or 2,

X⁴ is -O-, -NR¹⁷-, -S-, -S(O)-, -S(O)₂-, or a bond and preferable, X⁴ is -O- or a bond, and

10 R¹⁷ is hydrogen, C₁-C₆-alkyl or C₃-C₁₂-cycloalkyl. Preferably, R¹⁷ is hydrogen.

Particular combinations of q and r include moieties wherein q is 1 and r is 1, or q is 2 and r is 1.

15 Particular combinations of q, r and X⁴ include moieties where q is 1, r is 1 and X⁴ is -O- (oxetanyl); q is 1, r is 1 and X⁴ is a bond (cyclopropyl); or q is 2, r is 1 and X⁴ is a bond (cyclobutyl).

20 According to a preferred embodiment, R^{4b'} is hydrogen, halogen, C₁-C₆-alkyl, C₃-C₁₂-cycloalkyl-C₁-C₄-alkyl, halogenated C₁-C₆-alkyl, hydroxy-C₁-C₄-alkyl, C₁-C₆-alkoxy-C₁-C₄-alkyl, optionally substituted C₆-C₁₂-aryl-C₁-C₄-alkyl, C₂-C₆-alkenyl, C₂-C₆-alkynyl, optionally substituted C₆-C₁₂-aryl, cyano, hydroxy, C₁-C₆-alkoxy, halogenated C₁-C₆-alkoxy, C₁-C₆-hydroxyalkoxy, C₁-C₆-alkoxy-C₁-C₄-alkoxy, C₆-C₁₂-aryl-C₁-C₄-alkoxy, C₃-C₁₂-heterocycloxy or optionally substituted C₃-C₁₂-heterocyclyl.

25 According to a particular embodiment, R^{4b'} is hydrogen, halogen, C₁-C₆-alkyl, C₃-C₆-cycloalkyl-C₁-C₄-alkyl, halogenated C₁-C₆-alkyl, hydroxy-C₁-C₄-alkyl, C₁-C₆-alkoxy-C₁-C₄-alkyl, optionally substituted C₆-aryl-C₁-C₄-alkyl, optionally substituted C₆-aryl, cyano, hydroxy, C₁-C₆-alkoxy, halogenated C₁-C₆-alkoxy, C₁-C₆-hydroxyalkoxy, C₁-C₆-alkoxy-C₁-C₄-alkoxy or C₆-aryl-C₁-C₄-alkoxy.

30 According to a preferred embodiment, R^{4g} is hydrogen, halogen, C₁-C₆-alkyl, C₃-C₁₂-cycloalkyl-C₁-C₄-alkyl, halogenated C₁-C₆-alkyl, hydroxy-C₁-C₄-alkyl, C₁-C₆-alkoxy-C₁-C₄-alkyl, optionally substituted C₆-C₁₂-aryl-C₁-C₄-alkyl, C₂-C₆-alkenyl, C₂-C₆-alkynyl, optionally substituted C₆-C₁₂-aryl, cyano or optionally substituted C₃-C₁₂-heterocyclyl.

35 According to a particular embodiment, R^{4g} is hydrogen, halogen, C₁-C₆-alkyl, C₃-C₆-cycloalkyl-C₁-C₄-alkyl, halogenated C₁-C₆-alkyl, hydroxy-C₁-C₄-alkyl, C₁-C₆-alkoxy-C₁-C₄-alkyl, optionally substituted C₆-aryl-C₁-C₄-alkyl, optionally substituted C₆-aryl, cyano or C₆-aryl-C₁-C₄-alkoxy.

It is in particular preferred if R^{4g} is an electron withdrawing group.

40

In connection with R^{4a} substituted C₁-C₆-alkyl in particular includes C₁-C₆-alkyl, especially C₁-C₄-alkyl, substituted with 1, 2 or 3 substituents selected from the group consisting of hydroxy, C₁-C₆-


alkoxy, amino, C₁-C₆-alkylamino, di-C₁-C₆-alkylamino and M₃-M₁₂-heterocyclyl (e.g. morpholinyl or piperidinyl).

Preferably, R^{4a} is hydrogen, C₁-C₆-alkyl (e.g. methyl, ethyl, n-propyl, isopropyl, 2-methyl-but-4-yl, or 2-methyl-prop-3-yl), C₃-C₁₂-cycloalkyl-C₁-C₄-alkyl (e.g. cyclopropylmethyl, cyclobutylmethyl, cyclopentylmethyl, 1-cyclopropyl-eth-2-yl, 1-cyclopentyl-eth-2-yl, or cyclohexylmethyl), halogenated C₁-C₄-alkyl (e.g. 2-fluoroethyl or 2,2,2-trifluoroethyl), amino-C₁-C₄-alkyl, -CH₂CN, C₆-C₁₂-aryl-C₁-C₄-alkyl (e.g. benzyl), optionally substituted C₃-C₁₂-cycloalkyl (e.g. cyclopropyl or cyclobutyl), C₁-C₄-alkylcarbonyl (e.g. methylcarbonyl or isopropylcarbonyl), (halogenated C₁-C₄-alkyl)carbonyl (e.g. fluoromethylcarbonyl, difluoromethylcarbonyl or trifluoromethylcarbonyl), C₆-C₁₂-arylcarbonyl (e.g. phenylcarbonyl), C₁-C₄-alkoxycarbonyl (e.g. ethoxycarbonyl or tert-butylloxycarbonyl), C₆-C₁₂-aryloxycarbonyl (e.g. phenoxycarbonyl), -C(=NH)NH₂, -C(=NH)NHCN, C₁-C₆-alkylsulfonyl, amino, -NO or optionally substituted M₃-M₁₂-heterocyclyl (e.g. 3-oxetanyl).

More preferably, R^{4a} is hydrogen, C₁-C₆-alkyl (e.g. methyl, ethyl, n-propyl, isopropyl, 2-methyl-but-4-yl, or 2-methyl-prop-3-yl), optionally substituted C₃-C₁₂-cycloalkyl (e.g. cyclopropyl or cyclobutyl), C₃-C₁₂-cycloalkyl-C₁-C₄-alkyl (e.g. cyclopropylmethyl, cyclobutylmethyl, cyclopentylmethyl, 1-cyclopropyl-eth-2-yl, 1-cyclopentyl-eth-2-yl, or cyclohexylmethyl), or M₃-M₁₂-heterocyclyl (e.g. 3-oxetanyl).

In particular, R^{4a} is hydrogen, C₁-C₆-alkyl (e.g. methyl, ethyl, n-propyl, isopropyl, 2-methyl-but-4-yl, or 2-methyl-prop-3-yl), C₃-C₁₂-cycloalkyl-C₁-C₄-alkyl (e.g. cyclopropylmethyl, cyclobutylmethyl, cyclopentylmethyl, 1-cyclopropyl-eth-2-yl, 1-cyclopentyl-eth-2-yl, or cyclohexylmethyl), or optionally substituted C₃-C₁₂-cycloalkyl (e.g. cyclopropyl or cyclobutyl).

Alternatively, R^{4a} is optionally substituted C₁-C₄-alkylene (e.g. methylene, 1,2-ethylene, or 1,3-propylene) that is bound to a carbon atom in Y¹. In connection with R^{4a}, substituted C₁-C₄-alkylene in particular includes C₁-C₄-alkylene substituted with 1, 2 or 3 substituents selected from the group consisting of halogen, C₁-C₄-alkyl, C₁-C₄-haloalkyl, cyano, hydroxy and C₁-C₄-alkoxy. In particular, R^{4a} is C₁-C₄-alkylene (e.g. methylene or 1,2-ethylene) that is bound to a carbon atom in Y¹ with Y¹ being optionally substituted C₁-C₄-alkylene (e.g. 1,2-ethylene or 1,3-propylene) so that R^{4a} and at least part of Y¹ together with the nitrogen atom to which R^{4a} and Y¹ are bound form an N-containing heterocyclic ring having, in particular, 4, 5 or 6 ring member atoms (including the nitrogen atom). A derivative of the invention having such a ring may be represented by the following partial structure:



wherein A, R¹, W, A¹, Q, Y, R⁶, n1, n2, X¹, R², A², R³, R^{4b}, X², X³, R⁵ are as defined herein, s is 0, 1 or 2, and t is 0, 1, 2, or 3. Particular combinations of s and t include s=1, t=1; s=0, t=1; s=1, t=2; and s=0, t=2.

5

R^{4b} is hydrogen, C₁-C₆-alkyl (e.g. methyl, ethyl), halogenated C₁-C₄-alkyl, hydroxy-C₁-C₄-alkyl, C₁-C₆-alkoxy-C₁-C₄-alkyl, amino-C₁-C₄-alkyl, -CH₂CN, -CHO, C₁-C₄-alkylcarbonyl, (halogenated C₁-C₄-alkyl)carbonyl, C₆-C₁₂-arylcarbonyl, C₁-C₄-alkoxycarbonyl, C₆-C₁₂-aryloxycarbonyl, C₁-C₆-alkylaminocarbonyl, C₂-C₆-alkenyl, -C(=NH)NH₂, -C(=NH)NHCN, C₁-C₆-alkylsulfonyl, C₆-C₁₂-arylsulfonyl, amino, -NO or M₃-M₁₂-heterocyclyl.

10

Preferably, R^{4b} is hydrogen or C₁-C₆-alkyl (e.g. methyl or ethyl). In particular, R^{4b} is hydrogen.


Alternatively, R^{4a}, R^{4b} together are optionally substituted C₂-C₆-alkylene (e.g. 1,4-butylenes, 1,3-propylene, 2-fluoro-but-1,4-ylene, 1-oxo-but-1,4-ylene, 2-methyl-1,3-propylene, 2,2-dimethyl-1,3-propylene, or 2-methyl-2-hydroxy-1,3-propylene), wherein one -CH₂- of C₂-C₆-alkylene may be replaced by an oxygen atom (e.g. -CH₂-CH₂-O-CH₂-CH₂-) or -NR¹⁰.

15

In connection with R^{4a} and R^{4b}, substituted C₂-C₆-alkylene in particular includes C₂-C₆-alkylene substituted with 1 or more substituents R^{4c}, R^{4d} and/or R^{4e}.

20

The compounds of the invention may therefore be represented by the following formula:



wherein A, R¹, W, A¹, Q, Y, R⁶, n1, n2, X¹, R², A², R³, Y¹, X², X³ and R⁵ are as defined herein, R^{4c} and R^{4d} are hydrogen, or R^{4c}, R^{4d} together are C₁-C₅-alkylene optionally substituted with 1, 2 or 3 substituents R^{4f}, wherein one -CH₂- of C₁-C₅-alkylene may be replaced by an oxygen atom or -NR¹⁸,

25


- R^{4e} is hydrogen, halogen, C₁-C₆-alkyl, C₃-C₆-cycloalkyl-C₁-C₄-alkyl, halogenated C₁-C₆-alkyl, hydroxy-C₁-C₄-alkyl, C₁-C₆-alkoxy-C₁-C₄-alkyl, optionally substituted C₆-C₁₂-aryl-C₁-C₄-alkyl, optionally substituted C₆-C₁₂-aryl, cyano, hydroxy, C₁-C₆-alkoxy, halogenated C₁-C₆-alkoxy, C₁-C₆-hydroxyalkoxy, C₁-C₆-alkoxy-C₁-C₄-alkoxy or C₆-C₁₂-aryl-C₁-C₄-alkoxy; and in particular, R^{4e} is hydrogen,
- 5 t is 0, 1, 2 or 3; and according to a particular embodiment, t is 1, and R¹⁸ is hydrogen, C₁-C₆-alkyl or C₃-C₁₂-cycloalkyl. Preferably, R¹⁸ is hydrogen.

- In connection with R^{4a} and R^{4e} substituted C₆-C₁₂-aryl in particular includes C₆-C₁₂-aryl, such as phenyl, substituted with 1, 2 or 3 substituents selected from the group consisting of halogen, C₁-C₄-alkyl, C₁-C₄-haloalkyl, cyano, C₁-C₄-alkoxy and C₁-C₄-haloalkoxy. The same applies to substituted C₆-C₁₂-aryl in substituted C₆-C₁₂-aryl-C₁-C₄-alkyl.
- 10

- According to a particular embodiment R^{4c}, R^{4d} together are C₁-C₅-alkylene optionally substituted with 1, 2 or 3 substituents R^{4f}, wherein one -CH₂- of C₁-C₅-alkylene may be replaced by an oxygen atom or -NR¹⁸-.
- 15


- In connection with R^{4c}, R^{4d}, substituted C₁-C₅-alkylene in particular includes C₁-C₅-alkylene optionally substituted with 1, 2 or 3 substituents (R^{4f}) selected from the group consisting of hydrogen, halogen, C₁-C₆-alkyl, C₃-C₁₂-cycloalkyl-C₁-C₄-alkyl, halogenated C₁-C₆-alkyl, hydroxy-C₁-C₄-alkyl, C₁-C₆-alkoxy-C₁-C₄-alkyl, optionally substituted C₆-C₁₂-aryl-C₁-C₄-alkyl, C₂-C₆-alkenyl, C₂-C₆-alkynyl, optionally substituted C₆-C₁₂-aryl, cyano, hydroxy, C₁-C₆-alkoxy, halogenated C₁-C₆-alkoxy, C₁-C₆-hydroxyalkoxy, C₁-C₆-alkoxy-C₁-C₄-alkoxy, C₆-C₁₂-aryl-C₁-C₄-alkoxy, M₃-M₁₂-heterocycloxy or optionally substituted M₃-M₁₂-heterocyclyl, and more preferably hydrogen,
- 20 halogen; C₁-C₆-alkyl, C₃-C₆-cycloalkyl-C₁-C₄-alkyl, halogenated C₁-C₆-alkyl, hydroxy-C₁-C₄-alkyl, C₁-C₆-alkoxy-C₁-C₄-alkyl, optionally substituted C₁-C₆-aryl-C₁-C₄-alkyl, optionally substituted C₁-C₆-aryl, cyano, hydroxy, C₁-C₆-alkoxy, halogenated C₁-C₆-alkoxy, C₁-C₆-hydroxyalkoxy, C₁-C₆-alkoxy-C₁-C₄-alkoxy or C₁-C₆-aryl-C₁-C₄-alkoxy.
- 25

- According to a further particular embodiment, R^{4c}, R^{4d} together with the carbon atom or the carbon atoms to which they are bound form a 3-, 4-, 5- or 6-membered ring, for example a ring comprised by the formula:
- 30




wherein t is defined as herein and u is 0, 1, 2, or 3, and R^{4e} and R^{4f} are as defined herein. Particular combinations of u and t include $t=1$ and $u=0$.


- 5 In said formulae, there may be one or more than one radical R^{4e} and/or R^{4f} . More particularly, there may be up to 3 radicals R^{4e} and/or up to 3 radicals R^{4f} . Preferably there is one radical R^{4e} and/or one radical R^{4f} . Said formulae may thus also be depicted as follows:



- 10 In said formulae, e is 1, 2 or 3 and f is 1, 2, or 3, with R^{4e} , R^{4f} , t and u being as defined herein. If there is more than one radical R^{4e} , these may be the same or different radicals. If there is more than one radical R^{4f} , these may be the same or different radicals.


- 15 The following examples of bicyclic moieties illustrate particular combinations of t , u and R^{4e} , R^{4f} in the compounds of the present invention:





wherein R^{4e} , R^{4f} are as defined herein and in particular are both hydrogen.

Compounds of the invention having the following bicyclic moiety:



wherein R^{4e} , R^{4f} are as defined herein and in particular are both hydrogen, are particularly preferred.

- 5 X^2 is $-O-$, $-NR^{11a}-$, $-S-$, $>CR^{12a}R^{12b}$ or a bond. In particular, X^2 is not a bond. Preferably, X^2 is $>CR^{12a}R^{12b}$.

X^3 is $-O-$, $-NR^{11b}-$, $-S-$, $>CR^{13a}R^{13b}$ or a bond. Preferably, X^3 is a bond.

- 10 Thus, it is preferred if X^2 is $>CR^{12a}R^{12b}$ and X^3 is a bond.

R^{12a} is hydrogen, optionally substituted C_1 - C_6 -alkyl, C_1 - C_6 -alkylamino- C_1 - C_4 -alkyl, di- C_1 - C_6 -alkylamino- C_1 - C_4 -alkyl, M_3 - M_{12} -heterocycl- C_1 - C_6 -alkyl, optionally substituted C_6 - C_{12} -aryl or hydroxy. Preferably, R^{12a} is hydrogen or C_1 - C_6 -alkyl.

15

R^{13a} is hydrogen, optionally substituted C_1 - C_6 -alkyl, C_1 - C_6 -alkylamino- C_1 - C_4 -alkyl, di- C_1 - C_6 -alkylamino- C_1 - C_4 -alkyl, M_3 - M_{12} -heterocycl- C_1 - C_6 -alkyl, optionally substituted C_6 - C_{12} -aryl or hydroxy. Preferably, R^{13a} is hydrogen or C_1 - C_6 -alkyl.

- 20 In connection with R^{12a} and R^{13a} , substituted C_1 - C_6 -alkyl in particular includes C_1 - C_6 -alkyl substituted with 1, 2 or 3 substituents selected from the group consisting of halogen, hydroxy, C_1 - C_4 -alkoxy and amino.

- 25 In connection with R^{12a} and R^{13a} , substituted C_6 - C_{12} -aryl in particular includes C_6 - C_{12} -aryl, such as phenyl, substituted with 1, 2 or 3 substituents selected from the group consisting of C_1 - C_4 -alkyl, C_1 - C_4 -haloalkyl, cyano, C_1 - C_4 -alkoxy and C_1 - C_4 -haloalkoxy.

R^{12b} is hydrogen or C_1 - C_6 -alkyl. According to a particular embodiment, R^{12b} is hydrogen.

- 30 R^{13b} is hydrogen or C_1 - C_6 -alkyl. According to a particular embodiment, R^{13b} is hydrogen.

Alternatively, R^{12a} and R^{12b} , or R^{13a} and R^{13b} , together with the carbon atom to which they are attached form a carbonyl or, preferably, are optionally substituted C_2 - C_4 -alkylene (e.g. 1,3-propylene), wherein one $-CH_2-$ of C_2 - C_4 -alkylene may be replaced by an oxygen atom or $-NR^{14}-$.

35

In connection with R^{12a} and R^{12b} , or R^{13a} and R^{13b} , substituted C_2 - C_4 -alkylene in particular includes C_2 - C_4 -alkylene substituted with 1, 2 or 3 substituents selected from the group consisting of halogen, C_1 - C_4 -alkyl, C_1 - C_4 -haloalkyl, cyano, C_1 - C_4 -alkoxy and C_1 - C_4 -haloalkoxy.

According to a particular embodiment, R^{12a} is hydrogen or C_1 - C_6 -alkyl and R^{12b} is hydrogen or C_1 - C_6 -alkyl, or R^{13a} is hydrogen or C_1 - C_6 -alkyl and R^{13b} is hydrogen or C_1 - C_6 -alkyl.

- 5 According to a further particular embodiment, R^{12a} is hydrogen and R^{12b} is hydrogen, or R^{13a} is hydrogen and R^{13b} is hydrogen.

According to a further particular embodiment, R^{12a} and R^{12b} together are optionally substituted 1,3-propylene, or R^{13a} and R^{13b} together are optionally substituted 1,3-propylene.

10

R^5 is optionally substituted C_6 - C_{12} -aryl (e.g. phenyl, 2-fluorophenyl, 2-chlorophenyl, 3-fluorophenyl, 3-chlorophenyl; 3-cyanophenyl, 3-methylphenyl, 3-trifluoromethylphenyl, 3-methoxyphenyl, 4-fluorophenyl, 4-chlorophenyl, 4-methoxyphenyl, 3,4-difluorophenyl, 3,5-difluorophenyl, 3-fluoro-5-chlorophenyl, 3-chloro-4-fluorophenyl, 2,4-dichlorophenyl or 3,4-dichlorophenyl), optionally substituted C_3 - C_{12} -cycloalkyl (e.g. cyclohexyl) or optionally substituted M_3 - M_{12} -heterocyclyl.

15


- 20 In connection with R^5 , substituted C_3 - C_{12} -cycloalkyl in particular includes C_3 - C_{12} -cycloalkyl, such as cyclopropyl or cyclohexyl, substituted with 1, 2 or 3 substituents selected from the group consisting of halogen, optionally substituted C_1 - C_6 -alkyl, halogenated C_1 - C_6 -alkyl, CN, hydroxy, C_1 - C_6 -alkoxy, halogenated C_1 - C_6 -alkoxy, amino, C_1 - C_6 -alkylamino, di- C_1 - C_6 -alkylamino and M_3 - M_{12} -heterocyclyl.

- 25 In connection with R^5 , substituted C_6 - C_{12} -aryl in particular includes C_6 - C_{12} -aryl, such as phenyl, substituted with 1, 2 or 3 substituents selected from the group consisting of halogen (e.g. F, Cl, Br), optionally substituted C_1 - C_6 -alkyl (e.g. methyl), halogenated C_1 - C_6 -alkyl (e.g. trifluoromethyl), CN, hydroxy, C_1 - C_6 -alkoxy (e.g. methoxy), halogenated C_1 - C_6 -alkoxy, amino, C_1 - C_6 -alkylamino, di- C_1 - C_6 -alkylamino and M_3 - M_{12} -heterocyclyl.

- 30 In connection with R^5 , substituted M_3 - M_{12} -heterocyclyl in particular includes M_3 - M_{12} -heterocyclyl substituted with 1, 2 or 3 substituents selected from the group consisting of halogen, optionally substituted C_1 - C_6 -alkyl, halogenated C_1 - C_6 -alkyl, CN, hydroxy, C_1 - C_6 -alkoxy, halogenated C_1 - C_6 -alkoxy, amino, C_1 - C_6 -alkylamino, di- C_1 - C_6 -alkylamino and M_3 - M_{12} -heterocyclyl.

- 35 In connection with R^5 , M_3 - M_{12} -heterocyclyl in particular is M_3 - M_{12} -heteroaryl.

Preferably, R^5 is optionally substituted C_6 - C_{12} -aryl, in particular as in the compounds of the formula:




wherein A, R¹, W, A¹, Q, Y, R⁶, n1, n2, X¹, R², A², R³, Y¹, R^{4a}, R^{4b}, X², X³ are as defined herein, and

- 5 R^{16a}, R^{16b}, R^{16c}, R^{16d}, R^{16e} independently are hydrogen, halogen (e.g. F, Cl or Br), optionally substituted C₁-C₆-alkyl (e.g. methyl), halogenated C₁-C₆-alkyl (e.g. trifluoromethyl), CN, hydroxy, C₁-C₆-alkoxy (e.g. methoxy), amino, C₁-C₆-alkylamino, di-C₁-C₆-alkylamino or M₃-M₁₂-heterocyclyl. Preferably, R^{16a}, R^{16b}, R^{16c}, R^{16d}, R^{16e} independently are hydrogen, halogen (e.g. F, Cl or Br), or halogenated C₁-C₆-alkyl (e.g. trifluoromethyl).

10

It is also preferred if R⁵ is optionally substituted M₆-M₁₂-heteroaryl, in particular as in the compounds of the formula:




wherein A, R¹, W, A¹, Q, Y, R⁶, n1, n2, X¹, R², A², R³, Y¹, R^{4a}, R^{4b}, X², X³ are as defined herein, and

15

- R^{16b} , R^{16c} , R^{16d} , R^{16e} independently are hydrogen, halogen (e.g. F, Cl or Br), optionally substituted C_1 - C_6 -alkyl (e.g. methyl), halogenated C_1 - C_6 -alkyl (e.g. trifluoromethyl), CN, hydroxy, C_1 - C_6 -alkoxy (e.g. methoxy), amino, C_1 - C_6 -alkylamino, di- C_1 - C_6 -alkylamino or M_3 - M_{12} -heterocyclyl. Preferably, R^{16b} , R^{16c} , R^{16d} , R^{16e} independently are hydrogen, halogen (e.g. F, Cl or Br), or halogenated C_1 - C_6 -alkyl (e.g. trifluoromethyl).

According to a particular embodiment, the invention relates to compounds of the formula:



- wherein A, R, R^2 , A^2 , R^3 , Y^1 , R^{4a} , R^{4b} , R^5 are as defined herein, R^5 preferably being optionally substituted aryl and in particular optionally substituted phenyl or optionally substituted heteroaryl and in particular optionally substituted pyridinyl as disclosed herein.

- In connection with R^5 or R^{16a} , R^{16b} , R^{16c} , R^{16d} , R^{16e} , substituted C_1 - C_6 -alkyl in particular includes C_1 - C_6 -alkyl, especially C_1 - C_4 -alkyl, substituted with 1, 2 or 3 substituents selected from the group consisting of hydroxy, C_1 - C_6 -alkoxy, amino, C_1 - C_6 -alkylamino, di- C_1 - C_6 -alkylamino and M_3 - M_{12} -heterocyclyl (e.g. morpholinyl or piperidinyl).

- According to a particular embodiment, R^{16a} , R^{16b} , R^{16d} , R^{16e} are hydrogen and R^{16c} is different from hydrogen (para-mono-substitution).

According to a further particular embodiment, R^{16a} , R^{16c} , R^{16d} , R^{16e} are hydrogen and R^{16b} is different from hydrogen (meta-mono-substitution).


- In connection with R^{16a} , R^{16b} , R^{16c} , R^{16d} , R^{16e} , M_3 - M_{12} -heterocyclyl in particular includes morpholinyl, imidazolyl and pyrazolyl.

R^7 is hydrogen or C_1 - C_6 -alkyl. Preferably, R^7 is hydrogen.

- R^8 is hydrogen, C_1 - C_6 -alkyl (e.g. methyl or ethyl), C_3 - C_{12} -cycloalkyl (e.g. cyclopropyl), amino- C_1 - C_6 -alkyl, optionally substituted C_6 - C_{12} -aryl- C_1 - C_4 -alkyl or M_3 - M_{12} -heterocyclyl (e.g. 3-azetidiny). Preferably, R^8 is hydrogen or C_1 - C_6 -alkyl (e.g. methyl or ethyl). In particular, R^8 is hydrogen.

- According to a particular embodiment, R^8 and R^1 together are C_1 - C_4 -alkylene (e.g. 1,2-ethylene or 1,3-propylene) so as that R^8 and R^1 together with the atom in Q to which R^1 is bound and the nitrogen atom to which R^8 is bound form an heterocyclic ring having, in particular, 4, 5 or 6 ring mem-

ber atoms (including the nitrogen atom and Q). With W and A¹ both being a bond, such a ring may be represented by the following partial structure:



5

wherein Q, is as defined herein (e.g. S(O)₂) and n is 0, 1, 2, 3 or 4.

10 R⁹ is hydrogen, C₁-C₆-alkyl, C₃-C₁₂-cycloalkyl-C₁-C₄-alkyl, halogenated C₁-C₄-alkyl, hydroxy-C₁-C₄-alkyl, C₁-C₆-alkoxy-C₁-C₄-alkyl, amino-C₁-C₄-alkyl, -CH₂CN, C₆-C₁₂-aryl-C₁-C₄-alkyl, C₃-C₁₂-cycloalkyl, -CHO, C₁-C₄-alkylcarbonyl, (halogenated C₁-C₄-alkyl)carbonyl, C₆-C₁₂-arylcarbonyl, C₁-C₄-alkoxycarbonyl, C₆-C₁₂-aryloxycarbonyl, C₁-C₆-alkylaminocarbonyl, C₂-C₆-alkenyl, -C(=NH)NH₂, -C(=NH)NHCN, C₁-C₆-alkylsulfonyl, C₆-C₁₂-arylsulfonyl, amino, -NO or M₃-M₁₂-heterocyclyl. Preferably, R⁹ is hydrogen or C₁-C₆-alkyl. In particular, R⁹ is hydrogen.

15

R¹⁰ is hydrogen or C₁-C₆-alkyl. Preferably, R¹⁰ is hydrogen.

R^{11a} is hydrogen or C₁-C₆-alkyl. Preferably, R^{11a} is hydrogen.

20 R^{11b} is hydrogen or C₁-C₆-alkyl. Preferably, R^{11b} is hydrogen.

R¹⁴ is hydrogen or C₁-C₆-alkyl. Preferably, R¹⁴ is hydrogen.

R¹⁵ is hydrogen or C₁-C₆-alkyl. Preferably, R¹⁵ is hydrogen.

25

R¹⁷ is hydrogen, C₁-C₆-alkyl or C₃-C₁₂-cycloalkyl. Preferably, R¹⁸ is hydrogen.

R¹⁸ is hydrogen, C₁-C₆-alkyl or C₃-C₁₂-cycloalkyl. Preferably, R¹⁸ is hydrogen.

30 Particular embodiments of compounds of the invention result if

A is a benzene ring

35 R¹ is C₁-C₆-alkyl (e.g. methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, n-pentyl), C₃-C₁₂-cycloalkyl-C₁-C₄-alkyl (e.g. cyclopropylmethyl, cyclobutylmethyl, cyclopentylmethyl, cyclohexylmethyl, cyclopropylethyl), halogenated C₁-C₆-alkyl (e.g. 3-fluoroprop-1-yl, 3-chloroprop-1-yl or 3,3,3-trifluoroprop-1-yl), tri-(C₁-C₄-alkyl)-silyl-C₁-C₄-alkyl (e.g. trimethylsilylethyl), C₁-C₆-alkoxy-C₁-C₄-alkyl (e.g. ethoxyethyl), C₃-C₁₂-cycloalkyl (e.g. cyclopropyl, cyclobutyl, cyclohexyl), C₂-C₆-alkenyl (e.g. prop-1,2-en-1-yl), optionally substituted C₆-C₁₂-aryl (e.g. phenyl, 3-methylphenyl), or optionally substituted M₃-M₁₂-heterocyclyl (e.g. 2-pyridyl, 3-pyridyl, 2-F-pyridin-3-yl, 5-F-pyridin-3yl, pyridazin-3-yl, 1-methyl-pyrrol-

- 3-yl, 2-thienyl, 3-thienyl, 4-methyl-2-thienyl, 5-methyl-2-thienyl, 5-chloro-2-thienyl, 2,5-dimethyl-3-thienyl, 3-furanyl, 5-methyl-2-furanyl, 2,5-dimethyl-3-furanyl, 1,2-diazol-4-yl, 1-methyl-1,2-diazol-4-yl, 1,3-dimethyl-1,2-diazol-4-yl, 1-ethyl-1,2-diazol-4-yl, 1-difluoromethyl-1,2-diazol-4-yl, 1-methyl-3-trifluoromethyl-1,2-diazol-4-yl, 1-methyl-1,3-diazol-4-yl, 2-methyl-1,3-diazol-4-yl, 1, 2-dimethyl-1,3-diazol-4-yl, 2-methyl-1,3-thiazol-5-yl, 2,4-dimethyl-1,3-thiazol-5-yl, 5-methylisoxazol-3-yl, 1-methyl-1,2,3-triazol-4-yl, 1-ethyl-1,2,3-triazol-4-yl, 1-methyl-1,2,4-triazol-3-yl, 3-pyrrolidinyl, 3-oxenatyl, 3-methyl-piperidinyl, 4-morpholinyl, 2,2-difluoro-1,3-benzodioxol-5-yl);
- W is a bond or NR⁷;
- 10 A¹ is a bond;
- Q is -S(O)₂- or -C(O)-;
- Y is NR⁸ or a bond;
- n1 is 1 or 2;
- n2 is 1 or 2;
- 15 R⁶ is hydrogen, C₁-C₆-alkyl (e.g. methyl), or two radicals R⁶ together with the carbon atom to which they are attached form a carbonyl group;
- X¹ is >N- or >CH-;
- R² is hydrogen, halogen (e.g. fluorine), or -CN;
- A² is -O-;
- 20 R³ is hydrogen or C₁-C₆-alkyl (e.g. methyl);
- Y¹ is a bond or substituted C₁-C₄-alkylene (e.g. methylene, 1,2-ethylene);
- R^{4a} is hydrogen, C₁-C₆-alkyl (e.g. methyl, ethyl, n-propyl, isopropyl, 2-methyl-but-4-yl, 2-methyl-prop-3-yl), C₃-C₁₂-cycloalkyl-C₁-C₄-alkyl (e.g. cyclopropylmethyl, cyclobutylmethyl, cyclopentylmethyl, 1-cyclopropyleth-2-yl, 1-cyclopentyleth-2-yl, cyclohexylmethyl),
- 25 halogenated C₁-C₄-alkyl (e.g. 2-fluoroethyl, 2,2,2-trifluoroethyl), C₃-C₁₂-cycloalkyl (e.g. cyclopropyl, cyclobutyl), -CHO, C₁-C₄-alkylcarbonyl (e.g. methylcarbonyl, ethylcarbonyl, isopropylcarbonyl), (halogenated C₁-C₄-alkyl)carbonyl (e.g. fluoromethylcarbonyl, difluoromethylcarbonyl, trifluoromethylcarbonyl, 1,1,1-trifluoroeth-2-ylcarbonyl, 1,1,1-trifluoroprop-3-ylcarbonyl), C₆-C₁₂-arylcarbonyl (e.g. phenylcarbonyl), C₁-C₄-
- 30 alkoxycarbonyl (e.g. ethoxycarbonyl, tert-butyloxycarbonyl), C₆-C₁₂-aryloxycarbonyl (e.g. phenoxycarbonyl) or optionally substituted M₃-M₁₂-heterocyclyl (e.g. 3-oxetanyl, 3-cyano-3-oxetanyl); or
- R^{4a} is optionally substituted C₁-C₄-alkylene that is bound to a carbon atom in Y¹ (e.g. methylene, 1,2-ethylene, 1,3-propylene);
- 35 R^{4b} is hydrogen or C₁-C₆-alkyl (e.g. methyl, ethyl); or
- R^{4a}, R^{4b} together are optionally substituted C₂-C₆-alkylene (e.g. 1,3-propylene, 1,4-butylene, 2-methyl-1,3-propylene, 2,2-dimethyl-1,3-propylene, 2-methyl-2-hydroxy-1,3-propylene, 2-fluoro-but-1,4-ylene, 1-oxo-but-1,4-ylene, -CH₂-cycloprop-1,2-ylene-CH₂-) wherein one -CH₂- of C₂-C₆-alkylene may be replaced by an oxygen atom (e.g. -CH₂-CH₂-O-CH₂-CH₂-);
- 40 X² is >CR^{12a}R^{12b};
- X³ is a bond;

- R^5 is optionally substituted phenyl (e.g. phenyl, 2-fluorophenyl, 2-chlorophenyl, 3-fluorophenyl, 3-chlorophenyl, 3-trifluoromethylphenyl, 3-cyanophenyl, 3-methylphenyl, 3-methoxyphenyl, 4-fluorophenyl, 4-chlorophenyl, 4-methoxyphenyl, 3,4-difluorophenyl, 3,5-difluorophenyl, 3-fluoro-5-chlorophenyl, 3-chloro-4-fluorophenyl, 3,4-dichlorophenyl, 2,4-dichlorophenyl) or optionally substituted C_3 - C_{12} -cycloalkyl (e.g. cyclohexyl);
- R^7 is hydrogen or C_1 - C_6 -alkyl (e.g. methyl);
- R^8 is hydrogen, C_1 - C_6 -alkyl (e.g. methyl or ethyl), or C_3 - C_{12} -cycloalkyl (e.g. cyclopropyl); or R^8, R^1 together are C_1 - C_4 -alkylene (e.g. 1,3-propylene);
- R^{12a} is hydrogen, C_1 - C_6 -alkyl (e.g. methyl or ethyl);
- R^{12b} is hydrogen;
- R^{12a}, R^{12b} together are optionally substituted C_2 - C_4 -alkylene (e.g. 1,3-propylene).
- 15 Further particular embodiments of compounds of the invention result if
- A is a benzene ring
- R^1 is C_1 - C_6 -alkyl (e.g. ethyl, n-propyl), C_3 - C_{12} -cycloalkyl- C_1 - C_4 -alkyl (e.g. cyclopropylmethyl), or optionally substituted M_3 - M_{12} -heterocyclyl (e.g. 3 e.g. 1-methyl-1,2-diazol-4-yl, 1-methyl-1,3-diazol-4-yl, 1-methyl-1,2,3-triazol-4-yl, 1-ethyl-1,2,3-triazol-4-yl, 2-F-pyridin-3-yl, 5-F-pyridin-3yl, pyridazin-3-yl);
- W is a bond;
- A^1 is a bond;
- Q is $-S(O)_2-$;
- Y is NR^8 ;
- $n1$ is 1;
- $n2$ is 1;
- R^6 is hydrogen;
- X^1 is $>N-$ or $>CH-$;
- R^2 is hydrogen or halogen (e.g. fluorine);
- A^2 is $-O-$;
- R^3 is hydrogen;
- Y^1 is a bond;
- R^{4a} is hydrogen, C_1 - C_6 -alkyl (e.g. methyl, ethyl, n-propyl), C_3 - C_{12} -cycloalkyl- C_1 - C_4 -alkyl (e.g. cyclopropylmethyl) or C_3 - C_{12} -cycloalkyl (e.g. cyclobutyl);
- R^{4b} is hydrogen or C_1 - C_6 -alkyl (e.g. ethyl); or R^{4a}, R^{4b} together are C_2 - C_6 -alkylene (e.g. 1,3-propylene);
- X^2 is $>CR^{12a}R^{12b}$;
- X^3 is a bond;
- R^5 is optionally substituted phenyl (e.g. phenyl);
- R^8 is hydrogen, or
- R^{12a} is hydrogen;

R^{12b} is hydrogen.

Further particular compounds of the present invention are the individual aminochromane, amino-thiochromane and amino-1,2,3,4-tetrahydroquinoline derivatives of the formula (Id) as listed in the following tables 1 to 24 and physiologically tolerated salts thereof:




Table 1

Compounds of the formula (Ia) wherein R^6 is as defined herein and in particular represents H, $-A^2-$ is as defined herein and in particular represents $-O-$, $-Y^1-$ is as defined herein and in particular represents a bond, R^2 is hydrogen, R^3 is as defined herein and in particular represents hydrogen, R^{16} is hydrogen and the combination of R^1 , $-X^1-$, $n1$, $n2$, $>CR^{12a}R^{12b}$, R^{4a} , R^{4b} for a compound in each case corresponds to one line of Table A (A-1 to A-88).

Table 2

Compounds of the formula (Ia) wherein R^6 is as defined herein and in particular represents H, $-A^2-$ is as defined herein and in particular represents $-O-$, $-Y^1-$ is as defined herein and in particular represents a bond, R^2 is hydrogen, R^3 is as defined herein and in particular represents hydrogen, R^{16} is 3-F and the combination of R^1 , $-X^1-$, $n1$, $n2$, $>CR^{12a}R^{12b}$, R^{4a} , R^{4b} for a compound in each case corresponds to one line of Table A (A-1 to A-88).

Table 3

Compounds of the formula (Ia) wherein R^6 is as defined herein and in particular represents H, $-A^2-$ is as defined herein and in particular represents $-O-$, $-Y^1-$ is as defined herein and in particular represents a bond, R^2 is hydrogen, R^3 is as defined herein and in particular represents hydrogen, R^{16} is 3-Cl and the combination of R^1 , $-X^1-$, $n1$, $n2$, $>CR^{12a}R^{12b}$, R^{4a} , R^{4b} for a compound in each case corresponds to one line of Table A (A-1 to A-88).

Table 4

Compounds of the formula (Ia) wherein R^6 is as defined herein and in particular represents H, $-A^2-$ is as defined herein and in particular represents $-O-$, $-Y^1-$ is as defined herein and in particular represents a bond, R^2 is hydrogen, R^3 is as defined herein and in particular represents hydrogen, R^{16} is

3-CF₃ and the combination of R¹, -X¹-, n1, n2, >CR^{12a}R^{12b}, R^{4a}, R^{4b} for a compound in each case corresponds to one line of Table A (A-1 to A-88).

Table 5

- 5 Compounds of the formula (Ia) wherein R⁶ is as defined herein and in particular represents H, -A²- is as defined herein and in particular represents -O-, -Y¹- is as defined herein and in particular represents a bond, R² is hydrogen, R³ is as defined herein and in particular represents hydrogen, R¹⁶ is 4-F and the combination of R¹, -X¹-, n1, n2, >CR^{12a}R^{12b}, R^{4a}, R^{4b} for a compound in each case corresponds to one line of Table A (A-1 to A-88).

10

Table 6

- Compounds of the formula (Ia) wherein R⁶ is as defined herein and in particular represents H, -A²- is as defined herein and in particular represents -O-, -Y¹- is as defined herein and in particular represents a bond, R² is hydrogen, R³ is as defined herein and in particular represents hydrogen, R¹⁶ is 4-Cl and the combination of R¹, -X¹-, n1, n2, >CR^{12a}R^{12b}, R^{4a}, R^{4b} for a compound in each case corresponds to one line of Table A (A-1 to A-88).

15

Table 7

- Compounds of the formula (Ia) wherein R⁶ is as defined herein and in particular represents H, -A²- is as defined herein and in particular represents -O-, -Y¹- is as defined herein and in particular represents a bond, R² is 5-F, R³ is as defined herein and in particular represents hydrogen, R¹⁶ is hydrogen and the combination of R¹, -X¹-, n1, n2, >CR^{12a}R^{12b}, R^{4a}, R^{4b} for a compound in each case corresponds to one line of Table A (A-1 to A-88).

20

- 25 Table 8

- Compounds of the formula (Ia) wherein R⁶ is as defined herein and in particular represents H, -A²- is as defined herein and in particular represents -O-, -Y¹- is as defined herein and in particular represents a bond, R² is 5-F, R³ is as defined herein and in particular represents hydrogen, R¹⁶ is 3-F and the combination of R¹, -X¹-, n1, n2, >CR^{12a}R^{12b}, R^{4a}, R^{4b} for a compound in each case corresponds to one line of Table A (A-1 to A-88).

30

Table 9

- Compounds of the formula (Ia) wherein R⁶ is as defined herein and in particular represents H, -A²- is as defined herein and in particular represents -O-, -Y¹- is as defined herein and in particular represents a bond, R² is 5-F, R³ is as defined herein and in particular represents hydrogen, R¹⁶ is 3-Cl and the combination of R¹, -X¹-, n1, n2, >CR^{12a}R^{12b}, R^{4a}, R^{4b} for a compound in each case corresponds to one line of Table A (A-1 to A-88).

35

Table 10

- 40 Compounds of the formula (Ia) wherein R⁶ is as defined herein and in particular represents H, -A²- is as defined herein and in particular represents -O-, -Y¹- is as defined herein and in particular represents a bond, R² is 5-F, R³ is as defined herein and in particular represents hydrogen, R¹⁶ is 3-CF₃

and the combination of R^1 , $-X^1$ -, $n1$, $n2$, $>CR^{12a}R^{12b}$, R^{4a} , R^{4b} for a compound in each case corresponds to one line of Table A (A-1 to A-88).

Table 11

- 5 Compounds of the formula (Ia) wherein R^6 is as defined herein and in particular represents H, $-A^2$ - is as defined herein and in particular represents $-O-$, $-Y^1$ - is as defined herein and in particular represents a bond, R^2 is 5-F, R^3 is as defined herein and in particular represents hydrogen, R^{16} is 4-F and the combination of R^1 , $-X^1$ -, $n1$, $n2$, $>CR^{12a}R^{12b}$, R^{4a} , R^{4b} for a compound in each case corresponds to one line of Table A (A-1 to A-88).

10

Table 12

- 15 Compounds of the formula (Ia) wherein R^6 is as defined herein and in particular represents H, $-A^2$ - is as defined herein and in particular represents $-O-$, $-Y^1$ - is as defined herein and in particular represents a bond, R^2 is 5-F, R^3 is as defined herein and in particular represents hydrogen, R^{16} is 4-Cl and the combination of R^1 , $-X^1$ -, $n1$, $n2$, $>CR^{12a}R^{12b}$, R^{4a} , R^{4b} for a compound in each case corresponds to one line of Table A (A-1 to A-88).

Table 13

- 20 Compounds of the formula (Ia) wherein R^6 is as defined herein and in particular represents H, $-A^2$ - is as defined herein and in particular represents $-O-$, $-Y^1$ - is as defined herein and in particular represents a bond, R^2 is 7-F, R^3 is as defined herein and in particular represents hydrogen, R^{16} is hydrogen and the combination of R^1 , $-X^1$ -, $n1$, $n2$, $>CR^{12a}R^{12b}$, R^{4a} , R^{4b} for a compound in each case corresponds to one line of Table A (A-1 to A-88).

- 25 Table 14

- Compounds of the formula (Ia) wherein R^6 is as defined herein and in particular represents H, $-A^2$ - is as defined herein and in particular represents $-O-$, $-Y^1$ - is as defined herein and in particular represents a bond R^2 is 7-F, R^3 is as defined herein and in particular represents hydrogen, R^{16} is 3-F and the combination of R^1 , $-X^1$ -, $n1$, $n2$, $>CR^{12a}R^{12b}$, R^{4a} , R^{4b} for a compound in each case corresponds to one line of Table A (A-1 to A-88).

30

Table 15

- 35 Compounds of the formula (Ia) wherein R^6 is as defined herein and in particular represents H, $-A^2$ - is as defined herein and in particular represents $-O-$, $-Y^1$ - is as defined herein and in particular represents a bond, R^2 is 7-F, R^3 is as defined herein and in particular represents hydrogen, R^{16} is 3-Cl and the combination of R^1 , $-X^1$ -, $n1$, $n2$, $>CR^{12a}R^{12b}$, R^{4a} , R^{4b} for a compound in each case corresponds to one line of Table A (A-1 to A-88).

Table 16

- 40 Compounds of the formula (Ia) wherein R^6 is as defined herein and in particular represents H, $-A^2$ - is as defined herein and in particular represents $-O-$, $-Y^1$ - is as defined herein and in particular represents a bond, R^2 is 7-F, R^3 is as defined herein and in particular represents hydrogen, R^{16} is 3-CF₃

and the combination of R^1 , $-X^1$ -, $n1$, $n2$, $>CR^{12a}R^{12b}$, R^{4a} , R^{4b} for a compound in each case corresponds to one line of Table A (A-1 to A-88).

Table 17

- 5 Compounds of the formula (Ia) wherein R^6 is as defined herein and in particular represents H, $-A^2$ - is as defined herein and in particular represents $-O-$, $-Y^1$ - is as defined herein and in particular represents a bond, R^2 is 7-F, R^3 is as defined herein and in particular represents hydrogen, R^{16} is 4-F and the combination of R^1 , $-X^1$ -, $>CR^{12a}R^{12b}$, $n1$, $n2$, R^{4a} , R^{4b} for a compound in each case corresponds to one line of Table A (A-1 to A-88).

10

Table 18

- 15 Compounds of the formula (Ia) wherein R^6 is as defined herein and in particular represents H, $-A^2$ - is as defined herein and in particular represents $-O-$, $-Y^1$ - is as defined herein and in particular represents a bond, R^2 is 7-F, R^3 is as defined herein and in particular represents hydrogen, R^{16} is 4-Cl and the combination of R^1 , $-X^1$ -, $n1$, $n2$, $>CR^{12a}R^{12b}$, R^{4a} , R^{4b} for a compound in each case corresponds to one line of Table A (A-1 to A-88).

Table 19

- 20 Compounds of the formula (Ia) wherein R^6 is as defined herein and in particular represents H, $-A^2$ - is as defined herein and in particular represents $-O-$, $-Y^1$ - is as defined herein and in particular represents a bond, R^2 is 8-F, R^3 is as defined herein and in particular represents hydrogen, R^{16} is hydrogen and the combination of R^1 , $-X^1$ -, $n1$, $n2$, $>CR^{12a}R^{12b}$, R^{4a} , R^{4b} for a compound in each case corresponds to one line of Table A (A-1 to A-88).

- 25 Table 20

- Compounds of the formula (Ia) wherein R^6 is as defined herein and in particular represents H, $-A^2$ - is as defined herein and in particular represents $-O-$, $-Y^1$ - is as defined herein and in particular represents a bond, R^2 is 8-F, R^3 is as defined herein and in particular represents hydrogen, R^{16} is 3-F and the combination of R^1 , $-X^1$ -, $>CR^{12a}R^{12b}$, $n1$, $n2$, R^{4a} , R^{4b} for a compound in each case corresponds to one line of Table A (A-1 to A-88).

30

Table 21

- 35 Compounds of the formula (Ia) wherein R^6 is as defined herein and in particular represents H, $-A^2$ - is as defined herein and in particular represents $-O-$, $-Y^1$ - is as defined herein and in particular represents a bond, R^2 is 8-F, R^3 is as defined herein and in particular represents hydrogen, R^{16} is 3-Cl and the combination of R^1 , $-X^1$ -, $n1$, $n2$, $>CR^{12a}R^{12b}$, R^{4a} , R^{4b} for a compound in each case corresponds to one line of Table A (A-1 to A-88).

Table 22

- 40 Compounds of the formula (Ia) wherein R^6 is as defined herein and in particular represents H, $-A^2$ - is as defined herein and in particular represents $-O-$, $-Y^1$ - is as defined herein and in particular represents a bond, R^2 is 8-F, R^3 is as defined herein and in particular represents hydrogen, R^{16} is 3-CF₃

and the combination of R^1 , $-X^1$ -, n1, n2, $>CR^{12a}R^{12b}$, R^{4a} , R^{4b} for a compound in each case corresponds to one line of Table A (A-1 to A-88).



Table 23







- 5 Compounds of the formula (Ia) wherein R^6 is as defined herein and in particular represents H, $-A^2$ - is as defined herein and in particular represents $-O$ -, $-Y^1$ - is as defined herein and in particular represents a bond, R^2 is 8-F, R^3 is as defined herein and in particular represents hydrogen, R^{16} is 4-F and the combination of R^1 , $-X^1$ -, n1, n2, $>CR^{12a}R^{12b}$, R^{4a} , R^{4b} for a compound in each case corresponds to one line of Table A (A-1 to A-88).






10







Table 24







- 15 Compounds of the formula (Ia) wherein R^6 is as defined herein and in particular represents H, $-A^2$ - is as defined herein and in particular represents $-O$ -, $-Y^1$ - is as defined herein and in particular represents a bond, R^2 is 8-F, R^3 is as defined herein and in particular represents hydrogen, R^{16} is 4-Cl and the combination of R^1 , $-X^1$ -, n1, n2, $>CR^{12a}R^{12b}$, R^{4a} , R^{4b} for a compound in each case corresponds to one line of Table A (A-1 to A-88).






	R^1	$-X^1$ -	n1	n2	$>CR^{12a}R^{12b}$	R^{4a} , R^{4b}
A-1.		$>N$ -	1	1	$-CH_2-$	H, H
A-2.		$>N$ -	1	1	$-CH_2-$	H, H
A-3.		$>N$ -	1	1	$-CH_2-$	H, H
A-4.		$>N$ -	1	1	$-CH_2-$	H, H
A-5.		$>N$ -	1	1	$-CH_2-$	H, H







	R ¹	-X ¹ -	n1	n2	>CR ^{12a} R ^{12b}	R ^{4a} , R ^{4b}
A-6.		>N-	1	1	-CH ₂ -	H, H
A-7.		>N-	1	1	-CH ₂ -	H, H
A-8.		>N-	1	1	-CH ₂ -	H, H
A-9.		>N-	1	1	-CH ₂ -	H, H
A-10.		>N-	1	1	-CH ₂ -	H, H
A-11.		>N-	1	1	-CH ₂ -	H, H







	R ¹	-X ¹ -	n1	n2	>CR ^{12a} R ^{12b}	R ^{4a} , R ^{4b}
A-12.		>CH-	1	1	-CH ₂ -	H, H
A-13.		>CH-	1	1	-CH ₂ -	H, H
A-14.		>CH-	1	1	-CH ₂ -	H, H
A-15.		>CH-	1	1	-CH ₂ -	H, H
A-16.		>CH-	1	1	-CH ₂ -	H, H
A-17.		>CH-	1	1	-CH ₂ -	H, H
A-18.		>CH-	1	1	-CH ₂ -	H, H
A-19.		>CH-	1	1	-CH ₂ -	H, H






	R ¹	-X ¹ -	n1	n2	>CR ^{12a} R ^{12b}	R ^{4a} , R ^{4b}
A-20.		>CH-	1	1	-CH ₂ -	-H, H
A-21.		>CH-	1	1	-CH ₂ -	H, H
A-22.		>CH-	1	1	-CH ₂ -	H, H
A-23.		>N-	1	1	-CH ₂ -	-CH ₃ , H
A-24.		>N-	1	1	-CH ₂ -	-CH ₃ , H
A-25.		>N-	1	1	-CH ₂ -	-CH ₃ , H
A-26.		>N-	1	1	-CH ₂ -	-CH ₃ , H
A-27.		>N-	1	1	-CH ₂ -	-CH ₃ , H







	R ¹	-X ¹ -	n1	n2	>CR ^{12a} R ^{12b}	R ^{4a} , R ^{4b}
A-28.		>N-	1	1	-CH ₂ -	-CH ₃ , H
A-29.		>N-	1	1	-CH ₂ -	-CH ₃ , H
A-30.		>N-	1	1	-CH ₂ -	-CH ₃ , H
A-31.		>N-	1	1	-CH ₂ -	-CH ₃ , H
A-32.		>N-	1	1	-CH ₂ -	-CH ₃ , H
A-33.		>N-	1	1	-CH ₂ -	-CH ₃ , H







	R ¹	-X ¹ -	n1	n2	>CR ^{12a} R ^{12b}	R ^{4a} , R ^{4b}
A-34.		>CH-	1	1	-CH ₂ -	-CH ₃ , H
A-35.		>CH-	1	1	-CH ₂ -	-CH ₃ , H
A-36.		>CH-	1	1	-CH ₂ -	-CH ₃ , H
A-37.		>CH-	1	1	-CH ₂ -	-CH ₃ , H
A-38.		>CH-	1	1	-CH ₂ -	-CH ₃ , H
A-39.		>CH-	1	1	-CH ₂ -	-CH ₃ , H
A-40.		>CH-	1	1	-CH ₂ -	-CH ₃ , H
A-41.		>CH-	1	1	-CH ₂ -	-CH ₃ , H






	R ¹	-X ¹ -	n1	n2	>CR ^{12a} R ^{12b}	R ^{4a} , R ^{4b}
A-42.		>CH-	1	1	-CH ₂ -	-CH ₃ , H
A-43.		>CH-	1	1	-CH ₂ -	-CH ₃ , H
A-44.		>CH-	1	1	-CH ₂ -	-CH ₃ , H
A-45.		>N-	1	1	-CH ₂ -	-(CH ₂) ₃ -
A-46.		>N-	1	1	-CH ₂ -	-(CH ₂) ₃ -
A-47.		>N-	1	1	-CH ₂ -	-(CH ₂) ₃ -
A-48.		>N-	1	1	-CH ₂ -	-(CH ₂) ₃ -
A-49.		>N-	1	1	-CH ₂ -	-(CH ₂) ₃ -




	R ¹	-X ¹ -	n1	n2	>CR ^{12a} R ^{12b}	R ^{4a} , R ^{4b}
A-50.		>N-	1	1	-CH ₂ -	-(CH ₂) ₃ -
A-51.		>N-	1	1	-CH ₂ -	-(CH ₂) ₃ -
A-52.		>N-	1	1	-CH ₂ -	-(CH ₂) ₃ -
A-53.		>N-	1	1	-CH ₂ -	-(CH ₂) ₃ -
A-54.		>N-	1	1	-CH ₂ -	-(CH ₂) ₃ -
A-55.		>N-	1	1	-CH ₂ -	-(CH ₂) ₃ -

	R ¹	-X ¹ -	n1	n2	>CR ^{12a} R ^{12b}	R ^{4a} , R ^{4b}
A-56.		>CH-	1	1	-CH ₂ -	-(CH ₂) ₃ -
A-57.		>CH-	1	1	-CH ₂ -	-(CH ₂) ₃ -
A-58.		>CH-	1	1	-CH ₂ -	-(CH ₂) ₃ -
A-59.		>CH-	1	1	-CH ₂ -	-(CH ₂) ₃ -
A-60.		>CH-	1	1	-CH ₂ -	-(CH ₂) ₃ -
A-61.		>CH-	1	1	-CH ₂ -	-(CH ₂) ₃ -
A-62.		>CH-	1	1	-CH ₂ -	-(CH ₂) ₃ -
A-63.		>CH-	1	1	-CH ₂ -	-(CH ₂) ₃ -

	R ¹	-X ¹ -	n1	n2	>CR ^{12a} R ^{12b}	R ^{4a} , R ^{4b}
A-64.		>CH-	1	1	-CH ₂ -	-(CH ₂) ₃ -
A-65.		>CH-	1	1	-CH ₂ -	-(CH ₂) ₃ -
A-66.		>CH-	1	1	-CH ₂ -	-(CH ₂) ₃ -
A-67.		>N-	1	1	-CH ₂ -	-(CH ₂) ₄ -
A-68.		>N-	1	1	-CH ₂ -	-(CH ₂) ₄ -
A-69.		>N-	1	1	-CH ₂ -	-(CH ₂) ₄ -
A-70.		>N-	1	1	-CH ₂ -	-(CH ₂) ₄ -
A-71.		>N-	1	1	-CH ₂ -	-(CH ₂) ₄ -

	R ¹	-X ¹ -	n1	n2	>CR ^{12a} R ^{12b}	R ^{4a} , R ^{4b}
A-72.		>N-	1	1	-CH ₂ -	-(CH ₂) ₄ -
A-73.		>N-	1	1	-CH ₂ -	-(CH ₂) ₄ -
A-74.		>N-	1	1	-CH ₂ -	-(CH ₂) ₄ -
A-75.		>N-	1	1	-CH ₂ -	-(CH ₂) ₄ -
A-76.		>N-	1	1	-CH ₂ -	-(CH ₂) ₄ -
A-77.		>N-	1	1	-CH ₂ -	-(CH ₂) ₄ -


	R ¹	-X ¹ -	n1	n2	>CR ^{12a} R ^{12b}	R ^{4a} , R ^{4b}
A-78.		>CH-	1	1	-CH ₂ -	-(CH ₂) ₄ -
A-79.		>CH-	1	1	-CH ₂ -	-(CH ₂) ₄ -
A-80.		>CH-	1	1	-CH ₂ -	-(CH ₂) ₄ -
A-81.		>CH-	1	1	-CH ₂ -	-(CH ₂) ₄ -
A-82.		>CH-	1	1	-CH ₂ -	-(CH ₂) ₄ -
A-83.		>CH-	1	1	-CH ₂ -	-(CH ₂) ₄ -
A-84.		>CH-	1	1	-CH ₂ -	-(CH ₂) ₄ -
A-85.		>CH-	1	1	-CH ₂ -	-(CH ₂) ₄ -

	R ¹	-X ¹ -	n1	n2	>CR ^{12a} R ^{12b}	R ^{4a} , R ^{4b}
A-86.		>CH-	1	1	-CH ₂ -	-(CH ₂) ₄ -
A-87.		>CH-	1	1	-CH ₂ -	-(CH ₂) ₄ -
A-88.		>CH-	1	1	-CH ₂ -	-(CH ₂) ₄ -

- Still further particular compounds of the present invention are the compounds disclosed in preparation examples and physiologically tolerated salts thereof. These include for each preparation example the exemplified compound as well as the corresponding free base and any other physiologically tolerated salts of the free base (if the exemplified compound is a salt), or any physiologically tolerated salt of the free base (if the exemplified compound is a free base). These further include enantiomers, diastereomers, tautomers and any other isomeric forms of said compounds, be they explicitly or implicitly disclosed.
- 5 The compounds of the formula (I) can be prepared by analogy to methods which are well known in the art. Suitable methods for the preparation of compounds of formula (I) are outlined in the following schemes.
- 10

- The process depicted in scheme 1 is useful for obtaining aminochromanes of general formula 5, wherein X⁴ is -O-.
- 15

Scheme 1:




As shown in scheme 1, the compound of general formula 1 can be transferred into the corresponding hydroxylamine 2 (e.g. in presence of $\text{NH}_2\text{OH HCl}$). The hydroxyl group can be converted to a leaving group (e.g. tosyl or mesyl) to yield compounds of the general formula 3. Compounds 3 readily undergo Neber rearrangement in the presence of a base (e.g. NaOEt , J. Med. Chem. 1988, 31, 2178) followed by protection with a suitable protecting group L^2 (e.g. $\text{L}^2 = \text{COOEt}$) to give the compound of general formula 5.

- 10 In scheme 1, the variables R^2 and R^3 are as defined herein and L^1 is a suitable protecting group (e.g. $\text{L}^1 = \text{Me}$).

Compounds of the general formula 1 are also readily accessible from common bulk chemicals as described in scheme 2. The process depicted in scheme 2 is useful for obtaining aminochromanes of general formula 1, wherein X^4 is $-\text{O}-$ and L^1 is a suitable protecting group (e.g. $\text{L}^1 = \text{Me}$).

Scheme 2:




- 20 Phenols of the general formula 1-1 can be reacted with 3-halogenated carboxylic acids like 1-2 in presence of a base as described in the literature (e.g. potassium hydroxide, sodium hydrogencarbonate, J. Med. Chem. 1982, 25, 393) to give compounds of the general formula 1-3. In presence of an acid these compounds undergo acylation reactions to form compounds of the general formula 1 (e.g. polyphosphoric acid, J. Med. Chem. 1982, 25, 393).

25

The process depicted in scheme 3 is useful for obtaining aminochromanes, wherein X^1 is $>\text{N}-$, X^4 is $-\text{O}-$, Y is $-\text{NR}^8-$, and Q is $-\text{S}(\text{O})_2-$.

Scheme 3:




- Aminochromanes of the general formula 5 can be reacted with a Grignard reagent to give the alcohols of the general formula 6. In presence of an acid (e.g. aqueous hydrochloric acid) these alcohols undergo elimination to the corresponding alkenes of general formula 7. Reduction of compounds of type 7 (e.g. by hydrogenation with H_2 and Pd/C in presence of an acid (e.g. ammonium formate)) leads to aminochromanes of the general formula 8. The free phenols 9 can be accessed via removal of the protecting group L^1 (e.g. for $L^1 = Me$ by treating compounds 8 with boron tribromide). The phenols of general formula 9 can be transferred into the triflates 10 in presence of trifluoromethanesulfonic anhydride or 2-[N,N-Bis(trifluoromethylsulfonyl)amino]-5-chloropyridine. As described in the literature (e.g. Chem. Sci. 2011, 2, 27-50) triflate 10 can undergo a Buchwald-Hartwig amination with a cyclic amine in the presence of a palladium source (e.g. Pd(II) acetate), a ligand (e.g. dicyclohexyl(2',4',6'-triisopropyl-[1,1'-biphenyl]-2-yl)phosphine) and a base (e.g. cesium carbonate) to yield compounds of the general formula 11. Alternatively to the triflates 10 the corresponding nonaflates or bromides can be used to prepare compound 11. Deprotection of the Boc-group in presence of an acid (e.g. trifluoroacetic acid or formic acid) leads to the compounds of the general formula 12. Treatment with sulfonyl chlorides in presence of a base (e.g. N,N-dimethylaminopyridine or pyridine) yields compound 13. Deprotection of the protecting group L^2 (for $L^2 = ethylcarbamate$ e.g. ethanolic potassium hydroxide) will lead to the free amine of the general formula 14. Reductive amination using the corresponding ketones or aldehydes in presence of

a reduction reagent (e.g. sodiumcyanoborohydride and glacial acetic acid) or amide formation followed by subsequent reduction yields the corresponding higher alkylated amines of the general formula 15. Reduction of the ethylcarbamate (L^3) of compound 13 (e.g. using lithium aluminum hydride) leads directly to compounds of the general formula 15 with R^{4a} =methyl and R^{4b} =hydrogen or vice versa.

In scheme 3, the variables R^1 , W, A^1 , R^2 , R^3 , R^{4a} , R^{4b} , R^5 , R^6 , R^8 , X^2 , X^3 , n1 and n2 are as defined herein, and L^1 and L^2 are suitable protecting groups (e.g. L^1 = Me, and L^2 = COOEt).

The process depicted in scheme 4 is useful for obtaining aminochromanes, wherein X^1 is $-\text{CH}-$, Y is $-\text{NR}^8-$, and Q is $-\text{S}(\text{O})_2-$.

Scheme 4:



15

Triflates of the general formula 10 can be reacted with the corresponding alkyliodides in the presence of zinc and palladium (e.g. zinc, TMSCl, 1,2-dibromoethane, Pd(dba)₂, and dppf) to undergo a Negishi-coupling (Austr. J. Chem. 2004, 57, 107) and lead to compounds of the general formula 16. Alternatively, a Suzuki-coupling of the triflates 10 with the corresponding boron reagents (boronic acid, ester or trifluoroborates) in presence of a palladium source (e.g. palladiumdibenzylidene acetone), a ligand (e.g. 2-dicyclohexyl-phosphino-2',6'-diisopropoxybiphenyl) and a base (e.g. cesium carbonate) leads to the compounds of the general formula 16. Deprotection of the Boc-group in presence of an acid (e.g. trifluoroacetic acid or formic acid) leads to the compounds of the general formula 17. Treatment with sulfonyl chlorides in presence of a base (e.g. N,N-

20

dimethylaminopyridine or pyridine) yields compound 18. Deprotection of the protecting group L^3 (for L^3 =ethylcarbamate e.g. ethanolic potassium hydroxide) will lead to the free amine of the general formula 19. Reductive amination using the corresponding ketones or aldehydes in presence of a reduction reagent (e.g. sodiumcyanoborohydride and glacial acetic acid) or amide formation followed by subsequent reduction yields the corresponding higher alkylated amines of the general formula 20. Reduction of the ethylcarbamate (L^3) of compound 18 (e.g. using lithium aluminum hydride) leads directly to compounds of the general formula 20 with R^{4a} =methyl and R^{4b} =hydrogen or vice versa.

- 10 In scheme 4, the variables R^1 , W, A^1 , R^2 , R^3 , R^{4a} , R^{4b} , R^5 , R^6 , R^8 , X^2 , X^3 , n1 and n2 are as defined herein, and L^3 is a suitable protecting groups (e.g. L^3 = COOEt).

The compounds of the formula (I) are capable of inhibiting the activity of glycine transporter, in particular glycine transporter 1 (GlyT1).

15

The utility of the compounds in accordance with the present invention as inhibiting the glycine transporter activity, in particular GlyT1 activity, may be demonstrated by methodology known in the art. For instance, human GlyT1c expressing recombinant hGlyT1c_5_CHO cells can be used for measuring glycine uptake and its inhibition (IC_{50}) by a compound of formula (I).

20

Amongst the compounds of the formula (I) those are preferred which achieve effective inhibition at low concentrations. In particular, compounds of the formula (I) are preferred which inhibit glycine transporter 1 (GlyT1) at a level of $IC_{50} < 1 \mu\text{Mol}$, more preferably at a level of $IC_{50} < 0.5 \mu\text{Mol}$, particularly preferably at a level of $IC_{50} < 0.2 \mu\text{Mol}$ and most preferably at a level of $IC_{50} < 0.1 \mu\text{Mol}$.

25

Compounds of formula (I) combine high affinity with high metabolic stability.

- 30 The metabolic stability of a compound can be measured for example by incubating a solution of this compound with liver microsomes from particular species (for example rat, dog or human) and determining the half-life of the compound under these conditions (RS Obach, Curr Opin Drug Discov Devel. 2001, 4, 36-44). It is possible in this connection to conclude from an observed longer half-life that the metabolic stability of the compound is improved. The stability in the presence of human liver microsomes is of particular interest because it makes it possible to predict the metabolic degradation of the compound in the human liver. Compounds with increased metabolic stability (measured in the liver microsome test) are therefore probably also degraded more slowly in the liver. The slower metabolic degradation in the liver may lead to higher and/or longer-lasting concentrations (active levels) of the compound in the body, so that the elimination half-life of the compounds of the invention is increased. Increased and/or longer-lasting active levels may lead to a better activity of the compound in therapeutic treatment. In addition, an improved metabolic stability may lead to an increased bioavailability after oral administration, because the compound is subject, after absorption in the intestine, to less metabolic degradation in the liver (so-called first pass
- 35
- 40

effect). An increased oral bioavailability may, owing to an increased concentration (active level) of the compound, lead to a better activity of the compound after oral administration.

5 Amongst the compounds of the formula (I) those are particularly preferred which display good to moderate metabolic stability towards human liver microsomes. In particular, compounds of the formula (I) are preferred which display a microsomal clearance at a level of $mCl_{int,u} < 500 \text{ L/h/kg}$, more preferably at a level of $mCl_{int,u} < 100 \text{ L/h/kg}$, particularly preferably at a level of $mCl_{int,u} < 50 \text{ L/h/kg}$, and most preferably at a level of $mCl_{int,u} < 5 \text{ L/h/kg}$.

10 Further, compounds of formula (I) exhibit favorable efflux properties which may lead to enhanced oral bioavailability and/or increased brain availability. According to a particular embodiment, compounds of the invention combine high affinity and high metabolic stability with favorable efflux properties.

15 The efflux properties of a compound can be measured in well-known assays (e.g. Caco-2, MDCK assay).

The compounds of the formula (I) according to the present invention are thus useful as pharmaceuticals.

20 The present invention therefore also relates to pharmaceutical compositions which comprise an inert carrier and a compound of the formula (I).

25 The present invention also relates to the use of the compounds of the formula (I) in the manufacture of a medicament for inhibiting the glycine transporter GlyT1, and to corresponding methods of inhibiting the glycine transporter GlyT1.

30 The NMDA receptor is central to a wide range of CNS processes, and its role in a variety of diseases in humans or other species has been described. GlyT1 inhibitors slow the removal of glycine from the synapse, causing the level of synaptic glycine to rise. This in turn increases the occupancy of the glycine binding site on the NMDA receptor, which increases activation of the NMDA receptor following glutamate release from the presynaptic terminal. Glycine transport inhibitors and in particular inhibitors of the glycine transporter GlyT1 are thus known to be useful in treating a variety of neurologic and psychiatric disorders. Further, glycine A receptors play a role in a variety of
35 diseases in humans or other species. Increasing extracellular glycine concentrations by inhibiting glycine transport may enhance the activity of glycine A receptors. Glycine transport inhibitors and in particular inhibitors of the glycine transporter GlyT1 are thus useful in treating a variety of neurologic and psychiatric disorders.

40 The present invention thus further relates to the use of the compounds of the formula (I) for the manufacture of a medicament for treating a neurologic or psychiatric disorder, and to corresponding methods of treating said disorders.

According to a particular embodiment, the disorder is associated with glycinergic or glutamatergic neurotransmission dysfunction.

- 5 According to a further particular embodiment, the disorder is one or more of the following conditions or diseases: schizophrenia or a psychotic disorder including schizophrenia (paranoid, disorganized, catatonic or undifferentiated), schizophreniform disorder, schizoaffective disorder, delusional disorder, brief psychotic disorder, shared psychotic disorder, psychotic disorder due to a general medical condition and substance-induced psychotic disorder, including both the positive
10 and the negative symptoms of schizophrenia and other psychoses; cognitive disorders including dementia (associated with Alzheimer's disease, ischemia, multi-infarct dementia, trauma, vascular problems or stroke, HIV disease, Parkinson's disease, Huntington's disease, Pick's disease, Creutzfeldt-Jacob disease, perinatal hypoxia, other general medical conditions or substance abuse); delirium, amnesic disorders or cognitive impairment including age related cognitive decline; anxiety
15 disorders including acute stress disorder, agoraphobia, generalized anxiety disorder, obsessive-compulsive disorder, panic attack, panic disorder, post-traumatic stress disorder, separation anxiety disorder, social phobia, specific phobia, substance-induced anxiety disorder and anxiety due to a general medical condition; substance-related disorders and addictive behaviors (including substance-induced delirium, persisting dementia, persisting amnesic disorder, psychotic disorder or
20 anxiety disorder; tolerance, dependence or withdrawal from substances including alcohol, amphetamines, cannabis, cocaine, hallucinogens, inhalants, nicotine, opioids, phencyclidine, sedatives, hypnotics or anxiolytics); obesity, bulimia nervosa and compulsive eating disorders; bipolar disorders, mood disorders including depressive disorders; depression including unipolar depression, seasonal depression and post-partum depression, premenstrual syndrome (PMS) and premenstrual
25 dysphoric disorder (PDD), mood disorders due to a general medical condition, and substance-induced mood disorders; learning disorders, pervasive developmental disorder including autistic disorder, attention deficit disorders including attention-deficit hyperactivity disorder (ADHD) and conduct disorder; movement disorders, including akinesias and akinetic-rigid syndromes (including Parkinson's disease, drug-induced parkinsonism, postencephalitic parkinsonism, progressive supranuclear palsy, multiple system atrophy, corticobasal degeneration, parkinsonism-ALS dementia
30 complex and basal ganglia calcification), medication-induced parkinsonism (such as neuroleptic-induced parkinsonism, neuroleptic malignant syndrome, neuroleptic-induced acute dystonia, neuroleptic-induced acute akathisia, neuroleptic-induced tardive dyskinesia and medication-induced postural tremor), Gilles de la Tourette's syndrome, epilepsy, muscular spasms and disorders associated with muscular spasticity or weakness including tremors; dyskinesias [including tremor (such
35 as rest tremor, postural tremor and intention tremor), chorea (such as Sydenham's chorea, Huntington's disease, benign hereditary chorea, neuroacanthocytosis, symptomatic chorea, drug-induced chorea and hemiballism), myoclonus (including generalised myoclonus and focal myoclonus), tics (including simple tics, complex tics and symptomatic tics), and dystonia (including generalised
40 dystonia such as idiopathic dystonia, drug-induced dystonia, symptomatic dystonia and paroxysmal dystonia, and focal dystonia such as blepharospasm, oromandibular dystonia, spasmodic dysphonia, spasmodic torticollis, axial dystonia, dystonic writer's cramp and hemiplegic dystonia)]; uri-

nary incontinence; neuronal damage including ocular damage, retinopathy or macular degeneration of the eye, tinnitus, hearing impairment and loss, and brain edema; emesis; and sleep disorders including insomnia and narcolepsy.

- 5 According to a further particular embodiment, the disorder is pain, in particular chronic pain and especially neuropathic pain.

Pain can be classified as acute and chronic pain. Acute pain and chronic pain differ in their etiology, pathophysiology, diagnosis and treatment.

10

Acute pain, which occurs following tissue injury, is self-limiting, serves as an alert to ongoing tissue damage and following tissue repair it will usually subside. There are minimal psychological symptoms associated with acute pain apart from mild anxiety. Acute pain is nociceptive in nature and occurs following chemical, mechanical and thermal stimulation of A-delta and C-polymodal pain receptors.

15

- Chronic pain, on the other hand, serves no protective biological function. Rather than being the symptom of tissue damage it is a disease in its own right. Chronic pain is unrelenting and not self-limiting and can persist for years, perhaps decades after the initial injury. Chronic pain can be refractory to multiple treatment regimes. Psychological symptoms associated with chronic pain include chronic anxiety, fear, depression, sleeplessness and impairment of social interaction. Chronic non-malignant pain is predominantly neuropathic in nature and involves damage to either the peripheral or central nervous systems.

20

- 25 Acute pain and chronic pain are caused by different neuro-physiological processes and therefore tend to respond to different types of treatments. Acute pain can be somatic or visceral in nature. Somatic pain tends to be a well localised, constant pain and is described as sharp, aching, throbbing or gnawing. Visceral pain, on the other hand, tends to be vague in distribution, paroxysmal in nature and is usually described as deep, aching, squeezing or colicky in nature. Examples of acute pain include post-operative pain, pain associated with trauma and the pain of arthritis. Acute pain usually responds to treatment with opioids or non-steroidal anti-inflammatory drugs.

30

- Chronic pain, in contrast to acute pain, is described as burning, electric, tingling and shooting in nature. It can be continuous or paroxysmal in presentation. The hallmarks of chronic pain are chronic allodynia and hyperalgesia. Allodynia is pain resulting from a stimulus that normally does not elicit a painful response, such as a light touch. Hyperalgesia is an increased sensitivity to normally painful stimuli. Primary hyperalgesia occurs immediately within the area of the injury. Secondary hyperalgesia occurs in the undamaged area surrounding the injury. Examples of chronic pain include complex regional pain syndrome, pain arising from peripheral neuropathies, post-operative pain, chronic fatigue syndrome pain, tension-type headache, pain arising from mechanical nerve injury and severe pain associated with diseases such as cancer, metabolic disease, neuro-

35

40

tropic viral disease, neurotoxicity, inflammation, multiple sclerosis or any pain arising as a consequence of or associated with stress or depressive illness.

5 Although opioids are cheap and effective, serious and potentially life-threatening side effects occur with their use, most notably respiratory depression and muscle rigidity. In addition the doses of opioids which can be administered are limited by nausea, emesis, constipation, pruritis and urinary retention, often resulting in patients electing to receive sub-optimal pain control rather than suffer these distressing side-effects. Furthermore, these side-effects often result in patients requiring extended hospitalisation. Opioids are highly addictive and are scheduled drugs in many territories.

10 The compounds of formula (I) are particularly useful in the treatment of schizophrenia, bipolar disorder, depression including unipolar depression, seasonal depression and post-partum depression, premenstrual syndrome (PMS) and premenstrual dysphoric disorder (PDD), learning disorders, pervasive developmental disorder including autistic disorder, attention deficit disorders including Attention-Deficit/Hyperactivity Disorder, tic disorders including Tourette's disorder, anxiety disorders including phobia and post traumatic stress disorder, cognitive disorders associated with dementia, AIDS dementia, Alzheimer's, Parkinson's, Huntington's disease, spasticity, myoclonus, muscle spasm, tinnitus and hearing impairment and loss are of particular importance.

20 Particular cognitive disorders are dementia, delirium, amnesic disorders and cognitive impairment including age-related cognitive decline.

Particular anxiety disorders are generalized anxiety disorder, obsessive-compulsive disorder and panic attack.

25 Particular schizophrenia or psychosis pathologies are paranoid, disorganized, catatonic or undifferentiated schizophrenia and substance-induced psychotic disorder.

30 Particular neurologic disorders that can be treated with the compounds of of the formula (I) include in particular a cognitive disorder such as dementia, cognitive impairment, attention deficit hyperactivity disorder.

35 Particular psychiatric disorders that can be treated with the compounds of of the formula (I) include in particular an anxiety disorder, a mood disorder such as depression or a bipolar disorder, schizophrenia, a psychotic disorder.

40 Within the context of the treatment, the use according to the invention of the compounds of the formula (I) involves a method. In this method, an effective quantity of one or more compounds or the formula (I), as a rule formulated in accordance with pharmaceutical and veterinary practice, is administered to the individual to be treated, preferably a mammal, in particular a human being. Whether such a treatment is indicated, and in which form it is to take place, depends on the individual case and is subject to medical assessment (diagnosis) which takes into consideration signs,

symptoms and/or malfunctions which are present, the risks of developing particular signs, symptoms and/or malfunctions, and other factors.

5 As a rule, the treatment is effected by means of single or repeated daily administration, where appropriate together, or alternating, with other drugs or drug-containing preparations.

10 The invention also relates to the manufacture of pharmaceutical compositions for treating an individual, preferably a mammal, in particular a human being. Thus, the compounds of the formula (I) are customarily administered in the form of pharmaceutical compositions which comprise an inert carrier (e.g. a pharmaceutically acceptable excipient) together with at least one compound according to the invention and, where appropriate, other drugs. These compositions can, for example, be administered orally, rectally, transdermally, subcutaneously, intravenously, intramuscularly or intranasally.

15 Examples of suitable pharmaceutical formulations are solid medicinal forms, such as powders, granules, tablets, in particular film tablets, lozenges, sachets, cachets, sugar-coated tablets, capsules, such as hard gelatin capsules and soft gelatin capsules, suppositories or vaginal medicinal forms, semisolid medicinal forms, such as ointments, creams, hydrogels, pastes or plasters, and also liquid medicinal forms, such as solutions, emulsions, in particular oil-in-water emulsions, suspensions, for example lotions, injection preparations and infusion preparations, and eyedrops and eardrops. Implanted release devices can also be used for administering inhibitors according to the invention. In addition, it is also possible to use liposomes or microspheres.

25 When producing the compositions, the compounds according to the invention are optionally mixed or diluted with one or more carriers (excipients). Carriers (excipients) can be solid, semisolid or liquid materials which serve as vehicles, carriers or medium for the active compound.

30 Suitable carriers (excipients) are listed in the specialist medicinal monographs. In addition, the formulations can comprise pharmaceutically acceptable auxiliary substances, such as wetting agents; emulsifying and suspending agents; preservatives; antioxidants; antiirritants; chelating agents; coating auxiliaries; emulsion stabilizers; film formers; gel formers; odor masking agents; taste corrigents; resin; hydrocolloids; solvents; solubilizers; neutralizing agents; diffusion accelerators; pigments; quaternary ammonium compounds; refatting and overfatting agents; raw materials for ointments, creams or oils; silicone derivatives; spreading auxiliaries; stabilizers; sterilants; suppository bases; tablet auxiliaries, such as binders, fillers, glidants, disintegrants or coatings; propellants; drying agents; opacifiers; thickeners; waxes; plasticizers and white mineral oils. A formulation in this regard is based on specialist knowledge as described, for example, in Fiedler, H.P., Lexikon der Hilfsstoffe für Pharmazie, Kosmetik und angrenzende Gebiete [Encyclopedia of auxiliary substances for pharmacy, cosmetics and related fields], 4th edition, Aulendorf: ECV-Editio-Cantor-Verlag, 1996.

40 The compounds of formula (I) may also be suitable for combination with other therapeutic agents.

Thus, the present invention also provides:

- i) a combination comprising a compound of formula (I) with one or more further therapeutic agents;
- 5 ii) a pharmaceutical composition comprising a combination product as defined in i) above and at least one carrier, diluent or excipient;
- iii) the use of a combination as defined in i) above in the manufacture of a medicament for treating or preventing a disorder, disease or condition as defined herein;
- 10 iv) a combination as defined in i) above for use in treating or preventing a disorder, disease or condition as defined herein;
- v) a kit-of-parts for use in the treatment of a disorder, disease or condition as defined herein, comprising a first dosage form comprising a compound of formula (I) and one or more further dosage forms each comprising one or more further therapeutic agents for simultaneous therapeutic administration,
- 15 vi) a combination as defined in i) above for use in therapy;
- vii) a method of treatment or prevention of a disorder, disease or condition as defined herein comprising administering an effective amount of a combination as defined in i) above;
- viii) a combination as defined in i) above for treating or preventing a disorder, disease or condition as defined herein.

20

The combination therapies of the invention may be administered adjunctively. By adjunctive administration is meant the coterminous or overlapping administration of each of the components in the form of separate pharmaceutical compositions or devices. This regime of therapeutic administration of two or more therapeutic agents is referred to generally by those skilled in the art and
25 herein as adjunctive therapeutic administration; it is also known as add-on therapeutic administration. Any and all treatment regimes in which a patient receives separate but coterminous or overlapping therapeutic administration of the compounds of formula (I) and at least one further therapeutic agent are within the scope of the current invention. In one embodiment of adjunctive therapeutic administration as described herein, a patient is typically stabilised on a therapeutic administration of one or more of the components for a period of time and then receives administration of
30 another component.

The combination therapies of the invention may also be administered simultaneously. By simultaneous administration is meant a treatment regime wherein the individual components are administered together, either in the form of a single pharmaceutical composition or device comprising or
35 containing both components, or as separate compositions or devices, each comprising one of the components, administered simultaneously. Such combinations of the separate individual components for simultaneous combination may be provided in the form of a kit-of-parts.

40 In a further aspect, the invention provides a method of treatment of a psychotic disorder by adjunctive therapeutic administration of compounds of formula (I) to a patient receiving therapeutic administration of at least one antipsychotic agent. In a further aspect, the invention provides the use

of compounds of formula (I) in the manufacture of a medicament for adjunctive therapeutic administration for the treatment of a psychotic disorder in a patient receiving therapeutic administration of at least one antipsychotic agent. The invention further provides compounds of formula (I) for use for adjunctive therapeutic administration for the treatment of a psychotic disorder in a patient receiving therapeutic administration of at least one antipsychotic agent.

In a further aspect, the invention provides a method of treatment of a psychotic disorder by adjunctive therapeutic administration of at least one antipsychotic agent to a patient receiving therapeutic administration of compounds of formula (I). In a further aspect, the invention provides the use of at least one antipsychotic agent in the manufacture of a medicament for adjunctive therapeutic administration for the treatment of a psychotic disorder in a patient receiving therapeutic administration of compounds of formula (I). The invention further provides at least one antipsychotic agent for adjunctive therapeutic administration for the treatment of a psychotic disorder in a patient receiving therapeutic administration of compounds of formula (I).

In a further aspect, the invention provides a method of treatment of a psychotic disorder by simultaneous therapeutic administration of compounds of formula (I) in combination with at least one antipsychotic agent. The invention further provides the use of a combination of compounds of formula (I) and at least one antipsychotic agent in the manufacture of a medicament for simultaneous therapeutic administration in the treatment of a psychotic disorder. The invention further provides a combination of compounds of formula (I) and at least one antipsychotic agent for simultaneous therapeutic administration in the treatment of a psychotic disorder. The invention further provides the use of compounds of formula (I) in the manufacture of a medicament for simultaneous therapeutic administration with at least one antipsychotic agent in the treatment of a psychotic disorder. The invention further provides compounds of formula (I) for use for simultaneous therapeutic administration with at least one antipsychotic agent in the treatment of a psychotic disorder. The invention further provides the use of at least one antipsychotic agent in the manufacture of a medicament for simultaneous therapeutic administration with compounds of formula (I) in the treatment of a psychotic disorder. The invention further provides at least one antipsychotic agent for simultaneous therapeutic administration with compounds of formula (I) in the treatment of a psychotic disorder.

In further aspects, the invention provides a method of treatment of a psychotic disorder by simultaneous therapeutic administration of a pharmaceutical composition comprising compounds of formula (I) and at least one mood stabilising or antimanic agent, a pharmaceutical composition comprising compounds of formula (I) and at least one mood stabilising or antimanic agent, the use of a pharmaceutical composition comprising compounds of formula (I) and at least one mood stabilising or antimanic agent in the manufacture of a medicament for the treatment of a psychotic disorder, and a pharmaceutical composition comprising compounds of formula (I) and at least one mood stabilising or antimanic agent for use in the treatment of a psychotic disorder.

- Antipsychotic agents include both typical and atypical antipsychotic drugs. Examples of antipsychotic drugs that are useful in the present invention include, but are not limited to: butyrophenones, such as haloperidol, pimozide, and droperidol; phenothiazines, such as chlorpromazine, thioridazine, mesoridazine, trifluoperazine, perphenazine, fluphenazine, thiflupromazine, prochlorperazine, and acetophenazine; thioxanthenes, such as thiothixene and chlorprothixene; thienobenzodiazepines; dibenzodiazepines; benzisoxazoles; dibenzothiazepines; imidazolidinones; benzisothiazolyl-piperazines; triazine such as lamotrigine; dibenzoxazepines, such as loxapine; dihydroindolones, such as molindone; aripiprazole; and derivatives thereof that have antipsychotic activity.
- Examples of tradenames and suppliers of selected antipsychotic drugs are as follows: clozapine (available under the tradename CLOZARIL®, from Mylan, Zenith Goldline, UDL, Novartis); olanzapine (available under the tradename ZYPREX®, from Lilly); ziprasidone (available under the tradename GEODON®, from Pfizer); risperidone (available under the tradename RISPERDAL®, from Janssen); quetiapine fumarate (available under the tradename SEROQUEL®, from AstraZeneca); haloperidol (available under the tradename HALDOL®, from Ortho-McNeil); chlorpromazine (available under the tradename THORAZINE®, from SmithKline Beecham (GSK)); fluphenazine (available under the tradename PROLIXIN®, from Apothecon, Copley, Schering, Teva, and American Pharmaceutical Partners, Pasadena); thiothixene (available under the tradename NAVANE®, from Pfizer); trifluoperazine (10-[3-(4-methyl-1-piperazinyl)propyl]-2-(trifluoromethyl)phenothiazine dihydrochloride, available under the tradename STELAZINE®, from Smith Klein Beckman); perphenazine (available under the tradename TRILAFON®; from Schering); thioridazine (available under the tradename MELLARIL®; from Novartis, Roxane, HiTech, Teva, and Alparma); molindone (available under the tradename MOBAN®, from Endo); and loxapine (available under the tradename LOXITANE(D; from Watson). Furthermore, benperidol (Glianimon®), perazine (Taxilan®) or melperone (Eunerpan®) may be used. Other antipsychotic drugs include promazine (available under the tradename SPARINE®), triflupromazine (available under the tradename VESPRIN®), chlorprothixene (available under the tradename TARACTAN®), droperidol (available under the tradename INAPSINE®), acetophenazine (available under the tradename TINDAL®), prochlorperazine (available under the tradename COMPAZINE®), methotrimeprazine (available under the tradename NOZINAN®), pipotiazine (available under the tradename PIPOTRIL®), ziprasidone, and hoperidone.

- In a further aspect, the invention provides a method of treatment of a neurodegenerative disorder such as Alzheimer Disease by adjunctive therapeutic administration of compounds of formula (I) to a patient receiving therapeutic administration of at least one agent suitable for the treatment of a neurodegenerative disorder such as Alzheimer Disease. In a further aspect, the invention provides the use of compounds of formula (I) in the manufacture of a medicament for adjunctive therapeutic administration for the treatment of a neurodegenerative disorder such as Alzheimer Disease in a patient receiving therapeutic administration of at least one agent suitable for the treatment of a neurodegenerative disorder such as Alzheimer Disease. The invention further provides compounds of formula (I) for use for adjunctive therapeutic administration for the treatment of a neurodegenerative disorder.

tive disorder such as Alzheimer Disease in a patient receiving therapeutic administration of at least one agent suitable for the treatment of a neurodegenerative disorder such as Alzheimer Disease.

5 In a further aspect, the invention provides a method of treatment of a neurodegenerative disorder such as Alzheimer Disease by adjunctive therapeutic administration of at least one agent suitable for the treatment of a neurodegenerative disorder such as Alzheimer Disease to a patient receiving therapeutic administration of compounds of formula (I). In a further aspect, the invention provides the use of at least one agent suitable for the treatment of a neurodegenerative disorder such as Alzheimer Disease in the manufacture of a medicament for adjunctive therapeutic administration for
10 the treatment of a neurodegenerative disorder such as Alzheimer Disease in a patient receiving therapeutic administration of compounds of formula (I). The invention further provides at least one agent suitable for the treatment of a neurodegenerative disorder such as Alzheimer Disease for adjunctive therapeutic administration for the treatment of a neurodegenerative disorder such as Alzheimer Disease in a patient receiving therapeutic administration of compounds of formula (I).

15 In a further aspect, the invention provides a method of treatment of a neurodegenerative disorder such as Alzheimer Disease by simultaneous therapeutic administration of compounds of formula (I) in combination with at least one agent suitable for the treatment of a neurodegenerative disorder such as Alzheimer Disease. The invention further provides the use of a combination of compounds
20 of formula (I) and at least one agent suitable for the treatment of a neurodegenerative disorder such as Alzheimer Disease in the manufacture of a medicament for simultaneous therapeutic administration in the treatment of a neurodegenerative disorder such as Alzheimer Disease. The invention further provides a combination of compounds of formula (I) and at least one agent suitable for the treatment of a neurodegenerative disorder such as Alzheimer Disease for simultaneous therapeutic
25 administration in the treatment of a neurodegenerative disorder such as Alzheimer Disease. The invention further provides the use of compounds of formula (I) in the manufacture of a medicament for simultaneous therapeutic administration with at least one agent suitable for the treatment of a neurodegenerative disorder such as Alzheimer Disease in the treatment of a neurodegenerative disorder such as Alzheimer Disease. The invention further provides compounds of formula (I) for
30 use for simultaneous therapeutic administration with at least one agent suitable for the treatment of a neurodegenerative disorder such as Alzheimer Disease in the treatment of a neurodegenerative disorder such as Alzheimer Disease. The invention further provides the use of at least one agent suitable for the treatment of a neurodegenerative disorder such as Alzheimer Disease in the manufacture of a medicament for simultaneous therapeutic administration with compounds of formula
35 (I) in the treatment of a neurodegenerative disorder such as Alzheimer Disease. The invention further provides at least one agent suitable for the treatment of a neurodegenerative disorder such as Alzheimer Disease for simultaneous therapeutic administration with compounds of formula (I) in the treatment of a neurodegenerative disorder such as Alzheimer Disease.

40 Examples of agents suitable for the treatment of a neurodegenerative disorder such as Alzheimer Disease that are useful in the present invention include, but are not limited to: cholinesterase inhibitors, agents targeting nicotinic or muscarinic acetylcholine receptors, NMDA receptors, amyloid

formation, mitochondrial dysfunctions, disease associated calpain activity, neuroinflammation, tumor necrosis factor receptors, NF-kappaB, peroxisome proliferator activator receptor gamma, Apolipoprotein E variant 4 (ApoE4), disease-associated increase of the HPA axis, epileptic discharges, vascular dysfunction, vascular risk factors, and oxidative stress.

5

Suitable cholinesterase inhibitors which may be used in combination with the compounds of the inventions include for example tacrine, donepezil, galantamine and rivastigmine.

10

Suitable NMDA receptors targeting agents which may be used in combination with the compounds of the inventions include for example memantine.

Suitable agents affecting increased HPA axis activity which may be used in combination with the compounds of the inventions include for example CRF1 antagonists or V1b antagonists.

15

In a further aspect therefore, the invention provides a method of treatment of pain by adjunctive therapeutic administration of compounds of formula (I) to a patient receiving therapeutic administration of at least one agent suitable for the treatment of pain. In a further aspect, the invention provides the use of compounds of formula (I) in the manufacture of a medicament for adjunctive therapeutic administration for the treatment of pain in a patient receiving therapeutic administration of at least one agent suitable for the treatment of pain. The invention further provides compounds of formula (I) for use for adjunctive therapeutic administration for the treatment of pain in a patient receiving therapeutic administration of at least one agent suitable for the treatment of pain.

20

25

In a further aspect, the invention provides a method of treatment of pain by adjunctive therapeutic administration of at least one agent suitable for the treatment of pain to a patient receiving therapeutic administration of compounds of formula (I). In a further aspect, the invention provides the use of at least one agent suitable for the treatment of pain in the manufacture of a medicament for adjunctive therapeutic administration for the treatment of pain in a patient receiving therapeutic administration of compounds of formula (I). The invention further provides at least one agent suitable for the treatment of pain for adjunctive therapeutic administration for the treatment of pain in a patient receiving therapeutic administration of compounds of formula (I).

30

35

In a further aspect, the invention provides a method of treatment of pain by simultaneous therapeutic administration of compounds of formula (I) in combination with at least one agent suitable for the treatment of pain. The invention further provides the use of a combination of compounds of formula (I) and at least one agent suitable for the treatment of pain in the manufacture of a medicament for simultaneous therapeutic administration in the treatment of pain. The invention further provides a combination of compounds of formula (I) and at least one agent suitable for the treatment of pain for simultaneous therapeutic administration in the treatment of pain. The invention further provides the use of compounds of formula (I) in the manufacture of a medicament for simultaneous therapeutic administration with at least one agent suitable for the treatment of pain in the treatment of pain. The invention further provides compounds of formula (I) for use for simultane-

40

ous therapeutic administration with at least one agent suitable for the treatment of pain in the treatment of pain. The invention further provides the use of at least one agent suitable for the treatment of pain in the manufacture of a medicament for simultaneous therapeutic administration with compounds of formula (I) in the treatment of pain. The invention further provides at least one agent
5 suitable for the treatment of pain for simultaneous therapeutic administration with compounds of formula (I) in the treatment of pain.

Examples of agents suitable for the treatment of pain that are useful in the present invention include, but are not limited to: NSAIDs (Nonsteroidal Antiinflammatory Drugs), anticonvulsant
10 drugs such as carbamazepine and gabapentin, sodium channel blockers, antidepressant drugs, cannabinoids and local anaesthetics.

Suitable agents used in combination with the compounds of the inventions include for example celecoxib, etoricoxib, lumiracoxib, paracetamol, tramadol, methadone, venlafaxine, imipramine,
15 duloxetine, bupropion, gabapentin, pregabalin, lamotrigine, fentanyl, parecoxib, nefopam, remifentanyl, pethidine, diclofenac, rofecoxib, nalbuphine, sufentanil, pethidine, diamorphine and butorphanol.

It will be appreciated by those skilled in the art that the compounds according to the invention may
20 advantageously be used in conjunction with one or more other therapeutic agents, for instance, antidepressant agents such as 5HT₃ antagonists, serotonin agonists, NK-1 antagonists, selective serotonin reuptake inhibitors (SSRI), noradrenaline re-uptake inhibitors (SNRI), tricyclic antidepressants, dopaminergic antidepressants, H₃ antagonists, 5HT_{1A} antagonists, 5HT_{1B} antagonists, 5HT_{1D} antagonists, D₁ agonists, M₁ agonists and/or anticonvulsant agents, as well as cognitive
25 enhancers.

Suitable 5HT₃ antagonists which may be used in combination of the compounds of the inventions include for example ondansetron, granisetron, metoclopramide.

30 Suitable serotonin agonists which may be used in combination with the compounds of the invention include sumatriptan, rauwolscine, yohimbine, metoclopramide.

Suitable SSRIs which may be used in combination with the compounds of the invention include fluoxetine, citalopram, femoxetine, fluvoxamine, paroxetine, indalpine, sertraline, zimeldine.
35

Suitable SNRIs which may be used in combination with the compounds of the invention include venlafaxine and reboxetine.

Suitable tricyclic antidepressants which may be used in combination with a compound of the invention include imipramine, amitriptyline, chlomipramine and nortriptyline.
40

Suitable dopaminergic antidepressants which may be used in combination with a compound of the invention include bupropion and amineptine.


5 Suitable anticonvulsant agents which may be used in combination of the compounds of the invention include for example divalproex, carbamazepine and diazepam.

The following examples serve to explain the invention without limiting it.


10 The compounds were characterized by mass spectrometry, generally recorded via HPLC-MS in a fast gradient on C18-material (electrospray-ionisation (ESI) mode).

Preparation Examples

15 Example 1: N-[1-[3-(azetidin-1-yl)-4-benzyl-chroman-6-yl]azetidin-3-yl]-1-methyl-imidazole-4-sulfonamide



1.1 6-Methoxychroman-4-one oxime



20


5.2 g (29.2 mmol) of 6-methoxychroman-4-one were dissolved in ethanol and 2.53 g (36.5 mmol) hydroxylamine hydrochloride and 2.99 g (36.5 mmol) sodium acetate dissolved in 10 ml of water were added. The mixture was stirred at 65 °C for 1.5 hours. The mixture was allowed to cool to room temperature and concentrated. The residue was dissolved in methyl-tert-butylether. The organic phase was washed with water, dried over MgSO₄ and concentrated to give 5.6 g (29.4 mmol, quant.) of crude product, which was directly used in the next step.

25

ESI-MS [M+H]⁺ = 194

Calculated for C₁₀H₁₁NO₃ = 193.


1.2 6-Methoxychroman-4-one O-tosyl oxime



5.68 g (29.4 mmol) of 6-methoxychroman-4-one oxime were dissolved under argon atmosphere in 30 ml of dry pyridine. At 0°C 6.05 g (31.8 mmol) of 4-methylbenzene-1-sulfonyl chloride were added in small portions over 40 min. The mixture was stirred at 0 °C for an additional hour and then warmed to room temperature and stirred over night. The mixture was poured into 260 ml ice water, stirred, and the suspension was filtered. The solid residue was washed with a small amount of cold water (2x) and cold ethanol (1x), and dried to yield 8.96 g (25.8 mmol, 88 %) of desired product.

ESI-MS $[M+H^+] = 348$ Calculated for $C_{17}H_{17}NO_5S = 347$.


1.3 3-Amino-6-methoxychroman-4-one hydrochloride



To a solution of sodium ethoxide (10.5 ml, 28.1 mmol, 21 % in ethanol) under nitrogen atmosphere at 0°C was added a suspension of 8.96 g (25.8 mmol) of (Z)-6-methoxychroman-4-one O-tosyl oxime in toluene. The mixture was stirred over night and slowly warmed to room temperature. The suspension was filtered and rinsed with ether. 95 ml (190 mmol) of an aqueous solution of hydrogen chloride (2 N) was added to the filtrate and stirred at room temperature for 2 h. The suspension was diluted with 150 ml of water and phases were separated. The organic phase was extracted with aqueous hydrogen chloride solution (2x, 20-30 ml, 1 N) and water (1x, 30 ml). The combined aqueous layers were washed with ether (1x). The aqueous phase was stirred with a small amount of activated charcoal, filtered, and concentrated to a 1/5 of its volume until a crystalline precipitation was observed. The mixture was cooled to 0° C and the crystalline material was filtered off, washed with a small amount of cold ethanol, and dried in vacuo. The filtrate was also concentrated in vacuo. 3.67 g (15.98 mmol, 62 %) of combined crude desired product was obtained.

ESI-MS $[M+H^+] = 194$ Calculated for $C_{10}H_{11}NO_3 = 193$.

1.4 Ethyl 6-methoxy-4-oxochroman-3-ylcarbamate




2.82 g (12.3 mmol) of 6-methoxy-4-oxochroman-3-aminium chloride were dissolved in tetrahydrofuran under nitrogen atmosphere and cooled to 0 °C with an ice bath. Diisopropylethylamine and ethyl carbononochloridate were added. The mixture was allowed to warm to room temperature and stirred for 30 min. The mixture was diluted with ethyl acetate and washed with saturated ammonium chloride solution (2x) and water (1x). The organic phase was washed, dried over MgSO₄, and concentrated in vacuo to give 3.5 g (13.2 mmol, quant.) of crude material.

ESI-MS [M+H⁺] = 265

Calculated for C₁₃H₁₅NO₅ = 266.

10 1.5 Ethyl 4-benzyl-4-hydroxy-6-methoxychroman-3-ylcarbamate




26.4 ml (52.8 mmol) of benzylmagnesium chloride under nitrogen atmosphere were cooled to 0 °C with an ice bath and 3.5 g (13.2 mmol) ethyl 6-methoxy-4-oxochroman-3-ylcarbamate dissolved in 100 ml dry THF were slowly added. The mixture was stirred at 0 °C for 1 h. The cooling bath was removed and saturated ammonium chloride solution was added. Water was added until a clear solution was obtained. The phases were separated and the organic phase was washed with saturated ammonium chloride solution, dried over MgSO₄, and concentrated in vacuo to give 6.87 g (9.1 mmol, quant.) of crude material.

ESI-MS [M+Na⁺] = 380

Calculated for C₂₀H₂₃NO₅ = 357.

20

1.6 Ethyl 4-benzylidene-6-methoxychroman-3-ylcarbamate



6.87 g (12.5 mmol) of ethyl 4-benzyl-4-hydroxy-6-methoxychroman-3-ylcarbamate were added to 80 ml of half concentrated aqueous hydrochloric acid and stirred at 100 °C for 2.5 h. The mixture was cooled to 0 °C and diluted with water. Sodium hydroxide (50 % aqueous solution) was carefully added until pH >10. The aqueous phase was extracted with EtOAc (2x). The combined organic


25

phases were washed with water and brine, dried over MgSO_4 and the solvent was evaporated to give 5.7 g of crude material. The crude material was purified by flash chromatography to yield 3.1 g (9.1 mmol, 73 %) of the desired product.

ESI-MS $[\text{M}+\text{H}^+] = 339$ Calculated for $\text{C}_{20}\text{H}_{21}\text{NO}_4 = 340$.

5

1.7 Ethyl 4-benzyl-6-methoxychroman-3-ylcarbamate




3.1 g (9.1 mmol) of ethyl 4-benzylidene-6-methoxychroman-3-ylcarbamate were dissolved in 80 ml of EtOH and 910 mg (0.9 mmol) of Pd/C were added. Then, 5.8 g (91 mmol) of ammonium formate dissolved in 20 ml of water were added and the mixture was warmed to 70 °C and stirred for 1.5 h. The mixture was cooled to room temperature. The catalyst was filtered off and washed with EtOH/water. The filtrate was concentrated in vacuo to remove EtOH. The aqueous concentrate was extracted with ethyl acetate (2x). The combined organic phases were dried over MgSO_4 and the solvent was evaporated to yield 3.2 g (9.3 mmol, quant.) of the crude product (cis:trans ~ 7:1). The cis-isomer can be enriched (~26:1) by crystallization from hot heptane.

15

ESI-MS $[\text{M}+\text{H}^+] = 342$ Calculated for $\text{C}_{20}\text{H}_{23}\text{NO}_4 = 341$.

1.8 Ethyl 4-benzyl-6-hydroxychroman-3-ylcarbamate




3.19 g (9.3 mmol) of ethyl 4-benzyl-6-methoxychroman-3-ylcarbamate under nitrogen atmosphere were dissolved in 90 ml of methylene dichloride. At 0 °C 28.0 ml (28.0 mmol, 1 M in methylene dichloride) of boron tribromide were added. The reaction mixture was stirred at 0 °C for 2 hours. At 0 °C saturated sodium hydrogencarbonate solution was added to the reaction mixture. The phases were separated and the aqueous phase was extracted with methylene dichloride. The combined organic layers were washed with brine, dried over MgSO_4 and the solvent was evaporated to yield 3.0 g (9.2 mmol, 99%) of the crude product.

25

ESI-MS $[\text{M}+\text{H}^+] = 328$ Calculated for $\text{C}_{19}\text{H}_{21}\text{NO}_4 = 327$.

1.9 Cis-3-Amino-4-benzylchroman-6-ol and trans-3-Amino-4-benzylchroman-6-ol



2.3 g (7.0 mmol) of ethyl 4-benzyl-6-hydroxychroman-3-ylcarbamate were dissolved in ethanolic KOH 20% and stirred at 70°C over night. The solvent was evaporated, the residue partitioned between ethylacetate and water. The organic layer was washed twice with water and the combined water layer extracted another 2 times with ethyl acetate. The combined ethyl acetate extract was dried over MgSO₄, filtrated and evaporated. The crude material was purified by flash chromatography to yield 1.1 g (4.34 mmol, 62 %) of cis-Isomer and 0.25 g (0.97 mmol, 14 %) of the trans-Isomer.


Cis-Isomer: ESI-MS [M+H⁺] = 256

Calculated for C₁₆H₁₇NO₂ = 255.

10 Trans-Isomer: ESI-MS [M+H⁺] = 256

Calculated for C₁₆H₁₇NO₂ = 255.

1.10 Cis-3-(Azetidin-1-yl)-4-benzylchroman-6-ol




0.9 g (3.57 mmol) of cis-3-amino-4-benzylchroman-6-ol, 0.36 mL (3.55 mmol) 1,3-dibromopropane and 1.9 mL (10.88 mmol) N-ethyl-N-isopropylpropan-2-amine were combined with 18 mL acetonitrile and stirred at 130°C in the microwave (CEM) for 3 hours. Additional 75 µL 1,3-dibromopropane and 0.5 mL N-ethyl-N-isopropylpropan-2-amine were added to the reaction mixture (brown solution) and stirred at 130°C in the microwave (CEM) for an additional 1 hour. The reaction mixture was evaporated and the obtained residue partitioned between water and ethyl acetate. The organic phase was washed with water and brine and the combined aqueous phases extracted twice with ethyl acetate. Combined organic extracts were dried over MgSO₄, filtrated and evaporated to dryness to yield 1 g of crude material. The material was purified by flash chromatography to yield 0.6 g (2.01 mmol, 56 %) of desired product.

ESI-MS [M+H⁺] = 296

Calculated for C₁₉H₂₁NO₂ = 295.

1.11 Cis-3-(Azetidin-1-yl)-4-benzylchroman-6-yl trifluoromethanesulfonate




0.6 g (2.00 mmol) of cis-3-(Azetidin-1-yl)-4-benzylchroman-6-ol were dissolved in methylene chloride under nitrogen, 0.5 mL (6.18 mmol) pyridine were added and cooled with an ice bath to 0 °C. 2.2 mL (2.20 mmol) trifluoromethanesulfonic anhydride were added and the reaction mixture stirred under cooling for 1 hour. The reaction mixture was quenched with aqueous bicarbonate solution, and the aqueous phase separated and extracted with methylene chloride once. The combined organic layers were washed with water (2x) and brine(1x), dried over MgSO₄, filtrated and evaporated to dryness to yield 0.8 g of crude material.

ESI-MS [M+H⁺] = 427

Calculated for C₂₀H₂₀F₃NO₄S = 428.

10

1.12 Cis-tert-Butyl (1-(3-(azetidin-1-yl)-4-benzylchroman-6-yl)azetidin-3-yl)carbamate




0.2 g (0.36 mmol) of cis-3-(azetidin-1-yl)-4-benzylchroman-6-yl trifluoromethanesulfonate were dissolved in toluene under nitrogen, 0.01 g (0.05 mmol) Pd(II) acetate, 0.05 g (0.11 mmol) dicyclohexyl(2',4',6'-triisopropyl-[1,1'-biphenyl]-2-yl)phosphine (x-Phos) and 0.35 g (1.07 mmol) cesium carbonate were added to this solution and the resulting mixture stirred at 115 °C for 15 min. Then, 0.09 g (0.43 mmol) of 3-((tert-butoxycarbonyl)amino)azetidin-1-ium chloride was added and the reaction mixture stirred for 30 min. at 115 °C. The reaction mixture was allowed to cool at room temperature, the solvent evaporated, and the residue extracted between water and ethyl acetate. The organic phase was washed with brine, dried over MgSO₄, filtered and evaporated. The crude material was purified by flash chromatography to yield 0.15 g (0.33 mmol, 93 %).

ESI-MS [M+H⁺] = 450

Calculated for C₂₇H₃₅N₃O₃ = 449.

1.13 Cis-1-(3-(azetidin-1-yl)-4-benzylchroman-6-yl)azetidin-3-amine




0.15 g (0.33 mmol) of cis-tert-butyl (1-(3-(azetidin-1-yl)-4-benzylchroman-6-yl)azetidin-3-yl)carbamate were dissolved in methylene chloride, 0.25 mL (3.24 mmol) trifluoroacetic acid were added and the reaction mixture stirred at room temperature overnight. The solvents were evaporated. The residue was dissolved in water and washed twice with methyl-tert-butyl ether. The water layer was separated, aqueous sodium bicarbonate solution was added until pH 8 was reached, and extracted with methylene chloride (3x). The combined organic extracts were dried over MgSO_4 and concentrated to yield 0.10 g (0.29 mmol) of crude material.

ESI-MS $[\text{M}+\text{H}^+] = 350$

Calculated for $\text{C}_{22}\text{H}_{27}\text{N}_3\text{O} = 349$.

1.14 Cis-N-(1-(3-(azetidin-1-yl)-4-benzylchroman-6-yl)azetidin-3-yl)-1-methyl-1H-imidazole-4-sulfonamide




0.05 g (0.14 mmol) of 1-(3-(azetidin-1-yl)-4-benzylchroman-6-yl)azetidin-3-amine were dissolved in methylene chloride, 0.05 g (0.37 mmol) N,N-dimethylpyridin-4-amine and 0.03 g (0.19 mmol) 1-methyl-1H-imidazole-4-sulfonyl chloride were added. The reaction mixture was stirred at room temperature for 30 min. The solvent was evaporated and the residue partitioned between ethyl acetate and water. The organic layer was washed twice with water, dried over MgSO_4 , filtrated and evaporated. The crude material was purified by flash chromatography to yield 0.06 g (0.11 mmol, 81 %) of desired product.

ESI-MS $[\text{M}+\text{H}^+] = 494$

Calculated for $\text{C}_{26}\text{H}_{31}\text{N}_5\text{O}_3\text{S} = 494$.

Example 2: Cis-N-(1-(3-(azetidin-1-yl)-4-benzylchroman-6-yl)azetidin-3-yl)ethanesulfonamide




Cis-N-(1-(3-(azetidin-1-yl)-4-benzylchroman-6-yl)azetidin-3-yl)ethanesulfonamide was prepared in analogy to example 1.

ESI-MS $[M+H]^+$ = 442

Calculated for $C_{24}H_{31}N_3O_3S$ = 441.

5

Example 3: Cis-N-(1-(3-(azetidin-1-yl)-4-benzylchroman-6-yl)azetidin-3-yl)propane-1-sulfonamide




10 Cis-N-(1-(3-(azetidin-1-yl)-4-benzylchroman-6-yl)azetidin-3-yl)propane-1-sulfonamide was prepared in analogy to example 1.

ESI-MS $[M+H]^+$ = 456

Calculated for $C_{25}H_{33}N_3O_3S$ = 455.

Example 4: Cis-N-(1-(3-(azetidin-1-yl)-4-benzylchroman-6-yl)azetidin-3-yl)-1-

15 cyclopropylmethanesulfonamide




Cis-N-(1-(3-(azetidin-1-yl)-4-benzylchroman-6-yl)azetidin-3-yl)-1-cyclopropylmethane-sulfonamide was prepared in analogy to example 1.

ESI-MS $[M+H]^+$ = 468

Calculated for $C_{26}H_{33}N_3O_3S$ = 467.

- 5 Example 5: Cis-N-(1-(3-(azetidin-1-yl)-4-benzylchroman-6-yl)azetidin-3-yl)-1-methyl-1H-pyrazole-4-sulfonamide




- 10 Cis-N-(1-(3-(azetidin-1-yl)-4-benzylchroman-6-yl)azetidin-3-yl)-1-methyl-1H-pyrazole-4-sulfonamide was prepared in analogy to example 1.

ESI-MS $[M+H]^+$ = 494

Calculated for $C_{26}H_{31}N_5O_3S$ = 493.

Example 6: Cis-N-(1-(3-(azetidin-1-yl)-4-benzylchroman-6-yl)azetidin-3-yl)-1-methyl-1H-1,2,3-triazole-4-sulfonamide

15




- 20 Cis-N-(1-(3-(azetidin-1-yl)-4-benzylchroman-6-yl)azetidin-3-yl)-1-methyl-1H-1,2,3-triazole-4-sulfonamide was prepared in analogy to example 1.

ESI-MS $[M+H]^+$ = 495

Calculated for $C_{25}H_{30}N_6O_3S$ = 494.

Example 7: Cis-N-(1-(3-(azetidin-1-yl)-4-benzyl-7-fluorochroman-6-yl)azetidin-3-yl)-1-ethyl-1H-1,2,3-triazole-4-sulfonamide




Cis-N-(1-(3-(azetidin-1-yl)-4-benzyl-7-fluorochroman-6-yl)azetidin-3-yl)-1-ethyl-1H-1,2,3-triazole-4-sulfonamide was prepared in analogy to example 1.

ESI-MS $[M+H]^+$ = 527 Calculated for $C_{26}H_{31}FN_6O_3S$ = 526.

5

Example 8: Cis-N-(1-(3-(azetidin-1-yl)-4-benzyl-7-fluorochroman-6-yl)azetidin-3-yl)-2-fluoropyridine-3-sulfonamide




10 Cis-N-(1-(3-(azetidin-1-yl)-4-benzyl-7-fluorochroman-6-yl)azetidin-3-yl)-2-fluoropyridine-3-sulfonamide was prepared in analogy to example 1.

ESI-MS $[M+H]^+$ = 527 Calculated for $C_{27}H_{28}F_2N_4O_3S$ = 526.

Example 9: Cis-N-(1-(3-(azetidin-1-yl)-4-benzyl-7-fluorochroman-6-yl)azetidin-3-yl)pyridazine-3-sulfonamide


15



Cis-N-(1-(3-(azetidin-1-yl)-4-benzyl-7-fluorochroman-6-yl)azetidin-3-yl)pyridazine-3-sulfonamide was prepared in analogy to example 1.

20 ESI-MS $[M+H]^+$ = 510 Calculated for $C_{26}H_{28}FN_5O_3S$ = 509.

Example 10: Cis-N-(1-(3-(azetidin-1-yl)-4-benzylchroman-6-yl)azetidin-3-yl)-2-fluoropyridine-3-sulfonamide




- 5 Cis-N-(1-(3-(azetidin-1-yl)-4-benzylchroman-6-yl)azetidin-3-yl)-2-fluoropyridine-3-sulfonamide was prepared in analogy to example 1.


ESI-MS $[M+H]^+$ = 509

Calculated for $C_{27}H_{29}FN_4O_3S$ = 508.

- 10 Example 11: Cis-N-(1-(4-benzyl-3-(methylamino)chroman-6-yl)azetidin-3-yl)propane-1-sulfonamide



11.1 Cis-4-benzyl-3-((ethoxycarbonyl)amino)chroman-6-yl trifluoromethanesulfonate




- 15 Cis-4-benzyl-3-((ethoxycarbonyl)amino)chroman-6-yl trifluoromethanesulfonate was prepared starting from cis-ethyl 4-benzyl-6-hydroxychroman-3-ylcarbamate (Example 1, 1.8).
4.9 g (14.97 mmol) of 4-benzyl-3-((ethoxycarbonyl)amino)chroman-6-yl trifluoromethanesulfonate were dissolved in methylene chloride under nitrogen atmosphere, 3.0 mL (37.4 mmol) pyridine were added and cooled to 0 °C with an ice bath. Then, 16.5 mL (16.5 mmol) trifluoromethanesulfonic anhydride (1 M in methylene chloride) were added and the reaction mixture stirred under cooling for 0.5 hour. The reaction mixture was quenched with aqueous ammonium chloride solution, the phases were separated, and the aqueous phase was extracted with methylene chloride once. The combined organic layers were washed with aqueous ammonium chloride solution and
- 20

brine, dried over MgSO_4 , filtrated, and evaporated to dryness to yield 6.6 g (14.37 mmol, 96 %) of crude product.

ESI-MS $[\text{M}+\text{H}^+] = 460$

Calculated for $\text{C}_{20}\text{H}_{20}\text{F}_3\text{NO}_6\text{S} = 459$.

5 11.2 Cis-tert-Butyl (1-(3-(azetidin-1-yl)-4-benzyl-3-((ethoxycarbonyl)amino)chroman-6-yl carbamate




10 6.0 g (13.06 mmol) of cis-4-benzyl-3-((ethoxycarbonyl)amino)chroman-6-yl trifluoromethanesulfonate were dissolved in toluene under nitrogen atmosphere, 0.44 g (1.96 mmol) Pd(II) acetate, 1.87 g (3.92 mmol) dicyclohexyl(2',4',6'-triisopropyl-[1,1'-biphenyl]-2-yl)phosphine, and 10.64 g (32.6 mmol) cesium carbonate were added to this solution and the resulting mixture stirred at 115°C for 15 min. Then, 3-((tert-butoxycarbonyl)amino)azetidin-1-ium chloride was added and the reaction mixture stirred for 1.5 h at 115 °C. The mixture was cooled to room temperature, the solvent was evaporated, and the residue extracted between water and ethyl acetate. The organic phase was washed with water and brine, dried over MgSO_4 , filtered, and evaporated. The crude material was purified by flash chromatography to yield 7.8 g (12.15 mmol, 93 %) of the desired product.

ESI-MS $[\text{M}+\text{H}^+] = 482$

Calculated for $\text{C}_{27}\text{H}_{35}\text{N}_3\text{O}_5 = 481$.


20 11.3 Cis-ethyl (-6-(3-aminoazetidin-1-yl)-4-benzylchroman-3-yl)carbamate



25 7.8 g (12.15 mmol) of cis-tert-butyl (1-(3-(azetidin-1-yl)-4-benzyl-3-((ethoxycarbonyl)amino)chroman-6-yl carbamate were dissolved in methylene chloride, 9.0 mL (117 mmol) trifluoroacetic acid were added, and the reaction mixture stirred at room temperature over night. The solvent was evaporated, the residue dissolved in water and washed with methyl-tert-butylether. The organic layer was washed with water additional 3x. To the combined water layers aqueous sodium bicarbonate solution was added until pH 8 was reached and extracted with methylene chloride (3x). The combined methylene dichloride extracts were dried over MgSO_4 and concentrated to yield 4.33 g (11.35 mmol, 93 %).

ESI-MS $[M+H]^+$ = 382Calculated for $C_{22}H_{27}N_3O_5$ = 381.

11.4 Cis-ethyl (-4-benzyl-6-(3-(propylsulfonamido)azetidin-1-yl)chroman-3-yl)carbamate




5

1.5 g (3.93 mmol) of cis-ethyl (-6-(3-aminoazetidin-1-yl)-4-benzylchroman-3-yl)carbamate were dissolved in methylene dichloride, 1.2 g (9.83 mmol) N,N-dimethylpyridin-4-amine, and 0.60 mL (5.39 mmol) propane-1-sulfonyl chloride were added. The reaction mixture was stirred for 1 h at room temperature. The solvent was evaporated and the residue partitioned between ethyl acetate and water. The organic layer was washed twice with water and once with brine, dried over $MgSO_4$, filtered, and concentrated. The crude material was purified by flash chromatography to yield 1.74 g (3.57 mmol, 91 %) of the desired product.

10

ESI-MS $[M+H]^+$ = 382Calculated for $C_{22}H_{27}N_3O_5S$ = 381.

11.5 Cis-ethyl (-4-benzyl-6-(3-(propylsulfonamido)azetidin-1-yl)chroman-3-yl)carbamate



20


1.74 g (3.57 mmol) of cis-ethyl (-4-benzyl-6-(3-(propylsulfonamido)azetidin-1-yl)chroman-3-yl)carbamate were dissolved in tetrahydrofuran under nitrogen atmosphere, 18 mL (18 mmol) lithium aluminium hydride solution (1 M in tetrahydrofuran) were added, and the reaction mixture stirred at reflux for 1 h. The reaction mixture was cooled to room temperature, the excess of lithium aluminium hydride quenched with methanol, and the solvent was evaporated. The residue was partitioned between ethyl acetate and water. The resulting mixture was filtered through celite.

25

Filtrate: aqueous layer was separated and extracted with ethyl acetate. The combined organic layers were dried over $MgSO_4$, filtered, and concentrated. The crude material was purified by flash chromatography to yield 1.42 g (3.32 mmol, 93 %) of the desired product.

ESI-MS $[M+H]^+$ = 430Calculated for $C_{23}H_{31}N_3O_3S$ = 429.

Example 12: Cis-N-(1-(4-benzyl-3-(methylamino)chroman-6-yl)azetidin-3-yl)ethanesulfonamide




Cis-N-(1-(4-benzyl-3-(methylamino)chroman-6-yl)azetidin-3-yl)ethanesulfonamide was prepared in analogy to example 11.

ESI-MS $[M+H]^+$ = 416

Calculated for $C_{22}H_{29}N_3O_3S$ = 415.

5

Example 13: Cis-N-(1-(4-benzyl-3-(methylamino)chroman-6-yl)azetidin-3-yl)-1-cyclopropylmethanesulfonamide



Cis-N-(1-(4-benzyl-3-(methylamino)chroman-6-yl)azetidin-3-yl)-1-


10 cyclopropylmethanesulfonamide was prepared in analogy to example 11.

ESI-MS $[M+H]^+$ = 442

Calculated for $C_{24}H_{31}N_3O_3S$ = 441.

Example 14: Cis-N-(1-(4-benzyl-7-fluoro-3-(methylamino)chroman-6-yl)azetidin-3-

15 yl)ethanesulfonamide and trans-N-(1-(4-benzyl-7-fluoro-3-(methylamino)chroman-6-yl)azetidin-3-yl)ethanesulfonamide



Cis-N-(1-(4-benzyl-7-fluoro-3-(methylamino)chroman-6-yl)azetidin-3-yl)ethanesulfonamide and trans-N-(1-(4-benzyl-7-fluoro-3-(methylamino)chroman-6-yl)azetidin-3-yl)ethanesulfonamide were prepared in analogy to example 11.


20 14a. Cis-isomer: ESI-MS $[M+H]^+$ = 434

Calculated for $C_{22}H_{28}FN_3O_3S$ = 433.

14b. Trans-isomer: ESI-MS $[M+H]^+$ = 434

Calculated for $C_{22}H_{28}FN_3O_3S$ = 433.

Example 15: Cis-N-(1-(4-benzyl-7-fluoro-3-(methylamino)chroman-6-yl)azetidin-3-yl)-1-methyl-1H-imidazole-4-sulfonamide and trans-N-(1-(4-benzyl-7-fluoro-3-(methylamino)chroman-6-yl)azetidin-3-yl)-1-methyl-1H-imidazole-4-sulfonamide




5

Cis-N-(1-(4-benzyl-7-fluoro-3-(methylamino)chroman-6-yl)azetidin-3-yl)-1-methyl-1H-imidazole-4-sulfonamide and trans-N-(1-(4-benzyl-7-fluoro-3-(methylamino)chroman-6-yl)azetidin-3-yl)-1-methyl-1H-imidazole-4-sulfonamide were prepared in analogy to example 11.

15a. Cis-isomer: ESI-MS $[M+H]^+$ = 486 Calculated for $C_{24}H_{28}FN_5O_3S$ = 485.

10 15b. Trans-isomer: ESI-MS $[M+H]^+$ = 486 Calculated for $C_{24}H_{28}FN_5O_3S$ = 485.

Example 16: Cis-N-(1-(4-benzyl-7-fluoro-3-(methylamino)chroman-6-yl)azetidin-3-yl)propane-1-sulfonamide and trans-N-(1-(4-benzyl-7-fluoro-3-(methylamino)chroman-6-yl)azetidin-3-yl)propane-1-sulfonamide




15

Cis-N-(1-(4-benzyl-7-fluoro-3-(methylamino)chroman-6-yl)azetidin-3-yl)propane-1-sulfonamide and trans-N-(1-(4-benzyl-7-fluoro-3-(methylamino)chroman-6-yl)azetidin-3-yl)propane-1-sulfonamide were prepared in analogy to example 11.

16a. Cis-isomer: ESI-MS $[M+H]^+$ = 448 Calculated for $C_{23}H_{30}FN_3O_3S$ = 447.

20 16b. Trans-isomer: ESI-MS $[M+H]^+$ = 448 Calculated for $C_{23}H_{30}FN_3O_3S$ = 447.


Example 17: Cis-N-(1-(4-benzyl-7-fluoro-3-(methylamino)chroman-6-yl)azetidin-3-yl)-1-methyl-1H-pyrazole-4-sulfonamide and trans-N-(1-(4-benzyl-7-fluoro-3-(methylamino)chroman-6-yl)azetidin-3-yl)-1-methyl-1H-pyrazole-4-sulfonamide



Cis-N-(1-(4-benzyl-7-fluoro-3-(methylamino)chroman-6-yl)azetidin-3-yl)-1-methyl-1H-pyrazole-4-sulfonamide and trans-N-(1-(4-benzyl-7-fluoro-3-(methylamino)chroman-6-yl)azetidin-3-yl)-1-methyl-1H-pyrazole-4-sulfonamide were prepared in analogy to example 11.


- 5 17a. Cis-isomer: ESI-MS $[M+H]^+$ = 486 Calculated for $C_{24}H_{28}FN_5O_3S$ = 485.
 17b. Trans-isomer: ESI-MS $[M+H]^+$ = 486 Calculated for $C_{24}H_{28}FN_5O_3S$ = 485.

Example 18: Cis-N-(1-(4-benzyl-7-fluoro-3-(methylamino)chroman-6-yl)azetidin-3-yl)-1-methyl-1H-1,2,3-triazole-4-sulfonamide and trans-N-(1-(4-benzyl-7-fluoro-3-(methylamino)chroman-6-yl)azetidin-3-yl)-1-methyl-1H-1,2,3-triazole-4-sulfonamide



- 15 Cis-N-(1-(4-benzyl-7-fluoro-3-(methylamino)chroman-6-yl)azetidin-3-yl)-1-methyl-1H-1,2,3-triazole-4-sulfonamide and trans-N-(1-(4-benzyl-7-fluoro-3-(methylamino)chroman-6-yl)azetidin-3-yl)-1-methyl-1H-1,2,3-triazole-4-sulfonamide were prepared in analogy to example 11.
 18a. Cis-isomer: ESI-MS $[M+H]^+$ = 487 Calculated for $C_{23}H_{27}FN_6O_3S$ = 486.
 18b. Trans-isomer: ESI-MS $[M+H]^+$ = 487 Calculated for $C_{23}H_{27}FN_6O_3S$ = 486.

Example 19: Cis-N-(1-(4-benzyl-7-fluoro-3-(methylamino)chroman-6-yl)azetidin-3-yl)-1-cyclopropylmethanesulfonamide and trans-N-(1-(4-benzyl-7-fluoro-3-(methylamino)chroman-6-yl)azetidin-3-yl)-1-cyclopropylmethanesulfonamide



Cis-N-(1-(4-benzyl-7-fluoro-3-(methylamino)chroman-6-yl)azetidin-3-yl)-1-cyclopropylmethanesulfonamide and trans-N-(1-(4-benzyl-7-fluoro-3-(methylamino)chroman-6-yl)azetidin-3-yl)-1-cyclopropylmethanesulfonamide were prepared in analogy to example 11.


19a. Cis-isomer: ESI-MS $[M+H]^+$ = 460

Calculated for $C_{24}H_{30}FN_3O_3S$ = 459.

5 19b. Trans-isomer: ESI-MS $[M+H]^+$ = 460

Calculated for $C_{24}H_{30}FN_3O_3S$ = 459.

Example 20: Cis-N-(1-(3-amino-4-benzylchroman-6-yl)azetidin-3-yl)propane-1-sulfonamide




0.04 g (0.08 mmol) of cis-ethyl (-4-benzyl-6-(3-(propylsulfonamido)azetidin-1-yl)chroman-3-yl)carbamate (Example 11, 11.4) were dissolved in ethanolic potassium hydroxide solution (20%) and stirred at 90°C in the microwave for 15 min. The solvent was evaporated and the residue partitioned between ethyl acetate and water. The organic layer was washed twice with water and the combined water layers were extracted another 2x with ethyl acetate. The combined ethyl acetate extracts were dried over $MgSO_4$, filtrated, and concentrated. The crude material was purified by flash chromatography to yield 0.03 g (0.06mmol, 82 %) of the desired product.

15

ESI-MS $[M+H]^+$ = 416

Calculated for $C_{22}H_{29}N_3O_3S$ = 415.

Example 21: Cis-N-(1-(3-amino-4-benzylchroman-6-yl)azetidin-3-yl)-1-cyclopropylmethanesulfonamide




20

Cis-N-(1-(3-amino-4-benzylchroman-6-yl)azetidin-3-yl)-1-cyclopropylmethanesulfonamide was prepared in analogy to example 20.

ESI-MS $[M+H]^+$ = 428


Calculated for $C_{23}H_{29}N_3O_3S$ = 427.

25 Example 22: Cis-N-(1-(4-benzyl-3-(methylamino)chroman-6-yl)azetidin-3-yl)propane-1-sulfonamide – Isomer 1




- Cis-N-(1-(4-benzyl-3-(methylamino)chroman-6-yl)azetidin-3-yl)propane-1-sulfonamide was prepared by separation of the racemic mixture obtained in example 11 through chiral chromatography on Daicel Chiralpak (n-heptane ethanol 65:35 + 0,1% triethylamine) and isolation of the isomer as the first eluting peak.
- 5 ESI-MS $[M+H]^+$ = 430 Calculated for $C_{23}H_{31}N_3O_3S$ = 429.

Example 23: Cis-N-(1-(4-benzyl-3-(methylamino)chroman-6-yl)azetidin-3-yl)propane-1-sulfonamide – Isomer 2



- 10 Cis-N-(1-(4-benzyl-3-(methylamino)chroman-6-yl)azetidin-3-yl)propane-1-sulfonamide was prepared by separation of the racemic mixture obtained in example 11 through chiral chromatography on Daicel Chiralpak (n-heptane ethanol 65:35 + 0,1% triethylamine) and isolation of the isomer as the second eluting peak.
- 15 ESI-MS $[M+H]^+$ = 430 Calculated for $C_{23}H_{31}N_3O_3S$ = 429.


Example 24: Cis-N-(1-(4-benzyl-3-(methylamino)chroman-6-yl)azetidin-3-yl)-1-cyclopropylmethanesulfonamide – Isomer 1



- 20 Cis-N-(1-(4-benzyl-3-(methylamino)chroman-6-yl)azetidin-3-yl)-1-cyclopropylmethanesulfonamide was prepared by separation of the racemic mixture obtained in example 13 through chiral chromatography on Daicel Chiralpak (n-heptane ethanol 65:35 + 0,1% triethylamine) and isolation of the isomer as the first eluting peak.

ESI-MS $[M+H]^+$ = 442Calculated for $C_{24}H_{31}N_3O_3S$ = 441.

Example 25: Cis-N-(1-(4-benzyl-3-(methylamino)chroman-6-yl)azetidin-3-yl)-1-cyclopropylmethanesulfonamide – Isomer 2




5

Cis-N-(1-(4-benzyl-3-(methylamino)chroman-6-yl)azetidin-3-yl)-1-cyclopropylmethanesulfonamide was prepared by separation of the racemic mixture obtained in example 13 through chiral chromatography on Daicel Chiralpak (n-heptane ethanol 65:35 + 0,1% triethylamine) and isolation of the isomer as the second eluting peak.

10

ESI-MS $[M+H]^+$ = 442Calculated for $C_{24}H_{31}N_3O_3S$ = 441.

Example 26: Cis-1-(4-benzyl-7-fluoro-6-(3-(5-fluoropyridine-3-sulfonamido)azetidin-1-yl)chroman-3-yl)azetidin-1-ium (E)-3-carboxyacrylate




15

Cis-1-(4-benzyl-7-fluoro-6-(3-(5-fluoropyridine-3-sulfonamido)azetidin-1-yl)chroman-3-yl)azetidin-1-ium (E)-3-carboxyacrylate was prepared in analogy to example 1.

ESI-MS $[M+H]^+$ = 527Calculated for $C_{27}H_{28}F_2N_4O_3S$ = 526.

Example 27: Cis-N-(1-(4-benzyl-7-fluoro-3-(methylamino)chroman-6-yl)azetidin-3-yl)propane-1-sulfonamide – Isomer 1


20



Cis-N-(1-(4-benzyl-7-fluoro-3-(methylamino)chroman-6-yl)azetidin-3-yl)propane-1-sulfonamide was prepared by separation of the racemic mixture obtained in example 16 through chiral chromatography on Daicel Chiralpak (n-heptane ethanol 70:30 + 0,1% triethylamine) and isolation of the isomer as the first eluting peak.

5 ESI-MS $[M+H]^+$ = 448 Calculated for $C_{23}H_{30}FN_3O_3S$ = 429.

Example 28: Cis-N-(1-(4-benzyl-7-fluoro-3-(methylamino)chroman-6-yl)azetidin-3-yl)propane-1-sulfonamide – Isomer 2




10 Cis-N-(1-(4-benzyl-7-fluoro-3-(methylamino)chroman-6-yl)azetidin-3-yl)propane-1-sulfonamide was prepared by separation of the racemic mixture obtained in example 11 through chiral chromatography on Daicel Chiralpak (n-heptane ethanol 65:35 + 0,1% triethylamine) and isolation of the isomer as the second eluting peak.

ESI-MS $[M+H]^+$ = 448 Calculated for $C_{23}H_{30}FN_3O_3S$ = 429.

15

Example 29: Cis-N-(1-(4-benzyl-3-(propylamino)chroman-6-yl)azetidin-3-yl)propane-1-sulfonamide




0.17 g (0.41 mmol) of cis-N-(1-(3-amino-4-benzylchroman-6-yl)azetidin-3-yl)propane-1-sulfonamide was reacted with 0.03 mL (0.41 mmol) propionaldehyde in the presence of 0.02 mL (0.41 mol) glacial acetic acid in 20 mL dichloro ethylene. 0.03 g (0.51 mmol) sodiumcyanoborohydride were dissolved in 5 mL methanol, added, and the mixture was stirred at room temperature for 30 min. The reaction mixture was added to aqueous sodium bicarbonate solution and stirred for 15 min., then it was extracted three times with dichloro methylene. The combined organic phases were dried with $MgSO_4$ and evaporated to dryness. The residue was purified by flash chromatography to yield 0.09 g (0.19 mmol, 47 %) of the desired product.

25

ESI-MS $[M+H]^+$ = 458 Calculated for $C_{25}H_{35}N_3O_3S$ = 457.

30

Example 30: Cis-N-(1-(4-benzyl-3-(diethylamino)chroman-6-yl)azetidin-3-yl)propane-1-sulfonamide




Cis-N-(1-(4-benzyl-3-(diethylamino)chroman-6-yl)azetidin-3-yl)propane-1-sulfonamide was prepared in analogy to example 29.

ESI-MS $[M+H]^+$ = 472 Calculated for $C_{26}H_{37}N_3O_3S$ = 471.

5


Example 31: Cis-N-(1-(4-benzyl-3-((cyclopropylmethyl)amino)chroman-6-yl)azetidin-3-yl)propane-1-sulfonamide



10 Cis-N-(1-(4-benzyl-3-((cyclopropylmethyl)amino)chroman-6-yl)azetidin-3-yl)propane-1-sulfonamide was prepared in analogy to example 29.

ESI-MS $[M+H]^+$ = 470 Calculated for $C_{26}H_{35}N_3O_3S$ = 469.

Example 32: Cis-N-(1-(4-benzyl-3-(cyclobutylamino)chroman-6-yl)azetidin-3-yl)propane-1-sulfonamide



15

Cis-N-(1-(4-benzyl-3-(cyclobutylamino)chroman-6-yl)azetidin-3-yl)propane-1-sulfonamide was prepared in analogy to example 29.

ESI-MS $[M+H]^+$ = 470 Calculated for $C_{26}H_{35}N_3O_3S$ = 469.

20

Biological testing

1. $[^3H]$ -Glycine uptake into recombinant CHO cells expressing human GlyT1:

Human GlyT1c expressing recombinant hGlyT1c_5_CHO cells were plated at 20,000 cells per well in 96 well Cytostar-T scintillation microplates (Amersham Biosciences) and cultured to sub-confluency for 24 h. For glycine uptake assays the culture medium was aspirated and the cells were washed once with 100 μ l HBSS (Gibco BRL, #14025-050) with 5 mM L-Alanine (Merck #1007).

- 5 80 μ l HBSS buffer were added, followed by 10 μ l inhibitor or vehicle (10% DMSO) and 10 μ l [3 H]-glycine (TRK71, Amersham Biosciences) to a final concentration of 200 nM for initiation of glycine uptake. The plates were placed in a Wallac Microbeta (PerkinElmer) and continuously counted by solid phase scintillation spectrometry during up to 3 hours. Nonspecific uptake was determined in the presence of 10 μ M Org24598. IC₅₀ calculations were made by four-parametric logistic nonlinear regression analysis (GraphPad Prism) using determinations within the range of linear increase of [3 H]-glycine incorporation between 60 and 120 min.

2. Radioligand binding assays using recombinant CHO cell membranes expressing human GlyT1:

15 Radioligand binding to human GlyT1c transporter-expressing membranes was determined as described in Mezler et al., Molecular Pharmacology 74:1705-1715, 2008.

The following results were obtained with the compounds disclosed in the examples:

20

Table 1:

	radioligand binding
Example	K _{iapp} [μ M]
1	≤ 0.01
2*	≤ 1.0
3	≤ 0.1
4	≤ 0.1
5	≤ 0.1
6	≤ 0.01
7*	≤ 0.1
8*	≤ 1.0
9	≤ 1.0
10	≤ 0.1
11	≤ 0.01
12	≤ 0.1
13	≤ 0.01
14a	≤ 0.1
14b	≤ 0.1
15a	≤ 0.01
15b	≤ 0.01
16a	≤ 0.01

16b	≤ 0.01
17a	≤ 0.01
17b	≤ 0.01
18a	≤ 0.01
18b	≤ 0.01
19a	≤ 0.01
19b	≤ 0.01
20*	≤ 0.1
21*	≤ 0.01
22*	≤ 0.01
23*	≤ 1.0
24	≤ 0.01
25*	≤ 1.0
26*	≤ 0.1
27	≤ 0.01
28	≤ 0.1
29	≤ 0.1
30	≤ 0.1
31	≤ 1.0
32	≤ 1.0

* these compounds were tested in the form of the corresponding fumarate salts

3. Metabolic stability

5 Metabolic stability was determined as follows:

0.5 μ M test substance was preincubated together with human liver microsomes (0.25 mg of microsomal protein/ml) in 0.05 M potassium phosphate buffer of pH 7.4 in microtiter plates at 37°C for 5 min. The reaction was started by adding NADPH (1.0 mM). After 0, 5, 10, 15, 20 and 30 min the reaction was stopped and cooled with twice the amount of quench solution consisting of acetonitrile/methanol 1:1, and containing 0.2 μ M carbutamide. The samples were frozen until analyzed. The remaining concentration of undegraded test substance was determined by LC MSMS. The half-life ($T_{1/2}$) was determined from the gradient of the signal of test substance/unit time plot, allowing to calculate the half-life of the test substance, assuming first order kinetics, from the decrease in the concentration of the compound with time. The microsomal clearance (mClint) was calculated as follows: $mClint = ((\ln(2)/t_{1/2}) / \text{Microsomal Protein Concentration (mg/ml)}) * 1000$, leading to the unit of μ L/min/mg. The scaled clearance (mClin_scaled) was calculated as $mClin_scaled = mClint * (\text{Microsomal Yield (mg/kg BW)}) / 1000000 * 60$, leading to the units L/hr/kg. The Microsomal Yield is defined by the specifics of the used microsomes. Calculations

were modified from references: Di, The Society for Biomolecular Screening, 2003, 453-462; Obach, DMD, 1999 vol 27. N 11, 1350-1359.

The following results were obtained with the compounds disclosed in the examples:

5

Table 2:

Example	human mCl [L/h/Kg]
1	≤ 5
2*	≤ 5
3	≤ 5
4	≤ 5
5	≤ 5
6	≤ 5
7*	ND
8*	≤ 50
9	≤ 50
10	≤ 5
11	≤ 5
12	≤ 5
13	≤ 5
14a	≤ 5
14b	≤ 5
15a	≤ 5
15b	≤ 5
16a	≤ 5
16b	≤ 5
17a	≤ 5
17b	≤ 5
18a	≤ 50
18b	≤ 5
19a	≤ 50
19b	≤ 50
20*	ND
21*	≤ 5
22*	≤ 5
23*	≤ 50
24	≤ 50
25*	≤ 5
26*	≤ 50
27	≤ 50
28	≤ 5

29	≤ 5
30	≤ 50
31	≤ 5
32	≤ 5

* these compounds were tested in the form of the corresponding fumarate salts

4. Determination of efflux ratio using Madin-Darby Canine Kidney Type II cells


5 Bidirectional transport experiments were performed on Madin-Darby Canine Kidney Type II cells over-expressing multidrug resistance protein 1 (MDR1-MDCK) to evaluate the compounds as potential P-gp substrates.

10 Compounds were added at 1 μ M in HBSS-pH 7.4 (hanks balanced salt solution) to either the apical or basolateral side of MDR1-MDCK cell monolayers grown on Millicell 96-Cell polycarbonate filters. Samples were collected from both apical and basolateral sides at time 0 and after 1h incubation at 37C, compounds concentrations were measured by HPLC/MS/MS and permeability coefficients were then determined in both transport directions. The efflux ratio was subsequently calculated from the permeability coefficient.

15

We claim:

1. Compounds of the formula (I)



wherein

A is a 5- or 6-membered ring;

R^1 is hydrogen, C_1 - C_6 -alkyl, C_3 - C_{12} -cycloalkyl- C_1 - C_4 -alkyl, halogenated C_1 - C_6 -alkyl, tri-
(C_1 - C_4 -alkyl)-silyl- C_1 - C_4 -alkyl, hydroxy- C_1 - C_4 -alkyl, C_1 - C_6 -alkoxy- C_1 - C_4 -alkyl, amino-
 C_1 - C_4 -alkyl, C_1 - C_6 -alkylamino- C_1 - C_4 -alkyl, di- C_1 - C_6 -alkylamino- C_1 - C_4 -alkyl, C_1 - C_6 -
alkylcarbonylamino- C_1 - C_4 -alkyl, C_1 - C_6 -alkyloxycarbonylamino- C_1 - C_4 -alkyl, C_1 - C_6 -
alkylaminocarbonylamino- C_1 - C_4 -alkyl, di- C_1 - C_6 -alkylaminocarbonylamino- C_1 - C_4 -alkyl,
 C_1 - C_6 -alkylsulfonylamino- C_1 - C_4 -alkyl, (optionally substituted C_6 - C_{12} -aryl- C_1 - C_6 -
alkyl)amino- C_1 - C_4 -alkyl, optionally substituted C_6 - C_{12} -aryl- C_1 - C_4 -alkyl, optionally sub-
stituted M_3 - M_{12} -heterocyclyl- C_1 - C_4 -alkyl, C_3 - C_{12} -cycloalkyl, C_1 - C_6 -alkylcarbonyl, C_1 -
 C_6 -alkoxycarbonyl, halogenated C_1 - C_6 -alkoxycarbonyl, C_6 - C_{12} -aryloxycarbonyl, ami-
nocarbonyl, C_1 - C_6 -alkylaminocarbonyl, (halogenated C_1 - C_4 -alkyl)aminocarbonyl, C_6 -
 C_{12} -arylaminocarbonyl, C_2 - C_6 -alkenyl, C_2 - C_6 -alkynyl, optionally substituted C_6 - C_{12} -
aryl, hydroxy, C_1 - C_6 -alkoxy, halogenated C_1 - C_6 -alkoxy, C_1 - C_6 -hydroxyalkoxy, C_1 - C_6 -
alkoxy- C_1 - C_4 -alkoxy, amino- C_1 - C_4 -alkoxy, C_1 - C_6 -alkylamino- C_1 - C_4 -alkoxy, di- C_1 - C_6 -
alkylamino- C_1 - C_4 -alkoxy, C_1 - C_6 -alkylcarbonylamino- C_1 - C_4 -alkoxy, C_6 - C_{12} -
arylcarbonylamino- C_1 - C_4 -alkoxy, C_1 - C_6 -alkoxycarbonylamino- C_1 - C_4 -alkoxy, C_6 - C_{12} -
aryl- C_1 - C_4 -alkoxy, C_1 - C_6 -alkylsulfonylamino- C_1 - C_4 -alkoxy, (halogenated C_1 - C_6 -
alkyl)sulfonylamino- C_1 - C_4 -alkoxy, C_6 - C_{12} -arylsulfonylamino- C_1 - C_4 -alkoxy, (C_6 - C_{12} -
aryl- C_1 - C_6 -alkyl)sulfonylamino- C_1 - C_4 -alkoxy, M_3 - M_{12} -heterocyclylsulfonylamino- C_1 -
 C_4 -alkoxy, M_3 - M_{12} -heterocyclyl- C_1 - C_4 -alkoxy, C_6 - C_{12} -aryloxy, M_3 - M_{12} -
heterocyclyloxy, C_1 - C_6 -alkylthio, halogenated C_1 - C_6 -alkylthio, C_1 - C_6 -alkylamino, (hal-
ogenated C_1 - C_6 -alkyl)amino, di- C_1 - C_6 -alkylamino, di-(halogenated C_1 - C_6 -alkyl)amino,
 C_1 - C_6 -alkylcarbonylamino, (halogenated C_1 - C_6 -alkyl)carbonylamino, C_6 - C_{12} -


arylcarbonylamino, C₁-C₆-alkylsulfonylamino, (halogenated C₁-C₆-alkyl)sulfonylamino, C₆-C₁₂-arylsulfonylamino or optionally substituted M₃-M₁₂-heterocyclyl;

- 5 W is -NR⁷- or a bond;
- A¹ is optionally substituted C₁-C₄-alkylene or a bond;
- Q is -S(O)₂- or -C(O)-;
- 10 Y is -NR⁸- or a bond;
- n1 is 0, 1, 2, or 3;
- n2 is 0, 1, 2, or 3;
- 15 X¹ is >N- or >CH-;
- R⁶ is hydrogen, halogen, C₁-C₄-alkyl, halogenated C₁-C₄-alkyl, -CN, OH C₁-C₆-alkoxy or
20 halogenated C₁-C₆-alkoxy, or two radicals R⁶ together with the carbon atom to which
 they are attached form a carbonyl group;
- R² is hydrogen, halogen, C₁-C₆-alkyl, halogenated C₁-C₄-alkyl, -CN, C₂-C₆-alkenyl, C₂-C₆-
alkynyl, optionally substituted C₆-C₁₂-aryl, hydroxy, C₁-C₆-alkoxy, halogenated C₁-C₆-
alkoxy, C₁-C₆-alkoxycarbonyl, C₂-C₆-alkenyloxy, C₆-C₁₂-aryl-C₁-C₄-alkoxy, C₁-C₆-
25 alkylcarbonyloxy, C₁-C₆-alkylthio, C₁-C₆-alkylsulfinyl, C₁-C₆-alkylsulfonyl, aminosul-
fonyl, amino, C₁-C₆-alkylamino, C₂-C₆-alkenylamino, nitro or optionally substituted
M₃-M₁₂-heterocyclyl, or two radicals R² together with the ring atoms of A to which they
are bound form a 5- or 6 membered ring;
- 30 A² is -O-, -S- or -NR⁹-;
- R³ is hydrogen, halogen, C₁-C₆-alkyl or C₁-C₆-alkoxy, or two radicals R³ together with the
carbon atom to which they are attached form a carbonyl group;
- 35 Y¹ is a bond or optionally substituted C₁-C₄-alkylene;
- R^{4a} is hydrogen, C₁-C₆-alkyl, C₃-C₁₂-cycloalkyl-C₁-C₄-alkyl, halogenated C₁-C₄-alkyl, hy-
droxy-C₁-C₄-alkyl, C₁-C₆-alkoxy-C₁-C₄-alkyl, amino-C₁-C₄-alkyl, -CH₂CN, C₆-C₁₂-
aryl-C₁-C₄-alkyl, optionally substituted C₃-C₁₂-cycloalkyl, -CHO, C₁-C₄-alkylcarbonyl,
40 (halogenated C₁-C₄-alkyl)carbonyl, C₆-C₁₂-arylcarbonyl, C₁-C₄-alkoxycarbonyl, C₆-C₁₂-


aryloxycarbonyl, C₁-C₆-alkylaminocarbonyl, C₂-C₆-alkenyl, -C(=NH)NH₂, -C(=NH)NHCN, C₁-C₆-alkylsulfonyl, C₆-C₁₂-arylsulfonyl, amino, -NO or optionally substituted M₃-M₁₂-heterocyclyl; or

- 5 R^{4a} is optionally substituted C₁-C₄-alkylene that is bound to a carbon atom in Y¹;
- 10 R^{4b} is hydrogen, C₁-C₆-alkyl, halogenated C₁-C₄-alkyl, hydroxy-C₁-C₄-alkyl, C₁-C₆-alkoxy-C₁-C₄-alkyl, amino-C₁-C₄-alkyl, -CH₂CN, -CHO, C₁-C₄-alkylcarbonyl, (halogenated C₁-C₄-alkyl)carbonyl, C₆-C₁₂-arylcarbonyl, C₁-C₄-alkoxycarbonyl, C₆-C₁₂-aryloxycarbonyl, C₁-C₆-alkylaminocarbonyl, C₂-C₆-alkenyl, -C(=NH)NH₂, -C(=NH)NHCN, C₁-C₆-alkylsulfonyl, C₆-C₁₂-arylsulfonyl, amino, -NO or M₃-M₁₂-heterocyclyl; or
- 15 R^{4a}, R^{4b} together are optionally substituted C₂-C₆-alkylene, wherein one -CH₂- of C₂-C₆-alkylene may be replaced by an oxygen atom or -NR¹⁰;
- 20 X² is -O-, -NR^{11a}-, -S-, >CR^{12a}R^{12b} or a bond;
- 20 X³ is -O-, -NR^{11b}-, -S-, >CR^{13a}R^{13b} or a bond;
- 25 R⁵ is optionally substituted C₆-C₁₂-aryl, optionally substituted C₃-C₁₂-cycloalkyl or optionally substituted M₃-M₁₂-heterocyclyl;
- 25 R⁷ is hydrogen or C₁-C₆-alkyl;
- 30 R⁸ is hydrogen, C₁-C₆-alkyl, C₃-C₁₂-cycloalkyl, amino-C₁-C₆-alkyl, optionally substituted C₆-C₁₂-aryl-C₁-C₄-alkyl or M₃-M₁₂-heterocyclyl; or
- 30 R⁸, R¹ together are C₁-C₄-alkylene;
- 35 R⁹ is hydrogen, C₁-C₆-alkyl, C₃-C₁₂-cycloalkyl-C₁-C₄-alkyl, halogenated C₁-C₄-alkyl, hydroxy-C₁-C₄-alkyl, C₁-C₆-alkoxy-C₁-C₄-alkyl, amino-C₁-C₄-alkyl, CH₂CN, C₆-C₁₂-aryl-C₁-C₄-alkyl, C₃-C₁₂-cycloalkyl, -CHO, C₁-C₄-alkylcarbonyl, (halogenated C₁-C₄-alkyl)carbonyl, C₆-C₁₂-arylcarbonyl, C₁-C₄-alkoxycarbonyl, C₆-C₁₂-aryloxycarbonyl, C₁-C₆-alkylaminocarbonyl, C₂-C₆-alkenyl, -C(=NH)NH₂, -C(=NH)NHCN, C₁-C₆-alkylsulfonyl, C₆-C₁₂-arylsulfonyl, amino, -NO or M₃-M₁₂-heterocyclyl;
- 40 R¹⁰ is hydrogen or C₁-C₆-alkyl;

- R^{11a} is hydrogen or C_1 - C_6 -alkyl;
- R^{11b} is hydrogen or C_1 - C_6 -alkyl;
- R^{12a} is hydrogen, optionally substituted C_1 - C_6 -alkyl, C_1 - C_6 -alkylamino- C_1 - C_4 -alkyl, di- C_1 - C_6 -alkylamino- C_1 - C_4 -alkyl, M_3 - M_{12} -heterocyclyl- C_1 - C_6 -alkyl, optionally substituted C_6 - C_{12} -aryl or hydroxy;
- R^{12b} is hydrogen or C_1 - C_6 -alkyl, or
- R^{12a} , R^{12b} together with the carbon atom to which they are attached form a carbonyl or are optionally substituted C_2 - C_4 -alkylene, wherein one $-CH_2-$ of C_2 - C_4 -alkylene may be replaced by an oxygen atom or $-NR^{14}-$;
- R^{13a} is hydrogen, optionally substituted C_1 - C_6 -alkyl, C_1 - C_6 -alkylamino- C_1 - C_4 -alkyl, di- C_1 - C_6 -alkylamino- C_1 - C_4 -alkyl, M_3 - M_{12} -heterocyclyl- C_1 - C_6 -alkyl, optionally substituted C_6 - C_{12} -aryl or hydroxy;
- R^{13b} is hydrogen or C_1 - C_6 -alkyl, or
- R^{13a} , R^{13b} together with the carbon atom to which they are attached form a carbonyl or are optionally substituted C_2 - C_4 -alkylene, wherein one $-CH_2-$ of C_2 - C_4 -alkylene may be replaced by an oxygen atom or $-NR^{15}-$;
- R^{14} is hydrogen or C_1 - C_6 -alkyl; and
- R^{15} is hydrogen or C_1 - C_6 -alkyl,
- or a physiologically tolerated salt thereof.
2. Compound as claimed in claim 1, wherein A is a benzene ring or a ring selected from the group consisting of the following 5- or 6-membered heterocyclic rings:






3. Compound as claimed in claim 1 or 2, wherein R^1 is C_1 - C_6 -alkyl, C_3 - C_{12} -cycloalkyl- C_1 - C_4 -alkyl, or C_3 - C_{12} -cycloalkyl.
4. Compound as claimed in any one of claims 1 to 3, wherein W is a bond and A^1 is a bond.
5. Compound as claimed in any one of claims 1 to 4, wherein Q is $-S(O)_2-$.
6. Compound as claimed in any one of claims 1 to 5, wherein Y is $-NR^8-$.
7. Compound as claimed in any one of claims 1 to 6, wherein at least one of n_1 and n_2 is 1, 2, or 3.
8. Compound as claimed in any one of claims 1 to 7, wherein the sum of n_1+n_2 is 2, 3, or 4.
9. Compound as claimed in any one of claims 1 to 6, wherein X^1 is $>N-$, n_1 is 1, and n_2 is 1; or X^1 is $>CH-$, n_1 is 1, and n_2 is 1.
10. Compound as claimed in any one of claims 1 to 9, wherein R^6 is hydrogen or C_1 - C_4 -alkyl, or two radicals R^6 together with the carbon atom to which they are attached form a carbonyl group.
11. Compound as claimed in any one of claims 1 to 10, having the formula




wherein R^1 , W , A^1 , Q , Y , $n1$, $n2$, X^1 , R^6 , R^2 , A^2 , R^3 , Y^1 , R^{4a} , R^{4b} , X^2 , X^3 , R^5 are as defined in any one of claims 1 to 10.

- 5 12. Compound as claimed in any one of claims 1 to 11, wherein R^2 is hydrogen or halogen.
13. Compound as claimed in claim 11 or 12, having one of the formulae




wherein R^1 , W , A^1 , Q , Y , n_1 , n_2 , X^1 , R^6 , R^2 , A^2 , R^3 , Y^1 , R^{4a} , R^{4b} , X^2 , X^3 , R^5 are as defined in any of claims 1 to 12.

- 5 14. Compound as claimed in any one of claims 1 to 13, wherein A^2 is -O-.
15. Compound as claimed in any one of claims 1 to 14, wherein R^3 is hydrogen or C_1 - C_6 -alkyl.
16. Compound as claimed in any one of claims 1 to 15, having the formula



wherein R^{3a} , R^{3b} , R^{3c} , R^{3d} independently have the meaning of R^3 , and A, R^1 , W, A^1 , Q, Y, $n1$, $n2$, X^1 , R^6 , R^2 , A^2 , R^3 , Y^1 , R^{4a} , R^{4b} , X^2 , X^3 , R^5 are as defined in any one of claims 1 to 15.

- 5 17. Compound as claimed in claim 1 to 16, wherein Y^1 is a bond.
18. Compound as claimed in claim 1 to 17, wherein R^{4a} is hydrogen, C_1 - C_6 -alkyl, optionally substituted C_3 - C_{12} -cycloalkyl, C_3 - C_{12} -cycloalkyl- C_1 - C_4 -alkyl, or M_3 - M_{12} -heterocyclyl.
- 10 19. Compound as claimed in any one of claims 1 to 18, wherein R^{4b} is hydrogen or C_1 - C_6 -alkyl.
20. Compound as claimed in any one of claims 1 to 17, wherein R^{4a} , R^{4b} together are optionally substituted C_2 - C_6 -alkylene, wherein one $-CH_2-$ of C_2 - C_6 -alkylene may be replaced by an oxygen atom.
- 15 21. Compound as claimed in any one of claims 1 to 20, wherein X^2 is $>CR^{12a}R^{12b}$.
22. Compound as claimed in any one of claims 1 to 21, wherein X^3 is a bond.
- 20 23. Compound as claimed in any one of claims 1 to 22, wherein R^{12a} is hydrogen or C_1 - C_6 -alkyl and R^{12b} is hydrogen or C_1 - C_6 -alkyl.
24. Compound as claimed in any one of claims 1 to 23, wherein R^5 is optionally substituted aryl.
- 25 25. Compound as claimed in claim 24, having the formula



wherein A, R^1 , W, A^1 , Q, Y, n_1 , n_2 , X^1 , R^6 , R^2 , R^3 , A^2 , Y^1 , R^{4a} , R^{4b} , X^2 , X^3 are as defined in any one of claims 1 to 23; and

R^{16a} , R^{16b} , R^{16c} , R^{16d} , R^{16e}

5 independently are hydrogen, halogen, or halogenated C_1 - C_6 -alkyl.

26. Compound as claimed in any one of claims 1 to 25, wherein R^8 is hydrogen.

27. Compound as claimed in claim 1, wherein

10

A is a benzene ring;

R^1 is C_1 - C_6 -alkyl, C_3 - C_{12} -cycloalkyl- C_1 - C_4 -alkyl, or an optionally substituted M_3 - M_{12} -heterocyclyl;

15

W is a bond;

A^1 is a bond;

20

Q is $-S(O)_2-$;

Y is $-NR^8-$;

n_1 is 1;

25

n_2 is 1;

- R^6 is hydrogen;
- X^1 is $->N-$ or $>CH-$;
- 5 R^2 is hydrogen or halogen;
- A^2 is $-O-$;
- R^3 is hydrogen;
- 10 Y^1 is a bond;
- R^{4a} is hydrogen, C_1-C_6 -alkyl, C_3-C_{12} -cycloalkyl- C_1-C_4 -alkyl, or C_3-C_{12} -cycloalkyl;
- 15 R^{4b} is hydrogen or C_1-C_6 -alkyl; or
- R^{4a}, R^{4b}
together are C_2-C_6 -alkylene;
- 20 X^2 is $>CR^{12a}R^{12b}$;
- X^3 is a bond;
- R^5 is optionally substituted phenyl;
- 25 R^8 is hydrogen;
- R^{12a} is hydrogen; and
- 30 R^{12b} is hydrogen.
28. The compound as claimed in claim 1 which is:
- N-[1-[3-(azetidin-1-yl)-4-benzyl-chroman-6-yl]azetidin-3-yl]-1-methyl-imidazole-4-sulfonamide;
- 35 Cis-N-(1-(3-(azetidin-1-yl)-4-benzylchroman-6-yl)azetidin-3-yl)ethanesulfonamide;
- Cis-N-(1-(3-(azetidin-1-yl)-4-benzylchroman-6-yl)azetidin-3-yl)propane-1-sulfonamide;
- Cis-N-(1-(3-(azetidin-1-yl)-4-benzylchroman-6-yl)azetidin-3-yl)-1-cyclopropylmethanesulfonamide;
- Cis-N-(1-(3-(azetidin-1-yl)-4-benzylchroman-6-yl)azetidin-3-yl)-1-methyl-1H-pyrazole-4-sulfonamide;
- 40

- Cis-N-(1-(3-(azetidin-1-yl)-4-benzylchroman-6-yl)azetidin-3-yl)-1-methyl-1H-1,2,3-triazole-4-sulfonamide;
- Cis-N-(1-(3-(azetidin-1-yl)-4-benzyl-7-fluorochroman-6-yl)azetidin-3-yl)-1-ethyl-1H-1,2,3-triazole-4-sulfonamide;
- 5 Cis-N-(1-(3-(azetidin-1-yl)-4-benzyl-7-fluorochroman-6-yl)azetidin-3-yl)-2-fluoropyridine-3-sulfonamide;
- Cis-N-(1-(3-(azetidin-1-yl)-4-benzyl-7-fluorochroman-6-yl)azetidin-3-yl)pyridazine-3-sulfonamide;
- Cis-N-(1-(3-(azetidin-1-yl)-4-benzylchroman-6-yl)azetidin-3-yl)-2-fluoropyridine-3-sulfonamide;
- 10 Cis-N-(1-(4-benzyl-3-(methylamino)chroman-6-yl)azetidin-3-yl)propane-1-sulfonamide;
- Cis-N-(1-(4-benzyl-3-(methylamino)chroman-6-yl)azetidin-3-yl)ethanesulfonamide;
- Cis-N-(1-(4-benzyl-3-(methylamino)chroman-6-yl)azetidin-3-yl)-1-cyclopropylmethanesulfonamide;
- 15 Cis-N-(1-(4-benzyl-7-fluoro-3-(methylamino)chroman-6-yl)azetidin-3-yl)ethanesulfonamide;
- Trans-N-(1-(4-benzyl-7-fluoro-3-(methylamino)chroman-6-yl)azetidin-3-yl)ethanesulfonamide;
- Cis-N-(1-(4-benzyl-7-fluoro-3-(methylamino)chroman-6-yl)azetidin-3-yl)-1-methyl-1H-imidazole-4-sulfonamide;
- 20 Trans-N-(1-(4-benzyl-7-fluoro-3-(methylamino)chroman-6-yl)azetidin-3-yl)-1-methyl-1H-imidazole-4-sulfonamide;
- Cis-N-(1-(4-benzyl-7-fluoro-3-(methylamino)chroman-6-yl)azetidin-3-yl)propane-1-sulfonamide;
- Trans-N-(1-(4-benzyl-7-fluoro-3-(methylamino)chroman-6-yl)azetidin-3-yl)propane-1-sulfonamide;
- 25 Cis-N-(1-(4-benzyl-7-fluoro-3-(methylamino)chroman-6-yl)azetidin-3-yl)-1-methyl-1H-pyrazole-4-sulfonamide;
- Trans-N-(1-(4-benzyl-7-fluoro-3-(methylamino)chroman-6-yl)azetidin-3-yl)-1-methyl-1H-pyrazole-4-sulfonamide;
- 30 Cis-N-(1-(4-benzyl-7-fluoro-3-(methylamino)chroman-6-yl)azetidin-3-yl)-1-methyl-1H-1,2,3-triazole-4-sulfonamide;
- Trans-N-(1-(4-benzyl-7-fluoro-3-(methylamino)chroman-6-yl)azetidin-3-yl)-1-methyl-1H-1,2,3-triazole-4-sulfonamide;
- Cis-N-(1-(4-benzyl-7-fluoro-3-(methylamino)chroman-6-yl)azetidin-3-yl)-1-cyclopropylmethanesulfonamide;
- 35 Trans-N-(1-(4-benzyl-7-fluoro-3-(methylamino)chroman-6-yl)azetidin-3-yl)-1-cyclopropylmethanesulfonamide;
- Cis-N-(1-(3-amino-4-benzylchroman-6-yl)azetidin-3-yl)propane-1-sulfonamide;
- Cis-N-(1-(3-amino-4-benzylchroman-6-yl)azetidin-3-yl)-1-cyclopropylmethanesulfonamide;

Cis-1-(4-benzyl-7-fluoro-6-(3-(5-fluoropyridine-3-sulfonamido)azetidin-1-yl)chroman-3-yl)azetidin-1-ium (E)-3-carboxyacrylate;

Cis-N-(1-(4-benzyl-3-(propylamino)chroman-6-yl)azetidin-3-yl)propane-1-sulfonamide;

Cis-N-(1-(4-benzyl-3-(diethylamino)chroman-6-yl)azetidin-3-yl)propane-1-sulfonamide;

5 Cis-N-(1-(4-benzyl-3-((cyclopropylmethyl)amino)chroman-6-yl)azetidin-3-yl)propane-1-sulfonamide; and

Cis-N-(1-(4-benzyl-3-(cyclobutylamino)chroman-6-yl)azetidin-3-yl)propane-1-sulfonamide, or a physiologically tolerated salt thereof.

10 29. The compound as claimed in any one of claims 1 to 28 for use in therapy.

30. Pharmaceutical composition which comprises a carrier and a compound of any one of claims 1 to 28.

15 31. A method for inhibiting the glycine transporter GlyT1 in a mammal in need thereof which comprises the administration of an effective amount of a compound of any one of claims 1 to 28.

20 32. The use of a compound of any one of claims 1 to 28 in the manufacture of a medicament for inhibiting the glycine transporter GlyT1.

25 33. A method for treating a neurologic or psychiatric disorder or pain in a mammalian patient in need thereof which comprises administering to the patient a therapeutically effective amount of a compound of any one of claims 1 to 28.

34. The use of a compound of any one of claims 1 to 28 in the manufacture of a medicament for treating a neurologic or psychiatric disorder or pain.

30 35. The compound of any one of claims 1 to 28 for use in a method of treating a neurologic or psychiatric disorder or pain.

36. The method, use or compound as claimed in any one of claims 30 to 35, wherein the disorder is associated with glycinergic or glutamatergic neurotransmission dysfunction.

35 37. The method, use or compound as claimed in any one of claims 30 to 36, wherein the neurologic disorder is a cognitive disorder such as dementia, cognitive impairment, or attention deficit disorder.

40 38. The method, use or compound as claimed in claim 37, wherein the attention deficit disorder is an attention deficit disorder with hyperactivity.

39. The method, use or compound as claimed in any one of any one of claims 30 to 35, wherein the psychiatric disorder is an anxiety disorder, a mood disorder such as depression, a bipolar disorder, schizophrenia, or a psychotic disorder.