

The present invention relates to amylin analogues that are amylin receptor agonists, and to their medical use in the treatment and/or prevention of a variety of diseases, conditions or disorders, including treatment and/or prevention of excess food intake, obesity and excess body weight, metabolic diseases, and other conditions and disorders described herein. In particular, the present invention relates to stable amylin analogues that have a long duration of action and are well suited for use in the form of a liquid formulation.

BACKGROUND OF THE INVENTION

Amylin is one of a family of peptide hormones that includes amylin, calcitonin, calcitonin gene-related peptide, adrenomedullin and intermedin (intermedin also being known as AFP-6), and has been implicated in various metabolic diseases and disorders. Human amylin was first isolated, purified and characterized as the major component of amyloid deposits in the islets of pancreases from type 2 diabetes patients.

Native human amylin is a 37-amino acid peptide having the formula

H-KC()NTATC()ATQRLANFLVHSSNNFGAILSSTNVGSNTY-NH₂

wherein H- at the N-terminus designates a hydrogen atom, corresponding to the presence of a free amino group on the N-terminal amino acid residue [i.e. the lysine (K) residue at sequence position number 1 in the sequence shown above]; wherein -NH₂ at the C-terminus indicates that the C-terminal carboxyl group is in the amide form; and wherein the parentheses () associated with the two cysteine (C, Cys) residues at sequence positions 2 and 7 indicate the presence of an intramolecular disulfide bridge between the two Cys residues in question.

Amylin may be beneficial in treating metabolic disorders such as diabetes and/or obesity. Amylin is believed to regulate gastric emptying, and to suppress glucagon secretion and food intake, thereby regulating the rate of glucose release to the circulation. Amylin appears to complement the actions of insulin. Compared to healthy adults, type 1 diabetes patients have no circulating amylin, and type 2 diabetes patients exhibit reduced postprandial amylin concentrations. In human trials an amylin analogue known as pramlintide, described in WO 93/10146 and having the sequence Lys-Cys-Asn-Thr-Ala-Thr-Cys-Ala-Thr-Gln-Arg-Leu-Ala-Asn-Phe-Leu-Val-His-Ser-Ser-Asn-Asn-Phe-Gly-Pro-Ile-Leu-Pro-Pro-Thr-Asn-Val-Gly-Ser-Asn-Thr-Tyr, which also possesses a disulphide bridge between the Cys residues at positions 2 and 7, has been shown to reduce body weight or

reduce weight gain. An alternative amylin analogue incorporating N-methylated residues and having a reduced tendency to fibrillation, designated IAPP-GI, has been described by Yan et al. (PNAS, 103(7), 2046-2051, 2006). IAPP-GI appears to have lower activity than native amylin, however. Further analogues of amylin or pramlintide are described in
5 WO2013/156594, WO2012/168430, WO2012/168431 and WO2012/168432, as well as WO2006/042745.


Obesity is believed to be a major causal factor in development of type 2 diabetes, which constitutes a growing and worldwide major health problem. Diseases or disorders that
10 may develop as a consequence of untreated diabetes include cardiovascular and peripheral artery disease, micro- and macrovascular complications, stroke, and certain forms of cancer, particularly hematopoietic cancers.

There is a need in the art for further amylin analogues. For example, amylin analogues
15 that show a reduced tendency for fibrillation and/or high chemical stability at or around pH 7 might allow for a formulation at or near physiological pH. Amylin analogues having high levels of agonist activity at the amylin receptor and/or appropriately long plasma elimination half lives, may also enable longer intervals between dosing than is currently possible (e.g. once weekly, or even less frequently) and hence improve patient
20 compliance.

SUMMARY

The present invention relates to compounds which are analogues of human amylin. The subject matter of the present invention is as defined in the depending claims.
25

Also described herein is an amylin analogue which is a compound having the formula:



wherein

R¹ is hydrogen, C₁₋₄ acyl, benzoyl or C₁₋₄ alkyl, or a half-life extending moiety M, wherein
30 M is optionally linked to Z via a linker moiety L;

R² is OH or NHR³, wherein R³ is hydrogen or C₁₋₃-alkyl; and

Z is an amino acid sequence of formula I:

Arg-Cys-X3-Thr-Ala-Thr-Cys-Ala-Thr-X10-Arg-Leu-Ala-X14-Phe-Leu-X17-Arg-X19-X20-Phe-Gly(Me)-Ala-Ile(Me)-Leu-Ser-Ser-Thr-X31-Val-Gly-Ser-X35-Thr-X37 (I);

wherein

X3 is selected from the group consisting of Asn, Gly, Pro and Gln;

5 X10 is selected from the group consisting of Gln, Asp and Glu;

X14 is selected from the group consisting of Asp, His, Asn and Aad;

X17 is selected from the group consisting of His, Asn, Gln, Glu, Thr, Val, Lys and Aad;

X19-X20 is selected from the group consisting of Ser-Ser, Val-Val, Ser-Val and Val-Ser, or is absent;

10 X31 is selected from the group consisting of Asp, Glu and Asn;

X35 is selected from the group consisting of Asp, Glu, Asn, Ser, Phe, Orn, Aad, Gly and Thr; and

X37 is selected from the group consisting of Pro, Apr and Hyp;

and wherein the compound has at least one residue selected from:

15 X3 is Gln;

X14 is His, Asn or Aad;

X17 is Asn, Gln, Glu, Thr or Aad;

X19-X20 is Val-Ser or Ser-Val; and

X35 is Ser, Phe, Orn, Aad, Gly or Thr;

20 or a pharmaceutically acceptable salt or solvate thereof.

Throughout this specification, amino acid positions of the amylin analogues are numbered according to the corresponding position in human amylin having the sequence shown above. The sequence of Formula I (and other formulae herein) contains a two amino acid
25 deletion corresponding to the two residues Asn21 and Asn22 of human amylin. Thus, for ease of comparison with the amylin sequence, the Phe residue immediately C-terminal (downstream) of position X20 is designated as position 23, since it aligns with Phe23 of

human amylin. Thus, the numbering of any given residue in Formula I above, and in other formulae elsewhere in this specification, reflects the corresponding residue in human amylin when optimally aligned therewith and does not necessarily reflect its linear position in the particular sequence.

5

(It will be apparent that any of the relevant formulae presented in this specification could be written to include residues X21-X22 at the appropriate positions, wherein X21 and X22 are absent.)

10 It has surprisingly been found that simultaneous deletion of the residues at positions X21 and X22 (and even additionally at positions X19 and X20) leads to active and stable amylin analogues. Further, without wishing to be bound by any particular theory, it is believed that the deletion of these residues may enhance the chemical stability of the molecules, especially at neutral and/or alkaline pH. Fibrillation and precipitation of the
15 compounds may also be reduced. Thus the compounds may have superior properties for formulation as compared to existing amylin analogues.

Furthermore, the compounds described here show similar or even increased activity compared to wild type amylin (e.g. agonist activity at the hAMYR3 and/or hAMYR1 and/or
20 hCTR2 receptors), despite being methylated at the same positions as IAPP-GI (which has lower activity than wild type amylin).

In some examples of formula I it may be desirable that:

X31 is Glu;

25 X19-X20 is Ser-Ser or is absent; and/or

X37 is Hyp or Pro.

In one aspect, the present disclosure refers to an amylin analogue selected from the group consisting of: [19CD]-isoGlu-RCNTATCATQRLADFLHRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSETP-NH₂.

30 In another aspect, the disclosure refers to the use of an amylin analogue, pharmaceutically acceptable salt or solvate in the preparation of a medicament: (a) for treating obesity, morbid obesity, obesity prior to surgery, obesity-linked inflammation, obesity-linked gallbladder disease, obesity-induced sleep apnea and respiratory problems, degeneration of cartilage, osteoarthritis, or reproductive health complications of obesity or overweight such as infertility; or (b) for prevention or
35 treatment of Alzheimer's disease, diabetes, type 1 diabetes, type 2 diabetes, pre-diabetes, insulin resistance syndrome, impaired glucose tolerance (IGT), disease states associated with elevated blood glucose levels, metabolic disease including metabolic syndrome, hyperglycemia, hypertension, atherogenic dyslipidemia, hepatic steatosis ("fatty liver"), kidney failure, arteriosclerosis (e.g. atherosclerosis), macrovascular disease, microvascular disease, diabetic
40 heart (including diabetic cardiomyopathy and heart failure as a diabetic complication) coronary heart disease, peripheral artery disease or stroke, and combinations thereof.

It may be desirable that the amylin analogue contains at least one of His14, Asn14, Aad14, Gln17 and Thr17.

If X14 is Asp, then it may be desirable that X17 is Asn, Gln, Glu, Thr or Aad.

X17 is Gln may be particularly preferred.

- 5 In some circumstances, it may be desirable that X35 is not a hydrophobic residue, e.g. Phe. Such residues may increase tendency towards fibrillation in some formulations.

Z may be an amino acid sequence of formula II:

- 10 Arg-Cys-X3-Thr-Ala-Thr-Cys-Ala-Thr-X10-Arg-Leu-Ala-X14-Phe-Leu-X17-Arg-X19-X20-Phe-Gly(Me)-Ala-Ile(Me)-Leu-Ser-Ser-Thr-Glu-Val-Gly-Ser-X35-Thr-X37 (II);

wherein

X3 is selected from the group consisting of Asn, Gly and Gln;

X10 is selected from the group consisting of Gln, Asp and Glu;

X14 is selected from the group consisting of Asp, His and Aad;

- 15 X17 is selected from the group consisting of His, Asn, Gln, Glu, Lys and Aad;

X19-X20 is Ser-Ser or is absent;

X35 is selected from the group consisting of Asp, Glu, Asn, Ser, Orn, Aad, Gly and Thr; and

X37 is selected from the group consisting of Pro and Hyp;

and wherein the compound has at least one residue selected from:

- 20 X3 is Gln;

X14 is His or Aad;

X17 is Asn, Gln, Glu or Aad; and

X35 is Ser, Phe, Orn, Aad, Gly or Thr.

In some examples of formula II, X17 may be selected from His and Gln.

- 25 Z may be an amino acid sequence of formula III:

Arg-Cys-X3-Thr-Ala-Thr-Cys-Ala-Thr-X10-Arg-Leu-Ala-X14-Phe-Leu-X17-Arg-X19-X20-Phe-Gly(Me)-Ala-Ile(Me)-Leu-Ser-Ser-Thr-Glu-Val-Gly-Ser-X35-Thr-X37 (III);

wherein

X3 is selected from the group consisting of Asn, Gly and Gln;

- 30 X10 is selected from the group consisting of Gln, Asp and Glu;

X14 is selected from the group consisting of Asp, His and Aad;

X17 is selected from the group consisting of His and Gln;

X19-X20 is Ser-Ser or is absent;

X35 is selected from the group consisting of Asp, Glu, Asn, Aad and Gly; and

5 X37 is selected from the group consisting of Pro and Hyp;

and wherein the compound has at least one residue selected from:

X3 is Gln;

X14 is His or Aad;

X17 is Gln; and

10 X35 is Aad.

In any of the formulae described above, it may be desirable that:

X10 is selected from Gln and Glu;

and/or

15 X35 is selected from Asp, Glu, Asn and Aad, e.g. X35 is selected from Asp and Asn.

Additionally or alternatively, it may be that X3 is selected from Asn and Gly and/or X17 is Gln. The presence of Gln at position X17 is believed to correlate with good levels of chemical and physical stability.

20

Additionally or alternatively, X35 is Asn and/or X37 is Hyp.

In some examples of the formulae described above:

X3 is Gln;

25 X10 is selected from Glu and Asp;

X14 is selected from His and Aad;

X35 is selected from Gly and Asn; and

X37 is selected from Pro and Hyp.

In other examples of the formulae described above:

X3 is Gly;

5 X10 is selected from Glu and Asp;

X14 is selected from His and Aad;

X35 is selected from Gly and Asn; and

X37 is selected from Pro and Hyp.

10 Combinations of specific residues which may be present in any of the amylin analogues described include:

Gly3 + Glu10;

Asn3 + Glu10;

Gln3 + Glu10;

15 Asn3 + Gln10; or

Gln3 + Asp10.

In some examples, X14 is selected from His and Aad, and/or X17 is Gln.

20 For example, the combination of Aad14 and Gln17 appears to provide good activity.

Additionally or alternatively, X17 may be Gln, X31 may be Glu and X37 may be Hyp, i.e. the analogue may contain the combination Gln17 + Glu31 + Hyp37.

In some examples, X19-X20 is Ser-Ser. In other examples X19-X20 is absent.

Certain residue combinations which may be favourable for chemical stability include:

X3 is Gly, X10 is Glu and X14 is His;

X3 is selected from Asn and Gln, X10 is Glu and X14 is His;

5 X3 is Gly, X10 is Glu and X14 is selected from Aad and Asp;

X10 is Asp and X14 is Aad;

X14 is selected from Aad and His, X31 is Glu and X37 is selected from Pro and Hyp; and
X14 is Aad, X31 is Glu and X37 is Hyp.

10 Of these combinations, the following may additionally (or alternatively) have a favourable effect on activity:

X3 is Gly, X10 is Glu and X14 is selected from Aad and Asp;

X14 is selected from Aad and His, X31 is Glu and X37 is selected from Pro and Hyp;

X14 is Aad, X31 is Glu and X37 is Hyp.

15

For example, the compound may comprise the residues:

Gly3 + Glu10 + His14

Asn3 + Glu10 + His14;

Gln3 + Glu10 + His14;

20 Gly3 + Glu10 + Aad14;

Gly3 + Glu10 + Asp14;

Asp10 + Aad14;

His14 + Glu31 + Pro37

His14 + Glu31 + Hyp37

25 Aad14 + Glu31 + Pro37; and

Aad14 + Glu31 + Hyp37.

Yet further combinations of desirable residues include:

Gly3 + Glu31;

30 Gly3 + Ser19 + Ser20 + Glu31


Gly3 + Glu10 + Glu31 + Asn35 + Hyp37; and

Gly3 + Glu10 + Ser19 + Ser20 + Glu31 + Asn35 + Hyp37.

Any of the above-described residues and combinations of residues may be combined

35 except where inconsistent with one another.

In another aspect, the invention also provides an amylin analogue which is a compound having the formula:



wherein

- 5 R^1 is hydrogen, C_{1-4} acyl, benzoyl or C_{1-4} alkyl, or a half-life extending moiety M, wherein M is optionally linked to Z via a linker moiety L;

R^2 is OH or NHR^3 , wherein R^3 is hydrogen or C_{1-3} -alkyl; and

Z is an amino acid sequence selected from the group consisting of:

RCNTATCATQRLADFLHRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSETP;

- 10 RCNTATCATQRLADFLHRSSNNF-Gly(Me)-A-Ile(Me)-LSSTNVGSNT-Apr; and
RCNTATCATQRLAHFLHRSSNNF-Gly(Me)-A-Ile(Me)-LSSTNVGSNT-Apr;
or a pharmaceutically acceptable salt or solvate thereof.

Thus, the amylin analogue may have the formula:

- 15 R^1-Z-R^2

wherein

R^1 is hydrogen, C_{1-4} acyl, benzoyl or C_{1-4} alkyl, or a half-life extending moiety M, wherein M is optionally linked to Z via a linker moiety L;

R^2 is OH or NHR^3 , wherein R^3 is hydrogen or C_{1-3} -alkyl; and

- 20 Z is an amino acid sequence selected from the group consisting of:

RCNTATCATQRLAHFLHRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSETP

RCNTATCATQRLANFLHRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSETP

RCGTATCATERLAHFLNRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGsNT-Hyp

RCGTATCATERLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGsNT-Hyp

- 25 RCGTATCATERLAHFLERSSF-Gly(Me)-A-Ile(Me)-LSSTEVGsNT-Hyp

RCGTATCATERLAHFLTRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGsST-Hyp

RCGTATCATERLAHFLTRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGsFT-Hyp

- RCGTATCATERLAHFLTRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGGS-Orn-I-Hyp
- RCGTATCATERLAHFLTRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGGS-Aad-T-Hyp
- RCGTATCATERLAHFLTRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSGT-Hyp
- RCNTATCATQRLAHFLHRF-Gly(Me)-A-Ile(Me)-LSSTEVGSETP
- 5 RCGTATCATERLAHFLHRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSNTP
- RCNTATCATQRLAHFLHRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSNTP
- RCGTATCATERLANFLVRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSNTP-Hyp
- RCGTATCATERLAHFLTRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSNTP-Hyp
- RCGTATCATERLAHFLKRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSNTP-Hyp
- 10 RCNTATCATQRLAHFLHRVSF-Gly(Me)-A-Ile(Me)-LSSTEVGSETP
- RCNTATCATQRLAHFLHRVSF-Gly(Me)-A-Ile(Me)-LSSTEVGSETP
- RCGTATCATERLAHFLVRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSETP
- RCPTATCATDRLAHFLVRSSF-Gly(Me)-A-Ile(Me)-LSSTDVGSSTP
- RCNTATCATQRLAHFLVRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSETP
- 15 RCPTATCATDRLAHFLHRSSF-Gly(Me)-A-Ile(Me)-LSSTDVGSNTP
- RCGTATCATERLAHFLHRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSETP
- RCPTATCATDRLAHFLHRSSF-Gly(Me)-A-Ile(Me)-LSSTDVGSSTP
- RCGTATCATERLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSSST-Hyp
- RCGTATCATERLAHFLNRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSSST-Hyp
- 20 RCGTATCATERLAHFLHRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSSST-Hyp
- RCGTATCATERLAHFLVRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSSST-Hyp
- RCGTATCATERLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGST-Hyp
- RCGTATCATERLAHFLNRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGST-Hyp
- RCGTATCATERLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSSTP
- 25 RCGTATCATERLAHFLNRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSSTP
- RCGTATCATERLAHFLERSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSSTP
- RCGTATCATERLAHFLHRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSSTP
- RCGTATCATERLAHFLVRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSSTP
- RCGTATCATERLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSTP

- RCGTATCATERLAHFLNRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSTFP
- RCGTATCATERLADFLTRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSSST-Hyp
- RCGTATCATERLADFLTRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGST-Hyp
- RCGTATCATERLA-Aad-FLTRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSSST-Hyp
- 5 RCGTATCATERLA-Aad-FLTRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGST-Orn-T-Hyp
- RCGTATCATERLAHFLTRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSSST-Apr
- RCGTATCATERLAHFLTRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSSSTP
- RCGTATCATERLAHFLTRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGST-Orn-TP
- RCGTATCATERLAHFLTRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSTT-Hyp
- 10 RCNTATCATQRLADFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGST-Hyp
- RCNTATCATQRLAHFL-Aad-RSSF-Gly(Me)-A-Ile(Me)-LSSTEVGST-Hyp
- RCGTATCATERLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGST-Orn-T-Hyp
- RCGTATCATERLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSGT-Hyp
- RCGTATCATERLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGST-Aad-T-Hyp
- 15 RCGTATCATERLADFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGST-Hyp
- RCGTATCATERLA-Aad-FLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGST-Hyp
- RCNTATCATERLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGST-Hyp
- RCGTATCATERLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGST-Hyp
- RCNTATCATQRLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGST-Hyp
- 20 RCPTATCATDRLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTDVGST-Hyp
- RCNTATCATERLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSDT-Hyp
- RCNTATCATQRLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSTNP
- RCNTATCATERLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSTNP
- RCPTATCATERLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGST-Hyp
- 25 RCPTATCATQRLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGST-Hyp
- RCNTATCATQRLADFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSTNP
- RCNTATCATERLADFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSTNP
- RCPTATCATERLADFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSTNP
- RCNTATCATERLADFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGST-Hyp

- RCNTATCATQRLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSSST-Hyp
- RCPTATCATERLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSSST-Hyp
- RCNTATCATQRLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSGT-Hyp
- RCPTATCATERLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSGT-Hyp
- 5 RCNTATCATQRLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSTT-Hyp
- RCPTATCATERLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSTT-Hyp
- RCNTATCATQRLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSAad-T-Hyp
- RCPTATCATERLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSAad-T-Hyp
- RCGTATCATERLAHFLQRF-Gly(Me)-A-Ile(Me)-LSSTEVGSNTHyp
- 10 RCGTATCATERLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSNTP
- RCNTATCATQRLAHFLQRF-Gly(Me)-A-Ile(Me)-LSSTEVGSGT-Hyp
- RCNTATCATQRLADFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSGTP
- RCNTATCATQRLADFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSGT-Hyp
- RCNTATCATQRLADFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSTTP
- 15 RCNTATCATQRLADFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSTT-Hyp
- RCNTATCATQRLAHFLERSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSSSTP
- RCNTATCATQRLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSSSTP
- RCNTATCATQRLA-Aad-FLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSNTHyp
- RCNTATCATQRLA-Aad-FLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSSST-Hyp
- 20 RCQTATCATDRLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSNTHyp
- RCQTATCATDRLA-Aad-FLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSNTHyp
- RCQTATCATERLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSGT-Hyp
- RCQTATCATDRLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSGT-Hyp
- RCQTATCATDRLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSNTP
- 25 RCQTATCATERLA-Aad-FLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSNTP
- RCQTATCATERLA-Aad-FLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSGTP
- RCQTATCATDRLA-Aad-FLQRF-Gly(Me)-A-Ile(Me)-LSSTEVGSNTHyp
- RCQTATCATERLAHFLQRF-Gly(Me)-A-Ile(Me)-LSSTEVGSNTP

or a pharmaceutically acceptable salt or solvate thereof.

In some examples, R¹ is M or M-L-, and/or R² is NH₂.

Specific amylin analogues of the invention include:

[19CD]-isoGlu-RCNTATCATQRLADFLHRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSETP-NH₂

5 [19CD]-isoGlu-RCNTATCATQRLAHFLHRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSETP-NH₂

[19CD]-isoGlu-RCNTATCATQRLANFLHRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSETP-NH₂

[19CD]-isoGlu-RCNTATCATQRLADFLHRSSNNF-Gly(Me)-A-Ile(Me)-LSSTNVGSNT-Apr-NH₂

10 [19CD]-isoGlu-RCNTATCATQRLAHFLHRSSNNF-Gly(Me)-A-Ile(Me)-LSSTNVGSNT-Apr-NH₂

[19CD]-isoGlu-RCGTATCATERLAHFLNRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSNH-T-Hyp-NH₂

[19CD]-isoGlu-RCGTATCATERLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSNH-T-Hyp-NH₂

15 [19CD]-isoGlu-RCGTATCATERLAHFLERSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSNH-T-Hyp-NH₂

[19CD]-isoGlu-RCGTATCATERLAHFLTRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSS-T-Hyp-NH₂

[19CD]-isoGlu-RCGTATCATERLAHFLTRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGST-T-Hyp-NH₂

20 [19CD]-isoGlu-RCGTATCATERLAHFLTRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSO-T-Hyp-NH₂

[19CD]-isoGlu-RCGTATCATERLAHFLTRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSA-T-Hyp-NH₂

25 [19CD]-isoGlu-RCGTATCATERLAHFLTRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSG-T-Hyp-NH₂

[19CD]-isoGlu-RCNTATCATQRLAHFLHRF-Gly(Me)-A-Ile(Me)-LSSTEVGSETP-NH₂

[19CD]-isoGlu-RCGTATCATERLAHFLHRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSNTP-NH₂

[19CD]-isoGlu-RCNTATCATQRLAHFLHRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSNTP-NH₂

[19CD]-isoGlu-RCGTATCATERLANFLVRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSNTP-Hyp-NH₂

5 [19CD]-isoGlu-RCGTATCATERLAHFLTRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSNTP-Hyp-NH₂

[19CD]-isoGlu-RCGTATCATERLAHFLKRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSNTP-Hyp-NH₂

[19CD]-isoGlu-RCNTATCATQRLAHFLHRVSF-Gly(Me)-A-Ile(Me)-LSSTEVGSETP-NH₂

[19CD]-isoGlu-RCNTATCATQRLAHFLHRVSF-Gly(Me)-A-Ile(Me)-LSSTEVGSETP-NH₂

10 [19CD]-isoGlu-RCGTATCATERLAHFLVRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSETP-NH₂

[19CD]-isoGlu-RCPTATCATDRLAHFLVRSSF-Gly(Me)-A-Ile(Me)-LSSTDVGSDTP-NH₂

[19CD]-isoGlu-RCNTATCATQRLAHFLVRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSETP-NH₂

[19CD]-isoGlu-RCPTATCATDRLAHFLHRSSF-Gly(Me)-A-Ile(Me)-LSSTDVGSNTP-NH₂

[19CD]-isoGlu-RCGTATCATERLAHFLHRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSETP-NH₂

15 [19CD]-isoGlu-RCPTATCATDRLAHFLHRSSF-Gly(Me)-A-Ile(Me)-LSSTDVGSDTP-NH₂

[19CD]-isoGlu-RCGTATCATERLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSSST-Hyp-NH₂

[19CD]-isoGlu-RCGTATCATERLAHFLNRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSSST-Hyp-NH₂

20 [19CD]-isoGlu-RCGTATCATERLAHFLHRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSSST-Hyp-NH₂

[19CD]-isoGlu-RCGTATCATERLAHFLVRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSSST-Hyp-NH₂

25 [19CD]-isoGlu-RCGTATCATERLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGST-Hyp-NH₂

[19CD]-isoGlu-RCGTATCATERLAHFLNRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGST-Hyp-NH₂

- [19CD]-isoGlu-RCGTATCATERLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSSSTP-NH₂
- [19CD]-isoGlu-RCGTATCATERLAHFLNRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSSSTP-NH₂
- [19CD]-isoGlu-RCGTATCATERLAHFLERSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSSSTP-NH₂
- [19CD]-isoGlu-RCGTATCATERLAHFLHRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSSSTP-NH₂
- 5 [19CD]-isoGlu-RCGTATCATERLAHFLVRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSSSTP-NH₂
- [19CD]-isoGlu-RCGTATCATERLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSSFTP-NH₂
- [19CD]-isoGlu-RCGTATCATERLAHFLNRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSSFTP-NH₂
- [19CD]-isoGlu-RCGTATCATERLADFLTRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSSST-Hyp-NH₂
- 10 [19CD]-isoGlu-RCGTATCATERLADFLTRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSSFT-Hyp-NH₂
- [19CD]-isoGlu-RCGTATCATERLA-Aad-FLTRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSSST-Hyp-NH₂
- [19CD]-isoGlu-RCGTATCATERLA-Aad-FLTRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSS-Orn-T-Hyp-NH₂
- 15 [19CD]-isoGlu-RCGTATCATERLAHFLTRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSSST-Apr-NH₂
- [19CD]-isoGlu-RCGTATCATERLAHFLTRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSSSTP-NH₂
- [19CD]-isoGlu-RCGTATCATERLAHFLTRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSS-Orn-TP-NH₂
- [19CD]-isoGlu-RCGTATCATERLAHFLTRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSTT-Hyp-NH₂
- 20 [19CD]-isoGlu-RCNTATCATQRLADFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSSNT-Hyp-NH₂
- [19CD]-isoGlu-RCNTATCATQRLAHFL-Aad-RSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSSNT-Hyp-NH₂
- [19CD]-isoGlu-RCGTATCATERLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSS-Orn-T-
- 25 Hyp-NH₂

- [19CD]-isoGlu-RCGTATCATERLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSGT-Hyp-NH₂
- [19CD]-isoGlu-RCGTATCATERLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSAad-T-Hyp-NH₂
- 5 [19CD]-isoGlu-RCGTATCATERLADFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSNH-T-Hyp-NH₂
- [19CD]-isoGlu-RCGTATCATERLA-Aad-FLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSNH-T-Hyp-NH₂
- [19CD]-isoGlu-RCNTATCATERLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSNH-T-Hyp-NH₂
- 10 [19CD]-isoGlu-RCQTATCATERLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSNH-T-Hyp-NH₂
- [19CD]-isoGlu-RCNTATCATQRLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSNH-T-Hyp-NH₂
- 15 [19CD]-isoGlu-RCPTATCATDRLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTDVGSNH-T-Hyp-NH₂
- [19CD]-isoGlu-RCNTATCATERLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSDT-Hyp-NH₂
- [19CD]-isoGlu-RCNTATCATQRLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSNTP-NH₂
- 20 [19CD]-isoGlu-RCNTATCATERLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSNTP-NH₂
- [19CD]-isoGlu-RCPTATCATERLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSNH-T-Hyp-NH₂
- [19CD]-isoGlu-RCPTATCATQRLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSNH-T-Hyp-NH₂
- 25 [19CD]-isoGlu-RCNTATCATQRLADFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSNTP-NH₂
- [19CD]-isoGlu-RCNTATCATERLADFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSNTP-NH₂
- [19CD]-isoGlu-RCPTATCATERLADFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSNTP-NH₂

- [19CD]-isoGlu-RCNTATCATERLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGST-Hyp-NH₂
- [19CD]-isoGlu-RCNTATCATQRLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGST-Hyp-NH₂
- 5 [19CD]-isoGlu-RCPTATCATERLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGST-Hyp-NH₂
- [19CD]-isoGlu-RCNTATCATQRLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSGT-Hyp-NH₂
- [19CD]-isoGlu-RCPTATCATERLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSGT-Hyp-NH₂
- 10 [19CD]-isoGlu-RCNTATCATQRLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSTT-Hyp-NH₂
- [19CD]-isoGlu-RCPTATCATERLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSTT-Hyp-NH₂
- 15 [19CD]-isoGlu-RCNTATCATQRLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSAad-T-Hyp-NH₂
- [19CD]-isoGlu-RCPTATCATERLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSAad-T-Hyp-NH₂
- [19CD]-isoGlu-RCGTATCATERLAHFLQRF-Gly(Me)-A-Ile(Me)-LSSTEVGST-Hyp-NH₂
- 20 [19CD]-isoGlu-RCGTATCATERLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSTP-NH₂
- [19CD]-isoGlu-RCNTATCATQRLAHFLQRF-Gly(Me)-A-Ile(Me)-LSSTEVGSGT-Hyp-NH₂
- [19CD]-isoGlu-RCNTATCATQRLADFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSGTP-NH₂
- [19CD]-isoGlu-RCNTATCATQRLADFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSGT-Hyp-NH₂
- 25 [19CD]-isoGlu-RCNTATCATQRLADFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSTTP-NH₂
- [19CD]-isoGlu-RCNTATCATQRLADFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSTT-Hyp-NH₂

[19CD]-isoGlu-RCNTATCATQRLAHFLERSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSSSTP-NH₂

[19CD]-isoGlu-RCNTATCATQRLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSSSTP-NH₂

[19CD]-isoGlu-RCNTATCATQRLA-Aad-FLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSSNT-Hyp-NH₂

5 [19CD]-isoGlu-RCNTATCATQRLA-Aad-FLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSSST-Hyp-NH₂

[19CD]-isoGlu-RCQTATCATDRLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSSNT-Hyp-NH₂

10 [19CD]-isoGlu-RCQTATCATDRLA-Aad-FLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSSNT-Hyp-NH₂

[19CD]-isoGlu-RCQTATCATERLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSGT-Hyp-NH₂

[19CD]-isoGlu-RCQTATCATDRLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSGT-Hyp-NH₂

15 [19CD]-isoGlu-RCQTATCATDRLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSSNTP-NH₂

[19CD]-isoGlu-RCQTATCATERLA-Aad-FLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSSNTP-NH₂

[19CD]-isoGlu-RCQTATCATERLA-Aad-FLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSGTP-NH₂

20 [19CD]-isoGlu-RCQTATCATDRLA-Aad-FLQRF-Gly(Me)-A-Ile(Me)-LSSTEVGSSNT-Hyp-NH₂

[19CD]-isoGlu-RCQTATCATERLAHFLQRF-Gly(Me)-A-Ile(Me)-LSSTEVGSSNTP-NH₂

wherein [19CD] represents [19-carboxynonadecanoyl];

25 and pharmaceutically acceptable salts and solvates thereof.

Any of the sequences or compounds described above may possess an intramolecular disulphide bridge formed between the cysteine residues present at positions 2 and 7 of

the amino acid sequence (numbered from N- to C-terminus, and corresponding to the cysteine residues present at positions 2 and 7 of human amylin). In general, it may be desirable that compounds possess such a disulphide bridge at the time of administration to a subject, but it will be understood that the invention extends to compounds having the specified amino acid sequences before formation of the disulphide. The presence of the disulphide may be indicated by parentheses () following each relevant cysteine residue in the sequence. All of the generic and specific formulae provided above should be construed accordingly to include this possibility. Thus, for example, Formulae I, II and III may be shown as follows:

10 Arg-Cys()-X3-Thr-Ala-Thr-Cys()-Ala-Thr-X10-Arg-Leu-Ala-X14-Phe-Leu-X17-Arg-X19-X20-Phe-Gly(Me)-Ala-Ile(Me)-Leu-Ser-Ser-Thr-X31-Val-Gly-Ser-X35-Thr-X37 (I);

Arg-Cys()-X3-Thr-Ala-Thr-Cys()-Ala-Thr-X10-Arg-Leu-Ala-X14-Phe-Leu-X17-Arg-X19-X20-Phe-Gly(Me)-Ala-Ile(Me)-Leu-Ser-Ser-Thr-Glu-Val-Gly-Ser-X35-Thr-X37 (II); and

15 Arg-Cys()-X3-Thr-Ala-Thr-Cys()-Ala-Thr-X10-Arg-Leu-Ala-X14-Phe-Leu-X17-Arg-X19-X20-Phe-Gly(Me)-Ala-Ile(Me)-Leu-Ser-Ser-Thr-Glu-Val-Gly-Ser-X35-Thr-X37 (III);

while the specific compounds listed above may be designated as follows:

[19CD]-isoGlu-R-C()-NTAT-C()-ATQRLADFLHRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSETP-NH₂ (Compd. 1)

20 [19CD]-isoGlu-R-C()-NTAT-C()-ATQRLAHFLHRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSETP-NH₂ (Compd. 2)

[19CD]-isoGlu-R-C()-NTAT-C()-ATQRLANFLHRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSETP-NH₂ (Compd. 3)

[19CD]-isoGlu-R-C()-NTAT-C()-ATQRLADFLHRSSNNF-Gly(Me)-A-Ile(Me)-LSST-NVGSNT-Apr-NH₂ (Compd. 4)

25 [19CD]-isoGlu-R-C()-NTAT-C()-ATQRLAHFLHRSSNNF-Gly(Me)-A-Ile(Me)-LSST-NVGSNT-Apr-NH₂ (Compd. 5)

[19CD]-isoGlu-R-C()-GTAT-C()-ATERLAHFLNRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSNTHyp-NH₂ (Compd. 6)

30 [19CD]-isoGlu-R-C()-GTAT-C()-ATERLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSNTHyp-NH₂ (Compd. 7)

- [19CD]-isoGlu-R-C()-GTAT-C()-ATERLAHFLELSSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSNH₂-Hyp-NH₂ (Compd. 8)
- [19CD]-isoGlu-R-C()-GTAT-C()-ATERLAHFLTRSSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSSNH₂-Hyp-NH₂ (Compd. 9)
- 5 [19CD]-isoGlu-R-C()-GTAT-C()-ATERLAHFLTRSSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSTHyp-NH₂ (Compd. 10)
- [19CD]-isoGlu-R-C()-GTAT-C()-ATERLAHFLTRSSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSO-T-Hyp-NH₂ (Compd. 11)
- [19CD]-isoGlu-R-C()-GTAT-C()-ATERLAHFLTRSSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSA-T-Hyp-NH₂ (Compd. 12)
- 10 [19CD]-isoGlu-R-C()-GTAT-C()-ATERLAHFLTRSSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSGT-Hyp-NH₂ (Compd. 13)
- [19CD]-isoGlu-R-C()-NTAT-C()-ATQRLAHFLHRF-Gly(Me)-A-Ile(Me)-LSSTEVGSETP-NH₂ (Compd. 14)
- 15 [19CD]-isoGlu-R-C()-GTAT-C()-ATERLAHFLHRSSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSNTP-NH₂ (Compd. 15)
- [19CD]-isoGlu-R-C()-NTAT-C()-ATQRLAHFLHRSSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSNTP-NH₂ (Compd. 16).
- [19CD]-isoGlu-R-C()-GTAT-C()-ATERLANFLVRSSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSNHyp-NH₂ (Compd. 17)
- 20 [19CD]-isoGlu-R-C()-GTAT-C()-ATERLAHFLTRSSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSNHyp-NH₂ (Compd. 18)
- [19CD]-isoGlu-R-C()-GTAT-C()-ATERLAHFLKRSSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSNHyp-NH₂ (Compd. 19)
- 25 [19CD]-isoGlu-R-C()-NTAT-C()-ATQRLAHFLHRVSF-Gly(Me)-A-Ile(Me)-LSSTEVGSETP-NH₂ (Compd. 20)
- [19CD]-isoGlu-R-C()-NTAT-C()-ATQRLAHFLHRVSF-Gly(Me)-A-Ile(Me)-LSSTEVGSETP-NH₂ (Compd. 21)

- [19CD]-isoGlu-R-C()-GTAT-C()-ATERLAHFLVRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSETP-NH₂ (Compd. 22)
- [19CD]-isoGlu-R-C()-PTAT-C()-ATDRLAHFLVRSSF-Gly(Me)-A-Ile(Me)-LSSTDVGSDTP-NH₂ (Compd. 23)
- 5 [19CD]-isoGlu-R-C()-NTAT-C()-ATQRLAHFLVRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSETP-NH₂ (Compd. 24)
- [19CD]-isoGlu-R-C()-PTAT-C()-ATDRLAHFLHRSSF-Gly(Me)-A-Ile(Me)-LSSTDVGSNTP-NH₂ (Compd. 25)
- [19CD]-isoGlu-R-C()-GTAT-C()-ATERLAHFLHRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSETP-NH₂ (Compd. 26)
- 10 [19CD]-isoGlu-R-C()-PTAT-C()-ATDRLAHFLHRSSF-Gly(Me)-A-Ile(Me)-LSSTDVGSDTP-NH₂ (Compd. 27)
- [19CD]-isoGlu-R-C()-GTAT-C()-ATERLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSSST-Hyp-NH₂ (Compd. 28)
- 15 [19CD]-isoGlu-R-C()-GTAT-C()-ATERLAHFLNRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSSST-Hyp-NH₂ (Compd. 29)
- [19CD]-isoGlu-R-C()-GTAT-C()-ATERLAHFLHRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSSST-Hyp-NH₂ (Compd. 30)
- [19CD]-isoGlu-R-C()-GTAT-C()-ATERLAHFLVRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSSST-Hyp-NH₂ (Compd. 31)
- 20 [19CD]-isoGlu-R-C()-GTAT-C()-ATERLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSSFT-Hyp-NH₂ (Compd. 32)
- [19CD]-isoGlu-R-C()-GTAT-C()-ATERLAHFLNRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSSFT-Hyp-NH₂ (Compd. 33)
- 25 [19CD]-isoGlu-R-C()-GTAT-C()-ATERLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSSTP-NH₂ (Compd. 34)
- [19CD]-isoGlu-R-C()-GTAT-C()-ATERLAHFLNRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSSTP-NH₂ (Compd. 35)

- [19CD]-isoGlu-R-C()-GTAT-C()-ATERLAHFLELSSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSSSTP-NH₂ (Compd. 36)
- [19CD]-isoGlu-R-C()-GTAT-C()-ATERLAHFLHRSSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSSSTP-NH₂ (Compd. 37)
- 5 [19CD]-isoGlu-R-C()-GTAT-C()-ATERLAHFLVRSSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSSSTP-NH₂ (Compd. 38)
- [19CD]-isoGlu-R-C()-GTAT-C()-ATERLAHFLQRSSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSSFTP-NH₂ (Compd. 39)
- [19CD]-isoGlu-R-C()-GTAT-C()-ATERLAHFLNRSSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSSFTP-NH₂ (Compd. 40)
- 10 [19CD]-isoGlu-R-C()-GTAT-C()-ATERLADFLTRSSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSSST-Hyp-NH₂ (Compd. 41)
- [19CD]-isoGlu-R-C()-GTAT-C()-ATERLADFLTRSSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSSFT-Hyp-NH₂ (Compd. 42)
- 15 [19CD]-isoGlu-R-C()-GTAT-C()-ATERLA-Aad-FLTRSSSF-Gly(Me)-A-Ile(Me)-LSSTE-VGSST-Hyp-NH₂ (Compd. 43)
- [19CD]-isoGlu-R-C()-GTAT-C()-ATERLA-Aad-FLTRSSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSS-Orn-T-Hyp-NH₂ (Compd. 44)
- [19CD]-isoGlu-R-C()-GTAT-C()-ATERLAHFLTRSSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSSST-Apr-NH₂ (Compd. 45)
- 20 [19CD]-isoGlu-R-C()-GTAT-C()-ATERLAHFLTRSSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSSSTP-NH₂ (Compd. 46)
- [19CD]-isoGlu-R-C()-GTAT-C()-ATERLAHFLTRSSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSS-Orn-TP-NH₂ (Compd. 47)
- 25 [19CD]-isoGlu-R-C()-GTAT-C()-ATERLAHFLTRSSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSTT-Hyp-NH₂ (Compd. 48)
- [19CD]-isoGlu-R-C()-NTAT-C()-ATQRLADFLQRSSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSSNT-Hyp-NH₂ (Compd. 49)

- [19CD]-isoGlu-R-C()-NTAT-C()-ATQRLAHFL-Aad-RSSF-Gly(Me)-A-Ile(Me)-LSSTE-VGSNT-Hyp-NH₂ (Compd. 50)
- [19CD]-isoGlu-R-C()-GTAT-C()-ATERLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTE-VGS-Orn-T-Hyp-NH₂ (Compd. 51)
- 5 [19CD]-isoGlu-R-C()-GTAT-C()-ATERLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTE-VGS-GT-Hyp-NH₂ (Compd. 52)
- [19CD]-isoGlu-R-C()-GTAT-C()-ATERLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTE-VGS-Aad-T-Hyp-NH₂ (Compd. 53)
- [19CD]-isoGlu-R-C()-GTAT-C()-ATERLADFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTE-VGSNT-Hyp-NH₂ (Compd. 54)
- 10 [19CD]-isoGlu-R-C()-GTAT-C()-ATERLA-Aad-FLQRSSF-Gly(Me)-A-Ile(Me)-LSSTE-VGSNT-Hyp-NH₂ (Compd. 55)
- [19CD]-isoGlu-R-C()-NTAT-C()-ATERLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTE-VGSNT-Hyp-NH₂ (Compd. 56)
- 15 [19CD]-isoGlu-R-C()-QTAT-C()-ATERLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTE-VGSNT-Hyp-NH₂ (Compd. 57)
- [19CD]-isoGlu-R-C()-NTAT-C()-ATQRLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTE-VGSNT-Hyp-NH₂ (Compd. 58)
- [19CD]-isoGlu-R-C()-PTAT-C()-ATDRLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTD-VGSNT-Hyp-NH₂ (Compd. 59)
- 20 [19CD]-isoGlu-R-C()-NTAT-C()-ATERLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTE-VGS-DT-Hyp-NH₂ (Compd. 60)
- [19CD]-isoGlu-R-C()-NTAT-C()-ATQRLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTE-VGS-NTP-NH₂ (Compd. 61)
- 25 [19CD]-isoGlu-R-C()-NTAT-C()-ATERLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTE-VGS-NTP-NH₂ (Compd. 62)
- [19CD]-isoGlu-R-C()-PTAT-C()-ATERLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTE-VGSNT-Hyp-NH₂ (Compd. 63)

- [19CD]-isoGlu-R-C()-PTAT-C()-ATQRLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSNTHyp-NH₂ (Compd. 64)
- [19CD]-isoGlu-R-C()-NTAT-C()-ATQRLADFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSNTPNH₂ (Compd. 65)
- 5 [19CD]-isoGlu-R-C()-NTAT-C()-ATERLADFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSNTPNH₂ (Compd. 66)
- [19CD]-isoGlu-R-C()-PTAT-C()-ATERLADFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSNTPNH₂ (Compd. 67)
- [19CD]-isoGlu-R-C()-NTAT-C()-ATERLADFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSNTHyp-NH₂ (Compd. 68)
- 10 [19CD]-isoGlu-R-C()-NTAT-C()-ATQRLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSSTHyp-NH₂ (Compd. 69)
- [19CD]-isoGlu-R-C()-PTAT-C()-ATERLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSSTHyp-NH₂ (Compd. 70)
- 15 [19CD]-isoGlu-R-C()-NTAT-C()-ATQRLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSGTHyp-NH₂ (Compd. 71)
- [19CD]-isoGlu-R-C()-PTAT-C()-ATERLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSGTHyp-NH₂ (Compd. 72)
- [19CD]-isoGlu-R-C()-NTAT-C()-ATQRLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSTTHyp-NH₂ (Compd. 73)
- 20 [19CD]-isoGlu-R-C()-PTAT-C()-ATERLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSTTHyp-NH₂ (Compd. 74)
- [19CD]-isoGlu-R-C()-NTAT-C()-ATQRLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSAad-T-Hyp-NH₂ (Compd. 75)
- 25 [19CD]-isoGlu-R-C()-PTAT-C()-ATERLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSAad-T-Hyp-NH₂ (Compd. 76)
- [19CD]-isoGlu-R-C()-GTAT-C()-ATERLAHFLQRF-Gly(Me)-A-Ile(Me)-LSSTEVGSNTHyp-NH₂ (Compd. 77)

- [19CD]-isoGlu-R-C()-GTAT-C()-ATERLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSNTP-NH₂ (Compd. 78)
- [19CD]-isoGlu-R-C()-NTAT-C()-ATQRLAHFLQRF-Gly(Me)-A-Ile(Me)-LSSTEVGSGT-Hyp-NH₂ (Compd. 79)
- 5 [19CD]-isoGlu-R-C()-NTAT-C()-ATQRLADFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSGTP-NH₂ (Compd. 80)
- [19CD]-isoGlu-R-C()-NTAT-C()-ATQRLADFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSGT-Hyp-NH₂ (Compd. 81)
- [19CD]-isoGlu-R-C()-NTAT-C()-ATQRLADFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSTTP-NH₂ (Compd. 82)
- 10 [19CD]-isoGlu-R-C()-NTAT-C()-ATQRLADFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSTT-Hyp-NH₂ (Compd. 83)
- [19CD]-isoGlu-R-C()-NTAT-C()-ATQRLAHFLERSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSSTP-NH₂ (Compd. 84)
- 15 [19CD]-isoGlu-R-C()-NTAT-C()-ATQRLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSSTP-NH₂ (Compd. 85)
- [19CD]-isoGlu-R-C()-NTAT-C()-ATQRLA-Aad-FLQRSSF-Gly(Me)-A-Ile(Me)-LSSTE-VGSNT-Hyp-NH₂ (Compd. 86)
- [19CD]-isoGlu-R-C()-NTAT-C()-ATQRLA-Aad-FLQRSSF-Gly(Me)-A-Ile(Me)-LSSTE-VGSST-Hyp-NH₂ (Compd. 87)
- 20 [19CD]-isoGlu-R-C()-QTAT-C()-ATDRLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTE-VGSNT-Hyp-NH₂ (Compd. 88)
- [19CD]-isoGlu-R-C()-QTAT-C()-ATDRLA-Aad-FLQRSSF-Gly(Me)-A-Ile(Me)-LSSTE-VGSNT-Hyp-NH₂ (Compd. 89)
- 25 [19CD]-isoGlu-R-C()-QTAT-C()-ATERLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSGT-Hyp-NH₂ (Compd. 90)
- [19CD]-isoGlu-R-C()-QTAT-C()-ATDRLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSGT-Hyp-NH₂ (Compd. 91)

[19CD]-isoGlu-R-C()-QTAT-C()-ATDRLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSNTP-NH₂ (Compd. 92)

[19CD]-isoGlu-R-C()-QTAT-C()-ATERLA-Aad-FLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSNTP-NH₂ (Compd. 93)

5 [19CD]-isoGlu-R-C()-QTAT-C()-ATERLA-Aad-FLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSGTP-NH₂ (Compd. 94)


[19CD]-isoGlu-R-C()-QTAT-C()-ATDRLA-Aad-FLQRF-Gly(Me)-A-Ile(Me)-LSSTEVGSNTP-Hyp-NH₂ (Compd. 95)

10 [19CD]-isoGlu-R-C()-QTAT-C()-ATERLAHFLQRF-Gly(Me)-A-Ile(Me)-LSSTEVGSNTP-NH₂ (Compd. 96)

wherein [19CD] represents [19-carboxynonadecanoyl];

and pharmaceutically acceptable salts and solvates thereof.

15 In certain examples, the amylin analogue of the invention is not a compound having the formula:



wherein

R¹ is hydrogen, C₁₋₄ acyl, benzoyl or C₁₋₄ alkyl, or a half-life extending moiety M, wherein M is optionally linked to Z via a linker moiety L;

20 R² is OH or NHR³, wherein R³ is hydrogen or C₁₋₃-alkyl; and

Z is an amino acid sequence selected from the group consisting of:

RCNTATCATQRLADFLHRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSETP

RCNTATCATQRLAHFLHRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSETP

RCNTATCATQRLANFLHRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSETP

25 RCNTATCATQRLADFLHRSSNNF-Gly(Me)-A-Ile(Me)-LSSTNVGSNT-Apr

RCNTATCATQRLAHFLHRSSNNF-Gly(Me)-A-Ile(Me)-LSSTNVGSNT-Apr

RCGTATCATERLAHFLNRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSNTP-Hyp

RCGTATCATERLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSNTP-Hyp

RCGTATCATERLAHFLESSF-Gly(Me)-A-Ile(Me)-LSSTEVGSNTP-Hyp

- RCGTATCATERLAHFLTRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSSST-Hyp
 RCGTATCATERLAHFLTRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSTFT-Hyp
 RCGTATCATERLAHFLTRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSS-Orn-T-Hyp
 RCGTATCATERLAHFLTRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSS-Aad-T-Hyp
 5 RCGTATCATERLAHFLTRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSGT-Hyp
 RCNTATCATQRLAHFLHRF-Gly(Me)-A-Ile(Me)-LSSTEVGSETP
 RCGTATCATERLAHFLHRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSNTP
 and
 RCNTATCATQRLAHFLHRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSNTP
 10 or a pharmaceutically acceptable salt or solvate thereof.

For example, in such examples, the compound is not:

- [19CD]-isoGlu-RCNTATCATQRLADFLHRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSETP-NH₂
 [19CD]-isoGlu-RCNTATCATQRLAHFLHRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSETP-NH₂
 15 [19CD]-isoGlu-RCNTATCATQRLANFLHRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSETP-NH₂
 [19CD]-isoGlu-RCNTATCATQRLADFLHRSSNNF-Gly(Me)-A-Ile(Me)-LSSTNVGSNT-Apr-NH₂
 [19CD]-isoGlu-RCNTATCATQRLAHFLHRSSNNF-Gly(Me)-A-Ile(Me)-LSSTNVGSNT-Apr-NH₂
 20 [19CD]-isoGlu-RCGTATCATERLAHFLNRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSNTP-Hyp-NH₂
 [19CD]-isoGlu-RCGTATCATERLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSNTP-Hyp-NH₂
 [19CD]-isoGlu-RCGTATCATERLAHFLERSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSNTP-Hyp-NH₂
 25 [19CD]-isoGlu-RCGTATCATERLAHFLTRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSSST-Hyp-NH₂
 [19CD]-isoGlu-RCGTATCATERLAHFLTRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSTFT-Hyp-NH₂
 [19CD]-isoGlu-RCGTATCATERLAHFLTRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSS-Orn-T-Hyp-NH₂
 30 [19CD]-isoGlu-RCGTATCATERLAHFLTRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSS-Aad-T-Hyp-NH₂

[19CD]-isoGlu-RCGTATCATERLAHFLTRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSGT-Hyp-NH₂

[19CD]-isoGlu-RCNTATCATQRLAHFLHRF-Gly(Me)-A-Ile(Me)-LSSTEVGSETP-NH₂

[19CD]-isoGlu-RCGTATCATERLAHFLHRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSNTP-NH₂

or

- 5 [19CD]-isoGlu-RCNTATCATQRLAHFLHRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSNTP-NH₂;

(wherein [19CD] represents [19-carboxynonadecanoyl])

and pharmaceutically acceptable salts and solvates thereof;

or any such compounds comprising an intramolecular disulphide bridge formed between the cysteine residues present at positions 2 and 7 of the amino acid sequence.

- 10 In another aspect, the invention further provides a composition comprising an amylin analogue as described above. The composition may be a pharmaceutical composition, and may comprise a pharmaceutically acceptable carrier, excipient or vehicle.

- In another aspect, there is provided a device comprising an amylin analogue as described herein, for delivery of the amylin analogue to a subject, optionally further comprising an anti-diabetic agent, an anti-obesity agent, an agent for treatment of metabolic syndrome, an anti-dyslipidemia agent, an anti-hypertensive agent, a proton pump inhibitor, or an anti-inflammatory agent.
- 15

- The disclosure further provides a method for the synthesis of an amylin analogue as described above. The method may comprise the steps of synthesising the peptide by solid-phase or liquid-phase methodology, and optionally isolating and/or purifying the final product. The method may further comprise the step of forming a disulphide bond between the thiol groups of the cysteine side chains at positions 2 and 7.
- 20

Also described herein is an amylin analogue as disclosed herein for use in a method of medical treatment.

- 25 The amylin analogues are useful, inter alia, in the reduction of food intake, promotion of weight loss, and inhibition or reduction of weight gain. As a result, they may be used for treatment of a variety of conditions, diseases, or disorders in a subject, including, but not limited to, obesity and various obesity-related conditions, diseases, or disorders, such as diabetes (e.g. type 2 diabetes), hypertension, dyslipidemia, sleep apnea and cardiovascular disease. The subject may be affected by obesity accompanied by at least one weight-related co-morbid condition, such as diabetes (e.g. type 2 diabetes), hypertension, dyslipidemia, sleep apnea and cardiovascular disease. It will be understood that the amylin analogues may thus be administered to subjects affected by conditions characterised by inadequate control of appetite or otherwise over-feeding, such as binge-
- 30

eating disorder and Prader-Willi syndrome. It will be clear that the analogues can be used for treatment of combinations of the conditions described.

Thus, the disclosure provides an amylin analogue of the invention for use in a method of treating, inhibiting or reducing weight gain, promoting weight loss and/or reducing excess
5 body weight. Treatment may be achieved, for example, by control of appetite, feeding, food intake, calorie intake and/or energy expenditure.

The disclosure also provides an amylin analogue of the invention for use in a method of treating obesity as well as associated diseases, disorders and health conditions, including, but not limited to, morbid obesity, obesity prior to surgery, obesity-linked inflammation,
10 obesity-linked gallbladder disease and obesity-induced sleep apnea and respiratory problems, degeneration of cartilage, osteoarthritis, and reproductive health complications of obesity or overweight such as infertility. The subject may be affected by obesity accompanied by at least one weight-related co-morbid condition, such as diabetes (e.g. type 2 diabetes), hypertension, dyslipidemia, sleep apnea and cardiovascular disease.

15 The disclosure also provides an amylin analogue of the invention for use in a method of prevention or treatment of Alzheimer's disease, diabetes, type 1 diabetes, type 2 diabetes, pre-diabetes, insulin resistance syndrome, impaired glucose tolerance (IGT), disease states associated with elevated blood glucose levels, metabolic disease including metabolic syndrome, hyperglycemia, hypertension, atherogenic dyslipidemia, hepatic
20 steatosis ("fatty liver"; including non-alcoholic fatty liver disease (NAFLD), which itself includes non-alcoholic steatohepatitis (NASH)), kidney failure, arteriosclerosis (e.g. atherosclerosis), macrovascular disease, microvascular disease, diabetic heart (including diabetic cardiomyopathy and heart failure as a diabetic complication) coronary heart disease, peripheral artery disease or stroke, and combinations thereof.

25 The disclosure also provides an amylin analogue of the invention for use in a method of lowering circulating LDL levels and/or increasing HDL/LDL ratio.

Effects of amylin analogues on these conditions may be mediated in whole or in part via an effect on body weight, or may be independent thereof.

30 Also described herein is the use of an amylin analogue of the invention in the manufacture of a medicament for treating, inhibiting or reducing weight gain, promoting weight loss and/or reducing excess body weight.

In another aspect, the invention also provides use of an amylin analogue of the invention in the manufacture of a medicament for treating obesity as well as associated diseases, disorders and health conditions, including, but not limited to, morbid obesity, obesity prior to surgery, obesity-linked inflammation, obesity-linked gallbladder disease and obesity-induced sleep apnea and respiratory problems, degeneration of cartilage, osteoarthritis, and reproductive health complications of obesity or overweight such as infertility. The subject may be affected by obesity accompanied by at least one weight-related co-morbid condition, such as diabetes (e.g. type 2 diabetes), hypertension, dyslipidemia, sleep apnea and cardiovascular disease.

10 In another aspect, the invention also provides use of an amylin analogue of the invention in the manufacture of a medicament for the prevention or treatment of Alzheimer's disease, diabetes, type 1 diabetes, type 2 diabetes, pre-diabetes, insulin resistance syndrome, impaired glucose tolerance (IGT), disease states associated with elevated blood glucose levels, metabolic disease including metabolic syndrome, hyperglycemia, hypertension, atherogenic dyslipidemia, hepatic steatosis ("fatty liver"; including non-alcoholic fatty liver disease (NAFLD), which itself includes non-alcoholic steatohepatitis (NASH)), kidney failure, arteriosclerosis (e.g. atherosclerosis), macrovascular disease, microvascular disease, diabetic heart (including diabetic cardiomyopathy and heart failure as a diabetic complication) coronary heart disease, peripheral artery disease or stroke, and combinations thereof.

In another aspect, the invention also provides use of an amylin analogue of the invention in the manufacture of a medicament for lowering circulating LDL levels and/or increasing HDL/LDL ratio.

25 The disclosure further provides a method of treating, inhibiting or reducing weight gain, promoting weight loss and/or reducing excess body weight in a subject, comprising administering a therapeutically effective amount of an amylin analogue of the invention to the subject.

30 The disclosure also provides a method of treating obesity as well as associated diseases, disorders and health conditions, including, but not limited to, morbid obesity, obesity prior to surgery, obesity-linked inflammation, obesity-linked gallbladder disease and obesity-induced sleep apnea and respiratory problems, degeneration of cartilage, osteoarthritis, and reproductive health complications of obesity or overweight such as infertility in a subject, comprising administering a therapeutically effective amount of an amylin analogue of the invention to the subject. The subject may be affected by obesity

accompanied by at least one weight-related co-morbid condition, such as diabetes (e.g. type 2 diabetes), hypertension, dyslipidemia, sleep apnea and cardiovascular disease.

The disclosure also provides a method of prevention or treatment of Alzheimer's disease, diabetes, type 1 diabetes, type 2 diabetes, pre-diabetes, insulin resistance syndrome,
5 impaired glucose tolerance (IGT), disease states associated with elevated blood glucose levels, metabolic disease including metabolic syndrome, hyperglycemia, hypertension, atherogenic dyslipidemia, hepatic steatosis ("fatty liver"; including non-alcoholic fatty liver disease (NAFLD), which itself includes non-alcoholic steatohepatitis (NASH)), kidney failure, arteriosclerosis (e.g. atherosclerosis), macrovascular disease, microvascular
10 disease, diabetic heart (including diabetic cardiomyopathy and heart failure as a diabetic complication) coronary heart disease, peripheral artery disease or stroke, and combinations thereof, in a subject, comprising administering a therapeutically effective amount of an amylin analogue of the invention to the subject.

The disclosure further provides a method of lowering circulating LDL levels and/or
15 increasing HDL/LDL ratio in a subject, comprising administering a therapeutically effective amount of an amylin analogue of the invention to the subject.

The disclosure further provides the use of an amylin analogue as described above in a method of cosmetic (i.e. non-therapeutic) weight loss. It will be understood that references to therapeutic uses of amylin analogues and methods comprising
20 administration of amylin analogues may equally be taken to encompass uses and administration of such compositions.

Further aspects and embodiments of the present invention will become apparent from the disclosure below.

DETAILED DESCRIPTION OF THE INVENTION

25 Unless otherwise defined herein, scientific and technical terms used herein shall have the meanings that are commonly understood by those of ordinary skill in the art. Generally, nomenclature employed herein in connection with techniques of chemistry, molecular biology, cell and cancer biology, immunology, microbiology, pharmacology, and protein and nucleic acid chemistry, described herein, is that well known and commonly used in
30 the art.

All publications, patents and published patent applications referred to in this application are specifically incorporated by reference herein. In case of conflict, the present specification, including its specific definitions, will control.

Throughout this specification, the word “comprise” or variations such as “comprises” or “comprising” will be understood to imply the inclusion of a stated integer or component, or of a stated group of integers or components, but not the exclusion of any other integer or component or group of integers or components.

- 5 The singular forms “a,” “an,” and “the” include the plurals unless the context clearly dictates otherwise.

The term “including” is used to mean “including but not limited to.” “Including” and “including but not limited to” are used interchangeably.

- 10 The terms “patient,” “subject,” and “individual” may be used interchangeably and may refer to either a human or a non-human animal. Subjects are typically mammals, including humans, non-human primates (including great apes, Old World monkeys and New World monkeys), livestock animals (e.g., bovines, porcines), companion animals (e.g., canines, felines) and rodents (e.g., mice and rats).

- 15 As used herein, the term “pharmaceutically acceptable salt” is intended to indicate a salt which is not harmful to a patient or subject to which the salt in question is administered. It may suitably be a salt chosen, e.g., among acid addition salts and basic salts. Examples of acid addition salts include chloride salts, citrate salts and acetate salts. Examples of basic salts include salts where the cation is selected among alkali metal cations, such as sodium or potassium ions, alkaline earth metal cations, such as calcium or magnesium
20 ions, as well as substituted ammonium ions, such as ions of the type $N(R^1)(R^2)(R^3)(R^4)^+$, where R^1 , R^2 , R^3 and R^4 independently will typically designate hydrogen, optionally substituted C_{1-6} -alkyl or optionally substituted C_{2-6} -alkenyl. Examples of relevant C_{1-6} -alkyl groups include methyl, ethyl, 1-propyl and 2-propyl groups. Examples of C_{2-6} -alkenyl groups of possible relevance include ethenyl, 1-propenyl and 2-propenyl. Other examples
25 of pharmaceutically acceptable salts are described in “Remington’s Pharmaceutical Sciences”, 17th edition, Alfonso R. Gennaro (Ed.), Mark Publishing Company, Easton, PA, USA, 1985 (and more recent editions thereof), in the “Encyclopaedia of Pharmaceutical Technology”, 3rd edition, James Swarbrick (Ed.), Informa Healthcare USA (Inc.), NY, USA, 2007, and in *J. Pharm. Sci.* 66: 2 (1977).

- 30 The term “solvate” in the context of the present invention refers to a complex of defined stoichiometry formed between a solute (*in casu*, a peptide or pharmaceutically acceptable salt thereof according to the invention) and a solvent. The solvent in this connection may, for example, be water, ethanol or another pharmaceutically acceptable - typically small-

molecular - organic species, such as, but not limited to, acetic acid or lactic acid. When the solvent in question is water, such a solvate is normally referred to as a hydrate.

The term "agonist" as employed in the context of the invention refers to a substance that activates the receptor type in question, typically by binding to it (i.e. as a ligand).

- 5 Each embodiment of the invention described herein may be taken alone or in combination with one or more other embodiments of the invention.

Throughout the present specification, unless naturally occurring amino acids are referred to by their full name (e.g. alanine, arginine, etc.), they are designated by their conventional three-letter or single-letter abbreviations (e.g. Ala or A for alanine, Arg or R for arginine, etc.). In the case of certain less common or non-naturally occurring amino acids (i.e. amino acids other than the 20 encoded by the standard mammalian genetic code), unless they are referred to by their full name (e.g. sarcosine, ornithine, etc.), frequently employed three- or four-character codes are employed for residues thereof, including Orn (ornithine, i.e. 2,5-diaminopentanoic acid), Aib (α -aminoisobutyric acid), Dab (2,4-diaminobutanoic acid), Dap (2,3-diaminopropanoic acid), Har (homoarginine), γ -Glu (γ -glutamic acid), Gaba (γ -aminobutanoic acid), β -Ala (i.e. 3-aminopropanoic acid), 8Ado (8-amino-3,6-dioxaoctanoic acid).

Unless otherwise indicated, reference is made to the L-isomeric forms of the amino acids in question.

- 20 Additional abbreviations include the following:

Gly(Me): N-methylglycine [also known as sarcosine (Sar)]

Ile(Me): N-methylisoleucine

Aad: 2-aminoadipic acid, e.g. (2S)-2-aminoadipic acid [also (2S)-2-aminohexanedioic acid] , also known as homo-glutamic acid

- 25 Apr: 4-aminoproline, e.g. (2S,4R)-4-aminoproline

[also denoted (4R)-4-amino-L-proline]

Hyp: 4-hydroxyproline, e.g. (2S,4R)-4-hydroxyproline

[also denoted (4R)-4-hydroxy-L-proline]

The term "therapeutically effective amount" as used herein in the context of the above-described methods of treatment or other therapeutic interventions according to the invention refers to an amount that is sufficient to cure, ameliorate, alleviate or partially arrest the clinical manifestations of the particular disease, disorder or condition that is the object of the treatment or other therapeutic intervention in question e.g. as measured by established clinical endpoints or other biomarkers (established or experimental). A therapeutically relevant amount may be determined empirically by one skilled in the art based on the indication being treated or prevented and the subject to whom the therapeutically relevant amount is being administered. For example, the skilled worker may measure one or more of the clinically relevant indicators of bioactivity described herein, e.g. plasma lipid levels, blood glucose levels or insulin release. The skilled worker may determine a clinically relevant amount through *in vitro* or *in vivo* measurements. Other exemplary measures include weight gain, weight loss, and change in blood pressure.

An amount adequate to accomplish any or all of these effects is defined as a therapeutically effective amount. The administered amount and the method of administration can be tailored to achieve optimal efficacy. An amount effective for a given purpose will depend, *inter alia*, on the severity of the disease, disorder or condition that is the object of the particular treatment or other therapeutic intervention, on the body weight and general condition of the subject in question, on diet, on possible concurrent medication, and on other factors well known to those skilled in the medical arts. Determination of an appropriate dosage size and dosing regimen most appropriate for administration of a peptide or pharmaceutically acceptable salt or solvate thereof according to the invention to a human may be guided by the results obtained by the present invention, and may be confirmed in properly designed clinical trials. An effective dosage and treatment protocol may be determined by conventional means, starting with a low dose in laboratory animals and then increasing the dosage while monitoring the effects, and systematically varying the dosage regimen as well. Numerous factors may be taken into consideration by a clinician when determining an optimal dosage for a given subject. Such considerations are well known to the skilled person.

The terms "treatment" and grammatical variants thereof (e.g. "treated", "treating", "treat") as employed in the present context refer to an approach for obtaining beneficial or desired clinical results. For the purposes of this invention, beneficial or desired clinical results include, but are not limited to, alleviation of symptoms, diminishment of extent of disease, stabilization (i.e. not worsening) of state of disease, delay or slowing of disease progression, amelioration or palliation of the disease state, and remission (whether partial

or total), whether detectable or undetectable. "Treatment" can also mean prolonging survival relative to expected survival time if not receiving treatment. A subject (e.g. a human) in need of treatment may thus be a subject already afflicted with the disease or disorder in question. The term "treatment" includes inhibition or reduction of an increase in severity of a pathological state or symptoms (e.g. weight gain or hyperglycemia) relative to the absence of treatment, and is not necessarily meant to imply complete cessation of the relevant disease, disorder or condition.

The terms "prevention" and grammatical variants thereof (e.g., "prevented", "preventing", "prevent") as employed in the present context refer to an approach for hindering or preventing the development of, or altering the pathology of, a condition, disease or disorder. Accordingly, "prevention" may refer to prophylactic or preventive measures. For the purposes of this invention, beneficial or desired clinical results include, but are not limited to, prevention or slowing of symptoms, progression or development of a disease, whether detectable or undetectable. A subject (e.g. a human) in need of "prevention" may thus be a subject not yet afflicted with the disease or disorder in question. The term "prevention" thus includes inhibiting or slowing the onset of disease relative to the absence of treatment, and is not necessarily meant to imply permanent prevention of the relevant disease, disorder or condition.

Synthesis of amylin analogues

The invention further provides a method of synthesis of an amylin analogue of the invention. The amylin analogues (which may also be referred to as compounds or peptides) may suitably be manufactured by standard synthetic methods. Thus, the peptides may be synthesized by, e.g., methods comprising synthesizing the peptide by standard solid-phase or liquid-phase methodology, either stepwise or by fragment assembly, and optionally isolating and purifying the final peptide product. In this context, reference may be made to WO 98/11125 or, *inter alia*, Fields, G.B. et al., "Principles and Practice of Solid-Phase Peptide Synthesis"; in: *Synthetic Peptides*, Gregory A. Grant (ed.), Oxford University Press (2nd edition, 2002) and the synthesis examples herein. The method may further comprise the step of forming a disulfide bond between the thiol groups of the cysteine side chains at positions 2 and 7, e.g. by oxidative cyclisation. In the case of solid phase synthesis, cyclisation may be performed in situ on the solid phase (e.g. resin), i.e. before removal of the peptide from the solid phase.

C₁₋₄ acyl groups that may be present as a group R¹ in the context of compounds of the present invention include formyl (i.e. methanoyl), acetyl (i.e. ethanoyl), propanoyl, 1-butanoyl and 2-methylpropanoyl groups.

5

C₁₋₄ alkyl groups

C₁₋₄ alkyl groups that may be present as a group R¹ in the context of compounds of the present invention include, but are not limited to, C₁₋₃ alkyl groups, such as methyl, ethyl, 1-propyl or 2-propyl.

10

C₁₋₃ alkyl groups

C₁₋₃ alkyl groups that may be present as a group R³ in the context of compounds of the present invention include methyl, ethyl, 1-propyl and 2-propyl.

15 Half-life extending moieties M

As described herein, the N-terminal moiety R¹ in a compound of the invention may be a half-life extending moiety (sometimes referred to in the literature as, *inter alia*, a duration enhancing moiety or albumin binding moiety), optionally linked (covalently attached) to the peptide moiety Z via a linker moiety L. Among suitable half-life extending moieties are certain types of lipophilic substituents. Without wishing to be bound by any particular theory, it is thought that such lipophilic substituents (and other classes of half-life extending moieties) bind albumin in the blood stream, thereby shielding the compound of the invention from renal filtration as well as enzymatic degradation and thus possibly enhancing the half-life of the compound *in vivo*. The lipophilic substituent may also modulate the potency of the compound as an agonist to the amylin (calcitonin) receptor.

20

25

The lipophilic substituent may be attached to the N-terminal amino acid residue or to the linker L via an ester, a sulfonyl ester, a thioester, an amide, an amine or a sulfonamide. Accordingly it will be understood that preferably the lipophilic substituent includes an acyl group, a sulfonyl group, an N atom, an O atom or an S atom which forms part of the ester, sulfonyl ester, thioester, amide, amine or sulfonamide. Preferably, an acyl group in the lipophilic substituent forms part of an amide or ester with the amino acid residue or the linker.

30

35

The lipophilic substituent may comprise a hydrocarbon chain having from 10 to 24 C atoms, e.g. from 14 to 22 C atoms, e.g. from 16 to 20 C atoms. Preferably it has at least 14 C atoms, and preferably has 20 C atoms or fewer. For example, the hydrocarbon

chain may contain 14, 15, 16, 17, 18, 19 or 20 carbon atoms. The hydrocarbon chain may be linear or branched, and may be saturated or unsaturated. Furthermore, it can include a functional group at the end of the lipophilic chain, e.g. a carboxylic acid group which may or may not be protected during synthesis. From the discussion above it will be understood that the hydrocarbon chain is preferably substituted with a moiety which forms part of the attachment to the N-terminal amino acid residue of the peptide moiety Z or to the linker L, for example an acyl group, a sulfonyl group, an N atom, an O atom or an S atom.

Most preferably, the hydrocarbon chain is substituted with an acyl group, and accordingly the hydrocarbon chain may be part of an alkanoyl group, for example a dodecanoyl, 2-butyloctanoyl, tetradecanoyl, hexadecanoyl, heptadecanoyl, octadecanoyl, nonadecanoyl or eicosanoyl group. Examples of functionalized hydrocarbon chains are 15-carboxypentadecanoyl, 17-carboxyheptadecanoyl and 19-carboxynonadecanoyl.

As mentioned above, a lipophilic substituent M may be linked to the N-terminal amino acid residue of Z via a linker L. In embodiments, the linker moiety L may itself comprise one, two, three or more linked sub-moieties L^1 , L^2 , L^3 , ..etc. When the linker L comprises only one such moiety, it is attached to the lipophilic substituent and to the N-terminal amino acid residue of Z. The linker may then be attached to the lipophilic substituent and to the N-terminal amino acid residue of Z independently by means of an ester, a sulfonyl ester, a thioester, an amide, an amine or a sulfonamide bond. Accordingly, it may include two moieties independently selected from acyl, sulfonyl, an N atom, an O atom and an S atom. The linker may consist of a linear or branched C_{1-10} hydrocarbon chain or more preferably a linear C_{1-5} hydrocarbon chain. Furthermore the linker can be substituted with one or more substituents selected from C_{1-6} alkyl, amino C_{1-6} alkyl, hydroxy C_{1-6} alkyl and carboxy C_{1-6} alkyl.


In some embodiments the linker may be, for example, a residue of any naturally occurring or non-naturally occurring amino acid. For example, the linker may be a residue of Gly, Pro, Ala, Val, Leu, Ile, Met, Cys, Phe, Tyr, Trp, His, Lys, Arg, Gln, Asn, α -Glu, γ -Glu, ϵ -Lys, Asp, β -Asp, Ser, Thr, Gaba, Aib, β -Ala (i.e. 3-aminopropanoyl), 4-aminobutanoyl, 5-aminopentanoyl, 6-aminohexanoyl, 7-aminoheptanoyl, 8-aminooctanoyl, 9-aminononanoyl, 10-aminodecanoyl or 8Ado (i.e. 8-amino-3,6-dioxaoctanoyl).

References to γ -Glu, ϵ -Lys, and β -Asp indicate residues of amino acids which participate in bonds via their side chain carboxyl or amine functional groups. Thus γ -Glu, and β -Asp

participate in bonds via their amino and side chain carboxyl groups, while ϵ -Lys participates via its carboxyl and side chain amino groups.

In certain embodiments, the linker is a residue of Glu, γ -Glu, ϵ -Lys, β -Ala, 4-aminobutanoyl, 8-aminooctanoyl or 8Ado. In the context of the present invention, γ -Glu and isoGlu are used interchangeably.


An example of a lipophilic substituent comprising a lipophilic moiety M and linker L is shown in the formula below:



Here, the backbone nitrogen of an Arg residue is covalently attached to a γ -Glu linker (L) via an amide linkage. A 19-carboxy-nonadecanoyl group is covalently attached to the γ -Glu linker via an amide linkage. This combination of lipophilic moiety and linker, attached to an Arg residue, may be referred to by the shorthand notation [19CD]-isoGlu-R, e.g. when shown in formulae of specific compounds.

The skilled person will be well aware of suitable techniques for preparing the compounds employed in the context of the invention. For examples of suitable chemistry, see, e.g., WO98/08871, WO00/55184, WO00/55119, Madsen et al (*J. Med. Chem.* 2007, 50, 6126-32), and Knudsen et al. 2000 (*J. Med Chem.* 43, 1664-1669).


The hydrocarbon chain in a lipophilic substituent may be further substituted. For example, it may be further substituted with up to three substituents selected from NH_2 , OH and COOH. If the hydrocarbon chain is further substituted, it is preferably further substituted with only one substituent. Alternatively or additionally, the hydrocarbon chain may include a cycloalkane or heterocycloalkane moiety, for example as shown below:



In some embodiments, the cycloalkane or heterocycloalkane moiety is a six-membered ring, e.g. a piperidine ring.

- In alternative embodiments of the present invention, the N-terminal amino acid of Z in a compound of the invention may be linked (covalently attached) to a biotinyllic substituent, optionally via a linker moiety L. Without wishing to be bound by any particular theory, it is likewise believed that such biotinyllic substituents bind to albumin in the blood stream, thereby shielding the compound of the invention from enzymatic degradation and thus possibly enhancing the half-life of the compound *in vivo*. A linker, when present, may provide spacing between the peptide moiety Z and the biotinyllic substituent.
- 10 The biotinyllic substituent may be attached to the N-terminal amino acid residue or to the linker via an maleimide ester bond, a sulfonyl ester bond, a thioester bond, an amide bond, an amine bond or a sulfonamide bond. Accordingly it will be understood that the biotinyllic substituent preferably comprises an maleimido group, an acyl group, a sulfonyl group, an N atom, an O atom or an S atom which forms part of the ester, sulfonyl ester, thioester, amide, amine or sulfonamide bond in question.
- 15

Examples of biotinyllic substituents may include



Biotin is known as Vitamin H or Coenzyme R, and is a water-soluble B-complex vitamin (vitamin B7). It has been shown to increase oral uptake of certain drugs.

20 Efficacy of compounds

The compounds of the invention are amylin receptor agonists, i.e. they are capable of binding to, and inducing signalling by, one or more receptors or receptor complexes regarded as physiological receptors for human amylin. These include the human calcitonin receptor hCTR2, as well as complexes comprising the human calcitonin

receptor hCTR2 and at least one of the human receptor activity modifying proteins designated hRAMP1, hRAMP2 and hRAMP3. Complexes between hCTR2 and hRAMP1, hRAMP2 and hRAMP3 are designated hAMYR1, hAMYR2 and hAMYR3 (i.e. human amylin receptors 1, 2 and 3) respectively.

- 5 Without wishing to be bound by theory, a compound may be considered an amylin receptor agonist if it has agonist activity at one or more of hAMYR1, hAMYR2 and hAMYR3, e.g. against hAMYR1 and/or hAMYR3, e.g. at hAMYR3.

Typically an amylin receptor agonist will also have agonist activity at hCTR2 when expressed in the absence of hRAMP1, hRAMP2 and hRAMP3. Typically, the agonist will
10 have activity at hCTR2 (when expressed in the absence of hRAMP1, hRAMP2 and hRAMP3) which is less than 10-fold higher than its activity at any one of hAMYR1, hAMYR2 and hAMYR3 (i.e. its activity at all of these receptors) in a comparable assay. Agonist activity at hCTR2 may be less than 5-fold higher than agonist activity at hAMYR1, hAMYR2 and hAMYR3, substantially equal to (e.g. +/- 10%) agonist activity at hAMYR1,
15 hAMYR2 and hAMYR3, or less than agonist activity at hAMYR1, hAMYR2 and hAMYR3. In this regard, it may be sufficient just to compare activity between hCTR2 and hAMYR3.

The ability to induce cAMP formation (i.e. to induce adenylate cyclase activity) as a result of binding to the relevant receptor or receptor complex is typically regarded as indicative of agonist activity. Other intracellular signaling pathways or events may also be used as
20 read-outs for amylin receptor agonist activity. These may include calcium release, β -arrestin recruitment, receptor internalization, kinase activation or inactivation, lipase activation, inositol phosphate release, diacylglycerol release or nuclear transcription factor translocation.

A suitable comparable assay format would utilize cells which express hCTR2 and which
25 differ only in their expression of hRAMP1, 2 and 3. For example, a "base" cell line which does not express any of hCTR2, hRAMP1, hRAMP2 and hRAMP3 may be engineered to generate cells which express (i) hCTR2, and (ii) one of hAMYR1, hAMYR2 and hAMYR3 (i.e. hCTR2 plus one of hRAMP1, hRAMP2 and hRAMP3), e.g. hAMYR3. The base cells will typically be mammalian cells and may be primate cells. They may be non-human
30 primate cells. Preferably the base cell does not express any of CTR2, RAMP1, RAMP2 or RAMP3 (whether human, or native to the base cell if the base cell is non-human). The base cells may be fibroblast cells. Suitable non-human fibroblast base cells include COS7 cells, from African green monkey, which do not express native CTR2 or RAMPs.

Comparative activity may be measured by any suitable means, such as via determination of EC₅₀ values as described below. It will be apparent that the same biological read-out must be for both receptor types.

Compounds of the present invention may exhibit a number of advantageous properties in relation to human amylin and existing analogues thereof, such as pramlintide, IAPP-GI, and analogues described in WO2012/168430, WO2012/168431 and WO2012/168432. As compared to human amylin or any of those analogues, compounds of the invention may, for example, exhibit improved efficacy (e.g., in the form of improved *in vitro* activity or potency at one or more of the receptors hCTR2, hAMRY1, hAMRy2 or hAMYR3. Additionally or alternatively, compounds of the invention may exhibit improved solubility in aqueous media, especially at pH values in the range from 4 to 7.5, or at a range of pH values across that range. Moreover, compounds of the present invention may additionally or alternatively exhibit reduced tendency to undergo fibrillation in pharmaceutically relevant aqueous media, especially at pH values in the range from 4 to 7, or at a range of pH values across that range. Furthermore, compounds of the present invention may additionally or alternatively exhibit improved chemical stability (i.e. reduced tendency to undergo chemical degradation) in aqueous media, especially at pH values in the range from 4 to 9, or at a range of pH values across that range.

Compounds of the invention may thus be well suited for formulation in acidic media (e.g. pH 4) and in neutral or near-neutral media (e.g. pH 7 or 7.4). In contrast to pramlintide, for example, which generally exhibits poor chemical stability and rapid fibrillation in pharmaceutically relevant aqueous media at neutral pH, compounds of the invention may be thus well suited for co-formulation with, for example, insulin, various insulin analogues and/or other therapeutic (e.g. anti-diabetic or anti-obesity) agents that require a neutral or near-neutral formulation pH.

In general it is preferred to use a biological assay which measures intracellular signalling caused by binding of the compound to the relevant receptor, as discussed above. Activation of the calcitonin/amylin receptor by compounds of the invention (which behave as agonists of the receptor) induces cAMP formation and activation of other intracellular signaling pathways and events. Thus, production of cAMP or any other suitable parameter in suitable cells expressing the receptor can be used to monitor agonist activity towards the receptor.

The skilled person will be aware of suitable assay formats, and examples are provided below. For example, the assays may make use of the human calcitonin receptor (hCTR2) having primary accession number GI: 4502547 (NP_001733.1) or the hAMYR3 receptor

(see Example 2, below). Where sequences of precursor proteins are referred to, it should be understood that assays may make use of the mature protein, lacking the signal sequence.

EC₅₀ values may be used as a numerical measure of agonist potency at a given receptor.

- 5 An EC₅₀ value is a measure of the concentration of a compound required to achieve half of that compound's maximal activity in a particular assay. Thus, for example, a compound having EC₅₀ [hCTR2] lower than the EC₅₀ [hCTR2] of native amylin, or lower than that of pramlintide, in a particular assay may be considered to have higher potency or activity at the receptor than amylin, or higher than that of pramlintide, respectively.

- 10 In some embodiments of compounds of the present invention, the EC₅₀ towards hCTR2 is below 1.4 nM.

In some embodiments of compounds of the present invention, the EC₅₀ towards hCTR2 is below 0.8 nM.

- 15 In some embodiments of compounds of the present invention, the EC₅₀ towards hCTR2 is below 0.4 nM.

In some embodiments of compounds of the present invention, the EC₅₀ towards hCTR2 is below 0.2 nM.

In some embodiments of compounds of the present invention, the EC₅₀ towards hCTR2 is below 0.1 nM.

- 20 An EC₅₀ towards hCTR2 of approximately 0.2 (+/- 10%) or below may be desirable. The EC₅₀ at hCTR2 may be an indication of the effect of a compound on food intake, weight gain and/or weight loss. Compounds with lower EC₅₀ values at hCTR2 may have a greater effect on these parameters.

- 25 In some embodiments of compounds of the present invention, the EC₅₀ towards hAMYR3 is below 1 nM.

In some embodiments of compounds of the present invention, the EC₅₀ towards hAMYR3 is below 0.5 nM.

- 30 In some embodiments of compounds of the present invention, the EC₅₀ towards hAMYR3 is below 0.4 nM.

In some embodiments of compounds of the present invention, the EC_{50} towards hAMYR3 is below 0.3 nM.

In some embodiments of compounds of the present invention, the EC_{50} towards hAMYR3 is below 0.2 nM.

- 5 For example, the EC_{50} at hCTR2 (when expressed in the absence of hRAMP1, hRAMP2 and hRAMP3) may be less than 10-fold lower than the EC_{50} at any or all of hAMYR1, hAMYR2 and hAMYR3, e.g. at hAMYR3.

- The EC_{50} at hCTR2 (when expressed in the absence of hRAMP1, hRAMP2 and hRAMP3) may be less than 5-fold lower than the EC_{50} at any or all of hAMYR1, hAMYR2 and
10 hAMYR3, e.g. at hAMYR3.

The EC_{50} at hCTR2 (when expressed in the absence of hRAMP1, hRAMP2 and hRAMP3) may be substantially equal to (e.g. +/- 50%) the EC_{50} at any or all of hAMYR1, hAMYR2 and hAMYR3, e.g. at hAMYR3.

- The EC_{50} at hCTR2 (when expressed in the absence of hRAMP1, hRAMP2 and hRAMP3) may be higher than the EC_{50} at any or all of hAMYR1, hAMYR2 and hAMYR3, e.g. at
15 hAMYR3.

Such assays may be performed under the conditions described in Examples 2 and 3.

- Additionally or alternatively, compounds of the invention may show excellent solubility. For example, they may show solubility of greater than or equal to 1 mg/ml at pH 4, pH 5,
20 pH 6, pH 7 and/or pH 7.5, e.g. at 25°C, e.g. under the conditions described in Example 4.

Additionally or alternatively, compounds of the invention may show excellent resistance to fibrillation. For example, they may show no detectable fibrillation after 96 hours at pH 4.0 and/or pH 7.0, e.g. at 40°C, e.g. under the conditions described in Example 5.

- Additionally or alternatively, compounds of the invention may show excellent chemical
25 stability, i.e. resistance to degradation in solution. For example, they may retain at least 70% purity, at least 75% purity, at least 80% purity, at least 85% purity, at least 90% purity, or at least 95% purity after incubation at pH 4, pH 7.5, and/or pH 9 at 40°C for 7 days, e.g. under the conditions described in Example 6.

Therapeutic uses

The compounds of the invention are useful, *inter alia*, in the reduction of food intake, promotion of weight loss, and inhibition or reduction of weight gain. They may therefore
5 provide an attractive treatment option for, *inter alia*, obesity and metabolic diseases caused, characterised by, or associated with, excess body weight.

Thus, the compounds may be used in a method of treating, inhibiting or reducing weight gain, promoting weight loss, reducing food intake, and/or reducing excess body weight. Treatment may be achieved, for example, by control of appetite, feeding, food intake,
10 calorie intake and/or energy expenditure.

The compounds may be used in a method of treating obesity as well as associated diseases, disorders and health conditions, including, but not limited to, morbid obesity, obesity prior to surgery, obesity-linked inflammation, obesity-linked gallbladder disease and obesity-induced sleep apnea and respiratory problems, degeneration of cartilage,
15 osteoarthritis, and reproductive health complications of obesity or overweight such as infertility.

The compounds may also be used in a method of prevention or treatment of Alzheimer's disease, diabetes, type 1 diabetes, type 2 diabetes, pre-diabetes, insulin resistance syndrome, impaired glucose tolerance (IGT), disease states associated with
20 elevated blood glucose levels, metabolic disease including metabolic syndrome, hyperglycemia, hypertension, atherogenic dyslipidemia, hepatic steatosis ("fatty liver"; including non-alcoholic fatty liver disease (NAFLD), which itself includes non-alcoholic steatohepatitis (NASH)), kidney failure, arteriosclerosis (e.g. atherosclerosis), macrovascular disease, microvascular disease, diabetic heart (including diabetic
25 cardiomyopathy and heart failure as a diabetic complication) coronary heart disease, peripheral artery disease or stroke.

The compounds may also be useful in lowering circulating LDL levels and/or increasing HDL/LDL ratio.

The effects of the compounds described above may be mediated in whole or in part via an
30 effect on body weight, or may be independent thereof.

Metabolic syndrome is characterized by a group of metabolic risk factors in one person. They include abdominal obesity (excessive fat tissue around the abdominal internal

organs), atherogenic dyslipidemia (blood fat disorders including high triglycerides, low HDL cholesterol and/or high LDL cholesterol, which foster plaque buildup in artery walls), elevated blood pressure (hypertension), insulin resistance and glucose intolerance, prothrombotic state (e.g. high fibrinogen or plasminogen activator inhibitor-1 in the blood),
5 and proinflammatory state (e.g., elevated C-reactive protein in the blood).

Individuals with metabolic syndrome are at increased risk of coronary heart disease and other diseases related to other manifestations of arteriosclerosis (e.g. stroke and peripheral vascular disease). The dominant underlying risk factor for this syndrome appears to be abdominal obesity.

10 Pharmaceutical compositions

The invention also extends to compositions, such as pharmaceutical compositions, comprising amylin analogues. As with all aspects of the invention, it is to be understood that reference to an amylin analogue encompasses reference to pharmaceutically acceptable salts and solvates.

15 The amylin analogues of the present invention may be formulated as pharmaceutical compositions which are suited for administration with or without storage, and which typically comprise a therapeutically effective amount of at least one peptide of the invention, together with a pharmaceutically acceptable carrier, excipient or vehicle.

The term "pharmaceutically acceptable carrier" includes any of the standard
20 pharmaceutical carriers. Pharmaceutically acceptable carriers for therapeutic use are well known in the pharmaceutical art and are described, for example, in "Remington's Pharmaceutical Sciences", 17th edition, Alfonso R. Gennaro (Ed.), Mark Publishing Company, Easton, PA, USA, 1985. For example, sterile saline and phosphate-buffered saline at slightly acidic or physiological pH may be used. Suitable pH-buffering agents
25 may, e.g., be phosphate, citrate, acetate, tris(hydroxymethyl)aminomethane (TRIS), N-tris(hydroxymethyl)methyl-3-aminopropanesulfonic acid (TAPS), ammonium bicarbonate, diethanolamine, histidine, arginine, lysine or acetate (e.g. as sodium acetate), or mixtures thereof. The term further encompasses any carrier agents listed in the US Pharmacopeia for use in animals, including humans.

30 A pharmaceutical composition of the invention may be in unit dosage form. In such form, the composition is divided into unit doses containing appropriate quantities of the active component or components. The unit dosage form may be presented as a packaged preparation, the package containing discrete quantities of the preparation, for example, packaged tablets, capsules or powders in vials or ampoules. The unit dosage form may

also be, e.g., a capsule, cachet or tablet in itself, or it may be an appropriate number of any of these packaged forms. A unit dosage form may also be provided in single-dose injectable form, for example in the form of a pen device containing a liquid-phase (typically aqueous) composition. Compositions may be formulated for any suitable route and means of administration. Pharmaceutically acceptable carriers or diluents include those used in formulations suitable for e.g. oral, intravitreal, rectal, vaginal, nasal, topical, enteral or parenteral (including subcutaneous (sc), intramuscular (im), intravenous (iv), intradermal and transdermal) administration or administration by inhalation. The formulations may conveniently be presented in unit dosage form and may be prepared by any of the methods well known in the art of pharmaceutical formulation.

Subcutaneous or transdermal modes of administration may in some cases be suitable for peptides of the invention.

Further embodiments relate to devices, dosage forms and packages used to deliver the pharmaceutical formulations of the present invention. Thus, at least one peptide in a stable or preserved formulation or solution described herein can be administered to a patient in accordance with the present invention via a variety of delivery methods, including by sc or im injection, or by transdermal, pulmonary or transmucosal administration, or by implant, or by use of an osmotic pump, cartridge, micro-pump or other means recognized by a person of skill in the art.

Still further embodiments relate to oral formulations and oral administration. Formulations for oral administration may rely on the co-administration of adjuvants (e.g. resorcinols and/or nonionic surfactants such as polyoxyethylene oleyl ether and n-hexadecylpolyethylene ether) to artificially increase the permeability of the intestinal walls, and/or the co-administration of enzymatic inhibitors (e.g. pancreatic trypsin inhibitors, diisopropylfluorophosphate (DFF) or trasylol) to inhibit enzymatic degradation. The active constituent compound of a solid-type dosage form for oral administration can be mixed with at least one additive, such as sucrose, lactose, cellulose, mannitol, trehalose, raffinose, maltitol, dextran, starches, agar, alginates, chitins, chitosans, pectins, gum tragacanth, gum arabic, gelatin, collagen, casein, albumin, synthetic or semisynthetic polymer, or glyceride. These dosage forms can also contain other type(s) of additives, e.g. an inactive diluting agent, a lubricant (such as magnesium stearate), a paraben, a preserving agent (such as sorbic acid, ascorbic acid or alpha-tocopherol), an antioxidant (such as cysteine), a disintegrant, binder, thickener, buffering agent, pH-adjusting agent, sweetening agent, flavoring agent or perfuming agent.

Dosages

A typical dosage of an amylin analogue as employed in the context of the present invention may be in the range from about 0.0001 to about 100 mg/kg body weight per day, such as from about 0.0005 to about 50 mg/kg body weight per day, such as from about 0.001 to about 10 mg/kg body weight per day, e.g. from about 0.01 to about 1 mg/kg body weight per day, administered in one or more doses, such as from one to three doses. The exact dosage employed will depend, *inter alia*, on: the nature and severity of the disease or disorder to be treated, on the sex, age, body weight and general condition of the subject to be treated, on possible other, concomitant, disease or disorder that is undergoing or is to undergo treatment, as well as on other factors that will be known to a medical practitioner of skill in the art.

An amylin analogue of the invention may be administered continuously (e.g. by intravenous administration or another continuous drug administration method), or may be administered to a subject at intervals, typically at regular time intervals, depending on the desired dosage and the pharmaceutical composition selected by the skilled practitioner for the particular subject. Regular administration dosing intervals include, e.g., once daily, twice daily, once every two, three, four, five or six days, once or twice weekly, once or twice monthly, and the like. Such regular peptide administration regimens may, in certain circumstances such as, e.g., during chronic long-term administration, be advantageously interrupted for a period of time so that the medicated subject reduces the level of, or stops taking, the medication, often referred to as taking a "drug holiday." Drug holidays are useful for, e.g., maintaining or regaining sensitivity to a drug especially during long-term chronic treatment, or to reduce unwanted side-effects of long-term chronic treatment of the subject with the drug. The timing of a drug holiday depends on the timing of the regular dosing regimen and the purpose for taking the drug holiday (e.g., to regain drug sensitivity and/or to reduce unwanted side effects of continuous, long-term administration). In some embodiments, the drug holiday may be a reduction in the dosage of the drug (e.g. to below the therapeutically effective amount for a certain interval of time). In other embodiments, administration of the drug is stopped for a certain interval of time before administration is started again using the same or a different dosing regimen (e.g. at a lower or higher dose and/or frequency of administration). A drug holiday of the invention may thus be selected from a wide range of time-periods and dosage regimens. An exemplary drug holiday is two or more days, one or more weeks, or one or more months, up to about 24 months of drug holiday. So, for example, a regular daily dosing regimen with a peptide of the invention may, for example, be interrupted by a drug holiday

of a week, or two weeks, or four weeks, after which time the preceding, regular dosage regimen (e.g. a daily or a weekly dosing regimen) is resumed. A variety of other drug holiday regimens are envisioned to be useful for administering peptides of the invention.

Thus, the peptide may be delivered via an administration regime which comprises two or more administration phases separated by respective drug holiday phases.

During each administration phase, the peptide is administered to the recipient subject in a therapeutically effective amount according to a pre-determined administration pattern. The administration pattern may comprise continuous administration of the drug to the recipient subject over the duration of the administration phase. Alternatively, the administration pattern may comprise administration of a plurality of doses of the peptide to the recipient subject, wherein said doses are spaced by dosing intervals.

A dosing pattern may comprise at least two doses per administration phase, at least five doses per administration phase, at least 10 doses per administration phase, at least 20 doses per administration phase, at least 30 doses per administration phase, or more.

Said dosing intervals may be regular dosing intervals, which may be as set out above, including once daily, twice daily, once every two, three, four, five or six days, once or twice weekly, once or twice monthly, or a regular and even less frequent dosing interval, depending on the particular dosage formulation, bioavailability, and pharmacokinetic profile of the peptide.

An administration phase may have a duration of at least two days, at least a week, at least 2 weeks, at least 4 weeks, at least a month, at least 2 months, at least 3 months, at least 6 months, or more.

Where an administration pattern comprises a plurality of doses, the duration of a possible following drug holiday phase is longer than the dosing interval used in that administration pattern. Where the dosing interval is irregular, the duration of a drug holiday phase may be greater than the mean interval between doses over the course of the administration phase. Alternatively the duration of the drug holiday may be longer than the longest interval between consecutive doses during the administration phase.

The duration of a possible drug holiday phase may be at least twice that of the relevant dosing interval (or mean thereof), at least 3 times, at least 4 times, at least 5 times, at least 10 times, or at least 20 times that of the relevant dosing interval or mean thereof.

Within these constraints, a drug holiday phase may have a duration of at least two days, at least a week, at least 2 weeks, at least 4 weeks, at least a month, at least 2 months, at least 3 months, at least 6 months, or more, depending on the administration pattern during the previous administration phase.

An administration regime entailing the use of drug holiday comprises at least 2 administration phases. Consecutive administration phases are separated by respective drug holiday phases. Thus the administration regime may comprise at least 3, at least 4, at least 5, at least 10, at least 15, at least 20, at least 25, or at least 30 administration phases, or more, each separated by respective drug holiday phases.

Consecutive administration phases may utilise the same administration pattern, although this may not always be desirable or necessary. However, if other drugs or active agents are administered in combination with a peptide of the invention, then typically the same combination of drugs or active agents is given in consecutive administration phases. In certain embodiments, the recipient subject is a human.

Combination therapy

An amylin analogue of the invention may be administered as part of a combination therapy together with another active agent for the treatment of the disease or disorder in question, e.g. an anti-diabetic agent, an anti-obesity agent, an agent for treatment of metabolic syndrome, an anti-dyslipidemia agent, an anti-hypertensive agent, a proton pump inhibitor, or an anti-inflammatory agent. In such cases, the two active agents may be given together or separately, e.g. as constituents in the same pharmaceutical composition or formulation, or as separate formulations.

Thus a peptide of the invention may have some benefit if administered in combination with an anti-diabetic agent of known type, including, but not limited to, metformin, a sulfonylurea, a glinide, a dipeptidyl peptidase IV (DPP-IV) inhibitor, a glitazone, a Glucagon-like peptide-1 (GLP-1) receptor agonist (including GLP-1 or a GLP-1 analogue, an exendin-4 or an exendin-4 analogue, any other GLP-1 receptor agonist including liraglutide (Saxenda™, Victoza™), Dulaglutide or Albiglutide or a glucagon-GLP-1 dual agonist, e.g. as described in WO2008/101017, WO2008/152403, WO2010/070252, WO2010/070253, WO2010/070255, WO2010/070251, WO2011/006497, WO2011/160630, WO2011/160633, WO2013/092703, WO2014/041195), an Sodium-glucose co-transporter-2 (SGLT2) inhibitor (i.e. an inhibitor of sodium-glucose transport,

e.g. a gliflozin such as empagliflozin, canagliflozin, dapagliflozin or ipragliflozin), a G-protein coupled receptor 40 (GPR40) agonist (FFAR1/FFA1 agonist, e.g. fasiglifam), or an insulin or an insulin analogue. Examples of appropriate insulin analogues include, but are not limited to, Lantus™, Novorapid™, Humalog™, Novomix™, Actraphane™ HM, 5 Levemir™, Degludec™ and Apidra™. Other relevant anti-diabetic agents in this connection include GLP-1 receptor agonists, such as exenatide (Byetta™ and Bydureon™, exendin-4) and Byetta LAR™, lixisenatide (Lyxumia™) and liraglutide (Victoza™).

Moreover, a peptide of the invention may be used in combination with an anti-obesity 10 agent of known type, including, but not limited to, peptide YY or an analogue thereof, neuropeptide Y (NPY) or an analogue thereof, a cannabinoid receptor 1 antagonist, a lipase inhibitor, Human proinsulin Peptide (HIP), a melanocortin receptor 4 agonist, a GLP-1 receptor agonist (including GLP-1 or a GLP-1 analogue, an exendin-4 or an exendin-4 analogue, any other GLP-1 receptor agonist including liraglutide (Saxenda™, Victoza™), 15 Dulaglutide or Albiglutide or a glucagon-GLP-1 dual agonist, e.g. as described in WO2008/101017, WO2008/152403, WO2010/070252, WO2010/070253, WO2010/070255, WO2010/070251, WO2011/006497, WO2011/160630, WO2011/160633, WO2013/092703, WO2014/041195), Orlistat™, Sibutramine™, phentermine, a melanin concentrating hormone receptor 1 antagonist, Cholecystokinin 20 (CCK), amylin, pramlintide and leptin, as well as analogues thereof.

A peptide of the invention may further be used in combination with an anti-hypertension agent of a known type, including, but not limited to, an angiotensin-converting enzyme inhibitor, an angiotensin II receptor blocker, a diuretic, a beta-blocker and a calcium channel blocker.

25 A peptide of the invention may still further be used in combination with an anti-dyslipidemia agent of known type, including, but not limited to, a statin, a fibrate, a niacin, a PCSK9 (Proprotein convertase subtilisin/kexin type 9) inhibitor, and a cholesterol absorption inhibitor.

A peptide of the invention may also be used in combination with a proton pump inhibitor 30 (i.e. a pharmaceutical agent possessing pharmacological activity as an inhibitor of H⁺/K⁺-ATPase) of known type, including, but not limited to, an agent of the benzimidazole derivative type or of the imidazopyridine derivative type, such as Omeprazole™, Lansoprazole™, Dexlansoprazole™, Esomeprazole™, Pantoprazole™, Rabeprazole™, Zolpidem™, Alpidem™, Saripidem™ or Necopidem™.

In addition, with regard to anti-inflammatory treatment, a peptide of the invention may be beneficial if administered in combination with an anti-inflammatory agent of known type, including, but not limited to:

- steroids and corticosteroids, such as beclomethasone, methylprednisolone,
5 betamethasone, prednisone, dexamethasone, and hydrocortisone;
- non-steroidal anti-inflammatory agents (NSAIDs), such as propionic acid derivatives (e.g. alminoprofen, benoxaprofen, bucloxic acid, carprofen, fenbufen, fenoprofen, fluprofen, flurbiprofen, ibuprofen, indoprofen, ketoprofen, mioprofen, naproxen, oxaprozin, pirprofen, pranoprofen, suprofen, tiaprofenic acid and tioxaprofen); acetic acid derivatives
10 (e.g. indomethacin, acemetacin, alclofenac, clidanac, diclofenac, fenclofenac, fenclozic acid, fentiazac, furofenac, ibufenac, isoxepac, oxpinac, sulindac, tiopinac, tolmetin, zidometacin and zomepirac); fenamic acid derivatives (e.g. flufenamic acid, meclofenamic acid, mefenamic acid, niflumic acid and tolfenamic acid); biphenylcarboxylic acid derivatives (e.g. diflunisal and flufenisal); oxicams (e.g. isoxicam, piroxicam, sudoxicam
15 and tenoxicam); salicylates (e.g. acetylsalicylic acid and sulfasalazine); and pyrazolones (e.g. apazone, bezpiperylon, feprazone, mofebutazone, oxyphenbutazone and phenylbutazone);
- Cyclooxygenase-II (COX II) inhibitors, such as rofecoxib and celecoxib; preparations of
20 interferon beta (e.g. interferon beta-1a or interferon beta-1b);
- and certain other compounds, such as 5-aminosalicylic acid and prodrugs and pharmaceutically acceptable salts thereof.
- 25 Metformin has also been demonstrated to have anti-inflammatory properties (see, e.g., Haffner et al., *Diabetes* 54: 1566-1572 (2005)) and as such may also be useful in combination with compounds (peptides) of the invention.

Devices and kits

- 30 In some embodiments, the invention relates to a device comprising an amylin analogue or pharmaceutical composition of the invention, for delivery of the analogue to a subject. Via such devices, amylin analogues can be administered to a patient via a variety of delivery methods, including: intravenous, subcutaneous, intramuscular or intraperitoneal injection; oral administration; transdermal administration; pulmonary or transmucosal administration;

administration by implant, osmotic pump, cartridge or micro pump; or by other means recognized by a person of skill in the art.

In some embodiments, the invention relates to a kit comprising an amylin analogue of the invention or a pharmaceutical composition of the invention. In certain embodiments, the
5 kit further comprises packaging and/or instructions for use.

The device or kit may be useful for combination therapy as described above. Thus the device or kit may further comprise a further active agent, e.g. an anti-diabetic agent, an anti-obesity agent, an agent for treatment of metabolic syndrome, an anti-dyslipidemia agent, an anti-hypertensive agent, a proton pump inhibitor, or an anti-inflammatory agent
10 as described above, or a pharmaceutical composition comprising such an active agent.

EXAMPLES

The following examples demonstrate certain specific embodiments of the present invention. The following examples were carried out using standard techniques that are
15 well known and routine to those of skill in the art, except where otherwise described in detail. It is to be understood that these examples are for illustrative purposes only and do not purport to be wholly definitive as to conditions or scope of the invention. As such, they should not be construed as limiting the scope of the present invention in any way.

Abbreviations employed in the examples include:

20	Acm:	acetaminomethyl
	COMU TM :	(1-cyano-2-ethoxy-2-oxoethylidenaminoxy)dimethylamino-morpholino-carbenium hexafluorophosphate
	DCM:	dichloromethane
	DMF:	N,N-dimethylformamide
25	HATU:	2-(7-aza-1H-benzotriazole-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate
	DIPEA:	diisopropylethylamine
	DODT:	3,6-dioxa-1,8-octanedithiol
	EtOH:	ethanol
30	Et ₂ O:	diethyl ether

	TIS:	triisopropylsilane
	MeCN:	acetonitrile
	HPLC:	high performance liquid chromatography
5	RP-HPLC:	reverse phase high performance liquid chromatography
	MS:	mass spectrometry
	ESI-MS:	electron spray ionization mass spectrometry
	IBMX:	3-isobutyl-1-methylxanthine
	BSA:	bovine serum albumin
10	cAMP:	cyclic adenosine monophosphate
	DMEM:	Dulbecco's Modified Eagle Medium
	FCS:	fetal calf serum
	HEPES:	N-2-hydroxyethylpiperazine-N'-2-ethanesulfonic acid
	p-ERK:	phosphorylated extracellular regulated kinase
15	PBS:	phosphate-buffered saline
	Boc:	tert-butoxycarbonyl
	Fmoc:	9-fluorenylmethoxycarbonyl
	Trt:	trityl (i.e. triphenylmethyl)
	NEP:	N-ethylpyrrolidone
20	NMP	N-methylpyrrolidone
	v/v:	volume/volume
	w/v:	weight/volume

The following examples are provided to illustrate certain embodiments of the invention and are not intended to limit the scope of the invention.

Unless otherwise specified, whole-blood glucose levels were determined on tail-vein blood samples by the Biosen (EKF Diagnostic, Germany) enzyme-based electrode method.

- 5 Blood samples were analyzed for glycated hemoglobin (HbA1c) using a Cobas c111 analyzer (Roche Diagnostics, Mannheim, Germany). Plasma insulin levels were measured using a Meso Scale Discovery (MSD) system (Rockville, MD, USA). Liver fat content was determined by magnetic resonance (MR) scanning using an Echo systems MR scanner. Fat depots were measured by weighing of excised fat.

Example 1: Synthesis of compounds

- 10 The following compounds were synthesised:

[19CD]-isoGlu-R-C()-NTAT-C()-ATQRLADFLHRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSETP-NH₂ (Compd. 1)

[19CD]-isoGlu-R-C()-NTAT-C()-ATQRLAHFLHRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSETP-NH₂ (Compd.2)

- 15 [19CD]-isoGlu-R-C()-NTAT-C()-ATQRLANFLHRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSETP-NH₂ (Compd. 3)

[19CD]-isoGlu-R-C()-NTAT-C()-ATQRLADFLHRSSNNF-Gly(Me)-A-Ile(Me)-LSST-NVGSNT-Apr-NH₂ (Compd. 4)

- 20 [19CD]-isoGlu-R-C()-NTAT-C()-ATQRLAHFLHRSSNNF-Gly(Me)-A-Ile(Me)-LSST-NVGSNT-Apr-NH₂ (Compd. 5)

[19CD]-isoGlu-R-C()-GTAT-C()-ATERLAHFLNRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSNTHyp-NH₂ (Compd. 6)

[19CD]-isoGlu-R-C()-GTAT-C()-ATERLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSNTHyp-NH₂ (Compd. 7)

- 25 [19CD]-isoGlu-R-C()-GTAT-C()-ATERLAHFLELRRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSNTHyp-NH₂ (Compd. 8)

[19CD]-isoGlu-R-C()-GTAT-C()-ATERLAHFLTRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSSTHyp-NH₂ (Compd. 9)

- [19CD]-isoGlu-R-C()-GTAT-C()-ATERLAHFLTRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGST-Hyp-NH₂ (Compd. 10)
- [19CD]-isoGlu-R-C()-GTAT-C()-ATERLAHFLTRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGST-Orn-T-Hyp-NH₂ (Compd. 11)
- 5 [19CD]-isoGlu-R-C()-GTAT-C()-ATERLAHFLTRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGST-Aad-T-Hyp-NH₂ (Compd. 12)
- [19CD]-isoGlu-R-C()-GTAT-C()-ATERLAHFLTRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSGT-Hyp-NH₂ (Compd. 13)
- [19CD]-isoGlu-R-C()-NTAT-C()-ATQRLAHFLHRF-Gly(Me)-A-Ile(Me)-LSSTEVGSETP-NH₂ (Compd. 14)
- 10 [19CD]-isoGlu-R-C()-GTAT-C()-ATERLAHFLHRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSNTP-NH₂ (Compd. 15)
- [19CD]-isoGlu-R-C()-NTAT-C()-ATQRLAHFLHRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSNTP-NH₂ (Compd. 16).
- 15 [19CD]-isoGlu-R-C()-GTAT-C()-ATERLANFLVRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSNTHyp-NH₂ (Compd. 17)
- [19CD]-isoGlu-R-C()-GTAT-C()-ATERLAHFLTRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSNTHyp-NH₂ (Compd. 18)
- [19CD]-isoGlu-R-C()-GTAT-C()-ATERLAHFLKRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSNTHyp-NH₂ (Compd. 19)
- 20 [19CD]-isoGlu-R-C()-NTAT-C()-ATQRLAHFLHRVSF-Gly(Me)-A-Ile(Me)-LSSTEVGSETP-NH₂ (Compd. 20)
- [19CD]-isoGlu-R-C()-NTAT-C()-ATQRLAHFLHRVSF-Gly(Me)-A-Ile(Me)-LSSTEVGSETP-NH₂ (Compd. 21)
- 25 [19CD]-isoGlu-R-C()-GTAT-C()-ATERLAHFLVRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSETP-NH₂ (Compd. 22)
- [19CD]-isoGlu-R-C()-PTAT-C()-ATDRLAHFLVRSSF-Gly(Me)-A-Ile(Me)-LSSTDVGSDTP-NH₂ (Compd. 23)

- [19CD]-isoGlu-R-C()-NTAT-C()-ATQRLAHFLVRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSETP-NH₂ (Compd. 24)
- [19CD]-isoGlu-R-C()-PTAT-C()-ATDRLAHFLHRSSF-Gly(Me)-A-Ile(Me)-LSSTDVGSNTP-NH₂ (Compd. 25)
- 5 [19CD]-isoGlu-R-C()-GTAT-C()-ATERLAHFLHRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSETP-NH₂ (Compd. 26)
- [19CD]-isoGlu-R-C()-PTAT-C()-ATDRLAHFLHRSSF-Gly(Me)-A-Ile(Me)-LSSTDVGSSTP-NH₂ (Compd. 27)
- [19CD]-isoGlu-R-C()-GTAT-C()-ATERLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSSST-Hyp-NH₂ (Compd. 28)
- 10 [19CD]-isoGlu-R-C()-GTAT-C()-ATERLAHFLNRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSSST-Hyp-NH₂ (Compd. 29)
- [19CD]-isoGlu-R-C()-GTAT-C()-ATERLAHFLHRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSSST-Hyp-NH₂ (Compd. 30)
- 15 [19CD]-isoGlu-R-C()-GTAT-C()-ATERLAHFLVRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSSST-Hyp-NH₂ (Compd. 31)
- [19CD]-isoGlu-R-C()-GTAT-C()-ATERLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSSFT-Hyp-NH₂ (Compd. 32)
- [19CD]-isoGlu-R-C()-GTAT-C()-ATERLAHFLNRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSSFT-Hyp-NH₂ (Compd. 33)
- 20 [19CD]-isoGlu-R-C()-GTAT-C()-ATERLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSSTP-NH₂ (Compd. 34)
- [19CD]-isoGlu-R-C()-GTAT-C()-ATERLAHFLNRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSSTP-NH₂ (Compd. 35)
- 25 [19CD]-isoGlu-R-C()-GTAT-C()-ATERLAHFLELRRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSSTP-NH₂ (Compd. 36)
- [19CD]-isoGlu-R-C()-GTAT-C()-ATERLAHFLHRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSSTP-NH₂ (Compd. 37)

- [19CD]-isoGlu-R-C()-GTAT-C()-ATERLAHFLVRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSSSTP-NH₂ (Compd. 38)
- [19CD]-isoGlu-R-C()-GTAT-C()-ATERLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSSFTP-NH₂ (Compd. 39)
- 5 [19CD]-isoGlu-R-C()-GTAT-C()-ATERLAHFLNRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSSFTP-NH₂ (Compd. 40)
- [19CD]-isoGlu-R-C()-GTAT-C()-ATERLADFLTRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSSST-Hyp-NH₂ (Compd. 41)
- [19CD]-isoGlu-R-C()-GTAT-C()-ATERLADFLTRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSSFT-Hyp-NH₂ (Compd. 42)
- 10 [19CD]-isoGlu-R-C()-GTAT-C()-ATERLA-Aad-FLTRSSF-Gly(Me)-A-Ile(Me)-LSSTE-VGSST-Hyp-NH₂ (Compd. 43)
- [19CD]-isoGlu-R-C()-GTAT-C()-ATERLA-Aad-FLTRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSS-Orn-T-Hyp-NH₂ (Compd. 44)
- 15 [19CD]-isoGlu-R-C()-GTAT-C()-ATERLAHFLTRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSSST-Apr-NH₂ (Compd. 45)
- [19CD]-isoGlu-R-C()-GTAT-C()-ATERLAHFLTRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSSSTP-NH₂ (Compd. 46)
- [19CD]-isoGlu-R-C()-GTAT-C()-ATERLAHFLTRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSS-Orn-TP-NH₂ (Compd. 47)
- 20 [19CD]-isoGlu-R-C()-GTAT-C()-ATERLAHFLTRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSTT-Hyp-NH₂ (Compd. 48)
- [19CD]-isoGlu-R-C()-NTAT-C()-ATQRLADFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSSNT-Hyp-NH₂ (Compd. 49)
- 25 [19CD]-isoGlu-R-C()-NTAT-C()-ATQRLAHFL-Aad-RSSF-Gly(Me)-A-Ile(Me)-LSSTE-VGSNT-Hyp-NH₂ (Compd. 50)
- [19CD]-isoGlu-R-C()-GTAT-C()-ATERLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSS-Orn-T-Hyp-NH₂ (Compd. 51)

- [19CD]-isoGlu-R-C()-GTAT-C()-ATERLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSGT-Hyp-NH₂ (Compd. 52)
- [19CD]-isoGlu-R-C()-GTAT-C()-ATERLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSAad-T-Hyp-NH₂ (Compd. 53)
- 5 [19CD]-isoGlu-R-C()-GTAT-C()-ATERLADFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSNTHyp-NH₂ (Compd. 54)
- [19CD]-isoGlu-R-C()-GTAT-C()-ATERLA-Aad-FLQRSSF-Gly(Me)-A-Ile(Me)-LSSTE-VGSNT-Hyp-NH₂ (Compd. 55)
- [19CD]-isoGlu-R-C()-NTAT-C()-ATERLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSNTHyp-NH₂ (Compd. 56)
- 10 [19CD]-isoGlu-R-C()-QTAT-C()-ATERLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSNTHyp-NH₂ (Compd. 57)
- [19CD]-isoGlu-R-C()-NTAT-C()-ATQRLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSNTHyp-NH₂ (Compd. 58)
- 15 [19CD]-isoGlu-R-C()-PTAT-C()-ATDRLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTDVGSNTHyp-NH₂ (Compd. 59)
- [19CD]-isoGlu-R-C()-NTAT-C()-ATERLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSDTHyp-NH₂ (Compd. 60)
- [19CD]-isoGlu-R-C()-NTAT-C()-ATQRLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSNTP-NH₂ (Compd. 61)
- 20 [19CD]-isoGlu-R-C()-NTAT-C()-ATERLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSNTP-NH₂ (Compd. 62)
- [19CD]-isoGlu-R-C()-PTAT-C()-ATERLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSNTHyp-NH₂ (Compd. 63)
- 25 [19CD]-isoGlu-R-C()-PTAT-C()-ATQRLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSNTHyp-NH₂ (Compd. 64)
- [19CD]-isoGlu-R-C()-NTAT-C()-ATQRLADFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSNTP-NH₂ (Compd. 65)

- [19CD]-isoGlu-R-C()-NTAT-C()-ATERLADFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSNTP-NH₂ (Compd. 66)
- [19CD]-isoGlu-R-C()-PTAT-C()-ATERLADFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSNTP-NH₂ (Compd. 67)
- 5 [19CD]-isoGlu-R-C()-NTAT-C()-ATERLADFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSNTP-Hyp-NH₂ (Compd. 68)
- [19CD]-isoGlu-R-C()-NTAT-C()-ATQRLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSSST-Hyp-NH₂ (Compd. 69)
- [19CD]-isoGlu-R-C()-PTAT-C()-ATERLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSSST-Hyp-NH₂ (Compd. 70)
- 10 [19CD]-isoGlu-R-C()-NTAT-C()-ATQRLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSGT-Hyp-NH₂ (Compd. 71)
- [19CD]-isoGlu-R-C()-PTAT-C()-ATERLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSGT-Hyp-NH₂ (Compd. 72)
- 15 [19CD]-isoGlu-R-C()-NTAT-C()-ATQRLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSTT-Hyp-NH₂ (Compd. 73)
- [19CD]-isoGlu-R-C()-PTAT-C()-ATERLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSTT-Hyp-NH₂ (Compd. 74)
- [19CD]-isoGlu-R-C()-NTAT-C()-ATQRLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSAad-T-Hyp-NH₂ (Compd. 75)
- 20 [19CD]-isoGlu-R-C()-PTAT-C()-ATERLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSAad-T-Hyp-NH₂ (Compd. 76)
- [19CD]-isoGlu-R-C()-GTAT-C()-ATERLAHFLQRF-Gly(Me)-A-Ile(Me)-LSSTEVGSNTP-Hyp-NH₂ (Compd. 77)
- 25 [19CD]-isoGlu-R-C()-GTAT-C()-ATERLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSNTP-NH₂ (Compd. 78)
- [19CD]-isoGlu-R-C()-NTAT-C()-ATQRLAHFLQRF-Gly(Me)-A-Ile(Me)-LSSTEVGSGT-Hyp-NH₂ (Compd. 79)

- [19CD]-isoGlu-R-C()-NTAT-C()-ATQRLADFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSGTP-NH₂ (Compd. 80)
- [19CD]-isoGlu-R-C()-NTAT-C()-ATQRLADFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSGT-Hyp-NH₂ (Compd. 81)
- 5 [19CD]-isoGlu-R-C()-NTAT-C()-ATQRLADFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSTTP-NH₂ (Compd. 82)
- [19CD]-isoGlu-R-C()-NTAT-C()-ATQRLADFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSTT-Hyp-NH₂ (Compd. 83)
- [19CD]-isoGlu-R-C()-NTAT-C()-ATQRLAHFLERSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSSTP-NH₂ (Compd. 84)
- 10 [19CD]-isoGlu-R-C()-NTAT-C()-ATQRLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSSTP-NH₂ (Compd. 85)
- [19CD]-isoGlu-R-C()-NTAT-C()-ATQRLA-Aad-FLQRSSF-Gly(Me)-A-Ile(Me)-LSSTE-VGSNT-Hyp-NH₂ (Compd. 86)
- 15 [19CD]-isoGlu-R-C()-NTAT-C()-ATQRLA-Aad-FLQRSSF-Gly(Me)-A-Ile(Me)-LSSTE-VGSST-Hyp-NH₂ (Compd. 87)
- [19CD]-isoGlu-R-C()-QTAT-C()-ATDRLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTE-VGSNT-Hyp-NH₂ (Compd. 88)
- [19CD]-isoGlu-R-C()-QTAT-C()-ATDRLA-Aad-FLQRSSF-Gly(Me)-A-Ile(Me)-LSSTE-VGSNT-Hyp-NH₂ (Compd. 89)
- 20 [19CD]-isoGlu-R-C()-QTAT-C()-ATERLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTE-VGSGT-Hyp-NH₂ (Compd. 90)
- [19CD]-isoGlu-R-C()-QTAT-C()-ATDRLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTE-VGSGT-Hyp-NH₂ (Compd. 91)
- 25 [19CD]-isoGlu-R-C()-QTAT-C()-ATDRLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTE-VGSNTP-NH₂ (Compd. 92)
- [19CD]-isoGlu-R-C()-QTAT-C()-ATERLA-Aad-FLQRSSF-Gly(Me)-A-Ile(Me)-LSSTE-VGSNTP-NH₂ (Compd. 93)

[19CD]-isoGlu-R-C()-QTAT-C()-ATERLA-Aad-FLQRSSF-Gly(Me)-A-Ile(Me)-LSSIE-VGSGTP-NH₂ (Compd. 94)

[19CD]-isoGlu-R-C()-QTAT-C()-ATDRLA-Aad-FLQRF-Gly(Me)-A-Ile(Me)-LSSTEVGsNT-Hyp-NH₂ (Compd. 95)

- 5 [19CD]-isoGlu-R-C()-QTAT-C()-ATERLAHFLQRF-Gly(Me)-A-Ile(Me)-LSSTEVGsNTP-NH₂ (Compd. 96)

([19CD]-isoGlu represents a [19-carboxynonadecanoyl] group linked to the free alpha-amino group of the N-arginine residue via an iso-glutamic acid linker.)

- 10 Also synthesized, for use as controls, were:

Pramlintide: H-KC()NTATC()ATQRLANFLVHSSNFGPILPPTNVGSNTY-NH₂

- 15 NN96: N-α-[(S)-4-carboxy-4-(19-carboxynonadecanoylamino)butyryl]-[Arg1,Glu14,His17,Pro37]-pramlintide-NH₂

(NN96 is disclosed in WO 2012/168430 and has the amino acid sequence: RC()NTATC()ATQRLAEFLHHSSNFGPILPPTNVGSNTP)

- 20 [19CD]-isoGlu-R-C()-NTAT-C()-ATQRLAEFLHHSSFGPILPPTNVGSNTP-NH₂ (Compd. 97)

[19CD]-isoGlu-R-C()-GTAT-C()-ATERLA-Aad-FLQRSSNNF-Gly(Me)-A-Ile(Me)-LSSTEVGsNT-Hyp-NH₂ (Compd. 98)

[19CD]-isoGlu-R-C()-QTAT-C()-ATERLAHFLQRSSNNF-Gly(Me)-A-Ile(Me)-LSSTEVGsNT-Hyp-NH₂ (Compd. 99)

- 25 [19CD]-isoGlu-R-C()-QTAT-C()-ATDRLA-Aad-FLQRSSNNF-Gly(Me)-A-Ile(Me)-LSSTEVGsNT-Hyp-NH₂ (Compd. 100)

Compound 97 has the same sequence as NN96 except for a deletion of the Asn residues at positions 21 and 22.

- 30 Compounds 98, 99 and 100 have the same sequences as compound 55, compound 57 and compound 89 respectively, except for Asn residues inserted at positions 21 and 22.

Parentheses “()” indicate intramolecular disulphide bridges between the side chains of cysteine residues at positions 2 and 7 of the relevant amino acid sequences.

- 5 Unless otherwise specified, reagents and solvents employed in the following were available commercially in standard laboratory reagent or analytical grade, and were used without further purification.

Suitable general procedures for synthesis include the following:

General procedures for solid-phase synthesis of peptides

- 10 A CEM Liberty Peptide Synthesizer was employed, using standard Fmoc chemistry. TentaGel™ S Ram resin (1 g; 0.25 mmol/g) was swelled in DMF (10 ml) prior to use and transferred between tube and reaction vessel using DCM and DMF. Pseudoprolines [i.e. dipeptides employed to minimize aggregation during peptide synthesis, such as Fmoc-Phe-Thr(ψ -Me,Me-Pro)-OH and Fmoc-Asp-Ser(ψ -Me,Me-Pro)-OH and Fmoc-Ser-Ser(ψ -Me,Me-Pro)-OH] were used where appropriate, and non-naturally occurring amino acids and other suitable building blocks were employed without any changes to the general procedure.
- 15

The following optical isomers of particular amino acids (including non-naturally occurring amino acids) were employed in the synthesis of the compounds:

- 20 Apr: (2S,4R)-4-aminoproline [also denoted (4R)-4-amino-L-proline];
Hyp: (2S,4R)-4-hydroxyproline [also denoted (4R)-4-hydroxy-L-proline].
Aad: (2S)-2-aminoadipic acid

Coupling:

- 25 An Fmoc-amino acid in DMF/DCM (2:1; 0.2 M; 5 ml) was added to the resin in a CEM Discover microwave unit together with COMU/DMF (0.5 M; 2 ml) and DIPEA/DMF (2.0 M; 1 ml). The coupling mixture was heated to 75°C for 5 min. while nitrogen was bubbled through the mixture. The resin was then washed with DMF (4 x 10 ml). Alternatively the coupling was done without heating and the reaction time extended to 60 min in this case.

In the case of difficult couplings (e.g. coupling of a residue immediately after an N-methylated amino acid residue or other sterically hindered amino acid residue as recognized by a person of skill in the art) the coupling was repeated one or more times.

Deprotection:

- 5 Piperidine/DMF (1:4, i.e. 1 part piperidine to 4 parts DMF by volume; 10 ml) was added to the resin for initial deprotection, and the mixture was microwave-heated (40°C; 30 sec.). The reaction vessel was drained and a second portion of piperidine/DMF (1:4; 10 ml) was added and heated (75°C; 3 min) again. The resin was then washed with DMF (6 x 10 ml).

Oxidative cyclisation

- 10 Intramolecular ring formation (disulfide bridge formation) between the Cys residues in positions 2 and 7 (initially coupled in the form of Acm-protected cysteines) was performed with the peptide still attached to the resin, using 163 mg thallium(III) trifluoroacetate [Tl(TFA)₃] in 5 mL NMP in a simultaneous Acm-deprotection and disulfide-formation step.

Cleavage:

- 15 The resin was washed with EtOH (3 x 10 ml) and Et₂O (3 x 10 ml) and dried to constant weight at room temperature (r.t.). The crude peptide was cleaved from the resin by treatment with TFA/TIS/H₂O (90:5:5; 40 ml; 2 h; room temp.) or alternatively with TFA/DODT (95:5; 40 ml; 2h; room temp.). Most of the TFA was removed under reduced pressure, and the crude peptide was precipitated and washed three times with Et₂O and
20 dried to constant weight at room temperature.

Purification and characterisation:

- The crude peptide was purified by preparative reverse phase HPLC using a PerSeptive Biosystems VISION Workstation or a Gilson system (Pumps: "Pump 305", "331 Pump", "332 Pump", "402 Syringe Pump"; column changer "Valvemate® II" UV detector "UV/Vis-
25 155"; and the fraction collector "GX 281" equipped with a suitable column and a fraction collector, and run with a gradient of buffer A (0.1% aqueous TFA) and buffer B (0.1% TFA, 90% MeCN, water). Fractions were analysed by analytical HPLC and MS, and relevant fractions were pooled and lyophilised. The final product was characterized by HPLC and MS.
- 30 One of skill in the art will appreciate that standard methods of peptide synthesis may be used to generate the compounds of the invention.

Generation of cell lines expressing (i) human calcitonin receptor (hCTR2), and (ii) human amylin receptor 3 (hAMYR3) consisting of hCTR2 co-expressed with human Receptor Activity Modifying Protein 3 (hRAMP3)

- 5 The cell line COS7, originating from African Green Monkey kidney fibroblast cells, was used for establishment of the stable cell lines described below. The COS7 cellular background was used due to it being RAMP naïve, and as such it provides the best available system for testing of the monomeric hCTR and different heterodimeric amylin receptor subtypes functionally generated by introduction of the individual RAMPs.

10 Calcitonin receptor cell line:

A cell line expressing the human calcitonin receptor (hCTR2) was generated in the COS7 cellular background as stable clones. In brief, hCTR2 (GI: 4502547) was amplified by PCR using primers encoding terminal restriction sites for subcloning. The 5'-end primers additionally encoded a near Kozak consensus sequence to ensure efficient translation.

- 15 The fidelity of the DNA encoding the receptor was confirmed by DNA sequencing. The PCR products encoding the receptor were subcloned into a mammalian expression vector containing a neomycin (G418) resistance marker. The mammalian expression vector encoding the receptor was transfected into COS7 cells by a standard liposome transfection method. 48 hours post-transfection, cells were seeded for limited dilution
- 20 cloning and selected with 1 mg/ml G418 in the culture medium. After 3 weeks, surviving colonies of hCTR2-expressing cells were picked, propagated and tested in the amylin efficacy assay as described in Example 3 below. One hCTR2-expressing clone was chosen for compound profiling.

25 Amylin receptor cell line:

A cell line expressing the human amylin receptor 3 (hAMYR3) was generated in the COS7 cellular background as stable clones. In brief, hCTR2 (GI:4502547) and hRAMP3 (GI:118572586) were amplified by PCR using primers encoding terminal restriction sites for subcloning. The 5'-end primers additionally encoded a near Kozak consensus

30 sequence to ensure efficient translation. The fidelity of the DNA encoding the receptor was confirmed by DNA sequencing. The PCR products encoding the receptor were subcloned into a proprietary tri-cistronic mammalian expression vector with RAMP3, hCTR2 and neomycin (G418) resistance marker. The mammalian expression vector

encoding the receptor was transfected into COS7 cells by a standard liposome

transfection method. 48 hours post-transfection, cells were seeded for limited dilution

cloning and selected with 1 mg/ml G418 in the culture medium. After 3 weeks, surviving

colonies of hAMY3R-expressing cells were picked, propagated and tested in the amylin

5 efficacy assay as described in Example 3 below. The functional generation of the amylin

receptor phenotype was verified in the efficacy assay by a left-shifted efficacy response to

amylin relative to monomeric calcitonin receptor, and one hAMYR3-expressing clone was

chosen for compound profiling.

Example 3

10 hCTR2 and hAMYR3 assays

In vitro activity of test peptides at the hCTR2 and hAMYR3 receptors was assessed by

measuring the induction of cAMP following stimulation of the receptor using the

AlphaScreen® cAMP Assay kit from Perkin-Elmer.

Briefly, COS7 cells expressing hCTR2 (see Example 2, above) were seeded at 30-40,000

15 cells per well in 96-well microtiter plates coated with 0.01 % poly-L-lysine, and grown for 1

day in culture in 100 μ l growth medium [DMEM, 10% FCS, Penicillin (100 IU/ml),

Streptomycin (100 μ g/ml)]. On the day of analysis, growth medium was removed and the

cells were washed once with 200 μ l Tyrode buffer [Tyrode's Salts (9.6 g/l), 10 mM

HEPES, pH 7.4]. Cells were incubated in 100 μ l Tyrode buffer containing increasing

20 concentrations of test peptides, 100 μ M IBMX and 0.1% casein for 15 min at 37° C. The

reaction was stopped by carefully decanting off the compound/buffer medium and

replacing it with lysis/detection buffer (80 μ l 0.1 % w/v BSA, 5 mM HEPES, 0.3 % v/v

Tween-20 in deionized water). After incubation at room temperature for 10 min., the cAMP

content of the resulting cell lysate was estimated according to the AlphaScreen® cAMP

25 Assay manufacturer's instructions. EC₅₀ values were estimated by computer-aided curve

fitting using the 4-parameter logistic (4PL) non-linear model.

The *in vitro* activity results (expressed as EC₅₀ values) are summarized in Table 1, below.

Example 4

Assessment of solubility

30 A stock solution of the test peptide (2 mg/ml; determined from the weighed amount of

peptide) in demineralized water adjusted to pH 2.5 with HCl was prepared, and aliquots

were diluted 1:1 in 100 mM acetate buffer (pH 4.0 and pH 5.0), 100 mM histidine buffer (pH 6.0 and pH 7.0) and 100 mM phosphate buffer (pH 6.0, pH 7.0 and pH 7.5), respectively, and loaded in a standard flat-bottom, non-sterile 96-well UV Microplate. The absorbance of samples (single samples, n=1) at 280 and 325 nm was measured in an absorbance-based plate reader, which was preheated to ambient temperature (typically 25°C). The turbidity absorbance criterion for a peptide solubility of ≥ 1 mg/ml was an absorbance at 325 nm of ≤ 0.02 absorbance units (which is 5 to 6 times the standard deviation of 8 buffer samples in a plate).

Measurements were made on Compounds 1-17, 19, 27-38, 49-69 and 71-87. With the exception of Compound 10 at pH 6.0. Compound 31 at pH 7.0 and 7.5, and Compound 38 at pH 6.0 and 7.5, all of the tested compounds exhibited solubility ≥ 1 mg/ml in all the test buffers.

Example 5

Assessment of physical stability

Aggregation in the form of fibril formation was detected using the amyloid-specific dye Thioflavin T (ThT), which is frequently employed to demonstrate the presence of fibrils in solution (see, e.g., Groenning, M., J. Chem. Biol. **3**(1) (2010), pp. 1-18; Groenning et al., J. Struct. Biol. **158** (2007) pp. 358-369; and Levine, H., III, Protein Sci. **2** (1993) pp. 404-410) Test peptides (2 mg/ml) were dissolved in demineralized water adjusted to pH 2.5 with HCl, at ambient temperature (typically 25°C). Solutions containing (i) 1 mg/ml of test peptide, 40 μ M ThT and 50 mM phosphate (Ph) buffer (pH 7.0), (ii) 1 mg/ml of test peptide, 40 μ M ThT and 50 mM histidine (His) buffer (pH 7.0), and (iii) 1 mg/ml of test peptide, 40 μ M ThT and 50 mM acetate (Ac) buffer (pH 4.0), were loaded in a 96-well black fluorescence plate (clear bottom) in triplicate. Data were collected at fixed intervals of 10 min, each preceded by 300 s of automixing (agitation), over a period of 96 hours at 40 °C. Physical stability, expressed as lag-time of fibril formation (in hours), was defined as the intersection between two linear regressions representing the initial stable phase and the growth phase. Data are summarized in Table 1 below.

Example 6Assessment of chemical stability

5 Samples of each test peptide were dissolved in MilliQ™ water, and the pH of the solution was adjusted to pH 4, pH 6, pH 7.5 or pH 9, respectively, using either HCl or NaOH. The final peptide concentration was 0.2 mg/ml. Samples were placed in glass vials and incubated at 40 °C. The samples were analyzed by RP-HPLC on a C8 column with gradient elution using an ammonium formate/water eluent system, or on a C18 column with gradient elution using a trifluoroacetic acid/water eluent system. The area-percentage
10 (area-%) of the main peak after incubation time $T = t$ (relative to time $T = 0$) was determined by UV spectroscopy at 220 nm.

The purity was first determined as follows:

Purity (area-%) = (area of main peak/total area of all peaks) x 100.

The purity was then normalized between time points by setting purity at time 0 ($T=0$) to
15 100 for each pH value for a given peptide, as follows:

Normalised area-% at time t ($T=t$) = [area-% ($T=t$)/area-% ($T=0$)] x 100.

The chemical stability assessment results (in the form of normalized purity values) are summarized in Table 1 (below). The normalized purity values in Table 1 were determined after 7 days of incubation.

Table 1. EC₅₀, chemical stability and fibrillation data

Cmpd. No.	hCTR2 EC ₅₀ (nM)	hAMYR3 EC ₅₀ (nM)	Normalized purity after 7 days, 40°C**				Fibrillation ^{§§}		
			pH 4	pH 6	pH 7.5	pH 9	pH 4.0 (Ac)	pH 7.0 (His)	pH 7.0 (Phos.)
1	0.079	0.230					FND	FND	FND
2	0.076	0.120	A		A	B	FND	FD	FND
3	0.160	0.200	A		B	B	FD	FD	FD
4	0.210	0.220	B		C	C	FND	FND	FND
5	0.200	0.180	B		C	C	FND	FND	FND
6	0.077	0.220	A	B		B	FND	FND	FND
7	0.087	0.340	A	A	A	A	FND	FND	FND
8	0.180	0.360	A	B	A	A	FND	FND	FND
9	0.099	0.290	A	B	B	B	FND	FD	FD
10	0.160	0.350	A	A	B	B	FND	FD	FD
11	0.130	0.200	A	B	A	B	FND	FD	FD
12	0.120	0.250	A	A	B	B	FND	FD	FD
13	0.120	0.380	A	A	A	A	FND	FD	FD
14	0.150	0.260	A	A	A	A	FND	FD	FND
15	0.044	0.079	A	B	B	B	FND	FD	FND
16	0.073	0.110	A	A	A	B	FD	FD	FND
17	0.13	0.61							
18	0.12	0.41					FND	FD	FD
19	0.16	0.28	A	A	A	C	FND	FND	FND
20	0.19	0.33					FND	FD	FND
21	0.12	0.23					FND	FD	FD
22	0.078	0.59					FND	FD	FND
23	0.094	0.13							
24	0.085	0.19							
25	0.046	0.094							
26	0.057	0.12							
27	0.063	0.16					FD	FD	FD
28	0.11	0.13	A	B	B	B	FND	FND	FND
29	0.09	0.12	A		B	B			
30	0.07	0.12					FND	FD	FD
31	0.09	0.29							
32	0.17	0.23					FND	FD	FD
33	0.11	0.15							
34	0.09	0.12	A	B	B	B	FND	FND	FND
35	0.05	0.11	A	A	B	B	FND	FND	FND
36	0.21	0.36	A	A	B	B	FND	FND	FD

37	0.04	0.07							
38	0.06	0.12							
39	0.16	0.22							
40	0.12	0.19							
41	0.15	0.34	B	A	A	A			
42	0.12	0.34							
43	0.22	0.35	A	A	A	A			
44	0.2	0.3							
45	0.16	0.31							
46	0.1	0.14							
47	0.1	0.19							
48	0.12	0.26	A	A	A	A	FND	FND	FND
49	0.06	0.15	A	A	A	B	FD	FND	FND
50	0.12	0.19							
51	0.08	0.11	A	B	B	B	FND	FND	FND
52	0.07	0.09	A	B	B	B	FND	FND	FND
53	0.14	0.13	A	A	B	B	FND	FND	FND
54	0.08	0.14	A	A	B	B	FND	FND	FND
55	0.11	0.08	A	A	A	B	FND	FND	FND
56	0.06	0.08	A	A	A	B	FND	FND	FND
57	0.08	0.11	A	A	A	A	FND	FND	FND
58	0.05	0.1	A	A	B	B	FND	FD	FND
59	0.03	0.11	A	B	C	B	FND	FD	FD
60	0.08	0.09	A	A	A	B	FND	FND	FND
61	0.12	0.17							
62	0.18	0.22							
63	0.35	0.37							
64	0.11	0.15							
65	0.14	0.26	A	A	A	A			
66	0.19	0.30							
67	0.27	0.65							
68	0.32	0.29							
69	0.12	0.17	A	A	B	B			
70	0.35	0.36							
71	0.11	0.18	A	A	A	B	FND	FD	FD
72	0.35	0.27							
73	0.12	0.24							
74	0.58	0.39							
75	0.36	0.26							
76	0.44	0.49							
77	0.16	0.42	A	A	B	B	FND	FND	FND
78	0.15	0.24							
79	0.12	0.23	A	A	B	B	FND	FD	FD
80	0.13	0.28	A	A	A	A			
81	0.10	0.23	A	A	A	A			
82	0.21	0.53							

83	0.13	0.32							
84	0.15	0.54							
85	0.1	0.14							
86	0.1	0.15	A	A	A	A			
87	0.2	0.13							
88	0.06		A	A	A	A	FND	FND	FND
89	0.16	0.20	A	A	A	B	FND	FND	FND
90	0.20	0.24	A	A	A	A	FND	FND	FND
91	0.08	0.17	A	A	A	A	FND	FND	FND
92	0.10	0.22	A	A	A	A			
93		0.43							
95	0.18	0.20	A	A	A	A			
96	0.33	0.28	A	A	A	A	FND	FND	FND
Pramlin- tide	1.40	0.230					FD		FD
NN96	0.089	0.150	A	A	B	B	FND	FD	FD

** A = >90%; B = 70-90%; C = <70%; Blank = not determined.

§§ FND = fibrillation not detected; FD = fibrillation detected.

5 Example 7

Pharmacokinetic (PK) profiling in rats

Compounds 6, 7, 9, 11, 13, 15, 28, 34, 36, 49, 55, 57 and 71 were tested for their pharmacokinetic properties in rats.

10 Sprague Dawley male rats were given a single subcutaneous (sc) bolus of each peptide to be tested, as specified below.

30 nmol/kg doses of compound were administered. Blood samples were drawn from the tail vein prior to dosing and at 24, 48, 72, 96 and 168 hours after dosing. Blood samples were taken from two rats at each time point, and only 2 blood samples were drawn from each rat, i.e. 10 rats were used for each compound. The rats were euthanized
15 immediately after the last blood sampling by concussion and cervical dislocation.

The dosing vehicle used for each test peptide was a mannitol-containing histidine buffer (pH 7.0). Plasma samples were analyzed after precipitation with ethanol by liquid chromatography mass spectrometry (LC-MS/MS). Mean plasma concentrations were used for calculation of the pharmacokinetic parameters using the non-compartmental
20 approach in Phoenix WinNonlin 6.3.

Plasma terminal elimination half-life ($t_{1/2}$) was determined as $\ln(2)/\lambda_z$, where λ_z is the magnitude of the slope of the log linear regression of the log concentration versus time profile during the terminal phase. Apparent clearance (CL/F) was determined as dose/AUC_{inf} (sc), where AUC_{inf} is the area under the plasma concentration vs. time curve extrapolated to infinity (AUC_{inf} = AUC_{last} + C_{last}/λ_z, where C_{last} is the last observed plasma concentration).

Results

As shown in Table 2, below, the plasma terminal elimination half-lives for all tested peptides were determined to be in the range of 14.1 hours to 36.7 hours, while the apparent clearance values for all tested peptides were in the range of 0.00406 to 0.0146 liter/hour/kg.

Table 2. Plasma terminal elimination half-lives ($t_{1/2}$) and apparent clearance values (CL/F; l/h/kg).

Compound No.	CL/F	$t_{1/2}$ (hours)
6	0.0069	24.2
7	0.0087	25.4
9	0.0096	17.9
11	0.0118	14.1
13	0.0106	15.0
15	0.0116	18.9
28	0.0128	17.1
34	0.0148	17.2
36	0.0117	25.7
49	0.00406	36.7
55	0.0045	26.2
57	0.0078	17.9
71	0.0146	14.7

15 Example 8

Effect on acute food intake and body weight in normal Sprague Dawley rats

Compounds 6, 7, 9, 11, 13, 15, 28, 34, 36, 49, 55, 57, 59, 71 and 79 were tested for their effect on food intake and body weight in rats.

Sprague Dawley (SD) rats were obtained from Taconic A/S, Denmark. The animals arrived at least 7 days before the study start to allow acclimatization to experimental conditions. From arrival and throughout the study, the rats were housed in groups of 2 (n

= 2) in a light-, temperature- and humidity-controlled room (reversed 12/12 h light/dark cycle: lights turned off during day-time and on during night-time; temperature 20-22°C; relative humidity 50-80%). Animals had access ad libitum to food (Altromin™ 1324, Brogaarden A/S, Gentofte, Denmark) and water (domestic quality tap water with citric acid added to pH ~ 3.6) during the entire study. Stratification of rats was based on body weight (BW) on a cage-by-cage basis; mean BWs per cage were used as the basis for pairing four cages together in a group (n = 8 per group). A vehicle group and positive control group were included in each set of tests. Rats were dosed subcutaneously (sc) once in the morning immediately before turning off the lights, using a body weight-corrected dose (30 nmol/kg) of test peptide. Dosing volume was 5 ml/kg. Food intake was recorded manually at t = -24 and 0 hours (pre-dose) and at t = 24, 48, 72 and 96 hours after dosing. Body weight was measured daily.

Statistical analyses were performed using GraphPad™ Prism version 5. The measured parameters were compared using one-way ANOVA followed by Dunnett's multiple comparison tests. Differences were considered statistically significant at $p < 0.05$.

Results

As shown in Table 3, below, 48 hours after dosing, each of the tested compounds had given rise to a clear, statistically significant inhibition of food intake (vehicle-corrected, in %). This reduction in food intake was reflected by a decrease in body weight (vehicle-corrected, in %) observed on day 4 post-dosing (after 96 h) in the treated rats. (Not shown.) Normal feeding behavior was subsequently resumed.

Table 3. Acute food intake data

5	Compound No.	FI inhibition after 48 h (vehicle corrected) (%)
	6	97 ± 0.2
	7	94 ± 1.4
	9	91 ± 2.5
	11	88 ± 3.4
	13	88 ± 2.4
	15	93 ± 1.6
	28	74 ± 5.4
	34	70 ± 6.6
	36	20 ± 3.4
	49	98 ± 0.2
	55	90 ± 2.4
10	57	77 ± 8.7
	59	99 ± 0.1
	71	84 ± 5.7
	79	71 ± 5.6

Example 9Oral glucose tolerance test (OGTT) in rats

15 Male Sprague Dawley rats [CrI:CD(SD), weight range 260-280 g upon arrival] were housed in groups of 2 (n = 2) or 3 (n = 3). The animals were maintained on a 12h/12h light/dark cycle. Animals had access ad libitum to a standard chow diet and tap water (domestic quality tap water) during the initial one week acclimatization period.

20 Fasted rats (fasted for 7 hours) were employed in the test. Rats had previously been randomized into two groups, designated test group and control group, respectively. The total number of animals per group was 10 (n = 10).

A single subcutaneous injection of either vehicle (control group; vehicle: 50 mM histidine buffer, pH 7, with 200 mM mannitol) or Compound 7 in vehicle [test group; dose 30 nmol/kg body weight (2 ml/kg)] was subsequently administered to each animal. All rats
 25 then received an oral glucose bolus (2 g/kg) 16 hours after vehicle or compound administration. Blood glucose levels in tail blood samples taken 20 min. before glucose administration (time zero, baseline) and then 15 min., 30 min., 60 min., 90 min., 120 min. and 180 min after glucose administration were determined using a glucometer (GlukoSmart Swing™; MSP Bodmann GmbH, Germany).

- Baseline glucose levels were essentially the same in the test and control groups, but the elevation of glucose levels over the course of the 180 minute measurement time period was significantly lowered in the test group compared to the control group. Table 4 (below) summarizes the blood glucose AUC (area under curve) determined over the period from t = 15 min. to t = 180 min. from a plot of measured glucose levels as a function of time. The data are presented as mean $\pm$ S.E.M. The two-sided unpaired Student t-test was used for statistical comparison of the test group and the control group.

Table 4.

	Control group	Test group (Compound 7)	p-value
Glucose AUC, t = 15-180 min. (mmol/L); mean $\pm$ S.E.M.	270 $\pm$ 38	122 $\pm$ 29	p<0.0062

- 10 It is clear from the results that the tested compound significantly improves glucose tolerance in the test group compared to the control group.

Example 10

Effect of deletion at positions 21 and 22

- 15 The chemical stability assessment described in Example 6 was repeated for four pairs of compounds, each differing only by the presence or absence of an Asn-Asn (N-N) doublet at positions 21 and 22.

Table 5

Compound	Positions 21-22	pH 4	pH6	pH 7.5	pH 9
NN96	N-N	A	A	B	B
97	Δ	A	A	A	A
55	Δ	A	A	A	B
98	N-N	A	A	B	B
57	Δ	A	A	A	B
99	N-N	A	A	B	B
89	Δ	A	A	A	A
100	N-N	A	A	A	B

5 [Designations "A", "B" and "C" are the same as in Table 1. " Δ " signifies a deletion at positions 21 and 22.]

It can be seen that deletion of the Asn-Asn doublet at positions 21 and 22 tends to increase chemical stability in the neutral to alkaline pH range.

10 Activity at the human calcitonin (hCTR2) and amylin (hAMYR3) was also determined, as described in Example 3. As shown in Table 6, the deletion tends to increase the activity at both receptors, especially in compounds containing methylated amino acid residues.

Table 6

Compound	hCTR2 EC ₅₀ (nM)	hAMYR3 EC ₅₀ (nM)
NN96	0.089	0.150
97	0.10	0.13
55	0.11	0.08
98	7.2	0.38
57	0.08	0.11
99	3.8	0.59
89	0.16	0.20
100	2.2	0.53

CLAIMS

1. An amylin analogue selected from the group consisting of:

[19CD]-isoGlu-RCNTATCATQRLAHFLHRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSETP-NH₂

5 [19CD]-isoGlu-RCNTATCATQRLANFLHRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSETP-NH₂

[19CD]-isoGlu-RCGTATCATERLAHFLNRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSNTP-Hyp-NH₂

[19CD]-isoGlu-RCGTATCATERLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSNTP-Hyp-NH₂

[19CD]-isoGlu-RCGTATCATERLAHFLERSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSNTP-Hyp-NH₂

[19CD]-isoGlu-RCGTATCATERLAHFLTRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSSP-Hyp-NH₂

10 [19CD]-isoGlu-RCGTATCATERLAHFLTRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSP-Hyp-NH₂

[19CD]-isoGlu-RCGTATCATERLAHFLTRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSP-Orn-T-Hyp-NH₂

[19CD]-isoGlu-RCGTATCATERLAHFLTRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSP-Aad-T-Hyp-NH₂

15 [19CD]-isoGlu-RCGTATCATERLAHFLTRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSGP-Hyp-NH₂

[19CD]-isoGlu-RCNTATCATQRLAHFLHRF-Gly(Me)-A-Ile(Me)-LSSTEVGSETP-NH₂

[19CD]-isoGlu-RCGTATCATERLAHFLHRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSNTP-NH₂

[19CD]-isoGlu-RCNTATCATQRLAHFLHRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSNTP-NH₂

[19CD]-isoGlu-RCGTATCATERLANFLVRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSNTP-Hyp-NH₂

20 [19CD]-isoGlu-RCGTATCATERLAHFLTRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSNTP-Hyp-NH₂

[19CD]-isoGlu-RCGTATCATERLAHFLKRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSNTP-Hyp-NH₂

[19CD]-isoGlu-RCNTATCATQRLAHFLHRVSF-Gly(Me)-A-Ile(Me)-LSSTEVGSETP-NH₂

[19CD]-isoGlu-RCNTATCATQRLAHFLHRVSF-Gly(Me)-A-Ile(Me)-LSSTEVGSETP-NH₂

[19CD]-isoGlu-RCGTATCATERLAHFLVRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSETP-NH₂

- [19CD]-isoGlu-RCPTATCATDRLAHFLVRSSF-Gly(Me)-A-Ile(Me)-LSSTDVGS DTP-NH₂
- [19CD]-isoGlu-RCNTATCATQRLAHFLVRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSETP-NH₂
- [19CD]-isoGlu-RCPTATCATDRLAHFLHRSSF-Gly(Me)-A-Ile(Me)-LSSTDVGSNTP-NH₂
- [19CD]-isoGlu-RCGTATCATERLAHFLHRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSETP-NH₂
- 5 [19CD]-isoGlu-RCPTATCATDRLAHFLHRSSF-Gly(Me)-A-Ile(Me)-LSSTDVGS DTP-NH₂
- [19CD]-isoGlu-RCGTATCATERLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSSST-Hyp-NH₂
- [19CD]-isoGlu-RCGTATCATERLAHFLNRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSSST-Hyp-NH₂
- [19CD]-isoGlu-RCGTATCATERLAHFLHRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSSST-Hyp-NH₂
- [19CD]-isoGlu-RCGTATCATERLAHFLVRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSSST-Hyp-NH₂
- 10 [19CD]-isoGlu-RCGTATCATERLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVG SFT-Hyp-NH₂
- [19CD]-isoGlu-RCGTATCATERLAHFLNRSSF-Gly(Me)-A-Ile(Me)-LSSTEVG SFT-Hyp-NH₂
- [19CD]-isoGlu-RCGTATCATERLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVG S STP-NH₂
- [19CD]-isoGlu-RCGTATCATERLAHFLNRSSF-Gly(Me)-A-Ile(Me)-LSSTEVG S STP-NH₂
- [19CD]-isoGlu-RCGTATCATERLAHFLERSSF-Gly(Me)-A-Ile(Me)-LSSTEVG S STP-NH₂
- 15 [19CD]-isoGlu-RCGTATCATERLAHFLHRSSF-Gly(Me)-A-Ile(Me)-LSSTEVG S STP-NH₂
- [19CD]-isoGlu-RCGTATCATERLAHFLVRSSF-Gly(Me)-A-Ile(Me)-LSSTEVG S STP-NH₂
- [19CD]-isoGlu-RCGTATCATERLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVG S FTP-NH₂
- [19CD]-isoGlu-RCGTATCATERLAHFLNRSSF-Gly(Me)-A-Ile(Me)-LSSTEVG S FTP-NH₂
- [19CD]-isoGlu-RCGTATCATERLADFLTRSSF-Gly(Me)-A-Ile(Me)-LSSTEVG S STHyp-NH₂
- 20 [19CD]-isoGlu-RCGTATCATERLADFLTRSSF-Gly(Me)-A-Ile(Me)-LSSTEVG SFT-Hyp-NH₂
- [19CD]-isoGlu-RCGTATCATERLA-Aad-FLTRSSF-Gly(Me)-A-Ile(Me)-LSSTEVG S STHyp-NH₂

[19CD]-isoGlu-RCGTATCATERLA-Aad-FLTRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGGS-Orn-T-Hyp-NH₂

[19CD]-isoGlu-RCGTATCATERLAHFLTRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGGSST-Apr-NH₂

[19CD]-isoGlu-RCGTATCATERLAHFLTRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGGSSTP-NH₂

5 [19CD]-isoGlu-RCGTATCATERLAHFLTRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGGS-Orn-TP-NH₂

[19CD]-isoGlu-RCGTATCATERLAHFLTRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSTT-Hyp-NH₂

[19CD]-isoGlu-RCNTATCATQRLADFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGGSNT-Hyp-NH₂

[19CD]-isoGlu-RCNTATCATQRLAHFL-Aad-RSSF-Gly(Me)-A-Ile(Me)-LSSTEVGGSNT-Hyp-NH₂

10 [19CD]-isoGlu-RCGTATCATERLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGGS-Orn-T-Hyp-NH₂

[19CD]-isoGlu-RCGTATCATERLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSGT-Hyp-NH₂

[19CD]-isoGlu-RCGTATCATERLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGGS-Aad-T-Hyp-NH₂

15 [19CD]-isoGlu-RCGTATCATERLADFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGGSNT-Hyp-NH₂

[19CD]-isoGlu-RCGTATCATERLA-Aad-FLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGGSNT-Hyp-NH₂

[19CD]-isoGlu-RCNTATCATERLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGGSNT-Hyp-NH₂

[19CD]-isoGlu-RCQTATCATERLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGGSNT-Hyp-NH₂

20 [19CD]-isoGlu-RCNTATCATQRLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGGSNT-Hyp-NH₂

[19CD]-isoGlu-RCPTATCATDRLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTDVGSNT-Hyp-NH₂

[19CD]-isoGlu-RCNTATCATERLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSDT-Hyp-NH₂

[19CD]-isoGlu-RCNTATCATQRLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSNTP-NH₂

[19CD]-isoGlu-RCNTATCATERLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSNTP-NH₂

25 [19CD]-isoGlu-RCPTATCATERLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGGSNT-Hyp-NH₂

[19CD]-isoGlu-RCPTATCATQRLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSNTP-Hyp-NH₂

[19CD]-isoGlu-RCNTATCATQRLADFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSNTP-NH₂

[19CD]-isoGlu-RCNTATCATERLADFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSNTP-NH₂

[19CD]-isoGlu-RCPTATCATERLADFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSNTP-NH₂

5 [19CD]-isoGlu-RCNTATCATERLADFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSNTP-Hyp-NH₂

[19CD]-isoGlu-RCNTATCATQRLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSSST-Hyp-NH₂

[19CD]-isoGlu-RCPTATCATERLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSSST-Hyp-NH₂

[19CD]-isoGlu-RCNTATCATQRLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSGT-Hyp-NH₂

[19CD]-isoGlu-RCPTATCATERLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSGT-Hyp-NH₂

10 [19CD]-isoGlu-RCNTATCATQRLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSTT-Hyp-NH₂

[19CD]-isoGlu-RCPTATCATERLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSTT-Hyp-NH₂

[19CD]-isoGlu-RCNTATCATQRLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSAad-T-Hyp-NH₂

[19CD]-isoGlu-RCPTATCATERLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSAad-T-Hyp-NH₂

15 [19CD]-isoGlu-RCGTATCATERLAHFLQRF-Gly(Me)-A-Ile(Me)-LSSTEVGSNTP-Hyp-NH₂

[19CD]-isoGlu-RCGTATCATERLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSNTP-NH₂

[19CD]-isoGlu-RCNTATCATQRLAHFLQRF-Gly(Me)-A-Ile(Me)-LSSTEVGSGT-Hyp-NH₂

[19CD]-isoGlu-RCNTATCATQRLADFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSGTP-NH₂

20 [19CD]-isoGlu-RCNTATCATQRLADFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSGT-Hyp-NH₂

[19CD]-isoGlu-RCNTATCATQRLADFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSTTP-NH₂

[19CD]-isoGlu-RCNTATCATQRLADFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSTT-Hyp-NH₂

[19CD]-isoGlu-RCNTATCATQRLAHFLERSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSSSTP-NH₂

[19CD]-isoGlu-RCNTATCATQRLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSSSTP-NH₂

[19CD]-isoGlu-RCNTATCATQRLA-Aad-FLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSNTP-Hyp-NH₂

[19CD]-isoGlu-RCNTATCATQRLA-Aad-FLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSSST-Hyp-NH₂

[19CD]-isoGlu-RCQTATCATDRLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSNTP-Hyp-NH₂

[19CD]-isoGlu-RCQTATCATDRLA-Aad-FLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSNTP-Hyp-NH₂

[19CD]-isoGlu-RCQTATCATERLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSGTP-Hyp-NH₂

[19CD]-isoGlu-RCQTATCATDRLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSGTP-Hyp-NH₂

[19CD]-isoGlu-RCQTATCATDRLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSNTP-NH₂

[19CD]-isoGlu-RCQTATCATERLA-Aad-FLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSNTP-NH₂

[19CD]-isoGlu-RCQTATCATERLA-Aad-FLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSGTP-NH₂

[19CD]-isoGlu-RCQTATCATDRLA-Aad-FLQRF-Gly(Me)-A-Ile(Me)-LSSTEVGSNTP-Hyp-NH₂

[19CD]-isoGlu-RCQTATCATERLAHFLQRF-Gly(Me)-A-Ile(Me)-LSSTEVGSNTP-NH₂

wherein [19CD] represents [19-carboxynonadecanoyl];

or a pharmaceutically acceptable salt or solvate thereof.

2. An amylin analogue selected from the group consisting of:

[19CD]-isoGlu-RCNTATCATQRLADFLHRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSETP-NH₂;

or a pharmaceutically acceptable salt or solvate thereof.

3. The amylin analogue:

[19CD]-isoGlu-R-C()-GTAT-C()-ATERLAHFLNRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSNTP-Hyp-NH₂ (Compd. 6).

4. The amylin analogue:

[19CD]-isoGlu-R-C()-GTAT-C()-ATERLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSTNT-Hyp-NH₂ (Compd. 7).

5. The amylin analogue:

5 [19CD]-isoGlu-R-C()-GTAT-C()-ATERLAHFLTRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSGT-Hyp-NH₂ (Compd. 13).

6. The amylin analogue:

10 [19CD]-isoGlu-R-C()-GTAT-C()-ATERLAHFLHRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSTNTP-NH₂ (Compd. 15).

7. The amylin analogue:

15 [19CD]-isoGlu-R-C()-NTAT-C()-ATQRLADFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSTNT-Hyp-NH₂ (Compd. 49).

8. The amylin analogue:

[19CD]-isoGlu-R-C()-GTAT-C()-ATERLA-Aad-FLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSTNT-Hyp-NH₂ (Compd. 55).

20 9. The amylin analogue:

[19CD]-isoGlu-R-C()-NTAT-C()-ATERLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSTNT-Hyp-NH₂ (Compd. 56).

10. The amylin analogue:

25 [19CD]-isoGlu-R-C()-QTAT-C()-ATERLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSTNT-Hyp-NH₂ (Compd. 57).

11. The amylin analogue:

30 [19CD]-isoGlu-R-C()-NTAT-C()-ATERLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSDT-Hyp-NH₂ (Compd. 60).

12. The amylin analogue:

35 [19CD]-isoGlu-R-C()-QTAT-C()-ATDRLAHFLQRSSF-Gly(Me)-A-Ile(Me)-LSSTEVGSTNT-Hyp-NH₂ (Compd. 88).

13. A pharmaceutically acceptable salt or solvate of a compound according to any one of claims 3 to 12.

14. An amylin analogue, or pharmaceutically acceptable salt or solvate, according to any one of the preceding claims, comprising an intramolecular disulphide bridge formed between the cysteine residues present at positions 2 and 7 of the amino acid sequence.

15. A pharmaceutical composition comprising an amylin analogue, or pharmaceutically acceptable salt or solvate, according to any one of the preceding claims, in combination with a pharmaceutically acceptable carrier, excipient or vehicle.

16. A device comprising an amylin analogue, or pharmaceutically acceptable salt or solvate, according to any one of claims 1 to 14, for delivery of the amylin analogue to a subject, optionally further comprising an anti-diabetic agent, an anti-obesity agent, an agent for treatment of metabolic syndrome, an anti-dyslipidemia agent, an anti-hypertensive agent, a proton pump inhibitor, or an anti-inflammatory agent.

17. An amylin analogue, or pharmaceutically acceptable salt or solvate, according to any one of claims 1 to 14 for use in therapy.

18. Use of an amylin analogue, pharmaceutically acceptable salt or solvate according to any one of claims 1 to 14 in the preparation of a medicament:

(a) for treating obesity, morbid obesity, obesity prior to surgery, obesity-linked inflammation, obesity-linked gallbladder disease, obesity-induced sleep apnea and respiratory problems, degeneration of cartilage, osteoarthritis, or reproductive health complications of obesity or overweight such as infertility; or

(b) for prevention or treatment of Alzheimer's disease, diabetes, type 1 diabetes, type 2 diabetes, pre-diabetes, insulin resistance syndrome, impaired glucose tolerance (IGT), disease states associated with elevated blood glucose levels, metabolic disease including metabolic syndrome, hyperglycemia, hypertension, atherogenic dyslipidemia, hepatic steatosis ("fatty liver"), kidney failure, arteriosclerosis (e.g. atherosclerosis), macrovascular disease, microvascular disease, diabetic heart (including diabetic cardiomyopathy and heart failure as a diabetic complication) coronary heart disease, peripheral artery disease or stroke, and combinations thereof.

19. Use according to claim 18 wherein said obesity is accompanied by at least one weight-related co-morbid condition, such as diabetes (e.g. type 2 diabetes), hypertension, dyslipidemia, sleep apnea and cardiovascular disease.
- 5 20. Use according to claim 18 or claim 19 wherein said amylin analogue is for administration in conjunction with an anti-diabetic agent, an anti-obesity agent, an agent for treatment of metabolic syndrome, an anti-dyslipidemia agent, an anti-hypertensive agent, a proton pump inhibitor, or an anti-inflammatory agent.
- 10 21. Use according to claim 20 wherein:
- (a) said anti-diabetic agent is metformin, a sulfonylurea, a glinide, a dipeptidyl peptidase IV (DPP-IV) inhibitor, a glitazone, a Glucagon-like peptide-1(GLP-1) receptor agonist, an Sodium-glucose co-transporter-2 (SGLT2) inhibitor, a G-protein coupled receptor 40 (GPR40) agonist, an insulin, or an insulin analogue;
- 15 (b) said anti-obesity agent is peptide YY or an analogue thereof, neuropeptide Y (NPY) or an analogue thereof, a cannabinoid receptor 1 antagonist, a lipase inhibitor, Human proislet Peptide (HIP), a melanocortin receptor 4 agonist, a GLP-1 receptor agonist, phentermine, a melanin concentrating hormone receptor 1 antagonist, Cholecystokinin (CCK), amylin, pramlintide or leptin;
- 20 (c) said anti-hypertension agent is an angiotensin-converting enzyme inhibitor, an angiotensin II receptor blocker, a diuretic, a beta-blocker or a calcium channel blocker;
- (d) said anti-dyslipidemia agent is a statin, a fibrate, a niacin, a PCSK9 (Proprotein convertase subtilisin/kexin type 9) inhibitor or a cholesterol absorption inhibitor;
- (e) said proton pump inhibitor is an agent of the benzimidazole derivative type or of the
25 imidazopyridine derivative type; or
- (f) said anti-inflammatory agent is a steroid or corticosteroid, non-steroidal anti-inflammatory agent (NSAID), a Cyclooxygenase-II (COX II) inhibitor, 5-aminosalicylic acid or metformin.