

FACTOR XII (F12) iRNA COMPOSITIONS AND METHODS OF USE THEREOF**RELATED APPLICATIONS**

This application claims the benefit of priority to U.S. Provisional Application No. 63/232,840,
5 filed on August 13, 2021, the entire contents of which are incorporated herein by reference.

BACKGROUND OF THE INVENTION

Hereditary Angioedema (HAE) is a rare autosomal dominant disorder that causes recurrent edema and swelling of the extremities, face, larynx, upper respiratory tract, abdomen, trunk, and
10 genitals and a nonpruritic rash in one-third of patients. Untreated HAE patients experience an average of one-to-two angioedema attacks per month, but the frequency and severity of episodes can vary significantly. Edema swelling is often disfiguring and disabling, results in frequent hospitalization, and patients may require psychiatric care to treat disease-associated anxiety. Abdominal attacks can cause severe pain, nausea and vomiting, and may lead to inappropriate surgeries. Furthermore, over
15 half of HAE patients also experience life-threatening laryngeal edema during their lifetime that may require emergency tracheostomy to prevent asphyxiation. HAE affects an estimated 6,000 to 10,000 individuals of varying ethnic groups in the United States and causes significant economic harm to patients, accounting for 15,000 to 30,000 hospital visits and 20 to 100 sick days per year.

HAE results from a mutation of the C1 inhibitor (C1INH, SERPING1) gene that results in a
20 deficiency of C1INH protein. Over 250 different C1INH mutations have been demonstrated to cause an HAE clinical presentation. These C1INH mutations are typically inherited genetically, however, up to 25% of HAE cases result from *de novo* mutation of C1INH. HAE type I is caused by C1INH mutations that result in lower levels of truncated or misfolded proteins that are inefficiently secreted, and accounts for approximately 85% of HAE cases. HAE type II constitutes about 15% of cases and
25 is caused by mutations near the active site of C1INH that result in normal levels of dysfunctional C1INH protein. HAE type III, a rare third form the disease, occurs as a result of a gain-of-function mutation in coagulation factor XII (F12) (Hageman Factor).

C1 inhibitor is a serine protease inhibitor of the serpin family and a major inhibitor of proteases in the complement and contact activation pathways, as well as a minor inhibitor of
30 fibrinolytic protease plasmin. These plasma proteolytic cascades are activated during an HAE attack, generating substances that increase vascular permeability. Studies have shown that the bradykinin peptide, which activates proinflammatory signaling pathways that dilate vessels and induces chemotaxis of neutrophils, is the primary substance that enhances vascular permeability in an HAE attack by binding to the bradykinin receptor on vascular endothelial cells. The formation of
35 bradykinin in plasma generally requires interaction of three proteins, namely F12, prekallikrein and high-molecular-weight kininogen, collectively referred to as the “contact activation pathway” or the “Kallikrein-Kinin System.” Prekallikrein and high-molecular-weight kininogen circulate as a bi-

molecular complex. The contact activation pathway is initiated on binding of F12 to negatively charged surfaces (or macromolecules) which induces a conformational change in F12 resulting in active F12 (F12a). F12a cleaves prekallikrein to generate active kallikrein, which in turn reciprocally activates F12. The active kallikrein then digests high-molecular-weight kininogen to liberate
5 bradykinin.

Typically, C1INH inhibits the autoactivation of F12, the ability of F12a to activate prekallikrein, the activation of high molecular weight kininogen by kallikrein, and the feedback activation of F12 by kallikrein. Consequently, mutations causing C1INH deficiency or F12 gain-of-function result in excess production of bradykinin and onset of HAE angioedema.

10 Currently, HAE may be treated with 17α -alkylated androgens prophylactically to reduce to probability of recurrent episodes, or with disease-specific therapeutics to treat acute attacks. About 70% of individuals with HAE are treated with androgens or remain untreated, and about 30% receive therapeutics. However, androgens are unsuitable for short-term treatment of acute attacks because they take several days to become effective, and they can have significant side effects and may affect
15 growth and development adversely. As a result, androgens are used only for long-term prophylaxis and are typically not administered to pregnant women or children. Furthermore, current therapeutics used to treat acute attacks must be administered intravenously numerous times per week or may cause side-effects that require drug administration and subsequent patient monitoring in a hospital, thereby limiting their regular prophylactic use to manage the disease long-term.

20 Therefore, in the absence of regimens which be administered safely, effectively and by more convenient routes and regimens to treat acute angioedema attacks and prophylactically manage recurrent attacks in a large proportion of patients, including pregnant women and children, there is a need for alternative therapies for subjects suffering from HAE.

25 SUMMARY OF THE INVENTION

The present invention provides iRNA compositions which effect the RNA-induced silencing complex (RISC)-mediated cleavage of RNA transcripts of a gene encoding Factor XII (F12). The F12 gene may be within a cell, *e.g.*, a cell within a subject, such as a human subject. The present invention also provides methods of using the iRNA compositions of the invention for inhibiting the
30 expression of an F12 gene and/or for treating a subject who would benefit from inhibiting or reducing the expression of an F12 gene, *e.g.*, a subject suffering or prone to suffering from an F12-associated disorder, *e.g.*, hereditary angioedema (HAE) (such as hereditary angioedema type I; hereditary angioedema type II; hereditary angioedema type III; or any other hereditary angioedema caused by elevated levels of bradykinin); prekallikrein deficiency, hypertension, *e.g.*, malignant essential
35 hypertension, end stage renal disease, Fletcher Factor Deficiency, thromboembolic disease, inflammatory disease, or Alzheimer's Disease.

Accordingly, in an aspect, the invention provides a double stranded ribonucleic acid (dsRNA) agent for inhibiting expression of Factor XII (F12) in a cell, wherein the dsRNA agent comprises a sense strand and an antisense strand forming a double stranded region, wherein the sense strand comprises at least 15, *e.g.*, 15, 16, 17, 18, 19, 20, or 21, contiguous nucleotides differing by no more than 0, 1, 2, or 3 nucleotides from the nucleotide sequence of SEQ ID NO:1 and the antisense strand comprises at least 15, *e.g.*, 15, 16, 17, 18, 19, 20, 21, 22, or 23, contiguous nucleotides differing by no more than 1, 2, or 3 nucleotides from the corresponding portion of the nucleotide sequence of SEQ ID NO:2.

In another aspect, the present invention provides a double stranded ribonucleic acid (dsRNA) for inhibiting expression of Factor XII (F12) in a cell, wherein said dsRNA comprises a sense strand and an antisense strand forming a double stranded region, wherein the antisense strand comprises a region of complementarity to an mRNA encoding F12, and wherein the region of complementarity comprises at least 15, *e.g.*, 15, 16, 17, 18, 19, 20, 21, 22, or 23, contiguous nucleotides differing by no more than 0, 1, 2, or 3 nucleotides from any one of the antisense nucleotide sequences in any one of Tables 2-3.

In one embodiment, the dsRNA agent comprises a sense strand comprising a contiguous nucleotide sequence which has at least 85%, *e.g.*, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100%, nucleotide sequence identity over its entire length to any one of the nucleotide sequences of the sense strands in any one of Tables 2-3 and an antisense strand comprising a contiguous nucleotide sequence which has at least 85%, *e.g.*, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100%, nucleotide sequence identity over its entire length to any one of the nucleotide sequences of the antisense strands in any one of Tables 2-3.

In one embodiment, the dsRNA agent comprises a sense strand comprising at least 15, *e.g.*, 15, 16, 17, 18, 19, 20, or 21, contiguous nucleotides differing by no more than three nucleotides from any one of the nucleotide sequences of the sense strands in any one of Tables 2-3 and an antisense strand comprising at least 15, *e.g.*, 15, 16, 17, 18, 19, 20, 21, 22, or 23, contiguous nucleotides differing by no more than three nucleotides from any one of the nucleotide sequences of the antisense strands in any one of Tables 2-3.

In another embodiment, the dsRNA agent comprises a sense strand comprising at least 15, *e.g.*, 15, 16, 17, 18, 19, 20, or 21, contiguous nucleotides differing by no more than two nucleotides from any one of the nucleotide sequences of the sense strands in any one of Tables 2-3 and an antisense strand comprising at least 15, *e.g.*, 15, 16, 17, 18, 19, 20, 21, 22, or 23, contiguous nucleotides differing by no more than two nucleotides from any one of the nucleotide sequences of the antisense strands in any one of Tables 2-3.

In another embodiment, the dsRNA agent comprises a sense strand comprising at least 15, *e.g.*, 15, 16, 17, 18, 19, 20, or 21, contiguous nucleotides differing by no more than one nucleotide from any one of the nucleotide sequences of the sense strands in any one of Tables 2-3 and an

antisense strand comprising at least 15, *e.g.*, 15, 16, 17, 18, 19, 20, 21, 22, or 23, contiguous nucleotides differing by no more than one nucleotide from any one of the nucleotide sequences of the antisense strands in any one of Tables 2-3.

5 In another embodiment, the dsRNA agent comprises a sense strand comprising a nucleotide sequence selected from the group consisting of any one of the nucleotide sequences of the sense strands in any one of Tables 2-3 and an antisense strand comprising a nucleotide sequence selected from the group consisting of any one of the nucleotide sequences of the antisense strands in any one of Tables 2-3.

10 In one aspect, the present invention provides a double stranded ribonucleic acid (dsRNA) agent for inhibiting expression of Factor XII (F12) in a cell, wherein said dsRNA comprises a sense strand and an antisense strand forming a double stranded region, wherein the sense strand comprises at least 15, *e.g.*, 15, 16, 17, 18, 19, 20, or 21, contiguous nucleotides differing by no more than 0, 1, 2, or 3 nucleotides from any one of the nucleotide sequence of nucleotides 75-97, 76-98, 79-101, 115-137, 117-139, 123-145, 132-154, 135-157, 137-159, 144-166, 201-223, or 1462-1484 of SEQ ID NO: 1, and the antisense strand comprises at least 15, *e.g.*, 15, 16, 17, 18, 19, 20, 21, 22, or 23, contiguous nucleotides differing by no more than 0, 1, 2, or 3 nucleotides from the corresponding nucleotide sequence of SEQ ID NO:2.

20 In one embodiment, the sense strand comprises at least 15, *e.g.*, 15, 16, 17, 18, 19, 20, or 21, contiguous nucleotides differing by no more than 0, 1, 2, or 3 nucleotides from any one of the nucleotide sequence of nucleotides 135-157, 144-166, or 201-223 of SEQ ID NO: 1, and the antisense strand comprises at least 15, *e.g.*, 15, 16, 17, 18, 19, 20, 21, 22, or 23, contiguous nucleotides differing by no more than 0, 1, 2, or 3 nucleotides from the corresponding nucleotide sequence of SEQ ID NO:2.

25 In one embodiment, the antisense strand comprises at least 15, *e.g.*, 15, 16, 17, 18, 19, 20, 21, 22, or 23, contiguous nucleotides differing by no more than 0, 1, 2, or 3 nucleotides from any one of the antisense strand nucleotide sequences of a duplex selected from the group consisting of AD-1534528, AD-1534479, AD-1534582, AD-1534611, AD-1534506, AD-1631309, AD-1534471, AD-1534555, AD-1534678, AD-1534585, AD-1534502, and AD-1534491.

30 In one embodiment, the antisense strand comprises at least 15, *e.g.*, 15, 16, 17, 18, 19, 20, 21, 22, or 23, contiguous nucleotides differing by no more than 0, 1, 2, or 3 nucleotides from any one of the antisense strand nucleotide sequences of a duplex selected from the group consisting of AD-1534528, AD-1534479, and AD-1534582.

35 In one embodiment, the sense strand comprises at least 15, *e.g.*, 15, 16, 17, 18, 19, 20, or 21, contiguous nucleotides differing by no more than three nucleotides from any one of the sense strand nucleotide sequences of a duplex selected from the group consisting of AD-1534528, AD-1534479, and AD-1534582 and the antisense strand comprises at least 15, *e.g.*, 15, 16, 17, 18, 19, 20, 21, 22, or 23, contiguous nucleotides differing by no more than three nucleotides from any one of the antisense

strand nucleotide sequences of a duplex selected from the group consisting of AD-1534528, AD-1534479, and AD-1534582.

In one embodiment, the sense strand comprises at least 15, *e.g.*, 15, 16, 17, 18, 19, 20, or 21, contiguous nucleotides differing by no more than two nucleotides from any one of the sense strand nucleotide sequences of a duplex selected from the group consisting of AD-1534528, AD-1534479, and AD-1534582 and the antisense strand comprises at least 15, *e.g.*, 15, 16, 17, 18, 19, 20, 21, 22, or 23, contiguous nucleotides differing by no more than two nucleotides from any one of the antisense strand nucleotide sequences of a duplex selected from the group consisting of AD-1534528, AD-1534479, and AD-1534582.

In one embodiment, the sense strand comprises at least 15, *e.g.*, 15, 16, 17, 18, 19, 20, or 21, contiguous nucleotides differing by no more than one nucleotide from any one of the sense strand nucleotide sequences of a duplex selected from the group consisting of AD-1534528, AD-1534479, and AD-1534582 and the antisense strand comprises at least 15, *e.g.*, 15, 16, 17, 18, 19, 20, 21, 22, or 23, contiguous nucleotides differing by no more than one nucleotide from any one of the antisense strand nucleotide sequences of a duplex selected from the group consisting of AD-1534528, AD-1534479, and AD-1534582.

In one embodiment, the sense strand comprises any one of the sense strand nucleotide sequences of a duplex selected from the group consisting of AD-1534528, AD-1534479, and AD-1534582 and the antisense strand comprises any one of the antisense strand nucleotide sequences of a duplex selected from the group consisting of AD-1534528, AD-1534479, and AD-1534582.

In one embodiment, the dsRNA agent comprises at least one modified nucleotide.

In one embodiment, substantially all of the nucleotides of the sense strand are modified nucleotides; substantially all of the nucleotides of the antisense strand are modified nucleotides; or substantially all of the nucleotides of the sense strand and substantially all of the nucleotides of the antisense strand are modified nucleotides.

In one embodiment, all of the nucleotides of the sense strand are modified nucleotides; all of the nucleotides of the antisense strand are modified nucleotides; or all of the nucleotides of the sense strand and all of the nucleotides of the antisense strand are modified nucleotides.

In one embodiment, at least one of the modified nucleotides is selected from the group consisting of a deoxy-nucleotide, a 3'-terminal deoxythymidine (dT) nucleotide, a 2'-O-methyl modified nucleotide, a 2'-fluoro modified nucleotide, a 2'-deoxy-modified nucleotide, a 2'-5'-linked ribonucleotide (3'-RNA), a locked nucleotide, an unlocked nucleotide, a conformationally restricted nucleotide, a constrained ethyl nucleotide, an abasic nucleotide, a 2'-amino-modified nucleotide, a 2'-O-allyl-modified nucleotide, 2'-C-alkyl-modified nucleotide, 2'-hydroxly-modified nucleotide, a 2'-methoxyethyl modified nucleotide, a 2'-O-alkyl-modified nucleotide, a morpholino nucleotide, a phosphoramidate, a non-natural base comprising nucleotide, a tetrahydropyran modified nucleotide, a 1,5-anhydrohexitol modified nucleotide, a cyclohexenyl modified nucleotide, a nucleotide

comprising a phosphorothioate group, a nucleotide comprising a methylphosphonate group, a vinyl phosphonate nucleotide, a nucleotide comprising a 5'-phosphate, a nucleotide comprising a 5'-phosphate mimic, a thermally destabilizing nucleotide, a glycol modified nucleotide (GNA), a glycol nucleic acid S-Isomer (S-GNA), a nucleotide comprising a 2' phosphate, and a 2-O-(N-

5 methylacetamide) modified nucleotide; and combinations thereof.

In one embodiment, at least one of the modified nucleotides is selected from the group consisting of LNA, HNA, CeNA, 2'-methoxyethyl, 2'-O-alkyl, 2'-O-allyl, 2'-C-allyl, 2'-fluoro, 2'-deoxy, 2'-hydroxyl, and glycol; and combinations thereof.

10 In one embodiment, at least one of the modified nucleotides is selected from the group consisting of a deoxy-nucleotide, a 2'-O-methyl modified nucleotide, a 2'-fluoro modified nucleotide, a 2'-deoxy-modified nucleotide, a glycol modified nucleotide (GNA), *e.g.*, Ggn, Cgn, Tgn, or Agn, a nucleotide with a 2' phosphate, *e.g.*, G2p, C2p, A2p or U2p, and a nucleotide comprising a phosphorothioate group, and combinations thereof.

15 In another embodiment, at least one of the modified nucleotides is a nucleotide with a thermally destabilizing nucleotide modification.

In one embodiment, the thermally destabilizing nucleotide modification is selected from the group consisting of an abasic modification; a mismatch with the opposing nucleotide in the duplex; a destabilizing sugar modification, a 2'-deoxy modification, an acyclic nucleotide, an unlocked nucleic acid (UNA), a glycerol nucleic acid (GNA); and 2'-5'-linked ribonucleotides ("3'-RNA").

20 In some embodiments, the modified nucleotide comprises a short sequence of 3'-terminal deoxythymidine nucleotides (dT).

In some embodiments, the dsRNA agent further comprises at least one phosphorothioate internucleotide linkage. In some embodiments, the dsRNA agent comprises 6-8 phosphorothioate internucleotide linkages. In one embodiment, the phosphorothioate or methylphosphonate internucleotide linkage is at the 3'-terminus of one strand. Optionally, the strand is the antisense strand. In another embodiment, the strand is the sense strand. In a related embodiment, the phosphorothioate or methylphosphonate internucleotide linkage is at the 5'-terminus of one strand. Optionally, the strand is the antisense strand. In another embodiment, the strand is the sense strand. In another embodiment, the phosphorothioate or methylphosphonate internucleotide linkage is at the 30 both the 5'- and 3'-terminus of one strand. Optionally, the strand is the antisense strand. In another embodiment, the strand is the sense strand.

The double stranded region may be 19-30 nucleotide pairs in length; 19-25 nucleotide pairs in length; 19-23 nucleotide pairs in length; 23-27 nucleotide pairs in length; or 21-23 nucleotide pairs in length.

35 In one embodiment, each strand is independently no more than 30 nucleotides in length.

In one embodiment, the sense strand is 21 nucleotides in length and the antisense strand is 23 nucleotides in length.

The region of complementarity may be at least 17 nucleotides in length; between 19 and 23 nucleotides in length; or 19 nucleotides in length.

In one embodiment, at least one strand comprises a 3' overhang of at least 1 nucleotide. In another embodiment, at least one strand comprises a 3' overhang of at least 2 nucleotides.


5 In one embodiment, the dsRNA agent further comprises a ligand.

In one embodiment, the ligand is conjugated to the 3' end of the sense strand of the dsRNA agent.


In one embodiment, the ligand is an N-acetylgalactosamine (GalNAc) derivative.

10 In one embodiment, the ligand is one or more GalNAc derivatives attached through a monovalent, bivalent, or trivalent branched linker.

In one embodiment, the ligand is



In one embodiment, the dsRNA agent is conjugated to the ligand as shown in the following schematic



15 and, wherein X is O or S.

In one embodiment, the X is O.

In one embodiment, the dsRNA agent further comprises at least one phosphorothioate or methylphosphonate internucleotide linkage.

20 In one embodiment, the phosphorothioate or methylphosphonate internucleotide linkage is at the 3'-terminus of one strand, *e.g.*, the antisense strand or the sense strand.

In another embodiment, the phosphorothioate or methylphosphonate internucleotide linkage is at the 5'-terminus of one strand, *e.g.*, the antisense strand or the sense strand.

In one embodiment, the phosphorothioate or methylphosphonate internucleotide linkage is at the both the 5'- and 3'-terminus of one strand. In one embodiment, the strand is the antisense strand.

5 In one embodiment, the base pair at the 1 position of the 5'-end of the antisense strand of the duplex is an AU base pair.

The present invention also provides cells containing any of the dsRNA agents of the invention and pharmaceutical compositions comprising any of the dsRNA agents of the invention.

10 The pharmaceutical composition of the invention may include dsRNA agent in an unbuffered solution, *e.g.*, saline or water, or the pharmaceutical composition of the invention may include the dsRNA agent in a buffer solution, *e.g.*, a buffer solution comprising acetate, citrate, prolamine, carbonate, or phosphate or any combination thereof; or phosphate buffered saline (PBS).

In one aspect, the present invention provides a method of inhibiting expression of a Factor XII (F12) gene in a cell. The method includes contacting the cell with any of the dsRNAs of the invention or any of the pharmaceutical compositions of the invention, thereby inhibiting expression
15 of the F12 gene in the cell.

In one embodiment, the cell is within a subject, *e.g.*, a human subject, *e.g.*, a subject having a Factor XII (F12)-associated disorder, such as heredity angioedema (HAE) (*e.g.*, hereditary angioedema type I; hereditary angioedema type II; hereditary angioedema type III; or any other
20 hereditary angioedema caused by elevated levels of bradykinin); prekallikrein deficiency, hypertension, *e.g.*, malignant essential hypertension, end stage renal disease, Fletcher Factor Deficiency, thromboembolic disease, inflammatory disease, or Alzheimer's Disease.

In certain embodiments, the F12 expression is inhibited by at least about 30%, 40%, 50%, 60%, 70%, 80%, 90%, or 95%. In one embodiment, inhibiting expression of F12 decreases F12
25 protein level in serum of the subject by at least 30%, 40%, 50%, 60%, 70%, 80%, 90%, or 95%.

In one aspect, the present invention provides a method of treating a subject having a disorder that would benefit from reduction in Factor XII (F12) expression. The method includes administering to the subject a therapeutically effective amount of any of the dsRNAs of the invention or any of the pharmaceutical compositions of the invention, thereby treating the subject having the
30 disorder that would benefit from reduction in F12 expression.

In another aspect, the present invention provides a method of preventing at least one symptom in a subject having a disorder that would benefit from reduction in Factor XII (F12) expression. The method includes administering to the subject a prophylactically effective amount of any of the dsRNAs of the invention or any of the pharmaceutical compositions of the invention,
35 thereby preventing at least one symptom in the subject having the disorder that would benefit from reduction in F12 expression.

In certain embodiments, the disorder is a Factor XII (F12)-associated disorder.

In some embodiments, the F12-associated disorder is selected from the group consisting of heredity angioedema (HAE) (such as hereditary angioedema type I; hereditary angioedema type II; hereditary angioedema type III; or any other hereditary angioedema caused by elevated levels of bradykinin); prekallikrein deficiency, hypertension, *e.g.*, malignant essential hypertension, end stage renal disease, Fletcher Factor Deficiency, thromboembolic disease, inflammatory disease, and Alzheimer's Disease.

In some embodiments, the F12-associated disorder is HAE.

In other embodiments, the F12-associated disorder is prekallikrein deficiency.

In other embodiments, the F12-associated disorder is hypertension, *e.g.*, malignant essential hypertension.

In other embodiments, the F12-associated disorder is end stage renal disease.

In other embodiments, the F12-associated disorder is Fletcher Factor Deficiency.

In another embodiment, an F12-associated disorder is thromboembolic disease.

In another embodiment, an F12-associated disorder is inflammatory disease.

In another embodiment, an F12-associated disorder is Alzheimer's Disease.

In certain embodiments, administration of the dsRNA agent to the subject causes a decrease F12 protein accumulation in the subject.

In certain embodiments, administration of the dsRNA agent to the subject causes a decrease in one or more of elevated bradykinin, prekallikrein deficiency, malignant essential hypertension, hypertension, end stage renal disease, Fletcher Factor Deficiency, inflammation, thrombosis, edema swelling of the extremities, face, larynx, upper respiratory tract, abdomen, trunk, genitals, or prodrome; laryngeal swelling; nonpruritic rash; nausea; vomiting; or abdominal pain.

In a further aspect, the present invention also provides methods of inhibiting the expression of F12 in a subject. The methods include administering to the subject a therapeutically effective amount of any of the dsRNAs provided herein, thereby inhibiting the expression of F12 in the subject.

In one embodiment, the subject is human.

In one embodiment, administration of the dsRNA agent to the subject causes a decrease in F12 protein accumulation in the subject.

In one embodiment, the dsRNA agent is administered to the subject at a dose of about 0.01 mg/kg to about 50 mg/kg.

In one embodiment, the dsRNA agent is administered to the subject subcutaneously.

In one embodiment, the methods of the invention further include determining the level of F12 in a sample(s) from the subject.

In one embodiment, the level of F12 in the subject sample(s) is an F12 protein level in a blood or serum or liver tissue sample(s).

In some embodiments, the methods of the invention further include determining the level of bradykinin; prekallikrein, or blood pressure in the subject.

In certain embodiments, the methods of the invention further comprise administering to the subject an additional therapeutic agent.

5 In certain embodiments, the additional therapeutic agent is selected from the group consisting of an androgen, such as danazol or oxandrolone, Berinert®, Cinryze™, Rhuconest®, Ecallantide, Firazyr®, Kalbitor®, or a combination of any of the foregoing.

The present invention also provides kits comprising any of the dsRNAs of the invention or any of the pharmaceutical compositions of the invention, and optionally, instructions for use. In one
10 embodiment, the invention provides a kit for performing a method of inhibiting expression of F12 gene in a cell by contacting a cell with a double stranded RNAi agent of the invention in an amount effective to inhibit expression of the F12 in the cell. The kit comprises an RNAi agent and instructions for use and, optionally, means for administering the RNAi agent to a subject.

The present invention also provides vials comprising the dsRNA agent of the invention or the
15 pharmaceutical composition of the invention. The present invention further provides syringes comprising the dsRNA agent of the invention or the pharmaceutical composition of the invention.

The present invention further provides an RNA-induced silencing complex (RISC) comprising an antisense strand of any of the dsRNA agents of the invention.

In another embodiment, the RNAi agent is a pharmaceutically acceptable salt thereof.
20 “Pharmaceutically acceptable salts” of each of RNAi agents herein include, but are not limited to, a sodium salt, a calcium salt, a lithium salt, a potassium salt, an ammonium salt, a magnesium salt, a mixtures thereof. One skilled in the art will appreciate that the RNAi agent, when provided as a polycationic salt having one cation per free acid group of the optionally modified phosphodiester backbone and/or any other acidic modifications (*e.g.*, 5'-terminal phosphonate groups). For example,
25 an oligonucleotide of “n” nucleotides in length contains n-1 optionally modified phosphodiesters, so that an oligonucleotide of 21 nt in length may be provided as a salt having up to 20 cations (*e.g.*, 20 sodium cations). Similarly, an RNAi agent having a sense strand of 21 nt in length and an antisense strand of 23 nt in length may be provided as a salt having up to 42 cations (*e.g.*, 42 sodium cations). In the preceding example, where the RNAi agent also includes a 5'-terminal phosphate or a 5'-
30 terminal vinylphosphonate group, the RNAi agent may be provided as a salt having up to 44 cations (*e.g.*, 44 sodium cations).

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 is a graph depicting the percent of human Factor XII protein remaining in the serum
35 of humanized Factor XII mice administered a single 0.2. g/kg dose of the indicated duplexes at weekly intervals post-dose. The percent of human Factor XII protein remaining is shown relative to individual pre=dose levels of human Factor XII.

Figure 2 is a graph depicting the percent change of FXII protein levels compared to the levels of Factor XII on the indicated dosing day in the serum of *Cynomolgus* monkeys subcutaneously administered a single 1 mg/kg of the indicated duplexes.

DETAILED DESCRIPTION OF THE INVENTION

The present invention provides iRNA compositions which effect the RNA-induced silencing complex (RISC)-mediated cleavage of RNA transcripts of a Factor XII (F12) gene. The gene may be within a cell, *e.g.*, a cell within a subject, such as a human. The use of these iRNAs enables the targeted degradation of mRNAs of the corresponding gene (F12) in mammals.

The iRNAs of the invention have been designed to target the human Factor XII (F12) gene, including portions of the gene that are conserved in the F12 orthologs of other mammalian species. Without intending to be limited by theory, it is believed that a combination or sub-combination of the foregoing properties and the specific target sites or the specific modifications in these iRNAs confer to the iRNAs of the invention improved efficacy, stability, potency, durability, and safety.

Accordingly, the present invention provides methods for treating and preventing a Factor XII (F12)-associated disorder, *e.g.*, hereditary angioedema (HAE) (*e.g.*, hereditary angioedema type I; hereditary angioedema type II; hereditary angioedema type III; or any other hereditary angioedema caused by elevated levels of bradykinin); prekallikrein deficiency, hypertension, *e.g.*, malignant essential hypertension, end stage renal disease, Fletcher Factor Deficiency, , thromboembolic disease, inflammatory disease, or Alzheimer's Disease using iRNA compositions which effect the RNA-induced silencing complex (RISC)-mediated cleavage of RNA transcripts of an F12 gene.

The iRNAs of the invention include an RNA strand (the antisense strand) having a region which is up to about 30 nucleotides or less in length, *e.g.*, 19-30, 19-29, 19-28, 19-27, 19-26, 19-25, 19-24, 19-23, 19-22, 19-21, 19-20, 20-30, 20-29, 20-28, 20-27, 20-26, 20-25, 20-24, 20-23, 20-22, 20-21, 21-30, 21-29, 21-28, 21-27, 21-26, 21-25, 21-24, 21-23, or 21-22 nucleotides in length, which region is substantially complementary to at least part of an mRNA transcript of an F12 gene.

In certain embodiments, one or both of the strands of the double stranded RNAi agents of the invention is up to 66 nucleotides in length, *e.g.*, 36-66, 26-36, 25-36, 31-60, 22-43, 27-53 nucleotides in length, with a region of at least 19 contiguous nucleotides that is substantially complementary to at least a part of an mRNA transcript of an F12 gene. In some embodiments, such iRNA agents having longer length antisense strands may, for example, include a second RNA strand (the sense strand) of 20-60 nucleotides in length wherein the sense and antisense strands form a duplex of 18-30 contiguous nucleotides.

The use of iRNAs of the invention enables the targeted degradation of mRNAs of the corresponding gene (F12 gene) in mammals. Using *in vitro* assays, the present inventors have demonstrated that iRNAs targeting an F12 gene can potentially mediate RNAi, resulting in significant inhibition of expression of an F12 gene. Thus, methods and compositions including these iRNAs are

useful for treating a subject having an F12-associated disorder, *e.g.*, hereditary angioedema (HAE) (*e.g.*, hereditary angioedema type I; hereditary angioedema type II; hereditary angioedema type III; or any other hereditary angioedema caused by elevated levels of bradykinin); prekallikrein deficiency, hypertension, *e.g.*, malignant essential hypertension, end stage renal disease, Fletcher Factor Deficiency, thromboembolic disease, inflammatory disease, or Alzheimer's Disease.

Accordingly, the present invention provides methods and combination therapies for treating a subject having a disorder that would benefit from inhibiting or reducing the expression of an F12 gene, *e.g.*, a Factor XII (F12)-associated disorder, such as hereditary angioedema (HAE) (*e.g.*, hereditary angioedema type I; hereditary angioedema type II; hereditary angioedema type III; or any other hereditary angioedema caused by elevated levels of bradykinin); prekallikrein deficiency, hypertension, *e.g.*, malignant essential hypertension, end stage renal disease, Fletcher Factor Deficiency, thromboembolic disease, inflammatory disease, or Alzheimer's Disease using iRNA compositions which effect the RNA-induced silencing complex (RISC)-mediated cleavage of RNA transcripts of an F12 gene.

The present invention also provides methods for preventing at least one symptom in a subject having a disorder that would benefit from inhibiting or reducing the expression of an F12 gene, *e.g.*, a Factor XII (F12)-associated disorder, such as hereditary angioedema (HAE) (*e.g.*, hereditary angioedema type I; hereditary angioedema type II; hereditary angioedema type III; or any other hereditary angioedema caused by elevated levels of bradykinin); prekallikrein deficiency, hypertension, *e.g.*, malignant essential hypertension, end stage renal disease, Fletcher Factor Deficiency, thromboembolic disease, inflammatory disease, or Alzheimer's Disease.

The following detailed description discloses how to make and use compositions containing iRNAs to inhibit the expression of an F12 gene as well as compositions, uses, and methods for treating subjects that would benefit from inhibition and/or reduction of the expression of an F12 gene, *e.g.*, subjects susceptible to or diagnosed with an F12-associated disorder.

I. Definitions

In order that the present invention may be more readily understood, certain terms are first defined. In addition, it should be noted that whenever a value or range of values of a parameter are recited, it is intended that values and ranges intermediate to the recited values are also intended to be part of this invention.

The articles "a" and "an" are used herein to refer to one or to more than one (*i.e.*, to at least one) of the grammatical object of the article. By way of example, "an element" means one element or more than one element, *e.g.*, a plurality of elements.

The term "including" is used herein to mean, and is used interchangeably with, the phrase "including but not limited to".

The term "or" is used herein to mean, and is used interchangeably with, the term "and/or," unless context clearly indicates otherwise. For example, "sense strand or antisense strand" is understood as "sense strand or antisense strand or sense strand and antisense strand."

The term "about" is used herein to mean within the typical ranges of tolerances in the art. For example, "about" can be understood as about 2 standard deviations from the mean. In certain embodiments, about means $\pm 10\%$. In certain embodiments, about means $\pm 5\%$. When about is present before a series of numbers or a range, it is understood that "about" can modify each of the numbers in the series or range.

The term "at least", "no less than", or "or more" prior to a number or series of numbers is understood to include the number adjacent to the term "at least", and all subsequent numbers or integers that could logically be included, as clear from context. For example, the number of nucleotides in a nucleic acid molecule must be an integer. For example, "at least 19 nucleotides of a 21 nucleotide nucleic acid molecule" means that 19, 20, or 21 nucleotides have the indicated property. When at least is present before a series of numbers or a range, it is understood that "at least" can modify each of the numbers in the series or range.

As used herein, "no more than" or "or less" is understood as the value adjacent to the phrase and logical lower values or integers, as logical from context, to zero. For example, a duplex with an overhang of "no more than 2 nucleotides" has a 2, 1, or 0 nucleotide overhang. When "no more than" is present before a series of numbers or a range, it is understood that "no more than" can modify each of the numbers in the series or range. As used herein, ranges include both the upper and lower limit.

As used herein, methods of detection can include determination that the amount of analyte present is below the level of detection of the method.

In the event of a conflict between an indicated target site and the nucleotide sequence for a sense or antisense strand, the indicated sequence takes precedence.

In the event of a conflict between a sequence and its indicated site on a transcript or other sequence, the nucleotide sequence recited in the specification takes precedence.

As used herein, "Factor XII (Hageman Factor)," used interchangeably with the terms "coagulation factor XII," "FXII," "F12," active F12," and "F12a," refers to the naturally occurring gene that encodes the zymogen form of F12a. F12 a is an enzyme (EC 3.4.21.38) of the serine protease (or serine endopeptidase) class that cleaves prekallikrein to form kallikrein, which subsequently releases bradykinin from high-molecular weight kininogen and activates F12.

The sequence of a human F12 mRNA transcript can be found at, for example, GenBank Accession No. GI: 1813757520 (NM_000505.4; SEQ ID NO:1; reverse complement, SEQ ID NO: 2). The sequence of mouse F12 mRNA can be found at, for example, GenBank Accession No. GI: 805299477 (NM_021489.3; SEQ ID NO:3; reverse complement, SEQ ID NO:4). The sequence of rat F12 mRNA can be found at, for example, GenBank Accession No. GI: 62078740 (NM_001014006.1; SEQ ID NO:5; reverse complement, SEQ ID NO: 6). The sequence of Macaca

fascicularis F12 mRNA can be found at, for example, GenBank Accession No. GI: 982255999 (XM_015452310.1; SEQ ID NO:7; reverse complement, SEQ ID NO: 8). The sequence of *Macaca mulatta* F12 mRNA can be found at, for example, GenBank Accession No. GI: 1622948449 (XM_028849951.1; SEQ ID NO:9; reverse complement, SEQ ID NO:10).

5 Additional examples of F12 mRNA sequences are readily available through publicly available databases, *e.g.*, GenBank, UniProt, OMIM, and the *Macaca* genome project web site.

Further information on F12 can be found, for example, at
www.ncbi.nlm.nih.gov/gene/?term=F12.

10 The entire contents of each of the foregoing GenBank Accession numbers and the Gene database numbers are incorporated herein by reference as of the date of filing this application.

The term F12, as used herein, also refers to variations of the F12 gene including variants provided in the SNP database. Numerous sequence variations within the F12 gene have been identified and may be found at, for example, NCBI dbSNP and UniProt (see, *e.g.*,
www.ncbi.nlm.nih.gov/snp/?term=F12, the entire contents of which is incorporated herein by
15 reference as of the date of filing this application.

As used herein, "target sequence" refers to a contiguous portion of the nucleotide sequence of an mRNA molecule formed during the transcription of an F12 gene, including mRNA that is a product of RNA processing of a primary transcription product. In one embodiment, the target portion of the sequence will be at least long enough to serve as a substrate for iRNA-directed cleavage at or
20 near that portion of the nucleotide sequence of an mRNA molecule formed during the transcription of an F12 gene.

The target sequence may be from about 19-36 nucleotides in length, *e.g.*, about 19-30 nucleotides in length. For example, the target sequence can be about 19-30 nucleotides, 19-30, 19-29, 19-28, 19-27, 19-26, 19-25, 19-24, 19-23, 19-22, 19-21, 19-20, 20-30, 20-29, 20-28, 20-27, 20-26, 20-
25 25, 20-24, 20-23, 20-22, 20-21, 21-30, 21-29, 21-28, 21-27, 21-26, 21-25, 21-24, 21-23, or 21-22 nucleotides in length. In certain embodiments, the target sequence is 19-23 nucleotides in length, optionally 21-23 nucleotides in length. Ranges and lengths intermediate to the above recited ranges and lengths are also contemplated to be part of the disclosure.

As used herein, the term "strand comprising a sequence" refers to an oligonucleotide
30 comprising a chain of nucleotides that is described by the sequence referred to using the standard nucleotide nomenclature.

"G," "C," "A," "T," and "U" each generally stand for a nucleotide that contains guanine, cytosine, adenine, thymidine, and uracil as a base, respectively. However, it will be understood that the term "ribonucleotide" or "nucleotide" can also refer to a modified nucleotide, as further detailed
35 below, or a surrogate replacement moiety (see, *e.g.*, Table 1). The skilled person is well aware that guanine, cytosine, adenine, and uracil can be replaced by other moieties without substantially altering the base pairing properties of an oligonucleotide comprising a nucleotide bearing such replacement

moiety. For example, without limitation, a nucleotide comprising inosine as its base can base pair with nucleotides containing adenine, cytosine, or uracil. Hence, nucleotides containing uracil, guanine, or adenine can be replaced in the nucleotide sequences of dsRNA featured in the invention by a nucleotide containing, for example, inosine. In another example, adenine and cytosine anywhere
5 in the oligonucleotide can be replaced with guanine and uracil, respectively to form G-U Wobble base pairing with the target mRNA. Sequences containing such replacement moieties are suitable for the compositions and methods featured in the invention.

The terms “iRNA”, “RNAi agent,” “iRNA agent,” “RNA interference agent” as used interchangeably herein, refer to an agent that contains RNA as that term is defined herein, and which
10 mediates the targeted cleavage of an RNA transcript *via* an RNA-induced silencing complex (RISC) pathway. iRNA directs the sequence-specific degradation of mRNA through a process known as RNA interference (RNAi). The iRNA modulates, *e.g.*, inhibits, the expression of an F12 gene in a cell, *e.g.*, a liver cell within a subject, such as a mammalian subject.

In one embodiment, an RNAi agent of the invention includes a single stranded RNA that
15 interacts with a target RNA sequence, *e.g.*, an F12 target mRNA sequence, to direct the cleavage of the target RNA. Without wishing to be bound by theory it is believed that long double stranded RNA introduced into cells is broken down into siRNA by a Type III endonuclease known as Dicer (Sharp *et al.* (2001) *Genes Dev.* 15:485). Dicer, a ribonuclease-III-like enzyme, processes the dsRNA into 19-23 base pair short interfering RNAs with characteristic two base 3' overhangs (Bernstein, *et al.*,
20 (2001) *Nature* 409:363). The siRNAs are then incorporated into an RNA-induced silencing complex (RISC) where one or more helicases unwind the siRNA duplex, enabling the complementary antisense strand to guide target recognition (Nykanen, *et al.*, (2001) *Cell* 107:309). Upon binding to the appropriate target mRNA, one or more endonucleases within the RISC cleave the target to induce silencing (Elbashir, *et al.*, (2001) *Genes Dev.* 15:188). Thus, in one aspect the invention relates to a
25 single stranded RNA (siRNA) generated within a cell and which promotes the formation of a RISC complex to effect silencing of the target gene, *i.e.*, an F12 gene. Accordingly, the term “siRNA” is also used herein to refer to an iRNA as described above.

In certain embodiments, the RNAi agent may be a single-stranded siRNA (ssRNAi) that is introduced into a cell or organism to inhibit a target mRNA. Single-stranded RNAi agents bind to the
30 RISC endonuclease, Argonaute 2, which then cleaves the target mRNA. The single-stranded siRNAs are generally 15-30 nucleotides and are chemically modified. The design and testing of single-stranded siRNAs are described in U.S. Patent No. 8,101,348 and in Lima *et al.*, (2012) *Cell* 150:883-894, the entire contents of each of which are hereby incorporated herein by reference. Any of the antisense nucleotide sequences described herein may be used as a single-stranded siRNA as described
35 herein or as chemically modified by the methods described in Lima *et al.*, (2012) *Cell* 150:883-894.

In certain embodiments, an “iRNA” for use in the compositions, uses, and methods of the invention is a double stranded RNA and is referred to herein as a “double stranded RNA agent,”

“double stranded RNA (dsRNA) molecule,” “dsRNA agent,” or “dsRNA”. The term “dsRNA”, refers to a complex of ribonucleic acid molecules, having a duplex structure comprising two anti-parallel and substantially complementary nucleic acid strands, referred to as having “sense” and “antisense” orientations with respect to a target RNA, *i.e.*, an F12 gene. In some embodiments of the invention, a double stranded RNA (dsRNA) triggers the degradation of a target RNA, *e.g.*, an mRNA, through a post-transcriptional gene-silencing mechanism referred to herein as RNA interference or RNAi.

In general, the majority of nucleotides of each strand of a dsRNA molecule are ribonucleotides, but as described in detail herein, each or both strands can also include one or more non-ribonucleotides, *e.g.*, a deoxyribonucleotide or a modified nucleotide. In addition, as used in this specification, an “iRNA” may include ribonucleotides with chemical modifications; an iRNA may include substantial modifications at multiple nucleotides. As used herein, the term “modified nucleotide” refers to a nucleotide having, independently, a modified sugar moiety, a modified internucleotide linkage, or modified nucleobase, or any combination thereof. Thus, the term modified nucleotide encompasses substitutions, additions or removal of, *e.g.*, a functional group or atom, to internucleoside linkages, sugar moieties, or nucleobases. The modifications suitable for use in the agents of the invention include all types of modifications disclosed herein or known in the art. Any such modifications, as used in a siRNA type molecule, are encompassed by “iRNA” or “RNAi agent” for the purposes of this specification and claims.

In certain embodiments of the instant disclosure, inclusion of a deoxy-nucleotide if present within an RNAi agent can be considered to constitute a modified nucleotide.

The duplex region may be of any length that permits specific degradation of a desired target RNA through a RISC pathway, and may range from about 19 to 36 base pairs in length, *e.g.*, about 19-30 base pairs in length, for example, about 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, or 36 base pairs in length, such as about 19-30, 19-29, 19-28, 19-27, 19-26, 19-25, 19-24, 19-23, 19-22, 19-21, 19-20, 20-30, 20-29, 20-28, 20-27, 20-26, 20-25, 20-24, 20-23, 20-22, 20-21, 21-30, 21-29, 21-28, 21-27, 21-26, 21-25, 21-24, 21-23, or 21-22 base pairs in length. In certain embodiments, the duplex region is 19-21 base pairs in length, *e.g.*, 21 base pairs in length. Ranges and lengths intermediate to the above recited ranges and lengths are also contemplated to be part of the disclosure.

The two strands forming the duplex structure may be different portions of one larger RNA molecule, or they may be separate RNA molecules. Where the two strands are part of one larger molecule, and therefore are connected by an uninterrupted chain of nucleotides between the 3'-end of one strand and the 5'-end of the respective other strand forming the duplex structure, the connecting RNA chain is referred to as a “hairpin loop.” A hairpin loop can comprise at least one unpaired nucleotide. In some embodiments, the hairpin loop can comprise at least 2, 3, 4, 5, 6, 7, 8, 9, 10, 20, 23 or more unpaired nucleotides. In some embodiments, the hairpin loop can be 10 or fewer nucleotides. In some embodiments, the hairpin loop can be 8 or fewer unpaired nucleotides. In some

embodiments, the hairpin loop can be 4-10 unpaired nucleotides. In some embodiments, the hairpin loop can be 4-8 nucleotides.

Where the two substantially complementary strands of a dsRNA are comprised by separate RNA molecules, those molecules need not be, but can be covalently connected. Where the two strands are connected covalently by means other than an uninterrupted chain of nucleotides between the 3'-end of one strand and the 5'-end of the respective other strand forming the duplex structure, the connecting structure is referred to as a "linker." The RNA strands may have the same or a different number of nucleotides. The maximum number of base pairs is the number of nucleotides in the shortest strand of the dsRNA minus any overhangs that are present in the duplex. In addition to the duplex structure, an RNAi may comprise one or more nucleotide overhangs. In one embodiment of the RNAi agent, at least one strand comprises a 3' overhang of at least 1 nucleotide. In another embodiment, at least one strand comprises a 3' overhang of at least 2 nucleotides, *e.g.*, 2, 3, 4, 5, 6, 7, 9, 10, 11, 12, 13, 14, or 15 nucleotides. In other embodiments, at least one strand of the RNAi agent comprises a 5' overhang of at least 1 nucleotide. In certain embodiments, at least one strand comprises a 5' overhang of at least 2 nucleotides, *e.g.*, 2, 3, 4, 5, 6, 7, 9, 10, 11, 12, 13, 14, or 15 nucleotides. In still other embodiments, both the 3' and the 5' end of one strand of the RNAi agent comprise an overhang of at least 1 nucleotide.

In certain embodiments, an iRNA agent of the invention is a dsRNA, each strand of which comprises 19-23 nucleotides, that interacts with a target RNA sequence, *e.g.*, an F12 gene, to direct cleavage of the target RNA.

In some embodiments, an iRNA of the invention is a dsRNA of 24-30 nucleotides that interacts with a target RNA sequence, *e.g.*, an F12 target mRNA sequence, to direct the cleavage of the target RNA.

As used herein, the term "nucleotide overhang" refers to at least one unpaired nucleotide that protrudes from the duplex structure of a double stranded iRNA. For example, when a 3'-end of one strand of a dsRNA extends beyond the 5'-end of the other strand, or *vice versa*, there is a nucleotide overhang. A dsRNA can comprise an overhang of at least one nucleotide; alternatively the overhang can comprise at least two nucleotides, at least three nucleotides, at least four nucleotides, at least five nucleotides or more. A nucleotide overhang can comprise or consist of a nucleotide/nucleoside analog, including a deoxynucleotide/nucleoside. The overhang(s) can be on the sense strand, the antisense strand, or any combination thereof. Furthermore, the nucleotide(s) of an overhang can be present on the 5'-end, 3'-end, or both ends of either an antisense or sense strand of a dsRNA.

In one embodiment, the antisense strand of a dsRNA has a 1-10 nucleotide, *e.g.*, a 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10 nucleotide, overhang at the 3'-end or the 5'-end. In one embodiment, the sense strand of a dsRNA has a 1-10 nucleotide, *e.g.*, a 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10 nucleotide, overhang at the 3'-end or the 5'-end. In another embodiment, one or more of the nucleotides in the overhang is replaced with a nucleoside thiophosphate.

In certain embodiments, the antisense strand of a dsRNA has a 1-10 nucleotide, *e.g.*, 0-3, 1-3, 2-4, 2-5, 4-10, 5-10, *e.g.*, a 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10 nucleotide, overhang at the 3'-end or the 5'-end. In one embodiment, the sense strand of a dsRNA has a 1-10 nucleotide, *e.g.*, a 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10 nucleotide, overhang at the 3'-end or the 5'-end. In another embodiment, one or more of
 5 the nucleotides in the overhang is replaced with a nucleoside thiophosphate.

In certain embodiments, the antisense strand of a dsRNA has a 1-10 nucleotides, *e.g.*, a 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10 nucleotide, overhang at the 3'-end or the 5'-end. In certain embodiments, the overhang on the sense strand or the antisense strand, or both, can include extended lengths longer than 10 nucleotides, *e.g.*, 1-30 nucleotides, 2-30 nucleotides, 10-30 nucleotides, 10-25 nucleotides, 10-20
 10 nucleotides, or 10-15 nucleotides in length. In certain embodiments, an extended overhang is on the sense strand of the duplex. In certain embodiments, an extended overhang is present on the 3' end of the sense strand of the duplex. In certain embodiments, an extended overhang is present on the 5' end of the sense strand of the duplex. In certain embodiments, an extended overhang is on the antisense strand of the duplex. In certain embodiments, an extended overhang is present on the 3' end of the
 15 antisense strand of the duplex. In certain embodiments, an extended overhang is present on the 5' end of the antisense strand of the duplex. In certain embodiments, one or more of the nucleotides in the extended overhang is replaced with a nucleoside thiophosphate. In certain embodiments, the overhang includes a self-complementary portion such that the overhang is capable of forming a hairpin structure that is stable under physiological conditions.

“Blunt” or “blunt end” means that there are no unpaired nucleotides at that end of the double stranded RNA agent, *i.e.*, no nucleotide overhang. A “blunt ended” double stranded RNA agent is double stranded over its entire length, *i.e.*, no nucleotide overhang at either end of the molecule. The RNAi agents of the invention include RNAi agents with no nucleotide overhang at one end (*i.e.*, agents with one overhang and one blunt end) or with no nucleotide overhangs at either end. Most
 20 often such a molecule will be double-stranded over its entire length.

The term “antisense strand” or “guide strand” refers to the strand of an iRNA, *e.g.*, a dsRNA, which includes a region that is substantially complementary to a target sequence, *e.g.*, an F12 mRNA.

As used herein, the term “region of complementarity” refers to the region on the antisense strand that is substantially complementary to a sequence, for example a target sequence, *e.g.*, an F12
 30 nucleotide sequence, as defined herein. Where the region of complementarity is not fully complementary to the target sequence, the mismatches can be in the internal or terminal regions of the molecule. Generally, the most tolerated mismatches are in the terminal regions, *e.g.*, within 5, 4, or 3 nucleotides of the 5'- or 3'-end of the iRNA. In some embodiments, a double stranded RNA agent of the invention includes a nucleotide mismatch in the antisense strand. In some embodiments, the
 35 antisense strand of the double stranded RNA agent of the invention includes no more than 4 mismatches with the target mRNA, *e.g.*, the antisense strand includes 4, 3, 2, 1, or 0 mismatches with the target mRNA. In some embodiments, the antisense strand double stranded RNA agent of the

invention includes no more than 4 mismatches with the sense strand, *e.g.*, the antisense strand includes 4, 3, 2, 1, or 0 mismatches with the sense strand. In some embodiments, a double stranded RNA agent of the invention includes a nucleotide mismatch in the sense strand. In some embodiments, the sense strand of the double stranded RNA agent of the invention includes no more than 4 mismatches with the antisense strand, *e.g.*, the sense strand includes 4, 3, 2, 1, or 0 mismatches with the antisense strand. In some embodiments, the nucleotide mismatch is, for example, within 5, 4, 3 nucleotides from the 3'-end of the iRNA. In another embodiment, the nucleotide mismatch is, for example, in the 3'-terminal nucleotide of the iRNA agent. In some embodiments, the mismatch(s) is not in the seed region.

Thus, an RNAi agent as described herein can contain one or more mismatches to the target sequence. In one embodiment, an RNAi agent as described herein contains no more than 3 mismatches (*i.e.*, 3, 2, 1, or 0 mismatches). In one embodiment, an RNAi agent as described herein contains no more than 2 mismatches. In one embodiment, an RNAi agent as described herein contains no more than 1 mismatch. In one embodiment, an RNAi agent as described herein contains 0 mismatches. In certain embodiments, if the antisense strand of the RNAi agent contains mismatches to the target sequence, the mismatch can optionally be restricted to be within the last 5 nucleotides from either the 5'- or 3'-end of the region of complementarity. For example, in such embodiments, for a 23 nucleotide RNAi agent, the strand which is complementary to a region of an F12 gene, generally does not contain any mismatch within the central 13 nucleotides. The methods described herein or methods known in the art can be used to determine whether an RNAi agent containing a mismatch to a target sequence is effective in inhibiting the expression of an F12 gene. Consideration of the efficacy of RNAi agents with mismatches in inhibiting expression of an F12 gene is important, especially if the particular region of complementarity in an F12 gene is known to have polymorphic sequence variation within the population.

The term "sense strand" or "passenger strand" as used herein, refers to the strand of an iRNA that includes a region that is substantially complementary to a region of the antisense strand as that term is defined herein.

As used herein, "substantially all of the nucleotides are modified" are largely but not wholly modified and can include not more than 5, 4, 3, 2, or 1 unmodified nucleotides.

As used herein, the term "cleavage region" refers to a region that is located immediately adjacent to the cleavage site. The cleavage site is the site on the target at which cleavage occurs. In some embodiments, the cleavage region comprises three bases on either end of, and immediately adjacent to, the cleavage site. In some embodiments, the cleavage region comprises two bases on either end of, and immediately adjacent to, the cleavage site. In some embodiments, the cleavage site specifically occurs at the site bound by nucleotides 10 and 11 of the antisense strand, and the cleavage region comprises nucleotides 11, 12 and 13.

As used herein, and unless otherwise indicated, the term “complementary,” when used to describe a first nucleotide sequence in relation to a second nucleotide sequence, refers to the ability of an oligonucleotide or polynucleotide comprising the first nucleotide sequence to hybridize and form a duplex structure under certain conditions with an oligonucleotide or polynucleotide comprising the second nucleotide sequence, as will be understood by the skilled person. Such conditions can, for example, be stringent conditions, where stringent conditions can include: 400 mM NaCl, 40 mM PIPES pH 6.4, 1 mM EDTA, 50°C or 70°C for 12-16 hours followed by washing (see, *e.g.*, “Molecular Cloning: A Laboratory Manual, Sambrook, *et al.* (1989) Cold Spring Harbor Laboratory Press). Other conditions, such as physiologically relevant conditions as can be encountered inside an organism, can apply. The skilled person will be able to determine the set of conditions most appropriate for a test of complementarity of two sequences in accordance with the ultimate application of the hybridized nucleotides.

Complementary sequences within an iRNA, *e.g.*, within a dsRNA as described herein, include base-pairing of the oligonucleotide or polynucleotide comprising a first nucleotide sequence to an oligonucleotide or polynucleotide comprising a second nucleotide sequence over the entire length of one or both nucleotide sequences. Such sequences can be referred to as “fully complementary” with respect to each other herein. However, where a first sequence is referred to as “substantially complementary” with respect to a second sequence herein, the two sequences can be fully complementary, or they can form one or more, but generally not more than 5, 4, 3, or 2 mismatched base pairs upon hybridization for a duplex up to 30 base pairs, while retaining the ability to hybridize under the conditions most relevant to their ultimate application, *e.g.*, inhibition of gene expression, *in vitro* or *in vivo*. However, where two oligonucleotides are designed to form, upon hybridization, one or more single stranded overhangs, such overhangs shall not be regarded as mismatches with regard to the determination of complementarity. For example, a dsRNA comprising one oligonucleotide 21 nucleotides in length and another oligonucleotide 23 nucleotides in length, wherein the longer oligonucleotide comprises a sequence of 21 nucleotides that is fully complementary to the shorter oligonucleotide, can yet be referred to as “fully complementary” for the purposes described herein.

“Complementary” sequences, as used herein, can also include, or be formed entirely from, non-Watson-Crick base pairs or base pairs formed from non-natural and modified nucleotides, in so far as the above requirements with respect to their ability to hybridize are fulfilled. Such non-Watson-Crick base pairs include, but are not limited to, G:U Wobble or Hoogsteen base pairing.

The terms “complementary,” “fully complementary” and “substantially complementary” herein can be used with respect to the base matching between the sense strand and the antisense strand of a dsRNA, or between two oligonucleotides or polynucleotides, such as the antisense strand of a double stranded RNA agent and a target sequence, as will be understood from the context of their use.

As used herein, a polynucleotide that is “substantially complementary to at least part of” a messenger RNA (mRNA) refers to a polynucleotide that is substantially complementary to a

contiguous portion of the mRNA of interest (*e.g.*, an mRNA encoding an F12 gene). For example, a polynucleotide is complementary to at least a part of an F12 mRNA if the sequence is substantially complementary to a non-interrupted portion of an mRNA encoding an F12 gene.

Accordingly, in some embodiments, the antisense polynucleotides disclosed herein are fully complementary to the target F12 sequence. In other embodiments, the antisense polynucleotides disclosed herein are substantially complementary to the target F12 sequence and comprise a contiguous nucleotide sequence which is at least 80% complementary over its entire length to the equivalent region of the nucleotide sequence of any one of SEQ ID NOs:1, 3, 5, 7, or 9, or a fragment of any one of SEQ ID NOs:1, 3, 5, 7, or 9, such as about 85%, about 90%, about 91%, about 92%, about 93%, about 94%, about 95%, about 96%, about 97%, about 98%, or about 99% complementary.

In other embodiments, the antisense polynucleotides disclosed herein are substantially complementary to the target F12 sequence and comprise a contiguous nucleotide sequence which is at least about 80% complementary over its entire length to any one of the sense strand nucleotide sequences in any one of any one of Tables 2-3, or a fragment of any one of the sense strand nucleotide sequences in any one of Tables 2-3, such as about 85%, about 90%, about 91%, about 92%, about 93%, about 94%, about 95%, about 96%, about 97%, about 98%, about 99%, or 100% complementary.

In some embodiments, the antisense polynucleotides disclosed herein are substantially complementary to a fragment of a target F12 sequence and comprise a contiguous nucleotide sequence which is at least 80% complementary over its entire length to a fragment of SEQ ID NO: 1 selected from the group of nucleotides 75-97, 76-98, 79-101, 115-137, 117-139, 123-145, 132-154, 135-157, 137-159, 144-166, 201-223, or 1462-1484 of SEQ ID NO: 1, such as about 85%, about 90%, about 91%, about 92%, about 93%, about 94%, about 95%, about 96%, about 97%, about 98%, or about 99% complementary.

In some embodiments, the antisense polynucleotides disclosed herein are substantially complementary to a fragment of a target F12 sequence and comprise a contiguous nucleotide sequence which is at least 80% complementary over its entire length to a fragment of SEQ ID NO: 1 selected from the group of nucleotides 135-157, 144-166, or 201-223 of SEQ ID NO: 1, such as about 85%, about 90%, about 91%, about 92%, about 93%, about 94%, about 95%, about 96%, about 97%, about 98%, or about 99% complementary.

In one embodiment, an RNAi agent of the disclosure includes a sense strand that is substantially complementary to an antisense polynucleotide which, in turn, is the same as a target F12 sequence, and wherein the sense strand polynucleotide comprises a contiguous nucleotide sequence which is at least about 80% complementary over its entire length to the equivalent region of the nucleotide sequence of SEQ ID NOs: 2, 4, 6, 8, or 10, or a fragment of any one of SEQ ID NOs:2, 4, 6, 8, or 10, such as about 85%, about 90%, about 91%, about 92%, about 93%, about 94%, about 95%, about 96%, about 97%, about 98%, about 99%, or 100% complementary.

In some embodiments, an iRNA of the invention includes a sense strand that is substantially complementary to an antisense polynucleotide which, in turn, is complementary to a target F12 sequence, and wherein the sense strand polynucleotide comprises a contiguous nucleotide sequence which is at least about 80% complementary over its entire length to any one of the antisense strand nucleotide sequences in any one of any one of Tables 2-3, or a fragment of any one of the antisense strand nucleotide sequences in any one of Tables 2-3, such as about 85%, about 90%, about 91%, about 92%, about 93%, about 94%, about 95%, about 96%, about 97%, about 98%, about 99%, or 100% complementary.

In certain embodiments, the sense and antisense strands are selected from any one of duplexes AD-1534528, AD-1534479, AD-1534582, AD-1534611, AD-1534506, AD-1631309, AD-1534471, AD-1534555, AD-1534678, AD-1534585, AD-1534502, and AD-1534491.

In certain embodiments, the sense and antisense strands are selected from any one of duplexes AD-1534528, AD-1534479, and AD-1534582.

In general, an “iRNA” includes ribonucleotides with chemical modifications. Such modifications may include all types of modifications disclosed herein or known in the art. Any such modifications, as used in a dsRNA molecule, are encompassed by “iRNA” for the purposes of this specification and claims.

In certain embodiments of the instant disclosure, inclusion of a deoxy-nucleotide if present within an RNAi agent can be considered to constitute a modified nucleotide.

In an aspect of the invention, an agent for use in the methods and compositions of the invention is a single-stranded antisense oligonucleotide molecule that inhibits a target mRNA *via* an antisense inhibition mechanism. The single-stranded antisense oligonucleotide molecule is complementary to a sequence within the target mRNA. The single-stranded antisense oligonucleotides can inhibit translation in a stoichiometric manner by base pairing to the mRNA and physically obstructing the translation machinery, see Dias, N. *et al.*, (2002) *Mol Cancer Ther* 1:347-355. The single-stranded antisense oligonucleotide molecule may be about 14 to about 30 nucleotides in length and have a sequence that is complementary to a target sequence. For example, the single-stranded antisense oligonucleotide molecule may comprise a sequence that is at least about 14, 15, 16, 17, 18, 19, 20, or more contiguous nucleotides from any one of the antisense sequences described herein.

The phrase “contacting a cell with an iRNA,” such as a dsRNA, as used herein, includes contacting a cell by any possible means. Contacting a cell with an iRNA includes contacting a cell *in vitro* with the iRNA or contacting a cell *in vivo* with the iRNA. The contacting may be done directly or indirectly. Thus, for example, the iRNA may be put into physical contact with the cell by the individual performing the method, or alternatively, the iRNA may be put into a situation that will permit or cause it to subsequently come into contact with the cell.

Contacting a cell *in vitro* may be done, for example, by incubating the cell with the iRNA. Contacting a cell *in vivo* may be done, for example, by injecting the iRNA into or near the tissue where the cell is located, or by injecting the iRNA into another area, *e.g.*, the bloodstream or the subcutaneous space, such that the agent will subsequently reach the tissue where the cell to be
5 contacted is located. For example, the iRNA may contain or be coupled to a ligand, *e.g.*, GalNAc, that directs the iRNA to a site of interest, *e.g.*, the liver. Combinations of *in vitro* and *in vivo* methods of contacting are also possible. For example, a cell may also be contacted *in vitro* with an iRNA and subsequently transplanted into a subject.

In certain embodiments, contacting a cell with an iRNA includes “introducing” or “delivering
10 the iRNA into the cell” by facilitating or effecting uptake or absorption into the cell. Absorption or uptake of an iRNA can occur through unaided diffusion or active cellular processes, or by auxiliary agents or devices. Introducing an iRNA into a cell may be *in vitro* or *in vivo*. For example, for *in vivo* introduction, iRNA can be injected into a tissue site or administered systemically. *In vitro* introduction into a cell includes methods known in the art such as electroporation and lipofection.
15 Further approaches are described herein below or are known in the art.

The term “lipid nanoparticle” or “LNP” is a vesicle comprising a lipid layer encapsulating a pharmaceutically active molecule, such as a nucleic acid molecule, *e.g.*, an iRNA or a plasmid from which an iRNA is transcribed. LNPs are described in, for example, U.S. Patent Nos. 6,858,225, 6,815,432, 8,158,601, and 8,058,069, the entire contents of which are hereby incorporated herein by
20 reference.

As used herein, a “subject” is an animal, such as a mammal, including a primate (such as a human, a non-human primate, *e.g.*, a monkey, and a chimpanzee), a non-primate (such as a cow, a pig, a horse, a goat, a rabbit, a sheep, a hamster, a guinea pig, a cat, a dog, a rat, or a mouse), or a bird that expresses the target gene, either endogenously or heterologously. In an embodiment, the subject
25 is a human, such as a human being treated or assessed for a disease or disorder that would benefit from reduction in F12 expression; a human at risk for a disease or disorder that would benefit from reduction in F12 expression; a human having a disease or disorder that would benefit from reduction in F12 expression; or human being treated for a disease or disorder that would benefit from reduction in F12 expression as described herein. In some embodiments, the subject is a female human. In other
30 embodiments, the subject is a male human. In one embodiment, the subject is an adult subject. In another embodiment, the subject is a pediatric subject.

As used herein, the terms “treating” or “treatment” refer to a beneficial or desired result, such as reducing at least one sign or symptom of an F12-associated disorder in a subject. Treatment also includes a reduction of one or more sign or symptoms associated with unwanted F12 expression;
35 diminishing the extent of unwanted F12 activation or stabilization; amelioration or palliation of unwanted F12 activation or stabilization. “Treatment” can also mean prolonging survival as compared to expected survival in the absence of treatment.

The term “lower” in the context of the level of F12 in a subject or a disease marker or symptom refers to a statistically significant decrease in such level. The decrease can be, for example, at least 10%, 15%, 20%, 25%, 30%, %, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, or more. In certain embodiments, a decrease is at least 20%. In certain embodiments, the decrease is at least 50% in a disease marker, *e.g.*, protein or gene expression level. “Lower” in the context of the level of F12 in a subject is a decrease to a level accepted as within the range of normal for an individual without such disorder. In certain embodiments, “lower” is the decrease in the difference between the level of a marker or symptom for a subject suffering from a disease and a level accepted within the range of normal for an individual. The term “lower” can also be used in association with normalizing a symptom of a disease or condition, *i.e.* decreasing the difference between a level in a subject suffering from an F12-associated disorder towards or to a level in a normal subject not suffering from an F12-associated disorder. As used herein, if a disease is associated with an elevated value for a symptom, “normal” is considered to be the upper limit of normal. If a disease is associated with a decreased value for a symptom, “normal” is considered to be the lower limit of normal.

As used herein, “prevention” or “preventing,” when used in reference to a disease, disorder or condition thereof, may be treated or ameliorated by a reduction in expression of an F12 gene, refers to a reduction in the likelihood that a subject will develop a symptom associated with such a disease, disorder, or condition, *e.g.*, a symptom of an F12-associated disorder, *e.g.*, hereditary angioedema (HAE) (*e.g.*, hereditary angioedema type I; hereditary angioedema type II; hereditary angioedema type III; or any other hereditary angioedema caused by elevated levels of bradykinin); prekallikrein deficiency, hypertension, *e.g.*, malignant essential hypertension, end stage renal disease, Fletcher Factor Deficiency, thromboembolic disease, inflammatory disease, or Alzheimer’s Disease. The failure to develop a disease, disorder or condition, or the reduction in the development of a symptom associated with such a disease, disorder or condition (*e.g.*, by at least about 10% on a clinically accepted scale for that disease or disorder), or the exhibition of delayed symptoms delayed (*e.g.*, by days, weeks, months or years) is considered effective prevention.

As used herein, the term “Factor XII-associated disorder” or “F12-associated disorder,” is a disease or disorder that is caused by, or associated with, F12 gene expression or F12 protein production. The term “F12-associated disorder” includes a disease, disorder or condition that would benefit from a decrease in F12 gene expression, replication, or protein activity. In some embodiments, the F12-associated disorder is selected from the group consisting of hereditary angioedema (HAE) (*e.g.*, hereditary angioedema type I; hereditary angioedema type II; hereditary angioedema type III; or any other hereditary angioedema caused by elevated levels of bradykinin); prekallikrein deficiency, hypertension, *e.g.*, malignant essential hypertension, end stage renal disease, Fletcher Factor Deficiency, thromboembolic disease, inflammatory disease, and Alzheimer’s Disease.

In one embodiment, an F12-associated disorder is HAE. In another embodiment, an F12-associated disorder is prekallikrein deficiency. In another embodiment, an F12-associated disorder is hypertension, *e.g.*, malignant essential hypertension. In another embodiment, an F12-associated disorder is end stage renal disease. In another embodiment, an F12-associated disorder is Fletcher Factor deficiency. In another embodiment, an F12-associated disorder is thromboembolic disease. In another embodiment, an F12-associated disorder is inflammatory disease. In another embodiment, an F12-associated disorder is Alzheimer's Disease.

"Therapeutically effective amount," as used herein, is intended to include the amount of an RNAi agent that, when administered to a subject having an F12-associated disorder, is sufficient to effect treatment of the disease (*e.g.*, by diminishing, ameliorating, or maintaining the existing disease or one or more symptoms of disease). The "therapeutically effective amount" may vary depending on the RNAi agent, how the agent is administered, the disease and its severity and the history, age, weight, family history, genetic makeup, the types of preceding or concomitant treatments, if any, and other individual characteristics of the subject to be treated.

"Prophylactically effective amount," as used herein, is intended to include the amount of an RNAi agent that, when administered to a subject having an F12-associated disorder, is sufficient to prevent or ameliorate the disorder or one or more symptoms of the disorder, such as edema swelling of the extremities, face, larynx, upper respiratory tract, abdomen, trunk and genitals; prodrome; laryngeal swelling; nonpruritic rash; nausea; vomiting; or abdominal pain. Ameliorating the disease includes slowing the course of the disease or reducing the severity of later-developing disease. The "prophylactically effective amount" may vary depending on the RNAi agent, how the agent is administered, the degree of risk of disease, and the history, age, weight, family history, genetic makeup, the types of preceding or concomitant treatments, if any, and other individual characteristics of the patient to be treated.

A "therapeutically-effective amount" or "prophylactically effective amount" also includes an amount of an RNAi agent that produces some desired effect at a reasonable benefit/risk ratio applicable to any treatment. The iRNA employed in the methods of the present invention may be administered in a sufficient amount to produce a reasonable benefit/risk ratio applicable to such treatment.

The phrase "pharmaceutically acceptable" is employed herein to refer to those compounds, materials (including salts), compositions, or dosage forms which are, within the scope of sound medical judgment, suitable for use in contact with the tissues of human subjects and animal subjects without excessive toxicity, irritation, allergic response, or other problem or complication, commensurate with a reasonable benefit/risk ratio.

The phrase "pharmaceutically-acceptable carrier" as used herein means a pharmaceutically-acceptable material, composition, or vehicle, such as a liquid or solid filler, diluent, excipient, manufacturing aid (*e.g.*, lubricant, talc magnesium, calcium or zinc stearate, or steric acid), or solvent

encapsulating material, involved in carrying or transporting the subject compound from one organ, or portion of the body, to another organ, or portion of the body. Each carrier must be "acceptable" in the sense of being compatible with the other ingredients of the formulation and not injurious to the subject being treated. Such carriers are known in the art. Pharmaceutically acceptable carriers

5 include carriers for administration by injection.

The term "sample," as used herein, includes a collection of similar fluids, cells, or tissues isolated from a subject, as well as fluids, cells, or tissues present within a subject. Examples of biological fluids include blood, serum and serosal fluids, plasma, cerebrospinal fluid, ocular fluids, lymph, urine, saliva, and the like. Tissue samples may include samples from tissues, organs, or

10 localized regions. For example, samples may be derived from particular organs, parts of organs, or fluids or cells within those organs. In certain embodiments, samples may be derived from the liver (*e.g.*, whole liver or certain segments of liver or certain types of cells in the liver, such as, *e.g.*, hepatocytes). In some embodiments, a "sample derived from a subject" refers to urine obtained from the subject. A "sample derived from a subject" can refer to blood or blood derived serum or plasma

15 from the subject.

II. iRNAs of the Invention

The present invention provides iRNAs which inhibit the expression of an F12 gene. In certain embodiments, the iRNA includes double stranded ribonucleic acid (dsRNA) molecules for

20 inhibiting the expression of an F12 gene in a cell, such as a cell within a subject, *e.g.*, a mammal, such as a human susceptible to developing an F12-associated disorder, *e.g.*, hereditary angioedema (HAE) (*e.g.*, hereditary angioedema type I; hereditary angioedema type II; hereditary angioedema type III; or any other hereditary angioedema caused by elevated levels of bradykinin); prekallikrein deficiency, hypertension, *e.g.*, malignant essential hypertension, end stage renal disease, Fletcher Factor

25 Deficiency, thromboembolic disease, inflammatory disease, or Alzheimer's Disease. The dsRNAi agent includes an antisense strand having a region of complementarity which is complementary to at least a part of an mRNA formed in the expression of an F12 gene. The region of complementarity is about 19-30 nucleotides in length (*e.g.*, about 30, 29, 28, 27, 26, 25, 24, 23, 22, 21, 20, or 19 nucleotides in length).

30 Upon contact with a cell expressing the F12 gene, the iRNA inhibits the expression of the F12 gene (*e.g.*, a human, a primate, a non-primate, or a rat F12 gene) by at least about 50% as assayed by, for example, a PCR or branched DNA (bDNA)-based method, or by a protein-based method, such as by immunofluorescence analysis, using, for example, western blotting or flow cytometric techniques. In certain embodiments, inhibition of expression is determined by the qPCR method provided in the

35 examples herein with the siRNA at, *e.g.*, a 10 nM concentration, in an appropriate organism cell line provided therein. In certain embodiments, inhibition of expression *in vivo* is determined by knockdown of the human gene in a rodent expressing the human gene, *e.g.*, a mouse or an AAV-

infected mouse expressing the human target gene, *e.g.*, when administered as single dose, *e.g.*, at 3 mg/kg at the nadir of RNA expression.

A dsRNA includes two RNA strands that are complementary and hybridize to form a duplex structure under conditions in which the dsRNA will be used. One strand of a dsRNA (the antisense strand) includes a region of complementarity that is substantially complementary, and generally fully complementary, to a target sequence. The target sequence can be derived from the sequence of an mRNA formed during the expression of an F12 gene. The other strand (the sense strand) includes a region that is complementary to the antisense strand, such that the two strands hybridize and form a duplex structure when combined under suitable conditions. As described elsewhere herein and as known in the art, the complementary sequences of a dsRNA can also be contained as self-complementary regions of a single nucleic acid molecule, as opposed to being on separate oligonucleotides.

Generally, the duplex structure is 15 to 30 base pairs in length, *e.g.*, 15-29, 15-28, 15-27, 15-26, 15-25, 15-24, 15-23, 15-22, 15-21, 15-20, 15-19, 15-18, 15-17, 18-30, 18-29, 18-28, 18-27, 18-26, 18-25, 18-24, 18-23, 18-22, 18-21, 18-20, 19-30, 19-29, 19-28, 19-27, 19-26, 19-25, 19-24, 19-23, 19-22, 19-21, 19-20, 20-30, 20-29, 20-28, 20-27, 20-26, 20-25, 20-24, 20-23, 20-22, 20-21, 21-30, 21-29, 21-28, 21-27, 21-26, 21-25, 21-24, 21-23, or 21-22 base pairs in length. In certain embodiments, the duplex structure is 18 to 25 base pairs in length, *e.g.*, 18-25, 18-24, 18-23, 18-22, 18-21, 18-20, 19-25, 19-24, 19-23, 19-22, 19-21, 19-20, 20-25, 20-24, 20-23, 20-22, 20-21, 21-25, 21-24, 21-23, 21-22, 22-25, 22-24, 22-23, 23-25, 23-24 or 24-25 base pairs in length, for example, 19-21 basepairs in length. Ranges and lengths intermediate to the above recited ranges and lengths are also contemplated to be part of the disclosure.

Similarly, the region of complementarity to the target sequence is 15 to 30 nucleotides in length, *e.g.*, 15-29, 15-28, 15-27, 15-26, 15-25, 15-24, 15-23, 15-22, 15-21, 15-20, 15-19, 15-18, 15-17, 18-30, 18-29, 18-28, 18-27, 18-26, 18-25, 18-24, 18-23, 18-22, 18-21, 18-20, 19-30, 19-29, 19-28, 19-27, 19-26, 19-25, 19-24, 19-23, 19-22, 19-21, 19-20, 20-30, 20-29, 20-28, 20-27, 20-26, 20-25, 20-24, 20-23, 20-22, 20-21, 21-30, 21-29, 21-28, 21-27, 21-26, 21-25, 21-24, 21-23, or 21-22 nucleotides in length, for example 19-23 nucleotides in length or 21-23 nucleotides in length. Ranges and lengths intermediate to the above recited ranges and lengths are also contemplated to be part of the disclosure.

In some embodiments, the duplex structure is 19 to 30 base pairs in length. Similarly, the region of complementarity to the target sequence is 19 to 30 nucleotides in length.

In some embodiments, the dsRNA is about 19 to about 23 nucleotides in length, or about 25 to about 30 nucleotides in length. In general, the dsRNA is long enough to serve as a substrate for the Dicer enzyme. For example, it is well-known in the art that dsRNAs longer than about 21-23 nucleotides in length may serve as substrates for Dicer. As the ordinarily skilled person will also recognize, the region of an RNA targeted for cleavage will most often be part of a larger RNA molecule, often an mRNA molecule. Where relevant, a “part” of an mRNA target is a contiguous

sequence of an mRNA target of sufficient length to allow it to be a substrate for RNAi-directed cleavage (*i.e.*, cleavage through a RISC pathway).

One of skill in the art will also recognize that the duplex region is a primary functional portion of a dsRNA, *e.g.*, a duplex region of about 19 to about 30 base pairs, *e.g.*, about 19-30, 19-29, 19-28, 19-27, 19-26, 19-25, 19-24, 19-23, 19-22, 19-21, 19-20, 20-30, 20-29, 20-28, 20-27, 20-26, 20-25, 20-24, 20-23, 20-22, 20-21, 21-30, 21-29, 21-28, 21-27, 21-26, 21-25, 21-24, 21-23, or 21-22 base pairs. Thus, in one embodiment, to the extent that it becomes processed to a functional duplex, of *e.g.*, 15-30 base pairs, that targets a desired RNA for cleavage, an RNA molecule or complex of RNA molecules having a duplex region greater than 30 base pairs is a dsRNA. Thus, an ordinarily skilled artisan will recognize that in one embodiment, a miRNA is a dsRNA. In another embodiment, a dsRNA is not a naturally occurring miRNA. In another embodiment, an iRNA agent useful to target F12 gene expression is not generated in the target cell by cleavage of a larger dsRNA.

A dsRNA as described herein can further include one or more single-stranded nucleotide overhangs, *e.g.*, 1-4, 2-4, 1-3, 2-3, 1, 2, 3, or 4 nucleotides. dsRNAs having at least one nucleotide overhang can have superior inhibitory properties relative to their blunt-ended counterparts. A nucleotide overhang can comprise or consist of a nucleotide/nucleoside analog, including a deoxynucleotide/nucleoside. The overhang(s) can be on the sense strand, the antisense strand, or any combination thereof. Furthermore, the nucleotide(s) of an overhang can be present on the 5'-end, 3'-end, or both ends of an antisense or sense strand of a dsRNA.

A dsRNA can be synthesized by standard methods known in the art. Double stranded RNAi compounds of the invention may be prepared using a two-step procedure. First, the individual strands of the double stranded RNA molecule are prepared separately. Then, the component strands are annealed. The individual strands of the siRNA compound can be prepared using solution-phase or solid-phase organic synthesis or both. Organic synthesis offers the advantage that the oligonucleotide strands comprising unnatural or modified nucleotides can be easily prepared. Similarly, single-stranded oligonucleotides of the invention can be prepared using solution-phase or solid-phase organic synthesis or both.

In an aspect, a dsRNA of the invention includes at least two nucleotide sequences, a sense sequence and an anti-sense sequence. The sense strand is selected from the group of sequences provided in any one of Tables 2-3, and the corresponding antisense strand of the sense strand is selected from the group of sequences of any one of Tables 2-3. In this aspect, one of the two sequences is complementary to the other of the two sequences, with one of the sequences being substantially complementary to a sequence of an mRNA generated in the expression of an F12 gene. As such, in this aspect, a dsRNA will include two oligonucleotides, where one oligonucleotide is described as the sense strand in any one of Tables 2-3, and the second oligonucleotide is described as the corresponding antisense strand of the sense strand in any one of Tables 2-3.

In certain embodiments, the substantially complementary sequences of the dsRNA are contained on separate oligonucleotides. In other embodiments, the substantially complementary sequences of the dsRNA are contained on a single oligonucleotide.

It will be understood that, although the sequences in, for example, Table 3, are not described as modified or conjugated sequences, the RNA of the iRNA of the invention *e.g.*, a dsRNA of the invention, may comprise any one of the sequences set forth in any one of Tables 2-3 that is unmodified, un-conjugated, or modified or conjugated differently than described therein. In other words, the invention encompasses dsRNA of Tables 2-3 which are un-modified, un-conjugated, modified, or conjugated, as described herein.

The skilled person is well aware that dsRNAs having a duplex structure of about 20 to 23 base pairs, *e.g.*, 21, base pairs have been hailed as particularly effective in inducing RNA interference (Elbashir *et al.*, *EMBO* 2001, 20:6877-6888). However, others have found that shorter or longer RNA duplex structures can also be effective (Chu and Rana (2007) *RNA* 14:1714-1719; Kim *et al.* (2005) *Nat Biotech* 23:222-226). In the embodiments described above, by virtue of the nature of the oligonucleotide sequences provided in any one of Tables 2-3, dsRNAs described herein can include at least one strand of a length of minimally 21 nucleotides. It can be reasonably expected that shorter duplexes having any one of the sequences in any one of Tables 2-3 minus only a few nucleotides on one or both ends can be similarly effective as compared to the dsRNAs described above. Hence, dsRNAs having a sequence of at least 19, 20, or more contiguous nucleotides derived from any one of the sequences of any one of Tables 2-3, and differing in their ability to inhibit the expression of an F12 gene by not more than about 5, 10, 15, 20, 25, or 30 % inhibition from a dsRNA comprising the full sequence, are contemplated to be within the scope of the present invention.

In addition, the RNAs provided in Tables 2-3 identify a site(s) in an F12 transcript that is susceptible to RISC-mediated cleavage. As such, the present invention further features iRNAs that target within one of these sites. As used herein, an iRNA is said to target within a particular site of an RNA transcript if the iRNA promotes cleavage of the transcript anywhere within that particular site. Such an iRNA will generally include at least about 19 contiguous nucleotides from any one of the sequences provided in any one of Tables 2-3 coupled to additional nucleotide sequences taken from the region contiguous to the selected sequence in an F12 gene.

30

III. Modified iRNAs of the Invention

In certain embodiments, the RNA of the iRNA of the invention *e.g.*, a dsRNA, is unmodified, and does not comprise, *e.g.*, chemical modifications or conjugations known in the art and described herein. In other embodiments, the RNA of an iRNA of the invention, *e.g.*, a dsRNA, is chemically modified to enhance stability or other beneficial characteristics. In certain embodiments of the invention, substantially all of the nucleotides of an iRNA of the invention are modified. In other embodiments of the invention, all of the nucleotides of an iRNA or substantially all of the nucleotides

35

of an iRNA are modified, *i.e.*, not more than 5, 4, 3, 2, or 1 unmodified nucleotides are present in a strand of the iRNA.

The nucleic acids featured in the invention can be synthesized or modified by methods well established in the art, such as those described in “Current protocols in nucleic acid chemistry,”
5 Beaucage, S.L. *et al.* (Edrs.), John Wiley & Sons, Inc., New York, NY, USA, which is hereby incorporated herein by reference. Modifications include, for example, end modifications, *e.g.*, 5'-end modifications (phosphorylation, conjugation, inverted linkages) or 3'-end modifications (conjugation, DNA nucleotides, inverted linkages, *etc.*); base modifications, *e.g.*, replacement with stabilizing bases, destabilizing bases, or bases that base pair with an expanded repertoire of partners, removal of
10 bases (abasic nucleotides), or conjugated bases; sugar modifications (*e.g.*, at the 2'-position or 4'-position) or replacement of the sugar; or backbone modifications, including modification or replacement of the phosphodiester linkages. Specific examples of iRNA compounds useful in the embodiments described herein include, but are not limited to RNAs containing modified backbones or no natural internucleoside linkages. RNAs having modified backbones include, among others, those
15 that do not have a phosphorus atom in the backbone. For the purposes of this specification, and as sometimes referenced in the art, modified RNAs that do not have a phosphorus atom in their internucleoside backbone can also be considered to be oligonucleosides. In some embodiments, a modified iRNA will have a phosphorus atom in its internucleoside backbone.

Modified RNA backbones include, for example, phosphorothioates, chiral phosphorothioates,
20 phosphorodithioates, phosphotriesters, aminoalkylphosphotriesters, methyl and other alkyl phosphonates including 3'-alkylene phosphonates and chiral phosphonates, phosphinates, phosphoramidates including 3'-amino phosphoramidate and aminoalkylphosphoramidates, thionophosphoramidates, thionoalkylphosphonates, thionoalkylphosphotriesters, and boranophosphates having normal 3'-5' linkages, 2'-5'-linked analogs of these, and those having
25 inverted polarity wherein the adjacent pairs of nucleoside units are linked 3'-5' to 5'-3' or 2'-5' to 5'-2'. Various salts, mixed salts and free acid forms are also included. In some embodiments of the invention, the dsRNA agents of the invention are in a free acid form. In other embodiments of the invention, the dsRNA agents of the invention are in a salt form. In one embodiment, the dsRNA agents of the invention are in a sodium salt form. In certain embodiments, when the dsRNA agents of
30 the invention are in the sodium salt form, sodium ions are present in the agent as counterions for substantially all of the phosphodiester and/or phosphorothioate groups present in the agent. Agents in which substantially all of the phosphodiester and/or phosphorothioate linkages have a sodium counterion include not more than 5, 4, 3, 2, or 1 phosphodiester and/or phosphorothioate linkages without a sodium counterion. In some embodiments, when the dsRNA agents of the invention are in
35 the sodium salt form, sodium ions are present in the agent as counterions for all of the phosphodiester and/or phosphorothioate groups present in the agent.

Representative U.S. Patents that teach the preparation of the above phosphorus-containing linkages include, but are not limited to, U.S. Patent Nos. 3,687,808; 4,469,863; 4,476,301; 5,023,243; 5,177,195; 5,188,897; 5,264,423; 5,276,019; 5,278,302; 5,286,717; 5,321,131; 5,399,676; 5,405,939; 5,453,496; 5,455,233; 5,466,677; 5,476,925; 5,519,126; 5,536,821; 5,541,316; 5,550,111; 5,563,253; 5,571,799; 5,587,361; 5,625,050; 6,028,188; 6,124,445; 6,160,109; 6,169,170; 6,172,209; 6,239,265; 6,277,603; 6,326,199; 6,346,614; 6,444,423; 6,531,590; 6,534,639; 6,608,035; 6,683,167; 6,858,715; 6,867,294; 6,878,805; 7,015,315; 7,041,816; 7,273,933; 7,321,029; and U.S. Pat RE39464, the entire contents of each of which are hereby incorporated herein by reference.

Modified RNA backbones that do not include a phosphorus atom therein have backbones that are formed by short chain alkyl or cycloalkyl internucleoside linkages, mixed heteroatoms and alkyl or cycloalkyl internucleoside linkages, or one or more short chain heteroatomic or heterocyclic internucleoside linkages. These include those having morpholino linkages (formed in part from the sugar portion of a nucleoside); siloxane backbones; sulfide, sulfoxide and sulfone backbones; formacetyl and thioformacetyl backbones; methylene formacetyl and thioformacetyl backbones; alkene containing backbones; sulfamate backbones; methyleneimino and methylenehydrazino backbones; sulfonate and sulfonamide backbones; amide backbones; and others having mixed N, O, S, and CH₂ component parts.

Representative U.S. Patents that teach the preparation of the above oligonucleosides include, but are not limited to, U.S. Patent Nos. 5,034,506; 5,166,315; 5,185,444; 5,214,134; 5,216,141; 5,235,033; 5,64,562; 5,264,564; 5,405,938; 5,434,257; 5,466,677; 5,470,967; 5,489,677; 5,541,307; 5,561,225; 5,596,086; 5,602,240; 5,608,046; 5,610,289; 5,618,704; 5,623,070; 5,663,312; 5,633,360; 5,677,437; and 5,677,439, the entire contents of each of which are hereby incorporated herein by reference.

Suitable RNA mimetics are contemplated for use in iRNAs provided herein, in which both the sugar and the internucleoside linkage, *i.e.*, the backbone, of the nucleotide units are replaced with novel groups. The base units are maintained for hybridization with an appropriate nucleic acid target compound. One such oligomeric compound in which an RNA mimetic that has been shown to have excellent hybridization properties is referred to as a peptide nucleic acid (PNA). In PNA compounds, the sugar backbone of an RNA is replaced with an amide containing backbone, in particular an aminoethylglycine backbone. The nucleobases are retained and are bound directly or indirectly to aza nitrogen atoms of the amide portion of the backbone. Representative US patents that teach the preparation of PNA compounds include, but are not limited to, U.S. Patent Nos. 5,539,082; 5,714,331; and 5,719,262, the entire contents of each of which are hereby incorporated herein by reference. Additional PNA compounds suitable for use in the iRNAs of the invention are described in, for example, in Nielsen *et al.*, *Science*, 1991, 254, 1497-1500.

Some embodiments featured in the invention include RNAs with phosphorothioate backbones and oligonucleosides with heteroatom backbones, and in particular --CH₂--NH--CH₂--, --CH₂--

N(CH₃)--O--CH₂--[known as a methylene (methylimino) or MMI backbone], --CH₂--O--N(CH₃)--CH₂--, --CH₂--N(CH₃)--N(CH₃)--CH₂-- and --N(CH₃)--CH₂--CH₂-- of the above-referenced U.S. Patent No. 5,489,677, and the amide backbones of the above-referenced U.S. Patent No. 5,602,240. In some embodiments, the RNAs featured herein have morpholino backbone structures of the above-referenced U.S. Patent No. 5,034,506. The native phosphodiester backbone can be represented as O-P(O)(OH)-OCH₂-.

Modified RNAs can also contain one or more substituted sugar moieties. The iRNAs, *e.g.*, dsRNAs, featured herein can include one of the following at the 2'-position: OH; F; O-, S-, or N-alkyl; O-, S-, or N-alkenyl; O-, S- or N-alkynyl; or O-alkyl-O-alkyl, wherein the alkyl, alkenyl and alkynyl can be substituted or unsubstituted C₁ to C₁₀ alkyl or C₂ to C₁₀ alkenyl and alkynyl. Exemplary suitable modifications include O[(CH₂)_nO]_mCH₃, O(CH₂)_nOCH₃, O(CH₂)_nNH₂, O(CH₂)_nCH₃, O(CH₂)_nONH₂, and O(CH₂)_nON[(CH₂)_nCH₃]₂, where n and m are from 1 to about 10. In other embodiments, dsRNAs include one of the following at the 2' position: C₁ to C₁₀ lower alkyl, substituted lower alkyl, alkaryl, aralkyl, O-alkaryl or O-aralkyl, SH, SCH₃, OCN, Cl, Br, CN, CF₃, OCF₃, SOCH₃, SO₂CH₃, ONO₂, NO₂, N₃, NH₂, heterocycloalkyl, heterocycloalkaryl, aminoalkylamino, polyalkylamino, substituted silyl, an RNA cleaving group, a reporter group, an intercalator, a group for improving the pharmacokinetic properties of an iRNA, or a group for improving the pharmacodynamic properties of an iRNA, and other substituents having similar properties. In some embodiments, the modification includes a 2'-methoxyethoxy (2'-O--CH₂CH₂OCH₃, also known as 2'-O-(2-methoxyethyl) or 2'-MOE) (Martin *et al.*, *Helv. Chim. Acta*, 1995, 78:486-504) *i.e.*, an alkoxy-alkoxy group. Another exemplary modification is 2'-dimethylaminoethoxyethoxy, *i.e.*, a O(CH₂)₂ON(CH₃)₂ group, also known as 2'-DMAOE, as described in examples herein below, and 2'-dimethylaminoethoxyethyl (also known in the art as 2'-O-dimethylaminoethoxyethyl or 2'-DMAEOE), *i.e.*, 2'-O--CH₂--O--CH₂--N(CH₃)₂. Further exemplary modifications include : 5'-Me-2'-F nucleotides, 5'-Me-2'-OMe nucleotides, 5'-Me-2'-deoxynucleotides, (both R and S isomers in these three families); 2'-alkoxyalkyl; and 2'-NMA (N-methylacetamide).

Other modifications include 2'-methoxy (2'-OCH₃), 2'-aminopropoxy (2'-OCH₂CH₂CH₂NH₂) and 2'-fluoro (2'-F). Similar modifications can also be made at other positions on the RNA of an iRNA, particularly the 3' position of the sugar on the 3' terminal nucleotide or in 2'-5' linked dsRNAs and the 5' position of 5' terminal nucleotide. iRNAs can also have sugar mimetics such as cyclobutyl moieties in place of the pentofuranosyl sugar. Representative US patents that teach the preparation of such modified sugar structures include, but are not limited to, U.S. Patent Nos. 4,981,957; 5,118,800; 5,319,080; 5,359,044; 5,393,878; 5,446,137; 5,466,786; 5,514,785; 5,519,134; 5,567,811; 5,576,427; 5,591,722; 5,597,909; 5,610,300; 5,627,053; 5,639,873; 5,646,265; 5,658,873; 5,670,633; and 5,700,920, certain of which are commonly owned with the instant application,. The entire contents of each of the foregoing are hereby incorporated herein by reference.

An iRNA can also include nucleobase (often referred to in the art simply as “base”) modifications or substitutions. As used herein, “unmodified” or “natural” nucleobases include the purine bases adenine (A) and guanine (G), and the pyrimidine bases thymine (T), cytosine (C), and uracil (U). Modified nucleobases include other synthetic and natural nucleobases such as

5 deoxythymidine (dT), 5-methylcytosine (5-me-C), 5-hydroxymethyl cytosine, xanthine, hypoxanthine, 2-aminoadenine, 6-methyl and other alkyl derivatives of adenine and guanine, 2-propyl and other alkyl derivatives of adenine and guanine, 2-thiouracil, 2-thiothymine and 2-thiocytosine, 5-halouracil and cytosine, 5-propynyl uracil and cytosine, 6-azo uracil, cytosine and thymine, 5-uracil (pseudouracil), 4-thiouracil, 8-halo, 8-amino, 8-thiol, 8-thioalkyl, 8-hydroxyl and other 8-substituted

10 adenines and guanines, 5-halo, particularly 5-bromo, 5-trifluoromethyl and other 5-substituted uracils and cytosines, 7-methylguanine and 7-methyladenine, 8-azaguanine and 8-azaadenine, 7-deazaguanine and 7-daazaadenine and 3-deazaguanine and 3-deazaadenine. Further nucleobases include those disclosed in U.S. Pat. No. 3,687,808, those disclosed in Modified Nucleosides in Biochemistry, Biotechnology and Medicine, Herdewijn, P. ed. Wiley-VCH, 2008; those disclosed in

15 The Concise Encyclopedia Of Polymer Science And Engineering, pages 858-859, Kroschwitz, J. L, ed. John Wiley & Sons, 1990, those disclosed by Englisch *et al.*, Angewandte Chemie, International Edition, 1991, 30, 613, and those disclosed by Sanghvi, Y S., Chapter 15, dsRNA Research and Applications, pages 289-302, Crooke, S. T. and Lebleu, B., Ed., CRC Press, 1993. Certain of these nucleobases are particularly useful for increasing the binding affinity of the oligomeric compounds

20 featured in the invention. These include 5-substituted pyrimidines, 6-azapyrimidines and N-2, N-6 and 0-6 substituted purines, including 2-aminopropyladenine, 5-propynyluracil and 5-propynylcytosine. 5-methylcytosine substitutions have been shown to increase nucleic acid duplex stability by 0.6-1.2°C (Sanghvi, Y. S., Crooke, S. T. and Lebleu, B., Eds., dsRNA Research and Applications, CRC Press, Boca Raton, 1993, pp. 276-278) and are exemplary base substitutions, even more particularly when

25 combined with 2'-O-methoxyethyl sugar modifications.

Representative U.S. Patents that teach the preparation of certain of the above noted modified nucleobases as well as other modified nucleobases include, but are not limited to, the above noted U.S. Patent Nos. 3,687,808, 4,845,205; 5,130,30; 5,134,066; 5,175,273; 5,367,066; 5,432,272; 5,457,187; 5,459,255; 5,484,908; 5,502,177; 5,525,711; 5,552,540; 5,587,469; 5,594,121, 5,596,091;


30 5,614,617; 5,681,941; 5,750,692; 6,015,886; 6,147,200; 6,166,197; 6,222,025; 6,235,887; 6,380,368; 6,528,640; 6,639,062; 6,617,438; 7,045,610; 7,427,672; and 7,495,088, the entire contents of each of which are hereby incorporated herein by reference.

In some embodiments, an RNAi agent of the disclosure can also be modified to include one or more bicyclic sugar moieties. A “bicyclic sugar” is a furanosyl ring modified by a ring formed by the

35 bridging of two carbons, whether adjacent or non-adjacent. A “bicyclic nucleoside” (“BNA”) is a nucleoside having a sugar moiety comprising a ring formed by bridging two carbons, whether adjacent or non-adjacent, of the sugar ring, thereby forming a bicyclic ring system. In certain

embodiments, the bridge connects the 4'-carbon and the 2'-carbon of the sugar ring, optionally, via the 2'-acyclic oxygen atom. Thus, in some embodiments an agent of the invention may include one or more locked nucleic acids (LNA). A locked nucleic acid is a nucleotide having a modified ribose moiety in which the ribose moiety comprises an extra bridge connecting the 2' and 4' carbons. In other words, an LNA is a nucleotide comprising a bicyclic sugar moiety comprising a 4'-CH₂-O-2' bridge. This structure effectively "locks" the ribose in the 3'-endo structural conformation. The addition of locked nucleic acids to siRNAs has been shown to increase siRNA stability in serum, and to reduce off-target effects (Elmen, J. *et al.*, (2005) *Nucleic Acids Research* 33(1):439-447; Mook, OR. *et al.*, (2007) *Mol Canc Ther* 6(3):833-843; Grunweller, A. *et al.*, (2003) *Nucleic Acids Research* 31(12):3185-3193). Examples of bicyclic nucleosides for use in the polynucleotides of the invention include without limitation nucleosides comprising a bridge between the 4' and the 2' ribosyl ring atoms. In certain embodiments, the antisense polynucleotide agents of the invention include one or more bicyclic nucleosides comprising a 4' to 2' bridge.

A locked nucleoside can be represented by the structure (omitting stereochemistry),



wherein B is a nucleobase or modified nucleobase and L is the linking group that joins the 2'-carbon to the 4'-carbon of the ribose ring. Examples of such 4' to 2' bridged bicyclic nucleosides, include but are not limited to 4'-(CH₂)—O-2' (LNA); 4'-(CH₂)—S-2'; 4'-(CH₂)₂—O-2' (ENA); 4'-CH(CH₃)—O-2' (also referred to as "constrained ethyl" or "cEt") and 4'-CH(CH₂OCH₃)—O-2' (and analogs thereof; see, *e.g.*, U.S. Patent No. 7,399,845); 4'-C(CH₃)(CH₃)—O-2' (and analogs thereof; see *e.g.*, U.S. Patent No. 8,278,283); 4'-CH₂—N(OCH₃)-2' (and analogs thereof; see *e.g.*, U.S. Patent No. 8,278,425); 4'-CH₂—O—N(CH₃)-2' (see, *e.g.*, U.S. Patent Publication No. 2004/0171570); 4'-CH₂—N(R)—O-2', wherein R is H, C1-C12 alkyl, or a nitrogen protecting group (see, *e.g.*, U.S. Patent No. 7,427,672); 4'-CH₂—C(H)(CH₃)-2' (see, *e.g.*, Chattopadhyaya *et al.*, *J. Org. Chem.*, 2009, 74, 118-134); and 4'-CH₂—C(=CH₂)-2' (and analogs thereof; see, *e.g.*, U.S. Patent No. 8,278,426). The entire contents of each of the foregoing are hereby incorporated herein by reference.

Additional representative U.S. Patents and U.S. Patent Publications that teach the preparation of locked nucleic acid nucleotides include, but are not limited to, the following: U.S. Patent Nos. 6,268,490; 6,525,191; 6,670,461; 6,770,748; 6,794,499; 6,998,484; 7,053,207; 7,034,133; 7,084,125; 7,399,845; 7,427,672; 7,569,686; 7,741,457; 8,022,193; 8,030,467; 8,278,425; 8,278,426; 8,278,283; US 2008/0039618; and US 2009/0012281, the entire contents of each of which are hereby incorporated herein by reference.

Any of the foregoing bicyclic nucleosides can be prepared having one or more stereochemical sugar configurations including for example α -L-ribofuranose and β -D-ribofuranose (see WO 99/14226).

The RNA of an iRNA can also be modified to include one or more constrained ethyl nucleotides. As used herein, a "constrained ethyl nucleotide" or "cEt" is a locked nucleic acid comprising a bicyclic sugar moiety comprising a 4'-CH(CH₃)-O-2' bridge (*i.e.*, L in the preceding structure). In one embodiment, a constrained ethyl nucleotide is in the S conformation referred to herein as "S-cEt."

An iRNA of the invention may also include one or more "conformationally restricted nucleotides" ("CRN"). CRN are nucleotide analogs with a linker connecting the C2' and C4' carbons of ribose or the C3 and -C5' carbons of ribose. CRN lock the ribose ring into a stable conformation and increase the hybridization affinity to mRNA. The linker is of sufficient length to place the oxygen in an optimal position for stability and affinity resulting in less ribose ring puckering.

Representative publications that teach the preparation of certain of the above noted CRN include, but are not limited to, U.S. Patent Publication No. 2013/0190383; and PCT publication WO 2013/036868, the entire contents of each of which are hereby incorporated herein by reference.

In some embodiments, an iRNA of the invention comprises one or more monomers that are UNA (unlocked nucleic acid) nucleotides. UNA is unlocked acyclic nucleic acid, wherein any of the bonds of the sugar has been removed, forming an unlocked "sugar" residue. In one example, UNA also encompasses monomer with bonds between C1'-C4' have been removed (*i.e.* the covalent carbon-oxygen-carbon bond between the C1' and C4' carbons). In another example, the C2'-C3' bond (*i.e.* the covalent carbon-carbon bond between the C2' and C3' carbons) of the sugar has been removed (see *Nuc. Acids Symp. Series*, 52, 133-134 (2008) and Fluiters *et al.*, *Mol. Biosyst.*, 2009, 10, 1039 hereby incorporated by reference).

Representative U.S. publications that teach the preparation of UNA include, but are not limited to, U.S. Patent No. 8,314,227; and U.S. Patent Publication Nos. 2013/0096289; 2013/0011922; and 2011/0313020, the entire contents of each of which are hereby incorporated herein by reference.

Potentially stabilizing modifications to the ends of RNA molecules can include N-(acetylaminocaproyl)-4-hydroxyprolinol (Hyp-C6-NHAc), N-(caproyl-4-hydroxyprolinol (Hyp-C6), N-(acetyl-4-hydroxyprolinol (Hyp-NHAc), thymidine-2'-O-deoxythymidine (ether), N-(aminocaproyl)-4-hydroxyprolinol (Hyp-C6-amino), 2-docosanoyl-uridine-3'-phosphate, inverted 2'-deoxy-modified ribonucleotide, such as inverted dT(idT), inverted dA (idA), and inverted abasic 2'-deoxyribonucleotide (iAb) and others. Disclosure of this modification can be found in WO 2011/005861.

In one example, the 3' or 5' terminal end of a oligonucleotide is linked to an inverted 2'-deoxy-modified ribonucleotide, such as inverted dT(idT), inverted dA (idA), or a inverted abasic 2'-

deoxyribonucleotide (iAb). In one particular example, the inverted 2'-deoxy-modified ribonucleotide is linked to the 3'-end of an oligonucleotide, such as the 3'-end of a sense strand described herein, where the linking is *via* a 3'-3' phosphodiester linkage or a 3'-3'-phosphorothioate linkage.

In another example, the 3'-end of a sense strand is linked *via* a 3'-3'-phosphorothioate linkage to an inverted abasic ribonucleotide (iAb). In another example, the 3'-end of a sense strand is linked *via* a 3'-3'-phosphorothioate linkage to an inverted dA (idA).

In one particular example, the inverted 2'-deoxy-modified ribonucleotide is linked to the 3'-end of an oligonucleotide, such as the 3'-end of a sense strand described herein, where the linking is *via* a 3'-3' phosphodiester linkage or a 3'-3'-phosphorothioate linkage.

In another example, the 3'-terminal nucleotides of a sense strand is an inverted dA (idA) and is linked to the preceding nucleotide *via* a 3'-3'- linkage (*e.g.*, 3'-3'-phosphorothioate linkage).

Other modifications of the nucleotides of an iRNA of the invention include a 5' phosphate or 5' phosphate mimic, *e.g.*, a 5'-terminal phosphate or phosphate mimic on the antisense strand of an iRNA. Suitable phosphate mimics are disclosed in, for example U.S. Patent Publication No. 2012/0157511, the entire contents of which are incorporated herein by reference.

A. Modified iRNAs Comprising Motifs of the Invention

In certain aspects of the invention, the double stranded RNA agents of the invention include agents with chemical modifications as disclosed, for example, in WO2013/075035, the entire contents of each of which are incorporated herein by reference. As shown herein and in WO2013/075035, one or more motifs of three identical modifications on three consecutive nucleotides may be introduced into a sense strand or antisense strand of a dsRNAi agent, particularly at or near the cleavage site. In some embodiments, the sense strand and antisense strand of the dsRNAi agent may otherwise be completely modified. The introduction of these motifs interrupts the modification pattern, if present, of the sense or antisense strand. The dsRNAi agent may be optionally conjugated with a GalNAc derivative ligand, for instance on the sense strand.

More specifically, when the sense strand and antisense strand of the double stranded RNA agent are completely modified to have one or more motifs of three identical modifications on three consecutive nucleotides at or near the cleavage site of at least one strand of a dsRNAi agent, the gene silencing activity of the dsRNAi agent was observed.

Accordingly, the invention provides double stranded RNA agents capable of inhibiting the expression of a target gene (*i.e.*, F12 gene) *in vivo*. The RNAi agent comprises a sense strand and an antisense strand. Each strand of the RNAi agent may be, for example, 17-30 nucleotides in length, 25-30 nucleotides in length, 27-30 nucleotides in length, 19-25 nucleotides in length, 19-23 nucleotides in length, 19-21 nucleotides in length, 21-25 nucleotides in length, or 21-23 nucleotides in length.

The sense strand and antisense strand typically form a duplex double stranded RNA (“dsRNA”), also referred to herein as “dsRNAi agent.” The duplex region of a dsRNAi agent may be, for example, the duplex region can be 27-30 nucleotide pairs in length, 19-25 nucleotide pairs in length, 19-23 nucleotide pairs in length, 19- 21 nucleotide pairs in length, 21-25 nucleotide pairs in length, or 21-23 nucleotide pairs in length. In another example, the duplex region is selected from 19, 20, 21, 22, 23, 24, 25, 26, and 27 nucleotides in length.

In certain embodiments, the dsRNAi agent may contain one or more overhang regions or capping groups at the 3'-end, 5'-end, or both ends of one or both strands. The overhang can be, independently, 1-6 nucleotides in length, for instance 2-6 nucleotides in length, 1-5 nucleotides in length, 2-5 nucleotides in length, 1-4 nucleotides in length, 2-4 nucleotides in length, 1-3 nucleotides in length, 2-3 nucleotides in length, or 1-2 nucleotides in length. In certain embodiments, the overhang regions can include extended overhang regions as provided above. The overhangs can be the result of one strand being longer than the other, or the result of two strands of the same length being staggered. The overhang can form a mismatch with the target mRNA or it can be complementary to the gene sequences being targeted or can be another sequence. The first and second strands can also be joined, *e.g.*, by additional bases to form a hairpin, or by other non-base linkers.

In certain embodiments, the nucleotides in the overhang region of the dsRNAi agent can each independently be a modified or unmodified nucleotide including, but not limited to 2'-sugar modified, such as, 2'-F, 2'-O-methyl, thymidine (T), 2'-O-methoxyethyl-5-methyluridine (Teo), 2'-O-methoxyethyladenosine (Aeo), 2'-O-methoxyethyl-5-methylcytidine (m5Ceo), and any combinations thereof.

For example, TT can be an overhang sequence for either end on either strand. The overhang can form a mismatch with the target mRNA or it can be complementary to the gene sequences being targeted or can be another sequence.

The 5'- or 3'- overhangs at the sense strand, antisense strand, or both strands of the dsRNAi agent may be phosphorylated. In some embodiments, the overhang region(s) contains two nucleotides having a phosphorothioate between the two nucleotides, where the two nucleotides can be the same or different. In some embodiments, the overhang is present at the 3'-end of the sense strand, antisense strand, or both strands. In some embodiments, this 3'-overhang is present in the antisense strand. In some embodiments, this 3'-overhang is present in the sense strand.

The dsRNAi agent may contain only a single overhang, which can strengthen the interference activity of the RNAi, without affecting its overall stability. For example, the single-stranded overhang may be located at the 3'- end of the sense strand or, alternatively, at the 3'-end of the antisense strand. The RNAi may also have a blunt end, located at the 5'-end of the antisense strand (*i.e.*, the 3'-end of the sense strand) or *vice versa*. Generally, the antisense strand of the dsRNAi agent has a nucleotide overhang at the 3'-end, and the 5'-end is blunt. While not wishing to be bound by

theory, the asymmetric blunt end at the 5'-end of the antisense strand and 3'-end overhang of the antisense strand favor the guide strand loading into RISC process.

In certain embodiments, the dsRNAi agent is a double blunt-ended of 19 nucleotides in length, wherein the sense strand contains at least one motif of three 2'-F modifications on three consecutive nucleotides at positions 7, 8, 9 from the 5'-end. The antisense strand contains at least one motif of three 2'-O-methyl modifications on three consecutive nucleotides at positions 11, 12, and 13 from the 5'-end.

In other embodiments, the dsRNAi agent is a double blunt-ended of 20 nucleotides in length, wherein the sense strand contains at least one motif of three 2'-F modifications on three consecutive nucleotides at positions 8, 9, and 10 from the 5'-end. The antisense strand contains at least one motif of three 2'-O-methyl modifications on three consecutive nucleotides at positions 11, 12, and 13 from the 5'-end.

In yet other embodiments, the dsRNAi agent is a double blunt-ended of 21 nucleotides in length, wherein the sense strand contains at least one motif of three 2'-F modifications on three consecutive nucleotides at positions 9, 10, and 11 from the 5'-end. The antisense strand contains at least one motif of three 2'-O-methyl modifications on three consecutive nucleotides at positions 11, 12, and 13 from the 5'-end.

In certain embodiments, the dsRNAi agent comprises a 21 nucleotide sense strand and a 23 nucleotide antisense strand, wherein the sense strand contains at least one motif of three 2'-F modifications on three consecutive nucleotides at positions 9, 10, and 11 from the 5'-end; the antisense strand contains at least one motif of three 2'-O-methyl modifications on three consecutive nucleotides at positions 11, 12, and 13 from the 5'-end, wherein one end of the RNAi agent is blunt, while the other end comprises a 2 nucleotide overhang. In one embodiment, the 2 nucleotide overhang is at the 3'-end of the antisense strand.

When the 2 nucleotide overhang is at the 3'-end of the antisense strand, there may be two phosphorothioate internucleotide linkages between the terminal three nucleotides, wherein two of the three nucleotides are the overhang nucleotides, and the third nucleotide is a paired nucleotide next to the overhang nucleotide. In one embodiment, the RNAi agent additionally has two phosphorothioate internucleotide linkages between the terminal three nucleotides at both the 5'-end of the sense strand and at the 5'-end of the antisense strand. In certain embodiments, every nucleotide in the sense strand and the antisense strand of the dsRNAi agent, including the nucleotides that are part of the motifs are modified nucleotides. In certain embodiments each residue is independently modified with a 2'-O-methyl or 3'-fluoro, *e.g.*, in an alternating motif. Optionally, the dsRNAi agent further comprises a ligand (such as, GalNAc₃).

In certain embodiments, the dsRNAi agent comprises a sense and an antisense strand, wherein the sense strand is 25-30 nucleotide residues in length, wherein starting from the 5' terminal nucleotide (position 1) positions 1 to 23 of the first strand comprise at least 8 ribonucleotides; the

antisense strand is 36-66 nucleotide residues in length and, starting from the 3' terminal nucleotide, comprises at least 8 ribonucleotides in the positions paired with positions 1- 23 of sense strand to form a duplex; wherein at least the 3' terminal nucleotide of antisense strand is unpaired with sense strand, and up to 6 consecutive 3' terminal nucleotides are unpaired with sense strand, thereby forming a 3' single stranded overhang of 1-6 nucleotides; wherein the 5' terminus of antisense strand comprises from 10-30 consecutive nucleotides which are unpaired with sense strand, thereby forming a 10-30 nucleotide single stranded 5' overhang; wherein at least the sense strand 5' terminal and 3' terminal nucleotides are base paired with nucleotides of antisense strand when sense and antisense strands are aligned for maximum complementarity, thereby forming a substantially duplexed region between sense and antisense strands; and antisense strand is sufficiently complementary to a target RNA along at least 19 ribonucleotides of antisense strand length to reduce target gene expression when the double stranded nucleic acid is introduced into a mammalian cell; and wherein the sense strand contains at least one motif of three 2'-F modifications on three consecutive nucleotides, where at least one of the motifs occurs at or near the cleavage site. The antisense strand contains at least one motif of three 2'-O-methyl modifications on three consecutive nucleotides at or near the cleavage site.

In certain embodiments, the dsRNAi agent comprises sense and antisense strands, wherein the dsRNAi agent comprises a first strand having a length which is at least 25 and at most 29 nucleotides and a second strand having a length which is at most 30 nucleotides with at least one motif of three 2'-O-methyl modifications on three consecutive nucleotides at position 11, 12, 13 from the 5' end; wherein the 3' end of the first strand and the 5' end of the second strand form a blunt end and the second strand is 1-4 nucleotides longer at its 3' end than the first strand, wherein the duplex region which is at least 25 nucleotides in length, and the second strand is sufficiently complementary to a target mRNA along at least 19 nucleotide of the second strand length to reduce target gene expression when the RNAi agent is introduced into a mammalian cell, and wherein Dicer cleavage of the dsRNAi agent results in an siRNA comprising the 3'-end of the second strand, thereby reducing expression of the target gene in the mammal. Optionally, the dsRNAi agent further comprises a ligand.

In certain embodiments, the sense strand of the dsRNAi agent contains at least one motif of three identical modifications on three consecutive nucleotides, where one of the motifs occurs at the cleavage site in the sense strand.

In certain embodiments, the antisense strand of the dsRNAi agent can also contain at least one motif of three identical modifications on three consecutive nucleotides, where one of the motifs occurs at or near the cleavage site in the antisense strand.

For a dsRNAi agent having a duplex region of 19-23 nucleotides in length, the cleavage site of the antisense strand is typically around the 10, 11, and 12 positions from the 5'-end. Thus the motifs of three identical modifications may occur at the 9, 10, 11 positions; the 10, 11, 12 positions; the 11, 12, 13 positions; the 12, 13, 14 positions; or the 13, 14, 15 positions of the antisense strand, the count starting from the first nucleotide from the 5'-end of the antisense strand, or, the count starting

from the first paired nucleotide within the duplex region from the 5'-end of the antisense strand. The cleavage site in the antisense strand may also change according to the length of the duplex region of the dsRNAi agent from the 5'-end.

The sense strand of the dsRNAi agent may contain at least one motif of three identical
5 modifications on three consecutive nucleotides at the cleavage site of the strand; and the antisense strand may have at least one motif of three identical modifications on three consecutive nucleotides at or near the cleavage site of the strand. When the sense strand and the antisense strand form a dsRNA duplex, the sense strand and the antisense strand can be so aligned that one motif of the three
10 nucleotides on the sense strand and one motif of the three nucleotides on the antisense strand have at least one nucleotide overlap, *i.e.*, at least one of the three nucleotides of the motif in the sense strand forms a base pair with at least one of the three nucleotides of the motif in the antisense strand. Alternatively, at least two nucleotides may overlap, or all three nucleotides may overlap.

In some embodiments, the sense strand of the dsRNAi agent may contain more than one motif of three identical modifications on three consecutive nucleotides. The first motif may occur at or near
15 the cleavage site of the strand and the other motifs may be a wing modification. The term "wing modification" herein refers to a motif occurring at another portion of the strand that is separated from the motif at or near the cleavage site of the same strand. The wing modification is either adjacent to the first motif or is separated by at least one or more nucleotides. When the motifs are immediately adjacent to each other then the chemistries of the motifs are distinct from each other, and when the
20 motifs are separated by one or more nucleotide then the chemistries can be the same or different. Two or more wing modifications may be present. For instance, when two wing modifications are present, each wing modification may occur at one end relative to the first motif which is at or near cleavage site or on either side of the lead motif.

Like the sense strand, the antisense strand of the dsRNAi agent may contain more than one
25 motif of three identical modifications on three consecutive nucleotides, with at least one of the motifs occurring at or near the cleavage site of the strand. This antisense strand may also contain one or more wing modifications in an alignment similar to the wing modifications that may be present on the sense strand.

In some embodiments, the wing modification on the sense strand or antisense strand of the
30 dsRNAi agent typically does not include the first one or two terminal nucleotides at the 3'-end, 5'-end, or both ends of the strand.

In other embodiments, the wing modification on the sense strand or antisense strand of the dsRNAi agent typically does not include the first one or two paired nucleotides within the duplex region at the 3'-end, 5'-end, or both ends of the strand.

35 When the sense strand and the antisense strand of the dsRNAi agent each contain at least one wing modification, the wing modifications may fall on the same end of the duplex region, and have an overlap of one, two, or three nucleotides.

When the sense strand and the antisense strand of the dsRNAi agent each contain at least two wing modifications, the sense strand and the antisense strand can be so aligned that two modifications each from one strand fall on one end of the duplex region, having an overlap of one, two, or three nucleotides; two modifications each from one strand fall on the other end of the duplex region, having an overlap of one, two or three nucleotides; two modifications one strand fall on each side of the lead motif, having an overlap of one, two or three nucleotides in the duplex region.

In some embodiments, every nucleotide in the sense strand and antisense strand of the dsRNAi agent, including the nucleotides that are part of the motifs, may be modified. Each nucleotide may be modified with the same or different modification which can include one or more alteration of one or both of the non-linking phosphate oxygens or of one or more of the linking phosphate oxygens; alteration of a constituent of the ribose sugar, *e.g.*, of the 2'-hydroxyl on the ribose sugar; wholesale replacement of the phosphate moiety with "dephospho" linkers; modification or replacement of a naturally occurring base; and replacement or modification of the ribose-phosphate backbone.

As nucleic acids are polymers of subunits, many of the modifications occur at a position which is repeated within a nucleic acid, *e.g.*, a modification of a base, or a phosphate moiety, or a non-linking O of a phosphate moiety. In some cases the modification will occur at all of the subject positions in the nucleic acid but in many cases it will not. By way of example, a modification may only occur at a 3'- or 5' terminal position, may only occur in a terminal region, *e.g.*, at a position on a terminal nucleotide or in the last 2, 3, 4, 5, or 10 nucleotides of a strand. A modification may occur in a double strand region, a single strand region, or in both. A modification may occur only in the double strand region of an RNA or may only occur in a single strand region of a RNA. For example, a phosphorothioate modification at a non-linking O position may only occur at one or both termini, may only occur in a terminal region, *e.g.*, at a position on a terminal nucleotide or in the last 2, 3, 4, 5, or 10 nucleotides of a strand, or may occur in double strand and single strand regions, particularly at termini. The 5'-end or ends can be phosphorylated.

It may be possible, *e.g.*, to enhance stability, to include particular bases in overhangs, or to include modified nucleotides or nucleotide surrogates, in single strand overhangs, *e.g.*, in a 5'- or 3'-overhang, or in both. For example, it can be desirable to include purine nucleotides in overhangs. In some embodiments all or some of the bases in a 3'- or 5'-overhang may be modified, *e.g.*, with a modification described herein. Modifications can include, *e.g.*, the use of modifications at the 2' position of the ribose sugar with modifications that are known in the art, *e.g.*, the use of deoxyribonucleotides, 2'-deoxy-2'-fluoro (2'-F) or 2'-O-methyl modified instead of the ribosugar of the nucleobase, and modifications in the phosphate group, *e.g.*, phosphorothioate modifications.

Overhangs need not be homologous with the target sequence.

In some embodiments, each residue of the sense strand and antisense strand is independently modified with LNA, CRN, cET, UNA, HNA, CeNA, 2'-methoxyethyl, 2'-O-methyl, 2'-O-allyl, 2'-

C- allyl, 2'-deoxy, 2'-hydroxyl, or 2'-fluoro. The strands can contain more than one modification. In one embodiment, each residue of the sense strand and antisense strand is independently modified with 2'-O-methyl or 2'-fluoro.

At least two different modifications are typically present on the sense strand and antisense strand. Those two modifications may be the 2'-O-methyl or 2'-fluoro modifications, or others.

In certain embodiments, the N_a or N_b comprise modifications of an alternating pattern. The term "alternating motif" as used herein refers to a motif having one or more modifications, each modification occurring on alternating nucleotides of one strand. The alternating nucleotide may refer to one per every other nucleotide or one per every three nucleotides, or a similar pattern. For example, if A, B and C each represent one type of modification to the nucleotide, the alternating motif can be "ABABABABABAB...", "AABBAABBAABB...", "AABAABAABAAB...", "AAABAAABAAAB...", "AAABBBAAABBB...", or "ABCABCABCABC...", etc.

The type of modifications contained in the alternating motif may be the same or different. For example, if A, B, C, D each represent one type of modification on the nucleotide, the alternating pattern, *i.e.*, modifications on every other nucleotide, may be the same, but each of the sense strand or antisense strand can be selected from several possibilities of modifications within the alternating motif such as "ABABAB...", "ACACAC...", "BDBDBD..." or "CDCDCD...", etc.

In some embodiments, the dsRNAi agent of the invention comprises the modification pattern for the alternating motif on the sense strand relative to the modification pattern for the alternating motif on the antisense strand is shifted. The shift may be such that the modified group of nucleotides of the sense strand corresponds to a differently modified group of nucleotides of the antisense strand and *vice versa*. For example, the sense strand when paired with the antisense strand in the dsRNA duplex, the alternating motif in the sense strand may start with "ABABAB" from 5' to 3' of the strand and the alternating motif in the antisense strand may start with "BABABA" from 5' to 3' of the strand within the duplex region. As another example, the alternating motif in the sense strand may start with "AABBAABB" from 5' to 3' of the strand and the alternating motif in the antisense strand may start with "BBAABBAA" from 5' to 3' of the strand within the duplex region, so that there is a complete or partial shift of the modification patterns between the sense strand and the antisense strand.

In one particular example, the alternating motif in the sense strand is "ABABAB" from 5' to 3' of the strand, where each A is an unmodified ribonucleotide and each B is a 2'-O-methyl modified nucleotide.

In one particular example, the alternating motif in the sense strand is "ABABAB" from 5' to 3' of the strand, where each A is a 2'-deoxy-2'-fluoro modified nucleotide and each B is a 2'-O-methyl modified nucleotide.

In another particular example, the alternating motif in the antisense strand is "BABABA" from 3'-5' of the strand, where each A is a 2'-deoxy-2'-fluoro modified nucleotide and each B is a 2'-O-methyl modified nucleotide.

In one particular example, the alternating motif in the sense strand is “ABABAB” from 5’-3’ of the strand and the alternating motif in the antisense strand is “BABABA” from 3’-5’ of the strand, where each A is an unmodified ribonucleotide and each B is a 2’-O-methyl modified nucleotide.

In one particular example, the alternating motif in the sense strand is “ABABAB” from 5’-3’ of the strand and the alternating motif in the antisense strand is “BABABA” from 3’-5’ of the strand, where each A is a 2’-deoxy-2’-fluoro modified nucleotide and each B is a 2’-O-methyl modified nucleotide.

In some embodiments, the dsRNAi agent comprises the pattern of the alternating motif of 2’-O-methyl modification and 2’-F modification on the sense strand initially has a shift relative to the pattern of the alternating motif of 2’-O-methyl modification and 2’-F modification on the antisense strand initially, *i.e.*, the 2’-O-methyl modified nucleotide on the sense strand base pairs with a 2’-F modified nucleotide on the antisense strand and vice versa. The 1 position of the sense strand may start with the 2’-F modification, and the 1 position of the antisense strand may start with the 2’-O-methyl modification.

The introduction of one or more motifs of three identical modifications on three consecutive nucleotides to the sense strand or antisense strand interrupts the initial modification pattern present in the sense strand or antisense strand. This interruption of the modification pattern of the sense or antisense strand by introducing one or more motifs of three identical modifications on three consecutive nucleotides to the sense or antisense strand may enhance the gene silencing activity against the target gene.

In some embodiments, when the motif of three identical modifications on three consecutive nucleotides is introduced to any of the strands, the modification of the nucleotide next to the motif is a different modification than the modification of the motif. For example, the portion of the sequence containing the motif is “...N_aYYYN_b...,” where “Y” represents the modification of the motif of three identical modifications on three consecutive nucleotide, and “N_a” and “N_b” represent a modification to the nucleotide next to the motif “YYY” that is different than the modification of Y, and where N_a and N_b can be the same or different modifications. Alternatively, N_a or N_b may be present or absent when there is a wing modification present.

The iRNA may further comprise at least one phosphorothioate or methylphosphonate internucleotide linkage. The phosphorothioate or methylphosphonate internucleotide linkage modification may occur on any nucleotide of the sense strand, antisense strand, or both strands in any position of the strand. For instance, the internucleotide linkage modification may occur on every nucleotide on the sense strand or antisense strand; each internucleotide linkage modification may occur in an alternating pattern on the sense strand or antisense strand; or the sense strand or antisense strand may contain both internucleotide linkage modifications in an alternating pattern. The alternating pattern of the internucleotide linkage modification on the sense strand may be the same or different from the antisense strand, and the alternating pattern of the internucleotide linkage

modification on the sense strand may have a shift relative to the alternating pattern of the internucleotide linkage modification on the antisense strand. In one embodiment, a double-stranded RNAi agent comprises 6-8 phosphorothioate internucleotide linkages. In some embodiments, the antisense strand comprises two phosphorothioate internucleotide linkages at the 5'-end and two
5 phosphorothioate internucleotide linkages at the 3'-end, and the sense strand comprises at least two phosphorothioate internucleotide linkages at either the 5'-end or the 3'-end.

In some embodiments, the dsRNAi agent comprises a phosphorothioate or methylphosphonate internucleotide linkage modification in the overhang region. For example, the overhang region may contain two nucleotides having a phosphorothioate or methylphosphonate
10 internucleotide linkage between the two nucleotides. Internucleotide linkage modifications also may be made to link the overhang nucleotides with the terminal paired nucleotides within the duplex region. For example, at least 2, 3, 4, or all the overhang nucleotides may be linked through phosphorothioate or methylphosphonate internucleotide linkage, and optionally, there may be additional phosphorothioate or methylphosphonate internucleotide linkages linking the overhang
15 nucleotide with a paired nucleotide that is next to the overhang nucleotide. For instance, there may be at least two phosphorothioate internucleotide linkages between the terminal three nucleotides, in which two of the three nucleotides are overhang nucleotides, and the third is a paired nucleotide next to the overhang nucleotide. These terminal three nucleotides may be at the 3'-end of the antisense strand, the 3'-end of the sense strand, the 5'-end of the antisense strand, or the 5'-end of the antisense
20 strand.

In some embodiments, the 2-nucleotide overhang is at the 3'-end of the antisense strand, and there are two phosphorothioate internucleotide linkages between the terminal three nucleotides, wherein two of the three nucleotides are the overhang nucleotides, and the third nucleotide is a paired nucleotide next to the overhang nucleotide. Optionally, the dsRNAi agent may additionally have two
25 phosphorothioate internucleotide linkages between the terminal three nucleotides at both the 5'-end of the sense strand and at the 5'-end of the antisense strand.

In one embodiment, the dsRNAi agent comprises mismatch(es) with the target, within the duplex, or combinations thereof. The mismatch may occur in the overhang region or the duplex region. The base pair may be ranked on the basis of their propensity to promote dissociation or
30 melting (*e.g.*, on the free energy of association or dissociation of a particular pairing, the simplest approach is to examine the pairs on an individual pair basis, though next neighbor or similar analysis can also be used). In terms of promoting dissociation: A:U is preferred over G:C; G:U is preferred over G:C; and I:C is preferred over G:C (I=inosine). Mismatches, *e.g.*, non-canonical or other than canonical pairings (as described elsewhere herein) are preferred over canonical (A:T, A:U, G:C)
35 pairings; and pairings which include a universal base are preferred over canonical pairings.

In certain embodiments, the dsRNAi agent comprises at least one of the first 1, 2, 3, 4, or 5 base pairs within the duplex regions from the 5'-end of the antisense strand independently selected

from the group of: A:U, G:U, I:C, and mismatched pairs, *e.g.*, non-canonical or other than canonical pairings or pairings which include a universal base, to promote the dissociation of the antisense strand at the 5'-end of the duplex.

In certain embodiments, the nucleotide at the 1 position within the duplex region from the 5'-end in the antisense strand is selected from A, dA, dU, U, and dT. Alternatively, at least one of the first 1, 2, or 3 base pair within the duplex region from the 5'-end of the antisense strand is an AU base pair. For example, the first base pair within the duplex region from the 5'-end of the antisense strand is an AU base pair.

In other embodiments, the nucleotide at the 3'-end of the sense strand is deoxythymidine (dT) or the nucleotide at the 3'-end of the antisense strand is deoxythymidine (dT). For example, there is a short sequence of deoxythymidine nucleotides, for example, two dT nucleotides on the 3'-end of the sense, antisense strand, or both strands.

In certain embodiments, the sense strand sequence may be represented by formula (I):



wherein:

i and j are each independently 0 or 1;

p and q are each independently 0-6;

each N_a independently represents an oligonucleotide sequence comprising 0-25 modified nucleotides, each sequence comprising at least two differently modified nucleotides;

each N_b independently represents an oligonucleotide sequence comprising 0-10 modified nucleotides;

each n_p and n_q independently represent an overhang nucleotide;

wherein N_b and Y do not have the same modification; and

XXX, YYY, and ZZZ each independently represent one motif of three identical modifications on three consecutive nucleotides. In one embodiment, YYY is all 2'-F modified nucleotides.

In some embodiments, the N_a or N_b comprises modifications of alternating pattern.

In some embodiments, the YYY motif occurs at or near the cleavage site of the sense strand. For example, when the dsRNAi agent has a duplex region of 17-23 nucleotides in length, the YYY motif can occur at or the vicinity of the cleavage site (*e.g.*: can occur at positions 6, 7, 8; 7, 8, 9; 8, 9, 10; 9, 10, 11; 10, 11, 12; or 11, 12, 13) of the sense strand, the count starting from the first nucleotide, from the 5'-end; or optionally, the count starting at the first paired nucleotide within the duplex region, from the 5'-end.

In one embodiment, i is 1 and j is 0, or i is 0 and j is 1, or both i and j are 1. The sense strand can therefore be represented by the following formulas:



When the sense strand is represented by formula (Ib), N_b represents an oligonucleotide sequence comprising 0-10, 0-7, 0-5, 0-4, 0-2, or 0 modified nucleotides. Each N_a independently can represent an oligonucleotide sequence comprising 2-20, 2-15, or 2-10 modified nucleotides.

When the sense strand is represented as formula (Ic), N_b represents an oligonucleotide sequence comprising 0-10, 0-7, 0-10, 0-7, 0-5, 0-4, 0-2, or 0 modified nucleotides. Each N_a can independently represent an oligonucleotide sequence comprising 2-20, 2-15, or 2-10 modified nucleotides.

When the sense strand is represented as formula (Id), each N_b independently represents an oligonucleotide sequence comprising 0-10, 0-7, 0-5, 0-4, 0-2, or 0 modified nucleotides. In one embodiment, N_b is 0, 1, 2, 3, 4, 5, or 6. Each N_a can independently represent an oligonucleotide sequence comprising 2-20, 2-15, or 2-10 modified nucleotides.

Each of X, Y and Z may be the same or different from each other.

In other embodiments, i is 0 and j is 0, and the sense strand may be represented by the formula:

5' n_p - N_a -YYY- N_a - n_q 3' (Ia).

When the sense strand is represented by formula (Ia), each N_a independently can represent an oligonucleotide sequence comprising 2-20, 2-15, or 2-10 modified nucleotides.

In one embodiment, the antisense strand sequence of the RNAi may be represented by formula (II):

5' n_q - N_a' -(Z'Z'Z') $_k$ - N_b' -Y'Y'Y'- N_b' -(X'X'X') $_l$ - N_a' - n_p' 3' (II)

wherein:

k and l are each independently 0 or 1;

p' and q' are each independently 0-6;

each N_a' independently represents an oligonucleotide sequence comprising 0-25 modified nucleotides, each sequence comprising at least two differently modified nucleotides;

each N_b' independently represents an oligonucleotide sequence comprising 0-10 modified nucleotides;

each n_p' and n_q' independently represent an overhang nucleotide;

wherein N_b' and Y' do not have the same modification; and

X'X'X', Y'Y'Y', and Z'Z'Z' each independently represent one motif of three identical modifications on three consecutive nucleotides.

In some embodiments, the N_a' or N_b' comprises modifications of alternating pattern.

The Y'Y'Y' motif occurs at or near the cleavage site of the antisense strand. For example, when the dsRNAi agent has a duplex region of 17-23 nucleotides in length, the Y'Y'Y' motif can occur at positions 9, 10, 11; 10, 11, 12; 11, 12, 13; 12, 13, 14; or 13, 14, 15 of the antisense strand, with the count starting from the first nucleotide, from the 5'-end; or optionally, the count starting at

the first paired nucleotide within the duplex region, from the 5'-end. In one embodiment, the Y'Y'Y' motif occurs at positions 11, 12, 13.

In certain embodiments, Y'Y'Y' motif is all 2'-OMe modified nucleotides.

In certain embodiments, k is 1 and l is 0, or k is 0 and l is 1, or both k and l are 1.

5 The antisense strand can therefore be represented by the following formulas:

$5' n_q-N_a'-Z'Z'Z'-N_b'-Y'Y'Y'-N_a'-n_p, 3'$ (IIb);

$5' n_q-N_a'-Y'Y'Y'-N_b'-X'X'X'-n_p, 3'$ (IIc); or

$5' n_q-N_a'-Z'Z'Z'-N_b'-Y'Y'Y'-N_b'-X'X'X'-N_a'-n_p, 3'$ (IId).

10 When the antisense strand is represented by formula (IIb), N_b' represents an oligonucleotide sequence comprising 0-10, 0-7, 0-10, 0-7, 0-5, 0-4, 0-2, or 0 modified nucleotides. Each N_a' independently represents an oligonucleotide sequence comprising 2-20, 2-15, or 2-10 modified nucleotides.

15 When the antisense strand is represented as formula (IIc), N_b' represents an oligonucleotide sequence comprising 0-10, 0-7, 0-10, 0-7, 0-5, 0-4, 0-2, or 0 modified nucleotides. Each N_a' independently represents an oligonucleotide sequence comprising 2-20, 2-15, or 2-10 modified nucleotides.

20 When the antisense strand is represented as formula (IId), each N_b' independently represents an oligonucleotide sequence comprising 0-10, 0-7, 0-10, 0-7, 0-5, 0-4, 0-2, or 0 modified nucleotides. Each N_a' independently represents an oligonucleotide sequence comprising 2-20, 2-15, or 2-10 modified nucleotides. In one embodiment, N_b is 0, 1, 2, 3, 4, 5, or 6.

In other embodiments, k is 0 and l is 0 and the antisense strand may be represented by the formula:

$5' n_p-N_a'-Y'Y'Y'-N_a'-n_q, 3'$ (Ia).

25 When the antisense strand is represented as formula (IIa), each N_a' independently represents an oligonucleotide sequence comprising 2-20, 2-15, or 2-10 modified nucleotides.

Each of X', Y' and Z' may be the same or different from each other.

30 Each nucleotide of the sense strand and antisense strand may be independently modified with LNA, CRN, UNA, cEt, HNA, CeNA, 2'-methoxyethyl, 2'-O-methyl, 2'-O-allyl, 2'-C-allyl, 2'-hydroxyl, or 2'-fluoro. For example, each nucleotide of the sense strand and antisense strand is independently modified with 2'-O-methyl or 2'-fluoro. Each X, Y, Z, X', Y', and Z', in particular, may represent a 2'-O-methyl modification or a 2'-fluoro modification.


35 In some embodiments, the sense strand of the dsRNAi agent may contain YYY motif occurring at 9, 10, and 11 positions of the strand when the duplex region is 21 nt, the count starting from the first nucleotide from the 5'-end, or optionally, the count starting at the first paired nucleotide within the duplex region, from the 5'-end; and Y represents 2'-F modification. The sense strand may additionally contain XXX motif or ZZZ motifs as wing modifications at the opposite end of the

duplex region; and XXX and ZZZ each independently represents a 2'-OMe modification or 2'-F modification.

In some embodiments the antisense strand may contain Y'Y'Y' motif occurring at positions 11, 12, 13 of the strand, the count starting from the first nucleotide from the 5'-end, or optionally, the count starting at the first paired nucleotide within the duplex region, from the 5'-end; and Y' represents 2'-O-methyl modification. The antisense strand may additionally contain X'X'X' motif or Z'Z'Z' motifs as wing modifications at the opposite end of the duplex region; and X'X'X' and Z'Z'Z' each independently represents a 2'-OMe modification or 2'-F modification.

The sense strand represented by any one of the above formulas (Ia), (Ib), (Ic), and (Id) forms a duplex with an antisense strand being represented by any one of formulas (IIa), (IIb), (IIc), and (IId), respectively.

Accordingly, the dsRNAi agents for use in the methods of the invention may comprise a sense strand and an antisense strand, each strand having 14 to 30 nucleotides, the iRNA duplex represented by formula (III):



wherein:

i, j, k, and l are each independently 0 or 1;

p, p', q, and q' are each independently 0-6;

each N_a and N_a' independently represents an oligonucleotide sequence comprising 0-25 modified nucleotides, each sequence comprising at least two differently modified nucleotides;


each N_b and N_b' independently represents an oligonucleotide sequence comprising 0-10 modified nucleotides;


wherein each n_p , n_p , n_q , and n_q , each of which may or may not be present, independently represents an overhang nucleotide; and

XXX, YYY, ZZZ, X'X'X', Y'Y'Y', and Z'Z'Z' each independently represent one motif of three identical modifications on three consecutive nucleotides.



In one embodiment, i is 0 and j is 0; or i is 1 and j is 0; or i is 0 and j is 1; or both i and j are 0; or both i and j are 1. In another embodiment, k is 0 and l is 0; or k is 1 and l is 0; or k is 0 and l is 1; or both k and l are 0; or both k and l are 1.

Exemplary combinations of the sense strand and antisense strand forming an iRNA duplex include the formulas below:





(IIIb)



(IIIc)



(IIId)

When the dsRNAi agent is represented by formula (IIIa), each N_a independently represents an oligonucleotide sequence comprising 2-20, 2-15, or 2-10 modified nucleotides.

When the dsRNAi agent is represented by formula (IIIb), each N_b independently represents an oligonucleotide sequence comprising 1-10, 1-7, 1-5, or 1-4 modified nucleotides. Each N_a independently represents an oligonucleotide sequence comprising 2-20, 2-15, or 2-10 modified nucleotides.

When the dsRNAi agent is represented as formula (IIIc), each N_b, N_b' independently represents an oligonucleotide sequence comprising 0-10, 0-7, 0-10, 0-7, 0-5, 0-4, 0-2, or 0 modified nucleotides. Each N_a independently represents an oligonucleotide sequence comprising 2-20, 2-15, or 2-10 modified nucleotides.

When the dsRNAi agent is represented as formula (IIId), each N_b, N_b' independently represents an oligonucleotide sequence comprising 0-10, 0-7, 0-10, 0-7, 0-5, 0-4, 0-2, or 0 modified nucleotides. Each N_a, N_a' independently represents an oligonucleotide sequence comprising 2-20, 2-15, or 2-10 modified nucleotides. Each of N_a, N_a', N_b, and N_b' independently comprises modifications of alternating pattern.

Each of X, Y, and Z in formulas (III), (IIIa), (IIIb), (IIIc), and (IIId) may be the same or different from each other.

When the dsRNAi agent is represented by formula (III), (IIIa), (IIIb), (IIIc), and (IIId), at least one of the Y nucleotides may form a base pair with one of the Y' nucleotides. Alternatively, at least two of the Y nucleotides form base pairs with the corresponding Y' nucleotides; or all three of the Y nucleotides all form base pairs with the corresponding Y' nucleotides.

When the dsRNAi agent is represented by formula (IIIb) or (IIId), at least one of the Z nucleotides may form a base pair with one of the Z' nucleotides. Alternatively, at least two of the Z nucleotides form base pairs with the corresponding Z' nucleotides; or all three of the Z nucleotides all form base pairs with the corresponding Z' nucleotides.

When the dsRNAi agent is represented as formula (IIIc) or (IIId), at least one of the X nucleotides may form a base pair with one of the X' nucleotides. Alternatively, at least two of the X nucleotides form base pairs with the corresponding X' nucleotides; or all three of the X nucleotides all form base pairs with the corresponding X' nucleotides.

In certain embodiments, the modification on the Y nucleotide is different than the modification on the Y' nucleotide, the modification on the Z nucleotide is different than the modification on the Z' nucleotide, or the modification on the X nucleotide is different than the modification on the X' nucleotide.

5 In certain embodiments, when the dsRNAi agent is represented by formula (IIId), the N_a modifications are 2'-O-methyl or 2'-fluoro modifications. In other embodiments, when the RNAi agent is represented by formula (IIId), the N_a modifications are 2'-O-methyl or 2'-fluoro modifications and n_p' > 0 and at least one n_p' is linked to a neighboring nucleotide *via* phosphorothioate linkage. In yet other embodiments, when the RNAi agent is represented by formula (IIId), the N_a modifications are 2'-O-methyl or 2'-fluoro modifications, n_p' > 0 and at least one n_p' is linked to a neighboring nucleotide *via* phosphorothioate linkage, and the sense strand is conjugated to one or more GalNAc derivatives attached through a bivalent or trivalent branched linker (described below). In other
10 embodiments, when the RNAi agent is represented by formula (IIId), the N_a modifications are 2'-O-methyl or 2'-fluoro modifications, n_p' > 0 and at least one n_p' is linked to a neighboring nucleotide *via* phosphorothioate linkage, the sense strand comprises at least one phosphorothioate linkage, and the sense strand is conjugated to one or more GalNAc derivatives attached through a bivalent or trivalent
15 branched linker.

In some embodiments, when the dsRNAi agent is represented by formula (IIIa), the N_a modifications are 2'-O-methyl or 2'-fluoro modifications, n_p' > 0 and at least one n_p' is linked to a neighboring nucleotide *via* phosphorothioate linkage, the sense strand comprises at least one phosphorothioate linkage, and the sense strand is conjugated to one or more GalNAc derivatives attached through a bivalent or trivalent branched linker.
20

In some embodiments, the dsRNAi agent is a multimer containing at least two duplexes represented by formula (III), (IIIa), (IIIb), (IIIc), and (IIId), wherein the duplexes are connected by a linker. The linker can be cleavable or non-cleavable. Optionally, the multimer further comprises a ligand. Each of the duplexes can target the same gene or two different genes; or each of the duplexes can target same gene at two different target sites.
25

In some embodiments, the dsRNAi agent is a multimer containing three, four, five, six, or more duplexes represented by formula (III), (IIIa), (IIIb), (IIIc), and (IIId), wherein the duplexes are connected by a linker. The linker can be cleavable or non-cleavable. Optionally, the multimer further comprises a ligand. Each of the duplexes can target the same gene or two different genes; or each of the duplexes can target same gene at two different target sites.
30


In one embodiment, two dsRNAi agents represented by at least one of formulas (III), (IIIa), (IIIb), (IIIc), and (IIId) are linked to each other at the 5' end, and one or both of the 3' ends, and are optionally conjugated to a ligand. Each of the agents can target the same gene or two different genes; or each of the agents can target same gene at two different target sites.
35

In certain embodiments, an RNAi agent of the invention may contain a low number of nucleotides containing a 2'-fluoro modification, *e.g.*, 10 or fewer nucleotides with 2'-fluoro modification. For example, the RNAi agent may contain 10, 9, 8, 7, 6, 5, 4, 3, 2, 1 or 0 nucleotides with a 2'-fluoro modification. In a specific embodiment, the RNAi agent of the invention contains 10 nucleotides with a 2'-fluoro modification, *e.g.*, 4 nucleotides with a 2'-fluoro modification in the sense strand and 6 nucleotides with a 2'-fluoro modification in the antisense strand. In another specific embodiment, the RNAi agent of the invention contains 6 nucleotides with a 2'-fluoro modification, *e.g.*, 4 nucleotides with a 2'-fluoro modification in the sense strand and 2 nucleotides with a 2'-fluoro modification in the antisense strand.

In other embodiments, an RNAi agent of the invention may contain an ultra low number of nucleotides containing a 2'-fluoro modification, *e.g.*, 2 or fewer nucleotides containing a 2'-fluoro modification. For example, the RNAi agent may contain 2, 1 or 0 nucleotides with a 2'-fluoro modification. In a specific embodiment, the RNAi agent may contain 2 nucleotides with a 2'-fluoro modification, *e.g.*, 0 nucleotides with a 2'-fluoro modification in the sense strand and 2 nucleotides with a 2'-fluoro modification in the antisense strand.

Various publications describe multimeric iRNAs that can be used in the methods of the invention. Such publications include WO2007/091269, U.S. Patent No. 7,858,769, WO2010/141511, WO2007/117686, WO2009/014887, and WO2011/031520 the entire contents of each of which are hereby incorporated herein by reference.

In certain embodiments, the compositions and methods of the disclosure include a vinyl phosphonate (VP) modification of an RNAi agent as described herein. In exemplary embodiments, a 5'-vinyl phosphonate modified nucleotide of the disclosure has the structure:



wherein X is O or S;

R is hydrogen, hydroxy, fluoro, or C₁₋₂₀alkoxy (*e.g.*, methoxy or n-hexadecyloxy);

R^{5'} is =C(H)-P(O)(OH)₂ and the double bond between the C5' carbon and R^{5'} is in the E or Z orientation (*e.g.*, E orientation); and

B is a nucleobase or a modified nucleobase, optionally where B is adenine, guanine, cytosine, thymine, or uracil.

In one embodiment, R^{5'} is =C(H)-P(O)(OH)₂ and the double bond between the C5' carbon and R^{5'} is in the E orientation. In another embodiment, R is methoxy and R^{5'} is =C(H)-P(O)(OH)₂ and the double bond between the C5' carbon and R^{5'} is in the E orientation. In another embodiment,

X is S, R is methoxy, and R^{5'} is =C(H)-P(O)(OH)₂ and the double bond between the C5' carbon and R5' is in the E orientation.

A vinyl phosphonate of the instant disclosure may be attached to either the antisense or the sense strand of a dsRNA of the disclosure. In certain embodiments, a vinyl phosphonate of the instant disclosure is attached to the antisense strand of a dsRNA, optionally at the 5' end of the antisense strand of the dsRNA.

Vinyl phosphonate modifications are also contemplated for the compositions and methods of the instant disclosure. An exemplary vinyl phosphonate structure includes the preceding structure, where R5' is =C(H)-OP(O)(OH)₂ and the double bond between the C5' carbon and R5' is in the E or Z orientation (*e.g.*, E orientation).

As described in more detail below, the iRNA that contains conjugations of one or more carbohydrate moieties to an iRNA can optimize one or more properties of the iRNA. In many cases, the carbohydrate moiety will be attached to a modified subunit of the iRNA. For example, the ribose sugar of one or more ribonucleotide subunits of a iRNA can be replaced with another moiety, *e.g.*, a non-carbohydrate (such as, cyclic) carrier to which is attached a carbohydrate ligand. A ribonucleotide subunit in which the ribose sugar of the subunit has been so replaced is referred to herein as a ribose replacement modification subunit (RRMS). A cyclic carrier may be a carbocyclic ring system, *i.e.*, all ring atoms are carbon atoms, or a heterocyclic ring system, *i.e.*, one or more ring atoms may be a heteroatom, *e.g.*, nitrogen, oxygen, sulfur. The cyclic carrier may be a monocyclic ring system, or may contain two or more rings, *e.g.* fused rings. The cyclic carrier may be a fully saturated ring system, or it may contain one or more double bonds.

The ligand may be attached to the polynucleotide *via* a carrier. The carriers include (i) at least one "backbone attachment point," such as, two "backbone attachment points" and (ii) at least one "tethering attachment point." A "backbone attachment point" as used herein refers to a functional group, *e.g.* a hydroxyl group, or generally, a bond available for, and that is suitable for incorporation of the carrier into the backbone, *e.g.*, the phosphate, or modified phosphate, *e.g.*, sulfur containing, backbone, of a ribonucleic acid. A "tethering attachment point" (TAP) in some embodiments refers to a constituent ring atom of the cyclic carrier, *e.g.*, a carbon atom or a heteroatom (distinct from an atom which provides a backbone attachment point), that connects a selected moiety. The moiety can be, *e.g.*, a carbohydrate, *e.g.* monosaccharide, disaccharide, trisaccharide, tetrasaccharide, oligosaccharide, or polysaccharide. Optionally, the selected moiety is connected by an intervening tether to the cyclic carrier. Thus, the cyclic carrier will often include a functional group, *e.g.*, an amino group, or generally, provide a bond, that is suitable for incorporation or tethering of another chemical entity, *e.g.*, a ligand to the constituent ring.

The iRNA may be conjugated to a ligand *via* a carrier, wherein the carrier can be cyclic group or acyclic group. In one embodiment, the cyclic group is selected from pyrrolidinyl, pyrazolinyl,

pyrazolidinyl, imidazoliny, imidazolidinyl, piperidinyl, piperazinyl, [1,3]dioxolane, oxazolidinyl, isoxazolidinyl, morpholinyl, thiazolidinyl, isothiazolidinyl, quinoxaliny, pyridazinonyl, tetrahydrofuryl, and decalin. In one embodiment, the acyclic group is a serinol backbone or diethanolamine backbone.

5


i. Thermally Destabilizing Modifications

In certain embodiments, a dsRNA molecule can be optimized for RNA interference by incorporating thermally destabilizing modifications in the seed region of the antisense strand (*i.e.*, at positions 2-9 of the 5'-end of the antisense strand or at positions 2-8 of the 5'-end of the antisense strand) to reduce or inhibit off-target gene silencing.

The term "thermally destabilizing modification(s)" includes modification(s) that would result with a dsRNA with a lower overall melting temperature (T_m) than the T_m of the dsRNA without having such modification(s). For example, the thermally destabilizing modification(s) can decrease the T_m of the dsRNA by 1 – 4 °C, such as one, two, three or four degrees Celcius. And, the term "thermally destabilizing nucleotide" refers to a nucleotide containing one or more thermally destabilizing modifications.


It has been discovered that dsRNAs with an antisense strand comprising at least one thermally destabilizing modification of the duplex within the first 9 nucleotide positions, counting from the 5' end, of the antisense strand have reduced off-target gene silencing activity. Accordingly, in some embodiments, the antisense strand comprises at least one (*e.g.*, one, two, three, four, five or more) thermally destabilizing modification of the duplex within the first 9 nucleotide positions of the 5' region of the antisense strand. In some embodiments, one or more thermally destabilizing modification(s) of the duplex is/are located in positions 2-9, such as, positions 4-8, from the 5'-end of the antisense strand. In some further embodiments, the thermally destabilizing modification(s) of the duplex is/are located at position 6, 7 or 8 from the 5'-end of the antisense strand. In still some further embodiments, the thermally destabilizing modification of the duplex is located at position 7 from the 5'-end of the antisense strand. In some embodiments, the thermally destabilizing modification of the duplex is located at position 2, 3, 4, 5 or 9 from the 5'-end of the antisense strand.

An iRNA agent comprises a sense strand and an antisense strand, each strand having 14 to 40 nucleotides. The RNAi agent may be represented by formula (L):





In formula (L), B1, B2, B3, B1', B2', B3', and B4' each are independently a nucleotide containing a modification selected from the group consisting of 2'-O-alkyl, 2'-substituted alkoxy, 2'-substituted alkyl, 2'-halo, ENA, and BNA/LNA. In one embodiment, B1, B2, B3, B1', B2', B3', and B4' each contain 2'-OMe modifications. In one embodiment, B1, B2, B3, B1', B2', B3', and B4' each contain 2'-OMe or 2'-F modifications. In one embodiment, at least one of B1, B2, B3, B1', B2', B3', and B4' contain 2'-O-N-methylacetamido (2'-O-NMA, 2'-O-CH₂C(O)N(Me)H) modification.

C1 is a thermally destabilizing nucleotide placed at a site opposite to the seed region of the antisense strand (*i.e.*, at positions 2-8 of the 5'-end of the antisense strand or at positions 2-9 of the 5'-end of the antisense strand). For example, C1 is at a position of the sense strand that pairs with a nucleotide at positions 2-8 of the 5'-end of the antisense strand. In one example, C1 is at position 15 from the 5'-end of the sense strand. C1 nucleotide bears the thermally destabilizing modification which can include abasic modification; mismatch with the opposing nucleotide in the duplex; and sugar modification such as 2'-deoxy modification or acyclic nucleotide *e.g.*, unlocked nucleic acids (UNA) or glycerol nucleic acid (GNA). In one embodiment, C1 has thermally destabilizing modification selected from the group consisting of: i) mismatch with the opposing nucleotide in the antisense strand; ii) abasic modification selected from the group consisting of:




selected from the group consisting of:



20 , wherein B is a modified or unmodified nucleobase, R¹ and R² independently are H, halogen, OR₃, or alkyl; and R₃ is H, alkyl, cycloalkyl, aryl, aralkyl, heteroaryl or sugar. In one embodiment, the thermally destabilizing modification in C1 is a mismatch selected from the group consisting of G:G, G:A, G:U, G:T, A:A, A:C, C:C, C:U, C:T, U:U, T:T, and U:T; and optionally, at

least one nucleobase in the mismatch pair is a 2'-deoxy nucleobase. In one example, the thermally



destabilizing modification in C1 is GNA or

T1, T1', T2', and T3' each independently represent a nucleotide comprising a modification providing the nucleotide a steric bulk that is less or equal to the steric bulk of a 2'-OMe modification. A steric bulk refers to the sum of steric effects of a modification. Methods for determining steric effects of a modification of a nucleotide are known to one skilled in the art. The modification can be at the 2' position of a ribose sugar of the nucleotide, or a modification to a non-ribose nucleotide, acyclic nucleotide, or the backbone of the nucleotide that is similar or equivalent to the 2' position of the ribose sugar, and provides the nucleotide a steric bulk that is less than or equal to the steric bulk of a 2'-OMe modification. For example, T1, T1', T2', and T3' are each independently selected from DNA, RNA, LNA, 2'-F, and 2'-F-5'-methyl. In one embodiment, T1 is DNA. In one embodiment, T1' is DNA, RNA or LNA. In one embodiment, T2' is DNA or RNA. In one embodiment, T3' is DNA or RNA.

n^1 , n^3 , and q^1 are independently 4 to 15 nucleotides in length.

n^5 , q^3 , and q^7 are independently 1-6 nucleotide(s) in length.

n^4 , q^2 , and q^6 are independently 1-3 nucleotide(s) in length; alternatively, n^4 is 0.

q^5 is independently 0-10 nucleotide(s) in length.

n^2 and q^4 are independently 0-3 nucleotide(s) in length.

Alternatively, n^4 is 0-3 nucleotide(s) in length.

In one embodiment, n^4 can be 0. In one example, n^4 is 0, and q^2 and q^6 are 1. In another example, n^4 is 0, and q^2 and q^6 are 1, with two phosphorothioate internucleotide linkage modifications within position 1-5 of the sense strand (counting from the 5'-end of the sense strand), and two phosphorothioate internucleotide linkage modifications at positions 1 and 2 and two phosphorothioate internucleotide linkage modifications within positions 18-23 of the antisense strand (counting from the 5'-end of the antisense strand).

In one embodiment, n^4 , q^2 , and q^6 are each 1.

In one embodiment, n^2 , n^4 , q^2 , q^4 , and q^6 are each 1.

In one embodiment, C1 is at position 14-17 of the 5'-end of the sense strand, when the sense strand is 19-22 nucleotides in length, and n^4 is 1. In one embodiment, C1 is at position 15 of the 5'-end of the sense strand

In one embodiment, T3' starts at position 2 from the 5' end of the antisense strand. In one example, T3' is at position 2 from the 5' end of the antisense strand and q^6 is equal to 1.

In one embodiment, T1' starts at position 14 from the 5' end of the antisense strand. In one example, T1' is at position 14 from the 5' end of the antisense strand and q^2 is equal to 1.

In an exemplary embodiment, T3' starts from position 2 from the 5' end of the antisense strand and T1' starts from position 14 from the 5' end of the antisense strand. In one example, T3' starts from position 2 from the 5' end of the antisense strand and q^6 is equal to 1 and T1' starts from position 14 from the 5' end of the antisense strand and q^2 is equal to 1.

5 In one embodiment, T1' and T3' are separated by 11 nucleotides in length (*i.e.* not counting the T1' and T3' nucleotides).

In one embodiment, T1' is at position 14 from the 5' end of the antisense strand. In one example, T1' is at position 14 from the 5' end of the antisense strand and q^2 is equal to 1, and the modification at the 2' position or positions in a non-ribose, acyclic or backbone that provide less steric bulk than a 2'-OMe ribose.

10 In one embodiment, T3' is at position 2 from the 5' end of the antisense strand. In one example, T3' is at position 2 from the 5' end of the antisense strand and q^6 is equal to 1, and the modification at the 2' position or positions in a non-ribose, acyclic or backbone that provide less than or equal to steric bulk than a 2'-OMe ribose.

15 In one embodiment, T1 is at the cleavage site of the sense strand. In one example, T1 is at position 11 from the 5' end of the sense strand, when the sense strand is 19-22 nucleotides in length, and n^2 is 1. In an exemplary embodiment, T1 is at the cleavage site of the sense strand at position 11 from the 5' end of the sense strand, when the sense strand is 19-22 nucleotides in length, and n^2 is 1,

In one embodiment, T2' starts at position 6 from the 5' end of the antisense strand. In one example, T2' is at positions 6-10 from the 5' end of the antisense strand, and q^4 is 1.

20 In an exemplary embodiment, T1 is at the cleavage site of the sense strand, for instance, at position 11 from the 5' end of the sense strand, when the sense strand is 19-22 nucleotides in length, and n^2 is 1; T1' is at position 14 from the 5' end of the antisense strand, and q^2 is equal to 1, and the modification to T1' is at the 2' position of a ribose sugar or at positions in a non-ribose, acyclic or backbone that provide less steric bulk than a 2'-OMe ribose; T2' is at positions 6-10 from the 5' end of the antisense strand, and q^4 is 1; and T3' is at position 2 from the 5' end of the antisense strand, and q^6 is equal to 1, and the modification to T3' is at the 2' position or at positions in a non-ribose, acyclic or backbone that provide less than or equal to steric bulk than a 2'-OMe ribose.

In one embodiment, T2' starts at position 8 from the 5' end of the antisense strand. In one example, T2' starts at position 8 from the 5' end of the antisense strand, and q^4 is 2.

30 In one embodiment, T2' starts at position 9 from the 5' end of the antisense strand. In one example, T2' is at position 9 from the 5' end of the antisense strand, and q^4 is 1.

In one embodiment, B1' is 2'-OMe or 2'-F, q^1 is 9, T1' is 2'-F, q^2 is 1, B2' is 2'-OMe or 2'-F, q^3 is 4, T2' is 2'-F, q^4 is 1, B3' is 2'-OMe or 2'-F, q^5 is 6, T3' is 2'-F, q^6 is 1, B4' is 2'-OMe, and q^7 is 1; with two phosphorothioate internucleotide linkage modifications within positions 1-5 of the sense strand (counting from the 5'-end of the sense strand), and two phosphorothioate internucleotide linkage modifications at positions 1 and 2 and two phosphorothioate internucleotide linkage

modifications within positions 18-23 of the antisense strand (counting from the 5'-end of the antisense strand).

In one embodiment, n^4 is 0, B3 is 2'-OMe, n^5 is 3, B1' is 2'-OMe or 2'-F, q^1 is 9, T1' is 2'-F, q^2 is 1, B2' is 2'-OMe or 2'-F, q^3 is 4, T2' is 2'-F, q^4 is 1, B3' is 2'-OMe or 2'-F, q^5 is 6, T3' is 2'-F, q^6 is 1, B4' is 2'-OMe, and q^7 is 1; with two phosphorothioate internucleotide linkage modifications within positions 1-5 of the sense strand (counting from the 5'-end of the sense strand), and two phosphorothioate internucleotide linkage modifications at positions 1 and 2 and two phosphorothioate internucleotide linkage modifications within positions 18-23 of the antisense strand (counting from the 5'-end of the antisense strand).

In one embodiment, B1 is 2'-OMe or 2'-F, n^1 is 8, T1 is 2'-F, n^2 is 3, B2 is 2'-OMe, n^3 is 7, n^4 is 0, B3 is 2'-OMe, n^5 is 3, B1' is 2'-OMe or 2'-F, q^1 is 9, T1' is 2'-F, q^2 is 1, B2' is 2'-OMe or 2'-F, q^3 is 4, T2' is 2'-F, q^4 is 2, B3' is 2'-OMe or 2'-F, q^5 is 5, T3' is 2'-F, q^6 is 1, B4' is 2'-OMe, and q^7 is 1.

In one embodiment, B1 is 2'-OMe or 2'-F, n^1 is 8, T1 is 2'-F, n^2 is 3, B2 is 2'-OMe, n^3 is 7, n^4 is 0, B3 is 2'-OMe, n^5 is 3, B1' is 2'-OMe or 2'-F, q^1 is 9, T1' is 2'-F, q^2 is 1, B2' is 2'-OMe or 2'-F, q^3 is 4, T2' is 2'-F, q^4 is 2, B3' is 2'-OMe or 2'-F, q^5 is 5, T3' is 2'-F, q^6 is 1, B4' is 2'-OMe, and q^7 is 1; with two phosphorothioate internucleotide linkage modifications within positions 1-5 of the sense strand (counting from the 5'-end of the sense strand), and two phosphorothioate internucleotide linkage modifications at positions 1 and 2 and two phosphorothioate internucleotide linkage modifications within positions 18-23 of the antisense strand (counting from the 5'-end of the antisense strand).

In one embodiment, B1 is 2'-OMe or 2'-F, n^1 is 6, T1 is 2'-F, n^2 is 3, B2 is 2'-OMe, n^3 is 7, n^4 is 0, B3 is 2'-OMe, n^5 is 3, B1' is 2'-OMe or 2'-F, q^1 is 7, T1' is 2'-F, q^2 is 1, B2' is 2'-OMe or 2'-F, q^3 is 4, T2' is 2'-F, q^4 is 2, B3' is 2'-OMe or 2'-F, q^5 is 5, T3' is 2'-F, q^6 is 1, B4' is 2'-OMe, and q^7 is 1.

In one embodiment, B1 is 2'-OMe or 2'-F, n^1 is 6, T1 is 2'-F, n^2 is 3, B2 is 2'-OMe, n^3 is 7, n^4 is 0, B3 is 2'-OMe, n^5 is 3, B1' is 2'-OMe or 2'-F, q^1 is 7, T1' is 2'-F, q^2 is 1, B2' is 2'-OMe or 2'-F, q^3 is 4, T2' is 2'-F, q^4 is 2, B3' is 2'-OMe or 2'-F, q^5 is 5, T3' is 2'-F, q^6 is 1, B4' is 2'-OMe, and q^7 is 1; with two phosphorothioate internucleotide linkage modifications within positions 1-5 of the sense strand (counting from the 5'-end of the sense strand), and two phosphorothioate internucleotide linkage modifications at positions 1 and 2 and two phosphorothioate internucleotide linkage modifications within positions 18-23 of the antisense strand (counting from the 5'-end of the antisense strand).

In one embodiment, B1 is 2'-OMe or 2'-F, n^1 is 8, T1 is 2'-F, n^2 is 3, B2 is 2'-OMe, n^3 is 7, n^4 is 0, B3 is 2'-OMe, n^5 is 3, B1' is 2'-OMe or 2'-F, q^1 is 9, T1' is 2'-F, q^2 is 1, B2' is 2'-OMe or 2'-F, q^3 is 4, T2' is 2'-F, q^4 is 1, B3' is 2'-OMe or 2'-F, q^5 is 6, T3' is 2'-F, q^6 is 1, B4' is 2'-OMe, and q^7 is 1.

In one embodiment, B1 is 2'-OMe or 2'-F, n^1 is 8, T1 is 2'-F, n^2 is 3, B2 is 2'-OMe, n^3 is 7, n^4 is 0, B3 is 2'-OMe, n^5 is 3, B1' is 2'-OMe or 2'-F, q^1 is 9, T1' is 2'-F, q^2 is 1, B2' is 2'-OMe or 2'-F,

q³ is 4, T2' is 2'-F, q⁴ is 1, B3' is 2'-OMe or 2'-F, q⁵ is 6, T3' is 2'-F, q⁶ is 1, B4' is 2'-OMe, and q⁷ is 1; with two phosphorothioate internucleotide linkage modifications within positions 1-5 of the sense strand (counting from the 5'-end of the sense strand), and two phosphorothioate internucleotide linkage modifications at positions 1 and 2 and two phosphorothioate internucleotide linkage modifications within positions 18-23 of the antisense strand (counting from the 5'-end of the antisense strand).

In one embodiment, B1 is 2'-OMe or 2'-F, n¹ is 8, T1 is 2'-F, n² is 3, B2 is 2'-OMe, n³ is 7, n⁴ is 0, B3 is 2'-OMe, n⁵ is 3, B1' is 2'-OMe or 2'-F, q¹ is 9, T1' is 2'-F, q² is 1, B2' is 2'-OMe or 2'-F, q³ is 5, T2' is 2'-F, q⁴ is 1, B3' is 2'-OMe or 2'-F, q⁵ is 5, T3' is 2'-F, q⁶ is 1, B4' is 2'-OMe, and q⁷ is 1; optionally with at least 2 additional TT at the 3'-end of the antisense strand.

In one embodiment, B1 is 2'-OMe or 2'-F, n¹ is 8, T1 is 2'-F, n² is 3, B2 is 2'-OMe, n³ is 7, n⁴ is 0, B3 is 2'-OMe, n⁵ is 3, B1' is 2'-OMe or 2'-F, q¹ is 9, T1' is 2'-F, q² is 1, B2' is 2'-OMe or 2'-F, q³ is 5, T2' is 2'-F, q⁴ is 1, B3' is 2'-OMe or 2'-F, q⁵ is 5, T3' is 2'-F, q⁶ is 1, B4' is 2'-OMe, and q⁷ is 1; optionally with at least 2 additional TT at the 3'-end of the antisense strand; with two phosphorothioate internucleotide linkage modifications within positions 1-5 of the sense strand (counting from the 5'-end of the sense strand), and two phosphorothioate internucleotide linkage modifications at positions 1 and 2 and two phosphorothioate internucleotide linkage modifications within positions 18-23 of the antisense strand (counting from the 5'-end of the antisense strand).

In one embodiment, B1 is 2'-OMe or 2'-F, n¹ is 8, T1 is 2'-F, n² is 3, B2 is 2'-OMe, n³ is 7, n⁴ is 0, B3 is 2'-OMe, n⁵ is 3, B1' is 2'-OMe or 2'-F, q¹ is 9, T1' is 2'-F, q² is 1, B2' is 2'-OMe or 2'-F, q³ is 4, q⁴ is 0, B3' is 2'-OMe or 2'-F, q⁵ is 7, T3' is 2'-F, q⁶ is 1, B4' is 2'-OMe, and q⁷ is 1.

In one embodiment, B1 is 2'-OMe or 2'-F, n¹ is 8, T1 is 2'-F, n² is 3, B2 is 2'-OMe, n³ is 7, n⁴ is 0, B3 is 2'-OMe, n⁵ is 3, B1' is 2'-OMe or 2'-F, q¹ is 9, T1' is 2'-F, q² is 1, B2' is 2'-OMe or 2'-F, q³ is 4, q⁴ is 0, B3' is 2'-OMe or 2'-F, q⁵ is 7, T3' is 2'-F, q⁶ is 1, B4' is 2'-OMe, and q⁷ is 1; with two phosphorothioate internucleotide linkage modifications within positions 1-5 of the sense strand (counting from the 5'-end), and two phosphorothioate internucleotide linkage modifications at positions 1 and 2 and two phosphorothioate internucleotide linkage modifications within positions 18-23 of the antisense strand (counting from the 5'-end).


In one embodiment, B1 is 2'-OMe or 2'-F, n¹ is 8, T1 is 2'-F, n² is 3, B2 is 2'-OMe, n³ is 7, n⁴ is 0, B3 is 2'-OMe, n⁵ is 3, B1' is 2'-OMe or 2'-F, q¹ is 9, T1' is 2'-F, q² is 1, B2' is 2'-OMe or 2'-F, q³ is 4, T2' is 2'-F, q⁴ is 2, B3' is 2'-OMe or 2'-F, q⁵ is 5, T3' is 2'-F, q⁶ is 1, B4' is 2'-F, and q⁷ is 1. In one embodiment, B1 is 2'-OMe or 2'-F, n¹ is 8, T1 is 2'-F, n² is 3, B2 is 2'-OMe, n³ is 7, n⁴ is 0, B3 is 2'-OMe, n⁵ is 3, B1' is 2'-OMe or 2'-F, q¹ is 9, T1' is 2'-F, q² is 1, B2' is 2'-OMe or 2'-F, q³ is 4, T2' is 2'-F, q⁴ is 2, B3' is 2'-OMe or 2'-F, q⁵ is 5, T3' is 2'-F, q⁶ is 1, B4' is 2'-F, and q⁷ is 1; with two phosphorothioate internucleotide linkage modifications within positions 1-5 of the sense strand (counting from the 5'-end of the sense strand), and two phosphorothioate internucleotide linkage

modifications at positions 1 and 2 and two phosphorothioate internucleotide linkage modifications within positions 18-23 of the antisense strand (counting from the 5'-end of the antisense strand).


In one embodiment, B1 is 2'-OMe or 2'-F, n^1 is 8, T1 is 2'-F, n^2 is 3, B2 is 2'-OMe, n^3 is 7, n^4 is 0, B3 is 2'-OMe, n^5 is 3, B1' is 2'-OMe or 2'-F, q^1 is 9, T1' is 2'-F, q^2 is 1, B2' is 2'-OMe or 2'-F, q^3 is 4, q^4 is 0, B3' is 2'-OMe or 2'-F, q^5 is 7, T3' is 2'-F, q^6 is 1, B4' is 2'-F, and q^7 is 1.

In one embodiment, B1 is 2'-OMe or 2'-F, n^1 is 8, T1 is 2'-F, n^2 is 3, B2 is 2'-OMe, n^3 is 7, n^4 is 0, B3 is 2'-OMe, n^5 is 3, B1' is 2'-OMe or 2'-F, q^1 is 9, T1' is 2'-F, q^2 is 1, B2' is 2'-OMe or 2'-F, q^3 is 4, q^4 is 0, B3' is 2'-OMe or 2'-F, q^5 is 7, T3' is 2'-F, q^6 is 1, B4' is 2'-F, and q^7 is 1; with two phosphorothioate internucleotide linkage modifications within positions 1-5 of the sense strand (counting from the 5'-end of the sense strand), and two phosphorothioate internucleotide linkage modifications at positions 1 and 2 and two phosphorothioate internucleotide linkage modifications within positions 18-23 of the antisense strand (counting from the 5'-end of the antisense strand).


The RNAi agent can comprise a phosphorus-containing group at the 5'-end of the sense strand or antisense strand. The 5'-end phosphorus-containing group can be 5'-end phosphate (5'-P), 5'-end phosphorothioate (5'-PS), 5'-end phosphorodithioate (5'-PS₂), 5'-end vinylphosphonate (5'-



VP), 5'-end methylphosphonate (MePhos), or 5'-deoxy-5'-C-malonyl (). When the 5'-end phosphorus-containing group is 5'-end vinylphosphonate (5'-VP), the 5'-VP can be either



5'-E-VP isomer (*i.e.*, *trans*-vinylphosphonate,), 5'-Z-VP isomer (*i.e.*, *cis*-



vinylphosphonate,), or mixtures thereof.

In one embodiment, the RNAi agent comprises a phosphorus-containing group at the 5'-end of the sense strand. In one embodiment, the RNAi agent comprises a phosphorus-containing group at the 5'-end of the antisense strand.

In one embodiment, the RNAi agent comprises a 5'-P. In one embodiment, the RNAi agent comprises a 5'-P in the antisense strand.

In one embodiment, the RNAi agent comprises a 5'-PS. In one embodiment, the RNAi agent comprises a 5'-PS in the antisense strand.

In one embodiment, the RNAi agent comprises a 5'-VP. In one embodiment, the RNAi agent comprises a 5'-VP in the antisense strand. In one embodiment, the RNAi agent comprises a 5'-E-VP

in the antisense strand. In one embodiment, the RNAi agent comprises a 5'-Z-VP in the antisense strand.

In one embodiment, the RNAi agent comprises a 5'-PS₂. In one embodiment, the RNAi agent comprises a 5'-PS₂ in the antisense strand.

5 In one embodiment, the RNAi agent comprises a 5'-PS₂. In one embodiment, the RNAi agent comprises a 5'-deoxy-5'-C-malonyl in the antisense strand.

In one embodiment, B1 is 2'-OMe or 2'-F, n¹ is 8, T1 is 2'-F, n² is 3, B2 is 2'-OMe, n³ is 7, n⁴ is 0, B3 is 2'-OMe, n⁵ is 3, B1' is 2'-OMe or 2'-F, q¹ is 9, T1' is 2'-F, q² is 1, B2' is 2'-OMe or 2'-F, q³ is 4, T2' is 2'-F, q⁴ is 2, B3' is 2'-OMe or 2'-F, q⁵ is 5, T3' is 2'-F, q⁶ is 1, B4' is 2'-OMe, and q⁷ is 1. The

10 RNAi agent also comprises a 5'-PS.

In one embodiment, B1 is 2'-OMe or 2'-F, n¹ is 8, T1 is 2'-F, n² is 3, B2 is 2'-OMe, n³ is 7, n⁴ is 0, B3 is 2'-OMe, n⁵ is 3, B1' is 2'-OMe or 2'-F, q¹ is 9, T1' is 2'-F, q² is 1, B2' is 2'-OMe or 2'-F, q³ is 4, T2' is 2'-F, q⁴ is 2, B3' is 2'-OMe or 2'-F, q⁵ is 5, T3' is 2'-F, q⁶ is 1, B4' is 2'-OMe, and q⁷ is 1. The RNAi agent also comprises a 5'-P.

15 In one embodiment, B1 is 2'-OMe or 2'-F, n¹ is 8, T1 is 2'-F, n² is 3, B2 is 2'-OMe, n³ is 7, n⁴ is 0, B3 is 2'-OMe, n⁵ is 3, B1' is 2'-OMe or 2'-F, q¹ is 9, T1' is 2'-F, q² is 1, B2' is 2'-OMe or 2'-F, q³ is 4, T2' is 2'-F, q⁴ is 2, B3' is 2'-OMe or 2'-F, q⁵ is 5, T3' is 2'-F, q⁶ is 1, B4' is 2'-OMe, and q⁷ is 1. The RNAi agent also comprises a 5'-VP. The 5'-VP may be 5'-E-VP, 5'-Z-VP, or combination thereof.

20 In one embodiment, B1 is 2'-OMe or 2'-F, n¹ is 8, T1 is 2'-F, n² is 3, B2 is 2'-OMe, n³ is 7, n⁴ is 0, B3 is 2'-OMe, n⁵ is 3, B1' is 2'-OMe or 2'-F, q¹ is 9, T1' is 2'-F, q² is 1, B2' is 2'-OMe or 2'-F, q³ is 4, T2' is 2'-F, q⁴ is 2, B3' is 2'-OMe or 2'-F, q⁵ is 5, T3' is 2'-F, q⁶ is 1, B4' is 2'-OMe, and q⁷ is 1. The RNAi agent also comprises a 5'-PS₂.

In one embodiment, B1 is 2'-OMe or 2'-F, n¹ is 8, T1 is 2'-F, n² is 3, B2 is 2'-OMe, n³ is 7, n⁴ is 0, B3 is 2'-OMe, n⁵ is 3, B1' is 2'-OMe or 2'-F, q¹ is 9, T1' is 2'-F, q² is 1, B2' is 2'-OMe or 2'-F, q³ is 4, T2' is 2'-F, q⁴ is 2, B3' is 2'-OMe or 2'-F, q⁵ is 5, T3' is 2'-F, q⁶ is 1, B4' is 2'-OMe, and q⁷ is 1. The RNAi agent also comprises a 5'-deoxy-5'-C-malonyl.

25 In one embodiment, B1 is 2'-OMe or 2'-F, n¹ is 8, T1 is 2'-F, n² is 3, B2 is 2'-OMe, n³ is 7, n⁴ is 0, B3 is 2'-OMe, n⁵ is 3, B1' is 2'-OMe or 2'-F, q¹ is 9, T1' is 2'-F, q² is 1, B2' is 2'-OMe or 2'-F, q³ is 4, T2' is 2'-F, q⁴ is 2, B3' is 2'-OMe or 2'-F, q⁵ is 5, T3' is 2'-F, q⁶ is 1, B4' is 2'-OMe, and q⁷ is 1; with two phosphorothioate internucleotide linkage modifications within position 1-5 of the sense strand (counting from the 5'-end of the sense strand), and two phosphorothioate internucleotide linkage modifications at positions 1 and 2 and two phosphorothioate internucleotide linkage modifications within positions 18-23 of the antisense strand (counting from the 5'-end of the antisense strand). The RNAi agent also comprises a 5'-P.

35 In one embodiment, B1 is 2'-OMe or 2'-F, n¹ is 8, T1 is 2'-F, n² is 3, B2 is 2'-OMe, n³ is 7, n⁴ is 0, B3 is 2'-OMe, n⁵ is 3, B1' is 2'-OMe or 2'-F, q¹ is 9, T1' is 2'-F, q² is 1, B2' is 2'-OMe or 2'-F,

q³ is 4, T2' is 2'-F, q⁴ is 2, B3' is 2'-OMe or 2'-F, q⁵ is 5, T3' is 2'-F, q⁶ is 1, B4' is 2'-OMe, and q⁷ is 1; with two phosphorothioate internucleotide linkage modifications within position 1-5 of the sense strand (counting from the 5'-end of the sense strand), and two phosphorothioate internucleotide linkage modifications at positions 1 and 2 and two phosphorothioate internucleotide linkage modifications within positions 18-23 of the antisense strand (counting from the 5'-end of the antisense strand). The RNAi agent also comprises a 5'-PS.

In one embodiment, B1 is 2'-OMe or 2'-F, n¹ is 8, T1 is 2'-F, n² is 3, B2 is 2'-OMe, n³ is 7, n⁴ is 0, B3 is 2'-OMe, n⁵ is 3, B1' is 2'-OMe or 2'-F, q¹ is 9, T1' is 2'-F, q² is 1, B2' is 2'-OMe or 2'-F, q³ is 4, T2' is 2'-F, q⁴ is 2, B3' is 2'-OMe or 2'-F, q⁵ is 5, T3' is 2'-F, q⁶ is 1, B4' is 2'-OMe, and q⁷ is 1; with two phosphorothioate internucleotide linkage modifications within position 1-5 of the sense strand (counting from the 5'-end of the sense strand), and two phosphorothioate internucleotide linkage modifications at positions 1 and 2 and two phosphorothioate internucleotide linkage modifications within positions 18-23 of the antisense strand (counting from the 5'-end of the antisense strand). The RNAi agent also comprises a 5'-VP. The 5'-VP may be 5'-E-VP, 5'-Z-VP, or combination thereof.

In one embodiment, B1 is 2'-OMe or 2'-F, n¹ is 8, T1 is 2'-F, n² is 3, B2 is 2'-OMe, n³ is 7, n⁴ is 0, B3 is 2'-OMe, n⁵ is 3, B1' is 2'-OMe or 2'-F, q¹ is 9, T1' is 2'-F, q² is 1, B2' is 2'-OMe or 2'-F, q³ is 4, T2' is 2'-F, q⁴ is 2, B3' is 2'-OMe or 2'-F, q⁵ is 5, T3' is 2'-F, q⁶ is 1, B4' is 2'-OMe, and q⁷ is 1; with two phosphorothioate internucleotide linkage modifications within position 1-5 of the sense strand (counting from the 5'-end of the sense strand), and two phosphorothioate internucleotide linkage modifications at positions 1 and 2 and two phosphorothioate internucleotide linkage modifications within positions 18-23 of the antisense strand (counting from the 5'-end of the antisense strand). The RNAi agent also comprises a 5'-PS₂.

In one embodiment, B1 is 2'-OMe or 2'-F, n¹ is 8, T1 is 2'-F, n² is 3, B2 is 2'-OMe, n³ is 7, n⁴ is 0, B3 is 2'-OMe, n⁵ is 3, B1' is 2'-OMe or 2'-F, q¹ is 9, T1' is 2'-F, q² is 1, B2' is 2'-OMe or 2'-F, q³ is 4, T2' is 2'-F, q⁴ is 2, B3' is 2'-OMe or 2'-F, q⁵ is 5, T3' is 2'-F, q⁶ is 1, B4' is 2'-OMe, and q⁷ is 1; with two phosphorothioate internucleotide linkage modifications within position 1-5 of the sense strand (counting from the 5'-end of the sense strand), and two phosphorothioate internucleotide linkage modifications at positions 1 and 2 and two phosphorothioate internucleotide linkage modifications within positions 18-23 of the antisense strand (counting from the 5'-end of the antisense strand). The RNAi agent also comprises a 5'-deoxy-5'-C-malonyl.

In one embodiment, B1 is 2'-OMe or 2'-F, n¹ is 8, T1 is 2'-F, n² is 3, B2 is 2'-OMe, n³ is 7, n⁴ is 0, B3 is 2'-OMe, n⁵ is 3, B1' is 2'-OMe or 2'-F, q¹ is 9, T1' is 2'-F, q² is 1, B2' is 2'-OMe or 2'-F, q³ is 4, q⁴ is 0, B3' is 2'-OMe or 2'-F, q⁵ is 7, T3' is 2'-F, q⁶ is 1, B4' is 2'-OMe, and q⁷ is 1. The RNAi agent also comprises a 5'-P.

In one embodiment, B1 is 2'-OMe or 2'-F, n¹ is 8, T1 is 2'-F, n² is 3, B2 is 2'-OMe, n³ is 7, n⁴ is 0, B3 is 2'-OMe, n⁵ is 3, B1' is 2'-OMe or 2'-F, q¹ is 9, T1' is 2'-F, q² is 1, B2' is 2'-OMe or 2'-F,

q^3 is 4, q^4 is 0, $B3'$ is 2'-OMe or 2'-F, q^5 is 7, $T3'$ is 2'-F, q^6 is 1, $B4'$ is 2'-OMe, and q^7 is 1. The dsRNA agent also comprises a 5'-PS.

In one embodiment, B1 is 2'-OMe or 2'-F, n^1 is 8, T1 is 2'F, n^2 is 3, B2 is 2'-OMe, n^3 is 7, n^4 is 0, B3 is 2'-OMe, n^5 is 3, B1' is 2'-OMe or 2'-F, q^1 is 9, T1' is 2'-F, q^2 is 1, B2' is 2'-OMe or 2'-F, q^3 is 4, q^4 is 0, B3' is 2'-OMe or 2'-F, q^5 is 7, T3' is 2'-F, q^6 is 1, B4' is 2'-OMe, and q^7 is 1. The RNAi agent also comprises a 5'-VP. The 5'-VP may be 5'-E-VP, 5'-Z-VP, or combination thereof. In one embodiment, B1 is 2'-OMe or 2'-F, n^1 is 8, T1 is 2'F, n^2 is 3, B2 is 2'-OMe, n^3 is 7, n^4 is 0, B3 is 2'-OMe, n^5 is 3, B1' is 2'-OMe or 2'-F, q^1 is 9, T1' is 2'-F, q^2 is 1, B2' is 2'-OMe or 2'-F, q^3 is 4, q^4 is 0, B3' is 2'-OMe or 2'-F, q^5 is 7, T3' is 2'-F, q^6 is 1, B4' is 2'-OMe, and q^7 is 1. The RNAi agent also comprises a 5'-PS₂.

In one embodiment, B1 is 2'-OMe or 2'-F, n^1 is 8, T1 is 2'F, n^2 is 3, B2 is 2'-OMe, n^3 is 7, n^4 is 0, B3 is 2'-OMe, n^5 is 3, B1' is 2'-OMe or 2'-F, q^1 is 9, T1' is 2'-F, q^2 is 1, B2' is 2'-OMe or 2'-F, q^3 is 4, q^4 is 0, B3' is 2'-OMe or 2'-F, q^5 is 7, T3' is 2'-F, q^6 is 1, B4' is 2'-OMe, and q^7 is 1. The RNAi agent also comprises a 5'-deoxy-5'-C-malonyl.

In one embodiment, B1 is 2'-OMe or 2'-F, n^1 is 8, T1 is 2'F, n^2 is 3, B2 is 2'-OMe, n^3 is 7, n^4 is 0, B3 is 2'-OMe, n^5 is 3, B1' is 2'-OMe or 2'-F, q^1 is 9, T1' is 2'-F, q^2 is 1, B2' is 2'-OMe or 2'-F, q^3 is 4, q^4 is 0, B3' is 2'-OMe or 2'-F, q^5 is 7, T3' is 2'-F, q^6 is 1, B4' is 2'-OMe, and q^7 is 1; with two phosphorothioate internucleotide linkage modifications within position 1-5 of the sense strand (counting from the 5'-end), and two phosphorothioate internucleotide linkage modifications at positions 1 and 2 and two phosphorothioate internucleotide linkage modifications within positions 18-23 of the antisense strand (counting from the 5'-end). The RNAi agent also comprises a 5'-P.

In one embodiment, B1 is 2'-OMe or 2'-F, n^1 is 8, T1 is 2'F, n^2 is 3, B2 is 2'-OMe, n^3 is 7, n^4 is 0, B3 is 2'-OMe, n^5 is 3, B1' is 2'-OMe or 2'-F, q^1 is 9, T1' is 2'-F, q^2 is 1, B2' is 2'-OMe or 2'-F, q^3 is 4, q^4 is 0, B3' is 2'-OMe or 2'-F, q^5 is 7, T3' is 2'-F, q^6 is 1, B4' is 2'-OMe, and q^7 is 1; with two phosphorothioate internucleotide linkage modifications within position 1-5 of the sense strand (counting from the 5'-end), and two phosphorothioate internucleotide linkage modifications at positions 1 and 2 and two phosphorothioate internucleotide linkage modifications within positions 18-23 of the antisense strand (counting from the 5'-end). The RNAi agent also comprises a 5'-PS.

In one embodiment, B1 is 2'-OMe or 2'-F, n^1 is 8, T1 is 2'F, n^2 is 3, B2 is 2'-OMe, n^3 is 7, n^4 is 0, B3 is 2'-OMe, n^5 is 3, B1' is 2'-OMe or 2'-F, q^1 is 9, T1' is 2'-F, q^2 is 1, B2' is 2'-OMe or 2'-F, q^3 is 4, q^4 is 0, B3' is 2'-OMe or 2'-F, q^5 is 7, T3' is 2'-F, q^6 is 1, B4' is 2'-OMe, and q^7 is 1; with two phosphorothioate internucleotide linkage modifications within position 1-5 of the sense strand (counting from the 5'-end), and two phosphorothioate internucleotide linkage modifications at positions 1 and 2 and two phosphorothioate internucleotide linkage modifications within positions 18-23 of the antisense strand (counting from the 5'-end). The RNAi agent also comprises a 5'-VP. The 5'-VP may be 5'-E-VP, 5'-Z-VP, or combination thereof.

In one embodiment, B1 is 2'-OMe or 2'-F, n¹ is 8, T1 is 2'-F, n² is 3, B2 is 2'-OMe, n³ is 7, n⁴ is 0, B3 is 2'-OMe, n⁵ is 3, B1' is 2'-OMe or 2'-F, q¹ is 9, T1' is 2'-F, q² is 1, B2' is 2'-OMe or 2'-F, q³ is 4, q⁴ is 0, B3' is 2'-OMe or 2'-F, q⁵ is 7, T3' is 2'-F, q⁶ is 1, B4' is 2'-OMe, and q⁷ is 1; with two phosphorothioate internucleotide linkage modifications within position 1-5 of the sense strand

- 5 (counting from the 5'-end), and two phosphorothioate internucleotide linkage modifications at positions 1 and 2 and two phosphorothioate internucleotide linkage modifications within positions 18-23 of the antisense strand (counting from the 5'-end). The RNAi agent also comprises a 5'-PS₂.

- In one embodiment, B1 is 2'-OMe or 2'-F, n¹ is 8, T1 is 2'-F, n² is 3, B2 is 2'-OMe, n³ is 7, n⁴ is 0, B3 is 2'-OMe, n⁵ is 3, B1' is 2'-OMe or 2'-F, q¹ is 9, T1' is 2'-F, q² is 1, B2' is 2'-OMe or 2'-F, q³ is 4, q⁴ is 0, B3' is 2'-OMe or 2'-F, q⁵ is 7, T3' is 2'-F, q⁶ is 1, B4' is 2'-OMe, and q⁷ is 1; with two phosphorothioate internucleotide linkage modifications within position 1-5 of the sense strand (counting from the 5'-end), and two phosphorothioate internucleotide linkage modifications at positions 1 and 2 and two phosphorothioate internucleotide linkage modifications within positions 18-23 of the antisense strand (counting from the 5'-end). The RNAi agent also comprises a 5'-deoxy-5'-C-malonyl.

- In one embodiment, B1 is 2'-OMe or 2'-F, n¹ is 8, T1 is 2'-F, n² is 3, B2 is 2'-OMe, n³ is 7, n⁴ is 0, B3 is 2'-OMe, n⁵ is 3, B1' is 2'-OMe or 2'-F, q¹ is 9, T1' is 2'-F, q² is 1, B2' is 2'-OMe or 2'-F, q³ is 4, T2' is 2'-F, q⁴ is 2, B3' is 2'-OMe or 2'-F, q⁵ is 5, T3' is 2'-F, q⁶ is 1, B4' is 2'-F, and q⁷ is 1. The RNAi agent also comprises a 5'-P.

- 20 In one embodiment, B1 is 2'-OMe or 2'-F, n¹ is 8, T1 is 2'-F, n² is 3, B2 is 2'-OMe, n³ is 7, n⁴ is 0, B3 is 2'-OMe, n⁵ is 3, B1' is 2'-OMe or 2'-F, q¹ is 9, T1' is 2'-F, q² is 1, B2' is 2'-OMe or 2'-F, q³ is 4, T2' is 2'-F, q⁴ is 2, B3' is 2'-OMe or 2'-F, q⁵ is 5, T3' is 2'-F, q⁶ is 1, B4' is 2'-F, and q⁷ is 1. The RNAi agent also comprises a 5'-PS.

- In one embodiment, B1 is 2'-OMe or 2'-F, n¹ is 8, T1 is 2'-F, n² is 3, B2 is 2'-OMe, n³ is 7, n⁴ is 0, B3 is 2'-OMe, n⁵ is 3, B1' is 2'-OMe or 2'-F, q¹ is 9, T1' is 2'-F, q² is 1, B2' is 2'-OMe or 2'-F, q³ is 4, T2' is 2'-F, q⁴ is 2, B3' is 2'-OMe or 2'-F, q⁵ is 5, T3' is 2'-F, q⁶ is 1, B4' is 2'-F, and q⁷ is 1. The RNAi agent also comprises a 5'-VP. The 5'-VP may be 5'-E-VP, 5'-Z-VP, or combination thereof.

- 25 In one embodiment, B1 is 2'-OMe or 2'-F, n¹ is 8, T1 is 2'-F, n² is 3, B2 is 2'-OMe, n³ is 7, n⁴ is 0, B3 is 2'-OMe, n⁵ is 3, B1' is 2'-OMe or 2'-F, q¹ is 9, T1' is 2'-F, q² is 1, B2' is 2'-OMe or 2'-F, q³ is 4, T2' is 2'-F, q⁴ is 2, B3' is 2'-OMe or 2'-F, q⁵ is 5, T3' is 2'-F, q⁶ is 1, B4' is 2'-F, and q⁷ is 1. The dsRNAi RNA agent also comprises a 5'-PS₂.

- 30 In one embodiment, B1 is 2'-OMe or 2'-F, n¹ is 8, T1 is 2'-F, n² is 3, B2 is 2'-OMe, n³ is 7, n⁴ is 0, B3 is 2'-OMe, n⁵ is 3, B1' is 2'-OMe or 2'-F, q¹ is 9, T1' is 2'-F, q² is 1, B2' is 2'-OMe or 2'-F, q³ is 4, T2' is 2'-F, q⁴ is 2, B3' is 2'-OMe or 2'-F, q⁵ is 5, T3' is 2'-F, q⁶ is 1, B4' is 2'-F, and q⁷ is 1. The RNAi agent also comprises a 5'-deoxy-5'-C-malonyl.

In one embodiment, B1 is 2'-OMe or 2'-F, n¹ is 8, T1 is 2'-F, n² is 3, B2 is 2'-OMe, n³ is 7, n⁴ is 0, B3 is 2'-OMe, n⁵ is 3, B1' is 2'-OMe or 2'-F, q¹ is 9, T1' is 2'-F, q² is 1, B2' is 2'-OMe or 2'-F, q³ is 4, T2' is 2'-F, q⁴ is 2, B3' is 2'-OMe or 2'-F, q⁵ is 5, T3' is 2'-F, q⁶ is 1, B4' is 2'-F, and q⁷ is 1; with two phosphorothioate internucleotide linkage modifications within position 1-5 of the sense strand (counting from the 5'-end of the sense strand), and two phosphorothioate internucleotide linkage modifications at positions 1 and 2 and two phosphorothioate internucleotide linkage modifications within positions 18-23 of the antisense strand (counting from the 5'-end of the antisense strand). The RNAi agent also comprises a 5'-P.

In one embodiment, B1 is 2'-OMe or 2'-F, n¹ is 8, T1 is 2'-F, n² is 3, B2 is 2'-OMe, n³ is 7, n⁴ is 0, B3 is 2'-OMe, n⁵ is 3, B1' is 2'-OMe or 2'-F, q¹ is 9, T1' is 2'-F, q² is 1, B2' is 2'-OMe or 2'-F, q³ is 4, T2' is 2'-F, q⁴ is 2, B3' is 2'-OMe or 2'-F, q⁵ is 5, T3' is 2'-F, q⁶ is 1, B4' is 2'-F, and q⁷ is 1; with two phosphorothioate internucleotide linkage modifications within position 1-5 of the sense strand (counting from the 5'-end of the sense strand), and two phosphorothioate internucleotide linkage modifications at positions 1 and 2 and two phosphorothioate internucleotide linkage modifications within positions 18-23 of the antisense strand (counting from the 5'-end of the antisense strand). The RNAi agent also comprises a 5'-PS.

In one embodiment, B1 is 2'-OMe or 2'-F, n¹ is 8, T1 is 2'-F, n² is 3, B2 is 2'-OMe, n³ is 7, n⁴ is 0, B3 is 2'-OMe, n⁵ is 3, B1' is 2'-OMe or 2'-F, q¹ is 9, T1' is 2'-F, q² is 1, B2' is 2'-OMe or 2'-F, q³ is 4, T2' is 2'-F, q⁴ is 2, B3' is 2'-OMe or 2'-F, q⁵ is 5, T3' is 2'-F, q⁶ is 1, B4' is 2'-F, and q⁷ is 1; with two phosphorothioate internucleotide linkage modifications within position 1-5 of the sense strand (counting from the 5'-end of the sense strand), and two phosphorothioate internucleotide linkage modifications at positions 1 and 2 and two phosphorothioate internucleotide linkage modifications within positions 18-23 of the antisense strand (counting from the 5'-end of the antisense strand). The RNAi agent also comprises a 5'-VP. The 5'-VP may be 5'-E-VP, 5'-Z-VP, or combination thereof.

In one embodiment, B1 is 2'-OMe or 2'-F, n¹ is 8, T1 is 2'-F, n² is 3, B2 is 2'-OMe, n³ is 7, n⁴ is 0, B3 is 2'-OMe, n⁵ is 3, B1' is 2'-OMe or 2'-F, q¹ is 9, T1' is 2'-F, q² is 1, B2' is 2'-OMe or 2'-F, q³ is 4, T2' is 2'-F, q⁴ is 2, B3' is 2'-OMe or 2'-F, q⁵ is 5, T3' is 2'-F, q⁶ is 1, B4' is 2'-F, and q⁷ is 1; with two phosphorothioate internucleotide linkage modifications within position 1-5 of the sense strand (counting from the 5'-end of the sense strand), and two phosphorothioate internucleotide linkage modifications at positions 1 and 2 and two phosphorothioate internucleotide linkage modifications within positions 18-23 of the antisense strand (counting from the 5'-end of the antisense strand). The RNAi agent also comprises a 5'-PS₂.

In one embodiment, B1 is 2'-OMe or 2'-F, n¹ is 8, T1 is 2'-F, n² is 3, B2 is 2'-OMe, n³ is 7, n⁴ is 0, B3 is 2'-OMe, n⁵ is 3, B1' is 2'-OMe or 2'-F, q¹ is 9, T1' is 2'-F, q² is 1, B2' is 2'-OMe or 2'-F, q³ is 4, T2' is 2'-F, q⁴ is 2, B3' is 2'-OMe or 2'-F, q⁵ is 5, T3' is 2'-F, q⁶ is 1, B4' is 2'-F, and q⁷ is 1; with two phosphorothioate internucleotide linkage modifications within position 1-5 of the sense

strand (counting from the 5'-end of the sense strand), and two phosphorothioate internucleotide linkage modifications at positions 1 and 2 and two phosphorothioate internucleotide linkage modifications within positions 18-23 of the antisense strand (counting from the 5'-end of the antisense strand). The RNAi agent also comprises a 5'-deoxy-5'-C-malonyl.

5 In one embodiment, B1 is 2'-OMe or 2'-F, n^1 is 8, T1 is 2'-F, n^2 is 3, B2 is 2'-OMe, n^3 is 7, n^4 is 0, B3 is 2'-OMe, n^5 is 3, B1' is 2'-OMe or 2'-F, q^1 is 9, T1' is 2'-F, q^2 is 1, B2' is 2'-OMe or 2'-F, q^3 is 4, q^4 is 0, B3' is 2'-OMe or 2'-F, q^5 is 7, T3' is 2'-F, q^6 is 1, B4' is 2'-F, and q^7 is 1. The RNAi agent also comprises a 5'-P.

10 In one embodiment, B1 is 2'-OMe or 2'-F, n^1 is 8, T1 is 2'-F, n^2 is 3, B2 is 2'-OMe, n^3 is 7, n^4 is 0, B3 is 2'-OMe, n^5 is 3, B1' is 2'-OMe or 2'-F, q^1 is 9, T1' is 2'-F, q^2 is 1, B2' is 2'-OMe or 2'-F, q^3 is 4, q^4 is 0, B3' is 2'-OMe or 2'-F, q^5 is 7, T3' is 2'-F, q^6 is 1, B4' is 2'-F, and q^7 is 1. The RNAi agent also comprises a 5'-PS.

15 In one embodiment, B1 is 2'-OMe or 2'-F, n^1 is 8, T1 is 2'-F, n^2 is 3, B2 is 2'-OMe, n^3 is 7, n^4 is 0, B3 is 2'-OMe, n^5 is 3, B1' is 2'-OMe or 2'-F, q^1 is 9, T1' is 2'-F, q^2 is 1, B2' is 2'-OMe or 2'-F, q^3 is 4, q^4 is 0, B3' is 2'-OMe or 2'-F, q^5 is 7, T3' is 2'-F, q^6 is 1, B4' is 2'-F, and q^7 is 1. The RNAi agent also comprises a 5'-VP. The 5'-VP may be 5'-E-VP, 5'-Z-VP, or combination thereof.

20 In one embodiment, B1 is 2'-OMe or 2'-F, n^1 is 8, T1 is 2'-F, n^2 is 3, B2 is 2'-OMe, n^3 is 7, n^4 is 0, B3 is 2'-OMe, n^5 is 3, B1' is 2'-OMe or 2'-F, q^1 is 9, T1' is 2'-F, q^2 is 1, B2' is 2'-OMe or 2'-F, q^3 is 4, q^4 is 0, B3' is 2'-OMe or 2'-F, q^5 is 7, T3' is 2'-F, q^6 is 1, B4' is 2'-F, and q^7 is 1. The RNAi agent also comprises a 5'-PS₂.

In one embodiment, B1 is 2'-OMe or 2'-F, n^1 is 8, T1 is 2'-F, n^2 is 3, B2 is 2'-OMe, n^3 is 7, n^4 is 0, B3 is 2'-OMe, n^5 is 3, B1' is 2'-OMe or 2'-F, q^1 is 9, T1' is 2'-F, q^2 is 1, B2' is 2'-OMe or 2'-F, q^3 is 4, q^4 is 0, B3' is 2'-OMe or 2'-F, q^5 is 7, T3' is 2'-F, q^6 is 1, B4' is 2'-F, and q^7 is 1. The RNAi agent also comprises a 5'-deoxy-5'-C-malonyl.

25 In one embodiment, B1 is 2'-OMe or 2'-F, n^1 is 8, T1 is 2'-F, n^2 is 3, B2 is 2'-OMe, n^3 is 7, n^4 is 0, B3 is 2'-OMe, n^5 is 3, B1' is 2'-OMe or 2'-F, q^1 is 9, T1' is 2'-F, q^2 is 1, B2' is 2'-OMe or 2'-F, q^3 is 4, q^4 is 0, B3' is 2'-OMe or 2'-F, q^5 is 7, T3' is 2'-F, q^6 is 1, B4' is 2'-F, and q^7 is 1; with two phosphorothioate internucleotide linkage modifications within position 1-5 of the sense strand (counting from the 5'-end of the sense strand), and two phosphorothioate internucleotide linkage modifications at positions 1 and 2 and two phosphorothioate internucleotide linkage modifications within positions 18-23 of the antisense strand (counting from the 5'-end of the antisense strand). The RNAi agent also comprises a 5'-P.

30 In one embodiment, B1 is 2'-OMe or 2'-F, n^1 is 8, T1 is 2'-F, n^2 is 3, B2 is 2'-OMe, n^3 is 7, n^4 is 0, B3 is 2'-OMe, n^5 is 3, B1' is 2'-OMe or 2'-F, q^1 is 9, T1' is 2'-F, q^2 is 1, B2' is 2'-OMe or 2'-F, q^3 is 4, q^4 is 0, B3' is 2'-OMe or 2'-F, q^5 is 7, T3' is 2'-F, q^6 is 1, B4' is 2'-F, and q^7 is 1; with two phosphorothioate internucleotide linkage modifications within position 1-5 of the sense strand (counting from the 5'-end of the sense strand), and two phosphorothioate internucleotide linkage

modifications at positions 1 and 2 and two phosphorothioate internucleotide linkage modifications within positions 18-23 of the antisense strand (counting from the 5'-end of the antisense strand). The RNAi agent also comprises a 5'-PS.

In one embodiment, B1 is 2'-OMe or 2'-F, n^1 is 8, T1 is 2'-F, n^2 is 3, B2 is 2'-OMe, n^3 is 7, n^4 is 0, B3 is 2'-OMe, n^5 is 3, B1' is 2'-OMe or 2'-F, q^1 is 9, T1' is 2'-F, q^2 is 1, B2' is 2'-OMe or 2'-F, q^3 is 4, q^4 is 0, B3' is 2'-OMe or 2'-F, q^5 is 7, T3' is 2'-F, q^6 is 1, B4' is 2'-F, and q^7 is 1; with two phosphorothioate internucleotide linkage modifications within position 1-5 of the sense strand (counting from the 5'-end of the sense strand), and two phosphorothioate internucleotide linkage modifications at positions 1 and 2 and two phosphorothioate internucleotide linkage modifications within positions 18-23 of the antisense strand (counting from the 5'-end of the antisense strand). The RNAi agent also comprises a 5'-VP. The 5'-VP may be 5'-E-VP, 5'-Z-VP, or combination thereof.

In one embodiment, B1 is 2'-OMe or 2'-F, n^1 is 8, T1 is 2'-F, n^2 is 3, B2 is 2'-OMe, n^3 is 7, n^4 is 0, B3 is 2'-OMe, n^5 is 3, B1' is 2'-OMe or 2'-F, q^1 is 9, T1' is 2'-F, q^2 is 1, B2' is 2'-OMe or 2'-F, q^3 is 4, q^4 is 0, B3' is 2'-OMe or 2'-F, q^5 is 7, T3' is 2'-F, q^6 is 1, B4' is 2'-F, and q^7 is 1; with two phosphorothioate internucleotide linkage modifications within position 1-5 of the sense strand (counting from the 5'-end of the sense strand), and two phosphorothioate internucleotide linkage modifications at positions 1 and 2 and two phosphorothioate internucleotide linkage modifications within positions 18-23 of the antisense strand (counting from the 5'-end of the antisense strand). The RNAi agent also comprises a 5'-PS₂.

In one embodiment, B1 is 2'-OMe or 2'-F, n^1 is 8, T1 is 2'-F, n^2 is 3, B2 is 2'-OMe, n^3 is 7, n^4 is 0, B3 is 2'-OMe, n^5 is 3, B1' is 2'-OMe or 2'-F, q^1 is 9, T1' is 2'-F, q^2 is 1, B2' is 2'-OMe or 2'-F, q^3 is 4, q^4 is 0, B3' is 2'-OMe or 2'-F, q^5 is 7, T3' is 2'-F, q^6 is 1, B4' is 2'-F, and q^7 is 1; with two phosphorothioate internucleotide linkage modifications within position 1-5 of the sense strand (counting from the 5'-end of the sense strand), and two phosphorothioate internucleotide linkage modifications at positions 1 and 2 and two phosphorothioate internucleotide linkage modifications within positions 18-23 of the antisense strand (counting from the 5'-end of the antisense strand). The RNAi agent also comprises a 5'-deoxy-5'-C-malonyl.

In one embodiment, B1 is 2'-OMe or 2'-F, n^1 is 8, T1 is 2'-F, n^2 is 3, B2 is 2'-OMe, n^3 is 7, n^4 is 0, B3 is 2'-OMe, n^5 is 3, B1' is 2'-OMe or 2'-F, q^1 is 9, T1' is 2'-F, q^2 is 1, B2' is 2'-OMe or 2'-F, q^3 is 4, T2' is 2'-F, q^4 is 2, B3' is 2'-OMe or 2'-F, q^5 is 5, T3' is 2'-F, q^6 is 1, B4' is 2'-OMe, and q^7 is 1; with two phosphorothioate internucleotide linkage modifications within position 1-5 of the sense strand (counting from the 5'-end of the sense strand), and two phosphorothioate internucleotide linkage modifications at positions 1 and 2 and two phosphorothioate internucleotide linkage modifications within positions 18-23 of the antisense strand (counting from the 5'-end of the antisense strand). The RNAi agent also comprises a 5'-P and a targeting ligand. In one embodiment, the 5'-P is at the 5'-end of the antisense strand, and the targeting ligand is at the 3'-end of the sense strand.

In one embodiment, B1 is 2'-OMe or 2'-F, n^1 is 8, T1 is 2'-F, n^2 is 3, B2 is 2'-OMe, n^3 is 7, n^4 is 0, B3 is 2'-OMe, n^5 is 3, B1' is 2'-OMe or 2'-F, q^1 is 9, T1' is 2'-F, q^2 is 1, B2' is 2'-OMe or 2'-F, q^3 is 4, T2' is 2'-F, q^4 is 2, B3' is 2'-OMe or 2'-F, q^5 is 5, T3' is 2'-F, q^6 is 1, B4' is 2'-OMe, and q^7 is 1; with two phosphorothioate internucleotide linkage modifications within position 1-5 of the sense strand (counting from the 5'-end of the sense strand), and two phosphorothioate internucleotide linkage modifications at positions 1 and 2 and two phosphorothioate internucleotide linkage modifications within positions 18-23 of the antisense strand (counting from the 5'-end of the antisense strand). The RNAi agent also comprises a 5'-PS and a targeting ligand. In one embodiment, the 5'-PS is at the 5'-end of the antisense strand, and the targeting ligand is at the 3'-end of the sense strand.

In one embodiment, B1 is 2'-OMe or 2'-F, n^1 is 8, T1 is 2'-F, n^2 is 3, B2 is 2'-OMe, n^3 is 7, n^4 is 0, B3 is 2'-OMe, n^5 is 3, B1' is 2'-OMe or 2'-F, q^1 is 9, T1' is 2'-F, q^2 is 1, B2' is 2'-OMe or 2'-F, q^3 is 4, T2' is 2'-F, q^4 is 2, B3' is 2'-OMe or 2'-F, q^5 is 5, T3' is 2'-F, q^6 is 1, B4' is 2'-OMe, and q^7 is 1; with two phosphorothioate internucleotide linkage modifications within position 1-5 of the sense strand (counting from the 5'-end of the sense strand), and two phosphorothioate internucleotide linkage modifications at positions 1 and 2 and two phosphorothioate internucleotide linkage modifications within positions 18-23 of the antisense strand (counting from the 5'-end of the antisense strand). The RNAi agent also comprises a 5'-VP (*e.g.*, a 5'-E-VP, 5'-Z-VP, or combination thereof), and a targeting ligand.

In one embodiment, the 5'-VP is at the 5'-end of the antisense strand, and the targeting ligand is at the 3'-end of the sense strand.

In one embodiment, B1 is 2'-OMe or 2'-F, n^1 is 8, T1 is 2'-F, n^2 is 3, B2 is 2'-OMe, n^3 is 7, n^4 is 0, B3 is 2'-OMe, n^5 is 3, B1' is 2'-OMe or 2'-F, q^1 is 9, T1' is 2'-F, q^2 is 1, B2' is 2'-OMe or 2'-F, q^3 is 4, T2' is 2'-F, q^4 is 2, B3' is 2'-OMe or 2'-F, q^5 is 5, T3' is 2'-F, q^6 is 1, B4' is 2'-OMe, and q^7 is 1; with two phosphorothioate internucleotide linkage modifications within position 1-5 of the sense strand (counting from the 5'-end of the sense strand), and two phosphorothioate internucleotide linkage modifications at positions 1 and 2 and two phosphorothioate internucleotide linkage modifications within positions 18-23 of the antisense strand (counting from the 5'-end of the antisense strand). The RNAi agent also comprises a 5'-PS₂ and a targeting ligand. In one embodiment, the 5'-PS₂ is at the 5'-end of the antisense strand, and the targeting ligand is at the 3'-end of the sense strand.

In one embodiment, B1 is 2'-OMe or 2'-F, n^1 is 8, T1 is 2'-F, n^2 is 3, B2 is 2'-OMe, n^3 is 7, n^4 is 0, B3 is 2'-OMe, n^5 is 3, B1' is 2'-OMe or 2'-F, q^1 is 9, T1' is 2'-F, q^2 is 1, B2' is 2'-OMe or 2'-F, q^3 is 4, T2' is 2'-F, q^4 is 2, B3' is 2'-OMe or 2'-F, q^5 is 5, T3' is 2'-F, q^6 is 1, B4' is 2'-OMe, and q^7 is 1; with two phosphorothioate internucleotide linkage modifications within position 1-5 of the sense strand (counting from the 5'-end of the sense strand), and two phosphorothioate internucleotide linkage modifications at positions 1 and 2 and two phosphorothioate internucleotide linkage modifications within positions 18-23 of the antisense strand (counting from the 5'-end of the antisense strand). The RNAi agent also comprises a 5'-deoxy-5'-C-malonyl and a targeting ligand. In one

embodiment, the 5'-deoxy-5'-C-malonyl is at the 5'-end of the antisense strand, and the targeting ligand is at the 3'-end of the sense strand.

In one embodiment, B1 is 2'-OMe or 2'-F, n^1 is 8, T1 is 2'-F, n^2 is 3, B2 is 2'-OMe, n^3 is 7, n^4 is 0, B3 is 2'-OMe, n^5 is 3, B1' is 2'-OMe or 2'-F, q^1 is 9, T1' is 2'-F, q^2 is 1, B2' is 2'-OMe or 2'-F, q^3 is 4, q^4 is 0, B3' is 2'-OMe or 2'-F, q^5 is 7, T3' is 2'-F, q^6 is 1, B4' is 2'-OMe, and q^7 is 1; with two phosphorothioate internucleotide linkage modifications within position 1-5 of the sense strand (counting from the 5'-end), and two phosphorothioate internucleotide linkage modifications at positions 1 and 2 and two phosphorothioate internucleotide linkage modifications within positions 18-23 of the antisense strand (counting from the 5'-end). The RNAi agent also comprises a 5'-P and a targeting ligand. In one embodiment, the 5'-P is at the 5'-end of the antisense strand, and the targeting ligand is at the 3'-end of the sense strand.

In one embodiment, B1 is 2'-OMe or 2'-F, n^1 is 8, T1 is 2'-F, n^2 is 3, B2 is 2'-OMe, n^3 is 7, n^4 is 0, B3 is 2'-OMe, n^5 is 3, B1' is 2'-OMe or 2'-F, q^1 is 9, T1' is 2'-F, q^2 is 1, B2' is 2'-OMe or 2'-F, q^3 is 4, q^4 is 0, B3' is 2'-OMe or 2'-F, q^5 is 7, T3' is 2'-F, q^6 is 1, B4' is 2'-OMe, and q^7 is 1; with two phosphorothioate internucleotide linkage modifications within position 1-5 of the sense strand (counting from the 5'-end), and two phosphorothioate internucleotide linkage modifications at positions 1 and 2 and two phosphorothioate internucleotide linkage modifications within positions 18-23 of the antisense strand (counting from the 5'-end). The RNAi agent also comprises a 5'-PS and a targeting ligand. In one embodiment, the 5'-PS is at the 5'-end of the antisense strand, and the targeting ligand is at the 3'-end of the sense strand.

In one embodiment, B1 is 2'-OMe or 2'-F, n^1 is 8, T1 is 2'-F, n^2 is 3, B2 is 2'-OMe, n^3 is 7, n^4 is 0, B3 is 2'-OMe, n^5 is 3, B1' is 2'-OMe or 2'-F, q^1 is 9, T1' is 2'-F, q^2 is 1, B2' is 2'-OMe or 2'-F, q^3 is 4, q^4 is 0, B3' is 2'-OMe or 2'-F, q^5 is 7, T3' is 2'-F, q^6 is 1, B4' is 2'-OMe, and q^7 is 1; with two phosphorothioate internucleotide linkage modifications within position 1-5 of the sense strand (counting from the 5'-end), and two phosphorothioate internucleotide linkage modifications at positions 1 and 2 and two phosphorothioate internucleotide linkage modifications within positions 18-23 of the antisense strand (counting from the 5'-end). The RNAi agent also comprises a 5'-VP (*e.g.*, a 5'-E-VP, 5'-Z-VP, or combination thereof) and a targeting ligand. In one embodiment, the 5'-VP is at the 5'-end of the antisense strand, and the targeting ligand is at the 3'-end of the sense strand.

In one embodiment, B1 is 2'-OMe or 2'-F, n^1 is 8, T1 is 2'-F, n^2 is 3, B2 is 2'-OMe, n^3 is 7, n^4 is 0, B3 is 2'-OMe, n^5 is 3, B1' is 2'-OMe or 2'-F, q^1 is 9, T1' is 2'-F, q^2 is 1, B2' is 2'-OMe or 2'-F, q^3 is 4, q^4 is 0, B3' is 2'-OMe or 2'-F, q^5 is 7, T3' is 2'-F, q^6 is 1, B4' is 2'-OMe, and q^7 is 1; with two phosphorothioate internucleotide linkage modifications within position 1-5 of the sense strand (counting from the 5'-end), and two phosphorothioate internucleotide linkage modifications at positions 1 and 2 and two phosphorothioate internucleotide linkage modifications within positions 18-23 of the antisense strand (counting from the 5'-end). The RNAi agent also comprises a 5'-PS₂ and a

targeting ligand. In one embodiment, the 5'-PS₂ is at the 5'-end of the antisense strand, and the targeting ligand is at the 3'-end of the sense strand.

In one embodiment, B1 is 2'-OMe or 2'-F, n¹ is 8, T1 is 2'-F, n² is 3, B2 is 2'-OMe, n³ is 7, n⁴ is 0, B3 is 2'-OMe, n⁵ is 3, B1' is 2'-OMe or 2'-F, q¹ is 9, T1' is 2'-F, q² is 1, B2' is 2'-OMe or 2'-F, q³ is 4, q⁴ is 0, B3' is 2'-OMe or 2'-F, q⁵ is 7, T3' is 2'-F, q⁶ is 1, B4' is 2'-OMe, and q⁷ is 1; with two phosphorothioate internucleotide linkage modifications within position 1-5 of the sense strand (counting from the 5'-end), and two phosphorothioate internucleotide linkage modifications at positions 1 and 2 and two phosphorothioate internucleotide linkage modifications within positions 18-23 of the antisense strand (counting from the 5'-end). The RNAi agent also comprises a 5'-deoxy-5'-C-malonyl and a targeting ligand. In one embodiment, the 5'-deoxy-5'-C-malonyl is at the 5'-end of the antisense strand, and the targeting ligand is at the 3'-end of the sense strand.

In one embodiment, B1 is 2'-OMe or 2'-F, n¹ is 8, T1 is 2'-F, n² is 3, B2 is 2'-OMe, n³ is 7, n⁴ is 0, B3 is 2'-OMe, n⁵ is 3, B1' is 2'-OMe or 2'-F, q¹ is 9, T1' is 2'-F, q² is 1, B2' is 2'-OMe or 2'-F, q³ is 4, T2' is 2'-F, q⁴ is 2, B3' is 2'-OMe or 2'-F, q⁵ is 5, T3' is 2'-F, q⁶ is 1, B4' is 2'-F, and q⁷ is 1; with two phosphorothioate internucleotide linkage modifications within position 1-5 of the sense strand (counting from the 5'-end of the sense strand), and two phosphorothioate internucleotide linkage modifications at positions 1 and 2 and two phosphorothioate internucleotide linkage modifications within positions 18-23 of the antisense strand (counting from the 5'-end of the antisense strand). The RNAi agent also comprises a 5'-P and a targeting ligand. In one embodiment, the 5'-P is at the 5'-end of the antisense strand, and the targeting ligand is at the 3'-end of the sense strand.

In one embodiment, B1 is 2'-OMe or 2'-F, n¹ is 8, T1 is 2'-F, n² is 3, B2 is 2'-OMe, n³ is 7, n⁴ is 0, B3 is 2'-OMe, n⁵ is 3, B1' is 2'-OMe or 2'-F, q¹ is 9, T1' is 2'-F, q² is 1, B2' is 2'-OMe or 2'-F, q³ is 4, T2' is 2'-F, q⁴ is 2, B3' is 2'-OMe or 2'-F, q⁵ is 5, T3' is 2'-F, q⁶ is 1, B4' is 2'-F, and q⁷ is 1; with two phosphorothioate internucleotide linkage modifications within position 1-5 of the sense strand (counting from the 5'-end of the sense strand), and two phosphorothioate internucleotide linkage modifications at positions 1 and 2 and two phosphorothioate internucleotide linkage modifications within positions 18-23 of the antisense strand (counting from the 5'-end of the antisense strand). The RNAi agent also comprises a 5'-PS and a targeting ligand. In one embodiment, the 5'-PS is at the 5'-end of the antisense strand, and the targeting ligand is at the 3'-end of the sense strand.

In one embodiment, B1 is 2'-OMe or 2'-F, n¹ is 8, T1 is 2'-F, n² is 3, B2 is 2'-OMe, n³ is 7, n⁴ is 0, B3 is 2'-OMe, n⁵ is 3, B1' is 2'-OMe or 2'-F, q¹ is 9, T1' is 2'-F, q² is 1, B2' is 2'-OMe or 2'-F, q³ is 4, T2' is 2'-F, q⁴ is 2, B3' is 2'-OMe or 2'-F, q⁵ is 5, T3' is 2'-F, q⁶ is 1, B4' is 2'-F, and q⁷ is 1; with two phosphorothioate internucleotide linkage modifications within position 1-5 of the sense strand (counting from the 5'-end of the sense strand), and two phosphorothioate internucleotide linkage modifications at positions 1 and 2 and two phosphorothioate internucleotide linkage modifications within positions 18-23 of the antisense strand (counting from the 5'-end of the antisense strand). The RNAi agent also comprises a 5'-VP (e.g., a 5'-E-VP, 5'-Z-VP, or combination thereof)

and a targeting ligand. In one embodiment, the 5'-VP is at the 5'-end of the antisense strand, and the targeting ligand is at the 3'-end of the sense strand.

In one embodiment, B1 is 2'-OMe or 2'-F, n^1 is 8, T1 is 2'-F, n^2 is 3, B2 is 2'-OMe, n^3 is 7, n^4 is 0, B3 is 2'-OMe, n^5 is 3, B1' is 2'-OMe or 2'-F, q^1 is 9, T1' is 2'-F, q^2 is 1, B2' is 2'-OMe or 2'-F, q^3 is 4, T2' is 2'-F, q^4 is 2, B3' is 2'-OMe or 2'-F, q^5 is 5, T3' is 2'-F, q^6 is 1, B4' is 2'-F, and q^7 is 1; with two phosphorothioate internucleotide linkage modifications within position 1-5 of the sense strand (counting from the 5'-end of the sense strand), and two phosphorothioate internucleotide linkage modifications at positions 1 and 2 and two phosphorothioate internucleotide linkage modifications within positions 18-23 of the antisense strand (counting from the 5'-end of the antisense strand). The RNAi agent also comprises a 5'-PS₂ and a targeting ligand. In one embodiment, the 5'-PS₂ is at the 5'-end of the antisense strand, and the targeting ligand is at the 3'-end of the sense strand.

In one embodiment, B1 is 2'-OMe or 2'-F, n^1 is 8, T1 is 2'-F, n^2 is 3, B2 is 2'-OMe, n^3 is 7, n^4 is 0, B3 is 2'-OMe, n^5 is 3, B1' is 2'-OMe or 2'-F, q^1 is 9, T1' is 2'-F, q^2 is 1, B2' is 2'-OMe or 2'-F, q^3 is 4, T2' is 2'-F, q^4 is 2, B3' is 2'-OMe or 2'-F, q^5 is 5, T3' is 2'-F, q^6 is 1, B4' is 2'-F, and q^7 is 1; with two phosphorothioate internucleotide linkage modifications within position 1-5 of the sense strand (counting from the 5'-end of the sense strand), and two phosphorothioate internucleotide linkage modifications at positions 1 and 2 and two phosphorothioate internucleotide linkage modifications within positions 18-23 of the antisense strand (counting from the 5'-end of the antisense strand). The RNAi agent also comprises a 5'-deoxy-5'-C-malonyl and a targeting ligand. In one embodiment, the 5'-deoxy-5'-C-malonyl is at the 5'-end of the antisense strand, and the targeting ligand is at the 3'-end of the sense strand.

In one embodiment, B1 is 2'-OMe or 2'-F, n^1 is 8, T1 is 2'-F, n^2 is 3, B2 is 2'-OMe, n^3 is 7, n^4 is 0, B3 is 2'-OMe, n^5 is 3, B1' is 2'-OMe or 2'-F, q^1 is 9, T1' is 2'-F, q^2 is 1, B2' is 2'-OMe or 2'-F, q^3 is 4, q^4 is 0, B3' is 2'-OMe or 2'-F, q^5 is 7, T3' is 2'-F, q^6 is 1, B4' is 2'-F, and q^7 is 1; with two phosphorothioate internucleotide linkage modifications within position 1-5 of the sense strand (counting from the 5'-end of the sense strand), and two phosphorothioate internucleotide linkage modifications at positions 1 and 2 and two phosphorothioate internucleotide linkage modifications within positions 18-23 of the antisense strand (counting from the 5'-end of the antisense strand). The RNAi agent also comprises a 5'-P and a targeting ligand. In one embodiment, the 5'-P is at the 5'-end of the antisense strand, and the targeting ligand is at the 3'-end of the sense strand.

In one embodiment, B1 is 2'-OMe or 2'-F, n^1 is 8, T1 is 2'-F, n^2 is 3, B2 is 2'-OMe, n^3 is 7, n^4 is 0, B3 is 2'-OMe, n^5 is 3, B1' is 2'-OMe or 2'-F, q^1 is 9, T1' is 2'-F, q^2 is 1, B2' is 2'-OMe or 2'-F, q^3 is 4, q^4 is 0, B3' is 2'-OMe or 2'-F, q^5 is 7, T3' is 2'-F, q^6 is 1, B4' is 2'-F, and q^7 is 1; with two phosphorothioate internucleotide linkage modifications within position 1-5 of the sense strand (counting from the 5'-end of the sense strand), and two phosphorothioate internucleotide linkage modifications at positions 1 and 2 and two phosphorothioate internucleotide linkage modifications within positions 18-23 of the antisense strand (counting from the 5'-end of the antisense strand). The

RNAi agent also comprises a 5'-PS and a targeting ligand. In one embodiment, the 5'-PS is at the 5'-end of the antisense strand, and the targeting ligand is at the 3'-end of the sense strand.

In one embodiment, B1 is 2'-OMe or 2'-F, n^1 is 8, T1 is 2'-F, n^2 is 3, B2 is 2'-OMe, n^3 is 7, n^4 is 0, B3 is 2'-OMe, n^5 is 3, B1' is 2'-OMe or 2'-F, q^1 is 9, T1' is 2'-F, q^2 is 1, B2' is 2'-OMe or 2'-F, q^3 is 4, q^4 is 0, B3' is 2'-OMe or 2'-F, q^5 is 7, T3' is 2'-F, q^6 is 1, B4' is 2'-F, and q^7 is 1; with two phosphorothioate internucleotide linkage modifications within position 1-5 of the sense strand (counting from the 5'-end of the sense strand), and two phosphorothioate internucleotide linkage modifications at positions 1 and 2 and two phosphorothioate internucleotide linkage modifications within positions 18-23 of the antisense strand (counting from the 5'-end of the antisense strand). The RNAi agent also comprises a 5'-VP (*e.g.*, a 5'-E-VP, 5'-Z-VP, or combination thereof) and a targeting ligand. In one embodiment, the 5'-VP is at the 5'-end of the antisense strand, and the targeting ligand is at the 3'-end of the sense strand.

In one embodiment, B1 is 2'-OMe or 2'-F, n^1 is 8, T1 is 2'-F, n^2 is 3, B2 is 2'-OMe, n^3 is 7, n^4 is 0, B3 is 2'-OMe, n^5 is 3, B1' is 2'-OMe or 2'-F, q^1 is 9, T1' is 2'-F, q^2 is 1, B2' is 2'-OMe or 2'-F, q^3 is 4, q^4 is 0, B3' is 2'-OMe or 2'-F, q^5 is 7, T3' is 2'-F, q^6 is 1, B4' is 2'-F, and q^7 is 1; with two phosphorothioate internucleotide linkage modifications within position 1-5 of the sense strand (counting from the 5'-end of the sense strand), and two phosphorothioate internucleotide linkage modifications at positions 1 and 2 and two phosphorothioate internucleotide linkage modifications within positions 18-23 of the antisense strand (counting from the 5'-end of the antisense strand). The RNAi agent also comprises a 5'-PS₂ and a targeting ligand. In one embodiment, the 5'-PS₂ is at the 5'-end of the antisense strand, and the targeting ligand is at the 3'-end of the sense strand.

In one embodiment, B1 is 2'-OMe or 2'-F, n^1 is 8, T1 is 2'-F, n^2 is 3, B2 is 2'-OMe, n^3 is 7, n^4 is 0, B3 is 2'-OMe, n^5 is 3, B1' is 2'-OMe or 2'-F, q^1 is 9, T1' is 2'-F, q^2 is 1, B2' is 2'-OMe or 2'-F, q^3 is 4, q^4 is 0, B3' is 2'-OMe or 2'-F, q^5 is 7, T3' is 2'-F, q^6 is 1, B4' is 2'-F, and q^7 is 1; with two phosphorothioate internucleotide linkage modifications within position 1-5 of the sense strand (counting from the 5'-end of the sense strand), and two phosphorothioate internucleotide linkage modifications at positions 1 and 2 and two phosphorothioate internucleotide linkage modifications within positions 18-23 of the antisense strand (counting from the 5'-end of the antisense strand). The RNAi agent also comprises a 5'-deoxy-5'-C-malonyl and a targeting ligand. In one embodiment, the 5'-deoxy-5'-C-malonyl is at the 5'-end of the antisense strand, and the targeting ligand is at the 3'-end of the sense strand.

In a particular embodiment, an RNAi agent of the present invention comprises:

- (a) a sense strand having:
 - (i) a length of 21 nucleotides;
 - (ii) an ASGPR ligand attached to the 3'-end, wherein said ASGPR ligand comprises three GalNAc derivatives attached through a trivalent branched linker; and

(iii) 2'-F modifications at positions 1, 3, 5, 7, 9 to 11, 13, 17, 19, and 21, and 2'-OMe modifications at positions 2, 4, 6, 8, 12, 14 to 16, 18, and 20 (counting from the 5' end);
and

(b) an antisense strand having:

- 5 (i) a length of 23 nucleotides;
(ii) 2'-OMe modifications at positions 1, 3, 5, 9, 11 to 13, 15, 17, 19, 21, and 23, and 2'F modifications at positions 2, 4, 6 to 8, 10, 14, 16, 18, 20, and 22 (counting from the 5' end); and
(iii) phosphorothioate internucleotide linkages between nucleotide positions 21 and 22,
10 and between nucleotide positions 22 and 23 (counting from the 5' end);
wherein the dsRNA agents have a two nucleotide overhang at the 3'-end of the antisense strand, and a blunt end at the 5'-end of the antisense strand.

In another particular embodiment, an RNAi agent of the present invention comprises:

(a) a sense strand having:

- 15 (i) a length of 21 nucleotides;
(ii) an ASGPR ligand attached to the 3'-end, wherein said ASGPR ligand comprises three GalNAc derivatives attached through a trivalent branched linker;
(iii) 2'-F modifications at positions 1, 3, 5, 7, 9 to 11, 13, 15, 17, 19, and 21, and 2'-OMe modifications at positions 2, 4, 6, 8, 12, 14, 16, 18, and 20 (counting from the 5' end);
20 and
(iv) phosphorothioate internucleotide linkages between nucleotide positions 1 and 2, and between nucleotide positions 2 and 3 (counting from the 5' end);

and

(b) an antisense strand having:

- 25 (i) a length of 23 nucleotides;
(ii) 2'-OMe modifications at positions 1, 3, 5, 7, 9, 11 to 13, 15, 17, 19, and 21 to 23, and 2'F modifications at positions 2, 4, 6, 8, 10, 14, 16, 18, and 20 (counting from the 5' end);
and
(iii) phosphorothioate internucleotide linkages between nucleotide positions 1 and 2, between
30 nucleotide positions 2 and 3, between nucleotide positions 21 and 22, and between nucleotide positions 22 and 23 (counting from the 5' end);

wherein the RNAi agents have a two nucleotide overhang at the 3'-end of the antisense strand, and a blunt end at the 5'-end of the antisense strand.

In another particular embodiment, a RNAi agent of the present invention comprises:

35 (a) a sense strand having:

- (i) a length of 21 nucleotides;

- (ii) an ASGPR ligand attached to the 3'-end, wherein said ASGPR ligand comprises three GalNAc derivatives attached through a trivalent branched linker;
- (iii) 2'-OMe modifications at positions 1 to 6, 8, 10, and 12 to 21, 2'-F modifications at positions 7, and 9, and a deoxy-nucleotide (*e.g.* dT) at position 11 (counting from the 5' end); and
- (iv) phosphorothioate internucleotide linkages between nucleotide positions 1 and 2, and between nucleotide positions 2 and 3 (counting from the 5' end);
- and
- (b) an antisense strand having:
- (i) a length of 23 nucleotides;
- (ii) 2'-OMe modifications at positions 1, 3, 7, 9, 11, 13, 15, 17, and 19 to 23, and 2'-F modifications at positions 2, 4 to 6, 8, 10, 12, 14, 16, and 18 (counting from the 5' end); and
- (iii) phosphorothioate internucleotide linkages between nucleotide positions 1 and 2, between nucleotide positions 2 and 3, between nucleotide positions 21 and 22, and between nucleotide positions 22 and 23 (counting from the 5' end);
- wherein the RNAi agents have a two nucleotide overhang at the 3'-end of the antisense strand, and a blunt end at the 5'-end of the antisense strand.

In another particular embodiment, a RNAi agent of the present invention comprises:

- (a) a sense strand having:
- (i) a length of 21 nucleotides;
- (ii) an ASGPR ligand attached to the 3'-end, wherein said ASGPR ligand comprises three GalNAc derivatives attached through a trivalent branched linker;
- (iii) 2'-OMe modifications at positions 1 to 6, 8, 10, 12, 14, and 16 to 21, and 2'-F modifications at positions 7, 9, 11, 13, and 15; and
- (iv) phosphorothioate internucleotide linkages between nucleotide positions 1 and 2, and between nucleotide positions 2 and 3 (counting from the 5' end);
- and
- (b) an antisense strand having:
- (i) a length of 23 nucleotides;
- (ii) 2'-OMe modifications at positions 1, 5, 7, 9, 11, 13, 15, 17, 19, and 21 to 23, and 2'-F modifications at positions 2 to 4, 6, 8, 10, 12, 14, 16, 18, and 20 (counting from the 5' end); and
- (iii) phosphorothioate internucleotide linkages between nucleotide positions 1 and 2, between nucleotide positions 2 and 3, between nucleotide positions 21 and 22, and between nucleotide positions 22 and 23 (counting from the 5' end);

wherein the RNAi agents have a two nucleotide overhang at the 3'-end of the antisense strand, and a blunt end at the 5'-end of the antisense strand.

In another particular embodiment, a RNAi agent of the present invention comprises:

- (a) a sense strand having:
 - 5 (i) a length of 21 nucleotides;
 - (ii) an ASGPR ligand attached to the 3'-end, wherein said ASGPR ligand comprises three GalNAc derivatives attached through a trivalent branched linker;
 - (iii) 2'-OMe modifications at positions 1 to 9, and 12 to 21, and 2'-F modifications at positions 10, and 11; and
 - 10 (iv) phosphorothioate internucleotide linkages between nucleotide positions 1 and 2, and between nucleotide positions 2 and 3 (counting from the 5' end);
- and
- (b) an antisense strand having:
 - (i) a length of 23 nucleotides;
 - 15 (ii) 2'-OMe modifications at positions 1, 3, 5, 7, 9, 11 to 13, 15, 17, 19, and 21 to 23, and 2'-F modifications at positions 2, 4, 6, 8, 10, 14, 16, 18, and 20 (counting from the 5' end); and
 - (iii) phosphorothioate internucleotide linkages between nucleotide positions 1 and 2, between nucleotide positions 2 and 3, between nucleotide positions 21 and 22, and between
 - 20 nucleotide positions 22 and 23 (counting from the 5' end);

wherein the RNAi agents have a two nucleotide overhang at the 3'-end of the antisense strand, and a blunt end at the 5'-end of the antisense strand.

In another particular embodiment, a RNAi agent of the present invention comprises:

- (a) a sense strand having:
 - 25 (i) a length of 21 nucleotides;
 - (ii) an ASGPR ligand attached to the 3'-end, wherein said ASGPR ligand comprises three GalNAc derivatives attached through a trivalent branched linker;
 - (iii) 2'-F modifications at positions 1, 3, 5, 7, 9 to 11, and 13, and 2'-OMe modifications at positions 2, 4, 6, 8, 12, and 14 to 21; and
 - 30 (iv) phosphorothioate internucleotide linkages between nucleotide positions 1 and 2, and between nucleotide positions 2 and 3 (counting from the 5' end);
- and
- (b) an antisense strand having:
 - (i) a length of 23 nucleotides;
 - 35 (ii) 2'-OMe modifications at positions 1, 3, 5 to 7, 9, 11 to 13, 15, 17 to 19, and 21 to 23, and 2'-F modifications at positions 2, 4, 8, 10, 14, 16, and 20 (counting from the 5' end); and

- (iii) phosphorothioate internucleotide linkages between nucleotide positions 1 and 2, between nucleotide positions 2 and 3, between nucleotide positions 21 and 22, and between nucleotide positions 22 and 23 (counting from the 5' end);

wherein the RNAi agents have a two nucleotide overhang at the 3'-end of the antisense strand, and a blunt end at the 5'-end of the antisense strand.

In another particular embodiment, a RNAi agent of the present invention comprises:

- (a) a sense strand having:
 - (i) a length of 21 nucleotides;
 - (ii) an ASGPR ligand attached to the 3'-end, wherein said ASGPR ligand comprises three GalNAc derivatives attached through a trivalent branched linker;
 - (iii) 2'-OMe modifications at positions 1, 2, 4, 6, 8, 12, 14, 15, 17, and 19 to 21, and 2'-F modifications at positions 3, 5, 7, 9 to 11, 13, 16, and 18; and
 - (iv) phosphorothioate internucleotide linkages between nucleotide positions 1 and 2, and between nucleotide positions 2 and 3 (counting from the 5' end);

and

- (b) an antisense strand having:
 - (i) a length of 25 nucleotides;
 - (ii) 2'-OMe modifications at positions 1, 4, 6, 7, 9, 11 to 13, 15, 17, and 19 to 23, 2'-F modifications at positions 2, 3, 5, 8, 10, 14, 16, and 18, and deoxy-nucleotides (*e.g.* dT) at positions 24 and 25 (counting from the 5' end); and
 - (iii) phosphorothioate internucleotide linkages between nucleotide positions 1 and 2, between nucleotide positions 2 and 3, between nucleotide positions 21 and 22, and between nucleotide positions 22 and 23 (counting from the 5' end);

wherein the RNAi agents have a four nucleotide overhang at the 3'-end of the antisense strand, and a blunt end at the 5'-end of the antisense strand.

In another particular embodiment, a RNAi agent of the present invention comprises:

- (a) a sense strand having:
 - (i) a length of 21 nucleotides;
 - (ii) an ASGPR ligand attached to the 3'-end, wherein said ASGPR ligand comprises three GalNAc derivatives attached through a trivalent branched linker;
 - (iii) 2'-OMe modifications at positions 1 to 6, 8, and 12 to 21, and 2'-F modifications at positions 7, and 9 to 11; and
 - (iv) phosphorothioate internucleotide linkages between nucleotide positions 1 and 2, and between nucleotide positions 2 and 3 (counting from the 5' end);

and

- (b) an antisense strand having:
 - (i) a length of 23 nucleotides;

- (ii) 2'-OMe modifications at positions 1, 3 to 5, 7, 8, 10 to 13, 15, and 17 to 23, and 2'-F modifications at positions 2, 6, 9, 14, and 16 (counting from the 5' end); and
- (iii) phosphorothioate internucleotide linkages between nucleotide positions 1 and 2, between nucleotide positions 2 and 3, between nucleotide positions 21 and 22, and between nucleotide positions 22 and 23 (counting from the 5' end);
- wherein the RNAi agents have a two nucleotide overhang at the 3'-end of the antisense strand, and a blunt end at the 5'-end of the antisense strand.

In another particular embodiment, a RNAi agent of the present invention comprises:

- (a) a sense strand having:
- (i) a length of 21 nucleotides;
- (ii) an ASGPR ligand attached to the 3'-end, wherein said ASGPR ligand comprises three GalNAc derivatives attached through a trivalent branched linker;
- (iii) 2'-OMe modifications at positions 1 to 6, 8, and 12 to 21, and 2'-F modifications at positions 7, and 9 to 11;
- (iv) phosphorothioate internucleotide linkages between nucleotide positions 1 and 2, and between nucleotide positions 2 and 3 (counting from the 5' end);
- and
- (b) an antisense strand having:
- (i) a length of 23 nucleotides;
- (ii) 2'-OMe modifications at positions 1, 3 to 5, 7, 10 to 13, 15, and 17 to 23, and 2'-F modifications at positions 2, 6, 8, 9, 14, and 16 (counting from the 5' end); and
- (iii) phosphorothioate internucleotide linkages between nucleotide positions 1 and 2, between nucleotide positions 2 and 3, between nucleotide positions 21 and 22, and between nucleotide positions 22 and 23 (counting from the 5' end);
- wherein the RNAi agents have a two nucleotide overhang at the 3'-end of the antisense strand, and a blunt end at the 5'-end of the antisense strand.

In another particular embodiment, a RNAi agent of the present invention comprises:

- (a) a sense strand having:
- (i) a length of 19 nucleotides;
- (ii) an ASGPR ligand attached to the 3'-end, wherein said ASGPR ligand comprises three GalNAc derivatives attached through a trivalent branched linker;
- (iii) 2'-OMe modifications at positions 1 to 4, 6, and 10 to 19, and 2'-F modifications at positions 5, and 7 to 9; and
- (iv) phosphorothioate internucleotide linkages between nucleotide positions 1 and 2, and between nucleotide positions 2 and 3 (counting from the 5' end);
- and
- (b) an antisense strand having:

- (i) a length of 21 nucleotides;
- (ii) 2'-OMe modifications at positions 1, 3 to 5, 7, 10 to 13, 15, and 17 to 21, and 2'-F modifications at positions 2, 6, 8, 9, 14, and 16 (counting from the 5' end); and
- (iii) phosphorothioate internucleotide linkages between nucleotide positions 1 and 2, between
5 nucleotide positions 2 and 3, between nucleotide positions 19 and 20, and between
nucleotide positions 20 and 21 (counting from the 5' end);

wherein the RNAi agents have a two nucleotide overhang at the 3'-end of the antisense strand, and a blunt end at the 5'-end of the antisense strand.

In certain embodiments, the iRNA for use in the methods of the invention is an agent selected
10 from agents listed in any one of Tables 2-3. These agents may further comprise a ligand.

III. iRNAs Conjugated to Ligands

Another modification of the RNA of an iRNA of the invention involves chemically linking to
the iRNA one or more ligands, moieties or conjugates that enhance the activity, cellular distribution,
15 or cellular uptake of the iRNA *e.g.*, into a cell. Such moieties include but are not limited to lipid
moieties such as a cholesterol moiety (Letsinger *et al.*, *Proc. Natl. Acad. Sci. USA*, 1989, 86: 6553-
6556). In other embodiments, the ligand is cholic acid (Manoharan *et al.*, *Biorg. Med. Chem. Lett.*,
1994, 4:1053-1060), a thioether, *e.g.*, beryl-S-tritylthiol (Manoharan *et al.*, *Ann. N.Y. Acad. Sci.*, 1992,
660:306-309; Manoharan *et al.*, *Biorg. Med. Chem. Lett.*, 1993, 3:2765-2770), a thiocholesterol
20 (Oberhauser *et al.*, *Nucl. Acids Res.*, 1992, 20:533-538), an aliphatic chain, *e.g.*, dodecandiol or
undecyl residues (Saison-Behmoaras *et al.*, *EMBO J*, 1991, 10:1111-1118; Kabanov *et al.*, *FEBS
Lett.*, 1990, 259:327-330; Svinarchuk *et al.*, *Biochimie*, 1993, 75:49-54), a phospholipid, *e.g.*, di-
hexadecyl-rac-glycerol or triethyl-ammonium 1,2-di-O-hexadecyl-rac-glycero-3-phosphonate
(Manoharan *et al.*, *Tetrahedron Lett.*, 1995, 36:3651-3654; Shea *et al.*, *Nucl. Acids Res.*, 1990,
25 18:3777-3783), a polyamine or a polyethylene glycol chain (Manoharan *et al.*, *Nucleosides &
Nucleotides*, 1995, 14:969-973), or adamantane acetic acid (Manoharan *et al.*, *Tetrahedron Lett.*,
1995, 36:3651-3654), a palmityl moiety (Mishra *et al.*, *Biochim. Biophys. Acta*, 1995, 1264:229-237),
or an octadecylamine or hexylamino-carbonyloxycholesterol moiety (Crooke *et al.*, *J. Pharmacol.
Exp. Ther.*, 1996, 277:923-937).

30 In certain embodiments, a ligand alters the distribution, targeting, or lifetime of an iRNA
agent into which it is incorporated. In some embodiments a ligand provides an enhanced affinity for a
selected target, *e.g.*, molecule, cell or cell type, compartment, *e.g.*, a cellular or organ compartment,
tissue, organ or region of the body, as, *e.g.*, compared to a species absent such a ligand. In some
embodiments, ligands do not take part in duplex pairing in a duplexed nucleic acid.

35 Ligands can include a naturally occurring substance, such as a protein (*e.g.*, human serum
albumin (HSA), low-density lipoprotein (LDL), or globulin); carbohydrate (*e.g.*, a dextran, pullulan,
chitin, chitosan, inulin, cyclodextrin, N-acetylglucosamine, N-acetylgalactosamine, or hyaluronic

acid); or a lipid. The ligand can also be a recombinant or synthetic molecule, such as a synthetic polymer, *e.g.*, a synthetic polyamino acid. Examples of polyamino acids include polyamino acid is a polylysine (PLL), poly L-aspartic acid, poly L-glutamic acid, styrene-maleic acid anhydride copolymer, poly(L-lactide-co-glycolide) copolymer, divinyl ether-maleic anhydride copolymer, N-(2-hydroxypropyl)methacrylamide copolymer (HMPA), polyethylene glycol (PEG), polyvinyl alcohol (PVA), polyurethane, poly(2-ethylacrylic acid), N-isopropylacrylamide polymers, or polyphosphazene. Example of polyamines include: polyethylenimine, polylysine (PLL), spermine, spermidine, polyamine, pseudopeptide-polyamine, peptidomimetic polyamine, dendrimer polyamine, arginine, amidine, protamine, cationic lipid, cationic porphyrin, quaternary salt of a polyamine, or an alpha helical peptide.

Ligands can also include targeting groups, *e.g.*, a cell or tissue targeting agent, *e.g.*, a lectin, glycoprotein, lipid or protein, *e.g.*, an antibody, that binds to a specified cell type such as a kidney cell. A targeting group can be a thyrotropin, melanotropin, lectin, glycoprotein, surfactant protein A, Mucin carbohydrate, multivalent lactose, multivalent galactose, N-acetyl-galactosamine, N-acetyl-glucosamine multivalent mannose, multivalent fucose, glycosylated polyaminoacids, multivalent galactose, transferrin, bisphosphonate, polyglutamate, polyaspartate, a lipid, cholesterol, a steroid, bile acid, folate, vitamin B12, vitamin A, biotin, or an RGD peptide or RGD peptide mimetic. In certain embodiments, the ligand is a multivalent galactose, *e.g.*, an N-acetyl-galactosamine.

Other examples of ligands include dyes, intercalating agents (*e.g.* acridines), cross-linkers (*e.g.* psoralene, mitomycin C), porphyrins (TPPC4, texaphyrin, Sapphyrin), polycyclic aromatic hydrocarbons (*e.g.*, phenazine, dihydrophenazine), artificial endonucleases (*e.g.* EDTA), lipophilic molecules, *e.g.*, cholesterol, cholic acid, adamantane acetic acid, 1-pyrene butyric acid, dihydrotestosterone, 1,3-Bis-O(hexadecyl)glycerol, geranyloxyhexyl group, hexadecylglycerol, borneol, menthol, 1,3-propanediol, heptadecyl group, palmitic acid, myristic acid, O3-(oleoyl)lithocholic acid, O3-(oleoyl)cholenic acid, dimethoxytrityl, or phenoxazine) and peptide conjugates (*e.g.*, antennapedia peptide, Tat peptide), alkylating agents, phosphate, amino, mercapto, PEG (*e.g.*, PEG-40K), MPEG, [MPEG]₂, polyamino, alkyl, substituted alkyl, radiolabeled markers, enzymes, haptens (*e.g.* biotin), transport/absorption facilitators (*e.g.*, aspirin, vitamin E, folic acid), synthetic ribonucleases (*e.g.*, imidazole, bisimidazole, histamine, imidazole clusters, acridine-imidazole conjugates, Eu³⁺ complexes of tetraazamacrocycles), dinitrophenyl, HRP, or AP.

Ligands can be proteins, *e.g.*, glycoproteins, or peptides, *e.g.*, molecules having a specific affinity for a co-ligand, or antibodies *e.g.*, an antibody, that binds to a specified cell type such as a hepatic cell. Ligands can also include hormones and hormone receptors. They can also include non-peptidic species, such as lipids, lectins, carbohydrates, vitamins, cofactors, multivalent lactose, multivalent galactose, N-acetyl-galactosamine, N-acetyl-glucosamine multivalent mannose, or multivalent fucose. The ligand can be, for example, a lipopolysaccharide, an activator of p38 MAP kinase, or an activator of NF-κB.

The ligand can be a substance, *e.g.*, a drug, which can increase the uptake of the iRNA agent into the cell, for example, by disrupting the cell's cytoskeleton, *e.g.*, by disrupting the cell's microtubules, microfilaments, or intermediate filaments. The drug can be, for example, taxol, vincristine, vinblastine, cytochalasin, nocodazole, japlakinolide, latrunculin A, phalloidin, swinholide
5 A, indanocene, or myoservin.

In some embodiments, a ligand attached to an iRNA as described herein acts as a pharmacokinetic modulator (PK modulator). PK modulators include lipophiles, bile acids, steroids, phospholipid analogues, peptides, protein binding agents, PEG, vitamins, *etc.* Exemplary PK modulators include, but are not limited to, cholesterol, fatty acids, cholic acid, lithocholic acid,
10 dialkylglycerides, diacylglyceride, phospholipids, sphingolipids, naproxen, ibuprofen, vitamin E, biotin. Oligonucleotides that comprise a number of phosphorothioate linkages are also known to bind to serum protein, thus short oligonucleotides, *e.g.*, oligonucleotides of about 5 bases, 10 bases, 15 bases, or 20 bases, comprising multiple of phosphorothioate linkages in the backbone are also amenable to the present invention as ligands (*e.g.* as PK modulating ligands). In addition, aptamers
15 that bind serum components (*e.g.* serum proteins) are also suitable for use as PK modulating ligands in the embodiments described herein.

Ligand-conjugated iRNAs of the invention may be synthesized by the use of an oligonucleotide that bears a pendant reactive functionality, such as that derived from the attachment of a linking molecule onto the oligonucleotide (described below). This reactive oligonucleotide may be
20 reacted directly with commercially-available ligands, ligands that are synthesized bearing any of a variety of protecting groups, or ligands that have a linking moiety attached thereto.

The oligonucleotides used in the conjugates of the present invention may be conveniently and routinely made through the well-known technique of solid-phase synthesis. Equipment for such synthesis is sold by several vendors including, for example, Applied Biosystems® (Foster City,
25 Calif.). Any other methods for such synthesis known in the art may additionally or alternatively be employed. It is also known to use similar techniques to prepare other oligonucleotides, such as the phosphorothioates and alkylated derivatives.

In the ligand-conjugated iRNAs and ligand-molecule bearing sequence-specific linked nucleosides of the present invention, the oligonucleotides and oligonucleosides may be assembled on
30 a suitable DNA synthesizer utilizing standard nucleotide or nucleoside precursors, or nucleotide or nucleoside conjugate precursors that already bear the linking moiety, ligand-nucleotide or nucleoside-conjugate precursors that already bear the ligand molecule, or non-nucleoside ligand-bearing building blocks.

When using nucleotide-conjugate precursors that already bear a linking moiety, the synthesis
35 of the sequence-specific linked nucleosides is typically completed, and the ligand molecule is then reacted with the linking moiety to form the ligand-conjugated oligonucleotide. In some embodiments, the oligonucleotides or linked nucleosides of the present invention are synthesized by an automated

synthesizer using phosphoramidites derived from ligand-nucleoside conjugates in addition to the standard phosphoramidites and non-standard phosphoramidites that are commercially available and routinely used in oligonucleotide synthesis.

A. Lipid Conjugates

5 In certain embodiments, the ligand or conjugate is a lipid or lipid-based molecule. In one embodiment, such a lipid or lipid-based molecule binds a serum protein, *e.g.*, human serum albumin (HSA). An HSA binding ligand allows for distribution of the conjugate to a target tissue, *e.g.*, a non-kidney target tissue of the body. For example, the target tissue can be the liver, including parenchymal cells of the liver. Other molecules that can bind HSA can also be used as ligands. For
10 example, naproxen or aspirin can be used. A lipid or lipid-based ligand can (a) increase resistance to degradation of the conjugate, (b) increase targeting or transport into a target cell or cell membrane, or (c) can be used to adjust binding to a serum protein, *e.g.*, HSA.

A lipid based ligand can be used to inhibit, *e.g.*, control the binding of the conjugate to a target tissue. For example, a lipid or lipid-based ligand that binds to HSA more strongly will be less
15 likely to be targeted to the kidney and therefore less likely to be cleared from the body. A lipid or lipid-based ligand that binds to HSA less strongly can be used to target the conjugate to the kidney.

In certain embodiments, the lipid based ligand binds HSA. In one embodiment, it binds HSA with a sufficient affinity such that the conjugate will be distributed to a non-kidney tissue. However, it is preferred that the affinity not be so strong that the HSA-ligand binding cannot be reversed.

20 In other embodiments, the lipid based ligand binds HSA weakly or not at all. In one embodiment, the conjugate will be distributed to the kidney. Other moieties that target to kidney cells can also be used in place of, or in addition to, the lipid based ligand.

In another aspect, the ligand is a moiety, *e.g.*, a vitamin, which is taken up by a target cell, *e.g.*, a proliferating cell. These are particularly useful for treating disorders characterized by
25 unwanted cell proliferation, *e.g.*, of the malignant or non-malignant type, *e.g.*, cancer cells. Exemplary vitamins include vitamin A, E, and K. Other exemplary vitamins include are B vitamin, *e.g.*, folic acid, B12, riboflavin, biotin, pyridoxal or other vitamins or nutrients taken up by target cells such as liver cells. Also included are HSA and low density lipoprotein (LDL).

30 *B. Cell Permeation Agents*

In another aspect, the ligand is a cell-permeation agent, such as, a helical cell-permeation agent. In one embodiment, the agent is amphipathic. An exemplary agent is a peptide such as tat or antennopodia. If the agent is a peptide, it can be modified, including a peptidylmimetic, invertomers, non-peptide or pseudo-peptide linkages, and use of D-amino acids. In one embodiment, the helical
35 agent is an alpha-helical agent, which has a lipophilic and a lipophobic phase.

The ligand can be a peptide or peptidomimetic. A peptidomimetic (also referred to herein as an oligopeptidomimetic) is a molecule capable of folding into a defined three-dimensional structure

similar to a natural peptide. The attachment of peptide and peptidomimetics to iRNA agents can affect pharmacokinetic distribution of the iRNA, such as by enhancing cellular recognition and absorption. The peptide or peptidomimetic moiety can be about 5-50 amino acids long, *e.g.*, about 5, 10, 15, 20, 25, 30, 35, 40, 45, or 50 amino acids long.

5 A peptide or peptidomimetic can be, for example, a cell permeation peptide, cationic peptide, amphipathic peptide, or hydrophobic peptide (*e.g.*, consisting primarily of Tyr, Trp, or Phe). The peptide moiety can be a dendrimer peptide, constrained peptide or crosslinked peptide. In another alternative, the peptide moiety can include a hydrophobic membrane translocation sequence (MTS). An exemplary hydrophobic MTS-containing peptide is RFGF having the amino acid sequence
 10 AAVALLPAVLLALLAP (SEQ ID NO: 14). An RFGF analogue (*e.g.*, amino acid sequence AALLPVLLAAP (SEQ ID NO:15) containing a hydrophobic MTS can also be a targeting moiety. The peptide moiety can be a “delivery” peptide, which can carry large polar molecules including peptides, oligonucleotides, and protein across cell membranes. For example, sequences from the HIV Tat protein (GRKKRRQRRRPPQ (SEQ ID NO:16) and the Drosophila Antennapedia protein
 15 (RQIKIWFQNRRMKWKK (SEQ ID NO:17) have been found to be capable of functioning as delivery peptides. A peptide or peptidomimetic can be encoded by a random sequence of DNA, such as a peptide identified from a phage-display library, or one-bead-one-compound (OBOC) combinatorial library (Lam *et al.*, Nature, 354:82-84, 1991). Examples of a peptide or peptidomimetic tethered to a dsRNA agent *via* an incorporated monomer unit for cell targeting
 20 purposes is an arginine-glycine-aspartic acid (RGD)-peptide, or RGD mimic. A peptide moiety can range in length from about 5 amino acids to about 40 amino acids. The peptide moieties can have a structural modification, such as to increase stability or direct conformational properties. Any of the structural modifications described below can be utilized.

An RGD peptide for use in the compositions and methods of the invention may be linear or
 25 cyclic, and may be modified, *e.g.*, glycosylated or methylated, to facilitate targeting to a specific tissue(s). RGD-containing peptides and peptidomimetics may include D-amino acids, as well as synthetic RGD mimics. In addition to RGD, one can use other moieties that target the integrin ligand, *e.g.*, PECAM-1 or VEGF.

A “cell permeation peptide” is capable of permeating a cell, *e.g.*, a microbial cell, such as a
 30 bacterial or fungal cell, or a mammalian cell, such as a human cell. A microbial cell-permeating peptide can be, for example, an α -helical linear peptide (*e.g.*, LL-37 or Ceropin P1), a disulfide bond-containing peptide (*e.g.*, α -defensin, β -defensin or bactenecin), or a peptide containing only one or two dominating amino acids (*e.g.*, PR-39 or indolicidin). A cell permeation peptide can also include a nuclear localization signal (NLS). For example, a cell permeation peptide can be a bipartite
 35 amphipathic peptide, such as MPG, which is derived from the fusion peptide domain of HIV-1 gp41 and the NLS of SV40 large T antigen (Simeoni *et al.*, Nucl. Acids Res. 31:2717-2724, 2003).

C. Carbohydrate Conjugates

In some embodiments of the compositions and methods of the invention, an iRNA further comprises a carbohydrate. The carbohydrate conjugated iRNA is advantageous for the *in vivo* delivery of nucleic acids, as well as compositions suitable for *in vivo* therapeutic use, as described
5 herein. As used herein, “carbohydrate” refers to a compound which is either a carbohydrate *per se* made up of one or more monosaccharide units having at least 6 carbon atoms (which can be linear, branched or cyclic) with an oxygen, nitrogen or sulfur atom bonded to each carbon atom; or a compound having as a part thereof a carbohydrate moiety made up of one or more monosaccharide units each having at least six carbon atoms (which can be linear, branched or cyclic), with an oxygen,
10 nitrogen or sulfur atom bonded to each carbon atom. Representative carbohydrates include the sugars (mono-, di-, tri-, and oligosaccharides containing from about 4, 5, 6, 7, 8, or 9 monosaccharide units), and polysaccharides such as starches, glycogen, cellulose and polysaccharide gums. Specific monosaccharides include C5 and above (*e.g.*, C5, C6, C7, or C8) sugars; di- and trisaccharides include sugars having two or three monosaccharide units (*e.g.*, C5, C6, C7, or C8).

15 In certain embodiments, a carbohydrate conjugate for use in the compositions and methods of the invention is a monosaccharide.

In certain embodiments, the monosaccharide is an N-acetylgalactosamine (GalNAc). GalNAc conjugates, which comprise one or more N-acetylgalactosamine (GalNAc) derivatives, are described, for example, in US 8,106,022, the entire content of which is hereby incorporated herein by reference.
20 In some embodiments, the GalNAc conjugate serves as a ligand that targets the iRNA to particular cells. In some embodiments, the GalNAc conjugate targets the iRNA to liver cells, *e.g.*, by serving as a ligand for the asialoglycoprotein receptor of liver cells (*e.g.*, hepatocytes).

In some embodiments, the carbohydrate conjugate comprises one or more GalNAc derivatives. The GalNAc derivatives may be attached *via* a linker, *e.g.*, a bivalent or trivalent
25 branched linker. In some embodiments the GalNAc conjugate is conjugated to the 3' end of the sense strand. In some embodiments, the GalNAc conjugate is conjugated to the iRNA agent (*e.g.*, to the 3' end of the sense strand) *via* a linker, *e.g.*, a linker as described herein. In some embodiments the GalNAc conjugate is conjugated to the 5' end of the sense strand. In some embodiments, the GalNAc conjugate is conjugated to the iRNA agent (*e.g.*, to the 5' end of the sense strand) *via* a linker, *e.g.*, a
30 linker as described herein.


In certain embodiments of the invention, the GalNAc or GalNAc derivative is attached to an iRNA agent of the invention *via* a monovalent linker. In some embodiments, the GalNAc or GalNAc derivative is attached to an iRNA agent of the invention *via* a bivalent linker. In yet other
embodiments of the invention, the GalNAc or GalNAc derivative is attached to an iRNA agent of the
35 invention *via* a trivalent linker. In other embodiments of the invention, the GalNAc or GalNAc derivative is attached to an iRNA agent of the invention *via* a tetravalent linker.

In certain embodiments, the double stranded RNAi agents of the invention comprise one GalNAc or GalNAc derivative attached to the iRNA agent. In certain embodiments, the double stranded RNAi agents of the invention comprise a plurality (*e.g.*, 2, 3, 4, 5, or 6) GalNAc or GalNAc derivatives, each independently attached to a plurality of nucleotides of the double stranded RNAi agent through a plurality of monovalent linkers.


In some embodiments, for example, when the two strands of an iRNA agent of the invention are part of one larger molecule connected by an uninterrupted chain of nucleotides between the 3'-end of one strand and the 5'-end of the respective other strand forming a hairpin loop comprising, a plurality of unpaired nucleotides, each unpaired nucleotide within the hairpin loop may independently comprise a GalNAc or GalNAc derivative attached *via* a monovalent linker. The hairpin loop may also be formed by an extended overhang in one strand of the duplex.

In some embodiments, for example, when the two strands of an iRNA agent of the invention are part of one larger molecule connected by an uninterrupted chain of nucleotides between the 3'-end of one strand and the 5'-end of the respective other strand forming a hairpin loop comprising, a plurality of unpaired nucleotides, each unpaired nucleotide within the hairpin loop may independently comprise a GalNAc or GalNAc derivative attached *via* a monovalent linker. The hairpin loop may also be formed by an extended overhang in one strand of the duplex.


In one embodiment, a carbohydrate conjugate for use in the compositions and methods of the invention is selected from the group consisting of:




Formula II,




Formula III,




Formula IV,




Formula V,




Formula VI,




Formula VII,




Formula VIII,




Formula IX,




Formula X,




Formula XI,




Formula XII,




Formula XIII,




Formula XIV,




Formula XV,



Formula XVI,




Formula XVII,




Formula XVIII,


5




Formula XIX,




Formula XX,




Formula XXI,




Formula XXII,




Formula XXIII;




, wherein Y is O or S and n is 3 -6 (Formula XXIV);




, wherein Y is O or S and n is 3-6 (Formula XXV);




Formula XXVI;




, wherein X is O or S (Formula XXVII);




Formula XXVII; Formula XXIX;




Formula XXXI;



Formula XXX;




, and




Formula XXXII;

5 Formula XXXIII.




Formula XXXIV.

In another embodiment, a carbohydrate conjugate for use in the compositions and methods of the invention is a monosaccharide. In one embodiment, the monosaccharide is an N-acetylgalactosamine, such as




Formula II.


In some embodiments, the RNAi agent is attached to the carbohydrate conjugate *via* a linker as shown in the following schematic, wherein X is O or S.



In some embodiments, the RNAi agent is conjugated to L96 as defined in Table 1 and shown below:



5 Another representative carbohydrate conjugate for use in the embodiments described herein includes, but is not limited to,




(Formula XXXVI), when one of X or Y is an oligonucleotide, the other is a hydrogen.

10 In some embodiments, a suitable ligand is a ligand disclosed in WO 2019/055633, the entire contents of which are incorporated herein by reference. In one embodiment the ligand comprises the structure below:

5

10



In certain embodiments of the invention, the GalNAc or GalNAc derivative is attached to an iRNA agent of the invention *via* a monovalent linker. In some embodiments, the GalNAc or GalNAc derivative is attached to an iRNA agent of the invention *via* a bivalent linker. In yet other

15 embodiments of the invention, the GalNAc or GalNAc derivative is attached to an iRNA agent of the invention *via* a trivalent linker.

In one embodiment, the double stranded RNAi agents of the invention comprise one or more GalNAc or GalNAc derivative attached to the iRNA agent. The GalNAc may be attached to any nucleotide *via* a linker on the sense strand or antisense strand. The GalNAc may be attached to the

20 5'-end of the sense strand, the 3' end of the sense strand, the 5'-end of the antisense strand, or the 3' – end of the antisense strand. In one embodiment, the GalNAc is attached to the 3' end of the sense strand, *e.g.*, *via* a trivalent linker.

In other embodiments, the double stranded RNAi agents of the invention comprise a plurality (*e.g.*, 2, 3, 4, 5, or 6) GalNAc or GalNAc derivatives, each independently attached to a plurality of

25 nucleotides of the double stranded RNAi agent through a plurality of linkers, *e.g.*, monovalent linkers.

In some embodiments, for example, when the two strands of an iRNA agent of the invention is part of one larger molecule connected by an uninterrupted chain of nucleotides between the 3'-end of one strand and the 5'-end of the respective other strand forming a hairpin loop comprising, a plurality of unpaired nucleotides, each unpaired nucleotide within the hairpin loop may independently

30 comprise a GalNAc or GalNAc derivative attached *via* a monovalent linker.

In some embodiments, the carbohydrate conjugate further comprises one or more additional ligands as described above, such as, but not limited to, a PK modulator or a cell permeation peptide.

Additional carbohydrate conjugates and linkers suitable for use in the present invention include those described in PCT Publication Nos. WO 2014/179620 and WO 2014/179627, the entire

35 contents of each of which are incorporated herein by reference.

D. Linkers

In some embodiments, the conjugate or ligand described herein can be attached to an iRNA oligonucleotide with various linkers that can be cleavable or non-cleavable.

The term "linker" or "linking group" means an organic moiety that connects two parts of a compound, *e.g.*, covalently attaches two parts of a compound. Linkers typically comprise a direct bond or an atom such as oxygen or sulfur, a unit such as NR₈, C(O), C(O)NH, SO, SO₂, SO₂NH or a chain of atoms, such as, but not limited to, substituted or unsubstituted alkyl, substituted or unsubstituted alkenyl, substituted or unsubstituted alkynyl, arylalkyl, arylalkenyl, arylalkynyl, heteroarylalkyl, heteroarylalkenyl, heteroarylalkynyl, heterocyclalkyl, heterocyclalkenyl, heterocyclalkynyl, aryl, heteroaryl, heterocycl, cycloalkyl, cycloalkenyl, alkylarylalkyl, alkylarylalkenyl, alkylarylalkynyl, alkenylarylalkyl, alkenylarylalkenyl, alkenylarylalkynyl, alkynylarylalkyl, alkynylarylalkenyl, alkynylarylalkynyl, alkylheteroarylalkyl, alkylheteroarylalkenyl, alkylheteroarylalkynyl, alkenylheteroarylalkyl, alkenylheteroarylalkenyl, alkenylheteroarylalkynyl, alkynylheteroarylalkyl, alkynylheteroarylalkenyl, alkynylheteroarylalkynyl, alkylheterocyclalkyl, alkylheterocyclalkenyl, alkylheterocyclalkynyl, alkenylheterocyclalkyl, alkenylheterocyclalkenyl, alkenylheterocyclalkynyl, alkynylheterocyclalkyl, alkynylheterocyclalkenyl, alkynylheterocyclalkynyl, alkylaryl, alkenylaryl, alkynylaryl, alkylheteroaryl, alkenylheteroaryl, alkynylheteroaryl, which one or more methylenes can be interrupted or terminated by O, S, S(O), SO₂, N(R₈), C(O), substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, or substituted or unsubstituted heterocyclic; where R₈ is hydrogen, acyl, aliphatic, or substituted aliphatic. In one embodiment, the linker is about 1-24 atoms, 2-24, 3-24, 4-24, 5-24, 6-24, 6-18, 7-18, 8-18, 7-17, 8-17, 6-16, 7-17, or 8-16 atoms.

A cleavable linking group is one which is sufficiently stable outside the cell, but which upon entry into a target cell is cleaved to release the two parts the linker is holding together. In an exemplary embodiment, the cleavable linking group is cleaved at least about 10 times, 20, times, 30 times, 40 times, 50 times, 60 times, 70 times, 80 times, 90 times, or more, or at least 100 times faster in a target cell or under a first reference condition (which can, *e.g.*, be selected to mimic or represent intracellular conditions) than in the blood of a subject, or under a second reference condition (which can, *e.g.*, be selected to mimic or represent conditions found in the blood or serum).

Cleavable linking groups are susceptible to cleavage agents, *e.g.*, pH, redox potential, or the presence of degradative molecules. Generally, cleavage agents are more prevalent or found at higher levels or activities inside cells than in serum or blood. Examples of such degradative agents include: redox agents which are selected for particular substrates or which have no substrate specificity, including, *e.g.*, oxidative or reductive enzymes or reductive agents such as mercaptans, present in cells, that can degrade a redox cleavable linking group by reduction; esterases; endosomes or agents that can create an acidic environment, *e.g.*, those that result in a pH of five or lower; enzymes that can

hydrolyze or degrade an acid cleavable linking group by acting as a general acid, peptidases (which can be substrate specific), and phosphatases.

A cleavable linkage group, such as a disulfide bond can be susceptible to pH. The pH of human serum is 7.4, while the average intracellular pH is slightly lower, ranging from about 7.1-7.3. Endosomes have a more acidic pH, in the range of 5.5-6.0, and lysosomes have an even more acidic pH at around 5.0. Some linkers will have a cleavable linking group that is cleaved at a selected pH, thereby releasing a cationic lipid from the ligand inside the cell, or into the desired compartment of the cell.

A linker can include a cleavable linking group that is cleavable by a particular enzyme. The type of cleavable linking group incorporated into a linker can depend on the cell to be targeted. For example, a liver-targeting ligand can be linked to a cationic lipid through a linker that includes an ester group. Liver cells are rich in esterases, and therefore the linker will be cleaved more efficiently in liver cells than in cell types that are not esterase-rich. Other cell-types rich in esterases include cells of the lung, renal cortex, and testis.

Linkers that contain peptide bonds can be used when targeting cell types rich in peptidases, such as liver cells and synoviocytes.

In general, the suitability of a candidate cleavable linking group can be evaluated by testing the ability of a degradative agent (or condition) to cleave the candidate linking group. It will also be desirable to also test the candidate cleavable linking group for the ability to resist cleavage in the blood or when in contact with other non-target tissue. Thus, one can determine the relative susceptibility to cleavage between a first and a second condition, where the first is selected to be indicative of cleavage in a target cell and the second is selected to be indicative of cleavage in other tissues or biological fluids, *e.g.*, blood or serum. The evaluations can be carried out in cell free systems, in cells, in cell culture, in organ or tissue culture, or in whole animals. It can be useful to make initial evaluations in cell-free or culture conditions and to confirm by further evaluations in whole animals. In certain embodiments, useful candidate compounds are cleaved at least about 2, 4, 10, 20, 30, 40, 50, 60, 70, 80, 90, or 100 times faster in the cell (or under *in vitro* conditions selected to mimic intracellular conditions) as compared to blood or serum (or under *in vitro* conditions selected to mimic extracellular conditions).

i. Redox cleavable linking groups

In certain embodiments, a cleavable linking group is a redox cleavable linking group that is cleaved upon reduction or oxidation. An example of reductively cleavable linking group is a disulphide linking group (-S-S-). To determine if a candidate cleavable linking group is a suitable “reductively cleavable linking group,” or for example is suitable for use with a particular iRNA moiety and particular targeting agent one can look to methods described herein. For example, a candidate can be evaluated by incubation with dithiothreitol (DTT), or other reducing agent using

reagents known in the art, which mimic the rate of cleavage which would be observed in a cell, *e.g.*, a target cell. The candidates can also be evaluated under conditions which are selected to mimic blood or serum conditions. In one, candidate compounds are cleaved by at most about 10% in the blood. In other embodiments, useful candidate compounds are degraded at least about 2, 4, 10, 20, 30, 40, 50, 60, 70, 80, 90, or about 100 times faster in the cell (or under *in vitro* conditions selected to mimic intracellular conditions) as compared to blood (or under *in vitro* conditions selected to mimic extracellular conditions). The rate of cleavage of candidate compounds can be determined using standard enzyme kinetics assays under conditions chosen to mimic intracellular media and compared to conditions chosen to mimic extracellular media.

ii. Phosphate-based cleavable linking groups

In other embodiments, a cleavable linker comprises a phosphate-based cleavable linking group. A phosphate-based cleavable linking group is cleaved by agents that degrade or hydrolyze the phosphate group. An example of an agent that cleaves phosphate groups in cells are enzymes such as phosphatases in cells. Examples of phosphate-based linking groups are -O-P(O)(ORk)-O-, -O-P(S)(ORk)-O-, -O-P(S)(SRk)-O-, -S-P(O)(ORk)-O-, -O-P(O)(ORk)-S-, -S-P(O)(ORk)-S-, -O-P(S)(ORk)-S-, -S-P(S)(ORk)-O-, -O-P(O)(Rk)-O-, -O-P(S)(Rk)-O-, -S-P(O)(Rk)-O-, -S-P(S)(Rk)-O-, -S-P(O)(Rk)-S-, -O-P(S)(Rk)-S-, wherein Rk at each occurrence can be, independently, C1-C20 alkyl, C1-C20 haloalkyl, C6-C10 aryl, or C7-C12 aralkyl. Exemplary embodiments include -O-P(O)(OH)-O-, -O-P(S)(OH)-O-, -O-P(S)(SH)-O-, -S-P(O)(OH)-O-, -O-P(O)(OH)-S-, -S-P(O)(OH)-S-, -O-P(S)(OH)-S-, -S-P(S)(OH)-O-, -O-P(O)(H)-O-, -O-P(S)(H)-O-, -S-P(O)(H)-O-, -S-P(S)(H)-O-, -S-P(O)(H)-S-, and -O-P(S)(H)-S-. In certain embodiments a phosphate-based linking group is -O-P(O)(OH)-O-. These candidates can be evaluated using methods analogous to those described above.

iii. Acid cleavable linking groups

In other embodiments, a cleavable linker comprises an acid cleavable linking group. An acid cleavable linking group is a linking group that is cleaved under acidic conditions. In certain embodiments acid cleavable linking groups are cleaved in an acidic environment with a pH of about 6.5 or lower (*e.g.*, about 6.0, 5.5, 5.0, or lower), or by agents such as enzymes that can act as a general acid. In a cell, specific low pH organelles, such as endosomes and lysosomes can provide a cleaving environment for acid cleavable linking groups. Examples of acid cleavable linking groups include but are not limited to hydrazones, esters, and esters of amino acids. Acid cleavable groups can have the general formula -C=NN-, C(O)O, or -OC(O). An exemplary embodiment is when the carbon attached to the oxygen of the ester (the alkoxy group) is an aryl group, substituted alkyl group, or tertiary alkyl group such as dimethyl pentyl or t-butyl. These candidates can be evaluated using methods analogous to those described above.


iv. *Ester-based linking groups*

In other embodiments, a cleavable linker comprises an ester-based cleavable linking group. An ester-based cleavable linking group is cleaved by enzymes such as esterases and amidases in cells. Examples of ester-based cleavable linking groups include, but are not limited to, esters of alkylene, alkenylene and alkynylene groups. Ester cleavable linking groups have the general formula -C(O)O-, or -OC(O)-. These candidates can be evaluated using methods analogous to those described above.

v. *Peptide-based cleaving groups*


In yet other embodiments, a cleavable linker comprises a peptide-based cleavable linking group. A peptide-based cleavable linking group is cleaved by enzymes such as peptidases and proteases in cells. Peptide-based cleavable linking groups are peptide bonds formed between amino acids to yield oligopeptides (*e.g.*, dipeptides, tripeptides *etc.*) and polypeptides. Peptide-based cleavable groups do not include the amide group (-C(O)NH-). The amide group can be formed between any alkylene, alkenylene or alkynylene. A peptide bond is a special type of amide bond formed between amino acids to yield peptides and proteins. The peptide based cleavage group is generally limited to the peptide bond (*i.e.*, the amide bond) formed between amino acids yielding peptides and proteins and does not include the entire amide functional group. Peptide-based cleavable linking groups have the general formula -NHCHRAC(O)NHCHRBC(O)-, where RA and RB are the R groups of the two adjacent amino acids. These candidates can be evaluated using methods analogous to those described above.

In some embodiments, an iRNA of the invention is conjugated to a carbohydrate through a linker. Non-limiting examples of iRNA carbohydrate conjugates with linkers of the compositions and methods of the invention include, but are not limited to,




(Formula XXXVII),


25




(Formula XXXVIII),



(Formula XXXIX),




(Formula XL),




5


(Formula XLI),



(Formula XLII),



(Formula XLIII), and




(Formula XLIV), when one of X or Y is an oligonucleotide, the other is a hydrogen.



- 5 In certain embodiments of the compositions and methods of the invention, a ligand is one or more “GalNAc” (N-acetylgalactosamine) derivatives attached through a bivalent or trivalent branched linker.

In one embodiment, a dsRNA of the invention is conjugated to a bivalent or trivalent branched linker selected from the group of structures shown in any of formula (XLV) – (XLVIII):


10 Formula XLV



Formula XLVI



, or



;


Formula XLVII

Formula XLVIII


wherein:

- 15 q^2A , q^2B , q^3A , q^3B , q^4A , q^4B , q^5A , q^5B and q^5C represent independently for each occurrence 0-20 and wherein the repeating unit can be the same or different;

- $P^{2A}, P^{2B}, P^{3A}, P^{3B}, P^{4A}, P^{4B}, P^{5A}, P^{5B}, P^{5C}, T^{2A}, T^{2B}, T^{3A}, T^{3B}, T^{4A}, T^{4B}, T^{5A}, T^{5B}, T^{5C}$ are each independently for each occurrence absent, CO, NH, O, S, OC(O), NHC(O), CH₂, CH₂NH or CH₂O; $Q^{2A}, Q^{2B}, Q^{3A}, Q^{3B}, Q^{4A}, Q^{4B}, Q^{5A}, Q^{5B}, Q^{5C}$ are independently for each occurrence absent, alkylene, substituted alkylene wherein one or more methylenes can be interrupted or terminated by one or more of O, S, S(O), SO₂, N(R^N), C(R')=C(R''), C≡C or C(O);
- 5 $R^{2A}, R^{2B}, R^{3A}, R^{3B}, R^{4A}, R^{4B}, R^{5A}, R^{5B}, R^{5C}$ are each independently for each occurrence absent, NH, O,



- $L^{2A}, L^{2B}, L^{3A}, L^{3B}, L^{4A}, L^{4B}, L^{5A}, L^{5B}$ and L^{5C} represent the ligand; *i.e.* each independently for each occurrence a monosaccharide (such as GalNAc), disaccharide, trisaccharide, tetrasaccharide, oligosaccharide, or polysaccharide; and R^a is H or amino acid side chain. Trivalent conjugating GalNAc derivatives are particularly useful for use with RNAi agents for inhibiting the expression of a target gene, such as those of formula (XLIX):



- 15 wherein L^{5A}, L^{5B} and L^{5C} represent a monosaccharide, such as GalNAc derivative.

Examples of suitable bivalent and trivalent branched linker groups conjugating GalNAc derivatives include, but are not limited to, the structures recited above as formulas II, VII, XI, X, and XIII.

- 20 Representative U.S. Patents that teach the preparation of RNA conjugates include, but are not limited to, U.S. Patent Nos. 4,828,979; 4,948,882; 5,218,105; 5,525,465; 5,541,313; 5,545,730; 5,552,538; 5,578,717; 5,580,731; 5,591,584; 5,109,124; 5,118,802; 5,138,045; 5,414,077; 5,486,603; 5,512,439; 5,578,718; 5,608,046; 4,587,044; 4,605,735; 4,667,025; 4,762,779; 4,789,737; 4,824,941; 4,835,263; 4,876,335; 4,904,582; 4,958,013; 5,082,830; 5,112,963; 5,214,136; 5,082,830; 5,112,963;
- 25 5,214,136; 5,245,022; 5,254,469; 5,258,506; 5,262,536; 5,272,250; 5,292,873; 5,317,098; 5,371,241; 5,391,723; 5,416,203; 5,451,463; 5,510,475; 5,512,667; 5,514,785; 5,565,552; 5,567,810; 5,574,142; 5,585,481; 5,587,371; 5,595,726; 5,597,696; 5,599,923; 5,599,928; 5,688,941; 6,294,664; 6,320,017; 6,576,752; 6,783,931; 6,900,297; 7,037,646; and 8,106,022, the entire contents of each of which are hereby incorporated herein by reference.

It is not necessary for all positions in a given compound to be uniformly modified, and in fact more than one of the aforementioned modifications can be incorporated in a single compound or even at a single nucleoside within an iRNA. The present invention also includes iRNA compounds that are chimeric compounds.

5 “Chimeric” iRNA compounds or “chimeras,” in the context of this invention, are iRNA compounds, such as, dsRNAi agents, that contain two or more chemically distinct regions, each made up of at least one monomer unit, *i.e.*, a nucleotide in the case of a dsRNA compound. These iRNAs typically contain at least one region wherein the RNA is modified so as to confer upon the iRNA increased resistance to nuclease degradation, increased cellular uptake, or increased binding affinity
10 for the target nucleic acid. An additional region of the iRNA can serve as a substrate for enzymes capable of cleaving RNA:DNA or RNA:RNA hybrids. By way of example, RNase H is a cellular endonuclease which cleaves the RNA strand of an RNA:DNA duplex. Activation of RNase H, therefore, results in cleavage of the RNA target, thereby greatly enhancing the efficiency of iRNA inhibition of gene expression. Consequently, comparable results can often be obtained with shorter
15 iRNAs when chimeric dsRNAs are used, compared to phosphorothioate deoxy dsRNAs hybridizing to the same target region. Cleavage of the RNA target can be routinely detected by gel electrophoresis and, if necessary, associated nucleic acid hybridization techniques known in the art.

 In certain instances, the RNA of an iRNA can be modified by a non-ligand group. A number of non-ligand molecules have been conjugated to iRNAs in order to enhance the activity, cellular
20 distribution or cellular uptake of the iRNA, and procedures for performing such conjugations are available in the scientific literature. Such non-ligand moieties have included lipid moieties, such as cholesterol (Kubo, T. *et al.*, *Biochem. Biophys. Res. Comm.*, 2007, 365(1):54-61; Letsinger *et al.*, *Proc. Natl. Acad. Sci. USA*, 1989, 86:6553), cholic acid (Manoharan *et al.*, *Bioorg. Med. Chem. Lett.*, 1994, 4:1053), a thioether, *e.g.*, hexyl-S-tritylthiol (Manoharan *et al.*, *Ann. N.Y. Acad. Sci.*, 1992, 660:306; Manoharan *et al.*, *Bioorg. Med. Chem. Lett.*, 1993, 3:2765), a thiocholesterol (Oberhauser *et al.*, *Nucl. Acids Res.*, 1992, 20:533), an aliphatic chain, *e.g.*, dodecandiol or undecyl residues (Saison-
25 Behmoaras *et al.*, *EMBO J.*, 1991, 10:111; Kabanov *et al.*, *FEBS Lett.*, 1990, 259:327; Svinarchuk *et al.*, *Biochimie*, 1993, 75:49), a phospholipid, *e.g.*, di-hexadecyl-rac-glycerol or triethylammonium 1,2-di-O-hexadecyl-rac-glycero-3-H-phosphonate (Manoharan *et al.*, *Tetrahedron Lett.*, 1995, 36:3651; Shea *et al.*, *Nucl. Acids Res.*, 1990, 18:3777), a polyamine or a polyethylene glycol chain (Manoharan
30 *et al.*, *Nucleosides & Nucleotides*, 1995, 14:969), or adamantane acetic acid (Manoharan *et al.*, *Tetrahedron Lett.*, 1995, 36:3651), a palmityl moiety (Mishra *et al.*, *Biochim. Biophys. Acta*, 1995, 1264:229), or an octadecylamine or hexylamino-carbonyl-oxcholesterol moiety (Crooke *et al.*, *J. Pharmacol. Exp. Ther.*, 1996, 277:923). Representative United States patents that teach the
35 preparation of such RNA conjugates have been listed above. Typical conjugation protocols involve the synthesis of RNAs bearing an aminolinker at one or more positions of the sequence. The amino group is then reacted with the molecule being conjugated using appropriate coupling or activating

reagents. The conjugation reaction can be performed either with the RNA still bound to the solid support or following cleavage of the RNA, in solution phase. Purification of the RNA conjugate by HPLC typically affords the pure conjugate.

5 IV. Delivery of an iRNA of the Invention

The delivery of an iRNA of the invention to a cell *e.g.*, a cell within a subject, such as a human subject (*e.g.*, a subject in need thereof, such as a subject susceptible to or diagnosed with an F12-associated disorder, *e.g.*, hereditary angioedema (HAE) (*e.g.*, hereditary angioedema type I; hereditary angioedema type II; hereditary angioedema type III; or any other hereditary angioedema caused by elevated levels of bradykinin); prekallikrein deficiency, hypertension, *e.g.*, malignant essential hypertension, end stage renal disease, Fletcher Factor Deficiency, thromboembolic disease, inflammatory disease, or Alzheimer's Disease) can be achieved in a number of different ways. For example, delivery may be performed by contacting a cell with an iRNA of the invention either *in vitro* or *in vivo*. *In vivo* delivery may also be performed directly by administering a composition
10 comprising an iRNA, *e.g.*, a dsRNA, to a subject. Alternatively, *in vivo* delivery may be performed indirectly by administering one or more vectors that encode and direct the expression of the iRNA. These alternatives are discussed further below.

In general, any method of delivering a nucleic acid molecule (*in vitro* or *in vivo*) can be adapted for use with an iRNA of the invention (see *e.g.*, Akhtar S. and Julian RL. (1992) *Trends Cell Biol.* 2(5):139-144 and WO94/02595, which are incorporated herein by reference in their entireties).
20 For *in vivo* delivery, factors to consider in order to deliver an iRNA molecule include, for example, biological stability of the delivered molecule, prevention of non-specific effects, and accumulation of the delivered molecule in the target tissue. RNA interference has also shown success with local delivery to the CNS by direct injection (Dorn, G., *et al.* (2004) *Nucleic Acids* 32:e49; Tan, PH., *et al.* (2005) *Gene Ther.* 12:59-66; Makimura, H., *et al.* (2002) *BMC Neurosci.* 3:18; Shishkina, GT., *et al.* (2004) *Neuroscience* 129:521-528; Thakker, ER., *et al.* (2004) *Proc. Natl. Acad. Sci. U.S.A.* 101:17270-17275; Akaneya, Y., *et al.* (2005) *J. Neurophysiol.* 93:594-602). Modification of the RNA or the pharmaceutical carrier can also permit targeting of the iRNA to the target tissue and avoid undesirable off-target effects. iRNA molecules can be modified by chemical conjugation to lipophilic
25 groups such as cholesterol to enhance cellular uptake and prevent degradation. For example, an iRNA directed against ApoB conjugated to a lipophilic cholesterol moiety was injected systemically into mice and resulted in knockdown of apoB mRNA in both the liver and jejunum (Soutschek, J., *et al.* (2004) *Nature* 432:173-178).

In an alternative embodiment, the iRNA can be delivered using drug delivery systems such as
35 a nanoparticle, a dendrimer, a polymer, liposomes, or a cationic delivery system. Positively charged cationic delivery systems facilitate binding of an iRNA molecule (negatively charged) and also enhance interactions at the negatively charged cell membrane to permit efficient uptake of an iRNA

by the cell. Cationic lipids, dendrimers, or polymers can either be bound to an iRNA, or induced to form a vesicle or micelle (see *e.g.*, Kim SH, *et al* (2008) *Journal of Controlled Release* 129(2):107-116) that encases an iRNA. The formation of vesicles or micelles further prevents degradation of the iRNA when administered systemically. Methods for making and administering cationic- iRNA complexes are well within the abilities of one skilled in the art (see *e.g.*, Sorensen, DR, *et al* (2003) *J. Mol. Biol* 327:761-766; Verma, UN, *et al* (2003) *Clin. Cancer Res.* 9:1291-1300; Arnold, AS *et al* (2007) *J. Hypertens.* 25:197-205, which are incorporated herein by reference in their entirety). Some non-limiting examples of drug delivery systems useful for systemic delivery of iRNAs include DOTAP (Sorensen, DR., *et al* (2003), *supra*; Verma, UN, *et al* (2003), *supra*), "solid nucleic acid lipid particles" (Zimmermann, TS, *et al* (2006) *Nature* 441:111-114), cardiolipin (Chien, PY, *et al* (2005) *Cancer Gene Ther.* 12:321-328; Pal, A, *et al* (2005) *Int J. Oncol.* 26:1087-1091), polyethyleneimine (Bonnet ME, *et al* (2008) *Pharm. Res.* Aug 16 Epub ahead of print; Aigner, A. (2006) *J. Biomed. Biotechnol.* 71659), Arg-Gly-Asp (RGD) peptides (Liu, S. (2006) *Mol. Pharm.* 3:472-487), and polyamidoamines (Tomalia, DA, *et al* (2007) *Biochem. Soc. Trans.* 35:61-67; Yoo, H., *et al* (1999) *Pharm. Res.* 16:1799-1804). In some embodiments, an iRNA forms a complex with cyclodextrin for systemic administration. Methods for administration and pharmaceutical compositions of iRNAs and cyclodextrins can be found in U.S. Patent No. 7,427,605, which is herein incorporated by reference in its entirety. Certain aspects of the instant disclosure relate to a method of reducing the expression of an F12 gene in a cell, comprising contacting said cell with the double-stranded RNAi agent of the disclosure. In one embodiment, the cell is a hepatic cell, optionally a hepatocyte. In one embodiment, the cell is an extrahepatic cell.

A. Vector encoded iRNAs of the Invention

iRNA targeting the F12 gene can be expressed from transcription units inserted into DNA or RNA vectors (see, *e.g.*, Couture, A, *et al.*, *TIG.* (1996), 12:5-10; Skillern, A, *et al.*, International PCT Publication No. WO 00/22113, Conrad, International PCT Publication No. WO 00/22114, and Conrad, U.S. Patent No. 6,054,299). Expression can be transient (on the order of hours to weeks) or sustained (weeks to months or longer), depending upon the specific construct used and the target tissue or cell type. These transgenes can be introduced as a linear construct, a circular plasmid, or a viral vector, which can be an integrating or non-integrating vector. The transgene can also be constructed to permit it to be inherited as an extrachromosomal plasmid (Gassmann, *et al.*, *Proc. Natl. Acad. Sci. USA* (1995) 92:1292).

Viral vector systems which can be utilized with the methods and compositions described herein include, but are not limited to, (a) adenovirus vectors; (b) retrovirus vectors, including but not limited to lentiviral vectors, moloney murine leukemia virus, *etc.*; (c) adeno- associated virus vectors; (d) herpes simplex virus vectors; (e) SV 40 vectors; (f) polyoma virus vectors; (g) papilloma virus vectors; (h) picornavirus vectors; (i) pox virus vectors such as an orthopox, *e.g.*, vaccinia virus vectors

or avipox, *e.g.* canary pox or fowl pox; and (j) a helper-dependent or gutless adenovirus. Replication-defective viruses can also be advantageous. Different vectors will or will not become incorporated into the cells' genome. The constructs can include viral sequences for transfection, if desired.

Alternatively, the construct can be incorporated into vectors capable of episomal replication, *e.g.* EPV and EBV vectors. Constructs for the recombinant expression of an iRNA will generally require regulatory elements, *e.g.*, promoters, enhancers, *etc.*, to ensure the expression of the iRNA in target cells. Other aspects to consider for vectors and constructs are known in the art.

V. Pharmaceutical Compositions of the Invention

The present invention also includes pharmaceutical compositions and formulations which include the iRNAs of the invention. In one embodiment, provided herein are pharmaceutical compositions containing an iRNA, as described herein, and a pharmaceutically acceptable carrier. The pharmaceutical compositions containing the iRNA are useful for preventing or treating an F12-associated disorder, *e.g.*, heredity angioedema (HAE) (*e.g.*, hereditary angioedema type I; hereditary angioedema type II; hereditary angioedema type III; or any other hereditary angioedema caused by elevated levels of bradykinin); prekallikrein deficiency, hypertension, *e.g.*, malignant essential hypertension, end stage renal disease, Fletcher Factor Deficiency, thromboembolic disease, inflammatory disease, or Alzheimer's Disease.

Such pharmaceutical compositions are formulated based on the mode of delivery. One example is compositions that are formulated for systemic administration *via* parenteral delivery, *e.g.*, by subcutaneous (SC), intramuscular (IM), or intravenous (IV) delivery. The pharmaceutical compositions of the invention may be administered in dosages sufficient to inhibit expression of an F12 gene.

In some embodiments, the pharmaceutical compositions of the invention are sterile. In another embodiment, the pharmaceutical compositions of the invention are pyrogen free.

The pharmaceutical compositions of the invention may be administered in dosages sufficient to inhibit expression of an F12 gene. In general, a suitable dose of an iRNA of the invention will be in the range of about 0.001 to about 200.0 milligrams per kilogram body weight of the recipient per day, generally in the range of about 1 to 50 mg per kilogram body weight per day. Typically, a suitable dose of an iRNA of the invention will be in the range of about 0.1 mg/kg to about 5.0 mg/kg, such as, about 0.3 mg/kg and about 3.0 mg/kg. A repeat-dose regimen may include administration of a therapeutic amount of iRNA on a regular basis, such as every month, once every 3-6 months, or once a year. In certain embodiments, the iRNA is administered about once per month to about once per six months.

After an initial treatment regimen, the treatments can be administered on a less frequent basis. Duration of treatment can be determined based on the severity of disease.

In other embodiments, a single dose of the pharmaceutical compositions can be long lasting, such that doses are administered at not more than 1, 2, 3, or 4 month intervals. In some embodiments of the invention, a single dose of the pharmaceutical compositions of the invention is administered about once per month. In other embodiments of the invention, a single dose of the pharmaceutical compositions of the invention is administered quarterly (*i.e.*, about every three months). In other
5 embodiments of the invention, a single dose of the pharmaceutical compositions of the invention is administered twice per year (*i.e.*, about once every six months).

The skilled artisan will appreciate that certain factors can influence the dosage and timing required to effectively treat a subject, including but not limited to mutations present in the subject,
10 previous treatments, the general health or age of the subject, and other diseases present. Moreover, treatment of a subject with a prophylactically or therapeutically effective amount, as appropriate, of a composition can include a single treatment or a series of treatments.

The pharmaceutical compositions of the present disclosure can be administered in a number of ways depending upon whether local or systemic treatment is desired and upon the area to be
15 treated. Administration may be topical (including ophthalmic, vaginal, rectal, intranasal, transdermal), oral, or parenteral. Parenteral administration includes intravenous, intraarterial, subcutaneous, intraperitoneal or intramuscular injection or infusion; subdermal, *e.g.*, via an implanted device; or intracranial, *e.g.*, by intraparenchymal, intrathecal or intraventricular, administration.

The iRNA can be delivered in a manner to target a particular tissue, such as the liver.

Pharmaceutical compositions and formulations for topical administration can include
20 transdermal patches, ointments, lotions, creams, gels, drops, suppositories, sprays, liquids and powders. Conventional pharmaceutical carriers, aqueous, powder or oily bases, thickeners and the like can be necessary or desirable. Coated condoms, gloves and the like can also be useful. Suitable topical formulations include those in which the RNAi agents featured in the disclosure are in admixture with
25 a topical delivery agent such as lipids, liposomes, fatty acids, fatty acid esters, steroids, chelating agents and surfactants. Suitable lipids and liposomes include neutral (*e.g.*, dioleoylphosphatidyl DOPE ethanolamine, dimyristoylphosphatidyl choline DMPC, distearoylphosphatidyl choline) negative (*e.g.*, dimyristoylphosphatidyl glycerol DMPG) and cationic (*e.g.*,
dioleoyltetramethylaminopropyl DOTAP and dioleoylphosphatidyl ethanolamine DOTMA). RNAi
30 agents featured in the disclosure can be encapsulated within liposomes or can form complexes thereto, in particular to cationic liposomes. Alternatively, RNAi agents can be complexed to lipids, in particular to cationic lipids. Suitable fatty acids and esters include but are not limited to arachidonic acid, oleic acid, eicosanoic acid, lauric acid, caprylic acid, capric acid, myristic acid, palmitic acid, stearic acid, linoleic acid, linolenic acid, dicaprinate, tricaprinate, monoolein, dilaurin, glyceryl 1-
35 monocaprinate, 1-dodecylazacycloheptan-2-one, an acylcarnitine, an acylcholine, or a C1-20 alkyl ester (*e.g.*, isopropylmyristate IPM), monoglyceride, diglyceride or pharmaceutically acceptable salt

thereof. Topical formulations are described in detail in US 6,747,014, which is incorporated herein by reference.

In one embodiment, the siRNAs, double stranded RNA agents of the invention, are administered to a cell in a pharmaceutical composition by a topical route of administration.

- 5 In one embodiment, the pharmaceutical composition may include an siRNA compound mixed with a topical delivery agent. The topical delivery agent can be a plurality of microscopic vesicles. The microscopic vesicles can be liposomes. In some embodiments the liposomes are cationic liposomes.

- In another embodiment, the dsRNA agent is admixed with a topical penetration enhancer. In one embodiment, the topical penetration enhancer is a fatty acid. The fatty acid can be arachidonic acid, oleic acid, lauric acid, caprylic acid, capric acid, myristic acid, palmitic acid, stearic acid, 10 linoleic acid, linolenic acid, dicaprate, tricaprate, monolein, dilaurin, glyceryl 1-monocaprate, 1-dodecylazacycloheptan-2-one, an acylcarnitine, an acylcholine, or a C1-10 alkyl ester, monoglyceride, diglyceride or pharmaceutically acceptable salt thereof.

- In another embodiment, the topical penetration enhancer is a bile salt. The bile salt can be 15 cholic acid, dehydrocholic acid, deoxycholic acid, glucolic acid, glycholic acid, glycodeoxycholic acid, taurocholic acid, taurodeoxycholic acid, chenodeoxycholic acid, ursodeoxycholic acid, sodium tauro-24,25-dihydro-fusidate, sodium glycodihydrofusidate, polyoxyethylene-9-lauryl ether or a pharmaceutically acceptable salt thereof.

- In another embodiment, the penetration enhancer is a chelating agent. The chelating agent 20 can be EDTA, citric acid, a salicylate, a N-acyl derivative of collagen, laureth-9, an N-amino acyl derivative of a beta-diketone or a mixture thereof.

In another embodiment, the penetration enhancer is a surfactant, *e.g.*, an ionic or nonionic surfactant. The surfactant can be sodium lauryl sulfate, polyoxyethylene-9-lauryl ether, polyoxyethylene-20-cetyl ether, a perfluorchemical emulsion or mixture thereof.

- 25 In another embodiment, the penetration enhancer can be selected from a group consisting of unsaturated cyclic ureas, 1-alkyl-alkones, 1-alkenylazacyclo-alkanones, steroidal anti-inflammatory agents and mixtures thereof. In yet another embodiment the penetration enhancer can be a glycol, a pyrrol, an azone, or a terpenes.

- In one aspect, the invention features a pharmaceutical composition including an siRNA 30 compound, *e.g.*, a double-stranded siRNA compound, or ssiRNA compound, (*e.g.*, a precursor, *e.g.*, a larger siRNA compound which can be processed into a ssiRNA compound, or a DNA which encodes an siRNA compound, *e.g.*, a double-stranded siRNA compound, or ssiRNA compound, or precursor thereof) in an injectable dosage form. In one embodiment, the injectable dosage form of the pharmaceutical composition includes sterile aqueous solutions or dispersions and sterile powders. In 35 some embodiments the sterile solution can include a diluent such as water; saline solution; fixed oils, polyethylene glycols, glycerin, or propylene glycol.

The iRNA molecules of the invention can be incorporated into pharmaceutical compositions. Such compositions typically include one or more species of iRNA and a pharmaceutically acceptable carrier. As used herein the language “pharmaceutically acceptable carrier” is intended to include any and all solvents, dispersion media, coatings, antibacterial and antifungal agents, isotonic and
5 absorption delaying agents, and the like, compatible with pharmaceutical administration to a cell, *e.g.*, a liver cell. The use of such media and agents for pharmaceutically active substances is well known in the art. Except insofar as any conventional media or agent is incompatible with the active compound, use thereof in the compositions is contemplated. Supplementary active compounds can also be incorporated into the compositions.

10 Pharmaceutical compositions of the present invention include, but are not limited to, solutions, emulsions, and liposome-containing formulations. These compositions can be generated from a variety of components that include, but are not limited to, preformed liquids, self-emulsifying solids, and self-emulsifying semisolids. Formulations include those that target the liver.

The pharmaceutical formulations of the present invention, which can conveniently be
15 presented in unit dosage form, can be prepared according to conventional techniques well known in the pharmaceutical industry. Such techniques include the step of bringing into association the active ingredients with the pharmaceutical carrier(s) or excipient(s). In general, the formulations are prepared by uniformly and intimately bringing into association the active ingredients with liquid carriers.

20 *A. Additional Formulations*

i. Emulsions

The compositions of the present invention can be prepared and formulated as emulsions. Emulsions are typically heterogeneous systems of one liquid dispersed in another in the form of
25 droplets usually exceeding 0.1 μm in diameter (see *e.g.*, Ansel's Pharmaceutical Dosage Forms and Drug Delivery Systems, Allen, LV., Popovich NG., and Ansel HC., 2004, Lippincott Williams & Wilkins (8th ed.), New York, NY; Idson, in Pharmaceutical Dosage Forms, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 199; Rosoff, in Pharmaceutical Dosage Forms, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc.,
30 New York, N.Y., Volume 1, p. 245; Block in Pharmaceutical Dosage Forms, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 2, p. 335; Higuchi *et al.*, in Remington's Pharmaceutical Sciences, Mack Publishing Co., Easton, Pa., 1985, p. 301). Emulsions are often biphasic systems comprising two immiscible liquid phases intimately mixed and dispersed with each other. In general, emulsions can be of either the water-in-oil (w/o) or the oil-in-water (o/w)
35 variety. When an aqueous phase is finely divided into and dispersed as minute droplets into a bulk oily phase, the resulting composition is called a water-in-oil (w/o) emulsion. Alternatively, when an oily phase is finely divided into and dispersed as minute droplets into a bulk aqueous phase, the

resulting composition is called an oil-in-water (o/w) emulsion. Emulsions can contain additional components in addition to the dispersed phases, and the active drug which can be present as a solution either in the aqueous phase, oily phase or itself as a separate phase. Pharmaceutical excipients such as emulsifiers, stabilizers, dyes, and anti-oxidants can also be present in emulsions as needed.

- 5 Pharmaceutical emulsions can also be multiple emulsions that are comprised of more than two phases such as, for example, in the case of oil-in-water-in-oil (o/w/o) and water-in-oil-in-water (w/o/w) emulsions. Such complex formulations often provide certain advantages that simple binary emulsions do not. Multiple emulsions in which individual oil droplets of an o/w emulsion enclose small water droplets constitute a w/o/w emulsion. Likewise a system of oil droplets enclosed in globules of water
10 stabilized in an oily continuous phase provides an o/w/o emulsion.

- Emulsions are characterized by little or no thermodynamic stability. Often, the dispersed or discontinuous phase of the emulsion is well dispersed into the external or continuous phase and maintained in this form through the means of emulsifiers or the viscosity of the formulation. Other means of stabilizing emulsions entail the use of emulsifiers that can be incorporated into either phase
15 of the emulsion. Emulsifiers can broadly be classified into four categories: synthetic surfactants, naturally occurring emulsifiers, absorption bases, and finely dispersed solids (see *e.g.*, Ansel's Pharmaceutical Dosage Forms and Drug Delivery Systems, Allen, LV., Popovich NG., and Ansel HC., 2004, Lippincott Williams & Wilkins (8th ed.), New York, NY; Idson, in Pharmaceutical Dosage Forms, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y.,
20 volume 1, p. 199).

- Synthetic surfactants, also known as surface active agents, have found wide applicability in the formulation of emulsions and have been reviewed in the literature (see *e.g.*, Ansel's Pharmaceutical Dosage Forms and Drug Delivery Systems, Allen, LV., Popovich NG., and Ansel HC., 2004, Lippincott Williams & Wilkins (8th ed.), New York, NY; Rieger, in Pharmaceutical
25 Dosage Forms, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 285; Idson, in Pharmaceutical Dosage Forms, Lieberman, Rieger and Banker (Eds.), Marcel Dekker, Inc., New York, N.Y., 1988, volume 1, p. 199). Surfactants are typically amphiphilic and comprise a hydrophilic and a hydrophobic portion. The ratio of the hydrophilic to the hydrophobic nature of the surfactant has been termed the hydrophile/lipophile balance (HLB) and is a
30 valuable tool in categorizing and selecting surfactants in the preparation of formulations. Surfactants can be classified into different classes based on the nature of the hydrophilic group: nonionic, anionic, cationic, and amphoteric (see *e.g.*, Ansel's Pharmaceutical Dosage Forms and Drug Delivery Systems, Allen, LV., Popovich NG., and Ansel HC., 2004, Lippincott Williams & Wilkins (8th ed.), New York, NY Rieger, in Pharmaceutical Dosage Forms, Lieberman, Rieger and Banker (Eds.), 1988,
35 Marcel Dekker, Inc., New York, N.Y., volume 1, p. 285).

A large variety of non-emulsifying materials are also included in emulsion formulations and contribute to the properties of emulsions. These include fats, oils, waxes, fatty acids, fatty alcohols,

fatty esters, humectants, hydrophilic colloids, preservatives, and antioxidants (Block, in Pharmaceutical Dosage Forms, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 335; Idson, in Pharmaceutical Dosage Forms, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 199).

5 The application of emulsion formulations *via* dermatological, oral, and parenteral routes, and methods for their manufacture have been reviewed in the literature (see *e.g.*, Ansel's Pharmaceutical Dosage Forms and Drug Delivery Systems, Allen, LV., Popovich NG., and Ansel HC., 2004, Lippincott Williams & Wilkins (8th ed.), New York, NY; Idson, in Pharmaceutical Dosage Forms, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 199).

10

ii. Microemulsions

In one embodiment of the present invention, the compositions of iRNAs and nucleic acids are formulated as microemulsions. A microemulsion can be defined as a system of water, oil, and amphiphile which is a single optically isotropic and thermodynamically stable liquid solution (see
15 *e.g.*, Ansel's Pharmaceutical Dosage Forms and Drug Delivery Systems, Allen, LV., Popovich NG., and Ansel HC., 2004, Lippincott Williams & Wilkins (8th ed.), New York, NY; Rosoff, in Pharmaceutical Dosage Forms, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 245). Typically microemulsions are systems that are prepared by first dispersing an oil in an aqueous surfactant solution and then adding a sufficient amount of a fourth
20 component, generally an intermediate chain-length alcohol to form a transparent system. Therefore, microemulsions have also been described as thermodynamically stable, isotropically clear dispersions of two immiscible liquids that are stabilized by interfacial films of surface-active molecules (Leung and Shah, in: Controlled Release of Drugs: Polymers and Aggregate Systems, Rosoff, M., Ed., 1989, VCH Publishers, New York, pages 185-215).

25

iii. Microparticles

An iRNA of the invention may be incorporated into a particle, *e.g.*, a microparticle. Microparticles can be produced by spray-drying, but may also be produced by other methods including lyophilization, evaporation, fluid bed drying, vacuum drying, or a combination of these
30 techniques.

iv. Penetration Enhancers

In one embodiment, the present invention employs various penetration enhancers to effect the efficient delivery of nucleic acids, particularly iRNAs, to the skin of animals. Most drugs are present
35 in solution in both ionized and nonionized forms. However, usually only lipid soluble or lipophilic drugs readily cross cell membranes. It has been discovered that even non-lipophilic drugs can cross cell membranes if the membrane to be crossed is treated with a penetration enhancer. In addition to

aiding the diffusion of non-lipophilic drugs across cell membranes, penetration enhancers also enhance the permeability of lipophilic drugs.

Penetration enhancers can be classified as belonging to one of five broad categories, *i.e.*, surfactants, fatty acids, bile salts, chelating agents, and non-chelating non-surfactants (see *e.g.*,

- 5 Malmsten, M. Surfactants and polymers in drug delivery, Informa Health Care, New York, NY, 2002; Lee *et al.*, Critical Reviews in Therapeutic Drug Carrier Systems, 1991, p.92). Each of the above mentioned classes of penetration enhancers and their use in manufacture of pharmaceutical compositions and delivery of pharmaceutical agents are well known in the art.

10 *v. Excipients*

- In contrast to a carrier compound, a “pharmaceutical carrier” or “excipient” is a pharmaceutically acceptable solvent, suspending agent, or any other pharmacologically inert vehicle for delivering one or more nucleic acids to an animal. The excipient can be liquid or solid and is selected, with the planned manner of administration in mind, so as to provide for the desired bulk, consistency, etc., when combined with a nucleic acid and the other components of a given pharmaceutical composition. Such agent are well known in the art.
- 15

vi. Other Components

- The compositions of the present invention can additionally contain other adjunct components conventionally found in pharmaceutical compositions, at their art-established usage levels. Thus, for example, the compositions can contain additional, compatible, pharmaceutically-active materials such as, for example, antipruritics, astringents, local anesthetics or anti-inflammatory agents, or can contain additional materials useful in physically formulating various dosage forms of the compositions of the present invention, such as dyes, flavoring agents, preservatives, antioxidants, opacifiers, thickening agents and stabilizers. However, such materials, when added, should not unduly interfere with the biological activities of the components of the compositions of the present invention. The formulations can be sterilized and, if desired, mixed with auxiliary agents, *e.g.*, lubricants, preservatives, stabilizers, wetting agents, emulsifiers, salts for influencing osmotic pressure, buffers, colorings, flavorings, or aromatic substances, and the like which do not deleteriously interact with the nucleic acid(s) of the formulation.
- 20
- 25
- 30

Aqueous suspensions can contain substances which increase the viscosity of the suspension including, for example, sodium carboxymethylcellulose, sorbitol, or dextran. The suspension can also contain stabilizers.

- In some embodiments, pharmaceutical compositions featured in the invention include (a) one or more iRNA and (b) one or more agents which function by a non-iRNA mechanism and which are useful in treating an F12-associated disorder, *e.g.*, heredity angioedema (HAE) (*e.g.*, hereditary angioedema type I; hereditary angioedema type II; hereditary angioedema type III; or any other
- 35

hereditary angioedema caused by elevated levels of bradykinin); prekallikrein deficiency, hypertension, *e.g.*, malignant essential hypertension, end stage renal disease, Fletcher Factor Deficiency, thromboembolic disease, inflammatory disease, or Alzheimer's Disease.

Toxicity and prophylactic efficacy of such compounds can be determined by standard pharmaceutical procedures in cell cultures or experimental animals, *e.g.*, for determining the LD50 (the dose lethal to 50% of the population) and the ED50 (the dose prophylactically effective in 50% of the population). The dose ratio between toxic and therapeutic effects is the therapeutic index and it can be expressed as the ratio LD50/ED50. Compounds that exhibit high therapeutic indices are preferred.

The data obtained from cell culture assays and animal studies can be used in formulating a range of dosage for use in humans. The dosage of compositions featured herein in the invention lies generally within a range of circulating concentrations that include the ED50, such as, an ED80 or ED90, with little or no toxicity. The dosage can vary within this range depending upon the dosage form employed and the route of administration utilized. For any compound used in the methods featured in the invention, the prophylactically effective dose can be estimated initially from cell culture assays. A dose can be formulated in animal models to achieve a circulating plasma concentration range of the compound or, when appropriate, of the polypeptide product of a target sequence (*e.g.*, achieving a decreased concentration of the polypeptide) that includes the IC50 (*i.e.*, the concentration of the test compound which achieves a half-maximal inhibition of symptoms) or higher levels of inhibition as determined in cell culture. Such information can be used to more accurately determine useful doses in humans. Levels in plasma can be measured, for example, by high performance liquid chromatography.

In addition to their administration, as discussed above, the iRNAs featured in the invention can be administered in combination with other known agents used for the prevention or treatment of an F12-associated disorder, *e.g.*, hereditary angioedema (HAE) (*e.g.*, hereditary angioedema type I; hereditary angioedema type II; hereditary angioedema type III; or any other hereditary angioedema caused by elevated levels of bradykinin); prekallikrein deficiency, hypertension, *e.g.*, malignant essential hypertension, end stage renal disease, Fletcher Factor Deficiency, thromboembolic disease, inflammatory disease, or Alzheimer's Disease. In any event, the administering physician can adjust the amount and timing of iRNA administration on the basis of results observed using standard measures of efficacy known in the art or described herein.

VI. Methods For Inhibiting F12 Expression

The present invention also provides methods of inhibiting expression of an F12 gene in a cell. The methods include contacting a cell with an RNAi agent, *e.g.*, double stranded RNA agent, in an amount effective to inhibit expression of F12 in the cell, thereby inhibiting expression of F12 in the

cell. In some embodiments of the disclosure, expression of an F12 gene is inhibited preferentially in the liver (*e.g.*, hepatocytes).

Contacting of a cell with an iRNA, *e.g.*, a double stranded RNA agent, may be done *in vitro* or *in vivo*. Contacting a cell *in vivo* with the iRNA includes contacting a cell or group of cells within a subject, *e.g.*, a human subject, with the iRNA. Combinations of *in vitro* and *in vivo* methods of contacting a cell are also possible. Contacting a cell may be direct or indirect, as discussed above. Furthermore, contacting a cell may be accomplished *via* a targeting ligand, including any ligand described herein or known in the art. In some embodiments, the targeting ligand is a carbohydrate moiety, *e.g.*, a GalNAc₃ ligand, or any other ligand that directs the RNAi agent to a site of interest.

The term “inhibiting,” as used herein, is used interchangeably with “reducing,” “silencing,” “downregulating,” “suppressing”, and other similar terms, and includes any level of inhibition.

The phrase “inhibiting expression of an F12” is intended to refer to inhibition of expression of any F12 gene (such as, *e.g.*, a mouse F12 gene, a rat F12 gene, a monkey F12 gene, or a human F12 gene) as well as variants or mutants of an F12 gene. Thus, the F12 gene may be a wild-type F12 gene, a mutant F12 gene, or a transgenic F12 gene in the context of a genetically manipulated cell, group of cells, or organism.

“Inhibiting expression of an F12 gene” includes any level of inhibition of an F12 gene, *e.g.*, at least partial suppression of the expression of an F12 gene. The expression of the F12 gene may be assessed based on the level, or the change in the level, of any variable associated with F12 gene expression, *e.g.*, F12 mRNA level or F12 protein level. This level may be assessed in an individual cell or in a group of cells, including, for example, a sample derived from a subject.

Inhibition may be assessed by a decrease in an absolute or relative level of one or more variables that are associated with F12 expression compared with a control level. The control level may be any type of control level that is utilized in the art, *e.g.*, a pre-dose baseline level, or a level determined from a similar subject, cell, or sample that is untreated or treated with a control (such as, *e.g.*, buffer only control or inactive agent control).

In some embodiments of the methods of the invention, expression of an F12 gene is inhibited by at least 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, or 95%, or to below the level of detection of the assay. In some embodiments, expression of an F12 gene is inhibited by at least 70%.

It is further understood that inhibition of F12 expression in certain tissues, *e.g.*, in liver, without a significant inhibition of expression in other tissues, *e.g.*, brain, may be desirable. In some embodiments, expression level is determined using the assay method provided in Example 2 with a 10 nM siRNA concentration in the appropriate species matched cell line.

In certain embodiments, inhibition of expression *in vivo* is determined by knockdown of the human gene in a rodent expressing the human gene, *e.g.*, an AAV-infected mouse expressing the human target gene (*i.e.*, F12), *e.g.*, when administered as a single dose, *e.g.*, at 3 mg/kg at the nadir of RNA expression. Knockdown of expression of an endogenous gene in a model animal system can

also be determined, *e.g.*, after administration of a single dose at, *e.g.*, 3 mg/kg at the nadir of RNA expression. Such systems are useful when the nucleic acid sequence of the human gene and the model animal gene are sufficiently close such that the human iRNA provides effective knockdown of the model animal gene. RNA expression in liver is determined using the PCR methods provided in

5 Example 2.

Inhibition of the expression of an F12 gene may be manifested by a reduction of the amount of mRNA expressed by a first cell or group of cells (such cells may be present, for example, in a sample derived from a subject) in which an F12 gene is transcribed and which has or have been treated (*e.g.*, by contacting the cell or cells with an iRNA of the invention, or by administering an
10 iRNA of the invention to a subject in which the cells are or were present) such that the expression of an F12 gene is inhibited, as compared to a second cell or group of cells substantially identical to the first cell or group of cells but which has not or have not been so treated (control cell(s) not treated with an iRNA or not treated with an iRNA targeted to the gene of interest). In some embodiments, the inhibition is assessed by the method provided in Example 2 using a 10nM siRNA concentration in
15 the species matched cell line and expressing the level of mRNA in treated cells as a percentage of the level of mRNA in control cells, using the following formula:

$$\frac{(\text{mRNA in control cells}) - (\text{mRNA in treated cells})}{(\text{mRNA in control cells})} \bullet 100\%$$

In other embodiments, inhibition of the expression of an F12 gene may be assessed in terms of a reduction of a parameter that is functionally linked to F12 gene expression, *e.g.*, F12 protein level
20 in blood or serum from a subject. F12 gene silencing may be determined in any cell expressing F12, either endogenous or heterologous from an expression construct, and by any assay known in the art.

Inhibition of the expression of an F12 protein may be manifested by a reduction in the level of the F12 protein that is expressed by a cell or group of cells or in a subject sample (*e.g.*, the level of protein in a blood sample derived from a subject). As explained above, for the assessment of mRNA
25 suppression, the inhibition of protein expression levels in a treated cell or group of cells may similarly be expressed as a percentage of the level of protein in a control cell or group of cells, or the change in the level of protein in a subject sample, *e.g.*, blood or serum derived therefrom.

A control cell, a group of cells, or subject sample that may be used to assess the inhibition of the expression of an F12 gene includes a cell, group of cells, or subject sample that has not yet been
30 contacted with an RNAi agent of the invention. For example, the control cell, group of cells, or subject sample may be derived from an individual subject (*e.g.*, a human or animal subject) prior to treatment of the subject with an RNAi agent or an appropriately matched population control.

The level of F12 mRNA that is expressed by a cell or group of cells may be determined using any method known in the art for assessing mRNA expression. In one embodiment, the level of
35 expression of F12 in a sample is determined by detecting a transcribed polynucleotide, or portion thereof, *e.g.*, mRNA of the F12 gene. RNA may be extracted from cells using RNA extraction

techniques including, for example, using acid phenol/guanidine isothiocyanate extraction (RNAzol B; Biogenesis), RNeasyTM RNA preparation kits (Qiagen®) or PAXgeneTM (PreAnalytixTM, Switzerland). Typical assay formats utilizing ribonucleic acid hybridization include nuclear run-on assays, RT-PCR, RNase protection assays, northern blotting, *in situ* hybridization, and microarray analysis.

In some embodiments, the level of expression of F12 is determined using a nucleic acid probe. The term “probe”, as used herein, refers to any molecule that is capable of selectively binding to a specific F12. Probes can be synthesized by one of skill in the art, or derived from appropriate biological preparations. Probes may be specifically designed to be labeled. Examples of molecules that can be utilized as probes include, but are not limited to, RNA, DNA, proteins, antibodies, and organic molecules.

Isolated mRNA can be used in hybridization or amplification assays that include, but are not limited to, Southern or northern analyses, polymerase chain reaction (PCR) analyses and probe arrays. One method for the determination of mRNA levels involves contacting the isolated mRNA with a nucleic acid molecule (probe) that can hybridize to F12 mRNA. In one embodiment, the mRNA is immobilized on a solid surface and contacted with a probe, for example by running the isolated mRNA on an agarose gel and transferring the mRNA from the gel to a membrane, such as nitrocellulose. In an alternative embodiment, the probe(s) are immobilized on a solid surface and the mRNA is contacted with the probe(s), for example, in an Affymetrix® gene chip array. A skilled artisan can readily adapt known mRNA detection methods for use in determining the level of F12 mRNA.

An alternative method for determining the level of expression of F12 in a sample involves the process of nucleic acid amplification or reverse transcriptase (to prepare cDNA) of for example mRNA in the sample, *e.g.*, by RT-PCR (the experimental embodiment set forth in Mullis, 1987, U.S. Patent No. 4,683,202), ligase chain reaction (Barany (1991) *Proc. Natl. Acad. Sci. USA* 88:189-193), self sustained sequence replication (Guatelli *et al.* (1990) *Proc. Natl. Acad. Sci. USA* 87:1874-1878), transcriptional amplification system (Kwoh *et al.* (1989) *Proc. Natl. Acad. Sci. USA* 86:1173-1177), Q-Beta Replicase (Lizardi *et al.* (1988) *Bio/Technology* 6:1197), rolling circle replication (Lizardi *et al.*, U.S. Patent No. 5,854,033) or any other nucleic acid amplification method, followed by the detection of the amplified molecules using techniques well known to those of skill in the art. These detection schemes are especially useful for the detection of nucleic acid molecules if such molecules are present in very low numbers. In particular aspects of the invention, the level of expression of F12 is determined by quantitative fluorogenic RT-PCR (*i.e.*, the TaqManTM System). In some embodiments, expression level is determined by the method provided in Example 2 using, *e.g.*, a 10 nM siRNA concentration, in the species matched cell line.

The expression levels of F12 mRNA may be monitored using a membrane blot (such as used in hybridization analysis such as northern, Southern, dot, and the like), or microwells, sample tubes,

gels, beads or fibers (or any solid support comprising bound nucleic acids). See U.S. Patent Nos. 5,770,722, 5,874,219, 5,744,305, 5,677,195 and 5,445,934, which are incorporated herein by reference. The determination of F12 expression level may also comprise using nucleic acid probes in solution.

5 In some embodiments, the level of mRNA expression is assessed using branched DNA (bDNA) assays or real time PCR (qPCR). The use of these methods is described and exemplified in the Examples presented herein. In some embodiments, expression level is determined by the method provided in Example 2 using a 10nM siRNA concentration in the species matched cell line.

10 The level of F12 protein expression may be determined using any method known in the art for the measurement of protein levels. Such methods include, for example, electrophoresis, capillary electrophoresis, high performance liquid chromatography (HPLC), thin layer chromatography (TLC), hyperdiffusion chromatography, fluid or gel precipitin reactions, absorption spectroscopy, a colorimetric assays, spectrophotometric assays, flow cytometry, immunodiffusion (single or double), immunoelectrophoresis, western blotting, radioimmunoassay (RIA), enzyme-linked immunosorbent
15 assays (ELISAs), immunofluorescent assays, electrochemiluminescence assays, and the like.

 In some embodiments, the efficacy of the methods of the invention are assessed by a decrease in F12 mRNA or protein level (*e.g.*, in a liver biopsy).

 In some embodiments, the efficacy of the methods of the invention can be monitored by detecting or monitoring a reduction in a symptom of an F12-associate disorder, *e.g.*, reduction in
20 edema swelling of the extremities, face, larynx, upper respiratory tract, abdomen, trunk, and genitals, prodrome; laryngeal swelling; nonpruritic rash; nausea; vomiting; or abdominal pain. It is well within the ability of one skilled in the art to monitor efficacy of the methods by measuring any one of such parameters, or any combination of parameters.

 In some embodiments of the methods of the invention, the iRNA is administered to a subject
25 such that the iRNA is delivered to a specific site within the subject. The inhibition of expression of F12 may be assessed using measurements of the level or change in the level of F12 mRNA or F12 protein in a sample derived from fluid or tissue from the specific site within the subject (*e.g.*, liver or blood).

 As used herein, the terms detecting or determining a level of an analyte are understood to
30 mean performing the steps to determine if a material, *e.g.*, protein, RNA, is present. As used herein, methods of detecting or determining include detection or determination of an analyte level that is below the level of detection for the method used.

VII. Prophylactic and Treatment Methods of the Invention

35 The present invention also provides methods of using an iRNA of the invention or a composition containing an iRNA of the invention to inhibit expression of F12, thereby preventing or treating an F12-associated disorder, *e.g.*, heredity angioedema (HAE) (*e.g.*, hereditary angioedema

type I; hereditary angioedema type II; hereditary angioedema type III; or any other hereditary angioedema caused by elevated levels of bradykinin); prekallikrein deficiency, hypertension, *e.g.*, malignant essential hypertension, end stage renal disease, Fletcher Factor Deficiency, thromboembolic disease, inflammatory disease, or Alzheimer's Disease. In the methods of the invention the cell may be contacted with the siRNA *in vitro* or *in vivo*, *i.e.*, the cell may be within a subject.

A cell suitable for treatment using the methods of the invention may be any cell that expresses an F12 gene, *e.g.*, a liver cell. A cell suitable for use in the methods of the invention may be a mammalian cell, *e.g.*, a primate cell (such as a human cell, including human cell in a chimeric non-human animal, or a non-human primate cell, *e.g.*, a monkey cell or a chimpanzee cell), or a non-primate cell. In certain embodiments, the cell is a human cell, *e.g.*, a human liver cell. In the methods of the invention, F12 expression is inhibited in the cell by at least 50, 55, 60, 65, 70, 75, 80, 85, 90, or 95, or to a level below the level of detection of the assay.

The *in vivo* methods of the invention may include administering to a subject a composition containing an iRNA, where the iRNA includes a nucleotide sequence that is complementary to at least a part of an RNA transcript of the F12 gene of the mammal to which the RNAi agent is to be administered. The composition can be administered by any means known in the art including, but not limited to oral, intraperitoneal, or parenteral routes, including intracranial (*e.g.*, intraventricular, intraparenchymal, and intrathecal), intravenous, intramuscular, subcutaneous, transdermal, airway (aerosol), nasal, rectal, intraocular (*e.g.*, periocular, conjunctival, subtenon, intracameral, intravitreal, intraocular, anterior or posterior juxtasceral, subretinal, subconjunctival, retrobulbar, or intracanalicular injection), intravenous, intramuscular, subcutaneous, transdermal, airway (aerosol), and topical (including buccal and sublingual) administration.

In certain embodiments, the compositions are administered by intravenous infusion or injection. In certain embodiments, the compositions are administered by subcutaneous injection. In certain embodiments, the compositions are administered by intramuscular injection.

The mode of administration may be chosen based upon whether local or systemic treatment is desired and based upon the area to be treated. The route and site of administration may be chosen to enhance targeting.

In one aspect, the present invention also provides methods for inhibiting the expression of an F12 gene in a mammal. The methods include administering to the mammal a composition comprising a dsRNA that targets an F12 gene in a cell of the mammal and maintaining the mammal for a time sufficient to obtain degradation of the mRNA transcript of the F12 gene, thereby inhibiting expression of the F12 gene in the cell. Reduction in gene expression can be assessed by any methods known in the art and by methods, *e.g.* qRT-PCR, described herein, *e.g.*, in Example 2. Reduction in protein production can be assessed by any methods known in the art, *e.g.* ELISA. In certain embodiments, a puncture liver biopsy sample serves as the tissue material for monitoring the reduction in the F12 gene

or protein expression. In other embodiments, a blood sample serves as the subject sample for monitoring the reduction in the F12 protein expression.

The present invention further provides methods of treatment in a subject in need thereof, *e.g.*, a subject diagnosed with an F12-associated disorder, such as heredity angioedema (HAE) (*e.g.*,
5 hereditary angioedema type I; hereditary angioedema type II; hereditary angioedema type III; or any other hereditary angioedema caused by elevated levels of bradykinin); prekallikrein deficiency, hypertension, *e.g.*, malignant essential hypertension, end stage renal disease, Fletcher Factor Deficiency, thromboembolic disease, inflammatory disease, or Alzheimer's Disease.

The present invention further provides methods of prophylaxis in a subject in need thereof.
10 The treatment methods of the invention include administering an iRNA of the invention to a subject, *e.g.*, a subject that would benefit from a reduction of F12 expression, in a prophylactically effective amount of a dsRNA targeting an F12 gene or a pharmaceutical composition comprising a dsRNA targeting an F12 gene.

In one aspect, the present invention provides methods of treating a subject having a disorder that
15 would benefit from reduction in F12 expression, *e.g.*, an F12-associated disorder, such as heredity angioedema (HAE) (*e.g.*, hereditary angioedema type I; hereditary angioedema type II; hereditary angioedema type III; or any other hereditary angioedema caused by elevated levels of bradykinin); prekallikrein deficiency, hypertension, *e.g.*, malignant essential hypertension, end stage renal disease, Fletcher Factor Deficiency, thromboembolic disease, inflammatory disease, or Alzheimer's Disease.

20 Treatment of a subject that would benefit from a reduction and/or inhibition of F12 gene expression includes therapeutic treatment (*e.g.*, a subject is having an F12-associated disorder) and prophylactic treatment (*e.g.*, the subject is not having an F12-associated disorder or a subject may be at risk of developing an F12-associated disorder).

In some embodiments, the F12-associated disorder is selected from the group consisting of
25 heredity angioedema (HAE) (*e.g.*, hereditary angioedema type I; hereditary angioedema type II; hereditary angioedema type III; or any other hereditary angioedema caused by elevated levels of bradykinin); prekallikrein deficiency, hypertension, *e.g.*, malignant essential hypertension, end stage renal disease, Fletcher Factor Deficiency, thromboembolic disease, inflammatory disease, and Alzheimer's Disease.

In some embodiments, the F12-associated disorder is HAE.

30 In other embodiments, the F12-associated disorder is prekallikrein deficiency.

In other embodiments, the F12-associated disorder is hypertension, *e.g.*, malignant essential hypertension.

In other embodiments, the F12-associated disorder is end stage renal disease.

In other embodiments, the F12-associated disorder is Fletcher Factor Deficiency.

35 In another embodiment, an F12-associated disorder is thromboembolic disease.

In another embodiment, an F12-associated disorder is inflammatory disease.

In another embodiment, an F12-associated disorder is Alzheimer's Disease.

In some embodiments, the RNAi agent is administered to a subject in an amount effective to inhibit F12 expression in a cell within the subject. The amount effective to inhibit F12 expression in a cell within a subject may be assessed using methods discussed above, including methods that involve assessment of the inhibition of F12 mRNA, F12 protein, or related variables, such as a reduction in the severity of a symptom of an F12-associate disorder, *e.g.*, reduction in the severity of edema swelling of the extremities, face, larynx, upper respiratory tract, abdomen, trunk, and genitals, prodrome; laryngeal swelling; nonpruritic rash; nausea; vomiting; or abdominal pain.

An iRNA of the invention may be administered as a “free iRNA.” A free iRNA is administered in the absence of a pharmaceutical composition. The naked iRNA may be in a suitable buffer solution. The buffer solution may comprise acetate, citrate, prolamine, carbonate, or phosphate, or any combination thereof. In one embodiment, the buffer solution is phosphate buffered saline (PBS). The pH and osmolarity of the buffer solution containing the iRNA can be adjusted such that it is suitable for administering to a subject.

Alternatively, an iRNA of the invention may be administered as a pharmaceutical composition, such as a dsRNA liposomal formulation.

Subjects that would benefit from an inhibition of F12 gene expression are subjects susceptible to or diagnosed with an F12-associated disorder, such as heredity angioedema (HAE) (*e.g.*, hereditary angioedema type I; hereditary angioedema type II; hereditary angioedema type III; or any other hereditary angioedema caused by elevated levels of bradykinin); prekallikrein deficiency, hypertension, *e.g.*, malignant essential hypertension, end stage renal disease, Fletcher Factor Deficiency, thromboembolic disease, inflammatory disease, or Alzheimer’s Disease. In an embodiment, the method includes administering a composition featured herein such that expression of the target ab F12 gene is decreased, such as for about 1, 2, 3, 4, 5, 6, 1-6, 1-3, or 3-6 months per dose. In certain embodiments, the composition is administered once every 3-6 months.

In one embodiment, the iRNAs useful for the methods and compositions featured herein specifically target RNAs (primary or processed) of the target F12 gene. Compositions and methods for inhibiting the expression of these genes using iRNAs can be prepared and performed as described herein.

Administration of the iRNA according to the methods of the invention may result prevention or treatment of an F12-associated disorder, *e.g.*, heredity angioedema (HAE) (*e.g.*, hereditary angioedema type I; hereditary angioedema type II; hereditary angioedema type III; or any other hereditary angioedema caused by elevated levels of bradykinin); prekallikrein deficiency, hypertension, *e.g.*, malignant essential hypertension, end stage renal disease, Fletcher Factor Deficiency, thromboembolic disease, inflammatory disease, or Alzheimer’s Disease. Subjects can be administered a therapeutic amount of iRNA, such as about 0.01 mg/kg to about 200 mg/kg.

In one embodiment, the iRNA is administered subcutaneously, *i.e.*, by subcutaneous injection. One or more injections may be used to deliver the desired dose of iRNA to a subject. The injections may be repeated over a period of time.

The administration may be repeated on a regular basis. In certain embodiments, after an
5 initial treatment regimen, the treatments can be administered on a less frequent basis. A repeat-dose regimen may include administration of a therapeutic amount of iRNA on a regular basis, such as once per month to once a year. In certain embodiments, the iRNA is administered about once per month to about once every three months, or about once every three months to about once every six months.

The invention further provides methods and uses of an iRNA agent or a pharmaceutical
10 composition thereof for treating a subject that would benefit from reduction and/or inhibition of F12 gene expression, *e.g.*, a subject having an F12-associated disorder, in combination with other pharmaceuticals and/or other therapeutic methods, *e.g.*, with known pharmaceuticals and/or known therapeutic methods, such as, for example, those which are currently employed for treating these disorders.

15 Accordingly, in some aspects of the invention, the methods which include administration of an iRNA agent of the invention, further include administering to the subject one or more additional therapeutic agents. For example, in certain embodiments, an iRNA targeting F12 is administered in combination with, *e.g.*, an agent useful in treating an F12-associated disorder. Exemplary additional therapeutics and treatments for treating an F12-associated disorder include, an androgen, such as
20 danazol or oxandrolone, Berinert®, Cinryze™, Rhuconest®, Ecallantide, Firazyr®, Kalbitor®, and a combination of any of the foregoing.

The iRNA and additional therapeutic agents may be administered at the same time and/or in the same combination, *e.g.*, parenterally, or the additional therapeutic agent can be administered as part of a separate composition or at separate times and/or by another method known in the art or
25 described herein.

The iRNA agent and an additional therapeutic agent and/or treatment may be administered at the same time and/or in the same combination, *e.g.*, parenterally, or the additional therapeutic agent can be administered as part of a separate composition or at separate times and/or by another method known in the art or described herein.

30

VIII. Kits

In certain aspects, the instant disclosure provides kits that include a suitable container containing a pharmaceutical formulation of a siRNA compound, *e.g.*, a double-stranded siRNA compound, or siRNA compound, (*e.g.*, a precursor, *e.g.*, a larger siRNA compound which can be
35 processed into a siRNA compound, or a DNA which encodes an siRNA compound, *e.g.*, a double-stranded siRNA compound, or ssiRNA compound, or precursor thereof).

Such kits include one or more dsRNA agent(s) and instructions for use, *e.g.*, instructions for administering a prophylactically or therapeutically effective amount of a dsRNA agent(s). The dsRNA agent may be in a vial or a pre-filled syringe. The kits may optionally further comprise means for administering the dsRNA agent (*e.g.*, an injection device, such as a pre-filled syringe), or means for measuring the inhibition of F12 (*e.g.*, means for measuring the inhibition of F12 mRNA, F12 protein, and/or F12 activity). Such means for measuring the inhibition of F12 may comprise a means for obtaining a sample from a subject, such as, *e.g.*, a plasma sample. The kits of the invention may optionally further comprise means for determining the therapeutically effective or prophylactically effective amount.

In certain embodiments the individual components of the pharmaceutical formulation may be provided in one container, *e.g.*, a vial or a pre-filled syringe. Alternatively, it may be desirable to provide the components of the pharmaceutical formulation separately in two or more containers, *e.g.*, one container for a siRNA compound preparation, and at least another for a carrier compound. The kit may be packaged in a number of different configurations such as one or more containers in a single box. The different components can be combined, *e.g.*, according to instructions provided with the kit. The components can be combined according to a method described herein, *e.g.*, to prepare and administer a pharmaceutical composition. The kit can also include a delivery device.

This invention is further illustrated by the following examples which should not be construed as limiting. The entire contents of all references, patents and published patent applications cited throughout this application, as well as the informal Sequence Listing and Figures, are hereby incorporated herein by reference.

EXAMPLES

Example 1. iRNA Synthesis

Source of reagents

Where the source of a reagent is not specifically given herein, such reagent can be obtained from any supplier of reagents for molecular biology at a quality/purity standard for application in molecular biology.

siRNA Design

siRNAs targeting the human Factor XII (F12) gene (human: NCBI refseqID NM_000505.4, NCBI GeneID: 2161) were designed using custom R and Python scripts. The human NM_000505.4 REFSEQ mRNA, has a length of 2036 bases.

- 5 Detailed lists of the unmodified F12 sense and antisense strand nucleotide sequences are shown in Table 2. Detailed lists of the modified F12 sense and antisense strand nucleotide sequences are shown in Table 3.

It is to be understood that, throughout the application, a duplex name without a decimal is equivalent to a duplex name with a decimal which merely references the batch number of the duplex.

- 10 For example, AD-959917 is equivalent to AD-959917.1.

siRNA Synthesis

siRNAs were designed, synthesized, and prepared using methods known in the art.

- Briefly, siRNA sequences were synthesized on a 1 μ mol scale using a Mermade 192
15 synthesizer (BioAutomation) with phosphoramidite chemistry on solid supports. The solid support was controlled pore glass (500-1000 Å) loaded with a custom GalNAc ligand (3'-GalNAc conjugates), universal solid support (AM Chemicals), or the first nucleotide of interest. Ancillary synthesis reagents and standard 2-cyanoethyl phosphoramidite monomers (2'-deoxy-2'-fluoro, 2'-O-methyl, RNA, DNA) were obtained from Thermo-Fisher (Milwaukee, WI), Hongene (China), or
20 Chemgenes (Wilmington, MA, USA). Additional phosphoramidite monomers were procured from commercial suppliers, prepared in-house, or procured using custom synthesis from various CMOs. Phosphoramidites were prepared at a concentration of 100 mM in either acetonitrile or 9:1 acetonitrile:DMF and were coupled using 5-Ethylthio-1H-tetrazole (ETT, 0.25 M in acetonitrile) with a reaction time of 400 s. Phosphorothioate linkages were generated using a 100 mM solution of 3-
25 ((Dimethylamino-methylidene) amino)-3H-1,2,4-dithiazole-3-thione (DDTT, obtained from Chemgenes (Wilmington, MA, USA)) in anhydrous acetonitrile/pyridine (9:1 v/v). Oxidation time was 5 minutes. All sequences were synthesized with final removal of the DMT group ("DMT-Off").

- Upon completion of the solid phase synthesis, solid-supported oligoribonucleotides were treated with 300 μ L of Methylamine (40% aqueous) at room temperature in 96 well plates for
30 approximately 2 hours to afford cleavage from the solid support and subsequent removal of all additional base-labile protecting groups. For sequences containing any natural ribonucleotide linkages (2'-OH) protected with a tert-butyl dimethyl silyl (TBDMS) group, a second deprotection step was performed using TEA.3HF (triethylamine trihydrofluoride). To each oligonucleotide solution in aqueous methylamine was added 200 μ L of dimethyl sulfoxide (DMSO) and 300 μ L TEA.3HF and
35 the solution was incubated for approximately 30 mins at 60 °C. After incubation, the plate was allowed to come to room temperature and crude oligonucleotides were precipitated by the addition of 1 mL of 9:1 acetonitrile:ethanol or 1:1 ethanol:isopropanol. The plates were then centrifuged at 4 °C

for 45 mins and the supernatant carefully decanted with the aid of a multichannel pipette. The oligonucleotide pellet was resuspended in 20 mM NaOAc and subsequently desalted using a HiTrap size exclusion column (5 mL, GE Healthcare) on an Agilent LC system equipped with an autosampler, UV detector, conductivity meter, and fraction collector. Desalted samples were collected
5 in 96 well plates and then analyzed by LC-MS and UV spectrometry to confirm identity and quantify the amount of material, respectively.

Duplexing of single strands was performed on a Tecan liquid handling robot. Sense and antisense single strands were combined in an equimolar ratio to a final concentration of 10 μ M in 1x PBS in 96 well plates, the plate sealed, incubated at 100 °C for 10 minutes, and subsequently allowed
10 to return slowly to room temperature over a period of 2-3 hours. The concentration and identity of each duplex was confirmed and then subsequently utilized for *in vitro* screening assays.

Example 2. *In vitro* screening methods

Cell culture and 384-well transfections

15 Primary human hepatocytes (PHH) and primary *cynomolgus* hepatocytes (PCH) (ATCC, Manassas, VA) are grown to near confluence at 37°C in an atmosphere of 5% CO₂ in Eagle's Minimum Essential Medium (Gibco) supplemented with 10% FBS (ATCC) before being released from the plate by trypsinization. Transfection was carried out by adding 7.5 μ l of Opti-MEM plus 0.1
20 μ l of Lipofectamine RNAiMax per well (Invitrogen, Carlsbad CA. cat # 13778-150) to 2.5 μ l of each siRNA duplex to an individual well in a 384-well plate. The mixture was then incubated at room temperature for 15 minutes. Forty μ l of complete growth media without antibiotic containing $\sim 1.5 \times 10^4$ cells was then added to the siRNA mixture. Cells were incubated for 24 hours prior to RNA purification. Single dose experiments were performed in PHH cells at 10 nM, 1 nM, and 0.1 nM final duplex concentration and in PCH cells at 10 nM, 1 nM, and 0.1 nM final duplex concentration.

25

Free uptake transfection

A subset of the duplexes were also assessed by free uptake. Five μ l of each duplex in PBS was combined with 5 μ l of OptiMEM and 10,000 freshly thawed primary *cynomolgus* hepatocytes (BioIVT) resuspended in 45 μ l of In Vitro Gro CP media (In Vitro Technologies- Celsis, Baltimore,
30 MD) in each well of a 384 well plate. The mixture was incubated for about 48 hours at 37 °C in an atmosphere of 5% CO₂. Single dose experiments were performed at 250 nM, 100 nM, 10 nM, or 1 nM final duplex concentration.

DYNABEADS mRNA Isolation Kit (Invitrogen™, part #: 610-12)

35 Cells were lysed in 75 μ l of Lysis/Binding Buffer containing 3 μ L of beads per well and mixed for 10 minutes on an electrostatic shaker. The washing steps were automated on a Biotek EL406, using a magnetic plate support. Beads were washed (in 90 μ L) once in Buffer A, once in

Buffer B, and twice in Buffer E, with aspiration steps in between. Following a final aspiration, complete 10µL RT mixture was added to each well, as described below.

cDNA synthesis using ABI High capacity cDNA reverse transcription kit (Applied Biosystems, Foster City, CA, Cat #4368813)

A master mix of 1µl 10X Buffer, 0.4µl 25X dNTPs, 1µl Random primers, 0.5µl Reverse Transcriptase, 0.5µl RNase inhibitor and 6.6µl of H₂O per reaction was added per well. Plates were sealed, agitated for 10 minutes on an electrostatic shaker, and then incubated at 37 degrees C for 2 hours. Following this, the plates were agitated at 80 degrees C for 8 minutes.

Real time PCR

Two microlitre (µl) of cDNA was added to a master mix containing 0.5µl of human GAPDH TaqMan Probe (4326317E), 0.5µl human F12, 2µl nuclease-free water and 5µl Lightcycler 480 probe master mix (Roche Cat # 04887301001) per well in a 384 well plates (Roche cat # 04887301001).

Real time PCR was done in a LightCycler480 Real Time PCR system (Roche).

To calculate relative fold change, data were analyzed using the $\Delta\Delta C_t$ method and normalized to assays performed with cells transfected with 10nM AD-1955, or mock transfected cells. IC₅₀s were calculated using a 4 parameter fit model using XLFit and normalized to cells transfected with AD-1955 or mock-transfected. The sense and antisense sequences of AD-1955 are: sense:

cuuAcGcuGAGuAcuucGAdTsdT (SEQ ID NO:11) and antisense
UCGAAGuACUcAGCGuAAGdTsdT (SEQ ID NO:12).


The results of the single dose screens of the dsRNA agents listed in Tables 2 and 3 in PHH cells are shown in Table 4.



The results of the single dose transfection screens of the dsRNA agents listed in Tables 2 and 3 in PCH cells are shown in Table 5.

The results of the single dose free uptake screens of the dsRNA agents listed in Tables 2 and 3 in PCH cells are shown in Table 6.

Table 1. Abbreviations of nucleotide monomers used in nucleic acid sequence representation. It will be understood that these monomers, when present in an oligonucleotide, are mutually linked by 5'-3'-phosphodiester bonds; and it is understood that when the nucleotide contains a 2'-fluoro modification, then the fluoro replaces the hydroxy at that position in the parent nucleotide (*i.e.*, it is a 2'-deoxy-2'-fluoronucleotide). It is to be further understood that the nucleotide abbreviations in the table omit the 3'-phosphate (*i.e.*, they are 3'-OH) when placed at the 3'-terminal position of an oligonucleotide.

Abbreviation	Nucleotide(s)
A	Adenosine-3'-phosphate
Ab	beta-L-adenosine-3'-phosphate
Abs	beta-L-adenosine-3'-phosphorothioate

Abbreviation	Nucleotide(s)
Af	2'-fluoroadenosine-3'-phosphate
Afs	2'-fluoroadenosine-3'-phosphorothioate
As	adenosine-3'-phosphorothioate
C	cytidine-3'-phosphate
Cb	beta-L-cytidine-3'-phosphate
Cbs	beta-L-cytidine-3'-phosphorothioate
Cf	2'-fluorocytidine-3'-phosphate
Cfs	2'-fluorocytidine-3'-phosphorothioate
Cs	cytidine-3'-phosphorothioate
G	guanosine-3'-phosphate
Gb	beta-L-guanosine-3'-phosphate
Gbs	beta-L-guanosine-3'-phosphorothioate
Gf	2'-fluoroguanosine-3'-phosphate
Gfs	2'-fluoroguanosine-3'-phosphorothioate
Gs	guanosine-3'-phosphorothioate
T	5'-methyluridine-3'-phosphate
Tf	2'-fluoro-5-methyluridine-3'-phosphate
Tfs	2'-fluoro-5-methyluridine-3'-phosphorothioate
Ts	5-methyluridine-3'-phosphorothioate
U	Uridine-3'-phosphate
Uf	2'-fluorouridine-3'-phosphate
Ufs	2'-fluorouridine-3'-phosphorothioate
Us	uridine-3'-phosphorothioate
N	any nucleotide, modified or unmodified
a	2'-O-methyladenosine-3'-phosphate
as	2'-O-methyladenosine-3'-phosphorothioate
c	2'-O-methylcytidine-3'-phosphate
cs	2'-O-methylcytidine-3'-phosphorothioate
g	2'-O-methylguanosine-3'-phosphate
gs	2'-O-methylguanosine-3'-phosphorothioate
t	2'-O-methyl-5-methyluridine-3'-phosphate
ts	2'-O-methyl-5-methyluridine-3'-phosphorothioate
u	2'-O-methyluridine-3'-phosphate
us	2'-O-methyluridine-3'-phosphorothioate
s	phosphorothioate linkage
L10	N-(cholesterylcarboxamidocaproyl)-4-hydroxyprolinol (Hyp-C6-Chol) 

Abbreviation	Nucleotide(s)
L96	N-[tris(GalNAc-alkyl)-amidodecanoyl]-4-hydroxyprolinol (Hyp-(GalNAc-alkyl)3) (2S,4R)-1-[29-[[2-(acetylamino)-2-deoxy-β-D-galactopyranosyl]oxy]-14,14-bis[[3-[[3-[[5-[[2-(acetylamino)-2-deoxy-β-D-galactopyranosyl]oxy]-1-oxopentyl]amino]propyl]amino]-3-oxopropoxy]methyl]-1,12,19,25-tetraoxo-16-oxa-13,20,24-triazanonacos-1-yl]-4-hydroxy-2-hydroxymethylpyrrolidine 
uL96	2'-O-methyluridine-3'-phosphate ((2S,4R)-1-[29-[[2-(acetylamino)-2-deoxy-β-D-galactopyranosyl]oxy]-14,14-bis[[3-[[3-[[5-[[2-(acetylamino)-2-deoxy-β-D-galactopyranosyl]oxy]-1-oxopentyl]amino]propyl]amino]-3-oxopropoxy]methyl]-1,12,19,25-tetraoxo-16-oxa-13,20,24-triazanonacos-1-yl]-4-hydroxy-2-pyrrolidinyl)methyl ester 



Abbreviation	Nucleotide(s)
Y34	2-hydroxymethyl-tetrahydrofuran-4-methoxy-3-phosphate (abasic 2'-OMe furanose) 
Y44	inverted abasic DNA (2-hydroxymethyl-tetrahydrofuran-5-phosphate) 
(Agn)	Adenosine-glycol nucleic acid (GNA) S-Isomer
(Cgn)	Cytidine-glycol nucleic acid (GNA) S-Isomer
(Ggn)	Guanosine-glycol nucleic acid (GNA) S-Isomer
(Tgn)	Thymidine-glycol nucleic acid (GNA) S-Isomer
P	Phosphate
VP	Vinyl-phosphonate
dA	2'-deoxyadenosine-3'-phosphate
dAs	2'-deoxyadenosine-3'-phosphorothioate
dC	2'-deoxycytidine-3'-phosphate
dCs	2'-deoxycytidine-3'-phosphorothioate
dG	2'-deoxyguanosine-3'-phosphate
dGs	2'-deoxyguanosine-3'-phosphorothioate
dT	2'-deoxythymidine-3'-phosphate
dTs	2'-deoxythymidine-3'-phosphorothioate
dU	2'-deoxyuridine
dUs	2'-deoxyuridine-3'-phosphorothioate
(C2p)	cytidine-2'-phosphate
(G2p)	guanosine-2'-phosphate
(U2p)	uridine-2'-phosphate
(A2p)	adenosine-2'-phosphate
(Chd)	2'-O-hexadecyl-cytidine-3'-phosphate
(Ahd)	2'-O-hexadecyl-adenosine-3'-phosphate
(Ghd)	2'-O-hexadecyl-guanosine-3'-phosphate
(Uhd)	2'-O-hexadecyl-uridine-3'-phosphate

Table 2. Unmodified Sense and Antisense Strand Sequences of F12 dsRNA Agents

Duplex Name	Sense Strand Sequence 5' to 3'	SEQ ID NO:	Range in NM_000505.4	Antisense Strand Sequence 5' to 3'	SEQ ID NO:	Range in NM_000505.4
AD-1534462	AAAGAGAAAGUGCUUUUGAGCCU	21	431-451	AGGCTCAAAAGCACUUCUCUUUCU	285	429-451
AD-1534463	AAAUUAGCCACAAAGGGCCGU	22	215-235	ACGGCCUUGUGGGUACAUUUGU	286	213-235
AD-1534464	AACCCGGACAACGACAUCCGU	23	839-859	ACGGAUGUCGUTGUCGGGUUCC	287	837-859
AD-1534465	AACUCAAUAAAGUGCUUUGAU	24	2008-2028	ATCAAAGCACUTUAUUGAGUUCC	288	2006-2028
AD-1534466	AACUGAGCAAGCAGCUGUGGU	25	493-513	ACCACAGCUGTUGCUCAGUUUC	289	491-513
AD-1534467	AAGAAUGAGAUUGGUUAUAGU	26	473-493	ACUATACCAUATCUCAUUCUUUG	290	471-493
AD-1534468	AAGAGCACACAGUCGUUCUCU	27	138-158	AGAGAACGACUGUGUCUCUUA	291	136-158
AD-1534469	AAGCAGCUGUGGCCAGAUUCU	28	501-521	AGCATCTGGCCACAGCUCUUGC	292	499-521
AD-1534470	AAGCUGCUAUGAUGGCCCGCU	29	682-702	ACGCGGCCAUCAUAGCAGCUUGC	293	680-702
AD-1534471	AAGUACAAAGCUGAAGAGCAU	30	125-145	AUGCTCTUCAGCUUUUGACUUAU	294	123-145
AD-1534472	AAGUCUUUAGGCCUCAGCUU	31	437-457	AAGCTGAGGCUCAAAAGCAGCUU	295	435-457
AD-1534473	AAUAAAGUGCUUUUGAAAUGU	32	2013-2033	ACAUTUTCAAAGCACUUUAUUGA	296	2011-2033
AD-1534474	ACAAAUUGUACCCACAAAGGGCU	33	213-233	AGCCCUUGUGGGUACAUUUGUGG	297	211-233
AD-1534475	ACACAGUCGUUUCACUGUCU	34	144-164	AGACAGTGAGAACGACUGUGUGC	298	142-164
AD-1534476	ACACCGAUGUGGCCUACUACU	35	1833-1853	AGUAGUAGGCCACAUCCGGUGUAG	299	1831-1853
AD-1534477	ACACUUUCGAUUCACACCUUGU	36	86-106	ACAAGGTGGAATCGAAAAGUGUUG	300	84-106
AD-1534478	ACACUUUCGAUUCACACCUUGU	37	86-106	ACAAGGTGGAUUCGAAAGUGUUG	301	84-106
AD-1534479	ACAGUCGUUUCACACUGUCACU	38	146-166	AGUGACAGUGAGAACGACUGUGU	302	144-166
AD-1534480	ACAUCGCCGCGCUGUACUGGU	39	1194-1214	ACCAGUACAGCGCGCGAGUAG	303	1192-1214
AD-1534481	ACCAGCACGACCUUGGCUUGU	40	1410-1430	ACAGAGCCAGGTCGUGCUGGUAG	304	1408-1430
AD-1534482	ACCCGGACAACGACAUCGCGU	41	840-860	AGCGGATGUCGUUUGCCGGGUUC	305	838-860
AD-1534483	ACCCUGCAAGGCAUACUACAGU	42	1772-1792	ACUGAUGAUGCCUUGCAGGGUGA	306	1770-1792
AD-1534484	ACCGCAACAAGCCAGGCGUCU	43	1812-1832	AGACGCCUGGCTUGUUGCGGUCA	307	1810-1832
AD-1534485	ACCUGGCUCUGUUGCGCCUUU	44	1419-1439	AAAGGCGCAACAGAGCCAGGUGC	308	1417-1439
AD-1534486	ACGUUCAGCCGUGUGCCUGU	45	1479-1499	ACAGGCACACCGGCGUACGUAA	309	1477-1499
AD-1534487	ACUACCUUGGCCUGGAUCCGGU	46	1848-1868	ACCGGATCCAGGCCAGGUAGUAG	310	1846-1868

Duplex Name	Sense Strand Sequence 5' to 3'	SEQ ID NO:	Range in NM_000505.4	Antisense Strand Sequence 5' to 3'	SEQ ID NO:	Range in NM_000505.4
AD-1534488	ACUCAUAAAAGUGCUUUAGAAU	47	2009-2029	ATUCAAAAGCACTUUAUUAGAGUUC	311	2007-2029
AD-1534489	ACUGAGCAAGCAGCUGUGGCU	48	494-514	AGCCACAGCUGCUUGCUCAGUUC	312	492-514
AD-1534490	AGAGAAAGUGCUUUAGAGCCUCU	49	433-453	AGAGGCTCAAAAGCACUUCUCUUU	313	431-453
AD-1534491	AGAGCACACAGUCGUUUCUCAU	50	139-159	AUGAGAACGACUGUGUGCUCUUC	314	137-159
AD-1534492	AGAGUCUGUCUUCGAUGACCU	51	1131-1151	AGGUCATCGAAGACAGACUUCUUG	315	1129-1151
AD-1534493	AGAUGCCAGUGCAAGGGUCCU	52	515-535	AGGACCCUUGCACUGGCAUCUGG	316	513-535
AD-1534494	AGCAGCACCUGGUCUGUUGU	53	1413-1433	ACAACAGAGCCAGGUCGUGCUGG	317	1411-1433
AD-1534495	AGCCAGGCGUCUACACCGAAU	54	1821-1841	AUUGGTGUAGACGCCUGGCUUG	318	1819-1841
AD-1534496	AGCCAGGCGUCUACACCGAAU	55	1821-1841	AUUGGTGUAGACGCCUGGCUUG	319	1819-1841
AD-1534497	AGCCCAAGAAAGUGAAAGACU	56	306-326	AGUCTUTCACUUUCUUGGGCUC	320	304-326
AD-1534498	AGCUGAAGAGCACACAGUCGU	57	133-153	ACGACUGUGUGUCUUCAGCUUU	321	131-153
AD-1534499	AGCUGCUAUGAUGGCCCGCGU	58	683-703	ACCGCGGCCAUCAUAGCAGCUUG	322	681-703
AD-1534500	AGCUGUACCAACAAUUGUACCU	59	204-224	AGGUACAUUUGUGGUACAGCUGC	323	202-224
AD-1534501	AGGAGCAUAAAGUACAAAGCUU	60	117-137	AAGCTUTGUACTUAUGCUCCUUG	324	115-137
AD-1534502	AGGAGCAUAAAGUACAAAGCUU	61	117-137	AAGCTUTGUACUUAUGCUCCUUG	325	115-137
AD-1534503	AGGAUGGCGCAGGAACUCAU	62	1995-2015	AUUGAGTUCUUGCGCCAUCCUGG	326	1993-2015
AD-1534504	AGGCGUCUACACCGAUGUGGU	63	1825-1845	ACCACATCGGUGUAGACGCCUGG	327	1823-1845
AD-1534505	AGGCUUCAUGUCCCAUCUCAU	64	965-985	AAUGAGTGGGACAUAGAAAGCCUAG	328	963-985
AD-1534506	AGUCAACACUUUCGAUUCUCCAU	65	81-101	ATGGAATCGAAAGUGUUGACUCC	329	79-101
AD-1534507	AGUCAACACUUUCGAUUCUCCAU	66	81-101	AUGGAATCGAAAGUGUUGACUCC	330	79-101
AD-1534508	AUAAAAGUGCUUUUGAAAAUGCU	67	2014-2034	AGCATUTUCAAAAGCACUUUAUUG	331	2012-2034
AD-1534509	AUCCGCCCGUGGUGCUUCGUU	68	854-874	AACGAAGCACACCACGGGGCGAUGU	332	852-874
AD-1534510	AUCGGGCGUGUGUGACCCGCAU	69	1798-1818	AUGCGGTACCCACAGCCCGAUCC	333	1796-1818
AD-1534511	AUGCCAGUGCAAGGGUCCUUGU	70	517-537	ACAGGACCCUUGCACUGGCAUCU	334	515-537
AD-1534512	AUGCUCUGGCAGGGUUCUUCU	71	1673-1693	AAGGAACCCUUGCGCAGAGCAUGC	335	1671-1693
AD-1534513	AUGUACCCACAAGGGCCGCGU	72	217-237	AGCCGGCCCUUGUGGGUACAUUU	336	215-237
AD-1534514	AUUGCUCAGGACUACUUCU	73	1885-1905	AAAGAUGAGUCCUAGAGCAUCA	337	1883-1905
AD-1534515	CAAAGCUGAAGAGCACACAGU	74	130-150	ACUGTGTGCUCUUCAGCUUUUGUA	338	128-150

Duplex Name	Sense Strand Sequence 5' to 3'	SE Q ID NO:	Range in NM_000505.4	Antisense Strand Sequence 5' to 3'	SEQ ID NO:	Range in NM_000505.4
AD-1534516	CAAAUGUACCCACAAAGGGCCU	75	214-234	AGGCCCTUGUGGGUACAUUUGUG	339	212-234
AD-1534517	CAAGAAAAGUGAAAGACCACUU	76	310-330	AAGUGGTUUUACAUUUUCUUGGG	340	308-330
AD-1534518	CAAGAGUCUGUCUUCGAUGAU	77	1129-1149	ATCATCGAAGACAGACUCUUGCG	341	1127-1149
AD-1534519	CAAGGAGCAUAAAGUACAAAGU	78	115-135	ACUUTGTACUU AUGCUCCUUGGG	342	113-135
AD-1534520	CACAAAUGUACCCACAAAGGU	79	212-232	ACCCTUGUGGGTACAUUUGUGU	343	210-232
AD-1534521	CACCCUGCAAGGCAUCAUCAU	80	1771-1791	AUGATGAUGCCUUGCAGGGUGAG	344	1769-1791
AD-1534522	CACCGAUGUGGGCUACUACCU	81	1834-1854	AGGUAGTAGGCCACAUCCGUGUA	345	1832-1854
AD-1534523	CACGACCUUGGCUUGUUGCGU	82	1415-1435	ACGCAACAGAGCCAGGUUGCGU	346	1413-1435
AD-1534524	CACUGUCUCUGUCCACAACAU	83	389-409	AUGUTGTGGACAGAGACAGUGGG	347	387-409
AD-1534525	CACUUUCGAUUCACCUUGGU	84	87-107	ACCAAGGUGGAAUCGAAAGUGUU	348	85-107
AD-1534526	CAGAUCCAGUGCAAGGGUCU	85	514-534	AGACCTTUGCACUCUGGCAUCUGGC	349	512-534
AD-1534527	CAGCACGACCUGGUCUGUUU	86	1412-1432	AAACAGAGCCAGGUUGUUGGU	350	1410-1432
AD-1534528	CAGCUGUACCAACAAUUGUACU	87	203-223	AGUACATUUGUGGUACAGCUGCC	351	201-223
AD-1534529	CAGCUGUGGCCAGAUGCCAGU	88	504-524	ACUGGCAUCUGGCCACAGCUGCU	352	502-524
AD-1534530	CAGGAACUCAAUAAAGUGCUU	89	2004-2024	AAGCACTUUAUUGAGUUCUUGCG	353	2002-2024
AD-1534531	CAGGAUGGGCGAGGAACUCAU	90	1994-2014	AUGAGUTCCUGCGCCAUCCUGGC	354	1992-2014
AD-1534532	CAGGCGUCUACACCGAUGUGU	91	1824-1844	ACACAUCGGUGTAGACGCCUGGC	355	1822-1844
AD-1534533	CAGGGACUCAUGUUUCCUCUCU	92	1891-1911	AGAGGGAAGATGAGUCCUUGAG	356	1889-1911
AD-1534534	CAGUGCAAAGGUCCUGAUGCU	93	521-541	AGCATCAGGACCCUUGCACUGGC	357	519-541
AD-1534535	CCACAAAUGUACCCACAAGGU	94	211-231	ACCUTGTGGGUACAUUUGUGGUA	358	209-231
AD-1534536	CCACAAAUGUACCCACAAGGU	95	211-231	ACCUTGTGGGUACAUUUGUGGUA	359	209-231
AD-1534537	CCAGAUCCAGUGCAAGGGUU	96	513-533	AACCCUTGCACTGGCAUCUUGGCC	360	511-533
AD-1534538	CCAGGCGUCUACACCGAUGUU	97	1823-1843	AACATCGGUGUAGACGCCUUGGU	361	1821-1843
AD-1534539	CCCAAGAAAGUGAAAGACCAU	98	308-328	AUGGTCTUUCACUUUCUUGGGCU	362	306-328
AD-1534540	CCCGACAACGACAUCCGCCU	99	841-861	AGCGGAUGUCGUUUGUCCGGGUU	363	839-861
AD-1534541	CCCGACAACGACAUCCGCCU	100	841-861	AGCGGAUGUCGUUUGUCCGGGUU	364	839-861
AD-1534542	CCUGCAAGGCAUCAUCAGCU	101	1773-1793	AGCUGATGAUGCCUUGCAGGGUG	365	1771-1793
AD-1534543	CCGAUGUGGCCUACUACCUUGU	102	1836-1856	ACAGGUAGUAGGCCACAUCCGGUG	366	1834-1856

Duplex Name	Sense Strand Sequence 5' to 3'	SEQ ID NO:	Range in NM_000505.4	Antisense Strand Sequence 5' to 3'	SEQ ID NO:	Range in NM_000505.4
AD-1534544	CCGCAACAAAGCCAGGCGUCUU	103	1813-1833	AAGACGCCUGGCUUUGCGGUC	367	1811-1833
AD-1534545	CCGCAAGAGUCUGUCUUCGAU	104	1126-1146	ATCGAAGACAGACUCUUGCGGAG	368	1124-1146
AD-1534546	CCGGAACCCGGACAACGACAU	105	835-855	ATGUCGTUGUCCGGUUUCCGGCA	369	833-855
AD-1534547	CCGGAACCCGGACAACGACAU	106	835-855	AUGUCGTUGUCCGGUUUCCGGCA	370	833-855
AD-1534548	CCUGCAGGAGGCGCAGGUACU	107	1591-1611	AGUACCTGCGCCUCCUGCAGGAA	371	1589-1611
AD-1534549	CCUGUCGCCUUACGUUCAGCU	108	1468-1488	AGCUGAACGUAAGGCGACAGGAG	372	1466-1488
AD-1534550	CCUGUCGCCUUACGUUCAGCU	109	1468-1488	AGCUGAACGUAAGGCGACAGGAG	373	1466-1488
AD-1534551	CCUUACGUUCAGCCGGUGUGU	110	1475-1495	ACACCCGGCUGAACGUAAGGCG	374	1473-1495
AD-1534552	CGCAACAAGCCAGGCGUCUUAU	111	1814-1834	ATAGACGCCUGGCUUUGCGGU	375	1812-1834
AD-1534553	CGCAGGAACUCAUAAAGUGU	112	2002-2022	ACACTUTAUUGAGUUCUCCGCC	376	2000-2022
AD-1534554	CGCCUUACGUUCAGCCGGUGU	113	1473-1493	ACACCCGGCUGAACGUAAGGCGAC	377	1471-1493
AD-1534555	CGCUCCUGUCGCCUUACGUUU	114	1464-1484	AAACGUAAGGCGACAGGAGCGCG	378	1462-1484
AD-1534556	CGGAACCCGGACAACGACAUU	115	836-856	AAUATCGUUGUCCGGUUUCCGGC	379	834-856
AD-1534557	CGGCUCCGCAAGAGUCUGUCU	116	1121-1141	AGACAGACUCUUGCGGAGCCGCU	380	1119-1141
AD-1534558	CGUCUACACCGAUUGUGCCUU	117	1828-1848	AAGGCCACAUCCGUGUAGACGCC	381	1826-1848
AD-1534559	CGUGCCAGGGUGAUUCCGGAU	118	1710-1730	ATCCGGAAUCACCCUGGCACGCA	382	1708-1730
AD-1534560	CUACAUCCGCCGCGCUGUACUU	119	1192-1212	AAGUACAGCGCGGCGAUGUAGGG	383	1190-1212
AD-1534561	CUACCUUGGCCUGGAUCCCGGU	120	1849-1869	ACCCGGAUCCAGGCCAGGUAGUA	384	1847-1869
AD-1534562	CUCAAUAAAAGUGCUUUGAAAU	121	2010-2030	ATUUCAAAAGCACUUUAUUGAGUU	385	2008-2030
AD-1534563	CUCAAUAAAAGUGCUUUGAAAU	122	2010-2030	AUUUCAAAGCACUUUAUUGAGUU	386	2008-2030
AD-1534564	CUCAGGGACUACUUCUUCUCCU	123	1889-1909	AGGGAAGAUGAGUCCUUGAGCA	387	1887-1909
AD-1534565	CUCUGUCGCCUUACGUUACAU	124	1466-1486	ATGAACGUAAGGCGACAGGAGCG	388	1464-1486
AD-1534566	CUCUGUUGCGCCUUCAGGAGU	125	1425-1445	ACUCCUGAAGGCGCAACAGAGCC	389	1423-1445
AD-1534567	CUGAAGAGCACACAGUCUUAU	126	135-155	AAACGACUGUGTGCUUCUACGCU	390	133-155
AD-1534568	CUGAUUGCUCAGGGACUCAUU	127	1882-1902	AAUGAGTCCUGAGCAUUCAGGA	391	1880-1902
AD-1534569	CUGCAAGGCAUACAGCUGU	128	1775-1795	ACAGCUGAUGATGCCUUGCAGGG	392	1773-1795
AD-1534570	CUGCCGGAACCCGGACAACGU	129	832-852	ACGUTGTCCGGUUUCCGGCAGAA	393	830-852
AD-1534571	CUGUACCAACAUAUGUACCCAU	130	206-226	AUGGGUACAUAUUGUGGUACAGCU	394	204-226

Duplex Name	Sense Strand Sequence 5' to 3'	SEQ ID NO:	Range in NM_000505.4	Antisense Strand Sequence 5' to 3'	SEQ ID NO:	Range in NM_000505.4
AD-1534572	CUGUCGCCUUACGUUUCAGCCU	131	1469-1489	AGGCTGAACGUAAAGGCGACAGGA	395	1467-1489
AD-1534573	CUGUCUCUGUCCACAACACCU	132	391-411	AGGUGUTGUGGACACAGACAGUG	396	389-411
AD-1534574	CUGUGGUGACCGCAACAAGCU	133	1804-1824	AGCUTGTUGCGGUCACCAACAGCC	397	1802-1824
AD-1534575	CUUACGUUACAGCCGUGUGCU	134	1476-1496	AGCACACCGGCTGAACGUAAAGC	398	1474-1496
AD-1534576	CUUCGAUGACCCGCGUGGUU	135	1140-1160	AAACGACGCGGUAUCGAAAGAC	399	1138-1160
AD-1534577	CUUGGAGUCAACACUUUCGAU	136	76-96	ATCGAAAGUGUTGACUCCAAAGCU	400	74-96
AD-1534578	CUUUGAGCCUCAGCUUUCUCCU	137	442-462	AGGAGAAAGCUGAGGCUCAAAAGCA	401	440-462
AD-1534579	GAAAGUGAAAGACCACUGCAU	138	313-333	AUGCAGTGGUCUUUACUUUCUU	402	311-333
AD-1534580	GAACCCGGACAACGACAUCU	139	838-858	AGGATGTCGUUGUCCGGGUCCG	403	836-858
AD-1534581	GAACUCAUAAGUGCUUUGU	140	2007-2027	ACAAAGCACUUAUUGAGUUCU	404	2005-2027
AD-1534582	GAAGAGCACACAGUCGUUCU	141	137-157	AAGAACGACUGTGUGCUUCUAG	405	135-157
AD-1534583	GACCGCAACAAGCCAGGCGUU	142	1811-1831	AACGCTGGCUUUGUUGCGGUCAC	406	1809-1831
AD-1534584	GAGCACACAGUCGUUCUCACU	143	140-160	AGUGAGAACGACUGUGGUCUCUU	407	138-160
AD-1534585	GAGCAUAAAGUACAAAGCUGAU	144	119-139	AUCAGCTUUGUACUUAUGCUCCU	408	117-139
AD-1534586	GAGCCUCAGCUUCUCCGGUUU	145	446-466	AAACCGGAGAAAGCUGAGGCUCAA	409	444-466
AD-1534587	GAGUCUGUCUUCGUAUGACCCU	146	1132-1152	AGGGTCAUCGAAAGACAGACUCUU	410	1130-1152
AD-1534588	GAUGCCAGUGCAAGGGUCCUU	147	516-536	AAGGACCCUUGCACUGGCAUCUG	411	514-536
AD-1534589	GAUGGCGCAGGAACUCAUAU	148	1997-2017	AUAUTGAGUUCCUGCGCAUCCU	412	1995-2017
AD-1534590	GAUUGCUCAGGGACUCUAUCUU	149	1884-1904	AAGATGAGUCCUUGAGCAAUACAG	413	1882-1904
AD-1534591	GCAAGAGUCUGUCUUCGUAUGU	150	1128-1148	ACAUCGAAGACACAGACUUGCGG	414	1126-1148
AD-1534592	GCACACAGUCGUUCUCACUGU	151	142-162	ACAGTGAGAAACGACUGUGUCUC	415	140-162
AD-1534593	GCACGACCUGGCUUCUGUUGCU	152	1414-1434	AGCAACAGAGCCAGGUCGUGCUG	416	1412-1434
AD-1534594	GCACGACCUGGCUUCUGUUGCU	153	1414-1434	AGCAACAGAGCCAGGUCGUGCUG	417	1412-1434
AD-1534595	GCAGCUGUGGCCAGAUGCCAU	154	503-523	AUGGCATCUGGCCACAGCUGGUU	418	501-523
AD-1534596	GCAGGAACUCAUAAAGUGCU	155	2003-2023	AGCACUTUAUUGAGUUCUUGCGC	419	2001-2023
AD-1534597	GCCAGAUGCCAGUGCAAGGGU	156	512-532	ACCCTUGCACUGGCAUCUGGCCA	420	510-532
AD-1534598	GCCAGGCGUACACCGGAUGU	157	1822-1842	ACAUCGGUGUAGACGCCUUGGCUU	421	1820-1842
AD-1534599	GCCCAAGAAAGUGAAAGACCU	158	307-327	AGGUCUTUCACUUUUGGGCUC	422	305-327

Duplex Name	Sense Strand Sequence 5' to 3'	SE Q ID NO:	Range in NM_000505.4	Antisense Strand Sequence 5' to 3'	SEQ ID NO:	Range in NM_000505.4
AD-1534600	GCCUACGUAACGCGGUGUU	159	1474-1494	AACACCGGCGUAACGUAAGGCGA	423	1472-1494
AD-1534601	GCGCAGGAACUCAUAAAGUU	160	2001-2021	AACUTUAUUGAGUUCUGCGCCA	424	1999-2021
AD-1534602	GCGCCUUCAGGAGGAUGCGGU	161	1432-1452	ACCGCATCCUCCUGAAGGCGCAA	425	1430-1452
AD-1534603	GCGGCUCGCAAGAGUCUGUU	162	1120-1140	AACAGACUCUUGCGGAGCGGUG	426	1118-1140
AD-1534604	GCUCAGGACUCACUUCUUCU	163	1888-1908	AGAAAGAUAGUCCUAGGCAA	427	1886-1908
AD-1534605	GCUCCGCAAGAGUCUGUUCU	164	1123-1143	AAAGACAGACUUCUUGGGAGCGG	428	1121-1143
AD-1534606	GCUCUGUCGCCUUCAGUUCU	165	1465-1485	AGAACGTAGGCGACAGGAGCGC	429	1463-1485
AD-1534607	GCUCUGCGCAGGUUCCUCGU	166	1675-1695	ACGAGGAACCTTGGCAGAGCAU	430	1673-1695
AD-1534608	GCUCUGCGCAGGUUCCUCGU	167	1675-1695	ACGAGGAACCCUCCGAGAGCAU	431	1673-1695
AD-1534609	GCUCUGUUGCGCCUUCAGGAU	168	1424-1444	ATCCTGAAGGCGCAACAGAGCCA	432	1422-1444
AD-1534610	GCUCUGUUGCGCCUUCAGGAU	169	1424-1444	AUCCTGAAGGCGCAACAGAGCCA	433	1422-1444
AD-1534611	GCUGAAGAGCACACAGUCUGU	170	134-154	AACGACTGUGUGCUCUUCAGCUU	434	132-154
AD-1534612	GCUGCUAUGAUGGCCGCGGGU	171	684-704	ACCCGCGGCCATCAUAGCAGCUU	435	682-704
AD-1534613	GCUGGUGUGAGGACCAAGU	172	1735-1755	ACUUGGTCCUCACACACAGCGG	436	1733-1755
AD-1534614	GCUGUGGUGACCGCAACAAGU	173	1803-1823	ACUUGUTGCGGTACACAGCCC	437	1801-1823
AD-1534615	GCUGUGGUGACCGCAACAAGU	174	1803-1823	ACUUGUTGCGGUCACACAGCCC	438	1801-1823
AD-1534616	GCUUCAGUCCCAUCUAGCU	175	967-987	AGCATGAGUGGGAUAGAGCCU	439	965-987
AD-1534617	GCUUUGAGCCUCAGCUUCUCU	176	441-461	AGAGAAGCUGAGGCGCUCAAAGCAC	440	439-461
AD-1534618	GGAAACUCAAUAAAGUGCUUUU	177	2006-2026	AAAAGCACUUUAUUGAUUCCUG	441	2004-2026
AD-1534619	GGAGCAUAAGUACAAAGCUGU	178	118-138	ACAGCUTUGUACUUAUGCUCUUU	442	116-138
AD-1534620	GGAGCCCAAGAAAGUGAAAAGU	179	304-324	ACUUTCACUUUCUUGGGCUCCAA	443	302-324
AD-1534621	GGAGUCAACACUUUCGAUUCU	180	79-99	AGAATCGAAAGTGUUACUCCAA	444	77-99
AD-1534622	GGCGUCUACACCGAUUGGGCU	181	1826-1846	AGCCACAUCGGTGUAGACGCCUG	445	1824-1846
AD-1534623	GGCGUCUACACCGAUUGGGCU	182	1826-1846	AGCCACAUCGGUGUAGACGCCUG	446	1824-1846
AD-1534624	GGCUCGCAAGAGUCUGUCUU	183	1122-1142	AAGACAGACUUCTUGCGGAGCCG	447	1120-1142
AD-1534625	GGCUCUGUUGCGCCUUCAGGU	184	1423-1443	ACCUGAAGGCGCAACAGAGCCAG	448	1421-1443
AD-1534626	GGGUCCUGAUGCCCAUCUGCCU	185	529-549	AGGCAGTGGGCAUCAGGACCCUU	449	527-549
AD-1534627	GUCAACACUUUCGAUUCACU	186	82-102	AGUGGAAUCCGAAAGUGUUGACUC	450	80-102

Duplex Name	Sense Strand Sequence 5' to 3'	SE Q ID NO:	Range in NM_000505.4	Antisense Strand Sequence 5' to 3'	SEQ ID NO:	Range in NM_000505.4
AD-1534628	GUCAACACUUUCGUAUCCACU	187	82-102	AGUGAAUCGAAAGUGUUGACUC	451	80-102
AD-1534629	GUCGCCUUACGUUCAGCCGGU	188	1471-1491	ACCGGTGAACGUAAAGCGACAG	452	1469-1491
AD-1534630	GUCGCCUUACGUUCAGCCGGU	189	1471-1491	ACCGGTGAACGUAAAGCGACAG	453	1469-1491
AD-1534631	GUCGUUCUCACUGUACCCGGU	190	149-169	ACCGGUGACAGTGAGAACGACUG	454	147-169
AD-1534632	GUCUGUCUUCGUAUGACCCGGU	191	1134-1154	AGCGGTCAUCGAAAGACAGACUC	455	1132-1154
AD-1534633	GUCUUCGAUGACCCGGGUGU	192	1138-1158	ACGACGGGGUACUAGAACAGACAG	456	1136-1158
AD-1534634	GUGCAAGGUUCUUGAUGCCCU	193	523-543	AGGCATCAGGACCCUUGACUG	457	521-543
AD-1534635	GUGGUGACCCGCAACAAGCCAU	194	1806-1826	AUGGCTGUUGCGGUACACACAG	458	1804-1826
AD-1534636	GUUGCGCCUUCAGGAGGAGU	195	1429-1449	ACAUCCTCCUGAAGGCGAACAG	459	1427-1449
AD-1534637	GUUGCGCCUUCAGGAGGAGU	196	1429-1449	ACAUCCTCCUGAAGGCGAACAG	460	1427-1449
AD-1534638	UAAAGUACAAAGCUGAAGAGCU	197	124-144	AGCUCUTCAGCTUUGUACUUAUG	461	122-144
AD-1534639	UAAAGUACAAAGCUGAAGAGCU	198	124-144	AGCUCUTCAGCUUUGUACUUAUG	462	122-144
AD-1534640	UACAUCGCGCGCUGUACUGU	199	1193-1213	ACAGTACAGCGCGGCGAUGUAGG	463	1191-1213
AD-1534641	UACCUUGCCUGGAUCCGGGAU	200	1850-1870	ATCCCCGAUCCAGGCGGAGUAGU	464	1848-1870
AD-1534642	UACUACCUUGCCUGGAUCCCGU	201	1847-1867	ACGGAUCCAGGCGCAGGUAGUAGG	465	1845-1867
AD-1534643	UCAACACUUUCGAUUCACCU	202	83-103	AGGUGGAAUUCGAAAGUGUUGACU	466	81-103
AD-1534644	UCACCCUGCAAGGCAUCAUCU	203	1770-1790	AGAUGATGCCUUGCAGGGUGAGC	467	1768-1790
AD-1534645	UCAGCCGGUGUGCCUGGCCAAU	204	1483-1503	AUUGGCAGGCACACCGGCGUAAAC	468	1481-1503
AD-1534646	UCAGGAGGAUGCGGACGGCAU	205	1438-1458	AUGCCGTCCGCAUCCUCCUGAAG	469	1436-1458
AD-1534647	UCAGGGACUACUUCUCCCUU	206	1890-1910	AAGGGAAGAUAGAGUCCUAGAGC	470	1888-1910
AD-1534648	UCCGCAAGAGUCUGUUCUUCGU	207	1125-1145	ACGAAGACAGACUCUUGCGGAGC	471	1123-1145
AD-1534649	UCCUGUCGCCUUAACGUUCAGU	208	1467-1487	ACUGAACGUAAAGCGACAGGAGC	472	1465-1487
AD-1534650	UCGCCUUAACGUUCAGCCGGU	209	1472-1492	AACCGGUGAACGUAAAGCGGACA	473	1470-1492
AD-1534651	UCUCUGUCCACAACACCUCAU	210	394-414	AUGAGGTGUUGUGGACAGAGACA	474	392-414
AD-1534652	UCUGUUGCGCCUUCAGGAGGU	211	1426-1446	ACCUCTGAAGGCGCAACAGAGC	475	1424-1446
AD-1534653	UGAAGAGCACACAGUCGUUCU	212	136-156	AGAACGACUGUGUCUUCUACAGC	476	134-156
AD-1534654	UGAGCCUACGCUUCUCCGGU	213	445-465	AACCGGAGAACGUGAGGCUCAAA	477	443-465
AD-1534655	UGAGCUGGGAGUACUGCGACU	214	891-911	AGUCGCAGUACUCCAGCUCAGC	478	889-911

Duplex Name	Sense Strand Sequence 5' to 3'	SE Q ID NO:	Range in NM_000505.4	Antisense Strand Sequence 5' to 3'	SEQ ID NO:	Range in NM_000505.4
AD-1534656	UGAUGGCCGCGGGCUCAGCUU	215	691-711	AAGCTGAGCCCGCGGCCAUAUA	479	689-711
AD-1534657	UGCAAGGCAUCAACAGCUGGU	216	1776-1796	ACCAGCTGAUGAUGCCUUGCAGG	480	1774-1796
AD-1534658	UGCAAGGCAUCAACAGCUGGU	217	1776-1796	ACCAGCTGAUGAUGCCUUGCAGG	481	1774-1796
AD-1534659	UGCAAGGGUCCUGAUGCCCAU	218	524-544	AUGGGCAUCAGGACCCUUGCACU	482	522-544
AD-1534660	UGCCAGUGCAAGGGUCCUGAU	219	518-538	AUCAGGACCCUUGCACUUGGCAUC	483	516-538
AD-1534661	UGCCAUAGAGGGUCCUGCGU	220	34-54	AGCAGCAGAGCCUUCUUGGCAUC	484	32-54
AD-1534662	UGCGCCUUCAGGAGGAGUGCGU	221	1431-1451	ACGCAUCCUCCCTGAAGGCGCAAC	485	1429-1451
AD-1534663	UGCUCAGGGACUACUUCUUCU	222	1887-1907	AGAAAGAUAGAGTCCCUUGAGCAAU	486	1885-1907
AD-1534664	UGCUCAGGGACUACUUCUUCU	223	1887-1907	AGAAAGAUAGAGUCCCUUGAGCAAU	487	1885-1907
AD-1534665	UGGAGUCAACACUUCGUAUU	224	78-98	AAAUCCGAAAGUUGUAGACUCCAAAG	488	76-98
AD-1534666	UGGCCAGAUGCCAGUGCAAGU	225	510-530	ACUUGCACUUGGCAUCUUGGCCACA	489	508-530
AD-1534667	UGGCCUACUACCUUGGCCUGGU	226	1842-1862	ACCAGGCCAGGTAGUAGGCCACA	490	1840-1862
AD-1534668	UGGCUCUGUUGCGCCUUCAGU	227	1422-1442	ACUGAAGGCGCAACAGAGGCCAGG	491	1420-1442
AD-1534669	UGGUGACCGCAACAAGCCAGU	228	1807-1827	ACUGGCTUGUUGCGGUACCCACA	492	1805-1827
AD-1534670	UGUACCAACAAAUAGUACCCACU	229	207-227	AGUGGGTACAUUUGUGGUACAGC	493	205-227
AD-1534671	UGUCUCUGUCCACAAACCCU	230	392-412	AAGGTGTUGUGGACAGAGACAGU	494	390-412
AD-1534672	UGUGGCCAGAUGCCAGUGCAU	231	508-528	AUGCACTGGCAUCUGGCCACAGC	495	506-528
AD-1534673	UUACGUUCAGCCGGUGUGCCU	232	1477-1497	AGGCACACCCGGCUGAACGUAAGG	496	1475-1497
AD-1534674	UUCAGUCCCAACUACUGCCCU	233	969-989	AGGGCATGAGUGGGACAUAGAAGC	497	967-989
AD-1534675	UUCGAUGACCCGGUGUGUUGU	234	1141-1161	ACAACGACGCGGGUACUUGAAGA	498	1139-1161
AD-1534676	UUGAGCCUACAGCUUCUCCGGU	235	444-464	ACCGGAGAAAGTGTAGGCUCAAAAG	499	442-464
AD-1534677	UUGGAGCCCAAGAAAAGUGAAU	236	302-322	AUUCACTUUCUUGGGCUCCAAAC	500	300-322
AD-1534678	UUGGAGUCAACACUUCGUAUU	237	77-97	AAUCGAAAAGUGTUGACUCCAAAGC	501	75-97
AD-1534679	UUUGAGCCUACAGCUUCUCCGU	238	443-463	ACGGAGAAGCUGAGGUCUCAAAGC	502	441-463
AD-1631272	AAGAGCACACAGUCUUCUCU	239	138-158	AGAGAAGCAGUCUGUGUCUUCU	503	136-158
AD-1631273	AAUAAAGUGCUUUGAAAAUGU	240	2013-2033	ACAUTUTCAAAGCAGCUUUAUUCU	504	2011-2033
AD-1631274	ACAUCGCCGCGCUGUACUGGU	241	1194-1214	ACCAGUACAGCGCGCGGAGUUCU	505	1192-1214
AD-1631275	ACCAGCACGACCUUGGCUUCU	242	1410-1430	ACAGAGCCAGGTCGUGCUGGUCU	506	1408-1430

Duplex Name	Sense Strand Sequence 5' to 3'	SE Q ID NO:	Range in NM_000505.4	Antisense Strand Sequence 5' to 3'	SEQ ID NO:	Range in NM_000505.4
AD-1631276	ACCCUGCAAGGCAUCAACAGU	243	1772-1792	ACUGAUGAUGCCUUGCAGGGUCU	507	1770-1792
AD-1631277	ACCGCAACAAGCCAGGCGUCU	244	1812-1832	AGACGCCUGGCTUGUUUGCGUCU	508	1810-1832
AD-1631278	ACUACCCUGGCGGUAUCCGGU	245	1848-1868	ACCGGATCCAGGCCAGGUAGUCU	509	1846-1868
AD-1631279	AUUGCUCAGGGACUCACUUCU	246	1885-1905	AAAGAUGAGUCCUGAGCAAUCU	510	1883-1905
AD-1631280	CAGGGACUCACUUCUCCUCU	247	1891-1911	AGAGGAAAGATGAGUCCUCCUCU	511	1889-1911
AD-1631281	CCACAAUGUACCCACAAGGU	248	211-231	ACCUTGTGGGUACAUUUGUGGCU	512	209-231
AD-1631282	CCGCAAGAGUCUGUUCGUAU	249	1126-1146	ATCGAAGACAGACUCUUGCGGCU	513	1124-1146
AD-1631283	CCGGAACCCGGACAACGACAU	250	835-855	ATGUCGTUGUCCGGGUUCCGGCU	514	833-855
AD-1631284	CCUGUCGCCUUCGUUCAGCU	251	1468-1488	AGCUGAACGUAAGGCGACAGGCU	515	1466-1488
AD-1631285	CGCCUUCGUUCAGCCGGUGU	252	1473-1493	ACACCGGCUGAACGUAAGGCGCU	516	1471-1493
AD-1631286	CGUGCCAGGGUGAUUCCGGAU	253	1710-1730	ATCCGGAUUCACCCUGGCACGCU	517	1708-1730
AD-1631287	CUACCUGGCCUGGAUCCGGGU	254	1849-1869	ACCCGGAUCCAGGCCAGGUAGCU	518	1847-1869
AD-1631288	CUCAGGGACUCACUUCUCCCU	255	1889-1909	AGGGAAGAUGAGUCCUCCUAGCU	519	1887-1909
AD-1631289	CUUCGAUGACCCGCGUCGUUU	256	1140-1160	AAACGACGCGGUAUCGAAAGCU	520	1138-1160
AD-1631290	GAAGAGCACACAGUCGUUCUU	257	137-157	AAGAACGACUGTGUGCUUUCUU	521	135-157
AD-1631291	GAGCCUCAGCUUCUCCGGUUU	258	446-466	AAACCGGAGAAAGCUGAGGCUCCU	522	444-466
AD-1631292	GCCAGAUGCCAGUGCAAGGGU	259	512-532	ACCCTUGCACUGGCAUCUGGGCU	523	510-532
AD-1631293	GCCUUCGUUCAGCCGGUGUU	260	1474-1494	AACACCGGCUGAACGUAAGGCU	524	1472-1494
AD-1631294	GCGCAGGAACUCAUAAAGUU	261	2001-2021	AACUTUAUUGAGUUCUUGCGCCU	525	1999-2021
AD-1631295	GCGCCUUCAGGAGGAUGCGGU	262	1432-1452	ACCGCATCCUCCUUGAAGGCGCCU	526	1430-1452
AD-1631296	GCUCAGGGACUCACUUCUCCU	263	1888-1908	AGGAAAGAUGAGUCCUUGAGCCU	527	1886-1908
AD-1631297	GCUCUGCGCAGGGUUCUCCUGU	264	1675-1695	ACGAGGAACCCCTGCGCAGAGCCU	528	1673-1695
AD-1631298	GCUCUGUUGCGCCUUCAGGAU	265	1424-1444	ATCCTGAAGGCGCAACAGAGCCU	529	1422-1444
AD-1631299	GCUUUGAGCCUCAGCUUCUCU	266	441-461	AGAGAAGCUGAGGCUCAAAGCCU	530	439-461
AD-1631300	GGAGUCAACACUUCGUAUUCU	267	79-99	AGAATCGAAAGTGUUGACUCCCU	531	77-99
AD-1631301	GGCUCUGUUGCGCCUUCAGGU	268	1423-1443	ACCUGAAGGCGCAACAGAGCCCU	532	1421-1443
AD-1631302	GUCGCCUUCAGUUCAGCCGGU	269	1471-1491	ACCGGTGAACGUAAGGCGACCU	533	1469-1491
AD-1631303	GUCUUCGAUGACCCGCGUCCU	270	1138-1158	ACGACGCGGGUACUUGAAGACCU	534	1136-1158

Duplex Name	Sense Strand Sequence 5' to 3'	SEQ ID NO:	Range in NM_000505.4	Antisense Strand Sequence 5' to 3'	SEQ ID NO:	Range in NM_000505.4
AD-1631304	GUUGCGCCUUCAGGAGGAGU	271	1429-1449	ACAUCCTCCUGAAGGCGCAACCU	535	1427-1449
AD-1631305	UCGCCUUCAGUUCAGCCGGU	272	1472-1492	AACCGGCUGAACGUAAAGCGACU	536	1470-1492
AD-1631306	UGAGCCUCAGCUUCUCCGGU	273	445-465	AACCGGAGAAGCUGAGGCUCACU	537	443-465
AD-1631307	UGCGCCUUCAGGAGGAGUCGU	274	1431-1451	ACGCAUCCUCCTGAAGGCGCACU	538	1429-1451
AD-1631308	UGCUCAGGGACUACUUCU	275	1887-1907	AGAAAGAUAGTCCCUAGAGCACU	539	1885-1907
AD-1631309	UGGAGUCAACACUUCGAAU	276	78-98	AAUUCGAAAAGUUGAGCUCCACU	540	76-98
AD-1631310	UGGCCUACUACCUGGCCUGGU	277	1842-1862	ACCAGGCCAGGTAGUAGGCCACU	541	1840-1862
AD-1631311	UUCGAUGACCCGGUCGUUGU	278	1141-1161	ACAACGACGGGGUACUUGAACU	542	1139-1161
AD-1631312	UUGAGCCUCAGCUUCUCCGGU	279	444-464	ACCGGAGAAAGCTGAGGCUCACU	543	442-464
AD-85525	ACUCAUAAAGUGCUUUGAAA	280	2009-2029	UUUCAAAAGCACUUUAUUGAGUUC	544	2007-2029
AD-85531	UCAAUAAAGUGCUUUGAAAAU	281	2011-2031	AUUUCAAAAGCACUUUAUUGAGU	545	2009-2031
AD-85537	CUCAUAAAGUGCUUUGAAAA	282	2010-2030	UUUUCAAAAGCACUUUAUUGAGUU	546	2008-2030
AD-85541	GAACUCAUAAAGUGCUUUGA	283	2007-2027	UCAAAAGCACUUUAUUGAGUCCU	547	2005-2027
AD-67246	AAUAAAGUGCUUUGAAAAACGU	284	2013-2030	ACGUUUUCAAAAGCACUUUAUUGA	548	2011-2030

Table 3. Modified Sense and Antisense Strand Sequences of F12 dsRNA Agents

Duplex Name	Sense Strand Sequence 5' to 3'	SEQ ID NO:	Antisense Strand Sequence 5' to 3'	SEQ ID NO:	mRNA target sequence 5' to 3'	SEQ ID NO:
AD-1534462	asasagagaaGfUfGfcuuugagccuL96	549	asGfsgcdTc(Agn'aagcacUfuCfucuuuscsu	813	AGAAAGAGAAAGUGCUUUUGAGCCU	1077
AD-1534463	asasauguaeCfCfAfcagggccguL96	550	asdCsggdCcdCuugndGgGfuaaunusgsu	814	ACAAAUUGUACCCACAAAGGCCCGG	1078
AD-1534464	asaseceggaCfAfAfcgacauccguL96	551	asdCsggdAudGuegudTgUfcccggguiscsc	815	GGAACCCGGACAAACGACAUCCGC	1079
AD-1534465	asascucaauAfAfAfgugcuuugauL96	552	asdTscadAadGcaudTuAfuagguiscsc	816	GGAAACUCAAUAAAGUGCUUUUGAA	1080
AD-1534466	asascugagcAfAfGfcagcuugguL96	553	asdCscadCadGcugcdTuGfucagunuscsu	817	AGAAACUGAGCAAGCAGCUGUGGC	1081
AD-1534467	asasagaugaGfAfUfauugguuugauL96	554	asdCsuadTadCcaudTcUfcauucunusgsu	818	ACAAGAAUGAGAUUUGGUUAUAGA	1082

Duplex Name	Sense Strand Sequence 5' to 3'	SEQ ID NO:	Antisense Strand Sequence 5' to 3'	SEQ ID NO:	mRNA target sequence 5' to 3'	SEQ ID NO:
AD-1534468	asasgagcacAfcAfugcguucucL96	555	asdGsagdAadCgacudGuGfugcucucsa	819	UGAAGAGCACACAGUCGUCUUCUA	1083
AD-1534469	asasgcagcuGfUfGfgccagauGcuL96	556	asGfscadTc(Tgn)ggccacAfcGfugcucusc	820	GCAAAGCAGCUGUGGGCCAGAUGCC	1084
AD-1534470	asasgcugcuAfuGfGfagggccgcuL96	557	asdCsagdGcdCcaudAuAfcgagcucusc	821	GCAAAGCUGCUAUGAUGGGCCGCG	1085
AD-1534471	asasguacaaAfcGfCfugaagagcuL96	558	asUfsgcdTc(Tgn)ueagcuUfuGfuaucuas	822	AUAAGUACAAAGCUGAAGAGCAC	1086
AD-1534472	asasguGcuUfGfAfgccucagcuL96	559	asAfsagdTg(Agn)ggcucaAfaGfcaucusc	823	AGAAAGUGCUUUGAGCCUCAGCUU	1087
AD-1534473	asasuaaaguGfCfUfuaaagcuL96	560	asdCsaudTudTcaadGcAfcuuuauusga	824	UCAAUAAAGUGCUUUGAAAAUUGC	1088
AD-1534474	ascsaaauGuAfcCfcaaacggcuL96	561	asdGsacdCudTguggdGuAfcuuuugusg	825	CCACAAUGUACCCACAAAGGGCC	1089
AD-1534475	ascsacagucGfUfUfcacacugcuL96	562	asGfsacdAg(Tgn)ggaaacGfaCfugugusg	826	GCACACAGUCGUUUCACACUGUCA	1090
AD-1534476	ascsaccagauGfUfGfgccuacuL96	563	asGfsuadGu(Agn)ggccacAfuCfugugusag	827	CUACACCGAUGUGGGCCUACUACC	1091
AD-1534477	ascsacuucGfAfuUfuaacucguL96	564	asdCsacdGgdTggadTcGfaaagugusag	828	CAACACUUUCGAUUCACCUUGG	1092
AD-1534478	ascsacuucGfAfuUfuaacucguL96	565	asCfsaadGg(Tgn)ggaaacGfaAfaugugusg	829	CAACACUUUCGAUUCACCUUGG	1093
AD-1534479	ascsagucguUfcUfUfcacugcuL96	566	asGfsugdAc(Agn)guagagAfcGfagugusgu	830	ACACAGUCGUUUCACUGUCACC	1094
AD-1534480	ascsauegceGfCfGfuguaucugcuL96	567	asdCsacdGudAcagcdGcGfagcagugusag	831	CUACAUCCGCGCGCUGUACUGGG	1095
AD-1534481	ascsacagcacGfAfcUfcugugcugcuL96	568	asdCsagdAgdCcaagdTcGfugcugugusag	832	CUACACAGCAGCAGCUGGCUUCU	1096
AD-1534482	ascsccggacAfcAfcGfacaucgcuL96	569	asGfscgdGa(Tgn)guagcuGfuCfaggugusuc	833	GAACCCGGACAAACGACAUCCGCC	1097
AD-1534483	ascsccugcaAfcGfGfcaucaucagcuL96	570	asdCsugdAudGaugcdCuUfcagagugusga	834	UCACCCUGCAAGGCAUCAUCAGC	1098
AD-1534484	ascsccgaacAfcAfcGfcaagcugcuL96	571	asdGsacdGcdCuggdTuGfugcggusasa	835	UGACCCGCAACAAGCCAGGCGUCU	1099
AD-1534485	ascsccugcguCfUfGfugcggccuuL96	572	asdAsagdGcdGcaadAgAfcagcagucscg	836	CGACCCUGGCUUGUUGCGCCUUC	1100
AD-1534486	ascsguucagCfCfGfugugcugcuL96	573	asCfsagdGc(Agn)ccaggCfuGfcaagucasa	837	UUACGUUACGCCGGUGUGCCUGC	1101
AD-1534487	ascsuaccugGfCfCfuggaucgcuL96	574	asdCsagdGadTccagdGcCfaggaugusag	838	CUACUACCCUGGCUUGGAUCCCGG	1102
AD-1534488	ascsucaauaAfcAfcGfugcuuugaL96	575	asdTsuedAadAgcaadTuUfaugagucusc	839	GAACUCAUAUAAAGUGCUUUGAAA	1103
AD-1534489	ascsugagcaAfcGfCfagcugugcuL96	576	asGfscgdAc(Agn)gcuGcuUfcGfcaagucusc	840	GAACUGAGCAAGCAGCUGUGGGCC	1104
AD-1534490	asgsagaaguGfCfUfuaagcucucL96	577	asGfsagdGc(Tgn)caaacAfcUfucucucusc	841	AAAGAGAAAGUGCUUUGAGCCUCA	1105
AD-1534491	asgsagacacaCfAfcGfugcucucuaL96	578	asUfsgadGa(Agn)cgagcuUfgUfugcucusc	842	GAAGAGCACACAGCUGCUUCAC	1106
AD-1534492	asgsagucugUfcUfUfucgaugacucL96	579	asGfsugdCa(Tgn)cgaaagCfaGfcaucucusc	843	CAAGAGUCUGUCUUCUGAUGACCC	1107

Duplex Name	Sense Strand Sequence 5' to 3'	SEQ ID NO:	Antisense Strand Sequence 5' to 3'	SEQ ID NO:	mRNA target sequence 5' to 3'	SEQ ID NO:
AD-1534493	asgsaagccaGfUfGfcaagggucuuL96	580	asdGsgadCcdCuugcdAcUfggcaucussg	844	CCAGAUGCCAGUGCAAGGGUCCU	1108
AD-1534494	asgsacagacCfUfGfgeucuguuL96	581	asdCsaadCadGagcdAgGfueugucussg	845	CCAGCACGACCUUGUCUUGUUGC	1109
AD-1534495	asgsccaggeGfUfCfuacacgaauL96	582	asdAsuedGgdTguagcdAcGfueugucussg	846	CAAGCCAGGCGUUCACACCCGAUG	1110
AD-1534496	asgsccaggeGfUfCfuacacgaauL96	583	asAfsuedGg(Tgn)guagacGfcfueugucussg	847	CAAGCCAGGCGUUCACACCCGAUG	1111
AD-1534497	asgscccaagAfAfAfugaaagacuL96	584	asGfsuedTu(Tgn)cauuuCuUfgggucscsc	848	GGAGCCCAAGAAAGUGAAAGACC	1112
AD-1534498	asgsccgaagAfGfCfacacagucuuL96	585	asdCsgadCudGuugcdCuCfuacagucuu	849	AAAGCUGAAGAGCACACAGUCGU	1113
AD-1534499	asgsccgcuafGfAfugcccgccguL96	586	asdCscgdCgdGecandCaUfagecagucussg	850	CAAGCUGCUAUGAUGGCCCGCGGG	1114
AD-1534500	asgsccguuacCfAfCfaaanguuacuuL96	587	asGfsugdAc(Agn)uuugugGfuAfcagcuscsc	851	GCAGCUGUACCCACAAAUGUACCC	1115
AD-1534501	asgsccgacauAfAfGfuacaagaauL96	588	asdAsgcdTudTguacdTuAfuugcucussg	852	CAAGGAGCAUAAAGUACAAAGCUG	1116
AD-1534502	asgsccgacauAfAfGfuacaagaauL96	589	asAfsugdTu(Tgn)guacuuAfuGfueucussg	853	CAAGGAGCAUAAAGUACAAAGCUG	1117
AD-1534503	asgsccaggeGfCfAfggaacuaauL96	590	asUfsugdAg(Tgn)uucugeGfcCfuacucussg	854	CCAGGAUGGCGCAGGAACUCAAU	1118
AD-1534504	asgsccgcuafCfAfCfcgauguguuL96	591	asdCscadCadTcgguGdGuAfgagccucussg	855	CCAGGCGUCUACACCGAUGUGGC	1119
AD-1534505	asgsccuicaUfGfUfcccacuaauL96	592	asAfsugdAg(Tgn)gggacaUfgAfcagcusaag	856	CUAGGCUUCAUGUCCCCACUCAUG	1120
AD-1534506	asgsucaacaCfUfUfuegaauccauL96	593	asdTsggdAadTcggaadAgUfguuagucuscsc	857	GGAGUCAACACUUUCGAUUCAC	1121
AD-1534507	asgsucaacaCfUfUfuegaauccauL96	594	asUfsggdAa(Tgn)cgaaagUfgUfagucuscsc	858	GGAGUCAACACUUUCGAUUCAC	1122
AD-1534508	asusaaagugCfUfUfugaaaauuuL96	595	asdGscadTudTuaadAgCfauuuuauusug	859	CAAUAAAAGUGCUUUUGAAAAUGCU	1123
AD-1534509	asusccgcccGfUfGfugucucuuL96	596	asdAsecgdAadGecacdAcGfgegggaussu	860	ACAUCCGCCCGUGGUGCUUCCUG	1124
AD-1534510	asusccgggcuGfUfGfugagccgaauL96	597	asUfsugdGg(Tgn)caaccAcfGcfccgauscsc	861	GGAUCCGGGCUUGUGGUGACCGCAA	1125
AD-1534511	asusccagugGfCfAfcagggucuuL96	598	asdCsaadGadCccuudGcAfcuggaucscsu	862	AGAUGCCAGUGCAAGGGUCCUGA	1126
AD-1534512	asusccucugCfGfCfcagggucuuL96	599	asdAsggdAadCccuudGcCfcaggaucuscsc	863	GCAUGCUUGGCGCAGGGUCCUC	1127
AD-1534513	asusguacccAfCfAfagggcgccuuL96	600	asdGscgdGgdCccuudGuGfgegaucuu	864	AAAUUACCCACAAAGGGCGCGCC	1128
AD-1534514	asusgucaCaGfGfGfaucauuuL96	601	asdAsagdAudGaguedCcUfgegaucusa	865	UGAUUGCUCAGGACUCUACUUCU	1129
AD-1534515	csasaaagcugAfAfGfagcacacagaL96	602	asCfsugdTu(Tgn)guacuuCfaGfauuugsusa	866	UACAAAAGCUGAAGAGCACACAGU	1130
AD-1534516	csasaauguaCfCfCfacaaggccuL96	603	asGfsgcdCc(Tgn)uuguggUfaCfauuugsug	867	CACAAAUGUACCCACCAAGGGCGG	1131
AD-1534517	csasagaagUfGfAfaggaaccuuL96	604	asAfsugdGg(Tgn)cuuuaCfuUfueuugsgsg	868	CCCAAGAAAAGUGAAAGACACACUG	1132

Duplex Name	Sense Strand Sequence 5' to 3'	SEQ ID NO:	Antisense Strand Sequence 5' to 3'	SEQ ID NO:	mRNA target sequence 5' to 3'	SEQ ID NO:
AD-1534518	csasagagagcUfGfUfucucgaugauL96	605	asdTscadTcdGaagadCaGfacucugscsg	869	CGCAAGAGUCUGUCUUCGAUGAC	1133
AD-1534519	csasagagagcAfUfAfaguaaagauL96	606	asCfsuudTg(Tgn)acuuauGfcUfucucugsg	870	CCCAAGGAGCAUAAAGUACAAAGC	1134
AD-1534520	csascaaaugUfAfCfcacaaaggguL96	607	asdCscedTudGugggdTcCfaauuugugsgsu	871	ACCACAAUAGUACCCACCAAGGGC	1135
AD-1534521	csascccuvcAfAfGfgeucaucauL96	608	asUfsgadTg(Agn)ugccuGfcAfgggugsgasg	872	CUCACCCUGCAAGGCAUCAUCAG	1136
AD-1534522	csascggaugUfGfGfcucauacuuL96	609	asGfsgudAg(Tgn)agggcaCfaUfegguggsusa	873	UACACCGAUGUGGGCCUACUACCU	1137
AD-1534523	csascgaccuGfGfCfucugugcgulL96	610	asdCsgcdAadCagagdTcAfggucugsgscsu	874	AGCACGACCUUGGCUUCUUGCGC	1138
AD-1534524	csascugucuCfUfGfuccacaacauL96	611	asUfsgudTg(Tgn)ggacagAfGfAfcagugsgsg	875	CCCACUGUCUCUGUCCACAACAC	1139
AD-1534525	csascuucgAfUfUfcccacuuugguL96	612	asdCscadAgdGuggadAuCfgaagugsgsu	876	AACACUUUCGAUUCACCCUUGGG	1140
AD-1534526	csasgaugccAfGfUfgcgaaggguL96	613	asGfsgadCc(Tgn)ugcacuGfGfCfaucugsgsc	877	GCCAGAUGCCAGUGCAAGGGUCC	1141
AD-1534527	csasgcacgaCfCfUfggcuuguuuL96	614	asAfsadAg(Agn)ggcaggUfcGfucugsgsu	878	ACCAGCAGACCUUGGCUUCUUG	1142
AD-1534528	csasgcuguaCfCfAfcaaauguaucL96	615	asdGsuadCadTuuugdTgUfaccugsgscsc	879	GGCAGCUGUACCCACAAAUUGUACC	1143
AD-1534529	csasgcugugGfCfCfagauccaguuL96	616	asCfsugdGc(Agn)ucuggcCfaCfagucsgscsu	880	AGCAGCUGUGGGCCAGAUCCAGU	1144
AD-1534530	csasgggaacuCfAfAfuaaagugcuuL96	617	asAfsadAc(Tgn)uuuuugAfGfUfucugsgsg	881	CGCAGGAACUCAAUAAAGUGCUU	1145
AD-1534531	csasgggaugGfGfCfaggaacuuL96	618	asUfsgadGu(Tgn)ccugcgCfcAfucugsgsc	882	GCCAGGAUGGGCGCAGGAACUCAA	1146
AD-1534532	csasggcgucUfAfCfaccgauguguL96	619	asdCsacdAudCggugdTgGfaccugsgscsc	883	GCCAGGCGUCUACACCCGAUGUGG	1147
AD-1534533	csasggggacuCfAfUfcuuucccuL96	620	asdGsaadGgdAaagadTgAfucuccugsgasg	884	CUCAGGGACUCAUUUUCCCCUCC	1148
AD-1534534	csasgugcaGfGfGfucugauguuL96	621	asGfscadTc(Agn)ggacccUfuGfcaucugsgsc	885	GCCAGUGCAAGGGUCCUGAUGCC	1149
AD-1534535	csacacaaauGfUfAfccacaaaggguL96	622	asdCscudTgdTgggndAcAfuuuuguggsusa	886	UACCACAAAUUGUACCCACCAAGGG	1150
AD-1534536	csacacaaauGfUfAfccacaaaggguL96	623	asCfscudTg(Tgn)ggguacAfUfuguggsusa	887	UACCACAAAUUGUACCCACCAAGGG	1151
AD-1534537	csasagaugcCfAfGfugcaaggguuL96	624	asdAsecdCudTgcadTgGfcaucugsgscsc	888	GGCCAGAUGCCAGUGCAAGGGUCC	1152
AD-1534538	csasagggcgUfUfAfaccgauguuL96	625	asdAsecdTcdGugudAgAfccugcgscsu	889	AGCCAGGCGUCUACACCCGAUGUG	1153
AD-1534539	csascaagaaAfGfUfgaagaccuuL96	626	asUfsggdTc(Tgn)uucauUfuCfuugggscsu	890	AGCCCAAGAAAGUGAAAGACCAC	1154
AD-1534540	cscccgagacaAfCfGfacaucggcuL96	627	asdGsgcdGgdAugndGuUfucccgsggsusu	891	AACCCGGACAAACGACAUCGCCCC	1155
AD-1534541	cscccgagacaAfCfGfacaucggcuL96	628	asGfsgcdGg(Agn)ugucguUfgUfcccgggsusu	892	AACCCGGACAAACGACAUCGCCCC	1156
AD-1534542	cscccgcaagGfGfCfcaucaucagcuL96	629	asGfseudGa(Tgn)gaugccUfuGfaccgggsusg	893	CACCCUGCAAGGCAUCAUCAGCU	1157

Duplex Name	Sense Strand Sequence 5' to 3'	SEQ ID NO:	Antisense Strand Sequence 5' to 3'	SEQ ID NO:	mRNA target sequence 5' to 3'	SEQ ID NO:
AD-1534543	cscsgaugugGfCfCfuacuaccguL96	630	asdCsagdGudAguagGcCfacuagggusg	894	CACCGAUGUGGCCUACUACCCUGG	1158
AD-1534544	cscsgcaacaAfGfCfagggcgucuuL96	631	asdAsgadCgdCcuagGcuUfuuuagggusg	895	GACCGCAACAAGCCAGGCGUCUA	1159
AD-1534545	cscsgcaagaGfUfCfugucucgaulL96	632	asdTsgadAadGacagAcUfuuuagggusg	896	CUCGCAAGAGUCUUGUCUUCGAU	1160
AD-1534546	cscsggaaceCfGfGfacaacgacauL96	633	asdTsgudCgdTugudCgGfguuuagggusg	897	UGCCGGAACCCGGACAACGACAU	1161
AD-1534547	cscsggaaceCfGfGfacaacgacauL96	634	asUfsgudCg(Tgn)ugucgGfgUfuccggscsa	898	UGCCGGAACCCGGACAACGACAU	1162
AD-1534548	cscsugcaggAfGfGfagggagacuuL96	635	asGfsuadCc(Tgn)ggccuCfcUfagggscsa	899	UUCUUGCAGGAGGCGCAGGUACC	1163
AD-1534549	cscsugucgcCfUfUfagguuacgaulL96	636	asdGscudGadAagudAgGfagggaggsag	900	CUCUUGUGCCUUACGUUCAGCC	1164
AD-1534550	cscsugucgcCfUfUfagguuacgaulL96	637	asGfsuadGa(Agn)cguaagGfcGfagggaggsag	901	CUCUUGUGCCUUACGUUCAGCC	1165
AD-1534551	cscsuiuacguUfCfAfaggggugulL96	638	asdCsacdAcdCggcudGaAfagggaggsag	902	CGCCUUACGUUACGCCGGUGUGC	1166
AD-1534552	csgscaacaaGfCfCfaggggucuuL96	639	asdTsagAcdGcugGcUfuuuagggscsu	903	ACCGCAACAAGCCAGGCGUCUAC	1167
AD-1534553	csgsgcaggaaCfUfCfauuaagugulL96	640	asdCsacdTudTuuuagGfUfuccuagggscsc	904	GGCGCAGGAACUCUCAAUAAAGUGC	1168
AD-1534554	csgsccuuacGfUfUfaggggugulL96	641	asdCsacdCgdGcugAcGfuaagggscsc	905	GUCCGUUACGUUACGCCGGUGU	1169
AD-1534555	csgsccuucgUfCfGfaggggucuuL96	642	asdAsacdGudAagggGcCfagggaggsag	906	CGCGUCCUGUGCCUUACGUUUC	1170
AD-1534556	csgsgaaceGfGfAfcaacgacuuL96	643	asdAsugdTcdGuuudCcGfaggggscsc	907	GCCGGAACCCGGACAACGACAU	1171
AD-1534557	csgsgcuccgCfAfAfaggggucuuL96	644	asGfsuadAg(Agn)cuuagGfGfagggscsc	908	AGCGGUCCGCAAGAGUCUGUCU	1172
AD-1534558	csgsucuacaCfCfGfaggggucuuL96	645	asAfsuadCc(Agn)cuuagGfUfagggscsc	909	GGCGUUAACACCGAUGUGGCCUA	1173
AD-1534559	csgsugccagGfGfUfaggggucuuL96	646	asdTscdGgdAauadCcCfagggaggsag	910	UGCGUGCCAGGGUGAUUCCGGAG	1174
AD-1534560	cscsacuagCfCfGfaggggucuuL96	647	asAfsuadCc(Agn)ggcggGfGfaggggscsg	911	CCCUACAUCGCCCGCGUGUACUG	1175
AD-1534561	cscsaccugGfCfUfaggggucuuL96	648	asdCsacdGgdAuccadGgCfagggaggsa	912	UACUACCUUGGCCUGGAUCCGGGA	1176
AD-1534562	cscscauuaaAfGfUfaggggucuuL96	649	asdTsundCadAagadCuUfuuuagggscsu	913	AACUCAUAAAGUGCUUUUGAAA	1177
AD-1534563	cscscauuaaAfGfUfaggggucuuL96	650	asUfsuadCa(Agn)agacuuUfUfagggscsu	914	AACUCAUAAAGUGCUUUUGAAA	1178
AD-1534564	cscscaaggGfCfCfauuuccuuL96	651	asdGsggdAadAagudAgUfuccuaggsa	915	UGCUCAGGACUCUACUUCUCCU	1179
AD-1534565	cscsccuugGfCfCfauuuccuuL96	652	asdTsagadAadGuaagGcGfagggaggsag	916	CGCUCUUGUGCCUUACGUUCAG	1180
AD-1534566	cscsccuugGfCfCfauuuccuuL96	653	asdCsudCudGaaagGcCfagggaggsag	917	GGCUCUGUUGGCCUUUCAGGAGG	1181
AD-1534567	cscscaaggGfCfCfauuuccuuL96	654	asdAsacdGadCugudTgCfuuuaggsu	918	AGCUGAAAGAGCACACAGUCGUUC	1182

Duplex Name	Sense Strand Sequence 5' to 3'	SEQ ID NO:	Antisense Strand Sequence 5' to 3'	SEQ ID NO:	mRNA target sequence 5' to 3'	SEQ ID NO:
AD-1534568	csusgaungcUfCfAfgggacucauL96	655	asAfsugdAg(Tgn)cccugaGfcAfaucagsgsa	919	UCCUGAUUUGCUCAGGGACUCAUC	1183
AD-1534569	csusgcaaggCfAfUfcaucagcuL96	656	asdCsagdCudGaugdTgCfcuuigcagsgsg	920	CCCUGCAAGGCAUCAUCAGCUGG	1184
AD-1534570	csusgcccgaAfCfCfcggacaacguL96	657	asCfsgudTg(Tgn)ccggguUfcCfgcagsasa	921	UUCUGCCGGAACCCGGACAAACGA	1185
AD-1534571	csusguaccaCfAfAfauguacceauL96	658	asUfsggdGu(Agn)cauuugUfgGfuacagsesu	922	AGCUGUACCACAAAUGUACCCAC	1186
AD-1534572	csusgucgccUfUfAfcgmicagccuL96	659	asGfsggdTg(Agn)acguaaGfgCfgcagsgsa	923	UCCUGUCGCCUUACGUUUCAGCCG	1187
AD-1534573	csusgucucuGfUfCfcaacaacacuL96	660	asGfsgudGu(Tgn)guggacAfGfAfaucagsug	924	CACUGUCUCUGUCCACAAACACCU	1188
AD-1534574	csusgugugugAfCfCfcaacaagcuL96	661	asGfseudTg(Tgn)ugcgguCfaCfcaagscsc	925	GGCUGUGGUGACCGCAACAAGCC	1189
AD-1534575	csusuaeguuCfAfGfcggugugcuL96	662	asdGscadCadCcggedTgAfacguagsgsc	926	GCCUUAACGUUCAGCGCGGUGUGCC	1190
AD-1534576	csusugaugAfCfCfcccugcuuuL96	663	asdAsacdGadCcggedGuCfaucgaagsasc	927	GUCUUCGAUAGACCCGCGUCGUUG	1191
AD-1534577	csusuggaguCfAfAfaucuuuagauL96	664	asdTsegdAadAagudTgAfcucaaagsesu	928	AGCUUGGAGUCAACACUUUCGAU	1192
AD-1534578	csusuuagagcCfUfCfagcuuuccuL96	665	asGfsgadGa(Agn)geugagGfcUfcaaaagsca	929	UGC UUUGAGCCUCAGCUUCUCCG	1193
AD-1534579	gsasaauguAfAfGfaccacugcuL96	666	asUfsgcdAg(Tgn)ggucuuUfcAfcuuuesusu	930	AAGAAAGUGAAAGACCACUGCAG	1194
AD-1534580	gsasacccggAfCfAfaagacaucuuL96	667	asGfsgadTg(Tgn)eguuuGfcGfgguccscg	931	CGGAACCCGGACAACGACAUCGCG	1195
AD-1534581	gsasacuaaaUfAfAfagugcuuuuL96	668	asdCsaadAgdCacuudTaUfugaguucsesu	932	AGGAACUCAAUAAAGUGCUUUGA	1196
AD-1534582	gsasagagcaCfAfCfagucguuucuL96	669	asdAsgadAcgGacugdTgUfgcucucsasg	933	CUGAAGAGCACACAGUCGUGUUCUC	1197
AD-1534583	gsasccgcaaCfAfAfcccagcgcuL96	670	asAfsagdCc(Tgn)ggcuuUfuGfcggucscsc	934	GUGACCGCAACAAGCCAGGCGUC	1198
AD-1534584	gsasgcacacAfGfUfaguuuacuuL96	671	asdGsgudAgdAacgacCuGfugugucsesusu	935	AAGAGCACACAGUCGUGUUCACU	1199
AD-1534585	gsasgcauaaGfUfAfaaagcugauL96	672	asUfscadGc(Tgn)uuuguacUfuAfcucucsesu	936	AGGAGCAUAAAGUACAAAAGCUGAA	1200
AD-1534586	gsasgcccuaGfCfUfucuccgguuuL96	673	asdAsacdCgdGagaadGcUfagggucsesasa	937	UUUGAGCCUCAGCUUCUCCGGUUC	1201
AD-1534587	gsasgucuguCfUfUfagauagaccuuL96	674	asGfsggdTc(Agn)ucgaagAfcAfcucucsesu	938	AAGAGUCUGUCUUCGAUGACCCG	1202
AD-1534588	gsasugcccagUfGfCfaaggguuuccuL96	675	asdAsggdAdCcuugdCaCfugguacsesug	939	CAGAUCCAGUGCAAGGGUCCUG	1203
AD-1534589	gsasuggcgcAfGfGfaacucaauuL96	676	asUfsaudTg(Agn)guuucuGfcGfcaucsesu	940	AGGAUGGCGCAGGAACUCAAUAA	1204
AD-1534590	gsasuuugcucAfGfGfagacucauuuL96	677	asAfsagdTg(Agn)guccuuGfaGfcaucsesag	941	CUGAUUGUCUCAGGGACUCAUCU	1205
AD-1534591	gsascaagaguCfUfGfucuuuaguuL96	678	asdCsaudCgdAagacAdAgAfcuucugscsg	942	CCGCAAGAGUCUGUCUUCGAUGA	1206
AD-1534592	gscsacacagUfCfGfuuucacugcuL96	679	asCfsagdTg(Agn)gaacgaCfuGfugugscsc	943	GAGCACACAGUCGUUCUCACUGU	1207

Duplex Name	Sense Strand Sequence 5' to 3'	SEQ ID NO:	Antisense Strand Sequence 5' to 3'	SEQ ID NO:	mRNA target sequence 5' to 3'	SEQ ID NO:
AD-1534593	gscsacgaccUfGfGfcucuguuL96	680	asdGscadAcAdAagedCaGfncuguesug	944	CAGCACGACCCUGGCUUUGCG	1208
AD-1534594	gscsacgaccUfGfGfcucuguuL96	681	asGfscadAc(Agn)gagccaGfGfUfcuguesug	945	CAGCACGACCCUGGCUUUGCG	1209
AD-1534595	gscsacgaccUfGfGfCfagauL96	682	asUfsgdCa(Tgn)uuggccAfcAfcuguesu	946	AAGCAGCUGUGGCCAGAUCCAG	1210
AD-1534596	gscsacgaccUfCfAfauaaaguuL96	683	asdGscadCudTtuudGaGfucucugsc	947	GCGCAGGAACUCAAUAAAGUCU	1211
AD-1534597	gscsacgaccUfCfAfugcaaguuL96	684	asdCscadTudGeacudGgCfauucgscscsa	948	UGGCCAGAUGCCAGUGCAAGGU	1212
AD-1534598	gscsacgaccUfCfUfacaccuuL96	685	asdCsaudCgdGugadGaCfcccugscsu	949	AAGCAGGCGUUCACACCGAUGU	1213
AD-1534599	gscsacgaccUfCfUfacaccuuL96	686	asGfsgdCu(Tgn)uacuuUfcUfugscsc	950	GAGCCCAAGAAAGUGAAAGACCA	1214
AD-1534600	gscsacgaccUfCfUfacaccuuL96	687	asdAscadCcdGgcudAaCfuaagscscsa	951	UCGCCUACGUUACAGCCGGUGUG	1215
AD-1534601	gscsacgaccUfCfUfacaccuuL96	688	asdAscudTudAungadGuUfcucugscsu	952	UGGCGCAGGAACUCAAUAAAGUG	1216
AD-1534602	gscsacgaccUfCfUfacaccuuL96	689	asdCscgdCadTccudCuGfagagscscsa	953	UUGCGCCUUCAGGAGGAUGCGGA	1217
AD-1534603	gscsacgaccUfCfUfacaccuuL96	690	asdAscadGadCucudGcGfagagccscsu	954	CAGCGCCUCCGCAAGAGUCUCUC	1218
AD-1534604	gscsacgaccUfCfUfacaccuuL96	691	asdGsgadAadGaugadGuCfcccugscsa	955	UUGCUCAGGACUCACUCUUCUCCC	1219
AD-1534605	gscsacgaccUfCfUfacaccuuL96	692	asAfsadAc(Agn)gacuuUfcCfagagscsc	956	CGGCUCGCAAGAGUCUCUCUUC	1220
AD-1534606	gscsacgaccUfCfUfacaccuuL96	693	asdGsaadCgdTaaagdCgAfcagagscsc	957	GCGCUCUUGUGCCUACGUUCA	1221
AD-1534607	gscsacgaccUfCfUfacaccuuL96	694	asdCsgadGgdAaccddTgCfagagscscsu	958	AUGCUCUGCGCAGGCUUCCUGA	1222
AD-1534608	gscsacgaccUfCfUfacaccuuL96	695	asCfsgadGg(Agn)accuuCfCfagagscsu	959	AUGCUCUGCGCAGGCUUCCUGA	1223
AD-1534609	gscsacgaccUfCfUfacaccuuL96	696	asdTscddTgdAagadGcAfcagagscscsa	960	UGGCUCUUGUGCCUUCAGGAG	1224
AD-1534610	gscsacgaccUfCfUfacaccuuL96	697	asUfscddTg(Agn)agggcAfaCfagagscsa	961	UGGCUCUUGUGCCUUCAGGAG	1225
AD-1534611	gscsacgaccUfCfUfacaccuuL96	698	asdAscgdAcdTgugudGcUfcuucagcsu	962	AAGCUGAAGAGCACACAGUCGUU	1226
AD-1534612	gscsacgaccUfCfUfacaccuuL96	699	asdCscddGedGccadTcAfuagcagcsu	963	AAGCUGCUAUGAUGGCCCGCGGC	1227
AD-1534613	gscsacgaccUfCfUfacaccuuL96	700	asCfsuudGg(Tgn)ccuacAfcAfcagagscsc	964	CCGCUGGUGUGAGGACCAAGC	1228
AD-1534614	gscsacgaccUfCfUfacaccuuL96	701	asdCsuudGudTgagddTcAfcuagcagcsu	965	GGGCUGUGGUGACCCGCAACAAGC	1229
AD-1534615	gscsacgaccUfCfUfacaccuuL96	702	asCfsuudGu(Tgn)gagcAfcCfagagscsc	966	GGGCUGUGGUGACCCGCAACAAGC	1230
AD-1534616	gscsacgaccUfCfUfacaccuuL96	703	asGfscadTg(Agn)gugggAfaUfagagcsu	967	AGGCUCUACUUGUCCACUCAUGCC	1231
AD-1534617	gscsacgaccUfCfUfacaccuuL96	704	asdGsaadAadGeugadGgCfuaaagcsu	968	GUGCUUUGAGCCUACAGCUUCUCC	1232

Duplex Name	Sense Strand Sequence 5' to 3'	SEQ ID NO:	Antisense Strand Sequence 5' to 3'	SEQ ID NO:	mRNA target sequence 5' to 3'	SEQ ID NO:
AD-1534618	gsgsaacuaAfUfAfaagugcuuuL96	705	asAfsaadGc(Agn)cuuuuUfgAfguuccsug	969	CAGGAACUCAAUAAAAGUGCUUUG	1233
AD-1534619	gsgsagcauaAfGfUfacaagcugL96	706	asdCsagdCudTugnadCuUfauguccsusu	970	AAGGAGCAUAAGUACAAAAGCUGA	1234
AD-1534620	gsgsagcccaAfGfAfaagugaaagL96	707	asCfsuudTc(Agn)cuuuuUfgGfguccesasa	971	UUGGAGCCCAAGAAAGUGAAAAGA	1235
AD-1534621	gsgsagucuaAfAfCfuuucgaucL96	708	asdGsaadTcdGaaagdTgUfugacuccsasa	972	UUGGAGUCAACACUUUCGAUUC	1236
AD-1534622	gsgscgucuaAfAfCfcauguggcL96	709	asdGscddAcdaucggdTgUfagacgcsusg	973	CAGGCGUCUACACCGAUGUGGCC	1237
AD-1534623	gsgscgucuaAfAfCfcauguggcL96	710	asGfscddAc(Agn)ucgggUfaGfagcccsug	974	CAGGCGUCUACACCGAUGUGGCC	1238
AD-1534624	gsgscuccgcAfAfGfagucugucL96	711	asdAsgadCadGacudTuGfcggagccsusc	975	GCGGCUCCGCAAGAGUCUCUGUCU	1239
AD-1534625	gsgscucugUfGfCfcccucaggL96	712	asdCscudGadAagggdCaAfcaagcccsag	976	CUGGCUUGUUGCGCCUUCAGGA	1240
AD-1534626	gsgsguccugAfUfGfcccacugccL96	713	asGfsgddAg(Tgn)gggcauCfaGfagcccsu	977	AAGGUCUUGAUGGCCCCACUGCCA	1241
AD-1534627	gsgscaacaeUfUfUfcauguccacL96	714	asdGsugdGadAucgadAaGfuguuagcsusc	978	GAGUCAACACUUUCGAUUCACC	1242
AD-1534628	gsgscaacaeUfUfUfcauguccacL96	715	asGfsgddGa(Agn)ucgaaaGfuGfuagcsusc	979	GAGUCAACACUUUCGAUUCACC	1243
AD-1534629	gsgscgcccAfAfGfucagccggL96	716	asdCscgdGcdTgaacdGuAfagcgcccsag	980	CUGUCGCCUACGUUUCAGCCGGU	1244
AD-1534630	gsgscgcccAfAfGfucagccggL96	717	asCfsgddGc(Tgn)gaacguAfaGfagcccsag	981	CUGUCGCCUACGUUUCAGCCGGU	1245
AD-1534631	gsgscgcccAfAfCfugacccggL96	718	asdCscgdGudGacagdTgAfgaacgcsusc	982	CAGUCGUUCUACUUGUCACCCGGG	1246
AD-1534632	gsgscgcccUfCfGfagaccggL96	719	asGfscddGg(Tgn)caucgaAfgAfgaacgcsusc	983	GAGUCGUUCUUCGAUGACCCCGG	1247
AD-1534633	gsgscuccgaUfGfAfcccgcgucL96	720	asdCsgadCgdCgggudCaUfcaagcccsag	984	CUGUCUUCGAUGACCCCGGUCGU	1248
AD-1534634	gsgscaaggGfUfCfcaugagccL96	721	asGfsgddCa(Tgn)caagcaCfcUfagcccsusc	985	CAGUGCAAGGGUCCUGAUGCCCA	1249
AD-1534635	gsgsggugacCfGfCfaacaagccL96	722	asUfsgddCu(Tgn)guugcgGfuCfaccacsag	986	CUGUGGUGACCCGCAACAAGCCAG	1250
AD-1534636	gsgsgcggcUfUfCfagagggL96	723	asdCsandCcdTccugdAaGfagcccsag	987	CUGUUGCGCCUUCAGGAGGAUGC	1251
AD-1534637	gsgsgcggcUfUfCfagagggL96	724	asCfsandCc(Tgn)ccugaaGfgCfagcccsag	988	CUGUUGCGCCUUCAGGAGGAUGC	1252
AD-1534638	usasaguacaAfAfGfcaagagL96	725	asdGscudCudTcagdTuUfagcuuucL96	989	CAUAAAGUACAAAAGCUGAAGAGCA	1253
AD-1534639	usasaguacaAfAfGfcaagagL96	726	asGfscudCu(Tgn)caagcuUfUfagcuuucL96	990	CAUAAAGUACAAAAGCUGAAGAGCA	1254
AD-1534640	usascaucgcCfGfCfcaugagL96	727	asdCsagdTadCagagdCgGfcaugagL96	991	CCUACAUCGCCGCGCUGUACUGG	1255
AD-1534641	usasccuggcCfUfGfcauccggL96	728	asdTscddCgdGaucdAgGfcaagagL96	992	ACUACUUGGCCUGGAUCCGGGAG	1256
AD-1534642	usascuaccuGfGfCfcaugagL96	729	asdCsggdAudCaggdCcAfggagagL96	993	CCUACUACUUGGCCUGGAUCCGGG	1257

Duplex Name	Sense Strand Sequence 5' to 3'	SEQ ID NO:	Antisense Strand Sequence 5' to 3'	SEQ ID NO:	mRNA target sequence 5' to 3'	SEQ ID NO:
AD-1534643	uscsaacacuUfUfCfgaunuccacuuL96	730	asGfsugdGg(Agn)aucgaaAfgUfguugascsu	994	AGUCAAACAUUUUCGAUUCACCU	1258
AD-1534644	uscsaccucgCfAfAfgcaucauL96	731	asGfsaudGa(Tgn)gccuugCfaGfguugasgsc	995	GCUCACCCUGCAAGGCAUCAUCA	1259
AD-1534645	uscsagccggUfGfUfgecuuL96	732	asUfsugdGc(Agn)ggcacaCfcGfgeugasasc	996	GUUCAGCCGGUGUGCCUGCCAAAG	1260
AD-1534646	uscsagggaggAfUfGfGgagcgcauL96	733	asUfsugdGg(Tgn)ccgcauCfcUfgeugasag	997	CUUCAGGAGGAUGCGGACGGCAG	1261
AD-1534647	uscsaggggacUfCfAfucunuccuuL96	734	asdAsggdGadAagaudGaGfuccuugasgsc	998	GCUCAGGGACUCACUCUUUCCUC	1262
AD-1534648	uscsagcaagAfGfUfgecuuL96	735	asdCsgadAgdAcagadCuCfuugcgasgsc	999	GCUCCGCAAGAGUCUGUCUUCGA	1263
AD-1534649	uscsuugucCfCfUfuauguncaguL96	736	asdCsugdAadCguadGgCfagacagagasc	1000	GCUCUGUCGGCUUACGUUCAGC	1264
AD-1534650	uscsGCCUUAfGfUfucagcgguuL96	737	asdAsgcdGgdCugaadCgUfaagggcasca	1001	UGUCGCCUUAACGUUCAGCGCGUG	1265
AD-1534651	uscsuugucCfAfCfaacaccuL96	738	asUfsugdGg(Tgn)guuugGfaCfagagascsa	1002	UGUCUGUCCACACACACCUAC	1266
AD-1534652	uscsuugucGfCfCfuaagagagguL96	739	asdCsudCcdTgaagdGcGfcaacagagasc	1003	GCUCUGUCCGCCUUCAGGAGGA	1267
AD-1534653	usgsaagagcAfCfAfagucguuL96	740	asdGsaadCgdAeugndCuGfucucucagasc	1004	GCUGAAGAGCACACACAGUCGUUCU	1268
AD-1534654	usgsagccucAfGfCfucucggguL96	741	asdAsgcdGgdAgaadCuGfagggcasasa	1005	UUUGAGCCUCAGCUUCUCCGGU	1269
AD-1534655	usgsagcugGfAfGfuaucgagcuL96	742	asGfsudGc(Agn)guacucCfcAfgeucagasc	1006	GCUGAGCUGGGAGUACUGCGACC	1270
AD-1534656	usgsauggccGfCfGfgeucagcuL96	743	asAfsugdGg(Agn)ggcgcGfGfcaucacusa	1007	UAUGAUGGCCCGCGGGCUCAGCUA	1271
AD-1534657	usgscaaggcAfUfCfaucagcugguL96	744	asdCscadGcdTgaugdAuGfucucagagsg	1008	CCUGCAAGGCAUCAUCAGCUGGG	1272
AD-1534658	usgscaaggcAfUfCfaucagcugguL96	745	asCfsadGc(Tgn)gaugauGfCfuugcasgsg	1009	CCUGCAAGGCAUCAUCAGCUGGG	1273
AD-1534659	usgscaaggcUfCfCfuaugccccuL96	746	asUfsugdGc(Agn)ucaagGcCfuugcasasu	1010	AGUGCAAGGGUCCUGAUGCCCCAC	1274
AD-1534660	usgsccagugCfAfAfaggguccuL96	747	asUfsadGg(Agn)ccuugCfaCfugggcasusc	1011	GAUGCCAGUGCAAGGGUCCUGAU	1275
AD-1534661	usgscaugaGfGfGfucucugcuL96	748	asGfsadGc(Agn)ggcccUfcAfugggcasusc	1012	GAUGCCAUGAGGGCUCUCUGCU	1276
AD-1534662	usgsccuucCfAfGfaggaugcgguL96	749	asdCsgcdAudCcuudTgAfagggcasasc	1013	GUUGCGCCUUCAGGAGGGAUGCGG	1277
AD-1534663	usgsuucaggGfAfCfuaucuuuL96	750	asdGsaadAgdAugagdTcCfugagcasasu	1014	AUUGCUCAGGGACUCACUUCUCC	1278
AD-1534664	usgsuucaggGfAfCfuaucuuuL96	751	asGfsaadAg(Agn)ugagucCfcUfagagcasasu	1015	AUUGCUCAGGGACUCACUUCUCC	1279
AD-1534665	usgsagaucaAfCfAfucunuccuuL96	752	asdAasudCgdAaagudGuUfagucacasag	1016	CUUGGAGUCAAACACUUCGAUUC	1280
AD-1534666	usgsagcagaUfGfCfcaugcaaguL96	753	asCfsudGc(Agn)cuugcaUfcUfgeccasca	1017	UGUGGCCAGAUGCCAGUGCAAGG	1281
AD-1534667	usgsGCCUUAfAfCfucggcgguL96	754	asdCscadGgdCcaggdTgGfuaugggcasca	1018	UGUGGCCUACUACCUUGGCCUGGA	1282

Duplex Name	Sense Strand Sequence 5' to 3'	SEQ ID NO:	Antisense Strand Sequence 5' to 3'	SEQ ID NO:	mRNA target sequence 5' to 3'	SEQ ID NO:
AD-1534668	usgsugcucugUfUfGfGfGcGcUcagU96	755	asdCsugdAadGgGcgGdAaCfagagGccasgsg	1019	CCUGGCUCUGUUGCGCCUUCAGG	1283
AD-1534669	usgsugagaccGfCfAfacaagccagU96	756	asCfsugdGc(Tgn)uguuGcGfGfUfaccascsa	1020	UGUGGUGACCGCAACAAGCCAGG	1284
AD-1534670	usgsuaccacAfAfAfuguaaccacU96	757	asGfsugdGg(Tgn)acuuuGfuGfuaacasgsc	1021	GCUGUACCAACAAGUACCCACA	1285
AD-1534671	usgsucucugUfCfCfacaacacuuL96	758	asAfsugdGg(Tgn)uguggaCfaGfagacagssu	1022	ACUGUCUCUGUCCACAACACCUC	1286
AD-1534672	usgsuggcccaGfAfUfGccagugcauL96	759	asUfsgdAc(Tgn)ggcaucUfGfGccacagssc	1023	GCUGUGGCCAGAUGCCAGUGCAA	1287
AD-1534673	usugagcucAfGfCfGgugugccuL96	760	asGfsgdAc(Tgn)ccggcuGfaAfcguaasgsg	1024	CCUUACGUUCAGCCGGUGUGCCU	1288
AD-1534674	usugagcucCfCfAfGfGcGcUcagU96	761	asGfsgdCa(Tgn)gguggGfaCfauaasgsc	1025	GCUUAUGUCCCAUCUAUGCCCG	1289
AD-1534675	usugagcucCfCfAfGfGcGcUcagU96	762	asdCsaadCgdAegcgGgUfcaucgaasgsa	1026	UCUUCGAUGACCCCGCUGCUGUGG	1290
AD-1534676	usugagcucCfAfGfGcUcagUcagU96	763	asdCscgdGadGaagcdTgAfggcucaasag	1027	CUUUGAGCCUACGCUUCUCCCGU	1291
AD-1534677	usugagcucCfAfAfGfGcUcagUcagU96	764	asUfsudAc(Tgn)uuuGgGfGfCfuecaasac	1028	GUUUGGAGCCCAAGAAAGUGAAA	1292
AD-1534678	usugggagucAfAfCfGfGcUcagUcagU96	765	asdAsudCgdAagugdTgGfGfGcUcagUcagU96	1029	GCUUGGAGUCAACACUUCGGAU	1293
AD-1534679	usugggagucUfCfAfGfGcUcagUcagU96	766	asdCsggdAgdAagcdGgGfGfGcUcagUcagU96	1030	GCUUUGAGCCUCAGCUUCUCCCGG	1294
AD-1631272	asagagcaccAfCfAfGfGcUcagUcagU96	767	asdGsagdAadCgacudGuGfGfGcUcagUcagU96	1031	UGAAGAGCACACAGUCGUUCUCA	1295
AD-1631273	asasuaaagUfCfUfGfGcUcagUcagU96	768	asdCsandTudTcaadGcAfGfGcUcagUcagU96	1032	UCAAUAAAGUGCUUUGAAAAUGC	1296
AD-1631274	ascsaagcgcGfCfGfGcUcagUcagU96	769	asdCscadGudAcagcdGcGfGcGgagugscsu	1033	CUACAUCGCCGCGCUGUACUGGG	1297
AD-1631275	ascsaagcgcGfAfCfGfGcUcagUcagU96	770	asdCsagdAgdCagcdTcGfGfGcUcagUcagU96	1034	CUACCAAGCAGACCUUGGCUCUGU	1298
AD-1631276	ascsaagcgcAfGfGfGcUcagUcagU96	771	asdCsugdAudGaugcdCuUfGfGcUcagUcagU96	1035	UCACCCUGCAAGGCAUCAUCAGC	1299
AD-1631277	ascsaagcgcAfAfGfGcUcagUcagU96	772	asdGsacdGcdCugcdTgGfGfGcUcagUcagU96	1036	UGACCCGAACAAGCCAGGCGUCU	1300
AD-1631278	ascsaagcgcGfCfGfGcUcagUcagU96	773	asdCscgdGadTccagcdGcCfGfGcUcagUcagU96	1037	CUACUACCUUGGCCUGGAUCCCGGG	1301
AD-1631279	asusugcuaGfGfGfGcUcagUcagU96	774	asdAsagdAudGaugcdCcUfGfGcUcagUcagU96	1038	UGAUUGCUCAGGGACUCAUCUUU	1302
AD-1631280	csasgggagcuCfAfUfGfGcUcagUcagU96	775	asdGsagdGcdAaagdTgAfggcucaasgsu	1039	CUCAGGGACUCUUCUUUCCCUCC	1303
AD-1631281	csasaaagUfUfAfGfGcUcagUcagU96	776	asdCsandTgdTgggudAcAfGfGcUcagUcagU96	1040	UACCAACAAGUACCCACAAGGG	1304
AD-1631282	csasgaaagUfUfCfGfGcUcagUcagU96	777	asdTscgdAadGacagcdAcUfGfGcUcagUcagU96	1041	CUCCGCAAGAGUCUUCUUCGAU	1305
AD-1631283	csasgaaagCfGfGfGcUcagUcagU96	778	asdTsgudCgdTugcdCgGfGfGcUcagUcagU96	1042	UGCCGGAACCCGGACAACGACAU	1306
AD-1631284	csasugcgcCfUfUfGfGcUcagUcagU96	779	asdGscudGadAcagudAgGfGfGcUcagUcagU96	1043	CUCUCUGCGCUUACGUUCAGCC	1307

Duplex Name	Sense Strand Sequence 5' to 3'	SEQ ID NO:	Antisense Strand Sequence 5' to 3'	SEQ ID NO:	mRNA target sequence 5' to 3'	SEQ ID NO:
AD-1631285	csqscuuuacGfUfUfagccggguL96	780	asdCsacdCgdGeugadAcGfuuaagcgscsu	1044	GUCGCCUUACGUUACAGCCGGUGU	1308
AD-1631286	csqsguccagGfGfUfgauuacggauL96	781	asdTsscdGgdAueadCcCfugcgacscsu	1045	UGCGUGCCAGGGUGAUUCCGGAG	1309
AD-1631287	csusaccuugCfCfUfgauuacggguL96	782	asdCsccdGgdAueadGgCfcaaguuagscsu	1046	UACUACCUUGGCCUGGAUCCGGGA	1310
AD-1631288	csusacggggaCfUfCfauuuuuccuL96	783	asdGsggdAadAgaugdAgUfcccugagscsu	1047	UGCUCAGGGACUCAUUUUCCCU	1311
AD-1631289	csusucgaugAfCfCfcccgcguuuL96	784	asdAsacdGadCgCgggdGuCfaucgaagscsu	1048	GUCUUCGAUGACCCGCGUCGUUG	1312
AD-1631290	gsasagagcaCfAfCfagucguuuuL96	785	asdAsgadAcidGacugdTgUfgcucuucescu	1049	CUGAAGAGCACACAGUCGUUCUC	1313
AD-1631291	gsasgcucaGfCfUfucuceggguuL96	786	asdAsacdCgdGagaadGcUfgaggucescu	1050	UUGAGCCUCAGCUUUCUCCGGUUU	1314
AD-1631292	gscscgaugCfCfAfugcaagggguL96	787	asdCsccdTudGeacudGgCfaucggcgscsu	1051	UGGCCAGAUGCCAGUGCAAGGGU	1315
AD-1631293	gscscuuacGfUfCfagccggguuL96	788	asdAscacdCcdGgcugdAaCfuaagcgscsu	1052	UCGCCUUACGUUACAGCCGGUGUG	1316
AD-1631294	gscsgcaggAfcUfcauuuaaguuL96	789	asdAseudTudAuugadGuUfcccugcgscsu	1053	UGGCGCAGGAACUCACAAUAAAGUG	1317
AD-1631295	gscsgccuucAfGfGfaggaugcgguL96	790	asdCsccdCcdTccudCuGfaagggcgscsu	1054	UUGCGCCUUCAGGAGGAUGCGGA	1318
AD-1631296	gscsucaggAfcUfcauuuuccuL96	791	asdGsagdAadGaugadGuCfcccugcgscsu	1055	UUGCUCAGGGACUCACUUCUUUCCC	1319
AD-1631297	gscsucugcgCfAfGfguuuccugnuL96	792	asdCsagdGgdAaccddTgCfcaagagcscsu	1056	AUGCUCUGCGCAGGGUUCUCCGA	1320
AD-1631298	gscsucuguuGfCfGfcccuaagguL96	793	asdTsscdTgdAagggdGcAfacagagcscsu	1057	UGGCUCUGUUGCGCCUUCAGGAG	1321
AD-1631299	gscsuuugagCfCfUfagcuuuccuL96	794	asdGsagdAadGeugadGgCfuaaagcscsu	1058	GUGCUUUGAGCCUCAGCUUCUCC	1322
AD-1631300	gsgsagucaaCfAfCfuuuuaguuuL96	795	asdGsaadTcdGaaagdTgUfugacuccscsu	1059	UUGGAGUCAACACUUCUUGAUUCC	1323
AD-1631301	gsgscueuiguUfGfCfcccuaagguL96	796	asdCsacdGadAagcgdCaAfcagagcscsu	1060	CUGGCUCUGUUGCGCCUUCAGGA	1324
AD-1631302	gsuscgccuuAfCfGfucagccggguL96	797	asdCsagdGcdTgaacdGuAfagggcgscsu	1061	CUGUCGCCUUACGUUACAGCCGGU	1325
AD-1631303	gsuscuuigaUfGfAfcccgcguuL96	798	asdCsagdGgdCgggndCaUfcaagagcscsu	1062	CUGUCUUCGAUGACCCGCGUCGU	1326
AD-1631304	gsusugcgccUfUfCfagagagguuL96	799	asdCsacdCcdTccugdAaGfgecgacscsu	1063	CUGUUGCGCCUUCAGGAGGAUGC	1327
AD-1631305	uscsgecuuaCfGfUfucagccggguL96	800	asdAsccdGgdCuugaadCgUfaagggcgscsu	1064	UGUCGCCUUACGUUACAGCCGGUG	1328
AD-1631306	usgsagccucAfGfCfuuuuccggguL96	801	asdAsccdGgdAgaadCuGfagggcscsu	1065	UUUGAGCCUCAGCUUUCUCCGGUU	1329
AD-1631307	usgscccuuAfGfGfagagguuL96	802	asdCsagdAudCuccdTgAfagggcgscsu	1066	GUUGCGCCUUCAGGAGGAUGCGG	1330
AD-1631308	usgsuacaggGfAfCfucuuuuuL96	803	asdGsaadAadAagagdTcCfugagcscsu	1067	AUUGCUCAGGGACUCAUUUUCC	1331
AD-1631309	usgsagucuaAfCfAfccuuuaguuuL96	804	asdAsacdCgdAaagudGuUfaguccscsu	1068	CUUGGAGUCAACACAUUUUCGAUUC	1332

Duplex Name	Sense Strand Sequence 5' to 3'	SEQ ID NO:	Antisense Strand Sequence 5' to 3'	SEQ ID NO:	mRNA target sequence 5' to 3'	SEQ ID NO:
AD-1631310	usgsgccuacUfaAfCfcugggccugguL96	805	asdCscadGgdCcaggdTgGfuaggccascsu	1069	UGUGGCCUACUACCUAGGCCUGGA	1333
AD-1631311	ususcgaugaCfCfCfcgucgugguL96	806	asdCsaadCgdAcggdGgUfcaucgaascsu	1070	UCUUCGAUGACCCGCGUCGUUGG	1334
AD-1631312	usugagccuCfAfGfcuucuccggguL96	807	asdCsegdGadGaagdTgAfggcucuaascsu	1071	CUUUGAGCCUCAGCUUCUCCCGGU	1335
AD-85525	ascsucaaUfaAfAfGfugcuuuuuaaL96	808	usUfsucaaaagcacuuUfaUfugagusc	1072	GAACUCAAUAAAAGUGCUUUUGAAA	1336
AD-85531	uscuaaAfaGfUfGfcuuuuaaauL96	809	asUfsuuucaaaagcacUfuUfaungasgsu	1073	ACUCAAUAAAAGUGCUUUUGAAAAU	1337
AD-85537	csuscaauAfaAfGfUfGfcuuuuaaL96	810	usUfsuucuaagcacuuUfuAfuungasgsu	1074	AACUCAAUAAAAGUGCUUUUGAAAA	1338
AD-85541	gsasacucAfaUfAfAfagugcuuuuuaL96	811	usCfsaaagcacuuuaUfuGfagucscsu	1075	AGGAACUCAAUAAAAGUGCUUUUGA	1339
AD-67246	asasuuaaGfuGfCfUfuugaauaacguL96	812	asCfsguuUfuCfAfaagcAfcUfuuaungsa	1076	UCAATAAAAGTGCTTTGAAAA	1340

Table 4. Single Dose Screen in PHH Cells

Duplex ID	10 nM	ST DEVP	1 nM	ST DEVP	0.1 nM	ST DEVP
AD-1534661.1	44.05	4.93	69.18	4.10	85.65	8.36
AD-1534577.1	12.11	1.61	24.85	1.11	28.45	0.69
AD-1534678.1	10.93	1.19	22.47	2.79	26.15	3.02
AD-1534665.1	5.73	0.32	14.01	0.67	21.39	1.31
AD-1631309.1	7.03	0.66	14.67	0.88	21.63	2.28
AD-1534621.1	6.70	0.50	17.96	4.44	17.38	0.96
AD-1631300.1	10.06	1.77	17.60	1.48	24.79	2.03
AD-1534506.1	6.24	0.92	13.38	1.17	20.59	0.68
AD-1534507.1	10.74	0.98	15.73	1.15	24.57	3.14
AD-1534627.1	25.35	2.73	43.05	3.27	60.05	4.14
AD-1534628.1	41.16	3.47	63.30	4.02	73.10	3.82
AD-1534643.1	59.52	1.86	79.54	3.61	74.34	4.06
AD-1534477.1	9.74	1.31	21.58	1.03	33.46	3.63
AD-1534478.1	49.87	2.04	63.36	2.03	85.92	4.39
AD-1534525.1	25.71	2.68	48.60	7.18	60.80	3.95
AD-1534519.1	67.15	5.73	75.69	8.82	88.37	5.21
AD-1534501.1	7.52	1.75	15.11	1.03	22.92	1.42
AD-1534502.1	8.00	0.87	15.90	1.77	26.14	2.65
AD-1534619.1	11.28	0.97	25.24	2.79	30.78	3.45
AD-1534585.1	11.34	0.38	21.88	1.41	27.59	3.05
AD-1534638.1	17.72	1.81	27.67	0.63	33.57	2.93
AD-1534639.1	20.54	3.14	29.81	1.93	37.26	6.55
AD-1534471.1	6.51	0.69	12.63	1.00	22.48	2.27
AD-1534515.1	16.51	1.81	32.66	10.43	32.77	3.57
AD-1534498.1	10.26	1.63	19.40	2.23	32.84	4.14
AD-1534611.1	5.59	0.50	14.54	2.13	20.10	1.53
AD-1534567.1	3.99	0.33	11.49	2.11	13.54	1.29
AD-1534653.1	12.81	1.20	21.70	1.26	32.04	0.85
AD-1534582.1	4.41	0.68	15.00	3.15	18.57	2.10
AD-1631290.1	7.40	0.51	16.85	0.91	24.34	1.01
AD-1534468.1	12.06	1.11	25.04	1.77	34.73	3.53
AD-1631272.1	21.96	0.94	33.00	3.29	37.18	2.22
AD-1534491.1	5.19	0.37	13.84	1.96	21.60	1.19
AD-1534584.1	10.01	0.57	26.72	4.03	34.05	4.78
AD-1534475.1	34.84	2.54	55.91	4.77	80.78	1.98
AD-1534479.1	5.79	0.57	12.45	0.83	20.10	1.83
AD-1534631.1	9.40	1.46	18.99	0.61	29.08	2.89
AD-1534528.1	5.96	0.99	11.36	0.79	18.28	1.72
AD-1534500.1	54.63	8.31	61.60	0.80	78.36	3.61
AD-1534571.1	7.52	1.01	17.48	3.54	20.30	1.81
AD-1534670.1	34.73	3.06	44.58	6.06	52.88	1.88
AD-1534535.1	28.24	0.99	43.88	2.96	63.88	3.07
AD-1631281.1	49.87	6.19	69.66	4.09	82.29	8.76
AD-1534536.1	85.33	11.49	81.33	3.08	90.47	3.99
AD-1534520.1	54.45	4.62	64.37	4.70	78.68	15.12
AD-1534474.1	47.54	4.63	75.38	3.44	88.71	13.21

Duplex ID	10 nM	ST DEVP	1 nM	ST DEVP	0.1 nM	ST DEVP
AD-1534516.1	39.65	4.68	59.02	2.12	75.27	2.71
AD-1534463.1	69.01	3.89	53.48	2.10	63.99	6.25
AD-1534513.1	83.25	15.32	77.18	16.79	81.45	9.40
AD-1534677.1	88.76	4.69	76.94	2.75	86.09	3.80
AD-1534620.1	40.56	4.37	52.67	3.28	61.12	8.69
AD-1534497.1	20.56	1.70	39.51	2.12	61.93	3.33
AD-1534599.1	22.24	1.26	33.58	1.07	35.83	1.27
AD-1534539.1	15.85	1.61	28.76	6.80	35.78	2.40
AD-1534517.1	17.72	1.90	23.72	2.15	33.03	0.61
AD-1534579.1	16.26	2.41	24.17	3.87	28.24	3.91
AD-1534524.1	10.59	0.91	16.19	2.05	30.88	3.69
AD-1534573.1	31.99	3.60	47.90	1.48	46.22	5.02
AD-1534671.1	43.31	2.22	53.10	5.75	71.83	10.58
AD-1534651.1	70.56	7.43	64.51	4.91	67.31	4.94
AD-85525.11	14.68	2.89	25.29	0.71	29.25	2.79
AD-1534462.1	14.29	0.98	26.48	2.15	37.09	3.81
AD-1534490.1	11.99	0.21	25.71	1.39	39.22	2.55
AD-1534472.1	17.33	0.82	27.11	1.94	46.36	2.38
AD-1534617.1	23.19	1.41	42.01	11.05	54.20	5.66
AD-1631299.1	24.00	2.03	33.21	3.53	44.13	2.46
AD-1534578.1	76.95	9.07	96.34	10.63	86.34	4.96
AD-1534679.1	48.99	2.74	70.38	4.13	83.54	2.17
AD-1631312.1	32.27	2.68	45.09	5.43	57.26	1.81
AD-1534676.1	33.71	2.91	45.46	5.28	56.88	5.20
AD-1631306.1	23.23	0.77	32.22	2.00	42.10	5.65
AD-1534654.1	24.93	2.14	40.86	4.67	51.55	3.84
AD-1534586.1	9.45	1.12	20.64	3.59	27.92	1.46
AD-1631291.1	11.51	1.14	20.31	2.19	28.60	2.23
AD-1534467.1	5.30	0.63	12.78	0.40	20.64	1.07
AD-1534466.1	26.06	0.15	45.65	3.89	60.77	6.19
AD-1534489.1	33.87	3.66	66.95	5.67	87.84	9.58
AD-1534469.1	55.57	4.56	68.07	7.11	89.20	6.12
AD-1534595.1	55.23	2.75	69.66	7.29	70.81	6.57
AD-1534529.1	101.12	7.09	87.33	3.56	102.13	4.30
AD-1534672.1	72.47	5.94	64.77	3.99	83.59	3.66
AD-1534666.1	63.91	6.67	76.11	4.79	86.79	5.64
AD-1534597.1	49.56	3.29	89.70	12.87	96.87	8.59
AD-1631292.1	51.62	7.41	71.35	3.78	86.68	3.55
AD-1534537.1	29.30	2.87	40.25	4.60	62.01	2.78
AD-1534526.1	60.24	4.43	95.64	5.54	110.26	5.46
AD-1534493.1	21.43	1.40	34.04	3.79	56.91	5.89
AD-1534588.1	44.54	5.12	70.83	12.48	67.02	3.48
AD-1534511.1	22.85	1.44	44.49	8.24	61.39	4.77
AD-1534660.1	74.45	5.23	80.92	3.39	94.10	9.13
AD-1534534.1	22.93	0.28	39.25	2.24	61.05	2.64
AD-1534634.1	93.70	4.94	96.32	8.44	111.83	41.06
AD-1534659.1	37.46	5.93	53.76	6.38	64.44	4.47
AD-1534626.1	111.67	28.84	122.51	2.77	92.76	14.98
AD-1534470.1	41.78	7.90	44.45	4.38	62.80	6.14

Duplex ID	10 nM	ST DEVP	1 nM	ST DEVP	0.1 nM	ST DEVP
AD-1534499.1	69.72	3.78	62.83	11.66	82.50	8.57
AD-1534612.1	59.51	6.06	102.44	11.86	109.07	8.44
AD-1534656.1	95.76	9.13	102.62	7.53	110.72	12.77
AD-1534570.1	45.17	2.30	61.43	4.73	66.64	4.62
AD-1534546.1	30.42	2.00	44.54	2.58	63.28	2.03
AD-1631283.1	53.77	7.68	66.21	5.15	81.68	2.33
AD-1534547.1	73.39	6.72	69.85	3.20	83.71	6.41
AD-1534556.1	28.70	2.96	36.69	2.47	44.55	3.73
AD-1534580.1	46.19	5.80	70.43	6.94	79.09	5.28
AD-1534464.1	14.54	0.91	23.22	3.19	36.06	2.74
AD-1534482.1	105.24	4.35	99.89	10.63	113.69	7.62
AD-1534540.1	56.22	2.62	71.10	5.57	95.48	9.44
AD-1534541.1	58.76	1.47	78.48	10.61	87.69	3.78
AD-1534509.1	111.13	12.91	98.75	3.75	120.57	7.80
AD-1534655.1	75.99	10.47	82.03	2.89	94.00	6.37
AD-1534505.1	50.27	4.39	53.63	1.20	67.81	1.73
AD-1534616.1	63.20	4.69	90.91	9.30	87.50	6.44
AD-1534674.1	46.60	5.31	61.18	5.37	77.75	4.04
AD-1534603.1	57.38	3.04	71.07	8.17	76.83	11.06
AD-1534557.1	49.28	10.63	61.58	6.28	61.57	3.16
AD-1534624.1	37.25	4.11	46.55	3.61	58.11	11.13
AD-1534605.1	38.22	2.61	52.90	2.33	57.25	5.66
AD-1534648.1	23.28	0.90	36.75	0.48	43.91	5.12
AD-1534545.1	8.12	0.79	18.64	3.89	39.39	11.63
AD-1631282.1	17.87	1.46	25.82	1.70	33.53	1.13
AD-1534591.1	11.90	2.88	25.25	7.53	27.05	2.76
AD-1534518.1	14.37	0.54	23.94	2.38	39.01	3.34
AD-1534492.1	22.26	1.86	42.88	5.64	64.28	4.43
AD-1534587.1	31.44	6.31	64.88	7.94	67.09	4.43
AD-1534632.1	31.89	5.60	51.77	5.88	61.51	7.49
AD-1534633.1	25.37	1.39	39.39	1.51	50.81	7.80
AD-1631303.1	33.26	2.26	41.62	3.79	57.71	7.24
AD-1534576.1	29.90	4.81	48.65	5.92	56.26	2.84
AD-1631289.1	30.28	4.57	40.69	2.13	51.59	3.03
AD-1631311.1	33.11	1.82	45.93	3.49	57.99	2.57
AD-1534675.1	39.66	4.13	41.94	1.18	50.88	3.23
AD-1534560.1	52.00	5.20	80.40	16.47	64.60	2.84
AD-1534640.1	55.00	5.33	66.23	6.74	77.75	7.61
AD-1534480.1	51.85	0.44	71.51	1.59	90.26	2.48
AD-1631274.1	87.27	7.50	109.28	6.04	108.52	11.21
AD-1534481.1	67.88	8.36	72.24	9.31	98.31	4.10
AD-1631275.1	101.48	8.70	97.91	9.77	107.97	6.90
AD-1534527.1	30.30	1.72	34.31	2.33	59.43	6.24
AD-1534494.1	50.99	8.88	64.25	2.23	77.92	2.68
AD-1534593.1	42.30	4.13	70.83	4.22	84.34	2.97
AD-1534594.1	95.11	15.44	120.35	13.47	96.58	5.41
AD-1534523.1	20.73	2.38	35.28	4.19	52.64	4.37
AD-1534485.1	11.58	2.09	21.66	4.21	30.97	4.40
AD-1534668.1	62.38	4.22	69.11	4.86	78.53	3.23

Duplex ID	10 nM	ST DEVP	1 nM	ST DEVP	0.1 nM	ST DEVP
AD-1631301.1	55.40	2.99	76.96	3.34	93.86	6.65
AD-1534625.1	62.75	4.49	95.80	8.19	89.76	22.82
AD-1534609.1	31.94	6.22	49.22	5.02	81.17	11.88
AD-1631298.1	32.28	0.71	49.57	2.43	67.87	2.30
AD-1534610.1	66.61	10.90	85.27	7.31	83.15	7.74
AD-1534566.1	30.57	1.26	50.48	4.59	61.07	7.08
AD-1534652.1	56.37	4.36	80.74	2.00	101.95	2.26
AD-1534636.1	39.83	3.63	47.37	6.03	66.01	2.11
AD-1631304.1	58.79	5.17	71.58	5.61	89.88	6.49
AD-1534637.1	78.43	10.72	81.46	2.87	98.09	7.91
AD-1631307.1	32.14	4.23	40.25	4.31	53.35	2.05
AD-1534662.1	39.36	1.75	56.21	4.92	65.12	2.72
AD-1534602.1	71.84	2.85	87.49	7.15	96.69	21.16
AD-1631295.1	75.13	9.72	82.13	8.33	98.16	10.17
AD-1534646.1	74.39	8.40	79.91	2.36	88.50	8.43
AD-1534555.1	14.19	2.04	21.78	2.32	24.72	4.67
AD-1534606.1	22.38	2.44	41.68	6.48	48.18	5.88
AD-1534565.1	15.15	1.05	27.81	3.52	36.49	4.85
AD-1534649.1	32.03	2.33	42.71	2.10	55.30	1.28
AD-1534549.1	37.16	3.08	53.88	1.01	56.37	5.70
AD-1631284.1	58.27	4.92	73.77	4.58	89.86	12.68
AD-1534550.1	64.40	2.40	76.19	4.60	67.48	3.22
AD-1534572.1	70.78	8.37	101.43	14.08	82.58	5.38
AD-1534630.1	27.27	1.95	55.63	5.09	66.82	5.46
AD-1534629.1	30.89	2.17	44.31	4.68	53.39	2.22
AD-1631302.1	42.51	4.16	65.86	6.21	84.55	1.75
AD-1534650.1	24.92	3.62	44.74	3.33	49.68	1.53
AD-1631305.1	26.38	1.33	38.75	1.67	48.40	1.43
AD-1534554.1	36.30	3.47	49.15	4.29	54.07	2.93
AD-1631285.1	45.99	4.16	53.92	4.59	71.46	1.05
AD-1534600.1	21.88	1.47	36.18	6.87	40.92	0.57
AD-1631293.1	32.31	3.00	40.47	3.64	60.50	6.29
AD-1534551.1	27.52	1.50	51.79	5.47	50.75	4.67
AD-1534575.1	30.21	1.40	59.31	4.42	66.58	6.66
AD-1534673.1	88.45	7.15	96.79	5.22	97.59	5.35
AD-1534486.1	97.44	8.32	83.05	7.44	91.03	5.44
AD-1534645.1	92.81	4.34	97.75	7.27	94.46	1.87
AD-1534548.1	54.84	3.66	61.01	4.40	72.69	6.52
AD-1534512.1	41.38	4.05	49.95	5.89	72.18	11.00
AD-1534607.1	22.41	3.51	41.77	6.21	50.84	4.56
AD-1631297.1	34.17	0.45	45.39	2.50	65.66	1.88
AD-1534608.1	71.59	2.79	93.51	12.94	88.83	6.00
AD-1534559.1	67.09	14.06	88.92	9.40	67.72	9.79
AD-1631286.1	85.82	7.69	101.72	9.46	104.03	5.31
AD-1534613.1	48.97	1.67	81.30	3.49	89.54	4.96
AD-1534644.1	71.22	12.37	74.84	2.89	82.63	6.72
AD-1534521.1	95.41	5.25	91.24	10.80	87.94	7.81
AD-1534483.1	18.60	0.72	29.29	2.89	48.24	2.38
AD-1631276.1	28.16	3.11	38.22	4.72	46.80	4.46

Duplex ID	10 nM	ST DEVP	1 nM	ST DEVP	0.1 nM	ST DEVP
AD-1534542.1	32.16	4.33	42.12	3.99	76.81	4.74
AD-1534569.1	24.90	1.57	37.58	2.47	37.74	4.56
AD-1534657.1	68.88	9.23	83.63	3.46	86.67	3.28
AD-1534658.1	87.50	6.51	92.64	4.12	103.85	12.76
AD-1534510.1	66.95	12.63	76.26	6.81	109.70	8.76
AD-1534614.1	22.17	1.33	35.69	1.44	39.16	1.94
AD-1534615.1	48.68	5.86	58.64	0.34	56.03	5.90
AD-1534574.1	88.89	3.27	110.60	9.42	90.50	12.65
AD-1534635.1	97.55	11.72	99.79	11.64	79.32	6.76
AD-1534669.1	100.48	6.95	96.71	3.35	105.96	3.90
AD-1534583.1	40.12	6.39	53.52	6.93	74.22	3.31
AD-1534484.1	29.17	1.79	45.44	3.20	74.06	9.81
AD-1631277.1	45.53	6.63	51.88	3.05	64.54	4.20
AD-1534544.1	9.80	0.90	16.59	2.33	26.79	2.32
AD-1534552.1	11.79	0.84	18.76	1.61	21.40	1.21
AD-1534495.1	27.51	0.57	34.36	2.18	49.08	4.64
AD-1534496.1	59.63	5.55	71.87	4.27	85.96	9.83
AD-1534598.1	27.36	0.93	46.88	3.64	60.96	0.21
AD-1534538.1	16.64	0.97	28.96	7.93	36.95	3.66
AD-1534532.1	37.38	7.38	53.22	5.16	79.64	4.48
AD-1534504.1	33.37	2.43	63.89	10.64	78.46	2.75
AD-1534622.1	24.91	1.70	37.38	2.39	46.48	4.99
AD-1534623.1	38.76	3.45	59.33	3.08	63.85	6.29
AD-1534558.1	21.15	5.51	38.81	8.73	35.65	4.91
AD-1534476.1	100.42	3.72	103.32	7.53	112.28	15.47
AD-1534522.1	97.52	9.13	90.32	9.05	103.24	11.99
AD-1534543.1	23.40	3.30	31.89	5.31	62.63	5.62
AD-1631310.1	42.04	3.06	69.53	5.72	77.18	5.51
AD-1534667.1	79.92	5.34	85.29	6.91	89.36	1.42
AD-1534642.1	56.02	4.92	59.10	1.21	64.92	5.08
AD-1534487.1	57.71	2.85	82.81	4.42	101.67	8.02
AD-1631278.1	65.24	7.94	87.91	5.94	92.85	5.90
AD-1534561.1	21.42	0.17	45.34	4.28	52.27	6.74
AD-1631287.1	58.56	3.21	82.99	4.93	100.17	5.77
AD-1534641.1	73.86	8.50	84.35	7.79	86.67	8.05
AD-1534568.1	28.20	3.72	37.09	3.62	36.20	6.63
AD-1534590.1	45.61	4.96	65.21	2.09	64.08	8.72
AD-1534514.1	9.15	0.81	20.79	7.75	21.40	3.78
AD-1631279.1	15.09	1.83	24.40	1.84	29.12	3.14
AD-1534663.1	17.43	2.55	33.15	2.30	38.35	3.26
AD-1631308.1	18.85	2.28	26.52	1.35	32.51	3.61
AD-1534664.1	45.68	5.11	69.10	4.79	72.27	3.37
AD-1534604.1	22.49	2.74	37.81	4.80	43.88	1.36
AD-1631296.1	28.83	3.32	41.33	4.52	56.38	4.98
AD-1534564.1	11.94	0.63	26.23	3.96	30.49	2.09
AD-1631288.1	25.02	0.26	31.89	4.09	44.86	1.53
AD-1534647.1	24.19	3.90	36.83	2.48	47.37	4.84
AD-1534533.1	13.64	1.69	23.78	1.15	34.73	2.94
AD-1631280.1	22.71	3.94	33.78	1.19	40.60	5.98

Duplex ID	10 nM	ST DEVP	1 nM	ST DEVP	0.1 nM	ST DEVP
AD-1534531.1	56.59	2.28	56.07	2.84	77.08	1.45
AD-1534503.1	101.70	5.18	99.76	6.60	104.76	9.28
AD-1534589.1	31.00	2.96	39.68	4.48	33.08	4.27
AD-1534601.1	22.70	4.11	34.40	5.02	32.23	5.36
AD-1631294.1	29.61	4.32	31.06	5.07	41.95	4.70
AD-1534553.1	11.13	0.74	17.12	2.96	21.57	2.04
AD-1534596.1	37.73	2.95	59.86	3.33	70.31	2.58
AD-1534530.1	7.36	0.65	16.61	1.33	23.87	2.58
AD-1534618.1	12.56	1.50	27.00	2.97	29.22	2.76
AD-1534581.1	9.22	1.37	15.93	2.05	24.66	3.16
AD-85541.7	13.70	3.06	23.70	1.93	34.59	4.55
AD-85541.8	21.00	1.70	34.00	4.71	39.06	4.06
AD-85541.9	24.54	2.01	30.34	1.23	36.28	2.05
AD-1534465.1	10.13	1.67	21.35	3.86	27.79	2.47
AD-1534488.1	14.81	2.55	25.05	2.54	39.86	4.32
AD-1534562.1	12.89	0.79	23.06	1.66	30.15	2.35
AD-85537.26	16.08	2.94	30.40	7.02	31.69	0.91
AD-1534563.1	19.54	1.26	28.43	5.85	32.33	3.44
AD-85531.33	9.59	1.14	16.37	1.69	27.63	2.32
AD-85531.34	11.63	1.13	25.34	3.07	25.90	1.56
AD-85531.35	14.41	0.76	21.61	0.97	25.76	5.03
AD-1534473.1	8.34	1.93	19.25	1.09	24.74	3.16
AD-1631273.1	14.39	1.20	24.40	1.84	30.21	2.02
AD-67246.37	17.29	1.51	32.62	4.37	48.98	6.48
AD-67246.39	26.36	0.60	36.55	3.64	41.01	1.88
AD-67246.38	27.36	3.49	36.80	3.65	41.07	4.79
AD-1534508.1	16.93	2.35	29.16	1.21	37.18	2.49

Table 5. Single Dose Screen in PCH Cells (Transfection)

Duplex ID	10 nM	ST DEVP	1 nM	ST DEVP	0.1nM	ST DEVP
AD-1534465	18.28	2.59	15.45	4.14	45.78	2.05
AD-1534467	12.07	3.10	6.05	2.30	30.98	4.37
AD-1534471	16.33	1.87	26.32	5.91	33.02	5.65
AD-1534473	21.00	7.01	12.36	2.50	30.99	8.28
AD-1534479	14.34	2.96	17.49	2.45	23.12	4.90
AD-1534485	18.04	3.77	18.63	3.07	39.26	5.50
AD-1534491	15.63	3.10	14.30	0.62	27.53	3.54
AD-1534501	19.41	4.66	23.53	9.64	37.27	5.39
AD-1534502	22.47	5.56	24.66	3.53	42.18	6.77
AD-1534506	15.83	4.58	17.68	2.11	31.63	2.75
AD-1534507	22.94	2.67	18.38	1.38	47.06	11.33
AD-1534514	13.19	2.76	12.59	2.91	34.32	9.35
AD-1534528	9.97	0.51	7.92	1.46	32.13	3.41

Duplex ID	10 nM	ST DEVP	1 nM	ST DEVP	0.1nM	ST DEVP
AD-1534530	14.33	2.80	11.59	1.97	39.65	8.17
AD-1534544	22.56	1.46	21.96	5.92	40.59	5.23
AD-1534552	20.31	6.06	20.69	2.53	44.00	7.98
AD-1534553	19.61	6.51	14.53	2.92	37.40	3.63
AD-1534555	25.83	5.93	20.51	6.09	41.03	7.75
AD-1534562	23.80	5.39	21.31	3.20	43.33	7.54
AD-1534564	31.11	6.46	16.14	2.26	50.02	2.49
AD-1534567	13.12	2.13	14.23	2.49	32.72	1.77
AD-1534571	11.59	2.16	12.09	2.09	27.03	5.12
AD-1534577	23.92	4.59	25.94	6.11	35.58	6.86
AD-1534579	25.22	1.77	23.60	2.32	51.23	17.17
AD-1534581	16.65	3.40	20.77	1.77	36.46	8.19
AD-1534582	15.20	1.36	11.39	2.15	24.75	5.88
AD-1534585	27.97	5.53	21.50	5.56	41.44	5.78
AD-1534586	21.38	2.25	15.24	3.56	35.81	5.46
AD-1534591	22.21	4.44	11.47	2.83	39.13	14.38
AD-1534611	21.54	5.60	16.92	5.02	26.32	3.67
AD-1534618	12.24	1.49	17.18	1.28	23.62	3.38
AD-1534619	28.85	4.69	23.37	7.36	48.40	6.21
AD-1534621	19.40	3.69	18.97	3.45	34.18	10.17
AD-1534631	21.33	5.30	18.94	6.73	35.68	7.14
AD-1534653	19.84	2.61	10.18	1.16	37.95	7.51
AD-1534665	18.23	3.25	18.42	2.70	23.38	2.41
AD-1534678	18.35	3.52	21.08	2.66	53.65	27.11
AD-1631290	11.66	1.99	11.86	2.45	20.87	4.45
AD-1631291	20.88	6.70	14.34	3.15	32.19	7.09
AD-1631300	15.46	2.62	15.49	2.16	27.01	4.53
AD-1631309	22.31	5.30	16.15	2.09	29.91	6.11
AD-85525	20.67	1.13	18.92	2.46	31.87	7.27
AD-85531	18.32	4.78	14.89	2.49	35.11	5.44

Table 6. Single Dose Screen in PCH Cells (Free Uptake)

DuplexID	250 nM	ST DEVP	100 nM	ST DEVP	10 nM	ST DEVP	1 nM	ST DEVP
AD-1534465	15.49	4.60	11.75	0.66	23.94	3.21	72.39	5.65
AD-1534467	20.24	4.16	13.59	1.17	23.53	9.72	42.28	6.59
AD-1534471	12.66	5.63	7.71	0.77	19.57	2.37	39.78	1.01
AD-1534473	12.65	2.06	14.85	4.29	29.38	5.61	65.25	5.21
AD-1534479	8.80	2.22	7.78	1.45	19.10	3.76	33.74	3.53
AD-1534485	11.34	0.85	17.18	3.88	28.81	7.44	60.10	13.66
AD-1534491	15.43	5.04	9.75	1.79	27.15	7.17	54.43	9.67

DuplexID	250 nM	ST DEVP	100 nM	ST DEVP	10 nM	ST DEVP	1 nM	ST DEVP
AD-1534501	11.68	3.23	16.81	3.56	30.38	9.35	63.22	2.82
AD-1534502	18.95	4.92	19.95	3.55	35.53	5.83	72.11	10.93
AD-1534506	10.74	4.18	13.59	4.49	19.73	5.44	27.82	2.19
AD-1534507	17.45	3.38	17.10	2.07	26.46	4.47	90.06	17.23
AD-1534514	23.71	3.55	30.52	8.84	41.85	11.92	77.66	10.23
AD-1534528	11.03	1.87	11.12	0.94	25.24	6.79	36.92	2.42
AD-1534530	9.76	4.12	12.56	1.14	18.21	3.97	49.47	6.22
AD-1534544	32.95	12.99	21.42	1.92	34.90	6.86	66.88	6.19
AD-1534552	13.18	1.31	18.30	2.30	32.02	2.70	96.85	27.65
AD-1534553	13.62	1.42	14.67	1.69	28.77	4.34	64.03	6.70
AD-1534555	14.18	0.94	20.63	4.16	27.78	0.59	100.52	17.63
AD-1534562	27.51	5.92	15.55	1.93	35.28	7.24	87.15	23.59
AD-1534564	21.25	3.97	22.68	5.10	73.82	13.07	120.81	18.30
AD-1534567	8.54	1.44	8.92	2.37	13.74	2.39	43.37	3.72
AD-1534571	13.45	5.58	10.94	3.01	18.93	1.57	35.38	1.55
AD-1534577	19.69	5.02	15.07	5.24	33.22	8.93	55.75	10.30
AD-1534579	42.43	8.99	41.99	10.32	41.21	3.06	94.19	22.09
AD-1534581	18.28	5.74	17.64	2.32	20.20	4.12	47.15	5.06
AD-1534582	7.21	1.33	8.39	3.39	12.99	1.61	38.17	6.33
AD-1534585	20.27	4.42	19.78	3.81	31.81	4.61	74.16	8.03
AD-1534586	15.25	4.94	11.47	1.52	20.78	2.67	59.00	4.49
AD-1534591	17.17	5.83	16.29	5.06	25.53	2.44	75.89	9.01
AD-1534611	11.01	4.84	9.49	2.77	13.67	4.40	40.93	8.96
AD-1534618	10.02	0.99	12.89	3.09	21.95	2.61	47.87	7.74
AD-1534619	12.75	5.93	10.30	1.34	25.26	4.53	89.16	25.96
AD-1534621	7.79	1.98	8.33	2.89	13.57	1.24	41.04	10.75
AD-1534631	9.40	2.12	7.47	0.88	22.42	2.80	57.95	9.29
AD-1534653	9.46	1.32	18.66	7.80	18.66	7.90	75.10	4.95
AD-1534665	6.48	0.67	7.45	1.98	12.71	4.79	32.19	1.28
AD-1534678	10.38	2.80	16.52	8.97	27.79	8.28	117.53	7.03
AD-1631290	10.18	5.40	9.24	3.50	11.56	2.35	29.98	1.78
AD-1631291	17.98	5.19	19.35	5.09	26.38	3.97	57.54	12.13
AD-1631300	6.32	0.57	7.06	0.52	12.87	0.74	40.73	5.64
AD-1631309	8.01	1.29	6.06	1.13	11.91	2.02	39.44	10.42
AD-85525	28.43	7.95	27.87	9.26	30.92	7.75	84.19	18.23
AD-85531	18.70	9.18	15.06	5.36	28.70	5.06	73.62	7.07

Example 3. Single Dose Screen in Humanized Mice

A single 0.2 mg/kg dose of duplexes of interest identified in the above *in vitro* studies in 1 X PBS (AD-67244, AD-1534528, AD-1534582, AD-1534611, AD-1534479, AD-1534506, AD-1631309, AD-1534471, AD-1534555, or AD-1534678) were intravenously administered *via* tail vein injection to 36 week old mice (n=4) expressing human F12 mRNA (humanized F12 mice).

The unmodified nucleotide sequences of the sense and antisense strands of AD-67244 are:

5'-AACUCAAUAAAGUGCUUUGAA-3' (SEQ ID NO:13) and 5'-
UUCAAAGCACUUUAUUGAGUUUC-3' (SEQ ID NO:18); and the modified nucleotide sequences
of the sense and antisense strands of AD-67244 are:

5'-asascucaAfuAfAfAfgugcuuugaa-3' (SEQ ID NO:19) and 5'-
5 usUfscaaAfgCfAfcuuuAfuUfgaguususc-3' (SEQ ID NO:20).

At weekly intervals post-dose, up to six weeks post-dose, blood samples were collected with
3.8% sodium citrate (using a citrate-to-blood ratio of 1:9). Blood was processed for plasma within 45
minutes of collection, and plasma samples were frozen at -80°C until human FXII (hFXII) protein was
quantified using the Gryolab xP (BioAgilytix). The biotinylated capture and Alexa647-tagged
10 detection monoclonal anti-hFXII antibodies used on the Gyrolab platform were H4H26029P and
H4H26064P, respectively. Levels of human FXII protein in plasma samples were calibrated against a
standard curve generated using hFXII (HFXII 1212, Enzyme Research Laboratories) and plotted in
GraphPad Prism (Version 9.1.0) for each animal in each group of tested duplex as a function of
time.

15 The results of these assays are presented in the Tables below and Figure 1 and demonstrate
that the duplexes potently and durably inhibit human Factor XII protein expression.

Example 4. Effects of siRNA-GalNAC Conjugates in Non-Human Primates

Duplexes of interest from the *in vitro* and *in vivo* studies described above were further
20 investigated for their effectiveness in non-human primates. Specifically, a single 1 mg/kg or 3 mg/kg
dose of the duplexes of interest was subcutaneously administered to *Cynomolgus* monkeys. Plasma
samples were collected from the animals every week, and the plasma level of Factor XII (FXII)
protein was determined by ELISA. Plasma samples were collected from each animal pre-dose (Day -
3 and Day1) and on Days 8, 15, 22, 29, 36, 43, 50, 57, 64, 71, 78, 85, 92, 99, 106, 113, and 120 post-
25 dose. The data were normalized to the average of pre-dose (D-3 and D1). The D1 sample was taken
on the day of administration of the first dose, but before administration.

The results, presented as percent change of FXII protein level compared to the level of Factor
XII protein on dosing day, are shown in Figure 2 and demonstrate that the tested duplexes potently
and durably inhibit Factor XII protein expression.

30

week post dose	AD-67244				AD-1534528				AD-1534582			
	1	2	3	4	1	2	3	4	1	2	3	4
0	100	100	100	100	100	100	100	100	100	100	100	100
1	65.34895	20.50654	17.29683	27.21547	11.64461	3.554502	10.1083	14.98929	6.960784	7.714844	5.288673	9.657948
2	47.48658	15.84967	14.64581	14.79982	10.69943	8.395396	11.2515	12.95503	7.303922	5.615234	4.936095	7.243461
3	56.56418	21.07843	29.42199	21.1426	9.754253	7.989167	15.94465	15.63169	8.529412	8.886719	5.773468	7.696177
4	51.73255	39.29739	35.54976	28.78992	10.88847	13.30399	17.38869	21.19914	19.11765	19.33594	10.57735	14.78873
5	89.26305	56.0049	51.0213	45.56905	18.14745	56.93974	47.50301	45.61028	39.7549	37.64648	17.40855	29.67807
6	99.51196	67.81046	78.92221	77.10301	26.61626	24.64455	41.06498	44.48608	54.5098	63.08594	37.85809	45.57344

week post dose	AD-1534611				AD-1534479				AD-1534506			
	1	2	3	4	1	2	3	4	1	2	3	4
0	100	100	100	100	100	100	100	100	100	100	100	100
1	6.979543	7.963937	11.20084	18.43778	6.711409	18.90204	8.954584	12.01629	29.11985	45.15939	37.0155	27.28698
2	7.380666	8.226897	13.58848	17.93824	4.32047	19.00066	5.869752	8.044807	27.99625	52.89256	31.87984	35.18035
3	9.466506	11.26972	15.66011	27.65668	5.411074	14.49704	8.997429	11.0998	42.69663	63.10508	52.13178	62.25823
4	18.21099	21.93839	30.58287	47.00273	7.214765	14.2998	11.39674	13.30618	60.44007	87.7804	78.34302	99.94773
5	33.213	48.98573	74.92978	98.09264	12.41611	33.79356	19.66581	23.0482	107.912	159.3861	106.5891	134.2394
6	52.1059	68.67017	84.33989	97.32062	20.05034	29.91453	21.33676	29.9389	109.7846	174.144	109.5446	114.7412

week post dose	AD-1631309				AD-1534471				AD-1534555			
	1	2	3	4	1	2	3	4	1	2	3	4
0	100	100	100	100	100	100	100	100	100	100	100	100
1	50.47089	60.5	49.63441	39.58333	12.29314	17.51859	11.31222	17.60363	41.53754	37.8902	32.23109	39.03743
2	63.69863	65.6	71.91398	40.14757	12.19858	25.23234	16.20732	14.99148	45.29201	36.84966	33.86545	34.46767
3	75.21404	91.9	84.34409	69.57465	13.38061	25.78996	16.90662	17.54685	54.64839	41.15536	41.27708	44.57948
4	100.7705	100	82.92473	79.60069	17.68322	36.01301	23.44714	24.41794	77.89035	60.24399	55.72026	70.92854
5	133.476	137.7	91.69892	120.8333	23.92435	23.9777	20.19745	26.17831	138.4982	96.77072	88.93957	124.0156
6	135.8733	141.3	95.87097	108.7674	31.15839	38.61524	26.0798	51.50483	142.789	90.70685	102.8886	126.1546

week post dose	AD-1534678			
	1	2	3	4
0	100	100	100	100
1	44.48113	58.29255	32.05556	45.29486
2	43.06604	61.12017	34.05556	49.49812
3	47.0283	70.85372	49.88889	61.29235
4	67.59434	94.72539	69.38889	87.32748
5	105.8491	151.4954	122.3333	119.5734
6	125.9434	142.9581	139.3889	133.8143

EQUIVALENTS

Those skilled in the art will recognize, or be able to ascertain using no more than routine experimentation, many equivalents to the specific embodiments and methods described herein. Such equivalents are intended to be encompassed by the scope of the following claims.

5

We claim:

1. A double stranded ribonucleic acid (dsRNA) agent for inhibiting expression of Factor XII (F12) in a cell, wherein the dsRNA agent comprises a sense strand and an antisense strand forming a double stranded region, wherein the antisense strand comprises a region of complementarity to an mRNA encoding F12, and wherein the region of complementarity comprises at least 15 contiguous nucleotides differing by no more than 3 nucleotides from any one of the antisense nucleotide sequences in any one of Tables 2-3.
2. The dsRNA agent of claim 1, wherein the dsRNA agent comprises a sense strand comprising at least 15 contiguous nucleotides differing by no more than three nucleotides from any one of the nucleotide sequences of the sense strands in any one of Tables 2-3 and an antisense strand comprising at least 15 contiguous nucleotides differing by no more than three nucleotides from any one of the nucleotide sequences of the antisense strands in any one of Tables 2-3.
3. The dsRNA agent of claim 1, wherein the dsRNA agent comprises a sense strand comprising at least 15 contiguous nucleotides differing by no more than two nucleotides from any one of the nucleotide sequences of the sense strands in any one of Tables 2-3 and an antisense strand comprising at least 15 contiguous nucleotides differing by no more than two nucleotides from any one of the nucleotide sequences of the antisense strands in any one of Tables 2-3.
4. The dsRNA agent of claim 1, wherein the dsRNA agent comprises a sense strand comprising at least 15 contiguous nucleotides differing by no more than one nucleotide from any one of the nucleotide sequences of the sense strands in any one of Tables 2-3 and an antisense strand comprising at least 15 contiguous nucleotides differing by no more than one nucleotide from any one of the nucleotide sequences of the antisense strands in any one of Tables 2-3.
5. The dsRNA agent of claim 1, wherein the dsRNA agent comprises a sense strand comprising a nucleotide sequence selected from the group consisting of any one of the nucleotide sequences of the sense strands in any one of Tables 2-3 and an antisense strand comprising a nucleotide sequence selected from the group consisting of any one of the nucleotide sequences of the antisense strands in any one of Tables 2-3.
6. A double stranded ribonucleic acid (dsRNA) agent for inhibiting expression of Factor XII (F12) in a cell, wherein said dsRNA comprises a sense strand and an antisense strand forming a double stranded region, wherein the sense strand comprises at least 15 contiguous nucleotides

differing by no more than three nucleotides from any one of the nucleotide sequence of nucleotides 75-97, 76-98, 79-101, 115-137, 117-139, 123-145, 132-154, 135-157, 137-159, 144-166, 201-223, or 1462-1484 of SEQ ID NO: 1, and the antisense strand comprises at least 15 contiguous nucleotides differing by no more than 3 nucleotides from the corresponding nucleotide sequence of SEQ ID

5 NO:2.

7. The dsRNA agent of any one of claims 1-6, wherein the sense strand comprises at least 15 contiguous nucleotides differing by no more than 3 nucleotides from any one of the nucleotide sequence of nucleotides 135-157, 144-166, or 201-223 of SEQ ID NO: 1, and the antisense strand
10 comprises at least 15 contiguous nucleotides differing by no more than 3 nucleotides from the corresponding nucleotide sequence of SEQ ID NO:2.

8. The dsRNA agent of any one of claims 1-7, wherein the antisense strand comprises at least 15 contiguous nucleotides differing by no more than three nucleotides from any one of the antisense
15 strand nucleotide sequences of a duplex selected from the group consisting of AD-1534528, AD-1534479, AD-1534582, AD-1534611, AD-1534506, AD-1631309, AD-1534471, AD-1534555, AD-1534678, AD-1534585, AD-1534502, and AD-1534491.

9. The dsRNA agent of any one of claims 1-8, wherein the antisense strand comprises at least 15
20 contiguous nucleotides differing by no more than three nucleotides from any one of the antisense strand nucleotide sequences of a duplex selected from the group consisting of AD-1534528, AD-1534479, and AD-1534582.

10. The dsRNA agent of any one of claims 1-9, wherein the sense strand comprises at least 15
25 contiguous nucleotides differing by no more than three nucleotides from any one of the sense strand nucleotide sequences of a duplex selected from the group consisting of AD-1534528, AD-1534479, and AD-1534582 and the antisense strand comprises at least 15 contiguous nucleotides differing by no more than three nucleotides from any one of the antisense strand nucleotide sequences of a duplex selected from the group consisting of AD-1534528, AD-1534479, and AD-1534582.

30

11. The dsRNA agent of any one of claims 1-10, wherein the sense strand comprises at least 15 contiguous nucleotides differing by no more than two nucleotides from any one of the sense strand nucleotide sequences of a duplex selected from the group consisting of AD-1534528, AD-1534479, and AD-1534582 and the antisense strand comprises at least 15 contiguous nucleotides differing by

no more than two nucleotides from any one of the antisense strand nucleotide sequences of a duplex selected from the group consisting of AD-1534528, AD-1534479, and AD-1534582.

12. The dsRNA agent of any one of claims 1-11, wherein the sense strand comprises at least 15
5 contiguous nucleotides differing by no more than one nucleotide from any one of the sense strand
nucleotide sequences of a duplex selected from the group consisting of AD-1534528, AD-1534479,
and AD-1534582 and the antisense strand comprises at least 15 contiguous nucleotides differing by
no more than one nucleotide from any one of the antisense strand nucleotide sequences of a duplex
selected from the group consisting of AD-1534528, AD-1534479, and AD-1534582.
- 10 13. The dsRNA agent of any one of claims 1-12, wherein the sense strand comprises any one of
the sense strand nucleotide sequences of a duplex selected from the group consisting of AD-1534528,
AD-1534479, and AD-1534582 and the antisense strand comprises any one of the antisense strand
nucleotide sequences of a duplex selected from the group consisting of AD-1534528, AD-1534479,
15 and AD-1534582.
14. The dsRNA agent of any one of claims 1-13, wherein the dsRNA agent comprises at least
one modified nucleotide.
- 20 15. The dsRNA agent of any one of claims 1-14, wherein substantially all of the nucleotides of
the sense strand are modified nucleotides; substantially all of the nucleotides of the antisense strand
are modified nucleotides; or substantially all of the nucleotides of the sense strand and substantially
all of the nucleotides of the antisense strand are modified nucleotides.
- 25 16. The dsRNA agent of any one of claims 1-15, wherein all of the nucleotides of the sense
strand are modified nucleotides; all of the nucleotides of the antisense strand are modified
nucleotides; or all of the nucleotides of the sense strand and all of the nucleotides of the antisense
strand are modified nucleotides.
- 30 17. The dsRNA agent of any one of claims 14-16, wherein at least one of the modified
nucleotides is selected from the group consisting of a deoxy-nucleotide, a 3'-terminal
deoxythymidine (dT) nucleotide, a 2'-O-methyl modified nucleotide, a 2'-fluoro modified nucleotide,
a 2'-deoxy-modified nucleotide, a 2'-5'-linked ribonucleotide (3'-RNA), a locked nucleotide, an
unlocked nucleotide, a conformationally restricted nucleotide, a constrained ethyl nucleotide, an
35 abasic nucleotide, a 2'-amino-modified nucleotide, a 2'-O-allyl-modified nucleotide, 2'-C-alkyl-

modified nucleotide, 2'-hydroxly-modified nucleotide, a 2'-methoxyethyl modified nucleotide, a 2'-O-alkyl-modified nucleotide, a morpholino nucleotide, a phosphoramidate, a non-natural base comprising nucleotide, a tetrahydropyran modified nucleotide, a 1,5-anhydrohexitol modified nucleotide, a cyclohexenyl modified nucleotide, a nucleotide comprising a phosphorothioate group, a nucleotide comprising a methylphosphonate group, a nucleotide comprising a 5'-phosphate, a nucleotide comprising a 5'-phosphate mimic, a vinyl-phosphonate nucleotide, a thermally destabilizing nucleotide, a glycol modified nucleotide (GNA), a glycol nucleic acid S-Isomer (S-GNA), a nucleotide comprising a 2' phosphate, and a 2-O-(N-methylacetamide) modified nucleotide; and combinations thereof.

10

18. The dsRNA agent of any one of claims 14-16, wherein at least one of the modified nucleotides is selected from the group consisting of LNA, HNA, CeNA, 2'-methoxyethyl, 2'-O-alkyl, 2'-O-allyl, 2'-C- allyl, 2'-fluoro, 2'-deoxy, 2'-hydroxyl, and glycol; and combinations thereof.

15

19. The dsRNA agent of any one of claims 14-16, wherein at least one of the modified nucleotides is selected from the group consisting of a deoxy-nucleotide, a 2'-O-methyl modified nucleotide, a 2'-fluoro modified nucleotide, a 2'-deoxy-modified nucleotide, a 2'-5'-linked ribonucleotide (3'-RNA), a glycol modified nucleotide (GNA), a glycol nucleic acid S-Isomer (S-GNA), a nucleotide comprising a 2' phosphate, and a nucleotide comprising a phosphorothioate group; and combinations thereof.

20

20. The dsRNA agent of any one of claims 14-16, wherein at least one of the modified nucleotides is a nucleotide modified with a thermally destabilizing nucleotide modification.

25

21. The dsRNA agent of claim 20, wherein the thermally destabilizing nucleotide modification is selected from the group consisting of an abasic modification; a mismatch with the opposing nucleotide in the duplex; a destabilizing sugar modification, a 2'-deoxy modification, an acyclic nucleotide, an unlocked nucleic acid (UNA), a glycerol nucleic acid (GNA), and a glycol nucleic acid S-Isomer (S-GNA).

30

22. The dsRNA agent of any one of claims 1-21, wherein the double stranded region is 19-30 nucleotide pairs in length.

35


23. The dsRNA agent of claim 22, wherein the double stranded region is 19-25 nucleotide pairs in length.

24. The dsRNA agent of claim 22, wherein the double stranded region is 19-23 nucleotide pairs in length.
25. The dsRNA agent of claim 22, wherein the double stranded region is 23-27 nucleotide pairs in length.
26. The dsRNA agent of claim 22, wherein the double stranded region is 21-23 nucleotide pairs in length.
27. The dsRNA agent of any one of claims 1-26, wherein each strand is independently no more than 30 nucleotides in length.
28. The dsRNA agent of any one of claims 1-27, wherein the sense strand is 21 nucleotides in length and the antisense strand is 23 nucleotides in length.
29. The dsRNA agent of any one of claims 1-28 wherein the region of complementarity is at least 17 nucleotides in length.
30. The dsRNA agent of any one of claims 1-29, wherein the region of complementarity is between 19 and 23 nucleotides in length.
31. The dsRNA agent of any one of claims 1-30, wherein the region of complementarity is 19 nucleotides in length.
32. The dsRNA agent of any one of claims 1-31, wherein at least one strand comprises a 3' overhang of at least 1 nucleotide.
33. The dsRNA agent of any one of claims 1-31, wherein at least one strand comprises a 3' overhang of at least 2 nucleotides.
34. The dsRNA agent of any one of claims 1-33, further comprising a ligand.
35. The dsRNA agent of claim 34, wherein the ligand is conjugated to the 3' end of the sense strand of the dsRNA agent.


36. The dsRNA agent of claim 34 or 35, wherein the ligand is an N-acetylgalactosamine (GalNAc) derivative.

37. The dsRNA agent of any one of claims 34-36, wherein the ligand is one or more GalNAc derivatives attached through a monovalent, bivalent, or trivalent branched linker.

38. The dsRNA agent of any one of claims 34-37, wherein the ligand is



39. The dsRNA agent of claim 38, wherein the dsRNA agent is conjugated to the ligand as shown in the following schematic



and, wherein X is O or S.

40. The dsRNA agent of claim 39, wherein X is O.

41. The dsRNA agent of any one of claims 1-40, wherein the dsRNA agent further comprises at least one phosphorothioate or methylphosphonate internucleotide linkage.

42. The dsRNA agent of claim 41, wherein the phosphorothioate or methylphosphonate internucleotide linkage is at the 3'-terminus of one strand.
43. The dsRNA agent of claim 42, wherein the strand is the antisense strand.
- 5 44. The dsRNA agent of claim 42, wherein the strand is the sense strand.
45. The dsRNA agent of claim 41, wherein the phosphorothioate or methylphosphonate internucleotide linkage is at the 5'-terminus of one strand.
- 10 46. The dsRNA agent of claim 45, wherein the strand is the antisense strand.
47. The dsRNA agent of claim 45, wherein the strand is the sense strand.
- 15 48. The dsRNA agent of claim 41, wherein the phosphorothioate or methylphosphonate internucleotide linkage is at both the 5'- and 3'-terminus of one strand.
49. The dsRNA agent of claim 48, wherein the strand is the antisense strand.
- 20 50. The dsRNA agent of any one of claims 1-49, wherein the base pair at the 1 position of the 5'-end of the antisense strand of the duplex is an AU base pair.
51. A cell containing the dsRNA agent of any one of claims 1-50.
- 25 52. A pharmaceutical composition for inhibiting expression of a gene encoding Factor XII (F12) comprising the dsRNA agent of any one of claims 1-50 and a pharmaceutically acceptable carrier.
53. The pharmaceutical composition of claim 52, wherein dsRNA agent is in an unbuffered solution.
- 30 54. The pharmaceutical composition of claim 53, wherein the unbuffered solution is saline or water.
55. The pharmaceutical composition of claim 52, wherein said dsRNA agent is in a buffer solution.
- 35

56. The pharmaceutical composition of claim 55, wherein the buffer solution comprises acetate, citrate, prolamine, carbonate, or phosphate or any combination thereof.
57. The pharmaceutical composition of claim 56, wherein the buffer solution is phosphate buffered saline (PBS).
5
58. A method of inhibiting expression of a Factor XII (F12) gene in a cell, the method comprising contacting the cell with the dsRNA agent of any one of claims 1-50, or the pharmaceutical composition of any one of claims 52-57, thereby inhibiting expression of the F12 gene in the cell.
10
59. The method of claim 58, wherein the cell is within a subject.
60. The method of claim 59, wherein the subject is a human.
15
61. The method of claim 59 or 60, wherein the subject has an F12-associated disorder.
62. The method of claim 61, wherein the F12-associated disorder is selected from the group consisting of heredity angioedema (HAE), prekallikrein deficiency, malignant essential hypertension, hypertension, end stage renal disease, Fletcher Factor Deficiency, and , thromboembolic disease, inflammatory disease, or Alzheimer's Disease.
20
63. The method of claim 62, wherein the F12-associated disorder is heredity angioedema (HAE).
64. The method of claim 62, wherein the F12-associated disorder is prekallikrein deficiency.
25
65. The method of claim 62, wherein the F12-associated disorder is malignant essential hypertension.
66. The method of claim 62, wherein the F12-associated disorder is end stage renal disease.
30
67. The method of claim 62, wherein the F12-associated disorder is Fletcher Factor Deficiency.
68. The method of claim 62, wherein the F12-associated disorder is thromboembolic disease.
35

69. The method of claim 62, wherein the F12-associated disorder is inflammatory disease.
70. The method of claim 62, wherein the F12-associated disorder is Alzheimer's Disease.
- 5 71. The method of any one of claims 58-70, wherein contacting the cell with the dsRNA agent inhibits the expression of F12 by at least 50%, 60%, 70%, 80%, 90%, or 95%.
72. The method of any one of claims 58-71, wherein inhibiting expression of F12 decreases F12 protein level in serum of the subject by at least 50%, 60%, 70%, 80%, 90%, or 95%.
- 10 73. A method of treating a subject having a disorder that would benefit from reduction in Factor XII (F12) expression, comprising administering to the subject a therapeutically effective amount of the dsRNA agent of any one of claims 1-50, or the pharmaceutical composition of any one of claims 52-57, thereby treating the subject having the disorder that would benefit from reduction in F12 expression.
- 15 74. A method of preventing at least one symptom in a subject having a disorder that would benefit from reduction in Factor XII (F12) expression, comprising administering to the subject a prophylactically effective amount of the dsRNA agent of any one of claims 1-50, or the pharmaceutical composition of any one of claims 52-57, thereby preventing at least one symptom in the subject having the disorder that would benefit from reduction in F12 expression.
- 20 75. The method of claim 73 or 74, wherein the disorder is an F12-associated disorder.
- 25 76. The method of claim 75, wherein the F12-associated disorder is selected from the group consisting of heredity angioedema (HAE), prekallikrein deficiency, malignant essential hypertension, hypertension, end stage renal disease, Fletcher Factor Deficiency, thromboembolic disease, inflammatory disease, or Alzheimer's Disease.
- 30 77. The method of claim 76, wherein the F12-associated disorder is heredity angioedema (HAE).
78. The method of claim 76, wherein the F12-associated disorder is prekallikrein deficiency.
79. The method of claim 76, wherein the F12-associated disorder is malignant essential hypertension.
- 35

80. The method of claim 76, wherein the F12-associated disorder is end stage renal disease.
81. The method of claim 76, wherein the F12-associated disorder is Fletcher Factor Deficiency.
- 5 82. The method of claim 76, wherein the F12-associated disorder is thromboembolic disease.
83. The method of claim 76, wherein the F12-associated disorder is inflammatory disease.
84. The method of claim 76, wherein the F12-associated disorder is Alzheimer's Disease.
- 10 85. The method of any one of claims 73-84, wherein the subject is a human.
86. The method of any one of claims 73-85, wherein administration of the dsRNA agent to the subject causes a decrease in F12 protein accumulation in the subject.
- 15 87. The method of any one of claims 73-86, wherein administration of the dsRNA agent to the subject causes a decrease in one or more of elevated bradykinin, prekallikrein deficiency, malignant essential hypertension, hypertension, end stage renal disease, Fletcher Factor Deficiency, inflammation, thrombosis, edema swelling of the extremities, face, larynx, upper respiratory tract,
- 20 abdomen, trunk, genitals, or prodrome; laryngeal swelling; nonpruritic rash; nausea; vomiting; or abdominal pain.
88. The method of any one of claims 73-87, wherein the dsRNA agent is administered to the subject at a dose of about 0.01 mg/kg to about 50 mg/kg.
- 25 89. The method of any one of claims 73-88, wherein the dsRNA agent is administered to the subject subcutaneously.
90. The method of any one of claims 73-89, further comprising determining the level of F12 in a sample(s) from the subject.
- 30 91. The method of claim 90, wherein the level of F12 in the subject sample(s) is an F12 protein level in a blood or serum or liver tissue sample(s).
- 35 92. The method of any one of claims 73-91, further comprising determining the level of bradykinin, prekallikrein, or blood pressure in the subject.

93. The method of any one of claims 73-92, further comprising administering to the subject an additional therapeutic agent for treatment of an F12-associated disorder.

94. The method of claim 93, wherein the additional therapeutic agent is selected from the group consisting of an androgen, danazol or oxandrolone, Berinert®, Cinryze™, Rhuconest®, Ecallantide, Firazyr®, Kalbitor®, or a combination of any of the foregoing.

95. A kit comprising the dsRNA agent of any one of claims 1-45 or the pharmaceutical composition of any one of claims 52-57.

10

96. A vial comprising the dsRNA agent of any one of claims 1-45 or the pharmaceutical composition of any one of claims 52-57.

97. A syringe comprising the dsRNA agent of any one of claims 1-45 or the pharmaceutical composition of any one of claims 52-57.

15

98. An RNA-induced silencing complex (RISC) comprising an antisense strand of any of the dsRNA agents of claims 1-50.

20