

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
5 June 2003 (05.06.2003)

PCT

(10) International Publication Number
WO 03/045884 A1(51) International Patent Classification: C07C 39/16,
37/11, 37/62, 39/367, 39/42, 39/08, 37/18

(21) International Application Number: PCT/US02/35781

(22) International Filing Date:
7 November 2002 (07.11.2002)


(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
09/994,098 26 November 2001 (26.11.2001) US(71) Applicant: E.I. DU PONT DE NEMOURS AND COM-
PANY [US/US]; 1007 Market Street, Wilmington, DE
19898 (US).(72) Inventor: SHAPIRO, Rafael; 1415 Fresno Road, Wilm-
ington, DE 19803 (US).(74) Agent: DEITCH, Gerald, E.; E.I. du Pont de Nemours
and Company, Legal Patent Records Center, 4417 Lan-
caster Pike, Wilmington, DE 19805 (US).(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,
MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE,
SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ,
VC, VN, YU, ZA, ZM, ZW.(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE,
ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK,
TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ,
GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.(54) Title: PROCESS FOR PREPARING 3,3',6,6'-TETRAALKYL-2,2'-BIPHENOLS AND 3,3',6,6'-TETRAALKYL-5,5'-DI-
HALO-2,2'-BIPHENOLS

(I)

(57) Abstract: A process for making a
compound of the formula (I).

TITLE

PROCESS FOR PREPARING 3,3',6,6'-TETRAALKYL-2,2'-
5 BIPHENOLS AND 3,3',6,6'-TETRAALKYL-5,5'-DIHALO-2,2'-
BIPHENOLS

FIELD OF THE INVENTION

This invention relates to a process for preparing
10 3,3',6,6'-tetraalkyl-2,2'-biphenols and 3,3',6,6'-
tetraalkyl-5,5'-dihalo-2,2'-biphenols.

BACKGROUND OF THE INVENTION

Substituted biphenols such as 3,3',6,6'-
15 tetraalkyl-2,2'-biphenol; 3,3',4,4', 5,5',6,6'-
octaalkyl-2,2'-biphenols; 3,3',5,5',6,6'-hexaalkyl-
2,2'-biphenols; 3,3',5,5'-tetraalkyl-2,2'-biphenol; 3-
alkyl-5,5',6,6',7,7',8,8'-octahydro-2,2'-binaphthol;
3,3'-dialkyl-5,5',6,6',7,7',8,8'-octahydro-2,2'-
20 binaphthol and 3,3',6,6'-tetraalkyl-5,5'-dihalo-2,2'-
biphenol are compounds that can be used to make
phosphorus-based catalyst ligands. Such ligands
include phosphines, phosphinites, phosphonites, and
phosphites. Mono(phosphorous) ligands are compounds
25 that contain a single phosphorus atom which serves as a
donor to a transition metal, while bis(phosphorus)
ligands, in general, contain two phosphorus donor atoms
and typically form cyclic chelate structures with
transition metals.

30 In general, biphenols can be made by the oxidative
coupling of (mono)phenols, but often other types of
products, such as ketones, are obtained, and/or overall
yields are poor for other reasons.

Phenols can be oxidatively coupled to make the
35 corresponding biphenols by the use of a variety of

oxidizing agents, such as nitric acid, ferric chloride, potassium ferricyanide, chromic acid, 2,3-dichloro-5,6-dicyanobenzoquinone and di-t-butyl peroxide. 2,2'-

- 5 Dihydroxy-3,3'-di-isopropyl-6,6'-dimethylbiphenyl can be prepared from 2-isopropyl-5-methyl-phenol with 2,3-dichloro-5,6-dicyanobenzoquinone or di-t-butyl peroxide. See Tetrahedron, 1875, 1971 and J. Chem. Soc., Perkin Trans. II, 587, 1983. Some of the
- 10 oxidants and/or co-catalysts involve the use of relatively expensive and/or explosive (peroxides) compounds, which pose disadvantages for large scale commercial use.

- Phenols can also be oxidatively coupled using a
- 15 combination of a transition metal catalyst and an oxidizing agent such as persulfate anion or oxygen. See U.S. Patents 6,077,979; 4,139,544; 4,132,722; 4,354,048; and 4,108,908. See also J. Org. Chem. 1984, 49, 4456 and J. Org. Chem. 1983, 48, 4948. The cited
- 20 patents disclose the use of oxygen as an oxidizing agent with various copper complexes as catalysts (copper chromite; copper acetate with sodium mercaptoacetate; copper acetate with pentasodium/diethylenetriaminepentacetate; copper
- 25 acetate with 1,3-diamino-2-hydroxypropane-tetracetic acid). The examples in the patents disclose the use of 2,6-disubstituted phenol or 2,4-di-tert-butylphenol.


- The use of copper amine catalysts, with oxygen as an oxidizing agent, has been described in connection
- 30 with the oxidative coupling of 2,4-di-tert-butylphenol, 2-methyl-4-tert-butylphenol, 2-chlor-4-tert-butylphenol and 4-tert-butylphenol. See J. Org. Chem. 1984, 49, 4456 and J. Org. Chem. 1983, 48, 4948.

There is a continuing need in the art for methods for making with decent yields substituted biphenols suitable for making phosphorous-based catalyst ligands.

5

SUMMARY OF THE INVENTION

In its first aspect, the present invention is a process for making a compound of the formula I



I

10

wherein

R¹ is C₁ to C₁₀ primary or secondary alkyl or cycloalkyl,

R² is C₁ to C₁₀ primary or secondary alkyl or cycloalkyl, and

15

X is H, Cl, Br, or I,


comprising:

(1) when X is Cl

(a) chlorinating a compound of the formula

II

20




II

at the 4-position thereof to produce a compound of the formula III

WO 03/045884

PCT/US02/35781



wherein X is Cl,

- 5 (b) oxidatively coupling the compound of the formula III wherein X is Cl to produce a compound of the formula I wherein X is Cl;

(2) when X is H

- 10 (a) chlorinating a compound of the formula II at the 4-position thereof to produce a compound of the formula III wherein X is Cl,

- 15 (b) oxidatively coupling the compound of the formula III wherein X is Cl to produce a compound of the formula I wherein X is Cl, and

- 20 (c) dechlorinating the compound of the formula I wherein X is Cl to produce a compound of the formula I wherein X is H; or

(3) when X is Br or I

- 25 (a) chlorinating a compound of the formula II at the 4-position thereof to produce a compound of the formula III wherein X is Cl,

- (b) oxidatively coupling the compound of the formula III wherein X is Cl to produce a compound of the formula I wherein X is Cl,


WO 03/045884

PCT/US02/35781

(c) dechlorinating the compound of the formula I wherein X is Cl to produce a compound of the formula I wherein X is H, and

(d) substituting Br or I, respectively, for H at the 5 and 5' positions of the compound of the formula I wherein X is H.

10 In its second aspect, the present invention is a process for making a compound of the formula IV



wherein

15 R¹ is C₁ to C₁₀ primary or secondary alkyl or cycloalkyl,


R⁴ is C₁ to C₁₀ primary or secondary alkyl or cycloalkyl, and

X is H, Cl, Br, or I

20 comprising:

(1) when X is H


(a) alkylating a compound of the formula V



25 at the 4-position thereof to produce a compound of the formula VI

WO 03/045884

PCT/US02/35781




VI

wherein R^3 is C_4 to C_{20} tertiary alkyl,

5

- (b) oxidatively coupling the compound of the formula VI to produce a compound of the formula VII



VII

10

- (c) dealkylating a compound of the formula VII to produce a compound of the formula IV wherein X is H; or

- (2) when X is Cl, Br, or I

15

- (a) alkylating a compound of the formula V at the 4-position thereof to produce a compound of the formula VI,
- (b) oxidatively coupling the compound of the formula VI to produce a compound of the formula VII,
- (c) dealkylating a compound of the formula VII to produce a compound of the formula IV wherein X is H, and
- (d) substituting Cl, Br, or I, respectively, for H at the 5 and 5' positions of the compound of the formula IV wherein X is H.


20

25

WO 03/045884

PCT/US02/35781

In its third aspect, the present invention is a process for making a compound of the formula I



I

wherein


R¹ is C₁ to C₁₀ primary or secondary alkyl or cycloalkyl,

R² is C₁ to C₁₀ primary or secondary alkyl or cycloalkyl, and

X is H,

comprising:

- (a) oxidatively coupling a compound of the formula III



III

wherein X is Cl, to produce a compound of the formula I wherein X is Cl, and

- (b) dechlorinating the compound of the formula I wherein X is Cl to produce a compound of the formula I wherein X is H.


In another aspect the present invention is a compound selected from the group consisting of

3,3',6,6'-tetramethyl-2,2'-biphenol, and
3,3'-di-isopropyl-6,6'-dimentyl-2,2'-biphenol.

5

DETAILED DESCRIPTION OF THE INVENTION

The first aspect of the present invention provides a process for preparing 3,3',6,6'-tetraalkyl-2,2'-biphenol, comprising (1) substituting chlorine for hydrogen at the 4- position of 2,5-dialkylphenol, (2)
10 oxidatively coupling the resulting 2,5-dialkyl-4-chloro-phenol, and (3) removing chlorine from the resulting compound. The second step is carried out by analogy with the methods of Sartori, et al (Tetrahedron, 1992, 48, 9483), but using the free
15 phenol rather than its dichloroaluminate derivative. The three steps of the process are shown below.



wherein R¹ is C₁ to C₁₀ primary or secondary alkyl or
20 cycloalkyl; and R² is C₁ to C₁₀ primary or secondary
alkyl or cycloalkyl.

Preferred R^1 are methyl, ethyl, *n*-propyl, isopropyl, *n*-butyl, *sec*-butyl, cyclohexyl, and cyclopentyl. Preferred R^2 are methyl and ethyl. The
5 alkyl groups at the 2- and 5-positions may be the same or different.


In the first step of the process, a 2,5-dialkylphenol can be reacted with a chlorinating agent, such as chlorine or sulfuryl chloride, preferably in
10 the presence of 1 to 10 mol% of a catalyst such as aluminum chloride or a diaryl sulfide such as diphenyl sulfide or a mixture thereof. See Watson, *J. Org. Chem.*, 1985, 50, 2145. The reaction may be conducted
neat (without a solvent) or in a medium such as
15 dichloromethane, chlorobenzene, or other inert solvent at a temperature between -30 and 60°C, preferably at about 25 °C. The reaction is typically performed at or about atmospheric pressure for ease of operation.

In the second step of the process, the resulting
20 2,5-dialkyl-4-chlorophenol can be oxidatively coupled to give the corresponding dimeric chlorophenols (5,5'-dichloro-3,3',6,6'-tetraalkyl-2,2'-biphenol). The preferred method for oxidative coupling of the chlorinated phenols is by the use of an iron(III) salt,
25 preferably ferric chloride, in a suitable polar, aprotic solvent such as dichloromethane or nitromethane, preferably nitromethane at temperature between 0°C and 60°C, preferably about 35°C. The product is isolated by dilution with water and
30 filtration.

In the third step of the process, dechlorination of 5,5'-dichloro-3,3',6,6'-tetraalkyl-2,2'-biphenols can be accomplished by hydrogenolytic reduction to provide the required 3,3',6,6'-tetraalkyl-2,2'-

biphenols. The reduction is carried out in the presence of hydrogen gas, preferably at pressures between 1 and 50 atmospheres and temperature between 5° and 80°C, and a formate salt, such as sodium formate, and Raney® nickel or palladium catalyst such as palladium hydroxide on carbon. If a palladium catalyst is used, the reaction is generally carried out in a protic solvent such as methanol, containing 1.0 to 4.0 equivalents of an amine such as triethylamine to absorb the hydrogen chloride produced in the reaction.

The second aspect of the present invention provides a process for preparing a compound of the formula IV, comprising (1) substituting a tertiary alkyl group for hydrogen at the 4-position of 2,5-dialkylphenol, (2) oxidatively coupling the resulting 2,5-dialkyl-4-tert-alkyl-phenol, and (3) removing the tertiary alkyl group from the resulting compound. The three steps of the process are shown below.



wherein R^1 is C_1 to C_{10} primary or secondary alkyl or cycloalkyl; R^4 is C_1 to C_{10} primary or secondary alkyl or cycloalkyl; and R^3 is C_4 to C_{20} tertiary alkyl.

5 In the first step of the process, a 2,5-dialkylphenol can be reacted with a tert-alkyl halide in the presence of a Lewis Acid catalyst, such as zinc chloride or aluminum chloride, to give a 2,5-dialkyl-4-tert-alkylphenol. Alternatively, the 2,5-dialkyl-4-
10 tert-alkylphenol can be prepared from contacting 2,5-dialkylphenol with 2,2-dialkylethylene in the presence of an acid catalyst. An example of the alternative method is the incorporation of a tert-butyl group into the 4- position of 2,5-dialkylphenol by reacting 2,5-
15 dialkylphenol with isobutylene in the presence of sulfuric acid.

 In the second step of the process, 2,5-dialkyl-4-tert-alkylphenol can be oxidatively coupled using a copper diamine catalyst and oxygen as the oxidizing
20 agent.


 The copper diamine catalyst can be prepared using the procedure described in the Tetrahedron Letters, 1994, 35, 7983. A copper halide, such as $CuCl$, $CuBr$, CuI , or $CuCl_2$, is added to a mixture of alcohol, such
25 as methanol, and water and the diamine is slowly added. After the addition of the diamine, air is sparged through the mixture with vigorous stirring. The catalyst is filtered. Additional catalyst can be obtained by concentrating the filtrate and filtering
30 the desired catalyst. The catalyst can also be prepared in situ by contacting the copper halide and the diamine in the solvent for the coupling reaction. Suitable solvents for the oxidative coupling of tri and tetrasubstituted phenols are methylene chloride and

aromatic solvents such as xylene, benzene and toluene. Example of diamines include, but are not limited to, the following: N,N,N',N'-tetraethylethylene diamine, N,N,N',N'-tetraethyl-1,3-propanediamine, N,N,N',N'-tetraethylmethane diamine, N,N,N',N'-tetramethyl-1,6-hexanediamine, N,N,N',N'-tetramethyl-1,3-propanediamine, dipiperidinomethane, N,N,N',N'-tetramethylethylene diamine and 1,4-diazabicyclo-
 10 (2,2,2)-octane. Preferably, the diamine is N,N,N',N'-tetramethylethylene diamine.

In the third step of the process, the 3,3',6,6'-tetraalkyl-5,5'-di-tert-alkyl-2,2'-biphenol can be dealkylated by contacting it with a strongly acidic
 15 catalyst, such as an alkyl- or arylsulfonic acid, sulfuric acid, phosphoric acid, aluminum chloride, or the like, optionally in the presence of a solvent such as toluene, chlorobenzene, nitromethane, or xylene, typically at temperatures between 10 and 150°C.

The oxidative coupling can be carried out neat (without a solvent) or using one or more of a wide range of poorly oxidizable solvents including dichloromethane, chlorobenzene, toluene, xylenes, nitromethane, paraffins, etc. Static air, air-flow, or
 25 oxygen can be used as oxidants in the oxidative coupling.

The third aspect of the present invention provides a process for making a compound of the formula I



I

WO 03/045884

PCT/US02/35781

wherein


R^1 is C_1 to C_{10} primary or secondary alkyl or cycloalkyl,

5 R^2 is C_1 to C_{10} primary or secondary alkyl or cycloalkyl, and

X is H,

comprising:

10 (a) oxidatively coupling a compound of the formula III



III

15 wherein X is Cl, to produce a compound of the formula I wherein X is Cl, and


(b) dechlorinating the compound of the formula I wherein X is Cl to produce a compound of the formula I wherein X is H.

20 In the first step of the process, the resulting 2,5-dialkyl-4-chlorophenol can be oxidatively coupled to give the corresponding dimeric chlorophenols (5,5'-dichloro-3,3',6,6'-tetraalkyl-2,2'-biphenol). The preferred method for oxidative coupling of the
 25 chlorinated phenols is by the use of an iron(III) salt, preferably ferric chloride, in a suitable polar, aprotic solvent such as dichloromethane or nitromethane, preferably nitromethane at temperature between 0°C and 60°C, preferably about 35°C. The

product is isolated by dilution with water and filtration.

In the second step of the process, dechlorination
 5 of 5,5'-dichloro-3,3',6,6'-tetraalkyl-2,2'-biphenols
 can be accomplished by hydrogenolytic reduction to
 provide the required 3,3',6,6'-tetraalkyl-2,2'-
 biphenols. The reduction is carried out in the
 presence of hydrogen gas, preferably at pressures
 10 between 1 and 50 atmospheres and temperature between 5°
 and 80°C, and a formate salt, such as sodium formate,
 and Raney® nickel or palladium catalyst such as
 palladium hydroxide on carbon. If a palladium catalyst
 is used, the reaction is generally carried out in a
 15 protic solvent such as methanol, containing 1.0 to 4.0
 equivalents of an amine such as triethylamine to absorb
 the hydrogen chloride produced in the reaction.

In the first, second and third aspects of the
 present invention, a 3,3',6,6'-tetraalkyl-5,5'-dihalo-
 20 2,2'-biphenol may be halogenated at the para positions
 of 3,3',6,6'-tetraalkyl-2,2'-biphenol, as shown below,



wherein R¹ is C₁ to C₁₀ primary or secondary alkyl or
 25 cycloalkyl; R² is C₁ to C₁₀ primary or secondary alkyl
 or cycloalkyl; and X is Cl, Br or I.

Addition of Br to 3,3',6,6'-tetraalkyl-2,2'-
 biphenols can be accomplished by reaction of Br₂ in a
 suitable solvent. Typical solvents for bromination are
 30 low polarity solvents such as chloroform,
 dichloromethane, carbon tetrachloride, and carbon

disulfide. In some cases, aqueous bromine can be used. The preferred process is one carried out in a low polarity solvent. This reaction can be accomplished at
5 -10°C to 50°C, preferably at room temperature.

The compounds which are produced by the processes of the present invention can be used as reactants to make phosphorous-containing ligands that are useful to make catalysts that, in turn, are useful in both
10 hydrocyanation and hydroformylation reactions. Bidentate phosphite ligands are particularly useful.

Bidentate phosphite ligands can be prepared as described in U.S. Patent 5,235,113 by contacting phosphorochloridites with the biphenol compounds made
15 by the processes of the present invention. More recent U.S. Patents 6,031,120 and 6,069,267, incorporated herein by reference, describe selective synthesis of bidentate phosphite ligands in which a phosphorochloridite is prepared in-situ from phosphorus
20 trichloride and a phenol such as o-cresol and then treated in the same reaction vessel with an aromatic diol to give the bidentate phosphite ligand. The biphenols of the present invention are substituted for the aromatic diol.

25 The compounds which are produced by the processes of the present invention can be polymerized and then used as reactants to make phosphorous-containing ligands that are useful to make catalysts that, in turn, are useful in both hydrocyanation and
30 hydroformylation reactions.

The compounds made by the processes of the present invention, in which X is H, can be used to make polymeric ligands by a process which comprises:

(1) reacting the compounds made by the processes of the present invention, in which X is H, with a compound containing at least two benzyl chloride groups, in the presence of a Lewis acid catalyst, and (2) reacting the product of step (1) with at least one phosphorochloridite compound in the presence of an organic base. Preferably the Lewis acid catalyst is zinc chloride or aluminum chloride, and the organic base is a trialkylamine.

The compounds made by the processes of the present invention, in which X is Cl, Br, or I, can be used to make polymeric ligands by a process which comprises:

- (1) protecting the OH groups by substituting a lower alkyl protecting group for H on the OH groups to make a protected compound,
- (2) treating the protected compound with a compound containing at least two boronic groups in the presence of a Group VIII transition metal catalyst,
- (3) replacing the protecting group of the product from step (2) with hydrogen, and
- (4) reacting the product of step (3) with at least one phosphorochlorodite compound in the presence of an organic base.

Preferably, the Group VIII transition metal is palladium, nickel or copper and the organic base is a trialkylamine compound in which the alkyl group is a C₁ to C₁₂ branched or straight chain alkyl group. More preferably the organic base is triethylamine.

Two particularly important industrial catalytic reactions using phosphorus-containing ligands are olefin hydrocyanation and isomerization of branched nitriles to linear nitriles. See, for example,

U.S. Patents 5,512,695 and 5,512,696, and International Patent Application WO9514659. Phosphite ligands are particularly useful for both reactions. The

5 hydrocyanation of unactivated and activated ethylenically unsaturated compounds (olefins) using transition metal complexes with monodentate and bidentate phosphite ligands is well known. Bidentate phosphinite and phosphonite ligands are useful as part
10 of a catalyst system for the hydrocyanation of ethylenically unsaturated compounds. Bidentate phosphinite ligands are also useful as part of a catalyst system for the hydrocyanation of aromatic vinyl compounds.

15 Hydroformylation is another industrially useful process that utilizes catalysts made from phosphorus-containing ligands. The use of phosphine ligands, including diphosphines, is known for this purpose. The use of catalysts made from phosphite ligands is also
20 known. Such catalysts usually contain a Group VIII metal. See for example, U.S. Patent 5,235,113.

Two particularly useful compounds that can be made by the present processes are 3,3',6,6'-tetramethyl-2,2'-biphenol and 3,3'-di-isopropyl-6,6'-dimentyl-2,2'-
25 biphenol.

EXAMPLES

The following non-limiting examples illustrate the
30 present invention. All parts, proportions, and percentages are by weight, unless otherwise indicated.

Example 1

Process in accordance with the first aspect of the invention for preparing 3,3',6,6'-tetramethyl-2,2'-

5 biphenol

First step of the process: Preparation of 4-Chloro-2,5-xyleneol :

To a solution of 100 g (0.82 mol) of 2,5-xyleneol and 0.9 g of diphenyl sulfide in 700 mL of
10 dichloromethane was added a solution of 106 g (0.79 mol) of sulfuryl chloride in 100 mL of dichloromethane, maintaining the temperature at 5-15 °C. The mixture was stirred for an additional hour and then poured onto 400 g of ice-water containing 5 g of sodium bisulfite.
15 The layers were separated, and the organic phase was washed with water, dried (MgSO₄), and concentrated to dryness. The crude solids were slurried with a minimum amount of hexanes, filtered, and suction-dried to give 121 g (94%) of product, homogeneous by Thin Layer
20 Chromatography (TLC), GC, and ¹H-NMR analysis. ¹H-NMR (CDCl₃) δ 2.18 (s, 3H), 2.27 (s, 3H), 4.61 (s, 1H), 6.63 (s, 1H), 7.07 (s, 1H). Lit (Blackstock, Aust. J. Chem. 1973, 26, 775): mp 74-75°C.

25 Second step of the process: Preparation of 5,5'-dichloro-3,3',6,6'-tetramethyl-2,2'-biphenol :

To a mechanically-stirred mixture of 71.4 g (0.458 mol) of 4-chloro-2,5-xyleneol and 120 mL of nitromethane was added 94 g (0.59 mol) of anhydrous ferric chloride
30 in portions over about 20 minutes with cooling to maintain the temperature below 35 °C. The mixture was stirred for an additional 3 hours and then 300 mL of ice water containing 50 mL of concentrated HCl was added, followed by 300 mL of hexanes. The mixture was

filtered, and the solids were washed with water and hexanes and dried in vacuo to give 51.0 g of product. The organic phase was separated from the filtrate, washed with water, and concentrated; the residue was then slurried in hexanes and filtered to give another 11 g of product. The total yield was thus 62 g (87%) of a tan solid (mp 148-155 °C). Additional purification to remove traces of iron helps facilitate the subsequent reductive dechlorination. Purification could be accomplished by dissolving in ethyl acetate, washing this solution with aqueous ethylenediamine-tetraacetic acid disodium salt (EDTA-2Na, EDTA = ethylenediaminetetraacetic acid), drying (MgSO₄), concentration and washing with hexanes to afford off-white material with mp 164°C. ¹H-NMR (CDCl₃) δ 1.98 (s, 3H), 2.25 (s, 3H), 4.60 (s, 1H), 7.25 (s, 1H).

Third step of the process: Preparation of 3,3',6,6'-tetramethyl-2,2'-biphenol:

A sample of purified 5,5'-dichloro-3,3',6,6'-tetramethyl-2,2'-biphenol (15.0 g, 48.4 mmol) was dissolved in 100 mL of ethanol containing 10 mL of water and 20 mL of triethylamine. This solution was added to 1.0 g (dry weight basis) of moist 20% Pd(OH)₂/C (Pearlman's catalyst) and hydrogenated at 50 psig for 2 hours at ambient temperature. The product was isolated by filtration of catalyst, concentration, dissolution of the residue in EtOAc, washing with water, and concentration to dryness to give 11.0 g (94%) of product, mp 110-113°C. ¹H-NMR (CDCl₃) δ 1.95 (s, 3H), 2.25 (s, 3H), 4.71 (s, 1H), 6.81 (d, 1H, J = 7.5 Hz), 7.10 (d, 1H, J = 7.5 Hz).

The second and third steps of the foregoing example also illustrate the third aspect of the invention.

5

Example 2

Process in accordance with the third aspect of the invention for preparing 3,3'-di-isopropyl-6,6'-dimethyl-2,2'-biphenol

- 10 First step of the process: Preparation of 5,5'-dichloro-3,3'-diisopropyl-6,6'-dimethyl-2,2'-biphenol:

A well-stirred mixture of 36.0 g (0.195 mol) of chlorothymol and 50 mL of nitromethane was cooled to 5 °C and 40 g (0.25 mol) of anhydrous ferric chloride was added over 20 minutes. The mixture was allowed to warm to ambient temperature and held an additional hour. Ice-water (300 mL) was added all at once, and the mixture was concentrated at reduced pressure to remove about 100 mL of the nitromethane-water azeotrope. The solids were filtered and recrystallized from aqueous isopropanol to give 23.3 g of a first crop and 3.9 g of a second crop of solids, mp 98°C. ¹H-NMR (CDCl₃) δ 1.24 (two d, 6H, J = 7 Hz), 1.98 (s, 3H), 3.26 (septet, 1H, J = 7 Hz), 4.63 (s, 1H), 7.30 (s, 1H).

- 25 Second step of the process: 3,3'-di-isopropyl,6,6'-dimethyl-2,2'-biphenol:

This substituted biphenol was prepared similarly to the third step of Example 1 except 5,5'-dichloro-3,3'-diisopropyl-6,6'-dimethyl-2,2'-biphenol was used instead of 5,5'-dichloro-3,3',6,6'-tetramethyl-2,2'-biphenol, mp 89-92°C. ¹H-NMR (CDCl₃) δ 1.25 (d, 6H), 1.95 (s, 3H), 3.28 (septet, 1H), 4.76 (s, 1H), 6.88 (d, 1H, J = 7.5 Hz), 7.18 (d, 1H, J = 7.5 Hz).

Example 3

5 Process in accordance with the second aspect of the
invention for preparing 3,3',6,6'-tetramethyl-2,2'-
biphenol

First step of the process: Preparation of 4-t-butyl-
2,5-xyleneol

Preparation of 4-t-Butyl-2,5-xyleneol: 2,5-Xyleneol
10 (90 g, 0.73 mol) was melted at 80 °C, 1 mL of
concentrated sulfuric acid was added, and the mixture
was heated at 90 °C while isobutylene gas was
introduced subsurface over 4 hours. The reaction
appeared to stall at about 80% conversion. The
15 reaction mass was diluted with water and neutralized
with NaHCO₃, and some starting xyleneol was removed by
steam-distillation. Since the steam-distillation did
not completely remove the starting material, the
residue was dissolved in hot hexanes, separated from
20 the aqueous phase, and cooled in an ice-bath. The
precipitated product was filtered and washed with cold
hexanes to give 64 g (49%) of 4-t-Butyl-2,5-xyleneol ;
lit.(Stevens, *Ind. Eng. Chem.* 1943, 655; Parc, *Rev.*
Inst. Fr. Pet. 1960, 680) mp 70-72°C. ¹H-NMR (CDCl₃)
25 δ 1.37, (s, 9H), 2.20 (s, 3H), 2.43 (s, 3H), 4.85 (s,
1H), 6.53 (s, 1H), 7.08 (s, 1H).

Second step of the process: Preparation of 5,5'-Bis(t-
butyl)-3,3',6,6'-tetramethyl-2,2'-biphenol :

To a solution of 18.6 g (0.104 mol) of 4-t-butyl-
30 2,5-xyleneol in 20 mL of dichloromethane was added 0.6 g
(3 mmol) of copper chlorohydroxide-TMEDA complex (TMEDA
= tetramethylethylenediamine). The dark purple mixture
was stirred under ambient air overnight. Gas
chromatography (GC) analysis showed only 25%

conversion, so the mixture was diluted with dichloromethane, dried (MgSO_4) and concentrated to dryness. To the crude residue was added 20 mL of cyclohexane and 1.2 g (6 mmol) of the above copper chlorohydroxide-TMEDA catalyst, and the mixture was stirred under air at ambient temperature for three days (85% conversion). The purple solution was concentrated to dryness, and the residue was chromatographed on silica gel to give 10.2 g (55%) of pure 5,5'-Bis(t-butyl)-3,3',6,6'-tetramethyl-2,2'-biphenol, mp 103-105°C. $^1\text{H-NMR}$ (CDCl_3) δ 1.42, (s, 9H), 2.06 (s, 3H), 2.25 (s, 3H), 4.54 (s, 1H), 6.51 (s, 1H), 7.24 (s, 1H).

Third step of the process: Preparation of 3,3',6,6'-tetramethyl-2,2'-biphenol:

To a 50 mL flask were added 0.5 g of 5,5'-Bis(t-butyl)-3,3',6,6'-tetramethyl-2,2'-biphenol, 5 mL of xylenes and 0.05 g of p-toluenesulfonic acid. The mixture was refluxed for about 2 hours. The mixture was cooled and water added. The mixture was extracted with hexanes; the organic layer was washed with water and dried over MgSO_4 . After removing the solvent, the residue was recrystallized from petroleum ether.

Example 4

Process in accordance with the second aspect of the invention for preparing 5,5'-Di-t-butyl-3,3'-di-isopropyl,6,6'-dimethyl-2,2'-biphenol

First step of the process: Preparation of 4-t-Butylthymol:

To 30 g (0.20 mol) of thymol, heated at 60 °C under nitrogen, was added 1 g of concentrated sulfuric acid. After heating to 90 °C, a slow stream of isobutylene was introduced over about 2 hours. The

reaction stalled at about 50% conversion, so an additional charge of sulfuric acid was added and the reaction was monitored by GC-analysis until
5 approximately 70-80% conversion was achieved. The reaction was worked up as in Example 1 and the crude residue was recrystallized from hexanes to give 20 g of 4-t-butylthymol, mp 68-69°C, lit (United States Patent 4,880,775): mp 76-77°C. ¹H-NMR (CDCl₃) δ 1.25 (d, 6H, J = 7 Hz), 1.38, (s, 9H), 2.44 (s, 3H), 3.15 (septet, 1H), 4.49 (s, 1H), 6.51 (s, 1H), 7.18 (s, 1H).

Second step of the process: Preparation of 5,5'-Di-t-butyl-3,3'-di-isopropyl,6,6'-dimethyl-2,2'-biphenol :

15 To a solution of 20 g (0.104 mol) of 4-t-butylthymol in 50 mL of dichloromethane was added 1.0 g (5 mmol) of copper chlorohydroxide-TMEDA complex and the dark purple mixture was allowed to stir under ambient air for three days (50% conversion). The
20 mixture was diluted with hexanes, washed with aqueous EDTA solution, dried (MgSO₄) and concentrated to dryness. The residue was chromatographed on silica gel to give 3.6 g (34% based on conversion) of pure dimer 5,5'-Di-t-butyl-3,3'-di-isopropyl,6,6'-dimethyl-2,2'-
25 biphenol, mp 105-108 °C. ¹H-NMR (CDCl₃) δ 1.26 (d, 6H) 1.43, (s, 9H), 3.25 (septet, 1H), 4.58 (s, 1H), 7.30 (s, 1H).

Debutylating 5,5'-di-t-butyl-3,3'-diisopropyl-6,6'-dimethyl-2,2'-biphenol

30 A 500-mL resin kettle equipped with mechanical stirrer and condenser was placed in an oil bath and charged with 153 g of a mixture of 5,5'-di-t-butyl-3,3'-diisopropyl-6,6'-dimethyl-2,2'-biphenol in a

hydrocarbon solvent. By gas chromatography analysis, the mixture was 15.0% 5,5'-di-t-butyl-3,3'-di-isopropyl,6,6'-dimethyl-2,2'-biphenol. 1.5 g p-
5 toluenesulfonic acid was charged, and the mixture was heated to 130°C. After 7.5 hours, gas chromatography analysis showed the mixture contained 11.6% fully debutylated product, 3,3'-diisopropyl-6,6'-dimethyl-2,2'-biphenol; 2.7% mono-debutylated product, 5,-t-
10 butyl-3,3'-diisopropyl-6,6'-dimethyl-2,2'-biphenol; and 0.3% unreacted starting material.

Example 5


Bromination of 2,2'-dihydroxy-3,3'-diisopropyl-5,5'- 15 dimethylbiphenyl

Under an atmosphere of nitrogen, Br₂ (3.36 mL, 0.0652 mol) in CH₂Cl₂ (5 mL) was added dropwise to a CH₂Cl₂ (200 mL) solution of 2,2'-dihydroxy-3,3'-diisopropyl-5,5'-dimethylbiphenyl (6.488 g, 0.0217
20 mol). The resulting mixture was stirred at room temperature overnight. After the reaction was complete, the mixture was washed with 10% NaHSO₃ (3 x 50mL) followed by brine (2 x 50 mL) and dried over MgSO₄. The solvent was removed under vacuum to afford
25 an orange oil, which was purified by column chromatography (silica gel, 10% EtOAc/hexane). Yield of light-brown solid was 3.95 g (40%). ¹H NMR (C₆D₆): 1.07 (d, 6H), 1.89 (s, 3H), 3.17 (m, 1H) 4.30 (br s, 1H), 7.52 (s, 1H).

30

5 Claims:

1. A process for making a compound of the formula I



I

10 wherein

R¹ is C₁ to C₁₀ primary or secondary alkyl or cycloalkyl,


R² is C₁ to C₁₀ primary or secondary, alkyl or cycloalkyl, and

15 X is H, Cl, Br, or I,
comprising:

(1) when X is Cl


(a) chlorinating a compound of the formula

II



II

at the 4-position thereof to produce a
compound of the formula III



III

wherein X is Cl,

- (b) oxidatively coupling the compound of the formula III wherein X is Cl to produce a compound of the formula I wherein X is Cl;

(2) when X is H

- (a) chlorinating a compound of the formula II at the 4-position thereof to produce a compound of the formula III wherein X is Cl,

- (b) oxidatively coupling the compound of the formula III wherein X is Cl to produce a compound of the formula I wherein X is Cl, and

- (c) dechlorinating the compound of the formula I wherein X is Cl to produce a compound of the formula I wherein X is H; and

(3) when X is Br or I

- (a) chlorinating a compound of the formula II at the 4-position thereof to produce a compound of the formula III wherein X is Cl,


- (b) oxidatively coupling the compound of the formula III wherein X is Cl to produce a compound of the formula I wherein X is Cl,

- (c) dechlorinating the compound of the formula I wherein X is Cl to produce a compound of the formula I wherein X is H, and

- (d) substituting Br or I, respectively, for H at the 5 and 5' positions of the

compound of the formula I wherein X is
H.

- 5 2. The process of Claim 1 wherein the chlorinating
step is carried out by reacting the compound of formula
II with chlorine or sulfuryl chloride.
- 10 3. The process of claim 2 wherein the chlorinating
step is carried out in the presence of at least one
catalyst selected from the group consisting of aluminum
chloride and diaryl sulfide.
- 15 4. The process of claim 3 wherein the diaryl sulfide
is diphenyl sulfide.
- 20 5. The process of Claim 1 wherein the oxidative
coupling step is carried out by exposing the compound
of formula III to an iron (III) salt in a polar,
aprotic solvent.
6. The process of claim 5 wherein the iron (III) salt
is ferric chloride.
- 25 7. The process of Claim 1 wherein the dechlorinating
step is carried out by contacting the compound of
formula I wherein X is Cl with hydrogen, a formate salt
and a nickel or palladium-containing catalyst.
- 30 8. A process for making a compound of the formula IV



IV

wherein

5 R^1 is C_1 to C_{10} primary or secondary alkyl or cycloalkyl,


R^4 is C_1 to C_{10} primary or secondary alkyl or cycloalkyl, and

X is H, Cl, Br, or I

10 comprising:

(1) when X is H


(a) alkylating a compound of the formula V



V

15

at the 4-position thereof to produce a compound of the formula VI




VI

wherein R^3 is C_4 to C_{20} tertiary alkyl,

20

(b) oxidatively coupling the compound of the formula VI to produce a compound of the formula VII



VII

- 5 (c) dealkylating a compound of the formula VII to produce a compound of the formula IV wherein X is H; and
- (2) when X is Cl, Br, or I
- 10 (a) alkylating a compound of the formula V at the 4-position thereof to produce a compound of the formula VI,
- (b) oxidatively coupling the compound of the formula VI to produce a compound of the formula VII,
- 15 (c) dealkylating a compound of the formula VII to produce a compound of the formula IV wherein X is H, and
- (d) substituting Cl, Br, or I, respectively, for H at the 5 and 5' positions of the compound of the formula IV wherein X is
- 20 H.
9. The process of Claim 8 wherein the alkylating step is carried out by reacting the compound of the formula V with a tert-alkyl halide in the presence of a Lewis
- 25 Acid catalyst.
10. The process of claim 9 wherein the Lewis Acid catalyst is zinc chloride or aluminum chloride.

11. The process of claim 8 wherein the alkyating step
is carried out by reacting the compound of the formula
V with a 2,2-dialkylethylene in the presence of an acid
5 catalyst.

12. The process of claim 8 wherein the oxidative
coupling step is carried out by exposing the compound
of the formula VI to oxygen and a copper diamine
10 catalyst.


13. The process of claim 8 wherein the dealkylating
step is carried out by contacting the compound of the
formula VII with a strong acid.

15

14. The process of claim 13 wherein the strong acid is
selected from the group consisting of alkylsulfonic
acid, arylsulfonic acid, sulfuric acid, phosphoric
acid, and aluminum chloride.

20

15. A process for making a compound of the formula I



wherein


25 R^1 is C_1 to C_{10} primary or secondary alkyl or
cycloalkyl,

R^2 is C_1 to C_{10} primary or secondary alkyl or
cycloalkyl, and

X is H,

30 comprising:

- (a) oxidatively coupling a compound of the formula III



III

wherein X is Cl, to produce a compound of the formula I wherein X is Cl, and

- (b) dechlorinating the compound of the formula I wherein X is Cl to produce a compound of the formula I wherein X is H.

16. The process of Claim 15 wherein the oxidative coupling step is carried out by exposing the compound of formula III to an iron (III) salt in a polar, aprotic solvent.

17. The process of Claim 16 wherein the iron (III) salt is ferric chloride.

18. The process of Claim 15 wherein the dechlorinating step is carried out by contacting the compound of formula I wherein X is Cl with hydrogen, a formate salt and a nickel or palladium-containing catalyst.