## CLAIMS

What is claimed:

1. Use of a composition comprising a therapeutically effective amount of rapamycin, or a salt, solvate, enantiomer or diastereoisomer thereof in the manufacture of a medicament for the treatment of a dermal disorder in a mammalian subject in need thereof, wherein the dermal disorder is at least one selected form the group consisting of static wrinkles, fine wrinkles, loss of skin tone, ephelides, melasma, senile purpura, static rhytides, UV damage, prominent hand veins, field cancerization of epithelium such as in squamous cell carcinoma, keloids, and imbalance in ratio of collagen isoforms, and wherein the composition is to be topically administered to the subject.

2. The use of claim 1, wherein the field cancerization of epithelium comprises epidermal dysplasia and actinic damage.

3. The use of claim 1, wherein the subject is a human.

4. The use of claim 1, wherein the composition comprises about 0.1% to about 0.0001 % by weight of the rapamycin.

5. The use of claim 1, wherein the composition comprises about 0.001% to about 0.0001% by weight of the rapamycin.

6. The use of claim 1, wherein the composition further comprises a dermatologically acceptable carrier.

7. The use of claim 6, wherein the dermatologically acceptable carrier is at least one selected from the group consisting of a solvent, lubricant, emollient, emulsifier, moisturizer, thickening wax, softener, fragrance, preservative, and artificial color.

8. The use of claim 6, wherein the dermatologically acceptable carrier comprises petrolatum.

9. The use of claim 1, wherein the composition is applied to the site of the dermal disorder of the subject.

10. A kit for treating and/or preventing a dermal disorder in a mammalian subject in need thereof, the kit comprising: a composition comprising a therapeutically effective amount of rapamycin or a salt, solvate, enantiomer or diastereoisomer thereof, an applicator, and instructions for topically administering the composition to the subject, wherein the dermal disorder is at least one selected form the group consisting of: static wrinkles, fine wrinkles, loss of skin tone, ephelides, melasma, senile purpura, static rhytides, prominent hand veins, field cancerization of epithelium, keloids, and imbalance in ratio of collagen isoforms.

11. A topical composition comprising a therapeutically effective amount of rapamycin, or a salt, solvate, enantiomer or diastereoisomer thereof for use in therapy.

12. The topical composition for use of claim 11, wherein the therapeutically effective amount of rapamycin, or a salt, solvate, enantiomer or diastereoisomer thereof is 0.001% to 0.0001% by weight of the composition.

13. The topical composition for use of claim 11, further comprising a dermatologically acceptable carrier.