

HETEROCYCLIC COMPOUNDS AND THEIR USE AS GLYCOGEN SYNTHASE KINASE-3 INHIBITORS

Technical Field

5

The present invention relates to novel heterocyclic compounds which are useful for inhibiting glycogen synthase kinase 3 (GSK-3), methods of making the compounds, compositions containing the compounds, and methods of treatment using the compounds.

10

Background of the Invention

Glycogen synthase kinase-3 (GSK-3) is a serine/threonine kinase encoded by two isoforms, GSK-3 α and GSK-3 β , with molecular weights of 51 and 47 kDa, respectively.

15

These share 97% sequence similarity in their kinase catalytic domains. The GSK-3 α isoform has an extended glycine-rich N-terminal tail. A minor splice variant of GSK-3 β has been identified (expressed at ~15% of total) with a 13 amino acid insert within the kinase domain. This variant had a reduced activity towards tau. GSK-3 is highly conserved throughout evolution, and found in all mammals thus far with high homology in the kinase domain. Both isoforms are ubiquitously expressed in mammalian tissues, including the brain. Pharmacological GSK-3 inhibitors are not able to selectively inhibit one of the isoforms.

20

GSK-3 β plays an important role in the control of metabolism, differentiation and survival. It was initially identified as an enzyme able to phosphorylate and hence inhibit glycogen synthase. Subsequently, it was recognised that GSK-3 β was identical to tau protein kinase 1 (TPK1), an enzyme that phosphorylates tau protein in epitopes that are also found to be hyperphosphorylated in Alzheimer's disease and in several tauopathies.

25

30

Interestingly, protein kinase B (AKT) phosphorylation of GSK-3 β results in a loss of kinase activity, and it has been proposed that this inhibition may mediate some of the effects of neurotrophic factors. Moreover, phosphorylation of β -catenin (a protein involved in cell survival) by GSK-3 β , results in its degradation by an ubiquitination dependent proteasome pathway.

35

Therefore it appears that inhibition of GSK-3 β activity may result in neurotrophic activity. There is evidence that lithium, an uncompetitive inhibitor of GSK-3 β , enhances neurogenesis in some models and can also increase neuronal survival, through the induc-

tion of survival factors such as Bcl-2 and the inhibition of the expression of proapoptotic factors such as P53 and Bax.

Further studies have shown that β -amyloid increases GSK-3 β activity and tau protein phosphorylation. Moreover, this hyperphosphorylation as well as the neurotoxic effects of β -amyloid are blocked by lithium chloride and by a GSK-3 β antisense mRNA. These observations taken together suggest that GSK-3 β may be the link between the two major pathological processes in Alzheimer's disease: abnormal APP (Amyloid Precursor Protein) processing and tau protein hyperphosphorylation.

These experimental observations indicate that compounds which modulate the GSK-3 β activity may find application in the treatment of the neuropathological consequences and the cognitive and attention deficits associated with Alzheimer's disease, as well as other acute and chronic neurodegenerative diseases. These include, but are not limited to: behavioural and psychiatric symptoms of dementia, Parkinson's disease, tauopathies (e.g. frontotemporoparietal dementia, corticobasal degeneration, Pick's disease, progressive supranuclear palsy, argyophilic grain disease) and other dementia including vascular dementia; acute stroke and others traumatic injuries; cerebrovascular accidents (e.g. age related macular degeneration); brain and spinal cord trauma; peripheral neuropathies; bipolar disorders, retinopathies and glaucoma.

GSK-3 β may also have utility in the treatment of pain.

GSK-3 β may further have utility in the treatment of inflammatory diseases, such as rheumatoid arthritis and osteoarthritis.

GSK-3 β may also have utility in the treatment of other diseases such as: Non-insulin dependent diabetes and obesity; osteoporosis; manic depressive illness; schizophrenia; alopecia; cancers such as breast cancer, non-small cell lung carcinoma, thyroid cancer, T or B-cell leukemia and several virus-induced tumors.

A review on GSK-3, its functions, its therapeutic potential and its possible inhibitors is given in "Glycogen Synthase Kinase 3 (GSK-3) and its inhibitors: Drug Discovery and Developments" by A. Martinez et al. (editors), John Wiley and Sons, 2006.

WO 03/053330 describes 2-oxindoles substituted in the 3-position with a bicyclic hetaryl group and their use for treating conditions related to glycogen synthase kinase-3. WO 03/082853 describes substituted 2-oxindoles substituted in the 3-position with a monocyclic hetaryl group and their use for treating conditions related to glycogen syn-

thase kinase-3. WO 2005/123672 relates to 2-hydroxyindoles carrying in the 3-position an optionally fused pyrid-2-yl ring and their use for inhibiting kinases. WO 2005/061519 relates to 2-hydroxyindoles carrying in the 3-position a pyrid-2-yl ring fused to an aromatic or heteroaromatic ring and their use for inhibiting kinases.


5

Summary of the Invention

The object of the present invention is to provide compounds which modulate the GSK-3 β activity, in particular compounds which have an inhibitory activity on GSK-3 β and which thus are useful as an active ingredient of a composition for preventive and/or therapeutic treatment of a disease caused by abnormal GSK-3 β activity, especially of neurodegenerative and/or inflammatory diseases. More specifically, the goal is to provide novel compounds useful as an active ingredient of a composition that enables prevention and/or treatment of neurodegenerative diseases such as Alzheimer's disease.

15

It was surprisingly found that the problem is solved by providing a heterocyclic compound of the general formula I



20

the stereoisomers, prodrugs, N-oxides, tautomers and/or physiologically tolerated acid addition salts thereof, and the compounds of the general formula I, wherein at least one of the atoms has been replaced by its stable, non-radioactive isotope,

25

wherein

A is selected from the group consisting of CR^{A1}R^{A2} and NR^B; where

R^{A1} and R^{A2} are independently of each other selected from the group consisting of hydrogen, C₁-C₂-alkyl, C₁-C₂-haloalkyl, NH₂ and OH; and

30

R^B is selected from H, C₁-C₄-alkyl and C₁-C₄-haloalkyl;

X¹ and X² are independently of each other selected from the group consisting of CR² and N;

5 X³, X⁴, X⁵ and X⁶ are independently of each other selected from the group consisting of CR³, CR⁴ and N;

with the proviso that no more than two of X³, X⁴, X⁵ and X⁶ are CR⁴;

10 Y¹, Y², Y³ and Y⁴ are independently of each other selected from the group consisting of CR⁴, CR⁵ and N;

with the proviso that at most one of Y¹, Y², Y³ and Y⁴ is N and with the proviso that at most one of Y¹, Y², Y³ and Y⁴ is CR⁴; and

15 with the proviso that one of Y¹, Y², Y³ and Y⁴ is CR⁴ or C-CF₃, if none of X³, X⁴, X⁵ and X⁶ is CR⁴;

R¹ is selected from hydrogen, C₁-C₄-alkyl and C₁-C₄-haloalkyl;

20 each R² is independently selected from the group consisting of hydrogen, OH, halogen (preferably F or Cl, more preferably F), CN, C₁-C₆-alkyl, C₁-C₆-haloalkyl, C₁-C₆-alkoxy, C₁-C₆-haloalkoxy and NR^aR^b;
or two radicals R² bonded at the carbon atoms of groups X¹ and X², together with the carbon atoms to which they are bonded, form a 5- or 6-membered saturated
25 or unsaturated ring which may contain 1 or 2 heteroatoms as ring members selected from the group consisting of N, O and S and which optionally carries 1, 2 or 3 substituents R⁶;

30 each R³ is independently selected from the group consisting of hydrogen, CN, NR^aR^b, OH, halogen (preferably F or Cl, more preferably F), C₁-C₆-alkyl, C₁-C₆-haloalkyl, C₃-C₇-cycloalkyl, C₃-C₇-halocycloalkyl, C₂-C₄-alkenyl, C₂-C₄-haloalkenyl, C₁-C₆-alkoxy, C₁-C₆-haloalkoxy, formyl, C₁-C₆-alkylcarbonyl, C₁-C₆-haloalkylcarbonyl, C₁-C₆-alkoxycarbonyl, C₁-C₆-haloalkoxycarbonyl, C₁-C₆-alkyl-NR^aR^b and an aromatic radical Ar, which is selected from the group consisting of phenyl and a 5- or
35 6-membered N- or C-bound heteroaromatic radical comprising one nitrogen atom and optionally 1, 2 or 3 further heteroatoms independently selected from O, S and N as ring members, wherein Ar is unsubstituted or carries one or two radicals R⁷ and wherein Ar may also be bonded via a CH₂ group;

- 5 R^4 is a C-bound saturated or partially unsaturated monocyclic 3-, 4-, 5-, 6- or 7-membered heterocyclic ring containing 1, 2 or 3 heteroatoms or heteroatom-containing groups selected from O, N, S, NO, SO and SO_2 as ring members, where the heterocyclic ring optionally carries 1, 2 or 3 C- or N-bound substituents R^8 ;
- 10 R^5 is selected from the group consisting of hydrogen, CN, NR^aR^b , OH, halogen (preferably F or Cl, more preferably F), C_1 - C_6 -alkyl, C_1 - C_6 -haloalkyl, C_3 - C_7 -cycloalkyl, C_3 - C_7 -halocycloalkyl, C_2 - C_4 -alkenyl, C_2 - C_4 -haloalkenyl, C_1 - C_6 -alkoxy, C_1 - C_6 -haloalkoxy, formyl, C_1 - C_6 -alkylcarbonyl, C_1 - C_6 -haloalkylcarbonyl, C_1 - C_6 -alkoxycarbonyl, C_1 - C_6 -haloalkoxycarbonyl, C_1 - C_6 -alkyl- NR^aR^b and an aromatic radical Ar, which is selected from the group consisting of phenyl and a 5- or 6-membered N- or C-bound heteroaromatic radical comprising one nitrogen atom and optionally 1, 2 or 3 further heteroatoms independently selected from O, S and N as ring members, wherein Ar is unsubstituted or carries one or two radicals R^7 and wherein Ar may also be bonded via a CH_2 group;
- 20 R^6 and R^8 , independently of each other and independently of each occurrence, are selected from the group consisting of CN, NR^aR^b , OH, halogen (preferably F or Cl, more preferably F), C_1 - C_6 -alkyl, C_1 - C_6 -haloalkyl, C_3 - C_7 -cycloalkyl, C_3 - C_7 -halocycloalkyl, C_2 - C_4 -alkenyl, C_2 - C_4 -haloalkenyl, C_1 - C_6 -alkoxy, C_1 - C_6 -haloalkoxy, formyl, C_1 - C_6 -alkylcarbonyl, C_1 - C_6 -haloalkylcarbonyl, C_1 - C_6 -alkoxycarbonyl, C_1 - C_6 -haloalkoxycarbonyl, C_1 - C_6 -alkyl- NR^aR^b and an aromatic radical Ar, which is selected from the group consisting of phenyl and a 5- or 6-membered N- or C-bound heteroaromatic radical comprising one nitrogen atom and optionally 1, 2 or 3 further heteroatoms independently selected from O, S and N as ring members, wherein Ar is unsubstituted or carries one or two radicals R^7 and wherein Ar may also be bonded via a CH_2 group;
- 30 each R^7 is independently selected from the group consisting of halogen (preferably F or Cl, more preferably F), CN, C_1 - C_6 -alkyl, C_1 - C_6 -haloalkyl, C_3 - C_6 -cycloalkyl, C_3 - C_6 -halocycloalkyl, C_1 - C_6 -alkoxy, C_1 - C_6 -haloalkoxy, NR^aR^b , a phenyl group and a 5- or 6-membered heteroaromatic radical comprising one nitrogen atom and optionally 1, 2 or 3 further heteroatoms independently selected from O, S and N as ring members, wherein phenyl and the heteroaromatic radical are, independently of each other, unsubstituted or substituted by 1, 2, 3 or 4 radicals selected from halogen, cyano, C_1 - C_4 -alkyl, C_1 - C_4 -haloalkyl, C_1 - C_4 -alkoxy and C_1 - C_4 -haloalkoxy; and
- 35

R^a and R^b are independently of each other selected from the group consisting of hydrogen, C₁-C₆-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy, C₁-C₄-alkylcarbonyl and C₁-C₄-haloalkylcarbonyl;

5 or R^a and R^b form, together with the nitrogen atom to which they are bonded, a 3-, 4-, 5-, 6- or 7-membered saturated or unsaturated aromatic or non-aromatic N-heterocyclic ring, which may contain 1 further heteroatom or heteroatom containing group selected from the group consisting of O, S, SO, SO₂ and N as a ring member.

10 Thus, the present invention relates to compounds of the formula I as defined herein and in the claims, to the stereoisomers, tautomers, prodrugs and/or physiologically tolerated acid addition salts thereof, and also to compounds of the general formula I, wherein at least one of the atoms has been replaced by its stable, non-radioactive isotope.

15 According to a further aspect, the present invention relates to a pharmaceutical composition comprising at least one compound of the formula I as defined herein, a stereoisomer, a tautomer, a prodrug and/or a physiologically tolerated acid addition salt thereof or comprising at least one heterocyclic compound as defined above, wherein at least one of the atoms has been replaced by its stable, non-radioactive isotope, optionally together with at least one physiologically acceptable carrier and/or auxiliary substance.

25 According to a further aspect, the present invention relates to the use of at least one compound of the formula I as defined herein, the stereoisomers, tautomers, prodrugs and/or physiologically tolerated acid addition salts thereof, for the preparation of a medicament for the treatment of a medical disorder susceptible to treatment with a compound that modulates glycogen synthase kinase 3 β activity.

30 According to a further aspect, the present invention relates to a method for treating a medical disorder susceptible to treatment with a compound that modulates glycogen synthase kinase 3 β activity, said method comprising administering an effective amount of at least one compound of the formula I as defined herein, a stereoisomer, a tautomer, a prodrug and/or a physiologically tolerated acid addition salt thereof, to a subject in need thereof.

Detailed description of the invention


Provided the compounds of the formula I of a given constitution may exist in different spatial arrangements, for example if they possess one or more centers of asymmetry, polysubstituted rings or double bonds, or as different tautomers, it is also possible to use enantiomeric mixtures, in particular racemates, diastereomeric mixtures and
 5 tautomeric mixtures, preferably, however, the respective essentially pure enantiomers, diastereomers and tautomers of the compounds of formula I and/or of their salts.

It is likewise possible to use physiologically tolerated salts of the compounds of the formula I, especially acid addition salts with physiologically tolerated acids. Examples
 10 of suitable physiologically tolerated organic and inorganic acids are hydrochloric acid, hydrobromic acid, phosphoric acid, sulfuric acid, C₁-C₄-alkylsulfonic acids, such as methanesulfonic acid, aromatic sulfonic acids, such as benzenesulfonic acid and toluenesulfonic acid, oxalic acid, maleic acid, fumaric acid, lactic acid, tartaric acid, adipic acid and benzoic acid. Other utilizable acids are described in Fortschritte der Arzneimittelforschung [Advances in drug research], Volume 10, pages 224 et seq., Birkhäuser
 15 Verlag, Basel and Stuttgart, 1966.

In the terms of the present invention, "prodrugs" are compounds which are metabolized in vivo to give the compounds of the invention of formula I. Typical examples for prodrugs are for example described in C.G. Wermeth (editor): The Practice of Medicinal Chemistry, Academic Press, San Diego, 1996, pages 671-715. Examples are phosphates, carbamates, aminoacids, esters, amides, peptides, urea and the like. In the present case, suitable prodrugs can be compounds of formula I wherein an external nitrogen atom, for example a secondary nitrogen ring atom of the ring R⁴ or a nitrogen
 20 atom of a primary or secondary amino group being a substituent R², R³, R⁵, R⁶, R⁷ and/or R⁸ (= at least one of R², R³, R⁵, R⁶, R⁷ and R⁸ is NR^aR^b, wherein at least one of R^a and R^b is H), forms an amide/peptide bond in that this nitrogen atom is substituted by a C₁-C₄-alkylcarbonyl group, e.g. by acetyl, propionyl, n-propylcarbonyl, isopropylcarbonyl, n-butylcarbonyl or tert-butylcarbonyl (pivaloyl), by benzoyl, or by an amino-
 25 acid group bonded via CO, e.g. glycine, alanine, serine, phenylalanine and the like bonded via CO. Suitable prodrugs are furthermore alkylcarbonyloxyalkylcarbamates, wherein said nitrogen atom carries a group -C(=O)-O-CHR^x-O-C(=O)-R^y, wherein R^x und R^y independently of each other are C₁-C₄-alkyl. These carbamate compounds are for example described in J. Alexander, R. Cargill, S. R. Michelson, H. Schwam, J. Medicinal Chem. 1988, 31(2), 318-322. These groups can be removed under metabolic
 30 conditions and result in compounds I wherein said nitrogen atom carries a hydrogen atom instead. Also, R⁸, if bound to a nitrogen ring atom of R⁴, may be chosen so as to be hydrolysable under metabolic conditions and thus to be one of the above-listed

groups (i.e. a C₁-C₄-alkylcarbonyl group, an aminoacid group bonded via CO or a group -C(=O)-O-CHR^x-O-C(=O)-R^y).

- 5 The compounds of formula I may also be present in the form of the respective tautomers. Tautomerism may be present in compounds I wherein R² or R³ is OH and this substituent is bonded to a carbon atom which is in α -position to a nitrogen ring atom. This results for example in following tautomeric formulae:



The organic moieties mentioned in the above definitions of the variables are - like the term halogen - collective terms for individual listings of the individual group members.

The prefix C_n-C_m indicates in each case the possible number of carbon atoms in the group.

5 The term halogen denotes in each case fluorine, bromine, chlorine or iodine, in particular fluorine, chlorine or bromine.

C₁-C₂-Alkyl is methyl or ethyl; C₁-C₃-alkyl is additionally n-propyl or isopropyl.

10 C₁-C₄-Alkyl is a straight-chain or branched alkyl group having from 1 to 4 carbon atoms. Examples are methyl, ethyl, n-propyl, isopropyl, n-butyl, 2-butyl (sec-butyl), isobutyl and tert-butyl.

15 C₁-C₆-Alkyl is a straight-chain or branched alkyl group having from 1 to 6 carbon atoms. Examples include the residues mentioned above for C₁-C₄-alkyl and also pentyl, 1-methylbutyl, 2-methylbutyl, 3-methylbutyl, 2,2-dimethylpropyl, 1-ethylpropyl, hexyl, 1,1-dimethylpropyl, 1,2-dimethylpropyl, 1-methylpentyl, 2-methylpentyl, 3-methylpentyl, 4-methylpentyl, 1,1-dimethylbutyl, 1,2-dimethylbutyl, 1,3-dimethylbutyl, 2,2-dimethylbutyl, 2,3-dimethylbutyl, 3,3-dimethylbutyl, 1-ethylbutyl, 2-ethylbutyl, 1,1,2-trimethylpropyl, 1,2,2-trimethylpropyl, 1-ethyl-1-methylpropyl and 1-ethyl-2-methylpropyl.

20

C₁-C₂-Haloalkyl is an alkyl group having 1 or 2 carbon atoms (as mentioned above), where at least one of the hydrogen atoms, e.g. 1, 2, 3, 4 or 5 hydrogen atoms in these groups are replaced by halogen atoms as mentioned above, such as chloromethyl, dichloromethyl, trichloromethyl, fluoromethyl, difluoromethyl, trifluoromethyl, bromomethyl, chlorofluoromethyl, dichlorofluoromethyl, chlorodifluoromethyl, 1-chloroethyl, 1-bromoethyl, 1-fluoroethyl, 2-chloroethyl, 2-bromoethyl, 2-fluoroethyl, 2,2-difluoroethyl, 2,2,2-trifluoroethyl, 2-chloro-2-fluoroethyl, 2-chloro-2,2-difluoroethyl, 2,2-dichloro-2-fluoroethyl, 2,2,2-trichloroethyl or pentafluoroethyl.

25

30 C₁-C₄-Haloalkyl is a straight-chain or branched alkyl group having 1 to 4 carbon atoms (as mentioned above), where at least one of the hydrogen atoms, e.g. 1, 2, 3, 4 or 5 hydrogen atoms in these groups are replaced by halogen atoms as mentioned above. Examples are, apart those listed above for C₁-C₂-haloalkyl, 1-chloropropyl, 1-bromopropyl, 1-fluoropropyl, 2-chloropropyl, 2-bromopropyl, 2-fluoropropyl, 3-chloropropyl, 3-bromopropyl, 3-fluoropropyl, 1,1-dichloropropyl, 1,1-difluoropropyl, 2,2-dichloropropyl, 2,2-difluoropropyl, 2,3-dichloropropyl, 2,3-difluoropropyl, 1,3-dichloropropyl, 1,3-difluoropropyl, 3,3-dichloropropyl, 3,3-difluoropropyl, 1,1,2-trichloropropyl, 1,1,2-trifluoropropyl, 1,2,2-trichloropropyl, 1,2,2-trifluoropropyl, 1,2,3-

35

trichloropropyl, 1,2,3-trifluoropropyl, 2,2,3-trichloropropyl, 2,2,3-trifluoropropyl, 3,3,3-trichloropropyl, 3,3,3-trifluoropropyl, 1,1,1-trifluoroprop-2-yl, 1-chlorobutyl, 1-bromobutyl, 1-fluorobutyl, 2-chlorobutyl, 2-bromobutyl, 2-fluorobutyl, 3-chlorobutyl, 3-bromobutyl, 3-fluorobutyl, 4-chlorobutyl, 4-bromobutyl, 4-fluorobutyl, and the like.

5

C₁-C₆-Haloalkyl is a straight-chain or branched alkyl group having 1 to 6 carbon atoms (as mentioned above), where at least one of the hydrogen atoms in these groups is replaced by halogen atoms as mentioned above. Examples are, apart those listed above for C₁-C₄-haloalkyl, chloropentyl, bromopentyl, fluoropentyl, chlorohexyl, bromohexyl, fluorohexyl, and the like.

10

C₁-C₂-Fluoroalkyl (= fluorinated C₁-C₂-alkyl) is an alkyl group having 1 or 2 carbon atoms (as mentioned above), where at least one of the hydrogen atoms, e.g. 1, 2, 3, 4 or 5 hydrogen atoms in these groups are replaced by fluorine atoms, such as difluoromethyl, trifluoromethyl, 1-fluoroethyl, (R)-1-fluoroethyl, (S)-1-fluoroethyl, 2-fluoroethyl, 2,2-difluoroethyl, 2,2,2-trifluoroethyl, or pentafluoroethyl.

15

C₁-C₄-Fluoroalkyl (= fluorinated C₁-C₄-alkyl) is a straight-chain or branched alkyl group having 1 to 4 carbon atoms (as mentioned above), where at least one of the hydrogen atoms, e.g. 1, 2, 3, 4 or 5 hydrogen atoms in these groups are replaced by fluorine atoms. Examples are, apart those listed above for C₁-C₂-fluoroalkyl, 1-fluoropropyl, (R)-1-fluoropropyl, (S)-1-fluoropropyl, 2-fluoropropyl, (R)-2-fluoropropyl, (S)-2-fluoropropyl, 3-fluoropropyl, 1,1-difluoropropyl, 2,2-difluoropropyl, 1,2-difluoropropyl, 2,3-difluoropropyl, 1,3-difluoropropyl, 3,3-difluoropropyl, 1,1,2-trifluoropropyl, 1,2,2-trifluoropropyl, 1,2,3-trifluoropropyl, 2,2,3-trifluoropropyl, 3,3,3-trifluoropropyl, 1,1,1-trifluoroprop-2-yl, 2-fluoro-1-methylethyl, (R)-2-fluoro-1-methylethyl, (S)-2-fluoro-1-methylethyl, 2,2-difluoro-1-methylethyl, (R)-2,2-difluoro-1-methylethyl, (S)-2,2-difluoro-1-methylethyl, 1,2-difluoro-1-methylethyl, (R)-1,2-difluoro-1-methylethyl, (S)-1,2-difluoro-1-methylethyl, 2,2,2-trifluoro-1-methylethyl, (R)-2,2,2-trifluoro-1-methylethyl, (S)-2,2,2-trifluoro-1-methylethyl, 2-fluoro-1-(fluoromethyl)ethyl, 1-(difluoromethyl)-2,2-difluoroethyl, 1-(trifluoromethyl)-2,2,2-trifluoroethyl, 1-(trifluoromethyl)-1,2,2,2-tetrafluoroethyl, 1-fluorobutyl, (R)-1-fluorobutyl, (S)-1-fluorobutyl, 2-fluorobutyl, (R)-2-fluorobutyl, (S)-2-fluorobutyl, 3-fluorobutyl, (R)-3-fluorobutyl, (S)-3-fluorobutyl, 4-fluorobutyl, 1,1-difluorobutyl, 2,2-difluorobutyl, 3,3-difluorobutyl, 4,4-difluorobutyl, 4,4,4-trifluorobutyl and the like.

20

25

30

35

C₁-C₆-Fluoroalkyl (= fluorinated C₁-C₆-alkyl) is a straight-chain or branched alkyl group having 1 to 6 carbon atoms (as mentioned above), where at least one of the hydrogen atoms, e.g. 1, 2, 3, 4 or 5 hydrogen atoms in these groups are replaced by fluorine at-

- oms. Examples are, apart those listed above for C₁-C₄-fluoroalkyl, 1-fluoropentyl, (R)-1-fluoropentyl, (S)-1-fluoropentyl, 2-fluoropentyl, (R)-2-fluoropentyl, (S)-2-fluoropentyl, 3-fluoropentyl, (R)-3-fluoropentyl, (S)-3-fluoropentyl, 4-fluoropentyl, (R)-4-fluoropentyl, (S)-4-fluoropentyl, 5-fluoropentyl, (R)-5-fluoropentyl, (S)-5-fluoropentyl, 1-fluorohexyl, (R)-1-fluorohexyl, (S)-1-fluorohexyl, 2-fluorohexyl, (R)-2-fluorohexyl, (S)-2-fluorohexyl, 3-fluorohexyl, (R)-3-fluorohexyl, (S)-3-fluorohexyl, 4-fluorohexyl, (R)-4-fluorohexyl, (S)-4-fluorohexyl, 5-fluorohexyl, (R)-5-fluorohexyl, (S)-5-fluorohexyl, 6-fluorohexyl, (R)-6-fluorohexyl, (S)-6-fluorohexyl, and the like.
- 10 C₁-C₄-Alkoxy is a straight-chain or branched alkyl group having from 1 to 4 carbon atoms, which is bound to the remainder of the molecule via an oxygen atom. Examples include methoxy, ethoxy, n-propoxy, isopropoxy, n-butoxy, 2-butoxy, isobutoxy and tert-butoxy.
- 15 C₁-C₆-Alkoxy is a straight-chain or branched alkyl group having from 1 to 6 carbon atoms, which is bound to the remainder of the molecule via an oxygen atom. Examples include, apart those listed above for C₁-C₄-alkoxy, pentyloxy, 1-methylbutoxy, 2-methylbutoxy, 3-methylbutoxy, 2,2-dimethylpropoxy, 1-ethylpropoxy, hexyloxy, 1,1-dimethylpropoxy, 1,2-dimethylpropoxy, 1-methylpentyloxy, 2-methylpentyloxy, 3-methylpentyloxy, 4-methylpentyloxy, 1,1-dimethylbutoxy, 1,2-dimethylbutoxy, 1,3-dimethylbutoxy, 2,2-dimethylbutoxy, 2,3-dimethylbutoxy, 3,3-dimethylbutoxy, 1-ethylbutoxy, 2-ethylbutoxy, 1,1,2-trimethylpropoxy, 1,2,2-trimethylpropoxy, 1-ethyl-1-methylpropoxy and 1-ethyl-2-methylpropoxy.
- 25 Halogenated C₁-C₆-alkoxy (which is also termed C₁-C₆-haloalkoxy), in particular fluorinated C₁-C₆-alkoxy (also termed C₁-C₆-fluoroalkoxy) is a straight-chain or branched alkoxy group having from 1 to 6, in particular 1 to 4 carbon atoms (= fluorinated C₁-C₄-alkoxy), wherein at least one, e.g. 1, 2, 3, 4 or all of the hydrogen atoms are replaced by a halogen atoms, in particular fluorine atoms such as in fluoromethoxy, difluoromethoxy, trifluoromethoxy, (R)-1-fluoroethoxy, (S)-1-fluoroethoxy, 2-fluoroethoxy, 1,1-difluoroethoxy, 2,2-difluoroethoxy, 2,2,2-trifluoroethoxy, 1,1,2,2-tetrafluoroethoxy, (R)-1-fluoropropoxy, (S)-1-fluoropropoxy, (R)-2-fluoropropoxy, (S)-2-fluoropropoxy, 3-fluoropropoxy, 1,1-difluoropropoxy, 2,2-difluoropropoxy, 3,3-difluoropropoxy, 3,3,3-trifluoropropoxy, (R)-2-fluoro-1-methylethoxy, (S)-2-fluoro-1-methylethoxy, (R)-2,2-difluoro-1-methylethoxy, (S)-2,2-difluoro-1-methylethoxy, (R)-1,2-difluoro-1-methylethoxy, (S)-1,2-difluoro-1-methylethoxy, (R)-2,2,2-trifluoro-1-methylethoxy, (S)-2,2,2-trifluoro-1-methylethoxy, 2-fluoro-1-(fluoromethyl)ethoxy, 1-(difluoromethyl)-2,2-difluoroethoxy, (R)-1-fluorobutoxy, (S)-1-fluorobutoxy, 2-fluorobutoxy, 3-fluorobutoxy, 4-

fluorobutoxy, 1,1-difluorobutoxy, 2,2-difluorobutoxy, 3,3-difluorobutoxy, 4,4-difluorobutoxy, 4,4,4-trifluorobutoxy, and the like.

5 C₁-C₄-Alkylcarbonyl is a straight-chain or branched alkyl group having from 1 to 4 carbon atoms), which is bound to the remainder of the molecule via a carbonyl group (CO), such as in acetyl, propionyl, isopropylcarbonyl, butylcarbonyl, sec-butylcarbonyl, isobutylcarbonyl, and tert-butylcarbonyl.

10 C₁-C₆-Alkylcarbonyl is a straight-chain or branched alkyl group having from 1 to 6 carbon atoms, which is bound to the remainder of the molecule via a carbonyl group (CO). Examples include, apart those listed above for C₁-C₄-alkylcarbonyl, pentylcarbonyl, hexylcarbonyl and the constitutional isomers thereof.

15 C₁-C₄-Haloalkylcarbonyl is a straight-chain or branched haloalkyl group having from 1 to 4 carbon atoms as defined above, which is bound to the remainder of the molecule via a carbonyl group (CO)

20 C₁-C₆-Haloalkylcarbonyl is a straight-chain or branched haloalkyl group having from 1 to 6 carbon atoms as defined above, which is bound to the remainder of the molecule via a carbonyl group (CO)

25 C₁-C₄-Fluoroalkylcarbonyl is a straight-chain or branched fluoroalkyl group having from 1 to 4 carbon atoms as defined above, which is bound to the remainder of the molecule via a carbonyl group (CO)

C₁-C₆-fluoroalkylcarbonyl is a straight-chain or branched fluoroalkyl group having from 1 to 6 carbon atoms as defined above, which is bound to the remainder of the molecule via a carbonyl group (CO)

30 C₁-C₆-Alkoxy carbonyl is a straight-chain or branched alkoxy group having from 1 to 6, especially 1 to 4 carbon atoms (= C₁-C₄-alkoxy carbonyl), in particular 1 to 3 carbon atoms (= C₁-C₃-alkoxy carbonyl), which is bound to the remainder of the molecule via a carbonyl group (CO), such as in methoxy carbonyl, ethoxy carbonyl, propyloxy carbonyl, and isopropyloxy carbonyl.

35 C₁-C₆-Haloalkoxy carbonyl is a straight-chain or branched haloalkoxy group having from 1 to 6, especially 1 to 4 carbon atoms (= C₁-C₄-haloalkoxy carbonyl), in particular 1 to 3 carbon atoms (= C₁-C₃-haloalkoxy carbonyl) as defined above, which is bound to the remainder of the molecule via a carbonyl group (CO).

C₁-C₆-Fluoroalkoxycarbonyl is a straight-chain or branched fluoroalkoxy group having from 1 to 6, especially 1 to 4 carbon atoms (= C₁-C₄-fluoroalkoxycarbonyl), in particular 1 to 3 carbon atoms (= C₁-C₃-fluoroalkoxycarbonyl) as defined above, which is bound to the remainder of the molecule via a carbonyl group (CO).

C₃-C₆-Cycloalkyl is a cycloaliphatic radical having from 3 to 6 C atoms, such as cyclopropyl, cyclobutyl, cyclopentyl and cyclohexyl. C₃-C₄-cycloalkyl is a cycloaliphatic radical having from 3 to 4 C atoms, such as cyclopropyl and cyclobutyl.

C₃-C₇-Cycloalkyl is a cycloaliphatic radical having from 3 to 7 C atoms, such as cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and cycloheptyl.

C₃-C₆-Halocycloalkyl is a cycloaliphatic radical having from 3 to 6 C atoms, such as cyclopropyl, cyclobutyl, cyclopentyl and cyclohexyl, wherein at least one, e.g. 1, 2, 3, 4 or all of the hydrogen atoms are replaced by a halogen atoms, preferably by fluorine atoms such as in 1-fluorocyclopropyl, 2-fluorocyclopropyl, (S)- and (R)-2,2-difluorocyclopropyl, 1,2-difluorocyclopropyl, 2,3-difluorocyclopropyl, pentafluorocyclopropyl, 1-fluorocyclobutyl, 2-fluorocyclobutyl, 3-fluorocyclobutyl, 2,2-difluorocyclobutyl, 3,3-difluorocyclobutyl, 1,2-difluorocyclobutyl, 1,3-difluorocyclobutyl, 2,3-difluorocyclobutyl, 2,4-difluorocyclobutyl, or 1,2,2-trifluorocyclobutyl.

C₃-C₇-Halocycloalkyl is a cycloaliphatic radical having from 3 to 7 C atoms, such as cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and cycloheptyl, wherein at least one, e.g. 1, 2, 3, 4 or all of the hydrogen atoms are replaced by a halogen atoms, preferably by fluorine atoms. Examples include, apart those listed above for C₃-C₆-fluorocycloalkyl, 1-fluorocycloheptyl, 2-fluorocycloheptyl, 3-fluorocycloheptyl, 4-fluorocycloheptyl, 1,2-difluorocycloheptyl, 1,3-difluorocycloheptyl, 1,4-difluorocycloheptyl, 2,2-difluorocycloheptyl, 2,3-difluorocycloheptyl, 2,4-difluorocycloheptyl, 2,5-difluorocycloheptyl, 2,6-difluorocycloheptyl, 2,7-difluorocycloheptyl, 3,3-difluorocycloheptyl, 3,4-difluorocycloheptyl, 3,5-difluorocycloheptyl, 3,6-difluorocycloheptyl, 4,4-difluorocycloheptyl, 4,5-difluorocycloheptyl, and the like.

C₂-C₄-Alkenyl is a singly unsaturated hydrocarbon radical having 2, 3 or 4 C-atoms and one C-C double bond, e.g. vinyl, allyl (2-propen-1-yl), 1-propen-1-yl, 2-propen-2-yl, buten-1-yl, buten-2-yl, buten-3-yl, methallyl (2-methylprop-2-en-1-yl) and the like.

C₂-C₄-Haloalkenyl is a singly unsaturated hydrocarbon radical having 2, 3 or 4 C-atoms, wherein at least one, e.g. 1, 2, 3, 4 or all of the hydrogen atoms are replaced by halogen atoms, preferably by fluorine atoms such as in 1-fluorovinyl, 2-fluorovinyl, 2,2-fluorovinyl, 3,3,3-fluoropropenyl, 1,1-difluoro-2-propenyl, 1-fluoro-2-propenyl and the like.

Examples for 5- or 6-membered N- or C-bound heteroaromatic radicals comprising one nitrogen atom and optionally 1, 2 or 3 further heteroatoms independently selected from O, S and N as ring members are pyrrol-1-yl, pyrrol-2-yl, pyrrol-3-yl, pyrazol-1-yl, pyrazol-3-yl, pyrazol-4-yl, pyrazol-5-yl, imidazol-1-yl, imidazol-2-yl, imidazol-4-yl, imidazol-5-yl, oxazol-2-yl, oxazol-4-yl, oxazol-5-yl, isoxazol-3-yl, isoxazol-4-yl, isoxazol-5-yl, thiazol-2-yl, thiazol-4-yl, thiazol-5-yl, isothiazol-3-yl, isothiazol-4-yl, isothiazol-5-yl, [1,2,3]-1H-triazol-1-yl, [1,2,3]-1H-triazol-4-yl, [1,2,3]-1H-triazol-5-yl, [1,2,3]-2H-triazol-2-yl, [1,2,3]-2H-triazol-4-yl, [1,2,3]-2H-triazol-5-yl, [1,2,4]-1H-triazol-1-yl, [1,2,4]-1H-triazol-3-yl, [1,2,4]-1H-triazol-5-yl, [1,2,4]-4H-triazol-3-yl, [1,2,4]-4H-triazol-4-yl, oxadiazolyl, thiadiazolyl, [1,2,3,4]-1H-tetrazol-1-yl, [1,2,3,4]-1H-tetrazol-5-yl, [1,2,3,4]-2H-tetrazol-2-yl, [1,2,3,4]-2H-tetrazol-5-yl, pyridin-2-yl, pyridin-3-yl, pyridin-4-yl, pyridazin-3-yl, pyridazin-4-yl, pyrimidin-2-yl, pyrimidin-4-yl, pyrimidin-5-yl, pyrazin-2-yl and triazin-2-yl.


Examples for 5- or 6-membered N- or C-bound heteroaromatic radicals comprising 1, 2 or 3 heteroatoms independently selected from O, S and N as ring members are furan-2-yl, furan-3-yl, thien-2-yl, thien-3-yl, pyrrol-1-yl, pyrrol-2-yl, pyrrol-3-yl, pyrazol-1-yl, pyrazol-3-yl, pyrazol-4-yl, pyrazol-5-yl, imidazol-1-yl, imidazol-2-yl, imidazol-4-yl, imidazol-5-yl, oxazol-2-yl, oxazol-4-yl, oxazol-5-yl, isoxazol-3-yl, isoxazol-4-yl, isoxazol-5-yl, thiazol-2-yl, thiazol-4-yl, thiazol-5-yl, isothiazol-3-yl, isothiazol-4-yl, isothiazol-5-yl, [1,2,3]-1H-triazol-1-yl, [1,2,3]-1H-triazol-4-yl, [1,2,3]-1H-triazol-5-yl, [1,2,3]-2H-triazol-2-yl, [1,2,3]-2H-triazol-4-yl, [1,2,3]-2H-triazol-5-yl, [1,2,4]-1H-triazol-1-yl, [1,2,4]-1H-triazol-3-yl, [1,2,4]-1H-triazol-5-yl, [1,2,4]-4H-triazol-3-yl, [1,2,4]-4H-triazol-4-yl, oxadiazolyl, thiadiazolyl, [1,2,3,4]-1H-tetrazol-1-yl, [1,2,3,4]-1H-tetrazol-5-yl, [1,2,3,4]-2H-tetrazol-2-yl, [1,2,3,4]-2H-tetrazol-5-yl, pyridin-2-yl, pyridin-3-yl, pyridin-4-yl, pyridazin-3-yl, pyridazin-4-yl, pyrimidin-2-yl, pyrimidin-4-yl, pyrimidin-5-yl, pyrazin-2-yl and triazin-2-yl.


Examples for N-bound 3-, 4-, 5-, 6- or 7-membered saturated or unsaturated aromatic or non-aromatic N-heterocyclic rings, which may contain 1 further heteroatom or heteroatom-containing group selected from the group consisting of O, S, SO, SO₂ and N as a ring member (thus as rings formed by R^a and R^b together with the nitrogen atom to which they are bound), are aziridin-1-yl, azetidin-1-yl, pyrrolidin-1-yl, pyrazolidin-1-yl, imidazolidin-1-yl, oxazolidin-3-yl, isoxazolidin-2-yl, thiazolidin-3-yl, isothiazolidin-1-yl,

[1,2,3]-triazolidin-1-yl, [1,2,3]-triazolidin-2-yl, [1,2,4]-triazolidin-1-yl, [1,2,4]-triazolidin-4-yl, piperidin-1-yl, piperazin-1-yl, morpholin-4-yl, thiomorpholin-1-yl, 1-oxohiomorpholin-1-yl, 1,1-dioxothiomorpholin-1-yl, azepan-1-yl, azirin-1-yl, azetin-1-yl, pyrrolin-1-yl, pyrazolin-1-yl, imidazolin-1-yl, oxazolin-3-yl, isoxazolin-2-yl, thiazolin-3-yl, isothiazolin-1-yl, 1,2-dihydropyridin-1-yl, 1,2,3,4-tetrahydropyridin-1-yl, 1,2,5,6-tetrahydropyridin-1-yl, 1,2-dihydropyridazin, 1,6-dihydropyridazin, 1,2,3,4-tetrahydropyridazin-1-yl, 1,2,5,6-tetrahydropyridazin-1-yl, 1,2-dihydropyrimidin, 1,6-dihydropyrimidin, 1,2,3,4-tetrahydropyrimidin-1-yl, 1,2,5,6-tetrahydropyrimidin-1-yl, 1,2-dihydropyrazin-1-yl, 1,2,3,4-tetrahydropyrazin-1-yl, 1,2,5,6-tetrahydropyrazin-1-yl, pyrrol-1-yl, pyrazol-1-yl, imidazol-1-yl, [1,2,3]-1H-triazol-1-yl, [1,2,3]-2H-triazol-2-yl, [1,2,4]-1H-triazol-1-yl and [1,2,4]-4H-triazol-4-yl.


Examples for C-bound saturated or partially unsaturated monocyclic 3-, 4-, 5-, 6- or 7-membered heterocyclic ring containing 1, 2 or 3 heteroatoms or heteroatom-containing groups selected from O, N, S, NO, SO and SO₂ as ring members (i.e. for rings R⁴) are C-bound oxiranyl, thiiranyl, aziridinyl, oxetanyl, azetidiny, dihydrofuranyl, tetrahydrofuranyl, dihydrothienyl, tetrahydrothienyl, dihydrothienyl-1-oxide, tetrahydrothienyl-1-oxide, pyrrolinyl, pyrrolidinyl, pyrazolinyl, pyrazolidinyl, imidazolinyl, imidazolidinyl, triazolinyl, triazolidinyl, oxazolinyl, oxazolidinyl, isoxazolinyl isoxazolidinyl, oxadiazolinyl, oxadiazolidinyl, thiazolinyl, thiazolidinyl, isothiazolinyl, isothiazolidinyl, thiadiazolinyl, thiadiazolidinyl, dihydropyranyl, tetrahydropyranyl, dihydrothiopyranyl, tetrahydrothiopyranyl, dihydropyridinyl, tetrahydropyridinyl, piperidinyl, dihydropyridazinyl, tetrahydropyridazinyl, hexahydropyridazinyl, dihydropyrimidinyl, tetrahydropyrimidinyl, hexahydropyrimidinyl, dihydropyrazinyl, tetrahydropyrazinyl, piperazinyl, dihydro-[1,4]-oxazinyl, morpholinyl, dihydro-[1,4]-thiazinyl, thiomorpholinyl, thiomorpholinyl-1-oxide, thiomorpholinyl-1-dioxide, dihydroazepinyl, tetrahydroazepinyl, azepanyl, dihydrodiazepinyl, tetrahydrodiazepinyl, diazepanyl and the like.

More precisely, examples for C-bound saturated or partially unsaturated monocyclic 3-, 4-, 5-, 6- or 7-membered heterocyclic ring containing 1, 2 or 3 heteroatoms or heteroatom-containing groups selected from O, N, S, NO, SO and SO₂ as ring members (i.e. for rings R⁴) comprise the following structures:







5





10





15





19






5





10






15








5






10





15






21



5






10





15




22



5






10





15



23



5



10



where # is the attachment point to the remainder of the molecule.

15

The remarks made above and in the following with respect to preferred aspects of the invention, e.g. to preferred meanings of the variables X^1 , X^2 , X^3 , X^4 , X^5 , X^6 , Y^1 , Y^2 , Y^3 , Y^4 , R^1 , R^2 , R^3 , R^4 , R^5 , R^6 , R^7 , R^8 , R^a , R^b , R^{A1} , R^{A2} , R^B of compounds I, to preferred

compounds I and to preferred embodiments of the method or the use according to the invention, apply in each case on their own or in particular to combinations thereof.

5 In one embodiment of the invention one of Y^1 , Y^2 , Y^3 and Y^4 is CR^4 if none of X^3 , X^4 , X^5 and X^6 is CR^4 .

In another embodiment of the invention one of Y^1 , Y^2 , Y^3 and Y^4 is $C-CF_3$ if none of X^3 , X^4 , X^5 and X^6 is CR^4 .

10 In a preferred embodiment of the invention, either one of X^3 , X^4 , X^5 and X^6 is CR^4 and none of Y^1 , Y^2 , Y^3 and Y^4 is CR^4 , or one of Y^1 , Y^2 , Y^3 and Y^4 is CR^4 and none of X^3 , X^4 , X^5 and X^6 is CR^4 . In other words, either the condensed ring system (with X^1 - X^6 as ring members) carries one group R^4 or the ring with Y^1 - Y^4 as ring members carries one group R^4 .

15 In an alternatively preferred embodiment of the invention, one of Y^1 , Y^2 , Y^3 and Y^4 is $C-CF_3$ and none of X^3 , X^4 , X^5 and X^6 is CR^4 .


20 If the condensed ring system (with X^1 - X^6 as ring members) carries one group R^4 , this is preferably bound in the position of X^4 or X^5 .


If the ring with Y^1 - Y^4 as ring members carries one group R^4 , this is preferably bound in the position of Y^4 .


25 R^4 is preferably selected from a C-bound saturated or partially unsaturated monocyclic 4-, 5- or 6-membered heterocyclic ring containing 1 or 2 or 3 heteroatoms selected from O, N, S and SO, as ring members, where the heterocyclic ring optionally carries 1, 2 or 3 substituents R^8 .

30 More preferably, R^4 is selected from C-bound oxetanyl, thietanyl, azetidiny, tetrahydrofuranyl, tetrahydrothienyl, tetrahydrothienyl-1-oxide, pyrrolidiny, pyrrolinyl, pyrazolidinyl, pyrazolinyl, imidazolidinyl, imidazolinyl, tetrahydropyranyl, dihydropyranyl, piperidinyl, tetrahydropyridinyl, dihydropyridinyl, piperazinyl and morpholinyl, where the heterocyclic ring optionally carries 1, 2 or 3 substituents R^8 .


35 Preferably, these cyclic structures have the following formulae:








5



10




where the above cyclic structures optionally carry 1, 2 or 3 substituents R^8 , where R^8 may be bound to a carbon ring atom or to a nitrogen ring atom, and where # is the attachment point to the remainder of the molecule.


15

Among the above partially saturated rings preference is given to those having one or more C=C bonds and where one carbon atom of such a C=C double bond forms the attachment point to the remainder of the molecule.

20

Thus, even more preferably R^4 is selected from the following structures






15 where the above cyclic structures optionally carry 1, 2 or 3 substituents R^8 , where R^8 may be bound to a carbon ring atom or to a nitrogen ring atom, and where # is the attachment point to the remainder of the molecule.

Particularly preferably, R⁴ is selected from azetidin-3-yl, tetrahydrofuran-3-yl, pyrrolidin-
20 3-yl, pyrrolin-3-yl, tetrahydropyran-4-yl, tetrahydropyran-3-yl, dihydropyran-4-yl, dihy-

dropyran-3-yl, piperidin-4-yl, 1,2,5,6-tetrahydropyridin-4-yl and 1,2-dihydropyridin-4-yl, where the heterocyclic ring optionally carries 1, 2 or 3 substituents R^8 .

Preferably, these cyclic structures have the following formulae:

5



- 10 where the above cyclic structures optionally carry 1, 2 or 3 substituents R^8 , where R^8 may be bound to a carbon ring atom or to a nitrogen ring atom, and where # is the attachment point to the remainder of the molecule.

Among these, more preference is given to structures c, d, e, f, g, j, k and l.

15


- R^8 is preferably selected from C_1 - C_6 -alkyl, C_1 - C_6 -haloalkyl, C_3 - C_7 -cycloalkyl, C_3 - C_7 -halocycloalkyl, C_2 - C_4 -alkenyl, C_2 - C_4 -haloalkenyl, C_1 - C_6 -alkoxy, C_1 - C_6 -haloalkoxy, formyl, C_1 - C_6 -alkylcarbonyl, C_1 - C_6 -haloalkylcarbonyl, C_1 - C_6 -alkoxycarbonyl, C_1 - C_6 -haloalkoxycarbonyl and benzyl and is more preferably C_1 - C_6 -alkyl, C_1 - C_6 -haloalkyl or C_1 - C_6 -alkoxycarbonyl. Even more preferably, R^8 is C_1 - C_4 -alkyl, C_1 - C_4 -haloalkyl, especially fluorinated C_1 - C_4 -alkyl, or tert-butoxycarbonyl, and is in particular C_1 - C_2 -alkyl, C_1 - C_2 -haloalkyl, especially fluorinated C_1 - C_2 -alkyl, or tert-butoxycarbonyl.

20

Preferably, R^8 is N-bound; i.e. it is bound to a nitrogen ring atom.

25

Specifically, R^4 is selected from following structures:



where

R^{8a} is hydrogen or has one of the general or, in particular, one of the preferred meanings given above for R^8 ; and

is the attachment point to the remainder of the molecule.

Preferably, at most one of X^1 , X^2 , X^3 , X^4 , X^5 and X^6 is N. If one of X^1 , X^2 , X^3 , X^4 , X^5 and X^6 is N, this is preferably X^3 or X^4 and more preferably X^3 .

In one particular embodiment, none of X^1 , X^2 , X^3 , X^4 , X^5 and X^6 is N. In other words, more preferably X^1 and X^2 are CR^2 and X^3 , X^4 , X^5 and X^6 are CR^3 or CR^4 .

In another particular embodiment, X^1 and X^2 are CR^2 , X^3 is N and X^4 , X^5 and X^6 are CR^3 or CR^4 .

If one of Y^1 , Y^2 , Y^3 and Y^4 is N, this is preferably Y^2 . In other words, preferably Y^1 , Y^2 , Y^3 and Y^4 are CR^4 or CR^5 , or Y^2 is N and Y^1 , Y^3 and Y^4 are CR^4 or CR^5 , of course with the proviso that at most one of Y^1 , Y^2 , Y^3 and Y^4 is CR^4 .

A is preferably NR^B . R^B is preferably hydrogen or C_1 - C_4 -alkyl, more preferably hydrogen or methyl and specifically hydrogen. Thus, A is specifically NH.

R^1 is preferably hydrogen or C_1 - C_4 -alkyl, more preferably hydrogen or methyl and in particular hydrogen.

R^2 is preferably hydrogen.

R^3 is preferably selected from hydrogen, CN, halogen (preferably F or Cl, more preferably F), C_1 - C_6 -alkyl, C_1 - C_6 -haloalkyl, C_3 - C_7 -cycloalkyl, C_3 - C_7 -halocycloalkyl, C_1 - C_6 -alkoxy, C_1 - C_6 -haloalkoxy, formyl, C_1 - C_6 -alkylcarbonyl, C_1 - C_6 -haloalkylcarbonyl, C_1 - C_6 -alkoxycarbonyl and C_1 - C_6 -haloalkoxycarbonyl, more preferably from hydrogen, CN, F, Cl, C_1 - C_6 -alkyl, C_1 - C_6 -haloalkyl, C_1 - C_6 -alkoxy and C_1 - C_6 -haloalkoxy and is in particular selected from hydrogen, CN, F, C_1 - C_6 -alkyl, C_1 - C_6 -haloalkyl, C_1 - C_6 -alkoxy and C_1 - C_6 -

haloalkoxy, specifically hydrogen, F, C₁-C₂-alkyl, C₁-C₂-haloalkyl, C₁-C₂-alkoxy and C₁-C₂-haloalkoxy.

Preferably, 0, 1 or 2 of the radicals R³ is different from hydrogen.

5

Specifically, R³ is hydrogen if one or two of X³, X⁴, X⁵ and X⁶ are CR⁴.

In particular, at most two of the radicals R³ are different from hydrogen, if none of X³, X⁴, X⁵ and X⁶ are CR⁴, in particular if also none of Y¹, Y², Y³ and Y⁴ is CR⁴, provided, of course, that in this particular case one of Y¹, Y², Y³ and Y⁴ is C-CF₃.

10

R⁵ is preferably selected from hydrogen, CN, halogen (preferably F or Cl, more preferably F), C₁-C₆-alkyl, C₁-C₆-haloalkyl, C₃-C₇-cycloalkyl, C₃-C₇-halocycloalkyl, C₁-C₆-alkoxy, C₁-C₆-haloalkoxy, formyl, C₁-C₆-alkylcarbonyl, C₁-C₆-haloalkylcarbonyl, C₁-C₆-alkoxycarbonyl and C₁-C₆-haloalkoxycarbonyl, more preferably from hydrogen, halogen, C₁-C₄-haloalkyl, C₃-C₇-cycloalkyl and C₃-C₇-halocycloalkyl, even more preferably from hydrogen, F, Cl, C₁-C₄-haloalkyl, C₃-C₇-cycloalkyl and C₃-C₇-halocycloalkyl, in particular from hydrogen, F, C₁-C₄-haloalkyl, C₃-C₇-cycloalkyl and C₃-C₇-halocycloalkyl and specifically from hydrogen, fluorinated C₁-C₂-alkyl and C₃-C₆-cycloalkyl.

15

Preferably, at most one of the radicals R⁵ is different from hydrogen.

If none of X³, X⁴, X⁵ and X⁶ are CR⁴ and if also none of Y¹, Y², Y³ and Y⁴ is CR⁴, one of Y¹, Y², Y³ and Y⁴ is C-CF₃. In this particular embodiment, Y⁴ is preferably C-CF₃. Likewise, it is preferred that in this particular embodiment X¹ is CR², X² is CR², X³, X⁴, X⁵ and X⁶ are CR³ while Y¹, Y², Y³ and Y⁴ are independently of each other are selected from CR⁵ and N, where R², R³ and R⁵ are as defined herein, provided that at most one and in particular none of Y¹, Y², Y³ and Y⁴ is N and one of Y¹, Y², Y³ and Y⁴ is C-CF₃, preferably with Y⁴ being C-CF₃. In this particular embodiment, preferably 0, 1 or 2 radicals R³ are different from hydrogen. In this particular case, R² is preferably hydrogen.


20

25

30

In a particularly preferred embodiment of the invention the compounds of formula I are compounds of formula I-1

31



where

X^3 is N or CH;

Y^2 is N or CH;

- 5 R^{31} has one of the general or, in particular, one of the preferred meanings given above for R^3 except for hydrogen, and is preferably halogen, C_1 - C_4 -alkyl or C_1 - C_4 -alkoxy and more preferably F, C_1 - C_4 -alkyl or C_1 - C_4 -alkoxy;

R^4 has one of the general or, in particular, one of the preferred meanings given above;

- 10 R^{51} has one of the general or, in particular, one of the preferred meanings given above for R^5 except for hydrogen; and

a is 0, 1 or 2; and


b, c and d are independently of each other 0 or 1, with the proviso that one of b and c is 1.

15

Preferably, a is 0 if b is 1.

In another particularly preferred embodiment of the invention the compounds of formula I are compounds of formula I-2

20



where

Y^2 is N or in particular CH;

R^{31} if present, has one of the meanings given for R^3 except for hydrogen and has preferably one of the preferred meanings;

R^4 if present, has one of the meanings given herein;

a is 0, 1 or 2;

5 b is 0 or 1, in particular 0.

Amongst the compounds of formula I-2, a particular embodiment relates to those compounds, where Y^2 is CH.

10 Amongst the compounds of formula I-2, a particular embodiment relates to those compounds, where R^{31} , if present, is selected from the group consisting of halogen, trifluoromethyl, cyano and methoxy.


If n is 1 or 2, R^{31} is in particular located in the 6-, 7- or 8-position of formula I-2.

15


Amongst the compounds of formula I-2, a particular embodiment relates to those compounds, where b is 0. However, if b is 1, R^4 is preferably located in the 7-position as indicated.

20 Suitable compounds I are those of formulae I.a to I.l, the stereoisomers, N-oxides, prodrugs, tautomers and/or physiologically tolerated acid addition salts thereof, wherein Y^2 is N or CH, R^4 has the above-defined general or preferred meanings, R^{3a} , R^{3b} , R^{3c} and R^{3d} are hydrogen or have one of the above-defined general or preferred meanings given for R^3 and R^{5a} and R^{5b} are hydrogen or have one of the above-defined general or preferred meanings given for R^5 . Particularly preferred meanings of R^{3a} , R^{3b} , R^{3c} , R^{3d} , R^4 , R^{5a} , R^{5b} and Y^2 specifically in compounds of formulae I.a to I.l are as defined below.


25



I.a












I.b



I.c

33



5

Examples of preferred compounds which are represented by the formulae I.a to I.l are the individual compounds compiled in the tables 1 to 418 below, where the variable R^4 has the meanings given in one row of Table A. Moreover, the meanings mentioned for the individual variables in the tables are per se, independently of the combination in which they are mentioned, a particularly preferred embodiment of the substituents in question.

Table 1
Compounds of the formula I.a in which Y^2 is CH, R^{5a} and R^{5b} are H and R^4 for a compound corresponds in each case to one row of Table A.

15

Table 2

Compounds of the formula I.a in which Y^2 is CH, R^{5a} is F, R^{5b} is H and R^4 for a compound corresponds in each case to one row of Table A.

Table 3

- 5 Compounds of the formula I.a in which Y^2 is CH, R^{5a} is Cl, R^{5b} is H and R^4 for a compound corresponds in each case to one row of Table A.

Table 4

Compounds of the formula I.a in which Y^2 is CH, R^{5a} is Br, R^{5b} is H and R^4 for a compound corresponds in each case to one row of Table A.

- 10 Table 5

Compounds of the formula I.a in which Y^2 is CH, R^{5a} is CH_3 , R^{5b} is H and R^4 for a compound corresponds in each case to one row of Table A.

Table 6

- 15 Compounds of the formula I.a in which Y^2 is CH, R^{5a} is CF_3 , R^{5b} is H and R^4 for a compound corresponds in each case to one row of Table A.

Table 7

Compounds of the formula I.a in which Y^2 is CH, R^{5a} is OCH_3 , R^{5b} is H and R^4 for a compound corresponds in each case to one row of Table A.

Table 8

- 20 Compounds of the formula I.a in which Y^2 is CH, R^{5a} is OCF_3 , R^{5b} is H and R^4 for a compound corresponds in each case to one row of Table A.

Table 9

Compounds of the formula I.a in which Y^2 is CH, R^{5a} is cyclopropyl, R^{5b} is H and R^4 for a compound corresponds in each case to one row of Table A.

- 25 Table 10

Compounds of the formula I.a in which Y^2 is CH, R^{5a} is H, R^{5b} is F and R^4 for a compound corresponds in each case to one row of Table A.

Table 11

- 30 Compounds of the formula I.a in which Y^2 is CH, R^{5a} is H, R^{5b} is Cl and R^4 for a compound corresponds in each case to one row of Table A.

Table 12

Compounds of the formula I.a in which Y^2 is CH, R^{5a} is H, R^{5b} is Br and R^4 for a compound corresponds in each case to one row of Table A.

Table 13

- 35 Compounds of the formula I.a in which Y^2 is CH, R^{5a} is H, R^{5b} is CH_3 and R^4 for a compound corresponds in each case to one row of Table A.

Table 14

Compounds of the formula I.a in which Y^2 is CH, R^{5a} is H, R^{5b} is CF_3 and R^4 for a compound corresponds in each case to one row of Table A.

Table 15

Compounds of the formula I.a in which Y^2 is CH, R^{5a} is H, R^{5b} is OCH_3 and R^4 for a compound corresponds in each case to one row of Table A.

Table 16

- 5 Compounds of the formula I.a in which Y^2 is CH, R^{5a} is H, R^{5b} is OCF_3 and R^4 for a compound corresponds in each case to one row of Table A.

Table 17

Compounds of the formula I.a in which Y^2 is CH, R^{5a} is H, R^{5b} is cyclopropyl and R^4 for a compound corresponds in each case to one row of Table A.

10 Table 18

Compounds of the formula I.a in which Y^2 is N, R^{5a} and R^{5b} are H and R^4 for a compound corresponds in each case to one row of Table A.

Table 19

- 15 Compounds of the formula I.a in which Y^2 is N, R^{5a} is F, R^{5b} is H and R^4 for a compound corresponds in each case to one row of Table A.

Table 20

Compounds of the formula I.a in which Y^2 is N, R^{5a} is Cl, R^{5b} is H and R^4 for a compound corresponds in each case to one row of Table A.

Table 21

- 20 Compounds of the formula I.a in which Y^2 is N, R^{5a} is Br, R^{5b} is H and R^4 for a compound corresponds in each case to one row of Table A.

Table 22

Compounds of the formula I.a in which Y^2 is N, R^{5a} is CH_3 , R^{5b} is H and R^4 for a compound corresponds in each case to one row of Table A.

25 Table 23

Compounds of the formula I.a in which Y^2 is N, R^{5a} is CF_3 , R^{5b} is H and R^4 for a compound corresponds in each case to one row of Table A.

Table 24

- 30 Compounds of the formula I.a in which Y^2 is N, R^{5a} is OCH_3 , R^{5b} is H and R^4 for a compound corresponds in each case to one row of Table A.

Table 25

Compounds of the formula I.a in which Y^2 is N, R^{5a} is OCF_3 , R^{5b} is H and R^4 for a compound corresponds in each case to one row of Table A.

Table 26

- 35 Compounds of the formula I.a in which Y^2 is N, R^{5a} is cyclopropyl, R^{5b} is H and R^4 for a compound corresponds in each case to one row of Table A.

Table 27

Compounds of the formula I.a in which Y^2 is N, R^{5a} is H, R^{5b} is F and R^4 for a compound corresponds in each case to one row of Table A.

Table 28

Compounds of the formula I.a in which Y^2 is N, R^{5a} is H, R^{5b} is Cl and R^4 for a compound corresponds in each case to one row of Table A.

Table 29

- 5 Compounds of the formula I.a in which Y^2 is N, R^{5a} is H, R^{5b} is Br and R^4 for a compound corresponds in each case to one row of Table A.

Table 30

Compounds of the formula I.a in which Y^2 is N, R^{5a} is H, R^{5b} is CH_3 and R^4 for a compound corresponds in each case to one row of Table A.

- 10 Table 31

Compounds of the formula I.a in which Y^2 is N, R^{5a} is H, R^{5b} is CF_3 and R^4 for a compound corresponds in each case to one row of Table A.

Table 32

- 15 Compounds of the formula I.a in which Y^2 is N, R^{5a} is H, R^{5b} is OCH_3 and R^4 for a compound corresponds in each case to one row of Table A.

Table 33

Compounds of the formula I.a in which Y^2 is N, R^{5a} is H, R^{5b} is OCF_3 and R^4 for a compound corresponds in each case to one row of Table A.

Table 34

- 20 Compounds of the formula I.a in which Y^2 is N, R^{5a} is H, R^{5b} is cyclopropyl and R^4 for a compound corresponds in each case to one row of Table A.

Tables 35 to 68

- Compounds of the formula I.b in which the combination of Y^2 , R^{5a} and R^{5b} is as defined in tables 1 to 34 and R^4 for a compound corresponds in each case to one row of Table A.
- 25 A.

Tables 69 to 102

- Compounds of the formula I.c in which the combination of Y^2 , R^{5a} and R^{5b} is as defined in tables 1 to 34 and R^4 for a compound corresponds in each case to one row of Table A.

- 30 Tables 103 to 136

Compounds of the formula I.d in which the combination of Y^2 , R^{5a} and R^{5b} is as defined in tables 1 to 34 and R^4 for a compound corresponds in each case to one row of Table A.

Tables 137 to 170

- 35 Compounds of the formula I.e in which the combination of Y^2 , R^{5a} and R^{5b} is as defined in tables 1 to 34 and R^4 for a compound corresponds in each case to one row of Table A.

Tables 171 to 204

Compounds of the formula I.f in which the combination of Y^2 , R^{5a} and R^{5b} is as defined in tables 1 to 34 and R^4 for a compound corresponds in each case to one row of Table A.

Tables 205 to 238

- 5 Compounds of the formula I.g in which the combination of Y^2 , R^{5a} and R^{5b} is as defined in tables 1 to 34 and R^4 for a compound corresponds in each case to one row of Table A.

Tables 239 to 272

- 10 Compounds of the formula I.h in which the combination of Y^2 , R^{5a} and R^{5b} is as defined in tables 1 to 34 and R^4 for a compound corresponds in each case to one row of Table A.

Tables 273 to 306

- 15 Compounds of the formula I.i in which the combination of Y^2 , R^{5a} and R^{5b} is as defined in tables 1 to 34 and R^4 for a compound corresponds in each case to one row of Table A.

Tables 307 to 340

Compounds of the formula I.j in which the combination of Y^2 , R^{5a} and R^{5b} is as defined in tables 1 to 34 and R^4 for a compound corresponds in each case to one row of Table A.

- 20 Table 341

Compounds of the formula I.k in which Y^2 is CH, R^{3a} , R^{3b} , R^{3c} and R^{3d} are H and R^4 for a compound corresponds in each case to one row of Table A.

Table 342

- 25 Compounds of the formula I.k in which Y^2 is CH, R^{3a} is F, R^{3b} , R^{3c} and R^{3d} are H and R^4 for a compound corresponds in each case to one row of Table A.

Table 343

Compounds of the formula I.k in which Y^2 is CH, R^{3a} is Cl, R^{3b} , R^{3c} and R^{3d} are H and R^4 for a compound corresponds in each case to one row of Table A.

Table 344

- 30 Compounds of the formula I.k in which Y^2 is CH, R^{3a} is Br, R^{3b} , R^{3c} and R^{3d} are H and R^4 for a compound corresponds in each case to one row of Table A.

Table 345

Compounds of the formula I.k in which Y^2 is CH, R^{3a} is CH_3 , R^{3b} , R^{3c} and R^{3d} are H and R^4 for a compound corresponds in each case to one row of Table A.

- 35 Table 346

Compounds of the formula I.k in which Y^2 is CH, R^{3a} is CF_3 , R^{3b} , R^{3c} and R^{3d} are H and R^4 for a compound corresponds in each case to one row of Table A.

Table 347

Compounds of the formula I.k in which Y² is CH, R^{3a} is OCH₃, R^{3b}, R^{3c} and R^{3d} are H and R⁴ for a compound corresponds in each case to one row of Table A.

Table 348

Compounds of the formula I.k in which Y² is CH, R^{3a} is OCF₃, R^{3b}, R^{3c} and R^{3d} are H and R⁴ for a compound corresponds in each case to one row of Table A.

5

Table 349

Compounds of the formula I.k in which Y² is CH, R^{3a} is cyclopropyl, R^{3b}, R^{3c} and R^{3d} are H and R⁴ for a compound corresponds in each case to one row of Table A.

Table 350

Compounds of the formula I.k in which Y² is CH, R^{3b} is F, R^{3a}, R^{3c} and R^{3d} are H and R⁴ for a compound corresponds in each case to one row of Table A.

10

Table 351

Compounds of the formula I.k in which Y² is CH, R^{3b} is Cl, R^{3a}, R^{3c} and R^{3d} are H and R⁴ for a compound corresponds in each case to one row of Table A.

15

Table 352

Compounds of the formula I.k in which Y² is CH, R^{3b} is Br, R^{3a}, R^{3c} and R^{3d} are H and R⁴ for a compound corresponds in each case to one row of Table A.

Table 353

Compounds of the formula I.k in which Y² is CH, R^{3b} is CH₃, R^{3a}, R^{3c} and R^{3d} are H and R⁴ for a compound corresponds in each case to one row of Table A.

20

Table 354

Compounds of the formula I.k in which Y² is CH, R^{3b} is CF₃, R^{3a}, R^{3c} and R^{3d} are H and R⁴ for a compound corresponds in each case to one row of Table A.

Table 355

Compounds of the formula I.k in which Y² is CH, R^{3b} is OCH₃, R^{3a}, R^{3c} and R^{3d} are H and R⁴ for a compound corresponds in each case to one row of Table A.

25

Table 356

Compounds of the formula I.k in which Y² is CH, R^{3b} is OCF₃, R^{3a}, R^{3c} and R^{3d} are H and R⁴ for a compound corresponds in each case to one row of Table A.

30

Table 357

Compounds of the formula I.k in which Y² is CH, R^{3b} is cyclopropyl, R^{3a}, R^{3c} and R^{3d} are H and R⁴ for a compound corresponds in each case to one row of Table A.

Table 358

Compounds of the formula I.k in which Y² is CH, R^{3c} is F, R^{3a}, R^{3b} and R^{3d} are H and R⁴ for a compound corresponds in each case to one row of Table A.

35

Table 359

Compounds of the formula I.k in which Y² is CH, R^{3c} is Cl, R^{3a}, R^{3b} and R^{3d} are H and R⁴ for a compound corresponds in each case to one row of Table A.

Table 360

Compounds of the formula I.k in which Y^2 is CH, R^{3c} is Br, R^{3a} , R^{3b} and R^{3d} are H and R^4 for a compound corresponds in each case to one row of Table A.

Table 361

- 5 Compounds of the formula I.k in which Y^2 is CH, R^{3c} is CH_3 , R^{3a} , R^{3b} and R^{3d} are H and R^4 for a compound corresponds in each case to one row of Table A.

Table 362

Compounds of the formula I.k in which Y^2 is CH, R^{3c} is CF_3 , R^{3a} , R^{3b} and R^{3d} are H and R^4 for a compound corresponds in each case to one row of Table A.

Table 363

- 10 Compounds of the formula I.k in which Y^2 is CH, R^{3c} is OCH_3 , R^{3a} , R^{3b} and R^{3d} are H and R^4 for a compound corresponds in each case to one row of Table A.

Table 364

Compounds of the formula I.k in which Y^2 is CH, R^{3c} is OCF_3 , R^{3a} , R^{3b} and R^{3d} are H and R^4 for a compound corresponds in each case to one row of Table A.

- 15 Table 365

Compounds of the formula I.k in which Y^2 is CH, R^{3c} is cyclopropyl, R^{3a} , R^{3b} and R^{3d} are H and R^4 for a compound corresponds in each case to one row of Table A.

Table 366

- 20 Compounds of the formula I.k in which Y^2 is CH, R^{3d} is F, R^{3a} , R^{3b} and R^{3c} are H and R^4 for a compound corresponds in each case to one row of Table A.

Table 367

Compounds of the formula I.k in which Y^2 is CH, R^{3d} is Cl, R^{3a} , R^{3b} and R^{3c} are H and R^4 for a compound corresponds in each case to one row of Table A.

Table 368

- 25 Compounds of the formula I.k in which Y^2 is CH, R^{3d} is Br, R^{3a} , R^{3b} and R^{3c} are H and R^4 for a compound corresponds in each case to one row of Table A.

Table 369

Compounds of the formula I.k in which Y^2 is CH, R^{3d} is CH_3 , R^{3a} , R^{3b} and R^{3c} are H and R^4 for a compound corresponds in each case to one row of Table A.

- 30 Table 370

Compounds of the formula I.k in which Y^2 is CH, R^{3d} is CF_3 , R^{3a} , R^{3b} and R^{3c} are H and R^4 for a compound corresponds in each case to one row of Table A.

Table 371

- 35 Compounds of the formula I.k in which Y^2 is CH, R^{3d} is OCH_3 , R^{3a} , R^{3b} and R^{3c} are H and R^4 for a compound corresponds in each case to one row of Table A.

Table 372

Compounds of the formula I.k in which Y^2 is CH, R^{3d} is OCF_3 , R^{3a} , R^{3b} and R^{3c} are H and R^4 for a compound corresponds in each case to one row of Table A.

Table 373

Compounds of the formula I.k in which Y^2 is CH, R^{3d} is cyclopropyl, R^{3a} , R^{3b} and R^{3c} are H and R^4 for a compound corresponds in each case to one row of Table A.

Table 374

Compounds of the formula I.k in which Y^2 is CH, R^{3b} is CH_3 , R^{3d} is Cl, R^{3a} and R^{3c} are H and R^4 for a compound corresponds in each case to one row of Table A.

Table 375

Compounds of the formula I.k in which Y^2 is CH, R^{3b} is Cl, R^{3d} is Cl, R^{3a} and R^{3c} are H and R^4 for a compound corresponds in each case to one row of Table A.

Table 376

Compounds of the formula I.k in which Y^2 is CH, R^{3b} is F, R^{3d} is F, R^{3a} and R^{3c} are H and R^4 for a compound corresponds in each case to one row of Table A.

Table 377

Compounds of the formula I.k in which Y^2 is CH, R^{3b} is OCH_3 , R^{3d} is F, R^{3a} and R^{3c} are H and R^4 for a compound corresponds in each case to one row of Table A.

Table 378

Compounds of the formula I.k in which Y^2 is CH, R^{3a} is F, R^{3d} is F, R^{3b} and R^{3c} are H and R^4 for a compound corresponds in each case to one row of Table A.

Tables 379 to 416

Compounds of the formula I.k in which Y^2 is N, the combination of R^{3a} , R^{3b} , R^{3c} and R^{3d} is as defined in tables 341 to 378 and R^4 for a compound corresponds in each case to one row of Table A.

Table 417



Compounds of the formula I.l in which Y^2 is CH and R^4 for a compound corresponds in each case to one row of Table A.





Table 418



















Compounds of the formula I.l in which Y^2 is N and R^4 for a compound corresponds in each case to one row of Table A.

















Table A

30

No.	R^4
A-1	
A-2	
A-3	
A-4	

No.	R^4
A-5	
A-6	
A-7	
A-8	

No.	R ⁴
A-9	
A-10	
A-11	
A-12	
A-13	
A-14	
A-15	
A-16	
A-17	
A-18	
A-19	
A-20	
A-21	
A-22	
A-23	
A-24	
A-25	
A-26	

No.	R ⁴
A-27	
A-28	
A-29	
A-30	
A-31	
A-32	
A-33	
A-34	
A-35	
A-36	
A-37	
A-38	
A-39	
A-40	
A-41	
A-42	
A-43	

No.	R ⁴
A-44	
A-45	
A-46	
A-47	
A-48	
A-49	
A-50	
A-51	

No.	R ⁴
A-52	
A-53	
A-54	
A-55	
A-56	
A-57	

Among the above compounds, preference is given to compounds of formulae I.b, I.c, I.k and I.l.

- 5 Suitable compounds I are also those of formula I.m, the stereoisomers, N-oxides, prodrugs, tautomers and/or physiologically tolerated acid addition salts thereof, wherein Y² is N or CH, R^{3a}, R^{3b} and R^{3c} have one of the above-defined general or preferred meanings given for R³ and are in particular selected from the group consisting of hydrogen, halogen, trifluoromethyl, cyano and methoxy. Particularly preferred meanings of R^{3a},
 10 R^{3b} and R^{3c} and Y² specifically in compounds of formula I.m are as defined in the table below.




Table 417

Compounds of the formula I.m, in which Y² is CH and R^{3a}, R^{3b} and R^{3c} correspond in each case to one row of Table B.

Table 418

Compounds of the formula I.m, in which Y² is CH and R^{3a}, R^{3b} and R^{3c} correspond in each case to one row of Table B.

5

Table B


No.	R ^{3a}	R ^{3b}	R ^{3c}
B-1	H	H	H
B-2	F	H	H
B-3	H	F	H
B-4	H	H	F
B-5	F	F	H
B-6	F	H	F
B-7	H	F	F
B-8	Cl	H	H
B-9	H	Cl	H
B-10	H	H	Cl
B-11	Cl	Cl	H
B-12	Cl	H	Cl
B-13	H	Cl	Cl
B-14	Br	H	H
B-15	H	Br	H
B-16	H	H	Br
B-17	I	H	H
B-18	H	I	H
B-19	H	H	I
B-20	CF ₃	H	H
B-21	H	CF ₃	H
B-22	H	H	CF ₃
B-23	CF ₃	CF ₃	H
B-24	CF ₃	H	CF ₃
B-25	H	CF ₃	CF ₃
B-26	OCH ₃	H	H
B-27	H	OCH ₃	H
B-28	H	H	OCH ₃
B-29	OCH ₃	OCH ₃	H
B-30	OCH ₃	H	OCH ₃

No.	R ^{3a}	R ^{3b}	R ^{3c}
B-31	H	OCH ₃	OCH ₃
B-32	CN	H	H
B-33	H	CN	H
B-34	H	H	CN
B-35	H	H	I

The compounds of the present invention can be prepared by analogy to routine techniques a skilled person is familiar with. In particular, the compounds of the formula I can be prepared according to the following schemes, wherein the variables, if not
5 stated otherwise, are as defined above and Z indicates a halogen atom, especially Br or I.

Compounds I wherein A is NR^B can be prepared as described in schemes 1 to 4 and 6
10 to 8.


Scheme 1:



The bicyclic amine 1 can be reacted with amine 2 in the presence of triphosgene (or phosgene or other equivalent reagent), together with a non-alkylating base such as
15 triethylamine. The reaction is carried out in the presence of a suitable solvent such as toluene or N,N-dimethylformamide. The reaction is usually carried out at temperatures of from -30 to 50°C to give substituted ureas of general formula 3.

Disubstituted urea compounds of the general formula 3 can also be prepared according
20 to a route depicted in scheme 2.


Scheme 2:



The amine **1** can be acylated by reaction with an acyl azide **4** (prepared by reaction of the corresponding acyl halide with a metal azide salt according to standard methods of organic chemistry) to give disubstituted ureas of general formula **3**. The reaction is carried out in the presence of a suitable solvent such as toluene or N,N-dimethylformamide. The reaction is usually carried out at temperatures of from 20 - 120°C. Other conditions for this transformation (known as the Curtius rearrangement) are described in the following articles: Journal of Organic Chemistry, 1986, 51, 3007 & 5123; Journal of Organic Chemistry, 1987, 52, 4875; Tetrahedron Letters, 1984, 25, 3515; and Organic Reactions, 1947, 3, 337.


Disubstituted urea compounds of the general formula **3** can also be prepared according to a route depicted in scheme 3.

Scheme 3:



The amine **1** can be acylated by reaction with an isocyanate **5** to give disubstituted ureas of general formula **3**. The reaction is carried out in the presence of a suitable solvent such as toluene or N,N-dimethylformamide. The reaction is usually carried out at temperatures of from 20 - 120°C.

Scheme 4:




The bicyclic amine **1** can be converted to trichloroacetamide **6** by reaction with tri-chloroacetyl chloride. The reaction is carried out in the presence of a suitable solvent such as toluene or N,N-dimethylformamide. The reaction is usually carried out at temperatures of from 20 - 120°C. The trichloroacetamide **6** can be reacted with an amine **2** to give substituted ureas of general formula **3**.

Compounds I wherein R^B and/or R^1 are different from H can be prepared by N-alkylating compound **3**.

10

Compounds I wherein A is CH_2 can be prepared as described in schemes 5 to 8.

Scheme 5:



15

The carboxylic acid **7** can be converted into the amide **10** by reaction with an amine **8** using standard amide formation conditions familiar to those skilled in the art. The reaction is carried out in the presence of a suitable solvent such dimethylacetamide, N,N-

dimethylformamide or THF. The reaction is usually carried out at temperatures of from 20 to 120°C. Coupling reagents such as HOBT or carbonyl diimidazole are employed.


Compounds I, wherein A is CR^{A1}R^{A2} can be prepared analogously.

5

The introduction of the group R⁴ can be accomplished as shown in the following schemes.

Scheme 6

10




Substitution of an appropriate placed halogen Z can be accomplished by Suzuki coupling of 11 with an appropriately substituted boronic acid (R = H) or ester (R ≠ H) 12, wherein R⁴ is a C-bound partially (C=C) unsaturated monocyclic 3-, 4-, 5-, 6- or 7-membered heterocyclic ring containing 1, 2 or 3 heteroatoms or heteroatom-containing groups selected from O, N, S, NO, SO and SO₂ as ring members, where the heterocyclic ring optionally carries 1, 2 or 3 C- or N-bound substituents R⁸, where the ring has the C=C bond in α-position to the attachment point to the boron atom (so that the C=C double bond is a vinylic bond to B) to give substituted products of the general structure I'. The substitution may be conducted via a palladium-mediated coupling using a catalyst such as tetrakis(triphenylphosphine)palladium(0) in the presence of a base (e.g. Na₂CO₃) in a solvent such as DMF. The unsaturated compound I' may be reduced to the saturated product I'', wherein R⁴ is a C-bound saturated monocyclic 3-, 4-, 5-, 6- or 7-membered heterocyclic ring containing 1, 2 or 3 heteroatoms or heteroatom-

15

20

containing groups selected from O, N, S, NO, SO and SO₂ as ring members, where the heterocyclic ring optionally carries 1, 2 or 3 C- or N-bound substituents R⁸, by reduction of I' in the presence of a catalyst (e.g. Pd on carbon) in the presence of hydrogen.

5 Scheme 7:




A similar sequence may be accomplished by reaction of **11** in a Heck coupling with a partially unsaturated ring H-R^{4'} to give substituted products of general structure I' which can in turn be reduced to the saturated product I'' as already described.

Starting compound **11** in schemes 6 and 7 can be obtained by using in the reactions of schemes 1 to 5 compounds **1** or **7** yet carrying a halogen atom Z in the desired position.


15

Scheme 8



20 Groups R⁴ can moreover be introduced in an alkylation reaction, e.g. by reacting the compound **13** under the reaction conditions of a Friedel-Crafts alkylation with a compound Z-R⁴, where Z is a halogen atom, especially Cl or Br. Suitable Lewis acids are for example AlCl₃, FeCl₃, SbCl₅, SnCl₄, BF₃, TiCl₄ and ZnCl₂.

25 Compounds I where R⁴ is bound to the ring containing Y¹ to Y⁴ as ring members can be prepared in analogy to the reactions shown in schemes 6 to 8 by starting from a compound **14**, wherein Z is a halogen atom, especially Br or I.

**14**

Compound 14 can be prepared by using in the reactions of schemes 1 to 5 starting compounds 2, 4, 5 or 8 yet carrying a halogen atom Z in the desired position.

If not indicated otherwise, the above-described reactions are generally carried out in a solvent at temperatures between room temperature and the boiling temperature of the solvent employed. Alternatively, the activation energy which is required for the reaction can be introduced into the reaction mixture using microwaves, something which has proved to be of value, in particular, in the case of the reactions catalyzed by transition metals (with regard to reactions using microwaves, see Tetrahedron 2001, 57, p. 9199 ff. p. 9225 ff. and also, in a general manner, "Microwaves in Organic Synthesis", André Loupy (Ed.), Wiley-VCH 2002.

The acid addition salts of compounds I are prepared in a customary manner by mixing the free base with a corresponding acid, where appropriate in solution in an organic solvent, for example a lower alcohol, such as methanol, ethanol or propanol, an ether, such as methyl tert-butyl ether or diisopropyl ether, a ketone, such as acetone or methyl ethyl ketone, or an ester, such as ethyl acetate.

The present invention moreover relates to compounds of formula I as defined above, wherein at least one of the atoms has been replaced by its stable, non-radioactive isotope (e.g., hydrogen by deuterium, ^{12}C by ^{13}C , ^{14}N by ^{15}N , ^{16}O by ^{18}O) and preferably wherein at least one hydrogen atom has been replaced by a deuterium atom.

Of course, the compounds according to the invention contain more of the respective isotope than this naturally occurs and thus is anyway present in the compounds I.

Stable isotopes (e.g., deuterium, ^{13}C , ^{15}N , ^{18}O) are nonradioactive isotopes which contain one additional neutron than the normally abundant isotope of the respective atom. Deuterated compounds have been used in pharmaceutical research to investigate the

- in vivo metabolic fate of the compounds by evaluation of the mechanism of action and metabolic pathway of the non deuterated parent compound (Blake et al. *J. Pharm. Sci.* 64, 3, 367-391 (1975)). Such metabolic studies are important in the design of safe, effective therapeutic drugs, either because the in vivo active compound administered to the patient or because the metabolites produced from the parent compound prove to be toxic or carcinogenic (Foster et al., *Advances in Drug Research* Vol. 14, pp. 2-36, Academic press, London, 1985; Kato et al., *J. Labelled Comp. Radiopharmaceut.*, 36(10):927-932 (1995); Kushner et al., *Can. J. Physiol. Pharmacol.*, 77, 79-88 (1999).
- 5 incorporation of a heavy atom, particularly substitution of deuterium for hydrogen, can give rise to an isotope effect that could alter the pharmacokinetics of the drug. This effect is usually insignificant if the label is placed at a metabolically inert position of the molecule.
- 10 Stable isotope labeling of a drug can alter its physico-chemical properties such as pK_a and lipid solubility. These changes may influence the fate of the drug at different steps along its passage through the body. Absorption, distribution, metabolism or excretion can be changed. Absorption and distribution are processes that depend primarily on the molecular size and the lipophilicity of the substance. These effects and alterations
- 15 can affect the pharmacodynamic response of the drug molecule if the isotopic substitution affects a region involved in a ligand-receptor interaction.
- 20

Drug metabolism can give rise to large isotopic effect if the breaking of a chemical bond to a deuterium atom is the rate limiting step in the process. While some of the physical properties of a stable isotope-labeled molecule are different from those of the unlabeled one, the chemical and biological properties are the same, with one important exception: because of the increased mass of the heavy isotope, any bond involving the heavy isotope and another atom will be stronger than the same bond between the light isotope and that atom. In any reaction in which the breaking of this bond is the rate limiting step, the reaction will proceed slower for the molecule with the heavy isotope due to "kinetic isotope effect". A reaction involving breaking a C--D bond can be up to 700 percent slower than a similar reaction involving breaking a C--H bond. If the C--D bond is not involved in any of the steps leading to the metabolite, there may not be any effect to alter the behavior of the drug. If a deuterium is placed at a site involved in the metabolism of a drug, an isotope effect will be observed only if breaking of the C--D bond is the rate limiting step. There is evidence to suggest that whenever cleavage of an aliphatic C--H bond occurs, usually by oxidation catalyzed by a mixed-function oxidase, replacement of the hydrogen by deuterium will lead to observable isotope effect. It is also important to understand that the incorporation of deuterium at the site of me-

25

30

35

tabolism slows its rate to the point where another metabolite produced by attack at a carbon atom not substituted by deuterium becomes the major pathway a process called "metabolic switching".

- 5 Deuterium tracers, such as deuterium-labeled drugs and doses, in some cases repeatedly, of thousands of milligrams of deuterated water, are also used in healthy humans of all ages, including neonates and pregnant women, without reported incident (e.g. Pons G and Rey E, *Pediatrics* 1999 104: 633; Coward W A et al., *Lancet* 1979 7: 13; Schwarcz H P, *Control. Clin. Trials* 1984 5(4 Suppl): 573; Rodewald L E et al., *J. Pediatr.* 1989 114: 885; Butte N F et al. *Br. J. Nutr.* 1991 65: 3; MacLennan A H et al. *Am. J. Obstet Gynecol.* 1981 139: 948). Thus, it is clear that any deuterium released, for instance, during the metabolism of compounds of this invention poses no health risk.

- 15 The weight percentage of hydrogen in a mammal (approximately 9%) and natural abundance of deuterium (approximately 0.015%) indicates that a 70 kg human normally contains nearly a gram of deuterium. Furthermore, replacement of up to about 15% of normal hydrogen with deuterium has been effected and maintained for a period of days to weeks in mammals, including rodents and dogs, with minimal observed adverse effects (Czajka D M and Finkel A J, *Ann. N.Y. Acad. Sci.* 1960 84: 770; Thomson J F, *Ann. New York Acad. Sci* 1960 84: 736; Czajka D M et al., *Am. J. Physiol.* 1961 201: 357). Higher deuterium concentrations, usually in excess of 20%, can be toxic in animals. However, acute replacement of as high as 15%-23% of the hydrogen in humans' fluids with deuterium was found not to cause toxicity (Blagojevic N et al. in "Dosimetry & Treatment Planning for Neutron Capture Therapy", Zamenhof R, Solares G and Harling O Eds. 1994. Advanced Medical Publishing, Madison Wis. pp.125-134; *Diabetes Metab.* 23: 251 (1997)).

- 30 Increasing the amount of deuterium present in a compound above its natural abundance is called enrichment or deuterium-enrichment. Examples of the amount of enrichment include from about 0.5, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 12, 16, 21, 25, 29, 33, 37, 42, 46, 50, 54, 58, 63, 67, 71, 75, 79, 84, 88, 92, 96, to about 100 mol %.

- 35 The hydrogens present on a particular organic compound have different capacities for exchange with deuterium. Certain hydrogen atoms are easily exchangeable under physiological conditions and, if replaced by deuterium atoms, it is expected that they will readily exchange for protons after administration to a patient. Certain hydrogen atoms may be exchanged for deuterium atoms by the action of a deuteriic acid such as D_2SO_4/D_2O . Alternatively, deuterium atoms may be incorporated in various combinations during the synthesis of compounds of the invention. Certain hydrogen atoms are

not easily exchangeable for deuterium atoms. However, deuterium atoms at the remaining positions may be incorporated by the use of deuterated starting materials or intermediates during the construction of compounds of the invention.

- 5 Deuterated and deuterium-enriched compounds of the invention can be prepared by using known methods described in the literature. Such methods can be carried out utilizing corresponding deuterated and optionally, other isotope-containing reagents and/or intermediates to synthesize the compounds delineated herein, or invoking standard synthetic protocols known in the art for introducing isotopic atoms to a chemical
- 10 structure. Relevant procedures and intermediates are disclosed, for instance in Lizondo, J et al., *Drugs Fut*, 21(11), 1116 (1996); Brickner, S J et al., *J Med Chem*, 39(3), 673 (1996); Mallesham, B et al., *Org Lett*, 5(7), 963 (2003); PCT publications WO1997010223, WO2005099353, WO1995007271, WO2006008754; US Patent Nos. 7538189; 7534814; 7531685; 7528131; 7521421; 7514068; 7511013; and US Patent
- 15 Application Publication Nos. 20090137457; 20090131485; 20090131363; 20090118238; 20090111840; 20090105338; 20090105307; 20090105147; 20090093422; 20090088416; 20090082471, the methods are hereby incorporated by reference.
- 20 The present invention further relates to a pharmaceutical composition comprising at least one compound of formula I, a stereoisomer, prodrug, tautomer and/or physiologically tolerated acid addition salt thereof and optionally at least one physiologically acceptable carrier and/or auxiliary substance.
- 25 The invention also relates to compounds of formula I or the stereoisomers, N-oxides, prodrugs, tautomers or physiologically tolerated acid addition salts thereof for use as a medicament, and to compounds of formula I or the stereoisomers, N-oxides, prodrugs, tautomers or physiologically tolerated acid addition salts thereof for use in the treatment of a medical disorder susceptible to the treatment with a compound that modu-
- 30 lates, preferably inhibits, the activity of glycogen synthase kinase 3 β .

- The invention also relates to the use of the compounds of formula I or of a stereoisomer, prodrug, tautomer or physiologically tolerated acid addition salt thereof for the preparation of a medicament for the treatment of a disorder susceptible to the treatment with a compound that modulates, preferably inhibits, the activity of glycogen synthase kinase 3 β .
- 35

Furthermore, the invention relates to a method for treating a medical disorder susceptible to treatment with a compound that modulates glycogen synthase kinase 3 β activity,

said method comprising administering an effective amount of at least one compound of formula I or of a stereoisomer, prodrug, tautomer or physiologically tolerated acid addition salt thereof or of a pharmaceutical composition as defined above to a subject in need thereof.

5

The compounds of the of formula I according to the present invention, as well as the stereoisomers, the tautomers, the prodrugs and physiologically tolerated acid addition salts thereof, are capable of modulating the activity on glycogen synthase kinase 3 β . In particular, the compounds of the of formula I, as well as the stereoisomers, the tautomers, the prodrugs and physiologically tolerated acid addition salts thereof, have an inhibitory activity on glycogen synthase kinase 3 β . Amongst the compounds of formula I those are preferred which achieve effective inhibition at low concentrations. In particular, compounds of the formula I are preferred which inhibit glycogen synthase kinase 3 β at a level of $IC_{50} < 1 \mu\text{Mol}$, more preferably at a level of $IC_{50} < 0.5 \mu\text{Mol}$, particularly preferably at a level of $IC_{50} < 0.2 \mu\text{Mol}$ and most preferably at a level of $IC_{50} < 0.1 \mu\text{Mol}$.

Therefore the compounds of the of formula I according to the present invention, their stereoisomers, tautomers, their prodrugs and their physiologically tolerated acid addition salts are useful for the treatment of a medical disorder susceptible to treatment with a compound that modulates glycogen synthase kinase 3 β activity. As mentioned above, diseases caused by abnormal GSK-3 β activity and which thus can be treated by supplying the compound of the formula I, a stereoisomer, tautomer, prodrug and/or a physiologically tolerated acid addition salt thereof, include in particular neurodegenerative diseases such as Alzheimer's disease. In addition, the compounds of the present invention are also useful for treatment of other neurodegenerative diseases such as behavioural psychiatric symptoms of dementia, Parkinson's disease, tauopathies (e.g. frontotemporoparietal dementia, corticobasal degeneration, Pick's disease, progressive supranuclear palsy, argyophilic brain disease) and other dementia including vascular dementia; acute stroke and others traumatic injuries; cerebrovascular accidents (e.g. age related macular degeneration); brain and spinal cord trauma; peripheral neuropathies; bipolar disorders, retinopathies and glaucoma. In addition, the compounds of the present invention are also useful for treatment of schizophrenia. The compounds of the present invention are also useful for treatment of pain.

Diseases which can be treated by supplying the compound of the of formula I, a stereoisomer, tautomer, prodrug and/or a physiologically tolerated acid addition salt thereof, include furthermore inflammatory diseases, such as rheumatoid arthritis and osteoarthritis.

Within the meaning of the invention, a treatment also includes a preventive treatment (prophylaxis), in particular as relapse prophylaxis or phase prophylaxis, as well as the treatment of acute or chronic signs, symptoms and/or malfunctions. The treatment can be orientated symptomatically, for example as the suppression of symptoms. It can be effected over a short period, be orientated over the medium term or can be a long-term treatment, for example within the context of a maintenance therapy.

Within the context of the treatment, the use according to the invention of the compounds of the formula I involves a method. In this method, an effective quantity of one or more compounds I, a stereoisomer, tautomer, prodrug or physiologically tolerable acid addition salt thereof, as a rule formulated in accordance with pharmaceutical and veterinary practice, is administered to the individual to be treated, preferably a mammal, in particular a human being, productive animal or domestic animal. Whether such a treatment is indicated, and in which form it is to take place, depends on the individual case and is subject to medical assessment (diagnosis) which takes into consideration signs, symptoms and/or malfunctions which are present, the risks of developing particular signs, symptoms and/or malfunctions, and other factors.

As a rule, the treatment is effected by means of single or repeated daily administration, where appropriate together, or alternating, with other active compounds or active compound-containing preparations such that a daily dose of preferably from about 0.1 to 1000 mg/kg of bodyweight, in the case of oral administration, or of from about 0.1 to 100 mg/kg of bodyweight, in the case of parenteral administration, is supplied to an individual to be treated.

The invention also relates to pharmaceutical compositions for treating an individual, preferably a mammal, in particular a human being, productive animal or domestic animal. Thus, the compounds according to the invention are customarily administered in the form of pharmaceutical compositions which comprise a pharmaceutically acceptable excipient together with at least one compound according to the invention and, where appropriate, other active compounds. These compositions can, for example, be administered orally, rectally, transdermally, subcutaneously, intravenously, intramuscularly or intranasally.

Examples of suitable pharmaceutical formulations are solid medicinal forms, such as powders, granules, tablets, in particular film tablets, lozenges, sachets, cachets, sugar-coated tablets, capsules, such as hard gelatin capsules and soft gelatin capsules, suppositories or vaginal medicinal forms, semisolid medicinal forms, such as ointments, creams, hydrogels, pastes or plasters, and also liquid medicinal forms, such as solu-

tions, emulsions, in particular oil-in-water emulsions, suspensions, for example lotions, injection preparations and infusion preparations, and eyedrops and eardrops. Implanted release devices can also be used for administering inhibitors according to the invention. In addition, it is also possible to use liposomes or microspheres.

5

When producing the pharmaceutical compositions, the compounds according to the invention are optionally mixed or diluted with one or more excipients. Excipients can be solid, semisolid or liquid materials which serve as vehicles, carriers or medium for the active compound.

10

Suitable excipients are listed in the specialist medicinal monographs. In addition, the formulations can comprise pharmaceutically acceptable carriers or customary auxiliary substances, such as glidants; wetting agents; emulsifying and suspending agents; preservatives; antioxidants; antiirritants; chelating agents; coating auxiliaries; emulsion stabilizers; film formers; gel formers; odor masking agents; taste corrigents; resin; hydrocolloids; solvents; solubilizers; neutralizing agents; diffusion accelerators; pigments; quaternary ammonium compounds; refatting and overfatting agents; raw materials for ointments, creams or oils; silicone derivatives; spreading auxiliaries; stabilizers; sterilants; suppository bases; tablet auxiliaries, such as binders, fillers, glidants, disintegrants or coatings; propellants; drying agents; opacifiers; thickeners; waxes; plasticizers and white mineral oils. A formulation in this regard is based on specialist knowledge as described, for example, in Fiedler, H.P., *Lexikon der Hilfsstoffe für Pharmazie, Kosmetik und angrenzende Gebiete* [Encyclopedia of auxiliary substances for pharmacy, cosmetics and related fields], 4th edition, Aulendorf: ECV-Editio-Kantor-Verlag, 1996.

25

The following examples serve to explain the invention without limiting it.

Examples

30

The compounds were either characterized via proton-NMR in d₆-dimethylsulfoxide or d-chloroform on a 400 MHz or 500 MHz NMR instrument (Bruker AVANCE), or by mass spectrometry, generally recorded via HPLC-MS in a fast gradient on C18-material (electrospray-ionisation (ESI) mode), or melting point.

35

The magnetic nuclear resonance spectral properties (NMR) refer to the chemical shifts (δ) expressed in parts per million (ppm). The relative area of the shifts in the ¹H-NMR spectrum corresponds to the number of hydrogen atoms for a particular functional type in the molecule. The nature of the shift, as regards multiplicity, is indicated as

singlet (s), broad singlet (s. br.), doublet (d), broad doublet (d br.), triplet (t), broad triplet (t br.), quartet (q), quintet (quint.) and multiplet (m).

Abbreviations:


5	DCM	dichloromethane
	DMSO	dimethylsulfoxide
	DMF	dimethylformamide
	MeOH	methanol
	AcOH	acetic acid
10	TFA	trifluoroacetic acid
	RT	room temperature
	quant.	quantitative

I. Preparation Examples

15

Example 1

1-(6-(3,6-Dihydro-2H-pyran-4-yl)pyridin-2-yl)-3-(8-fluoroquinolin-4-yl)urea




1-(8-Fluoroquinolin-4-yl)-3-(6-iodopyridin-2-yl)urea (250 mg, 0.582 mmol), 2-(3,6-dihydro-2H-pyran-4-yl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (187 mg, 0.873 mmol), sodium carbonate solution (2M, 1.31 mL, 2.62 mmol) and tetrakis(triphenylphosphine)-palladium(0) (68 mg, 0.058 mmol) were dissolved in 5 ml of DMF under an argon atmosphere and the mixture was stirred at 80°C, in a microwave oven, for one hour. After the solvent had been evaporated down to dryness, the resulting residue was treated with water and the mixture was then extracted with ethyl acetate. After the combined organic phases had been dried over sodium sulfate and the solvent had been filtered and evaporated down to dryness, the resulting residue was purified by column chromatography to give the title product (50 mg, 19%).

30 ¹H-NMR (d₆-DMSO, 400 MHz) δ 3.32 (m, 5H), 3.95 (s, 2H), 4.21 (s, 2H), 6.65 (s, 1H), 7.19 (m, 1H), 7.60 (m, 4H), 7.78 (m, 1H), 8.01 (m, 1H), 8.32 (s, 1H), 8.80 (m, 1H), 9.98 (s, 1H), 10.70 (s, 1H); MS (APCI+) m/z 365.1 (M+H⁺, 100%).

Example 2

35 1-(8-Fluoroquinolin-4-yl)-3-(6-(tetrahydro-2H-pyran-4-yl)pyridin-2-yl)urea




A solution of 1-(6-(3,6-dihydro-2H-pyran-4-yl)pyridin-2-yl)-3-(8-fluoroquinolin-4-yl)urea (21 mg, 0.058 mmol) and 12.3 mg of palladium on charcoal in 20 mL of methanol together with 1 mL of DMF and 1 drop of AcOH was stirred under 1 atm of hydrogen for 18h. Filtration and evaporation to dryness afforded 21 mg (99%) of the desired title product.

¹H-NMR (d₆-DMSO, 400 MHz) δ 1.15 (m, 2H), 1.78 (m, 4H), 2.95 (m, 1H), 3.95 (d, 2H), 6.97 (d, 1H), 7.48 (br s, 1H), 7.65 (m, 2H), 7.75 (t, 1H), 8.08 (m, 1H), 8.35 (d, 1H), 9.82 (m, 1H), 10.11 (s, 1H); MS (APCI+) m/z 367.1 (M+H⁺, 100%).

Example 3


1-(6-(3,4-Dihydro-2H-pyran-5-yl)pyridin-2-yl)-3-(8-fluoroquinolin-4-yl)urea



Prepared by the method described for Example 1.
MS (APCI+) m/z 365.1 (M+H⁺, 100%).

Example 4


1-(6-(3,6-Dihydro-2H-pyran-4-yl)pyridin-2-yl)-3-(7-methoxyquinolin-4-yl)urea



Prepared by the method described for Example 1.
MS (APCI+) m/z 377.1 (M+H⁺, 100%).

Example 5

1-(6-(3,4-Dihydro-2H-pyran-5-yl)quinolin-4-yl)-3-(6-(trifluoromethyl)pyridin-2-yl)urea




Prepared by the method described for Example 1.

MS (APCI+) m/z 415.1 ($M+H^+$, 100%).

5

Example 6

1-(7-(3,6-Dihydro-2H-pyran-4-yl)quinolin-4-yl)-3-(6-(trifluoromethyl)pyridin-2-yl)urea




10 Prepared by the method described for Example 1.

MS (APCI+) m/z 415.1 ($M+H^+$, 100%).

Example 7

1-(6-(Tetrahydro-2H-pyran-4-yl)quinolin-4-yl)-3-(6-(trifluoromethyl)pyridin-2-yl)urea




15

Prepared by the method described for Example 2.

MS (APCI+) m/z 417.1 ($M+H^+$, 100%).

20 Example 8

1-(7-Methoxyquinolin-4-yl)-3-(6-(tetrahydro-2H-pyran-4-yl)pyridin-2-yl)urea




Prepared by the method described for Example 2.

MS (APCI+) m/z 379.1 ($M+H^+$, 100%).

5 Example 9

1-(7-(Tetrahydro-2H-pyran-4-yl)quinolin-4-yl)-3-(6-(trifluoromethyl)pyridin-2-yl)urea




Prepared by the method described for Example 2.

10 MS (APCI+) m/z 417.1 ($M+H^+$, 100%).

Example 10

1-(6-(Tetrahydro-2H-pyran-3-yl)quinolin-4-yl)-3-(6-(trifluoromethyl)pyridin-2-yl)urea




15

Prepared by the method described for Example 2.

MS (APCI+) m/z 417.1 ($M+H^+$, 100%).

20 Example 11

1-(7-(3,6-Dihydro-2H-pyran-4-yl)quinolin-4-yl)-3-(pyrazin-2-yl)urea




Prepared by the method described for Example 1.

25 MS (APCI+) m/z 348.1 ($M+H^+$, 100%).

Example 12

1-(6-(3,4-Dihydro-2H-pyran-5-yl)pyridin-2-yl)-3-(7-methoxyquinolin-4-yl)urea




Prepared by the method described for Example 1.

5 MS (APCI+) m/z 377.1 ($M+H^+$, 100%).

Example 13

1-(7-Methoxyquinolin-4-yl)-3-(6-(tetrahydro-2H-pyran-3-yl)pyridin-2-yl)urea




10

Prepared by the method described for Example 2.

MS (APCI+) m/z 379.2 ($M+H^+$, 100%).

Example 14

15 1-(Pyrazin-2-yl)-3-(7-(tetrahydro-2H-pyran-4-yl)quinolin-4-yl)urea




Prepared by the method described for Example 2.

MS (APCI+) m/z 350.2 ($M+H^+$, 100%).

20

Example 15


1-(7-(3,4-Dihydro-2H-pyran-5-yl)quinolin-4-yl)-3-(6-(trifluoromethyl)pyridin-2-yl)urea



25 Prepared by the method described for Example 1.

MS (APCI+) m/z 415.1 (M+H⁺, 100%).

Example 16: 1-(6-Bromo-quinolin-4-yl)-3-(6-trifluoromethyl-pyridin-2-yl)-urea




5

366 mg of bis(trichloromethyl)carbonate (1.234mmol) were dissolved in 7 ml of di-chloromethane (DCM) and cooled to 0°C. To this mixture a solution of 400 mg 6-(trifluoromethyl)pyridin-2-amine (2.467mmol) and 379 µl triethylamine (5.43 mmol) in 10 9.7 ml DCM was slowly added over 60 minutes at 0°C and stirring was then continued for additional 60 minutes at 23°C. 550 mg 6-bromoquinolin-4-amine (2.467mmol) and 379 µl triethylamine (5.43 mmol) were suspended in 9.7 ml DCM and the suspension was slowly added to the reaction mixture at 23°C, which became a clear solution after 45 minutes. After stirring over night the reaction mixture was poured into ice water and 15 stirred for 2 h. The precipitate formed was filtered off and dried in vacuo for 24 h. 810 mg of the desired product were obtained as off white powder (yield: 80%). The already quite pure raw material was used without further purification.

¹H NMR (DMSO-d₆, 400MHz): δ [ppm]: 10.26 (s, 1 H), 9.70 (s, 1 H), 8.79 (d, 1 H), 8.44 (d, 1 H), 8.28 (d, 1 H), 8.19 (d, 1 H), 8.09 (dd, 1 H), 7.88 (dd, 1H), 7.93 (d, 1H), 7.57 (d, 1H)
ESI-MS [M+H⁺]: 413.0

25


Example 17: 1-(6-(Trifluoromethyl)pyridin-2-yl)-3-quinolin-4-yl-urea



The title compound was prepared as described in example 16, substituting 6-bromoquinolin-4-amine by quinolin-4-ylamine.

ESI-MS $[M+H]^+$: 333.1


5 Example 18: 1-(6,8-Difluoroquinolin-4-yl)-3-(6-trifluoromethyl-pyridin-2-yl)-urea



10 The title compound was prepared as described in example 16, substituting 6-bromoquinolin-4-amine by 6,8-difluoro-quinolin-4-ylamine.

ESI-MS $[M+H]^+$: 369.0

Example 19: 1-(7-Bromoquinolin-4-yl)-3-(6-trifluoromethyl-pyridin-2-yl)-urea




The title compound was prepared as described in example 16, substituting 6-bromoquinolin-4-amine by 7-bromoquinolin-4-amine.

ESI-MS $[M+H]^+$: 410.9/412.9

20


Example 20: 1-(7-Trifluoromethylquinolin-4-yl)-3-(6-trifluoromethyl-pyridin-2-yl)-urea



The title compound was prepared as described in example 16, substituting 6-bromoquinolin-4-amine by 7-trifluoromethyl-quinolin-4-ylamine.

ESI-MS $[M+H]^+$: 401.1


5 Example 21: 1-(7-Methoxyquinolin-4-yl)-3-(6-trifluoromethyl-pyridin-2-yl)-urea



10 The title compound was prepared as described in example 16, substituting 6-bromoquinolin-4-amine by 7-methoxy-quinolin-4-ylamine.

ESI-MS $[M+H]^+$: 363.1

Example 22: 1-(8-Trifluoromethylquinolin-4-yl)-3-(6-trifluoromethyl-pyridin-2-yl)-urea




15

The title compound was prepared as described in example 16, substituting 6-bromoquinolin-4-amine by 8-trifluoromethyl-quinolin-4-ylamine.

ESI-MS $[M+H]^+$: 401.1

20


Example 23: 1-(8-Cyanoquinolin-4-yl)-3-(6-trifluoromethyl)pyridin-2-yl)-urea



The title compound was prepared as described in example 16, substituting 6-bromoquinolin-4-amine by 8-cyano-quinolin-4-ylamine.

5 ESI-MS [M+H⁺]: 358.1

Example 24: 1-(8-Iodoquinolin-4-yl)-3-(6-trifluoromethyl-pyridin-2-yl)-urea




10

The title compound was prepared as described in example 16, substituting 6-bromoquinolin-4-amine by 8-iodo-quinolin-4-ylamine.

ESI-MS [M+H⁺]: 459.0/459.9


15 Example 25: 1-(8-Cyanoquinolin-4-yl)-3-pyrazin-2-yl-urea



20 The title compound was prepared as described in example 16, substituting 6-bromoquinolin-4-amine by 8-cyano-quinolin-4-ylamine and 6-trifluoromethylpyridine-2-amine by pyrazine-2-ylamine.

ESI-MS [M+H⁺]: 291.1

Example 26: 1-(7-Methoxyquinolin-4-yl)-3-pyrazin-2-yl-urea




5

The title compound was prepared as described in example 16, substituting 6-bromoquinolin-4-amine by 7-methoxy-quinolin-4-ylamine and 6-trifluoromethylpyridine-2-amine by pyrazine-2-ylamine..

ESI-MS $[M+H]^+$: 296.1

10

Example 27: tert-Butyl 4-{4-[3-(6-trifluoromethyl-pyridin-2-yl)-ureido]-quinolin-6-yl}-3,6-dihydro-2H-pyridine-1-carboxylate



15

Step 1: 1-(6-Bromo-quinolin-4-yl)-3-(6-trifluoromethyl-pyridin-2-yl)-urea

366 mg of bis(trichloromethyl)carbonate (1,234 mmol) were dissolved in 7 ml of DCM and cooled to 0°C. To this mixture the solution of 400 mg of 6-(trifluoromethyl)pyridin-2-amine (2.467 mmol) and 379 µl of triethylamine (5.43 mmol) in 9.7ml DCM was slowly added over 60 minutes at 0°C, and stirring was continued for an additional 60 minutes at room temperature. 550 mg of 6-bromoquinolin-4-amine (2.467 mmol) and 379 µl of triethylamine (5.43 mmol) were suspended in 9.7 ml and subsequently slowly added to the reaction mixture at RT, which became a clear solution after 45 minutes. After stirring over night the reaction mixture was poured into ice water and stirred for 2h. The precipitation formed was filtered off and dried in vacuum for 24h. 810 mg of the desired product were obtained as an off white powder (yield: 80%). The quite pure raw material was used without further purification in the next step.

25

¹H NMR (DMSO-d₆, 400MHz): δ [ppm]: 10.26 (s, 1 H), 9.70 (s, 1 H), 8.79 (d, 1 H), 8.44 (d, 1 H), 8.28 (d, 1 H), 8.19 (d, 1 H), 8.09 (dd, 1 H), 7.88 (dd, 1H), 7.93 (d, 1H), 7.57 (d, 1H)

ESI-MS [M+H⁺]: 413.0


5

Step 2: tert-Butyl 4-{4-[3-(6-trifluoromethyl-pyridin-2-yl)-ureido]-quinolin-6-yl}-3,6-dihydro-2H-pyridine-1-carboxylate

1.500g of 1-(6-bromo-quinolin-4-yl)-3-(6-trifluoromethyl-pyridin-2-yl)-urea (3.65 mmol),
 10 1.692 g of tert-butyl 4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-5,6-dihydropyridine-1(2H)-carboxylate (5.47 mmol), 0.422g of tetrakis(triphenylphosphine)palladium(0) (0.365 mmol) and 8.21 ml of a 2M sodium carbonate solution (16.42 mmol) were dissolved/suspended in 32ml DMF in a microwave vessel, flushed with vacuum/argon, and subsequently heated in the microwave for 1h at 80°C. HPLC/MSD indicated almost
 15 complete formation of the desired product. The reaction mixture was poured on ice, stirred for 1h, and the precipitation formed was filtered off and dried in vacuum for 24h. 2 g of the desired product were obtained as an off white powder (yield: quant.). The quite pure raw material was used without further purification in the next step.
 ESI-MS [M+H⁺]: 514.2

20

Example 28: 1-[6-(1,2,3,6-Tetrahydro-pyridin-4-yl)-quinolin-4-yl]-3-(6-trifluoromethyl-pyridin-2-yl)-urea



25


0.2 g of tert-butyl 4-{4-[3-(6-trifluoromethyl-pyridin-2-yl)-ureido]-quinolin-6-yl}-3,6-dihydro-2H-pyridine-1-carboxylate (0.389 mmol) were dissolved in 0.6 ml DCM (yellow solution). 0.300 ml of TFA (3.89 mmol) were added and the reaction mixture was stirred for 2h at RT. After evaporation of the solvent and excess TFA in vacuum, water
 30 was added to the residue, the acidic mixture extracted with ethyl acetate, the water phase adjusted to pH 9 with 1M NaOH and subsequently extracted three times with DCM. The combined DCM solutions were washed with a small amount of brine, dried over sodium sulphate and the solvent was removed in vacuum. 130 mg of the desired product were obtained as light yellow powder (yield: 81%).

35

¹H NMR (DMSO-d₆, 500MHz): δ [ppm]: 10.81 (broad s, 1 H), 10.11 (broad s, 1 H), 8.67 (d, 1 H), 8.28 (d, 1 H), 8.23 (s, 1 H), 8.20 (d, 1 H), 8.06 (dd, 1 H), 7.88 and 7.91 (2d,

2H), 7.53 (d, 1H), 6.47 (broad s, 1H), 3.45 (broad s, 2H), 3.3 (very broad s), 3.00 (t, 2H), 2.56 (broad s, 2H)
ESI-MS [M+H⁺]: 414.1

5 Example 29: 1-(6-(Piperidin-4-yl)quinolin-4-yl)-3-(6-(trifluoromethyl)pyridin-2-yl)urea




0.59 g of 1-(6-(1,2,3,6-tetrahydropyridin-4-yl)quinolin-4-yl)-3-(6-(trifluoromethyl)pyridin-2-yl)urea (0.427 mmol) were dissolved in a mixture of 143 ml of dioxane and 10 ml of ethanol. After addition of 0.152 g of Pd/C (1.427 mmol) and 0.082 ml of acetic acid (1.427 mmol), hydrogenation was performed for 7h at 60°C and over night at room temperature, which led to complete double bond hydrogenation according to HPLC/MSD analysis. The catalyst was filtered off, the solvent removed under reduced pressure and the residue extracted three times with ethyl acetate after adding water and a small amount of NaOH (pH 9). The combined organic phases were washed with a small amount of brine, dried over sodium sulfate and the solvent was removed under reduced pressure. 593 mg of the desired product were obtained as light yellow powder (yield: 64%).

¹H NMR (DMSO-d₆, 500MHz): δ [ppm]: 10.64 (broad s, 1 H), 9.79 (broad s, 1 H), 8.70 (d, 1 H), 8.21-8.24 (m, 2H), 8.05-8.09 (m, 2H), 7.93 (d, 1H), 7.66 (d, 1H), 7.54 (d, 1H), 3.74 (very broad s), 3.16 (broad d, 2H), 2.82-2.89 (m, 1H), 2.69-2.77 (m, 2H), 1.75-1.89 (m, 4H),

ESI-MS [M+H⁺]: 416.2

Example 30: 1-[6-(1-Methyl-1,2,3,6-tetrahydro-pyridin-4-yl)-quinolin-4-yl]-3-(6-trifluoromethyl-pyridin-2-yl)-urea



30


To 0.12 g of 1-[6-(1,2,3,6-tetrahydro-pyridin-4-yl)-quinolin-4-yl]-3-(6-trifluoromethyl-pyridin-2-yl)-urea (0.290 mmol) were added subsequently 2 ml of methanol, 0.366 ml of formaldehyde (4.64 mmol) and a solution of 0.055 g of sodium borohydride (1.451

mmol) in 0.8 ml of water. The reaction was slightly exotherm. A small amount of 2M HCl was added to the reaction mixture up to a pH of 3 to destroy the excess of sodium bororhydride. After neutralization with sodium bicarbonate, methanol was evaporated under reduced pressure, water added to the residue, the reaction mixture extracted three times with ethylacetate, the organic phases combined and dried over sodium sulfate and the solvent evaporated under reduced pressure. 120 mg of the desired product were obtained as an off white powder (yield: 97%).

^1H NMR (DMSO- d_6 , 500MHz): δ [ppm]: 10.45 (broad s, 1 H), 9.47 (broad s, 1 H), 8.67 (d, 1 H), 8.28 (d, 1 H), 8.22 (d, 1 H), 8.10 (s, 1 H), 8.06 (dd, 1 H), 7.89 (dd, 2 H), 7.53 (d, 1 H), 6.39 (broad s, 1H), 3.40 (s, 3H), 3.09 (s, 2H), 2.68 (broad s, 2H), 2.64-2.66 (m, 2H).

ESI-MS $[\text{M}+\text{H}^+]$: 428.2

- Example 31: 1-[6-(1-Methyl-piperidin-4-yl)-quinolin-4-yl]-3-(6-trifluoromethyl-pyridin-2-yl)-urea




- The title compound was prepared as described in example 29, substituting 1-(6-(1,2,3,6-tetrahydropyridin-4-yl)quinolin-4-yl)-3-(6-(trifluoromethyl)pyridin-2-yl)urea by 1-[6-(1-Methyl-1,2,3,6-tetrahydro-pyridin-4-yl)-quinolin-4-yl]-3-(6-trifluoromethyl-pyridin-2-yl)-urea.

ESI-MS $[\text{M}+\text{H}^+]$: 430.1

25

Example 32: 1-[6-[1-(2-Fluoro-ethyl)-piperidin-4-yl]-quinolin-4-yl]-3-(6-trifluoromethyl-pyridin-2-yl)-urea




30

The reaction mixture containing 0.27 g of 1-(6-(piperidin-4-yl)quinolin-4-yl)-3-(6-(trifluoromethyl)pyridin-2-yl)urea (0.650 mmol), 0.073 ml of 1-bromo-2-fluoroethane (0.975 mmol), 1.059g of cesium carbonate (3.25 mmol) and 6.5 of dioxane was shaken

in a flask at 60°C for 4h. Due to non complete reaction, additional 0.049 ml of 1-bromo-2-fluoroethane (0.650 mmol) were added and the reaction mixture was shaken at 60°C for 18h. Saturated sodium bicarbonate solution was added and the mixture extracted three times with ethyl acetate. The combined organic phases were washed once with
 5 brine, dried over sodiumsulfat and the solvent removed under reduced pressure. After purifying of the raw material (190 mg) by column chromatography (silica gel, DCM/MeOH 95:5) 90mg of the desired product were obtained as beige powder (yield: 30%).

10 ESI-MS [M+H⁺]: 462.2

Example 33: 1-[6-[1-(2,2-Difluoro-ethyl)-piperidin-4-yl]-quinolin-4-yl]-3-(6-trifluoromethyl-pyridin-2-yl)-urea




15

The title compound was prepared as described in example 32, substituting 1-bromo-2-fluoroethane by 1,1-difluoro-2-iodoethane.

20 ESI-MS [M+H⁺]: 480.2

Example 34: 1-[6-(3,6-Dihydro-2H-pyran-4-yl)-quinolin-4-yl]-3-(6-trifluoromethyl-pyridin-2-yl)-urea




25

The title compound was prepared as described in example 27 step 2, substituting tert-butyl 4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-5,6-dihydropyridine-1(2H)-carboxylate by 2-(3,6-dihydro-2H-pyran-4-yl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane.

30

ESI-MS [M+H⁺]: 415.1

Example 35: 1-[6-(3,6-Dihydro-2H-pyran-4-yl)-quinolin-4-yl]-3-pyrazin-2-yl-urea



Step 1: 1-(6-Bromoquinolin-4-yl)-3-(pyrazin-2-yl)urea

5

The title compound was prepared as described in example 27 step 1, substituting 6-(trifluoromethyl)pyridin-2-amine by 6-bromoquinolin-4-amine and 6-bromoquinolin-4-amine by pyrazin-2-amine.


10 ESI-MS [M+H⁺]: 345.9

Step 2: 1-[6-(3,6-Dihydro-2H-pyran-4-yl)-quinolin-4-yl]-3-pyrazin-2-yl-urea

The title compound was prepared as described in example 27 step 2, substituting 1-(6-bromo-quinolin-4-yl)-3-(6-trifluoromethyl-pyridin-2-yl)-urea by 1-(6-bromoquinolin-4-yl)-3-(pyrazin-2-yl)urea and tert-butyl 4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-5,6-dihydropyridine-1(2H)-carboxylate by 2-(3,6-dihydro-2H-pyran-4-yl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane.

20 ESI-MS [M+H⁺]: 348.1

Example 36: tert-Butyl 4-[4-(3-pyrazin-2-ylureido)quinolin-6-yl]-5,6-dihydropyridine-1(2H)-carboxylate




25

The title compound was prepared as described in example 27 step 2, substituting 1-(6-bromo-quinolin-4-yl)-3-(6-trifluoromethyl-pyridin-2-yl)-urea by 1-(6-bromoquinolin-4-yl)-3-(pyrazin-2-yl)urea (example 35 step 1).

30

ESI-MS [M+H⁺]: 447.2


Example 37: 1-Pyrazin-2-yl-3-[6-(1,2,3,6-tetrahydro-pyridin-4-yl)-quinolin-4-yl]-urea



The title compound was prepared as described in example 28, substituting tert-butyl 4-{4-[3-(6-trifluoromethyl-pyridin-2-yl)-ureido]-quinolin-6-yl}-3,6-dihydro-2H-pyridine-1-carboxylate by tert-butyl 4-[4-(3-pyrazin-2-ylureido)quinolin-6-yl]-5,6-dihydropyridine-1(2H)-carboxylate (example 36).

ESI-MS [M+H⁺]: 347.1


10 Example 38: 1-(6-Piperidin-4-yl-quinolin-4-yl)-3-pyrazin-2-yl-urea



The title compound was prepared as described in example 29, substituting 1-(6-(1,2,3,6-tetrahydropyridin-4-yl)quinolin-4-yl)-3-(6-(trifluoromethyl)pyridin-2-yl)urea by 1-pyrazin-2-yl-3-[6-(1,2,3,6-tetrahydro-pyridin-4-yl)-quinolin-4-yl]-urea (example 37).

ESI-MS [M+H⁺]: 349.2


20 Example 39: 1-{6-[1-(2-Fluoro-ethyl)-piperidin-4-yl]-quinolin-4-yl}-3-pyrazin-2-yl-urea



The title compound was prepared as described in example 32, substituting 1-(6-(piperidin-4-yl)quinolin-4-yl)-3-(6-(trifluoromethyl)pyridin-2-yl)urea by 1-(6-piperidin-4-yl-quinolin-4-yl)-3-pyrazin-2-yl-urea (example 38).

ESI-MS [M+H⁺]: 395.2


30 Example 40: 1-[6-(1-Methyl-piperidin-4-yl)-quinolin-4-yl]-3-pyrazin-2-yl-urea



The title compound was prepared as described in example 30, substituting 1-[6-(1,2,3,6-tetrahydropyridin-4-yl)-quinolin-4-yl]-3-(6-trifluoromethyl-pyridin-2-yl)-urea by
 5 1-(6-piperidin-4-yl-quinolin-4-yl)-3-pyrazin-2-yl-urea (example 38).

ESI-MS $[M+H]^+$: 363.2

Example 41: tert-Butyl 4-{4-[3-(6-cyclopropyl-pyrazin-2-yl)-ureido]-quinolin-6-yl}-3,6-
 10 dihydro-2H-pyridine-1-carboxylate



Step 1: 1-(6-Bromoquinolin-4-yl)-3-(6-cyclopropylpyrazin-2-yl)urea
 15

The title compound was prepared as described in example 27 step 1, substituting 6-(trifluoromethyl)pyridin-2-amine by 6-bromoquinolin-4-amine and 6-bromoquinolin-4-amine by 6-cyclopropylpyrazin-2-amine.


20 ESI-MS $[M+H]^+$: 386.0

Step 2: Tert-butyl 4-{4-[3-(6-cyclopropyl-pyrazin-2-yl)-ureido]-quinolin-6-yl}-3,6-dihydro-2H-pyridine-1-carboxylate

25 The title compound was prepared as described in example 27 step 2, substituting 1-(6-bromo-quinolin-4-yl)-3-(6-trifluoromethyl-pyridin-2-yl)-urea by 1-(6-bromoquinolin-4-yl)-3-(6-cyclopropylpyrazin-2-yl)urea (example 15 step 1).

ESI-MS $[M+H]^+$: 487.2
 30


Example 42: tert-Butyl 4-{4-[3-(6-cyclopropyl-pyrazin-2-yl)-ureido]-quinolin-6-yl}-piperidine-1-carboxylate



The title compound was prepared as described in example 29, substituting 1-(6-(1,2,3,6-tetrahydropyridin-4-yl)quinolin-4-yl)-3-(6-(trifluoromethyl)pyridin-2-yl)urea by
 5 tert-butyl 4-{4-[3-(6-cyclopropyl-pyrazin-2-yl)-ureido]-quinolin-6-yl}-3,6-dihydro-2H-pyridine-1-carboxylate (example 41).

ESI-MS [M+H⁺]: 489.2

10 Example 43: 1-(6-Cyclopropyl-pyrazin-2-yl)-3-(6-piperidin-4-yl-quinolin-4-yl)-urea




The title compound was prepared as described in example 28, substituting tert-butyl 4-{4-[3-(6-trifluoromethyl-pyridin-2-yl)-ureido]-quinolin-6-yl}-3,6-dihydro-2H-pyridine-1-carboxylate by tert-butyl 4-{4-[3-(6-cyclopropyl-pyrazin-2-yl)-ureido]-quinolin-6-yl}-
 15 piperidine-1-carboxylate (example 42).

ESI-MS [M+H⁺]: 389.2

20

Example 44: 1-(6-Cyclopropyl-pyrazin-2-yl)-3-[6-(1-methyl-piperidin-4-yl)-quinolin-4-yl]-urea




25

The title compound was prepared as described in example 30, substituting 1-[6-(1,2,3,6-tetrahydro-pyridin-4-yl)-quinolin-4-yl]-3-(6-trifluoromethyl-pyridin-2-yl)-urea by
 1-(6-cyclopropyl-pyrazin-2-yl)-3-(6-piperidin-4-yl-quinolin-4-yl)-urea (example 43).

30 ESI-MS [M+H⁺]: 403.1

Example 45: tert-Butyl 3-{4-[3-(6-Trifluoromethyl-pyridin-2-yl)-ureido]-quinolin-6-yl}-2,5-dihydro-pyrrole-1-carboxylate



5


The title compound was prepared as described in example 27 step 2, substituting tert-butyl 4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-5,6-dihydropyridine-1(2H)-carboxylate by tert-butyl 3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-2,5-dihydro-1H-pyrrole-1-carboxylate.

10

ESI-MS [M+H⁺]: 500.2

Example 46: 1-[6-(2,5-Dihydro-1H-pyrrol-3-yl)-quinolin-4-yl]-3-(6-trifluoromethyl-pyridin-2-yl)-urea

15




The title compound was prepared as described in example 28, substituting tert-butyl 4-{4-[3-(6-trifluoromethyl-pyridin-2-yl)-ureido]-quinolin-6-yl}-3,6-dihydro-2H-pyridine-1-carboxylate by tert-butyl 3-{4-[3-(6-trifluoromethyl-pyridin-2-yl)-ureido]-quinolin-6-yl}-2,5-dihydro-pyrrole-1-carboxylate (example 45).

20

ESI-MS [M+H⁺]: 400.1

Example 47: 1-[6-(3,6-Dihydro-2H-pyran-4-yl)-pyridin-2-yl]-3-(8-fluoro-quinolin-4-yl)-urea

25



Step 1: 1-(8-Fluoroquinolin-4-yl)-3-(6-iodopyridin-2-yl)urea

30

The title compound was prepared as described in example 27 step 1, substituting 6-(trifluoromethyl)pyridin-2-amine by 4-amino-8-fluoroquinoline and 6-bromoquinolin-4-amine by 6-iodopyridin-2-amine .


5 ESI-MS [M+H⁺]: 409.0

Step 2: 1-[6-(3,6-Dihydro-2H-pyran-4-yl)-pyridin-2-yl]-3-(8-fluoro-quinolin-4-yl)-urea

10 The title compound was prepared as described in example 27 step 2, substituting 1-(6-bromo-quinolin-4-yl)-3-(6-trifluoromethyl-pyridin-2-yl)-urea by 1-(8-fluoroquinolin-4-yl)-3-(6-iodopyridin-2-yl)urea (example 47, step 1) and tert-butyl 4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-5,6-dihydropyridine-1(2H)-carboxylate by 2-(3,6-dihydro-2H-pyran-4-yl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane.

15 ESI-MS [M+H⁺]: 365.1

Example 48: 1-(8-Fluoro-quinolin-4-yl)-3-[6-(tetrahydro-pyran-4-yl)-pyridin-2-yl]-urea



20


The title compound was prepared as described in example 29, substituting 1-(6-(1,2,3,6-tetrahydropyridin-4-yl)quinolin-4-yl)-3-(6-(trifluoromethyl)pyridin-2-yl)urea by 1-[6-(3,6-dihydro-2H-pyran-4-yl)-pyridin-2-yl]-3-(8-fluoro-quinolin-4-yl)-urea (example 47) and by performing the hydrogenation at room temperature for at least 24h .

25

ESI-MS [M+H⁺]: 367.15

Example 49: 1-(8-Chloro-6-methyl-quinolin-4-yl)-3-[6-(tetrahydro-pyran-4-yl)-pyridin-2-yl]-urea


30



The title compound was prepared as described in example 27 step 1, substituting 6-(trifluoromethyl)pyridin-2-amine by 4-amino-8-chloro-6-methylquinoline and 6-bromoquinolin-4-amine by 6-(tetrahydro-2H-pyran-4-yl)pyridin-2-amine .

5 ESI-MS $[M+H]^+$: 397.1

Example 50: 1-(6,8-Dichloro-quinolin-4-yl)-3-[6-(tetrahydro-pyran-4-yl)-pyridin-2-yl]-urea




The title compound was prepared as described in example 27 step 1, substituting 6-(trifluoromethyl)pyridin-2-amine by 6-(tetrahydro-2H-pyran-4-yl)pyridin-2-amine and 6-bromoquinolin-4-amine by 4-amino-6,8-dichloroquinoline.

15

ESI-MS $[M+H]^+$: 418.1

Example 51: 1-(6,8-Difluoro-quinolin-4-yl)-3-[6-(tetrahydro-pyran-4-yl)-pyridin-2-yl]-urea




The title compound was prepared as described in example 27 step 1, substituting 6-(trifluoromethyl)pyridin-2-amine by 4-amino-6,8-difluoroquinoline and 6-bromoquinolin-4-amine by 6-(tetrahydro-2H-pyran-4-yl)pyridin-2-amine .

25

ESI-MS $[M+H]^+$: 385.1

Example 52: 1-(8-Chloro-quinolin-4-yl)-3-[6-(tetrahydro-pyran-4-yl)-pyridin-2-yl]-urea




The title compound was prepared as described in example 27 step 1, substituting 6-(trifluoromethyl)pyridin-2-amine by 6-(tetrahydro-2H-pyran-4-yl)pyridin-2-amine and 6-bromoquinolin-4-amine by 8-chloroquinolin-4-amine.

5

ESI-MS $[M+H]^+$: 383.2

Example 53: 1-[1,5]Naphthyridin-4-yl-3-[6-(tetrahydro-pyran-4-yl)-pyridin-2-yl]-urea




10

The title compound was prepared as described in example 27 step 1, substituting 6-(trifluoromethyl)pyridin-2-amine by 6-(tetrahydro-2H-pyran-4-yl)pyridin-2-amine and 6-bromoquinolin-4-amine by 1,5-naphthyridin-4-amine.

15

ESI-MS $[M+H]^+$: 350.2

Example 54: 1-(5,8-Difluoro-quinolin-4-yl)-3-[6-(tetrahydro-pyran-4-yl)-pyridin-2-yl]-urea



20


The title compound was prepared as described in example 1 step 1, substituting 6-(trifluoromethyl)pyridin-2-amine by 5,8-difluoroquinolin-4-amine and 6-bromoquinolin-4-amine by 6-(tetrahydro-2H-pyran-4-yl)pyridin-2-amine .

25

ESI-MS $[M+H]^+$: 385.1

Example 55: 1-(8-Fluoro-6-methoxy-quinolin-4-yl)-3-[6-(tetrahydro-pyran-4-yl)-pyridin-2-yl]-urea

30




The title compound was prepared as described in example 27 step 1, substituting 6-(trifluoromethyl)pyridin-2-amine by 8-fluoro-6-methoxyquinolin-4-amine and 6-bromoquinolin-4-amine by 6-(tetrahydro-2H-pyran-4-yl)pyridin-2-amine .

5

ESI-MS $[M+H]^+$: 397.2

Example 56: 1-[6-(5,6-Dihydro-4H-pyran-3-yl)-pyridin-2-yl]-3-(8-fluoro-quinolin-4-yl)-urea

10




The title compound was prepared as described in example 27 step 2, substituting 1-(6-bromo-quinolin-4-yl)-3-(6-trifluoromethyl-pyridin-2-yl)-urea by 1-(8-fluoroquinolin-4-yl)-3-(6-iodopyridin-2-yl)urea (example 47, step 1) and tert-butyl 4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-5,6-dihydropyridine-1(2H)-carboxylate by 2-(3,4-dihydro-2H-pyran-5-yl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane.

15

ESI-MS $[M+H]^+$: 365.1

20

Example 57: 1-(8-Fluoro-quinolin-4-yl)-3-[6-(tetrahydro-pyran-3-yl)-pyridin-2-yl]-urea




25

The title compound was prepared as described in example 29, substituting 1-(6-(1,2,3,6-tetrahydropyridin-4-yl)quinolin-4-yl)-3-(6-(trifluoromethyl)pyridin-2-yl)urea by 1-[6-(5,6-dihydro-4H-pyran-3-yl)-pyridin-2-yl]-3-(8-fluoro-quinolin-4-yl)-urea (example 56) and by performing the hydrogenation at room temperature for at least 24h .

30

ESI-MS $[M+H]^+$: 367.15

Example 58: 1-[6-(3,6-Dihydro-2H-pyran-4-yl)-pyridin-2-yl]-3-[1,5]naphthyridin-4-yl-urea



Step 1: 1-(6-iodopyridin-2-yl)-3-(1,5-naphthyridin-4-yl)urea

- 5 The title compound was prepared as described in example 27 step 1, substituting 6-(trifluoromethyl)pyridin-2-amine by 6-iodopyridin-2-amine and 6-bromoquinolin-4-amine by 1,5-naphthyridin-4-ylamine.

ESI-MS [M+H⁺]: 392.0

10

Step 2: 1-[6-(3,6-Dihydro-2H-pyran-4-yl)-pyridin-2-yl]-3-[1,5]naphthyridin-4-yl-urea


- The title compound was prepared as described in example 27 step 2, substituting 1-(6-bromo-quinolin-4-yl)-3-(6-trifluoromethyl-pyridin-2-yl)-urea by 1-(6-iodopyridin-2-yl)-3-(1,5-naphthyridin-4-yl)urea (example 58, step 1) and tert-butyl 4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-5,6-dihydropyridine-1(2H)-carboxylate by 2-(3,6-dihydro-2H-pyran-4-yl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane.

15

ESI-MS [M+H⁺]: 348.1

20

Example 59: tert-Butyl-6-[3-(8-fluoro-quinolin-4-yl)-ureido]-3',6'-dihydro-2'H-[2,4']bipyridinyl-1'-carboxylate




25

The title compound was prepared as described in example 27 step 2, substituting 1-(6-bromo-quinolin-4-yl)-3-(6-trifluoromethyl-pyridin-2-yl)-urea by 1-(8-fluoroquinolin-4-yl)-3-(6-iodopyridin-2-yl)urea (example 47 step 1).

30

ESI-MS [M+H⁺]: 464.0


Example 60: 1-(8-Fluoro-quinolin-4-yl)-3-(1',2',3',6'-tetrahydro-[2,4']bipyridinyl-6-yl)-urea



The title compound was prepared as described in example 28, substituting tert-butyl 4-{4-[3-(6-trifluoromethyl-pyridin-2-yl)-ureido]-quinolin-6-yl}-3,6-dihydro-2H-pyridine-1-carboxylate by tert-butyl-6-[3-(8-fluoro-quinolin-4-yl)-ureido]-3',6'-dihydro-2'H-[2,4']bipyridinyl-1'-carboxylate (example 59).

ESI-MS [M+H⁺]: 364.0

Example 61: tert-Butyl-6-[3-(8-fluoro-quinolin-4-yl)-ureido]-3',4',5',6'-tetrahydro-2'H-[2,4']bipyridinyl-1'-carboxylate




0.340 g of tert-butyl-6-[3-(8-fluoro-quinolin-4-yl)-ureido]-3',6'-dihydro-2'H-[2,4']bipyridinyl-1'-carboxylate (0.734 mmol) were dissolved in a mixture of 220 ml of DMF and 125 ml of methanol. After addition of 0.078 g of Pd/C (0.734 mmol) and 0.5 ml of acetic acid (8.73 mmol) hydrogenation was performed for 24h at room temperature, which led to complete double bond hydrogenation according to HPLC/MSD analysis. The catalyst was filtered off, the solvent removed under reduced pressure and the residue extracted three times with ethyl acetate after adding water and a small amount of NaOH (pH 9). The combined organic phases were washed with a small amount of brine, dried over sodium sulfate and the solvent was removed under reduced pressure. 300 mg of the desired product were obtained as a powder (yield: 88%).

ESI-MS [M+H⁺]: 466.2


Example 62: 1-(8-Fluoro-quinolin-4-yl)-3-(1',2',3',4',5',6'-hexahydro-[2,4']bipyridinyl-6-yl)-urea

30



The title compound was prepared as described in example 28, substituting tert-butyl 4-{4-[3-(6-trifluoromethyl-pyridin-2-yl)-ureido]-quinolin-6-yl}-3,6-dihydro-2H-pyridine-1-carboxylate by tert-butyl-6-[3-(8-fluoro-quinolin-4-yl)-ureido]-3',4',5',6'-tetrahydro-2'H-[2,4']bipyridinyl-1'-carboxylate (example 61).
ESI-MS [M+H⁺]: 366.2


Example 63: 1-(8-Fluoro-quinolin-4-yl)-3-(1'-methyl-1',2',3',4',5',6'-hexahydro-[2,4']bipyridinyl-6-yl)-urea



The title compound was prepared as described in example 30, substituting 1-[6-(1,2,3,6-tetrahydro-pyridin-4-yl)-quinolin-4-yl]-3-(6-(trifluoromethyl-pyridin-2-yl)-urea by 1-(8-fluoro-quinolin-4-yl)-3-(1',2',3',4',5',6'-hexahydro-[2,4']bipyridinyl-6-yl)-urea (example 62).

ESI-MS [M+H⁺]: 380.2


Example 64: 1-[1'-(2-Fluoro-ethyl)-1',2',3',4',5',6'-hexahydro-[2,4']bipyridinyl-6-yl]-3-(8-fluoro-quinolin-4-yl)-urea



The title compound was prepared as described in example 32, substituting 1-(6-(piperidin-4-yl)quinolin-4-yl)-3-(6-(trifluoromethyl)pyridin-2-yl)urea by 1-(8-fluoro-quinolin-4-yl)-3-(1',2',3',4',5',6'-hexahydro-[2,4']bipyridinyl-6-yl)-urea (example 62).

ESI-MS [M+H⁺]: 412.2

Example 65: 1-[1'-(2,2-Difluoro-ethyl)-1',2',3',4',5',6'-hexahydro-[2,4']bipyridinyl-6-yl]-3-(8-fluoro-quinolin-4-yl)-urea



5


The title compound was prepared as described in example 32, substituting 1-(6-(piperidin-4-yl)quinolin-4-yl)-3-(6-(trifluoromethyl)pyridin-2-yl)urea by 1-(8-fluoro-quinolin-4-yl)-3-(1',2',3',4',5',6'-hexahydro-[2,4']bipyridinyl-6-yl)-urea (example 62) and 1-bromo-2-fluoroethane by 1,1-difluoro-2-iodoethane.

10

ESI-MS $[M+H]^+$: 430.2

Example 66: 1-(8-Fluoro-quinolin-4-yl)-3-(1'-isopropyl-1',2',3',4',5',6'-hexahydro-[2,4']bipyridinyl-6-yl)-urea

15



20


To a solution of 50 mg of 1-(8-fluoro-quinolin-4-yl)-3-(1',2',3',4',5',6'-hexahydro-[2,4']bipyridinyl-6-yl)-urea (example 62) (0.137 mmol) in 50 ml of acetone were added 26 mg of sodium borohydride (0.684 mmol) in small portions. The reaction was slightly exotherm. After stirring the reaction mixture over night a small amount of 2M HCl was added to the reaction mixture up to a pH of 3 to destroy the excess of sodium borohydride. After neutralization with sodium bicarbonate, acetone was evaporated under reduced pressure, water added to the residue, the reaction mixture extracted three times with DCM, the organic phases combined and dried over sodium sulfate and the solvent evaporated under reduced pressure. 45 mg of the desired product were obtained as off white powder (yield: 81 %) after column chromatography (silica gel, DCM/MeOH 98/2).

25

30

ESI-MS $[M+H]^+$: 408.2

Example 67: 1-(8-Fluoro-quinolin-4-yl)-3-[5-(tetrahydro-pyran-4-yl)-pyridin-2-yl]-urea



Step 1: 5-(3,6-Dihydro-2H-pyran-4-yl)pyridin-2-amine

5

0.220 g of 5-iodopyridin-2-amine (1.0 mmol), 0.315 g of 2-(3,6-dihydro-2H-pyran-4-yl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (1.50 mmol), 0.116 g of tetrakis(triphenylphosphine)palladium(0) (0.100 mmol) and 2.25 ml of a 2M sodium carbonate solution (4.50 mmol) were dissolved/suspended in 8.6 ml of DMF in a microwave vessel, flushed with vacuum/argon, and subsequently heated in the microwave for 20 min at 120°C. HPLC/MSD indicated almost complete formation of the desired product. The solvent was evaporated under reduced pressure, 2M HCl was added to the residue and the mixture extracted three times with DCM. To the acidic water phase 2M NaOH was added up to pH 9 and the mixture was extracted three times with ethyl acetate. The combined ethyl acetate solutions were washed once with brine, dried over sodium sulfate and the solvent removed under reduced pressure. 195 mg of the desired product were obtained as yellow powder (yield: quant.). The quite pure raw material (about 90%) was used without further purification in the next step.

20 ESI-MS [M+H⁺]: 177.1

Step 2: 5-(Tetrahydro-2H-pyran-4-yl)pyridin-2-amine

The title compound was prepared as described in example 61, substituting tert-butyl-6-[3-(8-fluoro-quinolin-4-yl)-ureido]-3',6'-dihydro-2'H-[2,4'']bipyridinyl-1'-carboxylate by 5-(3,6-dihydro-2H-pyran-4-yl)pyridin-2-amine and by using ethanol as solvent.

ESI-MS [M+H⁺]: 179.1

30 Step 3: 1-(8-Fluoro-quinolin-4-yl)-3-[5-(tetrahydro-pyran-4-yl)-pyridin-2-yl]-urea


The title compound was prepared as described in example 27 step 1, substituting 6-(trifluoromethyl)pyridin-2-amine by 8-fluoroquinolin-4-amine and 6-bromoquinolin-4-amine by 5-(tetrahydro-2H-pyran-4-yl)pyridin-2-amine (example 67, step 2).

35

ESI-MS $[M+H]^+$: 367.1

Example 68: tert-Butyl-3-[4-(3-pyrazin-2-yl-ureido)-quinolin-6-yl]-2,5-dihydro-pyrrole-1-carboxylate

5




The title compound was prepared as described in example 27 step 2, substituting 1-(6-bromo-quinolin-4-yl)-3-(6-trifluoromethyl-pyridin-2-yl)-urea by 1-(6-bromoquinolin-4-yl)-3-(pyrazin-2-yl)urea (example 9, step 1) and tert-butyl 4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-5,6-dihydropyridine-1(2H)-carboxylate by tert-butyl 3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-2,5-dihydro-1H-pyrrole-1-carboxylate.

15

ESI-MS $[M+H]^+$: 433.2

Example 69: tert-Butyl-3-[4-(3-pyrazin-2-yl-ureido)-quinolin-6-yl]-pyrrolidine-1-carboxylate




20

The title compound was prepared as described in example 61, substituting tert-butyl-6-[3-(8-fluoro-quinolin-4-yl)-ureido]-3',6'-dihydro-2'H-[2,4']bipyridinyl-1'-carboxylate by tert-butyl-3-[4-(3-pyrazin-2-yl-ureido)-quinolin-6-yl]-2,5-dihydro-pyrrole-1-carboxylate (example 68).

25

ESI-MS $[M+H]^+$: 435.2

Example 70: 1-Pyrazin-2-yl-3-(6-pyrrolidin-3-yl-quinolin-4-yl)-urea



30

To a solution of 50 mg of tert-butyl-3-[4-(3-pyrazin-2-yl-ureido)-quinolin-6-yl]-pyrrolidine-1-carboxylate (example 43) (0,120 mmol) in 5 ml of DCM was added 1 ml of 6M HCl solution in propanol-2 with stirring at room temperature. After reaction over night the solvent and excess HCl was removed in vacuum, and the residue was co-distilled at least three times with propanol-2. 50 mg of the desired product were obtained as powder as the di-hydrochloride salt (yield: quant.).

ESI-MS $[M+H^+]$: 335.1

II. Biological tests

The compounds according to the invention exhibit very good affinities for GSK-3 ($< 1 \mu\text{M}$, frequently $< 100 \text{ nM}$) and exhibited good selectivity against multiple kinase targets.

Methods - biochemical hGSK-3 β assay

Compounds were tested for their ability to inhibit human Glycogen Synthase Kinase-3 β (hGSK-3 β) to phosphorylate biotin-YRRAAVPPSPSLSRHSSPHQ(pS)EDEEE.

Compounds were incubated with $0.5 \mu\text{Ci}$ ^{33}P -ATP, $10 \mu\text{M}$ ATP, 0.0125 U hGSK-3 β (Upstate cell signaling solutions) and $1 \mu\text{M}$ substrate (biotin-YRRAAVPPSPSLSRHSSPHQ(pS)EDEEE) in 50 mM HEPES, 10 mM MgCl_2 , 100 mM Na_3VO_4 , 1 mM DTT, 0.0075% Triton, 2% DMSO (total volume $50 \mu\text{L}$) for 30 minutes at room temperature. The incubation was stopped by addition of an equal volume of 100 mM EDTA, 4 M NaCl. $80 \mu\text{L}$ of this mixture was added to streptavidin-coated Flash-plates (PerkinElmer). Following a wash step, ^{33}P incorporation was quantified on a MicroBeta microplate liquid scintillation counter (PerkinElmer). IC_{50} 's were determined by fitting a sigmoidal dose-response curve to the counts obtained at the different concentrations in GraphPad Prism.

The results of the binding tests are given in the table below.

Example #	GSK-3 β IC ₅₀
1	+++
2	+++
3	+++
16	+++
17	+++
18	+++
19	+++
20	+++
21	+++
22	+++
23	+++
24	+++

n.d. not determined

GSK-3 β IC₅₀ :


+ > 10 μ M

5 ++ from 100nM to 10 μ M

+++ <100 nM

We claim:

1. A heterocyclic compound of the formula (I)



5

the stereoisomers, N-oxides, prodrugs, tautomers and/or physiologically tolerated acid addition salts thereof, and the compounds of the general formula I, wherein at least one of the atoms has been replaced by its stable, non-radioactive isotope, wherein

10

A is selected from the group consisting of CR^{A1}R^{A2} and NR^B; where

R^{A1} and R^{A2} are independently of each other selected from the group consisting of hydrogen, C₁-C₂-alkyl, C₁-C₂-haloalkyl, NH₂ and OH; and

15

R^B is selected from hydrogen, C₁-C₄-alkyl and C₁-C₄-haloalkyl;

X¹ and X² are independently of each other selected from the group consisting of CR² and N;

20

X³, X⁴, X⁵ and X⁶ are independently of each other selected from the group consisting of CR³, CR⁴ and N;

25

with the proviso that no more than two of X³, X⁴, X⁵ and X⁶ are CR⁴;

Y¹, Y², Y³ and Y⁴ are independently of each other selected from the group consisting of CR⁴, CR⁵ and N;

30

with the proviso that at most one of Y¹, Y², Y³ and Y⁴ is N and with the proviso that at most one of Y¹, Y², Y³ and Y⁴ is CR⁴; and

with the proviso that one of Y¹, Y², Y³ and Y⁴ is CR⁴ or C-CF₃ if none of X³, X⁴, X⁵ and X⁶ is CR⁴;

R¹ is selected from hydrogen, C₁-C₄-alkyl and C₁-C₄-haloalkyl;

5

each R² is independently selected from the group consisting of hydrogen, OH, halogen, CN, C₁-C₆-alkyl, C₁-C₆-haloalkyl, C₁-C₆-alkoxy, C₁-C₆-haloalkoxy and NR^aR^b;

10

or two radicals R² bonded at the carbon atoms of groups X¹ and X², together with the carbon atoms to which they are bonded, form a 5- or 6-membered saturated or unsaturated ring which may contain 1 or 2 heteroatoms as ring members selected from the group consisting of N, O and S and which optionally carries 1, 2 or 3 substituents R⁶;

15

each R³ is independently selected from the group consisting of hydrogen, CN, NR^aR^b, OH, halogen, C₁-C₆-alkyl, C₁-C₆-haloalkyl, C₃-C₇-cycloalkyl, C₃-C₇-halocycloalkyl, C₂-C₄-alkenyl, C₂-C₄-haloalkenyl, C₁-C₆-alkoxy, C₁-C₆-haloalkoxy, formyl, C₁-C₆-alkylcarbonyl, C₁-C₆-haloalkylcarbonyl, C₁-C₆-alkoxycarbonyl, C₁-C₆-haloalkoxycarbonyl, C₁-C₆-alkyl-NR^aR^b and an aromatic radical Ar, which is selected from the group consisting of phenyl and a 5- or 6-membered N- or C-bound heteroaromatic radical comprising one nitrogen atom and optionally 1, 2 or 3 further heteroatoms independently selected from O, S and N as ring members, wherein Ar is unsubstituted or carries one or two radicals R⁷ and wherein Ar may also be bonded via a CH₂ group;

20

25

R⁴ is a C-bound saturated or partially unsaturated monocyclic 3-, 4-, 5-, 6- or 7-membered heterocyclic ring containing 1, 2 or 3 heteroatoms or heteroatom-containing groups selected from O, N, S, NO, SO and SO₂ as ring members, where the heterocyclic ring optionally carries 1, 2 or 3 C- or N-bound substituents R⁸;

30

R⁵ is selected from the group consisting of hydrogen, CN, NR^aR^b, OH, halogen, C₁-C₆-alkyl, C₁-C₆-haloalkyl, C₃-C₇-cycloalkyl, C₃-C₇-halocycloalkyl, C₂-C₄-alkenyl, C₂-C₄-haloalkenyl, C₁-C₆-alkoxy, C₁-C₆-haloalkoxy, formyl, C₁-C₆-alkylcarbonyl, C₁-C₆-haloalkylcarbonyl, C₁-C₆-alkoxycarbonyl, C₁-C₆-haloalkoxycarbonyl, C₁-C₆-alkyl-NR^aR^b and an aromatic radical Ar, which is selected from the group consisting of phenyl and a 5- or 6-membered N- or C-bound heteroaromatic radical comprising one nitrogen atom and option-

35

ally 1, 2 or 3 further heteroatoms independently selected from O, S and N as ring members, wherein Ar is unsubstituted or carries one or two radicals R⁷ and wherein Ar may also be bonded via a CH₂ group;

5 R⁶ and R⁸, independently of each other and independently of each occurrence, are selected from the group consisting of CN, NR^aR^b, OH, halogen, C₁-C₆-alkyl, C₁-C₆-haloalkyl, C₃-C₇-cycloalkyl, C₃-C₇-halocycloalkyl, C₂-C₄-alkenyl, C₂-C₄-haloalkenyl, C₁-C₆-alkoxy, C₁-C₆-haloalkoxy, formyl, C₁-C₆-alkylcarbonyl, C₁-C₆-haloalkylcarbonyl, C₁-C₆-alkoxycarbonyl, C₁-C₆-haloalkoxycarbonyl, C₁-C₆-alkyl-NR^aR^b and an aromatic radical Ar, which is
10 selected from the group consisting of phenyl and a 5- or 6-membered N- or C-bound heteroaromatic radical comprising one nitrogen atom and optionally 1, 2 or 3 further heteroatoms independently selected from O, S and N as ring members, wherein Ar is unsubstituted or carries one or two radicals R⁷ and wherein Ar may also be bonded via a CH₂ group;

each R⁷ is independently selected from the group consisting of halogen, CN, C₁-C₆-alkyl, C₁-C₆-haloalkyl, C₃-C₆-cycloalkyl, C₃-C₆-halocycloalkyl, C₁-C₆-alkoxy, C₁-C₆-haloalkoxy, NR^aR^b, a phenyl group and a 5- or 6-membered heteroaromatic radical comprising one nitrogen atom and optionally 1, 2 or 3 further heteroatoms independently selected from O, S and N as ring members, wherein phenyl and the heteroaromatic radical are, independently of each other, unsubstituted or substituted by 1, 2, 3 or 4 radicals selected from halogen, cyano, C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy and C₁-C₄-haloalkoxy; and
20
25

R^a and R^b are independently of each other selected from the group consisting of hydrogen, C₁-C₆-alkyl, C₁-C₄-haloalkyl, C₁-C₄-alkoxy, C₁-C₄-haloalkoxy, C₁-C₄-alkylcarbonyl and C₁-C₄-haloalkylcarbonyl;
30 or R^a and R^b form, together with the nitrogen atom to which they are bonded, a 3-, 4-, 5-, 6- or 7-membered saturated or unsaturated aromatic or non-aromatic N-heterocyclic ring, which may contain 1 further heteroatom or heteroatom containing group selected from the group consisting of O, S, SO, SO₂ and N as a ring member.


35

2. The heterocyclic compound according to claim 1, wherein either one of X³, X⁴, X⁵ and X⁶ is CR⁴ and none of Y¹, Y², Y³ and Y⁴ is CR⁴, or one of Y¹, Y², Y³ and Y⁴ is CR⁴ and none of X³, X⁴, X⁵ and X⁶ is CR⁴.

3. The heterocyclic compound according to claim 1, wherein one of Y¹, Y², Y³ and Y⁴ is CR⁴ if none of X³, X⁴, X⁵ and X⁶ is CR⁴.
4. The heterocyclic compound according to claim 1, wherein one of Y¹, Y², Y³ and Y⁴ is C-CF₃ if none of X³, X⁴, X⁵ and X⁶ is CR⁴.
5. The heterocyclic compound according to any of the preceding claims, where R⁴ is selected from a C-bound saturated or partially unsaturated monocyclic 4-, 5- or 6-membered heterocyclic ring containing 1 or 2 or 3 heteroatoms selected from O, N, S and SO, as ring members, where the heterocyclic ring optionally carries 1, 2 or 3 substituents R⁸.
6. The heterocyclic compound according to claim 5, where R⁴ is selected from C-bound oxetanyl, thietanyl, azetidiny, tetrahydrofuranyl, tetrahydrothienyl, tetrahydrothienyl-1-oxide, pyrrolidinyl, pyrrolinyl, pyrazolidinyl, pyrazolinyl, imidazolidinyl, imidazoliny, tetrahydropyranyl, dihydropyranyl, piperidinyl, tetrahydropyridinyl, dihydropyridinyl, piperazinyl and morpholinyl, where the heterocyclic ring optionally carries 1, 2 or 3 substituents R⁸.
7. The heterocyclic compound according to claim 6, where R⁴ is selected from C-bound oxetanyl, azetidiny, tetrahydrofuranyl, tetrahydrothienyl, tetrahydrothienyl-1-oxide, pyrrolidinyl, pyrrolinyl, tetrahydropyranyl, dihydropyranyl, piperidinyl, tetrahydropyridinyl, dihydropyridinyl, piperazinyl and morpholinyl, where the heterocyclic ring optionally carries 1, 2 or 3 substituents R⁸.
8. The heterocyclic compound according to claim 7, where R⁴ is selected from azetidin-3-yl, tetrahydrofuran-3-yl, pyrrolidin-3-yl, pyrrolin-3-yl, tetrahydropyran-4-yl, tetrahydropyran-3-yl, dihydropyran-4-yl, dihydropyran-3-yl, piperidin-4-yl, 1,2,5,6-tetrahydropyridin-4-yl and 1,2-dihydropyridin-4-yl, where the heterocyclic ring optionally carries 1, 2 or 3 substituents R⁸.
9. The heterocyclic compound according to any of the preceding claims, where R⁸ is selected from C₁-C₆-alkyl, C₁-C₆-haloalkyl, C₃-C₇-cycloalkyl, C₃-C₇-halocycloalkyl, C₂-C₄-alkenyl, C₂-C₄-haloalkenyl, C₁-C₆-alkoxy, C₁-C₆-haloalkoxy, formyl, C₁-C₆-alkylcarbonyl, C₁-C₆-haloalkylcarbonyl, C₁-C₆-alkoxycarbonyl, C₁-C₆-haloalkoxycarbonyl and benzyl and is preferably C₁-C₆-alkyl, C₁-C₆-haloalkyl or C₁-C₆-alkoxycarbonyl.

10. The heterocyclic compound according to any of the preceding claims, where R^8 is N-bound.

5 11. The heterocyclic compound according to any of the preceding claims, where R^4 is selected from following structures:



10

where

R^{8a} is hydrogen or has one of the meanings given in claims 1 or 9 for R^8 ; and

is the attachment point to the remainder of the molecule.

15

12. The heterocyclic compound according to any of the preceding claims, where at most one of X^1 , X^2 , X^3 , X^4 , X^5 and X^6 is N.

20 13. The heterocyclic compound according to claim 12, where X^1 and X^2 are CR^2 and X^3 , X^4 , X^5 and X^6 are CR^3 or CR^4 , or where X^1 and X^2 are CR^2 , X^3 is N and X^4 , X^5 and X^6 are CR^3 or CR^4 .

14. The heterocyclic compound according to any of the preceding claims, where Y^1 , Y^2 , Y^3 and Y^4 are CR^4 or CR^5 or Y^2 is N and Y^1 , Y^3 and Y^4 are CR^4 or CR^5 .

25

15. The heterocyclic compound according to any of the preceding claims, where A is NR^B , preferably NH.

30 16. The heterocyclic compound according to any of the preceding claims, where R^1 is hydrogen or methyl, preferably hydrogen.

17. The heterocyclic compound according to any of the preceding claims, where R^2 is hydrogen.

35 18. The heterocyclic compound according to any of the preceding claims, where R^3 is selected from hydrogen, CN, halogen, C_1 - C_6 -alkyl, C_1 - C_6 -haloalkyl, C_3 - C_7 -

cycloalkyl, C₃-C₇-halocycloalkyl, C₁-C₆-alkoxy, C₁-C₆-haloalkoxy, formyl, C₁-C₆-alkylcarbonyl, C₁-C₆-haloalkylcarbonyl, C₁-C₆-alkoxycarbonyl and C₁-C₆-haloalkoxycarbonyl and is preferably selected from hydrogen, CN, halogen, C₁-C₆-alkyl, C₁-C₆-haloalkyl, C₁-C₆-alkoxy and C₁-C₆-haloalkoxy.

5

19. The heterocyclic compound according to any of the preceding claims, where 0, 1 or 2 of the radicals R³ are different from hydrogen.

10


20. The heterocyclic compound according to any of the preceding claims, where R⁵ is selected from hydrogen, CN, halogen, C₁-C₆-alkyl, C₁-C₆-haloalkyl, C₃-C₇-cycloalkyl, C₃-C₇-halocycloalkyl, C₁-C₆-alkoxy, C₁-C₆-haloalkoxy, formyl, C₁-C₆-alkylcarbonyl, C₁-C₆-haloalkylcarbonyl, C₁-C₆-alkoxycarbonyl and C₁-C₆-haloalkoxycarbonyl and is preferably selected from hydrogen, halogen, C₁-C₄-haloalkyl, C₃-C₇-cycloalkyl and C₃-C₇-halocycloalkyl and in particular from hydrogen, fluorinated C₁-C₂-alkyl and C₃-C₆-cycloalkyl.

15

21. The heterocyclic compound according to any of the preceding claims, where at most one of the radicals R⁵ is different from hydrogen.

20

22. The heterocyclic compound according to any of the preceding claims, of formula I-1



where

25

X³ is N or CH;

Y² is N or CH;

R³¹ has one of the meanings given in claims 1, 18 or 19 for R³ except for hydrogen, and is preferably halogen, C₁-C₄-alkyl or C₁-C₄-alkoxy;

R⁴ has one of the meanings given in claims 1 or 5 to 11;


30

R⁵¹ has one of the meanings given in claims 1, 20 or 21 for R⁵ except for hydrogen;

a is 0, 1 or 2; and

b, c and d are independently of each other 0 or 1, with the proviso that one of b and c is 1.

23. The heterocyclic compound according to any of the preceding claims, of formula I-2



where

Y^2 is N or preferably CH;

- 10 R^{31} has one of the meanings given in claims 1, 18 or 19 for R^3 except for hydrogen, and is preferably selected from the group consisting of halogen, trifluoromethyl, cyano or methoxy;

R^4 has one of the meanings given in claims 1 or 5 to 11;

a is 0, 1 or 2;

- 15 b is 0 or 1.

24. The heterocyclic compound according to claim 23, where Y^2 is CH, b is 0 and a is 0, 1 or 2 and where R^{31} , if present, is selected from the group consisting of halogen, trifluoromethyl, cyano and methoxy.

20

25. A heterocyclic compound selected from the group consisting of
 1-(6-(3,6-dihydro-2H-pyran-4-yl)pyridin-2-yl)-3-(8-fluoroquinolin-4-yl)urea,
 1-(8-fluoroquinolin-4-yl)-3-(6-(tetrahydro-2H-pyran-4-yl)pyridin-2-yl)urea,
 1-(6-(3,4-dihydro-2H-pyran-5-yl)pyridin-2-yl)-3-(8-fluoroquinolin-4-yl)urea,
 25 1-(6-(3,6-dihydro-2H-pyran-4-yl)pyridin-2-yl)-3-(7-methoxyquinolin-4-yl)urea,
 1-(6-(3,4-dihydro-2H-pyran-5-yl)quinolin-4-yl)-3-(6-(trifluoromethyl)pyridin-2-yl)-urea,
 1-(7-(3,6-dihydro-2H-pyran-4-yl)quinolin-4-yl)-3-(6-(trifluoromethyl)pyridin-2-yl)-urea,
 30 1-(6-(tetrahydro-2H-pyran-4-yl)quinolin-4-yl)-3-(6-(trifluoromethyl)pyridin-2-yl)-urea,
 1-(7-methoxyquinolin-4-yl)-3-(6-(tetrahydro-2H-pyran-4-yl)pyridin-2-yl)urea,

- 1-(7-(tetrahydro-2H-pyran-4-yl)quinolin-4-yl)-3-(6-(trifluoromethyl)pyridin-2-yl)-urea,
- 1-(6-(tetrahydro-2H-pyran-3-yl)quinolin-4-yl)-3-(6-(trifluoromethyl)pyridin-2-yl)-urea,
- 5 1-(7-(3,6-dihydro-2H-pyran-4-yl)quinolin-4-yl)-3-(pyrazin-2-yl)urea,
- 1-(6-(3,4-dihydro-2H-pyran-5-yl)pyridin-2-yl)-3-(7-methoxyquinolin-4-yl)urea,
- 1-(7-methoxyquinolin-4-yl)-3-(6-(tetrahydro-2H-pyran-3-yl)pyridin-2-yl)urea,
- 1-(pyrazin-2-yl)-3-(7-(tetrahydro-2H-pyran-4-yl)quinolin-4-yl)urea,
- 1-(7-(3,4-dihydro-2H-pyran-5-yl)quinolin-4-yl)-3-(6-(trifluoromethyl)pyridin-2-yl)-urea,
- 10 urea,
- 1-(6-bromo-quinolin-4-yl)-3-(6-trifluoromethyl-pyridin-2-yl)-urea,
- 1-(6-(trifluoromethyl)pyridin-2-yl)-3-quinolin-4-yl-urea,
- 1-(6,8-difluoroquinolin-4-yl)-3-(6-trifluoromethyl-pyridin-2-yl)-urea,
- 1-(7-bromoquinolin-4-yl)-3-(6-trifluoromethyl-pyridin-2-yl)-urea,
- 15 1-(7-trifluoromethylquinolin-4-yl)-3-(6-trifluoromethyl-pyridin-2-yl)-urea,
- 1-(7-methoxyquinolin-4-yl)-3-(6-trifluoromethyl-pyridin-2-yl)-urea,
- 1-(8-trifluoromethylquinolin-4-yl)-3-(6-trifluoromethyl-pyridin-2-yl)-urea,
- 1-(8-cyanoquinolin-4-yl)-3-(6-trifluoromethyl)pyridin-2-yl)-urea,
- 1-(8-iodoquinolin-4-yl)-3-(6-trifluoromethyl)pyridin-2-yl)-urea,
- 20 1-(8-cyanoquinolin-4-yl)-3-pyrazin-2-yl-urea,
- 1-(7-methoxyquinolin-4-yl)-3-pyrazin-2-yl-urea,
- tert-butyl 4-{4-[3-(6-trifluoromethyl-pyridin-2-yl)-ureido]-quinolin-6-yl}-3,6-dihydro-2H-pyridine-1-carboxylate,
- 1-[6-(1,2,3,6-tetrahydro-pyridin-4-yl)-quinolin-4-yl]-3-(6-trifluoromethyl-pyridin-2-yl)-urea,
- 25 1-(6-(piperidin-4-yl)quinolin-4-yl)-3-(6-(trifluoromethyl)pyridin-2-yl)urea,
- 1-[6-(1-methyl-1,2,3,6-tetrahydro-pyridin-4-yl)-quinolin-4-yl]-3-(6-trifluoromethyl-pyridin-2-yl)-urea,
- 1-[6-(1-methyl-piperidin-4-yl)-quinolin-4-yl]-3-(6-trifluoromethyl-pyridin-2-yl)-urea,
- 30 1-[6-[1-(2-fluoro-ethyl)-piperidin-4-yl]-quinolin-4-yl]-3-(6-trifluoromethyl-pyridin-2-yl)-urea,
- 1-[6-[1-(2,2-difluoro-ethyl)-piperidin-4-yl]-quinolin-4-yl]-3-(6-trifluoromethyl-pyridin-2-yl)-urea,
- 1-[6-(3,6-dihydro-2H-pyran-4-yl)-quinolin-4-yl]-3-(6-trifluoromethyl-pyridin-2-yl)-urea,
- 35 1-[6-(3,6-dihydro-2H-pyran-4-yl)-quinolin-4-yl]-3-pyrazin-2-yl-urea,
- tert-butyl 4-[4-(3-pyrazin-2-ylureido)quinolin-6-yl]-5,6-dihydropyridine-1(2H)-carboxylate,
- 1-pyrazin-2-yl-3-[6-(1,2,3,6-tetrahydro-pyridin-4-yl)-quinolin-4-yl]-urea,

- 1-(6-piperidin-4-yl-quinolin-4-yl)-3-pyrazin-2-yl-urea,
 1-{6-[1-(2-fluoro-ethyl)-piperidin-4-yl]-quinolin-4-yl}-3-pyrazin-2-yl-urea,
 1-[6-(1-methyl-piperidin-4-yl)-quinolin-4-yl]-3-pyrazin-2-yl-urea,
 tert-butyl 4-{4-[3-(6-cyclopropyl-pyrazin-2-yl)-ureido]-quinolin-6-yl}-3,6-dihydro-
 5 2H-pyridine-1-carboxylate,
 tert-butyl 4-{4-[3-(6-cyclopropyl-pyrazin-2-yl)-ureido]-quinolin-6-yl}-piperidine-1-carboxylate,
 1-(6-cyclopropyl-pyrazin-2-yl)-3-(6-piperidin-4-yl-quinolin-4-yl)-urea,
 1-(6-cyclopropyl-pyrazin-2-yl)-3-[6-(1-methyl-piperidin-4-yl)-quinolin-4-yl]-urea,
 10 tert-butyl 3-{4-[3-(6-Trifluoromethyl-pyridin-2-yl)-ureido]-quinolin-6-yl}-2,5-dihydro-pyrrole-1-carboxylate,
 1-[6-(2,5-dihydro-1H-pyrrol-3-yl)-quinolin-4-yl]-3-(6-trifluoromethyl-pyridin-2-yl)-urea,
 1-[6-(3,6-dihydro-2H-pyran-4-yl)-pyridin-2-yl]-3-(8-fluoro-quinolin-4-yl)-urea,
 15 1-(8-fluoro-quinolin-4-yl)-3-[6-(tetrahydro-pyran-4-yl)-pyridin-2-yl]-urea,
 1-(8-chloro-6-methyl-quinolin-4-yl)-3-[6-(tetrahydro-pyran-4-yl)-pyridin-2-yl]-urea,
 1-(6,8-dichloro-quinolin-4-yl)-3-[6-(tetrahydro-pyran-4-yl)-pyridin-2-yl]-urea,
 1-(6,8-difluoro-quinolin-4-yl)-3-[6-(tetrahydro-pyran-4-yl)-pyridin-2-yl]-urea,
 1-(8-chloro-quinolin-4-yl)-3-[6-(tetrahydro-pyran-4-yl)-pyridin-2-yl]-urea,
 20 1-[1,5]naphthyridin-4-yl-3-[6-(tetrahydro-pyran-4-yl)-pyridin-2-yl]-urea,
 1-(5,8-difluoro-quinolin-4-yl)-3-[6-(tetrahydro-pyran-4-yl)-pyridin-2-yl]-urea,
 1-(8-fluoro-6-methoxy-quinolin-4-yl)-3-[6-(tetrahydro-pyran-4-yl)-pyridin-2-yl]-urea,
 1-[6-(5,6-dihydro-4H-pyran-3-yl)-pyridin-2-yl]-3-(8-fluoro-quinolin-4-yl)-urea,
 1-(8-fluoro-quinolin-4-yl)-3-[6-(tetrahydro-pyran-3-yl)-pyridin-2-yl]-urea,
 25 1-[6-(3,6-dihydro-2H-pyran-4-yl)-pyridin-2-yl]-3-[1,5]naphthyridin-4-yl-urea,
 tert-butyl-6-[3-(8-fluoro-quinolin-4-yl)-ureido]-3',6'-dihydro-2'H-[2,4']bipyridinyl-1'-carboxylate,
 1-(8-fluoro-quinolin-4-yl)-3-(1',2',3',6'-tetrahydro-[2,4']bipyridinyl-6-yl)-urea,
 tert-butyl-6-[3-(8-fluoro-quinolin-4-yl)-ureido]-3',4',5',6'-tetrahydro-2'H-[2,4']bipyridinyl-1'-carboxylate,
 30 1-(8-fluoro-quinolin-4-yl)-3-(1',2',3',4',5',6'-hexahydro-[2,4']bipyridinyl-6-yl)-urea,
 1-(8-fluoro-quinolin-4-yl)-3-(1'-methyl-1',2',3',4',5',6'-hexahydro-[2,4']bipyridinyl-6-yl)-urea,
 1-[1'-(2-fluoro-ethyl)-1',2',3',4',5',6'-hexahydro-[2,4']bipyridinyl-6-yl]-3-(8-fluoro-quinolin-4-yl)-urea,
 35 1-[1'-(2,2-difluoro-ethyl)-1',2',3',4',5',6'-hexahydro-[2,4']bipyridinyl-6-yl]-3-(8-fluoro-quinolin-4-yl)-urea,
 1-(8-fluoro-quinolin-4-yl)-3-(1'-isopropyl-1',2',3',4',5',6'-hexahydro-[2,4']bipyridinyl-6-yl)-urea,

1-(8-fluoro-quinolin-4-yl)-3-[5-(tetrahydro-pyran-4-yl)-pyridin-2-yl]-urea,
 tert-butyl-3-[4-(3-pyrazin-2-yl-ureido)-quinolin-6-yl]-2,5-dihydro-pyrrole-1-
 carboxylate,
 tert-butyl-3-[4-(3-pyrazin-2-yl-ureido)-quinolin-6-yl]-pyrrolidine-1-carboxylate,
 1-pyrazin-2-yl-3-(6-pyrrolidin-3-yl-quinolin-4-yl)-urea,
 the stereoisomers, N-oxides, prodrugs, tautomers or physiologically tolerated
 acid addition salts thereof.

26. The heterocyclic compound according to any of the preceding claims, wherein at
 least one of the atoms has been replaced by its stable, non-radioactive isotope,
 and preferably wherein at least one hydrogen atom has been replaced by a deu-
 terium atom.
27. A pharmaceutical composition comprising at least one heterocyclic compound as
 defined in any of the preceding claims, a stereoisomer, N-oxide, prodrug,
 tautomer and/or physiologically tolerated acid addition salt thereof or comprising
 at least one heterocyclic compound as defined in any of the preceding claims
 wherein at least one of the atoms has been replaced by its stable, non-
 radioactive isotope, preferably wherein at least one hydrogen atom has been re-
 placed by a deuterium atom, and at least one physiologically acceptable carrier
 and/or auxiliary substance.
28. The heterocyclic compounds as defined in any of claims 1 to 26 or the stereoi-
 somers, N-oxides, prodrugs, tautomers or physiologically tolerated acid addition
 salts thereof for use as a medicament.
29. The heterocyclic compounds as defined in any of claims 1 to 26 or the stereoi-
 somers, N-oxides, prodrugs, tautomers or physiologically tolerated acid addition
 salts thereof for the treatment of a medical disorder susceptible to the treatment
 with a compound that modulates, preferably inhibits, the activity of glycogen syn-
 thase kinase 3 β .
30. The use of the heterocyclic compound as defined in any of claims 1 to 26 or of a
 stereoisomer, N-oxide, prodrug, tautomer or physiologically tolerated acid addi-
 tion salt thereof for the preparation of a medicament for the treatment of a medi-
 cal disorder susceptible to the treatment with a compound that modulates, pref-
 erably inhibits, the activity of glycogen synthase kinase 3 β .

31. A method for treating a medical disorder susceptible to treatment with a compound that modulates, preferably inhibits, glycogen synthase kinase 3 β activity, said method comprising administering an effective amount of at least one heterocyclic compound as defined in any of claims 1 to 26 or of a stereoisomer, prodrug, N-oxide, tautomer or physiologically tolerated acid addition salt thereof or of a pharmaceutical composition as defined in claim 27 to a subject in need thereof.
- 5
32. The heterocyclic compounds according to claim 29 or the use according to claim 30 or the method according to claim 31, where the medical disorder is a neurodegenerative disorder or an inflammatory disorder.
- 10
33. The heterocyclic compounds or the use or the method according to claim 32, where the medical disorder is selected from schizophrenia, Alzheimer's disease, behavioural and psychiatric symptoms of dementia, Parkinson's disease, tauopathies, vascular dementia, acute stroke and other traumatic injuries, cerebrovascular accidents, brain and spinal cord trauma, peripheral neuropathies, bipolar disorders, retinopathies, glaucoma, pain, rheumatoid arthritis and osteoarthritis.
- 15