

DESCRIPTION

MICROORGANISMS THAT DECOMPOSE HALOGENATED HYDROCARBONS AND
THEIR USE

5 Technical Field

The present invention relates to a biocleaning process for water or soil contaminated by halogenated hydrocarbon.

Background Art

10 Recently, organic solvents, and particularly halogenated hydrocarbons, have been used in large amounts as cleaners and so forth in advanced industries. Since growing attention is being focused on contamination of groundwater and soil caused by these substances or waste
15 water containing these substances, there is a desire to implement countermeasures against this contamination immediately.

Examples of known physical methods that have been employed in the past as countermeasures include an air
20 stripping method in which the contaminated soil is excavated and air is blown through the soil to volatilize the halogenated hydrocarbons and adsorb it with activated charcoal and so forth, and a vacuum extraction method in which the contaminated soil is pounded into a pipe after
25 which a vacuum is drawn to aerate the soil and remove the contaminants. These methods are considered to be able to be applied to decontamination of groundwater as well.

However, these methods have the disadvantage of requiring a large amount of energy, such as for blowing in
30 air. In addition, the former has the disadvantage of requiring that the soil be excavated, while the latter has the disadvantage of extraction efficiency decreasing as the concentration of contaminant decreases, thus making the cleaning difficult. Moreover, from the viewpoint of
35 preventing secondary contamination such as air pollution, these methods require that separate contaminants be

detoxified in order to adsorb onto activated charcoal and so forth.

On the other hand, research has been conducted in recent years on so-called biocleaning methods in which
5 contaminants are efficiently decomposed and detoxified by microorganisms. Since these methods utilize the decomposition mechanism of microorganisms, they do not require a large amount of energy as compared with the above-mentioned physical methods. They are also able to
10 completely decompose and detoxify contaminants without causing secondary contamination. Moreover, the cleaning can be performed even at low concentrations of contaminants, thus enabling decontamination to be performed over a wide area at the original location and
15 creating significant expectations of low costs.

Examples of methods used to purify contaminated soil by microorganisms include a solid phase treatment in which microorganisms are mixed into excavated soil with nutrient sources such as phosphorous and nitrogen to promote
20 decomposition of contaminants, a slurry treatment in which microorganisms are mixed into excavated soil with water and nutrient sources to treat the soil in the liquid state and promote decomposition of contaminants, and an original location treatment in which air, nutrient sources and so
25 forth are injected into contaminated soil without excavating to promote decomposition of contaminants by microorganisms present in the soil.

Among the above-mentioned biotreatment techniques, since soil excavation is required and the application
30 range is limited in the case of the solid phase treatment and slurry treatment method, treatment and equipment costs are relatively high.

On the other hand, the original location treatment method involves relatively low costs and allows treatment
35 over wide area. However, the cleaning rate is slow since the the absolute number of soil microorganisms is low. In the case of compounds that are difficult to decompose such

as halogenated hydrocarbons in particular, there is a possibility that microorganisms being able to decompose contaminants in the soil may not be present in the soil, thus making cleaning impossible. In this case, acquiring
5 microorganisms that are able to decompose halogenated hydrocarbons and inoculating them into the soil enables the cleaning rate to be improved and soil to be purified even though microorganisms being able to decompose the contaminants are not present in the soil.

10 A halogenated hydrocarbon contaminant, trichloroethylene (TCE), is widely used in the IC industry, in dry cleaning and so forth. It is particular important as a contaminant since it is reported to be carcinogenic. Known examples of microorganisms that
15 decompose TCE include the methane assimilating microorganisms Methyrosinus tricosporium OB3 (Japanese Unexamined Patent Publication No. 4-501667, Japanese Unexamined Patent Publication No. 5-212371) and
20 Methyrosinus tricosporium TUKUBA (Japanese Unexamined Patent Publication No. 2-92274 and Japanese Unexamined Patent Publication No. 3-292970),

Pseudomonas, such as Pseudomonas putida F1 (Japanese Unexamined Patent Publication No. 64-34499), Pseudomonas putida BH (Fujita, et al., Chemical Engineering, 39, (6),
25 p.494-498, 1994), Pseudomonas putida UC-R5 and UC-P2 (Japanese Unexamined Patent Publication No. 62-84780), Pseudomonas putida KWI-9 (Japanese Unexamined Patent Publication No. 6-70753), Pseudomonas mendocina KR-1 (Japanese Unexamined Patent Publication No. 2-503866 and
30 5-502593), Pseudomonas cepacia G4 (Japanese Unexamined Patent Publication No. 4-502277) and Pseudomonas cepacia KK01 (Japanese Unexamined Patent Publication No. 6-296711) and other microorganisms such as Alcaligenes eutropus JMP134 (A.R. Harker, Appl. Environ. Microbiol., 56, (4),
35 1179-1181, 1990), Alcaligenes eutropus KS01 (Japanese Unexamined Patent Publication No. 7-123976), and the ammonia bacteria Nitrosomonas europaea (D. Arciero, et

al., Biochem. Biophys. Res. Commun., 159, (2), 640-643, 1989) are known.

Pseudomonas cepacia KK01 in particular is reported to decompose TCE at an initial concentration of 30 ppm to a concentration of 15 ppm in liquid culture, and TCE in soil having an initial concentration of 5 ppm to a concentration of 1 ppm (Japanese Unexamined Patent Publication No. 6-296711). In addition, Alcaligenes eutropus KS01 is reported to have the ability to decompose TCE at a concentration of 50 ppm in a liquid culture to a concentration below the detection limit, and decompose TCE in the soil at 1 ppm below the detection limit (Japanese Unexamined Patent Publication No. 7-123976).

However, when testing the decomposing abilities of these microorganisms, decomposition is demonstrated at extremely high cell concentrations (1×10^8 cells/ml) in all cases. When considering that this concentration is unrealistic in the actual soil environment, the decomposing abilities of these microorganisms is not always considered to be high. Thus, in the case of using microorganisms for soil cleaning, the microorganisms have sufficient decomposing ability and are able to demonstrate that ability in the special environment of the soil, such as in the presence of wild microorganisms. In addition, it is preferable that the tolerance of the microorganisms to TCE, the target of decomposition, be high, and that they also have the ability to decompose dichloroethylene (DCE), which is a partial decomposition product of TCE.

Disclosure of Invention

The object of the present invention relates to microorganisms that efficiently decompose halogenated hydrocarbons, and particularly high concentrations of TCE, DCE and so forth, as well as a cleaning process for water or soil that uses those microorganisms.

The present invention provides microorganisms having the ability to decompose halogenated hydrocarbon and belonging to the genus Burkholderia, and some of those

microorganisms belonging to the species Burkholderia cepacia. Examples of these microorganisms include Burkholderia N16-1 (FERM BP-5504), Burkholderia cepacia N15-1 (FERM BP-5502) and Burkholderia cepacia N15-2 (FERM BP-5503). Moreover, the present invention provides a process for cleaning water or soil where the above-mentioned microorganisms are added to water or soil containing halogenated hydrocarbon.

The present invention is also characterized by the addition of microorganism activator in combination with the above-mentioned microorganisms. These microorganisms have the ability to decompose 50% or more of 100 ppm of trichloroethylene in 2 days, or to decompose 100% of 30 ppm of trichloroethylene in 18 hours.

BRIEF DESCRIPTION OF THE DRAWINGS

Fig. 1 is a graph indicating the change over time in the percentage of remaining TCE in Example 2.

Fig. 2 is a graph indicating the change over time in the percentage of remaining TCE in Example 3.

Fig. 3 is a graph indicating the change over time in the percentage of remaining cis-1,2-DCE in Example 4.

Fig. 4 is a graph indicating the percentage of remaining TCE in Example 6.

Fig. 5 is a graph indicating the percentage of remaining TCE in Example 7.

DETAILED DESCRIPTION

The microorganisms of the present invention should be microorganisms belonging to the genus Burkholderia or microorganisms belonging to the species Burkholderia cepacia, specific examples of these microorganisms include Burkholderia N16-1, Burkholderia cepacia N15-1 and Burkholderia cepacia N15-2. These microorganisms are new strains isolated from nature such as rivers and soil, and their isolation method and taxonomical characteristics are specifically described. These strains, Burkholderia N16-1, Burkholderia cepacia N15-1 and Burkholderia cepacia N15-2 were deposited on April 12, 1996 with the National

Institute of Bioscience and Human Technology Agency of Industrial Science and Technology as FERM BP-5504, FERM BP-5502 and FERM BP-5503, respectively.

5 The microorganisms of the present invention can be cultured in the presence of routinely used carbon sources and nitrogen sources in a medium containing inorganic salt, vitamins and other trace elements as necessary. Any carbon source can be used provided it is a carbon source that is preferentially assimilated by the microorganisms
10 of the present invention. Although varying according to the type of carbon source, the concentration of carbon source in the medium is preferably, for example, 0.1 to 0.5 g/L. Examples of nitrogen sources that can be used include organic nitrogen sources such as yeast extract,
15 peptone and meat extract, while examples of inorganic nitrogen sources include ammonium salts and nitrates.

Although varying according to the particular type, the concentration of the nitrogen source is preferably 0.1 to 1.4 g/L. Preferable examples of inorganic salts
20 include those composed of metal ions such as potassium, calcium, magnesium, iron (II), manganese, cobalt and nickel ions, and anions such as chloride, sulfate and phosphate ions. Culturing is preferably performed aerobically, and aeration and stirring are preferable in
25 the case of shake culturing or large-scale culturing. The culture temperature is 20 to 37°C and preferably around 30°C.

In addition, the present invention relates to a process for cleaning water or soil characterized by adding
30 the above-mentioned microorganisms to water or soil containing halogenated hydrocarbon. In this process, the microorganisms of the present invention cultured in the manner described above should be added to the water or soil to be treated. The microorganisms may be added in
35 the form of a culture liquid or added in the form of microorganisms after isolating from a culture liquid.

Moreover, the microorganisms may also be added after adsorbing onto a separate carrier.

Although the amount of microorganisms added varies according to the halogenated hydrocarbon decomposition ability of the microorganisms, the amount of halogenated hydrocarbon in the water or soil to be treated and so forth, it is within the range of 10^5 to 10^9 cells/g. Although the time required for treatment also varies according to the halogenated hydrocarbon decomposition ability of the microorganisms used, the amount of halogenated hydrocarbon in the water or soil to be treated and the amount of microorganisms added, it is approximately 1 to 10 days.

In addition, the cleaning process for water or soil of the present invention relates to a process for cleaning water or soil in which the microorganisms of the present invention as described above are inoculated into water or soil contaminated by halogenated hydrocarbon and mixed to decompose the halogenated hydrocarbon contained in the water or soil, and is characterized by the addition and mixing of at least one type of microorganism activator during inoculation and mixing of the above-mentioned microorganisms in the above-mentioned water or soil. In this case, the microorganism activator has the role of activating the halogenated hydrocarbon decomposition ability of the microorganisms, and is referred to as an inducer. Examples of inducers that can be used include compounds that can be assimilated and decomposed by the above-mentioned microorganisms, preferable examples of which include benzene, toluene, phenol, cresol and 3-hydroxybenzyl alcohol. In addition, cyclopentanol, hexanoic acid, trans-3-hexenoic acid and suberinic acid can be used for N16-1 in addition to the inducers indicated above.

Examples of halogenated hydrocarbons that can be decomposed by the process of the present invention are

particularly chlorinated hydrocarbons such as TCE, DCE and monochloroethylene.

Although the process of the present invention can be applied to the solid phase treatment method and slurry treatment method previously described, it is not always necessary to use these methods that require excavation of the soil, but rather cleaning can be performed simply by adding and inoculating the microorganisms of the present invention into the soil or water.

10 **Examples**

The following provides a detailed explanation of the present invention through Examples.

Example 1. Isolation and Identification of
Microorganisms

15 Microorganisms of the present invention were screened and isolated from the soil on the grounds of a chemical plant using the method described below. 0.1 g of sampled soil were inoculated into 5 ml of NMS medium or M9 medium in a 25 ml screw-top test tube. Moreover, 500 ppm of
20 phenol and vitamin cocktail was added, after which the tube was capped and cultured with shaking for a prescribed time at 30°C. Culturing was continued until culture medium became cloudy due to microorganism growth in the medium. After completion of 10 rounds of subculturing,
25 the culture liquid was suitably diluted and swabbed onto plate medium to which 1.5% agar was added to isolate the microorganism colonies that appeared. The microorganisms were isolated by repeating this procedure.

Table 1

<u>Composition of NMS Medium (in 1 Liter)</u>		
5	MgSO ₄ ·7H ₂ O	1.0 g
	CaCl ₂	0.01 g
	Na ₂ HPO ₄ ·12H ₂ O	0.717 g
	NH ₄ Cl	0.6 g
	KH ₂ PO ₄	0.272 g
10	Trace element solution (pH 6.8)	0.5 ml
	<u>Trace Element Solution (in 1 Liter)</u>	
15	EDTA	500 mg
	FeSO ₄ ·7H ₂ O	200 mg
	ZnSO ₄ ·7H ₂ O	10 mg
	MnCl ₂ ·4H ₂ O	3 mg
	H ₃ BO ₃	30 mg
20	CoCl ₂ ·6H ₂ O	20 mg
	NiCl ₂ ·6H ₂ O	2 mg
	CaCl ₂	1 mg
	Na ₂ MoO ₄ ·2H ₂ O	2 mg
25	<u>Vitamin Cocktail (in 1 Liter)</u>	
	Thiamine hydrochloride	3 mg
	p-aminobenzoic acid	13 mg
	Adenine	1.0 g
	NAD	0.25 g
30	Vitamin B ₁₂	10 mg
	Thiamine diphosphochloride	100 mg

After culturing the isolated microorganisms in the liquid medium for 2 days, 1/100th volume of the culture liquid was inoculated into a 30 ml vial containing 4 ml of NMS medium to which phenol, 0.02% yeast extract and 1 mM glucose, had been added, followed by the addition of 10 ppm of TCE. After promptly sealing the vial with a Teflon-coated septum cap and aluminum cap and shake culturing for 5 days at 30°C, the vapor phase in the vial was analyzed by gas chromatography. The morphological and physiological properties of the three strains of microorganisms having a high degree of TCE decomposition activity that were selected in this manner were investigated. Those results are shown in Table 2.

Table 2

Test Parameter	Test Results		
	N15-1	N15-2	N16-1
Morphology	Rod	Rod	Rod
Gram staining	-	-	-
Spores	-	-	-
Motility	+	+	+
Flagella	Polar, multi- flagellate	Polar, multi- flagellate	Polar, multi- flagellate
Response to oxygen	Aerobic	Aerobic	Aerobic
Oxidase	+	+	+
Catalase	+	+	+
OF	O	O	O
Colony color tone	NP	NP	NP
Fluorescent pigment formation	-	-	-
Water-soluble pigment formation	-	-	-
PHB accumulation	+	+	+
Cleavage of protocatechinic acid	Ortho form	Ortho form	Ortho form
Arginine dihydrolase	-	-	-
Growth at 40°C	+	+	-
Denitrification reaction	-	-	-
Nitrate reduction	+	+	-
Gelatin liquefaction	+	+	-
Starch decomposition	-	-	-
Assimilation			
Glucose	+	+	+
Xylose	+	+	+
Rhamnose	-	-	+
Levulinic acid	+	+	-
Mesaconic acid	-	-	-
D-tartaric acid	-	-	+
2,3-butylene glycol	+	+	-
Tryptamine	-	-	-
Quinone type	Q-8	Q-8	Q-8
GC content of intracellular DNA (mol%)	66	67	62

- As a result of identifying the above-mentioned
- 5 microorganisms from the above results according to the literature (N.R. Krieg and J.G. Holt, "Bergey's Manual of Systematic Bacteriology", Vol. 1 (1984) Williams & Wilkins; J.G. Holt, N.R. Krieg, P.H.A. Sneath, J.T. Staley and S.T. Williams, "Bergey's Manual of Determination
- 10 Bacteriology", Ninth Edition (1994) Williams & Wilkins; N. Zhao, C. Qu, E. Wang and W. Chen, Int. J. Syst. Bacteriol., 45, 600 (1995); E. Yabuuti, Y. Kosako, H. Oyaizu, I. Yano, H. Hotta, Y. Hashimoto, T. Ezaki and M. Arakawa, Microbiol. Immunol., 36, 1251 (1992)), one strain
- 15 was identified as Burkholderia species, while the other

two strains were identified as Burkholderia cepacia, and were named N16-1, N15-1 and N15-2, respectively.

Known microorganisms that decompose TCE belonging to the species Burkholderia cepacia are G4 (Japanese
5 Unexamined Patent Publication No. 4-502277) and KK01 (Japanese Unexamined Patent Publication No. 6-296711) as previously described. Since both N15-1 and N15-2 have motility as shown in Table 4, they are clearly different from G4 which is not motile. In addition, although G4 and
10 KK01 are induced by toluene, since N15-1 and N15-2 are not induced by toluene, they are clearly different based on this parameter as well. On the other hand, since N16-1 has an intracellular DNA GC content of 62 mol% and does not fall under Burkholderia putida, mendocina or cepacia
15 that are known to be microorganisms that decompose TCE (previously classified as a Pseudomonas species prior to 1992), it was certified as a novel microorganism.

Example 2. Decomposition of TCE in Liquid Medium

Strains N15-1, N15-2 and N16-1 were cultured for 1
20 day each in NMS liquid medium to which 500 ppm of phenol and vitamin cocktail had been added. After collecting the microorganisms by centrifugation, the microorganisms were resuspended in 4 ml of the same medium not containing phenol. 30 ml of this suspension were transferred to
25 vials followed by the addition of 100 ppm of TCE after which the vials were promptly sealed with a Teflon-coated silicon septum and aluminum cap (microorganism concentration: 10^8 cells/ml). The vials were then
30 cultured by allowing to stand undisturbed at 30°C. The vapor phase was periodically analyzed by gas chromatography. Those results are shown in Fig. 1. More than 70% of the TCE in the culture liquid was decomposed in 5 days by each of the microorganisms.

Example 3. Decomposition of TCE in Liquid Medium

35 Strains N15-1, N15-2 and N16-1 were cultured for 1 day each in NMS liquid medium to which 0.2% yeast extract and 4 mM glucose had been added. 4 ml of NMS liquid

medium to which 0.02% yeast extract, 500 ppm of phenol and 1 mM glucose had been added was placed in vials and inoculated with 40 μ l each of the above-mentioned culture liquid (microorganism count of roughly 10^6 cells/ml). 100 ppm of TCE were then added followed by promptly sealing the vials with a Teflon-coated silicon septum and aluminum cap. The vials were incubated with shaking at 30°C, and the vapor phase was periodically analyzed by gas chromatography. Those results are shown in Fig. 2.

More than 60% of the TCE in the culture liquid was decomposed by each of the microorganisms in 10 days. When this decomposing ability was compared with known TCE-decomposing microorganisms, there were only two reports describing TCE in excess of 30 ppm. Of those, Pseudomonas cepacia KK01 (Japanese Unexamined Patent Publication No. 6-296711) was reported to have the ability to decompose 30 ppm of TCE down to roughly 15 ppm (50%) in 2 days, while Alcaligenes eutropus KS01 was reported to be able to decompose 50 ppm of TCE to a level below the detection limit of gas chromatography in 4 days at a microorganism concentration of 10^8 cells/ml. In comparison, since N15-1, N15-2 and N16-1 are able to decompose TCE at a concentration as high as 100 ppm while also having a high decomposing ability per cell, in addition to being able to accommodate a broader range of contamination concentrations at the time of actual use, and since the amount of microorganisms required can be drastically reduced, they also offer the advantage of being able to decrease costs.

Example 4. Decomposition of DCE in Liquid Medium

Strains N15-1, N15-2 and N16-1 were each inoculated into 5 ml of NMS liquid medium to which 0.02% yeast extract, 500 ppm phenol and 1 mM glucose had been added. After culturing for 2 days at 30°C, 1/100th volume of the culture liquid was added to a vial containing 4 ml of the same culture liquid containing 30 ppm cis-1,2-DCE, followed by culturing for 5 days at 30°C. The results are

shown in Fig. 3. Decomposition of more than 99% of the DCE was observed for all three strains.

Example 5. Effect of Temperature during Decomposition

5 Strains N15-1, N15-2 and N16-1 were each inoculated into 5 ml of NMS liquid medium to which 0.02% yeast extract, 500 ppm phenol and 1 mM glucose had been added. After culturing for 2 days at 30°C, 1/100th volume of the culture liquid was added to vials containing 4 ml of the
10 same culture liquid to which had been added 30 ppm of TCE, followed by culturing for 8 days at 16 to 30°C. In the cases of N16-1 and N15-1, TCE was decomposed below the detection limit at each temperature. In the case of N15-2, however, although approximately 30% of the TCE remained
15 at 16°C, the amount was below the detection limit at 20°C and above. Accordingly, it was shown that these microorganisms are able to decompose TCE at the temperature (15 to 20°C) of the soil.

Example 6. Effect of pH During Decomposition

20 Strains N15-1, N15-2 and N16-1 were each inoculated into 5 ml of M9 liquid medium (pH 7.0) to which 0.02% yeast extract, 500 ppm phenol and 1 mM glucose had been added. After culturing for 2 days at 30°C, 1/100th volume of the culture liquid was added to vials containing 4 ml
25 of culture medium adjusted to a pH of 5-10, and to which had been added 30 ppm of TCE instead of phenol, followed by culturing for 5 days at 30°C. N15-1, N15-2 and N16-1 each decomposed 30 ppm TCE to a level below the detection limit at all pH levels. In addition, although the growth
30 of N16-1 was inhibited at pH level of 7.4 and above, it decomposed TCE to a level below the detection limit in the pH range of 5 to 7.

Example 7. Decomposition Test of TCE in Soil

35 10 g of Andsols (sampled from Aichi prefecture and air dried) were placed in a vial having a volume of 30 ml followed by the addition of TCE to a concentration of 20 ppm. Strains N15-1, N15-2 and N16-1 were each inoculated

into 20 ml of NMS liquid medium to which 0.02% yeast extract, 500 ppm phenol and 1 mM glucose had been added. After shake-culturing for 3 days at 30°C and collecting the microbial cells from the culture liquid by
5 centrifugation, the cells were resuspended in adequate volume of NMS medium not containing phenol, and the amount of inoculated cells in the above-mentioned vials was 10^8 to 10^9 cells/g and the moisture content after addition of the suspension was 25%. After capping the vials with
10 screw-on caps wrapped with Teflon-coated packings and shaking, the vials were incubated for 7 days at 30°C.

10 g of the soil were weighed in an Erlenmeyer flask provided with a stopper followed by the addition of 90 ml of ion exchange water aerated with air passed through
15 activated charcoal, 5 ml of phosphoric acid and 10 ml of n-hexane. After sealing the flask, ultrasonic treatment was performed for 20 minutes in an ultrasonic cleaner followed by shaking for 5 minutes with a shaker. Next, the aqueous phase and n-hexane phase were transferred to a
20 colorimetric tube provided with a stopper. After sealing the colorimetric tube and performing ultrasonic treatment, the separated n-hexane was analyzed by gas chromatography. Those results are shown in Fig. 4.

30 to 50% of the TCE at an initial concentration of
25 10 ppm was decomposed in 7 days. Thus, the microorganisms of the present invention demonstrated decomposing ability even in a natural environment. Accordingly, it is possible clean contaminated soil by bringing the soil in contact with the microorganisms of the present invention
30 and water. In addition, this technique enables soil to be cleaned without adding activator compounds such as phenol directly to the soil.

Example 8. Decomposition Test of TCE in Soil

10 g of Andsols (sampled from Aichi prefecture and
35 air dried) were placed in a vial having a volume of 30 ml followed by the addition of TCE to a concentration of 20 ppm. Strains N15-1, N15-2 and N16-1 were each inoculated

into 20 ml of NMS liquid medium to which 0.02% yeast extract, 500 ppm phenol and 1 mM glucose had been added, followed by shake-culturing for 1 day at 30°C.

0.2 ml of this liquid was removed and added to the
5 above-mentioned vial so that the amount of inoculated microbial cells was 10^6 cells/g, the moisture content following addition of medium was 40% and phenol concentration was 500 ppm. After capping the vial with a screw-on cap wrapped with a Teflon-coated packing and
10 shaking, the culture liquid was cultured for 3 weeks at 30°C. Those results are shown in Fig. 5. 40 to 50% of the TCE was decomposed, thus indicating that the microorganisms of the present invention are able to decompose TCE in soil taken from a natural environment
15 even at a low concentration of 10^6 cells/g.

Example 9. Induction of Decomposition Activity by an Activator

Strains N15-1, N15-2 and N16-1 were each cultured for 2 days at 30°C in 5 ml of NMS liquid medium in which 0.2%
20 yeast extract and 5 mM glucose had been added. 4 ml of NMS liquid medium to which 0.02% yeast extract, 1 mM glucose and one of the activators listed in Table 4 at a concentration of 100 ppm had been added were placed in vials that were then inoculated with 40 μ l of each of the
25 above-mentioned culture liquid (microorganism count of approximately 10^6 cells/g). 30 ppm TCE was then added to each vial followed by promptly sealing the vials with a Teflon-coated silicon septum and an aluminum cap. The vials were cultured with shaking for 5 days at 30°C and
30 the vapor phase was analyzed by gas chromatography. The result is shown in Table 5. The numbers shown indicate the concentration of remaining TCE during addition of the microorganisms as a percentage when the concentration of TCE when microorganisms are not added is taken to be 100%.
35 Those results are shown in the following Table 3.

Table 3

Compound	Microorganism Strain		
	N16-1	N15-1	N15-2
Cyclohexanol	8.6	57.4	64.1
Cyclopentanol	49.3	100	100
Anthranilic acid	13.3	105	107
Caffeinic acid	52.6	96.7	102
Suberinic acid	7.3	99.8	100
Maleic acid	53.0	100	98.8
Fumaric acid	63.4	100	100
Succinic acid	63.3	72.4	92.4
Malonic acid	45.2	100	100
Trans-3-hexenoic acid	2.7	100	100
Hexanoic acid	8.3	100	100
Benzene	0	98.8	97.3
Ethylbenzene	28.7	98.4	101
Benzyl alcohol	0	71.9	72.5
Saligenin	0	102	105
Allylphenol	0.5	100	102
Guaiacol	35.0	99.2	107
Toluene	0	100	100
Benzaldehyde	6.8	103	99.0
p-hydroxybenzoic acid	12.5	102	98.8

5 N15-1 and N15-2 were activated by aromatic compounds
 such as phenol, toluene and benzene, which are known to be
 conventional activation inducers, as well as by the non-
 aromatic compound, cyclohexanol. In addition, succinic
 acid also exhibited the ability to activate these
 microorganisms. Moreover, N16-1 was also activated by
 10 cyclopentanol, anthranilic acid, p-hydroxybenzoic acid,
 suberinic acid, trans-3-hexenoic acid and straight chain
 carboxylic acids such as hexanoic acid.

Furthermore, a comparison of the properties of
 strains N15-1, N15-2 and N16-1 of the present invention
 15 with those of known Burkholderia cepacia strains KK01 and
 G4 is shown in the following Table 4.

Table 4

Property	Burkholderia cepacia			Burkholderia genus
	KK01	G4	N15-1,2	N16-1
Decomposing ability	30->15/2 days, organism count unknown	1->0.17/1 day, 3×10^8 cells/ml	100->40/2 days, 10^8 cells/ml	100->45/day, 10^8 cells/ml
Max. TCE conc. (ppm)	30	2	100	100
Activators	Toluene, phenol, cresol	Toluene, phenol, cresol	Not toluene, phenol, cresol, cyclohexanol	Toluene, cresol, cyclohexanol suberinic acid, others
Mobility	+	-	+	+

According to the present invention, the microorganisms of the present invention are able to decompose high concentrations of halogenated hydrocarbons such as TCE and DCE contained in water or soil in the presence of preferably at least one type of activator and sugar or other nutrient.

According to the process for purifying water or soil in the present invention, in addition to offering the advantages of biotechnology such as not requiring a large amount of energy and inhibiting the occurrence of secondary contamination, the process of the present invention is able to industrially and efficiently purify water or soil contaminated by halogenated hydrocarbons in the natural environment.

Example 10. Decomposition of trichloroethylene in liquid culture

The strains N15-1, N15-2 and N16-1 were separately cultured in a liquid NMS medium supplemented with 0.2% yeast extract and 5 mM glucose for one day. 4 ml of liquid NMS medium supplemented with 0.02% yeast extract, 100 ppm phenol and 1 mM glucose was put into each vial, and 40 μ l of the culture as prepared above was inoculated into the vials (about 10^6 cells/ml medium). Trichloroethylene was added to vials to make a concentration of 30 ppm, and the vials were rapidly

- 18 -

shielded with a Teflon-coated silicone plug and aluminum cap. The vials were incubated at 30°C with shaking, and the gas phase in the vials was periodically analyzed by gas chromatography. As a result, 100% trichloroethylene
5 was decomposed in 18 hours.

CLAIMS

1. Microorganism belonging to the genus Burkholderia and having an ability to decompose halogenated hydrocarbon, which are able to decompose 50%
5 or more of 100 ppm of trichloroethylene in 2 days, or to decompose 100% of 30 ppm of trichloroethylene in 18 hours.
2. Microorganism according to claim 1 wherein said halogenated hydrocarbon is trichloroethylene.
3. Microorganism belonging to the species
10 Burkholderia cepacia and having the ability to decompose halogenated hydrocarbon.
4. Microorganism according to claim 3 wherein said halogenated hydrocarbon is trichloroethylene.
5. Microorganism according to claim 1 or 2 that are
15 Burkholderia N16-1 (FERM BP-5504).
6. Microorganism according to claim 3 or 4 that are Burkholderia cepacia N15-1 (FERM BP-5502).
7. Microorganism according to claim 3 or 4 that are Burkholderia cepacia N15-2 (FERM BP-5503).
- 20 8. A process for cleaning of water or soil containing halogenated hydrocarbon comprising the steps of one or all strains of the microorganisms according to any of claims 1 through 7 is added to water or soil containing halogenated hydrocarbon.
- 25 9. A process according to claim 8 wherein microorganism activator is added to the water or soil with said added microorganisms.


THIS SHEET HAS BEEN REPLACED

This Sheet Has Been Replaced




1/5

Fig.1




2/5

Fig.2




3/5

Fig.3




$\frac{4}{5}$

Fig.4



$\frac{5}{5}$


Fig.5





CLAIMS

1. Microorganism belonging to the genus Burkholderia and having an ability to decompose halogenated hydrocarbon, which are able to decompose 50%
5 or more of 100 ppm of trichloroethylene in 2 days, or to decompose 100% of 30 ppm of trichloroethylene in 18 hours.
2. Microorganism according to claim 1 wherein said halogenated hydrocarbon is trichloroethylene.
3. Microorganism according to claim 1, wherein the genus
10 Burkholderia is the species Burkholderia ceracia having the ability to decompose halogenated hydrocarbon.
4. Microorganism according to claim 3 wherein said halogenated hydrocarbon is trichloroethylene.
5. Microorganism according to claim 1 or 2 that are
15 Burkholderia N16-1 (FERM BP-5504).
6. Microorganism according to claim 3 or 4 that are Burkholderia cepacia N15-1 (FERM BP-5502).
7. Microorganism according to claim 3 or 4 that are Burkholderia cepacia N15-2 (FERM BP-5503).
- 20 8. A process for cleaning of water or soil containing halogenated hydrocarbon comprising the steps of one or all strains of the microorganisms according to any of claims 1 through 7 is added to water or soil containing halogenated hydrocarbon.
- 25 9. A process according to claim 8 wherein microorganism activator is added to the water or soil with said added microorganisms.



PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12N 1/20, C02F 3/34, A62D 3/00, B09C 1/10 // (C12N 1/20, C12R 1:01)		A1	(11) International Publication Number: WO 97/40136
			(43) International Publication Date: 30 October 1997 (30.10.97)
(21) International Application Number: PCT/JP97/01359		(81) Designated States: CA, CN, KR, <u>SG</u> , US, VN, European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).	
(22) International Filing Date: 18 April 1997 (18.04.97)			
(30) Priority Data: ① 8/100466 ✓ 22 April 1996 (22.04.96) JP		Published With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.	
(71) Applicant (for all designated States except US): TOYOTA JIDOSHA KABUSHIKI KAISHA [JP/JP]; 1, Toyota-cho, Toyota-shi, Aichi 471-71 (JP).			
(72) Inventors; and			
(75) Inventors/Applicants (for US only): NAKAYAMA, Mika [JP/JP]; 5-1215-316, Sakashita-cho, Kasugai-shi, Aichi 480-03 (JP); MIYAZAKI, Chie [JP/JP]; 973-223, Minamiyama, Komenogi-cho, Nisshin-shi, Aichi 470-01 (JP); SASAMI, Osamu [JP/JP]; 3-1, Aza Togo, Oaza Higashino, Konan-shi, Aichi 483 (JP); YAMADA, Yukio [JP/JP]; 2-65, Teramae-cho, Tsushima-shi, Aichi 496 (JP); NUMATA, Koichi [JP/JP]; 1-98, Horagai, Midori-ku, Nagoya-shi, Aichi 458 (JP); ODA, Yasushi [JP/JP]; 522-4217, Toyota-cho, Toyota-shi, Aichi 471 (JP).			
(74) Agents: ISHIDA, Takashi et al.; A. Aoki & Associates, Toranomon 37 Mori Building, 5-1, Toranomon 3-chome, Minato-ku, Tokyo 105 (JP).			
(54) Title: MICROORGANISMS THAT DECOMPOSE HALOGENATED HYDROCARBONS AND THEIR USE ✓			
(57) Abstract			
Microorganisms belonging to the genus Burkholderia and having the ability to decompose halogenated hydrocarbon, which are able to decompose 50 % or more of 100 ppm of trichloroethylene in 2 days, or decompose 100 % of 30 ppm of trichloroethylene in 18 hours, as well as providing a process for decomposing halogenated hydrocarbons in water or soil using those microorganisms.			