

QUINOLINE DERIVATIVES AS ANTIBACTERIAL AGENTS

5 The present invention relates to the use of quinoline derivatives for the manufacture of a medicament for the treatment of a bacterial infection.

Resistance to first-line antibiotic agents is an emerging problem. Some important examples include penicillin-resistant *Streptococcus pneumoniae*, vancomycin-resistant enterococci, methicillin-resistant *Staphylococcus aureus*, multi-resistant salmonellae.

10

The consequences of resistance to antibiotic agents are severe. Infections caused by resistant microbes fail to respond to treatment, resulting in prolonged illness and greater risk of death. Treatment failures also lead to longer periods of infectivity, which increase the numbers of infected people moving in the community and thus exposing the general population to the risk of contracting a resistant strain infection.

15

Hospitals are a critical component of the antimicrobial resistance problem worldwide. The combination of highly susceptible patients, intensive and prolonged antimicrobial use, and cross-infection has resulted in infections with highly resistant bacterial pathogens.

20

Self-medication with antimicrobials is another major factor contributing to resistance. Self-medicated antimicrobials may be unnecessary, are often inadequately dosed, or may not contain adequate amounts of active drug.

25

Patient compliance with recommended treatment is another major problem. Patients forget to take medication, interrupt their treatment when they begin to feel better, or may be unable to afford a full course, thereby creating an ideal environment for microbes to adapt rather than be killed.

30

Because of the emerging resistance to multiple antibiotics, physicians are confronted with infections for which there is no effective therapy. The morbidity, mortality, and financial costs of such infections impose an increasing burden for health care systems worldwide.

35



Therefore, there is a high need for new compounds to treat bacterial infections, especially for the treatment of infections caused by resistant strains.

-2-

WO 2004/011436 discloses substituted quinoline derivatives having activity against Mycobacteria, in particular against *Mycobacterium tuberculosis*. One particular compound of these substituted quinoline derivatives is described in Science (2005), 307, 223-227.

- 5 It has now been found that quinoline derivatives described in WO 2004/011436 also show activity against other bacteria than Mycobacteria.

Therefore, the present invention relates to the use of a compound for the manufacture of a medicament for the treatment of a bacterial infection, said compound being a
10 compound of formula (Ia) or (Ib)




a *N*-oxide thereof, a tautomeric form thereof or a stereochemically isomeric form thereof wherein

A^- is a pharmaceutically acceptable counter ion;

15 R^1 is hydrogen, halo, haloalkyl, cyano, hydroxy, Ar, Het, alkyl, alkyloxy, alkylthio, alkyloxyalkyl, alkylthioalkyl, Ar-alkyl or di(Ar)alkyl ;

p is an integer equal to 1, 2, 3 or 4 ;

-3-

- R^2 is hydrogen, hydroxy, mercapto, alkyloxy, alkyloxyalkyloxy, alkylthio, mono or di(alkyl)amino or a radical of formula  wherein Y is CH_2 , O, S, NH or N-alkyl ;
- R^3 is alkyl, Ar, Ar-alkyl, Het or Het-alkyl;
- 5 q is an integer equal to zero, 1, 2, 3 or 4 ;
- R^4 and R^5 each independently are hydrogen, alkyl or benzyl;
- R^4 and R^5 together and including the N to which they are attached may form a radical selected from the group of pyrrolidinyl, 2-pyrrolinyl, 3-pyrrolinyl, pyrrolyl, imidazolidinyl, pyrazolidinyl, 2-imidazolinyl, 2-pyrazolinyl, 10 imidazolyl, pyrazolyl, triazolyl, piperidinyl, pyridinyl, piperazinyl, pyridazinyl, pyrimidinyl, pyrazinyl, triazinyl, morpholinyl and thiomorpholinyl, each of said rings may optionally be substituted with alkyl, halo, haloalkyl, hydroxy, alkyloxy, amino, mono- or dialkylamino, alkylthio, alkyloxyalkyl, alkylthioalkyl, Ar-alkyl or pyrimidinyl;
- 15 R^6 is hydrogen, halo, haloalkyl, hydroxy, Ar, alkyl, alkyloxy, alkylthio, alkyloxyalkyl, alkylthioalkyl, Ar-alkyl or di(Ar)alkyl ; or two vicinal R^6 radicals may be taken together to form a bivalent radical of formula $-CH=CH-CH=CH-$;
- r is an integer equal to 1, 2, 3, 4 or 5 ;
- 20 R^7 is hydrogen, alkyl, Ar or Het ;
- R^8 is hydrogen or alkyl ;
- R^9 is oxo ; or
- R^8 and R^9 together form the radical $-CH=CH-N=$;
- R^{10} is alkyl, alkylcarbonyl, Ar, Ar-alkyl, Ar-carbonyl, Het¹-alkyl or 25 Het¹-carbonyl;
- alkyl is a straight or branched saturated hydrocarbon radical having from 1 to 6 carbon atoms ; or is a cyclic saturated hydrocarbon radical having from 3 to 6 carbon atoms ; or is a cyclic saturated hydrocarbon radical having from 3 to 6 carbon atoms attached to a straight or branched saturated hydrocarbon radical having from 1 to 6 carbon atoms ; wherein each carbon atom can be optionally 30 substituted with hydroxy, alkyloxy or oxo ;
- Ar is a homocycle selected from the group of phenyl, naphthyl, acenaphthyl, tetrahydronaphthyl, each homocycle optionally substituted with 1, 2 or 3 substituents, each substituent independently selected from the group of hydroxy, 35 halo, cyano, nitro, amino, mono- or dialkylamino, alkyl, haloalkyl, alkyloxy,

haloalkyloxy, carboxyl, alkyloxycarbonyl, aminocarbonyl, morpholinyl and mono- or dialkylaminocarbonyl ;

Het is a monocyclic heterocycle selected from the group of *N*-phenoxypiperidinyl, piperidinyl, pyrrolyl, pyrazolyl, imidazolyl, furanyl, thienyl, oxazolyl,

5 isoxazolyl, thiazolyl, isothiazolyl, pyridinyl, pyrimidinyl, pyrazinyl and pyridazinyl; or a bicyclic heterocycle selected from the group of quinolinyl, quinoxalinyl, indolyl, benzimidazolyl, benzoxazolyl, benzisoxazolyl, benzothiazolyl, benzisothiazolyl, benzofuranyl, benzothienyl, 2,3-dihydrobenzo[1,4]dioxinyl and benzo[1,3]dioxolyl ; each monocyclic and
10 bicyclic heterocycle may optionally be substituted with 1, 2 or 3 substituents, each substituent independently selected from the group of halo, hydroxy, alkyl, alkyloxy, and Ar-carbonyl;

Het¹ is a monocyclic heterocycle selected from furanyl or thienyl; or a bicyclic heterocycle selected from benzofuranyl or benzothienyl; each monocyclic and
15 bicyclic heterocycle may optionally be substituted with 1, 2 or 3 substituents, each substituent independently selected from the group of halo, alkyl and Ar;

halo is a substituent selected from the group of fluoro, chloro, bromo and iodo; and



haloalkyl is a straight or branched saturated hydrocarbon radical having from 1 to
20 6 carbon atoms or a cyclic saturated hydrocarbon radical having from 3 to 6 carbon atoms or a cyclic saturated hydrocarbon radical having from 3 to 6 carbon atoms attached to a straight or branched saturated hydrocarbon radical having from 1 to 6 carbon atoms; wherein one or more carbon atoms are substituted with one or more halo atoms;

provided that the bacterial infection is other than a Mycobacterial infection.

25 The present invention also concerns a method of treating a bacterial infection in a mammal, in particular a warm-blooded mammal, more in particular a human, comprising administering an effective amount of a compound of the invention to the mammal.

30 The present invention also concerns a compound of formula (Ia) or (Ib)

-5-




a *N*-oxide thereof, a tautomeric form thereof or a stereochemically isomeric form thereof wherein

A^- is a pharmaceutically acceptable counter ion;

R^1 is hydrogen, halo, haloalkyl, cyano, hydroxy, Ar, Het, alkyl, alkyloxy, alkylthio, alkyloxyalkyl, alkylthioalkyl, Ar-alkyl or di(Ar)alkyl ;

p is an integer equal to 1, 2, 3 or 4 ;

R^2 is hydrogen, hydroxy, mercapto, alkyloxy, alkyloxyalkyloxy, alkylthio,

mono or di(alkyl)amino or a radical of formula  wherein Y is CH_2 , O, S, NH or *N*-alkyl ;

R^3 is alkyl, Ar, Ar-alkyl, Het or Het-alkyl;


q is an integer equal to zero, 1, 2, 3 or 4 ;

R^4 and R^5 each independently are hydrogen, alkyl or benzyl;

R^4 and R^5 together and including the N to which they are attached may form a radical selected from the group of pyrrolidinyl, 2-pyrrolinyl, 3-pyrrolinyl, pyrrolyl, imidazolidinyl, pyrazolidinyl, 2-imidazolinyl, 2-pyrazolinyl, imidazolyl, pyrazolyl, triazolyl, piperidinyl, pyridinyl, piperazinyl, pyridazinyl, pyrimidinyl, pyrazinyl, triazinyl, morpholinyl and thiomorpholinyl, each of said rings may optionally be substituted with

- alkyl, halo, haloalkyl, hydroxy, alkyloxy, amino, mono- or dialkylamino, alkylthio, alkyloxyalkyl, alkylthioalkyl, Ar-alkyl or pyrimidinyl;
- R^6 is hydrogen, halo, haloalkyl, hydroxy, Ar, alkyl, alkyloxy, alkylthio, alkyloxyalkyl, alkylthioalkyl, Ar-alkyl or di(Ar)alkyl ; or
- 5 two vicinal R^6 radicals may be taken together to form a bivalent radical of formula $-CH=CH-CH=CH-$;
- r is an integer equal to 1, 2, 3, 4 or 5 ;
- R^7 is hydrogen, alkyl, Ar or Het ;
- R^8 is hydrogen or alkyl ;
- 10 R^9 is oxo ; or
- R^8 and R^9 together form the radical $-CH=CH-N=$;
- R^{10} is alkyl, alkylcarbonyl, Ar, Ar-alkyl, Ar-carbonyl, Het¹-alkyl or Het¹-carbonyl;
- alkyl is a straight or branched saturated hydrocarbon radical having from 1 to 6
- 15 carbon atoms ; or is a cyclic saturated hydrocarbon radical having from 3 to 6 carbon atoms ; or is a cyclic saturated hydrocarbon radical having from 3 to 6 carbon atoms attached to a straight or branched saturated hydrocarbon radical having from 1 to 6 carbon atoms ; wherein each carbon atom can be optionally substituted with hydroxy, alkyloxy or oxo ;
- 20 Ar is a homocycle selected from the group of phenyl, naphthyl, acenaphthyl, tetrahydronaphthyl, each homocycle optionally substituted with 1, 2 or 3 substituents, each substituent independently selected from the group of hydroxy, halo, cyano, nitro, amino, mono- or dialkylamino, alkyl, haloalkyl, alkyloxy, haloalkyloxy, carboxyl, alkyloxycarbonyl, aminocarbonyl, morpholinyl and
- 25 mono- or dialkylaminocarbonyl ;
- Het is a monocyclic heterocycle selected from the group of *N*-phenoxypiperidinyl, piperidinyl, pyrrolyl, pyrazolyl, imidazolyl, furanyl, thienyl, oxazolyl, isoxazolyl, thiazolyl, isothiazolyl, pyridinyl, pyrimidinyl, pyrazinyl and pyridazinyl; or a bicyclic heterocycle selected from the group of quinolinyl,
- 30 quinoxalinyl, indolyl, benzimidazolyl, benzoxazolyl, benzisoxazolyl, benzothiazolyl, benzisothiazolyl, benzofuranyl, benzothienyl, 2,3-dihydrobenzo[1,4]dioxinyl and benzo[1,3]dioxolyl ; each monocyclic and bicyclic heterocycle may optionally be substituted with 1, 2 or 3 substituents, each substituent independently selected from the group of halo, hydroxy, alkyl, alkyloxy, and Ar-carbonyl;
- 35 Het¹ is a monocyclic heterocycle selected from furanyl or thienyl; or a bicyclic heterocycle selected from benzofuranyl or benzothienyl; each monocyclic and

- bicyclic heterocycle may optionally be substituted with 1, 2 or 3 substituents, each substituent independently selected from the group of halo, alkyl and Ar; halo is a substituent selected from the group of fluoro, chloro, bromo and iodo; and haloalkyl is a straight or branched saturated hydrocarbon radical having from 1 to 6 carbon atoms or a cyclic saturated hydrocarbon radical having from 3 to 6 carbon atoms or a cyclic saturated hydrocarbon radical having from 3 to 6 carbon atoms attached to a straight or branched saturated hydrocarbon radical having from 1 to 6 carbon atoms; wherein one or more carbon atoms are substituted with one or more halo atoms; provided that when R¹⁰ is alkyl or benzyl, then R⁴ and R⁵ are other than hydrogen; and provided that the compound is other than



- a *N*-oxide thereof, a tautomeric form thereof or a stereochemically isomeric form thereof.

- The compounds according to Formula (Ia) and (Ib) are interrelated in that e.g. a compound according to Formula (Ib), with R⁹ equal to oxo is the tautomeric equivalent of a compound according to Formula (Ia) with R² equal to hydroxy (keto-enol tautomerism).

- In the framework of this application, alkyl is a straight or branched saturated hydrocarbon radical having from 1 to 6 carbon atoms ; or is a cyclic saturated hydrocarbon radical having from 3 to 6 carbon atoms ; or is a cyclic saturated hydrocarbon radical having from 3 to 6 carbon atoms attached to a straight or branched saturated hydrocarbon radical having from 1 to 6 carbon atoms ; wherein each carbon atom can be optionally substituted with hydroxy, alkyloxy or oxo.

- Preferably, alkyl is methyl, ethyl or cyclohexylmethyl, more preferably methyl or ethyl. An interesting embodiment of alkyl in all definitions used hereinbefore or hereinafter is C₁₋₆alkyl which represents a straight or branched saturated hydrocarbon radical having from 1 to 6 carbon atoms such as for example methyl, ethyl, propyl, 2-methyl-ethyl,

pentyl, hexyl and the like. A preferred subgroup of C₁₋₆alkyl is C₁₋₄alkyl which represents a straight or branched saturated hydrocarbon radical having from 1 to 4 carbon atoms such as for example methyl, ethyl, propyl, 2-methyl-ethyl and the like.

- 5 In the framework of this application, Ar is a homocycle selected from the group of phenyl, naphthyl, acenaphthyl, tetrahydronaphthyl, each optionally substituted with 1, 2 or 3 substituents, each substituent independently selected from the group of hydroxy, halo, cyano, nitro, amino, mono- or dialkylamino, alkyl, haloalkyl, alkyloxy, haloalkyloxy, carboxyl, alkyloxycarbonyl, aminocarbonyl, morpholinyl and mono- or
10 dialkylaminocarbonyl. Preferably, Ar is naphthyl or phenyl, each optionally substituted with 1 or 2 halo substituents.

- In the framework of this application, Het is a monocyclic heterocycle selected from the group of *N*-phenoxypiperidinyl, piperidinyl, pyrrolyl, pyrazolyl, imidazolyl, furanyl,
15 thienyl, oxazolyl, isoxazolyl, thiazolyl, isothiazolyl, pyridinyl, pyrimidinyl, pyrazinyl and pyridazinyl; or a bicyclic heterocycle selected from the group of quinolinyl, quinoxalinyl, indolyl, benzimidazolyl, benzoxazolyl, benzisoxazolyl, benzothiazolyl, benzisothiazolyl, benzofuranyl, benzothienyl, 2,3-dihydrobenzo[1,4]dioxinyl and benzo[1,3]dioxolyl ; each monocyclic and bicyclic heterocycle may optionally be
20 substituted with 1, 2 or 3 substituents, each substituent independently selected from the group of halo, hydroxy, alkyl, alkyloxy and Ar-carbonyl. Preferably, Het is thienyl.

- In the framework of this application, halo is a substituent selected from the group of fluoro, chloro, bromo and iodo and haloalkyl is a straight or branched saturated
25 hydrocarbon radical having from 1 to 6 carbon atoms or a cyclic saturated hydrocarbon radical having from 3 to 6 carbon atoms or a cyclic saturated hydrocarbon radical having from 3 to 6 carbon atoms attached to a straight or branched saturated hydrocarbon radical having from 1 to 6 carbon atoms; wherein one or more carbon atoms are substituted with one or more halo atoms. Preferably, halo is bromo, fluoro or
30 chloro and preferably, haloalkyl is polyhaloC₁₋₆alkyl which is defined as mono- or polyhalosubstituted C₁₋₆alkyl, for example, methyl with one or more fluoro atoms, for example, difluoromethyl or trifluoromethyl, 1,1-difluoro-ethyl and the like. In case more than one halo atom is attached to an alkyl group within the definition of haloalkyl or polyhaloC₁₋₆alkyl, they may be the same or different.

35

In the definition of Het, it is meant to include all the possible isomeric forms of the heterocycles, for instance, pyrrolyl comprises 1*H*-pyrrolyl and 2*H*-pyrrolyl.

The Ar, Het or Het¹ listed in the definitions of the substituents of the compounds of formula (I) (see for instance R³) as mentioned hereinbefore or hereinafter may be attached to the remainder of the molecule of formula (Ia) or (Ib) through any ring carbon or heteroatom as appropriate, if not otherwise specified. Thus, for example,
5 when Het is imidazolyl, it may be 1-imidazolyl, 2-imidazolyl, 4-imidazolyl and the like.

Lines drawn from substituents into ring systems indicate that the bond may be attached
10 to any of the suitable ring atoms.

When two vicinal R⁶ radicals are taken together to form a bivalent radical of formula -CH=CH-CH=CH-, this means that the two vicinal R⁶ radicals form together with the phenyl ring to which they are attached a naphthyl.
15

Pharmaceutically acceptable counterions (A⁻) include chloro, bromo, iodo, trifluoroacetate, acetate, triflate, sulfate, sulfonate. The counterion of choice can be introduced using ion exchange resins.

20 The *N*-oxide forms of the present compounds are meant to comprise the compounds of formula (Ia) or (Ib) wherein one or several tertiary nitrogen atoms are oxidized to the so-called *N*-oxide.

The compounds of formula (Ia) and (Ib) may be converted to the corresponding
25 *N*-oxide forms following art-known procedures for converting a trivalent nitrogen into its *N*-oxide form. Said *N*-oxidation reaction may generally be carried out by reacting the starting material of formula (I) with an appropriate organic or inorganic peroxide. Appropriate inorganic peroxides comprise, for example, hydrogen peroxide, alkali metal or earth alkaline metal peroxides, e.g. sodium peroxide, potassium peroxide;
30 appropriate organic peroxides may comprise peroxy acids such as, for example, benzenecarboperoxoic acid or halo substituted benzenecarboperoxoic acid, e.g. 3-chlorobenzenecarboperoxoic acid, peroxyalkanoic acids, e.g. peroxyacetic acid, alkylhydroperoxides, e.g. *t*-butyl hydro-peroxide. Suitable solvents are, for example, water, lower alcohols, e.g. ethanol and the like, hydrocarbons, e.g. toluene, ketones,
35 e.g. 2-butanone, halogenated hydrocarbons, e.g. dichloromethane, and mixtures of such solvents.

It will be appreciated that some of the compounds of formula (I) and their *N*-oxides or tautomeric forms may contain one or more centres of chirality and exist as stereochemically isomeric forms.

- 5 Compounds of either formula (Ia) and (Ib) and some of the intermediate compounds invariably have at least two stereogenic centers in their structure which may lead to at least 4 stereochemically different structures.

- 10 The term "stereochemically isomeric forms" as used hereinbefore or hereinafter defines all the possible stereoisomeric forms which the compounds of formula (Ia) and (Ib), and their *N*-oxides, addition salts or physiologically functional derivatives may possess. Unless otherwise mentioned or indicated, the chemical designation of compounds denotes the mixture of all possible stereochemically isomeric forms, said mixtures containing all diastereomers and enantiomers of the basic molecular structure.
- 15 In particular, stereogenic centers may have the *R*- or *S*-configuration; substituents on bivalent cyclic (partially) saturated radicals may have either the *cis*- or *trans*-configuration. Compounds encompassing double bonds can have an *E* (*entgegen*) or *Z* (*zusammen*) -stereochemistry at said double bond. The terms *cis*, *trans*, *R*, *S*, *E* and *Z* are well known to a person skilled in the art.
- 20 Stereochemically isomeric forms of the compounds of formula (Ia) and (Ib) are obviously intended to be embraced within the scope of this invention.

- Following CAS-nomenclature conventions, when two stereogenic centers of known absolute configuration are present in a molecule, an *R* or *S* descriptor is assigned (based
- 25 on Cahn-Ingold-Prelog sequence rule) to the lowest-numbered chiral center, the reference center. The configuration of the second stereogenic center is indicated using relative descriptors [*R**,*R**] or [*R**,*S**], where *R** is always specified as the reference center and [*R**,*R**] indicates centers with the same chirality and [*R**,*S**] indicates centers of unlike chirality. For example, if the lowest-numbered chiral center in the
- 30 molecule has an *S* configuration and the second center is *R*, the stereo descriptor would be specified as *S*-[*R**,*S**]. If "α" and "β" are used : the position of the highest priority substituent on the asymmetric carbon atom in the ring system having the lowest ring number, is arbitrarily always in the "α" position of the mean plane determined by the ring system. The position of the highest priority substituent on the other asymmetric
- 35 carbon atom in the ring system relative to the position of the highest priority substituent on the reference atom is denominated "α", if it is on the same side of the mean plane

determined by the ring system, or “ β ”, if it is on the other side of the mean plane determined by the ring system.

When a specific stereoisomeric form is indicated, this means that said form is substantially free, *i.e.* associated with less than 50 %, preferably less than 20 %, more preferably less than 10 %, even more preferably less than 5%, further preferably less than 2 % and most preferably less than 1 % of the other isomer(s). Thus, when a compound of formula (I) is for instance specified as (α S, β R), this means that the compound is substantially free of the (α R, β S) isomer.


The compounds of either formula (Ia) and (Ib) may be synthesized in the form of racemic mixtures of enantiomers which can be separated from one another following art-known resolution procedures. The racemic compounds of either formula (Ia) and (Ib) may be converted into the corresponding diastereomeric salt forms by reaction with a suitable chiral acid. Said diastereomeric salt forms are subsequently separated, for example, by selective or fractional crystallization and the enantiomers are liberated therefrom by alkali. An alternative manner of separating the enantiomeric forms of the compounds of either formula (Ia) and (Ib) involves liquid chromatography using a chiral stationary phase. Said pure stereochemically isomeric forms may also be derived from the corresponding pure stereochemically isomeric forms of the appropriate starting materials, provided that the reaction occurs stereospecifically. Preferably if a specific stereoisomer is desired, said compound will be synthesized by stereospecific methods of preparation. These methods will advantageously employ enantiomerically pure starting materials.

The tautomeric forms of the compounds of either formula (Ia) and (Ib) are meant to comprise those compounds of either formula (Ia) and (Ib) wherein e.g. an enol group is converted into a keto group (keto-enol tautomerism).

The invention also comprises derivative compounds (usually called “pro-drugs”) of the pharmacologically-active compounds according to the invention, which are degraded *in vivo* to yield the compounds according to the invention. Pro-drugs are usually (but not always) of lower potency at the target receptor than the compounds to which they are degraded. Pro-drugs are particularly useful when the desired compound has chemical or physical properties that make its administration difficult or inefficient. For example, the desired compound may be only poorly soluble, it may be poorly transported across the mucosal epithelium, or it may have an undesirably short plasma half-life. Further

discussion on pro-drugs may be found in Stella, V. J. *et al.*, "Prodrugs", *Drug Delivery Systems*, 1985, pp. 112-176, and *Drugs*, 1985, **29**, pp. 455-473.



- Pro-drugs forms of the pharmacologically-active compounds according to the invention will generally be compounds according to either Formula (Ia) and (Ib), the pharmaceutically acceptable acid or base addition salts thereof, the stereochemically isomeric forms thereof, the tautomeric forms thereof and the *N*-oxide forms thereof, having an acid group which is esterified or amidated. Included in such esterified acid groups are groups of the formula $-\text{COOR}^x$, where R^x is a C_{1-6} alkyl, phenyl, benzyl or one of the following groups :



- Amidated groups include groups of the formula $-\text{CONR}^y\text{R}^z$, wherein R^y is H, C_{1-6} alkyl, phenyl or benzyl and R^z is $-\text{OH}$, H, C_{1-6} alkyl, phenyl or benzyl.



- Compounds according to the invention having an amino group may be derivatised with a ketone or an aldehyde such as formaldehyde to form a Mannich base. This base will hydrolyze with first order kinetics in aqueous solution.
- Whenever used herein, the term "compounds of formula (Ia) or (Ib)" is meant to also include their *N*-oxide forms, their tautomeric forms or their stereochemically isomeric forms. Of special interest are those compounds of formula (Ia) or (Ib) which are stereochemically pure.
- A first interesting embodiment of the present invention relates to a compound of Formula (Ia-1) or (Ib-1)

-13-





a *N*-oxide thereof, a tautomeric form thereof or a stereochemically isomeric form thereof.

- 5 A second interesting embodiment of the present invention relates to a compound of Formula (Ia-2) or (Ib-2)




a *N*-oxide thereof, a tautomeric form thereof or a stereochemically isomeric form thereof.

A third interesting embodiment of the present invention relates to a compound of
5 Formula (Ia-3) or (Ib-3)




a *N*-oxide thereof, a tautomeric form thereof or a stereochemically isomeric form thereof.

10 A fourth interesting embodiment relates to a compound of formula (Ia) or (Ib) or any
subgroup thereof as mentioned hereinbefore as interesting embodiment wherein the
compound has the following formula



-15-



a *N*-oxide thereof, a tautomeric form thereof or a stereochemically isomeric form thereof wherein


A⁻ is a pharmaceutically acceptable counter ion;

5 R¹ is hydrogen, halo, haloalkyl, cyano, hydroxy, Ar, Het, alkyl, alkyloxy, alkylthio, alkyloxyalkyl, alkylthioalkyl, Ar-alkyl or di(Ar)alkyl ;

p is an integer equal to 1, 2, 3 or 4 ;

R² is hydrogen, hydroxy, mercapto, alkyloxy, alkyloxyalkyloxy, alkylthio,

mono or di(alkyl)amino or a radical of formula



wherein Y is

CH₂, O, S, NH or *N*-alkyl ;

10 R³ is alkyl, Ar, Ar-alkyl, Het or Het-alkyl;

q is an integer equal to zero, 1, 2, 3 or 4 ;

R⁴ and R⁵ each independently are hydrogen, alkyl or benzyl;

R⁴ and R⁵ together and including the N to which they are attached may form a radical selected from the group of pyrrolidinyl, 2-pyrrolinyl, 3-pyrrolinyl,

15 pyrrolyl, imidazolidinyl, pyrazolidinyl, 2-imidazolinyl, 2-pyrazolinyl, imidazolyl, pyrazolyl, triazolyl, piperidinyl, pyridinyl, piperazinyl, pyridazinyl, pyrimidinyl, pyrazinyl, triazinyl, morpholinyl and thiomorpholinyl, each of said rings may optionally be substituted with alkyl, halo, haloalkyl, hydroxy, alkyloxy, amino, mono- or dialkylamino, alkylthio, alkyloxyalkyl, alkylthioalkyl, Ar-alkyl or pyrimidinyl;

20 R⁶ is hydrogen, halo, haloalkyl, hydroxy, Ar, alkyl, alkyloxy, alkylthio, alkyloxyalkyl, alkylthioalkyl, Ar-alkyl or di(Ar)alkyl ; or

two vicinal R⁶ radicals may be taken together to form a bivalent radical of formula -CH=CH-CH=CH- ;

25 r is an integer equal to 1, 2, 3, 4 or 5 ;


R⁷ is hydrogen, alkyl, Ar or Het ;

R⁸ is hydrogen or alkyl ;

- R⁹ is oxo ;
- R¹⁰ is alkyl, alkylcarbonyl, Ar, Ar-alkyl, Ar-carbonyl, Het¹-alkyl or Het¹-carbonyl;
- alkyl is a straight or branched saturated hydrocarbon radical having from 1 to 6 carbon atoms ; or is a cyclic saturated hydrocarbon radical having from 3 to 6 carbon atoms ; or is a cyclic saturated hydrocarbon radical having from 3 to 6 carbon atoms attached to a straight or branched saturated hydrocarbon radical having from 1 to 6 carbon atoms ; wherein each carbon atom can be optionally substituted with hydroxy, alkyloxy or oxo ;
- Ar is a homocycle selected from the group of phenyl, naphthyl, acenaphthyl, tetrahydronaphthyl, each homocycle optionally substituted with 1, 2 or 3 substituents, each substituent independently selected from the group of hydroxy, halo, cyano, nitro, amino, mono- or dialkylamino, alkyl, haloalkyl, alkyloxy, haloalkyloxy, carboxyl, alkyloxycarbonyl, aminocarbonyl, morpholinyl and mono- or dialkylaminocarbonyl ;
- Het is a monocyclic heterocycle selected from the group of *N*-phenoxypiperidinyl, piperidinyl, pyrrolyl, pyrazolyl, imidazolyl, furanyl, thienyl, oxazolyl, isoxazolyl, thiazolyl, isothiazolyl, pyridinyl, pyrimidinyl, pyrazinyl and pyridazinyl; or a bicyclic heterocycle selected from the group of quinolinyl, quinoxalinyl, indolyl, benzimidazolyl, benzoxazolyl, benzisoxazolyl, benzothiazolyl, benzisothiazolyl, benzofuranyl, benzothienyl, 2,3-dihydrobenzo[1,4]dioxinyl and benzo[1,3]dioxolyl ; each monocyclic and bicyclic heterocycle may optionally be substituted with 1, 2 or 3 substituents, each substituent independently selected from the group of halo, hydroxy, alkyl, alkyloxy, and Ar-carbonyl;
- Het¹ is a monocyclic heterocycle selected from furanyl or thienyl; or a bicyclic heterocycle selected from benzofuranyl or benzothienyl; each monocyclic and bicyclic heterocycle may optionally be substituted with 1, 2 or 3 substituents, each substituent independently selected from the group of halo, alkyl and Ar;
- halo is a substituent selected from the group of fluoro, chloro, bromo and iodo; and haloalkyl is a straight or branched saturated hydrocarbon radical having from 1 to 6 carbon atoms or a cyclic saturated hydrocarbon radical having from 3 to 6 carbon atoms or a cyclic saturated hydrocarbon radical having from 3 to 6 carbon atoms attached to a straight or branched saturated hydrocarbon radical having from 1 to 6 carbon atoms; wherein one or more carbon atoms are substituted with one or more halo atoms;

optionally provided that when R^{10} is alkyl or benzyl, then R^4 and R^5 are other than hydrogen; and

optionally provided that the compound is other than



a *N*-oxide thereof, a tautomeric form thereof or a stereochemically isomeric form thereof.


A fifth interesting embodiment relates to a compound of formula (Ia) or (Ib) or any subgroup thereof as mentioned hereinbefore as interesting embodiment wherein

A^- is iodo;

10 R^1 is hydrogen, halo, cyano, Ar, Het, alkyl, and alkyloxy ;

p is an integer equal to 1, 2, 3 or 4 ; in particular 1 or 2;

R^2 is hydrogen, hydroxy, alkyloxy, alkyloxyalkyloxy, alkylthio or a radical of



formula wherein Y is O ;

R^3 is alkyl, Ar, Ar-alkyl or Het ;

15 q is an integer equal to zero, 1, 2, or 3 ;

R^4 and R^5 each independently are hydrogen, alkyl or benzyl;

R^6 is hydrogen, halo or alkyl ; or

two vicinal R^6 radicals may be taken together to form a bivalent radical of formula $-CH=CH-CH=CH-$;

20 r is an integer equal to 1 ;

R^7 is hydrogen ;

R^8 is hydrogen or alkyl ;

R^9 is oxo ; or

R^8 and R^9 together form the radical $-CH=CH-N=$;

25 R^{10} is alkyl, in particular C_{1-4} alkyl;

alkyl is a straight or branched saturated hydrocarbon radical having from 1 to 6 carbon atoms ; or is a cyclic saturated hydrocarbon radical having from 3 to 6 carbon atoms ; or is a a cyclic saturated hydrocarbon radical having from 3 to 6 carbon atoms attached to a straight or branched saturated hydrocarbon radical

- having from 1 to 6 carbon atoms ; wherein each carbon atom can be optionally substituted with hydroxy ;
- Ar is a homocycle selected from the group of phenyl, naphthyl, acenaphthyl and tetrahydronaphthyl, each homocycle optionally substituted with 1, 2 or 3 substituents, each substituent independently selected from the group of halo, haloalkyl, cyano, alkyloxy and morpholinyl ;
- Het is a monocyclic heterocycle selected from the group of *N*-phenoxypiperidinyl, piperidinyl, furanyl, thienyl, pyridinyl and pyrimidinyl; or a bicyclic heterocycle selected from the group of benzothienyl, 2,3-dihydrobenzo[1,4]dioxinyl and benzo[1,3]dioxolyl; each monocyclic and bicyclic heterocycle may optionally be substituted with 1, 2 or 3 alkyl or Ar-carbonyl substituents ; and
- halo is a substituent selected from the group of fluoro, chloro and bromo.

A sixth interesting embodiment relates to a compound of formula (Ia) or (Ib) or any subgroup thereof as mentioned hereinbefore as interesting embodiment wherein R^1 is hydrogen, halo, Ar, alkyl or alkyloxy; preferably, R^1 is halo; more preferably, R^1 is bromo.

A seventh interesting embodiment relates to a compound of formula (Ia) or (Ib) or any subgroup thereof as mentioned hereinbefore as interesting embodiment wherein p is equal to 1; preferably wherein p is equal to 1 and R^1 is other than hydrogen. Preferably, the R^1 substituent is placed in position 5, 6 or 7 of the quinoline ring, more preferably in position 6.

An eighth interesting embodiment relates to a compound of formula (Ia) or (Ib) or any subgroup thereof as mentioned hereinbefore as interesting embodiment wherein R^2 is hydrogen, alkyloxy or alkylthio; preferably, R^2 is alkyloxy, in particular C_{1-4} alkyloxy, more in particular methyloxy; or alkylthio, in particular C_{1-4} alkylthio, more in particular methylthio; more preferably, R^2 is alkyloxy, in particular C_{1-4} alkyloxy, more in particular methyloxy.

A ninth interesting embodiment relates to a compound of formula (Ia) or (Ib) or any subgroup thereof as mentioned hereinbefore as interesting embodiment wherein R^3 is Het, Ar or Ar-alkyl, each optionally substituted with 1 or 2 substituents, that substituent preferably being a halo or haloalkyl, most preferably being a halo; preferably, R^3 is Ar or Ar-alkyl, each optionally substituted with 1 or 2 substituents, that substituent preferably being a halo or haloalkyl, most preferably being a halo; more preferably, R^3

is naphthyl, phenyl, naphthylC₁₋₄alkyl or phenylC₁₋₄alkyl, each optionally substituted with halo, preferably 3-fluoro; more preferably, R³ is naphthyl, phenyl or phenylC₁₋₄alkyl; preferred, R³ is naphthyl, phenyl or phenylethyl.

- 5 A tenth interesting embodiment relates to a compound of formula (Ia) or (Ib) or any subgroup thereof as mentioned hereinbefore as interesting embodiment wherein q is equal to 1, 2 or 3; preferably, q is equal to 3.

- 10 An eleventh interesting embodiment relates to a compound of formula (Ia) or (Ib) or any subgroup thereof as mentioned hereinbefore as interesting embodiment wherein R⁴ and R⁵ each independently are hydrogen or alkyl, in particular hydrogen or C₁₋₄alkyl; preferably C₁₋₄alkyl; most preferably methyl or ethyl.

- 15 A twelfth interesting embodiment relates to a compound of formula (Ia) or (Ib) or any subgroup thereof as mentioned hereinbefore as interesting embodiment wherein R⁴ and R⁵ together and including the N to which they are attached may form a radical selected from the group of pyrrolidinyl, 2-pyrrolinyl, 3-pyrrolinyl, pyrrolyl, imidazolidinyl, pyrazolidinyl, 2-imidazolyl, 2-pyrazolyl, imidazolyl, pyrazolyl, triazolyl, piperidinyl, pyridinyl, piperazinyl, pyridazinyl, pyrimidinyl, pyrazinyl, triazinyl, morpholinyl and thiomorpholinyl, each of said rings may optionally be substituted with
20 alkyl, halo, haloalkyl, hydroxy, alkyloxy, amino, mono- or dialkylamino, alkylthio, alkyloxyalkyl, alkylthioalkyl, Ar-alkyl or pyrimidinyl; preferably R⁴ and R⁵ together and including the N to which they are attached may form a radical selected from the group of piperidinyl, morpholinyl or piperazinyl, each of said rings may optionally be
25 substituted with alkyl or Ar-alkyl; or R⁴ and R⁵ together and including the N to which they are attached may form a radical selected from imidazolyl or piperidinyl.

- A thirteenth interesting embodiment relates to a compound of formula (Ia) or (Ib) or any subgroup thereof as mentioned hereinbefore as interesting embodiment wherein R⁶
30 is hydrogen, alkyl or halo; preferably, R⁶ is hydrogen or halo; more preferably, R⁶ is hydrogen.

- A fourteenth interesting embodiment relates to a compound of formula (Ia) or (Ib) or any subgroup thereof as mentioned hereinbefore as interesting embodiment wherein r is
35 1 or 2; preferably r is 1.

A fifteenth interesting embodiment relates to a compound of formula (Ia) or (Ib) or any subgroup thereof as mentioned hereinbefore as interesting embodiment wherein R^7 is hydrogen or methyl; preferably R^7 is hydrogen.

- 5 A sixteenth interesting embodiment relates to a compound of formula (Ia) or (Ib) or any subgroup thereof as mentioned hereinbefore as interesting embodiment wherein, for compounds according to Formula (Ib) only, R^8 is alkyl, preferably methyl, and R^9 is oxygen.

- 10 A seventeenth interesting embodiment relates to a compound of formula (Ia) or (Ib) or any subgroup thereof as mentioned hereinbefore as interesting embodiment wherein R^{10} is alkyl; preferably C_{1-6} alkyl; more preferably C_{1-4} alkyl.

- 15 An eighteenth interesting embodiment relates to a compound of formula (Ia) or (Ib) or any subgroup thereof as mentioned hereinbefore as interesting embodiment wherein A⁻ is chloro, bromo, iodo, trifluoroacetate, acetate, triflate, sulfate, sulfonate; preferably chloro, bromo or iodo, more preferably iodo.

- 20 A nineteenth interesting embodiment relates to a compound of formula (Ia) or (Ib) or any subgroup thereof as mentioned hereinbefore as interesting embodiment wherein the compound is a compound according to formula (Ia).

- 25 A twentieth interesting embodiment relates to a compound of formula (Ia) or (Ib) or any subgroup thereof as mentioned hereinbefore as interesting embodiment wherein one or more, preferably all, of the following definitions apply :

R^1 is halo, in particular bromo;

$p = 1$;

R^2 is alkyloxy, in particular C_{1-4} alkyloxy, more in particular methoxy; or alkylthio, in particular C_{1-4} alkylthio, more in particular methylthio;

- 30 R^3 is naphthyl; phenyl; phenylethyl or Het, in particular thienyl;

$q = 1, 2$ or 3 ;

R^4 and R^5 each independently are alkyl, in particular C_{1-4} alkyl; or R^4 and R^5 together and including the N to which they are attached may form a radical selected from imidazolyl or piperidiny;

- 35 R^6 is hydrogen or halo;

r is equal to 1 ;

R^7 is hydrogen;

R¹⁰ is alkyl, in particular C₁₋₆alkyl, more in particular C₁₋₄alkyl.

A twenty first interesting embodiment is the use of a compound of formula (Ia) or (Ib) or any subgroup thereof as mentioned hereinbefore as interesting embodiment for the manufacture of a medicament for the treatment of an infection with a gram-positive and/or a gram-negative bacterium.


A twenty second interesting embodiment is the use of a compound of formula (Ia) or (Ib) or any subgroup thereof as mentioned hereinbefore as interesting embodiment for the manufacture of a medicament for the treatment of an infection with a gram-positive bacterium.


A twenty third interesting embodiment is the use of the compounds of formula (Ia) or (Ib) or any subgroup thereof as mentioned hereinbefore as interesting embodiment for the manufacture of a medicament for the treatment of an infection with a gram-negative bacterium.

A twenty fourth interesting embodiment is the use of a compound of formula (Ia) or (Ib) or any subgroup thereof as mentioned hereinbefore as interesting embodiment for the manufacture of a medicament for the treatment of a bacterial infection wherein the compound of formula (Ia) or (Ib) has a IC₉₀ < 15 µl/ml against at least one bacterium, in particular a gram-positive bacterium, preferably a IC₉₀ < 10 µl/ml, more preferably a IC₉₀ < 5 µl/ml; the IC₉₀ value being determined as described hereinafter.

Preferably, in the compounds of formula (Ia) and (Ib) or any subgroup thereof as mentioned hereinbefore as interesting embodiment, the term "alkyl" represents C₁₋₆alkyl, more preferably C₁₋₄alkyl.

Preferred compounds are selected from the following :





a *N*-oxide thereof or a stereochemically isomeric form thereof.

Especially preferred compounds are selected from compound 121, 102, 103, 10, A, E, K and R (see the Tables hereinbelow), a *N*-oxide thereof, a tautomeric form thereof or a stereochemically isomeric form thereof.

The present invention also relates to compounds A to C and E to U, a *N*-oxide thereof or a stereochemically isomeric form thereof, in particular to compounds A, E, K and R or a stereochemically isomeric form thereof.


10

The compounds of formula (I) can be prepared according to the methods described in WO 2004/011436, which is incorporated herein by reference.

In general, the compounds according to the invention can be prepared by a succession of steps, each of which is known to the skilled person.

15


In particular, compounds of formula (Ia) or (Ib) can be prepared by reacting an intermediate of formula (II-a) respectively (II-b) with an intermediate of formula (III) in the presence of a suitable solvent, such as for example acetone.



Intermediates according to formula (II-a) can be prepared by reacting an intermediate compound of formula (IV) with an intermediate compound of formula (V) according to the following reaction scheme (1) :

5

Scheme 1



using nBuLi in a mixture of diisopropyl amine and tetrahydrofuran, wherein all variables are defined as in formula (Ia). Stirring may enhance the rate of the reaction. The reaction may conveniently be carried out at a temperature ranging between -20 and -70 °C.


10

The same reaction procedure can be used to synthesize intermediates of formula (II-b).

15

The starting materials and the intermediate compounds of formula (IV) and (V) are compounds that are either commercially available or may be prepared according to conventional reaction procedures generally known in the art. For example, intermediate compounds of formula (IV-a) or (IV-b) may be prepared according to the following reaction scheme (2):


Scheme 2



wherein all variables are defined as in formula (Ia). Reaction scheme (2) comprises


- 5 step (a) in which an appropriately substituted aniline is reacted with an appropriate acylchloride such as 3-phenylpropionyl chloride, 3-fluorobenzenepropionyl chloride or *p*-chlorobenzenepropionyl chloride, in the presence of a suitable base, such as triethylamine and a suitable reaction-inert solvent, such as methylene chloride or ethylene dichloride. The reaction may conveniently be carried out at a temperature ranging between room temperature and reflux temperature. In a next step (b) the
- 10 adduct obtained in step (a) is reacted with phosphoryl chloride (POCl₃) in the presence of *N,N*-dimethylformamide (Vilsmeier-Haack formylation followed by cyclization). The reaction may conveniently be carried out at a temperature ranging between room temperature and reflux temperature. In a next step (c-1), a specific R²-group, wherein R² is for example a C₁₋₆alkyloxy radical is introduced by reacting the intermediate
- 15 compound obtained in step (b) with a $O-C_{1-6}$ alkyl in the presence of a suitable solvent, such as for example HO-C₁₋₆alkyl. The intermediate compound obtained in step (b) can also be converted into an intermediate compound wherein R² is for example a C₁₋₆alkylthio radical by reaction with S=C(NH₂)₂ in the presence of a suitable solvent, such as for example an alcohol, e.g. ethanol (step (c-2)) followed by reaction with
- 20 C₁₋₆alkyl-I in the presence of a suitable base, such as for example K₂CO₃ and a suitable solvent, such as for example 2-propanone.

Intermediate compounds according to formula (IV-c) may be prepared according to the following reaction scheme (3), wherein in a first step (a) an optionally substituted indole-2,3-dione is reacted with an optionally substituted 3-phenylpropionaldehyde in the presence of a suitable base such as sodium hydroxide (Pfitzinger reaction), after
5 which the carboxylic acid compound is decarboxylated in a next step (b) at high temperature in the presence of a suitable reaction-inert solvent such as diphenylether.

Scheme 3

It is evident that in the foregoing and in the following reactions, the reaction products
10 may be isolated from the reaction medium and, if necessary, further purified according to methodologies generally known in the art, such as extraction, crystallization and chromatography. It is further evident that reaction products that exist in more than one enantiomeric form, may be isolated from their mixture by known techniques, in particular preparative chromatography, such as preparative HPLC. Typically,
15 compounds of formula (Ia) or (Ib) may be separated into their isomeric forms.

The intermediate compounds of formula (V) are compounds that are either commercially available or may be prepared according to conventional reaction procedures generally known in the art. For example, intermediate compounds of
20 formula (V) may be prepared according to the following reaction scheme (4):

Scheme 4

- Reaction scheme (4) comprises step (a) in which R^3 , in particular an appropriately substituted Ar, more in particular an appropriately substituted phenyl, is reacted by Friedel-Craft reaction with an appropriate acylchloride such as 3-chloropropionyl chloride or 4-chlorobutyryl chloride, in the presence of a suitable Lewis acid, such as for example AlCl_3 , FeCl_3 , SnCl_4 , TiCl_4 or ZnCl_2 and a suitable reaction-inert solvent, such as methylene chloride or ethylene dichloride. The reaction may conveniently be carried out at a temperature ranging between room temperature and reflux temperature. In a next step (b) an amino group ($-\text{NR}^4\text{R}^5$) is introduced by reacting the intermediate compound obtained in step (a) with a primary or secondary amine (HNR^4R^5).

- In general, bacterial pathogens may be classified as either gram-positive or gram-negative pathogens. Antibiotic compounds with activity against both gram-positive and gram-negative pathogens are generally regarded as having a broad spectrum of activity. The compounds of the present invention are regarded as active against gram-positive and/or gram-negative bacterial pathogens. In particular, the present compounds are active against at least one gram-positive bacterium, preferably against several gram-positive bacteria, more preferably against one or more gram-positive bacteria and/or one or more gram-negative bacteria.

The present compounds have bactericidal or bacteriostatic activity.

- Examples of gram-positive and gram-negative aerobic and anaerobic bacteria, include Staphylococci, for example *S. aureus*; Enterococci, for example *E. faecalis*; Streptococci, for example *S. pneumoniae*, *S. mutans*, *S. pyogenes*; Bacilli, for example *Bacillus subtilis*; Listeria, for example *Listeria monocytogenes*; Haemophilus, for example *H. influenza*; Moraxella, for example *M. catarrhalis*; Pseudomonas, for example *Pseudomonas aeruginosa*; and Escherichia, for example *E. coli*. Gram-positive pathogens, for example Staphylococci, Enterococci and Streptococci are particularly important because of the development of resistant strains which are both difficult to treat and difficult to eradicate from for example a hospital environment once established. Examples of such strains are methicillin resistant *Staphylococcus aureus*

(MRSA), methicillin resistant coagulase negative staphylococci (MRCNS), penicillin resistant *Streptococcus pneumoniae* and multiple resistant *Enterococcus faecium*.

5 The compounds of the present invention also show activity against resistant bacterial strains.

The compounds of the present invention are especially active against *Staphylococcus aureus*, including resistant *Staphylococcus aureus* such as for example methicillin resistant *Staphylococcus aureus* (MRSA), and *Streptococcus pneumoniae*, in particular
10 against *Staphylococcus aureus*.

In particular, the compounds of the present invention are active on those bacteria of which the viability depends on proper functioning of F1F0 ATP synthase. Without being bound to any theory, it is taught that the activity of the present compounds lies in
15 inhibition of the F1F0 ATP synthase, in particular the inhibition of the F0 complex of the F1F0 ATP synthase, more in particular the inhibition of subunit c of the F0 complex of the F1F0 ATP synthase, leading to killing of the bacteria by depletion of the cellular ATP levels of the bacteria.

20 Whenever used hereinbefore or hereinafter, that the compounds can treat a bacterial infection it is meant that the compounds can treat an infection with one or more bacterial strains.

Whenever used hereinbefore or hereinafter, that the bacterial infection is other than a Mycobacterial infection it is meant that the bacterial infection is other than an infection
25 with one or more Mycobacteria strains.

The exact dosage and frequency of administration of the present compounds depends on the particular compound of formula (Ia) or (Ib) used, the particular condition being treated, the severity of the condition being treated, the age, weight, gender, diet, time of
30 administration and general physical condition of the particular patient, the mode of administration as well as other medication the individual may be taking, as is well known to those skilled in the art. Furthermore, it is evident that the effective daily amount may be lowered or increased depending on the response of the treated subject and/or depending on the evaluation of the physician prescribing the compounds of the
35 instant invention.

The compound of the present invention may be administered in a pharmaceutically acceptable form optionally in a pharmaceutically acceptable carrier. The compounds and compositions comprising the compounds can be administered by routes such as topically, locally or systemically. Systemic application includes any method of
5 introducing the compound into the tissues of the body, e.g., intrathecal, epidural, intramuscular, transdermal, intravenous, intraperitoneal, subcutaneous, sublingual, rectal, and oral administration. The specific dosage of antibacterial to be administered, as well as the duration of treatment, may be adjusted as needed.

- 10 Bacterial infections which may be treated by the present compounds include, for example, central nervous system infections, external ear infections, infections of the middle ear, such as acute otitis media, infections of the cranial sinuses, eye infections, infections of the oral cavity, such as infections of the teeth, gums and mucosa, upper respiratory tract infections, lower respiratory tract infections, genitourinary infections,
15 gastrointestinal infections, gynecological infections, septicemia, bone and joint infections, skin and skin structure infections, bacterial endocarditis, burns, antibacterial prophylaxis of surgery, and antibacterial prophylaxis in immunosuppressed patients, such as patients receiving cancer chemotherapy, or organ transplant patients.
- 20 Given the fact that the compounds of formula (Ia) or (Ib) are active against bacterial infections, the present compounds may be combined with other antibacterial agents in order to effectively combat bacterial infections.

Therefore, the present invention also relates to a combination of (a) a compound of
25 formula (Ia) or (Ib), and (b) one or more other antibacterial agents provided that the one or more other antibacterial agents are other than antimycobacterial agents.

The present invention also relates to a combination of (a) a compound of formula (Ia) or (Ib), and (b) one or more other antibacterial agents provided that the one or more
30 other antibacterial agents are other than antimycobacterial agents, for use as a medicine.

A pharmaceutical composition comprising a pharmaceutically acceptable carrier and, as active ingredient, a therapeutically effective amount of (a) a compound of formula
35 (Ia) or (Ib), and (b) one or more other antibacterial agents provided that the one or more other antibacterial agents are other than antimycobacterial agents, is also comprised by the present invention.

The present invention also relates to the use of a combination or pharmaceutical composition as defined above for the treatment of a bacterial infection or for the manufacture of a medicament for the treatment of a bacterial infection.

5

The present pharmaceutical composition may have various pharmaceutical forms for administration purposes. As appropriate compositions there may be cited all compositions usually employed for systemically administering drugs. To prepare the pharmaceutical compositions of this invention, an effective amount of the particular compounds, optionally in addition salt form, as the active ingredient is combined in intimate admixture with a pharmaceutically acceptable carrier, which carrier may take a wide variety of forms depending on the form of preparation desired for administration. These pharmaceutical compositions are desirable in unitary dosage form suitable, in particular, for administration orally or by parenteral injection. For example, in preparing the compositions in oral dosage form, any of the usual pharmaceutical media may be employed such as, for example, water, glycols, oils, alcohols and the like in the case of oral liquid preparations such as suspensions, syrups, elixirs, emulsions and solutions; or solid carriers such as starches, sugars, kaolin, diluents, lubricants, binders, disintegrating agents and the like in the case of powders, pills, capsules and tablets. Because of their ease in administration, tablets and capsules represent the most advantageous oral unit dosage forms in which case solid pharmaceutical carriers are obviously employed. For parenteral compositions, the carrier will usually comprise sterile water, at least in large part, though other ingredients, for example, to aid solubility, may be included. Injectable solutions, for example, may be prepared in which the carrier comprises saline solution, glucose solution or a mixture of saline and glucose solution. Injectable suspensions may also be prepared in which case appropriate liquid carriers, suspending agents and the like may be employed. Also included are solid form preparations which are intended to be converted, shortly before use, to liquid form preparations.

30

Depending on the mode of administration, the pharmaceutical composition will preferably comprise from 0.05 to 99 % by weight, more preferably from 0.1 to 70 % by weight of the active ingredients, and, from 1 to 99.95 % by weight, more preferably from 30 to 99.9 weight % of a pharmaceutically acceptable carrier, all percentages being based on the total composition.

35

The weight to weight ratio's of the compound of formula (Ia) or (Ib) and (b) the other antibacterial agent(s) when given as a combination may be determined by the person skilled in the art. Said ratio and the exact dosage and frequency of administration depends on the particular compound of formula (Ia) or (Ib) and the other antibacterial agent(s) used, the particular condition being treated, the severity of the condition being treated, the age, weight, gender, diet, time of administration and general physical condition of the particular patient, the mode of administration as well as other medication the individual may be taking, as is well known to those skilled in the art. Furthermore, it is evident that the effective daily amount may be lowered or increased depending on the response of the treated subject and/or depending on the evaluation of the physician prescribing the compounds of the instant invention.

The compounds of formula (Ia) or (Ib) and the one or more other antibacterial agents may be combined in a single preparation or they may be formulated in separate preparations so that they can be administered simultaneously, separately or sequentially. Thus, the present invention also relates to a product containing (a) a compound of formula (Ia) or (Ib), and (b) one or more other antibacterial agents provided that the one or more other antibacterial agents are other than antimycobacterial agents, as a combined preparation for simultaneous, separate or sequential use in the treatment of a bacterial infection.

The pharmaceutical composition may additionally contain various other ingredients known in the art, for example, a lubricant, stabilising agent, buffering agent, emulsifying agent, viscosity-regulating agent, surfactant, preservative, flavouring or colorant.

It is especially advantageous to formulate the aforementioned pharmaceutical compositions in unit dosage form for ease of administration and uniformity of dosage. Unit dosage form as used herein refers to physically discrete units suitable as unitary dosages, each unit containing a predetermined quantity of active ingredient calculated to produce the desired therapeutic effect in association with the required pharmaceutical carrier. Examples of such unit dosage forms are tablets (including scored or coated tablets), capsules, pills, powder packets, wafers, suppositories, injectable solutions or suspensions and the like, and segregated multiples thereof. The daily dosage of the compound according to the invention will, of course, vary with the compound employed, the mode of administration, the treatment desired and the bacterial disease indicated.

The other antibacterial agents which may be combined with the compounds of formula (I) are antibacterial agents known in the art. The other antibacterial agents comprise antibiotics of the β -lactam group such as natural penicillins, semisynthetic penicillins, natural cephalosporins, semisynthetic cephalosporins, cephamycins, 1-oxacephems, clavulanic acids, penems, carbapenems, nocardicins, monobactams; tetracyclines, anhydrotetracyclines, anthracyclines; aminoglycosides; nucleosides such as *N*-nucleosides, *C*-nucleosides, carbocyclic nucleosides, blasticidin S; macrolides such as 12-membered ring macrolides, 14-membered ring macrolides, 16-membered ring macrolides; ansamycins; peptides such as bleomycins, gramicidins, polymyxins, bacitracins, large ring peptide antibiotics containing lactone linkages, actinomycins, amphomycin, capreomycin, distamycin, enduracidins, mikamycin, neocarzinostatin, stendomycin, viomycin, virginiamycin; cycloheximide; cycloserine; variotin; sarkomycin A; novobiocin; griseofulvin; chloramphenicol; mitomycins; fumagillin; monensins; pyrrolnitrin; fosfomycin; fusidic acid; D-(*p*-hydroxyphenyl)glycine; D-phenylglycine; enediynes.

Specific antibiotics which may be combined with the present compounds of formula (Ia) or (Ib) are for example benzylpenicillin (potassium, procaine, benzathine), phenoxymethylpenicillin (potassium), phenethicillin potassium, propicillin, carbenicillin (disodium, phenyl sodium, indanyl sodium), sulbenicillin, ticarcillin disodium, methicillin sodium, oxacillin sodium, cloxacillin sodium, dicloxacillin, flucloxacillin, ampicillin, mezlocillin, piperacillin sodium, amoxicillin, ciclacillin, hecclacillin, sulbactam sodium, talampicillin hydrochloride, bacampicillin hydrochloride, pivmecillinam, cephalixin, cefaclor, cephaloglycin, cefadroxil, cephadrine, cefroxadine, cephapirin sodium, cephalothin sodium, cephradine sodium, cefsulodin sodium, cephaloridine, cefatrizine, cefoperazone sodium, cefamandole, vefotiam hydrochloride, cefazolin sodium, ceftizoxime sodium, cefotaxime sodium, cefmenoxime hydrochloride, cefuroxime, ceftriaxone sodium, ceftazidime, cefoxitin, cefmetazole, cefotetan, latamoxef, clavulanic acid, imipenem, aztreonam, tetracycline, chlortetracycline hydrochloride, demethylchlortetracycline, oxytetracycline, methacycline, doxycycline, rolitetracycline, minocycline, daunorubicin hydrochloride, doxorubicin, aclarubicin, kanamycin sulfate, bekanamycin, tobramycin, gentamycin sulfate, dibekacin, amikacin, micromomicin, ribostamycin, neomycin sulfate, paromomycin sulfate, streptomycin sulfate, dihydrostreptomycin, destomycin A, hygromycin B, apramycin, sisomicin, netilmicin sulfate, spectinomycin hydrochloride, astromicin sulfate, validamycin, kasugamycin, polyoxin, blasticidin S, erythromycin, erythromycin estolate, oleandomycin phosphate, tracetyloleandomycin, kitasamycin,

josamycin, spiramycin, tylosin, ivermectin, midecamycin, bleomycin sulfate, peplomycin sulfate, gramicidin S, polymyxin B, bacitracin, colistin sulfate, colistinmethanesulfonate sodium, enramycin, mikamycin, virginiamycin, capreomycin sulfate, viomycin, enviomycin, vancomycin, actinomycin D, neocarzinostatin, bestatin, 5 pepstatin, monensin, lasalocid, salinomycin, amphotericin B, nystatin, natamycin, trichomycin, mithramycin, lincomycin, clindamycin, clindamycin palmitate hydrochloride, flavophospholipol, cycloserine, pecilocin, griseofulvin, chloramphenicol, chloramphenicol palmitate, mitomycin C, pyrrolnitrin, fosfomycin, fusidic acid, bicozamycin, tiamulin, siccanin.


10

EXPERIMENTAL PART

Of some compounds the absolute stereochemical configuration of the stereogenic carbon atom(s) therein was not experimentally determined. In those cases the stereochemically isomeric form which was first isolated is designated as "A" and the 15 second as "B", without further reference to the actual stereochemical configuration. However, said "A" and "B" isomeric forms can be unambiguously characterized by a person skilled in the art, using art-known methods such as, for example, X-ray diffraction.

In case "A" and "B" are stereoisomeric mixtures, they can be further separated 20 whereby the respective first fractions isolated are designated "A1" respectively "B1" and the second as "A2" respectively "B2", without further reference to the actual stereochemical configuration. However, said "A1", "A2" and "B1", "B2" isomeric forms can be unambiguously characterized by a person skilled in the art, using art-known methods such as, for example, X-ray diffraction.

25 For some of the compounds, stereochemical configurations are indicated in the structures. The configurations are relative configurations indicating that the groups concerned are located in the same or opposite plane of the molecule



30 The present compounds which are also described in WO 2004/011436 bear the same compound number as in WO 2004/011436. The Ex. Nr. in the below Tables and in the synthesis protocols herein below refer to the Example numbers of WO 2004/011436 indicating according to which protocol the compounds were prepared.

35 Additional compounds are indicated by way of letters.

Synthesis of compounds A, B and C

A solution of compound 15 of WO 2004/011436 (prepared according to B7) (0.1g, 0.18 mmol) and ethyl iodide (0.02 ml, 0.19 mmol) in acetone (2 ml) was stirred at 60°C for
5 12 hours. The solvent was evaporated and the residue was crystallized from diisopropylether and acetone. Yield : 0.02 g of compound A (diastereoisomer B) (16 %, mp=244°C).

Compound B (diastereoisomer A) was prepared according to the above protocol but
10 starting from compound 14 of WO 2004/011436 (prepared according to B7). Yield : 71 %, mp=204°C.

Compound C (diastereoisomer B) was prepared according to the above protocol starting from compound 15 of WO 2004/011436 (prepared according to B7) and
15 reacting this compound 15 with butyl iodide. Yield : 50 %, mp=182°C.


Synthesis of compounds E and F

A solution of compound 95 (0.1 g, 0.187 mmol) of WO 2004/011436 (prepared according to B1) and methyl iodide (0.02 ml, 0.281 mmol) in acetone (2 ml) was stirred
20 at room temperature for 3 hours. The solvent was evaporated and the residue was crystallized from diisopropylether. Yield : 0.115 g of compound E (diastereoisomer A) (91 %, mp>250°C).

Compound F (diastereoisomer B) was prepared according to the above protocol but
25 starting from compound 96 of WO 2004/011436 (prepared according to B1). Yield : 87 %, mp>250°C.

Synthesis of compound G

Compound G was prepared according to the following scheme :



Synthesis of intermediate compound 1

Intermediate compound 1 was prepared in the same way as intermediate compound 12 of WO 2004/011436 (according to A8) with 4-chlorobutylchloride. The residue
 5 (70.7 g) was purified by column chromatography over silica gel (eluent: Cyclohexane/AcOEt; 70:30; 20-45 μ m). Two fractions (F1 and F2) were collected and the solvent was evaporated. F1 : 45.5 g of intermediate compound 1 (yield = 63 %).

Synthesis of intermediate compound 2

10 A solution of intermediate compound 1 (2 g, 0.0086 mol), dimethylamine hydrochloride (1.4 g, 0.0172 mol) and potassium carbonate (2.4 g, 0.0174 mol) in acetonitrile (30 ml) was stirred at 80°C for 12 hours. The solvent was evaporated and the residue (2.8 g) was purified by column chromatography over silica gel (eluent: CH₂Cl₂/CH₃OH/NH₄OH; 93/7/0.5; 20-45 μ m) yielding 1 g of intermediate compound 2
 15 as an oil (yield = 49 %).

Synthesis of intermediate compound 3

*n*BuLi 1.6M (3.3 ml, 0.0024 mol) was added slowly at -20°C under N₂ flow to a solution of diisopropylamine (0.33 ml, 0.0024 mol) in tetrahydrofuran (5 ml). The
 20 mixture was stirred at -20°C for 20 minutes, then cooled at -70°C. A solution of intermediate compound 3 of WO 2004/011436 (prepared according to A3) (0.6 g, 0.0018 mol) in tetrahydrofuran (5 ml) was added slowly. The mixture was stirred at -70°C for 1h30. A solution of present intermediate compound 2 (0.54 g, 0.0022 mol) in

tetrahydrofuran (5 ml) was added slowly. The mixture was stirred at -70°C for 3 hours, hydrolyzed at -30°C with ice water, and extracted with EtOAc. The organic layer was separated, dried over MgSO_4 , filtered, and the solvent was evaporated. The residue (5.3 g) was purified by column chromatography over silica gel (eluent:
5 $\text{CH}_2\text{Cl}_2/\text{CH}_3\text{OH}/\text{NH}_4\text{OH}$; 99/1/0.1; 20-45 μm). Two fractions were collected and the solvent was evaporated. Fractions were crystallized separately from diisopropylether and diethylether yielding 0.04 g of diastereoisomer A (4 %) and 0.04 g of present intermediate compound 3 (diastereoisomer B) (4 %).

10 *Synthesis of compound G*

A solution of intermediate compound 3 (0.04 g, 0.07 mmol) and methyl iodide (0.01 ml, 0.14 mmol) in acetone (2 ml) was stirred at room temperature for 2 hours. The solvent was evaporated and the residue was crystallized from diisopropylether and acetone yielding 0.047 g of compound G (diastereoisomer B) (93 %).

15

Synthesis of compounds H and I

A solution of compound 197 of WO 2004/01146 (0.15g, 0.257mmol) (prepared according to B7) and methyl iodide (0.02ml, 0.257mmol) in acetone (3ml) was stirred
20 at room temperature for 18 hours. The solvent was evaporated and the residue was crystallized from diethylether yielding 0.154g of compound I (diastereoisomer A) (83 %, $\text{mp}=188^{\circ}\text{C}$).

Compound H (diastereoisomer B) was prepared according to the above protocol but
25 starting from compound 191 of WO 2004/011436 (prepared according to B7).
Yield : 0.154 g of compound H (75 %, $\text{mp}=172^{\circ}\text{C}$).


Synthesis of compounds K and J

A solution of compound 66 of WO 2004/011436 (0.10 g, 0.187 mmol) (prepared
30 according to B1) and methyl iodide (0.018 ml, 0.281 mmol) in acetone (3 ml) was stirred at room temperature for 18 hours. The precipitate was filtered off, washed with acetone and dried at 70°C yielding 0.08 g of compound J (diastereoisomer B) (53 %, $\text{mp}=175^{\circ}\text{C}$).

Compound K (diastereoisomer A) was prepared according to the above protocol but starting from compound 65 of WO 2004/011436 (prepared according to B1). Yield : 64 % of compound K (245°C).


5 Synthesis of compound L

Compound L was synthesized according to the following scheme




Synthesis of intermediate compound 4

a) Preparation of intermediate compound 4a



- Benzenepropanoyl chloride (0.488 mol) was added dropwise at room temperature to a solution of 4-bromobenzenamine (0.407 mol) in Et₃N (70 ml) and CH₂Cl₂ (700 ml) and the mixture was stirred at room temperature overnight. The mixture was poured out into water and concentrated NH₄OH, and extracted with CH₂Cl₂. The organic layer was dried (MgSO₄), filtered, and the solvent was evaporated. The residue was crystallized from diethyl ether. The residue (119.67 g) was taken up in CH₂Cl₂ and washed with HCl 1N. The organic layer was dried (MgSO₄), filtered, and the solvent was evaporated. Yield: 107.67 g of intermediate compound 4a.


b) Preparation of intermediate compound 4b



The reaction was carried out twice. POCl₃ (1.225 mol) was added dropwise at 10°C to N,N-dimethylformamide (0.525 mol). Then intermediate compound 4a (0.175 mol) was


added at room temperature. The mixture was stirred overnight at 80°C, poured out on ice and extracted with CH₂Cl₂. The organic layer was dried (MgSO₄), filtered, and the solvent was evaporated. The product was used without further purification. Yield: 77.62 g (67 %) of intermediate compound 4b.

c) Preparation of intermediate compound 4c



- 5 A mixture of intermediate compound 4b (0.045 mol) and thiourea (0.05 mol) in ethanol (150 ml) was stirred and refluxed for 8 hours and then brought to room temperature. A solution of KOH (0.068 mol) in water (15 ml) was added. The mixture was stirred and refluxed for 1 hour and poured out on ice. The precipitate was filtered off, washed with H₂O and dried. Yield: 11 g (74 %) of intermediate compound 4c.


d) Preparation of intermediate compound 4



- 10 CH₃I (0.037 mol) was added slowly at room temperature to a mixture of intermediate compound 4c (0.033 mol) and K₂CO₃ (0.037 mol) in 2-propanone (150 ml). The mixture was stirred at room temperature for 8 hours, poured out into H₂O and extracted with CH₂Cl₂. The organic layer was separated, dried (MgSO₄), filtered and the solvent was evaporated. Yielding: 11.2 g. Part of this fraction (2 g) was crystallized from diethyl ether. The precipitate was filtered off and dried. Yield: 1.45 g (70 %) of intermediate compound 4.

Synthesis of intermediate compound 5

Preparation of intermediate compound 5




- 20 nBuLi 1.6M in hexane (0.0035 mol) was added dropwise at -20°C to a solution of N-(1-methylethyl)-2-propanamine (0.0035 mol) in tetrahydrofuran (7ml) under N₂ flow. The mixture was stirred at -20°C for 20 minutes, then cooled to -70°C. A solution of intermediate compound 4 (0.003 mol) in tetrahydrofuran (10 ml) was added. The mixture was stirred at -70°C for 1 hour. A solution of 5-(dimethylamino)-1-phenyl-1-pentanone (J.Am.Chem.Soc. 1972, 94(11), 3877-3883) (0.0035 mol) in tetrahydrofuran

- (10ml) was added. The mixture was stirred at -70°C for 3 hours. H_2O was added. The mixture was extracted with EtOAc. The organic layer was washed with saturated NaCl, dried (MgSO_4), filtered and the solvent was evaporated. The residue (2 g) was purified by column chromatography over silica gel (eluent: $\text{CH}_2\text{Cl}_2/\text{CH}_3\text{OH}/\text{NH}_4\text{OH}$ 94/6/0.3; 15-40 μm). Two fractions were collected and the solvent was evaporated. Residue 1 was crystallized from diisopropylether. The precipitate was filtered off and dried. Yield: (2 %) of intermediate compound 5 (diastereoisomer A).

Synthesis of compound L

- A solution of intermediate compound 5 (0.27g, 0.49 mmol) and methyl iodide (0.046 ml, 0.74 mmol) in acetone (5 ml) was stirred at room temperature for 24 hours. The solvent was evaporated, the residue was taken up in diisopropylether/ CH_2Cl_2 , the precipitate was filtered off, washed with diisopropylether and dried at 70°C yielding 0.17 g of compound L (diastereoisomer A) (51 %, $\text{mp}=233^{\circ}\text{C}$).

Synthesis of compound U



Synthesis of intermediate compound 6

- $n\text{BuLi}$ 1.6M (1.15 ml, 1.83 mmol) was added slowly at -20°C under N_2 flow to a solution of diisopropylamine (0.256 ml, 1.83 mmol) in tetrahydrofuran (4 ml). The mixture was stirred at -20°C for 20 minutes, then cooled at -70°C . A solution of intermediate compound 3 of WO 2004/011436 (0.5 g, 1.52 mmol) in tetrahydrofuran (5 ml) was added slowly. The mixture was stirred at -70°C for 1 hour. A solution of

1-phenyl-5-(1-piperidiny)-1-pentanone (0.45 g, 1.83 mmol) in tetrahydrofuran (5 ml) was added slowly. The mixture was stirred at -70°C for 1.5 hours, hydrolyzed at -70°C with water, and extracted with EtOAc. The organic layer was separated, washed with brine, dried over MgSO_4 , filtered, and the solvent was evaporated. The residue (0.9 g) was purified by column chromatography over silica gel (eluent: $\text{CH}_2\text{Cl}_2/\text{CH}_3\text{OH}/\text{NH}_4\text{OH}$; 97/3/0.1; kromasil $10\mu\text{m}$). Two fractions were collected and the solvent was evaporated. The first product fraction yielded intermediate compound 6 (A). Yield: 0.085 g of intermediate compound 6 (diastereoisomer A) (10 %, mp= 129°C). The second product fraction was crystallized from diisopropylether to give diastereoisomer B (yield = 6 %, mp= 166°C).

Synthesis of compound U

A solution of intermediate compound 6 (A) (0.020 g, 0.035 mmol) and methyl iodide (0.0032 ml, 0.052 mmol) in acetone (2 ml) was stirred at room temperature for 24 hours. The precipitate was filtered off, washed with diethylether and dried at 60°C . Yield: 0.014g of compound U (diastereoisomer A) (55%, mp= 170°C).

Synthesis of compounds R and S

A solution of compound 126 of WO 2004/011436 (0.2 g, 0.3 mmol) and methyl iodide (0.0558 g, 0.3 mmol) in acetone (5 ml) was stirred at room temperature for 24 hours. The mixture was evaporated till dryness and then crystallized from diisopropylether and acetone. Yielding : 0.147 g of compound R (B) (95 %, mp= 224°C).

Compound S (diastereoisomer A) was prepared according to the above protocol but starting from compound 125 of WO 2004/011436. Yield : 0.069 g of compound S (65 %, mp= 214°C).

Synthesis of compounds M, N, O, P, Q and T

Compound M (A) was prepared according to the protocol of compound R but starting from compound 24 of WO 2004/011436. Yield: 0.025 g of compound M (A) (43 %).

Compound N (B) was prepared according to the protocol of compound R but starting from compound 37 of WO 2004/011436. Yield: 0.048 g of compound N (B) (85 %).

Compound O (B) was prepared according to the protocol of compound R but starting from compound 39 of WO 2004/011436. Yield: 0.043 g of compound O (B) (75 %).


Compound P (B) was prepared according to the protocol of compound R but starting from compound 50 of WO 2004/011436. Yield: 0.032 g of compound P (B) (56 %).

Compound Q (A) was prepared according to the protocol of compound R but starting from compound 45 of WO 2004/011436. Yield: 0.149 g of compound Q (A) (91 %).











Compound T (A) was prepared according to the protocol of compound R but starting from compound 32 of WO 2004/011436. Yield: 0.038 g of compound T (A) (66 %).

Tables 1 and 2 list compounds of formula (Ia) or (Ib) according to the present invention.

Table 1:



Comp nr.	Ex. nr.	R ⁶	R ³	L	Stereochemis try and melting points
A		H	1-naphthyl		(B); 244°C
C		H	1-naphthyl		(B); 182°C

Comp nr.	Ex. nr.	R ⁶	R ³	L	Stereochemis try and melting points
121	B5	H	1-naphthyl		(A1); 210°C
B		H	1-naphthyl		(A); 204°C
103	B5	H	1-naphthyl		(B); >250°C
E		H	phenylCH ₂ -CH ₂ -		(A); >250°C
F		H	phenylCH ₂ -CH ₂ -		(B); >250°C
102	B5	H	1-naphthyl		(A2); 210°C
57	B5	H	phenyl		(A); 244°C
10	B5	H	phenyl		(B); 198°C
M		H			(A); 268°C









Comp nr.	Ex. nr.	R ⁶	R ³	L	Stereochemis try and melting points
N		Cl	phenyl		(B); 255°C
O		H	3-fluorophenyl		(B); 184°C
P		H	phenyl		(B); 246°C
Q		H	2-naphthyl		(A)

Table 2:



Comp. nr.	R ²	R ³	q	L	Stereochemistry and melting points
G	OCH ₃	1-naphthyl	2		(B)
H	OCH ₃	2-naphthyl	3		(B); 172°C
K	OCH ₃	phenyl	3		(A); 245°C
J	OCH ₃	phenyl	3		(B); 175°C
I	OCH ₃	2-naphthyl	3		(A); 188°C
L	SCH ₃	phenyl	3		(A); 233°C
R	OCH ₃	2-naphthyl	2		(B); 224°C

Comp. nr.	R ²	R ³	q	L	Stereochemistry and melting points
S	OCH ₃	2-naphthyl	2		(A); 214°C
T	SCH ₃	phenyl	1		(A); 266°C
U	OCH ₃	phenyl	3		(A); 170°C

Analytical methods

General method

- 5 The HPLC gradient was supplied by an Alliance HT 2795 (Waters) system consisting of a quaternary pump with degasser, an autosampler, and DAD detector. Flow from the column was split to the MS detector. MS detectors were configured with an electrospray ionization source. The capillary needle voltage was 3 kV and the source temperature was maintained at 100 °C. Nitrogen was used as the nebulizer gas. Data acquisition was performed with a Waters-Micromass MassLynx-Openlynx data system.

LCMS-method 1

- In addition to the general procedure: LCMS analysis was carried out (electrospray ionization in both positive and negative (pulsed) mode scanning from 100 to 1000 amu)
- 15 on a Sunfire C18 column (Waters, Milford, MA; 3.5 µm, 4.6 x 100 mm) with a flow rate of 0.8 ml/minute. Two mobile phases (mobile phase A: 35% 6.5 mM ammonium acetate + 30 % acetonitrile +35 % formic acid (2 ml/l); mobile phase B: 100 % acetonitrile) were employed to run a gradient condition from 100 % A for 1 minute to 100 % B in 4 minutes, 100 % B at a flow rate of 1.2 ml/minute for 4 minutes to 100 % A at 0.8 ml/minute in 3 minutes, and reequilibrate with 100 % A for 1.5 minute.

The mass of compound G (without counter ion) was recorded with LCMS-method 1 (liquid chromatography mass spectrometry). The parent peak (MH⁺) is 583.

LCMS-method 2

- 5 In addition to the general procedure: Reversed phase HPLC was carried out on an Kromasil C18 column (5 μ m, 4.6 x 150 mm) with a flow rate of 1.0 ml/min. Three mobile phases (mobile phase A: 100 % 7 mM ammonium acetate; mobile phase B: 100 % acetonitrile; mobile phase C: 0.2 % formic acid + 99.8 % ultra-pure Water) were employed to run a gradient condition from 30 % A , 40% B and 30% C (hold for 10 1 minute) to 100 % B in 4 minutes, 100% B for 5 minutes and reequilibrate with initial conditions for 3 minutes. An injection volume of 5 μ l was used. Cone voltage was 20 V for positive ionization mode. Mass spectra were acquired by scanning from 100 to 900 in 0.8 seconds using an interscan delay of 0.08 seconds.
- 15 The mass of compound Q (without counter ion) was recorded with LCMS-method 2 (liquid chromatography mass spectrometry). The parent peak (MH⁺) is 569. The retention time (R_t) is 6.20.

Pharmacological examples

20 ***Preparation of bacterial suspensions for susceptibility testing:***

The bacteria used in this study were grown overnight in flasks containing 100 ml Mueller-Hinton Broth (Becton Dickinson - cat. no. 275730) in sterile de-ionized water, with shaking, at 37 °C. Stocks (0.5 ml/tube) were stored at -70 °C until use. Bacteria titrations were performed in microtiter plates and colony forming units (CFUs) were 25 determined. In general, an inoculum level of approximately 100 CFUs was used for susceptibility testing.

Anti bacterial Susceptibility testing: IC₉₀ determination

Microtitre plate assay

- 30 Flat-bottom, sterile 96-well plastic microtiter plates were filled with 180 μ l of sterile deionized water, supplemented with 0.25 % BSA. Subsequently, stock solutions (7.8 x final test concentration) of compounds were added in 45 μ l volumes in column 2. Serial five-fold dilutions (45 μ l in 180 μ l) were made directly in the microtiter plates from column 2 to reach column 11. Untreated control samples with (column 1) and without 35 (column 12) inoculum were included in each microtiter plate. Depending on the bacteria type, approximately 10 to 60 CFU per well of bacteria inoculum (100 TCID₅₀), in a volume of 100 μ l in 2.8x Mueller-Hinton broth medium, was added to

the rows A to H, except column 12. The same volume of broth medium without inoculum was added to column 12 in row A to H. The cultures were incubated at 37°C for 24 hours under a normal atmosphere (incubator with open air valve and continuous ventilation). At the end of incubation, one day after inoculation, the bacterial growth was quantitated fluorometrically. Therefore resazurin (0.6 mg/ml) was added in a volume of 20 µl to all wells 3 hours after inoculation, and the plates were re-incubated overnight. A change in colour from blue to pink indicated the growth of bacteria. The fluorescence was read in a computer-controlled fluorometer (Cytofluor Biosearch) at an excitation wavelength of 530 nm and an emission wavelength of 590 nm. The % growth inhibition achieved by the compounds was calculated according to standard methods. The IC₉₀ (expressed in µg/ml) was defined as the 90 % inhibitory concentration for bacterial growth. The results are shown in Table 3.

Agar dilution method.

MIC₉₉ values (the minimal concentration for obtaining 99 % inhibition of bacterial growth) can be determined by performing the standard Agar dilution method according to NCCLS standards* wherein the media used includes Mueller-Hinton agar.

* Clinical laboratory standard institute. 2005. Methods for dilution Antimicrobial susceptibility tests for bacteria that grows Aerobically: approved standard -sixth edition

Time kill assays

Bactericidal or bacteriostatic activity of the compounds may be determined in a time kill assay using the broth microdilution method *. In a time kill assay on *Staphylococcus aureus* and methicillin resistant *S. aureus* (MRSA), the starting inoculum of *S. aureus* and MRSA is 10⁶ CFU / ml in Muller Hinton broth. The antibacterial compounds are used at the concentration of 0.1 to 10 times the MIC (i.e. IC₉₀ as determined in microtitre plate assay). Wells receiving no antibacterial agent constitute the culture growth control. The plates containing the microorganism and the test compounds are incubated at 37 °C. After 0, 4, 24, and 48 hrs of incubation samples are removed for determination of viable counts by serial dilution (10⁻¹ to 10⁻⁶) in sterile PBS and plating (200 µl) on Mueller Hinton agar. The plates are incubated at 37 °C for 24 hrs and the number of colonies are determined. Killing curves can be constructed by plotting the log₁₀CFU per ml versus time. A bactericidal effect is commonly defined as 3-log₁₀ decrease in number of CFU per ml as compared to untreated inoculum. The potential carryover effect of the drugs is removed by serial dilutions and counting the colonies at highest dilution used for plating. No carryover effect is observed at the

dilution of 10^{-2} used for plating. This results in limit of detection 5×10^2 CFU / ml or $<2.7 \log$ CFU/ml.

* Zurenko, G.E. *et al.* In vitro activities of U-100592 and U-100766, novel oxazolidinone antibacterial agents. *Antimicrob. Agents Chemother.* **40**, 839-845 (1996).

5

Results

A time kill assay was performed with compound 102 and the control drug ciprofloxacin.

Compound 102 demonstrated bactericidal activity on *S. aureus*, as did the control
10 antibiotic ciprofloxacin. Bactericidal activities were observed at 1 and 10 times MIC90 (1 and 10 X MIC equals to 2.3 and 23 ug/ml for compound 102). At 0.1 times the MIC, the treated samples followed the control in growth.

Also for MRSA, compound 12 demonstrated marked bactericidal activity as compared to ciprofloxacin for which these strains have developed resistance. MRSA is resistant
15 not only to methicillin but also to flouroquinolones like ciprofloxacin and as such no bactericidal effect was observed using this drug. On MRSA at 24 hours compound 12 was mostly bacteriostatic but after 48 hours it showed marked reduction in viable counts.

20 ***Determination of cellular ATP levels***

In order to analyse the change in the total cellular ATP concentration (using ATP bioluminescence Kit, Roche), assays are carried out by growing a culture of *S. aureus* (ATCC29213) stock in 100 ml Mueller Hinton flasks and incubate in a shaker-incubator for 24 hrs at 37 °C (300 rpm). Measure OD₄₀₅ nm and calculate the CFU/ml.
25 Dilute the cultures to 1×10^6 CFU/ml (final concentration for ATP measurement: 1×10^5 CFU/100 µl per well) and add test compound at 0.1 to 10 times the MIC (i.e. IC₉₀ as determined in microtitre plate assay). Incubate these tubes for 0, 30 and 60 minutes at 300 rpm and 37°C. Use 0.6 ml bacterial suspension from the snap-cap tubes and add to a new 2 ml eppendorf tubes. Add 0.6 ml cell lysis reagent (Roche kit), vortex at
30 max speed and incubate for 5 minutes at room temperature. Cool on ice. Let the luminometer warm up to 30°C (Luminoskan Ascent Labsystems with injector). Fill one column (= 6 wells) with 100 µl of the same sample. Add 100 µl Luciferase reagent to each well by using the injector system. Measure the luminescence for 1 sec.

Table 3 : IC₉₀ values (µg/ml) determined according to the Microtitre plate assay.

Comp. No.	IC90 (µg/ml)													
	BSU 43639	ECO 25922	ECO 35218	EFA 14506	EFA 29212	LMO 49594	PAE 27853	SMU 33402	SPN 6305	SPY 8668	STA 43300	STA 25923	STA 29213	STA RMETH
A													0.4	
B													0.4	
121	2.8	12.4		2.8	2.8	0.5	13.9	2.8	3.5	3.5	3.5	2.2	0.4	2.2
C													0.4	
G													0.4	
H													0.5	
103	2.8	11.1	17.5	2.8	2.8	0.5	13.9	2.8	3.5	2.8	2.8	2.2	2.0	2.2
E													2.1	
F													2.1	
K													2.1	
J													2.1	
102	2.8	13.9		2.8	2.8	2.2	13.9	2.5	3.1	2.5	3.5	2.2	2.8	2.8
I													2.9	
57	2.6			12.9	12.9	2.6		12.9	16.3	12.9	12.9	12.9	12.9	12.9
10	10.3			12.9	5.8	1.2		10.3	16.3	1.3	5.8	11.5	14.5	8.2
L	2.2		11		2.2	2.2	2.2	0.4	2.2	0.4			2.2	
M									8.34	8.34			1.7	
N			17.55		1.75		9.87		1.75	1.75			1.8	
O					8.53				1.7	1.7			1.9	
P					10.84		10.84		2.16	2.16			2.2	
Q					1.8		4.53		1.8	1.8			1.8	
R			9.27		1.17		2.07		0.46	0.74			0.4	
S					1.85		1.85		0.47	1.85			0.4	
U					1.86		9.33		2.09	1.86			1.9	
T					3.8		8.5		1.7	1.7			1.7	



BSU 43639 means *Bacillus subtilis* (ATCC43639); ECO 25922 means *Escherichia coli*

- 5 (ATCC25922); ECO 35218 means *Escherichia coli* (ATCC35218); EFA 14506 means *Enterococcus faecalis* (ATCC14506); EFA 29212 means *Enterococcus faecalis* (ATCC29212); LMO 49594 means *Listeria monocytogenes* (ATCC49594); PAE 27853 means *Pseudomonas aeruginosa* (ATCC27853); SMU 33402 means *Streptococcus mutans* (ATCC33402); SPN 6305 means *Streptococcus pneumoniae* (ATCC6305);
- 10 SPY 8668 means *Streptococcus pyogenes* (ATCC8668); STA 43300 means *Staphylococcus aureus* (ATCC43300); STA 25923 means *Staphylococcus aureus* (ATCC25923); STA 29213 means *Staphylococcus aureus* (ATCC29213); STA RMETH means methicilline resistant *Staphylococcus aureus* (MRSA) (a clinical isolate from the University of Antwerp).

- 15 ATCC means American type tissue culture.

Claims

1. Use of a compound for the manufacture of a medicament for the treatment of a bacterial infection, said compound being a compound of formula (Ia) or (Ib)



a *N*-oxide thereof, a tautomeric form thereof or a stereochemically isomeric form thereof wherein


A^- is a pharmaceutically acceptable counter ion;

R^1 is hydrogen, halo, haloalkyl, cyano, hydroxy, Ar, Het, alkyl, alkyloxy, alkylthio, alkyloxyalkyl, alkylthioalkyl, Ar-alkyl or di(Ar)alkyl ;

p is an integer equal to 1, 2, 3 or 4 ;

R^2 is hydrogen, hydroxy, mercapto, alkyloxy, alkyloxyalkyloxy, alkylthio,

mono or di(alkyl)amino or a radical of formula



wherein Y is

R^3 is alkyl, Ar, Ar-alkyl, Het or Het-alkyl;

q is an integer equal to zero, 1, 2, 3 or 4 ;

R^4 and R^5 each independently are hydrogen, alkyl or benzyl;

R^4 and R^5 together and including the N to which they are attached may form a radical selected from the group of pyrrolidinyl, 2-pyrrolinyl, 3-pyrrolinyl, pyrrolyl, imidazolidinyl, pyrazolidinyl, 2-imidazolyl, 2-pyrazolyl,

- imidazolyl, pyrazolyl, triazolyl, piperidinyl, pyridinyl, piperazinyl, pyridazinyl, pyrimidinyl, pyrazinyl, triazinyl, morpholinyl and thiomorpholinyl, each of said rings may optionally be substituted with alkyl, halo, haloalkyl, hydroxy, alkyloxy, amino, mono- or dialkylamino, alkylthio, alkyloxyalkyl, alkylthioalkyl, Ar-alkyl or pyrimidinyl;
- 5 R^6 is hydrogen, halo, haloalkyl, hydroxy, Ar, alkyl, alkyloxy, alkylthio, alkyloxyalkyl, alkylthioalkyl, Ar-alkyl or di(Ar)alkyl ; or two vicinal R^6 radicals may be taken together to form a bivalent radical of formula $-CH=CH-CH=CH-$;
- 10 r is an integer equal to 1, 2, 3, 4 or 5 ;
 R^7 is hydrogen, alkyl, Ar or Het ;
 R^8 is hydrogen or alkyl ;
 R^9 is oxo ; or
 R^8 and R^9 together form the radical $-CH=CH-N=$;
- 15 R^{10} is alkyl, alkylcarbonyl, Ar, Ar-alkyl, Ar-carbonyl, Het¹-alkyl or Het¹-carbonyl;
- alkyl is a straight or branched saturated hydrocarbon radical having from 1 to 6 carbon atoms ; or is a cyclic saturated hydrocarbon radical having from 3 to 6 carbon atoms ; or is a cyclic saturated hydrocarbon radical having from 3 to 6 carbon atoms attached to a straight or branched saturated hydrocarbon radical having from 1 to 6 carbon atoms ; wherein each carbon atom can be optionally substituted with hydroxy, alkyloxy or oxo ;
- 20 Ar is a homocycle selected from the group of phenyl, naphthyl, acenaphthyl, tetrahydronaphthyl, each homocycle optionally substituted with 1, 2 or 3 substituents, each substituent independently selected from the group of hydroxy, halo, cyano, nitro, amino, mono- or dialkylamino, alkyl, haloalkyl, alkyloxy, haloalkyloxy, carboxyl, alkyloxycarbonyl, aminocarbonyl, morpholinyl and mono- or dialkylaminocarbonyl ;
- 25 Het is a monocyclic heterocycle selected from the group of *N*-phenoxypiperidinyl, piperidinyl, pyrrolyl, pyrazolyl, imidazolyl, furanyl, thienyl, oxazolyl, isoxazolyl, thiazolyl, isothiazolyl, pyridinyl, pyrimidinyl, pyrazinyl and pyridazinyl; or a bicyclic heterocycle selected from the group of quinolinyl, quinoxalinyl, indolyl, benzimidazolyl, benzoxazolyl, benzisoxazolyl, benzothiazolyl, benzisothiazolyl, benzofuranyl, benzothieryl,
- 30 2,3-dihydrobenzo[1,4]dioxinyl and benzo[1,3]dioxolyl ; each monocyclic and bicyclic heterocycle may optionally be substituted with 1, 2 or 3 substituents,
- 35

each substituent independently selected from the group of halo, hydroxy, alkyl, alkyloxy, and Ar-carbonyl;



Het¹ is a monocyclic heterocycle selected from furanyl or thienyl; or a bicyclic heterocycle selected from benzofuranyl or benzothieryl; each monocyclic and bicyclic heterocycle may optionally be substituted with 1, 2 or 3 substituents,

halo is a substituent selected from the group of fluoro, chloro, bromo and iodo; and

haloalkyl is a straight or branched saturated hydrocarbon radical having from 1 to 6 carbon atoms or a cyclic saturated hydrocarbon radical having from 3 to 6 carbon atoms or a cyclic saturated hydrocarbon radical having from 3 to 6 carbon atoms attached to a straight or branched saturated hydrocarbon radical having from 1 to 6 carbon atoms; wherein one or more carbon atoms are substituted with one or more halo atoms;



provided that the bacterial infection is other than a Mycobacterial infection.

2. Use according to claim 1 wherein the compound of formula (Ia) or (Ib) is a compound having the following formula





a *N*-oxide thereof, a tautomeric form thereof or a stereochemically isomeric form thereof.

3. Use according to claim 1 wherein the compound of formula (Ia) or (Ib) is a compound having the following formula



a *N*-oxide thereof, a tautomeric form thereof or a stereochemically isomeric form thereof.

- 5 4. Use according to claim 1 wherein the compound of formula (Ia) or (Ib) is a compound having the following formula



a *N*-oxide thereof, a tautomeric form thereof or a stereochemically isomeric form thereof.

5. Use according to any one of the preceding claims wherein R^1 is halo.

5

6. Use according to any one of the preceding claims wherein p is equal to 1.

7. Use according to any one of the preceding claims wherein R^2 is alkyloxy or alkylthio.

10

8. Use according to claim 7 wherein R^2 is C_{1-4} alkyloxy.

9. Use according to any one of the preceding claims wherein R^3 is Het, Ar or Ar-alkyl.

15 10. Use according to claim 9 wherein R^3 is Ar or Ar-alkyl.

11. Use according to claim 9 wherein R^3 is thienyl, naphthyl, phenyl, naphthyl C_{1-4} alkyl or phenyl C_{1-4} alkyl.

20 12. Use according to claim 11 wherein R^3 is naphthyl, phenyl or phenyl C_{1-4} alkyl.

13. Use according to any one of the preceding claims wherein R^4 and R^5 are C_{1-4} alkyl or R^4 and R^5 together and including the N to which they are attached may form a radical selected from imidazolyl or piperidinyl.

25

14. Use according to claim 13 wherein R^4 and R^5 are C_{1-4} alkyl.

15. Use according to any one of the preceding claims wherein R^6 is hydrogen or halo.

30 16. Use according to any one of the preceding claims wherein r is equal to 1.

17. Use according to any one of the preceding claims wherein R^7 is hydrogen.

18. Use according to any one of the preceding claims wherein R^{10} is alkyl.

35

19. Use according to claim 18 wherein R^{10} is C_{1-6} alkyl.

20. Use according to any one of the preceding claims wherein A^- is iodo.


21. Use according to any one of the preceding claims wherein the compound is a compound according to formula (Ia).

5

22. Use of a compound of formula (Ia) according to claim 1 wherein R^1 is halo; $p = 1$; R^2 is alkyloxy or alkylthio; R^3 is naphthyl, phenyl, phenylethyl or thienyl; $q = 1, 2$ or 3 ; R^4 and R^5 are C_{1-4} alkyl or R^4 and R^5 together and including the N to which they are attached may form a radical selected from imidazolyl or piperidiny; R^6 is hydrogen or halo; r is equal to 1; R^7 is hydrogen; R^{10} is C_{1-6} alkyl; A^- is iodo.

10



23. Use according to claim 1 wherein the compound is selected from the following compounds




a *N*-oxide thereof or a stereochemically isomeric form thereof.

24. Use according to any one of the preceding claims wherein the bacterial infection is an infection with a gram-positive bacterium.

25. A compound of formula (Ia) or (Ib)




- 5 a *N*-oxide thereof, a tautomeric form thereof or a stereochemically isomeric form thereof wherein R^1 , p , R^2 , R^3 , q , R^4 , R^5 , R^6 , r , R^7 , R^8 , R^9 , R^{10} and A^- are defined as in claim 1;
provided that when R^{10} is alkyl or benzyl, then R^4 and R^5 are other than hydrogen; and
provided that the compound is other than




- 10 a *N*-oxide thereof, a tautomeric form thereof or a stereochemically isomeric form thereof.

26. A compound according to claim 25 wherein the compound is selected from:



a *N*-oxide thereof or a stereochemically isomeric form thereof.

27. A compound according to claim 26 wherein the compound is selected from :



5 or a stereochemically isomeric form thereof.

28. A combination of (a) a compound of formula (Ia) or (Ib) as defined in any one of the preceding claims, and (b) one or more other antibacterial agents provided that the one or more other antibacterial agents are other than antimycobacterial agents.

- 5 29. A pharmaceutical composition comprising a pharmaceutically acceptable carrier and, as active ingredient, a therapeutically effective amount of (a) a compound of formula (Ia) or (Ib) as defined in any one of claims 1 to 27, and (b) one or more other antibacterial agents provided that the one or more other antibacterial agents are other than antimycobacterial agents.

10

30. The use of a combination as claimed in claim 28 or a pharmaceutical composition as claimed in claim 29 for the manufacture of a medicament for the treatment of a bacterial infection.

- 15 31. A product containing (a) a compound of formula (Ia) or (Ib) as defined in any one of claims 1 to 27, and (b) one or more other antibacterial agents provided that the one or more other antibacterial agents are other than antimycobacterial agents, as a combined preparation for simultaneous, separate or sequential use in the treatment of a bacterial infection.

20