

5 **SUBSTITUTED-3-SULFONYLINDAZOLE DERIVATIVES AS**
 5-HYDROXYTRYPTAMINE-6 LIGANDS

BACKGROUND OF THE INVENTION

10 Serotonin (5-hydroxytryptamine) (5-HT) receptors play a critical role in many
physiological and behavioral functions in humans and animals. These functions are
mediated through various 5-HT receptors distributed throughout the body. There are
now approximately fifteen different human 5-HT receptor subtypes that have been
cloned, many with well-defined roles in humans. One of the most recently identified
15 5-HT receptor subtypes is the 5-HT6 receptor, first cloned from rat tissue in 1993
(Monsma, F. J.; Shen, Y.; Ward, R. P.; Hamblin, M. W. *Molecular Pharmacology*
1993, 43, 320-327) and subsequently from human tissue (Kohen, R.; Metcalf, M. A.;
Khan, N.; Druck, T.; Huebner, K.; Sibley, D. R. *Journal of Neurochemistry* 1996, 66,
47-56). The receptor is a G-protein coupled receptor (GPCR) positively coupled to
20 adenylyate cyclase (Ruat, M.; Traiffort, E.; Arrang, J-M.; Tardivel-Lacombe, L.; Diaz,
L.; Leurs, R.; Schwartz, J-C. *Biochemical Biophysical Research Communications*
1993, 193, 268-276). The receptor is found almost exclusively in the central nervous
system (CNS) areas both in rat and in human. *In situ* hybridization studies of the
5-HT6 receptor in rat brain using mRNA indicate principal localization in the areas of
25 5-HT projection including striatum, nucleus accumbens, olfactory tubercle, and
hippocampal formation (Ward, R. P.; Hamblin, M. W.; Lachowicz, J. E.; Hoffman, B.
J.; Sibley, D. R.; Dorsa, D. M. *Neuroscience* 1995, 64, 1105-1111).

 There are many potential therapeutic uses for 5-HT6 ligands in humans
based on direct effects and on indications from available scientific studies. These
30 studies provided information including the localization of the receptor, the affinity of
ligands with known *in vivo* activity, and results obtained from various animal studies
conducted so far (Woolley, M. L.; Marsden, C. A.; Fone, K. C. F. *Current Drug*
Targets: CNS & Neurological Disorders 2004, 3(1), 59-79).

One therapeutic use of modulators of 5-HT₆ receptor function is in the enhancement of cognition and memory in human diseases such as Alzheimer's. The high levels of receptor found in important structures in the forebrain, including the caudate/putamen, hippocampus, nucleus accumbens, and cortex indicate a role for the receptor in memory and cognition since these areas are known to play a vital role in memory (Gerard, C.; Martres, M.-P.; Lefevre, K.; Miquel, M.C.; Verge, D.; Lanfumey, R.; Doucet, E.; Hamon, M.; El Mestikawy, S. *Brain Research*, 1997, 746, 207-219). The ability of known 5-HT₆ receptor ligands to enhance cholinergic transmission also supported the cognition use (Bentley, J. C.; Boursson, A.; Boess, F. G.; Kone, F. C.; Marsden, C. A.; Petit, N.; Sleight, A. J. *British Journal of Pharmacology*, 1999, 126(7), 1537-1542). Studies have demonstrated that a known 5-HT₆ selective antagonist significantly increased glutamate and aspartate levels in the frontal cortex without elevating levels of noradrenaline, dopamine, or 5-HT. This selective elevation of neurochemicals known to be involved in memory and cognition indicates the role 5-HT₆ ligands play in cognition (Dawson, L. A.; Nguyen, H. Q.; Li, P. *British Journal of Pharmacology*, 2000, 130(1), 23-26). Animal studies of memory and learning with a known selective 5-HT₆ antagonist found positive effects (Rogers, D. C.; Hatcher, P. D.; Hagan, J. J. *Society of Neuroscience, Abstracts* 2000, 26, 680). More recent studies have supported this finding in several additional animal models of cognition and memory including in a novel object discrimination model (King, M. V.; Sleight, A. J.; Wooley, M. L.; Topham, I. A.; Marsden, C. A.; Fone, K. C. *Neuropharmacology* 2004, 47(2), 195-204 and Wooley, M. L.; Marsden, C. A.; Sleight, A. J.; Fone, K. C. *Psychopharmacology*, 2003, 170(4), 358-367) and in a water maze model (Rogers, D. C.; Hagan, J. J. *Psychopharmacology*, 2001, 158(2), 114-119 and Foley, A. G.; Murphy, K. J.; Hirst, W. D.; Gallagher, H. C.; Hagan, J. J.; Upton, N.; Walsh, F. S.; Regan, C. M. *Neuropsychopharmacology* 2004, 29(1), 93-100).

A related therapeutic use for 5-HT₆ ligands is the treatment of attention deficit disorders (ADD, also known as Attention Deficit Hyperactivity Disorder or ADHD) in both children and adults. Because 5-HT₆ antagonists enhance the activity of the nigrostriatal dopamine pathway and because ADHD has been linked to abnormalities in the caudate (Ernst, M.; Zametkin, A. J.; Matochik, J. H.; Jons, P. A.; Cohen, R. M.

Journal of Neuroscience 1998, 18(15), 5901-5907), 5-HT₆ antagonists attenuate attention deficit disorders.

Early studies examining the affinity of various CNS ligands with known therapeutic utility or a strong structural resemblance to known drugs implicates 5-HT₆ ligands in the treatment of schizophrenia and depression. For example, clozapine (an effective clinical antipsychotic) has high affinity for the 5-HT₆ receptor subtype. Also, several clinical antidepressants have high affinity for the receptor as well and act as antagonists at this site (Branchek, T. A.; Blackburn, T. P. *Annual Reviews in Pharmacology and Toxicology* 2000, 40, 319-334).

Further, recent *in vivo* studies in rats indicate that 5-HT₆ modulators are useful in the treatment of movement disorders including epilepsy (Stean, T.; Routledge, C.; Upton, N. *British Journal of Pharmacology* 1999, 127 Proc. Supplement 131P and Routledge, C.; Bromidge, S. M.; Moss, S. F.; Price, G. W.; Hirst, W.; Newman, H.; Riley, G.; Gager, T.; Stean, T.; Upton, N.; Clarke, S. E.; Brown, A. M. *British Journal of Pharmacology* 2000, 130(7), 1606-1612).


Therefore, it is an object of this invention to provide compounds which are useful as therapeutic agents in the treatment of a variety of central nervous system disorders related to or affected by the 5-HT₆ receptor.

It is another object of this invention to provide therapeutic methods and pharmaceutical compositions useful for the treatment of central nervous system disorders related to or affected by the 5-HT₆ receptor.

It is a feature of this invention that the compounds provided may also be used to further study and elucidate the 5-HT₆ receptor.

SUMMARY OF THE INVENTION

The present invention provides a 3-sulfonylindazole compound of formula I



(I)

wherein

X is O, S, NR, CH₂, CH₂Y, CH₂Z, CO, CONR or NRCO;

Y is O, S or NR;

Z is CO;

5 n is 0 or an integer of 1, 2, 3, 4, 5 or 6 when X is CH₂;

n is an integer of 1, 2, 3, 4, 5 or 6 when X is CH₂Z, CO or NRCO;

n is an integer of 2, 3, 4, 5 or 6 when X is O, S, NR, CH₂Y or CONR;

R is H or an optionally substituted alkyl group;

R₁ is H or an alkyl, cycloalkyl, aryl or heteroaryl group each optionally substituted;

10 R₂ is an optionally substituted alkyl, cycloalkyl, aryl or heteroaryl group or an optionally substituted 8- to 13-membered bicyclic or tricyclic ring system having a N atom at the bridgehead and optionally containing 1, 2 or 3 additional heteroatoms selected from N, O or S;

R₃ and R₄ are each independently H, or an optionally substituted alkyl group;

15 R₅ and R₆ are each independently H, or an alkyl, alkenyl, alkynyl, cycloalkyl, cycloheteroalkyl, aryl or heteroaryl group each optionally substituted, or R₅ and R₆ may be taken together with the atom to which they are attached to form an optionally substituted 3- to 7-membered ring optionally containing an additional heteroatom selected from O, N or S;

20 R₇ is H, halogen, CN, OR₈, CO₂R₉, CONR₁₀R₁₁, or an alkyl, alkenyl, alkynyl, cycloalkyl, cycloheteroalkyl, aryl or heteroaryl group each optionally substituted;

m is an integer of 1, 2 or 3;

25 R₈ is H, COR₁₂ or an alkyl, alkenyl, alkynyl, aryl or heteroaryl group each optionally substituted;

R₉ is H or a C₁-C₆alkyl, aryl or heteroaryl group each optionally substituted;

R₁₀ and R₁₁ are each independently H or an optionally substituted alkyl group; and

30 R₁₂ is an optionally substituted C₁-C₆alkyl, cycloalkyl, cycloheteroalkyl, aryl or heteroaryl group; or

a stereoisomer thereof or a pharmaceutically acceptable salt thereof.


The present invention also provides methods and compositions useful for the therapeutic treatment of central nervous system disorders related to or affected by the 5-HT6 receptor.

DETAILED DESCRIPTION OF THE INVENTION

The 5-hydroxytryptamine-6 (5-HT6) receptor has been identified by molecular cloning. Its ability to bind a wide range of therapeutic compounds used in psychiatry, coupled with its intriguing distribution in the brain has stimulated significant interest in new compounds which are capable of interacting with or affecting said receptor.

Significant efforts are being made to understand the role of the 5-HT6 receptor in psychiatry, cognitive dysfunction, motor function and control, memory, mood and the like. To that end, compounds which demonstrate a binding affinity for the 5-HT6 receptor are earnestly sought both as an aid in the study of the 5-HT6 receptor and as potential therapeutic agents in the treatment of central nervous system disorders, for example see C. Reavill and D. C. Rogers, *Current Opinion in Investigational Drugs*, 2001, 2(1):104-109, Pharma Press Ltd and Woolley, M. L.; Marsden, C. A.; Fone, K. C. F. *Current Drug Targets: CNS & Neurological Disorders* 2004, 3(1), 59-79.

Surprisingly, it has now been found that 3-sulfonylindazole compounds of formula I demonstrate 5-HT6 receptor affinity along with significant receptor sub-type selectivity. Advantageously, said formula I compounds are effective therapeutic agents for the treatment of central nervous system (CNS) disorders associated with or affected by the 5-HT6 receptor. Accordingly, the present invention provides a 3-sulfonylindazole compound of formula I



(I)

wherein

X is O, S, NR, CH₂, CH₂Y, CH₂Z, CO, CONR or NRCO;

Y is O, S or NR;

Z is CO;

n is 0 or an integer of 1, 2, 3, 4, 5 or 6 when X is CH₂;

n is an integer of 1, 2, 3, 4, 5 or 6 when X is CH₂Z, CO or NRCO;

5 n is an integer of 2, 3, 4, 5 or 6 when X is O, S, NR, CH₂Y or CONR;

R is H or an optionally substituted alkyl group;

R₁ is H or an alkyl, cycloalkyl, aryl or heteroaryl group each optionally substituted;

10 R₂ is an optionally substituted alkyl, cycloalkyl, aryl or heteroaryl group or an optionally substituted 8- to 13-membered bicyclic or tricyclic ring system having a N atom at the bridgehead and optionally containing 1, 2 or 3 additional heteroatoms selected from N, O or S;

R₃ and R₄ are each independently H, or an optionally substituted alkyl group;

15 R₅ and R₆ are each independently H, or an alkyl, alkenyl, alkynyl, cycloalkyl, cycloheteroalkyl, aryl or heteroaryl group each optionally substituted, or R₅ and R₆ may be taken together with the atom to which they are attached to form an optionally substituted 3- to 7-membered ring optionally containing an additional heteroatom selected from O, N or S;

20 R₇ is H, halogen, CN, OR₈, CO₂R₉, CONR₁₀R₁₁, or an alkyl, alkenyl, alkynyl, cycloalkyl, cycloheteroalkyl, aryl or heteroaryl group each optionally substituted;

m is an integer of 1, 2 or 3;

R₈ is H, COR₁₂ or an alkyl, alkenyl, alkynyl, aryl or heteroaryl group each optionally substituted;

R₉ is H or a C₁-C₆alkyl, aryl or heteroaryl group each optionally substituted;

25 R₁₀ and R₁₁ are each independently H or an optionally substituted alkyl group; and

R₁₂ is an optionally substituted C₁-C₆alkyl, cycloalkyl, cycloheteroalkyl, aryl or heteroaryl group; or

a stereoisomer thereof or a pharmaceutically acceptable salt thereof.

30 It is understood that the claims encompass all possible stereoisomers and prodrugs. Moreover, unless stated otherwise, each alkyl, alkenyl, alkynyl, cycloalkyl, cycloheteroalkyl, aryl or heteroaryl group is contemplated as being optionally substituted.

An optionally substituted moiety may be substituted with one or more substituents. The substituent groups, which are optionally present, may be one or more of those customarily employed in the development of pharmaceutical compounds or the modification of such compounds to influence their

5 structure/activity, persistence, absorption, stability or other beneficial property. Specific examples of such substituents include halogen atoms, nitro, cyano, thiocyanato, cyanato, hydroxyl, alkyl, haloalkyl, alkoxy, haloalkoxy, amino, alkylamino, dialkylamino, formyl, alkoxycarbonyl, carboxyl, alkanoyl, alkylthio, alkylsulphinyl, alkylsulphonyl, carbamoyl, alkylamido, phenyl, phenoxy, benzyl,
10 benzyloxy, heterocyclyl or cycloalkyl groups, preferably halogen atoms or lower alkyl or lower alkoxy groups. Unless otherwise specified, typically, 0-4 substituents may be present. When any of the foregoing substituents represents or contains an alkyl substituent group, this may be linear or branched and may contain up to 12 carbon atoms, preferably up to 6 carbon atoms, more preferably up to 4 carbon atoms.

15 As used herein, the term "alkyl" includes both (C_1 - C_{10}) straight chain and (C_3 - C_{12}) branched-chain (unless defined otherwise) monovalent saturated hydrocarbon moiety. Examples of saturated hydrocarbon alkyl moieties include, but are not limited to, chemical groups such as methyl, ethyl, *n*-propyl, isopropyl, *n*-butyl, *tert*-butyl, isobutyl, *sec*-butyl; higher homologs such as *n*-pentyl, *n*-hexyl, and the like.
20 Specifically included within the definition of "alkyl" are those alkyl groups that are optionally substituted. Suitable alkyl substitutions include, but are not limited to, CN, OH, $NR_{10}R_{11}$, halogen, phenyl, carbamoyl, carbonyl, alkoxy or aryloxy.

As used herein the term "haloalkyl" designates a C_nH_{2n+1} group having from one to $2n+1$ halogen atoms which may be the same or different. Examples of
25 haloalkyl groups include CF_3 , CH_2Cl , C_2H_3BrCl , $C_3H_5F_2$, or the like.


The term "halogen", as used herein, designates fluorine, chlorine, bromine, and iodine.

The term "alkenyl", as used herein, refers to either a (C_2 - C_8) straight chain or (C_3 - C_{10}) branched-chain monovalent hydrocarbon moiety containing at least one
30 double bond. Such hydrocarbon alkenyl moieties may be mono or polyunsaturated, and may exist in the E or Z configurations. The compounds of this invention are meant to include all possible E and Z configurations. Examples of mono or polyunsaturated hydrocarbon alkenyl moieties include, but are not limited to,

chemical groups such as vinyl, 2-propenyl, isopropenyl, crotyl, 2-isopentenyl, butadienyl, 2-(butadienyl), 2,4-pentadienyl, 3-(1,4-pentadienyl), and higher homologs, isomers, or the like.

The term "cycloalkyl", as used herein, refers to a monocyclic, bicyclic, tricyclic, fused, bridged, or spiro monovalent saturated hydrocarbon moiety of 3-10 carbon atoms, unless otherwise specified, wherein the carbon atoms are located inside or outside of the ring system. Any suitable ring position of the cycloalkyl moiety may be covalently linked to the defined chemical structure. Examples of cycloalkyl moieties include, but are not limited to, chemical groups such as cyclopropyl, cyclopropylmethyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclohexylmethyl, cyclohexylethyl, cycloheptyl, norbornyl, adamantyl, spiro[4.5]decanyl, and homologs, isomers, or the like.

The term "cycloheteroalkyl" as used herein designates a C₅-C₇cycloalkyl ring system containing 1, 2 or 3 heteroatoms, which may be the same or different, selected from N, O or S and optionally containing one double bond. Exemplary of the cycloheteroalkyl ring systems included in the term as designated herein are the following rings wherein X₁ is NR', O or S and R is H or an optional substituent as defined hereinbelow.




20

The term "aryl", as used herein, refers to an aromatic carbocyclic moiety of up to 20 carbon atoms, which may be a single ring (monocyclic) or multiple rings (bicyclic, up to three rings) fused together or linked covalently. Any suitable ring position of the aryl moiety may be covalently linked to the defined chemical structure. Examples of aryl moieties include, but are not limited to, chemical groups such as phenyl, 1-naphthyl, 2-naphthyl, biphenyl, anthryl, phenanthryl, fluorenyl, indanyl, biphenylenyl, acenaphthenyl, acenaphthylenyl, and the like.

The term "heteroaryl" as used herein designates an aromatic heterocyclic ring system, which may be a single ring (monocyclic) or multiple rings (bicyclic, up to three rings) fused together or linked covalently. Preferably, heteroaryl is a 5- to 6-membered ring. The rings may contain from one to four hetero atoms selected from nitrogen, oxygen, or sulfur, wherein the nitrogen or sulfur atom(s) are optionally oxidized, or the nitrogen atom(s) are optionally quarternized. Any suitable ring position of the heteroaryl moiety may be covalently linked to the defined chemical structure. Examples of heteroaryl moieties include, but are not limited to, heterocycles such as furan, thiophene, pyrrole, pyrazole, imidazole, oxazole, isoxazole, thiazole, isothiazole, oxadiazole, triazole, pyridine, pyrimidine, pyrazine, pyridazine, benzimidazole, benzoxazole, benzisoxazole, benzothiazole, benzofuran, benzothiophene, thianthrene, dibenzofuran, dibenzothiophene, indole, indazole, quinoline, isoquinoline, quinazoline, quinoxaline, purine, or the like.

Exemplary of the 8- to 13-membered bicyclic or tricyclic ring systems having a N atom at the bridgehead and optionally containing 1, 2 or 3 additional heteroatoms selected from N, O or S included in the term as designated herein are the following ring systems wherein W is NR', O or S; and R' is H or an optional substituent as described hereinbelow:



The compounds of this invention are limited to those that are chemically feasible and stable. Therefore, a combination of substituents or variables in the compounds described above is permissible only if such a combination results in a stable or chemically feasible compound.

Unless otherwise stated, structures depicted herein are also meant to include all stereochemical forms of the structure; i.e., the R and S configurations for each asymmetric center. Therefore, single stereochemical isomers as well as enantiomeric and diastereomeric mixtures of the present compounds are within the scope of the invention. Unless otherwise stated, structures depicted herein are also meant to include compounds which differ only in the presence of one or more isotopically enriched atoms. For example, compounds having the present structures except for the replacement of a hydrogen by a deuterium or tritium, or the replacement of a carbon by a ^{13}C - or ^{14}C -enriched carbon are within the scope of this invention.

The compounds of the present invention may be converted to salts, in particular pharmaceutically acceptable salts using art recognized procedures. Suitable salts with bases are, for example, metal salts, such as alkali metal or alkaline earth metal salts, for example sodium, potassium or magnesium salts, or salts with ammonia or an organic amine, such as morpholine, thiomorpholine, piperidine, pyrrolidine, a mono-, di- or tri-lower alkylamine, for example ethyl-tert-butyl-, diethyl-, diisopropyl-, triethyl-, tributyl- or dimethylpropylamine, or a mono-, di-, or trihydroxy lower alkylamine, for example mono-, di- or triethanolamine. Internal salts may furthermore be formed. Salts which are unsuitable for pharmaceutical uses but which can be employed, for example, for the isolation or purification of free compounds or their pharmaceutically acceptable salts, are also included. The term "pharmaceutically acceptable salt", as used herein, refers to salts derived from organic and inorganic acids such as, for example, acetic, propionic, lactic, citric, tartaric, succinic, fumaric, maleic, malonic, mandelic, malic, phthalic, hydrochloric, hydrobromic, phosphoric, nitric, sulfuric, methanesulfonic, naphthalenesulfonic, benzenesulfonic, toluenesulfonic, camphorsulfonic, and similarly known acceptable acids when a compound of this invention contains a basic moiety. Salts may also be formed from organic and inorganic bases, preferably alkali metal salts, for example, sodium, lithium, or potassium, when a compound of this invention contains a carboxylate or phenolic moiety, or similar moiety capable of forming base addition salts.

Compounds of the invention include esters, carbamates or other conventional prodrug forms, which in general, are functional derivatives of the compounds of the invention and which are readily converted to the inventive active moiety *in vivo*. Correspondingly, the method of the invention embraces the treatment of the various conditions described hereinabove with a compound of formula I or with a compound which is not specifically disclosed but which, upon administration, converts to a compound of formula I *in vivo*. Also included are metabolites of the compounds of the present invention defined as active species produced upon introduction of these compounds into a biological system.

Preferred compounds of the invention are those compounds of formula I wherein X is O, NR or CH₂.

Another group of preferred compounds is those formula I compounds wherein n is 2 or 3.

Also preferred are those formula I compounds wherein R₂ is an optionally substituted aryl or heteroaryl group or an optionally substituted 8- to 13-membered bicyclic or tricyclic ring system having a N atom at the bridgehead and optionally containing 1, 2 or 3 additional heteroatoms selected from N, O or S.

More preferred compounds of the invention are those compounds of formula I wherein X is O and R₅ and R₆ are each independently H or C₁-C₄ alkyl. Another group of more preferred compounds is those compounds of formula I wherein X is O and n is 3. A further group of more preferred compounds are those compounds of formula I wherein X is O; n is 3 and R₂ is naphthyl.

Among the preferred compounds of the invention are:

- N, N-Dimethyl-3-[[3-(1-naphthylsulfonyl)-1H-indazol-5-yl]oxy]propan-1-amine;
- N-Methyl-N-{2-[3-(phenylsulfonyl)-1H-indazol-5-yl]ethyl}amine;
- N,N-Dimethyl-N-{2-[3-(phenylsulfonyl)-1H-indazol-5-yl]ethyl}amine;
- 5 {2-[3-(Phenylsulfonyl)-1H-indazol-7-yl]ethyl}amine;
- N,N-Dimethyl-N-{2-[3-(phenylsulfonyl)-1H-indazol-7-yl]ethyl}amine;
- N-{2-[3-(Phenylsulfonyl)-1H-indazol-7-yl]ethyl}cyclopropanamine;
- N,N-Dimethyl-N-{3-[3-(phenylsulfonyl)-1H-indazol-5-yl]propyl}amine;
- N-{3-[3-(Phenylsulfonyl)-1H-indazol-5-yl]propyl}cyclopropanamine;
- 10 {3-[3-(Phenylsulfonyl)-1H-indazol-5-yl]propyl}amine;
- {4-[3-(Phenylsulfonyl)-1H-indazol-5-yl]butyl}amine;
- N-Methyl-N-{2-[3-(phenylsulfonyl)-1H-indazol-7-yl]ethyl}amine;
- N-[3-(1-Naphthylsulfonyl)-1H-indazol-5-yl]ethane-1,2-diamine;
- N,N-Dimethyl-2-[[3-(phenylsulfonyl)-1H-indazol-5-yl]oxy]ethanamine;
- 15 3-(Phenylsulfonyl)-5-(2-piperidin-1-ylethoxy)-1H-indazole;
- 3-(1-Naphthylsulfonyl)-5-(2-pyrrolidin-1-ylethoxy)-1H-indazole;
- N,N-Dimethyl-2-[[3-(1-naphthylsulfonyl)-1H-indazol-5-yl]oxy]ethanamine;
- N-(2-[[3-(1-Naphthylsulfonyl)-1H-indazol-5-yl]oxy]ethyl)cyclopentanamine;
- 5-(2-Morpholin-4-ylethoxy)-3-(1-naphthylsulfonyl)-1H-indazole;
- 20 N-Ethyl-N-methyl-2-[[3-(1-naphthylsulfonyl)-1H-indazol-5-yl]oxy]ethanamine;
- N-(2-[[3-(1-Naphthylsulfonyl)-1H-indazol-5-yl]oxy]ethyl)butan-1-amine;
- N~1~-{3-(1-Naphthylsulfonyl)-1H-indazol-5-yl]-beta-alaninamide;

- N-Ethyl-2-[[3-(phenylsulfonyl)-1H-indazol-5-yl]oxy]ethanamine;
 N-(2-[[3-(Phenylsulfonyl)-1H-indazol-5-yl]oxy]ethyl)propan-2-amine;
 N-(2-[[3-(1-Naphthylsulfonyl)-1H-indazol-5-yl]oxy]ethyl)propan-2-amine;
 N-Ethyl-2-[[3-(1-naphthylsulfonyl)-1H-indazol-5-yl]oxy]ethanamine;
 5 N-Methyl-2-[[3-(1-naphthylsulfonyl)-1H-indazol-5-yl]oxy]ethanamine;
 1-Methyl-3-(1-naphthylsulfonyl)-7-(2-piperidin-1-ylethoxy)-1H-indazole;
 3-(1-Naphthylsulfonyl)-5-(2-piperidin-1-ylethoxy)-1H-indazole;
 3-(2-Aminoethyl)-1-[(2,5-dimethoxyphenyl)sulfonyl]-1,3-dihydro-2H-imidazo[4,5-
 b]pyridin-2-one;
 10 N,N-Diethyl-2-[[3-(1-naphthylsulfonyl)-1H-indazol-5-yl]oxy]ethanamine;
 N-(2-[[3-(1-Naphthylsulfonyl)-1H-indazol-5-yl]oxy]ethyl)cyclopropanamine;
 1-(3-Chlorobenzyl)-3-(1-naphthylsulfonyl)-7-(2-piperidin-1-ylethoxy)-1H-indazole;
 1-(3-Chlorobenzyl)-3-(1-naphthylsulfonyl)-7-(2-pyrrolidin-1-ylethoxy)-1H-indazole;
 (2S)-3-Methyl-N~1~-[[3-(1-naphthylsulfonyl)-1H-indazol-5-yl]butane-1,2-diamine;
 15 (2-[[1-(3-Chlorobenzyl)-3-(1-naphthylsulfonyl)-1H-indazol-7-yl]oxy]ethyl)amine;
 N-(2-[[3-(Phenylsulfonyl)-1H-indazol-5-yl]oxy]ethyl)cyclopentanamine;
 3-(Phenylsulfonyl)-5-(2-pyrrolidin-1-ylethoxy)-1H-indazole;
 N-Methyl-2-[[3-(phenylsulfonyl)-1H-indazol-5-yl]oxy]ethanamine;
 N-Methyl-2-[[1-methyl-3-(phenylsulfonyl)-1H-indazol-7-yl]oxy]ethanamine;
 20 1-Methyl-3-(phenylsulfonyl)-7-(2-pyrrolidin-1-ylethoxy)-1H-indazole;
 (2-[[1-(3-Chlorobenzyl)-3-(phenylsulfonyl)-1H-indazol-7-yl]oxy]ethyl)methylamine;
 (2-[[1-(3-Chlorobenzyl)-3-(phenylsulfonyl)-1H-indazol-7-yl]oxy]ethyl)ethylamine;
 1-(3-Chlorobenzyl)-3-(phenylsulfonyl)-7-(2-pyrrolidin-1-ylethoxy)-1H-indazole;
 1-(3-Chlorobenzyl)-5-methoxy-3-(1-naphthylsulfonyl)-7-(2-piperidin-1-ylethoxy)-1H-
 25 indazole;
 N-Methyl-2-[[3-(phenylsulfonyl)-1H-indazol-7-yl]oxy]ethanamine;
 (2-[[1-(3-Chlorobenzyl)-3-(1-naphthylsulfonyl)-1H-indazol-7-yl]oxy]ethyl)ethylamine;
 (2-[[1-(3-Chlorobenzyl)-3-(1-naphthylsulfonyl)-1H-indazol-7-yl]oxy]ethyl)methylamine;
 N-Ethyl-2-[[3-(phenylsulfonyl)-1H-indazol-7-yl]oxy]ethanamine;
 30 N,N-Diethyl-2-[[3-(phenylsulfonyl)-1H-indazol-7-yl]oxy]ethanamine;
 N-(2-[[3-(Phenylsulfonyl)-1H-indazol-7-yl]oxy]ethyl)butan-1-amine;
 3-(Phenylsulfonyl)-7-(2-pyrrolidin-1-ylethoxy)-1H-indazole;
 3-(Phenylsulfonyl)-7-(2-piperidin-1-ylethoxy)-1H-indazole;

- N,N-Diethyl-2-[[1-methyl-3-(1-naphthylsulfonyl)-1H-indazol-7-yl]oxy]ethanamine;
1-Methyl-3-(1-naphthylsulfonyl)-7-(2-pyrrolidin-1-ylethoxy)-1H-indazole;
N-Ethyl-2-[[1-methyl-3-(1-naphthylsulfonyl)-1H-indazol-7-yl]oxy]ethanamine;
3-(1-Naphthylsulfonyl)-7-(2-piperidin-1-ylethoxy)-1H-indazole;
5 3-(1-Naphthylsulfonyl)-7-(2-pyrrolidin-1-ylethoxy)-1H-indazole;
N-Ethyl-2-[[3-(1-naphthylsulfonyl)-1H-indazol-7-yl]oxy]ethanamine;
(2-[[3-(1-Naphthylsulfonyl)-1H-indazol-7-yl]oxy]ethyl)amine;
(2-[[1-(3-Chlorobenzyl)-5-fluoro-3-(1-naphthylsulfonyl)-1H-indazol-7-yl]oxy]ethyl)-
dimethylamine;
10 (2-[[1-Benzyl-3-(1-naphthylsulfonyl)-1H-indazol-5-yl]oxy]ethyl)methylamine;
(2-[[1-Benzyl-3-(1-naphthylsulfonyl)-1H-indazol-5-yl]oxy]ethyl)dimethylamine;
(2-[[1-Benzyl-3-(1-naphthylsulfonyl)-1H-indazol-5-yl]oxy]ethyl)ethylamine;
N-Methyl-2-[[3-(1-naphthylsulfonyl)-1H-indazol-7-yl]oxy]ethanamine;
N,N-Dimethyl-2-[[3-(1-naphthylsulfonyl)-1H-indazol-7-yl]oxy]ethanamine;
15 5-Fluoro-3-(1-naphthylsulfonyl)-7-(2-pyrrolidin-1-ylethoxy)-1H-indazole;
5-Fluoro-3-(1-naphthylsulfonyl)-7-(2-piperidin-1-ylethoxy)-1H-indazole;
N,N-Diethyl-2-[[5-fluoro-3-(1-naphthylsulfonyl)-1H-indazol-7-yl]oxy]ethanamine;
(2-[[5-Fluoro-3-(1-naphthylsulfonyl)-1H-indazol-7-yl]oxy]ethyl)dimethylamine;
N-Ethyl-2-[[5-fluoro-3-(1-naphthylsulfonyl)-1H-indazol-7-yl]oxy]ethanamine;
20 (2-[[1-(3-Chlorobenzyl)-3-(1-naphthylsulfonyl)-1H-indazol-5-yl]oxy]ethyl)-
dimethylamine;
N-Methyl-3-[[3-(1-naphthylsulfonyl)-1H-indazol-5-yl]oxy]propan-1-amine;
N-Ethyl-N-methyl-3-[[3-(1-naphthylsulfonyl)-1H-indazol-5-yl]oxy]propan-1-amine;
3-(1-Naphthylsulfonyl)-5-(3-piperidin-1-ylpropoxy)-1H-indazole;
25 N,N-Dimethyl-3-[[3-(1-naphthylsulfonyl)-1H-indazol-5-yl]oxy]propan-1-amine;
N,N-Diethyl-3-[[3-(1-naphthylsulfonyl)-1H-indazol-5-yl]oxy]propan-1-amine;
N-(3-[[3-(1-Naphthylsulfonyl)-1H-indazol-5-yl]oxy]propyl)butan-1-amine;
3-(1-Naphthylsulfonyl)-5-(3-pyrrolidin-1-ylpropoxy)-1H-indazole;
(2-[[5-Methoxy-3-(1-naphthylsulfonyl)-1H-indazol-7-yl]oxy]ethyl)methylamine;
30 (2-[[5-Methoxy-3-(1-naphthylsulfonyl)-1H-indazol-7-yl]oxy]ethyl)dimethylamine;
5-Methoxy-3-(1-naphthylsulfonyl)-7-(2-pyrrolidin-1-ylethoxy)-1H-indazole;
5-Methoxy-3-(1-naphthylsulfonyl)-7-(2-piperidin-1-ylethoxy)-1H-indazole;
(2-[[1-(3-Chlorobenzyl)-3-(1-naphthylsulfonyl)-1H-indazol-5-yl]oxy]ethyl)ethylamine;

- (3-[[1-(3-Chlorobenzyl)-3-(1-naphthylsulfonyl)-1H-indazol-7-yl]oxy]propyl)-
diethylamine;
- 1-(3-Chlorobenzyl)-3-(1-naphthylsulfonyl)-7-(3-pyrrolidin-1-ylpropoxy)-1H-indazole;
- N-Methyl-3-[[3-(1-naphthylsulfonyl)-1H-indazol-7-yl]oxy]propan-1-amine;
- 5 N,N-Diethyl-3-[[3-(1-naphthylsulfonyl)-1H-indazol-7-yl]oxy]propan-1-amine;
- N-Methyl-2-[[1-methyl-3-(1-naphthylsulfonyl)-1H-indazol-5-yl]oxy]ethanamine;
- N,N-Dimethyl-2-[[1-methyl-3-(1-naphthylsulfonyl)-1H-indazol-5-yl]oxy]ethanamine;
- N-Ethyl-N-methyl-2-[[1-methyl-3-(1-naphthylsulfonyl)-1H-indazol-5-yl]oxy]ethan-
amine;
- 10 N-Ethyl-2-[[1-methyl-3-(1-naphthylsulfonyl)-1H-indazol-5-yl]oxy]ethanamine;
- N,N-Diethyl-2-[[1-methyl-3-(1-naphthylsulfonyl)-1H-indazol-5-yl]oxy]ethanamine;
- N-(2-[[1-Methyl-3-(1-naphthylsulfonyl)-1H-indazol-5-yl]oxy]ethyl)propan-2-amine;
- 1-Methyl-3-(1-naphthylsulfonyl)-5-(2-pyrrolidin-1-ylethoxy)-1H-indazole;
- {3-[[3-(1-Naphthylsulfonyl)-1H-indazol-5-yl]propyl]amine;
- 15 (2-[[1-Methyl-3-(1-naphthylsulfonyl)-1H-indazol-7-yl]oxy]ethyl)amine;
- N-Ethyl-3-[[3-(1-naphthylsulfonyl)-1H-indazol-5-yl]oxy]propan-1-amine;
- N-Isopropyl-3-[[3-(1-naphthylsulfonyl)-1H-indazol-5-yl]oxy]propan-1-amine;
- N-(3-[[3-(1-Naphthylsulfonyl)-1H-indazol-5-yl]oxy]propyl)cyclopentanamine;
- 5-(3-Morpholin-4-ylpropoxy)-3-(1-naphthylsulfonyl)-1H-indazole;
- 20 N-(3-[[3-(1-Naphthylsulfonyl)-1H-indazol-5-yl]oxy]propyl)cyclopropanamine;
- (3-[[3-(1-Naphthylsulfonyl)-1H-indazol-5-yl]oxy]propyl)amine;
- N-Methyl-4-[[3-(1-naphthylsulfonyl)-1H-indazol-5-yl]oxy]butan-1-amine
- N,N-Dimethyl-4-[[3-(1-naphthylsulfonyl)-1H-indazol-5-yl]oxy]butan-1-amine;
- N-Ethyl-4-[[3-(1-naphthylsulfonyl)-1H-indazol-5-yl]oxy]butan-1-amine;
- 25 N,N-Diethyl-4-[[3-(1-naphthylsulfonyl)-1H-indazol-5-yl]oxy]butan-1-amine;
- N-Methyl-4-[[3-(1-naphthylsulfonyl)-1H-indazol-5-yl]oxy]-N-propylbutan-1-amine;
- 3-(1-Naphthylsulfonyl)-5-(4-pyrrolidin-1-ylbutoxy)-1H-indazole;
- 3-(1-Naphthylsulfonyl)-5-(4-piperidin-1-ylbutoxy)-1H-indazole;
- (4-[[3-(1-Naphthylsulfonyl)-1H-indazol-5-yl]oxy]butyl)amine;
- 30 (2-[[5-Fluoro-3-(1-naphthylsulfonyl)-1H-indazol-7-yl]oxy]ethyl)methylamine;
- 5-[[4-Methylpiperazin-1-yl)methyl]-3-(1-naphthylsulfonyl)-1H-indazole;
- 3-(1-Naphthylsulfonyl)-5-(piperazin-1-ylmethyl)-1H-indazole;
- N-[[3-(1-Naphthylsulfonyl)-1H-indazol-5-yl]methyl]ethane-1,2-diamine;

- N-Methyl-3-[3-(1-naphthylsulfonyl)-1H-indazol-5-yl]propan-1-amine;
 N,N-Dimethyl-4-[3-(1-naphthylsulfonyl)-1H-indazol-5-yl]butan-1-amine;
 N,N-Dimethyl-3-[3-(1-naphthylsulfonyl)-1H-indazol-5-yl]propan-1-amine;
 N-Ethyl-N-methyl-3-[3-(1-naphthylsulfonyl)-1H-indazol-5-yl]propan-1-amine;
 5 N-Isopropyl-3-[3-(1-naphthylsulfonyl)-1H-indazol-5-yl]propan-1-amine;
 N-Ethyl-N-methyl-4-[3-(1-naphthylsulfonyl)-1H-indazol-5-yl]butan-1-amine;
 (2-[[3-(1-Naphthylsulfonyl)-1H-indazol-5-yl]oxy]ethyl)amine;
 3-(1-Naphthylsulfonyl)-5-(3-pyrrolidin-1-ylpropyl)-1H-indazole;
 N-Isopropyl-4-[3-(1-naphthylsulfonyl)-1H-indazol-5-yl]butan-1-amine;
 10 3-(1-Naphthylsulfonyl)-5-(4-pyrrolidin-1-ylbutyl)-1H-indazole;
 N-Ethyl-4-[3-(1-naphthylsulfonyl)-1H-indazol-5-yl]butan-1-amine;
 5-[(3-Methylpiperazin-1-yl)methyl]-3-(1-naphthylsulfonyl)-1H-indazole;
 5-[(3,5-Dimethylpiperazin-1-yl)methyl]-3-(1-naphthylsulfonyl)-1H-indazole;
 N-Ethyl-3-[3-(1-naphthylsulfonyl)-1H-indazol-5-yl]propan-1-amine;
 15 {4-[3-(1-Naphthylsulfonyl)-1H-indazol-5-yl]butyl}amine;
 5-[1-(4-Methylpiperazin-1-yl)ethyl]-3-(1-naphthylsulfonyl)-1H-indazole;
 N,N,N'-Trimethyl-N'-[[3-(1-naphthylsulfonyl)-1H-indazol-5-yl]methyl]ethane-1,2-
 diamine;
 N,N-Dimethyl-2-[[3-(1-naphthylsulfonyl)-1H-indazol-5-yl]methoxy]ethanamine;
 20 5-[[3-(3R)-3-Methylpiperazin-1-yl]methyl]-3-(1-naphthylsulfonyl)-1H-indazole;
 5-[[3-(3S)-3-Methylpiperazin-1-yl]methyl]-3-(1-naphthylsulfonyl)-1H-indazole;
 (3S)-N-[[3-(1-Naphthylsulfonyl)-1H-indazol-5-yl]methyl]pyrrolidin-3-amine;
 (3R)-1-[[3-(1-Naphthylsulfonyl)-1H-indazol-5-yl]methyl]pyrrolidin-3-amine;
 N-[2-(Dimethylamino)ethyl]-3-(1-naphthylsulfonyl)-1H-indazole-5-carboxamide;
 25 2-[[5-Fluoro-3-(phenylsulfonyl)-1H-indazol-7-yl]oxy]ethanamine;
 N-[3-(1-Naphthylsulfonyl)-1H-indazol-6-yl]-beta-alaninamide;
 N-[3-(1-Naphthylsulfonyl)-1H-indazol-7-yl]-3-piperidin-1-ylpropanamide;
 N-3~,N-3~-Dimethyl-N-[3-(1-naphthylsulfonyl)-1H-indazol-7-yl]-beta-alaninamide;
 2-[[3-(Phenylsulfonyl)-1H-indazol-7-yl]oxy]ethanamine;
 30 N-[3-(1-Naphthylsulfonyl)-1H-indazol-7-yl]-beta-alaninamide;
 N-[3-(1-Naphthylsulfonyl)-1H-indazol-7-yl]ethane-1,2-diamine;
 N-[3-(1-Naphthylsulfonyl)-1H-indazol-6-yl]-3-piperidin-1-ylpropanamide;
 N-[3-(1-Naphthylsulfonyl)-1H-indazol-6-yl]ethane-1,2-diamine;

N^3, N^3 -diethyl-N-[3-(1-naphthylsulfonyl)-1H-indazol-7-yl]-beta-alaninamide;
 N,N-Dimethyl-2-[[3-(1-naphthylsulfonyl)-1H-indazol-4-yl]oxy]ethanamine;
 3-(1-Naphthylsulfonyl)-4-(2-piperidin-1-ylethoxy)-1H-indazole;
 3-(1-Naphthylsulfonyl)-4-(2-pyrrolidin-1-ylethoxy)-1H-indazole;
 5 2-[[3-(1-Naphthylsulfonyl)-1H-indazol-4-yl]oxy]ethanamine;
 N-Methyl-2-[[3-(1-naphthylsulfonyl)-1H-indazol-6-yl]oxy]ethanamine;
 2-[[3-(1-Naphthylsulfonyl)-1H-indazol-6-yl]oxy]ethanamine;
 a stereoisomer thereof; or a pharmaceutically acceptable salt thereof.

In one embodiment X is O.

In one embodiment n is 3.

In one embodiment R_2 is an optionally substituted phenyl, naphthyl or imidazothiazolyl group. In another embodiment R_2 is naphthyl.

In one embodiment R_1 is H or optionally substituted alkyl.

In a further embodiment m is 0 or 1. In another embodiment m is 0.


In one embodiment R_5 and R_6 are each independently H or C_1 - C_4 alkyl.

Compounds of the invention may be prepared using conventional synthetic methods and, if required, standard isolation or separation techniques. In Schemes I through VII below, Z represents the group $(CR_3R_4)_n$ wherein R_3 , R_4 and n are as defined hereinabove for formula I.

- 10 Compounds of formula I wherein X is O (Ia) may be prepared as illustrated in Scheme I below wherein the chloromethylsulfone **2** is either commercially available or readily prepared using methods as described by M. Makosza and J. Golinski in J. Org. Chem., 1984, 49, 1488-1494; or by Antane, S.; Bernotas, R., Li, Y.; McDevitt, R.; Yan, Y. *Synthetic Communications* 2004, 34(13), 2443-2449 or by other known
- 15 methods. Reaction of a sulfonylchloride **1** with sodium sulfite under basic conditions followed by reaction with chloro-bromo-methane affords the chloromethylsulfone **2**. Reaction of **2** with a fluoronitrobenzene **3** under basic conditions gives the benzylsulfonyl derivative **4**. Compound **4** is reacted with the diol, $HO-(CR_3R_4)_n-OH$, under basic conditions to give **5**. Tosylation, followed by hydrogenation, of **5** gives
- 20 the aniline **6**. Compound **6** is reacted with sodium nitrite in the presence of an acid to give the indazole **7**. Alkylation of the indazole **7** yields the alkylated tosylate compound **8**. Displacement of the tosyl groups of either of compounds **7** or **8** with the appropriate amine then gives the desired compound of formula Ia. The reaction

is shown in Scheme I wherein Ts represents p-toluenesulfonyl and Hal represents Cl, Br or I.


SCHEME I




5

Similarly, compounds of formula I wherein X is O and R₅ and R₆ are H (lb), or compounds of formula Ia, may be prepared by reacting the nitrobenzene derivative 9 with chloromethylsulfone 2, followed by hydrogenation to give the aniline 10. The aniline 10 is converted to the indazole derivative 11, as described hereinabove in Scheme I. Alkylation or protection of the indazole 11 gives compound 12. Reaction of 12 with sodium azide followed by reduction of the azido group yields the desired primary amine lb. Alternatively, reaction of 11 or 12 with an amine, HNR₅R₆, gives the compound of Ia. The reaction is shown in Scheme II wherein Hal represents Cl, Br or I.

15


SCHEME II

- 5 Alternatively, formula Ia compounds may be prepared by the reaction of an amino alcohol **14** with either a fluoro-nitrobenzene **13** under basic conditions, or a nitrophenol **15** under Mitsunobu conditions, to give the compound **16**. Compound **16** is reacted with a chlorosulfone **2**, followed by hydrogenation, to give the aniline **17**. Subsequent formation of the desired formula Ia indazoles is carried out as described
- 10 hereinabove in Schemes I and II. The reaction is shown in Scheme III wherein Hal represents Cl, Br or I.


SCHEME III

- Compounds of formula I wherein X is NR and R and R₁ are H (Ib) or X is
- 5 NRCO and R and R₁ are H (Ic) may be prepared by reacting a nitroindazole 18 with iodine to give the corresponding 3-iodoindazole 19; coupling 19 with a thiol 20, followed by oxidation with a suitable oxidizing agent such as m-chloroperbenzoic acid (mCPBA) to give the sulfone 21; reducing the nitro group of 21 with Sn/HCl or SnCl₂/HCl to obtain the corresponding amine 22; and either reacting 22 with the
- 10 amino aldehyde 23 under reductive amination conditions to afford the desired compound of formula Ib, or coupling 22 with an amino acid 24 to give the desired compound of formula Ic. The reactions are shown in Scheme IV hereinbelow, wherein Ac represents COCH₃.

SCHEME IV




- 5 Compounds of formula I wherein X is CH_2 and R_5 and R_6 are other than H (Id) may be prepared by reacting the nitrobenzene compound 25 with the the chloromethylsulfonyl compound 2 to obtain the intermediate 26; reducing the nitro group of compound 26 to the corresponding amine and nitrosating said amine, as described in Schemes II and III, to form the desired compound of formula Id wherein
- 10 R_1 is H and optionally alkylating said compound to obtain the desired compound of formula Id wherein R_1 is other than H. The reactions are shown in Scheme V hereinbelow, wherein Hal represents Cl, Br or I.

SCHEME V

Compounds of formula Id wherein R₅ is H (Ie) may be prepared in a similar manner by protecting the compound 27 with a suitable protecting group such as t-butyloxycarbonyl (BOC), ethyloxycarbonyl, methyloxycarbonyl, benzyl (Bn), phthalimide, fluorenylmethylcarbonyl (Fmoc), acetyl, benzoyl, or the like to give the protected amine 28; reacting compound 28 with the chloromethylsulfonyl 2, followed by reduction and nitrosation as shown hereinabove in Scheme V to obtain the protected compound 29 and deprotecting to give the desired compound of formula Ie. The reaction is shown in Scheme VI hereinbelow, wherein Hal represents Cl, Br or I.

SCHEME VI



Compounds of formula Id may also be prepared by coupling the compound

5 **31** with an alkyne **32** using Sonagashira conditions to give the compound **33**;


reducing **33** to the fully saturated amine **34**; reacting **34** with NaNO_2 to give the

indazole **35** and converting the hydroxyl group to a leaving group and displacing the

leaving group with an amine, HNR_5R_6 , optionally alkylating the resultant product to

give the desired compound of formula Id. The reactions are shown in Scheme VII

10 hereinbelow wherein TsCl represents tosyl chloride and Hal represents Cl, Br or I.

SCHEME VII

- 5 Advantageously, the formula I compounds of the invention are useful for the treatment of CNS disorders related to or affected by the 5-HT6 receptor including motor, mood, personality, behavioral, psychiatric, cognitive, neurodegenerative, or the like disorders, for example Alzheimer's disease, Parkinson's disease, attention deficit disorder, anxiety, epilepsy, depression, obsessive compulsive disorder, sleep disorders, neurodegenerative disorders (such as head trauma or stroke), feeding disorders (such as anorexia or bulimia), schizophrenia, memory loss, disorders associated with withdrawal from drug or nicotine abuse, or the like or certain gastrointestinal disorders such as irritable bowel syndrome. Accordingly, the present invention provides a method for the treatment of a disorder of the central nervous system related to or affected by the 5-HT6 receptor in a patient in need thereof which comprises providing said patient a therapeutically effective amount of a compound of formula I as described hereinabove. The compounds may be provided by oral or parenteral administration or in any common manner known to be an effective administration of a therapeutic agent to a patient in need thereof.
- 10
- 15
- 20 The term "providing" as used herein with respect to providing a compound or substance embraced by the invention, designates either directly administering such a

compound or substance, or administering a prodrug, derivative or analog which forms an equivalent amount of the compound or substance within the body.

The inventive method includes: a method for the treatment of schizophrenia; a method for the treatment of a disease associated with a deficit in memory, cognition, and/or learning or a cognitive disorder such as Alzheimer's disease or attention deficit disorder; a method for the treatment of developmental disorders such as schizophrenia; Down's syndrome, Fragile X syndrome, autism or the like; a method for the treatment of behavioral disorders, e.g., anxiety, depression, or obsessive compulsive disorder; a method for the treatment of motion or motor disorders such as Parkinson's disease or epilepsy; a method for the treatment of a neurodegenerative disorder such as stroke or head trauma or withdrawal from drug addiction including addiction to nicotine, alcohol, or other substances of abuse, or any other CNS disease or disorder associated with or related to the 5-HT₆ receptor.

In one embodiment, the present invention provides a method for treating attention deficit disorders (ADD, also known as Attention Deficit Hyperactivity Disorder or ADHD) in both children and adults. Accordingly, in this embodiment, the present invention provides a method for treating attention deficit disorders in a pediatric patient.

The present invention therefore provides a method for the treatment of each of the conditions listed above in a patient, preferably in a human, said method comprises providing said patient a therapeutically effective amount of a compound of formula I as described hereinabove. The compounds may be provided by oral or parenteral administration or in any common manner known to be an effective administration of a therapeutic agent to a patient in need thereof.

The therapeutically effective amount provided in the treatment of a specific CNS disorder may vary according to the specific condition(s) being treated, the size, age and response pattern of the patient, the severity of the disorder, the judgment of the attending physician and the like. In general, effective amounts for daily oral administration may be about 0.01 to 1,000 mg/kg, preferably about 0.5 to 500 mg/kg and effective amounts for parenteral administration may be about 0.1 to 100 mg/kg, preferably about 0.5 to 50 mg/kg.

In actual practice, the compounds of the invention are provided by administering the compound or a precursor thereof in a solid or liquid form, either

neat or in combination with one or more conventional pharmaceutical carriers or excipients. Accordingly, the present invention provides a pharmaceutical composition which comprises a pharmaceutically acceptable carrier and an effective amount of a compound of formula I as described hereinabove.

5 In one embodiment, the invention relates to compositions comprising at least one compound of formula I, or a pharmaceutically acceptable salt thereof, and one or more pharmaceutically acceptable carriers, excipients, or diluents. Such compositions include pharmaceutical compositions for treating or controlling disease states or conditions of the central nervous system. In certain embodiments, the
10 compositions comprise mixtures of one or more compounds of formula I.

 In certain embodiments, the invention relates to compositions comprising at least one compound of formula I, or a pharmaceutically acceptable salt thereof, and one or more pharmaceutically acceptable carriers, excipients, or diluents. Such compositions are prepared in accordance with acceptable pharmaceutical procedures. Pharmaceutically acceptable carriers are those carriers that are compatible with the other ingredients in the formulation and are biologically acceptable.

 The compounds of formula I may be administered orally or parenterally, neat, or in combination with conventional pharmaceutical carriers. Applicable solid carriers can include one or more substances that can also act as flavoring agents, lubricants, solubilizers, suspending agents, fillers, glidants, compression aids, binders, tablet-disintegrating agents, or encapsulating materials. In powders, the carrier is a finely divided solid that is in admixture with the finely divided active ingredient. In tablets, the active ingredient is mixed with a carrier having the necessary compression properties in suitable proportions and compacted in the shape and size desired. The powders and tablets preferably contain up to 99% of the active ingredient. Suitable solid carriers include, for example, calcium phosphate, magnesium stearate, talc, sugars, lactose, dextrin, starch, gelatin, cellulose, methyl cellulose, sodium carboxymethyl cellulose, polyvinylpyrrolidone, low melting waxes and ion exchange resins.

 In certain embodiments, a compound of formula I is provided in a disintegrating tablet formulation suitable for pediatric administration.

Liquid carriers can be used in preparing solutions, suspensions, emulsions, syrups and elixirs. The active ingredient can be dissolved or suspended in a pharmaceutically acceptable liquid carrier such as water, an organic solvent, a mixture of both, or a pharmaceutically acceptable oil or fat. The liquid carrier can contain other suitable pharmaceutical additives such as, for example, solubilizers, emulsifiers, buffers, preservatives, sweeteners, flavoring agents, suspending agents, thickening agents, colors, viscosity regulators, stabilizers or osmo-regulators. Suitable examples of liquid carriers for oral and parenteral administration include water (particularly containing additives as above, e.g. cellulose derivatives, preferably sodium carboxymethyl cellulose solution), alcohols (including monohydric alcohols and polyhydric alcohols e.g. glycols) and their derivatives, and oils (e.g. fractionated coconut oil and arachis oil). For parenteral administration, the carrier can also be an oily ester such as ethyl oleate and isopropyl myristate. Sterile liquid carriers are used in sterile liquid form compositions for parenteral administration. The liquid carrier for pressurized compositions can be halogenated hydrocarbon or other pharmaceutically acceptable propellant.

In certain embodiments, a liquid pharmaceutical composition is provided wherein said composition is suitable for pediatric administration. In other embodiments, the liquid composition is a syrup or suspension.

Liquid pharmaceutical compositions that are sterile solutions or suspensions can be administered by, for example, intramuscular, intraperitoneal or subcutaneous injection. Sterile solutions can also be administered intravenously. Compositions for oral administration can be in either liquid or solid form.

The compounds of formula I may be administered rectally or vaginally in the form of a conventional suppository. For administration by intranasal or intrabronchial inhalation or insufflation, the compounds of formula I can be formulated into an aqueous or partially aqueous solution, which can then be utilized in the form of an aerosol. The compounds of formula I can also be administered transdermally through the use of a transdermal patch containing the active compound and a carrier that is inert to the active compound, is non-toxic to the skin, and allows delivery of the agent for systemic absorption into the blood stream via the skin. The carrier can take any number of forms such as creams and ointments, pastes, gels, and occlusive devices. The creams and ointments can be viscous liquid or semisolid emulsions of

either the oil-in-water or water-in-oil type. Pastes comprised of absorptive powders dispersed in petroleum or hydrophilic petroleum containing the active ingredient can also be suitable. A variety of occlusive devices can be used to release the active ingredient into the blood stream such as a semipermeable membrane covering a reservoir containing the active ingredient with or without a carrier, or a matrix containing the active ingredient. Other occlusive devices are known in the literature.


Preferably the pharmaceutical composition is in unit dosage form, e.g. as tablets, capsules, powders, solutions, suspensions, emulsions, granules, or suppositories. In such form, the composition is sub-divided in unit dose containing appropriate quantities of the active ingredient; the unit dosage forms can be packaged compositions, for example, packeted powders, vials, ampoules, prefilled syringes or sachets containing liquids. The unit dosage form can be, for example, a capsule or tablet itself, or it can be the appropriate number of any such compositions in package form.

The therapeutically effective amount of a compound of formula I provided to a patient will vary depending upon what is being administered, the purpose of the administration, such as prophylaxis or therapy, the state of the patient, the manner of administration, and the like. In therapeutic applications, compounds of formula I are provided to a patient suffering from a condition in an amount sufficient to treat or at least partially treat the symptoms of the condition and its complications. An amount adequate to accomplish this is a "therapeutically effective amount" as described previously herein. The dosage to be used in the treatment of a specific case must be subjectively determined by the attending physician. The variables involved include the specific condition and the size, age, and response pattern of the patient. The treatment of substance abuse follows the same method of subjective drug administration under the guidance of the attending physician. Generally, a starting dose is about 5 mg per day with gradual increase in the daily dose to about 150 mg per day, to provide the desired dosage level in the patient.

In certain embodiments, the present invention is directed to prodrugs of compounds of formula I. The term "prodrug," as used herein, means a compound that is convertible *in vivo* by metabolic means (e.g. by hydrolysis) to a compound of formula I. Various forms of prodrugs are known in the art such as those discussed in, for example, Bundgaard, (ed.), Design of Prodrugs, Elsevier (1985); Widder, et al.

(ed.), Methods in Enzymology, vol. 4, Academic Press (1985); Krogsgaard-Larsen, et al., (ed). "Design and Application of Prodrugs, Textbook of Drug Design and Development, Chapter 5, 113-191 (1991), Bundgaard, et al., Journal of Drug Delivery Reviews, 8:1-38(1992), Bundgaard, J. of Pharmaceutical Sciences, 77:285 et seq.
5 (1988); and Higuchi and Stella (eds.) Prodrugs as Novel Drug Delivery Systems, American Chemical Society (1975).

For a more clear understanding, and in order to illustrate the invention more clearly, specific examples thereof are set forth hereinbelow. The following examples are merely illustrative and are not to be understood as limiting the scope and
10 underlying principles of the invention in any way. The terms HNMR, MS and HPLC designate mass spectrum, high performance liquid chromatography and proton nuclear magnetic resonance, respectively. The terms THF, DMF and DMSO designate tetrahydrofuran, dimethyl formamide and dimethylsulfoxide, respectively. All column chromatography is performed using SiO₂ as support. Unless otherwise
15 noted, all parts are parts by weight.


Example 1**[2-(3-Benzenesulfonyl-1H-indazol-5-yloxy)-ethyl]-dimethyl-amine****Step 1****5 2-Benzenesulfonylmethyl-4-fluoro-1-nitrobenzene:**

To a solution of chloromethylphenyl sulfone (10.51 g, 55.13 mmol) in THF (110 mL) was added 5.9 mL (56 mmol) of 1-fluoro-4-nitrobenzene. The reaction mixture was chilled to 0 °C, and 1.0 M potassium *tert*-butoxide in THF (145 mL, 145 mmol) was added dropwise. The reaction mixture was stirred under nitrogen at ambient temperature for one hour. Acetic acid (9 mL, 160 mmol) was then added.

10 The reaction mixture was solvent evaporated and partitioned in brine and ethyl acetate. The organic phase was then dried with anhydrous magnesium sulfate, filtered and concentrated. The residue was stirred in diethyl ether, filtered and dried *in vacuo* at 73°C for 12 hours. 2-Benzenesulfonylmethyl-4-fluoro-1-nitrobenzene as

15 a light brown-yellow solid was obtained (10.24 g, 62.8 %): MP: 169-171 °C; Mass spectrum (-EI, [M-H]⁻) *m/z* 294. ¹HNMR (500 MHz, DMSO-*d*₆): δ 8.10-8.14 (m, 1H), 7.70-7.75 (m, 1H), 7.56-7.63 (m, 4H), 7.45-7.50 (m, 1H), 7.20 (dd, 1H, *J* = 9.27 Hz and 2.81 Hz), 5.12 ppm (s, 2H). Elemental Analysis for C₁₃H₁₀FNO₄S: Calcd: C, 52.88; H, 3.41; N, 4.74; Found: C, 52.63; H, 3.14; N, 4.66;


20

Step 2**2-(3-Benzenesulfonylmethyl-4-nitro-phenoxy)-ethanol**

A mixture of 2-benzenesulfonylmethyl-4-fluoro-1-nitrobenzene (10.2 g, 34.7 mmol), ethylene glycol (80 mL, 1.4 mol), and 1 M potassium *tert*-butoxide in THF (78 mL, 78 mmol) in THF (50 mL) was refluxed under nitrogen for 30 minutes. After cooling to about ambient temperature, the reaction mixture was solvent evaporated. Water was added to the residue, and it was poured into 2.0 N hydrochloric acid and ice. The mixture was then extracted with ethyl acetate and washed with water and

brine. The organic phase was dried with anhydrous magnesium sulfate, filtered, concentrated and dried *in vacuo* at 74 °C for 20 minutes. 2-(3-Benzenesulfonylmethyl-4-nitro-phenoxy)-ethanol as a dark brown gum (11.7 g, 100 % yield) was obtained; Mass spectrum (-EI, [M-H]⁻) *m/z* 336. ¹HNMR (500 MHz, DMSO-d₆): δ 8.01 (d, 1H, *J*=9.15 Hz), 7.67-7.73 (m, 1H), 7.54-7.63 (m, 4H), 7.10 (dd, 1H, *J*= 9.51 Hz and 2.93 Hz), 6.85 (d, 1H, *J*=2.81 Hz), 5.12 (s, 2H), 4.88-4.90 (m, 1H), 3.93-3.96 (m, 2H), 3.65 ppm (d, 2H, *J*=4.51 Hz). Elemental Analysis for C₁₅H₁₅NO₆S: Calcd: C, 53.41; H, 4.48; N, 4.15; Found: C, 53.47; H, 4.69; N, 4.04;


10 Step 3



Toluene-4-sulfonic acid 2-(3-benzenesulfonylmethyl-4-nitro-phenoxy)-ethyl ester

A mixture of 2-(3-benzenesulfonylmethyl-4-nitro-phenoxy)-ethanol (8.2 g, 24 mmol), *p*-toluenesulfonyl chloride (9.39 g, 49.3 mmol) and triethylamine (19 mL, 140 mmol) in methylene chloride (85 mL) was stirred at ambient temperature under nitrogen for 4.5 hours. It then was solvent evaporated and partitioned in methylene chloride and aqueous sodium bicarbonate. The organic phase was then washed with brine, dried with anhydrous magnesium sulfate, filtered, concentrated and dried *in vacuo* at 63°C for 30 minutes. The residue was purified by flash chromatography with 100% chloroform and 5% methanol in chloroform. It was dried *in vacuo* at 65 °C for 20 minutes to yield toluene-4-sulfonic acid 2-(3-benzenesulfonylmethyl-4-nitro-phenoxy)-ethyl ester as a yellow solid (8.1 g, 69 %); Mass spectrum (-EI, [M-H]⁻) *m/z* 490. ¹HNMR (300 MHz, DMSO-d₆): δ 7.99 (d, 1H, *J*=9.02 Hz), 7.67-7.75 (m, 3H), 7.54-7.62 (m, 4H), 7.43 (d, 2H, *J*=7.93 Hz), 7.00 (dd, 1H, *J*=9.15 Hz and 2.80 Hz), 6.78 (d, 1H, *J*=2.81 Hz), 5.09 (s, 2H), 4.29-4.31 (m, 2H), 4.13-4.15 (m, 2H), 2.37 ppm (s, 3H). Elemental Analysis for C₂₂H₂₁NO₈S 0.40 mol H₂O: Calcd: C, 52.98; H, 4.41; N, 2.81; Found: C, 52.67; H, 4.26; N, 2.61.


30 Step 4



Toluene-4-sulfonic acid 2-(4-amino-3-benzenesulfonylmethyl-phenoxy)-ethyl ester

A solution of toluene-4-sulfonic acid 2-(3-benzenesulfonylmethyl-4-nitro-phenoxy)-ethyl ester (13.1 g, 26.7 mmol) in THF (350 mL) and ethanol (250 mL) was added to 10 % palladium on carbon (5.8 g) and hydrogenated on the Parr apparatus with for one hour (starting pressure of 30-40 psi.). The mixture was filtered over Celite, concentrated and dried *in vacuo* at 80 °C for 30 minutes to yield toluene-4-sulfonic acid 2-(4-amino-3-benzenesulfonylmethyl-phenoxy)-ethyl ester as a light yellow solid (11.7 g, 95.1 %): MP: 144-6 °C; Mass spectrum (+EI, [M+H]⁺) *m/z* 462. ¹HNMR (500 MHz, DMSO-d₆): δ 7.72-7.75 (m, 4H), 7.63-7.66 (m, 1H), 7.52-7.56 (m, 2H), 7.44 (d, 2H, *J*=8.05 Hz), 6.52 (d, 2H, *J*=1.58 Hz), 6.29-6.30 (m, 1H), 4.65 (s, 2H), 4.46 (s, 2H), 4.16-4.18 (m, 2H), 3.79-3.82 (m, 2H), 2.38 ppm (s, 3H). Elemental Analysis for C₂₂H₂₃NO₆S: Calcd: C, 57.25; H, 5.02; N, 3.03; Found: C, 57.60; H, 4.98; N, 3.10.

Step 5




Toluene-4-sulfonic acid 2-(3-benzenesulfonyl-1H-indazol-5-yloxy)-ethyl ester

To a mixture of toluene-4-sulfonic acid 2-(4-amino-3-benzenesulfonylmethyl-phenoxy)-ethyl ester (11.7 g, 25.3 mmol) of in ethanol (350 mL) and 1.0 N hydrochloric acid (425 mL) was dropwise added sodium nitrite (2.67 g, 38.7 mmol) in water (50 mL). After stirring at ambient temperature for 1.5 hours, solid sodium carbonate was added to basic pH. The reaction mixture was stirred for an additional 2 hours. It was then solvent evaporated and extracted with warm ethyl acetate. The organic phase was washed with water and brine, dried with anhydrous magnesium sulfate, filtered and concentrated. The residue was purified by flash chromatography with 2 % methanol in chloroform to yield toluene-4-sulfonic acid 2-(3-


benzenesulfonyl-1H-indazol-5-yloxy)-ethyl ester as an orange solid (9.54 g, 79.5 %):
 MP: 174-7 °C; Mass Spectrum (-EI, [M-H]⁻) *m/z* 471. ¹HNMR (500 MHz, DMSO-d₆):
 δ14.11 (s, 1H), 7.96-7.98 (m, 2H), 7.74-7.77 (m, 2H), 7.62-7.66 (m, 1H), 7.53-7.59
 (m, 3H), 7.39 (d, 2H, *J*=8.05 Hz), 7.24 (d, 1H, *J*=2.32 Hz), 6.97-7.00 (m, 1H), 4.35-
 5 4.37 (m, 2H), 4.21-4.23 (m, 2H), 2.33 ppm (s, 3H).

Step 6




[2-(3-Benzenesulfonyl-1H-indazol-5-yloxy)-ethyl]-dimethyl-amine

10 A solution of toluene-4-sulfonic acid 2-(3-benzenesulfonyl-1H-indazol-5-yloxy)-ethyl ester (0.341 g, 0.722 mmol) in 2.0 N dimethylamine in THF (9 mL, 18 mmol) of was stirred at 70 °C for 6 hours in a sealed tube. After cooling somewhat, the reaction mixture was solvent evaporated and partitioned in ethyl acetate and aqueous sodium bicarbonate. The organic phase was washed with water and brine.
 15 It was dried with anhydrous magnesium sulfate, filtered and concentrated. The residue was partitioned in ethyl acetate/aqueous potassium carbonate. The organic phase was washed with water and brine, dried with anhydrous magnesium sulfate, filtered and concentrated. [2-(3-Benzenesulfonyl-1H-indazol-5-yloxy)-ethyl]-dimethyl-amine as a light yellow-brown solid (0.167 g, 67.1 %) resulted. The compound was
 20 dissolved in methanol, and ethereal hydrochloride was added. After concentrating and drying *in vacuo* for 12 hours at 70 °C, the hydrochloride as a light orange foam (94.0 mg) was obtained; Mass Spectrum (+EI, [M+H]⁺) *m/z* 346. ¹HNMR (500 MHz, DMSO-d₆): δ14.22 (s, 1H), 10.00 (br, 1H), 7.97-7.99 (m, 2H), 7.56-7.68 (m, 4H), 7.42 (d, 1H, *J*= 2.20 Hz), 7.17-7.20 (m, 1H), 4.39-4.42 (m, 2H), 3.52-3.53 (m, 2H), 2.84
 25 ppm (s, 6H).

Example 2**[2-(3-Benzenesulfonyl-1H-indazol-5-yloxy)-ethyl]-ethyl-methyl-amine**


A solution of toluene-4-sulfonic acid 2-(3-benzenesulfonyl-1H-indazol-5-yloxy)-ethyl ester (0.364 g, 0.770 mmol) and N-ethylmethylamine (0.95 mL, 11 mmol) in THF (8 mL) was stirred for 6 hours at 70 °C in a sealed tube. After cooling somewhat to ambient temperature, the reaction mixture was solvent evaporated and partitioned in ethyl acetate and aqueous sodium bicarbonate. The organic phase was washed with water and brine, dried with anhydrous magnesium sulfate, filtered and concentrated. After drying *in vacuo* at ambient temperature for 1.5 hours, the resulting buff solid (0.249 g, 89.9%), [2-(3-benzenesulfonyl-1H-indazol-5-yloxy)-ethyl]-ethyl-methyl-amine, was dissolved in methanol and methylene chloride, and ethereal hydrochloride was added. Concentration and drying at 80 °C for 12 hours yielded the hydrochloride as a light orange foam (0.253 g); Mass Spectrum (+EI, [M+H]⁺) *m/z* 360. ¹HNMR (500 MHz, DMSO-*d*₆): δ 14.23 (s, 1H), 9.95 (br, s, 1H), 7.97-7.99 (m, 2H), 7.56-7.68 (m, 4H), 7.42 (d, 1H, *J*=2.32 Hz), 7.19 (dd, 1H, *J*=9.15 Hz and 2.32 Hz), 4.41-4.43 (m, 2H), 3.55-3.60 (m, 1H), 3.43-3.49 (m, 1H), 3.11-3.17 (m, 1H), 2.82 (d, 3H, *J*=4.63 Hz), 1.22-1.25 ppm (m, 3H).

Example 3**3-Benzenesulfonyl-5-(2-piperidin-1-yl-ethoxy)-1H-indazole**

A solution of toluene-4-sulfonic acid 2-(3-benzenesulfonyl-1H-indazol-5-yloxy)-ethyl ester (0.386 g, 0.817 mmol) and piperidine (1.0 mL, 10 mmol) in THF (8 mL) was stirred at 70 °C for 6 hours in a sealed tube. After cooling somewhat to ambient temperature, the reaction mixture was solvent evaporated and partitioned in ethyl acetate and aqueous sodium bicarbonate. The organic phase was washed with


water and brine. It was dried with anhydrous magnesium sulfate, filtered and concentrated. Drying *in vacuo* at 80 °C for 20 minutes resulted in 3-benzenesulfonyl-5-(2-piperidin-1-yl-ethoxy)-1H-indazole as a light yellow solid (0.275 g, 87.3%). This was dissolved in methanol, and ethereal hydrochloride was added. This mixture was concentrated and dried *in vacuo* at 70°C for about 12 hours to yield the hydrochloride as a yellow solid (0.268 g): MP: 249-250 °C; Mass Spectrum (+EI, [M+H]⁺) *m/z* 386. ¹HNMR (500 MHz, DMSO-d₆): δ14.24 (s, 1H), 10.02-10.03 (br, 1H), 7.97-7.99 (m, 2H), 7.56-7.68 (m, 4H), 7.41 (d, 1H, *J*=2.32 Hz), 7.18 (dd, 1H, *J*=9.15 Hz and 2.33 Hz), 4.44-4.46 (m, 2H), 3.48-3.51 (m, 4H), 2.94-3.04 (m, 2H), 1.65-1.77 (m, 5H), 1.31-1.42 ppm (m, 1H). Elemental Analysis for C₂₀H₂₃N₃O₃S·1.00 mol HCl · 0.15 mol H₂O: Calcd: C, 56.57; H, 5.77; N, 9.90; Found: C, 56.34; H, 5.88; N, 9.57.

Example 4




15 [2-(3-Benzenesulfonyl-1H-indazol-5-yloxy)-ethyl]-ethyl-amine

A solution of toluene-4-sulfonic acid 2-(3-benzenesulfonyl-1H-indazol-5-yloxy)-ethyl ester (0.361 g, 0.764 mmol) in 2.0 N ethylamine in THF (9 mL, 18 mmol) was stirred at 70 °C for 6 hours in a sealed tube. After cooling somewhat, the reaction mixture was solvent evaporated and partitioned in ethyl acetate and aqueous sodium bicarbonate. The organic phase was then washed with water and brine, dried with anhydrous magnesium sulfate, filtered and concentrated. The residue was purified by flash chromatography using 10 % methanol in chloroform. After drying *in vacuo* at 68 °C for 20 minutes, [2-(3-benzenesulfonyl-1H-indazol-5-yloxy)-ethyl]-ethyl-amine as a light brown gum (0.116 g, 43.9 %) was obtained. This was dissolved in chloroform and methanol, and ethereal hydrochloride was added. Concentrating and drying at 74°C for 13.5 hours *in vacuo* yielded the hydrochloride as a buff solid (0.113 g): MP: 248-50 °C (dec). Mass Spectrum (+EI, [M+H]⁺) *m/z* 346. ¹HNMR (500 MHz, DMSO-d₆): δ14.23 (s, 1H), 8.81 (br, 2H), 7.96-7.98 (m, 2H), 7.56-7.68 (m, 4H), 7.40 (d, 1H, *J*=2.07 Hz), 7.19 (dd, 1H, *J*=9.15 Hz and 2.32 Hz), 4.31 (t, 2H, *J*=5.00 Hz), 3.35 (br, 2H), 3.00-3.05 (m, 2H), 1.18-1.22 ppm (m, 3H).

Example 5**[2-(3-Benzenesulfonyl-1H-indazol-5-yloxy)-ethyl]-diethyl-amine**

- 5 A solution of toluene-4-sulfonic acid 2-(3-benzenesulfonyl-1H-indazol-5-yloxy)-ethyl ester (0.420 g, 0.889 mmol) and diethylamine (1.0 mL, 9.7 mmol) in THF (8 mL) was stirred at 70 °C for 6 hours in a sealed tube. Additional diethylamine (1.0 mL, 9.7 mmol) was added, and the reaction mixture was stirred at 80 °C in a sealed tube for an 6 hours. After cooling somewhat, the reaction mixture was solvent
- 10 evaporated and partitioned in ethyl acetate and aqueous sodium bicarbonate. The organic phase was then washed with water and brine, dried with anhydrous magnesium sulfate, filtered and concentrated. The residue was purified by flash chromatography using 7.5 % methanol in chloroform. After drying *in vacuo* at 80 °C for 20 minutes, [2-(3-benzenesulfonyl-1H-indazol-5-yloxy)-ethyl]-diethyl-amine as a
- 15 pale yellow solid (0.155 g, 46.6 %) was obtained. This was dissolved in methanol and chloroform and ethereal hydrochloride was added. The mixture was concentrated and dried *in vacuo* at 74 °C for 13.5 hours to yield the hydrochloride as a light orange foam (0.163 g); Mass Spectrum (+EI, [M+H]⁺) *m/z* 374. ¹HNMR (500 MHz, DMSO-d₆): δ14.24 (s, 1H), 9.94 (br, 1H), 7.96-7.99 (m, 2H), 7.56-7.68 (m, 4H), 7.41 (d, 1H, *J*=2.20 Hz), 7.16-7.19 (m, 1H), 4.40-4.43 (m, 2H), 3.52 (br, 2H), 3.19-3.23 (m, 4H), 1.23 ppm (t, 6H, *J*= 7.20 Hz). Elemental Analysis for C₁₉H₂₃N₃O₃S · 1.00 mol HCl 0.20 mol H₂O: Calcd: C, 55.18; H, 5.95; N, 10.16; Found: C, 54.85; H, 5.85; N, 10.02.
- 20

Example 6**[2-(3-Benzenesulfonyl-1H-indazol-5-yloxy)-ethyl]-isopropyl-amine**

A solution of toluene-4-sulfonic acid 2-(3-benzenesulfonyl-1H-indazol-5-yloxy)-ethyl ester (0.378 g, 0.800 mmol) and isopropylamine (1.0 mL, 12 mmol) in THF (8 mL) was stirred at 70 °C for 6 hours in a sealed tube. Additional isopropylamine (1.0 mL, 12 mmol) was added, and the reaction mixture was stirred at 80 °C for 6 hours in a sealed tube. After cooling somewhat, the reaction mixture was solvent evaporated and partitioned in ethyl acetate and aqueous sodium bicarbonate. The organic phase was then washed with water and brine, dried with anhydrous magnesium sulfate, filtered and concentrated. The residue was purified by flash chromatography using 10 % methanol in chloroform. After drying *in vacuo* at 80 °C for 30 minutes, [2-(3-benzenesulfonyl-1H-indazol-5-yloxy)-ethyl]-isopropyl-amine as a pale yellow solid was obtained (0.110 g, 38.2 %). This was dissolved in methanol and chloroform, and ethereal hydrochloride was added. The mixture was concentrated and dried *in vacuo* at 74 °C for 13.5 hours to yield the hydrochloride as an off-white solid (0.117 g): MP: 275-7 °C (dec); Mass Spectrum (+EI, [M+H]⁺) *m/z* 360. ¹HNMR (500 MHz, DMSO-*d*₆): δ 14.22 (br, 1H), 8.75 (br, 2H), 7.96-7.98 (m, 2H), 7.56-7.68 (m, 4H), 7.41 (d, 1H, *J*=2.08 Hz), 7.19 (dd, 1H, *J*= 9.15 Hz and 2.32 Hz), 4.30-4.33 (m, 2H), 3.34-3.40 (m, 3H), 1.25 ppm (d, 6H, *J*=6.59 Hz). Elemental Analysis for C₁₈H₂₁N₃O₃S·1.00 mol HCl · 0.10 mol H₂O: Calcd: C, 54.36; H, 5.36; N, 10.57; Found: C, 54.02; H, 5.42; N, 10.22.

Example 7




[2-(3-Benzenesulfonyl-1H-indazol-5-yloxy)-ethyl]-butyl-amine

A solution of toluene-4-sulfonic acid 2-(3-benzenesulfonyl-1H-indazol-5-yloxy)-ethyl ester (0.348 g, 0.736 mmol) and *n*-butylamine (1.0 mL, 10 mmol) in THF (8 mL) was stirred at 70 °C for 6 hours in a sealed tube. After cooling somewhat, the reaction mixture was solvent evaporated and partitioned in ethyl acetate and aqueous sodium bicarbonate. The organic phase was then washed with water and brine, dried with anhydrous magnesium sulfate, filtered and concentrated. The residue was purified by flash chromatography using 7.5–10 % methanol in

chloroform. After drying *in vacuo* at 55 °C for 40 minutes, [2-(3-benzenesulfonyl-1H-indazol-5-yloxy)-ethyl]-butyl-amine as a light beige semi-solid (0.105 g, 38.2 %) was obtained. This was dissolved in chloroform, and ethereal hydrochloride was added. The resulting solid was filtered and dried *in vacuo* at 75 °C for 16.5 hours to yield the hydrochloride as a white solid (0.0813 g): MP: 271-3 °C dec; Mass Spectrum (+EI, [M+H]⁺) *m/z* 374. ¹HNMR (300 MHz, DMSO-d₆): δ14.27-14.35 (br, 1H), 8.98-9.00 (br, 2H), 8.03-8.06 (m, 2H), 7.62-7.76 (m, 4H), 7.46 (d, 1H, *J*=2.19 Hz) 7.25 (dd, 1H, *J*=9.15 Hz and 2.29 Hz), 4.38-4.41 (m, 2H), 3.41-3.44 (m, 2H), 3.03 (t, 2H, *J*=7.87 Hz), 1.62-1.72 (m, 2H), 1.33-1.45 (m, 2H), 0.94 ppm (t, 3H, *J*=7.32 Hz). Elemental Analysis for C₁₉H₂₃N₃O₃S · 1.00 mol HCl · 0.05 mol H₂O: Calcd: C, 55.55; H, 5.91; N, 10.23; Found: C, 55.23; H, 5.87; N, 10.09.

Example 8




15 [2-(3-Benzenesulfonyl-1H-indazol-5-yloxy)-ethyl]-cyclopropyl-amine

A solution of toluene-4-sulfonic acid 2-(3-benzenesulfonyl-1H-indazol-5-yloxy)-ethyl ester (0.336 g, 0.711 mmol) and cyclopropylamine (1.0 mL, 14 mmol) in THF (8 mL) was stirred at 70 °C for 6 hours in a sealed tube. Additional cyclopropylamine (1.0 mL, 14 mmol) was added, and the reaction mixture was stirred for 6 hours at 80 °C in a sealed tube. After cooling somewhat, the reaction mixture was solvent evaporated and partitioned in ethyl acetate and aqueous sodium bicarbonate. The organic phase was then washed with water and brine, dried with anhydrous magnesium sulfate, filtered and concentrated. The residue was purified by flash chromatography using 5 % methanol in chloroform. After drying *in vacuo* at 55 °C temperature for 25 minutes, [2-(3-benzenesulfonyl-1H-indazol-5-yloxy)-ethyl]-cyclopropyl-amine as a light yellow foam (0.127 g, 50.0 %) was obtained. This was dissolved in chloroform, and ethereal hydrochloride was added. The mixture was concentrated and dried *in vacuo* at 75 °C for 16.5 hours to yield the hydrochloride as a buff solid (0.131 g): MP: 184-5 °C dec.; Mass Spectrum (+EI, [M+H]⁺) *m/z* 358. ¹HNMR (300 MHz, DMSO-d₆): δ14.33 (s, 1H), 9.33 (br, 2H), 8.03-8.06 (m, 2H), 7.60-

7.76 (m, 4H), 7.47 (d, 1H, $J=2.11$ Hz), 7.25 (dd, 1H, $J=9.15$ Hz and 2.38 Hz), 4.40-4.43 (m, 2H), 3.49-3.52 (m, 2H), 2.81-2.88 (m, 1H), 1.62-1.72 (m, 2H), 0.92-0.97 (m, 2H), 0.77-0.87 ppm (m, 2H).


5

Example 9




3-Benzenesulfonyl-5-(2-cyclopentyl-1-yl-ethoxy)-1H-indazole

A solution of toluene-4-sulfonic acid 2-(3-benzenesulfonyl-1H-indazol-5-yloxy)-ethyl ester (0.345 g, 0.730 mmol) and cyclopentylamine (1.0 mL, 10 mmol) in THF (8 mL) was stirred in a sealed tube at 70 °C for 6 hours and 80 °C for 6 hours. After cooling somewhat, the reaction mixture was solvent evaporated and partitioned in ethyl acetate and aqueous sodium bicarbonate. The organic phase was then washed with water and brine, dried with anhydrous magnesium sulfate, filtered and concentrated. The residue was purified by flash chromatography using 7.5 % methanol in chloroform. After drying *in vacuo* at 57 °C for 25 minutes, 3-benzenesulfonyl-5-(2-cyclopentyl-1-yl-ethoxy)-1H-indazole as a yellow foam (0.137 g, 48.8 %) was obtained. This was dissolved in methanol and chloroform, and ethereal hydrochloride was added. The mixture was concentrated and dried *in vacuo* at 75 °C for 16.5 hours to yield the hydrochloride as a white solid (0.0956 g): MP: 255-8 °C(dec.); Mass Spectrum (+EI, $[M+H]^+$) m/z 386. ^1H NMR (300 MHz, DMSO- d_6): δ 14.31-14.41 (br, 1H), 9.13-9.14 (br, 2H), 8.03-8.06 (m, 2H), 7.62-7.76 (m, 4H), 7.46 (d, 1H, $J=2.10$ Hz) 7.25 (dd, 1H, $J=9.15$ Hz and 2.29 Hz), 4.39-4.42 (m, 2H), 3.58-3.63 (m, 1H), 3.41 (s, 2H), 1.99-2.05 (m, 2H), 1.67-1.82 (m, 4H), 1.53-1.62 ppm (m, 2H). Elemental Analysis for $\text{C}_{20}\text{H}_{23}\text{N}_3\text{O}_3\text{S} \cdot 1.00 \text{ mol HCl} \cdot 0.50 \text{ mol H}_2\text{O}$: Calcd: C, 55.74; H, 5.85; N, 9.75; Found: C, 55.39; H, 5.74; N, 9.62.

Example 10**3-Benzenesulfonyl-5-(2-morpholin-4-yl-ethoxy)-1H-indazole**


A solution of toluene-4-sulfonic acid 2-(3-benzenesulfonyl-1H-indazol-5-yloxy)-ethyl ester (0.359 g, 0.760 mmol) and morpholine (1.0 mL, 11 mmol) in THF (8 mL) was stirred at 70 °C for 6 hours in a sealed tube. After cooling somewhat, the reaction mixture was solvent evaporated and partitioned in ethyl acetate and aqueous sodium bicarbonate. The organic phase was then washed with water and brine, dried with anhydrous magnesium sulfate, filtered and concentrated. The residue was purified by flash chromatography using 5 % methanol in chloroform. After drying *in vacuo* at 57 °C for 20 minutes, 3-benzenesulfonyl-5-(2-morpholin-4-yl-ethoxy)-1H-indazole as an off-white foam (0.0443 g, 15.1 %) was obtained. This was dissolved in methanol and chloroform, and ethereal hydrochloride was added. The mixture was concentrated and dried *in vacuo* at 75 °C for 16.5 hours to yield the hydrochloride as a light orange foam (0.0411 g). Mass Spectrum (+EI, [M+H]⁺) *m/z* 388. ¹HNMR (300 MHz, DMSO-*d*₆): δ14.31 (s, 1H), 10.82-8.8 (br, 1H), 8.03-8.07 (m, 2H), 7.62-7.76 (m, 4H), 7.49 (d, 1H, *J*=2.10 Hz), 7.26 (dd, 1H, *J*=9.14 Hz and 2.28 Hz), 4.54 (s, 2H), 4.02 (d, 2H, *J*=11.71 Hz), 3.80-3.87 (m, 2H), 3.51-3.73 (m, 4H), 3.19-3.31 ppm (m, 2H).

Example 11**3-Benzenesulfonyl-5-(2-pyrrolidin-1-yl-ethoxy)-1H-indazole**

A solution of toluene-4-sulfonic acid 2-(3-benzenesulfonyl-1H-indazol-5-yloxy)-ethyl ester (0.347 g, 0.734 mmol) and pyrrolidine (1.0 mL, 12 mmol) in THF (8 mL) was stirred at 70 °C in a sealed tube for 6 hours. After cooling somewhat, the reaction mixture was solvent evaporated and partitioned in ethyl acetate and

aqueous sodium bicarbonate. The organic phase was then washed with water and brine, dried with anhydrous magnesium sulfate, filtered and concentrated. The residue was purified by flash chromatography using 7.5 % methanol in chloroform. After drying at 55 °C *in vacuo*, 3-benzenesulfonyl-5-(2-pyrrolidin-1-yl-ethoxy)-1H-indazole as a pale yellow solid (0.0595 g, 21.8 %) was obtained. This was dissolved in methanol and chloroform, and ethereal hydrochloride was added. The mixture was concentrated and dried *in vacuo* at 75 °C for 16.5 hours to yield the hydrochloride as a yellow foam (0.0598 g). Mass Spectrum (+EI, [M+H]⁺) *m/z* 372. ¹HNMR (300 MHz, DMSO-d₆): δ 14.30 (s, 1H), 10.35 (br, 1H), 8.05 (d, 2H, *J*=7.04 Hz), 7.63-7.75 (m, 4H), 7.48 (d, 1H, *J*=1.83 Hz) 7.25-7.29 (m, 1H), 4.45-4.48 (m, 2H), 3.66 (s, br, 4H), 3.18-3.19 (m, 2H), 1.94-2.06 ppm (m, 4H).

Example 12




15 [2-(3-Benzenesulfonyl-1H-indazol-5-yloxy)-ethyl]-methyl-amine

A solution of toluene-4-sulfonic acid 2-(3-benzenesulfonyl-1H-indazol-5-yloxy)-ethyl ester (0.340 g, 0.720 mmol) in 2.0 M methylamine in THF (8 mL, 16 mmol) was stirred at 70 °C in a sealed tube for about 15.5 hours. After cooling somewhat, the reaction mixture was solvent evaporated and partitioned in chloroform and aqueous sodium bicarbonate. The organic phase was then washed with brine, dried with anhydrous magnesium sulfate, filtered and concentrated. The residue was purified by flash chromatography using 10 % methanol in chloroform and 1.0 % ammonium hydroxide/10% methanol in chloroform. After drying at 57 °C *in vacuo* for 20 minutes, [2-(3-benzenesulfonyl-1H-indazol-5-yloxy)-ethyl]-methyl-amine as a light yellow semi-solid (0.0461 g, 19.4 %) was obtained. This was dissolved in methanol and chloroform, and ethereal hydrochloride was added. The mixture was concentrated and dried *in vacuo* at 73 °C for 20 hours to yield the hydrochloride as a tan foam (0.0422 g). Mass Spectrum (+EI, [M+H]⁺) *m/z* 332. ¹HNMR (500 MHz, DMSO-d₆): δ 14.24 (s, 1H), 8.89 (s, 2H), 7.96-7.99 (m, 2H), 7.53-7.68 (m, 4H), 7.40

(d, 1H, $J=2.19$ Hz), 7.25 (dd, 1H, $J= 9.15$ Hz and 2.32 Hz), 4.30-4.32 (m, 2H), 3.33-3.35 (m, 2H), 2.60-2.63 ppm (m, 3H).

Example 13




2-(3-Benzenesulfonyl-1H-indazol-5-yloxy)-ethylamine

Over five different reactions, liquid ammonia (about 10 mL) was added to a – 78 °C solution of toluene-4-sulfonic acid 2-(3-benzenesulfonyl-1H-indazol-5-yloxy)-ethyl ester (1.72 g, 3.63 mmol) in THF (48 mL). This was usually allowed to warm to ambient temperature. This was heated in sealed tube for about 16-35 hours at 70-100 °C. After cooling to about ambient temperature, the combined reaction mixtures were then poured into excess sodium bicarbonate solution and extracted with chloroform or methylene chloride. The organic phase was washed with brine, dried with anhydrous magnesium sulfate, filtered and concentrated. The residue was purified by flash chromatography with 0.75 % ammonium hydroxide/7.5% methanol in chloroform. The residue was further purified by HPLC using 30-80% (chloroform/methanol(8:2)/TEA) in heptane/TEA. Concentrating and drying yielded 2-(3-benzenesulfonyl-1H-indazol-5-yloxy)-ethylamine as a white solid to which was added methanol and ethereal hydrochloride. Concentration and drying for 30 hours *in vacuo* at 83 °C yielded the hydrochloride as a cream-colored solid (0.0951 g, 7.4 %): MP: >300 °C; Mass Spectrum (–EI, $[M-H]^+$) m/z 316. ^1H NMR (500 MHz, DMSO- d_6): δ 14.24(s, 1H), 8.09 (br,s, 3H), 7.96-7.98 (m, 2H), 7.56-7.68 (m, 4H), 7.39 (d, 1H, $J=2.20$ Hz), 7.17 (dd, 1H, $J= 9.15$ Hz and 2.32 Hz), 4.22-4.25 (m, 2H), 3.20-3.24 ppm (m, 2H). Elemental Analysis for $\text{C}_{15}\text{H}_{15}\text{N}_3\text{O}_3\text{S} \cdot 1.00$ mol HCl $\cdot 0.85$ mol H_2O : Calcd: C, 48.81; H, 4.83; N, 11.38; Found: C, 48.65; H, 4.77; N, 11.00.

Example 14

Dimethyl-{3-[3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-propyl}-amine

Step 1



1-Chloromethanesulfonyl-naphthalene

A mixture of 1-naphthalene sulfonyl chloride (20.2 g, 89.1 mmol), sodium sulfite (22.5 g, 178 mmol) and sodium bicarbonate (15.1 g, 180 mmol) in water (125 mL) was stirred at 100 °C for one hour. After cooling to ambient temperature for 40 minutes, bromochloromethane (90 mL, 1.4 mol) and tetrabutylammonium bromide (2.87 g, 8.91 mmol) were added. The reaction mixture was then stirred at 75 °C for 14.5 hours. After cooling to ambient temperature, the layers of the reaction mixture were separated, and the organic phase was concentrated. The residue was purified by flash chromatography with 100 % ethyl acetate. Hexane was added after concentration to help solidify, and the mixture was again concentrated. Drying at 80 °C *in vacuo* for 20 minutes yielded 1-chloromethanesulfonyl-naphthalene as a pale yellow solid (19.0 g, 88.8 %). MP 103-5 °C. Mass Spectrum (+EI, M⁺) *m/z* 240. ¹HNMR (500 MHz, DMSO-*d*₆): δ 8.64-5 (m, 1H), 8.41 (d, 1H, *J*=8.23 Hz), 8.27 (dd, 1H, *J*=7.33 Hz and 1.22 Hz), 8.16-8.18 (m, 1H), 7.71-7.81 (m, 3H), 5.40 ppm (s, 2H). Elemental Analysis for C₁₁H₉ClO₂S: Calcd: C, 54.89; H, 3.77; N, 0.00; Found: C, 54.98; H, 3.81; N, 0.00.

Step 2




1-(5-Fluoro-2-nitro-phenylmethanesulfonyl)-naphthalene

To a chilled mixture of 1-chloromethanesulfonyl-naphthalene (19.7 g, 81.8 mmol) and 1-fluoro-4-nitrobenzene (8.7 mL, 82 mmol) in dry THF (197 mL) was dropwise added 1.0M potassium *t*-butoxide in THF (205 mL, 205 mmol). The reaction mixture was stirred at ambient temperature under nitrogen for 1.5 hours. Glacial acetic acid (16 mL, 280 mmol) was then added. After stirring at ambient temperature for 1 hour, 40 minutes, the reaction mixture was concentrated and partitioned in warm ethyl acetate and brine. The organic phase was dried with anhydrous magnesium sulfate, filtered and concentrated. The solid residue was

stirred in ether, filtered and dried at 82 °C *in vacuo* for 1 hour. This yielded 1-(5-fluoro-2-nitro-phenylmethanesulfonyl)-naphthalene as a brown/rust-colored solid (19.9 g, 70.6 %): MP: 155-160 °C; Mass Spectrum (-EI, [M-H]⁻) *m/z* 344. ¹HNMR (300 MHz, DMSO-d₆): δ 8.50-8.52 (m, 1H), 8.33 (d, 1H, *J*=8.30 Hz), 8.06-8.14 (m, 2H), 7.97 (dd, 1H, *J*=7.32 Hz and 1.22 Hz), 7.62-7.74 (m, 3H), 7.44-7.48 (m, 1H), 7.23-7.26 (m, 1H), 5.23 ppm (s, 2H). Elemental Analysis for C₁₇H₁₂FNO₄S: Calcd: C, 59.12; H, 3.50; N, 4.06; Found: C, 58.77; H, 3.30; N, 3.92.


Step 3:



3-[3-(Naphthalene-1-sulfonylmethyl)-4-nitro-phenoxy]-propan-1-ol

To a mixture of 1-(5-fluoro-2-nitro-phenylmethanesulfonyl)-naphthalene (19.9g, 57.6 mmol) and 1,3-propanediol (49 mL, 680 mmol) in dry THF (17 mL) was dropwise added 1.0M potassium *tert*-butoxide in THF (123 mL, 123 mmol). The reaction mixture was refluxed under nitrogen for 1 hour. It was allowed to cool to ambient temperature. Water was added to the reaction mixture, and it was poured into a mixture of ice and 2.0N hydrochloric acid. It was then extracted with ethyl acetate. The organic phase was washed with brine, dried with anhydrous magnesium sulfate, filtered and concentrated. The residue was purified by flash chromatography using 60% ethyl acetate in hexane and 100% ethyl acetate. The resulting yellow solid (12.9 g, 55.8 % yield), 3-[3-(naphthalene-1-sulfonylmethyl)-4-nitro-phenoxy]-propan-1-ol, was obtained: MP: 134-5 °C; Mass Spectrum (-EI, [M-H]⁻) *m/z* 400. ¹HNMR (500 MHz, DMSO-d₆): δ 8.50-8.53 (m, 1H), 8.31 (d, 1H, *J*=8.17 Hz), 8.10-8.12 (m, 1H), 7.95-7.99 (m, 2H), 7.61-7.71 (m, 3H), 7.06 (dd, 1H, *J*=9.15 Hz and 2.80 Hz), 6.78 (d, 1H, *J*=2.81 Hz), 5.23 (s, 2H), 4.52-4.55 (m, 1H), 3.91-3.94 (m, 2H), 3.43-3.47 (m, 2H), 1.71-1.78 ppm (m, 2H). Elemental Analysis for C₂₀H₁₉NO₆S: Calcd: C, 59.84; H, 4.77; N, 3.49; Found: C, 59.78; H, 4.41; N, 3.43.

Step 4



Toluene-4-sulfonic acid 3-[3-(naphthalene-1-sulfonylmethyl)-4-nitro-phenoxy]-propyl ester

- 5 A solution of 3-[3-(naphthalene-1-sulfonylmethyl)-4-nitro-phenoxy]-propan-1-ol (12.9 g, 32.1 mmol), *p*-toluenesulfonyl chloride (24.6 g, 129 mmol) and triethylamine (37 mL, 270 mmol) in methylene chloride (300 mL) was stirred at ambient temperature under nitrogen. After 2 hours, a second portion of *p*-toluenesulfonyl chloride (12.1 g, 63.5 mmol) was added. The reaction was stirred for
- 10 2 more hours and then concentrated. The residue was partitioned in methylene chloride and aqueous sodium bicarbonate. The organic phase was washed with brine, dried with anhydrous magnesium sulfate, filtered and concentrated. The residue was purified by flash chromatography with 20-50% ethyl acetate in hexane, 100 % ethyl acetate and then 100 % methylene chloride (due to limited solubility). A
- 15 yellow solid (13.0 g, 73.0 %) was obtained as toluene-4-sulfonic acid 2-[3-(naphthalene-1-sulfonylmethyl)-4-nitro-phenoxy]-propyl ester: MP: 162-4 °C; Mass Spectrum (-EI, [M-H]⁺) *m/z* 554. ¹HNMR (500 MHz, DMSO-*d*₆): 8.52-8.54 (m, 1H), 8.31 (d, 1H, *J*=8.18 Hz), 8.09-8.11 (m, 1H), 7.97-8.00 (m, 2H), 7.62-7.73 (m, 5H), 7.35 (d, 2H, *J*=7.93 Hz), 6.92 (dd, 1H, *J*=9.15 Hz and 2.81 Hz), 6.73 (d, 1H, *J*=2.80
- 20 Hz), 5.22 (s, 2H), 4.09 (t, 2H, *J*= 5.98 Hz), 3.79-3.82 (m, 2H), 2.29 (s, 3H), 1.91-1.97 ppm (m, 2H). Elemental Analysis for C₂₇H₂₅NO₈S₂: Calcd: C, 58.37; H, 4.54; N, 2.52; Found: C, 58.13; H, 4.43; N, 2.41.


Step 5



Toluene-4-sulfonic acid 3-[4-amino-3-(naphthalene-1-sulfonylmethyl)-phenoxy]-propyl ester

Ethanol (30 mL) and 10 % palladium on carbon (0.439 g) were added to a hot solution of toluene-4-sulfonic acid 2-[3-(naphthalene-1-sulfonylmethyl)-4-nitro-phenoxy]-propyl ester (0.785 g, 1.41 mmol) in hot, anhydrous THF (40 mL). The reaction mixture was hydrogenated on the Parr apparatus for 2 hours (starting pressure of 43 psi). The reaction mixture was then filtered over Celite and concentrated. It was dried at 75 °C *in vacuo* for 20 minutes to yield toluene-4-sulfonic acid 3-[4-amino-3-(naphthalene-1-sulfonylmethyl)-phenoxy]-propyl ester as a brown-green semi-solid (0.627 g, 84.4 %). Mass Spectrum (+EI, [M+H]⁺) *m/z* 526. ¹HNMR (500 MHz, DMSO-d₆): δ 8.61 (d, 1H, *J*=8.42 Hz), 8.25 (d, 1H, *J*=8.17 Hz), 8.02-8.07 (m, 2H), 7.59-7.72 (m, 5H), 7.39 (d, 2H, *J*=7.93 Hz), 6.56 (d, 1H, *J*=8.17 Hz), 6.44-6.47 (m, 1H), 6.01 (d, 1H, *J*=2.81 Hz), 4.62 (s, 2H), 3.98-4.01 (m, 2H), 3.36-3.39 (m, 2H), 2.35 (s, 3H), 1.71-1.79 ppm (m, 2H).

Step 6




Toluene-4-sulfonic acid 3-[3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-propyl ester

Isopropanol (205 mL) and 1.0N hydrochloric acid (200 mL) were added to toluene-4-sulfonic acid 3-[4-amino-3-(naphthalene-1-sulfonylmethyl)-phenoxy]-propyl ester (8.1 g, 15 mmol). The reaction mixture was heated because of limited solubility. Sodium nitrite (1.6 g, 2.3 mmol) in water (22 mL) was then added. More isopropanol (100 mL) was added to reaction mixture for increased solubility. After stirring at ambient temperature for 1 hour, sodium carbonate was added to basic pH. This reaction mixture was then stirred at ambient temperature for 30 minutes. After concentrating, the residue was partitioned in ethyl acetate and water. The organic phase was then washed with brine, dried with anhydrous magnesium sulfate, filtered and concentrated. The residue was purified by flash chromatography using 50 % ethyl acetate in hexane. A light amber semi-solid (3.3 g, 41 %), toluene-4-sulfonic acid 3-[3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-propyl ester, was obtained. Mass Spectrum (-EI, [M-H]⁻) *m/z* 535. ¹HNMR (500 MHz, DMSO-d₆): δ 14.08 (s, 1H),


8.78 (d, 1H, $J=8.78$ Hz), 8.55 (dd, 1H, $J=7.44$ Hz and 1.22 Hz), 8.28 (d, 1H, $J=8.17$ Hz), 8.02-8.04 (m, 1H), 7.73-7.77 (m, 1H), 7.55-7.65 (m, 4H), 7.48 (d, 1H, $J=9.15$ Hz), 7.03-7.06 (m, 3H), 6.88-6.91 (m, 1H), 4.17 (t, 2H, $J=5.98$ Hz), 3.85-3.88 (m, 2H), 2.00-2.03 (m, 2H), 1.97 ppm (s, 3H). Elemental Analysis for $C_{27}H_{24}N_2O_6S_2$ 0.10 mol H_2O : Calcd: C, 60.23; H, 4.53; N, 5.20; Found: C, 59.92; H, 4.30; N, 5.17.

Step 7




Dimethyl-{3-[3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-propyl}-amine

A solution of toluene-4-sulfonic acid 3-[3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-propyl ester (1.02 g, 1.89 mmol) in 2.0 M dimethylamine in THF (20 mL, 40 mmol) was stirred for 16 hours at 70 °C in a sealed tube. After cooling to ambient temperature, the reaction mixture was concentrated and partitioned in chloroform and aqueous sodium bicarbonate. The organic phase was washed with brine, dried with anhydrous magnesium sulfate, filtered and concentrated. The residue was purified by flash chromatography using 0.5 % ammonium hydroxide/5.0 % methanol in chloroform. Dimethyl-{3-[3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-propyl}-amine was obtained as a light yellow solid (0.453 g, 58.5 %). Methanol and ethereal hydrochloride were then added to this compound. The resulting solution was concentrated and dried for 14 hours at 84 °C *in vacuo*. Dimethyl-{3-[3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-propyl}-amine hydrochloride as a very light yellow solid (0.455g) was obtained: MP: 264-5 °C; Mass Spectrum (-EI, $[M-H]^+$) m/z 408. 1H NMR (500 MHz, DMSO- d_6): δ 14.17(s, 1H), 10.19 (s, 1H), 8.75 (d, 1H, $J=8.67$ Hz), 8.53 (dd, 1H, $J=7.44$ Hz and 1.22 Hz), 8.27 (d, 1H, $J=8.30$ Hz), 8.02-8.04 (m, 1H), 7.71-7.75 (m, 1H), 7.54-7.65 (m, 3H), 7.29 (d, 1H, $J=2.08$ Hz), 7.07-7.10 (m, 1H), 4.07-4.10 (m, 2H), 3.19-3.23 (m, 2H), 2.76 (s, 6H), 2.11-2.18 ppm (m, 2H). Elemental Analysis for $C_{22}H_{23}N_3O_3S$ · 1.00 mol HCl: Calcd: C, 59.25; H, 5.42; N, 9.42; Found: C, 58.97; H, 5.58; N, 9.23.


Example 15**Methyl-{3-[3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-propyl}-amine**

A solution of toluene-4-sulfonic acid 3-[3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-propyl ester (0.422 g, 0.786 mmol) in 2.0 M methylamine in THF (8.0 mL, 16 mmol). was stirred for 15 hours at 70 °C in a sealed tube. After cooling to ambient temperature, the reaction mixture was concentrated and partitioned in ethyl acetate and aqueous sodium bicarbonate. The organic phase was washed with brine. The aqueous phases were extracted with chloroform, and both organic phases were combined and dried with anhydrous magnesium sulfate, filtered and concentrated. The residue was purified by flash chromatography using 0.75 % ammonium hydroxide/7.5 % methanol in chloroform. Methyl-{3-[3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-propyl}-amine was obtained as a light yellow foam (0.209 g, 67.2 %). Methanol and ethereal hydrochloride were added to the compound. The resulting solution was concentrated and dried for 14 hours at 84 °C *in vacuo*. The hydrochloride was obtained as a light yellow foam (0.217g). Mass Spectrum (-EI, [M-H]⁻) *m/z* 394. ¹HNMR (500 MHz, DMSO-d₆): δ 14.16 (s, 1H), 8.74-8.76 (m, 1H), 8.67 (s, br, 2H), 8.52 (dd, 1H, *J*=7.44 Hz and 1.22 Hz), 8.27 (d, 1H, *J*=8.30 Hz), 8.02-8.05 (m, 1H), 7.70-7.75 (m, 1H), 7.54-7.65 (m, 3H), 7.29 (d, 1H, *J*=2.19 Hz), 7.08-7.11 (m, 1H), 4.09 (t, 2H, *J*=5.98 Hz), 3.05 (s, br, 2H), 2.55 (s, 3H), 2.05-2.11 ppm (m, 2H). Elemental Analysis for C₂₁H₂₁N₃O₃S · 1.00 mol HCl · 0.45 mol H₂O: Calcd: C, 57.32; H, 5.25; N, 9.55; Found: C, 57.64; H, 5.33; N, 9.39.

Example 16**Ethyl-{3-[3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-propyl}-amine**

A solution of toluene-4-sulfonic acid 3-[3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-propyl ester (0.080 mg, 0.15 mmol) in 10 mL of 2.0 M ethylamine/THF (20 mmols) was stirred at 90 °C for about 2 hours in a sealed tube. After cooling to ambient temperature the reaction mixture was solvent evaporated. It was dissolved in ethyl acetate and washed twice with aqueous sodium bicarbonate. The organic phase was dried over anhydrous magnesium sulfate, filtered and concentrated. The residue was purified by flash chromatography using 0.75 % ammonium hydroxide/7.5 % methanol in chloroform. Drying at 66 °C *in vacuo* for 20 minutes yielded ethyl-
 5 [3-[3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-propyl]-amine as a light yellow foam (43.0 mg, 70.0%). This was dissolved in methanol and ethereal hydrochloride was added. Concentrating and drying *in vacuo* at 83 °C for 16 hours gave the hydrochloride as a pale yellow semi-solid (41.7 mg, 62.3%). Mass Spectrum (+EI, [M+H]⁺) *m/z* 410; ¹HNMR (500 MHz, DMSO-*d*₆): δ14.14-14.18 (br, 1H), 8.74-8.76 (m, 1H), 8.56-8.72 (br, 2H), 8.52 (dd, 1h, *J*=7.44 Hz and 1.22 Hz), 8.27 (d, 1H, *J*=8.29 Hz), 8.03-8.05 (m, 1H), 7.71-7.75 (m, 1H), 7.54-7.65 (m, 3H), 7.29 (d, 1H, *J*=2.08 Hz), 7.07-7.10 (m, 1H), 4.09-4.12 (m, 2H), 3.03-3.07 (m, 2H), 2.90-2.97 (m, 2H), 2.05-2.12 (m, 2H), 1.15-1.19 ppm (m, 3H). Elemental Analysis for C₂₂H₂₃N₃O₃S · 1.00 mol HCl · 1.20 mol H₂O: Calcd: C, 56.51; H, 5.69; N, 8.99; Found: C, 56.20; H, 5.36; N, 8.81.


20

Example 17**Ethyl-methyl-3-[3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-propyl-amine**

A solution of toluene-4-sulfonic acid 3-[3-(naphthalene-1-sulfonyl)-1H-indazol-
 25 5-yloxy]-propyl ester (0.080 mg, 0.15 mmol) and methylethylamine (0.45-0.75 mmol) was stirred at 90 °C for about 2 hours in a sealed tube. After cooling to ambient temperature the reaction mixture was solvent evaporated. It was dissolved in ethyl acetate and washed twice with aqueous sodium bicarbonate. The organic phase was dried over anhydrous magnesium sulfate, filtered and concentrated. The
 30 residue was dissolved in methanol, and ethereal hydrochloride was added. The


mixture was then concentrated and dried for 29 hours at 80°C *in vacuo*. Ethyl-methyl-3-[3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-propyl-amine hydrochloride was a tan semi-solid (53.2 mg, 77.1%); Mass Spectrum (+EI, [M+H]⁺) *m/z* 424. ¹HNMR (500 MHz, DMSO-d₆): δ 14.14-14.17 (br, 1H), 9.71-9.77 (br, 1H), 8.74-8.76 (m, 1H), 8.52 (dd, 1H, *J*=7.44 Hz and 1.22 Hz), 8.27 (d, 1H, *J*=8.30 Hz), 8.03-8.05 (m, 1H), 7.70-7.74 (m, 1H), 7.56-7.64 (m, 3H), 7.29 (d, 1H, *J*=2.19 Hz), 7.08-7.10 (m, 1H), 4.08-4.11 (m, 2H), 3.04-3.25 (m, 2H), 2.72 (s, 3H), 1.18-1.21 ppm (m, 3H). Elemental Analysis for C₂₃H₂₅N₃O₃S · 1.00 mol HCl · 0.90 mol H₂O: Calcd: C, 58.01; H, 5.88; N, 8.82; Found: C, 58.37; H, 5.55; N, 8.50.

Example 18




Diethyl-3-[3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-propyl-amine

A solution of toluene-4-sulfonic acid 3-[3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-propyl ester (0.080 mg, 0.15 mmol) and diethylamine (0.45-0.75 mmol) in THF (10 mL) was stirred at 90 °C for about 2 hours in a sealed tube. After cooling to ambient temperature the reaction mixture was solvent evaporated. It was dissolved in ethyl acetate and washed twice with aqueous sodium bicarbonate. The organic phase was dried over anhydrous magnesium sulfate, filtered and concentrated. The residue was dissolved in methanol and chloroform, and ethereal hydrochloride was added. The mixture was concentrated and dried *in vacuo* at 80 °C for 17 hours. Diethyl-3-[3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-propyl-amine hydrochloride was a tan foam (69.9 mg, 98.3 %); Mass Spectrum (+EI, [M+H]⁺) *m/z* 438. ¹HNMR (500 MHz, DMSO-d₆): δ 14.13-14.11 (br, 1H), 9.59-9.63 (br, 1H), 8.75 (d, 1H, *J*=8.42 Hz), 8.51 (dd, 1H, *J*=7.44 Hz and 1.10 Hz), 8.27 (d, 1H, *J*=8.29 Hz), 8.03-8.05 (m, 1H), 7.70-7.74 (m, 1H), 7.52-7.64 (m, 3H), 7.05-7.11 (m, 2H), 4.10 (t, 2H, *J*=5.98 Hz), 3.11-3.25 (m, 6H), 2.09-2.24 (m, 2H), 1.18 ppm (t, 6H, *J*=7.20 Hz).

Example 19**Butyl-{3-[3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-propyl}-amine**


A solution of toluene-4-sulfonic acid 3-[3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-propyl ester (0.080 mg, 0.15 mmol) and butylamine (0.45-0.75 mmol) in THF (10 mL) was stirred at 90 °C for about 2 hours in a sealed tube. After cooling to ambient temperature the reaction mixture was solvent evaporated. It was dissolved in ethyl acetate and washed twice with aqueous sodium bicarbonate. The organic phase was dried over anhydrous magnesium sulfate, filtered and concentrated. The residue was dissolved in methanol, and ethereal hydrochloride was added. Concentrating and drying *in vacuo* at 80 °C for 17 hours yielded butyl-{3-[3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-propyl}-amine hydrochloride as a light orange semi-solid (43.3 mg, 60.9 %); Mass Spectrum (+EI, [M+H]⁺) *m/z* 438. ¹HNMR (500 MHz, DMSO-*d*₆): δ 13.8-14.2 (br, 1H), 8.75 (d, 1H, *J*=8.66 Hz), 8.50-8.52 (m, 1H), 8.27 (d, 1H, *J*=8.30 Hz), 8.03-8.05 (m, 1H), 7.71-7.75 (m, 1H), 7.54-7.64 (m, 3H), 7.29 (d, 1H, *J*=2.07 Hz), 7.07-7.10 (m, 1H), 4.08-4.11 (m, 2H), 3.04-3.08 (m, 2H), 2.86-2.90 (m, 2H), 2.06-2.13 (m, 2H), 1.52-1.59 (m, 2H), 1.26-1.33 (m, 4H), 0.88-0.84 ppm (m, 3H).

Example 20**Cyclopropyl-{3-[3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-propyl}-amine**

A solution of toluene-4-sulfonic acid 3-[3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-propyl ester (0.080 mg, 0.15 mmol) and cyclopropylamine (0.45-0.75 mmol) in THF (10 mL) was stirred at 90 °C for about 2 hours in a sealed tube. After cooling to ambient temperature the reaction mixture was solvent evaporated. It was dissolved in ethyl acetate and washed twice with aqueous sodium bicarbonate. The

organic phase was dried over anhydrous magnesium sulfate, filtered and concentrated. The residue was purified twice by flash chromatography using 0.75% ammonium hydroxide/7.5 % methanol in chloroform and 0.5 % ammonium hydroxide/5.0 % methanol in chloroform to give cyclopropyl-{3-[3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-propyl}-amine as a clear yellow film (51.0 mg, 80.7%). This compound was dissolved in methanol, and ethereal hydrochloride was added. Concentrating and drying *in vacuo* for 24 hours at 83 °C gave the hydrochloride as a yellow foam (48.9 mg, 71.2%); Mass Spectrum (+EI, [M+H]⁺) *m/z* 422. ¹HNMR (500 MHz, DMSO-d₆): δ 14.15 (s, 1H), 9.01 (s, 2H), 8.74-8.76 (m, 1H), 8.51-8.53 (m, 1H), 8.27 (d, 1H, *J*=8.29 Hz), 8.02-8.05 (m, 1H), 7.71-7.75 (m, 1H), 7.51-7.65 (m, 3H), 7.29 (d, 1H, *J*=2.19 Hz), 7.08-7.11 (m, 1H), 4.11 (t, 2H, *J*=6.10 Hz), 3.11-3.20 (br, s, 2H), 2.65-2.75 (m, 1H), 2.08-2.15 (m, 2H), 0.83-0.87 (m, 2H), 0.70-0.77 ppm (m, 2H). Elemental Analysis for C₂₃H₂₃N₃O₃S · 1.00 mol HCl · 0.70 mol H₂O: Calcd: C, 58.70; H, 5.44; N, 8.93; Found: C, 58.70; H, 5.09; N, 8.68.

Example 21




Cyclopentyl-{3-[3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-propyl}-amine

A solution of toluene-4-sulfonic acid 3-[3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-propyl ester (0.080 mg, 0.15 mmol) and cyclopentylamine (0.45-0.75 mmol) in THF (10 mL) was stirred at 90 °C for 2 hours in a sealed tube. After cooling to ambient temperature, the reaction mixture was solvent evaporated. It was partitioned in ethyl acetate and washed twice with aqueous sodium bicarbonate. The organic phase was dried over anhydrous magnesium sulfate, filtered and concentrated. The residue was purified by flash chromatography using 0.75% ammonium hydroxide/7.5% methanol in chloroform to give cyclopentyl-{3-[3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-propyl}-amine as a beige foam (70.6 mg, 100%). This compound was dissolved in methanol, and ethereal hydrochloride was added. Concentrating and drying *in vacuo* for 12.5 hours at 83 °C gave the hydrochloride as a light brown foam (69.7 mg, 95.6%); Mass Spectrum (+EI, [M+H]⁺) *m/z* 450. ¹HNMR


(500 MHz, DMSO- d_6): δ 14.15-14.21 (br, 1H), 8.70-8.83 (m, 3H), 8.51-8.53 (m, 1H), 8.27 (d, 1H, $J=8.30$ Hz), 8.03-8.05 (m, 1H), 7.71-7.75 (m, 1H), 7.54-7.65 (m, 3H), 7.30 (d, 1H, $J=2.20$ Hz), 7.09 (dd, 1H, $J=9.15$ Hz and 2.31 Hz), 4.11 (t, 2H, $J=6.10$ Hz), 3.43-3.49 (m, 1H), 3.06 (br, s, 2H), 2.05-2.14 (m, 2H), 1.88-1.96 (m, 2H), 1.43-1.71 ppm (m, 6H). Elemental Analysis for $C_{25}H_{27}N_3O_3S \cdot 1.00$ mol HCl: 0.75 mol H_2O : Calcd: C, 60.11; H, 5.95; N, 8.41; Found: C, 59.71; H, 5.83; N, 8.25.

Example 22




10 Isopropyl-{3-[3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-propyl}-amine

A solution of toluene-4-sulfonic acid 3-[3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-propyl ester (0.080 mg, 0.15 mmol) and isopropylamine (0.45-0.75 mmol) in THF (10 mL) was stirred at 90 °C for about 2 hours in a sealed tube. After cooling to ambient temperature the reaction mixture was solvent evaporated. It was dissolved in ethyl acetate and washed twice with aqueous sodium bicarbonate. The organic phase was dried over anhydrous magnesium sulfate, filtered and concentrated. The residue was purified by flash chromatography using 0.75 % ammonium hydroxide/7.5 % methanol in chloroform. The compound was dissolved in methanol and ethereal hydrochloride was added. Concentrating and drying for *in vacuo* 16 hours 83 °C gave isopropyl-{3-[3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-propyl}-amine hydrochloride as a light yellow foam (48.6 mg, 70.4%); Mass Spectrum (+EI, $[M+H]^+$) m/z 424. 1H NMR (500 MHz, DMSO- d_6): δ 14.15-14.16 (br, 1H), 8.75 (d, 1H, $J=8.78$ Hz), 8.60-8.70 (br, 2H), 8.52 (dd, 1H, $J=7.45$ Hz and 1.22 Hz), 8.27 (d, 1H, $J=8.29$ Hz), 8.03-8.05 (m, 1H), 7.71-7.75 (m, 1H), 7.54-7.65 (m, 3H), 7.30 (d, 1H, $J=2.20$ Hz), 7.08-7.11 (m, 1H), 4.09-4.13 (m, 2H), 3.05 (s, br, 2H), 2.07-2.13 (m, 2H), 1.22 ppm (d, 6H, $J=6.46$ Hz) Elemental Analysis for $C_{23}H_{25}N_3O_3S \cdot 1.00$ mol HCl: 0.55 mol H_2O : Calcd: C, 58.79; H, 5.81; N, 8.94; Found: C, 58.46; H, 5.61; N, 8.72.

Example 23**3-(Naphthalene-1-sulfonyl)-5-(3-pyrrolidin-1-yl-propoxy)-1H-indazole**

A solution of toluene-4-sulfonic acid 3-[3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-propyl ester (0.080 mg, 0.15 mmol) and pyrrolidine (0.45-0.75 mmol) in THF (10 mL) was stirred at 90 °C for about 2 hours in a sealed tube. After cooling to ambient temperature the reaction mixture was solvent evaporated. It was dissolved in ethyl acetate and washed twice with aqueous sodium bicarbonate. The organic phase was dried over anhydrous magnesium sulfate, filtered and concentrated. The residue was dissolved in methanol and chloroform, and ethereal hydrochloride was added. Concentrating and drying *in vacuo* at 80 °C for 17 hours gave 3-(naphthalene-1-sulfonyl)-5-(3-pyrrolidin-1-yl-propoxy)-1H-indazole hydrochloride as a brown semi-solid (19.7 mg, 27.8%); Mass Spectrum (+EI, [M+H]⁺) *m/z* 436. ¹HNMR (500 MHz, DMSO-*d*₆): δ 14.13 (s, 1H), 10.00-10.03 (br, 1H), 8.75 (d, 1H, *J*=8.66 Hz), 8.51-8.53 (m, 1H), 8.27 (d, 1H, *J*=8.30 Hz), 8.03-8.05 (m, 1H), 7.71-7.74 (m, 1H), 7.54-7.64 (m, 3H), 7.29 (d, 1H, *J*=2.20 Hz), 7.07-7.10 (m, 1H), 4.08-4.11 (m, 2H), 3.51-3.58 (m, 2H), 2.95-3.06 (m, 2H), 2.10-2.17 (m, 2H), 1.89-1.99 (m, 2H), 1.77-1.87 ppm (m, 2H).


Example 24**3-(Naphthalene-1-sulfonyl)-5-(3-piperidin-1-yl-propoxy)-1H-indazole**

A solution of toluene-4-sulfonic acid 3-[3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-propyl ester (0.080 mg, 0.15 mmol) and piperidine (0.45-0.75 mmol) in THF (10 mL) was stirred at 90 °C for about 2 hours in a sealed tube. After cooling to ambient temperature the reaction mixture was solvent evaporated. It was dissolved in ethyl acetate and washed twice with aqueous sodium bicarbonate. The organic

phase was dried over anhydrous magnesium sulfate, filtered and concentrated. The residue was dissolved in methanol, and ethereal hydrochloride was added.

Concentrating and drying at 80 °C for 29 hours *in vacuo* gave 3-(naphthalene-1-sulfonyl)-5-(3-piperidin-1-yl-propoxy)-1H-indazole hydrochloride as a brown semi-solid (49.0 mg, 67.2%); Mass Spectrum (-EI, [M-H]⁻) *m/z* 448. ¹HNMR (500 MHz, DMSO-d₆): δ14.14 (s, 1H), 9.59-9.66 (br, 1H), 8.75 (d, 1H, *J*=8.54 Hz), 8.51-8.53 (m, 1H), 8.27 (d, 1H, *J*=8.30 Hz), 8.03-8.05 (m, 1H), 7.71-7.74 (m, 1H), 7.54-7.65 (m, 3H), 7.29 (d, 1H, *J*=2.20 Hz), 7.08 (dd, 1H, *J*=9.15 Hz and 2.32 Hz), 4.09 (t, 2H, *J*=5.98 Hz), 3.42-3.45 (m, 2H), 3.16-3.25 (m, 2H), 2.82-2.96 (m, 2H), 2.13-2.29 (m, 2H), 1.58-1.80 (m, 5H), 1.30-1.40 ppm (m, 1H). Elemental Analysis for C₂₅H₂₇N₃O₃S · 1.00 mol HCl · 1.00 mol H₂O: Calcd: C, 59.57; H, 6.00; N, 8.34; Found: C, 59.36; H, 5.82; N, 8.24.

Example 25



5-(3-Morpholin-4-yl-propoxy)-3-(naphthalene-1-sulfonyl)-1H-indazole


A solution of toluene-4-sulfonic acid 3-[3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-propyl ester (0.080 mg, 0.15 mmol) and morpholine (0.45-0.75 mmol) in THF (10 mL) was stirred at 90 °C for about 2 hours in a sealed tube. After cooling to ambient temperature the reaction mixture was solvent evaporated. It was dissolved in ethyl acetate and washed twice with aqueous sodium bicarbonate. The organic phase was dried over anhydrous magnesium sulfate, filtered and concentrated. The residue was purified by flash chromatography using 0.5 % ammonium hydroxide/5.0% methanol in chloroform. 5-(3-Morpholin-4-yl-propoxy)-3-(naphthalene-1-sulfonyl)-1H-indazole as a light yellow solid (74.4 mg, 100%) was obtained. This product was dissolved in methanol, and ethereal hydrochloride was added. Concentrating and drying *in vacuo* 83 °C for 24 hours yielded the hydrochloride as a yellow foam (74.5 mg, 100 %); Mass Spectrum (+EI, [M+H]⁺) *m/z* 452. ¹HNMR (500 MHz, DMSO-d₆): δ14.15 (s, 1H), 10.56-10.62 (br, 1H), 8.75 (d, 1H, *J*=8.66 Hz), 8.52 (dd, 1H, *J*=7.44 Hz and 1.22 Hz), 8.27 (d, 1H, *J*=8.30 Hz), 8.02-

8.05 (m, 1H), 7.71-7.75 (m, 1H), 7.54-7.65 (m, 3H), 7.29 (d, 1H, $J=2.32$ Hz), 7.07-7.10 (m, 1H), 4.10 (t, 2H, $J=5.86$ Hz), 3.93-3.96 (m, 2H), 3.71-3.77 (m, 2H), 3.43-3.46 (m, 2H), 2.99-3.15 (m, 2H), 2.14-2.22 ppm (m, 2H). Elemental Analysis for $C_{24}H_{25}N_3O_5S \cdot 1.00$ mol HCl $\cdot 1.30$ mol H_2O : Calcd: C, 56.36; H, 5.36; N, 8.22; Found: C, 56.05; H, 5.65; N, 8.09.

Example 26

3-[3-(Naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-propylamine


Step 1



5-(3-Azido-propoxy)-3-(naphthalene-1-sulfonyl)-1H-indazole

Sodium azide (0.425 g, 6.54 mmol) was added to a solution of toluene-4-sulfonic acid 3-[3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-propyl ester (1.16 g, 2.16 mmol) in DMF (20 mL). The reaction mixture was stirred at 80 °C for 15 hours in a sealed tube. After cooling to ambient temperature, it was poured into excess water and extracted with ethyl acetate. The organic phase was washed with 10% ammonium chloride/water solution, water and brine. It was dried with anhydrous magnesium sulfate, filtered, concentrated and dried at 73 °C *in vacuo* for 45 minutes to yield 5-(3-azido-propoxy)-3-(naphthalene-1-sulfonyl)-1H-indazole as an amber gum (0.770 g, 87.7 %); Mass Spectrum (-EI, $[M-H]^+$) m/z 406. 1H NMR (400 MHz, $DMSO-d_6$): δ 14.05 (br, 1H), 8.64-8.68 (m, 1H), 8.53 (dd, 1H, $J=7.42$ Hz and 1.16 Hz), 8.25-8.28 (m, 1H), 8.01-8.05 (m, 1H), 7.70-7.74 (m, 1H), 7.52-7.64 (m, 3H), 7.26 (d, 1H, $J=2.08$ Hz), 7.08 (dd, 1H, $J=9.16$ Hz and 2.32 Hz), 4.06 (t, 2H, $J=6.14$ Hz), 3.49-3.52 (m, 2H), 1.96-1.99 ppm (m, 2H).

Step 2



3-[3-(Naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-propylamine

5-(3-Azido-propoxy)-3-(naphthalene-1-sulfonyl)-1H-indazole (0.763 g, 1.87 mmol) was dissolved in hot ethanol (125 mL). 10% Palladium on carbon (0.16 g) was added, and the reaction mixture was shaken and hydrogenated on the Parr apparatus for 2 hours, starting pressure 51 psi. Additional 10% Palladium on carbon (0.32 g) was added, and the reaction mixture was again hydrogenated on the Parr apparatus for 2 hours. It was then filtered over Celite and concentrated. The residue was purified by flash chromatography with 0.75% ammonium hydroxide/7.5% methanol in chloroform. The product was further purified by HPLC using a gradient of chloroform/methanol in heptane/TFA. 3-[3-(Naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-propylamine as a light yellow foam (0.229 g, 32.1%) resulted. Methanol and ethereal hydrochloride were added. The mixture was concentrated and dried at 83 °C *in vacuo* for 12 hours to yield the hydrochloride as a light buff foam (0.235 g). Mass Spectrum (-EI, [M-H]⁺) *m/z* 380. ¹HNMR (500 MHz, DMSO-d₆): δ 14.15 (s, 1H), 8.75 (d, 1H, *J*=8.67 Hz), 8.50-8.52 (m, 1H), 8.27 (d, 1H, *J*=8.30 Hz), 8.02-8.05 (m, 1H), 7.86 (s, br, 3H), 7.71-7.75 (m, 1H), 7.54-7.64 (m, 3H), 7.29 (d, 1H, *J*=2.20 Hz), 7.08-7.11 (m, 1H), 4.09 (t, 2H, *J*=6.10 Hz), 2.94-2.99 (m, 2H), 2.00-2.07 ppm (m, 2H). Elemental Analysis for C₂₀H₁₉N₃O₃S · 1.00 mol HCl · 0.40 mol H₂O: Calcd: C, 56.51; H, 4.93; N, 9.88; Found: C, 56.21; H, 5.12; N, 9.56.

Example 27**Dimethyl-{2-[3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-ethyl}-amine****Step 1****1-(5-Fluoro-2-nitro-phenylmethanesulfonyl)-naphthalene**


To a chilled mixture of 1-chloromethanesulfonyl-naphthalene (21.7 g, 90.1 mmol) and 1-fluoro-4-nitrobenzene (9.6 mL, 90 mmol) in dry THF (220 mL) was dropwise added 1.0 M potassium *tert*-butoxide in THF (193 mL, 193 mmol). The reaction mixture was stirred at ambient temperature under nitrogen for 1 hour, 10 minutes. Glacial acetic acid (17 mL, 300 mmol) was then added. The reaction mixture was concentrated and partitioned in warm ethyl acetate and brine. The

organic phase was dried with anhydrous magnesium sulfate, filtered and concentrated. The solid residue was stirred in ether, filtered and dried at 65 °C *in vacuo* for 16 hours. This yielded a brown/rust-colored solid (22.2 g, 71.4 %) as 1-(5-fluoro-2-nitro-phenylmethanesulfonyl)-naphthalene: MP: 155-156 °C; Mass

- 5 Spectrum (-EI, [M-H]⁺) *m/z* 344. ¹HNMR (300 MHz, DMSO-d₆): δ 8.50-8.52 (m, 1H), 8.33 (d, 1H, *J*=8.30 Hz), 8.06-8.14 (m, 2H), 7.97 (dd, 1H, *J*=7.32 Hz and 1.22 Hz), 7.62-7.74 (m, 3H), 7.44-7.48 (m, 1H), 7.23-7.26 (m, 1H), 5.23 ppm (s, 2H). Elemental Analysis for C₁₇H₁₂FNO₄S: Calcd: C, 59.12; H, 3.50; N, 4.06; Found: C, 58.77; H, 3.30; N, 3.92.

10


Step 2



2-[3-(Naphthalene-1-sulfonylmethyl)-4-nitro-phenoxy]-ethanol

- A mixture of 1-(5-fluoro-2-nitro-phenylmethanesulfonyl)-naphthalene (14.6 g, 42.3 mmol) and ethylene glycol (35 mL, 630 mmol) in 1.0 N potassium *tert*-butoxide in THF (90 mL, 90 mmol) was refluxed under nitrogen for one hour. After cooling to ambient temperature, the reaction mixture was concentrated. Excess water was added to the residue, and the mixture was poured into ice/2N hydrochloric acid. It was extracted with ethyl acetate and washed with water and brine. It was then dried with anhydrous magnesium sulfate, filtered, concentrated and dried *in vacuo* at 80 °C for 25 minutes to yield 2-[3-(naphthalene-1-sulfonylmethyl)-4-nitro-phenoxy]-ethanol as a brown gum (14.3 g, 87.2 %). Mass Spectrum (-EI, [M-H]⁺) *m/z* 386. ¹HNMR (500 MHz, DMSO-d₆): δ 8.56-8.57 (m, 1H), 8.35 (d, 1H, *J*=8.24 Hz), 8.14-8.15 (m, 1H), 8.00-8.02 (m, 2H), 7.65-7.75 (m, 3H), 7.11-7.13 (m, 1H), 6.89 (d, 1H, *J*=2.90 Hz), 5.27 (s, 2H), 4.92 (t, 1H, *J*=5.49 Hz), 3.93-3.95 (m, 2H), 3.67 ppm (dd, 2H, *J*=9.91 Hz and 5.34 Hz). Elemental Analysis for C₁₉H₁₇NO₆S·0.25 mol H₂O: Calcd: C, 58.23; H, 4.50; N, 3.57; Found: C, 57.83; H, 4.25; N, 3.50.


Step 3



Toluene-4-sulfonic acid 2-[3-(naphthalene-1-sulfonylmethyl)-4-nitro-phenoxy]-ethyl ester

- 5 A solution of 2-[3-(naphthalene-1-sulfonylmethyl)-4-nitro-phenoxy]-ethanol (14.3 g, 36.9 mmol), *p*-toluenesulfonylchloride (14.9 g, 78.2 mmol) and triethylamine (23 mL, 170 mmol) in methylene chloride (400 mL) was stirred under nitrogen at ambient temperature for 2 hours, 45 minutes. The reaction mixture was then concentrated and partitioned in methylene chloride and aqueous sodium bicarbonate.
- 10 The organic phase was washed with brine, dried with anhydrous magnesium sulfate, filtered and concentrated. The residue was purified by flash chromatography with 25-75 % ethyl acetate in hexane and 100 % ethyl acetate. Drying for 25 minutes *in vacuo* at 80 °C yielded toluene-4-sulfonic acid 2-[3-(naphthalene-1-sulfonylmethyl)-4-nitro-phenoxy]-ethyl ester as a rust-colored solid (15.7 g, 78.5 %). Mass Spectrum (+EI, [M+H]⁺) *m/z* 542. ¹HNMR (500 MHz, DMSO-*d*₆): 8.53 (dd, 1H, *J*= 8.18 Hz and 1.10 Hz), 8.31 (d, 1H, *J*=8.30 Hz), 8.09-8.11 (m, 1H), 7.94-7.99 (m, 2H), 7.62-7.75 (m, 5H), 7.42 (d, 2H, *J*=7.93 Hz), 6.98 (dd, 1H, *J*=9.15 Hz and 2.93 Hz), 6.80 (d, 1H, *J*=2.81 Hz), 5.20 (s, 2H), 4.28-4.30 (m, 2H), 4.10-4.12 (m, 2H), 2.37 ppm (s, 3H).
- 15

20 **Step 4**




Toluene-4-sulfonic acid 2-[4-amino-3-(naphthalene-1-sulfonylmethyl)-phenoxy]-ethyl ester

- 25 Ethanol (180 mL) and 10 % palladium on carbon (5.2 g) were added to a hot solution of toluene-4-sulfonic acid 2-[3-(naphthalene-1-sulfonylmethyl)-4-nitro-phenoxy]-ethyl ester (11.5 g, 21.2 mmol) in THF (180 mL). This mixture was shaken on the Parr hydrogenation apparatus for 5 hours, with a starting pressure of 47 psi. It was then filtered over Celite, concentrated and dried *in vacuo* at 80 °C for 30 minutes

to give toluene-4-sulfonic acid 2-[4-amino-3-(naphthalene-1-sulfonylmethyl)-phenoxy]-ethyl ester as a dark brown gum (9.5 g, 88.0 %). Mass Spectrum (+EI, $[M+H]^+$) m/z 512. ^1H NMR (500 MHz, $\text{DMSO}-d_6$): δ 8.61 (d, 1H, $J=8.41$ Hz), 8.25 (d, 1H, $J=8.30$ Hz), 8.03-8.08 (m, 2H), 7.59-7.75 (m, 5H), 7.43 (d, 2H, $J=7.93$ Hz), 6.49-6.53 (m, 2H), 6.11 (d, 1H, $J=2.68$ Hz), 4.65-4.77 (br, 2H), 4.59 (s, 2H), 4.09-4.11 (m, 2H), 3.63-3.65 (m, 2H), 2.38 ppm (s, 3H).


Step 5




10 Toluene-4-sulfonic acid 2-[3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-ethyl ester

Ethanol (430 mL) in two portions and 1 N hydrochloric acid (200 mL) were added to toluene-4-sulfonic acid 2-[4-amino-3-(naphthalene-1-sulfonylmethyl)-phenoxy]-ethyl ester (6.5 g, 13 mmol). A solution of sodium nitrite (1.5 g, 22 mmol) in water was then added to the reaction mixture. The reaction was heated to aid in solubility. After stirring at ambient temperature for 2 hours, solid sodium carbonate was added to basic pH. The reaction mixture was stirred at ambient temperature for one hour. It was then solvent evaporated and partitioned in water and ethyl acetate. The organic phase was washed with brine, dried with anhydrous magnesium sulfate, filtered and concentrated. The residue was purified by flash chromatography with 2% methanol in chloroform. Drying at 63 °C *in vacuo* yielded toluene-4-sulfonic acid 2-[3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-ethyl ester a buff-colored foam (3.9 g, 59 %); Mass Spectrum (+EI, $[M+H]^+$) m/z 523. ^1H NMR (500 MHz, $\text{DMSO}-d_6$): δ 14.09 (s, 1H), 8.76 (d, 1H, $J=8.78$ Hz), 8.52-8.55 (m, 1H), 8.27 (d, 1H, $J=8.30$ Hz), 8.02-8.04 (m, 1H), 7.64-7.74 (m, 3H), 7.55-7.63 (m, 2H), 7.50 (d, 1H, $J=9.15$ Hz), 7.27 (d, 2H, $J=8.05$ Hz), 7.14 (d, 1H, $J=2.20$ Hz), 6.94 (dd, 1H, $J=9.15$ Hz and 2.44 Hz), 4.32-4.34 (m, 2H), 4.18-4.20 (m, 2H), 2.25 ppm (s, 3H). Elemental Analysis for $\text{C}_{26}\text{H}_{22}\text{N}_2\text{O}_6\text{S}_2$: Calcd: C, 59.76; H, 4.24; N, 5.36; Found: C, 59.69; H, 4.28; N, 5.14.

Step 6

**Dimethyl-2-[3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-ethylamine**


A solution of toluene-4-sulfonic acid 2-[3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-ethyl ester (0.382 g, 0.731 mmol) in 2.0 M dimethylamine in THF (8 mL, 16 mmol) was stirred for 4 hours at 70 °C in a sealed tube. After cooling to ambient temperature, the reaction mixture was concentrated and partitioned in ethyl acetate and aqueous sodium bicarbonate. The organic phase was washed with water and brine, dried with anhydrous magnesium sulfate, filtered and concentrated. Drying at 80 °C for 20 minutes *in vacuo* yielded dimethyl-2-[3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-ethylamine as a light yellow solid (0.212 g, 73.4 %). Methanol, chloroform and etheric hydrochloride were added. The resulting solution was concentrated and dried for 15 hours at 78 °C *in vacuo*. The hydrochloride as a buff foam (0.217 g) was obtained. Mass Spectrum (+EI, [M+H]⁺) *m/z* 396. ¹HNMR (500 MHz, DMSO-d₆): δ 14.20 (s, 1H), 10.05 (s, 1H), 8.74-8.77 (m, 1H), 8.52 (dd, 1H, *J*=7.44 Hz and 1.22 Hz), 8.27 (d, 1H, *J*=8.17 Hz), 8.03-8.05 (m, 1H), 7.70-7.74 (m, 1H), 7.56-7.65 (m, 3H), 7.37 (d, 1H, *J*=2.20 Hz), 7.15-7.18 (m, 1H), 4.37-4.39 (m, 2H), 3.51 (t, 2H, *J*=4.76 Hz), 2.83 ppm (s, 6H). Elemental Analysis for C₂₁H₂₁N₃O₃S · 1.00 mol HCl · 0.70 mol H₂O: Calcd: C, 56.74; H, 5.31; N, 9.45; Found: C, 56.72; H, 5.33; N, 9.06.

Example 28**Isopropyl-2-[3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-ethylamine**

A solution of step 6 (1.01 g, 1.93 mmol) and isopropylamine (2.0 mL, 23 mmol) in THF (25 mL) was stirred for 16 hours at 70 °C in a sealed tube. More isopropylamine (2.0 mL, 23 mmol) was added, and the reaction mixture was stirred at

80 °C for 20 hours in a sealed tube. After cooling to ambient temperature, the reaction mixture was concentrated and partitioned in ethyl acetate and aqueous sodium bicarbonate. The organic phase was washed with brine, dried with anhydrous magnesium sulfate, filtered and concentrated. The residue was purified by flash chromatography using 0.75 % ammonium hydroxide/7.5% methanol in chloroform. After concentrating and drying *in vacuo* at 72 °C for 25 minutes, isopropyl-{2-[3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-ethyl}-amine was obtained as a light orange solid (0.313 g, 39.6 %). Methanol, chloroform and ethereal hydrochloride were added to this compound. The resulting solution was concentrated and dried for 16 hours at 84 °C *in vacuo*. The hydrochloride was a light brown foam (0.332g). Mass Spectrum (+EI, [M+H]⁺) *m/z* 410. ¹HNMR (500 MHz, DMSO-*d*₆): δ14.22 (s, 1H), 8.86 (s, 2H), 8.74-8.77 (m, 1H), 8.53 (dd, 1H, *J*=7.44 Hz and 1.22 Hz), 8.27 (d, 1H, *J*=8.30 Hz), 8.03-8.05 (m, 1H), 7.71-7.75 (m, 1H), 7.56-7.65 (m, 3H), 7.34 (d, 1H, *J*=2.32 Hz), 7.16 (dd, 1H, *J*=9.15 Hz and 2.44 Hz), 4.29-4.32 (m, 2H), 1.25 ppm (d, 6H, *J*=6.59 Hz). Elemental Analysis for C₂₂H₂₃N₃O₃S · 1.00 mol HCl · 0.35 mol H₂O: Calcd: C, 58.43; H, 5.50; N, 9.29; Found: C, 58.03; H, 5.25; N, 8.94.

Example 29



Ethyl-methyl-{2-[3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-ethyl}-amine


A solution of toluene-4-sulfonic acid 2-[3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-ethyl ester (1.18 g, 2.26 mmol) and ethylmethylamine (2.0 mL, 23 mmol) in THF (20 mL) was stirred for 16 hours at 70 °C in a sealed tube. More ethylmethylamine (2.0 mL, 23 mmol) was added, and the reaction mixture was heated to 80 °C for 20 hours in a sealed tube. After cooling to ambient temperature, the reaction mixture was concentrated and partitioned in ethyl acetate and aqueous sodium bicarbonate. The organic phase was washed with brine, dried with anhydrous magnesium sulfate, filtered and concentrated. The residue was purified by flash chromatography using 0.75 % ammonium hydroxide/7.5% methanol in

chloroform. Drying *in vacuo* at 69 °C for 20 minutes yielded ethyl-methyl-{2-[3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-ethyl}-amine as a buff foam (0.409 g, 44.2 %). Methanol, chloroform and ethereal hydrochloride were added. The resulting solution was concentrated and dried for 16 hours at 84 °C *in vacuo*. The hydrochloride was obtained as a buff semi-solid (0.424g). Mass Spectrum (+EI, [M+H]⁺) *m/z* 410. ¹HNMR (500 MHz, DMSO-*d*₆): δ 14.21 (s, 1H), 10.00-10.01 (s, 1H), 8.74-8.77 (m, 1H), 8.52 (dd, 1H, *J*= 7.44 Hz and 1.22 Hz), 8.27 (d, 1H, *J*= 8.29 Hz), 8.03-8.05 (m, 1H), 7.73 (dd, 1H, *J*=8.06 Hz and 7.57 Hz), 7.56-7.65 (m, 3H), 7.37 (d, 1H, *J*=2.20 Hz), 7.15-7.18 (m, 1H), 4.40 (t, 2H, *J*=5.00 Hz), 3.53-3.59 (m, 1H), 3.39-3.49 (m, 1H), 3.21-3.25 (m, 1H), 3.10-3.20 (m, 1H), 2.81 (d, 3H, *J*=4.88 Hz), 1.21-1.25 ppm (m, 3H). Elemental Analysis for C₂₂H₂₃N₃O₃S · 1.00 mol HCl · 0.30 mol H₂O: Calcd: C, 58.54; H, 5.49; N, 9.31; Found: C, 58.30; H, 5.52; N, 8.91.

Example 30

2-[3-(Naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-ethylamine

Step 1




5-(2-Azido-ethoxy)-3-(naphthalene-1-sulfonyl)-1H-indazole

Sodium azide (0.79 g, 12 mmol) was added to a solution of toluene-4-sulfonic acid 2-[3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-ethyl ester (2.06 g, 3.94 mmol) in DMF (20 mL). The reaction mixture was stirred at 80 °C for 12 hours in a sealed tube. After cooling to ambient temperature, it was poured into excess water and extracted with ethyl acetate. The organic phase was washed with 10% ammonium chloride/water solution, water and brine. It was dried with anhydrous magnesium sulfate, filtered, concentrated and dried at 72 °C *in vacuo* for 40 minutes to yield 5-(2-azido-ethoxy)-3-(naphthalene-1-sulfonyl)-1H-indazole as a amber gum/foam (1.34 g, 86.5 %). Mass spectrum (-EI, [M-H]⁻) *m/z* 392. ¹HNMR (400 MHz, DMSO-*d*₆): δ 8.78 (d, 1H, *J*=8.70 Hz), 8.56 (dd, 1H, *J*=7.43 Hz and 1.16 Hz), 8.29 (d, 1H, *J*=8.35 Hz), 8.06 (d, 1H, *J*=7.66 Hz), 7.72-7.76 (m, 1H), 7.57-7.67 (m, 3H), 7.31 (d, 1H, *J*=2.32

Hz), 7.12 (dd, 1H, $J = 9.16$ Hz and 2.32 Hz), 4.22-4.24 (m, 2H), 3.67-3.69 ppm (m, 2H).

Step 2




5

2-[3-(Naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-ethylamine

To 5-(2-azido-ethoxy)-3-(naphthalene-1-sulfonyl)-1H-indazole (1.34 g, 3.41 mmol) in hot ethanol (210 mL) was added 10 % palladium on carbon (0.8 g). The reaction mixture was hydrogenated on the Parr apparatus for 1.25 hours, starting pressure 49 psi. It was then filtered over Celite and concentrated. The residue was purified by flash chromatography with 0.5 % ammonium hydroxide/5.0 % methanol in chloroform. Drying *in vacuo* at 63 °C for 20 minutes gave 2-[3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-ethylamine as a light yellow foam resulted (0.775 g, 57.0 %). Methanol and ethereal hydrochloride were added. Drying *in vacuo* for 16 hours 68 °C yielded the hydrochloride as a yellow foam (0.825 g). Mass spectrum (-EI, $[M-H]^-$) m/z 366. $^1\text{H-NMR}$ (500 MHz, DMSO- d_6): δ 14.22 (s, 1H), 8.75 (d, 1H, $J = 8.66$ Hz), 8.51-8.53 (m, 1H), 8.27 (d, 1H, $J = 8.29$ Hz), 8.10 (s, br, 3H), 8.03-8.05 (m, 1H), 7.71-7.75 (m, 1H), 7.56-7.65 (m, 3H), 7.33 (d, 1H, $J = 2.19$ Hz), 7.13-7.16 (m, 1H), 4.21 (t, 2H, $J = 5.00$ Hz), 3.23 ppm (s, 2H). Elemental Analysis for $\text{C}_{19}\text{H}_{17}\text{N}_3\text{O}_3\text{S}$ 1.00 mol HCl: 0.55 mol H_2O : Calcd: C, 55.15; H, 4.65; N, 10.15; Found: C, 55.54; H, 4.90; N, 10.12.

20

Example 31




25 Methyl-2-[3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-ethylamine

A solution of toluene-4-sulfonic acid 2-[3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-ethyl ester (200 mg, 0.383 mmol) in 2.0 M methylamine in THF (10 mL, 20

mmol) was stirred at 70 °C for 2-3 hours in a sealed tube. After cooling to ambient temperature, the reaction mixture was solvent evaporated and partitioned in ethyl acetate and aqueous sodium bicarbonate. The organic phase was dried with anhydrous magnesium sulfate, filtered and concentrated. It was dissolved in methanol, and ethereal hydrochloride was added. The mixture was concentrated and dried for about 16 hours at 70 °C *in vacuo* to give methyl-{2-[3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-ethyl}-amine hydrochloride as a pale yellow foam (100 mg, 62.5 %); Mass Spectrum (-EI, [M-H]⁺) *m/z* 380. ¹HNMR (400 MHz, DMSO-d₆): δ 14.21 (br, s, 1H), 8.85-8.91 (s, 2H), 8.74-8.75 (m, 1H), 8.51-8.53 (m, 1H), 8.26-8.28 (m, 1H), 8.03-8.05 (m, 1H), 7.71-7.74 (m, 1H), 7.56-7.64 (m, 3H), 7.34 (d, 1H, *J*=2.20 Hz), 7.14-7.17 (m, 1H), 4.27-4.30 (m, 2H), 3.32-3.34 (m, 2H), 2.61 ppm (s, 3H). Elemental Analysis for C₂₀H₁₉N₃O₃S · 1.00 mol HCl · 1.40 H₂O: Calcd: C, 54.21; H, 5.19; N, 9.48; Found: C, 54.31; H, 4.80; N, 9.10.

Example 32




Ethyl-{2-[3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-ethyl}-amine

A solution of toluene-4-sulfonic acid 2-[3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-ethyl ester (200 mg, 0.383 mmol) in 2.0 M ethylamine (10 mL, 20 mmol) was stirred at 70 °C for 2-3 hours in a sealed tube. After cooling to ambient temperature, the reaction mixture was solvent evaporated and partitioned in ethyl acetate and aqueous sodium bicarbonate. The organic phase was dried with anhydrous magnesium sulfate, filtered and concentrated. It was dissolved in methanol, and ethereal hydrochloride was added. The mixture was concentrated and dried for 16 hours *in vacuo* at 70 °C to give ethyl-{2-[3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-ethyl}-amine hydrochloride as a pale yellow foam (30 mg, 18.2%). Mass Spectrum (-EI, [M-H]⁺) *m/z* 394. ¹HNMR (500 MHz, DMSO-d₆): δ 14.18 (s, 1H), 8.75 (d, 3H, *J*=8.54 Hz), 8.51-8.53 (m, 1H), 8.27 (d, 1H, *J*=8.17 Hz), 8.03-8.05 (m, 1H), 7.70-7.74 (m, 1H), 7.56-7.65 (m, 3H), 7.34 (d, 1H, *J*=2.20 Hz), 7.16 (dd, 1H, *J*=9.15

Hz and 2.32 Hz), 4.26-4.29 (m, 2H), 3.31-3.34 (m, 2H), 3.01-3.05 (s, br, 2H), 1.18-1.21 ppm (m, 3H).


Example 33



Diethyl-2-[3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-ethylamine

A solution of toluene-4-sulfonic acid 2-[3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-ethyl ester (200 mg, 0.383 mmol) and diethylamine (1.1-1.5 mmol) in THF (10 mL) was stirred at 70 °C for 2-3 hours in a sealed tube. After cooling to ambient temperature, the reaction mixture was solvent evaporated and partitioned in ethyl acetate and aqueous sodium bicarbonate. The organic phase was dried with anhydrous magnesium sulfate, filtered and concentrated. It was dissolved in methylene chloride and methanol, and ethereal hydrochloride was added. The mixture was concentrated and dried *in vacuo* at 67 °C for 16 hours to give diethyl-2-[3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-ethylamine hydrochloride as an off-white foam (148 mg, 84.1%); Mass Spectrum (-EI, [M-H]⁺) *m/z* 422. ¹HNMR (400 MHz, DMSO-d₆): δ 14.20 (s, 1H), 9.91 (s, 1H), 8.75 (d, 1H, *J*=8.54 Hz), 8.51-8.53 (m, 1H), 8.27 (d, 1H, *J*=8.29 Hz), 8.03-8.05 (m, 1H), 7.71-7.75 (m, 1H), 7.55-7.65 (m, 3H), 7.36 (d, 1H, *J*=2.20 Hz), 7.15 (dd, 1H, *J*=9.15 Hz and 2.32 Hz), 4.39 (t, 2H, *J*=4.39 Hz), 3.51 (d, br, 2H, *J*=3.91 Hz), 3.12-3.24 (m, 4H), 1.23 ppm (t, 6H, *J*=7.20 Hz). Elemental Analysis for C₂₃H₂₅N₃O₃S · 1.00 mol HCl/0.80 mol H₂O: Calcd: C, 58.23; H, 5.86; N, 8.86; Found: C, 57.92; H, 5.52; N, 8.59.


Example 34



Butyl-2-[3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-ethylamine

A solution of toluene-4-sulfonic acid 2-[3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-ethyl ester (200 mg, 0.383 mmol) and butylamine (1.1-1.5 mmol) in THF (10 mL) was stirred at 70 °C for 2-3 hours in a sealed tube. After cooling to ambient temperature, the reaction mixture was solvent evaporated and partitioned in ethyl acetate and aqueous sodium bicarbonate. The organic phase was dried with anhydrous magnesium sulfate, filtered and concentrated. It was dissolved in methylene chloride and methanol, and ethereal hydrochloride was added. The mixture was concentrated and dried *in vacuo* at 67 °C to give butyl-{2-[3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-ethyl}-amine hydrochloride as a light orange foam (166 mg, 94.3%); Mass Spectrum (-EI, [M-H]⁺) *m/z* 422. ¹HNMR (500 MHz, DMSO-d₆): δ 8.74-8.76 (m, 1H), 8.51-8.53 (m, 1H), 8.27 (d, 1H, *J*=8.29 Hz), 8.03-8.05 (m, 1H), 7.71-7.74 (m, 1H), 7.56-7.65 (m, 3H), 7.34 (d, 1H, *J*=2.20 Hz), 7.16 (dd, 1H, *J*=9.15 Hz and 2.22 Hz), 4.28-4.30 (m, 2H), 3.32-3.35 (m, 2H), 2.93-2.97 (m, 2H), 1.55-1.63 (m, 2H), 1.27-1.36 (m, 2H), 0.85-0.89 ppm (m, 3H).

Example 35




Cyclopentyl-{2-[3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-ethyl}-amine

A solution of toluene-4-sulfonic acid 2-[3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-ethyl ester (200 mg, 0.383 mmol) and cyclopentylamine (1.1-1.5 mmol) in THF (10 mL) was stirred at 70 °C for 2-3 hours in a sealed tube. After cooling to ambient temperature, the reaction mixture was solvent evaporated and partitioned in ethyl acetate and aqueous sodium bicarbonate. The organic phase was dried with anhydrous magnesium sulfate, filtered and concentrated. It was dissolved in methylene chloride and methanol, and ethereal hydrochloride was added. The mixture was concentrated and dried *in vacuo* at 67 °C for 17 hours to give cyclopentyl-{2-[3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-ethyl}-amine hydrochloride as a brown foam (148 mg, 81.8 %); Mass Spectrum (-EI, [M-H]⁺) *m/z* 436. ¹HNMR (500 MHz, DMSO-d₆): δ 14.20 (s, 1H), 8.89-8.93 (s, br, 2H), 8.74-8.76 (m, 1H), 8.51-8.53 (m, 1H), 8.28 (d, 1H, *J*=8.29 Hz), 8.03-8.05 (m, 1H), 7.71-7.75 (m,


1H), 7.56-7.65 (m, 3H), 7.34 (d, 1H, $J=2.32$ Hz), 7.15-7.18 (m, 1H), 4.28-4.30 (m, 2H), 3.51-3.57 (m, 1H), 3.32-3.36 (m, 2H), 1.93-2.00 (m, 2H), 1.44-1.73 ppm (m, 6H).
 Elemental Analysis for $C_{24}H_{25}N_3O_3S \cdot 1.00$ HCl $\cdot 0.70$ mol HCl: Calcd: C, 59.48; H, 5.70; N, 8.67; Found: C, 59.67; H, 5.86; N, 8.32.

5

Example 36**Cyclopropyl-{2-[3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-ethyl}-amine**

A solution of toluene-4-sulfonic acid 2-[3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-ethyl ester (200 mg, 0.383 mmol) and cyclopropylamine (1.1-1.5 mmol) in THF (10 mL) was stirred at 70 °C for 2-3 hours in a sealed tube. After cooling to ambient temperature, the reaction mixture was solvent evaporated and partitioned in ethyl acetate and aqueous sodium bicarbonate. The organic phase was dried with anhydrous magnesium sulfate, filtered and concentrated. It was dissolved in methanol and ethereal hydrochloride was added. The mixture was concentrated and dried *in vacuo* at 70 °C for 16 hours to give cyclopropyl-{2-[3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-ethyl}-amine hydrochloride as a yellow foam (58 mg, 34 %); Mass spectrum ($-EI$, $[M-H]^+$) m/z 406. 1H NMR (400 MHz, DMSO- d_6): δ 14.20 (s, 1H), 9.19 (s, br, 2H), 8.74-8.76 (m, 1H), 8.52 (dd, 1H, $J=7.45$ Hz and 1.22 Hz), 8.27 (d, 1H, $J=8.30$ Hz), 8.02-8.05 (m, 1H), 7.71-7.75 (m, 1H), 7.56-7.65 (m, 3H), 7.34 (d, 1H, $J=2.19$ Hz), 7.16 (dd, 1H, $J=9.15$ Hz and 2.32 Hz), 4.30-4.33 (m, 2H), 3.43 (br, s, 2H), 2.76-2.77 (m, 1H), 0.84-0.88 (m, 2H), 0.71-0.76 ppm (m, 2H). Elemental Analysis for $C_{22}H_{21}N_3O_4S \cdot 1.00$ HCl $\cdot 0.60$ mol HCl: Calcd: C, 58.11; H, 5.14; N, 9.24; Found: C, 57.83; H, 4.81; N, 8.91.


25

Example 37

3-(Naphthalene-1-sulfonyl)-5-(2-pyrrolidin-1-yl-ethoxy)-1H-indazole

A solution of toluene-4-sulfonic acid 2-[3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-ethyl ester (200 mg, 0.383 mmol) and pyrrolidine (1.1-1.5 mmol) in THF (10 mL) was stirred at 70 °C for 2-3 hours in a sealed tube. After cooling to ambient temperature, the reaction mixture was solvent evaporated and partitioned in ethyl acetate and aqueous sodium bicarbonate. The organic phase was dried with anhydrous magnesium sulfate, filtered and concentrated. It was dissolved in methanol and ethereal hydrochloride was added. The mixture was concentrated and dried *in vacuo* at 67 °C for 20 hours to give 3-(naphthalene-1-sulfonyl)-5-(2-pyrrolidin-1-yl-ethoxy)-1H-indazole hydrochloride as a light brown foam (97.8 mg, 55.9 %);

Mass Spectrum (-EI, [M-H]⁺) *m/z* 420. ¹HNMR (500 MHz, DMSO-d₆): δ 14.19 (s, 1H), 10.26-10.29 (br, 1H), 8.74-8.77 (m, 1H), 8.53 (dd, 1H, *J*=7.44 Hz and 1.22 Hz), 8.27 (d, 1H, *J*=8.29 Hz), 8.03-8.05 (m, 1H), 7.71-7.74 (m, 1H), 7.55-7.65 (m, 3H), 7.36 (d, 1H, *J*=2.19 Hz), 7.16-7.19 (m, 1H), 4.36-4.38 (m, 2H), 3.58 (s, br, 4H), 3.06-3.19 (s, br, 2H), 1.81-2.05 ppm (br, m, 4H).


Example 38**3-(Naphthalene-1-sulfonyl)-5-(2-piperidin-1-yl-ethoxy)-1H-indazole**

A solution of the toluene-4-sulfonic acid 2-[3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-ethyl ester (200 mg, 0.383 mmol) and piperidine (1.1-1.5 mmol) in THF (10 mL) was stirred in a sealed tube for about 16 hours at 70 °C. After cooling to ambient temperature, the reaction mixture was solvent evaporated and partitioned in ethyl acetate and aqueous sodium bicarbonate. The organic phase was dried with anhydrous magnesium sulfate, filtered and concentrated. It was dissolved in methanol and chloroform, concentrated and dried *in vacuo* at 70 °C for 16 hours to give 3-(Naphthalene-1-sulfonyl)-5-(2-piperidin-1-yl-ethoxy)-1H-indazole hydrochloride as a pale yellow foam (106 mg, 58.6%).

Mass Spectrum (-EI, [M-H]⁺) *m/z* 434. ¹HNMR (500 MHz, DMSO-d₆): δ 14.18 (s, 1H), 9.77-9.84 (br, 1H), 8.75 (d, 1H, *J*=8.78 Hz), 8.51-8.53 (m, 1H), 8.27 (d, 1H, *J*=8.30 Hz), 8.03-8.05 (m, 1H), 7.70-7.74 (m,

1H), 7.56-7.65 (m, 3H), 7.35 (d, 1H, $J=2.20$ Hz), 7.15 (dd, 1H, $J=9.15$ Hz and 2.32 Hz), 4.40-4.41 (m, 2H), 3.44-3.50 (m, 3H), 2.90-3.06 (m, 2H), 1.61-1.76 (m, 5H), 1.31-1.41 ppm (m, 2H). Elemental Analysis for $C_{24}H_{25}N_3O_4S \cdot 1.00$ HCl $\cdot 0.70$ mol HCl: Calcd: C, 59.48; H, 5.70; N, 8.67; Found: C, 59.22; H, 5.63; N, 8.30.

5

Example 39**5-(2-Morpholin-4-yl-ethoxy)-3-(naphthalene-1-sulfonyl)-1H-indazole**


A solution of toluene-4-sulfonic acid 2-[3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-ethyl ester (200 mg, 0.383 mmol) and morpholine (1.1-1.5 mmol) in THF (10 mL) was stirred at 70 °C for 2-3 hours in a sealed tube. After cooling to ambient temperature, the reaction mixture was solvent evaporated and triturated with ether and ethyl acetate. The resulting solid was then dissolved in ethyl acetate and washed with aqueous sodium bicarbonate. The organic phase was dried with anhydrous magnesium sulfate, filtered, concentrated. The residue was dissolved in methylene chloride and methanol, and ethereal hydrochloride was added. The mixture was concentrated and dried *in vacuo* at 67 °C for 17 hours to give 5-(2-morpholin-4-yl-ethoxy)-3-(naphthalene-1-sulfonyl)-1H-indazole hydrochloride as a light brown semi-solid (0.176 g, 96.7%); Mass Spectrum (+EI, $[M+H]^+$) m/z 438.

$^1\text{H-NMR}$ (500 MHz, DMSO- d_6): δ 14.19 (s, 1H), 10.61-10.66 (br, 1H), 8.74-8.76 (m, 1H), 8.52 (dd, 1H, $J=7.44$ Hz and 1.22 Hz), 8.27 (d, 1H, $J=8.29$ Hz), 8.03-8.05 (m, 1H), 7.71-7.75 (m, 1H), 7.56-7.65 (m, 3H), 7.36 (d, 1H, $J=1.40$ Hz), 7.15-7.17 (dd, 1H, $J=9.15$ Hz and 2.32 Hz), 4.44 (s, br, 2H), 3.93-3.96 (m, 2H), 3.72-3.78 (m, 2H), 3.42-3.69 (m, 4H), 3.12-3.23 ppm (m, 2H). Elemental Analysis for $C_{23}H_{23}N_3O_4S \cdot 1.00$ HCl $\cdot 0.50$ mol HCl: Calcd: C, 57.20; H, 5.22; N, 8.70; Found: C, 57.28; H, 5.24; N, 8.45.

Example 40


Methyl-{2-[1-methyl-3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-ethyl}-amine

30

Step 1

Toluene-4-sulfonic acid 2-[1-methyl-3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-ethyl ester

- 5 Methyliodide (0.06 mL, 0.96 mmol) was added to a chilled mixture of toluene-4-sulfonic acid 2-[3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-ethyl ester (0.401 g, 0.767 mmol) and cesium carbonate (0.29 g, 0.89 mmol) in acetone (10 mL). The reaction mixture was stirred at ambient temperature under nitrogen for 3 hours. It was then partitioned in ethyl acetate and water. The organic phase was washed with
- 10 brine, dried with anhydrous magnesium sulfate, filtered and concentrated. The residue was purified by flash chromatography with 25-40% ethyl acetate in hexane, 100 % chloroform and 1% methanol in chloroform. Drying at 65 °C *in vacuo* for yielded toluene-4-sulfonic acid 2-[1-methyl-3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-ethyl ester as a buff foam (0.224 g, 54.4 %). ¹HNMR (400 MHz, DMSO-d₆):
- 15 δ 8.27 (d, 1H, *J*=8.54 Hz), 8.55 (dd, 1H, *J*=7.42 Hz and 1.16 Hz), 8.30 (d, 1H, *J*=8.35 Hz), 8.05-8.07 (m, 1H), 7.72-7.76 (m, 3H), 7.60-7.70 (m, 3H), 7.31 (d, 2H, *J*=8.01 Hz), 7.18 (d, 1H, *J*=2.20 Hz), 7.00-7.03 (m, 1H), 4.35-4.37 (m, 2H), 4.22-4.24 (m, 2H), 4.07 (s, 3H), 2.28 ppm (s, 3H).


Step 2

Methyl-{2-[1-methyl-3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-ethyl}-amine

- 25 A solution of toluene-4-sulfonic acid 2-[1-methyl-3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-ethyl ester (0.351 g, 0.654 mmol) in 2.0 M methylamine in THF (10 mL, 20 mmol) was heated and stirred in a sealed tube at 78 °C for 1 hour, 40 minutes. Additional methylamine in THF (2.0 mL, 4.0 mmol) of was added, and the

reaction mixture was stirred at 80 °C for 16.5 hours in a sealed tube. After cooling to ambient temperature, the reaction mixture was solvent evaporated. It was partitioned with ethyl acetate and aqueous sodium bicarbonate. It was washed with brine, dried with anhydrous magnesium sulfate, filtered, concentrated and dried at 80 °C *in vacuo* for 20 minutes, resulting in methyl-{2-[1-methyl-3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-ethyl}-amine as a yellow gum (0.198 g, 76.4 %). This was dissolved in chloroform, and ethereal hydrochloride was added. The solid was filtered and dried at 83 °C *in vacuo* for 25 hours. A buff solid (0.156 g) was obtained: MP: 272-4°C(dec.); Mass Spectrum (+EI, [M+H]⁺) *m/z* 396. ¹HNMR (500 MHz, DMSO-d₆): δ 8.89 (s, br, 2H), 8.73-8.75 (m, 1H), 8.50-8.52 (m, 1H), 8.26-8.28 (m, 1H), 8.03-8.05 (m, 1H), 7.71-7.74 (m, 2H), 7.56-7.66 (m, 2H), 7.37 (d, 1H, *J* = 2.07 Hz), 7.19-7.22 (m, 1H), 4.29-4.31 (m, 2H), 4.05 (s, 3H), 3.34 (t, 2H, *J* = 4.39 Hz), 2.62 ppm (s, 3H). Elemental Analysis for C₂₁H₂₁N₃O₃S · 1.00 mol HCl · 0.60 mol H₂O: Calcd: C, 56.97; H, 5.28; N, 9.49; Found: C, 56.91; H, 5.27; N, 9.14.

Example 41




Dimethyl-{2-[1-methyl-3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-ethyl}-amine

A solution of toluene-4-sulfonic acid 2-[1-methyl-3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-ethyl ester (0.39 g, 0.73 mmol) in 2.0 M dimethylamine in THF (10 mL, 20 mmol) was stirred for 16.5 hours at 80 °C in a sealed tube. After cooling to ambient temperature, the reaction mixture was solvent evaporated. The residue was partitioned in ethyl acetate and aqueous sodium bicarbonate. The organic phase was washed with brine, dried with anhydrous magnesium sulfate, filtered, concentrated and dried at 80 °C *in vacuo* for 35 minutes to give dimethyl-{2-[1-methyl-3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-ethyl}-amine a light brown-orange semi-solid (0.257 g, 86.2 %). This was dissolved in chloroform, and ethereal hydrochloride was added. The mixture was concentrated and dried for 25 hours at 83 °C *in vacuo* to give the hydrochloride as a buff-colored foam (0.254 g). Mass


Spectrum (+EI, [M+H]⁺) *m/z* 410. ¹HNMR (500 MHz, DMSO-d₆): δ10.25 (s, 1H), 8.74-8.76 (m, 1H), 8.51 (dd, 1H, *J*=7.44 Hz and 1.22 Hz), 8.26-8.28 (m, 1H), 8.03-8.05 (m, 1H), 7.71-7.74 (m, 2H), 7.56-7.66 (m, 2H), 7.39 (d, 1H, *J*=2.08 Hz), 7.22 (dd, 1H, *J*=9.28 Hz and 2.32 Hz), 4.39-4.42 (m, 2H), 4.05 (s, 3H), 3.50-3.52 (m, 2H), 2.83 ppm (s, 6H).

Example 42




Ethyl-methyl-{2-[1-methyl-3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-ethyl}-amine

A solution of toluene-4-sulfonic acid 2-[1-methyl-3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-ethyl ester (0.398 g, 0.742 mmol) and ethylmethylamine (2.0 mL, 23 mmol) in THF (10 mL) was stirred at 80 °C in a sealed tube for 16.5 hours. After cooling to ambient temperature, the reaction mixture was solvent evaporated. The residue was partitioned in ethyl acetate and aqueous sodium bicarbonate. The organic phase was washed with brine, dried with anhydrous magnesium sulfate, filtered, concentrated and dried at 80 °C *in vacuo* for 20 minutes to give ethyl-methyl-{2-[1-methyl-3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-ethyl}-amine as a light brown gum (0.227 g, 72.3 %). This was dissolved in chloroform, and etheric hydrochloride was added. The mixture was concentrated and dried for 25 hours at 83 °C *in vacuo* to give the hydrochloride as a buff-colored foam (0.254 g); Mass Spectrum (+EI, [M+H]⁺) *m/z* 424. ¹HNMR (500 MHz, DMSO-d₆): δ10.26-10.27 (s, br, 1H), 8.75 (d, 1H, *J*=8.66 Hz), 8.50-8.53 (m, 1H), 8.27 (d, 1H, *J*=8.17 Hz), 8.04 (d, 1H, *J*=7.56 Hz), 7.71-7.75 (m, 2H), 7.56-7.66 (m, 2H), 7.38 (d, 1H, *J*=2.19 Hz), 7.22 (dd, 1H, *J*=9.15 Hz and 2.32 Hz), 4.43 (t, 2H, *J*=4.88 Hz), 4.05 (s, 3H), 3.40-3.60 (m, 3H), 3.10-3.20 (br, 1H), 2.80 (s, 3H), 1.22-1.26 ppm (m, 3H).

Example 43**Diethyl-{2-[1-methyl-3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-ethyl}-amine**


- 5 A solution of toluene-4-sulfonic acid 2-[1-methyl-3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-ethyl ester (0.385 g, 0.717 mmol) and diethylamine (2.0 mL, 19 mmol) in THF (10 mL) was stirred at 80 °C in a sealed tube for 16.5 hours. After cooling to ambient temperature, the reaction mixture was solvent evaporated. The residue was partitioned in chloroform and aqueous sodium bicarbonate. The organic
- 10 phase was washed with brine, dried with anhydrous magnesium sulfate, filtered, concentrated and dried at 80 °C *in vacuo* for 20 minutes to give diethyl-{2-[1-methyl-3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-ethyl}-amine a clear amber gum (0.269 g, 85.7 %). This was dissolved in chloroform, and ethereal hydrochloride was added. The mixture was concentrated and dried for 25 hours at 83 °C *in vacuo* to
- 15 give the hydrochloride as a brown foam (0.253 g); Mass Spectrum (+EI, [M+H]⁺) *m/z* 438. ¹HNMR (500 MHz, DMSO-*d*₆): δ10.11-10.12 (br, 1H), 8.74 (d, 1H, *J*=8.66 Hz), 8.50-8.52 (m, 1H), 8.27 (d, 1H, *J*=8.29 Hz), 8.03-8.05 (m, 1H), 7.71-7.75 (m, 2H), 7.56-7.66 (m, 2H), 7.37 (d, 1H, *J*=2.20 Hz), 7.21 (dd, 1H, *J*=9.27 Hz and 2.32 Hz), 4.41-4.43 (m, 2H), 4.05 (s, 3H), 3.49-3.52 (m, 2H), 3.15-3.25 (m, 4H), 1.22-1.25 ppm
- 20 (m, 6H).

Example 44**Ethyl-{2-[1-methyl-3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-ethyl}-amine**

- 25 A solution of toluene-4-sulfonic acid 2-[1-methyl-3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-ethyl ester (0.413 g, 0.770 mmol) and 2.0 M ethylamine in THF (10 mL, 20 mmol) in THF (10 mL) was stirred at 80 °C in a sealed tube for 16.5

hours. After cooling to ambient temperature, the reaction mixture was solvent evaporated. The residue was partitioned in ethyl acetate and aqueous sodium bicarbonate. The organic phase was washed with brine, dried with anhydrous magnesium sulfate, filtered, concentrated and dried at 80 °C *in vacuo* for 20 minutes to give ethyl-{2-[1-methyl-3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-ethyl}-amine as a dark yellow solid (0.292 g, 92.7 %). This was dissolved in chloroform and methanol, and ethereal hydrochloride was added. The mixture was concentrated and dried for 25 hours at 83 °C *in vacuo* to give the hydrochloride as a beige solid (0.227 g) MP 276-7 °Cdec.; Mass Spectrum (+EI, [M+H]⁺) *m/z* 410. ¹HNMR (500 MHz, DMSO-d₆): δ8.95 (s, br, 2H), 8.75 (d, 1H, J=8.79 Hz), 8.52 (dd, 1H, J=7.45 Hz and 1.22 Hz), 8.27 (d, 1H, J=8.30 Hz), 8.04 (d, 1H, J=7.57 Hz), 7.71-7.75 (m, 2H), 7.56-7.66 (m, 2H), 7.36 (d, 1H, J=2.07 Hz), 7.22 (dd, 1H, J=9.15 Hz and 2.31 Hz), 4.30-4.33 (m, 2H), 4.05 (s, 3H), 3.33-3.34 (m, 2H), 2.99-3.05 (m, 2H), 1.19-1.23 ppm (m, 3H). Elemental Analysis for C₂₂H₂₃N₃O₃S · 1.00 mol HCl · 0.40 mol H₂O: Calcd: C, 58.31; H, 5.52; N, 9.27; Found: C, 58.62; H, 5.54; N, 9.11.

Example 45




Isopropyl-{2-[1-methyl-3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-ethyl}-amine

A solution of toluene-4-sulfonic acid 2-[1-methyl-3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-ethyl ester (0.433 g, 0.807 mmol) and isopropylamine (1.0 mL, 12 mmol) in THF (10 mL) was stirred at 80 °C in a sealed tube for 15 hours. Additional isopropylamine (1.0 mL, 12 mmol) was added, and the reaction mixture was stirred at 80 °C for 21 hours. It was allowed to cool to room temperature, and partitioned in ethyl acetate and aqueous sodium bicarbonate. The organic phase was washed with water and brine. It was dried with anhydrous magnesium sulfate, filtered and concentrated. The residue was purified by flash chromatography using 0.25 % ammonium hydroxide/2.5 % methanol in chloroform. Drying at 63 °C *in vacuo* for 30 minutes yielded isopropyl-{2-[1-methyl-3-(naphthalene-1-sulfonyl)-1H-

indazol-5-yloxy]-ethyl)-amine as a yellow semi-solid 0.198 g, 57.9 %). This was dissolved in chloroform, and ethereal hydrochloride was added. The precipitate was filtered and dried at 85 °C *in vacuo* for 13 hours to give the hydrochloride as an off-white solid (0.180 g): MP: 274-6 °C (dec).; Mass Spectrum (+EI, [M+H]⁺) *m/z* 424.

- 5 ¹HNMR (500 MHz, DMSO-d₆): δ8.83 (s, br, 2H), 8.74 (d, 1H, *J*=8.79 Hz), 8.51 (dd, 1H, *J*=7.44 Hz and 1.22 Hz), 8.27 (d, 1H, *J*=8.30 Hz), 8.03-8.05 (m, 1H), 7.71-7.75 (m, 2H), 7.56-7.66 (m, 2H), 7.36 (d, 1H, *J*=2.20 Hz), 7.20-7.23 (m, 1H), 4.30-4.33 (m, 2H), 4.06 (s, 3H), 3.30-3.38 (m, 3H), 1.25 ppm (d, 6H, *J*=6.46 Hz). Elemental Analysis for C₂₃H₂₅N₃O₃S .1.00 mol HCl. 0.10 mol H₂O: Calcd: C, 59.82; H, 5.72; N, 9.10; Found: C, 59.53; H, 5.66; N, 9.06.

Example 46



1-Methyl-3-(naphthalene-1-sulfonyl)-5-(2-pyrrolidin-1-ylethoxy)-1H-indazole


- 15 A solution of toluene-4-sulfonic acid 2-[1-methyl-3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-ethyl ester (0.421 g, 0.785 mmol) and pyrrolidine (1.0 mL, 12 mmol) in THF (10 mL) was stirred at 80 °C in a sealed tube for 15 hours. After cooling to ambient temperature, the reaction mixture was solvent evaporated and partitioned in ethyl acetate and aqueous sodium bicarbonate. The organic phase
- 20 was washed with brine, dried with anhydrous magnesium sulfate, filtered and concentrated. The residue was purified by flash chromatography with 0.25 % ammonium hydroxide/2.5 % methanol in chloroform. Drying at 68 °C *in vacuo* for 1 hour resulted in 1-methyl-3-(naphthalene-1-sulfonyl)-5-(2-pyrrolidin-1-ylethoxy)-1H-indazole as a light brown semi-solid (0.159 g, 46.5%). This was dissolved in
- 25 chloroform, and ethereal hydrochloride was added. The precipitate was filtered, dissolved in methanol and concentrated. Drying at 85 °C *in vacuo* for 13 hours yielded the hydrochloride as a light orange semi-solid (0.156 g); Mass Spectrum (+EI, [M+H]⁺) *m/z* 436. ¹HNMR (500 MHz, DMSO-d₆): δ10.60-10.61 (s, br, 1H), 8.75 (d, 1H, *J*=8.66 Hz), 8.51-8.53 (m, 1H), 8.27 (d, 1H, *J*=8.29 Hz), 8.04 (d, 1H, *J*=7.93 Hz), 7.71-7.74 (m, 2H), 7.56-7.67 (m, 2H), 7.37 (d, 1H, *J*=2.20 Hz), 7.22-7.25 (m,
- 30

1H), 4.39-4.41 (m, 2H), 4.05 (s, 3H), 3.57-3.60 (m, 4H), 3.06-3.14 (m, 2H), 1.95-2.01 (m, 2H), 1.86-1.92 ppm (m, 2H). Elemental Analysis for $C_{24}H_{25}N_3O_3S \cdot 1.00 \text{ mol HCl} \cdot 1.00 \text{ mol H}_2\text{O}$: Calcd: C, 58.82; H, 5.76; N, 8.58; Found: C, 58.77; H, 6.00; N, 8.47.

5

Example 47

{2-[1-Benzyl-3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-ethyl}-ethyl-methyl-amine

Step 1


10

Toluene-4-sulfonic acid 2-[1-benzyl-3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-ethyl ester

Benzyl bromide (0.46 mL, 3.9 mmol) was added to a stirring suspension of toluene-4-sulfonic acid 2-[3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-ethyl ester (1.85 g, 3.54 mmol) and cesium carbonate (1.28 g, 3.93 mmol) in acetone (60 mL). The reaction mixture was stirred under nitrogen at ambient temperature for 1.5 hours. It was then poured into excess water and extracted with chloroform. The organic phase was washed with brine, dried with anhydrous magnesium sulfate, filtered and concentrated. The residue was purified by flash chromatography with 40-50 % ethyl acetate in hexane. Drying at 80 °C *in vacuo* for 30 minutes yielded toluene-4-sulfonic acid 2-[1-benzyl-3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-ethyl ester as a light buff foam (1.39 g, 64.1 %); Mass Spectrum (+EI, $[M+H]^+$) m/z 613. ^1H NMR (500 MHz, DMSO- d_6): δ 8.74-8.77 (m, 1H), 8.55-8.57 (m, 1H), 8.25-8.27 (m, 1H), 8.01-8.05 (m, 1H), 7.64-7.76 (m, 4H), 7.58-7.64 (m, 2H), 7.25 (d, 2H, $J=7.93 \text{ Hz}$), 7.10-7.19 (m, 5H), 7.07 (d, 1H, $J=2.19 \text{ Hz}$), 6.94-6.97 (m, 1H), 5.69 (s, 2H), 4.30-4.32 (m, 2H), 4.14-4.16 (m, 2H), 2.18 ppm (s, 3H). Elemental Analysis for $C_{33}H_{28}N_2O_6S_2 \cdot 0.60 \text{ mol H}_2\text{O}$: Calcd: C, 63.57; H, 4.72; N, 4.49; Found: C, 63.17; H, 4.63; N, 4.30.

25


Step 2



{2-[1-Benzyl-3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-ethyl}-ethyl-methyl-amine

A solution of toluene-4-sulfonic acid 2-[1-benzyl-3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-ethyl ester (0.363 g, 0.592 mmol) and ethylmethylamine (2.0 mL, 23 mmol) in THF (8 mL) was stirred at 70 °C for 2.5 hours in a sealed tube. After cooling to ambient temperature, the residue was partitioned in ethyl acetate and aqueous sodium bicarbonate. The organic phase was washed with brine, dried with anhydrous magnesium sulfate, filtered and concentrated. Drying at 80 °C *in vacuo* for 30 minutes gave {2-[1-benzyl-3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-ethyl}-ethyl-methyl-amine as a yellow foam/gum (0.241 g, 81.4 %). This was dissolved in chloroform, and ethereal hydrochloride was added. The mixture was concentrated and dried *in vacuo* for 13 hours to yield the hydrochloride as a buff foam (0.242 g). Mass Spectrum (+EI, [M+H]⁺) *m/z* 500. ¹HNMR (500 MHz, DMSO-*d*₆): δ 10.04-10.10 (s, br, 1H), 8.74-8.77 (m, 1H), 8.54-8.56 (m, 1H), 8.27 (d, 1H, *J*=8.29 Hz), 8.02-8.05 (m, 1H), 7.79 (d, 1H, *J*=9.28 Hz), 7.70-7.74 (m, 1H), 7.55-7.67 (m, 2H), 7.28 (d, 1H, *J*=2.19 Hz), 7.04-7.20 (m, 6H), 5.70 (s, 2H), 4.35 (s, 2H), 3.33-3.41 (br, 2H), 3.01-3.19 (br, 2H), 2.73 (s, 3H), 1.17-1.21 ppm (m, 3H). Elemental Analysis for C₂₉H₂₉N₃O₃S·1.00 mol HCl·1.10 mol H₂O: Calcd: C, 62.66; H, 5.84; N, 7.56; Found: C, 62.73; H, 6.19; N, 7.17.


Example 48



{2-[1-Benzyl-3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-ethyl}-methyl-amine

A solution of toluene-4-sulfonic acid 2-[1-benzyl-3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-ethyl ester (0.476 g, 0.777 mmol) in 2.0 M methylamine in THF (8.0 mL, 16 mmol) was stirred at 70 °C in a sealed tube for 3 hours. After cooling to ambient temperature and concentrating, the residue was partitioned in ethyl acetate and aqueous sodium bicarbonate. The organic phase was washed with water and brine. It was dried with anhydrous magnesium sulfate, filtered and concentrated. Drying at 80 °C *in vacuo* for 20 minutes gave {2-[1-benzyl-3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-ethyl}-methyl-amine as a clear, dark yellow gum (0.324 g, 88.5 %). This was dissolved in chloroform, and ethereal hydrochloride was added. The mixture was concentrated and dried at 81 °C *in vacuo* for 13 hours. The hydrochloride as a light orange foam (0.330 g) resulted; Mass Spectrum (+EI, [M+H]⁺) *m/z* 472. ¹HNMR (500 MHz, DMSO-*d*₆): δ 8.86 (s, 2H), 8.74-8.77 (m, 1H), 8.54-8.56 (m, 1H), 8.27 (d, 1H, *J*=8.30 Hz), 8.02-8.05 (m, 1H), 7.79 (d, 1H, *J*=9.15 Hz), 7.70-7.74 (m, 1H), 7.58-7.65 (m, 2H), 7.27 (d, 1H, *J*=3.30 Hz), 7.09-7.18 (m, 6H), 5.70 (s, 2H), 4.24-4.27 (m, 2H), 2.59 ppm (s, 3H). Elemental Analysis for C₂₇H₂₅N₃O₃S₂·1.00 HCl·0.55 mol H₂O: Calcd: C, 62.61; H, 5.27; N, 8.11; Found: C, 62.23; H, 5.45; N, 7.72.

Example 49



{2-[1-Benzyl-3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-ethyl}-isopropyl-amine

A solution of toluene-4-sulfonic acid 2-[1-benzyl-3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-ethyl ester (0.444 g, 0.725 mmol) and isopropylamine (2.0 mL, 23 mmol) in THF (8 mL) was stirred at 70 °C in a sealed tube for 3 hours. More isopropylamine (2.0 mL, 23 mmol) was then added, and the reaction mixture was stirred at 80 °C in a sealed tube for 16.5 hours. After cooling somewhat, the reaction


mixture was solvent evaporated and partitioned in ethyl acetate and aqueous sodium bicarbonate. The organic phase was then washed with water and brine. It was dried with anhydrous magnesium sulfate, filtered and concentrated. Drying at 80 °C *in vacuo* for 20 minutes gave {2-[1-benzyl-3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-ethyl}-isopropyl-amine as a yellow semi-solid (0.290 g, 80.1% mmol). This was dissolved in chloroform, and ethereal hydrochloride was added. The mixture was concentrated and dried for 13 hours *in vacuo* at 81 °C. The result was the hydrochloride as a buff foam (0.273 g); Mass Spectrum (+EI, [M+H]⁺) *m/z* 500.

¹HNMR (500 MHz, DMSO-d₆): δ8.80 (s, 2H), 8.74-8.78 (m, 1H), 8.55 (dd, 1H, *J*=7.44 Hz and 1.22 Hz), 8.27 (d, 1H, *J*=8.29 Hz), 8.01-8.05 (m, 1H), 7.79 (d, 1H, *J*=9.15 Hz), 7.70-7.74 (m, 1H), 7.58-7.65 (m, 2H), 7.27 (d, 1H, *J*=2.20 Hz), 7.09-7.19 (m, 6H), 5.70 (s, 2H), 4.26-4.28 (m, 2H), 3.31 (s, 3H), 1.23 ppm (d, 6H, *J*=6.47 Hz).

Elemental Analysis for C₂₉H₂₉N₃O₃S· 1.00 HCl·0.7 mol H₂O: Calcd: C, 63.48; H, 5.77; N, 7.66;

Found: C, 63.10; H, 5.67; N, 7.37.

Example 50




{2-[1-Benzyl-3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-ethyl}-diethyl-amine

A solution of toluene-4-sulfonic acid 2-[1-benzyl-3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-ethyl ester (0.449 g, 0.733 mmol) and diethylamine (2.0 mL, 19 mmol) in THF (7.5 mL) was stirred at 70 °C in a sealed tube for 3 hours. More diethylamine (2.0 mL, 19 mmol) was then added, and the reaction mixture was stirred at 80 °C in a sealed tube for 16.5 hours. After cooling somewhat, the reaction mixture was solvent evaporated and partitioned in ethyl acetate and aqueous sodium bicarbonate. The organic phase was then washed with water and brine. It was dried with anhydrous magnesium sulfate, filtered and concentrated. Drying at 82°C *in vacuo* for 35 minutes gave {2-[1-benzyl-3-(naphthalene-1-sulfonyl)-1H-indazol-5-

yloxy]-ethyl)-diethyl-amine as a light brown-yellow semi-solid (0.365 g, 97.1% mmol). This was dissolved in chloroform, and ethereal hydrochloride was added. The mixture was concentrated and dried for 13 hours *in vacuo* at 81 °C. The result was the hydrochloride as a light orange semi-solid (0.332 g); Mass Spectrum (+EI, $[M+H]^+$) m/z 514. ¹HNMR (500 MHz, DMSO- d_6): δ 10.15-10.18 (s, 1H), 8.74-8.76 (m, 1H), 8.56 (dd, 1H, $J=7.44$ Hz and 1.22 Hz), 8.26 (d, 1H, $J=8.29$ Hz), 8.02-8.05 (m, 1H), 7.78 (d, 1H, $J=9.15$ Hz), 7.70-7.74 (m, 1H), 7.58-7.65 (m, 2H), 7.26 (s, 1H), 7.08-7.19 (m, 6H), 5.70 (s, 2H), 4.25 (s, br, 2H), 3.39-3.55 (br, s, 2H), 3.15 (br, s, 3H), 1.13-1.19 ppm (s, br, 6H). Elemental Analysis for $C_{30}H_{31}N_3O_3S \cdot 1.00 HCl \cdot 0.75$ mol H_2O : Calcd: C, 63.93; H, 5.99; N, 7.46; Found: C, 63.59; H, 5.94; N, 7.23.

Example 51




{2-[1-Benzyl-3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-ethyl}-dimethyl-amine

A solution of toluene-4-sulfonic acid 2-[1-benzyl-3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-ethyl ester (0.432 g, 0.705 mmol) in 2.0 M dimethylamine in THF (8.0 mL, 16 mmol) was stirred at 70 °C in a sealed tube for 3 hours. After cooling somewhat, the reaction mixture was solvent evaporated and partitioned in ethyl acetate and aqueous sodium bicarbonate. The organic phase was then washed with water and brine. It was dried with anhydrous magnesium sulfate, filtered and concentrated. Drying at 80 °C *in vacuo* for 20 minutes gave {2-[1-benzyl-3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-ethyl}-dimethyl-amine as a yellow solid (0.288 g, 84.2% mmol). This was dissolved in chloroform, and ethereal hydrochloride was added. The mixture was concentrated and dried for 13 hours *in vacuo* at 81 °C. The result was the hydrochloride as a buff foam (0.294 g); Mass Spectrum (+EI, $[M+H]^+$) m/z 486. ¹HNMR (500 MHz, DMSO- d_6): δ 10.11 (s, 1H), 8.74-8.77 (m, 1H), 8.54-8.56 (m, 1H), 8.27 (d, 1H, $J=8.30$ Hz), 8.03-8.05 (m, 1H), 7.79 (d, 1H, $J=9.15$ Hz), 7.70-7.74 (m, 1H), 7.58-7.65 (m, 2H), 7.30 (d, 1H, $J=2.20$

Hz), 7.08-7.19 (m, 6H), 5.70 (s, 2H), 4.34-4.37 (m, 2H), 3.48 (t, 2H, $J=4.76$ Hz), 2.80 ppm (s, 6H). Elemental Analysis for $C_{28}H_{27}N_3O_3S \cdot 1.00$ HCl $\cdot 1.55$ mol H_2O : Calcd: C, 61.15; H, 5.70; N, 7.64; Found: C, 61.33; H, 5.59; N, 7.24.


5

Example 52**1-Benzyl-3-(naphthalene-1-sulfonyl)-5-(2-pyrrolidin-1-ylethoxy)-1H-indazole**

A solution of toluene-4-sulfonic acid 2-[1-benzyl-3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-ethyl ester (0.379 g, 0.619 mmol) and pyrrolidine (1.0 mL, 12 mmol) in THF (80 mL) was stirred at 70 °C in a sealed tube for 3 hours. The reaction mixture was solvent evaporated and partitioned in ethyl acetate and aqueous sodium bicarbonate. The organic phase was then washed with water and brine. It was dried with anhydrous magnesium sulfate, filtered and concentrated. Drying at 80 °C *in vacuo* for 25 minutes gave 1-benzyl-3-(naphthalene-1-sulfonyl)-5-(2-pyrrolidin-1-ylethoxy)-1H-indazole as a yellow-orange solid (0.259 g, 82.0% mmol). This was dissolved in chloroform, and ethereal hydrochloride was added. The mixture was concentrated and dried for 13 hours *in vacuo* at 81 °C. The result was the hydrochloride as a buff foam (0.267g); Mass Spectrum (+EI, $[M+H]^+$) m/z 512. 1H NMR (500 MHz, $DMSO-d_6$): δ 10.42 (s, 1H), 8.74-8.77 (m, 1H), 8.56 (dd, 1H, $J=7.44$ Hz and 1.22 Hz), 8.27 (d, 1H, $J=8.30$ Hz), 8.02-8.05 (m, 1H), 7.79 (d, 1H, $J=9.27$ Hz), 7.70-7.74 (m, 1H), 7.55-7.65 (m, 2H), 7.28 (d, 1H, $J=2.07$ Hz), 7.08-7.20 (m, 6H), 5.70 (s, 2H), 4.33-4.36 (m, 2H), 3.54 (br, s, 4H), 3.07 (s, br, 2H), 1.79-2.01 ppm (m, 4H). Elemental Analysis for $C_{30}H_{29}N_3O_3S \cdot 1.00$ HCl $\cdot 1.25$ mol H_2O : Calcd: C, 63.15; H, 5.74; N, 7.36; Found: C, 62.82; H, 5.74; N, 6.99.

25

Example 53




{2-[1-Benzyl-3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-ethyl}-ethyl-amine

A solution of toluene-4-sulfonic acid 2-[1-benzyl-3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-ethyl ester (0.430 g, 0.702 mmol) and 2.0 M ethylamine in THF (4.8 mL, 9.6 mmol) was stirred at 70 °C in a sealed tube for 3 hours. After cooling to ambient temperature, the reaction mixture was solvent evaporated and partitioned in ethyl acetate and aqueous sodium bicarbonate. The organic phase was washed with water and brine. It was dried with anhydrous magnesium sulfate, filtered and concentrated. The residue was purified by flash chromatography with 5.0-7.5 % methanol in chloroform and by HPLC with 5-50 % (chloroform/methanol (8:2)/TEA) in heptane/TEA. The result was {2-[1-benzyl-3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-ethyl}-ethyl-amine as a yellow semi-solid (0.0528g, 15.5 %). This was dissolved in chloroform, and ethereal hydrochloride was added. The mixture was concentrated and dried for 14 hours at 82 °C *in vacuo* to give the hydrochloride as a light orange semi-solid (0.0546 g); Mass Spectrum (+EI, [M+H]⁺) *m/z* 486. ¹HNMR (500 MHz, DMSO-*d*₆): δ 8.74-8.77 (m, 1H), 8.65-8.72 (br, s, 2H), 8.55 (dd, 1H, *J*=7.32 Hz and 1.22 Hz), 8.26-8.28 (m, 1H), 8.03-8.05 (m, 1H), 7.79 (d, 1H, *J*=9.15 Hz), 7.70-7.74 (m, 1H), 7.58-7.65 (m, 2H), 7.27 (d, 1H, *J*=2.07 Hz), 7.09-7.18 (m, 6H), 5.70 (s, 2H), 4.23-4.26 (m, 2H), 3.29-3.32 (m, 2H), 2.97-3.02 (m, 2H), 1.16-1.19 ppm (m, 3H). Elemental Analysis for C₂₈H₂₇N₃O₃S·1.00 HCl·0.80 mol H₂O: Calcd: C, 62.69; H, 5.56; N, 7.83; Found: C, 62.38; H, 5.38; N, 7.58.

Example 54

{2-[1-(3-Chloro-benzyl)-3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-ethyl}-dimethyl-amine


Step 1



Toluene-4-sulfonic acid 2-[1-(3-chloro-benzyl)-3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-ethyl ester

- 5 3-Chlorobenzylbromide (0.90 mL, 6.8 mmol) was added to a stirring suspension of toluene-4-sulfonic acid 2-[3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-ethyl ester (3.24 g, 6.20 mmol) and cesium carbonate (2.21 g, 6.78 mmol) in acetone (80 mL). The mixture was stirred at ambient temperature under nitrogen for 2 hours. It was then poured into excess water and extracted with chloroform. The organic phase was washed with brine, dried with anhydrous magnesium sulfate, filtered and concentrated. The residue was purified by flash chromatography using 30-35 % ethyl acetate in hexane. Drying at 65 °C *in vacuo* for 30 minutes gave toluene-4-sulfonic acid 2-[1-(3-chloro-benzyl)-3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-ethyl ester as a dark yellow foam (2.55 g, 63.6 %); Mass Spectrum (+EI, [M+H]⁺) *m/z* 647. ¹HNMR (500 MHz, DMSO-*d*₆): δ 8.73-8.75 (m, 1H), 8.55-8.57 (m, 1H), 8.27 (d, 1H, *J*=8.30 Hz), 8.01-8.04 (m, 1H), 7.67-7.75 (m, 4H), 7.56-7.63 (m, 2H), 7.17-7.27 (m, 5H), 7.05-7.07 (m, 2H), 6.98 (dd, 1H, *J*=9.28 Hz and 2.32 Hz), 5.71 (s, 2H), 4.31-4.33 (m, 2H), 4.15-4.17 (m, 2H), 2.17 ppm (s, 3H). Elemental Analysis for C₃₃H₂₇ClN₂O₆S₂: Calcd: C, 61.25; H, 4.21; N, 4.33; Found: C, 60.93; H, 4.22; N, 4.21.
- 10
- 15
- 20


Step 2



{2-[1-(3-Chloro-benzyl)-3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-ethyl}-dimethyl-amine

A solution of toluene-4-sulfonic acid 2-[1-(3-chloro-benzyl)-3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-ethyl ester (0.344 g, 0.532 mmol) in 2.0 M dimethylamine in THF (8.0 mL, 16.0 mmol) was stirred at 70 °C in a sealed tube for 2 hours. After cooling to ambient temperature, the reaction mixture was solvent evaporated. It was partitioned with ethyl acetate and aqueous sodium bicarbonate. The organic phase was washed with brine, dried with anhydrous magnesium sulfate, filtered and concentrated. Drying at 80 °C *in vacuo* for 20 minutes gave {2-[1-(3-chloro-benzyl)-3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-ethyl}-dimethyl-amine as a dark yellow gum (0.261 g, 94.6 %). This was dissolved in chloroform, and ethereal hydrochloride was added. The mixture was concentrated and dried *in vacuo* at 82 °C for 14 hours. The result was the hydrochloride as a light orange foam (0.252 g); Mass Spectrum (+EI, [M+H]⁺) *m/z* 520. ¹HNMR (500 MHz, DMSO-*d*₆): 89.92-9.94 (br, s, 1H), 8.73-8.75 (m, 1H), 8.56 (dd, 1H, *J*=7.32 Hz and 1.10 Hz), 8.27 (d, 1H, *J*=8.29 Hz), 8.02-8.05 (m, 1H), 7.82 (d, 1H, *J*=9.15 Hz), 7.70-7.74 (m, 1H), 7.58-7.64 (m, 2H), 7.31 (d, 1H, *J*=2.32 Hz), 7.15-7.27 (m, 3H), 7.04 (d, 1H, *J*=7.68 Hz), 5.73 (s, 2H), 4.34-4.37 (m, 2H), 3.47-3.49 (m, 2H), 2.81 ppm (s, 6H).

Example 55




{2-[1-(3-Chloro-benzyl)-3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-ethyl}-methyl-amine

A solution of toluene-4-sulfonic acid 2-[1-(3-chloro-benzyl)-3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-ethyl ester (0.350 g, 0.541 mmol) and ethylmethylamine (1.0 mL, 12 mmol) in THF (8 mL) was stirred for 2 hours in a sealed tube at 70 °C. Additional ethylmethylamine (1.0 mL, 12 mmol) was added, and the reaction mixture was stirred at 80 °C in sealed tube for 2 hours. After cooling to ambient temperature, the reaction mixture was solvent evaporated and partitioned in ethyl acetate and

aqueous sodium bicarbonate. The organic phase was then washed with brine, dried with anhydrous magnesium sulfate, filtered and concentrated. It was dried at 80 °C for 20 minutes *in vacuo* to yield {2-[1-(3-chloro-benzyl)-3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-ethyl}-ethyl-methyl-amine as a yellow semi-solid (0.268 g, 92.7 %). This was dissolved in chloroform, and ethereal hydrochloride was added. The mixture was then concentrated and dried *in vacuo* at 82 °C for 14 hours. The result was the hydrochloride as a light orange foam (0.266 g); Mass Spectrum (+EI, [M+H]⁺) *m/z* 534. ¹HNMR (500 MHz, DMSO-d₆): δ 9.90-9.93 (s, br, 1H), 8.73-8.75 (m, 1H), 8.54-8.57 (m, 1H), 8.27 (d, 1H, *J*=8.29 Hz), 8.02-8.05 (m, 1H), 7.82 (d, 1H, *J*=9.27 Hz), 7.70-7.74 (m, 1H), 7.56-7.63 (m, 2H), 7.30 (d, 1H, *J*=2.20 Hz), 7.15-7.26 (m, 4H), 7.04 (d, 1H, *J*=7.57 Hz), 5.73 (s, 2H), 4.36-4.38 (m, 2H), 3.40-3.55 (m, 3H), 3.09-3.14 (br, 1H), 2.78 (s, 3H), 1.19-1.23 ppm (m, 3H). Elemental Analysis for C₂₉H₂₈ClN₃O₃S · 1.00 HCl · 0.75 mol H₂O: Calcd: C, 59.64; H, 5.26; N, 7.19; Found: C, 59.24; H, 5.17; N, 6.89.


Example 56



{2-[1-(3-Chloro-benzyl)-3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-ethyl}-diethyl-amine

A solution of toluene-4-sulfonic acid 2-[1-(3-chloro-benzyl)-3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-ethyl ester (0.338 g, 0.522 mmol) and diethylamine (1.0 mL, 9.7 mmol) in THF (8 mL) was stirred at 70 °C in a sealed tube for 2 hours. Additional diethylamine (1.0 mL, 9.7 mmol) added, and the reaction mixture was stirred at 80 °C in a sealed tube for 2 hours. A third portion of diethylamine (1.0 mL, 9.7 mmol) was added to the reaction mixture, and it was stirred at 80 °C in a sealed tube for 30 hours. The reaction mixture was then solvent evaporated and partitioned in ethyl acetate and brine. It was dried with anhydrous magnesium sulfate, filtered and concentrated. Drying at 80 °C *in vacuo* for 20 minutes yielded {2-[1-(3-chloro-benzyl)-3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-ethyl}-diethyl-amine as a light

brown semi-solid (0.278 g, 97.2%). It was dissolved in chloroform, and ethereal hydrochloride was added. The mixture was concentrated and dried *in vacuo* at 82 °C for 14 hours. The result was the hydrochloride as a brown foam (0.275 g); Mass Spectrum (+EI, [M+H]⁺) *m/z* 548. ¹HNMR (500 MHz, DMSO-d₆): δ 9.86-9.88 (s, 1H), 8.72-8.75 (m, 1H), 8.56 (dd, 1H, *J*=7.44 Hz and 1.22 Hz), 8.27 (d, 1H, *J*=8.30 Hz), 8.02-8.05 (m, 1H), 7.82 (d, 1H, *J*=9.15 Hz), 7.70-7.74 (m, 1H), 7.57-7.63 (m, 2H), 7.29 (d, 1H, *J*=2.20 Hz), 7.24-7.26 (m, 1H), 7.15-7.21 (m, 3H), 7.03 (d, 1H, *J*=7.68 Hz), 5.73 (s, 2H), 4.35-4.38 (m, 2H), 3.49 (d, br, 2H, *J*=4.52 Hz), 3.14-3.25 (m, 4H), 1.19-1.23 ppm (m, 6H). Elemental Analysis for C₃₀H₃₀ClN₃O₃S· 1.00 HCl·0.55 mol H₂O: Calcd: C, 60.61; H, 5.44; N, 7.07; Found: C, 60.21; H, 5.44; N, 6.73.


Example 57

{2-[1-(3-Chloro-benzyl)-3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-ethyl}-isopropyl-amine

A solution of toluene-4-sulfonic acid 2-[1-(3-chloro-benzyl)-3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-ethyl ester (0.339 g, 0.524 mmol) and isopropylamine (1.0 mL, 12 mmol) in THF (8 mL) was stirred at 70 °C in a sealed tube for 2 hours. Isopropylamine (1.0 mL, 12 mmol) was added, and the reaction mixture was stirred at 80 °C in a sealed tube for 2 hours. Isopropylamine (1.0 mL, 12 mmol) was added to the reaction mixture, and it was stirred at 80 °C in a sealed tube for 30 hours. After cooling to ambient temperature, the reaction mixture was solvent evaporated and partitioned in ethyl acetate and aqueous sodium bicarbonate. It was washed with brine, dried with anhydrous magnesium sulfate, filtered, concentrated and dried *in vacuo* for 20 minutes at 80 °C to give {2-[1-(3-chloro-benzyl)-3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-ethyl}-isopropyl-amine as a yellow solid (0.280 g, 100 %). It was dissolved in chloroform, and ethereal hydrochloride was added. The resulting solid was filtered and dried at 82 °C *in vacuo* for 14 hours. The result was the hydrochloride as a pale yellow solid (0.236 g); MP: 206-8 °C; Mass Spectrum


(+EI, $[M+H]^+$) m/z 534. ^1H NMR (500 MHz, DMSO- d_6): δ 8.65-8.76 (m, 3H), 8.54-8.56 (m, 1H), 8.27 (d, 1H, $J=8.17$ Hz), 8.02-8.05 (m, 1H), 7.82 (d, 1H, $J=9.27$ Hz), 7.72 (dd, 1H, $J=8.07$ Hz and 7.57 Hz), 7.57-7.63 (m, 2H), 7.24-7.28 (m, 2H), 7.16-7.21 (m, 3H), 7.05 (d, 1H, $J=7.81$ Hz), 5.73 (s, 2H), 4.27 (t, 2H, $J=5.00$ Hz), 3.31-3.37 (m, 3H), 1.23 ppm (d, 6H, $J=6.69$ Hz). Elemental Analysis for $\text{C}_{26}\text{H}_{28}\text{ClN}_3\text{O}_3\text{S} \cdot 1.00 \text{ HCl}$: Calcd: C, 61.05; H, 5.12; N, 7.36; Found: C, 60.68; H, 5.26; N, 7.14.

Example 58



10 1-(3-Chloro-benzyl)-3-(naphthalene-1-sulfonyl)-5-(2-pyrrolidin-1-yl-ethoxy)-1H-indazole

A solution of toluene-4-sulfonic acid 2-[1-(3-chloro-benzyl)-3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-ethyl ester (0.359 g, 0.555 mmol) and pyrrolidine (1.0 mL, 12 mmol) in THF (8 mL) was stirred at 70 °C in a sealed tube for 2 hours. After cooling to ambient temperature, the reaction mixture was solvent evaporated. It was then partitioned in ethyl acetate and aqueous sodium bicarbonate. The organic phase was washed with brine, dried with anhydrous magnesium sulfate, filtered, concentrated and dried *in vacuo* at 80 °C for 20 minutes to give 1-(3-chloro-benzyl)-3-(naphthalene-1-sulfonyl)-5-(2-pyrrolidin-1-yl-ethoxy)-1H-indazole as a yellow semi-solid (0.266 g, 87.8 %). This was dissolved in chloroform, and ethereal hydrochloride was added. The mixture was concentrated and dried at 82 °C *in vacuo* for 14 hours to yield the hydrochloride as a light orange foam (0.205 g); Mass Spectrum (+EI, $[M+H]^+$) m/z 546. ^1H NMR (500 MHz, DMSO- d_6): δ 10.18-10.19 (s, 1H), 8.73-8.75 (m, 1H), 8.55-8.57 (m, 1H), 8.27 (d, 1H, $J=8.30$ Hz), 8.02-8.04 (m, 1H), 7.82 (d, 1H, $J=9.15$ Hz), 7.70-7.74 (m, 1H), 7.57-7.64 (m, 2H), 7.15-7.29 (m, 5H), 7.04 (d, 1H, $J=7.69$ Hz), 5.73 (s, 2H), 4.33-4.35 (m, 2H), 3.55 (br, s, 4H), 3.07-3.15 (s, br, 2H), 1.81-2.02 ppm (br, m, 4H). Elemental Analysis for $\text{C}_{30}\text{H}_{28}\text{ClN}_3\text{O}_3\text{S} \cdot 1.00 \text{ HCl} \cdot 0.70 \text{ mole H}_2\text{O}$: Calcd: C, 60.54; H, 5.15; N, 7.06; Found: C, 60.16; H, 5.11; N, 6.74.


Example 59

{2-[1-(3-Chloro-benzyl)-3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-ethyl}-methyl-amine

- 5 A solution of toluene-4-sulfonic acid 2-[1-(3-chloro-benzyl)-3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-ethyl ester (0.367 g, 0.567 mmol) in 2.0 M methylamine in THF (8.0 mL, 16 mmol) was stirred at 70 °C in a sealed tube for 2 hours. After cooling to ambient temperature, the reaction mixture was solvent evaporated. The residue was partitioned in ethyl acetate and aqueous sodium bicarbonate. The
- 10 organic phase was then washed with brine, dried with anhydrous magnesium sulfate, filtered and concentrated. The residue was purified by HPLC with 25-60% (chloroform/methanol 8:2/TEA) in heptane/TEA. Concentration and drying at 60 °C *in vacuo* yielded {2-[1-(3-chloro-benzyl)-3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-ethyl}-methyl-amine as a light yellow foam (0.190 g, 66.2%). This was dissolved in
- 15 chloroform, and ethereal hydrochloride was added. The mixture was then concentrated and dried *in vacuo* for 29 hours at 80 °C. The hydrochloride as an off-white foam resulted (0.206 g); Mass Spectrum (+EI, [M+H]⁺) *m/z* 506. ¹HNMR (500 MHz, DMSO-*d*₆): δ 8.71-8.76 (m, 3H), 8.54-8.56 (m, 1H), 8.27 (d, 1H, *J*=8.30 Hz), 8.02-8.05 (m, 1H), 7.82 (d, 1H, *J*=9.27 Hz), 7.70-7.74 (m, 1H), 7.57-7.63 (m, 2H),
- 20 7.24-7.28 (m, 2H), 7.16-7.20 (m, 3H), 7.05 (d, 1H, *J*=7.69 Hz), 5.73 (s, 2H), 4.24-4.26 (m, 2H), 3.30-3.32 (m, 2H), 2.60 ppm (s, 4H). Elemental Analysis for C₂₇H₂₄ClN₃O₃S·1.00 HCl·1.50 mole H₂O: Calcd: C, 56.94; H, 4.96; N, 7.38; Found: C, 56.64; H, 4.88; N, 7.06.

25

Example 60




{2-[1-(3-Chloro-benzyl)-3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-ethyl}-ethyl-amine

- A solution of toluene-4-sulfonic acid 2-[1-(3-chloro-benzyl)-3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-ethyl ester (0.415 g, 0.641 mmol) in 2.0 M ethylamine in THF (8.0 mL, 16.0 mmol) was stirred at 70 °C for 3 hours and then at 80 °C for 19 hours. The reaction mixture was allowed to cool to ambient temperature and solvent evaporated. It was partitioned in ethyl acetate and aqueous sodium bicarbonate. The organic phase was washed with brine, dried with magnesium sulfate, filtered and concentrated. Drying *in vacuo* at 80 °C for 20 minutes resulted in {2-[1-(3-chloro-benzyl)-3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-ethyl}-ethyl-amine as an orange semi-solid (0.270 g, 81.1 %). This was dissolved in chloroform, and ethereal hydrochloride was added. The mixture was concentrated and dried *in vacuo* at 83 °C for 16 hours. The hydrochloride as a pale orange foam resulted (0.276 g); Mass Spectrum (+EI, [M+H]⁺) *m/z* 520. ¹HNMR (500 MHz, DMSO-*d*₆): δ 8.71-8.77 (m, 3H), 8.55 (dd, 1H, *J*=7.44 Hz and 1.22 Hz), 8.27 (d, 1H, *J*=8.29 Hz), 8.02-8.05 (m, 1H), 7.82 (d, 1H, *J*=9.15 Hz), 7.70-7.74 (m, 1H), 7.57-7.64 (m, 2H), 7.24-7.28 (m, 2H), 7.16-7.21 (m, 3H), 7.05 (d, 1H, *J*=7.56 Hz), 5.73 (s, 2H), 4.24-4.27 (m, 2H), 3.30-3.32 (m, 2H), 2.97-3.03 (m, 2H), 1.16-1.20 ppm (m, 3H). Elemental Analysis for C₂₈H₂₆ClN₃O₃S · 1.00 HCl · 0.60 mole H₂O: Calcd: C, 59.28; H, 5.01; N, 7.41; Found: C, 58.95; H, 5.06; N, 7.14.

Example 61

3-(1-naphthylsulfonyl)-5-(4-piperidin-1-ylbutoxy)-1H-indazole

**Step 1****1-(4-Chloro-butoxy)-4-nitro-benzene**

A mixture of *para*-nitrophenol (0.83 g, 6 mmol), 1-bromo-4-chlorobutane (1.23 g, 7.2 mmol), and K_2CO_3 (1.24 g, 9 mmol) was stirred together in DMF at 80 °C for 1 hour. Reaction mixture was diluted with H_2O , extracted with EtOAc, washed with water (2x), brine (1x), dried over Na_2SO_4 , and concentrated under vacuum. The crude product was purified by HPLC using as eluent 30%EtOAc/hexane to afford the title compound as an off-white solid (1.28 g, 5.6 mmol).

Step 2**1-[5-(4-Chloro-butoxy)-2-nitro-phenylmethanesulfonyl]-naphthalene**

A mixture of 1-(4-chloro-butoxy)-4-nitrobenzene (1.28 g, 5.6 mmol) and 1-chloromethanesulfonyl-naphthalene (1.6 g, 6.72 mmol) was stirred in THF (50 ml) at -78 °C, in a round bottom flask under nitrogen. A solution of 1M potassium *t*-butoxide was added dropwise (16.8 ml, 16.8 mmol) over a half an hour period. Temperature was allowed to rise to -40 °C, and the reaction mixture was stirred at this temperature for 5 hours. The reaction mixture was poured into cold 2N HCl, extracted with EtOAc, dried over Na_2SO_4 , and concentrated under vacuum. Compound was recrystallized from CH_2Cl_2 /hexane to afford the title compound as an off-white solid (1.94 g, 4.5 mmol).

Step 3**4-(4-Chloro-butoxy)-2-(naphthalene-1-sulfonylmethyl)-phenyl amine**

A mixture of 1-[5-(4-chloro-butoxy)-2-nitro-phenyl-methanesulfonyl]-naphthalene (1.94 g, 4.5 mmol) and 10% Pd/C in THF (20 mL), methanol (20 mL),

and formic acid (5 mL) was hydrogenated in a Parr hydrogenation bottle (250 mL) at 40 lb/in² for 20 hours. The mixture was filtered through Celite, and the filtrate was diluted with EtOAc, washed with water, dried over Na₂SO₄, and concentrated under vacuum. The crude product was purified by flash chromatography using as eluent 5% EtOAc/CH₂Cl₂ to afford the title compound as an off-white solid (1.54 g, 3.8 mmols).

Step 4


5-(4-Chloro-butoxy)-3-(naphthalene-1-sulfonyl)-1-H-indazole

A mixture of 4-(4-chloro-butoxy)-2-(naphthalene-1-sulfonylmethyl)-phenyl amine (1.54 g, 3.8 mmols) in THF (7 mL), and 4M HCl (15 mL) was stirred in a round bottom flask, under nitrogen, at 3 °C. A solution of sodium nitrite (0.34 g, 4.0 mmols) in H₂O (1 mL) was added dropwise. The reaction mixture was poured into a cold solution of saturated sodium bicarbonate (100 mL) and extracted with EtOAc. Compound was dried over Na₂SO₄, and concentrated under vacuum to afford the title compound as an off white solid (1.55 g, 3.75 mmols).


Step 5

A mixture of 5-(4-chloro-butoxy)-3-(naphthalene-1-sulfonyl)-1-H-indazole (0.065 g, 0.12 mmols) and piperidine (0.48 mmols) in DMF (1 mL) was stirred under nitrogen at 100 °C overnight. Mixture was cooled to room temperature, diluted with water, extracted with EtOAc, washed with water (2x), brine (1x), dried over Na₂SO₄, and concentrated under vacuum. Compound was purified by flash chromatography using as eluent 5% CH₃OH/EtOAc. The purified compound was dissolved in methanol, 1M HCl in ether (0.1 mL, 0.1 mmols) was added, and the compound was dried to afford the title compound 3-(1-naphthylsulfonyl)-5-(4-piperidin-1-ylbutoxy)-1H-indazole as the HCl salt, MS: (ES⁺) 463 [M+H]⁺

Examples 62-67



Using essentially the same procedure described in Example 61, Step 5 and employing an appropriate amine, the compounds shown on Table I were obtained and identified by HPLC and mass spectral analyses.


TABLE I

5

Ex. No.	R5	R6	[M+H] ⁺
62	CH ₃	H	465
63	C ₂ H ₅	H	424
64	CH ₃	CH ₃	424
65	CH ₃	<i>n</i> -propyl	452
66	C ₂ H ₅	C ₂ H ₅	452
67	-CH ₂ CH ₂ CH ₂ CH ₂ -		450

Example 68

10 (4-{[3-(1-naphthylsulfonyl)-1H-indazol-5-yl] oxy} butyl) amine

**Step 1**


15 A mixture of 5-(4-chloro-butoxy)-3-(naphthalene-1-sulfonyl)-1-H-indazole (0.065 g, 0.12 mmoles) and sodium azide (0.24 mmoles) in DMSO (1 mL) was stirred under nitrogen at 90°C for 5 hours. Mixture was cooled to room temperature, diluted with water, extracted with EtOAc, washed with water (2x), brine (1x), dried over Na₂SO₄, and concentrated under vacuum. Compound was purified by flash chromatography using as eluent 5%CH₃OH/EtOAc.

Step 2.

The azide prepared in step 1 was subjected to hydrogenation over 10 %Pd/C in THF (2 mL), and methanol (8 mL) in a Parr hydrogenation bottle (250 mL) at 52lb/in² for 2 hours. The mixture was filtered through Celite, and the filtrate was concentrated under vacuum. The crude product was recrystallized from CH₂Cl₂/hexane, 1M HCl in ether (0.9 ml, 0.9 mmoles) was added, then evaporated to afford the title compound - (4-[[3- (1-naphthylsulfonyl)-1H-indazol-5-yl] oxy} butyl) amine as an off-white HCl salt (0.04 g, 0.1 mmoles), MS: (ES⁺) 394[M-H]⁺

Example 69

(2-[[1-(3-Chlorobenzyl)-3-(1-naphthylsulfonyl)-1-H-indazol-7-yl] oxy} ethyl) amine



Step 1

1-(2-Chloro-ethoxy)-2-nitro-benzene

A mixture of *ortho*-nitrophenol (5.0 g, 35.8 mmoles) and 2-chloroethanol (9.5 ml, 143 mmoles) in THF (50 mL), in a round bottom flask, under nitrogen, was stirred at room temperature. Triphenylphosphine (14 g, 53.7 mmoles) was added, followed by diethylazodicarboxylate dropwise (8.5 mL, 53.7 mmoles). The mixture was stirred overnight at room temperature. The reaction mixture was diluted with water, extracted with EtOAc, washed with water (1x), brine (1x), dried (Na₂SO₄), and concentrated under vacuum. The crude product was purified by flash chromatography using as eluent 40 %EtOAc/hexane to give the title compound as an off-white solid (5.6 g, 28 mmoles).

Step 2

1-Chloromethane-sulfonyl-naphthalene

A mixture of naphthalene-1-sulfonyl chloride (10.0 g, 44 mmoles), sodium sulfite (11.12 g, 88 mmoles), and sodium bicarbonate (7.4 g, 88 mmoles) in water (50 mL) was heated to 100 °C for one hour. The crude sodium sulfinic acid solution was allowed to cool for 30 minutes, and then treated with bromochloromethane (43 mL, 5 661 mmoles) and tetra-N-butylammonium bromide (1.4 g, 4.4 mmoles). The resultant mixture was heated to 75 °C overnight. All solvents were removed under vacuum. Compound was recrystallized from CH₂Cl₂/hexane to give the title compound as an off white solid (10.62 g, 44 mmoles).

10 Step 3

1-[3-(2-Chloro-ethoxy)-2-nitro-phenyl-methanesulfonyl]-naphthalene

A mixture of 1-(2-chloro-ethoxy)-2-nitrobenzene (1.2 g, 6 mmoles) and 1-chloromethane-sulfonyl-naphthalene (2.16 g, 9 mmoles) was stirred in THF (50 ml) at -78 °C, in a round bottom flask under nitrogen. A solution of 1M potassium t-butoxide was added drop wise (18 mL, 18 mmoles) over a half an hour period. 15 Temperature was allowed to rise to -40 °C, and the reaction mixture was stirred at this temperature for 5 hours. The reaction mixture was poured into cold 2N HCl, extracted with EtOAc, dried over Na₂SO₄, and concentrated under vacuum. Compound was recrystallized from CH₂Cl₂/hexane to afford the title compound as an 20 off-white solid (1.4 g, 3 mmoles).

Step 4

2-(2-Chloro-ethoxy)-6-(naphthalene-1-sulfonylmethyl)-phenyl amine

A mixture of 1-[3-(2-chloro-ethoxy)-2-nitro-phenyl-methanesulfonyl]-naphthalene (1.24 g, 3.5 mmoles) and 10% Pd/C in THF (20 mL), methanol (5 mL), 25 and formic acid (5 mL) was hydrogenated in a Parr hydrogenation bottle (250 mL) at 40 lb/in² for 20 hours. The mixture was filtered through Celite, and the filtrate was diluted with EtOAc, washed with water, dried over Na₂SO₄, and concentrated under vacuum. The crude product was purified by flash chromatography using as eluent 30 5% EtOAc/CH₂Cl₂ to afford the title compound as an off-white solid (1.0 g, 3.1 mmoles).

Step 5

7-(2-Chloro-ethoxy)-3-(naphthalene-1-sulfonyl)-1-H-indazole

A mixture of 2-(2-chloro-ethoxy)-6-(naphthalene-1-sulfonylmethyl)-phenyl amine (0.97 g, 3 mmoles) in THF (7 mL), and 4M HCl (15 mL) was stirred in a round bottom flask, under nitrogen, at 3 °C. A solution of sodium nitrite (0.21 g, 3.15 mmoles) in H₂O (1 mL) was added dropwise. The reaction mixture was poured into a cold solution of saturated sodium bicarbonate (100 mL) and extracted with EtOAc. Compound was dried over Na₂SO₄, and concentrated under vacuum to afford the title compound as an off white solid (0.9 g, 2.7 mmoles).

Step 6**1-(3-Chloro-benzyl)-7-(2-chloro-ethoxy)-3-(naphthalene-1-sulfonyl)-1-H-indazole**

A mixture of 7-(2-chloro-ethoxy)-3-(naphthalene-1-sulfonyl)-1-H-indazole (0.7 g, 1.8 mmoles), 3-chloro-benzyl bromide (0.28 mL, 2.17 mmoles), and cesium carbonate (0.7 g, 2.17 mmoles) in DMF (5 mL) was stirred together in a round bottom flask at room temperature for 30 minutes. Reaction mixture was diluted with H₂O, extracted with EtOAc, washed with water (2x), brine (1x), dried over Na₂SO₄, and concentrated under vacuum. The crude product was purified by HPLC using as eluent 30% EtOAc/hexane to afford the title compound as an off-white solid (0.55 g, 1.1 mmoles).

Step 7**7-(2-Azido-ethoxy)-3-(naphthalene-1-sulfonyl)-1-H-indazole**

A mixture of 1-(3-chloro-benzyl)-7-(2-chloro-ethoxy)-3-(naphthalene-1-sulfonyl)-1-H-indazole (0.25 g, 0.49 mmoles) and sodium azide (0.04 g, 0.58 mmoles) in DMSO (3 mL) was stirred together in a round bottom flask under nitrogen at 90 °C for 3 hours. Reaction mixture was cooled to room temperature, diluted with water, extracted with EtOAc, washed with water (2x), brine (1x), dried over Na₂SO₄, and concentrated under vacuum to afford the title compound as an off white solid (0.23 g, 0.44 mmoles).

Step 8


2-[[1-(3-Chlorobenzyl)-3-(1-naphthylsulfonyl)-1-H-indazol-7-yl] oxy} ethyl) amine

A mixture of 7-(2-azido-ethoxy)-3-(naphthalene-1-sulfonyl)-1-*H*-indazole (0.23 g, 0.44 mmoles) and 10% Pd/C in THF (2 mL), and methanol (5 mL) was hydrogenated in a Parr hydrogenation bottle (250 mL) at 52 lb/in² for 2 hours. The mixture was filtered through Celite, and the filtrate was concentrated under vacuum.

- 5 The crude product was purified by flash chromatography using as eluent 5% CH₃OH/CH₂Cl₂ to afford an off-white solid (0.2 g, 0.4 mmoles), MS: (ES⁺) 493 [M+H]⁺

Example 70

(2-[[1-(3-Chlorobenzyl)-3-(1-naphthylsulfonyl)-1-*H*-indazol-7-yl] oxy} ethyl) methylamine




10


A mixture of 1-(3-chloro-benzyl)-7-(2-chloro-ethoxy)-3-(naphthalene-1-sulfonyl)-1-*H*-indazole (0.065 g, 0.12 mmoles) and methylamine (0.48 mmoles) in DMF (1 mL) was stirred under nitrogen at 100 °C overnight. Mixture was cooled to room temperature, diluted with water, extracted with EtOAc, washed with water (2x),
 15 brine (1x), dried over Na₂SO₄, and concentrated under vacuum. Compound was purified by flash chromatography using as eluent 5% CH₃OH/EtOAc. The purified compound was dissolved in methanol, 1M HCl in ether (0.1 mL, 0.1 mmoles) was added, and compound was dried to afford the title compound as the HCl salt, MS: (ES⁺) 507 [M+H]⁺.

20

Examples 71-75




Using essentially the same procedure described in Example 70 and employing an appropriate amine, the compounds shown in Table II were obtained and identified by HPLC and mass spectral analyses.

TABLE II

Ex. No.	R5	R6	[M+H] ⁺
71	C ₂ H ₅	H	522
72	C ₂ H ₅	C ₂ H ₅	549
73	<i>n</i> -butyl	H	549
74	-CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ -		561
75	-CH ₂ CH ₂ CH ₂ CH ₂ -		547

Example 76


(2-[[3-(1-naphthylsulfonyl)-1-H-indazol-7-yl]oxy]ethyl)amine



A mixture of 0.075 g of 1-(3-chlorobenzyl)-3-(1-naphthylsulfonyl)-7-(2-(2-aminoethoxy)-1-H-indazole, DMSO (1 mL) and *t*-BuOH (0.2 mL) was stirred at room temperature in a round bottom flask under oxygen atmosphere. A solution of potassium *t*-butoxide (0.98 mL, 0.98 mmoles) was added dropwise and the reaction mixture stirred for 1 hr. Reaction mixture was quenched with saturated ammonium chloride, extracted with EtOAc, dried over Na₂SO₄, and concentrated under vacuum. Crude compound was purified by flash chromatography using 10% CH₃OH/CH₂Cl₂. The purified compounds were dissolved in methanol, 1M HCl in ether (0.1 mL, 0.1

mmoles) was added, and compound was dried to afford the title compound as the HCl salt, MS: (ES⁺) 368 [M+H]⁺.

Examples 77-81




5


Using essentially the same procedure described in Example 76 and employing the appropriate indazol-7-yloxyethanamine substrate, the compounds shown in Table III were obtained and identified by HPLC and mass spectral analyses.

10

TABLE III




Ex. No.	R5	R6	[M+H] ⁺
77	CH ₃	H	382
78	CH ₃	CH ₃	396
79	C ₂ H ₅	H	396
80	-CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ -		436
81	-CH ₂ CH ₂ CH ₂ CH ₂ -		422

Examples 82-93**Step 1****5 1-Benzene-sulfonyl-methyl-3- (2-chloro-ethoxy)-2-nitro-benzene**


A mixture of 1-(2-chloro-ethoxy)-2-nitrobenzene (1.2 g, 6 mmoles) and chloromethyl phenyl sulfone (2.16 g, 9 mmoles) was stirred in THF (50 mL) at -78 °C, in a round bottom flask under nitrogen. A solution of 1M potassium *t*-butoxide was added dropwise (18 mL, 18 mmoles) over a half an hour period. Temperature was allowed to rise to -40 °C, and the reaction mixture was stirred at this temperature for 5 hours. The reaction mixture was poured into cold 2N HCl, extracted with EtOAc, dried over Na₂SO₄, and concentrated under vacuum. Compound was recrystallized from CH₂Cl₂/hexane to afford the title compound as an off-white solid (1.5 g, 4.2 mmoles).

15 Steps 2-6

Using essentially the same procedures described in Example 69 steps 3-4, and Examples 70 and 76, the compounds shown in Table IV were obtained and identified by HPLC and mass spectral analyses.

TABLE IV

Ex. No.	R1	R5	R6	[M+H] ⁺
82	3-Cl-benzyl	CH ₃	H	457
83	3-Cl-benzyl	C ₂ H ₅	H	471
84	3-Cl-benzyl	C ₂ H ₅	C ₂ H ₅	499
85	3-Cl-benzyl	<i>n</i> -butyl	H	499
86	3-Cl-benzyl	-CH ₂ CH ₂ CH ₂ CH ₂ -		497
87	3-Cl-benzyl	-CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ -		511
88	H	CH ₃	H	332
89	H	C ₂ H ₅	H	346
90	H	C ₂ H ₅	C ₂ H ₅	374
91	H	<i>n</i> -butyl	H	374
92	H	-CH ₂ CH ₂ CH ₂ CH ₂ -		372
93	H	-CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ -		386

Examples 94-105

5

Step 1**7-(2-Chloro-ethoxy)-1-methyl-3-(naphthalene-1-sulfonyl)-1H-indazole:**

A mixture of 7-(2-chloro-ethoxy)-3-(naphthalene-1-sulfonyl)-1-*H*-indazole (0.7 g, 1.8 mmoles), methyl iodide (0.28 mL, 2.17 mmoles), and potassium carbonate (0.29 g, 2.17 mmoles) in DMF (10 mL) was stirred together in a round bottom flask at room temperature for 2 hours. Reaction mixture was diluted with H₂O, extracted with


10

EtOAc, washed with water (2x), brine (1x), dried over Na₂SO₄, and concentrated under vacuum. The crude product was purified by HPLC using as eluent 30% EtOAc/hexane to afford the title compound as an off-white solid, 0.55 g, 1.375 mmoles.

5 Step 2

Using essentially the same procedure described in Example 70 and employing the appropriate 3-arylsulfonyl-1-methylindazole substrate and desired amine and conversion into HCl salts, the compounds shown in Table V are obtained and identified by HPLC and mass spectral analyses.


TABLE V



Ex. No.	R2	R5	R6	[M+H] ⁺
94	1-naphthyl	H	H	419
95	1-naphthyl	C ₂ H ₅	H	410
96	1-naphthyl	CH ₃	CH ₃	410
97	1-naphthyl	C ₂ H ₅	C ₂ H ₅	438
98	1-naphthyl	<i>n</i> -butyl	H	438
99	1-naphthyl	-CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ -		450
100	1-naphthyl	-CH ₂ CH ₂ CH ₂ CH ₂ -		436
101	phenyl	CH ₃	H	346
102	phenyl	C ₂ H ₅	H	360
103	phenyl	C ₂ H ₅	C ₂ H ₅	388
104	phenyl	-CH ₂ CH ₂ CH ₂ CH ₂ -		386
105	phenyl	-CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ -		400

Example 106

(2-[[1-(3-Chlorobenzyl)-5-fluoro-3- (1-naphthylsulfonyl)-1H-indazol-7-yl] oxy} ethyl) amine

5 **Step 1****2-(2-Chloro-ethoxy)-4-fluoro-1-nitro-benzene**

In a round bottom flask under nitrogen, 2-chloro-ethanol (8.3 mL, 120 mmol) in THF (40 mL) was cooled to 0 °C. LDA (60 mL, 120 mmol) was added dropwise, while maintaining the temperature constant at 0 °C. The mixture was stirred at this temperature for 15 minutes, followed by the addition of 2,4-difluoronitrobenzene (11 mL, 100 mmol). The mixture was stirred at room temperature overnight. Reaction mixture was diluted with water, extracted with EtOAc, washed with brine (1x), dried over Na₂SO₄, and concentrated under vacuum, to afford the title compound as an off-white solid (20.0 g, 91 mmol).

15 **Step 2****1-[3-(2-Chloro-ethoxy)-5-fluoro-2-nitro-phenylmethanesulfonyl]-naphthalene**

A mixture of 2-(2-chloro-ethoxy)-4-fluoro-1-nitrobenzene (1.3 g, 6 mmol) and 1-chloromethanesulfonyl-naphthalene (2.16 g, 9 mmol) was stirred in THF (50 mL) at -78 °C, in a round bottom flask under nitrogen. A solution of 1M potassium *t*-butoxide was added drop wise (18 mL, 18 mmol) over a half an hour period. Temperature was allowed to rise to -40 °C, and the reaction mixture was stirred at this temperature for 5 hours. The reaction mixture was poured into cold 2N HCl, extracted with EtOAc, dried over Na₂SO₄, and concentrated under vacuum.

Compound was recrystallized from CH₂Cl₂/hexane to afford the title compound as an off-white solid (2.25 g, 5.3 mmoles).

Step 3

2-(2-Chloro-ethoxy)-4-fluoro-6-(naphthalene-1-sulfonyl-methyl)-phenylamine

5 A mixture of 1-[3-(2-chloro-ethoxy)-5-fluoro-2-nitro-phenylmethanesulfonyl]-naphthalene (1.0 g, 2.36 mmoles) in ethanol (25 mL) was stirred under nitrogen in a round bottom flask at 60 °C. 10% Pd/C was added, and the temperature was increased to 80 °C. Hydrazine hydrate (2.0 mL) was added dropwise and the mixture was stirred at reflux for 3 hours. Reaction mixture was filtered off through Celite, and
10 the solution was washed with H₂O (3x), dried over Na₂SO₄, and concentrated under vacuum to afford the title compound as an off white solid (0.91 g, 2.31 mmoles).

Step 4

7-(2-Chloro-ethoxy)-5-fluoro-3-(naphthalene-1-sulfonyl)-1H-indazole

15 A mixture of 2-(2-chloro-ethoxy)-4-fluoro-6-(naphthalene-1-sulfonyl-methyl)-phenylamine (0.91 g, 2.31 mmoles) in THF (7 mL), and 4M HCl (15 mL) was stirred in a round bottom flask, under nitrogen, at 3 °C. A solution of sodium nitrite (0.16 g, 2.4 mmoles) in H₂O (1 mL) was added dropwise. The reaction mixture was poured into a cold solution of saturated sodium bicarbonate (100 mL) and extracted with EtOAc. Compound was dried over Na₂SO₄, and concentrated under vacuum to
20 afford the title compound as an off white solid (0.9 g, 2.2 mmoles).

Step 5

1-(3-Chloro-benzyl)-7-(2-chloro-ethoxy)-5-fluoro-3-(naphthalene-1-sulfonyl)-1H-indazole

25 A mixture of 7-(2-chloro-ethoxy)-5-fluoro-3-(naphthalene-1-sulfonyl)-1H-indazole (0.9 g, 2.2 mmoles), 3-chlorobenzyl bromide (0.35 mL, 2.7 mmoles), and cesium carbonate (0.87 g, 2.7 mmoles) in DMF (5 mL) was stirred together in a round bottom flask at room temperature for 30 minutes. Reaction mixture was diluted with H₂O, extracted with EtOAc, washed with water (2x), brine (1x), dried over Na₂SO₄, and concentrated under vacuum. The crude product was purified by HPLC
30 using as eluent 30% EtOAc/hexane to afford the title compound as an off-white solid (0.85 g, 2 mmoles).

Step 6

7-(2-Azido-ethoxy)-1-(3-chloro-benzyl)-5-fluoro-3-(naphthalene-1-sulfonyl)-1H-indazole

A mixture of 1-(3-chloro-benzyl)-7-(2-chloro-ethoxy)-5-fluoro-3-(naphthalene-1-sulfonyl)-1H-indazole (0.1 g, 0.19 mmoles) and sodium azide (0.014 g, 0.22 mmoles) in DMSO (3 mL) was stirred together in a round bottom flask under nitrogen at 90 °C for 3 hours. Reaction mixture was cooled to room temperature, diluted with water, extracted with EtOAc, washed with water (2x), brine (1x), dried over Na₂SO₄, and concentrated under vacuum to afford the title compound as an off white solid (0.09 g, 0.17 mmoles).


Step 7

(2-[[1-(3-Chlorobenzyl)-5-fluoro-3-(1-naphthylsulfonyl)-1H-indazol-7-yl] oxy} ethyl) amine

A mixture of 7-(2-azido-ethoxy)-1-(3-chloro-benzyl)-5-fluoro-3-(naphthalene-1-sulfonyl)-1H-indazole (0.09 g, 0.17 mmoles) and 10% Pd/C in THF (5 mL), and methanol (15 mL) was hydrogenated in a Parr hydrogenation bottle (250 mL) at 52 lb/in² for 2 hours. The mixture was filtered through Celite, and the filtrate was concentrated under vacuum. The crude product was purified by flash chromatography using as eluent 5 % CH₃OH/CH₂Cl₂ to afford the title compound as an off-white solid (0.08 g, 0.14 mmoles), MS: (ES⁺) 511 [M+H]⁺.

Example 107

(2-[[1-(3-chlorobenzyl)-5-fluoro-3-(1-naphthylsulfonyl)-1H-indazol-7-yl] oxy} ethyl) methylamine Hydrochloride




A mixture of 1-(3-chloro-benzyl)-7-(2-chloro-ethoxy)-5-fluoro-3-(naphthalene-1-sulfonyl)-1H-indazole (0.075 g, 0.14 mmoles) and methylamine (0.56 mmoles) in DMSO (1 mL) was stirred under nitrogen at 100 °C for 4 hours. Mixture was cooled to room temperature, diluted with water, extracted with EtOAc, washed with water

(2x), brine (1x), dried over Na_2SO_4 , and concentrated under vacuum. Compound was purified by flash chromatography using as eluent 5% $\text{CH}_3\text{OH}/\text{EtOAc}$. The purified compound was dissolved in methanol, 1M HCl in ether (0.1 mL, 0.1 mmoles) was added, and compound was dried to afford the title compound as the HCl salt.


5 MS: (ES^+) 525 $[\text{M}+\text{H}]^+$

Examples 108-118




10 Using essentially the same procedures described in Examples 107 and 76 and employing the desired amine, the compounds shown in Table VI were obtained and identified by HPLC and mass spectral analyses.

TABLE VI




Ex. No.	R1	R5	R6	$[\text{M}+\text{H}]^+$
108	3-Cl-benzyl	C_2H_5	H	539
109	3-Cl-benzyl	CH_3	CH_3	539
110	3-Cl-benzyl	C_2H_5	C_2H_5	567
111	3-Cl-benzyl	$-\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2-$		579
112	3-Cl-benzyl	$-\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2-$		565
113	H	CH_3	H	400
114	H	C_2H_5	H	414
115	H	CH_3	CH_3	414

TABLE VI, Cont.

Ex. No.	R1	R5	R6	[M+H] ⁺
116	H	C ₂ H ₅	C ₂ H ₅	442
117	H	-CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ -		454
118	H	-CH ₂ CH ₂ CH ₂ CH ₂ -		440

5

Examples 119-127**Step 1) 2-(2-Chloroethoxy)-4-methoxy-1-nitrobenzene**

To a mixture of 2-chloroethanol (0.7 mL, 10.5 mmoles) in THF (20 mL), cooled to 0 °C was added 2M LDA (5.25 mL, 10.5 mmoles) dropwise. When the addition of LDA was complete, the mixture was stirred for an additional 15 minutes, and then 2-fluoro-4-methoxy-1-nitrobenzene (1.5 g, 8.8 mmoles) was added. The mixture was allowed to warm to room temperature, and stirred at this temperature overnight. Reaction mixture was diluted with water, extracted with EtOAc, washed with brine (1x), dried over Na₂SO₄, and concentrated under vacuum. The crude

compound was recrystallized from CH_2Cl_2 /hexane to give the title compound (1.5 g, 6.5 mmoles).

Step 2) 1-[3-(2-Chloro-ethoxy)-5-methoxy-2-nitro-phenylmethanesulfonyl]-naphthalene

5 A mixture of 2-(2-chloro-ethoxy)-4-methoxy-1-nitrobenzene (1.4 g, 6 mmoles) and 1-chloromethane-sulfonyl-naphthalene (1.4 g, 6 mmoles) was stirred in THF (50 mL) at -78°C , in a round bottom flask under nitrogen. A solution of 1M potassium *t*-butoxide was added dropwise (18 mL, 18 mmoles) over a half an hour period. Temperature was allowed to rise to -40°C , and the reaction mixture was stirred at
10 this temperature for 4 hours. The reaction mixture was poured into cold 2N HCl, extracted with EtOAc, dried over Na_2SO_4 , and concentrated under vacuum. Compound was recrystallized from CH_2Cl_2 /hexane to afford the title compound as an off-white solid (2 g, 4.6 mmoles).

15 **Step 3) 2-(2-Chloro-ethoxy)-4-methoxy-6-(naphthalene-1-sulfonylmethyl)-phenylamine**

 A mixture of 1-[3-(2-chloro-ethoxy)-5-methoxy-2-nitro-phenylmethanesulfonyl]-naphthalene (1.0 g, 2.5 mmoles) in ethanol (25 mL) was stirred under nitrogen in a round bottom flask at 60°C . 10% Pd/C was added, and the temperature was increased to 80°C . Hydrazine hydrate (2.0 mL) was added
20 dropwise and the mixture was stirred at reflux for 3 hours. Reaction mixture was filtered off through Celite, and the solution was washed with H_2O (3x), dried over Na_2SO_4 , and concentrated under vacuum to afford the title compound as an off white solid (1.0 g, 2.46 mmoles).

Step 4) 7-(2-Chloro-ethoxy)-5-methoxy-3-(naphthalene-1-sulfonyl)-1H-indazole

25 A mixture of 2-(2-chloro-ethoxy)-4-methoxy-6-(naphthalene-1-sulfonylmethyl)-phenylamine (1.0 g, 2.46 mmoles) in THF (7 mL), and 4M HCl (15 mL) was stirred in a round bottom flask, under nitrogen, at 3°C . A solution of sodium nitrite (0.17 g, 2.6 mmoles) in H_2O (1 mL) was added dropwise. The reaction mixture was poured into a cold solution of saturated sodium bicarbonate (100 mL) and extracted with EtOAc.
30 Compound was dried over Na_2SO_4 , and concentrated under vacuum to afford the title compound as an off white solid (1.0 g, 2.39 mmoles).


Step 5) 1-(3-Chloro-benzyl)-7-(2-chloro-ethoxy-5-methoxy-3-(naphthalene-1-sulfonyl)-1H-indazole

A mixture of 7-(2-chloro-ethoxy)-5-methoxy-3-(naphthalene-1-sulfonyl)-1H-indazole (1.0 g, 2.39 mmoles), 3-chlorobenzyl bromide (0.35 mL, 2.7 mmoles), and cesium carbonate (0.87 g, 2.7 mmoles) in DMF (5 mL) was stirred together in a round bottom flask at room temperature for 10 minutes. Reaction mixture was diluted with H₂O, extracted with EtOAc, washed with water (2x), brine (1x), dried over Na₂SO₄, and concentrated under vacuum. The crude product was purified by HPLC using as eluent 30% EtOAc/hexane to afford the title compound as an off-white solid (1.1 g, 2 mmoles).


Steps 6 and 7) Amination and Debenzylation

Using essentially the same procedures described in Examples 107 and 76 and employing the 7-(2-chloroethoxy)-5-methoxy-3-naphthylsulfonylindazole substrate and the desired amine, the compounds shown on Table VII were obtained and identified by HPLC and mass spectral analyses.

TABLE VII



Ex. No.	R1	R5	R6	[M+H] ⁺
119	3-Cl-benzyl	CH ₃	H	537
120	3-Cl-benzyl	C ₂ H ₅	H	551
121	3-Cl-benzyl	CH ₃	CH ₃	551
122	3-Cl-benzyl	-CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ -		591
123	3-Cl-benzyl	-CH ₂ CH ₂ CH ₂ CH ₂ -		577
124	H	CH ₃	H	412
125	H	CH ₃	CH ₃	426
126	H	-CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ -		466
127	H	-CH ₂ CH ₂ CH ₂ CH ₂ -		452

Examples 128-138**Step 1) 1-(3-Chloropropoxy)-2-nitro-benzene**

A mixture of *ortho*-nitrophenol (0.83 g, 6 mmoles), 1-bromo-3-chloropropane (1.1 g, 7.2 mmoles), and K_2CO_3 (1.24 g, 9 mmoles) was stirred together in DMF at 80 °C for 1 hour. Reaction mixture was diluted with H_2O , extracted with EtOAc, washed with water (2x), brine (1x), dried over Na_2SO_4 , and concentrated under vacuum. The crude product was purified by HPLC using as eluent 30% EtOAc/hexane to afford the title compound as an off-white solid (1.2 g, 5.6 mmoles).

Step 2) 1-[3-(3-Chloropropoxy)-2-nitro-phenyl-methanesulfonyl]-naphthalene

A mixture of 1-(3-chloropropoxy)-2-nitrobenzene (1.2 g, 6 mmoles) and 1-chloromethane-sulfonyl-naphthalene (2.16 g, 9 mmoles) was stirred in THF (50 mL) at -78 °C, in a round bottom flask under nitrogen. A solution of 1M potassium *t*-butoxide was added dropwise (18 mL, 18 mmoles) over a half hour period.

Temperature was allowed to rise to -40 °C, and the reaction mixture was stirred at this temperature for 5 hours. The reaction mixture was poured into cold 2N HCl, extracted with EtOAc, dried over Na_2SO_4 , and concentrated under vacuum. Compound was recrystallized from CH_2Cl_2 /hexane to afford the title compound as an off-white solid (1.9 g, 4.5 mmoles).

Step 3) 2-(3-Chloropropoxy)-6-(naphthalene-1-sulfonylmethyl)-phenyl amine

A mixture of 1-[3-(3-chloropropoxy)-2-nitro-phenyl-methanesulfonyl]-naphthalene (1.9 g, 4.5 mmoles) and 10% Pd/C in THF (20 mL), methanol (20 mL), and formic acid (5 mL) was hydrogenated in a Parr hydrogenation bottle (250 mL) at 40 lb/in² for 20 hours. The mixture was filtered through Celite, and the filtrate was

diluted with EtOAc, washed with water, dried over Na₂SO₄, and concentrated under vacuum. The crude product was purified by flash chromatography using as eluent 5% EtOAc/CH₂Cl₂ to afford the title compound as an off-white solid (1.66 g, 4.25 mmol).

5 **Step 4) 7-(3-Chloro-propoxy)-3-(naphthalene-1-sulfonyl)-1-H-indazole**

A mixture of 2-(3-chloro-propoxy)-6-(naphthalene-1-sulfonylmethyl)-phenyl amine (1.66 g, 4.25 mmol) in THF (7 mL), and 4M HCl (15 mL) was stirred in a round bottom flask, under nitrogen, at 3 °C. A solution of sodium nitrite (0.3 g, 4.4 mmol) in H₂O (1 mL) was added dropwise. The reaction mixture was poured into a
10 cold solution of saturated sodium bicarbonate (100 mL) and extracted with EtOAc. Compound was dried over Na₂SO₄, and concentrated under vacuum to afford the title compound as an off white solid (1.6 g, 4 mmol).


Step 5) 1-(3-Chloro-benzyl)-7-(3-chloro-propoxy)-3-(naphthalene-1-sulfonyl)-1-H-indazole

15 A mixture of 7-(3-chloro-propoxy)-3-(naphthalene-1-sulfonyl)-1-H-indazole (0.7 g, 1.75 mmol), 3-chloro-benzylbromide (0.28 mL, 2.17 mmol), and cesium carbonate (0.7 g, 2.17 mmol) in DMF (5 mL) was stirred together in a round bottom flask at room temperature for 10 minutes. Reaction mixture was diluted with H₂O, extracted with EtOAc, washed with water (2x), brine (1x), dried over Na₂SO₄, and
20 concentrated under vacuum. The crude product was purified by HPLC using as eluent 30% EtOAc/hexane to afford the title compound as an off-white solid (0.87 g, 1.66 mmol).

Steps 6 and 7) Amination and Debenzylation


25 Using essentially the same procedures described in Examples 107 and 76 and employing the 7-(3-chloropropoxy)-3-naphthylsulfonylindazole substrate and the desired amine, the compounds shown in Table VIII were obtained and identified by HPLC and mass spectral analyses. (Amination was run in parallel manner on a heated carousel in 3-dram vials.)

TABLE VIII



Ex. No.	R1	R5	R6	[M+H] ⁺
128	3-Cl-benzyl	CH ₃	H	521
129	3-Cl-benzyl	C ₂ H ₅	H	535
130	3-Cl-benzyl	CH ₃	CH ₃	535
131	3-Cl-benzyl	C ₂ H ₅	C ₂ H ₅	563
132	3-Cl-benzyl	-CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ -		575
133	3-Cl-benzyl	-CH ₂ CH ₂ CH ₂ CH ₂ -		561
134	H	CH ₃	H	396
135	H	C ₂ H ₅	H	410
136	H	CH ₃	CH ₃	410
137	H	C ₂ H ₅	C ₂ H ₅	438
138	H	-CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ -		450

5

Example 139**2-{[3-Phenylsulfonyl)-1H-indazol-7-yl] oxy} ethanamine****Step 1) 1-Benzenesulfonylmethyl-3- (2-chloroethoxy)-2-nitrobenzene**

10

A mixture of 1-(2-chloro-ethoxy)-2-nitrobenzene (0.5 g, 2.5 mmoles) and 1-chloromethane-sulfonyl-benzene (0.56 g, 2.97 mmoles) was stirred in THF (10 mL) at -78 °C, in a round bottom flask under nitrogen. A solution of 1M potassium *t*-

butoxide was added dropwise (7.45 mL, 7.44 mmoles) over a half an hour period. Temperature was allowed to rise to $-40\text{ }^{\circ}\text{C}$, and the reaction mixture was stirred at this temperature for 5 hours. The reaction mixture was poured into cold 2N HCl, extracted with EtOAc, dried over Na_2SO_4 , and concentrated under vacuum.

- 5 Compound was recrystallized from CH_2Cl_2 /hexane to afford the title compound as an off-white solid (0.57 g, 1.6 mmoles).

Step 2) 2-Benzenesulfonylmethyl-6- (2-chloro-ethoxy)-phenylamine

- A mixture of 1-benzenesulfonylmethyl-3- (2-chloro-ethoxy)-2-nitrobenzene (0.57 g, 1.6 mmoles) in ethanol (10 mL) was stirred under nitrogen in a round bottom
10 flask at $60\text{ }^{\circ}\text{C}$. 10% Pd/C was added, and the temperature was increased to $80\text{ }^{\circ}\text{C}$. Hydrazine hydrate (1.5 mL) was added dropwise and the mixture was stirred at reflux for 3 hours. Reaction mixture was filtered off through Celite, and the solution was washed with H_2O (3x), dried over Na_2SO_4 , and concentrated under vacuum to afford the title compound as an off white solid (0.5 g, 1.53 mmoles).

- 15 **Step 3) 2-(2-Azido-ethoxy)-6-benzene-sulfonylmethyl-phenylamine**

- A mixture of 2-benzenesulfonylmethyl-6-(2-chloro-ethoxy)-phenylamine (0.5 g, 1.53 mmoles) and sodium azide (0.15 g, 2.29 mmoles) in DMSO (10 mL) was stirred together in a round bottom flask under nitrogen at $90\text{ }^{\circ}\text{C}$ for 3 hours. Reaction mixture was cooled to room temperature, diluted with water, extracted with EtOAc,
20 washed with water (2x), brine (1x), dried over Na_2SO_4 , and concentrated under vacuum. Crude compound was purified by normal phase HPLC using as eluent 40% EtOAc/hexane to afford the title compound as an off white solid (0.39 g, 1.17 mmoles).

Step 4) 7-(2-Azido-ethoxy)-3-benzenesulfonyl-1H-indazole


- 25 A mixture of 2-(2-azido-ethoxy-6-benzene-sulfonylmethyl-phenylamine (0.39 g, 1.17 mmoles) in THF (2 mL), and 4M HCl (10 mL) was stirred in a round bottom flask, under nitrogen, at $3\text{ }^{\circ}\text{C}$. A solution of sodium nitrite (0.08 g, 1.23 mmoles) in H_2O (1 mL) was added dropwise. The reaction mixture was poured into a cold solution of saturated sodium bicarbonate (100 mL) and extracted with EtOAc.
30 Compound was dried over Na_2SO_4 , and concentrated under vacuum to afford the title compound as an off white solid (0.3 g, 0.87 mmoles).

Step5) 2-[[3-Phenylsulfonyl]-1H-indazol-7-yl] oxy} ethanamine

A mixture of 7-(2-azido-ethoxy)-3-benzenesulfonyl-1*H*-indazole (0.3 g, 0.87 mmoles) and 10% Pd/C in THF (2 mL), and methanol (8 mL) was hydrogenated in a Parr hydrogenation bottle (250 mL) at 52 lb/in² for 2 hours. The mixture was filtered through Celite, and the filtrate was concentrated under vacuum. The crude product was recrystallized from CH₂Cl₂/hexane, 1M HCl in ether (0.9 mL, 0.9 mmoles) was added, then dried, to afford the title compound as an off-white HCl salt (0.2 g, 0.6 mmoles), MS: (ES⁺) 317 [M+H]⁺

Example 140

2-[[5-Fluoro-3-phenylsulfonyl)-1*H*-indazol-7-yl] oxy} ethanamine



Step 1) 1-Benzenesulfonylmethyl-3- (2-chloro-ethoxy)-5-fluoro-2-nitro-benzene

A mixture of 2-(2-chloroethoxy)-4-fluoro-1-nitrobenzene (1.3 g, 6 mmoles) and 1-chloromethane-sulfonyl-benzene (1.7 g, 9 mmoles) was stirred in THF (50 mL) at –78 °C, in a round bottom flask under nitrogen. A solution of 1M potassium *t*-butoxide was added dropwise (18 mL, 18 mmoles) over a half hour period. Temperature was allowed to rise to –40 °C, and the reaction mixture was stirred at this temperature for 5 hours. The reaction mixture was poured into cold 2N HCl, extracted with EtOAc, dried over Na₂SO₄, and concentrated under vacuum. Compound was recrystallized from CH₂Cl₂/hexane to afford the title compound as an off-white solid (1.98 g, 5.3 mmoles).

Step 2) 2-Benzenesulfonylmethyl-6- (2-chloro-ethoxy)-4-fluorophenylamine

A mixture of 1-benzenesulfonylmethyl-3- (2-chloro-ethoxy)-5-fluoro-2-nitrobenzene (1.98 g, 5.3 mmoles) in ethanol (25 mL) was stirred under nitrogen in a round bottom flask at 60 °C. 10% Pd/C was added, and the temperature was increased to 80 °C. Hydrazine hydrate (2.0 mL) was added dropwise and the mixture was stirred at reflux for 3 hours. Reaction mixture was filtered off through Celite, and the solution was washed with H₂O (3x), dried over Na₂SO₄, and concentrated under vacuum to afford the title compound as an off white solid (1.67 g, 4.87 mmoles).

Step 3) 2-(2-Azido-ethoxy)-6-benzenesulfonylmethyl-4-fluoro-phenylamine

A mixture of 2-benzenesulfonylmethyl-6- (2-chloro-ethoxy)-4-fluorophenylamine (1.67 g, 4.87 mmoles) and sodium azide (0.38 g, 5.84 mmoles) in DMSO (20 mL) was stirred together in a round bottom flask under nitrogen at 90 °C for 3 hours. Reaction mixture was cooled to room temperature, diluted with water, extracted with EtOAc, washed with water (2x), brine (1x), dried over Na₂SO₄, and concentrated under vacuum to afford the title compound as an off white solid (1.44 g, 4.14 mmoles).


Step 4) 7-(2-Azido-ethoxy)-3-benzenesulfonyl-5-fluoro-1H-indazole

A mixture of 2-(2-azido-ethoxy-6-benzene-sulfonylmethyl-4-fluoro-phenylamine (1.44 g, 4.14 mmoles) in THF (5 mL), and 4M HCl (20 mL) was stirred in a round bottom flask, under nitrogen, at 3 °C. A solution of sodium nitrite (0.28 g, 4.34 mmoles) in H₂O (2 mL) was added dropwise. The reaction mixture was poured into a cold solution of saturated sodium bicarbonate (100 mL) and extracted with EtOAc. Compound was dried over Na₂SO₄, and concentrated under vacuum to afford the title compound as an off white solid (1.34 g, 3.72 mmoles).

Step 5) 2-[[5-Fluoro-3-phenylsulfonyl]-1H-indazol-7-yl] oxy} ethanamine

A mixture of 7-(2-azido-ethoxy)-3-benzenesulfonyl-5-fluoro-1H-indazole (0.3 g, 0.83 mmoles) and 10% Pd/C in THF (2 mL), and methanol (8 mL) was hydrogenated in a Parr hydrogenation bottle (250 mL) at 52 lb/in² for 2 hours. The mixture was filtered through Celite, and the filtrate was concentrated under vacuum. The crude product was recrystallized from CH₂Cl₂/hexane, 1M HCl in ether (0.8 mL, 0.8 mmoles) was added, then dried, to afford the title compound as an off-white HCl salt (0.2 g, 0.59 mmoles), MS: (ES⁺) 336 [M+H]⁺

25

Example 141**2-[[3-(1-Naphthylsulfonyl)-1H-indazol-4-yl]oxy]ethanamine****Step 1) 1-(2-Chloroethoxy)-3-nitrobenzene**

5 A mixture of 3-nitrophenol (0.83 g, 6 mmoles), bromo-chloroethane (1.03 g, 7.2 mmoles), and K_2CO_3 (1.24 g, 9 mmoles) was stirred together in DMF at room temperature for 1 hour. Reaction mixture was diluted with H_2O , extracted with EtOAc, washed with water (2x), brine (1x), dried over Na_2SO_4 , and concentrated under vacuum. The crude product was purified by HPLC using as eluent 30%

10 EtOAc/hexane to afford the title compound as an off-white solid (1.12 g, 5.6 mmoles).

Step 2) 1-[2-(2-Chloroethoxy)-6-nitrophenylmethanesulfonyl]naphthalene

A mixture of 1-(2-chloro-ethoxy)-3-nitrobenzene (1.12 g, 5.6 mmoles) and 1-chloromethanesulfonyl-naphthalene (1.6 g, 6.72 mmoles) was stirred in THF (50 mL) at -78°C , in a round bottom flask under nitrogen. A solution of 1M potassium *t*-butoxide was added dropwise (16.8 mL, 16.8 mmoles) over a half an hour period.

15 Temperature was allowed to rise to -40°C , and the reaction mixture was stirred at this temperature for 5 hours. The reaction mixture was poured into cold 2N HCl, extracted with EtOAc, dried over Na_2SO_4 , and concentrated under vacuum. Compound was purified by normal phase HPLC using as eluent 40% EtOAc/hexane

20 to afford the title compound as an off-white solid (0.9 g, 2.24 mmoles), and 1-[4-(2-chloro-ethoxy)-2-nitro-phenylmethanesulfonyl]naphthalene (0.79 g, 2.1 mmoles).

Step 3) 3-(2-Chloro-ethoxy)-2-(naphthalene-1-sulfonylmethyl)-phenyl amine

A mixture of 1-[2-(2-Chloro-ethoxy)-6-nitro-phenylmethanesulfonyl]naphthalene (0.9 g, 2.24 mmoles) and 10% Pd/C in THF (10 mL), methanol (10 mL), and formic acid (2 mL) was hydrogenated in a Parr hydrogenation bottle (250 mL) at

25 40 lb/in^2 for 20 hours. The mixture was filtered through Celite, and the filtrate was

diluted with EtOAc, washed with water, dried over Na₂SO₄, and concentrated under vacuum to afford the title compound as an off-white solid (0.78 g, 2.1 mmoles).

Step 4) 4-(2-Chloro-ethoxy)-3-(naphthalene-1-sulfonyl)-1-H-indazole

5 A mixture of 3-(2-chloro-ethoxy)-2-(naphthalene-1-sulfonylmethyl)-phenyl amine (0.78 g, 2.1 mmoles) in THF (5 mL), and 4M HCl (10 mL) was stirred in a round bottom flask, under nitrogen, at 3 °C. A solution of sodium nitrite (0.15 g, 2.2 mmoles) in H₂O (1 mL) was added dropwise. The reaction mixture was poured into a cold solution of saturated sodium bicarbonate (100 mL) and extracted with EtOAc. Compound was dried over Na₂SO₄, and concentrated under vacuum to afford the title
10 compound as an off white solid (0.74 g, 1.93 mmoles).

Step 5) 4-(2-Azido-ethoxy)-3-(naphthalene-1-sulfonyl)-1H-indazole


A mixture of 4-(2-chloro-ethoxy)-3-(naphthalene-1-sulfonyl)-1-H-indazole (0.2 g, 0.5 mmoles) and sodium azide (0.04 g, 0.62 mmoles) in DMSO (2 mL) was stirred together in a round bottom flask under nitrogen at 90 °C for 3 hours. Reaction
15 mixture was cooled to room temperature, diluted with water, extracted with EtOAc, washed with water (2x), brine (1x), dried over Na₂SO₄, and concentrated under vacuum. Crude compound was purified by normal phase HPLC using as eluent 40% EtOAc/hexane to afford the title compound as an off white solid (0.17 g, 0.45 mmoles).

Step 6) 2-[[3-(1-Naphthylsulfonyl)-1H-indazol-4-yl] oxy} ethanamine

A mixture of 4-(2-azido-ethoxy)-3-(naphthalene-1-sulfonyl)-1H-indazole (0.17 g, 0.45 mmoles) and 10% Pd/C in THF (2 mL), and methanol (8 mL) was hydrogenated in a Parr hydrogenation bottle (250 mL) at 52 lb/in² for 2 hours. The mixture was filtered through Celite, and the filtrate was concentrated under vacuum.
25 The crude product was recrystallized from CH₂Cl₂/hexane, 1M HCl in ether (0.4 mL, 0.4 mmoles) was added, then dried to afford the title compound as an off-white HCl salt (0.15 g, 0.4 mmoles), MS: (ES⁺) 369 [M+H]⁺


Example 142

30 N-Methyl-2- {[3-naphthylsulfonyl)-1H-indazol-4-yl] oxy} ethanamine Hydrochloride



A mixture of 4-(2-chloro-ethoxy)-3-(naphthalene-1-sulfonyl)-1-*H*-indazole (0.075 g, 0.19 mmoles) and methylamine (0.28 mL, 0.56 mmoles) in DMSO (1 mL) was stirred under nitrogen at 100 °C for 4 hours. Mixture was cooled to room temperature, diluted with water, extracted with EtOAc, washed with water (2x), brine (1x), dried over Na₂SO₄, and concentrated under vacuum. Compound was purified by flash chromatography using as eluent 5% CH₃OH/EtOAc. The purified compound was dissolved in methanol, 1M HCl in ether (0.2 mL, 0.2 mmoles) was added, then dried to afford the title compound as the HCl salt (0.07 g, 0.19 mmoles), MS: (ES⁺) 417 [M+H]⁺


Examples 143-145



Using essentially the same procedure described in Example 142 and employing the desired amine, the compounds shown in Table IX were obtained and identified by HPLC and mass spectral analyses.

TABLE IX

Ex. No.	R5	R6	[M+H] ⁺
143	CH ₃	CH ₃	397
144	-CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ -		437
145	-CH ₂ CH ₂ CH ₂ CH ₂ -		423

Example 146**2-([3-(1-Naphthylsulfonyl)-1H-indazol-6-yl] oxy) ethanamine**

5

Step 1) 5-(2-Chloro-ethoxy-2- (naphthalene-1-sulfonylmethyl)-phenylamine

A mixture of 1-[4-(2-chloro-ethoxy)-2-nitro-phenylmethanesulfonyl]-naphthalene (0.79 g, 2.1 mmoles) and 10% Pd/C in THF (10 mL), methanol (10 mL), and formic acid (2 mL) was hydrogenated in a Parr hydrogenation bottle (250 mL) at 40 lb/in² for 20 hours. The mixture was filtered through Celite, and the filtrate was diluted with EtOAc, washed with water, dried over Na₂SO₄, and concentrated under vacuum to afford the title compound as an off-white solid (0.74 g, 1.99 mmoles).

10

Step 2) 6-(2-Chloro-ethoxy)-3-(naphthalene-1-sulfonyl)-1H-indazole

A mixture of 5-(2-chloro-ethoxy)-2-(naphthalene-1-sulfonylmethyl)-phenylamine (0.74 g, 1.99 mmoles) in THF (5 mL), and 4M HCl (10 mL) was stirred in a round bottom flask, under nitrogen, at 3 °C. A solution of sodium nitrite (0.14 g, 2.08 mmoles) in H₂O (1 mL) was added dropwise. The reaction mixture was poured into a cold solution of saturated sodium bicarbonate (100 mL) and extracted with EtOAc. Compound was dried over Na₂SO₄, and concentrated under vacuum to afford the title compound as an off white solid (0.74 g, 1.93 mmoles).

15

20


Step 3) 6-(2-Azido-ethoxy)-3-(naphthalene-1-sulfonyl)-1H-indazole

A mixture of 6-(2-chloro-ethoxy)-3-(naphthalene-1-sulfonyl)-1-H-indazole (0.19 g, 0.5 mmoles) and sodium azide (0.04 g, 0.62 mmoles) in DMSO (2 mL) was stirred together in a round bottom flask under nitrogen at 90 °C for 3 hours. Reaction mixture was cooled to room temperature, diluted with water, extracted with EtOAc, washed with water (2x), brine (1x), dried over Na₂SO₄, and concentrated under vacuum. Crude compound was purified by normal phase HPLC using as eluent 40% EtOAc/hexane to afford the title compound as an off white solid (0.17 g, 0.45 mmoles).


25

Step 4) 2-[[3-(1-Naphthylsulfonyl)-1H-indazol-6-yl]oxy} ethanamine


A mixture of 6-(2-azido-ethoxy)-3-(naphthalene-1-sulfonyl)-1H-indazole (0.17 g, 0.45 mmoles) and 10% Pd/C in THF (2 mL), and methanol (8 mL) was hydrogenated in a Parr hydrogenation bottle (250 mL) at 52 lb/in² for 2 hours. The mixture was filtered through Celite, and the filtrate was concentrated under vacuum. The crude product was recrystallized from CH₂Cl₂/hexane, 1M HCl in ether (0.4 mL, 0.4 mmoles) was added, then dried, to afford the title compound as an off-white HCl salt (0.15 g, 0.4 mmoles), MS: (ES⁺) 368 [M+H]⁺

Example 147**N-Methyl-2-[[3-(1-naphthylsulfonyl)-1H-indazol-6-yl]oxy}ethanamine Hydrochloride**

A mixture of 6-(2-chloro-ethoxy)-3-(naphthalene-1-sulfonyl)-1H-indazole (0.075 g, 0.19 mmoles) and methylamine (0.28 mL, 0.56 mmoles) in DMSO (1 mL) was stirred under nitrogen at 100 °C for 4 hours. Mixture was cooled to room temperature, diluted with water, extracted with EtOAc, washed with water (2x), brine (1x), dried over Na₂SO₄, and concentrated under vacuum. Compound was purified by flash chromatography using as eluent 5% CH₃OH/EtOAc. The purified compound was dissolved in methanol, 1M HCl in ether (0.2 mL, 0.2 mmoles) was added, then dried, to afford the title compound as the HCl salt (0.07 g, 0.19 mmoles), MS: (ES⁺) 381 [M+H]⁺

Examples 148-150


Using essentially the same procedure described in Example 147 and employing the desired amine, the compounds shown in Table X were obtained and identified by HPLC and mass spectral analyses.

TABLE X

Ex. No.	R5	R6	[M+H] ⁺
148	CH ₃	CH ₃	396
149	-CH ₂ CH ₂ CH ₂ CH ₂ CH ₂ -		436
150	-CH ₂ CH ₂ CH ₂ CH ₂ -		422

Example 151

N-[2-(Dimethylamino)ethyl]-3-(1-naphthylsulfonyl)-1H-indazole-5-carboxamide

**Step 1) 3-(Naphthalene 1-sulfonylmethyl)-4-nitro-benzoic acid methyl ester**

A mixture of 4-nitro-benzoic acid methyl ester (0.8 g, 4.4 mmoles) and 1-chloromethanesulfonyl-naphthalene (1.3 g, 5.3 mmoles) was stirred in THF (50 mL) at -78 °C, in a round bottom flask under nitrogen. A solution of 1M potassium *t*-butoxide was added dropwise (13 mL, 13 mmoles) over a half an hour period. Temperature was allowed to rise to -40 °C, and the reaction mixture was stirred at this temperature for 5 hours. The reaction mixture was poured into cold 2N HCl,

extracted with EtOAc, dried over Na₂SO₄, and concentrated under vacuum. Compound was purified by normal phase HPLC using as eluent 40% EtOAc/hexane to afford the title compound as an off-white solid (1.5 g, 3.9 mmoles).

Step 2) 4-Amino-3-(naphthalene-1-sulfonylmethyl)benzoic acid methyl ester

5 A mixture of 3-(naphthalene-1-sulfonylmethyl)-4-nitro-benzoic acid methyl ester (1.5 g, 3.9 mmoles) and 10% Pd/C in THF (10 mL), and methanol (20 mL) was hydrogenated in a Parr hydrogenation bottle (250 mL) at 52 lb/in² overnight. The mixture was filtered through Celite, and the filtrate was concentrated under vacuum to afford the title compound as an off-white solid (0.9 g, 2.5 mmoles).

10 **Step 3) 3-(Naphthalene-1-sulfonyl)-1H-indazole-5-carboxylic acid methyl ester**

 A mixture of 4-amino-3-(naphthalene-1-sulfonylmethyl)benzoic acid methyl ester (0.9 g, 2.5 mmoles) in THF (5 mL), and 4M HCl (10 mL) was stirred in a round bottom flask, under nitrogen, at 3 °C. A solution of sodium nitrite (0.18 g, 2.62 mmoles) in H₂O (1 mL) was added dropwise. The reaction mixture was poured into a
15 cold solution of saturated sodium bicarbonate (100 mL) and extracted with EtOAc. Compound was dried over Na₂SO₄, and concentrated under vacuum to afford the title compound as an off white solid (0.82 g, 2.25 mmoles).

Step 4) 1-(3-Chlorobenzyl)-2-(naphthalene-1-sulfonyl)-1H-indazole-5-carboxylic acid methyl ester

20 A mixture of 3-(naphthalene-1-sulfonyl)-1H-indazole-5-carboxylic acid methyl ester (0.82 g, 2.25 mmoles), 3-chlorobenzyl bromide (0.34 mL, 2.7 mmoles), and cesium carbonate (0.87 g, 2.7 mmoles) in DMF (5 mL) was stirred together in a round bottom flask at room temperature for 30 minutes. Reaction mixture was diluted with H₂O, extracted with EtOAc, washed with water (2x), brine (1x), dried over
25 Na₂SO₄, and concentrated under vacuum. The crude product was purified by HPLC using as eluent 30% EtOAc/hexane to afford the title compound as an off-white solid (1.01 g, 2.07 mmoles).

Step 5) 1-(3-Chloro-benzyl)-3-(naphthalene-1-sulfonyl)-1H-indazole-5-carboxylic acid (2-dimethyl-amino-ethyl)-amide

30 To a solution of dimethyl ethylene diamine (0.02 mL, 0.2 mmoles) in THF (2 mL), cooled to 0 °C was added LDA dropwise (0.15 mL, 0.3 mmoles). To this mixture was then added a solution of 1-(3-chloro-benzyl)-2-(naphthalene-1-sulfonyl)-1H-indazole-5-carboxylic acid methyl ester (0.05 g, 0.1 mmoles) in THF (1 mL).

Mixture was allowed to warm slowly to room temperature. Reaction mixture was diluted with water, extracted with EtOAc (1x), CH₂Cl₂ (1x); the organics were washed with brine (1x), and concentrated under vacuo to afford the title compound (0.3 g, 0.04 mmoles).


5 **Step 6) N-[2-(Dimethylamino) ethyl]-3-(1-naphthylsulfonyl)-1H-indazole-5-carboxamide**

A mixture of 1-(3-chloro-benzyl)-3-(naphthalene-1-sulfonyl)-1H-indazole-5-carboxylic acid (2-dimethyl-amino-ethyl)-amide (0.3 g, 0.04 mmoles), DMSO (1 mL) and *t*-BuOH (0.2 mL) was stirred at room temperature in a round bottom flask under oxygen atmosphere. A solution of potassium *t*-butoxide (0.05 mL, 0.05 mmoles) was added dropwise and the reaction mixture stirred for 30 min. Reaction mixture was quenched with saturated ammonium chloride, extracted with EtOAc, dried over Na₂SO₄, and concentrated under vacuum. Crude compound was purified by reverse phase chromatography to afford the title compound, MS: (ES⁺) 423 [M+H]⁺

15

Example 152

3-(1-Naphthylsulfonyl)-N-(2-piperidin-1-ylethyl)-1H-indazole-5-carboxamide




Using essentially the same procedure described in Example 151 and employing 1-(2-aminoethyl)piperidine in step 5, the title compound is obtained and identified by HPLC and mass spectral analyses, MS: (ES⁺) 463 [M+H]⁺.

20

Example 153

N, N, N'-Trimethyl-N'-{[3-(1-naphthylsulfonyl)-1H-indazol-5-yl]methyl}ethane-1,2-diamine

25



Step 1) 2-[3-(Naphthalene-1-sulfonylmethyl)-4-nitrophenyl]-[1,3]dioxolane

A mixture of 2-(4-Nitro-phenyl)-[1, 3]dioxolane (1.85 g, 9.5 mmol) and 1-chloromethanesulfonyl-naphthalene (2.74 g, 11.4 mmol) was stirred in THF (50 mL) at -78 °C, in a round bottom flask under nitrogen. A solution of 1M potassium *t*-butoxide was added dropwise (19 mL, 19 mmol) over a half an hour period. Temperature was allowed to rise to -40 °C, and the reaction mixture was stirred at this temperature for 5 hours. The reaction mixture was poured into cold 2N HCl, extracted with EtOAc, dried over Na₂SO₄, and concentrated under vacuum. Compound was purified by normal phase HPLC using as eluent 40% EtOAc/hexane to afford the title compound as an off-white solid (3.03 g, 7.6 mmol).

Step 2) 3-(Naphthalene-1-sulfonylmethyl)-4-nitrobenzaldehyde

A mixture of 2-[3-(naphthalene-1-sulfonylmethyl)-4-nitro-phenyl]-[1,3]dioxolane (3.03 g, 7.6 mmol), and 2N HCl (4 mL, 8 mmol) in THF (30 mL) was stirred at 40 °C for 4 hours. The reaction mixture was cooled to room temperature, diluted with water, extracted with EtOAc, dried over Na₂SO₄, and concentrated under vacuum to yield the title compound (2.56 g, 7.22 mmol).

Step 3) 4-Amino-3-(naphthalene-1-sulfonylmethyl)benzaldehyde

A mixture of 3-(naphthalene-1-sulfonylmethyl)-4-nitro-benzaldehyde (2.5 g, 7.22 mmol) and 10% Pd/C in THF (10 mL), and methanol (20 mL) was hydrogenated in a Parr hydrogenation bottle (250 mL) at 52 lb/in² overnight. The mixture was filtered through Celite, and the filtrate was concentrated under vacuum to afford the title compound as an off-white solid (2.4 g, 6.85 mmol).

Step 4) 3-(Naphthalene-1-sulfonyl)-1H-indazole-5-carbaldehyde

A mixture of 4-amino-3-(naphthalene-1-sulfonylmethyl)-benzaldehyde (2.4 g, 6.85 mmoles) in THF (10 mL) and 4M HCl (20 mL) was stirred in a round bottom flask at 3 °C. A solution of sodium nitrite (0.49 g, 7.19 mmoles in H₂O (2 mL) was added. The reaction mixture was poured into a cold solution of saturated sodium bicarbonate (100 mL) and extracted with EtOAc. Compound was dried over Na₂SO₄, and concentrated under vacuum to afford the title compound as an off white solid (1.84 g, 5.5 mmoles).


Step 5) N, N, N'-Trimethyl-N'-[[3-(1-naphthylsulfonyl)-1H-indazol-5-yl]methyl]ethane-1,2-diamine

3-(Naphthalene-1-sulfonyl)-1H-indazole-5-carbaldehyde (0.17 g, 0.5 mmol), trimethyl ethylene diamine (0.6 mmol) and sodium triacetoxyborohydride (0.7 mmol) in dichloroethane (5 mL) was stirred at room temperature for 24 hrs. After completion, the solvent was removed *in vacuo*, crude material dispersed in water and the pH brought to 3.4. Solid material was filtered off and washed with cold water to afford after drying the target material as a free base. The latter was converted into hydrochloride salt by dissolution in methanol, followed by treatment with the excess of 2N HCl and the evacuation of the volatiles *in vacuo* to afford the title compound hydrochloride salt, mp > 200 °C; MS (APPI) 423 [M+H]

20

Example 154


(3S)-N-[[3-(1-Naphthylsulfonyl)-1H-indazol-5-yl]methyl]pyrrolidin-3-amine Hydrochloride



25


Using essentially the same procedure described in Example 153, step 5, and employing Boc-protected S(-)-pyrrolidin-3-ylamine, followed by the removal of the Boc-group by TFA in methylene chloride and treatment of the free base with 2N HCl, the title product was obtained, mp > 200 °C; MS (ES) (M+H)⁺ 407.1; MS (ES) (M+H+Na)⁺ 429.1

30

Example 155**N-[[3-(1-Naphthylsulfonyl)-1H-indazol-5-yl]methyl]ethane-1,2-diamine**

5 3-(Naphthalene-1-sulfonyl)-1H-indazole-5-carbaldehyde (0.3 mmol) was stirred for 24 hrs with the excess of ethylene diamine (1 mmol) in methanol. Sodium borohydride (0.6 mmol) was added and stirring continued for another 24 hrs. After completion, the volatiles were removed *in vacuo*, crude material diluted with cold water, acidified to pH 3.4, filtered off, washed on a filter with cold water and dried to afford the target material as a free base. The latter was converted into hydrochloride salt by dissolution in methanol, followed by treatment with the excess of 2N HCl and the evacuation of the volatiles *in vacuo* to afford the title compound, mp > 200 °C; MS (ES⁺) 381[M+H]⁺

15

Example 156**N,N-Dimethyl-2-[[3-(1-naphthylsulfonyl)-1H-indazol-5-yl]methoxy]ethanamine****Step 1) 1-(3-Chloro-benzyl)-3-naphthalene-1-sulfonyl)-1H-indazole-5-carbaldehyde**

20 A mixture of 3-(1-naphthylsulfonyl)-1H-indazole-5-carbaldehyde (0.17 g, 0.5 mmoles), 3-chlorobenzyl bromide (0.07 mL, 0.6 mmoles), and cesium carbonate (0.19 g, 0.6 mmoles) in DMF (5 mL) was stirred together in a round bottom flask at room temperature for 30 minutes. Reaction mixture was diluted with H₂O, extracted with EtOAc, washed with water (2x), brine (1x), dried over Na₂SO₄, and concentrated under vacuum. The crude product was purified by normal phase HPLC using as

25

eluent 30% EtOAc/hexane to afford the title compound as an off-white solid (0.18 g, 0.4 mmoles).

Step 2) [1-(3-Chlorobenzyl)-3-(1-naphthylsulfonyl)-1H-indazol-5-yl]-methanol

1-(3-Chloro-benzyl)-3-(naphthalene-1-sulfonyl)-1H-indazole-5-carbaldehyde
5 (0.18g, 0.4 mmol) was dissolved in THF and treated under nitrogen with the excess of sodium borohydride (0.2 mmol). After 1 hr the reaction mixture was diluted with water and the product extracted with methylene chloride to afford after evaporation of the solvent the title compound as a colorless solid (0.18 g, 0.4 mmoles).

Step 3) 1-(3-Chlorobenzyl)-5-(2-chloroethoxymethyl)-3-(1-naphthylsulfonyl)-1H-indazole

[1-(3-Chlorobenzyl)-3-(1-naphthylsulfonyl)-1H-indazol-5-yl]-methanol (0.18 g, 0.4 mmol) and diisopropyl ethyl amine (1 mmol) in methylene chloride were treated at -20 °C with trifluoromethanesulfonic anhydride (0.14 g, 0.5 mmol). After stirring for 30 min an excess of 2-chloroethanol (2 mmol) was added and the mixture stirred for
15 additional 2 hrs at -20 °C and at room temperature for 12 hrs. The reaction mixture was diluted with water, extracted with ethyl acetate and after evaporation of solvent, chromatographed on silica gel using 40% ethyl acetate/hexane as an eluent to afford the target material (0.08 g, 0.16 mmoles).

Step 4) {2-[1-(3-Chlorobenzyl)-3-(naphthalene-1-sulfonyl)-1H-indazol-5-ylmethoxy]ethyl}dimethylamine

A mixture of 1-(3-chlorobenzyl)-5-(2-chloroethoxymethyl)-3-(1-naphthylsulfonyl)-1H-indazole (0.08 g, 0.16 mmoles) and dimethylamine (0.28 ml, 0.56 mmoles) in DMSO (1 mL) was stirred under nitrogen at 100 °C for 4 hours. Mixture was cooled to room temperature, diluted with water, extracted with EtOAc,
25 washed with water (2x), brine (1x), dried over Na₂SO₄, and concentrated under vacuum. Compound was purified by flash chromatography using as eluent 5% CH₃OH/EtOAc to afford the title compound (0.08 g, 0.15 mmoles).

Step 5) N,N-Dimethyl-2-{[3-(1-naphthylsulfonyl)-1H-indazol-5-yl]methoxy}ethanamine


30 A mixture of {2-[1-(3-chloro-benzyl)-3-(naphthalene-1-sulfonyl)-1H-indazol-5-ylmethoxy]-ethyl}-dimethyl-amine (0.08 g, 0.15 mmoles) in DMSO (1 mL) and *t*-BuOH (0.2 mL) was stirred at room temperature in a round bottom flask under oxygen atmosphere. A solution of potassium *t*-butoxide (0.05 mL, 0.05 mmoles) was added

dropwise and the reaction mixture stirred for 30 min. Reaction mixture was quenched with saturated ammonium chloride, extracted with EtOAc, dried over Na_2SO_4 , and concentrated under vacuum. Crude compound was converted into the HCl salt as previously described to afford the title compound, mp > 200 °C; MS (ES^+) 410 $[\text{M}+\text{H}]^+$.

Example 157


N1-[3-(Naphthalene-1-sulfonyl)-1H-indazol-5-yl]-ethane-1,2-diamine dihydrochloride

Step 1




3-Iodo-5-nitro-1H-indazole Iodine (26.46 g, 104.27 mmol) and potassium hydroxide pellets (11.70 g, 208.54 mmol) were successively added into a DMF (104 mL) solution of 5-nitroindazole (8.50 g, 52.13 mmol) at room temperature and stirred for 4 days. The reaction mixture was then poured into NaHSO_3 solution (11.06 g in 200 mL water). The brown color faded away, and the formed yellow precipitate was filtered and washed with water and dried *in vacuo* to provide the title compound as a yellow solid (14.74 g, 98% yield). MS (ES^+) m/e 290 (MH^+)


Step 2



3-(Naphthalen-1-ylsulfanyl)-5-nitro-1H-indazole A mixture of 3-iodo-5-nitro-1H-indazole (10.00 g, 34.60 mmol), 1-naphthylethiol (5.54 g, 34.60 mmol), CuI (0.659 g, 3.46 mmol), ethylene glycol (4.30 g, 69.20 mmol) in isopropanol (49.40 mL) was heated at 90 °C under nitrogen overnight, cooled, diluted with 30% MeOH in CH_2Cl_2 , and passed through a pad of silica gel. The solution was concentrated *in vacuo* and purified by chromatography with 1% MeOH in CH_2Cl_2 to provide the title compound (5.5 g, 49%). MS (ES^+) m/e 322 (MH^+).


Step 3

3-(Naphthalene-1-sulfonyl)-1H-indazol-5-ylamine A mixture of 3-(naphthalen-1-ylsulfanyl)-5-nitro-1H-indazole (5.50 g, 17.11 mmol) and 3-chloroperoxybenzoic acid (17.91 g, 103.80 mmol) in CHCl_3 (115 mL) was stirred at room temperature for 4 hr, diluted with EtOAc, washed with Na_2SO_3 solution, water, brine, dried over Na_2SO_4 , and concentrated *in vacuo* to afford the crude intermediate which was carried out directly for the next step reaction without further purification. The mixture of the crude sulfone intermediate, tin mossy (15.79 g, 133.01 mmol) in MeOH and conc. hydrochloric acid was heated at 60 °C, diluted with CH_2Cl_2 , and neutralized to basic with NaOH or Na_2CO_3 solution. The aqueous layer was extracted with CH_2Cl_2 . Combined organic layers were dried over Na_2SO_4 and concentrated *in vacuo* followed by chromatography purification to provide the title compound (2.50 g, 45% overall yield). MS (ES^+) m/e 324 (MH^+)

Step 4**N1-[3-(Naphthalene-1-sulfonyl)-1H-indazol-5-yl]-ethane-1,2-diamine**

dihydrochloride A mixture of 3-(naphthalen-1-sulfonyl)-1H-indazol-5-ylamine hydrochloride (334 mg, 0.93 mmol), 2-oxazolidone (81 mg, 0.93 mmol), and diethylene glycol monomethyl ether (0.16 mL) was heated at 170 °C overnight, diluted with MeOH, and purified by reverse phase HPLC followed by conversion to HCl salt by treatment with HCl solution to provide the title compound as a white solid (86 mg, 21% yield). MS (ES^+) m/e 367 (MH^+)

Example 158



3-Amino-N-[3-(naphthalene-1-sulfonyl)-1H-indazol-5-yl]-propionamide

hydrochloride A mixture of 3-(naphthalene-1-sulfonyl)-1H-indazol-5-ylamine (500


mg, 1.55 mmol), N-*t*-Boc-β-alanine (381 mg, 2.01 mmol), 1-[3-

- 5 (dimethylamino)propyl]-3-ethylcarbodiimide hydrochloride (386 mg, 2.01 mmol) in CH₃CN was stirred at room temperature overnight and concentrated to dryness. The resulting residue was subjected to TFA, concentrated, and purified by reverse phase HPLC followed by treatment with HCl solution to provide the title compound as a white solid (180 mg, 24% yield). MS (ES⁺) *m/e* 395 (MH⁺)

10

Example 159


Step 1



{1-[3-(Naphthalene-1-sulfonyl)-1H-indazol-5-ylcarbamoyl]-ethyl}-carbamic acid

- 15 **tert-butyl ester** A mixture of 3-(naphthalene-1-sulfonyl)-1H-indazol-5-ylamine (500 mg, 1.55 mmol), N-*t*-Boc-alanine (381 mg, 2.01 mmol), 1-[3-(dimethylamino)propyl]-3-ethylcarbodiimide hydrochloride (386 mg, 2.01 mmol) in CH₃CN was stirred at room temperature overnight, concentrated, and purified by chromatography with 3% MeOH in CH₂Cl₂ to provide the title compound (110 mg, 48%), characterized by NMR and
- 20 mass spectral analyses.

Step 2



N1-[3-(Naphthalene-1-sulfonyl)-1H-indazol-5-yl]-propane-1,2-diamine

- 25 **hydrochloride** {1-[3-(Naphthalene-1-sulfonyl)-1H-indazol-5-ylcarbamoyl]-ethyl}-

carbamic acid *tert*-butyl ester (120 mg, 0.37 mmol) was subjected to TFA at room temperature for 2 hr and concentrated to dryness. The resulting residue was heated with BH_3 in THF (1 M, 4.5 mL) at reflux overnight. To the mixture was slowly added HCl (6 M, 1 mL). The resulting solution was heated at 80 °C for 20 min, concentrated, and purified by reverse phase HPLC followed by treatment with HCl solution to provide the title compound (35 mg, 38%). MS (ES^+) m/e 381 (MH^+)


Example 160

(S)-3-Methyl-N1-[3-(naphthalene-1-sulfonyl)-1H-indazol-5-yl]-butane-1,2-diamine dihydrochloride

Using essentially the same procedure described in Example 159 and employing (S)-*t*-Boc-Valine as the starting material, the title compound was obtained, MS (ES^+) m/e 409 (MH^+)

Example 161


N1-[3-(Naphthalene-1-sulfonyl)-1H-indazol-7-yl]-ethane-1,2-diamine dihydrochloride




Using essentially the same procedure described in Example 157 (step 4) and employing product from Example 162 (step 1) as the starting material, the title compound was obtained, MS (ES^+) m/e 367 (MH^+)

Example 162


3-Dimethylamino-N-[3-(naphthalene-1-sulfonyl)-1H-indazol-7-yl]-propionamide hydrochloride

Step 1

3-(Naphthalene-1-sulfonyl)-1H-indazol-7-ylamine The title compound was prepared according to the procedure described in Example 157 (steps 1-3) and employing 7-nitro-indazole as the starting material.


Step 2

3-Dimethylamino-N-[3-(naphthalene-1-sulfonyl)-1H-indazol-7-yl]-propionamide dihydrochloride The title compound was prepared in a similar manner as described in Example 158 and employing the appropriate starting material. MS (ES⁺) *m/e* 423 (MH⁺)

Example 163

N-[3-(Naphthalene-1-sulfonyl)-1H-indazol-7-yl]-3-piperidin-1-yl-propionamide hydrochloride The title compound was prepared in a similar manner as described in Example 158 and employing the appropriate starting material. MS (ES⁺) *m/e* 463 (MH⁺)

Example 164




3-Amino-N-[3-(naphthalene-1-sulfonyl)-1H-indazol-7-yl]-propionamide

hydrochloride The title compound was prepared in a similar manner as described in Example 158 and employing the appropriate starting material. MS (ES⁺) *m/e* 395 (MH⁺)

5

Example 165



3-Amino-N-[3-(naphthalene-1-sulfonyl)-1H-indazol-6-yl]-propionamide


hydrochloride

10

Using essentially the same procedure described in Example 158 and employing 6-amino-3-(1-naphthylsulfonyl)-1H-indazole as the starting material, the title compound was obtained, MS (ES⁺) *m/e* 395 (MH⁺)

15


Example 166




3-Diethylamino-N-[3-(naphthalene-1-sulfonyl)-1H-indazol-7-yl]-propionamide hydrochloride

Using essentially the same procedure described in Example 158 and employing 7-amino-3-(1-naphthylsulfonyl)-1H-indazole and the desired amino acid as the starting materials, the title compound was obtained. MS (ES⁺) *m/e* 451 (MH⁺)


20

Example 167**N-[3-(Naphthalene-1-sulfonyl)-1H-indazol-6-yl]-3-piperidin-1-yl-propionamide
hydrochloride**


Using essentially the same procedure described in Example 158 and employing 6-amino-3-(1-naphthylsulfonyl)-1H-indazole and the desired amino acid as the starting materials, the title compound was obtained, MS (ES^+) m/e 463 (MH^+).

Example 168**3-Dimethylamino-N-[3-(naphthalene-1-sulfonyl)-1H-indazol-6-yl]-propionamide
hydrochloride**

Using essentially the same procedure described in Example 158 and employing 6-amino-3-(1-naphthylsulfonyl)-1H-indazole and the desired amino acid as starting materials, the title compound was obtained, MS (ES^+) m/e 423 (MH^+).

Example 169**3-Diethylamino-N-[3-(naphthalene-1-sulfonyl)-1H-indazol-6-yl]-propionamide
hydrochloride**

Using essentially the same procedure described in Example 158 and employing 6-amino-3-(1-naphthylsulfonyl)-1H-indazole and the desired amino acid as starting materials, the title compound was obtained, MS (ES^+) m/e 451 (MH^+).


Example 170

N1-[3-(Naphthalene-1-sulfonyl)-1H-indazol-6-yl]-ethane-1,2-diamine dihydrochloride


- 5 Using essentially the same procedure described in Example 158 and employing 6-amino-3-(1-naphthylsulfonyl)-1H-indazole as starting material, the title compound was obtained, MS (ES⁺) *m/e* 367 (MH⁺).

Example 171

10 **N,N-Dimethyl-N-{2-[3-(phenylsulfonyl)-1H-indazol-7-yl]ethyl}amine hydrochloride**

Step 1


- 3-Bromo-2-nitrobenzyl phenyl sulfone** To a stirred solution of 1-bromo-2-nitrobenzene (10.1 g, 50 mmol) and chloromethylphenyl sulfone (9.5 g, 50 mmol) in dry THF (100 mL) at -65°C under nitrogen is added 1.0M KOtBu in THF (110 mL, 110 mmol). The deep purple reaction is allowed to warm to 0 °C over 1.5 hours and then treated with glacial acetic acid (8 mL). The reaction is diluted with water (200 mL) and saturated aqueous NaHCO₃ (200 mL), and then extracted with CH₂Cl₂ (2 x 400 mL). The extracts are dried (MgSO₄) and concentrated *in vacuo* to a light orange solid. Trituration with ethyl acetate and hexanes affords the title compound as a pale yellow solid (13 g, 73%). Mp: 138 - 141 °C. MS (ES⁻): 354 (M-H)

Step 2

25


6-Bromo-2-[(phenylsulfonyl)methyl]aniline Catalytic hydrogenation of 6-bromo-2-nitrobenzyl phenyl sulfone (0.36 g, 1 mmol) in the presence of platinum on carbon, disulfided and hydrogen (45 psi) in ethyl alcohol (40 mL) for 1 hour gives a reaction mixture. The reaction mixture is filtered through celite and concentrated *in vacuo* to give the title compound as a light brown solid (0.32 g, 99%). Mp: 174 - 177 °C. MS (ES+): 326 (M+H)

Step 3



- 10 **7-Bromo-3-(phenylsulfonyl)-1H-indazole** A solution of NaNO₂ (0.91 g, 13.35 mmol) in H₂O (10 mL) was added to a solution of 6-bromo-2-[(phenylsulfonyl)methyl]aniline (2.9 g, 8.9 mmol) in 70 mL of 4 N HCl at approximately 5 °C. The reaction mixture was stirred for 30 min. at 0 °C, and neutralized with 10 % NaOH. The resulting solid was collected by filtration, washed with water and purified by flash chromatography
- 15 (25% ethyl acetate/petroleum ether) to afford the title compound as a pink solid (2.27g, 91 %) Mp: 173 -175 °C, MS (ES-) 335 (M-H)


Step 4



- 20 **3-(Phenylsulfonyl)-7-vinyl-1H-indazole** A mixture of 7-Bromo-3-(phenylsulfonyl)-1H-indazole (2.72 g, 8.07 mmol) and dichlorobis(tri-*o*-tolylphosphine)-palladium(II) (0.94 g, 1.2 mmol) were dissolved in toluene (220 mL) and stirred for 10 minutes at room temperature under a nitrogen atmosphere. Tributyl(vinyl)tin (3.3 mg, 10.55 mmol) was added and the mixture was refluxed for 15 min or until turned black. The
- 25 mixture was cooled to room temperature, diluted with ethyl acetate (150 mL), 1 M KF (25 mL) and stirred for 12 hours. The resulting tin salt precipitate was removed by suction filtration and the organic layer was washed with water (100 mL), then brine and dried over MgSO₄, filtered and concentrated *in vacuo*. Purification by flash


chromatography (40% ethyl acetate/petroleum ether) gave the title compound as a light yellow solid (2.02 g, 88%). Mp: 129 – 130 °C. MS (ES⁺): 285 (M+H)

Step 5



5 **2-[3-(Phenylsulfonyl)-1H-indazol-7-yl]ethanol** To a solution of 3-(phenylsulfonyl)-7-vinyl-1H-indazole (2.0 g, 7.04 mmol) in THF (40 mL) at 0 °C was added dropwise BH₃-THF (15 mL of 1 M THF solution, 15 mmol). The solution was stirred for 3 hours at 0 °C, and H₂O (20 mL) was added slowly. To this mixture was added 10% NaOH
10 (17 mL), 30% H₂O₂ (15 mL) and the mixture was stirred vigorously at room temperature overnight. The mixture was partitioned between ethyl acetate and H₂O, and the aqueous layer was extracted with ethyl acetate (2 x 50 mL). The combined organic layers were washed with brine, dried (MgSO₄), filtered and concentrated under reduced pressure. Silica gel chromatography (50% ethyl
15 acetate/dichloromethane) gave the title compound as a white solid (1.42 g, 66%). Mp: 137 – 138 °C. MS (ES⁺): 303 (M+H)


Step 6



20 **2-[3-(Phenylsulfonyl)-1H-indazol-7-yl]ethyl 4-methylbenzenesulfonate** To a solution of 2-[3-(phenylsulfonyl)-1H-indazol-7-yl]ethanol (0.64 g, 2.12 mmol) in anhydrous dichloromethane (30 mL) at 0 °C was added pyridine (0.43 mL) and toluenesulfonyl chloride (0.475 g, 2.5 mmol). The solution was stirred for 12 hours at room temperature. The mixture was concentrated *in vacuo* and taken up in ethyl
25 acetate (30 mL). The organic layer was washed with 2 M HCl, (2 x 25 mL), brine, dried (MgSO₄), filtered and concentrated under reduced pressure. Silica gel

chromatography (50% ethyl acetate/petroleum ether) gave the title compound as a white foam (0.9 g, 93%). Mp: 61 – 64 °C. MS (ES⁻): 455 (M-H)

Step 7




N,N-Dimethyl-N-{2-[3-(phenylsulfonyl)-1H-indazol-7-yl]ethyl}amine

hydrochloride A solution of 2-[3-(phenylsulfonyl)-1H-indazol-7-yl]ethyl 4-methylbenzenesulfonate (0.137 g, 0.3 mmol) in anhydrous THF (2 mL) was added an excess of dimethylamine (0.5 mL of 2 M THF solution, 1 mmol) and heated to 71 °C for 24 hours. The mixture was partitioned between ethyl acetate and H₂O, and the aqueous layer was extracted with ethyl acetate (10 mL). The combined organic layers were washed with 2 M NaOH, (2 x 15 mL), brine, dried (MgSO₄), filtered and concentrated under reduced pressure. The product was passed through a plug of silica gel eluting with (20% EtOH/2N ammonia/dichloromethane) to give the title compound as a white solid (0.049 g, 77%). This solid was dissolved in diethyl ether and treated with 1 N HCl in diethyl ether (0.12 mL, 0.12 mmol) to afford a white precipitate isolated by vacuum filtration. Mp: 76 – 80 °C. MS (ES⁻): 328 (M-H)


Example 172

N-{2-[3-(phenylsulfonyl)-1H-indazol-7-yl]ethyl}cyclopropanamine hydrochloride




Using essentially the same procedure described in Example 171, step 7, and employing cyclopropyl amine, the title compound was obtained as a white solid, mp 115 – 117 °C, MS: (M+H) 342.

Example 173

N-Methyl-N-{2-[3-(phenylsulfonyl)-1H-indazol-7-yl]ethyl}amine trifluoroacetate

Using essentially the same procedure described in Example 171, step 7, and
 5 employing methylamine and substituting trifluoroacetic acid for HCl, the title
 compound was obtained as a clear glass, MS: (M+H) 316.

Example 174**{2-[3-(Phenylsulfonyl)-1H-indazol-7-yl]ethyl}amine Hydrochloride**


10

Step 1) 7-(2-Azidoethyl)-3-(phenylsulfonyl)-1H-indazole

A solution of 2-[3-(phenylsulfonyl)-1H-indazol-7-yl]ethyl 4-methylbenzenesulfonate (0.14 g, 0.3 mmol) in anhydrous DMF (2.5 mL) was added sodium azide (0.06 g, 0.9 mmol) and heated to 100 °C for 6 hours. The mixture was partitioned
 15 between ethyl acetate and H₂O, and the aqueous layer was extracted with ethyl acetate (10 mL). The combined organic layers were washed with brine (1 x 15 mL), dried (MgSO₄), filtered and concentrated under reduced pressure. Silica gel chromatography (40% Ethyl acetate/petroleum ether) gave the azide as a white solid (0.08 g, 86%). Mp: 105 – 107 °C. MS (ES⁻): 326 (M-H)

Step 2) {2-[3-(Phenylsulfonyl)-1H-indazol-7-yl]ethyl}amine Hydrochloride


7-(2-Azidoethyl)-3-(phenylsulfonyl)-1H-indazole (0.08 g, 0.20 mmol) was reduced by catalytic hydrogenation in the presence of palladium on carbon and hydrogen (40 psi) in ethanol (30 mL) for 4 hours. The reaction mixture was filtered through Celite and concentrated *in vacuo* to give a white solid (0.06 g, 99%). This
 25 solid was dissolved in diethyl ether and treated with 1 N HCl in diethyl ether (0.21 mL, 0.21 mmol) to afford a tan precipitate isolated by vacuum filtration. Mp: 157 - 160 °C. MS (ES⁻): 300 (M-H)

Example 175**N-Methyl-N-{2-[3-(phenylsulfonyl)-1H-indazol-5-yl]ethyl}amine****Step 1**

5

Methyl [2-(4-nitrophenyl)ethyl]carbamate To a stirred solution of [2-(4-nitrophenyl)ethyl]amine (6.06 g, 30 mmol) in CH_2Cl_2 (75 ml), MeOH (5 mL), and TEA (9.5 mL) at 0 °C was added chloromethylformate (3.39 g, 36 mmol) dropwise. The reaction mixture was stirred at room temperature for 1 hr. concentrated *in vacuo*, and partitioned between ethyl acetate and water; the aqueous layer was extracted with ethyl acetate. The combined organic layers were washed with water (50 mL) and brine (50 mL). The organic layer was dried (MgSO_4), filtered and concentrated under reduced pressure to afford a yellow solid (6.6 g, 98%). Mp: 38 °C. MS (ES⁻): 223 (M-H).

10


Step 2

Methyl methyl[2-(4-nitrophenyl)ethyl]carbamate To a stirred solution of methyl [2-(4-nitrophenyl)ethyl]carbamate (224 mg, 1 mmol) in THF (2 mL) was added KO^tBu (201 mg, 1.8 mmol) and MeI (256 mg, 1.8 mmol) sequentially. The reaction mixture was stirred for 12 h at room temperature. The reaction was diluted with water (5 mL), extracted with EtOAc (2 x 15 mL). The organics were washed with water and brine, dried (MgSO_4), filtered and concentrated under reduced pressure. Silica gel chromatography (40% ethyl acetate/ petroleum ether) afforded the title compound as a semi solid (160 mg, 67.2 %). MS (ES⁺): 239 (M+H)⁺.

20


25

Step 3

**Methyl methyl (2-(4-amino-3-((phenylsulfonyl)methyl)phenyl)ethyl)carbamate**


Using substantially the same manner as described in Example 171, step 1 and
 5 employing methyl methyl[2-(4-nitrophenyl)ethyl]carbamate (2.47 g, 10.38 mmol), the
 title compound was obtained as a white solid, (2.06 g, 51%). Mp: 42 °C, MS (ES+):
 393 (M+H)⁺.

Step 4

**Methyl (2-(4-amino-3-((phenylsulfonyl)methyl)phenyl)ethyl)methylcarbamate**


Catalytic hydrogenation of methyl methyl (2-(4-nitro-3-((phenylsulfonyl)methyl)-
 phenyl)ethyl)carbamate (1.8 g, 4.6 mmol) in the presence of Raney nickel and
 hydrogen (45 psi) in ethyl acetate/ethanol (60 mL) for 2 h gives a reaction mixture.
 15 The reaction mixture is filtered through celite and concentrated *in vacuo* to give the
 title compound as a light brown solid (1.36 g, 82%). Mp: 95 - 96 °C. MS (ES+): 363
 (M+H)⁺.

Step 5




Methyl methyl{2-[3-(phenylsulfonyl)-1*H*-indazol-5-yl]ethyl}carbamate Using substantially the same procedure described in Example 171, step 3 and employing methyl (2-{4-amino-3-[(phenylsulfonyl)methyl]phenyl}ethyl)methylcarbamate (0.362 g, 1 mmol), the title compound was obtained as a white solid, (0.29 g, 78%). Mp: 145 - 147 °C, MS (ES⁺): 374 (M+H)⁺

Step 6



N-Methyl-N-{2-[3-(phenylsulfonyl)-1*H*-indazol-5-yl]ethyl}amine To a solution of methyl methyl{2-[3-(phenylsulfonyl)-1*H*-indazol-5-yl]ethyl}carbamate (250 mg, 0.67 mmol) in ethylene glycol (7 mL) was added H₂NNH₂ (110 mg, 3.35 mmol), KOH (1.38 g 10.05 mmol), and heated to 110 °C overnight. The reaction was cooled to room temperature, water was added and the mixture extracted with CH₂Cl₂. The organic layers were washed with water, brine, dried (MgSO₄) and concentrated under reduced pressure to give a residue which is triturated with CH₂Cl₂ to afford the title compound as a off white solid 210 mg (99%) Mp: 186-188 °C, MS (ES⁺): 316 (M+H)⁺


Example 176




N,N-Dimethyl-N-{2-[3-(phenylsulfonyl)-1*H*-indazol-5-yl]ethyl}amine hydrochloride To a solution of *N*-methyl-*N*-{2-[3-(phenylsulfonyl)-1*H*-indazol-5-yl]ethyl}amine (38 mg, 0.12 mmol) in acetonitrile (3 mL) was added NaBH(OAc)₃ (100 mg, 0.48 mmol), formaldehyde (0.027 mL, 0.36 mmol), and stirred at rt. for 3 hr. The reaction mixture was poured into ice water and MeCN was removed under reduced pressure. The resultant suspension was extracted with CH₂Cl₂. The organic extracts were washed with water, brine, dried (MgSO₄) and concentrated to a residue

which was dissolved in MeOH and treated with a slight excess of HCl (1N / Et₂O) to afford the desired product as the mono HCl salt (43 mg, 98%) as a white solid. Mp: 108-110 °C. MS (ES⁺): 330 (M+H)⁺

5

Example 177**N,N-Dimethyl-N-{3-[3-(phenylsulfonyl)-1H-indazol-5-yl]propyl}amine trifluoroacetate****Step 1**


- 10 **5-Bromo-2-nitrobenzyl phenyl sulfone** To a stirred solution of 1-bromo-4-nitrobenzene (5.05 g, 25 mmol) and chloromethylphenylsulfone (4.76 g, 25 mmol) in dry THF (50 mL) at -65 °C under nitrogen is added 1.0M KO^tBu in THF (55 mL, 55 mmol). The deep purple reaction is allowed to warm to 0°C over 1.5 h and then treated with glacial acetic acid (4 mL). The reaction is diluted with water (100 mL)
- 15 and saturated aqueous NaHCO₃ (100 mL), and then extracted with CH₂Cl₂ (2 x 200 mL). The extracts are dried (MgSO₄) and concentrated *in vacuo* to a light orange solid. Trituration with ethyl acetate and hexanes affords the title compound as a pale yellow solid (6.45 g, 72%). Mp: 143 - 144 °C. MS (ES⁻): 354 (M-H)⁺

20 **Step 2**

- Methyl (2E)-3-{4-nitro-3-[(phenylsulfonyl)methyl]phenyl}acrylate** A solution of 5-bromo-2-nitrobenzyl phenyl sulfone (0.356 g, 1 mmol) and methyl acrylate (0.172 g, 2 mmol) in dry DMF (3 mL) was heated at 100°C under nitrogen in the presence of diisopropyl amine (0.21 mL, 1.75 mmol), Pd(OAc)₂ (5 mg, 0.02 mmol), and PPh₃ (10 mg, 0.04 mmol) for 3 days. After being cooled to rt., water was added, and the solution was extracted with ethyl acetate (3 x 15 mL). The organic layers were washed with water then brine and dried (MgSO₄). Evaporation of the solvent under
- 25


reduced pressure gave a residue that was purified by column chromatography (40% ethyl acetate / hexane) to afford the title compound as a yellow solid (0.30 g, 83%). Mp: 165 - 166 °C. MS (ES-): 360 (M-H)⁺

5 **Step 3**




Methyl 3-{4-amino-3-[(phenylsulfonyl)methyl]phenyl}propanoate This compound was prepared by catalytic hydrogenation of methyl (2E)-3-{4-nitro-3-[(phenylsulfonyl)methyl]phenyl}acrylate (3.33 g, 9.22 mmol) in the presence of
 10 Palladium on carbon and hydrogen (45 psi) in ethyl acetate (100 ml) for 4h. The reaction mixture is filtered through celite and concentrated *in vacuo* to give the title compound as a white solid (3.07 g, 99%). Mp: 59 - 60 °C. MS (ES⁺): 334 (M+H)⁺

15 **Step 4**




Methyl 3-{3-[(phenylsulfonyl)-1H-indazol-5-yl]phenyl}propanoate Using substantially the same procedure described in Example 171, step 3, and employing methyl 3-{4-amino-3-[(phenylsulfonyl)methyl]phenyl}propanoate (2.8 g, 8.4 mmol), the title compound was obtained as a red solid (2.2 g, 73%). Mp: 115 - 117 °C. MS (ES⁺):
 20 345 (M+H)⁺

Step 5




3-{3-[(Phenylsulfonyl)-1H-indazol-5-yl]propan-1-ol} A solution of methyl 3-{3-[(phenylsulfonyl)-1H-indazol-5-yl]propanoate (2.0 g, 5.8 mmol) in anhydrous THF (120 ml) under a N₂ atmosphere was cooled to -78 °C, treated with DIBAL-H (1.0 M in
 25

THF; 23 mL), allowed to warm to room temperature, stirred for 12 h, cooled to 0 °C, quenched slowly with a saturated solution of Na₂SO₄ (12 mL) and filtered. The filtercake was washed with ethyl acetate. The combined filtrates were dried over MgSO₄ and concentrated under reduced pressure. Purification of the resultant residue by flash chromatography (60% ethyl acetate/ hexanes) gave the title compound as an off white solid (1.56 g, 83%), mp: 92 - 96 °C. MS (ES⁺): 317 (M+H)⁺

Step 6

3-[3-(Phenylsulfonyl)-1H-indazol-5-yl]propyl 4-methylbenzenesulfonate Using substantially the same procedure described in Example 171, step 6, and employing 3-[3-(phenylsulfonyl)-1H-indazol-5-yl]propan-1-ol (1.52 g, 4.8 mmol), the title compound was obtained as a white foam (1.28 g, 57%). Mp: 58 - 60°C. MS (ES⁺): 471 (M+H)⁺


Step 7

N,N-Dimethyl-N-{3-[3-(phenylsulfonyl)-1H-indazol-5-yl]propyl}amine

trifluoroacetate Using substantially the same procedure described in Example 171, step 7, and employing 3-[3-(phenylsulfonyl)-1H-indazol-5-yl]propyl 4-methylbenzenesulfonate (125 mg, 0.26 mmol), the title product was obtained as an off white solid (83 mg, 93%). Mp: 75 - 80 °C. MS (ES⁺): 344 (M+H)⁺

Example 178


N-{3-[3-(Phenylsulfonyl)-1H-indazol-5-yl]propyl}cyclopropanamine trifluoroacetate



Using essentially the same procedure described in Example 177, step 7 and employing cyclopropyl amine and substituting trifluoroacetic acid for HCl, the title compound was obtained as a white solid, Mp 68 - 169 °C, MS (M+H)⁺ 356

Example 179

N-Isopropyl-N-{3-[3-(phenylsulfonyl)-1H-indazol-5-yl]propyl}amine trifluoroacetate



10


Using essentially the same procedure described in Example 177, step 7 and employing isopropyl amine and substituting trifluoroacetic acid for HCl, the title compound was obtained as a white solid, mp 169 – 171 °C, MS, (M+H)⁺ 358

15

Example 180

{3-[3-(Phenylsulfonyl)-1H-indazol-5-yl]propyl}amine trifluoroacetate


Step 1



5-(3-Azidopropyl)-3-(phenylsulfonyl)-1H-indazole Using essentially the same procedure described in Example 174, step 1, and employing 3-[3-(phenylsulfonyl)-1H-indazol-5-yl]propyl 4-methylbenzenesulfonate (350 mg, 0.75 mmol), the title

compound was obtained as a white solid (220 mg, 86%). Mp: 133 - 135 °C. MS (ES⁻): 340 (M-H)⁺

Step 2




{3-[3-(Phenylsulfonyl)-1H-indazol-5-yl]propyl}amine trifluoroacetate Using essentially the same procedure described in Example 174, Step 2, and employing 5-(3-azidopropyl)-3-(phenylsulfonyl)-1H-indazole (200 mg, 0.59 mmol), the title product was obtained as an off white solid (120 mg, 67%). Mp: 92 - 94 °C. MS (ES⁺): 316 (M+H)⁺

Example 181


{4-[3-(Phenylsulfonyl)-1H-indazol-5-yl]butyl}amine hydrochloride

Step 1




4-{4-Nitro-3-[(phenylsulfonyl)methyl]phenyl}but-3-yn-1-ol A solution of 5-Bromo-2-nitrobenzyl phenyl sulfone (2.5 g, 7.02 mmol), diisopropyl amine (4.92 ml, 35.1 mmol) and homo propargol alcohol (0.59 mL, 7.72 mmol) in degassed toluene (30 mL) was treated simultaneously with CuI (66.8 mg, 0.35 mmol) and tetrakis(triphenylphosphine)-palladium(0) (742 mg, 0.35 mmol), heated to 90 °C, stirred for 15 minutes under a nitrogen atmosphere, cooled to room temperature, diluted with ethyl acetate (10 mL), and filtered through celite. The filtrate was washed with water (25 mL), then brine and dried over MgSO₄ and concentrated *in vacuo*. Purification of the resultant residue by flash chromatography (40% ethyl acetate/petroleum ether) gave the title compound as a white solid (2.15 g, 89%). Mp: 129 °C. MS (ES⁻): 344 (M-H)⁺

Step 2




5 **4-{4-Amino-3-[(phenylsulfonyl)methyl]phenyl}butan-1-ol** Catalytic hydrogenation of 4-{4-nitro-3-[(phenylsulfonyl)methyl]phenyl}but-3-yn-1-ol (1.0 g, 2.85 mmol) in the presence of Palladium on carbon and hydrogen (45 psi) in ethyl acetate (30 ml) for 24h gave a reaction mixture. The reaction mixture was filtered through celite and the filtrate was concentrated *in vacuo* to give the title compound as a yellow solid (0.907 g, 99%). Mp: 70 – 72.5 °C. MS (ES⁺): 320 (M+H)⁺

10 Step 3




15 **4-[3-(Phenylsulfonyl)-1H-indazol-5-yl]butan-1-ol** Using essentially the same described in Example 171, Step 3, and employing 4-{4-amino-3-[(phenylsulfonyl)-methyl]phenyl}butan-1-ol (0.725 g, 2.27 mmol), the title product was obtained as a red solid light pink solid (0.605 g, 81%). Mp: 215 - 216 °C. MS (ES⁺): 331 (M+H)⁺.

Step 4




20 **4-[3-(phenylsulfonyl)-1H-indazol-5-yl]butyl 4-methylbenzenesulfonate** Using substantially the same procedure described in Example 171, Step 6, and employing 4-[3-(phenylsulfonyl)-1H-indazol-5-yl]butan-1-ol (0.5 g, 1.53 mmol), the title compound was obtained as a light orange solid (0.703 g, 95%). Mp: 157 – 159 °C. MS (ES⁺): 485 (M+H)⁺

Step 5



- 5 **5-(4-Azidobutyl)-3-(phenylsulfonyl)-1H-indazole** Using substantially the same procedure described in Example 174, Step 1, and employing 4-[3-(phenylsulfonyl)-1H-indazol-5-yl]butyl 4-methylbenzenesulfonate (133 mg, 0.27 mmol), the title compound was obtained as a clear glass (94 mg, 96%). MS (ES⁺): 356 (M+H)⁺

Step 6




- 10 **{4-[3-(Phenylsulfonyl)-1H-indazol-5-yl]butyl}amine** Using substantially the same procedure described in Example 174, Step 2, and employing 5-(4-azidobutyl)-3-(phenylsulfonyl)-1H-indazole (94 mg, 0.264 mmol), the title compound was obtained as a white solid (65 mg, 86%). Mp: 120 - 125 °C. MS (ES⁺): 330 (M+H)⁺

15

Example 182

N,N-Dimethyl-N-{3-[3-(naphthylsulfonyl)-1H-indazol-5-yl]propyl}amine hydrochloride

Step 1




20

5-Bromo-2-nitrobenzyl naphthyl sulfone Using substantially the same procedure described in Example 177, Step 1, and employing 1-bromo-4-nitrobenzene (5.05 g, 25 mmol) and chloromethylnaphthyl sulfone (7.1 g, 30 mmol), the title product was obtained as a light tan solid (8.6 g, 85%). Mp: 165 - 168 °C. MS (ES⁻): 404 (M-H)⁺


25

Step 2




Methyl (2E)-3-{4-nitro-3-[(naphthylsulfonyl)methyl]phenyl}propanoate Using substantially the same procedure described in Example 177, Step 2, and employing 5-bromo-2-nitrobenzyl naphthyl sulfone (0.540 g, 1.33 mmol), the title product was obtained as a light tan solid (0.438 g, 80%). Mp: 178 °C. MS (ES⁻): 410 (M-H)⁺

Step 3




Methyl 3-{4-amino-3-[(naphthylsulfonyl)methyl]phenyl}propanoate Using substantially the same procedure described in Example 177, Step 3, and employing methyl (2E)-3-(4-nitro-3-[(naphthylsulfonyl)methyl]phenyl)acrylate (2.5 g, 6.0 mmol), the title compound was obtained as a brown gum (2.1 g, 90%). MS (ES⁺): 384 (M+H)⁺

Step 4




Methyl 3-[3-(Naphthylsulfonyl)-1H-indazol-5-yl]propanoate Using substantially the same procedure described in Example 171, Step 3, and employing methyl 3-(4-amino-3-[(naphthylsulfonyl)methyl]phenyl)propanoate (1.2 g, 3.11 mmol), the title compound was obtained as a light brown solid (1.02 g, 83%). Mp: 60 - 65 °C. MS (ES⁺): 395 (M+H)⁺

Step 5




- 5 **3-[3-(Naphthylsulfonyl)-1H-indazol-5-yl]propan-1-ol** Using substantially the same procedure described in Example 177, Step 5, and employing methyl 3-[3-(naphthylsulfonyl)-1H-indazol-5-yl]propanoate (1.0 g, 2.5 mmol), the title compound was obtained as a white solid (0.83 g, 89%). Mp: 170 °C. MS (ES⁺): 367 (M+H)⁺

Step 6




- 10 **3-[3-(Naphthylsulfonyl)-1H-indazol-5-yl]propyl 4-methylbenzenesulfonate** Using substantially the same procedure described in Example 171, Step 6, and employing 3-[3-(naphthylsulfonyl)-1H-indazol-5-yl]propan-1-ol (0.763 g, 2.08 mmol), the title product was obtained as a white foam (0.8 g, 74%). Mp: 66 – 68 °C. MS (ES⁺): 521 (M+H)⁺

15 Step 7



- 20 **N,N-Dimethyl-N-{3-[3-(naphthylsulfonyl)-1H-indazol-5-yl]propyl}amine hydrochloride** Using substantially the same procedure described in Example 171, Step 7, and employing 3-[3-(naphthylsulfonyl)-1H-indazol-5-yl]propyl 4-methylbenzenesulfonate (100 mg, 0.19 mmol), the title compound was obtained as an off white solid (73 mg, 99%). Mp: 247 – 248 °C. MS (ES⁺): 394 (M+H)⁺


Examples 183-187



Using essentially the same procedure described in Example 171, Step 7, and employing 3-[3-(naphthylsulfonyl)-1H-indazol-5-yl]propyl 4-methylbenzenesulfonate and the desired amine, the compounds shown in Table X were obtained and

5 identified by HPLC and mass spectral analyses.

TABLE X




Ex. No.	R5	R6	mp °C	[M+H] ⁺
183	CH ₃	H	155-160	380
184	C ₂ H ₅	H	228-232	393
185	isopropyl	H	160-163	408
186	C ₂ H ₅	CH ₃	205-207	408
187	-CH ₂ CH ₂ CH ₂ CH ₂ -		135-138	420

10

Example 188

{3-[3-(Naphthylsulfonyl)-1H-indazol-5-yl]propyl}amine

Step 1




5-(3-Azidopropyl)-3-(naphthylsulfonyl)-1H-indazole Using essentially the same

15 procedure described in Example 174, Step 1, and employing 3-[3-(naphthylsulfonyl)-

1H-indazol-5-yl]propyl 4-methylbenzenesulfonate (135 mg, 0.26 mmol), the title compound was obtained as a clear glass (97 mg, 96%). MS (ES⁻): 390 (M-H)⁺

Step 2




{3-[3-(Naphthylsulfonyl)-1H-indazol-5-yl]propyl}amine Using essentially the same procedure described in Example 174, Step 2, and employing 5-(3-azidopropyl)-3-(naphthylsulfonyl)-1H-indazole (96 mg, 0.25 mmol), the title compound was obtained as an off white solid (90 mg, 99%). Mp: 92 - 94 °C. MS (ES⁺): 366 (M+H)⁺

10

Example 189


{4-[3-(Naphthylsulfonyl)-1H-indazol-5-yl]butyl}amine hydrochloride

Step 1



4-{4-Nitro-3-[(naphthylsulfonyl)methyl]phenyl}but-3-yn-1-ol Using substantially the same procedure described in Example 181, Step 1, and employing 5-bromo-2-nitrobenzyl naphthyl sulfone (5000 mg, 12.3 mmol), the title compound was obtained as an off yellow solid (4381 mg, 90%). Mp: 155 °C. MS (ES⁻): 394 (M-H)⁺


Step 2



4-{4-Amino-3-[(naphthylsulfonyl)methyl]phenyl}butan-1-ol Using substantially the same procedure described in Example 181, Step 2, and employing 4-{4-nitro-3-[(naphthylsulfonyl)methyl]phenyl}but-3-yn-1-ol (1750 mg, 4.4 mmol), the title

compound was obtained as a yellow solid (1616 mg, 99%). Mp: 80 - 85 °C. MS (ES⁺): 370 (M+H)⁺


Step 3



4-[3-(Naphthylsulfonyl)-1H-indazol-5-yl]butan-1-ol Using substantially the same procedure described in Example 171, Step 3, and employing 4-[4-amino-3-[(naphthylsulfonyl)methyl]phenyl]butan-1-ol (1.6 g, 4.3 mmol), the title compound was obtained as a light pink solid (1.64 g, 99%). Mp: 138 - 139 °C. MS (ES⁻): 379 (M-H)⁺

10


Step 4



4-[3-(Naphthylsulfonyl)-1H-indazol-5-yl]butyl 4-methylbenzenesulfonate Using substantially the same procedure described in Example 171, Step 6, and employing 4-[3-(naphthylsulfonyl)-1H-indazol-5-yl]butan-1-ol (1.6 g, 4.2 mmol), the title compound was obtained as a white foam (2.02 g, 90%). Mp: 63 - 66 °C. MS (ES⁺): 535 (M+H)⁺

15


Step 5



5-(4-Azidobutyl)-3-(naphthylsulfonyl)-1H-indazole Using substantially the same procedure described in Example 174, Step 1, and employing 4-[3-(naphthylsulfonyl)-


1H-indazol-5-yl]butyl 4-methylbenzenesulfonate (150 mg, 0.28 mmol), the title compound was obtained as a clear glass (104 mg, 92%). MS (ES⁺): 406 (M+H)⁺

Step 6




10 **{4-[3-(Naphthylsulfonyl)-1H-indazol-5-yl]butyl}amine hydrochloride** Using substantially the same procedure described in Example 17 Step 2, and employing 5-(4-azidobutyl)-3-(naphthylsulfonyl)-1H-indazole (104 mg, 0.25 mmol), the title compound was obtained as a white solid (91mg, 94%). Mp: 150 - 152 °C. MS (ES⁺): 380 (M+H)⁺

Examples 190-194




15 Using essentially the same procedure described in Example 171, Step 7, and employing 3-[3-(naphthylsulfonyl)-1H-indazol-5-yl]butyl 4-methylbenzenesulfonate and the desired amine, the compounds shown in Table XI were obtained and identified by HPLC and mass spectral analyses.

TABLE XI

Ex. No.	R5	R6	mp °C	[M+H] ⁺
190	CH ₃	CH ₃	232-233	408
191	C ₂ H ₅	H	140-145	408
192	isopropyl	H	145-150	422
193	C ₂ H ₅	CH ₃	195-198	422
194	-CH ₂ CH ₂ CH ₂ CH ₂ -		238-240	434

5

Example 195**3-(1-Naphthylsulfonyl)-5-(piperazin-1-ylmethyl)-1H-indazole****Step 1:**

2-[3-(Naphthalene-1-sulfonylmethyl)-4-nitro-phenyl]-[1,3] dioxolane A mixture of 2-(4-nitro-phenyl)-[1, 3] dioxolane (1.85 g, 9.5 mmoles) and 1-chloromethane-sulfonyl-naphthalene (2.74 g, 11.4 mmoles) was stirred in THF (50 ml) at -78 °C, in a round bottom flask under nitrogen. A solution of 1M potassium t-butoxide was added dropwise (19 ml, 19 mmoles) over a half an hour period. Temperature was allowed to rise to -40 °C, and the reaction mixture was stirred at this temperature for 5 hours. The reaction mixture was poured into cold 2N HCl, extracted with EtOAc,

dried over Na₂SO₄, and concentrated under vacuum. Compound was purified by normal phase HPLC on silica column, using as eluent 40% EtOAc/hexane, to afford the title compound as an off-white solid (3.03 g, 7.6 mmol).

Step 2:

5 **3-(Naphthalene-1-sulfonylmethyl)-4-nitro-benzaldehyde** A mixture of 2-[3-(naphthalene-1-sulfonylmethyl)-4-nitro-phenyl]-[1,3] dioxolane (3.03 g, 7.6 mmol), and 2N HCl (4 ml, 8 mmol) in THF (30 mL) was stirred at 40 °C for 4 hours. The reaction mixture was cooled to room temperature, diluted with water, extracted with CH₂Cl₂, dried over Na₂SO₄, and concentrated under vacuum to yield the title
10 compound (2.56 g, 7.22 mmol).

Step 3:

4-Amino-3-(naphthalene-1-sulfonylmethyl)-benzaldehyde A mixture of 3-(naphthalene-1-sulfonylmethyl)-4-nitro-benzaldehyde (2.5 g, 7.22 mmol) and 10%Pd/C in THF (10 mL), and methanol (20 mL) was hydrogenated in a Parr
15 hydrogenation bottle (250 mL) at 52lb/in² overnight. The mixture was filtered through Celite, and the filtrate was concentrated under vacuum to afford the title compound as an off-white solid (2.4 g, 6.85 mmol).

Step 4:


3-(Naphthalene-1-sulfonyl)-1H-indazole-5-carbaldehyde A mixture of 4-amino-3-(naphthalene-1-sulfonylmethyl)-benzaldehyde (2.4 g, 6.85 mmol) in THF (10mL)
20 and 4M HCl (20 mL) was stirred in a round bottom flask at 3 °C. A solution of sodium nitrite (0.49 g, 7.19 mmol) in H₂O (2 mL) was added. The reaction mixture was poured into a cold solution of saturated sodium bicarbonate (100 mL) and extracted with EtOAc. Compound was dried over Na₂SO₄, and concentrated under vacuum to
25 afford the title compound as an off white solid (1.84 g, 5.5 mmol).

Step 5:

3-(1-Naphthylsulfonyl)-5-(piperazin-1-ylmethyl)-1H-indazole A mixture of 3-(naphthalene-1-sulfonyl)-1H-indazole-5-carbaldehyde (0.17 g, 0.5 mmol), piperazine (0.2 mL, 2.0 mmol) and sodium triacetoxyborohydride (0.15 g, 0.7 mmol) in
30 dichloroethane (5 mL) was stirred at room temperature for 24 hrs. After completion, the solvent was removed *in vacuo*, crude material dispersed in water and pH brought to 3.4. Solid material was filtered off and washed with cold water to afford after drying the target material as a free base. The latter was converted into hydrochloride


salt by dissolution in methanol, followed by treatment with the excess of 2N HCl and the evacuation of the volatiles in vacuo to afford the title compound as the hydrochloride salt, mp > 200 °C; MS (ES) *m/z* 406.

5


Examples 196-201

Using essentially the same reductive amination procedure described in Example 195, Step 5, and employing the desired amine and treatment of the free base with HCl; the compounds shown in Table XII were obtained and identified by HPLC and mass spectral analyses.

10

TABLE XII

Ex. No.	NR5R6	mp °C	MS <i>m/z</i>
196	4-methylpiperazin-1-yl	>200	419.1
197	3-methylpiperazin-1-yl	>200	421.1
198	3,5-dimethylpiperazin-1-yl	>200	433.1
199	(3S)-3-methylpiperazin-1-yl	178-181	419.2
200	(3R)-3-methylpiperazin-1-yl	179-181	419.2
201	(3R)-3-aminopyrrolidin-1-yl	>200	405.1

Example 202**3-(1-Naphthylsulfonyl)-5-(piperazin-1-ylcarbonyl)-1H-indazole****Step 1: 3-(Naphthalene-1-sulfonyl)-1H-indazole-5-carboxylic-acid**


5 A mixture of 3-(naphthalene-1-sulfonyl)-1H-indazole-5-carbaldehyde (0.15 g, 0.44 mmoles) and KMnO_4 (0.03 g, 0.29 moles) was stirred in $\text{CH}_3\text{CN}/\text{H}_2\text{O}$ (4:1) for 1 hour. Reaction mixture was acidified with 2N HCl, stirred with saturated sodium bisulfite for 10 minutes, then extracted with EtOAc, dried over Na_2SO_4 and concentrated *in vacuo* to afford the title compound (0.13 g, 3.9 mmoles).

10

Step 2: 3-(1-Naphthylsulfonyl)-5-(piperazin-1-ylcarbonyl)-1H-indazole

A mixture of 3-(naphthalene-1-sulfonyl)-1H-indazole-5-carboxylic-acid (0.13 g, 0.39 mmoles) piperazine (0.036 g, 0.42 mmoles), and 1-[3-(Dimethylamino) propyl-3-ethylcarbodiimide hydrochloride (0.08 g, 0.42mmoles) was stirred in CH_2Cl_2 for ½ hour. Reaction mixture was diluted with H_2O , extracted with EtOAc, washed with water (2x), brine (1x), dried over Na_2SO_4 , and concentrated under vacuum. The product was converted into the hydrochloride salt by dissolution in methanol, followed by treatment with the excess of 2N HCl and the evacuation of the volatiles *in vacuo* to afford the title compound hydrochloride salt; Mp: > 200 °C; MS (ES) m/z 419.1.

20

Example 203**5-[1-(4-Methylpiperazin-1-yl) ethyl]-3-(1-naphthylsulfonyl)-1H-indazole****Step 1: 1-[3-(Naphthalene-1-sulfonyl)-1H-indazol-5-yl]-ethanol**

A mixture of 3-(naphthalene-1-sulfonyl)-1*H*-indazole-5-carbaldehyde (0.15 g, 0.44 mmol) and MeMgBr (0.36 ml of 3M solution in ether, 1.1 mmol) was stirred in THF at -20 °C to 0 °C for 30 minutes. Reaction mixture was diluted with water, acidified to pH=3 with 2N HCl, then extracted with EtOAc, dried over Na₂SO₄ and concentrated in vacuo to afford the title compound (0.13 g, 3.9 mmol).

Step 2: 5-[1-(4-methylpiperazin-1-yl) ethyl]-3-(1-naphthylsulfonyl)-1*H*-indazole:

A mixture of 1-[3-(naphthalene-1-sulfonyl)-1*H*-indazol-5-yl]ethanol (0.13 g, 0.38 mmol), methane sulfonic anhydride (0.16 g, 0.95 mmol), and triethylamine (0.13 ml, 0.95 mmol) was stirred in CH₂Cl₂ from 0 °C to room temperature for 5 hours to afford methanesulfonic acid 1-[3-(naphthalene-1-sulfonyl)-1*H*-indazol-5-yl]-ethylester. This reaction mixture was treated with excess *N*-methyl-piperazine (0.22 g, 2 mmol). The crude product was purified by flash chromatography using 5% MeOH/CH₂Cl₂ to afford the title compound; Mp: 183-185 °C, MS (ES) *m/z* 433.1.

Example 204

Evaluation of 5-HT₆ Binding Affinity of Test Compounds

The affinity of test compounds for the serotonin 5-HT₆ receptor was evaluated in the following manner. Cultured Hela cells expressing human cloned 5-HT₆ receptors were harvested and centrifuged at low speed (1,000 x g) for 10.0 minutes to remove the culture media. The harvested cells were suspended in half volume of fresh physiological phosphate buffered saline solution and recentrifuged at the same speed. This operation was repeated. The collected cells were then homogenized in ten volumes of 50 mM Tris.HCl (pH 7.4) and 0.5 mM EDTA. The homogenate was centrifuged at 40,000 x g for 30.0 min and the precipitate was collected. The obtained pellet was resuspended in 10 volumes of Tris.HCl buffer and recentrifuged at the same speed. The final pellet was suspended in a small volume of Tris.HCl buffer and the tissue protein content was determined in aliquots of 10-25 µl volumes. Bovine Serum Albumin was used as the standard in the protein determination according to the method described in Lowry et al., J. Biol. Chem., 193: 265 (1951). The volume of the suspended cell membranes was adjusted to give a tissue protein concentration of 1.0 mg/ml of suspension. The prepared membrane suspension (10

times concentrated) was aliquoted in 1.0 ml volumes and stored at -70° C until used in subsequent binding experiments.

Binding experiments were performed in a 96 well microtiter plate format, in a total volume of 200 µl. To each well was added the following mixture: 80.0 µl of incubation buffer made in 50 mM Tris.HCl buffer (pH 7.4) containing 10.0 mM MgCl₂ and 0.5 mM EDTA and 20 µl of [³H]-LSD (S.A., 86.0 Ci/mmol, available from Amersham Life Science), 3.0 nM. The dissociation constant, K_D of the [³H]LSD at the human serotonin 5-HT₆ receptor was 2.9 nM, as determined by saturation binding with increasing concentrations of [³H]LSD. The reaction was initiated by the final addition of 100.0 µl of tissue suspension. Nonspecific binding was measured in the presence of 10.0 µM methiothepin. The test compounds were added in 20.0 µl volume.

The reaction was allowed to proceed in the dark for 120 minutes at room temperature, at which time, the bound ligand-receptor complex was filtered off on a 96 well unfilter with a Packard Filtermate® 196 Harvester. The bound complex caught on the filter disk was allowed to air dry and the radioactivity is measured in a Packard TopCount® equipped with six photomultiplier detectors, after the addition of 40.0µl Microscint®-20 scintillant to each shallow well. The unfilter plate was heat-sealed and counted in a PackardTopCount® with a tritium efficiency of 31.0%.

Specific binding to the 5-HT₆ receptor was defined as the total radioactivity bound less the amount bound in the presence of 10.0µM unlabeled methiothepin. Binding in the presence of varying concentrations of test compound was expressed as a percentage of specific binding in the absence of test compound. The results were plotted as log % bound versus log concentration of test compound. Nonlinear regression analysis of data points with a computer assisted program Prism® yielded both the IC₅₀ and the K_i values of test compounds with 95% confidence limits. A linear regression line of data points was plotted, from which the IC₅₀ value is determined and the K_i value is determined based upon the following equation:

$$K_i = IC_{50} / (1 + L/K_D)$$

where L was the concentration of the radioactive ligand used and K_D is the dissociation constant of the ligand for the receptor, both expressed in nM.

Using this assay, the following Ki values were determined. The data are shown in Table XIII, below.

For Table XIII

A = 0.01nM-10nM

B = 11nM-25nM

C = 26nM-35nM

D = 36nM-45nM

E = >45nM

5

TABLE XIII

<u>Test Compound (Example No.)</u>	<u>5-HT₆ Binding Ki (nM)</u>
1	B
2	E
3	C
4	A
5	D
6	B
7	C
8	C
9	B
10	E
11	A
12	B
13	D
14	A
15	A
16	A
17	A
18	A
19	A
20	A
21	A
22	A
23	A

Test Compound (Example No.)	5-HT₆ Binding K_i (nM)
24	A
25	A
26	A
27	A
28	A
29	A
30	A
31	A
32	A
33	A
34	A
35	A
36	A
37	A
38	A
39	B
40	A
41	A
42	A
43	A
44	A
45	A
46	A
47	C
48	C
49	C
50	D
51	B
52	E
53	B
54	B

<u>Test Compound (Example No.)</u>	<u>5-HT₆ Binding K_i (nM)</u>
55	E
56	E
57	D
58	C
59	E
60	B
61	A
62	A
63	A
64	A
65	A
66	A
67	A
68	A
69	C
70	B
71	B
72	D
73	E
74	A
75	B
76	A
77	A
78	A
79	B
80	A
81	A
82	B
83	C
84	E
85	E

<u>Test Compound (Example No.)</u>	<u>5-HT₆ Binding K_i (nM)</u>
86	C
87	E
88	A
89	A
90	A
91	A
92	A
93	A
94	B
95	A
96	E
97	A
98	C
99	A
100	B
101	B
102	C
103	E
104	C
105	E
106	E
107	E
108	E
109	B
110	E
111	E
112	E
113	A
114	A
115	B
116	B


<u>Test Compound (Example No.)</u>	<u>5-HT₆ Binding Ki (nM)</u>
117	A
118	A
119	E
120	E
121	E
122	A
123	E
124	B
125	B
126	A
127	B
128	E
129	E
130	E
131	B
132	C
133	B
134	B
135	D
136	D
137	C
138	D
139	A
140	A
141	B
142	--
143	A
144	A
145	B
146	B
147	B

Test Compound (Example No.)	5-HT₆ Binding K_i (nM)
148	E
149	E
150	E
151	A
152	E
153	B
154	B
155	B
156	A
157	A
158	A
159	B
160	A
161	B
162	A
163	B
164	A
165	B
166	C
167	B
168	D
169	E
170	B
171	A
172	B
173	A
174	A
175	A
176	B
177	B
178	B

Test Compound (Example No.)	5-HT₆ Binding K_i (nM)
179	E
180	B
181	C
182	A
183	A
184	A
185	A
186	A
187	A
188	A
189	A
190	A
191	A
192	A
193	A
194	B
195	A
196	A
197	A
198	A
199	B
200	A
201	A
202	E
203	A

Claims:

1. A compound of formula I



wherein

X is O or S;

n is an integer of 2, 3, 4, 5 or 6;

R₁ is H or an alkyl, cycloalkyl, aryl or heteroaryl group each optionally substituted;

R₂ is an optionally substituted cycloalkyl or aryl group;

R₃ and R₄ are each independently H, or an optionally substituted alkyl group;

R₅ and R₆ are each independently H, or an alkyl, alkenyl, alkynyl, or cycloalkyl, each optionally substituted;

R₇ is H, halogen, CN, OR₈, CO₂R₉, CONR₁₀R₁₁, or an alkyl, alkenyl, alkynyl, cycloalkyl, cycloheteroalkyl, aryl or heteroaryl group each optionally substituted;

m is an integer of 1, 2 or 3;

R₈ is H, COR₁₂ or an alkyl, alkenyl, alkynyl, aryl or heteroaryl group each optionally substituted;

R₉ is H or a C₁-C₆alkyl, aryl or heteroaryl group each optionally substituted;

R₁₀ and R₁₁ are each independently H or an optionally substituted alkyl group; and

R₁₂ is an optionally substituted C₁-C₆alkyl, cycloalkyl, cycloheteroalkyl, aryl or heteroaryl group; or

a stereoisomer thereof or a pharmaceutically acceptable salt thereof.


2. The compound according to claim 1 wherein X is O.

3. The compound according to claim 1 wherein n is 2 or 3.
4. The compound according to any one of claims 1 to 3 wherein R₂ is an optionally substituted aryl group.
5. The compound according to claim 3 wherein X is O.
6. The compound according to any one of claims 1 to 5 wherein R₅ and R₆ are each independently H or C₁-C₄ alkyl.
7. The compound according to any one of claims 1 to 6 wherein R₂ is an optionally substituted naphthyl group.
8. The compound according to any one of claims 1 to 7 wherein R₂ is naphthyl and n is 3.
9. The compound according to claim 1 selected from the group consisting of:
 N, N-Dimethyl-3-{{[3-(1-naphthylsulfonyl)-1H-indazol-5-yl]oxy}propan-1-amine;
 N,N-Dimethyl-2-{{[3-(phenylsulfonyl)-1H-indazol-5-yl]oxy}ethanamine;
 N,N-Dimethyl-2-{{[3-(1-naphthylsulfonyl)-1H-indazol-5-yl]oxy}ethanamine;
 N-(2-{{[3-(1-Naphthylsulfonyl)-1H-indazol-5-yl]oxy}ethyl)cyclopentanamine;
 N-Ethyl-N-methyl-2-{{[3-(1-naphthylsulfonyl)-1H-indazol-5-yl]oxy}ethanamine;
 N-(2-{{[3-(1-Naphthylsulfonyl)-1H-indazol-5-yl]oxy}ethyl)butan-1-amine;
 N-Ethyl-2-{{[3-(phenylsulfonyl)-1H-indazol-5-yl]oxy}ethanamine;
 N-(2-{{[3-(Phenylsulfonyl)-1H-indazol-5-yl]oxy}ethyl)propan-2-amine;
 N-(2-{{[3-(1-Naphthylsulfonyl)-1H-indazol-5-yl]oxy}ethyl)propan-2-amine;
 N-Ethyl-2-{{[3-(1-naphthylsulfonyl)-1H-indazol-5-yl]oxy}ethanamine;
 N-Methyl-2-{{[3-(1-naphthylsulfonyl)-1H-indazol-5-yl]oxy}ethanamine;
 N,N-Diethyl-2-{{[3-(1-naphthylsulfonyl)-1H-indazol-5-yl]oxy}ethanamine;
 N-(2-{{[3-(1-Naphthylsulfonyl)-1H-indazol-5-yl]oxy}ethyl)cyclopropanamine;
 (2-{{[1-(3-Chlorobenzyl)-3-(1-naphthylsulfonyl)-1H-indazol-7-yl]oxy}ethyl)amine;
 N-(2-{{[3-(Phenylsulfonyl)-1H-indazol-5-yl]oxy}ethyl)cyclopentanamine;

N-Methyl-2-[[3-(phenylsulfonyl)-1H-indazol-5-yl]oxy]ethanamine;
 N-Methyl-2-[[1-methyl-3-(phenylsulfonyl)-1H-indazol-7-yl]oxy]ethanamine;
 (2-[[1-(3-Chlorobenzyl)-3-(phenylsulfonyl)-1H-indazol-7-yl]oxy]ethyl)methylamine;
 (2-[[1-(3-Chlorobenzyl)-3-(phenylsulfonyl)-1H-indazol-7-yl]oxy]ethyl)ethylamine;
 N-Methyl-2-[[3-(phenylsulfonyl)-1H-indazol-7-yl]oxy]ethanamine;
 (2-[[1-(3-Chlorobenzyl)-3-(1-naphthylsulfonyl)-1H-indazol-7-yl]oxy]ethyl)ethylamine;
 (2-[[1-(3-Chlorobenzyl)-3-(1-naphthylsulfonyl)-1H-indazol-7-yl]oxy]ethyl)methylamine;
 N-Ethyl-2-[[3-(phenylsulfonyl)-1H-indazol-7-yl]oxy]ethanamine;
 N,N-Diethyl-2-[[3-(phenylsulfonyl)-1H-indazol-7-yl]oxy]ethanamine;
 N-(2-[[3-(Phenylsulfonyl)-1H-indazol-7-yl]oxy]ethyl)butan-1-amine;
 N,N-Diethyl-2-[[1-methyl-3-(1-naphthylsulfonyl)-1H-indazol-7-yl]oxy]ethanamine;
 N-Ethyl-2-[[1-methyl-3-(1-naphthylsulfonyl)-1H-indazol-7-yl]oxy]ethanamine;
 N-Ethyl-2-[[3-(1-naphthylsulfonyl)-1H-indazol-7-yl]oxy]ethanamine;
 (2-[[3-(1-Naphthylsulfonyl)-1H-indazol-7-yl]oxy]ethyl)amine;
 (2-[[1-(3-Chlorobenzyl)-5-fluoro-3-(1-naphthylsulfonyl)-1H-indazol-7-yl]oxy]ethyl)-
 dimethylamine;
 (2-[[1-Benzyl-3-(1-naphthylsulfonyl)-1H-indazol-5-yl]oxy]ethyl)methylamine;
 (2-[[1-Benzyl-3-(1-naphthylsulfonyl)-1H-indazol-5-yl]oxy]ethyl)dimethylamine;
 (2-[[1-Benzyl-3-(1-naphthylsulfonyl)-1H-indazol-5-yl]oxy]ethyl)ethylamine;
 N-Methyl-2-[[3-(1-naphthylsulfonyl)-1H-indazol-7-yl]oxy]ethanamine;
 N,N-Dimethyl-2-[[3-(1-naphthylsulfonyl)-1H-indazol-7-yl]oxy]ethanamine;
 N,N-Diethyl-2-[[5-fluoro-3-(1-naphthylsulfonyl)-1H-indazol-7-yl]oxy]ethanamine;
 (2-[[5-Fluoro-3-(1-naphthylsulfonyl)-1H-indazol-7-yl]oxy]ethyl)dimethylamine;
 N-Ethyl-2-[[5-fluoro-3-(1-naphthylsulfonyl)-1H-indazol-7-yl]oxy]ethanamine;
 (2-[[1-(3-Chlorobenzyl)-3-(1-naphthylsulfonyl)-1H-indazol-5-yl]oxy]ethyl)-
 dimethylamine;
 N-Methyl-3-[[3-(1-naphthylsulfonyl)-1H-indazol-5-yl]oxy]propan-1-amine;
 N-Ethyl-N-methyl-3-[[3-(1-naphthylsulfonyl)-1H-indazol-5-yl]oxy]propan-1-amine;
 N,N-Diethyl-3-[[3-(1-naphthylsulfonyl)-1H-indazol-5-yl]oxy]propan-1-amine;
 N-(3-[[3-(1-Naphthylsulfonyl)-1H-indazol-5-yl]oxy]propyl)butan-1-amine;
 (2-[[5-Methoxy-3-(1-naphthylsulfonyl)-1H-indazol-7-yl]oxy]ethyl)methylamine;
 (2-[[5-Methoxy-3-(1-naphthylsulfonyl)-1H-indazol-7-yl]oxy]ethyl)dimethylamine;
 (2-[[1-(3-Chlorobenzyl)-3-(1-naphthylsulfonyl)-1H-indazol-5-yl]oxy]ethyl)ethylamine;

(3-[[1-(3-Chlorobenzyl)-3-(1-naphthylsulfonyl)-1H-indazol-7-yl]oxy}propyl)-
 diethylamine;
 N-Methyl-3-[[3-(1-naphthylsulfonyl)-1H-indazol-7-yl]oxy}propan-1-amine;
 N,N-Diethyl-3-[[3-(1-naphthylsulfonyl)-1H-indazol-7-yl]oxy}propan-1-amine;
 N-Methyl-2-[[1-methyl-3-(1-naphthylsulfonyl)-1H-indazol-5-yl]oxy}ethanamine;
 N,N-Dimethyl-2-[[1-methyl-3-(1-naphthylsulfonyl)-1H-indazol-5-yl]oxy}ethanamine;
 N-Ethyl-N-methyl-2-[[1-methyl-3-(1-naphthylsulfonyl)-1H-indazol-5-yl]oxy}ethan-
 amine;
 N-Ethyl-2-[[1-methyl-3-(1-naphthylsulfonyl)-1H-indazol-5-yl]oxy}ethanamine;
 N,N-Diethyl-2-[[1-methyl-3-(1-naphthylsulfonyl)-1H-indazol-5-yl]oxy}ethanamine;
 N-(2-[[1-Methyl-3-(1-naphthylsulfonyl)-1H-indazol-5-yl]oxy}ethyl)propan-2-amine;
 (2-[[1-Methyl-3-(1-naphthylsulfonyl)-1H-indazol-7-yl]oxy}ethyl)amine;
 N-Ethyl-3-[[3-(1-naphthylsulfonyl)-1H-indazol-5-yl]oxy}propan-1-amine;
 N-Isopropyl-3-[[3-(1-naphthylsulfonyl)-1H-indazol-5-yl]oxy}propan-1-amine;
 N-(3-[[3-(1-Naphthylsulfonyl)-1H-indazol-5-yl]oxy}propyl)cyclopentanamine;
 N-(3-[[3-(1-Naphthylsulfonyl)-1H-indazol-5-yl]oxy}propyl)cyclopropanamine;
 (3-[[3-(1-Naphthylsulfonyl)-1H-indazol-5-yl]oxy}propyl)amine;
 N-Methyl-4-[[3-(1-naphthylsulfonyl)-1H-indazol-5-yl]oxy}butan-1-amine
 N,N-Dimethyl-4-[[3-(1-naphthylsulfonyl)-1H-indazol-5-yl]oxy}butan-1-amine;
 N-Ethyl-4-[[3-(1-naphthylsulfonyl)-1H-indazol-5-yl]oxy}butan-1-amine;
 N,N-Diethyl-4-[[3-(1-naphthylsulfonyl)-1H-indazol-5-yl]oxy}butan-1-amine;
 N-Methyl-4-[[3-(1-naphthylsulfonyl)-1H-indazol-5-yl]oxy}-N-propylbutan-1-amine;
 (4-[[3-(1-Naphthylsulfonyl)-1H-indazol-5-yl]oxy}butyl)amine;
 (2-[[5-Fluoro-3-(1-naphthylsulfonyl)-1H-indazol-7-yl]oxy}ethyl)methylamine;
 (2-[[3-(1-Naphthylsulfonyl)-1H-indazol-5-yl]oxy}ethyl)amine;
 2-[[5-Fluoro-3-(phenylsulfonyl)-1H-indazol-7-yl]oxy}ethanamine;
 2-[[3-(Phenylsulfonyl)-1H-indazol-7-yl]oxy}ethanamine;
 N,N-Dimethyl-2-[[3-(1-naphthylsulfonyl)-1H-indazol-4-yl]oxy}ethanamine;
 2-[[3-(1-Naphthylsulfonyl)-1H-indazol-4-yl]oxy}ethanamine;
 N-Methyl-2-[[3-(1-naphthylsulfonyl)-1H-indazol-6-yl]oxy}ethanamine;
 2-[[3-(1-Naphthylsulfonyl)-1H-indazol-6-yl]oxy}ethanamine;
 a stereoisomer thereof; and
 a pharmaceutically acceptable salt thereof.

10. A pharmaceutical composition which comprises a pharmaceutically acceptable carrier and an effective amount of a compound of formula I



(I)

wherein

X is O or S;

n is an integer of 2, 3, 4, 5 or 6;

R_1 is H or an alkyl, cycloalkyl, aryl or heteroaryl group each optionally substituted;

R_2 is an optionally substituted cycloalkyl or aryl group;

R_3 and R_4 are each independently H, or an optionally substituted alkyl group;

R_5 and R_6 are each independently H, or an alkyl, alkenyl, alkynyl, or cycloalkyl, each optionally substituted;

R_7 is H, halogen, CN, OR_8 , CO_2R_9 , $CONR_{10}R_{11}$, or an alkyl, alkenyl, alkynyl, cycloalkyl, cycloheteroalkyl, aryl or heteroaryl group each optionally substituted;

m is an integer of 1, 2 or 3;

R_8 is H, COR_{12} or an alkyl, alkenyl, alkynyl, aryl or heteroaryl group each optionally substituted;

R_9 is H or a C_1 - C_6 alkyl, aryl or heteroaryl group each optionally substituted;

R_{10} and R_{11} are each independently H or an optionally substituted alkyl group; and

R_{12} is an optionally substituted C_1 - C_6 alkyl, cycloalkyl, cycloheteroalkyl, aryl or heteroaryl group; or

a stereoisomer thereof or a pharmaceutically acceptable salt thereof.

11. The composition according to claim 10 having a formula I compound wherein X is O.

12. The composition according to claim 10 or claim 11, having a formula I compound wherein n is 2 or 3.

13. The composition according to any one of claims 10 to 12 having a formula I compound wherein X is O; R₂ is an optionally substituted phenyl or naphthyl group; and n is 3.

14. The composition according to claim 10 having a formula I compound selected from the group consisting of:

N, N-Dimethyl-3-{{3-(1-naphthylsulfonyl)-1H-indazol-5-yl}oxy}propan-1-amine;
 N,N-Dimethyl-2-{{3-(phenylsulfonyl)-1H-indazol-5-yl}oxy}ethanamine;
 N,N-Dimethyl-2-{{3-(1-naphthylsulfonyl)-1H-indazol-5-yl}oxy}ethanamine;
 N-(2-{{3-(1-Naphthylsulfonyl)-1H-indazol-5-yl}oxy}ethyl)cyclopentanamine;
 N-Ethyl-N-methyl-2-{{3-(1-naphthylsulfonyl)-1H-indazol-5-yl}oxy}ethanamine;
 N-(2-{{3-(1-Naphthylsulfonyl)-1H-indazol-5-yl}oxy}ethyl)butan-1-amine;
 N-Ethyl-2-{{3-(phenylsulfonyl)-1H-indazol-5-yl}oxy}ethanamine;
 N-(2-{{3-(Phenylsulfonyl)-1H-indazol-5-yl}oxy}ethyl)propan-2-amine;
 N-(2-{{3-(1-Naphthylsulfonyl)-1H-indazol-5-yl}oxy}ethyl)propan-2-amine;
 N-Ethyl-2-{{3-(1-naphthylsulfonyl)-1H-indazol-5-yl}oxy}ethanamine;
 N-Methyl-2-{{3-(1-naphthylsulfonyl)-1H-indazol-5-yl}oxy}ethanamine;
 N,N-Diethyl-2-{{3-(1-naphthylsulfonyl)-1H-indazol-5-yl}oxy}ethanamine;
 N-(2-{{3-(1-Naphthylsulfonyl)-1H-indazol-5-yl}oxy}ethyl)cyclopropanamine;
 (2-{{1-(3-Chlorobenzyl)-3-(1-naphthylsulfonyl)-1H-indazol-7-yl}oxy}ethyl)amine;
 N-(2-{{3-(Phenylsulfonyl)-1H-indazol-5-yl}oxy}ethyl)cyclopentanamine;
 N-Methyl-2-{{3-(phenylsulfonyl)-1H-indazol-5-yl}oxy}ethanamine;
 N-Methyl-2-{{1-methyl-3-(phenylsulfonyl)-1H-indazol-7-yl}oxy}ethanamine;
 (2-{{1-(3-Chlorobenzyl)-3-(phenylsulfonyl)-1H-indazol-7-yl}oxy}ethyl)methylamine;
 (2-{{1-(3-Chlorobenzyl)-3-(phenylsulfonyl)-1H-indazol-7-yl}oxy}ethyl)ethylamine;
 N-Methyl-2-{{3-(phenylsulfonyl)-1H-indazol-7-yl}oxy}ethanamine;
 (2-{{1-(3-Chlorobenzyl)-3-(1-naphthylsulfonyl)-1H-indazol-7-yl}oxy}ethyl)ethylamine;

(2-{{1-(3-Chlorobenzyl)-3-(1-naphthylsulfonyl)-1H-indazol-7-yl}oxy}ethyl)methylamine;
 N-Ethyl-2-{{3-(phenylsulfonyl)-1H-indazol-7-yl}oxy}ethanamine;
 N,N-Diethyl-2-{{3-(phenylsulfonyl)-1H-indazol-7-yl}oxy}ethanamine;
 N-(2-{{3-(Phenylsulfonyl)-1H-indazol-7-yl}oxy}ethyl)butan-1-amine;
 N,N-Diethyl-2-{{1-methyl-3-(1-naphthylsulfonyl)-1H-indazol-7-yl}oxy}ethanamine;
 N-Ethyl-2-{{1-methyl-3-(1-naphthylsulfonyl)-1H-indazol-7-yl}oxy}ethanamine;
 N-Ethyl-2-{{3-(1-naphthylsulfonyl)-1H-indazol-7-yl}oxy}ethanamine;
 (2-{{3-(1-Naphthylsulfonyl)-1H-indazol-7-yl}oxy}ethyl)amine;
 (2-{{1-(3-Chlorobenzyl)-5-fluoro-3-(1-naphthylsulfonyl)-1H-indazol-7-yl}oxy}ethyl)-
 dimethylamine;
 (2-{{1-Benzyl-3-(1-naphthylsulfonyl)-1H-indazol-5-yl}oxy}ethyl)methylamine;
 (2-{{1-Benzyl-3-(1-naphthylsulfonyl)-1H-indazol-5-yl}oxy}ethyl)dimethylamine;
 (2-{{1-Benzyl-3-(1-naphthylsulfonyl)-1H-indazol-5-yl}oxy}ethyl)ethylamine;
 N-Methyl-2-{{3-(1-naphthylsulfonyl)-1H-indazol-7-yl}oxy}ethanamine;
 N,N-Dimethyl-2-{{3-(1-naphthylsulfonyl)-1H-indazol-7-yl}oxy}ethanamine;
 N,N-Diethyl-2-{{5-fluoro-3-(1-naphthylsulfonyl)-1H-indazol-7-yl}oxy}ethanamine;
 (2-{{5-Fluoro-3-(1-naphthylsulfonyl)-1H-indazol-7-yl}oxy}ethyl)dimethylamine;
 N-Ethyl-2-{{5-fluoro-3-(1-naphthylsulfonyl)-1H-indazol-7-yl}oxy}ethanamine;
 (2-{{1-(3-Chlorobenzyl)-3-(1-naphthylsulfonyl)-1H-indazol-5-yl}oxy}ethyl)-
 dimethylamine;
 N-Methyl-3-{{3-(1-naphthylsulfonyl)-1H-indazol-5-yl}oxy}propan-1-amine;
 N-Ethyl-N-methyl-3-{{3-(1-naphthylsulfonyl)-1H-indazol-5-yl}oxy}propan-1-amine;
 N,N-Diethyl-3-{{3-(1-naphthylsulfonyl)-1H-indazol-5-yl}oxy}propan-1-amine;
 N-(3-{{3-(1-Naphthylsulfonyl)-1H-indazol-5-yl}oxy}propyl)butan-1-amine;
 (2-{{5-Methoxy-3-(1-naphthylsulfonyl)-1H-indazol-7-yl}oxy}ethyl)methylamine;
 (2-{{5-Methoxy-3-(1-naphthylsulfonyl)-1H-indazol-7-yl}oxy}ethyl)dimethylamine;
 (2-{{1-(3-Chlorobenzyl)-3-(1-naphthylsulfonyl)-1H-indazol-5-yl}oxy}ethyl)ethylamine;
 (3-{{1-(3-Chlorobenzyl)-3-(1-naphthylsulfonyl)-1H-indazol-7-yl}oxy}propyl)-
 diethylamine;
 N-Methyl-3-{{3-(1-naphthylsulfonyl)-1H-indazol-7-yl}oxy}propan-1-amine;
 N,N-Diethyl-3-{{3-(1-naphthylsulfonyl)-1H-indazol-7-yl}oxy}propan-1-amine;
 N-Methyl-2-{{1-methyl-3-(1-naphthylsulfonyl)-1H-indazol-5-yl}oxy}ethanamine;
 N,N-Dimethyl-2-{{1-methyl-3-(1-naphthylsulfonyl)-1H-indazol-5-yl}oxy}ethanamine;

N-Ethyl-N-methyl-2-[[1-methyl-3-(1-naphthylsulfonyl)-1H-indazol-5-yl]oxy]ethan-amine;
 N-Ethyl-2-[[1-methyl-3-(1-naphthylsulfonyl)-1H-indazol-5-yl]oxy]ethanamine;
 N,N-Diethyl-2-[[1-methyl-3-(1-naphthylsulfonyl)-1H-indazol-5-yl]oxy]ethanamine;
 N-(2-[[1-Methyl-3-(1-naphthylsulfonyl)-1H-indazol-5-yl]oxy]ethyl)propan-2-amine;
 (2-[[1-Methyl-3-(1-naphthylsulfonyl)-1H-indazol-7-yl]oxy]ethyl)amine;
 N-Ethyl-3-[[3-(1-naphthylsulfonyl)-1H-indazol-5-yl]oxy]propan-1-amine;
 N-Isopropyl-3-[[3-(1-naphthylsulfonyl)-1H-indazol-5-yl]oxy]propan-1-amine;
 N-(3-[[3-(1-Naphthylsulfonyl)-1H-indazol-5-yl]oxy]propyl)cyclopentanamine;
 N-(3-[[3-(1-Naphthylsulfonyl)-1H-indazol-5-yl]oxy]propyl)cyclopropanamine;
 (3-[[3-(1-Naphthylsulfonyl)-1H-indazol-5-yl]oxy]propyl)amine;
 N-Methyl-4-[[3-(1-naphthylsulfonyl)-1H-indazol-5-yl]oxy]butan-1-amine
 N,N-Dimethyl-4-[[3-(1-naphthylsulfonyl)-1H-indazol-5-yl]oxy]butan-1-amine;
 N-Ethyl-4-[[3-(1-naphthylsulfonyl)-1H-indazol-5-yl]oxy]butan-1-amine;
 N,N-Diethyl-4-[[3-(1-naphthylsulfonyl)-1H-indazol-5-yl]oxy]butan-1-amine;
 N-Methyl-4-[[3-(1-naphthylsulfonyl)-1H-indazol-5-yl]oxy]-N-propylbutan-1-amine;
 (4-[[3-(1-Naphthylsulfonyl)-1H-indazol-5-yl]oxy]butyl)amine;
 (2-[[5-Fluoro-3-(1-naphthylsulfonyl)-1H-indazol-7-yl]oxy]ethyl)methylamine;
 (2-[[3-(1-Naphthylsulfonyl)-1H-indazol-5-yl]oxy]ethyl)amine;
 2-[[5-Fluoro-3-(phenylsulfonyl)-1H-indazol-7-yl]oxy]ethanamine;
 2-[[3-(Phenylsulfonyl)-1H-indazol-7-yl]oxy]ethanamine;
 N,N-Dimethyl-2-[[3-(1-naphthylsulfonyl)-1H-indazol-4-yl]oxy]ethanamine;
 2-[[3-(1-Naphthylsulfonyl)-1H-indazol-4-yl]oxy]ethanamine;
 N-Methyl-2-[[3-(1-naphthylsulfonyl)-1H-indazol-6-yl]oxy]ethanamine;
 2-[[3-(1-Naphthylsulfonyl)-1H-indazol-6-yl]oxy]ethanamine;
 a stereoisomer thereof; and
 a pharmaceutically acceptable salt thereof.

15. The compound according to claim 1 selected from the group consisting of:

[2-(3-benzenesulfonyl-1H-indazol-5-yloxy)-ethyl]ethyl-methylamine;
 [2-(3-benzenesulfonyl-1H-indazol-5-yloxy)-ethyl]diethylamine;
 [2-(3-benzenesulfonyl-1H-indazol-5-yloxy)-ethyl]isopropylamine;

[2-(3-benzenesulfonyl-1H-indazol-5-yloxy)-ethyl]butylamine;
 [2-(3-benzenesulfonyl-1H-indazol-5-yloxy)-ethyl]cyclopropylamine;
 dimethyl-{3-[3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]propyl}amine;
 methyl-{3-[3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]propyl}amine;
 butyl-{3-[3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]propyl}amine;
 {2-[1-benzyl-3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]ethyl-isopropylamine};
 {2-[1-benzyl-3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-ethyl}diethylamine;
 {2-[1-(3-chloro-benzyl)-3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-ethyl}ethylmethylamine;
 {2-[1-(3-chloro-benzyl)-3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-ethyl}diethylamine;
 {2-[1-(3-chloro-benzyl)-3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-ethyl}isopropylamine;
 {2-[1-(3-chloro-benzyl)-3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]ethyl}methylamine;
 {2-[1-(3-chloro-benzyl)-3-(naphthalene-1-sulfonyl)-1H-indazol-7-yloxy]ethyl}ethylamine;
 {2-[1-(3-chloro-benzyl)-3-(naphthalene-1-sulfonyl)-1H-indazol-7-yloxy]ethyl}diethylamine;
 butyl-{2-[1-(3-chloro-benzyl)-3-(naphthalene-1-sulfonyl)-1H-indazol-7-yloxy]ethyl}methylamine;
 {2-[3-benzenesulfonyl-1-(3-chloro-benzyl)-1H-indazol-7-yloxy]-ethyl}diethylamine;
 {2-[3-benzenesulfonyl-1-(3-chloro-benzyl)-1H-indazol-7-yloxy]-ethyl}butylamine;
 dimethyl-{2-[1-methyl-3-(naphthalene-1-sulfonyl)-1H-indazol-7-yloxy]-ethyl}amine;
 butyl-{2-[1-methyl-3-(naphthalene-1-sulfonyl)-1H-indazol-7-yloxy]-ethyl}amine;
 [2-(3-benzenesulfonyl-1-methyl-1H-indazol-7-yloxy)-ethyl]ethylamine;
 [2-(3-benzenesulfonyl-1-methyl-1H-indazol-7-yloxy)-ethyl]diethylamine;
 (2-{[1-(3-chlorobenzyl)-5-fluoro-3-(1-naphthylsulfonyl)-1H-indazol-7-yl]oxy}ethyl)amine;
 (2-{[1-(3-chlorobenzyl)-5-fluoro-3-(1-naphthylsulfonyl)-1H-indazol-7-yl]oxy}ethyl)methylamine hydrochloride;
 {2-[1-(3-chloro-benzyl)-5-fluoro-3-(naphthalene-1-sulfonyl)-1H-indazol-7-yloxy]ethyl}ethylamine hydrochloride;

{2-[1-(3-chloro-benzyl)-5-fluoro-3-(naphthalene-1-sulfonyl)-1H-indazol-7-yloxy]ethyl}diethylamine hydrochloride;
 {2-[5-fluoro-3-(naphthalene-1-sulfonyl)-1H-indazol-7-yloxy]-ethyl}methylamine hydrochloride;
 {2-[1-(3-chlorobenzyl)-5-methoxy-3-(naphthalene-1-sulfonyl)-1H-indazol-7-yloxy]ethyl}methylamine hydrochloride;
 {2-[1-(3-chlorobenzyl)-5-methoxy-3-(naphthalene-1-sulfonyl)-1H-indazol-7-yloxy]ethyl}ethylamine hydrochloride;
 {2-[1-(3-chlorobenzyl)-5-methoxy-3-(naphthalene-1-sulfonyl)-1H-indazol-7-yloxy]ethyl}dimethylamine hydrochloride;
 {3-[1-(3-chlorobenzyl)-5-methoxy-3-(naphthalene-1-sulfonyl)-1H-indazol-7-yloxy]propyl}methylamine hydrochloride;
 {3-[1-(3-chlorobenzyl)-5-methoxy-3-(naphthalene-1-sulfonyl)-1H-indazol-7-yloxy]propyl}ethylamine hydrochloride;
 {3-[1-(3-chlorobenzyl)-5-methoxy-3-(naphthalene-1-sulfonyl)-1H-indazol-7-yloxy]propyl}dimethylamine hydrochloride;
 ethyl{3-[5-methoxy-3-(naphthalene-1-sulfonyl)-1H-indazol-7-yloxy]propyl}amine hydrochloride;
 {3-[5-methoxy-3-(naphthalene-1-sulfonyl)-1H-indazol-7-yloxy]propyl}dimethylamine hydrochloride;
 N-methyl-2-[[3-naphthylsulfonyl)-1H-indazol-4-yl]oxy}ethanamine hydrochloride;
 dimethyl{2-[3-(naphthalene-1-sulfonyl)-1H-indazol-6-yloxy]ethyl}amine hydrochloride;
 a stereoisomer thereof; and
 a pharmaceutically acceptable salt thereof.

16. The composition according to claim 10 having a formula I compound selected from the group consisting of:

[2-(3-benzenesulfonyl-1H-indazol-5-yloxy)-ethyl]ethyl-methylamine;
 [2-(3-benzenesulfonyl-1H-indazol-5-yloxy)-ethyl]diethylamine;
 [2-(3-benzenesulfonyl-1H-indazol-5-yloxy)-ethyl]isopropylamine;
 [2-(3-benzenesulfonyl-1H-indazol-5-yloxy)-ethyl]butylamine;
 [2-(3-benzenesulfonyl-1H-indazol-5-yloxy)-ethyl]cyclopropylamine;
 dimethyl-{3-[3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]propyl}amine;

methyl-{3-[3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]propyl}amine;
 butyl-{3-[3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]propyl}amine;
 {2-[1-benzyl-3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]ethyl-isopropylamine;
 {2-[1-benzyl-3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-ethyl}diethylamine;
 {2-[1-(3-chloro-benzyl)-3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-ethyl}ethyl-
 methylamine;
 {2-[1-(3-chloro-benzyl)-3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-
 ethyl}diethylamine;
 {2-[1-(3-chloro-benzyl)-3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]-
 ethyl}isopropylamine;
 {2-[1-(3-chloro-benzyl)-3-(naphthalene-1-sulfonyl)-1H-indazol-5-
 yloxy]ethyl}methylamine;
 {2-[1-(3-chloro-benzyl)-3-(naphthalene-1-sulfonyl)-1H-indazol-7-
 yloxy]ethyl}ethylamine;
 {2-[1-(3-chloro-benzyl)-3-(naphthalene-1-sulfonyl)-1H-indazol-7-
 yloxy]ethyl}diethylamine;
 butyl-{2-[1-(3-chloro-benzyl)-3-(naphthalene-1-sulfonyl)-1H-indazol-7-
 yloxy]ethyl}methylamine;
 {2-[3-benzenesulfonyl-1-(3-chloro-benzyl)-1H-indazol-7-yloxy]-ethyl}diethylamine;
 {2-[3-benzenesulfonyl-1-(3-chloro-benzyl)-1H-indazol-7-yloxy]-ethyl}butylamine;
 dimethyl-{2-[1-methyl-3-(naphthalene-1-sulfonyl)-1H-indazol-7-yloxy]-ethyl}amine;
 butyl-{2-[1-methyl-3-(naphthalene-1-sulfonyl)-1H-indazol-7-yloxy]-ethyl}amine;
 [2-(3-benzenesulfonyl-1-methyl-1H-indazol-7-yloxy)-ethyl]ethylamine;
 [2-(3-benzenesulfonyl-1-methyl-1H-indazol-7-yloxy)-ethyl]diethylamine;
 (2-{{1-(3-chlorobenzyl)-5-fluoro-3-(1-naphthylsulfonyl)-1H-indazol-7-
 yl}oxy}ethyl)amine;
 (2-{{1-(3-chlorobenzyl)-5-fluoro-3-(1-naphthylsulfonyl)-1H-indazol-7-
 yl}oxy}ethyl)methylamine hydrochloride;
 {2-[1-(3-chloro-benzyl)-5-fluoro-3-(naphthalene-1-sulfonyl)-1H-indazol-7-
 yloxy]ethyl}ethylamine hydrochloride;
 {2-[1-(3-chloro-benzyl)-5-fluoro-3-(naphthalene-1-sulfonyl)-1H-indazol-7-
 yloxy]ethyl}diethylamine hydrochloride;

{2-[5-fluoro-3-(naphthalene-1-sulfonyl)-1H-indazol-7-yloxy]-ethyl}methylamine hydrochloride;
 {2-[1-(3-chlorobenzyl)-5-methoxy-3-(naphthalene-1-sulfonyl)-1H-indazol-7-yloxy]ethyl}methylamine hydrochloride;
 {2-[1-(3-chlorobenzyl)-5-methoxy-3-(naphthalene-1-sulfonyl)-1H-indazol-7-yloxy]ethyl}ethylamine hydrochloride;
 {2-[1-(3-chlorobenzyl)-5-methoxy-3-(naphthalene-1-sulfonyl)-1H-indazol-7-yloxy]ethyl}dimethylamine hydrochloride;
 {3-[1-(3-chlorobenzyl)-5-methoxy-3-(naphthalene-1-sulfonyl)-1H-indazol-7-yloxy]propyl}methylamine hydrochloride;
 {3-[1-(3-chlorobenzyl)-5-methoxy-3-(naphthalene-1-sulfonyl)-1H-indazol-7-yloxy]propyl}ethylamine hydrochloride;
 {3-[1-(3-chlorobenzyl)-5-methoxy-3-(naphthalene-1-sulfonyl)-1H-indazol-7-yloxy]propyl}dimethylamine hydrochloride;
 ethyl{3-[5-methoxy-3-(naphthalene-1-sulfonyl)-1H-indazol-7-yloxy]propyl}amine hydrochloride;
 {3-[5-methoxy-3-(naphthalene-1-sulfonyl)-1H-indazol-7-yloxy]propyl}dimethylamine hydrochloride;
 N-methyl-2-[[3-naphthylsulfonyl)-1H-indazol-4-yl]oxy}ethanamine hydrochloride;
 dimethyl{2-[3-(naphthalene-1-sulfonyl)-1H-indazol-6-yloxy]ethyl}amine hydrochloride;
 a stereoisomer thereof; and
 a pharmaceutically acceptable salt thereof.

17. The compound according to claim 15, which is dimethyl-{3-[3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]propyl}amine or a pharmaceutically acceptable salt thereof.

18. The compound according to claim 17, which is dimethyl-{3-[3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]propyl}amine hydrochloride.

19. The compound according to claim 15, which is methyl-{3-[3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]propyl}amine or a pharmaceutically acceptable salt thereof.

20. The compound according to claim 19, which is methyl-{3-[3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]propyl}amine hydrochloride.
21. The composition according to claim 10 having a formula I compound, which is dimethyl-{3-[3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]propyl}amine or a pharmaceutically acceptable salt thereof.
22. The composition according to claim 21 which is dimethyl-{3-[3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]propyl}amine hydrochloride.
23. The composition according to claim 10 having a formula I compound, which is methyl-{3-[3-(naphthalene-1-sulfonyl)-1H-indazol-5-yloxy]propyl}amine or a pharmaceutically acceptable salt thereof.