

COMPOSITIONS AND METHODS FOR TCR REPROGRAMMING USING FUSION PROTEINS

CROSS-REFERENCE

[0001] This application claims the benefit of U.S. Provisional Application No. 62/163,342, filed May 18, 2015, which is incorporated herein by reference in its entirety.

BACKGROUND OF THE INVENTION

[0002] Most patients with hematological malignancies or with late-stage solid tumors are incurable with standard therapy. In addition, traditional treatment options often have serious side effects. Numerous attempts have been made to engage a patient's immune system for rejecting cancerous cells, an approach collectively referred to as cancer immunotherapy. However, several obstacles make it rather difficult to achieve clinical effectiveness. Although hundreds of so-called tumor antigens have been identified, these are often derived from self and thus can direct the cancer immunotherapy against healthy tissue, or are poorly immunogenic. Furthermore, cancer cells use multiple mechanisms to render themselves invisible or hostile to the initiation and propagation of an immune attack by cancer immunotherapies.

[0003] Recent developments using chimeric antigen receptor (CAR) modified autologous T-cell therapy, which relies on redirecting genetically engineered T-cells to a suitable cell-surface molecule on cancer cells, show promising results in harnessing the power of the immune system to treat B cell malignancies (see, *e.g.*, Sadelain et al., *Cancer Discovery* 3:388-398 (2013)). The clinical results with CD19-specific CAR T-cells (called CTL019) have shown complete remissions in patients suffering from chronic lymphocytic leukemia (CLL) as well as in childhood acute lymphoblastic leukemia (ALL) (see, *e.g.*, Kalos et al., *Sci Transl Med* 3:95ra73 (2011), Porter et al., *NEJM* 365:725-733 (2011), Grupp et al., *NEJM* 368:1509-1518 (2013)). An alternative approach is the use of T-cell receptor (TCR) alpha and beta chains selected for a tumor-associated peptide antigen for genetically engineering autologous T-cells. These TCR chains will form complete TCR complexes and provide the T-cells with a TCR for a second defined specificity. Encouraging results were obtained with engineered autologous T-cells expressing NY-ESO-1-specific TCR alpha and beta chains in patients with synovial carcinoma.

[0004] Besides the ability for genetically modified T-cells expressing a CAR or a second TCR to recognize and destroy respective target cells *in vitro/ex vivo*, successful patient therapy with engineered T-cells requires the T-cells to be capable of strong activation, expansion, persistence over time, and, in case of relapsing disease, to enable a 'memory' response. High and manageable clinical efficacy of CAR T-cells is currently limited to CD19-positive B cell malignancies and to NY-ESO-1-peptide expressing synovial sarcoma patients expressing HLA-

A2. There is a clear need to improve genetically engineered T-cells to more broadly act against various human malignancies. Described herein are novel fusion proteins of TCR subunits, including CD3 epsilon, CD3gamma and CD3 delta, and of TCR alpha and TCR beta chains with binding domains specific for cell surface antigens that have the potential to overcome limitations of existing approaches. Described herein are novel fusion proteins that more efficiently kill target cells than CARs, but release comparable or lower levels of pro-inflammatory cytokines. These fusion proteins and methods of their use represent an advantage for TFPs relative to CARs because elevated levels of these cytokines have been associated with dose-limiting toxicities for adoptive CAR-T therapies.

SUMMARY OF THE INVENTION

[0005] Provided herein are T-cell receptor (TCR) fusion proteins (TFPs), T-cells engineered to express one or more TFPs, and methods of use thereof for the treatment of diseases.

[0006] In one aspect, provided herein is an isolated recombinant nucleic acid molecule encoding a T-cell receptor (TCR) fusion protein (TFP) comprising a TCR subunit comprising at least a portion of a TCR extracellular domain, and a TCR intracellular domain comprising a stimulatory domain from an intracellular signaling domain of CD3 epsilon; and a human or humanized antibody domain comprising an antigen binding domain wherein the TCR subunit and the antibody domain are operatively linked, and wherein the TFP incorporates into a TCR when expressed in a T-cell.

[0007] In one aspect, provided herein is an isolated recombinant nucleic acid molecule encoding a T-cell receptor (TCR) fusion protein (TFP) comprising a TCR subunit comprising at least a portion of a TCR extracellular domain, and a TCR intracellular domain comprising a stimulatory domain from an intracellular signaling domain of CD3 gamma; and a human or humanized antibody domain comprising an antigen binding domain wherein the TCR subunit and the antibody domain are operatively linked, and wherein the TFP incorporates into a TCR when expressed in a T-cell.

[0008] In one aspect, provided herein is an isolated recombinant nucleic acid molecule encoding a T-cell receptor (TCR) fusion protein (TFP) comprising a TCR subunit comprising at least a portion of a TCR extracellular domain, and a TCR intracellular domain comprising a stimulatory domain from an intracellular signaling domain of CD3 delta; and a human or humanized antibody domain comprising an antigen binding domain wherein the TCR subunit and the antibody domain are operatively linked, and wherein the TFP incorporates into a TCR when expressed in a T-cell.

[0009] In one aspect, provided herein is an isolated recombinant nucleic acid molecule encoding a T-cell receptor (TCR) fusion protein (TFP) comprising a TCR subunit comprising at least a portion of a TCR extracellular domain, and a TCR intracellular domain comprising a stimulatory domain from an intracellular signaling domain of TCR alpha; and a human or humanized antibody domain comprising an antigen binding domain wherein the TCR subunit and the antibody domain are operatively linked, and wherein the TFP incorporates into a TCR when expressed in a T-cell.

[0010] In one aspect, provided herein is an isolated recombinant nucleic acid molecule encoding a T-cell receptor (TCR) fusion protein (TFP) comprising a TCR subunit comprising at least a portion of a TCR extracellular domain, and a TCR intracellular domain comprising a stimulatory domain from an intracellular signaling domain of TCR beta; and a human or humanized antibody domain comprising an antigen binding domain wherein the TCR subunit and the antibody domain are operatively linked, and wherein the TFP incorporates into a TCR when expressed in a T-cell.

[0011] In one aspect, provided herein is an isolated recombinant nucleic acid molecule encoding a T-cell receptor (TCR) fusion protein (TFP) comprising a TCR subunit and a human or humanized antibody domain comprising an antigen binding domain that is an anti-CD19 binding domain.

[0012] In one aspect, provided herein is an isolated recombinant nucleic acid molecule encoding a T-cell receptor (TCR) fusion protein (TFP) comprising a TCR subunit and a human or humanized antibody domain comprising an antigen binding domain that is an anti-B-cell maturation antigen (BCMA) binding domain.

[0013] In some instances, the TCR subunit and the antibody domain are operatively linked. In some instances, the TFP incorporates into a TCR when expressed in a T-cell. In some instances, the encoded antigen binding domain is connected to the TCR extracellular domain by a linker sequence. In some instances, the encoded linker sequence comprises $(G_4S)_n$, wherein $n=1$ to 4. In some instances, the TCR subunit comprises a TCR extracellular domain. In some instances, the TCR subunit comprises a TCR transmembrane domain. In some instances, the TCR subunit comprises a TCR intracellular domain. In some instances, the TCR subunit comprises (i) a TCR extracellular domain, (ii) a TCR transmembrane domain, and (iii) a TCR intracellular domain, wherein at least two of (i), (ii), and (iii) are from the same TCR subunit. In some instances, the TCR subunit comprises a TCR intracellular domain comprising a stimulatory domain selected from an intracellular signaling domain of CD3 epsilon, CD3 gamma or CD3 delta, or an amino acid sequence having at least one, two or three modifications thereto. In some instances, the TCR subunit comprises an intracellular domain comprising a stimulatory domain selected from

a functional signaling domain of 4-1BB and/or a functional signaling domain of CD3 zeta, or an amino acid sequence having at least one modification thereto. In some instances, the human or humanized antibody domain comprises an antibody fragment. In some instances, the human or humanized antibody domain comprises a scFv or a V_H domain. In some instances, the isolated nucleic acid molecule encodes (i) a light chain (LC) CDR1, LC CDR2 and LC CDR3 of an anti-CD19 light chain binding domain amino acid sequence with 70-100% sequence identity to SEQ ID NO: 25, SEQ ID NO: 27 and SEQ ID NO: 29, respectively, and/or (ii) a heavy chain (HC) CDR1, HC CDR2 and HC CDR3 of an anti-CD19 heavy chain binding domain amino acid sequence with 70-100% sequence identity to SEQ ID NO: 31, SEQ ID NO: 33 and SEQ ID NO: 35, respectively. In some instances, the isolated nucleic acid molecule encodes a light chain variable region, wherein the light chain variable region comprises an amino acid sequence having at least one but not more than 30 modifications of a light chain variable region amino acid sequence of SEQ ID NO: 49, or a sequence with 95-99% identity to a light chain variable region amino acid sequence of SEQ ID NO: 49. In some instances, the isolated nucleic acid molecule encodes a heavy chain variable region, wherein the heavy chain variable region comprises an amino acid sequence having at least one but not more than 30 modifications of a heavy chain variable region amino acid sequence of SEQ ID NO: 51, or a sequence with 95-99% identity to a heavy chain variable region amino acid sequence of SEQ ID NO: 51. In some instances, the isolated nucleic acid molecule encodes (i) a light chain (LC) CDR1, LC CDR2 and LC CDR3 of an anti-BCMA light chain binding domain amino acid sequence with 70-100% sequence identity to SEQ ID NO: 37, SEQ ID NO: 39 and SEQ ID NO: 41, respectively, and/or (ii) a heavy chain (HC) CDR1, HC CDR2 and HC CDR3 of an anti-BCMA heavy chain binding domain amino acid sequence with 70-100% sequence identity to SEQ ID NO: 43, SEQ ID NO: 45 and SEQ ID NO: 47, respectively. In some instances, the isolated nucleic acid molecule encodes a light chain variable region, wherein the light chain variable region comprises an amino acid sequence having at least one but not more than 30 modifications of a light chain variable region amino acid sequence of SEQ ID NO: 53, or a sequence with 95-99% identity to a light chain variable region amino acid sequence of SEQ ID NO: 53. In some instances, the isolated nucleic acid molecule encodes a heavy chain variable region, wherein the heavy chain variable region comprises an amino acid sequence having at least one but not more than 30 modifications of a heavy chain variable region amino acid sequence of SEQ ID NO: 55, or a sequence with 95-99% identity to a heavy chain variable region amino acid sequence of SEQ ID NO: 55. In some instances, the TFP includes an extracellular domain of a TCR subunit that comprises an extracellular domain or portion thereof of a protein selected from the group consisting of a TCR alpha chain, a TCR beta chain, a CD3 epsilon TCR subunit, a CD3 gamma

TCR subunit, a CD3 delta TCR subunit, functional fragments thereof, and amino acid sequences thereof having at least one but not more than 20 modifications. In some instances, the encoded TFP includes a transmembrane domain that comprises a transmembrane domain of a protein selected from the group consisting of a TCR alpha chain, a TCR beta chain, a CD3 epsilon TCR subunit, a CD3 gamma TCR subunit, a CD3 delta TCR subunit, functional fragments thereof, and amino acid sequences thereof having at least one but not more than 20 modifications. In some instances, the encoded TFP includes a transmembrane domain that comprises a transmembrane domain of a protein selected from the group consisting of a TCR alpha chain, a TCR beta chain, a TCR zeta chain, a CD3 epsilon TCR subunit, a CD3 gamma TCR subunit, a CD3 delta TCR subunit, CD45, CD4, CD5, CD8, CD9, CD16, CD22, CD33, CD28, CD37, CD64, CD80, CD86, CD134, CD137, CD154, functional fragments thereof, and amino acid sequences thereof having at least one but not more than 20 modifications. In some instances, the isolated nucleic acid molecule further comprises a sequence encoding a costimulatory domain. In some instances, the costimulatory domain is a functional signaling domain obtained from a protein selected from the group consisting of OX40, CD2, CD27, CD28, CD29, ICAM-1, LFA-1 (CD11a/CD18), ICOS (CD278), and 4-1BB (CD137), and amino acid sequences thereof having at least one but not more than 20 modifications thereto. In some instances, the isolated nucleic acid molecule further comprises a leader sequence. In some instances, the isolated nucleic acid molecule is mRNA.

[0014] In some instances, the TFP includes an immunoreceptor tyrosine-based activation motif (ITAM) of a TCR subunit that comprises an ITAM or portion thereof of a protein selected from the group consisting of CD3 zeta TCR subunit, CD3 epsilon TCR subunit, CD3 gamma TCR subunit, CD3 delta TCR subunit, TCR zeta chain, Fc epsilon receptor 1 chain, Fc epsilon receptor 2 chain, Fc gamma receptor 1 chain, Fc gamma receptor 2a chain, Fc gamma receptor 2b1 chain, Fc gamma receptor 2b2 chain, Fc gamma receptor 3a chain, Fc gamma receptor 3b chain, Fc beta receptor 1 chain, TYROBP (DAP12), CD5, CD16a, CD16b, CD22, CD23, CD32, CD64, CD79a, CD79b, CD89, CD278, CD66d, functional fragments thereof, and amino acid sequences thereof having at least one but not more than 20 modifications thereto. In some instances, the ITAM replaces an ITAM of CD3 gamma, CD3 delta, or CD3 epsilon. In some instances, the ITAM is selected from the group consisting of CD3 zeta TCR subunit, CD3 epsilon TCR subunit, CD3 gamma TCR subunit, and CD3 delta TCR subunit and replaces a different ITAM selected from the group consisting of CD3 zeta TCR subunit, CD3 epsilon TCR subunit, CD3 gamma TCR subunit, and CD3 delta TCR subunit.

[0015] In some instances, the nucleic acid comprises a nucleotide analog. In some instances, the nucleotide analog is selected from the group consisting of 2'-O-methyl, 2'-O-methoxyethyl (2'-

O-MOE), 2'-O-aminopropyl, 2'-deoxy, T-deoxy-2'-fluoro, 2'-O-aminopropyl (2'-O-AP), 2'-O-dimethylaminoethyl (2'-O-DMAOE), 2'-O-dimethylaminopropyl (2'-O-DMAP), T-O-dimethylaminoethoxyethyl (2'-O-DMAEOE), 2'-O-N-methylacetamido (2'-O-NMA) modified, a locked nucleic acid (LNA), an ethylene nucleic acid (ENA), a peptide nucleic acid (PNA), a 1',5'- anhydrohexitol nucleic acid (HNA), a morpholino, a methylphosphonate nucleotide, a thiolphosphonate nucleotide, and a 2'-fluoro N3-P5'-phosphoramidite

[0016] In one aspect, provided herein is an isolated polypeptide molecule encoded by a nucleic acid molecule provided herein.

[0017] In one aspect, provided herein is an isolated TFP molecule comprising a human or humanized anti-CD19 binding domain, a TCR extracellular domain, a transmembrane domain, and an intracellular domain.

[0018] In one aspect, provided herein is an isolated TFP molecule comprising a human or humanized anti-CD19 binding domain, a TCR extracellular domain, a transmembrane domain, and an intracellular signaling domain, wherein the TFP molecule is capable of functionally interacting with an endogenous TCR complex and/or at least one endogenous TCR polypeptide.

[0019] In one aspect, provided herein is an isolated TFP molecule comprising a human or humanized anti-CD19 binding domain, a TCR extracellular domain, a transmembrane domain, and an intracellular signaling domain, wherein the TFP molecule is capable of functionally integrating into an endogenous TCR complex

[0020] In some instances, the isolated TFP molecule comprises an antibody or antibody fragment comprising a human or humanized anti-CD19 binding domain, a TCR extracellular domain, a transmembrane domain, and an intracellular domain. In some instances, the anti-CD19 binding domain is a scFv or a V_H domain. In some instances, the anti-CD19 binding domain comprises a heavy chain with 95-100% identity to an amino acid sequence of SEQ ID NO: 51, a functional fragment thereof, or an amino acid sequence thereof having at least one but not more than 30 modifications. In some instances, the anti-CD19 binding domain comprises a light chain with 95-100% identity to an amino acid sequence of SEQ ID NO: 49, a functional fragment thereof, or an amino acid sequence thereof having at least one but not more than 30 modifications. In some instances, the isolated TFP molecule comprises a TCR extracellular domain that comprises an extracellular domain or portion thereof of a protein selected from the group consisting of a TCR alpha chain, a TCR beta chain, a CD3 epsilon TCR subunit, a CD3 gamma TCR subunit, a CD3 delta TCR subunit, functional fragments thereof, and amino acid sequences thereof having at least one but not more than 20 modifications. In some instances, the anti-CD19 binding domain is connected to the TCR extracellular domain by a linker sequence. In some instances, the linker region comprises (G₄S)_n, wherein n=1 to 4.

[0021] In one aspect, provided herein is an isolated TFP molecule comprising a human or humanized anti-BCMA binding domain, a TCR extracellular domain, a transmembrane domain, and an intracellular domain.

[0022] In one aspect, provided herein is an isolated TFP molecule comprising a human or humanized anti-BCMA binding domain, a TCR extracellular domain, a transmembrane domain, and an intracellular signaling domain, wherein the TFP molecule is capable of functionally interacting with an endogenous TCR complex and/or at least one endogenous TCR polypeptide.

[0023] In one aspect, provided herein is an isolated TFP molecule comprising a human or humanized anti-BCMA binding domain, a TCR extracellular domain, a transmembrane domain, and an intracellular signaling domain, wherein the TFP molecule is capable of functionally integrating into an endogenous TCR complex.

[0024] In some instances, the isolated TFP molecule comprises an antibody or antibody fragment comprising a human or humanized anti-BCMA binding domain, a TCR extracellular domain, a transmembrane domain, and an intracellular domain. In some instances, the anti-BCMA binding domain is a scFv or a V_H domain. In some instances, the anti-BCMA binding domain comprises a heavy chain with 95-100% identity to an amino acid sequence of SEQ ID NO: 55, a functional fragment thereof, or an amino acid sequence thereof having at least one but not more than 30 modifications. In some instances, the anti-BCMA binding domain comprises a light chain with 95-100% identity to an amino acid sequence of SEQ ID NO: 53, a functional fragment thereof, or an amino acid sequence thereof having at least one but not more than 30 modifications. In some instances, the isolated TFP molecule comprises a TCR extracellular domain that comprises an extracellular domain or portion thereof of a protein selected from the group consisting of a TCR alpha chain, a TCR beta chain, a CD3 epsilon TCR subunit, a CD3 gamma TCR subunit, a CD3 delta TCR subunit, functional fragments thereof, and amino acid sequences thereof having at least one but not more than 20 modifications. In some instances, the anti-BCMA binding domain is connected to the TCR extracellular domain by a linker sequence. In some instances, the linker region comprises (G₄S)_n, wherein n=1 to 4. In some instances, the isolated TFP molecule further comprises a sequence encoding a costimulatory domain. In some instances, the isolated TFP molecule further comprises a sequence encoding an intracellular signaling domain. In some instances, the isolated TFP molecule further comprises a leader sequence.

[0025] In one aspect, provided herein is a vector comprising a nucleic acid molecule encoding a TFP provided herein. In some instances, the vector is selected from the group consisting of a DNA, a RNA, a plasmid, a lentivirus vector, adenoviral vector, a Rous sarcoma viral (RSV) vector, or a retrovirus vector. In some instances, the vector further comprises a promoter. In

some instances, the vector is an *in vitro* transcribed vector. In some instances, a nucleic acid sequence in the vector further comprises a poly(A) tail. In some instances, a nucleic acid sequence in the vector further comprises a 3'UTR.

[0026] In one aspect, provided herein is a cell comprising a vector provided herein. In some instances, the cell is a human T-cell. In some instances, the T-cell is a CD8⁺ or CD4⁺ T-cell. In some instances, the cell further comprises a nucleic acid encoding an inhibitory molecule that comprises a first polypeptide that comprises at least a portion of an inhibitory molecule, associated with a second polypeptide that comprises a positive signal from an intracellular signaling domain. In some instances, the inhibitory molecule comprise first polypeptide that comprises at least a portion of PD1 and a second polypeptide comprising a costimulatory domain and primary signaling domain.

[0027] In one aspect, provided herein is a human CD8⁺ or CD4⁺ T-cell comprising at least two TFP molecules, the TFP molecules comprising a human or humanized anti-CD19 binding domain, a TCR extracellular domain, a transmembrane domain, and an intracellular domain, wherein the TFP molecule is capable of functionally interacting with an endogenous TCR complex and/or at least one endogenous TCR polypeptide in, at and/or on the surface of the human CD8⁺ or CD4⁺ T-cell.

[0028] In one aspect, provided herein is a protein complex comprising: a TFP molecule comprising a human or humanized anti-CD19 binding domain, a TCR extracellular domain, a transmembrane domain, and an intracellular domain; and at least one endogenous TCR complex.

[0029] In some instances, the TCR comprises an extracellular domain or portion thereof of a protein selected from the group consisting of TCR alpha chain, a TCR beta chain, a CD3 epsilon TCR subunit, a CD3 gamma TCR subunit, and a CD3 delta TCR subunit. In some instances, the anti-CD19 binding domain is connected to the TCR extracellular domain by a linker sequence. In some instances, the linker region comprises (G₄S)_n, wherein n=1 to 4.

[0030] In one aspect, provided herein is a protein complex comprising: a TFP molecule comprising a human or humanized anti-BCMA binding domain, a TCR extracellular domain, a transmembrane domain, and an intracellular domain; and at least one endogenous TCR complex.

[0031] In some instances, the TCR comprises an extracellular domain or portion thereof of a protein selected from the group consisting of TCR alpha chain, a TCR beta chain, a CD3 epsilon TCR subunit, a CD3 gamma TCR subunit, and a CD3 delta TCR subunit. In some instances, the anti-BCMA binding domain is connected to the TCR extracellular domain by a linker sequence. In some instances, the linker region comprises (G₄S)_n, wherein n=1 to 4.

[0032] In one aspect, provided herein is a human CD8⁺ or CD4⁺ T-cell comprising at least two different TFP proteins per a protein complex provided herein.

[0033] In one aspect, provided herein is a method of making a cell comprising transducing a T-cell with a vector provided herein.

[0034] In one aspect, provided herein is a method of generating a population of RNA-engineered cells comprising introducing an *in vitro* transcribed RNA or synthetic RNA into a cell, where the RNA comprises a nucleic acid encoding a TFP molecule provided herein.

[0035] In one aspect, provided herein is a method of providing an anti-tumor immunity in a mammal comprising administering to the mammal an effective amount of a cell expressing a TFP molecule provided herein, or expressing a polypeptide molecule provided herein.

[0036] In some instances, the cell is an autologous T-cell. In some instances, the cell is an allogeneic T-cell. In some instances, the mammal is a human.

[0037] In one aspect, provided herein is a method of treating a mammal having a disease associated with expression of CD19 or BCMA comprising administering to the mammal an effective amount of a TFP molecule provided herein, a cell provided herein, or a polypeptide molecule provided herein.

[0038] In some instances, the disease associated with CD19 or BCMA expression is selected from the group consisting of a proliferative disease, a cancer, a malignancy, myelodysplasia, a myelodysplastic syndrome, a preleukemia, a non-cancer related indication associated with expression of CD19. In some instances, the disease is a hematologic cancer selected from the group consisting of B-cell acute lymphoid leukemia (B-ALL), T-cell acute lymphoid leukemia (T-ALL), acute lymphoblastic leukemia (ALL); chronic myelogenous leukemia (CML), chronic lymphocytic leukemia (CLL), B cell prolymphocytic leukemia, blastic plasmacytoid dendritic cell neoplasm, Burkitt's lymphoma, diffuse large B cell lymphoma, follicular lymphoma, hairy cell leukemia, small cell-follicular lymphoma, large cell-follicular lymphoma, malignant lymphoproliferative conditions, MALT lymphoma, mantle cell lymphoma, Marginal zone lymphoma, multiple myeloma, myelodysplasia, myelodysplastic syndrome, non-Hodgkin's lymphoma, plasmablastic lymphoma, plasmacytoid dendritic cell neoplasm, Waldenstrom macroglobulinemia, preleukemia, a disease associated with CD19 or BCMA expression, and combinations thereof. In some instances, the cells expressing a TFP molecule are administered in combination with an agent that increases the efficacy of a cell expressing a TFP molecule. In some instances, less cytokines are released in the mammal compared a mammal administered an effective amount of a T-cell expressing an anti-CD19 chimeric antigen receptor (CAR) or an anti-BCMA CAR. In some instances, the cells expressing a TFP molecule are administered in combination with an agent that ameliorates one or more side effects associated with administration of a cell expressing a TFP molecule. In some instances, the cells expressing a

TFP molecule are administered in combination with an agent that treats the disease associated with CD19 or BCMA.

[0039] In one aspect, an isolated nucleic acid molecule provided herein, an isolated polypeptide molecule provided herein, an isolated TFP provided herein, a complex provided herein, a vector provided herein, or a cell provided herein, is for use as a medicament.

[0040] In one aspect, provided herein is a method of treating a mammal having a disease associated with expression of CD19 or BCMA comprising administering to the mammal an effective amount of a TFP molecule provided herein, a cell provided herein, or a polypeptide molecule provided herein, wherein less cytokines are released in the mammal compared a mammal administered an effective amount of a T-cell expressing an anti-CD19 chimeric antigen receptor (CAR) or an anti-BCMA CAR.

[0040a] In one aspect, provided herein is a pharmaceutical composition comprising (I) a T cell from a human subject, wherein the T cell comprises a recombinant nucleic acid molecule encoding a T-cell receptor (TCR) fusion protein (TFP) comprising

(a) a TCR subunit comprising

(i) at least a portion of a TCR extracellular domain,

(ii) a TCR transmembrane domain, and

(iii) a TCR intracellular domain comprising a stimulatory domain from an intracellular signaling domain of CD3 epsilon, CD3 gamma, or CD3 delta; and

(b) a murine, human or humanized scFv or single domain antibody comprising an antigen binding domain;

wherein the TCR subunit and the antibody domain are operatively linked,

wherein the TFP incorporates into a TCR when expressed in a T-cell, and

wherein the T cell exhibits increased cytotoxicity to a human cell expressing an antigen that specifically interacts with the antigen binding domain compared to a T cell not containing the TFP; and

(II) a pharmaceutically acceptable carrier.

[0040b] In one aspect, provided herein is a use of an effective amount of the pharmaceutical composition as disclosed herein in the manufacture of a medicament for treating a mammal having a disease.

INCORPORATION BY REFERENCE

[0041] All publications, patents, and patent applications mentioned in this specification are herein incorporated by reference to the same extent as if each individual publication, patent, or patent application was specifically and individually indicated to be incorporated by reference.

BRIEF DESCRIPTION OF THE DRAWINGS

The novel features of the invention are set forth with particularity in the appended claims. A better understanding of the features and advantages of the present invention will be obtained by reference to the following detailed description that sets forth illustrative embodiments, in which the principles of the invention are utilized, and the accompanying drawings of which:

[0042] FIG. 1 is a schematic illustration demonstrating the use of T-cell receptor fusion polypeptides (TFPs) of the invention. An exemplary TFP contains an anti-CD19 scFv and a full-length CD3 epsilon polypeptide fused via a $(G_4S)_3$ linker sequence. When produced by or introduced into a T-cell, the TFP associates with other polypeptides of the endogenous T-cell receptor (TCR) (shown to include two CD3 epsilon polypeptides, one CD3 gamma polypeptide, one CD3 delta polypeptide, two CD3 zeta polypeptides, one TCR alpha subunit and one TCR beta subunit, where the horizontal grey segment represents the plasma membrane) to form a reprogrammed TCR in which one or both of the endogenous CD3 epsilon polypeptides are substituted by the TFP.

[0043] FIG. 2A represents schematic illustrations demonstrating exemplary variations of reprogrammed T-cell receptor fusion polypeptides (TFPs) of the invention.

[0044] Figure 2A illustrates an exemplary reprogrammed TCR containing a TFP that contains an anti-CD19 scFv and a full-length TCR V α polypeptide fused via a $(G_4S)_3$ linker sequence.

[0046] **FIG. 2B** illustrates a series of exemplary reprogrammed TCRs that contain multiple TFPs including i) an anti-CD19 scFv and a full-length TCR V α polypeptide fused via a (G₄S)₃ linker sequence and ii) an anti-CD19 scFv and a full-length TCR V β polypeptide fused via a (G₄S)₃ linker sequence.

[0047] **FIG. 2C** illustrates an exemplary reprogrammed TCR that contains multiple TFPs including i) an anti-CD19 scFv and a truncated (Δ) TCR polypeptide fused via a (G₄S)₃ linker sequence and ii) an anti-CD19 scFv and a full-length CD3 epsilon polypeptide fused via a (G₄S)₃ linker sequence. The truncated (Δ) TCR polypeptide is truncated by the deletion of the V α .

[0048] **FIG. 2D** illustrates an exemplary reprogrammed TCR that contains multiple TFPs including i) an anti-CD19 scFv and a truncated (Δ) TCR V α polypeptide fused via a (G₄S)₃ linker sequence and ii) an anti-CD19 scFv and a truncated (Δ) TCR V β polypeptide fused via a (G₄S)₃ linker sequence. The truncated (Δ) TCR polypeptide is truncated by the deletion of the V β .

[0049] **FIG. 3** is a schematic illustration demonstrating the use of T-cell receptor fusion polypeptides (TFPs) of the invention. An exemplary TFP contains an anti-CD19 V_H domain and a full-length CD3 epsilon polypeptide fused via a (G₄S)₃ linker sequence. When produced by a T-cell or introduced into a T-cell, the TFP associates with other polypeptides of the endogenous T-cell receptor (TCR) (shown to include two CD3 epsilon polypeptides, one CD3 gamma polypeptide, one CD3 delta polypeptide, two CD3 zeta polypeptides, one TCR alpha subunit and one TCR beta subunit, where the horizontal grey segment represents the plasma membrane) to form a reprogrammed TCR in which one or both of the endogenous CD3 epsilon polypeptides are substituted by the TFP.

[0050] **FIG. 4** is a series of schematic illustrations demonstrating DNA constructs encoding various TFPs.

[0051] **FIG. 5** is an exemplary bar graph depicting surface expression of anti-CD19 LL (long linker) TFPs on T-cells after lentiviral transduction. Effector T-cells were either un-transduced or transduced with either anti-CD19-28 ζ CAR or the indicated anti-CD19 LL TFP constructs. After being expanded for 10 days in IL-2, their surface expression of the appropriate CAR or TFP construct was determined by flow cytometry.

[0052] **FIG. 6** is an exemplary bar graph depicting surface expression of anti-CD19 SL (short linker) TFPs on T-cells after lentiviral transduction. Effector T-cells were either un-transduced or transduced with either anti-CD19-28 ζ CAR or the indicated anti-CD19 SL TFP constructs. After being expanded for 7 days in IL-2, their surface expression of the appropriate CAR or TFP construct was determined by flow cytometry.

[0053] **FIG. 7** is an exemplary bar graph depicting surface expression of anti-BCMA TFPs on T-cells after lentiviral transduction. Effector T-cells were either un-transduced or transduced with either anti-BCMA-CD3 ϵ or anti-BCMA-CD3 γ TFP constructs. After being expanded for 10 days in IL-2, their surface TFP expression was determined by flow cytometry.

[0054] **FIG. 8** is an exemplary bar graph depicting killing of CD19-expressing Raji target cells by anti-CD19 LL TFPs. Transduced effector T-cells were expanded for 14 days prior to incubation for 18 hours with 1×10^4 Raji target cells at E:T ratios of 20:1, 10:1, or 5:1. The percentage cytotoxicity was determined in a flow-cytometric cytotoxicity assay.

[0055] **FIG. 9** is an exemplary bar graph depicting killing of BCMA-expressing RPMI8226 target cells by anti-BCMA TFPs. Transduced effector T-cells were expanded for 12 days prior to incubation for 4 hours with 1×10^4 RPMI8226 target cells at E:T ratios of 10:1, or 5:1. The percentage cytotoxicity was determined in a flow-cytometric cytotoxicity assay.

[0056] **FIG. 10A** is an exemplary graph depicting killing of CD19-transduced HeLa target cells by an anti-CD19-28 ζ CAR construct over time. Transduced effector T-cells were expanded for 14 days prior to incubation with 1×10^4 CD19-transduced HeLa target cells. The cell index, indicative of cytotoxicity, was determined in a RTCA assay.

[0057] **FIG. 10B** is an exemplary graph depicting killing of CD19-transduced HeLa target cells by an anti-CD19-CD3 ϵ LL TFP construct over time. Transduced effector T-cells were expanded for 14 days prior to incubation with 1×10^4 CD19-transduced HeLa target cells. The cell index, indicative of cytotoxicity, was determined in a RTCA assay.

[0058] **FIG. 10C** is an exemplary graph depicting killing of CD19-transduced HeLa target cells by an anti-CD19-CD3 γ LL TFP construct over time. Transduced effector T-cells were expanded for 14 days prior to incubation with 1×10^4 CD19-transduced HeLa target cells. The cell index, indicative of cytotoxicity, was determined in a RTCA assay.

[0059] **FIG. 10D** is an exemplary graph depicting killing of CD19-transduced HeLa target cells by anti-CD19-TCR $\alpha\epsilon$ LL TFP construct over time. Transduced effector T-cells were expanded for 14 days prior to incubation with 1×10^4 CD19-transduced HeLa target cells. The cell index, indicative of cytotoxicity, was determined in a RTCA assay.

[0060] **FIG. 10E** is an exemplary graph depicting killing of CD19-transduced HeLa target cells by anti-CD19-TCR $\beta\epsilon$ LL TFP construct over time. Transduced effector T-cells were expanded for 14 days prior to incubation with 1×10^4 CD19-transduced HeLa target cells. The cell index, indicative of cytotoxicity, was determined in a RTCA assay.

[0061] **FIG. 10F** is an exemplary graph depicting killing of CD19-transduced HeLa target cells by anti-CD19-TCR α LL TFP construct over time. Transduced effector T-cells were expanded

for 14 days prior to incubation with 1×10^4 CD19-transduced HeLa target cells. The cell index, indicative of cytotoxicity, was determined in a RTCA assay.

[0062] **FIG. 10G** is an exemplary graph depicting killing of CD19-transduced HeLa target cells by anti-CD19-TCR β LL TFP construct over time. Transduced effector T-cells were expanded for 14 days prior to incubation with 1×10^4 CD19-transduced HeLa target cells. The cell index, indicative of cytotoxicity, was determined in a RTCA assay.

[0063] **FIG. 11** is an exemplary graph depicting killing of CD19-transduced HeLa target cells by anti-CD19 TFPs. Transduced effector T-cells were expanded for 7 days prior to incubation with 1×10^4 CD19-transduced HeLa target cells. The cell index, indicative of cytotoxicity, was determined in a RTCA assay.

[0064] **FIG. 12** is an exemplary graph depicting killing of BCMA-transduced HeLa target cells by anti-BCMA TFPs over time. Effector T-cells that were either non-transduced or transduced with either anti-BCMA-CD3 ϵ or anti-BCMA-CD3 γ TFPs were expanded for 7 days prior to incubation with either 1×10^4 HeLa or HeLa-BCMA target cells. The cell index, indicative of cytotoxicity, was determined in a RTCA assay.

[0065] **FIG. 13** is an exemplary graph depicting killing activity of T-cells transduced with various amounts of lentivirus encoding anti-CD19-CD3 ϵ LL TFP over time. T-cells transduced with the indicated MOI of lentivirus encoding anti-CD19-CD3 ϵ LL TFP were expanded for 14 days prior to incubation with 1×10^4 CD19-transduced HeLa target cells. The cell index, indicative of cytotoxicity, was determined.

[0066] **FIG. 14** is an exemplary graph depicting killing activity of T-cells transfected by electroporation with *in vitro* transcribed (IVT) mRNA encoding anti-CD19-CD3 ϵ SL or anti-CD19-CD3 γ SL TRuCs. Effector T cells were transfected by electroporation of activated PBMCs with *in vitro* transcribed (IVT) mRNAs encoding either GFP control, anti-CD19-CD3 ϵ SL, or anti-CD19-CD3 γ SL TRuCs. After expansion for 3 days the effectors were incubated for 4 hours with 1×10^4 Raji cells or K562 cells at E:T ratios of 10:1. The percentage cytotoxicity was determined in a flow-cytometric cytotoxicity assay.

[0067] **FIG. 15A** is an exemplary graph depicting IL-2 release by T-cells transduced with anti-CD19 LL TFPs in response to CD19-bearing target cells. Effector T-cells that were either non-transduced, transduced with a control CAR, an anti-CD19-28 ζ CAT or the indicated anti-CD19 LL TFP, were expanded for 14 days prior to incubation with either 1×10^4 Raji or K562 target cells. IL-2 levels were determined by ELISA.

[0068] **FIG. 15B** is an exemplary graph depicting IFN- γ release by T-cells transduced with anti-CD19 LL TFPs in response to CD19-bearing target cells. Effector T-cells that were either non-transduced, transduced with a control CAR, an anti-CD19-28 ζ CAT or the indicated anti-CD19

LL TFP, were expanded for 14 days prior to incubation with either 1×10^4 Raji or K562 target cells. IFN- γ levels were determined by ELISA.

[0069] **FIG. 15C** is an exemplary graph depicting IL-2 release by T-cells transduced with anti-CD19 LL TFPs in response to CD19-bearing target cells. Effector T-cells that were either non-transduced, transduced with a control CAR, an anti-CD19-28 ζ CAT or the indicated anti-CD19 LL TFP, were expanded for 14 days prior to incubation with either 1×10^4 HeLa or CD19-HeLa target cells. IL-2 levels were determined by ELISA.

[0070] **FIG. 15D** is an exemplary graph depicting IFN- γ release by T-cells transduced with anti-CD19 LL TFPs in response to CD19-bearing target cells. Effector T-cells that were either non-transduced, transduced with a control CAR, an anti-CD19-28 ζ CAT or the indicated anti-CD19 LL TFP, were expanded for 14 days prior to incubation with either 1×10^4 HeLa or CD19-HeLa target cells. IFN- γ levels were determined by ELISA.

[0071] **FIG. 16** is an exemplary graph depicting IFN- γ release by T-cells transduced with anti-CD19 TFPs in response to CD19-bearing target cells. Effector T-cells that were either non-transduced or transduced with the indicated anti-CD19 TFP were expanded for 7 days prior to incubation with either 1×10^4 HeLa or CD19-HeLa target cells. IFN- γ levels were determined by ELISA.

[0072] **FIG. 17A** is an exemplary graph depicting IL-2 release by T-cells transduced with anti-BCMA TFPs in response to BCMA-bearing target cells. Effector T-cells that were either non-transduced or transduced with either anti-BCMA-CD3 ϵ or anti-BCMA-CD3 γ TFPs were expanded for 7 days prior to incubation with either 1×10^4 HeLa or HeLa-BCMA target cells. IL-2 production was determined by 2-plex Luminex.

[0073] **FIG. 17B** is an exemplary graph depicting IFN- γ release by T-cells transduced with anti-BCMA TFPs in response to BCMA-bearing target cells. Effector T-cells that were either non-transduced or transduced with either anti-BCMA-CD3 ϵ or anti-BCMA-CD3 γ TFPs were expanded for 7 days prior to incubation with either 1×10^4 HeLa or HeLa-BCMA target cells. IFN- γ production was determined by 2-plex Luminex.

[0074] **FIG. 18** is an exemplary graph depicting degranulation of T-cells transduced with anti-CD19 TFPs in response to CD19-bearing target cells. Effector T-cells that were either non-transduced or transduced with either anti-CD19-28 ζ CAR, anti-BCMA-CD3 ϵ LL TFP or anti-BCMA-CD3 γ LL TFP were expanded for 14 days prior to incubation with 1×10^4 of the indicated CD19 +ve or CD19 -ve target cells. The percentage of CD107+ cells in the CD3+CD8+ gate was determined. Target and effector cells were co-cultured in the presence of a fluorescently-labelled anti-CD107a antibody. The percentage of T-cells within CD3 and

CD4/CD8 gates that stained positively for cell surface CD107a was then determined by flow cytometry.

[0075] **FIG. 19** is an exemplary graph depicting degranulation of T-cells transduced with anti-BCMA TFPs in response to BCMA-bearing target cells. Effector T-cells that were either non-transduced or transduced with 50 MOI of either anti-BCMA-CD3 ϵ or anti-BCMA-CD3 γ TFPs were expanded for 13 days prior to incubation with 1×10^4 of the indicated BCMA +ve RPMI8226 target cells. The percentage of CD107+ cells in the CD3+CD8+ gate was determined.

[0076] **FIG. 20A** depicts exemplary graphs of the *in-vivo* efficacy of T-cells transduced with anti-CD19 LL TFPs in disseminated human leukemic xenograft models. NSG mice were challenged intravenously with either 5×10^5 Raji cells three days prior to adoptive transfer of 5×10^6 T-cells that were either non-transduced or transduced with either anti-CD19-28 ζ CAR, anti-CD19-CD3 ϵ LL TFP or anti-CD19-CD3 γ LL TFP.

[0077] **FIG. 20B** depicts exemplary graphs of the *in-vivo* efficacy of T-cells transduced with anti-CD19 LL TFPs in disseminated human leukemic xenograft models. NSG mice were challenged intravenously with either 1×10^6 Nalm-6 cells (right) three days prior to adoptive transfer of 5×10^6 T-cells that were either non-transduced or transduced with either anti-CD19-28 ζ CAR, anti-CD19-CD3 ϵ LL TFP or anti-CD19-CD3 γ LL TFP. Comparison of survival curves by the log-rank (Mantel-Cox) test showed a $p = 0.0001$ (Group 4 vs 1, 2, 3), $p = 0.0001$ (Group 1 vs 2, 3), and $p = 0.0004$ (Group 2 vs 3). Comparison of survival curves by the Gehan-Breslow-Wilcoxon test showed a $p = 0.0001$ (Group 4 vs 1, 2, 3), $p = 0.0001$ (Group 1 vs 2, 3), and $p = 0.0005$ (Group 2 vs 3).

DETAILED DESCRIPTION OF THE INVENTION

[0078] In one aspect, described herein are isolated nucleic acid molecules encoding a T-cell Receptor (TCR) fusion protein (TFP) that comprise a TCR subunit and a human or humanized antibody domain comprising an anti-CD19 binding domain. In some embodiments, the TCR subunit comprises a TCR extracellular domain. In other embodiments, the TCR subunit comprises a TCR transmembrane domain. In yet other embodiments, the TCR subunit comprises a TCR intracellular domain. In further embodiments, the TCR subunit comprises (i) a TCR extracellular domain, (ii) a TCR transmembrane domain, and (iii) a TCR intracellular domain, wherein at least two of (i), (ii), and (iii) are from the same TCR subunit. In yet further embodiments, the TCR subunit comprises a TCR intracellular domain comprising a stimulatory domain selected from an intracellular signaling domain of CD3 epsilon, CD3 gamma or CD3 delta, or an amino acid sequence having at least one, two or three modifications thereto. In yet

further embodiments, the TCR subunit comprises an intracellular domain comprising a stimulatory domain selected from a functional signaling domain of 4-1BB and/or a functional signaling domain of CD3 zeta, or an amino acid sequence having at least one, two or three modifications thereto.

[0079] In some embodiments, the human or humanized antibody domain comprises an antibody fragment. In some embodiments, the human or humanized antibody domain comprises a scFv or a V_H domain.

[0080] In some embodiments, the isolated nucleic acid molecules comprise (i) a light chain (LC) CDR1, LC CDR2 and LC CDR3 of any anti-CD19 light chain binding domain amino acid sequence provided herein, and/or (ii) a heavy chain (HC) CDR1, HC CDR2 and HC CDR3 of any anti-CD19 heavy chain binding domain amino acid sequence provided herein.

[0081] In some embodiments, the light chain variable region comprises an amino acid sequence having at least one, two or three modifications but not more than 30, 20 or 10 modifications of an amino acid sequence of a light chain variable region provided herein, or a sequence with 95-99% identity to an amino acid sequence provided herein. In other embodiments, the heavy chain variable region comprises an amino acid sequence having at least one, two or three modifications but not more than 30, 20 or 10 modifications of an amino acid sequence of a heavy chain variable region provided herein, or a sequence with 95-99% identity to an amino acid sequence provided herein.

[0082] In some embodiments, the TFP includes an extracellular domain of a TCR subunit that comprises an extracellular domain or portion thereof of a protein selected from the group consisting of the alpha or beta chain of the T-cell receptor, CD3 delta, CD3 epsilon, or CD3 gamma, or a functional fragment thereof, or an amino acid sequence having at least one, two or three modifications but not more than 20, 10 or 5 modifications thereto. In other embodiments, the encoded TFP includes a transmembrane domain that comprises a transmembrane domain of a protein selected from the group consisting of the alpha, beta chain of the TCR or TCR subunits CD3 epsilon, CD3 gamma and CD3 delta, or a functional fragment thereof, or an amino acid sequence having at least one, two or three modifications but not more than 20, 10 or 5 modifications thereto.

[0083] In some embodiments, the encoded TFP includes a transmembrane domain that comprises a transmembrane domain of a protein selected from the group consisting of the alpha, beta or zeta chain of the TCR or CD3 epsilon, CD3 gamma and CD3 delta CD45, CD4, CD5, CD8, CD9, CD16, CD22, CD33, CD28, CD37, CD64, CD80, CD86, CD134, CD137 and CD154, or a functional fragment thereof, or an amino acid sequence having at least one, two or three modifications but not more than 20, 10 or 5 modifications thereto.

[0084] In some embodiments, the encoded anti-CD19 binding domain is connected to the TCR extracellular domain by a linker sequence. In some instances, the encoded linker sequence comprises $(G_4S)_n$, wherein $n=1$ to 4. In some instances, the encoded linker sequence comprises a long linker (LL) sequence. In some instances, the encoded long linker sequence comprises $(G_4S)_n$, wherein $n=2$ to 4. In some instances, the encoded linker sequence comprises a short linker (SL) sequence. In some instances, the encoded short linker sequence comprises $(G_4S)_n$, wherein $n=1$ to 3.

[0085] In some embodiments, the isolated nucleic acid molecules further comprise a sequence encoding a costimulatory domain. In some instances, the costimulatory domain is a functional signaling domain obtained from a protein selected from the group consisting of OX40, CD2, CD27, CD28, CDS, ICAM-1, LFA-1 (CD11a/CD18), ICOS (CD278), and 4-1BB (CD137), or an amino acid sequence having at least one, two or three modifications but not more than 20, 10 or 5 modifications thereto.

[0086] In some embodiments, the isolated nucleic acid molecules further comprise a leader sequence.

[0087] Also provided herein are isolated polypeptide molecules encoded by any of the previously described nucleic acid molecules.

[0088] Also provided herein in another aspect, are isolated T-cell receptor fusion protein (TFP) molecules that comprise a human or humanized anti-CD19 binding domain, a TCR extracellular domain, a transmembrane domain, and an intracellular domain. In some embodiments, the isolated TFP molecules comprises an antibody or antibody fragment comprising a human or humanized anti-CD19 binding domain, a TCR extracellular domain, a transmembrane domain, and an intracellular domain.

[0089] In some embodiments, the anti-CD19 binding domain is a scFv or a V_H domain. In other embodiments, the anti-CD19 binding domain comprises a light chain and a heavy chain of an amino acid sequence provided herein, or a functional fragment thereof, or an amino acid sequence having at least one, two or three modifications but not more than 30, 20 or 10 modifications of an amino acid sequence of a light chain variable region provided herein, or a sequence with 95-99% identity with an amino acid sequence provided herein.

[0090] In some embodiments, the isolated TFP molecules comprise a TCR extracellular domain that comprises an extracellular domain or portion thereof of a protein selected from the group consisting of the alpha or beta chain of the T-cell receptor, CD3 delta, CD3 epsilon, or CD3 gamma, or an amino acid sequence having at least one, two or three modifications but not more than 20, 10 or 5 modifications thereto.

[0091] In some embodiments, the anti-CD19 binding domain is connected to the TCR extracellular domain by a linker sequence. In some instances, the linker region comprises $(G_4S)_n$, wherein $n=1$ to 4. In some instances, the linker sequence comprises a long linker (LL) sequence. In some instances, the long linker sequence comprises $(G_4S)_n$, wherein $n=2$ to 4. In some instances, the linker sequence comprises a short linker (SL) sequence. In some instances, the short linker sequence comprises $(G_4S)_n$, wherein $n=1$ to 3.

[0092] In some embodiments, the isolated TFP molecules further comprise a sequence encoding a costimulatory domain. In other embodiments, the isolated TFP molecules further comprise a sequence encoding an intracellular signaling domain. In yet other embodiments, the isolated TFP molecules further comprise a leader sequence.

[0093] Also provided herein are vectors that comprise a nucleic acid molecule encoding any of the previously described TFP molecules. In some embodiments, the vector is selected from the group consisting of a DNA, a RNA, a plasmid, a lentivirus vector, adenoviral vector, or a retrovirus vector. In some embodiments, the vector further comprises a promoter. In some embodiments, the vector is an *in vitro* transcribed vector. In some embodiments, a nucleic acid sequence in the vector further comprises a poly(A) tail. In some embodiments, a nucleic acid sequence in the vector further comprises a 3'UTR.

[0094] Also provided herein are cells that comprise any of the described vectors. In some embodiments, the cell is a human T-cell. In some embodiments, the cell is a CD8⁺ or CD4⁺ T-cell. In other embodiments, the cells further comprise a nucleic acid encoding an inhibitory molecule that comprises a first polypeptide that comprises at least a portion of an inhibitory molecule, associated with a second polypeptide that comprises a positive signal from an intracellular signaling domain. In some instances, the inhibitory molecule comprise first polypeptide that comprises at least a portion of PD1 and a second polypeptide comprising a costimulatory domain and primary signaling domain.

[0095] In another aspect, provided herein are isolated TFP molecules that comprise a human or humanized anti-CD19 binding domain, a TCR extracellular domain, a transmembrane domain, and an intracellular signaling domain, wherein the TFP molecule is capable of functionally interacting with an endogenous TCR complex and/or at least one endogenous TCR polypeptide.

[0096] In another aspect, provided herein are isolated TFP molecules that comprise a human or humanized anti-CD19 binding domain, a TCR extracellular domain, a transmembrane domain, and an intracellular signaling domain, wherein the TFP molecule is capable of functionally integrating into an endogenous TCR complex.

[0097] In another aspect, provided herein are human CD8⁺ or CD4⁺ T-cells that comprise at least two TFP molecules, the TFP molecules comprising a human or humanized anti-CD19

binding domain, a TCR extracellular domain, a transmembrane domain, and an intracellular domain, wherein the TFP molecule is capable of functionally interacting with an endogenous TCR complex and/or at least one endogenous TCR polypeptide in, at and/or on the surface of the human CD8⁺ or CD4⁺ T-cell.

[0098] In another aspect, provided herein are protein complexes that comprise i) a TFP molecule comprising a human or humanized anti-CD19 binding domain, a TCR extracellular domain, a transmembrane domain, and an intracellular domain; and ii) at least one endogenous TCR complex.

[0099] In some embodiments, the TCR comprises an extracellular domain or portion thereof of a protein selected from the group consisting of the alpha or beta chain of the T-cell receptor, CD3 delta, CD3 epsilon, or CD3 gamma. In some embodiments, the anti-CD19 binding domain is connected to the TCR extracellular domain by a linker sequence. In some instances, the linker region comprises (G₄S)_n, wherein n=1 to 4. In some instances, the linker sequence comprises a long linker (LL) sequence. In some instances, the long linker sequence comprises (G₄S)_n, wherein n=2 to 4. In some instances, the linker sequence comprises a short linker (SL) sequence. In some instances, the short linker sequence comprises (G₄S)_n, wherein n=1 to 3.

[0100] Also provided herein are human CD8⁺ or CD4⁺ T-cells that comprise at least two different TFP proteins per any of the described protein complexes.

[0101] In another aspect, provided herein is a population of human CD8⁺ or CD4⁺ T-cells, wherein the T-cells of the population individually or collectively comprise at least two TFP molecules, the TFP molecules comprising a human or humanized anti-CD19 or anti-BCMA binding domain, a TCR extracellular domain, a transmembrane domain, and an intracellular domain, wherein the TFP molecule is capable of functionally interacting with an endogenous TCR complex and/or at least one endogenous TCR polypeptide in, at and/or on the surface of the human CD8⁺ or CD4⁺ T-cell.

[0102] In another aspect, provided herein is a population of human CD8⁺ or CD4⁺ T-cells, wherein the T-cells of the population individually or collectively comprise at least two TFP molecules encoded by an isolated nucleic acid molecule provided herein.

[0103] In another aspect, provided herein are methods of making a cell that comprise transducing a T-cell with any of the described vectors.

[0104] In another aspect, provided herein are methods of generating a population of RNA-engineered cells that comprise introducing an *in vitro* transcribed RNA or synthetic RNA into a cell, where the RNA comprises a nucleic acid encoding any of the described TFP molecules.

[0105] In another aspect, provided herein are methods of providing an anti-tumor immunity in a mammal that comprise administering to the mammal an effective amount of a cell expressing

any of the described TFP molecules. In some embodiments, the cell is an autologous T-cell. In some embodiments, the cell is an allogeneic T-cell. In some embodiments, the mammal is a human.

[0106] In another aspect, provided herein are methods of treating a mammal having a disease associated with expression of CD19 that comprise administering to the mammal an effective amount of the cell of comprising any of the described TFP molecules. In some embodiments, the disease associated with CD19 expression is selected from a proliferative disease such as a cancer or malignancy or a precancerous condition such as a myelodysplasia, a myelodysplastic syndrome or a preleukemia, or is a non-cancer related indication associated with expression of CD19. In some embodiments, the disease is a hematologic cancer selected from the group consisting of one or more acute leukemias including but not limited to B-cell acute lymphoid leukemia ("B-ALL"), T-cell acute lymphoid leukemia ("T-ALL"), acute lymphoblastic leukemia (ALL); one or more chronic leukemias including but not limited to chronic myelogenous leukemia (CML), chronic lymphocytic leukemia (CLL); additional hematologic cancers or hematologic conditions including, but not limited to B cell prolymphocytic leukemia, blastic plasmacytoid dendritic cell neoplasm, Burkitt's lymphoma, diffuse large B cell lymphoma, follicular lymphoma, hairy cell leukemia, small cell- or a large cell-follicular lymphoma, malignant lymphoproliferative conditions, MALT lymphoma, mantle cell lymphoma, Marginal zone lymphoma, multiple myeloma, myelodysplasia and myelodysplastic syndrome, non-Hodgkin's lymphoma, plasmablastic lymphoma, plasmacytoid dendritic cell neoplasm, Waldenstrom macroglobulinemia, and "preleukemia" which are a diverse collection of hematological conditions united by ineffective production (or dysplasia) of myeloid blood cells, and to disease associated with CD19 expression include, but not limited to atypical and/or non-classical cancers, malignancies, precancerous conditions or proliferative diseases expressing CD19; and combinations thereof.

[0107] In some embodiments, the cells expressing any of the described TFP molecules are administered in combination with an agent that ameliorates one or more side effects associated with administration of a cell expressing a TFP molecule. In some embodiments, the cells expressing any of the described TFP molecules are administered in combination with an agent that treats the disease associated with CD19.

[0108] Also provided herein are any of the described isolated nucleic acid molecules, any of the described isolated polypeptide molecules, any of the described isolated TFPs, any of the described protein complexes, any of the described vectors or any of the described cells for use as a medicament

Definitions

[0109] Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which the invention pertains.

[0110] The term “a” and “an” refers to one or to more than one (i.e., to at least one) of the grammatical object of the article. By way of example, “an element” means one element or more than one element.

[0111] As used herein, “about” can mean plus or minus less than 1 or 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 25, 30, or greater than 30 percent, depending upon the situation and known or knowable by one skilled in the art.

[0112] As used herein the specification, “subject” or “subjects” or “individuals” may include, but are not limited to, mammals such as humans or non-human mammals, *e.g.*, domesticated, agricultural or wild, animals, as well as birds, and aquatic animals. “Patients” are subjects suffering from or at risk of developing a disease, disorder or condition or otherwise in need of the compositions and methods provided herein.

[0113] As used herein, “treating” or “treatment” refers to any indicia of success in the treatment or amelioration of the disease or condition. Treating can include, for example, reducing, delaying or alleviating the severity of one or more symptoms of the disease or condition, or it can include reducing the frequency with which symptoms of a disease, defect, disorder, or adverse condition, and the like, are experienced by a patient. As used herein, “treat or prevent” is sometimes used herein to refer to a method that results in some level of treatment or amelioration of the disease or condition, and contemplates a range of results directed to that end, including but not restricted to prevention of the condition entirely.

[0114] As used herein, “preventing” refers to the prevention of the disease or condition, *e.g.*, tumor formation, in the patient. For example, if an individual at risk of developing a tumor or other form of cancer is treated with the methods of the present invention and does not later develop the tumor or other form of cancer, then the disease has been prevented, at least over a period of time, in that individual.

[0115] As used herein, a “therapeutically effective amount” is the amount of a composition or an active component thereof sufficient to provide a beneficial effect or to otherwise reduce a detrimental non-beneficial event to the individual to whom the composition is administered. By “therapeutically effective dose” herein is meant a dose that produces one or more desired or desirable (*e.g.*, beneficial) effects for which it is administered, such administration occurring one or more times over a given period of time. The exact dose will depend on the purpose of the treatment, and will be ascertainable by one skilled in the art using known techniques (see, *e.g.*

Lieberman, *Pharmaceutical Dosage Forms* (vols. 1-3, 1992); Lloyd, *The Art, Science and Technology of Pharmaceutical Compounding* (1999); and Pickar, *Dosage Calculations* (1999))

[0116] As used herein, a “T-cell receptor (TCR) fusion protein” or “TFP” includes a recombinant polypeptide derived from the various polypeptides comprising the TCR that is generally capable of i) binding to a surface antigen on target cells and ii) interacting with other polypeptide components of the intact TCR complex, typically when co-located in or on the surface of a T-cell.

[0117] As used herein, the term “CD19” refers to the Cluster of Differentiation 19 protein, which is an antigenic determinant detectable on B cell leukemia precursor cells, other malignant B cells and most cells of the normal B cell lineage. The human and murine amino acid and nucleic acid sequences can be found in a public database, such as GenBank, UniProt and Swiss-Prot. For example, the amino acid sequence of human CD19 can be found as UniProt/Swiss-Prot Accession No. P15391. The human CD19 polypeptide canonical sequence is UniProt Accession No. P15391 (or P15391-1):

MPPPRLLFFLLFLTPMEVRPEEPLVVKVEEGDNAVLQCLKGTSDGPTQQLTWSRESPLK
PFLKLSLGLPGLGIHMRPLAIWLFIFNVSQQMGGFYLCQPGPPSEKAWQPGWTVNVEGS
GELFRWNVSDLGGLGCLKNRSSEGPSSPSGKLMSPKLYVWAKDRPEIWEGEPPCLPPR
DSL NQSL SQDL TMAPGSTLWLSCGVPPDSVSRGPLSWTHVHPKGPKSLLSLELKDDRPA
RDMWVMETGLLLPRATAQDAGKYYCHRGNL TMSFHLEITARPVLWHWLLRTGGWKV
SAVTLAYLIFCLCSLVGILHLQRALVLRKRKRMTDPTRRFFKVTPPPGSGPQNQYGNV
LSLPTPTSGLGRAQRWAAGLGGTAPSYGNPSSDVQADGALGSRPPGVGPEEEEEGEGYE
EPDSEEDSEFYENDSNLGQDQLSQDGSQYENPEDEPLGPEDEDSFSNAESYENEDEELTQ
PVARTMDFLSPHGS AWDP SREATSLGSQSYEDMRGILYAAPQLRSIRGQPGPNHEEDAD
SYENMDNPDGPDPAWGGGGRMGTWSTR (SEQ ID NO: 1).

[0118] The nucleotide sequence encoding of the human CD19 can be found at Accession No. NM001178098. CD19 is expressed on most B lineage cancers, including, *e.g.*, ALL, CLL and non-Hodgkin’s lymphoma (NHL). Other cells that express CD19 are provided below in the definition of “disease associated with expression of CD19.” It is also an early marker of normal B cell progenitors. See, *e.g.*, Nicholson et al. *Mol. Immun.* 34 (16-17): 1157-1165 (1997). In one example, the antigen-binding portion of TFPs recognizes and binds an epitope within the extracellular domain of the CD19 protein as expressed on a malignant and normal B cell.

[0119] As used herein, the term “BCMA” refers to the B-cell maturation antigen also known as tumor necrosis factor receptor superfamily member 17 (TNFRSF17) and Cluster of Differentiation 269 protein (CD269) is a protein that in humans is encoded by the TNFRSF17 gene. TNFRSF17 is a cell surface receptor of the TNF receptor superfamily which recognizes B-

cell activating factor (BAFF) (see, *e.g.*, Laabi et al., EMBO 11 (11): 3897–904 (1992). This receptor is expressed in mature B lymphocytes, and may be important for B-cell development and autoimmune response. The human and murine amino acid and nucleic acid sequences can be found in a public database, such as GenBank, UniProt and Swiss-Prot. For example, the amino acid sequence of human BCMA can be found as UniProt/Swiss-Prot Accession No. Q02223. The human BCMA polypeptide canonical sequence is UniProt Accession No. Q02223 (or Q02223-1):

MLQMAGQCSQNEYFDSLLHACIPCQLRCSSNTPPLTCQRYCNASVTNSVKGTNAILWT
CLGLSLIISLAVFVLMFLLRKINSEPLKDEFKNTGSGLLGMANIDLEKSRTGDEIILPRGLE
YTVEECTCEDCIKSKPKVDSHCFPLPAMEEGATILVTTKTNDYCKSLPAALSATEIEKSI
SAR (SEQ ID NO: 2).

[0120] The nucleotide sequence encoding of the human BCMA can be found at Accession No. NM001192. BCMA is expressed on most B-lineage cancers, including, *e.g.*, leukemia, lymphomas, and multiple myeloma. Other cells that express BCMA are provided below in the definition of “disease associated with expression of BCMA.” This receptor has been shown to specifically bind to the tumor necrosis factor (ligand) superfamily, member 13b (TNFSF13B/TALL-1/BAFF), and to lead to NF-kappaB and MAPK8/JNK activation. This receptor also binds to various TRAF family members, and thus may transduce signals for cell survival and proliferation (see, *e.g.*, Laabi et al., Nucleic Acids Research 22 (7): 1147–54 (1994). In one example, the antigen-binding portion of TFPs recognizes and binds an epitope within the extracellular domain of the BCMA protein as expressed on a malignant and normal B cell.

[0121] The term “antibody,” as used herein, refers to a protein, or polypeptide sequences derived from an immunoglobulin molecule, which specifically binds to an antigen. Antibodies can be intact immunoglobulins of polyclonal or monoclonal origin, or fragments thereof and can be derived from natural or from recombinant sources.

[0122] The terms “antibody fragment” or “antibody binding domain” refer to at least one portion of an antibody, or recombinant variants thereof, that contains the antigen binding domain, i.e., an antigenic determining variable region of an intact antibody, that is sufficient to confer recognition and specific binding of the antibody fragment to a target, such as an antigen and its defined epitope. Examples of antibody fragments include, but are not limited to, Fab, Fab’, F(ab’)₂, and Fv fragments, single-chain (sc)Fv (“scFv”) antibody fragments, linear antibodies, single domain antibodies such as sdAb (either V_L or V_H), camelid V_{HH} domains, and multi-specific antibodies formed from antibody fragments.

[0123] The term “scFv” refers to a fusion protein comprising at least one antibody fragment comprising a variable region of a light chain and at least one antibody fragment comprising a variable region of a heavy chain, wherein the light and heavy chain variable regions are contiguously linked via a short flexible polypeptide linker, and capable of being expressed as a single polypeptide chain, and wherein the scFv retains the specificity of the intact antibody from which it is derived.

[0124] “Heavy chain variable region” or “V_H” with regard to an antibody refers to the fragment of the heavy chain that contains three CDRs interposed between flanking stretches known as framework regions, these framework regions are generally more highly conserved than the CDRs and form a scaffold to support the CDRs.

[0125] Unless specified, as used herein a scFv may have the V_L and V_H variable regions in either order, *e.g.*, with respect to the N-terminal and C-terminal ends of the polypeptide, the scFv may comprise V_L-linker-V_H or may comprise V_H-linker-V_L.

[0126] The portion of the TFP composition of the invention comprising an antibody or antibody fragment thereof may exist in a variety of forms where the antigen binding domain is expressed as part of a contiguous polypeptide chain including, for example, a single domain antibody fragment (sdAb), a single chain antibody (scFv) derived from a murine, humanized or human antibody (Harlow et al., 1999, In: Using Antibodies: A Laboratory Manual, Cold Spring Harbor Laboratory Press, N.Y.; Harlow et al., 1989, In: Antibodies: A Laboratory Manual, Cold Spring Harbor, N.Y.; Houston et al., 1988, Proc. Natl. Acad. Sci. USA 85:5879-5883; Bird et al., 1988, Science 242:423-426). In one aspect, the antigen binding domain of a TFP composition of the invention comprises an antibody fragment. In a further aspect, the TFP comprises an antibody fragment that comprises a scFv or a sdAb.

[0127] The term “antibody heavy chain,” refers to the larger of the two types of polypeptide chains present in antibody molecules in their naturally occurring conformations, and which normally determines the class to which the antibody belongs.

[0128] The term “antibody light chain,” refers to the smaller of the two types of polypeptide chains present in antibody molecules in their naturally occurring conformations. Kappa (“κ”) and lambda (“λ”) light chains refer to the two major antibody light chain isotypes.

[0129] The term “recombinant antibody” refers to an antibody that is generated using recombinant DNA technology, such as, for example, an antibody expressed by a bacteriophage or yeast expression system. The term should also be construed to mean an antibody which has been generated by the synthesis of a DNA molecule encoding the antibody and which DNA molecule expresses an antibody protein, or an amino acid sequence specifying the antibody,

wherein the DNA or amino acid sequence has been obtained using recombinant DNA or amino acid sequence technology which is available and well known in the art.

[0130] The term “antigen” or “Ag” refers to a molecule that is capable of being bound specifically by an antibody, or otherwise provokes an immune response. This immune response may involve either antibody production, or the activation of specific immunologically-competent cells, or both.

[0131] The skilled artisan will understand that any macromolecule, including virtually all proteins or peptides, can serve as an antigen. Furthermore, antigens can be derived from recombinant or genomic DNA. A skilled artisan will understand that any DNA, which comprises a nucleotide sequences or a partial nucleotide sequence encoding a protein that elicits an immune response therefore encodes an “antigen” as that term is used herein. Furthermore, one skilled in the art will understand that an antigen need not be encoded solely by a full length nucleotide sequence of a gene. It is readily apparent that the present invention includes, but is not limited to, the use of partial nucleotide sequences of more than one gene and that these nucleotide sequences are arranged in various combinations to encode polypeptides that elicit the desired immune response. Moreover, a skilled artisan will understand that an antigen need not be encoded by a “gene” at all. It is readily apparent that an antigen can be generated synthesized or can be derived from a biological sample, or might be macromolecule besides a polypeptide. Such a biological sample can include, but is not limited to a tissue sample, a tumor sample, a cell or a fluid with other biological components.

[0132] The term “anti-tumor effect” refers to a biological effect which can be manifested by various means, including but not limited to, *e.g.*, a decrease in tumor volume, a decrease in the number of tumor cells, a decrease in the number of metastases, an increase in life expectancy, decrease in tumor cell proliferation, decrease in tumor cell survival, or amelioration of various physiological symptoms associated with the cancerous condition. An “anti-tumor effect” can also be manifested by the ability of the peptides, polynucleotides, cells and antibodies of the invention in prevention of the occurrence of tumor in the first place.

[0133] The term “autologous” refers to any material derived from the same individual to whom it is later to be re-introduced into the individual.

[0134] The term “allogeneic” refers to any material derived from a different animal of the same species or different patient as the individual to whom the material is introduced. Two or more individuals are said to be allogeneic to one another when the genes at one or more loci are not identical. In some aspects, allogeneic material from individuals of the same species may be sufficiently unlike genetically to interact antigenically.

[0135] The term “xenogeneic” refers to a graft derived from an animal of a different species.

[0136] The term “cancer” refers to a disease characterized by the rapid and uncontrolled growth of aberrant cells. Cancer cells can spread locally or through the bloodstream and lymphatic system to other parts of the body. Examples of various cancers are described herein and include but are not limited to, breast cancer, prostate cancer, ovarian cancer, cervical cancer, skin cancer, pancreatic cancer, colorectal cancer, renal cancer, liver cancer, brain cancer, lymphoma, leukemia, lung cancer and the like.

[0137] The phrase “disease associated with expression of CD19” and “disease associated with expression of BCMA” includes, but is not limited to, a disease associated with expression of CD19 or BCMA or condition associated with cells which express CD19 or BCMA including, *e.g.*, proliferative diseases such as a cancer or malignancy or a precancerous condition such as a myelodysplasia, a myelodysplastic syndrome or a preleukemia; or a noncancer related indication associated with cells which express CD19 or BCMA. In one aspect, a cancer associated with expression of CD19 or BCMA is a hematological cancer. In one aspect, the hematological cancer is a leukemia or a lymphoma. In one aspect, a cancer associated with expression of CD19 or BCMA includes cancers and malignancies including, but not limited to, *e.g.*, one or more acute leukemias including but not limited to, *e.g.*, B cell ALL, T-cell acute lymphoid leukemia (TALL), one or more chronic leukemias including but not limited to, *e.g.*, CLL or chronic myelogenous leukemia (CML). Additional cancers or hematologic conditions associated with expression of CD19 comprise, but are not limited to, *e.g.*, B cell prolymphocytic leukemia, blastic plasmacytoid dendritic cell neoplasm, Burkitt’s lymphoma, diffuse large B cell lymphoma, Follicular lymphoma, Hairy cell leukemia, small cell- or a large cell-follicular lymphoma, malignant lymphoproliferative conditions, MALT lymphoma, mantle cell lymphoma, Marginal zone lymphoma, multiple myeloma, myelodysplasia and myelodysplastic syndrome, non-Hodgkin’s lymphoma, plasmablastic lymphoma, plasmacytoid dendritic cell neoplasm, Waldenstrom macroglobulinemia, and “preleukemia” which are a diverse collection of hematological conditions united by ineffective production (or dysplasia) of myeloid blood cells, and the like. Further diseases associated with expression of CD19 or BCMA expression include, but are not limited to, *e.g.*, atypical and/or non-classical cancers, malignancies, precancerous conditions or proliferative diseases associated with expression of CD19 or BCMA. Non-cancer related indications associated with expression of CD19 or BCMA include, but are not limited to, *e.g.*, autoimmune disease, (*e.g.*, lupus, rheumatoid arthritis, colitis), inflammatory disorders (allergy and asthma), and transplantation.

[0138] The term “conservative sequence modifications” refers to amino acid modifications that do not significantly affect or alter the binding characteristics of the antibody or antibody fragment containing the amino acid sequence. Such conservative modifications include amino

acid substitutions, additions and deletions. Modifications can be introduced into an antibody or antibody fragment of the invention by standard techniques known in the art, such as site-directed mutagenesis and PCR-mediated mutagenesis. Conservative amino acid substitutions are ones in which the amino acid residue is replaced with an amino acid residue having a similar side chain. Families of amino acid residues having similar side chains have been defined in the art. These families include amino acids with basic side chains (*e.g.*, lysine, arginine, histidine), acidic side chains (*e.g.*, aspartic acid, glutamic acid), uncharged polar side chains (*e.g.*, glycine, asparagine, glutamine, serine, threonine, tyrosine, cysteine, tryptophan), nonpolar side chains (*e.g.*, alanine, valine, leucine, isoleucine, proline, phenylalanine, methionine), beta-branched side chains (*e.g.*, threonine, valine, isoleucine) and aromatic side chains (*e.g.*, tyrosine, phenylalanine, tryptophan, histidine). Thus, one or more amino acid residues within a TFP of the invention can be replaced with other amino acid residues from the same side chain family and the altered TFP can be tested using the functional assays described herein.

[0139] The term “stimulation” refers to a primary response induced by binding of a stimulatory domain or stimulatory molecule (*e.g.*, a TCR/CD3 complex) with its cognate ligand thereby mediating a signal transduction event, such as, but not limited to, signal transduction via the TCR/CD3 complex. Stimulation can mediate altered expression of certain molecules, and/or reorganization of cytoskeletal structures, and the like.

[0140] The term “stimulatory molecule” or “stimulatory domain” refers to a molecule or portion thereof expressed by a T-cell that provides the primary cytoplasmic signaling sequence(s) that regulate primary activation of the TCR complex in a stimulatory way for at least some aspect of the T-cell signaling pathway. In one aspect, the primary signal is initiated by, for instance, binding of a TCR/CD3 complex with an MHC molecule loaded with peptide, and which leads to mediation of a T-cell response, including, but not limited to, proliferation, activation, differentiation, and the like. A primary cytoplasmic signaling sequence (also referred to as a “primary signaling domain”) that acts in a stimulatory manner may contain a signaling motif which is known as immunoreceptor tyrosine-based activation motif or “ITAM”. Examples of an ITAM containing primary cytoplasmic signaling sequence that is of particular use in the invention includes, but is not limited to, those derived from TCR zeta, FcR gamma, FcR beta, CD3 gamma, CD3 delta, CD3 epsilon, CD5, CD22, CD79a, CD79b, CD278 (also known as “ICOS”) and CD66d.

[0141] The term “antigen presenting cell” or “APC” refers to an immune system cell such as an accessory cell (*e.g.*, a B-cell, a dendritic cell, and the like) that displays a foreign antigen complexed with major histocompatibility complexes (MHC's) on its surface. T-cells may

recognize these complexes using their T-cell receptors (TCRs). APCs process antigens and present them to T-cells.

[0142] An “intracellular signaling domain,” as the term is used herein, refers to an intracellular portion of a molecule. The intracellular signaling domain generates a signal that promotes an immune effector function of the TFP containing cell, *e.g.*, a TFP-expressing T-cell. Examples of immune effector function, *e.g.*, in a TFP-expressing T-cell, include cytolytic activity and T helper cell activity, including the secretion of cytokines. In an embodiment, the intracellular signaling domain can comprise a primary intracellular signaling domain. Exemplary primary intracellular signaling domains include those derived from the molecules responsible for primary stimulation, or antigen dependent stimulation. In an embodiment, the intracellular signaling domain can comprise a costimulatory intracellular domain. Exemplary costimulatory intracellular signaling domains include those derived from molecules responsible for costimulatory signals, or antigen independent stimulation.

[0143] A primary intracellular signaling domain can comprise an ITAM (“immunoreceptor tyrosine-based activation motif”). Examples of ITAM containing primary cytoplasmic signaling sequences include, but are not limited to, those derived from CD3 zeta, FcR gamma, FcR beta, CD3 gamma, CD3 delta, CD3 epsilon, CD5, CD22, CD79a, CD79b, and CD66d DAP10 and DAP12.

[0144] The term “costimulatory molecule” refers to the cognate binding partner on a T-cell that specifically binds with a costimulatory ligand, thereby mediating a costimulatory response by the T-cell, such as, but not limited to, proliferation. Costimulatory molecules are cell surface molecules other than antigen receptors or their ligands that are required for an efficient immune response. Costimulatory molecules include, but are not limited to an MHC class 1 molecule, BTLA and a Toll ligand receptor, as well as OX40, CD2, CD27, CD28, CDS, ICAM-1, LFA-1 (CD11a/CD18) and 4-1BB (CD137). A costimulatory intracellular signaling domain can be the intracellular portion of a costimulatory molecule. A costimulatory molecule can be represented in the following protein families: TNF receptor proteins, Immunoglobulin-like proteins, cytokine receptors, integrins, signaling lymphocytic activation molecules (SLAM proteins), and activating NK cell receptors. Examples of such molecules include CD27, CD28, 4-1BB (CD137), OX40, GITR, CD30, CD40, ICOS, BAFFR, HVEM, lymphocyte function-associated antigen-1 (LFA-1), CD2, CD7, LIGHT, NKG2C, SLAMF7, NKp80, CD160, B7-H3, and a ligand that specifically binds with CD83, and the like. The intracellular signaling domain can comprise the entire intracellular portion, or the entire native intracellular signaling domain, of the molecule from which it is derived, or a functional fragment thereof. The term “4-1BB” refers to a member of the TNFR superfamily with an amino acid sequence provided as GenBank Acc.

No. AAA62478.2, or the equivalent residues from a non-human species, *e.g.*, mouse, rodent, monkey, ape and the like; and a “4-1BB costimulatory domain” is defined as amino acid residues 214-255 of GenBank Acc. No. AAA62478.2, or the equivalent residues from a non-human species, *e.g.*, mouse, rodent, monkey, ape and the like.

[0145] The term “encoding” refers to the inherent property of specific sequences of nucleotides in a polynucleotide, such as a gene, a cDNA, or an mRNA, to serve as templates for synthesis of other polymers and macromolecules in biological processes having either a defined sequence of nucleotides (*e.g.*, rRNA, tRNA and mRNA) or a defined sequence of amino acids and the biological properties resulting therefrom. Thus, a gene, cDNA, or RNA, encodes a protein if transcription and translation of mRNA corresponding to that gene produces the protein in a cell or other biological system. Both the coding strand, the nucleotide sequence of which is identical to the mRNA sequence and is usually provided in sequence listings, and the non-coding strand, used as the template for transcription of a gene or cDNA, can be referred to as encoding the protein or other product of that gene or cDNA.

[0146] Unless otherwise specified, a “nucleotide sequence encoding an amino acid sequence” includes all nucleotide sequences that are degenerate versions of each other and that encode the same amino acid sequence. The phrase nucleotide sequence that encodes a protein or an RNA may also include introns to the extent that the nucleotide sequence encoding the protein may in some version contain one or more introns.

[0147] The term “effective amount” or “therapeutically effective amount” are used interchangeably herein, and refer to an amount of a compound, formulation, material, or composition, as described herein effective to achieve a particular biological or therapeutic result.

[0148] The term “endogenous” refers to any material from or produced inside an organism, cell, tissue or system.

[0149] The term “exogenous” refers to any material introduced from or produced outside an organism, cell, tissue or system.

[0150] The term “expression” refers to the transcription and/or translation of a particular nucleotide sequence driven by a promoter.

[0151] The term “transfer vector” refers to a composition of matter which comprises an isolated nucleic acid and which can be used to deliver the isolated nucleic acid to the interior of a cell. Numerous vectors are known in the art including, but not limited to, linear polynucleotides, polynucleotides associated with ionic or amphiphilic compounds, plasmids, and viruses. Thus, the term “transfer vector” includes an autonomously replicating plasmid or a virus. The term should also be construed to further include non-plasmid and non-viral compounds which facilitate transfer of nucleic acid into cells, such as, for example, a polylysine compound,

liposome, and the like. Examples of viral transfer vectors include, but are not limited to, adenoviral vectors, adeno-associated virus vectors, retroviral vectors, lentiviral vectors, and the like.

[0152] The term “expression vector” refers to a vector comprising a recombinant polynucleotide comprising expression control sequences operatively linked to a nucleotide sequence to be expressed. An expression vector comprises sufficient cis-acting elements for expression; other elements for expression can be supplied by the host cell or in an *in vitro* expression system. Expression vectors include all those known in the art, including cosmids, plasmids (*e.g.*, naked or contained in liposomes) and viruses (*e.g.*, lentiviruses, retroviruses, adenoviruses, and adeno-associated viruses) that incorporate the recombinant polynucleotide.

[0153] The term “lentivirus” refers to a genus of the Retroviridae family. Lentiviruses are unique among the retroviruses in being able to infect non-dividing cells; they can deliver a significant amount of genetic information into the DNA of the host cell, so they are one of the most efficient methods of a gene delivery vector. HIV, SIV, and FIV are all examples of lentiviruses.

[0154] The term “lentiviral vector” refers to a vector derived from at least a portion of a lentivirus genome, including especially a self-inactivating lentiviral vector as provided in Milone et al., Mol. Ther. 17(8): 1453-1464 (2009). Other examples of lentivirus vectors that may be used in the clinic, include but are not limited to, *e.g.*, the LENTIVECTOR™ gene delivery technology from Oxford BioMedica, the LENTIMAX™ vector system from Lentigen, and the like. Nonclinical types of lentiviral vectors are also available and would be known to one skilled in the art.

[0155] The term “homologous” or “identity” refers to the subunit sequence identity between two polymeric molecules, *e.g.*, between two nucleic acid molecules, such as, two DNA molecules or two RNA molecules, or between two polypeptide molecules. When a subunit position in both of the two molecules is occupied by the same monomeric subunit; *e.g.*, if a position in each of two DNA molecules is occupied by adenine, then they are homologous or identical at that position. The homology between two sequences is a direct function of the number of matching or homologous positions; *e.g.*, if half (*e.g.*, five positions in a polymer ten subunits in length) of the positions in two sequences are homologous, the two sequences are 50% homologous; if 90% of the positions (*e.g.*, 9 of 10), are matched or homologous, the two sequences are 90% homologous.

[0156] “Humanized” forms of non-human (*e.g.*, murine) antibodies are chimeric immunoglobulins, immunoglobulin chains or fragments thereof (such as Fv, Fab, Fab', F(ab')₂ or other antigen-binding subsequences of antibodies) which contain minimal sequence derived

from non-human immunoglobulin. For the most part, humanized antibodies and antibody fragments thereof are human immunoglobulins (recipient antibody or antibody fragment) in which residues from a complementary-determining region (CDR) of the recipient are replaced by residues from a CDR of a non-human species (donor antibody) such as mouse, rat or rabbit having the desired specificity, affinity, and capacity. In some instances, Fv framework region (FR) residues of the human immunoglobulin are replaced by corresponding non-human residues. Furthermore, a humanized antibody/antibody fragment can comprise residues which are found neither in the recipient antibody nor in the imported CDR or framework sequences. These modifications can further refine and optimize antibody or antibody fragment performance. In general, the humanized antibody or antibody fragment thereof will comprise substantially all of at least one, and typically two, variable domains, in which all or substantially all of the CDR regions correspond to those of a non-human immunoglobulin and all or a significant portion of the FR regions are those of a human immunoglobulin sequence. The humanized antibody or antibody fragment can also comprise at least a portion of an immunoglobulin constant region (Fc), typically that of a human immunoglobulin. For further details, see Jones et al., *Nature*, 321: 522-525, 1986; Reichmann et al., *Nature*, 332: 323-329, 1988; Presta, *Curr. Op. Struct. Biol.*, 2: 593-596, 1992.

[0157] “Human” or “fully human” refers to an immunoglobulin, such as an antibody or antibody fragment, where the whole molecule is of human origin or consists of an amino acid sequence identical to a human form of the antibody or immunoglobulin.

[0158] The term “isolated” means altered or removed from the natural state. For example, a nucleic acid or a peptide naturally present in a living animal is not “isolated,” but the same nucleic acid or peptide partially or completely separated from the coexisting materials of its natural state is “isolated.” An isolated nucleic acid or protein can exist in substantially purified form, or can exist in a non-native environment such as, for example, a host cell.

[0159] In the context of the present invention, the following abbreviations for the commonly occurring nucleic acid bases are used. “A” refers to adenosine, “C” refers to cytosine, “G” refers to guanosine, “T” refers to thymidine, and “U” refers to uridine.

[0160] The term “operably linked” or “transcriptional control” refers to functional linkage between a regulatory sequence and a heterologous nucleic acid sequence resulting in expression of the latter. For example, a first nucleic acid sequence is operably linked with a second nucleic acid sequence when the first nucleic acid sequence is placed in a functional relationship with the second nucleic acid sequence. For instance, a promoter is operably linked to a coding sequence if the promoter affects the transcription or expression of the coding sequence. Operably linked

DNA sequences can be contiguous with each other and, *e.g.*, where necessary to join two protein coding regions, are in the same reading frame.

[0161] The term “parenteral” administration of an immunogenic composition includes, *e.g.*, subcutaneous (s.c.), intravenous (i.v.), intramuscular (i.m.), or intrasternal injection, intratumoral, or infusion techniques.

[0162] The term “nucleic acid” or “polynucleotide” refers to deoxyribonucleic acids (DNA) or ribonucleic acids (RNA) and polymers thereof in either single- or double-stranded form. Unless specifically limited, the term encompasses nucleic acids containing known analogues of natural nucleotides that have similar binding properties as the reference nucleic acid and are metabolized in a manner similar to naturally occurring nucleotides. Unless otherwise indicated, a particular nucleic acid sequence also implicitly encompasses conservatively modified variants thereof (*e.g.*, degenerate codon substitutions), alleles, orthologs, SNPs, and complementary sequences as well as the sequence explicitly indicated. Specifically, degenerate codon substitutions may be achieved by generating sequences in which the third position of one or more selected (or all) codons is substituted with mixed-base and/or deoxyinosine residues (Batzner et al., *Nucleic Acid Res.* 19:5081 (1991); Ohtsuka et al., *J. Biol. Chem.* 260:2605-2608 (1985); and Rossolini et al., *Mol. Cell. Probes* 8:91-98 (1994)).

[0163] The terms “peptide,” “polypeptide,” and “protein” are used interchangeably, and refer to a compound comprised of amino acid residues covalently linked by peptide bonds. A protein or peptide must contain at least two amino acids, and no limitation is placed on the maximum number of amino acids that can comprise a protein’s or peptide’s sequence. Polypeptides include any peptide or protein comprising two or more amino acids joined to each other by peptide bonds. As used herein, the term refers to both short chains, which also commonly are referred to in the art as peptides, oligopeptides and oligomers, for example, and to longer chains, which generally are referred to in the art as proteins, of which there are many types. “Polypeptides” include, for example, biologically active fragments, substantially homologous polypeptides, oligopeptides, homodimers, heterodimers, variants of polypeptides, modified polypeptides, derivatives, analogs, fusion proteins, among others. A polypeptide includes a natural peptide, a recombinant peptide, or a combination thereof.

[0164] The term “promoter” refers to a DNA sequence recognized by the transcription machinery of the cell, or introduced synthetic machinery, required to initiate the specific transcription of a polynucleotide sequence.

[0165] The term “promoter/regulatory sequence” refers to a nucleic acid sequence which is required for expression of a gene product operably linked to the promoter/regulatory sequence. In some instances, this sequence may be the core promoter sequence and in other instances, this

sequence may also include an enhancer sequence and other regulatory elements which are required for expression of the gene product. The promoter/regulatory sequence may, for example, be one which expresses the gene product in a tissue specific manner.

[0166] The term “constitutive” promoter refers to a nucleotide sequence which, when operably linked with a polynucleotide which encodes or specifies a gene product, causes the gene product to be produced in a cell under most or all physiological conditions of the cell.

[0167] The term “inducible” promoter refers to a nucleotide sequence which, when operably linked with a polynucleotide which encodes or specifies a gene product, causes the gene product to be produced in a cell substantially only when an inducer which corresponds to the promoter is present in the cell.

[0168] The term “tissue-specific” promoter refers to a nucleotide sequence which, when operably linked with a polynucleotide encodes or specified by a gene, causes the gene product to be produced in a cell substantially only if the cell is a cell of the tissue type corresponding to the promoter.

[0169] The terms “linker” and “flexible polypeptide linker” as used in the context of a scFv refers to a peptide linker that consists of amino acids such as glycine and/or serine residues used alone or in combination, to link variable heavy and variable light chain regions together. In one embodiment, the flexible polypeptide linker is a Gly/Ser linker and comprises the amino acid sequence (Gly-Gly-Gly-Ser)_n, where n is a positive integer equal to or greater than 1. For example, n=1, n=2, n=3, n=4, n=5, n=6, n=7, n=8, n=9 and n=10. In one embodiment, the flexible polypeptide linkers include, but are not limited to, (Gly₄Ser)₄ or (Gly₄Ser)₃. In another embodiment, the linkers include multiple repeats of (Gly₂Ser), (GlySer) or (Gly₃Ser). Also included within the scope of the invention are linkers described in WO2012/138475 (incorporated herein by reference). In some instances, the linker sequence comprises a long linker (LL) sequence. In some instances, the long linker sequence comprises (G₄S)_n, wherein n=2 to 4. In some instances, the linker sequence comprises a short linker (SL) sequence. In some instances, the short linker sequence comprises (G₄S)_n, wherein n=1 to 3.

[0170] As used herein, a 5' cap (also termed an RNA cap, an RNA 7-methylguanosine cap or an RNA m⁷G cap) is a modified guanine nucleotide that has been added to the “front” or 5' end of a eukaryotic messenger RNA shortly after the start of transcription. The 5' cap consists of a terminal group which is linked to the first transcribed nucleotide. Its presence is critical for recognition by the ribosome and protection from RNases. Cap addition is coupled to transcription, and occurs co-transcriptionally, such that each influences the other. Shortly after the start of transcription, the 5' end of the mRNA being synthesized is bound by a cap-synthesizing complex associated with RNA polymerase. This enzymatic complex catalyzes the

chemical reactions that are required for mRNA capping. Synthesis proceeds as a multi-step biochemical reaction. The capping moiety can be modified to modulate functionality of mRNA such as its stability or efficiency of translation.

[0171] As used herein, “*in vitro* transcribed RNA” refers to RNA, preferably mRNA, which has been synthesized *in vitro*. Generally, the *in vitro* transcribed RNA is generated from an *in vitro* transcription vector. The *in vitro* transcription vector comprises a template that is used to generate the *in vitro* transcribed RNA.

[0172] As used herein, a “poly(A)” is a series of adenosines attached by polyadenylation to the mRNA. In the preferred embodiment of a construct for transient expression, the polyA is between 50 and 5000, preferably greater than 64, more preferably greater than 100, most preferably greater than 300 or 400. Poly(A) sequences can be modified chemically or enzymatically to modulate mRNA functionality such as localization, stability or efficiency of translation.

[0173] As used herein, “polyadenylation” refers to the covalent linkage of a polyadenylyl moiety, or its modified variant, to a messenger RNA molecule. In eukaryotic organisms, most messenger RNA (mRNA) molecules are polyadenylated at the 3’ end. The 3’ poly(A) tail is a long sequence of adenine nucleotides (often several hundred) added to the pre-mRNA through the action of an enzyme, polyadenylate polymerase. In higher eukaryotes, the poly(A) tail is added onto transcripts that contain a specific sequence, the polyadenylation signal. The poly(A) tail and the protein bound to it aid in protecting mRNA from degradation by exonucleases. Polyadenylation is also important for transcription termination, export of the mRNA from the nucleus, and translation. Polyadenylation occurs in the nucleus immediately after transcription of DNA into RNA, but additionally can also occur later in the cytoplasm. After transcription has been terminated, the mRNA chain is cleaved through the action of an endonuclease complex associated with RNA polymerase. The cleavage site is usually characterized by the presence of the base sequence AAUAAA near the cleavage site. After the mRNA has been cleaved, adenosine residues are added to the free 3’ end at the cleavage site.

[0174] As used herein, “transient” refers to expression of a non-integrated transgene for a period of hours, days or weeks, wherein the period of time of expression is less than the period of time for expression of the gene if integrated into the genome or contained within a stable plasmid replicon in the host cell.

[0175] The term “signal transduction pathway” refers to the biochemical relationship between a variety of signal transduction molecules that play a role in the transmission of a signal from one portion of a cell to another portion of a cell. The phrase “cell surface receptor” includes

molecules and complexes of molecules capable of receiving a signal and transmitting signal across the membrane of a cell.

[0176] The term “subject” is intended to include living organisms in which an immune response can be elicited (*e.g.*, mammals, human).

[0177] The term, a “substantially purified” cell refers to a cell that is essentially free of other cell types. A substantially purified cell also refers to a cell which has been separated from other cell types with which it is normally associated in its naturally occurring state. In some instances, a population of substantially purified cells refers to a homogenous population of cells. In other instances, this term refers simply to cell that have been separated from the cells with which they are naturally associated in their natural state. In some aspects, the cells are cultured *in vitro*. In other aspects, the cells are not cultured *in vitro*.

[0178] The term “therapeutic” as used herein means a treatment. A therapeutic effect is obtained by reduction, suppression, remission, or eradication of a disease state.

[0179] The term “prophylaxis” as used herein means the prevention of or protective treatment for a disease or disease state.

[0180] In the context of the present invention, “tumor antigen” or “hyperproliferative disorder antigen” or “antigen associated with a hyperproliferative disorder” refers to antigens that are common to specific hyperproliferative disorders. In certain aspects, the hyperproliferative disorder antigens of the present invention are derived from, cancers including but not limited to primary or metastatic melanoma, thymoma, lymphoma, sarcoma, lung cancer, liver cancer, NHL, leukemias, uterine cancer, cervical cancer, bladder cancer, kidney cancer and adenocarcinomas such as breast cancer, prostate cancer, ovarian cancer, pancreatic cancer, and the like.

[0181] The term “transfected” or “transformed” or “transduced” refers to a process by which exogenous nucleic acid is transferred or introduced into the host cell. A “transfected” or “transformed” or “transduced” cell is one which has been transfected, transformed or transduced with exogenous nucleic acid. The cell includes the primary subject cell and its progeny.

[0182] The term “specifically binds,” refers to an antibody, an antibody fragment or a specific ligand, which recognizes and binds a cognate binding partner (*e.g.*, CD19) present in a sample, but which does not necessarily and substantially recognize or bind other molecules in the sample.

[0183] Ranges: throughout this disclosure, various aspects of the invention can be presented in a range format. It should be understood that the description in range format is merely for convenience and brevity and should not be construed as an inflexible limitation on the scope of the invention. Accordingly, the description of a range should be considered to have specifically

disclosed all the possible subranges as well as individual numerical values within that range. For example, description of a range such as from 1 to 6 should be considered to have specifically disclosed subranges such as from 1 to 3, from 1 to 4, from 1 to 5, from 2 to 4, from 2 to 6, from 3 to 6 etc., as well as individual numbers within that range, for example, 1, 2, 2.7, 3, 4, 5, 5.3, and 6. As another example, a range such as 95-99% identity, includes something with 95%, 96%, 97%, 98% or 99% identity, and includes subranges such as 96-99%, 96-98%, 96-97%, 97-99%, 97-98% and 98-99% identity. This applies regardless of the breadth of the range.

DESCRIPTION

[0184] Provided herein are compositions of matter and methods of use for the treatment of a disease such as cancer, using T-cell receptor (TCR) fusion proteins. As used herein, a “T-cell receptor (TCR) fusion protein” or “TFP” includes a recombinant polypeptide derived from the various polypeptides comprising the TCR that is generally capable of i) binding to a surface antigen on target cells and ii) interacting with other polypeptide components of the intact TCR complex, typically when co-located in or on the surface of a T-cell. As provided herein, TFPs provide substantial benefits as compared to Chimeric Antigen Receptors. The term “Chimeric Antigen Receptor” or alternatively a “CAR” refers to a recombinant polypeptide comprising an extracellular antigen binding domain in the form of a scFv, a transmembrane domain, and cytoplasmic signaling domains (also referred to herein as “an intracellular signaling domains”) comprising a functional signaling domain derived from a stimulatory molecule as defined below. Generally, the central intracellular signaling domain of a CAR is derived from the CD3 zeta chain that is normally found associated with the TCR complex. The CD3 zeta signaling domain can be fused with one or more functional signaling domains derived from at least one co-stimulatory molecule such as 4-1BB (i.e., CD137), CD27 and/or CD28.

T-cell receptor (TCR) fusion proteins (TFP)

[0185] The present invention encompasses recombinant DNA constructs encoding TFPs, wherein the TFP comprises an antibody fragment that binds specifically to CD19, *e.g.*, human CD19, wherein the sequence of the antibody fragment is contiguous with and in the same reading frame as a nucleic acid sequence encoding a TCR subunit or portion thereof. The present invention encompasses recombinant DNA constructs encoding TFPs, wherein the TFP comprises an antibody fragment that binds specifically to BCMA, *e.g.*, human BCMA, wherein the sequence of the antibody fragment is contiguous with and in the same reading frame as a nucleic acid sequence encoding a TCR subunit or portion thereof. The TFPs provided herein are able to associate with one or more endogenous (or alternatively, one or more exogenous, or a combination of endogenous and exogenous) TCR subunits in order to form a functional TCR complex.

[0186] In one aspect, the TFP of the invention comprises a target-specific binding element otherwise referred to as an antigen binding domain. The choice of moiety depends upon the type and number of target antigen that define the surface of a target cell. For example, the antigen binding domain may be chosen to recognize a target antigen that acts as a cell surface marker on target cells associated with a particular disease state. Thus examples of cell surface markers that may act as target antigens for the antigen binding domain in a TFP of the invention include those associated with viral, bacterial and parasitic infections; autoimmune diseases; and cancerous diseases (*e.g.*, malignant diseases).

[0187] In one aspect, the TFP-mediated T-cell response can be directed to an antigen of interest by way of engineering an antigen-binding domain into the TFP that specifically binds a desired antigen.

[0188] In one aspect, the portion of the TFP comprising the antigen binding domain comprises an antigen binding domain that targets CD19. In one aspect, the antigen binding domain targets human CD19. In one aspect, the portion of the TFP comprising the antigen binding domain comprises an antigen binding domain that targets BCMA. In one aspect, the antigen binding domain targets human BCMA.

[0189] The antigen binding domain can be any domain that binds to the antigen including but not limited to a monoclonal antibody, a polyclonal antibody, a recombinant antibody, a human antibody, a humanized antibody, and a functional fragment thereof, including but not limited to a single-domain antibody such as a heavy chain variable domain (V_H), a light chain variable domain (V_L) and a variable domain (V_{HH}) of a camelid derived nanobody, and to an alternative scaffold known in the art to function as antigen binding domain, such as a recombinant fibronectin domain, anticalin, DARPIN and the like. Likewise a natural or synthetic ligand specifically recognizing and binding the target antigen can be used as antigen binding domain for the TFP. In some instances, it is beneficial for the antigen binding domain to be derived from the same species in which the TFP will ultimately be used in. For example, for use in humans, it may be beneficial for the antigen binding domain of the TFP to comprise human or humanized residues for the antigen binding domain of an antibody or antibody fragment.

[0190] Thus, in one aspect, the antigen-binding domain comprises a humanized or human antibody or an antibody fragment, or a murine antibody or antibody fragment. In one embodiment, the humanized or human anti-CD19 or anti-BCMA binding domain comprises one or more (*e.g.*, all three) light chain complementary determining region 1 (LC CDR1), light chain complementary determining region 2 (LC CDR2), and light chain complementary determining region 3 (LC CDR3) of a humanized or human anti-CD19 or anti-BCMA binding domain described herein, and/or one or more (*e.g.*, all three) heavy chain complementary determining

region 1 (HC CDR1), heavy chain complementary determining region 2 (HC CDR2), and heavy chain complementary determining region 3 (HC CDR3) of a humanized or human anti-CD19 binding domain described herein, *e.g.*, a humanized or human anti-CD19 or anti-BCMA binding domain comprising one or more, *e.g.*, all three, LC CDRs and one or more, *e.g.*, all three, HC CDRs. In one embodiment, the humanized or human anti-CD19 binding domain comprises one or more (*e.g.*, all three) heavy chain complementary determining region 1 (HC CDR1), heavy chain complementary determining region 2 (HC CDR2), and heavy chain complementary determining region 3 (HC CDR3) of a humanized or human anti-CD19 or anti-BCMA binding domain described herein, *e.g.*, the humanized or human anti-CD19 or anti-BCMA binding domain has two variable heavy chain regions, each comprising a HC CDR1, a HC CDR2 and a HC CDR3 described herein. In one embodiment, the humanized or human anti-CD19 or anti-BCMA binding domain comprises a humanized or human light chain variable region described herein and/or a humanized or human heavy chain variable region described herein. In one embodiment, the humanized or human anti-CD19 or anti-BCMA binding domain comprises a humanized heavy chain variable region described herein, *e.g.*, at least two humanized or human heavy chain variable regions described herein. In one embodiment, the anti-CD19 or anti-BCMA binding domain is a scFv comprising a light chain and a heavy chain of an amino acid sequence provided herein. In an embodiment, the anti-CD19 or anti-BCMA binding domain (*e.g.*, a scFv) comprises: a light chain variable region comprising an amino acid sequence having at least one, two or three modifications (*e.g.*, substitutions) but not more than 30, 20 or 10 modifications (*e.g.*, substitutions) of an amino acid sequence of a light chain variable region provided herein, or a sequence with 95-99% identity with an amino acid sequence provided herein; and/or a heavy chain variable region comprising an amino acid sequence having at least one, two or three modifications (*e.g.*, substitutions) but not more than 30, 20 or 10 modifications (*e.g.*, substitutions) of an amino acid sequence of a heavy chain variable region provided herein, or a sequence with 95-99% identity to an amino acid sequence provided herein. In one embodiment, the humanized or human anti-CD19 or anti-BCMA binding domain is a scFv, and a light chain variable region comprising an amino acid sequence described herein, is attached to a heavy chain variable region comprising an amino acid sequence described herein, via a linker, *e.g.*, a linker described herein. In one embodiment, the humanized anti-CD19 or anti-BCMA binding domain includes a (Gly₄-Ser)_n linker, wherein n is 1, 2, 3, 4, 5, or 6, preferably 3 or 4. The light chain variable region and heavy chain variable region of a scFv can be, *e.g.*, in any of the following orientations: light chain variable region-linker-heavy chain variable region or heavy chain variable region-linker-light chain variable region. In some instances, the linker sequence comprises a long linker (LL) sequence. In some instances, the long linker sequence

comprises $(G_4S)_n$, wherein $n=2$ to 4. In some instances, the linker sequence comprises a short linker (SL) sequence. In some instances, the short linker sequence comprises $(G_4S)_n$, wherein $n=1$ to 3.

[0191] In some aspects, a non-human antibody is humanized, where specific sequences or regions of the antibody are modified to increase similarity to an antibody naturally produced in a human or fragment thereof. In one aspect, the antigen binding domain is humanized.

[0192] A humanized antibody can be produced using a variety of techniques known in the art, including but not limited to, CDR-grafting (see, *e.g.*, European Patent No. EP 239,400; International Publication No. WO 91/09967; and U.S. Pat. Nos. 5,225,539, 5,530,101, and 5,585,089, each of which is incorporated herein in its entirety by reference), veneering or resurfacing (see, *e.g.*, European Patent Nos. EP 592,106 and EP 519,596; Padlan, 1991, *Molecular Immunology*, 28(4/5):489-498; Studnicka et al., 1994, *Protein Engineering*, 7(6):805-814; and Roguska et al., 1994, *PNAS*, 91:969-973, each of which is incorporated herein by its entirety by reference), chain shuffling (see, *e.g.*, U.S. Pat. No. 5,565,332, which is incorporated herein in its entirety by reference), and techniques disclosed in, *e.g.*, U.S. Patent Application Publication No. US2005/0042664, U.S. Patent Application Publication No. US2005/0048617, U.S. Pat. No. 6,407,213, U.S. Pat. No. 5,766,886, International Publication No. WO 9317105, Tan et al., *J. Immunol.*, 169:1119-25 (2002), Caldas et al., *Protein Eng.*, 13(5):353-60 (2000), Morea et al., *Methods*, 20(3):267-79 (2000), Baca et al., *J. Biol. Chem.*, 272(16):10678-84 (1997), Roguska et al., *Protein Eng.*, 9(10):895-904 (1996), Couto et al., *Cancer Res.*, 55 (23 Supp):5973s-5977s (1995), Couto et al., *Cancer Res.*, 55(8):1717-22 (1995), Sandhu J S, *Gene*, 150(2):409-10 (1994), and Pedersen et al., *J. Mol. Biol.*, 235(3):959-73 (1994), each of which is incorporated herein in its entirety by reference. Often, framework residues in the framework regions will be substituted with the corresponding residue from the CDR donor antibody to alter, for example improve, antigen binding. These framework substitutions are identified by methods well-known in the art, *e.g.*, by modeling of the interactions of the CDR and framework residues to identify framework residues important for antigen binding and sequence comparison to identify unusual framework residues at particular positions (see, *e.g.*, Queen et al., U.S. Pat. No. 5,585,089; and Riechmann et al., 1988, *Nature*, 332:323, which are incorporated herein by reference in their entireties.)

[0193] A humanized antibody or antibody fragment has one or more amino acid residues remaining in it from a source which is nonhuman. These nonhuman amino acid residues are often referred to as “import” residues, which are typically taken from an “import” variable domain. As provided herein, humanized antibodies or antibody fragments comprise one or more CDRs from nonhuman immunoglobulin molecules and framework regions wherein the amino

acid residues comprising the framework are derived completely or mostly from human germline. Multiple techniques for humanization of antibodies or antibody fragments are well-known in the art and can essentially be performed following the method of Winter and co-workers (Jones et al., *Nature*, 321:522-525 (1986); Riechmann et al., *Nature*, 332:323-327 (1988); Verhoeyen et al., *Science*, 239:1534-1536 (1988)), by substituting rodent CDRs or CDR sequences for the corresponding sequences of a human antibody, i.e., CDR-grafting (EP 239,400; PCT Publication No. WO 91/09967; and U.S. Pat. Nos. 4,816,567; 6,331,415; 5,225,539; 5,530,101; 5,585,089; 6,548,640, the contents of which are incorporated herein by reference in their entirety). In such humanized antibodies and antibody fragments, substantially less than an intact human variable domain has been substituted by the corresponding sequence from a nonhuman species.

Humanized antibodies are often human antibodies in which some CDR residues and possibly some framework (FR) residues are substituted by residues from analogous sites in rodent antibodies. Humanization of antibodies and antibody fragments can also be achieved by veneering or resurfacing (EP 592,106; EP 519,596; Padlan, 1991, *Molecular Immunology*, 28(4/5):489-498; Studnicka et al., *Protein Engineering*, 7(6):805-814 (1994); and Roguska et al., *PNAS*, 91:969-973 (1994)) or chain shuffling (U.S. Pat. No. 5,565,332), the contents of which are incorporated herein by reference in their entirety.

[0194] The choice of human variable domains, both light and heavy, to be used in making the humanized antibodies is to reduce antigenicity. According to the so-called “best-fit” method, the sequence of the variable domain of a rodent antibody is screened against the entire library of known human variable-domain sequences. The human sequence which is closest to that of the rodent is then accepted as the human framework (FR) for the humanized antibody (Sims et al., *J. Immunol.*, 151:2296 (1993); Chothia et al., *J. Mol. Biol.*, 196:901 (1987), the contents of which are incorporated herein by reference herein in their entirety). Another method uses a particular framework derived from the consensus sequence of all human antibodies of a particular subgroup of light or heavy chains. The same framework may be used for several different humanized antibodies (see, *e.g.*, Nicholson et al. *Mol. Immun.* 34 (16-17): 1157-1165 (1997); Carter et al., *Proc. Natl. Acad. Sci. USA*, 89:4285 (1992); Presta et al., *J. Immunol.*, 151:2623 (1993), the contents of which are incorporated herein by reference herein in their entirety). In some embodiments, the framework region, *e.g.*, all four framework regions, of the heavy chain variable region are derived from a V_H4-4-59 germline sequence. In one embodiment, the framework region can comprise, one, two, three, four or five modifications, *e.g.*, substitutions, *e.g.*, from the amino acid at the corresponding murine sequence. In one embodiment, the framework region, *e.g.*, all four framework regions of the light chain variable region are derived from a VK3-1.25 germline sequence. In one embodiment, the framework region can comprise,

one, two, three, four or five modifications, *e.g.*, substitutions, *e.g.*, from the amino acid at the corresponding murine sequence.

[0195] In some aspects, the portion of a TFP composition of the invention that comprises an antibody fragment is humanized with retention of high affinity for the target antigen and other favorable biological properties. According to one aspect of the invention, humanized antibodies and antibody fragments are prepared by a process of analysis of the parental sequences and various conceptual humanized products using three-dimensional models of the parental and humanized sequences. Three-dimensional immunoglobulin models are commonly available and are familiar to those skilled in the art. Computer programs are available which illustrate and display probable three-dimensional conformational structures of selected candidate immunoglobulin sequences. Inspection of these displays permits analysis of the likely role of the residues in the functioning of the candidate immunoglobulin sequence, *e.g.*, the analysis of residues that influence the ability of the candidate immunoglobulin to bind the target antigen. In this way, FR residues can be selected and combined from the recipient and import sequences so that the desired antibody or antibody fragment characteristic, such as increased affinity for the target antigen, is achieved. In general, the CDR residues are directly and most substantially involved in influencing antigen binding.

[0196] A humanized antibody or antibody fragment may retain a similar antigenic specificity as the original antibody, *e.g.*, in the present invention, the ability to bind human CD19. In some embodiments, a humanized antibody or antibody fragment may have improved affinity and/or specificity of binding to human CD19 or human BCMA.

[0197] In one aspect, the anti-CD19 or anti-BCMA binding domain is characterized by particular functional features or properties of an antibody or antibody fragment. For example, in one aspect, the portion of a TFP composition of the invention that comprises an antigen binding domain specifically binds human CD19 or human BCMA. In one aspect, the antigen binding domain has the same or a similar binding specificity to human CD19 as the FMC63 scFv described in Nicholson et al. Mol. Immun. 34 (16-17): 1157-1165 (1997). In one aspect, the invention relates to an antigen binding domain comprising an antibody or antibody fragment, wherein the antibody binding domain specifically binds to a CD19 or BCMA protein or fragment thereof, wherein the antibody or antibody fragment comprises a variable light chain and/or a variable heavy chain that includes an amino acid sequence provided herein. In certain aspects, the scFv is contiguous with and in the same reading frame as a leader sequence.

[0198] In one aspect, the anti-CD19 or anti-BCMA binding domain is a fragment, *e.g.*, a single chain variable fragment (scFv). In one aspect, the anti-CD19 binding domain is a Fv, a Fab, a (Fab')₂, or a bi-functional (*e.g.* bi-specific) hybrid antibody (*e.g.*, Lanzavecchia et al., Eur. J.

Immunol. 17, 105 (1987)). In one aspect, the antibodies and fragments thereof of the invention binds a CD19 protein with wild-type or enhanced affinity.

[0199] Also provided herein are methods for obtaining an antibody antigen binding domain specific for a target antigen (*e.g.*, CD19, BCMA or any target antigen described elsewhere herein for targets of fusion moiety binding domains), the method comprising providing by way of addition, deletion, substitution or insertion of one or more amino acids in the amino acid sequence of a V_H domain set out herein a V_H domain which is an amino acid sequence variant of the V_H domain, optionally combining the V_H domain thus provided with one or more V_L domains, and testing the V_H domain or V_H/V_L combination or combinations to identify a specific binding member or an antibody antigen binding domain specific for a target antigen of interest (*e.g.*, CD19 or BCMA) and optionally with one or more desired properties.

[0200] In some instances, V_H domains and scFvs can be prepared according to method known in the art (see, for example, Bird et al., (1988) Science 242:423-426 and Huston et al., (1988) Proc. Natl. Acad. Sci. USA 85:5879-5883). scFv molecules can be produced by linking V_H and V_L regions together using flexible polypeptide linkers. The scFv molecules comprise a linker (*e.g.*, a Ser-Gly linker) with an optimized length and/or amino acid composition. The linker length can greatly affect how the variable regions of a scFv fold and interact. In fact, if a short polypeptide linker is employed (*e.g.*, between 5-10 amino acids) intra-chain folding is prevented. Inter-chain folding is also required to bring the two variable regions together to form a functional epitope binding site. In some instances, the linker sequence comprises a long linker (LL) sequence. In some instances, the long linker sequence comprises $(G_4S)_n$, wherein $n=2$ to 4. In some instances, the linker sequence comprises a short linker (SL) sequence. In some instances, the short linker sequence comprises $(G_4S)_n$, wherein $n=1$ to 3. For examples of linker orientation and size see, *e.g.*, Hollinger et al. 1993 Proc Natl Acad. Sci. U.S.A. 90:6444-6448, U.S. Patent Application Publication Nos. 2005/0100543, 2005/0175606, 2007/0014794, and PCT publication Nos. WO2006/020258 and WO2007/024715, is incorporated herein by reference.

[0201] A scFv can comprise a linker of about 10, 11, 12, 13, 14, 15 or greater than 15 residues between its V_L and V_H regions. The linker sequence may comprise any naturally occurring amino acid. In some embodiments, the linker sequence comprises amino acids glycine and serine. In another embodiment, the linker sequence comprises sets of glycine and serine repeats such as $(Gly_4Ser)_n$, where n is a positive integer equal to or greater than 1. In one embodiment, the linker can be $(Gly_4Ser)_4$ or $(Gly_4Ser)_3$. Variation in the linker length may retain or enhance activity, giving rise to superior efficacy in activity studies. In some instances, the linker sequence comprises a long linker (LL) sequence. In some instances, the long linker sequence comprises $(G_4S)_n$, wherein $n=2$ to 4. In some instances, the linker sequence comprises a short

linker (SL) sequence. In some instances, the short linker sequence comprises $(G_4S)_n$, wherein $n=1$ to 3.

Stability and Mutations

[0202] The stability of an anti-CD19 or anti-BCMA binding domain, *e.g.*, scFv molecules (*e.g.*, soluble scFv) can be evaluated in reference to the biophysical properties (*e.g.*, thermal stability) of a conventional control scFv molecule or a full length antibody. In one embodiment, the humanized or human scFv has a thermal stability that is greater than about 0.1, about 0.25, about 0.5, about 0.75, about 1, about 1.25, about 1.5, about 1.75, about 2, about 2.5, about 3, about 3.5, about 4, about 4.5, about 5, about 5.5, about 6, about 6.5, about 7, about 7.5, about 8, about 8.5, about 9, about 9.5, about 10 degrees, about 11 degrees, about 12 degrees, about 13 degrees, about 14 degrees, or about 15 degrees Celsius than a parent scFv in the described assays.

[0203] The improved thermal stability of the anti-CD19 or anti-BCMA binding domain, *e.g.*, scFv is subsequently conferred to the entire CD19-TFP construct, leading to improved therapeutic properties of the anti-CD19 or anti-BCMA TFP construct. The thermal stability of the anti-CD19 or anti-BCMA binding domain, *e.g.*, scFv can be improved by at least about 2 °C or 3 °C as compared to a conventional antibody. In one embodiment, the anti-CD19 or anti-BCMA binding domain, *e.g.*, scFv has a 1 °C improved thermal stability as compared to a conventional antibody. In another embodiment, the anti-CD19 binding domain, *e.g.*, scFv has a 2 °C improved thermal stability as compared to a conventional antibody. In another embodiment, the scFv has a 4 °C, 5 °C, 6 °C, 7 °C, 8 °C, 9 °C, 10 °C, 11 °C, 12 °C, 13 °C, 14 °C, or 15 °C improved thermal stability as compared to a conventional antibody. Comparisons can be made, for example, between the scFv molecules disclosed herein and scFv molecules or Fab fragments of an antibody from which the scFv V_H and V_L were derived. Thermal stability can be measured using methods known in the art. For example, in one embodiment, T_M can be measured. Methods for measuring T_M and other methods of determining protein stability are described in more detail below.

[0204] Mutations in scFv (arising through humanization or direct mutagenesis of the soluble scFv) alter the stability of the scFv and improve the overall stability of the scFv and the anti-CD19 or anti-BCMA TFP construct. Stability of the humanized scFv is compared against the murine scFv using measurements such as T_M , temperature denaturation and temperature aggregation. In one embodiment, the anti-CD19 or anti-BCMA binding domain, *e.g.*, a scFv, comprises at least one mutation arising from the humanization process such that the mutated scFv confers improved stability to the Anti-CD19 TFP construct. In another embodiment, the anti-CD19 binding domain, *e.g.*, scFv comprises at least 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 mutations

arising from the humanization process such that the mutated scFv confers improved stability to the CD19-TFP or BCMA-TFP construct.

[0205] In one aspect, the antigen binding domain of the TFP comprises an amino acid sequence that is homologous to an antigen binding domain amino acid sequence described herein, and the antigen binding domain retains the desired functional properties of the anti-CD19 or anti-BCMA antibody fragments described herein. In one specific aspect, the TFP composition of the invention comprises an antibody fragment. In a further aspect, that antibody fragment comprises a scFv.

[0206] In various aspects, the antigen binding domain of the TFP is engineered by modifying one or more amino acids within one or both variable regions (*e.g.*, V_H and/or V_L), for example within one or more CDR regions and/or within one or more framework regions. In one specific aspect, the TFP composition of the invention comprises an antibody fragment. In a further aspect, that antibody fragment comprises a scFv.

[0207] It will be understood by one of ordinary skill in the art that the antibody or antibody fragment of the invention may further be modified such that they vary in amino acid sequence (*e.g.*, from wild-type), but not in desired activity. For example, additional nucleotide substitutions leading to amino acid substitutions at “non-essential” amino acid residues may be made to the protein. For example, a nonessential amino acid residue in a molecule may be replaced with another amino acid residue from the same side chain family. In another embodiment, a string of amino acids can be replaced with a structurally similar string that differs in order and/or composition of side chain family members, *e.g.*, a conservative substitution, in which an amino acid residue is replaced with an amino acid residue having a similar side chain, may be made.

[0208] Families of amino acid residues having similar side chains have been defined in the art, including basic side chains (*e.g.*, lysine, arginine, histidine), acidic side chains (*e.g.*, aspartic acid, glutamic acid), uncharged polar side chains (*e.g.*, glycine, asparagine, glutamine, serine, threonine, tyrosine, cysteine), nonpolar side chains (*e.g.*, alanine, valine, leucine, isoleucine, proline, phenylalanine, methionine, tryptophan), beta-branched side chains (*e.g.*, threonine, valine, isoleucine) and aromatic side chains (*e.g.*, tyrosine, phenylalanine, tryptophan, histidine).

[0209] Percent identity in the context of two or more nucleic acids or polypeptide sequences refers to two or more sequences that are the same. Two sequences are “substantially identical” if two sequences have a specified percentage of amino acid residues or nucleotides that are the same (*e.g.*, 60% identity, optionally 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% identity over a specified region, or, when not specified, over the entire

sequence), when compared and aligned for maximum correspondence over a comparison window, or designated region as measured using one of the following sequence comparison algorithms or by manual alignment and visual inspection. Optionally, the identity exists over a region that is at least about 50 nucleotides (or 10 amino acids) in length, or more preferably over a region that is 100 to 500 or 1000 or more nucleotides (or 20, 50, 200 or more amino acids) in length.

[0210] For sequence comparison, typically one sequence acts as a reference sequence, to which test sequences are compared. When using a sequence comparison algorithm, test and reference sequences are entered into a computer, subsequence coordinates are designated, if necessary, and sequence algorithm program parameters are designated. Default program parameters can be used, or alternative parameters can be designated. The sequence comparison algorithm then calculates the percent sequence identities for the test sequences relative to the reference sequence, based on the program parameters. Methods of alignment of sequences for comparison are well known in the art. Optimal alignment of sequences for comparison can be conducted, *e.g.*, by the local homology algorithm of Smith and Waterman, (1970) *Adv. Appl. Math.* 2:482c, by the homology alignment algorithm of Needleman and Wunsch, (1970) *J. Mol. Biol.* 48:443, by the search for similarity method of Pearson and Lipman, (1988) *Proc. Nat'l. Acad. Sci. USA* 85:2444, by computerized implementations of these algorithms (GAP, BESTFIT, FASTA, and TFASTA in the Wisconsin Genetics Software Package, Genetics Computer Group, 575 Science Dr., Madison, Wis.), or by manual alignment and visual inspection (see, *e.g.*, Brent et al., (2003) *Current Protocols in Molecular Biology*). Two examples of algorithms that are suitable for determining percent sequence identity and sequence similarity are the BLAST and BLAST 2.0 algorithms, which are described in Altschul et al., (1977) *Nuc. Acids Res.* 25:3389-3402; and Altschul et al., (1990) *J. Mol. Biol.* 215:403-410, respectively. Software for performing BLAST analyses is publicly available through the National Center for Biotechnology Information.

[0211] In one aspect, the present invention contemplates modifications of the starting antibody or fragment (*e.g.*, scFv) amino acid sequence that generate functionally equivalent molecules. For example, the V_H or V_L of an anti-CD19 or anti-BCMA binding domain, *e.g.*, scFv, comprised in the TFP can be modified to retain at least about 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% identity of the starting V_H or V_L framework region of the anti-CD19 binding domain, *e.g.*, scFv. The present invention contemplates modifications of the entire TFP construct, *e.g.*, modifications in one or more amino acid sequences of the various domains of the TFP construct in order to generate functionally equivalent molecules. The TFP construct can be modified to retain at least about 70%, 71%, 72%, 73%, 74%, 75%,

76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% identity of the starting TFP construct.

Extracellular domain

[0212] The extracellular domain may be derived either from a natural or from a recombinant source. Where the source is natural, the domain may be derived from any protein, but in particular a membrane-bound or transmembrane protein. In one aspect the extracellular domain is capable of associating with the transmembrane domain. An extracellular domain of particular use in this invention may include at least the extracellular region(s) of *e.g.*, the alpha, beta or zeta chain of the T-cell receptor, or CD3 epsilon, CD3 gamma, or CD3 delta, or in alternative embodiments, CD28, CD45, CD4, CD5, CD8, CD9, CD16, CD22, CD33, CD37, CD64, CD80, CD86, CD134, CD137, CD154.

Transmembrane Domain

[0213] In general, a TFP sequence contains an extracellular domain and a transmembrane domain encoded by a single genomic sequence. In alternative embodiments, a TFP can be designed to comprise a transmembrane domain that is heterologous to the extracellular domain of the TFP. A transmembrane domain can include one or more additional amino acids adjacent to the transmembrane region, *e.g.*, one or more amino acid associated with the extracellular region of the protein from which the transmembrane was derived (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or up to 15 amino acids of the extracellular region) and/or one or more additional amino acids associated with the intracellular region of the protein from which the transmembrane protein is derived (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or up to 15 amino acids of the intracellular region). In one aspect, the transmembrane domain is one that is associated with one of the other domains of the TFP is used. In some instances, the transmembrane domain can be selected or modified by amino acid substitution to avoid binding of such domains to the transmembrane domains of the same or different surface membrane proteins, *e.g.*, to minimize interactions with other members of the receptor complex. In one aspect, the transmembrane domain is capable of homodimerization with another TFP on the TFP-T-cell surface. In a different aspect the amino acid sequence of the transmembrane domain may be modified or substituted so as to minimize interactions with the binding domains of the native binding partner present in the same TFP.

[0214] The transmembrane domain may be derived either from a natural or from a recombinant source. Where the source is natural, the domain may be derived from any membrane-bound or transmembrane protein. In one aspect the transmembrane domain is capable of signaling to the intracellular domain(s) whenever the TFP has bound to a target. A transmembrane domain of particular use in this invention may include at least the transmembrane region(s) of *e.g.*, the

alpha, beta or zeta chain of the T-cell receptor, CD28, CD3 epsilon, CD45, CD4, CD5, CD8, CD9, CD16, CD22, CD33, CD37, CD64, CD80, CD86, CD134, CD137, CD154.

[0215] In some instances, the transmembrane domain can be attached to the extracellular region of the TFP, *e.g.*, the antigen binding domain of the TFP, via a hinge, *e.g.*, a hinge from a human protein. For example, in one embodiment, the hinge can be a human immunoglobulin (Ig) hinge, *e.g.*, an IgG4 hinge, or a CD8a hinge.

Linkers

[0216] Optionally, a short oligo- or polypeptide linker, between 2 and 10 amino acids in length may form the linkage between the transmembrane domain and the cytoplasmic region of the TFP. A glycine-serine doublet provides a particularly suitable linker. For example, in one aspect, the linker comprises the amino acid sequence of GGGGSGGGGS (SEQ ID NO: 3). In some embodiments, the linker is encoded by a nucleotide sequence of GGTGGCGGAGGTTCTGGAGGTGGAGGTTCC (SEQ ID NO: 4).

Cytoplasmic Domain

[0217] The cytoplasmic domain of the TFP can include an intracellular signaling domain, if the TFP contains CD3 gamma, delta or epsilon polypeptides; TCR alpha and TCR beta subunits are generally lacking in a signaling domain. An intracellular signaling domain is generally responsible for activation of at least one of the normal effector functions of the immune cell in which the TFP has been introduced. The term “effector function” refers to a specialized function of a cell. Effector function of a T-cell, for example, may be cytolytic activity or helper activity including the secretion of cytokines. Thus the term “intracellular signaling domain” refers to the portion of a protein which transduces the effector function signal and directs the cell to perform a specialized function. While usually the entire intracellular signaling domain can be employed, in many cases it is not necessary to use the entire chain. To the extent that a truncated portion of the intracellular signaling domain is used, such truncated portion may be used in place of the intact chain as long as it transduces the effector function signal. The term intracellular signaling domain is thus meant to include any truncated portion of the intracellular signaling domain sufficient to transduce the effector function signal.

[0218] Examples of intracellular signaling domains for use in the TFP of the invention include the cytoplasmic sequences of the T-cell receptor (TCR) and co-receptors that act in concert to initiate signal transduction following antigen receptor engagement, as well as any derivative or variant of these sequences and any recombinant sequence that has the same functional capability.

[0219] It is known that signals generated through the TCR alone are insufficient for full activation of naive T-cells and that a secondary and/or costimulatory signal is required. Thus,

naïve T-cell activation can be said to be mediated by two distinct classes of cytoplasmic signaling sequences: those that initiate antigen-dependent primary activation through the TCR (primary intracellular signaling domains) and those that act in an antigen-independent manner to provide a secondary or costimulatory signal (secondary cytoplasmic domain, *e.g.*, a costimulatory domain).

[0220] A primary signaling domain regulates primary activation of the TCR complex either in a stimulatory way, or in an inhibitory way. Primary intracellular signaling domains that act in a stimulatory manner may contain signaling motifs which are known as immunoreceptor tyrosine-based activation motifs (ITAMs).

[0221] Examples of ITAMs containing primary intracellular signaling domains that are of particular use in the invention include those of CD3 zeta, FcR gamma, FcR beta, CD3 gamma, CD3 delta, CD3 epsilon, CD5, CD22, CD79a, CD79b, and CD66d. In one embodiment, a TFP of the invention comprises an intracellular signaling domain, *e.g.*, a primary signaling domain of CD3-epsilon. In one embodiment, a primary signaling domain comprises a modified ITAM domain, *e.g.*, a mutated ITAM domain which has altered (*e.g.*, increased or decreased) activity as compared to the native ITAM domain. In one embodiment, a primary signaling domain comprises a modified ITAM-containing primary intracellular signaling domain, *e.g.*, an optimized and/or truncated ITAM-containing primary intracellular signaling domain. In an embodiment, a primary signaling domain comprises one, two, three, four or more ITAM motifs.

[0222] The intracellular signaling domain of the TFP can comprise the CD3 zeta signaling domain by itself or it can be combined with any other desired intracellular signaling domain(s) useful in the context of a TFP of the invention. For example, the intracellular signaling domain of the TFP can comprise a CD3 epsilon chain portion and a costimulatory signaling domain. The costimulatory signaling domain refers to a portion of the TFP comprising the intracellular domain of a costimulatory molecule. A costimulatory molecule is a cell surface molecule other than an antigen receptor or its ligands that is required for an efficient response of lymphocytes to an antigen. Examples of such molecules include CD27, CD28, 4-1BB (CD137), OX40, CD30, CD40, PD1, ICOS, lymphocyte function-associated antigen-1 (LFA-1), CD2, CD7, LIGHT, NKG2C, B7-H3, and a ligand that specifically binds with CD83, and the like. For example, CD27 costimulation has been demonstrated to enhance expansion, effector function, and survival of human TFP-T-cells *in vitro* and augments human T-cell persistence and antitumor activity *in vivo* (Song et al. Blood. 2012; 119(3):696-706).

[0223] The intracellular signaling sequences within the cytoplasmic portion of the TFP of the invention may be linked to each other in a random or specified order. Optionally, a short oligo-

or polypeptide linker, for example, between 2 and 10 amino acids (*e.g.*, 2, 3, 4, 5, 6, 7, 8, 9, or 10 amino acids) in length may form the linkage between intracellular signaling sequences.

[0224] In one embodiment, a glycine-serine doublet can be used as a suitable linker. In one embodiment, a single amino acid, *e.g.*, an alanine, a glycine, can be used as a suitable linker.

[0225] In one aspect, the TFP-expressing cell described herein can further comprise a second TFP, *e.g.*, a second TFP that includes a different antigen binding domain, *e.g.*, to the same target (CD19 or BCMA) or a different target (*e.g.*, CD123). In one embodiment, when the TFP-expressing cell comprises two or more different TFPs, the antigen binding domains of the different TFPs can be such that the antigen binding domains do not interact with one another. For example, a cell expressing a first and second TFP can have an antigen binding domain of the first TFP, *e.g.*, as a fragment, *e.g.*, a scFv, that does not form an association with the antigen binding domain of the second TFP, *e.g.*, the antigen binding domain of the second TFP is a V_{HH} .

[0226] In another aspect, the TFP-expressing cell described herein can further express another agent, *e.g.*, an agent which enhances the activity of a TFP-expressing cell. For example, in one embodiment, the agent can be an agent which inhibits an inhibitory molecule. Inhibitory molecules, *e.g.*, PD1, can, in some embodiments, decrease the ability of a TFP-expressing cell to mount an immune effector response. Examples of inhibitory molecules include PD1, PD-L1, CTLA4, TIM3, LAG3, VISTA, BTLA, TIGIT, LAIR1, CD160, 2B4 and TGFR beta. In one embodiment, the agent which inhibits an inhibitory molecule comprises a first polypeptide, *e.g.*, an inhibitory molecule, associated with a second polypeptide that provides a positive signal to the cell, *e.g.*, an intracellular signaling domain described herein. In one embodiment, the agent comprises a first polypeptide, *e.g.*, of an inhibitory molecule such as PD1, LAG3, CTLA4, CD160, BTLA, LAIR1, TIM3, 2B4 and TIGIT, or a fragment of any of these (*e.g.*, at least a portion of an extracellular domain of any of these), and a second polypeptide which is an intracellular signaling domain described herein (*e.g.*, comprising a costimulatory domain (*e.g.*, 4-1BB, CD27 or CD28, *e.g.*, as described herein) and/or a primary signaling domain (*e.g.*, a CD3 zeta signaling domain described herein). In one embodiment, the agent comprises a first polypeptide of PD1 or a fragment thereof (*e.g.*, at least a portion of an extracellular domain of PD1), and a second polypeptide of an intracellular signaling domain described herein (*e.g.*, a CD28 signaling domain described herein and/or a CD3 zeta signaling domain described herein). PD1 is an inhibitory member of the CD28 family of receptors that also includes CD28, CTLA-4, ICOS, and BTLA. PD-1 is expressed on activated B cells, T-cells and myeloid cells (Agata et al. 1996 Int. Immunol 8:765-75). Two ligands for PD1, PD-L1 and PD-L2 have been shown to downregulate T-cell activation upon binding to PD1 (Freeman et al. 2000 J Exp Med 192:1027-34; Latchman et al. 2001 Nat Immunol 2:261-8; Carter et al. 2002 Eur J Immunol 32:634-43).

PD-L1 is abundant in human cancers (Dong et al. 2003 J Mol Med 81:281-7; Blank et al. 2005 Cancer Immunol. Immunother 54:307-314; Konishi et al. 2004 Clin Cancer Res 10:5094). Immune suppression can be reversed by inhibiting the local interaction of PD1 with PD-L1.

[0227] In one embodiment, the agent comprises the extracellular domain (ECD) of an inhibitory molecule, *e.g.*, Programmed Death 1 (PD1) can be fused to a transmembrane domain and optionally an intracellular signaling domain such as 41BB and CD3 zeta (also referred to herein as a PD1 TFP). In one embodiment, the PD1 TFP, when used in combinations with an anti-CD19 TFP described herein, improves the persistence of the T-cell. In one embodiment, the TFP is a PD1 TFP comprising the extracellular domain of PD 1. Alternatively, provided are TFPs containing an antibody or antibody fragment such as a scFv that specifically binds to the Programmed Death-Ligand 1 (PD-L1) or Programmed Death-Ligand 2 (PD-L2).

[0228] In another aspect, the present invention provides a population of TFP-expressing T-cells, *e.g.*, TFP-T-cells. In some embodiments, the population of TFP-expressing T-cells comprises a mixture of cells expressing different TFPs. For example, in one embodiment, the population of TFP-T-cells can include a first cell expressing a TFP having an anti-CD19 or anti-BCMA binding domain described herein, and a second cell expressing a TFP having a different anti-CD19 or anti-BCMA binding domain, *e.g.*, an anti-CD19 or anti-BCMA binding domain described herein that differs from the anti-CD19 binding domain in the TFP expressed by the first cell. As another example, the population of TFP-expressing cells can include a first cell expressing a TFP that includes an anti-CD19 or anti-BCMA binding domain, *e.g.*, as described herein, and a second cell expressing a TFP that includes an antigen binding domain to a target other than CD19 or BCMA (*e.g.*, another tumor-associated antigen).

[0229] In another aspect, the present invention provides a population of cells wherein at least one cell in the population expresses a TFP having an anti-CD19 or anti-BCMA domain described herein, and a second cell expressing another agent, *e.g.*, an agent which enhances the activity of a TFP-expressing cell. For example, in one embodiment, the agent can be an agent which inhibits an inhibitory molecule. Inhibitory molecules, *e.g.*, can, in some embodiments, decrease the ability of a TFP-expressing cell to mount an immune effector response. Examples of inhibitory molecules include PD1, PD-L1, PD-L2, CTLA4, TIM3, LAG3, VISTA, BTLA, TIGIT, LAIR1, CD160, 2B4 and TGFR beta. In one embodiment, the agent that inhibits an inhibitory molecule comprises a first polypeptide, *e.g.*, an inhibitory molecule, associated with a second polypeptide that provides a positive signal to the cell, *e.g.*, an intracellular signaling domain described herein.

[0230] Disclosed herein are methods for producing *in vitro* transcribed RNA encoding TFPs. The present invention also includes a TFP encoding RNA construct that can be directly

transfected into a cell. A method for generating mRNA for use in transfection can involve *in vitro* transcription (IVT) of a template with specially designed primers, followed by polyA addition, to produce a construct containing 3' and 5' untranslated sequence ("UTR"), a 5' cap and/or Internal Ribosome Entry Site (IRES), the nucleic acid to be expressed, and a polyA tail, typically 50-2000 bases in length. RNA so produced can efficiently transfect different kinds of cells. In one aspect, the template includes sequences for the TFP.

[0231] In one aspect the anti-CD19 or anti-BCMA TFP is encoded by a messenger RNA (mRNA). In one aspect the mRNA encoding the anti-CD19 or anti-BCMA TFP is introduced into a T-cell for production of a TFP-T-cell. In one embodiment, the *in vitro* transcribed RNA TFP can be introduced to a cell as a form of transient transfection. The RNA is produced by *in vitro* transcription using a polymerase chain reaction (PCR)-generated template. DNA of interest from any source can be directly converted by PCR into a template for *in vitro* mRNA synthesis using appropriate primers and RNA polymerase. The source of the DNA can be, for example, genomic DNA, plasmid DNA, phage DNA, cDNA, synthetic DNA sequence or any other appropriate source of DNA. The desired template for *in vitro* transcription is a TFP of the present invention. In one embodiment, the DNA to be used for PCR contains an open reading frame. The DNA can be from a naturally occurring DNA sequence from the genome of an organism. In one embodiment, the nucleic acid can include some or all of the 5' and/or 3' untranslated regions (UTRs). The nucleic acid can include exons and introns. In one embodiment, the DNA to be used for PCR is a human nucleic acid sequence. In another embodiment, the DNA to be used for PCR is a human nucleic acid sequence including the 5' and 3' UTRs. The DNA can alternatively be an artificial DNA sequence that is not normally expressed in a naturally occurring organism. An exemplary artificial DNA sequence is one that contains portions of genes that are ligated together to form an open reading frame that encodes a fusion protein. The portions of DNA that are ligated together can be from a single organism or from more than one organism.

[0232] PCR is used to generate a template for *in vitro* transcription of mRNA which is used for transfection. Methods for performing PCR are well known in the art. Primers for use in PCR are designed to have regions that are substantially complementary to regions of the DNA to be used as a template for the PCR. "Substantially complementary," as used herein, refers to sequences of nucleotides where a majority or all of the bases in the primer sequence are complementary, or one or more bases are non-complementary, or mismatched. Substantially complementary sequences are able to anneal or hybridize with the intended DNA target under annealing conditions used for PCR. The primers can be designed to be substantially complementary to any portion of the DNA template. For example, the primers can be designed to amplify the portion

of a nucleic acid that is normally transcribed in cells (the open reading frame), including 5' and 3' UTRs. The primers can also be designed to amplify a portion of a nucleic acid that encodes a particular domain of interest. In one embodiment, the primers are designed to amplify the coding region of a human cDNA, including all or portions of the 5' and 3' UTRs. Primers useful for PCR can be generated by synthetic methods that are well known in the art. "Forward primers" are primers that contain a region of nucleotides that are substantially complementary to nucleotides on the DNA template that are upstream of the DNA sequence that is to be amplified. "Upstream" is used herein to refer to a location 5' to the DNA sequence to be amplified relative to the coding strand. "Reverse primers" are primers that contain a region of nucleotides that are substantially complementary to a double-stranded DNA template that are downstream of the DNA sequence that is to be amplified. "Downstream" is used herein to refer to a location 3' to the DNA sequence to be amplified relative to the coding strand.

[0233] Any DNA polymerase useful for PCR can be used in the methods disclosed herein. The reagents and polymerase are commercially available from a number of sources.

[0234] Chemical structures with the ability to promote stability and/or translation efficiency may also be used. The RNA preferably has 5' and 3' UTRs. In one embodiment, the 5' UTR is between one and 3000 nucleotides in length. The length of 5' and 3' UTR sequences to be added to the coding region can be altered by different methods, including, but not limited to, designing primers for PCR that anneal to different regions of the UTRs. Using this approach, one of ordinary skill in the art can modify the 5' and 3' UTR lengths required to achieve optimal translation efficiency following transfection of the transcribed RNA.

[0235] The 5' and 3' UTRs can be the naturally occurring, endogenous 5' and 3' UTRs for the nucleic acid of interest. Alternatively, UTR sequences that are not endogenous to the nucleic acid of interest can be added by incorporating the UTR sequences into the forward and reverse primers or by any other modifications of the template. The use of UTR sequences that are not endogenous to the nucleic acid of interest can be useful for modifying the stability and/or translation efficiency of the RNA. For example, it is known that AU-rich elements in 3'UTR sequences can decrease the stability of mRNA. Therefore, 3' UTRs can be selected or designed to increase the stability of the transcribed RNA based on properties of UTRs that are well known in the art.

[0236] In one embodiment, the 5' UTR can contain the Kozak sequence of the endogenous nucleic acid. Alternatively, when a 5' UTR that is not endogenous to the nucleic acid of interest is being added by PCR as described above, a consensus Kozak sequence can be redesigned by adding the 5' UTR sequence. Kozak sequences can increase the efficiency of translation of some RNA transcripts, but does not appear to be required for all RNAs to enable efficient translation.

The requirement for Kozak sequences for many mRNAs is known in the art. In other embodiments the 5' UTR can be 5'UTR of an RNA virus whose RNA genome is stable in cells. In other embodiments various nucleotide analogues can be used in the 3' or 5' UTR to impede exonuclease degradation of the mRNA.

[0237] To enable synthesis of RNA from a DNA template without the need for gene cloning, a promoter of transcription should be attached to the DNA template upstream of the sequence to be transcribed. When a sequence that functions as a promoter for an RNA polymerase is added to the 5' end of the forward primer, the RNA polymerase promoter becomes incorporated into the PCR product upstream of the open reading frame that is to be transcribed. In one preferred embodiment, the promoter is a T7 polymerase promoter, as described elsewhere herein. Other useful promoters include, but are not limited to, T3 and SP6 RNA polymerase promoters. Consensus nucleotide sequences for T7, T3 and SP6 promoters are known in the art.

[0238] In a preferred embodiment, the mRNA has both a cap on the 5' end and a 3' poly(A) tail which determine ribosome binding, initiation of translation and stability mRNA in the cell. On a circular DNA template, for instance, plasmid DNA, RNA polymerase produces a long concatameric product which is not suitable for expression in eukaryotic cells. The transcription of plasmid DNA linearized at the end of the 3' UTR results in normal sized mRNA which is not effective in eukaryotic transfection even if it is polyadenylated after transcription.

[0239] On a linear DNA template, phage T7 RNA polymerase can extend the 3' end of the transcript beyond the last base of the template (Schenborn and Mierendorf, Nuc Acids Res., 13:6223-36 (1985); Nacheva and Berzal-Herranz, Eur. J. Biochem., 270:1485-65 (2003)).

[0240] The conventional method of integration of polyA/T stretches into a DNA template is molecular cloning. However polyA/T sequence integrated into plasmid DNA can cause plasmid instability, which is why plasmid DNA templates obtained from bacterial cells are often highly contaminated with deletions and other aberrations. This makes cloning procedures not only laborious and time consuming but often not reliable. That is why a method which allows construction of DNA templates with polyA/T 3' stretch without cloning highly desirable.

[0241] The polyA/T segment of the transcriptional DNA template can be produced during PCR by using a reverse primer containing a polyT tail, such as 100 T tail (size can be 50-5000 T), or after PCR by any other method, including, but not limited to, DNA ligation or *in vitro* recombination. Poly(A) tails also provide stability to RNAs and reduce their degradation. Generally, the length of a poly(A) tail positively correlates with the stability of the transcribed RNA. In one embodiment, the poly(A) tail is between 100 and 5000 adenosines.

[0242] Poly(A) tails of RNAs can be further extended following *in vitro* transcription with the use of a poly(A) polymerase, such as E. coli polyA polymerase (E-PAP). In one embodiment,

increasing the length of a poly(A) tail from 100 nucleotides to between 300 and 400 nucleotides results in about a two-fold increase in the translation efficiency of the RNA. Additionally, the attachment of different chemical groups to the 3' end can increase mRNA stability. Such attachment can contain modified/artificial nucleotides, aptamers and other compounds. For example, ATP analogs can be incorporated into the poly(A) tail using poly(A) polymerase. ATP analogs can further increase the stability of the RNA.

[0243] 5' caps on also provide stability to RNA molecules. In a preferred embodiment, RNAs produced by the methods disclosed herein include a 5' cap. The 5' cap is provided using techniques known in the art and described herein (Cougot, et al., Trends in Biochem. Sci., 29:436-444 (2001); Stepinski, et al., RNA, 7:1468-95 (2001); Elango, et al., Biochim. Biophys. Res. Commun., 330:958-966 (2005)).

[0244] The RNAs produced by the methods disclosed herein can also contain an internal ribosome entry site (IRES) sequence. The IRES sequence may be any viral, chromosomal or artificially designed sequence which initiates cap-independent ribosome binding to mRNA and facilitates the initiation of translation. Any solutes suitable for cell electroporation, which can contain factors facilitating cellular permeability and viability such as sugars, peptides, lipids, proteins, antioxidants, and surfactants can be included.

[0245] RNA can be introduced into target cells using any of a number of different methods, for instance, commercially available methods which include, but are not limited to, electroporation (Amaza Nucleofector-II (Amaza Biosystems, Cologne, Germany)), (ECM 830 (BTX) (Harvard Instruments, Boston, Mass.) or the Gene Pulser II (BioRad, Denver, Colo.), Multiporator (Eppendorf, Hamburg Germany), cationic liposome mediated transfection using lipofection, polymer encapsulation, peptide mediated transfection, or biolistic particle delivery systems such as "gene guns" (see, for example, Nishikawa, et al. Hum Gene Ther., 12(8):861-70 (2001)).

Nucleic Acid Constructs Encoding a TFP

[0246] The present invention also provides nucleic acid molecules encoding one or more TFP constructs described herein. In one aspect, the nucleic acid molecule is provided as a messenger RNA transcript. In one aspect, the nucleic acid molecule is provided as a DNA construct.

[0247] The nucleic acid sequences coding for the desired molecules can be obtained using recombinant methods known in the art, such as, for example by screening libraries from cells expressing the gene, by deriving the gene from a vector known to include the same, or by isolating directly from cells and tissues containing the same, using standard techniques. Alternatively, the gene of interest can be produced synthetically, rather than cloned.

[0248] The present invention also provides vectors in which a DNA of the present invention is inserted. Vectors derived from retroviruses such as the lentivirus are suitable tools to achieve

long-term gene transfer since they allow long-term, stable integration of a transgene and its propagation in daughter cells. Lentiviral vectors have the added advantage over vectors derived from onco-retroviruses such as murine leukemia viruses in that they can transduce non-proliferating cells, such as hepatocytes. They also have the added advantage of low immunogenicity.

[0249] In another embodiment, the vector comprising the nucleic acid encoding the desired TFP of the invention is an adenoviral vector (A5/35). In another embodiment, the expression of nucleic acids encoding TFPs can be accomplished using of transposons such as sleeping beauty, crisper, CAS9, and zinc finger nucleases. See below June et al. 2009 Nature Reviews Immunology 9.10: 704-716, is incorporated herein by reference.

[0250] The expression constructs of the present invention may also be used for nucleic acid immunization and gene therapy, using standard gene delivery protocols. Methods for gene delivery are known in the art (see, *e.g.*, U.S. Pat. Nos. 5,399,346, 5,580,859, 5,589,466, incorporated by reference herein in their entireties). In another embodiment, the invention provides a gene therapy vector.

[0251] The nucleic acid can be cloned into a number of types of vectors. For example, the nucleic acid can be cloned into a vector including, but not limited to a plasmid, a phagemid, a phage derivative, an animal virus, and a cosmid. Vectors of particular interest include expression vectors, replication vectors, probe generation vectors, and sequencing vectors.

[0252] Further, the expression vector may be provided to a cell in the form of a viral vector. Viral vector technology is well known in the art and is described, for example, in Sambrook et al., 2012, Molecular Cloning: A Laboratory Manual, volumes 1-4, Cold Spring Harbor Press, NY), and in other virology and molecular biology manuals. Viruses, which are useful as vectors include, but are not limited to, retroviruses, adenoviruses, adeno-associated viruses, herpes viruses, and lentiviruses. In general, a suitable vector contains an origin of replication functional in at least one organism, a promoter sequence, convenient restriction endonuclease sites, and one or more selectable markers, (*e.g.*, WO 01/96584; WO 01/29058; and U.S. Pat. No. 6,326,193).

[0253] A number of virally based systems have been developed for gene transfer into mammalian cells. For example, retroviruses provide a convenient platform for gene delivery systems. A selected gene can be inserted into a vector and packaged in retroviral particles using techniques known in the art. The recombinant virus can then be isolated and delivered to cells of the subject either *in vivo* or *ex vivo*. A number of retroviral systems are known in the art. In some embodiments, adenovirus vectors are used. A number of adenovirus vectors are known in the art. In one embodiment, lentivirus vectors are used.

[0254] Additional promoter elements, *e.g.*, enhancers, regulate the frequency of transcriptional initiation. Typically, these are located in the region 30-110 bp upstream of the start site, although a number of promoters have been shown to contain functional elements downstream of the start site as well. The spacing between promoter elements frequently is flexible, so that promoter function is preserved when elements are inverted or moved relative to one another. In the thymidine kinase (tk) promoter, the spacing between promoter elements can be increased to 50 bp apart before activity begins to decline. Depending on the promoter, it appears that individual elements can function either cooperatively or independently to activate transcription.

[0255] An example of a promoter that is capable of expressing a TFP transgene in a mammalian T-cell is the EF1a promoter. The native EF1a promoter drives expression of the alpha subunit of the elongation factor-1 complex, which is responsible for the enzymatic delivery of aminoacyl tRNAs to the ribosome. The EF1a promoter has been extensively used in mammalian expression plasmids and has been shown to be effective in driving TFP expression from transgenes cloned into a lentiviral vector (see, *e.g.*, Milone et al., Mol. Ther. 17(8): 1453-1464 (2009)). Another example of a promoter is the immediate early cytomegalovirus (CMV) promoter sequence. This promoter sequence is a strong constitutive promoter sequence capable of driving high levels of expression of any polynucleotide sequence operatively linked thereto. However, other constitutive promoter sequences may also be used, including, but not limited to the simian virus 40 (SV40) early promoter, mouse mammary tumor virus (MMTV), human immunodeficiency virus (HIV) long terminal repeat (LTR) promoter, MoMuLV promoter, an avian leukemia virus promoter, an Epstein-Barr virus immediate early promoter, a Rous sarcoma virus promoter, as well as human gene promoters such as, but not limited to, the actin promoter, the myosin promoter, the elongation factor-1a promoter, the hemoglobin promoter, and the creatine kinase promoter. Further, the invention should not be limited to the use of constitutive promoters. Inducible promoters are also contemplated as part of the invention. The use of an inducible promoter provides a molecular switch capable of turning on expression of the polynucleotide sequence which it is operatively linked when such expression is desired, or turning off the expression when expression is not desired. Examples of inducible promoters include, but are not limited to a metallothionein promoter, a glucocorticoid promoter, a progesterone promoter, and a tetracycline-regulated promoter.

[0256] In order to assess the expression of a TFP polypeptide or portions thereof, the expression vector to be introduced into a cell can also contain either a selectable marker gene or a reporter gene or both to facilitate identification and selection of expressing cells from the population of cells sought to be transfected or infected through viral vectors. In other aspects, the selectable marker may be carried on a separate piece of DNA and used in a co-transfection procedure. Both

selectable markers and reporter genes may be flanked with appropriate regulatory sequences to enable expression in the host cells. Useful selectable markers include, for example, antibiotic-resistance genes, such as neo and the like.

[0257] Reporter genes are used for identifying potentially transfected cells and for evaluating the functionality of regulatory sequences. In general, a reporter gene is a gene that is not present in or expressed by the recipient organism or tissue and that encodes a polypeptide whose expression is manifested by some easily detectable property, *e.g.*, enzymatic activity. Expression of the reporter gene is assayed at a suitable time after the DNA has been introduced into the recipient cells. Suitable reporter genes may include genes encoding luciferase, beta-galactosidase, chloramphenicol acetyl transferase, secreted alkaline phosphatase, or the green fluorescent protein gene (*e.g.*, Ui-Tei et al., 2000 FEBS Letters 479: 79-82). Suitable expression systems are well known and may be prepared using known techniques or obtained commercially. In general, the construct with the minimal 5' flanking region showing the highest level of expression of reporter gene is identified as the promoter. Such promoter regions may be linked to a reporter gene and used to evaluate agents for the ability to modulate promoter-driven transcription.

[0258] Methods of introducing and expressing genes into a cell are known in the art. In the context of an expression vector, the vector can be readily introduced into a host cell, *e.g.*, mammalian, bacterial, yeast, or insect cell by any method in the art. For example, the expression vector can be transferred into a host cell by physical, chemical, or biological means.

[0259] Physical methods for introducing a polynucleotide into a host cell include calcium phosphate precipitation, lipofection, particle bombardment, microinjection, electroporation, and the like. Methods for producing cells comprising vectors and/or exogenous nucleic acids are well-known in the art. See, for example, Sambrook et al., 2012, *Molecular Cloning: A Laboratory Manual*, volumes 1-4, Cold Spring Harbor Press, NY). A preferred method for the introduction of a polynucleotide into a host cell is calcium phosphate transfection

[0260] Biological methods for introducing a polynucleotide of interest into a host cell include the use of DNA and RNA vectors. Viral vectors, and especially retroviral vectors, have become the most widely used method for inserting genes into mammalian, *e.g.*, human cells. Other viral vectors can be derived from lentivirus, poxviruses, herpes simplex virus I, adenoviruses and adeno-associated viruses, and the like (see, *e.g.*, U.S. Pat. Nos. 5,350,674 and 5,585,362).

[0261] Chemical means for introducing a polynucleotide into a host cell include colloidal dispersion systems, such as macromolecule complexes, nanocapsules, microspheres, beads, and lipid-based systems including oil-in-water emulsions, micelles, mixed micelles, and liposomes. An exemplary colloidal system for use as a delivery vehicle *in vitro* and *in vivo* is a liposome

(e.g., an artificial membrane vesicle). Other methods of state-of-the-art targeted delivery of nucleic acids are available, such as delivery of polynucleotides with targeted nanoparticles or other suitable sub-micron sized delivery system.

[0262] In the case where a non-viral delivery system is utilized, an exemplary delivery vehicle is a liposome. The use of lipid formulations is contemplated for the introduction of the nucleic acids into a host cell (*in vitro*, *ex vivo* or *in vivo*). In another aspect, the nucleic acid may be associated with a lipid. The nucleic acid associated with a lipid may be encapsulated in the aqueous interior of a liposome, interspersed within the lipid bilayer of a liposome, attached to a liposome via a linking molecule that is associated with both the liposome and the oligonucleotide, entrapped in a liposome, complexed with a liposome, dispersed in a solution containing a lipid, mixed with a lipid, combined with a lipid, contained as a suspension in a lipid, contained or complexed with a micelle, or otherwise associated with a lipid. Lipid, lipid/DNA or lipid/expression vector associated compositions are not limited to any particular structure in solution. For example, they may be present in a bilayer structure, as micelles, or with a “collapsed” structure. They may also simply be interspersed in a solution, possibly forming aggregates that are not uniform in size or shape. Lipids are fatty substances which may be naturally occurring or synthetic lipids. For example, lipids include the fatty droplets that naturally occur in the cytoplasm as well as the class of compounds which contain long-chain aliphatic hydrocarbons and their derivatives, such as fatty acids, alcohols, amines, amino alcohols, and aldehydes.

[0263] Lipids suitable for use can be obtained from commercial sources. For example, dimyristyl phosphatidylcholine (“DMPC”) can be obtained from Sigma, St. Louis, Mo.; dicetyl phosphate (“DCP”) can be obtained from K & K Laboratories (Plainview, N.Y.); cholesterol (“Choi”) can be obtained from Calbiochem-Behring; dimyristyl phosphatidylglycerol (“DMPG”) and other lipids may be obtained from Avanti Polar Lipids, Inc. (Birmingham, Ala.). Stock solutions of lipids in chloroform or chloroform/methanol can be stored at about -20 °C. Chloroform is used as the only solvent since it is more readily evaporated than methanol. “Liposome” is a generic term encompassing a variety of single and multilamellar lipid vehicles formed by the generation of enclosed lipid bilayers or aggregates. Liposomes can be characterized as having vesicular structures with a phospholipid bilayer membrane and an inner aqueous medium. Multilamellar liposomes have multiple lipid layers separated by aqueous medium. They form spontaneously when phospholipids are suspended in an excess of aqueous solution. The lipid components undergo self-rearrangement before the formation of closed structures and entrap water and dissolved solutes between the lipid bilayers (Ghosh et al., 1991 *Glycobiology* 5: 505-10). However, compositions that have different structures in solution than

the normal vesicular structure are also encompassed. For example, the lipids may assume a micellar structure or merely exist as nonuniform aggregates of lipid molecules. Also contemplated are lipofectamine-nucleic acid complexes.

[0264] Regardless of the method used to introduce exogenous nucleic acids into a host cell or otherwise expose a cell to the inhibitor of the present invention, in order to confirm the presence of the recombinant DNA sequence in the host cell, a variety of assays may be performed. Such assays include, for example, “molecular biological” assays well known to those of skill in the art, such as Southern and Northern blotting, RT-PCR and PCR; “biochemical” assays, such as detecting the presence or absence of a particular peptide, *e.g.*, by immunological means (ELISAs and Western blots) or by assays described herein to identify agents falling within the scope of the invention.

[0265] The present invention further provides a vector comprising a TFP encoding nucleic acid molecule. In one aspect, a TFP vector can be directly transduced into a cell, *e.g.*, a T-cell. In one aspect, the vector is a cloning or expression vector, *e.g.*, a vector including, but not limited to, one or more plasmids (*e.g.*, expression plasmids, cloning vectors, minicircles, minivectors, double minute chromosomes), retroviral and lentiviral vector constructs. In one aspect, the vector is capable of expressing the TFP construct in mammalian T-cells. In one aspect, the mammalian T-cell is a human T-cell.

Sources of T-cells

[0266] Prior to expansion and genetic modification, a source of T-cells is obtained from a subject. The term “subject” is intended to include living organisms in which an immune response can be elicited (*e.g.*, mammals). Examples of subjects include humans, dogs, cats, mice, rats, and transgenic species thereof. T-cells can be obtained from a number of sources, including peripheral blood mononuclear cells, bone marrow, lymph node tissue, cord blood, thymus tissue, tissue from a site of infection, ascites, pleural effusion, spleen tissue, and tumors. In certain aspects of the present invention, any number of T-cell lines available in the art, may be used. In certain aspects of the present invention, T-cells can be obtained from a unit of blood collected from a subject using any number of techniques known to the skilled artisan, such as FicollTM separation. In one preferred aspect, cells from the circulating blood of an individual are obtained by apheresis. The apheresis product typically contains lymphocytes, including T-cells, monocytes, granulocytes, B cells, other nucleated white blood cells, red blood cells, and platelets. In one aspect, the cells collected by apheresis may be washed to remove the plasma fraction and to place the cells in an appropriate buffer or media for subsequent processing steps. In one aspect of the invention, the cells are washed with phosphate buffered saline (PBS). In an alternative aspect, the wash solution lacks calcium and may lack magnesium or may lack many

if not all divalent cations. Initial activation steps in the absence of calcium can lead to magnified activation. As those of ordinary skill in the art would readily appreciate a washing step may be accomplished by methods known to those in the art, such as by using a semi-automated “flow-through” centrifuge (for example, the Cobe 2991 cell processor, the Baxter CytoMate, or the Haemonetics Cell Saver 5) according to the manufacturer’s instructions. After washing, the cells may be resuspended in a variety of biocompatible buffers, such as, for example, Ca-free, Mg-free PBS, PlasmaLyte A, or other saline solution with or without buffer. Alternatively, the undesirable components of the apheresis sample may be removed and the cells directly resuspended in culture media.

[0267] In one aspect, T-cells are isolated from peripheral blood lymphocytes by lysing the red blood cells and depleting the monocytes, for example, by centrifugation through a PERCOLL™ gradient or by counterflow centrifugal elutriation. A specific subpopulation of T-cells, such as CD3⁺, CD28⁺, CD4⁺, CD8⁺, CD45RA⁺, and CD45RO⁺ T-cells, can be further isolated by positive or negative selection techniques. For example, in one aspect, T-cells are isolated by incubation with anti-CD3/anti-CD28 (*e.g.*, 3x28)-conjugated beads, such as DYNABEADS™ M-450 CD3/CD28 T, for a time period sufficient for positive selection of the desired T-cells. In one aspect, the time period is about 30 minutes. In a further aspect, the time period ranges from 30 minutes to 36 hours or longer and all integer values there between. In a further aspect, the time period is at least 1, 2, 3, 4, 5, or 6 hours. In yet another preferred aspect, the time period is 10 to 24 hours. In one aspect, the incubation time period is 24 hours. Longer incubation times may be used to isolate T-cells in any situation where there are few T-cells as compared to other cell types, such in isolating tumor infiltrating lymphocytes (TIL) from tumor tissue or from immunocompromised individuals. Further, use of longer incubation times can increase the efficiency of capture of CD8⁺ T-cells. Thus, by simply shortening or lengthening the time T-cells are allowed to bind to the CD3/CD28 beads and/or by increasing or decreasing the ratio of beads to T-cells (as described further herein), subpopulations of T-cells can be preferentially selected for or against at culture initiation or at other time points during the process. Additionally, by increasing or decreasing the ratio of anti-CD3 and/or anti-CD28 antibodies on the beads or other surface, subpopulations of T-cells can be preferentially selected for or against at culture initiation or at other desired time points. The skilled artisan would recognize that multiple rounds of selection can also be used in the context of this invention. In certain aspects, it may be desirable to perform the selection procedure and use the “unselected” cells in the activation and expansion process. “Unselected” cells can also be subjected to further rounds of selection.

[0268] Enrichment of a T-cell population by negative selection can be accomplished with a combination of antibodies directed to surface markers unique to the negatively selected cells. One method is cell sorting and/or selection via negative magnetic immunoadherence or flow cytometry that uses a cocktail of monoclonal antibodies directed to cell surface markers present on the cells negatively selected. For example, to enrich for CD4⁺ cells by negative selection, a monoclonal antibody cocktail typically includes antibodies to CD14, CD20, CD11b, CD16, HLA-DR, and CD8. In certain aspects, it may be desirable to enrich for or positively select for regulatory T-cells which typically express CD4⁺, CD25⁺, CD62Lhi, GITR⁺, and FoxP3⁺. Alternatively, in certain aspects, T regulatory cells are depleted by anti-CD25 conjugated beads or other similar method of selection.

[0269] In one embodiment, a T-cell population can be selected that expresses one or more of IFN- γ , TNF-alpha, IL-17A, IL-2, IL-3, IL-4, GM-CSF, IL-10, IL-13, granzyme B, and perforin, or other appropriate molecules, *e.g.*, other cytokines. Methods for screening for cell expression can be determined, *e.g.*, by the methods described in PCT Publication No.: WO 2013/126712.

[0270] For isolation of a desired population of cells by positive or negative selection, the concentration of cells and surface (*e.g.*, particles such as beads) can be varied. In certain aspects, it may be desirable to significantly decrease the volume in which beads and cells are mixed together (*e.g.*, increase the concentration of cells), to ensure maximum contact of cells and beads. For example, in one aspect, a concentration of 2 billion cells/mL is used. In one aspect, a concentration of 1 billion cells/mL is used. In a further aspect, greater than 100 million cells/mL is used. In a further aspect, a concentration of cells of 10, 15, 20, 25, 30, 35, 40, 45, or 50 million cells/mL is used. In yet one aspect, a concentration of cells from 75, 80, 85, 90, 95, or 100 million cells/mL is used. In further aspects, concentrations of 125 or 150 million cells/mL can be used. Using high concentrations can result in increased cell yield, cell activation, and cell expansion. Further, use of high cell concentrations allows more efficient capture of cells that may weakly express target antigens of interest, such as CD28-negative T-cells, or from samples where there are many tumor cells present (*e.g.*, leukemic blood, tumor tissue, etc.). Such populations of cells may have therapeutic value and would be desirable to obtain. For example, using high concentration of cells allows more efficient selection of CD8⁺ T-cells that normally have weaker CD28 expression.

[0271] In a related aspect, it may be desirable to use lower concentrations of cells. By significantly diluting the mixture of T-cells and surface (*e.g.*, particles such as beads), interactions between the particles and cells is minimized. This selects for cells that express high amounts of desired antigens to be bound to the particles. For example, CD4⁺ T-cells express higher levels of CD28 and are more efficiently captured than CD8⁺ T-cells in dilute

concentrations. In one aspect, the concentration of cells used is $5 \times 10^6/\text{mL}$. In other aspects, the concentration used can be from about $1 \times 10^5/\text{mL}$ to $1 \times 10^6/\text{mL}$, and any integer value in between. In other aspects, the cells may be incubated on a rotator for varying lengths of time at varying speeds at either 2-10 °C or at room temperature.

[0272] T-cells for stimulation can also be frozen after a washing step. Wishing not to be bound by theory, the freeze and subsequent thaw step provides a more uniform product by removing granulocytes and to some extent monocytes in the cell population. After the washing step that removes plasma and platelets, the cells may be suspended in a freezing solution. While many freezing solutions and parameters are known in the art and will be useful in this context, one method involves using PBS containing 20% DMSO and 8% human serum albumin, or culture media containing 10% Dextran 40 and 5% Dextrose, 20% Human Serum Albumin and 7.5% DMSO, or 31.25% Plasmalyte-A, 31.25% Dextrose 5%, 0.45% NaCl, 10% Dextran 40 and 5% Dextrose, 20% Human Serum Albumin, and 7.5% DMSO or other suitable cell freezing media containing for example, Hespan and PlasmaLyte A, the cells then are frozen to -80 °C at a rate of 1 per minute and stored in the vapor phase of a liquid nitrogen storage tank. Other methods of controlled freezing may be used as well as uncontrolled freezing immediately at -20 °C or in liquid nitrogen. In certain aspects, cryopreserved cells are thawed and washed as described herein and allowed to rest for one hour at room temperature prior to activation using the methods of the present invention.

[0273] Also contemplated in the context of the invention is the collection of blood samples or apheresis product from a subject at a time period prior to when the expanded cells as described herein might be needed. As such, the source of the cells to be expanded can be collected at any time point necessary, and desired cells, such as T-cells, isolated and frozen for later use in T-cell therapy for any number of diseases or conditions that would benefit from T-cell therapy, such as those described herein. In one aspect a blood sample or an apheresis is taken from a generally healthy subject. In certain aspects, a blood sample or an apheresis is taken from a generally healthy subject who is at risk of developing a disease, but who has not yet developed a disease, and the cells of interest are isolated and frozen for later use. In certain aspects, the T-cells may be expanded, frozen, and used at a later time. In certain aspects, samples are collected from a patient shortly after diagnosis of a particular disease as described herein but prior to any treatments. In a further aspect, the cells are isolated from a blood sample or an apheresis from a subject prior to any number of relevant treatment modalities, including but not limited to treatment with agents such as natalizumab, efalizumab, antiviral agents, chemotherapy, radiation, immunosuppressive agents, such as cyclosporin, azathioprine, methotrexate, mycophenolate, and FK506, antibodies, or other immunoablative agents such as CAMPATH,

anti-CD3 antibodies, cytoxan, fludarabine, cyclosporin, FK506, rapamycin, mycophenolic acid, steroids, FR901228, and irradiation.

[0274] In a further aspect of the present invention, T-cells are obtained from a patient directly following treatment that leaves the subject with functional T-cells. In this regard, it has been observed that following certain cancer treatments, in particular treatments with drugs that damage the immune system, shortly after treatment during the period when patients would normally be recovering from the treatment, the quality of T-cells obtained may be optimal or improved for their ability to expand *ex vivo*. Likewise, following *ex vivo* manipulation using the methods described herein, these cells may be in a preferred state for enhanced engraftment and *in vivo* expansion. Thus, it is contemplated within the context of the present invention to collect blood cells, including T-cells, dendritic cells, or other cells of the hematopoietic lineage, during this recovery phase. Further, in certain aspects, mobilization (for example, mobilization with GM-CSF) and conditioning regimens can be used to create a condition in a subject wherein repopulation, recirculation, regeneration, and/or expansion of particular cell types is favored, especially during a defined window of time following therapy. Illustrative cell types include T-cells, B cells, dendritic cells, and other cells of the immune system.

Activation and Expansion of T Cells

[0275] T-cells may be activated and expanded generally using methods as described, for example, in U.S. Pat. Nos. 6,352,694; 6,534,055; 6,905,680; 6,692,964; 5,858,358; 6,887,466; 6,905,681; 7,144,575; 7,067,318; 7,172,869; 7,232,566; 7,175,843; 5,883,223; 6,905,874; 6,797,514; 6,867,041; and U.S. Patent Application Publication No. 20060121005.

[0276] Generally, the T-cells of the invention may be expanded by contact with a surface having attached thereto an agent that stimulates a CD3/TCR complex associated signal and a ligand that stimulates a costimulatory molecule on the surface of the T-cells. In particular, T-cell populations may be stimulated as described herein, such as by contact with an anti-CD3 antibody, or antigen-binding fragment thereof, or an anti-CD2 antibody immobilized on a surface, or by contact with a protein kinase C activator (*e.g.*, bryostatin) in conjunction with a calcium ionophore. For co-stimulation of an accessory molecule on the surface of the T-cells, a ligand that binds the accessory molecule is used. For example, a population of T-cells can be contacted with an anti-CD3 antibody and an anti-CD28 antibody, under conditions appropriate for stimulating proliferation of the T-cells. To stimulate proliferation of either CD4⁺ T-cells or CD8⁺ T-cells, an anti-CD3 antibody and an anti-CD28 antibody. Examples of an anti-CD28 antibody include 9.3, B-T3, XR-CD28 (Diaclone, Besancon, France) can be used as can other methods commonly known in the art (Berg et al., Transplant Proc. 30(8):3975-3977, 1998;

Haanen et al., J. Exp. Med. 190(9):1319-1328, 1999; Garland et al., J. Immunol. Meth. 227(1-2):53-63, 1999).

[0277] T-cells that have been exposed to varied stimulation times may exhibit different characteristics. For example, typical blood or apheresed peripheral blood mononuclear cell products have a helper T-cell population (TH, CD4+) that is greater than the cytotoxic or suppressor T-cell population (TC, CD8+). *Ex vivo* expansion of T-cells by stimulating CD3 and CD28 receptors produces a population of T-cells that prior to about days 8-9 consists predominately of TH cells, while after about days 8-9, the population of T-cells comprises an increasingly greater population of TC cells. Accordingly, depending on the purpose of treatment, infusing a subject with a T-cell population comprising predominately of TH cells may be advantageous. Similarly, if an antigen-specific subset of TC cells has been isolated it may be beneficial to expand this subset to a greater degree.

[0278] Further, in addition to CD4 and CD8 markers, other phenotypic markers vary significantly, but in large part, reproducibly during the course of the cell expansion process. Thus, such reproducibility enables the ability to tailor an activated T-cell product for specific purposes.

[0279] Once an anti-CD19 or anti-BCMA TFP is constructed, various assays can be used to evaluate the activity of the molecule, such as but not limited to, the ability to expand T-cells following antigen stimulation, sustain T-cell expansion in the absence of re-stimulation, and anti-cancer activities in appropriate *in vitro* and animal models. Assays to evaluate the effects of an anti-CD19 or anti-BCMA TFP are described in further detail below

[0280] Western blot analysis of TFP expression in primary T-cells can be used to detect the presence of monomers and dimers (see, *e.g.*, Milone et al., Molecular Therapy 17(8): 1453-1464 (2009)). Very briefly, T-cells (1:1 mixture of CD4⁺ and CD8⁺ T-cells) expressing the TFPs are expanded *in vitro* for more than 10 days followed by lysis and SDS-PAGE under reducing conditions. TFPs are detected by Western blotting using an antibody to a TCR chain. The same T-cell subsets are used for SDS-PAGE analysis under non-reducing conditions to permit evaluation of covalent dimer formation.

[0281] *In vitro* expansion of TFP⁺ T-cells following antigen stimulation can be measured by flow cytometry. For example, a mixture of CD4⁺ and CD8⁺ T-cells are stimulated with alphaCD3/alphaCD28 and APCs followed by transduction with lentiviral vectors expressing GFP under the control of the promoters to be analyzed. Exemplary promoters include the CMV IE gene, EF-1alpha, ubiquitin C, or phosphoglycerokinase (PGK) promoters. GFP fluorescence is evaluated on day 6 of culture in the CD4⁺ and/or CD8⁺ T-cell subsets by flow cytometry (see, *e.g.*, Milone et al., Molecular Therapy 17(8): 1453-1464 (2009)). Alternatively, a mixture of

CD4⁺ and CD8⁺ T-cells are stimulated with alphaCD3/alphaCD28 coated magnetic beads on day 0, and transduced with TFP on day 1 using a bicistronic lentiviral vector expressing TFP along with eGFP using a 2A ribosomal skipping sequence. Cultures are re-stimulated with either CD19⁺ K562 cells (K562-CD19), wild-type K562 cells (K562 wild type) or K562 cells expressing hCD32 and 4-1BBL in the presence of antiCD3 and anti-CD28 antibody (K562-BBL-3/28) following washing. Exogenous IL-2 is added to the cultures every other day at 100 IU/mL. GFP⁺ T-cells are enumerated by flow cytometry using bead-based counting (see, *e.g.*, Milone et al., Molecular Therapy 17(8): 1453-1464 (2009)).

[0282] Sustained TFP⁺ T-cell expansion in the absence of re-stimulation can also be measured (see, *e.g.*, Milone et al., Molecular Therapy 17(8): 1453-1464 (2009)). Briefly, mean T-cell volume (fl) is measured on day 8 of culture using a Coulter Multisizer III particle counter following stimulation with alphaCD3/alphaCD28 coated magnetic beads on day 0, and transduction with the indicated TFP on day 1.

[0283] Animal models can also be used to measure a TFP-T activity. For example, xenograft model using human CD19-specific TFP⁺ T-cells to treat a primary human pre-B ALL in immunodeficient mice can be used (see, *e.g.*, Milone et al., Molecular Therapy 17(8): 1453-1464 (2009)). Very briefly, after establishment of ALL, mice are randomized as to treatment groups. Different numbers of engineered T-cells are coinjected at a 1:1 ratio into NOD/SCID/ $\gamma^{-/-}$ mice bearing B-ALL. The number of copies of each vector in spleen DNA from mice is evaluated at various times following T-cell injection. Animals are assessed for leukemia at weekly intervals. Peripheral blood CD19⁺ B-ALL blast cell counts are measured in mice that are injected with alphaCD19-zeta TFP⁺ T-cells or mock-transduced T-cells. Survival curves for the groups are compared using the log-rank test. In addition, absolute peripheral blood CD4⁺ and CD8⁺ T-cell counts 4 weeks following T-cell injection in NOD/SCID/ $\gamma^{-/-}$ mice can also be analyzed. Mice are injected with leukemic cells and 3 weeks later are injected with T-cells engineered to express TFP by a bicistronic lentiviral vector that encodes the TFP linked to eGFP. T-cells are normalized to 45-50% input GFP⁺ T-cells by mixing with mock-transduced cells prior to injection, and confirmed by flow cytometry. Animals are assessed for leukemia at 1-week intervals. Survival curves for the TFP⁺ T-cell groups are compared using the log-rank test.

[0284] Dose dependent TFP treatment response can be evaluated (see, *e.g.*, Milone et al., Molecular Therapy 17(8): 1453-1464 (2009)). For example, peripheral blood is obtained 35-70 days after establishing leukemia in mice injected on day 21 with TFP T-cells, an equivalent number of mock-transduced T-cells, or no T-cells. Mice from each group are randomly bled for determination of peripheral blood CD19⁺ ALL blast counts and then killed on days 35 and 49. The remaining animals are evaluated on days 57 and 70.

[0285] Assessment of cell proliferation and cytokine production has been previously described, *e.g.*, at Milone et al., Molecular Therapy 17(8): 1453-1464 (2009). Briefly, assessment of TFP-mediated proliferation is performed in microtiter plates by mixing washed T-cells with K562 cells expressing CD19 (K19) or CD32 and CD137 (KT32-BBL) for a final T-cell:K562 ratio of 2:1. K562 cells are irradiated with gamma-radiation prior to use. Anti-CD3 (clone OKT3) and anti-CD28 (clone 9.3) monoclonal antibodies are added to cultures with KT32-BBL cells to serve as a positive control for stimulating T-cell proliferation since these signals support long-term CD8⁺ T-cell expansion *ex vivo*. T-cells are enumerated in cultures using CountBrightTM fluorescent beads (Invitrogen) and flow cytometry as described by the manufacturer. TFP⁺ T-cells are identified by GFP expression using T-cells that are engineered with eGFP-2A linked TFP-expressing lentiviral vectors. For TFP⁺ T-cells not expressing GFP, the TFP⁺ T-cells are detected with biotinylated recombinant CD19 protein and a secondary avidin-PE conjugate. CD4⁺ and CD8⁺ expression on T-cells are also simultaneously detected with specific monoclonal antibodies (BD Biosciences). Cytokine measurements are performed on supernatants collected 24 hours following re-stimulation using the human TH1/TH2 cytokine cytometric bead array kit (BD Biosciences) according to the manufacturer's instructions. Fluorescence is assessed using a FACScalibur flow cytometer, and data is analyzed according to the manufacturer's instructions.

[0286] Cytotoxicity can be assessed by a standard ⁵¹Cr-release assay (see, *e.g.*, Milone et al., Molecular Therapy 17(8): 1453-1464 (2009)). Briefly, target cells (K562 lines and primary pro-B-ALL cells) are loaded with ⁵¹Cr (as NaCrO₄, New England Nuclear) at 37 °C for 2 hours with frequent agitation, washed twice in complete RPMI and plated into microtiter plates. Effector T-cells are mixed with target cells in the wells in complete RPMI at varying ratios of effector cell:target cell (E:T). Additional wells containing media only (spontaneous release, SR) or a 1% solution of triton-X 100 detergent (total release, TR) are also prepared. After 4 hours of incubation at 37 °C, supernatant from each well is harvested. Released ⁵¹Cr is then measured using a gamma particle counter (Packard Instrument Co., Waltham, Mass.). Each condition is performed in at least triplicate, and the percentage of lysis is calculated using the formula: % Lysis=(ER-SR)/(TR-SR), where ER represents the average ⁵¹Cr released for each experimental condition.

[0287] Imaging technologies can be used to evaluate specific trafficking and proliferation of TFPs in tumor-bearing animal models. Such assays have been described, *e.g.*, in Barrett et al., Human Gene Therapy 22:1575-1586 (2011). Briefly, NOD/SCID/ γ c^{-/-} (NSG) mice are injected IV with Nalm-6 cells followed 7 days later with T-cells 4 hour after electroporation with the TFP constructs. The T-cells are stably transfected with a lentiviral construct to express firefly

luciferase, and mice are imaged for bioluminescence. Alternatively, therapeutic efficacy and specificity of a single injection of TFP+ T-cells in Nalm-6 xenograft model can be measured as the following: NSG mice are injected with Nalm-6 transduced to stably express firefly luciferase, followed by a single tail-vein injection of T-cells electroporated with CD19 TFP 7 days later. Animals are imaged at various time points post injection. For example, photon-density heat maps of firefly luciferase positive leukemia in representative mice at day 5 (2 days before treatment) and day 8 (24 hours post TFP+ PBLs) can be generated.

[0288] Other assays, including those described in the Example section herein as well as those that are known in the art can also be used to evaluate the anti-CD19 or anti-BCMA TFP constructs of the invention.

Therapeutic Applications

CD19 or BCMA Associated Diseases and/or Disorders

[0289] In one aspect, the invention provides methods for treating a disease associated with CD19 or BCMA expression. In one aspect, the invention provides methods for treating a disease wherein part of the tumor is negative for CD19 or BCMA and part of the tumor is positive for CD19 or BCMA. For example, the TFP of the invention is useful for treating subjects that have undergone treatment for a disease associated with elevated expression of CD19 or BCMA, wherein the subject that has undergone treatment for elevated levels of CD19 or BCMA exhibits a disease associated with elevated levels of CD19 or BCMA.

[0290] In one aspect, the invention pertains to a vector comprising anti-CD19 or BCMA TFP operably linked to promoter for expression in mammalian T-cells. In one aspect, the invention provides a recombinant T-cell expressing the CD19 or BCMA TFP for use in treating CD19- or BCMA- expressing tumors, wherein the recombinant T-cell expressing the CD19 or BCMA TFP is termed a CD19 or BCMA TFP-T. In one aspect, the CD19 or BCMA TFP-T of the invention is capable of contacting a tumor cell with at least one CD19 or BCMA TFP of the invention expressed on its surface such that the TFP-T targets the tumor cell and growth of the tumor is inhibited.

[0291] In one aspect, the invention pertains to a method of inhibiting growth of a CD19- or BCMA- expressing tumor cell, comprising contacting the tumor cell with a CD19 or BCMA TFP T-cell of the present invention such that the TFP-T is activated in response to the antigen and targets the cancer cell, wherein the growth of the tumor is inhibited.

[0292] In one aspect, the invention pertains to a method of treating cancer in a subject. The method comprises administering to the subject a CD19 or BCMA TFP T-cell of the present invention such that the cancer is treated in the subject. An example of a cancer that is treatable by the CD19 or BCMA TFP T-cell of the invention is a cancer associated with expression of

CD19 or BCMA. In one aspect, the cancer associated with expression of CD19 or BCMA is a hematological cancer. In one aspect, the hematological cancer is leukemia or lymphoma. In one aspect, a cancer associated with expression of CD19 includes cancers and malignancies including, but not limited to, *e.g.*, one or more acute leukemias including but not limited to, *e.g.*, B-cell acute Lymphoid Leukemia (“BALL”), T-cell acute Lymphoid Leukemia (“TALL”), acute lymphoid leukemia (ALL); one or more chronic leukemias including but not limited to, *e.g.*, chronic myelogenous leukemia (CML), Chronic Lymphoid Leukemia (CLL). Additional cancers or hematologic conditions associated with expression of CD19 or BCMA include, but are not limited to, *e.g.*, B cell prolymphocytic leukemia, blastic plasmacytoid dendritic cell neoplasm, Burkitt’s lymphoma, diffuse large B cell lymphoma, Follicular lymphoma, Hairy cell leukemia, small cell- or a large cell-follicular lymphoma, malignant lymphoproliferative conditions, MALT lymphoma, mantle cell lymphoma, Marginal zone lymphoma, multiple myeloma, myelodysplasia and myelodysplastic syndrome, non-Hodgkin’s lymphoma, plasmablastic lymphoma, plasmacytoid dendritic cell neoplasm, Waldenstrom macroglobulinemia, and “preleukemia” which are a diverse collection of hematological conditions united by ineffective production (or dysplasia) of myeloid blood cells, and the like. Further a disease associated with CD19 or BCMA expression include, but not limited to, *e.g.*, atypical and/or non-classical cancers, malignancies, precancerous conditions or proliferative diseases associated with expression of CD19 or BCMA.

[0293] In some embodiments, a cancer that can be treated with a CD19 or BCMA TFP, *e.g.*, described herein, is multiple myeloma. Multiple myeloma is a cancer of the blood, characterized by accumulation of a plasma cell clone in the bone marrow. Current therapies for multiple myeloma include, but are not limited to, treatment with lenalidomide, which is an analog of thalidomide. Lenalidomide has activities which include anti-tumor activity, angiogenesis inhibition, and immunomodulation. Generally, myeloma cells are thought to be negative for CD19 or BCMA expression by flow cytometry. The present invention encompasses the recognition that a small percent of myeloma tumor cells express CD19 or BCMA. Thus, in some embodiments, a C19 or BCMA TFP, *e.g.*, as described herein, may be used to target myeloma cells. In some embodiments, CD19 or BCMA TFP therapy can be used in combination with one or more additional therapies, *e.g.*, lenalidomide treatment.

[0294] The invention includes a type of cellular therapy where T-cells are genetically modified to express a TFP and the TFP-expressing T-cell is infused to a recipient in need thereof. The infused cell is able to kill tumor cells in the recipient. Unlike antibody therapies, TFP-expressing T-cells are able to replicate *in vivo* resulting in long-term persistence that can lead to sustained tumor control. In various aspects, the T-cells administered to the patient, or their progeny,

persist in the patient for at least four months, five months, six months, seven months, eight months, nine months, ten months, eleven months, twelve months, thirteen months, fourteen months, fifteen months, sixteen months, seventeen months, eighteen months, nineteen months, twenty months, twenty-one months, twenty-two months, twenty-three months, two years, three years, four years, or five years after administration of the T-cell to the patient.

[0295] The invention also includes a type of cellular therapy where T-cells are modified, *e.g.*, by *in vitro* transcribed RNA, to transiently express a TFP and the TFP-expressing T-cell is infused to a recipient in need thereof. The infused cell is able to kill tumor cells in the recipient. Thus, in various aspects, the T-cells administered to the patient, is present for less than one month, *e.g.*, three weeks, two weeks, or one week, after administration of the T-cell to the patient.

[0296] Without wishing to be bound by any particular theory, the anti-tumor immunity response elicited by the TFP-expressing T-cells may be an active or a passive immune response, or alternatively may be due to a direct vs indirect immune response. In one aspect, the TFP transduced T-cells exhibit specific proinflammatory cytokine secretion and potent cytolytic activity in response to human cancer cells expressing the CD19 or BCMA antigen, resist soluble CD19 or BCMA inhibition, mediate bystander killing and mediate regression of an established human tumor. For example, antigen-less tumor cells within a heterogeneous field of CD19-expressing or BCMA-expressing tumor may be susceptible to indirect destruction by CD19-redirection or BCMA-redirection T-cells that has previously reacted against adjacent antigen-positive cancer cells.

[0297] In one aspect, the human TFP-modified T-cells of the invention may be a type of vaccine for *ex vivo* immunization and/or *in vivo* therapy in a mammal. In one aspect, the mammal is a human.

[0298] With respect to *ex vivo* immunization, at least one of the following occurs *in vitro* prior to administering the cell into a mammal: i) expansion of the cells, ii) introducing a nucleic acid encoding a TFP to the cells or iii) cryopreservation of the cells.

[0299] *Ex vivo* procedures are well known in the art and are discussed more fully below. Briefly, cells are isolated from a mammal (*e.g.*, a human) and genetically modified (*i.e.*, transduced or transfected *in vitro*) with a vector expressing a TFP disclosed herein. The TFP-modified cell can be administered to a mammalian recipient to provide a therapeutic benefit. The mammalian recipient may be a human and the TFP-modified cell can be autologous with respect to the recipient. Alternatively, the cells can be allogeneic, syngeneic or xenogeneic with respect to the recipient.

[0300] The procedure for *ex vivo* expansion of hematopoietic stem and progenitor cells is described in U.S. Pat. No. 5,199,942, incorporated herein by reference, can be applied to the

cells of the present invention. Other suitable methods are known in the art, therefore the present invention is not limited to any particular method of *ex vivo* expansion of the cells. Briefly, *ex vivo* culture and expansion of T-cells comprises: (1) collecting CD34+ hematopoietic stem and progenitor cells from a mammal from peripheral blood harvest or bone marrow explants; and (2) expanding such cells *ex vivo*. In addition to the cellular growth factors described in U.S. Pat. No. 5,199,942, other factors such as flt3-L, IL-1, IL-3 and c-kit ligand, can be used for culturing and expansion of the cells.

[0301] In addition to using a cell-based vaccine in terms of *ex vivo* immunization, the present invention also provides compositions and methods for *in vivo* immunization to elicit an immune response directed against an antigen in a patient.

[0302] Generally, the cells activated and expanded as described herein may be utilized in the treatment and prevention of diseases that arise in individuals who are immunocompromised. In particular, the TFP-modified T-cells of the invention are used in the treatment of diseases, disorders and conditions associated with expression of CD19 or BCMA. In certain aspects, the cells of the invention are used in the treatment of patients at risk for developing diseases, disorders and conditions associated with expression of CD19 or BCMA. Thus, the present invention provides methods for the treatment or prevention of diseases, disorders and conditions associated with expression of CD19 or BCMA comprising administering to a subject in need thereof, a therapeutically effective amount of the TFP-modified T-cells of the invention.

[0303] In one aspect the TFP-T-cells of the inventions may be used to treat a proliferative disease such as a cancer or malignancy or is a precancerous condition such as a myelodysplasia, a myelodysplastic syndrome or a preleukemia. In one aspect, the cancer is a hematological cancer. In one aspect, the hematological cancer is leukemia or lymphoma. In one aspect, the TFP-T-cells of the invention may be used to treat cancers and malignancies such as, but not limited to, *e.g.*, acute leukemias including but not limited to, *e.g.*, B-cell acute lymphoid leukemia (“BALL”), T-cell acute lymphoid leukemia (“TALL”), acute lymphoid leukemia (ALL); one or more chronic leukemias including but not limited to, *e.g.*, chronic myelogenous leukemia (CML), chronic lymphocytic leukemia (CLL); additional hematologic cancers or hematologic conditions including, but not limited to, *e.g.*, B cell prolymphocytic leukemia, blastic plasmacytoid dendritic cell neoplasm, Burkitt’s lymphoma, diffuse large B cell lymphoma, Follicular lymphoma, Hairy cell leukemia, small cell- or a large cell-follicular lymphoma, malignant lymphoproliferative conditions, MALT lymphoma, mantle cell lymphoma, Marginal zone lymphoma, multiple myeloma, myelodysplasia and myelodysplastic syndrome, non-Hodgkin’s lymphoma, plasmablastic lymphoma, plasmacytoid dendritic cell neoplasm, Waldenstrom macroglobulinemia, and “preleukemia” which are a diverse collection

of hematological conditions united by ineffective production (or dysplasia) of myeloid blood cells, and the like. Further a disease associated with CD19 or BCMA expression include, but not limited to, *e.g.*, atypical and/or non-classical cancers, malignancies, precancerous conditions or proliferative diseases expressing CD19 or BCMA. Non-cancer related indications associated with expression of CD19 or BCMA include, but are not limited to, *e.g.*, autoimmune disease, (*e.g.*, lupus), inflammatory disorders (allergy and asthma) and transplantation.

[0304] The TFP-modified T-cells of the present invention may be administered either alone, or as a pharmaceutical composition in combination with diluents and/or with other components such as IL-2 or other cytokines or cell populations.

Hematologic Cancer

[0305] Hematological cancer conditions are the types of cancer such as leukemia and malignant lymphoproliferative conditions that affect blood, bone marrow and the lymphatic system.

[0306] Leukemia can be classified as acute leukemia and chronic leukemia. Acute leukemia can be further classified as acute myelogenous leukemia (AML) and acute lymphoid leukemia (ALL). Chronic leukemia includes chronic myelogenous leukemia (CML) and chronic lymphoid leukemia (CLL). Other related conditions include myelodysplastic syndromes (MDS, formerly known as “preleukemia”) which are a diverse collection of hematological conditions united by ineffective production (or dysplasia) of myeloid blood cells and risk of transformation to AML.

[0307] The present invention provides for compositions and methods for treating cancer. In one aspect, the cancer is a hematologic cancer including but is not limited to hematological cancer is a leukemia or a lymphoma. In one aspect, the TFP-T-cells of the invention may be used to treat cancers and malignancies such as, but not limited to, *e.g.*, acute leukemias including but not limited to, *e.g.*, B-cell acute lymphoid leukemia (“BALL”), T-cell acute lymphoid leukemia (“TALL”), acute lymphoid leukemia (ALL); one or more chronic leukemias including but not limited to, *e.g.*, chronic myelogenous leukemia (CML), chronic lymphocytic leukemia (CLL); additional hematologic cancers or hematologic conditions including, but not limited to, *e.g.*, B cell prolymphocytic leukemia, blastic plasmacytoid dendritic cell neoplasm, Burkitt’s lymphoma, diffuse large B cell lymphoma, Follicular lymphoma, Hairy cell leukemia, small cell- or a large cell-follicular lymphoma, malignant lymphoproliferative conditions, MALT lymphoma, mantle cell lymphoma, Marginal zone lymphoma, multiple myeloma, myelodysplasia and myelodysplastic syndrome, non-Hodgkin’s lymphoma, plasmablastic lymphoma, plasmacytoid dendritic cell neoplasm, Waldenstrom macroglobulinemia, and “preleukemia” which are a diverse collection of hematological conditions united by ineffective production (or dysplasia) of myeloid blood cells, and the like. Further a disease associated with CD19 or BCMA expression includes, but not limited to, *e.g.*, atypical and/or non-classical

cancers, malignancies, precancerous conditions or proliferative diseases expressing CD19 or BCMA.

[0308] The present invention also provides methods for inhibiting the proliferation or reducing a CD19- or BCMA-expressing cell population, the methods comprising contacting a population of cells comprising a CD19- or BCMA-expressing cell with an anti-CD19 or anti-BCMA TFP-T-cell of the invention that binds to the CD19- or BCMA-expressing cell. In a specific aspect, the present invention provides methods for inhibiting the proliferation or reducing the population of cancer cells expressing CD19 or BCMA, the methods comprising contacting the CD19- or BCMA-expressing cancer cell population with an anti-CD19 or anti-BCMA TFP-T-cell of the invention that binds to the CD19- or BCMA-expressing cell. In one aspect, the present invention provides methods for inhibiting the proliferation or reducing the population of cancer cells expressing CD19 or BCMA, the methods comprising contacting the CD19- or BCMA-expressing cancer cell population with an anti-CD19 or anti-BCMA TFP-T-cell of the invention that binds to the CD19- or BCMA-expressing cell. In certain aspects, the anti-CD19 or anti-BCMA TFP-T-cell of the invention reduces the quantity, number, amount or percentage of cells and/or cancer cells by at least 25%, at least 30%, at least 40%, at least 50%, at least 65%, at least 75%, at least 85%, at least 95%, or at least 99% in a subject with or animal model for myeloid leukemia or another cancer associated with CD19- or BCMA-expressing cells relative to a negative control. In one aspect, the subject is a human.

[0309] The present invention also provides methods for preventing, treating and/or managing a disease associated with CD19- or BCMA-expressing cells (*e.g.*, a hematologic cancer or atypical cancer expressing CD19 or BCMA), the methods comprising administering to a subject in need an anti-CD19 or anti-BCMA TFP-T-cell of the invention that binds to the CD19- or BCMA-expressing cell. In one aspect, the subject is a human. Non-limiting examples of disorders associated with CD19- or BCMA-expressing cells include autoimmune disorders (such as lupus), inflammatory disorders (such as allergies and asthma) and cancers (such as hematological cancers or atypical cancers expressing CD19 or BCMA).

[0310] The present invention also provides methods for preventing, treating and/or managing a disease associated with CD19- or BCMA-expressing cells, the methods comprising administering to a subject in need an anti-CD19 or anti-BCMA TFP-T-cell of the invention that binds to the CD19- or BCMA-expressing cell. In one aspect, the subject is a human.

[0311] The present invention provides methods for preventing relapse of cancer associated with CD19- or BCMA-expressing cells, the methods comprising administering to a subject in need thereof an anti-CD19 or anti-BCMA TFP-T-cell of the invention that binds to the CD19- or BCMA-expressing cell. In one aspect, the methods comprise administering to the subject in need

thereof an effective amount of an anti-CD19 or anti-BCMA TFP-T-cell described herein that binds to the CD19- or BCMA-expressing cell in combination with an effective amount of another therapy.

Combination Therapies

[0312] A TFP-expressing cell described herein may be used in combination with other known agents and therapies. Administered “in combination”, as used herein, means that two (or more) different treatments are delivered to the subject during the course of the subject’s affliction with the disorder, *e.g.*, the two or more treatments are delivered after the subject has been diagnosed with the disorder and before the disorder has been cured or eliminated or treatment has ceased for other reasons. In some embodiments, the delivery of one treatment is still occurring when the delivery of the second begins, so that there is overlap in terms of administration. This is sometimes referred to herein as “simultaneous” or “concurrent delivery”. In other embodiments, the delivery of one treatment ends before the delivery of the other treatment begins. In some embodiments of either case, the treatment is more effective because of combined administration. For example, the second treatment is more effective, *e.g.*, an equivalent effect is seen with less of the second treatment, or the second treatment reduces symptoms to a greater extent, than would be seen if the second treatment were administered in the absence of the first treatment or the analogous situation is seen with the first treatment. In some embodiments, delivery is such that the reduction in a symptom, or other parameter related to the disorder is greater than what would be observed with one treatment delivered in the absence of the other. The effect of the two treatments can be partially additive, wholly additive, or greater than additive. The delivery can be such that an effect of the first treatment delivered is still detectable when the second is delivered.

[0313] In some embodiments, the “at least one additional therapeutic agent” includes a TFP-expressing cell. Also provided are T-cells that express multiple TFPs, which bind to the same or different target antigens, or same or different epitopes on the same target antigen. Also provided are populations of T-cells in which a first subset of T-cells express a first TFP and a second subset of T-cells express a second TFP.

[0314] A TFP-expressing cell described herein and the at least one additional therapeutic agent can be administered simultaneously, in the same or in separate compositions, or sequentially. For sequential administration, the TFP-expressing cell described herein can be administered first, and the additional agent can be administered second, or the order of administration can be reversed.

[0315] In further aspects, a TFP-expressing cell described herein may be used in a treatment regimen in combination with surgery, chemotherapy, radiation, immunosuppressive agents, such

as cyclosporin, azathioprine, methotrexate, mycophenolate, and FK506, antibodies, or other immunoablative agents such as CAMPATH, anti-CD3 antibodies or other antibody therapies, cytoxin, fludarabine, cyclosporin, FK506, rapamycin, mycophenolic acid, steroids, FR901228, cytokines, and irradiation. peptide vaccine, such as that described in Izumoto et al. 2008 J Neurosurg 108:963-971.

[0316] In one embodiment, the subject can be administered an agent which reduces or ameliorates a side effect associated with the administration of a TFP-expressing cell. Side effects associated with the administration of a TFP-expressing cell include, but are not limited to cytokine release syndrome (CRS), and hemophagocytic lymphohistiocytosis (HLH), also termed Macrophage Activation Syndrome (MAS). Symptoms of CRS include high fevers, nausea, transient hypotension, hypoxia, and the like. Accordingly, the methods described herein can comprise administering a TFP-expressing cell described herein to a subject and further administering an agent to manage elevated levels of a soluble factor resulting from treatment with a TFP-expressing cell. In one embodiment, the soluble factor elevated in the subject is one or more of IFN- γ , TNF α , IL-2 and IL-6. Therefore, an agent administered to treat this side effect can be an agent that neutralizes one or more of these soluble factors. Such agents include, but are not limited to a steroid, an inhibitor of TNF α , and an inhibitor of IL-6. An example of a TNF α inhibitor is entanercept. An example of an IL-6 inhibitor is tocilizumab (toc).

[0317] In one embodiment, the subject can be administered an agent which enhances the activity of a TFP-expressing cell. For example, in one embodiment, the agent can be an agent which inhibits an inhibitory molecule. Inhibitory molecules, *e.g.*, Programmed Death 1 (PD1), can, in some embodiments, decrease the ability of a TFP-expressing cell to mount an immune effector response. Examples of inhibitory molecules include PD1, PD-L1, CTLA4, TIM3, LAG3, VISTA, BTLA, TIGIT, LAIR1, CD160, 2B4 and TGFR beta. Inhibition of an inhibitory molecule, *e.g.*, by inhibition at the DNA, RNA or protein level, can optimize a TFP-expressing cell performance. In embodiments, an inhibitory nucleic acid, *e.g.*, an inhibitory nucleic acid, *e.g.*, a dsRNA, *e.g.*, an siRNA or shRNA, can be used to inhibit expression of an inhibitory molecule in the TFP-expressing cell. In an embodiment the inhibitor is a shRNA. In an embodiment, the inhibitory molecule is inhibited within a TFP-expressing cell. In these embodiments, a dsRNA molecule that inhibits expression of the inhibitory molecule is linked to the nucleic acid that encodes a component, *e.g.*, all of the components, of the TFP. In one embodiment, the inhibitor of an inhibitory signal can be, *e.g.*, an antibody or antibody fragment that binds to an inhibitory molecule. For example, the agent can be an antibody or antibody fragment that binds to PD1, PD-L1, PD-L2 or CTLA4 (*e.g.*, ipilimumab (also referred to as MDX-010 and MDX-101, and marketed as YervoyTM, Bristol-Myers Squibb; Tremelimumab

(IgG2 monoclonal antibody available from Pfizer, formerly known as ticilimumab, CP-675,206)). In an embodiment, the agent is an antibody or antibody fragment that binds to TIM3. In an embodiment, the agent is an antibody or antibody fragment that binds to LAG3.

[0318] In some embodiments, the agent which enhances the activity of a TFP-expressing cell can be, *e.g.*, a fusion protein comprising a first domain and a second domain, wherein the first domain is an inhibitory molecule, or fragment thereof, and the second domain is a polypeptide that is associated with a positive signal, *e.g.*, a polypeptide comprising an intracellular signaling domain as described herein. In some embodiments, the polypeptide that is associated with a positive signal can include a costimulatory domain of CD28, CD27, ICOS, *e.g.*, an intracellular signaling domain of CD28, CD27 and/or ICOS, and/or a primary signaling domain, *e.g.*, of CD3 zeta, *e.g.*, described herein. In one embodiment, the fusion protein is expressed by the same cell that expressed the TFP. In another embodiment, the fusion protein is expressed by a cell, *e.g.*, a T-cell that does not express an anti-CD19 TFP.

Pharmaceutical Compositions

[0319] Pharmaceutical compositions of the present invention may comprise a TFP-expressing cell, *e.g.*, a plurality of TFP-expressing cells, as described herein, in combination with one or more pharmaceutically or physiologically acceptable carriers, diluents or excipients. Such compositions may comprise buffers such as neutral buffered saline, phosphate buffered saline and the like; carbohydrates such as glucose, mannose, sucrose or dextrans, mannitol; proteins; polypeptides or amino acids such as glycine; antioxidants; chelating agents such as EDTA or glutathione; adjuvants (*e.g.*, aluminum hydroxide); and preservatives. Compositions of the present invention are in one aspect formulated for intravenous administration.

[0320] Pharmaceutical compositions of the present invention may be administered in a manner appropriate to the disease to be treated (or prevented). The quantity and frequency of administration will be determined by such factors as the condition of the patient, and the type and severity of the patient's disease, although appropriate dosages may be determined by clinical trials.

[0321] In one embodiment, the pharmaceutical composition is substantially free of, *e.g.*, there are no detectable levels of a contaminant, *e.g.*, selected from the group consisting of endotoxin, mycoplasma, replication competent lentivirus (RCL), p24, VSV-G nucleic acid, HIV gag, residual anti-CD3/anti-CD28 coated beads, mouse antibodies, pooled human serum, bovine serum albumin, bovine serum, culture media components, vector packaging cell or plasmid components, a bacterium and a fungus. In one embodiment, the bacterium is at least one selected from the group consisting of *Alcaligenes faecalis*, *Candida albicans*, *Escherichia coli*,

Haemophilus influenza, Neisseria meningitides, Pseudomonas aeruginosa, Staphylococcus aureus, Streptococcus pneumonia, and Streptococcus pyogenes group A.

[0322] When “an immunologically effective amount,” “an anti-tumor effective amount,” “a tumor-inhibiting effective amount,” or “therapeutic amount” is indicated, the precise amount of the compositions of the present invention to be administered can be determined by a physician with consideration of individual differences in age, weight, tumor size, extent of infection or metastasis, and condition of the patient (subject). It can generally be stated that a pharmaceutical composition comprising the T-cells described herein may be administered at a dosage of 10^4 to 10^9 cells/kg body weight, in some instances 10^5 to 10^6 cells/kg body weight, including all integer values within those ranges. T-cell compositions may also be administered multiple times at these dosages. The cells can be administered by using infusion techniques that are commonly known in immunotherapy (see, *e.g.*, Rosenberg et al., New Eng. J. of Med. 319:1676, 1988).

[0323] In certain aspects, it may be desired to administer activated T-cells to a subject and then subsequently redraw blood (or have an apheresis performed), activate T-cells therefrom according to the present invention, and reinfuse the patient with these activated and expanded T-cells. This process can be carried out multiple times every few weeks. In certain aspects, T-cells can be activated from blood draws of from 10 cc to 400 cc. In certain aspects, T-cells are activated from blood draws of 20 cc, 30 cc, 40 cc, 50 cc, 60 cc, 70 cc, 80 cc, 90 cc, or 100 cc.

[0324] The administration of the subject compositions may be carried out in any convenient manner, including by aerosol inhalation, injection, ingestion, transfusion, implantation or transplantation. The compositions described herein may be administered to a patient trans arterially, subcutaneously, intradermally, intratumorally, intranodally, intramedullary, intramuscularly, by intravenous (i.v.) injection, or intraperitoneally. In one aspect, the T-cell compositions of the present invention are administered to a patient by intradermal or subcutaneous injection. In one aspect, the T-cell compositions of the present invention are administered by i.v. injection. The compositions of T-cells may be injected directly into a tumor, lymph node, or site of infection.

[0325] In a particular exemplary aspect, subjects may undergo leukapheresis, wherein leukocytes are collected, enriched, or depleted *ex vivo* to select and/or isolate the cells of interest, *e.g.*, T-cells. These T-cell isolates may be expanded by methods known in the art and treated such that one or more TFP constructs of the invention may be introduced, thereby creating a TFP-expressing T-cell of the invention. Subjects in need thereof may subsequently undergo standard treatment with high dose chemotherapy followed by peripheral blood stem cell transplantation. In certain aspects, following or concurrent with the transplant, subjects receive

an infusion of the expanded TFP T-cells of the present invention. In an additional aspect, expanded cells are administered before or following surgery.

[0326] The dosage of the above treatments to be administered to a patient will vary with the precise nature of the condition being treated and the recipient of the treatment. The scaling of dosages for human administration can be performed according to art-accepted practices. The dose for CAMPATH, for example, will generally be in the range 1 to about 100 mg for an adult patient, usually administered daily for a period between 1 and 30 days. The preferred daily dose is 1 to 10 mg per day although in some instances larger doses of up to 40 mg per day may be used (described in U.S. Pat. No. 6,120,766).

[0327] In one embodiment, the TFP is introduced into T-cells, *e.g.*, using *in vitro* transcription, and the subject (*e.g.*, human) receives an initial administration of TFP T-cells of the invention, and one or more subsequent administrations of the TFP T-cells of the invention, wherein the one or more subsequent administrations are administered less than 15 days, *e.g.*, 14, 13, 12, 11, 10, 9, 8, 7, 6, 5, 4, 3, or 2 days after the previous administration. In one embodiment, more than one administration of the TFP T-cells of the invention are administered to the subject (*e.g.*, human) per week, *e.g.*, 2, 3, or 4 administrations of the TFP T-cells of the invention are administered per week. In one embodiment, the subject (*e.g.*, human subject) receives more than one administration of the TFP T-cells per week (*e.g.*, 2, 3 or 4 administrations per week) (also referred to herein as a cycle), followed by a week of no TFP T-cells administrations, and then one or more additional administration of the TFP T-cells (*e.g.*, more than one administration of the TFP T-cells per week) is administered to the subject. In another embodiment, the subject (*e.g.*, human subject) receives more than one cycle of TFP T-cells, and the time between each cycle is less than 10, 9, 8, 7, 6, 5, 4, or 3 days. In one embodiment, the TFP T-cells are administered every other day for 3 administrations per week. In one embodiment, the TFP T-cells of the invention are administered for at least two, three, four, five, six, seven, eight or more weeks.

[0328] In one aspect, CD19 TFP T-cells are generated using lentiviral viral vectors, such as lentivirus. TFP-T-cells generated that way will have stable TFP expression.

[0329] In one aspect, TFP T-cells transiently express TFP vectors for 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15 days after transduction. Transient expression of TFPs can be effected by RNA TFP vector delivery. In one aspect, the TFP RNA is transduced into the T-cell by electroporation.

[0330] A potential issue that can arise in patients being treated using transiently expressing TFP T-cells (particularly with murine scFv bearing TFP T-cells) is anaphylaxis after multiple treatments.

[0331] Without being bound by this theory, it is believed that such an anaphylactic response might be caused by a patient developing humoral anti-TFP response, i.e., anti-TFP antibodies having an anti-IgE isotype. It is thought that a patient's antibody producing cells undergo a class switch from IgG isotype (that does not cause anaphylaxis) to IgE isotype when there is a ten to fourteen day break in exposure to antigen.

[0332] If a patient is at high risk of generating an anti-TFP antibody response during the course of transient TFP therapy (such as those generated by RNA transductions), TFP T-cell infusion breaks should not last more than ten to fourteen days.

EXAMPLES

[0333] The invention is further described in detail by reference to the following experimental examples. These examples are provided for purposes of illustration only, and are not intended to be limiting unless otherwise specified. Thus, the invention should in no way be construed as being limited to the following examples, but rather, should be construed to encompass any and all variations which become evident as a result of the teaching provided herein. Without further description, it is believed that one of ordinary skill in the art can, using the preceding description and the following illustrative examples, make and utilize the compounds of the present invention and practice the claimed methods. The following working examples specifically point out various aspects of the present invention, and are not to be construed as limiting in any way the remainder of the disclosure.

Example 1: TFP Constructs

[0334] Anti-CD19 TFP constructs were engineered by cloning an anti-CD19 scFv DNA fragment linked to a CD3 or TCR DNA fragment by either a DNA sequence encoding a short linker (SL): AAAGGGGSGGGGSGGGGSLE (SEQ ID NO: 5) or a long linker (LL): AAAIEVMYPPPYLGGGGSGGGGSGGGGSLE (SEQ ID NO: 6) into p510 vector ((System Biosciences (SBI)) at XbaI and EcoRI sites.

[0335] The anti-CD19 TRuC constructs generated were p510_antiCD19_LL_TCR α (anti-CD19 scFv – long linker- human full length T cell receptor α chain), p510_antiCD19_LL_TCR α C (anti-CD19 scFv – long linker- human T cell receptor α constant domain chain), p510_antiCD19_LL_TCR β (anti-CD19 scFv – long linker- human full length T cell receptor β chain), p510_antiCD19_LL_TCR β C (anti-CD19 scFv – long linker- human T cell receptor β constant domain chain), p510_antiCD19_LL_CD3 γ (anti-CD19 scFv – long linker- human CD3 γ chain), p510_antiCD19_LL_CD3 δ (anti-CD19 scFv – long linker- human CD3 δ chain), p510_antiCD19_LL_CD3 ϵ (anti-CD19 scFv – long linker- human CD3 ϵ chain), p510_antiCD19_SL_TCR β (anti-CD19 scFv – short linker- human full length T cell receptor β chain), p510_antiCD19_SL_CD3 γ (anti-CD19 scFv – short linker- human CD3 γ chain),

p510_antiCD19_SL_CD3 δ (anti-CD19 scFv – short linker- human CD3 δ chain),

p510_antiCD19_SL_CD3 ϵ (anti-CD19 scFv – short linker- human CD3 ϵ chain).

[0336] The anti-CD19 CAR construct, p510_antiCD19_28 ζ was generated by cloning synthesized DNA encoding anti-CD19, partial CD28 extracellular domain, CD28 transmembrane domain, CD28 intracellular domain and CD3 zeta into p510 vector at XbaI and EcoR1 sites.

[0337] Anti-BCMA TFP constructs were engineered by cloning an anti-BCMA scFv DNA fragment linked to a CD3 DNA fragment by a DNA sequence encoding the linker: GGGGSGGGGSGGGGSLE (SEQ ID NO: 7) into p510 vector (SBI) at XbaI and EcoR1 sites. The anti-BCMA TFP constructs generated were p510_antiBCMA_CD3 γ (anti-BCMA scFv – linker- human CD3 γ chain) and p510_antiBCMA_CD3 ϵ (anti-BCMA scFv – linker- human CD3 ϵ chain).

[0338] Full length BCMA was synthesized and cloned into p514 (SBI) at BamHI and NheI sites to generate the construct p514_BCMA, used to generate stable target cell lines.

[0339] Anti-Fibroblast activation protein (FAP) and anti-Carboanhydrase-9 (CAIX) TFP constructs are engineered by cloning an anti-FAP or anti-CAIX scFv DNA fragment linked to a CD3 DNA fragment by a DNA sequence encoding the linker: GGGGSGGGGSGGGGSLE (SEQ ID NO: 7) into p510 vector (SBI) at XbaI and EcoR1 sites. The anti-FAP or anti-CAIX TFP constructs that can be generated include p510_antiFAP_CD3 γ (anti-FAP scFv – linker- human CD3 γ chain) and p510_antiFAP_CD3 ϵ (anti-FAP scFv – linker- human CD3 ϵ chain) and p510_antiCAIX_CD3 γ (anti-CAIX scFv – linker-human CD3 γ chain) and p510_antiCAIX_CD3 ϵ (anti-CAIX scFv – linker- human CD3 ϵ chain).

[0340] Full length FAP and CAIX can be synthesized and cloned into p514 (SBI) at BamHI and NheI sites to generate the constructs p514_FAP and p514_CAIX, that can be used to generate stable target cell lines.

[0341] Exemplary construct sequences are shown below:

CONSTRUCT SEQUENCES

Target Construct

***P514_BCMA* (SEQ ID NO: 8)**

```

1      acgcgtgtag tcttatgcaa tactcttgta gtcttgcaac atggtaacga tgagttagca
61     acatgcctta caaggagaga aaaagcaccg tgcattgcga ttggtggaag taagggtgga
121    cgatcgtagc ttattaggaa ggcaacagac gggcttgaca tggattggac gaaccactga
181    attgccgcat tgcagagata ttgtatttaa gtgcctagct cgatacataa acgggtctct
241    ctgggttagac cagatctgag cctgggagct ctctggctaa ctagggaacc cactgcttaa
301    gcctcaataa agcttgccct gagtgcttca agtagtgtgt gcccgctctgt tgtgtgactc
361    tggttaactag agatccctca gaccctttta gtcagtgtgg aaaatctcta gcagtggcgc
421    ccgaacaggg acctgaaagc gaaagggaaa ccagagctct ctgcagcgag gactcggcct
481    gctgaagcgc gcacggcaag aggcgagggg cggcgactgg tgagtacgcc aaaaattttg
541    actagcggag gctagaagga gagagatggg tgcgagagcg tcagtattaa gcggggggaga

```

601 attagatcgc gatgggaaaa aattcgggta aggccagggg gaaagaaaaa atataaatta
 661 aaacatatag tatgggcaag caggagagcta gaacgattcg cagttaatcc tggcctgtta
 721 gaaacatcag aaggctgtag acaaatactg ggacagctac aaccatccct tcagacagga
 781 tcagaagaac ttagatcatt atataatata gtagcaaccc tctatttgtt gcatcaaagg
 841 atagagataa aagacaccaa ggaagcttta gacaagatag aggaagagca aaacaaaagt
 901 aagaccaccg cacagcaagc ggccactgat cttcagacct ggaggaggag atatgaggga
 961 caattggaga agtgaattat ataaatataa agtagtaaaa attgaacct taggagtagc
 1021 acccaccaag gcaaagagaa gagtgggtgca gagagaaaaa agagcagtgg gaataggagc
 1081 tttgttcctt ggggttccttg gagcagcagg aagcactatg ggccgcagcgt caatgacgct
 1141 gacgggtacag gccagacaat tattgtctgg tatagtgcag cagcagaaca atttgctgag
 1201 ggctattgag gcgcaacagc atctgttgca actcacagtc tggggcatca agcagctcca
 1261 ggcaagaatc ctggctgtgg aaagatacct aaaggatcaa cagctcctgg ggatttgggg
 1321 ttgctctgga aaactcattt gcaccactgc tgtgccttgg aatgctagtt ggagtaataa
 1381 atctctggaa cagatttgga atcacacgac ctggatggag tgggacagag aaattaacaa
 1441 ttacacaagc ttaatacact ccttaattga agaatcgcaa aaccagcaag aaaagaatga
 1501 acaagaatta ttggaattag ataaatgggc aagtttgtgg aattgggtta acataacaaa
 1561 ttggctgtgg tatataaaat tattcataat gatagtagga ggcttggtag gtttaagaat
 1621 agtttttgct gtactttcta tagtgaatag agttaggcag ggatattcac cattatcggt
 1681 tcagaccacac ctcccaaccc cgaggggacc cgacaggccc gaaggaatag aagaagaagg
 1741 tggagagaga gacagagaca gatccattcg attagtgaac ggatctcgac ggtatcgggt
 1801 aactttttaa agaaaagggg ggattggggg gtacagtgcg ggggaaagaa tagtagacat
 1861 aatagcaaca gacatacaaa ctaaagaatt acaaaaaaaa attacaaaat tcaaaatttt
 1921 atcgatacta gtattatgcc cagtacatga ccttatggga ctttcctact tggcagtaca
 1981 tctacgtatt agtcacgct attaccatgg tgatgcgggt ttggcagtac atcaatgggc
 2041 gtggatagcg gtttgactca cggggatttc caagtctcca cccattgac gtcaatggga
 2101 gtttggtttg gcacccaaaat caacgggact ttccaaaatg tcgtaacaac tccgccccat
 2161 tgacgcaaat gggcggtagg cgtgtacggg gggagggtta tataagcaga gctcgtttag
 2221 tgaaccgtca gatcgctgg agacgccatc cagctgtttt tgacctccat agaagattct
 2281 agagctagcg ccgccaccat gctccagatg gctggccagt gcagccagaa cgagtacttc
 2341 gacagcctgc tgcacgcctg catcccttgc cagctgcggg gcagcagcaa caccaccacc
 2401 ctgacctgcc agcgggtact caacgccagc gtgaccaaca gcgtgaaggg caccaacgcc
 2461 atcctgtgga cctgcctggg cctgagcctg atcatcagcc tggccgtgtt cgtgctgatg
 2521 ttcctgctgc ggaagatcaa cagcagagcc ctgaaggacg agttcaagaa caccggcagc
 2581 ggcctgctgg gcatggccaa catcgacctg gaaaagagcc ggaccggcga cgagatcatc
 2641 ctgcccagag gcctggagta caccgtggaa gagtgtacct gcgaggactg catcaagagc
 2701 aagcccaagg tggacagcga ccactgcttc cctctgcccg ccatggaaga gggcgccacc
 2761 atcctggtga caacaaagac caacgactac tgcaagagcc tgcctgcgcg cctgagcgcc
 2821 accgagatcg agaagtccat cagcgccaga tgaggatccg cggccgcaag gatctgcgat
 2881 cgctccggtg cccgtcagtg ggcagagcgc acatcgccca cagtccccga gaagtgggg
 2941 ggaggggctg gcaattgaac ggggtgcctag agaagggtggc gcggggtaaa ctgggaaagt
 3001 gatgtcgtgt actggctccg cctttttccc gaggggtgggg gagaaccgta tataagtga
 3061 gtagtcgccc tgaacgttct ttttcgcaac ggggttgccg ccagaacaca gctgaagctt
 3121 cgaggggctc gcatctctcc ttcacgcgcc cgcgcgcceta cctgaggccg ccatccacgc
 3181 cggttgagtc gcgttctgcc gcctcccgcc tgtggtgcct cctgaactgc gtccgcccgc
 3241 taggtaagtt taaagctcag gtcgagaccg ggctttgtc cggcgctccc ttggagccta
 3301 cctagactca gccggctctc cacgctttgc ctgacctgc ttgctcaact ctacgtcttt
 3361 gtttcgtttt ctgttctgcg ccgttacaga tccaagctgt gaccggcgcc tacgtcgaga
 3421 tgattgaaca agatggattg cacgcagggt ctccggccgc ttgggtggag aggctattcg
 3481 gctatgactg ggcacaacag acaatcggct gctctgatgc cgccgtgttc cggctgtcag
 3541 cgcagggggc cccggttctt tttgtcaaga ccgacctgtc cgggtgccctg aatgaactgc
 3601 aggacgaggc agegcggcta tegtggctgg ccgcgacggg cgttccttgc gcagctgtgc
 3661 tcgacgttgt cactgaagcg ggaagggact ggctgctatt gggcgaaagt ccggggcagg
 3721 atctcctgtc atctcacctt gctcctgccg agaaagtatc catcatggct gatgcaatgc
 3781 ggcggctgca tacgcttgat ccggctacct gccattcga ccaccaagcg aaacatcgca
 3841 tcgagcgagc acgtactcgg atggaagcgg gtcttgtcga tcaggatgat ctggacgaag
 3901 agcatcaggg gctcgcgcc a gccgaactgt tcgccaggct caaggcgccg atgcccagc
 3961 gcgaggatct cgtcgtgacc catggcgatg cctgcttgcc gaatatcatg gtggaaaaatg
 4021 gccgcttttc tggattcctc gactgtggcc ggctgggtgt ggccgaccgc tatcaggaca
 4081 tagcgttggc taccctgat attgctgaag agcttggcgg cgaatgggct gaccgcttcc

4141 tcgtgcttta cggatatcgcc gctcccgatt cgcagcgcac cgccttctat cgccttcttg
 4201 acgagttcct ctgactcgac aatcaacctc tggattacaa aatttgtgaa agattgactg
 4261 gtattcttaa ctatgttgct ccttttacgc tatgtggata cgctgcttta atgcctttgt
 4321 atcatgctat tgcttcccgt atggctttca ttttctctc cttgtataaa tcttggttgc
 4381 tgtctcttta tgaggagtgt tggcccgttg tcaggcaacg tggcgtggtg tgcactgtgt
 4441 ttgctgacgc aacccccact gggtggggca ttgccaccac ctgtcagctc ctttccggga
 4501 ctttctgctt cccctccctt attgccacgg cggaaactcat cgcgcctgc cttgcccgct
 4561 gctggacagg ggctcggttg ttgggcaactg acaattccgt ggtgttgtcg gggaaatcat
 4621 cgtcctttcc ttggctgtct gcctgtgttg ccacctggat tctgcgcggg acgtccttct
 4681 gctacgtccc ttccggcctc aatccagcgg accttcttc ccgcggcctg ctgccggctc
 4741 tgcggcctct tccgctctt cgccttcgcc ctacagacgag tcggatctcc ctttggggcg
 4801 cctccccgcc tgggtacctt aagaccaatg acttacaagg cagctgtaga tcttagccac
 4861 tttttaaaag aaaagggggg actggaaggg ctaattcact cccaacgaag ataagatctg
 4921 ctttttgctt gtactgggtc tctctggtta gaccagatct gagcctggga gctctctggc
 4981 taactaggga acccactgct taagcctcaa taaagcttg cttgagtgt tcaagtagtg
 5041 tgtgcccgtc tgttgtgtga ctctggtaac tagagatccc tcagaccctt ttagtcagt
 5101 tggaaaatct ctacgagtag tagttcatgt catcttatta ttcagtattt ataacttgca
 5161 aagaaatgaa tatcagagag tgagaggaac ttgtttattg cagcttataa tggttacaaa
 5221 taaagcaata gcatcacaaa tttcacaaat aaagcatttt tttcactgca ttctagtgt
 5281 ggtttgtcca aactcatcaa tgtatcttat catgtctggc tctagctatc ccgcccctaa
 5341 ctccgcccat cccgcccta actccgccca gttccgccca ttctccgcc catggttgac
 5401 taattttttt tatttatgca gaggccgagg ccgcctcggc ctctgagcta ttccagaagt
 5461 agtgaggagg cttttttgga ggcctagact tttgcagaga cggcccaaat tcgtaatcat
 5521 ggtcatagct gtttctgtg tgaaattgtt atccgctcac aattccacac aacatacgag
 5581 ccggaagcat aaagtgtaaa gcctgggggtg cctaattgagt gagctaactc acattaattg
 5641 cgttgcgctc actgcccgct ttccagtcgg gaaacctgtc gtgccagctg cattaatgaa
 5701 tcggccaacg cgcggggaga ggcggtttgc gtattggggc ctcttccgct tctcgtccta
 5761 ctgactcgct gcgctcggtc gttcggctgc ggcgagcgg atcagctcac tcaaaggcgg
 5821 taatacggtt atccacagaa tcaggggata acgcaggaaa gaacatgtga gcaaaaggcc
 5881 agcaaaaggc caggaaccgt aaaaaggccg cgttgctggc gtttttccat aggctccgcc
 5941 cccctgacga gcatcacaaa aatcgacgct caagtcagag gtggcgaaac ccgacaggac
 6001 tataaagata ccaggcggtt cccctggaa gctccctcgt gcgctctcct gttccgacct
 6061 tgccgcttac cggatacctg tccgccttcc tcccttcggg aagcgtggcg ctttctcata
 6121 gctcacgctg taggtatctc agttcggtgt aggtcggttcg ctccaagctg ggctgtgtgc
 6181 acgaaccccc cgttcagccc gaccgctgcg cttatccgg taactatcgt cttgagtcca
 6241 acccggttag acacgactta tcgccactgg cagcagccac tggtaacagg attagcagag
 6301 cgaggatagt aggcggtgct acagagttct tgaagtgggt gcctaactac ggctacacta
 6361 gaaggacagt atttggatc tgcgctctgc tgaagccagt taccttcgga aaaagagtgt
 6421 gtagctcttg atccggcaaa caaaccaccg ctggtagcgg tggttttttt gtttgcaagc
 6481 agcagattac gcgcagaaaa aaaggatctc aagaagatcc tttgatcttt tctacggggg
 6541 ctgacgctca gtggaacgaa aactcacgtt aagggaattt ggtcatgaga ttatcaaaaa
 6601 ggatcttcac ctagatcctt ttaaattaaa aatgaagtgt taaatcaatc taaagtatat
 6661 atgagtaaac ttggtctgac agttaccaat gcttaatcag tgaggcacct atctcagcga
 6721 tctgtctatt tcgttcaccc atagttgcct gactccccgt cgtgtagata actacgatac
 6781 gggagggctt accatctggc cccagtgtcg caatgatacc gcgagaccca cgctcaccgg
 6841 ctccagattt atcagcaata aaccagccag ccggaagggg cgagcgcaga agtggtcctg
 6901 caactttatc cgctccatc cagtctatta attgttgccg ggaagctaga gtaagtgtt
 6961 cgccagttaa tagtttgccg aacgttggtg ccattgctac aggcctcgtg gtgtcacgct
 7021 cgtcgttttg tatggcttca ttcagctccg gttcccaacg atcaaggcga gttacatgat
 7081 ccccatgtt gtgcaaaaaa gcggttagct ccttcggtcc tccgatcgtt gtcagaagta
 7141 agttggccgc agtggttatca ctcatgggtta tggcagcaact gcataattct cttactgtca
 7201 tgccatccgt aagatgcttt tctgtgactg gtgagtactc aaccaagtca ttctgagaat
 7261 agtggtatgc gcgaccgagt tgccttgcgc cggcgtcaat acgggataat accgcgccac
 7321 atagcagaac tttaaaagtgc ctcatcattg gaaaacgttc ttcggggcga aaactctcaa
 7381 ggatcttacc gctgttgaga tccagttcga tgtaaaccac tegtgcaccc aactgatctt
 7441 cagcatcttt tactttcacc agcgtttctg ggtgagcaaa aacaggaagg caaaatgccg
 7501 caaaaaaggg aataaggggcg acacggaaat gttgaatact catactcttc ctttttcaat
 7561 attattgaag catttatcag gggtattgtc tcatgagcgg atacatattt gaatgtattt
 7621 agaaaaataa acaaataggg gttccgcgca catttccccg aaaagtgccca cctgacgtct

7681 aagaaacccat tattatcatg acattaacct ataaaaatag gcgtatcacg aggccctttc
 7741 gtctcgcgcg tttcgggtgat gacgggtgaaa acctctgaca catgcagctc ccggagacgg
 7801 tcacagcttg tctgtaagcg gatgccggga gcagacaagc ccgtcagggc gcgtcagcgg
 7861 gtgttggcgg gtgtcggggc tggcttaact atgcggcatc agagcagatt gtactgagag
 7921 tgcaccatat gcggtgtgaa ataccgcaca gatgcgtaag gagaaaatac cgcacaggc
 7981 gccattcgcc attcaggctg cgcaactgtt gggaagggcg atcgggtcgg gcctcttcgc
 8041 tattacgcca gctggcgaaa gggggatgtg ctgcaaggcg attaagttgg gtaacgccag
 8101 ggttttccca gtcacgacgt tgtaaacga cggccagtgc caagctg.

CAR Constructs***p510_antiCD19_28z*** (SEQ ID NO: 9)

1 acgcgtgtag tcttatgcaa tactcttgta gtcttgcaac atggtaacga tgagttagca
 61 acatgcctta caaggagaga aaaagcaccg tgcattgcga ttggtggaag taagggtgta
 121 cgatcgtgcc ttattaggaa ggcaacagac gggcttgaca tggattggac gaaccactga
 181 attgccgcat tgcagagata ttgtatttaa gtgcctagct cgatacaata aacgggtctc
 241 tctgggttaga ccagatctga gcctgggagc tctctggcta actagggaaac ccaactgctta
 301 agcctcaata aagcttgcc ttagtgcttc aagtagtgtg tgcccgtctg ttgtgtgact
 361 ctggtaacta gagatccctc agaccctttt agtcagtgtg gaaaatctct agcagtggcg
 421 cccgaacagg gacctgaaag cgaaagggaa accagagctc tctcgacgca ggactcggct
 481 tgctgaagcg cgcacggcaa gaggcgaggg gcggcgactg gtgagtacgc caaaaatttt
 541 gactagcggg ggctagaagg agagagatgg gtgcgagagc gtcagtatta agcgggggag
 601 aattagatcg cgatgggaaa aaattcgggtt aaggccaggg ggaaagaaaa aatataaatt
 661 aaaacatata gtatgggcaa gcaggagct agaacgattc gcagttaatc ctggcctgtt
 721 agaaacatca gaaggctgta gacaaatact gggacagcta caaccatccc ttcagacagg
 781 atcagaagaa cttagatcat tatataatac agtagcaacc ctctattgtg tgcacaaag
 841 gatagagata aaagacacca aggaagcttt agacaagata gaggaagagc aaaacaaaag
 901 taagaccacc gcacagcaag cggccactga tcttcagacc tggaggagga gatatgaggg
 961 acaattggag aagtgaatta tataaatata aagtagtaaa aattgaacca ttaggagtag
 1021 caccaccaa ggcaagaga agagtgggtg agagagaaaa aagagcagtg ggaataggag
 1081 ctttgttctt tgggttcttg ggagcagcag gaagcactat gggcgagcc tcaatgacgc
 1141 tgacggtaca ggccagacaa ttattgtctg gtatagtga gcagcagaac aatttgtctga
 1201 gggctattga ggcgcaacag catctgttgc aactcacagt ctggggcatc aagcagctcc
 1261 aggcaagaat cctggctgtg gaaagatacc taaaggatca acagctcctg gggatttggg
 1321 gttgctctgg aaaactcatt tgcaccactg ctgtgccttg gaatgctagt tggagtaata
 1381 aatctctgga acagattgga atcacacgac ctggatggag tgggacagag aaattaacaa
 1441 ttacacaagc ttaatacact ccttaattga agaatcgcaa aaccagcaag aaaagaatga
 1501 acaagaatta ttggaattag ataaatgggc aagtttgttg aattggttta acataacaaa
 1561 ttggctgttg tatataaaat tattcataat gatagtagga ggcttggtag gtttaagaat
 1621 agtttttctg gtactttcta tagtgaatag agttaggcag ggatattcac cattatcggt
 1681 tcagaccac cteccaaccc cgaggggacc cgacaggccc gaaggaatag aagaagaagg
 1741 tggagagaga gacagagaca gatccattcg attagtgaac ggatctcgac ggtatcgggt
 1801 aactttttaa agaaaagggg ggattggggg gtacagtga ggggaaagaa tagtagacat
 1861 aatagcaaca gacatacaaa cttaaagaatt acaaaaacaa attacaaaat tcaaaatttt
 1921 atcgatacta gtattatgcc cagtacatga ccttatggga ctttcctact tggcagtaca
 1981 tctacgtatt agtcatcgct attaccatgg tgatgcgggt ttggcagtag atcaatgggc
 2041 gtggatagcg gtttgactca cggggatttc caagtctcca cccattgac gtcaatggga
 2101 gtttgttttg gcaccaaatt caacgggact ttccaaaatg tcgtaacaac tccgcccat
 2161 tgacgcaaatt gggcggtagg cgtgtacggg gggaggttta tataagcaga gctcgtttag
 2221 tgaaccgtca gatcgcttg agacgccatc cagctgttt tgacctcat agaagattct
 2281 agagccgcca ccatgcttct cctggtgaca agccttctgc tctgtgagtt accacacca
 2341 gcattcctcc tgatcccaga catccagatg acacagacta catcctccct gtctgcctct
 2401 ctgggagaca gagtcccat cagttgcagg gcaagtcagg acattagtaa atatttaaat
 2461 tggatatcagc agaaaccaga tggaaactgt aaactcctga tctaccatac atcaagatta
 2521 cactcaggag tcccatcaag gttcagtggc agtgggtctg gaacagatta ttctctcacc
 2581 attagcaacc tggagcaaga agatattgcc acttactttt gccaacaggg taatacgctt
 2641 ccgtacacgt tcggaggggg gactaagttg gaaataacag gctccacctc tggatccggc
 2701 aagcccggat ctggcgaggg atccaccaag ggcgaggtga aactgcagga gtcaggacct
 2761 ggcctgggtg cgccctcaca gagcctgtcc gtcacatgca ctgtctcagg ggtctcatta
 2821 cccgactatg gtgtaagctg gattcgccag cctccacgaa agggctctgga gtggctggga
 2881 gtaatatggg gtagtgaaac cacatactat aattcagctc tcaaatccag actgaccatc

2941 atcaaggaca actccaagag ccaagttttc ttaaaaatga acagtctgca aactgatgac
 3001 acagccattt actactgtgc caaacattat tactacgggtg gtagctatgc tatggactac
 3061 tgggggtcaag gaacctcagt caccgtctcc tcagcggccg caattgaagt tatgtatcct
 3121 cctccttacc tagacaatga gaagagcaat ggaaccatta tccatgtgaa agggaaacac
 3181 ctttgtccaa gtcccttatt tcccggacct tctaagccct tttgggtgct ggtggtggtt
 3241 gggggaggtcc tggcttgcta tagcttgcta gtaacagtgg cttttattat tttctgggtg
 3301 aggagtaaga ggagcaggct cctgcacagt gactacatga acatgactcc ccgcccgc
 3361 gggcccaccc gcaagcatta ccagccctat gcccaccac gcgacttcgc agcctatcgc
 3421 tccagagtga agttcagcag gagcgcagac gccccgcgt accagcagg ccagaaccag
 3481 ctctataacg agctcaatct aggacgaaga gaggagtacg atgttttgga caagagacgt
 3541 ggccgggacc ctgagatggg gggaaagccg agaaggaaga accctcagga aggcctgtac
 3601 aatgaactgc agaaagataa gatggcggag gcctacagtg agattgggat gaaaggcgag
 3661 cgccggaggg gcaaggggca cgatggcctt taccagggtc tcagtacagc caccaaggac
 3721 acctacgacg cccttcacat gcaggccctg cccctcgtc aagaattcgg atccgcggcc
 3781 gcgaaggatc tgcgatcgtc ccggtgcccg tcagtgggca gagcgcacat cgcccacagt
 3841 ccccgagaag ttggggggag gggtcggcaa ttgaacgggt gcctagagaa ggtggcgagg
 3901 ggtaaactgg gaaagtgatg tcgtgtactg gctccgctt tttccgagg gtgggggaga
 3961 accgtatata agtgcagtag tcgccgtgaa cgttcttttt cgcaacgggt ttgcccag
 4021 aacacagctg aagcttcgag gggctcgcac ctctccttca cgcgcccgcc gccctacctg
 4081 aggcgcgcat ccacgcgggt tgagtcgctt tctgcccgt cccgctgtg gtgctcctg
 4141 aactgcgtcc gccgtctagg taagtttaaa gctcaggteg agaccgggccc tttgtccggc
 4201 gctcccttgg agcctaccta gactcagccg gctctccacg ctttgctga ccctgcttgc
 4261 tcaactctac gtctttgttt cgttttctgt tctgcgcgt tacagatcca agctgtgacc
 4321 ggcgcctacg ctagatgacc gagtacaagc ccacgggtgc cctgcgccac cgcgacgacg
 4381 tccccagggc cgtacgcacc ctgcgcgcgc cgttcgcgca ctaccccgcc acgcgccaca
 4441 ccgtcgatcc ggaccgccac atcgagcggg tcaccgagct gcaagaactc ttcctcacgc
 4501 gcgtcgggct cgacatcggc aaggtgtggg tcgcggacga cggcgccgcg gtggcggtct
 4561 ggaccacgcc ggagagcgtc gaagcggggg cgggtgttcg cgagatcggc ccgcgcagtg
 4621 ccgagttgag cggttcccgc ctggccgcgc agcaacagat ggaaggcctc ctggcgccgc
 4681 accggcccaa ggagcccgcg tggttcctgg ccaccgtcgg cgtctcgccc gaccaccagg
 4741 gcaaggggtc gggcagcgcc gtcgtgctcc ccggagtggg ggccggcgag cgcgcggggg
 4801 tgcccgctt cctggagacc tccgcgcccc gcaacctccc cttctacgag cggctcggt
 4861 tcaccgtcac cgcgcagctc gaggtgcccg aaggaccgcg cacctggtgc atgaccgca
 4921 agcccggtgc ctgagtcgac aatcaacctc tggattacaa aatttgtgaa agattgactg
 4981 gtattcttaa ctatgttgc ccttttacgc tatgtggata cgctgcttta atgcctttgt
 5041 atcatgctat tgcttcccgt atggctttca tttctcctc cttgtataaa tctgggttgc
 5101 tgtctcttta tgaggagtgt tggcccggtg tcaggcaacg tggcggtgtg tgcactgtgt
 5161 ttgttgacgc aacccccact ggttggggca ttgccaccac ctgtcagctc ctttcggga
 5221 ctttcgcttt cccctccctt attgccacgg cggaaactcat cgcgcctgc cttgcgcgt
 5281 gctggacagg ggctcggctg ttgggcactg acaattcgt ggtgttgtcg gggaaatcat
 5341 cgtcctttcc ttggctgtc gctgtgttg ccacctggat tctgcgcggg acgtccttct
 5401 gctacgtccc ttcggccctc aatccagcgg accttcttc ccgcggcctg ctgcgggctc
 5461 tgcggcctct tccgcgtctt cgccttcgce ctcagacgag tcggatctcc ctttgggccc
 5521 ctcccccgcc tggtagcttt aagaccaatg acttacaagg cagctgtaga tcttagccac
 5581 tttttaaaag aaaagggggg actggaaggg ctaattcact cccaacgaaa ataagatctg
 5641 ctttttgctt gtactgggtc tctctggtta gaccagatct gagcctggga gctctctggc
 5701 taactaggga acccactgct taagcctcaa taaagcttgc cttgagtgt tcaagtagtg
 5761 tgtgcccgtc tgttgtgtga ctctggtaac tagagatccc tcagaccctt ttagtcagtg
 5821 tggaaaatct ctagcagtag tagttcatgt catcttatta ttcagtattt ataacttgca
 5881 aagaaatgaa tatcagagag tgagaggaac ttgtttattg cagcttataa tgggtacaaa
 5941 taaagcaata gcatcacaaa tttcacaaat aaagcatttt tttcaactgca tctagttgt
 6001 ggtttgtcca aactcatcaa tgtatcttat catgtctggc tctagctatc ccgcccctaa
 6061 ctccgcccag ttcgcccctt tctccgcccc atggctgact aatttttttt atttatgcag
 6121 aggcgcaggc cgcctcggcc tctgagctat tccagaagta gtgaggaggc ttttttgag
 6181 gcctagactt ttgcagagac ggcccaaatt cgtaatcatg gtcatagetg tttcctgtgt
 6241 gaaattgtta tccgctcaca attccacaca acatacgagc cggaagcata aagtgtaaag
 6301 cctgggggtgc ctaatgagtg agctaactca cattaattgc gttgcgctca ctgcccgtt
 6361 tccagtcggg aaacctgtcg tgccagctgc attaataaat cggccaacgc gcggggagag
 6421 gcggtttgcg tattgggcgc tcttccgctt cctcgtctac tgactcgtg cgctcggtcg

6481 ttcggctgcg gcgagcggtta tcagctcact caaaggcggt aatacgggta tccacagaat
 6541 caggggataa cgcaggaaaag aacatgtgag caaaaggcca gcaaaaggcc aggaaccgta
 6601 aaaaggccgc gttgctggcg tttttccata ggctccgccc ccctgacgag catcacaaaa
 6661 atcgacgctc aagtcagagg tggcgaaaacc cgacaggact ataaagatac caggcggttc
 6721 cccctggaag ctccctcgtg cgctctcctg ttccgacctt gccgcttacc ggatacctgt
 6781 ccgcctttct cccttcggga agcgtggcgc tttctcatag ctacgctgtt aggtatctca
 6841 gttcgggtgta ggtcgttcgc tccaagctgg gctgtgtgca cgaaccccc gttcagcccg
 6901 accgctgcgc cttatccggg aactatcgtc ttgagtccaa cccggtaaga cagcacttat
 6961 cgccactggc agcagccact ggtaacagga ttagcagagc gaggtatgta ggcgggtgcta
 7021 cagagttctt gaagtgggtg cctaactacg gctacactag aaggacagta tttgggtatct
 7081 gcgctctgct gaagccagtt accttcggaa aaagagttgg tagctcttga tccggcaaac
 7141 aaaccaccgc tggtagcggg ggtttttttg tttgcaagca gcagattacg cgcagaaaaa
 7201 aaggatctca agaagatcct ttgatctttt ctacgggggtc tgacgctcag tggaaacgaaa
 7261 actcacgtta agggattttg gtcattgagat tatcaaaaag gatcttcacc tagatccttt
 7321 taaattaaaa atgaagtttt aaatcaatct aaagtatata tgagtaaact tgggtctgaca
 7381 gttaccaatg cttaatcagt gaggcaccta tctcagcgat ctgtctattt cgttcatcca
 7441 tagttgcctg actccccgtc gtgtagataa ctacgatacg ggagggtcta ccatctggcc
 7501 ccagtgtgctc aatgataccg cgagacccac gctcacgggc tccagattta tcagcaataa
 7561 accagccagc cgggaagggcc gagcgcagaa gtggtcctgc aactttatcc gcctccatcc
 7621 agtctattaa ttggtgcggg gaagctagag taagtagttc gccagttaat agtttgcgca
 7681 acgttggtgc cattgctaca ggcacgtggg tgtcacgctc gtcgtttggt atggttcat
 7741 tcagctccgg ttcccaacga tcaaggcgag ttacatgata ccccatgttg tgcaaaaaag
 7801 cggttagctc cttcggctct ccgacgttg tccagaagtaa gttggccgca gtgttatcac
 7861 tcatgggttat ggcagcactg cataattctc ttactgtcat gccatccgta agatgctttt
 7921 ctgtgactgg tgagtactca accaagtcac tctgagaata gtgtatgcgg cgaccgagtt
 7981 gctcttgccc ggcgtaataa cgggataata ccgcgccaca tagcagaact taaaagtgc
 8041 tcatcattgg aaaacgttct tcggggcgaa aactctcaag gatcttaccg ctgttgagat
 8101 ccagttcgat gtaaccactc cgtgcaccca actgatcttc agcatctttt actttcacca
 8161 gcgtttctgg gtgagcaaaa acaggaaggc aaaatgccgc aaaaaaggga ataaggcgca
 8221 cacggaaatg ttgaatactc atactcttcc tttttcaata ttattgaagc atttatcagg
 8281 gttattgtct catgagcgga tacatatattg aatgtattta gaaaaataaa caaatagggg
 8341 ttccgcgcac atttccccga aaagtgccac ctgacgtcta agaaaccatt attatcatga
 8401 cattaaccta taaaaatagg cgtatcacga ggccctttcg tctcgcgctt ttcggtgatg
 8461 acggtgaaaa cctctgacac atgcagctcc cggagacggg cacagcttgt ctgtaagcgg
 8521 atgccgggag cagacaagcc cgtcagggcg cgtcagcggg tgttgggcgg tgctggggct
 8581 ggcttaacta tgcggcatca gagcagattg tactgagagt gcaccatatg cgggtgtgaaa
 8641 taccgcacag atgcgtaagg agaaaatacc gcatcaggcg ccattcgcca ttcaggctgc
 8701 gcaactgttg ggaaggcgga tcggtgcggg cctcttcgct attacgccag ctggcgaaaag
 8761 ggggatgtgc tgcaaggcga ttaagttggg taacgccagg gttttccag tcacgacgtt
 8821 gtaaaacgac ggccagtgc aagctg.

p526A_19BBZ (SEQ ID NO: 10)

1 acgctgtag tcttatgcaa tactcttgta gtcttgcaac atggtaacga tgagttagca
 61 acatgcctta caaggagaga aaaagcaccg tgcatgccga ttgggtggaag taagggtgga
 121 cgatcgtgcc ttattaggaa ggcaacagac gggcttgaca tggattggac gaaccactga
 181 attgccgcat tgcagagata ttgtatttaa gtgcctagct cgatacaata aacgggtctc
 241 tctgggttaga ccagatctga gcctgggagc tctctggcta actaggggaa ccactgctta
 301 agcctcaata aagcttgccct tgagtgtctt aagtagtgtg tgcccgtctg ttgtgtgact
 361 ctggtaacta gagatccctc agaccctttt agtcagtgtg gaaaatctct agcagtggcg
 421 cccgaacagg gacctgaaag cgaaagggaa accagagctc tctcgacgca ggactcggct
 481 tgctgaagcg cgcacggcaa gaggcgaggg gcggcgactg gtgagtacgc caaaaatttt
 541 gactagcgga ggctagaagg agagagatgg gtgcgagagc gtcagtatta agcgggggag
 601 aattagatcg cgatgggaaa aaattcgggt aaggccaggg ggaaagaaaa aatataaatt
 661 aaaacatata gtatgggcaa gcaggagact agaacgattc gcagttaatc ctggcctgtt
 721 agaaacatca gaaggctgta gacaaatact gggacagcta caaccatccc ttcagacagg
 781 atcagaagaa cttagatcat tatataatac agtagcaacc ctctattgtg tgcataaag
 841 gatagagata aaagacacca aggaagcttt agacaagata gaggaagagc aaaacaaaag
 901 taagaccacc gcacagcaag cggccactga tcttcagacc tggaggagga gatatgaggg
 961 acaattggag aagtgaatta tataaatata aagtagtaaa aattgaacca ttaggagtag
 1021 caccaccaa ggcaagaga agagtgggtgc agagagaaaa aagagcagtg ggaataggag

```

1081 ctttgttctt tgggttcttg ggagcagcag gaagcactat gggcgagcgc tcaatgacgc
1141 tgacggtaca ggccagacaa ttattgtctg gtatagtcca gcagcagaac aatttgctga
1201 gggctattga ggcgcaacag catctgttgc aactcacagt ctggggcatc aagcagctcc
1261 aggcaagaat cctggctgtg gaaagatacc taaaggatca acagctcctg gggatttggg
1321 gttgctctgg aaaactcatt tgcaccactg ctgtgccttg gaatgctagt tggagtaata
1381 aatctctgga acagatttgg aatcacacga cctggatgga gtgggacaga gaaattaaca
1441 attacacaag cttaatacac tccttaattg aagaatcgca aaaccagcaa gaaaagaatg
1501 aacaagaatt attggaatta gataaatggg caagtttgtg gaattgggtt aacataacaa
1561 attggctgtg gtatataaaa ttattcataa tgatagtagg aggcttggta ggtttaagaa
1621 tagtttttgc tgtactttct atagtgaata gagttaggca gggatattca ccattatcgt
1681 ttcagaccca cctcccaacc ccgaggggac ccgacaggcc cgaaggaata gaagaagaag
1741 gtggagagag agacagagac agatccattc gattagtcaa cggatctcga cggtatcggg
1801 taacttttaa aagaaaaggg gggattgggg ggtacagtgc aggggaaaga atagtagaca
1861 taatagcaac agacatacaa actaaagaat taaaaaaca aattacaaaa ttcaaaattt
1921 tatcgatact agtggatctg cgatcgctcc ggtgcccgtc agtgggcaga gcgcacatcg
1981 cccacagtcc ccgagaagtt ggggggaggg gtccggcaatt gaacgggtgc ctagagaagg
2041 tggcgcgggg taaactggga aagtgatgtc gtgtactggc tccgcctttt tcccgagggt
2101 ggggggagaac cgtatataag tgcagtagtc gccgtgaacg ttcttttttc caacgggttt
2161 gccgccagaa cacagctgaa gcttcgaggg gctcgcattc ctcttcacg cgcccgccgc
2221 cctacctgag gccgccatcc acgcgggttg agtcgcgttc tgccgcctcc cgctgtgggt
2281 gcctcctgaa ctgcgtccgc cgtctaggta agtttaaagc tcaggtcgag accgggcctt
2341 tgcctcgccg tcccttggag cctacctaga ctacgcggc tctccacgct ttgctgacc
2401 ctgcttgctc aactctacgt ctttgttttc tttctgttc tgccgcgtta cagatccaag
2461 ctgtgaccgg cgcctactct agagccgcca ccatggccct gctgtgaca gctctgctgc
2521 tgcctctggc cctgctgctc catgccgcca gaccgatata ccagatgacc cagaccacca
2581 gcagcctgag cgcagcctg ggcgatagag tgaccatcag ctgccgggcc agccaggaca
2641 tcagcaagta cctgaactgg tatcagcaga aaccgcagcg caccgtgaag ctgctgatct
2701 accacaccag cagactgcac agcggcgtgc ccagcagatt ttctggcagc ggctccggca
2761 ccgactacag cctgaccatc tccaacctgg aacaggaaga tatcgctacc tacttctgtc
2821 agcaaggcaa caccctgccc tacaccttcg gcggaggcac caagctggaa atcacaggcg
2881 gcggaggatc tggcgagggt ggaagtggcg gaggcggcag cgaagtgaag ctgcaggaaa
2941 gcggccctgg cctgggtggc ccttctcagt ctctgtccgt gacctgtacc gtgtccggcg
3001 tgtccctgcc cgattatggc gtgtcctgga tccggcagcc tccagaaaag ggctcggaat
3061 ggctgggcgt gatctggggc agcgagacaa cctactacaa cagcgccctg aagtccggc
3121 tgaccatcat caaggacaa tccaagagcc aggtgttcct gaagatgaac agcctgcaga
3181 ccgacgacac cgcctctac tactgcgcca agcactacta ctacggcggc agctacgcca
3241 tggactactg gggccagggc accagcgtga ccgtgtctag cacaaccacc cctgcccta
3301 gacctccac cccagccca acaattgcca gccagcctct gtctctcgcg cccgaagctt
3361 gtagacctgc tgcggcgga gccgtgcaca ccagaggact ggatttcgcc tgcgacatct
3421 acatctgggc cctctgggc ggcacatgtg gcgtgtgtct cctcagcctg gtcacaccc
3481 tgtactgcaa gcggggcaga aagaaactgc tctacatctt caagcagccc ttcagcggc
3541 ccgtgcagac cacacaggaa gaggacggct gctcctgcag attccccgag gaagaagaag
3601 gcggctgcga gctgagagtg aagttcagca gatccgcga cgcctctgcc taccagcagg
3661 gacagaacca gctgtacaac gagctgaacc tgggcagacg ggaagagtac gacgtgctgg
3721 acaagcggag aggcagagat cccgagatgg gcggcaagcc cagacggaag aatccccagg
3781 aaggcctgta taacgaactg cagaaagaca agatggccga ggctacagc gagatcgga
3841 tgaaggcgca gcggagaaga ggcaagggcc acgatggcct gtaccagggc ctgagcaccg
3901 ccaccaagga cacctacgat gccctgcaca tgaggccct gccaccaga gaattcgaag
3961 gatccgcggc cgtgaggggc agaggaagtc ttctaactat cggtgacgtg gaggagaatc
4021 ccggcccttc cggaatggag agcgacgaga gcggcctgcc cgccatggag atcgagtgcc
4081 gcatcaccgg caccctgaac ggcgtggagt tcgagctggg gggcggcgga gagggcaccc
4141 ccaagcaggg ccgcatgacc aacaagatga agagcaccaa aggcgccctg accttcagcc
4201 cctacctgct gagccacgtg atgggctacg gcttctacca ctteggcacc taccaccagc
4261 gctacgagaa ccccttcttg cacgccatca acaacggcgg ctacaccaac accgcacatc
4321 agaagtacga ggacggcggc gtgctgcacg tgagcttcag ctaccgctac gaggcgggcc
4381 gcgtgatcgg cgacttcaag gtggtgggca ccggcttccc cgaggacagc gtgatcttca
4441 ccgacaagat catccgcagc aacgccaccg tggagcacct gcaccccatg ggcgataacg
4501 tgctgggtgg cagcttcgcc cgcaccttca gcctgcgcga cggcggttac tacagcttcg
4561 tgggtggacag ccacatgcac ttcaagagcg ccatccaccc cagcatcctg cagaacgggg

```

4621 gccccatggt cgccttccgc cgcgtggagg agctgcacag caacaccgag ctgggcatcg
 4681 tggagtacca gcacgccttc aagaccccca tcgccttcgc cagatcccg cgtcagtcgt
 4741 ccaattctgc cgtggacggc accgccggac ccggctccac cggatctcgc tagagctgaa
 4801 tctaagtcga caatcaacct ctggattaca aaatttgtga aagattgact ggtattctta
 4861 actatgttgc tccttttacg ctatgtggat acgctgcttt aatgcctttg tatcatgcta
 4921 ttgcttcccg tatggctttc attttctcct ccttgataaa atcctgggtg ctgtctcttt
 4981 atgaggagtt gtggcccggt gtcaggcaac gtggcgtggg gtgcactgtg tttgctgacg
 5041 caacccccac tggttggggc attgccacca cctgtcagct cctttccggg actttcgcct
 5101 tccccctccc tattgccacg gcggaactca tcgccgcctg ccttgcccg cgtgagacag
 5161 gggctcggct gttgggcact gacaattccg tgggtgtgtc ggggaaatca tcgtcctttc
 5221 cttggctgct cgcctgtgtt gccacctgga ttctgcgcgg gacgtccttc tgctacgtcc
 5281 cttcggccct caatccagcg gaccttcctt cccgcggcct gctgcgggct ctgcggcctc
 5341 ttccgcgtct tcgccttcgc cctcagacga gtccgatctc cctttgggcc gcctccccgc
 5401 ctggtacctt taagaccaat gacttacaag gcagctgtag atcttagcca ctttttaaaa
 5461 gaaaaggggg gactggaagg gctaattcac tcccaacgaa aataagatct gctttttgct
 5521 tgtactgggt ctctctgggt agaccagatc tgagcctggg agctctctgg ctaactaggg
 5581 aacccactgc ttaagcctca ataaagcttg ccttgagtgc ttcaagtagt gtgtgcccg
 5641 ctggtgtgtg actctggtaa ctagagatcc ctacagacct ttagtcagt gtggaaaatc
 5701 tctagcagta gtatgtcatg tcatcttatt attcagtatt tataacttgc aaagaaatga
 5761 atatcagaga gtgagaggaa cttgtttatt gcagcttata atggttacaa ataaagcaat
 5821 agcatcacia atttcacaaa taaagcattt ttttcaactg attctagttg tggtttgtcc
 5881 aaactcatca atgtatctta tcatgtctgg ctctagctat cccgccctta actccgccca
 5941 gtcccgccca ttctccgccc catggctgac taattttttt tatatatgca gaggccgagg
 6001 ccgcctcggc ctctgagcta ttccagaagt agtgaggagg cttttttgga ggccctagact
 6061 tttgcagaga cggcccaaat tcgtaatcat ggtcatagct gtttcctgtg tgaaattgtt
 6121 atccgctcac aattccacac aacatacgag ccggaagcat aaagtgtaaa gcctggggtg
 6181 cctaatagag gagctaactc acattaattg cgttgcgctc actgcccgct ttccagtcgg
 6241 gaaacctgtc gtgccagctg cattaatgaa tcggccaacg cgcggggaga ggcgggttgc
 6301 gtattgggcg ctcttcgctt tccctcgctca ctgactcgct gcgctcggtc gttcggctgc
 6361 ggcgagcggg atcagctcac tcaaaggcgg taatacgggt atccacagaa tcaggggata
 6421 acgcaggaaa gaacatgtga gcaaaaaggc agcaaaaagg caggaaacct aaaaaggccg
 6481 cgttgctggc gtttttccat aggctccgcc cccctgacga gcatcaciaa aatcgacgct
 6541 caagtcagag gtggcgaaac ccgacaggac tataaagata ccaggcgttt cccctggaa
 6601 gctccctcgt gcgctctcct gttccgacce tgccgcttac cggataacct tccgcctttc
 6661 tcccttcggg aagcgtggcg ctttctcata gctcacgctg taggtatctc agttcggtgt
 6721 aggtcgttcg ctccaagctg ggctgtgtgc acgaaccccc cgttcagccc gaccgctgcg
 6781 ccttatccgg taactatcgt cttgagtcca acccggttag acacgactta tcgccactgg
 6841 cagcagccac tggtaacagg attagcagag cgaggtatgt aggcgggtgct acagagtctt
 6901 tgaagtgggt gcctaactac ggctacacta gaaggacagt atttgggtatc tgcgctctgc
 6961 tgaagccagt taccttcgga aaaagagttg gtatctcttg atccggcaaa caaaccaccg
 7021 ctggtagcgg tggttttttt gtttgcaagc agcagattac gcgcagaaaa aaaggatctc
 7081 aagaagatcc tttgatcttt tctacggggg ctgacgctca gtggaacgaa aactcacgtt
 7141 aagggaattt ggtcatgaga ttatcaaaaa ggatcttcac ctagatcctt ttaaattaaa
 7201 aatgaagttt taaatcaatc taaagtatat atgagtaaac ttggtctgac agttaccaat
 7261 gcttaatcag tgaggcacct atctcagcga tctgtctatt tcgttcaccc atagttgcct
 7321 gactccccgt cgtgtagata actacgatac gggagggctt accatctggc cccagtgcgt
 7381 caatgatacc gcgagacca cgtcaccggc ctccagattt atcagcaata aaccagccag
 7441 ccggaagggc cgagcgacga agtggtcctg caactttatc cgcctccatc cagtctatta
 7501 attgttgccg ggaagctaga gtaagtagtt cgccagttaa tagtttgccg aacgttgttg
 7561 ccattgctac aggcacgtg gtgtcacgct cgtcgtttgg tatggcttca ttcagctccg
 7621 gttcccaacg atcaaggcga gttacatgat ccccatggtt gtgcaaaaaa gcggttagct
 7681 cttcgggtcc tccgatcgtt gtcagaagta agttggccgc agtggttatca ctcatggtta
 7741 tggcagcact gcataattct cttactgtca tgccatccgt aagatgcttt tctgtgactg
 7801 gtgagtactc aaccaagtca ttctgagaat agtgtatgcg gcgaccgagt tgctcttgcc
 7861 cggcgtcaat acgggataat accgcgccac atagcagaac tttaaaagtg ctcatcattg
 7921 gaaaacgttc ttcggggcga aaactctcaa ggatcttacc gctgttgaga tccagttcga
 7981 tgtaaccacac tcgtgcaccc aactgatctt cagcatcttt tactttcacc agcgtttctg
 8041 ggtgagcaaa aacaggaagg caaaatgccg caaaaaaggg aataaggggc acacggaaat
 8101 gttgaatact catactcttc ctttttcaat attattgaag catttatcag ggttattgtc

8161 tcatgagcgg atacatatattt gaatgtattt agaaaaataa acaaataggg gtccgcgca
 8221 catttccccg aaaagtgcc aatgacgtct aagaaacat tattatcatg acattaacct
 8281 ataaaaatag gcgtatcacg aggccctttc gtctcgcgcg ttccggtgat gacggtgaaa
 8341 acctctgaca catgcagctc ccggagacgg tcacagcttg tctgtaagcg gatgcgggga
 8401 gcagacaagc ccgtcagggc gcgtcagcgg gtgttgccgg gtgtcggggc tggcttaact
 8461 atgcggcatc agagcagatt gtactgagag tgcaccatat gcggtgtgaa ataccgcaca
 8521 gatgcgtaag gagaaaaatac cgcacagggc gccattcggc attcaggctg cgcaactgtt
 8581 gggaagggcg atcgggtgcgg gcctcttcgc tattacgcca gctggcgaaa gggggatgtg
 8641 ctgcaaggcg attaagttgg gtaacgccag ggttttccca gtcacgacgt tgtaaaacga
 8701 cggccagtgc caagctg.

TFP (TRuC) constructs***p510_antiCD19_LL_TCRalpha* (SEQ ID NO: 11)**

1 acgcgtgtag tcttatgcaa tactcttgta gtcttgcaac atggtaacga tgagttagca
 61 acatgcctta caaggagaga aaaagcaccg tgcattgccga ttgggtggaag taagggtgga
 121 cgatcgtgcc ttattaggaa ggcaacagac gggctctgaca tggattggac gaaccactga
 181 attgccgcat tgcagagata ttgtatttaa gtgcctagct cgatacaata aacgggtctc
 241 tctggttaga ccagatctga gcctgggagc tctctggcta actagggaac ccactgctta
 301 agcctcaata aagcttgcc ttagtgcttc aagtagtgtg tgcccgctctg ttgtgtgact
 361 ctggttaacta gagatccctc agaccctttt agtcagtgtg gaaaatctct agcagtggcg
 421 cccgaacagg gacctgaaag cgaaagggaa accagagctc tctcgacgca ggactcggct
 481 tgctgaagcg cgcacggcaa gaggcgaggg gcggcgactg gtgagtacgc caaaaaattt
 541 gactagcggg ggctagaagg agagagatgg gtgcgagagc gtcagtatta agcgggggag
 601 aattagatcg cgatgggaaa aaattcgggt aaggccaggg ggaaagaaaa aatataaatt
 661 aaaacatata gtatgggcaa gcagggagct agaacgattc gcagttaatc ctggcctgtt
 721 agaaacatca gaaggctgta gacaaatact gggacagcta caaccatccc ttcagacagg
 781 atcagaagaa cttagatcat tatataatac agtagcaacc ctctattgtg tgcataaag
 841 gatagagata aaagacacca aggaagcttt agacaagata gaggaagagc aaaacaaaag
 901 taagaccacc gcacagcaag cggccactga tcttcagacc tggaggagga gatattgagg
 961 acaattggag aagtgaatta tataaatata aagtagtaaa aattgaacca ttaggagtag
 1021 caccaccaa ggcaagaga agagtgggtg agagagaaaa aagagcagtg ggaataggag
 1081 ctttgttctc tgggttcttg ggagcagcag gaagcactat gggcgagcc tcaatgacgc
 1141 tgacggtaca ggccagacaa ttattgtctg gtatagtga gcagcagaac aatttgctga
 1201 gggctattga ggcgcaacag catctgttgc aactcacagt ctggggcatc aagcagctcc
 1261 aggcaagaat cctggctgtg gaaagatacc taaaggatca acagctcctg gggatttggg
 1321 gttgctctgg aaaactcatt tgcaccactg ctgtgccttg gaatgctagt tggagtaata
 1381 aatctctgga acagattgga atcacacgac ctggatggag tgggacagag aaattaacaa
 1441 ttacacaagc ttaatacact ccttaattga agaatcgcaa aaccagcaag aaaagaatga
 1501 acaagaatta ttggaattag ataaatgggc aagtttgttg aattgggtta acataacaaa
 1561 ttggctgttg tatataaaat tattcataat gatagtagga ggcttggtag gtttaagaat
 1621 agtttttctg gtactttcta tagtgaatag agttaggcag ggatattcac cattatcggt
 1681 tcagaccac ctcccaacc cgaggggacc cgacaggccc gaaggaatag aagaagaagg
 1741 tggagagaga gacagagaca gatccattcg attagtgaac ggatctcgac ggtatcggtt
 1801 aactttttaa agaaaagggg ggattggggg gtacagtga ggggaaagaa tagtagacat
 1861 aatagcaaca gacatacaaa ctaaagaatt acaaaaacaa attacaaaat tcaaaatttt
 1921 atcgatacta gtattatgcc cagtacatga ccttatggga ctttctact tggcagtaca
 1981 tctacgtatt agtcacgtct attaccatgg tgatgcgggt ttggcagtag atcaatgggc
 2041 gtggatagcg gtttgactca cggggatttc caagtctcca cccattgac gtcaatggga
 2101 gtttggtttg gcacccaaat caacgggact ttccaaaatg tcgtaacaac tccgccccat
 2161 tgacgcaaat gggcggtagg cgtgtacggg gggaggttta tataagcaga gctcgtttag
 2221 tgaaccgtca gatcgcttg agacgccatc cacgtgttt tgacctccat agaagattct
 2281 agagccgcca ccatgcttct cctgggtgaca agccttctgc tctgtgagtt accacaccca
 2341 gcattcctcc tgatcccaga catccagatg acacagacta catcctccct gtctgcctct
 2401 ctgggagaca gagtcccat cagttgcagg gcaagtcagg acattagtaa atatttaaat
 2461 tggatatcagc agaaaccaga tggaaactgt aaactcctga tctaccatac atcaagatta
 2521 cactcaggag tcccatcaag gttcagtggc agtgggtctg gaacagatta ttctctcacc
 2581 attagcaacc tggagcaaga agatattgcc acttactttt gccaacaggg taatacgtct
 2641 ccgtacacgt tccgaggggg gactaagttg gaaataacag gctccacctc tggatccggc
 2701 aagcccgat ctggcgaggg atccaccaag ggcgaggtga aactgcagga gtcaggacct
 2761 ggcctgggtg cgcctcaca gagcctgtcc gtcacatgca ctgtctcagg ggtctcatta

2821 cccgactatg gtgtaagctg gattcgccag cctccacgaa agggctctgga gtggctggga
 2881 gtaatatggg gtagtgaaac cacatactat aattcagctc tcaaatccag actgaccatc
 2941 atcaaggaca actccaagag ccaagttttc ttaaaaatga acagtctgca aactgatgac
 3001 acagccattt actactgtgc caaacattat tactacgggtg gtagctatgc tatggactac
 3061 tgggggtcaag gaacctcagt caccgtctcc tcagcggccg caattgaagt tatgtatcct
 3121 cctccttacc taggtggcgg cggttctggg ggcgccgggt ctgggtggcgg cggttctctc
 3181 gaggtgaatg gagagaatgt ggagcagcat cttcaaccc tgagtgtcca ggaggagac
 3241 agcgtgttta tcaagtgtac ttattcagac agtgectcaa actacttccc ttggtataag
 3301 caagaacttg gaaaaagacc tcagcttatt atagacattc gttcaaatgt gggcgaaaag
 3361 aaagaccaac gaattgctgt tacattgaac aagacagcca aacatttctc cctgcacatc
 3421 acagagaccc aacctgaaga ctcggtgtgc tacttctgtg cagcaagtag gaaggactct
 3481 ggggggttacc agaaagttaac ctttggaact ggaacaaagc tccaagtcac cccaaatatac
 3541 cagaaccctg accctgccgt gtaccagctg agagactcta aatccagtga caagtctgtc
 3601 tgcctattca ccgattttga ttctcaaaca aatgtgtcac aaagtaagga ttctgatgtg
 3661 tatatcacag acaaaaactgt gctagacatg aggtctatgg acttcaagag caacagtgtc
 3721 gtggcctgga gcaacaaatc tgactttgca tgtgcaaacg cttcaacaa cagcattatt
 3781 ccagaagaca ccttcttccc cagcccagaa agttcctgtg atgtcaagct ggtcgagaaa
 3841 agctttgaaa cagatacgaa cctaaacttt caaaacctgt cagtgattgg gttccgaatc
 3901 ctctctctga aagtggccgg gtttaatctg ctcatgacgc tgcggtgtg gtccagctga
 3961 taagaattcg atccgcggcc gcgaaggatc tgcgctcgt ccggtgcccg tcagtgggca
 4021 gagegcacat cgcaccagat ccccgagaag ttggggggag gggctggcaa ttgaacgggt
 4081 gcctagagaa ggtggcgcgg ggtaaactgg gaaagtgatg tctgtactg gctccgcctt
 4141 tttcccgagg gtgggggaga accgtatata agtgacgtag tgcgctgaa cgttctttt
 4201 cgcaacgggt ttgcccagag aacacagctg aagcttcgag gggctcgcat ctctccttca
 4261 cgcgcccggc gccctacctg aggcggccat ccacgcgggt tgagtcgct tctgcccgt
 4321 cccgctgtg gtgctctctg aactgcgtcc gccgtctagg taagttttaa gctcaggtcg
 4381 agaccgggccc tttgtccggc gctcccttgg agcctaccta gactcagccg gctctccacg
 4441 ctttgctga ccctgcttgc tcaactctac gtctttgttt cgttttctgt tctgcccgt
 4501 tacagatcca agctgtgacc ggcgcctacg ctatagatgacc gagtacaagc ccacgggtgcg
 4561 cctcgccacc cgcgacgacg tcccagggc cgtacgcacc ctgcgcgcg cgttcgcca
 4621 ctaccccgcc acgcgccaca ccgtcgatcc ggaccgccac atcgagcggg tcaccgagct
 4681 gcaagaactc ttcctcacgc gcgtcgggct cgacatcggc aaggtgtggg tgcgggacga
 4741 cggcgccgcg gtggcggtct ggaccacgce ggagagcgtc gaagcggggg cgggtgtcgc
 4801 cgagatcggc ccgcgcattg ccgagttgag cggttcccg ctggccgcgc agcaacagat
 4861 ggaaggcctc ctggcgccgc accggcccaa ggagcccgcg tggttcctgg ccaccgtcg
 4921 cgtctcgccc gaccaccagg gcaagggtct gggcagcgc gtcgtgctcc ccggagtga
 4981 ggcggccgag cgcgcggggg tgcccgcctt cctggagacc tccgcgccc gcaacctccc
 5041 cttctacgag cggctcggct tcaccgtcac cgcgacgtc gaggtgccc aaggaccgcg
 5101 cactgggtgc atgacccgca agcccgggtg ctgagtcgac aatcaacctc tggattacaa
 5161 aatttgtgaa agattgactg gtattcttaa ctatgttgc cttttacgc tatgtggata
 5221 cgctgcttta atgcctttgt atcatgctat tgcttccgt atggctttca ttttctctc
 5281 cttgtataaa tccgtggtgc tgtctcttta tgaggagtgt tggeccgttg tcaggcaacg
 5341 tggcgtggtg tgcactgtgt ttgctgacgc aaccccact ggttggggca ttgccaccac
 5401 ctgtcagctc ctttccggga ctttcgcttt cccctccct attgccacgg cggaaactcat
 5461 cgccgcctgc cttgcccgct gctggacagg ggctcggctg ttgggcactg acaattccgt
 5521 ggtgttgtcg gggaaatcat cgtcctttcc ttggtgctc gctgtgttg ccacctggat
 5581 tctgcgcggg acgtccttct gctacgtccc ttcggccctc aatccagcgg accttcttc
 5641 ccgcggcctg ctgcgcgctc tgcggcctct tccgcgtct cgccttcgcc ctacagcag
 5701 tcgcatctcc ctttgggcgg cctcccgcgc tggtaacctt aagaccaatg acttacaagg
 5761 cagctgtaga tcttagccac tttttaaaag aaaagggggg actggaaggg ctaattcact
 5821 cccaacgaaa ataagatctg ctttttgcct gtactgggtc tctctggtta gaccagatct
 5881 gagcctggga gctctctggc taactagggg acccactgct taagcctcaa taaagcttgc
 5941 cttgagtgtc tcaagtagtg tgtgcccgtc tgttgtgtga ctctggtaac tagagatccc
 6001 tcagaccctt ttagtcagtgt tggaaaatct ctagcagtag tagttcatgt catcttatta
 6061 ttcagtattt ataacttgca aagaaatgaa tatcagagag tgagaggaaac ttgtttattg
 6121 cagcttataa tggttacaaa taaagcaata gcatcacaaa tttcacaaat aaagcatttt
 6181 tttcactgca ttctagttgt ggtttgtcca aactcatcaa tgtatcttat catgtctggc
 6241 tctagctatc ccgcccctaa ctccgcccag ttcgcccct tctccgccc atggctgact
 6301 aatttttttt atttatgcag aggcggagge cgctcggcc tctgagctat tccagaagta

6361 gtgaggaggc ttttttggag gcctagactt ttgcagagac ggcccaaatt cgtaatcatg
6421 gtcatagtctg tttcctgtgt gaaattgtta tccgtcaca attccacaca acatacgagc
6481 cggaagcata aagtgtaaag cctgggggtgc ctaatgagtg agctaactca cattaattgc
6541 gttgcgctca ctgcccgtt tccagtcggg aaacctgtcg tgccagctgc attaatgaat
6601 cggccaacgc gcggggagag gcggtttgctg tattgggcgc tcttcgctt cctcgctcac
6661 tgactcgctg cgctcggtcg ttcggctgctg gcgagcggta tcagctcact caaaggcggg
6721 aatacgggta tccacagaat caggggataa cgcaggaaag aacatgtgag caaaaggcca
6781 gcaaaaggcc aggaaccgta aaaaggccgc gttgctggcg tttttccata ggctccgccc
6841 ccctgacgag catcacaaaa atcgacgctc aagtcagagg tggcgaaacc cgacaggact
6901 ataaagatac caggcgtttc cccctggaag ctccctcgtg cgctctcctg ttccgaccct
6961 gccgcttacc ggatacctgt ccgcctttct ccttcggga agcgtggcg tttctcatag
7021 ctcaogctgt aggtatctca gttcgggtgta ggtcgttcgc tccaagctgg gctgtgtgca
7081 cgaaccccc gttcagcccg accgctgcgc cttatccggg aactatcgtc ttgagtccea
7141 cccggtaaga cagacttat cgccactggc agcagccact ggtaacagga ttagcagagc
7201 gaggtatgta ggcgggtgcta cagagttctt gaagtgggtg cctaactacg gctacactag
7261 aaggacagta tttgggtatct gcgctctgct gaagccagtt accttcggaa aaagagttgg
7321 tagctcttga tccggcaaac aaaccaccgc tggtagcggg gggttttttg tttgcaagca
7381 gcagattacg cgcagaaaaa aaggatctca agaagatcct ttgatctttt ctacggggctc
7441 tgacgctcag tggaaacgaaa actcacgtta agggattttg gtcatgagat tatcaaaaag
7501 gatcttcacc tagatccttt taaattaaaa atgaagtttt aaatcaatct aaagtatata
7561 tgagtaaaact tgggtctgaca gttaccaatg cttaatcagt gaggcacct tctcagcgat
7621 ctgtctatctt cgttcatcca tagttgcctg actccccgtc gtgtagataa ctacgatacg
7681 ggagggtctta ccatctggcc ccagtgcctgc aatgataccg cgagaccac gctcaccggc
7741 tccagattta tcagcaataa accagccagc cggaaggggc gagcgcagaa gtggctcctgc
7801 aacttttatcc gcctccatcc agtctattaa ttgttgccgg gaagctagag taagtagttc
7861 gccagttaat agtttgcgca acgttggtgc cattgctaca ggcacgtgg tgtcacgctc
7921 gtcgtttggg atggcttcat tcagctccgg tcccaacga tcaaggcgag ttacatgatc
7981 ccccatgttg tgcaaaaaag cgggttagctc cttcggctct ccgatcggtg tcagaagtaa
8041 gttggccgca gtgttatcac tcatggttat ggcagcactg cataattctc ttactgtcat
8101 gccatccgta agatgctttt ctgtgactgg tgagtactca accaagtcac tctgagaata
8161 gtgtatgcgg cgaccgagtt gctcttgccc ggcgtcaata cgggataata ccgcgccaca
8221 tagcagaact ttaaaagtgc tcatcattgg aaaacgttct tcggggcgaa aactctcaag
8281 gatcttaccg ctggtgagat ccagttcgat gtaaccact cgtgcacca actgatcttc
8341 agcatctttt actttcacca gcgtttctgg gtgagcaaaa acaggaaggc aaaatgccgc
8401 aaaaaaggga ataaggcgca cacggaaatg ttgaatactc atactcttcc tttttcaata
8461 ttattgaagc atttatcagg gttattgtct catgagcgga tacatatttg aatgtattta
8521 gaaaaataaa caaatagggg ttccgcgcac atttccccga aaagtgcac ctgacgtcta
8581 agaaaccatt attatcatga cattaacctc taaaaatagg cgtatcacga ggccctttcg
8641 tctcgcgcgt ttcgggtgat acggtgaaaa cctctgacac atgcagctcc cggagacggg
8701 cacagcttgt ctgtaagcgg atgccgggag ggccttaacta tgcggcatca gagcagattg tactgagagt
8761 tgttgccggg tgtcggggct ggccttaacta atgcgtaagg agaaaatacc gcatcaggcg
8821 gcaccatatt cgggtgtgaa taccgcacag ggaaggcgga tcgggtgcgg cctcttcgct
8881 ccattcgcca ttcaggctgc gcaactgttg ggaaggcgga tgaaggcgga ttaagttggg taacgccagg
8941 attacgccag ctggcgaaag ggggatgtgc tgcaaggcgga aagctg.

p510_antiCD19_LL_TCRalphaC (SEQ ID NO: 12)

1 acgctgttag tcttatgcaa tactcttgta gtcttgcaac atggtaacga tgagttagca
61 acatgcctta caaggagaga aaaagcacccg tgcattgcga ttgggtggaag taagggtgga
121 cgatcggtcc ttattaggaa ggcaacagac gggctctgaca tggattggac gaaccactga
181 attgccgcat tgcagagata ttgtatttaa gtgcctagct cgatacaata aacgggtctc
241 tctgggttaga ccagatctga gcctgggagc tctctggcta actagggaac ccactgctta
301 agcctcaata aagcttgctt tgagtgtctt aagtagtgtg tgcccgtctg ttgtgtgact
361 ctggtaacta gagatccctc agaccctttt agtcagtggt gaaaatctct agcagtgggc
421 cccgaacagg gacctgaaag cgaaagggaa accagagctc tctcgacgca ggactcggct
481 tgctgaagcg cgcacggcaa gaggcgaggg gcggcgactg gtgagtacgc caaaaatttt
541 gactagcgga ggctagaagg agagagatgg gtgcgagagc gtcagtatta agcgggggag
601 aattagatcg cgatgggaaa aaattcgggt aaggccaggg ggaaagaaaa aatataaatt
661 aaaacatata gtatgggcaa gcaggagctc agaacgattc gcagttaatc ctggcctgtt
721 agaaacatca gaaggctgta gacaaatact gggacagcta caaccatccc ttcagacagg

```

781  atcagaagaa  cttagatcat  tatataatac  agtagcaacc  ctctattgtg  tgcacaaag
841  gatagagata  aaagacacca  aggaagcttt  agacaagata  gaggaagagc  aaaacaaaag
901  taagaccacc  gcacagcaag  cggccactga  tcttcagacc  tggaggagga  gatatgaggg
961  acaattggag  aagtgaatta  tataaatata  aagtagtaaa  aattgaacca  ttaggagtag
1021 caccacacaa  ggcaaagaga  agagtgggtg  agagagaaaa  aagagcagtg  ggaataggag
1081 ctttgttctt  tgggttcttg  ggagcagcag  gaagcactat  gggcgagcc  tcaatgacgc
1141 tgacggtaca  ggccagacaa  ttattgtctg  gtatagtgc  gcagcagaac  aatttgctga
1201 gggctattga  ggcgcaacag  catctgttgc  aactcacagt  ctggggcatc  aagcagctcc
1261 aggcaagaat  cctggctgtg  gaaagatacc  taaaggatca  acagctcctg  gggatttggg
1321 gttgctctgg  aaaactcatt  tgcaccactg  ctgtgccttg  gaatgctagt  tggagtaata
1381 aatctctgga  acagattgga  atcacacgac  ctggatggag  tgggacagag  aaattaacaa
1441 ttacacaagc  ttaatacact  ccttaattga  agaatcgcaa  aaccagcaag  aaaagaatga
1501 acaagaatta  ttggaattag  ataaatgggc  aagtttgtgg  aattggttta  acataacaaa
1561 ttggctgtgg  tatataaaat  tattcataat  gatagtagga  ggcttggtag  gtttaagaat
1621 agtttttgct  gtacttttcta  tagtgaatag  agttaggcag  ggatattcac  cattatcggt
1681 tcagacccac  ctcccaaccc  cgaggggacc  cgacaggccc  gaaggaatag  aagaagaagg
1741 tggagagaga  gacagagaca  gatccattcg  attagtgaac  ggatctcgac  ggtatcgggt
1801 aactttttaa  agaaaagggg  ggattggggg  gtacagtgc  ggggaaagaa  tagtagacat
1861 aatagcaaca  gacatacaaa  ctaaagaatt  acaaaaacaa  attacaaaat  tcaaaatttt
1921 atcgatacta  gtattatgcc  cagtacatga  ccttatggga  ctttcctact  tggcagtaca
1981 tctacgtatt  agtcatcgct  attaccatgg  tgatgcgggt  ttggcagtac  atcaatgggc
2041 gtggatagcg  gtttgactca  cggggatttc  caagtctcca  cccattgac  gtcaatggga
2101 gtttgttttg  gcacccaaat  caacgggact  ttccaaaatg  tcgtaacaac  tccgccccat
2161 tgacgcaaat  gggcggtagg  cgtgtacgg  gggaggttta  tataagcaga  gctcgtttag
2221 tgaaccgtca  gatcgcttg  agacgccatc  cacgtgttt  tgacctccat  agaagattct
2281 agagccgcca  ccatgcttct  cctggtgaca  agccttctgc  tctgtgagtt  accacacca
2341 gcattcctcc  tgatcccaga  catccagatg  acacagacta  catcctccct  gtctgcctct
2401 ctgggagaca  gagtcaccat  cagttgcagg  gcaagtcagg  acattagtaa  atattttaat
2461 tgggtatcagc  agaaaccaga  tggaaactgt  aaactcctga  tctaccatac  atcaagatta
2521 cactcaggag  tcccatcaag  gttcagtggc  agtgggtctg  gaacagatta  ttctctcacc
2581 attagcaacc  tggagcaaga  agatattgcc  acttactttt  gccaacaggg  taatacgtct
2641 ccgtacacgt  tcggaggggg  gactaagttg  gaaataacag  gctccacctc  tggatccggc
2701 aagcccggat  ctggcgaggg  atccaccaag  ggcgaggtga  aactgcagga  gtcaggacct
2761 ggcctgggtg  cgccctcaca  gagcctgtcc  gtcacatgca  ctgtctcagg  ggtctcatta
2821 cccgactatg  gtgtaagctg  gattcgccag  cctccacgaa  agggctctga  gtggctggga
2881 gtaatatggg  gtagtgaaac  cacatactat  aattcagctc  tcaaatccag  actgaccatc
2941 atcaaggaca  actccaagag  ccaagtttct  ttaaaaatga  acagtctgca  aactgatgac
3001 acagccattt  actactgtgc  caaacattat  tactacgggt  gtagctatgc  tatggactac
3061 tgggggtcaag  gaacctcagt  caccgtctcc  tcagcgccg  caattgaagt  tatgtatcct
3121 cctcettacc  taggtggcgg  cgggtctggg  ggcggcggtt  ctgggtggcg  cgggtctctc
3181 gagccaaata  tccagaaccc  tgacctgccc  gtgtaccagc  tgagagactc  taaatccagt
3241 gacaagtctg  tctgcctatt  caccgatttt  gattctcaaa  caaatgtgtc  acaaagtaag
3301 gattctgatg  tgtatatcac  agacaaaact  gtgctagaca  tgaggtctat  ggacttcaag
3361 agcaacagtg  ctgtggcctg  gagcaacaaa  tctgactttg  catgtgcaaa  cgcttcaac
3421 aacagcatta  ttccagaaga  caccttcttc  ccagcccag  aaagtctctg  tgatgtcaag
3481 ctggctcgaga  aaagctttga  aacagatacg  aacctaaact  ttcaaaacct  gtcagtgatt
3541 gggttccgaa  tcctcctcct  gaaagtggcc  gggtttaate  tgctcatgac  gctgcggctg
3601 tggctccagct  gataagaatt  cgatccgcgg  ccgcgaagga  tctgcgatcg  ctccgggtgc
3661 cgtcagtggg  cagagcgcac  atcgcccaca  gtccccgaga  agttgggggg  aggggtcggc
3721 aattgaacgg  gtgcctagag  aaggtggcgc  ggggtaaaact  gggaaagtga  tgtcgtgtac
3781 tggctccgcc  tttttcccg  ggggtggggg  gaaccgtata  taagtgcagt  agtcgccgtg
3841 aacgttcttt  ttcgcaacgg  gtttgccgcc  agaacacagc  tgaagcttcg  aggggctcgc
3901 atctctcctt  caecgcgccg  ccgccttacc  tgaggccgcc  atccacgccg  gttgagtcgc
3961 gttctgccgc  ctcccgctg  tgggtgcctc  tgaactgcgt  ccgcctctta  ggtaagttta
4021 aagctcaggt  cgagaccggg  cctttgtccg  gcgtccctt  ggagcctacc  tagactcagc
4081 cggctctcca  cgctttgcct  gacctgtct  gctcaactct  acgtctttgt  ttcgttttct
4141 gttctgccgc  gttacagatc  caagctgtga  ccggcgccct  cgctagatga  ccgagtacaa
4201 gccacgggtg  cgctcgcca  ccgcgcagca  cgtccccagg  gccgtacgca  ccctcgccgc
4261 cgcgttcgcc  gactaccccg  ccacgcgcca  caccgtcgat  ccggaccgcc  acatcgagcg

```

4321 ggtcaccgag ctgcaagaac tcttcctcac gcgcgtcggg ctcgacatcg gcaaggtgtg
 4381 ggtcgcggac gacggcgccg cgggtggcgg ctggaccacg ccggagagcg tcgaagcggg
 4441 ggcgggtgttc gccgagatcg gccgcgcgat ggccgagttg agcggttccc ggctggccgc
 4501 gcagcaacag atggaaggcc tcttggcgcg gcaccggccc aaggagcccc cgtggttcct
 4561 ggccaccgtc ggcgctctcg ccgaccacca gggcaagggt ctgggcagcg ccgtcgtgct
 4621 ccccgagtg gaggcggccg agcgcgcggg ggtgcccgcc ttcctggaga cctccgcgcc
 4681 ccgcaacctc cccttctacg agcggctcgg cttaccgctc accgccgacg tcgaggtgcc
 4741 cgaaggaccg cgcacctggt gcatgaccgg caagcccggt gcctgagtcg acaatcaacc
 4801 tctggattac aaaatttgtg aaagattgac tggattctt aactatgttg ctcttttac
 4861 gctatgtgga tacgctgctt taatgccttt gtatcatgct attgcttccc gtatggcttt
 4921 cattttctcc tcttgtata aatcctgggt gctgtctctt tatgaggagt tgtggcccg
 4981 tgtcaggcaa cgtggcgtgg tgtgactgt gtttgctgac gcaaccccc ctgggtggg
 5041 cattgccacc acctgtcagc tctttccgg gacttctgct tccccctcc ctattgccac
 5101 ggcggaactc atcgccgctt gccttgcccg ctgctggaca ggggctcggc tgttgggcac
 5161 tgacaattcc gtggtgttgt cggggaatc atcgctcttt ccttggctgc tcgcctgtgt
 5221 tgccacctgg attctgcgcg ggacgtcctt ctgctacgtc ccttcggccc tcaatccagc
 5281 ggaccttctt tcccgcgccc tgetgcgggc tctgcggcct cttccgcgtc ttcgccttcg
 5341 cctcagacg agtcggatct cctttggggc cgctccccg cctggtacct ttaagaccaa
 5401 tgacttacaa ggcagctgta gatcttagcc actttttaaa agaaaagggg ggactggaag
 5461 ggctaattca ctcccaacga aaataagatc tgetttttgc ttgtaactgg tctctctggt
 5521 tagaccagat ctgagcctgg gagctctctg gctaactagg gaacctactg cttagacctc
 5581 aataaagctt gccttgagtg cttcaagtag tgtgtgccg tctgttgtgt gactctggta
 5641 actagagatc cctcagaccc ttttagtcag tgtggaaaat ctctagcagt agtagttcat
 5701 gtcactctat tattcagtat ttataacttg caaagaaatg aatatcagag agtgagagga
 5761 acttgtttat tgcagcttat aatggttaca aataaagcaa tagcatcaca aatttcacaa
 5821 ataaagcatt tttttcactg cattctagtt gtggtttgtc caaactcatc aatgtatctt
 5881 atcatgtctg gctctagcta tcccgccctt aactccgccc agttccgccc attctccgcc
 5941 ccatggctga ctaatttttt ttatttatgc agaggccgag gccgcctcgg cctctgagct
 6001 attccagaag tagtgaggag gctttttttg aggcctagac ttttgagag acggcccaaa
 6061 ttcgtaatca tgggtcatagc tgtttcctgt gtgaaattgt tatccgctca caattccaca
 6121 caacatacga gccggaagca taaagtgtaa agcctggggg gcctaataag tgagctaaact
 6181 cacattaatt gcgttgcgct cactgcccgc tttccagtcg ggaaacctgt cgtgccagct
 6241 gcattaatga atcgccaac gcgcggggag aggcggtttg cgtattgggc gctcttcgc
 6301 ttcctcgctc actgactcgc tgcgctcggg cgctcggctg cggcgagcgg tatcagctca
 6361 ctcaaaggcg gtaatacggg tatccacaga atcaggggat aacgcaggaa agaacatgtg
 6421 agcaaaaggc cagcaaaagg ccaggaaccg taaaaaggcc gcgttgctgg cgtttttcca
 6481 taggctccgc cccctgacg agcatcacaa aaatcgacgc tcaagtcaga ggtggcgaaa
 6541 cccgacagga ctataaagat accaggcggt tccccctgga agctccctcg tgcgctctcc
 6601 tgttccgacc ctgcccctta ccggatacct gtccgccttt cctccctcgg gaagcgtggc
 6661 gctttctcat agctcacgct gtaggtatct cagttcgggt taggtcgctt gctccaagct
 6721 gggctgtgtg cacgaacccc ccgttcagcc cgaccgctgc gccttatccg gtaactatcg
 6781 tcttgagtcc aacccggtaa gacacgactt atcgccactg gcagcagcca ctggtaacag
 6841 gattagcaga gcgaggtatg taggcgggtg tacagagttc ttgaagtggg ggcctaacta
 6901 cggctacact agaaggacag tatttgggtat ctgcgctctg ctgaagccag ttaccttcgg
 6961 aaaaagagtt ggtagctctt gatccggcaa acaaaccacc gctggtagcg gtggtttttt
 7021 tgtttgcaag cagcagatta cgcgcagaaa aaaaggatct caagaagatc ctttgatctt
 7081 ttctacgggg tctgacgctc agtggaacga aaactcacgt taagggattt tgggtcatgag
 7141 attatcaaaa aggatcttca cctagatcct tttaaattaa aatgaagtt ttaaataaat
 7201 ctaaagtata tatgagtaaa cttgggtctga cagttaccaaa tgcttaataca gtgaggcacc
 7261 tatctcagcg atctgtctat ttcgttcate catagttgcc tgactccccg tcgtgtagat
 7321 aactacgata cgggagggct taccatctgg cccagtgct gcaatgatac cgcgagaccc
 7381 acgtcaccg gctccagatt tatcagcaat aaaccagcca gccggaaggg ccgagcgcag
 7441 aagtggctct gcaactttat ccgcctccat ccagtcctatt aattgttgcc gggaagctag
 7501 agtaagtagt tcgccagtta atagtttgcg caacgttggt gccattgcta caggcatcgt
 7561 ggtgtcacgc tcgtcgtttg gtatggcttc attcagctcc ggttcccaac gatcaaggcg
 7621 agttacatga tccccatgt tgtgcaaaaa agcggtttagc tcttcgggtc ctccgatcgt
 7681 tgtcagaagt aagttggccg cagtgttatc actcatgggt atggcagcac tgcataattc
 7741 tcttactgtc atgccatccg taagatgctt ttctgtgact ggtgagtaact caaccaagtc
 7801 attctgagaa tagtgtatgc ggcgaccgag ttgctcttgc ccggcgctcaa tacgggataa

```

7861 taccgcgcc catagcagaa ctttaaaagt gctcatcatt ggaaaacgtt ctccggggcg
7921 aaaactctca aggatctttac cgctgttgag atccagttcg atgtaaccca ctctgtgcacc
7981 caactgatct tcagcatctt ttacttttcac cagcgtttct ggggtgagcaa aaacaggaag
8041 gcaaaatgcc gcaaaaaagg gaataagggc gacacggaaa tgttgaatac tcatactctt
8101 cctttttcaa tattattgaa gcatttatca ggggttattgt ctcatgagcg gatacatatt
8161 tgaatgtatt tagaaaaata aacaaatagg gggtccgcgc acatttcccc gaaaagtgcc
8221 acctgacgtc taagaaacca ttattatcat gacattaacc tataaaaaata ggcgtatcac
8281 gagggccctt cgtctcgcgc gtttcgggtga tgacgggtgaa aacctctgac acatgcagct
8341 cccggagacg gtcacagctt gtctgtgaagc ggatgccggg agcagacaag cccgtcaggg
8401 cgcgtcagcg ggtgttgagg ggtgtcgggg ctggcttaac tatgcggcat cagagcagat
8461 tgtactgaga gtgcaccata tgcgggtgtga aataccgcac agatgcgtaa ggagaaaaata
8521 ccgcatcagg cgccattcgc cattcaggct gcgcaactgt tgggaagggc gatcgggtgcg
8581 ggccctcttc ctattacgcc agctggcgaa agggggatgt gctgcaaggc gattaagttg
8641 ggtaacgcc ggggtttccc agtcacgacg ttgtaaaacg acggccagtg ccaagctg.

```

p510_antiCD19_LL_TCRbeta (SEQ ID NO: 13)

```

1 acgcgtgtag tcttatgcaa tactcttgta gtcttgcaac atggtaacga tgagtttagca
61 acatgcctta caaggagaga aaaagcacccg tgcattgccga ttgggtggaag taagggtgga
121 cgatcgtgcc ttattaggaa ggcaacagac gggctctgaca tggattggac gaaccactga
181 attgccgcat tgcagagata ttgtatttaa gtgcctagct cgatacaata aacgggtctc
241 tctgggttaga ccagatctga gcctggggagc tctctggcta actagggaac ccactgctta
301 agcctcaata aagcttgccct tgagtgcctc aagtagtgtg tgcccgtctg ttgtgtgact
361 ctggtaacta gagatccctc agaccctttt agtcagtgtg gaaaatctct agcagtggcg
421 cccgaacagg gacctgaaag cgaaagggaa accagagctc tctcgacgca ggactcggct
481 tgctgaagcg cgcacggcaa gaggcgaggg gcggcgactg gtgagtacgc caaaaaatctt
541 gactagcggg ggctagaagg agagagatgg gtgcgagagc gtcagtatta agcgggggag
601 aattagatcg cgatgggaaa aaattcgggt aaggccaggg ggaaagaaaa aatataaatt
661 aaaacatata gtatgggcaa gcaggagact agaacgattc gcagttaatc ctggcctgtt
721 agaaacatca gaaggctgta gacaaatact gggacagcta caaccatccc ttcagacagg
781 atcagaagaa cttagatcat tatataatac agtagcaacc ctctattgtg tgcataaag
841 gatagagata aaagacacca aggaagcttt agacaagata gaggaagagc aaaacaaaag
901 taagaccacc gcacagcaag cggccactga tcttcagacc tggaggagga gatatgaggg
961 acaattggag aagtgaatta tataaatata aagtagtaaa aattgaacca ttaggagtag
1021 caccaccaa ggcaagaga agagtgggtg agagagaaaa aagagcagtg ggaataggag
1081 ctttgttctt tgggttcttg ggagcagcag gaagcactat gggcgcagcc tcaatgacgc
1141 tgacggtaca ggccagacaa ttattgtctg gtatagtgca gcagcagaac aatttgctga
1201 gggctattga ggcgcaacag catctgttgc aactcacagt ctggggcatc aagcagctcc
1261 aggcaagaat cctggctgtg gaaagatacc taaaggatca acagctcctg gggatttggg
1321 gttgctctgg aaaactcatt tgcaccactg ctgtgccttg gaatgctagt tggagtaata
1381 aatctctgga acagattgga atcacacgac ctggatggag tgggacagag aaattaacaa
1441 ttacacaagc ttaatacact ccttaattga agaatcgcaa aaccagcaag aaaagaatga
1501 acaagaatta ttggaattag ataaatgggc aagtttgttg aattgggtta acataacaaa
1561 ttggctgtgg tatataaaat tattcataat gatagtagga ggcttggtag gtttaagaat
1621 agtttttgct gtactttcta tagtgaatag agttaggcag ggatattcac cattatcggt
1681 tcagaccacac ctcccaaccc cgaggggacc cgacaggccc gaaggaatag aagaagaagg
1741 tggagagaga gacagagaca gatccattcg attagtgaac ggatctcgac ggtatcgggt
1801 aactttttaa agaaaagggg ggattggggg gtacagtgca ggggaaagaa tagtagacat
1861 aatagcaaca gacatacaaa cttaaagaatt acaaaaacaa attacaaaat tcaaaatctt
1921 atcgatacta gtattatgcc cagtacatga ccttatggga ctttcctact tggcagtaga
1981 tctacgtatt agtcatcgct attaccatgg tgatgcgggt ttggcagtag atcaatgggc
2041 gtggatagcg gtttgactca cggggatttc caagtctcca cccattgac gtcaatggga
2101 gtttgttttg gcaccaaatt caacgggact ttccaaaatg tcgtaacaac tccgccccat
2161 tgacgcaaatt gggcggtagg cgtgtacggt gggaggttta tataagcaga gctcgtttag
2221 tgaaccgtca gatcgcttg agacgccatc cagctgttt tgacctcat agaagattct
2281 agagccgcc ccatgcttct cctggtgaca agccttctgc tctgtgagtt accacacca
2341 gcattcctcc tgatcccaga catccagatg acacagacta catcctcct gtctgcctct
2401 ctgggagaca gagtcccat cagttgcagg gcaagtcagg acattagtaa atatttaaat
2461 tggatatcagc agaaaccaga tggaaactgt aaactcctga tctaccatac atcaagatta
2521 cactcaggag tcccatcaag gttcagtggc agtgggtctg gaacagatta ttctctcacc
2581 attagcaacc tggagcaaga agatattgcc acttactttt gccaacaggg taatacgtt

```

2641 ccgtacacgt tcggaggggg gactaagttg gaaataacag gctccacctc tggatccggc
 2701 aagccccgat ctggcgaggg atccaccaag ggcgaggtga aactgcagga gtcaggacct
 2761 ggccctggtg cgccctcaca gagcctgtcc gtcacatgca ctgtctcagg ggtctcatta
 2821 cccgactatg gtgtaagctg gattcgccag cctccacgaa agggctctgga gtggctggga
 2881 gtaatatggg gtagtgaaac cacatactat aattcagctc tcaaattccag actgaccatc
 2941 atcaaggaca actccaagag ccaagttttc ttaaaaatga acagtctgca aactgatgac
 3001 acagccatth actactgtgc caaacattat tactacgggtg gtagctatgc tatggactac
 3061 tgggggtcaag gaacctcagt caccgtctcc tcagcggccg caattgaagt tatgtatcct
 3121 cctccttacc taggtggcgg cggttctggt ggcgggcggtt ctgggtggcgg cggttctctc
 3181 gagctgggag caggcccagt ggattctgga gtcacacaaa ccccaaagca cctgatcaca
 3241 gcaactggac agcagtgac gctgagatgc tcccttaggt ctggagacct ctctgtgtca
 3301 tggtagcaac agagcctgga ccagggcctc cagttcctca ttcagtatta taatggagaa
 3361 gagagagcaa aaggaaacat tcttgaacga ttctccgcac aacagttccc tgacttgac
 3421 tctgaactaa acctgagctc tctggagctg ggggactcag ctttgtatth ctgtgccagc
 3481 agcccccgga caggcctgaa cactgaagct ttctttggac aaggcaccag actcacagtt
 3541 gtagaggacc tgaacaaggt gttcccaccc gaggtcgtctg tgtttgagcc atcagaagca
 3601 gagatctccc acacccaaaa ggccacactg gtgtgcctgg ccacaggett cttccccgac
 3661 cacgtggagc tgagctgggt ggtgaatggg aaggaggtgc acagtggggg cagcacggac
 3721 ccgcagcccc tcaaggagca gccgcctcct aatgactcca gatactgcct gacgagccgc
 3781 ctgaggggtc cgccacctt ctggcagaac ccccgcaacc acttcgctg tcaagtccag
 3841 ttctacgggc tctcgagaa tgacgagtgg acccaggata gggccaaacc cgtcaccag
 3901 atcgtcagcg ccgaggcctg gggtagagca gactgtgggt ttacctcggg gtccaccag
 3961 caaggggtcc tgtctgccac catcctctat gagatcctgc tagggaaggc caccctgtat
 4021 gctgtgctgg tcagcgccct tgtgttgatg gccatggtca agagaaagga tttctgataa
 4081 gaattcgatc cgcggccgcg aaggatctgc gatcgctccg gtgcccgtca gtgggcagag
 4141 cgcacatcgc ccacagtcct cgagaagttg gggggagggg tcggcaattg aacgggtgcc
 4201 tagagaaggt ggcgcggggt aaactgggaa agtgatgtcg tgtactggct ccgccttttt
 4261 cccgaggggt ggggagaacc gtatataagt gcagtagtcg ccgtgaacgt tctttttcgc
 4321 aacgggtttg ccgccagaac acagctgaag cttcgagggg ctcgcatctc tccctcacgc
 4381 gccgcgcgcc ctacctgagg ccgccatcca cgccggttga gtcgcgttct gccgcctccc
 4441 gcctgtgggt cctcctgaac tgcgtccgcc gtctaggtaa gtttaaagct cagggtcgaga
 4501 ccgggccttt gtccggcgct cccttgagac ctacctagac tcagccgggt ctcacagctt
 4561 tgcctgaccc tgcctgtca actctacgtc tttgtttcgt tttctgttct gcgcggttac
 4621 agatccaagc tgtgaccggc gcctacgcta gatgaccgag tacaagccca cgggtgcgct
 4681 cgccaccgcg gacgacgtcc ccagggcctg acgcaccctc gccgcgcgt tcgcgcacta
 4741 cccgcgcacg cgccacaccg tcgatccgga ccgccacatc gagcgggtca ccgagctgca
 4801 agaactcttc ctacgcgcg tcgggctcga catcggaag gtgtgggtcg cggacgacgg
 4861 cgccgcggtg gcggtctgga ccacgcgga gagcgtcgaa gcggggcgcg tgttcgcca
 4921 gatcggcccg cgcagtgccg agttgagcgg ttcccggctg gccgcgcagc aacagatgga
 4981 aggcctcctg gcgcgcgacc ggcccaagga gccgcgctg gtcctggcca ccgtcggcgt
 5041 ctgcgccgac caccagggca aggtctggtg cagcgcgctc gtgctccccg gactggaggc
 5101 ggccgagcgc gccgggggtg ccgccttctt ggagacctcc gcgcgccgca acctccctt
 5161 ctacgagcgg ctgcgcttca ccgtcacgcg cgacgtcgag gtgcccgaag gacgcgcac
 5221 ctggtgcatg acccgcaagc ccggtgcttg agtcgacaat caacctctgg attacaaaat
 5281 ttgtgaaaga ttgactggta ttcttaacta tgttgctcct ttacgctat gtggatacgc
 5341 tgctttaaatg cctttgtatc atgctattgc ttcccgtatg gctttcatth tctcctcctt
 5401 gtataaatcc tgggtgctgt ctctttatga ggagttgtgg cccgttgtca ggcaacgtgg
 5461 cgtgggtgtg actgtgtttg ctgacgcaac cccactggt tggggcattg ccaccacctg
 5521 tcagctcctt tcggggactt tcgctttccc cctccctatt gccacggcgg aactcatcgc
 5581 cgctgcctt gcccgctgct ggacaggggc tcggctgttg ggcactgaca attcgtggt
 5641 gttgtcgggg aaatcatcgt cctttccttg gctgctcgcc tgtgttgcca cctggattct
 5701 gcgcgggacg tcttctgtct acgtcccttc ggccctcaat ccagcggacc tctctccc
 5761 cggcctgctg ccggtctgct ggctcttccc gcgtcttcgc cttcgcctc agacgagtcg
 5821 gatctccctt tgggcgcctt cccgcctggg tacctttaag accaatgact tacaaggcag
 5881 ctgtagatct tagccactth ttaaaagaaa aggggggact ggaagggcta attcactccc
 5941 aacgaaaata agatctgctt tttgcttgta ctgggtctct ctggttagac cagatctgag
 6001 cctgggagct ctctggctaa ctaggaagcc cactgcttaa gcctcaataa agcttgccct
 6061 gactgcttca agtagtgtgt gccgctctgt tgtgtgactc tggtaactag agatccctca
 6121 gaccctttta gtcagtgtgg aaaatctcta gcagtagtag ttcagtcat cttattattc

```

6181 agtattttata acttgcaaag aaatgaatat cagagagtga gaggaacttg tttattgcag
6241 cttataatgg ttacaaataa agcaatagca tcacaaattt cacaaataaa gcattttttt
6301 cactgcattc tagttgtggt ttgtccaaac tcacaaatgt atcttatcat gtctggctct
6361 agctatccc cccctaactc cgcccagttc cgcccattct ccgcccctatg gctgactaat
6421 tttttttatt tatgcagagg ccgaggccgc ctccgctctc gagctattcc agaagtagtg
6481 aggaggcttt tttggaggcc tagacttttg cagagacggc ccaaattcgt aatcatggtc
6541 atagctgttt cctgtgtgaa attgttatcc gctcacaatt ccacacaaca tacgagccgg
6601 aagcataaag tgtaaagcct ggggtgccta atgagtgage taactcacat taattgcgtt
6661 gcgctcactg cccgctttcc agtcgggaaa cctgtcgtgc cagctgcatt aatgaatcgg
6721 ccaacgcgcg gggagaggcg gtttgcgat tgggcgctct tccgcttctc cgctcactga
6781 ctccgctgcg tcggtcgttc ggctgcggcg agcggtatca gctcactcaa aggcggtaat
6841 acggttatcc acagaatcag gggataacgc aggaagaac atgtgagcaa aaggccagca
6901 aaaggccagg aaccgtaaaa aggcgcgctt gctggcgttt tcccataggc tccgcccccc
6961 tgacgagcat cacaaaaatc gacgctcaag tcagaggtgg cgaaaccgca caggactata
7021 aagataccag gcgtttcccc ctggaagctc cctcgtgcgc tctcctgttc cgaccctgcc
7081 gcttaccgga tacctgtccg cctttctccc ttcgggaagc gtggcgcttt ctcatagctc
7141 acgctgtagg tatctcagtt cgggtgtagg gctgcgcctt atccggtaac tatcgtcttg agtccaacc
7201 acccccgctt cagcccgacc gctgcgcctt agccactggt aacaggatta gcagagcgag
7261 ggtaagacac gacttatcgc cactggcagc agccactggt aacaggatta gcagagcgag
7321 gtatgtaggc ggtgctacag agttcttgaa gtggtggcct aactacggct acactagaag
7381 gacagtattt ggtatctgcg ctctgctgaa gccagttacc ttcggaaaaa gagttggtag
7441 ctcttgatcc ggcaaaaaa ccaccgctgg tagcgggtgg tttttgttt gcaagcagca
7501 gattacgcgc agaaaaaaag gatctcaaga agatcctttg atcttttcta cggggtctga
7561 cgctcagtgg aacgaaaact cacgttaagg gattttggtc atgagattat caaaaaggat
7621 cttcacctag atccttttaa attaaaaatg aagtttttaa tcaatctaaa gtatatatga
7681 gtaaaacttg tctgacagtt accaatgctt aatcagtgag gcacctatct cagcgatctg
7741 tctatttcgt tcatccatag ttgcctgact ccccgctcgt tagataacta cgatacggga
7801 gggcttacca tctggcccca gtgctgcaat gataccgcga gaccacgct caccggctcc
7861 agatttatca gcaataaacc agccagccgg aaggcccgag cgcagaagtg gtctgcaac
7921 tttatccgcc tccatccagt ctattaattg ttgccgggaa gctagagtaa gtagttcgcc
7981 agttaatagt ttgcgcaacg ttgttgccat tgctacaggc atcgtggtgt cacgctcgtc
8041 gtttggtatg gcttcattca gctccggttc ccaacgatca aggcgagtta catgatcccc
8101 catgttgtgc aaaaaagcgg ttagctcctt cggctcctcc atcgttgtca gaagtaagtt
8161 ggccgcagtg ttatcactca tgggtatggc agcactgcat aattctctta ctgtcatgcc
8221 atccgtaaga tgcttttctg tgactggtga gtactcaacc aagtcattct gagaatagtg
8281 tatgcggcga ccgagttgct cttgcccggc gtcaatacgg gataataccg cgccacatag
8341 cagaacttta aaagtgtca tcattggaaa acgttcttcg gggcgaaaac tctcaaggat
8401 cttaccgctg ttgagatcca gttcgatgta accactcgt gcaccaact gatcttcagc
8461 atcttttact ttcaccagcg tttctgggtg agcaaaaaa ggaaggcaaa atgccgcaa
8521 aaagggaata agggcgacac ggaaatgttg aatactcata ctcttctttt tccaatatta
8581 ttgaagcatt tatcagggtt attgtctcat gagcggatag atatttgaat gtatttagaa
8641 aaataaacia ataggggttc cgcgcacatt tcccgaaaa gtgccacctg acgtctaaga
8701 aaccattatt atcatgacat taacctataa aaataggcgt atcacgaggc cctttcgtct
8761 cgcgcgtttc ggtgatgacg gtgaaaacct ctgacacatg cagctcccgg agacggctac
8821 agcttgctct taagcggatg ccgggagcag acaagcccgt cagggcgctg cagcgggtgt
8881 tggcgggtgt cggggctggc ttaactatgc ggcatcagag cagattgtac tgagagtga
8941 ccatatgcgg tgtgaaatac cgcacagatg cgtaaggaga aaataccgca tcaggcgcca
9001 ttcgccattc aggtcgcgca actgttgagg agggcgatcg gtgcgggctt cttcgtctat
9061 acgccagctg gcgaaagggg gatgtgctgc aaggcgatta agttgggtaa cgccagggtt
9121 ttcccagtcg cgacgttgta aaacgacggc cagtgccaaag ctg

```

p510_antiCD19_LL_TCRbetaC (SEQ ID NO: 14)

```

1 acgctgttag tcttatgcaa tactcttgta gtcttgcaac atggtaacga tgagttagca
61 acatgcctta caaggagaga aaaagcaccg tgcatgcga ttgggtggaag taagggtgga
121 cgatcgtgcc ttattaggaa ggcaacagac gggctctgaca tggattggac gaaccactga
181 attgccgcat tgcagagata ttgtatttaa gtgcctagct cgatacaata aacgggtctc
241 tctgggttaga ccagatctga gcctgggagc tctctggcta actagggaa ccactgctta
301 agcctcaata aagcttgccct tgagtgtctc aagtagtgtg tgcccgtctg ttgtgtgact
361 ctggtaacta gagatccctc agaccctttt agtcagtggt gaaaatctct agcagtggtg
421 cccgaacagg gacctgaaag cgaaagggaa accagagctc tctcgacgca ggactcggct

```

```

481  tgctgaagcg  cgcacggcaa  gaggcgaggg  gcggcgactg  gtgagtacgc  caaaaatttt
541  gactagcgga  ggctagaagg  agagagatgg  gtgcgagagc  gtcagtatta  agcgggggag
601  aattagatcg  cgatgggaaa  aaattcgggt  aaggccaggg  ggaaagaaaa  aatataaatt
661  aaaacatata  gtatgggcaa  gcagggagct  agaacgattc  gcagttaatc  ctggcctgtt
721  agaaacatca  gaaggctgta  gacaaatact  gggacagcta  caaccatccc  ttcagacagg
781  atcagaagaa  cttagatcat  tatataatac  agtagcaacc  ctctattgtg  tgcatacaaag
841  gatagagata  aaagacacca  aggaagcttt  agacaagata  gaggaagagc  aaaacaaaag
901  taagaccacc  gcacagcaag  cggccactga  tcttcagacc  tggaggagga  gatatgaggg
961  acaattggag  aagtgaatta  tataaatata  aagtagtaaa  aattgaacca  ttaggagtag
1021  caccaccaa  ggcaaagaga  agagtgggtg  agagagaaaa  aagagcagtg  ggaataggag
1081  ctttgttctt  tgggttcttg  ggagcagcag  gaagcactat  gggcgagcc  tcaatgacgc
1141  tgacggtaca  ggccagacaa  ttattgtctg  gtatagtgca  gcagcagaac  aatttgctga
1201  gggctattga  ggcgcaacag  catctgttgc  aactcacagt  ctggggcatc  aagcagctcc
1261  aggcaagaat  cctggctgtg  gaaagatacc  taaaggatca  acagctcctg  gggatttggg
1321  gttgctctgg  aaaactcatt  tgcaccactg  ctgtgccttg  gaatgctagt  tggagtaata
1381  aatctctgga  acagattgga  atcacacgac  ctggatggag  tgggacagag  aaattaacaa
1441  ttacacaagc  ttaatacact  ccttaattga  agaatcgcaa  aaccagcaag  aaaagaatga
1501  acaagaatta  ttggaattag  ataaatgggc  aagtttgttg  aattggttta  acataacaaa
1561  ttggctgtgg  tatataaaat  tattcataat  gatagtagga  ggcttggtag  gtttaagaat
1621  agtttttgc  gtactttcta  tagtgaatag  agttaggcag  ggatattcac  cattatcggt
1681  tcagaccac  ctcccaaccc  cgaggggacc  cgacaggccc  gaaggaatag  aagaagaagg
1741  tggagagaga  gacagagaca  gatccattcg  attagtgaac  ggatctcgac  ggtatcggtt
1801  aactttttaa  agaaaagggg  ggattggggg  gtacagtgca  ggggaaagaa  tagtagacat
1861  aatagcaaca  gacatacaaa  ctaaagaatt  acaaaaacaa  attacaaaat  tcaaaatttt
1921  atcgatacta  gtattatgcc  cagtacatga  ccttatggga  ctttcctact  tggcagtaca
1981  tctacgtatt  agtcacgcgt  attaccatgg  tgatgcgggt  ttggcagtac  atcaatgggc
2041  gtggatagcg  gtttgactca  cggggatttc  caagtctcca  cccattgac  gtcaatggga
2101  gtttgttttg  gcacccaaaat  caacgggact  ttccaaaatg  tcgtaacaac  tccgccccat
2161  tgacgcaaat  gggcggtagg  cgtgtacggt  gggaggttta  tataagcaga  gctcgtttag
2221  tgaaccgtca  gatcgcttgg  agacgccatc  cacgctgttt  tgacctccat  agaagattct
2281  agagccgcca  ccatgcttct  cctggtgaca  agccttctgc  tctgtgagtt  accacacca
2341  gcattcctcc  tgatcccaga  catccagatg  acacagacta  catcctccct  gtctgcctct
2401  ctgggagaca  gagtcaccat  cagttgcagg  gcaagtcagg  acattagtaa  atattttaat
2461  tggtatcagc  agaaaccaga  tggaaactgt  aaactcctga  tctaccatac  atcaagatta
2521  cactcaggag  tcccatcaag  gttcagtggc  agtgggtctg  gaacagatta  ttctctcacc
2581  attagcaacc  tggagcaaga  agatattgcc  acttactttt  gccaacaggg  taatacgctt
2641  ccgtacacgt  tcggaggggg  gactaagttg  gaaataacag  gctccacctc  tggatccggc
2701  aagcccggat  ctggcgaggg  atccaccaag  ggcgaggtga  aactgcagga  gtcaggacct
2761  ggcctggtgg  cgccctcaca  gagcctgtcc  gtcacatgca  ctgtctcagg  ggtctcatta
2821  cccgactatg  gtgtaagctg  gattcgccag  cctccacgaa  agggctctga  gtggctggga
2881  gtaatatggg  gtagtgaaac  cacatactat  aattcagctc  tcaaattccag  actgaccatc
2941  atcaaggaca  actccaagag  ccaagttttc  ttaaaaatga  acagtctgca  aactgatgac
3001  acagccattt  actactgtgc  caaacattat  tactacgggt  gtagctatgc  tatggactac
3061  tggggtcaag  gaacctcagt  caccgtctcc  tcagcggccg  caattgaagt  tatgtatcct
3121  cctccttacc  taggtggcgg  cggttctggt  ggcggcgggt  ctggtggcgg  cggttctctc
3181  gaggaggacc  tgaacaaggt  gttcccaccc  gaggtcgctg  tgtttgagcc  atcagaagca
3241  gagatctccc  acacccaaaa  ggccacactg  gtgtgcctgg  ccacaggctt  ctccccgac
3301  cagtgaggac  tgagctggtg  ggtgaatggg  aaggaggtgc  acagtggggt  cagcacagac
3361  ccgcagcccc  tcaaggagca  gccgcctc  aatgactcca  gatactgcct  gagcagccgc
3421  ctgaggggtc  cggccacctt  ctggcagaac  ccccgcaacc  acttcgctg  tcaagtccag
3481  ttctacgggc  tctcggagaa  tgacgagtgg  acccaggata  gggccaaaacc  cgtcaccag
3541  atcgtcagcg  ccgaggcctg  gggtagagca  gactgtgggt  ttacctcggt  gtectaccag
3601  caaggggtcc  tgtctgccac  catcctctat  gagatcctgc  tagggaaggc  caccctgtat
3661  gctgtgctgg  tcagcgccct  tgtgttgatg  gccatggtca  agagaaagga  tttctgataa
3721  gaattcgatc  cgcggccgcg  aaggatctgc  gategctccg  gtgcccgtca  gtgggcagag
3781  cgcacatcgc  ccacagtc  cgagaagttg  gggggagggg  tcggcaattg  aacgggtgcc
3841  tagagaaggt  ggcgcggggg  aaactgggaa  agtgatgtcg  tgtactgggt  ccgccttttt
3901  cccgaggggt  ggggagaacc  gtatataagt  gcagtagtcg  ccgtgaacgt  tctttttcgc
3961  aacgggtttg  ccgccagaac  acagctgaag  cttcgagggg  ctgcgatctc  tccttcacgc

```

4021 gccccgcgcgc ctacctgagg ccgccatcca cgccgggttga gtcgcggttct gccgcctccc
 4081 gcctgtggtg cctcctgaac tgcgtccgcc gtctaggttaa gtttaaagct caggctcgaga
 4141 ccgggccttt gtccggcgct cccttggagc ctacctagac tcagccggct ctccacgctt
 4201 tgcttgaccc tgettgtctca actctacgtc tttgtttcgt tttctgttct gcgcggttac
 4261 agatccaagc tgtgaccggc gcctacgcta gatgaccgag tacaagccca cgggtgcgcct
 4321 cgccacccgc gacgacgtcc ccagggccgt acgcaccctc gccgcgcgct tcgccgacta
 4381 cccccccacg cgccacaccg tcgatccgga ccgccacatc gagcgggtca ccgagctgca
 4441 agaactcttc ctacgcgcgc tcgggtctga catcggaag gtgtgggtcg cggacgacgg
 4501 cgccgcggtg gcggtctgga ccacgcggga gagcgtcgaa gcggggggcg tgttcgccga
 4561 gatcgggccg cgcatggccg agttgagcgg ttcccggtcg gccgcgcagc aacagatgga
 4621 aggcctcctg gcgcgcgacc ggcccaagga gccgcggtgg ttcttgcca ccgtcggcgt
 4681 ctgcgccgac caccagggca agggctctggg cagcgcgcgc gtgctccccg gaggggaggg
 4741 ggccgagcgc gccggggtgc ccgccttctt ggagacctcc gcgccccgca acctccctt
 4801 ctacgagcgg ctcggttca ccgtcaccgc cgacgtcgag gtgcccgaag gaccgcgcac
 4861 ctggtgcatg acccgcaagc ccggtgcctg agtcgacaat caacctctgg attacaaaat
 4921 ttgtgaaaga ttgactggta ttcttaacta tgttgctcct tttacgctat gtggatacgc
 4981 tgctttaatg cctttgtatc atgctattgc ttcccgatg gctttcattt tctctcctt
 5041 gtataaatcc tgggtgctgt ctctttatga ggagttgtgg ccggttgta ggcaacgtgg
 5101 cgtggtgtgc actgtgtttg ctgacgcaac cccactggt tggggcattg ccaccacctg
 5161 tcagctcctt tcgggactt tcgctttccc cctccctatt gccacggcgg aactcatcgc
 5221 cgctgcctt gccgcgtgct ggacaggggc tcggctgttg ggcaactgaca attcgcgtgt
 5281 gttgtcgggg aaatcatcgt cctttccttg gctgctcgcc tgtgttgcca cctggattct
 5341 gcgcgggacg tccttctgct acgtcccttc ggccctcaat ccagcggacc ttcttcccg
 5401 cggcctgctg ccggctctgc ggctcttccc gctcttcgc cttcgcctc agacgagtcg
 5461 gatctccctt tgggcccgcct cccgcctgg tacctttaag accaatgact tacaaggcag
 5521 ctgtagatct tagccacttt ttaaaagaaa aggggggact ggaagggcta attcactccc
 5581 aacgaaaata agatctgctt tttgcttgta ctgggtctct ctggttagac cagatctgag
 5641 cctgggagct ctctggctaa ctagggaacc cactgcttaa gcctcaataa agcttgctt
 5701 gagtgttca agtagtgtgt gccgctctgt tgtgtgactc tggtaactag agatccctca
 5761 gaccctttta gtcagtgtgg aaaatctcta gcagtagtag tcatgtcat cttattattc
 5821 agtatattata acttgcaaag aaatgaatat cagagagtga gaggaacttg tttattgcag
 5881 cttataatgg ttacaaataa agcaatagca tcacaaattt cacaaataaa gcattttttt
 5941 cactgcattc tagttgtggg ttgtccaaac tcatcaatgt atcttatcat gtctggctct
 6001 agctatcccg cccctaactc cgcccagttc cgccattct ccgccccatg gctgactaat
 6061 tttttttatt tatgcagagg ccgaggccgc ctcgccctct gagctattcc agaagtagtg
 6121 aggaggcttt tttggaggcc tagacttttg cagagacggc ccaaatctgt aatcatggtc
 6181 atagctgttt cctgtgtgaa attgttatcc gtcacaatt ccacacaaca tacgagccgg
 6241 aagcataaag tgtaaagcct ggggtgccta atgagtgagc taactcacat taattgcgtt
 6301 gcgctcactg ccgcctttcc agtcgggaaa cctgtcgtgc cagctgcatt aatgaatcgg
 6361 ccaacgcgcg gggagaggcg gtttgcgat tgggcgctct tccgcttctt cgctcactga
 6421 ctgcgtgcgc tcggtcgttc ggctgcggcg agcgggtatc gctcactcaa aggcggtaat
 6481 acggttatcc acagaatcag gggataacgc aggaagaac atgtgagcaa aaggccagca
 6541 aaaggccagg aaccgtaaaa aggcgcggtt gctggcggtt ttccataggc tccgcccccc
 6601 tgacgagcat cacaaaaatc gacgctcaag tcagaggtgg cgaaaccoga caggactata
 6661 aagataccag gcgtttcccc ctggaagctc cctcgtgcgc tctcctgttc cgaccctgcc
 6721 gcttaccgga tacctgtccg cctttctccc ttcgggaagc gtggcgcttt ctcatagctc
 6781 acgctgtagg tatctcagtt cgggttaggt cgttcgctcc aagctgggct gtgtgcacga
 6841 acccccgtt cagcccagcc gctgcgcctt atccggtaac tatcgtcttg agtccaacc
 6901 ggtaagacac gacttatcgc cactggcagc agccactggt aacaggatta gcagagcgag
 6961 gtatgtaggc ggtgctacag agttcttgaa gtggtggcct aactacggct acactagaag
 7021 gacagtattt ggtatctgcg ctctgtgaa gccagttacc ttcggaaaaa gagtgtgtag
 7081 ctcttgatcc ggcaaaacaa ccaccgctgg tagcgggtgg tttttgttt gcaagcagca
 7141 gattacgcgc agaaaaaaag gatctcaaga agatcctttg atcttttcta cggggtctga
 7201 cgctcagtggt aacgaaaact cacgttaagg gattttggct atgagattat caaaaaggat
 7261 cttcacctag atccttttaa attaaaaatg aagttttaaa tcaatctaaa gtatatatga
 7321 gtaaaccttg tctgacagtt accaatgctt aatcagtgag gcacctatct cagcgatctg
 7381 tctatttctg tcatccatag ttgcctgact ccccgctcgt tagataacta cgatacggga
 7441 gggcttacca tctggcccca gtgctgcaat gataccgoga gaccacgct caccggtcc
 7501 agatttatca gcaataaacc agccagccgg aaggggcgag cgcagaagtg gtctgcaac

```

7561 tttatccgcc tccatccagt ctattaattg ttgccgggaa gctagagtaa gtagttcgcc
7621 agttaatatg ttgcgcaacg ttgttgccat tgctacaggc atcgtggtgt cacgctcgtc
7681 gtttggtatg gcttcattca gctccggttc ccaacgatca aggcgagtta catgatcccc
7741 catgttggtg aaaaaagcgg ttagctcctt cggctctccg atcgttggtc gaagtaagtt
7801 ggccgcagtg ttatcactca tgggtatggc agcactgcat aattctctta ctgtcatgcc
7861 atccgtaaga tgcttttctg tgactgggtga gtactcaacc aagtcattct gagaatagtg
7921 tatgcggcga ccgagttgct cttgcccggc gtcaatacgg gataataccg cgccacatag
7981 cagaacttta aaagtgtctc tcattggaaa acgttcttcg gggcgaaaac tctcaaggat
8041 cttaccgctg ttgagatcca gttcgatgta acccactcgt gcacccaact gatcttcagc
8101 atcttttact ttcaccagcg tttctgggtg agcaaaaaca ggaaggcaaa atgccgcaaa
8161 aaagggaata agggcgacac ggaaatgttg aatactcata ctcttctttt ttcaatatta
8221 ttgaagcatt tatcagggtt attgtctcat gagcggatag atatttgaat gtatttagaa
8281 aaataaacia ataggggttc cgcgcacatt tccccgaaaa gtgccacctg acgtctaaga
8341 aaccattatt atcatgacat taacctataa aaataggcgt atcacgaggc ctttctgtct
8401 cgcgcgtttc ggtgatgacg gtgaaaacct ctgacacatg cagctcccgg agacgggtcac
8461 agcttggtct taagcggatg ccgggagcag acaagcccggt cagggcgctg cagcgggtgt
8521 tggcgggtgt cggggctggc ttaactatgc ggcatcagag cagattgtac tgagagtgcg
8581 ccataatgcg tgtgaaatac cgcacagatg cgtaaggaga aaataccgca tcaggcgcca
8641 ttcgccattc aggtctgcga actgttggga agggcgatcg gtgcgggcct ctctgctatt
8701 acgccagctg gcgaaagggg gatgtgctgc aaggcgatta agttgggtaa cgccagggtt
8761 ttcccagtcg cgacgttgta aaacgacggc cagtgccaaag ctg.

```

p510_antiCD19_LL_CD3gamma (SEQ ID NO: 15)

```

1 acgcgtgtag tcttatgcaa tactcttgta gtcttgcaac atggtaacga tgagttagca
61 acatgcctta caaggagaga aaaagcaccg tgcatgccga ttggtggaag taaggtggtg
121 cgatcgtgcc ttattaggaa ggcaacagac gggctctgaca tggattggac gaaccactga
181 attgccgcat tgcagagata ttgtatttaa gtgcctagct cgatacaata aacgggtctc
241 tctggttaga ccagatctga gcctgggagc tctctggcta actagggaac ccactgctta
301 agcctcaata aagcttgctt tgagtgtctc aagtagtgtg tgcccgtctg ttgtgtgact
361 ctggtaaacta gagatccctc agaccctttt agtcagtgtg gaaaatctct agcagtggcg
421 cccgaacagg gacctgaaag cgaaagggaa accagagctc tctcgacgca ggactcggct
481 tgctgaagcg cgcacggcaa gaggcgaggg gcggcgactg gtgagtacgc caaaaatttt
541 gactagcggg ggcctagaagg agagagatgg gtgcgagagc gtcagtatta agcgggggag
601 aattagatcg cgatgggaaa aaattcgggt aaggccaggg ggaaagaaaa aatataaatt
661 aaaacatata gtatgggcaa gcaggagact agaacgattc gcagttaatc ctggcctgtt
721 agaaacatca gaaggctgta gacaaatact gggacagcta caaccatccc ttcagacagg
781 atcagaagaa cttagatcat tatataatac agtagcaacc ctctattgtg tgcatcaaag
841 gatagagata aaagacacca aggaagcttt agacaagata gaggaagagc aaaacaaaag
901 taagaccacc gcacagcaag cggccactga tcttcagacc tggaggagga gatattgagg
961 acaattggag aagtgaatta tataaatata aagtagtaaa aattgaacca ttaggagtag
1021 caccaccaa ggcaaaagaga agagtgggtg agagagaaaa aagagcagtg ggaataggag
1081 ctttggttct tgggttcttg ggagcagcag gaagcactat gggcgagcc tcaatgacgc
1141 tgacgggtaca ggccagacaa ttattgtctg gtatagtgcg gcagcagaac aatttgctga
1201 gggctattga ggcgcaacag catctgttgc aactcacagt ctggggcatc aagcagctcc
1261 aggcaagaat cctggctgtg gaaagatacc taaaggatca acagctcctg gggatttggg
1321 gttgctctgg aaaactcatt tgcaccactg ctgtgccttg gaatgctagt tggagtaata
1381 aatctctgga acagattgga atcacacgac ctggatggag tgggacagag aaattaacaa
1441 ttacacaagc ttaatacact ccttaattga agaatcgcaa aaccagcaag aaaagaatga
1501 acaagaatta ttggaattag ataaatgggc aagtttgtgg aattgggtta acataacaaa
1561 ttggctgtgg tatataaaat tattcataat gatagtagga ggcttggtag gtttaagaat
1621 agtttttctg gtactttcta tagtgaatag agttaggcag ggatattcac cattatcggt
1681 tcagaccacac ctcccacccc cgaggggacc cgacaggccc gaaggaatag aagaagaagg
1741 tggagagaga gacagagaca gatccattcg attagtgaac ggatctcgac ggtatcggtt
1801 aactttttaa agaaaagggg ggattggggg gtacagtgcg ggggaaagaa tagtagacat
1861 aatagcaaca gacatacaaa cttaaagaatt acaaaaacaa attacaaaat tcaaaatttt
1921 atcgatacta gtattatgcc cagtacatga ccttatggga ctttctact tggcagtaca
1981 tctacgtatt agtcatcgct attaccatgg tgatgcgggt ttggcagtac atcaatgggc
2041 gtggatagcg gtttgactca cggggatttc caagtctcca cccattgac gtcaatggga
2101 gtttggtttg gcaccaaatt caacgggact ttccaaaatg tcgtaacaac tccgccccat
2161 tgacgcaaat gggcggtagg cgtgtacggt gggaggttta tataagcaga gctcgtttag

```

2221 tgaaccgtca gatcgcttgg agacgccatc cacgctgttt tgacctccat agaagattct
 2281 agagccgcca ccatgcttct cctggtgaca agccttctgc tctgtgagtt accacaccca
 2341 gcattcctcc tgatcccaga catccagatg acacagacta catcctccct gtctgcctct
 2401 ctgggagaca gagtcacat cagttgcagg gcaagtcagg acattagtaa atattttaat
 2461 tgggtatcagc agaaaccaga tggaaactgt aaactcctga tctaccatac atcaagatta
 2521 cactcaggag tcccatcaag gttcagtggt agtgggtctg gaacagatta ttctctcacc
 2581 attagcaacc tggagcaaga agatattgcc acttactttt gccaacaggg taatacgtct
 2641 ccgtacacgt tcggaggggg gactaagttg gaaataacag gctccacctc tggatccggc
 2701 aagcccggat ctggcgaggg atccaccaag ggcgaggtga aactgcagga gtcaggacct
 2761 ggcctgggtg cgccctcaca gagcctgtcc gtcacatgca ctgtctcagg ggtctcatta
 2821 cccgactatg gtgtaagctg gattcgccag cctccacgaa agggctctgga gtggctggga
 2881 gtaatatggg gtagtgaaac cacatactat aattcagctc tcaaattccag actgaccatc
 2941 atcaaggaca actccaagag ccaagttttc ttaaaaatga acagtctgca aactgatgac
 3001 acagccattt actactgtgc caaacattat tactacggtg gtagctatgc tatggactac
 3061 tgggggtcaag gaacctcagt caccgtctcc tcagcggccg caattgaagt tatgtatcct
 3121 cctccttacc taggtggcgg cggttctggt ggcgggcggt ctgggtggcg cggttctctc
 3181 gagcagtc aa tcaaaggaaa ccacttggtt aaggtgtatg actatcaaga agatgggttcg
 3241 gtacttctga cttgtgatgc agaagccaaa aatatcacat ggtttaaaga tgggaagatg
 3301 atcggtctcc taactgaaga taaaaaaaaa tggaaatctg gaagtaatgc caaggaccca
 3361 cgagggatgt atcagtgtaa aggatcacag aacaagtcaa aaccactcca agtgattac
 3421 agaattgtgc agaactgcat tgaactaaat gcagccacca tatctggctt tctctttgct
 3481 gaaatcgtca gcattttctg ccttgtctgt ggggtctact tcattgtctg acaggatgga
 3541 gttegccagt cgagagcttc agacaagcag actctgttgc ccaatgacca gctctaccag
 3601 cccctcaagg atcgagaaga tgaccagtac agccacctc aaggaaacca gttgaggagg
 3661 aattgataag aattcgatcc gcggccgcga aggatctgcg atcgctccgg tgcctcgag
 3721 tgggcagagc gcacatcgcc cacagtcccc gagaagttgg ggggaggggt cggcaattga
 3781 acgggtgcct agagaagggt gcgcggggta aactgggaaa gtgatgtcgt gtactggctc
 3841 cgcctttttc ccgaggggtg gggagaaccg tatataagtg cagtagtcgc cgtgaacgtt
 3901 ctttttctga acgggtttgc cgccagaaca cagctgaagc ttcgaggggc tgcctctct
 3961 ccttcaacgc cccgcgcgcc tacttgaggc cgccatccac gccggttgag tgcggttctg
 4021 ccgcctcccg cctgtggtgc ctctgaact gcgtccgcgg tctaggtaag tttaaagctc
 4081 aggtcgagac cgggcctttg tccggcgctc cctggagcc tacctagact cagccggctc
 4141 tccacgcttt gcctgacctt gcttgcctca ctctacgtct ttgtttcgtt ttctgttctg
 4201 ccgcgttaca gatccaagct gtgaccggcg cctacgctag atgaccgagt acaagccac
 4261 ggtgcgcctc gccaccgcgc acgacgtccc cagggccgta cgcacctcg ccgcgcgtt
 4321 ccgcgactac ccgcgcacgc gccacaccgt cgatccggac cgccacatcg agcgggtcac
 4381 cgagctgcaa gaactcttcc tcacgcgcgt cgggtctgac atcggaagg tgtgggtcgc
 4441 ggacgacggc gccgcggtgg cggctctggc cagcccgag agcgtcgaag cgggggcggt
 4501 gttegcgcag atcggcccg ccatggccga gttgagcgg tcccggctgg ccgcgcagca
 4561 acagatggaa ggcctcctgg cgcgcaccc gccaaggag cccgcgtggt tccctggcac
 4621 cgtcggcgct tcgcccagc accagggcaa gggctctggg agcgcgctg tgcctcccgg
 4681 agtggaggcg gccgagcgcg ccgggggtgc cgccttctg gagacctcc cgccccgcaa
 4741 cctccccctc tacgagcggc tgggttcac cgtcacccgc cgtcagagg tgcccgaagg
 4801 accgcgcacc tgggtgcatga ccgcgaagc cgggtgctga gtcgacaatc aacctctgga
 4861 ttacaaaatt tgtgaaagat tgactgggtat tcttaactat gttgctcctt ttacgctatg
 4921 tggatacgct gctttaatgc ctttgtatca tgcatttgc tcccgtatgg ctttcatttt
 4981 ctctccttg tataaatcct ggttgcgtgc tctttatgag gattgtggc ccgttgtcag
 5041 gcaacgtggc gtggtgtgca ctgtgtttgc tgacgcaacc cccactgggt ggggcattgc
 5101 caccacctgt cagctccttt ccgggacttt cgctttcccc ctccctattg ccacggcgga
 5161 actcatcgcc gcctgccttg ccgcgtgctg gacaggggt cggctgttgg gcactgacaa
 5221 ttccgtggtg ttgtcgggga aatcatcgct ctttctgcta cgtcccttcg gccctcaatc cagcggacct
 5281 ctggattctg ccggggacgt ccttctgcta cgtcccttcg gccctcaatc cagcggacct
 5341 tcttccccgc ggctgctgc cggctctgcg gcctcttcgg cgtcttcgcc ttccgctca
 5401 gacgagtcgg atctcccttt gggccgcctc ccgcctgggt acctttaaga ccaatgactt
 5461 acaaggcagc tgtagatctt agccactttt taaaagaaaa ggggggactg gaagggttaa
 5521 ttactccca acgaaaataa gatctgcttt ttgctgtac tgggtctctc tgggttagacc
 5581 agatctgagc ctgggagctc tctggctaac tagggaacct actgcttaag cctcaataaa
 5641 gcttgccctg agtgcttcaa gtagtgtgtg ccgctctgtt gtgtgactct ggtaactaga
 5701 gatecctcag accttttag tcagtgtgga aaatctctag cagtagtagt tcatgtcatc

5761 ttattatttca gtattttataa cttgcaaaga aatgaatatc agagagtgag aggaacttgt
 5821 ttattgcagc ttataatgggt tacaaataaa gcaatagcat cacaaatttc acaaataaag
 5881 cttttttttc actgcattct agttgtgggt tgtccaaact catcaatgta tcttatcatg
 5941 tctggctcta gctatcccgc ccctaactcc gccagttcc gccattctc cgccccatgg
 6001 ctgactaatt ttttttattt atgcagaggg cgaggccgcc tcggcctctg agctattcca
 6061 gaagtagtga ggaggctttt ttggaggcct agacttttgc agagacggcc caaattcgta
 6121 atcatgggtca tagctgtttc ctgtgtgaaa ttgttatccg ctcacaattc cacacaacat
 6181 acgagccgga agcataaagt gtaaagcctg ggggtgcctaa tgagttagct aactcacatt
 6241 aattgcgttg cgtcactgac ccgctttcca gtcgggaaac ctgtcgtgcc agctgcatta
 6301 atgaatcggc caacgcgcgg ggagaggcgg tttgcgtatt gggcgctctt ccgcttcctc
 6361 gctcactgac tcgctgcgct cggctgttcg gctcggcgga gcggtatcag ctcactcaaa
 6421 ggcggtaata cggttatcca cagaatcagg ggataacgca ggaaagaaca tgtgagcaaa
 6481 agggcagcaa aaggccagga accgtaaaaa ggccgcgttg ctggcgttt tccataggct
 6541 ccgccccctc gacgagcatc acaaaaatcg acgtcaagt cagaggtggc gaaacccgac
 6601 aggactataa agataaccagg cgtttccccc tggaagctcc ctctgcgct ctctgttcc
 6661 gacctgccc cttaccggat acctgtccgc ctttctccct tcgggaagcg tggcgcttcc
 6721 tcatagctca cgtgttaggt atctcagttc ggtgttaggtc gttcgtcca agctgggctg
 6781 tgtgcacgaa cccccgttc agcccgaccg ctgcgcctta tccgtaact atcgtcttga
 6841 gtccaacccg gtaagacacg acttatcgcc actggcagca gccactggta acaggattag
 6901 cagagcgagg tatgtaggcg gtgctacaga gttcttgaag tgggtggccta actacggcta
 6961 cactagaagg acagtatttg gtatctgcgc tctgctgaag ccagttacct tcggaaaaag
 7021 agttggtagc tcttgatccg gcaaacaac caccgctggg agcgggtgggt tttttgtttg
 7081 caagcagcag attacgcgca gaaaaaaagg atctcaagaa gatcctttga tcttttctac
 7141 ggggtctgac gctcagtggg acgaaaactc acgttaagggt attttgggtca tgagattatc
 7201 aaaaaggatc ttacacctaga tctttttaa ttaaaaatga agttttaaat caatctaaag
 7261 tatatatgag taaacttgggt ctgacagtta ccaatgctta atcagttagg cacctatctc
 7321 agcgatctgt ctatttcgtt catccatagt tgctgactc ccgctcgtgt agataactac
 7381 gatacgggag ggcttaccat ctggccccag tgctgcaatg ataccgcgag acccacgctc
 7441 accggctcca gatttatcag caataaacca gccagccgga agggccgagc gcagaagtgg
 7501 tcttgcaact ttatccgcct ccattccagtc tattaattgt tgccgggaag ctagagtaag
 7561 tagttcgcca gttaatagtt tgcgcaacgt tgttgccatt gctacaggca tcgtgggtgc
 7621 acgctcgtcg tttgggtatgg cttcattcag ctccgggtcc caacgatcaa ggcgagttac
 7681 atgatcccc atgttgtgca aaaaagcgggt tagctccttc ggtcctccga tcgttgtcag
 7741 aagtaagtgt gccgcagtggt tatcactcat ggttatggca gcaactgcata attctcttac
 7801 tgtcatgcca tccgtaagat gcttttctgt gactgggtgag tactcaacca agtcattctg
 7861 agaatagtgt atgcggcgac cgagttgctc ttgcccggcg tcaatacggg ataataccgc
 7921 gccacatagc agaactttaa aagtgtcat cattggaaaa cgttcttcgg ggcgaaaact
 7981 ctcaaggatc ttaccgctgt tgagatccag ttctggtgaa cccactcgtg caccactg
 8041 atcttcagca tcttttactt tcaccagcgt ttctgggtga gcaaaaacag gaaggcaaaa
 8101 tgccgcaaaa aagggaataa gggcgacacg gaaatgttga atactcatac tcttctttt
 8161 tcaatattat tgaagcattt atcagggtta ttgtctcatg agcggatata tatttgaatg
 8221 tatttagaaa aataaacaaa taggggttcc gcgcacattt ccccgaaaag tgccacctga
 8281 cgtctaagaa accattatta tcatgacatt aacctataaa aataggcgta tcacgaggcc
 8341 ctttcgtctc gcgcgtttcg gtgatgacgg tgaaaacctc tgacacatgc agctcccgga
 8401 gacggtcaca gcttgtctgt aagcggatgc cgggagcaga caagcccgtc agggcgcgctc
 8461 agcgggtggt ggcgggtgct ggggctggct taactatgcg gcatcagagc agattgtact
 8521 gagagtgcac catatgcggt gtgaaatacc gcacagatgc gtaaggagaa aataccgc
 8581 caggcgccat tcgccattca ggctgcgcaa ctgttgggaa gggcgatcgg tcggggcctc
 8641 ttcgctatta cgccagctgg cgaaaggggg atgtgctgca aggcgattaa gttgggtaac
 8701 gccagggttt tcccagtcac gacgttgtaa aacgacggcc agtgccaagc tg.

p510_antiCD19_LL_CD3delta (SEQ ID NO: 16)

1 acgcgtgtag tcttatgcaa tactcttgta gtcttgcaac atggtaacga tgagtttagca
 61 acatgcctta caaggagaga aaaagcaccg tgcattgccga ttgggtggaag taagggtggtg
 121 cgatcgtgcc ttattaggaa ggcaacagac ggggtctgaca tggattggac gaaccactga
 181 attgccgcat tgcagagata ttgtatttaa gtgcctagct cgatacaata aacgggtctc
 241 tctgggttaga ccagatctga gcctgggagc tctctggcta actagggaac ccaactgctta
 301 agcctcaata aagcttgccct tgagtgtctc aagtagtgtg tgcccgtctg ttgtgtgact
 361 ctggtaacta gagatccctc agaccctttt agtcagtgtg gaaaatctct agcagtggcg
 421 cccgaacagg gacctgaaag cgaaagggaa accagagctc tctcgacgca ggactcggct

```

481  tgctgaagcg  cgcacggcaa  gaggcgaggg  gcggcgactg  gtgagtacgc  caaaaatttt
541  gactagcgga  ggctagaagg  agagagatgg  gtgcgagagc  gtcagtatta  agcgggggag
601  aattagatcg  cgatgggaaa  aaattcgggt  aaggccaggg  ggaaagaaaa  aatataaatt
661  aaaacatata  gtatgggcaa  gcagggagct  agaacgattc  gcagttaatc  ctggcctgtt
721  agaaacatca  gaaggctgta  gacaaatact  gggacagcta  caaccatccc  ttcagacagg
781  atcagaagaa  cttagatcat  tatataatac  agtagcaacc  ctctattgtg  tgcatacaaag
841  gatagagata  aaagacacca  aggaagcttt  agacaagata  gaggaagagc  aaaacaaaag
901  taagaccacc  gcacagcaag  cggccactga  tcttcagacc  tggaggagga  gatatgaggg
961  acaattggag  aagtgaatta  tataaatata  aagtagtaaa  aattgaacca  ttaggagtag
1021 caccacacaa  ggcaaagaga  agagtgggtg  agagagaaaa  aagagcagtg  ggaataggag
1081 ctttgttcct  tgggttcttg  ggagcagcag  gaagcactat  gggcgagcc  tcaatgacgc
1141 tgacggtaca  ggccagacaa  ttattgtctg  gtatagtgca  gcagcagaac  aatttgctga
1201 gggctattga  ggcgcaacag  catctgttgc  aactcacagt  ctggggcatc  aagcagctcc
1261 aggcaagaat  cctggctgtg  gaaagatacc  taaaggatca  acagctcctg  gggatttggg
1321 gttgctctgg  aaaactcatt  tgcaccactg  ctgtgccttg  gaatgctagt  tggagtaata
1381 aatctctgga  acagattgga  atcacacgac  ctggatggag  tgggacagag  aaattaacaa
1441 ttacacaagc  ttaatacact  ccttaattga  agaatcgcaa  aaccagcaag  aaaagaatga
1501 acaagaatta  ttggaattag  ataaatgggc  aagtttgtgg  aattggttta  acataacaaa
1561 ttggctgtgg  tatataaaat  tattcataat  gatagtagga  ggcttggtag  gtttaagaat
1621 agtttttgc  gtactttcta  tagtgaatag  agttaggcag  ggatattcac  cattatcggt
1681 tcagaccac  ctcccaaccc  cgaggggacc  cgacaggccc  gaaggaatag  aagaagaagg
1741 tggagagaga  gacagagaca  gatccattcg  attagtgaac  ggatctcgac  ggtatcggtt
1801 aacttttaaa  agaaaagggg  ggattggggg  gtacagtgca  ggggaaagaa  tagtagacat
1861 aatagcaaca  gacatacaaa  ctaaagaatt  acaaaaacaa  attacaaaat  tcaaaatttt
1921 atcgatacta  gtattatgcc  cagtacatga  ccttatggga  ctttcctact  tggcagtaca
1981 tctacgtatt  agtcacgcgt  attaccatgg  tgatgcgggt  ttggcagtac  atcaatgggc
2041 gtggatagcg  gtttgactca  cggggatttc  caagtctcca  cccattgac  gtcaatggga
2101 gtttgttttg  gcacccaaaat  caacgggact  ttccaaaatg  tcgtaacaac  tccgccccat
2161 tgacgcaaat  gggcggtagg  cgtgtacggt  gggaggttta  tataagcaga  gctcgtttag
2221 tgaaccgtca  gatcgcttgg  agacgccatc  cacgtgtttt  tgacctccat  agaagattct
2281 agagccgcc  ccatgcttct  cctggtgaca  agccttctgc  tctgtgagtt  accacacca
2341 gcattcctcc  tgatcccaga  catccagatg  acacagacta  catcctccct  gtctgcctct
2401 ctgggagaca  gagtcaccat  cagttgcagg  gcaagtcagg  acattagtaa  atattttaat
2461 tggtatcagc  agaaaccaga  tggaaactgt  aaactcctga  tctaccatac  atcaagatta
2521 cactcaggag  tcccatcaag  gttcagtggc  agtgggtctg  gaacagatta  ttctctcacc
2581 attagcaacc  tggagcaaga  agatattgcc  acttactttt  gccaacaggg  taatacgctt
2641 ccgtacacgt  tcggaggggg  gactaagttg  gaaataacag  gctccacctc  tggatccggc
2701 aagcccggat  ctggcgaggg  atccaccaag  ggcgaggtga  aactgcagga  gtcaggacct
2761 ggcctggtgg  cgccctcaca  gagcctgtcc  gtcacatgca  ctgtctcagg  ggtctcatta
2821 cccgactatg  gtgtaagctg  gattcgccag  cctccacgaa  agggctctga  gtggctggga
2881 gtaatatggg  gtagtgaaac  cacatactat  aattcagctc  tcaaattccag  actgaccatc
2941 atcaaggaca  actccaagag  ccaagttttc  ttaaaaatga  acagtctgca  aactgatgac
3001 acagccattt  actactgtgc  caaacattat  tactacgggt  gtagctatgc  tatggactac
3061 tggggtcaag  gaacctcagt  caccgtctcc  tcagcggccg  caattgaagt  tatgtatcct
3121 cctccttacc  taggtggcgg  cggttctggt  ggcggcgggt  ctggtggcgg  cggttctctc
3181 gagttcaaga  tacctataga  ggaacttgag  gacagagtgt  ttgtgaattg  caataccagc
3241 atcacatggg  tagagggaac  ggtgggaaca  ctgctctcag  acattacaag  actggacctg
3301 ggaaaacgca  tcctggaccc  acgaggaata  tataggtgta  atgggacaga  tatatacaag
3361 gacaaagaat  ctaccgtgca  agttcattat  cgaatgtgcc  agagctgtgt  ggagctggat
3421 ccagccaccg  tggttgcat  cattgtcact  gatgtcattg  ccactctgct  ccttgctttg
3481 ggagtcttct  gctttgctgg  acatgagact  ggaaggctgt  ctggggctgc  cgacacacaa
3541 gctctgttga  ggaatgacca  ggtctatcag  cccctccgag  atcgagatga  tgctcagtac
3601 agccaccttg  gaggaaactg  ggctcggaac  aagtgataag  aattcgatcc  gcggccgcga
3661 aggatctgcg  atcgctccgg  tgcccgctcag  tgggcagagc  gcacatcgcc  cacagtcacc
3721 gagaagttgg  ggggaggggt  cggcaattga  acgggtgcct  agagaagggt  gcgcggggta
3781 aactgggaaa  gtgatgtcgt  gtactggctc  cgcttttttc  ccgaggggtg  gggagaaccg
3841 tatataagtg  cagtagtcgc  cgtgaacgtt  ctttttcgca  acgggtttgc  cgccagaaca
3901 cagctgaagc  ttcgaggggc  tcgcatctct  ccttcacgcg  cccgcgcgcc  tacctgaggc
3961 cgccatccac  gccggttag  tcgcgttctg  ccgcctcccg  cctgtggtgc  ctctgaact

```

4021 gcgtccgccg tctaggtaag tttaaagctc aggtcgagac cgggcctttg tccggcgctc
 4081 ccttgagacc tacctagact cagccggctc tccacgcttt gcctgaccct gcttgctcaa
 4141 ctctacgtct ttgtttcggt ttctgttctg cgcggttaca gatccaagct gtgaccggcg
 4201 cctacgctag atgaccgagt acaagccac ggtgcgcctc gccaccgcg acgacgtccc
 4261 cagggccgta cgcaccctcg ccgccgcgtt cgcgactac cccgccacgc gccacaccgt
 4321 cgatccggac cgcacacatcg agcgggtcac cgagctgcaa gaactcttcc tcacgcgcgt
 4381 cgggctcgac atcggcaagg tgtgggtcgc ggacgacggc gccgcggtgg cggctctggac
 4441 cacgccggag agcgtcgaag cggggcggtt gttcgccgag atcgcccgcg gcatggccga
 4501 gttgagcggg tcccggctgg ccgcgagca acagatggaa ggctctctgg cgcgcaccg
 4561 gcccaaggag cccgcgtggt tcttgccac cgtcggcgct tcgcccagac accagggcaa
 4621 ggggtctgggc agcgcctcg tgcctcccg cgtggaggcg gccgagcgcg ccggggtgcc
 4681 cgccttcttg gagacctccg cgcgccgcaa cctcccttc tacgagcgcg tcggttcac
 4741 cgtcaccgcc gacgtcgagg tgccgaagg accgcgcacc tgggtcatga cccgcaagcc
 4801 cgggtgcctga gtcgacaatc aacctctgga ttacaaaatt tgtgaaagat tgactggtat
 4861 tcttaactat gttgctcctt ttacgctatg tggatacgt gctttaatgc ctttgtatca
 4921 tgctattgct tcccgatagg ctttcatttt ctcctccttg tataaatcct ggttgctgtc
 4981 tctttatgag gaggttgtgg ccgttgctag gcaacgtggc gtgggtgtgca ctgtgtttgc
 5041 tgacgcaacc cccactggtt ggggcattgc caccacctgt cagctccttt ccgggacttt
 5101 cgctttcccc ctccctattg ccacggcgga actcatcgcc gcctgccttg ccgctgctg
 5161 gacaggggct cggtctgttg gcactgacaa ttcggtggtg ttgtcgggga aatcatcgct
 5221 ctttcccttg ctgctgcctt gtgttgccac ctggattctg cgcgggacgt ccttctgcta
 5281 cgtecccttg gccctcaatc cagcggacct tcttcccgcg gccctgctgc cggctctgcg
 5341 gcctcttccg cgtcttcgcc ttccgctca gacgagtcgg atctcccttt gggccgcctc
 5401 ccgcctgggt acctttaaga ccaatgactt acaaggcagc tgtagatctt agccactttt
 5461 taaaagaaaa ggggggactg gaagggttaa ttcactccca acgaaaataa gatctgcttt
 5521 ttgcttgtag tgggtctctc tgggttagacc gcttgccctg agtgcttcaa gtagtggtg
 5581 tagggaaacc actgcttaag ggtaactaga gatccctcag acccttttag tcagtggtga
 5641 ccgctctggt gtgtgactct tcatgtcatc ttattattca gtatttataa cttgcaaaga
 5701 aaatctctag agagagttag aggaacttgt ttattgcagc ttataatggt taaaaataaa
 5761 aatgaatatc gcaatagcat cacaaatttc acaaataaag catttttttc actgcattct agttgtggtt
 5821 tgtccaaact catcaatgta tcttatcatg tctggctcta gctatcccg ccctaactcc
 5881 gcccagttcc gcccatcttc cgccccatgg ctgactaatt ttttttattt atgcagaggc
 6001 cgaggccgcc tcggcctctg agctattcca gaagtagtga ggaggctttt ttggaggcct
 6061 agacttttgc agagacggcc caaattcgta atcatggtca tagctgtttc ctgtgtgaaa
 6121 ttgttatccg ctcaacaatc cacacaacat acgagccgga agcataaagt gtaaagcctg
 6181 ggggtgcctaa tgagttagct aactcacatt aattgcgttg cgctcactgc ccgctttcca
 6241 gtcgggaaac ctgtcgtgcc agctgcatta atgaatcggc caacgcgcgg ggagaggcgg
 6301 tttgcgtatt gggcgctctt ccgcttcctc gctcactgac tcgctgcgct cggctgttcg
 6361 gctgcggcga gcggtatcag ctactcaaaa ggcggttaata cggttatcca cagaatcagg
 6421 ggataacgca ggaaagaaca tgtgagcaaa aggccagcaa aaggccagga accgtaaaaa
 6481 ggccgcgttg ctggcgtttt tccataggct ccgccccctt gacgagcatc aaaaaaatcg
 6541 acgctcaagt cagaggtggc gaaaccgcac aggactataa agataccagg cgtttccccc
 6601 tggaagctcc ctctgctgct ctctgttcc gacctgcgc cttaccggat acctgtccgc
 6661 ctttctccct tcgggaagcg tggcgctttc tcatagctca cgctgtagg atctcagttc
 6721 ggtgtaggtc gttcgtcca agctgggctg tgtgcacgaa ccccccgttc agcccgaccg
 6781 ctgcgcctta tccggttaact atcgtcttga gtccaaccgc gtaagacacg acttatcgcc
 6841 actggcagca gccactggta acaggattag cactagaagg acagtatttg gtatctgcgc
 6901 gttcttgaag ccagttacct tcggaaaaag agttggtagc tcttgatccg gcaaaacaac
 6961 tctgctgaag agcgggtggt tttttgtttg caagcagcag attacgcgca gaaaaaaagg
 7021 caccgctggt gatcctttga tcttttctac ggggtctgac gctcagtgga acgaaaactc
 7081 atctcaagaa attttggtca tgagattatc aaaaaggatc ttcacctaga tctttttaa
 7141 acgttaaggg agtttttaaat caatctaaag tatatatgag taaacttggt ctgacagtta
 7201 ttaaaaaatga atcagtgagg cacctatctc agcgtctgt ctatttcgtt catccatagt
 7261 ccaatgctta cccgtcgtgt agataactac gatacgggag ggcttaccat ctggccccag
 7321 tgctgcaatg ataccgcgag acccacgctc accggctcca gatttatcag caataaacca
 7381 gccagccgga agggccgagc gcagaagtgg tctgcaact ttatccgcct ccatccagtc
 7441 tattaattgt tgccgggaag ctagagtaag tagttcgcca gttaatagtt tgcgcaacgt

7561 tgttgccatt gctacaggca tegtgggtgtc acgctcgtcg tttgggtatgg cttcattcag
 7621 ctccgggtcc caacgatcaa ggcgagttac atgatcccc atgttggtgca aaaaagcggg
 7681 tagctccttc ggtcctccga tegtgtgtcag aagtaagttg gccgcagtg taccactcat
 7741 gggttatggca gcaactgcata attctctttac tgtcatgcc tccgtaagat gcttttctgt
 7801 gactgggtgag tactcaacca agtcattctg agaatagtgt atgcggcgac cgagttgctc
 7861 ttgcccggcg tcaatacggg ataataccgc gccacatagc agaactttaa aagtgtcat
 7921 cattggaaaa cgttcttcgg ggcgaaaact ctcaaggatc ttaccgctgt tgagatccag
 7981 ttcgatgtaa cccactcgtg caccactg atcttcagca tcttttactt tcaccagcgt
 8041 ttctgggtga gcaaaaacag gaaggcaaaa tgccgcaaaa aagggaataa gggcgacacg
 8101 gaaatgttga atactcatac tcttcctttt tcaatattat tgaagcattt atcagggtta
 8161 ttgtctcatg agcggataca tatttgaatg tatttagaaa aataaaca aa taggggttcc
 8221 gcgcacattt ccccgaaaag tgccacctga cgtctaagaa accattatta tcatgacatt
 8281 aacctataaa aataggcgta tcacgaggcc ctttcgtctc gcgcgtttcg gtgatgacgg
 8341 tgaaaacctc tgacacatgc agctcccggg gacggtcaca gcttgctgt aagcggatgc
 8401 cgggagcaga caagcccgctc agggcgcgctc agcgggtgtt ggcgggtgtc ggggctggct
 8461 taactatgcy gcatcagagc agattgtact gagagtgcac catatgcggt gtgaaatacc
 8521 gcacagatgc gtaaggagaa aataccgcac caggcgccat tcgccattca ggctgcgcaa
 8581 ctggtgggaa gggcgatcgg tgcgggcctc ttcgctatta cgccagctgg cgaaaggggg
 8641 atgtgctgca aggcgattaa gttgggtaac gccagggttt tccagtcac gacgttgtaa
 8701 aacgacggcc agtgccaagc tg.

p510_antiCD19_LL_CD3epsilon (SEQ ID NO: 17)

1 acgcgtgtag tcttatgcaa tactcttgta gtcttgcaac atggtaacga tgagttagca
 61 acatgcctta caaggagaga aaaagcaccg tgcatgccga ttgggtggaag taaggtggta
 121 cgatcgtgcc ttattaggaa ggcaacagac gggctctgaca tggattggac gaaccactga
 181 attgccgcat tgcagagata ttgtatttaa gtgcctagct cgatacaata aacgggtctc
 241 tctgggttaga ccagatctga gcctgggagc tctctggcta actagggaac ccactgctta
 301 agcctcaata aagcttgctt tgagtgtctc aagtagtgtg tgcccgtctg ttgtgtgact
 361 ctggttaacta gagatccctc agaccctttt agtcagtgtg gaaaatctct agcagtggcg
 421 cccgaacagg gacctgaaag cgaaagggaa accagagctc tctcgacgca ggactcggct
 481 tgctgaagcg cgcacggcaa gaggcgaggg gcggcgactg gtgagtacgc caaaaatttt
 541 gactagcggg ggctagaagg agagagatgg gtgcgagagc gtcagtatta agcgggggag
 601 aattagatcg cgatgggaaa aaattcgggt aaggccaggg ggaaagaaaa aatataaatt
 661 aaaacatata gtatgggcaa gcaggagact agaacgattc gcagttaatc ctggcctgtt
 721 agaaacatca gaaggctgta gacaaatact gggacagcta caaccatccc ttcagacagg
 781 atcagaagaa cttagatcat tatataatac agtagcaacc ctctattgtg tgcataaag
 841 gatagagata aaagacacca aggaagcttt agacaagata gaggaagagc aaaacaaaag
 901 taagaccacc gcacagcaag cggccactga tcttcagacc tggaggagga gatatgaggg
 961 acaattggag aagtgaatta tataaatata aagtagtaaa aattgaacca ttaggagtag
 1021 caccaccaa ggcaaaagaga agagtgggtg agagagaaaa aagagcagtg ggaataggag
 1081 ctttgttcct tgggttcttg ggagcagcag gaagcactat gggcgagcc tcaatgacgc
 1141 tgacggtaca ggccagacaa ttattgtctg gtatagtgca gcagcagaac aatttgctga
 1201 gggctattga ggcgcaacag catctgttgc aactcacagt ctggggcatc aagcagctcc
 1261 aggcaagaat cctggctgtg gaaagatacc taaaggatca acagctcctg gggatttggg
 1321 gttgctctgg aaaactcatt tgcaccactg ctgtgccttg gaatgctagt tggagtaata
 1381 aatctctgga acagattgga atcacacgac ctggatggag tgggacagag aaattaacaa
 1441 ttacacaagc ttaatacact ccttaattga agaactgcaa aaccagcaag aaaagaatga
 1501 acaagaatta ttggaattag ataaatgggc aagtttggtg aattgggtta acataacaaa
 1561 ttggctgtgg tatataaaat tattcataat gatagtagga ggcttggtag gtttaagaat
 1621 agtttttgct gtacttttcta tagtgaatag agttaggcag ggatattcac cattatcggt
 1681 tcagaccac ccccaaccc cgaggggacc cgacaggccc gaaggaatag aagaagaagg
 1741 tggagagaga gacagagaca gatccattcg attagtgaac ggatctcgac ggtatcggtt
 1801 aacttttaaa agaaaagggg ggattggggg gtacagtgca ggggaaagaa tagtagacat
 1861 aatagcaaca gacatacaaa cttaaagaatt acaaaaacaa attacaaaat tcaaaatttt
 1921 atcgatacta gtattatgcc cagtacatga ccttatggga ctttctact tggcagtaca
 1981 tctacgtatt agtcacgct attaccatgg tgatgcgggt ttggcagtac atcaatgggc
 2041 gtggatagcg gtttgactca cggggatttc caagtctcca cccattgac gtcaatggga
 2101 gtttgttttg gcaccaaatt caacgggact ttccaaaatg tcgtaacaac tccgccccat
 2161 tgacgcaaat gggcggtagg cgtgtacggt gggaggttta tataagcaga gctcgtttag
 2221 tgaaccgtca gatcgcttgg agacgccatc cacgtgttt tgacctccat agaagattct

2281 agagccgcca ccatgcttct cctggtgaca agccttctgc tctgtgagtt accacaccca
 2341 gcattcctcc tgatcccaga catccagatg acacagacta catcctccct gtctgcctct
 2401 ctgggagaca gagtcccat cagttgcagg gcaagtcagg acattagtaa atattttaat
 2461 tgggtatcagc agaaaccaga tggaaactgtt aaactcctga tctaccatac atcaagatta
 2521 cactcaggag tcccatcaag gttcagtgge agtgggtctg gaacagatta ttctctcacc
 2581 attagcaacc tggagcaaga agatattgcc acttactttt gccaacaggg taatacgctt
 2641 ccgtacacgt tcggaggggg gactaagttg gaaataacag gctccacctc tggatccggc
 2701 aagcccggat ctggcgaggg atccaccaag ggcgaggtga aactgcagga gtcaggacct
 2761 ggcctggtgg cgccctcaca gagcctgtcc gtcacatgca ctgtctcagg ggtctcatta
 2821 cccgactatg gtgtaagctg gattcgccag cctccacgaa agggctctgga gtggctggga
 2881 gtaatatggg gtagtgaaac cacatactat aattcagctc tcaaattccag actgaccatc
 2941 atcaaggaca actccaagag ccaagttttc ttaaaaatga acagtctgca aactgatgac
 3001 acagccattt actactgtgc caaacattat tactacggtg gtagctatgc tatggactac
 3061 tgggggtcaag gaacctcagt caccgtctcc tcagcgcccg caattgaagt tatgtatcct
 3121 cctccttacc taggtggcgg cggttctggt ggcgggcggt ctggtggcgg cggttctctc
 3181 gaggatggta atgaagaaat ggggtggtatt acacagacac catataaagt ctccatctct
 3241 ggaaccacag taatattgac atgcccctcag tatcctggat ctgaaatact atggcaacac
 3301 aatgataaaa acataggcgg tgatgaggat gataaaaaca taggcagtga tgaggatcac
 3361 ctgtcactga aggaattttc agaattggag caaagtgggt attatgtctg ctaccccaga
 3421 ggaagcaaac cagaagatgc gaacttttat ctctacctga gggcaagagt gtgtgagaac
 3481 tgcattggaga tggatgtgat gtcggtggcc acaattgtca tagtggacat ctgcatcact
 3541 gggggcttgc tgetgctggt ttactactgg agcaagaata gaaaggccaa ggccaagcct
 3601 gtgacacgag gagcgggtgc tggcggcagg caaaggggac aaaacaagga gaggccacca
 3661 cctgttccca acccagacta tgagcccatc cggaaggcc agcgggacct gtattctggc
 3721 ctgaatcaga gacgcactctg ataagaattc gatccgcggc cggaaggat ctgcgatcgc
 3781 tccgggtgcc gtcagtgggc agagcgcaca tcgcccacag tcccgcagaa gttgggggga
 3841 ggggtcggca attgaacggg tgccatagaga aggtggcgcg gggtaaaactg ggaaagtgat
 3901 gtcgtgtact ggctccgcct ttttcccagag ggtgggggag aaccgtatat aagtgcagta
 3961 gtcgccgtga acgttctttt tcgcaacggg tttgcccga gaacacagct gaagcttcga
 4021 ggggctcgca tctctccttc acgcgccgc cgccctacct gaggcgcga tccacgcgg
 4081 ttgagtcgcy ttctgcgcgc tccgcctgt ggtgcctcct gaactgcgtc cgccgtctag
 4141 gtaagtttaa agctcaggtc gagaccgggc ctttgtccgg cgctcccttg gagcctacct
 4201 agactcagcc ggctctccac gctttgcctg accctgcttg ctcaactcta cgtctttggt
 4261 tcgttttctg ttctgcgcgc ttacagatcc aagctgtgac cggcgccctac gctagatgac
 4321 cgagtacaag cccacgggtgc gcctcgccac ccgcgacgac gtcccaggg ccgtacgcac
 4381 cctcgccgcc gcgttcgccc actacccgc cagcgccac accgtcgatc cggaccgcca
 4441 catcgagcgg gtcaccgagc tgcaagaact cttcctcacg cgcgtcgggc tcgacatcgg
 4501 caaggtgtgg gtcgcgagc acggcgccgc ggtggcggtc tggaccagc cggagagcgt
 4561 cgaagcgggg gcgggtgttc ccgagatcgg ccgcgcgatg gccgagttga gcgggtccc
 4621 gctggcgcgc cagcaacaga tggaaaggct cctggcgccg caccggccca aggagcccgc
 4681 gtggttcttg gccaccgtcg gctctcgcgc cgaccaccag ggcaagggtc tgggcagcgc
 4741 cgtcgtgtc cccggagtgg aggcggccga gcgcgcgggg gtgcccgcct tctggagac
 4801 ctccgcgcgc cgcaacctcc ccttctacga gcggtcggc ttcaccgtca ccgcgacgt
 4861 cgaggtgccc gaaggaccgc gcacctggtg catgaccgc aagcccgggt cctgagtcga
 4921 caatcaacct ctggattaca aaatttgtga aagattgact ggtattctta actatgttgc
 4981 tctttttacg ctatgtggat acgctgcttt aatgcctttg tatcatgcta ttgcttccc
 5041 tatggcttct attttctcct ccttgtataa atcctggttg ctgtctcttt atgaggagtt
 5101 gtggcccgtt gtcaggcaac gtggcgtggt gtgcactgtg tttgctgacg caacccccac
 5161 tgggtggggc attgccacca cctgtcagct cctttccggg actttcgtt tccccctccc
 5221 tattgccacg gcggaactca tcgcgcctg ccttgcccgc tgcaggacag gggctcggt
 5281 gttgggcact gacaattccg tgggtgtgtc ggggaaatca tcgtcctttc cttggctgct
 5341 cgctgtgtt gccacctgga ttctgcgcgg gacgtccttc tgctacgtcc ctteggccct
 5401 caatccagcg gaccttctt cccgcggcct gctgccggct ctgcggcctc ttcgcgctct
 5461 tcgccttcgc cctcagacga gtcggatctc cctttgggac gctccccgc ctggtacctt
 5521 taagaccaat gacttacaag gcagctgtag atcttagcca ctttttataa gaaaaggggg
 5581 gactggaagg gctaattcac tcccacgaa aataagatct gctttttgct tgtactgggt
 5641 ctctctggtt agaccagatc tgagcctggg agctctctgg ctaactaggg aaccactgc
 5701 ttaagcctca ataaagcttg ccttgagtgc ttcaagtagt gtgtgcccgt ctgttgtgtg
 5761 actctggtaa ctagagatcc ctacagacct tttagtcagt gtggaaaatc tctagcagta

5821 gtagttcatg tcatcttatt attcagtatt tataacttgc aaagaaatga atatcagaga
 5881 gtgagaggaa cttgtttatt gcagcttata atggttacaa ataaagcaat agcatcacaa
 5941 atttcacaaa taaagcattt ttttactgct attctagtgt tggtttgtcc aaactcatca
 6001 atgtatctta tcatgtctgg ctctagctat cccgccctta actccgcca gtccgcca
 6061 ttctccgccc catggctgac taattttttt tatttatgca gaggcgagg ccgctcggc
 6121 ctctgagcta ttccagaagt agtgaggagg cttttttgga ggcctagact tttgcagaga
 6181 cggcccaaatt tcgtaatcat ggtcatagct gtttcctgtg tgaaattgtt atccgctcac
 6241 aattccacac aacatacgag ccggaagcat aaagtgtaaa gcttggggtg cctaagtagt
 6301 gagctaactc acattaattg cgttgcgctc actgccgct ttcagtcgg gaaacctgtc
 6361 gtgccagctg cattaatgaa tcggccaacg cgcggggaga ggcggtttgc gtattgggcg
 6421 ctcttcgctc tcctcgctca ctgactcgct gcgctcggtc gttcggctgc ggcgagcggg
 6481 atcagctcac tcaaaggcgg taatacgggt atccacagaa tcaggggata acgcaggaaa
 6541 gaacatgtga gcaaaaggcc agcaaaaggc caggaaccgt aaaaaggccg cgttgctggc
 6601 gtttttccat aggcctccgc cccctgacga gcatcacaaa aatcgacgct caagtcagag
 6661 gtggcgaaac ccgacaggac tataaagata ccaggcgttt cccctggaa gctccctcgt
 6721 gcgctctcct gttccgaccc tgccgcttac cggatacctg tccgcctttc tccctcggg
 6781 aagcgtggcg ctttctcata gctcacgctg taggtatctc agttcgggtg aggtcggtcg
 6841 ctccaagctg ggctgtgtgc acgaaccccc cgttcagccc gaccgctgcg cttatccgg
 6901 taactatcgt cttgagtcca acccggttaag acacgactta tcgccactgg cagcagccac
 6961 tggtaacagg attagcagag cgaggtatgt aggcgggtgt acagagttct tgaagtgggtg
 7021 gcctaactac ggctacacta gaaggacagt atttgggtatc tgcgctctgc tgaagccagt
 7081 taccttcgga aaaagagttg gtagctcttg atccggcaaa caaacaccg ctggtagcgg
 7141 tggttttttt gtttgcaagc agcagattac gcgcagaaaa aaaggatctc aagaagatcc
 7201 tttgatcttt tctacggggg ctgacgctca gtggaacgaa aactcacgtt aagggtttt
 7261 ggtcatgaga ttatcaaaaa ggatcttcac ctagatcctt ttaaattaaa aatgaagttt
 7321 taaatcaatc taaagtatat atgagtaaac ttggctctgac agttaccaat gcttaatcag
 7381 tgaggcacct atctcagcga tctgtctatt tcgttcaccc atagttgcct gactccccgt
 7441 cgtgtagata actacgatac gggagggtct accatctggc ccagtgctg caatgatacc
 7501 gcgagaccca cgctcaccgg ctccagattt atcagcaata aaccagccag ccggaagggc
 7561 cgagcgcaga agtggtcctg caactttatc cgctccatc cagtctatta attggtgccg
 7621 ggaagctaga gtaagtagtt cgccagttaa tagtttgccg aacgttggtg ccattgctac
 7681 aggcacgtg gtgtcacgct cgtcggttggt tatggcttca ttcagctccg gttcccaacg
 7741 atcaaggcga gttacatgat ccccatggtt gtgcaaaaaa gcggttagct cttcggtcc
 7801 tccgatcgtt gtcagaagta agttggccgc agtggtatca ctcatggtta tggcagcact
 7861 gcataattct ctactgtca tgccatccgt aagatgcttt tctgtgactg gtgagtactc
 7921 aaccaagtca ttctgagaat agtgtatgcg gcgaccgagt tgctcttgcc cggcgtaaat
 7981 acgggataat accgcgccac atagcagaac tttaaaagtg ctcatcattg gaaaacgttc
 8041 ttcggggcga aaactctcaa ggatcttacc gctgttgaga tccagttcga tgaaccac
 8101 tcgtgcaccc aactgatctt cagcatcttt tactttcacc agcgtttctg ggtgagcaaa
 8161 aacaggaagg caaaatgccg caaaaaaggg aataaggcg acacggaaat gttgaatact
 8221 catactcttc ctttttcaat attattgaag cattatcag ggttattgtc tcatgagcgg
 8281 atacataatt gaatgtattt agaaaaataa acaaataggg gttccgcgca catttccccg
 8341 aaaagtgcc cctgacgtct aagaaacat tattatcatg acattaacct ataaaaatag
 8401 gcgtatcacg aggccttttc gtctcgcgcg tttcgggtgat gacggtgaaa acctctgaca
 8461 catgcagctc ccggagacgg tcacagcttg tctgtaagcg gatgccggga gcagacaagc
 8521 ccgtcagggc gcgtcagcgg gtgttgccgg gtgtcggggc tggcttaact atgcggcatc
 8581 agagcagatt gtactgagag tgcaccatat gcggtgtgaa ataccgcaca gatgcgtaag
 8641 gagaaaatac cgcacagggc gccattcgcc attcaggctg cgcaactgtt gggaaggcg
 8701 atcgggtcgg gcctcttcgc tattacgcca gctggcgaaa ggggatgtg ctgcaaggcg
 8761 attagttgg gtaacgccag ggttttccca gtcacgacgt tgtaaacga cggccagtgc
 8821 caagctg.

p510_antiCD19_SL_CD3epsilon (SEQ ID NO: 18)

1 acgcgtgtag tcttatgcaa tactcttgta gtcttgcaac atggtaacga tgagttagca
 61 acatgcctta caaggagaga aaaagcaccg tgcattgccg ttgggtggaag taagggtgga
 121 cgatcggtgc ttattaggaa ggcaacagac gggctctgaca tggattggac gaaccactga
 181 attgccgcat tgcagagata ttgtatttaa gtgcctagct cgatacaata aacgggtctc
 241 tctgggttaga ccagatctga gcctgggagc tctctggcta actaggaac ccactgctta
 301 agcctcaata aagcttgctt tgagtgtctc aagtagtgtg tgcccgtctg ttgtgtgact
 361 ctggtaacta gagatccctc agaccctttt agtcagtgtg gaaaatctct agcagtggcg

421 cccgaacagg gacctgaaag cgaaagggaa accagagctc tctcgacgca ggactcggct
 481 tgctgaagcg cgcacggcaa gaggcgaggg gcggcgactg gtgagtacgc caaaaatttt
 541 gactagcggg ggctagaagg agagagatgg gtgcgagagc gtcagtatta agcgggggag
 601 aattagatcg cgatgggaaa aaattcgggtt aaggccaggg ggaaagaaaa aatataaatt
 661 aaaacatata gtatgggcaa gcaggggagct agaacgattc gcagttaatc ctggcctgtt
 721 agaaacatca gaaggctgta gacaaatact gggacagcta caaccatccc ttcagacagg
 781 atcagaagaa cttagatcat tatataatac agtagcaacc ctctattgtg tgcatacaag
 841 gatagagata aaagacacca aggaagcttt agacaagata gaggaagagc aaaacaaaag
 901 taagaccacc gcacagcaag cggccactga tcttcagacc tggaggagga gatatgaggg
 961 acaattggag aagtgaatta tataaatata aagtagtaaa aattgaacca ttaggagtag
 1021 caccaccaa ggcaaagaga agagtgggtgc agagagaaaa aagagcagtg ggaataggag
 1081 ctttgttctt tgggttcttg ggagcagcag gaagcactat gggcgagcc tcaatgacgc
 1141 tgacggtaca ggccagacaa ttattgtctg gtatagtga gcagcagaac aatttgctga
 1201 gggctattga ggcgcaacag catctgttgc aactcacagt ctggggcatc aagcagctcc
 1261 aggcaagaat cctggctgtg gaaagatacc taaaggatca acagctcctg gggatttggg
 1321 gttgctctgg aaaactcatt tgcaccactg ctgtgccttg gaatgctagt tggagtaata
 1381 aatctctgga acagattgga atcacacgac ctggatggag tgggacagag aaattaacaa
 1441 ttacacaagc ttaatacact ccttaattga agaatcgcaa aaccagcaag aaaagaatga
 1501 acaagaatta ttggaattag ataaatgggc aagtttgttg aattggttta acataacaaa
 1561 ttggctgtgg tatataaaat tattcataat gatagtagga ggcttggtag gtttaagaat
 1621 agtttttgc gtactttcta tagtgaatag agttaggcag ggatattcac cattatcggt
 1681 tcagaccac ctcccaaccc cgaggggacc cgacaggccc gaaggaatag aagaagaagg
 1741 tggagagaga gacagagaca gatccattcg attagtgaac ggatctcgac ggtatcgggt
 1801 aactttttaa agaaaagggg ggattggggg gtacagtga ggggaaagaa tagtagacat
 1861 aatagcaaca gacatacaaa cttaaagaatt acaaaaacaa attacaaaat tcaaaatttt
 1921 atcgatacta gtattatgcc cagtacatga ccttatggga ctttcctact tggcagtaca
 1981 tctacgtatt agtcacgct attaccatgg tgatgcgggt ttggcagtag atcaatgggc
 2041 gtggatagcg gtttgactca cggggatttc caagtctcca cccattgac gtcaatggga
 2101 gtttgttttg gcaccaaatt caacgggact ttccaaaatg tcgtaacaac tccgccccat
 2161 tgacgcaaatt gggcggtagg cgtgtacggt gggaggttta tataagcaga gctcgtttag
 2221 tgaaccgtca gatcgcttg agacgccatc cacgctgttt tgacctccat agaagattct
 2281 agagccgcca ccatgcttct cctggtgaca agccttctgc tctgtgagtt accacacca
 2341 gcattcctcc tgatcccaga catccagatg acacagacta catcctccct gtctgcctct
 2401 ctgggagaca gagtcacat cagttgcagg gcaagtcagg acattagtaa atatttaaat
 2461 tggatcagc agaaaccaga tggaaactgt aaactcctga tctaccatac atcaagatta
 2521 cactcaggag tcccatcaag gttcagtggt agtgggtctg gaacagatta ttctctcacc
 2581 attagcaacc tggagcaaga agatattgcc acttactttt gccaacaggg taatacgtt
 2641 ccgtacacgt tcggaggggg gactaagtgt gaaataacag gctccacctc tggatccggc
 2701 aagcccggt ctggcgaggg atccaccaag ggcgaggtga aactgcagga gtcaggacct
 2761 ggcttgggtg cgcctcaca gagcctgtcc gtcacatgca ctgtctcagg ggtctcatta
 2821 ccgactatg gtgtaagctg gattcgccag cctccacgaa agggctctga gtggctggga
 2881 gtaatatggg gtagtgaac cacatactat aattcagctc tcaaattccag actgaccatc
 2941 atcaaggaca actccaagag ccaagtttct ttaaaaatga acagtctgca aactgatgac
 3001 acagccattt actactgtgc caaacattat tactacgggt gtagctatgc tatggactac
 3061 tgggggtcaag gaacctcagt caccgtctcc tcagcggccg caggtggcgg cggttctggg
 3121 ggcggcggtt ctggtggcgg cggttctctc gaggatggta atgaagaaat ggggtggtatt
 3181 acacagacac catataaagt ctccatctct ggaaccacag taatattgac atgccctcag
 3241 tatcctggat ctgaaatact atggcaacac aatgataaaa acataggcgg tgatgaggat
 3301 gataaaaaca taggcagtga tgaggatcac ctgtcactga aggaattttc agaattggag
 3361 caaagtgtt attatgtctg ctaccccaga ggaagcaaac cagaagatgc gaacttttat
 3421 ctctacctga gggcaagagt gtgtgagaac tgcattggaga tggatgtgat gtcgggtggc
 3481 acaattgtca tagtggacat ctgcatcact gggggcttgc tgctgtggt ttactactgg
 3541 agcaagaata gaaaggccaa ggccaagcct gtgacacgag gagcgggtgc tggcggcagg
 3601 caaaggggac aaaacaagga gaggccacca cctgttccca acccagacta tgagcccatc
 3661 cggaaaggcc agcgggacct gtattctggc ctgaatcaga gacgcactg ataagaattc
 3721 gatccgcggc cgcgaaggat ctgcatcgc tccggtgcc gtcagtgggc agagcgcaca
 3781 tcgcccacag tccccgagaa gttgggggga ggggtcggca attgaacggg tgccatagaga
 3841 aggtggcgcg gggtaaactg ggaaagtgat gtcgtgtact ggctccgct ttttcccgag
 3901 ggtgggggag aaccgtatat aagtgcagta gtcgccgtga acgttctttt tcgcaacggg

3961 tttgccgcca gaacacagct gaagcttcga ggggctcgca tctctccttc acgcgcccgc
 4021 cgccctacct gaggcgcgcca tccacgcggg ttgagtcgcg ttctgcgcgc tcccgctgt
 4081 ggtgcctcct gaactgcgctc cgccgtctag gtaagtttaa agctcaggctc gagaccgggc
 4141 ctttgtccgg cgctcccttg gagcctacct agactcagcc ggctctccac gctttgcctg
 4201 accctgcttg ctcaactcta cgtctttgtt tcgtttctg ttctgcgcgc ttacagatcc
 4261 aagctgtgac cggcgcctac gctagatgac cgagtacaag cccacgggtgc gcctcgccac
 4321 ccgcgacgac gtccccaggg ccgtacgcac cctcgccgcg gcgttcgccc actaccccgc
 4381 cacgcgccac accgtcgatc cggaccgcca catcgagcgg gtcaccgagc tgcaagaact
 4441 cttcctcacg cgcgtcgggc tcgacatcgg caaggtgtgg gtcgcggacg acggcgccgc
 4501 ggtggcggtc tggaccacgc cggagagcgt cgaagcgggg gcggtgttcg ccgagatcgg
 4561 cccgcgcatg gccgagttga gcggttccc gctggcgcg cagcaacaga tggaggcct
 4621 cctggcgccg caccggccca aggagccgc gtggttctg gccaccgtcg gcgtctcgcc
 4681 cgaccaccag ggcaagggtc tgggcagcgc cgtcgtgctc cccggagtgg aggcggccga
 4741 gcgcgcggg gtgcccgcct tcctggagac ctccgcgcc cgcaacctcc cttctacga
 4801 gcggctcggc ttcaccgtca ccgcgacgt cgaggtgcc gaaggaccgc gcacctggtg
 4861 catgaccgcg aagcccggtg cctgagtcga caatcaacct ctggattaca aaatttgtga
 4921 aagattgact ggtattctta actatgttgc tccttttaag ctatgtggat acgctgcttt
 4981 aatgcctttg tatcatgcta ttgcttccg tatggctttc atttctctc cttgtataa
 5041 atcctggttg ctgtctcttt atgaggagtt gtggcccgtt gtcaggcaac gtggcggtg
 5101 gtgcactgtg tttgctgacg caacccccac tggttggggc attgccacca cctgtcagct
 5161 cctttccggg actttcgctt tccccctccc tattgccacg gcggaactca tcgcgcctg
 5221 ccttgcccgc tgetggacag gggctcggct gttgggcaat gacaattccg tgggtgtgtc
 5281 ggggaaatca tcgtcctttc cttggctgct cgctgtgtt gccacctgga ttctgcgcgc
 5341 gacgtccttc tgetacgtcc cttcgccct caatccagcg gaccttctt cccgcggcct
 5401 gctgcgcgct ctgcggcctc ttccgcgtct tcgccttcgc cctcagacga gtcggatctc
 5461 cctttggggc gcctccccgc ctggtacctt taagaccaat gacttacaag gcagctgtag
 5521 atccttagcca ctttttaaaa gaaaaggggg gactggaagg gctaattcac tccaacgaa
 5581 aataagatct gctttttgtc tgtactgggt ctctctgggt agaccagatc tgagcctggg
 5641 agctctctgg ctaactaggg aaccactgct ttaagcctca ataaagcttg ccttgagtgc
 5701 ttcaagtagt gtgtgcccgt ctgttggtg actctggtaa ctagagatcc ctcagacct
 5761 tttagtcagt gtggaaaatc tctagcagta gtagttcatg tcatcttatt attcagtatt
 5821 tataacttgc aaagaaatga atatcagaga gtgagaggaa cttgtttatt gcagcttata
 5881 atggttacaa ataaagcaat agcatcacia atttcacaaa taaagcattt ttttactgc
 5941 attctagttg tggtttgtcc aaactcatca atgtatctta tcatgtctgg ctctagctat
 6001 cccgccccta actccgcccc gtccgcccc ttctccgccc catggctgac taatttttt
 6061 tatttatgca gaggcgcgag ccgcctcggc ctctgagcta tccagaagt agtgaggagg
 6121 cttttttgga ggcctagact tttgcagaga cggcccaaat tcgtaatcat ggtcatagct
 6181 gtttctctgt tgaaattgtt atccgctcac aattccacac aacatacgag ccggaagcat
 6241 aaagtgtaaa gcctgggggt cctaattgag gaaacctgtc gagctaactc acattaattg cgttgcgctc
 6301 actgcccgtc ttccagtcgg gaaacctgtc gtgccagctg cattaatgaa tcggccaacg
 6361 cgcggggaga ggcgggtttgc gtattggggc ctcttcgctc tctcgcctca ctgactcgct
 6421 gcgctcggtc gttcggctgc ggcgagcgg atcagctcac tcaaaggcgg taatacgggt
 6481 atccacagaa tcaggggata acgcaggaaa gaacatgtga gcaaaaggcc agcaaaaggc
 6541 caggaaccgt aaaaaggcgg cgttgctggc gtttttccat aggtccgccc cccctgacga
 6601 gcatcacaaa aatcgacgct caagtcagag gtggcgaaac ccgacaggac tataaagata
 6661 ccaggcggtt cccctggaa gctccctcgt gcgctctct gttccgacct tgccgcttac
 6721 cggtacactg tcgccttttc tccttcggg aagcgtggcg cttctctata gctcacgctg
 6781 taggtatctc agttcggtgt aggtcgttcg ctccaagctg ggctgtgtgc acgaacccc
 6841 cgttcagccc gaccgtgctg ccttatccgg taactatcgt cttgagtcca acccggtag
 6901 acacgactta tcgccactgg cagcagccac tggtaacagg attagcagag cgaggatagt
 6961 aggcgggtgt acagagttct tgaagtgggt gcctaactac ggctacacta gaaggacagt
 7021 atttggtatc tgcgctctgc tgaagccagt taccttcgga aaaagagttg gtagctcttg
 7081 atccggcaaa caaaccaccg ctggtagcgg tggttttttt gtttgcaagc agcagattac
 7141 gcgcagaaaa aaaggatctc aagaagatcc tttgatcttt tctacgggt ctgacgctca
 7201 gtggaacgaa aactcacgtt aagggatttt ggtcatgaga ttatcaaaaa ggatcttcac
 7261 ctagatcctt ttaaattaaa aatgaagttt taaatcaatc taaagtatat atgagtaaac
 7321 ttggtctgac agttaccaat gcttaatcag tgaggcacct atctcagcga tctgtctatt
 7381 tcgttccatc atagttgcct gactccccgt cgtgtagata actacgatac gggagggctt
 7441 accatctggc cccagtgctg caatgatacc gcgagaccca cgctcaccgg ctccagattt

7501 atcagcaata aaccagccag ccggaagggc cgagcgcaga agtgggtcctg caactttatc
 7561 cgcctccatc cagtctatta attgttgccg ggaagctaga gtaagtagtt cgccaggttaa
 7621 tagtttgccg aacgttggtg ccattgctac aggcacgtg gtgtcacgct cgtcggttg
 7681 tatggcttca ttcagctccg gttcccaacg atcaaggcga gttacatgat ccccatgtt
 7741 gtgcaaaaaa gcggttagct ccttcgggtc tccgatcggt gtcagaagta agttggccgc
 7801 agtggttatca ctcatggtta tggcagcact gcataattct cttactgtca tgccatccgt
 7861 aagatgcttt tctgtgactg gtgagtactc aaccaagtca ttctgagaat agtgatgctg
 7921 gcgaccgagt tgcctctgcc cggcgtcaat acgggataat accgcgccac atagcagaac
 7981 tttaaaagtg ctcatcattg gaaaacgttc ttcggggcga aaactctcaa ggatcttacc
 8041 gctgttgaga tccagttcga tgtaacccac tctgtcacc aactgatctt cagcatcttt
 8101 tacttttacc agcgtttctg ggtgagcaaa aacaggaagg caaaatgccg caaaaaagg
 8161 aataagggcg acacggaaat gttgaatact catactcttc ctttttcaat attattgaag
 8221 catttatcag ggttattgtc tcatgagcgg atacatattt gaatgtattt agaaaaataa
 8281 acaaataagg gttccgcgca catttccccg aaaagtgcc cctgacgtct aagaaacat
 8341 tattatcatg acattaacct ataaaaatag gcgtatcacg aggccctttc gtctcgcgcg
 8401 tttcgggtgat gacggtgaaa acctctgaca catgcagctc cgggagacgg tcacagcttg
 8461 tctgtaagcg gatgccggga gcagacaagc ccgtcagggc gcgtcagcgg gtgttgccgg
 8521 gtgtcggggc tggcttaact atgcggcatc agagcagatt gtactgagag tgcaccatat
 8581 gcggtgtgaa ataccgcaca gatgcgtaag gagaaaatac cgcacagggc gccattcgcc
 8641 attcaggtcg cgcaactgtt ggggaagggcg atcgggtcgg gcctcttcgc tattacgcca
 8701 gctggcgaaa gggggatgtg ctgcaaggcg attaagttgg gtaacgccag ggttttccca
 8761 gtcacgacgt tgtaaaacga cggccagtgc caagctg.

p510_antiCD19_SL_CD3gamma (SEQ ID NO: 19)

1 acgcgtgtag tcttatgcaa tactcttgta gtcttgcaac atggtaacga tgagttagca
 61 acatgcctta caaggagaga aaaagcaccg tgcatgccga ttggtggaag taaggtggtta
 121 cgatcggtgc ttattaggaa ggcaacagac gggctctgaca tggattggac gaaccactga
 181 attgccgcat tgcagagata ttgtatttaa gtgcctagct cgatacaata aacgggtctc
 241 tctgggttaga ccagatctga gcctgggagc tctctggcta actagggaac ccactgctta
 301 agcctcaata aagcttgccct tgagtgtctc aagtagtgtg tgcccgctctg ttgtgtgact
 361 ctggttaacta gagatccctc agaccctttt agtcagtgtg gaaaatctct agcagtggcg
 421 cccgaacagg gacctgaaag cgaaagggaa accagagctc tctcgacgca ggactcggct
 481 tgctgaagcg cgcacggcaa gaggcgaggg gcggcgactg gtgagtacgc caaaaaattt
 541 gactagcggg ggctagaagg agagagatgg gtgcgagagc gtcagtatta agcgggggag
 601 aattagatcg cgatgggaaa aaattcgggt aaggccaggg ggaaagaaaa aatataaatt
 661 aaaacatata gtatgggcaa gcagggagct agaacgattc gcagttaatc ctggcctgtt
 721 agaaacatca gaaggctgta gacaaatact gggacagcta caaccatccc ttcagacagg
 781 atcagaagaa cttagatcat tatataatac agtagcaacc ctctattgtg tgcatacaag
 841 gatagagata aaagacacca aggaagcttt agacaagata gaggaagagc aaaacaaaag
 901 taagaccacc gcacagcaag cggccactga tcttcagacc tggaggagga gatatgaggg
 961 acaattggag aagtgaatta tataaatata aagtagtaaa aattgaacca ttaggagtag
 1021 caccaccaa ggcaagaga agagtgggtg agagagaaaa aagagcagtg ggaataggag
 1081 ctttggttct tgggttcttg ggagcagcag gaagcactat gggcgagcc tcaatgacgc
 1141 tgacggtaca ggccagacaa ttattgtctg gtatagtgca gcagcagaac aatttgctga
 1201 gggctattga ggcgcaacag catctgttgc aactcacagt ctggggcatc aagcagctcc
 1261 aggcaagaat cctggctgtg gaaagatacc taaaggatca acagctcctg gggatttggg
 1321 gttgctctgg aaaactcatt tgcaccactg ctgtgccttg gaatgctagt tggagtaata
 1381 aatctctgga acagattgga atcacacgac ctggatggag tgggacagag aaattaacaa
 1441 ttacacaagc ttaatacact ccttaattga agaactgcaa aaccagcaag aaaagaatga
 1501 acaagaatta ttggaattag ataaatgggc aagtttggtg aattgggtta acataacaaa
 1561 ttggctgtgg tatataaaat tattcataat gatagtagga ggcttggtag gtttaagaat
 1621 agtttttgcgt gtactttctc tagtgaatag agttaggcag ggatattcac cattatcggt
 1681 tcagaccacac ctcccaaccc cgaggggacc cgacaggccc gaaggaatag aagaagaagg
 1741 tggagagaga gacagagaca gatccattcg attagtgaac ggatctcgac ggtatcggtt
 1801 aactttttaa agaaaagggg ggattggggg gtacagtgca ggggaaagaa tagtagacat
 1861 aatagcaaca gacatacaaa cttaaagaatt acaaaaaaaa attacaaaat tcaaaatttt
 1921 atcgatacta gtattatgcc cagtacatga ccttatggga ctttctact tggcagtaca
 1981 tctacgtatt agtcacgtct attaccatgg tgatgcgggt ttggcagtag atcaatgggc
 2041 gtggatagcg gtttgactca cggggatttc caagtctcca cccattgac gtcaatggga
 2101 gtttggtttg gcacccaaat caacgggact ttccaaaatg tcgtaacaac tccgccccat

2161 tgacgcaaatt gggcggttagg cgtgtacggt gggaggttta tataagcaga gctcgttttag
 2221 tgaaccgtca gatcgcttgg agacgccatc cacgctgttt tgacctcoat agaagatttct
 2281 agagccgcca ccatgcttct cctggtgaca agccttctgc tctgtgagtt accacaccca
 2341 gcattcctcc tgatcccaga catccagatg acacagacta catcctccct gtctgcctct
 2401 ctgggagaca gagtcacat cagttgcagg gcaagtcagg acattagtaa atattttaat
 2461 tgggtatcagc agaaaccaga tggaaactgtt aaactcctga tctaccatac atcaagatta
 2521 cactcaggag tcccatcaag gttcagtggtc agtgggtctg gaacagatta ttctctcacc
 2581 attagcaacc tggagcaaga agatattgcc acttactttt gccaacaggg taatacgtct
 2641 ccgtacacgt tcggaggggg gactaagttg gaaataacag gctccacctc tggatccggc
 2701 aagcccggat ctggcgaggg atccaccaag ggcgaggtga aactgcagga gtcaggacct
 2761 ggcctggtgg cgccctcaca gagcctgtcc gtcacatgca ctgtctcagg ggtctcatta
 2821 cccgactatg gtgtaagctg gattcgccag cctccacgaa aggggtctgga gtggctggga
 2881 gtaatatggg gtagtgaaac cacatactat aattcagctc tcaaataccag actgaccatc
 2941 atcaaggaca actccaagag ccaagttttc ttaaaaatga acagtctgca aactgatgac
 3001 acagccattt actactgtgc caaacattat tactacggtg gtagctatgc tatggactac
 3061 tgggggtcaag gaacctcagt caccgtctcc tcagcggccg caggtggcgg cggttctggt
 3121 ggcggcggtt ctggtggcgg cggttctctc gagcagtcga tcaaaggaaa ccacttggtt
 3181 aaggtgtatg actatcaaga agatggttcg gtacttctga cttgtgatgc agaagccaaa
 3241 aatatcacat ggttttaaaga tgggaagatg atcggcttcc taactgaaga taaaaaaaaa
 3301 tggaatctgg gaagtaatgc caaggaccca cgagggatgt atcagtgtaa aggatcacag
 3361 aacaagtcaa aaccactcca agtgtattac agaattgtgc agaactgcat tgaactaat
 3421 gcagccacca tatctggctt tctctttgct gaaatcgtca gcattttcgt ccttgctgtt
 3481 ggggtctact tcattgctgg acaggatgga gttcgccagt cgagagcttc agacaagcag
 3541 actctgttgc ccaatgacca gctctaccag cccctcaagg atcgagaaga tgaccagtac
 3601 agccaccttc aaggaaacca gttgaggagg aattgataag aattcgatcc gcggccgcca
 3661 aggatctgcg atcgctccgg tgcctcgtag tgggcagagc gcacatcgcc cacagtcctc
 3721 gagaagttgg ggggaggggt cggcaattga acgggtgcct agagaaggtg gcgcggggta
 3781 aactgggaaa gtgatgtcgt gtactggctc cgcttttttc ccgaggggtg gggagaaccg
 3841 tatataagtg cagtagtcgc cgtgaacgtt ctttttcgca acgggtttgc cgccagaaca
 3901 cagctgaagc ttcgaggggc tcgcatctct ccttcacgcg cccgcgcgcc tacctgaggc
 3961 cgccatccac gccgggttag tgcggttctg ccgcctcccg cctgtggtgc ctctgaact
 4021 gcgtccgcgg tctaggttag tttaaagctc aggtcgagac cgggcctttg tccggcgctc
 4081 ccttgaggcc tacctagact cagccggctc tccacgcttt gctgacctt gcttgctcaa
 4141 ctctacgtct ttgtttcgtt ttctgttctg cgccgttaca gatccaagct gtgaccggcg
 4201 cctacgctag atgaccgagt acaagccac ggtgcgcctc gccaccgcgg acgacgtccc
 4261 cagggccgta cgcacctcgc ccgcgcggtt cgccgactac cccgccacgc gccacaccgt
 4321 cgatccggac cgccacatcg agcgggtcac cgagctgcaa gaactcttcc tcacgcgcgt
 4381 cgggctcgac atcggcaagg tgtgggtcgc ggacgacggc gccgcggtgg cggctgggac
 4441 cagcccgag agcgtcgaag cgggggcggt gttcgccgag atcggccgcg gcatggccga
 4501 gttgagcggg tcccggctgg ccgcgcagca acagatggaa ggctccttgg cgcgcaccg
 4561 gcccaaggag ccgcgctggt tccgtggccac cgtcggcgct tcgcccagacc accagggcaa
 4621 ggggtctgggc agcgcctcgc tgcctcccgg agtgaggcg gccgagcgcg ccgggggtgc
 4681 cgcttctctg gagacctcgc cgcccgcgaa cctccccttc tacgagcggc tcggcttcac
 4741 cgtcacgcgc gacgtcgagg tgcccgaagg accgcgcacc tgggtgcatga cccgcaagcc
 4801 cgggtgcctga gtcgacaatc aacctctgga ttacaaaatt tgtgaaagat tgactggtat
 4861 tcttaactat gttgctcctt ttacgtatg tggatacgt gctttaatgc ctttgtatca
 4921 tgctattgct tcccgtatgg ctttcatctt cctcctcttg tataaatcct ggttgctgtc
 4981 tctttatgag gagttgtggc ccgttgtcag gcaacgtggc caccacctgt cagctccttt ccgggacttt
 5041 tgacgcaacc cccactggtt ggggcattgc caccacctgt actcctcgc cctgctctg
 5101 cgctttcccc ctccctattg ccacggcgga actcctcgc cctgctctg cccgctgctg
 5161 gacaggggct cggtctgttg gcaactgaca ttccgtgggt ttgtcgggga aatcatcgct
 5221 ctttctcttg ctgctcgctt gtgttgccac ctggattctg cgcgggacgt ccttctgcta
 5281 cgtcccttcg gccctcaatc cagcggacct tccttccgc ggctgctgc cggtctgctg
 5341 gcctcttccg cgtcttcgcc ttccgacctc gacgagtcgg atctcccttt ggccgcctc
 5401 ccgcctgggt acctttaaga ccaatgactt acaaggcagc tgtagatctt agccactttt
 5461 taaaagaaaa ggggggactg gaagggttaa ttcactccca acgaaaataa gatctgcttt
 5521 ttgcttgtag tgggtctctc tgggttagacc agatctgagc ctgggagctc tctggctaac
 5581 tagggaaacc actgcttaag cctcaataaa gcttgcttgg agtgcttcaa gtagtggtg
 5641 ccgctctggt gtgtgactct ggtaactaga gatccctcag acccttttag tcagtgtgga

5701 aaatctctag cagtagtagt tcatgtcatc ttattattca gtatttataa cttgcaaaga
 5761 aatgaatatc agagagtgag aggaacttgt ttattgcagc ttataatggg taaaaataaa
 5821 gcaatagcat cacaaatttc acaaataaag cttttttttc actgcattct agttgtgggt
 5881 tgtccaaact catcaatgta tcttatcatg tctggctcta gctatccgc ccctaactcc
 5941 gccagttcc gccattctc cgcctcatgg ctgactaatt ttttttattt atgcagaggc
 6001 cgaggccgcc tcggcctctg agctattcca gaagtagtga ggaggctttt ttggaggcct
 6061 agacttttgc agagacggcc caaatcgtat atcatgggtc tagctgtttc ctgtgtgaaa
 6121 ttgttatccg ctccacaattc cacacaacat acgagccgga agcataaagt gtaaagcctg
 6181 ggggtgcctaa tgagttagct aactcacatt aattgcgttg cgctcactgc ccgctttcca
 6241 gtcgggaaac ctgtcgtgcc agctgcatta atgaatcggc caacgcgcgg ggagaggcgg
 6301 tttgcgtatt gggcgctctt ccgcttcctc gctcactgac tcgctgcgct cggctcgttcg
 6361 gctgcggcga gcggtatcag ctactcaaaa ggcggttaata cgggttatcca cagaatcagg
 6421 ggataacgca ggaaagaaca tgtgagcaaa aggccagcaa aaggccagga accgtaaaaa
 6481 ggccgcgttg ctggcgtttt tccataggct ccgccccctt gacgagcatc aaaaaaatcg
 6541 acgctcaagt cagagggtggc gaaacccgac aggactataa agataccagg cgtttccccc
 6601 tggaaagctc ctctgctgct ctctgtttcc gacctgccc cttaccggat acctgtccgc
 6661 ctttctccct tcgggaagcg tggcgctttc tcatagctca cgctgtagggt atctcagttc
 6721 ggtgtagggt gttcgtctca agctgggctg tgtgcacgaa cccccggtt agcccgaccg
 6781 ctgcgcctta tccggttaact atcgtcttga gtccaacccg gtaagacacg acttatcgcc
 6841 actggcagca gccactggta acaggattag cagagcgagg tatgtaggcg gtgctacaga
 6901 gttcttgaag tgggtggccta actacggcta cactagaagg acagtatttg gtatctgcgc
 6961 tctgctgaag ccagttacct tcggaaaaag agttggtagc tcttgatccg gcaaacaaac
 7021 caccgctggt agcgggtggt tttttgtttg caagcagcag attacgcgca gaaaaaaagg
 7081 atctcaagaa gatcctttga tcttttctac ggggtctgac gctcagtgga acgaaaactc
 7141 acgttaaggg attttgggtca tgagattatc aaaaaggatc ttcacctaga tctttttaa
 7201 ttaaaaaatga agtttttaaat caatctaaag tatatatgag taaacttggt ctgacagtta
 7261 ccaatgctta atcagtgagg cacctatctc agcatctgt ctatttcggt catccatagt
 7321 tgcctgactc cccgctggtg agataactac gatacgggag ggcttaccat ctggccccag
 7381 tgctgcaatg ataccgcgag acccacgctc accggctcca gatttatcag caataaacca
 7441 gccagccgga agggccgagc gcagaagtgg tctgcaact ttatccgctt ccatccagtc
 7501 tattaattgt tgccgggaag ctagagtaag tagttcgcca gttaaatagtt tgcgcaacgt
 7561 tggtgccatt gctacaggca tcgtggtgtc acgctcgctc tttggtagtg cttcattcag
 7621 ctccggttcc caacgatcaa ggcgagttac atgatcccc atgttggtgca aaaaagcggg
 7681 tagctccttc ggtcctccga tcgttgtcag aagtaagttg gccgcagtg tactactcat
 7741 ggttatggca gcactgcata attctcttac tgtcatgcca tccgtaagat gcttttctgt
 7801 gactggtgag tactcaacca agtcattctg agaatagtg atgcggcgac cgagttgtct
 7861 ttgcccggcg tcaatacggg ataataccgc gccacatagc agaactttaa aagtgtcat
 7921 cattggaaaa cgttcttcgg ggcgaaaact ctcaaggatc ttaccgctgt tgagatccag
 7981 ttcgatgtaa cccactcgtg caccactg atcttcagca tcttttactt tcaccagcgt
 8041 ttctgggtga gcaaaaacag gaaggcaaaa tgcgcgcaaaa aagggaataa gggcgacacg
 8101 gaaatgttga atactcatat tcttcttttt tcaatattat tgaagcattt atcagggtta
 8161 ttgtctcatg agcggatata tatttgaatg tatttagaaa aataaacaata taggggttcc
 8221 gcgcacattt ccccgaaaag tgccacctga cgtctaagaa accattatta tcatgacatt
 8281 aacctataaa aataggcgta tcacgaggcc ctttcgtctc gcgcgtttcg gtgatgacgg
 8341 tgaaaacctc tgacacatgc agctcccgga gacgggtcaca gcttgctgtt aagcggatgc
 8401 cgggagcaga caagcccgct agggcgcgct agcgggtgtt ggccgggtgtc ggggctggct
 8461 taactatgct gcatcagagc agattgtact gagagtgcac catatgcggt gtgaaatacc
 8521 gcacagatgc gtaaggagaa aataccgcat caggcgccat tcgcccattc ggctgcgcaa
 8581 ctgttgaggaa gggcgatcgg tgccggcctc ttcgctatta cgccagctgg cgaaaggggg
 8641 atgtgctgca aggcgattaa gttgggtaac gccagggttt tcccagtcac gacgttgtaa
 8701 aacgacggcc agtgccaagc tg.

p510_antiCD19_SL_CD3delta (SEQ ID NO: 20)

1 acgctgttag tcttatgcaa tactcttgta gtcttgcaac atggtaacga tgagtttagca
 61 acatgcctta caaggagaga aaaagcaccg tgcatgccga ttgggtggaag taagggtggt
 121 cgatcgtgcc ttattaggaa ggcaacagac ggggtctgaca tggattggac gaaccactga
 181 attgccgcat tgcagagata ttgtatttaa gtgcctagct cgatacaata aacgggtctc
 241 tctgggttaga ccagatctga gcctgggagc tctctggcta actagggaa ccactgctta
 301 agcctcaata aagcttgctt tgagtgtctc aagtagtgtg tgcccgtctg ttgtgtgact
 361 ctggtaacta gagatccctc agaccctttt agtcagtggt gaaaatctct agcagtgggc

421 cccgaacagg gacctgaaag cgaaagggaa accagagctc tctcgacgca ggactcggct
 481 tgctgaagcg cgcacggcaa gaggcgaggg gcggcgactg gtgagtagcg caaaaatttt
 541 gactagcggg ggctagaagg agagagatgg gtgcgagagc gtcagtatta agcgggggag
 601 aattagatcg cgatgggaaa aaattcgggtt aaggccaggg ggaaagaaaa aatataaatt
 661 aaaacatata gtatgggcaa gcaggggagct agaacgattc gcagttaatc ctggcctgtt
 721 agaaacatca gaaggctgta gacaaatact gggacagcta caaccatccc ttcagacagg
 781 atcagaagaa cttagatcat tatataatac agtagcaacc ctctattgtg tgcatacaag
 841 gatagagata aaagacacca aggaagcttt agacaagata gaggaagagc aaaacaaaag
 901 taagaccacc gcacagcaag cggccactga tcttcagacc tggaggagga gatatgaggg
 961 acaattggag aagtgaatta tataaatata aagtagtaaa aattgaacca ttaggagtag
 1021 caccacacaa ggcaaagaga agagtgggtgc agagagaaaa aagagcagtg ggaataggag
 1081 ctttgttctt tgggttcttg ggagcagcag gaagcactat gggcgagcc tcaatgacgc
 1141 tgacggtaca ggccagacaa ttattgtctg gtatagtga gcagcagaac aatttgctga
 1201 gggctattga ggcgcaacag catctgttgc aactcacagt ctggggcatc aagcagctcc
 1261 aggcaagaat cctggctgtg gaaagatacc taaaggatca acagctcctg gggatttggg
 1321 gttgctctgg aaaactcatt tgcaccactg ctgtgccttg gaatgctagt tggagtaata
 1381 aatctctgga acagattgga atcacacgac ctggatggag tgggacagag aaattaacaa
 1441 ttacacaagc ttaatacact ccttaattga agaatcgcaa aaccagcaag aaaagaatga
 1501 acaagaatta ttggaattag ataaatgggc aagtttgttg aattggttta acataacaaa
 1561 ttggctgtgg tatataaaat tattcataat gatagtagga ggcttggtag gtttaagaat
 1621 agtttttgc tacttttcta tagtgaatag agttaggcag ggatattcac cattatcggt
 1681 tcagaccac ccccaaccc cgaggggacc cgacaggccc gaaggaatag aagaagaagg
 1741 tggagagaga gacagagaca gatccattcg attagtgaac ggatctcgac ggtatcgggt
 1801 aactttttaa agaaaagggg ggattggggg gtacagtga ggggaaagaa tagtagacat
 1861 aatagcaaca gacatacaaa ctaaagaatt acaaaaacaa attacaaaat tcaaaatttt
 1921 atcgatacta gtattatgcc cagtacatga ccttatggga ctttcctact tggcagtaca
 1981 tctacgtatt agtcacgct attaccatgg tgatgcgggt ttggcagtag atcaatgggc
 2041 gtggatagcg gtttgactca cggggatttc caagtctcca cccattgac gtcaatggga
 2101 gtttgttttg gcacccaaat caacgggact ttccaaaatg tcgtaacaac tccgccccat
 2161 tgacgcaa at gggcggtagg cgtgtacggt gggaggttta tataagcaga gctcgtttag
 2221 tgaaccgtca gatcgctgg agacgccatc cacgctgttt tgacctccat agaagattct
 2281 agagccgcca ccatgcttct cctggtgaca agccttctgc tctgtgagtt accacacca
 2341 gcattcctcc tgatcccaga catccagatg acacagacta catcctccct gtctgcctct
 2401 ctgggagaca gagtcacat cagttgcagg gcaagtcagg acattagtaa atatttaaat
 2461 tggatcagc agaaaccaga tggaaactgt aaactcctga tctaccatac atcaagatta
 2521 cactcaggag tcccatcaag gttcagtggt agtggtctg gaacagatta ttctctcacc
 2581 attagcaacc tggagcaaga agatattgcc acttactttt gccaacaggg taatacgctt
 2641 ccgtacacgt tcggaggggg gactaagttg gaaataacag gctccacctc tggatccggc
 2701 aagcccggt ctggcgaggg atccaccaag ggcgaggtga aactgcagga gtcaggacct
 2761 ggcttgggtg cgcctcaca gagcctgtcc gtcacatgca ctgtctcagg ggtctcatta
 2821 cccgactatg gtgtaagctg gattcgccag cctccacgaa agggctctga gtggctggga
 2881 gtaatatggg gtagtgaac cacatactat aattcagctc tcaaateccag actgaccatc
 2941 atcaaggaca actccaagag ccaagtttct ttaaaaatga acagtctgca aactgatgac
 3001 acagccattt actactgtgc caaacattat tactacgggt gtagctatgc tatggactac
 3061 tgggggtcaag gaacctcagt caccgtctcc tcagcggccg caggtggcgg cgggtctggt
 3121 ggcggcggtt ctggtggcgg cggttctctc gagttcaaga tacctataga ggaacttgag
 3181 gacagagtgt ttgtgaattg caataccagc atcacatggg tagagggaac ggtgggaaca
 3241 ctgctctcag acattacaag actggacctg ggaaaacgca tcttggaacc acgaggaata
 3301 tataggtgta atgggacaga tatatacaag gacaaagaat ctaccgtgca agttcattat
 3361 cgaatgtgcc agagctgtgt ggagctggat ccagccaccg tggctggcat cattgtcact
 3421 gatgtcattg ccactctgct cettgtcttg ggagtcttct gcttctgctg acatgagact
 3481 ggaaggctgt ctggggctgc cgacacacaa gctctgttga ggaatgacca ggtctatcag
 3541 cccctccgag atcgagatga tgcctcagtag agccaccttg gaggaactg ggctcggaac
 3601 aagtgataag aattcgatcc gcggccgcga aggatctgcg atcgctccgg tgcccgtcag
 3661 tgggcagagc gcacatcgcc cacagtcccc gagaagttgg ggggaggggt cggcaattga
 3721 acgggtgcct agagaagggt gcgcggggta aactgggaaa gtgatgtcgt gtactggctc
 3781 cgcttttttc ccgaggggtg gggagaaccg tatataagtg cagtagtcgc cgtgaacgtt
 3841 ctttttcgca acgggtttgc cgccagaaca cagctgaagc ttcgaggggc tcgcatctct
 3901 ccttcacgcg cccgccgccc tacctgaggg cgccatccac gccggttgag tcgcgttctg

3961 ccgcctcccg cctgtggtgc ctctgaact gcgtccgcgc tctaggttag tttaaagctc
 4021 aggtcgagac cgggcctttg tccggcgctc ccttgagacc tacctagact cagccggctc
 4081 tccacgcttt gcctgaccct gcttgctcaa ctctacgtct ttgtttcggt ttctgttctg
 4141 cgccgttaca gatccaagct gtgaccggcg cctacgctag atgaccgagt acaagcccac
 4201 ggtgcgcctc gccacccgcg acgacgtccc cagggccgta cgcacctcg ccgcgcgtt
 4261 cgccgactac cccgccacgc gccacaccgt cgatccggac cgccacatcg agcgggtcac
 4321 cgagctgcaa gaactcttcc tcacgcgcgt cgggtcgcac atcggcaagg tgtgggtcgc
 4381 ggacgacggc gccgcggtgg cgggtctggac cagccggag agcgtcgaag cgggggcggt
 4441 gttcgccgag atcggcccgc gcatggccga gttgagcggg tcccggctgg ccgcgcagca
 4501 acagatggaa ggctcctgg cgccgcaccg gcccaaggag cccgcgtggg tcttggccac
 4561 cgtcggcgct tcgcccgaac accagggcaa ggggtctggg agcgcgctcg tgctccccgg
 4621 agtggaggcg gccgagcgcg ccggggtgcc cgcttctctg gagacctccg cgccccgcaa
 4681 cctccccctt taegagcggc tgggttcac cgtcacccgc gacgtcgagg tgcccgaagg
 4741 accgcgcacc tgggtgatga cccgcaagcc cgggtgcctga gtcgacaatc aacctctgga
 4801 ttacaaaatt tgtgaaagat tgactgggat tcttaactat gttgctcctt ttacgctatg
 4861 tggatacgct gctttaatgc ctttgtatca tgctattgct tcccgatagg ctttcaattt
 4921 ctctccttg tataaatcct gggtgctgtc tctttatgag gagggtggc ccgttgctag
 4981 gcaacgtggc gtgggtgtgca ctgtgtttgc tgacgcaacc cccactgggt ggggcattgc
 5041 caccacctgt cagctccttt ccgggacttt cgctttcccc ctccctattg ccacggcgga
 5101 actcatcgcc gcctgccttg ccgctgctg gacagggggt cggctggttg gcactgacaa
 5161 ttccgtgggt ttgtcgggga aatcatcgct ctttcttggt ctgctcgct gtgttgccac
 5221 ctggattctg cgcgggacgt ccttctgcta cgtcccttcg gccctcaatc cagcggacct
 5281 tccttccgcg ggctgctgc cggtctgctg gctcttccg cgtcttcgcc ttcgccctca
 5341 gacgagtcgg atctcccttt gggccgcctc ccgctggt acctttaaga ccaatgactt
 5401 acaaggcagc tgtagatctt agccactttt taaaagaaaa ggggggactg gaagggctaa
 5461 ttcactccca acgaaaataa gatctgcttt ttgcttgtag tgggtctctc tgggttagacc
 5521 agatctgagc ctgggagctc tctggctaac tagggaaccc actgcttaag cctcaataaa
 5581 gcttgccctg agtgcttcaa gtagtgtgtg ccgctctggt gtgtgactct ggtaactaga
 5641 gatccctcag acccttttag tcagtgtgga aaatctctag cagtagtagt tcatgtcatc
 5701 ttattattca gtatttataa cttgcaaaga aatgaatata agagagttag aggaacttgt
 5761 ttattgcagc ttataatggt tacaaataaa gcaatagcat cacaaatttc acaataaag
 5821 catttttttc actgcattct agttgtgggt tgtccaaact catcaatgta tcttatcatg
 5881 tctggctcta gctatccgc ccctaactcc gccagttcc gccattctc cgccccatgg
 5941 ctgactaatt ttttttattt atgcagagge cgaggccgc tcggcctctg agctattcca
 6001 gaagtagtga ggaggctttt ttggaggcct agacttttgc agagacggcc caaattcgta
 6061 atcatggtca tagctgtttc ctgtgtgaaa ttgttatccg ctcaaatc cacacaacat
 6121 acgagccgga agcataaagt gtaaagcctg ggggtgcctaa tgagttagct aactcacatt
 6181 aattgcgttg cgtcactgc ccgctttcca gtcgggaaac ctgtcgtgcc agctgcatta
 6241 atgaatcggc caacgcgcgg ggagaggcgg tttgcgtatt gggcgtctt ccgcttctc
 6301 gctcactgac tcgctgcgct cggctgttcg gctgcggcga gcggtatcag ctactcaaa
 6361 ggcggtaata cggttatcca cagaatcagg ggataacgca ggaaagaaca tgtgagcaaa
 6421 aggccagcaa aaggccagga accgtaaaaa ggccgcgttg ctggcgttt tccataggct
 6481 ccgccccctc gacgagcatc acaaaaatcg acgtcaagt cagaggtggc gaaacccgac
 6541 aggactataa agataccagg cgtttcccc tggaagctcc ctgctgcgct ctctgttcc
 6601 gacctgccc cttaccggat acctgtccgc ctttctccct tcgggaagcg tggcgcttcc
 6661 tcatagctca cgtgttaggt atctcagttc ggtgtagggt gttcgtcca agctgggctg
 6721 tgtgcacgaa cccccgttc agcccgaccg ctgcgcctta tccggttaact atcgtcttga
 6781 gtccaacccg gtaagacacg acttatcgcc actggcagca gccactggta acaggattag
 6841 cagagcgagg tatgtaggcg gtgctacaga gttcttgaag tgggtggccta actacggcta
 6901 cactagaagg acagtatttg gtatctgcgc tctgctgaag ccagttacct tcggaaaaag
 6961 agttggtagc tcttgatccg gcaaacaaac caccgctggg agcgggtgggt tttttgtttg
 7021 caagcagcag attacgcgca gaaaaaaagg atctcaagaa gatcctttga tcttttctac
 7081 ggggtctgac gctcagtggg acgaaaactc acgttaaggg attttgggtca tgagattatc
 7141 aaaaaggatc ttacactaga tcctttttaa ttaaaaatga agtttttaaat caatctaaag
 7201 tatatatgag taaacttggt ctgacagtta ccaatgctta atcagttagg cacctatctc
 7261 agcgatctgt ctatttcgtt catccatagt tgctgactc ccgctcgtgt agataactac
 7321 gatacgggag ggcttaccat ctggccccag tgctgcaatg ataccgcgag acccacgctc
 7381 accggtcca gatttatcag caataaacca gccagccgga agggccgagc gcagaagtgg
 7441 tcctgcaact ttatccgcct ccattccagtc tattaattgt tgccgggaag ctagagtagg

7501 tagttcgcca gttaatatgtt tgcgcaacgt tgttgccatt gctacaggca tctgtggtgtc
 7561 acgctcgctcg tttgggtatgg cttcattcag ctccgggtcc caacgatcaa ggcgagttac
 7621 atgatccccc atgttgtgca aaaaagcggg tagctccttc ggtcctccga tctgtgtcag
 7681 aagtaagttg gccgcagtgt tatcactcat gggtatggca gcactgcata attctcttac
 7741 tgtcatgccca tccgtaagat gcttttctgt gactgggtgag tactcaacca agtcattctg
 7801 agaatagtgt atgcggcgac cgagttgctc ttgcccggcg tcaatacggg ataataccgc
 7861 gccacatagc agaactttaa aagtgtctcat cattggaaaa cgttcttcgg ggcgaaaact
 7921 ctcaaggatc ttaccgctgt tgagatccag ttcgatgtaa cccactcgtg cacccaactg
 7981 atcttcagca tcttttactt tcaccagcgt ttctgggtga gcaaaaacag gaaggcaaaa
 8041 tgccgcaaaa aagggaataa gggcgacacg gaaatgttga atactcatac tcttcctttt
 8101 tcaatattat tgaagcattt atcagggtta ttgtctcatg agcggatata tatttgaaatg
 8161 tattttagaaa aataaacaata taggggttcc gcgcacattt ccccgaaaag tgccacctga
 8221 cgtctaagaa accattatta tcatgacatt aacctataaa aataggcgta tcacgaggcc
 8281 ctttcgtctc gcgcgtttcg gtgatgacgg tgaaaacctc tgacacatgc agctcccgga
 8341 gacggtcaca gcttgtctgt aagcggatgc cgggagcaga caagcccgtc agggcgctc
 8401 agcgggtgtt ggcgggtgtc ggggctggct taactatgcg gcatcagagc agattgtact
 8461 gagagtgcac catatgcggg gtgaaatacc gcacagatgc gtaaggagaa aataccgcat
 8521 caggcgccat tcgccattca ggctgcgcaa ctgttgggaa gggcgatcgg tgcgggcctc
 8581 ttcgctatta cgccagctgg cgaaaggggg atgtgctgca aggcgattaa gttgggtaac
 8641 gccagggttt tcccagtcac gacgttgtaa aacgacggcc agtgccaagc tg.

p510_antiCD19_SL_TCRbeta (SEQ ID NO: 21)

1 acgcgtgtag tcttatgcaa tactcttgta gtcttgcaac atggtaacga tgagttagca
 61 acatgcctta caaggagaga aaaagcaccg tgcatgccga ttggtggaag taaggtggta
 121 cgatcggtgcc ttattaggaa ggcaacagac gggctctgaca tggattggac gaaccactga
 181 attgccgcat tgcagagata ttgtatttaa gtgcctagct cgatacaata aacgggtctc
 241 tctgggttaga ccagatctga gcctgggagc tctctggcta actagggaac ccactgctta
 301 agcctcaata aagcttgctt tgagtgtctc aagtagtgtg tgcccgtctg ttgtgtgact
 361 ctggttaacta gagatccctc agaccctttt agtcagtgtg gaaaatctct agcagtggcg
 421 cccgaacagg gacctgaaag cgaaagggaa accagagctc tctcgacgca ggactcggct
 481 tgctgaagcg cgcacggcaa gaggcgaggg gcggcgactg gtgagtacgc caaaaatttt
 541 gactagcggg ggctagaagg agagagatgg gtgcgagagc gtcagtatta agcgggggag
 601 aattagatcg cgatgggaaa aaattcgggt aaggccaggg ggaaagaaaa aatataaatt
 661 aaaacatata gtatgggcaa gcaggagact agaacgattc gcagttaatc ctggcctgtt
 721 agaaacatca gaaggctgta gacaaatact gggacagcta caaccatccc ttcagacagg
 781 atcagaagaa cttagatcat tatataatac agtagcaacc ctctattgtg tgcataaag
 841 gatagagata aaagacacca aggaagcttt agacaagata gaggaagagc aaaacaaaag
 901 taagaccacc gcacagcaag cggccactga tcttcagacc tggaggagga gatatgaggg
 961 acaattggag aagtgaatta tataaatata aagtagtaaa aattgaacca ttaggagtag
 1021 caccaccaa ggcaaaagaga agagtgggtg agagagaaaa aagagcagtg ggaataggag
 1081 ctttgttctt tgggttcttg ggagcagcag gaagcactat gggcgagcc tcaatgacgc
 1141 tgacggtaca ggccagacaa ttattgtctg gtatagtgca gcagcagaac aatttgcctga
 1201 gggctattga ggcgcaacag catctgttgc aactcacagt ctggggcatc aagcagctcc
 1261 aggcaagaat cctggctgtg gaaagatacc taaaggatca acagctcctg gggatttggg
 1321 gttgctctgg aaaactcatt tgcaccactg ctgtgccttg gaatgctagt tggagtaata
 1381 aatctctgga acagattgga atcacacgac ctggatggag tgggacagag aaattaacaa
 1441 ttacacaagc ttaatacact ccttaattga agaatcgcaa aaccagcaag aaaagaatga
 1501 acaagaatta ttggaattag ataaatgggc aagtttgtgg aattgggtta acataacaaa
 1561 ttggctgtgg tatataaaat tattcataat gatagtagga ggcttggtag gtttaagaat
 1621 agtttttgcg gtacttttcta tagtgaatag agttaggcag ggatattcac cattatcggt
 1681 tcagaccacac ctcccaaccc cgaggggacc cgacaggccc gaaggaatag aagaagaagg
 1741 tggagagaga gacagagaca gatccattcg attagtgaac ggatctcgac ggtatcggtt
 1801 aactttttaa agaaaagggg ggattggggg gtacagtgca ggggaaagaa tagtagacat
 1861 aatagcaaca gacatacaaa cttaaagaatt acaaaaacaa attacaaaat tcaaaatttt
 1921 atcgatacta gtattatgcc cagtacatga ccttatggga ctttctact tggcagtaca
 1981 tctacgtatt agtcacgct attaccatgg tgatgcgggt ttggcagtac atcaatgggc
 2041 gtggatagcg gtttgactca cggggatttc caagtctcca cccattgac gtcaatggga
 2101 gtttgttttg gcaccaaatt caacgggact ttccaaaatg tcgtaacaac tccgccccat
 2161 tgacgcaaat gggcggtagg cgtgtacggg gggaggttta tataagcaga gctcgtttag
 2221 tgaaccgtca gatcgcttgg agacgccatc cacgtgtttt tgacctccat agaagattct

2281 agagccgccca ccatgcttct cctggtgaca agccttctgc tctgtgagtt accacaccca
 2341 gcattcctcc tgatcccaga catccagatg acacagacta catcctccct gtctgcctct
 2401 ctgggagaca gagtcccat cagttgcagg gcaagtcagg acattagtaa atatttaa
 2461 tgggtatcagc agaaaccaga tggaaactgtt aaactcctga tctaccatac atcaagatta
 2521 cactcaggag tcccatcaag gttcagtgge agtgggtctg gaacagatta ttctctcacc
 2581 attagcaacc tggagcaaga agatattgcc acttactttt gccaacaggg taatacgtct
 2641 ccgtacacgt tcggaggggg gactaagttg gaaataacag gctccacctc tggatccggc
 2701 aagcccggat ctggcgaggg atccaccaag ggcgaggtga aactgcagga gtcaggacct
 2761 ggcctggtgg cgccctcaca gagcctgtcc gtcacatgca ctgtctcagg ggtctcatta
 2821 cccgactatg gtgtaagctg gattcgccag cctccacgaa agggctctgga gtggctggga
 2881 gtaatatggg gtagtgaaac cacatactat aattcagctc tcaaattccag actgaccatc
 2941 atcaaggaca actccaagag ccaagttttc ttaaaaatga acagtctgca aactgatgac
 3001 acagccattt actactgtgc caaacattat tactacggtg gtagctatgc tatggactac
 3061 tgggggtcaag gaacctcagt caccgtctcc tcagcggccg cagggtggcg cggttctggt
 3121 ggcggcggtt ctggtggcgg cggttctctc gagctgggag caggcccagt ggattctgga
 3181 gtcacacaaa ccccaaagca cctgatcaca gcaactggac agcgagtgc gctgagatgc
 3241 tcccctaggt ctggagacct ctctgtgtca tggtagcaac agagcctgga ccagggcctc
 3301 cagttcctca ttcagtatta taatggagaa gagagagcaa aaggaaacat tcttgaacga
 3361 ttctccgcac aacagttccc tgacttgca tctgaactaa acctgagctc tctggagctg
 3421 ggggactcag ctttgtattt ctgtgccagc agccccgga caggcctgaa cactgaagct
 3481 ttctttggac aaggcaccag actcacagtt gtagaggacc tgaacaaggt gtccccacc
 3541 gaggtcgctg tgtttgagcc atcagaagca gagatctccc acacccaaaa ggccacactg
 3601 gtgtgcctgg ccacaggctt cttccccgac cagtgaggc cgcagcccc tcaaggagca gcccgccctc
 3661 aaggaggtgc acagtggggt cagcacggac ctgaggggtc cgccacctt ctggcagaac
 3721 aatgactcca gatactgcct gagcagccgc tcaagtccag ttctacgggc tctcgagaa tgacgagtg
 3781 ccccgcaacc acttcgctg tcaagtccag atcgtcagcg ccgaggcctg gggtagagca
 3841 acccaggata gggccaaacc cgtcaccag atcgtcagcg caaggggtcc tgtctgccac catcctctat
 3901 gactgtggct ttacctcggt gtcctaccag gctgtgctgg tcagcgcct tgtgttgatg
 3961 gagatcctgc tagggaaagg caccctgtat gctgtgctgg tcagcgcct tgtgttgatg
 4021 gccatggtca agagaaagga tttctgataa gaattcgatc cgcggcccg aaggatctgc
 4081 gatcgctccg gtgcccgtca gtgggcagag cgcacatcgc ccacagtccc cgagaagttg
 4141 gggggagggg tcggcaattg aacgggtgcc tagagaaggt ggcgcggggt aaactgggaa
 4201 agtgatgtcg tgtactggct ccgccttttt cccgaggggt ggggagaacc gtatataagt
 4261 gcagtagtcg ccgtgaacgt tctttttcgc aacgggtttg ccgcagaaac acagctgaag
 4321 cttcgagggg ctgcacatct tcttcacgc gcccgccgc ctacctgagg ccgccatcca
 4381 cgccggttga gtcgcgttct gccgcctccc gctgtggtg cctcctgaac tgcgtccgcc
 4441 gtctaggtaa gtttaaagct caggtcgaga ccgggccttt gtccggcgct ccctgggagc
 4501 ctacctagac tcagccggct ctccacgctt tgctgacct tgcctgtcct actctacgtc
 4561 tttgtttcgt tttctgttct gcgcggttac agatccaagc tgtgaccggc gcctacgcta
 4621 gatgaccgag tacaagccca cgggtgcgct cgcacccgc gacgacgtcc ccagggccgt
 4681 acgcaccctc gccgcgcgt tcgccgacta cccgcacag cgccacaccg tcgatccgga
 4741 ccgccacatc gagcgggtca ccgagctgca agaactcttc ctacgcgcg tcgggctcga
 4801 catcggaag gtgtgggtcg cggacgacgg cgcgcgggtg gcggtctgga ccacgcggga
 4861 gagcgtcgaa gcggggcggt tgttcgccga gatcgcccg cgcattggcg agttgagcgg
 4921 ttcccggtg gccgcgcagc aacagatgga aggcctcctg gcgcgcacc ggcccaagga
 4981 gccgcggtg ttcttgcca ccgtcgcggt ctgcgccgac caccagggca agggctctggg
 5041 cagcgcgctc gtgctccccg gagtggaggc ggccgagcgc gccggggtgc ccgccttct
 5101 ggagacctcc gcgccccgca acctccctt ctacgagcgc ctcggttca ccgtcaccgc
 5161 cgacgtcgag gtgcccgaag gaccgcgcac ctggtgcatg acccgcaagc ccggtgctg
 5221 agtcgacaat caacctctgg attacaaaat ttgtgaaaga ttgactggta ttcttaacta
 5281 tgttgcctct tttacgtat gtggatacgc tgetttaatg cttttgtatc atgctattgc
 5341 ttcccgatg gctttcattt tctcctcctt gtataaatcc tggttgctgt ctctttatga
 5401 ggagttgtgg ccggtgttca ggcaacgtgg cgtggtgtgc actgtgtttg ctgacgcaac
 5461 ccccaactgg tggggcattg ccaccacctg tcagctcctt tccgggaact tcgctttccc
 5521 cctccctatt gccacggcgg aactcategc cgctgcctt gccgctgct ggacaggggc
 5581 tcggctgttg ggcactgaca attccgtggg gttgtcgggg aaatcatcgt cctttccttg
 5641 gctgctcgcc tgtgttgcca cctggattct gcgcgggacg tcttctgct acgtcccttc
 5701 ggccctcaat ccagcggacc ttccttcccg cggcctgctg ccggctctgc ggcccttcc
 5761 gcgtcttcgc cttcgccctc agacgagtc gatctcctt tgggcgcct ccccgccctg

5821 tacctttaag accaatgact tacaaggcag ctgtagatct tagccacttt ttaaaagaaa
 5881 aggggggact ggaagggcta attcactccc aacgaaaata agatctgctt tttgcttgta
 5941 ctgggtctct ctgggttagac cagatctgag cctgggagct ctctggctaa ctaggggaacc
 6001 cactgcttaa gcctcaataa agcttgccctt gagtgcctca agtagtgtgt gcccgctctgt
 6061 tgtgtgactc tggttaactag agatccctca gaccctttta gtcagtgtgg aaaatctcta
 6121 gcagtagtag ttcatgtcat cttattattc agtatttata acttgcaaaag aaatgaatat
 6181 cagagagtga gaggaacttg tttattgcag cttataatgg ttacaaataa agcaatagca
 6241 tcacaaattht cacaaataaa gcattttttt cactgcattc tagttgtggg ttgtccaaac
 6301 tcatcaatgt atcttatcat gtctggctct agctatcccg cccctaactc cgcccagttc
 6361 cgcccattht ccgccccatg gctgactaat tttttttatt tatgcagagg ccgaggccgc
 6421 ctcggcctct gagctattcc agaagtagtg aggaggcttt tttggaggcc tagacttttg
 6481 cagagacggc ccaaattcgt aatcatgggc atagctgttt cctgtgtgaa attgttatcc
 6541 gctcacaatt ccacacaaca tacgagccgg aagcataaag tgtaaagcct ggggtgccta
 6601 atgagtgagc taactcacat taattgcgtt gcgctcactg cccgctttcc agtcgggaaa
 6661 cctgtcgtgc cagctgcatt aatgaatcgg ccaacgcgcg gggagaggcg gtttgcgtat
 6721 tgggcgctct tccgcttctt cgtcactga ctcgctgcgc tcggctgttc ggctgcggcg
 6781 agcgggtatc gctcactcaa aggcggtaat acggttatcc acagaatcag gggataacgc
 6841 aggaaagaac atgtgagcaa aaggccagca aaaggccagg aaccgtaaaa aggcgcgctt
 6901 gctggcgctt tcccataggc tccgcccccc tgacgagcat cacaaaaatc gacgtcaag
 6961 tcagaggtgg cgaaaccgca caggactata aagataccag gcgtttcccc ctggaagctc
 7021 cctcgtgcgc tctcctgttc cgacctgcc gcttaccgga tacctgtccg cttttctccc
 7081 ttcgggaagc gtggcgcttt ctcatagctc acgctgtagg tatctcagtt cgggtgtaggt
 7141 cgttcgctcc aagctgggct gtgtgcacga acccccctt cagcccgacc gctgcgcctt
 7201 atccggtaac tatcgtcttg agtccaaccc ggtaagacac gacttatcgc cactggcagc
 7261 agccactggg aacaggatta gcagagcgag gtatgtaggc ggtgctacag agttcttgaa
 7321 gtgggtggct aactacggct acactagaag gacagtattt ggtatctgcg ctctgctgaa
 7381 gccagttacc ttcggaaaaa gaggttggtag ctcttgatcc ggcaaacaaa ccaccgctgg
 7441 tagcgggtgg ttttttgttt gcaagcagca gattacgcgc agaaaaaaag gatctcaaga
 7501 agatcctttg atcttttcta cggggtctga cgtcagtggt aacgaaaact cacgttaagg
 7561 gatttttggtc atgagattat caaaaaggat cttcacctag atccttttaa attaaaaatg
 7621 aagtttttaa tcaatctaaa gtatatatga gtaaacttgg tctgacagtt accaatgctt
 7681 aatcagtgag gcacctatct cagcgatctg tctatttcgt tcatccatag ttgcctgact
 7741 ccccgctcgt tagataacta cgatacggga gggcttacca tctggcccca gtgctgcaat
 7801 gataccgcga gaccacgct caccggctcc agatttatca gcaataaacc agccagccgg
 7861 aagggccgag cgcagaagtg gtcctgcaac ttatccgcc tccatccagt ctattaattg
 7921 ttgcgggaa gctagagtaa gtagttcgcc agttaatagt ttgcgcaacg ttgttgccat
 7981 tgctacaggc atcgtggtgt cagcctcgtc gtttggtatg gcttcattca gctccggttc
 8041 ccaacgatca aggcgagtta catgatcccc catgttgtgc aaaaaagcgg ttagctcctt
 8101 cggctcctcc atcgttgtca gaagtaagtt ggccgcagtg ttatcactca tgggtatggc
 8161 agcactgcat aattctctta ctgtcatgcc atccgtaaga tgcttttctg tgactggtga
 8221 gtactcaacc aagtcattct gagaatagt tatgcccga ccgagttgct cttgcccgcc
 8281 gtcaatacgg gataataccg cgccacatag cagaacttta aaagtgtca tcattggaaa
 8341 acgttcttcg gggcgaaaac tctcaaggat cttaccgctg ttgagatcca gttcgatgta
 8401 acccactcgt gcacccaact gatcttcagc atcttttact ttcaccagcg tttctgggtg
 8461 agcaaaaaca ggaaggcaaa atgcccgaaa aaagggaata agggcgacac ggaaatgttg
 8521 aatactcata ctcttccttt ttcaatatta ttgaagcatt tatcagggtt attgtctcat
 8581 gagcggatac atatttgaat gtatttagaa aaataaacia ataggggttc cgcgcacatt
 8641 tccccgaaaa gtgccacctg acgtctaaga aaccattatt atcatgacat taacctataa
 8701 aaataggcgt atcacgaggc cctttcgtct cgcgcgtttc ggtgatgacg gtgaaaacct
 8761 ctgacacatg cagctcccgg agacggtcac agcttgtctg taagcggatg ccgggagcag
 8821 acaagcccggt cagggcgcgt cagcgggtgt tggcgggtgt cggggctggc ttaactatgc
 8881 ggcatcagag cagattgtac tgagagtgca ccatatgcgg tgtgaaatac cgcacagatg
 8941 cgtaaggaga aaataaccgca tcaggcgcca ttcgccattc aggtctgcga actgttggga
 9001 agggcgatcg gtgcgggcct cttegtatt acgccagctg gcgaaagggg gatgtgctgc
 9061 aaggcgatta agttgggtaa cgccagggtt ttcccagtea cgacgttgta aaacgacggc
 9121 cagtgccaaag ctg.

p510_antiBCMA_CD3epsilon (SEQ ID NO: 22)

1 acgcgtgtag tcttatgcaa tactcttgta gtcttgcaac atggtaacga tgagttagca
 61 acatgcctta caaggagaga aaaagcaccg tgcatgccga ttgggtggaag taagggtgta

```

121  cgatcgtgcc ttattaggaa ggcaacagac gggctctgaca tggattggac gaaccactga
181  attgccgcat tgcagagata ttgtatttaa gtgcctagct cgatacaata aacgggtctc
241  tctgggttaga ccagatctga gcctgggagc tctctggcta actagggAAC ccactgctta
301  agcctcaata aagcttgccct tgagtgtctc aagtagtgtg tgcccgtctg ttgtgtgact
361  ctggtaacta gagatccctc agaccctttt agtcagtgtg gaaaatctct agcagtggcg
421  cccgaacagg gacctgaaag cgaaagggaa accagagctc tctcgacgca ggactcggct
481  tgctgaagcg cgcacggcaa gaggcgaggg gcggcgactg gtgagtacgc caaaaatttt
541  gactagcgga ggctagaagg agagagatgg gtgcgagagc gtcagtatta agcgggggag
601  aattagatcg cgatgggaaa aaattcgggt aaggccaggg ggaaagaaaa aatataaatt
661  aaaacatata gtatgggcaa gcaggagctc agaacgattc gcagttaatc ctggcctgtt
721  agaaacatca gaaggctgta gacaaatact gggacagcta caaccatccc ttcagacagg
781  atcagaagaa cttagatcat tatataatac agtagcaacc ctctattgtg tgcatacaag
841  gatagagata aaagacacca aggaagcttt agacaagata gaggaagagc aaaacaaaag
901  taagaccacc gcacagcaag cggccactga tcttcagacc tggaggagga gatatgaggg
961  acaattggag aagtgaatta tataaatata aagtagtaaa aattgaacca ttaggagtag
1021 caccaccaa ggcaaagaga agagtgggtg agagagaaaa aagagcagtg ggaataggag
1081 ctttgttctt tgggttcttg ggagcagcag gaagcactat gggcgagcc tcaatgacgc
1141 tgacggtaca ggccagacaa ttattgtctg gtatagtgca gcagcagaac aatttgctga
1201 gggctattga ggcgcaacag catctgttgc aactcacagt ctggggcatc aagcagctcc
1261 aggcaagaat cctggctgtg gaaagatacc taaaggatca acagctcctg gggatttggg
1321 gttgctctgg aaaactcatt tgcaccactg ctgtgccttg gaatgctagt tggagtaata
1381 aatctctgga acagattgga atcacacgac ctggatggag tgggacagag aaattaacaa
1441 ttacacaagc ttaatacact ccttaattga agaatcgcaa aaccagcaag aaaagaatga
1501 acaagaatta ttggaattag ataaatgggc aagtttgttg aattggttta acataacaaa
1561 ttggctgtgg tatataaaat tattcataat gatagtagga ggcttggtag gtttaagaat
1621 agtttttgc gtactttcta tagtgaatag agttaggcag ggatattcac cattatcggt
1681 tcagaccac cteccaaccc cgaggggacc cgacaggccc gaaggaatag aagaagaagg
1741 tggagagaga gacagagaca gatccattcg attagtgaac ggatctcgac ggtatcgggt
1801 aactttttaa agaaaagggg ggattggggg gtacagtgca ggggaaagaa tagtagacat
1861 aatagcaaca gacatacaaa ctaaagaatt acaaaaacaa attacaaaat tcaaaatttt
1921 atcgatacta gtattatgcc cagtacatga ccttatggga ctttcctact tggcagtaca
1981 tctacgtatt agtcacgctc attaccatgg tgatgcgggt ttggcagtag atcaatgggc
2041 gtggatagcg gtttgactca cggggatttc caagtctcca cccattgac gtcaatggga
2101 gtttgttttg gcaccaaatt caacgggact ttccaaaatg tcgtaacaac tccgccccat
2161 tgacgcaaat gggcggtagg cgtgtacggt gggaggttta tataagcaga gctcgtttag
2221 tgaaccgtca gatcgctgg agacgccatc cacgctgttt tgacctccat agaagattct
2281 agagccgcca ccatgcttct cctggtgaca agccttctgc tctgtgagtt accacacca
2341 gcattcctcc tgatccaca ggtgcagctg gtgcagagcg gcgcggaagt gaaaaaacg
2401 ggcgcgagcg tgaaagtga ctgcaaagcg agcggctata gctttccgga ttattatatt
2461 aactgggtgc gccaggcgcc gggccagggc ctggaatgga tgggctggat ttattttgcg
2521 agcggcaaca gcgaatataa ccagaaattt accggccgcg tgacctgac ccgcgatacc
2581 agcagcagca ccgcgtatat ggaactgagc agcctgcgca gcgaagatac cgcggtgat
2641 ttttgcgcg gctgtatga ttatgattgg tattttgatg tgtggggcca gggcaccatg
2701 gtgaccgtga gcagcggcgg cggcggcagc ggcgcgcgcg gcagcggcgg cggcggcagc
2761 gatattgtga tgaccagac ccgcgtgagc ctgagcgtga ccccgggcga accggcgagc
2821 attagctgca aaagcagcca gagcctggtg catagcaacg gcaacaccta tctgcattgg
2881 tatctgcaga aaccgggcca gagcccgag ctgctgattt ataaagtga caaccgcttt
2941 agcggcgtgc cggatcgctt tagcggcagc ggcagcgcg cggattttac cctgaaaatt
3001 agccgcgtgg aagcgggaaga tgtgggctg tattattgcg cggaaccag ccatgtgccg
3061 tggacctttg gccagggcac caaactggaa attaaaagcg gtggcgcgcg ttctgggtggc
3121 ggcggttctg gtggcgcgcg ttctctcgag gatggtaatg aagaaatggg tggattaca
3181 cagacaccat ataaagtctc catctctgga accacagtaa tattgacatg ccctcagtat
3241 cctggatctg aaatactatg gcaacacaaat gataaaaaca taggcggtga tgaggatgat
3301 aaaaacatag gcagtgatga ggatcacctg tcaactgaagg aattttcaga attggagcaa
3361 agtggttatt atgtctgcta ccccagagga agcaaaccag aagatgcgaa cttttatctc
3421 tacctgaggg caagagtgtg tgagaactgc atggagatgg atgtgatgtc ggtggccaca
3481 attgtcatag tggacatctg catcactggg ggcttgcctg tgctggttta ctactggagc
3541 aagaatagaa aggccaaggc caagcctgtg acacgaggag cgggtgctgg cggcaggcaa
3601 aggggacaaa acaaggagag gccaccacct gttcccaacc cagactatga gcccatccgg

```

```

3661 aaaggccagc gggacctgta ttctggcctg aatcagagac gcatctgata agaattcggg
3721 tccgcggccg cgaaggatct gcgatcgctc cggtgcccgt cagtgggcag agcgcacatc
3781 gccacagtc cccgagaagt tggggggagg ggtcggcaat tgaacgggtg cctagagaag
3841 gtggcgcggg gtaaactggg aaagtgatgt cgtgtactgg ctccgccttt tcccagaggg
3901 tgggggagaa ccgtatataa gtgcagtagt cgccgtgaac gttctttttc gcaacggggt
3961 tgccgccaga acacagctga agcttcgagg ggctcgcata tctccttcac gcgcccgcg
4021 ccctacctga ggccgccatc cacgccgggt gagtcgcgtt ctgccgcctc ccgcctgtgg
4081 tgccctctga actgcgtccg ccgtctaggt aagtttaag ctcaggtcga gaccgggcct
4141 ttgtccggcg ctcccttgga gcctacctag actcagccgg ctctccacgc tttgcctgac
4201 cctgcttgct caactctacg tctttgtttc gttttctgtt ctgcgccgtt acagatccaa
4261 gctgtgaccg gcgcctacgc tagatgaccg agtacaagcc cacggtgcgc ctgcgccacc
4321 gcgacgacgt ccccagggcc gtacgcaccc tcgccgcgcg gttcgcgcgac taccgcccca
4381 cgcgccacac cgtcgatccg gaccgccaca tcgagcgggt caccgagctg caagaactct
4441 tcctcacgcg cgtcgggctc gacatcgga aggtgtgggt cgcggacgac ggcgcgcg
4501 tggcgggtct gaccacgccg gagagcgtcg aagcgggggc ggtgttcgcc gagatcggcc
4561 cgcgcatggc cgagttgagc ggttcccggc tggccgcgca gcaacagatg gaaggcctcc
4621 tggcgccgca ccggcccaag gagcccgctt ggttctggc caccgtcggc gtctcgcccg
4681 accaccaggg caagggtctg ggcagcgccg tcgtgctccc cggagtggag gcggccgagc
4741 gcgcgggggt gcccgccttc ctggagacct ccgcgccccg caacctcccc ttctacgagc
4801 ggctcggtt caccgtcacc gccgacgtcg aggtgcccgaggaccgcgc acctgggtgca
4861 tgaccgcaa gcccgtgccc tgagtcgaca atcaacctct ggattacaaa atttgtgaaa
4921 gattgactgg tattcttaac tatgttgctc cttttacgct atgtggatac gctgctttaa
4981 tgcccttgta tcatgctatt gcttcccgtg tggctttcat tttctctccc ttgtataaat
5041 cctgggtgct gtctctttat gaggagtgtt ggcccggtgt caggcaacgt ggcgtgggtg
5101 gcactgtggt tgetgacgca acccccactg gttggggcat tgccaccacc tgtcagctcc
5161 tttccgggac tttcgctttc cccctcccta ttgccacggc ggaactcacc gccgcctgcc
5221 ttgcccgctg ctggacaggg gctcggtgtt tgggactga caattccgtg gtgttgtcgg
5281 ggaaatcacc gtcctttcct tggctgctcg cctgtgttgc cacctggatt ctgcgcggga
5341 cgtccttctg ctacgtccct tcggccctca atccagcgga cttccttccc cgcggcctgc
5401 tgccggctct gcggcctctt ccgctcttcc gcttcgccc tcagacgagt cggatctccc
5461 tttggggcgc ctccccgcct ggtaccttta agaccaatga cttacaaggc agctgtagat
5521 cttagccact ttttaaaaga aaagggggga ctggaagggc taattcactc ccaacgaaaa
5581 taagatctgc tttttgcttg tactgggtct ctctgggttag accagatctg agcctgggag
5641 ctctctggct aactagggaa cccactgctt aagcctcaat aaagcttgcc ttgagtgtt
5701 caagtagtgt gtgcccgtct gttgtgtgac tctggtaact agagatccct cagaccctt
5761 tagtcagtgt ggaaaatctc tagcagtagt agttcatgtc atcttattat tcagtattta
5821 taacttgcaa agaaatgaat atcagagagt gagaggaaact tgtttattgc agcttataat
5881 ggttacaaat aaagcaatag catcacaat ttcacaaata aagcattttt ttcactgcat
5941 tctagttgtg gtttgtccaa actcatcaat gtatcttata atgtctggct ctagctatcc
6001 cgcccctaac tcgcccagt tcgcccatt ctccgcccc cagctgacta atttttttta
6061 tttatgcaga ggccgaggcc gcctcggcct ctgagctatt ccagaagtag tgaggaggct
6121 tttttggagg cctagacttt tgcagagacg gcccaaattc gtaatcatgg tcatagctgt
6181 ttctgtgtg aaattgttat ccgctcaciaa ttccacaciaa catacagacc ggaagcataa
6241 agtgtaaagc ctggggtgcc taatgagtga gctaactcac attaatgctg ttgcgctcac
6301 tgcccgcttt ccagtcggga aacctgtcgt gccagctgca ttaatgaatc ggccaacgcg
6361 cggggagagg cggtttgctt attgggcgct ctccgcttc ctgcctcact gactcgctgc
6421 gctcggtcgt tcggctgcgg cgagcgggat cagctcactc aaaggcggtg atacggttat
6481 ccacagaatc aggggataac gcaggaaaga acatgtgagc aaaaggccag caaaaggcca
6541 ggaaccgtaa aaaggccgcg ttgctggcgt ttttccatag gctccgcccc cctgacgagc
6601 atcacaaaaa tcgacgctca agtcagaggt ggcgaaaccc gacaggacta taaagatacc
6661 aggcgtttcc ccctggaagc tccctcgtgc gctctcctgt tccgaccctg ccgcttaccg
6721 gatacctgtc cgcttttctc ccttcgggaa gcgtggcgct ttctcatagc tcacgctgta
6781 ggtatctcag ttcgggtgtg gtcgttcgct ccaagctggg ctgtgtgcac gaaccccccg
6841 ttcagcccca ccgctgcgcc ttatccggta actatcgtct tgagtcacac ccggtaaagc
6901 acgacttata gccactggca gcagccactg gtaacaggat tagcagagcg aggtatgtag
6961 gcggtgctac agagtctctg aagtgggtgg ctaactacgg ctacactaga aggacagtat
7021 ttggtatctg cgtctgtctg aagccagtta ccttcggaaa aagagttggg agctcttgat
7081 ccggcaaaaa aaccaccgct ggtagcgggt gtttttttgt ttgcaagcag cagattacgc
7141 gcagaaaaaa aggatctcaa gaagatcctt tgatcttttc tacggggtct gacgctcagt

```

```

7201 ggaacgaaaa ctcacgttaa gggatttttgg tcatgagatt atcaaaaagg atcttcacct
7261 agatcctttt aaattaaaaa tgaagtttta aatcaatcta aagtatatat gagtaaactt
7321 ggtctgacag ttaccaatgc ttaatcagtg aggcacctat ctcagegatc tgtctatttc
7381 gttcatccat agttgcctga ctccccgtcg tgtagataac tacgatacgg gagggccttac
7441 catctggccc cagtgtctga atgataccgc gagaccacag ctcaccggct ccagatttat
7501 cagcaataaa ccagccagcc ggaagggccg agcgcagaag tggtcctgca actttatccg
7561 cctccatcca gtctattaat tgttgccggg aagctagagt aagtagttcg ccagttaata
7621 gtttgcgcaa cgttggttgc attgtctacag gcatcgtggg gtcacgctcg tcgtttggta
7681 tggcttcatt cagctccggg tcccaacgat caaggcgagt tacatgatcc cccatgttgt
7741 gcaaaaaagc ggtagctcc ttcggctctc cgatcgttgt cagaagtaag ttggccgcag
7801 tgttatcact catggttatg gcagcactgc ataattctct tactgtcatg ccatccgtaa
7861 gatgcttttc tgtgactggg gagtactcaa ccaagtcatt ctgagaatag tgtatgcggc
7921 gaccgagttg ctcttgcccg gcgtcaatac gggataatac cgcgccacat agcagaactt
7981 taaaagtgtc catcattgga aaacgttctt cggggcgaaa actctcaagg atcttaccgc
8041 tgttgagatc cagttcgatg taaccctact gtgcacccaa ctgatcttca gcatctttta
8101 ctttcaccag cgtttctggg tgagcaaaaa caggaaggca aaatgccgca aaaaagggaa
8161 taagggcgac acggaaatgt tgaatactca tactcttctt ttttcaatat tattgaagca
8221 tttatcaggg ttattgtctc atgagcggat acatatttga atgtatttag aaaaataaac
8281 aaataggggt tccgcgcaca tttccccgaa aagtgccacc tgacgtctaa gaaaccatta
8341 ttatcatgac attaacctat aaaaataggc gtatcacgag gccctttcgt ctgcgcgctt
8401 tcggtgatga cggtgaaaac ctctgacaca tgcagctccc ggagacggtc acagcttgctc
8461 tgtaagcgga tgccgggagc agacaagccc gtcagggcgc gtcagegggt gttggcgggt
8521 gtcggggctg gcttaactat gcggcatcag agcagattgt actgagagtg caccatatgc
8581 ggtgtgaaat accgcacaga tgcgtaagga gaaaataccg catcaggcgc cattcgccat
8641 tcaggctgcg caactgttgg gaagggcgat cgggtgcgggc ctcttcgcta ttacgccagc
8701 tggcgaaaagg gggatgtgct gcaaggcgat taagttgggt aacgccaggg ttttcccagt
8761 cacgacgttg taaaacgacg gccagtgcc aagctg

```

***p510_antiBCMA_CD3gamma* (SEQ ID NO: 23)**

```

1 acgcgtgtag tcttatgcaa tactcttgta gtcttgcaac atggtaacga tgagttagca
61 acatgcctta caaggagaga aaaagcacccg tgcattgccga ttggtggaag taaggtggta
121 cgatcgtgcc ttattaggaa ggcaacagac gggctctgaca tggattggac gaaccactga
181 attgccgcat tgcagagata ttgtatttaa gtgcctagct cgatacaata aacgggtctc
241 tctggttaga ccagatctga gcctgggagc tctctggcta actagggaac ccactgctta
301 agcctcaata aagcttgcc ttagtgcttc aagtagtgtg tgcccgtctg ttgtgtgact
361 ctggtaacta gagatccctc agaccctttt agtcagtgtg gaaaatctct agcagtggcg
421 cccgaacagg gacctgaaag cgaaagggaa accagagctc tctcgacgca ggactcggct
481 tgctgaagcg cgcacggcaa gaggcgaggg gcggcgactg gtgagtacgc caaaaatttt
541 gactagcgga ggctagaagg agagagatgg gtgcgagagc gtcagtatta agcgggggag
601 aattagatcg cgatgggaaa aaattcgggt aaggccaggg ggaaagaaaa aatataaatt
661 aaaacatata gtatgggcaa gcaggagctc agaacgattc gcagttaatc ctggcctgtt
721 agaaacatca gaaggctgta gacaaatact gggacagcta caacctccc ttcagacagg
781 atcagaagaa cttagatcat tatataatac agtagcaacc ctctattgtg tgcataaag
841 gatagagata aaagacacca aggaagcttt agacaagata gaggaagagc aaaacaaaag
901 taagaccacc gcacagcaag cggccactga tcttcagacc tggaggagga gatatgaggg
961 acaattggag aagtgaatta tataaatata aagtagtaaa aattgaacca ttaggagtag
1021 caccaccaa ggcaagaga agagtgggtg agagagaaaa aagagcagtg ggaataggag
1081 ctttgttcct tgggttcttg ggagcagcag gaagcactat gggcgcagcc tcaatgacgc
1141 tgacggtaca ggccagacaa ttattgtctg gtatagtgca gcagcagaac aatttgctga
1201 gggctattga ggcgcaacag catctgttgc aactcacagt ctggggcatc aagcagctcc
1261 aggcaagaat cctggctgtg gaaagatacc taaaggatca acagctcctg gggatttggg
1321 gttgctctgg aaaactcatt tgcaccactg ctgtgccttg gaatgctagt tggagtaata
1381 aatctctgga acagattgga atcacacgac ctggatggag tgggacagag aaattaacaa
1441 ttacacaagc ttaatacact ccttaattga agaatcgcaa aaccagcaag aaaagaatga
1501 acaagaatta ttggaattag ataaatgggc aagtttgtgg aattggttta acataacaaa
1561 ttggctgtgg tatataaaat tattcataat gatagtagga ggcttggtag gtttaagaat
1621 agtttttgc gtactttcta tagtgaatag agttaggcag ggatattcac cattatcggt
1681 tcagaccacac ctcccacccc cgaggggacc cgacaggccc gaaggaatag aagaagaagg
1741 tggagagaga gacagagaca gatccattcg attagtgaac ggatctcgac ggtatcgggt
1801 aactttttaa agaaaagggg ggattggggg gtacagtgca ggggaaagaa tagtagacat

```

1861	aatagcaaca	gacatacaaa	ctaaagaatt	acaaaaacaa	attacaaaat	tcaaaatttt
1921	atcgatacta	gtattatgcc	cagtacatga	ccttatggga	ctttcctact	tggcagtaga
1981	tctacgtatt	agtcacgcgt	attaccatgg	tgatgcgggt	ttggcagtag	atcaatgggc
2041	gtggatagcg	gtttgactca	cggggatttc	caagtctcca	ccccattgac	gtcaatggga
2101	gtttgttttg	gcacccaaaat	caacgggact	ttccaaaatg	tcgtaacaac	tccgccccat
2161	tgacgcaaat	gggcggtagg	cgtgtacggg	gggaggttta	tataagcaga	gctcgttttag
2221	tgaaccgtca	gatcgcttgg	agacgccatc	cacgctgttt	tgacctccat	agaagattct
2281	agagccgcca	ccatgcttct	cctgggtgaca	agccttctgc	tctgtgagtt	accacaccca
2341	gcatttctcc	tgatcccaca	ggtgcagctg	gtgcagagcg	gcgcggaagt	gaaaaaacgg
2401	ggcgcgagcg	tgaaagttag	ctgcaaagcg	agcggctata	gctttccgga	ttattatatt
2461	aactgggtgc	gccaggcgcc	gggccagggc	ctggaatgga	tgggctggat	ttatttttgcg
2521	agcggcaaca	gcgaatataa	ccagaaattt	accggccgcg	tgaccatgac	ccgcgatacc
2581	agcagcagca	ccgcgtatat	ggaactgagc	agcctgcgca	gcgaagatac	cgcggtgtat
2641	ttttgcgcga	gcctgtatga	ttatgattgg	tattttgatg	tgtggggcca	gggcaccatg
2701	gtgaccgtga	gcagcggcgg	cggcggcagc	ggcggcggcg	gcagcggcgg	cggcggcagc
2761	gatattgtga	tgaccagac	cccgcctgagc	ctgagcgtga	ccccgggcca	accggcgagc
2821	attagctgca	aaagcagcca	gagcctgggtg	catagcaacg	gcaacaccta	tctgcattgg
2881	tatctgcaga	aaccgggcca	gagcccgcag	ctgctgattt	ataaagttag	caaccgcttt
2941	agcggcgtgc	cggatcgctt	tagcggcagc	ggcagcggcg	cggattttac	cctgaaaatt
3001	agccgcgtgg	aagcgggaaga	tgtgggcgtg	tattattgcg	cggaaaccag	ccatgtgccg
3061	tggacctttg	gccagggcac	caaactggaa	attaaaagcg	gtggcggcgg	ttctgggtggc
3121	ggcggttctg	gtggcggcgg	ttctctcgag	cagtcaatca	aaggaaacca	cttgggttaag
3181	gtgtatgact	atcaagaaga	tgggttcggta	cttctgactt	gtgatgcaga	agccaaaaat
3241	atcacatggt	ttaaagatgg	gaagatgate	ggcttcceta	ctgaagataa	aaaaaaatgg
3301	aatctgggaa	gtaatgccaa	ggaccacaga	gggatgtatc	agtgtaaagg	atcacagaac
3361	aagtcaaaac	cactccaagt	gtattacaga	atgtgtcaga	actgcattga	actaaatgca
3421	gccaccatat	ctggctttct	ctttgctgaa	atcgtcagca	ttttcgtcct	tgtgtttggg
3481	gtctacttca	ttgctggaca	ggatggagtt	cgcagtcga	gagcttcaga	caagcagact
3541	ctggttgccca	atgaccagct	ctaccagccc	ctcaaggatc	gagaagatga	ccagtacagc
3601	caccttcaag	gaaaccagtt	gaggaggaat	tgataagaat	tcggatccgc	ggccgcgaag
3661	gatctgcgat	cgctccgggtg	cccgtcagtg	ggcagagcgc	acatcgccca	cagtcgccga
3721	gaagttgggg	ggaggggtcg	gcaattgaac	gggtgcctag	agaaggtggc	gcggggtaaa
3781	ctgggaaagt	gatgtcgtgt	actggctccg	cctttttccc	gaggggtggg	gagaaccgta
3841	tataagtgcg	gtagtgcgg	tgaacgttct	ttttcgcaac	gggtttgccc	ccagaacaca
3901	gctgaagctt	cgaggggctc	gcatctctcc	ttcacgcgcc	cgcgcgccca	cctgaggccg
3961	ccatccacgc	cggttgagtc	gcgttctgcc	gcctcccgcc	tgtggtgcct	cctgaactgc
4021	gtccgcgcgc	taggtaagtt	taaagctcag	gtcgagaccg	ggcctttgtc	cggcgctccc
4081	ttggagccta	cctagactca	gccggctctc	cacgctttgc	ctgaccttgc	ttgctcaact
4141	ctacgtcttt	gtttcgtttt	ctgttctgcg	ccgttacaga	tccaagctgt	gaccggcgcc
4201	tacgctagat	gaccgagtag	aagcccacgg	tgcgcctcgc	caccgcgcgc	gacgtcccca
4261	gggcgcgtacg	caccctcgcc	gcgcggttcg	ccgactaccc	cgccacgcgc	cacaccgtcg
4321	atccggaccg	ccacatcgag	cgggtcacccg	agctgcaaga	actcttcctc	acgcgcgtcg
4381	ggctcgacat	cggcaagggtg	tgggtcgcgg	acgacggcgc	cgcgggtggcg	gtctggacca
4441	cgcgcgagag	cgtcgaaagcg	ggggcggtgt	tcgcgcgagat	cggcccgcg	atggccgagt
4501	tgagcgggttc	ccggctggcc	gcgcagcaac	agatggaagg	cctcctggcg	ccgcaccggc
4561	ccaaggagcc	cgcgtggttc	ctggccaccg	tcggcgtctc	gcccgaaccac	cagggcaagg
4621	gtctgggcag	cgcgcgtcgtg	ctccccggag	tggaggcggc	cgagcgcgc	gggggtgccc
4681	ccttctctgga	gacctccgcg	ccccgcaacc	tccccttcta	cgagcggctc	ggcttcaccg
4741	tcaccgcccga	cgtcgaggtg	cccgaaggac	cgcgcacctg	gtgcatgacc	cgcaagcccg
4801	gtgcctgagt	cgacaatcaa	cctctggatt	acaaaatttg	tgaaagattg	actggtattc
4861	ttaaactatgt	tgtctctttt	acgctatgtg	gatacgtcgc	tttaatgcct	ttgtatcatg
4921	ctattgcttc	ccgtatggct	ttcattttct	cctccttgta	taaatcctgg	ttgctgtctc
4981	tttatgagga	gttgtggccc	gttgtcaggg	aacgtggcgt	ggtgtgcact	gtgtttgctg
5041	acgcaacccc	cactggttgg	ggcattgccca	ccacctgtca	gctcctttcc	gggaactttcg
5101	ctttccccct	ccctattggc	acggcggaac	tcategcgcg	ctgccttgcc	cgtctgtgga
5161	caggggctcg	gctgttgggc	actgacaatt	ccgtgggtgt	gtcggggaaa	tcategctct
5221	ttccttggct	gctcgcctgt	gttgccacct	ggattctgcg	cgggaacgtcc	ttctgctacg
5281	tcccttcggc	cctcaatcca	gcggaccttc	cttcccgcgg	cctgctgcgc	gctctgcggc
5341	ctcttcgcgc	tcttcgcctt	cgccttcaga	cgagtcggat	ctcccttttg	gcgcctccc

5401 cgcctggtac ctttaagacc aatgacttac aaggcagctg tagatccttag ccacttttta
 5461 aaagaaaagg ggggactgga agggctaatt cactcccaac gaaaataaga tctgcttttt
 5521 gcttgtagtg ggtctctctg gtttagaccag atctgagcct gggagctctc tggctaacta
 5581 gggaacccac tgettaagcc tcaataaagc ttgccttgag tgcttcaagt agtgtgtgcc
 5641 cgtctgttgt gtgactctgg taactagaga tcctcagac ctttttagtc agtgtggaaa
 5701 atctctagca gtagtagttc atgtcatctt attattcagt atttataact tgcaaagaaa
 5761 tgaatatcag agagtgagag gaacttgttt attgcagctt ataatggtta caaataaagc
 5821 aatagcatca caaatttcac aaataaagca tttttttcac tgcattctag ttgtggtttg
 5881 tccaaactca tcaatgtatc ttatcatgtc tggctctagc tatcccgccc ctaactccgc
 5941 ccagttccgc ccattctccg ccccatggct gactaatttt ttttatttat gcagaggccg
 6001 aggccgcctc ggccctctgag ctattccaga agtagtgagg aggccttttt ggaggcctag
 6061 actttttgcag agacggccca aattcgtaat catggtcata gctgtttcct gtgtgaaatt
 6121 gttatccgct cacaattcca cacaacatac gagccggaag cataaagtgt aaagcctggg
 6181 gtgcctaata agtgagctaa ctacacattaa ttgcgttgcg ctactgccc gctttccagt
 6241 cgggaaacct gtcgtgccag ctgcattaat gaatcggcc aacgcgggg agaggcggtt
 6301 tgcgtattgg gcgctcttcc gcttctctgc tcaactgact gctgcgctcg gtcgttcggc
 6361 tgcggcgagc ggtatcagct cactcaaagg cggtaataac gttatccaca gaatcagggg
 6421 ataacgcagg aaagaacatg tgagcaaaag gccagcaaaa ggccagggaac cgtaaaaagg
 6481 ccgcggttgc ggcggttttt cataggctcc gccccctga cgagcatcac aaaaatcgac
 6541 gctcaagtca gaggtggcga aaccgcagac gactataaag ataccaggcg tttccccctg
 6601 gaagctccct cgtgcgctct cctgttccga cctgcgct taccggatac ctgtccgctt
 6661 ttctcccttc ggggaagcgtg gcgctttctc atagctcacg ctgtagggtat ctgagttcgg
 6721 tgtagggtcg tcgctccaag ctgggctgtg tgcacgaacc ccccgttcag cccgaccgct
 6781 gcgccttate cggtaactat cgtcttgagt ccaaccgggt aagacacgac ttatcgccac
 6841 tggcagcagc cactggtaac aggattagca gagcgaggta tgtaggcggt gctacagagt
 6901 tcttgaagtg gtggcctaac tacggctaca ctagaaggac agtatttggt atctgcgctc
 6961 tgctgaagcc agttaccttc ggaaaaagag ttggtagctc ttgatccggc aaacaaacca
 7021 ccgctggtag cgggtggtttt tttgtttgca agcagcagat tacgcgcaga aaaaaaggat
 7081 ctcaagaaga tcctttgatc ttttctacgg ggtctgacgc tcagtggaaac gaaaactcac
 7141 gttaagggat tttgggtcatg agattatcaa aaaggatctt cacctagatc cttttaaatt
 7201 aaaaatgaag ttttaaatac atctaaagta tatatgagta aacttggtct gacagttacc
 7261 aatgcttaat cagtgaggca cctatctcag cgatctgtct atttcggtca tccatagttg
 7321 cctgactccc cgtcgtgtag ataactacga tacgggaggg cttaccatct ggccccagtg
 7381 ctgcaatgat accgcgagac ccacgctcac cggctccaga tttatcagca ataaaccagc
 7441 cagccggaag ggccgagcgc agaagtggtc ctgcaacttt atccgcctcc atccagtcta
 7501 ttaattgttg ccgggaagct agagtaagta gttcgccagt taatagtttg cgcaacggtg
 7561 ttgccattgc tacaggcatc gtggtgtcac gctcgtcgtt tggtaggct tcattcagct
 7621 ccggttccca acgatcaagg cgagttacat gatccccat gttgtgcaaa aaagcggtta
 7681 gtccttccg tcctccgac gttgtcagaa gtaagttggc cgcagtgtta tcaactatgg
 7741 ttatggcagc actgcataat tctcttactg tcatgccatc cgtaagatgc ttttctgtga
 7801 ctgggtgagta ctcaaccaag tcattctgag aatagtgtat gcggcgaccg agttgctctt
 7861 gccggcgctc aatacgggat aataccgcgc cacatagcag aactttaaaa gtgctcatca
 7921 ttggaaaacg ttcttcgggg cgaaaactct caaggatctt accgctgttg agatccagtt
 7981 cgatgtaacc cactcgtgca cccaactgat cttcagcatc ttttactttc accagcgttt
 8041 ctgggtgagc aaaaacagga aggcaaaatg ccgcaaaaaa gggaataagg gcgacacgga
 8101 aatggtgaat actcatactc ttcttttttc aatattattg aagcatttat cagggttatt
 8161 gtctcatgag cggatacata tttgaatgta tttagaaaaa taaacaaata ggggttccgc
 8221 gcacatttcc ccgaaaagtg ccacctgacg tctaagaaac cattattatc atgacattaa
 8281 cctataaaaa taggcgtatc acgaggccct ttcgtctcgc gcgtttcggg gatgacgggtg
 8341 aaaacctctg acacatgcag ctcccgagga cggtcacagc ttgtctgtaa gcggatgccg
 8401 ggagcagaca agcccgctcag ggcgcgtcag cgggtgttgg cgggtgtcgg ggctggctta
 8461 actatgcggc atcagagcag attgtactga gagtgcacca tatgcgggtg gaaataaccgc
 8521 acagatgcgt aaggagaaaa taccgcatca ggcccatc gccattcagg ctgcgcaact
 8581 gttgggaagg gcgatcgggt cgggcctctt cgctattacg ccagctggcg aaagggggat
 8641 gtgctgcaag gcgattaagt tgggtaacgc cagggttttc ccagtcacga cgttgtaaaa
 8701 cgacggccag tgccaagctg.

Example 2: Antibody Sequences***Generation of Antibody Sequences***

[0342] The human CD19 polypeptide canonical sequence is UniProt Accession No. P15391 (or P15391-1). The human BCMA polypeptide canonical sequence is UniProt Accession No. Q02223 (or Q02223-1). Provided are antibody polypeptides that are capable of specifically binding to the human CD19 polypeptide or human BCMA polypeptide or human FAP polypeptide or human BCMA polypeptide, and fragments or domains thereof. Anti-CD19, anti-FAP, anti-CAIX and anti-BCMA antibodies can be generated using diverse technologies (see, e.g., (Nicholson et al, 1997). Where murine anti-CD19, anti-FAP, anti-CAIX or anti-BCMA antibodies are used as a starting material, humanization of murine anti-CD19, anti-FAP, anti-CAIX or anti-BCMA antibodies is desired for the clinical setting, where the mouse-specific residues may induce a human-anti-mouse antigen (HAMA) response in subjects who receive T-cell receptor (TCR) fusion protein (TFP) treatment, i.e., treatment with T-cells transduced with the TFP.CD19, TFP.FAP, TFP.CAIX, or TFP.BCMA construct. Humanization is accomplished by grafting CDR regions from murine anti-CD19, anti-FAP, anti-CAIX or anti-BCMA antibody onto appropriate human germline acceptor frameworks, optionally including other modifications to CDR and/or framework regions. As provided herein, antibody and antibody fragment residue numbering follows Kabat (Kabat E. A. et al, 1991; Chothia et al, 1987).

Generation of scFvs

[0343] Human or humanized anti-CD19, anti-FAP, anti-CAIX or anti-BCMA IgGs are used to generate scFv sequences for TFP constructs. DNA sequences coding for human or humanized V_L and V_H domains are obtained, and the codons for the constructs are, optionally, optimized for expression in cells from Homo sapiens. The order in which the V_L and V_H domains appear in the scFv is varied (i.e., V_L-V_H, or V_H-V_L orientation), and three copies of the “G₄S” or “G₄S” subunit (G₄S)₃ connect the variable domains to create the scFv domain. Anti-CD19, anti-FAP, anti-CAIX and anti-BCMA scFv plasmid constructs can have optional Flag, His or other affinity tags, and are electroporated into HEK293 or other suitable human or mammalian cell lines and purified. Validation assays include binding analysis by FACS, kinetic analysis using Proteon, and staining of CD19-expressing cells.

[0344] Exemplary anti-CD19 or anti-BCMA CDRs of V_L and V_H domains and the nucleotide sequences encoding them, respectively, are shown below:

Anti-CD19**[0345] Anti-CD19 light chain CDR1**

Coding Sequence: AGGGCAAGTCAGGACATTAGTAAA (SEQ ID NO: 24).

Amino acid sequence: RASQDISK (SEQ ID NO: 25).

[0346] Anti-CD19 light chain CDR2

Coding Sequence: ATCTACCATACATCAAGATTA (SEQ ID NO: 26).

Amino acid sequence: IYHTSRL (SEQ ID NO: 27).

[0347] Anti-CD19 light chain CDR3

Coding Sequence: CAACAGGGTAATACGCTTCCGTACACG (SEQ ID NO: 28).

Amino acid sequence: QQGNTLPYT (SEQ ID NO: 29).

[0348] Anti-CD19 heavy chain CDR1

Coding Sequence: GGGGTCTCATTACCCGACTATGGTGTAAGC (SEQ ID NO: 30).

Amino acid sequence: GVSLPDYGV (SEQ ID NO: 31).

[0349] Anti-CD19 heavy chain CDR2

Coding Sequence: GTAATATGGGGTAGTGAAACCACATACTATAATTCAGCTCTC (SEQ ID NO: 32).

Amino acid sequence: VIWGSETTYNSAL (SEQ ID NO: 33).

[0350] Anti-CD19 heavy chain CDR3

Coding Sequence: CATTATTACTACGGTGGTAGCTATGCTATGGACTAC (SEQ ID NO: 34).

Amino acid sequence: HYYYGGSYAMDY (SEQ ID NO: 35).

Anti-BCMA**[0351] Anti-BCMA light chain CDR1**

Coding Sequence:

AAAAGCAGCCAGAGCCTGGTGCATAGCAACGGCAACACCTATCTGCAT (SEQ ID NO: 36).

Amino acid sequence: KSSQSLVHSNGNTYLH (SEQ ID NO: 37).

[0352] Anti-BCMA light chain CDR2

Coding Sequence: AAAGTGAGCAACCGCTTTAGC (SEQ ID NO: 38).

Amino acid sequence: KVSNRFS (SEQ ID NO: 39).

[0353] Anti-BCMA light chain CDR3

Coding Sequence: GCGGAAACCAGCCATGTGCCGTGGACC (SEQ ID NO: 40)

Amino acid sequence: AETSHVPWT (SEQ ID NO: 41).

[0354] Anti-BCMA heavy chain CDR1

Coding Sequence: AAAGCGAGCGGCTATAGCTTTCCGGATTATTATATTAAC (SEQ ID NO: 42).

Amino acid sequence: KASGYSFPDYYIN (SEQ ID NO: 43).

[0355] Anti-BCMA heavy chain CDR2

Coding Sequence:

TGGATTTATTTTGCAGCGGCAACAGCGAATATAACCAGAAATTTACCGGC (SEQ ID NO: 44).

Amino acid sequence: WIYFASGNSEYNQKFTG (SEQ ID NO: 45).

[0356] Anti-BCMA heavy chain CDR3

Coding Sequence: CTGTATGATTATGATTGGTATTTTGATGTG (SEQ ID NO: 46).

Amino acid sequence: LYDYDWYFDV (SEQ ID NO: 47).

[0357] Anti-CD19 light chain variable region*Coding Sequence:*

GACATCCAGATGACACAGACTACATCCTCCCTGTCTGCCTCTCTGGGAGACAGAGTC
 ACCATCAGTTGCAGGGCAAGTCAGGACATTAGTAAATATTTAAATTGGTATCAGCA
 GAAACCAGATGGAAGTGTAACTCCTGATCTACCATACATCAAGATTACACTCAG
 GAGTCCCATCAAGGTTCAAGTGGCAGTGGGTCTGGAACAGATTATTCTCTCACCATTA
 GCAACCTGGAGCAAGAAGATATTGCCACTTACTTTTGCCAACAGGGTAATACGCTT
 CCGTACACGTTTCGGAGGGGGGACTAAGTTGGAAATAACA (SEQ ID NO: 48).

Amino acid sequence:

DIQMTQTTSSLSASLGDRVTISCRASQDISKYLNWYQQKPDGTVKLLIYHTSRLHSGVPS
 RFSGSGSGTDYSLTISNLEQEDIATYFCQQGNTLPYTFGGGKLEIT (SEQ ID NO: 49).

[0358] Anti-CD19 heavy chain variable region*Coding Sequence:*

GAGGTGAAACTGCAGGAGTCAGGACCTGGCCTGGTGGCGCCCTCACAGAGCCTGTC
 CGTCACATGCACTGTCTCAGGGGTCTCATTACCCGACTATGGTGTAAGCTGGATTGCG
 CCAGCCTCCACGAAAGGGTCTGGAGTGGCTGGGAGTAATATGGGGTAGTGAAACCA
 CATACTATAATTCAGCTCTCAAATCCAGACTGACCATCATCAAGGACAACTCCAAG
 AGCCAAGTTTTCTTAAAAATGAACAGTCTGCAAAGTATGACACAGCCATTTACTAC
 TGTGCCAAACATTATTACTACGGTGGTAGCTATGCTATGGACTACTGGGGTCAAGG
 AACCTCAGTCACCGTCTCCTCA (SEQ ID NO: 50).

Amino acid sequence:

EVKLQESGPGLVAPSQSLSVTCTVSGVSLPDYGVSWIRQPPRKGLEWLGVWGSETTY
 NSALKSRLTIKDNSKSQVFLKMNSLQTDDTAIYYCAKHYYYGGSYAMDYWGQGT
 SVTVSS (SEQ ID NO: 51).

[0359] Anti-BCMA light chain variable region*Coding Sequence:*

GATATTGTGATGACCCAGACCCCGCTGAGCCTGAGCGTGACCCCGGGCGAACCGGC
 GAGCATTAGCTGCAAAAGCAGCCAGAGCCTGGTGCATAGCAACGGCAACACCTATC

TGCATTGGTATCTGCAGAAACCGGGCCAGAGCCCGCAGCTGCTGATTATAAAGTG
AGCAACCGCTTTAGCGGCGTGCCGGATCGCTTTAGCGGCAGCGGCAGCGGCGCGGA
TTTTACCCTGAAAATTAGCCGCGTGGAAGCGGAAGATGTGGGCGTGTATTATTGCG
CGGAAACCAGCCATGTGCCGTGGACCTTTGGCCAGGGCACCAAACCTGGAAATTAAGC
(SEQ ID NO: 52).

Amino acid sequence:

DIVMTQTPLSLSVTPGEPASISCKSSQSLVHSNGNTYLHWYLQKPGQSPQLLIYKVSNR
SGVPDRFSGSGSGADFTLKISRVEAEDVGVYYCAETSHVPWTFGQGTKLEIKS(SEQ ID
NO: 53).

[0360] Anti-BCMA heavy chain variable region

Coding Sequence:

CAGGTGCAGCTGGTGCAGAGCGGCGCGGAAGTGAAAAACCGGGCGCGAGCGTG
AAGTGAGCTGCAAAGCGAGCGGCTATAGCTTCCGGATTATTATATTAAGTGGGTG
CGCCAGGCGCCGGGCGAGGGCCTGGAATGGATGGGCTGGATTTATTTTGCAGCGG
CAACAGCGAATATAACCAGAAATTTACCGGCCGCGTGACCATGACCCGCGATACCA
GCAGCAGCACCGCGTATATGGAAGTGAAGCAGCCTGCGCAGCGAAGATACCGCGGTG
TATTTTTCGCGAGCCTGTATGATTATGATTGGTATTTTGATGTGTGGGGCCAGGGC
ACCATGGTGACCGTGAGCAGC (SEQ ID NO: 54).

Amino acid sequence:

QVQLVQSGAEVKKPGASVKVSKASGYSFPDYYINWVRQAPGQGLEWMGWIYFASGNSE
YNQKFTGRVTMTRDTSSSTAYMELSSLRSEDTAVYFCASLYDYDWYFDVWGQGTMTVTS
S (SEQ ID NO: 55).

Source of TCR Subunits

[0361] Subunits of the human T Cell Receptor (TCR) complex all contain an extracellular domain, a transmembrane domain, and an intracellular domain. A human TCR complex contains the CD3-epsilon polypeptide, the CD3-gamma polypeptide, the CD3-delta polypeptide, the CD3-zeta polypeptide, the TCR alpha chain polypeptide and the TCR beta chain polypeptide. The human CD3-epsilon polypeptide canonical sequence is Uniprot Accession No. P07766. The human CD3-gamma polypeptide canonical sequence is Uniprot Accession No. P09693. The human CD3-delta polypeptide canonical sequence is Uniprot Accession No. P043234. The human CD3-zeta polypeptide canonical sequence is Uniprot Accession No. P20963. The human TCR alpha chain canonical sequence is Uniprot Accession No. Q6ISU1. The human TCR beta chain C region canonical sequence is Uniprot Accession No. P01850, a human TCR beta chain V region sequence is P04435.

[0362] The human CD3-epsilon polypeptide canonical sequence is:

MQSGTHWRVLGLCLLSVGWVGQDGNEEMGGITQTPYKVSISGTTVILTCPQYPGSEIL
WQHNDKNIGGDEDDKNIGSDEDHLSLKEFSELEQSGYYVCYPRGSKPEDANFYLYLRA
RVCENCMEMDVMSVATIVIVDICITGGLLLL VYYWSKNRKAKAKPVTRGAGAGGRQR
GQNKERPPPVPNPDIYEPIRKGQRDLYSGLNQRRRI (SEQ ID NO: 56).

[0363] The human CD3-gamma polypeptide canonical sequence is:

MEQGKGLAVLILAILLQGTLAQSIKGNHLVKVYDYQEDGSVLLTCDAEAKNITWFKD
GKMIGFLTEDKKKWNLGSNAKDPRGMYQCKGSQNKSKPLQVYYRMCQNCIELNAATI
SGFLFAEIVSIFVLAVGVYFIAGQDGVQRQSRASDKQTLLPNDQLYQPLKDREDDQYSHL
QGNQLRRN (SEQ ID NO: 57).

[0364] The human CD3-delta polypeptide canonical sequence is:

MEHSTFLSGLVLATLLSQVSPFKIPIEELEDVRFVNCNTSITWVEGTVGTLLSDITRLDLG
KRILDPRGIYRCNGTDIYKDKESTVQVHYRMCQSCVELDPATVAGIIVTDVIATLLLALG
VFCFAGHETGRLSGAADTQALLRNDQVYQPLRDRDDAQYSHLGGNWARNK (SEQ ID
NO: 58).

[0365] The human CD3-zeta polypeptide canonical sequence is:

MKWKALFTAAILQAQLPITEAQSFGLLDPKLCYLLDGILFIYGVILTALFLRVKFSRSAD
APAYQQGQNQLYNELNLGRREEYDVLDKRRGRDPGEMGGKPKRRKNPQEGLYNELQK
DKMAEAYSEIGMKGERRRGKGHDGLYQGLSTATKDTYDALHMQUALPPR (SEQ ID NO:
59).

[0366] The human TCR alpha chain canonical sequence is:

MAGTWLLLLLALGCPALPTGVGGTPFPSLAPPIMLLVDGKQQMVVVCLVLDVAPPGLD
SPIWFSAGNGSALDAFTYGPSPATDGTWTNLAHLSLPSEELASWEPLVCHTGPAGEGHS
RSTQPMHLSGEASTARTCPQEPLRGTPGGALWLGVLRLLLFKLLLFDLLLTCSCLCDPA
GPLPSPATTTLRALGSHRLHPATETGGREATSSPRPQPRDRRWGDTTPGRKPGSPVWG
EGSYLSSYPTCPAQAWCSRSALRAPSSSLGAFFAGDLPPPLQAGAA (SEQ ID NO: 60).

[0367] The human TCR alpha chain C region canonical sequence is:

PNIQNPDPAVYQLRDSKSSDKSVCLFTDFDSQTNVSQSKSDSVYITDKTVLDMRSMDFK
SNSAVAWSNKSDFACANAFNNIIPEDTFFPSPESSCDVKLVEKSFETDTNLFQNLSVIG
FRILLLKVAGFNLLMTLRLWSS (SEQ ID NO: 61).

[0368] The human TCR alpha chain V region CTL-L17 canonical sequence is:

MAMLLGASVLILWLQPDWVNSQQKNDDQQVKQNSPSLSVQEGRISILNCDYTNSMFD
YFLWYKKYPAGEPTFLISSIKDKNEDGRFTVFLNKS AKHLSLHIVPSQPGDSAVYFCA
AKGAGTASKLTFGTGTRLQVTL (SEQ ID NO: 62).

[0369] The human TCR beta chain C region canonical sequence is:

EDLNKVFPEVAVFEPSEAEISHTQKATLVCLATGFFPDHVELSWWVNGKEVHSGVSTD
PQPLKEQPALNDSRYCLSSRLRVSAFTWQNPRNHFRCQVQFYGLSENDEWTQDRAKPV
TQIVSAEAWGRADCGFTSVSYQQGVLSATILYEILLGKATLYAVLVSAVLMLMAMVKRK
DF (SEQ ID NO: 63).

[0370] The human TCR beta chain V region CTL-L17 canonical sequence is:

MGTSLLCWMALCLLGADHADTGVSQNPRHNITKRGQNVTFRCDPISSEHNRLYWYRQT
LGQGPEFLTYFQNEAQLEKSRLLSDRFSAERPKGSFSTLEIQRTEQGDSAMLYLCASSLAG
LNQPQHFGDGTRLISIL (SEQ ID NO: 64).

[0371] The human TCR beta chain V region YT35 canonical sequence is:

MDSWTFCCVSLCILVAKHTDAGVIQSPRHEVTEMGQEVTLRCKPISGHNSLFWYRQTM
MRGLELLIYFNNNPIDDSGMPEDRFSKMPNASFSTLKIQPSEPRDSAVYFCASSFSTCS
ANYGYTFGSGTRLTVV (SEQ ID NO: 65).

[0372] An exemplary anti-BCMA heavy chain sequence is:

Generation of TFPs from TCR Domains and scFvs

[0373] The CD19 or BCMA scFvs are recombinantly linked to CD3-epsilon or other TCR subunits (see 1C) using a linker sequence, such as G₄S, (G₄S)₂ (G₄S)₃ or (G₄S)₄. Various linkers and scFv configurations are utilized. TCR alpha and TCR beta chains were used for generation of TFPs either as full length polypeptides or only their constant domains. Any variable sequence of TCR alpha and TCR beta chains is allowed for making TFPs.

TFP Expression Vectors

[0374] Expression vectors are provided that include: a promoter (Cytomegalovirus (CMV) enhancer-promoter), a signal sequence to enable secretion, a polyadenylation signal and transcription terminator (Bovine Growth Hormone (BGH) gene), an element allowing episomal replication and replication in prokaryotes (e.g., SV40 origin and ColE1 or others known in the art) and elements to allow selection (ampicillin resistance gene and zeocin marker).

[0375] Preferably, the TFP-encoding nucleic acid construct is cloned into a lentiviral expression vector and expression validated based on the quantity and quality of the effector T-cell response of TFP.CD19-transduced T-cells ("CD19.TFP" or "CD19.TFP T-cells" or "TFP.CD19" or "TFP.CD19 T-cells") in response to CD19+ target cells, TFP.FAP-transduced T-cells ("FAP.TFP" or "FAP.TFP T-cells" or "TFP.FAP" or "TFP.FAP T-cells") in response to FAP+ target cells, TFP.CAIX-transduced T-cells ("CAIX.TFP" or "CAIX.TFP T-cells" or "TFP.CAIX" or "TFP.CAIX T-cells") in response to CAIX+ target cells, or TFP.BCMA-transduced T-cells ("BCMA.TFP" or "BCMA.TFP T-cells" or "TFP.BCMA" or "TFP.BCMA T-cells") in response to BCMA + target cells. Effector T-cell responses include, but are not

limited to, cellular expansion, proliferation, doubling, cytokine production and target cell lysis or cytolytic activity (i.e., degranulation).

[0376] The TFP.CD19, TFP.FAP, TFP.CAIX or TFP.BCMA lentiviral transfer vectors are used to produce the genomic material packaged into the VSVg pseudotyped lentiviral particles. Lentiviral transfer vector DNA is mixed with the three packaging components of VSVg, gag/pol and rev in combination with Lipofectamine reagent to transfect them together into 293 cells. After 24 and 48 hours, the media is collected, filtered and concentrated by ultracentrifugation. The resulting viral preparation is stored at -80C. The number of transducing units is determined by titration on SupT1 cells. Redirected TFP.CD19, TFP.FAP, TFP.CAIX or TFP.BCMA T-cells are produced by activating fresh naive T-cells with anti-CD3x anti-CD28 beads for 24 hrs and then adding the appropriate number of transducing units to obtain the desired percentage of transduced T-cells. These modified T-cells are allowed to expand until they become rested and come down in size at which point they are cryopreserved for later analysis. The cell numbers and sizes are measured using a coulter multisizer III. Before cryopreserving, percentage of cells transduced (expressing the TFP.CD19, TFP.FAP, TFP.CAIX or TFP.BCMA on the cell surface) and their relative fluorescence intensity of that expression are determined by flow cytometric analysis. From the histogram plots, the relative expression levels of the TFPs are examined by comparing percentage transduced with their relative fluorescent intensity.

[0377] In some embodiments multiple TFPs are introduced by T-cell transduction with multiple viral vectors.

Evaluating Cytolytic Activity, Proliferation Capabilities and Cytokine Secretion of Humanized TFP Redirected T Cells

[0378] The functional abilities of TFP.CD19, TFP.FAP, TFP.CAIX or TFP.BCMA T-cells to produce cell-surface expressed TFPs, and to kill target tumor cells, proliferate and secrete cytokines are determined using assays known in the art.

[0379] Human PBMCs (e.g., blood from a normal apheresed donor whose naive T-cells are obtained by negative selection for T-cells, CD4+ and CD8+ lymphocytes) are treated with human interleukin-2 (IL-2) then activated with anti-CD3x anti-CD28 beads, e.g., in 10% RPMI at 37 °C, 5% CO₂ prior to transduction with the TFP-encoding lentiviral vectors. Flow cytometry assays are utilized to confirm cell surface presence of a TFP, such as by an anti-FLAG antibody or an anti-murine variable domain antibody. Cytokine (e.g., IFN- γ) production is measured using ELISA or other assays.

Example 3: Human TFP T-cell Efficacy in a Human ALL Mouse Model

[0380] Primary human ALL cells can be grown in immune compromised mice (e.g., NSG or NOD) without having to culture them *in vitro*. Likewise, cultured human ALL cell lines can

induce leukemia in such mice. ALL-bearing mice can be used to test the efficacy of human TFP.CD19, TFP.FAP, TFP.CAIX or TFP.BCMA T-cells, for instance, in the model HALLX5447. The readout in this model is the survival of mice after intravenous (i.v.) infusion of ALL cells in the absence and presence of i.v. administered human TFP.CD19, TFP.FAP, TFP.CAIX or TFP.BCMA T-cells.

Example 4: Human TFP T-cell Treatment in an *In Vivo* Solid Tumor Xenograft Mouse Model

[0381] The efficacy of human TFP.CD19 or TFP.BCMA T-cells can also be tested in immune compromised mouse models bearing subcutaneous solid tumors derived from human CD19- or BCMA-expressing ALL, CLL or NHL human cell lines. Tumor shrinkage in response to human TFP.CD19, TFP.FAP, TFP.CAIX or TFP.BCMA T-cell treatment can be either assessed by caliper measurement of tumor size, or by following the intensity of a GFP fluorescence signal emitted by GFP-expressing tumor cells.

[0382] Primary human solid tumor cells can be grown in immune compromised mice without having to culture them *in vitro*. Exemplary solid cancer cells include solid tumor cell lines, such as provided in The Cancer Genome Atlas (TCGA) and/or the Broad Cancer Cell Line Encyclopedia (CCLE, see Barretina et al., Nature 483:603 (2012)). Exemplary solid cancer cells include primary tumor cells isolated from renal cell carcinoma, breast cancer, lung cancer, ovarian cancer, prostate cancer, colon cancer, cervical cancer, brain cancer, liver cancer, pancreatic cancer, kidney or stomach cancer. These mice can be used to test the efficacy of TFP.CD19, TFP.FAP, TFP.CAIX or TFP.BCMA T-cells in the human tumor xenograft models (see, e.g., Morton et al., Nat. Protoc. 2:247 (2007)). Following an implant or injection of 1×10^6 - 1×10^7 primary cells (collagenase-treated bulk tumor suspensions in EC matrix material) or tumor fragments (primary tumor fragments in EC matrix material) subcutaneously, tumors are allowed to grow to 200-500 mm³ prior to initiation of treatment.

Example 5: Demonstration of Multiplexed TFP polypeptides, and use of multiplexed humanized TFP Redirected T Cells

[0383] The TFP polypeptides provided herein are capable of functionally associating with endogenous TCR subunit polypeptides to form functional TCR complexes. Here, multiple TFPS in lentiviral vectors are used to transduce T-cells in order to create a functional, multiplexed recombinant TCR complex. For example, provided is T-cell containing i) a first TFP having an extracellular domain, a transmembrane domain, and an intracellular domain from the CD3-delta polypeptide and an CD19-, FAP-, CAIX-, or BCMA-specific scFv antibody fragment, and ii) a second TFP having an extracellular domain, a transmembrane domain, and an intracellular domain from the CD3-gamma polypeptide and a CD19-, FAP-, CAIX-, or BCMA-specific

antibody fragment. The first TFP and second TFP are capable of interacting with each other and with endogenous TCR subunit polypeptides, thereby forming a functional TCR complex.

[0384] The use of these multiplexed humanized TFP.CD19, TFP.FAP, TFP.CAIX or TFP.BCMA T-cells is demonstrated in liquid and solid tumors as provided in Examples 2 and 3 above.

Example 6: Preparation of T-cells transduced with TFPs

Lentiviral production

[0385] Lentivirus encoding the appropriate constructs were prepared as follows. 5×10^6 HEK293FT-cells were seeded into a 100 mm dish and allowed to reach 70-90% confluency overnight. 2.5 μ g of the indicated DNA plasmids and 20 μ L Lentivirus Packaging Mix (ALSTEM, cat# VP100; see Appendix B3) were diluted in 0.5 mL DMEM or Opti-MEM I Medium without serum and mixed gently. In a separate tube, 30 μ L of NanoFect transfection reagent (ALSTEM, cat.no. NF100) was diluted in 0.5 mL DMEM or Opti-MEM I Medium without serum and mixed gently. The NanoFect/DMEM and DNA/DMEM solutions were then mixed together and vortexed for 10-15 seconds prior to incubation of the DMEM-plasmid-NanoFect mixture at room temperature for 15 minutes. The complete transfection complex from the previous step was added dropwise to the plate of cells and rocked to disperse the transfection complex evenly in the plate. The plate was then incubated overnight at 37 °C in a humidified 5% CO₂ incubator. The following day, the supernatant was replaced with 10 mL fresh media and supplemented with 20 μ L of ViralBoost (500x, ALSTEM, cat.no. VB100). The plates were then incubated at 37 °C for an additional 24 hours. The lentivirus containing supernatant was then collected into a 50 mL sterile, capped conical centrifuge tube and put on ice. After centrifugation at 3000 rpm for 15 minutes at 4 °C, the cleared supernatant was filtered with a low-protein binding 0.45 μ m sterile filter and virus was subsequently isolated by ultracentrifugation at 25,000 rpm (Beckmann, L8-70M) for 1.5 hours, at 4 °C. The pellet was removed and re-suspended in DMEM media and Lentivirus concentrations/titers were established by quantitative RT-PCR, using the Lenti-X qRT-PCR Titration kit (Clontech; catalog number 631235). Any residual plasmid DNA was removed by treatment with DNaseI. The virus stock preparation was either used for infection immediately or aliquoted and stored at -80 °C for future use.

[0386] Similar experiments can be carried out with FAP.TFP and CAIX.TFP constructs.

PBMC isolation

[0387] Peripheral Blood Mononuclear Cells (PBMCs) were prepared from either whole blood or buffy coat. Whole blood was collected in 10 mL Heparin vacutainers and either processed immediately or stored overnight at 4 °C. Approximately 10 mL of whole anti-coagulated blood was mixed with sterile phosphate buffered saline (PBS) buffer for a total volume of 20 mL in a

50 mL conical centrifuge tube (PBS, pH 7.4, without $\text{Ca}^{2+}/\text{Mg}^{2+}$). 20 mL of this blood/PBS mixture was then gently overlaid onto the surface of 15 mL of Ficoll-Paque PLUS (GE Healthcare, 17-1440-03) prior to centrifugation at 400g for 30-40 min at room temperature with no brake application.

[0388] Buffy coat was purchased from Research Blood Components (Boston, MA). Leucosep tubes (Greiner bio-one) were prepared by adding 15 mL Ficoll-Paque (GE Health Care) and centrifuged at 1000g for 1 minute. Buffy coat was diluted 1:3 in PBS (pH 7.4, without Ca^{2+} or Mg^{2+}). The diluted buffy coat was transferred to Leucosep tube and centrifuged at 1000g for 15 minutes with no brake application. The layer of cells containing peripheral blood mononuclear cells (PBMC), seen at the diluted plasma/Ficoll interface, was removed carefully to minimize contamination by Ficoll. Residual Ficoll, platelets, and plasma proteins were then removed by washing the PBMCs three times with 40 mL of PBS by centrifugation at 200g for 10 minutes at room temperature. The cells were then counted with a hemocytometer. The washed PBMC were washed once with CAR-T media (AIM V-AlbuMAX (BSA) (Life Technologies), with 5% AB serum and 1.25 $\mu\text{g/mL}$ amphotericin B (Gemini Bioproducts, Woodland, CA), 100 U/mL penicillin, and 100 $\mu\text{g/mL}$ streptomycin). Alternatively, the washed PBMC's were transferred to insulated vials and frozen at -80°C for 24 hours before storing in liquid nitrogen for later use.

[0389] Similar experiments can be carried out with FAP.TFP and CAIX.TFP constructs.

T-cell activation

[0390] Peripheral Blood Mononuclear Cells (PBMCs) prepared from either whole blood or buffy coat were stimulated with anti-human CD28 and CD3 antibody-conjugated magnetic beads for 24 hours prior to viral transduction. Freshly isolated PBMC were washed once in CAR-T media (AIM V-AlbuMAX(BSA)(Life Technologies), with 5% AB serum and 1.25 $\mu\text{g/mL}$ amphotericin B (Gemini Bioproducts), 100 U/mL penicillin, and 100 $\mu\text{g/mL}$ streptomycin) without huIL-2, before being re-suspended at a final concentration of 1×10^6 cells/mL in CAR-T medium with 300 IU/mL human IL-2 (from a 1000x stock; Invitrogen). If the PBMCs had previously been frozen they were thawed and re-suspended at 1×10^7 cells/mL in 9 mL of pre-warmed (37°C) cDMEM media (Life Technologies), in the presence of 10% FBS, 100 U/mL penicillin, and 100 $\mu\text{g/mL}$ streptomycin, at a concentration of 1×10^6 cells/mL prior to washing once in CART medium, re-suspension at 1×10^6 cells/mL in CAR-T medium, and addition of IL-2 as described above.

[0391] Prior to activation, anti-human CD28 and CD3 antibody-conjugated magnetic beads (Invitrogen) were washed three times with 1 mL of sterile 1x PBS (pH7.4), using a magnetic rack to isolate beads from the solution, before re-suspension in CAR-T medium, with 300 IU/mL human IL-2, to a final concentration of 4×10^7 beads/mL. PBMC and beads were then

mixed at a 1:1 bead-to-cell ratio, by transferring 25 μL (1×10^6 beads) of beads to 1 mL of PBMC. The desired number of aliquots were then dispensed to single wells of a 12-well low-attachment, or non-treated cell culture plate, and incubated at 37 °C, with 5% CO_2 , for 24 hours before viral transduction.

[0392] Similar experiments can be carried out with FAP.TFP and CAIX.TFP constructs.

T-cell transduction/transfection and expansion

[0393] Following activation of PBMC cells were incubated for 24 hours at 37 °C, 5% CO_2 . Lentivirus was thawed on ice and 5×10^6 lentivirus, along with 2 μL of Transplus (Alstem) per mL of media (a final dilution of 1:500) was added to each well of 1×10^6 cells. Cells were incubated for an additional 24 hours before repeating addition of virus. Alternatively, lentivirus was thawed on ice and the respective virus was added at 5 or 50 MOI in presence of 5 $\mu\text{g/mL}$ Polybrene (Sigma). Cells were spinoculated at 100g for 100 minutes at room temperature. Cells were then grown in the continued presence of 300 IU/mL of human IL-2 for a period of 6-14 days (total incubation time is dependent on the final number of CAR-T-cells required). Cell concentrations were analyzed every 2-3 days, with media being added at that time to maintain the cell suspension at 1×10^6 cells/mL.

[0394] In some instances, activated PBMCs were electroporated with *in vitro* transcribed (IVT) mRNA (**FIG. 14**). Human PBMCs were stimulated with Dyna beads (ThermoFisher) at 1-to-1 ratio for 3 days in the presence of 300 IU/ml recombinant human IL-2 (R&D System). The beads were removed before electroporation. The cells were washed and re-suspended in OPTI-MEM medium (ThermoFisher) at the concentration of 2.5×10^7 cells/ mL. 200 μL of the cell suspension (5×10^6 cells) were transferred to the 2 mm gap Electroporation Cuvettes Plus™ (Harvard Apparatus BTX) and prechilled on ice. 10 μg of IVT TFP mRNA was added to the cell suspension. The mRNA/cell mixture was then electroporated at 200 V for 20 milliseconds using ECM830 Electro Square Wave Porator (Harvard Apparatus BTX). Immediately after the electroporation, the cells were transferred to fresh cell culture medium (AIM V AlbuMAX (BSA) serum free medium + 5% human AB serum + 300 IU/ml IL-2) and incubated at 37 °C.

[0395] Similar experiments can be carried out with FAP.TFP and CAIX.TFP constructs.

Verification of TFP expression by cell staining

[0396] Following lentiviral transduction or mRNA electroporation, expression of anti-CD19, anti-FAP, anti-CAIX and anti-BCMA CARs and TFPs was confirmed by flow cytometry, using an anti-mouse Fab antibody to detect the murine anti-CD19, anti-FAP, anti-CAIX or anti-BCMA scFv. T-cells were washed three times in 3 mL staining buffer (PBS, 4% BSA) and re-suspended in PBS at 1×10^6 cells per well. For dead cell exclusion, cells were incubated with Live dead aqua (Invitrogen) for 30 minutes on ice. Cells were washed twice with PBS and re-

suspended in 50 μ L staining buffer. To block Fc receptors, 1 μ L of 1:100 diluted normal goat IgG (Life Technologies) was added to each tube and incubated in ice for 10 minutes. 1.0 mL FACS buffer was added to each tube, mixed well, and cells were pelleted by centrifugation at 300g for 5 min. Surface expression of scFv TFPs was detected by biotin-labeled polyclonal goat anti-mouse-F(ab)₂ antibodies (Life Technologies) with biotin-labeled normal polyclonal goat IgG antibodies (Life Technologies) serving as an isotype control. Both antibodies were added at 10 μ g/mL in a reaction volume of 100 μ L. Cells were then incubated at 4 °C for 45 minutes, washed once, re-suspended in FACS buffer, and blocked with normal mouse IgG (Invitrogen) by adding 100 μ L 1:1000 diluted normal mouse IgG to each tube. The cells were then incubated on ice for 10 minutes, washed with stain buffer and re-suspended in 100 μ L stain buffer. The cells were then stained by the addition of 1.0 μ L phycoerythrin (PE)-labeled streptavidin (BD Biosciences) and APC anti-human CD3 antibody (Clone-UCHT1, BD Biosciences), PerCP/Cy5.5 anti-human CD8 antibody (Clone-SK1, BD Biosciences) and Pacific Blue anti-human CD4 antibody (Clone-RPA-T4, BD Biosciences) were added to each tube. Flow cytometry was performed using LSRFortessa™ X20 (BD Biosciences) and data was acquired using FACS diva software and was analyzed with FlowJo (Treestar, Inc. Ashland, OR). Between 20% and 40% of the transduced T-cells expressed anti-CD19 CAR, anti-CD19 LL TFP, anti-CD19 SL TFP or anti-BCMA TFP, indicating comparable levels of transduction and surface expression of CAR and TFP constructs (**FIGs. 5-7**).

[0397] Similar experiments can be carried out with FAP.TFP and CAIX.TFP constructs.

Example 7: Cytotoxicity assay by Flow Cytometry

[0398] Target cells that were either positive or negative for the respective CD19, FAP, CAIX or BCMA targets, were labelled with the fluorescent dye, carboxyfluorescein diacetate succinimidyl ester (CFSE). These target cells were mixed with effector T-cells that were either un-transduced, transduced with control CAR-T constructs, or transduced with TFPs. After the indicated incubation period, the percentage of dead to live CFSE-labeled target cells and negative control target cells was determined for each effector/target cell culture by flow cytometry. The percent survival of target cells in each T-cell + target cell culture was calculated relative to wells containing target cells alone.

[0399] The cytotoxic activity of effector T-cells was measured by comparing the number of surviving target cells in target cells without or with effector T-cells, following co-incubation of effector and target cells, using flow cytometry. In experiments with CD19 TFPs or CAR-T-cells, the target cells were CD19-positive Raji Burkitt lymphoma cells (ATCC, CCL-86), while cells used as a negative control were CD19-negative K562 cells (ATCC, CCL-243). In experiments with BCMA TFP T-cells, the target cells were BCMA-positive RPMI-8226

plasmacytoma/myeloma cells (ATCC, CCL-155), while cells used as a negative control were BCMA-negative Raji Burkitt's lymphoma cells (ATCC, CCL-86).

[0400] Target cells were washed once, and re-suspended in PBS at 1×10^6 cells/mL. The fluorescent dye carboxyfluorescein diacetate succinimidyl ester (CFSE) (ThermoFisher) was added to the cell suspension at a concentration of 0.03 μ M and the cells were incubated for 20 minutes at room temperature. The labeling reaction was stopped, by adding to the cell suspension with complete cell culture medium (RPMI-1640 + 10% HI-FBS) at the volume 5 times of the reaction volume, and the cells were incubated for an additional 2 minutes at room temperature. The cells were pelleted by centrifugation and re-suspended in cytotoxicity medium (Phenol red-free RPMI1640 (Invitrogen) plus 5% AB serum (Gemini Bioproducts) at 2×10^5 cells/mL. Fifty microliters of CFSE labelled-target cell suspension (equivalent to 10,000 cells) were added to each well of the 96-well U-bottom plate (Corning).

[0401] Effector T-cells transduced with BCMA TFP constructs, together with non-transduced T-cells as negative controls, were washed and suspended at 2×10^6 cells/mL, or 1×10^6 cells/mL in cytotoxicity medium. 50 μ L of effector T-cell suspensions (equivalent to 100,000 or 50,000 cells) were added to the plated target cells to reach the effector-to-target ratio of 10-to-1 or 5-to-1, respectively, in a total volume of 100 μ L. The cultures were then mixed, spin down, and incubated for 4 hours at 37 °C, 5% CO₂. Immediately following this incubation, 7AAD (7-aminoactinomycin D) (BioLegend) was added to the cultured cells as recommended by the manufacturer, and flow cytometry was performed with a BD Fortessa X-20 (BD Biosciences). Analysis of flow cytometric data was performed using FlowJo software (TreeStar, Inc.).

[0402] The percentage of survival for RPMI-8226 target cells was calculated by dividing the number of alive RPMI-8226 target cells (CFSE+7-AAD-) in sample with effector T-cells and target cells, by the number of alive RPMI-8226 (CFSE+7-AAD-) cells in the sample with target cells alone. The Cytotoxicity for effector cells was calculated as the percentage of killing for RPMI-8226 = 100% - percentage of survival for RPMI-8226 cells.

[0403] As previously described, T-cells transduced with an anti-CD19 28 ζ CAR construct demonstrated cytotoxicity against CD19-expressing Raji B cells, when compared to T-cells that were either non-transduced or were transduced with a non-CD19-specific CAR control (**FIG. 8**). However, T-cells transduced with anti-CD19-CD3 ϵ induced more efficient cytotoxicity against the Raji targets than the anti-CD19 CAR control at all effector:target ratios tested. Anti-CD19-CD3 γ TFPs also mediated robust cytotoxicity that was greater than that observed with anti-CD19-CAR at effector:target ratios between 5 and 10:1 (**FIG. 8**). Some cytotoxicity was observed with anti-CD19-TCR α and anti-CD19-TCR β TFPs. Similar results were obtained with anti-CD19 TFPs constructed with an alternative hinge region. Once again, cytotoxicity against

CD19-expressing Raji target cells was greater with anti-CD19-CD3 ϵ or anti-CD19-CD3 γ TFP-transduced T-cells than with anti-CD19-CAR-transduced T-cells.

[0404] T-cells electroporated with mRNA encoding TFPs specific for CD-19 also demonstrated robust cytotoxicity against CD19-expressing Raji cells. While no significant killing of the CD19-negative K562 cells was seen with either control or anti-CD19 TRuC constructs, CD19-specific killing of Raji was observed by T cells transduced with either anti-CD19-CD3 ϵ SL, or anti-CD19-CD3 γ SL TRuCs (**FIG. 14**).

[0405] T-cells transduced with TFPs specific for B-cell maturation antigen (BCMA) also demonstrated robust cytotoxicity against BCMA-expressing RPMI8226 cells. T-cells transduced with anti-BCMA-CD3 ϵ or anti-BCMA-CD3 γ TFPs efficiently mediated cytotoxicity against the BCMA-expressing RPMI8226 target cells. At 10:1 ratio of effectors to target cells, almost 100% of the target cells were killed (**FIG. 9**).

[0406] Similar experiments can be carried out with FAP.TFP and CAIX.TFP constructs.

Example 8: Cytotoxicity by Real Time Cytotoxicity Assay

[0407] Anti-CD19 and anti-BCMA TFPs also demonstrated superior cytotoxicity to anti-CD19 CARs in the real-time cytotoxicity assay (RTCA) format. The RTCA assay measures the electrical impedance of an adherent target cell monolayer, in each well of a specialized 96-well plate, in real time and presents the final readout as a value called the cell index. Changes in cell index indicate disruption of the target cell monolayer as a result of killing of target cells by co-incubated T-cell effectors. Thus the cytotoxicity of the effector T-cells can be evaluated as the change in cell index of wells with both target cells and effector T-cells compared to that of wells with target cells alone.

[0408] Target cells for RTCA were HeLa cells expressing either CD19 (CD19-HeLa) or BCMA (BCMA-HeLa) with parental, non-transduced, HeLa cells as negative controls. The DNA encoding full-length human CD19 or BCMA was synthesized by GeneArt (ThermoFisher) and inserted into the multiple cloning site of dual-promoter lentiviral vector pCDH514B (System Bioscience) carrying neomycin as selection marker, under the control of EF1 α promoter. Lentivirus carrying either the CD19 or BCMA encoding vector was then packaged. HeLa cells were transduced with either CD19- or BCMA-lentivirus for 24 hours and then selected with G418 (1 mg/mL). The expression of CD19 or BCMA by the transduced CD19-HeLa or BCMA-HeLa was confirmed by FACS analysis with anti-human CD19 or BCMA antibodies (BioLegend, clone# 19A2; Miltenyi, clone# REA315).

[0409] Adherent target cells were cultured in DMEM, 10% FBS, 1% Antibiotic-Antimycotic (Life Technologies). To prepare the RTCA, 50 μ L of RPMI medium was added into the appropriate wells of an E-plate (ACEA Biosciences, Inc, Catalog#: JL-10-156010-1A). The

plate was then placed into a RTCA MP instrument (ACEA Biosciences, Inc.) and the appropriate plate layout and assay schedule entered into the RTCA 2.0 software as described in the manufacturers manual. Baseline measurement was performed every 15 minutes for 100 measurements. 1×10^4 target cells in a 100 μ L volume were then added to each assay well and the cells were allowed to settle for 15 minutes. The plate was returned to the reader and readings were resumed.

[0410] The next day, effector T-cells were washed and re-suspended in cytotoxicity media (Phenol red-free RPMI1640 (Invitrogen) plus 5% AB serum (Gemini Bioproducts; 100-318)). The plate was then removed from the instrument and the effector T-cells, suspended in cytotoxicity medium (Phenol red-free RPMI1640 + 5% AB serum), were added to each well at 100,000 cells or 50,000 cells to reach the effector-to-target ratio of 10-to-1 or 5-to-1, respectively. The plate was then placed back to the instrument. The measurement was carried out for every 2 minutes for 100 measurements, and then every 15 minutes for 1000 measurements.

[0411] In the RTCA assay, killing of CD19-transduced HeLa was observed by T-cells transduced with anti-CD19-28 ζ CAR-transduced T-cells, as demonstrated by a time-dependent decrease in the cell index following addition of the effector cells relative to HeLa alone or HeLa co-incubated with T-cells transduced with a control CAR construct (**FIG. 11**). However, target cell killing by anti-CD19-CD3 ϵ or anti-BCMA-CD3 γ TFP-expressing T-cells was deeper and more rapid than that observed with the anti-CD19 CAR. For example, within 4 hours of addition of T-cells transduced with anti-CD19-CD3 ϵ TFP, killing of the CD19-expressing target cells was essentially complete. Little or no killing was observed with T-cells transduced with a number of TFP constructs comprising other CD3 and TCR constructs. Similar results were obtained with anti-CD19 TFPs constructed with an alternative hinge region. Cytotoxicity against CD19-transduced HeLa target cells was again greater with anti-CD19-CD3 ϵ or anti-CD19-CD3 γ TFP-transduced T-cells than with anti-CD19-CAR-transduced T-cells.

[0412] T-cells transduced with anti-BCMA TFPs also demonstrated robust cytotoxicity against BCMA-expressing RPMI8226 cells. As shown in **FIG. 9**, T-cells transduced with anti-BCMA-CD3 ϵ or anti-BCMA-CD3 γ TFPs efficiently mediated cytotoxicity against the BCMA-expressing RPMI8226 target cells. At an effector to target ratio of 10:1, almost 100% of the target cells were killed (**FIG. 12**).

[0413] The cytotoxic activity of TFP-transduced T-cells was dose-dependent with respect to the amount of virus (MOI) used for transduction. Increased killing of CD19-HeLa was observed with increasing MOI of anti-CD19-CD3 ϵ TFP lentivirus, further reinforcing the relationship between TFP transduction and cytotoxic activity (**FIG. 13**).

[0414] Similar experiments can be carried out with FAP.TFP and CAIX.TFP constructs.

Example 9: IL-2 and IFN- γ Secretion by ELISA

[0415] Another measure of effector T-cell activation and proliferation associated with the recognition of cells bearing cognate antigen is the production of effector cytokines such as interleukin-2 (IL-2) and interferon-gamma (IFN- γ).

[0416] ELISA assays for Human IL-2 (catalog #EH2IL2, Thermo Scientific) and IFN- γ catalog #KHC4012, Invitrogen) were performed as described in the product inserts. Briefly, 50 μ L of reconstituted standards or samples in duplicate were added to each well of a 96 well plate followed by 50 μ L of Biotinylated Antibody Reagent. Samples were mixed by gently tapping the plate several times. 50 μ L of Standard Diluent was then added to all wells that did not contain standards or samples and the plate was carefully sealed with an adhesive plate cover prior to incubation for 3 hours at room temperature (20-25 °C). The plate cover was then removed, plate contents were emptied, and each well was filled with Wash Buffer. This wash procedure was repeated a total of 3 times and the plate was blotted onto paper towels or other absorbent material. 100 μ L of prepared Streptavidin-HRP Solution was added to each well and a new plate cover was attached prior to incubation for 30 minutes at room temperature. The plate cover was again removed, the plate contents were discarded, and 100 μ L of TMB Substrate Solution was added into each well. The reaction was allowed to develop at room temperature in the dark for 30 minutes, after which 100 μ L of Stop Solution was added to each well. Evaluate the plate. Absorbance was measured on an ELISA plate reader set at 450 nm and 550 nm within 30 minutes of stopping the reaction. 550 nm values were subtracted from 450 nm values and IL-2 amounts in unknown samples were calculated relative to values obtained from an IL-2 standard curve.

[0417] Alternatively, 2-Plex assays were performed using the Human Cytokine Magnetic Buffer Reagent Kit (Invitrogen, LHB0001M) with the Human IL-2 Magnetic Bead Kit (Invitrogen, LHC0021M) and the Human IFN- γ Magnetic Bead Kit (Invitrogen, LHC4031M). Briefly, 25 μ L of Human IL-2 and IFN- γ antibody beads were added to each well of a 96 well plate and washed using the following guidelines: two washes of 200 μ L 1x wash solution, placing the plate in contact with a Magnetic 96-well plate Separator (Invitrogen, A14179), letting the beads settle for 1 minute and decanting the liquid. Then, 50 μ L of Incubation Buffer was added to each well of the plate with 100 μ L of reconstituted standards in duplicates or 50 μ L of samples (supernatants from cytotoxicity assays) and 50 μ L of Assay Diluent, in triplicate, for a total volume of 150 μ L. Samples were mixed in the dark at 600 rpm with an orbital shaker with a 3 mm orbital radius for 2 hours at room temperature. The plate was washed following the same washing guidelines and 100 μ L of human IL-2 and IFN- γ biotinylated detector antibody was

added to each well. Samples were mixed in the dark at 600 rpm with an orbital shaker with a 3 mm orbital radius for 1 hour at room temperature. The plate was washed following the same washing guidelines and 100 μ L of Streptavidin-R-Phycoerythrin was added to each well. Samples were mixed in the dark at 600 rpm with an orbital shaker with a 3 mm orbital radius for 30 minutes at room temperature. The plate was washed 3 times using the same washing guidelines and after decanting the liquid the samples were re-suspended in 150 μ L of 1x wash solution. The samples were mixed at 600 rpm with an orbital shaker with a 3 mm orbital radius for 3 minutes and stored over night at 4 °C. Afterwards, the plate was washed following the same washing guidelines and the samples were re-suspended in 150 μ L of 1x wash solution. [0418] The plate was read using the MAGPIX System (Luminex) and xPONENT software. Analysis of the data was performed using MILLIPLEX Analyst software, which provides the standard curve and cytokine concentrations.

[0419] **FIG. 15** shows that, relative to non-transduced or control CAR-transduced T-cells, T-cells transduced with anti-CD19 TFPs produce higher levels of both IL-2 and IFN- γ when co-cultured with either Raji cells that endogenously express CD19 or CD19-transduced HeLa cells. In contrast, co-culture with CD19 negative K562 cells or non-transduced HeLa cells, results in little or no cytokine release from TFP-transduced T-cells. Consistent with the previous cytotoxicity data, anti-CD19 TFPs constructed with an alternative hinge region generated similar results upon co-culture with CD19-bearing target cells (**FIG. 16**).

[0420] In agreement with the previous cytotoxicity data, anti-CD19-CD3 ϵ and anti-CD19-CD3 γ produced the highest IL-2 and IFN- γ levels of the TFP constructs (**FIGs. 15 and 16**). However, cytokine production by T-cells transduced with anti-CD19-CD3 ϵ and anti-CD19-CD3 γ TFPs was comparable to that of T-cells expressing anti-CD19-28 ζ CAR, despite the TFPs demonstrating much higher levels of target cell killing (**FIGs. 8 and 11**). The possibility that TFPs may more efficiently kill target cells than CARs, but release comparable or lower levels of pro-inflammatory cytokines, represents a potential advantage for TFPs relative to CARs since elevated levels of these cytokines have been associated with dose-limiting toxicities for adoptive CAR-T therapies.

[0421] T-cells transduced with anti-BCMA-CD3 ϵ or anti-BCMA-CD3 γ TFPs also produced IL-2 and IFN- γ upon co-culture with BCMA-HeLa but not control HeLa cells that did not express BCMA (**FIG. 17**).

[0422] Similar experiments can be carried out with FAP.TFP and CAIX.TFP constructs.

Example 10: CD107a Exposure by Flow Cytometry

[0423] An additional assay for T-cell activation is surface expression of CD107a, a lysosomal associated membrane protein (LAMP-1) that is located in the membrane of cytoplasmic

cytolytic granules in resting cells. Degranulation of effector T-cells, a prerequisite for cytolytic activity, results in mobilization of CD107a to the cell surface following activation-induced granule exocytosis. Thus, CD107a exposure provides an additional measure of T-cell activation, in addition to cytokine production, that correlates closely with cytotoxicity.

[0424] Target and effector cells were separately washed and re-suspended in cytotoxicity medium (RPMI+5% human AB serum + 1% antibiotic antimycotic). The assay was performed by combining 2×10^5 effectors cells with 2×10^5 target cells in a 100 μ L final volume in U-bottom 96-well plates (Corning), in the presence of 0.5 μ L/well of PE/Cy7-labelled anti-human CD107a (LAMP-1) antibody (Clone-H4A3, BD Biosciences). The cultures were then incubated for an hour at 37 °C, 5% CO₂. Immediately following this incubation, 10 μ L of a 1:10 dilution of the secretion inhibitor monensin (1000x solution, BD GolgiStop™) was carefully added to each well without disturbing the cells. The plates were then incubated for a further 2.5 hours at 37 °C, 5% CO₂. Following this incubation, the cells were stained with APC anti-human CD3 antibody (Clone-UCHT1, BD Biosciences), PerCP/Cy5.5 anti-human CD8 antibody (Clone-SK1, BD Biosciences) and Pacific Blue anti-human CD4 antibody (Clone-RPA-T4, BD Biosciences) and then incubated for 30 minutes at 37 °C, 5% CO₂. The cells were then washed 2x with FACS buffer (and resuspended in 100 μ L FACS buffer and 100ul IC fix buffer prior to analysis.

[0425] Exposure of CD107a on the surface of T-cells was detected by flow cytometry. Flow cytometry was performed with a LSRFortessa™ X20 (BD Biosciences) and analysis of flow cytometric data was performed using FlowJo software (Treestar, Inc. Ashland, OR). The percentage of CD8+ effector cells, within the CD3 gate, that were CD107 +ve was determined for each effector/target cell culture.

[0426] Consistent with the previous cytotoxicity and cytokine data, co-culture of CD19-expressing target cells, such as Raji or Nalm-6 cells, with effector T-cells transduced with anti-CD19-28 ζ CAR induced a 3 to 5-fold increase in surface CD107a expression relative to effectors incubated with CD19 –ve target cells (**FIG. 18**). In comparison, under the same conditions, anti-CD19-CD3 ϵ LL or anti-CD19-CD3 γ LL TFP-expressing effectors exhibited a 5 to 7-fold induction of CD107a expression. Anti-CD19 TFPs constructed with an alternative hinge region generated similar results upon co-culture with CD19-bearing target cells.

[0427] Relative to non-transduced T-cells, cells transduced with anti-BCMA-CD3 ϵ or anti-BCMA-CD3 γ TFPs also exhibited an increase in surface expression of CD107a upon co-culture with BCMA +ve RPMI8226 cells (**FIG. 19**). These results indicate that TFP-transduced effector T-cells become activated and degranulate upon exposure to target cells expressing their cognate antigen.

[0428] Similar experiments can be carried out with FAP.TFP and CAIX.TFP constructs.

Example 11: *In Vivo* Mouse Efficacy Studies

[0429] To assess the ability of effector T-cells transduced with anti-CD19 TFPs to achieve anti-tumor responses *in vivo*, effector T-cells transduced with either anti-CD19-28 ζ CAR, anti-CD19-CD3 ϵ LL TFP or anti-CD19-CD3 γ LL TFP were adoptively transferred into NOD/SCID/IL-2R $\gamma^{-/-}$ (NSG-JAX) mice that had previously been inoculated with CD19⁺ Raji or Nalm6 human leukemic cell lines.

[0430] Female NOD/SCID/IL-2R $\gamma^{-/-}$ (NSG-JAX) mice, at least 6 weeks of age prior to the start of the study, were obtained from The Jackson Laboratory (stock number 005557) and acclimated for 3 days before experimental use. Raji and Nalm-6 human leukemic cell lines for inoculation were maintained in log-phase culture prior to harvesting and counting with trypan blue to determine a viable cell count. On the day of tumor challenge, the cells were centrifuged at 300g for 5 minutes and re-suspended in pre-warmed sterile PBS at either 1×10^6 cells/100 μ L (Nalm-6) or 5×10^5 cells/100 μ L (Raji). T-cells for adoptive transfer, either non-transduced or transduced with anti-CD19-28 ζ CAR, anti-CD19-CD3 ϵ LL TFP or anti-CD3 γ LL TFP constructs were prepared. On day 0 of the study, 10 animals per experimental group were challenged intravenously with either 5×10^5 Raji or 1×10^6 Nalm-6 cells. 3 days later, 5×10^6 of the indicated effector T-cell populations were intravenously transferred to each animal in 100 μ L of sterile PBS. Detailed clinical observations on the animals were recorded daily until euthanasia. Body weight measurements were made on all animals weekly until death or euthanasia. All animals were euthanized 35 days after adoptive transfer of test and control articles. Any animals appearing moribund during the study were euthanized at the discretion of the study director in consultation with a veterinarian.

[0431] Relative to non-transduced T-cells, adoptive transfer of T-cell transduced with either anti-CD19-28 ζ CAR, anti-CD19-CD3 ϵ LL TFP or anti-CD19-CD3 γ LL TFP prolonged survival of both Raji (**FIG. 20A**) and Nalm6 (**FIG. 20B**) tumor-bearing mice, indicating that both anti-CD19 CAR and TFP-transduced T-cells were capable of mediating target cell killing with corresponding increased survival in these mouse models. Collectively, these data indicate that TFPs represent an alternative platform for engineering chimeric receptors that demonstrate superior antigen-specific killing to first generation CARs both *in vitro* and *in vivo*.

[0432] Similar experiments can be carried out with FAP.TFP and CAIX.TFP constructs.

[0433] While preferred embodiments of the present invention have been shown and described herein, it will be obvious to those skilled in the art that such embodiments are provided by way of example only. Numerous variations, changes, and substitutions will now occur to those skilled in the art without departing from the invention. It should be understood that various alternatives to the embodiments of the invention described herein may be employed in practicing

the invention. It is intended that the following claims define the scope of the invention and that methods and structures within the scope of these claims and their equivalents be covered thereby.

CLAIMS

1. A pharmaceutical composition comprising
 - (I) a T cell from a human subject, wherein the T cell comprises a recombinant nucleic acid molecule encoding a T-cell receptor (TCR) fusion protein (TFP) comprising
 - (a) a TCR subunit comprising
 - (i) at least a portion of a TCR extracellular domain,
 - (ii) a TCR transmembrane domain, and
 - (iii) a TCR intracellular domain comprising a stimulatory domain from an intracellular signaling domain of CD3 epsilon, CD3 gamma, or CD3 delta; and
 - (b) a murine, human or humanized scFv or single domain antibody comprising an antigen binding domain;wherein the TCR subunit and the antibody domain are operatively linked, wherein the TFP incorporates into a TCR when expressed in a T-cell, and wherein the T cell exhibits increased cytotoxicity to a human cell expressing an antigen that specifically interacts with the antigen binding domain compared to a T cell not containing the TFP; and
 - (II) a pharmaceutically acceptable carrier.
2. The pharmaceutical composition of claim 1, wherein the antigen binding domain is an anti-CD19 binding domain or an anti-B-cell maturation antigen (BCMA) binding domain.
3. The pharmaceutical composition of claim 1 or 2, wherein the encoded antigen binding domain is connected to the TCR extracellular domain by a linker sequence, optionally, wherein the encoded linker sequence comprises $(G_4S)_n$, wherein $n=1$ to 4.
4. The pharmaceutical composition of any one of claims 1-3, wherein the TCR subunit comprises (i) a TCR extracellular domain, (ii) a TCR transmembrane domain, and (iii) a TCR intracellular domain, wherein at least two of (i), (ii), and (iii) are from the same TCR subunit, optionally, wherein (i), (ii), and (iii) are from the same TCR subunit.
5. The pharmaceutical composition of any one of claims 1-4, wherein the human or humanized antibody domain comprises a scFv or a single domain antibody, wherein the single domain antibody is optionally a V_H domain.
6. The pharmaceutical composition of any one of claims 1-5, wherein the encoded antigen binding domain comprises (i) a light chain (LC) CDR1, LC CDR2 and LC CDR3 of an anti-CD19 light chain binding domain amino acid sequence with 70-100% sequence identity to SEQ ID NO: 25, SEQ ID NO: 27 and SEQ ID NO: 29, respectively, and/or (ii) a heavy chain (HC) CDR1, HC CDR2 and HC CDR3 of an anti-CD19 heavy chain binding domain amino acid sequence with 70-100% sequence identity to SEQ ID NO: 31, SEQ ID NO: 33 and SEQ ID NO: 35, respectively.

7. The pharmaceutical composition of any one of claims 1-6, wherein the TFP includes an extracellular domain of a TCR subunit that comprises an extracellular domain or portion thereof of a protein selected from the group consisting of a TCR alpha chain, a TCR beta chain, a CD3 epsilon TCR subunit, a CD3 gamma TCR subunit, a CD3 delta TCR subunit, functional fragments thereof, and amino acid sequences thereof having at least one but not more than 20 modifications.

8. The pharmaceutical composition of any one of claims 1-7, wherein the TFP includes a transmembrane domain that comprises a transmembrane domain of a protein selected from the group consisting of a TCR alpha chain, a TCR beta chain, a CD3 epsilon TCR subunit, a CD3 gamma TCR subunit, a CD3 delta TCR subunit, functional fragments thereof, and amino acid sequences thereof having at least one but not more than 20 modifications.
9. The pharmaceutical composition of any one of claims 1-8, wherein the TFP includes a transmembrane domain that comprises a transmembrane domain of a protein selected from the group consisting of a TCR alpha chain, a TCR beta chain, a TCR zeta chain, a CD3 epsilon TCR subunit, a CD3 gamma TCR subunit, a CD3 delta TCR subunit, CD45, CD4, CD5, CD8, CD9, CD16, CD22, CD33, CD28, CD37, CD64, CD80, CD86, CD134, CD137, CD154, functional fragments thereof, and amino acid sequences thereof having at least one but not more than 20 modifications.
10. The pharmaceutical composition of any one of claims 1-9, wherein the recombinant nucleic acid molecule further comprises a sequence encoding a costimulatory domain, optionally, wherein the costimulatory domain is a functional signaling domain obtained from a protein selected from the group consisting of OX40, CD2, CD27, CD28, CD30, ICAM-1, LFA-1 (CD11a/CD18), ICOS (CD278), and 4-1BB (CD137), and amino acid sequences thereof having at least one but not more than 20 modifications thereto.
11. The pharmaceutical composition of any one of claims 1-10, wherein the TFP includes an immunoreceptor tyrosine-based activation motif (ITAM) of a TCR subunit that comprises an ITAM or portion thereof of a protein selected from the group consisting of CD3 zeta TCR subunit, CD3 epsilon TCR subunit, CD3 gamma TCR subunit, CD3 delta TCR subunit, TCR zeta chain, Fc epsilon receptor 1 chain, Fc epsilon receptor 2 chain, Fc gamma receptor 1 chain, Fc gamma receptor 2a chain, Fc gamma receptor 2b1 chain, Fc gamma receptor 2b2 chain, Fc gamma receptor 3a chain, Fc gamma receptor 3b chain, Fc beta receptor 1 chain, TYROBP (DAP12), CD5, CD16a, CD16b, CD22, CD23, CD32, CD64, CD79a, CD79b, CD89, CD278, CD66d, functional fragments thereof, and amino acid sequences thereof having at least one but not more than 20 modifications thereto.
12. The pharmaceutical composition of any one of claims 1-11, wherein the T cell is a primary T-cell.
13. The pharmaceutical composition of any one of claims 1-12, wherein the T cell is a CD8+ or CD4+ T cell.
14. The pharmaceutical composition of any one of claims 1-13, wherein the T cell further comprises a nucleic acid encoding at least a portion of an inhibitory molecule, wherein the at least a portion of an inhibitory molecule associates with a second polypeptide that comprises a positive signal from an intracellular signaling domain, optionally, wherein the inhibitory molecule comprise first polypeptide that comprises at least a portion of PD1 and a second polypeptide comprising a costimulatory domain and primary signaling domain.
15. The pharmaceutical composition of any one of claims 1-14, wherein the T cell forms a conjugate with a human cell expressing an antigen that specifically interacts with the antigen binding domain, production

- of IL-2 by the T cell is increased in the presence of a human cell expressing an antigen that specifically interacts with the antigen binding domain, or wherein the T cell exhibits increased cytotoxicity to a human cell expressing an antigen that specifically interacts with the antigen binding domain.
16. The pharmaceutical composition of any one of claims 1-15, wherein the TCR intracellular domain comprises a stimulatory domain from an intracellular signaling domain of CD3 epsilon, CD3 gamma or CD3 delta.
 17. Use of an effective amount of the pharmaceutical composition of any one of claims 1-16 in the manufacture of a medicament for treating a mammal having a disease.
 18. The use of claim 17, wherein the disease is selected from the group consisting of a proliferative disease, a cancer, a malignancy, myelodysplasia, a myelodysplastic syndrome, and a preleukemia.
 19. The use of claim 17, wherein the disease is a hematologic cancer selected from the group consisting of B-cell acute lymphoid leukemia (B-ALL), T-cell acute lymphoid leukemia (T-ALL), acute lymphoblastic leukemia (ALL); chronic myelogenous leukemia (CML), chronic lymphocytic leukemia (CLL), B cell prolymphocytic leukemia, blastic plasmacytoid dendritic cell neoplasm, Burkitt's lymphoma, diffuse large B cell lymphoma, follicular lymphoma, hairy cell leukemia, small cell-follicular lymphoma, large cell-follicular lymphoma, malignant lymphoproliferative conditions, MALT lymphoma, mantle cell lymphoma, Marginal zone lymphoma, multiple myeloma, myelodysplasia, myelodysplastic syndrome, non-Hodgkin's lymphoma, plasmablastic lymphoma, plasmacytoid dendritic cell neoplasm, Waldenstrom macroglobulinemia, preleukemia, a disease associated with CD19 or BCMA expression, and combinations thereof.
 20. The use of any one of claims 17-19, wherein the pharmaceutical composition is to be administered in combination with an agent that treats the disease associated with CD19 or BCMA.