

DESCRIPTION

Title of the Invention: ISOINDOLINE-1-ONE DERIVATIVES
AS CHOLINERGIC MUSCARINIC M1 RECEPTOR POSITIVE ALLOESTERIC
MODULATOR ACTIVITY FOR THE TREATMENT OF ALZHEIMERS DISEASE

5

Technical Field

[0001]

The present invention relates to a nitrogen-containing heterocyclic compound which has a cholinergic muscarinic M1
10 receptor positive allosteric modulator activity and is useful as a medicament such as an agent for the prophylaxis or treatment of Alzheimer's disease, schizophrenia, pain, sleep disorder, Parkinson's disease dementia, dementia with Lewy bodies and the like. As used herein, the positive allosteric
15 modulator activity refers to an action to potentiate receptor function by binding at a different site from that of an endogenous activator (acetylcholine for this receptor).

[0002]

(Background of the Invention)


20 Acetylcholine is a neurotransmitter that induces signal transduction in the central nervous system and the neuromuscular connections (the parasympathetic nerve and motor nerve). In the central nervous system, nuclei of origin of the acetylcholine neuron are in the brain stem and forebrain, and
25 those acetylcholine neurons project to cerebral cortex, hippocampus, and limbic area. In addition, some interneurons in some brain areas such as striatum utilize acetylcholine as a neurotransmitter. Acetylcholine receptor is classified into a ligand dependent-ion channel (cholinergic nicotinic receptor)
30 and a G-protein-coupled receptor (cholinergic muscarinic receptor). The cholinergic muscarinic receptor is one kind of receptor for excitatory neurotransmitter, acetylcholine, and was named based on the selective activation of the receptor by muscarine. The muscarinic receptor is further classified into
35 subtypes of M1 to M5. The M1 receptor is known to be mainly

distributed in the brain, and deeply involved particularly in learning, memory, sleep, neuropathic pain, and the like. The importance of cholinergic muscarinic M1 receptor in brain physiology is well known, and a compound which enhances M1
5 receptor function is expected to be useful as an agent for the prophylaxis or treatment of mental diseases, neurodegenerative diseases, memory disorders, pain, sleep disorders, Parkinson's disease dementia, dementia with Lewy bodies and the like (non-patent document 1).

10 [0003]

WO 02/081447 A1 (Patent Document 1) discloses the following compound as a compound having a tumor necrosis factor- α (TNF- α) or a cAMP phosphodiesterase IV (PDE4) inhibitory activity and useful for the prophylaxis or treatment
15 of inflammation and autoimmune disease.

[0004]




[0005]

wherein each symbol is as defined in the document.

20 [0006]

WO 02/081446 A1 (Patent Document 2) discloses the following compound as a compound having a tumor necrosis factor- α (TNF- α) or a cAMP phosphodiesterase IV (PDE4) inhibitory activity and useful for the prophylaxis or treatment of
25 inflammation and autoimmune disease.

[0007]




[0008]

wherein each symbol is as defined in the document.

[0009]

5 WO 2006/020879 A1 (Patent Document 3) discloses the following compound as a glutamic acid receptor potentiator useful for the prophylaxis or treatment of psychoneurotic disorder associated with glutamate dysfunction.

[0010]



10


[0011]

wherein each symbol is as defined in the document.

[0012]

15 WO 2013/063549 A1 (Patent Document 4) discloses the following compound as a compound useful for the prophylaxis or treatment of psychoneurotic disorder associated with muscarinic acetylcholine receptor dysfunction.

[0013]




[0014]

wherein each symbol is as defined in the document.

[0015]

5 Bioorganic & Medicinal Chemistry Letters, 20 (2010) 1792-1795 (Non-Patent Document 2) discloses the following compound as an M1 receptor positive allosteric modulator.


[0016]



10 [0017]

Gordon, C. P., Byrne, N., McCluskey, A. Green Chem., 2010, 12, 1000-1006. (Non-Patent Document 3) discloses the following compound similar to the compound of the present invention.

[0018]




15

[0019]

WO 2010/096338 A1 (Patent Document 5) discloses the

following compound as an M1 receptor positive allosteric modulator useful for the prophylaxis or treatment of a Alzheimer's disease, schizophrenia, pain or sleep disorder.

[0020]



5


[0021]

wherein each symbol is as defined in the document.

[0022]

WO 95/030647 A1 (Patent Document 6) discloses the following compound similar to the compound of the present invention.

[0023]



[0024]


15 wherein each symbol is as defined in the document.

[0025]

WO 2007/139464 A1 (Patent Document 7) discloses the following compound as a CB₁ receptor ligand useful for the prophylaxis or treatment of pain, cancer, multiple sclerosis, Parkinson's disease, Huntington's disease, Alzheimer's disease,

anxiety disorder, gastrointestinal disorder and cardiovascular disorder.

[0026]




5 [0027]

wherein each symbol is as defined in the document.

[0028]

US 2008/0108659 A1 (Patent Document 8) discloses the following compound as a compound having poly(ADP ribose)
 10 polymerase (PARP) activity and useful for the prophylaxis or treatment of cancer, central nervous system disease, inflammation disease and the like.

[0029]



15 [0030]


wherein each symbol is as defined in the document.

[0031]

WO 2011/006794 A1 (Patent Document 9) discloses the following compound as a compound selectively inhibiting an
 20 activity of poly (ADP-ribose) polymerase PARP-1 with respect to poly (ADP-ribose) polymerase PARP-2 and useful for the

prophylaxis or treatment of cancer, cardiovascular disorder, central nervous system disorder and the like.

[0032]




5 [0033]

wherein each symbol is as defined in the document.

[0034]

WO 2012/003147 A1 (Patent Document 10) discloses the following compound as a compound having an M1PAM activity and
10 useful for the prophylaxis or treatment of Alzheimer's disease, schizophrenia, pain, sleep disorder and the like.

[0035]




[0036]

15 wherein each symbol is as defined in the document.

[0037]

WO 2012/158475 A1 (Patent Document 11) discloses the following compound as a compound having an M1PAM activity and useful for the prophylaxis or treatment of Alzheimer's disease
20 and other diseases.

[0038]




[0039]

wherein each symbol is as defined in the document.

5 [0040]

JP-B-S44-16647 (Patent Document 12) discloses the following compound similar to the compound of the present invention.

[0041]



10

Document List

Patent Document

[0042]

- Patent Document 1: WO 02/081447 A1
- 15 Patent Document 2: WO 02/081446 A1
- Patent Document 3: WO 2006/020879 A1
- Patent Document 4: WO 2013/063549 A1
- Patent Document 5: WO 2010/096338 A1
- Patent Document 6: WO 95/030647 A1
- 20 Patent Document 7: WO 2007/139464 A1
- Patent Document 8: US 2008/0108659 A1
- Patent Document 9: WO 2011/006794 A1
- Patent Document 10: WO 2012/003147 A1
- Patent Document 11: WO 2012/158475 A1

Patent Document 12: JP-B-S44-16647

Non-Patent Document

[0043]

Non-Patent Document 1: Nature Reviews Drug Discovery, 2007, 6,
5 721-733.

Non-Patent Document 2: Bioorganic & Medicinal Chemistry
Letters, 20 (2010) 1792-1795.

Non-Patent Document 3: Gordon, C. P., Byrne, N., McCluskey, A.
Green Chem., 2010, 12, 1000-1006.

10 Summary of the Invention

Problems to be Solved by the Invention

[0044]

The development of a compound having a cholinergic
muscarinic M1 receptor (M1 receptor) positive allosteric
15 modulator activity and useful as an agent for the prophylaxis
or treatment of for Alzheimer's disease, schizophrenia, pain,
sleep disorder, Parkinson's disease dementia, dementia with
Lewy bodies and the like is desired. As used herein, the
positive allosteric modulator activity refers to an action to
20 potentiate receptor function by binding at a different site
from that of an endogenous activator (acetylcholine for this
receptor).

Means of Solving the Problems

[0045]


25 The present inventors have conducted intensive studies in
an attempt to solve the aforementioned problems and found that
a compound represented by the following formula (I) has a
cholinergic muscarinic M1 receptor positive allosteric
modulator activity, which resulted in the completion of the
30 present invention.

[0046]

Accordingly, the present invention relates to the
following.

[1] A compound represented by the formula (I):

35 [0047]



[0048]

wherein

R^1 is an optionally substituted 5- or 6-membered cyclic
5 group or an optionally substituted C_{1-6} alkyl group;

R^2 and R^3 are the same or different and each is a
hydrogen atom, a halogen atom, a cyano group, an optionally
substituted C_{1-6} alkyl group, an optionally substituted C_{1-6}
alkoxy group or an optionally substituted C_{3-6} cycloalkyl group;

10 R^4 is a halogen atom, a cyano group, an optionally
substituted C_{1-6} alkyl group, an optionally substituted C_{1-6}
alkoxy group, an optionally substituted carbamoyl group or an
optionally substituted 3- to 8-membered cyclic group; and

Ring A is an optionally further substituted 6-membered
15 aromatic ring,
or a salt thereof (in the present specification, to be referred
as compound (1)).

[0049]

[2] The compound of the above-mentioned [1], wherein R^1 is
20 (1) an optionally substituted phenyl group,
(2) an optionally substituted C_{5-6} cycloalkyl group,
(3) an optionally substituted 5- or 6-membered non-aromatic
heterocyclic group,
(4) an optionally substituted 5- or 6-membered monocyclic
25 aromatic heterocyclic group, or
(5) an optionally substituted C_{1-6} alkyl group,

or a salt thereof.

[3] The compound of the above-mentioned [1] or [2], wherein R⁴ is

- (1) a halogen atom,
 - 5 (2) a cyano group,
 - (3) a C₁₋₆ alkyl group,
 - (4) an optionally substituted C₁₋₆ alkoxy group,
 - (5) an optionally substituted carbamoyl group, or
 - (6) an optionally substituted 5- or 6-membered monocyclic
 - 10 aromatic heterocyclic group,
- or a salt thereof.

[4] The compound of any of the above-mentioned [1] to [3], wherein Ring A is a 6-membered aromatic ring optionally further substituted by 1 to 3 substituents, in addition to R⁴, selected

15 from

- (a) a halogen atom,
 - (b) a C₁₋₆ alkyl group, and
 - (c) a C₁₋₆ alkoxy group,
- or a salt thereof.

20 [0050]

[5] The compound of any of the above-mentioned [1] to [4], wherein

R¹ is

- (1) an optionally substituted phenyl group,
- 25 (2) an optionally substituted C₅₋₆ cycloalkyl group,
- (3) an optionally substituted 5- or 6-membered non-aromatic heterocyclic group,
- (4) an optionally substituted 5- or 6-membered monocyclic aromatic heterocyclic group, or
- 30 (5) an optionally substituted C₁₋₆ alkyl group;

R² is

- (1) a hydrogen atom,
- (2) a halogen atom, or
- (3) an optionally substituted C₁₋₆ alkyl group;

35 R³ is

- (1) a hydrogen atom,
- (2) a halogen atom,
- (3) a cyano group,
- (4) an optionally substituted C₁₋₆ alkyl group,
- 5 (5) a C₁₋₆ alkoxy group, or
- (6) a C₃₋₆ cycloalkyl group;

R⁴ is

- (1) a halogen atom,
- (2) a cyano group,
- 10 (3) a C₁₋₆ alkyl group,
- (4) an optionally substituted C₁₋₆ alkoxy group,
- (5) an optionally substituted carbamoyl group, or
- (6) an optionally substituted 5- or 6-membered monocyclic aromatic heterocyclic group; and

15 Ring A is a 6-membered aromatic ring optionally further substituted by 1 to 3 substituents, in addition to R⁴, selected from

- (a) a halogen atom,
- (b) a C₁₋₆ alkyl group, and
- 20 (c) a C₁₋₆ alkoxy group,

or a salt thereof.

[0051]

[6] The compound of any of the above-mentioned [1] to [5], wherein


25 R¹ is

- (1) a phenyl group optionally substituted by 1 to 3 substituents selected from
 - (i) a halogen atom, and
 - (ii) a cyano group,
- 30 (2) a C₅₋₆ cycloalkyl group optionally substituted by 1 to 3 substituents selected from
 - (i) a hydroxy group,
 - (ii) a C₁₋₆ alkyl group optionally substituted by 1 to 3 hydroxy groups, and
 - 35 (iii) a C₁₋₆ alkoxy group,

- (3) a 5- or 6-membered monocyclic non-aromatic heterocyclic group optionally substituted by 1 to 3 hydroxy groups,
(4) a 5- or 6-membered monocyclic aromatic heterocyclic group optionally substituted by 1 to 3 halogen atoms, or
5 (5) a C₁₋₆ alkyl group optionally substituted by 1 to 3 substituents selected from
 (i) a hydroxy group, and
 (ii) a 3- to 8-membered monocyclic non-aromatic heterocyclic group;
- 10 R² is
 (1) a hydrogen atom,
 (2) a halogen atom, or
 (3) a C₁₋₆ alkyl group;
 R³ is
15 (1) a hydrogen atom,
 (2) a halogen atom,
 (3) a cyano group,
 (4) a C₁₋₆ alkyl group optionally substituted by 1 to 3 halogen atoms,
20 (5) a C₁₋₆ alkoxy group, or
 (6) a C₃₋₆ cycloalkyl group;
 R⁴ is
 (1) a halogen atom,
 (2) a cyano group,
25 (3) a C₁₋₆ alkyl group,
 (4) a C₁₋₆ alkoxy group optionally substituted by 1 to 3 halogen atoms,
 (5) a carbamoyl group,
 (6) a mono- or di-C₁₋₆ alkyl-carbamoyl group, or
30 (7) a 5- or 6-membered monocyclic aromatic heterocyclic group optionally substituted by 1 to 3 C₁₋₆ alkyl groups; and
 Ring A is a 6-membered aromatic ring optionally further substituted by 1 to 3 substituents, in addition to R⁴, selected from
35 (a) a halogen atom,

(b) a C₁₋₆ alkyl group, and
(c) a C₁₋₆ alkoxy group,
or a salt thereof.


[7] The compound of the above-mentioned [6], wherein the
5 partial structure represented by the following formula:
[0052]



[0053]

in the formula (I) is a partial structure represented by the
10 following formula:

[0054]



[0055]

or a salt thereof.

15 [0056]

[8] The compound of any of the above-mentioned [1] to [7],
wherein

R¹ is

- (1) a phenyl group optionally substituted by 1 to 3
20 substituents selected from
(i) a halogen atom, and
(ii) a cyano group,
(2) a C₅₋₆ cycloalkyl group optionally substituted by 1 to 3
hydroxy groups,
25 (3) a 5- or 6-membered monocyclic non-aromatic heterocyclic
group optionally substituted by 1 to 3 hydroxy groups, or
(4) a C₁₋₆ alkyl group optionally substituted by 1 to 3
substituents selected from
(i) a hydroxy group, and
30 (ii) a 3- to 8-membered monocyclic non-aromatic

heterocyclic group;

R^2 is

- (1) a hydrogen atom,
- (2) a halogen atom, or
- 5 (3) a C_{1-6} alkyl group;

R^3 is

- (1) a hydrogen atom,
- (2) a halogen atom,
- (3) a cyano group,
- 10 (4) a C_{1-6} alkyl group optionally substituted by 1 to 3 halogen atoms,
- (5) a C_{1-6} alkoxy group, or
- (6) a C_{3-6} cycloalkyl group;

R^4 is

- 15 (1) a halogen atom,
- (2) a cyano group,
- (3) a C_{1-6} alkyl group,
- (4) a C_{1-6} alkoxy group optionally substituted by 1 to 3 halogen atoms,
- 20 (5) a mono- or di- C_{1-6} alkyl-carbamoyl group, or
- (6) a 5- or 6-membered monocyclic aromatic heterocyclic group optionally substituted by 1 to 3 C_{1-6} alkyl groups; and

Ring A is a 6-membered aromatic ring optionally further substituted by 1 to 3 substituents, in addition to R^4 , selected
25 from


(a) a halogen atom, and

(b) a C_{1-6} alkoxy group,

or a salt thereof.

[9] The compound of the above-mentioned [8], wherein the
30 partial structure represented by the following formula:


[0057]



[0058]

in the formula (I) is a partial structure represented by the following formula:

[0059]



5 [0060]

or a salt thereof.

[0061]

[10] The compound of any of the above-mentioned [1] to [9], wherein

10 R^1 is

(1) a C_{5-6} cycloalkyl group optionally substituted by 1 to 3 hydroxy groups, or

(2) a 5- or 6-membered monocyclic non-aromatic heterocyclic group optionally substituted by 1 to 3 hydroxy groups;

15 R^2 is

(1) a halogen atom, or

(2) a C_{1-6} alkyl group;

R^3 is a C_{1-6} alkyl group;

R^4 is

20 (1) a C_{1-6} alkyl group,

(2) a C_{1-6} alkoxy group, or


(3) a 5- or 6-membered monocyclic aromatic heterocyclic group;

and

Ring A is a benzene ring or a pyridine ring, each of which is unsubstituted, in addition to R^4 ,
25 or a salt thereof.

[11] The compound of the above-mentioned [10], wherein the partial structure represented by the following formula:

[0062]




30

[0063]

in the formula (I) is a partial structure represented by the following formula:

[0064]



[0065]

or a salt thereof.

[0066]

[12] The compound of any of the above-mentioned [1] to [11],
10 wherein

R^1 is

- (1) a cyclohexyl group substituted by one hydroxy group, or
- (2) a tetrahydropyranyl group substituted by one hydroxy group;

R^2 is

- 15
- (1) a halogen atom, or
 - (2) a C_{1-6} alkyl group;

R^3 is a C_{1-6} alkyl group;


R^4 is

- 20
- (1) a C_{1-6} alkyl group,
 - (2) a C_{1-6} alkoxy group, or
 - (3) a pyrazolyl group; and

Ring A is a benzene ring or a pyridine ring, each of which is unsubstituted, in addition to R^4 , or a salt thereof.

25 [13] The compound of the above-mentioned [12], wherein the partial structure represented by the following formula:

[0067]




[0068]

30 in the formula (I) is a partial structure represented by the

following formula:

[0069]



[0070]

5 or a salt thereof.

[0071]

[14] 2-[(3S,4S)-4-Hydroxytetrahydro-2H-pyran-3-yl]-6-(4-methoxybenzyl)-4,5-dimethyl-2,3-dihydro-1H-isoindol-1-one, or a salt thereof.

10 [15] 4-Fluoro-2-[(3S,4S)-4-hydroxytetrahydro-2H-pyran-3-yl]-5-methyl-6-[4-(1H-pyrazol-1-yl)benzyl]-2,3-dihydro-1H-isoindol-1-one, or a salt thereof.

[16] 2-((1S,2S)-2-Hydroxycyclohexyl)-4,5-dimethyl-6-((6-methylpyridin-3-yl)methyl)isoindolin-1-one, or a salt thereof.

15 [17] A medicament comprising the compound of any of the above-mentioned [1] to [16] or a salt thereof.

[18] The medicament of the above-mentioned [17], which is a cholinergic muscarinic M1 receptor positive allosteric modulator.

20 [19] The medicament of the above-mentioned [17], which is an agent for the prophylaxis or treatment of Alzheimer's disease, schizophrenia, pain, sleep disorder, Parkinson's disease dementia or dementia with Lewy bodies.

[20] The compound of any of the above-mentioned [1] to [16] or
25 a salt thereof for use in the prophylaxis or treatment of Alzheimer's disease, schizophrenia, pain, sleep disorder, Parkinson's disease dementia or dementia with Lewy bodies.

[21] A method of cholinergic muscarinic M1 receptor positive allosteric modulation in a mammal, which comprises
30 administering an effective amount of the compound of any of the above-mentioned [1] to [16] or a salt thereof to the mammal.

[22] A method for the prophylaxis or treatment of Alzheimer's

disease, schizophrenia, pain, sleep disorder, Parkinson's disease dementia or dementia with Lewy bodies in a mammal, which comprises administering an effective amount of the compound of any of the above-mentioned [1] to [16] or a salt thereof to the mammal.

[23] Use of the compound of any of the above-mentioned [1] to [16] or a salt thereof for the production of an agent for the prophylaxis or treatment of Alzheimer's disease, schizophrenia, pain, sleep disorder, Parkinson's disease dementia or dementia with Lewy bodies.

Effect of the Invention

[0072]

The compound of the present invention has a cholinergic muscarinic M1 receptor positive allosteric modulator activity, and is useful as an agent for the prophylaxis or treatment of, for example, Alzheimer's disease, schizophrenia, pain, sleep disorder, Parkinson's disease dementia, dementia with Lewy bodies and the like.

[0073]

(Detailed Description of the Invention)

The definition of each substituent used in the present specification is described in detail in the following. Unless otherwise specified, each substituent has the following definition.

In the present specification, examples of the "halogen atom" include fluorine, chlorine, bromine and iodine.

In the present specification, examples of the "C₁₋₆ alkyl group" include methyl, ethyl, propyl, isopropyl, butyl, isobutyl, sec-butyl, tert-butyl, pentyl, isopentyl, neopentyl, 1-ethylpropyl, hexyl, isohexyl, 1,1-dimethylbutyl, 2,2-dimethylbutyl, 3,3-dimethylbutyl and 2-ethylbutyl.

In the present specification, examples of the "optionally halogenated C₁₋₆ alkyl group" include a C₁₋₆ alkyl group optionally having 1 to 7, preferably 1 to 5 halogen atoms. Specific examples thereof include methyl, chloromethyl,

difluoromethyl, trichloromethyl, trifluoromethyl, ethyl, 2-bromoethyl, 2,2,2-trifluoroethyl, tetrafluoroethyl, pentafluoroethyl, propyl, 2,2-difluoropropyl, 3,3,3-trifluoropropyl, isopropyl, butyl, 4,4,4-trifluorobutyl, 5 isobutyl, sec-butyl, tert-butyl, pentyl, isopentyl, neopentyl, 5,5,5-trifluoropentyl, hexyl and 6,6,6-trifluorohexyl.

In the present specification, examples of the "C₂₋₆ alkenyl group" include ethenyl, 1-propenyl, 2-propenyl, 2-methyl-1-propenyl, 1-butenyl, 2-butenyl, 3-butenyl, 3-methyl-2- 10 butenyl, 1-pentenyl, 2-pentenyl, 3-pentenyl, 4-pentenyl, 4-methyl-3-pentenyl, 1-hexenyl, 3-hexenyl and 5-hexenyl.

In the present specification, examples of the "C₂₋₆ alkynyl group" include ethynyl, 1-propynyl, 2-propynyl, 1-butynyl, 2-butynyl, 3-butynyl, 1-pentynyl, 2-pentynyl, 3- 15 pentynyl, 4-pentynyl, 1-hexynyl, 2-hexynyl, 3-hexynyl, 4-hexynyl, 5-hexynyl and 4-methyl-2-pentynyl.

In the present specification, examples of the "C₃₋₁₀ cycloalkyl group" include cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, cyclooctyl, bicyclo[2.2.1]heptyl, 20 bicyclo[2.2.2]octyl, bicyclo[3.2.1]octyl and adamantyl.

In the present specification, examples of the "optionally halogenated C₃₋₁₀ cycloalkyl group" include a C₃₋₁₀ cycloalkyl group optionally having 1 to 7, preferably 1 to 5 halogen atoms. Specific examples thereof include cyclopropyl, 2,2- 25 difluorocyclopropyl, 2,3-difluorocyclopropyl, cyclobutyl, difluorocyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl and cyclooctyl.

In the present specification, examples of the "C₃₋₁₀ cycloalkenyl group" include cyclopropenyl, cyclobutenyl, 30 cyclopentenyl, cyclohexenyl, cycloheptenyl and cyclooctenyl.

In the present specification, examples of the "C₆₋₁₄ aryl group" include phenyl, 1-naphthyl, 2-naphthyl, 1-anthryl, 2-anthryl and 9-anthryl.

In the present specification, examples of the "C₇₋₁₆ aralkyl group" include benzyl, phenethyl, naphthylmethyl and 35 aralkyl group

phenylpropyl.

[0074]

In the present specification, examples of the "C₁₋₆ alkoxy group" include methoxy, ethoxy, propoxy, isopropoxy, butoxy, isobutoxy, sec-butoxy, tert-butoxy, pentyloxy and hexyloxy.

In the present specification, examples of the "optionally halogenated C₁₋₆ alkoxy group" include a C₁₋₆ alkoxy group optionally having 1 to 7, preferably 1 to 5 halogen atoms. Specific examples thereof include methoxy, difluoromethoxy, trifluoromethoxy, ethoxy, 2,2,2-trifluoroethoxy, propoxy, isopropoxy, butoxy, 4,4,4-trifluorobutoxy, isobutoxy, sec-butoxy, pentyloxy and hexyloxy.

In the present specification, examples of the "C₃₋₁₀ cycloalkyloxy group" include cyclopropyloxy, cyclobutyloxy, cyclopentyloxy, cyclohexyloxy, cycloheptyloxy and cyclooctyloxy.

In the present specification, examples of the "C₁₋₆ alkylthio group" include methylthio, ethylthio, propylthio, isopropylthio, butylthio, sec-butylthio, tert-butylthio, pentylthio and hexylthio.

In the present specification, examples of the "optionally halogenated C₁₋₆ alkylthio group" include a C₁₋₆ alkylthio group optionally having 1 to 7, preferably 1 to 5 halogen atoms. Specific examples thereof include methylthio, difluoromethylthio, trifluoromethylthio, ethylthio, propylthio, isopropylthio, butylthio, 4,4,4-trifluorobutylthio, pentylthio and hexylthio.

In the present specification, examples of the "C₁₋₆ alkyl-carbonyl group" include acetyl, propanoyl, butanoyl, 2-methylpropanoyl, pentanoyl, 3-methylbutanoyl, 2-methylbutanoyl, 2,2-dimethylpropanoyl, hexanoyl and heptanoyl.

In the present specification, examples of the "optionally halogenated C₁₋₆ alkyl-carbonyl group" include a C₁₋₆ alkyl-carbonyl group optionally having 1 to 7, preferably 1 to 5 halogen atoms. Specific examples thereof include acetyl, chloroacetyl, trifluoroacetyl, trichloroacetyl, propanoyl,

butanoyl, pentanoyl and hexanoyl.

In the present specification, examples of the "C₁₋₆ alkoxy-carbonyl group" include methoxycarbonyl, ethoxycarbonyl, propoxycarbonyl, isopropoxycarbonyl, butoxycarbonyl, 5 isobutoxycarbonyl, sec-butoxycarbonyl, tert-butoxycarbonyl, pentyloxycarbonyl and hexyloxycarbonyl.

In the present specification, examples of the "C₆₋₁₄ aryl-carbonyl group" include benzoyl, 1-naphthoyl and 2-naphthoyl.

In the present specification, examples of the "C₇₋₁₆ 10 aralkyl-carbonyl group" include phenylacetyl and phenylpropionyl.

In the present specification, examples of the "5- to 14-membered aromatic heterocyclylcarbonyl group" include nicotinoyl, isonicotinoyl, thenoyl and furoyl.

15 In the present specification, examples of the "3- to 14-membered non-aromatic heterocyclylcarbonyl group" include morpholinylcarbonyl, piperidinylcarbonyl and pyrrolidinylcarbonyl.

[0075]

20 In the present specification, examples of the "mono- or di-C₁₋₆ alkyl-carbamoyl group" include methylcarbamoyl, ethylcarbamoyl, dimethylcarbamoyl, diethylcarbamoyl and N-ethyl-N-methylcarbamoyl.

In the present specification, examples of the "mono- or 25 di-C₇₋₁₆ aralkyl-carbamoyl group" include benzylcarbamoyl and phenethylcarbamoyl.

In the present specification, examples of the "C₁₋₆ alkylsulfonyl group" include methylsulfonyl, ethylsulfonyl, propylsulfonyl, isopropylsulfonyl, butylsulfonyl, sec- 30 butylsulfonyl and tert-butylsulfonyl.

In the present specification, examples of the "optionally halogenated C₁₋₆ alkylsulfonyl group" include a C₁₋₆ alkylsulfonyl group optionally having 1 to 7, preferably 1 to 5 halogen atoms. Specific examples thereof include 35 methylsulfonyl, difluoromethylsulfonyl, trifluoromethylsulfonyl,

ethylsulfonyl, propylsulfonyl, isopropylsulfonyl, butylsulfonyl, 4,4,4-trifluorobutylsulfonyl, pentylsulfonyl and hexylsulfonyl.

In the present specification, examples of the "C₆₋₁₄ arylsulfonyl group" include phenylsulfonyl, 1-naphthylsulfonyl and 2-naphthylsulfonyl.

[0076]

In the present specification, examples of the "substituent" include a halogen atom, a cyano group, a nitro group, an optionally substituted hydrocarbon group, an optionally substituted heterocyclic group, an acyl group, an optionally substituted amino group, an optionally substituted carbamoyl group, an optionally substituted thiocarbamoyl group, an optionally substituted sulfamoyl group, an optionally substituted hydroxy group, an optionally substituted sulfanyl (SH) group and an optionally substituted silyl group.

In the present specification, examples of the "hydrocarbon group" (including "hydrocarbon group" of "optionally substituted hydrocarbon group") include a C₁₋₆ alkyl group, a C₂₋₆ alkenyl group, a C₂₋₆ alkynyl group, a C₃₋₁₀ cycloalkyl group, a C₃₋₁₀ cycloalkenyl group, a C₆₋₁₄ aryl group and a C₇₋₁₆ aralkyl group.

[0077]

In the present specification, examples of the "optionally substituted hydrocarbon group" include a hydrocarbon group optionally having substituent(s) selected from the following Substituent Group A.

[Substituent Group A]

- (1) a halogen atom,
- (2) a nitro group,
- 30 (3) a cyano group,
- (4) an oxo group,
- (5) a hydroxy group,
- (6) an optionally halogenated C₁₋₆ alkoxy group,
- (7) a C₆₋₁₄ aryloxy group (e.g., phenoxy, naphthoxy),
- 35 (8) a C₇₋₁₆ aralkyloxy group (e.g., benzyloxy),

- (9) a 5- to 14-membered aromatic heterocycloxy group (e.g., pyridyloxy),
- (10) a 3- to 14-membered non-aromatic heterocycloxy group (e.g., morpholinyl, piperidinyl),
- 5 (11) a C₁₋₆ alkyl-carbonyloxy group (e.g., acetoxy, propanoyloxy),
- (12) a C₆₋₁₄ aryl-carbonyloxy group (e.g., benzoyloxy, 1-naphthoyloxy, 2-naphthoyloxy),
- (13) a C₁₋₆ alkoxy-carbonyloxy group (e.g., methoxycarbonyloxy, 10 ethoxycarbonyloxy, propoxycarbonyloxy, butoxycarbonyloxy),
- (14) a mono- or di-C₁₋₆ alkyl-carbamoyloxy group (e.g., methylcarbamoyloxy, ethylcarbamoyloxy, dimethylcarbamoyloxy, diethylcarbamoyloxy),
- (15) a C₆₋₁₄ aryl-carbamoyloxy group (e.g., phenylcarbamoyloxy, 15 naphthylcarbamoyloxy),
- (16) a 5- to 14-membered aromatic heterocyclylcarbonyloxy group (e.g., nicotinoyloxy),
- (17) a 3- to 14-membered non-aromatic heterocyclylcarbonyloxy group (e.g., morpholinylcarbonyloxy, piperidinylcarbonyloxy),
- 20 (18) an optionally halogenated C₁₋₆ alkylsulfonyloxy group (e.g., methylsulfonyloxy, trifluoromethylsulfonyloxy),
- (19) a C₆₋₁₄ arylsulfonyloxy group optionally substituted by a C₁₋₆ alkyl group (e.g., phenylsulfonyloxy, toluenesulfonyloxy),
- (20) an optionally halogenated C₁₋₆ alkylthio group,
- 25 (21) a 5- to 14-membered aromatic heterocyclic group,
- (22) a 3- to 14-membered non-aromatic heterocyclic group,
- (23) a formyl group,
- (24) a carboxy group,
- (25) an optionally halogenated C₁₋₆ alkyl-carbonyl group,
- 30 (26) a C₆₋₁₄ aryl-carbonyl group,
- (27) a 5- to 14-membered aromatic heterocyclylcarbonyl group,
- (28) a 3- to 14-membered non-aromatic heterocyclylcarbonyl group,
- (29) a C₁₋₆ alkoxy-carbonyl group,
- 35 (30) a C₆₋₁₄ aryloxy-carbonyl group (e.g., phenyloxycarbonyl, 1-

- naphthyloxycarbonyl, 2-naphthyloxycarbonyl),
(31) a C₇₋₁₆ aralkyloxy-carbonyl group (e.g., benzyloxycarbonyl, phenethyloxycarbonyl),
(32) a carbamoyl group,
5 (33) a thiocarbamoyl group,
(34) a mono- or di-C₁₋₆ alkyl-carbamoyl group,
(35) a C₆₋₁₄ aryl-carbamoyl group (e.g., phenylcarbamoyl),
(36) a 5- to 14-membered aromatic heterocyclylcarbamoyl group (e.g., pyridylcarbamoyl, thienylcarbamoyl),
10 (37) a 3- to 14-membered non-aromatic heterocyclylcarbamoyl group (e.g., morpholinylcarbamoyl, piperidinylcarbamoyl),
(38) an optionally halogenated C₁₋₆ alkylsulfonyl group,
(39) a C₆₋₁₄ arylsulfonyl group,
(40) a 5- to 14-membered aromatic heterocyclylsulfonyl group
15 (e.g., pyridylsulfonyl, thienylsulfonyl),
(41) an optionally halogenated C₁₋₆ alkylsulfinyl group,
(42) a C₆₋₁₄ arylsulfinyl group (e.g., phenylsulfinyl, 1-naphthylsulfinyl, 2-naphthylsulfinyl),
(43) a 5- to 14-membered aromatic heterocyclylsulfinyl group
20 (e.g., pyridylsulfinyl, thienylsulfinyl),
(44) an amino group,
(45) a mono- or di-C₁₋₆ alkylamino group (e.g., methylamino, ethylamino, propylamino, isopropylamino, butylamino, dimethylamino, diethylamino, dipropylamino, dibutylamino, N-ethyl-N-methylamino),
25 (46) a mono- or di-C₆₋₁₄ arylamino group (e.g., phenylamino),
(47) a 5- to 14-membered aromatic heterocyclylamino group (e.g., pyridylamino),
(48) a C₇₋₁₆ aralkylamino group (e.g., benzylamino),
30 (49) a formylamino group,
(50) a C₁₋₆ alkyl-carbonylamino group (e.g., acetylamino, propanoylamino, butanoylamino),
(51) a (C₁₋₆ alkyl)(C₁₋₆ alkyl-carbonyl)amino group (e.g., N-acetyl-N-methylamino),
35 (52) a C₆₋₁₄ aryl-carbonylamino group (e.g., phenylcarbonylamino,

- naphthylcarbonylamino),
- (53) a C₁₋₆ alkoxy-carbonylamino group (e.g., methoxycarbonylamino, ethoxycarbonylamino, propoxycarbonylamino, butoxycarbonylamino, tert-butoxycarbonylamino),
- 5 (54) a C₇₋₁₆ aralkyloxy-carbonylamino group (e.g., benzyloxycarbonylamino),
- (55) a C₁₋₆ alkylsulfonylamino group (e.g., methylsulfonylamino, ethylsulfonylamino),
- (56) a C₆₋₁₄ arylsulfonylamino group optionally substituted by a
- 10 C₁₋₆ alkyl group (e.g., phenylsulfonylamino, toluenesulfonylamino),
- (57) an optionally halogenated C₁₋₆ alkyl group,
- (58) a C₂₋₆ alkenyl group,
- (59) a C₂₋₆ alkynyl group,
- 15 (60) a C₃₋₁₀ cycloalkyl group,
- (61) a C₃₋₁₀ cycloalkenyl group and
- (62) a C₆₋₁₄ aryl group.
- [0078]

The number of the above-mentioned substituents in the

20 "optionally substituted hydrocarbon group" is, for example, 1 to 5, preferably 1 to 3. When the number of the substituents is two or more, the respective substituents may be the same or different.

In the present specification, examples of the

25 "heterocyclic group" (including "heterocyclic group" of "optionally substituted heterocyclic group") include (i) an aromatic heterocyclic group, (ii) a non-aromatic heterocyclic group and (iii) a 7- to 10-membered bridged heterocyclic group, each containing, as a ring-constituting atom besides carbon

30 atom, 1 to 4 hetero atoms selected from a nitrogen atom, a sulfur atom and an oxygen atom.

[0079]

In the present specification, examples of the "aromatic heterocyclic group" (including "5- to 14-membered aromatic

35 heterocyclic group") include a 5- to 14-membered (preferably 5-

to 10-membered) aromatic heterocyclic group containing, as a ring-constituting atom besides carbon atom, 1 to 4 hetero atoms selected from a nitrogen atom, a sulfur atom and an oxygen atom.

Preferable examples of the "aromatic heterocyclic group" include 5- or 6-membered monocyclic aromatic heterocyclic groups such as thienyl, furyl, pyrrolyl, imidazolyl, pyrazolyl, thiazolyl, isothiazolyl, oxazolyl, isoxazolyl, pyridyl, pyrazinyl, pyrimidinyl, pyridazinyl, 1,2,4-oxadiazolyl, 1,3,4-oxadiazolyl, 1,2,4-thiadiazolyl, 1,3,4-thiadiazolyl, triazolyl, tetrazolyl, triazinyl and the like; and 8- to 14-membered fused polycyclic (preferably bi or tricyclic) aromatic heterocyclic groups such as benzothiophenyl, benzofuranyl, benzimidazolyl, benzoxazolyl, benzisoxazolyl, benzothiazolyl, benzisothiazolyl, benzotriazolyl, imidazopyridinyl, thienopyridinyl, furopyridinyl, pyrrolopyridinyl, pyrazolopyridinyl, oxazolopyridinyl, thiazolopyridinyl, imidazopyrazinyl, imidazopyrimidinyl, thienopyrimidinyl, furopyrimidinyl, pyrrolopyrimidinyl, pyrazolopyrimidinyl, oxazolopyrimidinyl, thiazolopyrimidinyl, pyrazolotriazinyl, naphtho[2,3-b]thienyl, phenoxathiinyl, indolyl, isoindolyl, 1H-indazolyl, purinyl, isoquinolyl, quinolyl, phthalazinyl, naphthyridinyl, quinoxalinyl, quinazolinyl, cinnolinyl, carbazolyl, β -carbolinyl, phenanthridinyl, acridinyl, phenazinyl, phenothiazinyl, phenoxazinyl and the like.

[0080]

In the present specification, examples of the "non-aromatic heterocyclic group" (including "3- to 14-membered non-aromatic heterocyclic group") include a 3- to 14-membered (preferably 4- to 10-membered) non-aromatic heterocyclic group containing, as a ring-constituting atom besides carbon atom, 1 to 4 hetero atoms selected from a nitrogen atom, a sulfur atom and an oxygen atom.

Preferable examples of the "non-aromatic heterocyclic group" include 3- to 8-membered monocyclic non-aromatic

heterocyclic groups such as aziridinyl, oxiranyl, thiiranyl, azetidiny, oxetanyl, thietanyl, tetrahydrothienyl, tetrahydrofuranyl, pyrrolinyl, pyrrolidinyl, imidazolinyl, imidazolidinyl, oxazolinyl, oxazolidinyl, pyrazolinyl, 5 pyrazolidinyl, thiazolinyl, thiazolidinyl, tetrahydroisothiazolyl, tetrahydrooxazolyl, tetrahydroisooxazolyl, piperidinyl, piperazinyl, tetrahydropyridinyl, dihydropyridinyl, dihydrothiopyranyl, tetrahydropyrimidinyl, tetrahydropyridazinyl, dihydropyranyl, 10 tetrahydropyranyl, tetrahydrothiopyranyl, morpholinyl, thiomorpholinyl, azepanyl, diazepanyl, azepinyl, oxepanyl, azocanyl, diazocanyl and the like; and 9- to 14-membered fused polycyclic (preferably bi or tricyclic) non-aromatic heterocyclic groups such as dihydrobenzofuranyl, 15 dihydrobenzimidazolyl, dihydrobenzoxazolyl, dihydrobenzothiazolyl, dihydrobenzisoctiazolyl, dihydronaphtho[2,3-b]thienyl, tetrahydroisoquinolyl, tetrahydroquinolyl, 4H-quinoliziny, indoliny, isoindoliny, tetrahydrothieno[2,3-c]pyridiny, tetrahydrobenzazepiny, 20 tetrahydroquinoxaliny, tetrahydrophenanthridiny, hexahydrophenothiaziny, hexahydrophenoxaziny, tetrahydrophthalaziny, tetrahydronaphthyridiny, tetrahydroquinazoliny, tetrahydrocinnoliny, tetrahydrocarbazolyl, tetrahydro- β -carboliny, 25 tetrahydroacrydiny, tetrahydrophenaziny, tetrahydrothioxantheny, octahydroisoquinolyl and the like.

[0081]

In the present specification, preferable examples of the "7- to 10-membered bridged heterocyclic group" include 30 quinuclidiny and 7-azabicyclo[2.2.1]heptanyl.

In the present specification, examples of the "nitrogen-containing heterocyclic group" include a "heterocyclic group" containing at least one nitrogen atom as a ring-constituting atom.

35 In the present specification, examples of the "optionally

substituted heterocyclic group" include a heterocyclic group optionally having substituent(s) selected from the above-mentioned Substituent Group A.

The number of the substituents in the "optionally substituted heterocyclic group" is, for example, 1 to 3. When the number of the substituents is two or more, the respective substituents may be the same or different.

[0082]

In the present specification, examples of the "acyl group" include a formyl group, a carboxy group, a carbamoyl group, a thiocarbamoyl group, a sulfinio group, a sulfo group, a sulfamoyl group and a phosphono group, each optionally having "1 or 2 substituents selected from a C₁₋₆ alkyl group, a C₂₋₆ alkenyl group, a C₃₋₁₀ cycloalkyl group, a C₃₋₁₀ cycloalkenyl group, a C₆₋₁₄ aryl group, a C₇₋₁₆ aralkyl group, a 5- to 14-membered aromatic heterocyclic group and a 3- to 14-membered non-aromatic heterocyclic group, each of which optionally has 1 to 3 substituents selected from a halogen atom, an optionally halogenated C₁₋₆ alkoxy group, a hydroxy group, a nitro group, a cyano group, an amino group and a carbamoyl group".

Examples of the "acyl group" also include a hydrocarbon-sulfonyl group, a heterocyclylsulfonyl group, a hydrocarbon-sulfinyl group and a heterocyclylsulfinyl group.

Here, the hydrocarbon-sulfonyl group means a hydrocarbon group-bonded sulfonyl group, the heterocyclylsulfonyl group means a heterocyclic group-bonded sulfonyl group, the hydrocarbon-sulfinyl group means a hydrocarbon group-bonded sulfinyl group and the heterocyclylsulfinyl group means a heterocyclic group-bonded sulfinyl group.

Preferable examples of the "acyl group" include a formyl group, a carboxy group, a C₁₋₆ alkyl-carbonyl group, a C₂₋₆ alkenyl-carbonyl group (e.g., crotonoyl), a C₃₋₁₀ cycloalkyl-carbonyl group (e.g., cyclobutanecarbonyl, cyclopentanecarbonyl, cyclohexanecarbonyl, cycloheptanecarbonyl), a C₃₋₁₀ cycloalkenyl-carbonyl group (e.g., 2-cyclohexenecarbonyl), a

C₆₋₁₄ aryl-carbonyl group, a C₇₋₁₆ aralkyl-carbonyl group, a 5- to 14-membered aromatic heterocyclylcarbonyl group, a 3- to 14-membered non-aromatic heterocyclylcarbonyl group, a C₁₋₆ alkoxy-carbonyl group, a C₆₋₁₄ aryloxy-carbonyl group (e.g., phenyloxycarbonyl, naphthyloxycarbonyl), a C₇₋₁₆ aralkyloxy-carbonyl group (e.g., benzyloxycarbonyl, phenethyloxycarbonyl), a carbamoyl group, a mono- or di-C₁₋₆ alkyl-carbamoyl group, a mono- or di-C₂₋₆ alkenyl-carbamoyl group (e.g., diallylcarbamoyl), a mono- or di-C₃₋₁₀ cycloalkyl-carbamoyl group (e.g., cyclopropylcarbamoyl), a mono- or di-C₆₋₁₄ aryl-carbamoyl group (e.g., phenylcarbamoyl), a mono- or di-C₇₋₁₆ aralkyl-carbamoyl group, a 5- to 14-membered aromatic heterocyclylcarbamoyl group (e.g., pyridylcarbamoyl), a thiocarbamoyl group, a mono- or di-C₁₋₆ alkyl-thiocarbamoyl group (e.g., methylthiocarbamoyl, N-ethyl-N-methylthiocarbamoyl), a mono- or di-C₂₋₆ alkenyl-thiocarbamoyl group (e.g., diallylthiocarbamoyl), a mono- or di-C₃₋₁₀ cycloalkyl-thiocarbamoyl group (e.g., cyclopropylthiocarbamoyl, cyclohexylthiocarbamoyl), a mono- or di-C₆₋₁₄ aryl-thiocarbamoyl group (e.g., phenylthiocarbamoyl), a mono- or di-C₇₋₁₆ aralkyl-thiocarbamoyl group (e.g., benzylthiocarbamoyl, phenethylthiocarbamoyl), a 5- to 14-membered aromatic heterocyclylthiocarbamoyl group (e.g., pyridylthiocarbamoyl), a sulfinio group, a C₁₋₆ alkylsulfinyl group (e.g., methylsulfinyl, ethylsulfinyl), a sulfo group, a C₁₋₆ alkylsulfonyl group, a C₆₋₁₄ arylsulfonyl group, a phosphono group and a mono- or di-C₁₋₆ alkylphosphono group (e.g., dimethylphosphono, diethylphosphono, diisopropylphosphono, dibutylphosphono).

[0083]

In the present specification, examples of the "optionally substituted amino group" include an amino group optionally having "1 or 2 substituents selected from a C₁₋₆ alkyl group, a C₂₋₆ alkenyl group, a C₃₋₁₀ cycloalkyl group, a C₆₋₁₄ aryl group, a C₇₋₁₆ aralkyl group, a C₁₋₆ alkyl-carbonyl group, a C₆₋₁₄ aryl-carbonyl group, a C₇₋₁₆ aralkyl-carbonyl group, a 5- to 14-

membered aromatic heterocyclylcarbonyl group, a 3- to 14-membered non-aromatic heterocyclylcarbonyl group, a C₁₋₆ alkoxy-carbonyl group, a 5- to 14-membered aromatic heterocyclic group, a carbamoyl group, a mono- or di-C₁₋₆ alkyl-carbamoyl group, a
5 mono- or di-C₇₋₁₆ aralkyl-carbamoyl group, a C₁₋₆ alkylsulfonyl group and a C₆₋₁₄ arylsulfonyl group, each of which optionally has 1 to 3 substituents selected from Substituent Group A".

Preferable examples of the optionally substituted amino group include an amino group, a mono- or di-(optionally
10 halogenated C₁₋₆ alkyl)amino group (e.g., methylamino, trifluoromethylamino, dimethylamino, ethylamino, diethylamino, propylamino, dibutylamino), a mono- or di-C₂₋₆ alkenylamino group (e.g., diallylamino), a mono- or di-C₃₋₁₀ cycloalkylamino group (e.g., cyclopropylamino, cyclohexylamino), a mono- or di-
15 C₆₋₁₄ arylamino group (e.g., phenylamino), a mono- or di-C₇₋₁₆ aralkylamino group (e.g., benzylamino, dibenzylamino), a mono- or di-(optionally halogenated C₁₋₆ alkyl)-carbonylamino group (e.g., acetylamino, propionylamino), a mono- or di-C₆₋₁₄ aryl-carbonylamino group (e.g., benzoylamino), a mono- or di-C₇₋₁₆
20 aralkyl-carbonylamino group (e.g., benzylcarbonylamino), a mono- or di-5- to 14-membered aromatic heterocyclylcarbonylamino group (e.g., nicotinoylamino, isonicotinoylamino), a mono- or di-3- to 14-membered non-aromatic heterocyclylcarbonylamino group (e.g.,
25 piperidiny carbonylamino), a mono- or di-C₁₋₆ alkoxy-carbonylamino group (e.g., tert-butoxycarbonylamino), a 5- to 14-membered aromatic heterocyclylamino group (e.g., pyridylamino), a carbamoylamino group, a (mono- or di-C₁₋₆ alkyl-carbamoyl)amino group (e.g., methylcarbamoylamino), a
30 (mono- or di-C₇₋₁₆ aralkyl-carbamoyl)amino group (e.g., benzylcarbamoylamino), a C₁₋₆ alkylsulfonylamino group (e.g., methylsulfonylamino, ethylsulfonylamino), a C₆₋₁₄ arylsulfonylamino group (e.g., phenylsulfonylamino), a (C₁₋₆ alkyl)(C₁₋₆ alkyl-carbonyl)amino group (e.g., N-acetyl-N-
35 methylamino) and a (C₁₋₆ alkyl)(C₆₋₁₄ aryl-carbonyl)amino group

(e.g., N-benzoyl-N-methylamino).

[0084]

In the present specification, examples of the "optionally substituted carbamoyl group" include a carbamoyl group
5 optionally having "1 or 2 substituents selected from a C₁₋₆ alkyl group, a C₂₋₆ alkenyl group, a C₃₋₁₀ cycloalkyl group, a C₆₋₁₄ aryl group, a C₇₋₁₆ aralkyl group, a C₁₋₆ alkyl-carbonyl group, a C₆₋₁₄ aryl-carbonyl group, a C₇₋₁₆ aralkyl-carbonyl group, a 5- to 14-membered aromatic heterocyclylcarbonyl group, a 3- to 14-
10 membered non-aromatic heterocyclylcarbonyl group, a C₁₋₆ alkoxy-carbonyl group, a 5- to 14-membered aromatic heterocyclic group, a carbamoyl group, a mono- or di-C₁₋₆ alkyl-carbamoyl group and a mono- or di-C₇₋₁₆ aralkyl-carbamoyl group, each of which optionally has 1 to 3 substituents selected from Substituent
15 Group A".

Preferable examples of the optionally substituted carbamoyl group include a carbamoyl group, a mono- or di-C₁₋₆ alkyl-carbamoyl group, a mono- or di-C₂₋₆ alkenyl-carbamoyl group (e.g., diallylcarbamoyl), a mono- or di-C₃₋₁₀ cycloalkyl-
20 carbamoyl group (e.g., cyclopropylcarbamoyl, cyclohexylcarbamoyl), a mono- or di-C₆₋₁₄ aryl-carbamoyl group (e.g., phenylcarbamoyl), a mono- or di-C₇₋₁₆ aralkyl-carbamoyl group, a mono- or di-C₁₋₆ alkyl-carbonyl-carbamoyl group (e.g., acetylcarbamoyl, propionylcarbamoyl), a mono- or di-C₆₋₁₄ aryl-
25 carbonyl-carbamoyl group (e.g., benzoylcarbamoyl) and a 5- to 14-membered aromatic heterocyclylcarbamoyl group (e.g., pyridylcarbamoyl).

[0085]

In the present specification, examples of the "optionally substituted thiocarbamoyl group" include a thiocarbamoyl group
30 optionally having "1 or 2 substituents selected from a C₁₋₆ alkyl group, a C₂₋₆ alkenyl group, a C₃₋₁₀ cycloalkyl group, a C₆₋₁₄ aryl group, a C₇₋₁₆ aralkyl group, a C₁₋₆ alkyl-carbonyl group, a C₆₋₁₄ aryl-carbonyl group, a C₇₋₁₆ aralkyl-carbonyl group, a 5- to 14-membered aromatic heterocyclylcarbonyl group, a 3- to 14-
35

membered non-aromatic heterocyclylcarbonyl group, a C₁₋₆ alkoxy-carbonyl group, a 5- to 14-membered aromatic heterocyclic group, a carbamoyl group, a mono- or di-C₁₋₆ alkyl-carbamoyl group and a mono- or di-C₇₋₁₆ aralkyl-carbamoyl group, each of which
5 optionally has 1 to 3 substituents selected from Substituent Group A".

Preferable examples of the optionally substituted thiocarbamoyl group include a thiocarbamoyl group, a mono- or di-C₁₋₆ alkyl-thiocarbamoyl group (e.g., methylthiocarbamoyl, ethylthiocarbamoyl, dimethylthiocarbamoyl, diethylthiocarbamoyl, N-ethyl-N-methylthiocarbamoyl), a mono- or di-C₂₋₆ alkenyl-thiocarbamoyl group (e.g., diallylthiocarbamoyl), a mono- or di-C₃₋₁₀ cycloalkyl-thiocarbamoyl group (e.g., cyclopropylthiocarbamoyl, cyclohexylthiocarbamoyl), a mono- or di-C₆₋₁₄ aryl-thiocarbamoyl group (e.g., phenylthiocarbamoyl), a mono- or di-C₇₋₁₆ aralkyl-thiocarbamoyl group (e.g., benzylthiocarbamoyl, phenethylthiocarbamoyl), a mono- or di-C₁₋₆ alkyl-carbonyl-thiocarbamoyl group (e.g., acetylthiocarbamoyl, propionylthiocarbamoyl), a mono- or di-C₆₋₁₄ aryl-carbonyl-thiocarbamoyl group (e.g., benzoylthiocarbamoyl) and a 5- to 14-membered aromatic heterocyclylthiocarbamoyl group (e.g., pyridylthiocarbamoyl).

[0086]

In the present specification, examples of the "optionally substituted sulfamoyl group" include a sulfamoyl group
25 optionally having "1 or 2 substituents selected from a C₁₋₆ alkyl group, a C₂₋₆ alkenyl group, a C₃₋₁₀ cycloalkyl group, a C₆₋₁₄ aryl group, a C₇₋₁₆ aralkyl group, a C₁₋₆ alkyl-carbonyl group, a C₆₋₁₄ aryl-carbonyl group, a C₇₋₁₆ aralkyl-carbonyl group, a 5- to 14-membered aromatic heterocyclylcarbonyl group, a 3- to 14-membered non-aromatic heterocyclylcarbonyl group, a C₁₋₆ alkoxy-carbonyl group, a 5- to 14-membered aromatic heterocyclic group, a carbamoyl group, a mono- or di-C₁₋₆ alkyl-carbamoyl group and a mono- or di-C₇₋₁₆ aralkyl-carbamoyl group, each of which
35 optionally has 1 to 3 substituents selected from Substituent

Group A".

Preferable examples of the optionally substituted sulfamoyl group include a sulfamoyl group, a mono- or di-C₁₋₆ alkyl-sulfamoyl group (e.g., methylsulfamoyl, ethylsulfamoyl, dimethylsulfamoyl, diethylsulfamoyl, N-ethyl-N-methylsulfamoyl), a mono- or di-C₂₋₆ alkenyl-sulfamoyl group (e.g., diallylsulfamoyl), a mono- or di-C₃₋₁₀ cycloalkyl-sulfamoyl group (e.g., cyclopropylsulfamoyl, cyclohexylsulfamoyl), a mono- or di-C₆₋₁₄ aryl-sulfamoyl group (e.g., phenylsulfamoyl), a mono- or di-C₇₋₁₆ aralkyl-sulfamoyl group (e.g., benzylsulfamoyl, phenethylsulfamoyl), a mono- or di-C₁₋₆ alkyl-carbonyl-sulfamoyl group (e.g., acetylsulfamoyl, propionylsulfamoyl), a mono- or di-C₆₋₁₄ aryl-carbonyl-sulfamoyl group (e.g., benzoylsulfamoyl) and a 5- to 14-membered aromatic heterocyclylsulfamoyl group (e.g., pyridylsulfamoyl).

[0087]

In the present specification, examples of the "optionally substituted hydroxy group" include a hydroxyl group optionally having "a substituent selected from a C₁₋₆ alkyl group, a C₂₋₆ alkenyl group, a C₃₋₁₀ cycloalkyl group, a C₆₋₁₄ aryl group, a C₇₋₁₆ aralkyl group, a C₁₋₆ alkyl-carbonyl group, a C₆₋₁₄ aryl-carbonyl group, a C₇₋₁₆ aralkyl-carbonyl group, a 5- to 14-membered aromatic heterocyclylcarbonyl group, a 3- to 14-membered non-aromatic heterocyclylcarbonyl group, a C₁₋₆ alkoxy-carbonyl group, a 5- to 14-membered aromatic heterocyclic group, a carbamoyl group, a mono- or di-C₁₋₆ alkyl-carbamoyl group, a mono- or di-C₇₋₁₆ aralkyl-carbamoyl group, a C₁₋₆ alkylsulfonyl group and a C₆₋₁₄ arylsulfonyl group, each of which optionally has 1 to 3 substituents selected from Substituent Group A".

Preferable examples of the optionally substituted hydroxy group include a hydroxy group, a C₁₋₆ alkoxy group, a C₂₋₆ alkenyloxy group (e.g., allyloxy, 2-butenyloxy, 2-pentenylloxy, 3-hexenyloxy), a C₃₋₁₀ cycloalkyloxy group (e.g., cyclohexyloxy), a C₆₋₁₄ aryloxy group (e.g., phenoxy, naphthyloxy), a C₇₋₁₆ aralkyloxy group (e.g., benzyloxy, phenethylloxy), a C₁₋₆ alkyl-

carbonyloxy group (e.g., acetyloxy, propionyloxy, butyryloxy, isobutyryloxy, pivaloyloxy), a C₆₋₁₄ aryl-carbonyloxy group (e.g., benzoyloxy), a C₇₋₁₆ aralkyl-carbonyloxy group (e.g., benzylcarbonyloxy), a 5- to 14-membered aromatic
5 heterocyclylcarbonyloxy group (e.g., nicotinoyloxy), a 3- to 14-membered non-aromatic heterocyclylcarbonyloxy group (e.g., piperidinylcarbonyloxy), a C₁₋₆ alkoxy-carbonyloxy group (e.g., tert-butoxycarbonyloxy), a 5- to 14-membered aromatic
heterocyclyloxy group (e.g., pyridyloxy), a carbamoyloxy group,
10 a C₁₋₆ alkyl-carbamoyloxy group (e.g., methylcarbamoyloxy), a C₇₋₁₆ aralkyl-carbamoyloxy group (e.g., benzylcarbamoyloxy), a C₁₋₆ alkylsulfonyloxy group (e.g., methylsulfonyloxy, ethylsulfonyloxy) and a C₆₋₁₄ arylsulfonyloxy group (e.g., phenylsulfonyloxy).

15 [0088]

In the present specification, examples of the "optionally substituted sulfanyl group" include a sulfanyl group optionally having "a substituent selected from a C₁₋₆ alkyl group, a C₂₋₆ alkenyl group, a C₃₋₁₀ cycloalkyl group, a C₆₋₁₄ aryl group, a C₇₋₁₆ aralkyl group, a C₁₋₆ alkyl-carbonyl group, a C₆₋₁₄ aryl-carbonyl group and a 5- to 14-membered aromatic heterocyclic group, each of which optionally has 1 to 3 substituents
20 selected from Substituent Group A" and a halogenated sulfanyl group.

25 Preferable examples of the optionally substituted sulfanyl group include a sulfanyl (-SH) group, a C₁₋₆ alkylthio group, a C₂₋₆ alkenylthio group (e.g., allylthio, 2-butenylthio, 2-pentenylthio, 3-hexenylthio), a C₃₋₁₀ cycloalkylthio group (e.g., cyclohexylthio), a C₆₋₁₄ arylthio group (e.g., phenylthio, naphthylthio), a C₇₋₁₆ aralkylthio group (e.g., benzylthio, phenethylthio), a C₁₋₆ alkyl-carbonylthio group (e.g., acetylthio, propionylthio, butyrylthio, isobutyrylthio, pivaloylthio), a C₆₋₁₄ aryl-carbonylthio group (e.g., benzoylthio), a 5- to 14-membered aromatic heterocyclylthio
30 group (e.g., pyridylthio) and a halogenated thio group (e.g.,

pentafluorothio).

[0089]

In the present specification, examples of the "optionally substituted silyl group" include a silyl group optionally
5 having "1 to 3 substituents selected from a C₁₋₆ alkyl group, a C₂₋₆ alkenyl group, a C₃₋₁₀ cycloalkyl group, a C₆₋₁₄ aryl group and a C₇₋₁₆ aralkyl group, each of which optionally has 1 to 3 substituents selected from Substituent Group A".

Preferable examples of the optionally substituted silyl
10 group include a tri-C₁₋₆ alkylsilyl group (e.g., trimethylsilyl, tert-butyl(dimethyl)silyl).

[0090]

Each symbol in formula (I) is explained below.

[0091]

15 R¹ is an optionally substituted 5- or 6-membered cyclic group or an optionally substituted C₁₋₆ alkyl group.

[0092]

Examples of the "5- or 6-membered cyclic group" of the "optionally substituted 5- or 6-membered cyclic group" for R¹
20 include a phenyl group, a C₅₋₆ cycloalkyl group (cyclopentyl, cyclohexyl), a C₅₋₆ cycloalkenyl group (cyclopentenyl, cyclohexenyl), a 5- or 6-membered monocyclic aromatic heterocyclic group, a 5- or 6-membered monocyclic non-aromatic heterocyclic group and the like.

25 [0093]

Examples of the "5- or 6-membered monocyclic aromatic heterocyclic group" exemplified as the "5- or 6-membered cyclic group" include a 5- or 6-membered monocyclic aromatic heterocyclic group containing, as a ring-constituting atom
30 besides carbon atom, 1 to 4 heteroatoms selected from a nitrogen atom, a sulfur atom and an oxygen atom, and specific examples thereof include those exemplified as the "5- to 6-membered monocyclic aromatic heterocyclic group", from among the above-mentioned preferable examples of the "aromatic
35 heterocyclic group".

[0094]

Examples of the "5- or 6-membered monocyclic non-aromatic heterocyclic group" exemplified as the "5- or 6-membered cyclic group" include a 5- or 6-membered monocyclic non-aromatic
5 heterocyclic group containing, as a ring-constituting atom besides carbon atom, 1 to 4 heteroatoms selected from a nitrogen atom, a sulfur atom and an oxygen atom, and specific examples thereof include 5- or 6-membered monocyclic non-aromatic heterocyclic groups such as tetrahydrothienyl,
10 tetrahydrofuranyl, pyrrolinyl, pyrrolidinyl, imidazolinyl, imidazolidinyl, oxazolinyl, oxazolidinyl, pyrazolinyl, pyrazolidinyl, thiazolinyl, thiazolidinyl, tetrahydroisothiazolyl, tetrahydrooxazolyl, tetrahydroisoxazolyl, piperidinyl, piperazinyl,
15 tetrahydropyridinyl, dihydropyridinyl, dihydrothiopyranyl, tetrahydropyrimidinyl, tetrahydropyridazinyl, dihydropyranyl, tetrahydropyranyl, tetrahydrothiopyranyl, morpholinyl, thiomorpholinyl and the like.

[0095]

20 The "C₁₋₆ alkyl group" of the "optionally substituted C₁₋₆ alkyl group" and the "5- or 6-membered cyclic group" of the "optionally substituted 5- or 6-membered cyclic group" for R¹ each optionally has 1 to 5 (preferably 1 to 3) substituents at substitutable position(s). Examples of the substituent include
25 the above-mentioned Substituent Group A. When the number of the substituents is plural, the respective substituents may be the same or different.

[0096]

R¹ is preferably

- 30 (1) an optionally substituted phenyl group,
(2) an optionally substituted C₅₋₆ cycloalkyl group,
(3) an optionally substituted 5- or 6-membered non-aromatic heterocyclic group,
(4) an optionally substituted 5- or 6-membered monocyclic
35 aromatic heterocyclic group, or

(5) an optionally substituted C₁₋₆ alkyl group.

[0097]

R¹ is more preferably

(1) a phenyl group, a C₅₋₆ cycloalkyl group (cyclopentyl,
5 cyclohexyl), a 5- or 6-membered monocyclic non-aromatic
heterocyclic group (e.g., tetrahydropyranyl) or a 5- or 6-
membered monocyclic aromatic heterocyclic group (e.g., pyridyl),
each of which is optionally substituted by 1 to 3 substituents
selected from

10 (i) a halogen atom (e.g., a fluorine atom, a chlorine atom),
(ii) a cyano group,
(iii) a hydroxy group,
(iv) a C₁₋₆ alkyl group (e.g., methyl) optionally substituted
by 1 to 3 hydroxy groups, and

15 (v) a C₁₋₆ alkoxy group (e.g., methoxy), or
(2) a C₁₋₆ alkyl group (e.g., methyl, isobutyl, 1,2-
dimethylpropyl) optionally substituted by 1 to 3 substituents
selected from

(i) a hydroxy group, and
20 (ii) a 3- to 14-membered non-aromatic heterocyclic group
(preferably a 3- to 8-membered monocyclic non-aromatic
heterocyclic group) (e.g., tetrahydrofuryl).

[0098]

R¹ is further more preferably

25 (1) a phenyl group optionally substituted by 1 to 3
substituents selected from

(i) a halogen atom (e.g., a fluorine atom), and
(ii) a cyano group,

(2) a C₅₋₆ cycloalkyl group (cyclopentyl, cyclohexyl) optionally
30 substituted by 1 to 3 substituents selected from

(i) a hydroxy group,
(ii) a C₁₋₆ alkyl group (e.g., methyl) optionally substituted
by 1 to 3 hydroxy groups, and
(iii) a C₁₋₆ alkoxy group (e.g., methoxy),

35 (3) a 5- or 6-membered monocyclic non-aromatic heterocyclic

group (e.g., tetrahydropyranyl) optionally substituted by 1 to 3 hydroxy groups,

(4) a 5- or 6-membered monocyclic aromatic heterocyclic group (e.g., pyridyl) optionally substituted by 1 to 3 halogen atoms (e.g., a fluorine atom, a chlorine atom), or

(5) a C₁₋₆ alkyl group (e.g., methyl, isobutyl, 1,2-dimethylpropyl) optionally substituted by 1 to 3 substituents selected from

- (i) a hydroxy group, and
- (ii) a 3- to 8-membered monocyclic non-aromatic heterocyclic group (e.g., tetrahydrofuryl).

[0099]

R¹ is still more preferably

(1) a phenyl group optionally substituted by 1 to 3 substituents selected from

- (i) a halogen atom (e.g., a fluorine atom), and
 - (ii) a cyano group,
- (2) a C₅₋₆ cycloalkyl group (cyclopentyl, cyclohexyl) optionally substituted by 1 to 3 substituents selected from

- (i) a hydroxy group,
- (ii) a C₁₋₆ alkyl group (e.g., methyl) optionally substituted by 1 to 3 hydroxy groups, and
- (iii) a C₁₋₆ alkoxy group (e.g., methoxy),

(3) a tetrahydropyranyl group optionally substituted by 1 to 3 hydroxy groups,

- (4) a pyridyl group optionally substituted by 1 to 3 halogen atoms (e.g., a fluorine atom, a chlorine atom), or
- (5) a C₁₋₆ alkyl group (e.g., methyl, isobutyl, 1,2-dimethylpropyl) optionally substituted by 1 to 3 substituents selected from

- (i) a hydroxy group, and
- (ii) a tetrahydrofuryl group.

[0100]

In another embodiment, R¹ is more preferably

(1) a phenyl group, a C₅₋₆ cycloalkyl group (cyclopentyl,

cyclohexyl), a 5- or 6-membered monocyclic non-aromatic heterocyclic group (e.g., tetrahydropyranyl) or a 5- or 6-membered monocyclic aromatic heterocyclic group (e.g., pyridyl), each of which is optionally substituted by 1 to 3 substituents
5 selected from

- (i) a halogen atom (e.g., a fluorine atom, a chlorine atom),
 - (ii) a cyano group, and
 - (iii) a hydroxy group, or
- (2) a C₁₋₆ alkyl group (e.g., methyl, isobutyl, 1,2-dimethylpropyl) optionally substituted by 1 to 3 substituents
10 selected from
- (i) a hydroxy group, and
 - (ii) a 3- to 14-membered non-aromatic heterocyclic group (preferably a 3- to 8-membered monocyclic non-aromatic
15 heterocyclic group) (e.g., tetrahydrofuryl).

[0101]

In this embodiment, R¹ is further more preferably

- (1) a phenyl group optionally substituted by 1 to 3 substituents selected from
- 20 (i) a halogen atom (e.g., a fluorine atom), and
 - (ii) a cyano group,
- (2) a C₅₋₆ cycloalkyl group (cyclopentyl, cyclohexyl) optionally substituted by 1 to 3 hydroxy groups,
- (3) a 5- or 6-membered monocyclic non-aromatic heterocyclic
25 group (e.g., tetrahydropyranyl) optionally substituted by 1 to 3 hydroxy groups,
- (4) a 5- or 6-membered monocyclic aromatic heterocyclic group (e.g., pyridyl) optionally substituted by 1 to 3 halogen atoms (e.g., a fluorine atom, a chlorine atom), or
- 30 (5) a C₁₋₆ alkyl group (e.g., methyl, isobutyl, 1,2-dimethylpropyl) optionally substituted by 1 to 3 substituents selected from
- (i) a hydroxy group, and
 - (ii) a 3- to 8-membered monocyclic non-aromatic
35 heterocyclic group (e.g., tetrahydrofuryl).

[0102]

In this embodiment, R^1 is still more preferably

(1) a phenyl group optionally substituted by 1 to 3 substituents selected from

5 (i) a halogen atom (e.g., a fluorine atom), and

(ii) a cyano group,

(2) a C_{5-6} cycloalkyl group (cyclopentyl, cyclohexyl) optionally substituted by 1 to 3 hydroxy groups,

(3) a tetrahydropyranyl group optionally substituted by 1 to 3
10 hydroxy groups,

(4) a pyridyl group optionally substituted by 1 to 3 halogen atoms (e.g., a fluorine atom, a chlorine atom), or

(5) a C_{1-6} alkyl group (e.g., methyl, isobutyl, 1,2-dimethylpropyl) optionally substituted by 1 to 3 substituents
15 selected from

(i) a hydroxy group, and

(ii) a tetrahydrofuryl group.

[0103]

In another embodiment, R^1 is preferably

20 (1) an optionally substituted phenyl group,

(2) an optionally substituted C_{5-6} cycloalkyl group,

(3) an optionally substituted 5- or 6-membered non-aromatic heterocyclic group, or

(4) an optionally substituted C_{1-6} alkyl group.

25 [0104]

In this embodiment, R^1 is more preferably

(1) a phenyl group, a C_{5-6} cycloalkyl group (cyclopentyl, cyclohexyl) or a 5- or 6-membered monocyclic non-aromatic heterocyclic group (e.g., tetrahydropyranyl), each of which is

30 optionally substituted by 1 to 3 substituents selected from

(i) a halogen atom (e.g., a fluorine atom),

(ii) a cyano group, and

(iii) a hydroxy group, or

(2) a C_{1-6} alkyl group (e.g., methyl, isobutyl) optionally
35 substituted by 1 to 3 substituents selected from

- (i) a hydroxy group, and
- (ii) a 3- to 14-membered non-aromatic heterocyclic group (preferably a 3- to 8-membered monocyclic non-aromatic heterocyclic group) (e.g., tetrahydrofuryl).

5 [0105]

In this embodiment, R^1 is further more preferably

- (1) a phenyl group optionally substituted by 1 to 3 substituents selected from

- (i) a halogen atom (e.g., a fluorine atom), and
- 10 (ii) a cyano group,
- (2) a C_{5-6} cycloalkyl group (cyclopentyl, cyclohexyl) optionally substituted by 1 to 3 hydroxy groups,
- (3) a 5- or 6-membered monocyclic non-aromatic heterocyclic group (e.g., tetrahydropyranyl) optionally substituted by 1 to
- 15 3 hydroxy groups, or
- (4) a C_{1-6} alkyl group (e.g., methyl, isobutyl) optionally substituted by 1 to 3 substituents selected from
- (i) a hydroxy group, and
- (ii) a 3- to 8-membered monocyclic non-aromatic
- 20 heterocyclic group (e.g., tetrahydrofuryl).

[0106]

In this embodiment, R^1 is still more preferably

- (1) a phenyl group optionally substituted by 1 to 3 substituents selected from

- (i) a halogen atom (e.g., a fluorine atom), and
- 25 (ii) a cyano group,
- (2) a C_{5-6} cycloalkyl group (cyclopentyl, cyclohexyl) optionally substituted by 1 to 3 hydroxy groups,
- (3) a tetrahydropyranyl group optionally substituted by 1 to 3
- 30 hydroxy groups, or
- (4) a C_{1-6} alkyl group (e.g., methyl, isobutyl) optionally substituted by 1 to 3 substituents selected from
- (i) a hydroxy group, and
- (ii) a tetrahydrofuryl group.

35 [0107]

In another embodiment, R^1 is still more preferably
(1) a C_{5-6} cycloalkyl group (e.g., cyclohexyl) optionally
substituted by 1 to 3 hydroxy groups, or

(2) a 5- or 6-membered monocyclic non-aromatic heterocyclic
5 group (e.g., tetrahydropyranyl) optionally substituted by 1 to
3 hydroxy groups.

[0108]

In this embodiment, R^1 is even more preferably

(1) a C_{5-6} cycloalkyl group (e.g., cyclohexyl) substituted by 1
10 to 3 hydroxy groups, or

(2) a 5- or 6-membered monocyclic non-aromatic heterocyclic
group (e.g., tetrahydropyranyl) substituted by 1 to 3 hydroxy
groups.

[0109]

15 In this embodiment, R^1 is particularly preferably

(1) a cyclohexyl group substituted by one hydroxy group, or

(2) a tetrahydropyranyl group substituted by one hydroxy group.

[0110]

R^2 and R^3 are the same or different and each is a
20 hydrogen atom, a halogen atom, a cyano group, an optionally
substituted C_{1-6} alkyl group, an optionally substituted C_{1-6}
alkoxy group or an optionally substituted C_{3-6} cycloalkyl group.

[0111]

Examples of the " C_{3-6} cycloalkyl group" of the "optionally
25 substituted C_{3-6} cycloalkyl group" for R^2 or R^3 include
cyclopropyl, cyclobutyl, cyclopentyl and cyclohexyl.

[0112]

The " C_{1-6} alkyl group" of the "optionally substituted C_{1-6}
alkyl group", the " C_{1-6} alkoxy group" of the "optionally
30 substituted C_{1-6} alkoxy group" and the " C_{3-6} cycloalkyl group" of
the "optionally substituted C_{3-6} cycloalkyl group" for R^2 or R^3
each optionally has 1 to 5 (preferably 1 to 3) substituents at
substitutable position(s). Examples of the substituent include
the above-mentioned Substituent Group A. When the number of
35 the substituents is plural, the respective substituents may be

the same or different.

[0113]

R^2 is preferably

- (1) a hydrogen atom,
- 5 (2) a halogen atom (e.g., a fluorine atom, a chlorine atom), or
- (3) an optionally substituted C_{1-6} alkyl group (e.g., methyl).

[0114]

R^2 is more preferably

- (1) a hydrogen atom,
- 10 (2) a halogen atom (e.g., a chlorine atom, a fluorine atom), or
- (3) a C_{1-6} alkyl group (e.g., methyl).

[0115]

R^2 is further more preferably

- (1) a halogen atom (e.g., a fluorine atom), or
- 15 (2) a C_{1-6} alkyl group (e.g., methyl).

[0116]

R^3 is preferably

- (1) a hydrogen atom,
- (2) a halogen atom (e.g., a chlorine atom),
- 20 (3) a cyano group,
- (4) an optionally substituted C_{1-6} alkyl group (e.g., methyl, ethyl),
- (5) a C_{1-6} alkoxy group (e.g., methoxy), or
- (6) a C_{3-6} cycloalkyl group (e.g., cyclopropyl).

25 [0117]

R^3 is more preferably

- (1) a hydrogen atom,
- (2) a halogen atom (e.g., a chlorine atom),
- (3) a cyano group,
- 30 (4) a C_{1-6} alkyl group (e.g., methyl, ethyl) optionally substituted by 1 to 3 halogen atoms (e.g., a fluorine atom),
- (5) a C_{1-6} alkoxy group (e.g., methoxy), or
- (6) a C_{3-6} cycloalkyl group (e.g., cyclopropyl).

[0118]

35 R^3 is further more preferably

- (1) a hydrogen atom,
(2) a halogen atom (e.g., a chlorine atom),
(3) a cyano group,
(4) a C₁₋₆ alkyl group (e.g., methyl) optionally substituted by
5 1 to 3 halogen atoms (e.g., a fluorine atom),
(5) a C₁₋₆ alkoxy group (e.g., methoxy), or
(6) a C₃₋₆ cycloalkyl group (e.g., cyclopropyl).

[0119]

R³ is particularly preferably a C₁₋₆ alkyl group (e.g.,
10 methyl).

[0120]

R⁴ is a halogen atom, a cyano group, an optionally substituted C₁₋₆ alkyl group, an optionally substituted C₁₋₆ alkoxy group, an optionally substituted carbamoyl group or an
15 optionally substituted 3- to 8-membered cyclic group.

[0121]

Examples of the "3- to 8-membered cyclic group" of the "optionally substituted 3- to 8-membered cyclic group" for R⁴ include a phenyl group, a C₃₋₈ cycloalkyl group, a C₃₋₈
20 cycloalkenyl group, a 5- or 6-membered monocyclic aromatic heterocyclic group, a 3- to 8-membered monocyclic non-aromatic heterocyclic group and the like, and a 5- or 6-membered monocyclic aromatic heterocyclic group is preferable.

[0122]

Examples of the "C₃₋₈ cycloalkyl group" exemplified as the above-mentioned "3- to 8-membered cyclic group" include cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, cyclooctyl, bicyclo[2.2.1]heptyl, bicyclo[2.2.2]octyl, bicyclo[3.2.1]octyl and the like.

30 [0123]

Examples of the "C₃₋₈ cycloalkenyl group" exemplified as the above-mentioned "3- to 8-membered cyclic group" include cyclopropenyl, cyclobutenyl, cyclopentenyl, cyclohexenyl, cycloheptenyl and cyclooctenyl.

35 [0124]

Examples of the "5- or 6-membered monocyclic aromatic heterocyclic group" exemplified as the above-mentioned "3- to 8-membered cyclic group" include a 5- or 6-membered monocyclic aromatic heterocyclic group containing, as a ring-constituting atom besides carbon atom, 1 to 4 heteroatoms selected from a nitrogen atom, a sulfur atom and an oxygen atom, and specific examples thereof include those exemplified as the "5- to 6-membered monocyclic aromatic heterocyclic group", from among the above-mentioned preferable examples of the "aromatic heterocyclic group".

[0125]

Examples of the "3- to 8-membered monocyclic non-aromatic heterocyclic group" exemplified as the above-mentioned "3- to 8-membered cyclic group" include a 3- to 8-membered monocyclic non-aromatic heterocyclic group containing, as a ring-constituting atom besides carbon atom, 1 to 4 heteroatoms selected from a nitrogen atom, a sulfur atom and an oxygen atom, and specific examples thereof include those exemplified as the "3- to 8-membered monocyclic non-aromatic heterocyclic group", from among the above-mentioned preferable examples of the "non-aromatic heterocyclic group".

[0126]

The "C₁₋₆ alkyl group" of the "optionally substituted C₁₋₆ alkyl group", the "C₁₋₆ alkoxy group" of the "optionally substituted C₁₋₆ alkoxy group" and the "3- to 8-membered cyclic group" of the "optionally substituted 3- to 8-membered cyclic group" for R⁴ each optionally has 1 to 5 (preferably 1 to 3) substituents at substitutable position(s). Examples of the substituent include the above-mentioned Substituent Group A. When the number of the substituents is plural, the respective substituents may be the same or different.

[0127]


R⁴ is preferably bonded to the carbon atom at the p-position (4-position) in the "6-membered aromatic ring" of the "optionally further substituted 6-membered aromatic ring" for

Ring A (i.e., at the p-position (4-position) relative to the binding site of the oxoisoindolinylmethyl in the formula (I)).

[0128]

That is, the partial structure represented by the following formula:


[0129]



[0130]

in the formula (I) is preferably a partial structure represented by the following formula:

[0131]



[0132]

R⁴ is preferably

- (1) a halogen atom,
- (2) a cyano group,
- (3) a C₁₋₆ alkyl group,
- (4) an optionally substituted C₁₋₆ alkoxy group,
- (5) an optionally substituted carbamoyl group, or
- (6) an optionally substituted 5- or 6-membered monocyclic aromatic heterocyclic group.

[0133]

R⁴ is more preferably

- (1) a halogen atom (e.g., a fluorine atom, a chlorine atom),
- (2) a cyano group,
- (3) a C₁₋₆ alkyl group (e.g., methyl, ethyl),
- (4) a C₁₋₆ alkoxy group (e.g., methoxy, ethoxy, isopropoxy) optionally substituted by 1 to 3 halogen atoms (e.g., a fluorine atom),
- (5) a carbamoyl group,

(6) a mono- or di-C₁₋₆ alkyl-carbamoyl group (e.g., methylcarbamoyl), or

(7) a 5- or 6-membered monocyclic aromatic heterocyclic group (e.g., pyrazolyl, pyridyl, triazolyl, pyridazinyl) optionally substituted by 1 to 3 C₁₋₆ alkyl groups (e.g., methyl).

[0134]

R⁴ is further more preferably

(1) a halogen atom (e.g., a fluorine atom, a chlorine atom),

(2) a cyano group,

10 (3) a C₁₋₆ alkyl group (e.g., methyl, ethyl),

(4) a C₁₋₆ alkoxy group (e.g., methoxy, ethoxy, isopropoxy) optionally substituted by 1 to 3 halogen atoms (e.g., a fluorine atom),

(5) a carbamoyl group,

15 (6) a mono- or di-C₁₋₆ alkyl-carbamoyl group (e.g., methylcarbamoyl),

(7) a pyrazolyl group optionally substituted by 1 to 3 C₁₋₆ alkyl groups (e.g., methyl),

(8) a pyridyl group optionally substituted by 1 to 3 C₁₋₆ alkyl groups (e.g., methyl),

(9) a triazolyl group optionally substituted by 1 to 3 C₁₋₆ alkyl groups (e.g., methyl), or

(10) a pyridazinyl group optionally substituted by 1 to 3 C₁₋₆ alkyl groups (e.g., methyl).

25 [0135]

In another embodiment, R⁴ is more preferably

(1) a halogen atom (e.g., a fluorine atom, a chlorine atom),

(2) a cyano group,

(3) a C₁₋₆ alkyl group (e.g., methyl, ethyl),

30 (4) a C₁₋₆ alkoxy group (e.g., methoxy, ethoxy) optionally substituted by 1 to 3 halogen atoms (e.g., a fluorine atom),

(5) a carbamoyl group,

(6) a mono- or di-C₁₋₆ alkyl-carbamoyl group (e.g., methylcarbamoyl), or

35 (7) a 5- or 6-membered monocyclic aromatic heterocyclic group

(e.g., pyrazolyl, pyridyl, triazolyl, pyridazinyl) optionally substituted by 1 to 3 C₁₋₆ alkyl groups (e.g., methyl).

[0136]

In this embodiment, R⁴ is further more preferably

- 5 (1) a halogen atom (e.g., a fluorine atom, a chlorine atom),
- (2) a cyano group,
- (3) a C₁₋₆ alkyl group (e.g., methyl, ethyl),
- (4) a C₁₋₆ alkoxy group (e.g., methoxy, ethoxy) optionally substituted by 1 to 3 halogen atoms (e.g., a fluorine atom),
- 10 (5) a carbamoyl group,
- (6) a mono- or di-C₁₋₆ alkyl-carbamoyl group (e.g., methylcarbamoyl),
- (7) a pyrazolyl group optionally substituted by 1 to 3 C₁₋₆ alkyl groups (e.g., methyl),
- 15 (8) a pyridyl group optionally substituted by 1 to 3 C₁₋₆ alkyl groups (e.g., methyl),
- (9) a triazolyl group optionally substituted by 1 to 3 C₁₋₆ alkyl groups (e.g., methyl), or
- (10) a pyridazinyl group optionally substituted by 1 to 3 C₁₋₆ alkyl groups (e.g., methyl).

[0137]

In another embodiment, R⁴ is more preferably

- (1) a halogen atom (e.g., a fluorine atom, a chlorine atom),
- (2) a cyano group,
- 25 (3) a C₁₋₆ alkyl group (e.g., methyl),
- (4) a C₁₋₆ alkoxy group (e.g., methoxy) optionally substituted by 1 to 3 halogen atoms (e.g., a fluorine atom),
- (5) a mono- or di-C₁₋₆ alkyl-carbamoyl group (e.g., methylcarbamoyl), or
- 30 (6) a 5- or 6-membered monocyclic aromatic heterocyclic group (e.g., pyrazolyl, pyridyl, triazolyl) optionally substituted by 1 to 3 C₁₋₆ alkyl groups (e.g., methyl).

[0138]

In this embodiment, R⁴ is further more preferably

- 35 (1) a halogen atom (e.g., a fluorine atom, a chlorine atom),

- (2) a cyano group,
(3) a C₁₋₆ alkyl group (e.g., methyl),
(4) a C₁₋₆ alkoxy group (e.g., methoxy) optionally substituted
by 1 to 3 halogen atoms (e.g., a fluorine atom),
5 (5) a mono- or di-C₁₋₆ alkyl-carbamoyl group (e.g.,
methylcarbamoyl),
(6) a pyrazolyl group optionally substituted by 1 to 3 C₁₋₆
alkyl groups (e.g., methyl),
(7) a pyridyl group optionally substituted by 1 to 3 C₁₋₆ alkyl
10 groups (e.g., methyl), or
(8) a triazolyl group optionally substituted by 1 to 3 C₁₋₆
alkyl groups (e.g., methyl).

[0139]

In another embodiment, R⁴ is further more preferably

- 15 (1) a C₁₋₆ alkyl group (e.g., methyl),
(2) a C₁₋₆ alkoxy group (e.g., methoxy), or
(3) a 5- or 6-membered monocyclic aromatic heterocyclic group
(e.g., pyrazolyl).

[0140]

20 R⁴ is particularly preferably

- (1) a C₁₋₆ alkyl group (e.g., methyl),
(2) a C₁₋₆ alkoxy group (e.g., methoxy), or
(3) a pyrazolyl group.

[0141]

25 Ring A is an optionally further substituted 6-membered
aromatic ring.

Examples of the "6-membered aromatic ring" of the
"optionally further substituted 6-membered aromatic ring" for
Ring A include a benzene ring, a pyridine ring, a pyridazine
30 ring, a pyrimidine ring, a pyrazine ring, a triazine ring and
the like, a benzene ring and a pyridine ring are preferable.

[0142]

The "6-membered aromatic ring" of the "optionally further
substituted 6-membered aromatic ring" for Ring A optionally has
35 1 to 4 (preferably 1 to 3) substituents, in addition to R⁴, at

substitutable position(s). Examples of the substituent include the above-mentioned Substituent Group A. When the number of the substituents is plural, the respective substituents may be the same or different.

5 [0143]

Ring A is preferably a 6-membered aromatic ring (e.g., a benzene ring, a pyridine ring) optionally further substituted by 1 to 3 substituents, in addition to R⁴, selected from

- (a) a halogen atom (e.g., a fluorine atom),
- 10 (b) a C₁₋₆ alkyl group (e.g., methyl),
- (c) a C₁₋₆ alkoxy group (e.g., methoxy),
- (d) a cyano group,
- (e) a carbamoyl group,
- (f) a mono- or di-C₁₋₆ alkyl-carbamoyl group (e.g.,
- 15 methylcarbamoyl), and
- (g) a 5- to 14-membered aromatic heterocyclic group (preferably a 5- or 6-membered monocyclic aromatic heterocyclic group) (e.g., pyrazolyl, triazolyl) optionally substituted by 1 to 3 C₁₋₆ alkyl groups (e.g., methyl).

20 [0144]

Ring A is more preferably a 6-membered aromatic ring (e.g., a benzene ring, a pyridine ring) optionally further substituted by 1 to 3 substituents, in addition to R⁴, selected from

- 25 (a) a halogen atom (e.g., a fluorine atom),
- (b) a C₁₋₆ alkyl group (e.g., methyl),
- (c) a C₁₋₆ alkoxy group (e.g., methoxy),
- (d) a cyano group,
- (e) a carbamoyl group,
- 30 (f) a mono- or di-C₁₋₆ alkyl-carbamoyl group (e.g., methylcarbamoyl),
- (g) a pyrazolyl group optionally substituted by 1 to 3 C₁₋₆ alkyl groups (e.g., methyl), and
- (h) a triazolyl group optionally substituted by 1 to 3 C₁₋₆
- 35 alkyl groups (e.g., methyl).

[0145]

Ring A is further more preferably

(1) a benzene ring optionally further substituted by 1 to 3 substituents, in addition to R⁴, selected from

- 5 (a) a halogen atom (e.g., a fluorine atom),
- (b) a C₁₋₆ alkyl group (e.g., methyl),
- (c) a C₁₋₆ alkoxy group (e.g., methoxy),
- (d) a cyano group,
- (e) a carbamoyl group,
- 10 (f) a mono- or di-C₁₋₆ alkyl-carbamoyl group (e.g., methylcarbamoyl),
- (g) a pyrazolyl group optionally substituted by 1 to 3 C₁₋₆ alkyl groups (e.g., methyl), and
- (h) a triazolyl group optionally substituted by 1 to 3 C₁₋₆
- 15 alkyl groups (e.g., methyl), or
- (2) a pyridine ring.

[0146]

In another embodiment, Ring A is preferably a 6-membered aromatic ring (e.g., a benzene ring, a pyridine ring)

20 optionally further substituted by 1 to 3 substituents, in addition to R⁴, selected from

- (a) a halogen atom (e.g., a fluorine atom),
- (b) a C₁₋₆ alkyl group (e.g., methyl),
- (c) a C₁₋₆ alkoxy group (e.g., methoxy),
- 25 (d) a cyano group,
- (e) a mono- or di-C₁₋₆ alkyl-carbamoyl group (e.g., methylcarbamoyl), and
- (f) a 5- to 14-membered aromatic heterocyclic group (preferably a 5- or 6-membered monocyclic aromatic
- 30 heterocyclic group) (e.g., pyrazolyl).

[0147]

In this embodiment, Ring A is more preferably a 6-membered aromatic ring (e.g., a benzene ring, a pyridine ring) optionally further substituted by 1 to 3 substituents, in

35 addition to R⁴, selected from

- (a) a halogen atom (e.g., a fluorine atom),
- (b) a C₁₋₆ alkoxy group (e.g., methoxy),
- (c) a cyano group,
- (d) a mono- or di-C₁₋₆ alkyl-carbamoyl group (e.g., methylcarbamoyl), and
- (e) a 5- to 14-membered aromatic heterocyclic group (preferably a 5- or 6-membered monocyclic aromatic heterocyclic group) (e.g., pyrazolyl).

[0148]

10 In this embodiment, Ring A is further more preferably a 6-membered aromatic ring (e.g., a benzene ring, a pyridine ring) optionally further substituted by 1 to 3 substituents, in addition to R⁴, selected from

- (a) a halogen atom (e.g., a fluorine atom),
- 15 (b) a C₁₋₆ alkoxy group (e.g., methoxy),
- (c) a cyano group,
- (d) a mono- or di-C₁₋₆ alkyl-carbamoyl group (e.g., methylcarbamoyl), and
- (e) a pyrazolyl group.

20 [0149]

In this embodiment, Ring A is still more preferably (1) a benzene ring optionally further substituted by 1 to 3 substituents, in addition to R⁴, selected from

- (a) a halogen atom (e.g., a fluorine atom),
- 25 (b) a C₁₋₆ alkoxy group (e.g., methoxy),
- (c) a cyano group,
- (d) a mono- or di-C₁₋₆ alkyl-carbamoyl group (e.g., methylcarbamoyl), and
- (e) a pyrazolyl group, or

30 (2) a pyridine ring.

[0150]

In another embodiment, Ring A is preferably a 6-membered aromatic ring (e.g., a benzene ring, a pyridine ring) optionally further substituted by 1 to 3 substituents, in addition to R⁴, selected from

35

- (a) a halogen atom (e.g., a fluorine atom),
- (b) a C₁₋₆ alkyl group (e.g., methyl), and
- (c) a C₁₋₆ alkoxy group (e.g., methoxy).

[0151]

- 5 In this embodiment, Ring A is more preferably
- (1) a benzene ring optionally further substituted by 1 to 3 substituents, in addition to R⁴, selected from
 - (a) a halogen atom (e.g., a fluorine atom),
 - (b) a C₁₋₆ alkyl group (e.g., methyl), and
 - 10 (c) a C₁₋₆ alkoxy group (e.g., methoxy), or
 - (2) a pyridine ring.

[0152]

- In another embodiment, Ring A is preferably a 6-membered aromatic ring (e.g., a benzene ring, a pyridine ring)
- 15 optionally further substituted by 1 to 3 substituents, in addition to R⁴, selected from
- (a) a halogen atom (e.g., a fluorine atom), and
 - (b) a C₁₋₆ alkoxy group (e.g., methoxy).

[0153]

- 20 In this embodiment, Ring A is more preferably
- (1) a benzene ring optionally further substituted by 1 to 3 substituents, in addition to R⁴, selected from
 - (a) a halogen atom (e.g., a fluorine atom), and
 - (b) a C₁₋₆ alkoxy group (e.g., methoxy), or
 - 25 (2) a pyridine ring.

[0154]

Ring A is particularly preferably a benzene ring or a pyridine ring, each of which is unsubstituted, in addition to R⁴.

30 [0155]

Preferable examples of compound (I) include the following compounds.

[0156]

[Compound A-1]

35 Compound (I) wherein

R¹ is

- (1) an optionally substituted phenyl group,
- (2) an optionally substituted C₅₋₆ cycloalkyl group,
- (3) an optionally substituted 5- or 6-membered non-aromatic
5 heterocyclic group,
- (4) an optionally substituted 5- or 6-membered monocyclic
aromatic heterocyclic group, or
- (5) an optionally substituted C₁₋₆ alkyl group;

R² is

- 10 (1) a hydrogen atom,
- (2) a halogen atom (e.g., a fluorine atom, a chlorine atom), or
- (3) an optionally substituted C₁₋₆ alkyl group (e.g., methyl);

R³ is

- (1) a hydrogen atom,
- 15 (2) a halogen atom (e.g., a chlorine atom),
- (3) a cyano group,
- (4) an optionally substituted C₁₋₆ alkyl group (e.g., methyl,
ethyl),
- (5) a C₁₋₆ alkoxy group (e.g., methoxy), or
- 20 (6) a C₃₋₆ cycloalkyl group (e.g., cyclopropyl);

R⁴ is

- (1) a halogen atom,
- (2) a cyano group,
- (3) a C₁₋₆ alkyl group,
- 25 (4) an optionally substituted C₁₋₆ alkoxy group,
- (5) an optionally substituted carbamoyl group, or
- (6) an optionally substituted 5- or 6-membered monocyclic
aromatic heterocyclic group; and


Ring A is a 6-membered aromatic ring (e.g., a benzene
30 ring, a pyridine ring) optionally further substituted by 1 to 3
substituents, in addition to R⁴, selected from

- (a) a halogen atom (e.g., a fluorine atom),
- (b) a C₁₋₆ alkyl group (e.g., methyl), and
- (c) a C₁₋₆ alkoxy group (e.g., methoxy).

35 In [Compound A-1], the partial structure represented by

the following formula:


[0157]



[0158]

- 5 in the formula (I) is preferably a partial structure represented by the following formula:

[0159]



[0160]

- 10 [Compound A-2]

Compound (I) wherein

R^1 is

- (1) a phenyl group, a C_{5-6} cycloalkyl group (cyclopentyl, cyclohexyl), a 5- or 6-membered monocyclic non-aromatic
 15 heterocyclic group (e.g., tetrahydropyranyl) or a 5- or 6-membered monocyclic aromatic heterocyclic group (e.g., pyridyl), each of which is optionally substituted by 1 to 3 substituents selected from
- (i) a halogen atom (e.g., a fluorine atom, a chlorine atom),
 - 20 (ii) a cyano group,
 - (iii) a hydroxy group,
 - (iv) a C_{1-6} alkyl group (e.g., methyl) optionally substituted by 1 to 3 hydroxy groups, and
 - (v) a C_{1-6} alkoxy group (e.g., methoxy), or
- 25 (2) a C_{1-6} alkyl group (e.g., methyl, isobutyl, 1,2-dimethylpropyl) optionally substituted by 1 to 3 substituents selected from
- (i) a hydroxy group, and
 - (ii) a 3- to 14-membered non-aromatic heterocyclic group
 - 30 (preferably a 3- to 8-membered monocyclic non-aromatic

heterocyclic group) (e.g., tetrahydrofuryl);

R² is

- (1) a hydrogen atom,
- (2) a halogen atom (e.g., a chlorine atom, a fluorine atom), or
- 5 (3) a C₁₋₆ alkyl group (e.g., methyl);

R³ is

- (1) a hydrogen atom,
- (2) a halogen atom (e.g., a chlorine atom),
- (3) a cyano group,
- 10 (4) a C₁₋₆ alkyl group (e.g., methyl, ethyl) optionally substituted by 1 to 3 halogen atoms (e.g., a fluorine atom),
- (5) a C₁₋₆ alkoxy group (e.g., methoxy), or
- (6) a C₃₋₆ cycloalkyl group (e.g., cyclopropyl);

R⁴ is


- 15 (1) a halogen atom (e.g., a fluorine atom, a chlorine atom),
- (2) a cyano group,
- (3) a C₁₋₆ alkyl group (e.g., methyl, ethyl),
- (4) a C₁₋₆ alkoxy group (e.g., methoxy, ethoxy, isopropoxy) optionally substituted by 1 to 3 halogen atoms (e.g., a
- 20 fluorine atom),
- (5) a carbamoyl group,
- (6) a mono- or di-C₁₋₆ alkyl-carbamoyl group (e.g., methylcarbamoyl), or
- (7) a 5- or 6-membered monocyclic aromatic heterocyclic group
- 25 (e.g., pyrazolyl, pyridyl, triazolyl, pyridazinyl) optionally substituted by 1 to 3 C₁₋₆ alkyl groups (e.g., methyl); and

Ring A is a 6-membered aromatic ring (e.g., a benzene ring, a pyridine ring) optionally further substituted by 1 to 3 substituents, in addition to R⁴, selected from

- 30 (a) a halogen atom (e.g., a fluorine atom),
- (b) a C₁₋₆ alkyl group (e.g., methyl), and
- (c) a C₁₋₆ alkoxy group (e.g., methoxy).

In [Compound A-2], the partial structure represented by the following formula:


35 [0161]



[0162]

in the formula (I) is preferably a partial structure represented by the following formula:

5 [0163]



[0164]

[Compound A-3]

Compound (I) wherein

10 R^1 is

(1) a phenyl group optionally substituted by 1 to 3 substituents selected from

- (i) a halogen atom (e.g., a fluorine atom), and
- (ii) a cyano group,

15 (2) a C_{5-6} cycloalkyl group (cyclopentyl, cyclohexyl) optionally substituted by 1 to 3 substituents selected from

- (i) a hydroxy group,
- (ii) a C_{1-6} alkyl group (e.g., methyl) optionally substituted by 1 to 3 hydroxy groups, and

20 (iii) a C_{1-6} alkoxy group (e.g., methoxy),

(3) a 5- or 6-membered monocyclic non-aromatic heterocyclic group (e.g., tetrahydropyranyl) optionally substituted by 1 to 3 hydroxy groups,

25 (4) a 5- or 6-membered monocyclic aromatic heterocyclic group (e.g., pyridyl) optionally substituted by 1 to 3 halogen atoms (e.g., a fluorine atom, a chlorine atom), or

(5) a C_{1-6} alkyl group (e.g., methyl, isobutyl, 1,2-dimethylpropyl) optionally substituted by 1 to 3 substituents selected from

30 (i) a hydroxy group, and

(ii) a 3- to 8-membered monocyclic non-aromatic heterocyclic group (e.g., tetrahydrofuryl);

R^2 is

- (1) a hydrogen atom,
- 5 (2) a halogen atom (e.g., a chlorine atom, a fluorine atom), or
- (3) a C_{1-6} alkyl group (e.g., methyl);

R^3 is

- (1) a hydrogen atom,
- (2) a halogen atom (e.g., a chlorine atom),
- 10 (3) a cyano group,
- (4) a C_{1-6} alkyl group (e.g., methyl, ethyl) optionally substituted by 1 to 3 halogen atoms (e.g., a fluorine atom),
- (5) a C_{1-6} alkoxy group (e.g., methoxy), or
- (6) a C_{3-6} cycloalkyl group (e.g., cyclopropyl);

15 R^4 is


- (1) a halogen atom (e.g., a fluorine atom, a chlorine atom),
- (2) a cyano group,
- (3) a C_{1-6} alkyl group (e.g., methyl, ethyl),
- (4) a C_{1-6} alkoxy group (e.g., methoxy, ethoxy, isopropoxy)
- 20 optionally substituted by 1 to 3 halogen atoms (e.g., a fluorine atom),
- (5) a carbamoyl group,
- (6) a mono- or di- C_{1-6} alkyl-carbamoyl group (e.g., methylcarbamoyl), or
- 25 (7) a 5- or 6-membered monocyclic aromatic heterocyclic group (e.g., pyrazolyl, pyridyl, triazolyl, pyridazinyl) optionally substituted by 1 to 3 C_{1-6} alkyl groups (e.g., methyl); and

Ring A is a 6-membered aromatic ring (e.g., a benzene ring, a pyridine ring) optionally further substituted by 1 to 3
30 substituents, in addition to R^4 , selected from

- (a) a halogen atom (e.g., a fluorine atom),
- (b) a C_{1-6} alkyl group (e.g., methyl), and
- (c) a C_{1-6} alkoxy group (e.g., methoxy).

In [Compound A-3], the partial structure represented by
35 the following formula:


[0165]



[0166]

in the formula (I) is preferably a partial structure
 5 represented by the following formula:

[0167]



[0168]

[Compound A-4]

10 Compound (I) wherein

R^1 is

(1) a phenyl group optionally substituted by 1 to 3
 substituents selected from

(i) a halogen atom (e.g., a fluorine atom), and

15 (ii) a cyano group,

(2) a C_{5-6} cycloalkyl group (cyclopentyl, cyclohexyl) optionally
 substituted by 1 to 3 substituents selected from

(i) a hydroxy group,

(ii) a C_{1-6} alkyl group (e.g., methyl) optionally substituted
 20 by 1 to 3 hydroxy groups, and

(iii) a C_{1-6} alkoxy group (e.g., methoxy),

(3) a tetrahydropyranyl group optionally substituted by 1 to 3
 hydroxy groups,

(4) a pyridyl group optionally substituted by 1 to 3 halogen
 25 atoms (e.g., a fluorine atom, a chlorine atom), or

(5) a C_{1-6} alkyl group (e.g., methyl, isobutyl, 1,2-
 dimethylpropyl) optionally substituted by 1 to 3 substituents
 selected from

(i) a hydroxy group, and

30 (ii) a tetrahydrofuryl group;

R² is

- (1) a hydrogen atom,
- (2) a halogen atom (e.g., a chlorine atom, a fluorine atom), or
- (3) a C₁₋₆ alkyl group (e.g., methyl);

5 R³ is

- (1) a hydrogen atom,
- (2) a halogen atom (e.g., a chlorine atom),
- (3) a cyano group,
- (4) a C₁₋₆ alkyl group (e.g., methyl, ethyl) optionally
- 10 substituted by 1 to 3 halogen atoms (e.g., a fluorine atom),
- (5) a C₁₋₆ alkoxy group (e.g., methoxy), or
- (6) a C₃₋₆ cycloalkyl group (e.g., cyclopropyl);

R⁴ is

- (1) a halogen atom (e.g., a fluorine atom, a chlorine atom),
- 15 (2) a cyano group,
- (3) a C₁₋₆ alkyl group (e.g., methyl, ethyl),
- (4) a C₁₋₆ alkoxy group (e.g., methoxy, ethoxy, isopropoxy)
- optionally substituted by 1 to 3 halogen atoms (e.g., a
- fluorine atom),
- 20 (5) a carbamoyl group,
- (6) a mono- or di-C₁₋₆ alkyl-carbamoyl group (e.g.,
- methylcarbamoyl),
- (7) a pyrazolyl group optionally substituted by 1 to 3 C₁₋₆
- alkyl groups (e.g., methyl),
- 25 (8) a pyridyl group optionally substituted by 1 to 3 C₁₋₆ alkyl
- groups (e.g., methyl),
- (9) a triazolyl group optionally substituted by 1 to 3 C₁₋₆
- alkyl groups (e.g., methyl), or
- (10) a pyridazinyl group optionally substituted by 1 to 3 C₁₋₆
- 30 alkyl groups (e.g., methyl); and


Ring A is

- (1) a benzene ring optionally further substituted by 1 to 3
- substituents, in addition to R⁴, selected from
- (a) a halogen atom (e.g., a fluorine atom),
- 35 (b) a C₁₋₆ alkyl group (e.g., methyl), and

- (c) a C₁₋₆ alkoxy group (e.g., methoxy), or
(2) a pyridine ring.

In [Compound A-4], the partial structure represented by the following formula:


5 [0169]



[0170]

in the formula (I) is preferably a partial structure represented by the following formula:

10 [0171]



[0172]

[Compound B-1]

Compound (I) wherein

15 R¹ is

- (1) a phenyl group, a C₅₋₆ cycloalkyl group (cyclopentyl, cyclohexyl), a 5- or 6-membered monocyclic non-aromatic heterocyclic group (e.g., tetrahydropyranyl) or a 5- or 6-membered monocyclic aromatic heterocyclic group (e.g., pyridyl),
20 each of which is optionally substituted by 1 to 3 substituents selected from

- (i) a halogen atom (e.g., a fluorine atom, a chlorine atom),
(ii) a cyano group, and
(iii) a hydroxy group, or

- 25 (2) a C₁₋₆ alkyl group (e.g., methyl, isobutyl, 1,2-dimethylpropyl) optionally substituted by 1 to 3 substituents selected from

- (i) a hydroxy group, and
(ii) a 3- to 14-membered non-aromatic heterocyclic group
30 (preferably a 3- to 8-membered monocyclic non-aromatic

heterocyclic group) (e.g., tetrahydrofuryl);

R^2 is

- (1) a hydrogen atom,
- (2) a halogen atom (e.g., a chlorine atom, a fluorine atom), or
- 5 (3) a C_{1-6} alkyl group (e.g., methyl);

R^3 is

- (1) a hydrogen atom,
- (2) a halogen atom (e.g., a chlorine atom),
- (3) a cyano group,
- 10 (4) a C_{1-6} alkyl group (e.g., methyl, ethyl) optionally substituted by 1 to 3 halogen atoms (e.g., a fluorine atom),
- (5) a C_{1-6} alkoxy group (e.g., methoxy), or
- (6) a C_{3-6} cycloalkyl group (e.g., cyclopropyl);

R^4 is


- 15 (1) a halogen atom (e.g., a fluorine atom, a chlorine atom),
- (2) a cyano group,
- (3) a C_{1-6} alkyl group (e.g., methyl, ethyl),
- (4) a C_{1-6} alkoxy group (e.g., methoxy, ethoxy) optionally substituted by 1 to 3 halogen atoms (e.g., a fluorine atom),
- 20 (5) a carbamoyl group,
- (6) a mono- or di- C_{1-6} alkyl-carbamoyl group (e.g., methylcarbamoyl), or
- (7) a 5- or 6-membered monocyclic aromatic heterocyclic group (e.g., pyrazolyl, pyridyl, triazolyl, pyridazinyl) optionally
- 25 substituted by 1 to 3 C_{1-6} alkyl groups (e.g., methyl); and

Ring A is a 6-membered aromatic ring (e.g., a benzene ring, a pyridine ring) optionally further substituted by 1 to 3 substituents, in addition to R^4 , selected from

- (a) a halogen atom (e.g., a fluorine atom),
- 30 (b) a C_{1-6} alkyl group (e.g., methyl), and
- (c) a C_{1-6} alkoxy group (e.g., methoxy).

In [Compound B-1], the partial structure represented by the following formula:


[0173]



[0174]

in the formula (I) is preferably a partial structure represented by the following formula:

5 [0175]



[0176]

[Compound B-2]

Compound (I) wherein

10 R^1 is

(1) a phenyl group optionally substituted by 1 to 3 substituents selected from

- (i) a halogen atom (e.g., a fluorine atom), and
- (ii) a cyano group,

15 (2) a C_{5-6} cycloalkyl group (cyclopentyl, cyclohexyl) optionally substituted by 1 to 3 hydroxy groups,

(3) a 5- or 6-membered monocyclic non-aromatic heterocyclic group (e.g., tetrahydropyranyl) optionally substituted by 1 to 3 hydroxy groups,

20 (4) a 5- or 6-membered monocyclic aromatic heterocyclic group (e.g., pyridyl) optionally substituted by 1 to 3 halogen atoms (e.g., a fluorine atom, a chlorine atom), or

(5) a C_{1-6} alkyl group (e.g., methyl, isobutyl, 1,2-dimethylpropyl) optionally substituted by 1 to 3 substituents

25 selected from

- (i) a hydroxy group, and
- (ii) a 3- to 8-membered monocyclic non-aromatic heterocyclic group (e.g., tetrahydrofuryl);

 R^2 is

30 (1) a hydrogen atom,

- (2) a halogen atom (e.g., a chlorine atom, a fluorine atom), or
 (3) a C₁₋₆ alkyl group (e.g., methyl);

R³ is

- (1) a hydrogen atom,
 5 (2) a halogen atom (e.g., a chlorine atom),
 (3) a cyano group,
 (4) a C₁₋₆ alkyl group (e.g., methyl, ethyl) optionally
 substituted by 1 to 3 halogen atoms (e.g., a fluorine atom),
 (5) a C₁₋₆ alkoxy group (e.g., methoxy), or
 10 (6) a C₃₋₆ cycloalkyl group (e.g., cyclopropyl);

R⁴ is


- (1) a halogen atom (e.g., a fluorine atom, a chlorine atom),
 (2) a cyano group,
 (3) a C₁₋₆ alkyl group (e.g., methyl, ethyl),
 15 (4) a C₁₋₆ alkoxy group (e.g., methoxy, ethoxy) optionally
 substituted by 1 to 3 halogen atoms (e.g., a fluorine atom),
 (5) a carbamoyl group,
 (6) a mono- or di-C₁₋₆ alkyl-carbamoyl group (e.g.,
 methylcarbamoyl), or
 20 (7) a 5- or 6-membered monocyclic aromatic heterocyclic group
 (e.g., pyrazolyl, pyridyl, triazolyl, pyridazinyl) optionally
 substituted by 1 to 3 C₁₋₆ alkyl groups (e.g., methyl); and

Ring A is a 6-membered aromatic ring (e.g., a benzene
 ring, a pyridine ring) optionally further substituted by 1 to 3
 25 substituents, in addition to R⁴, selected from

- (a) a halogen atom (e.g., a fluorine atom),
 (b) a C₁₋₆ alkyl group (e.g., methyl), and
 (c) a C₁₋₆ alkoxy group (e.g., methoxy).

In [Compound B-2], the partial structure represented by
 30 the following formula:


[0177]



[0178]

in the formula (I) is preferably a partial structure represented by the following formula:

[0179]



5 [0180]

[Compound B-3]

Compound (I) wherein

R^1 is

- (1) a phenyl group optionally substituted by 1 to 3
10 substituents selected from
 - (i) a halogen atom (e.g., a fluorine atom), and
 - (ii) a cyano group,
- (2) a C_{5-6} cycloalkyl group (cyclopentyl, cyclohexyl) optionally substituted by 1 to 3 hydroxy groups,
- 15 (3) a tetrahydropyranyl group optionally substituted by 1 to 3 hydroxy groups,
- (4) a pyridyl group optionally substituted by 1 to 3 halogen atoms (e.g., a fluorine atom, a chlorine atom), or
- (5) a C_{1-6} alkyl group (e.g., methyl, isobutyl, 1,2-
20 dimethylpropyl) optionally substituted by 1 to 3 substituents selected from

- (i) a hydroxy group, and
- (ii) a tetrahydrofuryl group;

R^2 is

- 25 (1) a hydrogen atom,
- (2) a halogen atom (e.g., a chlorine atom, a fluorine atom), or
- (3) a C_{1-6} alkyl group (e.g., methyl);

R^3 is

- (1) a hydrogen atom,
- 30 (2) a halogen atom (e.g., a chlorine atom),
- (3) a cyano group,
- (4) a C_{1-6} alkyl group (e.g., methyl, ethyl) optionally

substituted by 1 to 3 halogen atoms (e.g., a fluorine atom),
(5) a C₁₋₆ alkoxy group (e.g., methoxy), or
(6) a C₃₋₆ cycloalkyl group (e.g., cyclopropyl);

R⁴ is


- 5 (1) a halogen atom (e.g., a fluorine atom, a chlorine atom),
(2) a cyano group,
(3) a C₁₋₆ alkyl group (e.g., methyl, ethyl),
(4) a C₁₋₆ alkoxy group (e.g., methoxy, ethoxy) optionally
substituted by 1 to 3 halogen atoms (e.g., a fluorine atom),
10 (5) a carbamoyl group,
(6) a mono- or di-C₁₋₆ alkyl-carbamoyl group (e.g.,
methylcarbamoyl),
(7) a pyrazolyl group optionally substituted by 1 to 3 C₁₋₆
alkyl groups (e.g., methyl),
15 (8) a pyridyl group optionally substituted by 1 to 3 C₁₋₆ alkyl
groups (e.g., methyl),
(9) a triazolyl group optionally substituted by 1 to 3 C₁₋₆
alkyl groups (e.g., methyl), or
(10) a pyridazinyl group optionally substituted by 1 to 3 C₁₋₆
20 alkyl groups (e.g., methyl); and

Ring A is

- (1) a benzene ring optionally further substituted by 1 to 3
substituents, in addition to R⁴, selected from
(a) a halogen atom (e.g., a fluorine atom),
25 (b) a C₁₋₆ alkyl group (e.g., methyl), and
(c) a C₁₋₆ alkoxy group (e.g., methoxy), or
(2) a pyridine ring.

In [Compound B-3], the partial structure represented by
the following formula:

30 [0181]




[0182]

in the formula (I) is preferably a partial structure

represented by the following formula:

[0183]



[0184]

5 [Compound C-1]

Compound (I) wherein

R^1 is

- (1) an optionally substituted phenyl group,
- (2) an optionally substituted C_{5-6} cycloalkyl group,
- 10 (3) an optionally substituted 5- or 6-membered non-aromatic heterocyclic group, or
- (4) an optionally substituted C_{1-6} alkyl group;

R^2 is

- (1) a hydrogen atom,
- 15 (2) a halogen atom (e.g., a fluorine atom, a chlorine atom), or
- (3) an optionally substituted C_{1-6} alkyl group (e.g., methyl);

R^3 is

- (1) a hydrogen atom,
- (2) a halogen atom (e.g., a chlorine atom),
- 20 (3) a cyano group,
- (4) an optionally substituted C_{1-6} alkyl group (e.g., methyl),
- (5) a C_{1-6} alkoxy group (e.g., methoxy), or
- (6) a C_{3-6} cycloalkyl group (e.g., cyclopropyl);

R^4 is

- 25 (1) a halogen atom,
- (2) a cyano group,
- (3) a C_{1-6} alkyl group,
- (4) an optionally substituted C_{1-6} alkoxy group,
- (5) an optionally substituted carbamoyl group, or
- 30 (6) an optionally substituted 5- or 6-membered monocyclic aromatic heterocyclic group; and

Ring A is a 6-membered aromatic ring (e.g., a benzene

ring, a pyridine ring) optionally further substituted by 1 to 3 substituents, in addition to R⁴, selected from

- (a) a halogen atom (e.g., a fluorine atom), and
- (b) a C₁₋₆ alkoxy group (e.g., methoxy).

5 [0185]

[Compound Ca-1]

Compound (I) wherein

R¹ is

- (1) an optionally substituted phenyl group,
- 10 (2) an optionally substituted C₅₋₆ cycloalkyl group,
- (3) an optionally substituted 5- or 6-membered non-aromatic heterocyclic group, or
- (4) an optionally substituted C₁₋₆ alkyl group;

R² is

- 15 (1) a hydrogen atom,
- (2) a halogen atom (e.g., a fluorine atom, a chlorine atom), or
- (3) an optionally substituted C₁₋₆ alkyl group (e.g., methyl);

R³ is

- (1) a hydrogen atom,
- 20 (2) a halogen atom (e.g., a chlorine atom),
- (3) a cyano group,
- (4) an optionally substituted C₁₋₆ alkyl group (e.g., methyl),
- (5) a C₁₋₆ alkoxy group (e.g., methoxy), or
- (6) a C₃₋₆ cycloalkyl group (e.g., cyclopropyl);

25 R⁴ is

- (1) a halogen atom,
- (2) a cyano group,
- (3) a C₁₋₆ alkyl group,
- (4) an optionally substituted C₁₋₆ alkoxy group,
- 30 (5) an optionally substituted carbamoyl group, or
- (6) an optionally substituted 5- or 6-membered monocyclic aromatic heterocyclic group;


Ring A is a 6-membered aromatic ring (e.g., a benzene ring, a pyridine ring) optionally further substituted by 1 to 3
35 substituents, in addition to R⁴, selected from

(a) a halogen atom (e.g., a fluorine atom), and

(b) a C₁₋₆ alkoxy group (e.g., methoxy); and

the partial structure represented by the following formula:


5 [0186]



[0187]

in the formula (I) is a partial structure represented by the following formula:

10 [0188]



[0189]

[Compound C-2]

Compound (I) wherein

15 R¹ is

(1) a phenyl group, a C₅₋₆ cycloalkyl group (cyclopentyl, cyclohexyl) or a 5- or 6-membered monocyclic non-aromatic heterocyclic group (e.g., tetrahydropyranyl), each of which is optionally substituted by 1 to 3 substituents selected from

20 (i) a halogen atom (e.g., a fluorine atom),

(ii) a cyano group, and

(iii) a hydroxy group, or

(2) a C₁₋₆ alkyl group (e.g., methyl, isobutyl) optionally substituted by 1 to 3 substituents selected from

25 (i) a hydroxy group, and

(ii) a 3- to 14-membered non-aromatic heterocyclic group (preferably a 3- to 8-membered monocyclic non-aromatic heterocyclic group) (e.g., tetrahydrofuryl);

R² is

30 (1) a hydrogen atom,

(2) a halogen atom (e.g., a chlorine atom, a fluorine atom), or

(3) a C₁₋₆ alkyl group (e.g., methyl);

R³ is

(1) a hydrogen atom,

5 (2) a halogen atom (e.g., a chlorine atom),

(3) a cyano group,

(4) a C₁₋₆ alkyl group (e.g., methyl) optionally substituted by
1 to 3 halogen atoms (e.g., a fluorine atom),

(5) a C₁₋₆ alkoxy group (e.g., methoxy), or

10 (6) a C₃₋₆ cycloalkyl group (e.g., cyclopropyl);

R⁴ is

(1) a halogen atom (e.g., a chlorine atom, a fluorine atom),

(2) a cyano group,

(3) a C₁₋₆ alkyl group (e.g., methyl),

15 (4) a C₁₋₆ alkoxy group (e.g., methoxy) optionally substituted
by 1 to 3 halogen atoms (e.g., a fluorine atom),

(5) a mono- or di-C₁₋₆ alkyl-carbamoyl group (e.g.,
methylcarbamoyl), or

(6) a 5- or 6-membered monocyclic aromatic heterocyclic group
20 (e.g., pyrazolyl, pyridyl, triazolyl) optionally substituted by
1 to 3 C₁₋₆ alkyl groups (e.g., methyl); and

Ring A is a 6-membered aromatic ring (e.g., a benzene
ring, a pyridine ring) optionally further substituted by 1 to 3
substituents, in addition to R⁴, selected from

25 (1) a halogen atom (e.g., a fluorine atom), and

(2) a C₁₋₆ alkoxy group (e.g., methoxy).

[0190]

[Compound Ca-2]

Compound (I) wherein

30 R¹ is

(1) a phenyl group, a C₅₋₆ cycloalkyl group (cyclopentyl,
cyclohexyl) or a 5- or 6-membered monocyclic non-aromatic
heterocyclic group (e.g., tetrahydropyranyl), each of which is
optionally substituted by 1 to 3 substituents selected from

35 (i) a halogen atom (e.g., a fluorine atom),

(ii) a cyano group, and
(iii) a hydroxy group, or
(2) a C₁₋₆ alkyl group (e.g., methyl, isobutyl) optionally substituted by 1 to 3 substituents selected from

- 5 (i) a hydroxy group, and
(ii) a 3- to 14-membered non-aromatic heterocyclic group (preferably a 3- to 8-membered monocyclic non-aromatic heterocyclic group) (e.g., tetrahydrofuryl);

R² is

- 10 (1) a hydrogen atom,
(2) a halogen atom (e.g., a chlorine atom, a fluorine atom), or
(3) a C₁₋₆ alkyl group (e.g., methyl);

R³ is

- (1) a hydrogen atom,
15 (2) a halogen atom (e.g., a chlorine atom),
(3) a cyano group,
(4) a C₁₋₆ alkyl group (e.g., methyl) optionally substituted by 1 to 3 halogen atoms (e.g., a fluorine atom),
(5) a C₁₋₆ alkoxy group (e.g., methoxy), or
20 (6) a C₃₋₆ cycloalkyl group (e.g., cyclopropyl);

R⁴ is


- (1) a halogen atom (e.g., a chlorine atom, a fluorine atom),
(2) a cyano group,
(3) a C₁₋₆ alkyl group (e.g., methyl),
25 (4) a C₁₋₆ alkoxy group (e.g., methoxy) optionally substituted by 1 to 3 halogen atoms (e.g., a fluorine atom),
(5) a mono- or di-C₁₋₆ alkyl-carbamoyl group (e.g., methylcarbamoyl), or
(6) a 5- or 6-membered monocyclic aromatic heterocyclic group
30 (e.g., pyrazolyl, pyridyl, triazolyl) optionally substituted by 1 to 3 C₁₋₆ alkyl groups (e.g., methyl);

Ring A is a 6-membered aromatic ring (e.g., a benzene ring, a pyridine ring) optionally further substituted by 1 to 3 substituents, in addition to R⁴, selected from

- 35 (1) a halogen atom (e.g., a fluorine atom), and

(2) a C₁₋₆ alkoxy group (e.g., methoxy); and
the partial structure represented by the following
formula:


[0191]



[0192]

in the formula (I) is a partial structure represented by the
following formula:

[0193]



[0194]

[Compound C-3]

Compound (I) wherein

R¹ is

15 (1) a phenyl group optionally substituted by 1 to 3
substituents selected from

- (i) a halogen atom (e.g., a fluorine atom), and
- (ii) a cyano group,

(2) a C₅₋₆ cycloalkyl group (cyclopentyl, cyclohexyl) optionally
20 substituted by 1 to 3 hydroxy groups,

(3) a 5- or 6-membered monocyclic non-aromatic heterocyclic
group (e.g., tetrahydropyranyl) optionally substituted by 1 to
3 hydroxy groups, or

(4) a C₁₋₆ alkyl group (e.g., methyl, isobutyl) optionally
25 substituted by 1 to 3 substituents selected from

- (i) a hydroxy group, and
- (ii) a 3- to 8-membered monocyclic non-aromatic
heterocyclic group (e.g., tetrahydrofuryl);

R² is

30 (1) a hydrogen atom,

- (2) a halogen atom (e.g., a chlorine atom, a fluorine atom), or
(3) a C₁₋₆ alkyl group (e.g., methyl);

R³ is

- (1) a hydrogen atom,
5 (2) a halogen atom (e.g., a chlorine atom),
(3) a cyano group,
(4) a C₁₋₆ alkyl group (e.g., methyl) optionally substituted by
1 to 3 halogen atoms (e.g., a fluorine atom),
(5) a C₁₋₆ alkoxy group (e.g., methoxy), or
10 (6) a C₃₋₆ cycloalkyl group (e.g., cyclopropyl);

R⁴ is

- (1) a halogen atom (e.g., a fluorine atom, a chlorine atom),
(2) a cyano group,
(3) a C₁₋₆ alkyl group (e.g., methyl),
15 (4) a C₁₋₆ alkoxy group (e.g., methoxy) optionally substituted
by 1 to 3 halogen atoms (e.g., a fluorine atom),
(5) a mono- or di-C₁₋₆ alkyl-carbamoyl group (e.g.,
methylcarbamoyl), or
(6) a 5- or 6-membered monocyclic aromatic heterocyclic group
20 (e.g., pyrazolyl, pyridyl, triazolyl) optionally substituted by
1 to 3 C₁₋₆ alkyl groups (e.g., methyl); and

Ring A is a 6-membered aromatic ring (e.g., a benzene
ring, a pyridine ring) optionally further substituted by 1 to 3
substituents, in addition to R⁴, selected from

- 25 (a) a halogen atom (e.g., a fluorine atom), and
(b) a C₁₋₆ alkoxy group (e.g., methoxy).

[0195]

[Compound Ca-3]

Compound (I) wherein

30 R¹ is

- (1) a phenyl group optionally substituted by 1 to 3
substituents selected from

- (i) a halogen atom (e.g., a fluorine atom), and
(ii) a cyano group,

- 35 (2) a C₅₋₆ cycloalkyl group (cyclopentyl, cyclohexyl) optionally

substituted by 1 to 3 hydroxy groups,

(3) a 5- or 6-membered monocyclic non-aromatic heterocyclic group (e.g., tetrahydropyranyl) optionally substituted by 1 to 3 hydroxy groups, or

5 (4) a C₁₋₆ alkyl group (e.g., methyl, isobutyl) optionally substituted by 1 to 3 substituents selected from

(i) a hydroxy group, and

(ii) a 3- to 8-membered monocyclic non-aromatic heterocyclic group (e.g., tetrahydrofuryl);

10 R² is

(1) a hydrogen atom,

(2) a halogen atom (e.g., a chlorine atom, a fluorine atom), or

(3) a C₁₋₆ alkyl group (e.g., methyl);

R³ is

15 (1) a hydrogen atom,

(2) a halogen atom (e.g., a chlorine atom),

(3) a cyano group,

(4) a C₁₋₆ alkyl group (e.g., methyl) optionally substituted by 1 to 3 halogen atoms (e.g., a fluorine atom),

20 (5) a C₁₋₆ alkoxy group (e.g., methoxy), or

(6) a C₃₋₆ cycloalkyl group (e.g., cyclopropyl);

R⁴ is

(1) a halogen atom (e.g., a fluorine atom, a chlorine atom),

(2) a cyano group,

25 (3) a C₁₋₆ alkyl group (e.g., methyl),

(4) a C₁₋₆ alkoxy group (e.g., methoxy) optionally substituted by 1 to 3 halogen atoms (e.g., a fluorine atom),

(5) a mono- or di-C₁₋₆ alkyl-carbamoyl group (e.g., methylcarbamoyl), or

30 (6) a 5- or 6-membered monocyclic aromatic heterocyclic group (e.g., pyrazolyl, pyridyl, triazolyl) optionally substituted by 1 to 3 C₁₋₆ alkyl groups (e.g., methyl);

Ring A is a 6-membered aromatic ring (e.g., a benzene ring, a pyridine ring) optionally further substituted by 1 to 3


35 substituents, in addition to R⁴, selected from

(a) a halogen atom (e.g., a fluorine atom), and

(b) a C₁₋₆ alkoxy group (e.g., methoxy); and

the partial structure represented by the following formula:


5 [0196]



[0197]

in the formula (I) is a partial structure represented by the following formula:

10 [0198]



[0199]

[Compound C-4]

Compound (I) wherein

15 R¹ is

(1) a phenyl group optionally substituted by 1 to 3 substituents selected from

(i) a halogen atom (e.g., a fluorine atom), and

(ii) a cyano group,

20 (2) a C₅₋₆ cycloalkyl group (cyclopentyl, cyclohexyl) optionally substituted by 1 to 3 hydroxy groups,

(3) a tetrahydropyranyl group optionally substituted by 1 to 3 hydroxy groups, or

(4) a C₁₋₆ alkyl group (e.g., methyl, isobutyl) optionally substituted by 1 to 3 substituents selected from

(i) a hydroxy group, and

(ii) a tetrahydrofuryl group;

R² is

(1) a hydrogen atom,

30 (2) a halogen atom (e.g., a chlorine atom, a fluorine atom), or

(3) a C₁₋₆ alkyl group (e.g., methyl);

R³ is

(1) a hydrogen atom,

(2) a halogen atom (e.g., a chlorine atom),

5 (3) a cyano group,

(4) a C₁₋₆ alkyl group (e.g., methyl) optionally substituted by
1 to 3 halogen atoms (e.g., a fluorine atom),

(5) a C₁₋₆ alkoxy group (e.g., methoxy), or

(6) a C₃₋₆ cycloalkyl group (e.g., cyclopropyl);

10 R⁴ is

(1) a halogen atom (e.g., a fluorine atom, a chlorine atom),

(2) a cyano group,

(3) a C₁₋₆ alkyl group (e.g., methyl),

(4) a C₁₋₆ alkoxy group (e.g., methoxy) optionally substituted
15 by 1 to 3 halogen atoms (e.g., a fluorine atom),

(5) a mono- or di-C₁₋₆ alkyl-carbamoyl group (e.g.,
methylcarbamoyl),

(6) a pyrazolyl group optionally substituted by 1 to 3 C₁₋₆
alkyl groups (e.g., methyl),

20 (7) a pyridyl group optionally substituted by 1 to 3 C₁₋₆ alkyl
groups (e.g., methyl), or

(8) a triazolyl group optionally substituted by 1 to 3 C₁₋₆
alkyl groups (e.g., methyl); and

Ring A is

25 (1) a benzene ring optionally further substituted by 1 to 3
substituents, in addition to R⁴, selected from

(a) a halogen atom (e.g., a fluorine atom), and

(b) a C₁₋₆ alkoxy group (e.g., methoxy), or

(2) a pyridine ring.

30 [0200]

[Compound Ca-4]

Compound (I) wherein

R¹ is

(1) a phenyl group optionally substituted by 1 to 3
35 substituents selected from

- (i) a halogen atom (e.g., a fluorine atom), and
(ii) a cyano group,
(2) a C₅₋₆ cycloalkyl group (cyclopentyl, cyclohexyl) optionally substituted by 1 to 3 hydroxy groups,
5 (3) a tetrahydropyranyl group optionally substituted by 1 to 3 hydroxy groups, or
(4) a C₁₋₆ alkyl group (e.g., methyl, isobutyl) optionally substituted by 1 to 3 substituents selected from
(i) a hydroxy group, and
10 (ii) a tetrahydrofuryl group;
R² is
(1) a hydrogen atom,
(2) a halogen atom (e.g., a chlorine atom, a fluorine atom), or
(3) a C₁₋₆ alkyl group (e.g., methyl);
15 R³ is
(1) a hydrogen atom,
(2) a halogen atom (e.g., a chlorine atom),
(3) a cyano group,
(4) a C₁₋₆ alkyl group (e.g., methyl) optionally substituted by
20 1 to 3 halogen atoms (e.g., a fluorine atom),
(5) a C₁₋₆ alkoxy group (e.g., methoxy), or
(6) a C₃₋₆ cycloalkyl group (e.g., cyclopropyl);
R⁴ is
(1) a halogen atom (e.g., a fluorine atom, a chlorine atom),
25 (2) a cyano group,
(3) a C₁₋₆ alkyl group (e.g., methyl),
(4) a C₁₋₆ alkoxy group (e.g., methoxy) optionally substituted by 1 to 3 halogen atoms (e.g., a fluorine atom),
(5) a mono- or di-C₁₋₆ alkyl-carbamoyl group (e.g.,
30 methylcarbamoyl),
(6) a pyrazolyl group optionally substituted by 1 to 3 C₁₋₆ alkyl groups (e.g., methyl),
(7) a pyridyl group optionally substituted by 1 to 3 C₁₋₆ alkyl groups (e.g., methyl), or
35 (8) a triazolyl group optionally substituted by 1 to 3 C₁₋₆

alkyl groups (e.g., methyl);

Ring A is

(1) a benzene ring optionally further substituted by 1 to 3 substituents, in addition to R⁴, selected from


5 (a) a halogen atom (e.g., a fluorine atom), and

(b) a C₁₋₆ alkoxy group (e.g., methoxy), or

(2) a pyridine ring; and

the partial structure represented by the following formula:


10 [0201]



[0202]

in the formula (I) is a partial structure represented by the following formula:

15 [0203]



[0204]

[Compound D-1]

Compound (I) wherein

20 R¹ is

(1) a C₅₋₆ cycloalkyl group (e.g., cyclohexyl) optionally substituted by 1 to 3 hydroxy groups, or

(2) a 5- or 6-membered monocyclic non-aromatic heterocyclic group (e.g., tetrahydropyranyl) optionally substituted by 1 to

25 3 hydroxy groups;

R² is

(1) a halogen atom (e.g., a fluorine atom), or

(2) a C₁₋₆ alkyl group (e.g., methyl);

R³ is a C₁₋₆ alkyl group (e.g., methyl);

30 R⁴ is

- (1) a C₁₋₆ alkyl group (e.g., methyl),
- (2) a C₁₋₆ alkoxy group (e.g., methoxy), or
- (3) a 5- or 6-membered monocyclic aromatic heterocyclic group (e.g., pyrazolyl); and

5 Ring A is a benzene ring or a pyridine ring, each of which is unsubstituted, in addition to R⁴.

[0205]

[Compound Da-1]

Compound (I) wherein

10 R¹ is

- (1) a C₅₋₆ cycloalkyl group (e.g., cyclohexyl) optionally substituted by 1 to 3 hydroxy groups, or
 - (2) a 5- or 6-membered monocyclic non-aromatic heterocyclic group (e.g., tetrahydropyranyl) optionally substituted by 1 to
- 15 3 hydroxy groups;

R² is

- (1) a halogen atom (e.g., a fluorine atom), or
- (2) a C₁₋₆ alkyl group (e.g., methyl);

R³ is a C₁₋₆ alkyl group (e.g., methyl);


20 R⁴ is

- (1) a C₁₋₆ alkyl group (e.g., methyl),
- (2) a C₁₋₆ alkoxy group (e.g., methoxy), or
- (3) a 5- or 6-membered monocyclic aromatic heterocyclic group (e.g., pyrazolyl);

25 Ring A is a benzene ring or a pyridine ring, each of which is unsubstituted, in addition to R⁴; and

the partial structure represented by the following formula:


[0206]



[0207]

in the formula (I) is a partial structure represented by the following formula:

[0208]



[0209]

[Compound D-2]

5 Compound (I) wherein

R^1 is

- (1) a cyclohexyl group substituted by one hydroxy group, or
- (2) a tetrahydropyranyl group substituted by one hydroxy group;

R^2 is

- 10 (1) a halogen atom (e.g., a fluorine atom), or
- (2) a C_{1-6} alkyl group (e.g., methyl);

R^3 is a C_{1-6} alkyl group (e.g., methyl);

R^4 is

- (1) a C_{1-6} alkyl group (e.g., methyl),
- 15 (2) a C_{1-6} alkoxy group (e.g., methoxy), or
- (3) a pyrazolyl group; and

Ring A is a benzene ring or a pyridine ring, each of which is unsubstituted, in addition to R^4 .

[0210]

20 [Compound Da-2]

Compound (I) wherein

R^1 is

- (1) a cyclohexyl group substituted by one hydroxy group, or
- (2) a tetrahydropyranyl group substituted by one hydroxy group;

25 R^2 is

- (1) a halogen atom (e.g., a fluorine atom), or
- (2) a C_{1-6} alkyl group (e.g., methyl);

R^3 is a C_{1-6} alkyl group (e.g., methyl);


R^4 is

- 30 (1) a C_{1-6} alkyl group (e.g., methyl),
- (2) a C_{1-6} alkoxy group (e.g., methoxy), or
 - (3) a pyrazolyl group;

Ring A is a benzene ring or a pyridine ring, each of which is unsubstituted, in addition to R⁴; and

the partial structure represented by the following formula:


5 [0211]



[0212]

in the formula (I) is a partial structure represented by the following formula:

10 [0213]



[0214]

[Compound E]

2-[(3S,4S)-4-Hydroxytetrahydro-2H-pyran-3-yl]-6-(4-methoxybenzyl)-4,5-dimethyl-2,3-dihydro-1H-isoindol-1-one, or a salt thereof

4-Fluoro-2-[(3S,4S)-4-hydroxytetrahydro-2H-pyran-3-yl]-5-methyl-6-[4-(1H-pyrazol-1-yl)benzyl]-2,3-dihydro-1H-isoindol-1-one, or a salt thereof

20 2-((1S,2S)-2-Hydroxycyclohexyl)-4,5-dimethyl-6-((6-methylpyridin-3-yl)methyl)isoindolin-1-one, or a salt thereof

[0215]

When compound (I) is in a form of a salt, examples of such salt include salts with inorganic base, an ammonium salt, salts with organic base, salts with inorganic acid, salts with organic acid, salts with basic or acidic amino acid, and the like.

[0216]

Preferable examples of the salt with inorganic base include alkali metal salts such as sodium salt, potassium salt

and the like; alkaline earth metal salts such as calcium salt, magnesium salt, barium salt and the like; an aluminum salt, and the like.

[0217]

5 Preferable examples of the salt with organic base include salts with trimethylamine, triethylamine, pyridine, picoline, ethanolamine, diethanolamine, triethanolamine, dicyclohexylamine, N,N'-dibenzylethylenediamine and the like.

[0218]

10 Preferable examples of the salt with inorganic acid include salts with hydrochloric acid, hydrobromic acid, nitric acid, sulfuric acid, phosphoric acid and the like.

[0219]

 Preferable examples of the salt with organic acid include
15 salts with formic acid, acetic acid, trifluoroacetic acid, fumaric acid, oxalic acid, tartaric acid, maleic acid, citric acid, succinic acid, malic acid, methanesulfonic acid, benzenesulfonic acid, p-toluenesulfonic acid and the like.

[0220]

20 Preferable examples of the salt with basic amino acid include salts with arginine, lysine, ornithine and the like.

[0221]

 Preferable examples of the salt with acidic amino acid include salts with aspartic acid, glutamic acid and the like.

25 [0222]

 Among these salts, a pharmaceutically acceptable salt is preferable. When a compound has a basic functional group, preferable examples of the pharmaceutically acceptable salt include salts with inorganic acid such as hydrochloric acid,
30 hydrobromic acid, nitric acid, sulfuric acid, phosphoric acid and the like, and salts with organic acid such as acetic acid, phthalic acid, fumaric acid, oxalic acid, tartaric acid, maleic acid, citric acid, succinic acid, methanesulfonic acid, p-toluenesulfonic acid and the like. In addition, when a
35 compound has an acidic functional group, examples thereof

include inorganic salts such as alkali metal salts (e.g., sodium salt, potassium salt etc.), alkaline earth metal salts (e.g., calcium salt, magnesium salt, barium salt etc.) and the like, ammonium salt and the like.

5 [0223]

Compound (I) may be a crystal, and both a single crystal and crystal mixtures are encompassed in the compound (I).

Compound (I) may be a pharmaceutically acceptable cocrystal or cocrystal salt. Here, the cocrystal or cocrystal
10 salt means a crystalline substance consisting of two or more particular substances which are solids at room temperature, each having different physical properties (e.g., structure, melting point, heat of melting, hygroscopicity, solubility, stability etc.). The cocrystal and cocrystal salt can be
15 produced by cocrystallization method known *per se*.

Compound (I) encompasses solvates (e.g., hydrate) and non-solvates within the scope thereof. Compound (I) may be a compound labeled or substituted with an isotope (e.g., ^2H , ^3H , ^{11}C , ^{14}C , ^{18}F , ^{35}S , ^{125}I). A compound labeled with or substituted
20 by an isotope can be used, for example, as a tracer used for Positron Emission Tomography (PET) (PET tracer), and is useful in the field of medical diagnosis and the like.

[0224]

When compound (I) of the present invention has an
25 asymmetric center, isomers such as enantiomer, diastereomer and the like may be present. Such isomers and a mixture thereof are all encompassed within the scope of the present invention. When an isomer is formed due to the conformation or tautomerism, such isomers and a mixture thereof are all encompassed in
30 compound (I) of the present invention.

[0225]

The production methods of the compound of the present invention are explained below.

[0226]

35 The raw material compound and reagent used and the

compound obtained in each step in the following production method may be each in a form of a salt, and examples of such salt include those similar to the salts of the compound of the present invention and the like.

5 [0227]

When the compound obtained in each step is a free form, it can be converted to the objective salt according to a method known per se. When the compound obtained in each step is a salt, it can be converted to the objective free form or the
10 other salt according to a method known per se.

[0228]

The compound obtained in each step can be used directly as the reaction mixture or as a crude product for the next reaction. Alternatively, the compound obtained in each step
15 can be isolated and purified from a reaction mixture according to a method known per se, for example, a separation means such as concentration, crystallization, recrystallization, distillation, solvent extraction, fractional distillation, column chromatography and the like.

20 [0229]

When the raw material compound and reagent used in each step are commercially available, the commercially available product can also be used directly.

[0230]

25 In the reaction in each step, while the reaction time varies depending on the kind of the reagent and solvent to be used, it is generally 1 min - 48 hr, preferably 10 min - 8 hr, unless otherwise specified.

[0231]

30 In the reaction in each step, while the reaction temperature varies depending on the kind of the reagent and solvent to be used, it is generally -78°C - 300°C, preferably -78°C - 150°C, unless otherwise specified.

[0232]

35 In the reaction in each step, while the pressure varies

depending on the kind of the reagent and solvent to be used, it is generally 1 atm - 20 atm, preferably 1 atm - 3 atm, unless otherwise specified.

[0233]

5 Microwave synthesizer such as Initiator manufactured by Biotage and the like may be used for the reaction in each step. While the reaction temperature varies depending on the kind of the reagent and solvent to be used, it is generally room temperature - 300°C, preferably 50°C - 250°C, unless otherwise
10 specified. While the reaction time varies depending on the kind of the reagent and solvent to be used, it is generally 1 min - 48 hr, preferably 1 min - 8 hr, unless otherwise specified.

[0234]

15 In the reaction in each step, the reagent is used in an amount of 0.5 equivalents - 20 equivalents, preferably 0.8 equivalents - 5 equivalents, relative to the substrate, unless otherwise specified. When the reagent is used as a catalyst, the reagent is used in an amount of 0.001 equivalents - 1
20 equivalents, preferably 0.01 equivalents - 0.2 equivalents, relative to the substrate. When the reagent is used as a reaction solvent, the reagent is used in a solvent amount.

[0235]

Unless otherwise specified, the reaction in each step is
25 carried out without solvent, or by dissolving or suspending the raw material compound in a suitable solvent. Examples of the solvent include those described in Examples and the following solvents.

alcohols: methanol, ethanol, tert-butyl alcohol, 2-
30 methoxyethanol and the like;

ethers: diethyl ether, diphenyl ether, tetrahydrofuran, 1,2-dimethoxyethane and the like;

aromatic hydrocarbons: chlorobenzene, toluene, xylene and the like;

35 saturated hydrocarbons: cyclohexane, hexane and the like;

amides: N,N-dimethylformamide, N-methylpyrrolidone and the like;

halogenated hydrocarbons: dichloromethane, carbon tetrachloride and the like;

5 nitriles: acetonitrile and the like;

sulfoxides: dimethyl sulfoxide and the like;

aromatic organic bases: pyridine and the like;

anhydrides: acetic anhydride and the like;

organic acids: formic acid, acetic acid, trifluoroacetic acid

10 and the like;

inorganic acids: hydrochloric acid, sulfuric acid and the like;

esters: ethyl acetate and the like;

ketones: acetone, methyl ethyl ketone and the like;

water.

15 The above-mentioned solvent can be used in a mixture of two or more kinds thereof in an appropriate ratio.

[0236]

When a base is used for the reaction in each step, examples thereof include those described in Examples and the
20 following bases.

inorganic bases: sodium hydroxide, magnesium hydroxide and the like;

basic salts: sodium carbonate, calcium carbonate, sodium hydrogen carbonate and the like;

25 organic bases: triethylamine, diethylamine, pyridine, 4-dimethylaminopyridine, N,N-dimethylaniline, 1,4-diazabicyclo[2.2.2]octane, 1,8-diazabicyclo[5.4.0]-7-undecene, imidazole, piperidine and the like;

metal alkoxides: sodium ethoxide, potassium tert-butoxide and
30 the like;

alkali metal hydrides: sodium hydride and the like;

metal amides: sodium amide, lithium diisopropylamide, lithium hexamethyldisilazide and the like;

organic lithiums: n-butyllithium and the like.

35 [0237]

When an acid or an acid catalyst is used for the reaction in each step, examples thereof include those described in Examples and the following acids and acid catalysts.

inorganic acids: hydrochloric acid, sulfuric acid, nitric acid,
5 hydrobromic acid, phosphoric acid and the like;
organic acids: acetic acid, trifluoroacetic acid, citric acid,
p-toluenesulfonic acid, 10-camphorsulfonic-acid and the like;
Lewis acid: boron trifluoride diethyl ether complex, zinc
iodide, anhydrous aluminium chloride, anhydrous zinc chloride,
10 anhydrous iron chloride and the like.

[0238]

Unless otherwise specified, the reaction in each step is carried out according to a method known per se, for example, the method described in Jikken Kagaku Kouza, 5th Edition,
15 vol.13-19 (the Chemical Society of Japan ed.); Shin Jikken Kagaku Kouza, vol.14-15 (the Chemical Society of Japan ed.); Fine Organic Chemistry, Revised 2nd Edition (L. F. Tietze, Th. Eicher, Nankodo); Organic Name Reactions, the Reaction Mechanism and Essence, Revised Edition (Hideo Togo, Kodansha);
20 ORGANIC SYNTHESSES Collective Volume I-VII (John Wiley & Sons Inc); Modern Organic Synthesis in the Laboratory A Collection of Standard Experimental Procedures (Jie Jack Li, OXFORD UNIVERSITY); Comprehensive Heterocyclic Chemistry III, Vol.1 -Vol.14 (Elsevier Japan); Strategic Applications of Named
25 Reactions in Organic Synthesis (translated by Kiyoshi Tomioka, Kagakudojin); Comprehensive Organic Transformations (VCH Publishers Inc.), 1989, or the like, or the method described in Examples.

[0239]

30 In each step, protection or deprotection of functional groups is performed according to a method known per se, for example, the methods described in Wiley-Interscience, 2007, "Protective Groups in Organic Synthesis, 4th Ed." (Theodora W. Greene, Peter G. M. Wuts); Thieme, 2004, "Protecting Groups 3rd
35 Ed." (P. J. Kocienski) and the like, or the methods described

in the Examples.

Examples of the protecting group for hydroxyl group of alcohol and the like and phenolic hydroxyl group include ether-type protecting groups such as methoxymethyl ether, benzyl
5 ether, t-butyldimethylsilyl ether, tetrahydropyranyl ether and the like; carboxylate-type protecting groups such as acetate and the like; sulfonate-type protecting groups such as methanesulfonate and the like; carbonate-type protecting groups such as t-butyl carbonate and the like; and the like.

10 Examples of the protecting group for carbonyl group of aldehyde include acetal-type protecting groups such as dimethyl acetal and the like; cyclic acetal-type protecting groups such as cyclic 1,3-dioxane and the like; and the like.

Examples of the protecting group for carbonyl group of
15 ketone include ketal-type protecting groups such as dimethyl ketal and the like; cyclic ketal-type protecting groups such as cyclic 1,3-dioxane and the like; oxime-type protecting groups such as O-methyloxime and the like; hydrazone-type protecting groups such as N,N-dimethylhydrazone and the like; and the like.

20 Examples of the protecting group for carboxyl group include ester-type protecting groups such as methyl ester and the like; amide-type protecting groups such as N,N-dimethylamide and the like; and the like.

Examples of the protecting group for thiol include ether-
25 type protecting groups such as benzyl thioether and the like; ester-type protecting groups such as thioacetate, thiocarbonate, thiocarbamate and the like; and the like.

Examples of the protecting group for amino group, and aromatic heterocycle such as imidazole, pyrrole, indole and the
30 like include carbamate-type protecting groups such as benzyl carbamate, tert-butyl carbamate and the like; amide-type protecting groups such as acetamide and the like; alkylamine-type protecting groups such as N-triphenylmethylaniline and the like; sulfonamide-type protecting groups such as
35 methanesulfonamide and the like; and the like.

Protecting groups can be removed by a method known per se, for example, methods using acid, base, ultraviolet rays, hydrazine, phenylhydrazine, sodium N-methyldithiocarbamate, tetrabutylammonium fluoride, palladium acetate, trialkylsilyl
5 halide (e.g., trimethylsilyl iodide, trimethylsilyl bromide), reduction methods and the like.

[0240]

When reduction reaction is carried out in each step, examples of the reducing agent to be used include metal
10 hydrides such as lithium aluminium hydride, sodium triacetoxyborohydride, sodium cyanoborohydride, diisobutylaluminium hydride (DIBAL-H), sodium borohydride, tetramethylammonium triacetoxyborohydride and the like; boranes such as borane tetrahydrofuran complex and the like; Raney
15 nickel; Raney cobalt; hydrogen; formic acid; triethylsilane and the like. When carbon-carbon double bond or triple bond is reduced, a method using a catalyst such as palladium-carbon, Lindlar's catalyst and the like may be employed.

[0241]

20 When oxidation reaction is carried out in each step, examples of the oxidizing agent to be used include peroxides such as m-chloroperbenzoic acid (mCPBA), hydrogen peroxide, t-butylhydroperoxide and the like; perchlorates such as tetrabutylammonium perchlorate and the like; chlorates such as
25 sodium chlorate and the like; chlorites such as sodium chlorite and the like; periodic acids such as sodium periodate and the like; hypervalent iodine reagents such as iodosylbenzene and the like; reagents containing manganese such as manganese dioxide, potassium permanganate and the like; leads such as
30 lead tetraacetate and the like; reagents containing chromium such as pyridinium chlorochromate (PCC), pyridinium dichromate (PDC), Jones reagent and the like; halogen compounds such as N-bromosuccinimide (NBS) and the like; oxygen; ozone; sulfur trioxide-pyridine complex; osmium tetroxide; selenium dioxide;
35 2,3-dichloro-5,6-dicyano-1,4-benzoquinone (DDQ) and the like.

[0242]

When radical cyclization reaction is carried out in each step, examples of the radical initiator to be used include azo compounds such as azobisisobutyronitrile (AIBN) and the like; 5 water-soluble radical initiators such as 4-4'-azobis-4-cyanopentanoic acid (ACPA) and the like; triethylboron in the presence of air or oxygen; benzoyl peroxide and the like. Examples of the radical reagent to be used include tributylstannane, tris(trimethylsilyl)silane, 1,1,2,2- 10 tetraphenyldisilane, diphenylsilane, samarium iodide and the like.

[0243]

When Wittig reaction is carried out in each step, examples of the Wittig reagent to be used include alkylidene 15 phosphoranes and the like. The alkylidene phosphoranes can be prepared according to a method known per se, for example, by reacting a phosphonium salt with a strong base.

[0244]

When Horner-Emmons reaction is carried out in each step, 20 examples of the reagent to be used include phosphonoacetates such as methyl dimethylphosphonoacetate, ethyl diethylphosphonoacetate and the like; and bases such as alkali metal hydrides, organic lithiums and the like.

[0245]

25 When Friedel-Crafts reaction is carried out in each step, a combination of a Lewis acid and an acid chloride or a combination of a Lewis acid and an alkylating agent (e.g., an alkyl halide, an alcohol, an olefin etc.) is used as a reagent. Alternatively, an organic acid or an inorganic acid can also be 30 used instead of a Lewis acid, and an anhydride such as acetic anhydride and the like can also be used instead of an acid chloride.

[0246]

When aromatic nucleophilic substitution reaction is 35 carried out in each step, a nucleophile (e.g., an amine,

imidazole etc.) and a base (e.g., a basic salt, an organic base etc.) are used as a reagent.

[0247]

When nucleophilic addition reaction by a carbo anion,
5 nucleophilic 1,4-addition reaction (Michael addition reaction)
by a carbo anion or nucleophilic displacement reaction by a
carbo anion is carried out in each step, examples of the base
to be used for generation of the carbo anion include organic
lithiums, metal alkoxides, inorganic bases, organic bases and
10 the like.

[0248]

When Grignard reagent is carried out in each step,
examples of the Grignard reagent to be used include
arylmagnesium halides such as phenylmagnesium bromide and the
15 like; and alkylmagnesium halides such as methylmagnesium
bromide and the like. The Grignard reagent can be prepared
according to a method known per se, for example, by reacting an
alkyl halide or an aryl halide with a metal magnesium in an
ether or tetrahydrofuran as a solvent.

20 [0249]

When Knoevenagel condensation reaction is carried out in
each step, a compound having an activated methylene group with
two electron withdrawing groups (e.g., malonic acid, diethyl
malonate, malononitrile etc.) and a base (e.g., an organic base,
25 a metal alkoxide, an inorganic base) are used as a reagent.

[0250]

When Vilsmeier-Haack reaction is carried out in each step,
phosphoryl chloride and an amide derivative (e.g., N,N-
dimethylformamide etc.) are used as a reagent.

30 [0251]

When azidation reaction of an alcohol, an alkyl halide or
a sulfonate is carried out in each step, examples of the
azidating agent to be used include diphenylphosphorylazide
(DPPA), trimethylsilylazide, sodium azide and the like. For
35 example, for the azidation reaction of an alcohol, a method

using diphenylphosphorylazide and 1,8-diazabicyclo[5.4.0]undec-7-ene (DBU), a method using trimethylsilylazide and a Lewis acid, and the like are employed.

[0252]

5 When reductive amination reaction is carried out in each step, examples of the reducing agent to be used include sodium triacetoxyborohydride, sodium cyanoborohydride, hydrogen, formic acid and the like. When the substrate is an amine compound, examples of the carbonyl compound to be used include
10 paraformaldehyde, aldehydes such as acetaldehyde and the like, and ketones such as cyclohexanone and the like. When the substrate is a carbonyl compound, examples of the amine to be used include ammonia, primary amines such as methylamine and the like; secondary amines such as dimethylamine and the like,
15 and the like.

[0253]

When Mitsunobu reaction is carried out in each step, an azodicarboxylate (e.g., diethyl azodicarboxylate (DEAD), diisopropyl azodicarboxylate (DIAD) etc.) and
20 triphenylphosphine are used as a reagent.

[0254]

When esterification reaction, amidation reaction or ureation reaction is carried out in each step, examples of the reagent to be used include acyl halides such as acid chlorides,
25 acid bromides and the like; activated carboxylic acids such as anhydrides, activated esters, sulfates and the like. Examples of the activating agent of the carboxylic acid include carbodiimide condensing agents such as 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide hydrochloride (WSCD) and the
30 like; triazine condensing agents such as 4-(4,6-dimethoxy-1,3,5-triazin-2-yl)-4-methylmorpholinium chloride n-hydrate (DMT-MM) and the like; carbonate condensing agents such as 1,1-carbonyldiimidazole (CDI) and the like; diphenylphosphoryl azide (DPPA); benzotriazol-1-yloxy-trisdimethylaminophosphonium
35 salt (BOP reagent); 2-chloro-1-methyl-pyridinium iodide

(Mukaiyama reagent); thionyl chloride; lower alkyl haloformates such as ethyl chloroformate and the like; O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate (HATU); sulfuric acid; combinations thereof and the like. When
5 carbodiimide condensing agent is used, an additive such as 1-hydroxybenzotriazole (HOBt), N-hydroxysuccinimide (HOSu), dimethylaminopyridine (DMAP) and the like may be added to the reaction system.

[0255]

10 When coupling reaction is carried out in each step, examples of the metal catalyst to be used include palladium compounds such as palladium(II) acetate, tetrakis(triphenylphosphine)palladium(0), dichlorobis(triphenylphosphine)palladium(II),
15 dichlorobis(triethylphosphine)palladium(II), tris(dibenzylideneacetone)dipalladium(0), 1,1'-bis(diphenylphosphino)ferrocene palladium(II) chloride, palladium(II) acetate and the like; nickel compounds such as tetrakis(triphenylphosphine)nickel(0) and the like; rhodium
20 compounds such as tris(triphenylphosphine)rhodium(III) chloride and the like; cobalt compounds; copper compounds such as copper oxide, copper(I) iodide and the like; platinum compounds and the like. In addition, a base can be added to the reaction system, and examples thereof include inorganic bases, basic
25 salts and the like.

[0256]

When thiocarbonylation reaction is carried out in each step, phosphorus pentasulfide is typically used as the thiocarbonylating agent. Alternatively, a reagent having a
30 1,3,2,4-dithiadiphosphetane-2,4-disulfide structure (e.g., 2,4-bis(4-methoxyphenyl)-1,3,2,4-dithiadiphosphetane-2,4-disulfide (Lawesson reagent) etc.) can also be used instead of phosphorus pentasulfide.

[0257]

35 When Wohl-Ziegler reaction is carried out in each step,

examples of the halogenating agent to be used include N-iodosuccinimide, N-bromosuccinimide (NBS), N-chlorosuccinimide (NCS), bromine, sulfuryl chloride and the like. In addition, the reaction can be accelerated by subjecting a radical
5 initiator such as heat, light, benzoyl peroxide, azobisisobutyronitrile and the like to the reaction system reaction.

[0258]

When halogenation reaction of a hydroxy group is carried
10 out in each step, examples of the halogenating agent to be used include hydrohalic acids and acid halides of inorganic acids, specifically, hydrochloric acid, thionyl chloride, phosphorus oxychloride and the like for chlorination, 48% hydrobromic acid and the like for bromination. In addition, a method of
15 producing an alkyl halide by reacting an alcohol with triphenylphosphine and carbon tetrachloride or carbon tetrabromide or the like can be employed. Alternatively, a method of producing an alkyl halide via two step comprising converting an alcohol to the corresponding sulfonate, and then
20 reacting the sulfonate with lithium bromide, lithium chloride or sodium iodide can also be employed.

[0259]

When Arbuzov reaction is carried out in each step, examples of the reagent to be used include alkyl halides such
25 as ethyl bromoacetate and the like; and phosphites such as triethyl phosphite, tri(isopropyl) phosphite and the like.

[0260]

When sulfonate esterification reaction is carried out in each step, examples of the sulfonating agent to be used include
30 methanesulfonyl chloride, p-toluenesulfonyl chloride, methanesulfonic anhydride, p-toluenesulfonic anhydride, N-phenylbis(trifluoromethanesulfonimide) and the like.

[0261]

When hydrolysis reaction is carried out in each step, an
35 acid or a base is used as a reagent. For acid hydrolysis

reaction of t-butyl ester, formic acid, triethylsilane and the like may be added to reductively-trap t-butyl cation which is by-produced.

[0262]


5 When dehydration reaction is carried out in each step, examples of the dehydrating agent to be used include sulfuric acid, diphosphorus pentoxide, phosphorus oxychloride, N,N'-dicyclohexylcarbodiimide, alumina, polyphosphoric acid and the like..

10 [0263]

Compound (I) can be produced from compound (1) or compound (2) according to the following method.

[0264]

Scheme (a)



15 [0265]

wherein X¹ is a C₁₋₆ alkyl group (e.g., methyl, ethyl), X² is a halogen atom (e.g., a fluorine atom), and the other symbols are as defined above.

[0266]

20 In Scheme (a), compound (3) can be produced by subjecting compound (1) to an oxidation reaction using an oxidant containing manganese such as potassium permanganate and the like, or subjecting compound (2) to a hydrolysis reaction.

Compound (4) can be produced by esterification reaction
 25 between compound (3) and alcohols under the acidic conditions with sulfuric acid.

Compound (5) can be produced by subjecting compound (4) to a halogenation reaction using a halogenating agent such as bromine and the like.

Compound (6) can be produced by subjecting compound (5) to a coupling reaction in the presence of a metal catalyst. The reaction may be a two step-reaction via a borate. The halide to be reacted with a borate can be produced according to a method known per se.

Compound (7) can be produced by subjecting compound (6) to a sulfonate esterification reaction using a sulfonating agent.

Compound (8) can be produced by subjecting compound (7) to a coupling reaction in the presence of a palladium compound.

Compound (9) can be produced by subjecting compound (8) to an oxidation reaction.


Compound (I) can be produced by subjecting compound (9) to a reductive amination reaction with an amine using a reducing agent.

[0267]

Compound (I) can also be produced from compound (6) according to the following method.

[0268]

Scheme (b)



[0269]

wherein each symbol is as defined above.

[0270]

In Scheme (b), compound (10) can be produced by subjecting compound (6) to a hydrolysis reaction using an acid or a base.

5 Compound (11) can be produced by activating compound (10) using an activating agent of a carboxylic acid, and then subjecting the resulting compound to an amidation reaction with an amine.

10 Compound (12) can be produced by subjecting compound (11) to a sulfonate esterification reaction using a sulfonating agent.

Compound (13) can be produced by subjecting compound (12) to a coupling reaction in the presence of a palladium compound.

15 Compound (14) can be produced by subjecting compound (13) to an oxidation reaction. In the reaction, the resulting aldehyde is cyclized. Examples of the oxidant to be used include those similar to the oxidant used in the step of producing compound (9) from compound (8) in Scheme (a).

20 Compound (I) can be produced by subjecting compound (14) to a reduction reaction.

[0271]

When compound (I) has an optical isomer, a stereoisomer, a regioisomer or a rotamer, these are also encompassed in compound (I), and can be obtained as a single product according to synthesis and separation methods known *per se*. For example, 25 when compound (I) contains an optical isomer, an optical isomer resolved from this compound is also encompassed in compound (I).

[0272]

The optical isomer can be produced according to a method 30 known *per se*. To be specific, the optical isomer is obtained using an optically active synthetic intermediate, or subjecting the final racemate product to an optical resolution according to a conventional method.

[0273]

35 For example, the method of optical resolution may be a

method known *per se*, such as a fractional recrystallization method, a chiral column method, a diastereomer method etc.

1) Fractional recrystallization method

A method wherein a salt with a racemate with an optically active compound (e.g., (+)-mandelic acid, (-)-mandelic acid, (+)-tartaric acid, (-)-tartaric acid, (+)-1-phenethylamine, (-)-1-phenethylamine, cinchonine, (-)-cinchonidine, brucine etc.) is formed, which is separated by a fractional recrystallization method, and if desired, a neutralization step to give a free optical isomer.

[0274]

2) Chiral column method

A method wherein a racemate or a salt thereof is applied to a column for separation of an optical isomer (a chiral column) to allow separation. In the case of a liquid chromatography, for example, a mixture of the optical isomers is applied to a chiral column such as ENANTIO-OVM (manufactured by Tosoh Corporation), CHIRAL series (manufactured by Daicel Corporation) and the like, and developed with water, various buffers (e.g., phosphate buffer, etc.) and organic solvents (e.g., ethanol, methanol, isopropanol, acetonitrile, trifluoroacetic acid, diethylamine etc.), solely or as a mixed solution thereof to separate the optical isomer.

[0275]

3) Diastereomer method

A method wherein a racemic mixture is prepared into a diastereomeric mixture by chemical reaction with an optically active reagent, which is made into a single substance by a typical separation means (e.g., a fractional recrystallization method, a chromatography method etc.) and the like, and is subjected to a chemical treatment such as hydrolysis and the like to remove an optically active reagent moiety, whereby an optical isomer is obtained. For example, when compound (I) contains hydroxy group, or primary or secondary amino group within a molecule, the compound and an optically active organic

acid (e.g., MTPA [α -methoxy- α -(trifluoromethyl)phenylacetic acid], (-)-menthoxyacetic acid etc.) and the like are subjected to condensation reaction to give diastereomers of the ester compound or the amide compound, respectively. When compound (I) 5 has a carboxylic acid group, this compound and an optically active amine or an optically active alcohol reagent are subjected to condensation reaction to give diastereomers of the amide compound or the ester compound, respectively. The separated diastereomer is converted to an optical isomer of the 10 original compound by acid hydrolysis or base hydrolysis.

[0276]

When compound (I) is obtained as a free compound, the compound can be converted to an objective salt according to a method known *per se* or a method analogous thereto. Conversely, 15 when it is obtained in the form of a salt, the salt can be converted to a free form or other objective salt according to a method known *per se* or a method analogous thereto.

[0277]

Compound (I) may be a prodrug, and the prodrug of 20 compound (I) refers to a compound which is converted to compound (I) as a result of a reaction with an enzyme, gastric acid, etc. under physiological conditions *in vivo*, thus a compound that undergoes enzymatic oxidation, reduction, hydrolysis etc. to convert to compound (I) and a compound that 25 undergoes hydrolysis and the like by gastric acid, etc. to convert to compound (I).

[0278]

Examples of the prodrug for compound (I) include a compound obtained by subjecting an amino group in compound 30 (I) to acylation, alkylation or phosphorylation (e.g., a compound obtained by subjecting an amino group in compound (I) to eicosanoylation, alanylation, pentylaminocarbonylation, (5-methyl-2-oxo-1,3-dioxolen-4-yl)methoxycarbonylation, tetrahydrofurylation, pyrrolidylmethylation, 35 pivaloyloxymethylation or t-butylation, and the like);

a compound obtained by subjecting a hydroxy group in compound (I) to acylation, alkylation, phosphorylation or boration (e.g., a compound obtained by subjecting a hydroxy group in compound (I) to acetylation, palmitoylation, propanoylation, 5 pivaloylation, succinylation, fumarylation, alanylation or dimethylaminomethylcarbonylation, and the like);
a compound obtained by subjecting a carboxyl group in compound (I) to esterification or amidation (e.g., a compound obtained by subjecting a carboxyl group in compound (I) to ethyl 10 esterification, phenyl esterification, carboxymethyl esterification, dimethylaminomethyl esterification, pivaloyloxymethyl esterification, ethoxycarbonyloxyethyl esterification, phthalidyl esterification, (5-methyl-2-oxo-1,3-dioxolen-4-yl)methyl esterification, cyclohexyloxycarbonylethyl 15 esterification or methylamidation, and the like) and the like.
Any of these compounds can be produced from compound (I) according to a method known *per se*.

A prodrug of compound (I) may also be one which is converted to compound (I) under physiological conditions as 20 described in "IYAKUJIN no KAIHATSU (Development of Pharmaceuticals)", Vol. 7, Design of Molecules, p. 163-198 (HIROKAWA SHOTEN).

[0279]

Compound (I) may be a crystal, and a single crystal form 25 and a mixture of crystal forms are both encompassed in compound (I) of the present invention. The crystal can be produced by crystallizing according to a crystallization method known *per se*.

[0280]

30 Compound (I) is useful for mammals (e.g., mouse, rat, hamster, rabbit, cat, dog, bovine, sheep, monkey, human etc.) as an agent for the prophylaxis or treatment of diseases, for example,

(1) psychiatric diseases [e.g., depression, major depression, 35 bipolar depression, dysthymic disorder, emotional disorder

(seasonal affective disorder and the like), recurrent depression, postpartum depression, stress disorder, depression symptom, mania, anxiety, generalized anxiety disorder, anxiety syndrome, panic disorder, phobia, social phobia, social anxiety disorder, obsessive disorder, post-traumatic stress syndrome, post-traumatic stress disorder, Tourette syndrome, autism, autism spectrum syndrome, fragile X syndrome, Rett syndrome, adjustment disorder, bipolar disorder, neurosis, schizophrenia (e.g., positive symptom, negative symptom, and cognitive impairment), cognitive impairment associated with schizophrenia, chronic fatigue syndrome, anxiety neurosis, compulsive neurosis, epilepsy, anxiety symptom, anxious mental state, emotional abnormality, cyclothymia, nervous erethism, faint, addiction, low sex drive, attention deficit hyperactivity disorder (ADHD), psychotic major depression, refractory major depression, treatment-resistant depression],

(2) neurodegenerative diseases [e.g., Alzheimer's disease, Alzheimer-type senile dementia, Parkinson's disease, Parkinson's disease dementia, Huntington's disease, multi-infarct dementia, frontotemporal dementia, frontotemporal dementia Parkinson's Type, progressive supranuclear palsy, Pick's syndrome, Niemann-Pick syndrome, corticobasal degeneration, Down's syndrome, vascular dementia, postencephalitic parkinsonism, dementia with Lewy bodies, HIV dementia, amyotrophic lateral sclerosis (ALS), motor neurogenesis disease (MND), Creutzfeldt-Jakob disease or prion disease, cerebral palsy, multiple sclerosis],

(3) age-related cognition and memory disorders [e.g., age-related memory disorders, senile dementia]

(4) sleep disorders [e.g., intrinsic sleep disorders (e.g., psychophysiological insomnia and the like), extrinsic sleep disorder, circadian rhythm disorders (e.g., time zone change syndrome (jet lag), shift work sleep disorder, irregular sleep-wake pattern, delayed sleep phase syndrome, advanced sleep phase syndrome, non-24-hour sleep-wake and the like),

parasomnia, sleep disorders associated with internal medical or psychiatric disorder (e.g., chronic obstructive pulmonary diseases, Alzheimer's disease, Parkinson's disease, cerebrovascular dementia, schizophrenia, depression, anxiety neurosis), stress insomnia, insomnia, insomniac neurosis, sleep apnea syndrome],

(5) respiratory depression caused by anesthetics, traumatic disease, or neurodegenerative disease and the like,

(6) traumatic brain injury, cerebral apoplexy, neurotic anorexia, eating disorder, anorexia nervosa, hyperorexia, other eating disorder, alcohol dependence, alcohol abuse, alcoholic amnesia, alcohol paranoia, alcohol preference, alcohol withdrawal, alcoholic insanity, alcohol poisoning, alcoholic jealousy, alcoholic mania, alcohol-dependent psychiatric disorder, alcoholic insanity, pharmacophilia, pharmacophobia, pharmacomania, drug withdrawal, migraine, stress headache, catatonic headache, diabetic neuropathy, obesity, diabetes, muscular spasm, Meniere's disease, autonomic ataxia, alopecia, glaucoma, hypertension, cardiac disease, tachycardia, congestive cardiac failure, hyperventilation, bronchial asthma, apnea, sudden infant death syndrome, inflammatory disease, allergic disease, impotence, climacteric disorder, infertility, cancer, immunodeficiency syndrome caused by HIV infection, immunodeficiency syndrome caused by stress, cerebrospinal meningitis, acromegaly, incontinence, metabolic syndrome, osteoporosis, peptic ulcer, irritable bowel syndrome, inflammatory bowel disease, ulcerative colitis, Crohn's disease, stress gastrointestinal disorder, neurological vomiting, peptic ulcer, diarrhea, constipation, postoperative ileus, stress gastrointestinal disorder, and

(7) pain.

A cholinergic muscarinic M1 receptor positive allosteric modulator is particularly preferably useful for the prophylaxis or treatment of Alzheimer's disease, schizophrenia, pain, sleep disorder, Parkinson's disease dementia, dementia with Lewy

bodies and the like.

[0281]

Since compound (I) has a high cholinergic muscarinic M1 receptor positive allosteric modulator activity, it is expected to provide an excellent prophylactic or therapeutic effect for the above-mentioned diseases.

[0282]

Compound (I) shows excellent solubility in water, the second solution of Japanese Pharmacopeia Elution Test, or the second solution of Japanese Pharmacopoeia Disintegration Test, shows excellent in vivo kinetics (e.g., plasma drug half-life, intracerebral migration, metabolic stability, CYP inhibition), shows low toxicity (e.g., more excellent as a medicament in terms of acute toxicity, chronic toxicity, genetic toxicity, reproductive toxicity, cardiotoxicity, drug interaction, carcinogenicity, phototoxicity, and the like), and also has excellent properties as a pharmaceutical product such as a few side effects. Therefore, compound (I) can be safely administered orally or parenterally to a mammal (e.g., mouse, rat, hamster, rabbit, cat, dog, bovine, sheep, monkey, human and the like). Examples of the "parenteral" include intravenous, intramuscular, subcutaneous, intra-organ, intranasal, intradermal, instillation, intracerebral, intrarectal, intravaginal, intraperitoneal and intratumor administrations, administration to the vicinity of tumor etc. and direct administration to the lesion.

[0283]

A preparation containing compound (I) may be any of a solid preparation such as powder, granule, tablet, capsule, orally disintegrable film and the like, or a liquid agent such as syrup, emulsion, injection and the like.

[0284]

The medicament of the present invention can be produced by a conventional method such as blending, kneading, granulation, tableting, coating, sterilization treatment,

emulsification and the like according to the form of the preparation. As for the production of the preparation, for example, each item of the Japanese Pharmacopoeia Preparation General Rules and the like can be referred to. In addition,
5 the medicament of the present invention may be formed into a sustained-release preparation containing an active ingredient and a biodegradable polymer compound. The sustained-release preparation can be produced according to the method described in JP-A-H9-263545.

10 [0285]

In the preparation of the present invention, the content of compound (I) varies depending on the form of the preparation, but is generally 0.01 to 100 % by weight, preferably 0.1 to 50 % by weight, more preferably 0.5 to 20 % by weight, as the
15 amount of compound (I) relative to the whole preparation.

[0286]

When compound (I) is used as the above-mentioned pharmaceutical products, it may be used alone or in admixture with a suitable, pharmacologically acceptable carrier, for
20 example, excipients (e.g., starch, lactose, sucrose, calcium carbonate, calcium phosphate, etc.), binders (e.g., starch, arabic gum, carboxymethyl cellulose, hydroxypropyl cellulose, crystalline cellulose, alginic acid, gelatin, polyvinylpyrrolidone, etc.), lubricants (e.g., stearic acid,
25 magnesium stearate, calcium stearate, talc, etc.), disintegrants (e.g., calcium carboxymethylcellulose, talc, etc.), diluents (e.g., water for injection, physiological saline, etc.) and if desired, with the additives (e.g., a stabilizer, a preservative, a colorant, a fragrance, a
30 solubilizing agent, an emulsifier, a buffer, an isotonic agent, etc.) and the like, by a conventional method, which is processed into a dosage form of a solid agent such as powder, fine granule, granule, tablet, capsule and the like or a liquid form such as injection and the like, and safely administered
35 orally or parenterally. When compound (I) is formed as a

preparation for topical administration, it can also be directly administered to the affected part of an articular disease. In this case, an injection is preferable. The compound can also be administered as a parenteral agent for topical

5 administration (e.g., intramuscular injection, subcutaneous injection, organ injection, injection to the vicinity of a joint and the like, solid preparation such as implant, granule, powder and the like, liquid such as suspension and the like, ointment etc.) and the like.

10 [0287]

For formulation into an injection, for example, compound (I) is formulated into an aqueous suspension with a dispersing agent (e.g., surfactant such as Tween 80, HCO-60 and the like, polysaccharides such as carboxymethylcellulose, sodium alginate, 15 hyaluronic acid and the like, polysorbate etc.), preservative (e.g., methylparaben, propylparaben etc.), isotonic agent (e.g., sodium chloride, mannitol, sorbitol, glucose etc.), buffer (e.g., calcium carbonate etc.), pH adjuster (e.g., sodium phosphate, potassium phosphate etc.) and the like to give a 20 practical preparation for injection. In addition, an oily suspension can be obtained by dispersing the compound together with vegetable oil such as sesame oil, corn oil and the like or a mixture thereof with a phospholipid such as lecithin and the like, or medium-chain triglyceride (e.g., miglyol 812 etc.) to 25 give an injection to be actually used.

[0288]

The dose of compound (I) varies depending on the subject of administration, administration route and symptoms and is not particularly limited. For example, for oral administration to 30 adult patients (body weight adult 40 to 80 kg, for example, 60 kg) with Alzheimer's disease, the dose is, for example, 0.001 to 1000 mg/kg body weight/day, preferably 0.01 to 100 mg/kg body weight/day, more preferably 0.1 to 10 mg/kg body weight/day, as compound (I). This amount can be administered 35 in one to three portions per day.

[0289]

A medicament containing the compound of the present invention can be safely administered solely or by mixing with a pharmaceutically acceptable carrier according to a method known *per se* (e.g., the method described in the Japanese Pharmacopoeia etc.) as the production method of a pharmaceutical preparation, and in the form of, for example, tablet (including sugar-coated tablet, film-coated tablet, sublingual tablet, orally disintegrating tablet, buccal and the like), pill, powder, granule, capsule (including soft capsule, microcapsule), troche, syrup, liquid, emulsion, suspension, release control preparation (e.g., immediate-release preparation, sustained-release preparation, sustained-release microcapsule), aerosol, film (e.g., orally disintegrating film, oral mucosa-adhesive film), injection (e.g., subcutaneous injection, intravenous injection, intramuscular injection, intraperitoneal injection), drip infusion, transdermal absorption type preparation, ointment, lotion, adhesive preparation, suppository (e.g., rectal suppository, vaginal suppository), pellet, nasal preparation, pulmonary preparation (inhalant), eye drop and the like, orally or parenterally (e.g., intravenous, intramuscular, subcutaneous, intraorgan, intranasal, intradermal, instillation, intracerebral, intrarectal, intravaginal, intraperitoneal administrations, and administration to the lesion).

[0290]

As the aforementioned "pharmaceutically acceptable carrier", various organic or inorganic carriers conventionally used as preparation materials (starting materials) can be used. For example, excipient, lubricant, binder, disintegrant and the like are used for solid preparations, and solvent, solubilizing agent, suspending agent, isotonic agent, buffer, soothing agent and the like are used for liquid preparations. Where necessary, preparation additives such as preservative, antioxidant, colorant, sweetening agent and the like can also be used.

[0291]

Examples of the excipient include lactose, sucrose, D-mannitol, starch, corn starch, crystalline cellulose, light anhydrous silicic acid and the like.

5 [0292]

Examples of the lubricant include magnesium stearate, calcium stearate, talc, colloidal silica and the like.

[0293]

Examples of the binder include crystalline cellulose,
10 white sugar, D-mannitol, dextrin, hydroxypropylcellulose, hydroxypropylmethylcellulose, polyvinylpyrrolidone, starch, sucrose, gelatin, methylcellulose, carboxymethylcellulose sodium and the like.

[0294]

15 Examples of the disintegrant include starch, carboxymethylcellulose, carboxymethylcellulose calcium, sodium carboxymethyl starch, L-hydroxypropylcellulose and the like.

[0295]

Examples of the solvent include water for injection,
20 alcohol, propylene glycol, macrogol, sesame oil, corn oil, olive oil and the like.

[0296]

Examples of the solubilizing agent include polyethylene glycol, propylene glycol, D-mannitol, benzyl benzoate, ethanol,
25 trisaminomethane, cholesterol, triethanolamine, sodium carbonate, sodium citrate and the like.

[0297]

Examples of the suspending agent include surfactants such as stearyl triethanolamine, sodium lauryl sulfate,
30 laurylaminopropionic acid, lecithin, benzalkonium chloride, benzetonium chloride, glycerin monostearate and the like; hydrophilic polymers such as polyvinyl alcohol, polyvinylpyrrolidone, carboxymethylcellulose sodium, methylcellulose, hydroxymethylcellulose, hydroxyethylcellulose,
35 hydroxypropylcellulose and the like; and the like.

[0298]

Examples of the isotonic agent include glucose, D-sorbitol, sodium chloride, glycerin, D-mannitol and the like.

[0299]

5 Examples of the buffer include buffer solutions such as phosphates, acetates, carbonates, citrates and the like.

[0300]

Examples of the soothing agent include benzyl alcohol and the like.

10 [0301]

Examples of the preservative include p-oxybenzoates, chlorobutanol, benzyl alcohol, phenylethyl alcohol, dehydroacetic acid, sorbic acid and the like.

[0302]

15 Examples of the antioxidant include sulfite, ascorbic acid, α -tocopherol and the like.

[0303]

While the pharmaceutical composition varies according to the dosage form, administration method, carrier and the like, 20 it can be produced according to a conventional method by adding the compound of the present invention in a proportion of generally 0.01 - 100%(w/w), preferably 0.1 - 95%(w/w), of the total amount of the preparation.

[0304]

25 The compound of the present invention can be used in combination with other active ingredients (hereinafter to be abbreviated as concomitant drug).

[0305]

Examples of the concomitant drug include the following.
30 benzodiazepine (chlordiazepoxide, diazepam, potassium clorazepate, lorazepam, clonazepam, alprazolam etc.), L-type calcium channel inhibitor (pregabalin etc.), tricyclic or tetracyclic antidepressant (imipramine hydrochloride, amitriptyline hydrochloride, desipramine hydrochloride, 35 clomipramine hydrochloride etc.), selective serotonin reuptake

inhibitor (fluvoxamine maleate, fluoxetine hydrochloride, citalopram hydrobromide, sertraline hydrochloride, paroxetine hydrochloride, escitalopram oxalate etc.), serotonin-noradrenaline reuptake inhibitor (venlafaxine hydrochloride, duloxetine hydrochloride, desvenlafaxine hydrochloride etc.), noradrenaline reuptake inhibitor (reboxetine mesylate etc.), noradrenaline-dopamine reuptake inhibitor (bupropion hydrochloride etc.), mirtazapine, trazodone hydrochloride, nefazodone hydrochloride, bupropion hydrochloride, setiptiline maleate, 5-HT_{1A} agonist (buspirone hydrochloride, tandospirone citrate, osetozotan hydrochloride etc.), 5-HT₃ antagonist (cyamemazine etc.), heart non-selective β inhibitor (propranolol hydrochloride, oxprenolol hydrochloride etc.), histamine H₁ antagonist (hydroxyzine hydrochloride etc.), therapeutic drug for schizophrenia (chlorpromazine, haloperidol, sulpiride, clozapine, trifluoperazine hydrochloride, fluphenazine hydrochloride, olanzapine, quetiapine fumarate, risperidone, aripiprazole etc.), CRF antagonist, other antianxiety drug (meprobamate etc.), tachykinin antagonist (MK-869, saredutant etc.), medicament that acts on metabotropic glutamate receptor, CCK antagonist, β 3 adrenaline antagonist (amibegron hydrochloride etc.), GAT-1 inhibitor (tiagabine hydrochloride etc.), N-type calcium channel inhibitor, carbonic anhydrase II inhibitor, NMDA glycine moiety agonist, NMDA antagonist (memantine etc.), peripheral benzodiazepine receptor agonist, vasopressin antagonist, vasopressin V1b antagonist, vasopressin V1a antagonist, phosphodiesterase inhibitor, opioid antagonist, opioid agonist, uridine, nicotinic acid receptor agonist, thyroid hormone (T3, T4), TSH, TRH, MAO inhibitor (phenelzine sulfate, tranylcypromine sulfate, moclobemide etc.), 5-HT_{2A} antagonist, 5-HT_{2A} inverse agonist, COMT inhibitor (entacapone etc.), therapeutic drug for bipolar disorder (lithium carbonate, sodium valproate, lamotrigine, riluzole, felbamate etc.), cannabinoid CB1 antagonist (rimonabant etc.), FAAH inhibitor, sodium channel inhibitor, anti-ADHD drug

(methylphenidate hydrochloride, methamphetamine hydrochloride etc.), therapeutic drug for alcoholism, therapeutic drug for autisma, therapeutic drug for chronic fatigue syndrome, therapeutic drug for spasm, therapeutic drug for fibromyalgia syndrome, therapeutic drug for headache, therapeutic drug for insomnia (etizolam, zopiclone, triazolam, zolpidem, ramelteon, indiplon etc.), therapeutic drug for quitting smoking, therapeutic drug for myasthenia gravis, therapeutic drug for cerebral infarction, therapeutic drug for mania, therapeutic drug for hypersomnia, therapeutic drug for pain, therapeutic drug for dysthymia, therapeutic drug for autonomic ataxia, therapeutic drug for male and female sexual dysfunction, therapeutic drug for migraine, therapeutic drug for pathological gambler, therapeutic drug for restless legs syndrome, therapeutic drug for substance addiction, therapeutic drug for alcohol-related syndrome, therapeutic drug for irritable bowel syndrome, therapeutic drug for Alzheimer's disease (donepezil, galanthamine, memantine, rivastigmine etc.), therapeutic drug for Parkinson's disease (levodopa, carbidopa, benserazide, selegiline, rasagiline, zonisamide, entacapone, amantadine, talipexole, pramipexole, ropinirole, rotigotine, apomorphine, cabergoline, pergolide, bromocriptine, istradefylline, trihexyphenidyl, biperiden, piroheptine, profenamine, promethazine, droxidopa, combination of those drugs etc.), therapeutic drug for Parkinson's disease dementia (rivastigmine), therapeutic drug for dementia with Lewy bodies (donepezil), therapeutic drug for ALS (riluzole, neurotrophic factor etc.), therapeutic drug for lipid abnormality such as cholesterol-lowering drug (statin series (pravastatin sodium, atorvastatin, simvastatin, rosuvastatin etc.), fibrate (clofibrate etc.), squalene synthetase inhibitor), therapeutic drug for abnormal behavior or suppressant of dromomania due to dementia (sedatives, antianxiety drug etc.), apoptosis inhibitor, antiobesity drug, therapeutic drug for diabetes, therapeutic drug for hypertension, therapeutic drug for

hypotension, therapeutic drug for rheumatism (DMARD), anti-cancer agent, therapeutic drug for hypothyroidism (PTH), calcium receptor antagonist, sex hormone or a derivative thereof (progesterone, estradiol, estradiol benzoate etc.),
5 neuronal differentiation promoter, nerve regeneration promoter, non-steroidal anti-inflammatory drug (meloxicam, tenoxicam, indomethacin, ibuprofen, celecoxib, rofecoxib, aspirin etc.), steroid (dexamethasone, cortisone acetate etc.), anti-cytokine drug (TNF inhibitor, MAP kinase inhibitor etc.), antibody
10 medicament, nucleic acid or nucleic acid derivative, aptamer drug and the like.

[0306]

By combining the compound of the present invention and a concomitant drug, a superior effect such as
15 (1) the dose can be reduced as compared to single administration of the compound of the present invention or a concomitant drug,
(2) the drug to be combined with the compound of the present invention can be selected according to the condition of
20 patients (mild case, severe case and the like),
(3) the period of treatment can be set longer by selecting a concomitant drug having different action and mechanism from the compound of the present invention,
(4) a sustained treatment effect can be designed by selecting a
25 concomitant drug having different action and mechanism from the compound of the present invention,
(5) a synergistic effect can be afforded by a combined use of the compound of the present invention and a concomitant drug, and the like, can be achieved.

30 [0307]

Hereinafter the compound of the present invention and a concomitant drug used in combination are referred to as the "combination agent of the present invention".

[0308]

35 When using the combination agent of the present invention,

the administration time of the compound of the present invention and the concomitant drug is not restricted, and the compound of the present invention or a pharmaceutical composition thereof and the concomitant drug or a pharmaceutical composition thereof can be administered to an administration subject simultaneously, or may be administered at different times. The dosage of the concomitant drug may be determined according to the dose clinically used, and can be appropriately selected depending on an administration subject, administration route, disease, combination and the like.

[0309]

The administration mode of the combination drug of the present invention is not particularly restricted, and it is sufficient that the compound of the present invention and the concomitant drug are combined in administration. Examples of such administration mode include the following methods:

(1) administration of a single preparation obtained by simultaneously processing the compound of the present invention and the concomitant drug, (2) simultaneous administration of two kinds of preparations of the compound of the present invention and the concomitant drug, which have been separately produced, by the same administration route, (3) administration of two kinds of preparations of the compound of the present invention and the concomitant drug, which have been separately produced, by the same administration route in a staggered manner, (4) simultaneous administration of two kinds of preparations of the compound of the present invention and the concomitant drug, which have been separately produced, by different administration routes, (5) administration of two kinds of preparations of the compound of the present invention and the concomitant drug, which have been separately produced, by different administration routes in a staggered manner (e.g., administration in the order of the compound of the present invention and the concomitant drug, or in the reverse order) and the like.

[0310]

The combination drug of the present invention exhibits low toxicity. For example, the compound of the present invention or (and) the aforementioned concomitant drug can be
5 combined with a pharmacologically acceptable carrier according to the known method to prepare a pharmaceutical composition such as tablets (including sugar-coated tablet and film-coated tablet), powders, granules, capsules (including soft capsule), liquids, injections, suppositories, sustained-release agents,
10 etc. These compositions can be administered safely orally or non-orally (e.g., topical, rectal, intravenous administration etc.). Injection can be administered intravenously, intramuscularly, subcutaneously, or by intraorgan administration or directly to the lesion.

15 [0311]

Examples of the pharmacologically acceptable carriers usable for the production of a combination agent in the present invention, various organic or inorganic carrier substances conventionally used as preparation materials can be mentioned.
20 For solid preparations, for example, excipient, lubricant, binder and disintegrant can be used. For liquid preparations, for example, solvent, solubilizing agent, suspending agent, isotonic agent, buffering agent, soothing agent and the like can be used. Where necessary, conventional preservative,
25 antioxidant, colorant, sweetening agent, adsorbent, wetting agent and the like can be used as appropriate.

[0312]

Examples of the excipient include lactose, sucrose, D-mannitol, starch, corn starch, crystalline cellulose, light
30 anhydrous silicic acid and the like.

[0313]

Examples of the lubricant include magnesium stearate, calcium stearate, talc, colloidal silica and the like.

[0314]

35 Examples of the binder include crystalline cellulose,

white sugar, D-mannitol, dextrin, hydroxypropylcellulose, hydroxypropylmethylcellulose, polyvinylpyrrolidone, starch, sucrose, gelatin, methylcellulose, carboxymethylcellulose sodium and the like.

5 [0315]

Examples of the disintegrant include starch, carboxymethylcellulose, carboxymethylcellulose calcium, sodium carboxymethyl starch, L-hydroxypropylcellulose and the like.

10 [0316]

Examples of the solvent include water for injection, alcohol, propylene glycol, macrogol, sesame oil, corn oil, olive oil and the like.

[0317]

15 Examples of the solubilizing agent include polyethylene glycol, propylene glycol, D-mannitol, benzyl benzoate, ethanol, trisaminomethane, cholesterol, triethanolamine, sodium carbonate, sodium citrate and the like.

[0318]

20 Examples of the suspending agent include surfactants such as stearyl triethanolamine, sodium lauryl sulfate, laurylaminopropionic acid, lecithin, benzalkonium chloride, benzetonium chloride, glycerin monostearate and the like; hydrophilic polymers such as polyvinyl alcohol,
25 polyvinylpyrrolidone, carboxymethylcellulose sodium, methylcellulose, hydroxymethylcellulose, hydroxyethylcellulose, hydroxypropylcellulose and the like; and the like.

[0319]

30 Examples of the isotonic agent include glucose, D-sorbitol, sodium chloride, glycerin, D-mannitol and the like.

[0320]

Examples of the buffer include buffer solutions such as phosphates, acetates, carbonates, citrates and the like.

[0321]

35 Examples of the soothing agent include benzyl alcohol and

the like.

[0322]

Examples of the preservative include p-oxybenzoates, chlorobutanol, benzyl alcohol, phenylethyl alcohol, dehydroacetic acid, sorbic acid and the like.

[0323]

Examples of the antioxidant include sulfite, ascorbic acid, α -tocopherol and the like.

[0324]

The mixing ratio of the compound of the present invention to the concomitant drug in the combination agent of the present invention can be appropriately selected depending on an administration subject, administration route, diseases and the like.

[0325]

For example, the content of the compound of the present invention in the combination agent of the present invention differs depending on the form of a preparation, and usually from about 0.01 to about 100 wt%, preferably from about 0.1 to about 50 wt%, further preferably from about 0.5 to about 20 wt%, based on the preparation.

[0326]

The content of the concomitant drug in the combination agent of the present invention differs depending on the form of a preparation, and usually from about 0.01 to about 100 wt%, preferably from about 0.1 to about 50 wt%, further preferably from about 0.5 to about 20 wt%, based on the preparation.

[0327]

The content of additives such as a carrier and the like in the combination agent of the present invention differs depending on the form of a preparation, and usually from about 1 to about 99.99 wt%, preferably from about 10 to about 90 wt%, based on the preparation.

[0328]

When the compound of the present invention and a

concomitant drug are separately formulated into preparations, the contents thereof are similar to the above.

Examples

[0329]

5 The present invention is explained in detail in the following by referring to Examples, Experimental Examples and Formulation Examples, which are not to be construed as limitative, and the invention may be changed within the scope of the present invention.

10 In the following Examples, the "room temperature" generally means about 10°C to about 35°C. The ratios indicated for mixed solvents are volume mixing ratios, unless otherwise specified. % means wt%, unless otherwise specified.

15 In silica gel column chromatography, NH means use of aminopropylsilane-bound silica gel. In HPLC (high performance liquid chromatography), C18 means use of octadecyl-bound silica gel. The ratios of elution solvents are volume mixing ratios, unless otherwise specified.

20 The "osmium oxide (fixed catalyst I)" in Example means osmium oxide (VIII) (about 7% content) fixed to high solvent resistance polymer, which is commercially available from Wako Pure Chemical Industries, Ltd., unless otherwise specified. In addition, "sodium hydride" means a 60% oil dispersion (mineral mixture).

25 [0330]

 In the following Examples, the following abbreviations are used.

THF: tetrahydrofuran

DMF: N,N-dimethylformamide

30 DMSO: dimethyl sulfoxide

NBS: N-bromosuccinimide

AIBN: 2,2'-azobis(isobutyronitrile)

DME: 1,2-dimethoxyethane

[M+H]⁺: molecular ion peak

35 M: mol concentration

N: normal concentration

HPLC: high-performance liquid chromatography

tRn (n=1-4): retention time in high-performance liquid chromatography (the number means elution order)

5 [0331]

¹H NMR (proton nuclear magnetic resonance spectrum) was measured by Fourier-transform NMR. For the analysis, ACD/SpecManager (trade name) and the like were used. Peaks with very mild protons such as a hydroxy group, an amino group
10 and the like are not sometimes described.

MS (mass spectrum) was measured by LC/MS (liquid chromatography mass spectrometer). As Ionization, ESI (Electro Spray Ionization) method, or APCI (Atmospheric Pressure Chemical Ionization) method was used. The data indicates those
15 actual measured value (found). Generally, molecular ion peaks are observed. In the case of a compound having a tert-butoxycarbonyl group (-Boc), a peak after elimination of a tert-butoxycarbonyl group or tert-butyl group may be observed as a fragment ion. In the case of a compound having a hydroxy
20 group (-OH), a peak after elimination of H₂O may be observed as a fragment ion. In the case of a salt, a molecular ion peak or fragment ion peak of free form is generally observed.

[0332]

Example 1

25 rac-2-(trans-2-hydroxycyclohexyl)-6-((6-(1-methyl-1H-pyrazol-4-yl)pyridin-3-yl)methyl)isoindolin-1-one

A) methyl 5-bromo-2-(bromomethyl)benzoate

To a solution of methyl 5-bromo-2-methylbenzoate (5.27 g) in trifluoromethylbenzene (50.0 mL) were added AIBN (0.04 g)
30 and N-bromosuccinimide (4.50 g), and the mixture was stirred at 90°C for 4 hr under argon atmosphere. The reaction mixture was diluted with ethyl acetate, and the mixture was washed with saturated brine. The organic layer was dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced
35 pressure. The residue was purified by silica gel column

chromatography (ethyl acetate/hexane) to give the title compound (4.89 g).

¹H NMR (300 MHz, CDCl₃) δ 3.95 (3H, s), 4.90 (2H, s), 7.34 (1H, d, J = 8.3 Hz), 7.62 (1H, dd, J = 8.2, 2.2 Hz), 8.11 (1H, d, J = 2.1 Hz).

[0333]

B) rac-6-bromo-2-(trans-2-hydroxycyclohexyl)isoindolin-1-one

To a solution of methyl 5-bromo-2-(bromomethyl)benzoate (0.50 g) in DMF (5.00 mL) were added trans-2-aminocyclohexanol hydrochloride (0.37 g) and N-ethyldiisopropylamine (1.42 mL), and the mixture was stirred at 80°C for 3 hr. To the reaction mixture was added 5% aqueous sodium bicarbonate, and the mixture was extracted with ethyl acetate. The organic layer was washed with saturated brine, and dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (ethyl acetate/hexane) to give the title compound (0.22 g).

MS: [M+H]⁺ 312.0.

[0334]

C) rac-6-((6-chloropyridin-3-yl)methyl)-2-(trans-2-hydroxycyclohexyl)isoindolin-1-one

To a solution of rac-6-bromo-2-(trans-2-hydroxycyclohexyl)isoindolin-1-one (0.23 g) in THF (3.0 mL) were added ((6-chloropyridin-3-yl)methyl)zinc(II) chloride (3.66 mL) and bis(tri-tert-butylphosphine)palladium(0) (0.07 g) under ice-cooling, and the mixture was stirred at room temperature for 3 hr under argon atmosphere. To the reaction mixture were added 5% aqueous sodium bicarbonate and ethyl acetate, the mixture was stirred, and the insoluble substance was removed by filtration. The filtrate was extracted with ethyl acetate, the organic layer was washed with saturated brine, and dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (ethyl

acetate/hexane) to give the title compound (0.15 g).

MS: $[M+H]^+$ 357.2.

[0335]

D) rac-2-(trans-2-hydroxycyclohexyl)-6-((6-(1-methyl-1H-pyrazol-4-yl)pyridin-3-yl)methyl)isoindolin-1-one

To a solution of rac-6-((6-chloropyridin-3-yl)methyl)-2-(trans-2-hydroxycyclohexyl)isoindolin-1-one (0.09 g) in a mixed solvent of THF (9.00 mL)-water (3.00 mL) were added 1-methyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-pyrazole (0.11 g), cesium carbonate (0.33 g) and bis(tri-tert-butylphosphine)palladium(0) (0.03 g), and the mixture was stirred overnight at 85°C under argon atmosphere. To the reaction mixture was added water, and the mixture was extracted with ethyl acetate. The organic layer was washed with saturated brine, and dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (methanol/ethyl acetate) to give the title compound (0.03 g). ^1H NMR (300 MHz, DMSO- d_6) δ 1.27 (3H, brs), 1.53 (1H, brs), 1.66 (3H, brs), 1.94 (1H, brs), 3.54 (1H, brs), 3.78 (1H, d, J = 7.0 Hz), 3.86 (3H, s), 4.05 (2H, s), 4.39 (2H, s), 4.71 (1H, d, J = 5.5 Hz), 7.43-7.68 (5H, m), 7.92 (1H, s), 8.21 (1H, s), 8.45 (1H, s).

[0336]

Example 2

rac-5-chloro-2-(trans-2-hydroxycyclohexyl)-6-(4-(1H-pyrazol-1-yl)benzyl)isoindolin-1-one

A) methyl 4-chloro-2-hydroxy-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate

To a solution of methyl 5-bromo-4-chloro-2-hydroxybenzoate (2.40 g) in toluene (75.0 mL) were added bis(pinacolato)diboron (3.44 g), potassium acetate (2.66 g) and trans-dichlorobis(triphenylphosphine)palladium(II) (0.32 g), and the mixture was stirred at 110°C for 14 hr under argon atmosphere. The reaction mixture was allowed to be cooled to

room temperature, water and ethyl acetate were added thereto, and the mixture was partitioned. The organic layer was washed with water and saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced
5 pressure. The residue was purified by silica gel column chromatography (ethyl acetate/hexane) to give the title compound (1.34 g).

MS: $[M-H]^+$ 311.1.

[0337]

10 B) methyl 5-(4-(1H-pyrazol-1-yl)benzyl)-4-chloro-2-hydroxybenzoate

To a solution of methyl 4-chloro-2-hydroxy-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (0.15 g) in a mixed solvent of toluene (4.00 mL)-ethanol (0.80 mL)-water
15 (0.80 mL) were added 1-(4-(bromomethyl)phenyl)-1H-pyrazole (0.11 g), tetrakis(triphenylphosphine)palladium(0) (0.06 g) and tripotassium phosphate (0.26 g), and the mixture was stirred overnight at 100°C under argon atmosphere. The reaction mixture was allowed to be cooled to room temperature, water and
20 ethyl acetate were added thereto, and the mixture was partitioned. The organic layer was washed with water and saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (ethyl
25 acetate/hexane) to give the title compound (0.06 g).

^1H NMR (300 MHz, CDCl_3) δ 3.90-3.93 (3H, m), 4.06 (2H, s), 6.41-6.48 (1H, m), 7.07 (1H, s), 7.23 (1H, s), 7.26 (1H, s), 7.57-7.66 (3H, m), 7.71 (1H, d, $J=1.5$ Hz), 7.89 (1H, d, $J=3.0$ Hz), 10.69 (1H, s).

30 [0338]

C) methyl 5-(4-(1H-pyrazol-1-yl)benzyl)-4-chloro-2-(((trifluoromethyl)sulfonyl)oxy)benzoate

To a solution of methyl 5-(4-(1H-pyrazol-1-yl)benzyl)-4-chloro-2-hydroxybenzoate (0.18 g) in DMF (3.00 mL) were added
35 sodium hydride (0.03 g) and N-

phenylbis(trifluoromethanesulfonimide) (0.21 g) under ice-cooling, and the mixture was stirred at room temperature for 4 hr. To the reaction mixture was added 1N hydrochloric acid, and the mixture was extracted with ethyl acetate. The organic layer was washed with saturated aqueous sodium bicarbonate solution and saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (ethyl acetate/hexane) to give the title compound (0.18 g).

MS: $[M+H]^+$ 475.0.

[0339]

D) methyl 5-(4-(1H-pyrazol-1-yl)benzyl)-4-chloro-2-vinylbenzoate

To a solution of methyl 5-(4-(1H-pyrazol-1-yl)benzyl)-4-chloro-2-(((trifluoromethyl)sulfonyl)oxy)benzoate (0.18 g) in DMF (3.50 mL) were added tributylvinyltin (0.17 mL), trans-dichlorobis(triphenylphosphine)palladium(II) (0.01 g) and lithium chloride (0.12 g), and the mixture was stirred at 90°C for 1 hr under argon atmosphere. To the reaction mixture was added aqueous potassium fluoride solution, and the precipitated insoluble substance was removed by filtration through Celite. The filtrate was diluted with ethyl acetate, and the mixture was washed with water and saturated brine. The organic layer was dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (ethyl acetate/hexane) to give the title compound (0.13 g).

MS: $[M+H]^+$ 353.1.

[0340]

E) methyl 5-(4-(1H-pyrazol-1-yl)benzyl)-4-chloro-2-formylbenzoate

To a solution of methyl 5-(4-(1H-pyrazol-1-yl)benzyl)-4-chloro-2-vinylbenzoate (0.12 g) in a mixed solvent of acetone (2.00 mL)-acetonitrile (2.00 mL)-water (2.00 mL) were added

osmium oxide (fixed catalyst I) (0.04 g) and sodium periodate (0.36 g), and the mixture was stirred overnight at room temperature. The reaction mixture was filtered, and the filtrate was extracted with ethyl acetate. The organic layer
5 was washed with water and saturated brine, and dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure to give the title compound as a crude product. This compound was used in the next step without an additional purification.

10 [0341]

F) rac-5-chloro-2-(trans-2-hydroxycyclohexyl)-6-(4-(1H-pyrazol-1-yl)benzyl)isoindolin-1-one

A solution of methyl 5-(4-(1H-pyrazol-1-yl)benzyl)-4-chloro-2-formylbenzoate (0.12 g), trans-2-aminocyclohexanol
15 hydrochloride (0.05 g), triethylamine (0.05 mL) and anhydrous magnesium sulfate (0.08 g) in THF (2.50 mL) was stirred at room temperature for 1 hr. The insoluble substance was removed by filtration, and the filtrate was concentrated under reduced pressure. The residue was diluted with methanol (2.50 mL) and
20 THF (2.50 mL), sodium triacetoxyborohydride (0.15 g) was added thereto, and the mixture was stirred at room temperature for 3 hr. The reaction mixture was diluted with ethyl acetate, and the mixture was washed with water and saturated brine. The organic layer was dried over anhydrous magnesium sulfate, and
25 the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (ethyl acetate/hexane) to give the title compound (0.05 mg).

¹H NMR (300 MHz, DMSO-d₆) δ 1.23-1.36 (3H, m), 1.48-1.72 (4H, m), 1.89-2.00 (1H, m), 3.49-3.62 (1H, m), 3.74-3.85 (1H, m),
30 4.20 (2H, s), 4.42 (2H, s), 4.76 (1H, d, J = 5.3 Hz), 6.48-6.54 (1H, m), 7.32 (2H, d, J = 8.7 Hz), 7.66 (1H, s), 7.69-7.78 (4H, m), 8.43 (1H, d, J = 2.6 Hz).

[0342]

Example 3

35 rac-2-(trans-2-hydroxycyclohexyl)-5-methoxy-6-(4-(1H-pyrazol-1-

yl)benzyl)isoindolin-1-one

A) methyl 5-bromo-2-hydroxy-4-methoxybenzoate

To a solution of 5-bromo-2-hydroxy-4-methoxybenzoic acid (2.00 g) in methanol (10.0 mL) was added 0.6M (diazomethyl)trimethylsilane/hexane solution (14.8 mL) under ice-cooling, and the mixture was stirred for 3 hr. To the reaction mixture was added acetic acid (0.12 mL), and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (ethyl acetate/hexane) to give the title compound (2.08 g).
¹H NMR (300 MHz, CDCl₃) δ 3.91 (3H, s), 3.93 (H, s), 6.49 (1H, s), 7.99 (1H, s), 10.93 (1H, s).

[0343]

B) methyl 2-hydroxy-4-methoxy-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate

To a solution of methyl 5-bromo-2-hydroxy-4-methoxybenzoate (2.08 g) in toluene (70.0 mL) were added bis(pinacolato)diboron (3.03 g), potassium acetate (2.35 g) and trans-dichlorobis(triphenylphosphine)palladium(II) (0.28 g), and the mixture was stirred at 110°C for 14 hr under argon atmosphere. The reaction mixture was allowed to be cooled to room temperature, water and ethyl acetate were added thereto, and the mixture was partitioned. The organic layer was washed with water and saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (ethyl acetate/hexane) to give the title compound (1.93 g).

MS: [M-H]⁺ 309.2.

[0344]

C) methyl 5-(4-(1H-pyrazol-1-yl)benzyl)-2-hydroxy-4-methoxybenzoate

To a solution of methyl 2-hydroxy-4-methoxy-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (0.80 g) in a mixed solvent of DME (12.0 mL)-water (4.00 mL) were added 1-(4-

(bromomethyl)phenyl)-1H-pyrazole (0.62 g),
tetrakis(triphenylphosphine)palladium(0) (0.15 g) and sodium
carbonate (0.55 g), and the mixture was stirred overnight at
80°C under argon atmosphere. The reaction mixture was allowed
5 to be cooled to room temperature, water and ethyl acetate were
added thereto, and the mixture was partitioned. The organic
layer was washed with water and saturated brine, and dried over
anhydrous magnesium sulfate, and the solvent was evaporated
under reduced pressure. The residue was purified by silica gel
10 column chromatography (ethyl acetate/hexane) to give the title
compound (0.48 g).

MS: $[M-H]^+$ 339.1.

[0345]

D) methyl 5-(4-(1H-pyrazol-1-yl)benzyl)-4-methoxy-2-
15 (((trifluoromethyl)sulfonyl)oxy)benzoate

To a solution of methyl 5-(4-(1H-pyrazol-1-yl)benzyl)-2-
hydroxy-4-methoxybenzoate (0.18 g) in DMF (4.00 mL) were added
sodium hydride (0.03 g) and N-
phenylbis(trifluoromethanesulfonimide) (0.21 g) under ice-
20 cooling, and the mixture was stirred at room temperature for 2
hr. To the reaction mixture was added 1N hydrochloric acid,
and the mixture was extracted with ethyl acetate. The organic
layer was washed with saturated aqueous sodium bicarbonate
solution and saturated brine, and dried over anhydrous
25 magnesium sulfate, and the solvent was evaporated under reduced
pressure. The residue was purified by silica gel column
chromatography (ethyl acetate/hexane) to give the title
compound (0.24 g).

MS: $[M+H]^+$ 471.1.

30 [0346]

E) methyl 5-(4-(1H-pyrazol-1-yl)benzyl)-4-methoxy-2-
vinylbenzoate

To a solution of methyl 5-(4-(1H-pyrazol-1-yl)benzyl)-4-
methoxy-2-(((trifluoromethyl)sulfonyl)oxy)benzoate (0.23 g) in
35 DMF (4.50 mL) were added tributylvinyltin (0.22 mL), trans-

dichlorobis(triphenylphosphine)palladium(II) (0.02 g) and lithium chloride (0.16 g), and the mixture was stirred at 90°C for 1 hr under argon atmosphere. To the reaction mixture was added aqueous potassium fluoride solution, and the precipitated
5 insoluble substance was removed by filtration through Celite. The filtrate was diluted with ethyl acetate, and the mixture was washed with water and saturated brine. The organic layer was dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by
10 silica gel column chromatography (ethyl acetate/hexane) to give the title compound (0.16 g).

MS: $[M+H]^+$ 349.1.

[0347]

F) methyl 5-(4-(1H-pyrazol-1-yl)benzyl)-2-formyl-4-
15 methoxybenzoate

To a solution of methyl 5-(4-(1H-pyrazol-1-yl)benzyl)-4-methoxy-2-vinylbenzoate (0.16 g) in a mixed solvent of acetone (2.20 mL)-acetonitrile (2.20 mL)-water (2.20 mL) were added osmium oxide (fixed catalyst I) (0.06 g) and sodium periodate
20 (0.48 g), and the mixture was stirred overnight at room temperature. The reaction mixture was filtered, and the filtrate was extracted with ethyl acetate. The organic layer was washed with water and saturated brine, and dried over anhydrous sodium sulfate, and the solvent was evaporated under
25 reduced pressure to give the title compound (0.16 g) as a crude product. This compound was used in the next step without an additional purification.

MS: $[M+H]^+$ 351.1.

[0348]

30 G) rac-2-(trans-2-hydroxycyclohexyl)-5-methoxy-6-(4-(1H-pyrazol-1-yl)benzyl)isoindolin-1-one

A solution of methyl 5-(4-(1H-pyrazol-1-yl)benzyl)-2-formyl-4-methoxybenzoate (0.16 g), trans-2-aminocyclohexanol hydrochloride (0.07 g), triethylamine (0.06 mL) and anhydrous
35 magnesium sulfate (0.10 g) in THF (3.50 mL) was stirred at room

temperature for 2 hr. The insoluble substance was removed by filtration, and the filtrate was concentrated under reduced pressure. The residue was diluted with methanol (3.50 mL) and THF (3.50 mL), sodium triacetoxyborohydride (0.19 g) was added
5 thereto, and the mixture was stirred at room temperature for 3.5 hr. The reaction mixture was diluted with ethyl acetate, and the mixture was washed with water and saturated brine. The organic layer was dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue
10 was purified by silica gel column chromatography (ethyl acetate/hexane) to give the title compound (0.04 g).

^1H NMR (300 MHz, DMSO- d_6) δ 1.16-1.36 (3H, m), 1.42-1.74 (4H, m), 1.88-2.00 (1H, m), 3.49-3.63 (1H, m), 3.70-3.83 (1H, m), 3.87 (3H, s), 3.99 (2H, s), 4.37 (2H, s), 4.70 (1H, d, $J = 5.7$
15 Hz), 6.48-6.53 (1H, m), 7.20 (1H, s), 7.31 (2H, d, $J = 8.7$ Hz), 7.41 (1H, s), 7.67-7.77 (3H, m), 8.42 (1H, d, $J = 2.6$ Hz).

[0349]

Example 4

2-((1S,2S)-2-hydroxycyclopentyl)-5-methyl-6-(4-(1H-pyrazol-1-yl)benzyl)isoindolin-1-one
20

A) 5-bromo-2-hydroxy-4-methylbenzoic acid

To a solution of 2-hydroxy-4-methylbenzoic acid (5.00 g) in acetic acid (70.0 mL) was added dropwise bromine (1.68 mL), and the mixture was stirred at room temperature for 5.5 hr. To
25 the reaction mixture was added water, and the precipitate was collected by filtration, and dried under reduced pressure to give the title compound (6.78 g).

MS: $[\text{M}-\text{H}]^+$ 229.0.

[0350]

30 B) methyl 5-bromo-2-hydroxy-4-methylbenzoate

To a solution of 5-bromo-2-hydroxy-4-methylbenzoic acid (6.78 g) in methanol (200 mL) was added dropwise sulfuric acid (6.77 mL), and the mixture was stirred overnight at 70°C. The reaction mixture was concentrated under reduced pressure, the
35 residue was neutralized with saturated aqueous sodium

bicarbonate solution under ice-cooling, and the mixture was extracted with ethyl acetate. The organic layer was washed with water and saturated brine, and dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure.

5 The residue was purified by silica gel column chromatography (ethyl acetate/hexane) to give the title compound (6.56 g).

MS: $[M-H]^+$ 243.0.

[0351]

C) methyl 2-hydroxy-4-methyl-5-(4,4,5,5-tetramethyl-1,3,2-
10 dioxaborolan-2-yl)benzoate

To a solution of methyl 5-bromo-2-hydroxy-4-methylbenzoate (3.20 g) in toluene (95.0 mL) were added bis(pinacolato)diboron (4.97 g), potassium acetate (3.84 g) and trans-dichlorobis(triphenylphosphine)palladium(II) (0.46 g),
15 and the mixture was stirred at 110°C for 15 hr under argon atmosphere. The reaction mixture was allowed to be cooled to room temperature, water and ethyl acetate were added thereto, and the mixture was partitioned. The organic layer was washed with water and saturated brine, and dried over anhydrous
20 magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (ethyl acetate/hexane) to give the title compound (3.84 g).

MS: $[M+H]^+$ 293.1.

25 [0352]

D) methyl 5-(4-(1H-pyrazol-1-yl)benzyl)-2-hydroxy-4-methylbenzoate

To a solution of methyl 2-hydroxy-4-methyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (1.40 g) in a
30 mixed solvent of DME (21.0 mL)-water (7.00 mL) were added 1-(4-(bromomethyl)phenyl)-1H-pyrazole (1.14 g), tetrakis(triphenylphosphine)palladium(0) (0.28 g) and sodium carbonate (1.02 g), and the mixture was stirred overnight at
80°C under argon atmosphere. The reaction mixture was allowed
35 to be cooled to room temperature, water and ethyl acetate were

added thereto, and the mixture was partitioned. The organic layer was washed with water and saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (ethyl acetate/hexane) to give the title compound (0.85 g).

MS: $[M+H]^+$ 323.1.

[0353]

E) methyl 5-(4-(1H-pyrazol-1-yl)benzyl)-4-methyl-2-
10 (((trifluoromethyl)sulfonyl)oxy)benzoate

To a solution of methyl 5-(4-(1H-pyrazol-1-yl)benzyl)-2-hydroxy-4-methylbenzoate (0.85 g) in DMF (17.0 mL) were added sodium hydride (0.13 g) and N-phenylbis(trifluoromethanesulfonimide) (1.03 g) under ice-cooling, and the mixture was stirred at room temperature for 3 hr. To the reaction mixture was added 1N hydrochloric acid, and the mixture was extracted with ethyl acetate. The organic layer was washed with saturated aqueous sodium bicarbonate solution and saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (ethyl acetate/hexane) to give the title compound (1.13 g).

MS: $[M+H]^+$ 455.1.

25 [0354]

F) methyl 5-(4-(1H-pyrazol-1-yl)benzyl)-4-methyl-2-vinylbenzoate

To a solution of methyl 5-(4-(1H-pyrazol-1-yl)benzyl)-4-methyl-2-(((trifluoromethyl)sulfonyl)oxy)benzoate (1.12 g) in DMF (25.0 mL) were added tributylvinyltin (1.08 mL), trans-dichlorobis(triphenylphosphine)palladium(II) (0.09 g) and lithium chloride (0.77 g), and the mixture was stirred at 90°C 1.5 hr under argon atmosphere. To the reaction mixture was added aqueous potassium fluoride solution, and the precipitated insoluble substance was removed by filtration through Celite.

The filtrate was diluted with ethyl acetate, and the mixture was washed with water and saturated brine. The organic layer was dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (ethyl acetate/hexane) to give the title compound (0.73 g).

MS: $[M+H]^+$ 333.1.

[0355]

G) methyl 5-(4-(1H-pyrazol-1-yl)benzyl)-2-formyl-4-methylbenzoate

To a solution of methyl 5-(4-(1H-pyrazol-1-yl)benzyl)-4-methyl-2-vinylbenzoate (0.31 g) in a mixed solvent of acetone (6.20 mL)-acetonitrile (6.20 mL)-water (6.20 mL) were added osmium oxide (fixed catalyst I) (0.12 g) and sodium periodate (1.00 g), and the mixture was stirred overnight at room temperature. The reaction mixture was filtered, and the filtrate was extracted with ethyl acetate. The organic layer was washed with water and saturated brine, and dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure to give the title compound (0.16 g) as a crude product. This compound was used in the next step without an additional purification.

MS: $[M+H]^+$ 335.1.

[0356]

H) 2-((1S,2S)-2-hydroxycyclopentyl)-5-methyl-6-(4-(1H-pyrazol-1-yl)benzyl)isoindolin-1-one

A solution of methyl 5-(4-(1H-pyrazol-1-yl)benzyl)-2-formyl-4-methylbenzoate (0.16 g), (1S,2S)-2-aminocyclopentanol hydrochloride (0.06 g), triethylamine (0.07 mL) and anhydrous magnesium sulfate (0.11 g) in THF (3.10 mL) was stirred at room temperature for 3 hr. The insoluble substance was removed by filtration, and the filtrate was concentrated under reduced pressure. The residue was diluted with methanol (3.10 mL) and THF (3.10 mL), sodium triacetoxyborohydride (0.20 g) was added thereto, and the mixture was stirred overnight at room

temperature. The reaction mixture was diluted with ethyl acetate, and the mixture was washed with water and saturated brine. The organic layer was dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure.

5 The residue was purified by silica gel column chromatography (ethyl acetate/hexane) to give the title compound (71 mg).

^1H NMR (300 MHz, DMSO- d_6) δ 1.45-1.97 (6H, m), 2.32 (3H, s), 4.00-4.28 (4H, m), 4.40 (2H, s), 4.90 (1H, d, J = 4.9 Hz), 6.49-6.56 (1H, m), 7.25 (2H, d, J = 8.5 Hz), 7.41 (2H, d, J =
10 12.8 Hz), 7.67-7.81 (3H, m), 8.44 (1H, d, J = 2.5 Hz).

[0357]

Example 5

rac-6-((6-(1,3-dimethyl-1H-pyrazol-4-yl)pyridin-3-yl)methyl)-2-(trans-2-hydroxycyclohexyl)-5-methylisoindolin-1-one

15 A) methyl 5-((6-chloropyridin-3-yl)methyl)-2-hydroxy-4-methylbenzoate

To a solution of methyl 5-bromo-2-hydroxy-4-methylbenzoate (0.61 g) in THF (20.0 mL) were added 0.5M (2-chloro-5-pyridyl)methylzinc chloride/THF solution (12.5 mL) and
20 bis(tri-tert-butylphosphine)palladium(0) (0.13 g), and the mixture was stirred overnight at room temperature under argon atmosphere. The reaction mixture was concentrated under reduced pressure, and the residue was purified by silica gel column chromatography (ethyl acetate/hexane) to give the title
25 compound (0.52 g).

MS: $[\text{M}+\text{H}]^+$ 292.0.

[0358]

B) methyl 5-((6-(1,3-dimethyl-1H-pyrazol-4-yl)pyridin-3-yl)methyl)-2-hydroxy-4-methylbenzoate

30 To a solution of methyl 5-((6-chloropyridin-3-yl)methyl)-2-hydroxy-4-methylbenzoate (0.09 g) in a mixed solvent of THF (2.40 mL)-water (0.80 mL) were added 1,3-dimethyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-pyrazole (0.14 g), potassium carbonate (0.17 g) and
35 tetrakis(triphenylphosphine)palladium(0) (0.03 g), and the

mixture was stirred overnight at 85°C under argon atmosphere. To the reaction mixture was added water, and the mixture was extracted with ethyl acetate. The organic layer was washed with water and saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (ethyl acetate/hexane) to give the title compound (0.09 g).

MS: $[M+H]^+$ 352.1.

10 [0359]

C) methyl 5-((6-(1,3-dimethyl-1H-pyrazol-4-yl)pyridin-3-yl)methyl)-4-methyl-2-(((trifluoromethyl)sulfonyl)oxy)benzoate

To a solution of methyl 5-((6-(1,3-dimethyl-1H-pyrazol-4-yl)pyridin-3-yl)methyl)-2-hydroxy-4-methylbenzoate (0.27 g) in DMF (6.00 mL) were added sodium hydride (0.04 g) and N-phenylbis(trifluoromethanesulfonimide) (0.30 g) under ice-cooling, and the mixture was stirred at room temperature for 2 hr. To the reaction mixture was added 1N hydrochloric acid, and the mixture was extracted with ethyl acetate. The organic layer was washed with saturated aqueous sodium bicarbonate solution and saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (ethyl acetate/hexane) to give the title compound (0.26 g).

MS: $[M+H]^+$ 484.1.

[0360]

D) methyl 5-((6-(1,3-dimethyl-1H-pyrazol-4-yl)pyridin-3-yl)methyl)-4-methyl-2-vinylbenzoate

To a solution of methyl 5-((6-(1,3-dimethyl-1H-pyrazol-4-yl)pyridin-3-yl)methyl)-4-methyl-2-(((trifluoromethyl)sulfonyl)oxy)benzoate (0.26 g) in DMF (5.5 mL) were added tributylvinyltin (0.24 mL), trans-dichlorobis(triphenylphosphine)palladium(II) (0.02 g) and lithium chloride (0.17 g), and the mixture was stirred at 90°C

for 1.5 hr under argon atmosphere. To the reaction mixture was added aqueous potassium fluoride solution, and the precipitated insoluble substance was removed by filtration through Celite. The filtrate was diluted with ethyl acetate, and the mixture
5 was washed with water and saturated brine. The organic layer was dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (ethyl acetate/hexane) to give the title compound (0.18 g).

10 MS: $[M+H]^+$ 362.1.

[0361]

E) methyl 5-((6-(1,3-dimethyl-1H-pyrazol-4-yl)pyridin-3-yl)methyl)-2-formyl-4-methylbenzoate

To a solution of methyl 5-((6-(1,3-dimethyl-1H-pyrazol-4-yl)pyridin-3-yl)methyl)-4-methyl-2-vinylbenzoate (0.18 g) in a
15 mixed solvent of acetone (2.80 mL)-acetonitrile (2.80 mL)-water (2.80 mL) were added osmium oxide (fixed catalyst I) (0.06 g) and sodium periodate (0.53 g), and the mixture was stirred overnight at room temperature. The reaction mixture was
20 filtered, and the filtrate was extracted with ethyl acetate. The organic layer was washed with water and saturated brine, and dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure to give the title compound (0.18 g) as a crude product. This compound was used in the
25 next step without an additional purification.

MS: $[M+H]^+$ 364.2.

[0362]

F) rac-6-((6-(1,3-dimethyl-1H-pyrazol-4-yl)pyridin-3-yl)methyl)-2-(trans-2-hydroxycyclohexyl)-5-methylisoindolin-1-
30 one

A solution of methyl 5-((6-(1,3-dimethyl-1H-pyrazol-4-yl)pyridin-3-yl)methyl)-2-formyl-4-methylbenzoate (0.09 g),
trans-2-aminocyclohexanol hydrochloride (0.04 g), triethylamine (0.04 mL) and anhydrous magnesium sulfate (0.06 g) in THF (2.00
35 mL) was stirred at room temperature for 1.5 hr. The insoluble

substance was removed by filtration, and the filtrate was concentrated under reduced pressure. The residue was diluted with methanol (2.00 mL) and THF (2.00 mL), sodium triacetoxyborohydride (0.11 g) was added thereto, and the mixture was stirred overnight at room temperature. The reaction mixture was diluted with ethyl acetate, and the mixture was washed with water and saturated brine. The organic layer was dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (ethyl acetate/hexane) to give the title compound (0.05 g).

¹H NMR (300 MHz, CDCl₃) δ 1.27-1.59 (5H, m), 1.81 (2H, d, J = 11.7 Hz), 1.92 (1H, d, J = 13.2 Hz), 2.13-2.24 (1H, m), 2.35 (3H, s), 2.49 (3H, s), 3.61-3.73 (1H, m), 3.87 (3H, s), 4.03 (2H, s), 4.06-4.15 (1H, m), 4.29-4.47 (2H, m), 7.27-7.39 (3H, m), 7.64 (1H, s), 7.76 (1H, s), 8.42 (1H, d, J = 1.5 Hz).

[0363]

Example 6

rac-2-(trans-2-hydroxycyclohexyl)-5-methyl-6-((2'-methyl-2,4'-bipyridin-5-yl)methyl)isoindolin-1-one

A) methyl 2-hydroxy-4-methyl-5-((2'-methyl-[2,4'-bipyridine]-5-yl)methyl)benzoate

To a solution of methyl 5-((6-chloropyridin-3-yl)methyl)-2-hydroxy-4-methylbenzoate (0.35 g) in a mixed solvent of THF (8.40 mL)-water (2.80 mL) were added 2-methyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)pyridine (0.53 g), potassium carbonate (0.66 g) and tetrakis(triphenylphosphine)palladium(0) (0.14 g), and the mixture was stirred overnight at 85°C under argon atmosphere. To the reaction mixture was added water, and the mixture was extracted with ethyl acetate. The organic layer was washed with water and saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (ethyl acetate/hexane) to give the title

compound (0.35 g).

MS: $[M+H]^+$ 349.1.

[0364]

B) methyl 4-methyl-5-((2'-methyl-[2,4'-bipyridine]-5-yl)methyl)-2-(((trifluoromethyl)sulfonyl)oxy)benzoate

To a solution of methyl 2-hydroxy-4-methyl-5-((2'-methyl-[2,4'-bipyridine]-5-yl)methyl)benzoate (0.34 g) in DMF (7.00 mL) were added sodium hydride (0.05 g) and N-phenylbis(trifluoromethanesulfonimide) (0.39 g) under ice-cooling, and the mixture was stirred at room temperature for 2.5 hr. To the reaction mixture was added 1N hydrochloric acid, and the mixture was extracted with ethyl acetate. The organic layer was washed with saturated aqueous sodium bicarbonate solution and saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (ethyl acetate/hexane) to give the title compound (0.22 g).

MS: $[M+H]^+$ 481.1.

[0365]

C) methyl 4-methyl-5-((2'-methyl-[2,4'-bipyridine]-5-yl)methyl)-2-vinylbenzoate

To a solution of methyl 4-methyl-5-((2'-methyl-[2,4'-bipyridine]-5-yl)methyl)-2-(((trifluoromethyl)sulfonyl)oxy)benzoate (0.21 g) in DMF (4.00 mL) were added tributylvinyltin (0.19 mL), trans-dichlorobis(triphenylphosphine)palladium(II) (0.02 g) and lithium chloride (0.14 g), and the mixture was stirred at 90°C for 1.5 hr under argon atmosphere. To the reaction mixture was added aqueous potassium fluoride solution, and the precipitated insoluble substance was removed by filtration through Celite. The filtrate was diluted with ethyl acetate, and the mixture was washed with water and saturated brine. The organic layer was dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by

silica gel column chromatography (ethyl acetate/hexane) to give the title compound (0.11 g).

MS: $[M+H]^+$ 359.1.

[0366]

- 5 D) methyl 2-formyl-4-methyl-5-((2'-methyl-[2,4'-bipyridine]-5-yl)methyl)benzoate

To a solution of methyl 4-methyl-5-((2'-methyl-[2,4'-bipyridine]-5-yl)methyl)-2-vinylbenzoate (0.11 g) in a mixed solvent of acetone (2.00 mL)-acetonitrile (2.00 mL)-water (2.00
10 mL) were added osmium oxide (fixed catalyst I) (0.04 g) and sodium periodate (0.33 g), and the mixture was stirred overnight at room temperature. The reaction mixture was filtered, and the filtrate was extracted with ethyl acetate. The organic layer was washed with water and saturated brine,
15 and dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure to give the title compound (0.11 g) as a crude product. This compound was used in the next step without an additional purification.

MS: $[M+H]^+$ 361.2.

20 [0367]

- E) rac-2-(trans-2-hydroxycyclohexyl)-5-methyl-6-((2'-methyl-2,4'-bipyridin-5-yl)methyl)isoindolin-1-one

A solution of methyl 2-formyl-4-methyl-5-((2'-methyl-[2,4'-bipyridine]-5-yl)methyl)benzoate (0.06 g), trans-2-
25 aminocyclohexanol hydrochloride (0.02 g), triethylamine (0.02 mL) and anhydrous magnesium sulfate (0.04 g) in THF (1.20 mL) was stirred at room temperature for 1.5 hr. The insoluble substance was removed by filtration, and the filtrate was concentrated under reduced pressure. The residue was diluted
30 with methanol (1.20 mL) and THF (1.20 mL), sodium triacetoxyborohydride (0.06 g) was added thereto, and the mixture was stirred overnight at room temperature. The reaction mixture was diluted with ethyl acetate, and the mixture was washed with water and saturated brine. The organic
35 layer was dried over anhydrous magnesium sulfate, and the

solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (ethyl acetate/hexane) to give the title compound (0.02 g).

¹H NMR (300 MHz, DMSO-d₆) δ 1.23-1.36 (3H, m), 1.45-1.72 (4H, m), 1.90-2.01 (1H, m), 2.35 (3H, s), 2.54 (3H, s), 3.58 (1H, dt, J = 11.8, 5.6 Hz), 3.74-3.85 (1H, m), 4.16 (2H, s), 4.38 (2H, s), 4.73 (1H, d, J = 5.7 Hz), 7.41 (1H, s), 7.47 (1H, s), 7.66 (1H, dd, J = 7.9, 2.3 Hz), 7.81 (1H, d, J = 5.3 Hz), 7.90 (1H, s), 8.01 (1H, d, J = 8.3 Hz), 8.53 (1H, d, J = 5.3 Hz), 8.61 (1H, d, J = 1.5 Hz).

[0368]

Example 7

3-fluoro-2-(5-methyl-1-oxo-6-(4-(1H-pyrazol-1-yl)benzyl)-1,3-dihydro-2H-isoindol-2-yl)benzonitrile

A) 6-(4-(1H-pyrazol-1-yl)benzyl)-2-(2,4-dimethoxybenzyl)-5-methylisoindolin-1-one

A solution of methyl 5-(4-(1H-pyrazol-1-yl)benzyl)-2-formyl-4-methylbenzoate (0.30 g), (2,4-dimethoxyphenyl)methanamine (0.02 g) and anhydrous magnesium sulfate (0.21 g) in THF (6.00 mL) was stirred at room temperature for 1.5 hr. The insoluble substance was removed by filtration, and the filtrate was concentrated under reduced pressure. The residue was diluted with methanol (6.00 mL) and THF (6.00 mL), sodium triacetoxyborohydride (0.38 g) was added thereto, and the mixture was stirred overnight at room temperature. The reaction mixture was diluted with ethyl acetate, and the mixture was washed with water and saturated brine. The organic layer was dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (ethyl acetate/hexane) to give the title compound (0.25 g).

MS: [M+H]⁺ 454.2.

[0369]

B) 6-(4-(1H-pyrazol-1-yl)benzyl)-5-methylisoindolin-1-one

To a solution of 6-(4-(1H-pyrazol-1-yl)benzyl)-2-(2,4-

dimethoxybenzyl)-5-methylisoindolin-1-one (0.24 g) in trifluoroacetic acid (4.34 mL) was added anisole (0.23 mL), and the mixture was stirred at 80°C for 2 hr. The reaction mixture was concentrated under reduced pressure, and the residue was
5 purified by silica gel column chromatography (ethyl acetate/hexane) to give the title compound (0.15 g).

MS: $[M+H]^+$ 304.1.

[0370]

C) 3-fluoro-2-(5-methyl-1-oxo-6-(4-(1H-pyrazol-1-yl)benzyl)-
10 1,3-dihydro-2H-isoindol-2-yl)benzonitrile

To a solution of 6-(4-(1H-pyrazol-1-yl)benzyl)-5-methylisoindolin-1-one (0.05 g) in DMF (1.00 mL) were added potassium carbonate (0.07 g) and 2,3-difluorobenzonitrile (0.05 g), and the mixture was stirred overnight at 150°C under argon
15 atmosphere. The reaction mixture was diluted with water and ethyl acetate, and the mixture was partitioned. The organic layer was washed with water and saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel
20 column chromatography (ethyl acetate/hexane) and then HPLC (water/methanol) to give the title compound (7.80 mg).

^1H NMR (300 MHz, DMSO- d_6) δ 2.39 (3H, s), 4.16 (2H, s), 4.90 (2H, s), 6.50-6.55 (1H, m), 7.31 (2H, d, $J = 8.5$ Hz), 7.54 (1H, s), 7.62 (1H, s), 7.65-7.73 (2H, m), 7.75-7.91 (4H, m), 8.45 (1H,
25 d, $J = 2.4$ Hz).

[0371]

Example 8

5-methyl-6-(4-(1H-pyrazol-1-yl)benzyl)-2-(tetrahydro-2H-pyran-4-yl)isoindolin-1-one

30 A solution of methyl 5-(4-(1H-pyrazol-1-yl)benzyl)-2-formyl-4-methylbenzoate (0.10 g), tetrahydro-2H-pyran-4-amine (0.03 g) and anhydrous magnesium sulfate (0.07 g) in THF (2.00 mL) was stirred at room temperature for 1 hr. The insoluble substance was removed by filtration, and the filtrate was
35 concentrated under reduced pressure. The residue was diluted

with methanol (2.00 mL) and THF (2.00 mL), sodium triacetoxyborohydride (0.13 g) was added thereto, and the mixture was stirred overnight at room temperature. The reaction mixture was diluted with ethyl acetate, and the mixture was washed with water and saturated brine. The organic layer was dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (ethyl acetate/hexane) to give the title compound (0.04 g).

¹H NMR (300 MHz, DMSO-d₆) δ 1.59-1.70 (2H, m), 1.80 (2H, qd, J = 12.1, 4.3 Hz), 2.32 (3H, s), 3.3-3.49 (2H, m), 3.94 (2H, dd, J = 11.1, 4.0 Hz), 4.09 (2H, s), 4.18-4.29 (1H, m), 4.41 (2H, s), 6.49-6.55 (1H, m), 7.25 (2H, d, J = 8.3 Hz), 7.42 (2H, d, J = 13.9 Hz), 7.70-7.78 (3H, m), 8.44 (1H, d, J = 2.6 Hz).

[0372]

Example 9

rac-5-cyclopropyl-2-(trans-2-hydroxycyclopentyl)-6-(4-(1H-pyrazol-1-yl)benzyl)isoindolin-1-one

A) methyl 4-bromo-2-hydroxybenzoate

To a solution of 4-bromo-2-hydroxybenzoic acid (15.0 g) in methanol (150 mL) was added dropwise thionyl chloride (10.1 mL) under ice-cooling, and the mixture was stirred overnight at 70°C under argon atmosphere. The reaction mixture was concentrated under reduced pressure, and the residue was purified by silica gel column chromatography (ethyl acetate/hexane) to give the title compound (14.2 g).

¹H NMR (300 MHz, DMSO-d₆) δ 3.88 (3H, s), 7.10-7.18 (1H, m), 7.21-7.28 (1H, m), 7.69 (1H, d, J = 8.3 Hz), 10.65 (1H, s).

[0373]

B) methyl 4-cyclopropyl-2-hydroxybenzoate

To a solution of methyl 4-bromo-2-hydroxybenzoate (3.00 g) in toluene (30.0 mL) were added tris(dibenzylideneacetone)dipalladium(0) (0.60 g), 2-dicyclohexylphosphino-2',6'-dimethoxybiphenyl (0.53 g), cyclopropylboronic acid (2.79 g) and sodium carbonate (3.44 g),

and the mixture was stirred overnight at 100°C under argon atmosphere. The reaction mixture was diluted with ethyl acetate, and the mixture was washed with water and saturated brine. The organic layer was dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (ethyl acetate/hexane) to give the title compound (1.80 g). ¹H NMR (300 MHz, DMSO-d₆) δ 0.70-0.80 (2H, m), 0.97-1.07 (2H, m), 1.85-1.99 (1H, m), 3.87 (3H, s), 6.61-6.73 (2H, m), 7.65 (1H, d, J = 7.9 Hz), 10.50 (1H, s).

[0374]

C) methyl 5-bromo-4-cyclopropyl-2-hydroxybenzoate

To a solution of methyl 4-cyclopropyl-2-hydroxybenzoate (1.80 g) in acetic acid (15.0 mL) was added dropwise bromine (1.57 g) under ice-cooling. The mixture was stirred at room temperature for 2 hr, to the reaction mixture was added water, and the resulting solid was collected by filtration. The obtained solid was dried under reduced pressure to give the title compound (2.28 g). ¹H NMR (300 MHz, DMSO-d₆) δ 0.69-0.82 (2H, m), 1.02-1.12 (2H, m), 2.04-2.18 (1H, m), 3.87 (3H, s), 6.58 (1H, s), 7.88 (1H, s), 10.38 (1H, s).

[0375]

D) methyl 4-cyclopropyl-2-hydroxy-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate

To a solution of methyl 5-bromo-4-cyclopropyl-2-hydroxybenzoate (1.05 g) in toluene (30 mL) were added bis(pinacolato)diboron (1.48 g), potassium acetate (1.14 g) and trans-dichlorobis(triphenylphosphine)palladium(II) (0.14 g), and the mixture was stirred at 110°C for 15 hr under argon atmosphere. The reaction mixture was allowed to be cooled to room temperature, water and ethyl acetate were added thereto, and the mixture was partitioned. The organic layer was washed with water and saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced

pressure. The residue was purified by silica gel column chromatography (ethyl acetate/hexane) to give the title compound (0.72 g).

MS: $[M+H]^+$ 319.2.

5 [0376]

E) methyl 5-(4-(1H-pyrazol-1-yl)benzyl)-4-cyclopropyl-2-hydroxybenzoate

To a solution of methyl 4-cyclopropyl-2-hydroxy-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (0.36 g) in a mixed solvent of DME (5.70 mL)-water (1.90 mL) were added
10 1-(4-(bromomethyl)phenyl)-1H-pyrazole (0.27 g), tetrakis(triphenylphosphine)palladium(0) (0.07 g) and sodium carbonate (0.24 g), and the mixture was stirred overnight at 80°C under argon atmosphere. The reaction mixture was allowed
15 to be cooled to room temperature, water and ethyl acetate were added thereto, and the mixture was partitioned. The organic layer was washed with water and saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel
20 column chromatography (ethyl acetate/hexane) to give the title compound (0.26 g).

MS: $[M+H]^+$ 349.1.

[0377]

F) methyl 5-(4-(1H-pyrazol-1-yl)benzyl)-4-cyclopropyl-2-
25 (((trifluoromethyl)sulfonyl)oxy)benzoate

To a solution of methyl 5-(4-(1H-pyrazol-1-yl)benzyl)-4-cyclopropyl-2-hydroxybenzoate (0.25 g) in DMF (5.00 mL) were added sodium hydride (0.04 g) and N-phenylbis(trifluoromethanesulfonimide) (0.29 g) under ice-
30 cooling, and the mixture was stirred at room temperature for 3 hr. To the reaction mixture was added 1N hydrochloric acid, and the mixture was extracted with ethyl acetate. The organic layer was washed with saturated aqueous sodium bicarbonate solution and saturated brine, and dried over anhydrous
35 magnesium sulfate, and the solvent was evaporated under reduced

pressure. The residue was purified by silica gel column chromatography (ethyl acetate/hexane) to give the title compound (0.33 g).

MS: $[M+H]^+$ 481.1.

5 [0378]

G) methyl 5-(4-(1H-pyrazol-1-yl)benzyl)-4-cyclopropyl-2-vinylbenzoate

To a solution of methyl 5-(4-(1H-pyrazol-1-yl)benzyl)-4-cyclopropyl-2-(((trifluoromethyl)sulfonyl)oxy)benzoate (0.33 g) in DMF (6.60 mL) were added tributylvinyltin (0.30 mL), trans-dichlorobis(triphenylphosphine)palladium(II) (0.02 g) and lithium chloride (0.22 g), and the mixture was stirred at 90°C for 1.5 hr under argon atmosphere. To the reaction mixture was added aqueous potassium fluoride solution, and the precipitated insoluble substance was removed by filtration through Celite. The filtrate was diluted with ethyl acetate, and the mixture was washed with water and saturated brine. The organic layer was dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (ethyl acetate/hexane) to give the title compound (0.22 g).

MS: $[M+H]^+$ 359.2.

[0379]

H) methyl 5-(4-(1H-pyrazol-1-yl)benzyl)-4-cyclopropyl-2-formylbenzoate

To a solution of methyl 5-(4-(1H-pyrazol-1-yl)benzyl)-4-cyclopropyl-2-vinylbenzoate (0.22 g) in a mixed solvent of acetone (4.40 mL)-acetonitrile (4.40 mL)-water (4.40 mL) were added osmium oxide (fixed catalyst I) (0.08 g) and sodium periodate (0.65 g), and the mixture was stirred overnight at room temperature. The reaction mixture was filtered, and the filtrate was extracted with ethyl acetate. The organic layer was washed with water and saturated brine, and dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure to give the title compound (0.22 g) as a crude

product. This compound was used in the next step without an additional purification.

MS: $[M+H]^+$ 361.2.

[0380]

- 5 I) rac-5-cyclopropyl-2-(trans-2-hydroxycyclopentyl)-6-(4-(1H-pyrazol-1-yl)benzyl)isoindolin-1-one

A solution of methyl 5-(4-(1H-pyrazol-1-yl)benzyl)-4-cyclopropyl-2-formylbenzoate (0.11 g), trans-2-aminocyclopentanol hydrochloride (0.04 g), triethylamine (0.04 mL) and anhydrous magnesium sulfate (0.07 g) in THF (2.20 mL) was stirred at room temperature for 5 hr. The insoluble substance was removed by filtration, and the filtrate was concentrated under reduced pressure. The residue was diluted with methanol (2.20 mL) and THF (2.20 mL), sodium triacetoxymethylborohydride (0.13 g) was added thereto, and the mixture was stirred overnight at room temperature. The reaction mixture was diluted with ethyl acetate, and the mixture was washed with water and saturated brine. The organic layer was dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (ethyl acetate/hexane) to give the title compound (0.06 g).

^1H NMR (300 MHz, DMSO- d_6) δ 0.62-0.70 (2H, m), 0.89-0.98 (2H, m), 1.47-1.75 (4H, m), 1.79-2.07 (3H, m), 4.06-4.25 (2H, m), 4.27 (2H, s), 4.38 (2H, s), 4.90 (1H, d, $J = 4.9$ Hz), 6.49-6.54 (1H, m), 7.19 (1H, s), 7.28 (2H, d, $J = 8.3$ Hz), 7.44 (1H, s), 7.70 - 7.78 (3H, m), 8.44 (1H, d, $J = 2.3$ Hz).

[0381]

Example 10

- 30 rac-4-chloro-2-(trans-2-hydroxycyclohexyl)-6-(4-(1H-pyrazol-1-yl)benzyl)-5-(trifluoromethyl)isoindolin-1-one

A) 2-chloro-1-(methoxymethoxy)-3-(trifluoromethyl)benzene

To a suspension of sodium hydride (3.78 g) in THF (150 mL) were added dropwise 2-chloro-3-hydroxybenzotrifluoride (12.4 g) and chloromethyl methyl ether (6.10 g) under ice-

cooling, and the mixture was stirred at 12°C for 16 hr. The reaction mixture was poured into saturated aqueous sodium chloride solution, and the mixture was extracted with ethyl acetate. The organic layer was dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure to give the title compound (16.8 g) as a crude product.

¹H NMR (400 MHz, CDCl₃) δ 3.53 (3H, s), 5.28 (2H, s), 7.27-7.32 (1H, m), 7.34-7.39 (2H, m).

[0382]

10 B) methyl 3-chloro-2-[(methoxycarbonyl)oxy]-4-(trifluoromethyl)benzoate

To a solution of 2-chloro-1-(methoxymethoxy)-3-(trifluoromethyl)benzene (11.8 g) in THF (100 mL) was added dropwise n-butyllithium (2.5M hexane solution) (21.0 mL) at 10°C, and the mixture was stirred for 2 hr. Then, to the reaction mixture was added dropwise a solution of methyl chloroformate (23.0 g) in THF (50.0 mL), and the mixture was stirred at 10°C for 16 hr. The reaction mixture was poured into saturated aqueous sodium bicarbonate solution, and the mixture was extracted with ethyl acetate (x 2). The organic layer was dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by flash silica gel column chromatography (ethyl acetate/petroleum ether) to give the title compound (6.00 g).

25 ¹H NMR (400 MHz, CDCl₃) δ 3.93 (3H, s), 3.99 (3H, s), 7.69 (1H, d, J = 8.4 Hz), 8.00 (1H, d, J = 8.4 Hz).

[0383]

C) methyl 3-chloro-2-hydroxy-4-(trifluoromethyl)benzoate

To a solution of methyl 3-chloro-2-[(methoxycarbonyl)oxy]-4-(trifluoromethyl)benzoate (6.00 g) in methanol (60.0 mL) was added potassium carbonate (8.00 g), and the mixture was stirred at 15°C for 16 hr. The reaction mixture was filtered, the filtrate was neutralized with 1N hydrochloric acid, and the mixture was extracted with ethyl acetate. The organic layer was washed with saturated brine,

and dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure to give the title compound (3.50 g).

¹H NMR (400 MHz, CDCl₃) δ 4.06 (3H, s), 7.29 (1H, d, J = 5.2 Hz), 7.90 (1H, d, J = 8.0 Hz), 11.55 (1H, brs).

[0384]

D) methyl 5-bromo-3-chloro-2-hydroxy-4-(trifluoromethyl)benzoate

To a solution of methyl 3-chloro-2-hydroxy-4-(trifluoromethyl)benzoate (6.30 g) in DMF (65.0 mL) was added N-bromosuccinimide (4.41 g), and the mixture was stirred at 15°C for 16 hr. The reaction mixture was poured into water, and the mixture was extracted with ethyl acetate. The organic layer was washed with saturated brine, and dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure to give the title compound (7.00 g).

¹H NMR (400 MHz, CDCl₃) δ 4.03 (3H, s), 8.13 (1H, s), 11.43 (1H, brs).

[0385]

E) methyl 3-chloro-2-hydroxy-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-4-(trifluoromethyl)benzoate

To a solution of methyl 5-bromo-3-chloro-2-hydroxy-4-(trifluoromethyl)benzoate (4.80 g) in toluene (70.0 mL) were added bis(pinacolato)diboron (5.48 g), potassium acetate (4.20 g) and trans-dichlorobis(triphenylphosphine)palladium(II) (0.51 g), and the mixture was stirred at 110°C for 16 hr under argon atmosphere. The reaction mixture was allowed to be cooled to room temperature, and filtered, and the filtrate was extracted with ethyl acetate. The organic layer was dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by flash silica gel column chromatography (ethyl acetate/petroleum ether) to give the title compound (2.10 g).

¹H NMR (400 MHz, CDCl₃) δ 1.37 (12H, s), 4.02 (3H, s), 7.88 (1H, s), 11.57 (1H, brs):

[0386]

F) ethyl 4-(1H-pyrazol-1-yl)benzoate

To a solution of ethyl 4-fluorobenzoate (105 g) in DMSO (250 mL) were added pyrazole (34.0 g) and potassium carbonate (138 g), and the mixture was heated with stirring at 130°C for 16 hr. The reaction mixture was diluted with water, and the mixture was extracted with ethyl acetate. The organic layer was washed with saturated brine, and dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure to give the title compound (73.7 g).

¹H NMR (400 MHz, CDCl₃) δ 1.40 (3H, t, J = 7.2 Hz), 4.38 (2H, q, J = 7.2 Hz), 6.49 (1H, t, J = 2.0 Hz), 7.75 (1H, d, J = 1.6 Hz), 7.77 (2H, d, J = 8.8 Hz), 7.99 (1H, d, J = 2.4 Hz), 8.12 (2H, d, J = 8.8 Hz).

[0387]

G) [4-(1H-pyrazol-1-yl)phenyl]methanol

To a solution of ethyl 4-(1H-pyrazol-1-yl)benzoate (73.7 g) in THF (500 mL) were added sodium borohydride (19.5 g) and calcium chloride (56.8 g) under ice-cooling, and the mixture was stirred at room temperature for 16 hr, and then heated with reflux for 2 days. The reaction mixture was diluted with 1N hydrochloric acid, and the mixture was extracted with ethyl acetate (x 4). The organic layer was washed with saturated brine, and dried over anhydrous sodium sulfate. The solvent was evaporated under reduced pressure, and the residue was washed with tert-butyl methyl ether to give the title compound (49.6 g).

¹H NMR (400 MHz, CDCl₃) δ 2.79 (1H, brs), 4.68 (2H, s), 6.45 (1H, t, J = 2.0 Hz), 7.38 (2H, d, J = 7.6 Hz), 7.61 (2H, d, J = 8.4 Hz), 7.70 (1H, s), 7.89 (1H, d, J = 1.6 Hz).

[0388]

H) 1-[4-(chloromethyl)phenyl]-1H-pyrazole

To a solution of [4-(1H-pyrazol-1-yl)phenyl]methanol (24.0 g) in 1,2-dichloroethane (200 mL) was added dropwise thionyl chloride (26.3 g) under ice-cooling, and the mixture

was stirred at room temperature for 16 hr. The reaction mixture was concentrated under reduced pressure, and the residue was washed with tert-butyl methyl ether to give the title compound (23.5 g).

5 ^1H NMR (400 MHz, CDCl_3) δ 4.62 (2H, s), 6.48 (1H, t, $J = 2.0$ Hz), 7.47 (2H, d, $J = 8.8$ Hz), 7.69 (2H, d, $J = 8.8$ Hz), 7.73 (1H, d, $J = 2.0$ Hz), 7.93 (1H, d, $J = 2.4$ Hz).

[0389]

I) methyl 3-chloro-2-hydroxy-5-[4-(1H-pyrazol-1-yl)benzyl]-4-(trifluoromethyl)benzoate and 3-chloro-2-hydroxy-5-[4-(1H-pyrazol-1-yl)benzyl]-4-(trifluoromethyl)benzoic acid

To a solution of methyl 3-chloro-2-hydroxy-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-4-(trifluoromethyl)benzoate (1.50 g) in 1,4-dioxane (30.0 mL) were added 1-(4-(chloromethyl)phenyl)-1H-pyrazole (0.76 g), tetrakis(triphenylphosphine)palladium(0) (0.23 g) and tripotassium phosphate trihydrate (2.10 g), and the mixture was stirred at 90°C for 16 hr under nitrogen atmosphere. The reaction mixture was allowed to be cooled to room temperature, water and ethyl acetate were added thereto, and the mixture was partitioned. The organic layer was dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by flash silica gel column chromatography (ethyl acetate/petroleum ether) to give methyl 3-chloro-2-hydroxy-5-[4-(1H-pyrazol-1-yl)benzyl]-4-(trifluoromethyl)benzoate (0.12 g) and 3-chloro-2-hydroxy-5-[4-(1H-pyrazol-1-yl)benzyl]-4-(trifluoromethyl)benzoic acid (0.70 g).

30 ^1H NMR (400 MHz, CDCl_3) δ 3.98 (3H, s), 4.21 (2H, d, $J = 2.0$ Hz), 6.40 (1H, t, $J = 2.0$ Hz), 7.12 (2H, d, $J = 8.4$ Hz), 7.62 (2H, d, $J = 8.4$ Hz), 7.68 (1H, s), 7.72 (1H, d, $J = 1.6$ Hz), 7.90 (1H, d, $J = 2.4$ Hz), 11.40 (1H, brs).

^1H NMR (400 MHz, CDCl_3) δ 3.49 (2H, s), 4.18 (1H, s), 6.40-6.52 (1H, m), 7.05-7.24 (2H, m), 7.44-7.58 (2H, m), 7.78-7.95 (3H, m). An active proton was not observed.

[0390]

J) methyl 3-chloro-5-[4-(1H-pyrazol-1-yl)benzyl]-4-(trifluoromethyl)-2-[[trifluoromethyl)sulfonyl]oxy}benzoate

To a solution of methyl 3-chloro-2-hydroxy-5-[4-(1H-pyrazol-1-yl)benzyl]-4-(trifluoromethyl)benzoate (0.15 g) in dichloromethane (5.00 mL) were added triethylamine (0.05 g) and trifluoromethanesulfonic anhydride (0.14 g) under ice-cooling, and the mixture was stirred at room temperature for 16 hr. To the reaction mixture was added water, and the mixture was extracted with ethyl acetate. The organic layer was dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure to give the title compound (0.24 g) as a crude product. This compound was used in the next step without an additional purification.

MS: [M+H]⁺ 542.9.

[0391]

K) methyl 3-chloro-2-ethenyl-5-[4-(1H-pyrazol-1-yl)benzyl]-4-(trifluoromethyl)benzoate

To a solution of methyl 3-chloro-5-[4-(1H-pyrazol-1-yl)benzyl]-4-(trifluoromethyl)-2-[[trifluoromethyl)sulfonyl]oxy}benzoate (0.24 g) in THF (5.00 mL)-water (5.00 mL) were added potassium vinyltrifluoroborate (0.07 g), cesium carbonate (0.24 g) and tetrakis(triphenylphosphine)palladium(0) (0.01 g), and the mixture was stirred at 90°C for 16 hr under nitrogen atmosphere. To the reaction mixture was added water, and the mixture was extracted with ethyl acetate. The organic layer was dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by prep-thin layer chromatography (ethyl acetate/petroleum ether) to give the title compound (0.08 g).

¹H NMR (400 MHz, CDCl₃) δ 3.83 (3H, s), 4.28 (2H, d, J = 2.0 Hz), 5.32 (1H, d, J = 18.0 Hz), 5.54 (1H, d, J = 12.4 Hz), 6.43-6.48 (1H, t, J = 2.4 Hz), 6.94 (1H, dd, J = 17.6, 11.2 Hz), 7.15 (2H, d, J = 8.4 Hz), 7.37 (1H, s), 7.63 (2H, d, J = 8.8

Hz), 7.71 (1H, d, $J = 1.2$ Hz), 7.90 (1H, d, $J = 2.4$ Hz).

[0392]

L) ethyl 3-chloro-2-hydroxy-5-[4-(1H-pyrazol-1-yl)benzyl]-4-(trifluoromethyl)benzoate

5 To a solution of 3-chloro-2-hydroxy-5-[4-(1H-pyrazol-1-yl)benzyl]-4-(trifluoromethyl)benzoic acid (0.60 g) in dichloromethane (20.0 mL) was added oxalyl chloride (0.29 g) under ice-cooling, and the mixture was stirred at 15°C for 16 hr. The reaction mixture was added to a solution of
10 triethylamine (2.00 mL) in ethanol (40.0 mL), and the mixture was stirred for 1 hr. The mixture was concentrated under reduced pressure, and the residue was diluted with ethyl acetate. The mixture was washed with 1N hydrochloric acid and saturated aqueous sodium bicarbonate solution, and the organic
15 layer was concentrated under reduced pressure. The residue was purified by flash silica gel column chromatography (ethyl acetate/petroleum ether) to give the title compound (0.20 g).
 ^1H NMR (400 MHz, CDCl_3) δ 1.41 (3H, t, $J = 7.2$ Hz), 4.22 (2H, d, $J = 2.0$ Hz), 4.45 (2H, q, $J = 7.2$ Hz), 6.42-6.49 (1H, m), 7.11
20 (2H, d, $J = 8.8$ Hz), 7.59-7.64 (2H, m), 7.66-7.73 (2H, m), 7.89 (1H, d, $J = 2.4$ Hz), 11.49 (1H, brs).

[0393]

M) ethyl 3-chloro-5-[4-(1H-pyrazol-1-yl)benzyl]-4-(trifluoromethyl)-2-[(trifluoromethyl)sulfonyl]oxybenzoate

25 To a solution of ethyl 3-chloro-2-hydroxy-5-[4-(1H-pyrazol-1-yl)benzyl]-4-(trifluoromethyl)benzoate (0.20 g) in dichloromethane (15.0 mL) were added triethylamine (0.10 g) and trifluoromethanesulfonic anhydride (0.27 g) under ice-cooling, and the mixture was stirred at room temperature for 16 hr. To
30 the reaction mixture was added water, and the mixture was extracted with ethyl acetate. The organic layer was dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure to give the title compound (0.25 g) as a crude product. This compound was used in the next step without an
35 additional purification.

MS: $[M+H]^+$ 557.0.

[0394]

N) ethyl 3-chloro-2-ethenyl-5-[4-(1H-pyrazol-1-yl)benzyl]-4-(trifluoromethyl)benzoate

5 To a solution of ethyl 3-chloro-5-[4-(1H-pyrazol-1-yl)benzyl]-4-(trifluoromethyl)-2-
{[(trifluoromethyl)sulfonyl]oxy}benzoate (0.25 g) in THF (8.00 mL)-water (5.00 mL) were added potassium vinyltrifluoroborate (0.13 g), cesium carbonate (0.46 g) and
10 tetrakis(triphenylphosphine)palladium(0) (0.03 g), and the mixture was stirred at 80°C for 16 hr under nitrogen atmosphere. To the reaction mixture was added water, and the mixture was extracted with ethyl acetate. The organic layer was dried over anhydrous sodium sulfate, and the solvent was evaporated under
15 reduced pressure. The residue was purified by thin layer chromatography (ethyl acetate/petroleum ether) to give the title compound (0.08 g).

^1H NMR (400 MHz, CDCl_3) δ 1.32 (3H, t, $J = 7.2$ Hz), 4.25-4.34 (4H, m), 5.34 (1H, d, $J = 17.6$ Hz), 5.54 (1H, d, $J = 12.4$ Hz),
20 6.46 (1H, t, $J = 2.0$ Hz), 6.94 (1H, dd, $J = 17.2, 11.2$ Hz), 7.15 (2H, d, $J = 8.0$ Hz), 7.36 (1H, s), 7.63 (2H, d, $J = 8.8$ Hz), 7.71 (1H, d, $J = 1.6$ Hz), 7.90 (1H, d, $J = 2.4$ Hz).

[0395]

O) 3-chloro-2-ethenyl-5-[4-(1H-pyrazol-1-yl)benzyl]-4-(trifluoromethyl)benzoic acid
25

To a solution of methyl 3-chloro-2-ethenyl-5-[4-(1H-pyrazol-1-yl)benzyl]-4-(trifluoromethyl)benzoate (0.80 g) and ethyl 3-chloro-2-ethenyl-5-[4-(1H-pyrazol-1-yl)benzyl]-4-(trifluoromethyl)benzoate (0.80 g) in THF (5.08 mL)-water (5.08
30 mL) were added lithium hydroxide monohydrate (0.20 g) and methanol (0.50 mL), and the mixture was stirred at 15°C for 16 hr. The reaction mixture was poured into 1N hydrochloric acid, and the mixture was extracted with ethyl acetate. The organic layer was dried over anhydrous sodium sulfate, and the solvent
35 was evaporated under reduced pressure to give the title

compound (0.15 g).

¹H NMR (400 MHz, CDCl₃) δ 4.28 (2H, s), 5.44 (1H, d, J = 17.2 Hz), 5.59 (1H, d, J = 12.0 Hz), 6.49 (1H, t, J = 2.0 Hz), 6.96 (1H, dd, J = 18.0, 11.6 Hz), 7.16 (2H, d, J = 8.4 Hz), 7.51 (1H, s), 7.59 (2H, d, J = 8.4 Hz), 7.80 (1H, d, J = 1.6 Hz), 7.90 (1H, d, J = 2.4 Hz).

[0396]

P) rac-3-chloro-2-ethenyl-N-(trans-2-hydroxycyclohexyl)-5-[4-(1H-pyrazol-1-yl)benzyl]-4-(trifluoromethyl)benzamide

To a solution of 3-chloro-2-ethenyl-5-[4-(1H-pyrazol-1-yl)benzyl]-4-(trifluoromethyl)benzoic acid (0.05 g), trans-2-aminocyclohexanol hydrochloride (0.03 g), 1-hydroxybenzotriazole (0.03 g) and 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (0.04 g) in dichloromethane (5.00 mL) was added triethylamine (0.04 g), and the mixture was stirred for 16 hr. The reaction mixture was poured into water, and the mixture was extracted with ethyl acetate. The organic layer was washed with 1N hydrochloric acid and saturated aqueous sodium bicarbonate solution, and dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure to give the title compound (0.07 g).

¹H NMR (400 MHz, CDCl₃) δ 1.25-1.40 (4H, m), 1.70-1.77 (2H, m), 1.90-2.00 (1H, m), 2.05-2.10 (1H, m), 3.32-3.40 (1H, m), 3.60-3.80 (1H, m), 4.26 (2H, s), 5.50-5.75 (3H, m), 6.45 (1H, t, J = 2.0 Hz), 6.85-6.98 (1H, m), 7.14 (2H, d, J = 8.4 Hz), 7.30 (1H, s), 7.61 (2H, d, J = 8.4 Hz), 7.71 (1H, d, J = 1.2 Hz), 7.89 (1H, d, J = 2.4 Hz).

[0397]

Q) rac-4-chloro-2-(trans-2-hydroxycyclohexyl)-6-(4-(1H-pyrazol-1-yl)benzyl)-5-(trifluoromethyl)isoindolin-1-one

To a solution of rac-3-chloro-2-ethenyl-N-(trans-2-hydroxycyclohexyl)-5-[4-(1H-pyrazol-1-yl)benzyl]-4-(trifluoromethyl)benzamide (0.07 g) in a mixed solvent of acetone (2.00 mL)-acetonitrile (2.00 mL)-water (2.00 mL) were added potassium osmate(VI) dihydrate (4.00 mg) and sodium

periodate (0.08 g), and the mixture was stirred for 16 hr. The reaction mixture was diluted with aqueous sodium thiosulfate solution, and the mixture was extracted with ethyl acetate. The organic layer was dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure to give rac-4-chloro-3-hydroxy-2-[trans-2-hydroxycyclohexyl]-6-[4-(1H-pyrazol-1-yl)benzyl]-5-(trifluoromethyl)-2,3-dihydro-1H-isoindol-1-one (0.05 g) as a crude product. This compound was used in the next step without an additional purification.

MS: $[M+H]^+$ 505.9.

To a solution of the above-mentioned compound (0.05 g) in dichloromethane (2.00 mL) was added trifluoroacetic acid (0.30 mL) under ice-cooling. After 15 min, triethylsilane (0.40 mL) was added thereto, and the mixture was stirred at 15°C for 16 hr. The reaction mixture was poured into saturated aqueous sodium bicarbonate solution, and the mixture was extracted with ethyl acetate (x 2). The organic layer was dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by prep-HPLC, then lyophilized to give the title compound (0.01 g).

^1H NMR (400 MHz, CDCl_3) δ 1.36-1.45 (2H, m), 1.54-1.62 (1H, m, overlapped with water signal), 1.83 (2H, d, $J = 12.0$ Hz), 1.92 (1H, d, $J = 12.0$ Hz), 2.13-2.26 (2H, m), 3.64-3.77 (1H, m), 4.03-4.13 (1H, m), 4.33 (2H, s), 4.37-4.55 (2H, m), 6.45 (1H, t, $J = 2.0$ Hz), 7.14 (2H, d, $J = 8.4$ Hz), 7.60 (2H, d, $J = 8.4$ Hz), 7.64 (1H, s), 7.70 (1H, d, $J = 1.6$ Hz), 7.89 (1H, d, $J = 2.4$ Hz).

[0398]

Example 11

rac-2-(trans-2-hydroxycyclohexyl)-4-methyl-1-oxo-6-(4-(1H-pyrazol-1-yl)benzyl)isoindoline-5-carbonitrile

A) 3-hydroxy-2-methylbenzonitrile

To a solution of 3-bromo-2-methylphenol (20.0 g) in DMF (250 mL) were added copper(I) cyanide (19.0 g) and tetrakis(triphenylphosphine)palladium(0) (3.70 g), and the

mixture was heated with stirring at 120°C for 16 hr under nitrogen atmosphere. The reaction mixture was poured into water, and the mixture was filtered. The filtrate was extracted with ethyl acetate, and the organic layer was dried over anhydrous sodium sulfate. The solvent was evaporated under reduced pressure, and the residue was washed with tert-butyl methyl ether to give the title compound (7.00 g).
¹H NMR (400 MHz, CDCl₃) δ 2.28 (3H, s), 7.00-7.15 (1H, m), 7.14-7.24 (2H, m), 10.11 (1H, brs).

10 [0399]

B) 4-formyl-3-hydroxy-2-methylbenzonitrile

To a solution of 3-hydroxy-2-methylbenzonitrile (7.00 g) in THF (100 mL) were added triethylamine (13.1 g), magnesium chloride (12.4 g) and paraformaldehyde (6.60 g), and the mixture was heated with reflux for 16 hr under nitrogen atmosphere. The reaction mixture was poured into 1N hydrochloric acid, and the mixture was extracted with ethyl acetate. The organic layer was dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure.
20 The residue was purified by flash silica gel column chromatography (ethyl acetate/petroleum ether) to give the title compound (1.30 g).

¹H NMR (400 MHz, CDCl₃) δ 2.50 (3H, s), 7.24-7.28 (2H, m, overlapped with CDCl₃ signal), 7.53 (1H, d, J = 8.0 Hz), 9.98 (1H, s), 11.42 (1H, brs).

25 [0400]

C) 4-cyano-2-hydroxy-3-methylbenzoic acid

To a solution of 4-formyl-3-hydroxy-2-methylbenzonitrile (3.90 g) in DMSO (21.0 mL) were added sodium dihydrogenphosphate (7.26 g) and an aqueous solution (16.0 mL) of sodium chlorite (5.46 g) under ice-cooling, and the mixture was stirred at 17°C for 16 hr. The reaction mixture was poured into saturated aqueous sodium carbonate solution, and the mixture was extracted with tert-butyl methyl ether. The aqueous layer was acidified with 1N hydrochloric acid, and the
35

mixture was extracted with ethyl acetate. The combined organic layers were dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure to give the title compound (4.00 g: containing DMSO).

5 ^1H NMR (400 MHz, DMSO- d_6) δ 2.36 (3H, s), 7.29 (1H, d, J = 8.0 Hz), 7.77 (1H, d, J = 8.0 Hz). An active proton was not observed.

[0401]

D) 5-bromo-4-cyano-2-hydroxy-3-methylbenzoic acid

10 To a solution of 4-cyano-2-hydroxy-3-methylbenzoic acid (5.50 g) in DMF (50.0 mL) was added N-bromosuccinimide (5.50 g) under ice-cooling, and the mixture was stirred at 15°C for 16 hr. The reaction mixture was poured into water, and the mixture was extracted with ethyl acetate. The organic layer
15 was dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure to give the title compound (7.40 g: containing DMF).

^1H NMR (400 MHz, CDCl_3) δ 2.51 (3H, s), 8.10 (1H, s), 11.48 (1H, brs).

20 [0402]

E) methyl 5-bromo-4-cyano-2-hydroxy-3-methylbenzoate

To a solution of 5-bromo-4-cyano-2-hydroxy-3-methylbenzoic acid (7.40 g) in dichloromethane (70.0 mL) were added oxalyl chloride (3.91 g) and DMF (0.20 mL) under ice-
25 cooling, and the mixture was stirred for 3 hr. The reaction mixture was concentrated under reduced pressure, and the residue was diluted with THF. To this mixture was added dropwise a solution of triethylamine (5.66 g) in methanol (50.0 mL) under ice-cooling, and the mixture was stirred for 20 min.
30 The reaction mixture was poured into 1N hydrochloric acid, and the mixture was extracted with ethyl acetate. The organic layer was dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (ethyl
35 acetate/petroleum ether) to give the title compound (4.20 g).

¹H NMR (400 MHz, CDCl₃) δ 2.51 (3H, s), 4.00 (3H, s), 7.97 (1H, s), 10.14 (1H, brs).

[0403]

F) methyl 4-cyano-2-hydroxy-3-methyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate

To a solution of methyl 5-bromo-4-cyano-2-hydroxy-3-methylbenzoate (2.00 g) in toluene (30.0 mL) were added bis(pinacolato)diboron (2.82 g), potassium acetate (2.18 g) and trans-dichlorobis(triphenylphosphine)palladium(II) (0.26 g), and the mixture was stirred at 110°C for 16 hr under argon atmosphere. The reaction mixture was poured into water, and the mixture was extracted with ethyl acetate. The organic layer was washed with saturated brine, and dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by flash silica gel column chromatography (ethyl acetate/petroleum ether) to give the title compound (2.50 g: containing bis(pinacolato)diboron). ¹H NMR (400 MHz, CDCl₃) δ 1.38 (12H, s), 2.49 (3H, s), 3.99 (3H, s), 8.20 (1H, s), 11.43 (1H, brs).

[0404]

G) methyl 4-cyano-2-hydroxy-3-methyl-5-[4-(1H-pyrazol-1-yl)benzyl]benzoate

To a solution of methyl 4-cyano-2-hydroxy-3-methyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (3.50 g) in 1,4-dioxane (30.0 mL) were added 1-(4-(chloromethyl)phenyl)-1H-pyrazole (2.12 g), tripotassium phosphate trihydrate (5.90 g) and [1,1'-bis(diphenylphosphino)ferrocene]dichloropalladium(II) dichloromethane adduct (0.34 g), and the mixture was stirred at 90°C for 16 hr under nitrogen atmosphere. The reaction mixture was poured into water, and the mixture was extracted with ethyl acetate. The organic layer was dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by flash silica gel column chromatography (ethyl acetate/petroleum ether) to give the

title compound (0.86 g).

¹H NMR (400 MHz, CDCl₃) δ 2.49 (3H, s), 3.95 (3H, s), 4.15 (2H, s), 6.46 (1H, s), 7.30 (2H, d, J = 8.4 Hz), 7.59-7.60 (1H, m), 7.64 (2H, d, J = 8.4 Hz), 7.71 (1H, s), 7.90 (1H, d, J = 2.4 Hz), 11.07 (1H, brs).

[0405]

H) methyl 4-cyano-3-methyl-5-[4-(1H-pyrazol-1-yl)benzyl]-2-[[trifluoromethyl)sulfonyl]oxy}benzoate

To a solution of methyl 4-cyano-2-hydroxy-3-methyl-5-[4-(1H-pyrazol-1-yl)benzyl]benzoate (0.86 g) in THF (20.0 mL) was added sodium hydride (0.12 g) under ice-cooling, and the mixture was stirred for 20 min. To this reaction mixture was added N-phenylbis(trifluoromethanesulfonylimide) (1.77 g), and the mixture was stirred at 15°C for 16 hr. The reaction mixture was poured into saturated aqueous ammonium chloride solution, and the mixture was extracted with ethyl acetate. The organic layer was dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure to give the title compound (3.00 g) as a crude product. This compound was used in the next step without an additional purification.

MS: [M+H]⁺ 479.8.

[0406]

I) methyl 4-cyano-2-ethenyl-3-methyl-5-[4-(1H-pyrazol-1-yl)benzyl]benzoate

To a solution of methyl 4-cyano-3-methyl-5-[4-(1H-pyrazol-1-yl)benzyl]-2-[[trifluoromethyl)sulfonyl]oxy}benzoate (2.80 g) in THF (25.0 mL)-water (5.00 mL) were added potassium vinyltrifluoroborate (0.85 g), cesium carbonate (6.23 g) and tetrakis(triphenylphosphine)palladium(0) (0.38 g), and the mixture was stirred at 80°C for 16 hr under nitrogen atmosphere. To the reaction mixture was added water, and the mixture was extracted with ethyl acetate. The organic layer was dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by flash silica gel column chromatography (ethyl acetate/petroleum ether) to give

the title compound (0.80 g).

¹H NMR (400 MHz, CDCl₃) δ 2.56 (3H, s), 3.83 (3H, s), 4.23 (2H, s), 5.21 (1H, d, J = 17.6 Hz), 5.46-5.60 (1H, m), 6.46 (1H, s), 6.88 (1H, dd, J = 17.6, 11.4 Hz), 7.32 (2H, d, J = 8.4 Hz),
5 7.41 (1H, s), 7.64 (2H, d, J = 8.4 Hz), 7.71 (1H, s), 7.90 (1H, d, J = 2.4 Hz).

[0407]

J) 4-cyano-2-ethenyl-3-methyl-5-[4-(1H-pyrazol-1-yl)benzyl]benzoic acid

10 To a solution of methyl 4-cyano-2-ethenyl-3-methyl-5-[4-(1H-pyrazol-1-yl)benzyl]benzoate (0.70 g) in THF (5.00 mL)-water (5.00 mL) were added lithium hydroxide monohydrate (0.25 g) and methanol (1.00 mL), and the mixture was stirred at 10°C for 16 hr. The reaction mixture was diluted with water, and
15 the mixture was extracted with tert-butyl methyl ether. The aqueous layer was acidified with 2N hydrochloric acid, and the mixture was extracted with ethyl acetate. The organic layer was dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure to give the title compound
20 (0.60 g).

¹H NMR (400 MHz, CDCl₃) δ 2.55 (3H, s), 4.22 (2H, s), 5.25 (1H, dd, J = 17.8, 1.4 Hz), 5.55 (1H, dd, J = 11.6, 1.2 Hz), 6.47 (1H, t, J = 2.0 Hz), 6.93 (1H, dd, J = 17.8, 11.4 Hz), 7.34 (2H, d, J = 8.4 Hz), 7.58 (1H, s), 7.62 (2H, d, J = 8.8 Hz), 7.79
25 (1H, d, J = 1.6 Hz), 7.90 (1H, d, J = 2.4 Hz). One active proton was not observed.

[0408]

K) rac-4-cyano-2-ethenyl-N-(trans-2-hydroxycyclohexyl)-3-methyl-5-[4-(1H-pyrazol-1-yl)benzyl]benzamide

30 To a solution of 4-cyano-2-ethenyl-3-methyl-5-[4-(1H-pyrazol-1-yl)benzyl]benzoic acid (0.10 g), trans-2-aminocyclohexanol hydrochloride (0.06 g), 1-hydroxybenzotriazole (0.06 g) and 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (0.08 g) in dichloromethane
35 (5.00 mL) was added triethylamine (0.11 g), and the mixture was

stirred for 16 hr. The reaction mixture was poured into 1N hydrochloric acid, and the mixture was extracted with ethyl acetate. The organic layer was washed with 1N hydrochloric acid and saturated aqueous sodium bicarbonate solution, and
5 dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure to give the title compound (0.10 g).

^1H NMR (400 MHz, CDCl_3) δ 1.11-1.41 (6H, m), 1.95-2.10 (2H, m), 2.55 (3H, s), 3.21-3.30 (1H, m), 3.36 (1H, td, $J = 10.0, 4.4$
10 Hz), 3.68-3.80 (1H, m), 4.20 (2H, s), 5.47 (1H, dd, $J = 17.8, 1.2$ Hz), 5.61 (1H, dd, $J = 11.6, 1.2$ Hz), 5.70 (1H, d, $J = 7.2$ Hz), 6.45 (1H, t, $J = 2.0$ Hz), 6.78 (1H, dd, $J = 17.8, 11.6$ Hz), 7.25-7.26 (1H, m, overlapped with CDCl_3 signal), 7.32 (2H, d, $J = 8.4$ Hz), 7.62 (2H, d, $J = 8.4$ Hz), 7.70 (1H, d, $J = 2.0$ Hz),
15 7.89 (1H, d, $J = 2.4$ Hz).

[0409]

L) rac-3-hydroxy-2-(trans-2-hydroxycyclohexyl)-4-methyl-1-oxo-6-[4-(1H-pyrazol-1-yl)benzyl]-2,3-dihydro-1H-isoindole-5-carbonitrile

20 To a solution of rac-4-cyano-2-ethenyl-N-(trans-2-hydroxycyclohexyl)-3-methyl-5-[4-(1H-pyrazol-1-yl)benzyl]benzamide (0.09 g) in a mixed solvent of acetone (2.00 mL)-acetonitrile (2.00 mL)-water (2.00 mL) were added potassium osmate(VI) dihydrate (7.00 mg) and sodium periodate
25 (0.18 g), and the mixture was stirred for 16 hr. The reaction mixture was diluted with saturated aqueous sodium thiosulfate solution, and the mixture was extracted with ethyl acetate. The organic layer was dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure to give the
30 title compound (0.10 g) as a crude product. This compound was used in the next step without an additional purification.

MS: $[\text{M}+\text{H}]^+$ 443.0.

[0410]

M) rac-2-(trans-2-hydroxycyclohexyl)-4-methyl-1-oxo-6-(4-(1H-pyrazol-1-yl)benzyl)isoindoline-5-carbonitrile
35

To a solution of rac-3-hydroxy-2-(trans-2-hydroxycyclohexyl)-4-methyl-1-oxo-6-[4-(1H-pyrazol-1-yl)benzyl]-2,3-dihydro-1H-isoindole-5-carbonitrile (0.10 g) in dichloromethane (2.00 mL) was added trifluoroacetic acid (0.31 mL) under ice-cooling. After 15 min, triethylsilane (0.50 mL) was added thereto, and the mixture was stirred at 15°C for 16 hr. The reaction mixture was poured into saturated aqueous sodium bicarbonate solution, and the mixture was extracted with ethyl acetate (x 2). The organic layer was dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by prep-HPLC, then lyophilized to give the title compound (0.02 g).

¹H NMR (400 MHz, CDCl₃) δ 1.35-1.52 (3H, m), 1.85 (2H, d, J = 12.4 Hz), 1.94 (1H, d, J = 13.6 Hz), 2.10 (1H, d, J = 6.8 Hz), 2.20 (1H, d, J = 12.0 Hz), 2.57 (3H, s), 3.55-3.65 (1H, m), 4.07-4.16 (1H, m), 4.30-4.48 (4H, m), 6.41-6.51 (1H, m), 7.35 (2H, d, J = 8.4 Hz), 7.61-7.69 (3H, m), 7.72 (1H, d, J = 1.6 Hz), 7.91 (1H, d, J = 2.4 Hz). One active proton was observed. [0411]

Example 12

rac-4-chloro-2-(trans-2-hydroxycyclohexyl)-5-methoxy-6-(4-(1H-pyrazol-1-yl)benzyl)isoindolin-1-one

A) 2-chloro-3-methoxyphenol

To an aqueous solution (100 mL) of potassium hydroxide (11.0 g) was added 2-chlorobenzene-1,3-diol (22.8 g). Then, dimethyl sulfate (19.9 g) was slowly added to the reaction mixture while keeping the reaction mixture at 10°C to 20°C, and the mixture was stirred at 100°C for 2 days. The reaction mixture was acidified with 2N hydrochloric acid, and the mixture was extracted with dichloromethane. The organic layer was dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (ethyl acetate/petroleum ether) to give the title compound (18.0 g).

¹H NMR (400 MHz, DMSO-d₆) δ 3.78 (3H, s), 6.54-6.58 (2H, m),

7.05 (1H, t, J = 8.4 Hz), 10.04 (1H, brs).

[0412]

B) 3-chloro-2-hydroxy-4-methoxybenzaldehyde

To a solution of 2-chloro-3-methoxyphenol (19.3 g) in
5 1,2-dichloroethane (150 mL) were added triethylamine (73.7 g),
magnesium chloride (57.8 g) and paraformaldehyde (36.5 g), and
the mixture was stirred at 70°C for 4 hr under nitrogen
atmosphere. The reaction mixture was poured into 1N
hydrochloric acid, and the precipitate was removed by
10 filtration, and washed with dichloromethane. The organic layer
was washed with 1N hydrochloric acid and saturated brine, and
dried over anhydrous sodium sulfate, and the solvent was
evaporated under reduced pressure to give the title compound
(21.5 g).

15 ¹H NMR (400 MHz, DMSO-d₆) δ 3.97 (3H, s), 6.92 (1H, d, J = 8.8
Hz), 7.76 (1H, d, J = 9.2 Hz), 9.95 (1H, s), 11.47 (1H, brs).

[0413]

C) 3-chloro-2-hydroxy-4-methoxybenzoic acid

To a solution of 3-chloro-2-hydroxy-4-methoxybenzaldehyde
20 (16.6 g) and sodium dihydrogenphosphate (34.7 g) in a mixed
solvent of DMSO (180 mL)-water (45.0 mL) was added an aqueous
solution (35.0 mL) of sodium chlorite (27.2 g) under ice-
cooling, and the mixture was stirred at 15°C for 16 hr. The
reaction mixture was poured into saturated aqueous sodium
25 carbonate solution, the mixture was diluted with water, and
filtered, and the filtrate was extracted with petroleum
ether/ethyl acetate=5:1. The aqueous layer was acidified with
conc. hydrochloric acid to adjusted pH=1, and the mixture was
extracted with ethyl acetate. The organic layer was washed
30 with 1N hydrochloric acid, and dried over anhydrous sodium
sulfate, and the solvent was evaporated under reduced pressure
to give the title compound (11.1 g).

¹H NMR (400 MHz, DMSO-d₆) δ 3.93 (3H, s), 6.77 (1H, d, J = 9.2
Hz), 7.78 (1H, d, J = 9.2 Hz), 12.11 (1H, brs). One active
35 proton was not observed.

[0414]

D) 5-bromo-3-chloro-2-hydroxy-4-methoxybenzoic acid

To a solution of 3-chloro-2-hydroxy-4-methoxybenzoic acid (11.1 g) in acetic acid (275 mL) was added bromine (8.79 g) at room temperature, and the mixture was stirred for 16 hr. To the reaction mixture was added water, and the precipitate was collected by filtration. The precipitate was washed with water, and dissolved in ethyl acetate, and the solution was washed with water and saturated brine. The organic layer was dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure to give the title compound (10.4 g). ^1H NMR (400 MHz, DMSO- d_6) δ 3.87 (3H, s), 7.94 (1H, s). Two active protons were not observed.

[0415]

E) methyl 5-bromo-3-chloro-2-hydroxy-4-methoxybenzoate

To a solution of 5-bromo-3-chloro-2-hydroxy-4-methoxybenzoic acid (10.4 g) in dichloromethane (150 mL) were added a solution of oxalyl chloride (7.05 g) in dichloromethane (10 mL) and DMF (3 drops) under ice-cooling, and the mixture was stirred at 13°C for 1 hr. The solvent was evaporated under reduced pressure, and the residue was dissolved in THF (50 mL). This solution was added dropwise to a solution of triethylamine (1.20 g) in methanol (150 mL) under ice-cooling, and the mixture was stirred at 13°C for 16 hr. The solvent was evaporated under reduced pressure from the reaction mixture, and the residue was diluted with ethyl acetate and water. 1N Hydrochloric acid was added thereto, and the mixture was partitioned. The organic layer was washed with water and saturated brine, and dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure to give the title compound (10.9 g).

^1H NMR (400 MHz, DMSO- d_6) δ 3.86 (3H, s), 3.91 (3H, s), 7.94 (1H, s), 11.11 (1H, brs).

[0416]

F) methyl 5-bromo-3-chloro-4-methoxy-2-(methoxymethoxy)benzoate

To a solution of methyl 5-bromo-3-chloro-2-hydroxy-4-methoxybenzoate (5.00 g) in THF (120 mL) was added sodium hydride (1.01 g) under ice-cooling, and the mixture was stirred for 1 hr under nitrogen atmosphere. To this reaction mixture
5 was added dropwise chloromethyl methyl ether (1.63 g), and the mixture was stirred at 20°C for 16 hr. To the reaction mixture was added water, and the mixture was extracted with ethyl acetate. The organic layer was washed with saturated brine, and dried over anhydrous sodium sulfate. The solvent was
10 evaporated under reduced pressure, and the residue was purified by flash silica gel column chromatography (ethyl acetate/petroleum ether) to give the title compound (3.08 g).
¹H NMR (400 MHz, CDCl₃) δ 3.65 (3H, s), 3.93 (3H, s), 3.96 (3H, s), 5.15 (2H, s), 8.00 (1H, s).

15 [0417]

G) 1-{4-[(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)methyl]phenyl}-1H-pyrazole

To a solution of 1-[4-(chloromethyl)phenyl]-1H-pyrazole (2.20 g) in 1,4-dioxane (50.0 mL) were added
20 bis(pinacolato)diboron (3.47 g), potassium acetate (3.35 g) and [1,1'-bis(diphenylphosphino)ferrocene]dichloropalladium(II) dichloromethane adduct (0.47 g), and the mixture was stirred at 90°C for 16 hr under argon atmosphere. The reaction mixture was allowed to be cooled to room temperature, and filtered, and
25 the solvent was evaporated under reduced pressure from the filtrate. The residue was purified by silica gel column chromatography (ethyl acetate/petroleum ether) to give the title compound (3.10 g).

¹H NMR (400 MHz, CDCl₃) δ 1.22 (12H, s), 2.32 (2H, s), 6.43 (1H, t, J = 2.0 Hz), 7.23-7.29 (2H, m, overlap with CDCl₃ signal),
30 7.54 (2H, d, J = 8.4 Hz), 7.69 (1H, d, J = 1.2 Hz), 7.87 (1H, d, J = 2.8 Hz).

[0418]

H) methyl 3-chloro-2-hydroxy-4-methoxy-5-[4-(1H-pyrazol-1-yl)benzyl]benzoate
35

To a solution of methyl 5-bromo-3-chloro-4-methoxy-2-(methoxymethoxy)benzoate (3.00 g) in 1,4-dioxane (50.0 mL)-water (5.00 mL) were added 1-{4-[(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)methyl]phenyl}-1H-pyrazole (7.50 g),
5 potassium carbonate (2.44 g) and tetrakis(triphenylphosphine)palladium(0) (1.02 g), and the mixture was stirred at 90°C for 16 hr under nitrogen atmosphere. The reaction mixture was poured into water, and the mixture was extracted with ethyl acetate. The organic layer was dried over
10 anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by flash silica gel column chromatography (ethyl acetate/petroleum ether) to give methyl 3-chloro-4-methoxy-2-(methoxymethoxy)-5-[4-(1H-pyrazol-1-yl)benzyl]benzoate (0.40 g).

15 To a solution of the above-mentioned compound (0.40 g) in ethyl acetate (5.00 mL) was added 4N hydrogen chloride/ethyl acetate (25.0 mL), and the mixture was stirred at room temperature for 16 hr. The reaction mixture was poured into saturated aqueous sodium bicarbonate solution, and the mixture
20 was extracted with ethyl acetate. The organic layer was dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by flash silica gel column chromatography (ethyl acetate/petroleum ether) to give the title compound (0.23 g).

25 ¹H NMR (400 MHz, CDCl₃) δ 3.78 (3H, s), 3.93 (3H, s), 3.97 (2H, s), 6.45 (1H, t, J = 2.0 Hz), 7.24 (1H, s), 7.27 (1H, s, overlapped with CDCl₃ signal), 7.58-7.63 (3H, m), 7.71 (1H, d, J = 1.6 Hz), 7.89 (1H, d, J = 2.4 Hz), 11.36 (1H, brs).

[0419]

30 I) methyl 3-chloro-4-methoxy-5-[4-(1H-pyrazol-1-yl)benzyl]-2-[[trifluoromethyl)sulfonyl]oxy]benzoate

To a solution of methyl 3-chloro-2-hydroxy-4-methoxy-5-[4-(1H-pyrazol-1-yl)benzyl]benzoate (0.23 g) in dichloromethane (10.0 mL) were added triethylamine (0.13 g) and
35 trifluoromethanesulfonic anhydride (0.35 g) under ice-cooling,

and the mixture was stirred at room temperature for 16 hr. To the reaction mixture was added water, and the mixture was extracted with ethyl acetate. The organic layer was dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure to give the title compound (0.36 g) as a crude product. This compound was used in the next step without an additional purification.

MS: $[M+H]^+$ 505.0.

[0420]

10 J) methyl 3-chloro-2-ethenyl-4-methoxy-5-[4-(1H-pyrazol-1-yl)benzyl]benzoate

To a solution of methyl 3-chloro-4-methoxy-5-[4-(1H-pyrazol-1-yl)benzyl]-2-[[trifluoromethyl)sulfonyl]oxy]benzoate (0.36 g) in THF (8.00 mL)-water (3.00 mL) were added potassium vinyltrifluoroborate (0.17 g), cesium carbonate (0.60 g) and tetrakis(triphenylphosphine)palladium(0) (0.04 g), and the mixture was heated with reflux for 16 hr under nitrogen atmosphere. To the reaction mixture was added water, and the mixture was extracted with ethyl acetate. The organic layer was dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by thin layer chromatography (ethyl acetate/petroleum ether) to give the title compound (0.08 g).

^1H NMR (400 MHz, CDCl_3) δ 3.76 (3H, s), 3.81 (3H, s), 4.05 (2H, s), 5.33 (1H, d, $J = 17.6$ Hz), 5.52 (1H, dd, $J = 11.2, 1.2$ Hz), 6.45 (1H, t, $J = 2.0$ Hz), 6.93 (1H, dd, $J = 17.8, 11.2$ Hz), 7.28 (2H, d, $J = 8.4$ Hz), 7.43 (1H, s), 7.62 (2H, d, $J = 8.4$ Hz), 7.71 (1H, d, $J = 1.4$ Hz), 7.89 (1H, d, $J = 2.4$ Hz).

[0421]

30 K) 3-chloro-2-ethenyl-4-methoxy-5-[4-(1H-pyrazol-1-yl)benzyl]benzoic acid

To a solution of methyl 3-chloro-2-ethenyl-4-methoxy-5-[4-(1H-pyrazol-1-yl)benzyl]benzoate (0.10 g) in THF (3.00 mL)-water (3.00 mL) and methanol (0.50 mL) was added lithium hydroxide monohydrate (0.10 g), and the mixture was stirred at

20°C for 16 hr. The reaction mixture was poured into 1N hydrochloric acid, and the mixture was extracted with ethyl acetate. The organic layer was dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure
5 to give the title compound (0.08 g).

¹H NMR (400 MHz, CDCl₃) δ 3.78 (3H, s), 4.22 (2H, s), 4.06 (2H, s), 5.42 (1H, d, J = 17.6 Hz), 5.58 (1H, d, J = 11.2 Hz), 6.46 (1H, t, J = 2.0 Hz), 6.94 (1H, dd, J = 18.0, 11.6 Hz), 7.25-7.40 (2H, m, overlap with CDCl₃ signal), 7.50-7.65 (3H, m),
10 7.72 (1H, s), 7.89 (1H, d, J = 2.4 Hz). An active proton was not observed.

[0422]

L) rac-3-chloro-2-ethenyl-N-(trans-2-hydroxycyclohexyl)-4-methoxy-5-[4-(1H-pyrazol-1-yl)benzyl]benzamide

15 To a solution of 3-chloro-2-ethenyl-4-methoxy-5-[4-(1H-pyrazol-1-yl)benzyl]benzoic acid (0.06 g), trans-2-aminocyclohexanol hydrochloride (0.04 g), 1-hydroxybenzotriazole (0.03 g) and 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (0.05 g) in dichloromethane
20 (10.0 mL) was added triethylamine (0.03 g), and the mixture was stirred at room temperature for 16 hr. The reaction mixture was poured into water, and the mixture was extracted with ethyl acetate. The organic layer was dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure
25 to give the title compound (0.08 g).

¹H NMR (400 MHz, CDCl₃) δ 1.31-1.39 (3H, m), 1.70-1.76 (2H, m), 1.97-2.12 (3H, m), 3.23 (1H, brs), 3.30-3.40 (1H, m), 3.65-3.75 (4H, m), 4.03 (2H, s), 5.56-5.65 (2H, m), 5.69 (1H, d, J = 7.6 Hz), 6.45 (1H, t, J = 2.0 Hz), 6.88 (1H, dd, J = 18.0, 11.2 Hz),
30 7.27-7.30 (2H, m, overlapped with CDCl₃ signal), 7.61 (2H, d, J = 8.4 Hz), 7.70 (1H, d, J = 1.6 Hz), 7.89 (1H, d, J = 2.0 Hz).

[0423]

M) rac-4-chloro-3-hydroxy-2-(trans-2-hydroxycyclohexyl)-5-methoxy-6-[4-(1H-pyrazol-1-yl)benzyl]-2,3-dihydro-1H-isoindol-
35 1-one

To a solution of rac-3-chloro-2-ethenyl-N-(trans-2-hydroxycyclohexyl)-4-methoxy-5-[4-(1H-pyrazol-1-yl)benzyl]benzamide (0.08 g) in a mixed solvent of acetone (2.00 mL)-acetonitrile (2.00 mL)-water (2.00 mL) were added
5 potassium osmate(VI) dihydrate (5.00 mg) and sodium periodate (0.12 g), and the mixture was stirred at room temperature for 16 hr. The reaction mixture was diluted with saturated aqueous sodium thiosulfate solution, and the mixture was extracted with ethyl acetate. The organic layer was dried over anhydrous
10 sodium sulfate, and the solvent was evaporated under reduced pressure to give the title compound (0.07 g) as a crude product. This compound was used in the next step without an additional purification.

MS: $[M+Na]^+$ 450.1.

15 [0424]

N) rac-4-chloro-2-(trans-2-hydroxycyclohexyl)-5-methoxy-6-(4-(1H-pyrazol-1-yl)benzyl)isoindolin-1-one

To a solution of rac-4-chloro-3-hydroxy-2-(trans-2-hydroxycyclohexyl)-5-methoxy-6-[4-(1H-pyrazol-1-yl)benzyl]-2,3-
20 dihydro-1H-isoindol-1-one (0.07 g) in dichloromethane (2.00 mL) was added trifluoroacetic acid (0.50 mL) under ice-cooling, and the mixture was stirred for 15 min. Then, triethylsilane (0.40 mL) was added thereto, and the mixture was stirred at room temperature for 1 hr. The reaction mixture was poured into
25 saturated aqueous sodium bicarbonate solution, and the mixture was extracted with ethyl acetate (x 2). The organic layer was dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by prep-HPLC, then lyophilized to give the title compound (0.02 g).
30 ^1H NMR (400 MHz, CDCl_3) δ 1.35-1.45 (3H, m), 1.50-1.60 (1H, m, overlapped with water signal), 1.81 (2H, d, $J = 12.0$ Hz), 1.91 (1H, d, $J = 12.0$ Hz), 2.13-2.33 (2H, m), 3.60-3.70 (1H, m), 3.78 (3H, s), 4.03-4.12 (3H, m), 4.37 (2H, q, $J = 17.2$ Hz), 6.44 (1H, t, $J = 2.0$ Hz), 7.20-7.30 (2H, m, overlapped with
35 CDCl_3 signal), 7.57-7.62 (3H, m), 7.70 (1H, d, $J = 1.2$ Hz),

7.88 (1H, d, J = 2.4 Hz).

[0425]

Example 13

2-((1S,2S)-2-hydroxycyclopentyl)-4,5-dimethyl-6-(4-(1-methyl-
5 1H-pyrazol-3-yl)benzyl)isoindolin-1-one

A) 1-methyl-1H-pyrazol-3-yl trifluoromethanesulfonate

To a solution of 1-methyl-1H-pyrazol-3-ol (2.92 g) in
pyridine (50.0 mL) was added trifluoromethanesulfonic anhydride
(6.03 mL) under ice-cooling, and the mixture was stirred for 1
10 hr. The reaction mixture was diluted with saturated aqueous
sodium bicarbonate solution, and the mixture was extracted with
ethyl acetate. The organic layer was washed with saturated
brine, and dried over anhydrous sodium sulfate, and the solvent
was evaporated under reduced pressure. The residue was
15 purified by silica gel column chromatography (ethyl
acetate/hexane) to give the title compound (5.44 g).

¹H NMR (300MHz, CDCl₃) δ 3.87 (3H, s), 6.11 (1H, d, J = 2.3 Hz),
7.32 (1H, d, J = 2.3 Hz).

[0426]

20 B) [4-(1-methyl-1H-pyrazol-3-yl)phenyl]methanol

A mixture of [4-(hydroxymethyl)phenyl]boronic acid (4.46
g), 1-methyl-1H-pyrazol-3-yl trifluoromethanesulfonate (4.50 g),
cesium carbonate (19.1 g) and [1,1'-
bis(diphenylphosphino)ferrocene]dichloropalladium(II)
25 dichloromethane adduct (0.80 g) in toluene (15.0 mL)-ethanol
(1.00 mL)-water (1.00 mL) was subjected to microwave
irradiation at 150°C for 1 hr. The reaction mixture was
diluted with ethyl acetate, and the insoluble substance was
removed by filtration. The filtrate was washed successively
30 with saturated aqueous sodium bicarbonate solution and
saturated brine, and dried over anhydrous sodium sulfate, and
the solvent was evaporated under reduced pressure. The residue
was purified by silica gel column chromatography (ethyl
acetate/hexane) to give the title compound (1.00 g).

35 ¹H NMR (300MHz, CDCl₃) δ 3.95 (3H, s), 4.71 (2H, d, J = 6.0 Hz),

6.54 (1H, d, $J = 2.3$ Hz), 7.33-7.45 (1H, m), 7.73-7.87 (1H, m).
[0427]

C) 3-[4-(chloromethyl)phenyl]-1-methyl-1H-pyrazole

To a solution of [4-(1-methyl-1H-pyrazol-3-yl)phenyl]methanol (1.00 g) in THF (15.0 mL) was added dropwise
5 thionyl chloride (0.58 mL) under ice-cooling, and the mixture
was stirred at 17°C for 16 hr. The reaction mixture was
diluted with water and ethyl acetate, and saturated aqueous
sodium bicarbonate was added thereto. The organic layer was
10 washed with saturated brine, and dried over anhydrous sodium
sulfate, and the solvent was evaporated under reduced pressure.
The residue was purified by silica gel column chromatography
(ethyl acetate/hexane) to give the title compound (0.65 g).
 ^1H NMR (300MHz, CDCl_3) δ 3.95 (3H, s), 4.61 (2H, s), 6.49-6.57
15 (1H, m), 7.33-7.44 (3H, m), 7.74-7.82 (2H, m).
[0428]

D) 2-amino-5-bromo-3,4-dimethylbenzoic acid

To a solution of 2-amino-3,4-dimethylbenzoic acid (50.0
g) in DMSO (500 mL) was added hydrobromic acid (174 mL) while
20 keeping the internal temperature at 25 to 30°C, and the mixture
was stirred overnight at room temperature. To the reaction
mixture was added water (500 mL), and the mixture was stirred
for 30 min. The precipitate was collected by filtration, and
washed with water to give the title compound (102 g: containing
25 DMSO).

MS: $[\text{M}-\text{H}]^+$ 241.9.

[0429]

E) methyl 2-amino-5-bromo-3,4-dimethylbenzoate

To a solution of 2-amino-5-bromo-3,4-dimethylbenzoic acid
30 (73.9 g) in DMF (750 mL) was added cesium carbonate (148 g),
and the mixture was stirred at room temperature for 30 min. To
this reaction mixture was added dropwise methyl iodide (22.7
mL) at room temperature, and the mixture was stirred overnight.
To the reaction mixture was added water, and the mixture was
35 extracted with ethyl acetate. The organic layer was washed

with water and saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (ethyl acetate/hexane) to give the title

5 compound (56.8 g).

MS: $[M+H]^+$ 258.1.

[0430]

F) methyl 5-bromo-2-hydroxy-3,4-dimethylbenzoate

To an aqueous solution (160 mL) of methyl 2-amino-5-bromo-3,4-dimethylbenzoate (28.0 g) in 25% sulfuric acid was added dropwise an aqueous solution (80.0 mL) of sodium nitrite (11.2 g) over 50 min at the internal temperature of 2-3°C under ice-cooling. Then, 5% aqueous sulfuric acid solution (1600 mL) was added dropwise thereto over 50 min at 0-15°C. The reaction mixture was stirred at 100°C for 2 hr. The reaction mixture was allowed to be cooled to room temperature, and the precipitate was collected by filtration, and washed with water to give the title compound (25.7 g).

MS: $[M-H]^+$ 256.9.

20 [0431]

G) methyl 2-hydroxy-3,4-dimethyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate

To a solution of methyl 5-bromo-2-hydroxy-3,4-dimethylbenzoate (25.7 g) in toluene (500 mL) were added bis(pinacolato)diboron (37.8 g), potassium acetate (29.2 g) and trans-dichlorobis(triphenylphosphine)palladium(II) (0.70 g), and the mixture was stirred under argon atmosphere at 100°C for 2 hr. To the reaction mixture was again added trans-dichlorobis(triphenylphosphine)palladium(II) (0.70 g), and the mixture was stirred overnight at 100°C. The reaction mixture was allowed to be cooled to room temperature, water was added thereto, and the mixture was extracted with ethyl acetate. The organic layer was washed with water and saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was collected

by filtration, and washed with ethyl acetate to give the title compound (18.3 g).

MS: $[M-H]^+$ 305.1.

[0432]

- 5 H) methyl 2-hydroxy-3,4-dimethyl-5-(4-(1-methyl-1H-pyrazol-3-yl)benzyl)benzoate

To a solution of methyl 2-hydroxy-3,4-dimethyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (0.97 g), 3-[4-(chloromethyl)phenyl]-1-methyl-1H-pyrazole (0.65 g) and
10 sodium carbonate (0.67 g) in a mixed solvent of 1,2-dimethoxyethane (12.0 mL)-water (4.00 mL) was added tetrakis(triphenylphosphine)palladium(0) (0.18 g) under argon atmosphere, and the mixture was stirred overnight at 80°C. To the reaction mixture was added water, and the mixture was
15 extracted with ethyl acetate. The organic layer was washed with saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (ethyl acetate/hexane) to give the title compound (0.65 g).
20 ^1H NMR (300MHz, CDCl_3) δ 2.14 (3H, s), 2.20 (3H, s), 3.91 (3H, s), 3.94 (3H, s), 3.97 (2H, s), 6.49 (1H, d, $J = 2.3$ Hz), 7.10 (2H, d, $J = 8.3$ Hz), 7.35 (1H, d, $J = 2.3$ Hz), 7.53 (1H, s), 7.65-7.72 (2H, m), 10.99 (1H, s).

[0433]

- 25 I) methyl 3,4-dimethyl-5-(4-(1-methyl-1H-pyrazol-3-yl)benzyl)-2-(((trifluoromethyl)sulfonyl)oxy)benzoate

To a solution of methyl 2-hydroxy-3,4-dimethyl-5-(4-(1-methyl-1H-pyrazol-3-yl)benzyl)benzoate (1.00 g) in DMF (12.0 mL) were added sodium hydride (0.14 g) and N-
30 phenylbis(trifluoromethanesulfonimide) (1.12 g) under ice-cooling, and the mixture was stirred for 2 hr. To the reaction mixture was added 1N hydrochloric acid, and the mixture was extracted with ethyl acetate. The organic layer was washed with saturated aqueous sodium bicarbonate solution and
35 saturated brine, and dried over anhydrous magnesium sulfate,

and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (ethyl acetate/hexane) to give the title compound (1.30 g).

¹H NMR (300MHz, CDCl₃) δ 2.21 (3H, s), 2.30 (3H, s), 3.91 (3H, s), 3.94 (3H, s), 4.05 (2H, s), 6.50 (1H, d, J = 2.3 Hz), 7.10 (1H, d, J = 8.1 Hz), 7.36 (1H, d, J = 2.3 Hz), 7.65-7.74 (1H, m).

[0434]

J) methyl 3,4-dimethyl-5-(4-(1-methyl-1H-pyrazol-3-yl)benzyl)-2-vinylbenzoate

A solution of methyl 3,4-dimethyl-5-(4-(1-methyl-1H-pyrazol-3-yl)benzyl)-2-(((trifluoromethyl)sulfonyl)oxy)benzoate (1.30 g), tributylvinyltin (1.28 g), trans-dichlorobis(triphenylphosphine)palladium(II) (0.10 g) and lithium chloride (0.80 g) in DMF (12.0 mL) was stirred at 90°C for 2 hr under argon atmosphere. To the reaction mixture was added aqueous potassium fluoride solution, and the precipitated insoluble substance was removed by filtration through Celite. The filtrate was diluted with ethyl acetate, and the mixture was washed with water and saturated brine. The organic layer was dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (ethyl acetate/hexane) to give the title compound (0.96 g).

¹H NMR (300MHz, CDCl₃) δ 2.17 (3H, s), 2.26 (3H, s), 3.81 (3H, s), 3.94 (3H, s), 4.04 (2H, s), 5.11 (1H, dd, J = 17.8, 1.9 Hz), 5.44 (1H, dd, J = 11.2, 1.8 Hz), 6.49 (1H, d, J = 2.3 Hz), 7.04 (1H, dd, J = 17.8, 11.3 Hz), 7.12 (2H, d, J = 8.5 Hz), 7.32-7.39 (2H, m), 7.44 (1H, s), 7.65-7.72 (1H, m).

[0435]

K) methyl 2-formyl-3,4-dimethyl-5-(4-(1-methyl-1H-pyrazol-3-yl)benzyl)benzoate

To a solution of methyl 3,4-dimethyl-5-(4-(1-methyl-1H-pyrazol-3-yl)benzyl)-2-vinylbenzoate (0.96 g) in a mixed solvent of acetone (10.0 mL)-acetonitrile (10.0 mL)-water (10.0

mL) were added osmium oxide (fixed catalyst I) (0.34 g) and sodium periodate (2.86 g), and the mixture was stirred overnight at room temperature. The reaction mixture was concentrated under reduced pressure, and the residue was
5 diluted with ethyl acetate and water. The insoluble substance was removed by filtration, and the filtrate was washed with saturated aqueous sodium bicarbonate solution, water and saturated brine, and dried over anhydrous sodium sulfate. The solvent was evaporated under reduced pressure, and the residue
10 was purified by silica gel column chromatography (ethyl acetate/hexane) to give the title compound (0.17 g).

MS: $[M+H]^+$ 363.2.

[0436]

L) 2-((1S,2S)-2-hydroxycyclopentyl)-4,5-dimethyl-6-(4-(1-methyl-1H-pyrazol-3-yl)benzyl)isoindolin-1-one
15

A solution of methyl 2-formyl-3,4-dimethyl-5-(4-(1-methyl-1H-pyrazol-3-yl)benzyl)benzoate (0.09 g), (1S,2S)-2-aminocyclopentanol hydrochloride (0.03 g), triethylamine (0.03 mL) and anhydrous magnesium sulfate (0.06 g) in THF (1.80 mL)
20 was stirred at room temperature for 46 hr under nitrogen atmosphere. The insoluble substance was removed by filtration, and the filtrate was concentrated under reduced pressure. The residue was diluted with methanol (1.80 mL) and THF (1.80 mL), sodium triacetoxyborohydride (0.10 g) was added thereto, and
25 the mixture was stirred at room temperature for 7.5 hr. The reaction mixture was diluted with ethyl acetate, and the mixture was washed with water and saturated brine. The organic layer was dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was
30 purified by silica gel column chromatography (ethyl acetate/hexane) to give the title compound (0.04 g).

^1H NMR (300 MHz, DMSO- d_6) δ 1.47-1.77 (4H, m), 1.81-1.97 (2H, m), 2.20 (3H, s), 2.23 (3H, s), 3.86 (3H, s), 4.07-4.30 (4H, m), 4.39 (2H, s), 4.89 (1H, d, J = 4.9 Hz), 6.61 (1H, d, J = 2.3
35 Hz), 7.12 (2H, d, J = 7.9 Hz), 7.33 (1H, s), 7.65-7.71 (3H, m).

[0437]

Example 14

4,5-dimethyl-6-(4-(1-methyl-1H-1,2,3-triazol-4-yl)benzyl)-2-
((2S)-tetrahydrofuran-2-ylmethyl)isoindolin-1-one

5 A) [4-(1-methyl-1H-1,2,3-triazol-4-yl)phenyl]methanol

A mixture of [4-(hydroxymethyl)phenyl]boronic acid (7.10 g), 4-bromo-1-methyl-1H-1,2,3-triazole (5.00 g), sodium carbonate (6.59 g) and tetrakis(triphenylphosphine)palladium(0) (3.59 g) in a mixed solvent of water (30.0 mL)-1,4-dioxane (100
10 mL) was heated with reflux for 16 hr under nitrogen atmosphere. The reaction mixture was allowed to be cooled to room temperature, the organic layer was concentrated under reduced pressure, and the residue was extracted with dichloromethane (x 3). The combined extracts were dried over anhydrous sodium
15 sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (ethyl acetate/petroleum ether) to give the title compound (3.40 g).

MS: [M+H]⁺ 189.9.

20 [0438]

B) 4-[4-(bromomethyl)phenyl]-1-methyl-1H-1,2,3-triazole

To a solution of [4-(1-methyl-1H-1,2,3-triazol-4-yl)phenyl]methanol (3.40 g) in dichloromethane (130 mL) was added dropwise phosphorus tribromide (23.0 g) under ice-cooling,
25 and the mixture was stirred at 20°C for 16 hr. The reaction mixture was concentrated under reduced pressure, the residue was poured into saturated aqueous sodium bicarbonate solution, and the mixture was extracted with dichloromethane (x 3). The combined organic layers were washed with water and saturated
30 brine, and dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (ethyl acetate/petroleum ether) to give the title compound (3.50 g) as a mixture with phosphine oxide. This mixture was diluted with
35 tert-butyl methyl ether, and the mixture was stirred at 20°C

for 16 hr. The precipitate was collected by filtration, and dried under reduced pressure to give the title compound (3.40 g).

MS: $[M+H]^+$ 251.8.

5 [0439]

C) methyl 2-hydroxy-3,4-dimethyl-5-(4-(1-methyl-1H-1,2,3-triazol-4-yl)benzyl)benzoate

To a mixture of methyl 2-hydroxy-3,4-dimethyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (0.76 g), 4-[4-(bromomethyl)phenyl]-1-methyl-1H-1,2,3-triazole (0.63 g) and
10 sodium carbonate (0.53 g) in 1,2-dimethoxyethane (12.0 mL)-water (4.00 mL) was added

tetrakis(triphenylphosphine)palladium(0) (0.29 g), and the mixture was stirred overnight at 80°C under argon atmosphere.

15 To the reaction mixture were added water and ethyl acetate, and the mixture was partitioned. The organic layer was washed with water and saturated brine, and dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography
20 (ethyl acetate/hexane) to give the title compound (0.60 g).

MS: $[M+H]^+$ 352.2.

[0440]

D) methyl 3,4-dimethyl-5-(4-(1-methyl-1H-1,2,3-triazol-4-yl)benzyl)-2-(((trifluoromethyl)sulfonyl)oxy)benzoate

25 To a solution of methyl 2-hydroxy-3,4-dimethyl-5-(4-(1-methyl-1H-1,2,3-triazol-4-yl)benzyl)benzoate (0.60 g) in DMF (12.0 mL) were added sodium hydride (0.08 g) and N-phenylbis(trifluoromethanesulfonimide) (0.67 g) under ice-cooling, and the mixture was stirred for 2 hr. To the reaction
30 mixture was added 1N hydrochloric acid, and the mixture was extracted with ethyl acetate. The organic layer was washed with saturated aqueous sodium bicarbonate solution and saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The
35 residue was purified by silica gel column chromatography (ethyl

acetate/hexane) to give the title compound (0.60 g).

MS: $[M+H]^+$ 484.1.

[0441]

E) methyl 3,4-dimethyl-5-(4-(1-methyl-1H-1,2,3-triazol-4-yl)benzyl)-2-vinylbenzoate

To a solution of methyl 3,4-dimethyl-5-(4-(1-methyl-1H-1,2,3-triazol-4-yl)benzyl)-2-(((trifluoromethyl)sulfonyl)oxy)benzoate (0.60 g) in DMF (12.0 mL) were added tributylvinyltin (1.28 g), trans-
10 dichlorobis(triphenylphosphine)palladium(II) (0.10 g) and lithium chloride (0.80 g), and the mixture was stirred at 90°C for 1.5 hr under argon atmosphere. To the reaction mixture was added aqueous potassium fluoride solution, and the precipitated insoluble substance was removed by filtration through Celite.
15 The filtrate was diluted with ethyl acetate, and the mixture was washed with water and saturated brine. The organic layer was dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (ethyl acetate/hexane) to give
20 the title compound (0.42 g).

MS: $[M+H]^+$ 362.2.

[0442]

F) methyl 2-formyl-3,4-dimethyl-5-(4-(1-methyl-1H-1,2,3-triazol-4-yl)benzyl)benzoate

To a solution of methyl 3,4-dimethyl-5-(4-(1-methyl-1H-1,2,3-triazol-4-yl)benzyl)-2-vinylbenzoate (0.42 g) in a mixed solvent of acetone (10.0 mL)-acetonitrile (10.0 mL)-water (10.0 mL) were added osmium oxide (fixed catalyst I) (0.15 g) and sodium periodate (1.24 g), and the mixture was stirred
30 overnight at room temperature. The reaction mixture was filtered, and the filtrate was extracted with ethyl acetate. The organic layer was washed with water and saturated brine, and dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure to give the title compound
35 (0.26 g) as a crude product.

MS: $[M+H]^+$ 364.2.

[0443]

G) 3-hydroxy-4,5-dimethyl-6-(4-(1-methyl-1H-1,2,3-triazol-4-yl)benzyl)-2-(((2S)-tetrahydrofuran-2-yl)methyl)isoindolin-1-one

To a solution of methyl 2-formyl-3,4-dimethyl-5-(4-(1-methyl-1H-1,2,3-triazol-4-yl)benzyl)benzoate (0.13 g) in THF (3.00 mL) were added 1-((2S)-tetrahydrofuran-2-yl)methanamine (0.04 mL) and anhydrous magnesium sulfate (0.04 g), and the mixture was stirred overnight at room temperature. The reaction mixture was diluted with ethyl acetate, and the mixture was washed with water and saturated brine. The organic layer was dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (ethyl acetate/hexane) to give the title compound (0.11 g).

MS: $[M+H]^+$ 433.3.

[0444]

H) 4,5-dimethyl-6-(4-(1-methyl-1H-1,2,3-triazol-4-yl)benzyl)-2-((2S)-tetrahydrofuran-2-ylmethyl)isoindolin-1-one

To a solution of 3-hydroxy-4,5-dimethyl-6-(4-(1-methyl-1H-1,2,3-triazol-4-yl)benzyl)-2-(((2S)-tetrahydrofuran-2-yl)methyl)isoindolin-1-one (0.11 g) in trifluoroacetic acid (1.50 mL) was added triethylsilane (0.08 mL), and the mixture was stirred at room temperature for 15 min. The reaction mixture was concentrated under reduced pressure, and the residue was purified by silica gel column chromatography (ethyl acetate/hexane) to give the title compound (0.08 g).

^1H NMR (300 MHz, DMSO- d_6) δ 1.49-1.62 (1H, m), 1.74-1.98 (3H, m), 2.21 (6H, s), 3.45-3.56 (1H, m), 3.57-3.68 (2H, m), 3.74-3.84 (1H, m), 4.01-4.14 (6H, m), 4.39-4.54 (2H, m), 7.19 (2H, d, $J = 7.9$ Hz), 7.34 (1H, s), 7.73 (2H, d, $J = 8.3$ Hz), 8.44 (1H, s).

[0445]

Example 15

2-(2-hydroxy-2-methylpropyl)-4,5-dimethyl-6-(4-(1H-pyrazol-1-yl)benzyl)isoindolin-1-one

A) methyl 5-(4-(1H-pyrazol-1-yl)benzyl)-2-hydroxy-3,4-dimethylbenzoate

5 To a solution of methyl 2-hydroxy-3,4-dimethyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (0.60 g) in DME (12.0 mL) were added 1-(4-(bromomethyl)phenyl)-1H-pyrazole (0.51 g), [1,1'-bis(diphenylphosphino)ferrocene]dichloropalladium(II) dichloromethane adduct (0.08 g) and 2mol/L aqueous sodium carbonate solution (1.96 mL), and the mixture was stirred overnight at 80°C under argon atmosphere. The reaction mixture was allowed to be cooled to room temperature, water and ethyl acetate were added thereto, and the mixture was partitioned.

10 The organic layer was washed with water and saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (ethyl acetate/hexane) to give the title compound (0.54 g).

20 MS: [M+H]⁺ 337.2.

[0446]

B) methyl 5-(4-(1H-pyrazol-1-yl)benzyl)-3,4-dimethyl-2-vinylbenzoate

To a solution of methyl 5-(4-(1H-pyrazol-1-yl)benzyl)-2-hydroxy-3,4-dimethylbenzoate (0.54 g) in DMF (10 mL) was added sodium hydride (0.08 g) under ice-cooling, and the mixture was stirred for 30 min. To this reaction mixture was added N-phenylbis(trifluoromethanesulfonimide) (0.63 g), and the mixture was stirred at room temperature for 1 hr. To the reaction mixture was added 1N hydrochloric acid, and the mixture was extracted with ethyl acetate. The organic layer was washed with water and saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The obtained crude methyl 3,4-dimethyl-5-[4-(1H-pyrazol-1-yl)benzyl]-2-[[trifluoromethyl)sulfonyl]oxy]benzoate

30

35

was used in the next step without an additional purification.
MS: $[M+H]^+$ 469.1.

The above-mentioned compound was dissolved in DMF (10 mL), tributylvinyltin (0.71 mL), trans-
5 dichlorobis(triphenylphosphine)palladium(II) (0.06 g) and lithium chloride (0.50 g) were added thereto, and the mixture was stirred overnight at 90°C under argon atmosphere. To the reaction mixture was again added trans-
dichlorobis(triphenylphosphine)palladium(II) (0.11 g), and the
10 mixture was stirred at 90°C for 2 hr under argon atmosphere. To the reaction mixture was added aqueous potassium fluoride solution, and the precipitated insoluble substance was removed by filtration through Celite. The filtrate was diluted with ethyl acetate, and the mixture was washed with water and
15 saturated brine. The organic layer was dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (ethyl acetate/hexane) to give the title compound (0.19 g).

20 MS: $[M+H]^+$ 347.2.

[0447]

C) methyl 5-(4-(1H-pyrazol-1-yl)benzyl)-2-formyl-3,4-dimethylbenzoate

To a solution of methyl 5-(4-(1H-pyrazol-1-yl)benzyl)-
25 3,4-dimethyl-2-vinylbenzoate (0.40 g) in a mixed solvent of acetone (8.00 mL)-acetonitrile (8.00 mL)-water (8.00 mL) were added osmium oxide (fixed catalyst I) (0.15 g) and sodium periodate (1.24 g), and the mixture was stirred overnight at room temperature. The reaction mixture was filtered, and the
30 filtrate was concentrated under reduced pressure. To the residue was added water, and the mixture was extracted with ethyl acetate. The organic layer was washed with water and saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The
35 residue was purified by silica gel column chromatography (ethyl

acetate/hexane) to give the title compound (0.11 g).

MS: $[M+H]^+$ 349.1.

[0448]

D) 2-(2-hydroxy-2-methylpropyl)-4,5-dimethyl-6-(4-(1H-pyrazol-1-yl)benzyl)isoindolin-1-one

To a solution of methyl 5-(4-(1H-pyrazol-1-yl)benzyl)-2-formyl-3,4-dimethylbenzoate (0.11 g) in THF (4.00 mL) was added 1-amino-2-methylpropan-2-ol (0.03 g), and the mixture was stirred at room temperature for 2 hr. The reaction mixture was concentrated, and the residue was diluted with methanol (4.0 mL). Sodium triacetoxyborohydride (0.11 g) was added thereto under argon atmosphere, and the mixture was stirred overnight at room temperature. To the reaction mixture was added water, and the mixture was extracted with ethyl acetate. The organic layer was washed with water and saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by NH silica gel column chromatography (ethyl acetate/hexane) to give the title compound (0.04 g).

^1H NMR (300 MHz, CDCl_3) δ 1.31 (6H, s), 2.23 (3H, s), 2.26 (3H, s), 3.27 (1H, s), 3.63 (2H, s), 4.13 (2H, s), 4.51 (2H, s), 6.43-6.47 (1H, m), 7.19 (2H, d, $J = 8.5$ Hz), 7.56-7.62 (3H, m), 7.71 (1H, d, $J = 1.5$ Hz), 7.88 (1H, d, $J = 2.5$ Hz).

[0449]

Example 16

6-(2-fluoro-4-(1H-pyrazol-1-yl)benzyl)-2-((1S,2S)-2-hydroxycyclopentyl)-4,5-dimethylisoindolin-1-one

A) methyl 2-fluoro-4-(1H-pyrazol-1-yl)benzoate

To a solution of (3-fluoro-4-(methoxycarbonyl)phenyl)boronic acid (2.40 g) and 1H-pyrazole (0.99 g) in methanol (54.0 mL) was added copper(I) oxide (0.10 g), and the mixture was stirred overnight at 50°C. The reaction mixture was concentrated under reduced pressure, and the residue was purified by silica gel column chromatography (ethyl acetate/hexane) to give the title compound (0.89 g).

MS: $[M+H]^+$ 221.1.

[0450]

B) (2-fluoro-4-(1H-pyrazol-1-yl)phenyl)methanol

To a solution of lithium aluminium hydride (0.16 g) in
5 THF (5.50 mL) was added a solution of methyl 2-fluoro-4-(1H-
pyrazol-1-yl)benzoate (0.89 g) in THF (5.50 mL) under ice-
cooling, and the mixture was stirred for 1 hr. To the reaction
solution was added water, the insoluble substance was removed
by filtration, and the filtrate was concentrated under reduced
10 pressure to give the title compound (0.71 g) as a crude product.

MS: $[M+H]^+$ 193.1.

[0451]

C) 1-(4-(chloromethyl)-3-fluorophenyl)-1H-pyrazole

To a solution of (2-fluoro-4-(1H-pyrazol-1-
15 yl)phenyl)methanol (0.71 g) in THF (15.0 mL) was added thionyl
chloride (0.40 mL) under ice-cooling, and the mixture was
stirred at room temperature for 16 hr. The reaction mixture
was diluted with ethyl acetate, saturated aqueous sodium
bicarbonate was added thereto, and the mixture was partitioned.
20 The organic layer was washed with saturated brine, and dried
over anhydrous magnesium sulfate, and the solvent was
evaporated under reduced pressure. The residue was purified by
silica gel column chromatography (ethyl acetate/hexane) to give
the title compound (0.50 g).

25 MS: $[M+H]^+$ 211.1.

[0452]

D) methyl 5-(2-fluoro-4-(1H-pyrazol-1-yl)benzyl)-2-hydroxy-3,4-
dimethylbenzoate

To a mixture of methyl 2-hydroxy-3,4-dimethyl-5-(4,4,5,5-
30 tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (0.51 g), 1-(4-
(chloromethyl)-3-fluorophenyl)-1H-pyrazole (0.35 g) and sodium
carbonate (0.35 g) in 1,2-dimethoxyethane (7.80 mL)-water (2.60
mL) was added tetrakis(triphenylphosphine)palladium(0) (0.10 g),
and the mixture was stirred overnight at 80°C under argon
35 atmosphere. To the reaction mixture were added water and ethyl

acetate, and the mixture was partitioned. The organic layer was washed with water and saturated brine, and dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (ethyl acetate/hexane) to give the title compound (0.54 g).

MS: $[M+H]^+$ 355.2.

[0453]

E) methyl 5-(2-fluoro-4-(1H-pyrazol-1-yl)benzyl)-3,4-dimethyl-2-(((trifluoromethyl)sulfonyl)oxy)benzoate

To a solution of methyl 5-(2-fluoro-4-(1H-pyrazol-1-yl)benzyl)-2-hydroxy-3,4-dimethylbenzoate (0.54 g) in DMF (11.0 mL) were added sodium hydride (0.07 g) and N-phenylbis(trifluoromethanesulfonimide) (0.60 g) under ice-cooling, and the mixture was stirred for 2 hr. To the reaction mixture was added 1N hydrochloric acid, and the mixture was extracted with ethyl acetate. The organic layer was washed with saturated aqueous sodium bicarbonate solution and saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (ethyl acetate/hexane) to give the title compound (0.50 g).

MS: $[M+H]^+$ 487.1.

[0454]

F) methyl 5-(2-fluoro-4-(1H-pyrazol-1-yl)benzyl)-3,4-dimethyl-2-vinylbenzoate

To a solution of methyl 5-(2-fluoro-4-(1H-pyrazol-1-yl)benzyl)-3,4-dimethyl-2-(((trifluoromethyl)sulfonyl)oxy)benzoate (0.50 g) in DMF (10.0 mL) were added tributylvinyltin (0.50 g), trans-dichlorobis(triphenylphosphine)palladium(II) (0.04 g) and lithium chloride (0.32 g), and the mixture was stirred at 90°C for 1.5 hr under argon atmosphere. To the reaction mixture was added aqueous potassium fluoride solution, and the precipitated insoluble substance was removed by filtration through Celite.

The filtrate was diluted with ethyl acetate, and the mixture was washed with water and saturated brine. The organic layer was dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (ethyl acetate/hexane) to give the title compound (0.40 g).

MS: $[M+H]^+$ 365.2.

[0455]

G) methyl 5-(2-fluoro-4-(1H-pyrazol-1-yl)benzyl)-2-formyl-3,4-dimethylbenzoate

To a solution of methyl 5-(2-fluoro-4-(1H-pyrazol-1-yl)benzyl)-3,4-dimethyl-2-vinylbenzoate (0.40 g) in a mixed solvent of acetone (5.00 mL)-acetonitrile (5.00 mL)-water (5.00 mL) were added osmium oxide (fixed catalyst I) (0.14 g) and sodium periodate (1.17 g), and the mixture was stirred overnight at room temperature. The reaction mixture was filtered, and the filtrate was extracted with ethyl acetate. The organic layer was washed with water and saturated brine, and dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure to give the title compound (0.40 g) as a crude product. This compound was used in the next step without an additional purification.

[0456]

H) 6-(2-fluoro-4-(1H-pyrazol-1-yl)benzyl)-2-((1S,2S)-2-hydroxycyclopentyl)-4,5-dimethylisoindolin-1-one

A solution of methyl 5-(2-fluoro-4-(1H-pyrazol-1-yl)benzyl)-2-formyl-3,4-dimethylbenzoate (0.17 g), (1S,2S)-2-aminocyclopentanol hydrochloride (0.06 g), triethylamine (0.06 mL) and anhydrous magnesium sulfate (0.10 g) in THF (3.40 mL) was stirred at room temperature for 10 hr. The insoluble substance was removed by filtration, and the filtrate was concentrated under reduced pressure. The residue was diluted with methanol (3.40 mL) and THF (3.40 mL), sodium triacetoxyborohydride (0.19 g) was added thereto, and the mixture was stirred overnight at room temperature. The

reaction mixture was diluted with ethyl acetate, and the mixture was washed with water and saturated brine. The organic layer was dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was
5 purified by silica gel column chromatography (ethyl acetate/hexane) to give the title compound (0.07 g).

¹H NMR (300 MHz, DMSO-d₆) δ 1.45-1.60 (1H, m), 1.62-1.79 (3H, m), 1.80-1.98 (2H, m), 2.24 (3H, s), 2.25 (3H, s), 4.08-4.29 (4H, m), 4.39 (2H, s), 4.88 (1H, d, J = 4.9 Hz), 6.52-6.58 (1H,
10 m), 7.15 (1H, t, J = 8.4 Hz), 7.23 (1H, s), 7.63 (1H, dd, J = 8.3, 2.1 Hz), 7.68-7.77 (2H, m), 8.52 (1H, d, J = 2.5 Hz).

[0457]

Example 17

1,5-anhydro-2-(6-(4-chlorobenzyl)-4,5-dimethyl-1-oxo-1,3-
15 dihydro-2H-isoindol-2-yl)-2,4-dideoxy-L-threo-pentitol

A) methyl 5-(4-chlorobenzyl)-2-hydroxy-3,4-dimethylbenzoate

To a solution of methyl 2-hydroxy-3,4-dimethyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (0.70 g) in DME (14.0 mL) were added 1-(bromomethyl)-4-chlorobenzene
20 (0.49 g), [1,1'-

bis(diphenylphosphino)ferrocene]dichloropalladium(II) dichloromethane adduct (0.09 g) and 2mol/L aqueous sodium carbonate solution (2.29 mL), and the mixture was stirred overnight at 80°C under argon atmosphere. The reaction mixture
25 was allowed to be cooled to room temperature, water and ethyl acetate were added thereto, and the mixture was partitioned. The organic layer was washed with water and saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by
30 silica gel column chromatography (ethyl acetate/hexane) to give the title compound (0.50 g).

MS: [M+H]⁺ 305.1.

[0458]

B) methyl 5-(4-chlorobenzyl)-3,4-dimethyl-2-vinylbenzoate

35 To a solution of methyl 5-(4-chlorobenzyl)-2-hydroxy-3,4-

dimethylbenzoate (0.50 g) in DMF (10.0 mL) was added sodium hydride (0.08 g) under ice-cooling, and the mixture was stirred at room temperature for 30 min. To this reaction mixture was added N-phenylbis(trifluoromethanesulfonimide) (0.65 g) under
5 ice-cooling, and the mixture was stirred at room temperature for 1 hr. To the reaction mixture was added 1N hydrochloric acid, and the mixture was extracted with ethyl acetate. The organic layer was washed with water and saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was
10 evaporated under reduced pressure. The obtained crude methyl 5-(4-chlorobenzyl)-3,4-dimethyl-2-
{[(trifluoromethyl)sulfonyl]oxy}benzoate was used in the next step without an additional purification.

The above-mentioned compound was dissolved in DMF (10.0
15 mL), to the solution were added tributylvinyltin (0.72 mL), trans-dichlorobis(triphenylphosphine)palladium(II) (0.23 g) and lithium chloride (0.52 g), and the mixture was stirred overnight at 90°C under argon atmosphere. To the reaction mixture was added aqueous potassium fluoride solution, and the
20 precipitated insoluble substance was removed by filtration through Celite. The filtrate was diluted with ethyl acetate, and the mixture was washed with water and saturated brine. The organic layer was dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue
25 was purified by silica gel column chromatography (ethyl acetate/hexane) to give the title compound (0.17 g).

MS: $[M+H]^+$ 315.2.

[0459]

C) methyl 5-(4-chlorobenzyl)-2-formyl-3,4-dimethylbenzoate

30 To a solution of methyl 5-(4-chlorobenzyl)-3,4-dimethyl-2-vinylbenzoate (0.17 g) in a mixed solvent of acetone (4.00 mL)-acetonitrile (4.00 mL)-water (4.00 mL) were added osmium oxide (fixed catalyst I) (0.07 g) and sodium periodate (0.58 g), and the mixture was stirred overnight at room temperature under
35 argon atmosphere. The reaction mixture was filtered, the

filtrate was concentrated under reduced pressure, and the residue was extracted with ethyl acetate. The organic layer was washed with water and saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (ethyl acetate/hexane) to give the title compound (0.02 g).

MS: $[M+H]^+$ 317.1.

[0460]

10 D) 1,5-anhydro-2-(6-(4-chlorobenzyl)-4,5-dimethyl-1-oxo-1,3-dihydro-2H-isindol-2-yl)-2,4-dideoxy-L-threo-pentitol

To a solution of methyl 5-(4-chlorobenzyl)-2-formyl-3,4-dimethylbenzoate (0.02 g) in THF (1.00 mL) was added (3S,4S)-3-aminotetrahydro-2H-pyran-4-ol (8.14 mg), and the mixture was stirred at room temperature for 2 hr. The reaction mixture was concentrated, and the residue was diluted with methanol (1.00 mL). Sodium triacetoxyborohydride (0.02 g) was added thereto under argon atmosphere, and the mixture was stirred overnight at room temperature. The reaction mixture was diluted with ethyl acetate, and the mixture was washed with water and saturated brine. The organic layer was dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by reverse-phase HPLC. The fractions were combined, saturated aqueous sodium hydrogencarbonate solution was added thereto, and the mixture was extracted with ethyl acetate. The organic layer was dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure to give the title compound (5.60 mg).

30 ^1H NMR (300 MHz, CDCl_3) δ 1.72-1.88 (1H, m), 2.09-2.17 (1H, m), 2.20 (3H, s), 2.25 (3H, s), 3.45-3.62 (2H, m), 4.01-4.20 (6H, m), 4.26-4.50 (2H, m), 7.02 (2H, d, $J = 8.5$ Hz), 7.19-7.25 (2H, m), 7.51 (1H, s). 1H undetected.

[0461]

35 Example 18-1

2-[(3S,4S)-4-hydroxytetrahydro-2H-pyran-3-yl]-6-(4-methoxybenzyl)-4,5-dimethyl-2,3-dihydro-1H-isoindol-1-one
Alias; 1,5-anhydro-2,4-dideoxy-2-(6-(4-methoxybenzyl)-4,5-dimethyl-1-oxo-1,3-dihydro-2H-isoindol-2-yl)-L-threo-pentitol

5 A) methyl 2-hydroxy-5-(4-methoxybenzyl)-3,4-dimethylbenzoate

To a mixture of methyl 2-hydroxy-3,4-dimethyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (18.3 g), 1-(chloromethyl)-4-methoxybenzene (9.38 g) and sodium carbonate (12.7 g) in 1,2-dimethoxyethane (255 mL)-water (85.0 mL) was
10 added tetrakis(triphenylphosphine)palladium(0) (3.46 g), and the mixture was stirred overnight at 80°C under argon atmosphere. To the reaction mixture were added water and ethyl acetate, and the organic layer was separated, washed with saturated brine. The organic layer was dried over anhydrous
15 sodium sulfate, and the solvent was evaporated under reduced pressure. To the residue was added a mixed solvent of ethyl acetate-diisopropyl ether, and the solid was collected by filtration to give the title compound (11.12 g). The filtrate was concentrated, and the residue was purified by silica gel
20 column chromatography (ethyl acetate/hexane) to give the title compound (4.99 g).

MS: [M+H]⁺ 301.1.

[0462]

B) methyl 5-(4-methoxybenzyl)-3,4-dimethyl-2-
25 (((trifluoromethyl)sulfonyl)oxy)benzoate

To a solution of methyl 2-hydroxy-5-(4-methoxybenzyl)-3,4-dimethylbenzoate (16.1 g) in DMF (300 mL) were added sodium hydride (2.57 g) and N-phenylbis(trifluoromethanesulfonimide) (21.1 g) under ice-
30 cooling, and the mixture was stirred at room temperature for 2.5 hr. To the reaction mixture was added 1N hydrochloric acid, and the mixture was extracted with ethyl acetate. The organic layer was washed with saturated aqueous sodium bicarbonate solution and saturated brine, and dried over anhydrous
35 magnesium sulfate, and the solvent was evaporated under reduced

pressure. The residue was purified by silica gel column chromatography (ethyl acetate/hexane) to give the title compound (26.1 g).

¹H NMR (300 MHz, CDCl₃) δ 2.22 (3H, s), 2.31 (3H, s), 3.79 (3H, s), 3.92 (3H, s), 3.99 (2H, s), 6.83 (2H, d, J = 8.7 Hz), 7.01 (2H, d, J = 8.7 Hz), 7.64 (1H, s).

[0463]

C) methyl 5-(4-methoxybenzyl)-3,4-dimethyl-2-vinylbenzoate

To a solution of methyl 5-(4-methoxybenzyl)-3,4-dimethyl-2-(((trifluoromethyl)sulfonyl)oxy)benzoate (23.2 g) in DMF (360 mL) were added tributylvinyltin (25.5 g), trans-dichlorobis(triphenylphosphine)palladium(II) (1.88 g) and lithium chloride (16.8 g), and the mixture was stirred at 90°C for 1.5 hr under argon atmosphere. To the reaction mixture was added aqueous potassium fluoride solution, and the precipitated insoluble substance was removed by filtration through Celite. The filtrate was diluted with ethyl acetate, and the mixture was washed with water and saturated brine. The organic layer was dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (ethyl acetate/hexane) to give the title compound (13.6 g).

MS: [M+H]⁺ 311.2.

[0464]

D) methyl 2-formyl-5-(4-methoxybenzyl)-3,4-dimethylbenzoate

To a solution of methyl 5-(4-methoxybenzyl)-3,4-dimethyl-2-vinylbenzoate (13.6 g) in a mixed solvent of acetone (135 mL)-acetonitrile (135 mL)-water (135 mL) were added osmium oxide (fixed catalyst I) (5.57 g) and sodium periodate (46.9 g), and the mixture was stirred overnight at room temperature. The reaction mixture was filtered, and the filtrate was extracted with ethyl acetate. The organic layer was washed with water and saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure to give the title compound (13.5 g) as a crude product. This compound

was used in the next step without an additional purification.

MS: $[M+H]^+$ 313.2.

[0465]

E) 2-[(3S,4S)-4-hydroxytetrahydro-2H-pyran-3-yl]-6-(4-methoxybenzyl)-4,5-dimethyl-2,3-dihydro-1H-isoindol-1-one
5 Alias; 1,5-anhydro-2,4-dideoxy-2-(6-(4-methoxybenzyl)-4,5-dimethyl-1-oxo-1,3-dihydro-2H-isoindol-2-yl)-L-threo-pentitol

To a solution of methyl 2-formyl-5-(4-methoxybenzyl)-3,4-dimethylbenzoate (13.5 g) in THF (270 mL) were added (3S,4S)-3-aminotetrahydro-2H-pyran-4-ol (5.06 g) and anhydrous magnesium sulfate (9.99 g), and the mixture was stirred at room temperature for 5 hr. The insoluble substance was removed by filtration, and the filtrate was concentrated. The residue was diluted with methanol (220 mL)-THF (250 mL), sodium triacetoxymethylborohydride (18.3 g) was added thereto, and the mixture was stirred at room temperature for 15 hr. The reaction mixture was diluted with ethyl acetate, the mixture was washed with water and saturated brine, and the organic layer was dried over anhydrous magnesium sulfate. The organic layer was concentrated under reduced pressure. The resulting solid was washed with ethyl acetate to give the crude title compound (6.4 g). The filtrate was concentrated under reduced pressure, and the residue was purified by silica gel column chromatography (ethyl acetate/hexane) to give the title compound (0.85 g). The crude title compound and the title compound purified by column were combined, and recrystallized from ethanol to give the title compound (6.48 g).

^1H NMR (300 MHz, DMSO- d_6) δ 1.44-1.65 (1H, m), 1.95 (1H, d, J = 11.3 Hz), 2.19 (3H, s), 2.21 (3H, s), 3.33-3.49 (2H, m), 3.71 (4H, s), 3.82-3.96 (3H, m), 4.00 (2H, s), 4.33-4.50 (2H, m), 5.05 (1H, d, J = 4.9 Hz), 6.84 (2H, d, J = 8.7 Hz), 7.03 (2H, d, J = 8.7 Hz), 7.28 (1H, s).

X-ray powder diffraction pattern with specific peaks at d value (or d -spacing) = 18.3, 9.8, 9.2, 6.8, 6.1, 5.2, 4.6, 4.2 and 3.8 Å.

[0466]

Example 18-2

2-[(3S,4S)-4-hydroxytetrahydro-2H-pyran-3-yl]-6-(4-methoxybenzyl)-4,5-dimethyl-2,3-dihydro-1H-isoindol-1-one

5 Alias; 1,5-anhydro-2,4-dideoxy-2-(6-(4-methoxybenzyl)-4,5-dimethyl-1-oxo-1,3-dihydro-2H-isoindol-2-yl)-L-threo-pentitol

This compound was also synthesized by the following method.

[0467]

10 A) methyl 2-hydroxy-5-(4-methoxybenzyl)-3,4-dimethylbenzoate

To a mixture of methyl 2-hydroxy-3,4-dimethyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (0.40 g), 1-(chloromethyl)-4-methoxybenzene (0.21 g) and sodium carbonate (0.28 g) in 1,2-dimethoxyethane (6.00 mL)-water (2.00 mL) was
15 added tetrakis(triphenylphosphine)palladium(0) (0.08 g), and the mixture was stirred overnight at 80°C under argon atmosphere. To the reaction mixture was added water, and the mixture was extracted with ethyl acetate. The organic layer was separated, washed with saturated brine, and dried over
20 anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (ethyl acetate/hexane) to give the title compound (0.30 g).

¹H NMR (300 MHz, CDCl₃) δ 2.13 (3H, s), 2.19 (3H, s), 3.77 (3H, s), 3.90 (2H, s), 3.91 (3H, s), 6.80 (2H, d, J = 8.7 Hz), 6.99 (2H, d, J = 8.7 Hz), 7.50 (1H, s), 10.97 (1H, s).

[0468]

B) methyl 5-(4-methoxybenzyl)-3,4-dimethyl-2-(((trifluoromethyl)sulfonyl)oxy)benzoate

30 To a mixture of methyl 2-hydroxy-5-(4-methoxybenzyl)-3,4-dimethylbenzoate (0.30 g), sodium hydride (0.05 g) and DMF (6.00 mL) was added N-phenylbis(trifluoromethanesulfonimide) (0.39 g) under ice-cooling, and the mixture was stirred at room temperature for 2
35 hr. To the reaction mixture was added 1N hydrochloric acid,

and the mixture was extracted with ethyl acetate. The organic layer was washed with saturated aqueous sodium bicarbonate solution and saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced
5 pressure. The residue was purified by silica gel column chromatography (ethyl acetate/hexane) to give the title compound (0.34 g).

^1H NMR (300 MHz, CDCl_3) δ 2.21 (3H, s), 2.30 (3H, s), 3.78 (3H, s), 3.91 (3H, s), 3.98 (2H, s), 6.78-6.87 (2H, m), 7.00 (2H, d,
10 $J = 8.7$ Hz), 7.63 (1H, s).

[0469]

C) methyl 5-(4-methoxybenzyl)-3,4-dimethyl-2-vinylbenzoate

To a solution of methyl 5-(4-methoxybenzyl)-3,4-dimethyl-2-(((trifluoromethyl)sulfonyl)oxy)benzoate (0.34 g) in DMF
15 (7.00 mL) were added tributylvinyltin (0.37 g), trans-dichlorobis(triphenylphosphine)palladium(II) (0.03 g) and lithium chloride (0.25 g), and the mixture was stirred at 90°C for 2 hr under argon atmosphere. To the reaction mixture was added aqueous potassium fluoride solution, and the precipitated
20 insoluble substance was removed by filtration through Celite. The filtrate was diluted with ethyl acetate, and the mixture was washed with saturated brine. The organic layer was dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by
25 silica gel column chromatography (ethyl acetate/hexane) to give the title compound (0.20 g).

^1H NMR (300 MHz, CDCl_3) δ 2.17 (3H, s), 2.26 (3H, s), 3.77 (3H, s), 3.80 (3H, s), 3.97 (2H, s), 5.10 (1H, dd, $J = 17.7, 1.9$ Hz), 5.43 (1H, dd, $J = 11.3, 1.9$ Hz), 6.76-6.86 (2H, m), 6.96-7.09
30 (3H, m), 7.41 (1H, s).

[0470]

D) methyl 2-formyl-5-(4-methoxybenzyl)-3,4-dimethylbenzoate

To a solution of methyl 5-(4-methoxybenzyl)-3,4-dimethyl-2-vinylbenzoate (0.19 g) in a mixed solvent of acetone (2.30
35 mL)-acetonitrile (2.30 mL)-water (2.30 mL) were added osmium

oxide (fixed catalyst I) (0.08 g) and sodium periodate (0.67 g), and the mixture was stirred overnight at room temperature. The reaction mixture was filtered, and the filtrate was extracted with ethyl acetate. The organic layer was washed with water
5 and saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure to give the title compound (0.19 g) as a crude product. This compound was used in the next step without an additional purification.
MS: $[M+H]^+$ 313.2.

10 [0471]

E) 2-[(3S,4S)-4-hydroxytetrahydro-2H-pyran-3-yl]-6-(4-methoxybenzyl)-4,5-dimethyl-2,3-dihydro-1H-isoindol-1-one
Alias; 1,5-anhydro-2,4-dideoxy-2-(6-(4-methoxybenzyl)-4,5-dimethyl-1-oxo-1,3-dihydro-2H-isoindol-2-yl)-L-threo-pentitol

15 To a solution of methyl 2-formyl-5-(4-methoxybenzyl)-3,4-dimethylbenzoate (0.10 g) in THF (1.90 mL) were added (3S,4S)-3-aminotetrahydro-2H-pyran-4-ol (0.04 g) and anhydrous magnesium sulfate (0.07 g), and the mixture was stirred at room temperature for 5 hr. The insoluble substance was removed by
20 filtration, and the filtrate was concentrated. The residue was diluted with methanol (1.90 mL)-THF (1.90 mL), sodium triacetoxyborohydride (0.13 g) was added thereto, and the mixture was stirred at room temperature for 2.5 days. The reaction mixture was diluted with ethyl acetate, and the
25 mixture was washed with water and saturated brine. The organic layer was concentrated under reduced pressure, and the residue was purified by silica gel column chromatography (ethyl acetate/hexane) to give the title compound (0.04 g).

^1H NMR (300 MHz, DMSO- d_6) δ 1.48-1.64 (1H, m), 1.94 (1H, d, J = 14.7 Hz), 2.19 (3H, s), 2.21 (3H, s), 3.35-3.44 (2H, m), 3.65-3.74 (4H, m), 3.82-3.95 (3H, m), 4.00 (2H, s), 4.34-4.49 (2H, m), 5.04 (1H, d, J = 5.3 Hz), 6.84 (2H, d, J = 8.7 Hz), 7.03 (2H, d, J = 8.3 Hz), 7.28 (1H, s).

[0472]

35 Example 19

1,5-anhydro-2,4-dideoxy-2-(6-(3-fluoro-4-(methylcarbamoyl)benzyl)-4,5-dimethyl-1-oxo-1,3-dihydro-2H-isoindol-2-yl)-L-threo-pentitol

A) 2-fluoro-N-methyl-4-vinylbenzamide

5 A mixture of 4-bromo-2-fluoro-N-methylbenzamide (0.80 g), vinylboronic acid pinacol cyclic ester (0.80 g) and 2M aqueous sodium carbonate solution (3.45 mL) in DME (17.3 mL) was argon-purged. Bis(triphenylphosphine)palladium(II) dichloride (0.12 g) was added thereto, and the mixture was stirred overnight at
10 80°C. The reaction solution was diluted with ethyl acetate, and the mixture was washed with water and saturated brine. The organic layer was dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was subjected to short silica gel column chromatography, and
15 the solvent was evaporated under reduced pressure to give the title compound (0.74 g).

MS: $[M+H]^+$ 180.1.

[0473]

B) 2-fluoro-4-formyl-N-methylbenzamide

20 To a solution of 2-fluoro-N-methyl-4-vinylbenzamide (0.62 g) in a mixed solvent of acetone (23 mL)-acetonitrile (23 mL)-water (23 mL) were added osmium oxide (fixed catalyst I) (0.44 g) and sodium periodate (3.69 g), and the mixture was stirred overnight at room temperature. The insoluble substance was
25 removed by filtration, and the filtrate was diluted with ethyl acetate and water. The insoluble substance was removed by filtration, the filtrate was concentrated under reduced pressure, and the residue was diluted with ethyl acetate. The mixture was washed with water and saturated brine, the organic
30 layer was dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was subjected to short silica gel column chromatography, and the solvent was evaporated under reduced pressure to give the title compound (0.62 g).

35 MS: $[M+H]^+$ 182.1.

[0474]

C) 2-fluoro-4-(hydroxymethyl)-N-methylbenzamide

To a solution of 2-fluoro-4-formyl-N-methylbenzamide (0.62 g) in methanol (17 mL) was added sodium borohydride (0.16 g) in small portions, and the mixture was stirred at room temperature for 3 days. To the reaction mixture was added aqueous ammonium chloride, and the mixture was extracted with ethyl acetate. The organic layer was washed with water and saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (ethyl acetate/hexane) to give the title compound (0.44 g).

MS: $[M+H]^+$ 184.1.

[0475]

D) 4-(chloromethyl)-2-fluoro-N-methylbenzamide

To a solution of 2-fluoro-4-(hydroxymethyl)-N-methylbenzamide (2.11 g) in THF (46.1 mL) was added dropwise thionyl chloride (1.01 mL) under ice-cooling, and the mixture was stirred overnight at room temperature. To the reaction solution was added saturated aqueous sodium bicarbonate, and the mixture was extracted with ethyl acetate. The organic layer was washed with saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was subjected to silica gel column chromatography (ethyl acetate/hexane). The objective fractions were collected, and washed with diisopropyl ether, and the precipitate was collected by filtration. The filtrate was concentrated, and the residue was washed with diisopropyl ether. This procedure was repeated three times to give the title compound (0.96 g).

MS: $[M+H]^+$ 202.1.

[0476]

E) methyl 5-(3-fluoro-4-(methylcarbamoyl)benzyl)-2-hydroxy-3,4-dimethylbenzoate

To a mixture of methyl 2-hydroxy-3,4-dimethyl-5-(4,4,5,5-

tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (0.35 g), 4-(chloromethyl)-2-fluoro-N-methylbenzamide (0.23 g) and sodium carbonate (0.24 g) in 1,2-dimethoxyethane (5.40 mL)-water (1.80 mL) was added tetrakis(triphenylphosphine)palladium(0) (0.07 g),
5 and the mixture was stirred overnight at 80°C under argon atmosphere. To the reaction mixture were added water and ethyl acetate, and the mixture was partitioned. The organic layer was washed with water and saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated
10 under reduced pressure. The residue was purified by silica gel column chromatography (ethyl acetate/hexane) to give the title compound (0.37 g).

MS: $[M+H]^+$ 346.2.

[0477]

15 F) methyl 5-(3-fluoro-4-(methylcarbamoyl)benzyl)-3,4-dimethyl-2-(((trifluoromethyl)sulfonyl)oxy)benzoate

To a solution of methyl 5-(3-fluoro-4-(methylcarbamoyl)benzyl)-2-hydroxy-3,4-dimethylbenzoate (0.36 g) in DMF (7.00 mL) were added sodium hydride (0.05 g) and N-phenylbis(trifluoromethanesulfonimide) (0.41 g) under ice-cooling, and the mixture was stirred for 2 hr. To the reaction mixture was added 1N hydrochloric acid, and the mixture was extracted with ethyl acetate. The organic layer was washed with saturated aqueous sodium bicarbonate solution and
20 saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (ethyl acetate/hexane) to give the title compound (0.29 g).

MS: $[M+H]^+$ 478.0.

30 [0478]

G) methyl 5-(3-fluoro-4-(methylcarbamoyl)benzyl)-3,4-dimethyl-2-vinylbenzoate

To a solution of methyl 5-(3-fluoro-4-(methylcarbamoyl)benzyl)-3,4-dimethyl-2-(((trifluoromethyl)sulfonyl)oxy)benzoate (0.28 g) in DMF (6.00
35

mL) were added tributylvinyltin (0.29 g), trans-dichlorobis(triphenylphosphine)palladium(II) (0.02 g) and lithium chloride (0.19 g), and the mixture was stirred at 90°C for 2 hr under argon atmosphere. To the reaction mixture was
5 added aqueous potassium fluoride solution, and the precipitated insoluble substance was removed by filtration through Celite. The filtrate was diluted with ethyl acetate, and the mixture was washed with water and saturated brine. The organic layer was dried over anhydrous magnesium sulfate, and the solvent was
10 evaporated under reduced pressure. The residue was purified by silica gel column chromatography (ethyl acetate/hexane) to give the title compound (0.19 g).

MS: $[M+H]^+$ 356.1.

[0479]

15 H) methyl 5-(3-fluoro-4-(methylcarbamoyl)benzyl)-2-formyl-3,4-dimethylbenzoate

To a solution of methyl 5-(3-fluoro-4-(methylcarbamoyl)benzyl)-3,4-dimethyl-2-vinylbenzoate (0.19 g) in a mixed solvent of acetone (2.20 mL)-acetonitrile (2.20 mL)-
20 water (2.20 mL) were added osmium oxide (fixed catalyst I) (0.07 g) and sodium periodate (0.56 g), and the mixture was stirred overnight at room temperature. The reaction mixture was filtered, and the filtrate was extracted with ethyl acetate. The organic layer was washed with water and saturated brine,
25 and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure to give the title compound (0.19 g) as a crude product. This compound was used in the next step without an additional purification.

MS: $[M+H]^+$ 358.2.

30 [0480]

I) 1,5-anhydro-2,4-dideoxy-2-(6-(3-fluoro-4-(methylcarbamoyl)benzyl)-4,5-dimethyl-1-oxo-1,3-dihydro-2H-isoindol-2-yl)-L-threo-pentitol

To a solution of methyl 5-(3-fluoro-4-(methylcarbamoyl)benzyl)-2-formyl-3,4-dimethylbenzoate (0.09 g)
35

in THF (2.00 mL) were added (3S,4S)-3-aminotetrahydro-2H-pyran-4-ol (0.03 g) and anhydrous magnesium sulfate (0.06 g), and the mixture was stirred at room temperature for 5 hr. The reaction mixture was concentrated, and the residue was diluted with
5 methanol (2.00 mL)-THF (2.00 mL). Sodium triacetoxyborohydride (0.11 g) was added thereto, and the mixture was stirred overnight at room temperature. The reaction mixture was diluted with ethyl acetate, and the mixture was washed with water and saturated brine. The organic layer was dried over
10 anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (methanol/ethyl acetate) to give the title compound (0.04 g).

¹H NMR (300 MHz, DMSO-d₆) δ 1.47-1.63 (1H, m), 1.89-1.99 (1H, m), 2.18 (3H, s), 2.22 (3H, s), 2.75 (3H, d, J = 4.5 Hz), 3.40 (2H, d, J = 10.5 Hz), 3.70 (1H, dd, J = 10.9, 3.4 Hz), 3.81-3.97 (3H, m), 4.14 (2H, s), 4.35-4.51 (2H, m), 5.05 (1H, d, J = 5.3 Hz), 6.96-7.05 (2H, m), 7.36 (1H, s), 7.54 (1H, t, J = 7.9 Hz), 8.14 (1H, brs).

20 [0481]

Example 20-1

4-fluoro-2-[(3S,4S)-4-hydroxytetrahydro-2H-pyran-3-yl]-5-methyl-6-[4-(1H-pyrazol-1-yl)benzyl]-2,3-dihydro-1H-isoindol-1-one

25 Alias; 1,5-anhydro-2,4-dideoxy-2-(4-fluoro-5-methyl-1-oxo-6-(4-(1H-pyrazol-1-yl)benzyl)-1,3-dihydro-2H-isoindol-2-yl)-L-threo-pentitol

A) 3-fluoro-2-hydroxy-4-methylbenzoic acid

To an aqueous solution (50.0 mL) of sodium chlorite (22.2
30 g) was added a mixture of 3-fluoro-2-hydroxy-4-methylbenzaldehyde (9.47 g), sodium dihydrogenphosphate (33.2 g) and 2-methyl-2-butene (32.5 mL) in tert-butanol (200 mL)-water (100 mL) under ice-cooling, and the mixture was stirred at the same temperature for 3 hr. The pH of the reaction
35 mixture was adjusted to 2-3 with 2N hydrochloric acid. To the

reaction mixture was added water, and the mixture was extracted with ethyl acetate. The organic layer was washed with water and saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure to give the title compound (10.5 g) as a crude product. This compound was used in the next step without an additional purification. [0482]

B) methyl 3-fluoro-2-hydroxy-4-methylbenzoate

To a solution of 3-fluoro-2-hydroxy-4-methylbenzoic acid (10.5 g) in methanol (50.0 mL) was added sulfuric acid (5.00 mL) at room temperature, and the mixture was stirred at 60°C for 24 hr. The solvent was evaporated under reduced pressure, to the residue were added water and ethyl acetate, and the mixture was extracted with ethyl acetate. The organic layer was washed with water and saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (ethyl acetate/hexane) to give the title compound (6.40 g).

MS: $[M+H]^+$ 185.0.

[0483]

C) methyl 5-bromo-3-fluoro-2-hydroxy-4-methylbenzoate

To a solution of methyl 3-fluoro-2-hydroxy-4-methylbenzoate (6.40 g) in acetic acid (120 mL) was added bromine (1.87 mL) at room temperature, and the mixture was stirred at the same temperature for 2 hr. To the reaction mixture was added 10% aqueous sodium thiosulfate solution, and the mixture was extracted with ethyl acetate. The organic layer was washed with water and saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure to give a mixture (7.99 g) of the title compound and the raw material (about 2:1). The obtained product was used in the next step without an additional purification.

^1H NMR (300 MHz, CDCl_3) δ 2.38 (3H, d, $J = 2.8$ Hz), 3.98 (3H,

s), 7.82 (1H, d, J = 2.1 Hz), 10.67 (1H, s).

[0484]

D) methyl 3-fluoro-2-hydroxy-4-methyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate

5 trans-Dichlorobis(triphenylphosphine)palladium(II) (1.07 g) was added to a mixture of the 2:1 mixture (7.99 g) of methyl 5-bromo-3-fluoro-2-hydroxy-4-methylbenzoate and methyl 3-fluoro-2-hydroxy-4-methylbenzoate, bis(pinacolato)diboron (11.6 g), potassium acetate (8.94 g) and toluene (160 mL) at room
10 temperature under argon atmosphere, and the mixture was stirred at 100°C for 2 hr. To the reaction mixture was again added trans-dichlorobis(triphenylphosphine)palladium(II) (1.07 g), and the mixture was stirred at 100°C for 3 days. To the reaction mixture was again added trans-
15 dichlorobis(triphenylphosphine)palladium(II) (1.07 g), and the mixture was stirred overnight at 100°C. The reaction mixture was allowed to be cooled to room temperature, water was added thereto, and the precipitate was removed by filtration. The filtrate was extracted with ethyl acetate, the organic layer
20 was washed with water and saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (ethyl acetate/hexane) to give the title compound (2.20 g). In addition, the title compound (4.03 g)
25 was obtained from second fraction of column chromatography.
MS: [M+H]⁺ 311.1.

[0485]

E) methyl 5-(4-(1H-pyrazol-1-yl)benzyl)-3-fluoro-2-hydroxy-4-methylbenzoate

30 Tetrakis(triphenylphosphine)palladium(0) (0.40 g) was added to a mixture of methyl 3-fluoro-2-hydroxy-4-methyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (2.13 g), 1-(4-(chloromethyl)phenyl)-1H-pyrazole (1.32 g), sodium carbonate (1.46 g), DME (30.0 mL) and water (10.0 mL) under
35 argon atmosphere, and the mixture was stirred overnight at 80°C.

The reaction mixture was allowed to be cooled to room temperature, water was added thereto, and the mixture was extracted with ethyl acetate. The organic layer was washed with saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (ethyl acetate/hexane) to give the title compound (0.33 g). MS: $[M+H]^+$ 341.1.

[0486]

10 F) methyl 5-(4-(1H-pyrazol-1-yl)benzyl)-3-fluoro-4-methyl-2-(((trifluoromethyl)sulfonyl)oxy)benzoate

To a solution of methyl 5-(4-(1H-pyrazol-1-yl)benzyl)-3-fluoro-2-hydroxy-4-methylbenzoate (0.33 g) in DMF (7.0 mL) were added sodium hydride (0.047 g) and N-phenylbis(trifluoromethanesulfonimide) (0.38 g) under ice-cooling, and the mixture was stirred at room temperature for 2 hr. To the reaction mixture was added 1N hydrochloric acid, and the mixture was extracted with ethyl acetate. The organic layer was washed with saturated aqueous sodium bicarbonate solution and saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (ethyl acetate/hexane) to give the title compound (0.36 g).

25 MS: $[M+H]^+$ 473.1.

[0487]

G) methyl 5-(4-(1H-pyrazol-1-yl)benzyl)-3-fluoro-4-methyl-2-vinylbenzoate

A mixture of methyl 5-(4-(1H-pyrazol-1-yl)benzyl)-3-fluoro-4-methyl-2-(((trifluoromethyl)sulfonyl)oxy)benzoate (0.36 g), tributylvinyltin (0.33 mL), trans-dichlorobis(triphenylphosphine)palladium(II) (0.03 g), lithium chloride (0.24 g) and DMF (7.2 mL) was stirred at 90°C for 1 hr under argon atmosphere. To the reaction mixture was added aqueous potassium fluoride solution, and the insoluble

substance was removed by filtration through Celite. The filtrate was diluted with ethyl acetate, and the mixture was washed with saturated brine. The organic layer was dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (ethyl acetate/hexane) to give the title compound (0.21 g).

MS: $[M+H]^+$ 351.2.

[0488]

10 H) methyl 5-(4-(1H-pyrazol-1-yl)benzyl)-3-fluoro-2-formyl-4-methylbenzoate

To a mixture of methyl 5-(4-(1H-pyrazol-1-yl)benzyl)-3-fluoro-4-methyl-2-vinylbenzoate (0.21 g), acetone (2.3 mL), acetonitrile (2.3 mL) and water (2.3 mL) were added osmium oxide (fixed catalyst I) (0.08 g) and sodium periodate (0.63 g) at room temperature, and the mixture was stirred overnight at the same temperature. The insoluble substance was removed by filtration, and the filtrate was diluted with ethyl acetate. The solution was washed with saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure to give the title compound (0.21 g) as a crude product. This compound was used in the next step without an additional purification.

MS: $[M+H]^+$ 353.1.

25 [0489]

I) 4-fluoro-2-[(3S,4S)-4-hydroxytetrahydro-2H-pyran-3-yl]-5-methyl-6-[4-(1H-pyrazol-1-yl)benzyl]-2,3-dihydro-1H-isoindol-1-one

Alias; 1,5-anhydro-2,4-dideoxy-2-(4-fluoro-5-methyl-1-oxo-6-(4-(1H-pyrazol-1-yl)benzyl)-1,3-dihydro-2H-isoindol-2-yl)-L-threo-pentitol

A mixture of methyl 5-(4-(1H-pyrazol-1-yl)benzyl)-3-fluoro-2-formyl-4-methylbenzoate (0.21 g), (3S,4S)-3-aminotetrahydro-2H-pyran-4-ol (0.07 g), anhydrous magnesium sulfate (0.14 g) and THF (4.00 mL) was stirred at room

temperature for 6 hr under nitrogen atmosphere. The insoluble substance was removed by filtration, and the filtrate was concentrated. The residue was dissolved in a mixed solvent of methanol (4.00 mL)-THF (4.00 mL), sodium triacetoxyborohydride
5 (0.25 g) was added thereto, and the mixture was stirred at room temperature for 16 hr. The reaction mixture was diluted with ethyl acetate, and the mixture was washed with water and saturated brine. The organic layer was dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced
10 pressure. The residue was purified by silica gel column chromatography (ethyl acetate/hexane) to give the title compound (0.11 g).

¹H NMR (300 MHz, DMSO-d₆) δ 1.45-1.64 (1H, m), 1.88-1.99 (1H, m), 2.23 (3H, d, J = 2.1 Hz), 3.33-3.47 (2H, m), 3.64-3.95 (4H, m), 4.15 (2H, s), 4.55 (2H, s), 5.08 (1H, d, J = 5.1 Hz), 6.49-
15 6.56 (1H, m), 7.27 (2H, d, J = 8.7 Hz), 7.38 (1H, s), 7.68-7.81 (3H, m), 8.44 (1H, d, J = 2.1 Hz).

[0490]

Example 20-2

20 4-fluoro-2-[(3S,4S)-4-hydroxytetrahydro-2H-pyran-3-yl]-5-methyl-6-[4-(1H-pyrazol-1-yl)benzyl]-2,3-dihydro-1H-isoindol-1-one

Alias; 1,5-anhydro-2,4-dideoxy-2-(4-fluoro-5-methyl-1-oxo-6-(4-(1H-pyrazol-1-yl)benzyl)-1,3-dihydro-2H-isoindol-2-yl)-L-threo-
25 pentitol

This compound was also synthesized by the following method.

[0491]

A) 3-fluoro-2-hydroxy-4-methylbenzoic acid

30 A mixture of 2,3-difluoro-4-methylbenzoic acid (25.0 g), sodium hydroxide (23.2 g) and DMSO (250 mL) was stirred at 140°C for 12 hr, and then overnight at room temperature, under argon atmosphere. To the reaction mixture was added 6M hydrochloric acid (100 mL) under ice-cooling. Ethyl acetate
35 and water were added thereto at room temperature, the organic

layer was separated, and the aqueous layer was extracted with ethyl acetate. The organic layers were combined, washed with water and saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure
5 to give the title compound (24.3 g). This compound was used in the next step without an additional purification.

^1H NMR (300 MHz, CDCl_3) δ 2.35 (3H, d, $J = 2.3$ Hz), 6.74 (1H, dd, $J = 7.9, 6.8$ Hz), 7.58 (1H, dd, $J = 8.3, 1.5$ Hz), 10.37 (1H, s), 1H undetected.

10 [0492]

B) methyl 3-fluoro-2-hydroxy-4-methylbenzoate

To a solution of 3-fluoro-2-hydroxy-4-methylbenzoic acid (24.3 g) in methanol (500 mL) was added sulfuric acid (7.60 mL) at room temperature, and the mixture was stirred at 60°C for 3
15 days. The solvent was evaporated under reduced pressure, and ethyl acetate and saturated brine were added thereto. The organic layer was separated, and the aqueous layer was extracted with ethyl acetate. The combined organic layers were washed with water and saturated brine, and dried over anhydrous
20 magnesium sulfate, and the solvent was evaporated under reduced pressure. To the residue was added diisopropyl ether (50 mL), and the insoluble substance was removed by filtration. The filtrate was concentrated under reduced pressure to give the title compound (25.4 g).

25 ^1H NMR (300 MHz, CDCl_3) δ 2.32 (3H, d, $J = 2.5$ Hz), 3.95 (3H, s), 6.68 (1H, dd, $J = 7.9, 6.8$ Hz), 7.49 (1H, dd, $J = 8.3, 1.7$ Hz), 10.74 (1H, s).

[0493]

C) methyl 5-bromo-3-fluoro-2-hydroxy-4-methylbenzoate

30 To a solution of methyl 3-fluoro-2-hydroxy-4-methylbenzoate (25.4 g) in acetic acid (250 mL) was added bromine (7.78 mL), and the mixture was stirred at room temperature for 3 hr. To the reaction mixture was added 5% aqueous sodium thiosulfate solution (250 mL) at room
35 temperature, and the precipitate was collected by filtration,

and washed with water to give the title compound (21.1 g).

¹H NMR (300 MHz, CDCl₃) δ 2.37 (3H, d, J = 2.6 Hz), 3.97 (3H, s), 7.81 (1H, d, J = 1.7 Hz), 10.65 (1H, s).

[0494]

- 5 D) methyl 3-fluoro-2-hydroxy-4-methyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate

trans-Dichlorobis(triphenylphosphine)palladium(II) (2.0 g) was added to a mixture of methyl 5-bromo-3-fluoro-2-hydroxy-4-methylbenzoate (15.0 g), bis(pinacolato)diboron (21.7 g),
10 potassium acetate (16.8 g) and toluene (290 mL) under argon atmosphere, and the mixture was stirred at 110°C for 15 hr. To the reaction mixture was added water, and the mixture was extracted with ethyl acetate. The organic layer was washed with water and saturated brine, and dried over anhydrous
15 magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was washed with diisopropyl ether to give the title compound (12.3 g). The filtrate was purified by silica gel column chromatography (ethyl acetate/hexane) to give the title compound (1.99 g).

- 20 MS: [M+H]⁺ 311.2.

[0495]

- E) methyl 5-(4-(1H-pyrazol-1-yl)benzyl)-3-fluoro-2-hydroxy-4-methylbenzoate

Tetrakis(triphenylphosphine)palladium(0) (2.52 g) was
25 added to a mixture of methyl 3-fluoro-2-hydroxy-4-methyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (13.5 g), 1-(4-(chloromethyl)phenyl)-1H-pyrazole (8.39 g), sodium carbonate (9.23 g), DME (195 mL) and water (65.0 mL) under argon atmosphere, and the mixture was stirred overnight at 80°C.
30 To the reaction mixture was added water, and the mixture was extracted with ethyl acetate. The organic layer was washed with saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was washed with a mixed solvent of diisopropyl
35 ether-ethyl acetate to give a crude product. The filtrate was

concentrated under reduced pressure, and the residue was purified by silica gel column chromatography (ethyl acetate/methanol). The obtained crude products are combined, and washed with diisopropyl ether-ethyl acetate to give the
5 title compound (12.4 g).

MS: $[M+H]^+$ 341.1.

[0496]

F) methyl 5-(4-(1H-pyrazol-1-yl)benzyl)-3-fluoro-4-methyl-2-(((trifluoromethyl)sulfonyl)oxy)benzoate

10 To a solution of methyl 5-(4-(1H-pyrazol-1-yl)benzyl)-3-fluoro-2-hydroxy-4-methylbenzoate (12.4 g) in DMF (250 mL) were added sodium hydride (1.74 g) and N-phenylbis(trifluoromethanesulfonimide) (14.3 g) under ice-cooling, and the mixture was stirred at room temperature for 2
15 hr. To the reaction mixture was added 1N hydrochloric acid, and the mixture was extracted with ethyl acetate. The organic layer was washed with saturated aqueous sodium bicarbonate solution and saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced
20 pressure. The residue was purified by silica gel column chromatography (ethyl acetate/hexane) to give the title compound (12.5 g).

MS: $[M+H]^+$ 473.1.

[0497]

25 G) methyl 5-(4-(1H-pyrazol-1-yl)benzyl)-3-fluoro-4-methyl-2-vinylbenzoate

A mixture of methyl 5-(4-(1H-pyrazol-1-yl)benzyl)-3-fluoro-4-methyl-2-(((trifluoromethyl)sulfonyl)oxy)benzoate (11.8 g), tributylvinyltin (11.9 g), trans-
30 dichlorobis(triphenylphosphine)palladium(II) (0.88 g), lithium chloride (7.84 g) and DMF (240 mL) was stirred at 90°C for 1.5 hr under argon atmosphere. To the reaction mixture was added aqueous potassium fluoride solution, and the precipitated insoluble substance was removed by filtration through Celite.
35 The filtrate was diluted with ethyl acetate, and the mixture

was washed with saturated brine. The organic layer was dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (ethyl acetate/hexane) to give
5 the title compound (7.98 g).

MS: $[M+H]^+$ 351.2.

[0498]

H) methyl 5-(4-(1H-pyrazol-1-yl)benzyl)-3-fluoro-2-formyl-4-methylbenzoate

10 To a mixture of methyl 5-(4-(1H-pyrazol-1-yl)benzyl)-3-fluoro-4-methyl-2-vinylbenzoate (7.97 g), acetone (93.0 mL), acetonitrile (93.0 mL) and water (93.0 mL) were added osmium oxide (fixed catalyst I) (2.89 g) and sodium periodate (24.3 g) at room temperature, and the mixture was stirred overnight at
15 the same temperature. The insoluble substance was removed by filtration, and the filtrate was diluted with ethyl acetate. The solution was washed with saturated brine, the organic layer was and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure to give the title
20 compound (7.97 g) as a crude product. This compound was used in the next step without an additional purification.

MS: $[M+H]^+$ 353.2.

[0499]

I) 4-fluoro-2-[(3S,4S)-4-hydroxytetrahydro-2H-pyran-3-yl]-5-methyl-6-[4-(1H-pyrazol-1-yl)benzyl]-2,3-dihydro-1H-isoindol-1-one
25

Alias; 1,5-anhydro-2,4-dideoxy-2-(4-fluoro-5-methyl-1-oxo-6-(4-(1H-pyrazol-1-yl)benzyl)-1,3-dihydro-2H-isoindol-2-yl)-L-threo-pentitol

30 A mixture of methyl 5-(4-(1H-pyrazol-1-yl)benzyl)-3-fluoro-2-formyl-4-methylbenzoate (7.97 g), (3S,4S)-3-aminotetrahydro-2H-pyran-4-ol (2.65 g), anhydrous magnesium sulfate (5.23 g) and THF (160 mL) was stirred at room temperature for 5 hr under nitrogen atmosphere. The insoluble
35 substance was removed by filtration, and the filtrate was

concentrated. The residue was dissolved in a mixed solvent of methanol (120 mL)-THF (150 mL), sodium triacetoxyborohydride (9.59 g) was added thereto, and the mixture was stirred at room temperature for 15 hr. Sodium triacetoxyborohydride (9.59 g) was again added thereto, and the mixture was stirred for 3 hr. The reaction mixture was diluted with ethyl acetate, and the mixture was washed with water and saturated brine. The organic layer was dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was dissolved in DMSO-toluene, and the solution was purified by NH silica gel column chromatography (ethyl acetate/hexane). The obtained crude product was dissolved in THF, the solution was washed with water and saturated brine, and the solvent was evaporated under reduced pressure. The residue was washed with a mixed solvent of diisopropyl ether-ethyl acetate to give a crude product. The crude product (5.28 g) was dissolved in hot ethanol (60 mL), and recrystallized over 4 hr under ice-cooling to give the title compound (4.88 g).

^1H NMR (300 MHz, CDCl_3) δ 1.71-1.87 (1H, m), 2.08-2.17 (1H, m), 2.22 (3H, d, $J = 2.3$ Hz), 2.47-2.57 (1H, m), 3.42-3.62 (2H, m), 3.97-4.15 (6H, m), 4.31-4.63 (2H, m), 6.39-6.50 (1H, m), 7.17 (2H, d, $J = 8.7$ Hz), 7.47 (1H, s), 7.59 (2H, d, $J = 8.7$ Hz), 7.70 (1H, d, $J = 1.5$ Hz), 7.88 (1H, d, $J = 2.3$ Hz).

X-ray powder diffraction pattern with specific peaks at d value (or d -spacing) = 12.8, 8.0, 7.5, 6.2, 6.0, 5.6, 5.0, 4.6, 4.5 and 4.2 Å.

[0500]

Example 21

1,5-anhydro-2,4-dideoxy-2-(5-methyl-1-oxo-6-(4-(1H-pyrazol-1-yl)benzyl)-1,3-dihydro-2H-isoindol-2-yl)-L-threo-pentitol

To a solution of methyl 5-(4-(1H-pyrazol-1-yl)benzyl)-2-formyl-4-methylbenzoate (0.20 g) in THF (4.00 mL) was added (3S,4S)-3-aminotetrahydro-2H-pyran-4-ol (0.07 g), and the mixture was stirred at room temperature for 4 hr under argon atmosphere. The reaction mixture was concentrated, and the

residue was diluted with acetic acid (4.00 mL). Sodium triacetoxyborohydride (0.19 g) was added thereto, and the mixture was stirred overnight at room temperature. The reaction mixture was diluted with saturated aqueous sodium bicarbonate solution and extracted with ethyl acetate, the organic layer was washed with water and saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by NH silica gel column chromatography (ethyl acetate/hexane) to give the title compound (0.11 g).

¹H NMR (300 MHz, CDCl₃) δ 1.71-1.90 (1H, m), 2.07-2.18 (1H, m), 2.31 (3H, s), 2.69 (1H, d, J = 5.5 Hz), 3.43-3.59 (2H, m), 3.98-4.15 (6H, m), 4.29-4.55 (2H, m), 6.45 (1H, t, J = 2.1 Hz), 7.18 (2H, d, J = 8.5 Hz), 7.24 (1H, s), 7.55-7.61 (2H, m), 7.63 (1H, s), 7.70 (1H, d, J = 1.7 Hz), 7.88 (1H, d, J = 2.5 Hz). [0501]

Example 22

1,5-anhydro-2,4-dideoxy-2-(6-(4-(difluoromethoxy)benzyl)-4,5-dimethyl-1-oxo-1,3-dihydro-2H-isoindol-2-yl)-L-threo-pentitol
A) methyl 5-(4-(difluoromethoxy)benzyl)-2-hydroxy-3,4-dimethylbenzoate

To a solution of methyl 2-hydroxy-3,4-dimethyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (0.60 g) in DME (12.0 mL) were added 1-(bromomethyl)-4-(difluoromethoxy)benzene (0.47 g), [1,1'-bis(diphenylphosphino)ferrocene]dichloropalladium(II) dichloromethane adduct (0.16 g) and 2mol/L aqueous sodium carbonate solution (1.96 mL), and the mixture was stirred overnight at 80°C under argon atmosphere. Water and ethyl acetate were added thereto, the mixture was allowed to be cooled to room temperature, and the precipitate was removed by filtration. The filtrate was extracted with ethyl acetate, the organic layer was washed with water and saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by

silica gel column chromatography (ethyl acetate/hexane) to give the title compound (0.60 g).

MS: $[M+H]^+$ 337.0.

[0502]

- 5 B) methyl 5-(4-(difluoromethoxy)benzyl)-3,4-dimethyl-2-vinylbenzoate

To a solution of methyl 5-(4-(difluoromethoxy)benzyl)-2-hydroxy-3,4-dimethylbenzoate (0.60 g) in DMF (12.0 mL) was added sodium hydride (0.08 g) under ice-cooling, and the mixture was stirred at room temperature for 30 min. To this reaction mixture was added N-phenylbis(trifluoromethanesulfonimide) (0.77 g) under ice-cooling, and the mixture was stirred at room temperature for 1 hr. To the reaction mixture was added 1N hydrochloric acid under ice-cooling, and the mixture was extracted with ethyl acetate. The organic layer was washed with water and saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The obtained crude methyl 5-[4-(difluoromethoxy)benzyl]-3,4-dimethyl-2-
20 {[(trifluoromethyl)sulfonyl]oxy}benzoate was used in the next step without an additional purification.

The above-mentioned compound was dissolved in DMF (12.0 mL), tributylvinyltin (0.78 mL), trans-dichlorobis(triphenylphosphine)palladium(II) (0.13 g) and lithium chloride (0.53 g) were added thereto, and the mixture was stirred overnight at 90°C under argon atmosphere. To the reaction mixture was added aqueous potassium fluoride solution, and the precipitated insoluble substance was removed by filtration through Celite. The filtrate was diluted with ethyl acetate, and the mixture was washed with water and saturated brine. The organic layer was dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (ethyl acetate/hexane) to give the title compound (0.16 g).
35 MS: $[M+H]^+$ 347.2.

[0503]

C) methyl 5-(4-(difluoromethoxy)benzyl)-2-formyl-3,4-dimethylbenzoate

To a solution of methyl 5-(4-(difluoromethoxy)benzyl)-
5 3,4-dimethyl-2-vinylbenzoate (0.16 g) in a mixed solvent of
acetone (3.00 mL)-acetonitrile (3.00 mL)-water (3.00 mL) were
added osmium oxide (fixed catalyst I) (0.06 g) and sodium
periodate (0.49 g), and the mixture was stirred overnight at
room temperature. The reaction mixture was filtered, and the
10 filtrate was extracted with ethyl acetate. The organic layer
was washed with water and saturated brine, and dried over
anhydrous sodium sulfate, and the solvent was evaporated under
reduced pressure. The residue was purified by silica gel
column chromatography (ethyl acetate/hexane) to give the title
15 compound (0.04 g).

MS: [M+H]⁺ 349.1.

[0504]

D) 1,5-anhydro-2,4-dideoxy-2-(6-(4-(difluoromethoxy)benzyl)-
4,5-dimethyl-1-oxo-1,3-dihydro-2H-isoindol-2-yl)-L-threo-
20 pentitol

To a solution of methyl 5-(4-(difluoromethoxy)benzyl)-2-
formyl-3,4-dimethylbenzoate (0.04 g) in THF (2.00 mL) was added
(3S,4S)-3-aminotetrahydro-2H-pyran-4-ol (0.01 g) under argon
atmosphere, and the mixture was stirred at room temperature for
25 3 hr. The reaction mixture was concentrated, and the residue
was diluted with acetic acid (2.00 mL). Sodium
triacetoxymethylborohydride (0.03 g) was added thereto under argon
atmosphere, and the mixture was stirred overnight at room
temperature. The reaction mixture was diluted with saturated
30 aqueous sodium bicarbonate solution, and extracted with ethyl
acetate. The organic layer was washed with water and saturated
brine, and dried over anhydrous magnesium sulfate, and the
solvent was evaporated under reduced pressure. The residue was
purified by NH silica gel column chromatography (ethyl
35 acetate/hexane) to give the title compound (0.02 g).

¹H NMR (300 MHz, CDCl₃) δ 1.73-1.88 (1H, m), 2.09-2.18 (1H, m), 2.21 (3H, s), 2.25 (3H, s), 2.52 (1H, d, J = 5.3 Hz), 3.45-3.63 (2H, m), 4.00-4.18 (6H, m), 4.26-4.50 (2H, m), 6.19-6.75 (1H, m), 6.99-7.04 (2H, m), 7.06-7.11 (2H, m), 7.51 (1H, s).

5 [0505]

Example 23

1,5-anhydro-2,4-dideoxy-2-(6-(4-fluoro-3-methoxybenzyl)-4,5-dimethyl-1-oxo-1,3-dihydro-2H-isoindol-2-yl)-L-threo-pentitol
A) methyl 5-(4-fluoro-3-methoxybenzyl)-2-hydroxy-3,4-

10 dimethylbenzoate

To a solution of methyl 2-hydroxy-3,4-dimethyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (0.50 g) in DME (12.0 mL) were added 4-(bromomethyl)-1-fluoro-2-methoxybenzene (0.39 g), [1,1'-

15 bis(diphenylphosphino)ferrocene]dichloropalladium(II)

dichloromethane adduct (0.07 g) and 2mol/L aqueous sodium carbonate solution (1.63 mL), and the mixture was stirred

overnight at 80°C under argon atmosphere. The reaction mixture was allowed to be cooled to room temperature, and diluted with
20 ethyl acetate, and the mixture was washed with saturated brine. The organic layer was dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (ethyl acetate/hexane) to give the title compound (0.45 g).

25 ¹H NMR (300 MHz, DMSO-d₆) δ 2.13 (6H, s), 3.78 (3H, s), 3.88 (3H, s), 3.94 (2H, s), 6.53 (1H, ddd, J = 8.3, 4.4, 2.1 Hz), 6.97 (1H, dd, J = 8.5, 1.9 Hz), 7.07 (1H, dd, J = 11.5, 8.3 Hz), 7.49 (1H, s), 10.90 (1H, s).

[0506]

30 B) methyl 5-(4-fluoro-3-methoxybenzyl)-3,4-dimethyl-2-(((trifluoromethyl)sulfonyl)oxy)benzoate

To a solution of methyl 5-(4-fluoro-3-methoxybenzyl)-2-hydroxy-3,4-dimethylbenzoate (0.45 g) in DMF (5.0 mL) were added sodium hydride (0.07 g) and N-

35 phenylbis(trifluoromethanesulfonimide) (0.55 g) under ice-

cooling, and the mixture was stirred at room temperature for 2 hr under argon atmosphere. The reaction mixture was diluted with ethyl acetate, and the mixture was washed with saturated brine. The organic layer was dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (ethyl acetate/hexane) to give the title compound (0.63 g). ¹H NMR (300 MHz, DMSO-d₆) δ 2.19-2.32 (6H, m), 3.76-3.85 (6H, m), 4.08 (2H, s), 6.59 (1H, ddd, J = 8.3, 4.3, 2.1 Hz), 6.99-7.16 (2H, m), 7.61 (1H, s).

[0507]

C) methyl 5-(4-fluoro-3-methoxybenzyl)-3,4-dimethyl-2-vinylbenzoate

To a solution of methyl 5-(4-fluoro-3-methoxybenzyl)-3,4-dimethyl-2-(((trifluoromethyl)sulfonyl)oxy)benzoate (0.63 g) in DMF (7.00 mL) were added tributylvinyltin (0.61 mL), trans-dichlorobis(triphenylphosphine)palladium(II) (0.05 g) and lithium chloride (0.44 g), and the mixture was stirred at 90°C for 2 hr under argon atmosphere. To the reaction mixture was added aqueous potassium fluoride solution, and the precipitated insoluble substance was removed by filtration through Celite. The filtrate was diluted with ethyl acetate, and the mixture was washed with saturated brine. The organic layer was dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (ethyl acetate/hexane) to give the title compound (0.39 g).

MS: [M+H]⁺ 329.1.

[0508]

D) methyl 5-(4-fluoro-3-methoxybenzyl)-2-formyl-3,4-dimethylbenzoate

To a solution of methyl 5-(4-fluoro-3-methoxybenzyl)-3,4-dimethyl-2-vinylbenzoate (0.39 g) in a mixed solvent of acetone (9.00 mL)-acetonitrile (9.00 mL)-water (9.00 mL) were added osmium oxide (fixed catalyst I) (0.15 g) and sodium periodate

(1.27 g), and the mixture was stirred at room temperature for 2.5 days. The reaction mixture was filtered, and the filtrate was extracted with ethyl acetate. The organic layer was washed with saturated brine, and dried over anhydrous magnesium sulfate to give the title compound (0.39 g). This compound was used in the next step without an additional purification.

MS: $[M+H]^+$ 331.1.

[0509]

E) 1,5-anhydro-2,4-dideoxy-2-(6-(4-fluoro-3-methoxybenzyl)-4,5-dimethyl-1-oxo-1,3-dihydro-2H-isoindol-2-yl)-L-threo-pentitol

To a solution of methyl 5-(4-fluoro-3-methoxybenzyl)-2-formyl-3,4-dimethylbenzoate (0.35 g) in THF (4.00 mL) were added (3S,4S)-3-aminotetrahydro-2H-pyran-4-ol (0.12 g) and anhydrous magnesium sulfate (0.26 g), and the mixture was stirred at room temperature for 5 hr under nitrogen atmosphere. The insoluble substance was removed by filtration, the filtrate was concentrated, and the residue was diluted with methanol (2.00 mL)-THF (4.00 mL). Sodium triacetoxyborohydride (0.45 g) was added thereto, and the mixture was stirred overnight at room temperature. The reaction mixture was diluted with ethyl acetate, and the mixture was washed with water and saturated brine. The organic layer was dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (ethyl acetate/hexane) to give the title compound (0.13 g).

^1H NMR (300 MHz, DMSO- d_6) δ 1.47-1.66 (1H, m), 1.89-2.00 (1H, m), 2.18-2.26 (6H, m), 3.34-3.45 (2H, m), 3.69 (1H, dd, J = 10.9, 3.4 Hz), 3.78 (3H, s), 3.82-3.96 (3H, m), 4.05 (2H, s), 4.34-4.50 (2H, m), 5.05 (1H, d, J = 4.5 Hz), 6.49-6.61 (1H, m), 6.97-7.13 (2H, m), 7.30 (1H, s).

[0510]

Example 24

1,5-anhydro-2-(4-chloro-6-(4-methoxybenzyl)-5-methyl-1-oxo-1,3-dihydro-2H-isoindol-2-yl)-2,4-dideoxy-L-threo-pentitol

A) methyl 5-bromo-3-chloro-2-hydroxy-4-methylbenzoate

To a solution of methyl 5-bromo-2-hydroxy-4-methylbenzoate (4.54 g) in DMF (34.0 mL) was added N-chlorosuccinimide (2.47 g), and the mixture was stirred overnight at room temperature. To the reaction mixture was added water, and the mixture was extracted with ethyl acetate. The organic layer was washed with water and saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure to give the title compound (5.20 g).

MS: $[M-H]^+$ 276.7.

[0511]

B) methyl 3-chloro-2-hydroxy-4-methyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate

To a solution of methyl 5-bromo-3-chloro-2-hydroxy-4-methylbenzoate (5.20 g) in toluene (140 mL) were added bis(pinacolato)diboron (7.09 g), potassium acetate (5.48 g) and trans-dichlorobis(triphenylphosphine)palladium(II) (0.65 g), and the mixture was stirred at 110°C for 15 hr under argon atmosphere. The reaction mixture was allowed to be cooled to room temperature, water and ethyl acetate were added thereto, and the mixture was partitioned. The organic layer was washed with water and saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (ethyl acetate/hexane) to give the title compound (3.54 g).

MS: $[M+H]^+$ 327.1.

[0512]

C) methyl 3-chloro-2-hydroxy-5-(4-methoxybenzyl)-4-methylbenzoate

To a solution of methyl 3-chloro-2-hydroxy-4-methyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (0.90 g) in a mixed solvent of DME (13.5 mL)-water (4.50 mL) were added 1-(chloromethyl)-4-methoxybenzene (0.43 g), tetrakis(triphenylphosphine)palladium(0) (0.16 g) and sodium

carbonate (0.58 g), and the mixture was stirred at 80°C overnight under argon atmosphere. The reaction mixture was allowed to be cooled to room temperature, water and ethyl acetate were added thereto, and the mixture was partitioned.

5 The organic layer was washed with water and saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (ethyl acetate/hexane) to give the title compound (0.37 g).

10 MS: $[M-H]^+$ 318.9.

[0513]

D) methyl 3-chloro-5-(4-methoxybenzyl)-4-methyl-2-
(((trifluoromethyl)sulfonyl)oxy)benzoate

To a solution of methyl 3-chloro-2-hydroxy-5-(4-methoxybenzyl)-4-methylbenzoate (0.36 g) in DMF (7.50 mL) were added sodium hydride (0.05 g) and N-phenylbis(trifluoromethanesulfonimide) (0.36 g) under ice-cooling, and the mixture was stirred at room temperature for 2.5 hr. To the reaction mixture was added 1N hydrochloric acid, and the mixture was extracted with ethyl acetate. The organic layer was washed with saturated aqueous sodium bicarbonate solution and saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (ethyl acetate/hexane) to give the title compound (0.51 g).

MS: $[M-H]^+$ 451.8.

[0514]

E) methyl 3-chloro-5-(4-methoxybenzyl)-4-methyl-2-vinylbenzoate

30 To a solution of methyl 3-chloro-5-(4-methoxybenzyl)-4-methyl-2-(((trifluoromethyl)sulfonyl)oxy)benzoate (0.51 g) in DMF (10.0 mL) were added tributylvinyltin (0.54 g), trans-dichlorobis(triphenylphosphine)palladium(II) (0.04 g) and lithium chloride (0.36 g), and the mixture was stirred at 90°C for 1 hr under argon atmosphere. To the reaction mixture was

added aqueous potassium fluoride solution, and the precipitated insoluble substance was removed by filtration through Celite. The filtrate was diluted with ethyl acetate, and the mixture was washed with water and saturated brine. The organic layer
5 was dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (ethyl acetate/hexane) to give the title compound (0.31 g).

MS: $[M+H]^+$ 331.1.

10 [0515]

F) methyl 3-chloro-2-formyl-5-(4-methoxybenzyl)-4-methylbenzoate

To a solution of methyl 3-chloro-5-(4-methoxybenzyl)-4-methyl-2-vinylbenzoate (0.31 g) in a mixed solvent of acetone
15 (3.60 mL)-acetonitrile (3.60 mL)-water (3.60 mL) were added osmium oxide (fixed catalyst I) (0.12 g) and sodium periodate (1.00 g), and the mixture was stirred overnight at room temperature. The reaction mixture was filtered, and the filtrate was extracted with ethyl acetate. The organic layer
20 was washed with water and saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure to give the title compound (0.31 g) as a crude product. This compound was used in the next step without an additional purification.

25 [0516]

G) 1,5-anhydro-2-(4-chloro-6-(4-methoxybenzyl)-5-methyl-1-oxo-1,3-dihydro-2H-isoindol-2-yl)-2,4-dideoxy-L-threo-pentitol

To a solution of methyl 3-chloro-2-formyl-5-(4-methoxybenzyl)-4-methylbenzoate (0.10 g) in THF (2.00 mL) were
30 added (3S,4S)-3-aminotetrahydro-2H-pyran-4-ol (0.04 g) and anhydrous magnesium sulfate (0.07 g), and the mixture was stirred at room temperature for 6 hr. The insoluble substance was removed by filtration, the filtrate was concentrated, and the residue was diluted with methanol (2.00 mL)-THF (2.00 mL).
35 Sodium triacetoxyborohydride (0.19 g) was added thereto, and

the mixture was stirred overnight at room temperature. The reaction mixture was diluted with ethyl acetate, and the mixture was washed with water and saturated brine. The organic layer was dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (ethyl acetate/hexane) to give the title compound (0.05 g).

¹H NMR (300 MHz, DMSO-d₆) δ 1.47-1.62 (1H, m), 1.89-1.99 (1H, m), 2.35 (3H, s), 3.34-3.48 (2H, m), 3.64-3.74 (4H, m), 3.80-3.97 (3H, m), 4.07 (2H, s), 4.39-4.54 (2H, m), 5.11 (1H, d, J = 5.3 Hz), 6.82-6.90 (2H, m), 7.05 (2H, d, J = 8.7 Hz), 7.42 (1H, s).

[0517]

Example 25-1

2-((1S,2S)-2-hydroxycyclohexyl)-4,5-dimethyl-6-((6-methylpyridin-3-yl)methyl)isoindolin-1-one

A) 5-(chloromethyl)-2-methylpyridine

To a solution of (6-methylpyridin-3-yl)methanol (1.08 g) in THF (15.0 mL) was added thionyl chloride (1.57 g) under ice-cooling, and the mixture was stirred overnight at room temperature. The reaction mixture was diluted with water and ethyl acetate, saturated aqueous sodium bicarbonate was added thereto, and the mixture was partitioned. The organic layer was washed with saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (ethyl acetate/hexane) to give the title compound (0.81 g).

¹H NMR (300 MHz, CDCl₃) δ 2.57 (3H, s), 4.57 (2H, s), 7.17 (1H, d, J = 7.9 Hz), 7.63 (1H, dd, J = 7.9, 2.3 Hz), 8.50 (1H, d, J = 2.3 Hz).

[0518]

B) methyl 2-hydroxy-3,4-dimethyl-5-((6-methylpyridin-3-yl)methyl)benzoate

To a solution of methyl 2-hydroxy-3,4-dimethyl-5-

(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (0.80 g) in a mixed solvent of DME (12.0 mL)-water (4.00 mL) were added 5-(chloromethyl)-2-methylpyridine (0.41 g), tetrakis(triphenylphosphine)palladium(0) (0.15 g) and sodium carbonate (0.55 g), and the mixture was stirred overnight at 80°C under argon atmosphere. The reaction mixture was allowed to be cooled to room temperature, water and ethyl acetate were added thereto, and the mixture was partitioned. The organic layer was washed with water and saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (ethyl acetate/hexane) to give the title compound (0.48 g).

MS: $[M-H]^+$ 286.1.

[0519]

C) methyl 3,4-dimethyl-5-((6-methylpyridin-3-yl)methyl)-2-(((trifluoromethyl)sulfonyl)oxy)benzoate

To a solution of methyl 2-hydroxy-3,4-dimethyl-5-((6-methylpyridin-3-yl)methyl)benzoate (0.48 g) in DMF (9.50 mL) were added sodium hydride (0.08 g) and N-phenylbis(trifluoromethanesulfonimide) (0.66 g) under ice-cooling, and the mixture was stirred at room temperature for 2.5 hr. To the reaction mixture was added 1N hydrochloric acid, and the mixture was extracted with ethyl acetate. The organic layer was washed with saturated aqueous sodium bicarbonate solution and saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (ethyl acetate/hexane) to give the title compound (0.70 g).

MS: $[M+H]^+$ 418.1.

[0520]

D) methyl 3,4-dimethyl-5-((6-methylpyridin-3-yl)methyl)-2-vinylbenzoate

To a solution of methyl 3,4-dimethyl-5-((6-methylpyridin-

3-yl)methyl)-2-(((trifluoromethyl)sulfonyl)oxy)benzoate (0.70 g) in DMF (14.0 mL) were added tributylvinyltin (0.80 g), trans-dichlorobis(triphenylphosphine)palladium(II) (0.06 g) and lithium chloride (0.53 g), and the mixture was stirred at 90°C for 1 hr under argon atmosphere. To the reaction mixture was added aqueous potassium fluoride solution, and the precipitated insoluble substance was removed by filtration through Celite. The filtrate was diluted with ethyl acetate, and the mixture was washed with water and saturated brine. The organic layer was dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (ethyl acetate/hexane) to give the title compound (0.44 g).

MS: $[M+H]^+$ 296.1.

[0521]

E) methyl 2-formyl-3,4-dimethyl-5-((6-methylpyridin-3-yl)methyl)benzoate

To a solution of methyl 3,4-dimethyl-5-((6-methylpyridin-3-yl)methyl)-2-vinylbenzoate (0.44 g) in a mixed solvent of acetone (5.40 mL)-acetonitrile (5.40 mL)-water (5.40 mL) were added osmium oxide (fixed catalyst I) (0.19 g) and sodium periodate (1.59 g), and the mixture was stirred overnight at room temperature. The reaction mixture was filtered, and the filtrate was extracted with ethyl acetate. The organic layer was washed with water and saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure to give the title compound (0.44 g) as a crude product. This compound was used in the next step without an additional purification.

MS: $[M+H]^+$ 298.1.

[0522]

F) 2-((1S,2S)-2-hydroxycyclohexyl)-4,5-dimethyl-6-((6-methylpyridin-3-yl)methyl)isoindolin-1-one

To a solution of methyl 2-formyl-3,4-dimethyl-5-((6-methylpyridin-3-yl)methyl)benzoate (0.15 g) in THF (2.90 mL)

were added (1S,2S)-aminocyclohexanol (0.06 g) and anhydrous magnesium sulfate (0.11 g), and the mixture was stirred at room temperature for 6 hr. The insoluble substance was removed by filtration, the filtrate was concentrated, and the residue was
5 diluted with methanol (2.90 mL)-THF (2.90 mL). Sodium triacetoxyborohydride (0.21 g) was added thereto, and the mixture was stirred overnight at room temperature. The reaction mixture was diluted with ethyl acetate, and the mixture was washed with water and saturated brine. The organic
10 layer was dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (methanol/ethyl acetate) to give the title compound (0.05 g).

¹H NMR (300 MHz, DMSO-d₆) δ 1.22-1.38 (3H, m), 1.48-1.72 (4H, m), 1.90-2.01 (1H, m), 2.19 (3H, s), 2.22 (3H, s), 2.41 (3H, s),
15 3.55-3.66 (1H, m), 3.81 (1H, td, J = 10.7, 4.1 Hz), 4.05 (2H, s), 4.35 (2H, s), 4.71 (1H, d, J = 5.7 Hz), 7.14 (1H, d, J = 7.9 Hz), 7.27-7.38 (2H, m), 8.29 (1H, d, J = 1.9 Hz).

[0523]

20 Example 25-2

2-((1S,2S)-2-Hydroxycyclohexyl)-4,5-dimethyl-6-((6-methylpyridin-3-yl)methyl)isoindolin-1-one was also synthesized by the following method.

[0524]

25 A) 5-(chloromethyl)-2-methylpyridine hydrochloride

To a solution of (6-methylpyridin-3-yl)methanol (1.24 g) in THF (12.4 mL) was added thionyl chloride (1.10 mL) under ice-cooling, and the mixture was stirred overnight at room temperature. The reaction mixture was concentrated under
30 reduced pressure, and the obtained residue was suspended in ethyl acetate. The precipitate was collected by filtration, and washed with ethyl acetate-hexane to give the title compound (1.64 g).

¹H NMR (300 MHz, DMSO-d₆) δ 2.72 (3H, s), 4.93 (2H, s), 7.86
35 (1H, d, J = 8.1 Hz), 8.44 (1H, dd, J = 8.2, 2.0 Hz), 8.85 (1H,

d, $J = 1.9$ Hz).

[0525]

B) methyl 2-hydroxy-3,4-dimethyl-5-((6-methylpyridin-3-yl)methyl)benzoate

5 To a solution of methyl 2-hydroxy-3,4-dimethyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (2.00 g) in a mixed solvent of DME (30.0 mL)-water (10.0 mL) were added 5-(chloromethyl)-2-methylpyridine hydrochloride (1.22 g), tetrakis(triphenylphosphine)palladium(0) (0.38 g) and sodium
10 carbonate (2.11 g), and the mixture was stirred overnight at 80°C under argon atmosphere. The reaction mixture was allowed to be cooled to room temperature, water and ethyl acetate were added thereto, and the mixture was partitioned. The organic layer was washed with saturated brine, and dried over anhydrous
15 magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (ethyl acetate/hexane) to give the title compound (1.57 g).

MS: $[M-H]^+$ 286.1.

20 [0526]

C) methyl 3,4-dimethyl-5-((6-methylpyridin-3-yl)methyl)-2-(((trifluoromethyl)sulfonyl)oxy)benzoate

To a mixture of methyl 2-hydroxy-3,4-dimethyl-5-((6-methylpyridin-3-yl)methyl)benzoate (1.57 g), sodium hydride
25 (0.26 g) and DMF (32.0 mL) was added N-phenylbis(trifluoromethanesulfonimide) (2.16 g) under ice-cooling, and the mixture was stirred at room temperature for 2 hr. To the reaction mixture was added 1N hydrochloric acid, and the mixture was extracted with ethyl acetate. The organic
30 layer was washed with saturated aqueous sodium bicarbonate solution and saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (ethyl acetate/hexane) to give the title
35 compound (0.76 g).

MS: $[M+H]^+$ 418.1.

[0527]

D) methyl 3,4-dimethyl-5-((6-methylpyridin-3-yl)methyl)-2-vinylbenzoate

5 To a solution of methyl 3,4-dimethyl-5-((6-methylpyridin-3-yl)methyl)-2-(((trifluoromethyl)sulfonyl)oxy)benzoate (0.76 g) in DMF (16.0 mL) were added tributylvinyltin (0.87 g), bis(triphenylphosphine)palladium(II) dichloride (0.06 g) and lithium chloride (0.57 g), and the mixture was stirred at 90°C
10 for 1.5 hr under argon atmosphere. To the reaction mixture was added aqueous potassium fluoride solution, and the precipitated insoluble substance was removed by filtration through Celite. The filtrate was diluted with ethyl acetate, and the mixture was washed with saturated brine. The organic layer was dried
15 over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (ethyl acetate/hexane) to give the title compound (0.50 g).

MS: $[M+H]^+$ 296.2.

20 [0528]

E) methyl 2-formyl-3,4-dimethyl-5-((6-methylpyridin-3-yl)methyl)benzoate

To a solution of methyl 3,4-dimethyl-5-((6-methylpyridin-3-yl)methyl)-2-vinylbenzoate (0.49 g) in a mixed solvent of
25 acetone (6.10 mL)-acetonitrile (6.10 mL)-water (6.10 mL) were added osmium oxide (fixed catalyst I) (0.21 g) and sodium periodate (1.79 g), and the mixture was stirred overnight at room temperature. The reaction mixture was filtered, and the filtrate was extracted with ethyl acetate. The organic layer
30 was washed with saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure to give the title compound (0.47 g) as a crude product. This compound was used in the next step without an additional purification.

35 MS: $[M+H]^+$ 298.1.

[0529]

F) 2-((1S,2S)-2-hydroxycyclohexyl)-4,5-dimethyl-6-((6-methylpyridin-3-yl)methyl)isoindolin-1-one

To a solution of methyl 2-formyl-3,4-dimethyl-5-((6-methylpyridin-3-yl)methyl)benzoate (0.47 g) in THF (9.40 mL) were added (1S,2S)-aminocyclohexanol (0.18 g) and anhydrous magnesium sulfate (0.37 g), and the mixture was stirred at room temperature for 6 hr. The insoluble substance was removed by filtration, the filtrate was concentrated, and the residue was diluted with methanol (9.40 mL)-THF (9.40 mL). Sodium triacetoxyborohydride (0.67 g) was added thereto, and the mixture was stirred overnight at room temperature. The reaction mixture was diluted with ethyl acetate, and the mixture was washed with water and saturated brine. The organic layer was dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (methanol/ethyl acetate) to give the title compound (0.20 g).

¹H NMR (300 MHz, DMSO-d₆) δ 1.22-1.39 (3H, m), 1.47-1.75 (4H, m), 1.89-2.02 (1H, m), 2.20 (3H, s), 2.22 (3H, s), 2.41 (3H, s), 3.61 (1H, dd, J = 10.0, 5.1 Hz), 3.74-3.88 (1H, m), 4.05 (2H, s), 4.35 (2H, s), 4.69 (1H, d, J = 5.5 Hz), 7.14 (1H, d, J = 7.9 Hz), 7.29 (1H, s), 7.35 (1H, dd, J = 7.9, 2.3 Hz), 8.29 (1H, d, J = 1.9 Hz).

X-ray powder diffraction pattern with specific peaks at d value (or d-spacing) = 18.5, 10.3, 9.2, 7.0, 5.3, 5.1, 4.7, 4.4, 4.3 and 4.2 Å.

[0530]

Example 26

1,5-anhydro-2-(6-(4-cyano-3-fluorobenzyl)-4,5-dimethyl-1-oxo-1,3-dihydro-2H-isoindol-2-yl)-2,4-dideoxy-L-threo-pentitol
A) methyl 5-(4-cyano-3-fluorobenzyl)-2-hydroxy-3,4-dimethylbenzoate

To a mixture of methyl 2-hydroxy-3,4-dimethyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (1.07 g), 4-

(chloromethyl)-2-fluorobenzonitrile (0.89 g), 2mol/L aqueous sodium carbonate solution (3.50 mL) and DME (20.0 mL) was added [1,1'-bis(diphenylphosphino)ferrocene]dichloropalladium(II) dichloromethane adduct (0.14 g) under argon atmosphere, and the mixture was stirred overnight at 80°C, and then at room temperature for weekend. To the reaction mixture was added water at room temperature, and the mixture was extracted with ethyl acetate. The organic layer was separated, washed with water and saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (ethyl acetate/hexane) to give the title compound (1.10 g).

MS: $[M+H]^+$ 312.0.

[0531]

B) methyl 5-(4-cyano-3-fluorobenzyl)-3,4-dimethyl-2-vinylbenzoate

A solution of methyl 5-(4-cyano-3-fluorobenzyl)-2-hydroxy-3,4-dimethylbenzoate (1.10 g) in DMF (20.0 mL) was ice-cooled under argon atmosphere, sodium hydride (0.15 g) was added thereto, and the mixture was stirred at room temperature for 30 min. To this reaction mixture was added N-phenylbis(trifluoromethanesulfonimide) (1.51 g) under ice-cooling, and the mixture was stirred at room temperature for 1.5 hr. To the reaction mixture was added 1N hydrochloric acid under ice-cooling, and the mixture was extracted with ethyl acetate. The organic layer was separated, washed with water and saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The obtained crude methyl 5-(4-cyano-3-fluorobenzyl)-3,4-dimethyl-2-[[trifluoromethyl)sulfonyl]oxy]benzoate was used in the next step without an additional purification.

To a mixture of the above-mentioned compound, lithium chloride (1.04 g), tributylvinyltin (1.54 mL) and DMF (20.0 mL) was added trans-dichlorobis(triphenylphosphine)palladium(II) (0.12 g) under argon atmosphere, and the mixture was stirred at

90°C for 2 hr, and then overnight at room temperature. To the reaction mixture was added tributylvinyltin (1.54 mL) under argon atmosphere, and the mixture was stirred at 90°C for 2 hr. To the reaction mixture were added ethyl acetate and 10% aqueous potassium fluoride solution, and the precipitated insoluble substance was removed by filtration through Celite. The filtrate was extracted with ethyl acetate, the organic layer was separated, washed with water and saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (ethyl acetate/hexane) to give the title compound (0.67 g).

MS: $[M+H]^+$ 322.1.

[0532]

C) methyl 5-(4-cyano-3-fluorobenzyl)-2-formyl-3,4-dimethylbenzoate

To a mixture of methyl 5-(4-cyano-3-fluorobenzyl)-3,4-dimethyl-2-vinylbenzoate (0.67 g) and sodium periodate (2.22 g) in acetone (15.0 mL)-acetonitrile (15.0 mL)-water (15.0 mL) was added osmium oxide (fixed catalyst I) (0.26 g), and the mixture was stirred overnight at room temperature under argon atmosphere. The reaction mixture was filtered, and the filtrate was extracted with ethyl acetate. The organic layer was washed with water and saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (ethyl acetate/hexane) to give the title compound (0.25 g).

MS: $[M+H]^+$ 326.1.

[0533]

D) 1,5-anhydro-2-(6-(4-cyano-3-fluorobenzyl)-4,5-dimethyl-1-oxo-1,3-dihydro-2H-isoindol-2-yl)-2,4-dideoxy-L-threo-pentitol

To a solution of methyl 5-(4-cyano-3-fluorobenzyl)-2-formyl-3,4-dimethylbenzoate (0.25 g) in THF (5.00 mL) was added (3S,4S)-3-aminotetrahydro-2H-pyran-4-ol (0.09 g), and the

mixture was stirred overnight at room temperature under argon atmosphere. The reaction mixture was concentrated, and the residue was diluted with acetic acid (5.00 mL). Then, sodium triacetoxyborohydride (0.24 g) was added thereto and the
5 mixture was stirred at room temperature for 2 hr under argon atmosphere. To the reaction mixture was added saturated aqueous sodium bicarbonate, and the mixture was extracted with ethyl acetate. The organic layer was washed with water and saturated brine, and dried over anhydrous magnesium sulfate,
10 and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (ethyl acetate/hexane, methanol/ethyl acetate) to give the title compound (0.16 g).

^1H NMR (300 MHz, CDCl_3) δ 1.73–1.89 (1H, m), 2.08–2.18 (4H, m),
15 2.26 (3H, s), 2.56 (1H, d, $J = 5.7$ Hz), 3.45–3.63 (2H, m),
3.99–4.19 (6H, m), 4.27–4.53 (2H, m), 6.89 (1H, d, $J = 10.0$ Hz),
7.00 (1H, d, $J = 7.9$ Hz), 7.47–7.55 (2H, m).

[0534]

Example 38

20 rac-2-(trans-2-methoxycyclohexyl)-5-methyl-6-(4-(1H-pyrazol-1-yl)benzyl)isoindolin-1-one

To a solution of rac-2-(trans-2-hydroxycyclohexyl)-5-methyl-6-(4-(1H-pyrazol-1-yl)benzyl)isoindolin-1-one (0.06 g) (obtained in the same manner as in Example 8) in DMF (1.20 mL)
25 was added sodium hydride (9.0 mg) under ice-cooling, and the mixture was stirred for 20 min. Methyl iodide (0.05 mL) was added thereto at the same temperature, and the mixture was stirred at room temperature for two nights under nitrogen atmosphere. To this reaction mixture was added sodium hydride
30 (9.0 mg) under ice-cooling, and the mixture was stirred at room temperature for 7 hr. The reaction mixture was poured into water under ice-cooling, and the mixture was extracted with ethyl acetate. The organic layer was separated, washed with saturated brine, and dried over anhydrous magnesium sulfate,
35 and the solvent was evaporated under reduced pressure. The

residue was purified by silica gel column chromatography (ethyl acetate/hexane) and then HPLC (water/acetonitrile) to give the title compound (0.004 g).

¹H NMR (300 MHz, DMSO-d₆) δ 1.29-1.46 (3H, m), 1.63-1.96 (4H, m), 2.19-2.30 (1H, m), 2.31 (3H, s), 3.27 (3H, s), 3.38-3.51 (1H, m), 4.04-4.17 (3H, m), 4.34 (2H, s), 6.44 (1H, t, J = 2.1 Hz), 7.17-7.25 (3H, m), 7.58 (2H, d, J = 8.7 Hz), 7.66-7.74 (2H, m), 7.88 (1H, d, J = 2.4 Hz).

[0535]

10 Example 57

5-ethyl-6-(4-(1-methyl-1H-pyrazol-3-yl)benzyl)-2-(tetrahydrofuran-2-ylmethyl)isoindolin-1-one

A) methyl 4-bromo-2-hydroxybenzoate

To a solution of 4-bromo-2-hydroxybenzoic acid (15.0 g) in methanol (150 mL) was added dropwise thionyl chloride (10.1 mL) under ice-cooling. The reaction solution was stirred at 70°C overnight under argon atmosphere. The reaction solution was concentrated under reduced pressure, and the residue was purified by silica gel column chromatography (ethyl acetate/hexane) to give the title compound (14.2 g).

¹H NMR (300 MHz, DMSO-d₆) δ 3.88 (3H, s), 7.10-7.18 (1H, m), 7.21-7.28 (1H, m), 7.69 (1H, d, J = 8.3 Hz), 10.65 (1H, s).

[0536]

B) methyl 2-hydroxy-4-vinylbenzoate

25 To a solution of methyl 4-bromo-2-hydroxybenzoate (3.0 g) in DMF (50.0 mL) were added tributylvinyltin (6.18 g), bis(triphenylphosphine)palladium(II) chloride (0.46 g) and lithium chloride (4.07 g), and the mixture was stirred at 90°C for 2 hr under argon atmosphere. To the reaction mixture was added aqueous potassium fluoride solution, and the precipitated insoluble substance was removed by filtration through Celite. The filtrate was diluted with ethyl acetate, and the mixture was washed with water and saturated brine. The organic layer was dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by

silica gel column chromatography (ethyl acetate/hexane) to give the title compound. This compound was used in the next step without an additional purification.

[0537]

5 C) methyl 4-ethyl-2-hydroxybenzoate

To a solution of methyl 2-hydroxy-4-vinylbenzoate (2.31 g) in ethanol (25.0 mL) was added palladium-carbon (1.38 g), and the mixture was stirred overnight at room temperature under hydrogen atmosphere. The insoluble substance was removed by
10 filtration, and the filtrate was concentrated under reduced pressure. The obtained residue was purified by NH silica gel column chromatography (ethyl acetate/hexane) to give the title compound (1.43 g).

MS: $[M+H]^+$ 181.1.

15 [0538]

D) methyl 5-bromo-4-ethyl-2-hydroxybenzoate

To a solution of methyl 4-ethyl-2-hydroxybenzoate (1.43 g) in acetic acid (15.0 mL) was added bromine (1.40 g) under ice-cooling, and the mixture was stirred at room temperature
20 for 2 hr under argon atmosphere. To the reaction mixture was added water, the resulting solid was collected by filtration, and dried under reduced pressure to give the title compound (2.21 g) as a mixture with methyl 3,5-dibromo-4-ethyl-2-hydroxybenzoate (2:1). This compound was used in the next step
25 without an additional purification.

[0539]

E) methyl 4-ethyl-2-hydroxy-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate

A mixture of the mixture (1.00 g) of methyl 5-bromo-4-ethyl-2-hydroxybenzoate and methyl 3,5-dibromo-4-ethyl-2-hydroxybenzoate, bis(pinacolato)diboron (1.47 g), potassium acetate (1.14 g), trans-dichlorobis(triphenylphosphine)palladium(II) (0.14 g) and toluene (20.0 mL) was stirred overnight at 100°C under argon
35 atmosphere. The reaction mixture was diluted with ethyl

acetate, and the mixture was washed with saturated brine. The organic layer was dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (ethyl

5 acetate/hexane) to give the title compound (0.82 g). This compound was used in the next step without an additional purification.

MS: $[M+H]^+$ 307.2.

[0540]

10 F) methyl 4-ethyl-2-hydroxy-5-(4-(1-methyl-1H-pyrazol-3-yl)benzyl)benzoate

A mixture of methyl 4-ethyl-2-hydroxy-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (0.41 g), 3-(4-(bromomethyl)phenyl)-1-methyl-1H-pyrazole (0.44 g), sodium
15 carbonate (0.28 g), tetrakis(triphenylphosphine)palladium(0) (0.15 g), DME (15.0 mL) and water (5.0 mL) was stirred overnight at 90°C under argon atmosphere. The reaction mixture was diluted with ethyl acetate, and the mixture was washed with saturated brine. The organic layer was dried over anhydrous
20 sodium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (ethyl acetate/hexane) to give the title compound (0.26 g).

MS: $[M+H]^+$ 351.1.

25 [0541]

G) methyl 4-ethyl-5-(4-(1-methyl-1H-pyrazol-3-yl)benzyl)-2-(((trifluoromethyl)sulfonyl)oxy)benzoate

To a solution of methyl 4-ethyl-2-hydroxy-5-(4-(1-methyl-1H-pyrazol-3-yl)benzyl)benzoate (0.26 g) and N-
30 phenylbis(trifluoromethanesulfonimide) (0.29 g) in DMF (3.00 mL) was added sodium hydride (0.04 g) under ice-cooling, and the mixture was stirred at room temperature for 1 hr under argon atmosphere. The reaction mixture was diluted with ethyl acetate, and the mixture was washed with saturated brine. The
35 organic layer was dried over anhydrous sodium sulfate, and the

solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (ethyl acetate/hexane) to give the title compound (0.32 g).

MS: $[M+H]^+$ 483.1.

5 [0542]

H) methyl 4-ethyl-5-(4-(1-methyl-1H-pyrazol-3-yl)benzyl)-2-vinylbenzoate

A mixture of methyl 4-ethyl-5-(4-(1-methyl-1H-pyrazol-3-yl)benzyl)-2-(((trifluoromethyl)sulfonyl)oxy)benzoate (0.32 g),
10 tributylvinyltin (0.32 g), trans-dichlorobis(triphenylphosphine)palladium(II) (0.02 g), lithium chloride (0.21 g) and DMF (6.00 mL) was stirred at 90°C for 2 hr under argon atmosphere. To the reaction mixture was added aqueous potassium fluoride solution, and the precipitated
15 insoluble substance was removed by filtration. The filtrate was diluted with ethyl acetate, and the mixture was washed with saturated brine. The organic layer was dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by NH silica gel column
20 chromatography (ethyl acetate/hexane) to give the title compound (0.21 g). This compound was used in the next step without an additional purification.

[0543]

I) methyl 4-ethyl-2-formyl-5-(4-(1-methyl-1H-pyrazol-3-yl)benzyl)benzoate
25

A mixture of methyl 4-ethyl-5-(4-(1-methyl-1H-pyrazol-3-yl)benzyl)-2-vinylbenzoate (0.21 g), osmium oxide (fixed catalyst I) (0.08 g) and sodium periodate (0.64 g) in acetone (4.00 mL)-acetonitrile (4.00 mL)-water (4.00 mL) was stirred
30 overnight at room temperature. The insoluble substance was removed by filtration, the filtrate was diluted with ethyl acetate, and the mixture was washed with saturated brine. The organic layer was dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure to give the title
35 compound (0.21 g) as a crude product. This compound was used

in the next step without an additional purification.

[0544]

J) 5-ethyl-6-(4-(1-methyl-1H-pyrazol-3-yl)benzyl)-2-
((tetrahydrofuran-2-yl)methyl)isoindolin-1-one

5 A mixture of methyl 4-ethyl-2-formyl-5-(4-(1-methyl-1H-pyrazol-3-yl)benzyl)benzoate (0.11 g), (tetrahydrofuran-2-yl)methanamine (0.03 g) and anhydrous magnesium sulfate (0.07 g) in THF (3.00 mL) was stirred overnight at room temperature. The insoluble substance was removed by filtration, and the
10 filtrate was concentrated under reduced pressure. The residue was diluted with acetic acid (3.00 mL), sodium triacetoxyborohydride (0.09 g) was added thereto, and the mixture was stirred overnight at room temperature. The reaction mixture was neutralized with saturated aqueous sodium
15 bicarbonate solution, and the mixture was extracted with ethyl acetate. The organic layer was washed with water and saturated brine, and dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by NH silica gel column chromatography (ethyl
20 acetate/hexane), and crystallized from ethyl acetate/hexane to give the title compound (0.03 g).

¹H NMR (300 MHz, DMSO-d₆) δ 1.11 (3H, t, J = 7.6 Hz), 1.47-1.63 (1H, m), 1.74-2.00 (3H, m), 2.69 (2H, q, J = 7.7 Hz), 3.44-3.67 (3H, m), 3.72-3.82 (1H, m), 3.86 (3H, s), 3.98-4.06 (1H, m),
25 4.09 (2H, s), 4.50 (2H, d, J = 4.5 Hz), 6.62 (1H, d, J = 2.3 Hz), 7.15 (2H, d, J = 8.3 Hz), 7.42 (2H, s), 7.63-7.75 (3H, m).

[0545]

Example 96

4-fluoro-5-methoxy-6-(4-(1H-pyrazol-1-yl)benzyl)-2-
30 (tetrahydrofuran-2-ylmethyl)isoindolin-1-one
A) 2-fluoro-3-methoxyphenol

To a solution of 2-fluoro-3-methoxyphenylboronic acid (30.4 g) in THF (300 mL) was added dropwise aqueous hydrogen peroxide (100 mL, 30% wt in water), and the mixture was heated
35 with reflux for 1 hr. The reaction mixture was allowed to be

cooled to room temperature, saturated aqueous sodium sulfite was added thereto, and the mixture was extracted with ethyl acetate. The organic layer was dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure.

5 The residue was purified by combi flash (petroleum ether/ethyl acetate) to give the title compound (24.8 g).

^1H NMR (400 MHz, CDCl_3) δ 3.88 (3H, s), 5.23 (1H, brs), 6.53 (1H, t, $J = 8.4$ Hz), 6.62 (1H, t, $J = 8.4$ Hz), 6.93 (1H, td, $J = 8.4, 2.0$ Hz).

10 [0546]

B) 3-fluoro-2-hydroxy-4-methoxybenzaldehyde

To a solution of 2-fluoro-3-methoxyphenol (22.0 g) and triethylamine (93.9 g) in dichloroethane (250 mL) was added magnesium chloride (71.7 g), and the mixture was stirred at
15 40°C for 1 hr. To this mixture was added paraformaldehyde (46.5 g), and the mixture was stirred for 16 hr. The reaction solution was allowed to be cooled to room temperature, 1N hydrochloric acid was added thereto, and the mixture was extracted with dichloromethane. The organic layer was dried
20 over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by combi flash (petroleum ether/ethyl acetate) to give the title compound (26.0 g).

^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 3.98 (3H, s), 6.64 (1H, dd, $J = 8.8, 6.8$ Hz), 7.32 (1H, dd, $J = 8.8, 1.6$ Hz), 9.77 (1H, d, $J = 2.0$ Hz). One active proton was not observed.

[0547]

C) 3-fluoro-2-hydroxy-4-methoxybenzoic acid

To a mixture of 3-fluoro-2-hydroxy-4-methoxybenzaldehyde
30 (10.0 g) and sodium dihydrogenphosphate (22.9 g) in DMSO (100 mL) and water (25.0 mL) was added dropwise an aqueous solution (30.0 mL) of sodium chlorite (14.5 g), and the mixture was stirred at 20°C for 16 hr. The solvent was evaporated under reduced pressure, the residue was diluted with water, and the
35 mixture was extracted with ethyl acetate (x 6). The organic

layer was washed with saturated brine, and dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure to give the title compound (7.42 g).

¹H NMR (400 MHz, DMSO-d₆) δ 3.89 (3H, s), 6.75 (1H, t, J = 8.4 Hz), 7.59 (1H, dd, J = 9.2, 2.0 Hz). Two active protons were not observed.

[0548]

D) 5-bromo-3-fluoro-2-hydroxy-4-methoxybenzoic acid

To a solution of 3-fluoro-2-hydroxy-4-methoxybenzoic acid (7.30 g) in DMF (70.0 mL) was added NBS, and the mixture was stirred at 25°C for 2 hr. The solvent was evaporated under reduced pressure, and the residue was diluted with ethyl acetate. The mixture was washed with water and saturated brine, and dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure to give the title compound (8.86 g).

¹H NMR (400 MHz, DMSO-d₆) δ 4.00 (3H, s), 7.74 (1H, d, J = 2.0 Hz). Two active protons were not observed and it contained some impurity.

[0549]

E) methyl 5-bromo-3-fluoro-2-hydroxy-4-methoxybenzoate

To a solution of 5-bromo-3-fluoro-2-hydroxy-4-methoxybenzoic acid (0.70 g) in methanol (15.0 mL) was added dropwise thionyl chloride (0.39 mL) under ice-cooling. The reaction solution was stirred at 70°C for 15 hr under argon atmosphere, and then overnight at room temperature. To the reaction mixture was added water, and the mixture was extracted with ethyl acetate. The organic layer was washed with water and saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (ethyl acetate/hexane) to give the title compound (0.52 g).

¹H NMR (300 MHz, CDCl₃) δ 3.97 (3H, s), 4.12 (3H, d, J = 3.0 Hz), 7.82 (1H, d, J = 2.3 Hz), 10.84 (1H, s).

[0550]

F) methyl 3-fluoro-2-hydroxy-4-methoxy-5-[4-(1H-pyrazol-1-yl)benzyl]benzoate

To a solution of methyl 5-bromo-3-fluoro-2-hydroxy-4-methoxybenzoate (0.52 g) in DME (10.0 mL) were added
5 bis(pinacolato)diboron (0.71 g), potassium acetate (0.55 g) and [1,1'-bis(diphenylphosphino)ferrocene]palladium(II) dichloride dichloromethane adduct (0.08 g), and the mixture was stirred at 80°C for 5 hr under argon atmosphere. The reaction mixture was allowed to be cooled to room temperature, water was added
10 thereto, and the mixture was diluted with ethyl acetate. The organic layer was washed with water and saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure to give methyl 3-fluoro-2-hydroxy-4-methoxy-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate as a crude product. To a solution of the above-
15 mentioned crude product in DME (10.0 mL) were added 1-(4-(bromomethyl)phenyl)-1H-pyrazole (0.44 g), [1,1'-bis(diphenylphosphino)ferrocene]palladium(II) dichloride dichloromethane adduct (0.08 g) and 2M aqueous sodium carbonate
20 solution (1.86 mL), and the mixture was stirred at 80°C overnight under argon atmosphere. The reaction mixture was allowed to be cooled to room temperature, water was added thereto, and the mixture was extracted with ethyl acetate. The organic layer was washed with water and saturated brine, and
25 dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (ethyl acetate/hexane) to give the title compound (0.21 g).

¹H NMR (300 MHz, CDCl₃) δ 3.92 (2H, s), 3.93 (3H, s), 3.94-3.97
30 (3H, m), 6.46 (1H, t, J = 2.2 Hz), 7.24 (2H, s), 7.41 (1H, d, J = 2.1 Hz), 7.61 (2H, d, J = 8.5 Hz), 7.72 (1H, d, J = 1.5 Hz), 7.88-7.92 (1H, m), 10.81 (1H, s).

[0551]

G) methyl 2-ethenyl-3-fluoro-4-methoxy-5-[4-(1H-pyrazol-1-yl)benzyl]benzoate
35

To a solution of methyl 3-fluoro-2-hydroxy-4-methoxy-5-[4-(1H-pyrazol-1-yl)benzyl]benzoate (0.21 g) in DMF (5.00 mL) was added sodium hydride (0.03 g) under ice-cooling, and the mixture was stirred at room temperature for 0.5 hr under argon atmosphere. To the reaction mixture was added N-phenylbis(trifluoromethanesulfonimide) (0.23 g), and the mixture was stirred at room temperature for 0.5 hr under argon atmosphere. To the reaction mixture was added 1N hydrochloric acid, and the mixture was extracted with ethyl acetate. The organic layer was washed with water and saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure to give methyl 3-fluoro-4-methoxy-5-[4-(1H-pyrazol-1-yl)benzyl]-2-[[trifluoromethyl)sulfonyl]oxy]benzoate as a crude product.

To a mixture of the obtained crude product, tributylvinyltin (0.26 mL) and lithium chloride (0.19 g) in DMF (5.00 mL) was added bis(triphenylphosphine)palladium(II) dichloride (0.02 g), and the mixture was stirred at 90°C for 2 hr under argon atmosphere. To the reaction mixture was added 10% aqueous potassium fluoride solution, and the precipitated insoluble substance was removed by filtration through Celite. To the filtrate was added water, and the mixture was extracted with ethyl acetate. The organic layer was washed with water and saturated brine, and dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (ethyl acetate/hexane) to give the title compound (0.12 g).

MS: $[M+H]^+$ 367.2.

[0552]

H) 4-fluoro-5-methoxy-6-(4-(1H-pyrazol-1-yl)benzyl)-2-(tetrahydrofuran-2-ylmethyl)isoindolin-1-one

To a solution of methyl 2-ethenyl-3-fluoro-4-methoxy-5-[4-(1H-pyrazol-1-yl)benzyl]benzoate (0.12 g) in a mixed solvent of acetone (2.00 mL)-acetonitrile (2.00 mL)-water (2.00 mL) were added osmium oxide (fixed catalyst I) (0.04 g) and sodium

periodate (0.35 g), and the mixture was stirred at room temperature for 3 hr under argon atmosphere. The reaction mixture was filtered, and the filtrate was concentrated under reduced pressure. To the residue was added water, and the mixture was extracted with ethyl acetate. The organic layer was washed with saturated brine, and dried over anhydrous sodium sulfate, and the solvent was evaporated under reduced pressure to give methyl 3-fluoro-2-formyl-4-methoxy-5-[4-(1H-pyrazol-1-yl)benzyl]benzoate as a crude product. The obtained crude product (20% v/v) was dissolved in THF (2.00 mL), (tetrahydrofuran-2-yl)methanamine (0.006 g) was added thereto, and the mixture was stirred at room temperature for 1 hr. The reaction solution was concentrated under reduced pressure, and the residue was diluted with acetic acid (1.00 mL). Sodium triacetoxymethylborohydride (0.03 g) was added thereto, and the mixture was stirred overnight at room temperature. The reaction mixture was neutralized with saturated aqueous sodium hydrogencarbonate solution, and the mixture was extracted with ethyl acetate. The organic layer was washed with water and saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was purified by HPLC (water/acetonitrile, containing 0.1% TFA). The fractions were combined, saturated aqueous sodium hydrogencarbonate solution was added thereto, and the mixture was extracted with ethyl acetate. The organic layer was dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure to give the title compound (2.90 mg).

MS: $[M+H]^+$ 422.2.

[0553]

Example 113

rac-3-fluoro-4-((2-(trans-4-hydroxytetrahydro-2H-pyran-3-yl)-6,7-dimethyl-3-oxoisindolin-5-yl)methyl)benzamide

To a mixture of rac-3-fluoro-4-((2-(trans-4-hydroxytetrahydro-2H-pyran-3-yl)-6,7-dimethyl-3-oxoisindolin-

5-yl)methyl)benzonitrile (0.08 g) and potassium carbonate (0.08 g) in DMSO (2.00 mL) was added 35% aqueous hydrogen peroxide (0.17 mL) under ice-cooling, and the mixture was stirred at room temperature for 3 hr. The reaction solution was diluted
5 with water, and the mixture was extracted with ethyl acetate-THF. The organic layer was washed with saturated brine, and dried over anhydrous magnesium sulfate, and the solvent was evaporated under reduced pressure. The residue was washed with ethyl acetate to give the title compound (0.07 g).

10 MS: $[M+H]^+$ 413.2.

[0554]

The compounds of Examples 27 to 37, 39 to 56, 58 to 95, 97 to 112 and 114 to 168 in Table 1 were synthesized according to the method shown in the above-mentioned Examples or a method
15 analogous thereto. The Example compounds are shown in Table 1. MS in the tables means actual measured value.

[0555]

Table 1-1

Ex. No.	IUPAC Name	structure formula	MS
1	rac-2-(trans-2-hydroxycyclohexyl)-6-((6-(1-methyl-1H-pyrazol-4-yl)pyridin-3-yl)methyl)isoindolin-1-one		403.2
2	rac-5-chloro-2-(trans-2-hydroxycyclohexyl)-6-(4-(1H-pyrazol-1-yl)benzyl)isoindolin-1-one		422.1
3	rac-2-(trans-2-hydroxycyclohexyl)-5-methoxy-6-(4-(1H-pyrazol-1-yl)benzyl)isoindolin-1-one		418.2
4	2-((1S,2S)-2-hydroxycyclopentyl)-5-methyl-6-(4-(1H-pyrazol-1-yl)benzyl)isoindolin-1-one		388.2
5	rac-6-((6-(1,3-dimethyl-1H-pyrazol-4-yl)pyridin-3-yl)methyl)-2-(trans-2-hydroxycyclohexyl)-5-methylisoindolin-1-one		431.2

[0556]

Table 1-2

Ex. No.	IUPAC Name	structure formula	MS
6	rac-2-(trans-2-hydroxycyclohexyl)-5-methyl-6-((2'-methyl-2,4'-bipyridin-5-yl)methyl)isoindolin-1-one		428.2
7	3-fluoro-2-(5-methyl-1-oxo-6-(4-(1H-pyrazol-1-yl)benzyl)-1,3-dihydro-2H-isoindol-2-yl)benzonitrile		423.1
8	5-methyl-6-(4-(1H-pyrazol-1-yl)benzyl)-2-(tetrahydro-2H-pyran-4-yl)isoindolin-1-one		388.2
9	rac-5-cyclopropyl-2-(trans-2-hydroxycyclopentyl)-6-(4-(1H-pyrazol-1-yl)benzyl)isoindolin-1-one		414.2
10	rac-4-chloro-2-(trans-2-hydroxycyclohexyl)-6-(4-(1H-pyrazol-1-yl)benzyl)-5-(trifluoromethyl)isoindolin-1-one		490.1






[0557]

Table 1-3

Ex. No.	IUPAC Name	structure formula	MS
11	rac-2-(trans-2-hydroxycyclohexyl)-4-methyl-1-oxo-6-(4-(1H-pyrazol-1-yl)benzyl)isoindoline-5-carbonitrile		427.2
12	rac-4-chloro-2-(trans-2-hydroxycyclohexyl)-5-methoxy-6-(4-(1H-pyrazol-1-yl)benzyl)isoindolin-1-one		452.1
13	2-((1S,2S)-2-hydroxycyclopentyl)-4,5-dimethyl-6-(4-(1-methyl-1H-pyrazol-3-yl)benzyl)isoindolin-1-one		416.2
14	4,5-dimethyl-6-(4-(1-methyl-1H-1,2,3-triazol-4-yl)benzyl)-2-((2S)-tetrahydrofuran-2-ylmethyl)isoindolin-1-one		417.1
15	2-(2-hydroxy-2-methylpropyl)-4,5-dimethyl-6-(4-(1H-pyrazol-1-yl)benzyl)isoindolin-1-one		390.1






[0558]

Table 1-4

Ex. No.	IUPAC Name	structure formula	MS
16	6-(2-fluoro-4-(1H-pyrazol-1-yl)benzyl)-2-((1S,2S)-2-hydroxycyclopentyl)-4,5-dimethylisoindolin-1-one		420.1
17	1,5-anhydro-2-(6-(4-chlorobenzyl)-4,5-dimethyl-1-oxo-1,3-dihydro-2H-isoindol-2-yl)-2,4-dideoxy-L-threo-pentitol		386.0
18	2-[(3S,4S)-4-hydroxytetrahydro-2H-pyran-3-yl]-6-(4-methoxybenzyl)-4,5-dimethyl-2,3-dihydro-1H-isoindol-1-one Alias; 1,5-anhydro-2,4-dideoxy-2-(6-(4-methoxybenzyl)-4,5-dimethyl-1-oxo-1,3-dihydro-2H-isoindol-2-yl)-L-threo-pentitol		382.2
19	1,5-anhydro-2,4-dideoxy-2-(6-(3-fluoro-4-(methylcarbamoyl)benzyl)-4,5-dimethyl-1-oxo-1,3-dihydro-2H-isoindol-2-yl)-L-threo-pentitol		427.1
20	4-fluoro-2-[(3S,4S)-4-hydroxytetrahydro-2H-pyran-3-yl]-5-methyl-6-[4-(1H-pyrazol-1-yl)benzyl]-2,3-dihydro-1H-isoindol-1-one Alias; 1,5-anhydro-2,4-dideoxy-2-(4-fluoro-5-methyl-1-oxo-6-(4-(1H-pyrazol-1-yl)benzyl)-1,3-dihydro-2H-isoindol-2-yl)-L-threo-pentitol		422.1

[0559]

Table 1-5

Ex. No.	IUPAC Name	structure formula	MS
21	1,5-anhydro-2,4-dideoxy-2-(5-methyl-1-oxo-6-(4-(1H-pyrazol-1-yl)benzyl)-1,3-dihydro-2H-isoindol-2-yl)-L-threo-pentitol		404.1
22	1,5-anhydro-2,4-dideoxy-2-(6-(4-(difluoromethoxy)benzyl)-4,5-dimethyl-1-oxo-1,3-dihydro-2H-isoindol-2-yl)-L-threo-pentitol		418.0
23	1,5-anhydro-2,4-dideoxy-2-(6-(4-fluoro-3-methoxybenzyl)-4,5-dimethyl-1-oxo-1,3-dihydro-2H-isoindol-2-yl)-L-threo-pentitol		400.1
24	1,5-anhydro-2-(4-chloro-6-(4-methoxybenzyl)-5-methyl-1-oxo-1,3-dihydro-2H-isoindol-2-yl)-2,4-dideoxy-L-threo-pentitol		402.1
25	2-((1S,2S)-2-hydroxycyclohexyl)-4,5-dimethyl-6-((6-methylpyridin-3-yl)methyl)isoindolin-1-one		365.2

[0560]

Table 1-6

Ex. No.	IUPAC Name	structure formula	MS
26	1,5-anhydro-2-(6-(4-cyano-3-fluorobenzyl)-4,5-dimethyl-1-oxo-1,3-dihydro-2H-isoindol-2-yl)-2,4-dideoxy-L-threo-pentitol		395.1
27	2-(2-fluorophenyl)-6-((6-(1-methyl-1H-pyrazol-4-yl)pyridin-3-yl)methyl)isoindolin-1-one		399.1
28	rac-2-(trans-2-hydroxycyclohexyl)-4,5-dimethyl-6-((6-(1-methyl-1H-pyrazol-4-yl)pyridin-3-yl)methyl)isoindolin-1-one		431.2
29	rac-2-(trans-2-hydroxycyclohexyl)-4,5-dimethyl-6-(4-(1H-pyrazol-1-yl)benzyl)isoindolin-1-one		416.2
30	rac-2-(trans-2-hydroxycyclohexyl)-4-methyl-6-(4-(1H-pyrazol-1-yl)benzyl)isoindolin-1-one		402.2






[0561]

Table 1-7

Ex. No.	IUPAC Name	structure formula	MS
31	rac-2-(trans-2-hydroxycyclopentyl)-4-methyl-6-(4-(1H-pyrazol-1-yl)benzyl)isoindolin-1-one		388.1
32	rac-2-(trans-2-hydroxycyclohexyl)-5-methyl-6-(4-(1H-pyrazol-1-yl)benzyl)isoindolin-1-one		402.2
33	rac-2-(trans-2-hydroxycyclopentyl)-5-methyl-6-(4-(1H-pyrazol-1-yl)benzyl)isoindolin-1-one		388.2
34	rac-2-(trans-2-hydroxycyclopentyl)-4,5-dimethyl-6-(4-(1H-pyrazol-1-yl)benzyl)isoindolin-1-one		402.1
35	rac-2-(trans-4-hydroxytetrahydro-2H-pyran-3-yl)-4,5-dimethyl-6-(4-(1H-pyrazol-1-yl)benzyl)-2,3-dihydro-1H-isoindol-1-one		418.2






[0562]

Table 1-8

Ex. No.	IUPAC Name	structure formula	MS
36	5-methyl-6-(4-(1H-pyrazol-1-yl)benzyl)-2-(tetrahydrofuran-2-ylmethyl)isoindolin-1-one		388.2
37	2-(1-(hydroxymethyl)cyclopentyl)-5-methyl-6-(4-(1H-pyrazol-1-yl)benzyl)isoindolin-1-one		402.2
38	rac-2-(trans-2-methoxycyclohexyl)-5-methyl-6-(4-(1H-pyrazol-1-yl)benzyl)isoindolin-1-one		416.2
39	2-((1R,2R)-2-hydroxycyclopentyl)-5-methyl-6-(4-(1H-pyrazol-1-yl)benzyl)isoindolin-1-one		388.2
40	rac-6-((6-(1,3-dimethyl-1H-pyrazol-4-yl)pyridin-3-yl)methyl)-2-(trans-2-hydroxycyclopentyl)-5-methylisoindolin-1-one		417.1

[0563]

Table 1-9

Ex. No.	IUPAC Name	structure formula	MS
41	rac-2-(trans-2-hydroxycyclopentyl)-5-methyl-6-((2'-methyl-2,4'-bipyridin-5-yl)methyl)isoindolin-1-one		414.2
42	rac-2-(trans-2-hydroxycyclohexyl)-5-methyl-6-((6-(1-methyl-1H-pyrazol-3-yl)pyridin-3-yl)methyl)isoindolin-1-one		417.1
43	rac-2-(trans-2-hydroxycyclopentyl)-5-methyl-6-((6-(1-methyl-1H-pyrazol-3-yl)pyridin-3-yl)methyl)isoindolin-1-one		403.2
44	rac-6-(2,4-difluorobenzyl)-2-(trans-2-hydroxycyclohexyl)-5-methylisoindolin-1-one		372.2
45	rac-6-(2,4-difluorobenzyl)-2-(trans-2-hydroxycyclopentyl)-5-methylisoindolin-1-one		358.1

[0564]

Table 1-10

Ex. No.	IUPAC Name	structure formula	MS
46	2-((1R,2S)-2-hydroxycyclopentyl)-5-methyl-6-(4-(1H-pyrazol-1-yl)benzyl)isoindolin-1-one		388.2
47	rac-2-(trans-2-hydroxycyclohexyl)-5-methyl-6-(4-(1-methyl-1H-1,2,3-triazol-4-yl)benzyl)isoindolin-1-one		417.1
48	rac-2-(trans-2-hydroxycyclopentyl)-5-methyl-6-(4-(1-methyl-1H-1,2,3-triazol-4-yl)benzyl)isoindolin-1-one		403.2
49	5-methyl-6-(4-(1-methyl-1H-1,2,3-triazol-4-yl)benzyl)-2-(tetrahydrofuran-2-ylmethyl)isoindolin-1-one		403.2
50	5-methyl-6-(4-(1H-pyrazol-1-yl)benzyl)-2-(tetrahydro-2H-pyran-3-yl)isoindolin-1-one		388.2






[0565]

Table 1-11

Ex. No.	IUPAC Name	structure formula	MS
51	rac-2-(trans-2-hydroxycyclopentyl)-5-methyl-6-(4-(1-methyl-1H-pyrazol-3-yl)benzyl)isoindolin-1-one		402.1
52	5-methyl-6-(4-(1-methyl-1H-pyrazol-3-yl)benzyl)-2-(tetrahydrofuran-2-ylmethyl)isoindolin-1-one		402.1
53	rac-6-(2-fluoro-4-(1-methyl-1H-1,2,3-triazol-4-yl)benzyl)-2-(trans-2-hydroxycyclopentyl)-5-methylisoindolin-1-one		421.1
54	6-(2-fluoro-4-(1-methyl-1H-1,2,3-triazol-4-yl)benzyl)-5-methyl-2-(tetrahydrofuran-2-ylmethyl)isoindolin-1-one		421.2
55	rac-6-(4-(1,3-dimethyl-1H-pyrazol-4-yl)benzyl)-2-(trans-2-hydroxycyclopentyl)-5-methylisoindolin-1-one		416.2

[0566]

Table 1-12

Ex. No.	IUPAC Name	structure formula	MS
56	6-(4-(1,3-dimethyl-1H-pyrazol-4-yl)benzyl)-5-methyl-2-(tetrahydrofuran-2-ylmethyl)isoindolin-1-one		416.2
57	5-ethyl-6-(4-(1-methyl-1H-pyrazol-3-yl)benzyl)-2-(tetrahydrofuran-2-ylmethyl)isoindolin-1-one		416.1
58	5-ethyl-6-(4-(1H-pyrazol-1-yl)benzyl)-2-(tetrahydrofuran-2-ylmethyl)isoindolin-1-one		402.1
59	rac-5-ethyl-2-(trans-2-hydroxycyclopentyl)-6-(4-(1H-pyrazol-1-yl)benzyl)isoindolin-1-one		402.1
60	5-chloro-6-(4-(1-methyl-1H-1,2,3-triazol-4-yl)benzyl)-2-(tetrahydrofuran-2-ylmethyl)isoindolin-1-one		423.1






[0567]

Table 1-13

Ex. No.	IUPAC Name	structure formula	MS
61	5-cyclopropyl-6-(4-(1-methyl-1H-1,2,3-triazol-4-yl)benzyl)-2-(tetrahydrofuran-2-ylmethyl)isoindolin-1-one		429.2
62	5-cyclopropyl-6-(4-(1H-pyrazol-1-yl)benzyl)-2-(tetrahydrofuran-2-ylmethyl)isoindolin-1-one		414.1
63	rac-5-ethyl-2-(trans-2-hydroxycyclopentyl)-6-(4-(1-methyl-1H-pyrazol-3-yl)benzyl)isoindolin-1-one		416.2
64	rac-5-cyclopropyl-2-(trans-2-hydroxycyclopentyl)-6-(4-(1-methyl-1H-1,2,3-triazol-4-yl)benzyl)isoindolin-1-one		429.2
65	rac-6-(2-fluoro-4-(1-methyl-1H-pyrazol-3-yl)benzyl)-2-(trans-2-hydroxycyclopentyl)-5-methylisoindolin-1-one		420.1

[0568]

Table 1-14

Ex. No.	IUPAC Name	structure formula	MS
66	6-(2-fluoro-4-(1-methyl-1H-pyrazol-3-yl)benzyl)-5-methyl-2-(tetrahydrofuran-2-ylmethyl)isoindolin-1-one		420.1
67	rac-2-(trans-2-hydroxycyclopentyl)-5-methyl-6-(4-(1-methyl-1H-pyrazol-4-yl)benzyl)isoindolin-1-one		402.1
68	5-methyl-6-(4-(1-methyl-1H-pyrazol-4-yl)benzyl)-2-(tetrahydrofuran-2-ylmethyl)isoindolin-1-one		402.1
69	rac-2-(trans-2-hydroxycyclopentyl)-4,5-dimethyl-6-(4-(1-methyl-1H-pyrazol-4-yl)benzyl)isoindolin-1-one		416.2
70	6-(2-fluoro-4-(1-methyl-1H-pyrazol-3-yl)benzyl)-4,5-dimethyl-2-(tetrahydrofuran-2-ylmethyl)isoindolin-1-one		434.2

[0569]

Table 1-15

Ex. No.	IUPAC Name	structure formula	MS
71	rac-6-(2-fluoro-4-(1-methyl-1H-pyrazol-3-yl)benzyl)-2-(trans-2-hydroxycyclopentyl)-4,5-dimethylisoindolin-1-one		434.2
72	4-fluoro-5-methyl-6-(4-(1H-pyrazol-1-yl)benzyl)-2-(tetrahydrofuran-2-ylmethyl)isoindolin-1-one		406.1
73	rac-5-chloro-2-(trans-2-hydroxycyclopentyl)-6-(4-(1H-pyrazol-1-yl)benzyl)isoindolin-1-one		408.1
74	5-chloro-6-(4-(1H-pyrazol-1-yl)benzyl)-2-(tetrahydrofuran-2-ylmethyl)isoindolin-1-one		408.1
75	rac-5-chloro-2-(trans-2-hydroxycyclohexyl)-6-(4-(1-methyl-1H-pyrazol-4-yl)benzyl)isoindolin-1-one		436.1

[0570]

Table 1-16

Ex. No.	IUPAC Name	structure formula	MS
76	rac-5-chloro-2-(trans-2-hydroxycyclopentyl)-6-(4-(1-methyl-1H-pyrazol-4-yl)benzyl)isoindolin-1-one		422.0
77	5-chloro-6-(4-(1-methyl-1H-pyrazol-4-yl)benzyl)-2-(tetrahydrofuran-2-ylmethyl)isoindolin-1-one		422.0
78	rac-4-fluoro-2-(trans-2-hydroxycyclohexyl)-5-methyl-6-(4-(1H-pyrazol-1-yl)benzyl)isoindolin-1-one		420.1
79	rac-4-fluoro-2-(trans-2-hydroxycyclopentyl)-5-methyl-6-(4-(1H-pyrazol-1-yl)benzyl)isoindolin-1-one		406.1
80	4,5-dimethyl-6-(4-(1H-pyrazol-1-yl)benzyl)-2-(tetrahydrofuran-2-ylmethyl)isoindolin-1-one		402.1

[0571]

Table 1-17

Ex. No.	IUPAC Name	structure formula	MS
81	4-methyl-1-oxo-6-(4-(1H-pyrazol-1-yl)benzyl)-2-(tetrahydrofuran-2-ylmethyl)isoindoline-5-carbonitrile		413.2
82	rac-2-(trans-2-hydroxycyclohexyl)-4,5-dimethyl-6-(4-(1-methyl-1H-pyrazol-3-yl)benzyl)isoindolin-1-one		430.2
83	rac-2-(trans-2-hydroxycyclopentyl)-4,5-dimethyl-6-(4-(1-methyl-1H-pyrazol-3-yl)benzyl)isoindolin-1-one		416.2
84	4,5-dimethyl-6-(4-(1-methyl-1H-pyrazol-4-yl)benzyl)-2-(tetrahydrofuran-2-ylmethyl)isoindolin-1-one		416.2
85	4,5-dimethyl-6-(4-(1-methyl-1H-pyrazol-3-yl)benzyl)-2-(tetrahydrofuran-2-ylmethyl)isoindolin-1-one		416.2

[0572]

Table 1-18 ■

Ex. No.	IUPAC Name	structure formula	MS
86	4-chloro-6-(4-(1H-pyrazol-1-yl)benzyl)-2-(tetrahydrofuran-2-ylmethyl)-5-(trifluoromethyl)isoindolin-1-one		476.1
87	rac-2-(trans-2-hydroxycyclopentyl)-4-methyl-1-oxo-6-(4-(1H-pyrazol-1-yl)benzyl)isoindoline-5-carbonitrile		413.2
88	rac-4-chloro-2-(trans-2-hydroxycyclopentyl)-6-(4-(1H-pyrazol-1-yl)benzyl)-5-(trifluoromethyl)isoindolin-1-one		476.2
89	rac-2-(trans-2-hydroxycyclohexyl)-4,5-dimethyl-6-(4-(1-methyl-1H-pyrazol-4-yl)benzyl)isoindolin-1-one		430.2
90	4-chloro-5-methoxy-6-(4-(1H-pyrazol-1-yl)benzyl)-2-(tetrahydrofuran-2-ylmethyl)isoindolin-1-one		438.1

[0573]

Table 1-19

Ex. No.	IUPAC Name	structure formula	MS
91	rac-2-(trans-2-hydroxycyclopentyl)-5-methyl-6-(4-(6-methylpyridazin-4-yl)benzyl)isoindolin-1-one		414.2
92	rac-4-chloro-2-(trans-2-hydroxycyclopentyl)-5-methoxy-6-(4-(1H-pyrazol-1-yl)benzyl)isoindolin-1-one		438.1
93	rac-4-chloro-2-(trans-2-hydroxycyclohexyl)-5-methyl-6-(4-(1H-pyrazol-1-yl)benzyl)isoindolin-1-one		436.1
94	rac-6-(4-(1,3-dimethyl-1H-pyrazol-4-yl)-2-fluorobenzyl)-2-(trans-2-hydroxycyclohexyl)-4,5-dimethylisoindolin-1-one		462.2
95	rac-6-(4-(1,3-dimethyl-1H-pyrazol-4-yl)-2-fluorobenzyl)-2-(trans-2-hydroxycyclopentyl)-4,5-dimethylisoindolin-1-one		448.1






[0574]

Table 1-20

Ex. No.	IUPAC Name	structure formula	MS
96	4-fluoro-5-methoxy-6-(4-(1H-pyrazol-1-yl)benzyl)-2-(tetrahydrofuran-2-ylmethyl)isoindolin-1-one		422.1
97	rac-4-fluoro-2-(trans-2-hydroxycyclohexyl)-5-methoxy-6-(4-(1H-pyrazol-1-yl)benzyl)isoindolin-1-one		436.2
98	rac-4-fluoro-2-(trans-2-hydroxycyclopentyl)-5-methoxy-6-(4-(1H-pyrazol-1-yl)benzyl)isoindolin-1-one		422.1
99	rac-2-(trans-2-hydroxycyclohexyl)-5-methyl-6-(4-(2-methylpyridin-4-yl)benzyl)isoindolin-1-one		427.2
100	rac-2-(trans-2-hydroxycyclopentyl)-5-methyl-6-(4-(2-methylpyridin-4-yl)benzyl)isoindolin-1-one		413.2

[0575]

Table 1-21

Ex. No.	IUPAC Name	structure formula	MS
101	5-methyl-6-(4-(2-methylpyridin-4-yl)benzyl)-2-(tetrahydrofuran-2-ylmethyl)isoindolin-1-one		413.2
102	rac-4-chloro-2-(trans-2-hydroxycyclopentyl)-5-methyl-6-(4-(1H-pyrazol-1-yl)benzyl)isoindolin-1-one		422.1
103	6-(2-fluoro-4-(1-methyl-1H-1,2,3-triazol-4-yl)benzyl)-4,5-dimethyl-2-(tetrahydrofuran-2-ylmethyl)isoindolin-1-one		435.1
104	2-(2-hydroxy-2-methylpropyl)-5-methyl-6-(4-(1H-pyrazol-1-yl)benzyl)isoindolin-1-one		376.1
105	5-ethyl-6-(4-(1-methyl-1H-1,2,3-triazol-4-yl)benzyl)-2-(tetrahydrofuran-2-ylmethyl)isoindolin-1-one		417.1

[0576]

Table 1-22

Ex. No.	IUPAC Name	structure formula	MS
106	rac-5-ethyl-2-(trans-2-hydroxycyclopentyl)-6-(4-(1-methyl-1H-1,2,3-triazol-4-yl)benzyl)isoindolin-1-one		417.1
107	4-fluoro-2-((1S,2S)-2-hydroxycyclopentyl)-5-methyl-6-(4-(1H-pyrazol-1-yl)benzyl)isoindolin-1-one		406.1
108	4,5-dimethyl-6-(4-(1-methyl-1H-pyrazol-3-yl)benzyl)-2-((2R)-tetrahydrofuran-2-ylmethyl)isoindolin-1-one		416.2
109	4,5-dimethyl-6-(4-(1-methyl-1H-pyrazol-3-yl)benzyl)-2-((2S)-tetrahydrofuran-2-ylmethyl)isoindolin-1-one		416.2
110	4,5-dimethyl-6-(4-(1-methyl-1H-1,2,3-triazol-4-yl)benzyl)-2-((2R)-tetrahydrofuran-2-ylmethyl)isoindolin-1-one		417.1

[0577]

Table 1-23

Ex. No.	IUPAC Name	structure formula	MS
111	4-fluoro-2-((1S,2S)-2-hydroxycyclopentyl)-5-methyl-6-(4-(1-methyl-1H-pyrazol-3-yl)benzyl)isoindolin-1-one		420.2
112	rac-3-fluoro-4-((2-(trans-4-hydroxytetrahydro-2H-pyran-3-yl)-6,7-dimethyl-3-oxo-2,3-dihydro-1H-isoindol-5-yl)methyl)benzonitrile		395.2
113	rac-3-fluoro-4-((2-(trans-4-hydroxytetrahydro-2H-pyran-3-yl)-6,7-dimethyl-3-oxo-2,3-dihydro-1H-isoindol-5-yl)methyl)benzamide		413.2
114	2-(2-hydroxy-2-methylpropyl)-4,5-dimethyl-6-(4-(1-methyl-1H-pyrazol-3-yl)benzyl)isoindolin-1-one		404.2
115	6-(2-fluoro-4-(1H-pyrazol-1-yl)benzyl)-4,5-dimethyl-2-((2S)-tetrahydrofuran-2-ylmethyl)isoindolin-1-one		420.2

[0578]

Table 1-24

Ex. No.	IUPAC Name	structure formula	MS
116	5-ethyl-2-[(3S,4S)-4-hydroxytetrahydro-2H-pyran-3-yl]-6-[4-(1-methyl-1H-pyrazol-3-yl)benzyl]-2,3-dihydro-1H-isoindol-1-one Alias; 1,5-anhydro-2,4-dideoxy-2-(5-ethyl-6-(4-(1-methyl-1H-pyrazol-3-yl)benzyl)-1-oxo-1,3-dihydro-2H-isoindol-2-yl)-L-threo-pentitol		432.1
117	5-ethyl-2-[(3R,4R)-4-hydroxytetrahydro-2H-pyran-3-yl]-6-[4-(1-methyl-1H-pyrazol-3-yl)benzyl]-2,3-dihydro-1H-isoindol-1-one Alias; 1,5-anhydro-2,4-dideoxy-2-(5-ethyl-6-(4-(1-methyl-1H-pyrazol-3-yl)benzyl)-1-oxo-1,3-dihydro-2H-isoindol-2-yl)-D-threo-pentitol		432.1
118	5-ethyl-2-(2-hydroxy-2-methylpropyl)-6-(4-(1-methyl-1H-1,2,3-triazol-4-yl)benzyl)isoindolin-1-one		405.1
119	5-ethyl-2-((1S,2S)-2-hydroxycyclopentyl)-6-(4-(1-methyl-1H-1,2,3-triazol-4-yl)benzyl)isoindolin-1-one		417.0
120	5-ethyl-2-((1S,2S)-2-hydroxycyclopentyl)-6-(4-(1-methyl-1H-pyrazol-3-yl)benzyl)isoindolin-1-one		416.1

[0579]

Table 1-25

Ex. No.	IUPAC Name	structure formula	MS
121	6-(2-fluoro-4-(1H-pyrazol-1-yl)benzyl)-2-(2-hydroxy-2-methylpropyl)-4,5-dimethylisoindolin-1-one		408.1
122	2-((1S,2S)-2-hydroxycyclopentyl)-6-(4-methoxybenzyl)-4,5-dimethylisoindolin-1-one		366.1
123	2-fluoro-4-((2-((1S,2S)-2-hydroxycyclopentyl)-6,7-dimethyl-3-oxo-2,3-dihydro-1H-isoindol-5-yl)methyl)-N-methylbenzamide		411.1
124	3-fluoro-4-((2-[(3S,4S)-4-hydroxytetrahydro-2H-pyran-3-yl]-6,7-dimethyl-3-oxo-2,3-dihydro-1H-isoindol-5-yl)methyl)benzonitrile Alias; 1,5-anhydro-2-(6-(4-cyano-2-fluorobenzyl)-4,5-dimethyl-1-oxo-1,3-dihydro-2H-isoindol-2-yl)-2,4-dideoxy-L-threo-pentitol		395.1
125	3-fluoro-4-((2-((1S,2S)-2-hydroxycyclopentyl)-6,7-dimethyl-3-oxo-2,3-dihydro-1H-isoindol-5-yl)methyl)benzonitrile		379.2





[0580]

Table 1-26

Ex. No.	IUPAC Name	structure formula	MS
126	3-fluoro-4-((2-(2-hydroxy-2-methylpropyl)-6,7-dimethyl-3-oxo-2,3-dihydro-1H-isoindol-5-yl)methyl)benzonitrile		367.1
127	4-fluoro-2-(2-hydroxy-2-methylpropyl)-5-methyl-6-(4-(1H-pyrazol-1-yl)benzyl)isoindolin-1-one		394.1
128	5-ethyl-2-[(3S,4S)-4-hydroxytetrahydro-2H-pyran-3-yl]-6-[4-(1H-pyrazol-1-yl)benzyl]-2,3-dihydro-1H-isoindol-1-one Alias; 1,5-anhydro-2,4-dideoxy-2-(5-ethyl-1-oxo-6-(4-(1H-pyrazol-1-yl)benzyl)-1,3-dihydro-2H-isoindol-2-yl)-L-threo-pentitol		418.0
129	5-ethyl-2-(2-hydroxy-2-methylpropyl)-6-(4-(1H-pyrazol-1-yl)benzyl)isoindolin-1-one		390.1
130	4-chloro-2-[(3S,4S)-4-hydroxytetrahydro-2H-pyran-3-yl]-5-methyl-6-[4-(1H-pyrazol-1-yl)benzyl]-2,3-dihydro-1H-isoindol-1-one Alias; 1,5-anhydro-2-(4-chloro-5-methyl-1-oxo-6-(4-(1H-pyrazol-1-yl)benzyl)-1,3-dihydro-2H-isoindol-2-yl)-2,4-dideoxy-L-threo-pentitol		438.0

[0581]

Table 1-27

Ex. No.	IUPAC Name	structure formula	MS
131	4-chloro-2-((1S,2S)-2-hydroxycyclopentyl)-5-methyl-6-(4-(1H-pyrazol-1-yl)benzyl)isoindolin-1-one		422.0
132	4-chloro-2-(2-hydroxy-2-methylpropyl)-5-methyl-6-(4-(1H-pyrazol-1-yl)benzyl)isoindolin-1-one		410.0
133	2-(3-hydroxy-3-methylbutan-2-yl)-5-methyl-6-(4-(1H-pyrazol-1-yl)benzyl)isoindolin-1-one		390.1
134	2-((1S,2S)-2-hydroxycyclopentyl)-4,5-dimethyl-6-((6-methylpyridin-3-yl)methyl)isoindolin-1-one		351.2
135	2-(2-fluorophenyl)-5-methyl-6-(4-(1H-pyrazol-1-yl)benzyl)isoindolin-1-one		398.0






[0582]

Table 1-28

Ex. No.	IUPAC Name	structure formula	MS
136	4,5-dimethyl-6-((6-(1H-pyrazol-1-yl)pyridin-3-yl)methyl)-2-((2S)-tetrahydrofuran-2-ylmethyl)isoindolin-1-one		403.1
137	5-ethyl-2-(2-hydroxy-2-methylpropyl)-6-(4-(1-methyl-1H-pyrazol-3-yl)benzyl)isoindolin-1-one		404.1
138	2-fluoro-4-((2-(2-hydroxy-2-methylpropyl)-6,7-dimethyl-3-oxo-2,3-dihydro-1H-isoindol-5-yl)methyl)-N-methylbenzamide		399.0
139	4-((6,7-dimethyl-3-oxo-2-((2S)-tetrahydrofuran-2-ylmethyl)-2,3-dihydro-1H-isoindol-5-yl)methyl)-2-fluoro-N-methylbenzamide		411.1
140	4-fluoro-2-[(3S,4S)-4-hydroxytetrahydro-2H-pyran-3-yl]-5-methyl-6-[4-(1-methyl-1H-pyrazol-3-yl)benzyl]-2,3-dihydro-1H-isoindol-1-one Alias; 1,5-anhydro-2,4-dideoxy-2-(4-fluoro-5-methyl-6-(4-(1-methyl-1H-pyrazol-3-yl)benzyl)-1-oxo-1,3-dihydro-2H-isoindol-2-yl)-L-threo-pentitol		436.0

[0583]

Table 1-29

Ex. No.	IUPAC Name	structure formula	MS
141	2-((1S,2S)-2-hydroxycyclohexyl)-6-(4-methoxybenzyl)-4,5-dimethylisoindolin-1-one		380.1
142	6-(4-ethoxybenzyl)-2-[(3S,4S)-4-hydroxytetrahydro-2H-pyran-3-yl]-4,5-dimethyl-2,3-dihydro-1H-isoindol-1-one Alias; 1,5-anhydro-2,4-dideoxy-2-(6-(4-ethoxybenzyl)-4,5-dimethyl-1-oxo-1,3-dihydro-2H-isoindol-2-yl)-L-threo-pentitol		396.1
143	2-[(3S,4S)-4-hydroxytetrahydro-2H-pyran-3-yl]-4,5-dimethyl-6-[4-(propan-2-yloxy)benzyl]-2,3-dihydro-1H-isoindol-1-one Alias; 1,5-anhydro-2,4-dideoxy-2-(6-(4-isopropoxybenzyl)-4,5-dimethyl-1-oxo-1,3-dihydro-2H-isoindol-2-yl)-L-threo-pentitol		410.1
144	5-ethyl-6-[2-fluoro-4-(1H-pyrazol-1-yl)benzyl]-2-[(3S,4S)-4-hydroxytetrahydro-2H-pyran-3-yl]-2,3-dihydro-1H-isoindol-1-one Alias; 1,5-anhydro-2,4-dideoxy-2-(5-ethyl-6-(2-fluoro-4-(1H-pyrazol-1-yl)benzyl)-1-oxo-1,3-dihydro-2H-isoindol-2-yl)-L-threo-pentitol		436.2
145	5-ethyl-6-(2-fluoro-4-(1H-pyrazol-1-yl)benzyl)-2-((1S,2S)-2-hydroxycyclopentyl)isoindolin-1-one		420.3






[0584]

Table 1-30

Ex. No.	IUPAC Name	structure formula	MS
146	5-ethyl-6-(2-fluoro-4-(1H-pyrazol-1-yl)benzyl)-2-(2-hydroxy-2-methylpropyl)isoindolin-1-one		408.1
147	2-[(3S,4S)-4-hydroxytetrahydro-2H-pyran-3-yl]-4,5-dimethyl-6-[4-(trifluoromethoxy)benzyl]-2,3-dihydro-1H-isoindol-1-one Alias; 1,5-anhydro-2,4-dideoxy-2-(4,5-dimethyl-1-oxo-6-(4-(trifluoromethoxy)benzyl)-1,3-dihydro-2H-isoindol-2-yl)-L-threo-pentitol		436.1
148	5-ethyl-2-[(3S,4S)-4-hydroxytetrahydro-2H-pyran-3-yl]-6-(4-methoxybenzyl)-2,3-dihydro-1H-isoindol-1-one Alias; 1,5-anhydro-2,4-dideoxy-2-(5-ethyl-6-(4-methoxybenzyl)-1-oxo-1,3-dihydro-2H-isoindol-2-yl)-L-threo-pentitol		382.2
149	6-((6-ethylpyridin-3-yl)methyl)-2-((1S,2S)-2-hydroxycyclohexyl)-4,5-dimethylisoindolin-1-one		379.3
150	4-fluoro-2-[(3S,4S)-4-hydroxytetrahydro-2H-pyran-3-yl]-6-(4-methoxybenzyl)-5-methyl-2,3-dihydro-1H-isoindol-1-one Alias; 1,5-anhydro-2,4-dideoxy-2-(4-fluoro-6-(4-methoxybenzyl)-5-methyl-1-oxo-1,3-dihydro-2H-isoindol-2-yl)-L-threo-pentitol		386.1






[0585]

Table 1-31

Ex. No.	IUPAC Name	structure formula	MS
151	4-chloro-2-((1S,2S)-2-hydroxycyclohexyl)-6-(4-methoxybenzyl)-5-methylisoindolin-1-one		400.1
152	4-chloro-2-((1S,2S)-2-hydroxycyclopentyl)-6-(4-methoxybenzyl)-5-methylisoindolin-1-one		386.1
153	2-(3-fluoropyridin-2-yl)-4,5-dimethyl-6-(4-(1H-pyrazol-1-yl)benzyl)isoindolin-1-one		413.1
154	4-fluoro-2-((1S,2S)-2-hydroxycyclohexyl)-6-(4-methoxybenzyl)-5-methylisoindolin-1-one		384.2
155	4-((2-((1S,2S)-2-hydroxycyclohexyl)-6,7-dimethyl-3-oxo-2,3-dihydro-1H-isoindol-5-yl)methyl)benzonitrile		375.2






[0586]

Table 1-32

Ex. No.	IUPAC Name	structure formula	MS
156	4-({2-[(3S,4S)-4-hydroxytetrahydro-2H-pyran-3-yl]-6,7-dimethyl-3-oxo-2,3-dihydro-1H-isoindol-5-yl)methyl}benzonitrile Alias; 1,5-anhydro-2-(6-(4-cyanobenzyl)-4,5-dimethyl-1-oxo-1,3-dihydro-2H-isoindol-2-yl)-2,4-dideoxy-L-threo-pentitol		377.3
157	4-fluoro-2-[(3S,4S)-4-hydroxytetrahydro-2H-pyran-3-yl]-5-methoxy-6-(4-methoxybenzyl)-2,3-dihydro-1H-isoindol-1-one Alias; 1,5-anhydro-2,4-dideoxy-2-(4-fluoro-5-methoxy-6-(4-methoxybenzyl)-1-oxo-1,3-dihydro-2H-isoindol-2-yl)-L-threo-pentitol		402.1
158	6-(3-fluoro-4-methoxybenzyl)-2-[(3S,4S)-4-hydroxytetrahydro-2H-pyran-3-yl]-4,5-dimethyl-2,3-dihydro-1H-isoindol-1-one Alias; 1,5-anhydro-2,4-dideoxy-2-(6-(3-fluoro-4-methoxybenzyl)-4,5-dimethyl-1-oxo-1,3-dihydro-2H-isoindol-2-yl)-L-threo-pentitol		400.1
159	2-(2-fluorophenyl)-4,5-dimethyl-6-(4-(1H-pyrazol-1-yl)benzyl)isoindolin-1-one		412.2
160	2-[(3S,4S)-4-hydroxytetrahydro-2H-pyran-3-yl]-6-(4-methoxy-3-methylbenzyl)-4,5-dimethyl-2,3-dihydro-1H-isoindol-1-one Alias; 1,5-anhydro-2,4-dideoxy-2-(6-(4-methoxy-3-methylbenzyl)-4,5-dimethyl-1-oxo-1,3-dihydro-2H-isoindol-2-yl)-L-threo-pentitol		396.2

[0587]

Table 1-33

Ex. No.	IUPAC Name	structure formula	MS
161	4-fluoro-2-((1S,2S)-2-hydroxycyclohexyl)-5-methoxy-6-(4-methoxybenzyl)isoindolin-1-one		400.2
162	2-(5-chloro-3-fluoropyridin-2-yl)-4,5-dimethyl-6-(4-(1H-pyrazol-1-yl)benzyl)isoindolin-1-one		447.1
163	6-(2-fluoro-4-methoxybenzyl)-2-[(3S,4S)-4-hydroxytetrahydro-2H-pyran-3-yl]-4,5-dimethyl-2,3-dihydro-1H-isoindol-1-one Alias; 1,5-anhydro-2,4-dideoxy-2-(6-(2-fluoro-4-methoxybenzyl)-4,5-dimethyl-1-oxo-1,3-dihydro-2H-isoindol-2-yl)-L-threo-pentitol		400.1
164	2-[(3S,4S)-4-hydroxytetrahydro-2H-pyran-3-yl]-6-(4-methoxybenzyl)-5-methyl-2,3-dihydro-1H-isoindol-1-one Alias; 1,5-anhydro-2,4-dideoxy-2-(6-(4-methoxybenzyl)-5-methyl-1-oxo-1,3-dihydro-2H-isoindol-2-yl)-L-threo-pentitol		368.2
165	rac-2-(trans-4-hydroxytetrahydro-2H-pyran-3-yl)-6-(4-methoxybenzyl)-4,5-dimethyl-2,3-dihydro-1H-isoindol-1-one		382.2

[0588]

Table 1-34

Ex. No.	IUPAC Name	structure formula	MS
166	2-[(3S,4S)-4-hydroxytetrahydro-2H-pyran-3-yl]-5-methoxy-6-(4-methoxybenzyl)-4-methyl-2,3-dihydro-1H-isoindol-1-one Alias; 1,5-anhydro-2,4-dideoxy-2-(5-methoxy-6-(4-methoxybenzyl)-4-methyl-1-oxo-1,3-dihydro-2H-isoindol-2-yl)-L-threo-pentitol		398.2
167	2-((1S,2S)-2-hydroxycyclohexyl)-6-(4-methoxybenzyl)-5-methylisoindolin-1-one		366.1
168	2-(3-hydroxy-3-methylbutan-2-yl)-6-(4-methoxybenzyl)-4,5-dimethylisoindolin-1-one		368.2

[0589]

5 Formulation Example 1

(1) Compound obtained in Example 1	10.0 g
(2) Lactose	60.0 g
(3) Cornstarch	35.0 g
(4) Gelatin	3.0 g
10 (5) Magnesium stearate	2.0 g

A mixture of the compound (10.0 g) obtained in Example 1, lactose (60.0 g) and cornstarch (35.0 g) is passed through a 1 mm mesh sieve by using 10 wt% aqueous gelatin solution (30 mL) (3.0 g as gelatin) and the granules are dried at 40°C and
5 sieved again. The obtained granules are mixed with magnesium stearate (2.0 g) and the mixture is compressed. The obtained core tablets are coated with a sugar coating of an aqueous suspension of saccharose, titanium dioxide, talc and gum arabic. The coated tablets are glazed with beeswax to give
10 1000 coated tablets.

[0590]

Formulation Example 2

(1) Compound obtained in Example 1	10.0 g
(2) Lactose	70.0 g
15 (3) Cornstarch	50.0 g
(4) Soluble starch	7.0 g
(5) Magnesium stearate	3.0 g

The compound (10.0 g) obtained in Example 1 and magnesium stearate (3.0 g) are granulated using aqueous soluble starch
20 solution (70 mL) (7.0 g as soluble starch), and the obtained granules are dried, and mixed with lactose (70.0 g) and cornstarch (50.0 g). The mixture is compressed to give 1000 tablets.

[0591]

25 Experimental Example 1

Measurement of M1 receptor positive allosteric modulator (M1PAM) activity

The activity of a test compound in the presence of acetylcholine at EC20 concentration (final concentration 0.6-
30 0.8 nM), which affords an action corresponding to about 20% of the maximum activity, was measured as PAM activity. The method is as follows. CHO-K1 cells stably expressing a human M1 receptor (hCHRM1) were plated on a 384-well black clear bottom plate (BD Falcon) at 5,000 cells/well, and cultured in an
35 incubator at 37°C, 5% CO₂ for 1 day. The medium in the cell

plate was removed, and assay buffer A (Recording medium (DOJINDO LABORATORIES), 0.1% BSA (Wako Pure Chemical Industries, Ltd.), 2.5 µg/mL Fluo-4 AM (DOJINDO LABORATORIES), 0.08% Pluronic F127 (DOJINDO LABORATORIES), 1.25 mM probenecid (DOJINDO LABORATORIES)) containing a calcium indicator was added at 30 µL/well. The cells were left standing in the incubator at 37°C, 5% CO₂ for 30 min, and further left standing at room temperature for 30 min. A test compound prepared by diluting with assay buffer B (HBSS (Invitrogen), 20 mM HEPES (Invitrogen), 0.1% BSA) containing 2.4-3.2 nM acetylcholine was added at 10 µL/well, and the fluorescence was measured by FLIPRtetra (Molecular Devices) for 1 min every 1 second. With the definition that the amount of change in the fluorescence on addition of acetylcholine (final concentration 1 µM) is 100% and that on addition of DMSO instead of a test compound is 0%, the activity (%) of the test compound was calculated, and the inflection point in the concentration-dependent curve of the test compound was calculated as IP values. The results are shown in Table 2.

20

[0592]

Table 2-1

Example No.	IP value (nM)	activity (%) at 10 μ M
1	750	111
2	86	109
3	15	106
4	19	98
5	8.2	102
6	5.3	101
7	180	107
8	290	90
9	43	102
10	66	106
11	13	105
12	7.7	91
13	7.9	104
14	1.2	94
15	11	103
16	12	94
17	18	90
18	10	88
19	2	101
20	5.5	109
21	6.2	106
22	32	110
23	46	100
24	4.3	88
25	10	90
26	9.2	99
28	4.8	106
29	9.5	102
32	5.6	99
33	37	95
34	18	108

WO 2015/163485

PCT/JP2015/062912

35	3.9	103
36	83	107
40	37	97

[0593]

Table 2-2

Example No.	IP value (nM)	activity (%) at 10 μ M
41	30	97
42	9.2	115
43	98	110
47	3	99
48	31	102
49	31	97
51	42	86
53	28	103
54	37	92
55	98	101
58	95	100
59	34	100
60	72	94
61	35	97
62	95	103
63	45	99
64	22	104
67	26	93
68	96	95
69	13	95
71	56	100
75	23	97
78	8.8	95
79	72	93
80	14	90

[0594]

Table 2-3

Example No.	IP value (nM)	activity (%) at 10 μ M
82	5.7	91
83	13	92
84	52	104
85	41	105
88	85	98
89	5.7	94
91	38	95
92	60	91
93	5.6	94
94	9.5	92
95	26	92
97	5.3	96
99	16	103
100	94	101
102	13	91
103	6	94
104	54	101
105	10	92
106	11	92
107	14	102
109	23	100
110	24	77
111	26	103
112	30	93
113	1.5	82
114	15	96
115	25	92
116	6.1	111
118	25	106
119	3.1	91
120	25	93

[0595]

Table 2-4

Example No.	IP value (nM)	activity (%) at 10 μ M
121	22	89
123	2.1	90
124	16	86
125	73	93
127	59	104
128	5	97
129	53	100
130	2.2	94
131	12	100
132	7.3	87
133	35	87
136	75	100
138	25	106
139	14	102
140	2	93
141	18	106
142	31	106
144	23	97
148	63	89
149	16	99
150	14	89
151	16	92
152	47	91
153	57	92
154	22	100
155	7.1	100
156	4.3	80
157	84	97
158	28	111
160	47	112
161	32	94

162	66	95
163	51	108
164	62	102
165	36	103
167	30	95

[0596]

Experimental Example 2

Measurement of myo-Inositol 1 phosphate (IP1)

5 Animals used were male Long-Evans rats. They were used after acclimation for at least 1 week. Test compounds were suspended in 0.5% aqueous methylcellulose solution, and the suspension was orally administered to the rats. At a certain period of time after the oral administration, the solution

10 prepared by dissolving lithium chloride in saline was subcutaneously administered into the rats. At a certain period of time after the subcutaneously administration, their bilateral hippocampi were isolated from the rats, and the wet weight thereof was measured. The hippocampi were homogenized

15 with HEPES buffer, followed by centrifugation. The IP1 and protein concentrations in the supernatant were measured by IP-One HTRF assay kit (Cisbio Bioassays) and BCA protein assay kit (Thermo Scientific), respectively. The level of the IP1 production was expressed as the ratio of the concentration of

20 IP1 to that of protein. The increase rate of the IP1 production was shown as a relative value when Vehicle administration group as 100%. The results are shown in Table 3.

[0597]

Table 3

test compound	increase rate at 10 mg/kg (% of Vehicle)
Example 18	40
Example 20	108
Example 25	32

[0598]

Experimental Example 3

Novel object recognition test

Novel object recognition test is comprised of two trials
5 called the acquisition and the retention trials. Scopolamine-
induced memory deficits models were used for the test, and
animals used were male Long-Evans rats (7-week-old).
On the day before the test, for acclimation, the rats were
allowed to freely move about the test box (40×40×50 cm) for 10
10 minutes. On the test day, the rats were acclimated to the test
room for about 1 hr prior to the test. The test compounds were
orally administered to the rats in a single dose 2 hr before
the acquisition trial. For induction of learning and memory
deficits, scopolamine (0.1 mg/kg) was subcutaneously
15 administered into the rats 30 min before the acquisition trial.
For the acquisition trial, two identical objects (A1, A2) were
placed in the test box. The rats were put in the test box for
3 min, and the duration exploring each object was measured.
The retention trial was performed 4 hr after the acquisition
20 trial. For the retention trial, the familiar object (A3) used
for the acquisition trial and the novel object (B) different
shape from A3 were placed in the test box. The rats were put
in the test box for 3 min. The duration exploring each object
in the acquisition trial and the retention trial, and the
25 exploration rate (%) of novel object was calculated. The
exploration rate (%) of novel object was expressed as (the
duration exploring the novel object)/[(the duration exploring
the novel object)+(the duration exploring the familiar
object)]×100 (%) at mean ± standard error. The results are
30 shown below.

exploration rate (%) of novel object

control: 63.35±1.59%

solvent-scopolamine group: 52.08±2.47%

35 Example 18 (3 mg/kg)-scopolamine group: 59.11±3.87%

control :66.4±2.3%

solvent-scopolamine group: 50.4±2.4%

Example 20 (3 mg/kg)-scopolamine group: 61.2±3.1%

5

control: 62.37±3.24 %

solvent-scopolamine group: 48.39±2.01 %

Example 25 (10 mg/kg)-scopolamine group: 57.55±5.03 %

Industrial Applicability

10 [0599]


The compound of the present invention is useful as a cholinergic muscarinic M1 receptor positive allosteric modulator, or a medicament such as an agent for the prophylaxis or treatment of Alzheimer's disease, schizophrenia, pain, sleep
15 disorder, Parkinson's disease dementia, dementia with Lewy bodies and the like.

[0600]

This application is based on patent application No. 2014-089585 filed on April 23, 2014 in Japan, the contents of which
20 are encompassed in full herein.

CLAIMS

1. A compound represented by the formula (I):



5 wherein

R¹ is

(1) a phenyl group optionally substituted by 1 to 3 substituents selected from

(i) a halogen atom, and

10 (ii) a cyano group,

(2) a C₅₋₆ cycloalkyl group optionally substituted by 1 to 3 substituents selected from

(i) a hydroxy group,

(ii) a C₁₋₆ alkyl group optionally substituted by 1 to 3

15 hydroxy groups, and

(iii) a C₁₋₆ alkoxy group,

(3) a 5- or 6-membered monocyclic non-aromatic heterocyclic group optionally substituted by 1 to 3 hydroxy groups,

(4) a 5- or 6-membered monocyclic aromatic heterocyclic group

20 optionally substituted by 1 to 3 halogen atoms, or

(5) a C₁₋₆ alkyl group optionally substituted by 1 to 3 substituents selected from

(i) a hydroxy group, and

(ii) a 3- to 8-membered monocyclic non-aromatic

25 heterocyclic group;

R² is

- (1) a hydrogen atom,
- (2) a halogen atom, or
- (3) a C₁₋₆ alkyl group;

R³ is

- 5 (1) a hydrogen atom,
- (2) a halogen atom,
- (3) a cyano group,
- (4) a C₁₋₆ alkyl group optionally substituted by 1 to 3 halogen atoms,
- 10 (5) a C₁₋₆ alkoxy group, or
- (6) a C₃₋₆ cycloalkyl group;

R⁴ is


- (1) a halogen atom,
- (2) a cyano group,
- 15 (3) a C₁₋₆ alkyl group,
- (4) a C₁₋₆ alkoxy group optionally substituted by 1 to 3 halogen atoms,
- (5) a carbamoyl group,
- (6) a mono- or di-C₁₋₆ alkyl-carbamoyl group, or
- 20 (7) a 5- or 6-membered monocyclic aromatic heterocyclic group optionally substituted by 1 to 3 C₁₋₆ alkyl groups; and

Ring A is a 6-membered aromatic ring optionally further substituted by 1 to 3 substituents, in addition to R⁴, selected from

- 25 (a) a halogen atom,
- (b) a C₁₋₆ alkyl group, and
- (c) a C₁₋₆ alkoxy group,


or a salt thereof.

- 30 2. The compound according to claim 1, wherein the partial structure represented by the following formula:



in the formula (I) is a partial structure represented by the

following formula:



or a salt thereof.

5 3. The compound according to claim 1, wherein

R¹ is

(1) a phenyl group optionally substituted by 1 to 3
substituents selected from

(i) a halogen atom, and

10 (ii) a cyano group,

(2) a C₅₋₆ cycloalkyl group optionally substituted by 1 to 3
hydroxy groups,

(3) a 5- or 6-membered monocyclic non-aromatic heterocyclic
group optionally substituted by 1 to 3 hydroxy groups,

15 (4) a 5- or 6-membered monocyclic aromatic heterocyclic group
optionally substituted by 1 to 3 halogen atoms, or

(5) a C₁₋₆ alkyl group optionally substituted by 1 to 3
substituents selected from

(i) a hydroxy group, and

20 (ii) a 3- to 8-membered monocyclic non-aromatic
heterocyclic group;

R² is

(1) a hydrogen atom,

(2) a halogen atom, or

25 (3) a C₁₋₆ alkyl group;

R³ is

(1) a hydrogen atom,

(2) a halogen atom,

(3) a cyano group,

30 (4) a C₁₋₆ alkyl group optionally substituted by 1 to 3 halogen
atoms,

(5) a C₁₋₆ alkoxy group, or

(6) a C₃₋₆ cycloalkyl group;

R⁴ is

(1) a halogen atom,

(2) a cyano group,

5 (3) a C₁₋₆ alkyl group,

(4) a C₁₋₆ alkoxy group optionally substituted by 1 to 3 halogen atoms,

(5) a carbamoyl group,

(6) a mono- or di-C₁₋₆ alkyl-carbamoyl group, or

10 (7) a 5- or 6-membered monocyclic aromatic heterocyclic group optionally substituted by 1 to 3 C₁₋₆ alkyl groups; and

Ring A is a 6-membered aromatic ring optionally further substituted by 1 to 3 substituents, in addition to R⁴, selected from


15 (a) a halogen atom,

(b) a C₁₋₆ alkyl group, and


(c) a C₁₋₆ alkoxy group,

or a salt thereof.

20 4. The compound according to claim 3, wherein the partial structure represented by the following formula:



in the formula (I) is a partial structure represented by the following formula:



25

or a salt thereof.

5. The compound according to claim 1, wherein

R¹ is

(1) a phenyl group optionally substituted by 1 to 3 substituents selected from

5 (i) a halogen atom, and

(ii) a cyano group,

(2) a C₅₋₆ cycloalkyl group optionally substituted by 1 to 3 hydroxy groups,

(3) a 5- or 6-membered monocyclic non-aromatic heterocyclic
10 group optionally substituted by 1 to 3 hydroxy groups, or

(4) a C₁₋₆ alkyl group optionally substituted by 1 to 3 substituents selected from

(i) a hydroxy group, and

(ii) a 3- to 8-membered monocyclic non-aromatic
15 heterocyclic group;

R² is

(1) a hydrogen atom,

(2) a halogen atom, or

(3) a C₁₋₆ alkyl group;

20 R³ is

(1) a hydrogen atom,

(2) a halogen atom,

(3) a cyano group,

(4) a C₁₋₆ alkyl group optionally substituted by 1 to 3 halogen
25 atoms,

(5) a C₁₋₆ alkoxy group, or

(6) a C₃₋₆ cycloalkyl group;

R⁴ is

(1) a halogen atom,

30 (2) a cyano group,

(3) a C₁₋₆ alkyl group,

(4) a C₁₋₆ alkoxy group optionally substituted by 1 to 3 halogen
atoms,

(5) a mono- or di-C₁₋₆ alkyl-carbamoyl group, or


35 (6) a 5- or 6-membered monocyclic aromatic heterocyclic group

optionally substituted by 1 to 3 C₁₋₆ alkyl groups; and


Ring A is a 6-membered aromatic ring optionally further substituted by 1 to 3 substituents, in addition to R⁴, selected from

- 5 (a) a halogen atom, and
(b) a C₁₋₆ alkoxy group,
or a salt thereof.

6. The compound according to claim 5, wherein the partial
10 structure represented by the following formula:



in the formula (I) is a partial structure represented by the following formula:



15 or a salt thereof.

7. The compound according to claim 1, wherein

R¹ is

- (1) a C₅₋₆ cycloalkyl group optionally substituted by 1 to 3
20 hydroxy groups, or
(2) a 5- or 6-membered monocyclic non-aromatic heterocyclic
group optionally substituted by 1 to 3 hydroxy groups;

R² is

- (1) a halogen atom, or
25 (2) a C₁₋₆ alkyl group;

R³ is a C₁₋₆ alkyl group;

R⁴ is


- (1) a C₁₋₆ alkyl group,
(2) a C₁₋₆ alkoxy group, or
30 (3) a 5- or 6-membered monocyclic aromatic heterocyclic group;

and


Ring A is a benzene ring or a pyridine ring, each of which is unsubstituted, in addition to R⁴, or a salt thereof.

5

8. The compound according to claim 7, wherein the partial structure represented by the following formula:



in the formula (I) is a partial structure represented by the
10 following formula:



or a salt thereof.

9. The compound according to claim 1, wherein

15

R¹ is

- (1) a cyclohexyl group substituted by one hydroxy group, or
- (2) a tetrahydropyranyl group substituted by one hydroxy group;

R² is

- (1) a halogen atom, or
- 20 (2) a C₁₋₆ alkyl group;

R³ is a C₁₋₆ alkyl group;


R⁴ is

- (1) a C₁₋₆ alkyl group,
- (2) a C₁₋₆ alkoxy group, or
- 25 (3) a pyrazolyl group; and


Ring A is a benzene ring or a pyridine ring, each of which is unsubstituted, in addition to R⁴, or a salt thereof.

30

10. The compound according to claim 9, wherein the partial structure represented by the following formula:



in the formula (I) is a partial structure represented by the
5 following formula:



or a salt thereof.

11. 2-[(3S,4S)-4-Hydroxytetrahydro-2H-pyran-3-yl]-6-(4-methoxybenzyl)-4,5-dimethyl-2,3-dihydro-1H-isoindol-1-one, or a
10 salt thereof.

12. 4-Fluoro-2-[(3S,4S)-4-hydroxytetrahydro-2H-pyran-3-yl]-5-methyl-6-[4-(1H-pyrazol-1-yl)benzyl]-2,3-dihydro-1H-isoindol-1-
15 one, or a salt thereof.

13. 2-((1S,2S)-2-Hydroxycyclohexyl)-4,5-dimethyl-6-((6-methylpyridin-3-yl)methyl)isoindolin-1-one, or a salt thereof.

20 14. A medicament comprising the compound according to claim 1 or a salt thereof.

15. The medicament according to claim 14, which is a cholinergic muscarinic M1 receptor positive allosteric
25 modulator.

16. The compound according to claim 1 or a salt thereof for use in the prophylaxis or treatment of Alzheimer's disease, schizophrenia, pain, sleep disorder, Parkinson's disease
30 dementia or dementia with Lewy bodies.

17. Use of the compound according to claim 1 or a salt thereof
for the production of an agent for the prophylaxis or treatment
of Alzheimer's disease, schizophrenia, pain, sleep disorder,
5 Parkinson's disease dementia or dementia with Lewy bodies.