

DESCRIPTION

Title of Invention: SELECTIVE INHIBITOR OF EXON 18 AND/OR EXON 21
MUTANT EGFR

5 Technical Field
[0001]

The present invention relates to an antitumor agent against
cancers, comprising an exon 18 and/or exon 21 mutant epidermal
growth factor receptor (hereinafter also referred to as "EGFR").

10

Background Art
[0002]

EGFR is a receptor-type tyrosine kinase, exerts its
physiological function in normal tissue by being bound to
15 Epidermal Growth Factor (hereinafter also referred to as EGF),
which is a ligand, and contributes to growth and apoptosis
inhibition in epithelial tissues (NPL 1). Further, somatic
mutation of EGFR gene has been known as a cancer-causing gene;
for example, EGFR in which codons 746 to 750 in exon 19 are
20 deleted (hereinafter also referred to as "exon 19 deletion
mutation") and EGFR in which leucine encoded by codon 858 in exon
21 is mutated to arginine (hereinafter also referred to as "L858R
mutation") constantly induces EGF-independent kinase activity,
and contributes to the growth and survival of cancer cells (NPL
25 2). These mutations are observed, for example, in 30 to 50% of
non-small-cell lung cancer in East Asia. The mutations are also
observed in about 10% of non-small-cell lung cancer in Europe and
the United States, and are regarded as one of the causes of
cancers (NPL 3).

30 [0003]

Therefore, research and development of EGFR inhibitor
as an antitumor agent have actively been conducted, and
introduced into the treatment of various EGFR mutation-positive
lung cancers (NPL 2 and NPL 4). Gefitinib, erlotinib, and
35 afatinib have been used as a therapeutic agent against exon 19

deletion mutant and L858R mutant EGFR-positive lung cancers. Exon 19 deletion mutation and L858R mutation account for 90% of EGFR mutation. Further, occurrence of acquired resistance in the process of the treatment using these agents has been known, and 50% thereof is caused by resistance mutation EGFR in which codon 790 of exon 20 is changed from threonine to methionine (hereinafter also referred to as "T790M mutation"). To treat lung cancers having this mutation, osimertinib has been used as a therapeutic agent. Therefore, treatments using EGFR inhibitors are in the process of being established for lung cancer patients having major EGFR mutations.

[0004]

On the other hand, at present, no treatments using EGFR inhibitors have been established with respect to some rare EGFR mutations, such as point mutation or deletion mutation of exon 18, point mutation of exon 21, or the like; and there are reports that the drug sensitivity of these mutant EGFR varies depending on the mutation type (NPL 4). For example, the sensitivity of the lung cancer having point mutation in which glycine encoded by codon 719 of exon 18 is substituted with an arbitrary amino acid (hereinafter also referred to as "G719X mutation") or the lung cancer in which leucine encoded by codon 861 of exon 21 is substituted with glutamine (hereinafter also referred to as "L861Q mutation") with respect to gefitinib, erlotinib, and afatinib is lower than those of exon 19 deletion mutation and L858R mutation, which are drug-sensitive mutations. Further, there are reports of skin disorders and digestive tract disorders as common side effects by the administration of therapeutic doses of afatinib, gefitinib, and erlotinib. It is widely thought that these side effects are attributable to inhibition of the function of wild-type EGFR expressed in normal tissues, such as skin or digestive tract, by the therapeutic agent (NPL 1); the development of an inhibitor characterized by lower inhibitory activity with respect to wild-type EGFR of normal tissues, compared with mutant EGFR expressed in tumor tissues, has been

desired in view of reduction of side effects.

[0005]

Therefore, development of a drug having high inhibitory activity with respect to exon 18 and exon 21 mutant EGFR, as well as high selectivity with respect to mutant EGFR compared with wild-type EGFR, is expected to enable inhibition of growth of lung cancer cells having mutant EGFR with a dose lower than the dose causing the side effects in the skin or digestive tract, thereby contributing to life prolongation or increase in QOL of patients with mutant EGFR-positive cancers for which treatment methods have not been established. Further, a drug having high inhibitory activity with respect to T790M, which is acquired resistance mutation against treatments using EGFR inhibitors, is expected to reduce expression frequency of acquired resistance during the treatments using EGFR inhibitors against exon 18 or exon 21 mutant EGFR, which is *de novo* mutation; and is therefore expected to contribute to the life prolongation of cancer patients.

Citation List

Patent Literature

[0006]

PTL 1: WO2015/175632A1

PTL 2: WO2015/025936A1

25

Non-Patent Literature

[0007]

NPL 1: Nat. Rev. Cancer, Vol. 6, pp. 803-812 (2006)

NPL 2: Nature Medicine, Vol. 19, pp. 1389-1400 (2013)

30 NPL 3: Nat. Rev. Cancer, Vol. 7, pp. 169-181 (2007)

NPL 4: Lancet Oncol. Vol. 13, e. 23-31 (2012)

Summary of Invention

Technical Problem

35 [0008]

An object of the present invention is to provide an antitumor agent that does not cause inhibition of wild-type EGFR and thus causes smaller side effects, serving as an inhibitor that can ensure high selectivity with respect to exon 18 and/or
5 exon 21 mutant EGFR for which the therapeutic effects of the previously known EGFR inhibitors are insufficient.

Solution to Problem

[0009]

10 The inventors of the present invention conducted extensive research, and found that exon 18 and/or exon 21 mutant EGFR is an appropriate target in treating cancers, and that EGFR inhibitors conventionally used for the treatments have inferior selectivity between exon 18 and/or exon 21 mutant EGFR. Further,
15 the inventors also confirmed that the compound of the present invention exerts superior selectivity and tumor growth inhibitory effects with respect to exon 18 and/or exon 21 mutant EGFR. With this finding, the inventors accomplished the present invention.

[0010]

20 The present invention encompasses the following embodiments.

[0011]

Item 1.

 An antitumor agent for treating a malignant tumor
25 patient expressing EGFR having at least one mutation selected from the group consisting of G719X mutation of exon 18, E709X mutation of exon 18, and L861X mutation of exon 21, wherein X represents an arbitrary amino-acid residue, the antitumor agent comprising (S)-N-(4-amino-6-methyl-5-(quinolin-3-yl)-8,9-
30 dihydropyrimido[5,4-b]indolizin-8-yl)acrylamide or a salt thereof.

[0012]

Item 2.

 The antitumor agent according to Item 1, wherein the exon 18 mutation is at least one mutation selected from the group
35 consisting of G719A, G719S, G719C, E709K and E709A.

[0013]

Item 3.

The antitumor agent according to Item 1 or 2, wherein the exon 21 mutation is L861Q.

5 [0014]

Item 4.

The antitumor agent according to any one of Items 1 to 3, wherein the EGFR further has T790M mutation.

[0015]

10 Item 5.

A method for treating a malignant tumor patient, comprising the step of administering (S)-N-(4-amino-6-methyl-5-(quinolin-3-yl)-8,9-dihydropyrimido[5,4-b]indolizin-8-yl)acrylamide or a salt thereof to a malignant tumor patient expressing EGFR having at least one mutation selected from the group consisting of G719X mutation of exon 18, E709X mutation of exon 18, and L861X mutation of exon 21, wherein X represents an arbitrary amino-acid residue.

[0016]

20 Item 6.

The method according to Item 5, wherein the exon 18 mutation is at least one mutation selected from the group consisting of G719A, G719S, G719C, E709K and E709A.

[0017]

25 Item 7.

The method according to Item 5 or 6, wherein the exon 21 mutation is L861Q.

[0018]

Item 8.

30 The method according to any one of Items 5 to 7, wherein the EGFR further has T790M mutation.

[0019]

Item 9.

(S)-N-(4-amino-6-methyl-5-(quinolin-3-yl)-8,9-dihydropyrimido[5,4-b]indolizin-8-yl)acrylamide or a salt thereof

35

for treating a malignant tumor patient expressing EGFR having at least one mutation selected from the group consisting of G719X mutation of exon 18, E709X mutation of exon 18, and L861X mutation of exon 21, wherein X represents an arbitrary amino-acid residue.

[0020]

Item 10.

The (S)-N-(4-amino-6-methyl-5-(quinolin-3-yl)-8,9-dihydropyrimido[5,4-b]indolizin-8-yl)acrylamide or a salt thereof according to Item 9, wherein the exon 18 mutation is at least one mutation selected from the group consisting of G719A, G719S, G719C, E709K and E709A.

[0021]

Item 11.

The (S)-N-(4-amino-6-methyl-5-(quinolin-3-yl)-8,9-dihydropyrimido[5,4-b]indolizin-8-yl)acrylamide or a salt thereof according to Item 9 or 10, wherein the exon 21 mutation is L861Q.

[0022]

Item 12.

The (S)-N-(4-amino-6-methyl-5-(quinolin-3-yl)-8,9-dihydropyrimido[5,4-b]indolizin-8-yl)acrylamide or a salt thereof according to any one of Items 9 to 11, wherein the EGFR further has T790M mutation.

[0023]

Item 13.

Use of (S)-N-(4-amino-6-methyl-5-(quinolin-3-yl)-8,9-dihydropyrimido[5,4-b]indolizin-8-yl)acrylamide or a salt thereof for treating a malignant tumor patient expressing EGFR having at least one mutation selected from the group consisting of G719X mutation of exon 18, E709X mutation of exon 18, and L861X mutation of exon 21, wherein X represents an arbitrary amino-acid residue.

[0024]

Item 14.

The use according to Item 13, wherein the exon 18

mutation is at least one mutation selected from the group consisting of G719A, G719S, G719C, E709K and E709A.

[0025]

Item 15.

5 The use according to Item 13 or 14, wherein the exon 21 mutation is L861Q.

[0026]

Item 16.

10 The use according to any one of Items 13 to 15, wherein the EGFR further has T790M mutation.

[0027]

Item 17.

15 Use of (S)-N-(4-amino-6-methyl-5-(quinolin-3-yl)-8,9-dihydropyrimido[5,4-b]indolizin-8-yl)acrylamide or a salt thereof for the production of a pharmaceutical agent for treating a malignant tumor patient expressing EGFR having at least one mutation selected from the group consisting of G719X mutation of exon 18, E709X mutation of exon 18, and L861X mutation of exon 21, wherein X represents an arbitrary amino-acid residue.

20 [0028]

Item 18.

 The use according to Item 17, wherein the exon 18 mutation is at least one mutation selected from the group consisting of G719A, G719S, G719C, E709K and E709A.

25 [0029]

Item 19.

 The use according to Item 17 or 18, wherein the exon 21 mutation is L861Q.

[0030]

30 Item 20.

 The use according to any one of Items 17 to 19, wherein the EGFR further has T790M mutation.

[0031]

Item 21.

35 A pharmaceutical composition comprising (S)-N-(4-amino-

6-methyl-5-(quinolin-3-yl)-8,9-dihydropyrimido[5,4-b]indolizin-8-yl)acrylamide or a salt thereof and a pharmaceutically acceptable carrier, for treating a malignant tumor patient expressing EGFR having at least one mutation selected from the group consisting of G719X mutation of exon 18, E709X mutation of exon 18, and L861X mutation of exon 21, wherein X represents an arbitrary amino-acid residue.

[0032]

Item 22.

10 The pharmaceutical composition according to Item 21, wherein the exon 18 mutation is at least one mutation selected from the group consisting of G719A, G719S, G719C, E709K and E709A.

[0033]

Item 23.

15 The pharmaceutical composition according to Item 21 or 22, wherein the exon 21 mutation is L861Q.

[0034]

Item 24.

20 The pharmaceutical composition according to any one of Items 21 to 23, wherein the EGFR further has T790M mutation.

[0035]

Item 25.

25 A method for predicting therapeutic effects of chemotherapy using an antitumor agent comprising, as an active ingredient, (S)-N-(4-amino-6-methyl-5-(quinolin-3-yl)-8,9-dihydropyrimido[5,4-b]indolizin-8-yl)acrylamide or a salt thereof in a malignant tumor patient, the method comprising steps (1) and (2) below:

30 (1) a step of detecting the presence or absence of mutation of EGFR gene contained in a biological sample obtained from the patient; and

35 (2) a step of predicting that the chemotherapy is highly likely to exhibit sufficient therapeutic effects with respect to the patient when the results of the detection in step (1) found that the EGFR gene has at least one mutation selected

from the group consisting of G719X mutation of exon 18, E709X mutation of exon 18, and L861X mutation of exon 21, wherein X represents an arbitrary amino-acid residue.

[0036]

5 Item 26.

The method according to Item 25, wherein the exon 18 mutation is at least one mutation selected from the group consisting of G719A, G719S, G719C, E709K and E709A.

[0037]

10 Item 27.

The method according to Item 25 or 26, wherein the exon 21 mutation is L861Q.

[0038]

Item 28.

15 The method according to any one of Items 25 to 27, wherein the EGFR further has T790M mutation.

[0039]

Item 29.

20 A method for treating a malignant tumor patient, comprising steps (1) to (3) below:

(1) a step of detecting the presence or absence of mutation of EGFR gene contained in a biological sample obtained from the patient;

25 (2) a step of predicting that chemotherapy using an antitumor agent comprising, as an active ingredient, (S)-N-(4-amino-6-methyl-5-(quinolin-3-yl)-8,9-dihydropyrimido[5,4-b]indolizin-8-yl)acrylamide or a salt thereof is highly likely to exhibit sufficient therapeutic effects with respect to the patient when the results of the detection in step (1) found that
30 the EGFR gene has at least one mutation selected from the group consisting of G719X mutation of exon 18, E709X mutation of exon 18, and L861X mutation of exon 21, wherein X represents an arbitrary amino-acid residue; and

35 (3) a step of administering (S)-N-(4-amino-6-methyl-5-(quinolin-3-yl)-8,9-dihydropyrimido[5,4-b]indolizin-8-

yl)acrylamide or a salt thereof to a malignant tumor patient who was predicted highly likely to sufficiently respond to the chemotherapy using an antitumor agent comprising, as an active ingredient, (S)-N-(4-amino-6-methyl-5-(quinolin-3-yl)-8,9-dihydropyrimido[5,4-b]indolizin-8-yl)acrylamide or a salt thereof, in step (2).

[0040]

Item 30.

The method according to Item 29, wherein the exon 18 mutation is at least one mutation selected from the group consisting of G719A, G719S, G719C, E709K and E709A.

[0041]

Item 31.

The method according to Item 29 or 30, wherein the exon 21 mutation is L861Q.

[0042]

Item 32.

The method according to any one of Items 29 to 31, wherein the EGFR further has T790M mutation.

20

Advantageous Effects of Invention

[0043]

The antitumor agent of the present invention exerts high selectivity with respect to exon 18 and/or exon 21 mutant EGFR. Therefore, the antitumor agent of the present invention is useful in view of providing an antitumor agent that exerts superior therapeutic effects for a malignant tumor patient expressing EGFR having exon 18 and/or exon 21 mutation, for which the therapeutic effects of the previously known EGFR inhibitors are insufficient.

30

[0044]

The present invention is also useful in terms of providing a method for treating a malignant tumor patient expressing EGFR having exon 18 and/or exon 21 mutation.

35

[0045]

The previously known EGFR inhibitors have low selectivity with respect to exon 18 and exon 21 mutant EGFR, compared with wild-type EGFR; therefore, the difference between the dosage for ensuring the antitumor effects and the dosage causing the side effects (skin disorders, digestive tract disorders, etc.) derived from wild-type EGFR was small. Accordingly, the previously known EGFR inhibitors have difficulty in exerting sufficient therapeutic effects. In contrast, since the antitumor agent of the present invention has high selectivity with respect to exon 18 and exon 21 mutant EGFR, it is possible to increase the dosage without causing side effects derived from wild-type EGFR. Therefore, the antitumor agent of the present invention exerts superior therapeutic effects for a malignant tumor patient expressing EGFR having exon 18 and/or exon 21 mutation.

[0046]

In addition, the antitumor agent of the present invention exhibited high inhibitory activity with respect to exon 18 and exon 21 mutant EGFR in the presence of T790M mutation, which is acquired resistance mutation in the exon 20 region. Therefore, the antitumor agent of the present invention exerts superior therapeutic effects with respect to a malignant tumor patient whose response to the existing drug is low due to acquired resistance mutation caused by the use of existing antitumor agent.

[0047]

Further, the antitumor agent of the present invention is also useful in terms of reducing expression frequency of acquired resistance during the treatments using EGFR inhibitors against exon 18 or exon 21 mutant EGFR, which is *de novo* mutation, because of its high inhibitory activity against exon 18 and exon 21 mutant EGFR even under the presence of T790M mutation, which is acquired resistance mutation in the exon 20 region.

Brief Description of Drawings

[0048]

Fig. 1 illustrates the tumor volume (which may be hereinafter referred to as "TV") of mouse models subcutaneously transplanted with G719A mutant EGFR-expressing cell line to measure the
5 antitumor effect of compound A.

Fig. 2 illustrates the body weight during a dosing period of compounds of mouse models subcutaneously transplanted with G719A mutant EGFR-expressing cell line to measure the toxicity of
10 compound A.

Fig. 3 illustrates the tumor volume of mouse models subcutaneously transplanted with G719A+T790M mutant EGFR-expressing cell line to measure the antitumor effect of compound
15 A.

Fig. 4 illustrates the body weight during a dosing period of compounds of mouse models subcutaneously transplanted with G719A+T790M mutant EGFR-expressing cell line to measure the
20 toxicity of compound A.

Description of Embodiments

[0049]

Preferable examples of various definitions in the scope
25 of the present invention used in this specification are explained below in detail.

[0050]

In this specification, "EGFR" refers to a human epidermal growth factor receptor protein, and is also referred to
30 as ErbB-1 or HER1.

[0051]

In this specification, "wild-type EGFR" refers to EGFR free of somatic mutation, which is a protein comprising the amino acid sequence represented by SEQ ID NO: 1 (GenBank accession
35 number: NP_005219.2).

[0052]

In this specification, "exon 18" refers to 688-728 region in the amino acid sequence of wild-type EGFR (SEQ ID NO: 1).

5 [0053]

In this specification, "exon 18 mutation" refers to point mutation in amino acid in the exon 18 region of wild-type EGFR (SEQ ID NO: 1). Preferable exon 18 mutation is point mutation or deletion mutation with 1 amino acid substitution in
10 the exon 18 region. More preferably, the exon 18 mutation is E709X, which is point mutation in which glutamic acid encoded by codon 709 of exon 18 is substituted with an arbitrary amino acid; or G719X, which is point mutation in which glycine encoded by codon 719 of exon 18 is substituted with an arbitrary amino acid.
15 More specifically, preferable examples of E709X include E709K, which is point mutation in which glutamic acid encoded by codon 709 in the exon 18 region is substituted with lysine; and E709A, which is point mutation in which glutamic acid encoded by codon 709 in the exon 18 region is substituted with alanine. Preferable
20 examples of G719X include G719A, which is point mutation in which glycine encoded by codon 719 in the exon 18 region is substituted with alanine; G719S, which is point mutation in which glycine encoded by codon 719 in the exon 18 region is substituted with serine; and G719C, which is point mutation in which glycine
25 encoded by codon 719 in the exon 18 region is substituted with cysteine. Among these, G719A is particularly preferable.

[0054]

In the present invention, "exon 21" refers to 824-875 region in the amino acid sequence of wild-type EGFR (SEQ ID NO:
30 1).

[0055]

In this specification, "exon 21 mutation" refers to point mutation in amino acid in the exon 21 region of wild-type EGFR (SEQ ID NO: 1). Preferable exon 21 mutation is point
35 mutation with 1 amino acid substitution in the exon 21 region.

More preferably, the exon 21 mutation is L861X, which is point mutation in which leucine encoded by codon 861 in the exon 21 region is substituted with an arbitrary amino acid. More specifically, L861Q, which is point mutation in which leucine
5 encoded by codon 861 in the exon 21 region is substituted with glutamine, is preferable.

[0056]

In the present invention, "exon 18 and/or exon 21 mutation" encompasses "exon 18 mutation," "exon 21 mutation," and
10 "exon 18 and exon 21 mutation."

[0057]

In the present invention, "point mutation" refers to mutation causing substitution, insertion, or deletion of one or more (e.g., about 1 to 10, preferably about 1 to 5, more
15 preferably about 1, 2, or 3) amino-acid residues; and may include in-frame insertion and/or deletion mutation as nucleic acid.

[0058]

"EGFR having exon 18 and/or exon 21 mutation" encompasses "EGFR having exon 18," "EGFR having exon 21 mutation,"
20 and "EGFR having exon 18 and exon 21 mutation".

[0059]

In this specification, the "EGFR having exon 18 mutation" refers to EGFR having at least one exon 18 mutation. The EGFR may have two or more different exon 18 mutations, and
25 preferably has a single exon 18 mutation. Further, the EGFR may also have a mutation other than exon 18 mutation (such as exon 19 deletion mutation, L858R mutation, or L790M mutation).

[0060]

In this specification, the "EGFR having exon 21 mutation" refers to EGFR having at least one exon 21 mutation. The EGFR may have two or more different exon 21 mutations, and
30 preferably has a single exon 21 mutation. Further, the EGFR may also have a mutation other than exon 21 mutation (such as exon 19 deletion mutation, L858R mutation, or L790M mutation).

35 [0061]

Further, EGFR having exon 18 and/or exon 21 mutation may further have T790M mutation. T790M is acquired resistance mutation in the exon 20 region. T790M is known to be generated by the use of existing EGFR inhibitors. The acquisition of T790M often decreases the effects of existing drug with respect to malignant tumor patients.

[0062]

In the present invention, specifically, the EGFR having exon 18 and/or exon 21 mutation further having T790M mutation is preferably one of EGFR having E709X and/or G719X mutation in the exon 18 region further having T790M mutation, and EGFR having L861X mutation in the exon 21 region further having T790M mutation. In the present invention, more specifically, the EGFR having exon 18 and/or exon 21 mutation further having T790M mutation is preferably one of EGFR having E709K or E709A mutation further having T790M mutation, EGFR having G719A, G719S or G719C mutation further having T790M mutation, and EGFR having L861Q mutation further having T790M mutation. Among these, EGFR having G719A mutation further having T790M mutation and EGFR having L861Q mutation further having T790M mutation are particularly preferable.

[0063]

In the present invention, the method for detecting exon 18 and/or exon 21 mutation of EGFR expressed by a malignant tumor patient is not particularly limited insofar as the method is capable of detecting the mutations, and any known detection methods may be used.

[0064]

The sample used in the detection of exon 18 and/or exon 21 mutation is not particularly limited as long as the sample is a biological sample isolated from a malignant tumor patient, in particular, a sample obtained from a malignant tumor patient, and contains malignant tumor cells. Examples of biological samples include body fluids (e.g., blood, urine, etc.), tissues, the extracts thereof, and the cultures of obtained tissues. The

method for obtaining a biological sample can be suitably selected depending on the type of biological sample.

[0065]

The biological sample is prepared by being
5 appropriately treated according to the measurement method. Further, the reagent comprising primer or probe used for the detection may be prepared by a conventional method according to the measurement method therefor.

[0066]

10 In one embodiment of the present invention, a step of detecting the presence of exon 18 and/or exon 21 mutation of EGFR expressed by a malignant tumor patient may be performed before the administration of an antitumor agent to a malignant tumor patient.


15 [0067]

A malignant tumor may include two or more different kinds of malignant tumor cells. Further, two or more malignant tumors may be generated in a single patient. Therefore, a single patient may have different mutations in the same amino acid
20 position of EGFR (for example, the exon 18 mutation is G719A, G719S and G719C exon 18 mutation; E709K and E709A exon 18 mutation) at the same time.

[0068]

The antitumor agent of the present invention comprises,
25 as an active ingredient, (S)-N-(4-amino-6-methyl-5-(quinolin-3-yl)-8,9-dihydropyrimido[5,4-b]indolizin-8-yl)acrylamide (Compound (A)) or a salt thereof. Compound (A) is represented by the following chemical formula.

[0069]



[0070]

The method for producing the compound of the present invention is explained below.

5 Compound A of the present invention may be produced, for example, through the production method disclosed in WO2015/025936A1, the methods described in the Examples, and the like. However, the production method of the compound of the present invention is not limited to these reaction examples.

10 [0071]

When Compound A of the present invention has isomers such as optical isomers, stereoisomers, and tautomers, any of the isomers and mixtures thereof are included within the scope of the compound of the present invention, unless otherwise specified.

15 For example, when Compound A of the present invention has optical isomers, racemic mixtures and the optical isomers separated from a racemic mixture are also included within the scope of the compound of the present invention, unless otherwise specified.

[0072]

20 The salts of Compound A refer to any pharmaceutically acceptable salts; examples include base addition salts and acid addition salts.

[0073]

25 Examples of base addition salts include alkali metal salts such as sodium salts and potassium salts; alkaline earth

metal salts such as calcium salts and magnesium salts; ammonium salts; and organic amine salts such as trimethylamine salts, triethylamine salts, dicyclohexylamine salts, ethanolamine salts, diethanolamine salts, triethanolamine salts, procaine salts, and
5 N,N'-dibenzylethylenediamine salts.
[0074]

Examples of acid addition salts include inorganic acid salts such as hydrochlorides, sulfates, nitrates, phosphates, and perchlorates; organic acid salts such as acetates, formates,
10 maleates, fumarates, tartrates, citrates, ascorbates, and trifluoroacetates; and sulfonates such as methanesulfonates, isethionates, benzenesulfonates, and p-toluenesulfonates.
[0075]

The compound of the present invention and salts thereof
15 also encompass prodrugs thereof. A prodrug refers to a compound that can be converted to the compound of the present invention or a salt thereof through a reaction with an enzyme, gastric acid, or the like, under physiological conditions *in vivo*, i.e., a compound that can be converted to the compound of the present
20 invention or a salt thereof by enzymatic oxidation, reduction, hydrolysis, or the like; or a compound that can be converted to the compound of the present invention or a salt thereof by hydrolysis or the like with gastric acid or the like. Further, the prodrug may be compounds that can be converted to the
25 compound of the present invention or a salt thereof under physiological conditions, such as those described in "*Iyakuhi no Kaihatsu* [Development of Pharmaceuticals]," Vol. 7, Molecular Design, published in 1990 by Hirokawa Shoten Co., pp. 163-198.
[0076]

30 Description of Diseases

Specific examples of tumors targeted in the present invention include, but are not particularly limited to, head and neck cancer, gastrointestinal cancer (esophageal cancer, stomach cancer, duodenal cancer, liver cancer, biliary cancer (e.g.,
35 gallbladder and bile duct cancer), pancreatic cancer, colorectal

cancer (e.g., colon cancer, and rectal cancer), etc.), lung cancer (e.g., non-small-cell lung cancer, small-cell lung cancer, and mesothelioma), breast cancer, genital cancer (ovarian cancer, uterine cancer (e.g., cervical cancer, and endometrial cancer), etc.), urological cancer (e.g., kidney cancer, bladder cancer, prostate cancer, and testicular tumor), hematopoietic tumor (e.g., leukemia, malignant lymphoma, and multiple myeloma), osteosarcoma, soft-tissue sarcoma, skin cancer, brain tumor, and the like. Preferable examples include lung cancer, breast cancer, head and neck cancer, brain tumor, uterine cancer, digestive organ cancer, hematopoietic tumor, or skin cancer. Lung cancer is particularly preferable.

[0077]

When Compound A or a salt thereof are used as a pharmaceutical agent, a pharmaceutical carrier can be added, if required, thereby forming a suitable dosage form according to prevention and treatment purposes. Examples of the dosage form include oral preparations, injections, suppositories, ointments, patches, and the like. Oral preparations are preferable. Such dosage forms can be formed by methods conventionally known to persons skilled in the art.

[0078]

In one embodiment, the antitumor agent of the present invention is provided as a pharmaceutical composition comprising Compound A or a salt thereof, and a pharmaceutically acceptable carrier.

[0079]

As the pharmaceutically acceptable carrier, various conventional organic or inorganic carrier materials used as preparation materials may be blended as an excipient, binder, disintegrant, lubricant, or colorant in solid preparations; or as a solvent, solubilizing agent, suspending agent, isotonicizing agent, buffer, or soothing agent in liquid preparations. Moreover, pharmaceutical preparation additives, such as antiseptics, antioxidants, colorants, sweeteners, and stabilizers, may also be

used, if required.

[0080]

Oral solid preparations are prepared as follows. After an excipient is added optionally with an excipient, binder,
5 disintegrant, lubricant, colorant, taste-masking or flavoring agent, etc., to the compound of the present invention, the resulting mixture is formulated into tablets, coated tablets, granules, powders, capsules, or the like by ordinary methods.

[0081]

10 Examples of excipients include lactose, sucrose, D-mannitol, glucose, starch, calcium carbonate, kaolin, microcrystalline cellulose, and silicic acid anhydride. Examples of binders include water, ethanol, 1-propanol, 2-propanol, simple syrup, liquid glucose, liquid α -starch, liquid gelatin, D-
15 mannitol, carboxymethyl cellulose, hydroxypropyl cellulose, hydroxypropyl starch, methyl cellulose, ethyl cellulose, shellac, calcium phosphate, polyvinylpyrrolidone, and the like. Examples of disintegrators include dry starch, sodium alginate, powdered agar, sodium hydrogen carbonate, calcium carbonate, sodium lauryl
20 sulfate, stearic acid monoglyceride, lactose, and the like. Examples of lubricants include purified talc, sodium stearate, magnesium stearate, borax, polyethylene glycol, and the like. Examples of colorants include titanium oxide, iron oxide, and the like. Examples of taste-masking or flavoring agents include
25 sucrose, bitter orange peel, citric acid, tartaric acid, and the like.

[0082]

When a liquid preparation for oral administration is prepared, a taste-masking agent, a buffer, a stabilizer, a
30 flavoring agent, and the like may be added to the compound of the present invention; and the resulting mixture may be formulated into an oral liquid preparation, syrup, elixir, etc., according to an ordinary method.

[0083]

35 When an injection agent is prepared, a pH regulator, a

buffer, a stabilizer, an isotonizing agent, a local anesthetic, and the like, may be added to the compound of the present invention; and the mixture may be formulated into a subcutaneous, intramuscular, or intravenous injection according to an ordinary method.

[0084]

Examples of the pH adjuster and the buffer used herein include sodium citrate, sodium acetate, and sodium phosphate. Examples of the stabilizer include sodium pyrosulfite, EDTA, thioglycolic acid, and thiolactic acid. Examples of the local anesthetic include procaine hydrochloride and lidocaine hydrochloride. Examples of the tonicity agent include sodium chloride, glucose, D-mannitol, and glycerol.

[0085]

When a suppository is prepared, pharmaceutically acceptable carriers known in the related field, such as polyethylene glycol, lanolin, cacao butter, and fatty acid triglyceride; and as necessary, surfactants such as Tween 80 (registered trademark), may be added to Compound A, and the resulting mixture may be formulated into a suppository according to an ordinary method.

[0086]

When an ointment is prepared, a commonly used base, stabilizer, wetting agent, preservative, and the like, may be blended into Compound A, as necessary; and the obtained mixture is mixed and formulated into an ointment according to an ordinary method.

[0087]

Examples of the base include liquid paraffin, white petrolatum, white beeswax, octyl dodecyl alcohol, and paraffin.

[0088]

Examples of the preservative include methyl paraoxybenzoate, ethyl paraoxybenzoate, and propyl paraoxybenzoate.

[0089]

When a patch is prepared, the above-described ointment, cream, gel, paste, or the like, may be applied to an ordinary substrate according to an ordinary method.

[0090]

5 Examples of substrates include woven fabrics or non-woven fabrics comprising cotton, staple fibers, or chemical fibers; and films or foam sheets of soft vinyl chloride, polyethylene, polyurethane, etc., may also be used.

[0091]

10 The amount of Compound A to be incorporated in each of such dosage unit forms depends on the condition of the patient to whom the compound is administered, the dosage form thereof, etc. In general, in the case of an oral agent, the amount of the compound is preferably 0.05 to 1000 mg per dosage unit form. In
15 the case of an injection, the amount of the compound is preferably 0.01 to 500 mg per dosage unit form; and in the case of a suppository, the amount of the compound is preferably 1 to 1000 mg per dosage unit form.

[0092]

20 Further, the daily dose of the medicine in such a dosage form depends on the condition, body weight, age, sex, etc., of the patient, and cannot be generalized or limited. Usually, the daily dose for an adult (body weight: 50 kg) of Compound A may generally be 0.05 to 5000 mg, and preferably 0.1 to 1000 mg;
25 and is preferably administered in one dose, or in two to three divided doses, per day.

[0093]

 The present invention also provides a method for treating a malignant tumor patient, comprising the step of
30 administering Compound A or a salt thereof to a malignant tumor patient expressing EGFR having exon 18 and/or exon 21 mutation.

[0094]

 The present invention also provides Compound A or a salt thereof for treating a malignant tumor patient expressing
35 EGFR having exon 18 and/or exon 21 mutation.

[0095]

The present invention also provides use of Compound A or a salt thereof for treating a malignant tumor patient expressing EGFR having exon 18 and/or exon 21 mutation.

5 [0096]

The present invention also provides use of Compound A or a salt thereof for the production of a pharmaceutical agent for treating a malignant tumor patient expressing EGFR having exon 18 and/or exon 21 mutation.

10 [0097]

The present invention also provides a pharmaceutical composition for treating a malignant tumor patient expressing EGFR having exon 18 and/or exon 21 mutation, the pharmaceutical composition comprising Compound A or a salt thereof, and a pharmaceutically acceptable carrier.

[0098]

The present invention also provides a method for predicting therapeutic effects of chemotherapy using an antitumor agent comprising, as an active ingredient, Compound A or a salt thereof in a malignant tumor patient, the method comprising steps (1) and (2) below:

- (1) a step of detecting the presence or absence of mutation of EGFR gene contained in a biological sample obtained from the patient; and
- 25 (2) a step of predicting that the chemotherapy is highly likely to exhibit sufficient therapeutic effects to the patient when the results of the detection in step (1) found that the EGFR gene has exon 18 and/or exon 21 mutation.

[0099]

30 The present invention also provides a method for treating a malignant tumor patient, comprising steps (1) to (3) below:

- (1) a step of detecting the presence or absence of mutation of EGFR gene contained in a biological sample obtained from the patient; and
- 35

(2) a step of predicting that chemotherapy using an antitumor agent comprising, as an active ingredient, (S)-N-(4-amino-6-methyl-5-(quinolin-3-yl)-8,9-dihydropyrimido[5,4-b]indolizin-8-yl)acrylamide or a salt thereof is highly likely to exhibit sufficient therapeutic effects with respect to the patient when the results of the detection in step (1) found that the EGFR gene has exon 18 and/or exon 21 mutation; and

(3) a step of administering (S)-N-(4-amino-6-methyl-5-(quinolin-3-yl)-8,9-dihydropyrimido[5,4-b]indolizin-8-yl)acrylamide or a salt thereof to a malignant tumor patient who was predicted highly likely to sufficiently respond to the chemotherapy using an antitumor agent comprising, as an active ingredient, (S)-N-(4-amino-6-methyl-5-(quinolin-3-yl)-8,9-dihydropyrimido[5,4-b]indolizin-8-yl)acrylamide or a salt thereof, in step (2).

[0100]

The base sequence of EGFR gene is publicly known. The GenBank accession number of the base sequence of cDNA is NM_005228.4.

[0101]

The "therapeutic effects" can be evaluated by tumor shrinkage effects, relapse-suppressing effects, life-prolonging effects, and the like. The relapse-suppressing effects may be shown as degree of the extension of non-relapse period or the degree of the improvement in relapse rate; and the life-prolonging effects may be shown as the degree of the entire survival time or the degree of the extension of the median of progression-free survival, or the like. The "sufficient therapeutic effects" of the chemotherapy using an antitumor agent comprising, as an active ingredient, Compound A or a salt thereof means, for example, that superior therapeutic effects are obtained by the administration of the antitumor agent comprising, as an active ingredient, Compound A or a salt thereof, such as extension of survival time, suppression of relapse, and the like, compared with non-administration.

Examples

[0102]

The following describes the present invention in more
5 detail with reference to the following Test Examples. However,
the present invention is not limited to these Examples (Test
Examples).

[0103]

Test Example 1: In Vitro Drug Efficacy Test

10 Evaluation Results of Intracellular Phosphorylation in Mutant
EGFR Forced Expression System Using HEK293 Cells (Inhibitory
Activity)

The intracellular target inhibitory activity of
compounds was evaluated based on the following as an index:
15 intracellular EGFR phosphorylation in a mutant EGFR forced
expression system using Jump-In (trademark) Grip (trademark)
HEK293 cells (Thermo Fisher Scientific Inc.) (which may be
hereinafter referred to as "HEK293 cells").

HEK293 cells were maintained in D-MEM with GlutaMAX
20 (trademark)-I (high-glucose) (Thermo Fisher Scientific Inc.) that
contained 10% dialyzed FBS and 100 U/mL penicillin/100 µg/mL
streptomycin (Thermo Fisher Scientific Inc.); and a pJTI™ R4 DEST
CMV pA vector to which a human EGFR gene (G719A, G719S, G719C,
E709K, E709A, L861Q, G719A+T790M, or L861Q+T790M; the symbol "+"
25 indicates that both mutations are contained) was encoded was
introduced into HEK293 cells, together with Opti-MEM (trademark)
I (Thermo Fisher Scientific Inc.), using a ViaFect (trademark)
Transfection Reagent (Promega Corporation).

HEK293 cells expressing mutant human EGFR were seeded
30 in each well of a 384-well flat-bottom microplate such that the
cell count per well was 10,000, and incubated in a 5% CO₂ gas-
containing incubator at 37°C for 1 day. Compound A, erlotinib,
afatinib, and osimertinib (erlotinib, afatinib, and osimertinib
may be each hereinafter referred to as a "comparative compound")
35 were individually dissolved in DMSO, and diluted with DMSO or the

medium used for suspending the cells. The solutions were then individually added to each well of the culture plate of the cells, and the cells were incubated in a 5% CO₂ gas-containing incubator at 37°C for 6 hours. After incubation, the cells were immobilized
5 using 20% neutral buffered formalin (Wako Pure Chemical Industries, Ltd.), and blocked by an Odyssey (trademark) blocking buffer (PBS) (M&S TechnoSystems Inc.). The cells were then reacted with a primary antibody (EGFR Monoclonal Antibody (R19/48MIX) #AHR5062) (Thermo Fisher Scientific Inc.) that was
10 diluted with an Odyssey (trademark) blocking buffer (PBS) to 1/200, and a Phospho-EGFR Receptor (Tyr1068) Antibody #2234L (CST)), followed by subjecting the cells to permeation at 4°C overnight. The following day, the cells were reacted with a secondary antibody (IRDye 800CW Goat aRabbit #926-32211 and IRDye
15 680RD Goat aMouse #926-68070 (M&S TechnoSystems Inc.)) diluted with an Odyssey (trademark) blocking buffer (PBS) to 1/800, and subjected to permeation at room temperature for 1 hour. The fluorescence intensity (which may be hereinafter referred to as "FI") was detected with an Odyssey Infrared Imaging System (LI-
20 COR Bioscience) at a fluorescence wavelength of 800 nm and 700 nm.
[0104]

The value obtained by deducting the FI of a well without the cells from the FI detected at a fluorescence wavelength of 800 nm or 700 nm is referred to as FI (800, EGFR)-
25 Blank (for 800 nm) and FI (700, p-EGFR)-Blank (for 700 nm). The value obtained by dividing FI (700, p-EGFR)-Blank of each well by FI (800, EGFR)-Blank was determined to be FI (p-EGFR/EGFR). The phosphorylated EGFR inhibitory rate was calculated using the following formula to determine the concentration of the test
30 compounds at which phosphorylated EGFR was inhibited by 50% (IC₅₀ (μM)). Table 1 illustrates the results.
[0105]

Phosphorylated EGFR Inhibitory Rate (%) = $T/C \times 100$
T: FI (p-EGFR/EGFR) of a well to which a test compound was added.
35 C: FI (p-EGFR/EGFR) of a well to which a test compound was not

added.

[0106]

As is clear from Table 1, compound A exhibited high inhibitory activity against intracellular phosphorylation of exon 18 or exon 21 mutant EGFR; and the activity was higher than that of erlotinib or osimertinib, and was equivalent to that of afatinib. The inhibitory activity of compound A against exon 18 or exon 21 mutant EGFR was higher than that of afatinib in the presence of T790M mutation, which is an acquired resistant mutation.

[0107]

Table 1

	Compound A of the Present Invention	Osimertinib	Afatinib	Erlotinib
G719A	20.5	237.0	20.0	>1000
G719S	22.7	251.9	16.9	539.7
G719C	9.3	102.0	15.3	98.3
E709K	86.0	424.2	14.0	451.4
E709A	103.5	548.9	19.2	478.1
L861Q	40.5	143.1	24.7	608.1
G719A+T790M	31.3	132.1	110.2	>1000
L861Q+T790M	81.8	132.4	177.6	>1000

[0108]

Test Example 2

Evaluation of Cell Growth Inhibitory Effect Against Wild-Type EGFR- and Mutant EGFR-Expressing Cell Line (In Vitro)

The inhibitory activity of compounds against wild-type EGFR and mutant EGFR was evaluated using Ba/F3 cells, which are mouse B-lymphocyte precursor cell line to which a human EGFR gene was introduced. Ba/F3 cells were maintained in an RPMI-1640 medium (Thermo Fisher Scientific Inc.) containing 10% fetal bovine serum (FBS), 100 U/mL penicillin/100 µg/mL streptomycin (Thermo Fisher Scientific Inc.), and 1 ng/mL mouse interleukin-3 (mIL-3) (CST). A PB-CMV-MCS-EF1-GFP+Puro vector or PB-CMV-MCS-EF1-RFP+Puro vector into which a human EGFR gene (wild-type (WT), G719A, or L861Q) was encoded was introduced into the cells,

together with a Super PiggyBac Transposase expression vector, by electroporation using an Amaxa (trademark) Cell Line Nucleofector (trademark) Kit V, followed by selection using puromycin (SIGMA). Ba/F3 cells expressing wild-type EGFR (which may be hereinafter referred to as "Ba/F3-EGFR_WT") exhibited mIL-3-independent growth in the presence of 50 ng/mL EGF (R&D Systems); and Ba/F3 cells expressing exon 18 or exon 21 active mutant EGFR (which may be hereinafter referred to as "Ba/F3-EGFR G719A," and "Ba/F3-EGFR L861Q") exhibited mIL-3-independent growth in the absence of EGF.

[0109]

To evaluate a cell growth inhibitory effect, Ba/F3-EGFR_WT cells were suspended in an RPMI-1640 medium containing 10% FBS, 100 U/mL penicillin, 100 µg/mL streptomycin, and 50 ng/mL EGF; and the cell suspension was seeded in each well of a 96-well flat-bottom microplate such that the cell count per well was 30,000. Ba/F3-EGFR G719A cells and Ba/F3-EGFR L861Q cells were suspended in respective RPMI-1640 mediums containing 10% FBS, 100 U/mL penicillin, and 100 µg/mL streptomycin; and the cell suspensions were individually seeded in each well of a 96-well flat-bottom microplate such that the cell count per well was 15,000. Subsequently, compound A, gefitinib, erlotinib, afatinib, and osimertinib (gefitinib, erlotinib, afatinib, and osimertinib may be each hereinafter referred to as a "comparative compound") were individually dissolved in DMSO, and diluted with DMSO or the medium used for suspending the cells. The solutions were then added to each well of the culture plate of the cells, and the cells were incubated in a 5% CO₂ gas-containing incubator at 37°C for 3 days. The cell count after incubation was measured by a CellTiter-Glo (trademark) Luminescent Cell Viability Assay (Promega Corporation) in accordance with the manufacturer's recommended protocol. The growth inhibition rate was calculated using the following formula to determine the concentration of each test compound for 50% inhibition (IC₅₀ (µM)).

[0110]

Growth Inhibitory Rate (%) = T/C × 100

T: the luminescence intensity of a well to which a test compound was added.

C: the luminescence intensity of a well to which a test compound was not added.

5 [0111]

The IC₅₀ ratio between wild-type EGFR and G719A mutant EGFR, or between wild-type EGFR and L861Q mutant EGFR was determined using the following formula. Table 2 illustrates the results.

10 [0112]

IC₅₀ Ratio = IC₅₀ (WT)/IC₅₀ (G719A or L861Q)

[0113]

As is clear from Table 2, compound A exhibited selective inhibitory activity against G719A mutation and L861Q mutation.

15

[0114]

Table 2

	IC ₅₀ (nM)			IC ₅₀ Ratio	
	WT	G719A	L861Q	WT/G719A	WT/L861Q
Compound A of the Present Invention	597.3	9.0	19.0	66.4	31.4
Osimertinib	378.7	209.0	47.0	1.8	8.1
Afatinib	17.7	3.3	2.3	5.3	7.6
Erlotinib	778.3	336.3	353.0	2.3	2.2

[0115]

20 Test Example 3: In Vivo Drug Efficacy Test

Evaluation of Antitumor Effect on Mouse Model Subcutaneously Transplanted with G719A Mutant EGFR-Expressing Cell Line

The evaluation on mouse models subcutaneously transplanted with G719A mutant EGFR-expressing cell line was performed using NIH-3T3 cells, which are mouse fibroblast cell line to which a human EGFR gene was introduced. NIH-3T3 cells were maintained in a D-MEM (high-glucose) medium (Wako Pure Chemical Industries, Ltd.) containing 10% newborn calf serum (NBCS), 1,500 mg/L sodium hydrogen carbonate, and 100 U/mL

25

penicillin/100 µg/mL streptomycin (Thermo Fisher Scientific Inc.); and a PB-CMV-MCS-EF1-RFP+Puro vector into which a human EGFR gene (G719A) was encoded was introduced to the cells, together with a Super PiggyBac Transposase expression vector, by
5 electroporation using an Amaxa (trademark) Cell Line Nucleofector (trademark) Kit R, followed by selection using puromycin (SIGMA). NIH-3T3 cells expressing exon 18 mutant EGFR (which may be hereinafter referred to as "NIH3T3-EGFR G719A") exhibited growth in the absence of EGF under 1% NBCS conditions.

10 [0116]

In evaluation using mouse models subcutaneously transplanted with G719A mutant EGFR-expressing cell line, nude mice were subcutaneously transplanted with NIH3T3-EGFR G719A cells into which a mutant human EGFR was introduced. At the point
15 at which the tumor volume of the tumor engrafted in the mice grew to about 100 to 200 mm³, the mice were allocated into groups, 5 or 6 mice for each group, by stratified randomization such that the average tumor volume between the groups was uniform. The mice were then orally administered compound A, afatinib, or
20 osimertinib once daily for 14 consecutive days.

[0117]

The dose of afatinib was 20 mg/kg/day, which is the maximum tolerated dose (the maximum dose at which the weight loss during a dosing period is less than 20%), for 14 days, the dosing
25 period of this test. The dose of osimertinib was 25 mg/kg/day, which is a clinically efficacious dose. For compound A, three types of doses were set: 200 mg/kg/day (maximum tolerated dose), 100 mg/kg/day, and 50 mg/kg/day. The maximum tolerated dose was determined in accordance with the "Guidelines Involving
30 Experimental Neoplasia Proposals in Mice and Rats" of the National Cancer Institute (NCI), from a humanitarian perspective.

[0118]

To compare the changes in growth of tumor over time due to administration of the individual test compounds, the tumor
35 volume (which may be hereinafter referred to as "TV") was used as

an index. For a toxicity index, the body weight was measured over time, and the body weight change (which may be hereinafter referred to as "BWC (%)") from the day on which the mice were divided into groups was calculated in accordance with the following formula.

[0119]

$$\text{BWC (\%)} = (\text{the body weight measured on body weight measurement day}) / (\text{the body weight on the day mice were divided into groups})$$

[0120]

When the difference in the average TV between the control group and the group administered with a test compound on the final evaluation day was statistically significant (Dunnett's test, $p < 0.05$), and the value of treatment/control (T/C) calculated using the following formula was less than 100, the test compound was determined to be effective. Such a case is indicated by the symbol "*" in the figures.

[0121]

$$\text{T/C (\%)} = (\text{the average TV of the group administered with a test compound}) / (\text{the average TV of the control group}) \times 100$$

[0122]

As is clear from the results illustrated in Fig. 1, compound A of the present invention exhibited a remarkable antitumor effect on G719A mutant EGFR-expressing cell line subcutaneously transplanted into nude mice, accompanied by tumor regression. The effect was also higher than that of afatinib or osimertinib. The mice did not show symptoms, such as serious weight loss, as illustrated in Fig. 2, abnormal feces, or abnormal skin.

[0123]

Test Example 4: In Vitro Drug Efficacy Test

Evaluation of Cell Growth Inhibitory Effect on T790M Mutant EGFR-Expressing Cell Line (In Vitro)

The inhibitory activity of a compound on T790M mutant EGFR was evaluated using Ba/F3 cells, which are mouse B-lymphocyte precursor cell line into which a human EGFR gene was

introduced. Ba/F3 cells were maintained in an RPMI-1640 medium (Thermo Fisher Scientific Inc.) containing 10% fetal bovine serum (FBS), 100 U/mL penicillin/100 µg/mL streptomycin (Thermo Fisher Scientific Inc.), and 1 ng/mL mouse interleukin-3 (mIL-3) (CST);
5 and a PB-CMV-MCS-EF1-GFP+Puro vector or PB-CMV-MCS-EF1-RFP+Puro vector into which a human EGFR gene (wild-type (WT), G719A+T790M, or L861Q+T790M) was encoded was introduced into the cells, together with a Super PiggyBac Transposase expression vector, by electroporation using an Amaxa (trademark) Cell Line Nucleofector
10 (trademark) Kit V, followed by selection using puromycin (SIGMA). Ba/F3 cells expressing G719A+T790M mutant EGFR or L861Q+T790M mutant EGFR (which may be hereinafter referred to as "Ba/F3-EGFR G719A+T790M," and "Ba/F3-EGFR L861Q+T790M") exhibited mIL-3-independent growth in the absence of EGF.

15 [0124]

To evaluate a cell growth inhibitory effect, Ba/F3-EGFR G719A+T790M cells and Ba/F3-EGFR L861Q+T790M cells were individually suspended in an RPMI-1640 medium containing 10% FBS, 100 U/mL penicillin, and 100 µg/mL streptomycin; and the cell
20 suspensions were individually seeded in each well of a 96-well flat-bottom microplate such that the cell count per well was 15,000. Subsequently, compound A, gefitinib, erlotinib, afatinib, and osimertinib (gefitinib, erlotinib, afatinib, and osimertinib may be each hereinafter referred to as a "comparative compound")
25 were individually dissolved in DMSO, and diluted with DMSO or the medium used for suspending the cells. These solutions were then added to each well of the culture plate of the cells, and incubated in a 5% CO₂ gas-containing incubator at 37°C for 3 days. The cell count after incubation was measured using a CellTiter-
30 Glo (trademark) Luminescent Cell Viability Assay (Promega Corporation), in accordance with the manufacturer's recommended protocol. The growth inhibition rate was calculated using the following formula to determine the concentration of each test compound for 50% inhibition (IC₅₀ (µM)).

35 [0125]

Growth Inhibition Rate (%) = $T/C \times 100$

T: the luminescence intensity of a well to which a test compound was added.

C: the luminescence intensity of a well to which the test compound was not added.

[0126]

Additionally, the IC₅₀ ratio between wild-type EGFR and G719A+T790M mutant EGFR, or between wild-type EGFR and L861Q+T790M mutant EGFR was calculated using the following formula. Table 3 illustrates the results.

[0127]

IC₅₀ Ratio = IC₅₀ (WT)/IC₅₀ (G719A+T790M or L861Q+T790M)

[0128]

As is clear from Table 3, compound A exhibited selective inhibitory activity against G719A+T790M mutation and L861Q+T790M mutation.

[0129]

Table 3

	IC ₅₀ (nM)		IC ₅₀ Ratio	
	G719A+T790M	L861Q+T790M	WT/ G719A+T790M	WT/ L861Q+T790M
Compound A of the Present Invention	15.8	37.5	37.9	15.9
Osimertinib	89.4	36.4	4.2	10.4
Afatinib	68.2	168.7	0.3	0.1
Erlotinib	8030.0	6424.2	0.1	0.1

[0130]

Test Example 5: In Vivo Drug Efficacy Test

Evaluation of Antitumor Effect on Mouse Model Subcutaneously Transplanted with G719A+T790M Mutant EGFR-Expressing Cell Line

The evaluation of mouse models subcutaneously transplanted with G719A+T790M mutant EGFR-expressing cell line was performed using NIH-3T3 cells, which are mouse fibroblast cell line into which a human EGFR gene was introduced. NIH-3T3 cells were maintained in a D-MEM (high-glucose) medium (Wako Pure Chemical Industries, Ltd.) containing 10% newborn calf serum

(NBCS), 1,500 mg/L sodium hydrogen carbonate, and 100 U/mL penicillin/100 µg/mL streptomycin (Thermo Fisher Scientific Inc.); and a PB-CMV-MCS-EF1-RFP+Puro vector into which a human EGFR gene (G719A+T790M) was encoded was introduced to the cells, together with a Super PiggyBac Transposase expression vector, by electroporation using an Amaxa (trademark) Cell Line Nucleofector (trademark) Kit R, followed by selection using puromycin (SIGMA). NIH-3T3 cells expressing exon 18 mutant EGFR (which may be hereinafter referred to as "NIH3T3-EGFR G719A+T790M") exhibited growth in the absence of EGF under 1% NBCS conditions.

[0131]

In evaluation using mouse models subcutaneously transplanted with G719A+T790M mutant EGFR-expressing cell line, nude mice were subcutaneously transplanted with NIH3T3-EGFR G719A+T790M cells into which mutant human EGFR was introduced. At the point at which the tumor volume of the tumor engrafted in the mice grew to about 100 to 300 mm³, the mice were allocated into groups, 5 mice for each group, by stratified randomization such that the average tumor volume between the groups was uniform. The mice were then orally administered compound A of the present invention or afatinib once daily on consecutive days.

[0132]

The dose of afatinib was 20 mg/kg/day, which is the maximum tolerated dose (the maximum dose at which the weight loss during a dosing period is less than 20%), for 14 days, the dosing period of this test. For compound A of the present invention, three types of doses were set: 200 mg/kg/day (maximum tolerated dose), 100 mg/kg/day, and 50 mg/kg/day. The maximum tolerated dose was determined in accordance with the "Guidelines Involving Experimental Neoplasia Proposals in Mice and Rats" of the National Cancer Institute (NCI), from a humanitarian perspective.

[0133]

To compare the changes in growth of tumor over time due to administration of the individual test compounds, the tumor volume (which may be hereinafter referred to as "TV") was used as

an index. Fig. 3 illustrates changes in TV over time. For a toxicity index, the body weight was measured over time, and the body weight change (which may be hereinafter referred to as "BWC (%)") from the day on which the mice were divided into groups was calculated in accordance with the following formula. Fig. 4 illustrates changes in body weight over time.

$$\text{BWC (\%)} = (\text{the body weight measured on body weight measurement day}) / (\text{the body weight on the day mice were divided into groups})$$

[0134]

A Dunnett's test was performed using the average TV on the final evaluation day as an index. When the difference in the average TV between the control group and the group administered with a test compound was statistically significant ($p < 0.05$), and the value of treatment/control (T/C) calculated using the following formula was less than 100, the test compound was determined to be effective. Such a case is indicated by the symbol "*" in Fig. 3 and Table 4 (*: $p < 0.05$, **: $p < 0.01$, ***: $p < 0.001$). Additionally, when the difference in the average TV between the group administered with compound A of the present invention and the group administered with afatinib was statistically significant ($p < 0.05$), and the T/C value of the group administered with compound A of the present invention was smaller than the T/C value of the group administered with afatinib, compound A was determined to have a higher antitumor effect than afatinib. Such a case is indicated by the symbol "#" in Fig. 3 and Table 4 (#: $p < 0.05$, ##: $p < 0.01$, ###: $p < 0.001$).

[0135]

$$\text{T/C (\%)} = (\text{the average TV of the group administered with a test compound}) / (\text{the average TV of the control group}) \times 100$$

[0136]

As is clear from the results illustrated in Fig. 3, compound A of the present invention exhibited a significant antitumor effect on G719A+T790M mutant EGFR-expressing cell line subcutaneously transplanted into nude mice. Additionally, as

shown in Table 4, the effect was higher than that of afatinib, without symptoms such as serious weight loss (as illustrated in Fig. 4), abnormal feces, or abnormal skin in mice.

[0137]

5 Table 4

Compound	Dose (mg/kg)	T/C (%)	<i>p</i> value vs	
			Control	Afatinib
Control	—	100.0	—	—
Compound A of the Present Invention	50	47.7	***	N.S.
Compound A of the Present Invention	100	36.4	***	##
Compound A of the Present Invention	200	28.0	***	###
Afatinib	20	57.4	***	—

N.S.: No Significant Difference

***: $p < 0.001$ (Dunnett's test vs Control Group)

##: $p < 0.01$ (Dunnett's test vs Afatinib Group)

10 ###: $p < 0.001$ (Dunnett's test vs Afatinib Group)

SEQUENCE LISTING

<110> TAIHO PHARMACEUTICAL CO., LTD.

5 <120> Exon 18 and/or 21 EGFR Mutant-selective Inhibitor

<130> P18-145WO

<150> JP 2017-168606

10 <151> 2017-09-01

<160> 1

<170> PatentIn version 3.5

15

<210> 1

<211> 1210

<212> PRT

<213> Homo sapiens

20

<400> 1

Met Arg Pro Ser Gly Thr Ala Gly Ala Ala Leu Leu Ala Leu Leu Ala
1 5 10 15

25

Ala Leu Cys Pro Ala Ser Arg Ala Leu Glu Glu Lys Lys Val Cys Gln
20 25 30

30

Gly Thr Ser Asn Lys Leu Thr Gln Leu Gly Thr Phe Glu Asp His Phe
35 40 45

35 Leu Ser Leu Gln Arg Met Phe Asn Asn Cys Glu Val Val Leu Gly Asn

	Ser	Cys	Pro	Asn	Gly	Ser	Cys	Trp	Gly	Ala	Gly	Glu	Glu	Asn	Cys	Gln
	195					200					205					
5	Lys	Leu	Thr	Lys	Ile	Ile	Cys	Ala	Gln	Gln	Cys	Ser	Gly	Arg	Cys	Arg
	210					215					220					
10	Gly	Lys	Ser	Pro	Ser	Asp	Cys	Cys	His	Asn	Gln	Cys	Ala	Ala	Gly	Cys
	225		230					235					240			
15	Thr	Gly	Pro	Arg	Glu	Ser	Asp	Cys	Leu	Val	Cys	Arg	Lys	Phe	Arg	Asp
	245					250					255					
20	Glu	Ala	Thr	Cys	Lys	Asp	Thr	Cys	Pro	Pro	Leu	Met	Leu	Tyr	Asn	Pro
	260					265					270					
25	Thr	Thr	Tyr	Gln	Met	Asp	Val	Asn	Pro	Glu	Gly	Lys	Tyr	Ser	Phe	Gly
	275					280					285					
30	Ala	Thr	Cys	Val	Lys	Lys	Cys	Pro	Arg	Asn	Tyr	Val	Val	Thr	Asp	His
	290					295					300					
35	Gly	Ser	Cys	Val	Arg	Ala	Cys	Gly	Ala	Asp	Ser	Tyr	Glu	Met	Glu	Glu
	305		310					315					320			
40	Asp	Gly	Val	Arg	Lys	Cys	Lys	Lys	Cys	Glu	Gly	Pro	Cys	Arg	Lys	Val
	325					330					335					

	Cys	Asn	Gly	Ile	Gly	Ile	Gly	Glu	Phe	Lys	Asp	Ser	Leu	Ser	Ile	Asn	
			340				345				350						
5	Ala	Thr	Asn	Ile	Lys	His	Phe	Lys	Asn	Cys	Thr	Ser	Ile	Ser	Gly	Asp	
			355				360				365						
10	Leu	His	Ile	Leu	Pro	Val	Ala	Phe	Arg	Gly	Asp	Ser	Phe	Thr	His	Thr	
			370				375				380						
	Pro	Pro	Leu	Asp	Pro	Gln	Glu	Leu	Asp	Ile	Leu	Lys	Thr	Val	Lys	Glu	
15	385				390				395				400				
	Ile	Thr	Gly	Phe	Leu	Leu	Ile	Gln	Ala	Trp	Pro	Glu	Asn	Arg	Thr	Asp	
				405				410					415				
20	Leu	His	Ala	Phe	Glu	Asn	Leu	Glu	Ile	Ile	Arg	Gly	Arg	Thr	Lys	Gln	
			420					425				430					
25	His	Gly	Gln	Phe	Ser	Leu	Ala	Val	Val	Ser	Leu	Asn	Ile	Thr	Ser	Leu	
			435					440				445					
30	Gly	Leu	Arg	Ser	Leu	Lys	Glu	Ile	Ser	Asp	Gly	Asp	Val	Ile	Ile	Ser	
			450					455				460					
	Gly	Asn	Lys	Asn	Leu	Cys	Tyr	Ala	Asn	Thr	Ile	Asn	Trp	Lys	Lys	Leu	
35	465				470				475					480			

Phe Gly Thr Ser Gly Gln Lys Thr Lys Ile Ile Ser Asn Arg Gly Glu
485 490 495
5

Asn Ser Cys Lys Ala Thr Gly Gln Val Cys His Ala Leu Cys Ser Pro
500 505 510
10

Glu Gly Cys Trp Gly Pro Glu Pro Arg Asp Cys Val Ser Cys Arg Asn
515 520 525

15 Val Ser Arg Gly Arg Glu Cys Val Asp Lys Cys Asn Leu Leu Glu Gly
530 535 540

Glu Pro Arg Glu Phe Val Glu Asn Ser Glu Cys Ile Gln Cys His Pro
20 545 550 555 560

Glu Cys Leu Pro Gln Ala Met Asn Ile Thr Cys Thr Gly Arg Gly Pro
565 570 575
25

Asp Asn Cys Ile Gln Cys Ala His Tyr Ile Asp Gly Pro His Cys Val
580 585 590
30

Lys Thr Cys Pro Ala Gly Val Met Gly Glu Asn Asn Thr Leu Val Trp
595 600 605

35 Lys Tyr Ala Asp Ala Gly His Val Cys His Leu Cys His Pro Asn Cys

	610		615			620											
	Thr	Tyr	Gly	Cys	Thr	Gly	Pro	Gly	Leu	Glu	Gly	Cys	Pro	Thr	Asn	Gly	
5	625				630				635					640			
	Pro	Lys	Ile	Pro	Ser	Ile	Ala	Thr	Gly	Met	Val	Gly	Ala	Leu	Leu	Leu	
10				645					650				655				
	Leu	Leu	Val	Val	Ala	Leu	Gly	Ile	Gly	Leu	Phe	Met	Arg	Arg	Arg	His	
15				660					665				670				
	Ile	Val	Arg	Lys	Arg	Thr	Leu	Arg	Arg	Leu	Leu	Gln	Glu	Arg	Glu	Leu	
			675					680					685				
20	Val	Glu	Pro	Leu	Thr	Pro	Ser	Gly	Glu	Ala	Pro	Asn	Gln	Ala	Leu	Leu	
		690						695					700				
	Arg	Ile	Leu	Lys	Glu	Thr	Glu	Phe	Lys	Lys	Ile	Lys	Val	Leu	Gly	Ser	
25	705				710					715				720			
	Gly	Ala	Phe	Gly	Thr	Val	Tyr	Lys	Gly	Leu	Trp	Ile	Pro	Glu	Gly	Glu	
30				725						730				735			
	Lys	Val	Lys	Ile	Pro	Val	Ala	Ile	Lys	Glu	Leu	Arg	Glu	Ala	Thr	Ser	
35				740						745				750			

	Pro	Lys	Ala	Asn	Lys	Glu	Ile	Leu	Asp	Glu	Ala	Tyr	Val	Met	Ala	Ser
	755			760			765									
5	Val	Asp	Asn	Pro	His	Val	Cys	Arg	Leu	Leu	Gly	Ile	Cys	Leu	Thr	Ser
	770			775			780									
10	Thr	Val	Gln	Leu	Ile	Thr	Gln	Leu	Met	Pro	Phe	Gly	Cys	Leu	Leu	Asp
	785			790			795			800						
15	Tyr	Val	Arg	Glu	His	Lys	Asp	Asn	Ile	Gly	Ser	Gln	Tyr	Leu	Leu	Asn
				805			810			815						
	Trp	Cys	Val	Gln	Ile	Ala	Lys	Gly	Met	Asn	Tyr	Leu	Glu	Asp	Arg	Arg
				820			825			830						
20																
	Leu	Val	His	Arg	Asp	Leu	Ala	Ala	Arg	Asn	Val	Leu	Val	Lys	Thr	Pro
	835			840			845									
25	Gln	His	Val	Lys	Ile	Thr	Asp	Phe	Gly	Leu	Ala	Lys	Leu	Leu	Gly	Ala
	850			855			860									
	Glu	Glu	Lys	Glu	Tyr	His	Ala	Glu	Gly	Gly	Lys	Val	Pro	Ile	Lys	Trp
30	865			870			875			880						
	Met	Ala	Leu	Glu	Ser	Ile	Leu	His	Arg	Ile	Tyr	Thr	His	Gln	Ser	Asp
				885			890			895						
35																

	Val	Trp	Ser	Tyr	Gly	Val	Thr	Val	Trp	Glu	Leu	Met	Thr	Phe	Gly	Ser	
	900					905					910						
5	Lys	Pro	Tyr	Asp	Gly	Ile	Pro	Ala	Ser	Glu	Ile	Ser	Ser	Ile	Leu	Glu	
	915					920					925						
10	Lys	Gly	Glu	Arg	Leu	Pro	Gln	Pro	Pro	Ile	Cys	Thr	Ile	Asp	Val	Tyr	
	930					935					940						
	Met	Ile	Met	Val	Lys	Cys	Trp	Met	Ile	Asp	Ala	Asp	Ser	Arg	Pro	Lys	
15	945		950					955					960				
	Phe	Arg	Glu	Leu	Ile	Ile	Glu	Phe	Ser	Lys	Met	Ala	Arg	Asp	Pro	Gln	
	965					970					975						
20	Arg	Tyr	Leu	Val	Ile	Gln	Gly	Asp	Glu	Arg	Met	His	Leu	Pro	Ser	Pro	
	980					985					990						
25	Thr	Asp	Ser	Asn	Phe	Tyr	Arg	Ala	Leu	Met	Asp	Glu	Glu	Asp	Met	Asp	
	995					1000					1005						
30	Asp	Val	Val	Asp	Ala	Asp	Glu	Tyr	Leu	Ile	Pro	Gln	Gln	Gly	Phe		
	1010					1015					1020						
	Phe	Ser	Ser	Pro	Ser	Thr	Ser	Arg	Thr	Pro	Leu	Leu	Ser	Ser	Leu		
35	1025					1030					1035						

	Ser	Ala	Thr	Ser	Asn	Asn	Ser	Thr	Val	Ala	Cys	Ile	Asp	Arg	Asn
	1040					1045					1050				
5															
	Gly	Leu	Gln	Ser	Cys	Pro	Ile	Lys	Glu	Asp	Ser	Phe	Leu	Gln	Arg
	1055					1060				1065					
10															
	Tyr	Ser	Ser	Asp	Pro	Thr	Gly	Ala	Leu	Thr	Glu	Asp	Ser	Ile	Asp
	1070					1075				1080					
15	Asp	Thr	Phe	Leu	Pro	Val	Pro	Glu	Tyr	Ile	Asn	Gln	Ser	Val	Pro
	1085					1090				1095					
	Lys	Arg	Pro	Ala	Gly	Ser	Val	Gln	Asn	Pro	Val	Tyr	His	Asn	Gln
20	1100					1105				1110					
	Pro	Leu	Asn	Pro	Ala	Pro	Ser	Arg	Asp	Pro	His	Tyr	Gln	Asp	Pro
	1115					1120				1125					
25															
	His	Ser	Thr	Ala	Val	Gly	Asn	Pro	Glu	Tyr	Leu	Asn	Thr	Val	Gln
	1130					1135				1140					
30															
	Pro	Thr	Cys	Val	Asn	Ser	Thr	Phe	Asp	Ser	Pro	Ala	His	Trp	Ala
	1145					1150				1155					
35	Gln	Lys	Gly	Ser	His	Gln	Ile	Ser	Leu	Asp	Asn	Pro	Asp	Tyr	Gln

1160

1165

1170

5 Gln Asp Phe Phe Pro Lys Glu Ala Lys Pro Asn Gly Ile Phe Lys
 1175 1180 1185

10 Gly Ser Thr Ala Glu Asn Ala Glu Tyr Leu Arg Val Ala Pro Gln
 1190 1195 1200

 Ser Ser Glu Phe Ile Gly Ala
 1205 1210

15

CLAIMS

[Claim 1]

An antitumor agent for treating a malignant tumor patient expressing EGFR having at least one mutation selected from the group consisting of G719X mutation of exon 18, E709X mutation of exon 18, and L861X mutation of exon 21, wherein X represents an arbitrary amino-acid residue, the antitumor agent comprising (S)-N-(4-amino-6-methyl-5-(quinolin-3-yl)-8,9-dihydropyrimido[5,4-b]indolizin-8-yl)acrylamide or a salt thereof.

[Claim 2]

The antitumor agent according to claim 1, wherein the exon 18 mutation is at least one mutation selected from the group consisting of G719A, G719S, G719C, E709K and E709A.

[Claim 3]

The antitumor agent according to claim 1 or 2, wherein the exon 21 mutation is L861Q.

[Claim 4]

The antitumor agent according to any one of claims 1 to 3, wherein the EGFR further has T790M mutation.

[Claim 5]

A method for treating a malignant tumor patient, comprising the step of administering (S)-N-(4-amino-6-methyl-5-(quinolin-3-yl)-8,9-dihydropyrimido[5,4-b]indolizin-8-yl)acrylamide or a salt thereof to a malignant tumor patient expressing EGFR having at least one mutation selected from the group consisting of G719X mutation of exon 18, E709X mutation of exon 18, and L861X mutation of exon 21, wherein X represents an arbitrary amino-acid residue.

[Claim 6]

The method according to claim 5, wherein the exon 18

mutation is at least one mutation selected from the group consisting of G719A, G719S, G719C, E709K and E709A.

[Claim 7]

5 The method according to claim 5 or 6, wherein the exon 21 mutation is L861Q.

[Claim 8]

 The method according to any one of claims 5 to 7,
10 wherein the EGFR further has T790M mutation.

[Claim 9]

 (S)-N-(4-amino-6-methyl-5-(quinolin-3-yl)-8,9-dihydropyrimido[5,4-b]indolizin-8-yl)acrylamide or a salt thereof
15 for treating a malignant tumor patient expressing EGFR having at least one mutation selected from the group consisting of G719X mutation of exon 18, E709X mutation of exon 18, and L861X mutation of exon 21, wherein X represents an arbitrary amino-acid residue.

20

[Claim 10]

 The (S)-N-(4-amino-6-methyl-5-(quinolin-3-yl)-8,9-dihydropyrimido[5,4-b]indolizin-8-yl)acrylamide or a salt thereof according to claim 9, wherein the exon 18 mutation is at least
25 one mutation selected from the group consisting of G719A, G719S, G719C, E709K and E709A.

[Claim 11]

 The (S)-N-(4-amino-6-methyl-5-(quinolin-3-yl)-8,9-dihydropyrimido[5,4-b]indolizin-8-yl)acrylamide or a salt thereof according to claim 9 or 10, wherein the exon 21 mutation is L861Q.

[Claim 12]

 The (S)-N-(4-amino-6-methyl-5-(quinolin-3-yl)-8,9-dihydropyrimido[5,4-b]indolizin-8-yl)acrylamide or a salt thereof
35

according to any one of claims 9 to 11, wherein the EGFR further has T790M mutation.

[Claim 13]

5 Use of (S)-N-(4-amino-6-methyl-5-(quinolin-3-yl)-8,9-dihydropyrimido[5,4-b]indolizin-8-yl)acrylamide or a salt thereof for treating a malignant tumor patient expressing EGFR having at least one mutation selected from the group consisting of G719X mutation of exon 18, E709X mutation of exon 18, and L861X
10 mutation of exon 21, wherein X represents an arbitrary amino-acid residue.

[Claim 14]

 The use according to claim 13, wherein the exon 18
15 mutation is at least one mutation selected from the group consisting of G719A, G719S, G719C, E709K and E709A.

[Claim 15]

 The use according to claim 13 or 14, wherein the exon
20 21 mutation is L861Q.

[Claim 16]

 The use according to any one of claims 13 to 15,
wherein the EGFR further has T790M mutation.

25

[Claim 17]

 Use of (S)-N-(4-amino-6-methyl-5-(quinolin-3-yl)-8,9-dihydropyrimido[5,4-b]indolizin-8-yl)acrylamide or a salt thereof for the production of a pharmaceutical agent for treating a
30 malignant tumor patient expressing EGFR having at least one mutation selected from the group consisting of G719X mutation of exon 18, E709X mutation of exon 18, and L861X mutation of exon 21, wherein X represents an arbitrary amino-acid residue.

35

[Claim 18]

The use according to claim 17, wherein the exon 18 mutation is at least one mutation selected from the group consisting of G719A, G719S, G719C, E709K and E709A.

5 [Claim 19]

The use according to claim 17 or 18, wherein the exon 21 mutation is L861Q.

[Claim 20]

10 The use according to any one of claims 17 to 19, wherein the EGFR further has T790M mutation.

[Claim 21]

A pharmaceutical composition comprising (S)-N-(4-amino-
15 6-methyl-5-(quinolin-3-yl)-8,9-dihydropyrimido[5,4-b]indolizin-8-yl)acrylamide or a salt thereof and a pharmaceutically acceptable carrier, for treating a malignant tumor patient expressing EGFR having at least one mutation selected from the group consisting of G719X mutation of exon 18, E709X mutation of exon 18, and
20 L861X mutation of exon 21, wherein X represents an arbitrary amino-acid residue.

[Claim 22]

The pharmaceutical composition according to claim 21,
25 wherein the exon 18 mutation is at least one mutation selected from the group consisting of G719A, G719S, G719C, E709K and E709A.

[Claim 23]

The pharmaceutical composition according to claim 21 or
30 22, wherein the exon 21 mutation is L861Q.

[Claim 24]

The pharmaceutical composition according to any one of claims 21 to 23, wherein the EGFR further has T790M mutation.

35

[Claim 25]

A method for predicting therapeutic effects of chemotherapy using an antitumor agent comprising, as an active ingredient, (S)-N-(4-amino-6-methyl-5-(quinolin-3-yl)-8,9-dihydropyrimido[5,4-b]indolizin-8-yl)acrylamide or a salt thereof in a malignant tumor patient, the method comprising steps (1) and (2) below:

(1) a step of detecting the presence or absence of mutation of EGFR gene contained in a biological sample obtained from the patient; and

(2) a step of predicting that the chemotherapy is highly likely to exhibit sufficient therapeutic effects with respect to the patient when the results of the detection in step (1) found that the EGFR gene has at least one mutation selected from the group consisting of G719X mutation of exon 18, E709X mutation of exon 18, and L861X mutation of exon 21, wherein X represents an arbitrary amino-acid residue.

[Claim 26]

The method according to claim 25, wherein the exon 18 mutation is at least one mutation selected from the group consisting of G719A, G719S, G719C, E709K and E709A.

[Claim 27]

The method according to claim 25 or 26, wherein the exon 21 mutation is L861Q.

[Claim 28]

The method according to any one of claims 25 to 27, wherein the EGFR further has T790M mutation.

[Claim 29]

A method for treating a malignant tumor patient, comprising steps (1) to (3) below:

(1) a step of detecting the presence or absence of

mutation of EGFR gene contained in a biological sample obtained from the patient;

(2) a step of predicting that chemotherapy using an antitumor agent comprising, as an active ingredient, (S)-N-(4-amino-6-methyl-5-(quinolin-3-yl)-8,9-dihydropyrimido[5,4-b]indolizin-8-yl)acrylamide or a salt thereof is highly likely to exhibit sufficient therapeutic effects with respect to the patient when the results of the detection in step (1) found that the EGFR gene has at least one mutation selected from the group consisting of G719X mutation of exon 18, E709X mutation of exon 18, and L861X mutation of exon 21, wherein X represents an arbitrary amino-acid residue; and

(3) a step of administering (S)-N-(4-amino-6-methyl-5-(quinolin-3-yl)-8,9-dihydropyrimido[5,4-b]indolizin-8-yl)acrylamide or a salt thereof to a malignant tumor patient who was predicted highly likely to sufficiently respond to the chemotherapy using an antitumor agent comprising, as an active ingredient, (S)-N-(4-amino-6-methyl-5-(quinolin-3-yl)-8,9-dihydropyrimido[5,4-b]indolizin-8-yl)acrylamide or a salt thereof, in step (2).

[Claim 30]

The method according to claim 29, wherein the exon 18 mutation is at least one mutation selected from the group consisting of G719A, G719S, G719C, E709K and E709A.

[Claim 31]

The method according to claim 29 or 30, wherein the exon 21 mutation is L861Q.

[Claim 32]

The method according to any one of claims 29 to 31, wherein the EGFR further has T790M mutation.