

Arylamine-Substituted Quinazolinone Compounds

This invention relates to quinazolinone compounds, more particularly, to 2-arylamine substituted quinazolinone compounds and salts thereof which are useful as alpha-1-adrenergic receptor antagonists. The invention further relates to pharmaceutical compositions containing said compounds, to methods for their use as therapeutic agents, and to processes for making said compounds.

Alpha-1-adrenergic receptors are G-protein coupled transmembrane receptors that mediate various actions of the sympathetic nervous system through the binding of the catecholamines, epinephrine, and norepinephrine (NE). Currently, several subtypes of the alpha-1 adrenergic receptors are known to exist for which the genes have been cloned: alpha-1A (previously known as alpha-1C), alpha-1B and alpha-1D.

Alpha-1 adrenoceptor antagonists have been shown in numerous clinical studies to be effective in relieving the symptoms associated with benign prostatic hypertrophy, also known as benign prostatic hyperplasia (BPH), an illness typically affecting men over fifty. The symptoms of the condition include, but are not limited to, increased difficulty in urination and sexual dysfunction. Drugs such as prazosin, indoramin, doxazosin and tamsulosin are in common clinical use for BPH, and are effective in reducing both "obstructive" symptoms (*e.g.* weak stream) and "irritative" symptoms (*e.g.* nocturia, urinary urge and frequency). However, these compounds are all non-subtype-selective and have the potential to cause significant side-effects, particularly cardiovascular effects such as postural hypotension, dizziness, and syncope, and CNS effects such as aesthenia (tiredness). These effects can limit dosing and the clinical efficacy in reducing symptoms associated with BPH.

Pharmacological studies resulting in the subdivision of alpha-1 adrenoceptors into alpha-1A, alpha-1B, and alpha-1D adrenoceptors have led to the suggestion that development of subtype-selective antagonists may allow for an improved symptomatic treatment of BPH with a lower incidence of dose-limiting side-effects. Recently, much interest has been focused on the role of the alpha-1A adrenoceptor subtype in BPH, as studies have shown that this subtype predominates in the urethra and prostate of man and appears to


be the receptor mediating NE-induced smooth muscle contraction in these tissues. See, e.g., Price et al., J. Urol. 150:546-551 (1993); Faure et al., Life Sci. 54:1595-1605 (1994); Taniguchi et al., Naunyn Schmiedeberg's Arch. Pharmacol. 355:412-416 (1997); Forray et al., Mol. Pharmacol. 45:703-708 (1994); Hatano et al., Br. J. Pharmacol. 113:723-728 (1994); and Marshall et al., Br J. Pharmacol. 115:781-786 (1995). Smooth muscle tone is believed to contribute substantially to the total urinary outflow obstruction observed in patients with BPH [Furuya et al., J. Urol. 128:836-839 (1982)]. Increased prostate mass is also a contributing factor. These observations have fuelled the hypothesis that an alpha-1A subtype-selective antagonist may, via a selective and significant decrease in outlet resistance, lead to improved pharmacotherapy for BPH.

However, in BPH, it is often the irritative symptoms which prompt the patient to seek treatment, and these irritative symptoms may be present in patients with no demonstrable obstruction (*i.e.* normal urine flow rates). Thus, it would be advantageous to provide a therapy for treating patients exhibiting obstructive symptoms and/or irritative symptoms. It is believed that reduction of obstructive and irritative symptoms in patients with BPH may be achieved via a combination of alpha-1A and alpha-1B subtype selectivity in a drug molecule. The lack of alpha-1D adrenoceptor antagonism is expected to lead to reduced or fewer side effects than those associated with the use of non-subtype-selective agents.

The instant invention provides arylamine-substituted quinazolinone compounds that are useful as alpha 1a/b adrenergic receptor antagonists. Certain arylamine quinazoline compounds useful for other purposes are disclosed in Hess et al., Anti-hypertensive 2-Amino-4(3H)-quinazolinones, Medical Research Laboratories, Pfizer & Co. (Jan. 1968); Klopman et al., Molecular Pharmacology 34, No. 6 (December 1988); DeRuiter et al., J. Med. Chem. 29:627-29 (1986).

All publications, patents, and patent applications cited herein, whether *supra* or *infra*, are each hereby incorporated by reference in its entirety.

The present invention is directed to compounds having the Formula (I)



wherein

Y is optionally-substituted C₁₋₄alkylene, C₂₋₄alkenylene, heterocyclylene, or heterocyclC₁₋₄alkylene;

Z is -C(=O)- or -S(=O)₂-;

5 R and R' are alkyl;

R⁵ is selected from hydrogen, halogen, cyano, hydroxy, -R⁶, and -OR⁶;

R⁶ is selected from alkyl, aryl, heteroaryl, cycloalkyl, and heterocycl;

R¹⁰ is selected from hydrogen, alkyl, hydroxyalkyl, aminoalkyl and aralkyl;

Ar is optionally-substituted aryl or heteroaryl,

10 provided that if R⁵ is hydrogen, then Y-Ar considered together are not (i) unsubstituted benzyl, (ii) benzyl having a para substituent selected from hydroxy and CO₂H, or (iii) benzyl having a hydroxy meta substituent;
or an isomer or pharmaceutically-acceptable salt, hydrate, or prodrug thereof.

In one embodiment the present invention provides a compound according to Formula

15 (I)

wherein

Y is optionally-substituted C₁₋₄alkylene, C₂₋₄alkenylene, heterocyclylene, or heterocyclC₁₋₄alkylene;

Z is -C(=O)- or -S(=O)₂-;

20 R and R' are lower alkyl;

R⁵ is selected from hydrogen, halogen, cyano, hydroxy, -R⁶, and -OR⁶;

R⁶ is selected from alkyl, substituted alkyl, aryl, substituted aryl, heteroaryl, substituted heteroaryl, cycloalkyl, substituted cycloalkyl, heterocyclo, and substituted heterocyclo;

25 R¹⁰ is selected from hydrogen, alkyl, substituted alkyl, and optionally-substituted heterocyclo or cycloalkyl;

Ar is optionally-substituted aryl or heteroaryl,

provided that if R⁵ is hydrogen, then Y-Ar considered together are not (i) unsubstituted benzyl, (ii) benzyl having a para substituent selected from hydroxy and CO₂H, or (iii)

30 benzyl having a hydroxy meta substituent;

or an isomer or pharmaceutically-acceptable salt, hydrate, or prodrug thereof.

The compounds of Formula (I) above are surprisingly advantageous in selectively antagonizing the alpha-1A and alpha-1B subtype receptors with selectively lesser activity in antagonizing the alpha-1D adrenergic receptor, particularly compounds where R⁵ is

35 other than hydrogen. Accordingly, said compounds of Formula (I) are surprisingly

advantageous in methods for treating diseases responsive to alpha-1A and alpha-1B receptor antagonism with reduced side effects. The compounds having the Formula (I) are useful as alpha 1A/B adrenergic receptor antagonists.

Another aspect of this invention relates to the methods of treating a subject having a disease state that is alleviated by treatment with an alpha 1A/B adrenergic receptor antagonist, which comprises administering to such a subject in need of treatment therefore, a therapeutically effective amount of at least one compound of Formula I.

Still another aspect of the invention relates to a process for making the claimed compounds. In particular, the inventors herein have discovered a process for making intermediate diones of the formula (II)



wherein R, R' and R⁵ are as described above, from nitrobenzoic acid compounds having the formula (8)



via reduction of the nitro group to an amino group in a water-based solvent, by addition of base, heterogenous catalyst, and exposure to a hydrogen atmosphere. This process is amendable to large-scale synthesis and is high-yielding and environmentally friendly.

Unless otherwise stated, the following terms used in this application, including the specification and claims, have the definitions given below. As used in the specification and the appended claims, the singular forms "a", "an," and "the" include plural referents unless the context clearly dictates otherwise.

In general, the nomenclature used in this Application is based on AutoNom[®], a Beilstein Institute computerized system for the generation of IUPAC systematic nomenclature. Chemical structures shown herein were prepared using ISIS[®] version 2.2. Any open valency shown on a carbon, nitrogen or oxygen in the structures herein indicates the presence of a hydrogen.

As used herein, the term "alkyl" means a linear or branched, saturated monovalent hydrocarbon moiety of one to eight carbon atoms (preferably one to six carbon atoms), e.g., methyl, ethyl, *n*-propyl, 2-propyl, *tert*-butyl, pentyl, and the like. "Lower alkyl" means an alkyl of one to four carbon atoms. When a subscript is used herein following a carbon atom, the subscript refers to the number of carbon atoms that the named group may contain. Thus, e.g., C₁₋₄alkyl means an alkyl of one to four carbon atoms (*i.e.*, lower alkyl) including methyl, ethyl, propyl, *iso*-propyl, butyl, and *tert*-butyl; hydroxy(C₁₋₄)alkyl means an alkyl of one to four carbon atoms substituted with a hydroxy group; C₁₋₄alkoxyalkyl means an alkyl group substituted with an alkoxy group wherein the alkoxy group has one to four carbon atoms; C₁₋₄alkoxy(C₁₋₄)alkyl means an alkyl group of one to four carbon atoms substituted with an alkoxy group wherein the alkoxy group has one to four carbon atoms; and so forth.

"Alkylene" means a linear or branched, saturated bivalent hydrocarbon moiety of one to eight (preferably one to six) carbon atoms, e.g., methylene, ethylene, propylene, and the like.

When the term "alkyl" is used as a suffix following another term, as in "phenylalkyl," or "hydroxyalkyl," this is intended to refer to an alkyl group, as defined above, being substituted with one or two (preferably one) substituent(s) selected from the other, specifically-named group, also as defined herein. For example, "phenylalkyl" includes benzyl, phenylethyl, 2-phenylbutyl, and so forth. "Hydroxyalkyl" includes 2-hydroxyethyl, 1-(hydroxymethyl)-2-methylpropyl, 3,4-dihydroxybutyl, and so forth. "Alkoxyalkyl" refers to an alkyl group substituted with one to two of OR', wherein R' is alkoxy as defined below.

The term "substituted alkyl" refers to an alkyl group as defined above having one, two, three, or four substituents (preferably one to two substituents), independently selected from the group consisting of halo, haloalkoxy, trifluoromethyl, cyano, nitro, -OR^a, -SR^a, -S(O)R^c, -S(O)₂R^c, -C(=O)R^a, -C(=O)NR^aR^b, -C(O)₂R^a, -C(O)₂NR^aR^b, -S(O)₂NR^aR^b, -NR^aR^b, -NR^a(C=O)R^b, aryl, heteroaryl, cycloalkyl, and/or heterocyclo, wherein R^a and R^b are independently selected from hydrogen, C₁₋₆alkyl, aryl, heteroaryl, cycloalkyl, and heterocyclo, and R^c is selected from alkyl, aryl, heteroaryl, cycloalkyl, and heterocyclo, and each of R^a, R^b, and R^c in turn is optionally substituted with one, two, or three of alkyl, halo, haloalkyl, OR^e, haloalkoxy, cyano, -NR^eR^f, -SO₂(alkyl), -CO₂R^e, -C(=O)R^e, and/or -NR^eC(=O)R^f, and/or a C₁₋₆alkyl substituted with one to two of halo, OR^e, haloalkoxy, cyano, -NR^eR^f, -SO₂(alkyl), -CO₂R^e, -C(=O)R^e, and/or -NR^eC(=O)R^f,

wherein R^e and R^f are independently selected from hydrogen, alkyl, aminoalkyl, hydroxyalkyl, alkoxyalkyl, cyanoalkyl, haloalkyl, and alkylaminoalkyl. Additionally, when a substituted alkyl group is substituted with a cyclic group such as aryl, heteroaryl, cycloalkyl, or heterocyclo, it is understood said cyclic group in turn may be substituted
5 with one, two or three groups selected from alkyl, halo, haloalkyl, OR^e , haloalkoxy, cyano, $-NR^eR^f$, $-SO_2(alkyl)$, $-CO_2R^e$, $-C(=O)R^e$, and/or $-NR^eC(=O)R^f$, and/or a $C_{1-6}alkyl$ substituted with one to two of halo, OR^e , haloalkoxy, cyano, $-NR^eR^f$, $-SO_2(alkyl)$, $-CO_2R^e$, $-C(=O)R^e$, and/or $-NR^eC(=O)R^f$, wherein R^e and R^f are as defined immediately above.

The term "substituted alkylene" means an alkylene group as defined above wherein one,
10 two or three carbon atoms of the alkylene straight or branched chain is substituted with a group selected from those recited above for substituted alkyl groups.

The term "substituted lower alkyl" means an alkyl of one to four carbon atoms having one, two, or three substituents selected from those recited above for substituted alkyl.

The term "alkenyl" means a linear or branched, unsaturated monovalent hydrocarbon
15 moiety of two to eight carbon atoms (preferably two to six carbon atoms), having at least one double bond, *e.g.*, ethenyl, propenyl, butenyl, and the like. "Lower alkenyl" means an alkenyl of two to four carbon atoms.

"Alkenylene" means a linear or branched, unsaturated bivalent hydrocarbon moiety of two to eight (preferably two to six) carbon atoms, having at least one double bond, *e.g.*,
20 ethenylene, propenylene, and the like.

The term "substituted alkenyl" refers to an alkenyl group as defined above having one, two, or three substituents, as valence permits (preferably one substituent), independently selected from the group of substituents recited above for "substituted alkyl." A
"substituted alkenylene" has one, two, or three substituents, as valence permits
25 (preferably one substituent), independently selected from the group of substituents recited for "substituted alkyl."

"Alkoxy" refers to the group OR , wherein R is alkyl or substituted alkyl. A "lower alkoxy" is a group $-OR'$ wherein R' is $C_{1-4}alkyl$.

When the term "oxy" is used as a suffix following another specifically-named group, as in
30 "aryloxy", "heteroaryloxy", or "arylalkyloxy", this means that an oxygen atom is present as a linker to the other, specifically-named group. Thus, *e.g.*, "aryloxy" refers to the group $-OR$, wherein R is aryl; "heteroaryloxy" refers to the group $-OR'$, wherein R' is

heteroaryl; and "arylalkyloxy" refers to the group $-OR$, wherein R is arylalkyl such as benzyl. Similarly, a "substituted aryloxy" means the group $-OR$, wherein R is substituted aryl, and a "substituted heteroaryloxy" means the group $-OR$, wherein R is substituted heteroaryl.

- 5 "Amino" refers to the group NH_2 . Thus, an aminoalkyl refers to an alkyl group having an amino substituent, e.g., $-CH_2-NH_2$, $-CH_2-CH_2-NH_2$, $-CH_2-CH(NH_2)-CH_3$, and so forth.

"Alkylamino" as used herein refers to monoalkylamino groups having the formula $-NHR$, as well as dialkylamino groups having the formula $-NRR'$, wherein each R and R' are selected from alkyl and substituted alkyl groups as defined above.

- 10 An "alkylaminoalkyl" refers to an alkyl group substituted by one to two of $-NHR$ and/or $-NRR'$, wherein each R and R' is as defined above. A "lower alkylamino" refers to a group $-NHR'$ or $-NR'R'$, wherein each R' is C_{1-4} alkyl.

"Aminoalkoxy" means a group $-O-R-NHR'$ or $-O-R-NR'R''$ wherein R is alkylene as defined herein and R' and R'' each independently are alkyl as defined herein.

- 15 As used herein, the term "alkoxyalkylamino" refers to the group $-NR^fR^g$, wherein R^f is hydrogen, alkyl, or alkoxyalkyl, and R^g is an alkoxyalkyl (i.e., an alkyl group substituted with an alkoxy).

"Alkoxyalkylaminoalkyl" refers to an alkyl group as defined above that is substituted with an alkoxyalkylamino group as defined above.

- 20 A "hydroxyalkylamino" refers to the group $-NR^hR^i$, wherein R^h is hydrogen, alkyl, or hydroxyalkyl, and R^i is hydroxyalkyl (i.e., an alkyl group substituted with hydroxy).

"Hydroxyalkylaminoalkyl" refers to an alkyl group as defined above that is substituted with a hydroxyalkylamino group as defined above.


- 25 "Aminoalkoxy" means a group $-O-R-NHR'$ or $-O-R-NR'R''$ wherein R is alkylene as defined above and R' and R'' each independently are alkyl as defined above.

The term "alkylsulfonyl" refers to the group $-SO_2R$, wherein R is alkyl or substituted alkyl, and "alkylsulfinyl" refers to the group $-S(=O)R$, wherein R is alkyl or substituted alkyl. Thus, accordingly, methylsulfonyl refers to $-SO_2CH_3$, and methylsulfinyl refers to the group $-S(=O)CH_3$.

"Alkylsulfonylalkyl" means an alkyl group as defined above substituted with an alkylsulfonyl group as defined herein.

"Alkylsulfonylaminoalkyl" means a group $-R-NR'-SO_2-R''$ wherein R is alkylene as defined above and R' and R'' each independently are alkyl as defined herein.

- 5 "Acetamidinyl" means a group of the formula



wherein each R is independently hydrogen or alkyl as defined herein.

"Alkoxyalkyl" means a group $-R-OR'$ wherein R is alkylene as defined above and R' is alkyl as defined herein.

- 10 "Alkoxyalkoxy" means a group $-O-R-OR'$ wherein R is alkylene as defined above and R' is alkyl as defined herein.

The term "carboxy" refers to the group CO_2H . Thus, a carboxyalkyl is an alkyl group as defined above having at least one substituent that is $-CO_2H$.

- 15 The term alkoxy carbonyl refers to the group $-C(=O)R'$, wherein R' is alkoxy as defined above, i.e., alkoxy carbonyl is CO_2R , wherein R' is alkyl or substituted alkyl, as defined above. A "lower alkoxy carbonyl" refers to the group CO_2R' , wherein R' is lower alkyl. Thus, an alkylalkoxy carbonyl is an alkyl group as defined above having at least one substituent that is $-CO_2R'$, wherein R' is alkyl or substituted alkyl, as defined above.

- 20 "Alkoxyalkylaminocarbonyl" means a group $-(C=O)-NR-R'-OR''$ wherein R is hydrogen or alkyl as defined herein, R' is alkylene as defined herein, and R'' is alkyl as defined herein.

"Hydroxyalkylaminocarbonyl" means a group $-(C=O)-NR-R'-OH$ wherein R is hydrogen or alkyl as defined herein and R' is alkylene as defined herein.

- 25 "Aminoalkylaminocarbonyl" means a group $-(C=O)-NR-R'-NR''R'''$ wherein R is hydrogen or alkyl as defined herein, R' is alkylene as defined herein, and each R'' and R''' independently is hydrogen or alkyl as defined herein.

"Heterocyclalkylaminocarbonyl" means a group $-(C=O)-NR-R'-R''$ wherein R is hydrogen or alkyl as defined herein, R' is alkylene as defined herein, and R'' is heterocycl as defined herein.


The term alkylamidyl or alkylamide refers to the group $-NH(C=O)R$ or $-NR'(C=O)R$,
 5 wherein R is alkyl or substituted alkyl, and R' is lower alkyl. A lower alkylamidyl is a group $-NH(C=O)R'$ or $-NR'(C=O)R'$, where R' is lower alkyl.

The term "aryl" refers to a monovalent, monocyclic or bicyclic moiety in which at least one of the rings is an aromatic, carbocyclic moiety. Thus, the term "aryl" includes phenyl, 1-naphthyl, and 2-naphthyl. The term "aryl" also includes phenyl rings having
 10 fused thereto a second non-aromatic carbocyclic ring, or second fused heteroaryl or heterocyclic ring (thus, the term aryl includes groups such as benzothienyl, benzopyrazolyl, benzopiperadiny, benzocyclohexyl, and the like), with the understanding, however, that in the case of bicyclic aryl groups, the point of attachment will be to the phenyl ring.

15 A "substituted aryl" is an aryl group as defined above having one to four (preferably one to two) substituents independently selected from the group consisting of halo, haloalkyl, haloalkoxy, cyano, hydroxy, alkoxy, alkyl, substituted alkyl, $-SO_2R^f$, $-NR^pSO_2R^f$, $-NR^pC(=O)R^q$, $-NR^pR^q$, $-C(=O)NR^pR^q$, $-SO_2NR^pR^q$, $-NR^pC(=NR^s)R^q$, $-N=C(R^t)R^u$, $-SO_2N=C(R^t)R^u$, $-C(=O)R^p$, $-CO_2R^p$, and $-OR^p$, heterocyclo, heteroaryl, phenyl and
 20 cycloalkyl, wherein each R^f , R^p , and R^q is selected from hydrogen, alkyl, substituted alkyl, cycloalkyl, heterocyclo, aryl, and heteroaryl, except R^f is not hydrogen, R^s is selected from hydrogen, alkyl, substituted alkyl, cycloalkyl, and heterocyclo, or alternatively, R^p and R^q when attached to the same nitrogen atom may be taken together to form a heterocycl or heteroaryl; R^t is selected from hydrogen, alkyl, substituted alkyl, cycloalkyl, and hetero-
 25 cycl, and R^u is amino or alkylamino, or alternatively, R^t and R^u are taken together to form a cycloalkyl or heterocycl ring. It should be understood that when an aryl group is substituted by a further ring, and/or when any of R^p , R^q , R^f , R^s , R^t and R^u are selected from and/or form a ring, said cyclic groups in turn are optionally substituted with one to three groups selected from alkyl, substituted alkyl, halogen, haloalkoxy, trifluoromethyl,
 30 cyano, nitro, $-OR^a$, $-SR^a$, $-S(O)R^c$, $-S(O)_2R^c$, $-C(=O)R^a$, $-C(=O)NR^aR^b$, $-C(O)_2R^a$, $-C(O)_2NR^aR^b$, $-S(O)_2NR^aR^b$, $-NR^aR^b$, and/or $-NR^a(C=O)R^b$, wherein R^a , R^b and R^c are as defined above in the definition of substituted alkyl groups.

The term "carbocyclic" means a cyclic moiety in which all ring atoms are carbon atoms, including saturated, partially unsaturated, and unsaturated rings.

"Carbamylalkyl" means a group of the formula




- 5 wherein each R is independently hydrogen or alkyl.

The term "cycloalkyl" as used herein refers to saturated or partially unsaturated, monovalent, monocyclic carbocyclic moieties of three to seven ring carbon atoms and further includes such rings having a carbon-carbon bridge of one, two, or three bridgehead carbon atoms, and/or having a second ring fused thereto, with the
 10 understanding that the point of attachment will be to the non-aromatic carbocyclic ring moiety. Thus, the term "cycloalkyl" includes such rings as cyclopropyl, cyclopentyl, cyclopentenyl, cyclohexyl, cyclohexenyl, and the like. Additionally, one or two carbon atoms of a cycloalkyl group may optionally contain a carbonyl oxygen group, *e.g.*, one or two atoms in the ring may be a moiety of the formula $-C(=O)-$.

- 15 A "substituted cycloalkyl" is a cycloalkyl group as defined above having one to four (preferably one to two) substituents independently selected from the group consisting of substituents recited above for substituted aryl.

"Formamidinyl" means a group of the formula



- 20 wherein R is hydrogen or alkyl as defined herein. The term "N,N-dimethyl formamidine" or "N,N-dimethyl formamidinyl" refers to the above group wherein R is methyl.

The term "halo," "halide" or "halogen," when referring to a substituent means fluoro, chloro, bromo, or iodo (preferably fluoro or chloro).

- 25 The term "haloalkyl" means alkyl substituted with one or more same or different halo atoms, *e.g.*, $-CH_2Cl$, $-CF_3$, $-CH_2CF_3$, $-CH_2CCl_3$, and the like, and further includes those

alkyl groups such as perfluoroalkyl in which all alkyl hydrogen atoms are replaced by fluorine atoms.

The term "haloalkoxy" means a haloalkyl group as defined above linked through an oxygen atom, *e.g.*, it includes $-O-CH_2Cl$, $-O-CF_3$, $-O-CH_2CF_3$, $-O-CH_2CCl_3$, and the like.

"Haloalkylamino" means a group $-NH-R''$ or $-NR'-R''$ wherein R' is alkyl or haloalkyl as defined herein and R'' is haloalkyl as defined herein.

"Haloalkylaminoalkyl" means a group $-R-NH-R''$ or $-NR'-R''$ wherein R is alkylene as defined herein, R' is alkyl or haloalkyl as defined herein and R'' is haloalkyl as defined herein.


"Hydroxyalkyl" means a group $-R-OH$ wherein R is alkylene as defined herein.

"Hydroxyalkoxy" means a group $-O-R-OH$ wherein R is alkylene as defined herein.

"Heterocyclo," "heterocyclyl," or "heterocyclic" refers to a saturated or partially-unsaturated non-aromatic monocyclic or bicyclic moiety in which one or two ring atoms are heteroatoms selected from N, O, or $S(O)_x$ (where x is an integer from 0 to 2), the remaining ring atoms being carbon atoms, and additionally, one or two carbon atoms may optionally contain a carbonyl oxygen group, *e.g.*, one or two atoms in the ring may be a moiety of the formula $-C(=O)-$. Thus, the term heterocyclo includes rings such as tetrahydropyranyl, tetrahydrofuryl, piperidinyl, piperazinyl, morpholinyl, pyrrolidinyl, pyrrolinyl, imidazolidinyl, imidazolynyl, and the like, as well as such rings having a carbonyl oxygen atom in the ring. In the case of a bicyclic heterocyclo, one of the two rings may be a carbocyclic ring with the understanding that in such cases the point of attachment will be to the heterocyclic ring.

"Heterocyclylalkylamino" means a group $-NR'-R''$ wherein R is hydrogen or alkyl as defined herein, R' is alkylene as defined herein, and R'' is heterocyclyl as defined herein.

When reference is made to an imidazolynyl group, this is intended to refer to both



Imidazolynyl groups as defined herein may be optionally substituted with alkyl.

"Imidazolinyllalkyl" means a group -R-R' wherein R is alkylene as defined herein and R' is imidazolinyll as defined herein.

A "substituted heterocyclo" or "substituted heterocycle" refers to a heterocyclo group as defined above having one to four substituents (preferably one to two substituents)
 5 selected from the group of substituents recited above for substituted aryl.


A "heterocyclylene" means a bivalent heterocyclyl group as defined above, i.e., a heterocyclyl attached to two other groups (e.g., -NR¹⁰ and Ar in compounds of Formula I).

A "heterocyclylC₁₋₄alkylene" means a group X-R, wherein X is a heterocyclyl as defined
 10 above and R is a C₁₋₄alkylene as defined above. An optionally-substituted "heterocyclyl-C₁₋₄alkylene" means a group X-R, wherein X is an optionally-substituted heterocyclyl as defined herein and R is an optionally substituted C₁₋₄alkylene as defined herein.

"Heteroaryl" means a monovalent, monocyclic aromatic moiety of 5 to 6 ring atoms containing one, two, three, or four ring heteroatoms, each independently selected from N, O,
 15 or S, the remaining ring atoms being carbon, and it also includes such rings having a second ring fused thereto of five to six ring atoms, wherein the second fused ring may be aromatic or nonaromatic and may be carbocyclic, heterocyclic, or a heteroaryl ring, with the understanding, however, that in such cases the point of attachment will be to an aromatic ring containing at least one heteroatom. Thus, the term heteroaryl includes, but is
 20 not limited to, pyridyl, furyl, thiophenyl, thiazolyl, isothiazolyl, triazolyl, imidazolyl, isoxazolyl, pyrrolyl, pyrazolyl, pyrimidinyl, benzofuryl, isobenzofuryl, benzothiazolyl, benzoisothiazolyl, benzotriazolyl, indolyl, isoindolyl, benzoxazolyl, quinolyl, isoquinolyl, benzimidazolyl, benzisoxazolyl, benzothiophenyl, dibenzofuran, and benzodiazepin-2-one-5-yl, and derivatives thereof.

25 A "substituted heteroaryl" is a heteroaryl ring as defined above having one to four (preferably one or two) substituents selected from the group of substituents recited above for substituted aryl.


"Optionally substituted pyrrolidinyll" means a group



30 wherein E is -CH₂- or -(C=O)- and R is alkyl, alkoxy or hydroxy.

"Optionally substituted pyrrolidinylalkyl" means a group -R-R' wherein R is alkylene as defined herein and R' is imidazolinyl as defined herein.

"Ureidylalkyl" means a group



5 wherein each R is independently hydrogen or methyl.

"Leaving group" has the meaning conventionally associated with it in synthetic organic chemistry, *i.e.*, an atom or a group capable of being displaced by a nucleophile and includes halo (such as chloro, bromo, and iodo), alkanesulfonyloxy, arenesulfonyloxy, alkylcarbonyloxy (*e.g.*, acetoxy), arylcarbonyloxy, mesyloxy, tosyloxy,
 10 trifluoromethanesulfonyloxy, aryloxy (*e.g.*, 2,4-dinitrophenoxy), methoxy, N,O-dimethylhydroxylamino, and the like.


"Optional" or "optionally" means that the subsequently described event may but need not occur, and it includes instances where the event occurs and instances in which it does not. For example, "optionally-substituted cycloalkyl" refers to both cycloalkyl groups
 15 and substituted cycloalkyl groups, as defined above. When the term "optionally-substituted" precedes a number of different types of rings in one line or string, *e.g.*, as in "optionally-substituted cycloalkyl or heterocyclo", or "optionally-substituted carbocyclic or heterocyclic ring," or "optionally-substituted aryl, heteroaryl, cycloalkyl, or heterocyclo," it is intended that the term "optionally-substituted" modifies each of the
 20 rings identified in the line or string.

When the term "optionally-substituted" is used with respect to a particularly-named cyclic group, such as "optionally-substituted imidazolyl," or "optionally-substituted imidazolidinyl," it should be understood that the optional substituents for such particularly-named rings may be selected from the group of substituents recited above
 25 with respect to which the genus of which the particularly-named group is a member. Thus, *e.g.*, an "optionally-substituted imidazolyl" may be an unsubstituted imidazolyl or an imidazolyl group having one, two, or three substituents selected from those recited above for substituted heteroaryl groups. An optionally-substituted phenyl or benzyl ring will include an unsubstituted phenyl or benzyl group, and a phenyl or benzyl group
 30 having substituents selected from those recited above for substituted aryl groups.

It should be understood that when reference is made to a specific heterocyclo or cycloalkyl group, such as cyclopentyl, pyrrolidinyl, pyrrolinyl, and imidazoliny, such reference is intended to include such rings wherein optionally one to two carbon atoms of the named ring contain a carbonyl oxygen group, *e.g.*, one or two atoms in the ring
5 may be a moiety of the formula $-C(=O)-$, as set forth above in the definition of cycloalkyl and heterocyclo.


An optionally-substituted benzyl group means a benzyl group wherein the phenyl portion of the group is unsubstituted or substituted as defined above in the definition for substituted aryl.

- 10 A benzyl group having a para substituent selected from hydroxy and CO_2H , means a group having the formula




wherein R is hydroxy or CO_2H , and benzyl having a hydroxy meta substituent means a group having the formula

15



When reference is made herein to substituents on the quinazolinone core, *e.g.*, the “five position substituent,” the numbering of the ring atoms is intended to be as follows



- “Pharmaceutically acceptable excipient” means an excipient that is useful in preparing a
20 pharmaceutical composition that is generally safe, non-toxic and neither biologically nor otherwise undesirable. The term includes excipients that are acceptable for veterinary use as well as human pharmaceutical use. A “pharmaceutically acceptable excipient” as used in the specification and claims includes both one and more than one such excipient.

- “Pharmaceutically acceptable salt” of a compound means a salt that is generally safe,
25 non-toxic and neither biologically nor otherwise undesirable, and which possesses the

desired pharmacological activity of the parent compound. Such salts include: (1) acid addition salts, formed with inorganic acids such as hydrochloric acid, hydrobromic acid, sulfuric acid, nitric acid, phosphoric acid, and the like; or formed with organic acids such as acetic acid, trifluoroacetic acid, propionic acid, hexanoic acid, cyclopentanepropionic acid, glycolic acid, pyruvic acid, lactic acid, malonic acid, succinic acid, malic acid, maleic acid, fumaric acid, tartaric acid, citric acid, benzoic acid, 3-(4-hydroxybenzoyl)benzoic acid, cinnamic acid, mandelic acid, methanesulfonic acid, ethanesulfonic acid, 1,2-ethane-disulfonic acid, 2-hydroxyethanesulfonic acid, benzenesulfonic acid, 4-chlorobenzenesulfonic acid, 2-naphthalenesulfonic acid, 4-toluenesulfonic acid, camphorsulfonic acid, 4-methylbicyclo[2.2.2]-oct-2-ene-1-carboxylic acid, glucoheptonic acid, 4,4'-methylenebis-(3-hydroxy-2-ene-1-carboxylic acid), 3-phenylpropionic acid, trimethylacetic acid, tertiary butylacetic acid, lauryl sulfuric acid, gluconic acid, glutamic acid, hydroxynaphthoic acid, salicylic acid, stearic acid, muconic acid, and the like; or (2) salts formed when an acidic proton present in the parent compound either is replaced by a metal ion, e.g., an alkali metal ion, an alkaline earth ion, or an aluminum ion; or coordinates with an organic base such as ethanolamine, diethanolamine, triethanolamine, N-methylglucamine, and the like.

The terms "pro-drug" and "prodrug" are used interchangeably herein and refer to any compound which releases an active parent drug according to Formula I *in vivo* when such prodrug is administered to a mammalian subject. Prodrugs of a compound of Formula I are prepared by modifying one or more functional group(s) present in the compound of Formula I in such a way that the modification(s) may be cleaved *in vivo* to release the parent compound. Prodrugs include compounds of Formula I wherein a hydroxy, amino, or sulfhydryl group in a compound of Formula I is bonded to any group that may be cleaved *in vivo* to regenerate the free hydroxyl, amino, or sulfhydryl group, respectively. Examples of prodrugs include, but are not limited to, esters (e.g. acetate, dialkylaminoacetates, formates, phosphates, sulfates and benzoate derivatives) and carbamates of hydroxy functional groups (e.g. *N,N*-dimethylcarbonyl), esters of carboxyl functional groups (e.g. ethyl esters, morpholinoethanol esters), *N*-acyl derivatives (e.g. *N*-acetyl), *N*-Mannich bases, Schiff bases and enamines of amino functional groups, oximes, acetals, ketals, and enol esters of ketones and aldehyde functional groups in compounds of Formula I, and the like.

The prodrug can be metabolized before absorption, during absorption, after absorption, or at a specific site. Although metabolism occurs for many compounds primarily in the liver, almost all other tissues and organs, especially the lung, are able to carry out varying

degrees of metabolism. Prodrug forms of compounds may be utilized, e.g., to improve bioavailability, improve subject acceptability such as by masking or reducing unpleasant characteristics such as bitter taste or gastrointestinal irritability, alter solubility such as for intravenous use, provide for prolonged or sustained release or delivery, improve ease of formulation, or provide site-specific delivery of the compound. Reference to a compound herein includes prodrug forms of a compound. Prodrugs are described, e.g., in *The Organic Chemistry of Drug Design and Drug Action*, by Silverman, Academic Press, San Diego (1992), pp.352-401; *Design of Prodrugs*, edited by Bundgaard, Elsevier Science, Amsterdam (1985); *Design of Biopharmaceutical Properties through Prodrugs and Analogs*, Ed. by Roche, American Pharmaceutical Association, Washington (1977); and *Drug Delivery Systems*, ed. by Juliano, Oxford Univ. Press, Oxford (1980).


"Solvate" means solvent addition form that contains either stoichiometric or non-stoichiometric amounts of solvent. Some compounds have a tendency to trap a fixed molar ratio of solvent molecules in the crystalline solid state, thus forming a solvate. If the solvent is water the solvate formed is a hydrate, when the solvent is alcohol, the solvate formed is an alcoholate.

"Protecting group" refers to an atom or group of atoms that is attached to a reactive group in a molecule and masks, reduces, or prevents the reactivity of the group to which it is attached. Examples of protecting groups can be found in *Green and Wuts, Protective Groups in Organic Chemistry* (Wiley, 2nd ed. 1991), and *Harrison and Harrison et al., Compendium of Synthetic Organic Methods*, Vols. 1-8 (John Wiley and Sons, 1971-1996). Representative amino protecting groups include formyl, acetyl, trifluoroacetyl, benzyl, benzyloxycarbonyl (CBZ), *tert*-butoxycarbonyl (Boc), trimethyl silyl (TMS), 2-trimethylsilyl-ethanesulfonyl (SES), trityl and substituted trityl groups, allyloxycarbonyl, 9-fluorenylmethyloxycarbonyl (Fmoc), nitro-veratryloxycarbonyl (NVOC), and the like. Representative hydroxy protecting groups include those where the hydroxy group is either acylated or alkylated such as with benzyl or lower alkyl and trityl ethers as well as alkyl ethers, tetrahydropyranyl ethers, trialkylsilyl ethers, and allyl ethers.

Compounds that have the same molecular formula but differ in the nature or sequence of bonding of their atoms or the arrangement of their atoms in space are termed "isomers." Isomers that differ in the arrangement of their atoms in space are termed "stereoisomers". Stereoisomers that are not mirror images of one another are termed "diastereomers," and those that are non-superimposable mirror images of each other are termed "enantiomers". When a compound has an asymmetric center, e.g., if a carbon atom is bonded to four different groups, a pair of enantiomers is possible. An enantiomer can be

characterized by the absolute configuration of its asymmetric center and is described by the (R) and (S) sequencing rules of Cahn and Prelog, or by the manner in which the molecule rotates the plane of polarized light and designated as dextrorotatory or levorotatory (*i.e.*, as (+) or (-)-isomers respectively). A chiral compound can exist as
 5 either an individual enantiomer or as a mixture thereof. A mixture containing different enantiomers is called a "racemic mixture". The compounds of this invention may possess one or more asymmetric centers; such compounds can therefore be produced as individual (R)- or (S)-stereoisomers or as mixtures thereof. Unless indicated otherwise, the description or naming of a particular compound in the specification and claims is
 10 intended to include both individual enantiomers and mixtures (racemic or otherwise) thereof. The methods for the determination of stereochemistry and the separation of stereoisomers are well-known in the art [e.g. March, Advanced Organic Chemistry, Chap. 4, 4th edition, John Wiley and Sons, New York (1992)].

"Tautomers" refers to compounds whose structures differ markedly in arrangement of
 15 atoms, but which exist in easy and rapid equilibrium. It is to be understood that compounds of Formula I may be depicted as different tautomers. For example, compounds of Formula I wherein Z is -C(O)-, may be depicted in the following tautomer forms



20 Compounds of Formula I may also contain other groups that exist in tautomeric equilibrium. For example some of the compounds contain an imidazolin-2-yl amino group which may be in equilibrium with an imidazolin-2-ylidenamino group



It should also be understood that when compounds have tautomeric forms, all tautomeric forms are intended to be within the scope of the invention, and the naming of the compounds does not exclude any tautomer form.

5 "Treating" or "treatment" of a disease includes: (1) preventing the disease, *i.e.*, causing the clinical symptoms of the disease not to develop in a mammal that may be exposed to or predisposed to the disease but does not yet experience or display symptoms of the disease; (2) inhibiting the progression of the disease, *i.e.*, arresting or reducing the development of the disease or its symptoms; and (3) relieving the disease, *i.e.*, causing regression of the disease or its symptoms.

10 "A therapeutically effective amount" means the amount of a compound that, when administered to a mammal for treating a disease, is sufficient to effect a treatment for the disease. The "therapeutically effective amount" will vary depending on such factors as the compound being administered, the type of disease being treated, the progression or severity of the disease state, and the age, weight, and general health of the mammal being
15 treated.

"Patient" means mammals and non-mammals. Mammals means any member of the mammalia class including, but not limited to, humans, non-human primates such as chimpanzees and other apes and monkey species; farm animals such as cattle, horses, sheep, goats, and swine; domestic animals such as rabbits, dogs, and cats; and laboratory
20 animals such as rats, mice, and guinea pigs. Examples of non-mammals include, but are not limited to, birds, reptiles, and the like.

"Pharmacological effect" as used herein encompasses effects produced in the patient that achieve the intended purpose of a therapy. In one preferred embodiment, a pharmacological effect means that primary indications of the patient being treated are
25 prevented, alleviated, or reduced. For example, a pharmacological effect would be one that results in the prevention, alleviation or reduction of primary indications in a treated patient. In another preferred embodiment, a pharmacological effect means that disorders or symptoms of the primary indications of the patient being treated are prevented, alleviated, or reduced.

30 "Disease state" means any disease, condition, symptom, or indication.

"Disorders of the urinary tract" or "uropathy" refer to pathologic changes in the urinary tract and symptoms thereof. Disorders of the urinary tract include overactive bladder (also known as detrusor hyperactivity), outlet obstruction, outlet insufficiency, pelvic

hypersensitivity; incontinence, benign prostatic hypertrophy or hyperplasia (BPH), prostatitis, detrusor hyperreflexia, urinary frequency, nocturia, urinary urgency, pelvic hypersensitivity, urge incontinence, urethritis, prostatodynia, cystitis, and idiopathic bladder hypersensitivity.

- 5 "Overactive bladder" or "Detrusor hyperactivity" includes, but is not limited to, changes symptomatically manifested as urgency, frequency, reduced bladder capacity, and incontinence episodes; changes urodynamically manifested as changes in bladder capacity, micturition threshold, unstable bladder contractions, and sphincteric spasticity; and symptoms usually manifested in detrusor hyperreflexia (neurogenic bladder), in
10 conditions such as outlet obstruction, outlet insufficiency, pelvic hypersensitivity, or in idiopathic conditions such as detrusor instability.

"Outlet obstruction" includes, but is not limited to, benign prostatic hypertrophy or benign prostatic hyperplasia (BPH), urethral stricture disease, tumors and the like. It is usually symptomatically manifested as obstructive (low flow rates, difficulty in initiating
15 urination, and the like), or irritative (urgency, suprapubic pain, and the like).

"Outlet insufficiency" includes, but is not limited to, urethral hypermobility, intrinsic sphincteric deficiency, or mixed incontinence. It is usually symptomatically manifested as stress incontinence.

"Pelvic Hypersensitivity" includes but is not limited to, pelvic pain, interstitial (cell)
20 cystitis, prostatic pain, prostatitis, vulvodynia, urethritis, orchialgia, and the like. It is symptomatically manifested as pain, inflammation or discomfort referred to the pelvic region, and usually includes symptoms of overactive bladder.

"Sexual dysfunction" means the inability to achieve a normal sexual response and includes such conditions in males and females. Thus, it includes male erectile dysfunction
25 (MED) and female sexual dysfunction (FSD).

"Disease states associated with the Central Nervous System (CNS)" or "CNS disease states" mean neurological and/or psychiatric changes in the CNS, e.g., brain and spinal cord, which manifest in a variety of symptoms. Examples of CNS disease states include, but are not limited to, migraine headache; cerebrovascular deficiency; psychoses
30 including paranoia, schizophrenia, attention deficiency, and autism; obsessive/compulsive disorders including anorexia and bulimia; posttraumatic stress disorders, sleep disorders, convulsive disorders including epilepsy and withdrawal from addictive substances; cognitive diseases including Parkinson's disease and dementia; and

anxiety/depression disorders such as anticipatory anxiety (e.g., prior to surgery, dental work and the like), depression, mania, seasonal affective disorder (SAD), and convulsions and anxiety caused by withdrawal from addictive substances such as opiates, benzodiazepines, nicotine, alcohol, cocaine, and other substances of abuse; and improper thermoregulation.

The compounds of this invention demonstrate selectivity for the alpha-1A/B subtype over the alpha-1D subtype. The compounds of this invention may reduce both obstructive and irritative symptoms in patients with BPH. The attenuated antagonism of alpha 1D-adrenoceptor is expected to lead to reduced or fewer side effects than those associated with the use of non-subtype-selective agents.

In one embodiment the present invention provides compounds of formula I wherein Y is $-(\text{CHR}^1)-$ or $-(\text{CHR}^2-\text{CHR}^3)-$; and R^1 , R^2 and R^3 are independently selected from hydrogen, alkyl, and hydroxy, or R^2 and R^3 may together form a bond so that Y is ethenylene.

In one embodiment the present invention provides compounds of formula I wherein Y is heterocyclylene. In another embodiment the present invention provides compounds of formula I wherein said heterocyclene is piperazin-1,4-di-yl or piperidin-1-3-di-yl.

In one embodiment the present invention provides compounds of formula I wherein Z is $-\text{C}(=\text{O})-$.

In one embodiment the present invention provides compounds of formula I wherein R and R' are both CH_3 .

In one embodiment the present invention provides compounds of formula I wherein R^5 is selected from hydrogen, halogen, cyano, hydroxy, $-\text{R}^6$, and $-\text{OR}^6$; and R^6 is alkyl. In another embodiment the present invention provides compounds of formula I wherein R^5 is selected from hydrogen, methyl, ethyl, n-propyl, isopropyl, halogen, cyano, methoxy, and ethoxy.

In one embodiment the present invention provides compounds of formula I wherein R^{10} is selected from hydrogen, alkyl, hydroxyalkyl, and benzyl.


In one embodiment the present invention provides compounds of formula I wherein Z is $-\text{C}(=\text{O})-$; R and R' are both CH_3 ; R^5 is selected from methyl, ethyl, n-propyl, isopropyl, halogen, cyano, methoxy, and ethoxy; and R^{10} is selected from hydrogen and alkyl.

In another embodiment the present invention provides compounds of formula I wherein Z is $-C(=O)-$; R and R' are both CH_3 ; R^5 is selected from methyl, ethyl, n-propyl, isopropyl, halogen, cyano, methoxy, and ethoxy; R^{10} is selected from hydrogen and alkyl; and Ar is substituted phenyl or substituted pyridyl.

- 5 In another embodiment the present invention provides compounds of formula I wherein Z is $-C(=O)-$; R and R' are both CH_3 ; R^5 is selected from methyl, ethyl, n-propyl, isopropyl, halogen, cyano, methoxy, and ethoxy; R^{10} is selected from hydrogen and alkyl; and Ar is phenyl substituted with a group selected from a) optionally substituted alkenyl; b) optionally-substituted alkyl having one two or three substituents independently
 10 selected from $-NR^d(C=O)NR^aR^b$, $-NR^a(C=O)OR^c$ and $-NR^aS(O)_2R^c$ wherein R^a , R^b and R^d are independently hydrogen, C_{1-6} alkyl, aryl, heteroaryl, cycloalkyl, or heterocyclo, and R^c is alkyl, aryl, heteroaryl, cycloalkyl, or heterocyclo, and each of R^a , R^b , R^c , and R^d in turn is optionally substituted with one, two, or three of alkyl, halo, haloalkyl, OR^e , haloalkoxy, cyano, $-NR^eR^f$, $-SO_2(alkyl)$, $-CO_2R^e$, $-C(=O)R^e$, and/or $-NR^eC(=O)R^f$, and/or
 15 a C_{1-6} alkyl substituted with one to two of halo, OR^e , haloalkoxy, cyano, $-NR^eR^f$, $-SO_2(alkyl)$, $-CO_2R^e$, $-C(=O)R^e$, and/or $-NR^eC(=O)R^f$, wherein R^e and R^f are independently hydrogen, alkyl, aminoalkyl, hydroxyalkyl, alkoxyalkyl, cyanoalkyl, haloalkyl, or alkylaminoalkyl; and c) $-NR^jS(O)_2NR^pR^q$, $-NR^jC(=O)NR^pR^q$, $-NR^pC(=O)OR^r$, $-NR^pC(=O)C(=O)R^q$ and $-NR^jC(=O)NR^pOR^q$ wherein each R^j , R^p , R^q ,
 20 and R^r is independently hydrogen, alkyl, substituted alkyl, cycloalkyl, heterocyclo, aryl, or heteroaryl, except R^r is not hydrogen, or alternatively, R^p and R^q when attached to the same nitrogen atom may be taken together to form a heterocyclo or heteroaryl.

It should be understood that when any of R^j , R^p , R^q , and R^r is and/or forms a ring, said cyclic groups in turn are optionally substituted with one to three substituents selected
 25 from the group consisting of alkyl, substituted alkyl, halogen, haloalkoxy, trifluoromethyl, cyano, nitro, $-OR^a$, $-SR^a$, $-S(O)R^c$, $-S(O)_2R^c$, $-C(=O)R^a$, $-C(=O)NR^aR^b$, $-C(O)_2R^a$, $-C(O)_2NR^aR^b$, $-S(O)_2NR^aR^b$, $-NR^aR^b$, and $-NR^a(C=O)R^b$.

In another embodiment the present invention provides compounds of formula I wherein Z is $-C(=O)-$; R and R' are both CH_3 ; R^5 is selected from methyl, ethyl, n-propyl, isopropyl, halogen, cyano, methoxy, and ethoxy; R^{10} is selected from hydrogen and alkyl;
 30 and Ar is



and R⁷ and R⁸ are selected from

- (i) hydroxyalkoxy, optionally substituted pyrrolodiny, alkylaminoalkyl, alkylsulfonyl-aminoalkyl, haloalkylaminoalkyl, alkoxyalkylaminoalkyl, hydroxyalkylaminoalkyl, aminoalkoxy, aminoalkyl, ureidylalkyl, carbamylalkyl, acetamidinyl, formamidinyl,
- 5 optionally substituted imidazoliny, optionally substituted pyrrolodinylmethyl, optionally substituted imidazolinylmethyl, alkoxyalkylaminocarbonyl, hydroxyalkylaminocarbonyl, aminoalkylaminocarbonyl, heterocyclalkylaminocarbonyl, and heterocyclalkylamino;
- (ii) methyl-(3,3,3-trifluoro-propyl)-amino-methyl-, 2-hydroxy-3-methoxy-propoxy-, methoxy-, 3-hydroxy-2-hydroxymethyl-propoxy-, 1-(dimethylamino-carbonyl-methyl-
- 10 amino)-ethyl-, 1-[(methoxycarbonyl)-methylamino]-ethyl-, 3-hydroxy-2-hydroxymethyl-propoxy-, 2,3-dihydroxypropoxy-, (S)-4-methoxy-2-oxo-pyrrolidin-1-yl-, 2-methoxyethoxy-carbonylaminomethyl-, methanesulfonyl-N-methylamino-methyl-, 4,5-dihydro-oxazol-2-yl-, 2-oxo-pyrrolidin-1-yl-, 2-[(2-hydroxy-ethyl)-methyl-amino]-ethoxy-, 3-hydroxy-2-oxo-pyrrolidin-1-yl-, 2-methylamino-ethyl-, (S)-4-hydroxy-2-oxo-
- 15 pyrrolidin-1-yl-, 2-(N-ethyl-N-methyl-amino)-ethoxy-, 1-(methanesulfonyl-N-methylamino)-ethyl-, 2-(N-ethyl-N-methylamino)-ethyl-, 2-methylamino-ethoxy-, ethyl-(2-hydroxyethyl)-amino-methyl-, 2-hydroxy-ethoxy-, ethyl-(2-methoxyethyl)-aminomethyl-, 2-hydroxy-ethyl-methylamino-methyl-, 3-methoxy-pyrrolidin-1-ylmethyl-, N-ethyl-N-methylamino-methyl-, 2-methoxyethyl-aminomethyl-, 4-hydroxy-
- 20 piperidin-1-ylmethyl-, hydroxy-pyrrolidin-1-ylmethyl-, aminomethyl-, ethylamino-methyl-, 3-dimethylamino-propoxy-, pyrrolidin-1-ylmethyl-, azetidin-1-ylmethyl-, 1,3-dimethyl-imidazolidin-2-ylideneamino-, 1-(methylamino-methyl-carbonyl-methylamino)-ethyl-, N,N-dimethylacetamidinyl-, 1-[(2-methoxyethyl)-methylamino]-ethyl-, methylamino-methyl-, dimethylamino-methyl-, 3,4-dimethylimidazoline-2,4-
- 25 dione-1-yl-, 3-methylimidazoline-2,4-dione-1-yl-, 1-dimethylamino-ethyl-, 1-methylamino-ethyl-, cyclopropylamino-methyl-, isopropylamino-methyl-, N-methyl-N-propylamino-methyl-, methylamino-methyl-, N-ethyl-N-methyl-amino-methyl-, 3-methyl-2-oxo-pyrrolidin-1-yl-, 2-oxo-imidazolidin-1-yl-, 2-methoxyethyl-amino-methyl-, 1-methyl-piperidin-4-yl-, pyrrolidin-1-yl-, 4-methyl-piperazin-1-yl-, 4-methyl-
- 30 piperazin-1-yl-, N,N-dimethyl-formamidinyl-, methyl-, 4,5-dihydro-3H-pyrrol-2-ylamino-, acetamidinyl-, 4,5-Dihydro-1H-imidazol-2-ylamino-, 4,4-dimethyl-4,5-dihydro-imidazol-1-yl-, pyrimidin-2-ylamino-, (1H-Imidazol-2-ylmethyl)-amino-, amino-, cyano-, bromo-, fluoro, methylamino-, methylaminomethyl-, aminomethyl-, cyanomethyl-, hydroxymethyl-, morpholin-4-ylmethyl-, ethylaminomethyl-, (2,2,2-
- 35 trifluoro-ethylamino)-methyl-, [(2-hydroxy-ethyl)-methylamino]-methyl-, [(2-methoxy-ethyl)-methylamino]-methyl-, [ethyl-(2-methoxy-ethyl)-amino]-methyl-, methoxymethylcarbonyl-N-methylaminomethyl-, 1-methyl-4,5-dihydro-1H-imidazol-2-

ylmethyl-, 1-isopropyl-4,5-dihydro-1*H*-imidazol-2-ylmethyl-, 1-methyl-piperidin-4-ylmethyl-, methanesulfonyl-, dimethylaminosulfonyl-, 2-methoxyethylaminosulfonyl-, 3-methoxypropylaminosulfonyl-, carboxy-, methy-amino-carbonyl-, 2-dimethylamino-ethyl)-aminocarbonyl-, 2-dimethylaminoethyl-methylaminocarbonyl-, 3-

5 dimethylaminopropyl-ethylaminocarbonyl-, 2-hydroxyethylaminocarbonyl-, 2-hydroxypropylaminocarbonyl-, 2-hydroxyethyl-methylaminocarbonyl-, 2-methoxyethylaminocarbonyl-, 3-methoxypropylaminocarbonyl-, cyanomethylaminocarbonyl-, tetrahydrofuran-2-ylmethylaminocarbonyl-, furan-2-ylmethylaminocarbonyl-, (2-morpholin-4-yl)-ethylaminocarbonyl-, (4-hydroxymethyl)-

10 piperidin-1-ylaminocarbonyl-, 2-hydroxyethoxy-, methanesulfinylmethoxy-, pyrimidin-2-yloxy-, hydroxy-, methanesulfonylmethoxy-, carboxymethoxy-, 2-methoxy-ethoxy-, 3-methoxy-propyloxy-, 2-dimethylamino-ethoxy-, 2-ethylamino-ethoxy-, methanesulfonylamino-, dimethylaminosulfonylamino-, dimethylaminosulfonyl-N-methylamino-, methylcarbonylamino-, chloro-, dimethylaminocarbonylamino-,

15 methoxycarbonylamino-, methoxycarbonyl-carbonyl-methylamino-, methoxyaminocarbonylamino- (183), 2-hydroxyethyl-amino-, 3-hydroxypropyl-methylamino-, 3-hydroxypropyl-amino-, 2-methoxyethyl)-amino-, 2-methoxyethyl)-methylamino-, 3-hydroxypropyl)-methylamino-, (3-methoxypropyl)-amino-, 3-methoxypropyl)-methylamino-, bis-(2-hydroxyethyl)-amino-, (2-

20 methylcarbonylamino)-ethylamino-, acetamidinyl-, N,N'-dimethylacetamidinyl-, N-methylacetamidinyl-, N,N-dimethylaminoformamidyl-, N,N-dimethyl-aminoisobutyramidinyl-, N,N-dimethylamino-(3-methoxy)-propionamidyl-, N,N-dimethylcyclobutane-carboxamidinyl-, imidazolidin-2-ylideneamino-, 1-methyl-pyrrolidin-(2*Z*)-ylideneamino-, 2-oxo-tetrahydro-furan-3-ylamino-, (4,5-Dihydro-3*H*-


25 pyrrol-2-yl)-methyl-amino-, 2-chloro-pyrimidin-4-ylamino-, 4-chloro-pyrimidin-2-ylamino-, 2-chloro-5-methyl-pyrimidin-4-ylamino-, 2-oxo-pyrrolidin-1-yl-, 1-methyl-4,5-dihydro-1*H*-imidazol-2-yl-, 2-methyl-4,5-dihydro-imidazol-1-yl-, 4,4-dimethyl-4,5-dihydro-imidazol-1-yl-, 2,4,4-trimethyl-4,5-dihydro-imidazol-1-yl-, 4-methyl-imidazol-1-yl-, 4-methyl-imidazol-1-yl-, 2-methyl-imidazol-1-yl-, pyrazol-1-yl-, 3-methyl-

30 pyrazol-1-yl-, 3,5-dimethyl-pyrazol-1-yl-, pyrrolidin-2-yl-, pyrrolidin-3-yl-, 1-methyl-4,5-dihydro-1*H*-imidazol-2-yl-, 1-ethyl-4,5-dihydro-1*H*-imidazol-2-yl-, 1-isopropyl-4,5-dihydro-1*H*-imidazol-2-yl-, 1-isopropyl-4,5-dihydro-1*H*-imidazol-2-yl-, 1-(2-hydroxy-ethyl)-4,5-dihydro-1*H*-imidazol-2-yl-, 1-(2-methoxy-ethyl)-4,5-dihydro-1*H*-imidazol-2-yl-, 1*H*-Imidazol-2-yl-, 1-methyl-1*H*-imidazol-2-yl-, 1-(2-hydroxy-ethyl)-1*H*-imidazol-2-

35 yl-, 1,1-dioxo-1*λ*6-isothiazolidin-2-yl-, 2-oxo-oxazolidin-3-yl-, 3-methyl-2-oxo-pyrrolidin-1-yl-, ethyl-2-oxo-pyrrolidin-1-yl-), pyrrolidine-2,5-dione-1-yl-, 2-oxo-

imidazolidin-1-yl-, 3-methyl-2-oxo-imidazolidin-1-yl-, 3-ethyl-2-oxo-imidazolidin-1-yl-, 3-(2-methoxy-ethyl)-2-oxo-imidazolidin-1-yl-, 1*H*-tetrazol-5-yl-, ethoxy-, 1-pyrrolidin-1-yl-ethyl-, 1-Azetidin-1-yl-ethyl-, aminosulfonyl-, N,N-dimethylamino-acetamidinesulfonyl-, and 1-methanesulfonyl-methylamino)-ethyl-; or

5 (iii)



wherein

R^{32} is hydrogen or alkyl;

R^{33} is hydrogen, alkyl, hydroxy or alkoxy;

10 R^{34} is hydroxy or alkoxy; and

E is $(C=O)$ or CH_2 .

In another embodiment the present invention provides compounds of formula I wherein Z is $-C(=O)-$; R and R' are both CH_3 ; R^5 is selected from methyl, ethyl, n-propyl, isopropyl, halogen, cyano, methoxy, and ethoxy; R^{10} is selected from hydrogen and alkyl; and Ar is 2,3-dihydro-1*H*-isoindol-5-yl optionally substituted at the 2-position with methyl or 2-methoxyethyl.

In another embodiment the present invention provides compounds of formula I wherein Z is $-C(=O)-$; R and R' are both CH_3 ; R^5 is selected from methyl, ethyl, n-propyl, isopropyl, halogen, cyano, methoxy, and ethoxy; R^{10} is selected from hydrogen and alkyl; and Ar is 1*H*-indol-3-yl.

In another embodiment the present invention provides compounds of formula I wherein Z is $-C(=O)-$; R and R' are both CH_3 ; R^5 is selected from methyl, ethyl, n-propyl, isopropyl, halogen, cyano, methoxy, and ethoxy; R^{10} is selected from hydrogen and alkyl; and Ar is pyridin 2-yl optionally substituted at the 4-position or 6-position with

hydroxyalkoxy, optionally substituted pyrrolodiny, alkylaminoalkyl, alkylsulfonylaminoalkyl, haloalkylaminoalkyl, alkoxyalkylaminoalkyl, hydroxyalkylaminoalkyl, aminoalkoxy, aminoalkyl, ureidylalkyl, carbamylalkyl, acetamidinyl, formamidinyl, optionally substituted imidazoliny, optionally substituted
 5 pyrrolodinylmethyl, optionally substituted imidazolinylmethyl, alkoxyalkylaminocarbonyl, hydroxyalkylaminocarbonyl, aminoalkylaminocarbonyl, heterocyclalkylaminocarbonyl, or heterocyclalkylamino.

In one embodiment the present invention provides compounds of formula I wherein Ar is



X is N or CR⁸;

R⁷ is selected from alkyl, substituted alkyl, halogen, cyano, -SO₂R¹¹, -NR¹²SO₂R¹¹, -NR¹²CO₂R^{11a}, -NR¹²C(=O)R^{11a}, -NR¹²R¹⁴, -C(=O)NR¹²R¹⁴, -SO₂NR¹²R¹⁴, -NR¹²C(=NR¹³)R¹⁴, -N=C(R¹⁵)R¹⁶, -SO₂N=C(R¹⁵)R¹⁶, -C(=O)R¹⁷, -CO₂R¹⁷, -OR¹⁷,
 15 heterocyclo, heteroaryl, phenyl, and cycloalkyl, wherein said heterocyclo, heteroaryl, phenyl, and cycloalkyl groups are optionally substituted with up to three R²⁵;

R⁸ is selected from hydrogen, alkyl, lower alkyl, halogen, cyano, halo(C₁₋₄)alkyl, hydroxy, lower alkoxy, amino, (C₁₋₄)alkylamino, (C₁₋₄)alkylamino(C₁₋₄)alkyl, hydroxy(C₁₋₄)alkyl, (C₁₋₄)alkoxy(C₁₋₄)alkyl, (C₁₋₄)alkylsulfonyl, pyrrolidinyl, and
 20 piperidinyl (said R⁸ groups in turn being optionally substituted with one to two of lower alkyl, lower alkoxy, cyano, and/or halogen);

R¹¹, R¹², R¹³ and R¹⁵ are independently selected from hydrogen, alkyl, hydroxyalkyl, and cycloalkyl, except R¹¹ is not hydrogen;

R^{11a} is selected from alkyl, alkylamino, aminoalkyl, -C(=O)alkyl, N(O)alkyl, and
 25 hydroxyalkyl;

R¹⁶ is independently amino or alkylamino;

or alternatively, R¹⁵ and R¹⁶ may be taken together to form a heterocyclo ring optionally substituted with up to three groups selected from R²⁵;


R¹⁴ and R¹⁷ are independently selected from hydrogen, alkyl, substituted alkyl, and
 30 heterocyclo or heteroaryl optionally substituted with up to three groups selected from R²⁵;

or alternatively, R^{12} and R^{14} when attached to the same nitrogen atom (as in $-NR^{12}R^{14}$, $-C(=O)NR^{12}R^{14}$, and $-SO_2NR^{12}R^{14}$), may be taken together to form a heterocyclo ring optionally substituted with up to three groups selected from R^{25} ;

R^{25} is at each occurrence independently selected as valence permits from alkyl,


- 5 substituted alkyl, halogen, haloalkoxy, trifluoromethyl, cyano, nitro, $-OR^{26}$, $-SR^{26}$, $-S(O)R^{26}$, $-S(O)_2R^{28}$, $-C(=O)R^{26}$, $-C(=O)NR^{26}R^{27}$, $-C(O)_2R^{26}$, $-C(O)_2NR^{26}R^{27}$, $-S(O)_2NR^{26}R^{27}$, $-NR^{26}R^{27}$, and/or $-NR^{26}(C=O)R^{27}$, wherein R^{26} and R^{27} are selected from hydrogen, C_{1-6} alkyl, aryl, heteroaryl, cycloalkyl and heterocyclo, and R^{28} is selected from C_{1-6} alkyl, aryl, heteroaryl, cycloalkyl and heterocyclo; and
- 10 m is 0, 1, or 2.

In another embodiment the present invention provides compounds of formula I wherein Ar is




R^7 is selected from

- 15 (i) pyrrolidinyl, pyrrolinyl, imidazoliny, $-NR^{12}R^{14}$, $-NR^{12}C(=NR^{13})R^{14}$, $-N=C(R^{15})(R^{16})$, $-OR^{17}$, and C_{1-4} alkyl optionally substituted with pyrrolidinyl, pyrrolinyl, imidazoliny, $-NR^{12a}R^{14a}$, and/or $-OR^{17a}$, wherein said pyrrolidinyl, pyrrolinyl and imidazoliny groups in turn optionally may be substituted with one to three of lower alkyl, halogen, cyano, and/or hydroxy(C_{1-4})alkyl;
- 20 R^{12} , R^{12a} , R^{13} , R^{14} , R^{14a} , R^{17} , and R^{17a} are independently selected from hydrogen, lower alkyl, hydroxy(C_{1-4} alkyl), $-C_{1-4}$ alkylene-O(lower alkyl), $-C_{1-4}$ alkylene- NH_2 , $-C_{1-4}$ alkylene- NH (lower alkyl), $-C_{1-4}$ alkylene- N (lower alkyl) $_2$, and $-C_{1-4}$ alkylene-(C_{3-6} cycloalkyl);
- R^{15} is independently selected from hydrogen, lower alkyl, hydroxy(C_{1-4})alkyl, and
- 25 C_{3-6} cycloalkyl, and R^{16} is independently amino or (C_{1-4})alkylamino; or alternatively, R^{15} and R^{16} may be taken together to form a pyrrolidinyl, pyrrolinyl, or imidazoliny ring, wherein said pyrrolidinyl, pyrrolinyl and imidazoliny groups optionally may be substituted with one to three of lower alkyl, hydroxy, lower alkoxy, halogen, cyano, amino, (C_{1-4})alkylamino, hydroxy(C_{1-4})alkyl, amino(C_{1-4})alkyl, and/or (C_{1-4})alkylamino(C_{1-4})alkyl; or
- 30



wherein R^{30} and R^{31} each independently is selected from hydrogen, methyl, ethyl, methoxyethyl and hydroxyethyl, and R^{32} is hydrogen, methyl or ethyl.

- 5 Preferred compounds are those compounds having the Formula (Ia)



and isomers or pharmaceutically-acceptable salts, hydrates, or prodrugs thereof,
wherein

Y is $-(CHR^1)-$ or $-(CHR^2-CHR^3)-$;

- 10 R^1 , R^2 and R^3 are independently selected from hydrogen, lower alkyl, and hydroxy, or R^2 and R^3 may together form a bond so that Y is ethenylene;

R and R' are lower alkyl;

R^5 is selected from halogen, cyano, hydroxy, $-R^6$, and $-OR^6$;


R^6 is selected from alkyl, aryl, heteroaryl, and cycloalkyl;

- 15 R^{10} is selected from hydrogen, alkyl, hydroxyalkyl and aminoalkyl; and

Ar is optionally-substituted aryl or heteroaryl.

In one embodiment the present invention provides compounds of formula Ia wherein R and R' are CH_3 ; R^5 is selected from C_{1-4} alkyl, halogen, cyano, hydroxy and C_{1-4} alkoxy; and R^{10} is selected from hydrogen, C_{1-4} alkyl, and hydroxy(C_{1-4})alkyl. In another


- 20 embodiment the present invention provides compounds of formula Ia wherein R and R' are CH_3 ; R^5 is methyl or methoxy; and R^{10} is methyl. In another embodiment the present invention provides compounds of formula Ia wherein R and R' are CH_3 ; R^5 is methyl or methoxy; R^{10} is methyl; and Y-Ar considered together are




R^1 is hydrogen, hydroxy, or lower alkyl; and

R^7 is selected from

- (i) one pyrrolidinyl, pyrrolinyl, imidazolinyl, $-NR^{12}R^{14}$, $-NR^{12}C(=NR^{13})R^{14}$, $-N=C(R^{15})(R^{16})$, $-OR^{17}$, and C_{1-4} alkyl optionally substituted with pyrrolidinyl, pyrrolinyl, imidazolinyl, $-NR^{12a}R^{14a}$, and/or $-OR^{17a}$, wherein said pyrrolidinyl, pyrrolinyl and imidazolinyl groups in turn optionally may be substituted with one to three groups selected from R^{25} ;
- R^{12a} , R^{14a} , R^{17} , and R^{17a} are independently selected from hydrogen, lower alkyl, hydroxy(C_{1-4} alkyl), $-C_{1-4}$ alkylene-O(lower alkyl), $-C_{1-4}$ alkylene- NH_2 , $-C_{1-4}$ alkylene-NH(lower alkyl), $-C_{1-4}$ alkylene-N(lower alkyl) $_2$, and $-C_{1-4}$ alkylene-(C_{3-6} cycloalkyl);
- R^{12} is hydrogen, alkyl, or hydroxy(C_{1-4} alkyl);
- R^{14} is selected from hydrogen, alkyl, hydroxyalkyl, alkoxyalkyl, aminoalkyl, alkyl-aminoalkyl, alkylamidylalkyl, cyanoalkyl, pyrrolidinyl, pyrrolinyl, imidazolinyl, pyrimidinyl, tetrahydrofuranyl, furanylalkyl, isothiazolidinyl, oxazolindinyl, tetrahydrofuranylalkyl, morpholinylalkyl, and imidazolylalkyl, said R^{14} cyclic groups in turn being optionally substituted with one, two or three groups where valence allows selected from R^{25} ;
- or alternatively, R^{12} and R^{14} may be taken together to form pyrrolidinyl, piperidinyl, imidazolinyl, imidazolidinyl, pyrazolyl, or imidazolyl, said rings optionally substituted where valence allows with one, two or three groups selected from R^{25} ;
- R^{15} is independently selected from hydrogen, lower alkyl, hydroxy(C_{1-4} alkyl), and C_{3-6} cycloalkyl;
- R^{16} is independently amino or (C_{1-4})alkylamino;
- or alternatively, R^{15} and R^{16} may be taken together to form a pyrrolidinyl or imidazolinyl ring, wherein said pyrrolidinyl, pyrrolinyl and imidazolinyl groups optionally may be substituted with one to three groups selected from R^{25} ; and
- R^{25} at each occurrence is independently selected from lower alkyl, halogen, cyano, hydroxy(C_{1-4} alkyl), and (C_{1-4})alkoxy(C_{1-4})alkyl; or





wherein R^{30} and R^{31} each independently is selected from hydrogen, methyl, ethyl, methoxyethyl and hydroxyethyl, and R^{32} is hydrogen, methyl or ethyl.

In compounds of Formula (I) as set forth in the description and claims herein, and in
 5 Formula (Ia) as set forth above, preferred compounds are those in which R^5 is not hydrogen. Even more preferred are compounds where R^5 is selected from halogen, hydroxy, cyano, $-R^6$, and $-OR^6$, wherein R^6 is selected from C_{1-4} alkyl, aminoalkyl, C_{1-4} alkylamino(C_{1-4} alkyl), hydroxyalkyl, alkoxyalkyl, phenoxyalkyl, benzyloxyalkyl, cycloalkylalkyl, phenyl, benzyl, and cycloalkyl, wherein each of said phenyl, benzyl, and
 10 cycloalkyl groups is optionally substituted with one to two of lower alkyl, substituted lower alkyl, cyano, and/or halogen. Even more preferred are compounds wherein R^5 is selected from methyl, ethyl, n-propyl, isopropyl, cyano, halogen, methoxy, and ethoxy. Most preferred are compounds wherein R^5 is methyl or methoxy.

In compounds of Formulae (I) and (Ia), preferably R and R' are both CH_3 .


15 In compounds of Formulae (I) and (Ia), preferably R^{10} is selected from hydrogen, alkyl, amino(C_{1-6})alkyl, (C_{1-6})alkylamino(C_{1-6})alkyl, hydroxy(C_{1-6})alkyl, and an optionally-substituted five or six membered heterocyclo or C_{3-7} cycloalkyl. More preferred are compounds wherein R^{10} is selected from hydrogen, lower alkyl, amino(C_{1-4})alkyl, (C_{1-4})alkyl-amino(C_{1-2})alkyl, hydroxy(C_{1-4})alkyl, and piperidinyl optionally-substituted with lower
 20 alkyl or benzyl. Even more preferred are compounds where R^{10} is selected from hydrogen, lower alkyl, hydroxymethyl, hydroxyethyl, $-C_{1-4}$ alkylene- NH_2 , $-C_{1-4}$ alkylene- $NH(CH_3)$, and/or $-C_{1-4}$ alkylene- $N(CH_3)_2$, and most preferred are compounds wherein R^{10} is methyl or ethyl.

In certain embodiments of Formulae (I) and (Ia), Ar is is phenyl substituted with a group
 25 selected from:

- a) optionally substituted alkenyl;
- b) optionally-substituted alkyl having one two or three substituents independently selected from $-NR^d(C=O)NR^aR^b$, $-NR^a(C=O)OR^c$ and $-NR^aS(O)_2R^c$ wherein R^a , R^b , and R^d are independently hydrogen, C_{1-6} alkyl, aryl, heteroaryl, cycloalkyl, or heterocyclo, and
 30 R^c is alkyl, aryl, heteroaryl, cycloalkyl, or heterocyclo, and each of R^a , R^b , R^c , and R^d in


- turn is optionally substituted with one, two, or three of alkyl, halo, haloalkyl, OR^e, haloalkoxy, cyano, -NR^eR^f, -SO₂(alkyl), -CO₂R^e, -C(=O)R^e, and/or -NR^eC(=O)R^f, and/or a C₁₋₆alkyl substituted with one to two of halo, OR^e, haloalkoxy, cyano, -NR^eR^f, -SO₂(alkyl), -CO₂R^e, -C(=O)R^e, and/or -NR^eC(=O)R^f, wherein R^e and R^f are
- 5 independently hydrogen, alkyl, aminoalkyl, hydroxyalkyl, alkoxyalkyl, cyanoalkyl, haloalkyl, or alkylaminoalkyl; and
- c) -NR^jS(O)₂NR^pR^q, -NR^jC(=O)NR^pR^q, -NR^pC(=O)OR^r, -NR^pC(=O)C(=O)R^q, and -NR^jC(=O)NR^pOR^q wherein each R^j, R^p, R^q, and R^r is independently hydrogen, alkyl, substituted alkyl, cycloalkyl, heterocyclo, aryl, or heteroaryl, except R^r is not hydrogen, or
- 10 alternatively, R^p and R^q when attached to the same nitrogen atom may be taken together to form a heterocyclo or heteroaryl. It should be understood that when any of R^j, R^p, R^q, and R^r is and/or forms a ring, said cyclic groups in turn are optionally substituted with one to three substituents selected from the group consisting of alkyl, substituted alkyl, halogen, haloalkoxy, trifluoromethyl, cyano, nitro, -OR^a, -SR^a, -S(O)R^c, -S(O)₂R^c,
- 15 -C(=O)R^a, -C(=O)NR^aR^b, -C(O)₂R^a, -C(O)₂NR^aR^b, -S(O)₂NR^aR^b, -NR^aR^b, and -NR^a(C=O)R^b.

In certain embodiments of Formulae (I) and (Ia), Ar is



- and R⁷ may be selected from hydroxyalkoxy, optionally substituted pyrrolodiny, alkyl- aminoalkyl, alkylsulfonylaminoalkyl, haloalkylaminoalkyl, alkoxyalkylaminoalkyl, hydroxyalkylaminoalkyl, aminoalkoxy, aminoalkyl, ureidylalkyl, carbamylalkyl, acetamidinyl, formamidinyl, optionally substituted imidazoliny, optionally substituted pyrrolodinylmethyl, optionally substituted imidazolinylmethyl, alkoxyalkylaminocarbonyl, hydroxyalkylaminocarbonyl, aminoalkylaminocarbonyl, heterocyclalkylaminocarbonyl, and heterocyclalkylamino.
- 20
- 25

In other embodiments of Formulae (I) and (Ia) where Ar is




R⁷ may be selected from methyl-(3,3,3-trifluoro-propyl)-amino-methyl-, 2-hydroxy-3-methoxy-propoxy-, methoxy-, 3-hydroxy-2-hydroxymethyl-propoxy-, 1-


(dimethylamino-carbonyl-methylamino)-ethyl-, 1-[(methoxycarbonyl)-methylamino]-ethyl-, 3-hydroxy-2-hydroxymethyl-propoxy-, 2,3-dihydroxypropoxy-, (S)-4-methoxy-2-oxo-pyrrolidin-1-yl-, 2-methoxyethoxy-carbonylaminomethyl-, methanesulfonyl-N-methylamino-methyl-, 4,5-dihydro-oxazol-2-yl-, 2-oxo-pyrrolidin-1-yl-, 2-[(2-hydroxy-ethyl)-methyl-amino]-ethoxy-, 3-hydroxy-2-oxo-pyrrolidin-1-yl-, 2-methylamino-ethyl-, (S)-4-hydroxy-2-oxo-pyrrolidin-1-yl-, 2-(N-ethyl-N-methyl-amino)-ethoxy-, 1-(methanesulfonyl-N-methylamino)-ethyl-, 2-(N-ethyl-N-methylamino)-ethyl-, 2-methylamino-ethoxy-, ethyl-(2-hydroxyethyl)-amino-methyl-, 2-hydroxy-ethoxy-, ethyl-(2-methoxyethyl)-aminomethyl-, 2-hydroxy-ethyl-methylamino-methyl-, 3-methoxy-pyrrolidin-1-ylmethyl-, N-ethyl-N-methylamino-methyl-, 2-methoxyethyl-aminomethyl-, 4-hydroxy-piperidin-1-ylmethyl-, hydroxy-pyrrolidin-1-ylmethyl-, aminomethyl-, ethylamino-methyl-, 3-dimethylamino-propoxy-, pyrrolidin-1-ylmethyl-, azetidin-1-ylmethyl-, 1,3-dimethyl-imidazolidin-2-ylideneamino)-, 1-(methylamino-methyl-carbonyl-methylamino)-ethyl-, N,N-dimethylacetamidinyl-, 1-[(2-methoxyethyl)-methylamino]-ethyl-, methylamino-methyl-, dimethylamino-methyl-, 3,4-dimethylimidazoline-2,4-dione-1-yl-, 3-methylimidazoline-2,4-dione-1-yl-, 1-dimethylamino-ethyl-, 1-methylamino-ethyl-, cyclopropylamino-methyl-, isopropylamino-methyl-, N-methyl-N-propylamino-methyl-, methylamino-methyl-, N-ethyl-N-methyl-amino-methyl-, 3-methyl-2-oxo-pyrrolidin-1-yl-, 2-oxo-imidazolidin-1-yl-, 2-methoxyethyl-amino-methyl-, 1-methyl-piperidin-4-yl-, pyrrolidin-1-yl-, 4-methyl-piperazin-1-yl-, 4-methyl-piperazin-1-yl-, N,N-dimethyl-formamidinyl-, methyl-, 4,5-dihydro-3H-pyrrol-2-ylamino-, acetamidinyl-, 4,5-Dihydro-1H-imidazol-2-ylamino)-, 4,4-dimethyl-4,5-dihydro-imidazol-1-yl-, pyrimidin-2-ylamino-, (1H-Imidazol-2-ylmethyl)-amino-, amino-, cyano-, bromo-, fluoro, methylamino-, methylaminomethyl-, aminomethyl-, cyanomethyl-, hydroxymethyl-, morpholin-4-ylmethyl-, ethylaminomethyl-, (2,2,2-trifluoro-ethylamino)-methyl-, [(2-hydroxy-ethyl)-methylamino]-methyl-, [(2-methoxy-ethyl)-methylamino]-methyl-, [ethyl-(2-methoxy-ethyl)-amino]-methyl-, methoxymethylcarbonyl-N-methylaminomethyl-, 1-methyl-4,5-dihydro-1H-imidazol-2-ylmethyl-, 1-isopropyl-4,5-dihydro-1H-imidazol-2-ylmethyl-, 1-methyl-piperidin-4-ylmethyl-, methanesulfonyl-, dimethylaminosulfonyl-, 2-methoxyethylaminosulfonyl-, 3-methoxypropylaminosulfonyl-, carboxy-, methy-amino-carbonyl-, 2-dimethylamino-ethyl)-aminocarbonyl-, 2-dimethylaminoethyl-methylaminocarbonyl-, 3-dimethylaminopropyl-ethylaminocarbonyl-, 2-hydroxyethylaminocarbonyl-, 2-hydroxypropylaminocarbonyl-, 2-hydroxyethyl-methylaminocarbonyl-, 2-methoxyethylaminocarbonyl-, 3-methoxypropylaminocarbonyl-, cyanomethylaminocarbonyl-, tetrahydrofuran-2-ylmethylaminocarbonyl-, furan-2-ylmethylaminocarbonyl-, (2-morpholin-4-yl)-ethylaminocarb-

onyl-, (4-hydroxymethyl)-piperidin-1-ylaminocarbonyl-, 2-hydroxyethoxy-,
methanesulfinylmethoxy-, pyrimidin-2-yloxy-, hydroxy-, methanesulfonylmethoxy-,
carboxymethoxy-, 2-methoxy-ethoxy-, 3-methoxy-propyloxy-, 2-dimethylamino-
ethoxy-, 2-ethylamino-ethoxy-, methanesulfonylamino-, dimethylaminosulfonylamino-,
5 dimethylaminosulfonyl-N-methylamino-, methylcarbonylamino-, chloro-,
dimethylaminocarbonylamino-, methoxycarbonylamino-, methoxycarbonyl-carbonyl-
methylamino-, methoxyaminocarbonylamino- (183), 2-hydroxyethyl-amino-, 3-
hydroxypropyl-methylamino-, 3-hydroxypropyl-amino-, 2-methoxyethyl)-amino-, 2-
methoxyethyl)-methylamino-, 3-hydroxypropyl)-methylamino-, (3-methoxypropyl)-
10 amino-, 3-methoxypropyl)-methylamino-, bis-(2-hydroxyethyl)-amino-, (2-
methylcarbonylamino)-ethylamino-, acetamidinyl-, N,N'-dimethylacetamidinyl-, N-
methylacetamidinyl-, N,N-dimethylaminoformamidyl-, N,N-
dimethylaminoisobutyramidinyl-, N,N-dimethylamino-(3-methoxy)-propionamidyl-,
N,N-dimethylcyclobutane-carboxamidinyl-, imidazolidin-2-ylideneamino-, 1-methyl-
15 pyrrolidin-(2Z)-ylideneamino-, 2-oxo-tetrahydro-furan-3-ylamino-, (4,5-dihydro-3H-
pyrrol-2-yl)-methyl-amino-, 2-chloro-pyrimidin-4-ylamino-, 4-chloro-pyrimidin-2-
ylamino-, 2-chloro-5-methyl-pyrimidin-4-ylamino-, 2-oxo-pyrrolidin-1-yl-, 1-methyl-
4,5-dihydro-1H-imidazol-2-yl-, 2-methyl-4,5-dihydro-imidazol-1-yl-, 4,4-dimethyl-4,5-
dihydro-imidazol-1-yl-, 2,4,4-trimethyl-4,5-dihydro-imidazol-1-yl-, 4-methyl-imidazol-
20 1-yl-, 4-methyl-imidazol-1-yl-, 2-methyl-imidazol-1-yl-, pyrazol-1-yl-, 3-methyl-
pyrazol-1-yl-, 3,5-Dimethyl-pyrazol-1-yl-, pyrrolidin-2-yl-, pyrrolidin-3-yl-, 1-methyl-
4,5-dihydro-1H-imidazol-2-yl-, 1-ethyl-4,5-dihydro-1H-imidazol-2-yl-, 1-isopropyl-4,5-
dihydro-1H-imidazol-2-yl-, 1-isopropyl-4,5-dihydro-1H-imidazol-2-yl-, 1-(2-hydroxy-
ethyl)-4,5-dihydro-1H-imidazol-2-yl-, 1-(2-methoxy-ethyl)-4,5-dihydro-1H-imidazol-2-
25 yl-, 1H-Imidazol-2-yl-, 1-methyl-1H-imidazol-2-yl-, 1-(2-hydroxy-ethyl)-1H-imidazol-2-
yl-, 1,1-Dioxo-1λ6-isothiazolidin-2-yl-, 2-oxo-oxazolidin-3-yl-, 3-methyl-2-oxo-
pyrrolidin-1-yl-, ethyl-2-oxo-pyrrolidin-1-yl-, pyrrolidine-2,5-dione-1-yl-, 2-oxo-
imidazolidin-1-yl-, 3-methyl-2-oxo-imidazolidin-1-yl-, 3-ethyl-2-oxo-imidazolidin-1-yl-,
3-(2-methoxy-ethyl)-2-oxo-imidazolidin-1-yl-, 1H-tetrazol-5-yl-, ethoxy-, 1-
30 pyrrolidin-1-yl-ethyl-, 1-azetidin-1-yl-ethyl-, aminosulfonyl-, N,N-dimethylamino-
acetamidinesulfonyl-, and 1-methanesulfonyl-methylamino)-ethyl-.

In still other embodiments of Formulae (I) and (Ia) where Ar is




R^7 may be selected from:




wherein R^{30} and R^{31} each independently is selected from hydrogen, methyl, ethyl, methoxyethyl and hydroxyethyl, and R^{32} is hydrogen, methyl or ethyl.

5 In yet other embodiments of Formulae (I) and (Ia) where Ar is



R^7 may be selected from



10 wherein


R^{32} is hydrogen or alkyl;

R^{33} is hydrogen, alkyl, hydroxy or alkoxy;

R³⁴ is hydroxy or alkoxy; and


E is (C=O) or CH₂.

In certain embodiments of Formulae (I) and (Ia), Ar is



- 5 and R⁸ may be selected from hydroxyalkoxy, optionally substituted pyrrolodiny, alkyl-aminoalkyl, alkylsulfonylaminoalkyl, haloalkylaminoalkyl, alkoxyalkylaminoalkyl, hydroxyalkylaminoalkyl, aminoalkoxy, aminoalkyl, ureidylalkyl, carbamylalkyl, acetamidiny, formamidiny, optionally substituted imidazoliny, optionally substituted pyrrolodinylmethyl, optionally substituted imidazolinylmethyl,
- 10 alkoxyalkylaminocarbonyl, hydroxyalkylaminocarbonyl, aminoalkylaminocarbonyl, heterocyclalkylaminocarbonyl, and heterocyclalkylamino.

In other embodiments of Formulae (I) and (Ia) where Ar is




- R⁸ may be selected from methyl-(3,3,3-trifluoro-propyl)-amino-methyl-, 2-hydroxy-3-
- 15 methoxy-propoxy-, methoxy-, 3-hydroxy-2-hydroxymethyl-propoxy-, 1-(dimethylamino-carbonyl-methylamino)-ethyl-, 1-[(methoxycarbonyl)-methylamino]-ethyl-, 3-hydroxy-2-hydroxymethyl-propoxy-, 2,3-dihydroxypropoxy-, (S)-4-methoxy-2-oxo-pyrrolidin-1-yl-, 2-methoxyethoxy-carbonylaminomethyl-, methanesulfonyl-N-methylamino-methyl-, 4,5-dihydro-oxazol-2-yl-, 2-oxo-pyrrolidin-1-yl-, 2-[(2-hydroxy-
- 20 ethyl)-methyl-amino]-ethoxy-, 3-hydroxy-2-oxo-pyrrolidin-1-yl-, 2-methylamino-ethyl-, (S)-4-hydroxy-2-oxo-pyrrolidin-1-yl-, 2-(N-ethyl-N-methyl-amino)-ethoxy-, 1-(methanesulfonyl-N-methylamino)-ethyl-, 2-(N-ethyl-N-methylamino)-ethyl-, 2-methylamino-ethoxy-, ethyl-(2-hydroxyethyl)-amino-methyl-, 2-hydroxy-ethoxy-, ethyl-(2-methoxyethyl)-aminomethyl-, 2-hydroxy-ethyl-methylamino-methyl-, 3-methoxy-
- 25 pyrrolidin-1-ylmethyl-, N-ethyl-N-methylamino-methyl-, 2-methoxyethylaminomethyl-, 4-hydroxy-piperidin-1-ylmethyl-, hydroxy-pyrrolidin-1-ylmethyl-, aminomethyl-, ethylamino-methyl-, 3-dimethylamino-propoxy-, pyrrolidin-1-ylmethyl-, azetidin-1-ylmethyl-, 1,3-dimethyl-imidazolidin-2-ylideneamino-, 1-(methylamino-methyl-carbonyl-methylamino)-ethyl-, N,N-dimethylacetamidiny, 1-[(2-
- 30 methoxyethyl)-methylamino]-ethyl-, methylamino-methyl-, dimethylamino-methyl-,


3,4-dimethylimidazoline-2,4-dione-1-yl-, 3-methylimidazoline-2,4-dione-1-yl)-, 1-dimethylamino-ethyl-, 1-methylamino-ethyl-, cyclopropylamino-methyl-, isopropylamino-methyl-, N-methyl-N-propylamino-methyl-, methylamino-methyl-, N-ethyl-N-methyl-amino-methyl-, 3-methyl-2-oxo-pyrrolidin-1-yl-, 2-oxo-imidazolidin-1-yl-, 2-methoxyethyl-amino-methyl-, 1-methyl-piperidin-4-yl-, pyrrolidin-1-yl-, 4-methyl-piperazin-1-yl-, 4-methyl-piperazin-1-yl-, N,N-dimethyl-formamidinyl)-, methyl-, 4,5-dihydro-3H-pyrrol-2-ylamino-, acetamidinyl-, 4,5-Dihydro-1H-imidazol-2-ylamino)-, 4,4-dimethyl-4,5-dihydro-imidazol-1-yl)-, pyrimidin-2-ylamino-, (1H-Imidazol-2-ylmethyl)-amino-, amino-, cyano-, bromo-, fluoro, methylamino-, methylaminomethyl-, aminomethyl-, cyanomethyl-, hydroxymethyl-, morpholin-4-ylmethyl-, ethylaminomethyl-, (2,2,2-trifluoro-ethylamino)-methyl-, [(2-hydroxyethyl)-methylamino]-methyl-, [(2-methoxy-ethyl)-methylamino]-methyl-, [ethyl-(2-methoxy-ethyl)-amino]-methyl-, methoxymethylcarbonyl-N-methylaminomethyl-, 1-methyl-4,5-dihydro-1H-imidazol-2-ylmethyl-, 1-isopropyl-4,5-dihydro-1H-imidazol-2-ylmethyl-, 1-methyl-piperidin-4-ylmethyl-, methanesulfonyl-, dimethylaminosulfonyl-, 2-methoxyethylaminosulfonyl-, 3-methoxypropylaminosulfonyl-, carboxy-, methyamino-carbonyl-, 2-dimethylamino-ethyl)-aminocarbonyl-, 2-dimethylaminoethyl-methylaminocarbonyl-, 3-dimethylaminopropyl-ethylaminocarbonyl-, 2-hydroxyethylaminocarbonyl-, 2-hydroxypropylaminocarbonyl-, 2-hydroxyethyl-methylaminocarbonyl-, 2-methoxyethylaminocarbonyl-, 3-methoxypropylaminocarbonyl-, cyanomethylaminocarbonyl-, tetrahydrofuran-2-ylmethylaminocarbonyl-, furan-2-ylmethylaminocarbonyl-, (2-morpholin-4-yl)-ethylaminocarbonyl-, (4-hydroxymethyl)-piperidin-1-ylaminocarbonyl-, 2-hydroxyethoxy-, methanesulfinylmethoxy-, pyrimidin-2-yloxy-, hydroxy-, methanesulfonylmethoxy-, carboxymethoxy-, 2-methoxy-ethoxy-, 3-methoxy-propyloxy-, 2-dimethylamino-ethoxy-, 2-ethylamino-ethoxy-, methanesulfonylamino-, dimethylaminosulfonylamino-, dimethylaminosulfonyl-N-methylamino-, methylcarbonylamino-, chloro-, dimethylaminocarbonylamino-, methoxycarbonylamino-, methoxycarbonyl-carbonyl-methylamino-, methoxyaminocarbonylamino-, 2-hydroxyethyl-amino-, 3-hydroxypropyl-methylamino-, 3-hydroxypropyl-amino-, 2-methoxyethyl)-amino-, 2-methoxyethyl)-methylamino-, 3-hydroxypropyl)-methylamino-, (3-methoxypropyl)-amino-, 3-methoxypropyl)-methylamino-, bis-(2-hydroxyethyl)-amino-, (2-methylcarbonylamino)-ethylamino-, acetamidinyl-, N,N'-dimethylacetamidinyl-, N-methylacetamidinyl-, N,N-dimethylaminoformamidyl-, N,N-dimethylaminoisobutyramidinyl-, N,N-dimethylamino-(3-methoxy)-propionamidyl-, N,N-dimethylcyclobutane-carboxamidinyl-, imidazolidin-2-ylideneamino-, 1-methyl-pyrrolidin-(2Z)-ylideneamino-, 2-oxo-tetrahydro-furan-3-ylamino-, (4,5-Dihydro-3H-

pyrrol-2-yl)-methyl-amino-, 2-chloro-pyrimidin-4-ylamino-, 4-chloro-pyrimidin-2-ylamino-, 2-chloro-5-methyl-pyrimidin-4-ylamino-, 2-oxo-pyrrolidin-1-yl-, 1-methyl-4,5-dihydro-1*H*-imidazol-2-yl-, 2-methyl-4,5-dihydro-imidazol-1-yl-, 4,4-dimethyl-4,5-dihydro-imidazol-1-yl-, 2,4,4-trimethyl-4,5-dihydro-imidazol-1-yl-, 4-methyl-imidazol-1-yl-, 4-methyl-imidazol-1-yl-, 2-methyl-imidazol-1-yl-, pyrazol-1-yl-, 3-methyl-pyrazol-1-yl-, 3,5-dimethyl-pyrazol-1-yl-, pyrrolidin-2-yl-, pyrrolidin-3-yl-, 1-methyl-4,5-dihydro-1*H*-imidazol-2-yl-, 1-ethyl-4,5-dihydro-1*H*-imidazol-2-yl-, 1-isopropyl-4,5-dihydro-1*H*-imidazol-2-yl-, 1-isopropyl-4,5-dihydro-1*H*-imidazol-2-yl-, 1-(2-hydroxy-ethyl)-4,5-dihydro-1*H*-imidazol-2-yl-, 1-(2-methoxy-ethyl)-4,5-dihydro-1*H*-imidazol-2-yl-, 1*H*-Imidazol-2-yl-, 1-methyl-1*H*-imidazol-2-yl-, 1-(2-hydroxy-ethyl)-1*H*-imidazol-2-yl-, 1,1-Dioxo-1*λ*6-isothiazolidin-2-yl-, 2-oxo-oxazolidin-3-yl-, 3-methyl-2-oxo-pyrrolidin-1-yl-, ethyl-2-oxo-pyrrolidin-1-yl-, pyrrolidine-2,5-dione-1-yl-, 2-oxo-imidazolidin-1-yl-, 3-methyl-2-oxo-imidazolidin-1-yl-, 3-ethyl-2-oxo-imidazolidin-1-yl-, 3-(2-methoxy-ethyl)-2-oxo-imidazolidin-1-yl-, 1*H*-tetrazol-5-yl-, ethoxy-, 1-pyrrolidin-1-yl-ethyl-, 1-Azetidin-1-yl-ethyl-, aminosulfonyl-, N,N-dimethylamino-acetamidinesulfonyl-, and 1-methanesulfonyl-methylamino)-ethyl-.

In still other embodiments of Formulae (I) and (Ia) where Ar is




R^8 may be selected from:




wherein R^{30} and R^{31} each independently is selected from hydrogen, methyl, ethyl, methoxyethyl and hydroxyethyl, and R^{32} is hydrogen, methyl or ethyl.

In yet other embodiments of Formulae (I) and (Ia) where Ar is




R^8 may be selected from:




wherein

- 5 R^{32} is hydrogen or alkyl;
- R^{33} is hydrogen, alkyl, hydroxy or alkoxy;
- R^{34} is hydroxy or alkoxy; and
- E is (C=O) or CH₂.


In certain embodiments of Formulae (I) and (Ia), Y and Ar considered together may be
 10 selected from one of (A), (B), (C), or (D):



(A)




(B)



(C)

and



(D)

wherein

X is N or CH (wherein when X is CH, said hydrogen atom of X is optionally replaced by a substituent R^8);

5 R^1 , R^2 and R^3 are individually independently selected from hydrogen, lower alkyl, and hydroxy, or R^2 and R^3 together form a bond to define an ethylene group;

R^7 is selected from

a) alkyl, halogen, and cyano;

10 b) alkyl substituted with one to two of halogen, hydroxy, cyano, $-NR^{12a}R^{14a}$, $-NR^{12a}C(=O)R^{14a}$, azetidiny, morpholinyl, pyrrolidinyl, piperidinyl, piperazinyl, imidazoliny, said rings in turn being optionally substituted with one to two R^{25} ;

c) $-SO_2R^{11}$, $-NR^{12}SO_2R^{11}$, $-NR^{12}CO_2R^{11a}$, $-NR^{12}C(=O)R^{11a}$, $-NR^{12}R^{14}$, $-C(=O)NR^{12}R^{14}$, $-SO_2NR^{12}R^{14}$, $-NR^{12}C(=NR^{13})R^{14}$, $-N=C(R^{15})R^{16}$, $-SO_2N=C(R^{15})R^{16}$, $-C(=O)R^{17}$, $-CO_2R^{17}$, and $-OR^{17}$;

15 d) heterocyclo or heteroaryl selected from pyrrolidin-2-yl, pyrrolidin-3-yl, imidazolin-2-yl, imidazol-2-yl, piperidinyl, and tetrazolyl, said heterocyclic and heteroaryl rings in turn being optionally substituted as valance permits with one to three of R^{25} ;

20 R^8 is selected from alkyl, halogen, cyano, alkoxy, amino, alkylamino, alkylsulfonyl, and piperidinyl, each of said R^8 groups in turn being optionally substituted with one to two of lower alkyl, lower alkoxy, cyano, and/or halogen;

R^{11} is alkyl, alkylamino, aminoalkyl, or hydroxy(C_{1-4})alkyl;

R^{11a} is alkyl, alkylamino, aminoalkyl, $-C(=O)$ alkyl, $N(O)$ alkyl, or hydroxy(C_{1-4})alkyl;

25 R^{12a} and R^{14a} are independently selected from hydrogen, alkyl, hydroxyalkyl, lower alkoxyalkyl, haloalkyl, and lower alkylamino(alkyl);

R^{12} and R^{13} are selected from hydrogen, alkyl, and hydroxy(C_{1-4})alkyl;


- R^{14} is selected from hydrogen, alkyl, hydroxyalkyl, alkoxyalkyl, aminoalkyl, alkylamino-alkyl, alkylamidylalkyl, cyanoalkyl, pyrrolidinyl, pyrrolinyl, imidazoliny, pyrimidinyl, tetrahydrofuranyl, furanylalkyl, isothiazolidinyl, oxazolidinyl, tetrahydrofuranylalkyl, morpholinylalkyl, and imidazolylalkyl, said R^{14} cyclic groups
 5 in turn being optionally substituted with one, two or three groups where valence allows selected from R^{25} ;
 or alternatively, R^{12} and R^{14} when attached to the same nitrogen atom (as in $-NR^{12}R^{14}$, $-C(=O)NR^{12}R^{14}$, or $-SO_2NR^{12}R^{14}$), may be taken together to form pyrrolidinyl, piperidinyl, imidazoliny, imidazolidinyl, pyrazolyl, or imidazolyl, said rings
 10 optionally substituted where valence allows with one, two or three groups selected from R^{25} ;
 R^{15} is selected from hydrogen, alkyl, alkoxyalkyl, and C_{3-6} cycloalkyl, and
 R^{16} is amino or alkylamino, or alternatively, R^{15} and R^{16} are taken together to form a pyrrolidinyl or imidazolidinyl ring optionally substituted with one, two or three
 15 groups selected from R^{25} ;
 R^{17} is selected from hydrogen, alkyl, methylsulfinylalkyl, methylsulfonylalkyl, carboxy-alkyl, alkoxycarbonylalkyl, hydroxyalkyl, alkoxyalkyl, aminoalkyl, alkylaminoalkyl, furanyl(C_{1-4})alkyl, tetrahydrofuranyl(C_{1-4})alkyl, morpholinyl(C_{1-4})alkyl, and pyrimidinyl, each R^{17} in turn being optionally substituted where valence allows with one to
 20 two groups selected from R^{25} ;
 R^{25} at each occurrence is selected independently of each other R^{25} from lower alkyl, halogen, cyano, alkyl, hydroxy(C_{1-4})alkyl, (C_{1-4})alkoxy(C_{1-4})alkyl, halo(C_{1-4})alkyl, cyano(C_{1-4})alkyl, amino(C_{1-4})alkyl, (C_{1-4})alkylamino(C_{1-4})alkyl, halogen, haloalkoxy, trifluoromethyl, cyano, nitro, $-OR^{26}$, $-SR^{26}$, $-S(O)R^{26}$, $-S(O)_2R^{28}$,
 25 $-C(=O)R^{26}$, $-C(=O)NR^{26}R^{27}$, $-C(O)_2R^{26}$, $-C(O)_2NR^{26}R^{27}$, $-S(O)_2NR^{26}R^{27}$, $-NR^{26}R^{27}$, and/or $-NR^{26}(C=O)R^{27}$, wherein R^{26} and R^{27} are selected from hydrogen and C_{1-4} alkyl, and R^{28} is C_{1-4} alkyl; and
 m is 0, 1, or 2.

In certain preferred embodiments, Y and Ar considered together may be selected from
 30 one of (A) and (B);

- R^5 is selected from methyl, methoxy, cyano and fluoro; and
 R^7 is selected from one of pyrrolidinyl, pyrrolinyl, imidazoliny, $-NHC(=NH)R^{14}$, $-N=C(H)(R^{16})$, $-N=C(CH_3)(R^{16})$, $-OR^{17}$, and C_{1-2} alkyl optionally substituted with $-NR^{12a}R^{14a}$, wherein said pyrrolidinyl, pyrrolinyl and imidazoliny groups optionally
 35 may be substituted with one or two of lower alkyl, hydroxy, lower alkoxy, halogen,

- cyano, amino, (C₁₋₄)alkylamino, hydroxy(C₁₋₄)alkyl, amino(C₁₋₄)alkyl, and/or (C₁₋₄)alkylamino(C₁₋₄)alkyl;
- R¹⁰ is methyl;
- R¹⁴ is C₃₋₆cycloalkyl or -C₁₋₂alkylene-(C₃₋₆cycloalkyl);
- 5 R¹⁶ is -NH₂, -NH(C₁₋₂alkyl), or N(C₁₋₂alkyl)₂;
- R¹⁷ is -C₁₋₂alkylene-NH₂, -C₁₋₂alkylene-NH(lower alkyl), or -C₁₋₂alkylene-N(lower alkyl)₂; and
- R^{12a} and R^{14a} are selected from hydrogen, lower alkyl, hydroxy(C₁₋₄alkyl), and -C₁₋₄alkylene-O(lower alkyl); and
- 10 m is 0 or 1.

In certain embodiments of Formulae (I) and (Ia), Y and Ar considered together may be



wherein

- R¹ is hydrogen, hydroxy, or lower alkyl; and
- 15 R⁷ is selected from one pyrrolidinyl, pyrrolinyl, imidazoliny, -NR¹²R¹⁴, -NR¹²C(=NR¹³)R¹⁴, -N=C(R¹⁵)(R¹⁶), -OR¹⁷, and C₁₋₄alkyl optionally substituted with pyrrolidinyl, pyrrolinyl, imidazoliny, -NR^{12a}R^{14a}, and/or -OR^{17a}, wherein said pyrrolidinyl, pyrrolinyl and imidazoliny groups in turn optionally may be substituted with one to three groups selected from R²⁵;
- 20 R^{12a}, R^{14a}, R¹⁷, and R^{17a} are independently selected from hydrogen, lower alkyl, hydroxy-(C₁₋₄alkyl), -C₁₋₄alkylene-O(lower alkyl), -C₁₋₄alkylene-NH₂, -C₁₋₄alkylene-NH(lower alkyl), -C₁₋₄alkylene-N(lower alkyl)₂, and -C₁₋₄alkylene-(C₃₋₆cycloalkyl);
- R¹² is hydrogen, alkyl, or hydroxy(C₁₋₄)alkyl;
- R¹⁴ is selected from hydrogen, alkyl, hydroxyalkyl, alkoxyalkyl, aminoalkyl, alkylamino-
- 25 alkyl, alkylamidylalkyl, cyanoalkyl, pyrrolidinyl, pyrrolinyl, imidazoliny, pyrimidinyl, tetrahydrofuranyl, furanylalkyl, isothiazolidinyl, oxazolindinyl, tetrahydrofuranylalkyl, morpholinylalkyl, and imidazolylalkyl, said R¹⁴ cyclic groups in turn being optionally substituted with one, two or three groups where valence allows selected from R²⁵;
- 30 or alternatively, R¹² and R¹⁴ may be taken together to form pyrrolidinyl, piperidinyl, imidazoliny, imidazolidinyl, pyrazolyl, or imidazolyl, said rings optionally substituted where valence allows with one, two or three groups selected from R²⁵;

R^{15} is independently selected from hydrogen, lower alkyl, hydroxy(C_{1-4})alkyl, and C_{3-6} cycloalkyl;

R^{16} is independently amino or (C_{1-4})alkylamino;

or alternatively, R^{15} and R^{16} may be taken together to form a pyrrolidinyl or imidazolinyll ring, wherein said pyrrolidinyl, pyrrolinyl and imidazolinyll groups optionally may be substituted with one to three groups selected from R^{25} ; and

R^{25} at each occurrence is independently selected from lower alkyl, halogen, cyano, hydroxy(C_{1-4})alkyl, and (C_{1-4})alkoxy(C_{1-4})alkyl.


In certain embodiments of Formulae (I) and (Ia), Ar is 2,3-dihydro-1*H*-isoindol-5-yl optionally substituted at the 2-position with methyl or 2-methoxyethyl.

In certain embodiments of Formulae (I) and (Ia), Ar is 1*H*-indol-3-yl.

In certain embodiments of Formulae (I) and (Ia), Y is heterocyclylene such as piperazin-1,4-di-yl or piperidin-1-3-di-yl.

In certain embodiments of Formulae (I) and (Ia), Ar is pyridin-2-yl optionally substituted at the 4-position or 6-position with hydroxyalkoxy, optionally substituted pyrrolodinyll, alkylaminoalkyl, alkylsulfonylaminoalkyl, haloalkylaminoalkyl, alkoxyalkylaminoalkyl, hydroxyalkylaminoalkyl, aminoalkoxy, aminoalkyl, ureidylalkyl, carbamylalkyl, acetamidinyll, formamidinyll, optionally substituted imidazolinyll, optionally substituted pyrrolodinyllmethyl, optionally substituted imidazolinyllmethyl, alkoxyalkylaminocarbonyl, hydroxyalkylaminocarbonyl, aminoalkylaminocarbonyl, heterocyclylalkylaminocarbonyl, or heterocyclylalkylamino.

According to another aspect of the invention, preferred compounds are those compounds having the Formula (Ib)



and isomers or pharmaceutically-acceptable salts, hydrates, or prodrugs thereof, wherein

Y, R^1 , R^2 , R^3 , R, R' , R^5 , and R^6 are as defined above for compounds of Formula (Ia);

- X is N or CH (wherein when X is CH, said hydrogen atom of X is optionally replaced by a substituent R⁸);
- R⁷ is selected from alkyl, substituted alkyl, halogen, cyano, -SO₂R¹¹, -NR¹²SO₂R¹¹, -NR¹²CO₂R^{11a}, -NR¹²C(=O)R^{11a}, -NR¹²R¹⁴, -C(=O)NR¹²R¹⁴, -SO₂NR¹²R¹⁴, -NR¹²C(=NR¹³)R¹⁴, -N=C(R¹⁵)R¹⁶, -SO₂N=C(R¹⁵)R¹⁶, -C(=O)R¹⁷, -CO₂R¹⁷, -OR¹⁷, heterocyclo, heteroaryl, phenyl, and cycloalkyl, wherein said heterocyclo, heteroaryl, phenyl, and cycloalkyl groups are optionally substituted with up to three R²⁵;
- R⁸ is selected from alkyl, lower alkyl, halogen, cyano, halo(C₁₋₄)alkyl, hydroxy, lower alkoxy, amino, (C₁₋₄)alkylamino, (C₁₋₄)alkylamino(C₁₋₄)alkyl, hydroxy(C₁₋₄)alkyl, (C₁₋₄)alkoxy(C₁₋₄)alkyl, pyrrolidinyl, and piperidinyl (said pyrrolidinyl and piperidinyl in turn being optionally substituted with one to two of lower alkyl, lower alkoxy, cyano, and/or halogen);
- R¹⁰ is selected from hydrogen, C₁₋₄alkyl, hydroxy(C₁₋₄)alkyl, and carboxy(C₁₋₄)alkyl;
- R¹¹, R¹², R¹³ and R¹⁵ are independently selected from hydrogen, alkyl, hydroxyalkyl, alkoxyalkyl, and cycloalkyl, except R¹¹ is not hydrogen;
- R^{11a} is selected from alkyl, alkylamino, aminoalkyl, -C(=O)alkyl, N(O)alkyl, and hydroxyalkyl;
- R¹⁶ is independently amino or alkylamino; or alternatively, R¹⁵ and R¹⁶ may be taken together to form a heterocyclo ring optionally substituted with up to three groups selected from R²⁵;
- R¹⁴ and R¹⁷ are independently selected from hydrogen, alkyl, substituted alkyl, and heterocyclo or heteroaryl optionally substituted with up to three groups selected from R²⁵;
- or alternatively, R¹² and R¹⁴ when attached to the same nitrogen atom (as in -NR¹²R¹⁴, -C(=O)NR¹²R¹⁴, and -SO₂NR¹²R¹⁴), may be taken together to form a heterocyclo or heteroaryl ring optionally substituted with up to three groups selected from R²⁵;
- R²⁵ is at each occurrence independently selected as valence permits from alkyl, substituted alkyl, halogen, haloalkoxy, trifluoromethyl, cyano, nitro, -OR²⁶, -SR²⁶, -S(O)R²⁶, -S(O)₂R²⁶, -C(=O)R²⁶, -C(=O)NR²⁶R²⁷, -C(O)₂R²⁶, -C(O)₂NR²⁶R²⁷, -S(O)₂NR²⁶R²⁷, -NR²⁶R²⁷, and/or -NR²⁶(C=O)R²⁷, wherein R²⁶ and R²⁷ are selected from hydrogen, C₁₋₆alkyl, aryl, heteroaryl, cycloalkyl and heterocyclo, and R²⁸ is selected from C₁₋₆alkyl, aryl, heteroaryl, cycloalkyl and heterocyclo; and
- m is 0, 1, or 2.

Even more preferred are compounds of Formula (Ib), as immediately defined above, in which X is CH, m is 0, and R⁷ is selected from pyrrolidinyl, pyrrolinyl, imidazoliny,

- NR¹²R¹⁴, -NR¹²C(=NR¹³)R¹⁴, -N=C(R¹⁵)(R¹⁶), -OR¹⁷, and C₁₋₄alkyl optionally substituted with pyrrolidinyl, pyrrolinyl, imidazoliny, -NR^{12a}R^{14a}, and/or -OR^{17a}, wherein said pyrrolidinyl, pyrrolinyl and imidazoliny groups in turn optionally may be substituted with one to three of lower alkyl, halogen, cyano, and/or hydroxy(C₁₋₄)alkyl;
- 5 R¹², R^{12a}, R¹³, R¹⁴, R^{14a}R¹⁷, and R^{17a} are independently selected from hydrogen, lower alkyl, hydroxy(C₁₋₄alkyl), -C₁₋₄alkylene-O(lower alkyl), -C₁₋₄alkylene-NH₂, -C₁₋₄alkylene-NH(lower alkyl), -C₁₋₄alkylene-N(lower alkyl)₂, and -C₁₋₄alkylene-(C₃₋₆cycloalkyl); R¹⁵ is independently selected from hydrogen, lower alkyl, hydroxy(C₁₋₄)alkyl, and C₃₋₆cycloalkyl; R¹⁶ is independently amino or (C₁₋₄)alkylamino; or alternatively, R¹⁵ and R¹⁶
- 10 may be taken together to form a pyrrolidinyl, pyrrolinyl, or imidazoliny ring, wherein said pyrrolidinyl, pyrrolinyl and imidazoliny groups optionally may be substituted with one to three of lower alkyl, hydroxy, lower alkoxy, halogen, cyano, amino, (C₁₋₄)alkylamino, hydroxy(C₁₋₄)alkyl, amino(C₁₋₄)alkyl, and/or (C₁₋₄)alkylamino(C₁₋₄)alkyl.


In certain embodiments of Formula (Ib)

- 15 Y is methylene;
R and R' are methyl;
R⁵ is methoxy or methyl;
one of R⁷ and R⁸ is hydrogen, or methoxy, and the other is selected from methyl-(3,3,3-trifluoro-propyl)-amino-methyl-, 2-hydroxy-3-methoxy-propoxy-, methoxy-, 3-
- 20 hydroxy-2-hydroxymethyl-propoxy-, 1-(dimethylamino-carbonyl-methylamino)-ethyl-, 1-[(methoxycarbonyl)-methylamino]-ethyl-, 3-hydroxy-2-hydroxymethyl-propoxy-, 2,3-dihydroxypropoxy-, (S)-4-methoxy-2-oxo-pyrrolidin-1-yl-, 2-methoxyethoxy-carbonylaminomethyl-, methanesulfonyl-N-methylamino-methyl-, 4,5-dihydro-oxazol-2-yl-, 2-oxo-pyrrolidin-1-yl-, 2-[(2-hydroxy-ethyl)-methyl-amino]-ethoxy-, 3-hydroxy-
- 25 2-oxo-pyrrolidin-1-yl-, 2-methylamino-ethyl-, (S)-4-hydroxy-2-oxo-pyrrolidin-1-yl-, 2-(N-ethyl-N-methyl-amino)-ethoxy-, 1-(methanesulfonyl-N-methylamino)-ethyl-, 2-(N-ethyl-N-methylamino)-ethyl-, 2-methylamino-ethoxy-, ethyl-(2-hydroxyethyl)-amino-methyl-, 2-hydroxy-ethoxy-, ethyl-(2-methoxyethyl)-aminomethyl-, 2-hydroxy-ethyl-methylamino-methyl-, 3-methoxy-pyrrolidin-1-ylmethyl-, N-ethyl-N-methylamino-
- 30 methyl-, 2-methoxyethylaminomethyl-, 4-hydroxy-piperidin-1-ylmethyl-, hydroxy-pyrrolidin-1-ylmethyl-, aminomethyl-, ethylamino-methyl-, 3-dimethylamino-propoxy-, pyrrolidin-1-ylmethyl-, azetidin-1-ylmethyl-, 1,3-dimethyl-imidazolidin-2-ylideneamino)-, 1-(methylamino-methyl-carbonyl-methylamino)-ethyl-, N,N-dimethylacetamidinyl-, 1-[(2-methoxyethyl)-methylamino]-ethyl-, methylamino-
- 35 methyl-, dimethylamino-methyl-, 3,4-dimethylimidazoline-2,4-dione-1-yl-, 3-methylimidazoline-2,4-dione-1-yl-, 1-dimethylamino-ethyl-, 1-methylamino-ethyl-,

cyclopropylamino-methyl-, isopropylamino-methyl-, N-methyl-N-propylamino-methyl-, methylamino-methyl-, N-ethyl-N-methyl-amino-methyl-, 3-methyl-2-oxo-pyrrolidin-1-yl-, 2-oxo-imidazolidin-1-yl-, 2-methoxyethyl-amino-methyl-, 1-methyl-piperidin-4-yl-, pyrrolidin-1-yl-, 4-methyl-piperazin-1-yl-, 4-methyl-piperazin-1-yl-, *N,N*-dimethyl-
 5 formamidinyl-, methyl-, 4,5-dihydro-3*H*-pyrrol-2-ylamino-, acetamidinyl-, 4,5-Dihydro-1*H*-imidazol-2-ylamino-, 4,4-dimethyl-4,5-dihydro-imidazol-1-yl-, pyrimidin-2-ylamino-, (1*H*-Imidazol-2-ylmethyl)-amino-, amino-, cyano-, bromo-, fluoro, methylamino-, methylaminomethyl-, aminomethyl-, cyanomethyl-, hydroxymethyl-, morpholin-4-ylmethyl-, ethylaminomethyl-, (2,2,2-trifluoro-
 10 ethylamino)-methyl-, [(2-hydroxy-ethyl)-methylamino]-methyl-, [(2-methoxy-ethyl)-methylamino]-methyl-, [ethyl-(2-methoxy-ethyl)-amino]-methyl-, methoxymethylcarbonyl-N-methylaminomethyl-, 1-methyl-4,5-dihydro-1*H*-imidazol-2-ylmethyl-, 1-isopropyl-4,5-dihydro-1*H*-imidazol-2-ylmethyl-, 1-methyl-piperidin-4-ylmethyl-, methanesulfonyl-, dimethylaminosulfonyl-, 2-methoxyethylaminosulfonyl-,
 15 3-methoxypropylaminosulfonyl-, carboxy-, methyamino-carbonyl-, 2-dimethylamino-ethyl)-aminocarbonyl-, 2-dimethylaminoethyl-methylaminocarbonyl-, 3-dimethylaminopropyl-ethylaminocarbonyl-, 2-hydroxyethylaminocarbonyl-, 2-hydroxypropylaminocarbonyl-, 2-hydroxyethyl-methylaminocarbonyl-, 2-methoxyethylaminocarbonyl-, 3-methoxypropylaminocarbonyl-, cyanomethyl-
 20 aminocarbonyl-, tetrahydrofuran-2-ylmethylaminocarbonyl-, furan-2-ylmethylamino-carbonyl-, (2-morpholin-4-yl)-ethylaminocarbonyl-, (4-hydroxymethyl)-piperidin-1-yl-aminocarbonyl-, 2-hydroxyethoxy-, methanesulfinylmethoxy-, pyrimidin-2-yloxy-, hydroxy-, methanesulfonylmethoxy-, carboxymethoxy-, 2-methoxy-ethoxy-, 3-methoxy-propyloxy-, 2-dimethylamino-ethoxy-, 2-ethylamino-ethoxy-, methanesulfonylamino-,
 25 dimethylaminosulfonylamino-, dimethylaminosulfonyl-N-methylamino-, methylcarbonylamino-, chloro-, dimethylaminocarbonylamino-, methoxycarbonylamino-, methoxycarbonyl-carbonyl-methylamino-, methoxyaminocarbonylamino-, 2-hydroxyethyl-amino-, 3-hydroxypropyl-methylamino-, 3-hydroxypropyl-amino-, 2-methoxyethyl)-amino-, 2-methoxyethyl)-
 30 methylamino-, 3-hydroxypropyl)-methylamino-, (3-methoxypropyl)-amino-, 3-methoxypropyl)-methylamino-, bis-(2-hydroxyethyl)-amino-, (2-methylcarbonylamino)-ethylamino-, acetamidinyl-, *N,N'*-dimethylacetamidinyl-, *N*-methylacetamidinyl-, *N,N*-dimethylaminoformamidyl-, *N,N*-dimethylaminoisobutyramidinyl-, *N,N*-dimethylamino-(3-methoxy)-propionamidyl-,
 35 *N,N*-dimethylcyclobutane-carboxamidinyl-, imidazolidin-2-ylideneamino-, 1-methyl-pyrrolidin-(2*Z*)-ylideneamino-, 2-oxo-tetrahydro-furan-3-ylamino-, (4,5-Dihydro-3*H*-pyrrol-2-yl)-methyl-amino-, 2-chloro-pyrimidin-4-ylamino-, 4-chloro-pyrimidin-2-

ylamino-, 2-chloro-5-methyl-pyrimidin-4-ylamino-, 2-oxo-pyrrolidin-1-yl-, 1-methyl-4,5-dihydro-1*H*-imidazol-2-yl-, 2-methyl-4,5-dihydro-imidazol-1-yl-, 4,4-dimethyl-4,5-dihydro-imidazol-1-yl-, 2,4,4-trimethyl-4,5-dihydro-imidazol-1-yl-, 4-methyl-imidazol-1-yl-, 4-methyl-imidazol-1-yl-, 2-methyl-imidazol-1-yl-, pyrazol-1-yl-, 3-methyl-
 5 pyrazol-1-yl-, 3,5-Dimethyl-pyrazol-1-yl-, pyrrolidin-2-yl-, pyrrolidin-3-yl-, 1-methyl-4,5-dihydro-1*H*-imidazol-2-yl-, 1-ethyl-4,5-dihydro-1*H*-imidazol-2-yl-, 1-isopropyl-4,5-dihydro-1*H*-imidazol-2-yl-, 1-isopropyl-4,5-dihydro-1*H*-imidazol-2-yl-, 1-(2-hydroxy-ethyl)-4,5-dihydro-1*H*-imidazol-2-yl-, 1-(2-methoxy-ethyl)-4,5-dihydro-1*H*-imidazol-2-yl-, 1*H*-Imidazol-2-yl-, 1-methyl-1*H*-imidazol-2-yl-, 1-(2-hydroxy-ethyl)-1*H*-imidazol-2-yl-,
 10 1,1-dioxo-1λ6-isothiazolidin-2-yl-, 2-oxo-oxazolidin-3-yl-, 3-methyl-2-oxo-pyrrolidin-1-yl-, ethyl-2-oxo-pyrrolidin-1-yl-, pyrrolidine-2,5-dione-1-yl-, 2-oxo-imidazolidin-1-yl-, 3-methyl-2-oxo-imidazolidin-1-yl-, 3-ethyl-2-oxo-imidazolidin-1-yl-, 3-(2-methoxy-ethyl)-2-oxo-imidazolidin-1-yl-, 1*H*-tetrazol-5-yl-, ethoxy-, 1-pyrrolidin-1-yl-ethyl-, 1-Azetidin-1-yl-ethyl-, aminosulfonyl-, N,N-dimethylamino-acetamidinesulfonyl-, and 1-methanesulfonyl-methylamino)-ethyl-; and
 15 R¹⁰ is hydrogen or methyl.

According to another aspect of the invention, preferred are those compounds having the Formula (Ic)



20 and isomers or pharmaceutically-acceptable salts, hydrates, or prodrugs thereof, wherein
 Y is $-(\text{CHR}^1)-$ or $-(\text{CHR}^2-\text{CHR}^3)-$;
 R¹, R² and R³ are independently selected from hydrogen, lower alkyl, and hydroxy;
 R⁵ is selected from C₁₋₄alkyl, halogen, cyano, hydroxy, C₁₋₄alkoxy, $-\text{O}(\text{CH}_2)_r\text{NH}_2$,
 $-\text{O}(\text{CH}_2)_r\text{NH}(\text{C}_{1-4}\text{alkyl})$, $-\text{O}(\text{CH}_2)_r\text{N}(\text{C}_{1-4}\text{alkyl})_2$, $-\text{O}(\text{CH}_2)_r\text{OH}$ and
 25 $-\text{O}(\text{CH}_2)_r\text{O}(\text{C}_{1-4}\text{alkyl})$;
 R⁷ is selected from pyrrolidinyl, pyrrolinyl, imidazolynyl, $-\text{NR}^{12}\text{R}^{14}$, $-\text{NR}^{12}\text{C}(=\text{NR}^{13})\text{R}^{14}$,
 $-\text{N}=\text{C}(\text{R}^{15})(\text{R}^{16})$, $-\text{OR}^{17}$, and C₁₋₄alkyl optionally substituted with pyrrolidinyl,
 pyrrolinyl, imidazolynyl, $-\text{NR}^{12a}\text{R}^{14a}$, and/or $-\text{OR}^{17a}$, wherein said pyrrolidinyl,
 pyrrolinyl and imidazolynyl groups optionally may be substituted with one to two of
 30 lower alkyl, hydroxy, lower alkoxy, halogen, cyano, amino, (C₁₋₄)alkylamino,
 hydroxy(C₁₋₄)alkyl, amino(C₁₋₄)alkyl, and/or (C₁₋₄)alkylamino(C₁₋₄)alkyl;

R¹⁰ is methyl;

R¹², R^{12a}, R¹³, R¹⁴, R^{14a}, R¹⁷, and R^{17a} are independently selected from hydrogen, lower alkyl, hydroxy(C₁₋₄alkyl), -C₁₋₄alkylene-O(lower alkyl), -C₁₋₄alkylene-NH₂, -C₁₋₄alkylene-NH(lower alkyl), -C₁₋₄alkylene-N(lower alkyl)₂, C₃₋₆cycloalkyl and -C₁₋₄alkylene-(C₃₋₆cycloalkyl);


R¹⁵ is hydrogen, lower alkyl, hydroxyC₁₋₄alkyl or C₃₋₆cycloalkyl;

R¹⁶ is independently amino or (C₁₋₄)alkylamino;

or alternatively, R¹⁵ and R¹⁶ may be taken together to form a pyrrolidinyl, pyrrolinyl or imidazoliny ring, wherein said pyrrolidinyl, pyrrolinyl and imidazoliny groups optionally may be substituted with one to three of lower alkyl, halogen, cyano, -C(=O)H, -C(=O) (C₁₋₄)alkyl, amino, (C₁₋₄)alkylamino, hydroxy(C₁₋₄)alkyl, amino(C₁₋₄)alkyl and (C₁₋₄)alkylamino(C₁₋₄)alkyl; and

r is 1, 2, 3, or 4.

Within the above group of compounds according to Formula (Ic), more preferred are compounds having the Formula (Id)



(Id)

wherein

R⁵ is selected from methyl, methoxy, cyano and fluoro; and

R⁷ is selected from one of pyrrolidinyl, pyrrolinyl, imidazoliny, -NHC(=NH)R¹⁴,

-N=C(H)(R¹⁶), -N=C(CH₃)(R¹⁶), -OR¹⁷, and C₁₋₂alkyl optionally substituted with -NR^{12a}R^{14a}, wherein said pyrrolidinyl, pyrrolinyl and imidazoliny groups optionally may be substituted with one to two of lower alkyl, hydroxy, lower alkoxy, halogen, cyano, amino, (C₁₋₄)alkylamino, hydroxy(C₁₋₄)alkyl, amino(C₁₋₄)alkyl, and/or (C₁₋₄)alkylamino(C₁₋₄)alkyl;


R¹⁴ is C₃₋₆cycloalkyl or -C₁₋₂alkylene-(C₃₋₆cycloalkyl);

R¹⁶ is -NH₂, -NH(C₁₋₂alkyl), or N(C₁₋₂alkyl)₂;

R¹⁷ is -C₁₋₂alkylene-NH₂, -C₁₋₂alkylene-NH(lower alkyl), or -C₁₋₂alkylene-N(lower alkyl)₂; and

R^{12a} and R^{14a} are selected from hydrogen, lower alkyl, hydroxy(C₁₋₄alkyl), and -C₁₋₄alkylene-O(lower alkyl).


Within the above group of preferred compounds, more preferred are those compounds as immediately defined above wherein R^7 is selected from one of





wherein R^{30} , R^{31} , and R^{32} are independently selected from hydrogen and methyl.


- 5 According to another aspect of the invention, combinations of the preferred groups described herein form other preferred embodiments. In this manner, a variety of preferred compounds are embodied within the present invention. For example, another group of preferred compounds, selected from a combination of preferred groups recited above, are those compounds having the formula (Ic), as immediately defined above,
- 10 wherein R^5 is fluoro, cyano, methyl or methoxy, and R^{10} is methyl. Thus, further combinations of preferred compounds may be selected from the preferred groups recited above.

In one embodiment the present invention provides a compound of formula I having a formula selected from






and




- Examples for compounds of formula Ih are the compounds of Table 1. Examples for
 5 compounds of formula Ii are the compounds of Table 2. Examples for compounds of
 formula Ij are the compounds of Tables 3, 4, 5 or 6. Examples for compounds of formula
 Im are the compounds of Table 7. Examples for compounds of formula In are the
 compounds of Table 8. Examples for compounds of formula Io are the compounds of
 Table 9. Examples for compounds of formula Ip are the compounds of Table 10.
- 10 In another aspect the present invention is directed to a process of making an intermediate
 quinazolinindione of formula (II)




wherein R, R' and R⁵ are as recited above, comprising:

- (a) dissolving in water with the addition of base, a nitro-benzoic compound of formula
 15 (8),



wherein R, R' and R⁵ are as recited above;


(b) adding to the aqueous, basic solution of compound (4), a noble metal catalyst on an inert support, and exposing the solution to a hydrogen environment to provide an amine compound of the formula (9),



(9)

;

- 5 (c) reacting the amine compound of formula (9), with a cyanate salt and an acid to provide a urea compound having the formula (10),



(10)

and,

- (d) cyclizing the urea of formula (10) with addition of base and heat to provide the
 10 quinazolidione of formula (II),
 wherein the process of making dione of quinazolidione of formula (II) from the nitrobenzoic compound of formula (8), is performed in a water-based solvent system.

In another embodiment the present invention provides a process as described above wherein in step (a), the base is selected from NaOH, KOH, and LiOH. In another
 15 embodiment the present invention provides a process as described above wherein in step (b), the heterogenous catalyst on an inert support comprises palladium on carbon, and the hydrogen environment is exposure to hydrogen gas. In still another embodiment the present invention provides a process as described above wherein in step (b), the heterogenous catalyst on an inert support comprises palladium on carbon, and the
 20 hydrogen environment is transfer hydrogenation using a hydrogen source selected from hydrazine and a formate salt. In still another embodiment the present invention provides a process as described above wherein in step (b), the cyanate salt is selected from KOCN and NaOCN, and the acid is selected from HCl and HOAc.

Alpha-1 adrenoceptors mediate the contractile state of smooth muscle tissue and are pre-
 25 sent in the human prostate, bladder neck and urethra. Alpha-1 adrenoceptor stimulation also produces contraction of urethral and bladder neck smooth muscle, leading to in-

creased resistance in urinary outflow. Thus, alpha-1 adrenoceptor antagonists may be useful in treating disorders of the urinary tract, as previously defined.

Alpha-1B adrenoceptors are present in the liver, heart and cerebral cortex and are believed to be involved in mediating vascular contractile and blood pressure responses.

5 Alpha-1B adrenoceptors are also present in areas of the spinal cord which receive input from sympathetic neurons originating in the pontine micturition center and are presumed to be involved in the regulation of bladder function. Additionally, alpha-1B adrenoceptor antagonists are useful as analgesic/antihyperalgesic therapies for treating pain, including symptoms of acute pain, inflammatory pain, neuropathic pain (including
10 thermal and mechanical hyperalgesia as well as thermal and mechanical allodynia), complex regional pain syndromes (including reflex sympathetic dystrophy, causalgia and sympathetically maintained pain and the like).

However, it must be noted that in BPH, it is often the irritative symptoms which prompt the patient to seek treatment, and that these irritative symptoms may be present even in
15 patients with no demonstrable obstruction (*i.e.* normal urine flow rates). By combining both alpha-1A and alpha-1B subtype selectivity in a compound, a reduction of both obstructive and irritative symptoms in patients with BPH may be achieved. Lower levels or lack of alpha-1D adrenoceptor antagonism should lead to reduced or fewer side effects than those associated with the use of non-subtype-selective agents.

20 In a preferred embodiment, the compounds of this invention are useful for treating disorders and symptoms which can be ameliorated by blockade of alpha1A/B adrenoceptors, such as reduction or alleviation of urinary tract disorders, e.g., pelvic hypersensitivity (including interstitial cystitis, prostatitis, pelvic pain syndrome, infectious cystitis, prostatodynia, and the like), overactive bladder, urinary frequency,
25 nocturia, urinary urgency, detrusor hyperreflexia, outlet obstruction, BPH, prostatitis, urge incontinence, urethritis, idiopathic bladder hypersensitivity, sexual dysfunction, and the like.

In another preferred embodiment, the compounds of this invention are useful for treating disorders and symptoms which can be ameliorated by blockade of alpha-1A/B
30 adrenoceptors, such as reduction or alleviation of pain disorders, e.g., inflammatory pain, neuropathic pain, cancer pain, acute pain, chronic pain or complex regional pain syndromes.

In a more preferred embodiment, the compounds of this invention are useful for treating disorders and symptoms which can be ameliorated by blockade of both alpha-1A and alpha-1B adrenoceptors with diminished blockade of alpha-1D adrenoceptors, such as reduction or alleviation of both outlet obstruction, such as benign prostatic hypertrophy, and irritative symptoms associated with it, such as pain.

In another preferred embodiment, the compounds of this invention are useful for the improvement of sexual dysfunction including male erectile dysfunction (MED) and female sexual dysfunction (FSD).

These and other therapeutic uses are described, for example, in Goodman & Gilman's, The Pharmacological Basis of Therapeutics, ninth edition, McGraw-Hill, New York, 1996, Chapter 26, 601-616; and Coleman, Pharmacological Reviews 46:205-229 (1994).

The present invention includes pharmaceutical compositions comprising at least one compound of the present invention, or an individual isomer, racemic or non-racemic mixture of isomers or a pharmaceutically acceptable salt or solvate thereof, together with at least one pharmaceutically acceptable carrier, and optionally other therapeutic and/or prophylactic ingredients.

In general, the compounds of the present invention will be administered in a therapeutically effective amount by any of the accepted modes of administration for agents that serve similar utilities. Suitable dosage ranges are typically 1-500 mg daily, preferably 1-100 mg daily, and most preferably 1-30 mg daily, depending upon numerous factors such as the severity of the disease to be treated, the age and relative health of the subject, the potency of the compound used, the route and form of administration, the indication towards which the administration is directed, and the preferences and experience of the medical practitioner involved. One of ordinary skill in the art of treating such diseases will be able, without undue experimentation and in reliance upon personal knowledge and the disclosure of this Application, to ascertain a therapeutically effective amount of the compounds of the present invention for a given disease.

In general, compounds of the present invention will be administered as pharmaceutical formulations including those suitable for oral (including buccal and sublingual), rectal, nasal, topical, vaginal, or parenteral (including intramuscular, intraarterial, intrathecal, subcutaneous and intravenous) administration or pulmonary in a form suitable for administration by inhalation or insufflation. The preferred manner of administration is

generally oral using a convenient daily dosage regimen which can be adjusted according to the degree of affliction.

A compound or compounds of the present invention, together with one or more conventional adjuvants, carriers, or diluents, may be placed into the form of pharmaceutical compositions and unit dosages. The pharmaceutical compositions and unit dosage forms may be comprised of conventional ingredients in conventional proportions, with or without additional active compounds or principles, and the unit dosage forms may contain any suitable effective amount of the active ingredient commensurate with the intended daily dosage range to be employed. The pharmaceutical compositions may be employed as solids, such as tablets or filled capsules, semisolids, powders, sustained release formulations, or liquids such as solutions, suspensions, emulsions, elixirs, or filled capsules for oral use; or in the form of suppositories for rectal or vaginal administration; or in the form of sterile injectable solutions for parenteral use. Formulations containing about one (1) to about 20 milligram of active ingredient or, more broadly, about 0.01 to about one hundred (100) milligrams, per tablet, are accordingly suitable representative unit dosage forms.

The compounds of the present invention may be formulated in a wide variety of oral administration dosage forms. The pharmaceutical compositions and dosage forms may comprise a compound or compounds of the present invention or pharmaceutically acceptable salts thereof as the active component. The pharmaceutically acceptable carriers may be either solid or liquid. Solid form preparations include powders, tablets, pills, capsules, cachets, suppositories, and dispersible granules. A solid carrier may be one or more substances which may also act as diluents, flavoring agents, solubilizers, lubricants, suspending agents, binders, preservatives, tablet disintegrating agents, or an encapsulating material. In powders, the carrier generally is a finely divided solid which is a mixture with the finely divided active component. In tablets, the active component generally is mixed with the carrier having the necessary binding capacity in suitable proportions and compacted in the shape and size desired. The powders and tablets preferably contain from about one (1) to about seventy (70) percent of the active compound. Suitable carriers include but are not limited to magnesium carbonate, magnesium stearate, talc, sugar, lactose, pectin, dextrin, starch, gelatin, tragacanth, methylcellulose, sodium carboxymethylcellulose, a low melting wax, cocoa butter, and the like. The term "preparation" is intended to include the formulation of the active compound with encapsulating material as carrier, providing a capsule in which the active component, with or without carriers, is surrounded by a carrier, which is in association

with it. Similarly, cachets and lozenges are included. Tablets, powders, capsules, pills, cachets, and lozenges may be as solid forms suitable for oral administration.

Other forms suitable for oral administration include liquid form preparations including emulsions, syrups, elixirs, aqueous solutions, aqueous suspensions, or solid form
5 preparations which are intended to be converted shortly before use to liquid form preparations. Emulsions may be prepared in solutions, e.g., in aqueous propylene glycol solutions or may contain emulsifying agents, e.g., such as lecithin, sorbitan monooleate, or acacia. Aqueous solutions can be prepared by dissolving the active component in water and adding suitable colorants, flavors, stabilizing, and thickening agents. Aqueous
10 suspensions can be prepared by dispersing the finely divided active component in water with viscous material, such as natural or synthetic gums, resins, methylcellulose, sodium carboxymethylcellulose, and other well known suspending agents. Solid form preparations include solutions, suspensions, and emulsions, and may contain, in addition to the active component, colorants, flavors, stabilizers, buffers, artificial and natural
15 sweeteners, dispersants, thickeners, solubilizing agents, and the like.

The compounds of the present invention may be formulated for parenteral administration (e.g., by injection, e.g. bolus injection or continuous infusion) and may be presented in unit dose form in ampoules, pre-filled syringes, small volume infusion or in
20 multi-dose containers with an added preservative. The compositions may take such forms as suspensions, solutions, or emulsions in oily or aqueous vehicles, e.g. solutions in aqueous polyethylene glycol. Examples of oily or nonaqueous carriers, diluents, solvents or vehicles include propylene glycol, polyethylene glycol, vegetable oils (e.g., olive oil), and injectable organic esters (e.g., ethyl oleate), and may contain formulatory agents such as preserving, wetting, emulsifying or suspending, stabilizing and/or dispersing agents.
25 Alternatively, the active ingredient may be in powder form, obtained by aseptic isolation of sterile solid or by lyophilisation from solution for constitution before use with a suitable vehicle, e.g., sterile, pyrogen-free water.

The compounds of the present invention may be formulated for topical administration to the epidermis as ointments, creams or lotions, or as a transdermal patch. Ointments
30 and creams may, e.g., be formulated with an aqueous or oily base with the addition of suitable thickening and/or gelling agents. Lotions may be formulated with an aqueous or oily base and will in general also contain one or more emulsifying agents, stabilizing agents, dispersing agents, suspending agents, thickening agents, or coloring agents. Formulations suitable for topical administration in the mouth include lozenges
35 comprising active agents in a flavored base, usually sucrose and acacia or tragacanth;

pastilles comprising the active ingredient in an inert base such as gelatin and glycerin or sucrose and acacia; and mouthwashes comprising the active ingredient in a suitable liquid carrier.

The compounds of the present invention may be formulated for administration as
5 suppositories. A low melting wax, such as a mixture of fatty acid glycerides or cocoa butter is first melted and the active component is dispersed homogeneously, e.g., by stirring. The molten homogeneous mixture is then poured into convenient sized molds, allowed to cool, and to solidify.

The compounds of the present invention may be formulated for vaginal administration.
10 Pessaries, tampons, creams, gels, pastes, foams or sprays may contain in addition to the active ingredient, such carriers as are known in the art to be appropriate.

The compounds of the present invention may be formulated for nasal administration. The solutions or suspensions are applied directly to the nasal cavity by conventional means, e.g., with a dropper, pipette or spray. The formulations may be provided in a
15 single or multidose form. In the case of a dropper or pipette, dosing may be achieved by the patient administering an appropriate, predetermined volume of the solution or suspension. In the case of a spray, this may be achieved e.g. by means of a metering atomizing spray pump.

The compounds of the present invention may be formulated for aerosol administration,
20 particularly to the respiratory tract and including intranasal administration. The compound will generally have a small particle size e.g. on the order of five (5) microns or less. Such a particle size may be obtained by means known in the art, e.g. by micronization. The active ingredient is provided in a pressurized pack with a suitable propellant such as a chlorofluorocarbon (CFC), e.g., dichlorodifluoromethane, trichlorofluoromethane, or
25 dichlorotetrafluoroethane, or carbon dioxide or other suitable gas. The aerosol may conveniently also contain a surfactant such as lecithin. The dose of drug may be controlled by a metered valve. Alternatively the active ingredients may be provided in a form of a dry powder, e.g. a powder mix of the compound in a suitable powder base such as lactose, starch, and starch derivatives such as hydroxypropylmethyl cellulose, and
30 polyvinylpyrrolidone (PVP). The powder carrier will form a gel in the nasal cavity. The powder composition may be presented in unit dose form e.g. in capsules or cartridges of e.g., gelatin or blister packs from which the powder may be administered by means of an inhaler.

When desired, formulations can be prepared with enteric coatings adapted for sustained or controlled release administration of the active ingredient. For example, the compounds of the present invention can be formulated in transdermal or subcutaneous drug delivery devices. These delivery systems are advantageous when sustained release of the compound is necessary and when patient compliance with a treatment regimen is crucial. Compounds in transdermal delivery systems are frequently attached to a skin-adhesive solid support. The compound of interest can also be combined with a penetration enhancer, e.g., azone (1-dodecylazacycloheptan-2-one). Sustained release delivery systems are inserted subcutaneously into the subdermal layer by surgery or injection. The subdermal implants encapsulate the compound in a lipid soluble membrane, e.g., silicone rubber, or a biodegradable polymer, e.g., polylactic acid.

The pharmaceutical preparations are preferably in unit dosage forms. In such form, the preparation is subdivided into unit doses containing appropriate quantities of the active component. The unit dosage form can be a packaged preparation, the package containing discrete quantities of preparation, such as packeted tablets, capsules, and powders in vials or ampoules. Also, the unit dosage form can be a capsule, tablet, cachet, or lozenge itself, or it can be the appropriate number of any of these in packaged form.

Other suitable pharmaceutical carriers and their formulations are described in Remington: The Science and Practice of Pharmacy 1995, edited by Martin, Mack Publishing Company, 19th edition, Easton, Pennsylvania. Representative pharmaceutical formulations containing a compound of the present invention are described in Examples below.


Throughout the disclosure, and in the following Schemes and Examples herein, the following abbreviations are used for ease of reference: BOC: tert-butoxycarbonyl; BPH: Benign prostatic hypertrophy or benign prostatic hyperplasia; CBZ: carbobenzyloxy; CNS: Central nervous system; DCM: dichloromethane; DMF: *N,N*-Dimethylformamide; DMSO: dimethylsulfoxide; EtOH: ethanol; EtOAc: Ethyl Acetate; Hal: Halogen or halide; KOH: potassium hydroxide; L: Leaving group; MeOH: methanol; P: Protective group; Pd/C: palladium on carbon; THF: tetrahydrofuran; RT: room temperature

Compounds of the present invention may be made by the methods depicted in the illustrative synthetic reaction schemes shown and described below.

The starting materials and reagents used in preparing these compounds generally are either available from commercial suppliers, such as Aldrich Chemical Co., or are

- prepared by methods known to those skilled in the art following procedures set forth in references such as Fieser and Fieser's Reagents for Organic Synthesis; Wiley & Sons: New York, 1991, Volumes 1-15; Rodd's Chemistry of Carbon Compounds, Elsevier Science Publishers, 1989, Volumes 1-5 and Supplementals; and Organic Reactions, Wiley & Sons: New York, 1991, Volumes 1-40. The following synthetic reaction schemes are merely illustrative of some methods by which the compounds of the present invention may be synthesized, and various modifications to these synthetic reaction schemes may be made and will be suggested to one skilled in the art having referred to the disclosure contained in this Application.
- The starting materials and the intermediates of the synthetic reaction schemes may be isolated and purified if desired using conventional techniques including but not limited to filtration, distillation, crystallization, chromatography, and the like. Such materials may be characterized using conventional means, including physical constants and spectral data.
- Unless specified to the contrary, the reactions described herein preferably take place at atmospheric pressure over a temperature range from about -78°C to about 150°C , more preferably from about 0°C to reflux, and most preferably and conveniently at about room (or ambient) temperature, e.g., about 20°C .


Scheme 1



- Compounds of formula (Ia) can be prepared by coupling 2-chloro-quinazoline-4-one compounds (1) with amine compounds (2), in a solvent such as EtOH, with or without additives such as diisopropylethylamine. Quinazolin-4-ones can be prepared as described in Cronin et al., J. Med. Chem. 11:130-136 (1968) and WO 02/053558 A1, which are incorporated herein by reference. Alternatively, 2-chloro-quinazoline-4-one compounds (1) can be prepared as described below using the method described in Scheme 3.

Scheme 2

- 57 -



Compounds of formula (Ie), (If), and (Ig) can be prepared from 2-chloro-quinazolin-4-one compounds (1) using the process shown in Scheme 2.

Compounds of formula (Ie), (If), and (Ig) can be also prepared from 2-chloro-quinazoline compounds (1) using the process shown in Scheme 2.

2-Chloro-quinazolin-4-one compounds (1) can be converted to compounds of formula (Ie) by combining amines such as (3) (or a suitable salt of such amines, e.g. HCl salt thereof) with compounds (1) in a solvent, such as isopropanol, and heating the reaction mixture in the presence or absence of an additive like diisopropylethylamine.

- 10 Compounds (Ie) can be isolated by cooling and filtering the reaction mixture. In some cases, the recovery of product can be increased by adding an anti-solvent (e.g. water) prior to performing the filtration.


- 15 Compounds of formula (If) can be prepared by combining compounds (Ie), a heterogeneous catalyst on an inert support (e.g. palladium on carbon), an organic solvent (e.g. ethanol), water, and base (e.g. NaOH). Then, to this reaction mixture, hydrogen is introduced either directly or indirectly (e.g., via transfer hydrogenation using formate salts, hydrazine, etc.), and the nitro group is reduced to an amino group. The catalyst and support are removed, and the reaction mixture is neutralized by addition of acid (e.g. acetic, trifluoroacetic, hydrochloric acid, etc.). The precipitated compound of formula (If) is then collected via filtration.
- 20

Compounds of formula (Ig) can be prepared by combining compounds of formula (If) in an organic solvent (methylene chloride, acetonitrile, etc) and a solution of iminium salt (4) [prepared by activation of the appropriate acetamide or formamide with

phosphorous oxychloride or trifluoromethanesulfonic anhydride as described by Fabio et al, J. Med. Chem. 21:273-276 (1978); Campbell et al, Tetrahedron Letters 25:4813-4816 (1984), and Sforza et al, Tetrahedron Letters 39:711-714 (1998)]. The reaction is quenched by the addition of water, cooled, and then the pH is adjusted to ~8.5 with a
 5 base (e.g. NaOH, KOH, etc). The aqueous layer is extracted with an organic solvent (e.g. methylene chloride), then the organic solvent is converted to an alcohol (e.g. isopropanol), heated; cooled then filtered to provide compounds of formula (Ig).

Quinazolin-4-ones (1) can be prepared as described in Cronin et al., J. Med. Chem. 11:130-136 (1968) or WO 02/053558 A1 or using the method described in Scheme 3.

10 Scheme 3



Scheme 3 illustrates an alternate method for making 2-chloro-quinazolin-4-one compounds (1), used as starting material in Schemes 1 and 2.


Dione intermediate (5) is converted to dichloroquinazoline (6) by reacting (5) with
 15 phosphorous oxychloride in acetonitrile and heating the reaction mixture. The dichloroquinazoline (6) is isolated by quenching the reaction mixture into water and filtering the precipitated product, or by quenching the reaction mixture into a mixture of water and a water-immiscible solvent (e.g. methylene chloride), and extracting the product into the organic solvent. The solvent is evaporated to provide compound (6).

20 Compound (6) is then combined with a base (e.g., KOH, NaOH) in a mixture of water and a solvent like THF. At the end of the reaction, the organic solvent is partially removed by distillation, an acid (e.g. HOAc) is added, and the compound (1) is collected via filtration.

Dione intermediates (5) are commercially available or can be readily prepared by one
 25 skilled in the field, e.g., as described in Mizuno et al., Heteroatom Chemistry 11:428-433 (2000); Mizuno et al., Tetrahedron Letters 41:1051-1051 (2000); US 6,376,667; US 6,048,864; WO 97/23462; EP 775697-A1; and so forth.

Alternatively, dione intermediates (5) may be prepared as described in Scheme 4.

Scheme 4



Scheme 4 illustrates an alternative method for making of diones (5), used in Scheme 3 to prepare 2-chloro-quinazolin-4-one compounds (1).

- 5 Nitro-acids (8) are commercially available, or can be readily prepared by one skilled in the field from carboxylic acids (7) using several methods, including that of Kowalczyk et al., Organic Process Research and Development 1:355-358 (1997). Carboxylic acids (7) are commercially available.

Nitro acids (8) are dissolved in water by addition of base (e.g., NaOH, KOH, LiOH).

- 10 Typically, the amount of base used to dissolve the acids is in the range of a 1.3 to 1.5 molar equivalent, e.g., at a pH ≥ 12 . A heterogenous catalyst on an inert support is added (e.g., palladium on carbon), and the reaction mixture is exposed to a hydrogen atmosphere either directly (hydrogen gas) or indirectly (via transfer hydrogenation technique using, e.g., formate salts, hydrazine, etc., as the hydrogen source). The nitro-
15 group is thereby converted to an amino group to provide compounds (9).

- Compounds (9) can be converted to urea (10) by addition of a cyanate salt (e.g., KOCN, NaOCN) and an acid (e.g., HCl, HOAc). For this urea-forming step, the pH is preferably maintained in the range of 6.0 to 8.0, more preferably between 6.8 and 8.0. At pH greater than 8, the urea formation step may slow down, and when the pH is between 5.6 and 6.0,
20 an undesirable side reaction may occur. Degradation occurs at lower pH, e.g., at pH of ~ 3.9 . The urea (10) is then cyclized to a dione derivative (5) by adding a base (e.g., NaOH, KOH), wherein the pH is typically maintained at ≥ 12 , and heating the reaction mixture. The results of this cyclization step may be improved as the pH is increased. The dione (5)

- 60 -

is precipitated by adding an acid (e.g., HCl, HOAc) to the reaction mixture, to achieve a pH of about less than 8.2, more preferably in the range of 6.5 to 7.5, and the dione (5) may be isolated such as by filtration. Other acids also may be used, e.g., any acid that will generate HOCN in-situ from the cyanate salt and the acid.


- 5 The process of Scheme 4 is particularly advantageous in that the water-insoluble compounds (8)-(5) are converted into water-soluble compounds by conversion to their respective salts (by addition of base) and then the pH is adjusted to allow the desired reaction to occur in a water-based solvent system.

EXAMPLES


- 10 The following preparations and examples are provided to enable those skilled in the art to more clearly understand and to practice the present invention. However, these Examples should not be considered as limiting the scope of the invention, but merely as being illustrative and representative thereof.

Example A-1: 6,7-Dimethoxy-2-[methyl-(3-pyrrolidin-1-yl-benzyl)-amino]-1*H*-quin-
15 azolin-4-one

Step 1: Benzylamine (1) was prepared following the procedure of Sznaidman et al., Bioorg. Med. Chem. Lett. 6:565-568 (1996).




- Step 2: Benzylamine (1) (4.8 g, 27.3 mmol) in 100 mL DCM was reacted with 6.1 g (28
20 mmol) of di-*tert*-butyl dicarbonate for 2 h. The solvent was evaporated under reduced pressure and the remaining solid was dried at 60°C *in vacuo* to afford 7.7 g of the carbamate (2) (Mp = 106.8-107.5 °C).



- Step 3: The carbamate (2) from Step 2 (7.6 g, 27.5 mmol) was dissolved in 70 mL THF
25 and added over a period of 30 min. to a stirred slurry of 3 g (75 mmol) of LAH in 300 mL of diethyl ether. The mixture was stirred at reflux for 8 h and allowed to stand at ambient


temperature for 16 h. The reaction was quenched while stirring by sequential addition of 1- water; 2- 15% NaOH and 3-water until all solids appeared to be white. The solids were removed by filtration and washed once with THF. The filtrates were combined, concentrated under reduced pressure, and the remaining oil was distilled to yield 4.1 g (78%) of the N-methylbenzylamine (3). Bp 140 °C/1 mm.



Step 4: A sample of 2-chloro-6,7-dimethoxyquinazolin-4-one (300 mg, 1.25 mmol) in 4-5 mL of 1,2-dihydroxypropane was allowed to react with (400 mg, 2.1 mmol) of benzylamine (3) for 16 h at 110°C. The solvent was removed *in vacuo* and the remaining solid was chromatographed (EM Science silica gel 60; 2% MeOH in DCM) to yield 300 mg (60%) of the above titled-compound Example A-1. Mp 238-40°C; MS *m/z* 395 (M+H). The bis HCl salt was obtained when the free base in hot EtOH was treated with an excess of 10% HCl in EtOH, and was crystallized by addition of diethyl ether. Mp 220-223°C. Anal. (C₂₂H₂₆N₄O₃•2HCl), Calcd.: C, 56.54; H, 6.04; N, 11.99. Found: C, 56.20; H, 6.01; N, 11.95.

Example A-2: 2-[Benzyl-(1-benzyl-piperidin-4-yl)-amino]-6,7-dimethoxy-1H-quinazolin-4-one; compound with trifluoro-acetic acid

Step 1: Benzylamine (1) was prepared following the procedure of Harper et al., J. Medicinal Chemistry 7:729-32 (1964).



Step 2: The benzylamine (1) (14 mg) and 2-chloro-6,7-dimethoxyquinazolin-4-one (12 mg) were heated together in NMP at 80°C for 3 d. Purification by RP-HPLC provided the title compound: *m/z* 485 (M+H)⁺.

Examples A-3 to A-16

Following the same or similar method as described above for Example A-1, the compounds of formula Ia wherein R' and R are methyl and R⁵ is hydrogen, according to Table 1, were prepared, except the appropriately-substituted amine was used in place of *N*-methyl-(3-pyrrolidin-1-yl-benzyl)-amine.

5 **TABLE 1:**


Ex.	Compound Name	Mp°C	MS (M+H) ⁺
A-3	6,7-dimethoxy-2-[methyl-(6-methyl-pyridin-2-ylmethyl)-amino]-1H-quinazolin-4-one		341
A-4	2-[[2-(1H-indol-3-yl)-ethyl]-methyl-amino]-6,7-dimethoxy-1H-quinazolin-4-one		379
A-5	6,7-dimethoxy-2-{methyl-[3-(4-methyl-piperazin-1-yl)-benzyl]-amino}-1H-quinazolin-4-one		424
A-6	6,7-dimethoxy-5-methyl-2-[2-(4-methyl-piperazin-1-yl)-benzylamino]-1H-quinazolin-4-one		424
A-7	2-(benzyl-isopropyl-amino)-6,7-dimethoxy-1H-quinazolin-4-one	168-170.5	354
A-8	N'-(3-[[[(6,7-dimethoxy-4-oxo-1,4-dihydro-quinazolin-2-yl)-methyl-amino]-methyl]-phenyl]-N,N-dimethyl-formamidine, hydrochloride salt	242-245	395.4 M ⁺
A-9	2-[[3-(4,5-dihydro-3H-pyrrol-2-ylamino)-benzyl]-methyl-amino]-6,7-dimethoxy-1H-quinazolin-4-one	211-214	407.5 M ⁺
A-10	N-(3-[[[(6,7-dimethoxy-4-oxo-1,4-dihydro-quinazolin-2-yl)-methyl-amino]-methyl]-phenyl]-acetamidine, hydrogen bromide salt	263-266	381.4 M ⁺
A-11	2-[[3-(4,5-dihydro-1H-imidazol-2-ylamino)-benzyl]-methyl-amino]-6,7-dimethoxy-1H-quinazolin-4-one		409
A-12	2-[[3-(4,4-dimethyl-4,5-dihydro-imidazol-1-yl)-benzyl]-methyl-amino]-6,7-dimethoxy-1H-quinazolin-4-one	240-244	422
A-13	N'-(3-{2-[(6,7-dimethoxy-4-oxo-1,4-dihydro-quinazolin-2-yl)-methyl-amino]-ethyl}-phenyl)-N,N-dimethyl-acetamidine	135-138.1	423.5 M ⁺
A-14	6,7-dimethoxy-2-(methyl-{2-[3-(pyrimidin-2-yl-amino)-phenyl]-ethyl}-amino)-1H-quinazolin-4-one		433

A-15	2-[(2-{3-[(1H-imidazol-2-ylmethyl)-amino]-phenyl}-ethyl)-methyl-amino]-6,7-dimethoxy-1H-quinazolin-4-one	435
A-16	2-[[2-(3,4-Dimethoxy-phenyl)-ethyl]-methyl-amino]-6,7-dimethoxy-1H-quinazolin-4-one	400

Additional compounds prepared according to the procedure of Example A-1 using various aralkylamines with 2-chloro-6,7-dimethoxyquinazolin-4-one are shown in Table 11.


5 **Example B-1:** 2-[(3-Amino-benzyl)-methyl-amino]-6,7-dimethoxy-5-methyl-1H-quinazolin-4-one

Step 1: 2-Chloro-6,7-dimethoxy-5-methyl-quinazolin-4-one



A solution of 3 g (11 mmol) of the dichloroquinazoline (1A) in 50 mL THF was stirred
 10 with 30 mL of 40% aqueous KOH and 3 mL of 40% aqueous tetrabutylammonium hydroxide for 3 days. The layers were allowed to separate and the upper layer was concentrated to give a white solid. This was treated with dilute acetic acid and the solid was collected and washed with water, then dried at 65°C *in vacuo* to afford 2.4 g (86%) of the 2-chloroquinazolin-4-one (1). Mp 251-251.7 °C; MS (ES+) *m/z* 255 (M+H)⁺. Anal.
 15 (C₁₁H₁₁ClN₂O₃) Calcd.: C, 51.88; H, 4.35; N, 11.00. Found: C, 51.94; H, 4.34; N, 10.95.

Step 2:



- 64 -

A mixture consisting of the 2-chloroquinazolin-4-one (1) from Step 1 (2.95 g, 11.6 mmol), the above benzylamine (2) (1.65 g, 12.2 mmol), and *N,N*-diisopropylethylamine (1.5 g, 11.6 mmol) in 45 mL of EtOH was placed in a heavy-walled glass reaction vessel. A magnetic stirring bar was added and the reaction vessel was closed. The reaction mixture formed a slurry, and it was stirred while the reaction vessel was kept partially immersed in a 120°C bath for 2 h. The mixture was allowed to remain at ambient temperature without stirring for 16 h. The solid product was collected, washed twice with EtOH, and dried at 65°C *in vacuo* furnishing 3.7 g (90%) of Example B-1 as an off-white solid. Mp 268-269.5°C; Anal. (C₁₉H₂₂N₄O₃) Calcd.: C, 64.39; H, 6.26; N, 15.81. Found: C, 64.05; H, 6.25; N, 15.49.

Example B-2: 3-[[[(6,7-Dimethoxy-5-methyl-4-oxo-1,4-dihydro-quinazolin-2-yl)-methyl-amino]-methyl]-benzonitrile

Diisopropylethylamine (0.82 mL, 4.71 mmol) was added to a mixture of 2-chloro-6,7-dimethoxy-5-methylquinazolin-4-one (1.0 g, 3.92 mmol) and *N*-methyl-(3-cyanobenzyl)-amine (0.68 g, 4.71 mmol) in EtOH (50 mL). The mixture was heated at 120°C for 1.5 h in a sealed tube. After cooling, the resulting white solid was filtered and washed with MeOH to provide the aryl nitrile of Example B-2. Mp 279.9-281.5 °C; ¹H NMR (DMSO-d₆) δ 2.6 (s, 3H), 3.05 (s, 3H), 3.62 (s, 3H), 3.84 (s, 3H), 4.86 (s, 2H), 6.62 (s, 1H), 7.5-7.7.78 (m, 4H); ¹³C NMR (DMSO-d₆) δ 13.89, 35.58, 51.52, 55.93, 60.22, 104.79, 111.78, 119.15, 130.12, 131.19, 131.28, 131.96, 132.51, 140.04, 142.96, 150.51, 157.71. MW = 365.

Examples B-3 to B-10

Following the same or similar method as described for Examples B-1 and B-2 compounds of formula (Ic) wherein Y is -CH₂- according to Table 2 were prepared.

TABLE 2


Ex.	Compound Name	Mp°C	MS (M+H) ⁺
B-3	2-[benzyl-(2-dimethylamino-ethyl)-amino]-6,7-dimethoxy-5-methyl-1H-quinazolin-4-one		397
B-4	2-[(3-bromo-benzyl)-methyl-amino]-6,7-dimethoxy-5-methyl-1H-quinazolin-4-one		419
B-5	2-[(3-bromo-benzyl)-methyl-amino]-5,6,7-		435

	trimethoxy-1H-quinazolin-4-one	
B-6	2-[(2-fluoro-benzyl)-(2-hydroxy-ethyl)-amino]-6,7-dimethoxy-5-methyl-1H-quinazolin-4-one	388
B-7	3-{[methyl-(5,6,7-trimethoxy-4-oxo-1,4-dihydro-quinazolin-2-yl)-amino]-methyl}-benzonitrile	381
B-8	6,7-dimethoxy-5-methyl-2-[methyl-(3-methylamino-benzyl)-amino]-1H-quinazolin-4-one	368.4 M ⁺
B-9	2-[(3-amino-benzyl)-methyl-amino]-5,6,7-trimethoxy-1H-quinazolin-4-one	225-226.5 370.4 M ⁺
B-10	2-[(2-amino-benzyl)-methyl-amino]-5,6,7-trimethoxy-1H-quinazolin-4-one	370.4 M ⁺


Additional compounds prepared according to the procedure of Examples B-2 and B-3 are shown in Table 11.

Example C-1: 6,7-Dimethoxy-5-methyl-2-[methyl-(3-methylaminomethyl-benzyl)-amino]-1H-quinazolin-4-one

- 5 Step 1: 3-(Bromomethyl)benzoate (5.0 g, 21.83 mmol) and 40% methylamine in water (40.0 mL, 515.2 mmol) in EtOH (17mL) were heated at 80°C for 6 h. The solvent was concentrated to dryness. Purification by flash chromatography (CH₂Cl₂:MeOH: NH₄OH/300:10:1) gave 2.71 g (70%) of compound (1) as a clear oil.



- 10 Step 2: 1.0 M LiAlH₄ in THF (30.32 mL, 30.42 mmol) was added to compound (1) (2.71 g, 15.01 mmol) in THF (20 mL), and the mixture was heated at reflux for 16 h. After cooling to RT, Na₂SO₄·8H₂O (ca. 5 g) was added, and the heterogeneous solution was stirred at RT for 0.5 h. The white solid was filtered off and the filtrate was concentrated to dryness. Purification by flash chromatography (CH₂Cl₂: MeOH: NH₄OH/60:10:1)
- 15 gave 1.17 g (48%) of compound (2) as a clear oil.



Step 3: 2-Chloro-6,7-dimethoxy-5-methyl-1*H*-quinazolin-4-one (0.805 g, 3.16 mmol) and compound (2) (0.675 g, 4.109 mmol) in EtOH (10 mL) were heated in a sealed tube at 120°C for 3 h. The solvent was evaporated to dryness under reduced pressure.

Purification by flash chromatography (CH₂Cl₂: MeOH: NH₄OH/320:10:1) followed by
 5 recrystallization (CH₂Cl₂/ether) gave 0.468 g (39%) of the above Example C-1 as a white solid. mp 179.0-181.3°C; ¹H NMR (DMSO-*d*₆, 2.49) δ 1.74 (s, 3 H), 2.54 (t, 2 H), 2.80 (s, 3 H), 2.97 (s, 6 H), 3.62 (s, 3 H), 3.75 (t, 2H), 3.86 (2, 3H), 4.72 (s, 2H), 6.40 (d, 1H), 6.67 (s, 1H), 6.75 (d, 2H), 7.05 (t, 1H), 10.99 (s, 1H); IR (KBr) ν_{max} 1589 cm⁻¹; MS (ES+) *m/z* 383(M + H); Anal. (C₂₁H₂₆N₄O₃) C: calcd, 65.95; found, 65.75; H: calcd, 6.85;
 10 found, 6.76; N: calcd, 14.65; found, 14.66.

Examples C-2 to C-24

Compounds of formula Ic wherein Y is -CH₂- of Table 3 were prepared following the same or similar method as described for Example C-1.

TABLE 3

Ex.	Compound Name	Mp°C	MS (M+H)
C-2	2-[(3-Aminomethyl-benzyl)-methyl-amino]-6,7-dimethoxy-5-methyl-1 <i>H</i> -quinazolin-4-one		369
C-3	(3-{[Methyl-(5,6,7-trimethoxy-4-oxo-1,4-dihydroquinazolin-2-yl)-amino]-methyl}-phenyl)-acetonitrile		395
C-4	2-[(3-Hydroxymethyl-benzyl)-methyl-amino]-5,6,7-trimethoxy-1 <i>H</i> -quinazolin-4-one		386
C-5	5-Hydroxy-2-[(3-hydroxymethyl-benzyl)-methyl-amino]-6,7-dimethoxy-1 <i>H</i> -quinazolin-4-one		372
C-6	5,6,7-Trimethoxy-2-[methyl-(3-morpholin-4-ylmethyl-benzyl)-amino]-1 <i>H</i> -quinazolin-4-one		455
C-7	5,6,7-Trimethoxy-2-[methyl-(3-methylaminomethyl-benzyl)-amino]-1 <i>H</i> -quinazolin-4-one	124.9-127.6	398.5 M ⁺
C-8	2-[Ethyl-(3-methylaminomethyl-benzyl)-amino]-6,7-dimethoxy-5-methyl-1 <i>H</i> -quinazolin-4-one		396.5 M ⁺
C-9	2-[(3-Ethylaminomethyl-benzyl)-methyl-amino]-6,7-dimethoxy-5-methyl-1 <i>H</i> -quinazolin-4-one		396.5 M ⁺

C-10	2-({3-[(Ethyl-methyl-amino)-methyl]-benzyl}-methyl-amino)-5,6,7-trimethoxy-1H-quinazolin-4-one		426.5 M ⁺
C-11	6,7-Dimethoxy-5-methyl-2-(methyl-{3-[(2,2,2-trifluoro-ethylamino)-methyl]-benzyl}-amino)-1H-quinazolin-4-one		450.5 M ⁺
C-12	2-[(3-{[(2-Hydroxy-ethyl)-methyl-amino]-methyl}-benzyl)-methyl-amino]-5,6,7-trimethoxy-1H-quinazolin-4-one hydrochloride salt	169.0-171.9	456.5 M ⁺
C-13	6,7-dimethoxy-5-methyl-2-[(3-{[(2-methoxy-ethyl)-methyl-amino]-methyl}-benzyl)-methyl-amino]-1H-quinazolin-4-one		426.5 M ⁺
C-14	5,6,7-Trimethoxy-2-[(3-{[(2-methoxy-ethyl)-methyl-amino]-methyl}-benzyl)-methyl-amino]-1H-quinazolin-4-one hydrogen chloride salt	162.9-164.5	470.6 M ⁺
C-15	2-[(3-{[Ethyl-(2-methoxy-ethyl)-amino]-methyl}-benzyl)-methyl-amino]-5,6,7-trimethoxy-1H-quinazolin-4-one hydrogen chloride salt		470.6 M ⁺
C-16	N-(3-{[(6,7-Dimethoxy-5-methyl-4-oxo-1,4-dihydro-quinazolin-2-yl)-methyl-amino]-methyl}-benzyl)-2-methoxy-N-methyl-acetamide		454.5 M ⁺
C-17	2-Methoxy-N-methyl-N-(3-{[methyl-(5,6,7-trimethoxy-4-oxo-1,4-dihydro-quinazolin-2-yl)-amino]-methyl}-benzyl)-acetamide	115-117	470.5 M ⁺
C-18	2-[(3-{[Ethyl-(2-methoxy-ethyl)-amino]-methyl}-benzyl)-methyl-amino]-6,7-dimethoxy-5-methyl-1H-quinazolin-4-one		454.6 M ⁺
C-19	6,7-Dimethoxy-5-methyl-2-[methyl-(3-pyrrolidin-1-ylmethyl-benzyl)-amino]-1H-quinazolin-4-one		422.5M ⁺
C-20	5-Hydroxy-6,7-dimethoxy-2-{methyl-[3-(1-methyl-4,5-dihydro-1H-imidazol-2-ylmethyl)-benzyl]-amino}-1H-quinazolin-4-one		438
C-21	5,6,7-Trimethoxy-2-{methyl-[3-(1-methyl-4,5-dihydro-1H-imidazol-2-ylmethyl)-benzyl]-amino}-1H-quinazolin-4-one		452
C-22	5-Hydroxy-2-[[3-(1-isopropyl-4,5-dihydro-1H-		466

	imidazol-2-ylmethyl)-benzyl]-methyl-amino}-6,7-dimethoxy-1H-quinazolin-4-one	
C-23	2-[[3-(1-Isopropyl-4,5-dihydro-1H-imidazol-2-ylmethyl)-benzyl]-methyl-amino]-5,6,7-trimethoxy-1H-quinazolin-4-one	480
C-24	5,6,7-Trimethoxy-2-{methyl-[3-(1-methyl-piperidin-4-ylmethyl)-benzyl]-amino}-1H-quinazolin-4-one	466.6 M ⁺

Additional compounds prepared according to the procedure of Example C-1 are shown in Table 11.

Examples D-1 to D-8, E-1 to E-15

Compounds of formula Ic wherein Y is -CH₂- of Tables 4 and 5 were prepared following the same or similar method as described for the above Examples.

TABLE 4

Ex.	Compound Name	Mp°C	MS (M ⁺)
D-1	2-[(3-Methanesulfonyl-benzyl)-methyl-amino]-6,7-dimethoxy-5-methyl-1H-quinazolin-4-one	269.5-271.5	417.5
D-2	2-[(3-Methanesulfonyl-benzyl)-methyl-amino]-5,6,7-trimethoxy-1H-quinazolin-4-one	231.5-232.9	433.5
D-3	N,N-Dimethyl-3-{[methyl-(5,6,7-trimethoxy-4-oxo-1,4-dihydro-quinazolin-2-yl)-amino]-methyl}-benzenesulfonamide		462.5
D-4	3-[[[(6,7-Dimethoxy-5-methyl-4-oxo-1,4-dihydro-quinazolin-2-yl)-methyl-amino]-methyl]-N-(2-methoxy-ethyl)-benzenesulfonamide		476.6
D-5	N-(2-Methoxy-ethyl)-3-{[methyl-(5,6,7-trimethoxy-4-oxo-1,4-dihydro-quinazolin-2-yl)-amino]-methyl}-benzenesulfonamide		492.6
D-6	3-[[[(6,7-Dimethoxy-5-methyl-4-oxo-1,4-dihydro-quinazolin-2-yl)-methyl-amino]-methyl]-N-(3-methoxy-propyl)-benzenesulfonamide		490.6
D-7	3-[[[(6,7-dimethoxy-5-methyl-4-oxo-1,4-dihydro-quinazolin-2-yl)-methyl-amino]-methyl]-benzoic acid		383.4

D-8	3-[[Methyl-(5,6,7-trimethoxy-4-oxo-1,4-dihydro-quinazolin-2-yl)-amino]-methyl]-benzoic acid	400 (M+H) ⁺
-----	---	------------------------

TABLE 5

Ex.	Compound Name	Mp °C	MS (M+H) ⁺
E-1	3-[[[(6,7-Dimethoxy-5-methyl-4-oxo-1,4-dihydro-quinazolin-2-yl)-methyl-amino]-methyl]-N-methyl-benzamide	243.3-245.1	396.4 M ⁺
E-2	N-Methyl-3-[[methyl-(5,6,7-trimethoxy-4-oxo-1,4-dihydro-quinazolin-2-yl)-amino]-methyl]-benzamide	217.9-219.1	412.4 M ⁺
E-3	N-(2-Dimethylamino-ethyl)-3-[[methyl-(5,6,7-trimethoxy-4-oxo-1,4-dihydro-quinazolin-2-yl)-amino]-methyl]-benzamide		470
E-4	N-(2-Dimethylamino-ethyl)-N-methyl-3-[[methyl-(5,6,7-trimethoxy-4-oxo-1,4-dihydro-quinazolin-2-yl)-amino]-methyl]-benzamide		484
E-5	N-(3-Dimethylamino-propyl)-N-methyl-3-[[methyl-(5,6,7-trimethoxy-4-oxo-1,4-dihydro-quinazolin-2-yl)-amino]-methyl]-benzamide		498
E-6	N-(2-Hydroxy-ethyl)-3-[[methyl-(5,6,7-trimethoxy-4-oxo-1,4-dihydro-quinazolin-2-yl)-amino]-methyl]-benzamide		443
E-7	N-(2-Hydroxy-propyl)-3-[[methyl-(5,6,7-trimethoxy-4-oxo-1,4-dihydro-quinazolin-2-yl)-amino]-methyl]-benzamide		457
E-8	N-(2-Hydroxy-ethyl)-N-methyl-3-[[methyl-(5,6,7-trimethoxy-4-oxo-1,4-dihydro-quinazolin-2-yl)-amino]-methyl]-benzamide		457
E-9	N-(2-Methoxy-ethyl)-3-[[methyl-(5,6,7-trimethoxy-4-oxo-1,4-dihydro-quinazolin-2-yl)-amino]-methyl]-benzamide		457
E-10	N-(3-Methoxy-propyl)-3-[[methyl-(5,6,7-trimethoxy-4-oxo-1,4-dihydro-quinazolin-2-yl)-amino]-methyl]-benzamide		471

E-11	N-Cyanomethyl-3-{[methyl-(5,6,7-trimethoxy-4-oxo-1,4-dihydro-quinazolin-2-yl)-amino]-methyl}-benzamide	438
E-12	3-{[Methyl-(5,6,7-trimethoxy-4-oxo-1,4-dihydro-quinazolin-2-yl)-amino]-methyl}-N-(tetrahydrofuran-2-ylmethyl)-benzamide	483
E-13	N-Furan-2-ylmethyl-3-{[methyl-(5,6,7-trimethoxy-4-oxo-1,4-dihydro-quinazolin-2-yl)-amino]-methyl}-benzamide	479
E-14	3-{[Methyl-(5,6,7-trimethoxy-4-oxo-1,4-dihydro-quinazolin-2-yl)-amino]-methyl}-N-(2-morpholin-4-yl-ethyl)-benzamide	512
E-15	2-{{3-(4-Hydroxymethyl-piperidine-1-carbonyl)-benzyl}-methyl-amino}-5,6,7-trimethoxy-1H-quinazolin-4-one	497

Example F-1: 2-{{3-(2-Hydroxy-ethoxy)-benzyl}-methyl-amino}-5,6,7-trimethoxy-1H-quinazolin-4-one

Step 1: 3-[2-(tert-Butyl-dimethyl-silanyloxy)-ethoxy]-benzaldehyde: TBDMSCl

5 silylation reagent in DMF (60 mL) was added to 3-(2-hydroxy-ethoxy)-benzaldehyde (4.985 g, 30.00 mmol). The mixture was stirred at RT for 72 h. The mixture was diluted with water and extracted with diethyl ether. The ethereal extracts were dried over Na₂SO₄ and concentrated *in vacuo*. The residue was purified by SiO₂ eluted with 95:5 CH₂Cl₂/MeOH to give 2.38 g (28%) of a clear oil.

10 Step 2: {3-[2-(tert-Butyl-dimethyl-silanyloxy)-ethoxy]-benzylidene}-methyl-amine: 3-[2-(tert-Butyl-dimethyl-silanyloxy)-ethoxy]-benzaldehyde (2.375 g, 8.47 mmol) was dissolved in 10 mL EtOH. MeNH₂ (1.2 mL, 40% in H₂O) was added and the mixture was heated to 40°C for 30 min. The mixture was then concentrated *in vacuo*. The aqueous residue was diluted with isopropanol and then concentrated *in vacuo*. This was
15 repeated twice more giving 2.184 g (87.89%) of a dry white powder.

Step 3: {3-[2-(tert-Butyl-dimethyl-silanyloxy)-ethoxy]-benzyl}-methyl-amine: {3-[2-(tert-Butyl-dimethyl-silanyloxy)-ethoxy]-benzylidene}-methyl-amine (1.174 g, 4.00 mmol) was dissolved in EtOH and cooled to 0°C. NaBH₄ (189 mg, 5.00 mmol) was added and the mixture was allowed to stir for 4 h. The reaction was carefully quenched

with water. The EtOH was removed *in vacuo*. The aqueous residue was extracted with CH₂Cl₂. The organic extracts were dried over Na₂SO₄ and concentrated *in vacuo* to give 1.185 g (quant) of a white powder.

Step 4: 2-([3-[2-(tert-Butyl-dimethyl-silanyloxy)-ethoxy]-benzyl]-methyl-amino)-5,6,7-trimethoxy-1*H*-quinazolin-4-one: {3-[2-(tert-Butyl-dimethyl-silanyloxy)-ethoxy]-benzyl}-methyl-amine from Step 3 (294 mg, 1.00 mmol) and 2-chloro-5,6,7-trimethoxy-1*H*-quinazolin-4-one (270 mg, 1.00 mmol) were dissolved in dry DMSO. Et₃N (0.14 mL, 1.00 mmol) was added and the mixture was heated to 120°C overnight. Upon cooling, the mixture was poured into water and extracted with diethyl ether. The ethereal extracts were dried over Na₂SO₄ and concentrated *in vacuo*. The residue was purified by SiO₂ eluted with 99:1 CH₂Cl₂/MeOH to give 435 mg (82.12%) of a white crystalline solid.

Step 5: 2-([3-(2-Hydroxy-ethoxy)-benzyl]-methyl-amino)-5,6,7-trimethoxy-1*H*-quinazolin-4-one: Compound from step 4 (431 mg, 0.82 mmol) was dissolved in dry THF (15 mL). Bu₄NF (0.82 mL, 1.0 M in THF) was added and the mixture was stirred overnight. The mixture was concentrated *in vacuo*. The residue was dissolved in hot MeOH and allowed to stand for crystallization. A small fraction of diethyl ether was added to induce crystallization. The solid was filtered and dried to give 56 mg (16.57%) of a white crystalline solid. mp 146.0-147.5 °C; ¹H NMR (D₂O-*d*₂) δ 3.19 (s, 3 H), 3.86 (s, 3 H), 3.88 (s, 3 H), 3.89 (t, 2 H, *J* = 4.4), 3.92 (s, 3 H), 4.04 (t, 2 H, *J* = 4.4), 4.85 (s, 2 H), 6.63 (s, 1 H), 6.83 (m, 3 H), 7.23 (t, 1 H, *J* = 7.8); ¹³C NMR (DMSO-*d*₆, 39.50) δ 35.57, 52.90, 56.37, 61.81, 61.83, 62.16, 69.55, 102.77, 104.74, 113.65, 114.35, 120.55, 130.24, 138.86, 139.30, 150.96, 151.03, 153.47, 159.45, 159.56, 162.81; MS (ES+) *m/z* 416 (*M* + *H*); Anal. (C₂₁H₂₅N₃O₆) C: calcd, 60.71; found, 60.39; H: calcd, 6.07; found, 5.97; N: calcd, 10.11; found, 10.01.

Example F-2: 2-([3-(Methanesulfinylmethoxy-benzyl)-methyl-amino]-6,7-dimethoxy-5-methyl-1*H*-quinazolin-4-one

Step 1: 3-Methylsulfinylmethoxy-benzaldehyde: 3-Hydroxybenzaldehyde (6.106 g, 50.00 mmol) was dissolved in acetone at RT. K₂CO₃ (13.821 g, 100.00 mmol), NaI (7.569 g, 50.00 mmol) and chloromethyl methyl sulfide (4.19 mL, 50.00 mmol) were added, and the mixture was allowed to stir for 80 h. The mixture was filtered, the filtrate concentrated to an oil, and the residue purified by SiO₂ eluted with 99:1 CH₂Cl₂/MeOH to give 3.309 g (36.31%) of a yellow oil.

Step 2: Methyl-(3-methylsulfanylmethoxy-benzylidene)-amine: 3-Methylsulfanylmethoxy-benzaldehyde from step 1 (3.309 g, 18.16 mmol) was dissolved in EtOH (9 mL). MeNH₂ (2.55 mL, 40% in H₂O) was added and the mixture was heated to 40°C for 30 min. Upon cooling, the mixture was concentrated *in vacuo*. The aqueous residue was
5 diluted with isopropanol and then concentrated *in vacuo*. This was repeated twice more giving 3.55g (quant) of a pale yellow oil.


Step 3: Methyl-(3-methylsulfanylmethoxy-benzyl)-amine: Compound from step 2 (3.546 g, 18.16 mmol) was dissolved in EtOH (50 mL) and cooled to 0°C. NaBH₄ (859 mg, 22.70 mmol) was added and the mixture was allowed to stir for 4 h. The reaction
10 was carefully quenched with water. The EtOH was removed *in vacuo*. The aqueous residue was extracted with CH₂Cl₂. The organic extracts were dried over Na₂SO₄ and concentrated *in vacuo* to give 2.044 g (57%) of a clear yellow oil.

Step 4: (3-Methanesulfinylmethoxy-benzyl)-methyl-amine: Compound from step 3 (592 mg, 3.00 mmol) was dissolved in glacial acetic acid (4 mL) and cooled to 0°C. H₂O₂ (0.34
15 mL, 3.00 mmol) was added drop-wise. The mixture was allowed to stir for 6 h. The mixture was then diluted with CH₂Cl₂ and neutralized with K₂CO₃. The solids were filtered off. The filtrate was dried over Na₂SO₄ and concentrated *in vacuo* to give 640 mg (quant) of a yellow crystalline solid.


Step 5: 2-[(3-Methanesulfinylmethoxy-benzyl)-methyl-amino]-6,7-dimethoxy-5-methyl-1H-quinazolin-4-one: Compound from step 4 (213 mg, 1.00 mmol) and 2-chloro-6,7-dimethoxy-5-methyl-1H-quinazolin-4-one (254 mg, 1.00 mmol) were dissolved in dry DMSO. Et₃N (0.14 mL, 1.00 mmol) was added, and the mixture was heated to 120°C overnight. Upon cooling, the mixture was poured into water and extracted with diethyl ether. The ethereal extracts were dried over Na₂SO₄ and
25 concentrated *in vacuo*. The residue was recrystallized from MeOH to give 54 mg (12.53%) of the above Example F-2 as a white crystalline powder. mp 191.0-193.5 °C; ¹H NMR (DMSO-*d*₆, 2.49) δ 2.59 (s, 3 H), 2.60 (s, 3 H), 3.03 (s, 3 H), 3.62 (s, 3 H), 3.84 (s, 3 H), 4.79 (s, 2 H), 5.02 (d, 1 H, *J* = 10.6), 5.22 (d, 1 H, *J* = 10.6), 6.64 (s, 1 H), 6.90 (d, 1 H, *J* = 7.6), 7.01 (m, 2 H), 7.30 (t, 1 H, *J* = 7.9); ¹³C NMR (DMSO-*d*₆, 39.50) δ 13.90, 34.66,
30 35.59, 51.90, 55.93, 60.23, 84.92, 104.81, 108.46, 114.18, 115.02, 121.22, 130.13, 131.94, 140.10, 142.90, 150.63, 150.97, 157.71, 157.97, 163.67; MS (ES+) *m/z* 432 (M + H); Anal. (C₂₁H₂₅N₃O₅S) C: calcd, 58.45; found, 58.32; H: calcd, 5.84; found, 5.74; N: calcd, 9.74; found, 9.77.

Example F-3: 6,7-Dimethoxy-5-methyl-2-{methyl-[3-(pyrimidin-2-yloxy)-benzyl]-amino}-1*H*-quinazolin-4-one

Step 1: A mixture of 3-hydroxybenzaldehyde (2.00 g, 16.38 mmol), 2-chloropyrimidine (1.88 g, 16.38 mmol) and ⁱPr₂NEt (5.71 mL, 32.75 mmol), under N₂ atmosphere was
5 heated at 105°C for 24 h. The reaction mixture was allowed to cool to RT. The crude was subjected to column chromatography (silica gel, 20% EtOAc in hexanes) to furnish 1.62 g of the desired aryl ether (1) as an off-white solid.



Step 2: To a solution of the aryl ether (1) from Step 1 (1.00 g, 5.00 mmol) and MeNH₂
10 (3.00 mL, 2.0 M in THF, 5.99 mmol) in MeOH (75 mL) at RT was added NaCNBH₃ (377 mg, 5.99 mmol). The reaction was stirred for 7 h before it was concentrated under reduced pressure. The residue was taken-up in a mixture of 1 N aq. HCl (20 mL) and CH₂Cl₂ (5 mL). The aq. layer was separated, adjusted to pH 10, and extracted with CH₂Cl₂. The CH₂Cl₂ extract was washed with brine, dried over MgSO₄, and
15 concentrated. The residue was purified by column chromatography (silica gel, 10% MeOH in CH₂Cl₂) to furnish 363 mg of the benzylamine (2) as a faintly-colored oil.



Step 3: A mixture of the benzylamine (2) from Step 2 (152 mg, 0.71 mmol), 2-chloro-6,7-dimethoxy-5-methyl-quinazolin-4-one (150 mg, 0.59 mmol) and ⁱPr₂NEt (205 μ L, 1.18
20 mmol) in anhydrous EtOH (5 mL) in a sealed tube was heated at 65°C for 16 h before it was allowed to cool to RT slowly. The precipitate was collected by suction filtration, washed with EtOH, and dried in the oven at 60°C to provide 57 mg of Example F-3 as an off-white solid. Mp 224.0-227.8 °C; TLC R_f 0.52 (5% MeOH in CH₂Cl₂); ¹H NMR (DMSO-*d*₆, 2.49) δ 2.60 (s, 3 H), 3.05 (s, 3 H), 3.62 (s, 3 H), 3.83 (s, 3 H), 4.85 (s, 2 H),
25 6.62 (br s, 1 H), 7.08-7.13 (m, 3 H), 7.24 (t, 1 H, *J* = 4.8 Hz), 7.40 (t, 1 H, *J* = 7.7 Hz), 8.60 (d, 2 H, *J* = 4.8 Hz), 10.84 (br s, 1 H); MS (ES+) *m/z* 434 (M + H); Anal. (C₂₃H₂₃N₅O₄·2HCl·0.5H₂O) C: calcd, 53.60; found, 53.55; H: calcd, 13.58; found, 13.52; N: calcd, 5.08; found, 4.48.

Examples F-4 to F-16

Compounds of formula Ic wherein Y is $-\text{CH}_2-$ of Table 6 were prepared following the same or similar method as described for Examples F-1 through F-3.

TABLE 6

Ex.	Compound Name	Mp°C	MSM ⁺
F-4	2-[(3-Hydroxy-benzyl)-methyl-amino]-6,7-dimethoxy-5-methyl-1H-quinazolin-4-one		355.4
F-5	2-[(3-Hydroxy-benzyl)-methyl-amino]-5,6,7-trimethoxy-1H-quinazolin-4-one		371.4
F-6	2-[(3-Methanesulfonylmethoxy-benzyl)-methyl-amino]-5,6,7-trimethoxy-1H-quinazolin-4-one	183.9-185.4	463.5
F-7	(3-[[6,7-Dimethoxy-5-methyl-4-oxo-1,4-dihydroquinazolin-2-yl)-methyl-amino]-methyl]-phenoxy)-acetic acid		413.4
F-8	(3-[[Methyl-(5,6,7-trimethoxy-4-oxo-1,4-dihydroquinazolin-2-yl)-amino]-methyl]-phenoxy)-acetic acid		429.4
F-9	2-{[3-(2-Hydroxy-ethoxy)-benzyl]-methyl-amino}-6,7-dimethoxy-5-methyl-1H-quinazolin-4-one	180.5-183.1	399.4
F-10	6,7-Dimethoxy-2-{[3-(2-methoxy-ethoxy)-benzyl]-methyl-amino}-5-methyl-1H-quinazolin-4-one	144.8-145.9	413.5
F-11	5,6,7-Trimethoxy-2-{[3-(2-methoxy-ethoxy)-benzyl]-methyl-amino}-1H-quinazolin-4-one	144.8-145.9	429.5
F-12	6,7-Dimethoxy-2-{[3-(3-methoxy-propoxy)-benzyl]-methyl-amino}-5-methyl-1H-quinazolin-4-one		427.5
F-13	5,6,7-Trimethoxy-2-{[3-(3-methoxy-propoxy)-benzyl]-methyl-amino}-1H-quinazolin-4-one		443.5
F-14	2-{[3-(2-Dimethylamino-ethoxy)-benzyl]-methyl-amino}-6,7-dimethoxy-5-methyl-1H-quinazolin-4-one		426.5
F-15	2-{[3-(2-Ethylamino-ethoxy)-benzyl]-methyl-amino}-6,7-dimethoxy-5-methyl-1H-quinazolin-4-one		426.5
F-16	5,6,7-Trimethoxy-2-{methyl-[3-(pyrimidin-2-yloxy)-benzyl]-amino}-1H-quinazolin-4-one	191-192.9	449.5

5 Additional compounds prepared according to the procedure of Examples F-1 through F-3 are shown in Table 11.

Example G-1: N'-(3-[[(6,7-Dimethoxy-5-methyl-4-oxo-1,4-dihydro-quinazolin-2-yl)-methyl-amino]-methyl]-phenyl)-N,N-dimethyl-acetamidine

The aniline of Example B-1 (600 mg, 1.7 mmol), the dimethylacetal of DMF (330 mg, 2.23 mmol), and DMF (2 mL) was kept at 100°C for 35 min and then stored at ambient temperature for 16 h. The precipitated product was collected and washed with diethyl ether affording 340 mg of product. This material was further purified by chromatography (EM Science silica gel 60; 3% MeOH in DCM containing 0.5 % of ammonium hydroxide) to yield 250 mg (35%) of the above compound as a white solid. Mp 174.4-174.8°C; ms 424 (M+H). The HCl salt of the above compound was prepared mixing 600 mg (1.4 mmol) of the free base with 8 mL hot EtOH, and then adding to this solution 2.7 mL of a 1.4 M solution of HCl in EtOH (3.7 mmol). The HCl salt was crystallized by addition of diethyl ether. The collected product was dried at 80°C *in vacuo* to furnish 620 mg. Mp 170.6-172.5°C; Anal. (C₂₃H₂₉N₅O₃•2HCl•0.5 H₂O) Calcd.: C, 54.65; H, 6.38; N, 13.86. Found: C, 54.40; H, 6.14; N, 13.81.

Examples G-2 to G-55

Compounds of formula I wherein R and R' and R¹⁰ are methyl of Table 7 were prepared following the same or similar method as described for Example G-1.

TABLE 7

Ex. No	Compound Name	Mp°C	MW
G-2	N-(3-[[(5-Isopropyl-6,7-dimethoxy-4-oxo-1,4-dihydro-quinazolin-2-yl)-methyl-amino]-methyl]-phenyl)-methanesulfonamide	215-217	460.55
G-3	6,7-Dimethoxy-5-methyl-2-[methyl-(3-[(dimethylamino)sulfonyl]amino)-benzyl)-amino]-1H-quinazolin-4-one		461.54
G-4	6,7-Dimethoxy-5-methyl-2-[methyl-(3-[(dimethylamino)sulfonyl]methylamino)-benzyl)-amino]-1H-quinazolin-4-one		475.57
G-5	5,6,7-Trimethoxy-2-[methyl-(3-[(dimethylamino)sulfonyl]methylamino)-benzyl)-amino]-1H-quinazolin-4-one		477.54
G-6	5,6,7-Trimethoxy-2-[methyl-(3-[(dimethylamino)sulfonyl]methylamino)-benzyl)-amino]-1H-quinazolin-4-		491.57

	one		
G-7	N-(3-[[[(6,7-Dimethoxy-5-methyl-4-oxo-1,4-dihydro-quinazolin-2-yl)-methyl-amino]-methyl]-phenyl]-acetamide		396.45
G-8	N-(3-[[Methyl-(5,6,7-trimethoxy-4-oxo-1,4-dihydro-quinazolin-2-yl)-amino]-methyl]-phenyl)-acetamide		412.44
G-9	N-(4-Chloro-3-[[methyl-(5,6,7-trimethoxy-4-oxo-1,4-dihydro-quinazolin-2-yl)-amino]-methyl]-phenyl)-acetamide		446.89
G-10	1,1-Dimethyl-3-(3-[[methyl-(5,6,7-trimethoxy-4-oxo-1,4-dihydro-quinazolin-2-yl)-amino]-methyl]-phenyl)-urea		441.49
G-11	(3-[[Methyl-(5,6,7-trimethoxy-4-oxo-1,4-dihydro-quinazolin-2-yl)-amino]-methyl]-phenyl)-carbamic acid methyl ester		428.44
G-12	(3-[[[(6,7-Dimethoxy-5-methyl-4-oxo-1,4-dihydro-quinazolin-2-yl)-methyl-amino]-methyl]-phenyl]-methyl-carbamic acid methyl ester		426.47
G-13	N-(3-[[[(6,7-dimethoxy-5-methyl-4-oxo-1,4-dihydro-quinazolin-2-yl)-methyl-amino]-methyl]-phenyl]-N-methyl-2-oxo-propionamide		438.48
G-14	3-(3-[[[(6,7-Dimethoxy-5-methyl-4-oxo-1,4-dihydro-quinazolin-2-yl)-methyl-amino]-methyl]-phenyl]-1,1-dimethyl-urea		425.49
G-15	1-Methoxy-3-(3-[[methyl-(5,6,7-trimethoxy-4-oxo-1,4-dihydro-quinazolin-2-yl)-amino]-methyl]-phenyl)-urea		443.46
G-16	2-[[3-(2-Hydroxy-ethylamino)-benzyl]-methyl-amino]-5,6,7-trimethoxy-1H-quinazolin-4-one	60.5- 61.4	414.46
G-17	5-Hydroxy-2-([3-[(3-hydroxy-propyl)-methyl-amino]-benzyl]-methyl-amino)-6,7-dimethoxy-1H-quinazolin-4-one		428.49
G-18	2-[[3-(3-Hydroxy-propylamino)-benzyl]-methyl-amino]-5,6,7-trimethoxy-1H-quinazolin-4-one	47-48.5	428.49
G-19	5,6,7-Trimethoxy-2-[[3-(2-methoxy-ethylamino)-benzyl]-methyl-amino]-1H-quinazolin-4-one	178.9- 179.9	428.49
G-20	5,6,7-Trimethoxy-2-([3-[(2-methoxy-ethyl)-methyl-	179.3-	442.51

	amino]-benzyl}-methyl-amino)-1H-quinazolin-4-one	181.2	
G-21	2-({3-[(3-Hydroxy-propyl)-methyl-amino]-benzyl}-methyl-amino)-5,6,7-trimethoxy-1H-quinazolin-4-one	156-157.2	442.51
G-22	5,6,7-Trimethoxy-2-{{3-(3-methoxy-propylamino)-benzyl}-methyl-amino}-1H-quinazolin-4-one, hydrochloride salt		442.51
G-23	(3-{{[(6,7-Dimethoxy-5-methyl-4-oxo-1,4-dihydro-quinazolin-2-yl)-methyl-amino]-methyl}-phenyl)-methyl-carbamic acid methyl ester		455.5
G-24	2-({3-[Bis-(2-hydroxy-ethyl)-amino]-benzyl}-methyl-amino)-5,6,7-trimethoxy-1H-quinazolin-4-one		458.51
G-25	N-[2-(3-{1-[(6,7-Dimethoxy-5-methyl-4-oxo-1,4-dihydro-quinazolin-2-yl)-methyl-amino]-ethyl}-phenylamino)-ethyl]-acetamide		453.54
G-26	N-(3-{{[(6,7-Dimethoxy-5-methyl-4-oxo-1,4-dihydro-quinazolin-2-yl)-methyl-amino]-methyl}-phenyl)-acetamidine	155-58	395.46
G-27	N-(3-{{[Methyl-(5,6,7-trimethoxy-4-oxo-1,4-dihydro-quinazolin-2-yl)-amino]-methyl}-phenyl)-acetamidine		411.46
G-28	N-(3-{{[(6,7-Dimethoxy-5-methyl-4-oxo-1,4-dihydro-quinazolin-2-yl)-methyl-amino]-methyl}-phenyl)-N,N'-dimethyl-acetamidine		423.51
G-29	N-(3-{{[(6,7-Dimethoxy-5-methyl-4-oxo-1,4-dihydro-quinazolin-2-yl)-methyl-amino]-methyl}-phenyl)-N'-methyl-acetamidine		409.49
G-30	N'-(3-{{[(6,7-Dimethoxy-5-methyl-4-oxo-1,4-dihydro-quinazolin-2-yl)-methyl-amino]-methyl}-phenyl)-N,N-dimethyl-formamidine	185-86	409.49
G31	N,N-Dimethyl-N'-(3-{{[methyl-(5,6,7-trimethoxy-4-oxo-1,4-dihydro-quinazolin-2-yl)-amino]-methyl}-phenyl)-acetamidine	199-201	439.51
G-32	N,N-Dimethyl-N'-(3-{{[methyl-(5,6,7-trimethoxy-4-oxo-1,4-dihydro-quinazolin-2-yl)-amino]-methyl}-phenyl)-isobutyramidine	148-50	467.57
G-33	N'-(3-{{[(5-Ethyl-6,7-dimethoxy-4-oxo-1,4-dihydro-quinazolin-2-yl)-methyl-amino]-methyl}-phenyl)-N,N-	224.6-225.5	437.54

	dimethyl-acetamidine		
G-34	N'-(3-{[(5-Isopropyl-6,7-dimethoxy-4-oxo-1,4-dihydro-quinazolin-2-yl)-methyl-amino]-methyl}-phenyl)-N,N-dimethyl-acetamidine		451.57
G-35	N'-(3-{[(5-Hydroxy-6,7-dimethoxy-4-oxo-1,4-dihydro-quinazolin-2-yl)-methyl-amino]-methyl}-phenyl)-N,N-dimethyl-acetamidine		425.49
G-36	3-Methoxy-N,N-dimethyl-N'-(3-{[methyl-(5,6,7-trimethoxy-4-oxo-1,4-dihydro-quinazolin-2-yl)-amino]-methyl}-phenyl)-propionamidine	145	483.57
G-37	N,N-Dimethyl-N'-(3-{[methyl-(5,6,7-trimethoxy-4-oxo-1,4-dihydro-quinazolin-2-yl)-amino]-methyl}-phenyl)-cyclobutanecarboxamidine		479.57
G-38	2-{[3-(Imidazolidin-2-ylideneamino)-benzyl]-methyl-amino}-6,7-dimethoxy-5-methyl-1H-quinazolin-4-one	141.3- 144.7	422.49
G-39	5,6,7-Trimethoxy-2-{methyl-[3-(1-methyl-pyrrolidin-2-ylideneamino)-benzyl]-amino}-1H-quinazolin-4-one	130-133	451.52
G-40	2-{[2-Chloro-5-(1,3-dimethyl-imidazolidin-2-ylideneamino)-benzyl]-methyl-amino}-5,6,7-trimethoxy-1H-quinazolin-4-one		500.98
G-41	2-{[3-(1,3-dimethyl-imidazolidin-2-ylideneamino)-benzyl]-methyl-amino}-6,7-diimethoxy-5-ethyl-1H-quinazolin-4-one, hexafluorophosphate	212.5- 219.9	464.57
G-42	2-{[3-(4,5-Dihydro-3H-pyrrol-2-ylamino)-benzyl]-methyl-amino}-6,7-dimethoxy-5-methyl-1H-quinazolin-4-one	175.4- 177.2	421.50
G-43	2-{[3-(4,5-Dihydro-3H-pyrrol-2-ylamino)-benzyl]-methyl-amino}-5,6,7-trimethoxy-1H-quinazolin-4-one	207-209	437.50
G-44	5,6,7-Trimethoxy-2-{methyl-[3-(2-oxo-tetrahydrofuran-3-ylamino)-benzyl]-amino}-1H-quinazolin-4-one		454.48
G-45	2-({5-[(4,5-Dihydro-3H-pyrrol-2-yl)-methyl-amino]-2-fluoro-benzyl}-methyl-amino)-5-hydroxy-6,7-dimethoxy-1H-quinazolin-4-one		455.49
G-46	6,7-Dimethoxy-5-methyl-2-{methyl-[3-(pyrimidin-2-ylamino)-benzyl]-amino}-1H-quinazolin-4-one		432.48
G-47	5,6,7-Trimethoxy-2-{methyl-[3-(pyrimidin-2-ylamino)-		448.48


	benzyl]-amino}-1H-quinazolin-4-one	
G-48	2-{[3-(2-Chloro-pyrimidin-4-ylamino)-benzyl]-methyl-amino}-5,6,7-trimethoxy-1H-quinazolin-4-one	482.93
G-49	2-{[3-(4-Chloro-pyrimidin-2-ylamino)-benzyl]-methyl-amino}-5,6,7-trimethoxy-1H-quinazolin-4-one	482.93
G-50	2-{[3-(4-Chloro-pyrimidin-2-ylamino)-benzyl]-methyl-amino}-6,7-dimethoxy-5-methyl-1H-quinazolin-4-one	466.93
G-51	2-{[3-(2-Chloro-5-methyl-pyrimidin-4-ylamino)-benzyl]-methyl-amino}-5,6,7-trimethoxy-1H-quinazolin-4-one	496.95
G-52	2-{[3-(2-Chloro-6-methyl-pyrimidin-4-ylamino)-benzyl]-methyl-amino}-5,6,7-trimethoxy-1H-quinazolin-4-one	496.95
G-53	2-{[3-(4-Chloro-6-methyl-pyrimidin-2-ylamino)-benzyl]-methyl-amino}-5,6,7-trimethoxy-1H-quinazolin-4-one	496.95
G-54	2-({3-[(1H-Imidazol-2-ylmethyl)-amino]-benzyl}-methyl-amino)-5,6,7-trimethoxy-1H-quinazolin-4-one	450.50
G-55	N'-(3-[[5-Fluoro-6,7-dimethoxy-4-oxo-1,4-dihydro-quinazolin-2-yl)-methyl-amino]-methyl}-phenyl)-N,N-dimethyl-acetamide	427.48

Additional compounds prepared according to the procedure of Example G-1 are shown in Table 11.


Example H-1: 5,6,7-Trimethoxy-2-{methyl-[3-(2-oxo-pyrrolidin-1-yl)-benzyl]-amino}-1H-quinazolin-4-one

- 5 Step 1: To a solution of aniline (1) (1.39 g, 5.88 mmol) and ⁱPr₂N⁺Et (1.13 mL, 6.42 mmol) in THF (7 mL) at 0°C was added dropwise a solution of 4-chlorobutyl chloride (659 μL, 5.88 mmol) in THF (3 mL). The reaction mixture was then allowed to warm to RT and stirred for 1 h before it was poured into a mixture of EtOAc and saturated aq. NH₄Cl. The organic layer was separated and concentrated under reduced pressure to
- 10 provide the crude amide intermediate (2).

- 80 -



Step 2: To a solution of the amide intermediate (2) from Step 1 in MeCN (150 mL) at RT was added dropwise a solution of 50% aq. NaOH (0.94 mL, 11.76 mmol). The reaction mixture was stirred for 1 h before it was concentrated under reduced pressure. The residue was partitioned between EtOAc and H₂O. The organic layer was washed with brine, dried over MgSO₄, and concentrated. The residue was purified (silica gel, EtOAc) to furnish 1.80 g of lactam (3) as colorless oil.



Step 3: Trifluoroacetic acid (375 μ L, 4.87 mmol) was added via syringe to a stirred solution of (3) from Step 2 (189 mg, 0.62 mmol) at 0°C. After 3 hours, the solution was concentrated under reduced pressure, then pumped under vacuum for 1 hour. The crude residue was dissolved in EtOH (2 mL), and added to a sealed tube containing 2-chloro-5,6,7-trimethoxy-1*H*-quinazolin-4-one (132 mg, 0.49 mmol), diisopropylethylamine (425 μ L, 2.44 mmol), and EtOH (3 mL). The suspension was heated at 110°C for 2 hours, then cooled to RT and stirred overnight. The reaction mixture was concentrated under reduced pressure, and purified by chromatography to give 128 mg (60%) of the above Example H-1 as a cream colored solid. mp 165.5-166.5°C; TLC R_f 0.5 (5% MeOH/DCM); IR (KBr) ν_{max} 3429, 2933, 1657, 1592, 1481, 1452, 1120, 1047, 933; ¹H NMR (DMSO-*d*₆, 2.49) 2.04 (m, 2H), 2.50 (t, 2H, *J* = 1.9), 3.03 (s, 3H), 3.70 (s, 3H), 3.77 (s, 1H), 3.81 (t, 2H, *J* = 6.9), 3.84 (s, 3H), 4.81 (s, 2H), 6.57 (s, 2H), 6.99 (d, 1H, *J* = 7.7), 7.32 (t, 1H, *J* = 7.8), 7.49 (dd, 1H, *J* = 2.1, 7.4), 7.65 (s, 1H), 10.83 (s, 1H); MS (ES+) *m/z* 439 (*M* + H); Anal. (C₂₃H₂₆N₄O₅ w/ 0.25M CH₂Cl₂): calcd 60.75; found: 60.65; H: calcd, 5.81; found: 5.79; N: calcd, 12.19; found, 11.98).

Example H-2: 6,7-Dimethoxy-5-methyl-2-{methyl-[3-(1-methyl-4,5-dihydro-1*H*-imidazol-2-yl)-benzyl]-amino}-1*H*-quinazolin-4-one

Step 1: Diisopropylethylamine (0.82 mL, 4.71 mmol) was added to the mixture of 2-chloro-6,7-dimethoxy-5-methyl-quinazolin-4-one (1.0 g, 3.92 mmol) and *N*-methyl-(3-cyano-benzyl)-amine (0.68 g, 4.71 mmol) in EtOH (50 mL). It was heated at 120°C for 1.5h in the sealed tube. After cooling down, the resulting white solid was filtered and
5 washed with MeOH: mp 279.9-281.5°C; ¹H NMR (DMSO-d₆) δ 2.6 (s, 3H), 3.05 (s, 3H), 3.62 (s, 3H), 3.84 (s, 3H), 4.86 (s, 2H), 6.62 (s, 1H), 7.5-7.7.78 (m, 4H); ¹³C NMR (DMSO-d₆) δ 13.89, 35.58, 51.52, 55.93, 60.22, 104.79, 111.78, 119.15, 130.12, 131.19, 131.28, 131.96, 132.51, 140.04, 142.96, 150.51, 157.71;

Step 2: α-Bromo-m-cyanotoluene (1.0 g, 5.1 mmol) in 2M MeNH₂/MeOH (25.5 mL, 51
10 mmol) was refluxed for 1h. After evaporating most of solvent, the resulting solid was filtered. The mother liquor was concentrated *in vacuo* and it was purified by chromatography.

Step 3: A few drops of carbondisulfide were added to a solution of the aryl nitrile of Example B-2, above (0.08 g, 0.219 mol) in 1 mL of *N*-methylethylenediamine. The
15 mixture was allowed to stir at 120°C for 2 h. The excess amine was evaporated under reduced pressure and purified by flash column chromatography to provide the above Example H-2. Mp 145-148°C; IR (KBr) ν_{max} 2931, 1660, 1306 cm⁻¹; ¹H NMR (DMSO-d₆) δ 2.6 (s, 3H), 2.7 (s, 3H), 3.05 (s, 3H), 3.45 (t, 2H), 3.62 (s, 3H), 3.68 (t, 2H), 3.84 (s, 3H), 4.86 (s, 2H), 6.65 (s, 1H), 7.3-7.45 (m, 4H); ¹³C NMR (DMSO-d₆) δ 13.89, 35.60,
20 36.48, 39.05, 39.33, 39.60, 39.88, 51.82, 52.54, 53.90, 55.93, 60.23, 127.01, 127.10, 128.97, 129.04, 131.40, 131.98, 138.36, 142.94, 167.08; MS (ES+) *m/z* 422 (M + H); Anal. (C₂₃H₂₇N₅O₃·0.30M CH₂Cl₂) C; calcd, 62.61; found, 62.71; H; calcd, 6.22; found, 6.28; N; calcd, 15.67; found, 15.83.

Example H-3: 6,7-Dimethoxy-5-methyl-2-{methyl-[3-(1-methyl-piperidin-4-yl)-benzyl]-amino}-1*H*-quinazolin-4-one
25

Step 1: 4-(3-Methylcarbamoylphenyl)piperidine-1-methylcarboxamide: To a solution of 4-(3-carboxyphenyl)piperidine-1-carboxylic acid *tert*-butyl ester (1.0 g, 3.27 mmol) in 4 mL of DMF, was added CDI (0.63 g, 3.92 mmol) and the reaction mixture warmed to 60°C for two hours. After that time, the mixture was cooled to 0-5°C and, a 2 M solution
30 of methylamine in THF (4.91 mL, 9.82 mmol) was added via syringe and the mixture allowed to warm to RT overnight. The mixture was poured into a saturated solution of ammonium chloride, and the product was extracted with ethyl ether (3x50 mL). Organics were washed with water (3x30 mL) and brine (1x15 mL), then dried with

magnesium sulfate and concentrated. The dense liquid, crude product, was used into the next step without further treatment. Yield 1.05 g (100%). ^1H NMR (CDCl_3 , 7.26) δ 1.48 (s, 9 H), 1.63 (m, 2 H), 1.75 (m, 2 H), 2.69 (m, 1 H), 2.80 (m, 2 H), 3.01 (d, 3 H), 4.23 (br s, 2H), 6.15 (br s, D_2O , 1H), 7.32 (dt, 1H, $J = 6.87, 1.71$), 7.34 (t, 1H, $J = 6.72$), 7.55 (dt, 1H, $J = 6.87, 1.71$), 7.64 (t, 1H, $J = 1.71$); MS (ES+) m/z 319 (M + H), 341.2 (M + Na).

Step 2: Methyl-[3-(1-methylpiperidin-4-yl)benzyl]amine: To a solution of 4-(3-methyl-carbamoylphenyl)piperidine-1-methylcarboxamide (1.05 g, 3.29 mmol) in 50 mL of THF, was added portion-wise LiAlH_4 (0.625 g, 16.48 mmol). The reaction mixture was stirred at RT for 18 hours and then warmed to 50°C for 24 hours. Then, the mixture was cooled to RT and $\text{Na}_2\text{SO}_4 \cdot 10\text{H}_2\text{O}$ (10 g) was added portion-wise (very slowly), and the mixture stirred for two hours. Solids were removed by filtration (celite pad), rinsed with 10% MeOH/ CH_2Cl_2 (100 mL), and the filtrate concentrated. The residue was purified by flash chromatography, eluting with CH_2Cl_2 :MeOH: NH_4OH (60:10:1). Yield 0.59 g (81.9%). ^1H NMR (CDCl_3 , 7.26) δ 1.64 (br s, D_2O , 1H), 1.82 (m, 4 H), 2.04 (m, 2 H), 2.32 (s, 3 H), 2.45 (s, 3 H), 2.47 (m, 1 H), 2.97 (m, 2H), 3.72 (s, 2H), 7.13-7.18 (m, 3H), 7.26 (t, 1H, $J = 7.08$); MS (ES+) m/z 219.2 (M + H).

Step 3: 6,7-Dimethoxy-5-methyl-2-{methyl-[3-(1-methylpiperidin-4-yl)benzyl]amino}-1H-quinazolin-4-one: In a heavy-walled pressure tube, a mixture of methyl-[3-(1-methylpiperidin-4-yl)benzyl]amine (0.14 g, 0.64 mmol) and 2-chloro-6,7-dimethoxy-5-methyl-1H-quinazolin-4-one (0.155 g, 0.60 mmol) in 4 mL of EtOH, was heated to 120°C for 5 hours. After solvent removal, the residue was purified by flash chromatography eluting with CH_2Cl_2 :MeOH: NH_4OH (94.5:5:0.5), and the above-titled Example H-3 crystallized from CH_2Cl_2 /cyclohexane. mp $155.7\text{--}158^\circ\text{C}$; IR (KBr) ν_{max} 2933, 1651, 1590 cm^{-1} ; ^1H NMR ($\text{DMSO}-d_6$, 2.49) δ 1.59-1.71 (m, 4 H), 1.93 (m, 2 H), 2.16 (s, 3H), 2.42 (m, 1 H), 2.60 (s, 3H), 2.82 (m, 2 H), 3.01 (s, 3H), 3.62 (s, 3H), 3.84 (s, 3H), 4.78 (s, 2H), 6.63 (br s, 1H), 7.03 (d, 1H, $J = 7.44$), 7.11-7.13 (m, 2 H), 7.25 (t, 1H, $J = 7.74$), 10.81 (br s, D_2O , 1H); ^{13}C NMR ($\text{DMSO}-d_6$, 39.50) δ 13.44, 32.94, 35.12, 41.08, 46.09, 51.64, 55.44, 55.65, 59.76, 124.61, 125.32, 125.65, 128.44, 137.54, 146.43; MS (ES+) m/z 437 (M + H); Anal. ($\text{C}_{25}\text{H}_{32}\text{N}_4\text{O}_3 \cdot 0.25\text{H}_2\text{O}$) C: calcd, 68.08; found, 68.02; H: calcd, 7.43; found, 7.19; N: calcd, 12.70; found, 12.69.

Examples H-4 to H-47

Compounds of formula I wherein R, R' and R¹⁰ are methyl and Y is -CH₂- as in Table 8, were prepared following the same or similar method as described for Examples H-1 to H-3, except using an appropriately substituted benzyl amine.

TABLE 8

Ex. No	Compound Name	Mp°C	MW
H-4	6,7-dimethoxy-5-methyl-2-[methyl-(3-pyrrolidin-1-yl-benzyl)-amino]-1H-quinazolin-4-one		408.50
H-5	5,6,7-trimethoxy-2-[methyl-(3-pyrrolidin-1-yl-benzyl)-amino]-1H-quinazolin-4-one		424.50
H-6	6,7-dimethoxy-5-methyl-2-{methyl-[3-(2-methyl-4,5-dihydro-imidazol-1-yl)-benzyl]-amino}-1H-quinazolin-4-one	177-180	421.50
H-7	2-{[3-(4,4-dimethyl-4,5-dihydro-imidazol-1-yl)-benzyl]-methyl-amino}-6,7-dimethoxy-5-methyl-1H-quinazolin-4-one		435.53
H-8	6,7-Dimethoxy-5-methyl-2-{methyl-[3-(2,4,4-trimethyl-4,5-dihydro-imidazol-1-yl)-benzyl]-amino}-1H-quinazolin-4-one		449.55
H-9	6,7-Dimethoxy-5-methyl-2-{methyl-[3-(4-methyl-imidazol-1-yl)-benzyl]-amino}-1H-quinazolin-4-one		419.48
H-10	6,7-Dimethoxy-5-methyl-2-{methyl-[3-(2-methyl-imidazol-1-yl)-benzyl]-amino}-1H-quinazolin-4-one		419.48
H-11	6,7-Dimethoxy-5-methyl-2-[methyl-(3-pyrazol-1-yl-benzyl)-amino]-1H-quinazolin-4-one		405.46
H-12	6,7-Dimethoxy-5-methyl-2-{methyl-[3-(3-methyl-pyrazol-1-yl)-benzyl]-amino}-1H-quinazolin-4-one	220-223.5	419.48
H-13	2-{[3-(3,5-Dimethyl-pyrazol-1-yl)-benzyl]-methyl-amino}-6,7-dimethoxy-5-methyl-1H-quinazolin-4-one		433.51
H-14	6,7-Dimethoxy-5-methyl-2-[methyl-(3-pyrrolidin-2-yl-benzyl)-amino]-1H-quinazolin-4-one		408.50
H-15	5,6,7-Trimethoxy-2-[methyl-(3-pyrrolidin-3-yl-benzyl)-amino]-1H-quinazolin-4-one		424.50
H-16	6,7-dimethoxy-5-methyl-2-{methyl-[4-(1-methyl-4,5-dihydro-1H-imidazol-2-yl)-benzyl]-amino}-1H-quinazolin-		421.50

	4-one	
H-17	5,6,7-Trimethoxy-2-{methyl-[3-(1-methyl-4,5-dihydro-1H-imidazol-2-yl)-benzyl]-amino}-1H-quinazolin-4-one	437.50
H-18	2-{{3-(1-ethyl-4,5-dihydro-1H-imidazol-2-yl)-benzyl]-methyl-amino}-6,7-dimethoxy-5-methyl-1H-quinazolin-4-one	435.52
H-19	2-{{3-(1-Ethyl-4,5-dihydro-1H-imidazol-2-yl)-benzyl]-methyl-amino}-5,6,7-trimethoxy-1H-quinazolin-4-one	451.52
H-20	2-{{3-(1-Isopropyl-4,5-dihydro-1H-imidazol-2-yl)-benzyl]-methyl-amino}-5,6,7-trimethoxy-1H-quinazolin-4-one	465.55
H-21	2-{{4-(1-Isopropyl-4,5-dihydro-1H-imidazol-2-yl)-benzyl]-methyl-amino}-6,7-dimethoxy-5-methyl-1H-quinazolin-4-one	449.55
H-22	5-Hydroxy-2-{{3-(1-isopropyl-4,5-dihydro-1H-imidazol-2-yl)-benzyl]-methyl-amino}-6,7-dimethoxy-1H-quinazolin-4-one	451.52
H-23	2-{{4-(1-Isopropyl-4,5-dihydro-1H-imidazol-2-yl)-benzyl]-methyl-amino}-6,7-dimethoxy-5-methyl-1H-quinazolin-4-one	449.55
H-24	2-({3-[1-(2-Hydroxy-ethyl)-4,5-dihydro-1H-imidazol-2-yl]-benzyl]-methyl-amino)-6,7-dimethoxy-5-methyl-1H-quinazolin-4-one	451.52
H-25	2-({3-[1-(2-Hydroxy-ethyl)-4,5-dihydro-1H-imidazol-2-yl]-benzyl]-methyl-amino)-5,6,7-trimethoxy-1H-quinazolin-4-one	467.52
H-26	6,7-Dimethoxy-2-({3-[1-(2-methoxy-ethyl)-4,5-dihydro-1H-imidazol-2-yl]-benzyl]-methyl-amino)-5-methyl-1H-quinazolin-4-one	465.55
H-27	5,6,7-Trimethoxy-2-({3-[1-(2-methoxy-ethyl)-4,5-dihydro-1H-imidazol-2-yl]-benzyl]-methyl-amino)-1H-quinazolin-4-one	481.55
H-28	2-{{3-(1H-Imidazol-2-yl)-benzyl]-methyl-amino}-6,7-dimethoxy-5-methyl-1H-quinazolin-4-one	405.46
H-29	6,7-Dimethoxy-5-methyl-2-{methyl-[3-(1-methyl-1H-imidazol-2-yl)-benzyl]-amino}-1H-quinazolin-4-one	419.48

H-30	2-({3-[1-(2-Hydroxy-ethyl)-1H-imidazol-2-yl]-benzyl}-methyl-amino)-6,7-dimethoxy-5-methyl-1H-quinazolin-4-one	449.51
H-31	6,7-Dimethoxy-5-methyl-2-{methyl-[3-(2-oxo-pyrrolidin-1-yl)-benzyl]-amino}-1H-quinazolin-4-one	422.48
H-32	2-{{3-(1,1-Dioxo-1,6-isothiazolidin-2-yl)-benzyl}-methyl-amino}-6,7-dimethoxy-5-methyl-1H-quinazolin-4-one	458.54
H-33	2-{{3-(1,1-Dioxo-1,6-isothiazolidin-2-yl)-benzyl}-methyl-amino}-5,6,7-trimethoxy-1H-quinazolin-4-one	474.54
H-34	6,7-Dimethoxy-5-methyl-2-{methyl-[3-(2-oxo-oxazolidin-3-yl)-benzyl]-amino}-1H-quinazolin-4-one	424.45
H-35	5,6,7-Trimethoxy-2-{methyl-[3-(2-oxo-oxazolidin-3-yl)-benzyl]-amino}-1H-quinazolin-4-one	440.45
H-36	6,7-Dimethoxy-5-methyl-2-{methyl-[3-(3-methyl-2-oxo-pyrrolidin-1-yl)-benzyl]-amino}-1H-quinazolin-4-one	436.51
H-37	2-{{3-(3-Ethyl-2-oxo-pyrrolidin-1-yl)-benzyl}-methyl-amino}-6,7-dimethoxy-5-methyl-1H-quinazolin-4-one	450.54
H-38	2-{{3-(3-Ethyl-2-oxo-pyrrolidin-1-yl)-benzyl}-methyl-amino}-5,6,7-trimethoxy-1H-quinazolin-4-one	466.54
H-39	1-(3-{{(6,7-Dimethoxy-5-methyl-4-oxo-1,4-dihydro-quinazolin-2-yl)-methyl-amino}-methyl}-phenyl)-pyrrolidine-2,5-dione	436.47
H-40	1-(3-{{[Methyl-(5,6,7-trimethoxy-4-oxo-1,4-dihydro-quinazolin-2-yl)-amino]-methyl}-phenyl)-pyrrolidine-2,5-dione	452.47
H-41	5,6,7-Trimethoxy-2-{methyl-[3-(2-oxo-imidazolidin-1-yl)-benzyl]-amino}-1H-quinazolin-4-one	439.47
H-42	6,7-Dimethoxy-5-methyl-2-{methyl-[3-(3-methyl-2-oxo-imidazolidin-1-yl)-benzyl]-amino}-1H-quinazolin-4-one	437.50
H-43	5,6,7-Trimethoxy-2-{methyl-[3-(3-methyl-2-oxo-imidazolidin-1-yl)-benzyl]-amino}-1H-quinazolin-4-one	453.49
H-44	2-{{3-(3-Ethyl-2-oxo-imidazolidin-1-yl)-benzyl}-methyl-amino}-6,7-dimethoxy-5-methyl-1H-quinazolin-4-one	451.52
H-45	2-{{3-(3-Ethyl-2-oxo-imidazolidin-1-yl)-benzyl}-methyl-amino}-5,6,7-trimethoxy-1H-quinazolin-4-one	467.52
H-46	5,6,7-Trimethoxy-2-({3-[3-(2-methoxy-ethyl)-2-oxo-imid-	497.55

azolidin-1-yl]-benzyl}-methyl-amino)-1H-quinazolin-4-one

H-47 6,7-Dimethoxy-5-methyl-2-{methyl-[3-(1H-tetrazol-5-yl)-benzyl]-amino}-1H-quinazolin-4-one 407.43

Additional compounds prepared according to the procedure of Example H-1 through H-3 are shown in Table 11.

Example I-1: 6,7-Dimethoxy-5-methyl-2-[methyl-(4-pyrrolidin-1-yl-pyridin-2-ylmethyl)-amino]-1H-quinazolin-4-one

- 5 Step 1: To a 2 N solution of methylamine (6.15 mL, 12.3 mmol) in MeOH was added 4-chloro-2-chloromethylpyridine (0.2 g, 1.23 mmol). The resulting solution was allowed to stir for overnight at RT. After evaporation of the solvent, 4-chloro-2-methylaminomethylpyridine was purified by flash column chromatography. The starting material (4-chloro-2-chloromethylpyridine) was prepared according to Tamura et al.,
10 Chem. Pharm.Bull. 48:1514-1518 (2000).

- Step 2: Diisopropylethylamine (0.15 mL, 0.86 mmol) was added to a mixture of 2-chloro-5,6,7-trimethoxy-1H-quinazolin-4-one (0.2 g, 0.78 mmol) and 4-chloro-2-methylamino-methylpyridine (0.13 g, 0.86 mmol) in EtOH (8 mL). The mixture was heated at 90°C for 5 h. After cooling, the resulting white solid was filtered and washed with MeOH to
15 provide quinazolino-pyridine compound.

- Step 3: 4-chloropyridine compound from Step 2 (20 mg, 0.05 mmol) in pyrrolidine (0.5 mL) was heated at 130°C overnight in a sealed tube. The mixture was filtered, concentrated, and purified by preparative reversed-phase HPLC to afford the above Example I-1.
¹H NMR (DMSO-*d*₆) δ: 2.0 (br, 2H), 2.62 (s, 3H), 3.15 (s, 3H), 3.50 (m, 2H), 3.62 (s, 3H), 3.84 (s, 3H), 4.90 (s, 2H), 6.72 (m, 1H), 6.79 (dd, 1H), 7.54 (s, 1H), 8.17 (dd, 1H)
20 (as the TFA salt). MS (ES+) *m/z* 410 (M + H)⁺.

Examples I-2 To I-13

- Compounds I-3 through I-8 and I-13 of formula I wherein R, R' and R¹⁰ are methyl and Y is -CH₂- as in Table 9, were prepared following the same or similar method as
25 described for Example I-1, replacing the pyrrolidine in step 3 with the desired substituent. The remaining compounds were prepared by condensation of the appropriate amine and carbonyl compounds according to procedures in Abdel-Magid et al. J. Organic Chemistry 61:3849-3862 (1996).

TABLE 9

Ex. No	Compound Name	MW
I-2	5,6,7-Trimethoxy-2-[(6-{1-[(2-methoxy-ethyl)-methyl-amino]-ethyl}-pyridin-2-ylmethyl)-methyl-amino]-1H-quinazolin-4-one	471.56
I-3	2-[(4-Chloro-pyridin-2-ylmethyl)-methyl-amino]-5,6,7-trimethoxy-1H-quinazolin-4-one	390.83
I-4	5-Hydroxy-6,7-Dimethoxy-2-({4-[(2-methoxy-ethyl)-methyl-amino]-pyridin-2-ylmethyl}-methyl-amino)-5-methyl-1H-quinazolin-4-one	429.47
I-5	6,7-Dimethoxy-2-({4-[(2-methoxy-ethyl)-methyl-amino]-pyridin-2-ylmethyl}-methyl-amino)-5-methyl-1H-quinazolin-4-one	427.50
I-6	5,6,7-trimethoxy-2-({4-[(2-methoxy-ethyl)-methyl-amino]-pyridin-2-ylmethyl}-methyl-amino)-1H-quinazolin-4-one	443.50
I-7	5,6,7-Trimethoxy-2-{[4-(2-methoxy-ethoxy)-pyridin-2-ylmethyl]-methyl-amino}-1H-quinazolin-4-one	430.46
I-8	2-[(4-Ethoxy-pyridin-2-ylmethyl)-methyl-amino]-5,6,7-trimethoxy-1H-quinazolin-4-one	400.43
I-9	6,7-Dimethoxy-5-methyl-2-{methyl-[6-(1-pyrrolidin-1-yl-ethyl)-pyridin-2-ylmethyl]-amino}-1H-quinazolin-4-one	437.54
I-10	5,6,7-Trimethoxy-2-{methyl-[6-(1-pyrrolidin-1-yl-ethyl)-pyridin-2-ylmethyl]-amino}-1H-quinazolin-4-one	453.54
I-11	2-{[6-(1-Azetidin-1-yl-ethyl)-pyridin-2-ylmethyl]-methyl-amino}-6,7-dimethoxy-5-methyl-1H-quinazolin-4-one	423.51
I-12	2-{[6-(1-Azetidin-1-yl-ethyl)-pyridin-2-ylmethyl]-methyl-amino}-5,6,7-trimethoxy-1H-quinazolin-4-one	439.51
I-13	6,7-Dimethoxy-2-{[4-(2-methoxy-ethoxy)-pyridin-2-ylmethyl]-methyl-amino}-5-methyl-1H-quinazolin-4-one	414.46

Additional compounds prepared according to the procedure of Example I-1 are shown in Table 11.

Examples J-1 to J-7


- Compounds of formula I wherein R and R' are methyl as in Table 10 were prepared following the same or similar method as described above, e.g., upon coupling compounds having the desired aryl or heteroaryl groups with 2-chloro-6,7-dimethoxy-5-methyl-1H-quinazolin-4-one, or 2-chloro-5, 6,7-trimethoxy-1H-quinazolin-4-one.

TABLE 10

Ex. No		MW
J-1	N,N-Dimethyl-N'-(3-{2-[methyl-(5,6,7-trimethoxy-4-oxo-1,4-dihydro-quinazolin-2-yl)-amino]-ethyl}-phenyl)-acetamidine	453.54
J-2	5-[2-(6,7-Dimethoxy-5-methyl-4-oxo-1,4-dihydro-quinazolin-2-ylamino)-propyl]-2-methoxy-benzenesulfonamide	462.52
J-3	5-[2-(6,7-Dimethoxy-5-methyl-4-oxo-1,4-dihydro-quinazolin-2-ylamino)-propyl]-2-methoxy-N-(1-dimethylamino-ethylidene)-benzenesulfonamide	531.63
J-4	N'-(3-{2-[(6,7-Dimethoxy-5-methyl-4-oxo-1,4-dihydro-quinazolin-2-yl)-methyl-amino]-1-hydroxy-ethyl}-phenyl)-N,N-dimethyl-acetamidine	453.54
J-5	N'-(3-{2-[(6,7-Dimethoxy-5-methyl-4-oxo-1,4-dihydro-quinazolin-2-yl)-methyl-amino]-vinyl}-phenyl)-N,N-dimethyl-acetamidine	435.53
J-6	2-[[2-(3H-Imidazol-4-yl)-ethyl]-methyl-amino]-6,7-dimethoxy-5-methyl-1H-quinazolin-4-one	343.39
J-7	2-[[2-(3H-Imidazol-4-yl)-ethyl]-methyl-amino]-5,6,7-trimethoxy-1H-quinazolin-4-one	359.38
J-8	2-[(1-Benzyl-piperidin-3-yl)-methyl-amino]-6,7-dimethoxy-5-methyl-1H-quinazolin-4-one	422.53


Example K-1: *N,N*-Dimethyl-*N'*-(3-{[methyl-(6,7,8-trimethoxy-1,1-dioxo-1,4-dihydro-1*H*-6-benzo[1,2,4]thiadiazin-3-yl)-amino]-methyl}-phenyl)-acetamidine

- 5 Step 1: 6,7,8-Trimethoxy-1,1-dioxo-1,4-dihydro-2*H*-1*λ*^{6*}-benzo[1,2,4]thiadiazin-3-one



To 45 mL nitroethane was added chlorosulfonylisocyanate (3 mL, 34 mmol). The solution was cooled to -70°C. A solution of trimethoxyaniline (4.5 g, 24.5 mmol) in nitroethane (15 mL) was added over 5 minutes. The temperature of the solution rose to -64°C. The solution was removed from the ice bath and allowed to warm to -5°C while a heavy precipitate formed. In one portion, AlCl₃ (5 g, 37.6 mmol) was added. The precipitate went into solution as the temperature rose to 25°C. The resulting solution was placed in a 110°C oil bath and stirred for 30 min. The resulting mixture was cooled to RT and poured into ice water. The aqueous layer was removed from the resulting black gum, which contained some of the desired material, as well as starting material and side products. The aqueous layer was shaken with EtOAc and the layers separated. The organic layer was evaporated *in vacuo* and the residue crystallized from a small amount of EtOAc to yield 6,7,8-trimethoxy benzothiadiazine-3-one-1,1-dioxide. (3.4 g): mp 220-223°C; MS (ES+) *m/z* 289 (M + H); ¹H NMR (DMSO-*d*₆, 2.49), 3.75(s, 3H), 3.83(s, 3H), 3.89 (s, 3H), 6.58 (s, 1H), 11 (s, 1H).

Step 2: 6,7,8-trimethoxy-3-chloro-benzothiadiazine-3-1,1-dioxide



To 6,7,8-trimethoxy benzothiadiazine-3-one-1,1-dioxide (2.4 g, 8.3 mmol) was added phosphorous oxychloride (22 mL). The resulting mixture was cooled to 5°C and *N,N*-diethylaniline (2.6 mL, 20 mmol) was added. The mixture was allowed to warm to RT, and then heated at 80°C for 8 h. The solvent was removed by distillation under reduced pressure. The resulting syrup was poured into ice water. After rapid stirring the precipitate was filtered. The aqueous mother liquors were extracted with methylene chloride. The resulting organic layer was dried over Na₂SO₄, the drying agent removed by filtration and the solvent removed under reduced pressure, to afford 6,7,8-trimethoxy-3-chloro-benzothiadiazine-3-1,1-dioxide 920 mg. MS (ES+) *m/z* 307 (M+H); ¹H NMR (DMSO-*d*₆, 2.49), 3.75(s, 3H), 3.83(s, 3H), 3.89 (s, 3H), 6.58 (s, 1H), 11 (s, 1H).


Step 3: The chlorobenzothiadiazine dioxide from Step B (130 mg, 0.42 mmol) and *N,N*-dimethyl-*N'*-(3-methylaminomethyl-phenyl)-acetamidine (90 mg, 0.43 mmol) in EtOH (5 mL) were heated in a sealed tube at 100°C for 18 h. The mixture was cooled and the

suspension filtered. The desired product was found in both the filter cake and the mother liquors so these were combined and the solvent removed under reduced pressure. The resulting material was chromatographed on silica gel to yield a partially purified fraction which was then further purified by reverse phase HPLC to provide the above-

5 titled Example K-1. (M+H)⁺ 477.

Example K-2: *N'*-(3-[[[(6,7-Dimethoxy-1,1-dioxo-1,4-dihydrobenzo[1,2,4]thiadiazin-3-yl)-methyl-amino]-methyl]-phenyl)-*N,N*-dimethyl-acetamide

Step 1: 6,7-Dimethoxy-3-(methylaminomethyl-3-nitrophenyl) benzothiadiazine-3-one-1,1-dioxide



10 A solution of 3-chloro-6,7-dimethoxy-4*H*-benzo[1,2,4]thiadiazine 1,1-dioxide (404 mg, 1.5 mmol), methyl-(3-nitrobenzyl)-amine hydrochloride (400 mg, 2 mmol), DBU (0.25 mL, excess) and dimethoxyethanol (5 mL) was heated at 80°C 18 h. The solvent was removed under reduced pressure. The resulting gum was chromatographed on silica gel in

15 3 % MeOH in DCM to yield white crystals (250 mg): MS (ES⁺) *m/z* 407 (M+H).

Substituted 2*H*-benzo[1,2,4]thiadiazine-1,1-dioxides were prepared as described in WO 02/053558 A1. The nitro compound (250 mg, 0.6 mmol), 10% Pd/C (32 mg) and EtOH (60 mL) were hydrogenated on a Parr apparatus at 45 psi for 18 h. The reaction mixture was then filtered under nitrogen through a glass fiber paper and the solvent removed

20 under reduced pressure to yield the aniline as a white solid (210 mg). MS (ES⁺) *m/z* 377 (M+H).

Step 2: The aniline from Step 1 (210 mg, 0.56 mmol), dimethylacetamide dimethyl acetal (80 μ L, excess) and dimethylacetamide (2 mL) were heated at 80°C for 18 h. The solvent was removed under reduced pressure. The resulting gum was dissolved in DCM and

25 chromatographed over silica gel in 4 % MeOH in DCM to yield the above-titled Example K-2 as an off white solid.

Representative compounds in accordance with the invention are shown in Table 11, together with corresponding melting point or low resolution Mass Spectrum data.

TABLE 11

#	Systematic Name	MP°C	M+H
1	5,6,7-Trimethoxy-2-[methyl-(4-[[methyl-(3,3,3-trifluoro-propyl)-amino]-methyl]-benzyl)-amino]-1 <i>H</i> -quinazolin-4-one	97.8-101.9	
2	2-[[4-(2-Hydroxy-3-methoxy-propoxy)-benzyl]-methyl-amino]-5,6,7-trimethoxy-1 <i>H</i> -quinazolin-4-one	108.1-111.5	
3	2-[[3-(3-Hydroxy-2-hydroxymethyl-propoxy)-4-methoxy-benzyl]-methyl-amino]-5,6,7-trimethoxy-1 <i>H</i> -quinazolin-4-one	139.7-146.6 (HCl salt)	
4	1-[1-(3-[[[(6,7-Dimethoxy-5-methyl-4-oxo-1,4-dihydro-quinazolin-2-yl)-methyl-amino]-methyl]-phenyl)-ethyl]-1,3,3-trimethyl-urea	125.9-128.0	
5	[1-(3-[[[(6,7-Dimethoxy-5-methyl-4-oxo-1,4-dihydro-quinazolin-2-yl)-methyl-amino]-methyl]-phenyl)-ethyl]-methyl-carbamic acid methyl ester	117.9-120.1	
6	2-[[3-(3-Hydroxy-2-hydroxymethyl-propoxy)-4-methoxy-benzyl]-methyl-amino]-6,7-dimethoxy-5-methyl-1 <i>H</i> -quinazolin-4-one	210.8-215.4	
7	2-[[3-(2,3-Dihydroxy-propoxy)-benzyl]-methyl-amino]-6,7-dimethoxy-5-methyl-1 <i>H</i> -quinazolin-4-one	220-224 (HCl salt)	
8	5,6,7-Trimethoxy-2-[[3-((S)-4-methoxy-2-oxo-pyrrolidin-1-yl)-benzyl]-methyl-amino]-1 <i>H</i> -quinazolin-4-one	142.5-147.4	
9	(4-[[[(6,7-Dimethoxy-5-methyl-4-oxo-1,4-dihydro-quinazolin-2-yl)-methyl-amino]-methyl]-benzyl)-methyl-carbamic acid 2-methoxy-ethyl ester	159.4-160.4	
10	<i>N</i> -(4-[[[(6,7-Dimethoxy-5-methyl-4-oxo-1,4-dihydro-quinazolin-2-yl)-methyl-amino]-methyl]-benzyl)- <i>N</i> -methyl-methanesulfonamide	238-240	
11	2-[[3-(4,5-Dihydro-oxazol-2-yl)-benzyl]-methyl-amino]-5,6,7-trimethoxy-1 <i>H</i> -quinazolin-4-one	202.3-203.4	
12	5,6,7-Trimethoxy-2-{methyl-[4-(2-oxo-pyrrolidin-1-yl)-benzyl]-amino}-1 <i>H</i> -quinazolin-4-one	195.9-197.4	
13	2-[(2-Allyl-2,3-dihydro-1 <i>H</i> -isoindol-5-yl)methyl]-methyl-amino]-5,6,7-trimethoxy-1 <i>H</i> -quinazolin-4-one	133.7-136.6	
14	2-[[4-(2,3-Dihydroxy-propoxy)-benzyl]-methyl-amino]-6,7-dimethoxy-5-methyl-1 <i>H</i> -quinazolin-4-one	199.7-201.9	

15	2-[[4-(3-Hydroxy-2-hydroxymethyl-propoxy)-benzyl]-methyl-amino]-6,7-dimethoxy-5-methyl-1 <i>H</i> -quinazolin-4-one	197.5-199 (HCl salt)
16	2-[(3-{2-[(2-Hydroxy-ethyl)-methyl-amino]-ethoxy}-4-methoxy-benzyl)-methyl-amino]-6,7-dimethoxy-5-methyl-1 <i>H</i> -quinazolin-4-one	144.8-146.2
17	<i>N</i> -(3-[[[(6,7-Dimethoxy-5-methyl-4-oxo-1,4-dihydro-quinazolin-2-yl)-methyl-amino]-methyl]-benzyl])- <i>N</i> -methyl-methanesulfonamide	223.9-225.9
18	5,6,7-Trimethoxy-2-[[2-(2-methoxy-ethyl)-2,3-dihydro-1 <i>H</i> -isoindol-5-ylmethyl]-methyl-amino]-1 <i>H</i> -quinazolin-4-one	133.3-136.6
19	6,7-Dimethoxy-2-[[2-(2-methoxy-ethyl)-2,3-dihydro-1 <i>H</i> -isoindol-5-ylmethyl]-methyl-amino]-5-methyl-1 <i>H</i> -quinazolin-4-one	173.3-177.9
20	5,6,7-Trimethoxy-2-[[3-(3-methoxy-2-oxo-pyrrolidin-1-yl)-benzyl]-methyl-amino]-1 <i>H</i> -quinazolin-4-one	90.2-94.7
21	<i>N</i> -Methyl- <i>N</i> -(4-[[methyl-(5,6,7-trimethoxy-4-oxo-1,4-dihydro-quinazolin-2-yl)-amino]-methyl]-benzyl)-methanesulfonamide	148.8-151
22	2-[[3-(3-Hydroxy-2-oxo-pyrrolidin-1-yl)-benzyl]-methyl-amino]-5,6,7-trimethoxy-1 <i>H</i> -quinazolin-4-one	172.9-175.9 (HCl salt)
23	6,7-Dimethoxy-5-methyl-2-{methyl-[4-(2-methylamino-ethyl)-benzyl]-amino}-1 <i>H</i> -quinazolin-4-one	154.6-159.0
24	2-[[3-((<i>S</i>)-4-Hydroxy-2-oxo-pyrrolidin-1-yl)-benzyl]-methyl-amino]-5,6,7-trimethoxy-1 <i>H</i> -quinazolin-4-one	207.5-209.9
25	2-([4-[2-(Ethyl-methyl-amino)-ethoxy]-3-methoxy-benzyl]-methyl-amino)-6,7-dimethoxy-5-methyl-1 <i>H</i> -quinazolin-4-one	158.1-159.2
26	<i>N</i> -[1-(3-[[[(6,7-Dimethoxy-5-methyl-4-oxo-1,4-dihydro-quinazolin-2-yl)-methyl-amino]-methyl]-phenyl]-ethyl])- <i>N</i> -methyl-methanesulfonamide	130.0-133.8
27	2-[[3-((<i>S</i>)-4-Hydroxy-2-oxo-pyrrolidin-1-yl)-benzyl]-methyl-amino]-6,7-dimethoxy-5-methyl-1 <i>H</i> -quinazolin-4-one	230.8-232.5
28	2-([4-[2-(Ethyl-methyl-amino)-ethyl]-benzyl]-methyl-	172.4-173.2

	amino)-6,7-dimethoxy-5-methyl-1 <i>H</i> -quinazolin-4-one		
29	6,7-dimethoxy-2-{{3-methoxy-4-(2-methylamino-ethoxy)-benzyl}-methyl-amino}-5-methyl-1 <i>H</i> -quinazolin-4-one	168.9-169.8	
30	2-[(4-{{Ethyl-(2-hydroxy-ethyl)-amino}-methyl}-benzyl)-methyl-amino]-6,7-dimethoxy-5-methyl-1 <i>H</i> -quinazolin-4-one	142-5	
31	2-[[4-(2-Hydroxy-ethoxy)-benzyl]-methyl-amino]-6,7-dimethoxy-5-methyl-1 <i>H</i> -quinazolin-4-one	198.5-199.5	
32	2-[(4-{{Ethyl-(2-methoxy-ethyl)-amino}-methyl}-benzyl)-methyl-amino]-5,6,7-trimethoxy-1 <i>H</i> -quinazolin-4-one	120-122.4	
33	2-[(3-{{(2-Hydroxy-ethyl)-methyl-amino}-methyl}-benzyl)-methyl-amino]-6,7-dimethoxy-5-methyl-1 <i>H</i> -quinazolin-4-one	123.8-125.6	
34	5,6,7-Trimethoxy-2-{{3-(3-methoxy-pyrrolidin-1-ylmethyl)-benzyl}-methyl-amino}-1 <i>H</i> -quinazolin-4-one	89.6-90.2	
35	2-({4-[(Ethyl-methyl-amino)-methyl]-benzyl}-methyl-amino)-5,6,7-trimethoxy-1 <i>H</i> -quinazolin-4-one	172.3-174.4	427
36	6,7-Dimethoxy-2-({4-[(2-methoxy-ethylamino)-methyl]-benzyl}-methyl-amino)-5-methyl-1 <i>H</i> -quinazolin-4-one	265-268	(HCl salt)
37	6,7-Dimethoxy-2-{{3-(3-methoxy-pyrrolidin-1-ylmethyl)-benzyl}-methyl-amino}-5-methyl-1 <i>H</i> -quinazolin-4-one	151.2-152.9	
38	6,7-Dimethoxy-5-methyl-2-[methyl-(2-methyl-2,3-dihydro-1 <i>H</i> -isoindol-5-ylmethyl)-amino]-1 <i>H</i> -quinazolin-4-one	214.9-224.8	
39	2-{{3-(4-Hydroxy-piperidin-1-ylmethyl)-benzyl}-methyl-amino}-5,6,7-trimethoxy-1 <i>H</i> -quinazolin-4-one	97.5-104.7	
40	2-{{3-(4-Hydroxy-piperidin-1-ylmethyl)-benzyl}-methyl-amino}-6,7-dimethoxy-5-methyl-1 <i>H</i> -quinazolin-4-one	159.2-159.6	
41	2-[[4-(3-Hydroxy-pyrrolidin-1-ylmethyl)-benzyl]-methyl-amino]-6,7-dimethoxy-5-methyl-1 <i>H</i> -quinazolin-4-one	166-169	
42	2-[(4-Aminomethyl-benzyl)-methyl-amino]-6,7-dimethoxy-5-methyl-1 <i>H</i> -quinazolin-4-one	248.2-249.4	(HCl salt)
43	2-[(3-Ethylaminomethyl-benzyl)-methyl-amino]-5,6,7-trimethoxy-1 <i>H</i> -quinazolin-4-one	133.9-134.7	
44	2-{{4-(3-Dimethylamino-propoxy)-benzyl}-methyl-amino}-6,7-dimethoxy-5-methyl-1 <i>H</i> -quinazolin-4-one	191-192.2	(HCl salt)

45	5,6,7-Trimethoxy-2-[methyl-(2-methyl-2,3-dihydro-1 <i>H</i> -isindol-5-ylmethyl)-amino]-1 <i>H</i> -quinazolin-4-one	165.5-166.5
46	5,6,7-Trimethoxy-2-[methyl-(3-pyrrolidin-1-ylmethyl-benzyl)-amino]-1 <i>H</i> -quinazolin-4-one	127.0-129.9
47	2-[(3-Azetidin-1-ylmethyl-benzyl)-methyl-amino]-5,6,7-trimethoxy-1 <i>H</i> -quinazolin-4-one	131.3-133.1
48	2-[[4-(3-Dimethylamino-propoxy)-benzyl]-methyl-amino]-5,6,7-trimethoxy-1 <i>H</i> -quinazolin-4-one	108-111
49	2-[[3-(3-Hydroxy-pyrrolidin-1-ylmethyl)-benzyl]-methyl-amino]-6,7-dimethoxy-5-methyl-1 <i>H</i> -quinazolin-4-one	148.0-149.8
50	2-[[3-(1,3-Dimethyl-imidazolidin-2-ylideneamino)-benzyl]-methyl-amino]-6,7-dimethoxy-5-methyl-1 <i>H</i> -quinazolin-4-one	200.2-204.4
51	2-[[3-(3-Hydroxy-pyrrolidin-1-ylmethyl)-benzyl]-methyl-amino]-5,6,7-trimethoxy-1 <i>H</i> -quinazolin-4-one	174.3-176.4
52	<i>N</i> -[1-(3-[[[(6,7-Dimethoxy-5-methyl-4-oxo-1,4-dihydro-quinazolin-2-yl)-methyl-amino]-methyl]-phenyl)-ethyl]- <i>N</i> -methyl-2-methylamino-acetamide	118.1-120.8
53	<i>N'</i> -(3-[[[(5-Cyano-6,7-dimethoxy-4-oxo-1,4-dihydro-quinazolin-2-yl)-methyl-amino]-methyl]-phenyl]- <i>N,N</i> -dimethyl-acetamidine	222.1-226.9
54	6,7-Dimethoxy-2-[(3-{1-[(2-methoxy-ethyl)-methyl-amino]-ethyl}-benzyl)-methyl-amino]-5-methyl-1 <i>H</i> -quinazolin-4-one	114.5-116.9
55	6,7-Dimethoxy-5-methyl-2-[methyl-(4-methylamino-methyl-benzyl)-amino]-1 <i>H</i> -quinazolin-4-one	264-268 (HCl salt)
56	2-[(3-Dimethylaminomethyl-benzyl)-methyl-amino]-6,7-dimethoxy-5-methyl-1 <i>H</i> -quinazolin-4-one	174.9-176.0
57	2-[(3-Dimethylaminomethyl-benzyl)-methyl-amino]-5,6,7-trimethoxy-1 <i>H</i> -quinazolin-4-one	110.0-113.1
58	1,5-Dimethyl-3-(3-[[methyl-(5,6,7-trimethoxy-4-oxo-1,4-dihydro-quinazolin-2-yl)-amino]-methyl]-phenyl)-imidazolidine-2,4-dione	179.1-179.9
59	1-Methyl-3-(3-[[methyl-(5,6,7-trimethoxy-4-oxo-1,4-dihydro-quinazolin-2-yl)-amino]-methyl]-phenyl)-imidazolidine-2,4-dione	130.0-137.0

60	3-(3-{[(6,7-Dimethoxy-5-methyl-4-oxo-1,4-dihydro-quinazolin-2-yl)-methyl-amino]-methyl}-phenyl)-1-methyl-imidazolidine-2,4-dione	209.3-210.8
61	2-[[3-(1-Dimethylamino-ethyl)-benzyl]-methyl-amino]-6,7-dimethoxy-5-methyl-1 <i>H</i> -quinazolin-4-one	137.0-140.1
62	2-[[3-(1-Dimethylamino-ethyl)-benzyl]-methyl-amino]-5,6,7-trimethoxy-1 <i>H</i> -quinazolin-4-one	122.6-125.5
63	5,6,7-Trimethoxy-2-{methyl-[3-(1-methylamino-ethyl)-benzyl]-amino}-1 <i>H</i> -quinazolin-4-one	131.8-134.4
64	6,7-Dimethoxy-5-methyl-2-{methyl-[3-(1-methylamino-ethyl)-benzyl]-amino}-1 <i>H</i> -quinazolin-4-one	161.6-164.1
65	1-(3-{[(6,7-Dimethoxy-5-methyl-4-oxo-1,4-dihydro-quinazolin-2-yl)-methyl-amino]-methyl}-phenyl)-3-methyl-pyrrolidine-2,5-dione	217.8-219.2
66	2-[(3-Cyclopropylaminomethyl-benzyl)-methyl-amino]-6,7-dimethoxy-5-methyl-1 <i>H</i> -quinazolin-4-one	144.5-147.6
67	2-[[3-(Isopropylamino-methyl)-benzyl]-methyl-amino]-6,7-dimethoxy-5-methyl-1 <i>H</i> -quinazolin-4-one	143.3-144.4
68	6,7-Dimethoxy-5-methyl-2-(methyl-[3-[(methyl-propyl-amino)-methyl]-benzyl]-amino)-1 <i>H</i> -quinazolin-4-one	128.1-129.1
69	6,7-Dimethoxy-5-methyl-2-(3-methylaminomethyl-benzylamino)-1 <i>H</i> -quinazolin-4-one	142.5-142.9
70	<i>N'</i> -(3-{[(6,7-Dimethoxy-4-oxo-1,4-dihydro-quinazolin-2-yl)-methyl-amino]-methyl}-phenyl)- <i>N,N</i> -dimethyl-acetamide	221.8-223
71	2-({3-[(Ethyl-methyl-amino)-methyl]-benzyl}-methyl-amino)-6,7-dimethoxy-5-methyl-1 <i>H</i> -quinazolin-4-one	146.7-150.8
72	<i>N'</i> -{3-[(6,7-Dimethoxy-5-methyl-4-oxo-1,4-dihydro-quinazolin-2-ylamino)-methyl]-phenyl}- <i>N,N</i> -dimethyl-acetamide	173-175
73	5,6,7-Trimethoxy-2-[4-(4-methoxy-pyridin-2-yl)-piperazin-1-yl]-1 <i>H</i> -quinazolin-4-one	229-231
74	<i>N'</i> -(3-{[(7-Chloro-6-methoxy-4-oxo-1,4-dihydro-quinazolin-2-yl)-methyl-amino]-methyl}-phenyl)- <i>N,N</i> -dimethyl-acetamide	185.4-191.9
75	5,6,7-Trimethoxy-2-{methyl-[3-(3-methyl-2-oxo-	156.4-162.0

	pyrrolidin-1-yl)-benzyl]-amino}-1 <i>H</i> -quinazolin-4-one		
76	6,7-Dimethoxy-5-methyl-2-{methyl-[3-(2-oxo-imidazolidin-1-yl)-benzyl]-amino}-1 <i>H</i> -quinazolin-4-one	265.0-266.0	
77	6,7-Dimethoxy-2-({3-[(2-methoxy-ethylamino)-methyl]-benzyl}-methyl-amino)-5-methyl-1 <i>H</i> -quinazolin-4-one	150.0-152.1	
78	5,6,7-Trimethoxy-2-{methyl-[3-(1-methyl-piperidin-4-yl)-benzyl]-amino}-1 <i>H</i> -quinazolin-4-one	181.5-189.4	
79	6,7-Dimethoxy-2-[methyl-(3-pyrrolidin-1-yl-benzyl)-amino]-1 <i>H</i> -quinazolin-4-one	220-223	
80	2-[Benzyl-(1-benzyl-piperidin-4-yl)-amino]-6,7-dimethoxy-1 <i>H</i> -quinazolin-4-one		485
81	6,7-Dimethoxy-2-[methyl-(6-methyl-pyridin-2-ylmethyl)-amino]-1 <i>H</i> -quinazolin-4-one		341
82	2-[[2-(1 <i>H</i> -Indol-3-yl)-ethyl]-methyl-amino]-6,7-dimethoxy-1 <i>H</i> -quinazolin-4-one		379
83	6,7-Dimethoxy-2-{methyl-[3-(4-methyl-piperazin-1-yl)-benzyl]-amino}-1 <i>H</i> -quinazolin-4-one		424
84	6,7-Dimethoxy-2-[2-(4-methyl-piperazin-1-yl)-benzylamino]-1 <i>H</i> -quinazolin-4-one		
85	2-(Benzyl-isopropyl-amino)-6,7-dimethoxy-1 <i>H</i> -quinazolin-4-one	168-170.5	354
86	<i>N'</i> -(3-[[[(6,7-Dimethoxy-4-oxo-1,4-dihydro-quinazolin-2-yl)-methyl-amino]-methyl]-phenyl]- <i>N,N</i> -dimethyl-formamidine	242-245 (HCl salt)	395.4 <i>M</i> ⁺
87	2-[[3-(4,5-Dihydro-3 <i>H</i> -pyrrol-2-ylamino)-benzyl]-methyl-amino]-6,7-dimethoxy-1 <i>H</i> -quinazolin-4-one	211-14	407.5 <i>M</i> ⁺
88	<i>N</i> -(3-[[[(6,7-Dimethoxy-4-oxo-1,4-dihydro-quinazolin-2-yl)-methyl-amino]-methyl]-phenyl]-acetamidine	263-66 (HBr salt)	381.4 <i>M</i> ⁺
89	2-[[3-(4,5-Dihydro-1 <i>H</i> -imidazol-2-ylamino)-benzyl]-methyl-amino]-6,7-dimethoxy-1 <i>H</i> -quinazolin-4-one	163.8-176.9	409
90	2-[[3-(4,4-Dimethyl-4,5-dihydro-imidazol-1-yl)-benzyl]-methyl-amino]-6,7-dimethoxy-1 <i>H</i> -quinazolin-4-one	240-244	422
91	<i>N'</i> -(3-{2-[[[(6,7-Dimethoxy-4-oxo-1,4-dihydro-quinazolin-2-yl)-methyl-amino]-ethyl]-phenyl]- <i>N,N</i> -dimethyl-acetamidine	135-138.1	423.5 <i>M</i> ⁺
92	6,7-Dimethoxy-2-(methyl-{2-[3-(pyrimidin-2-ylamino)-		433

	phenyl}-ethyl}-amino)-1 <i>H</i> -quinazolin-4-one		
93	2-[(2-{3-[(1 <i>H</i> -Imidazol-2-ylmethyl)-amino]-phenyl}-ethyl)-methyl-amino]-6,7-dimethoxy-1 <i>H</i> -quinazolin-4-one		435
94	2-[[2-(3,4-Dimethoxy-phenyl)-ethyl]-methyl-amino]-6,7-dimethoxy-1 <i>H</i> -quinazolin-4-one		400
95	2-[(3-Amino-benzyl)-methyl-amino]-6,7-dimethoxy-5-methyl-1 <i>H</i> -quinazolin-4-one	268-269.5	
96	3-[[[(6,7-Dimethoxy-5-methyl-4-oxo-1,4-dihydro-quinazolin-2-yl)-methyl-amino]-methyl]-benzonitrile	279.9-281.5	365
		(TFA salt)	
97	2-[Benzyl-(2-dimethylamino-ethyl)-amino]-6,7-dimethoxy-5-methyl-1 <i>H</i> -quinazolin-4-one		397
98	2-[(3-Bromo-benzyl)-methyl-amino]-6,7-dimethoxy-5-methyl-1 <i>H</i> -quinazolin-4-one	232.9-234	419
99	2-[(3-Bromo-benzyl)-methyl-amino]-5,6,7-trimethoxy-1 <i>H</i> -quinazolin-4-one		435
100	2-[(2-Fluoro-benzyl)-(2-hydroxy-ethyl)-amino]-6,7-dimethoxy-5-methyl-1 <i>H</i> -quinazolin-4-one		388
101	3-[[Methyl-(5,6,7-trimethoxy-4-oxo-1,4-dihydro-quinazolin-2-yl)-amino]-methyl]-benzonitrile		381
102	6,7-Dimethoxy-5-methyl-2-[methyl-(3-methylamino-benzyl)-amino]-1 <i>H</i> -quinazolin-4-one	195.9-197.9	368.4
			M ⁺
103	2-[(3-Amino-benzyl)-methyl-amino]-5,6,7-trimethoxy-1 <i>H</i> -quinazolin-4-one	225-226.5	370.4
			M ⁺
104	2-[(2-Amino-benzyl)-methyl-amino]-5,6,7-trimethoxy-1 <i>H</i> -quinazolin-4-one	>300	370.4
		(decomp)	M ⁺
		(HCl Salt)	
105	6,7-Dimethoxy-5-methyl-2-[methyl-(3-methylamino-methyl-benzyl)-amino]-1 <i>H</i> -quinazolin-4-one	179.0-181.3	383
106	2-[(3-Aminomethyl-benzyl)-methyl-amino]-6,7-dimethoxy-5-methyl-1 <i>H</i> -quinazolin-4-one		369
107	3-[[Methyl-(5,6,7-trimethoxy-4-oxo-1,4-dihydro-quinazolin-2-yl)-amino]-methyl]-phenyl)-acetonitrile	149.5-153.3	395
108	2-[(3-Hydroxymethyl-benzyl)-methyl-amino]-5,6,7-trimethoxy-1 <i>H</i> -quinazolin-4-one		386
109	5-Hydroxy-2-[(3-hydroxymethyl-benzyl)-methyl-amino]-6,7-dimethoxy-1 <i>H</i> -quinazolin-4-one		372

110	5,6,7-Trimethoxy-2-[methyl-(3-morpholin-4-ylmethyl-benzyl)-amino]-1 <i>H</i> -quinazolin-4-one	160.0-162.1	455
111	5,6,7-Trimethoxy-2-[methyl-(3-methylaminomethyl-benzyl)-amino]-1 <i>H</i> -quinazolin-4-one	124.9-127.6	398.5 M ⁺
112	2-[Ethyl-(3-methylaminomethyl-benzyl)-amino]-6,7-dimethoxy-5-methyl-1 <i>H</i> -quinazolin-4-one	159.3-164.1	396.5 M ⁺
113	2-[(3-Ethylaminomethyl-benzyl)-methyl-amino]-6,7-dimethoxy-5-methyl-1 <i>H</i> -quinazolin-4-one	158.2-158.9	396.5 M ⁺
114	2-({3-[(Ethyl-methyl-amino)-methyl]-benzyl}-methyl-amino)-5,6,7-trimethoxy-1 <i>H</i> -quinazolin-4-one	106.5-108.9	426.5 M ⁺
115	6,7-Dimethoxy-5-methyl-2-(methyl-{3-[(2,2,2-trifluoroethylamino)-methyl]-benzyl}-amino)-1 <i>H</i> -quinazolin-4-one	180.4-181.5	450.5 M ⁺
116	2-[(3-{[(2-Hydroxy-ethyl)-methyl-amino]-methyl}-benzyl)-methyl-amino]-5,6,7-trimethoxy-1 <i>H</i> -quinazolin-4-one	169.0-171.9	443 (HCl Salt)
117	5,6,7-Trimethoxy-2-[(3-{[(2-methoxy-ethyl)-methyl-amino]-methyl}-benzyl)-methyl-amino]-1 <i>H</i> -quinazolin-4-one	162.9-164.5	457 (bishydrochloride salt)
118	2-[(3-{[Ethyl-(2-methoxy-ethyl)-amino]-methyl}-benzyl)-methyl-amino]-5,6,7-trimethoxy-1 <i>H</i> -quinazolin-4-one	164.5-166.0	470.6 (HCl Salt) M ⁺
119	<i>N</i> -(3-{[(6,7-Dimethoxy-5-methyl-4-oxo-1,4-dihydroquinazolin-2-yl)-methyl-amino]-methyl}-benzyl)-2-methoxy- <i>N</i> -methyl-acetamide	150.0-151.2	454.5 M ⁺
120	2-Methoxy- <i>N</i> -methyl- <i>N</i> -(3-{[methyl-(5,6,7-trimethoxy-4-oxo-1,4-dihydroquinazolin-2-yl)-amino]-methyl}-benzyl)-acetamide	115-117	470.5 M ⁺
121	2-[(3-{[Ethyl-(2-methoxy-ethyl)-amino]-methyl}-benzyl)-methyl-amino]-6,7-dimethoxy-5-methyl-1 <i>H</i> -quinazolin-4-one	121.8-122.3	454.6 M ⁺
122	6,7-Dimethoxy-5-methyl-2-[methyl-(3-pyrrolidin-1-ylmethyl-benzyl)-amino]-1 <i>H</i> -quinazolin-4-one	163.9-164.5	422.5 M ⁺
123	5-Hydroxy-6,7-dimethoxy-2-{methyl-[3-(1-methyl-4,5-dihydro-1 <i>H</i> -imidazol-2-ylmethyl)-benzyl]-amino}-1 <i>H</i> -quinazolin-4-one		438
124	5,6,7-Trimethoxy-2-{methyl-[3-(1-methyl-4,5-dihydro-1 <i>H</i> -imidazol-2-ylmethyl)-benzyl]-amino}-1 <i>H</i> -quinazolin-4-		452

	one		
125	5-Hydroxy-2-[[3-(1-isopropyl-4,5-dihydro-1 <i>H</i> -imidazol-2-ylmethyl)-benzyl]-methyl-amino]-6,7-dimethoxy-1 <i>H</i> -quinazolin-4-one		466
126	2-[[3-(1-Isopropyl-4,5-dihydro-1 <i>H</i> -imidazol-2-ylmethyl)-benzyl]-methyl-amino]-5,6,7-trimethoxy-1 <i>H</i> -quinazolin-4-one		480
127	5,6,7-Trimethoxy-2-{methyl-[3-(1-methyl-piperidin-4-ylmethyl)-benzyl]-amino}-1 <i>H</i> -quinazolin-4-one	175-195 (HCl Salt)	466.6 M ⁺
128	2-[(3-Methanesulfonyl-benzyl)-methyl-amino]-6,7-dimethoxy-5-methyl-1 <i>H</i> -quinazolin-4-one	269.5-271.5	417.5 M ⁺
129	2-[(3-Methanesulfonyl-benzyl)-methyl-amino]-5,6,7-trimethoxy-1 <i>H</i> -quinazolin-4-one	231.5-232.9	433.5 M ⁺
130	<i>N,N</i> -Dimethyl-3-[[methyl-(5,6,7-trimethoxy-4-oxo-1,4-dihydro-quinazolin-2-yl)-amino]-methyl]-benzene-sulfonamide	199.9 to 204.0	462.5 M ⁺
131	3-[[[(6,7-Dimethoxy-5-methyl-4-oxo-1,4-dihydro-quinazolin-2-yl)-methyl-amino]-methyl]- <i>N</i> -(2-methoxy-ethyl)-benzenesulfonamide	163.0-164.9	476.6 M ⁺
132	<i>N</i> -(2-Methoxy-ethyl)-3-[[methyl-(5,6,7-trimethoxy-4-oxo-1,4-dihydro-quinazolin-2-yl)-amino]-methyl]-benzenesulfonamide	176.9-179.1	492.6 M ⁺
133	3-[[[(6,7-Dimethoxy-5-methyl-4-oxo-1,4-dihydro-quinazolin-2-yl)-methyl-amino]-methyl]- <i>N</i> -(3-methoxy-propyl)-benzenesulfonamide	169.0-171.9	490.6 M ⁺
134	3-[[[(6,7-Dimethoxy-5-methyl-4-oxo-1,4-dihydro-quinazolin-2-yl)-methyl-amino]-methyl]-benzoic acid	283.1-286.2	383.4 M ⁺
135	3-[[Methyl-(5,6,7-trimethoxy-4-oxo-1,4-dihydro-quinazolin-2-yl)-amino]-methyl]-benzoic acid		400
136	3-[[[(6,7-Dimethoxy-5-methyl-4-oxo-1,4-dihydro-quinazolin-2-yl)-methyl-amino]-methyl]- <i>N</i> -methyl-benzamide	243.3-245.1	396.4 M ⁺
137	<i>N</i> -Methyl-3-[[methyl-(5,6,7-trimethoxy-4-oxo-1,4-dihydro-quinazolin-2-yl)-amino]-methyl]-benzamide	217.9-219.1	412.4 M ⁺
138	<i>N</i> -(2-Dimethylamino-ethyl)-3-[[methyl-(5,6,7-trimethoxy-4-oxo-1,4-dihydro-quinazolin-2-yl)-amino]-methyl]-benzamide		470

139	<i>N</i> -(2-Dimethylamino-ethyl)- <i>N</i> -methyl-3-[[methyl-(5,6,7-trimethoxy-4-oxo-1,4-dihydro-quinazolin-2-yl)-amino]-methyl]-benzamide	484
140	<i>N</i> -(3-Dimethylamino-propyl)- <i>N</i> -methyl-3-[[methyl-(5,6,7-trimethoxy-4-oxo-1,4-dihydro-quinazolin-2-yl)-amino]-methyl]-benzamide	498
141	<i>N</i> -(2-Hydroxy-ethyl)-3-[[methyl-(5,6,7-trimethoxy-4-oxo-1,4-dihydro-quinazolin-2-yl)-amino]-methyl]-benzamide	443
142	<i>N</i> -(2-Hydroxy-propyl)-3-[[methyl-(5,6,7-trimethoxy-4-oxo-1,4-dihydro-quinazolin-2-yl)-amino]-methyl]-benzamide	457
143	<i>N</i> -(2-Hydroxy-ethyl)- <i>N</i> -methyl-3-[[methyl-(5,6,7-trimethoxy-4-oxo-1,4-dihydro-quinazolin-2-yl)-amino]-methyl]-benzamide	457
144	<i>N</i> -(2-Methoxy-ethyl)-3-[[methyl-(5,6,7-trimethoxy-4-oxo-1,4-dihydro-quinazolin-2-yl)-amino]-methyl]-benzamide	457
145	<i>N</i> -(3-Methoxy-propyl)-3-[[methyl-(5,6,7-trimethoxy-4-oxo-1,4-dihydro-quinazolin-2-yl)-amino]-methyl]-benzamide	471
146	<i>N</i> -Cyanomethyl-3-[[methyl-(5,6,7-trimethoxy-4-oxo-1,4-dihydro-quinazolin-2-yl)-amino]-methyl]-benzamide	438
147	3-[[Methyl-(5,6,7-trimethoxy-4-oxo-1,4-dihydro-quinazolin-2-yl)-amino]-methyl]- <i>N</i> -(tetrahydro-furan-2-ylmethyl)-benzamide	483
148	<i>N</i> -Furan-2-ylmethyl-3-[[methyl-(5,6,7-trimethoxy-4-oxo-1,4-dihydro-quinazolin-2-yl)-amino]-methyl]-benzamide	479
149	3-[[Methyl-(5,6,7-trimethoxy-4-oxo-1,4-dihydro-quinazolin-2-yl)-amino]-methyl]- <i>N</i> -(2-morpholin-4-yl-ethyl)-benzamide	512
150	2-[[3-(4-Hydroxymethyl-piperidine-1-carbonyl)-benzyl]-methyl-amino]-5,6,7-trimethoxy-1 <i>H</i> -quinazolin-4-one	497
151	2-[[3-(2-Hydroxy-ethoxy)-benzyl]-methyl-amino]-5,6,7-trimethoxy-1 <i>H</i> -quinazolin-4-one	146.0-147.5 416
152	2-[(3-Methanesulfinylmethoxy-benzyl)-methyl-amino]-6,7-dimethoxy-5-methyl-1 <i>H</i> -quinazolin-4-one	191.0-193.5 432
153	6,7-Dimethoxy-5-methyl-2-{methyl-[3-(pyrimidin-2-	224.0-227.8 434

	yl oxy)-benzyl]-amino}-1 <i>H</i> -quinazolin-4-one		
154	2-[(3-Hydroxy-benzyl)-methyl-amino]-6,7-dimethoxy-5-methyl-1 <i>H</i> -quinazolin-4-one	252.9-254.35	355.4 M+
155	2-[(3-Hydroxy-benzyl)-methyl-amino]-5,6,7-trimethoxy-1 <i>H</i> -quinazolin-4-one	212.3-215.1	371.4 M+
156	2-[(3-Methanesulfonylmethoxy-benzyl)-methyl-amino]-5,6,7-trimethoxy-1 <i>H</i> -quinazolin-4-one	183.9-185.4	463.5 M+
157	(3-{[(6,7-Dimethoxy-5-methyl-4-oxo-1,4-dihydro-quinazolin-2-yl)-methyl-amino]-methyl}-phenoxy)-acetic acid	>300	413.4 M+
158	(3-{[Methyl-(5,6,7-trimethoxy-4-oxo-1,4-dihydro-quinazolin-2-yl)-amino]-methyl}-phenoxy)-acetic acid	>300	429.4 M+
159	2-[[3-(2-Hydroxy-ethoxy)-benzyl]-methyl-amino]-6,7-dimethoxy-5-methyl-1 <i>H</i> -quinazolin-4-one	180.5-183.1	399.4 M+
160	6,7-Dimethoxy-2-[[3-(2-methoxy-ethoxy)-benzyl]-methyl-amino]-5-methyl-1 <i>H</i> -quinazolin-4-one	176.6-177.9	413.5 M+
161	5,6,7-Trimethoxy-2-[[3-(2-methoxy-ethoxy)-benzyl]-methyl-amino]-1 <i>H</i> -quinazolin-4-one	144.8-145.9	429.5 M+
162	6,7-Dimethoxy-2-[[3-(3-methoxy-propoxy)-benzyl]-methyl-amino]-5-methyl-1 <i>H</i> -quinazolin-4-one	150.1-152.0	427.5 M+
163	5,6,7-Trimethoxy-2-[[3-(3-methoxy-propoxy)-benzyl]-methyl-amino]-1 <i>H</i> -quinazolin-4-one	147.5-149.1	443.5 M+
164	2-[[3-(2-Dimethylamino-ethoxy)-benzyl]-methyl-amino]-6,7-dimethoxy-5-methyl-1 <i>H</i> -quinazolin-4-one	137.9-141.9	426.5 M+
165	2-[[3-(2-Ethylamino-ethoxy)-benzyl]-methyl-amino]-6,7-dimethoxy-5-methyl-1 <i>H</i> -quinazolin-4-one	264.6-266.5	426.5 M+
166	5,6,7-Trimethoxy-2-{methyl-[3-(pyrimidin-2-yloxy)-benzyl]-amino}-1 <i>H</i> -quinazolin-4-one	191-192.9	449.5 M+
167	<i>N'</i> -(3-{[(6,7-Dimethoxy-5-methyl-4-oxo-1,4-dihydro-quinazolin-2-yl)-methyl-amino]-methyl}-phenyl)- <i>N,N</i> -dimethyl-acetamide	174.4-174.8	424
168	<i>N</i> -(3-{[(5-Isopropyl-6,7-dimethoxy-4-oxo-1,4-dihydro-quinazolin-2-yl)-methyl-amino]-methyl}-phenyl)-methanesulfonamide	215-17	
169	<i>N'</i> -(3-{[(6,7-dimethoxy-5-methyl-4-oxo-1,4-dihydro-quinazolin-2-yl)-methyl-amino]-methyl}-phenyl)- <i>N,N</i> -dimethylsulfamide	195.3-199.0	

170	<i>N</i> -(3-{[(6,7-dimethoxy-5-methyl-4-oxo-1,4-dihydro-quinazolin-2-yl)-methyl-amino]-methyl}-phenyl)- <i>N,N,N'</i> -trimethylsulfamide	160.0-161.7	
171	<i>N,N</i> -Dimethyl- <i>N'</i> -(3-{[methyl-(5,6,7-trimethoxy-4-oxo-1,4-dihydro-quinazolin-2-yl)-amino]-methyl}-phenyl)-sulfamide	176.5-179.7	
172	<i>N,N,N'</i> -Trimethyl- <i>N'</i> -(3-{[methyl-(5,6,7-trimethoxy-4-oxo-1,4-dihydro-quinazolin-2-yl)-amino]-methyl}-phenyl)-sulfamide	170.3-172.3	
173	<i>N</i> -(3-{[(6,7-Dimethoxy-5-methyl-4-oxo-1,4-dihydro-quinazolin-2-yl)-methyl-amino]-methyl}-phenyl)-acetamide	250-2	
174	<i>N</i> -(3-{[Methyl-(5,6,7-trimethoxy-4-oxo-1,4-dihydro-quinazolin-2-yl)-amino]-methyl}-phenyl)-acetamide	166.5-172.9	
175	<i>N</i> -(4-Chloro-3-{[methyl-(5,6,7-trimethoxy-4-oxo-1,4-dihydro-quinazolin-2-yl)-amino]-methyl}-phenyl)-acetamide		447, M+
176	1,1-Dimethyl-3-(3-{[methyl-(5,6,7-trimethoxy-4-oxo-1,4-dihydro-quinazolin-2-yl)-amino]-methyl}-phenyl)-urea		442
177	(3-{[Methyl-(5,6,7-trimethoxy-4-oxo-1,4-dihydro-quinazolin-2-yl)-amino]-methyl}-phenyl)-carbamic acid methyl ester	230.9-232.1	
178	(3-{[(6,7-Dimethoxy-5-methyl-4-oxo-1,4-dihydro-quinazolin-2-yl)-methyl-amino]-methyl}-phenyl)-methyl-carbamic acid methyl ester	191.9-194.1	
179	<i>N</i> -(3-{[(6,7-Dimethoxy-5-methyl-4-oxo-1,4-dihydro-quinazolin-2-yl)-methyl-amino]-methyl}-phenyl)- <i>N</i> -methyl-2-oxo-propionamide	198.9-201.9	
180	3-(3-{[(6,7-Dimethoxy-5-methyl-4-oxo-1,4-dihydro-quinazolin-2-yl)-methyl-amino]-methyl}-phenyl)-1,1-dimethyl-urea	131.5-168.0	
181	<i>N</i> -Methoxy- <i>N'</i> -(3-{[methyl-(5,6,7-trimethoxy-4-oxo-1,4-dihydro-quinazolin-2-yl)-amino]-methyl}-phenyl)-urea	165-7	
182	2-{[3-(2-Hydroxy-ethylamino)-benzyl]-methyl-amino}-5,6,7-trimethoxy-1 <i>H</i> -quinazolin-4-one	60.5-61.4	
183	5-Hydroxy-2-({3-[(3-hydroxy-propyl)-methyl-amino]-		429

	benzyl}-methyl-amino)-6,7-dimethoxy-1 <i>H</i> -quinazolin-4-one	
184	2-([3-(3-Hydroxy-propylamino)-benzyl]-methyl-amino)-5,6,7-trimethoxy-1 <i>H</i> -quinazolin-4-one	47-48.5
185	5,6,7-Trimethoxy-2-([3-(2-methoxy-ethylamino)-benzyl]-methyl-amino)-1 <i>H</i> -quinazolin-4-one	178.9-179.9 (HCl Salt)
186	5,6,7-Trimethoxy-2-([3-[(2-methoxy-ethyl)-methyl-amino]-benzyl]-methyl-amino)-1 <i>H</i> -quinazolin-4-one	179.3-181.2 (HCl Salt)
187	2-([3-[(3-Hydroxy-propyl)-methyl-amino]-benzyl]-methyl-amino)-5,6,7-trimethoxy-1 <i>H</i> -quinazolin-4-one	156-157.2 (HCl Salt)
188	5,6,7-Trimethoxy-2-([3-(3-methoxy-propylamino)-benzyl]-methyl-amino)-1 <i>H</i> -quinazolin-4-one	207.0-209.1 (HCl Salt)
189	5,6,7-Trimethoxy-2-([3-[(3-methoxy-propyl)-methyl-amino]-benzyl]-methyl-amino)-1 <i>H</i> -quinazolin-4-one bi	166.7-168.1 (HCl Salt)
190	2-([3-[Bis-(2-hydroxy-ethyl)-amino]-benzyl]-methyl-amino)-5,6,7-trimethoxy-1 <i>H</i> -quinazolin-4-one	138.1-141
191	<i>N</i> -[2-(3-[1-[(6,7-Dimethoxy-5-methyl-4-oxo-1,4-dihydro-quinazolin-2-yl)-methyl-amino]-ethyl]-phenylamino)-ethyl]-acetamide	220.6-222.4
192	<i>N</i> -(3-[(6,7-Dimethoxy-5-methyl-4-oxo-1,4-dihydro-quinazolin-2-yl)-methyl-amino]-methyl)-phenyl)-acetamidine	155-58
193	<i>N</i> -(3-[(Methyl-(5,6,7-trimethoxy-4-oxo-1,4-dihydro-quinazolin-2-yl)-amino)-methyl]-phenyl)-acetamidine	412
194	<i>N</i> -(3-[(6,7-Dimethoxy-5-methyl-4-oxo-1,4-dihydro-quinazolin-2-yl)-methyl-amino]-methyl)-phenyl)- <i>N,N'</i> -dimethyl-acetamidine	181.2-186.6
195	<i>N</i> -(3-[(6,7-Dimethoxy-5-methyl-4-oxo-1,4-dihydro-quinazolin-2-yl)-methyl-amino]-methyl)-phenyl)- <i>N'</i> -methyl-acetamidine	233.1-233.9
196	<i>N'</i> -(3-[(6,7-Dimethoxy-5-methyl-4-oxo-1,4-dihydro-quinazolin-2-yl)-methyl-amino]-methyl)-phenyl)- <i>N,N</i> -dimethyl-formamidine	185-86
197	<i>N,N</i> -Dimethyl- <i>N'</i> -(3-([methyl-(5,6,7-trimethoxy-4-oxo-1,4-dihydro-quinazolin-2-yl)-amino]-methyl)-phenyl)-acetamidine	199-201

198	<i>N,N</i> -Dimethyl- <i>N'</i> -(3-{[methyl-(5,6,7-trimethoxy-4-oxo-1,4-dihydro-quinazolin-2-yl)-amino]-methyl}-phenyl)-isobutyramidine	148-50	
199	<i>N'</i> -(3-{[(5-Ethyl-6,7-dimethoxy-4-oxo-1,4-dihydro-quinazolin-2-yl)-methyl-amino]-methyl}-phenyl)- <i>N,N</i> -dimethyl-acetamide	224.6-225.5	
200	<i>N'</i> -(3-{[(5-Isopropyl-6,7-dimethoxy-4-oxo-1,4-dihydro-quinazolin-2-yl)-methyl-amino]-methyl}-phenyl)- <i>N,N</i> -dimethyl-acetamide	235.6-236.5	
201	<i>N'</i> -(3-{[(5-Hydroxy-6,7-dimethoxy-4-oxo-1,4-dihydro-quinazolin-2-yl)-methyl-amino]-methyl}-phenyl)- <i>N,N</i> -dimethyl-acetamide		426
202	3-Methoxy- <i>N,N</i> -dimethyl- <i>N'</i> -(3-{[methyl-(5,6,7-trimethoxy-4-oxo-1,4-dihydro-quinazolin-2-yl)-amino]-methyl}-phenyl)-propionamide	145	
203	<i>N,N</i> -Dimethyl- <i>N'</i> -(3-{[methyl-(5,6,7-trimethoxy-4-oxo-1,4-dihydro-quinazolin-2-yl)-amino]-methyl}-phenyl)-cyclobutanecarboxamide	164.2-165.7	
204	2-{[3-(Imidazolidin-2-ylideneamino)-benzyl]-methyl-amino}-6,7-dimethoxy-5-methyl-1 <i>H</i> -quinazolin-4-one	141.3-144.7	
205	5,6,7-Trimethoxy-2-(methyl-{3-[1-methyl-pyrrolidin-(2 <i>Z</i>)-ylideneamino]-benzyl}-amino)-1 <i>H</i> -quinazolin-4-one	130-133	
206	2-{[2-Chloro-5-(1,3-dimethyl-imidazolidin-2-ylideneamino)-benzyl]-methyl-amino}-5,6,7-trimethoxy-1 <i>H</i> -quinazolin-4-one		501, M+
207	2-{[3-(1,3-dimethyl-imidazolidin-2-ylideneamino)-benzyl]-methyl-amino}-5-ethyl-6,7-dimethoxy-1 <i>H</i> -quinazolin-4-one	212.5-219.9 (PF ₆ salt)	
208	2-{[3-(4,5-Dihydro-3 <i>H</i> -pyrrol-2-ylamino)-benzyl]-methyl-amino}-6,7-dimethoxy-5-methyl-1 <i>H</i> -quinazolin-4-one	175.4-177.2	
209	2-{[3-(4,5-Dihydro-3 <i>H</i> -pyrrol-2-ylamino)-benzyl]-methyl-amino}-5,6,7-trimethoxy-1 <i>H</i> -quinazolin-4-one	207-209	
210	5,6,7-Trimethoxy-2-{methyl-[3-(2-oxo-tetrahydro-furan-3-ylamino)-benzyl]-amino}-1 <i>H</i> -quinazolin-4-one		455
211	2-({5-[(4,5-Dihydro-3 <i>H</i> -pyrrol-2-yl)-methyl-amino]-2-fluoro-benzyl}-methyl-amino)-5-hydroxy-6,7-dimethoxy-	264.9-267	

	1 <i>H</i> -quinazolin-4-one		
212	6,7-Dimethoxy-5-methyl-2-{methyl-[3-(pyrimidin-2-ylamino)-benzyl]-amino}-1 <i>H</i> -quinazolin-4-one	433	
213	5,6,7-Trimethoxy-2-{methyl-[3-(pyrimidin-2-ylamino)-benzyl]-amino}-1 <i>H</i> -quinazolin-4-one	190.3-191.1	
214	2-[[3-(2-Chloro-pyrimidin-4-ylamino)-benzyl]-methyl-amino]-5,6,7-trimethoxy-1 <i>H</i> -quinazolin-4-one	205.9-210.1	
215	2-[[3-(4-Chloro-pyrimidin-2-ylamino)-benzyl]-methyl-amino]-5,6,7-trimethoxy-1 <i>H</i> -quinazolin-4-one	128.9-133.4	
216	2-[[3-(2-Chloro-pyrimidin-4-ylamino)-benzyl]-methyl-amino]-6,7-dimethoxy-5-methyl-1 <i>H</i> -quinazolin-4-one	467	
217	2-[[3-(2-Chloro-5-methyl-pyrimidin-4-ylamino)-benzyl]-methyl-amino]-5,6,7-trimethoxy-1 <i>H</i> -quinazolin-4-one	225.9-227.0	
218	2-[[3-(2-Chloro-6-methyl-pyrimidin-4-ylamino)-benzyl]-methyl-amino]-5,6,7-trimethoxy-1 <i>H</i> -quinazolin-4-one	185.5-187.1	
219	2-[[3-(4-Chloro-6-methyl-pyrimidin-2-ylamino)-benzyl]-methyl-amino]-5,6,7-trimethoxy-1 <i>H</i> -quinazolin-4-one	191.3-192.8	
220	2-({3-[(1 <i>H</i> -Imidazol-2-ylmethyl)-amino]-benzyl}-methyl-amino)-5,6,7-trimethoxy-1 <i>H</i> -quinazolin-4-one	125.0-131.0	
221	<i>N'</i> -(3-[(5-Fluoro-6,7-dimethoxy-4-oxo-1,4-dihydro-quinazolin-2-yl)-methyl-amino]-methyl)-phenyl)- <i>N,N</i> -dimethyl-acetamidine	199.9-201.9	
222	5,6,7-Trimethoxy-2-{methyl-[3-(2-oxo-pyrrolidin-1-yl)-benzyl]-amino}-1 <i>H</i> -quinazolin-4-one	165.5-166.5	439
223	6,7-Dimethoxy-5-methyl-2-{methyl-[3-(1-methyl-4,5-dihydro-1 <i>H</i> -imidazol-2-yl)-benzyl]-amino}-1 <i>H</i> -quinazolin-4-one	145-148	422
224	6,7-Dimethoxy-5-methyl-2-{methyl-[3-(1-methyl-piperidin-4-yl)-benzyl]-amino}-1 <i>H</i> -quinazolin-4-one	155.7-158.0	437
225	6,7-Dimethoxy-5-methyl-2-[methyl-(3-pyrrolidin-1-yl-benzyl)-amino]-1 <i>H</i> -quinazolin-4-one		409
226	5,6,7-Trimethoxy-2-[methyl-(3-pyrrolidin-1-yl-benzyl)-amino]-1 <i>H</i> -quinazolin-4-one		425
227	6,7-Dimethoxy-5-methyl-2-{methyl-[3-(2-methyl-4,5-dihydro-imidazol-1-yl)-benzyl]-amino}-1 <i>H</i> -quinazolin-4-one	177-180	

228	2-{{3-(4,4-Dimethyl-4,5-dihydro-imidazol-1-yl)-benzyl}-methyl-amino}-6,7-dimethoxy-5-methyl-1 <i>H</i> -quinazolin-4-one	437
229	6,7-Dimethoxy-5-methyl-2-{methyl-[3-(2,4,4-trimethyl-4,5-dihydro-imidazol-1-yl)-benzyl]-amino}-1 <i>H</i> -quinazolin-4-one	450
230	6,7-Dimethoxy-5-methyl-2-{methyl-[3-(4-methyl-imidazol-1-yl)-benzyl]-amino}-1 <i>H</i> -quinazolin-4-one	245-249.5
231	6,7-Dimethoxy-5-methyl-2-{methyl-[3-(2-methyl-imidazol-1-yl)-benzyl]-amino}-1 <i>H</i> -quinazolin-4-one	420
232	6,7-Dimethoxy-5-methyl-2-[methyl-(3-pyrazol-1-yl-benzyl)-amino]-1 <i>H</i> -quinazolin-4-one	220.8-224.1
233	6,7-Dimethoxy-5-methyl-2-{methyl-[3-(3-methyl-pyrazol-1-yl)-benzyl]-amino}-1 <i>H</i> -quinazolin-4-one	220-223.5
234	2-{{3-(3,5-Dimethyl-pyrazol-1-yl)-benzyl}-methyl-amino}-6,7-dimethoxy-5-methyl-1 <i>H</i> -quinazolin-4-one	434
235	6,7-Dimethoxy-5-methyl-2-[methyl-(3-pyrrolidin-2-yl-benzyl)-amino]-1 <i>H</i> -quinazolin-4-one	409
236	5,6,7-Trimethoxy-2-[methyl-(3-pyrrolidin-3-yl-benzyl)-amino]-1 <i>H</i> -quinazolin-4-one	425
237	6,7-Dimethoxy-5-methyl-2-{methyl-[4-(1-methyl-4,5-dihydro-1 <i>H</i> -imidazol-2-yl)-benzyl]-amino}-1 <i>H</i> -quinazolin-4-one	179-181
238	5,6,7-Trimethoxy-2-{methyl-[3-(1-methyl-4,5-dihydro-1 <i>H</i> -imidazol-2-yl)-benzyl]-amino}-1 <i>H</i> -quinazolin-4-one	438, M+
239	2-{{3-(1-Ethyl-4,5-dihydro-1 <i>H</i> -imidazol-2-yl)-benzyl}-methyl-amino}-6,7-dimethoxy-5-methyl-1 <i>H</i> -quinazolin-4-one	436
240	2-{{3-(1-Ethyl-4,5-dihydro-1 <i>H</i> -imidazol-2-yl)-benzyl}-methyl-amino}-5,6,7-trimethoxy-1 <i>H</i> -quinazolin-4-one	452
241	2-{{3-(1-Isopropyl-4,5-dihydro-1 <i>H</i> -imidazol-2-yl)-benzyl}-methyl-amino}-5,6,7-trimethoxy-1 <i>H</i> -quinazolin-4-one	466
242	2-{{4-(1-Isopropyl-4,5-dihydro-1 <i>H</i> -imidazol-2-yl)-benzyl}-methyl-amino}-6,7-dimethoxy-5-methyl-1 <i>H</i> -quinazolin-4-one	450
243	5-Hydroxy-2-{{3-(1-isopropyl-4,5-dihydro-1 <i>H</i> -imidazol-2-	453

	yl)-benzyl]-methyl-amino}-6,7-dimethoxy-1 <i>H</i> -quinazolin-4-one	
244	2-({3-[1-(2-Hydroxy-ethyl)-4,5-dihydro-1 <i>H</i> -imidazol-2-yl]-benzyl}-methyl-amino)-6,7-dimethoxy-5-methyl-1 <i>H</i> -quinazolin-4-one	253-254.9
245	2-({3-[1-(2-Hydroxy-ethyl)-4,5-dihydro-1 <i>H</i> -imidazol-2-yl]-benzyl}-methyl-amino)-5,6,7-trimethoxy-1 <i>H</i> -quinazolin-4-one	106.1-116.6 (TFA Salt)
246	6,7-Dimethoxy-2-({3-[1-(2-methoxy-ethyl)-4,5-dihydro-1 <i>H</i> -imidazol-2-yl]-benzyl}-methyl-amino)-5-methyl-1 <i>H</i> -quinazolin-4-one	161-164.4
247	5,6,7-Trimethoxy-2-({3-[1-(2-methoxy-ethyl)-4,5-dihydro-1 <i>H</i> -imidazol-2-yl]-benzyl}-methyl-amino)-1 <i>H</i> -quinazolin-4-one	483
248	2-{{3-(1 <i>H</i> -Imidazol-2-yl)-benzyl}-methyl-amino}-6,7-dimethoxy-5-methyl-1 <i>H</i> -quinazolin-4-one	255.5-258.9
249	6,7-Dimethoxy-5-methyl-2-{methyl-[3-(1-methyl-1 <i>H</i> -imidazol-2-yl)-benzyl]-amino}-1 <i>H</i> -quinazolin-4-one	179-182.1
250	2-({3-[1-(2-Hydroxy-ethyl)-1 <i>H</i> -imidazol-2-yl]-benzyl}-methyl-amino)-6,7-dimethoxy-5-methyl-1 <i>H</i> -quinazolin-4-one	451
251	6,7-Dimethoxy-5-methyl-2-{methyl-[3-(2-oxo-pyrrolidin-1-yl)-benzyl]-amino}-1 <i>H</i> -quinazolin-4-one	210.9-212.1
252	2-{{3-(1,1-Dioxo-1λ6-isothiazolidin-2-yl)-benzyl}-methyl-amino}-6,7-dimethoxy-5-methyl-1 <i>H</i> -quinazolin-4-one	226.0-228.5
253	{{3-(1,1-Dioxo-1λ6-isothiazolidin-2-yl)-benzyl}-methyl-amino}-5,6,7-trimethoxy-1 <i>H</i> -quinazolin-4-one	181.1-183.5
254	6,7-Dimethoxy-5-methyl-2-{methyl-[3-(2-oxo-oxazolidin-3-yl)-benzyl]-amino}-1 <i>H</i> -quinazolin-4-one	254.8-257.1
255	5,6,7-Trimethoxy-2-{methyl-[3-(2-oxo-oxazolidin-3-yl)-benzyl]-amino}-1 <i>H</i> -quinazolin-4-one	226.0-229.5
256	6,7-Dimethoxy-5-methyl-2-{methyl-[3-(3-methyl-2-oxo-pyrrolidin-1-yl)-benzyl]-amino}-1 <i>H</i> -quinazolin-4-one	210.8-211.6
257	2-{{3-(3-Ethyl-2-oxo-pyrrolidin-1-yl)-benzyl}-methyl-amino}-6,7-dimethoxy-5-methyl-1 <i>H</i> -quinazolin-4-one	200.1-202.1
258	2-{{3-(3-Ethyl-2-oxo-pyrrolidin-1-yl)-benzyl}-methyl-	467

	amino}-5,6,7-trimethoxy-1 <i>H</i> -quinazolin-4-one	
259	1-(3-{[(6,7-Dimethoxy-5-methyl-4-oxo-1,4-dihydro-quinazolin-2-yl)-methyl-amino]-methyl}-phenyl)-pyrrolidine-2,5-dione	198.6-199.8
260	1-(3-{[Methyl-(5,6,7-trimethoxy-4-oxo-1,4-dihydro-quinazolin-2-yl)-amino]-methyl}-phenyl)-pyrrolidine-2,5-dione	125.5-128.5
261	5,6,7-Trimethoxy-2-{methyl-[3-(2-oxo-imidazolidin-1-yl)-benzyl]-amino}-1 <i>H</i> -quinazolin-4-one	130.0-133.0
262	6,7-Dimethoxy-5-methyl-2-{methyl-[3-(3-methyl-2-oxo-imidazolidin-1-yl)-benzyl]-amino}-1 <i>H</i> -quinazolin-4-one	230.0-230.5
263	5,6,7-Trimethoxy-2-{methyl-[3-(3-methyl-2-oxo-imidazolidin-1-yl)-benzyl]-amino}-1 <i>H</i> -quinazolin-4-one	205.4-207.8
264	2-[[3-(3-Ethyl-2-oxo-imidazolidin-1-yl)-benzyl]-methyl-amino]-6,7-dimethoxy-5-methyl-1 <i>H</i> -quinazolin-4-one	201.4-207.4
265	2-[[3-(3-Ethyl-2-oxo-imidazolidin-1-yl)-benzyl]-methyl-amino]-5,6,7-trimethoxy-1 <i>H</i> -quinazolin-4-one	200.3-200.9
266	5,6,7-Trimethoxy-2-({3-[3-(2-methoxy-ethyl)-2-oxo-imidazolidin-1-yl]-benzyl}-methyl-amino)-1 <i>H</i> -quinazolin-4-one	199.5-202.1
267	6,7-Dimethoxy-5-methyl-2-{methyl-[3-(1 <i>H</i> -tetrazol-5-yl)-benzyl]-amino}-1 <i>H</i> -quinazolin-4-one	408
268	6,7-Dimethoxy-5-methyl-2-[methyl-(4-pyrrolidin-1-yl-pyridin-2-ylmethyl)-amino]-1 <i>H</i> -quinazolin-4-one	410
269	5,6,7-Trimethoxy-2-[(6-{1-[(2-methoxy-ethyl)-methyl-amino]-ethyl}-pyridin-2-ylmethyl)-methyl-amino]-1 <i>H</i> -quinazolin-4-one	472
270	2-[(4-Chloro-pyridin-2-ylmethyl)-methyl-amino]-5,6,7-trimethoxy-1 <i>H</i> -quinazolin-4-one	391, 393
271	5-Hydroxy-6,7-dimethoxy-2-({4-[(2-methoxy-ethyl)-methyl-amino]-pyridin-2-ylmethyl}-methyl-amino)-1 <i>H</i> -quinazolin-4-one	430.1
272	6,7-Dimethoxy-2-({4-[(2-methoxy-ethyl)-methyl-amino]-pyridin-2-ylmethyl}-methyl-amino)-5-methyl-1 <i>H</i> -quinazolin-4-one	428
273	5,6,7-Trimethoxy-2-{{4-(2-methoxy-ethoxy)-pyridin-2-ylmethyl}-methyl-amino}-1 <i>H</i> -quinazolin-4-one	431

274	2-[(4-Ethoxy-pyridin-2-ylmethyl)-methyl-amino]-5,6,7-trimethoxy-1 <i>H</i> -quinazolin-4-one	401
275	6,7-Dimethoxy-5-methyl-2-{methyl-[6-(1-pyrrolidin-1-yl-ethyl)-pyridin-2-ylmethyl]-amino}-1 <i>H</i> -quinazolin-4-one	75.4-85.6
276	5,6,7-Trimethoxy-2-{methyl-[6-(1-pyrrolidin-1-yl-ethyl)-pyridin-2-ylmethyl]-amino}-1 <i>H</i> -quinazolin-4-one	454
277	2-[[6-(1-Azetidin-1-yl-ethyl)-pyridin-2-ylmethyl]-methyl-amino]-6,7-dimethoxy-5-methyl-1 <i>H</i> -quinazolin-4-one	424
278	2-[[6-(1-Azetidin-1-yl-ethyl)-pyridin-2-ylmethyl]-methyl-amino]-5,6,7-trimethoxy-1 <i>H</i> -quinazolin-4-one	70.8-78.8
279	6,7-Dimethoxy-2-{[4-(2-methoxy-ethoxy)-pyridin-2-ylmethyl]-methyl-amino}-5-methyl-1 <i>H</i> -quinazolin-4-one	415
280	<i>N,N</i> -Dimethyl- <i>N'</i> -(3-{2-[methyl-(5,6,7-trimethoxy-4-oxo-1,4-dihydro-quinazolin-2-yl)-amino]-ethyl}-phenyl)-acetamidine	454
281	5-[2-(6,7-Dimethoxy-5-methyl-4-oxo-1,4-dihydro-quinazolin-2-ylamino)-propyl]-2-methoxy-benzenesulfonamide	463
282	5-[2-(6,7-Dimethoxy-5-methyl-4-oxo-1,4-dihydro-quinazolin-2-ylamino)-propyl]- <i>N</i> -[1-dimethylamino-eth-(<i>E</i>)-ylidene]-2-methoxy-benzenesulfonamide	532
283	<i>N'</i> -(3-{2-[(6,7-Dimethoxy-5-methyl-4-oxo-1,4-dihydro-quinazolin-2-yl)-methyl-amino]-1-hydroxy-ethyl}-phenyl)- <i>N,N</i> -dimethyl-acetamidine	455
284	<i>N'</i> -(3-{(<i>E</i>)-2-[(6,7-Dimethoxy-5-methyl-4-oxo-1,4-dihydro-quinazolin-2-yl)-methyl-amino]-vinyl}-phenyl)- <i>N,N</i> -dimethyl-acetamidine	436
285	2-[[2-(3 <i>H</i> -Imidazol-4-yl)-ethyl]-methyl-amino]-6,7-dimethoxy-5-methyl-1 <i>H</i> -quinazolin-4-one	344
286	2-[[2-(3 <i>H</i> -Imidazol-4-yl)-ethyl]-methyl-amino]-5,6,7-trimethoxy-1 <i>H</i> -quinazolin-4-one	360
287	2-[(1-Benzyl-piperidin-3-yl)-methyl-amino]-6,7-dimethoxy-5-methyl-1 <i>H</i> -quinazolin-4-one	423
288	<i>N,N</i> -Dimethyl- <i>N'</i> -(3-{[methyl-(6,7,8-trimethoxy-1,1-dioxo-1,4-dihydro-1 <i>λ</i> 6-benzo[1,2,4]thiadiazin-3-yl)-amino]-methyl}-phenyl)-acetamidine	477
289	<i>N'</i> -(3-{[(6,7-Dimethoxy-1,1-dioxo-1,4-dihydro-1 <i>λ</i> 6-benzo-	446

	[1,2,4]thiadiazin-3-yl)-methyl-amino]-methyl}-phenyl)- N,N-dimethyl-acetamide	
290	6,7-Dimethoxy-2-[3-(4-methyl-piperazin-1-yl)-benzyl- amino]-1H-quinazolin-4-one	410
291	6,7-Dimethoxy-5-methyl-2-[2-(4-methyl-piperazin-1-yl)- benzylamino]-1H-quinazolin-4-one	424
292	N-Methyl-N-[1-(3-{[methyl-(5,6,7-trimethoxy-4-oxo-1,4- dihydro-quinazolin-2-yl)-amino]-methyl}-phenyl)-ethyl]- methanesulfonamide	171.0-172. 4
293	1,1,3-Trimethyl-3-[1-(3-{[methyl-(5,6,7-trimethoxy-4-oxo- 1,4-dihydro-quinazolin-2-yl)-amino]-methyl}-phenyl)- ethyl]-urea	68-70
294	(3-{[(6,7-Dimethoxy-5-methyl-4-oxo-1,4-dihydro- quinazolin-2-yl)-methyl-amino]-methyl}-benzyl)-methyl- carbamic acid 2-methoxy-ethyl ester	145.8-146.9

Assay Examples

EXAMPLE L-1

The potency and selectivity of the inventive compounds as α 1A/B antagonists was deter-
 5 mined with CHO-K1 cells expressing adrenoceptor subtype α 1A-215, α 1B or α 1D by
 measuring cAMP accumulation using AlphaScreen.

Cell preparation was accomplished by culturing CHO- α 1 cloned cells in Ham's F12
 nutrient media supplemented with 10% FBS and G418 (25mg/mL), harvested at 80%
 confluence, washed with warmed PBS x2, and detached with versene for 5 min. at 37°C.
 10 The cultured cells were then resuspended in 40 mL of stimulation buffer (HBSS with 5
 mM hepes, 0.1% BSA) and centrifuged at 500-100 rpm for 5 min. The obtained pellet
 was resuspended in stimulation buffer (with 0.5M IBMX), and the cells were counted.
 Cells were diluted to the desired number of cells/mL (α 1A at 3×10^6 /mL, α 1B 15×10^6 /mL,
 and α 1D 20×10^6 /mL).

15 The compounds being tested were diluted in stimulation buffer (with 0.5M IBMX), from
 10^{-5} to 10^{-11} (final) dilution, 11 points. 5 μ L of each compound was dispensed to 96 well
 $\frac{1}{2}$ area plates in triplicate. 5 μ L of stimulation buffer was dispensed to a norepinephrine
 (NE) plate. 10 μ L of cells were added with anti-cAMP Acceptor beads in stimulation buffer
 to each plate and incubated for 15 min. at RT (in dark or covered with black plate). Then

5 μ l of NE was added to the antagonist plates, at 1 μ M for α 1A and 1B and at 100nM for α 1D, and then 5 μ l serial dilution of NE was added to NE plate. Plates were incubated for 30 min. at RT (in dark or covered with black plate) and 10 μ l Donor beads + biotin-cAMP in lysis buffer (5mM Hepes, 0.54% Tween-20, 0.1% BSA) was added. Plates were
5 incubated for 3h. at RT with gentle shaking (in dark or covered with black plate). Plates were read on an AlphaScreen Fusion analyzer, using reagent pursuant to AlphaScreen cAMP detection kit (PerkinElmer Cat#6760600).
Reference compounds norepinephrine, prazosin and vehicle were run in every experiment. In each plate, on column 12, A-D were loaded with only cells to define total
10 count and E-H were loaded with NE 1 μ M, (total – NE 1 μ M), to define 100% NE activity. The values determined for each experimental well on the plate were divided by (total count – NE 1 μ M), to determine % of NE activity. All data was plotted using non-linear curve fitting by GraphPad Prism to get pEC₅₀ (pEC₅₀ being the negative logarithm of EC₅₀, i.e., the molar concentration of an agonist which produces 50% of the maximum
15 possible response for that agonist) for norepinephrine and pK_b (wherein K_b is the equilibrium dissociation for a competitive antagonist, determined in a functional assay, and being equal to the concentration of antagonist which would occupy 50% of the receptors at equilibrium, units=mol l⁻¹, and pK_b is the negative logarithm of K_b) for prazosin and tested compounds. (AlphaScreen cAMP Detection Kit from PerkinElmer
20 Life Sciences).

Example L-2: [³H]prazosin Binding (Alpha1-Adrenoceptor) Assay

Alpha1A, alpha1B, and alpha1D adrenoceptor transfected CHO-K1 cells, prepared using the methods described by Chang et al., FEBS Lett. 422:279-283 (1998) were grown to confluence in T-162 tissue culture flasks in Ham's F-12 culture medium supplemented
25 with 10% fetal bovine serum, geneticin (150 μ g/mL) and streptomycin/penicillin (30 μ g/mL/30 μ g/mL) at 37° C in 7% CO₂. Cells were harvested by incubating with phosphate-buffered saline (PBS) containing 30 μ M EDTA for 5-10 min at 37° C. Cells were pelleted by centrifuging at 500 x g for 5min, and the pelleted cells were homogenized (Polytron homogenizer) in 10 vols (w/v) of 50 mM Tris, 1 mM EDTA,
30 (homogenisation buffer, pH 7.4 at 4° C). The homogenate was centrifuged at 45,000 x g for 20 min. The pellet was resuspended in the homogenizing buffer and rehomogenized. The resulting homogenate was centrifuged at 45,000 x g for 20 min. The pellet was resuspended in 50 mM Tris buffer (pH 7.4 at 4°C), aliquoted, frozen, and stored at -80° C for further use.

The membranes were thawed at RT and diluted in assay buffer (50 mM Tris buffer at pH 4) at 37°C and homogenized using the Polytron tissue disrupter. The membranes were incubated with the radioligand ($[^3\text{H}]$ prazosin, NEN, 0.1 – 0.5 nM) and test compound at 37°C for 30 min. The membranes were then filtered over polyethyleneimine-treated GF/B unifilter plates using a Packard Filtermate Harvester and washed with ice-cold 50 mM Tris-HCl, 1 mM EDTA buffer (3 x 3 sec. washes). Scintillation cocktail was added to the filter plates and bound radioligand determined by liquid scintillation spectrophotometry.

For each experiment, total binding (in the absence of any test or reference compounds) and non-specific binding (10 μM phentolamine) were determined. For each sample tested, the concentration producing 50% inhibition of binding (IC_{50}) and Hill Slope (n_{H}) was determined using iterative non-linear curve fitting techniques with Kaleidagraph (Synergy Software) or other appropriate software. If the radioligand K_{D} was known, the inhibition dissociation constant (K_{I}) of each ligand was determined according to the method of Cheng and Prusoff [Biochem. Pharmacol. 22:3099-3108 (1973)].

Proceeding as in Example L-2, compounds of Formula I were tested and found to be selective $\alpha_1\text{A/B}$ -adrenoceptor antagonists.

Example L-3: Dog *In Vivo* Intraurethral and Blood Pressure Assay

The following describes an *in vivo* assay for measuring the relative effect of test compounds on hypogastric nerve stimulation-induced increases in intraurethral pressure and phenylephrine-induced increases in diastolic blood pressure in anesthetized dog. Male Mongrel dogs (10 to 20 kg) were fasted for 12 to 18 hours and anesthetized with phenobarbital sodium (36 mg/kg, *i.v.*). An endotracheal tube was inserted and thereafter the lungs were mechanically ventilated with room air. The right femoral vein was isolated and cannulated with two polyethylene cannulae, one for the administration of a continuous infusion of phenobarbital sodium (5 to 10 mg/kg/hr) and the other for bolus administration of test substances. The right femoral artery was isolated and cannulated to the abdominal aorta with a fluid filled polyethylene cannula connected to an external pressure transducer for monitoring diastolic aortic pressure (DAP). The bladder was exposed via a ventral midline abdominal incision and emptied of urine through a 22 gauge needle. The bladder was cannulated through a stab incision with a water filled balloon catheter connected to an external pressure transducer for monitoring prostatic intraurethral pressure (IUP). The right hypogastric nerve (HGN) was carefully isolated and attached to a Dastre's electrode for nerve stimulation.

The preparation was allowed to stabilize for at least 20-30 minutes and must have had a stable basal IUP for not less than 15 minutes prior to commencement of the assay protocol. The HGN was stimulated (20-50V, 10Hz, 10 msec pulse train for 10 sec) to induce a measurable increase in IUP and then phenylephrine (PE) was administered by bolus injection (6 µg/kg, *i.v.*) to induce a measurable increase in DAP. The HGN stimulation and PE bolus injection were repeated every 5 minutes until three consecutive reproducible increases in IUP and DAP were achieved. Test compound was administered and 10 minutes later the HGN stimulation and PE bolus injection were repeated. Test compound was administered approximately every 20 minutes, increasing the dose until maximal or near maximal inhibition of the increases in IUP and DAP is attained. Proceeding as in Example L-3, compounds of Formula I were tested and found to selectively inhibit the HGN stimulation-induced increases in IUP. In contrast, prazosin inhibited increases in IUP and DAP in similar fashion.

Compounds of formula I are active in the above assays. For some exemplary compounds the following table shows corresponding data.


Compound	pKi		
	alpha 1A	alpha 1B	alpha 1D
<i>N'</i> -{3-[(6,7-dimethoxy-5-methyl-4-oxo-1,4-dihydro-quinazolin-2-ylamino)-methyl]-phenyl}- <i>N,N</i> -dimethyl-acetamidine	8.13	9.05	7.16
2-{[4-(2,3-dihydroxy-propoxy)-benzyl]-methyl-amino}-6,7-dimethoxy-5-methyl-1 <i>H</i> -quinazolin-4-one	8.15	8.16	6.10
2-{[3-(4,5-dihydro-oxazol-2-yl)-benzyl]-methyl-amino}-5,6,7-trimethoxy-1 <i>H</i> -quinazolin-4-one	6.09	7.83	5.58
2-({4-[(ethyl-methyl-amino)-methyl]-benzyl}-methyl-amino)-5,6,7-trimethoxy-1 <i>H</i> -quinazolin-4-one	8.50	8.51	6.26
5,6,7-trimethoxy-2-{[3-(3-methoxy-pyrrolidin-1-ylmethyl)-benzyl]-methyl-amino}-1 <i>H</i> -quinazolin-4-one	8.27	8.13	5.27
6,7-Dimethoxy-5-methyl-2-[methyl-(2-methyl-2,3-dihydro-1 <i>H</i> -isoindol-5-ylmethyl)-amino]-1 <i>H</i> -quinazolin-4-one	8.49	7.83	5.58
2-{[3-(1-Dimethylamino-ethyl)-benzyl]-methyl-	8.81	8.43	6.85

amino}-5,6,7-trimethoxy-1*H*-quinazolin-4-one

While the present invention has been described with reference to the specific embodiments thereof, it should be understood by those skilled in the art that various changes may be made and equivalents may be substituted without departing from the true spirit and scope of the invention. In addition, many modifications may be made to
5 adapt a particular situation, material, composition of matter, process, process step or steps, to the objective spirit and scope of the present invention. All such modifications are intended to be within the scope of the claims appended hereto.

Claims

1. A compound having the Formula (I)



wherein

- 5 Y is optionally-substituted C₁₋₄alkylene, C₂₋₄alkenylene, heterocyclylene, or heterocyclC₁₋₄alkylene;
 Z is -C(=O)- or -S(=O)₂-;
 R and R' are alkyl;
 R⁵ is selected from hydrogen, halogen, cyano, hydroxy, -R⁶, and -OR⁶;
 10 R⁶ is selected from alkyl, aryl, heteroaryl, cycloalkyl, and heterocyclyl;
 R¹⁰ is selected from hydrogen, alkyl, hydroxyalkyl, aminoalkyl and aralkyl;
 Ar is optionally-substituted aryl or heteroaryl,
 provided that if R⁵ is hydrogen, then Y-Ar considered together are not (i) unsubstituted benzyl, (ii) benzyl having a para substituent selected from hydroxy and CO₂H, or (iii)
 15 benzyl having a hydroxy meta substituent;
 or an isomer or pharmaceutically-acceptable salt, hydrate, or prodrug thereof.

2. The compound of claim 1 selected from

- N'-{3-[(6,7-dimethoxy-5-methyl-4-oxo-1,4-dihydro-quinazolin-2-ylamino)-methyl]-phenyl}-N,N-dimethyl-acetamidine,
 20 2-{[4-(2,3-dihydroxy-propoxy)-benzyl]-methyl-amino}-6,7-dimethoxy-5-methyl-1H-quinazolin-4-one,
 2-{[3-(4,5-dihydro-oxazol-2-yl)-benzyl]-methyl-amino}-5,6,7-trimethoxy-1H-quinazolin-4-one,
 2-(4-{[4-(ethyl-methyl-amino)-methyl]-benzyl}-methyl-amino)-5,6,7-trimethoxy-1H-quinazolin-4-one,
 25 5,6,7-trimethoxy-2-{[3-(3-methoxy-pyrrolidin-1-ylmethyl)-benzyl]-methyl-amino}-1H-quinazolin-4-one,
 6,7-dimethoxy-5-methyl-2-[methyl-(2-methyl-2,3-dihydro-1H-isoindol-5-ylmethyl)-amino]-1H-quinazolin-4-one, and
 30 2-{[3-(1-dimethylamino-ethyl)-benzyl]-methyl-amino}-5,6,7-trimethoxy-1H-quinazolin-4-one.

3. The compound of claim 1 selected from
- 5,6,7-trimethoxy-2-[methyl-(4-{[methyl-(3,3,3-trifluoro-propyl)-amino]-methyl}-benzyl)-amino]-1*H*-quinazolin-4-one;
- 2-{[4-(2-hydroxy-3-methoxy-propoxy)-benzyl]-methyl-amino}-5,6,7-trimethoxy-1*H*-quinazolin-4-one;
- 2-{[3-(3-hydroxy-2-hydroxymethyl-propoxy)-4-methoxy-benzyl]-methyl-amino}-5,6,7-trimethoxy-1*H*-quinazolin-4-one;
- 1-[1-(3-{[(6,7-dimethoxy-5-methyl-4-oxo-1,4-dihydro-quinazolin-2-yl)-methyl-amino]-methyl}-phenyl)-ethyl]-1,3,3-trimethyl-urea;
- 10 [1-(3-{[(6,7-dimethoxy-5-methyl-4-oxo-1,4-dihydro-quinazolin-2-yl)-methyl-amino]-methyl}-phenyl)-ethyl]-methyl-carbamic acid methyl ester;
- 2-{[3-(3-hydroxy-2-hydroxymethyl-propoxy)-4-methoxy-benzyl]-methyl-amino}-6,7-dimethoxy-5-methyl-1*H*-quinazolin-4-one;
- 2-{[3-(2,3-dihydroxy-propoxy)-benzyl]-methyl-amino}-6,7-dimethoxy-5-methyl-1*H*-quinazolin-4-one ;
- 15 5,6,7-trimethoxy-2-{[3-((*S*)-4-methoxy-2-oxo-pyrrolidin-1-yl)-benzyl]-methyl-amino}-1*H*-quinazolin-4-one;
- (4-{[(6,7-dimethoxy-5-methyl-4-oxo-1,4-dihydro-quinazolin-2-yl)-methyl-amino]-methyl}-benzyl)-methyl-carbamic acid 2-methoxy-ethyl ester;
- 20 *N*-(4-{[(6,7-dimethoxy-5-methyl-4-oxo-1,4-dihydro-quinazolin-2-yl)-methyl-amino]-methyl}-benzyl)-*N*-methyl-methanesulfonamide;
- 2-{[3-(4,5-dihydro-oxazol-2-yl)-benzyl]-methyl-amino}-5,6,7-trimethoxy-1*H*-quinazolin-4-one;
- 5,6,7-trimethoxy-2-{methyl-[4-(2-oxo-pyrrolidin-1-yl)-benzyl]-amino}-1*H*-quinazolin-4-one;
- 25 2-[(2-allyl-2,3-dihydro-1*H*-isoindol-5-ylmethyl)-methyl-amino]-5,6,7-trimethoxy-1*H*-quinazolin-4-one;
- 2-{[4-(2,3-dihydroxy-propoxy)-benzyl]-methyl-amino}-6,7-dimethoxy-5-methyl-1*H*-quinazolin-4-one;
- 30 2-{[4-(3-hydroxy-2-hydroxymethyl-propoxy)-benzyl]-methyl-amino}-6,7-dimethoxy-5-methyl-1*H*-quinazolin-4-one ;
- 2-[(3-{2-[(2-hydroxy-ethyl)-methyl-amino]-ethoxy}-4-methoxy-benzyl)-methyl-amino]-6,7-dimethoxy-5-methyl-1*H*-quinazolin-4-one;
- N*-(3-{[(6,7-dimethoxy-5-methyl-4-oxo-1,4-dihydro-quinazolin-2-yl)-methyl-amino]-methyl}-benzyl)-*N*-methyl-methanesulfonamide;
- 35 5,6,7-trimethoxy-2-{[2-(2-methoxy-ethyl)-2,3-dihydro-1*H*-isoindol-5-ylmethyl]-methyl-amino}-1*H*-quinazolin-4-one;


- 6,7-dimethoxy-2-[[2-(2-methoxy-ethyl)-2,3-dihydro-1*H*-isoindol-5-ylmethyl]-methyl-amino]-5-methyl-1*H*-quinazolin-4-one;
- 5,6,7-trimethoxy-2-[[3-(3-methoxy-2-oxo-pyrrolidin-1-yl)-benzyl]-methyl-amino]-1*H*-quinazolin-4-one;
- 5 *N*-methyl-*N*-(4-[[methyl-(5,6,7-trimethoxy-4-oxo-1,4-dihydro-quinazolin-2-yl)-amino]-methyl]-benzyl)-methanesulfonamide;
- 2-[[3-(3-hydroxy-2-oxo-pyrrolidin-1-yl)-benzyl]-methyl-amino]-5,6,7-trimethoxy-1*H*-quinazolin-4-one;
- 6,7-dimethoxy-5-methyl-2-{methyl-[4-(2-methylamino-ethyl)-benzyl]-amino}-1*H*-
- 10 quinazolin-4-one;
- 2-[[3-((*S*)-4-hydroxy-2-oxo-pyrrolidin-1-yl)-benzyl]-methyl-amino]-5,6,7-trimethoxy-1*H*-quinazolin-4-one;
- 2-({4-[2-(ethyl-methyl-amino)-ethoxy]-3-methoxy-benzyl}-methyl-amino)-6,7-dimethoxy-5-methyl-1*H*-quinazolin-4-one;
- 15 *N*-[1-(3-[[6,7-dimethoxy-5-methyl-4-oxo-1,4-dihydro-quinazolin-2-yl)-methyl-amino]-methyl)-phenyl]-ethyl]-*N*-methyl-methanesulfonamide;
- 2-[[3-((*S*)-4-hydroxy-2-oxo-pyrrolidin-1-yl)-benzyl]-methyl-amino]-6,7-dimethoxy-5-methyl-1*H*-quinazolin-4-one;
- 2-({4-[2-(ethyl-methyl-amino)-ethyl]-benzyl}-methyl-amino)-6,7-dimethoxy-5-methyl-
- 20 1*H*-quinazolin-4-one;
- 6,7-dimethoxy-2-[[3-methoxy-4-(2-methylamino-ethoxy)-benzyl]-methyl-amino]-5-methyl-1*H*-quinazolin-4-one;
- 2-[(4-{[ethyl-(2-hydroxy-ethyl)-amino]-methyl}-benzyl)-methyl-amino]-6,7-dimethoxy-5-methyl-1*H*-quinazolin-4-one;
- 25 2-[[4-(2-hydroxy-ethoxy)-benzyl]-methyl-amino]-6,7-dimethoxy-5-methyl-1*H*-quinazolin-4-one;
- 2-[(4-{[ethyl-(2-methoxy-ethyl)-amino]-methyl}-benzyl)-methyl-amino]-5,6,7-trimethoxy-1*H*-quinazolin-4-one;
- 2-[(3-[(2-hydroxy-ethyl)-methyl-amino]-methyl)-benzyl]-methyl-amino]-6,7-dimethoxy-5-methyl-1*H*-quinazolin-4-one;
- 30 5,6,7-trimethoxy-2-[[3-(3-methoxy-pyrrolidin-1-ylmethyl)-benzyl]-methyl-amino]-1*H*-quinazolin-4-one;
- 2-({4-[(ethyl-methyl-amino)-methyl]-benzyl}-methyl-amino)-5,6,7-trimethoxy-1*H*-quinazolin-4-one;
- 35 6,7-dimethoxy-2-({4-[(2-methoxy-ethylamino)-methyl]-benzyl}-methyl-amino)-5-methyl-1*H*-quinazolin-4-one;

- 6,7-dimethoxy-2-[[3-(3-methoxy-pyrrolidin-1-ylmethyl)-benzyl]-methyl-amino]-5-methyl-1*H*-quinazolin-4-one;
- 6,7-dimethoxy-5-methyl-2-[methyl-(2-methyl-2,3-dihydro-1*H*-isoindol-5-ylmethyl)-amino]-1*H*-quinazolin-4-one;
- 5 2-[[3-(4-hydroxy-piperidin-1-ylmethyl)-benzyl]-methyl-amino]-5,6,7-trimethoxy-1*H*-quinazolin-4-one;
- 2-[[3-(4-hydroxy-piperidin-1-ylmethyl)-benzyl]-methyl-amino]-6,7-dimethoxy-5-methyl-1*H*-quinazolin-4-one;
- 2-[[4-(3-hydroxy-pyrrolidin-1-ylmethyl)-benzyl]-methyl-amino]-6,7-dimethoxy-5-methyl-1*H*-quinazolin-4-one;
- 10 2-[(4-aminomethyl-benzyl)-methyl-amino]-6,7-dimethoxy-5-methyl-1*H*-quinazolin-4-one;
- 2-[(3-ethylaminomethyl-benzyl)-methyl-amino]-5,6,7-trimethoxy-1*H*-quinazolin-4-one;
- 15 2-[[4-(3-dimethylamino-propoxy)-benzyl]-methyl-amino]-6,7-dimethoxy-5-methyl-1*H*-quinazolin-4-one;
- 5,6,7-trimethoxy-2-[methyl-(2-methyl-2,3-dihydro-1*H*-isoindol-5-ylmethyl)-amino]-1*H*-quinazolin-4-one;
- 5,6,7-trimethoxy-2-[methyl-(3-pyrrolidin-1-ylmethyl-benzyl)-amino]-1*H*-quinazolin-4-one;
- 20 2-[(3-azetidin-1-ylmethyl-benzyl)-methyl-amino]-5,6,7-trimethoxy-1*H*-quinazolin-4-one;
- 2-[[4-(3-dimethylamino-propoxy)-benzyl]-methyl-amino]-5,6,7-trimethoxy-1*H*-quinazolin-4-one;
- 25 2-[[3-(3-hydroxy-pyrrolidin-1-ylmethyl)-benzyl]-methyl-amino]-6,7-dimethoxy-5-methyl-1*H*-quinazolin-4-one;
- 2-[[3-(1,3-dimethyl-imidazolidin-2-ylideneamino)-benzyl]-methyl-amino]-6,7-dimethoxy-5-methyl-1*H*-quinazolin-4-one;
- 2-[[3-(3-hydroxy-pyrrolidin-1-ylmethyl)-benzyl]-methyl-amino]-5,6,7-trimethoxy-1*H*-quinazolin-4-one;
- 30 *N*-[1-(3-[[[(6,7-dimethoxy-5-methyl-4-oxo-1,4-dihydro-quinazolin-2-yl)-methyl-amino]-methyl]-phenyl)-ethyl]-*N*-methyl-2-methylamino-acetamide;
- N'*-(3-[[[(5-cyano-6,7-dimethoxy-4-oxo-1,4-dihydro-quinazolin-2-yl)-methyl-amino]-methyl]-phenyl)-*N,N*-dimethyl-acetamidine;
- 35 6,7-dimethoxy-2-[(3-[1-[(2-methoxy-ethyl)-methyl-amino]-ethyl]-benzyl)-methyl-amino]-5-methyl-1*H*-quinazolin-4-one;

- 6,7-dimethoxy-5-methyl-2-[methyl-(4-methylaminomethyl-benzyl)-amino]-1H-quinazolin-4-one;
- 2-[(3-dimethylaminomethyl-benzyl)-methyl-amino]-6,7-dimethoxy-5-methyl-1H-quinazolin-4-one;
- 5 2-[(3-dimethylaminomethyl-benzyl)-methyl-amino]-5,6,7-trimethoxy-1H-quinazolin-4-one;
- 1,5-dimethyl-3-(3-[[methyl-(5,6,7-trimethoxy-4-oxo-1,4-dihydro-quinazolin-2-yl)-amino]-methyl]-phenyl)-imidazolidine-2,4-dione;
- 1-methyl-3-(3-[[methyl-(5,6,7-trimethoxy-4-oxo-1,4-dihydro-quinazolin-2-yl)-amino]-methyl]-phenyl)-imidazolidine-2,4-dione;
- 10 3-(3-[[[(6,7-dimethoxy-5-methyl-4-oxo-1,4-dihydro-quinazolin-2-yl)-methyl-amino]-methyl]-phenyl]-1-methyl-imidazolidine-2,4-dione;
- 2-[[3-(1-dimethylamino-ethyl)-benzyl]-methyl-amino]-6,7-dimethoxy-5-methyl-1H-quinazolin-4-one;
- 15 2-[[3-(1-dimethylamino-ethyl)-benzyl]-methyl-amino]-5,6,7-trimethoxy-1H-quinazolin-4-one;
- 5,6,7-trimethoxy-2-{methyl-[3-(1-methylamino-ethyl)-benzyl]-amino}-1H-quinazolin-4-one;
- 6,7-dimethoxy-5-methyl-2-{methyl-[3-(1-methylamino-ethyl)-benzyl]-amino}-1H-quinazolin-4-one;
- 20 1-(3-[[[(6,7-dimethoxy-5-methyl-4-oxo-1,4-dihydro-quinazolin-2-yl)-methyl-amino]-methyl]-phenyl]-3-methyl-pyrrolidine-2,5-dione;
- 2-[(3-cyclopropylaminomethyl-benzyl)-methyl-amino]-6,7-dimethoxy-5-methyl-1H-quinazolin-4-one;
- 25 2-[[3-(isopropylamino-methyl)-benzyl]-methyl-amino]-6,7-dimethoxy-5-methyl-1H-quinazolin-4-one;
- 6,7-dimethoxy-5-methyl-2-(methyl-{3-[(methyl-propyl-amino)-methyl]-benzyl}-amino)-1H-quinazolin-4-one;
- 6,7-dimethoxy-5-methyl-2-(3-methylaminomethyl-benzylamino)-1H-quinazolin-4-one;
- 30 N'-(3-[[[(6,7-dimethoxy-4-oxo-1,4-dihydro-quinazolin-2-yl)-methyl-amino]-methyl]-phenyl]-N,N-dimethyl-acetamidine;
- 2-([3-[(ethyl-methyl-amino)-methyl]-benzyl]-methyl-amino)-6,7-dimethoxy-5-methyl-1H-quinazolin-4-one;
- N'-{3-[(6,7-dimethoxy-5-methyl-4-oxo-1,4-dihydro-quinazolin-2-ylamino)-methyl]-phenyl}-N,N-dimethyl-acetamidine;
- 35 5,6,7-trimethoxy-2-{methyl-[3-(3-methyl-2-oxo-pyrrolidin-1-yl)-benzyl]-amino}-1H-quinazolin-4-one;


- 6,7-dimethoxy-5-methyl-2-{methyl-[3-(2-oxo-imidazolidin-1-yl)-benzyl]-amino}-1*H*-quinazolin-4-one;
- 6,7-dimethoxy-2-({3-[(2-methoxy-ethylamino)-methyl]-benzyl}-methyl-amino)-5-methyl-1*H*-quinazolin-4-one;
- 5 5,6,7-trimethoxy-2-{methyl-[3-(1-methyl-piperidin-4-yl)-benzyl]-amino}-1*H*-quinazolin-4-one hydrochloride;
- 6,7-dimethoxy-2-[methyl-(3-pyrrolidin-1-yl-benzyl)-amino]-1*H*-quinazolin-4-one bishydrochloride;
- 2-[benzyl-(1-benzyl-piperidin-4-yl)-amino]-6,7-dimethoxy-1*H*-quinazolin-4-one;
- 10 6,7-dimethoxy-2-[methyl-(6-methyl-pyridin-2-ylmethyl)-amino]-1*H*-quinazolin-4-one;
- 2-[[2-(1*H*-indol-3-yl)-ethyl]-methyl-amino]-6,7-dimethoxy-1*H*-quinazolin-4-one;
- 6,7-dimethoxy-2-{methyl-[3-(4-methyl-piperazin-1-yl)-benzyl]-amino}-1*H*-quinazolin-4-one;
- 6,7-dimethoxy-2-[2-(4-methyl-piperazin-1-yl)-benzylamino]-1*H*-quinazolin-4-one;
- 15 2-(benzyl-isopropyl-amino)-6,7-dimethoxy-1*H*-quinazolin-4-one;
- N'*-(3-[[6,7-dimethoxy-4-oxo-1,4-dihydro-quinazolin-2-yl)-methyl-amino]-methyl}-phenyl)-*N,N*-dimethyl-formamidine ;
- 2-[[3-(4,5-dihydro-3*H*-pyrrol-2-ylamino)-benzyl]-methyl-amino]-6,7-dimethoxy-1*H*-quinazolin-4-one;
- 20 *N*-(3-[[6,7-dimethoxy-4-oxo-1,4-dihydro-quinazolin-2-yl)-methyl-amino]-methyl}-phenyl)-acetamidine;
- 2-[[3-(4,5-dihydro-1*H*-imidazol-2-ylamino)-benzyl]-methyl-amino]-6,7-dimethoxy-1*H*-quinazolin-4-one;
- 2-[[3-(4,4-dimethyl-4,5-dihydro-imidazol-1-yl)-benzyl]-methyl-amino]-6,7-dimethoxy-1*H*-quinazolin-4-one;
- 25 *N'*-(3-{2-[[6,7-dimethoxy-4-oxo-1,4-dihydro-quinazolin-2-yl)-methyl-amino]-ethyl}-phenyl)-*N,N*-dimethyl-acetamidine;
- 6,7-dimethoxy-2-(methyl-{2-[3-(pyrimidin-2-ylamino)-phenyl]-ethyl}-amino)-1*H*-quinazolin-4-one;
- 30 2-[(2-{3-[(1*H*-imidazol-2-ylmethyl)-amino]-phenyl}-ethyl)-methyl-amino]-6,7-dimethoxy-1*H*-quinazolin-4-one;
- 2-[[2-(3,4-dimethoxy-phenyl)-ethyl]-methyl-amino]-6,7-dimethoxy-1*H*-quinazolin-4-one;
- 2-[(3-amino-benzyl)-methyl-amino]-6,7-dimethoxy-5-methyl-1*H*-quinazolin-4-one;
- 35 3-[[6,7-dimethoxy-5-methyl-4-oxo-1,4-dihydro-quinazolin-2-yl)-methyl-amino]-methyl}-benzonitrile;

- 2-[benzyl-(2-dimethylamino-ethyl)-amino]-6,7-dimethoxy-5-methyl-1*H*-quinazolin-4-one; and
- 2-[(3-bromo-benzyl)-methyl-amino]-6,7-dimethoxy-5-methyl-1*H*-quinazolin-4-one; and
- 5 *N'*-(3-[(7-chloro-6-methoxy-4-oxo-1,4-dihydro-quinazolin-2-yl)-methyl-amino]-methyl}-phenyl)-*N,N*-dimethyl-acetamide.
4. The compound of claim 1 or claim 3 for use as therapeutic active substance.
5. A pharmaceutical composition comprising a therapeutically effective amount of at least one compound of claim 1 or claim 3 and at least one pharmaceutically acceptable
- 10 carrier.
6. A method of treating a disease state in a patient modulated via an alpha-1A/B adrenoceptor wherein said method comprises administering to the subject in need of such treatment a therapeutically effective amount of at least one compound of claim 1 or claim 3.
- 15 7. The method of claim 6 wherein the disease state is selected from disorders and symptoms of the urinary tract, sexual dysfunction, benign prostatic hypertrophy, and pain.
8. Use of a compound of claim 1 or claim 3 for the manufacture of a medicament comprising a compound according to claim 1 or claim 3 for treating a disease state in a
- 20 patient modulated via an alpha-1A/B adrenoceptor.
9. A process of making an intermediate quinazolidione of formula (II)



wherein R, R' and R⁵ are as recited in Claim 1, comprising:


- (a) dissolving in water with the addition of base, a nitro-benzoic compound of formula
- 25 (8),



(8) ;


wherein R, R' and R⁵ are as recited in Claim 1;

- (b) adding to the aqueous, basic solution of compound (4), a noble metal catalyst on an inert support, and exposing the solution to a hydrogen environment to provide an amine compound of the formula (9),



(9) ;


- (c) reacting the amine compound of formula (9), with a cyanate salt and an acid to provide a urea compound having the formula (10),



(10)

- and,

- (d) cyclizing the urea of formula (10) with addition of base and heat to provide the quinazolinodione of formula (II),



(II),

- wherein the process of making dione of quinazolinodione of formula (II) from the nitrobenzoic compound of formula (8), is performed in a water-based solvent system.

10. The invention as hereinbefore described.