

PROCESS FOR PREPARING MATRIX METALLOPROTEINASE INHIBITORS AND CHIRAL AUXILIARY THEREFOR

Field of the Invention

The present invention relates to chiral auxiliaries and their use.

5 The present invention also relates to the preparation of certain β -hydroxy- and amino-substituted carboxylic acids which are matrix metalloprotease inhibitors, particularly diastereomerically pure β -hydroxy-carboxylic acids.

Background of the Invention

10 Metalloproteinases (MMPs) are a naturally occurring superfamily of proteinases (enzymes) found in most mammals. The superfamily is composed of at least 26 members of zinc-containing enzymes produced by many cell types and sharing structural and functional features. Based on structural and functional considerations proteinases have been classified into different families and subfamilies (Vartak et al., (2007), *J. Drug Targeting*, 15, 1-20; and Hopper, (1994), *FEBS*, 354, 1-6), such as collagenases (MMP-1, -8 and -13), gelatinases (MMP-2, and -9),
15 metalloelastases (MMP-12), the MT-MMPs (MMP-14, -15, -16, -17, -24 and 25), matrilysins (MMP-7 and -26), stromelysins (MMP-3, -10 and -11) and sheddases such as TNF-converting enzymes (TACE, and ACE).

20 Metalloproteinases are believed to be important in physiological disease processes that involve remodeling such as embryonic development, bone formation and uterine remodeling during menstruation. One major biological function of MMPs is to catalyze the breakdown of connective tissues or extra-cellular matrix by their ability to hydrolyze various components of tissue or matrix. Apart from their role in degrading connective tissue, MMPs are involved in the activation of zymogen (pro) forms of other MMPs thereby inducing MMP activation. They are
25 also involved in the biosynthesis of TNF-alpha which is implicated in many pathological conditions.

MMP-12, also known as macrophage elastase or metalloelastase, is expressed in activated macrophages and has been shown to be secreted from alveolar macrophages from smokers as well as in foam cells in atherosclerotic lesions. MMP-12 knockout mouse studies have shown the development of significant emphysema, thus supporting its role in COPD. MMP-9 (gelatinase B, 92 kDa type IV collagenase) is one member of the MMP family that is released as a proenzyme and subsequently activated *via* a protease cascade *in vivo*.

The concentration of MMP-9 is increased in diseases like asthma, interstitial pulmonary fibrosis (IPF), adult respiratory distress syndrome (ARDS), and in chronic obstructive pulmonary disease (COPD). Because of its proteolytic ability, MMP-9 has been implicated in tissue remodelling of the airways and lungs in chronic inflammatory diseases such as severe asthma and COPD. MMP-9 is also likely to be physiologically important because of its ability to regulate the digestion of components of the extracellular matrix as well as the activity of other proteases and cytokines. MMP-9 is secreted in neutrophils, macrophages, osteoclasts, which are easily induced by cytokines and growth factors, and plays a role in various physiological and pathological processes.

Over-expression or over-activation of an MMP, or an imbalance between an MMP and a natural (i.e., endogenous) tissue inhibitor of a matrix metalloproteinase (TIMP) has been linked to a pathogenesis of diseases characterized by the breakdown of connective tissue or extracellular matrix.

Inhibition of the activity of one or more MMPs may be of benefit in treatment of various inflammatory, autoimmune and allergic diseases such as, inflammation of the joint, inflammation of the GI tract, inflammation of the skin, collagen remodeling, wound healing disorders etc.

The design and therapeutic application of MMP inhibitors has revealed that the requirement of a molecule to be an effective inhibitor of MMP class of enzymes is a functional group (e.g. carboxylic acid, hydroxamic acid or sulphhydryl) capable of chelating to the active site Zn^{2+} ion (Whittaker *et al.*, *Chem. Rev.*, (1999), 99; 2735-76).

WO 2004/110974 discloses compounds and their physiologically functional derivatives described as inhibitors of matrix metalloproteinase enzymes. WO 2004/113279 discloses alleged inhibitors of matrix metalloproteinase. WO 2005/026120

discloses compounds also described as inhibitors of matrix metalloproteinase. U.S. Patent No. 6,350,885 discloses tricyclic heteroaromatic compounds and their derivatives believed to be inhibitors of matrix metalloproteinases. WO 98/09940 discloses biphenyl butyric acids and their derivatives described as inhibitors of matrix metalloproteinases. *J. Med. Chem.*, (1968), vol. 11(6), 1139-1144 discloses the synthesis and anti-inflammatory activity of 4-(p-biphenyl)-3-hydroxybutyric acid and related compounds. WO 96/15096 discloses substituted 4-biarylbutyric or 5-biarylpentanoic acids and derivatives as alleged matrix metalloproteinase inhibitors. WO 2006/090235 describes 5-phenyl-pentanoic acid derivatives described as matrix metalloproteinase inhibitors for the treatment of asthma and other diseases.

Research has been carried out into the identification of inhibitors that are selective, *e.g.*, for a few of the MMP subtypes. An MMP inhibitor of improved selectivity would avoid potential side effects associated with inhibition of MMPs that are not involved in the pathogenesis of the disease being treated.

Further, use of more selective MMP inhibitors would require administration of a lower amount of the inhibitor for treatment of disease than would otherwise be required and, after administration, partitioned *in vivo* among multiple MMPs. Still further, the administration of a lower amount of compound would improve the margin of safety between the dose of the inhibitor required for therapeutic activity and the dose of the inhibitor at which toxicity is observed.

Many drugs exist as asymmetric three-dimensional molecules i.e., chiral and will therefore have several stereoisomers depending upon the number of chiral centers present. The importance of evaluating new chemical entities having chiral centers as single isomers is to understand their effect on pharmacological and toxicological aspects. There are often pharmacodynamic, pharmacokinetic and/or toxicological differences between enantiomers / diastereomers. Even if natural physiological mediators are achiral, based on their target environment, their receptors/enzymes may demonstrate a preference for only one optically pure enantiomer of agonists, antagonists or inhibitors. From a pharmacokinetics point of view, chirality can have an influence on drug absorption, distribution, metabolism and elimination. Pure single isomers may also offer advantages in terms of these pharmacokinetic parameters thus enabling better developability of such molecules as drug candidates. It is also known that chirality has a significant effect of the physicochemical

properties and crystallinity of a chiral molecule which in turn have profound effects on the pharmacokinetics and developability of the molecule. Besides those mentioned above, regulatory principles guide one to preferably develop single isomers as drug candidates in order to avoid any pharmacological, pharmacokinetic and toxicological problems that may arise due to interactions of an unwanted isomer with undesirable molecular targets.

In this context, synthetic strategies to produce pure single isomers offer advantages over analytical techniques of separation of isomer not only in terms of cost and efficiency but larger amounts of compound can be prepared for elaborate pharmaceutical testing. Thus, compounds prepared according to the present invention, which are single chiral isomers, have improved potency, improved pharmacokinetics and /or improved physicochemical properties as compared to racemic compounds.

The present invention is directed to overcoming problems encountered in the art.

Summary of the Invention

According to one aspect, the present invention relates to a chiral auxiliary (4S)-4-benzyl-1,3-thiazolidin-2-one and its use.

The present invention also relates to the preparation of β -hydroxy and amino substituted carboxylic acids, which act as matrix metalloprotease inhibitors, particularly diastereomerically pure β -hydroxy carboxylic acids and pharmaceutical compositions containing them. The compounds prepared are matrix metalloproteinase inhibitors useful as effective therapeutic or prophylactic agents in treatment of various inflammatory, autoimmune, and allergic diseases and other inflammatory disorders characterized by the over-expression and over-activation of a matrix metalloproteinase using the compounds.

The compounds prepared according to the invention act as dual MMP-9/12 inhibitors, which have desirable activity profiles and beneficial potency, selectivity and/or pharmacokinetic properties.

The present invention prepares chemical entities having chiral centers as single isomers. Synthetic strategies to produce pure single isomers that offer advantages over analytical techniques of separation of isomer not only in terms of cost and efficiency but also larger amount of compound can be prepared for elaborate pharmaceutical testing are also provided.

Other aspects will be set forth in the description which follows, and in part will be apparent from the description or may be learnt by the practice of the invention.

Detailed Description of the Invention

5 The present invention relates to a chiral auxiliary (4S)-4-benzyl-1,3-thiazolidin-2-one which would be of versatile utility for asymmetric synthesis. Chiral auxiliaries are utilized in a wide variety of synthetic transformations which include, but not limited to, asymmetric aldol condensation, stereoselective alkylation, stereoselective Diels-Alder reaction, stereoselective Michael reactions, stereoselective differentiation of enantiotopic
10 groups in molecules bearing prochiral centers. The chiral auxiliary is used in stoichiometric amounts to induce the stereoselective formation of stereogenic centers.

DEFINITIONS

The following definitions apply to terms as used herein

15 The term "alkyl," unless otherwise specified, refers to a monoradical branched or unbranched saturated hydrocarbon chain having from 1 to 20 carbon atoms. Alkyl groups can be optionally interrupted by atom(s) or group(s) independently selected from oxygen, sulfur, a phenylene, sulphinyl, sulphonyl group or $-NR_\alpha-$, wherein R_α can be hydrogen, alkyl, cycloalkyl, alkenyl, cycloalkenyl, alkynyl, aryl, acyl, aralkyl, $-C(=O)OR_\lambda$, SO_mR_ψ or
20 $-C(=O)NR_\lambda R_\pi$. This term can be exemplified by groups such as methyl, ethyl, n-propyl, iso-propyl, n-butyl, iso-butyl, sec-butyl, t-butyl, n-pentyl, isopentyl, neopentyl, n-hexyl, n-decyl, tetradecyl, and the like. Alkyl groups may be substituted further with one or more substituents selected from alkenyl, alkynyl, alkoxy, cycloalkyl, cycloalkenyl, acyl, acylamino, acyloxy, alkoxycarbonylamino, azido, cyano, halogen, hydroxy, keto, oxo,
25 thiocarbonyl, carboxy, carboxyalkyl, aryl, heterocyclyl, heteroaryl, (heterocyclyl)alkyl, cycloalkoxy, $-CH=N-O(C_{1-6}alkyl)$, $-CH=N-NH(C_{1-6}alkyl)$, $-CH=N-NH(C_{1-6}alkyl)-C_{1-6}alkyl$, arylthio, thiol, alkylthio, aryloxy, nitro, aminosulfonyl, aminocarbonylamino, $-NHC(=O)R_\lambda$, $-NR_\lambda R_\pi$, $-C(=O)NR_\lambda R_\pi$, $-NHC(=O)NR_\lambda R_\pi$, $-C(=O)heteroaryl$,

C(=O)heterocyclyl, -O-C(=O)NR_λR_π {wherein R_λ and R_π are independently selected from hydrogen, halogen, hydroxy, alkyl, alkenyl, alkynyl, alkenyl, alkoxy, cycloalkyl, cycloalkenyl, aryl, aralkyl, heterocyclyl, heteroaryl, heterocyclalkyl, heteroarylalkyl or carboxy}, nitro or -SO_mR_ψ (wherein m is an integer from 0-2 and R_ψ is hydrogen, alkyl, alkenyl, alkynyl, cycloalkyl, aralkyl, aryl, heterocyclyl, heteroaryl, heteroarylalkyl or heterocyclalkyl). Unless otherwise constrained by the definition, alkyl substituents may be further substituted by 1-3 substituents selected from alkyl, alkenyl, alkynyl, carboxy, -NR_λR_π, -C(=O)NR_λR_π, -OC(=O)NR_λR_π, -NHC(=O)NR_λR_π, hydroxy, alkoxy, halogen, CF₃, cyano, and -SO_mR_ψ; or an alkyl group also may be interrupted by 1-5 atoms of groups independently selected from oxygen, sulfur or -NR_α- (wherein R_α, R_λ, R_π, m and R_ψ are the same as defined earlier). Unless otherwise constrained by the definition, all substituents may be substituted further by 1-3 substituents selected from alkyl, alkenyl, alkynyl, carboxy, carboxyalkyl, -NR_λR_π, -C(=O)NR_λR_π, -O-C(=O)NR_λR_π, hydroxy, alkoxy, halogen, CF₃, cyano, and -SO_mR_ψ (wherein R_λ, R_π, m and R_ψ are the same as defined earlier); or an alkyl group as defined above that has both substituents as defined above and is also interrupted by 1-5 atoms or groups as defined above.

The term “alkenyl,” unless otherwise specified, refers to a monoradical of a branched or unbranched unsaturated hydrocarbon group having from 2 to 20 carbon atoms with cis, trans or geminal geometry. Alkenyl groups can be optionally interrupted by atom(s) or group(s) independently chosen from oxygen, sulfur, phenylene, sulphinyl, sulphonyl and -NR_α- (wherein R_α is the same as defined earlier). In the event that alkenyl is attached to a heteroatom, the double bond cannot be alpha to the heteroatom. Alkenyl groups may be substituted further with one or more substituents selected from alkyl, alkenyl, alkynyl, alkoxy, cycloalkyl, cycloalkenyl, acyl, acylamino, acyloxy, -NHC(=O)R_λ, -NR_λR_π, -C(=O)NR_λR_π, -NHC(=O)NR_λR_π, -O-C(=O)NR_λR_π, alkoxycarbonylamino, azido, cyano, halogen, hydroxy, oxo, keto, carboxyalkyl, thiocarbonyl, carboxy, arylthio, thiol, alkylthio, aryl, aralkyl, aryloxy, heterocyclyl, heteroaryl, heterocyclalkyl, heteroaryl alkyl, aminosulfonyl, aminocarbonylamino, alkoxyamino, hydroxyamino, alkoxyamino, nitro or SO_mR_ψ (wherein R_λ, R_π, m and R_ψ are as defined earlier). Unless otherwise constrained by the definition, alkenyl substituents

optionally may be substituted further by 1-3 substituents selected from alkyl, alkenyl, alkynyl, carboxy, hydroxy, alkoxy, halogen, $-\text{CF}_3$, cyano, $-\text{NR}_\lambda\text{R}_\pi$, $-\text{C}(=\text{O})\text{NR}_\lambda\text{R}_\pi$, $-\text{O}-\text{C}(=\text{O})\text{NR}_\lambda\text{R}_\pi$ and $-\text{SO}_m\text{R}_\psi$ (wherein R_λ , R_π , m and R_ψ are as defined earlier). Groups, such as ethenyl or vinyl ($\text{CH}=\text{CH}_2$), 1-propylene or allyl ($-\text{CH}_2\text{CH}=\text{CH}_2$), iso-propylene ($-\text{C}(\text{CH}_3)=\text{CH}_2$), bicyclo[2.2.1]heptene, and the like, exemplify this term.

The term “alkynyl,” unless otherwise specified, refers to a monoradical of an unsaturated hydrocarbon, having from 2 to 20 carbon atoms. Alkynyl groups can be optionally interrupted by atom(s) or group(s) independently chosen from oxygen, sulfur, phenylene, sulphinyl, sulphonyl and $-\text{NR}_\alpha-$ (wherein R_α is the same as defined earlier). In the event that alkynyl groups are attached to a heteroatom, the triple bond cannot be alpha to the heteroatom. Alkynyl groups may be substituted further with one or more substituents selected from alkyl, alkenyl, alkoxy, cycloalkyl, cycloalkenyl, acyl, acylamino, acyloxy, alkoxy-carbonylamino, azido, cyano, halogen, hydroxy, keto, oxo, thiocarbonyl, carboxy, carboxyalkyl, arylthio, thiol, alkylthio, aryl, aralkyl, aryloxy, aminosulfonyl, aminocarbonylamino, hydroxyamino, alkoxyamino, nitro, heterocyclyl, heteroaryl, heterocyclylalkyl, heteroarylalkyl, $-\text{NHC}(=\text{O})\text{R}_\lambda$, $-\text{NR}_\lambda\text{R}_\pi$, $-\text{NHC}(=\text{O})\text{NR}_\lambda\text{R}_\pi$, $-\text{C}(=\text{O})\text{NR}_\lambda\text{R}_\pi$, $-\text{O}-\text{C}(=\text{O})\text{NR}_\lambda\text{R}_\pi$ or $-\text{SO}_m\text{R}_\psi$ (wherein R_λ , R_π , m and R_ψ are the same as defined earlier). Unless otherwise constrained by the definition, alkynyl substituents optionally may be substituted further by 1-3 substituents selected from alkyl, alkenyl, alkynyl, carboxy, carboxyalkyl, hydroxy, alkoxy, halogen, CF_3 , $-\text{NR}_\lambda\text{R}_\pi$, $-\text{C}(=\text{O})\text{NR}_\lambda\text{R}_\pi$, $-\text{NHC}(=\text{O})\text{NR}_\lambda\text{R}_\pi$, $-\text{C}(=\text{O})\text{NR}_\lambda\text{R}_\pi$, cyano or $-\text{SO}_m\text{R}_\psi$ (wherein R_λ , R_π , m and R_ψ are the same as defined earlier).

The term “cycloalkyl,” unless otherwise specified, refers to cyclic alkyl groups of from 3 to 20 carbon atoms having a single cyclic ring or multiple condensed rings, which may optionally contain one or more olefinic bonds, unless otherwise constrained by the definition. Such cycloalkyl groups can include, for example, single ring structures, including cyclopropyl, cyclobutyl, cyclooctyl, cyclopentenyl, and the like or multiple ring structures, including adamantanyl, and bicyclo [2.2.1] heptane or cyclic alkyl groups to which is fused an aryl group, for example, indane, and the like. Spiro and fused ring structures can also be included. Cycloalkyl groups may be substituted further with one or

- more substituents selected from alkyl, alkenyl, alkynyl, alkoxy, cycloalkyl, cycloalkenyl, acyl, acylamino, acyloxy, alkoxycarbonylamino, azido, cyano, halogen, hydroxy, oxo, thiocarbonyl, carboxy, carboxyalkyl, arylthio, thiol, alkylthio, aryl, aralkyl, aryloxy, aminosulfonyl, aminocarbonylamino, $-NR_{\lambda}R_{\pi}$, $-NHC(=O)NR_{\lambda}R_{\pi}$, $-NHC(=O)R_{\lambda}$, -
- 5 $C(=O)NR_{\lambda}R_{\pi}$, $-O-C(=O)NR_{\lambda}R_{\pi}$, nitro, heterocyclyl, heteroaryl, heterocyclalkyl, heteroarylalkyl or SO_mR_{ψ} (wherein R_{λ} , R_{π} , m and R_{ψ} are the same as defined earlier). Unless otherwise constrained by the definition, cycloalkyl substituents optionally may be substituted further by 1-3 substituents selected from alkyl, alkenyl, alkynyl, carboxy, hydroxy, alkoxy, halogen, CF_3 , $-NR_{\lambda}R_{\pi}$, $-C(=O)NR_{\lambda}R_{\pi}$, $-NHC(=O)NR_{\lambda}R_{\pi}$, -
- 10 $OC(=O)NR_{\lambda}R_{\pi}$, cyano or $-SO_mR_{\psi}$ (wherein R_{λ} , R_{π} , m and R_{ψ} are the same as defined earlier). "Cycloalkylalkyl" refers to alkyl-cycloalkyl group linked through alkyl portion, wherein the alkyl and cycloalkyl are the same as defined earlier.

- The term "aralkyl," unless otherwise specified, refers to alkyl-aryl linked through an alkyl portion (wherein alkyl is as defined above) and the alkyl portion contains 1-6
- 15 carbon atoms and aryl is as defined below. Examples of aralkyl groups include benzyl, ethylphenyl, propylphenyl, naphthylmethyl and the like.

- The term "aryl," unless otherwise specified, refers to aromatic system having 6 to 14 carbon atoms, wherein the ring system can be mono-, bi- or tricyclic and are carbocyclic aromatic groups. For example, aryl groups include, but are not limited to,
- 20 phenyl, biphenyl, anthryl or naphthyl ring and the like, optionally substituted with 1 to 3 substituents selected from halogen (*e.g.*, F, Cl, Br, I), hydroxy, alkyl, alkenyl, alkynyl, cycloalkyl, alkoxy, acyl, aryloxy, CF_3 , cyano, nitro, $COOR_{\psi}$, $NHC(=O)R_{\lambda}$, $-NR_{\lambda}R_{\pi}$, - $C(=O)NR_{\lambda}R_{\pi}$, $-NHC(=O)NR_{\lambda}R_{\pi}$, $-O-C(=O)NR_{\lambda}R_{\pi}$, $-SO_mR_{\psi}$, carboxy, heterocyclyl, heteroaryl, heterocyclalkyl, heteroarylalkyl or amino carbonyl amino, mercapto,
- 25 haloalkyl, optionally substituted aryl, optionally substituted heterocyclalkyl, thioalkyl, - $CONHR_{\pi}$, $-OCOR_{\pi}$, $-COR_{\pi}$, $-NH SO_2R_{\pi}$ or $-SO_2NHR_{\pi}$ (wherein R_{λ} , R_{π} , m and R_{ψ} are the same as defined earlier). Aryl groups optionally may be fused with a cycloalkyl group, wherein the cycloalkyl group may optionally contain heteroatoms selected from O, N or S. Groups such as phenyl, naphthyl, anthryl, biphenyl, and the like exemplify this term.

The term “aryloxy” denotes the group O-aryl wherein aryl is the same as defined above.

The term “heteroaryl,” unless otherwise specified, refers to an aromatic ring structure containing 5 or 6 ring atoms or a bicyclic or tricyclic aromatic group having from 8 to 10 ring atoms, with one or more heteroatom(s) independently selected from N, O or S optionally substituted with 1 to 4 substituent(s) selected from halogen (*e.g.*, F, Cl, Br, I), hydroxy, alkyl, alkenyl, alkynyl, cycloalkyl, acyl, carboxy, aryl, alkoxy, aralkyl, cyano, nitro, heterocyclyl, heteroaryl, $-NR_{\lambda}R_{\pi}$, $CH=NOH$, $-(CH_2)_wC(=O)R_{\eta}$ {wherein *w* is an integer from 0-4 and R_{η} is hydrogen, hydroxy, OR_{λ} , $NR_{\lambda}R_{\pi}$, $-NHOR_{\omega}$ or $-NHOH$ }, $-C(=O)NR_{\lambda}R_{\pi}$, $-NHC(=O)NR_{\lambda}R_{\pi}$, $-SO_mR_{\psi}$, $-O-C(=O)NR_{\lambda}R_{\pi}$, $-O-C(=O)R_{\lambda}$, or $-O-C(=O)OR_{\lambda}$ (wherein *m*, R_{ψ} , R_{λ} and R_{π} are as defined earlier and R_{ω} is alkyl, cycloalkyl, aryl, heteroaryl, heterocyclyl, heteroarylalkyl or heterocyclylalkyl). Unless otherwise constrained by the definition, the substituents are attached to a ring atom, *i.e.*, carbon or heteroatom in the ring. Examples of heteroaryl groups include oxazolyl, imidazolyl, pyrrolyl, 1,2,3-triazolyl, 1,2,4-triazolyl, tetrazolyl, thiazolyl, oxadiazolyl, benzoimidazolyl, thiadiazolyl, pyridinyl, pyridazinyl, pyrimidinyl, pyrazinyl, thienyl, isoxazolyl, triazinyl, furanyl, benzofuranyl, indolyl, benzthiazinyl, benzthiazinonyl, benzoxazinyl, benzoxazinonyl, quinazonyl, carbazolyl phenothiazinyl, phenoxazinyl, benzothiazolyl or benzoxazolyl, and the like.

The term “heterocyclyl,” unless otherwise specified, refers to a non-aromatic monocyclic or bicyclic cycloalkyl group having 5 to 10 atoms wherein 1 to 4 carbon atoms in a ring are replaced by heteroatoms selected from O, S or N, and optionally are benzofused or fused heteroaryl having 5-6 ring members and/or optionally are substituted, wherein the substituents are selected from halogen (*e.g.*, F, Cl, Br, I), hydroxy, alkyl, alkenyl, alkynyl, cycloalkyl, acyl, optionally substituted aryl, alkoxy, alkaryl, cyano, nitro, oxo, carboxy, optionally substituted heterocyclyl, optionally substituted heterocyclylalkyl, optionally substituted heteroaryl, $-O-C(=O)R_{\lambda}$, $-O-C(=O)OR_{\lambda}$, $-C(=O)NR_{\lambda}R_{\pi}$, SO_mR_{ψ} , $-O-C(=O)NR_{\lambda}R_{\pi}$, $-NHC(=O)NR_{\lambda}R_{\pi}$, $-NR_{\lambda}R_{\pi}$, mercapto, haloalkyl, thioalkyl, $-COOR_{\psi}$, $-COONHR_{\lambda}$, $-COR_{\lambda}$, $-NHSO_2R_{\lambda}$ or SO_2NHR_{λ} (wherein *m*, R_{ψ} , R_{λ} and R_{π} are as defined earlier) or guanidine. Heterocyclyl can optionally include rings having one or more

double bonds. Such ring systems can be mono-, bi- or tricyclic. Carbonyl or sulfonyl group can replace carbon atom(s) of heterocyclyl. Unless otherwise constrained by the definition, the substituents are attached to the ring atom, *i.e.*, carbon or heteroatom in the ring. Also, unless otherwise constrained by the definition, the heterocyclyl ring optionally
5 may contain one or more olefinic bond(s). Examples of heterocyclyl groups include benzotriazinone, isoindoledione, pyrimidinedione, aza-spiro[4.5]decanedione, benzo-oxazinedione, imidazolidinedione, phthalazinone, oxazolidinyl, tetrahydrofuranyl, dihydrofuranyl, benzoxazinyl, benzthiazinyl, imidazolyl, benzimidazolyl, tetrazolyl, carbaxolyl, indolyl, phenoxazinyl, phenothiazinyl, dihydropyridinyl, dihydroisoxazolyl,
10 dihydrobenzofuryl, azabicyclohexyl, thiazolidinyl, dihydroindolyl, pyridinyl, isoindole 1,3-dione, piperidinyl, tetrahydropyranyl, piperazinyl, 3H-imidazo[4,5-b]pyridine, isoquinolinyl, 1H-pyrrolo[2,3-b]pyridine or piperazinyl and the like.

The term “*cycloalkylalkyl*” refers to cycloalkyl group linked through alkyl portion, wherein the alkyl having 1 to 6 carbon atoms and cycloalkyl are the same as defined
15 earlier.

The term “*Heteroarylalkyl*” refers to heteroaryl group linked through alkyl portion, wherein the alkyl having 1 to 6 carbon atoms and heteroaryl are the same as defined earlier.

The term “*Heterocyclylalkyl*” refers to heterocyclyl group linked through alkyl
20 portion, wherein the alkyl having 1 to 6 carbon atoms and heterocyclyl are the same as defined earlier.

The term “*amino*” refers to $-\text{NH}_2$

The term “*acyl*” refers to $-\text{C}(=\text{O})\text{R}_4$ wherein R_4 is the same as defined earlier.

The term “*thioacyl*” refers to $-\text{C}(=\text{S})\text{R}_4$ wherein R_4 is the same as defined above;

25 The term “*halogen*” refers to fluorine, chlorine, bromine or iodine;

The term “leaving group” refers to groups that exhibit or potentially exhibit the properties of being labile under the synthetic conditions and also, of being readily separated from synthetic products under defined conditions. Examples of leaving groups

include, but are not limited to, halogen (*e.g.*, F, Cl, Br, I), triflates, tosylate, mesylates, alkoxy, thioalkoxy, or hydroxy radicals and the like.

5 The term “protecting groups” refers to moieties that prevent chemical reaction at a location of a molecule intended to be left unaffected during chemical modification of such molecule. Unless otherwise specified, protecting groups may be used on groups, such as hydroxy, amino, or carboxy. Examples of protecting groups are found in T.W. Greene and P.G.M. Wuts, “Protective Groups in Organic Synthesis”, 2nd Ed., John Wiley and Sons, New York, N.Y., which is incorporated herein by reference. The species of the carboxylic protecting groups, amino protecting groups or hydroxy protecting groups employed are
10 not critical, as long as the derivatised moieties/moiety is/are stable to conditions of subsequent reactions and can be removed without disrupting the remainder of the molecule.

Compounds described herein can contain one or more asymmetric carbon atoms and thus occur as diastereomers. These compounds can also exist as conformers/rotamers.
15 All such isomeric forms of these compounds are included herein. Each stereogenic carbon may be of the *R* or *S* configuration. Although the specific compounds exemplified in this application may be depicted in a particular stereochemical configuration, compounds having either the opposite stereochemistry at any given chiral center or mixtures thereof are envisioned.

20 The term “*pharmaceutically acceptable salts*” includes the salts of carboxylic acid moiety, which may be prepared by reacting the compound with appropriate base to provide corresponding base addition salts. Examples of such bases are alkali metal hydroxide including potassium hydroxide, sodium hydroxide and lithium hydroxide; alkaline earth metal hydroxides such as magnesium hydroxide and calcium hydroxide.
25 Further the salts of organic bases such as lysine, arginine, guanidine, ethanolamine, choline and the like; inorganic bases *e.g.*, ammonium or substituted ammonium salts are also included. Wherever appropriate, compounds of the present invention may also form the acid addition salts by treating the said compounds with pharmaceutically acceptable organic and inorganic acids, *e.g.*, hydrohalides such as

hydrochloride, hydrobromide, hydroiodide; other mineral acids and their corresponding salts such as sulphate, nitrate, phosphate etc.; and alkyl and mono-arylsulphonates such as ethane sulphonate, toluene sulphonate and benzene sulphonate; and other organic acids and their corresponding salts such as acetate, tartarate, maleate, succinate, citrate etc.

5 The salt forms differ from the compound described herein in certain physical properties such as solubility, but the salts are otherwise equivalent for the purpose of this invention.

 The term “*pharmaceutically acceptable solvates*” refers to solvates with water (i.e., hydrates) or pharmaceutically acceptable solvents, for example solvates with ethanol and the
10 like. Such solvates are also encompassed within the scope of the disclosure. Furthermore, some of the crystalline forms for compounds described herein may exist as *polymorphs* and as such are intended to be included in the scope of the disclosure.

 The term “polymorphs” includes all crystalline form as well as amorphous forms for compounds described herein and as such are included in the present invention.

15 The phrase “pharmaceutically acceptable carriers” is intended to include non-toxic, inert solid, semi-solid or liquid filler, diluent, encapsulating material or formulation auxiliary of any type.

 The term “pharmaceutically acceptable” means approved by regulatory agency of the federal or a state government or listed in the U.S. Pharmacopoeia or other generally
20 recognized pharmacopoeia for use in animals, and more particularly in humans.

 Examples of inflammatory conditions and autoimmune disorders in which the compounds prepared according to the invention have potentially beneficial effects in treatment methods may include, but are not limited to diseases of the respiratory tract such as asthma (including allergen-induced asthmatic reactions), cystic fibrosis, bronchitis (including chronic
25 bronchitis), chronic obstructive pulmonary disease (COPD), adult respiratory distress syndrome (ARDS), chronic pulmonary inflammation, rhinitis and upper respiratory tract inflammatory disorders (URID), ventilator induced lung injury, silicosis, pulmonary sarcoidosis, idiopathic pulmonary fibrosis, bronchopulmonary dysplasia, arthritis, *e.g.*,


rheumatoid arthritis, osteoarthritis, infectious arthritis, psoriatic arthritis, traumatic arthritis, rubella arthritis, Reiter's syndrome, gouty arthritis and prosthetic joint failure, gout, acute synovitis, spondylitis and non-articular inflammatory conditions, *e.g.*, herniated/ruptured/prolapsed intervertebral disk syndrome, bursitis, tendonitis,

5 tenosynovitic, fibromyalgic syndrome and other inflammatory conditions associated with ligamentous sprain and regional musculoskeletal strain, inflammatory disorders of the gastrointestinal tract, *e.g.*, ulcerative colitis, diverticulitis, Crohn's disease, inflammatory bowel diseases, irritable bowel syndrome and gastritis, multiple sclerosis, systemic lupus erythematosus, scleroderma, autoimmune exocrinopathy, autoimmune encephalomyelitis, 10 diabetes, tumor angiogenesis and metastasis, cancer including carcinoma of the breast, colon, rectum, lung, kidney, ovary, stomach, uterus, pancreas, liver, oral, laryngeal and prostate, melanoma, acute and chronic leukemia, periodontal disease, neurodegenerative disease, Alzheimer's disease, Parkinson's disease, epilepsy, muscle degeneration, inguinal hernia, retinal degeneration, diabetic retinopathy, macular degeneration, inguinal hernia, 15 retinal degeneration, diabetic retinopathy, macular degeneration, ocular inflammation, bone resorption diseases, osteoporosis, osteopetrosis, graft vs. host reaction, allograft rejections, sepsis, endotoxemia, toxic shock syndrome, tuberculosis, usual interstitial and cryptogenic organizing pneumonia, bacterial meningitis, systemic cachexia, cachexia secondary to infection or malignancy, cachexia secondary to acquired immune deficiency 20 syndrome (AIDS), malaria, leprosy, leishmaniasis, Lyme disease, glomerulonephritis, glomerulosclerosis, renal fibrosis, liver fibrosis, pancreatitis, hepatitis, endometriosis, pain, *e.g.*, that associated with inflammation and/or trauma, inflammatory diseases of the skin, *e.g.*, dermatitis, dermatosis, skin ulcers, psoriasis, eczema, systemic vasculitis, vascular dementia, thrombosis, atherosclerosis, restenosis, reperfusion injury, plaque 25 calcification, myocarditis, aneurysm, stroke, pulmonary hypertension, left ventricular remodeling and heart failure. It will be appreciated by those skilled in the art that reference herein to treatment extends to prophylaxis as well as the treatment of established conditions.

Compounds disclosed herein may be prepared, for example, by techniques well 30 known in the organic synthesis and familiar to a practitioner ordinarily skilled in art of this

invention. In addition, the processes described herein may enable the synthesis of the compounds of the present invention. However, these may not be the only means by which the compounds described in the invention may be synthesized. Further, the various synthetic steps described herein may be performed in alternate sequences in order to furnish the desired compounds.

Scheme-I



Compounds of Formulae VIIIa and VIIIb can be prepared by following, for example, synthetic routes as depicted in Scheme I. Thus, a compound of Formula II can be converted to a compound of Formula III (wherein P₁ is a silyl protecting group for example, *tert*-butyldimethylsilane, *tert*-butyldiphenylsilane or triisopropylsilane), which can then be converted to a compound of Formula IV (wherein P₁ is defined as above, R' and R'' together form an acetal protecting group, for example, isopropylidene or cyclohexylidene acetal). The compound of Formula IV can be oxidized to form a compound of Formula V. The compound of Formula V can then be converted to a compound of Formula VI (wherein E is an alkyl group such as methyl, ethyl and the like), which can then be hydrogenated to

form compounds of Formulae VIIa and VIIb. Compounds of Formulae VIIa and VIIb can be reduced to form compounds of Formulae VIIIa and VIIIb.

Silylation of a compound of Formula II to form a compound of Formula III can be carried out with silylating reagents such as, for example, tert-butyldimethylchlorosilane, triphenylchlorosilane, t-butyldiphenylchlorosilane in presence of a base, for example, imidazole or triethylamine in an organic solvent for example, dimethyl formamide, dimethylsulphoxide or acetonitrile.

Acetonation of a compound of Formula III to form a compound of Formula IV can be carried out with excess acetone as a solvent, in the presence of a mild acid catalyst, for example, anhydrous copper (II) sulphate and camphorsulphonic acid, anhydrous zinc chloride and small amount of phosphoric acid, or anhydrous ferric chloride. Alternately, one may utilize trans-acetonation with dimethoxypropane in the presence of acid catalysts such as p-toluenesulfonic acid, sulfuric acid or montmorillonite-K.

The compound of Formula IV can be oxidized to form a compound of Formula V, for example, by using Swern oxidation (Dimethylsulphoxide and oxalyl chloride) or Corey-Kim oxidation (N-chlorosuccinimide and Dimethylsulphide) in presence of a base, for example, triethylamine, diisopropylethylamine in a solvent, for example, dichloromethane or toluene. Alternatively, the compound of Formula IV can be oxidized to a compound of Formula V in, for example, dichloromethane or chloroform with an oxidizing agent, such as Dess-Martin reagent, pyridinium chlorochromate (PCC) or pyridinium dichromate (PDC). Oxidation of the compound of Formula IV to form a compound of Formula V can also be carried out catalytically using, for example, 2,2,6,6,-tetramethylpiperidine N-oxyl (TEMPO) and the 4-substituted derivatives thereof, including for example, 4-methoxy-TEMPO, 4-ethoxy-TEMPO, 4-acetoxy-TEMPO, 4-acetamino-TEMPO, 4-hydroxy-TEMPO, 4-benzoyloxy-TEMPO, 4-amino -TEMPO, N,N-dimethylamino-TEMPO, or 4-oxo-TEMPO as a catalyst, in the presence of a co-catalyst, for example, potassium bromide or sodium bromide, with an oxidant, for example, sodium hypochlorite, potassium hypochlorite, calcium hypochlorite, sodium hypobromite or potassium hypobromite in a solvent, for example, methylene chloride, chloroform, ethyl


acetate, butyl acetate, acetonitrile, tetrahydrofuran, toluene, acetone, diethyl ether, methyl tert-butyl ether, pentane, hexane or mixtures of such solvents.

The compound of Formula V can be converted to a compound of Formula VI, for example, via a Horner-Wadsworth-Emmons reaction, thus a compound of Formula V can
5 be reacted with phosphonate carbanions (produced *in situ* by treating trimethylphosphonoacetate or triethylphosphonoacetate with a base, for example, sodium hydride, potassium hydride, potassium tert-butoxide, sodium tert-butoxide, potassium carbonate, triethylamine in an organic solvent, for example, tetrahydrofuran or dimethoxyethane to give a compound of Formula VI. Alternately, the Wittig reaction can
10 be carried out using the preformed Wittig reagent such as (carboethoxymethylene)-triphenylphosphorane.

Hydrogenation of a compound of Formula VI to form compounds of Formulae VIIa and VIIb can be carried with palladium on carbon in presence of hydrogen, in a suitable solvent, for example, methanol, ethanol, propanol, tetrahydrofuran, ethyl acetate
15 or mixture(s) thereof.

Compounds of Formulae VIIa and VIIb can be reduced to give compounds of Formula VIIIa and VIIIb in the presence of a reducing agent, for example, lithium aluminium hydride, lithium triethyl borohydride or sodium borohydride, in the presence of an additive, for example, lithium chloride or aluminium chloride, in an organic solvent, for
20 example, tetrahydrofuran, diethylether or diglyme.

Scheme-II



Compounds of Formula XX can be prepared, for example, by following synthetic routes as depicted in Scheme II. Thus, a compound of Formula VIIIa (wherein P₁, R' and R'' are same as defined earlier), can react with a compound of Formula IX (wherein R₂ is an N-containing heterocyclyl or heteroaryl) to give a compound of Formula X. The compound of Formula X can undergo deprotection to form a compound of Formula XI which on oxidation can give a compounds of Formula XII. The compound of Formula XII on reaction with a compound of Formula XIII (wherein Q is same as defined earlier, hal is Cl, Br or I and Ar is phenyl) can form a compound of Formula XIV. The compound of Formula XIV can then react with a compound of Formula XV (wherein R₁ is same as

defined earlier) to form a compound of Formula XVI which can then be hydrogenated to form a compound of Formula XVII. The compound of Formula XVII can undergo deprotection to form a compound of Formula XVIII which can then be oxidatively cleaved to form a compound of Formula XIX. The compound of Formula XIX can be
5 deformylated to form a compound of Formula XX.

The reaction of a compound of Formula VIIIa with a compound of Formula IX to give a compound of Formula X can be carried out using triphenylphosphine or tributylphosphine and diethyl azodicarboxylate, diisopropyl azodicarboxylate or 1,1'-azodicarbonyldipiperidine in an organic solvent, for example, tetrahydrofuran,
10 dimethylformamide or toluene.

The compound of Formula X can be deprotected to form a compound of Formula XI with deprotecting agents, for example, tetrabutylammonium fluoride or potassium fluoride in an organic solvent, for example, tetrahydrofuran, dimethylformamide, diethyl ether or dioxane, optionally in the presence of crown ethers such as, for example, 18-crown-6. The oxidation of a compound of Formula XI to give a compound of Formula
15 XII can be carried out similarly as that of conversion of a compound of Formula IV to form a compound of Formula V.

The compound of Formula XII can be converted to a compound of Formula XIV by reacting with a compound of Formula XIII (Wittig reagent, *i.e.*, an ylide, prepared by
20 reacting a phosphonium salt, in turn prepared from triphenylphosphine and alkyl halide, in a solvent for example, tetrahydrofuran, dimethyl sulphoxide or diethyl ether, with a strong base, for example, *n*-butyllithium, sodium hydride or potassium *tert*-butoxide).

The reaction of a compound of Formula XIV with a compound of Formula XV can be carried out in the presence of a metal catalyst, for example,
25 tetrakis(triphenylphosphine) palladium (0), tetrakis(tricyclohexylphosphine) palladium (0), tetrakis(tri-*tert*-butylphosphine) palladium (0) or palladium acetate and triphenylphosphine in the presence of a base for example, potassium carbonate or cesium carbonate, in an organic solvent, for example, toluene, dimethyl sulphoxide, dimethylformamide, tetrahydrofuran, dioxane or diethyl ether.


Hydrogenation of a compound of Formula XVI to form a compound of Formula XVII can be carried out similarly as conversion of a compound of Formula VI to compounds of Formulae VIIa and VIIb.

- 5 The compound of Formula XVII can be deprotected to form a compound of Formula XVIII with perchloric acid, acetic acid or hydrochloric acid in solvent(s), for example, acetonitrile, water, tetrahydrofuran or mixture(s) thereof.

- 10 Conversion of a compound of Formula XVIII to form a compound of Formula XIX can be carried out by diol cleavage in presence of, for example, sodium metaperiodate, lead tetraacetate, pyridinium chlorochromate or manganese dioxide, in co-solvents, for example, tert-butanol-water, methanol-tetrahydrofuran or tert-butanol-tetrahydrofuran, followed by oxidation with, for example, potassium permanganate or with a mixture of sodium dihydrogen phosphate, sodium chlorite and hydrogen peroxide.

- 15 The compound of Formula XIX can be deformylated to form a compound of Formula XX in presence of a base for example, potassium carbonate, sodium carbonate or triethylamine in a solvent, for example, methanol, tetrahydrofuran or mixture(s) thereof.

Scheme-III




Compounds of Formula XXVI can be prepared by following synthetic routes, for example, as depicted in Scheme III. Thus, a compound of Formula XII can react with a compound of Formula XXI (wherein Q is same as defined earlier and Ar is phenyl) to form a compound of Formula XXII (wherein R₂ is an N-containing heterocyclyl or heteroaryl), which can further be hydrogenated to form a compound of Formula XXIII. The compound of Formula XXIII can be deprotected to form a compound of Formula XXIV, which can then be oxidatively cleaved to give a compound of Formula XXV. The compound of Formula XXV can then be deformylated to form a compound of Formula XXVI.

The reaction of a compound of Formula XII with a compound of Formula XXI to form a compound of Formula XXII can be carried out similarly as conversion of a compound of Formula XII to a compound of Formula XIV. Hydrogenation of a compound of Formula XXII to give a compound of Formula XXIII can be carried out under similar conditions as that of conversion of a compound of Formula VI to compounds of Formulae VIIa and VIIb.

Deprotection of a compound of Formula XXIII to give a compound of Formula XXIV can be carried out similarly as conversion of a compound of Formula XVII to a compound of Formula XVIII. The oxidative cleavage of a compound of Formula XXIV to form a compound of Formula XXV can be carried out under similar condition as that of conversion of a compound of Formula XVIII to a compound of Formula XIX.

Deformylation of a compound of Formula XXV to form a compound of Formula XXVI can be carried out similarly as conversion of a compound of Formula XIX to give a compound of Formula XX.

Scheme-IV



Compounds of Formula XXXV can be prepared by, for example, following synthetic routes as depicted in Scheme IV. Thus, a compound of Formula VIIIb (wherein

5 P₁, R' and R'' are same as defined earlier), can react with a compound of Formula IX (wherein R₂ is an N-containing heterocyclyl or heteroaryl to give a compound of Formula XXVII. The compound of Formula XXVII can be deprotected to form a compound of Formula XXVIII, which on oxidation can give a compound of Formula XXIX. The compound of Formula XXIX on reaction with a compound of Formula XIII can form a

10 compound of Formula XXX which can then react with a compound of Formula XV (wherein R₁ is same as defined earlier) to form a compound of Formula XXXI. The compound of Formula XXXI can then be hydrogenated to form a compound of Formula XXXII which can then undergo deprotection to form a compound of Formula XXXIII. The compound of Formula XXXIII can be oxidatively cleaved to form

a compound of Formula XXXIV. The compound of Formula XXXIV can then be deformylated to form a compound of Formula XXXV.

5 The reaction of a compound of Formula VIIIb with a compound of Formula IX to give a compound of Formula XXVII can be carried out similarly as that of conversion of a compound of Formula VIIa to form a compound of Formula X.

Deprotection of a compound of Formula XXVII to give a compound of Formula XXVIII can be carried out similarly as that of deprotection of a compound of Formula X to form a compound of Formula XI.

10 Oxidation of a compound of Formula XXVIII to give a compound of Formula XXIX can be carried out similarly as that of conversion of a compound of Formula IV to form a compound of Formula V.

The reaction of a compound of Formula XXIX with a compound of Formula XIII to form a compound of Formula XXX can be carried out under similar condition as that of conversion of a compound of Formula XII to form a compound of Formula XIV.


15 Coupling of a compound of Formula XXX with a compound of Formula XV to form a compound of Formula XXXI can be carried out similarly as conversion of a compound of Formula XIV to form a compound of Formula XVI.

20 Hydrogenation of a compound of Formula XXXI to form a compound of Formula XXXII can be carried out similarly as conversion of a compound of Formula VI to form compound of Formulae VIIa and VIIb. The compound of Formula XXXII can be deprotected to form a compound of Formula XXXIII under similar condition as that of conversion of a compound of Formula XVII to form a compound of Formula XVIII.

25 Oxidative cleavage of a compound of Formula XXXIII to form a compound of Formula XXXIV can be carried out similarly as conversion of a compound of Formula XVIII to give a compound of Formula XIX.

Deformylation of a compound of Formula XXXIV to form a compound of Formula XXXV can be carried out under similar condition as that of conversion of a compound of Formula XIX to form a compound of Formula XX.

Scheme-V



Compounds of Formula LIII can be prepared, for example, by following synthetic routes as depicted in Scheme V. Thus, a compound of Formula XXXVI (wherein R₇ and R₈ together form a acetal protecting group, for example, isopropylidene acetal, R' and R'' are same as described earlier) can be oxidized to form a compound of Formula XXXVII which can then be converted to form a compound of Formula XXXVIII (wherein R₉ is alkyl or aryl). The compound of Formula XXXVIII can be hydrogenated to give a compound of Formula XXXIX which can then be hydrolyzed to give a compound of Formula XL. The compound of Formula XL can be oxidized to give a compound of Formula XLI which can then be converted to form a compound of Formula XLII. The compound of Formula XLII can be hydrogenated to form a compound of Formula XLIII which can then be reduced to form a compound of Formula XLIV. The compound of Formula XLIV can then be reacted with a compound of Formula IX (wherein R₂ is an N-containing heterocyclyl or heteroaryl) to afford a compound of Formula XLV which can then be deprotected to form a compound of Formula XLVI. The compound of Formula XLVI can be oxidized to form a compound of Formula XLVII which can then react with a compound of Formula XIII to form a compound of Formula XLVIII. The compound of Formula XLVIII can then be reacted with a compound of Formula XV (wherein R₁ is same as defined earlier) to give a compound of Formula XLIX which can then be hydrogenated to form a compound of Formula L. The compound of Formula L is deprotected to form a compound of Formula LI which can then be oxidatively cleaved to form a compound of Formula LII. The compound of Formula LII is then deformylated to form a compound of Formula LIII.

Oxidation of a compound of Formula XXXVI to form a compound of Formula XXXVII can be carried out under similar condition as that of oxidation of a compound of Formula IV to form a compound of Formula V. The compound of Formula XXXVII can be converted to a compound of Formula XXXVIII in presence of, for example, acetic anhydride or benzoic anhydride in presence of base, such as for example, pyridine, triethylamine or morpholine.

The compound of Formula XXXVIII can be hydrogenated to form a compound of Formula XXXIX similarly as that of hydrogenation of a compound of Formula VI to give

compounds of Formulae VIIa and VIIb. The hydrolysis of a compound of Formula XXXIX to form a compound of Formula XL can be carried out with a base, for example, sodium hydroxide, potassium hydroxide, sodium carbonate, potassium carbonate or sodium methoxide in the presence of a solvent, for example, methanol, ethanol or isopropanol. The compound of Formula XL can be oxidized to form a compound of Formula XLI similarly as that of conversion of a compound of Formula IV to form a compound of Formula V.

Conversion of a compound of Formula XLI to form a compound of Formula XLII can be carried out under similar condition as that of conversion of a compound of Formula V to form a compound of Formula VI. The compound of Formula XLII can be hydrogenated to form a compound of Formula XLIII similarly as that of conversion of compound of Formula VI to compounds of Formulae VIIa and VIIb. The reduction of a compound of Formula XLIII to form a compound of Formula XLIV can be carried out under similar condition as that of reduction of compounds of Formulae VIIa and VIIb to form compounds of Formulae VIIIa and VIIIb. The reaction of a compound of Formula XLIV with a compound of Formula IX to form compound of Formula XLV can be carried out similarly as that of conversion of a compound of Formula VIIIa to form a compound of Formula X.

Selective acetonide deprotection of a compound of Formula XLV to form a compound of Formula XLVI can be carried out with perchloric acid in a solvent, for example, tetrahydrofuran or diethyl ether or trifluoroacetic acid in dichloromethane. Oxidation of a compound of Formula XLVI to form a compound of Formula XLVII can be carried out with, for example, sodium metaperiodate, lead tetraacetate, pyridinium chlorochromate or manganese dioxide in a solvent, for example, acetone, methanol, ethanol or tert-butanol.


The reaction of a compound of Formula XLVII with a compound of Formula XIII to form a compound of Formula XLVIII can be carried out similarly as that of conversion of compound of Formula XII to form a compound of Formula XIV. Coupling of a compound of Formula XLVIII with a compound of Formula XV to form a compound of

Formula XLIX can be carried out similarly as conversion of a compound of Formula XIV to form a compound of Formula XVI. The compound of Formula XLIX can be hydrogenated to form a compound of Formula L under similar condition as that of conversion of compound of Formula VI to compound of Formulae VIIa and VIIb.

- 5 Deprotection of a compound of Formula L to give a compound of Formula LI can be carried out under similar condition as that of conversion of a compound of Formula XVII to form a compound of Formula XVIII.

- 10 The oxidative cleavage of a compound of Formula LI to form a compound of Formula LII can be done similarly as that of conversion of a compound of Formula XVIII to form a compound of Formula XIX. The compound of Formula LII can be deformylated to a compound of Formula LIII under similar condition as that of conversion of compound of Formula XIX to form compound of Formula XX.

Scheme-VI



Compounds of Formula LXVI can be prepared, for example, by following synthetic routes as depicted in Scheme VI. Thus, a compound of Formula XXXVII (wherein R_7 , R_8 , R' and R'' are same as described earlier) can be converted to form a compound of Formula LIV which can then be hydrogenated to form a compound of Formula LV. The compound of Formula LV can be selectively deprotected to form a compound of Formula LVI which on oxidation, can form a compound of Formula LVII. The compound of Formula LVII can be reacted with a compound of Formula XIII to form a compound of Formula LVIII which can then be coupled with a compound of Formula XV (wherein R_1 is same as defined earlier) to give a compound of Formula LIX. The compound of Formula LIX can be hydrogenated to form a compound of Formula LX which can then be reduced to form a compound of Formula LXI. The compound of Formula LXI can be activated to form a compound of Formula LXII (wherein U is an O-

activating group, for example, mesyl, tosyl or triflate), which can then be reacted with a compound of Formula IXa (wherein R₂ is an N-containing heterocyclyl or heteroaryl and M is a metal, for example, potassium, lithium or sodium) to form a compound of Formula LXIII. The compound of Formula LXIII can then be deprotected to form a compound of
5 Formula LXIV which can be oxidatively cleaved to form a compound of Formula LXV. The compound of Formula LXV can be deformylated to form a compound of Formula LXVI.

The compound of Formula XXXVII can be converted to form a compound of Formula LIV under similar condition as that of conversion of a compound of Formula V to
10 form a compound of Formula VI. Hydrogenation of a compound of Formula LIV to form a compound of Formula LV can be carried out similarly as that of conversion of a compound of Formula VI to form compounds of Formulae VIIa and VIIb.

The compound of Formula LV can be selectively deprotected to form a compound of Formula LVI under similar conditions as that of conversion of a compound of Formula
15 XLV to give a compound of Formula XLVI. The compound of Formula LVI can be oxidized to form a compound of Formula LVII similarly as that of conversion of a compound of Formula XLVI to form a compound of Formula XLVII.

The reaction of a compound of Formula LVII with a compound of Formula XIII to form a compound of Formula LVIII can be carried out under similar conditions as that of
20 conversion of a compound of Formula XII to form a compound of Formula XIV. The compound of Formula LVIII can be coupled with a compound of Formula XV to give a compound of Formula LIX under similar condition as that of conversion of a compound of Formula XIV to form a compound of Formula XVI. Hydrogenation of a compound of Formula LIX to form a compound of Formula LX can be carried out under similar
25 condition as that of conversion of a compound of Formula VI to form compounds of Formulae VIIa and VIIb.


The compound of Formula LX can be reduced to form a compound of Formula LXI under similar condition as that of conversion of compound of Formulae VIIa and VIIb to form compounds of Formulae VIIIa and VIIIb.

A compound of Formula LXI can be activated to form a compound of Formula LXII in a solvent, for example, dichloromethane, toluene or dichloroethane, using a base, for example, triethylamine, diisopropylamine or N-methylmorpholine, using a suitable sulphonyl chloride, for example, methanesulphonyl chloride or p-toluene sulphonyl chloride. The reaction of a compound of Formula LXII with a compound of Formula IXa to yield a compound of Formula LXIII can be carried out in an organic solvent, for example, tetrahydrofuran, dimethyl sulphoxide, dimethylformamide, acetonitrile, dioxane, dimethylacetamide. Alternatively, the reaction of a compound of Formula LXII with a compound of Formula IX to yield a compound of Formula LXIII can be carried out in the presence of a base, for example, sodium hydride, potassium tert-butoxide, sodium (m)ethoxide in an organic solvent, for example, tetrahydrofuran, dimethyl sulphoxide, dimethylformamide, acetonitrile, dioxane, or dimethylacetamide. Alternatively, a compound of Formula LXI can be converted to a compound of Formula LXIII following similar protocols as that of the reaction of a compound of Formula VIIIa with a compound of Formula IX to give a compound of Formula X.

The compound of Formula LXIII can be deprotected to form a compound of Formula LXIV similarly as that of conversion of a compound of Formula XVII to form a compound of Formula XVIII.

The oxidative cleavage of a compound of Formula LXIV to give a compound of Formula LXV can be done similarly as that of conversion of a compound of Formula XVIII to form a compound of Formula XIX. The compound of Formula LXV can be deformylated to form a compound of Formula LXVI under similar condition as that of conversion of a compound of Formula XIX to form a compound of Formula XX.

Scheme-VII



Compounds of Formula XX can also be prepared, for example, by following alternate synthetic routes as depicted in Scheme VII. Thus, a compound of Formula LXVII can be coupled with a compound of Formula LXVIII to form a compound of Formula LXIX (wherein R_1 and Q are same as defined earlier), which can then be converted to a compound of Formula LXX (Path A)(wherein E is same as defined earlier).

Alternately, compound of Formula LXVIIIa can undergo esterification to gives a compound of Formula LXIXa, which can be coupled with compound of Formula XV to form compound of Formula LXX (PathB) (wherein E is same as defined earlier). The compound of Formula LXX can be hydrogenated to form compound of Formula LXXI, which can then be reduced to form a compound of Formula LXXII. The compound of Formula LXXII can be oxidized to form a compound of Formula LXXIII, which can then react with a compound of Formula LXXIV (wherein, when R₂ is N-containing heterocyclyl or heteroaryl, Y and W can be oxygen or sulphur, R₁₀ can be alkyl, aryl or aralkyl and n is as defined earlier) to form compound of Formula LXXV. The compound of Formula LXXII can be further hydrolysed to form a compound of Formula XX.

Coupling of a compound of Formula LXVII with a compound of Formula LXVIII to form a compound of Formula LXIX can be carried out similarly as conversion of a compound of Formula XIV to form a compound of Formula XVI.

Conversion of a compound of Formula LXIX to form a compound of Formula LXX can be carried out under similar condition as that of conversion of a compound of Formula V to form a compound of Formula VI. Esterification of compound of Formula LXVIIIa to gives a compound of Formula LXIXa can be carried out in a solvent, for example, methanol, ethanol, tert-butanol or benzyl alcohol with a halogenating agent, for example, thionyl chloride or oxalyl chloride. Coupling of a compound of Formula LXIXa with a compound of Formula XV to form a compound of Formula LXX can be carried out similarly as conversion of a compound of Formula XIV to form a compound of Formula XVI.

The compound of Formula LXX can be hydrogenated to form a compound of Formula LXXI similarly as that of conversion of compound of Formula VI to compounds of Formulae VIIa and VIIb. The reduction of a compound of Formula LXXI to form a compound of Formula LXXII can be carried out under similar condition as that of reduction of compounds of Formulae VIIa and VIIb to form compounds of Formulae VIIIa and VIIIb. The compound of Formula LXXII can be oxidized to form a compound

of Formula LXXIII similarly as that of oxidation of a compound of Formula IV to form a compound of Formula V.

The asymmetric aldol addition of a compound of Formula LXXIII with a compound of Formula LXXIV to form compound of Formula LXXIV can be carried out
5 by generating the enolates with titanium chloride, dibutyl boron triflate, dialkyl boron chloride or tin(II) triflate, in the presence of a base, for example, diisopropylethylamine, tetramethylethylenediamine, tributylamine, N-ethylpiperidine, 1,4-diazabicyclo[2.2.2]octane, 1,8-Diazabicyclo[5.4.0]undec-7-ene, tetramethylpropylenediamine, or (-) sparteine, in a solvent, for example, dichloromethane,
10 or diethyl ether.

Hydrolysis of a compound of Formula LXXIV to form a compound of Formula XX can be carried out with hydrogen peroxide and lithium hydroxide, in the presence of a solvent, for example, tetrahydrofuran, water or mixture(s) thereof.

Compound Nos. 1 to 81, 86-88, 90-95, 97-119, 121-142, 144-203, 211-213 and
15 226-232 were prepared following Schemes I and II.

(2S,3R)-3-hydroxy-2-[2-(4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-5-(4-pyrimidin-5-ylphenyl)pentanoic acid (Compound No.1),

(2S,3R)-2-[2-(5-*tert*-butyl-1,3-dioxo-1,3-dihydro-2*H*-isoindol-2-yl)ethyl]-3-hydroxy-5-[4-(6-methoxypyridin-3-yl)phenyl]pentanoic acid (Compound No. 2),

20 (2S,3R)-2-[2-(1,3-dioxo-1,3-dihydro-2*H*-isoindol-2-yl)ethyl]-3-hydroxy-5-[4-(6-methylpyridin-3-yl)phenyl]pentanoic acid (Compound No. 3),

(2S,3R)-2-[2-(1,3-dioxo-1,3-dihydro-2*H*-isoindol-2-yl)ethyl]-5-(4'-fluorobiphenyl-4-yl)-3-hydroxypentanoic acid (Compound No. 4),

(2S,3R)-5-(3',4'-difluorobiphenyl-4-yl)-2-[2-(1,3-dioxo-1,3-dihydro-2*H*-isoindol-2-yl)ethyl]-3-hydroxypentanoic acid (Compound No. 5),
25

(2S,3R)-3-hydroxy-5-[4-(6-methylpyridin-3-yl)phenyl]-2-[2-(4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)ethyl]pentanoic acid (Compound No. 6),

(2*S*,3*R*)-3-hydroxy-2-[2-(5-methyl-1,3-dioxo-1,3-dihydro-2*H*-isoindol-2-yl)ethyl]-5-[4-(6-methylpyridin-3-yl)phenyl]pentanoic acid (Compound No. 7),

(2*S*,3*R*)-5-(4'-fluorobiphenyl-4-yl)-3-hydroxy-2-[2-(5-methyl-1,3-dioxo-1,3-dihydro-2*H*-isoindol-2-yl)ethyl]pentanoic acid (Compound No. 8),

5 (2*S*,3*R*)-5-(4'-fluorobiphenyl-4-yl)-3-hydroxy-2-[2-(4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)ethyl]pentanoic acid (Compound No. 9),

(2*S*,3*R*)-3-hydroxy-5-(4'-methoxybiphenyl-4-yl)-2-[2-(4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)ethyl]pentanoic acid (Compound No. 10),

10 (2*S*,3*R*)-5-(3',4'-difluorobiphenyl-4-yl)-3-hydroxy-2-[2-(5-methyl-1,3-dioxo-1,3-dihydro-2*H*-isoindol-2-yl)ethyl]pentanoic acid (Compound No. 11),

(2*S*,3*R*)-5-(3',4'-difluorobiphenyl-4-yl)-3-hydroxy-2-[2-(4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)ethyl]pentanoic acid (Compound No. 12),

(2*S*,3*R*)-2-[2-(1,3-dioxo-1,3-dihydro-2*H*-isoindol-2-yl)ethyl]-3-hydroxy-5-(4'-methoxybiphenyl-4-yl)pentanoic acid (Compound No. 13),

15 (2*S*,3*R*)-2-[2-(1,3-dioxo-1,3-dihydro-2*H*-isoindol-2-yl)ethyl]-3-hydroxy-5-[4'-(trifluoromethoxy)biphenyl-4-yl]pentanoic acid (Compound No. 14),

(2*S*,3*R*)-5-(4'-chloro-3'-fluorobiphenyl-4-yl)-3-hydroxy-2-[2-(4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)ethyl]pentanoic acid (Compound No. 15),

20 (2*S*,3*R*)-2-[2-(5-*tert*-butyl-1,3-dioxo-1,3-dihydro-2*H*-isoindol-2-yl)ethyl]-5-(3',4'-difluorobiphenyl-4-yl)-3-hydroxypentanoic acid (Compound No. 16),

(2*S*,3*R*)-2-[2-(5-*tert*-butyl-1,3-dioxo-1,3-dihydro-2*H*-isoindol-2-yl)ethyl]-3-hydroxy-5-(4'-methoxybiphenyl-4-yl)pentanoic acid (Compound No. 17),

(2*S*,3*R*)-3-hydroxy-2-[2-(4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)ethyl]-5-[4'-(trifluoromethyl)biphenyl-4-yl]pentanoic acid (Compound No. 18),

25 (2*S*,3*R*)-2-[2-(5-*tert*-butyl-1,3-dioxo-1,3-dihydro-2*H*-isoindol-2-yl)ethyl]-5-(4'-fluorobiphenyl-4-yl)-3-hydroxypentanoic acid (Compound No. 19),

(2*S*,3*R*)-2-[2-(5-*tert*-butyl-1,3-dioxo-1,3-dihydro-2*H*-isoindol-2-yl)ethyl]-5-(4'-chlorobiphenyl-4-yl)-3-hydroxypentanoic acid (Compound No. 20),

(2*S*,3*R*)-2-[2-(1,3-dioxo-1,3-dihydro-2*H*-isoindol-2-yl)ethyl]-3-hydroxy-5-[4'-(trifluoromethyl)biphenyl-4-yl]pentanoic acid (Compound No. 21),

5 (2*S*,3*R*)-2-[2-(1,3-dioxo-1,3-dihydro-2*H*-isoindol-2-yl)ethyl]-3-hydroxy-5-(4'-methylbiphenyl-4-yl)pentanoic acid (Compound No. 22),

(2*S*,3*R*)-2-[2-(1,3-dioxo-1,3-dihydro-2*H*-isoindol-2-yl)ethyl]-5-(3'-fluoro-4'-methoxybiphenyl-4-yl)-3-hydroxypentanoic acid (Compound No. 23),

10 (2*S*,3*R*)-3-hydroxy-5-[4-(6-methoxypyridin-3-yl)phenyl]-2-[2-(5-methyl-1,3-dioxo-1,3-dihydro-2*H*-isoindol-2-yl)ethyl]pentanoic acid (Compound No. 24),

(2*S*,3*R*)-3-hydroxy-5-(4'-methylbiphenyl-4-yl)-2-[2-(4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)ethyl]pentanoic acid (Compound No. 25),

(2*S*,3*R*)-5-(4'-ethylbiphenyl-4-yl)-3-hydroxy-2-[2-(4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)ethyl]pentanoic acid (Compound No. 26),

15 (2*S*,3*R*)-5-(4'-chlorobiphenyl-4-yl)-3-hydroxy-2-[2-(5-methyl-1,3-dioxo-1,3-dihydro-2*H*-isoindol-2-yl)ethyl]pentanoic acid (Compound No. 27),

(2*S*,3*R*)-5-(4'-chlorobiphenyl-4-yl)-3-hydroxy-2-[2-(4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)ethyl]pentanoic acid (Compound No. 28),

20 (2*S*,3*R*)-2-[2-(1,3-dioxo-1,3-dihydro-2*H*-isoindol-2-yl)ethyl]-3-hydroxy-5-(4-pyrimidin-5-ylphenyl)pentanoic acid (Compound No. 29),

(2*S*,3*R*)-2-[2-(1,3-dioxo-1,3-dihydro-2*H*-isoindol-2-yl)ethyl]-3-hydroxy-5-[4-(1-methyl-1*H*-pyrazol-4-yl)phenyl]pentanoic acid (Compound No. 30),

(2*S*,3*R*)-2-[2-(1,3-dioxo-1,3-dihydro-2*H*-isoindol-2-yl)ethyl]-3-hydroxy-5-(4-pyridin-3-ylphenyl)pentanoic acid (Compound No. 31),

25 (2*S*,3*R*)-5-[4-(2,3-dihydro-1,4-benzodioxin-6-yl)phenyl]-3-hydroxy-2-[2-(4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)ethyl]pentanoic acid (Compound No. 32),

(2*S*,3*R*)-3-hydroxy-5-[4-(6-methoxypyridin-3-yl)phenyl]-2-[2-(8-methyl-4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)ethyl]pentanoic acid (Compound No. 33),

(2*S*,3*R*)-3-hydroxy-2-[2-(8-methyl-4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)ethyl]-5-[4-(6-methylpyridin-3-yl)phenyl]pentanoic acid (Compound No. 34),

5 (2*S*,3*R*)-3-hydroxy-5-[4-(1-methyl-1*H*-pyrazol-4-yl)phenyl]-2-[2-(4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)ethyl]pentanoic acid (Compound No. 35),

(2*S*,3*R*)-3-hydroxy-2-[2-(8-methoxy-4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)ethyl]-5-[4-(6-methoxypyridin-3-yl)phenyl]pentanoic acid (Compound No. 36),

10 (2*S*,3*R*)-2-[2-(7-chloro-4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)ethyl]-3-hydroxy-5-[4-(6-methoxypyridin-3-yl)phenyl]pentanoic acid (Compound No. 37),

(2*S*,3*R*)-3-hydroxy-2-{2-[7-(6-methoxypyridin-3-yl)-4-oxo-1,2,3-benzotriazin-3(4*H*)-yl]ethyl}-5-[4-(6-methoxypyridin-3-yl)phenyl]pentanoic acid (Compound No. 38),

(2*S*,3*R*)-2-[2-(6,7-difluoro-4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)ethyl]-3-hydroxy-5-[4-(6-methoxypyridin-3-yl)phenyl]pentanoic acid (Compound No. 39),

15 (2*S*,3*R*)-2-[2-(5-fluoro-4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)ethyl]-3-hydroxy-5-[4-(6-methoxypyridin-3-yl)phenyl]pentanoic acid (Compound No. 40),

(2*S*,3*R*)-3-hydroxy-2-[2-(8-methoxy-4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)ethyl]-5-[4-(6-methylpyridin-3-yl)phenyl]pentanoic acid (Compound No. 41),

20 (2*S*,3*R*)-3-hydroxy-2-{2-[5-(6-methoxypyridin-3-yl)-4-oxo-1,2,3-benzotriazin-3(4*H*)-yl]ethyl}-5-[4-(6-methoxypyridin-3-yl)phenyl]pentanoic acid (Compound No. 42),

(2*S*,3*R*)-5-(4'-chloro-3'-fluorobiphenyl-4-yl)-2-[2-(1,3-dioxo-1,3-dihydro-2*H*-isoindol-2-yl)ethyl]-3-hydroxypentanoic acid (Compound No. 43),

(2*S*,3*R*)-2-[2-(1,3-dioxo-1,3-dihydro-2*H*-isoindol-2-yl)ethyl]-3-hydroxy-5-[4-(1-isobutyl-1*H*-pyrazol-4-yl)phenyl]pentanoic acid (Compound No. 44),

25 (2*S*,3*R*)-5-biphenyl-4-yl-2-[2-(1,3-dioxo-1,3-dihydro-2*H*-isoindol-2-yl)ethyl]-3-hydroxypentanoic acid (Compound No. 45),

(2*S*,3*R*)-2-[2-(1,3-dioxo-1,3-dihydro-2*H*-isoindol-2-yl)ethyl]-5-(3'-fluoro-4'-methylbiphenyl-4-yl)-3-hydroxypentanoic acid (Compound No. 46),

(2*S*,3*R*)-5-(3,3'-difluoro-4'-methoxybiphenyl-4-yl)-2-[2-(1,3-dioxo-1,3-dihydro-2*H*-isoindol-2-yl)ethyl]-3-hydroxypentanoic acid (Compound No. 47),

5 (2*S*,3*R*)-5-[4-(2,3-dihydro-1,4-benzodioxin-6-yl)phenyl]-2-[2-(1,3-dioxo-1,3-dihydro-2*H*-isoindol-2-yl)ethyl]-3-hydroxypentanoic acid (Compound No. 48),

(2*S*,3*R*)-2-[2-(1,3-dioxo-1,3-dihydro-2*H*-isoindol-2-yl)ethyl]-3-hydroxy-5-[4-(1*H*-tetrazol-1-yl)phenyl]pentanoic acid (Compound No. 49),

10 (2*S*,3*R*)-2-[2-(1,3-dioxo-1,3-dihydro-2*H*-isoindol-2-yl)ethyl]-5-[2-fluoro-4-(6-methoxypyridin-3-yl)phenyl]-3-hydroxypentanoic acid (Compound No. 50),

(2*S*,3*R*)-5-(3'-fluoro-4'-methylbiphenyl-4-yl)-3-hydroxy-2-[2-(4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)ethyl]pentanoic acid (Compound No. 51),

(2*S*,3*R*)-3-hydroxy-5-[4-(1-isobutyl-1*H*-pyrazol-4-yl)phenyl]-2-[2-(4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)ethyl]pentanoic acid (Compound No. 52),

15 (2*S*,3*R*)-5-biphenyl-4-yl-3-hydroxy-2-[2-(4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)ethyl]pentanoic acid (Compound No. 53),

(2*S*,3*R*)-3-hydroxy-5-[4-(6-methoxypyridin-3-yl)phenyl]-2-[2-(7-methyl-4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)ethyl]pentanoic acid (Compound No. 54),

20 (2*S*,3*R*)-2-[2-(1,3-dioxo-1,3-dihydro-2*H*-isoindol-2-yl)ethyl]-5-[6-(3-fluoro-4-methoxyphenyl)pyridin-3-yl]-3-hydroxypentanoic acid (Compound No. 55),

(2*S*,3*R*)-5-(4'-chloro-3-fluorobiphenyl-4-yl)-2-[2-(1,3-dioxo-1,3-dihydro-2*H*-isoindol-2-yl)ethyl]-3-hydroxypentanoic acid (Compound No. 56),

(2*S*,3*R*)-2-[2-(1,3-dioxo-1,3-dihydro-2*H*-isoindol-2-yl)ethyl]-3-hydroxy-5-[6-(4-methoxyphenyl)pyridin-3-yl]pentanoic acid (Compound No. 57),

25 (2*S*,3*R*)-2-[2-(1,3-dioxo-1,3-dihydro-2*H*-isoindol-2-yl)ethyl]-5-(3-fluoro-4'-methoxybiphenyl-4-yl)-3-hydroxypentanoic acid (Compound No. 58),

(2*S*,3*R*)-5-[6-(4-chlorophenyl)pyridin-3-yl]-2-[2-(1,3-dioxo-1,3-dihydro-2*H*-isoindol-2-yl)ethyl]-3-hydroxypentanoic acid (Compound No. 59),

(2*S*,3*R*)-2-[2-(1,3-dioxo-1,3-dihydro-2*H*-isoindol-2-yl)ethyl]-3-hydroxy-5-(6'-methoxy-2,3'-bipyridin-5-yl)pentanoic acid (Compound No. 60),

5 (2*S*,3*R*)-5-[4-(4-chlorophenyl)-2-thienyl]-2-[2-(1,3-dioxo-1,3-dihydro-2*H*-isoindol-2-yl)ethyl]-3-hydroxypentanoic acid (Compound No. 61),

(2*S*,3*R*)-2-[2-(1,3-dioxo-1,3-dihydro-2*H*-isoindol-2-yl)ethyl]-3-hydroxy-5-[4-(6-methoxypyridin-3-yl)-2-thienyl]pentanoic acid (Compound No. 62),

10 (2*S*,3*R*)-2-[2-(1,3-dioxo-1,3-dihydro-2*H*-isoindol-2-yl)ethyl]-3-hydroxy-5-{4-[4-(trifluoromethyl)phenyl]-2-thienyl}pentanoic acid (Compound No. 63),

(2*S*,3*R*)-2-[2-(1,3-dioxo-1,3-dihydro-2*H*-isoindol-2-yl)ethyl]-5-[4-(3-fluoro-4-methoxyphenyl)-2-thienyl]-3-hydroxypentanoic acid (Compound No. 64),

(2*S*,3*R*)-3-hydroxy-2-[2-(7-methyl-4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)ethyl]-5-[4'-(trifluoromethyl)biphenyl-4-yl]pentanoic acid (Compound No. 65),

15 (2*S*,3*R*)-3-hydroxy-2-[2-(6-methyl-4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)ethyl]-5-[4'-(trifluoromethyl)biphenyl-4-yl]pentanoic acid (Compound No. 66),

(2*S*,3*R*)-2-[2-(7-chloro-4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)ethyl]-3-hydroxy-5-(4'-methylbiphenyl-4-yl)pentanoic acid (Compound No. 67),

20 (2*S*,3*R*)-3-hydroxy-2-[2-(8-methoxy-4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)ethyl]-5-[4'-(trifluoromethyl)biphenyl-4-yl]pentanoic acid (Compound No. 68),

(2*S*,3*R*)-2-[2-(6,7-difluoro-4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)ethyl]-3-hydroxy-5-[4'-(trifluoromethyl)biphenyl-4-yl]pentanoic acid (Compound No. 69),

(2*S*,3*R*)-2-[2-(1,3-dioxo-1,3-dihydro-2*H*-isoindol-2-yl)ethyl]-5-[2-fluoro-5-(6-methoxypyridin-3-yl)phenyl]-3-hydroxypentanoic acid (Compound No. 70),

25 (2*S*,3*R*)-5-(4'-chloro-4-fluorobiphenyl-3-yl)-2-[2-(1,3-dioxo-1,3-dihydro-2*H*-isoindol-2-yl)ethyl]-3-hydroxypentanoic acid (Compound No. 71),

(2S,3R)-2-[2-(1,3-dioxo-1,3-dihydro-2*H*-isoindol-2-yl)ethyl]-5-[6-fluoro-4'-(trifluoromethyl)biphenyl-3-yl]-3-hydroxypentanoic acid (Compound No. 72),

(2S,3R)-2-[2-(1,3-dioxo-1,3-dihydro-2*H*-isoindol-2-yl)ethyl]-5-[4-fluoro-3-(6-methoxypyridin-3-yl)phenyl]-3-hydroxypentanoic acid (Compound No. 73),

5 (2S,3R)-5-(4'-chloro-6-fluorobiphenyl-3-yl)-2-[2-(1,3-dioxo-1,3-dihydro-2*H*-isoindol-2-yl)ethyl]-3-hydroxypentanoic acid (Compound No. 74),

(2S,3R)-5-(3',6-difluoro-4'-methoxybiphenyl-3-yl)-2-[2-(1,3-dioxo-1,3-dihydro-2*H*-isoindol-2-yl)ethyl]-3-hydroxypentanoic acid (Compound No. 75),

10 (2S,3R)-2-[2-(5-fluoro-4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)ethyl]-3-hydroxy-5-(4'-methoxybiphenyl-4-yl)pentanoic acid (Compound No. 76),

(2S,3R)-5-(4'-fluorobiphenyl-4-yl)-2-[2-(5-fluoro-4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)ethyl]-3-hydroxypentanoic acid (Compound No. 77),

(2S,3R)-3-hydroxy-2-[2-(6-methyl-4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)ethyl]-5-[4-(1-methyl-1*H*-pyrazol-4-yl)phenyl]pentanoic acid (Compound No. 78),

15 (2S,3R)-2-[2-(5-chloro-1,3-dioxo-1,3-dihydro-2*H*-isoindol-2-yl)ethyl]-3-hydroxy-5-[4'-(trifluoromethyl)biphenyl-4-yl]pentanoic acid (Compound No. 79),

(2S,3R)-2-[2-(4-fluoro-1,3-dioxo-1,3-dihydro-2*H*-isoindol-2-yl)ethyl]-3-hydroxy-5-[4'-(trifluoromethyl)biphenyl-4-yl]pentanoic acid (Compound No. 80),

20 (2S,3R)-5-(3',4'-difluorobiphenyl-4-yl)-2-[2-(5-fluoro-4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)ethyl]-3-hydroxypentanoic acid (Compound No. 81),

(2S,3R)-3-hydroxyl-5-[4-(6-methoxypyridin-3-yl)phenyl]-2-[2-(4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)ethyl]pentanoic acid (Compound No. 86),

(2S,3R)-5-(4'-chlorobiphenyl-4-yl)-2-[2-(1,3-dioxo-1,3-dihydro-2*H*-isoindol-2-yl)ethyl]-3-hydroxypentanoic acid (Compound No. 87).

25 (2S,3R)-2-[2-(1,3-dioxo-1,3-dihydro-2*H*-isoindol-2-yl)ethyl]-3-hydroxyl-5-[4-(6-methoxypyridin-3-yl)phenyl]pentanoic acid (Compound No. 88),

(2S,3R)-3-hydroxy-2-[2-(7-methyl-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-5-[4-(1-methyl-1H-pyrazol-4-yl)phenyl]pentanoic acid (Compound No. 90),

(2S,3R)-3-hydroxy-2-[2-(8-methoxy-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-5-[4-(1-methyl-1H-pyrazol-4-yl)phenyl]pentanoic acid (Compound No. 91),

5 (2S,3R)-2-[2-(7-chloro-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-5-(3',4'-difluorobiphenyl-4-yl)-3-hydroxypentanoic acid (Compound No. 92),

(2S,3R)-2-[2-(7-chloro-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-3-hydroxy-5-(4'-methoxybiphenyl-4-yl)pentanoic acid (Compound No. 93),

10 (2S,3R)-2-[2-(7-chloro-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-5-(4'-fluorobiphenyl-4-yl)-3-hydroxypentanoic acid (Compound No. 94),

(2S,3R)-2-[2-(7-chloro-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-3-hydroxy-5-[4-(6-methylpyridin-3-yl)phenyl]pentanoic acid (Compound No. 95),

(2S,3R)-5-(2',4'-difluorobiphenyl-4-yl)-2-[2-(1,3-dioxo-1,3-dihydro-2H-isoindol-2-yl)ethyl]-3-hydroxypentanoic acid (Compound No. 97),

15 (2S,3R)-2-[2-(6-chloro-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-3-hydroxy-5-[4-(1-methyl-1H-pyrazol-4-yl)phenyl]pentanoic acid (Compound No. 98),

(2S,3R)-2-[2-(5-chloro-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-3-hydroxy-5-[4-(1-methyl-1H-pyrazol-4-yl)phenyl]pentanoic acid (Compound No. 99),

20 (2S,3R)-2-[2-(1,3-dioxo-1,3-dihydro-2H-isoindol-2-yl)ethyl]-3-hydroxy-5-(4'-isopropylbiphenyl-4-yl)pentanoic acid (Compound No. 100),

(2S,3R)-5-(3'-chloro-4'-fluorobiphenyl-4-yl)-2-[2-(1,3-dioxo-1,3-dihydro-2H-isoindol-2-yl)ethyl]-3-hydroxypentanoic acid (Compound No. 101),

(2S,3R)-5-(4'-butylbiphenyl-4-yl)-3-hydroxy-2-[2-(4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 102),

25 (2S,3R)-5-(2'-fluorobiphenyl-4-yl)-3-hydroxy-2-[2-(4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 103),

(2S,3R)-2-[2-(5-fluoro-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-3-hydroxy-5-[4-(1-methyl-1H-pyrazol-4-yl)phenyl]pentanoic acid (Compound No. 104),

(2S,3R)-3-hydroxy-2-[2-(8-methyl-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-5-[4-(1-methyl-1H-pyrazol-4-yl)phenyl]pentanoic acid (Compound No. 105),

5 (2S,3R)-2-[2-(1,3-dioxo-1,3-dihydro-2H-isoindol-2-yl)ethyl]-3-hydroxy-5-{6-[4-(trifluoromethyl)phenyl]pyridin-3-yl}pentanoic acid (Compound No. 106),

(2S,3R)-5-[6-(3,4-difluorophenyl)pyridin-3-yl]-2-[2-(1,3-dioxo-1,3-dihydro-2H-isoindol-2-yl)ethyl]-3-hydroxypentanoic acid (Compound No. 107),

10 (2S,3R)-5-[6-(4-chloro-3-fluorophenyl)pyridin-3-yl]-2-[2-(1,3-dioxo-1,3-dihydro-2H-isoindol-2-yl)ethyl]-3-hydroxypentanoic acid (Compound No. 108),

(2S,3R)-2-[2-(1,3-dioxo-1,3-dihydro-2H-isoindol-2-yl)ethyl]-5-[6-(4-fluorophenyl)pyridin-3-yl]-3-hydroxypentanoic acid (Compound No. 109),

(2S,3R)-2-[2-(1,3-dioxo-1,3-dihydro-2H-isoindol-2-yl)ethyl]-5-[6-(3-fluoro-4-methylphenyl)pyridin-3-yl]-3-hydroxypentanoic acid (Compound No. 110),

15 (2S,3R)-2-[2-(7-chloro-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-3-hydroxy-5-[4-(1-methyl-1H-pyrazol-4-yl)phenyl]pentanoic acid (Compound No. 111),

(2S,3R)-2-[2-(8-chloro-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-3-hydroxy-5-[4-(1-methyl-1H-pyrazol-4-yl)phenyl]pentanoic acid (Compound No. 112),

20 (2S,3R)-2-[2-(6,7-difluoro-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-3-hydroxy-5-[4-(1-methyl-1H-pyrazol-4-yl)phenyl]pentanoic acid (Compound No. 113),

(2S,3R)-3-hydroxy-2-[2-(7-methyl-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-5-[4'-(trifluoromethoxy)biphenyl-4-yl]pentanoic acid (Compound No. 114),

(2S,3R)-3-hydroxy-5-(4'-methylbiphenyl-4-yl)-2-[2-(7-methyl-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 115),

25 (2S,3R)-5-(4'-fluorobiphenyl-4-yl)-3-hydroxy-2-[2-(7-methyl-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 116),

(2S,3R)-5-(3',4'-difluorobiphenyl-4-yl)-3-hydroxy-2-[2-(7-methyl-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 117),

(2S,3R)-3-hydroxy-2-[2-(7-methyl-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-5-[4-(6-methylpyridin-3-yl)phenyl]pentanoic acid (Compound No. 118),

5 (2S,3R)-5-[4-(2,3-dihydro-1,4-benzodioxin-6-yl)phenyl]-3-hydroxy-2-[2-(7-methyl-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 119),

(2S,3R)-2-[2-(5-chloro-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-3-hydroxy-5-(6'-methoxy-2,3'-bipyridin-5-yl)pentanoic acid (Compound No. 121),

10 (2S,3R)-2-[2-(7-chloro-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-3-hydroxy-5-(6'-methoxy-2,3'-bipyridin-5-yl)pentanoic acid (Compound No. 122),

(2S,3R)-2-[2-(6,7-difluoro-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-3-hydroxy-5-(6'-methoxy-2,3'-bipyridin-5-yl)pentanoic acid (Compound No. 123),

(2S,3R)-5-(3'-fluoro-4'-methylbiphenyl-4-yl)-3-hydroxy-2-[2-(7-methyl-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 124),

15 (2S,3R)-3-hydroxy-5-[4-(1-isobutyl-1H-pyrazol-4-yl)phenyl]-2-[2-(7-methyl-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 125),

(2S,3R)-2-[2-(5-fluoro-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-3-hydroxy-5-(6'-methoxy-2,3'-bipyridin-5-yl)pentanoic acid (Compound No. 126),

20 (2S,3R)-5-(4'-fluorobiphenyl-4-yl)-3-hydroxy-2-[2-(8-methoxy-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 127),

(2S,3R)-5-(3',4'-difluorobiphenyl-4-yl)-3-hydroxy-2-[2-(8-methoxy-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 128),

(2S,3R)-3-hydroxy-5-(4'-methoxybiphenyl-4-yl)-2-[2-(8-methoxy-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 129),

25 (2S,3R)-2-[2-(7-chloro-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-3-hydroxy-5-{6-[4-(trifluoromethyl)phenyl]pyridin-3-yl}pentanoic acid (Compound No. 130),

(2S,3R)-2-[2-(6,7-difluoro-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-3-hydroxy-5-{6-[4-(trifluoromethyl)phenyl]pyridin-3-yl}pentanoic acid (Compound No. 131),

(2S,3R)-2-[2-(5-chloro-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-3-hydroxy-5-{6-[4-(trifluoromethyl)phenyl]pyridin-3-yl}pentanoic acid (Compound No. 132),

5 (2S,3R)-3-hydroxy-2-[2-(7-methyl-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-5-{6-[4-(trifluoromethyl)phenyl]pyridin-3-yl}pentanoic acid (Compound No. 133),

(2S,3R)-2-[2-(5-fluoro-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-3-hydroxy-5-{6-[4-(trifluoromethyl)phenyl]pyridin-3-yl}pentanoic acid (Compound No. 134),

10 (2S,3R)-3-hydroxy-2-[2-(8-methoxy-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-5-(4'-methylbiphenyl-4-yl)pentanoic acid (Compound No. 135),

(2S,3R)-5-(3'-fluoro-4'-methylbiphenyl-4-yl)-2-[2-(5-fluoro-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-3-hydroxypentanoic acid (Compound No. 136),

(2S,3R)-2-[2-(6,7-difluoro-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-3-hydroxy-5-(4'-methylbiphenyl-4-yl)pentanoic acid (Compound No. 137),

15 (2S,3R)-2-[2-(5-fluoro-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-3-hydroxy-5-(4'-methylbiphenyl-4-yl)pentanoic acid (Compound No. 138),

(2S,3R)-5-(4'-chlorobiphenyl-4-yl)-2-[2-(5-fluoro-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-3-hydroxypentanoic acid (Compound No. 139),

20 (2S,3R)-3-hydroxy-5-(6'-methoxy-2,3'-bipyridin-5-yl)-2-[2-(4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 140),

(2S,3R)-3-hydroxy-5-(6'-methoxy-2,3'-bipyridin-5-yl)-2-[2-(6-methyl-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 141),

(2S,3R)-3-hydroxy-5-(6'-methoxy-2,3'-bipyridin-5-yl)-2-[2-(7-methyl-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 142),

25 (2S,3R)-2-[2-(6,7-difluoro-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-5-(3'-fluoro-4'-methylbiphenyl-4-yl)-3-hydroxypentanoic acid (Compound No. 144),

(2S,3R)-5-(4'-chlorobiphenyl-4-yl)-2-[2-(6,7-difluoro-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-3-hydroxypentanoic acid (Compound No. 145),

(2S,3R)-5-(3'-fluoro-4'-methylbiphenyl-4-yl)-3-hydroxy-2-[2-(8-methoxy-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 146),

5 (2S,3R)-5-[4-(2,3-dihydro-1,4-benzodioxin-6-yl)phenyl]-3-hydroxy-2-[2-(8-methoxy-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 147),

(2S,3R)-5-[4-(6-chloropyridin-3-yl)phenyl]-3-hydroxy-2-[2-(4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 148),

10 (2S,3R)-3-hydroxy-2-[2-(6-methoxy-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-5-[4'-(trifluoromethyl)biphenyl-4-yl]pentanoic acid (Compound No. 149),

(2S,3R)-3-hydroxy-2-[2-(7-methoxy-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-5-[4'-(trifluoromethyl)biphenyl-4-yl]pentanoic acid (Compound No. 150),

(2S,3R)-3-hydroxy-2-[2-(6-methoxy-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-5-(4'-methylbiphenyl-4-yl)pentanoic acid (Compound No. 151),

15 (2S,3R)-3-hydroxy-5-(4'-methoxybiphenyl-4-yl)-2-[2-(6-methoxy-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 152),

(2S,3R)-5-[2-fluoro-4-(6-methoxypyridin-3-yl)phenyl]-3-hydroxy-2-[2-(4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 153),

20 (2S,3R)-5-[2-fluoro-4-(6-methoxypyridin-3-yl)phenyl]-3-hydroxy-2-[2-(7-methoxy-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 154),

(2S,3R)-5-[2-fluoro-4-(6-methoxypyridin-3-yl)phenyl]-2-[2-(5-fluoro-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-3-hydroxypentanoic acid (Compound No. 155),

(2S,3R)-5-[2-fluoro-4-(6-methoxypyridin-3-yl)phenyl]-3-hydroxy-2-[2-(7-methyl-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 156),

25 (2S,3R)-5-[2-fluoro-4-(6-methoxypyridin-3-yl)phenyl]-3-hydroxy-2-[2-(6-methyl-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 157),

(2S,3R)-3-hydroxy-5-(4'-methoxybiphenyl-4-yl)-2-[2-(7-methoxy-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 158),

(2S,3R)-5-(4'-chlorobiphenyl-4-yl)-3-hydroxy-2-[2-(6-methoxy-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 159),

5 (2S,3R)-3-hydroxy-2-[2-(7-methoxy-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-5-[4-(6-methoxypyridin-3-yl)phenyl]pentanoic acid (Compound No. 160),

(2S,3R)-5-(3',4'-difluorobiphenyl-4-yl)-3-hydroxy-2-[2-(7-methoxy-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 161),

10 (2S,3R)-3-hydroxy-2-[2-(6-methoxy-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-5-[4-(6-methoxypyridin-3-yl)phenyl]pentanoic acid (Compound No. 162),

(2S,3R)-2-[2-(5-chloro-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-5-[2-fluoro-4-(6-methoxypyridin-3-yl)phenyl]-3-hydroxypentanoic acid (Compound No. 163),

(2S,3R)-2-[2-(6,7-difluoro-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-5-[2-fluoro-4-(6-methoxypyridin-3-yl)phenyl]-3-hydroxypentanoic acid (Compound No. 164),

15 (2S,3R)-5-[2-fluoro-4-(6-methoxypyridin-3-yl)phenyl]-3-hydroxy-2-[2-(5-methyl-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 165),

(2S,3R)-5-(4'-fluorobiphenyl-4-yl)-3-hydroxy-2-[2-(5-methyl-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 166),

20 (2S,3R)-3-hydroxy-5-[4-(6-methoxypyridin-3-yl)phenyl]-2-[2-(5-methyl-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 167),

(2S,3R)-3-hydroxy-5-(4'-methoxybiphenyl-4-yl)-2-[2-(5-methyl-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 168),

(2S,3R)-3-hydroxy-5-(4'-methylbiphenyl-4-yl)-2-[2-(5-methyl-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 169),

25 (2S,3R)-5-(4'-chlorobiphenyl-4-yl)-3-hydroxy-2-[2-(5-methyl-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 170),

(2S,3R)-5-(4'-fluorobiphenyl-4-yl)-3-hydroxy-2-[2-(6-methoxy-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 171),

(2S,3R)-5-(3',4'-difluorobiphenyl-4-yl)-3-hydroxy-2-[2-(6-methoxy-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 172),

5 (2S,3R)-3-hydroxy-2-[2-(6-methoxy-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-5-[4'-(trifluoromethoxy)biphenyl-4-yl]pentanoic acid (Compound No. 173),

(2S,3R)-5-(3'-fluoro-4'-methylbiphenyl-4-yl)-3-hydroxy-2-[2-(6-methoxy-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 174),

10 (2S,3R)-2-[2-(7-chloro-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-5-[2-fluoro-4-(6-methoxypyridin-3-yl)phenyl]-3-hydroxypentanoic acid (Compound No. 175),

(2S,3R)-3-hydroxy-5-[4-(1-methyl-1H-pyrazol-4-yl)-2-thienyl]-2-[2-(4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 176),

(2S,3R)-5-[2-fluoro-4-(6-methoxypyridin-3-yl)phenyl]-3-hydroxy-2-[2-(6-methoxy-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 177),

15 (2S,3R)-3-hydroxy-5-(6'-methoxy-2,3'-bipyridin-5-yl)-2-[2-(6-methoxy-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 178),

(2S,3R)-3-hydroxy-2-[2-(6-methoxy-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-5-[4-(6-methylpyridin-3-yl)phenyl]pentanoic acid (Compound No. 179),

20 (2S,3R)-5-[4-(2-chloropyridin-3-yl)phenyl]-3-hydroxy-2-[2-(6-methoxy-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 180),

(2S,3R)-2-[2-(6-fluoro-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-3-hydroxy-5-[4-(6-methoxypyridin-3-yl)phenyl]pentanoic acid (Compound No. 181),

(2S,3R)-5-(4'-chlorobiphenyl-4-yl)-2-[2-(6-fluoro-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-3-hydroxypentanoic acid (Compound No. 182),

25 (2S,3R)-5-(3',4'-difluorobiphenyl-4-yl)-2-[2-(6-fluoro-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-3-hydroxypentanoic acid (Compound No. 183),

(2S,3R)-2-[2-(6-fluoro-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-3-hydroxy-5-(4'-methoxybiphenyl-4-yl)pentanoic acid (Compound No. 184),

(2S,3R)-3-hydroxy-5-[4-(2-methoxypyrimidin-5-yl)phenyl]-2-[2-(4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 185),

5 (2S,3R)-5-[2-fluoro-4-(1-methyl-1H-pyrazol-4-yl)phenyl]-3-hydroxy-2-[2-(6-methyl-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 186),

(2S,3R)-5-[2-fluoro-4-(1-methyl-1H-pyrazol-4-yl)phenyl]-3-hydroxy-2-[2-(7-methyl-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 187),

10 (2S,3R)-2-[2-(5-chloro-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-5-[2-fluoro-4-(1-methyl-1H-pyrazol-4-yl)phenyl]-3-hydroxypentanoic acid (Compound No. 188),

(2S,3R)-2-[2-(7-chloro-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-5-[2-fluoro-4-(1-methyl-1H-pyrazol-4-yl)phenyl]-3-hydroxypentanoic acid (Compound No. 189),

(2S,3R)-5-[2-fluoro-4-(1-methyl-1H-pyrazol-4-yl)phenyl]-2-[2-(5-fluoro-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-3-hydroxypentanoic acid (Compound No. 190),

15 (2S,3R)-2-[2-(6,7-difluoro-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-5-[2-fluoro-4-(1-methyl-1H-pyrazol-4-yl)phenyl]-3-hydroxypentanoic acid (Compound No. 191),

(2S,3R)-5-[2-fluoro-4-(1-methyl-1H-pyrazol-4-yl)phenyl]-2-[2-(6-fluoro-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-3-hydroxypentanoic acid (Compound No. 192),

20 (2S,3R)-5-[2-fluoro-4-(1-methyl-1H-pyrazol-4-yl)phenyl]-3-hydroxy-2-[2-(8-methyl-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 193),

(2S,3R)-3-hydroxy-2-[2-(6-methoxy-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-5-[4-(2-methoxypyrimidin-5-yl)phenyl]pentanoic acid (Compound No. 194),

(2S,3R)-2-[2-(1,3-dioxo-1,3-dihydro-2H-isoindol-2-yl)ethyl]-3-hydroxy-5-[4-(2-methoxypyrimidin-5-yl)phenyl]pentanoic acid (Compound No. 195),

25 (2S,3R)-5-[2-fluoro-4-(1-methyl-1H-pyrazol-4-yl)phenyl]-3-hydroxy-2-[2-(1-oxophthalazin-2(1H)-yl)ethyl]pentanoic acid (Compound No. 196),

(2*S*,3*R*)-5-[2-fluoro-4-(1-methyl-1*H*-pyrazol-4-yl)phenyl]-3-hydroxy-2-[2-(3-methyl-2,6-dioxo-3,6-dihydropyrimidin-1(2*H*)-yl)ethyl]pentanoic acid (Compound No. 197),

5 (2*S*,3*R*)-3-hydroxy-5-[4-(6-methoxypyridin-3-yl)phenyl]-2-{2-[4-oxo-7-(trifluoromethyl)-1,2,3-benzotriazin-3(4*H*)-yl]ethyl} pentanoic acid (Compound No. 198),

(2*S*,3*R*)-3-hydroxy-5-[4-(6-methoxypyridin-3-yl)phenyl]-2-[2-(1-oxophthalazin-2(1*H*)-yl)ethyl]pentanoic acid (Compound No. 199),

(2*S*,3*R*)-3-hydroxy-5-[4-(6-methoxypyridin-3-yl)phenyl]-2-[2-(3-methyl-2,6-dioxo-3,6-dihydropyrimidin-1(2*H*)-yl)ethyl]pentanoic acid (Compound No. 200),

10 (2*S*,3*R*)-2-[2-(7,9-dioxo-8-azaspiro[4.5]dec-8-yl)ethyl]-3-hydroxy-5-[4-(6-methoxypyridin-3-yl)phenyl]pentanoic acid (Compound No. 201),

(2*S*,3*R*)-2-[2-(2,4-dioxo-2*H*-1,3-benzoxazin-3(4*H*)-yl)ethyl]-3-hydroxy-5-[4-(6-methoxypyridin-3-yl)phenyl]pentanoic acid (Compound No. 202),

15 (2*S*,3*R*)-3-hydroxy-5-[4-(6-methoxypyridin-3-yl)phenyl]-2-[2-(3,4,4-trimethyl-2,5-dioxoimidazolidin-1-yl)ethyl]pentanoic acid (Compound No. 203),

(2*S*,3*R*)-2-[2-(5-fluoro-4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)ethyl]-3-hydroxy-5-[4'-(trifluoromethyl)biphenyl-4-yl]pentanoic acid (Compound No. 211),

(2*S*,3*R*)-5-(4'-chlorobiphenyl-4-yl)-3-hydroxy-2-[2-(7-methyl-4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)ethyl]pentanoic acid (Compound No. 212),

20 (2*S*,3*R*)-3-hydroxy-5-(6'-methoxy-2,3'-bipyridin-5-yl)-2-[2-(8-methyl-4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)ethyl]pentanoic acid (Compound No. 213),

(2*S*,3*R*)-3-hydroxy-5-[4-(6-hydroxypyridin-3-yl)phenyl]-2-[2-(4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)ethyl]pentanoic acid (Compound No. 226),

25 (2*S*,3*R*)-3-hydroxy-5-(4'-methylbiphenyl-4-yl)-2-{2-[4-oxo-7-(trifluoromethyl)-1,2,3-benzotriazin-3(4*H*)-yl]ethyl} pentanoic acid (Compound No. 227),

(2*S*,3*R*)-2-[2-(2,4-dioxo-1,4-dihydroquinazolin-3(2*H*)-yl)ethyl]-3-hydroxy-5-[4-(6-methoxypyridin-3-yl)phenyl]pentanoic acid (Compound No. 228),

(2*S*,3*R*)-3-(acetyloxy)-5-[4-(6-methoxypyridin-3-yl)phenyl]-2-[2-(4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)ethyl]pentanoic acid (Compound No. 229),

(2*S*,3*R*)-2-[2-(8-chloro-4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)ethyl]-3-hydroxy-5-[4-(6-methoxypyridin-3-yl)phenyl]pentanoic acid (Compound No. 230),

5 (2*S*,3*R*)-5-(4'-chlorobiphenyl-4-yl)-2-[2-(2,4-dioxo-2*H*-1,3-benzoxazin-3(4*H*)-yl)ethyl]-3-hydroxypentanoic acid (Compound No. 231), and

(2*S*,3*R*)-5-(4'-chlorobiphenyl-4-yl)-3-hydroxy-2-(2-[(2-hydroxyphenyl)carbonyl]amino)ethyl) pentanoic acid (Compound No. 232) .

Compound Nos. 82 to 85, and 204 were prepared following Schemes I, II and III.

10 (2*S*,3*R*)-3-hydroxy-2-[2-(4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)ethyl]-5-phenylpentanoic acid (Compound No. 82),

(2*S*,3*R*)-2-[2-(1,3-dioxo-1,3-dihydro-2*H*-isoindol-2-yl)ethyl]-3-hydroxy-5-phenylpentanoic acid (Compound No. 83),

15 (2*S*,3*R*)-2-[2-(1,3-dioxo-1,3-dihydro-2*H*-isoindol-2-yl)ethyl]-3-hydroxy-5-[4(trifluoromethyl)phenyl]pentanoic acid (Compound No. 84),

(2*S*,3*R*)-5-(4-*tert*-butylphenyl)-2-[2-(1,3-dioxo-1,3-dihydro-2*H*-isoindol-2-yl)ethyl]-3-hydroxypentanoic acid (Compound No.85), and

(2*S*,3*R*)-5-(4-chloro-3-fluorophenyl)-2-[2-(1,3-dioxo-1,3-dihydro-2*H*-isoindol-2-yl)ethyl]-3-hydroxypentanoic acid (Compound No. 204),

20 Compound Nos. 143, and 208 were prepared following Scheme IV.

(2*S*,3*S*)-3-hydroxy-5-[4-(6-methoxypyridin-3-yl)phenyl]-2-[2-(4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)ethyl]pentanoic acid (Compound No. 143), and

(2*S*,3*S*)-3-hydroxy-5-(4'-methoxybiphenyl-4-yl)-2-[2-(4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)ethyl]pentanoic acid (Compound No. 208),

25 Compound Nos. 120, and 205-207 were prepared following Scheme V.

(2*R*,3*S*)-3-hydroxy-5-[4-(6-methoxypyridin-3-yl)phenyl]-2-[2-(4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)ethyl]pentanoic acid (Compound No. 120)

(2*R*,3*S*)-2-[2-(1,3-dioxo-1,3-dihydro-2*H*-isoindol-2-yl)ethyl]-5-(3'-fluoro-4'-methylbiphenyl-4-yl)-3-hydroxypentanoic acid (Compound No. 205),

- 5 (2*R*,3*S*)-2-[2-(1,3-dioxo-1,3-dihydro-2*H*-isoindol-2-yl)ethyl]-5-(4'-fluorobiphenyl-4-yl)-3-hydroxypentanoic acid (Compound No. 206), and

(2*R*,3*S*)-5-(3',4'-difluorobiphenyl-4-yl)-2-[2-(1,3-dioxo-1,3-dihydro-2*H*-isoindol-2-yl)ethyl]-3-hydroxypentanoic acid (Compound No. 207),

Compound Nos. 89, and 96 were prepared following Scheme VI.

- 10 (2*R*,3*R*)-5-(4'-chlorobiphenyl-4-yl)-2-[2-(1,3-dioxo-1,3-dihydro-2*H*-isoindol-2-yl)ethyl]-3-hydroxypentanoic acid (Compound No. 89),

(2*R*,3*R*)-3-hydroxyl-5-[4-(6-methoxypyridin-3-yl)phenyl]-2-[2-(4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)ethyl]pentanoic acid (Compound No. 96),

Compound Nos. 86, 209-210 and 214-225 were prepared following Scheme VII.

- 15 (2*S*,3*R*)-3-hydroxy-5-[4-(6-methoxypyridin-3-yl)phenyl]-2-[2-(4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)ethyl]pentanoic acid (Compound No. 86),

(2*S*,3*R*)-3-hydroxy-5-[4-(5-methylpyridin-2-yl)phenyl]-2-[2-(4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)ethyl]pentanoic acid (Compound No. 209),

- 20 (2*S*,3*R*)-5-[4-(6-fluoropyridin-3-yl)phenyl]-3-hydroxy-2-[2-(4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)ethyl]pentanoic acid (Compound No. 210),

(2*S*,3*R*)-5-(3'-fluoro-4'-methylbiphenyl-4-yl)-3-hydroxy-2-[(4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)methyl]pentanoic acid (Compound No. 214),

(2*S*,3*R*)-3-hydroxy-2-[(4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)methyl]-5-[4'-(trifluoromethyl)biphenyl-4-yl]pentanoic acid (Compound No. 215),

- 25 (2*S*,3*R*)-5-(2',4'-difluorobiphenyl-4-yl)-3-hydroxy-2-[(4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)methyl]pentanoic acid (Compound No. 216),

((2*S*,3*R*)-5-(4'-fluorobiphenyl-4-yl)-3-hydroxy-2-[(4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)methyl]pentanoic acid (Compound No. 217),

(2*S*,3*R*)-5-(3'-fluoro-4'-methoxybiphenyl-4-yl)-3-hydroxy-2-[(4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)methyl]pentanoic acid (Compound No. 218),

5 (2*S*,3*R*)-3-hydroxy-5-[4-(2-methoxypyrimidin-5-yl)phenyl]-2-[(4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)methyl]pentanoic acid(Compound No. 219),

(2*S*,3*R*)-3-hydroxy-5-[4-(6-methoxypyridin-3-yl)phenyl]-2-[(4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)methyl]pentanoic acid(Compound No. 220),

10 (2*S*,3*R*)-3-hydroxy-5-(4'-methylbiphenyl-4-yl)-2-[(4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)methyl]pentanoic acid(Compound No. 221),

(2*S*,3*R*)-3-hydroxy-2-[(4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)methyl]-5-[4'-(trifluoromethoxy)biphenyl-4-yl]pentanoic acid (Compound No. 222),

(2*S*,3*R*)-3-hydroxy-5-[4-(6-methylpyridin-3-yl)phenyl]-2-[(4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)methyl]pentanoic acid(Compound No. 223),

15 (2*S*,3*R*)-5-(4'-chlorobiphenyl-4-yl)-3-hydroxy-2-[(4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)methyl]pentanoic acid (Compound No. 224),

(2*S*,3*R*)-5-(3',4'-difluorobiphenyl-4-yl)-3-hydroxy-2-[(4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)methyl]pentanoic acid (Compound No. 225),

In the above schemes, where specific bases, acids, solvents, condensing agents, reducing agent, deprotecting agent, hydrolyzing agents, metal catalysts etc., are mentioned, it is to be understood that other acids, bases, solvents, condensing agents, reducing agent, deprotecting agent, hydrolyzing agents, metal catalysts etc., known to those skilled in the art may also be used. Similarly, the reaction temperature and duration of the reactions may be adjusted according to the requirements that arise during the process.

The following examples are set forth to demonstrate general synthetic procedures for the preparation of representative compounds of the present invention. The examples are provided to illustrate particular aspect of the disclosure and do not limit the scope of the present invention.

EXAMPLES

Synthesis of starting materials:

15 Synthesis of 6-methyl-1,2,3-benzotriazin-4(3H)-one

The title compound was prepared following the procedure outlined in *J. Med. Chem.*, (1992), 35(14), 2626-2630.

The following analogues of 6-methyl-1,2,3-benzotriazin-4(3H)-one were prepared analogously,

- 8-methyl-1,2,3-benzotriazin-4(3H)-one;
- 7-methyl-1,2,3-benzotriazin-4(3H)-one;
- 5 6-methyl-1,2,3-benzotriazin-4(3H)-one;
- 8-methoxy-1,2,3-benzotriazin-4(3H)-one;
- 6-methoxy-1,2,3-benzotriazin-4(3H)-one;
- 8-chloro-1,2,3-benzotriazin-4(3H)-one;
- 7-chloro-1,2,3-benzotriazin-4(3H)-one;
- 10 6-chloro-1,2,3-benzotriazin-4(3H)-one;
- 5-chloro-1,2,3-benzotriazin-4(3H)-one;
- 6,7-difluoro-1,2,3-benzotriazin-4(3H)-one;
- 8-fluoro-1,2,3-benzotriazin-4(3H)-one;
- 5-fluoro-1,2,3-benzotriazin-4(3H)-one;
- 15 6-fluoro-1,2,3-benzotriazin-4(3H)-one;
- 5-(6-methoxypyridin-3-yl)-1,2,3-benzotriazin-4(3H)-one;
- 7-(6-methoxypyridin-3-yl)-1,2,3-benzotriazin-4(3H)-one; and
- 7-(trifluoromethyl)-1,2,3-benzotriazin-4(3H)-one.

Synthesis of 5-tert-butyl-1H-isoindole-1,3(2H)-dione

- 20 The title compound was prepared following the procedure outlined in *Chem. Ber.*, (1877), 10, 1163.

Mass (m/z): 204.12(M⁺ + 1)

The following analogues of 5-tert-butyl-1H-isoindole-1,3(2H)-dione were prepared analogously:

- 25 4-fluoro-1H-isoindole-1,3(2H)-dione; and
- 5-chloro-1H-isoindole-1,3(2H)-dione.

Synthesis of (4-bromobenzyl)(triphenyl)phosphonium bromide

A mixture of 1-bromo-4-(bromomethyl)benzene (5g) and triphenylphosphine(5.24 g) in xylene (20 mL) were heated to reflux for 18 hours. The reaction mixture was cooled, filtered, washed with hexane and dried under vacuum to afford the title compound (8 g).

5 Mass (m/z):433.9 ($M^+ + 1$)

The following Wittig salts were prepared analogously:

[(4-bromo-2-thienyl)methyl]triphenyl phosphonium bromide;

[(6-bromopyridin-3-yl)methyl](triphenyl)phosphonium bromide;

(4-bromo-2-fluorobenzyl)(triphenyl)phosphonium bromide;

10 4-tert-butylbenzyl)(triphenyl)phosphonium bromide;

triphenyl[4-(trifluoromethyl)benzyl]phosphonium bromide;

benzyl(triphenyl)phosphonium bromide;

(4-bromo-3-fluorobenzyl)(triphenyl)phosphonium bromide;

(4-bromo-2-fluorobenzyl)(triphenyl)phosphonium bromide; and

15 (4-chloro-3-fluorobenzyl)(triphenyl)phosphonium bromide.

Synthesis of 4-(4-oxo-1,2,3-benzotriazin-3(4H)-yl)butanoic acid**Step a: Synthesis of ethyl 4-(4-oxo-1,2,3-benzotriazin-3(4H)-yl)butanoate**

In a dry round bottom flask, 1,2,3-benzotriazin-4(3H)-one (50 g), ethyl 4-bromobutanoate

20 (86.2 g) and potassium carbonate (141 g) were taken and dissolved in dimethylformamide (350 mL) and heated to 60-70 °C for 3-4 hours. After cooling to room temperature, water was added to the reaction mixture and extracted with ethyl acetate. The combined organic layer was washed with water, brine and dried over anhydrous sodium sulfate. Solvents were evaporated under reduced pressure and the crude product
25 was purified by silica gel flash column chromatography over silica gel using 20% ethylacetate in hexane as eluant to afford the title compound (68 g).

Step b: Synthesis of 4-(4-oxo-1,2,3-benzotriazin-3(4H)-yl)butanoic acid

To a stirred solution compound (40 g), obtained from step a above, in a solvent mixture of tetrahydrofuran/methanol:water (3:1:1, 400 mL) was added lithium hydroxide monohydrate (6.43 g) at 0 °C under a nitrogen atmosphere. The reaction mixture was stirred at 0 °C until completion of hydrolysis. The solvents were evaporated, diluted with water and extracted with ethyl acetate. The aqueous layers were acidified with saturated aqueous sodium hydrogen sulfate solution and extracted with ethyl acetate. The combined layers were washed with water, brine and dried over anhydrous sodium sulfate. The solvents were evaporated under reduced pressure to afford the title compound (28 g).

The following analogue of 4-(4-oxo-1,2,3-benzotriazin-3(4H)-yl)butanoic acid was prepared analogously:

3-(4-oxo-1,2,3-benzotriazin-3(4H)-yl)propanoic acid

Synthesis of 3-{4-[(4S)-4-benzyl-2-oxo-1,3-thiazolidin-3-yl]-4-oxobutyl}-1,2,3-benzotriazin-4(3H)-one

Step a: Synthesis of (2S)-2-amino-3-phenylpropan-1-ol

(Ref: *J. Org. Chem.* **1993**, *58*, 3568-3571)

To a suspension of sodium borohydride (16.5 g) in dry tetrahydrofuran (600 mL), (L)-phenylalanine (30g) was added at one portion. The flask was cooled to 0 °C under nitrogen atmosphere. Iodine (46.18 g) solution in tetrahydrofuran (150 mL) was added slowly in dropwise manner over 40 minutes, resulting in vigorous evolution of H₂. After complete addition of iodine, the reaction mixture was heated to reflux for 18 hours and cooled to room temperature. The reaction mixture was quenched with methanol until the reaction mixture become clear. Further, the solution was stirred for 30 minutes at room temperature. The solvents were removed by rotary evaporation to obtain a white paste which was dissolved by 20% aqueous potassium hydroxide (450 mL). The solution was stirred for 4 hours and then dichloromethane was added. The organic layers were separated and the aqueous layer was extracted with dichloromethane. The combined organic layers were washed with brine and dried over anhydrous sodium sulfate. The solvents were evaporated under reduced pressure to afford 32.5 g of (L)-phenylalaninol as a viscous liquid. The crude product was used as such for the next step.

Step b: Synthesis of (4S)-4-benzyl-1,3-thiazolidine-2-thione

(Ref: *J. Org. Chem.* **1995**, *60*, 6604-6607)

To a solution of compound (32.5 g) obtained in step a above, in 1N aqueous potassium hydroxide (1 lit), carbon disulphide (68 mL, 5.0 equiv.) was added and the reaction mixture was refluxed for 16 hours. After cooling to room temperature, the aqueous solution was extracted with dichloromethane. The organic layer was washed with brine, dried over anhydrous sodium sulfate and concentrated under reduced pressure. The crude product was purified by silica gel flash column chromatography using 15% ethyl acetate in hexane to get the title compound (22.5 g).

10 Step c: Synthesis of (4S)-4-benzyl-1,3-thiazolidine-2-one

To a solution (4S)-4-benzyl-1,3-thiazolidine-2-thione (38 g) in dichloromethane (350 mL), cooled to 0°C was added propylene oxide (12.7 mL) and trifluoroacetic acid (14 mL). After stirring the reaction mixture for 2 hours, the solvents were evaporated under reduced pressure to obtain a residue which was purified by column chromatography over silica gel using 20% ethylacetate in hexane as eluant to afford the title compound (0.9 g).
Mass (m/z): 194.18

Step d: Synthesis of 3-{4-[(4S)-4-benzyl-2-oxo-1,3-thiazolidin-3-yl]-4-oxobutyl}-1,2,3-benzotriazin-4(3H)-one

To a solution of compound (2.1g), obtained from step c above, in dichloromethane (25 mL) cooled to 0°C, 4-dimethylaminopridine (0.334 g) and triethylamine (5.7 mL) were added. After stirring the reaction mixture for 10 minutes, 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (3.9g) and 4-(4-oxo-1,2,3-benzotriazin-3(4H)-yl)butanoic acid (3 g) were added and the reaction mixture was stirred at room temperature for 14 hours. Dichloromethane and water were added to the reaction mixture. The organic layer was separated, washed with water and brine solution and dried over anhydrous sodium sulphate. Solvent was evaporated under reduced pressure to obtain a residue which was purified by column chromatography over silica gel using 25% ethylacetate in hexane as eluant to afford the title compound (4.4 g). Mass (m/z): 409.16

The following analogue of 3-{4-[(4S)-4-benzyl-2-oxo-1,3-thiazolidin-3-yl]-4-oxobutyl}-1,2,3-benzotriazin-4(3H)-one was prepared analogously:

3-{3-[(4*S*)-4-benzyl-2-oxo-1,3-thiazolidin-3-yl]-3-oxopropyl}-1,2,3-benzotriazin-4(3*H*)-one

Example 1: Synthesis of (2*S*,3*R*)-3-hydroxy-2-[2-(4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)ethyl]-5-(4-pyrimidin-5-ylphenyl)pentanoic acid (Compound No.1)

5 Step a: Synthesis of 5-O-[tert-butyl(diphenyl)silyl]-D-arabinofuranose

D-Arabinose (200 g) and imidazole (199 g) were placed in a three-neck round bottom flask and connected to high vacuum for 30 minutes. The vacuum was released under nitrogen atmosphere and dimethylformamide (1.8 L) was added to the above mixture at room temperature followed by drop wise addition of *t*-butyldiphenylchlorosilane (443 mL) for 10 minutes under a nitrogen atmosphere. The resulting mixture was stirred for 16 hours at the same temperature. Dimethylformamide was evaporated under the reduced pressure. The residue was taken up in ethyl acetate and washed with water. The organic layer was dried over anhydrous sodium sulphate. The solvent was evaporated under the reduced pressure to obtain a crude residue which was purified by column chromatography over silica gel using 80% ethyl acetate in hexane as eluant to afford the title compound (273 g). Mass (m/z): 389.37(M⁺ +1)

Step b: Synthesis of 5-O-[tert-butyl(diphenyl)silyl]-1,2-O-isopropylidene-β-D-arabinofuranose

To the solution of compound (273 g), obtained from step a above, in acetone (2.5 L) was charged DL-camphorsulphonic acid (16 g) and anhydrous copper sulphate (346 g) under a nitrogen atmosphere at room temperature. The reaction mixture was stirred for 16 hours at the same temperature. A saturated solution of sodium bicarbonate (2 L) was added dropwise till a basic pH was attained, and the reaction mixture was further stirred for 2 hours at the same temperature. The resulting mixture was filtered using a Buchner funnel and the residue was washed with acetone. The filtrate was concentrated, dissolved in ethyl acetate and washed with water. The organic layer was dried over anhydrous sodium sulphate, filtered and evaporated under reduced pressure to furnish the title compound (295 g). Mass (m/z): 429.41(M⁺ +1)

Step c: Synthesis of 5-O-[tert-butyl(diphenyl)silyl]-1,2-O-isopropylidene-β-D-threopentofuranos-3-ulose

Oxalyl chloride (145 ml) and dichloromethane (1 L) were taken in a three neck round bottom flask and cooled to -75 °C under a nitrogen atmosphere. Dimethylsulfoxide (212 mL) was added dropwise to the above solution maintaining the reaction temperature at -70 °C. The reaction mixture was stirred for 30 minutes at the same temperature, then a solution of the compound (285 g) obtained from step b above, in dichloromethane (1 L), was added slowly to the above mixture, maintaining the reaction temperature -70 °C. After 20 minutes of the above addition, triethylamine (560 mL) was added drop wise at the same temperature. Saturated solution of ammonium chloride in water (1.5 L) was then added after 30 minutes and the reaction temperature was allowed to rise to room temperature. The reaction mixture was extracted with dichloromethane. Organic extracts were dried over anhydrous sodium sulphate, filtered and evaporated under reduced pressure to afford the title compound (285 g)

Step d: Synthesis of 5-O-[tert-butyl(diphenyl)silyl]-3-deoxy-3-(2-ethoxy-2-oxoethylidene)-1,2-O-isopropylidene-β-D-arabinofuranose

To the suspension of sodium hydride (29.4 g, 60% in oil.) in tetrahydrofuran (1L) at 0 °C was charged triethyl phosphonoacetate (200 mL). After 20 min, a solution of compound (285 g), obtained from step c above, in tetrahydrofuran (2 L) was added dropwise and the reaction mixture was stirred for 1 hour at the same temperature. A saturated solution of ammonium chloride in water (1.5 L) was added to reaction mixture. The resulting mixture was extracted with ethyl acetate. Combined extracts were dried over anhydrous sodium sulphate and evaporated under reduced pressure to yield a residue which was purified by column chromatography over silica gel using 15% ethyl acetate in hexane as eluant to afford the title compound (275 g). Mass (m/z): 497.43 (M⁺ +1)

Step e: Synthesis of 5-O-[tert-butyl(diphenyl)silyl]-3-deoxy-3-(2-ethoxy-2-oxoethyl)-1,2-O-isopropylidene-β-D-lyxofuranose and 5-O-[tert-butyl(diphenyl)silyl]-3-deoxy-3-(2-ethoxy-2-oxoethyl)-1,2-O-isopropylidene-α-L-ribofuranose

10% Palladium on charcoal (100 g) was added to the solution of compound (275 g), obtained from step d above, in tetrahydrofuran (2 L) and methanol (1 L) at room temperature and hydrogen was supplied at 50 psi (Paar apparatus) for 2 hours. The

reaction mixture was filtered through a celite pad and the residue was washed with ethyl acetate. The filtrate was concentrated to afford a mixture of 5-O-[tert-butyl(diphenyl)silyl]-3-deoxy-3-(2-ethoxy-2-oxoethyl)-1,2-O-isopropylidene- β -D-lyxofuranose and 5-O-[tert-butyl(diphenyl)silyl]-3-deoxy-3-(2-ethoxy-2-oxoethyl)-1,2-O-isopropylidene- α -L-ribofuranose (275 g). Mass (m/z): 499.42 ($M^+ + 1$)

Step f: Synthesis of 5-O-[tert-butyl(diphenyl)silyl]-3-deoxy-3-(2-hydroxyethyl)-1,2-O-isopropylidene- β -D-lyxofuranose and 5-O-[tert-butyl(diphenyl)silyl]-3-deoxy-3-(2-hydroxyethyl)-1,2-O-isopropylidene- α -L-ribofuranose

To the suspension of lithium aluminium hydride (48 g) in tetrahydrofuran (1 L) was added a solution of compound (275 g), obtained from step e above, in tetrahydrofuran (1.5 L) at -50°C . The resulting mixture was allowed to attain 0°C . The reaction mixture was stirred for 30 minutes at same temperature and again cooled to -50°C . Ethyl acetate (2 L) was added slowly while maintaining -50°C . An aqueous solution of ammonium chloride (100 g) in water (2.5 L) was added at the same temperature. The reaction mixture was slowly allowed to warm to room temperature and the reaction mixture was stirred for 12 hours at same temperature. The reaction mixture was then filtered through a celite pad and the residue was washed with ethyl acetate. The organic layer was dried over anhydrous sodium sulphate, filtered, concentrated under reduced pressure and the residue thus obtained was purified by column chromatography over silica gel using 50% ethyl acetate in hexane as eluant to afford the 5-O-[tert-butyl(diphenyl)silyl]-3-deoxy-3-(2-hydroxyethyl)-1,2-O-isopropylidene- β -D-lyxofuranose (130 g) and 5-O-[tert-butyl(diphenyl)silyl]-3-deoxy-3-(2-hydroxyethyl)-1,2-O-isopropylidene- α -L-ribofuranose (40.0 g). Mass (m/z): 457.39($M^+ + 1$).

Step g: Synthesis of 5-O-[tert-butyl(diphenyl)silyl]-3-deoxy-1,2-O-isopropylidene-3-[2-(4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]- β -D-lyxofuranose

5-O-[tert-butyl(diphenyl)silyl]-3-deoxy-3-(2-hydroxyethyl)-1,2-O-isopropylidene- β -D-lyxofuranose (35g), triphenylphosphine (30.8g) and 1,2,3-benzotriazin-4(3H)-one (12.6g) were dried in high vacuum in a round bottom flask for 10 minutes. Then the vacuum was released under a nitrogen atmosphere and tetrahydrofuran (70mL) was added to the above reaction mixture. The reaction mixture was cooled to 0°C and diisopropyl azodicarboxylate (17mL) was added slowly. The reaction mixture was stirred for 30

minutes at same temperature, and then quenched with water and extracted with ethyl acetate. The organic layer was washed with water, brine solution, dried over anhydrous sodium sulphate and concentrated to obtain a residue which was purified by column chromatography over silica gel using 30% ethyl acetate in hexane as eluant to afford the
5 title compound (50g). Mass (m/z):586.05 ($M^+ + 1$)

Step h: Synthesis of 3-deoxy-1,2-O-isopropylidene-3-[2-(4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]- β -D-lyxofuranose

To a solution of compound (50g), obtained from step g above, in dry tetrahydrofuran (400mL) at 0°C, was added tetra-butyl ammonium fluoride (170mL). The
10 resulting mixture was initially stirred at 0°C for 1hour, and then at room temperature for 4 hours. The reaction mixture was cooled to 0°C, quenched with saturated ammonium chloride and extracted with ethyl acetate. The organic layer was washed with water and brine, dried over anhydrous sodium sulphate and concentrated. The residue thus obtained was purified by column chromatography over silica gel using 80% ethyl acetate in hexane
15 as eluant to furnish the title compound (22 g).

Mass (m/z):369.98 ($M^+ + 23$)

Step i: Synthesis of (5S)-3-deoxy-4,5-O-isopropylidene-3-[2-(4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-D-arabino-pentodialdo-5,2-furanose

Oxalyl chloride (13.7mL) and dichloromethane (300mL) were taken in a three
20 neck round bottom flask and cooled to -78°C. Dimethylsulfoxide (20.2mL) was added dropwise to the reaction mixture. The reaction mixture was warmed to -35°C for 5 to 10 minutes and again cooled to -78°C. A solution of compound (22 g), obtained from step h above, in dichloromethane (200mL) was added slowly while maintaining the same temperature. The reaction mixture was stirred for 45 minutes till the reaction temperature
25 reached -35°C. The reaction mixture was again cooled to -78°C and triethylamine (53mL) was added. The reaction mixture was stirred for an additional 30 minutes, and the temperature was allowed to reach -35°C. The reaction mixture was quenched with saturated solution of ammonium chloride and extracted with dichloromethane. The combined organic layers were washed with water and brine solution, dried over anhydrous
30 sodium sulphate and concentrated to furnish the title compound (22g).

Step j: Synthesis of 3-(2-((3aS,5R,6S,6aS)-5-[(E)-2-(4-bromophenyl)vinyl]-2,2-dimethyltetrahydrofuro[2,3-d][1,3]dioxol-6-yl)ethyl)-1,2,3-benzotriazin-4(3H)-one

To the suspension of sodium hydride (3.3g, 60% in oil) in tetrahydrofuran (50mL) cooled to 0°C, was added (4-bromobenzyl)triphenylphosphonium bromide (48.8g) in tetrahydrofuran (100mL). A solution of compound (22g), obtained from step i above, in tetrahydrofuran (100mL) was added dropwise after 20 minutes, and reaction mixture was stirred for 1 hour at the same temperature. The reaction mixture was quenched with water and extracted with ethyl acetate. Organic layer was washed with water and brine solution and dried over anhydrous sodium sulphate. The solvent was evaporated under reduced pressure and the residue thus obtained was purified by column chromatography using 30% ethyl acetate in hexane as eluant to furnish the title compound (24.5g). Mass (m/z): 498.84 ($M^+ + 1$)

Step k: Synthesis of 3-(2-((3aS,5R,6S,6aS)-2,2-dimethyl-5-[(E)-2-(4-pyrimidin-5-ylphenyl)vinyl]tetrahydrofuro[2,3-d][1,3]dioxol-6-yl)ethyl)-1,2,3-benzotriazin-4(3H)-one

A mixture of compound (0.35g), obtained from step j above, pyrimidin-5-ylboronic acid (0.174g), tetrakis(triphenylphosphine)palladium (0) (0.081g) and potassium carbonate (0.291g) was dried under high vacuum for 10 minutes and dry dimethylformamide (5mL) was added at room temperature. The reaction mixture was heated at 120°C for 2 hours, and then quenched with water and extracted with ethyl acetate. The organic layer was washed with water and brine solution and dried over anhydrous sodium sulphate. Solvent was evaporated under the reduced pressure to obtain a residue which was purified by column chromatography over silica gel using 40% ethyl acetate in hexane as eluant to afford the title compound (0.3g). Mass (m/z): 498.0 ($M^+ + 1$)

Step l: Synthesis of 3-(2-((3aS,5R,6S,6aS)-2,2-dimethyl-5-[2-(4-pyrimidin-5-ylphenyl)ethyl]tetrahydrofuro[2,3-d][1,3]dioxol-6-yl)ethyl)-1,2,3-benzotriazin-4(3H)-one

To the solution of compound (0.3g), obtained from step k above, in a solvent mixture of tetrahydrofuran: methanol (10mL, 1:1) was added 10% Palladium on charcoal (0.15g) at room temperature and the reaction mixture was hydrogenated with hydrogen at the pressure of 35 psi for 4 hours in a Paar apparatus. The reaction mixture was filtered

through a celite pad and the residue was washed with methanol. The filtrate was concentrated to afford the title compound (0.3g).

Step m: Synthesis of 3-(2-((2R,3R,4S,5R)-4,5-dihydroxy-2-[2-(4-pyrimidin-5-ylphenyl)ethyl]tetrahydrofuran-3-yl)ethyl)-1,2,3-benzotriazin-4(3H)-one

5 Perchloric acid (0.2mL) was added to a solution of compound (0.3g), obtained from step l above, in acetonitrile (4mL) and water (0.2mL) at room temperature. The reaction mixture was heated to 55°C for 30 minutes. The reaction mixture was then quenched using sodium bicarbonate solution. The solvents were evaporated under reduced pressure. The residue thus obtained was taken up in ethyl acetate and water. The
10 organic layer was separated and washed with water and brine solution and dried over anhydrous sodium sulphate. Solvent was concentrated under reduced pressure to yield the title compound (0.25g).

Step n: Synthesis of (2S,3R)-3-(formyloxy)-2-[2-(4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-5-(4-pyrimidin-5-ylphenyl)pentanoic acid

15 To a solution of compound (0.25g), obtained from step m above, in *tert*-butanol: tetrahydrofuran (5mL:5mL) at 0°C was added a solution of sodium metaperiodate (0.465 g in 5 mL of water). The reaction mixture was stirred for 2 hours at the same temperature and potassium permanganate (0.017g) was added at 0°C. After stirring the reaction mixture for an additional 6 hours at room temperature, the reaction mixture was evaporated on
20 rotary evaporator. The residue was taken into water and extracted with ethyl acetate. The organic layer was washed with water and brine solution and dried over anhydrous sodium sulphate. The solvent was evaporated under the reduced pressure, to afford the title compound (0.25g).

Step o: Synthesis of (2S,3R)-3-hydroxy-2-[2-(4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-5-(4-pyrimidin-5-ylphenyl)pentanoic acid

25 Potassium carbonate (0.081g) was added to a solution of compound (0.25g), obtained from step n above, in methanol (5mL) and tetrahydrofuran (5mL) at 0°C. The reaction mixture was stirred at room temperature for 3 hours. Solvents were evaporated and the residue was taken into water and ethyl acetate. The organic layer was washed with
30 water and brine solution and dried over anhydrous sodium sulphate. Solvent was evaporated under reduced pressure to obtain a residue which was purified by column

chromatography over silica gel using 60% ethyl acetate in hexane as eluant to afford the title compound (0.030g). Mass (m/z):446.0 ($M^+ + 1$).

^1H NMR (CD_3OD): δ 9.09 (s, 1H), 9.03 (s, 1H), 8.30 (d, 1H, $J = 8\text{Hz}$), 8.14 (d, 1H, $J = 8\text{Hz}$), 8.05- 8.03 (m, 1H), 7.89-7.87 (m, 1H), 7.61 (d, 2H, $J = 8\text{Hz}$), 7.53- 7.59 (m, 1H),
 5 7.35 (d, 2H, $J = 8\text{Hz}$), 4.57- 4.52 (m, 2H), 3.79-3.77 (m, 1H), 2.89 -2.87 (m, 1H), 2.69 - 2.67 (m, 1H), 2.47- 2.46 (m, 1H), 2.29- 2.25 (m, 2H), 1.82- 1.77 (m, 2H).

Example 1A: Synthesis of (2S,3R)-3-hydroxy-5-[4-(6-methoxypyridin-3-yl)phenyl]-2-[2-(4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 86)

Step a: Synthesis of 3-(2-[(3aS,5R,6S,6aS)-5-[(E)-2-(4-bromophenyl)ethenyl]-2,2-dimethyltetrahydrofuro[2,3-d][1,3]dioxol-6-yl)ethyl)-1,2,3-benzotriazin-4(3H)-one
 10

To the suspension of potassium t-butoxide (3.3g, 60% in oil) in dimethyl sulfoxide (50mL) cooled to 0°C , was added 4-bromobenzyl triphenyl-phosphonium bromide (48.8g) in dimethyl sulfoxide (100mL). After 20 minutes, compound (22g) obtained from step i of example I above, was added in dimethyl sulfoxide (100mL) drop wise and reaction
 15 mixture was stirred for 1 hour at the same temperature. Reaction mixture was quenched with water and extracted with ethyl acetate. Organic layer was washed with water and brine solution and dried over anhydrous sodium sulphate. Solvent was evaporated under the reduced pressure. Purification was performed on silica gel column by using 30% ethyl acetate in hexane as eluent to get the title compound (24.5g).

Step b: Synthesis of 3-{2-[(3aS,5R,6S,6aS)-5-{(E)-2-[4-(6-methoxypyridin-3-yl)phenyl]ethenyl}-2,2-dimethyltetrahydrofuro[2,3-d][1,3]dioxol-6-yl]ethyl}-1,2,3-benzotriazin-4(3H)-one
 20

A mixture of compound (12gms), obtained from step a above, 2-methoxy-5-pyridine boronic acid (7.4gms), tetrakis(triphenylphosphine)palladium (0) (2.8gms) and
 25 potassium carbonate (10gms) was dried under high vacuum for 10 minutes and dry dimethylformamide (60ml) was added at room temperature. The reaction mixture was heated at 120°C for 2 hours, and then quenched with water and extracted with ethyl acetate. The organic layer was washed with water and brine solution and dried over anhydrous sodium sulphate. Solvent was evaporated under the reduced pressure to obtain a
 30 residue which was purified by column chromatography over silica gel using 40% ethyl acetate in hexane as eluant to afford the title compound (8.5gms).

Step c: Synthesis of 3-{2-[(3*aS*,5*R*,6*S*,6*aS*)-5-{2-[4-(6-methoxypyridin-3-yl)phenyl]ethyl}-2,2-dimethyltetrahydrofuro[2,3-*d*][1,3]dioxol-6-yl]ethyl}-1,2,3-benzotriazin-4(3*H*)-one

To the solution of compound (8.5g), obtained from step b above, in a solvent mixture of tetrahydrofuran: methanol (60mL:40mL) was added 10% Palladium on charcoal (4g) at room temperature and the reaction mixture was hydrogenated with hydrogen at the pressure of 35 psi for 4 hours in a Paar apparatus. The reaction mixture was filtered through a celite pad and the residue was washed with methanol. The filtrate was concentrated to afford the title compound (8.1g).

Step d: Synthesis of 3-{2-[(2*R*,3*R*,4*S*,5*R*)-4,5-dihydroxy-2-{2-[4-(6-methoxypyridin-3-yl)phenyl]ethyl}tetrahydrofuran-3-yl]ethyl}-1,2,3-benzotriazin-4(3*H*)-one

Perchloric acid (4.8ml) was added to a solution of compound (8.1gms), obtained from step c above, in acetonitrile (50ml) and water (8ml) at room temperature. The reaction mixture was heated to 55°C for 30 minutes. The reaction mixture was then quenched using sodium bicarbonate solution. The solvents were evaporated under reduced pressure. The residue thus obtained was taken up in ethyl acetate and water. The organic layer was separated and washed with water and brine solution and dried over anhydrous sodium sulphate. Solvent was concentrated under reduced pressure to yield the title compound (8gm).

Step e: Synthesis of (2*S*,3*R*)-3-(formyloxy)-5-[4-(6-methoxypyridin-3-yl)phenyl]-2-[2-(4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)ethyl]pentanoic acid

To a solution of compound (8g), obtained from step d above, in *tert*-butanol: tetrahydrofuran (40mL:40mL) at 0°C was added a solution of sodium metaperiodate (14g in 40 mL of water). The reaction mixture was stirred for 2 hours at the same temperature and potassium permanganate (518mg) was added at 0°C. After stirring the reaction mixture for an additional 6 hours at room temperature, the reaction mixture was evaporated on rotary evaporator. The residue was taken into water and extracted with ethyl acetate. The organic layer was washed with water and brine solution and dried over anhydrous sodium sulphate. The solvent was evaporated under the reduced pressure, to afford the title compound (8g).

Step f: Synthesis of (2*S*,3*R*)-3-hydroxy-5-[4-(6-methoxypyridin-3-yl)phenyl]-2-[2-(4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)ethyl]pentanoic acid

Potassium carbonate (2.3g) was added to a solution of compound (8g), obtained from step e above, in methanol (40mL) and tetrahydrofuran (30mL) at 0°C. The reaction mixture was stirred at room temperature for 3 hours. Solvents were evaporated and the residue was taken into water and ethyl acetate. The organic layer was washed with water and brine solution and dried over anhydrous sodium sulphate. Solvent was evaporated under reduced pressure to obtain a residue which was purified by column chromatography over silica gel using 60% ethyl acetate in hexane as eluant to afford the title compound (3g). **Mass (m/z):** 474.87(M⁺ +1); **¹H NMR (400 MHz, MeOD):** δ 8.32- 8.30 (2H, m), 8.16- 8.14 (1H, m), 8.04- 8.03 (1H, m), 7.92- 7.88 (2H, m), 7.46 (2H, d, J= 8Hz), 7.24 (2H, m, J= 8Hz), 6.86 (1H, d, J= 8Hz), 4.57- 4.53 (2H, m), 3.93 (3H, s), 3.78-3.83 (1H, m), 2.83- 2.80 (1H, m), 2.63- 2.60 (1H, m), 2.52- 2.49 (1H, m), 2.32- 2.28 (2H, m), 1.80- 1.76 (2H, m).

Example 1B: Synthesis of (2*S*,3*R*)-3-hydroxy-5-[4-(6-hydroxypyridin-3-yl)phenyl]-2-[2-(4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)ethyl]pentanoic acid (Compound No. 226)

To a solution of compound(0.2 gm), obtained from Example1A above in dry toluene (5ml), cooled to -78 °C, was added boron tribromide (0.5 ml) and the contents were stirred at room temperature for 4 hours. Ethyl acetate and water were added to the reaction mixture. Organic layer was separated, washed with water and brine and dried over anhydrous sodium sulfate. Solvent was evaporated to obtain a residue which was purified by preparatory thin layer chromatography (2 mm thickness) using 15% methanol in dichloromethane as eluent to get the title compound (60mg). **Mass (m/z):** 461.23 (M⁺ +1); **¹H NMR (CD₃OD):** 8.30 (d, 1H, J = 8Hz), 8.15 (d, 1H, J = 8Hz), 8.06-8.01 (m, 2H), 7.88 (t, 1H, J = 8Hz), 7.79-7.76 (m, 1H), 7.43 (d, 2H, J = 12Hz), 7.25 (d, 2H, J = 12Hz), 6.75 (d, 1H, J = 12Hz), 4.58- 4.51 (m, 2H), 3.82-3.76 (m, 1H), 2.88 -2.78 (m, 1H), 2.66 - 2.57 (m, 1H), 2.54- 2.47 (m, 1H), 2.33- 2.25 (m, 2H), 1.84- 1.70 (m, 2H).

Example 1C: Synthesis of (2*S*,3*R*)-3-(acetyloxy)-5-[4-(6-methoxypyridin-3-yl)phenyl]-2-[2-(4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)ethyl]pentanoic acid (Compound No. 229)

Diisopropyl azodicarboxylate (63 mg) was added to a solution of compound (100 mg) obtained from Example1A above, tri-*n*-butylphosphine (93 mg) and acetic acid (1 ml)

in dry tetrahydrofuran (3 ml) at 0 °C. Reaction mixture was stirred for 1 hour and concentrated. Residue was taken in ethyl acetate and washed with water and brine. Organic layer was concentrated to get oily residue which was purified on preparative thin layer chromatography (2 mm thickness) using 10% methanol in dichloromethane to get the title compound (80 mg) (formed probably by esterification of acetic acid (used in large excess) and not due to Mitsunobu reaction). **Mass (m/z):** 517.06 ($M^+ + 1$); **¹H NMR:** δ 8.37 (m, 2H), 8.15 (d, 1H, J = 8Hz), 7.97- 7.93 (m, 1H), 7.82- 7.80 (m, 1H), 7.41 (d, 2H, J = 8Hz), 7.20 (d, 2H, J = 8Hz), 6.83 (d, 1H, J = 8Hz), 5.28- 5.27 (m, 1H), 4.56- 4.58 (m, 2H), 4.01 (s, 3H), 2.85- 2.80 (m, 1H), 2.55- 2.75 (m, 2H), 2.31- 2.29 (m, 2H), 2.25- 2.10 (m, 2H), 2.063 (s, 3H).

Example 1D: Synthesis of (2S,3R)-2-[2-(1,3-dioxo-1,3-dihydro-2H-isoindol-2-yl)ethyl]-3-hydroxy-5-[4-(6-methoxypyridin-3-yl)phenyl]pentanoic acid (Compound No.88)

Step a: Synthesis of 5-O-[tert-butyl(diphenyl)silyl]-3-deoxy-3-[2-(1,3-dioxo-1,3-dihydro-2H-isoindol-2-yl)ethyl]-1,2-O-(1-methylethylidene)- β -D-lyxofuranose

5-O-[tert-butyl(diphenyl)silyl]-3-deoxy-3-(2-hydroxyethyl)-1,2-O-isopropylidene- β -D-lyxofuranose (14g), triphenylphosphine (12g) and phthalimide (5g) were dried in high vacuum in a round bottom flask for 10 minutes. Then the vacuum was released under a nitrogen atmosphere and tetrahydrofuran (100mL) was added to the above reaction mixture. The reaction mixture was cooled to 0 °C and diethyl azodicarboxylate (7.8mL) was added slowly. The reaction mixture was stirred for 30 minutes at same temperature, and then quenched with water and extracted with ethyl acetate. The organic layer was washed with water, brine solution, dried over anhydrous sodium sulphate and concentrated to obtain a residue which was purified by column chromatography over silica gel using 30% ethyl acetate in hexane as eluant to afford the title compound (11g).

Step b: Synthesis of 3-deoxy-3-[2-(1,3-dioxo-1,3-dihydro-2H-isoindol-2-yl)ethyl]-1,2-O-(1-methylethylidene)- β -D-lyxofuranose

To a solution of compound (11g), obtained from step a above, in dry tetrahydrofuran (50mL) at 0 °C, was added tetra-butyl ammonium fluoride (41.36mL). The resulting mixture was initially stirred at 0 °C for 1 hour, and then at room temperature for 4 hours. The reaction mixture was cooled to 0 °C, quenched with saturated ammonium chloride and extracted with ethyl acetate. The organic layer was washed with water and

brine, dried over anhydrous sodium sulphate and concentrated. The residue thus obtained was purified by column chromatography over silica gel using 80% ethyl acetate in hexane as eluant to furnish the title compound (5 g).

Step c: Synthesis of (5*S*)-3-deoxy-3-[2-(1,3-dioxo-1,3-dihydro-2*H*-isoindol-2-yl)ethyl]-4,5-*O*-(1-methylethylidene)-D-*arabino*-pentodialdo-5,2-furanose

Oxalyl chloride (4.5mL) and dichloromethane (20mL) were taken in a three neck round bottom flask and cooled to -78°C. Dimethylsulfoxide (6.6mL) was added dropwise to the reaction mixture. The reaction mixture was warmed to -35°C for 5 to 10 minutes and again cooled to -78°C. A solution of compound (7.2 g), obtained from step b above, in dichloromethane (20mL) was added slowly while maintaining the same temperature. The reaction mixture was stirred for 45 minutes till the reaction temperature reached -35°C. The reaction mixture was again cooled to -78°C and triethylamine (17.3mL) was added. The reaction mixture was stirred for an additional 30 minutes, and the temperature was allowed to reach -35°C. The reaction mixture was quenched with saturated solution of ammonium chloride and extracted with dichloromethane. The combined organic layers were washed with water and brine solution, dried over anhydrous sodium sulphate and concentrated to furnish the title compound (7.3g).

Step d: Synthesis of 2-(2-((3*aS*,5*R*,6*S*,6*aS*)-5-[(*E*)-2-(4-bromophenyl)ethenyl]-2,2-dimethyltetrahydrofuro[2,3-*d*][1,3]dioxol-6-yl)ethyl)-1*H*-isoindole-1,3(2*H*)-dione

To the suspension of potassium *t*-butoxide (2.3g) in dimethyl sulfoxide (30ml) cooled to 0°C, was added (4-bromobenzyl)triphenylphosphonium bromide (12.2gm) in dimethyl sulfoxide (20ml). After 20 minutes, a solution of compound (4.1gms) obtained from step c above, in dimethyl sulfoxide (10ml) was added drop wise and reaction mixture was stirred for 1 hour at the same temperature. Reaction mixture was quenched with water and extracted with ethyl acetate. Organic layer was washed with water and brine solution and dried over anhydrous sodium sulphate. Solvent was evaporated under the reduced pressure. Purification was performed on silica gel column by using 40% ethyl acetate in hexane as eluant to furnish the title compound (4gms).

Step e: Synthesis of 2-{2-[(3*aS*,5*R*,6*S*,6*aS*)-5-[(*E*)-2-[4-(6-methoxypyridin-3-yl)phenyl]ethenyl]-2,2-dimethyltetrahydrofuro[2,3-*d*][1,3]dioxol-6-yl]ethyl}-1*H*-isoindole-1,3(2*H*)-dione

A mixture of compound (1g), obtained from step d above, 2-methoxy-5-pyridine
5 boronic acid (0.614g), tetrakis(triphenyl)phosphinepalladium (0) (0.115g) and potassium carbonate (0.832g) was dried under high vacuum for 10 minutes and dry dimethylformamide (8mL) was added at room temperature. The reaction mixture was heated at 120°C for 2 hours, and then quenched with water and extracted with ethyl acetate. The organic layer was washed with water and brine solution and dried over
10 anhydrous sodium sulphate. Solvent was evaporated under the reduced pressure to obtain a residue which was purified by column chromatography over silica gel using 30% ethyl acetate in hexane as eluant to afford the title compound (0.62g).

Step f: Synthesis of 2-{2-[(3*aS*,5*R*,6*S*,6*aS*)-5-{2-[4-(6-methoxypyridin-3-yl)phenyl]ethyl}-2,2-dimethyltetrahydrofuro[2,3-*d*][1,3]dioxol-6-yl]ethyl}-1*H*-isoindole-1,3(2*H*)-dione

To the solution of compound (0.6g), obtained from step e above, in a methanol (10mL), was added 10% Palladium on charcoal (0.05g) at room temperature and the reaction mixture was hydrogenated with hydrogen at the pressure of 35 psi for 4 hours in a Paar apparatus. The reaction mixture was filtered through a celite pad and the residue was
20 washed with methanol. The filtrate was concentrated to afford the title compound (0.56g).

Step g: Synthesis of 2-{2-[(2*R*,3*R*,4*S*,5*R*)-4,5-dihydroxy-2-{2-[4-(6-methoxypyridin-3-yl)phenyl]ethyl}tetrahydrofuran-3-yl]ethyl}-1*H*-isoindole-1,3(2*H*)-dione

Perchloric acid (0.4mL) was added to a solution of compound (0.5g), obtained from step f above, in acetonitrile (4mL) and water (0.8mL) at room temperature. The
25 reaction mixture was heated to 55°C for 30 minutes. The reaction mixture was then quenched using sodium bicarbonate solution. The solvents were evaporated under reduced pressure. The residue thus obtained was taken up in ethyl acetate and water. The organic layer was separated and washed with water and brine solution and dried over anhydrous sodium sulphate. Solvent was concentrated under reduced pressure to yield the
30 title compound (0.51g).

Step h: Synthesis of (1*R*,2*S*)-4-(1,3-dioxo-1,3-dihydro-2*H*-isoindol-2-yl)-2-formyl-1-{2-[4-(6-methoxypyridin-3-yl)phenyl]ethyl}butyl formate

To a solution of compound (0.56g), obtained from step g above, in methanol (4mL) at 0°C was added a solution of sodium metaperiodate (0.736 g in 1 mL of water).

- 5 The reaction mixture was stirred for 2 hours at the same temperature. After stirring the reaction mixture for an additional 1hour at room temperature, the reaction mixture was evaporated on rotary evaporator. The residue was taken into water and extracted with ethyl acetate. The organic layer was washed with water and brine solution and dried over anhydrous sodium sulphate. The solvent was evaporated under the reduced pressure to
10 obtained the title compound (0.54g).

Step i: Synthesis of (2*S*,3*R*)-2-[2-(1,3-dioxo-1,3-dihydro-2*H*-isoindol-2-yl)ethyl]-3-(formyloxy)-5-[4-(6-methoxypyridin-3-yl)phenyl]pentanoic acid

The compound (0.54 g) obtained was taken in acetonitrile (6mL) and water (1mL). To this solution, sodium dihydrogen phosphate (0.054g) was added. Reaction
15 mixture was cooled to 0°C and hydrogen peroxide (1mL) and sodium chlorite (0.208 g) were added. After stirring the reaction mixture for additional 1hour, solvents were evaporated on rotaevaporator; the residue was taken into water and extracted with ethyl acetate. Organic layer was washed with water and brine solution and dried over anhydrous sodium sulphate. Solvent was evaporated under the reduced pressure, to get the title
20 compound (0.5 g)

Step j: Synthesis of (2*S*,3*R*)-2-[2-(1,3-dioxo-1,3-dihydro-2*H*-isoindol-2-yl)ethyl]-3-hydroxy-5-[4-(6-methoxypyridin-3-yl)phenyl]pentanoic acid

- Potassium carbonate (0.165g) was added to a solution of compound (0.4g), obtained from step i above, in methanol (6mL) at 0°C. The reaction mixture was stirred at
25 room temperature for 3 hours. Solvents were evaporated and the residue was taken into water and ethyl acetate. The organic layer was washed with water and brine solution and dried over anhydrous sodium sulphate. Solvent was evaporated under reduced pressure to obtain a residue which was purification with preparatory TLC using 7% methanol in ethylacetate as eluent to get the title compound (0.13gm). **Mass (m/z):** 474.88(M⁺ +1);
30 ¹HNMR (CDCl₃):- δ 8.35 (s, 1H), 7.83-7.81 (m, 2H), 7.77-7.69 (m, 3H), 7.42 (d, 2H, J =

3Hz), 7.25 (d, 2H, J = 6Hz), 6.80 (d, 1H, J = 6Hz), 3.97 (s, 3H), 3.87-3.78 (m, 3H), 2.89(m, 1H), 2.69 (m, 1H), 2.57(m, 1H), 2.15-1.98 (m, 2H), 1.84-1.79 (m, 2H);

Example 1E: Synthesis of (2S,3R)-5-(2',4'-difluorobiphenyl-4-yl)-2-[2-(1,3-dioxo-1,3-dihydro-2H-isoindol-2-yl)ethyl]-3-hydroxypentanoic acid (Compound No. 97)

5 Step a: Synthesis of 2-(2-((3aS,5R,6S,6aS)-5-[(E)-2-(3',5'-difluorobiphenyl-4-yl)ethenyl]-2,2-dimethyltetrahydrofuro[2,3-d][1,3]dioxol-6-yl)ethyl)-1H-isoindole-1,3(2H)-dione

A mixture of compound (0.2g), obtained from step d of Example 1D above, 3,5-difluorophenyl boronic acid (0.135g), tetrakis(triphenyl)phosphinepalladium (0) (0.023g) and potassium carbonate (0.2g) was dried under high vacuum for 10 minutes and dry dimethylformamide (3mL) was added at room temperature. The reaction mixture was heated at 120°C for 2 hours, and then quenched with water and extracted with ethyl acetate. The organic layer was washed with water and brine solution and dried over anhydrous sodium sulphate. Solvent was evaporated under the reduced pressure to obtain a residue which was purified by column chromatography over silica gel using 30% ethyl acetate in hexane as eluant to afford the title compound (0.205g).

Step b: Synthesis of 2-(2-((3aS,5R,6S,6aS)-5-[2-(3',5'-difluorobiphenyl-4-yl)ethyl]-2,2-dimethyltetrahydrofuro[2,3-d][1,3]dioxol-6-yl)ethyl)-1H-isoindole-1,3(2H)-dione

To the solution of compound (0.2g), obtained from step a above, in a tetrahydrofuran (10mL), was added 10% Palladium on charcoal (0.1g) at room temperature and the reaction mixture was hydrogenated with hydrogen at the pressure of 35 psi for 4 hours in a Paar apparatus. The reaction mixture was filtered through a celite pad and the residue was washed with methanol. The filtrate was concentrated to afford the title compound (0.2g).

25 Step c: Synthesis of 2-(2-((2R,3R,4S,5R)-2-[2-(3',5'-difluorobiphenyl-4-yl)ethyl]-4,5-dihydroxytetrahydrofuran-3-yl)ethyl)-1H-isoindole-1,3(2H)-dione

Perchloric acid (0.3mL) was added to a solution of compound (0.2g), obtained from step b above, in acetonitrile (6mL) and water (2mL) at room temperature. The reaction mixture was heated to 55°C for 30 minutes. The reaction mixture was then quenched using sodium bicarbonate solution. The solvents were evaporated under

reduced pressure. The residue thus obtained was taken up in ethyl acetate and water. The organic layer was separated and washed with water and brine solution and dried over anhydrous sodium sulphate. Solvent was concentrated under reduced pressure to yield the title compound (0.15g).

5 **Step d: Synthesis of (1*R*,2*S*)-1-[2-(3',5'-difluorobiphenyl-4-yl)ethyl]-4-(1,3-dioxo-1,3-dihydro-2*H*-isoindol-2-yl)-2-formylbutyl formate**

To a solution of compound (0.15g), obtained from step c above, in methanol (6mL) at 0°C was added a solution of sodium metaperiodate (0.15 g in 1 mL of water). The reaction mixture was stirred for 2 hours at the same temperature. After stirring the reaction mixture for an additional 1 hour at room temperature, the reaction mixture was evaporated on rotary evaporator. The residue was taken into water and extracted with ethyl acetate. The organic layer was washed with water and brine solution and dried over anhydrous sodium sulphate. The solvent was evaporated under the reduced pressure to obtained the title compound (0.12g).

15 **Step e: Synthesis of (2*S*,3*R*)-5-(3',5'-difluorobiphenyl-4-yl)-2-[2-(1,3-dioxo-1,3-dihydro-2*H*-isoindol-2-yl)ethyl]-3-(formyloxy)pentanoic acid**

The compound (0.12 g) obtained from step d was taken in acetonitrile (6mL) and water (1mL), to this solution sodium dihydrogen phosphate (0.05g) was added. Reaction mixture was cooled to 0°C and hydrogen peroxide (0.5mL) and sodium chlorite (0.15 g) were added. After stirring the reaction mixture for additional 1 hour, solvents were evaporated on rotaevaporator, the residue was taken into water and extracted with ethyl acetate. Organic layer was washed with water and brine solution and dried over anhydrous sodium sulphate. Solvent was evaporated under the reduced pressure, to get the title compound (0.12 g)

25 **Step f: Synthesis of (2*S*,3*R*)-5-(3',5'-difluorobiphenyl-4-yl)-2-[2-(1,3-dioxo-1,3-dihydro-2*H*-isoindol-2-yl)ethyl]-3-hydroxypentanoic acid**

Potassium carbonate (0.06g) was added to a solution of compound (0.12g), obtained from step e above, in methanol (10mL) at 0°C. The reaction mixture was stirred at room temperature for 3 hours. Solvents were evaporated and the residue was taken into water and ethyl acetate. The organic layer was washed with water and brine solution and dried over anhydrous sodium sulphate. Solvent was evaporated under reduced pressure to

obtain a residue which was purification with preparatory thin layer chromatography using 10% methanol in ethylacetate as eluent to get the title compound (0.025gm). **Mass (m/z):** 480.13($M^+ + 1$); **¹HNMR (CD₃OD):** δ 7.48-7.76 (m, 4H), 7.47-7.37 (m, 3H), 7.25-7.23 (d, 2H), 7.03-6.98 (m, 2H), 3.78-3.72 (m, 3H), 2.85-2.81 (m 1H), 2.65-2.63 (m, 1H), 2.422.40 (m, 1H), 2.10-2.05 (m, 2H), 1.79-1.75 (m, 2H);

Example 2: Synthesis of (2S,3R)-3-hydroxy-2-[2-(4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-5-phenylpentanoic acid (Compound No. 82)

Step a: Synthesis of 3-{2-[(3aS,5R,6S,6aS)-2,2-dimethyl-5-(2-phenylethyl)tetrahydrofuro[2,3-d][1,3]dioxol-6-yl]ethyl}-1,2,3-benzotriazin-4(3H)-one

To the solution of compound (0.3g), obtained from step j of Example 1, in a solvent mixture of tetrahydrofuran: methanol (10mL:10 mL) was added palladium/carbon (0.2 g, 10%) at room temperature and the reaction mixture was hydrogenated at the pressure of 35 psi for 4 hours in a Paar apparatus. The reaction mixture was filtered through a celite pad and the residue was washed with methanol. The filtrate was concentrated to afford the title compound (0.3 g). **Mass (m/z):**421 (M^+)

Step b: Synthesis of 3-{2-[(2R,3R,4S,5R)-4,5-dihydroxy-2-(2-phenylethyl)tetrahydrofuran-3-yl]ethyl}-1,2,3-benzotriazin-4(3H)-one

Perchloric acid (0.2 mL) was added to a solution of compound (0.3g), obtained from step a above, in acetonitrile (4 mL) and water (0.3 mL) at room temperature. The reaction mixture was heated to 55°C for 30 minutes. The reaction mixture was then quenched using a sodium bicarbonate solution. The solvents were evaporated at reduced pressure. The residue thus obtained was taken up in ethyl acetate and water. The organic layer was separated and washed with water and brine solution and dried over anhydrous sodium sulphate. The solvent was concentrated under reduced pressure to yield the title compound (0.3 g).

Step c: Synthesis of (2S,3R)-3-(formyloxy)-2-[2-(4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-5-phenylpentanoic acid

To a solution of compound (0.3 g), obtained from step b above, in a *tert*-butanol:tetrahydrofuran (3mL:3mL) at 0°C was added a solution of sodium metaperiodate

(0.673 g in 3mL of water). The reaction mixture was stirred for 2 hours at same temperature and potassium permanganate (0.025 g) was added at 0°C. After stirring the reaction mixture for an additional 6 hours at room temperature, the reaction mixture was evaporated, the residue was taken into water and extracted with ethyl acetate. The organic layer was washed with water and brine solution and dried over anhydrous sodium sulphate. The solvent was evaporated under reduced pressure, to afford the title compound (0.3 g).

Step d: Synthesis of (2S,3R)-3-hydroxy-2-[2-(4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-5-phenylpentanoic acid

Potassium carbonate (0.11 g) was added to a solution of compound (0.3 g), obtained from step c above, in methanol (4mL) and tetrahydrofuran (4mL) at 0°C. The reaction mixture was stirred at room temperature for 3 hours. The solvents were evaporated and the residue was taken into water and ethyl acetate. The organic layer was washed with water and brine solution and dried over anhydrous sodium sulphate. The solvent was evaporated under reduced pressure to obtain a residue which was purified by column chromatography over silica gel using 10% methanol in dichloromethane as eluant to afford the title compound (0.020g). Mass (m/z):368.0 ($M^+ + 1$); ^1H NMR (CD_3OD): δ 8.31 (d, 1H, J = 8Hz), 8.15 (d, 1H, J = 8Hz), 8.06- 8.02 (m, 1H), 7.91- 7.89 (m, 2H), 7.62 (d, 2H, J = 8Hz), 7.36 (d, 1H, J = 8Hz), 7.21-7.07 (m, 1H), 4.57- 4.52 (m, 2H), 3.78-3.76 (m, 1H), 2.76 -2.75 (m, 1H), 2.57 - 2.49 (m, 2H), 2.29- 2.27 (m, 2H), 1.74- 1.71 (m, 2H).

Example 3: Synthesis of (2S,3R)-2-[2-(1,3-dioxo-1,3-dihydro-2H-isoindol-2-yl)ethyl]-3-hydroxy-5-[4(trifluoromethyl)phenyl]pentanoic acid (Compound No.84)

Step a: Synthesis of 5-O-[tert-butyl(diphenyl)silyl]-3-deoxy-3-[2-(1,3-dioxo-1,3-dihydro-2H-isoindol-2-yl)ethyl]-1,2-O-isopropylidene- β -D-lyxofuranose

A mixture of compound (0.14g), obtained from step f of Example 1, triphenylphosphine (0.080g) and phthalimide (0.045g) were taken in a round bottom flask and dried in high vacuum for 10 minutes. The vacuum was released under a nitrogen atmosphere and tetrahydrofuran (5mL) was added to the reaction mixture. The reaction mixture was cooled to 0°C and diethyl azodicarboxylate (0.1 mL) was added slowly. The reaction mixture was stirred for 30 minutes at same temperature and then quenched with water and extracted with ethyl acetate. The organic layer was washed with water, brine

solution, dried over anhydrous sodium sulphate and concentrated. The residue thus obtained was purified by column chromatography over silica gel using 30 % ethyl acetate in hexane as eluant to afford the title compound (0.11g). Mass (m/z): 608 ($M^+ + 23$)

Step b: Synthesis of 3-deoxy-3-[2-(1,3-dioxo-1,3-dihydro-2H-isoindol-2-yl)ethyl]-1,2-O-isopropylidene- β -D-lyxofuranose

To a solution of compound (0.060g), obtained from step a above, in dry tetrahydrofuran (2 mL) at 0°C, was added tetra-butyl ammonium fluoride (0.2 mL). The resulting mixture was initially stirred at 0°C for 1 hour, and then at room temperature for 4 hours. The reaction mixture was cooled to 0°C, quenched with saturated ammonium chloride and extracted with ethyl acetate. The organic layer was washed with water and brine, dried over anhydrous sodium sulphate and concentrated. The residue thus obtained was purified by column chromatography over silica gel using 60% ethyl acetate in hexane as eluant to furnish the title compound (0.4 g).

Step c: Synthesis of (5S)-3-deoxy-3-[2-(1,3-dioxo-1,3-dihydro-2H-isoindol-2-yl)ethyl]-4,5-O-isopropylidene-D-arabino-pentodialdo-5,2-furanose

In a three-neck round bottom flask, oxalyl chloride (0.6 mL) and dichloromethane (20 mL) were placed and cooled to -78°C. Dimethylsulfoxide (1mL) was added dropwise to the reaction mixture. The reaction mixture was warmed to -35°C for 5 to 10 minutes and again cooled to -78°C. A solution of compound (1 g), obtained from step b above, in dichloromethane (5mL) was added slowly while maintaining the same temperature. The reaction mixture was stirred for 45 minutes till reaction temperature reached -35°C. The reaction mixture was again cooled to -78°C and triethylamine (2.4 mL) was added. The reaction mixture was stirred for an additional 30 minutes and the temperature was allowed to reach -35°C. The reaction mixture was quenched with saturated solution of ammonium chloride and extracted with dichloromethane. The combined organic layer was washed with water and brine solution, dried over anhydrous sodium sulphate and concentrated to furnish the title compound (1.0 g).

Step d: Synthesis of 2-{2-[(3aS,5R,6S,6aS)-2,2-dimethyl-5-{(E)-2-[4-(trifluoromethyl)phenyl]vinyl}tetrahydrofuro[2,3-d][1,3]dioxol-6-yl]ethyl}-1H-isoindole-1,3(2H)-dione

To the suspension of triphenyl[4-(trifluoromethyl)benzyl]phosphonium bromide (0.795 g) in dimethylsulphoxide (5 mL), potassium tert-butoxide (0.292g) was added. A solution of compound (0.5 g), obtained from step c above, in tetrahydrofuran (7 mL) was added dropwise after 20 minutes, and the reaction mixture was stirred for 1 hour at the same temperature. The reaction mixture was quenched with water and extracted with ethyl acetate. The organic layer was washed with water and brine solution and dried over anhydrous sodium sulphate. The solvent was evaporated under reduced pressure and the residue thus obtained was purified by column chromatography using 30% ethyl acetate in hexane as eluant to furnish the title compound (0.25g). Mass (m/z):487 (M⁺)

Step e: Synthesis of 2-{2-[(3aS,5R,6S,6aS)-2,2-dimethyl-5-{2-[4-(trifluoromethyl)phenyl]ethyl}tetrahydrofuro[2,3-d][1,3]dioxol-6-yl]ethyl}-1H-isoindole-1,3(2H)-dione

10% Palladium on charcoal (0.2 g) was added to the solution of compound (0.25g), obtained from step d above, in ethyl acetate (20 mL) at room temperature and the reaction mixture was hydrogenated at 50 psi for 4 hours in a Paar apparatus. The reaction mixture was filtered through celite pad and the residue was washed with methanol. The filtrate was concentrated to afford the title compound (0.225 g).

Step f: Synthesis of 2-{2-[(2R,3R,4S,5R)-4,5-dihydroxy-2-{2-[4-(trifluoromethyl)phenyl]ethyl}tetrahydrofuran-3-yl]ethyl}-1H-isoindole-1,3(2H)-dione

Perchloric acid (0.2 mL) was added to a solution of compound (0.225 g), obtained from step e above, in acetonitrile (4mL), water (1 mL) and tetrahydrofuran (0.5 mL) at room temperature. The reaction mixture was heated to 55°C for 30 minutes. The reaction mixture was then quenched using sodium bicarbonate solution. The solvents were evaporated at reduced pressure. The residue thus obtained was taken in ethyl acetate and water. The organic layer was separated and washed with water and brine solution and dried over anhydrous sodium sulphate. The solvent was concentrated under reduced pressure to yield the title compound (0.22g).

Step g: Synthesis of (1R,2S)-4-(1,3-dioxo-1,3-dihydro-2H-isoindol-2-yl)-2-formyl-1-{2-[4-(trifluoromethyl)phenyl]ethyl}butyl formate

A solution of sodium metaperiodate (0.313 g in 1 mL of water) was added to a solution of compound (0.22g), obtained from step f above, in a methanol:tetrahydrofuran(3 mL:1 mL) at 0°C. The reaction mixture was stirred for 2 hours at same temperature. After stirring the reaction mixture for an additional 6 hours at room temperature, the reaction mixture was evaporated on a rotary-evaporator, and the residue was taken into water and extracted with ethyl acetate. The organic layer was washed with water and brine solution and dried over anhydrous sodium sulphate. The solvent was evaporated under the reduced pressure, to afford the title compound (0.22 g).

Step h: Synthesis of (2S,3R)-2-[2-(1,3-dioxo-1,3-dihydro-2H-isoindol-2-yl)ethyl]-3-(formyloxy)-5-[4-(trifluoromethyl)phenyl]pentanoic acid

To a solution of compound (0.22g), obtained from step g above, in a solvent mixture of acetonitrile:water (3mL:1mL) at 0°C, were added sodium dihydrogenphosphate (0.021g), sodium chlorite (0.081g) and hydrogen peroxide (1mL, 30% in water). The reaction mixture was stirred for 2 hours at room temperature. The solvents were evaporated under reduced pressure to obtain a residue. Ethyl acetate and water were added to the resulting residue. The organic layer was separated, washed with water and brine, dried over anhydrous sodium sulphate. The solvent was evaporated under the reduced pressure, to afford the title compound (0.2 g).

Step i: Synthesis of (2S,3R)-2-[2-(1,3-dioxo-1,3-dihydro-2H-isoindol-2-yl)ethyl]-3-hydroxy-5-[4-(trifluoromethyl)phenyl]pentanoic acid

Potassium carbonate (0.178 g) was added to a solution of compound (0.2g), obtained from step h above, in methanol (7 mL) at 0°C. The reaction mixture was stirred at room temperature for 3 hours. The solvents were evaporated and the residue was taken into water and ethyl acetate. The organic layer was washed with water and brine solution and dried over anhydrous sodium sulphate. The solvent was evaporated under reduced pressure obtain a residue which was purified by column chromatography over silica gel using 10 % methanol in dichloromethane as eluant to afford the title compound (0.090 g). Mass (m/z): 458.0(M⁺ +23)

¹HNMR (CD₃OD): δ 7.83 (q, 2H, J = 3.1 Hz), 7.71 (t, 2H, J = 3.2 Hz), 7.51 (d, 2H, J = 7.6 Hz), 7.28 (d, 2H, J = 7.9 Hz), 3.95-3.76 (m, 3H), 2.93-2.55 (m, 3H), 2.14-1.75 (m, 4H)

Example 4: Synthesis of (2R,3R)-5-(4'-chlorobiphenyl-4-yl)-2-[2-(1,3-dioxo-1,3-dihydro-2H-isoindol-2-yl)ethyl]-3-hydroxypentanoic acid (Compound No. 89)

5 Step a: Synthesis of 1,2:5,6-Di-O-isopropyliden-α-D-ribo-3-hexulo-furanose

Oxalyl chloride (25.16 mL) and dichloromethane (200 mL) were taken in a three-neck round bottom flask and cooled to -75 °C under a nitrogen atmosphere. Then dimethylsulfoxide (27.3 mL) was added dropwise slowly maintaining the temperature at -70 °C. The reaction mixture was stirred for 30 minutes at the same temperature, and then
10 diacetone-α-D-glucose (50 g) in dichloromethane (500 mL) was charged slowly maintaining the temperature at -70 °C. After 20 minutes, triethylamine (80 mL) was added dropwise to the above mixture at the same temperature. Saturated solution of ammonium chloride in water (500 mL) was charged to the reaction mixture after 30 minutes and the temperature was allowed to rise to room temperature. The reaction mixture was extracted
15 with dichloromethane. The organic layer was dried over anhydrous sodium sulphate. The solvent was evaporated under reduced pressure to afford the title compound (45 g).

Step b : Synthesis of 1,2:5,6-Di-O-isopropylidene-3-deoxy-3-(2-ethoxy-2-oxoethylidene)- α -D-ribo-hexofuranose

To the suspension of sodium hydride (0.155 g, 60% in oil.) in tetrahydrofuran (5
20 mL) at 0 °C was charged triethyl phosphonoacetate (1.55 mL). After 20 minutes, a solution of compound (1g), obtained from step a above, in tetrahydrofuran (3 mL) was added dropwise and the reaction mixture was stirred for 1 hour at the same temperature. A saturated solution of ammonium chloride was added to the reaction mixture. The resulting mixture was extracted with ethyl acetate. Combined extracts were dried over anhydrous
25 sodium sulphate and evaporated under reduced pressure to yield a residue which was purified by column chromatography over silica gel using 30 % ethyl acetate in hexane as eluant to afford the title compound (1.3 g). Mass (m/z): 351.1(M⁺ +23)

Step c: Synthesis of 1,2:5,6-Di-O-isopropylidene-3-deoxy-3-(2-ethoxy-2-oxoethyl)- α -D-allofuranose

30 10% Palladium on charcoal (0.05 g) was added to the solution of compound (0.15 g), obtained from step b above, in ethyl acetate (10 mL) at room temperature and hydrogen

was supplied at 50 psi for 4 hours. The reaction mixture was filtered through a celite pad and the residue was washed with ethyl acetate. The filtrate was concentrated to afford the title compound (0.12g). Mass (m/z): 353.2 ($M^+ + 1$).

Step d: Synthesis of 3-deoxy-3-(2-ethoxy-2-oxoethyl)-1,2-O-isopropylidene- α -D-allofuranose

30 % Perchloric acid (4 mL) was added to a solution of compound (2.0 g), obtained from step c above, in tetrahydrofuran (20 mL) at -5°C - 0°C . The reaction mixture was stirred for 5 hours at -5°C - 0°C and then quenched with saturated solution of sodium bicarbonate (20 mL). Solvents were evaporated, ethyl acetate and water were added to the resulting residue. The organic layer was separated, washed with brine, dried over anhydrous sodium sulphate and concentrated to afford 1.5 g of the title compound. Mass (m/z): 313.2 ($M^+ + 1$)

Step e: Synthesis of 3-deoxy-3-(2-ethoxy-2-oxoethyl)-1,2-O-isopropylidene- α -D-ribo-pentodialdo-1,4-furanose

To a solution of compound (1.5 g), obtained from step d above, in methanol (15 mL) was added aqueous solution of sodium periodate (1.65 g in 10 mL of water) at 0°C . The reaction mixture was stirred for 3 hours from 0°C to room temperature. Solvents were evaporated. Ethyl acetate and water were added to the residue. The organic layer was separated, washed with brine, dried over anhydrous sodium sulphate and concentrated to afford the title compound (0.9g). Mass (m/z): 259.2 ($M^+ + 1$)

Step f: Synthesis of ethyl {(3aR,5R,6R,6aR)-5-[(E)-2-(4-bromophenyl)vinyl]-2,2-dimethyltetrahydrofuro[2,3-d][1,3]dioxol-6-yl}acetate

To the suspension of (4-bromobenzyl)triphenylphosphonium bromide (2.67g) in dimethylsulphoxide (15 mL), potassium *tert*-butoxide (0.508g) was added at 0°C . After stirring the reaction mixture for 30 minutes at room temperature, a solution of compound (0.9 g), obtained from step e above, in tetrahydrofuran (5 mL) was added dropwise at 0°C . The reaction mixture was stirred for 2 hours from 0°C to room temperature, and quenched with ice cold water. Solvents were evaporated, and ethyl acetate and water were added to resulting residue. The organic layer was separated, washed with water and brine solution and dried over anhydrous sodium sulphate. Solvent was evaporated under reduced pressure and the residue thus obtained was purified by column chromatography using 10

% ethyl acetate in hexane as eluant to furnish the title compound (1.0 g). Mass (m/z): 413.3 ($M^+ + 1$)

Step g: Synthesis of ethyl {(3aR,5R,6R,6aR)-5-[(E)-2-(4'-chlorobiphenyl-4-yl)vinyl]-2,2-dimethyltetrahydrofuro[2,3-d][1,3]dioxol-6-yl}acetate

5 A mixture of compound (1.0g), obtained from step f above, (4-chlorophenyl)boronic acid (0.76 g), tetrakis(triphenylphosphine)palladium (0) (0.14 g) and potassium carbonate (1.0 g) in dry dimethylformamide (10 mL) was heated at 110°C for 4 hours. Ethyl acetate and water were added to the reaction mixture. The organic layer was separated, washed with water and brine solution and dried over anhydrous sodium
10 sulphate. Solvent was evaporated under reduced pressure to obtain a residue which was purified by column chromatography over silica gel using 12 % ethyl acetate in hexane as eluant to afford the title compound (0.9 g). Mass (m/z): 465.2 ($M^+ + 23$)

Step h: Synthesis of ethyl {(3aR,5R,6R,6aR)-5-[2-(4'-chlorobiphenyl-4-yl)ethyl]-2,2-dimethyltetrahydrofuro[2,3-d][1,3]dioxol-6-yl}acetate

15 To the solution of compound (0.8 g), obtained from step g above, in ethyl acetate (15 mL) was added 10% Palladium on charcoal (0.25 g) at room temperature. The system was evacuated with hydrogen and the reaction mixture was stirred for 4 hours at room temperature under a hydrogen atmosphere. The reaction mixture was filtered through celite pad and concentrated to afford the title compound (0.75 g). Mass (m/z): 415.4 ($M^+ -$
20 28)

Step i: Synthesis of 2-[(3aR,5R,6R,6aR)-5-[2-(4'-chlorobiphenyl-4-yl)ethyl]-2,2-dimethyltetrahydrofuro[2,3-d][1,3]dioxol-6-yl]ethanol

 To a solution of compound (0.75g), obtained from step h above, in tetrahydrofuran (15 mL) was added, lithium aluminium hydride (0.096 g) at 0°C. The resulting mixture
25 was stirred for 3 hours at same temperature, and a saturated solution of ammonium chloride was then added. The reaction mixture was then filtered through silica gel (100-200 mesh) and concentrated. Ethyl acetate and water was added to the resulting residue. The organic layer was separated, washed with water and brine, dried over anhydrous sodium sulphate, filtered, concentrated under reduced pressure to afford the title
30 compound (0.65 g). Mass (m/z): 425.2 ($M^+ + 23$)

Step j: Synthesis of 2-((3aR,5R,6R,6aR)-5-[2-(4'-chlorobiphenyl-4-yl)ethyl]-2,2-dimethyltetrahydrofuro[2,3-d][1,3]dioxol-6-yl)ethyl methanesulfonate

Triethylamine (0.43 ml) and methanesulfonyl chloride (0.21 mL) were added to a solution of the compound (0.6 g) obtained from step i above, in dichloromethane (15 mL) at 0°C. The reaction mixture was stirred at room temperature for 2 hours. Dichloromethane and water were added to reaction mixture. The organic layer was separated, washed with water, dried and concentrated under reduced pressure to furnish the title compound (0.6g).

Step k: Synthesis of 2-(2-((3aR,5R,6R,6aR)-5-[2-(4'-chlorobiphenyl-4-yl)ethyl]-2,2-dimethyltetrahydrofuro[2,3-d][1,3]dioxol-6-yl)ethyl)-1H-isoindole-1,3(2H)-dione

Potassium phthalimide (0.051 g) was added in one portion to a stirred solution of the compound (0.12 g), obtained from step j above, in dimethylformamide (5 mL) at room temperature under a nitrogen atmosphere. The resulting solution was heated at 50 °C for about 14 hours and then cooled to room temperature. Ethyl acetate and water were added to the reaction mixture. The organic layer was separated, washed with water and brine, dried over anhydrous sodium sulphate. The organic phase was evaporated to dryness under reduced pressure. The residue thus obtained was purified by column chromatography using 30 % ethyl acetate in hexane as eluant to furnish the title compound (0.1 g). Mass (m/z): 549.5 ($M^+ + NH_4^+$)

Step l: Synthesis of 2-(2-((2R,3S,4R,5S)-2-[2-(4'-chlorobiphenyl-4-yl)ethyl]-4,5-dihydroxytetrahydrofuran-3-yl)ethyl)-1H-isoindole-1,3(2H)-dione

Trifluoroacetic acid (1mL) and water (0.5 mL) were added to compound (0.04g), obtained from step k above. The reaction mixture was stirred at room temperature for 2 hours. The solvents were evaporated at reduced pressure. The residue thus obtained was taken in ethyl acetate and water. The organic layer was separated and washed with water and brine solution and dried over anhydrous sodium sulphate. The solvent was concentrated under the reduced pressure to yield the title compound (0.028 g).

Step m: Synthesis of (2R,3R)-5-(4'-chlorobiphenyl-4-yl)-2-[2-(1,3-dioxo-1,3-dihydro-2H-isoindol-2-yl)ethyl]-3-(formyloxy)pentanoic acid

To a solution of compound (0.028g), obtained from step l above, in a *tert*-butanol:water (2:3; 1.3 mL) was added sodium metaperiodate (0.051 g) at room

temperature. The reaction mixture was stirred for 2 hours at same temperature and potassium permanganate (0.001 g) was added at 0°C. After stirring, the reaction mixture for additional 6 hours at room temperature, reaction mixture was evaporated on rotary evaporator, the residue was taken into water and extracted with ethyl acetate. The organic layer was washed with water and brine solution and dried over anhydrous sodium sulphate. The solvent was evaporated under the reduced pressure, to afford the title compound (0.025 g).

Step n: Synthesis of (2R,3R)-5-(4'-chlorobiphenyl-4-yl)-2-[2-(1,3-dioxo-1,3-dihydro-2H-isoindol-2-yl)ethyl]-3-hydroxypentanoic acid

Potassium carbonate (0.007g) was added to a solution of compound (0.025 g), obtained from step m)-above, in methanol (1 mL) at 0°C. The reaction mixture was stirred at room temperature for 3 hours. The solvents were evaporated and the residue was taken into water and ethyl acetate. The organic layer was washed with water and brine solution and dried over anhydrous sodium sulphate. The solvent was evaporated under the reduced pressure obtain a residue which was purified by column chromatography over silica gel using 10 % methanol in dichloromethane as eluant to afford the title compound (0.014 g). Mass (m/z):478.2 (M⁺ +1)

¹HNMR (CD₃OD): δ 7.83 (q, 2H, J = 4Hz), 7.77 (q, 2H, J = 4Hz), 7.56 (d, 2H, J = 8Hz), 7.48 (d, 2H, J = 8Hz), 7.40 (d, 2H, J = 8Hz), 7.26 (d, 1H, J = 8Hz), 3.83-3.70 (m, 3H), 2.85-2.49 (m, 3H), 1.92-1.70(m, 4H).

Example 5: Synthesis of (2R,3S)-2-[2-(1,3-dioxo-1,3-dihydro-2H-isoindol-2-yl)ethyl]-5-(3'-fluoro-4'-methylbiphenyl-4-yl)-3-hydroxypentanoic acid (Compound No. 205)

Step a: Synthesis of 3-O-Acetyl-1,2:5,6-di-O-isopropylidene-α-D-erythrohexofuran-3-enose

Acetic anhydride (65.77 mL) was added to a solution of compound (45 g), obtained from step a of Example 4, in pyridine (500 mL) and the reaction mixture was heated at 60°C for overnight. Reaction mixture was then concentrated to obtain a residue. Ethyl acetate and water were added to the residue. Organic layer was separated, washed with dilute hydrochloric acid, water and brine. The solvent was evaporated under reduced pressure and the residue thus obtained was purified by silica gel column chromatography

using 8% ethyl acetate in hexane as eluant to furnish the title compound (21 g). Mass (m/z): 301.19($M^+ + 1$)

Step b: Synthesis of 3-O-acetyl-1,2:5,6-di-O-isopropylidene- α -D-gulofuranose

To a solution of compound (26 g), obtained from step a above, in ethyl acetate
5 (250 mL), 10% Palladium on charcoal (6g) was added and the reaction mixture was shaken under hydrogen atmosphere at 60 psi for 4 hours on Paar apparatus. The reaction mixture was filtered through a celite bed. The solvents were evaporated to obtain a crude residue which was purified using column chromatography over silica gel using 15 % ethyl acetate in hexane as eluant to furnish the title compound (17 g). Mass (m/z): 324.97 (M^+
10 +Na)

Step c: Synthesis of 1,2:5,6-di-O-isopropylidene- α -D-gulofuranose

Sodium methoxide (12.5 g) was added to a solution of compound (64 g), obtained from step b above, in methanol (10 mL). The reaction mixture was stirred for 3 hours at 0°C. The reaction mixture was then concentrated. The residue thus obtained was purified
15 by column chromatography over silica gel using 30 % ethyl acetate in hexane as eluant to afford the title compound (31 g). Mass (m/z): 261.26 ($M^+ + 1$)

Step d: Synthesis of 1,2:5,6-di-O-isopropylidene- α -D-xylo-hexofuranos-3-ulose

Sodium hypochlorite (225 mL, 4% solution) was added dropwise to a solution compound (30 g), obtained from step c above, followed by addition of 2,2,6,6,-
20 tetramethylpiperidine N-oxyl (0.18 g), potassium bromide (10.62 g) and sodium acetate (14.19 g) in ethyl acetate (300 mL) and water (100 mL). After 20 min, triethylamine (3.3 mL) was added dropwise to the above mixture at the same temperature. The reaction mixture was extracted with ethyl acetate after 30 min. The organic layer was dried over anhydrous sodium sulphate. The solvent was evaporated under the reduced pressure to
25 afford the title compound (10.4 g).

Step e: Synthesis of 1,2:5,6-Di-O-isopropylidene- 3-deoxy-3-(2-ethoxy-2-oxoethylidene)- α -D-xylo-hexofuranose

To the solution of compound (10.5 g), obtained from step d above, in tetrahydrofuran, carboethoxymethylene triphenyl-phosphorane (27 g) was added. The
30 reaction mixture was refluxed for 2 hours and concentrated to afford crude compound,

which was purified by column chromatography over silica gel using 8% ethyl acetate in hexane as eluant to yield the title compound (5.0 g). Mass (m/z): 350.28 ($M^+ + Na$)

Step f: Synthesis of 1,2:5,6-Di-O-isopropylidene- 3-deoxy-3-(2-ethoxy-2-oxoethyl)- α - D-gulofuranose

5 10% Palladium on charcoal (3 g) was added to the solution of compound (12 g), obtained from step e above, in methanol (50 mL) and the reaction mixture was stirred under hydrogen atmosphere at room temperature for 4 hours. The reaction mixture was filtered through a celite pad and the residue was washed using ethyl acetate. The filtrate was concentrated to furnish the title compound (12 g). Mass (m/z): 330.9 ($M^+ + 1$).

10 **Step g: Synthesis of 1,2:5,6-Di-O-isopropylidene-3-deoxy-3-(2-hydroxyethyl)- α -D-gulofuranose**

To the suspension of lithium aluminium hydride (2.37 g) in tetrahydrofuran (120 mL) was added a solution of compound (12 g), obtained from step f above, in tetrahydrofuran (100 mL) at -50°C . The reaction mixture was allowed to attain the
15 temperature 0°C and mixture was stirred for 30 minutes at 0°C . The reaction mixture was quenched using aqueous solution of ammonium chloride (25 mL). The reaction mixture was slowly allowed to attain the room temperature and further stirred for 12 hours at same temperature. The reaction mixture was filtered through a celite pad, residue was washed with ethyl acetate. The organic layer was dried over anhydrous sodium sulphate. The
20 solvent was evaporated under the reduced pressure to afford the title compound (8.5 g).

Step h: Synthesis of 1,2:5,6-Di-O-isopropylidene- 3-deoxy-3-[2-(1,3-dioxo1,3-dihydro-2H-isoindol-2-yl)ethyl]- α -D-gulofuranose

A mixture of compound (8.5 g), obtained from step g above, triphenyl phosphine (21.83g) and phthalimide (6.73 g) were taken in a round bottom flask and dried in high
25 vacuum for 10 minutes. Then vacuum was released under a nitrogen atmosphere and tetrahydrofuran (100 mL) was added to the reaction mixture. The reaction mixture was cooled to 0°C and diisopropyl azodicarboxylate (12.625 g) was added slowly. The reaction mixture was stirred for 30 minutes at same temperature and then quenched with water and extracted with ethyl acetate. The organic layer was washed with water, brine
30 solution, dried over anhydrous sodium sulphate and concentrated. The residue thus

obtained was purified by column chromatography over silica gel using 40% ethyl acetate in hexane as eluant to afford the title compound (8.5 g).

Step i: Synthesis of 3-deoxy-3-[2-(1,3-dioxo-1,3-dihydro-2H-isoindol-2-yl)ethyl]-1,2-O-(isopropylidene)- α -D-gulofuranose

- 5 30% Perchloric acid (8.15 mL) was added to a solution of compound (0.8 g) obtained from step h above, in tetrahydrofuran (200 mL). The reaction mixture was stirred for 2 hours at 0-5°C and then quenched with saturated solution of sodium hydrogen carbonate. Ethyl acetate and water were added to the resulting mixture. The organic layer was separated, washed with brine, dried over anhydrous sodium sulphate and
- 10 concentrated. The residue thus obtained was purified by column chromatography over silica gel using 60% ethyl acetate in hexane to afford 5.5 g of the title compound. Mass (m/z): 377.24 ($M^+ + 1$)

Step j: Synthesis of (5R)-3-deoxy-3-[2-(1,3-dioxo-1,3-dihydro-2H-isoindol-2-yl)ethyl]-4,5-O-(1-methylethylidene)-L-arabino-pentodialdo-5,2-furanose

- 15 To a solution of compound (5.5 g), obtained from step i above, in acetone (100 mL) was added aqueous solution of sodium periodate (9.33 g in 100 mL) at 0°C. The reaction mixture was stirred for 2 hours, then filtered and concentrated. The residue thus obtained was taken in distilled water and extracted with ethyl acetate. The organic layer was dried over sodium sulphate and concentrated to afford the title compound (5.2 g).
- 20 **Step k: Synthesis of 2-(2-((3aR,5S,6R,6aR)-5-[(E)-2-(4-bromophenyl)ethenyl]-2,2-dimethyltetrahydrofuro[2,3-d][1,3]dioxol-6-yl)ethyl)-1H-isoindole-1,3(2H)-dione**

- A solution of (4-bromobenzyl)triphenylphosphonium bromide (13.3 g) in dimethyl sulphoxide (10 mL) was added dropwise to the suspension of sodium hydride (1 g of 50% suspension) in dimethyl sulphoxide (20 mL) at 0°C. After 20 minutes, a solution of
- 25 compound (6 g), obtained from step j above, in dimethylsulphoxide (100 mL) was added dropwise and reaction mixture was stirred for 1 hour at the same temperature. The reaction mixture was quenched with water and extracted with ethyl acetate. The organic layer was washed with water and brine solution and dried over anhydrous sodium sulphate. The solvent was evaporated under the reduced pressure to obtain a residue
- 30 which was purified by column chromatography over silica gel using 40% ethyl acetate in hexane to furnish the title compound (5.5 g). Mass (m/z): 499.10 ($M^+ + 1$)

Step l: Synthesis of 2-(2-((3aR,5S,6R,6aR)-5-[(E)-2-(3'-fluoro-4'-methylbiphenyl-4-yl)ethenyl]-2,2-dimethyltetrahydrofuro[2,3-d][1,3]dioxol-6-yl)ethyl)-1H-isoindole-1,3(2H)-dione

A mixture of compound (0.5 g), obtained from step k above, 3-fluoro-4-methylphenylboronic acid (0.314 g), tetrakis(triphenylphosphine)palladium (0) (0.057 g) and potassium carbonate (0.414 g) was dried under high vacuum for 10 minutes. The vacuum was released under nitrogen atmosphere and dry dimethylformamide (5 mL) was added at room temperature. The reaction mixture was heated at 120°C for 2 hours, and then quenched with water and extracted with ethyl acetate. The organic layer was washed with water and brine solution and dried over anhydrous sodium sulphate. The solvent was evaporated under the reduced pressure to obtain a residue which was purified by column chromatography over silica gel using 30% ethylacetate in hexane as eluant to afford the title compound (0.37 g). Mass (m/z): 527.23 ($M^+ + 1$)

Step m: Synthesis of 2-(2-((3aR,5S,6R,6aR)-5-[2-(3'-fluoro-4'-methylbiphenyl-4-yl)ethyl]-2,2-dimethyltetrahydrofuro[2,3-d][1,3]dioxol-6-yl)ethyl)-1H-isoindole-1,3(2H)-dione

10% Palladium on charcoal (100 g) was added to the solution of compound (0.37g), obtained from step l above, in ethyl acetate (10 mL) at room temperature and the reaction mixture was hydrogenated at the pressure of 35 psi for 1 hour in a Paar apparatus. The reaction mixture was filtered through celite pad and the residue was washed with methanol. The filtrate was concentrated to afford the title compound (0.28 g).

Step n: Synthesis of 2-(2-((2S,3S,4R,5S)-2-[2-(3'-fluoro-4'-methylbiphenyl-4-yl)ethyl]-4,5-dihydroxytetrahydrofuran-3-yl)ethyl)-1H-isoindole-1,3(2H)-dione

To a solution of compound (0.28 g), obtained from step m above, in acetonitrile (20 mL) and water (2 mL) was added 30% perchloric acid (0.4 mL) at room temperature. The reaction mixture was heated to 55°C for 30 minutes. The reaction mixture was then quenched using sodium bicarbonate solution. The solvents were evaporated at reduced pressure. The residue thus obtained was taken in ethyl acetate and water. The organic layer was separated and washed with water and brine solution and dried over anhydrous sodium sulphate. The solvent was concentrated under the reduced pressure to yield the

title compound (0.418 g) as crude mass which was used further without any purification and characterization.

Step o: Synthesis of (2R,3S)-2-[2-(1,3-dioxo-1,3-dihydro-2H-isoindol-2-yl)ethyl]-5-(3'-fluoro-4'-methylbiphenyl-4-yl)-3-(formyloxy)pentanoic acid

5 A solution of sodium metaperiodate (0.489 g in 5 mL of water) was added to a solution of compound (0.4 g), obtained from step n above, in a *tert*-butanol:tetrahydrofuran (5 mL: 5 mL) at 0°C. The reaction mixture was stirred for 2 hours at same temperature and potassium permanganate (0.033 g) was added at 0°C. After stirring, the reaction mixture for additional 6 hours at room temperature, the reaction
10 mixture was evaporated on rotary evaporator, the residue was taken into water and extracted with ethyl acetate. The organic layer was washed with water and brine solution and dried over anhydrous sodium sulphate. The solvent was evaporated under the reduced pressure, to afford the title compound (0.31 g) as crude mass which was used further without any purification and characterization.

15 **Step p: Synthesis of (2R,3S)-2-[2-(1,3-dioxo-1,3-dihydro-2H-isoindol-2-yl)ethyl]-5-(3'-fluoro-4'-methylbiphenyl-4-yl)-3-hydroxypentanoic acid**

 To a solution of compound (0.31g), obtained from step o above, in methanol (5 mL), potassium carbonate (0.094 g) was added at 0°C. The reaction mixture was stirred at room temperature for 3 hours. Reaction mixture was diluted with ethyl acetate (50 mL),
20 acidified with sodium hydrogen sulphate and washed with water (20 mL). The organic layer was dried over anhydrous sodium sulphate and concentrated under the reduced pressure obtain a residue which was purified by preparative thin layer chromatography using 10% methanol in dichloromethane as eluant to afford the title compound (0.020 g).
Mass (m/z): 476.07 ($M^+ + 1$)

25 ^1H NMR (CD_3OD): δ 7.72-7.64 (m, 4H), 7.36-7.34 (m, 2H), 7.18-7.10 (m, 5H), 3.68-3.61 (m, 3H), 2.75-2.71 (m, 1H), 2.55-2.49 (s, 1H), 2.32-2.30 (m, 1H), 2.16 (s, 3H), 1.99-1.94 (m, 2H), 1.70-1.62 (m, 2H).

Example 6: Synthesis of (2*S*,3*S*)-3-hydroxy-5-(4'-methoxybiphenyl-4-yl)-2-[2-(4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)ethyl]pentanoic acid (Compound No. 208)

Step a: Synthesis of 5-O-[tert-butyl(diphenyl)silyl]-3-deoxy-1,2-O-isopropylidene-3-[2-(4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)ethyl]- α -L-ribofuranose

- 5 A mixture of compound (20 g), obtained from step f of Example 1, triphenyl phosphine (17.6 g) and 1,2,3-benzotriazin-4(3*H*)-one (7.2 g) were dried in high vacuum in a round bottom flask for 10 minutes. The vacuum was released under a nitrogen atmosphere and tetrahydrofuran (200 mL) was added to the above reaction mixture. The reaction mixture was cooled to 0°C and diisopropyl azodicarboxylate (9.8 mL) was added slowly.
- 10 The reaction mixture was stirred for 30 minutes at same temperature, and then quenched with water and extracted with ethyl acetate. The organic layer was washed with water, brine solution, dried over anhydrous sodium sulphate and concentrated to obtain a residue which was purified by column chromatography over silica gel using 30% ethylacetate in hexane as eluant to afford the title compound (22 g). Mass (m/z): 586.24 ($M^+ + 1$).

15 **Step b: Synthesis of 3-deoxy-1,2-O-isopropylidene-3-[2-(4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)ethyl]- α -L-ribofuranose**

- To a solution of compound (22 g), obtained from step a above, in dry tetrahydrofuran (200 mL) at 0°C, was added 1 M solution of tetra-butylammonium fluoride (75 mL). The resulting mixture was initially stirred at the 0°C for 1 hour and then
- 20 at room temperature for 4 hours. The reaction mixture was cooled to 0°C, quenched with saturated ammonium chloride and extracted with ethyl acetate. The organic layer was washed with water and brine, dried over anhydrous sodium sulphate and concentrated. The residue thus obtained was purified by column chromatography over silica gel using 50% ethyl acetate in hexane as eluant to furnish the title compound (6.5 g). Mass (m/z):
- 25 348.25($M^+ + 1$)

Step c: Synthesis of (5*S*)-3-deoxy-4,5-O-isopropylidene-3-[2-(4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)ethyl]-D-ribo-pentodialdo-5,2-furanose

- To the solution of compound (6.5 g), obtained from step b above, in dry dichloromethane (100 mL), cooled to 0°C, was added Dess-Martin periodinane reagent
- 30 (11.1 g). The reaction mixture was allowed to stir for 2 hours. The reaction mixture was quenched with sodium thiosulphate and sodium hydrogen carbonate. Dichloromethane

was added to the reaction mixture. Organic layer was separated, washed with water and brine solution, dried over sodium sulphate and evaporated to afford the title compound. (6.5g)

Step d: Synthesis of 3-(2-((3aS,5S,6S,6aS)-5-[(E)-2-(4-bromophenyl)vinyl]-2,2-dimethyltetrahydrofuro[2,3-d][1,3]dioxol-6-yl)ethyl)-1,2,3-benzotriazin-4(3H)-one

To the suspension of sodium hydride (0.979 g, 60% in oil) in dimethyl sulphoxide (60 mL), cooled to 0°C, was added (4-bromobenzyl)triphenylphosphonium bromide (14.5 g). Compound (6.5 g), obtained from step c above, in tetrahydrofuran (60 mL) was added dropwise after 20 minutes, and reaction mixture was stirred for 1 hour at the same temperature. The reaction mixture was quenched with water and extracted with ethyl acetate. The organic layer was washed with water and brine solution and dried over anhydrous sodium sulphate. The solvent was evaporated under the reduced pressure and the residue thus obtained was purified by column chromatography using 40 % ethyl acetate in hexane as eluant to furnish the title compound (4 g). Mass (m/z): 498.19(M⁺ +1)

Step e: Synthesis of 3-(2-((3aS,5S,6S,6aS)-5-[(E)-2-(4'-methoxybiphenyl-4-yl)vinyl]-2,2-dimethyltetrahydrofuro[2,3-d][1,3]dioxol-6-yl)ethyl)-1,2,3-benzotriazin-4(3H)-one

A mixture of compound (1g), obtained from step d above, 4-methoxyphenyl boronic acid (0.61 g), tetrakis(triphenyl)phosphine palladium (0) (0.232 g) and potassium carbonate (0.832 g) was dried under high vacuum for 10 minutes and dry dimethylformamide (20 mL) was added at room temperature. The reaction mixture was heated at 120°C for 2 hours, and then quenched with water and extracted with ethyl acetate. The organic layer was washed with water and brine solution and dried over anhydrous sodium sulphate. The solvent was evaporated under the reduced pressure to obtain a residue which was purified by column chromatography over silica gel using 50% ethyl acetate in hexane as eluant to afford the title compound (0.62 g). Mass (m/z): 526.41(M⁺ +1)

Step f: Synthesis of 3-(2-((3aS,5S,6S,6aS)-5-[2-(4'-methoxybiphenyl-4-yl)ethyl]-2,2-dimethyltetrahydrofuro[2,3-d][1,3]dioxol-6-yl)ethyl)-1,2,3-benzotriazin-4(3H)-one

To the solution of compound (0.62 g), obtained from step e above, in a solvent mixture of ethyl acetate (20 mL) was added 10% Palladium on charcoal (0.3 g) at room temperature and the reaction mixture was hydrogenated at the pressure of 35 psi for 4 hours in a Paar apparatus. The reaction mixture was filtered through a celite pad and the residue was washed with methanol. The filtrate was concentrated to afford the title compound (0.6 g).

Step g: Synthesis 3-(2-((2S,3R,4S,5R)-4,5-dihydroxy-2-[2-(4'-methoxybiphenyl-4-yl)ethyl]tetrahydrofuran-3-yl)ethyl)-1,2,3-benzotriazin-4(3H)-one

Perchloric acid (0.2 mL) was added to a solution of compound (0.6 g), obtained from step f above, in acetonitrile (50 mL) and water (10 mL) at room temperature. The reaction mixture was heated to 55°C for 30 minutes. The reaction mixture was then quenched using sodium hydrogen carbonate solution. The solvents were evaporated at reduced pressure. The residue thus obtained was taken in ethyl acetate and water. The organic layer was separated and washed with water and brine solution and dried over anhydrous sodium sulphate. The solvent was concentrated under the reduced pressure to yield the title compound (0.6 g).

Step h: Synthesis of (2S,3S)-3-(formyloxy)-2-[2-(4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-5-(4-pyrimidin-5-ylphenyl)pentanoic acid

To a solution of compound (0.6 g), obtained from step g above, in a *tert*-butanol:tetrahydrofuran (7 mL: 7 mL) at 0°C was added a solution of sodium metaperiodate (0.973 g in 7 mL of water). The reaction mixture was stirred for 2 hours at same temperature and potassium permanganate (0.036 g) was added at 0°C. After stirring the reaction mixture for an additional 6 hours at room temperature, reaction mixture was evaporated, the residue was taken into water and extracted with ethyl acetate. The organic layer was washed with water and brine solution and dried over anhydrous sodium sulphate. The solvent was evaporated under the reduced pressure, to afford the title compound (0.5 g).

Step i: Synthesis of (2S,3S)-3-hydroxy-5-(4'-methoxybiphenyl-4-yl)-2-[2-(4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]pentanoic acid

Potassium carbonate (0.151 g) was added to a solution of compound (0.5 g), obtained from step h above, in methanol (5 mL) and tetrahydrofuran (5 mL) at 0°C. The

reaction mixture was stirred at room temperature for 3h. Solvents were evaporated and the residue was taken into water and ethyl acetate. Organic layer was washed with water and brine solution and dried over anhydrous sodium sulphate. Solvent was evaporated under the reduced pressure obtain a residue which was purified by preparative thin layer chromatography using 10% methanol in dichloromethane as eluent to afford the title compound (0.008g). Mass (m/z): 474.31 ($M^+ + 1$); ^1H NMR (CD_3OD): δ 8.32-7.87 (m, 5H), 7.50-7.42 (m, 4H), 7.24-7.20 (m, 2H), 6.97-6.94 (m, 1H), 4.60-4.50 (m, 2H), 3.81 (s, 3H), 3.59 (br t, 1H), 3.0-1.8 (m, 7H).

Example 7: Synthesis of (2*S*,3*R*)-3-hydroxy-5-[4-(5-methylpyridin-2-yl)phenyl]-2-[2-(4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)ethyl]pentanoic acid (Compound No. 209)

Step a: Synthesis of 4-(5-methylpyridin-2-yl)benzaldehyde

A mixture of 2-bromo-5-methylpyridine (2g), (4-formylphenyl)boronic acid (3.5 g), tetrakis(triphenylphosphine)palladium (0) (0.672 g) and potassium carbonate (4.8 g) was dried under high vacuum for 10 minutes and dry dimethylformamide (15 mL) was added at room temperature. The reaction mixture was heated at 110°C for 6 hours, and then quenched with water and extracted with ethyl acetate. The organic layer was washed with water and brine solution and dried over anhydrous sodium sulphate. The solvent was evaporated under the reduced pressure to obtain a residue which was purified by column chromatography over silica gel using 20% ethylacetate in hexane as eluant to afford the title compound (2.8 g). Mass (m/z): 198.20

Step b: Synthesis of ethyl (2*E*)-3-[4-(5-methylpyridin-2-yl)phenyl]prop-2-enoate

To the suspension of sodium hydride (0.682 g, 60% in oil.) in tetrahydrofuran (20 mL) at 0°C was charged triethyl phosphonoacetate (3.82 g). After 15 minutes, a solution of compound (2.8 g), obtained from step a above, in tetrahydrofuran (5mL) was added dropwise and the reaction mixture was stirred for 30 minutes at room temperature. A saturated solution of ammonium chloride in water was added to reaction mixture. The solvent was evaporated and the resulting mixture was extracted with ethyl acetate. The combined extracts were dried over anhydrous sodium sulphate and evaporated under reduced pressure to yield a residue which was purified by column chromatography over silica gel using 15% ethylacetate in hexane as eluant to afford the title compound (1.8 g).

Step c: Synthesis of ethyl 3-[4-(5-methylpyridin-2-yl)phenyl]propanoate

10% Palladium on charcoal (0.8 g) was added to the solution of compound (1.8 g), obtained from step b above, in tetrahydrofuran (15 mL) at room temperature and hydrogen was supplied at the pressure of 50 psi (Paar apparatus) for 4 hours. The reaction mixture
5 was filtered through celite pad and concentrated to afford the title compound (1.8 g).

Step d: Synthesis of 3-[4-(5-methylpyridin-2-yl)phenyl]propan-1-ol

To the suspension of lithium aluminium hydride (0.424 g) in tetrahydrofuran (20 mL) was added a solution of compound (2 g), obtained from step c above, in tetrahydrofuran (10 mL) at -20 °C. The reaction mixture was stirred for 2 hours at -30 °C
10 and saturated solution of sodium sulphate was added at the same temperature. The reaction mixture was then filtered through a celite pad and the residue was washed with ethyl acetate. The organic layer was dried over anhydrous sodium sulphate, filtered, concentrated under reduced pressure and the residue thus obtained was purified by column chromatography over silica gel using 50% ethylacetate in hexane as eluant to afford the
15 title compound (1.2 g).

Step e: Synthesis of 3-[4-(5-methylpyridin-2-yl)phenyl]propanal

To a stirred solution of compound (1.4g), obtained from step d above, in dichloromethane (20 mL), 2,2,6,6,-tetramethylpiperidine N-oxyl (9.6 mg), and potassium bromide (73.4 mg) was added at 0 °C under nitrogen atmosphere. Sodium hypochlorite
20 (13.7 mL, 4% solution) was added at once at pH 8-9 (maintained by adding aqueous sodium bicarbonate solution). The reaction was stirred for 20 min at 0 °C. The organic layer was separated and the aqueous layer was extracted with dichloromethane. The combined organic layers were washed with saturated aqueous sodium bicarbonate solution, water and brine. The organic layer was dried over anhydrous sodium sulphate,
25 filtered, concentrated under reduced pressure to afford crude aldehyde (0.9 g).

Step f: Synthesis of 3-[(3S,4R)-3-[(4S)-4-benzyl-2-oxo-1,3-thiazolidin-3-yl]carbonyl]-4-hydroxy-6-[4-(5-methylpyridin-2-yl)phenyl]hexyl]-1,2,3-benzotriazin-4(3H)-one

In a flame-dried flask, 3-{4-[(4S)-4-benzyl-2-oxo-1,3-thiazolidin-3-yl]-4-oxobutyl}-1,2,3-benzotriazin-4(3H)-one (1.18 g) was taken up in dichloromethane (10
30 mL) and cooled to 0 °C. Titanium tetrachloride (3.4 mL) in dichloromethane (6 mL) was added dropwise and reaction mixture was stirred for 10-15 minutes. (-)-Sparteine (1.7 g)

was added slowly to the reaction mixture and stirred at 0 °C for 45 minutes. A solution of compound (0.65 g), obtained from step e above, in dichloromethane (10 mL) was added slowly and stirring was continued at 0 °C. After 3 hours, the reaction was quenched with dropwise addition of saturated ammonium chloride solution and dichloromethane was added. The organic layer was separated and the aqueous layer was extracted with dichloromethane. The organic layer was washed with brine, and dried over anhydrous sodium sulphate, and concentrated under reduced pressure. The crude product thus obtained was purified by silica gel flash column chromatography using 30% ethyl acetate in hexanes as eluant to afford the aldol adduct (0.56 g).

Step g: Synthesis of (2S,3R)-3-hydroxy-5-[4-(5-methylpyridin-2-yl)phenyl]-2-[2-(4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]pentanoic acid

To a stirred solution of the compound (0.2 g), obtained from step f above, in tetrahydrofuran: water (3:1, 10 mL) at 0 °C was added aqueous hydrogen peroxide solution (30%, 0.16 mL), followed by addition of lithium hydroxide monohydrate (0.02g) in water (5 mL). The reaction mixture was stirred at 0 °C for 30 minutes. The reaction mixture was concentrated and the residue was extracted with ethyl acetate. The aqueous layer was acidified with sodium hydrogen sulphate and extracted with ethyl acetate. The combined layers were washed with water, brine and dried over anhydrous sodium sulphate. The solvents were evaporated under reduced pressure and the crude residue was purified by silica gel flash column chromatography using 3% methanol in dichloromethane as eluant to afford the title compound (0.035 g). Mass (m/z): 459.21(M⁺ +1); ¹HNMR (CD₃OD): δ 8.40 (s, 1H), 8.30 (d, 1H, J = 7.88 Hz), 8.15 (d, 1H, J = 8Hz), 8.05-7.70 (m, 6H), 7.27 (d, 2H, J = 7.6 Hz), 4.40-4.0 (m, 2H), 3.80-3.60 (m, 1H), 2.80-2.40 (m, 5H), 2.38(s, 3H), 1.80-1.60 (m, 2H).

Example 7A: Synthesis of (2S,3R)-3-hydroxy-5-[4-(6-methoxypyridin-3-yl)phenyl]-2-[2-(4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 86)

Step a: Synthesis of methyl (2E)-3-(4-bromophenyl)prop-2-enoate

To a stirred solution of 4-bromocinnamic acid (16g) in methanol (150 mL) was added thionyl chloride (30 mL) at 0 °C. The reaction mixture was warmed to room temperature and refluxed for 3 hours. After cooling to room temperature the solvents were evaporated to dryness. The crude compound obtained was used as such for the next step.

Step b: Synthesis of methyl (2E)-3-[4-(6-methoxypyridin-3-yl)phenyl]prop-2-enoate

Compound (7.5 g) obtained from step a above, pyridine-2-methoxy-5-boronic acid (9.48 g), tetrakis(triphenylphosphine)palladium (0) (1.79 g) and potassium carbonate (12.83 g) were taken in dimethylformamide (60 mL) under nitrogen atmosphere. The reaction mixture was refluxed under nitrogen atmosphere for 6 hours. After cooling to room temperature, water was added and the reaction mixture was extracted with ethyl acetate. The combined organic layers were washed with water and brine, dried over anhydrous sodium sulfate and the solvents were evaporated under reduced pressure. The crude product was purified by silica gel flash column chromatography using 20-30 % ethyl acetate in hexane to get the title compound (8.09 g).

Step c: Synthesis of methyl 3-[4-(6-methoxypyridin-3-yl)phenyl]propanoate

To the solution of compound (8.1 g), obtained from step b above, in a mixture of ethyl acetate /methanol /tetrahydrofuran (~300 mL), 10% Palladium on charcoal (1.6 g) was added and stirred under hydrogen atmosphere at the pressure of 30 psi (Paar apparatus) for 2.5 hours. The reaction mixture was filtered through celite pad washed with ethyl acetate and the filtrate was concentrated under reduced pressure to obtain the title compound as white solid which was used as such for the next step.

Step d: Synthesis of 3-[4-(6-methoxypyridin-3-yl)phenyl]propanoic acid

To a stirred solution of lithium aluminium hydride (2.28 g) in tetrahydrofuran (120 mL) was added the compound (8g), obtained from step c above in tetrahydrofuran (50 mL) at 0°C. The reaction mixture was stirred at 0°C for 3 hours. The reaction mixture was quenched carefully with saturated ammonium chloride solution, filtered through celite, washed with ethyl acetate. The organic layers were separated, washed with brine and dried over anhydrous sodium sulfate and solvents were evaporated under reduced pressure. The crude product was purified by silica gel flash column chromatography using 20-30 % ethyl acetate in hexane to get the title compound as a white crystalline solid (5.1 g).

Step e: Synthesis of 3-[4-(6-methoxypyridin-3-yl)phenyl]propanal

To a stirred solution of compound (9.0 g) obtained from step d above, in dichloromethane (90 ml), 2,2,6,6,-tetramethylpiperidine N-oxyl (58.77 mg) and potassium bromide (447.57 mg) was added at 0°C under nitrogen atmosphere. A 4% aqueous sodium hypochlorite (3.497 g) was added at once at pH 8-9 (maintained by adding

saturated aqueous sodium bicarbonate solution). The reaction was stirred for 20 minutes at 0°C. Then, the organic layer was separated and the aqueous layer was extracted with dichloromethane. The combined organic layers were washed with saturated aqueous sodium bicarbonate solution, water and brine. Finally, the organic layers were dried over anhydrous sodium sulfate and concentrated under reduced pressure to afford crude aldehyde (8.1g) which was used as such in the next step.

Step f: Synthesis of 3-{(3*S*,4*S*)-3-[(4*S*)-4-benzyl-2-oxo-1,3-thiazolidin-3-yl]carbonyl}-4-hydroxy-6-[4-(5-methoxypyridin-2-yl)phenyl]hexyl}-1,2,3-benzotriazin-4(3*H*)-one

In a flame-dried flask, 3-{4-[(4*S*)-4-benzyl-2-oxo-1,3-thiazolidin-3-yl]-4-oxobutyl}-1,2,3-benzotriazin-4(3*H*)-one (2.0 g) was taken up in dichloromethane (10 mL) and cooled to 0°C. Titanium tetrachloride (0.619 mL) in dichloromethane (6 mL) was added dropwise and reaction mixture was stirred for 10-15 minutes. (-)-Sparteine (2.7 mL) was added slowly to the reaction mixture and stirred at 0°C for 45 minutes. A solution of compound (1.34 g), obtained from step e above, in dichloromethane (10 mL) was added slowly and stirring was continued at 0°C. After 3 hours, the reaction was quenched with dropwise addition of saturated ammonium chloride solution and dichloromethane was added. The organic layer was separated and the aqueous layer was extracted with dichloromethane. The organic layer was washed with brine, and dried over anhydrous sodium sulphate, and concentrated under reduced pressure. The crude product thus obtained was purified by silica gel flash column chromatography using 30% ethyl acetate in hexanes as eluants to afford the aldol adduct (1.21g).

Step g: Synthesis of (2*S*,3*R*)-3-hydroxy-5-[4-(5-methoxypyridin-2-yl)phenyl]-2-[2-(4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)ethyl]pentanoic acid

To a stirred solution of the compound (1.21 g), obtained from step f above, in tetrahydrofuran: water (3:1, 15 mL) at 0°C was added aqueous hydrogen peroxide solution (30%, 1.02 mL), followed by addition of lithium hydroxide monohydrate (114.5mg) in water (5 mL). The reaction mixture was stirred at 0°C until the completion of the hydrolysis. The reaction mixture was concentrated and the residue was extracted with ethyl acetate. The aqueous layer was acidified with sodium hydrogen sulphate and extracted with ethyl acetate. The combined layers were washed with water, brine and dried over anhydrous sodium sulphate. The solvents were evaporated under reduced

pressure and the crude residue was purified by silica gel flash column chromatography using 3% methanol in dichloromethane as eluant to afford the title compound (0.61 g).

Mass (m/z): 474.87(M⁺ +1); ¹H NMR (400 MHz, MeOD): δ 8.32- 8.30 (2H, m), 8.16- 8.14 (1H, m), 8.04- 8.03 (1H, m), 7.92- 7.88 (2H, m), 7.46 (2H, d, J = 8Hz), 7.24 (2H, m, J = 8Hz), 6.86 (1H, d, J = 8Hz), 4.57- 4.53 (2H, m), 3.93 (3H, s), 3.78-3.83 (1H, m), 2.83- 2.80 (1H, m), 2.63- 2.60 (1H, m), 2.52- 2.49 (1H, m), 2.32- 2.28 (2H, m), 1.80- 1.76 (2H, m).

Example 7B: Synthesis of (2*S*,3*R*)-5-(2',4'-difluorobiphenyl-4-yl)-3-hydroxy-2-[(4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)methyl]pentanoic acid (Compound No. 216)

10 **Step a: Synthesis of methyl (2*E*)-3-(4-bromophenyl)prop-2-enoate**

To a stirred solution of 4-bromocinnamic acid (16g) in methanol (150 mL) was added thionyl chloride (30 mL) at 0°C. The reaction mixture was warmed to room temperature and refluxed for 3 hours. After cooling to room temperature the solvents were evaporated to dryness. The crude compound obtained was used as such for the next step.

15 **Step b: Synthesis of methyl (2*E*)-3-(3',5'-difluorobiphenyl-4-yl)prop-2-enoate**

Compound (0.75 g) obtained from step a above, 2,4difluorophenyl boronic acid (0.737 g), tetrakis(triphenyl)phosphinepalladium (0) (0.07 g) and potassium carbonate (1.28 g) were taken in dimethylformamide (6 mL) under nitrogen atmosphere. The reaction mixture was refluxed under nitrogen atmosphere for 6 hours. After cooling to room temperature, water was added and the reaction mixture was extracted with ethyl acetate. The combined organic layers were washed with water and brine, dried over anhydrous sodium sulfate and the solvents were evaporated under reduced pressure. The crude product was purified by silica gel flash column chromatography using 20-30 % ethyl acetate in hexane to get the title compound (0.810 g).

25 **Step c: Synthesis of methyl 3-(3',5'-difluorobiphenyl-4-yl)propanoate**

To the solution of compound (0.79 g), obtained from step b above, in a mixture of ethyl acetate (~10 mL), 10% Palladium on charcoal (0.2g) was added and stirred under hydrogen atmosphere at the pressure of 30 psi for 2.5 hours. The reaction mixture was filtered through celite pad washed with ethyl acetate and the filtrate was concentrated under reduced pressure to obtain the title compound (0.77g) as white solid which was used as such for the next step.

Step d: Synthesis of 3-(3',5'-difluorobiphenyl-4-yl)propanoic acid

To a stirred solution of lithium aluminium hydride (0.2 g) in tetrahydrofuran (15 mL) was added the compound (0.77g), obtained from step c above in tetrahydrofuran (50 mL) at 0 °C. The reaction mixture was stirred at 0 °C for 3 hours. The reaction mixture
5 was quenched carefully with saturated ammonium chloride solution, filtered through celite, washed with ethyl acetate. The organic layers were separated, washed with brine and dried over anhydrous sodium sulfate and solvents were evaporated under reduced pressure. The crude product was purified by silica gel flash column chromatography using 20-30 % ethyl acetate in hexane to get the title compound as a white crystalline solid
10 (0.70 g).

Step e: Synthesis of 3-(3',5'-difluorobiphenyl-4-yl)propanal

To a stirred solution of compound (0.7g), obtained from step d above, in dichloromethane (10 mL), 2,2,6,6-tetramethylpiperidine N-oxyl (4.36 mg), and potassium bromide (33.32 mg) was added at 0 °C under nitrogen atmosphere. Sodium
15 hypochlorite (6.5mL, 4% solution) was added at once at pH 8-9 (maintained by adding aqueous sodium bicarbonate solution). The reaction was stirred for 20 min at 0 °C. The organic layer was separated and the aqueous layer was extracted with dichloromethane. The combined organic layers were washed with saturated aqueous sodium bicarbonate solution, water and brine. The organic layer was dried over anhydrous sodium sulphate,
20 filtered, concentrated under reduced pressure to afford crude aldehyde (0.57 g).

Step f: Synthesis of 3-[(2*R*,3*S*)-2-[(4*S*)-4-benzyl-2-oxo-1,3-thiazolidin-3-yl]carbonyl]-5-(3',5'-difluorobiphenyl-4-yl)-3-hydroxypentyl]-1,2,3-benzotriazin-4(3*H*)-one

In a flame-dried flask, 3-{3-[(4*S*)-4-benzyl-2-oxo-1,3-thiazolidin-3-yl]-3-oxopropyl}-1,2,3-benzotriazin-4(3*H*)-one (0.25 g) was taken up in dichloromethane (10
25 mL) and cooled to 0 °C. Titanium tetrachloride (0.76 mL, 1M solution) was added dropwise and reaction mixture was stirred for 10-15 minutes. (-)-Sparteine (0.36 mL) was added slowly to the reaction mixture and stirred at 0 °C for 20 minutes. A solution of compound (0.187 g), obtained from step e above, in dichloromethane (10 mL) was added slowly and stirring was continued at 0 °C. After 3 hours, the reaction was quenched with
30 dropwise addition of saturated ammonium chloride solution and dichloromethane was added. The organic layer was separated and the aqueous layer was extracted with

dichloromethane. The organic layer was washed with brine, and dried over anhydrous sodium sulphate, and concentrated under reduced pressure. The crude product thus obtained was purified by silica gel flash column chromatography using 30% ethyl acetate in hexane as eluants to afford the aldol adduct (0.24g).

5 Step g: Synthesis of (2*S*,3*R*)-5-(3',5'-difluorobiphenyl-4-yl)-3-hydroxy-2-[(4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)methyl]pentanoic acid

To a stirred solution of the compound (0.2g), obtained from step f above, in tetrahydrofuran: water (3:1, 10 mL) at 0 °C was added aqueous hydrogen peroxide solution (30%, 0.16 mL), followed by addition of lithium hydroxide monohydrate (19.53mg) in
10 water (2 mL). The reaction mixture was stirred at 0 °C until the completion of the hydrolysis. The reaction mixture was concentrated and the residue was extracted with ethyl acetate. The aqueous layer was acidified with sodium hydrogen sulphate and extracted with ethyl acetate. The combined layers were washed with water, brine and dried over anhydrous sodium sulphate. The solvents were evaporated under reduced
15 pressure and the crude residue was purified by silica gel flash column chromatography using 10% methanol in dichloromethane as eluant to afford the title compound (0.090g).
Mass (m/z): 466.06 ($M^+ + 1$); ¹HNMR (CD₃OD): δ 8.17-8.8.15 (m, 1H), 8.05-8.01(m, 1H), 7.90-7.88 (m, 1H), 7.86-7.84 (m, 1H), 7.46-7.42 (m, 2H), 7.40-7.29 (m, 2H), 7.03-6.98 (m, 2H), 4.82-4.84 (m, 2H), 3.94 (s, 1H), 3.22 – 3.21 (m, 1H), 2.91-2.89 (m, 1H), 2.78-
20 2.75 (m, 1H), 1.94-1.90 (m, 2H).

Example 7C: Synthesis of (2*S*,3*R*)-5-(4'-fluorobiphenyl-4-yl)-3-hydroxy-2-[(4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)methyl]pentanoic acid (Compound No. 217)

Step a: Synthesis of methyl (2*E*)-3-(4-bromophenyl)prop-2-enoate

To a stirred solution of 4-bromocinnamic acid (16g) in methanol (150 mL) was
25 added thionyl chloride (30 mL) at 0 °C. The reaction mixture was warmed to room temperature and refluxed for 3 hours. After cooling to room temperature the solvents were evaporated to dryness. The crude compound obtained was used as such for the next step.

Step b: Synthesis of methyl (2*E*)-3-(4'-fluorobiphenyl-4-yl)prop-2-enoate

Compound (0.75 g) obtained from step a above, 4-fluorophenyl boronic acid (0.65
30 g), tetrakis(triphenylphosphine)palladium (0) (0.07 g) and potassium carbonate (1.28 g) were taken in dimethylformamide (6 mL) under nitrogen atmosphere. The reaction

mixture was refluxed under nitrogen atmosphere for 6 hours. After cooling to room temperature, water was added and the reaction mixture was extracted with ethyl acetate. The combined organic layers were washed with water and brine, dried over anhydrous sodium sulfate and the solvents were evaporated under reduced pressure. The crude
5 product was purified by silica gel flash column chromatography using 20-30 % ethyl acetate in hexane to get the title compound (0.720 g).

Step c: Synthesis of methyl 3-(4'-fluorobiphenyl-4-yl) propanoate

To the solution of compound (0.7 g), obtained from step b above, in a mixture of ethyl acetate (~20 mL), 10% Palladium on charcoal (0.2 g) was added and stirred under
10 hydrogen atmosphere at the pressure of 30 psi (Paar apparatus) for 2.5 hours. The reaction mixture was filtered through celite pad washed with ethyl acetate and the filtrate was concentrated under reduced pressure to obtain the title compound (0.7g) as white solid which was used as such for the next step.

Step d: Synthesis of 3-(4'-fluorobiphenyl-4-yl)propanoic acid

To a stirred solution of lithium aluminium hydride (0.2 g) in tetrahydrofuran (15 mL) was added the compound (0.7g), obtained from step c above in tetrahydrofuran (50 mL) at 0°C. The reaction mixture was stirred at 0°C for 3 hours. The reaction mixture was quenched carefully with saturated ammonium chloride solution, filtered through celite, washed with ethyl acetate. The organic layers were separated, washed with brine
20 and dried over anhydrous sodium sulfate and solvents were evaporated under reduced pressure. The crude product was purified by silica gel flash column chromatography using 20-30 % ethyl acetate in hexane to get the title compound as a white crystalline solid (0.70 g).

Step e: Synthesis of 3-(4'-fluorobiphenyl-4-yl) propanal

To a stirred solution of compound (0.7g), obtained from step d above, in dichloromethane (10 mL), 2,2,6,6-tetramethylpiperidine N-oxyl (4.74 mg), and potassium bromide (36.05 mg) was added at 0°C under nitrogen atmosphere. Sodium hypochlorite (0.28g, 4% solution) was added at once at pH 8-9 (maintained by adding aqueous sodium bicarbonate solution). The reaction was stirred for 20 min at 0°C. The
30 organic layer was separated and the aqueous layer was extracted with dichloromethane. The combined organic layers were washed with saturated aqueous sodium bicarbonate

solution, water and brine. The organic layer was dried over anhydrous sodium sulphate, filtered, concentrated under reduced pressure to afford crude aldehyde (0.55 g).

Step f: Synthesis of 3-[(2*R*,3*S*)-2-[(4*S*)-4-benzyl-2-oxo-1,3-thiazolidin-3-yl]carbonyl]-5-(4'-fluorobiphenyl-4-yl)-3-hydroxypentyl]-1,2,3-benzotriazin-4(3*H*)-one

5 In a flame-dried flask, 3-{3-[(4*S*)-4-benzyl-2-oxo-1,3-thiazolidin-3-yl]-3-oxopropyl}-1,2,3-benzotriazin-4(3*H*)-one (0.25 g) was taken up in dichloromethane (10 mL) and cooled to 0 °C. Titanium tetrachloride (0.76 mL, 1M solution) was added dropwise and reaction mixture was stirred for 10-15 minutes. (-)-Sparteine (0.36 mL) was added slowly to the reaction mixture and stirred at 0 °C for 20 minutes. A solution of
10 compound (0.17 g), obtained from step e above, in dichloromethane (10 mL) was added slowly and stirring was continued at 0 °C. After 3 hours, the reaction was quenched with dropwise addition of saturated ammonium chloride solution and dichloromethane was added. The organic layer was separated and the aqueous layer was extracted with dichloromethane. The organic layer was washed with brine, and dried over anhydrous
15 sodium sulphate, and concentrated under reduced pressure. The crude product thus obtained was purified by silica gel flash column chromatography using 30% ethyl acetate in hexanes as eluants to afford the aldol adduct (0.2g).

Step g: Synthesis of (2*S*,3*R*)-5-(4'-fluorobiphenyl-4-yl)-3-hydroxy-2-[(4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)methyl]pentanoic acid

20 To a stirred solution of the compound (0.18 g), obtained from step f above, in tetrahydrofuran: water (3:1, 10 mL) at 0 °C was added aqueous hydrogen peroxide solution (30%, 0.15 mL), followed by addition of lithium hydroxide monohydrate (17.6mg) in water (2 mL). The reaction mixture was stirred at 0 °C until the completion of the hydrolysis. The reaction mixture was concentrated and the residue was extracted with
25 ethyl acetate. The aqueous layer was acidified with sodium hydrogen sulphate and extracted with ethyl acetate. The combined layers were washed with water, brine and dried over anhydrous sodium sulphate. The solvents were evaporated under reduced pressure and the crude residue was purified by silica gel flash column chromatography using 10% methanol in dichloromethane as eluant to afford the title compound (0.075 g).
30 Mass (m/z): 448.07 (M⁺ + 1); ¹HNMR (CD₃OD): δ 8.25-8.8.30 (d, 1H, J = 8Hz), 8.15-8.12 (d, 1H, J = 8Hz), 8.05-8.00 (t, 1H, J = 8Hz), 7.89-7.85 (t, 1H, J = 8Hz), 7.58-7.55 (m,

1H), 7.47-7.45 (d, 1H $J = 8\text{Hz}$), 7.28- 7.26 (d, 1H, $J = 8\text{Hz}$), 7.89-7.87 (m, 1H), 7.61 (m, 2H), 7.53- 7.59 (m, 1H), 7.30-7.25 (d, 2H $J = 8\text{Hz}$), 7.15- 7.11 (t, 2H, $J = 8\text{Hz}$), 4.81-4.79 (m, 2H), 3.92 (s, 1H), 3.22-3.16 (m, 1H), 2.92- 2.90 (m, 1H), 2.75- 2.73 (m, 1H), 1.93- 1.87 (m, 2H).

5 The following compounds were prepared employing procedures as provided in Examples 1 to 7C described above:

Compound Nos. 1 to 81, 86-88, 90-95, 97-119, 121-142, 144-203, 211-213 and 226-232 were prepared following Example 1; compound Nos. 82-83 were prepared following Example 2; compound Nos 84-85, and 204 were prepared following Example 3;
10 compound Nos. 89 and 96 were prepared following Example 4; compound Nos 120, and 205-207 were prepared following Example 5; compound Nos. 143 and 208 were prepared following Example 6; and compound Nos 86, 209-210 and 214-225 were prepared following Example 7.

Specific compounds, suitable for use, prepared in the present invention, are listed
15 below:

(2S,3R)-3-hydroxy-2-[2-(4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-5-(4-pyrimidin-5-ylphenyl)pentanoic acid (Compound No.1), Mass (m/z):446.0 ($M^+ + 1$);

(2S,3R)-2-[2-(5-tert-butyl-1,3-dioxo-1,3-dihydro-2H-isoindol-2-yl)ethyl]-3-hydroxy-5-[4-(6-methoxypyridin-3-yl)phenyl]pentanoic acid (Compound No. 2), Mass
20 (m/z): 531.09 ($M^+ + 1$);

(2S,3R)-2-[2-(1,3-dioxo-1,3-dihydro-2H-isoindol-2-yl)ethyl]-3-hydroxy-5-[4-(6-methylpyridin-3-yl)phenyl]pentanoic acid (Compound No. 3), Mass (m/z): 458.82($M^+ + 1$);

(2S,3R)-2-[2-(1,3-dioxo-1,3-dihydro-2H-isoindol-2-yl)ethyl]-5-(4'-fluorobiphenyl-4-yl)-3-hydroxypentanoic acid (Compound No. 4),
25

(2S,3R)-5-(3',4'-difluorobiphenyl-4-yl)-2-[2-(1,3-dioxo-1,3-dihydro-2H-isoindol-2-yl)ethyl]-3-hydroxypentanoic acid (Compound No. 5), Mass (m/z): 479.70($M^+ + 1$);

(2S,3R)-3-hydroxy-5-[4-(6-methylpyridin-3-yl)phenyl]-2-[2-(4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 6), Mass (m/z):458.94 ($M^+ + 1$);
30

(2S,3R)-3-hydroxy-2-[2-(5-methyl-1,3-dioxo-1,3-dihydro-2H-isoindol-2-yl)ethyl]-5-[4-(6-methylpyridin-3-yl)phenyl]pentanoic acid (Compound No. 7), Mass (m/z): 472.87 ($M^+ + 1$);

5 (2S,3R)-5-(4'-fluorobiphenyl-4-yl)-3-hydroxy-2-[2-(5-methyl-1,3-dioxo-1,3-dihydro-2H-isoindol-2-yl)ethyl]pentanoic acid (Compound No. 8),

(2S,3R)-5-(4'-fluorobiphenyl-4-yl)-3-hydroxy-2-[2-(4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 9), Mass (m/z): 461.87 ($M^+ + 1$);

(2S,3R)-3-hydroxy-5-(4'-methoxybiphenyl-4-yl)-2-[2-(4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 10), Mass (m/z): 473.79 ($M^+ + 1$);

10 (2S,3R)-5-(3',4'-difluorobiphenyl-4-yl)-3-hydroxy-2-[2-(5-methyl-1,3-dioxo-1,3-dihydro-2H-isoindol-2-yl)ethyl]pentanoic acid (Compound No. 11), Mass (m/z): 493.81 ($M^+ + 1$);

(2S,3R)-5-(3',4'-difluorobiphenyl-4-yl)-3-hydroxy-2-[2-(4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 12), Mass (m/z): 479.77 ($M^+ + 1$);

15 (2S,3R)-2-[2-(1,3-dioxo-1,3-dihydro-2H-isoindol-2-yl)ethyl]-3-hydroxy-5-(4'-methoxybiphenyl-4-yl)pentanoic acid (Compound No. 13), Mass (m/z): 473.86 ($M^+ + 1$);

(2S,3R)-2-[2-(1,3-dioxo-1,3-dihydro-2H-isoindol-2-yl)ethyl]-3-hydroxy-5-[4'-(trifluoromethoxy)biphenyl-4-yl]pentanoic acid (Compound No. 14), Mass (m/z): 510.66 ($M^+ - 18$);

20 (2S,3R)-5-(4'-chloro-3'-fluorobiphenyl-4-yl)-3-hydroxy-2-[2-(4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 15), Mass (m/z): 496.10 ($M^+ + 1$);

(2S,3R)-2-[2-(5-tert-butyl-1,3-dioxo-1,3-dihydro-2H-isoindol-2-yl)ethyl]-5-(3',4'-difluorobiphenyl-4-yl)-3-hydroxypentanoic acid (Compound No. 16), Mass (m/z): 536.11
25 ($M^+ + 1$);

(2S,3R)-2-[2-(5-tert-butyl-1,3-dioxo-1,3-dihydro-2H-isoindol-2-yl)ethyl]-3-hydroxy-5-(4'-methoxybiphenyl-4-yl)pentanoic acid (Compound No. 17), Mass (m/z): 530.09 ($M^+ + 1$);

(2S,3R)-3-hydroxy-2-[2-(4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-5-[4'-(trifluoromethyl)biphenyl-4-yl]pentanoic acid (Compound No. 18), Mass (m/z): 512.04($M^+ + 1$);

5 (2S,3R)-2-[2-(5-tert-butyl-1,3-dioxo-1,3-dihydro-2H-isoindol-2-yl)ethyl]-5-(4'-fluorobiphenyl-4-yl)-3-hydroxypentanoic acid (Compound No. 19), Mass (m/z): 518.14 ($M^+ + 1$);

(2S,3R)-2-[2-(5-tert-butyl-1,3-dioxo-1,3-dihydro-2H-isoindol-2-yl)ethyl]-5-(4'-chlorobiphenyl-4-yl)-3-hydroxypentanoic acid (Compound No. 20), Mass (m/z): 534.09 (M^+);

10 (2S,3R)-2-[2-(1,3-dioxo-1,3-dihydro-2H-isoindol-2-yl)ethyl]-3-hydroxy-5-[4'-(trifluoromethyl)biphenyl-4-yl]pentanoic acid (Compound No. 21), Mass (m/z): 512.08 ($M^+ + 1$);

(2S,3R)-2-[2-(1,3-dioxo-1,3-dihydro-2H-isoindol-2-yl)ethyl]-3-hydroxy-5-(4'-methylbiphenyl-4-yl)pentanoic acid (Compound No. 22), Mass (m/z): 458.14($M^+ + 1$);

15 (2S,3R)-2-[2-(1,3-dioxo-1,3-dihydro-2H-isoindol-2-yl)ethyl]-5-(3'-fluoro-4'-methoxybiphenyl-4-yl)-3-hydroxypentanoic acid (Compound No. 23), Mass (m/z): 492.07($M^+ + 1$);

(2S,3R)-3-hydroxy-5-[4-(6-methoxypyridin-3-yl)phenyl]-2-[2-(5-methyl-1,3-dioxo-1,3-dihydro-2H-isoindol-2-yl)ethyl]pentanoic acid (Compound No. 24), Mass
20 (m/z): 489.11 ($M^+ + 1$);

(2S,3R)-3-hydroxy-5-(4'-methylbiphenyl-4-yl)-2-[2-(4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 25), Mass (m/z): 458.14 ($M^+ + 1$);

(2S,3R)-5-(4'-ethylbiphenyl-4-yl)-3-hydroxy-2-[2-(4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 26), Mass (m/z): 472.15($M^+ + 1$);

25 (2S,3R)-5-(4'-chlorobiphenyl-4-yl)-3-hydroxy-2-[2-(5-methyl-1,3-dioxo-1,3-dihydro-2H-isoindol-2-yl)ethyl]pentanoic acid (Compound No. 27), Mass (m/z): 492.08 ($M^+ + 1$);

(2S,3R)-5-(4'-chlorobiphenyl-4-yl)-3-hydroxy-2-[2-(4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 28), Mass (m/z): 478.09 ($M^+ + 1$);

(2S,3R)-2-[2-(1,3-dioxo-1,3-dihydro-2H-isoindol-2-yl)ethyl]-3-hydroxy-5-(4-pyrimidin-5-ylphenyl)pentanoic acid (Compound No. 29), Mass (m/z):446.12 ($M^+ + 1$);

(2S,3R)-2-[2-(1,3-dioxo-1,3-dihydro-2H-isoindol-2-yl)ethyl]-3-hydroxy-5-[4-(1-methyl-1H-pyrazol-4-yl)phenyl]pentanoic acid (Compound No. 30), Mass (m/z): 448.25
5 ($M^+ + 1$);

(2S,3R)-2-[2-(1,3-dioxo-1,3-dihydro-2H-isoindol-2-yl)ethyl]-3-hydroxy-5-(4-pyridin-3-ylphenyl)pentanoic acid (Compound No. 31), Mass (m/z): 445.23($M^+ + 1$);

(2S,3R)-5-[4-(2,3-dihydro-1,4-benzodioxin-6-yl)phenyl]-3-hydroxy-2-[2-(4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 32), Mass (m/z):502.19
10 ($M^+ + 1$);

(2S,3R)-3-hydroxy-5-[4-(6-methoxypyridin-3-yl)phenyl]-2-[2-(8-methyl-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 33), Mass
(m/z):489.0($M^+ + 1$);

(2S,3R)-3-hydroxy-2-[2-(8-methyl-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-5-[4-
15 (6-methylpyridin-3-yl)phenyl]pentanoic acid (Compound No. 34), Mass (m/z):473.0($M^+ + 1$);

(2S,3R)-3-hydroxy-5-[4-(1-methyl-1H-pyrazol-4-yl)phenyl]-2-[2-(4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 35), Mass (m/z): 448.0 ($M^+ + 1$);

(2S,3R)-3-hydroxy-2-[2-(8-methoxy-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-5-
20 [4-(6-methoxypyridin-3-yl)phenyl]pentanoic acid (Compound No. 36), Mass (m/z): 505.0 ($M^+ + 1$);

(2S,3R)-2-[2-(7-chloro-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-3-hydroxy-5-[4-
(6-methoxypyridin-3-yl)phenyl]pentanoic acid (Compound No. 37), Mass (m/z):509.0
25 ($M^+ + 1$);

(2S,3R)-3-hydroxy-2-{2-[7-(6-methoxypyridin-3-yl)-4-oxo-1,2,3-benzotriazin-3(4H)-yl]ethyl}-5-[4-(6-methoxypyridin-3-yl)phenyl]pentanoic acid (Compound No. 38),
Mass (m/z):582.0 ($M^+ + 1$);

(2S,3R)-2-[2-(6,7-difluoro-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-3-hydroxy-5-[4-(6-methoxypyridin-3-yl)phenyl]pentanoic acid (Compound No. 39), Mass (m/z): 511.0 ($M^+ + 1$);

5 (2S,3R)-2-[2-(5-fluoro-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-3-hydroxy-5-[4-(6-methoxypyridin-3-yl)phenyl]pentanoic acid (Compound No. 40), Mass (m/z): 493.3 ($M^+ + 1$);

(2S,3R)-3-hydroxy-2-[2-(8-methoxy-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-5-[4-(6-methylpyridin-3-yl)phenyl]pentanoic acid (Compound No. 41), Mass (m/z): 489.0 ($M^+ + 1$);

10 (2S,3R)-3-hydroxy-2-{2-[5-(6-methoxypyridin-3-yl)-4-oxo-1,2,3-benzotriazin-3(4H)-yl]ethyl}-5-[4-(6-methoxypyridin-3-yl)phenyl]pentanoic acid (Compound No. 42), Mass (m/z): 582.0 ($M^+ + 1$);

15 (2S,3R)-5-(4'-chloro-3'-fluorobiphenyl-4-yl)-2-[2-(1,3-dioxo-1,3-dihydro-2H-isoindol-2-yl)ethyl]-3-hydroxypentanoic acid (Compound No. 43), Mass (m/z): 519.0 ($M^+ + 23$);

(2S,3R)-2-[2-(1,3-dioxo-1,3-dihydro-2H-isoindol-2-yl)ethyl]-3-hydroxy-5-[4-(1-isobutyl-1H-pyrazol-4-yl)phenyl]pentanoic acid (Compound No. 44), Mass (m/z): 490.0 ($M^+ + 1$);

20 (2S,3R)-5-biphenyl-4-yl-2-[2-(1,3-dioxo-1,3-dihydro-2H-isoindol-2-yl)ethyl]-3-hydroxypentanoic acid (Compound No. 45), Mass (m/z): 444.0 ($M^+ + 1$);

(2S,3R)-2-[2-(1,3-dioxo-1,3-dihydro-2H-isoindol-2-yl)ethyl]-5-(3'-fluoro-4'-methylbiphenyl-4-yl)-3-hydroxypentanoic acid (Compound No. 46), Mass (m/z): 476.0 ($M^+ + 1$);

25 (2S,3R)-5-(3,3'-difluoro-4'-methoxybiphenyl-4-yl)-2-[2-(1,3-dioxo-1,3-dihydro-2H-isoindol-2-yl)ethyl]-3-hydroxypentanoic acid (Compound No. 47), Mass (m/z): 510.0 ($M^+ + 1$);

(2S,3R)-5-[4-(2,3-dihydro-1,4-benzodioxin-6-yl)phenyl]-2-[2-(1,3-dioxo-1,3-dihydro-2H-isoindol-2-yl)ethyl]-3-hydroxypentanoic acid (Compound No. 48), Mass (m/z): 502.0 ($M^+ + 1$);

(2S,3R)-2-[2-(1,3-dioxo-1,3-dihydro-2H-isoindol-2-yl)ethyl]-3-hydroxy-5-[4-(1H-tetrazol-1-yl)phenyl]pentanoic acid (Compound No. 49), Mass (m/z): 458.0 ($M^+ + 23$);

(2S,3R)-2-[2-(1,3-dioxo-1,3-dihydro-2H-isoindol-2-yl)ethyl]-5-[2-fluoro-4-(6-methoxypyridin-3-yl)phenyl]-3-hydroxypentanoic acid (Compound No. 50), Mass
5 (m/z): 493.0 ($M^+ + 1$);

(2S,3R)-5-(3'-fluoro-4'-methylbiphenyl-4-yl)-3-hydroxy-2-[2-(4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 51), Mass (m/z): 498.0 ($M^+ + 23$);

(2S,3R)-3-hydroxy-5-[4-(1-isobutyl-1H-pyrazol-4-yl)phenyl]-2-[2-(4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 52), Mass (m/z): 490.0 ($M^+ + 1$);
10

(2S,3R)-5-biphenyl-4-yl-3-hydroxy-2-[2-(4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 53), Mass (m/z): 444.0 ($M^+ + 1$);

(2S,3R)-3-hydroxy-5-[4-(6-methoxypyridin-3-yl)phenyl]-2-[2-(7-methyl-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 54), Mass (m/z):
15 489.34 ($M^+ + 1$)

(2S,3R)-2-[2-(1,3-dioxo-1,3-dihydro-2H-isoindol-2-yl)ethyl]-5-[6-(3-fluoro-4-methoxyphenyl)pyridin-3-yl]-3-hydroxypentanoic acid (Compound No. 55), Mass (m/z): 493.16 ($M^+ + 1$);

(2S,3R)-5-(4'-chloro-3-fluorobiphenyl-4-yl)-2-[2-(1,3-dioxo-1,3-dihydro-2H-isoindol-2-yl)ethyl]-3-hydroxypentanoic acid (Compound No. 56), Mass (m/z): 496.13 ($M^+ + 1$);
20

(2S,3R)-2-[2-(1,3-dioxo-1,3-dihydro-2H-isoindol-2-yl)ethyl]-3-hydroxy-5-[6-(4-methoxyphenyl)pyridin-3-yl]pentanoic acid (Compound No. 57), Mass (m/z): 475.13 ($M^+ + 1$);
25

(2S,3R)-2-[2-(1,3-dioxo-1,3-dihydro-2H-isoindol-2-yl)ethyl]-5-(3-fluoro-4'-methoxybiphenyl-4-yl)-3-hydroxypentanoic acid (Compound No. 58), Mass (m/z): 492.13 ($M^+ + 1$);

(2S,3R)-5-[6-(4-chlorophenyl)pyridin-3-yl]-2-[2-(1,3-dioxo-1,3-dihydro-2H-isoindol-2-yl)ethyl]-3-hydroxypentanoic acid (Compound No. 59), Mass (m/z):479.12 (M^+ +1);

5 (2S,3R)-2-[2-(1,3-dioxo-1,3-dihydro-2H-isoindol-2-yl)ethyl]-3-hydroxy-5-(6'-methoxy-2,3'-bipyridin-5-yl)pentanoic acid (Compound No. 60), Mass (m/z): 476.16 (M^+ +1);

(2S,3R)-5-[4-(4-chlorophenyl)-2-thienyl]-2-[2-(1,3-dioxo-1,3-dihydro-2H-isoindol-2-yl)ethyl]-3-hydroxypentanoic acid (Compound No. 61), Mass (m/z):484.0 (M^+ +1);

10 (2S,3R)-2-[2-(1,3-dioxo-1,3-dihydro-2H-isoindol-2-yl)ethyl]-3-hydroxy-5-[4-(6-methoxypyridin-3-yl)-2-thienyl]pentanoic acid (Compound No. 62), Mass (m/z): 481.0(M^+ +1);

(2S,3R)-2-[2-(1,3-dioxo-1,3-dihydro-2H-isoindol-2-yl)ethyl]-3-hydroxy-5-[4-[4-(trifluoromethyl)phenyl]-2-thienyl]pentanoic acid (Compound No. 63), Mass (m/z):518.0 (M^+ +1);

15

(2S,3R)-2-[2-(1,3-dioxo-1,3-dihydro-2H-isoindol-2-yl)ethyl]-5-[4-(3-fluoro-4-methoxyphenyl)-2-thienyl]-3-hydroxypentanoic acid (Compound No. 64), Mass (m/z): 498.0 (M^+ +1);

(2S,3R)-3-hydroxy-2-[2-(7-methyl-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-5-[4'-(trifluoromethyl)biphenyl-4-yl]pentanoic acid (Compound No. 65), Mass (m/z):526.0 (M^+ +1);

20

(2S,3R)-3-hydroxy-2-[2-(6-methyl-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-5-[4'-(trifluoromethyl)biphenyl-4-yl]pentanoic acid (Compound No. 66), Mass (m/z):526.0 (M^+ +1);

25 (2S,3R)-2-[2-(7-chloro-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-3-hydroxy-5-(4'-methylbiphenyl-4-yl)pentanoic acid (Compound No. 67), Mass (m/z): 492.0(M^+ +1);

(2S,3R)-3-hydroxy-2-[2-(8-methoxy-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-5-[4'-(trifluoromethyl)biphenyl-4-yl]pentanoic acid (Compound No. 68), Mass (m/z): 542.29(M^+ +1);

(2S,3R)-2-[2-(6,7-difluoro-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-3-hydroxy-5-[4'-(trifluoromethyl)biphenyl-4-yl]pentanoic acid (Compound No. 69), Mass (m/z):548.54 ($M^+ + 1$);

5 (2S,3R)-2-[2-(1,3-dioxo-1,3-dihydro-2H-isoindol-2-yl)ethyl]-5-[2-fluoro-5-(6-methoxypyridin-3-yl)phenyl]-3-hydroxypentanoic acid (Compound No. 70), Mass (m/z): 493.30($M^+ + 1$);

(2S,3R)-5-(4'-chloro-4-fluorobiphenyl-3-yl)-2-[2-(1,3-dioxo-1,3-dihydro-2H-isoindol-2-yl)ethyl]-3-hydroxypentanoic acid (Compound No. 71), Mass (m/z):495.72 ($M^+ + 1$);

10 (2S,3R)-2-[2-(1,3-dioxo-1,3-dihydro-2H-isoindol-2-yl)ethyl]-5-[6-fluoro-4'-(trifluoromethyl)biphenyl-3-yl]-3-hydroxypentanoic acid (Compound No. 72), Mass (m/z):530.13 ($M^+ + 1$);

(2S,3R)-2-[2-(1,3-dioxo-1,3-dihydro-2H-isoindol-2-yl)ethyl]-5-[4-fluoro-3-(6-methoxypyridin-3-yl)phenyl]-3-hydroxypentanoic acid (Compound No. 73), Mass
15 (m/z):493.37 ($M^+ + 1$);

(2S,3R)-5-(4'-chloro-6-fluorobiphenyl-3-yl)-2-[2-(1,3-dioxo-1,3-dihydro-2H-isoindol-2-yl)ethyl]-3-hydroxypentanoic acid (Compound No. 74), Mass (m/z):496.26 ($M^+ + 1$);

(2S,3R)-5-(3',6-difluoro-4'-methoxybiphenyl-3-yl)-2-[2-(1,3-dioxo-1,3-dihydro-2H-isoindol-2-yl)ethyl]-3-hydroxypentanoic acid (Compound No. 75), Mass (m/z):509.77
20 ($M^+ + 1$);

(2S,3R)-2-[2-(5-fluoro-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-3-hydroxy-5-(4'-methoxybiphenyl-4-yl)pentanoic acid (Compound No. 76), Mass (m/z): 492.56($M^+ + 1$);

(2S,3R)-5-(4'-fluorobiphenyl-4-yl)-2-[2-(5-fluoro-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-3-hydroxypentanoic acid (Compound No. 77), Mass (m/z):480.0 ($M^+ + 1$);
25

(2S,3R)-3-hydroxy-2-[2-(6-methyl-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-5-[4-(1-methyl-1H-pyrazol-4-yl)phenyl]pentanoic acid (Compound No. 78), Mass (m/z): 462.19 ($M^+ + 1$);

(2S,3R)-2-[2-(5-chloro-1,3-dioxo-1,3-dihydro-2H-isoindol-2-yl)ethyl]-3-hydroxy-5-[4'-(trifluoromethyl)biphenyl-4-yl]pentanoic acid (Compound No. 79), Mass (m/z):546.32 ($M^+ + 1$);

5 (2S,3R)-2-[2-(4-fluoro-1,3-dioxo-1,3-dihydro-2H-isoindol-2-yl)ethyl]-3-hydroxy-5-[4'-(trifluoromethyl)biphenyl-4-yl]pentanoic acid (Compound No. 80), Mass (m/z):529.25 ($M^+ + 1$);

(2S,3R)-5-(3',4'-difluorobiphenyl-4-yl)-2-[2-(5-fluoro-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-3-hydroxypentanoic acid (Compound No. 81), Mass (m/z): 498.27($M^+ + 1$);

10 (2S,3R)-3-hydroxy-2-[2-(4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-5-phenylpentanoic acid (Compound No. 82); Mass (m/z):368.0 ($M^+ + 1$);

(2S,3R)-2-[2-(1,3-dioxo-1,3-dihydro-2H-isoindol-2-yl)ethyl]-3-hydroxy-5-phenylpentanoic acid (Compound No. 83), Mass (m/z): 368.07 ($M^+ + 1$);

15 (2S,3R)-2-[2-(1,3-dioxo-1,3-dihydro-2H-isoindol-2-yl)ethyl]-3-hydroxy-5-[4-(trifluoromethyl)phenyl]pentanoic acid (Compound No.84), Mass (m/z): 458.0($M^+ + 23$);

(2S,3R)-5-(4-tert-butylphenyl)-2-[2-(1,3-dioxo-1,3-dihydro-2H-isoindol-2-yl)ethyl]-3-hydroxypentanoic acid (Compound No.85), Mass (m/z):424.0 ($M^+ + 1$);

20 (2S,3R)-3-hydroxy-5-[4-(6-methoxypyridin-3-yl)phenyl]-2-[2-(4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 86), Mass (m/z): 474.87($M^+ + 1$);

(2S,3R)-5-(4'-chlorobiphenyl-4-yl)-2-[2-(1,3-dioxo-1,3-dihydro-2H-isoindol-2-yl)ethyl]-3-hydroxypentanoic acid (Compound No. 87), Mass (m/z): 478.09($M^+ + 1$);

25 (2S,3R)-2-[2-(1,3-dioxo-1,3-dihydro-2H-isoindol-2-yl)ethyl]-3-hydroxy-5-[4-(6-methoxypyridin-3-yl)phenyl]pentanoic acid (Compound No.88), Mass (m/z): 474.88($M^+ + 1$);

(2R,3R)-5-(4'-chlorobiphenyl-4-yl)-2-[2-(1,3-dioxo-1,3-dihydro-2H-isoindol-2-yl)ethyl]-3-hydroxypentanoic acid (Compound No. 89), Mass (m/z):478.2 ($M^+ + 1$);

(2*S*,3*R*)-3-hydroxy-2-[2-(7-methyl-4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)ethyl]-5-[4-(1-methyl-1*H*-pyrazol-4-yl)phenyl]pentanoic acid (Compound No. 90), Mass (*m/z*): 462.32($M^+ + 1$);

5 (2*S*,3*R*)-3-hydroxy-2-[2-(8-methoxy-4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)ethyl]-5-[4-(1-methyl-1*H*-pyrazol-4-yl)phenyl]pentanoic acid (Compound No. 91), Mass (*m/z*): 478.38($M^+ + 1$);

(2*S*,3*R*)-2-[2-(7-chloro-4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)ethyl]-5-(3',4'-difluorobiphenyl-4-yl)-3-hydroxypentanoic acid (Compound No. 92), Mass (*m/z*): 514.20($M^+ + 1$);

10 (2*S*,3*R*)-2-[2-(7-chloro-4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)ethyl]-3-hydroxy-5-(4'-methoxybiphenyl-4-yl)pentanoic acid (Compound No. 93), Mass (*m/z*): 508.29($M^+ + 1$);

(2*S*,3*R*)-2-[2-(7-chloro-4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)ethyl]-5-(4'-fluorobiphenyl-4-yl)-3-hydroxypentanoic acid (Compound No. 94),

15 (2*S*,3*R*)-2-[2-(7-chloro-4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)ethyl]-3-hydroxy-5-[4-(6-methylpyridin-3-yl)phenyl]pentanoic acid (Compound No. 95), Mass (*m/z*): 493.37($M^+ + 1$);

(2*R*,3*R*)-3-hydroxy-5-[4-(6-methoxypyridin-3-yl)phenyl]-2-[2-(4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)ethyl]pentanoic acid (Compound No. 96), Mass (*m/z*): 474.87($M^+ + 1$);

20 (2*S*,3*R*)-5-(2',4'-difluorobiphenyl-4-yl)-2-[2-(1,3-dioxo-1,3-dihydro-2*H*-isoindol-2-yl)ethyl]-3-hydroxypentanoic acid (Compound No. 97), Mass (*m/z*): 480.13($M^+ + 1$);

(2*S*,3*R*)-2-[2-(6-chloro-4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)ethyl]-3-hydroxy-5-[4-(1-methyl-1*H*-pyrazol-4-yl)phenyl]pentanoic acid (Compound No. 98), Mass (*m/z*): 482.15 ($M^+ + 1$);

25 (2*S*,3*R*)-2-[2-(5-chloro-4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)ethyl]-3-hydroxy-5-[4-(1-methyl-1*H*-pyrazol-4-yl)phenyl]pentanoic acid (Compound No. 99), Mass (*m/z*): 482.15($M^+ + 1$);

(2S,3R)-2-[2-(1,3-dioxo-1,3-dihydro-2H-isoindol-2-yl)ethyl]-3-hydroxy-5-(4'-isopropylbiphenyl-4-yl)pentanoic acid (Compound No. 100), Mass (m/z): 486.18(M⁺ +1);

(2S,3R)-5-(3'-chloro-4'-fluorobiphenyl-4-yl)-2-[2-(1,3-dioxo-1,3-dihydro-2H-isoindol-2-yl)ethyl]-3-hydroxypentanoic acid (Compound No. 101), Mass (m/z):
5 496.12(M⁺ +1);

(2S,3R)-5-(4'-butylbiphenyl-4-yl)-3-hydroxy-2-[2-(4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 102), Mass (m/z): 500.42(M⁺ +1);

(2S,3R)-5-(2'-fluorobiphenyl-4-yl)-3-hydroxy-2-[2-(4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 103), Mass (m/z): 462.32(M⁺ +1);

10 (2S,3R)-2-[2-(5-fluoro-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-3-hydroxy-5-[4-(1-methyl-1H-pyrazol-4-yl)phenyl]pentanoic acid (Compound No. 104), Mass (m/z): 466.31(M⁺ +1);

(2S,3R)-3-hydroxy-2-[2-(8-methyl-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-5-[4-(1-methyl-1H-pyrazol-4-yl)phenyl]pentanoic acid (Compound No. 105), Mass (m/z):
15 462.38(M⁺ +1);

(2S,3R)-2-[2-(1,3-dioxo-1,3-dihydro-2H-isoindol-2-yl)ethyl]-3-hydroxy-5-[6-[4-(trifluoromethyl)phenyl]pyridin-3-yl]pentanoic acid (Compound No. 106), Mass (m/z): 513.35(M⁺ +1);

(2S,3R)-5-[6-(3,4-difluorophenyl)pyridin-3-yl]-2-[2-(1,3-dioxo-1,3-dihydro-2H-isoindol-2-yl)ethyl]-3-hydroxypentanoic acid (Compound No. 107), Mass (m/z):
20 481.32(M⁺ +1);

(2S,3R)-5-[6-(4-chloro-3-fluorophenyl)pyridin-3-yl]-2-[2-(1,3-dioxo-1,3-dihydro-2H-isoindol-2-yl)ethyl]-3-hydroxypentanoic acid (Compound No. 108), Mass (m/z): 497.30(M⁺ +1);

25 (2S,3R)-2-[2-(1,3-dioxo-1,3-dihydro-2H-isoindol-2-yl)ethyl]-5-[6-(4-fluorophenyl)pyridin-3-yl]-3-hydroxypentanoic acid (Compound No. 109), Mass (m/z): 463.35(M⁺ +1);

(2S,3R)-2-[2-(1,3-dioxo-1,3-dihydro-2H-isoindol-2-yl)ethyl]-5-[6-(3-fluoro-4-methylphenyl)pyridin-3-yl]-3-hydroxypentanoic acid (Compound No. 110), Mass (m/z):
30 477.34(M⁺ +1);

(2S,3R)-2-[2-(7-chloro-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-3-hydroxy-5-[4-(1-methyl-1H-pyrazol-4-yl)phenyl]pentanoic acid (Compound No. 111), Mass (m/z): 482.38(M⁺ +1);

5 (2S,3R)-2-[2-(8-chloro-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-3-hydroxy-5-[4-(1-methyl-1H-pyrazol-4-yl)phenyl]pentanoic acid (Compound No. 112),

(2S,3R)-2-[2-(6,7-difluoro-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-3-hydroxy-5-[4-(1-methyl-1H-pyrazol-4-yl)phenyl]pentanoic acid (Compound No. 113), Mass (m/z): 484.39(M⁺ +1);

10 (2S,3R)-3-hydroxy-2-[2-(7-methyl-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-5-[4'-(trifluoromethoxy)biphenyl-4-yl]pentanoic acid (Compound No. 114), Mass (m/z): 542.36(M⁺ +1);

(2S,3R)-3-hydroxy-5-(4'-methylbiphenyl-4-yl)-2-[2-(7-methyl-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 115), Mass (m/z): 472.12(M⁺ +1);

15 (2S,3R)-5-(4'-fluorobiphenyl-4-yl)-3-hydroxy-2-[2-(7-methyl-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 116), Mass (m/z): 476.23(M⁺ +1);

20 (2S,3R)-5-(3',4'-difluorobiphenyl-4-yl)-3-hydroxy-2-[2-(7-methyl-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 117), Mass (m/z): 494.24(M⁺ +1);

(2S,3R)-3-hydroxy-2-[2-(7-methyl-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-5-[4-(6-methylpyridin-3-yl)phenyl]pentanoic acid (Compound No. 118), Mass (m/z): 473.38(M⁺ +1);

25 (2S,3R)-5-[4-(2,3-dihydro-1,4-benzodioxin-6-yl)phenyl]-3-hydroxy-2-[2-(7-methyl-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 119), Mass (m/z): 516.28(M⁺ +1);

(2R,3S)-3-hydroxy-5-[4-(6-methoxypyridin-3-yl)phenyl]-2-[2-(4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 120), Mass (m/z): 474.87(M⁺ +1);

(2S,3R)-2-[2-(5-chloro-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-3-hydroxy-5-(6'-methoxy-2,3'-bipyridin-5-yl)pentanoic acid (Compound No. 121), Mass (m/z): 510.23(M^+ +1);

5 (2S,3R)-2-[2-(7-chloro-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-3-hydroxy-5-(6'-methoxy-2,3'-bipyridin-5-yl)pentanoic acid (Compound No. 122), Mass (m/z): 510.23(M^+ +1);

(2S,3R)-2-[2-(6,7-difluoro-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-3-hydroxy-5-(6'-methoxy-2,3'-bipyridin-5-yl)pentanoic acid (Compound No. 123), Mass (m/z): 512.31(M^+ +1);

10 (2S,3R)-5-(3'-fluoro-4'-methylbiphenyl-4-yl)-3-hydroxy-2-[2-(7-methyl-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 124), Mass (m/z): 490.30(M^+ +1);

(2S,3R)-3-hydroxy-5-[4-(1-isobutyl-1H-pyrazol-4-yl)phenyl]-2-[2-(7-methyl-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 125), Mass (m/z): 504.33(M^+ +1);

15

(2S,3R)-2-[2-(5-fluoro-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-3-hydroxy-5-(6'-methoxy-2,3'-bipyridin-5-yl)pentanoic acid (Compound No. 126), Mass (m/z): 510.23(M^+ +1);

(2S,3R)-5-(4'-fluorobiphenyl-4-yl)-3-hydroxy-2-[2-(8-methoxy-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 127), Mass (m/z): 492.26 (M^+ +1);

20

(2S,3R)-5-(3',4'-difluorobiphenyl-4-yl)-3-hydroxy-2-[2-(8-methoxy-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 128), Mass (m/z): 510.30 (M^+ +1);

25 (2S,3R)-3-hydroxy-5-(4'-methoxybiphenyl-4-yl)-2-[2-(8-methoxy-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 129), Mass (m/z): 504.38 (M^+ +1);

(2S,3R)-2-[2-(7-chloro-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-3-hydroxy-5-[6-[4-(trifluoromethyl)phenyl]pyridin-3-yl]pentanoic acid (Compound No. 130), Mass (m/z): 547.29 (M^+ +1);

30

(2S,3R)-2-[2-(6,7-difluoro-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-3-hydroxy-5-{6-[4-(trifluoromethyl)phenyl]pyridin-3-yl}pentanoic acid (Compound No. 131), Mass (m/z): 549.30 ($M^+ + 1$);

5 (2S,3R)-2-[2-(5-chloro-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-3-hydroxy-5-{6-[4-(trifluoromethyl)phenyl]pyridin-3-yl}pentanoic acid (Compound No. 132), Mass (m/z): 547.29 ($M^+ + 1$);

(2S,3R)-3-hydroxy-2-[2-(7-methyl-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-5-{6-[4-(trifluoromethyl)phenyl]pyridin-3-yl}pentanoic acid (Compound No. 133), Mass (m/z): 527.40 ($M^+ + 1$);

10 (2S,3R)-2-[2-(5-fluoro-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-3-hydroxy-5-{6-[4-(trifluoromethyl)phenyl]pyridin-3-yl}pentanoic acid (Compound No. 134), Mass (m/z): 531.36 ($M^+ + 1$);

(2S,3R)-3-hydroxy-2-[2-(8-methoxy-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-5-(4'-methylbiphenyl-4-yl)pentanoic acid (Compound No. 135), Mass (m/z): 488.34 ($M^+ + 1$);

15

(2S,3R)-5-(3'-fluoro-4'-methylbiphenyl-4-yl)-2-[2-(5-fluoro-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-3-hydroxypentanoic acid (Compound No. 136), Mass (m/z): 494.33 ($M^+ + 1$);

(2S,3R)-2-[2-(6,7-difluoro-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-3-hydroxy-5-(4'-methylbiphenyl-4-yl)pentanoic acid (Compound No. 137), Mass (m/z): 494.33($M^+ + 1$);

20

(2S,3R)-2-[2-(5-fluoro-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-3-hydroxy-5-(4'-methylbiphenyl-4-yl)pentanoic acid (Compound No. 138), Mass (m/z): 476.30 ($M^+ + 1$);

(2S,3R)-5-(4'-chlorobiphenyl-4-yl)-2-[2-(5-fluoro-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-3-hydroxypentanoic acid (Compound No. 139), Mass (m/z): 496.30 ($M^+ + 1$);

25 (2S,3R)-3-hydroxy-5-(6'-methoxy-2,3'-bipyridin-5-yl)-2-[2-(4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 140), Mass (m/z): 476.35 ($M^+ + 1$);

(2S,3R)-3-hydroxy-5-(6'-methoxy-2,3'-bipyridin-5-yl)-2-[2-(6-methyl-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 141),

(2S,3R)-3-hydroxy-5-(6'-methoxy-2,3'-bipyridin-5-yl)-2-[2-(7-methyl-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 142), Mass (m/z): 490.34($M^+ + 1$);

5 (2S,3S)-3-hydroxy-5-[4-(6-methoxypyridin-3-yl)phenyl]-2-[2-(4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 143), Mass (m/z): 475.21($M^+ + 1$);

(2S,3R)-2-[2-(6,7-difluoro-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-5-(3'-fluoro-4'-methylbiphenyl-4-yl)-3-hydroxypentanoic acid (Compound No. 144), Mass (m/z): 512.32 ($M^+ + 1$);

10 (2S,3R)-5-(4'-chlorobiphenyl-4-yl)-2-[2-(6,7-difluoro-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-3-hydroxypentanoic acid (Compound No. 145), Mass (m/z): 514.37($M^+ + 1$);

(2S,3R)-5-(3'-fluoro-4'-methylbiphenyl-4-yl)-3-hydroxy-2-[2-(8-methoxy-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 146), Mass (m/z): 506.38 ($M^+ + 1$);

15

(2S,3R)-5-[4-(2,3-dihydro-1,4-benzodioxin-6-yl)phenyl]-3-hydroxy-2-[2-(8-methoxy-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 147), Mass (m/z): 532.38($M^+ + 1$);

(2S,3R)-5-[4-(6-chloropyridin-3-yl)phenyl]-3-hydroxy-2-[2-(4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 148), Mass (m/z): 479.36 ($M^+ + 1$);

20

(2S,3R)-3-hydroxy-2-[2-(6-methoxy-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-5-[4'-(trifluoromethyl)biphenyl-4-yl]pentanoic acid (Compound No. 149), Mass (m/z): 542.11($M^+ + 1$);

25 (2S,3R)-3-hydroxy-2-[2-(7-methoxy-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-5-[4'-(trifluoromethyl)biphenyl-4-yl]pentanoic acid (Compound No. 150), Mass (m/z): 542.11($M^+ + 1$);

(2S,3R)-3-hydroxy-2-[2-(6-methoxy-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-5-(4'-methylbiphenyl-4-yl)pentanoic acid (Compound No. 151), Mass (m/z): 488.09 ($M^+ + 1$);

30

(2S,3R)-3-hydroxy-5-(4'-methoxybiphenyl-4-yl)-2-[2-(6-methoxy-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 152), Mass (m/z): 504.06 (M^+ +1);

5 (2S,3R)-5-[2-fluoro-4-(6-methoxypyridin-3-yl)phenyl]-3-hydroxy-2-[2-(4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 153), Mass (m/z): 493.01(M^+ +1);

(2S,3R)-5-[2-fluoro-4-(6-methoxypyridin-3-yl)phenyl]-3-hydroxy-2-[2-(7-methoxy-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 154), Mass (m/z): 523.02 (M^+ +1);

10 (2S,3R)-5-[2-fluoro-4-(6-methoxypyridin-3-yl)phenyl]-2-[2-(5-fluoro-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-3-hydroxypentanoic acid (Compound No. 155), Mass (m/z): 511.04(M^+ +1);

(2S,3R)-5-[2-fluoro-4-(6-methoxypyridin-3-yl)phenyl]-3-hydroxy-2-[2-(7-methyl-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 156), Mass (m/z):
15 507.05 (M^+ +1);

(2S,3R)-5-[2-fluoro-4-(6-methoxypyridin-3-yl)phenyl]-3-hydroxy-2-[2-(6-methyl-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 157), Mass (m/z):507.05 (M^+ +1);

(2S,3R)-3-hydroxy-5-(4'-methoxybiphenyl-4-yl)-2-[2-(7-methoxy-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 158), Mass (m/z): 504.09 (M^+ +1);
20

(2S,3R)-5-(4'-chlorobiphenyl-4-yl)-3-hydroxy-2-[2-(6-methoxy-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 159), Mass (m/z): 508.03(M^+ +1);

25 (2S,3R)-3-hydroxy-2-[2-(7-methoxy-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-5-[4-(6-methoxypyridin-3-yl)phenyl]pentanoic acid (Compound No. 160), Mass (m/z): 505.11(M^+ +1);

(2S,3R)-5-(3',4'-difluorobiphenyl-4-yl)-3-hydroxy-2-[2-(7-methoxy-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 161), Mass (m/z): 510.04(M^+ +1);
30

(2S,3R)-3-hydroxy-2-[2-(6-methoxy-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-5-[4-(6-methoxypyridin-3-yl)phenyl]pentanoic acid (Compound No. 162), Mass (m/z): 505.11($M^+ + 1$);

5 (2S,3R)-2-[2-(5-chloro-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-5-[2-fluoro-4-(6-methoxypyridin-3-yl)phenyl]-3-hydroxypentanoic acid (Compound No. 163), Mass (m/z): 526.95($M^+ + 1$);

(2S,3R)-2-[2-(6,7-difluoro-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-5-[2-fluoro-4-(6-methoxypyridin-3-yl)phenyl]-3-hydroxypentanoic acid (Compound No. 164), Mass (m/z): 529.00 ($M^+ + 1$);

10 (2S,3R)-5-[2-fluoro-4-(6-methoxypyridin-3-yl)phenyl]-3-hydroxy-2-[2-(5-methyl-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 165), Mass (m/z): 507.03($M^+ + 1$);

(2S,3R)-5-(4'-fluorobiphenyl-4-yl)-3-hydroxy-2-[2-(5-methyl-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 166), Mass (m/z): 476.01($M^+ + 1$);

15

(2S,3R)-3-hydroxy-5-[4-(6-methoxypyridin-3-yl)phenyl]-2-[2-(5-methyl-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 167), Mass (m/z): 489.05($M^+ + 1$);

(2S,3R)-3-hydroxy-5-(4'-methoxybiphenyl-4-yl)-2-[2-(5-methyl-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 168), Mass (m/z): 488.06 ($M^+ + 1$);

20

(2S,3R)-3-hydroxy-5-(4'-methylbiphenyl-4-yl)-2-[2-(5-methyl-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 169), Mass (m/z): 472.03($M^+ + 1$);

25 (2S,3R)-5-(4'-chlorobiphenyl-4-yl)-3-hydroxy-2-[2-(5-methyl-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 170), Mass (m/z): 491.98 ($M^+ + 1$);

(2S,3R)-5-(4'-fluorobiphenyl-4-yl)-3-hydroxy-2-[2-(6-methoxy-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 171), Mass (m/z): 492.00($M^+ + 1$);

30

(2S,3R)-5-(3',4'-difluorobiphenyl-4-yl)-3-hydroxy-2-[2-(6-methoxy-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 172), Mass (m/z): 509.98(M^+ +1);

5 (2S,3R)-3-hydroxy-2-[2-(6-methoxy-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-5-[4'-(trifluoromethoxy)biphenyl-4-yl]pentanoic acid (Compound No. 173), Mass (m/z): 557.99 (M^+ +1);

(2S,3R)-5-(3'-fluoro-4'-methylbiphenyl-4-yl)-3-hydroxy-2-[2-(6-methoxy-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 174), Mass (m/z): 506.04(M^+ +1);

10 (2S,3R)-2-[2-(7-chloro-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-5-[2-fluoro-4-(6-methoxypyridin-3-yl)phenyl]-3-hydroxypentanoic acid (Compound No. 175), Mass (m/z): 526.91(M^+ +1);

(2S,3R)-3-hydroxy-5-[4-(1-methyl-1H-pyrazol-4-yl)-2-thienyl]-2-[2-(4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 176),

15 (2S,3R)-5-[2-fluoro-4-(6-methoxypyridin-3-yl)phenyl]-3-hydroxy-2-[2-(6-methoxy-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 177), Mass (m/z): 523.01(M^+ +1);

(2S,3R)-3-hydroxy-5-(6'-methoxy-2,3'-bipyridin-5-yl)-2-[2-(6-methoxy-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 178),

20 (2S,3R)-3-hydroxy-2-[2-(6-methoxy-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-5-[4-(6-methylpyridin-3-yl)phenyl]pentanoic acid (Compound No. 179), Mass (m/z): 489.07(M^+ +1);

(2S,3R)-5-[4-(2-chloropyridin-3-yl)phenyl]-3-hydroxy-2-[2-(6-methoxy-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 180), Mass (m/z):

25 508.99(M^+ +1);

(2S,3R)-2-[2-(6-fluoro-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-3-hydroxy-5-[4-(6-methoxypyridin-3-yl)phenyl]pentanoic acid (Compound No. 181), Mass (m/z): 493.19(M^+ +1);

30 (2S,3R)-5-(4'-chlorobiphenyl-4-yl)-2-[2-(6-fluoro-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-3-hydroxypentanoic acid (Compound No. 182), Mass (m/z): 496.17(M^+ +1);

(2S,3R)-5-(3',4'-difluorobiphenyl-4-yl)-2-[2-(6-fluoro-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-3-hydroxypentanoic acid (Compound No. 183), Mass (m/z): 498.15(M^+ +1);

5 (2S,3R)-2-[2-(6-fluoro-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-3-hydroxy-5-(4'-methoxybiphenyl-4-yl)pentanoic acid (Compound No. 184), Mass (m/z): 492.16 (M^+ +1);

(2S,3R)-3-hydroxy-5-[4-(2-methoxypyrimidin-5-yl)phenyl]-2-[2-(4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 185), Mass (m/z):476.13 (M^+ +1);

10 (2S,3R)-5-[2-fluoro-4-(1-methyl-1H-pyrazol-4-yl)phenyl]-3-hydroxy-2-[2-(6-methyl-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 186), Mass (m/z):480.08 (M^+ +1);

(2S,3R)-5-[2-fluoro-4-(1-methyl-1H-pyrazol-4-yl)phenyl]-3-hydroxy-2-[2-(7-methyl-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 187), Mass (m/z):480.15 (M^+ +1);

15 (2S,3R)-2-[2-(5-chloro-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-5-[2-fluoro-4-(1-methyl-1H-pyrazol-4-yl)phenyl]-3-hydroxypentanoic acid (Compound No. 188), Mass (m/z):500.07 (M^+ +1);

20 (2S,3R)-2-[2-(7-chloro-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-5-[2-fluoro-4-(1-methyl-1H-pyrazol-4-yl)phenyl]-3-hydroxypentanoic acid (Compound No. 189), Mass (m/z):500.13 (M^+ +1);

(2S,3R)-5-[2-fluoro-4-(1-methyl-1H-pyrazol-4-yl)phenyl]-2-[2-(5-fluoro-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-3-hydroxypentanoic acid (Compound No. 190), Mass (m/z):484.15 (M^+ +1);

25 (2S,3R)-2-[2-(6,7-difluoro-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-5-[2-fluoro-4-(1-methyl-1H-pyrazol-4-yl)phenyl]-3-hydroxypentanoic acid (Compound No. 191), Mass (m/z):502.16 (M^+ +1);

(2S,3R)-5-[2-fluoro-4-(1-methyl-1H-pyrazol-4-yl)phenyl]-2-[2-(6-fluoro-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-3-hydroxypentanoic acid (Compound No. 192), Mass (m/z):507.22(M^+ +23);

(2S,3R)-5-[2-fluoro-4-(1-methyl-1H-pyrazol-4-yl)phenyl]-3-hydroxy-2-[2-(8-methyl-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 193),

(2S,3R)-3-hydroxy-2-[2-(6-methoxy-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-5-[4-(2-methoxypyrimidin-5-yl)phenyl]pentanoic acid (Compound No. 194), Mass
5 (m/z):506.12 ($M^+ + 1$);

(2S,3R)-2-[2-(1,3-dioxo-1,3-dihydro-2H-isoindol-2-yl)ethyl]-3-hydroxy-5-[4-(2-methoxypyrimidin-5-yl)phenyl]pentanoic acid (Compound No. 195), Mass (m/z):476.18 ($M^+ + 1$);

(2S,3R)-5-[2-fluoro-4-(1-methyl-1H-pyrazol-4-yl)phenyl]-3-hydroxy-2-[2-(1-oxophthalazin-2(1H)-yl)ethyl]pentanoic acid (Compound No. 196), Mass (m/z):465.26 ($M^+ + 1$);
10

(2S,3R)-5-[2-fluoro-4-(1-methyl-1H-pyrazol-4-yl)phenyl]-3-hydroxy-2-[2-(3-methyl-2,6-dioxo-3,6-dihydropyrimidin-1(2H)-yl)ethyl]pentanoic acid (Compound No. 197), Mass (m/z):445.25 ($M^+ + 1$);

(2S,3R)-3-hydroxy-5-[4-(6-methoxypyridin-3-yl)phenyl]-2-[2-[4-oxo-7-(trifluoromethyl)-1,2,3-benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 198),
15 Mass (m/z):543.22 ($M^+ + 1$);

(2S,3R)-3-hydroxy-5-[4-(6-methoxypyridin-3-yl)phenyl]-2-[2-(1-oxophthalazin-2(1H)-yl)ethyl]pentanoic acid (Compound No. 199), Mass (m/z):474.25 ($M^+ + 1$);

(2S,3R)-3-hydroxy-5-[4-(6-methoxypyridin-3-yl)phenyl]-2-[2-(3-methyl-2,6-dioxo-3,6-dihydropyrimidin-1(2H)-yl)ethyl]pentanoic acid (Compound No. 200), Mass
20 (m/z):454.24 ($M^+ + 1$);

(2S,3R)-2-[2-(7,9-dioxo-8-azaspiro[4.5]dec-8-yl)ethyl]-3-hydroxy-5-[4-(6-methoxypyridin-3-yl)phenyl]pentanoic acid (Compound No. 201), Mass (m/z):495.27 ($M^+ + 1$);
25

(2S,3R)-2-[2-(2,4-dioxo-2H-1,3-benzoxazin-3(4H)-yl)ethyl]-3-hydroxy-5-[4-(6-methoxypyridin-3-yl)phenyl]pentanoic acid (Compound No. 202), Mass (m/z):491.17 ($M^+ + 1$);

(2S,3R)-3-hydroxy-5-[4-(6-methoxypyridin-3-yl)phenyl]-2-[2-(3,4,4-trimethyl-2,5-dioximidazolidin-1-yl)ethyl]pentanoic acid (Compound No. 203), Mass (m/z):470.12($M^+ + 1$);

5 (2S,3R)-5-(4-chloro-3-fluorophenyl)-2-[2-(1,3-dioxo-1,3-dihydro-2H-isoindol-2-yl)ethyl]-3-hydroxypentanoic acid (Compound No. 204), Mass (m/z): 420.06 ($M^+ + 1$);

(2R,3S)-2-[2-(1,3-dioxo-1,3-dihydro-2H-isoindol-2-yl)ethyl]-5-(3'-fluoro-4'-methylbiphenyl-4-yl)-3-hydroxypentanoic acid (Compound No. 205), Mass (m/z): 476.07 ($M^+ + 1$);

10 (2R,3S)-2-[2-(1,3-dioxo-1,3-dihydro-2H-isoindol-2-yl)ethyl]-5-(4'-fluorobiphenyl-4-yl)-3-hydroxypentanoic acid (Compound No. 206), Mass (m/z):462.03 ($M^+ + 1$);

(2R,3S)-5-(3',4'-difluorobiphenyl-4-yl)-2-[2-(1,3-dioxo-1,3-dihydro-2H-isoindol-2-yl)ethyl]-3-hydroxypentanoic acid (Compound No. 207), Mass (m/z):480.03 ($M^+ + 1$);

(2S,3S)-3-hydroxy-5-(4'-methoxybiphenyl-4-yl)-2-[2-(4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 208), Mass (m/z): 474.31 ($M^+ + 1$);

15 (2S,3R)-3-hydroxy-5-[4-(5-methylpyridin-2-yl)phenyl]-2-[2-(4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 209), Mass (m/z): 459.21($M^+ + 1$);

(2S,3R)-5-[4-(6-fluoropyridin-3-yl)phenyl]-3-hydroxy-2-[2-(4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 210), Mass (m/z): 463.17($M^+ + 1$);

20

(2S,3R)-2-[2-(5-fluoro-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-3-hydroxy-5-[4'-(trifluoromethyl)biphenyl-4-yl]pentanoic acid (Compound No. 211), Mass (m/z): 530.33($M^+ + 1$);

(2S,3R)-5-(4'-chlorobiphenyl-4-yl)-3-hydroxy-2-[2-(7-methyl-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 212), Mass (m/z): 492.29($M^+ + 1$);

25

(2S,3R)-3-hydroxy-5-(6'-methoxy-2,3'-bipyridin-5-yl)-2-[2-(8-methyl-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 213), Mass (m/z): 490.34 ($M^+ + 1$);

(2*S*,3*R*)-5-(3'-fluoro-4'-methylbiphenyl-4-yl)-3-hydroxy-2-[(4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)methyl]pentanoic acid (Compound No. 214),

(2*S*,3*R*)-3-hydroxy-2-[(4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)methyl]-5-[4'-(trifluoromethyl)biphenyl-4-yl]pentanoic acid (Compound No. 215), Mass (m/z): 498.09
5 (M⁺ +1);

(2*S*,3*R*)-5-(2',4'-difluorobiphenyl-4-yl)-3-hydroxy-2-[(4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)methyl]pentanoic acid (Compound No. 216), Mass (m/z): 466.06 (M⁺ +1);

((2*S*,3*R*)-5-(4'-fluorobiphenyl-4-yl)-3-hydroxy-2-[(4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)methyl]pentanoic acid (Compound No. 217), Mass (m/z): 448.07 (M⁺ +1);

10 (2*S*,3*R*)-5-(3'-fluoro-4'-methoxybiphenyl-4-yl)-3-hydroxy-2-[(4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)methyl]pentanoic acid (Compound No. 218), Mass (m/z): 478.22 (M⁺ +1);

(2*S*,3*R*)-3-hydroxy-5-[4-(2-methoxypyrimidin-5-yl)phenyl]-2-[(4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)methyl]pentanoic acid (Compound No. 219), Mass (m/z): 462.23
15 (M⁺ +1);

(2*S*,3*R*)-3-hydroxy-5-[4-(6-methoxypyridin-3-yl)phenyl]-2-[(4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)methyl]pentanoic acid (Compound No. 220), Mass (m/z): 461.23 (M⁺ +1);

(2*S*,3*R*)-3-hydroxy-5-(4'-methylbiphenyl-4-yl)-2-[(4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)methyl]pentanoic acid (Compound No. 221), Mass (m/z): 444.24 (M⁺ +1);
20

(2*S*,3*R*)-3-hydroxy-2-[(4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)methyl]-5-[4'-(trifluoromethoxy)biphenyl-4-yl]pentanoic acid (Compound No. 222), Mass (m/z): 514.14 (M⁺ +1);

(2*S*,3*R*)-3-hydroxy-5-[4-(6-methylpyridin-3-yl)phenyl]-2-[(4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)methyl]pentanoic acid (Compound No. 223), Mass (m/z): 445.18
25 (M⁺ +1);

(2*S*,3*R*)-5-(4'-chlorobiphenyl-4-yl)-3-hydroxy-2-[(4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)methyl]pentanoic acid (Compound No. 224), Mass (m/z): 464.18 (M⁺ +1);

(2*S*,3*R*)-5-(3',4'-difluorobiphenyl-4-yl)-3-hydroxy-2-[(4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)methyl]pentanoic acid (Compound No. 225), Mass (m/z): 466.20 ($M^+ + 1$);

(2*S*,3*R*)-3-hydroxy-5-[4-(6-hydroxypyridin-3-yl)phenyl]-2-[2-(4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)ethyl]pentanoic acid (Compound No. 226), Mass (m/z): 461.23 ($M^+ + 1$);

(2*S*,3*R*)-3-hydroxy-5-(4'-methylbiphenyl-4-yl)-2-{2-[4-oxo-7-(trifluoromethyl)-1,2,3-benzotriazin-3(4*H*)-yl]ethyl}pentanoic acid (Compound No. 227), Mass (m/z): 526.16 ($M^+ + 1$);

(2*S*,3*R*)-2-[2-(2,4-dioxo-1,4-dihydroquinazolin-3(2*H*)-yl)ethyl]-3-hydroxy-5-[4-(6-methoxypyridin-3-yl)phenyl]pentanoic acid (Compound No. 228), Mass (m/z): 490.17 ($M^+ + 1$)

(2*S*,3*R*)-3-(acetyloxy)-5-[4-(6-methoxypyridin-3-yl)phenyl]-2-[2-(4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)ethyl]pentanoic acid (Compound No. 229), Mass (m/z): 517.06 ($M^+ + 1$);

(2*S*,3*R*)-2-[2-(8-chloro-4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)ethyl]-3-hydroxy-5-[4-(6-methoxypyridin-3-yl)phenyl]pentanoic acid (Compound No. 230) Mass (m/z): 509.17 ($M^+ + 1$);

(2*S*,3*R*)-5-(4'-chlorobiphenyl-4-yl)-2-[2-(2,4-dioxo-2*H*-1,3-benzoxazin-3(4*H*)-yl)ethyl]-3-hydroxypentanoic acid (Compound No. 231), Mass (m/z): 493.93 ($M^+ + 1$)

(2*S*,3*R*)-5-(4'-chlorobiphenyl-4-yl)-3-hydroxy-2-(2-{[(2-hydroxyphenyl) carbonyl] amino}ethyl)pentanoic acid (Compound No. 232), Mass (m/z): 468.14 ($M^+ + 1$)

Example: Assay for Matrix Metallo Proteinases (MMPs)

New chemical entities prepared according to the present invention and corresponding standards used in the present invention were prepared (stock 10 mM) in 100% DMSO and subsequent dilutions were made in 50% DMSO-50% TCNB (50 mM Tris, 10 mM CaCl₂, 150 mM NaCl, 0.05% Brij-35, pH 7.5). 1 μ l of the compound and 88 μ l of TCNB was added to wells of 96 well plate to achieve the desired final concentration of NCE (final DMSO concentration should not exceed 0.5%). 1 μ l of activated, recombinant MMPs was added to each well (20-100 ng/100 μ l reaction mixture) except the "negative well." (MMP-1, 9 & 14 enzymes require prior activation. For this, supplied enzyme was incubated with

either APMA, final concentration 1 mM, for a time period of 1 hr at 37°C). Incubation was done at room temperature for 4-5 minutes. Reaction was initiated with 10 µl of 100 µM substrate (ES001: Aliquots were freshly diluted in TCNB; stock: 2 mM) and increase in florescence was monitored at excitation wavelength 320 nm followed by emission at 405 nm for 25-30 cycles. Increase in florescence (RFU) was calculated for positive, negative and NCE/standard wells. The percent inhibition compared to controls was calculated and IC₅₀ values determined using Graph-prism software.

The present invention relates to compounds that act as dual MMP-9/12 inhibitors, which have desirable activity profiles and beneficial potency, selectivity and/or pharmacokinetic properties.

In particular, compounds disclosed herein exhibited activity in MMP9 assays from ≤ 0.02 nM to about 40 µM, or from ≤ 0.02 nM to about 200 nM, or from ≤ 0.02 nM to about 20 nM, or from ≤ 0.02 nM to about 1.0 nM, or from ≤ 0.02 nM to about 0.3 nM. Compounds disclosed herein exhibited activity in MMP12 assays from ≤ 0.02 nM to about 3.8 µM, or from ≤ 0.02 nM to about 200 nM, or from ≤ 0.02 nM to about 20 nM, or from ≤ 0.02 nM to about 1.0 nM, or from ≤ 0.02 nM to about 0.3 nM. Particular compounds tested (Nos. 2-28, 30, 32-43, 46, 50-51, 55-58, 60-61, 63, 65-69, 76, 79-80, 86-89, 93-96, 98, 106-107, 113-122, 126, 130-132, 134, 136-138, 143-144, 148-150, 155, 157, 159, 173-175, 177, 179, 181-182, 184-186, 190, 192-195, 198 and 210-211) exhibited activity in MMP1 assays from about 100 nM to about 10 µM, for example from about 100 nM to about 5 µM, or from about 100 nM to about 2 µM, or from about 100 nM to about 1 µM indicating that compounds of the present invention can be selective over MMP-1 by ≥ 100 fold.

Assay for *in vivo* LPS induced rat neutrophilia model

Male wistar rats were treated with vehicle / NCEs (new chemical entities) or standard drug and 2 hrs later challenged with Lipopolysaccharide (LPS) in Phosphate buffered saline (PBS), by oro-intra tracheal route (400µL of 50µg/ml). Negative control animals received PBS alone. Intratracheal instillation was done under Ketamine and Xylazine anaesthesia. Two hours post-LPS challenge, rats were euthanised and bronchoalveolar lavage (BAL) performed. The TLC, DLC was done to enumerate

neutrophil count in the BAL fluid and results were expressed as percent inhibition using the following formula

$$\text{NeuLPS} - \text{NeuTEST}$$

$$\% \text{ Inhibition} = \frac{\text{NeuLPS} - \text{NeuTEST}}{\text{NeuLPS} - \text{NeuPBS}} \times 100$$

5 NeuLPS – NeuPBS

Where,

NeuLPS = Neutrophil count in vehicle treated LPS challenged group

NeuTEST = Neutrophil count in group treated with a given dose of test compound

NeuPBS = Neutrophil in vehicle treated group challenged with PBS

10 **Solubility Assessment**

Equilibrium Solubility:

 The pH-solubility profile of a compound is determined at 37°C in aqueous media with a pH in the range of 1-7.5. A sufficient number of pH conditions are evaluated to accurately define the pH-solubility profile. Standard buffer solutions described in the USP
15 are considered appropriate for use in solubility studies.

 The compound is weighed and transferred to the flasks. Media are added to each conical flask, the flask is sealed with stopper and paraffin film. The volume added is dependant upon the volume required for analysis of the content of the compound. The pH of the solution is measured after addition of the compound. The
20 flask is observed intermittently. If the drug substance is completely dissolved, additional amount of the compound is added until saturation (undissolved residue is observed) and pH is measured. Flasks are removed from water bath after equilibrium is achieved. The saturated solution is filtered through 0.45 µm membrane filter and the samples are analyzed to estimate the content of the test compound.

Pharmacokinetic screening assays for Matrix Metallo Proteinase (MMP 9/12) inhibitors

Intrinsic clearance:

Intrinsic clearance (or metabolism stability) is assessed by estimating the rate of initial decay of parent compound in a suitable biological matrix like human liver microsomes.

The study reaction consists of NADPH regeneration system and liver microsomes of the various species of interest (human, rat dog, mouse), added into buffer at a concentration of 0.5 mg/mL. After a short preincubation the metabolic reaction is initiated by the addition of 5 μ L of the substrate stock (100 μ M) to yield a final concentration of 0.5 μ M in the reaction. Periodic aliquots are drawn every three minutes until 30 minutes, quenched and the test compound concentration is estimated by LCMS. The rate of disappearance is estimated as the first order slope of the % parent remaining vs. time graph. The rate of decay is normalized to unit concentration of test compound and protein and extrapolated to 1g liver by using scaling factors (52.5 mg of CYP microsomal protein per gram of liver).

***Invitro* glucuronidation:**

A comparative assay in the form of intrinsic clearance (see above) with the addition of glucuronic acid and alamethacin to compare primarily the parent disappearance due to glucuronidation. Expressed as rate of clearance and normalized to per gram of liver.

Plasma protein binding:

Assessed by the equilibrium dialysis method where the unbound compound diffuses across a semi-permeable membrane and equilibrates with phosphate buffer (pH 7.4) is estimated, and is subtracted from the total drug in plasma to determine the percentage bound.


Equilibrium dialysis membranes are soaked overnight and the assembly is prepared. The test drug is spiked into plasma (100 and 1000 ng/mL) and incubated at 37°C and is transferred into the equilibrium apparatus with plasma added in one compartment and buffer in the other. The unit is rotated at constant rpm at 37°C for four hours to allow the unbound compound to dialyse and distribute with the buffer chamber. After four hours

the plasma and the buffer are removed from the respective compartments and the test compound concentrations are estimated. The percentage bound is estimated from the test compound concentrations.

- 5 Alternate methods include ultra filtration method where the compound spiked in plasma (100 and 1000 ng/mL) is filtered with Centricon filters (Millipore) having molecular weight cut off of (30000 DA) to prepare the retentate and ultra filtrate. The test compound is estimated in both and the percentage bound is calculated.


CLAIMS:

1. A chiral auxiliary which is (4S)-4-benzyl-1,3-thiazolidin-2-one.
2. Use of a chiral auxiliary of claim 1 in chemical synthetic transformation reactions selected from asymmetric aldol condensation, stereoselective alkylation, stereoselective Diels-Alder reaction, stereoselective Michael reactions, stereoselective differentiation reactions of enantiotopic groups in molecules bearing prochiral centers
3. A process for preparing a compound of Formula XX comprising:
 - a. reacting a compound of Formula VIIIa




Formula VIIIa

with a compound of Formula IX




to give a compound of Formula X;




Formula X

- b. deprotecting the compound of Formula X to give a compound of Formula XI;




Formula XI

- c. oxidizing the compound of Formula XI to give a compound of Formula XII;




Formula XII

- d. reacting the compound of Formula XII with a compound of Formula XIII




Formula XIII

- to give a compound of Formula XIV;




Formula XIV

- e. reacting the compound of Formula XIV with a compound of Formula XV




Formula XV

- to give a compound of Formula XVI;




Formula XVI

- f. hydrogenating a compound of Formula XVI to give a compound of Formula XVII




Formula XVII

- g. deprotecting the compound of Formula XVII to give a compound of Formula XVIII;




Formula XVIII

- h. converting the compound of Formula XVIII to a compound of Formula XIX; and



Formula XIX

- i. deformylating the compound of Formula XIX to give the compound of Formula XX.



Formula XX

wherein:

P₁ is a silyl protecting group;

R' and R'' together form an acetal protecting group;

R₂ is an N-containing heterocyclyl or heteroaryl;

hal is Cl, Br or I;

Ar is phenyl;

Q is optionally substituted aryl or heteroaryl; and

R₁ is hydrogen, optionally substituted alkyl, alkenyl, alkynyl, cycloalkyl, aryl, heterocyclyl, heteroaryl, aralkyl, alkoxy, aryloxy, alkenyloxy or alkynyloxy.

4. The process according to claim 3, wherein compounds of Formula XX are selected from:

(2S,3R)-3-hydroxy-2-[2-(4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-5-(4-pyrimidin-5-ylphenyl)pentanoic acid (Compound No.1),

(2S,3R)-2-[2-(5-*tert*-butyl-1,3-dioxo-1,3-dihydro-2H-isoindol-2-yl)ethyl]-3-hydroxy-5-[4-(6-methoxypyridin-3-yl)phenyl]pentanoic acid (Compound No. 2),

(2S,3R)-2-[2-(1,3-dioxo-1,3-dihydro-2H-isoindol-2-yl)ethyl]-3-hydroxy-5-[4-(6-methylpyridin-3-yl)phenyl]pentanoic acid (Compound No. 3),

(2S,3R)-2-[2-(1,3-dioxo-1,3-dihydro-2H-isoindol-2-yl)ethyl]-5-(4'-fluorobiphenyl-4-yl)-3-hydroxypentanoic acid (Compound No. 4),

(2*S*,3*R*)-5-(3',4'-difluorobiphenyl-4-yl)-2-[2-(1,3-dioxo-1,3-dihydro-2*H*-isoindol-2-yl)ethyl]-3-hydroxypentanoic acid (Compound No. 5),

(2*S*,3*R*)-3-hydroxy-5-[4-(6-methylpyridin-3-yl)phenyl]-2-[2-(4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)ethyl]pentanoic acid (Compound No. 6),

(2*S*,3*R*)-3-hydroxy-2-[2-(5-methyl-1,3-dioxo-1,3-dihydro-2*H*-isoindol-2-yl)ethyl]-5-[4-(6-methylpyridin-3-yl)phenyl]pentanoic acid (Compound No. 7),

(2*S*,3*R*)-5-(4'-fluorobiphenyl-4-yl)-3-hydroxy-2-[2-(5-methyl-1,3-dioxo-1,3-dihydro-2*H*-isoindol-2-yl)ethyl]pentanoic acid (Compound No. 8),

(2*S*,3*R*)-5-(4'-fluorobiphenyl-4-yl)-3-hydroxy-2-[2-(4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)ethyl]pentanoic acid (Compound No. 9),

(2*S*,3*R*)-3-hydroxy-5-(4'-methoxybiphenyl-4-yl)-2-[2-(4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)ethyl]pentanoic acid (Compound No. 10),

(2*S*,3*R*)-5-(3',4'-difluorobiphenyl-4-yl)-3-hydroxy-2-[2-(5-methyl-1,3-dioxo-1,3-dihydro-2*H*-isoindol-2-yl)ethyl]pentanoic acid (Compound No. 11),

(2*S*,3*R*)-5-(3',4'-difluorobiphenyl-4-yl)-3-hydroxy-2-[2-(4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)ethyl]pentanoic acid (Compound No. 12),

(2*S*,3*R*)-2-[2-(1,3-dioxo-1,3-dihydro-2*H*-isoindol-2-yl)ethyl]-3-hydroxy-5-(4'-methoxybiphenyl-4-yl)pentanoic acid (Compound No. 13),

(2*S*,3*R*)-2-[2-(1,3-dioxo-1,3-dihydro-2*H*-isoindol-2-yl)ethyl]-3-hydroxy-5-[4'-(trifluoromethoxy)biphenyl-4-yl]pentanoic acid (Compound No. 14),

(2*S*,3*R*)-5-(4'-chloro-3'-fluorobiphenyl-4-yl)-3-hydroxy-2-[2-(4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)ethyl]pentanoic acid (Compound No. 15),

(2*S*,3*R*)-2-[2-(5-*tert*-butyl-1,3-dioxo-1,3-dihydro-2*H*-isoindol-2-yl)ethyl]-5-(3',4'-difluorobiphenyl-4-yl)-3-hydroxypentanoic acid (Compound No. 16),

(2*S*,3*R*)-2-[2-(5-*tert*-butyl-1,3-dioxo-1,3-dihydro-2*H*-isoindol-2-yl)ethyl]-3-hydroxy-5-(4'-methoxybiphenyl-4-yl)pentanoic acid (Compound No. 17),

(2*S*,3*R*)-3-hydroxy-2-[2-(4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)ethyl]-5-[4'-(trifluoromethyl)biphenyl-4-yl]pentanoic acid (Compound No. 18),

39 (2*S*,3*R*)-2-[2-(5-*tert*-butyl-1,3-dioxo-1,3-dihydro-2*H*-isoindol-2-yl)ethyl]-5-(4'-
40 fluorobiphenyl-4-yl)-3-hydroxypentanoic acid (Compound No. 19),

41 (2*S*,3*R*)-2-[2-(5-*tert*-butyl-1,3-dioxo-1,3-dihydro-2*H*-isoindol-2-yl)ethyl]-5-(4'-
42 chlorobiphenyl-4-yl)-3-hydroxypentanoic acid (Compound No. 20),

43 (2*S*,3*R*)-2-[2-(1,3-dioxo-1,3-dihydro-2*H*-isoindol-2-yl)ethyl]-3-hydroxy-5-[4'-
44 (trifluoromethyl)biphenyl-4-yl]pentanoic acid (Compound No. 21),

45 (2*S*,3*R*)-2-[2-(1,3-dioxo-1,3-dihydro-2*H*-isoindol-2-yl)ethyl]-3-hydroxy-5-(4'-
46 methylbiphenyl-4-yl)pentanoic acid (Compound No. 22),

47 (2*S*,3*R*)-2-[2-(1,3-dioxo-1,3-dihydro-2*H*-isoindol-2-yl)ethyl]-5-(3'-fluoro-4'-
48 methoxybiphenyl-4-yl)-3-hydroxypentanoic acid (Compound No. 23),

49 (2*S*,3*R*)-3-hydroxy-5-[4-(6-methoxypyridin-3-yl)phenyl]-2-[2-(5-methyl-1,3-
50 dioxo-1,3-dihydro-2*H*-isoindol-2-yl)ethyl]pentanoic acid (Compound No. 24),

51 (2*S*,3*R*)-3-hydroxy-5-(4'-methylbiphenyl-4-yl)-2-[2-(4-oxo-1,2,3-benzotriazin-
52 3(4*H*)-yl)ethyl]pentanoic acid (Compound No. 25),

53 (2*S*,3*R*)-5-(4'-ethylbiphenyl-4-yl)-3-hydroxy-2-[2-(4-oxo-1,2,3-benzotriazin-
54 3(4*H*)-yl)ethyl]pentanoic acid (Compound No. 26),

55 (2*S*,3*R*)-5-(4'-chlorobiphenyl-4-yl)-3-hydroxy-2-[2-(5-methyl-1,3-dioxo-1,3-
56 dihydro-2*H*-isoindol-2-yl)ethyl]pentanoic acid (Compound No. 27),

57 (2*S*,3*R*)-5-(4'-chlorobiphenyl-4-yl)-3-hydroxy-2-[2-(4-oxo-1,2,3-benzotriazin-
58 3(4*H*)-yl)ethyl]pentanoic acid (Compound No. 28),

59 (2*S*,3*R*)-2-[2-(1,3-dioxo-1,3-dihydro-2*H*-isoindol-2-yl)ethyl]-3-hydroxy-5-(4-
60 pyrimidin-5-ylphenyl)pentanoic acid (Compound No. 29),

61 (2*S*,3*R*)-2-[2-(1,3-dioxo-1,3-dihydro-2*H*-isoindol-2-yl)ethyl]-3-hydroxy-5-[4-(1-
62 methyl-1*H*-pyrazol-4-yl)phenyl]pentanoic acid (Compound No. 30),

63 (2*S*,3*R*)-2-[2-(1,3-dioxo-1,3-dihydro-2*H*-isoindol-2-yl)ethyl]-3-hydroxy-5-(4-
64 pyridin-3-ylphenyl)pentanoic acid (Compound No. 31),

65 (2*S*,3*R*)-5-[4-(2,3-dihydro-1,4-benzodioxin-6-yl)phenyl]-3-hydroxy-2-[2-(4-oxo-
66 1,2,3-benzotriazin-3(4*H*)-yl)ethyl]pentanoic acid (Compound No. 32),

- 67 (2*S*,3*R*)-3-hydroxy-5-[4-(6-methoxypyridin-3-yl)phenyl]-2-[2-(8-methyl-4-oxo-
68 1,2,3-benzotriazin-3(4*H*)-yl)ethyl]pentanoic acid (Compound No. 33),
- 69 (2*S*,3*R*)-3-hydroxy-2-[2-(8-methyl-4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)ethyl]-5-[4-
70 (6-methylpyridin-3-yl)phenyl]pentanoic acid (Compound No. 34),
- 71 (2*S*,3*R*)-3-hydroxy-5-[4-(1-methyl-1*H*-pyrazol-4-yl)phenyl]-2-[2-(4-oxo-1,2,3-
72 benzotriazin-3(4*H*)-yl)ethyl]pentanoic acid (Compound No. 35),
- 73 (2*S*,3*R*)-3-hydroxy-2-[2-(8-methoxy-4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)ethyl]-5-
74 [4-(6-methoxypyridin-3-yl)phenyl]pentanoic acid (Compound No. 36),
- 75 (2*S*,3*R*)-2-[2-(7-chloro-4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)ethyl]-3-hydroxy-5-[4-
76 (6-methoxypyridin-3-yl)phenyl]pentanoic acid (Compound No. 37),
- 77 (2*S*,3*R*)-3-hydroxy-2-{2-[7-(6-methoxypyridin-3-yl)-4-oxo-1,2,3-benzotriazin-
78 3(4*H*)-yl]ethyl}-5-[4-(6-methoxypyridin-3-yl)phenyl]pentanoic acid (Compound No. 38),
- 79 (2*S*,3*R*)-2-[2-(6,7-difluoro-4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)ethyl]-3-hydroxy-5-
80 [4-(6-methoxypyridin-3-yl)phenyl]pentanoic acid (Compound No. 39),
- 81 (2*S*,3*R*)-2-[2-(5-fluoro-4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)ethyl]-3-hydroxy-5-[4-
82 (6-methoxypyridin-3-yl)phenyl]pentanoic acid (Compound No. 40),
- 83 (2*S*,3*R*)-3-hydroxy-2-[2-(8-methoxy-4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)ethyl]-5-
84 [4-(6-methylpyridin-3-yl)phenyl]pentanoic acid (Compound No. 41),
- 85 (2*S*,3*R*)-3-hydroxy-2-{2-[5-(6-methoxypyridin-3-yl)-4-oxo-1,2,3-benzotriazin-
86 3(4*H*)-yl]ethyl}-5-[4-(6-methoxypyridin-3-yl)phenyl]pentanoic acid (Compound No. 42),
- 87 (2*S*,3*R*)-5-(4'-chloro-3'-fluorobiphenyl-4-yl)-2-[2-(1,3-dioxo-1,3-dihydro-2*H*-
88 isoindol-2-yl)ethyl]-3-hydroxypentanoic acid (Compound No. 43),
- 89 (2*S*,3*R*)-2-[2-(1,3-dioxo-1,3-dihydro-2*H*-isoindol-2-yl)ethyl]-3-hydroxy-5-[4-(1-
90 isobutyl-1*H*-pyrazol-4-yl)phenyl]pentanoic acid (Compound No. 44),
- 91 (2*S*,3*R*)-5-biphenyl-4-yl-2-[2-(1,3-dioxo-1,3-dihydro-2*H*-isoindol-2-yl)ethyl]-3-
92 hydroxypentanoic acid (Compound No. 45),
- 93 (2*S*,3*R*)-2-[2-(1,3-dioxo-1,3-dihydro-2*H*-isoindol-2-yl)ethyl]-5-(3'-fluoro-4'-
94 methylbiphenyl-4-yl)-3-hydroxypentanoic acid (Compound No. 46),

- 95 (2*S*,3*R*)-5-(3,3'-difluoro-4'-methoxybiphenyl-4-yl)-2-[2-(1,3-dioxo-1,3-dihydro-
96 2*H*-isoindol-2-yl)ethyl]-3-hydroxypentanoic acid (Compound No. 47),
- 97 (2*S*,3*R*)-5-[4-(2,3-dihydro-1,4-benzodioxin-6-yl)phenyl]-2-[2-(1,3-dioxo-1,3-
98 dihydro-2*H*-isoindol-2-yl)ethyl]-3-hydroxypentanoic acid (Compound No. 48),
- 99 (2*S*,3*R*)-2-[2-(1,3-dioxo-1,3-dihydro-2*H*-isoindol-2-yl)ethyl]-3-hydroxy-5-[4-(1*H*-
100 tetrazol-1-yl)phenyl]pentanoic acid (Compound No. 49),
- 101 (2*S*,3*R*)-2-[2-(1,3-dioxo-1,3-dihydro-2*H*-isoindol-2-yl)ethyl]-5-[2-fluoro-4-(6-
102 methoxypyridin-3-yl)phenyl]-3-hydroxypentanoic acid (Compound No. 50),
- 103 (2*S*,3*R*)-5-(3'-fluoro-4'-methylbiphenyl-4-yl)-3-hydroxy-2-[2-(4-oxo-1,2,3-
104 benzotriazin-3(4*H*)-yl)ethyl]pentanoic acid (Compound No. 51),
- 105 (2*S*,3*R*)-3-hydroxy-5-[4-(1-isobutyl-1*H*-pyrazol-4-yl)phenyl]-2-[2-(4-oxo-1,2,3-
106 benzotriazin-3(4*H*)-yl)ethyl]pentanoic acid (Compound No. 52),
- 107 (2*S*,3*R*)-5-biphenyl-4-yl-3-hydroxy-2-[2-(4-oxo-1,2,3-benzotriazin-3(4*H*)-
108 yl)ethyl]pentanoic acid (Compound No. 53),
- 109 (2*S*,3*R*)-3-hydroxy-5-[4-(6-methoxypyridin-3-yl)phenyl]-2-[2-(7-methyl-4-oxo-
110 1,2,3-benzotriazin-3(4*H*)-yl)ethyl]pentanoic acid (Compound No. 54),
- 111 (2*S*,3*R*)-2-[2-(1,3-dioxo-1,3-dihydro-2*H*-isoindol-2-yl)ethyl]-5-[6-(3-fluoro-4-
112 methoxyphenyl)pyridin-3-yl]-3-hydroxypentanoic acid (Compound No. 55),
- 113 (2*S*,3*R*)-5-(4'-chloro-3-fluorobiphenyl-4-yl)-2-[2-(1,3-dioxo-1,3-dihydro-2*H*-
114 isoindol-2-yl)ethyl]-3-hydroxypentanoic acid (Compound No. 56),
- 115 (2*S*,3*R*)-2-[2-(1,3-dioxo-1,3-dihydro-2*H*-isoindol-2-yl)ethyl]-3-hydroxy-5-[6-(4-
116 methoxyphenyl)pyridin-3-yl]pentanoic acid (Compound No. 57),
- 117 (2*S*,3*R*)-2-[2-(1,3-dioxo-1,3-dihydro-2*H*-isoindol-2-yl)ethyl]-5-(3-fluoro-4'-
118 methoxybiphenyl-4-yl)-3-hydroxypentanoic acid (Compound No. 58),
- 119 (2*S*,3*R*)-5-[6-(4-chlorophenyl)pyridin-3-yl]-2-[2-(1,3-dioxo-1,3-dihydro-2*H*-
120 isoindol-2-yl)ethyl]-3-hydroxypentanoic acid (Compound No. 59),
- 121 (2*S*,3*R*)-2-[2-(1,3-dioxo-1,3-dihydro-2*H*-isoindol-2-yl)ethyl]-3-hydroxy-5-(6'-
122 methoxy-2,3'-bipyridin-5-yl)pentanoic acid (Compound No. 60),

- 123 (2*S*,3*R*)-5-[4-(4-chlorophenyl)-2-thienyl]-2-[2-(1,3-dioxo-1,3-dihydro-2*H*-isoindol-
124 2-yl)ethyl]-3-hydroxypentanoic acid (Compound No. 61),
- 125 (2*S*,3*R*)-2-[2-(1,3-dioxo-1,3-dihydro-2*H*-isoindol-2-yl)ethyl]-3-hydroxy-5-[4-(6-
126 methoxypyridin-3-yl)-2-thienyl]pentanoic acid (Compound No. 62),
- 127 (2*S*,3*R*)-2-[2-(1,3-dioxo-1,3-dihydro-2*H*-isoindol-2-yl)ethyl]-3-hydroxy-5-{4-[4-
128 (trifluoromethyl)phenyl]-2-thienyl}pentanoic acid (Compound No. 63),
- 129 (2*S*,3*R*)-2-[2-(1,3-dioxo-1,3-dihydro-2*H*-isoindol-2-yl)ethyl]-5-[4-(3-fluoro-4-
130 methoxyphenyl)-2-thienyl]-3-hydroxypentanoic acid (Compound No. 64),
- 131 (2*S*,3*R*)-3-hydroxy-2-[2-(7-methyl-4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)ethyl]-5-[4'-
132 (trifluoromethyl)biphenyl-4-yl]pentanoic acid (Compound No. 65),
- 133 (2*S*,3*R*)-3-hydroxy-2-[2-(6-methyl-4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)ethyl]-5-[4'-
134 (trifluoromethyl)biphenyl-4-yl]pentanoic acid (Compound No. 66),
- 135 (2*S*,3*R*)-2-[2-(7-chloro-4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)ethyl]-3-hydroxy-5-(4'-
136 methylbiphenyl-4-yl)pentanoic acid (Compound No. 67),
- 137 (2*S*,3*R*)-3-hydroxy-2-[2-(8-methoxy-4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)ethyl]-5-
138 [4'-(trifluoromethyl)biphenyl-4-yl]pentanoic acid (Compound No. 68),
- 139 (2*S*,3*R*)-2-[2-(6,7-difluoro-4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)ethyl]-3-hydroxy-5-
140 [4'-(trifluoromethyl)biphenyl-4-yl]pentanoic acid (Compound No. 69),
- 141 (2*S*,3*R*)-2-[2-(1,3-dioxo-1,3-dihydro-2*H*-isoindol-2-yl)ethyl]-5-[2-fluoro-5-(6-
142 methoxypyridin-3-yl)phenyl]-3-hydroxypentanoic acid (Compound No. 70),
- 143 (2*S*,3*R*)-5-(4'-chloro-4-fluorobiphenyl-3-yl)-2-[2-(1,3-dioxo-1,3-dihydro-2*H*-
144 isoindol-2-yl)ethyl]-3-hydroxypentanoic acid (Compound No. 71),
- 145 (2*S*,3*R*)-2-[2-(1,3-dioxo-1,3-dihydro-2*H*-isoindol-2-yl)ethyl]-5-[6-fluoro-4'-
146 (trifluoromethyl)biphenyl-3-yl]-3-hydroxypentanoic acid (Compound No. 72),
- 147 (2*S*,3*R*)-2-[2-(1,3-dioxo-1,3-dihydro-2*H*-isoindol-2-yl)ethyl]-5-[4-fluoro-3-(6-
148 methoxypyridin-3-yl)phenyl]-3-hydroxypentanoic acid (Compound No. 73),
- 149 (2*S*,3*R*)-5-(4'-chloro-6-fluorobiphenyl-3-yl)-2-[2-(1,3-dioxo-1,3-dihydro-2*H*-
150 isoindol-2-yl)ethyl]-3-hydroxypentanoic acid (Compound No. 74),

- 151 (2*S*,3*R*)-5-(3',6-difluoro-4'-methoxybiphenyl-3-yl)-2-[2-(1,3-dioxo-1,3-dihydro-
152 2*H*-isoindol-2-yl)ethyl]-3-hydroxypentanoic acid (Compound No. 75),
- 153 (2*S*,3*R*)-2-[2-(5-fluoro-4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)ethyl]-3-hydroxy-5-(4'-
154 methoxybiphenyl-4-yl)pentanoic acid (Compound No. 76),
- 155 (2*S*,3*R*)-5-(4'-fluorobiphenyl-4-yl)-2-[2-(5-fluoro-4-oxo-1,2,3-benzotriazin-3(4*H*)-
156 yl)ethyl]-3-hydroxypentanoic acid (Compound No. 77),
- 157 (2*S*,3*R*)-3-hydroxy-2-[2-(6-methyl-4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)ethyl]-5-[4-
158 (1-methyl-1*H*-pyrazol-4-yl)phenyl]pentanoic acid (Compound No. 78),
- 159 (2*S*,3*R*)-2-[2-(5-chloro-1,3-dioxo-1,3-dihydro-2*H*-isoindol-2-yl)ethyl]-3-hydroxy-
160 5-[4'-(trifluoromethyl)biphenyl-4-yl]pentanoic acid (Compound No. 79),
- 161 (2*S*,3*R*)-2-[2-(4-fluoro-1,3-dioxo-1,3-dihydro-2*H*-isoindol-2-yl)ethyl]-3-hydroxy-
162 5-[4'-(trifluoromethyl)biphenyl-4-yl]pentanoic acid (Compound No. 80),
- 163 (2*S*,3*R*)-5-(3',4'-difluorobiphenyl-4-yl)-2-[2-(5-fluoro-4-oxo-1,2,3-benzotriazin-
164 3(4*H*)-yl)ethyl]-3-hydroxypentanoic acid (Compound No. 81),
- 165 (2*S*,3*R*)-3-hydroxy-5-[4-(6-methoxypyridin-3-yl)phenyl]-2-[2-(4-oxo-1,2,3-
166 benzotriazin-3(4*H*)-yl)ethyl]pentanoic acid (Compound No. 86),
- 167 (2*S*,3*R*)-5-(4'-chlorobiphenyl-4-yl)-2-[2-(1,3-dioxo-1,3-dihydro-2*H*-isoindol-2-
168 yl)ethyl]-3-hydroxypentanoic acid (Compound No. 87),
- 169 (2*S*,3*R*)-2-[2-(1,3-dioxo-1,3-dihydro-2*H*-isoindol-2-yl)ethyl]-3-hydroxy-5-[4-(6-
170 methoxypyridin-3-yl)phenyl]pentanoic acid (Compound No.88),
- 171 (2*S*,3*R*)-3-hydroxy-2-[2-(7-methyl-4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)ethyl]-5-[4-
172 (1-methyl-1*H*-pyrazol-4-yl)phenyl]pentanoic acid (Compound No. 90),
- 173 (2*S*,3*R*)-3-hydroxy-2-[2-(8-methoxy-4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)ethyl]-5-
174 [4-(1-methyl-1*H*-pyrazol-4-yl)phenyl]pentanoic acid (Compound No. 91),
- 175 (2*S*,3*R*)-2-[2-(7-chloro-4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)ethyl]-5-(3',4'-
176 difluorobiphenyl-4-yl)-3-hydroxypentanoic acid (Compound No. 92),
- 177 (2*S*,3*R*)-2-[2-(7-chloro-4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)ethyl]-3-hydroxy-5-(4'-
178 methoxybiphenyl-4-yl)pentanoic acid (Compound No. 93),

- 179 (2S,3R)-2-[2-(7-chloro-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-5-(4'-
180 fluorobiphenyl-4-yl)-3-hydroxypentanoic acid (Compound No. 94),
- 181 (2S,3R)-2-[2-(7-chloro-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-3-hydroxy-5-[4-
182 (6-methylpyridin-3-yl)phenyl]pentanoic acid (Compound No. 95),
- 183 (2S,3R)-5-(2',4'-difluorobiphenyl-4-yl)-2-[2-(1,3-dioxo-1,3-dihydro-2H-isoindol-2-
184 yl)ethyl]-3-hydroxypentanoic acid (Compound No. 97),
- 185 (2S,3R)-2-[2-(6-chloro-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-3-hydroxy-5-[4-
186 (1-methyl-1H-pyrazol-4-yl)phenyl]pentanoic acid (Compound No. 98),
- 187 (2S,3R)-2-[2-(5-chloro-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-3-hydroxy-5-[4-
188 (1-methyl-1H-pyrazol-4-yl)phenyl]pentanoic acid (Compound No. 99),
- 189 (2S,3R)-2-[2-(1,3-dioxo-1,3-dihydro-2H-isoindol-2-yl)ethyl]-3-hydroxy-5-(4'-
190 isopropylbiphenyl-4-yl)pentanoic acid (Compound No. 100),
- 191 (2S,3R)-5-(3'-chloro-4'-fluorobiphenyl-4-yl)-2-[2-(1,3-dioxo-1,3-dihydro-2H-
192 isoindol-2-yl)ethyl]-3-hydroxypentanoic acid (Compound No. 101),
- 193 (2S,3R)-5-(4'-butylbiphenyl-4-yl)-3-hydroxy-2-[2-(4-oxo-1,2,3-benzotriazin-
194 3(4H)-yl)ethyl]pentanoic acid (Compound No. 102),
- 195 (2S,3R)-5-(2'-fluorobiphenyl-4-yl)-3-hydroxy-2-[2-(4-oxo-1,2,3-benzotriazin-
196 3(4H)-yl)ethyl]pentanoic acid (Compound No. 103),
- 197 (2S,3R)-2-[2-(5-fluoro-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-3-hydroxy-5-[4-
198 (1-methyl-1H-pyrazol-4-yl)phenyl]pentanoic acid (Compound No. 104),
- 199 (2S,3R)-3-hydroxy-2-[2-(8-methyl-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-5-[4-
200 (1-methyl-1H-pyrazol-4-yl)phenyl]pentanoic acid (Compound No. 105),
- 201 (2S,3R)-2-[2-(1,3-dioxo-1,3-dihydro-2H-isoindol-2-yl)ethyl]-3-hydroxy-5-{6-[4-
202 (trifluoromethyl)phenyl]pyridin-3-yl}pentanoic acid (Compound No. 106),
- 203 (2S,3R)-5-[6-(3,4-difluorophenyl)pyridin-3-yl]-2-[2-(1,3-dioxo-1,3-dihydro-2H-
204 isoindol-2-yl)ethyl]-3-hydroxypentanoic acid (Compound No. 107),
- 205 (2S,3R)-5-[6-(4-chloro-3-fluorophenyl)pyridin-3-yl]-2-[2-(1,3-dioxo-1,3-dihydro-
206 2H-isoindol-2-yl)ethyl]-3-hydroxypentanoic acid (Compound No. 108),

- 207 (2S,3R)-2-[2-(1,3-dioxo-1,3-dihydro-2H-isoindol-2-yl)ethyl]-5-[6-(4-
208 fluorophenyl)pyridin-3-yl]-3-hydroxypentanoic acid (Compound No. 109),
209 (2S,3R)-2-[2-(1,3-dioxo-1,3-dihydro-2H-isoindol-2-yl)ethyl]-5-[6-(3-fluoro-4-
210 methylphenyl)pyridin-3-yl]-3-hydroxypentanoic acid (Compound No. 110),
211 (2S,3R)-2-[2-(7-chloro-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-3-hydroxy-5-[4-
212 (1-methyl-1H-pyrazol-4-yl)phenyl]pentanoic acid (Compound No. 111),
213 (2S,3R)-2-[2-(8-chloro-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-3-hydroxy-5-[4-
214 (1-methyl-1H-pyrazol-4-yl)phenyl]pentanoic acid (Compound No. 112),
215 (2S,3R)-2-[2-(6,7-difluoro-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-3-hydroxy-5-
216 [4-(1-methyl-1H-pyrazol-4-yl)phenyl]pentanoic acid (Compound No. 113),
217 (2S,3R)-3-hydroxy-2-[2-(7-methyl-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-5-[4'-
218 (trifluoromethoxy)biphenyl-4-yl]pentanoic acid (Compound No. 114),
219 (2S,3R)-3-hydroxy-5-(4'-methylbiphenyl-4-yl)-2-[2-(7-methyl-4-oxo-1,2,3-
220 benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 115),
221 (2S,3R)-5-(4'-fluorobiphenyl-4-yl)-3-hydroxy-2-[2-(7-methyl-4-oxo-1,2,3-
222 benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 116),
223 (2S,3R)-5-(3',4'-difluorobiphenyl-4-yl)-3-hydroxy-2-[2-(7-methyl-4-oxo-1,2,3-
224 benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 117),
225 (2S,3R)-3-hydroxy-2-[2-(7-methyl-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-5-[4-
226 (6-methylpyridin-3-yl)phenyl]pentanoic acid (Compound No. 118),
227 (2S,3R)-5-[4-(2,3-dihydro-1,4-benzodioxin-6-yl)phenyl]-3-hydroxy-2-[2-(7-
228 methyl-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 119),
229 (2S,3R)-2-[2-(5-chloro-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-3-hydroxy-5-(6'-
230 methoxy-2,3'-bipyridin-5-yl)pentanoic acid (Compound No. 121),
231 (2S,3R)-2-[2-(7-chloro-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-3-hydroxy-5-(6'-
232 methoxy-2,3'-bipyridin-5-yl)pentanoic acid (Compound No. 122),
233 (2S,3R)-2-[2-(6,7-difluoro-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-3-hydroxy-5-
234 (6'-methoxy-2,3'-bipyridin-5-yl)pentanoic acid (Compound No. 123),

- 235 (2S,3R)-5-(3'-fluoro-4'-methylbiphenyl-4-yl)-3-hydroxy-2-[2-(7-methyl-4-oxo-
236 1,2,3-benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 124),
- 237 (2S,3R)-3-hydroxy-5-[4-(1-isobutyl-1H-pyrazol-4-yl)phenyl]-2-[2-(7-methyl-4-
238 oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 125),
- 239 (2S,3R)-2-[2-(5-fluoro-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-3-hydroxy-5-(6'-
240 methoxy-2,3'-bipyridin-5-yl)pentanoic acid (Compound No. 126),
- 241 (2S,3R)-5-(4'-fluorobiphenyl-4-yl)-3-hydroxy-2-[2-(8-methoxy-4-oxo-1,2,3-
242 benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 127),
- 243 (2S,3R)-5-(3',4'-difluorobiphenyl-4-yl)-3-hydroxy-2-[2-(8-methoxy-4-oxo-1,2,3-
244 benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 128),
- 245 (2S,3R)-3-hydroxy-5-(4'-methoxybiphenyl-4-yl)-2-[2-(8-methoxy-4-oxo-1,2,3-
246 benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 129),
- 247 (2S,3R)-2-[2-(7-chloro-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-3-hydroxy-5-{6-
248 [4-(trifluoromethyl)phenyl]pyridin-3-yl}pentanoic acid (Compound No. 130),
- 249 (2S,3R)-2-[2-(6,7-difluoro-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-3-hydroxy-5-
250 {6-[4-(trifluoromethyl)phenyl]pyridin-3-yl}pentanoic acid (Compound No. 131),
- 251 (2S,3R)-2-[2-(5-chloro-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-3-hydroxy-5-{6-
252 [4-(trifluoromethyl)phenyl]pyridin-3-yl}pentanoic acid (Compound No. 132),
- 253 (2S,3R)-3-hydroxy-2-[2-(7-methyl-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-5-{6-
254 [4-(trifluoromethyl)phenyl]pyridin-3-yl}pentanoic acid (Compound No. 133),
- 255 (2S,3R)-2-[2-(5-fluoro-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-3-hydroxy-5-{6-
256 [4-(trifluoromethyl)phenyl]pyridin-3-yl}pentanoic acid (Compound No. 134),
- 257 (2S,3R)-3-hydroxy-2-[2-(8-methoxy-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-5-
258 (4'-methylbiphenyl-4-yl)pentanoic acid (Compound No. 135),
- 259 (2S,3R)-5-(3'-fluoro-4'-methylbiphenyl-4-yl)-2-[2-(5-fluoro-4-oxo-1,2,3-
260 benzotriazin-3(4H)-yl)ethyl]-3-hydroxypentanoic acid (Compound No. 136),
- 261 (2S,3R)-2-[2-(6,7-difluoro-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-3-hydroxy-5-
262 (4'-methylbiphenyl-4-yl)pentanoic acid (Compound No. 137),

- 263 (2S,3R)-2-[2-(5-fluoro-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-3-hydroxy-5-(4'-
264 methylbiphenyl-4-yl)pentanoic acid (Compound No. 138),
- 265 (2S,3R)-5-(4'-chlorobiphenyl-4-yl)-2-[2-(5-fluoro-4-oxo-1,2,3-benzotriazin-3(4H)-
266 yl)ethyl]-3-hydroxypentanoic acid (Compound No. 139),
- 267 (2S,3R)-3-hydroxy-5-(6'-methoxy-2,3'-bipyridin-5-yl)-2-[2-(4-oxo-1,2,3-
268 benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 140),
- 269 (2S,3R)-3-hydroxy-5-(6'-methoxy-2,3'-bipyridin-5-yl)-2-[2-(6-methyl-4-oxo-1,2,3-
270 benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 141),
- 271 (2S,3R)-3-hydroxy-5-(6'-methoxy-2,3'-bipyridin-5-yl)-2-[2-(7-methyl-4-oxo-1,2,3-
272 benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 142),
- 273 (2S,3R)-2-[2-(6,7-difluoro-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-5-(3'-fluoro-
274 4'-methylbiphenyl-4-yl)-3-hydroxypentanoic acid (Compound No. 144),
- 275 (2S,3R)-5-(4'-chlorobiphenyl-4-yl)-2-[2-(6,7-difluoro-4-oxo-1,2,3-benzotriazin-
276 3(4H)-yl)ethyl]-3-hydroxypentanoic acid (Compound No. 145),
- 277 (2S,3R)-5-(3'-fluoro-4'-methylbiphenyl-4-yl)-3-hydroxy-2-[2-(8-methoxy-4-oxo-
278 1,2,3-benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 146),
- 279 (2S,3R)-5-[4-(2,3-dihydro-1,4-benzodioxin-6-yl)phenyl]-3-hydroxy-2-[2-(8-
280 methoxy-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 147),
- 281 (2S,3R)-5-[4-(6-chloropyridin-3-yl)phenyl]-3-hydroxy-2-[2-(4-oxo-1,2,3-
282 benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 148),
- 283 (2S,3R)-3-hydroxy-2-[2-(6-methoxy-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-5-
284 [4'-(trifluoromethyl)biphenyl-4-yl]pentanoic acid (Compound No. 149),
- 285 (2S,3R)-3-hydroxy-2-[2-(7-methoxy-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-5-
286 [4'-(trifluoromethyl)biphenyl-4-yl]pentanoic acid (Compound No. 150),
- 287 (2S,3R)-3-hydroxy-2-[2-(6-methoxy-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-5-
288 (4'-methylbiphenyl-4-yl)pentanoic acid (Compound No. 151),
- 289 (2S,3R)-3-hydroxy-5-(4'-methoxybiphenyl-4-yl)-2-[2-(6-methoxy-4-oxo-1,2,3-
290 benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 152),

- 291 (2S,3R)-5-[2-fluoro-4-(6-methoxypyridin-3-yl)phenyl]-3-hydroxy-2-[2-(4-oxo-
292 1,2,3-benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 153),
293 (2S,3R)-5-[2-fluoro-4-(6-methoxypyridin-3-yl)phenyl]-3-hydroxy-2-[2-(7-
294 methoxy-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 154),
295 (2S,3R)-5-[2-fluoro-4-(6-methoxypyridin-3-yl)phenyl]-2-[2-(5-fluoro-4-oxo-1,2,3-
296 benzotriazin-3(4H)-yl)ethyl]-3-hydroxypentanoic acid (Compound No. 155),
297 (2S,3R)-5-[2-fluoro-4-(6-methoxypyridin-3-yl)phenyl]-3-hydroxy-2-[2-(7-methyl-
298 4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 156),
299 (2S,3R)-5-[2-fluoro-4-(6-methoxypyridin-3-yl)phenyl]-3-hydroxy-2-[2-(6-methyl-
300 4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 157),
301 (2S,3R)-3-hydroxy-5-(4'-methoxybiphenyl-4-yl)-2-[2-(7-methoxy-4-oxo-1,2,3-
302 benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 158),
303 (2S,3R)-5-(4'-chlorobiphenyl-4-yl)-3-hydroxy-2-[2-(6-methoxy-4-oxo-1,2,3-
304 benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 159),
305 (2S,3R)-3-hydroxy-2-[2-(7-methoxy-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-5-
306 [4-(6-methoxypyridin-3-yl)phenyl]pentanoic acid (Compound No. 160),
307 (2S,3R)-5-(3',4'-difluorobiphenyl-4-yl)-3-hydroxy-2-[2-(7-methoxy-4-oxo-1,2,3-
308 benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 161),
309 (2S,3R)-3-hydroxy-2-[2-(6-methoxy-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-5-
310 [4-(6-methoxypyridin-3-yl)phenyl]pentanoic acid (Compound No. 162),
311 (2S,3R)-2-[2-(5-chloro-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-5-[2-fluoro-4-(6-
312 methoxypyridin-3-yl)phenyl]-3-hydroxypentanoic acid (Compound No. 163),
313 (2S,3R)-2-[2-(6,7-difluoro-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-5-[2-fluoro-4-
314 (6-methoxypyridin-3-yl)phenyl]-3-hydroxypentanoic acid (Compound No. 164),
315 (2S,3R)-5-[2-fluoro-4-(6-methoxypyridin-3-yl)phenyl]-3-hydroxy-2-[2-(5-methyl-
316 4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 165),
317 (2S,3R)-5-(4'-fluorobiphenyl-4-yl)-3-hydroxy-2-[2-(5-methyl-4-oxo-1,2,3-
318 benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 166),

- 319 (2S,3R)-3-hydroxy-5-[4-(6-methoxypyridin-3-yl)phenyl]-2-[2-(5-methyl-4-oxo-
320 1,2,3-benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 167),
- 321 (2S,3R)-3-hydroxy-5-(4'-methoxybiphenyl-4-yl)-2-[2-(5-methyl-4-oxo-1,2,3-
322 benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 168),
- 323 (2S,3R)-3-hydroxy-5-(4'-methylbiphenyl-4-yl)-2-[2-(5-methyl-4-oxo-1,2,3-
324 benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 169),
- 325 (2S,3R)-5-(4'-chlorobiphenyl-4-yl)-3-hydroxy-2-[2-(5-methyl-4-oxo-1,2,3-
326 benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 170),
- 327 (2S,3R)-5-(4'-fluorobiphenyl-4-yl)-3-hydroxy-2-[2-(6-methoxy-4-oxo-1,2,3-
328 benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 171),
- 329 (2S,3R)-5-(3',4'-difluorobiphenyl-4-yl)-3-hydroxy-2-[2-(6-methoxy-4-oxo-1,2,3-
330 benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 172),
- 331 (2S,3R)-3-hydroxy-2-[2-(6-methoxy-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-5-
332 [4'-(trifluoromethoxy)biphenyl-4-yl]pentanoic acid (Compound No. 173),
- 333 (2S,3R)-5-(3'-fluoro-4'-methylbiphenyl-4-yl)-3-hydroxy-2-[2-(6-methoxy-4-oxo-
334 1,2,3-benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 174),
- 335 (2S,3R)-2-[2-(7-chloro-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-5-[2-fluoro-4-(6-
336 methoxypyridin-3-yl)phenyl]-3-hydroxypentanoic acid (Compound No. 175),
- 337 (2S,3R)-3-hydroxy-5-[4-(1-methyl-1H-pyrazol-4-yl)-2-thienyl]-2-[2-(4-oxo-1,2,3-
338 benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 176),
- 339 (2S,3R)-5-[2-fluoro-4-(6-methoxypyridin-3-yl)phenyl]-3-hydroxy-2-[2-(6-
340 methoxy-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 177),
- 341 (2S,3R)-3-hydroxy-5-(6'-methoxy-2,3'-bipyridin-5-yl)-2-[2-(6-methoxy-4-oxo-
342 1,2,3-benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 178),
- 343 (2S,3R)-3-hydroxy-2-[2-(6-methoxy-4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-5-
344 [4-(6-methylpyridin-3-yl)phenyl]pentanoic acid (Compound No. 179),
- 345 (2S,3R)-5-[4-(2-chloropyridin-3-yl)phenyl]-3-hydroxy-2-[2-(6-methoxy-4-oxo-
346 1,2,3-benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 180),

- 347 (2*S*,3*R*)-2-[2-(6-fluoro-4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)ethyl]-3-hydroxy-5-[4-
348 (6-methoxypyridin-3-yl)phenyl]pentanoic acid (Compound No. 181),
- 349 (2*S*,3*R*)-5-(4'-chlorobiphenyl-4-yl)-2-[2-(6-fluoro-4-oxo-1,2,3-benzotriazin-3(4*H*)-
350 yl)ethyl]-3-hydroxypentanoic acid (Compound No. 182),
- 351 (2*S*,3*R*)-5-(3',4'-difluorobiphenyl-4-yl)-2-[2-(6-fluoro-4-oxo-1,2,3-benzotriazin-
352 3(4*H*)-yl)ethyl]-3-hydroxypentanoic acid (Compound No. 183),
- 353 (2*S*,3*R*)-2-[2-(6-fluoro-4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)ethyl]-3-hydroxy-5-(4'-
354 methoxybiphenyl-4-yl)pentanoic acid (Compound No. 184),
- 355 (2*S*,3*R*)-3-hydroxy-5-[4-(2-methoxypyrimidin-5-yl)phenyl]-2-[2-(4-oxo-1,2,3-
356 benzotriazin-3(4*H*)-yl)ethyl]pentanoic acid (Compound No. 185),
- 357 (2*S*,3*R*)-5-[2-fluoro-4-(1-methyl-1*H*-pyrazol-4-yl)phenyl]-3-hydroxy-2-[2-(6-
358 methyl-4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)ethyl]pentanoic acid (Compound No. 186),
- 359 (2*S*,3*R*)-5-[2-fluoro-4-(1-methyl-1*H*-pyrazol-4-yl)phenyl]-3-hydroxy-2-[2-(7-
360 methyl-4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)ethyl]pentanoic acid (Compound No. 187),
- 361 (2*S*,3*R*)-2-[2-(5-chloro-4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)ethyl]-5-[2-fluoro-4-(1-
362 methyl-1*H*-pyrazol-4-yl)phenyl]-3-hydroxypentanoic acid (Compound No. 188),
- 363 (2*S*,3*R*)-2-[2-(7-chloro-4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)ethyl]-5-[2-fluoro-4-(1-
364 methyl-1*H*-pyrazol-4-yl)phenyl]-3-hydroxypentanoic acid (Compound No. 189),
- 365 (2*S*,3*R*)-5-[2-fluoro-4-(1-methyl-1*H*-pyrazol-4-yl)phenyl]-2-[2-(5-fluoro-4-oxo-
366 1,2,3-benzotriazin-3(4*H*)-yl)ethyl]-3-hydroxypentanoic acid (Compound No. 190),
- 367 (2*S*,3*R*)-2-[2-(6,7-difluoro-4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)ethyl]-5-[2-fluoro-4-
368 (1-methyl-1*H*-pyrazol-4-yl)phenyl]-3-hydroxypentanoic acid (Compound No. 191),
- 369 (2*S*,3*R*)-5-[2-fluoro-4-(1-methyl-1*H*-pyrazol-4-yl)phenyl]-2-[2-(6-fluoro-4-oxo-
370 1,2,3-benzotriazin-3(4*H*)-yl)ethyl]-3-hydroxypentanoic acid (Compound No. 192),
- 371 (2*S*,3*R*)-5-[2-fluoro-4-(1-methyl-1*H*-pyrazol-4-yl)phenyl]-3-hydroxy-2-[2-(8-
372 methyl-4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)ethyl]pentanoic acid (Compound No. 193),
- 373 (2*S*,3*R*)-3-hydroxy-2-[2-(6-methoxy-4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)ethyl]-5-
374 [4-(2-methoxypyrimidin-5-yl)phenyl]pentanoic acid (Compound No. 194),

- 375 (2*S*,3*R*)-2-[2-(1,3-dioxo-1,3-dihydro-2*H*-isoindol-2-yl)ethyl]-3-hydroxy-5-[4-(2-
376 methoxypyrimidin-5-yl)phenyl]pentanoic acid (Compound No. 195),
- 377 (2*S*,3*R*)-5-[2-fluoro-4-(1-methyl-1*H*-pyrazol-4-yl)phenyl]-3-hydroxy-2-[2-(1-
378 oxophthalazin-2(1*H*)-yl)ethyl]pentanoic acid (Compound No. 196),
- 379 (2*S*,3*R*)-5-[2-fluoro-4-(1-methyl-1*H*-pyrazol-4-yl)phenyl]-3-hydroxy-2-[2-(3-
380 methyl-2,6-dioxo-3,6-dihydropyrimidin-1(2*H*)-yl)ethyl]pentanoic acid (Compound No.
381 197),
- 382 (2*S*,3*R*)-3-hydroxy-5-[4-(6-methoxypyridin-3-yl)phenyl]-2-{2-[4-oxo-7-
383 (trifluoromethyl)-1,2,3-benzotriazin-3(4*H*)-yl]ethyl}pentanoic acid (Compound No. 198),
- 384 (2*S*,3*R*)-3-hydroxy-5-[4-(6-methoxypyridin-3-yl)phenyl]-2-[2-(1-oxophthalazin-
385 2(1*H*)-yl)ethyl]pentanoic acid (Compound No. 199),
- 386 (2*S*,3*R*)-3-hydroxy-5-[4-(6-methoxypyridin-3-yl)phenyl]-2-[2-(3-methyl-2,6-
387 dioxo-3,6-dihydropyrimidin-1(2*H*)-yl)ethyl]pentanoic acid (Compound No. 200),
- 388 (2*S*,3*R*)-2-[2-(7,9-dioxo-8-azaspiro[4.5]dec-8-yl)ethyl]-3-hydroxy-5-[4-(6-
389 methoxypyridin-3-yl)phenyl]pentanoic acid (Compound No. 201),
- 390 (2*S*,3*R*)-2-[2-(2,4-dioxo-2*H*-1,3-benzoxazin-3(4*H*)-yl)ethyl]-3-hydroxy-5-[4-(6-
391 methoxypyridin-3-yl)phenyl]pentanoic acid (Compound No. 202),
- 392 (2*S*,3*R*)-3-hydroxy-5-[4-(6-methoxypyridin-3-yl)phenyl]-2-[2-(3,4,4-trimethyl-2,5-
393 dioxoimidazolidin-1-yl)ethyl]pentanoic acid (Compound No. 203),
- 394 (2*S*,3*R*)-2-[2-(5-fluoro-4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)ethyl]-3-hydroxy-5-[4'-
395 (trifluoromethyl)biphenyl-4-yl]pentanoic acid (Compound No. 211),
- 396 (2*S*,3*R*)-5-(4'-chlorobiphenyl-4-yl)-3-hydroxy-2-[2-(7-methyl-4-oxo-1,2,3-
397 benzotriazin-3(4*H*)-yl)ethyl]pentanoic acid (Compound No. 212),
- 398 (2*S*,3*R*)-3-hydroxy-5-(6'-methoxy-2,3'-bipyridin-5-yl)-2-[2-(8-methyl-4-oxo-1,2,3-
399 benzotriazin-3(4*H*)-yl)ethyl]pentanoic acid (Compound No. 213),
- 400 (2*S*,3*R*)-3-hydroxy-5-[4-(6-hydroxypyridin-3-yl)phenyl]-2-[2-(4-oxo-1,2,3-
401 benzotriazin-3(4*H*)-yl)ethyl]pentanoic acid (Compound No. 226),
- 402 (2*S*,3*R*)-3-hydroxy-5-(4'-methylbiphenyl-4-yl)-2-{2-[4-oxo-7-(trifluoromethyl)-
403 1,2,3-benzotriazin-3(4*H*)-yl]ethyl}pentanoic acid (Compound No. 227),

404 (2*S*,3*R*)-2-[2-(2,4-dioxo-1,4-dihydroquinazolin-3(2*H*)-yl)ethyl]-3-hydroxy-5-[4-
405 (6-methoxypyridin-3-yl)phenyl]pentanoic acid (Compound No. 228),

406 (2*S*,3*R*)-3-(acetyloxy)-5-[4-(6-methoxypyridin-3-yl)phenyl]-2-[2-(4-oxo-1,2,3-
407 benzotriazin-3(4*H*)-yl)ethyl]pentanoic acid (Compound No. 229),


408 (2*S*,3*R*)-2-[2-(8-chloro-4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)ethyl]-3-hydroxy-5-[4-
409 (6-methoxypyridin-3-yl)phenyl]pentanoic acid(Compound No. 230),

410 (2*S*,3*R*)-5-(4'-chlorobiphenyl-4-yl)-2-[2-(2,4-dioxo-2*H*-1,3-benzoxazin-3(4*H*)-
411 yl)ethyl]-3-hydroxypentanoic acid (Compound No. 231), and


412 (2*S*,3*R*)-5-(4'-chlorobiphenyl-4-yl)-3-hydroxy-2-(2-[(2-hydroxyphenyl)
413 carbonyl]amino}ethyl) pentanoic acid (Compound No. 232) .

1 5. A process for preparing a compound of Formula XXVI comprising:


2 a. reacting a compound of Formula XII with a compound of Formula XXI



4 to give a compound of Formula XXII;




6 b. hydrogenating the compound of Formula XXII to give a compound of
7 Formula XXIII;




Formula XXIII

- c. deprotecting the compound of Formula XXIII to give a compound of Formula XXIV;




Formula XXIV

- d. converting the compound of Formula XXIV to a compound of Formula XXV; and



Formula XXV

- e. deformylating the compound of Formula XXV to give the compound of Formula XXVI,



Formula XXVI

wherein

R' and R'' together form an acetal protecting group;

R₂ is an N-containing heterocyclcyl or heteroaryl;

Ar is phenyl; and

Q is optionally substituted aryl or heteroaryl.

6. The process according to claim 5, wherein compounds of Formula XXVI are selected from:

(2S,3R)-3-hydroxy-2-[2-(4-oxo-1,2,3-benzotriazin-3(4H)-yl)ethyl]-5-phenylpentanoic acid (Compound No. 82),

(2S,3R)-2-[2-(1,3-dioxo-1,3-dihydro-2H-isoindol-2-yl)ethyl]-3-hydroxy-5-phenylpentanoic acid (Compound No. 83),


(2S,3R)-2-[2-(1,3-dioxo-1,3-dihydro-2H-isoindol-2-yl)ethyl]-3-hydroxy-5-[4(trifluoromethyl)phenyl]pentanoic acid (Compound No. 84),

(2S,3R)-5-(4-*tert*-butylphenyl)-2-[2-(1,3-dioxo-1,3-dihydro-2H-isoindol-2-yl)ethyl]-3-hydroxypentanoic acid (Compound No.85), and


(2S,3R)-5-(4-chloro-3-fluorophenyl)-2-[2-(1,3-dioxo-1,3-dihydro-2H-isoindol-2-yl)ethyl]-3-hydroxypentanoic acid (Compound No. 204).

7. A process for preparing a compound of Formula XXXV comprising:


a. reacting a compound of Formula VIIIb



with a compound of Formula IX




to give a compound of Formula XXVII;




Formula XXVII

b. deprotecting a compound of Formula XXVII to give a compound of Formula XXVIII




Formula XXVIII

c. oxidizing the compound of Formula XXVIII to give a compound of Formula XXIX;




Formula XXIX

d. reacting the compound of Formula XXIX with a compound of Formula XIII




Formula XIII


to give a compound of Formula XXX;




e. reacting the compound of Formula XXX with a compound of Formula XV




to give a compound of Formula XXXI;



f. hydrogenating the compound of Formula XXXI to give a compound of Formula XXXII;




g. deprotecting the compound of Formula XXXII to give a compound of Formula XXXIII;




Formula XXXIII

h. converting the compound of Formula XXXIII to a compound of Formula XXXIV; and



Formula XXXIV

i. deformylating the compound of Formula XXXIV to give a compound of Formula XXXV,



Formula XXXV

wherein

P₁ is a silyl protecting group;

R' and R'' together form an acetal protecting group;

R₂ is N-containing heterocyclyl or heteroaryl;

hal is Cl, Br or I;

Ar is phenyl;

Q is optionally substituted aryl or heteroaryl; and

R₁ is hydrogen, optionally substituted alkyl, alkenyl, alkynyl, cycloalkyl, aryl, heterocyclyl, heteroaryl, aralkyl, alkoxy, aryloxy, alkenyloxy or alkynyloxy.


8. The process according to claim 7, wherein compounds of Formula XXXV are selected from:

(2*S*,3*S*)-3-hydroxy-5-[4-(6-methoxypyridin-3-yl)phenyl]-2-[2-(4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)ethyl]pentanoic acid (Compound No. 143),

(2*S*,3*S*)-3-hydroxy-5-(4'-methoxybiphenyl-4-yl)-2-[2-(4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)ethyl]pentanoic acid (Compound No. 208).


9. A process for preparing a compound of Formula LIII comprising:

a. converting a compound of Formula XLI to a compound of Formula XLII




by reaction with a Wittig reagent;

b. hydrogenating the compound of Formula XLII to give a compound of Formula XLIII;



c. reducing the compound of Formula XLIII to give a compound of Formula XLIV;




Formula XLIV

d. reacting the compound of Formula XLIV with a compound of Formula IX




Formula IX

to give a compound of Formula XLV;




Formula XLV

e. deprotecting the compound of Formula XLV to give a compound of Formula XLVI;




Formula XLVI

f. oxidizing the compound of Formula XLVI to give a compound of Formula XLVII;




Formula XLVII


g. reacting the compound of Formula XLVII with a compound of Formula XIII




to give a compound of Formula XLVIII;




h. reacting the compound of Formula XLVIII with a compound of Formula XV




to give a compound of Formula XLIX;



i. hydrogenating the compound of Formula XLIX to give a compound of Formula L;




- j. deprotecting the compound of Formula L to give a compound of Formula LI;




Formula LI

- k. converting the compound of Formula LI to a compound of Formula LII; and



Formula LII

- l. deformylating the compound of Formula LII to give the compound of Formula LIII,



Formula LIII

wherein

R_7 and R_8 together form acetal protecting group;

R' and R'' together form an acetal protecting group;

E is an alkyl group;

R_2 is an N-containing heterocyclyl or heteroaryl;

hal is Cl, Br or I;

Ar is phenyl;

Q is optionally substituted aryl or heteroaryl; and

R_1 is hydrogen, optionally substituted alkyl, alkenyl, alkynyl, cycloalkyl, aryl, heterocyclyl, heteroaryl, aralkyl, alkoxy, aryloxy, alkenyloxy or alkynyloxy.

R_1 is hydrogen, optionally substituted alkyl, alkenyl, alkynyl, cycloalkyl, aryl, heterocyclyl, heteroaryl, aralkyl, alkoxy, aryloxy, alkenyloxy or alkynyloxy.

10. The process according to claim 9, wherein compounds of Formula LIII are selected from:

(2*R*,3*S*)-3-hydroxy-5-[4-(6-methoxypyridin-3-yl)phenyl]-2-[2-(4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)ethyl]pentanoic acid (Compound No. 120),


(2*R*,3*S*)-2-[2-(1,3-dioxo-1,3-dihydro-2*H*-isoindol-2-yl)ethyl]-5-(3'-fluoro-4'-methylbiphenyl-4-yl)-3-hydroxypentanoic acid (Compound No. 205),

(2*R*,3*S*)-2-[2-(1,3-dioxo-1,3-dihydro-2*H*-isoindol-2-yl)ethyl]-5-(4'-fluorobiphenyl-4-yl)-3-hydroxypentanoic acid (Compound No. 206), and

(2*R*,3*S*)-5-(3',4'-difluorobiphenyl-4-yl)-2-[2-(1,3-dioxo-1,3-dihydro-2*H*-isoindol-2-yl)ethyl]-3-hydroxypentanoic acid (Compound No. 207).


11. A process for preparing a compound of Formula LXVI comprising:

a. converting a compound of Formula XXXVII to a compound of Formula LIV




by reaction with a Wittig reagent;

b. hydrogenating the compound of Formula LIV to give a compound of Formula LV;




c. deprotecting the compound of Formula LV to give a compound of Formula LVI;




Formula LVI

- d. oxidizing the compound of Formula LVI to give a compound of Formula LVII;




Formula LVII

- e. reacting the compound of Formula LVII with a compound of Formula XIII




Formula XIII

- to give a compound of Formula LVIII;




Formula LVIII

- f. reacting the compound of Formula LVIII with a compound of Formula XV




Formula XV

- to give a compound of Formula LIX;




Formula LIX

- g. hydrogenating the compound of Formula LIX to give a compound of Formula LX;




Formula LX

- h. deprotecting the compound of Formula LX to give a compound of Formula LXI;




Formula LXI

- i. activating the compound of Formula LXI to a compound of Formula LXII;




Formula LXII

- j. reacting the compound of Formula LXII with a compound of Formula IXa.




to give a compound of Formula LXIII;




Formula LXIII

- k. deprotecting the compound of Formula LXIII to give a compound of Formula LXIV;




Formula LXIV

- l. converting the compound of Formula LXIV to a compound of Formula LXV; and



Formula LXV

- m. deformylating the compound of Formula LXV to give the compound of Formula LXVI,



Formula LXVI

wherein

R₇ and R₈ together form a acetal protecting group;

R' and R'' together form an acetal protecting group;

R₂ is an N-containing heterocyclyl or heteroaryl;

U is an O-activating group;

E is an alkyl group;

hal is Cl, Br or I;

- 51 Ar is phenyl;
 52 M is a metal;
 53 Q is an optionally substituted aryl or heteroaryl; and
 54 R₁ is hydrogen, optionally substituted alkyl, alkenyl, alkynyl, cycloalkyl, aryl,
 55 heterocyclyl, heteroaryl, aralkyl, alkoxy, aryloxy, alkenyloxy or alkynyloxy.


56 12. The process according to claim 11, wherein compounds of Formula LXVI are
 57 selected from:

58 (2R,3R)-5-(4'-chlorobiphenyl-4-yl)-2-[2-(1,3-dioxo-1,3-dihydro-2H-isoindol-2-
 59 yl)ethyl]-3-hydroxypentanoic acid (Compound No.89),and


60 (2R,3R)-3-hydroxy-5-[4-(6-methoxypyridin-3-yl)phenyl]-2-[2-(4-oxo-1,2,3-
 61 benzotriazin-3(4H)-yl)ethyl]pentanoic acid (Compound No. 96).

1 13. A process for preparing a compound of Formula XX comprising:


2 a. reacting a compound of Formula LXVII




4 with a compound of Formula LXVIII




6 to give a compound of Formula LXIX;



8 b. converting the compound of Formula LXIX to a compound of Formula LXX;




10 c. hydrogenating the compound of Formula LXX to give a compound of Formula
 11 LXXI;




Formula LXXI

- d. reducing the compound of Formula LXXI to give a compound of Formula LXXII;




Formula LXXII

- e. oxidizing the compound of Formula LXXII to give a compound of Formula LXXIII;




Formula LXXIII

- f. reacting the compound of Formula LXXIII with a compound of Formula LXXIV




Formula LXXIV

- to give a compound of Formula LXXV; and



Formula LXXV

g. hydrolyzing the compound of Formula LXXV to give a compound of Formula XX,



Formula XX

wherein

R₂ is an N-containing heterocyclyl or heteroaryl,

E is an alkyl group;

hal is Cl, Br or I;

Q is an optionally substituted aryl or heteroaryl;

R₁ is hydrogen, optionally substituted alkyl, alkenyl, alkynyl, cycloalkyl, aryl, heterocyclyl, heteroaryl, aralkyl, alkoxy, aryloxy, alkenyloxy or alkynyloxy;

Y and W are independently oxygen or sulphur; and

R₁₀ is alkyl, aryl or aralkyl.

14. The process according to claim 13 wherein compounds of Formula XX are selected from:

(2*S*,3*R*)-3-hydroxy-5-[4-(6-methoxypyridin-3-yl)phenyl]-2-[2-(4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)ethyl]pentanoic acid (Compound No. 86),

(2*S*,3*R*)-3-hydroxy-5-[4-(5-methylpyridin-2-yl)phenyl]-2-[2-(4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)ethyl]pentanoic acid (Compound No. 209),

(2*S*,3*R*)-5-[4-(6-fluoropyridin-3-yl)phenyl]-3-hydroxy-2-[2-(4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)ethyl]pentanoic acid (Compound No. 210),

(2*S*,3*R*)-5-(3'-fluoro-4'-methylbiphenyl-4-yl)-3-hydroxy-2-[(4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)methyl]pentanoic acid (Compound No. 214),

(2*S*,3*R*)-3-hydroxy-2-[(4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)methyl]-5-[4'-(trifluoromethyl)biphenyl-4-yl]pentanoic acid (Compound No. 215),

(2*S*,3*R*)-5-(2',4'-difluorobiphenyl-4-yl)-3-hydroxy-2-[(4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)methyl]pentanoic acid (Compound No. 216),

((2*S*,3*R*)-5-(4'-fluorobiphenyl-4-yl)-3-hydroxy-2-[(4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)methyl]pentanoic acid (Compound No. 217),

(2*S*,3*R*)-5-(3'-fluoro-4'-methoxybiphenyl-4-yl)-3-hydroxy-2-[(4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)methyl]pentanoic acid (Compound No. 218),

(2*S*,3*R*)-3-hydroxy-5-[4-(2-methoxypyrimidin-5-yl)phenyl]-2-[(4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)methyl]pentanoic acid (Compound No. 219),

(2*S*,3*R*)-3-hydroxy-5-[4-(6-methoxypyridin-3-yl)phenyl]-2-[(4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)methyl]pentanoic acid (Compound No. 220),

(2*S*,3*R*)-3-hydroxy-5-(4'-methylbiphenyl-4-yl)-2-[(4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)methyl]pentanoic acid (Compound No. 221),

(2*S*,3*R*)-3-hydroxy-2-[(4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)methyl]-5-[4'-(trifluoromethoxy)biphenyl-4-yl]pentanoic acid (Compound No. 222),

(2*S*,3*R*)-3-hydroxy-5-[4-(6-methylpyridin-3-yl)phenyl]-2-[(4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)methyl]pentanoic acid (Compound No. 223),

(2*S*,3*R*)-5-(4'-chlorobiphenyl-4-yl)-3-hydroxy-2-[(4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)methyl]pentanoic acid (Compound No. 224), and

(2*S*,3*R*)-5-(3',4'-difluorobiphenyl-4-yl)-3-hydroxy-2-[(4-oxo-1,2,3-benzotriazin-3(4*H*)-yl)methyl]pentanoic acid (Compound No. 225).