activity on multiple kinases or multiple kinase pathways. Accordingly, there remains a need for new kinase modulators.

[0008] The protein named mTOR (mammalian target of rapamycin), which is also called FRAP, RAFTI or RAPT1), is a 2549-amino acid Ser/Thr protein kinase, that has been shown to be one of the most critical proteins in the mTOR/PI3K/Akt pathway that regulates cell growth and proliferation. Georgakis and Younes *Expert Rev. Anticancer Ther.* 6(1):131-140 (2006). mTOR exists within two complexes, mTORC1 and mTORC2. While mTORC1 is sensitive to rapamycin analogs (such as temsirolimus or everolimus), mTORC2 is largely rapamycin-insensitive. Notably, rapamycin is not a TOR kinase inhibitor. Several mTOR inhibitors have been or are being evaluated in clinical trials for the treatment of cancer. Temsirolimus was approved for use in renal cell carcinoma in 2007 and sirolimus was approved in 1999 for the prophylaxis of renal transplant rejection. Everolimus was approved in 2009 for renal cell carcinoma patients that have progressed on vascular endothelial growth factor receptor inhibitors, in 2010 for subependymal giant cell astrocytoma (SEGA) associated with tuberous sclerosis (TS) in patients who require therapy but are not candidates for surgical resection, and in 2011 for progressive neuroendocrine tumors of pancreatic origin (PNET) in patients with unresectable, locally advanced or metastatic disease. There remains a need for additional TOR kinase inhibitors.

[0009] Citation or identification of any reference in Section 2 of this application is not to be construed as an admission that the reference is prior art to the present application.

3. SUMMARY

[0010] In the first aspect, the present invention refers to the use of a TOR kinase inhibitor in the manufacture of a medicament for treating triple negative breast cancer or hormone receptor positive breast cancer in a patient.

In the second aspect, the present invention refers to the use of a TOR kinase inhibitor in the manufacture of a medicament for improving the Response Evaluation Criteria in Solid Tumors (RECIST 1.1) of a patient, wherein said patient is having triple negative breast cancer or hormone receptor positive breast cancer.

In the third aspect, the present invention refers to an *in vitro* method for inhibiting phosphorylation of S6RP, 4E-BP1 and/or AKT in a biological sample isolated of a patient having triple negative breast cancer or hormone receptor positive breast cancer, wherein said patient was administered an effective amount of a TOR kinase inhibitor, wherein the method comprises comparing the amount of phosphorylated S6RP, 4E-BP1 and/or AKT in a biological sample isolated of a patient obtained prior to and after administration of said TOR kinase inhibitor, wherein less phosphorylated S6RP, 4E-BP1 and/or

AKT in said biological sample obtained after administration of said TOR kinase inhibitor relative to the amount of phosphorylated S6RP, 4E-BP1 and/or AKT in said biological sample obtained prior to administration of said TOR kinase inhibitor indicates inhibition.

In the fourth aspect, the present invention refers to an *in vitro* method for inhibiting DNA-dependent protein kinase (DNA_PK) activity in a skin sample of a patient having triple negative breast cancer or hormone receptor positive breast cancer, wherein said patient was administered an effective amount of a TOR kinase inhibitor, wherein the method comprises comparing the amount of phosphorylated DNA-PK in a biological sample isolated of a patient obtained prior to and after administration of said TOR kinase inhibitor, wherein less phosphorylated DNA-PK in said biological sample obtained after administration of said TOR kinase inhibitor relative to the amount of phosphorylated DNA-PK in said biological sample obtained prior to administration of said TOR kinase inhibitor indicates inhibition.

In the fifth aspect, the present invention refers to an *in vitro* method for measuring inhibition of of phosphorylation of S6RP, 4E-BP1 or AKT in a patient having triple negative breast cancer or hormone receptor positive breast cancer, comprising measuring the amount of phosphorylated S6RP, 4E-BP1 or AKT in said patient who was administered an effective amount of a TOR kinase inhibitor, and comparing said amount of phosphorylated S6RP, 4E-BP1 or AKT to that of said patient prior to administration of an effective amount of a TOR kinase inhibitor.

In the sixth aspect, the present invention refers to a method for measuring inhibition of phosphorylation of DNA-PK S2056 in a skin sample of a patient having triple negative breast cancer or hormone receptor positive breast cancer, comprising irradiating a skin sample of said patient, who was administered an effective amount of a TOR kinase inhibitor, with UV light, measuring the amount of phosphorylated DNA-PK S2056 present in the UV light-irradiated skin sample isolated from said patient and comparing said amount of phosphorylated DNA-PK S2056 to that in a UV light-irradiated skin sample from said patient prior to administration of an effective amount of a TOR kinase inhibitor.

In the seventh aspect, the present invention refers to a kit comprising a TOR kinase inhibitor and means for monitoring patient response to administration of said TOR kinase inhibitor, wherein said patient has triple negative breast cancer or hormone receptor positive breast cancer.

~~Provided~~Disclosed herein are methods for treating or preventing triple negative breast cancer or hormone receptor positive breast cancer, comprising administering an effective amount of a TOR kinase inhibitor to a patient having triple negative breast cancer or hormone receptor positive breast cancer.

[0011] In certain ~~embodiments~~examples, provided herein are methods for improving the Response Evaluation Criteria in Solid Tumors (RECIST 1.1) of a patient, comprising

administering an effective amount of a TOR kinase inhibitor to a patient having triple negative breast cancer or hormone receptor positive breast cancer.

[0012] In some ~~embodiments~~examples, the TOR kinase inhibitor is a compound as described herein.

[0013] The present embodiments can be understood more fully by reference to the detailed description and examples, which are intended to exemplify non-limiting embodiments.

4. DETAILED DESCRIPTION

4.1 DEFINITIONS

[0014] An "alkyl" group is a saturated, partially saturated, or unsaturated straight chain or branched non-cyclic hydrocarbon having from 1 to 10 carbon atoms, typically from 1 to 8 carbons or, in some embodiments, from 1 to 6, 1 to 4, or 2 to 6 or carbon atoms. Representative alkyl groups include -methyl, -ethyl, -n-propyl, -n-butyl, -n-pentyl and -n-hexyl; while saturated branched alkyls include -isopropyl, -*sec*-butyl, -isobutyl, -*tert*-butyl, -isopentyl, 2-methylpentyl, 3-methylpentyl, 4-methylpentyl, 2,3-dimethylbutyl and the like. Examples of unsaturared alkyl groups include, but are not limited to, vinyl, allyl, $-CH=CH(CH_3)$, $-CH=C(CH_3)_2$, $-C(CH_3)=CH_2$, $-C(CH_3)=CH(CH_3)$, $-C(CH_2CH_3)=CH_2$, $-C\equiv CH$, $-C\equiv C(CH_3)$, $-C\equiv C(CH_2CH_3)$, $-CH_2C\equiv CH$, $-CH_2C\equiv C(CH_3)$ and $-CH_2C\equiv C(CH_7CH_3)$, among others. An alkyl group can be substituted or unsubstituted. In certain embodiments, when the alkyl groups described herein are said to be "substituted," they may be substituted with any substituent or substituents as those found in the exemplary compounds and embodiments disclosed herein, as well as halogen (chloro, iodo, bromo, or fluoro); hydroxyl; alkoxy; alkoxyalkyl; amino; alkylamino; carboxy; nitro; cyano; thiol; thioether; imine; imide; amidine; guanidine; enamine; aminocarbonyl; acylamino; phosphonato; phosphine; thiocarbonyl; sulfonyl; sulfone; sulfonamide; ketone; aldehyde; ester; urea; urethane; oxime; hydroxyl amine; alkoxyamine; aralkoxyamine; N-oxide;

CLAIMS

What is claimed is:

1. Use of a TOR kinase inhibitor ~~A method~~in the manufacture of a medicament for treating triple negative breast cancer or hormone receptor positive breast cancer in a patient~~, comprising administering an effective amount of a TOR kinase inhibitor to a patient having triple negative breast cancer or hormone receptor positive breast cancer~~.

2. The ~~method~~ use of claim 1, wherein the patient has triple negative breast cancer.

3. The ~~method~~ use of claim 1, wherein the patient has hormone receptor positive breast cancer.

4. The ~~method~~ use of claim 1, wherein the triple negative breast cancer or hormone receptor positive breast cancer is that in which the PI3K/mTOR pathway is activated.

5. The ~~method of~~use claim 4, wherein the triple negative breast cancer or hormone receptor positive breast cancer is that in which the PI3K/mTOR pathway is activated due to PTEN loss, a PIK3Ca mutation or EGFR overexpression, or a combination thereof.

6. The ~~method~~ use of claim 1, wherein said patient is to be administered about 0.5 mg/day to about 128 mg/day of a TOR kinase inhibitor.

7. The ~~method~~ use of claim 6, wherein said patient is to be administered 0.5 mg/day, 1 mg/day, 2 mg/day, 4 mg/day, 8 mg/day, 16 mg/day, 20 mg/day, 30 mg/day, 45 mg/day, 60 mg/day, 90 mg/day, 120 mg/day or 128 mg/day of a TOR kinase inhibitor.

8. The ~~method~~ use of claim 1, wherein said patient is to be administered a unit dosage form comprising 0.25 mg, 1.0 mg, 5.0 mg, 7.5 mg, or 10 mg of a TOR kinase inhibitor.

9. Use of a TOR kinase inhibitor ~~A method~~in the manufacture of a medicament for improving the Response Evaluation Criteria in Solid Tumors (RECIST 1.1) of a patient, ~~comprising administering an effective amount of a TOR kinase inhibitor to a~~wherein said patient is having triple negative breast cancer or hormone receptor positive breast cancer

10. The ~~method~~ use of claim 9, wherein the triple negative breast cancer or hormone receptor positive breast cancer is that in which the PI3K/mTOR pathway is activated.

11. The ~~method~~ use of claim 10, wherein the triple negative breast cancer or hormone receptor positive breast cancer is that in which the PI3K/mTOR pathway is activated due to PTEN loss, a PIK3Ca mutation or EGFR overexpression, or a combination thereof.

12. An *in vitro* method for inhibiting phosphorylation of S6RP, 4E-BP1 and/or AKT in a biological sample isolated of a patient having triple negative breast cancer or hormone receptor positive breast cancer, ~~comprising administering an effective amount of a TOR kinase inhibitor to~~wherein said patient was administered an effective amount of a TOR kinase inhibitor, ~~and~~ wherein the method comprises comparing the amount of phosphorylated S6RP, 4E-BP1 and/or AKT in a biological sample isolated of a patient obtained prior to and after administration of said TOR kinase inhibitor, wherein less phosphorylated S6RP, 4E-BP1 and/or AKT in said biological sample obtained after administration of said TOR kinase inhibitor relative to the amount of phosphorylated S6RP, 4E-BP1 and/or AKT in said biological sample obtained prior to administration of said TOR kinase inhibitor indicates inhibition.

13. The method of claim 12, wherein the triple negative breast cancer or hormone receptor positive breast cancer is that in which the PI3K/mTOR pathway is activated.

14. The method of claim 13, wherein the triple negative breast cancer or hormone receptor positive breast cancer is that in which the PI3K/mTOR pathway is

activated due to PTEN loss, a PIK3Ca mutation or EGFR overexpression, or a combination thereof.

15. An *in vitro* method for inhibiting DNA-dependent protein kinase (DNA-PK) activity in a skin sample of a patient having triple negative breast cancer or hormone receptor positive breast cancer, ~~comprising administering~~wherein said patient was administered an effective amount of a TOR kinase inhibitor ~~to said patient and~~, wherein the method comprises comparing the amount of phosphorylated DNA-PK in a biological sample isolated of a patient obtained prior to and after administration of said TOR kinase inhibitor, wherein less phosphorylated DNA-PK in said biological sample obtained after administration of said TOR kinase inhibitor relative to the amount of phosphorylated DNA-PK in said biological sample obtained prior to administration of said TOR kinase inhibitor indicates inhibition.

16. The method of claim 15, wherein the triple negative breast cancer or hormone receptor positive breast cancer is that in which the PI3K/mTOR pathway is activated.

17. The method of claim 16, wherein the triple negative breast cancer or hormone receptor positive breast cancer is that in which the PI3K/mTOR pathway is activated due to PTEN loss, a PIK3Ca mutation or EGFR overexpression, or a combination thereof.

18. An *in vitro* method for measuring inhibition of phosphorylation of S6RP, 4E-BP1 or AKT in a patient having triple negative breast cancer or hormone receptor positive breast cancer, comprising ~~administering an effective amount of a TOR kinase inhibitor to said patient,~~ measuring the amount of phosphorylated S6RP, 4E-BP1 or AKT in said patient who was administered an effective amount of a TOR kinase inhibitor, and comparing said amount of phosphorylated S6RP, 4E-BP1 or AKT to that of said patient prior to administration of an effective amount of a TOR kinase inhibitor.

19. The method of claim 16, wherein the triple negative breast cancer or hormone receptor positive breast cancer is that in which the PI3K/mTOR pathway is activated.

20. The method of claim 17, wherein the triple negative breast cancer or hormone receptor positive breast cancer is that in which the PI3K/mTOR pathway is activated due to PTEN loss, a PIK3Ca mutation or EGFR overexpression, or a combination thereof.

21. A method for measuring inhibition of phosphorylation of DNA-PK S2056 in a skin sample of a patient having triple negative breast cancer or hormone receptor positive breast cancer, comprising ~~administering an effective amount of a TOR kinase inhibitor to said patient,~~ irradiating a skin sample of said patient, who was administered an effective amount of a TOR kinase inhibitor, with UV light, measuring the amount of phosphorylated DNA-PK S2056 present in the UV light-irradiated skin sample isolated from said patient and comparing said amount of phosphorylated DNA-PK S2056 to that in a UV light-irradiated skin sample from said patient prior to administration of an effective amount of a TOR kinase inhibitor.

22. The method of claim 18, wherein the triple negative breast cancer or hormone receptor positive breast cancer is that in which the PI3K/mTOR pathway is activated.

23. The method of claim 19, wherein the triple negative breast cancer or hormone receptor positive breast cancer is that in which the PI3K/mTOR pathway is activated due to PTEN loss, a PIK3Ca mutation or EGFR overexpression, or a combination thereof.

24. A kit comprising a TOR kinase inhibitor and means for monitoring patient response to administration of said TOR kinase inhibitor, wherein said patient has triple