

ENGINEERED IGA ANTIBODIES AND METHODS OF USE

5

CROSS-REFERENCE

[0001] This application claims the benefit of U.S. Provisional Application No. 62/824,864, filed March 27, 2019; which is incorporated herein by reference in their entirety.

BACKGROUND OF THE DISCLOSURE

10 [0002] Monoclonal antibodies of IgG isotype targeting tumor antigens have proven to be an effective treatment of various cancers. Over the years an increasing number of monoclonal antibodies targeting different tumor antigens have been approved for use in cancer therapies. However, their clinical efficacy and side effects, especially as a monotherapy, are still insufficient. Therefore, it is of interest to develop new antibody therapies with increased clinical efficacy, novel targeting modalities or modes of action and/or
15 decreased number and severity of side effects.

SUMMARY OF THE DISCLOSURE

[0003] Provided herein is an antibody or a functional fragment thereof that comprises: an antigen binding domain; and a constant domain, wherein the constant domain comprises an immunoglobulin A (IgA) heavy
20 chain constant region, wherein the IgA heavy chain constant region comprises an IgA CH2 region and an IgA CH3 region, wherein the IgA heavy chain constant region comprises a modification of at least two naturally occurring glycosylation sites, as compared to a corresponding wild type (WT) IgA heavy chain constant region, and wherein each of the at least two naturally occurring glycosylation sites is in the IgA CH2 region or in the IgA CH3 region.

25 [0004] In some embodiments, the at least two naturally occurring glycosylation sites are two naturally occurring N-linked glycosylation sites. In some embodiments, one or more of the at least two naturally occurring glycosylation sites comprise a naturally occurring asparagine (N) amino acid residue which are modified in the engineered antibodies disclosed herein as compared to a corresponding wild type IgA. In some embodiments, the modification comprises an amino acid substitution, or an amino acid deletion of one,
30 or both of the at least two naturally occurring glycosylation sites. In some embodiments, the amino acid substitution is a non-conservative amino acid substitution. In some embodiments, the IgA heavy chain constant region comprises an amino acid substitution at: i. N114 and N135, ii. N114 and N15.2, or iii. N135 and N15.2, relative to a corresponding WT IgA heavy chain constant region comprising an amino acid sequence of SEQ ID NO: 1, numbering according to IMGT scheme. In some embodiments, the IgA heavy
35 chain constant region comprises: i. a N114T amino acid substitution and a N135Q amino acid substitution, ii. a N114T amino acid substitution and an amino acid substitution selected from the group consisting of

N15.2G, N15.2Q, and N15.2T, or iii. a N135Q amino acid substitution and an amino acid substitution from the group consisting of N15.2G, N15.2Q, and N15.2T, relative to a corresponding WT IgA heavy chain constant region comprising an amino acid sequence of SEQ ID NO: 1, numbering according to IMGT scheme.

- 5 [0005] In some embodiments, the IgA heavy chain constant region comprises a modification of at least three naturally occurring glycosylation sites, as compared to a corresponding wild type IgA. In some embodiments, the at least three naturally occurring glycosylation sites are three N-linked glycosylation sites. In some embodiments, the three naturally occurring glycosylation sites each comprise an asparagine (N) amino acid residue which is modified in the antibodies described herein.
- 10 [0006] In some embodiments, the modification is an amino acid substitution or an amino acid deletion. In some embodiments, the amino acid substitution is a non-conservative substitution. In some embodiments, the IgA heavy chain constant region comprises an amino acid substitution at: N114, N135, and N15.2, relative to the WT IgA heavy chain constant region comprising an amino acid sequence of SEQ ID NO: 1, numbering according to IMGT scheme. In some embodiments, the IgA heavy chain constant region comprises: i. a
- 15 N114T amino acid substitution, ii. a N135Q amino acid substitution, and iii. an amino acid substitution selected from the group consisting of N15.2G, N15.2Q, and N15.2T, relative to a WT IgA heavy chain constant region comprising an amino acid sequence of SEQ ID NO: 1, numbering according to the IMGT scheme.
- [0007] In some embodiments, an IgA heavy chain constant region of an antibody described herein further
- 20 comprises an IgA CH1 region. In some embodiments, the heavy chain constant region comprises a modification of at least one naturally occurring N-linked glycosylation site in the IgA CH2 region, at least one naturally occurring glycosylation site in the IgA CH3 regions, and at least one naturally occurring glycosylation site within the IgA CH1 region, as compared to a corresponding wild type IgA. In some
- 25 embodiments, the IgA heavy chain constant region comprises an amino acid substitution at: i. N45.2, N114, and N135, or ii. N45.2, N15.2, and N135, relative to a WT IgA heavy chain constant region comprising an amino acid sequence of SEQ ID NO: 1, numbering according to IMGT scheme.
- [0008] In some embodiments, the IgA heavy chain constant region comprises: i. a N45.2G amino acid substitution, a N114T amino acid substitution, and a N135Q amino acid substitution; or ii. the N45.2G amino acid substitution, the N135Q amino acid substitution, and an amino acid substitution selected from a group
- 30 consisting of N15.2G, N15.2Q, and N15.2T, relative to a WT IgA heavy chain constant region comprising an amino acid sequence of SEQ ID NO: 1, numbering according to IMGT scheme.
- [0009] In some embodiments, the heavy chain constant region of an antibody described herein comprises a modification of at least two naturally occurring N-linked glycosylation site in the IgA CH2 region, and at least one naturally occurring glycosylation site within the IgA CH1 region, as compared to a corresponding

wild type IgA. In some embodiments, the IgA heavy chain constant region comprises an amino acid substitution at: i. N45.2, N114, and N15.2G, numbering according to IMGT scheme. In some embodiments, described herein the IgA heavy chain constant region comprises: i. a N45.2G amino acid substitution, ii. a N114T amino acid substitution, and iii. an amino acid substitution selected from the group consisting of
5 N15.2G, N15.2Q, and N15.2T, relative to the WT IgA heavy chain constant region comprising an amino acid sequence of SEQ ID NO: 1, numbering according to IMGT scheme

[0010] In some embodiments, an antibody described herein or functional fragment thereof exhibits a greater circulating half-life compared to a corresponding WT IgA antibody. In some embodiments, the antibody or the functional fragment thereof exhibits decreased aggregation compared to a corresponding WT IgA
10 antibody. In some embodiments, the antibody or the functional fragment thereof exhibits decreased aggregation with serum proteins, compared to a corresponding WT IgA antibody. In some embodiments, the antibody or the functional fragment thereof induces increased antibody dependent cell mediated cytotoxicity (ADCC), compared to a comparable antibody comprising an IgG heavy chain constant region. In some
15 embodiments, the antibody or the functional fragment thereof exhibits increased thermostability, compared to a corresponding WT IgA antibody. In some embodiments, the antibody or the functional fragment thereof exhibits decreased glycosylation, compared to a corresponding WT IgA antibody.

[0011] In some embodiments, the IgA heavy chain constant region exhibits binding to a FcαR expressed on an immune effector cell with increased affinity, compared to a corresponding WT IgA antibody. In some
20 embodiments, the IgA heavy chain constant region comprises a modification of at least four naturally occurring glycosylation sites, as compared to a corresponding wild type IgA. In some embodiments, the at least four naturally occurring glycosylation sites are four naturally occurring N-linked glycosylation sites. In
25 some embodiments, the at least four naturally occurring glycosylation sites each comprise a naturally occurring asparagine (N) amino acid residue. In some embodiments, the heavy chain constant region comprises a modification of at least two naturally occurring N-linked glycosylation sites within the IgA CH2
30 region, at least one naturally occurring N-linked glycosylation sites within the IgA CH3 region, and at least one naturally occurring N-linked glycosylation site within the IgA CH1 region, as compared to a
corresponding wild type IgA. In some embodiments, the IgA heavy chain constant region comprises an amino acid substitution at amino acid residues: N45.2, N114, N135, and N15.2, relative to a WT IgA heavy chain
constant region comprising an amino acid sequence of SEQ ID NO: 1, numbering according to IMGT
scheme.

[0012] In some embodiments, the IgA heavy chain constant region of an antibody described herein comprises a non-conservative amino acid substitution at amino acid residues: N45.2, N114, N135, and N15.2, numbering according to IMGT scheme.

[0013] In some embodiments, the IgA heavy chain constant region comprises: i. a N45.2G amino acid substitution, ii. a N114T amino acid substitution, iii. a N135Q amino acid substitution, and iv an amino acid substitution selected from the group consisting of N15.2G, N15.2Q, and N15.2T, relative to the WT IgA heavy chain constant region comprising an amino acid sequence of SEQ ID NO: 1, numbering according to IMGT scheme. In some embodiments, the antibody or the functional fragment thereof exhibits a greater circulating half-life compared to a corresponding IgA antibody comprising at least one naturally occurring N-linked glycosylation site. In some embodiments, the antibody or the functional fragment thereof induces increased antibody dependent cell mediated cytotoxicity (ADCC), compared to a corresponding comparable antibody comprising an IgG CH2 domain and an IgG CH3 domain. In some embodiments, the antibody or the functional fragment thereof exhibits increased thermostability, compared to a corresponding IgA antibody comprising at least one naturally occurring N-linked glycosylation site. In some embodiments, the antibody or the functional fragment thereof exhibits decreased glycosylation, compared to a corresponding IgA antibody comprising at least one naturally occurring N-linked glycosylation site. In some embodiments, the IgA heavy chain constant region exhibits binding to a Fc α R expressed on an immune effector cell with increased affinity, compared to a corresponding IgA antibody comprising at least one naturally occurring N-linked glycosylation site.

[0014] In some embodiments, the IgA heavy chain constant region of an antibody described herein comprises a modification of at least one naturally occurring cysteine (C) amino acid residue, as compared to a corresponding wild type IgA. In some embodiments, the modification is an amino acid substitution or an amino acid deletion of the at least one naturally occurring cysteine (C) amino acid residue. In some embodiments, the amino acid substitution is a non-conservative amino acid substitution of the at least one naturally occurring cysteine (C) amino acid residue. In some embodiments, the at least one naturally occurring cysteine amino acid residue is C147 or C86, relative to a WT IgA heavy chain constant region comprising an amino acid sequence of SEQ ID NO: 1, numbering according to IMGT scheme. In some embodiments, the IgA heavy chain constant region comprises a C86S amino acid substitution, relative to the WT IgA heavy chain constant region comprising an amino acid sequence of SEQ ID NO: 1, numbering according to IMGT scheme. In some embodiments, the IgA heavy chain constant region comprises a deletion of C147, relative to the WT IgA heavy chain constant region comprising an amino acid sequence of SEQ ID NO: 1, numbering according to the IMGT scheme. In some embodiments, the antibody or the functional fragment thereof exhibits decreased aggregation compared to a corresponding WT IgA antibody or the functional fragment thereof. In some embodiments, the antibody or the functional fragment thereof exhibits decreased aggregation with serum proteins compared to a corresponding WT IgA construct.

[0015] In some embodiments, the IgA heavy chain constant region of an antibody described herein comprises a modification of at least two naturally occurring cysteine (C) amino acid residues, as compared to

a corresponding wild type IgA. In some embodiments, the modification comprises an amino acid substitution of one, or both of the at least two naturally occurring cysteine (C) amino acid residues. In some embodiments, the modification comprises a deletion of one, or both of the at least two naturally occurring cysteine (C) amino acid residues.

- 5 [0016] In some embodiments, the IgA heavy chain constant region comprises an amino acid substitution one of the at least two naturally occurring cysteine (C) amino acid residues, and a deletion of one of the at least two naturally occurring cysteine (C) amino acid residues. In some embodiments, the at least two naturally occurring cysteine amino acid residue are C147 and C86, relative to the WT IgA heavy chain constant region comprising an amino acid sequence of SEQ ID NO: 1, numbering according to IMGT scheme. In some
- 10 embodiments, the IgA heavy chain constant region comprises a deletion of C147, relative to the WT IgA heavy chain constant region comprising an amino acid sequence of SEQ ID NO: 1, numbering according to IMGT scheme. In some embodiments, the IgA heavy chain constant region comprises an amino acid substitution of C86, relative to the WT IgA heavy chain constant region comprising an amino acid sequence of SEQ ID NOs: 1, numbering according to IMGT scheme.
- 15 [0017] In some embodiments, the IgA heavy chain constant region of an antibody described herein comprises an amino acid substitution of the naturally occurring C86 amino acid residue, and a deletion of the naturally occurring C147 amino acid residue, relative to a WT IgA heavy chain constant region comprising an amino acid sequence of SEQ ID NO: 1, numbering according to IMGT scheme. In some embodiments, the IgA heavy chain constant region comprises an C86S amino acid substitution, relative to the WT IgA heavy
- 20 chain constant region comprising an amino acid sequence of SEQ ID NO: 1, numbering according to IMGT scheme. In some embodiments, the antibody or the functional fragment thereof exhibits decreased aggregation compared to a corresponding WT IgA antibody. In some embodiments, the antibody or the functional fragment thereof exhibits decreased aggregation with serum proteins compared to a corresponding WT IgA antibody.
- 25 [0018] In some embodiments, the IgA heavy chain constant region of an antibody described herein comprises a modification of at least one naturally occurring tyrosine (Y) amino acid residue, as compared to a corresponding wild type IgA. In some embodiments, the modification is an amino acid substitution or a deletion of the at least one naturally occurring tyrosine (Y) amino acid residue. In some embodiments, the amino acid substitution is a non-conservative amino acid mutation of the at least one naturally occurring
- 30 tyrosine (Y) amino acid residue, as compared to a WT IgA antibody. In some embodiments, the at least one tyrosine residue is Y148, relative to a WT IgA heavy chain constant region comprising an amino acid sequence of SEQ ID NO: 1, numbering according to IMGT scheme. In some embodiments, the amino acid Y148 is deleted, relative to the WT IgA heavy chain constant region comprising an amino acid sequence of SEQ ID NO: 1, numbering according to IMGT scheme. In some embodiments, the antibody or the functional

fragment thereof exhibits decreased aggregation compared to a corresponding WT IgA antibody. In some embodiments, the antibody or the functional fragment thereof exhibits decreased aggregation with serum proteins compared to a corresponding WT IgA antibody.

- [0019]** In some embodiments, the IgA heavy chain constant region of an antibody described herein comprises a modification of at least one naturally occurring threonine (T) amino acid residue, as compared to a corresponding wild type IgA antibody. In some embodiments, the modification comprises an amino acid substitution or deletion of the at least one naturally occurring threonine (T) amino acid residue, as compared to a corresponding wild type IgA antibody. In some embodiments, the amino acid substitution is a non-conservative amino acid substitution of the at least one naturally occurring threonine (T) amino acid residue.
- 10 In some embodiments, the at least one naturally occurring threonine amino acid residue is T116 or T16, relative to the WT IgA heavy chain constant region comprising an amino acid sequence of SEQ ID NO: 1, numbering according to IMGT scheme. In some embodiments, the antibody or the functional fragment thereof exhibits decreased aggregation compared to a corresponding WT IgA antibody. In some
- 15 embodiments, the antibody or the functional fragment thereof exhibits decreased aggregation with serum proteins compared to a corresponding WT IgA antibody. In some embodiments, the IgA heavy chain constant region comprises a modification of at least two naturally occurring threonine (T) amino acid residues, as compared to a corresponding wild type IgA. In some embodiments, the modification comprises an amino acid substitution or a deletion of one, or both the at least two naturally occurring threonine (T) amino acid residues. In some embodiments, the amino acid substitution is a non-conservative amino acid substitution of
- 20 one, or both of the at least two naturally occurring threonine (T) amino acid residues. In some embodiments, the at least two naturally occurring threonine (T) amino acid residue are T116 or T16, relative to a WT IgA heavy chain constant region comprising an amino acid sequence of SEQ ID NO: 1, numbering according to IMGT scheme. In some embodiments, the IgA heavy chain constant region comprises a T116S amino acid substitution, relative to a WT IgA heavy chain constant region comprising an amino acid sequence of SEQ ID
- 25 NO: 1, numbering according to IMGT scheme. In some embodiments, the IgA heavy chain constant region comprises a T16S amino acid substitution, relative to the WT IgA heavy chain constant region comprising an amino acid sequence of SEQ ID NO: 1, numbering according to IMGT scheme.

- [0020]** In some embodiments, the IgA heavy chain constant region of an antibody described herein comprises a modification of at least one naturally occurring isoleucine (I) amino acid residue, as compared to a corresponding wild type IgA. In some embodiments, the modification comprises an amino acid substitution or a deletion of the at least one naturally occurring isoleucine (I) amino acid residue. In some embodiments, the amino acid substitution comprises a non-conservative amino acid substitution of the at least one naturally occurring isoleucine (I) amino acid residue. In some embodiments, the at least one naturally occurring
- 30 isoleucine (I) residue is I115, relative to the WT IgA heavy chain constant region comprising an amino acid

sequence of SEQ ID NO: 1, numbering according to IMGT scheme. In some embodiments, the IgA heavy chain constant region comprises a I115L amino acid substitution, relative to a WT IgA heavy chain constant region comprising an amino acid sequence of SEQ ID NO: 1, numbering according to IMGT scheme. In some embodiments, the IgA heavy chain constant region comprises a modification of at least one naturally occurring leucine (L) amino acid residue, as compared to a corresponding wild type IgA. In some
5 embodiments, the modification comprises an amino acid substitution or a deletion of the at least one naturally occurring leucine (L) amino acid residue. In some embodiments, the amino acid substitution is a non-conservative amino acid substitution of the at least one naturally occurring leucine (L) amino acid residue. In some embodiments, the at least one naturally occurring leucine (L) residue is L15.3, relative to the WT IgA
10 heavy chain constant region comprising an amino acid sequence of SEQ ID NO: 1, numbering according to IMGT scheme. In some embodiments, the IgA heavy chain constant region comprises a L15.3I amino acid substitution, relative to a WT IgA heavy chain constant region comprising an amino acid sequence of SEQ ID NO: 1, numbering according to IMGT scheme.

[0021] In some embodiments, the IgA heavy chain constant region of an antibody described herein
15 comprises a modification of at least one naturally occurring proline (P) amino acid residue, as compared to a corresponding wild type IgA. In some embodiments, the IgA heavy chain constant region comprises an amino acid substitution or a deletion of the at least one naturally occurring proline (P) amino acid residue. In some embodiments, the amino acid substitution is a non-conservative amino acid substitution of the at least one naturally occurring proline (P) amino acid residue. In some embodiments, the at least one naturally occurring
20 proline (P) residue is P124, relative to a WT IgA heavy chain constant region comprising an amino acid sequence of SEQ ID NO: 1, numbering according to IMGT scheme. In some embodiments, the IgA heavy chain constant region comprises a P124R amino acid substitution, relative to the WT IgA heavy chain constant region comprising an amino acid sequence of SEQ ID NO: 1, numbering according to IMGT scheme. In some embodiments, the antibody or functional fragment thereof exhibits greater stability,
25 compared to a corresponding WT IgA antibody. In some embodiments, the antibody or the functional fragment thereof exhibits greater stability between a heavy chain and a light chain, compared to a corresponding WT IgA antibody. In some embodiments, the antibody or functional fragment thereof has a covalent linkage between the heavy chain and the light chain. In some embodiments, the antibody or the functional fragment thereof comprises a disulfide bond between a cysteine (C) amino acid residue of the
30 heavy chain and a cysteine (C) amino acid residue of the light chain. In some embodiments, the IgA heavy chain constant region comprises an amino acid sequence selected from any one of SEQ ID NOs: 16-21.

[0022] In some embodiments, antibody described herein comprises a heavy chain region of a modified IgA of an allotype IgA2m(1) antibody or IgA2m(2). In some embodiments, the antibody is an allotype Caucasian IgA2m(1) antibody. In some embodiments, the IgA heavy chain constant region comprises a modification of a

C-terminal IgA tail piece, as compared to a corresponding wild type IgA. In some embodiments, the modification comprises a deletion of the C-terminal 25, 24, 23, 22, 21, 20, 19, 18, 17, 16, 15, 12, 10, 8, 6, 4, or 2 amino acids. In some embodiments, the modification comprises a deletion of the C-terminal 18 amino acids. In some embodiments, the modification comprises a deletion of C-terminal amino acid residues 131-148, relative to the WT IgA heavy chain constant region comprising an amino acid sequence of SEQ ID NO: 1, numbering according to IMGT scheme. In some embodiments, the IgA heavy chain constant region comprises a deletion of the C-terminal amino acid residues P131-Y148 relative to the WT IgA heavy chain constant region comprising an amino acid sequence of SEQ ID NO: 1, numbering according to IMGT scheme.

- 10 **[0023]** In some embodiments, the IgA heavy chain constant region of an antibody described herein comprises an IgA1 constant region comprising an IgA1 CH2 region and an IgA1 CH3 region. In some embodiments, the IgA1 constant region further comprises an IgA1 CH1 region. In some embodiments, the IgA heavy chain constant region comprises an IgA2 constant region comprising an IgA2 CH2 region and an IgA2 CH3 region. In some embodiments, the IgA2 constant region further comprises an IgA2 CH1 region. In some embodiments, the antibody or the functional fragment thereof exhibits a circulating half-life within 1%, 5%, 10%, 20%, or 30% of the circulating half-life of a corresponding antibody comprising the antigen binding domain and an IgA heavy chain constant region comprising an IgG CH2 region and an IgG CH3 region.

- 15 **[0024]** Provided herein is an antibody or a functional fragment thereof comprising: an antigen binding domain; and a constant domain, wherein the constant domain comprises an IgA2 heavy chain constant region comprising an IgA2 CH1 region, IgA2 CH2 region and an IgA2 CH3 region; wherein the IgA2 heavy chain constant region comprises a N135Q amino acid substitution, numbering according to IMGT scheme, as compared to a corresponding relative to a WT IgA heavy chain constant region comprising an amino acid sequence of SEQ ID NO: 1.

- 20 **[0025]** In some embodiments, the heavy chain constant region further comprises: a N45.2G amino acid substitution, a N114T amino acid substitution, a I115L amino acid substitution, a T116S amino acid substitution, and an amino acid substitution selected from the group consisting of N15.2G, N15.2Q, and N15.2T, relative to a WT IgA2 heavy chain constant region comprising an amino acid sequence of SEQ ID NO: 1, numbering according to IMGT scheme. In some embodiments, the heavy chain constant region further comprises: a C86S amino acid substitution, a P124R amino acid substitution, a deletion of C147, a deletion of Y148, a L15.3I amino acid substitution, a T16S amino acid substitution, or a combination thereof, relative to the WT IgA2 heavy chain constant region comprising an amino acid sequence of SEQ ID NO: 1, numbering according to IMGT scheme.

- 30 **[0026]** Provided herein is an antibody or a functional fragment thereof comprising: an antigen binding domain; and a constant domain, wherein the constant domain comprises an IgA2 heavy chain constant region

comprising an IgA2 CH2 region and an IgA2 CH3 region, and wherein the IgA2 heavy chain constant region comprises: a C86S amino acid substitution, a N114T amino acid substitution, a I115L amino acid substitution, a T116S amino acid substitution, and a N135Q amino acid substitution, relative to WT IgA2 heavy chain constant region comprising an amino acid sequence of SEQ ID NO: 1, numbering according to IMGT scheme.

5 [0027] Provided herein is an antibody or a functional fragment thereof comprising: an antigen binding domain; and a constant domain, wherein the constant domain comprises an IgA2 heavy chain constant region comprising an IgA2 CH1 region, an IgA2 CH2 region, and an IgA2 CH3 region, and wherein the IgA2 heavy chain constant region comprises: a N45.2G amino acid substitution, a C86S amino acid substitution, a N114T
10 amino acid substitution, a I115L amino acid substitution, a T116S amino acid substitution, a N135Q amino acid substitution, a deletion of C147, and a deletion of Y148, relative to the WT IgA2 heavy chain constant region comprising an amino acid sequence of SEQ ID NO: 1, numbering according to IMGT scheme. In some embodiments, the IgA2 heavy chain constant region comprises an amino acid sequence that is at least 95% identical to SEQ ID NO: 16.

15 [0028] Provided herein is an antibody or a functional fragment thereof comprising: an antigen binding domain; and a constant domain, wherein the constant domain comprises an IgA2 heavy chain constant region comprising an IgA2 CH1 region, an IgA2 CH2 region, and an IgA2 CH3 region, and wherein the IgA2 heavy chain constant region comprises: a N45.2G amino acid substitution, a C86S amino acid substitution, a N114T amino acid substitution, a I115L amino acid substitution, a T116S amino acid substitution, and a deletion of
20 C-terminal tail piece, relative to the WT IgA2 heavy chain constant region comprising an amino acid sequence of SEQ ID NO: 1, numbering according to IMGT scheme.

[0029] In some embodiments, the deletion of C-terminal tail piece comprises a deletion of C-terminal amino acid residues P131-Y148, relative to the WT IgA heavy chain constant region comprising an amino acid sequence of SEQ ID NO: 1, numbering according to the IMGT scheme.

25 [0030] In some embodiments, an IgA2 heavy chain constant region of an antibody described herein comprises an amino acid sequence that is at least 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, 99%, 99.1%, 99.2%, 99.3%, 99.4%, 99.5%, 99.6, 99.7%, 99.8%, 99.9% or 100% identical to SEQ ID NO: 17.

[0031] Provided herein is an antibody or a functional fragment thereof comprising: an antigen binding domain; and a constant domain, wherein the constant domain comprises an IgA2 heavy chain constant region
30 comprising an IgA2 CH1 region, an IgA2 CH2 region, and an IgA2 CH3 region; wherein the IgA2 heavy chain constant region comprises: a N45.2G amino acid substitution, a N114T amino acid substitution, a I115L amino acid substitution, a T116S amino acid substitution, a N135Q amino acid substitution, an amino acid substitution selected from the group consisting of N15.2G, N15.2Q, and N15.2T, a L15.3I amino acid

substitution, and a T16S amino acid substitution, relative to the WT IgA2 heavy chain constant region comprising an amino acid sequence of SEQ ID NO: 1, numbering according to the IMGT scheme.

[0032] Provided herein is an antibody or a functional fragment thereof comprising: an antigen binding domain; and a constant domain, wherein the constant domain comprises an IgA2 heavy chain constant region comprising an IgA2 CH1 region, an IgA2 CH2 region, and an IgA2 CH3 region, and wherein the IgA2 heavy chain constant region comprises: a N45.2G amino acid substitution, a N114T amino acid substitution, a I115L amino acid substitution, a T116S amino acid substitution, an amino acid substitution selected from the group consisting of N15.2G, N15.2Q, and N15.2T, a L15.3I amino acid substitution, a T16S amino acid substitution, and a deletion of C-terminal amino acids P131-Y148, relative to the WT IgA2 heavy chain constant region comprising an amino acid sequence of SEQ ID NO: 1, numbering according to the IMGT scheme.

[0033] In some embodiments, an antibody or the functional fragment thereof disclosed herein comprising: a C86S amino acid substitution, a P124R amino acid substitution, a deletion of C147, a deletion of Y148, or a combination thereof, relative to the WT IgA2 heavy chain constant region comprising an amino acid sequence of SEQ ID NO: 1, numbering according to IMGT scheme. In some embodiments, the IgA2 heavy chain constant region comprises an amino acid sequence that is at least 95% identical to a sequence selected from any one of SEQ ID NOs: 18-21. In some embodiments, the antibody or the functional fragment thereof is aglycosylated. In some embodiments, the antibody or the functional fragment thereof has increased circulatory half life than a corresponding IgA comprising one or more of wild type amino acid residues N45.2, N114, I115, T116, N15.2, L15.3, T16, or a C-terminal amino acid residues P131-Y148, numbering according to the IMGT scheme.

[0034] In some embodiments, an antibody or functional fragment thereof described herein induces increased antibody dependent cell mediated cytotoxicity (ADCC), compared to a corresponding antibody comprising an IgG heavy chain constant domain. In some embodiments, the antibody or the functional fragment thereof exhibits increased thermostability, compared to a corresponding IgA comprising one or more of wild type amino acid residues N45.2, N114, I115, T116, N15.2, L15.3, T16, or a C-terminal amino acid residues P131-Y148, numbering according to the IMGT scheme, numbering according to the IMGT scheme. In some embodiments, the antibody or the functional fragment thereof exhibits decreased glycosylation, compared to a corresponding IgA comprising one or more of wild type amino acid residues N45.2, N114, I115, T116, N15.2, L15.3, T16, or a C-terminal amino acid residues P131-Y148, numbering according to the IMGT scheme. In some embodiments, the IgA heavy chain constant region exhibits binding to FcαR expressed on an immune effector cell with increased affinity, compared to a comprising one or more of wild type amino acid residues N45.2, N114, I115, T116, N15.2, L15.3, T16, or a C-terminal amino acid residues P131-Y148, numbering according to the IMGT scheme. In some embodiments, the IgA heavy chain constant region

further comprises a hinge region. In some embodiments, the hinge region comprises an IgA hinge amino acid sequence or a variant or a fragment thereof. In some embodiments, the hinge region comprises a human IgA hinge amino acid sequence or a variant or a fragment thereof. In some embodiments, the hinge is an IgA1 hinge or an IgA2 hinge, or a variant or a fragment thereof. In some embodiments, the constant domain further comprises a light chain constant region. In some embodiments, the light chain constant region is a kappa light chain constant region, wherein the kappa light chain constant region comprises a sequence of SEQ ID NO: 31. In some embodiments, the IgA heavy chain constant region further comprises one or more albumin binding regions. In some embodiments, the one or more albumin binding domains are fused to the C-terminus of the CH3 region. In some embodiments, the constant region comprises the light chain constant region and the one or more albumin binding domain is fused to the light chain constant region. In some embodiments, the antibody or the functional fragment thereof has a greater circulating half-life compared to a corresponding IgA antibody that does not comprise the one or more albumin binding domains. In some embodiments, the antibody or the functional fragment thereof exhibits decreased complement-dependent cytotoxicity (CDC) compared to a corresponding antibody comprising an IgG heavy chain constant domain when measured in a suitable in vitro CDC assay.

[0035] In some embodiments, an antibody or functional fragment thereof described herein exhibits increased antibody-dependent cell-mediated cytotoxicity (ADCC) compared to a corresponding WT IgA antibody when measured in a suitable in vitro ADCC assay. In some embodiments, the antigen binding domain comprises a heavy chain variable region and a light chain variable region. In some embodiments, the heavy chain variable region of an antibody described herein comprises at least one of complementarity determining regions (CDRs): HC-CDR1 comprising an amino acid sequence selected from any one of SEQ ID NOs: 33-40, HC-CDR2 comprising an amino acid sequence selected from any one of SEQ ID NOs: 41-48; and HC-CDR3 comprising an amino acid sequence selected from any one of SEQ ID NOs: 49-56.

[0036] In some embodiments, the light chain variable region of an antibody described herein comprises at least one of complementarity determining regions (CDRs): LC-CDR1 comprising an amino acid sequence selected from any one of SEQ ID NOs: 57-64; LC-CDR2 comprising an amino acid sequence selected from any one of SEQ ID NOs: 65-72; and LC-CDR3 comprising an amino acid sequence selected from any one of SEQ ID NOs: 73-80.

[0037] In some embodiments, the heavy chain variable region of an antibody described herein comprises a sequence having at least 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, 99%, 99.1%, 99.2%, 99.3%, 99.4%, 99.5%, 99.6%, 99.7%, 99.8%, 99.9% or 100% identity to an amino acid sequence selected from any one of SEQ ID NOs: 4, 7, 81-86. In some embodiments, the light chain variable region comprises a sequence having at least 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, 99%, 99.1%, 99.2%, 99.3%, 99.4%, 99.5%, 99.6%, 99.7%, 99.8%, 99.9% or 100% identity to an amino acid sequence selected from any one of SEQ ID NOs: 5, 8, 95-

100. In some embodiments, each of the heavy chain and light chain CDRs are derived from an IgG antibody. In some embodiments, the heavy chain variable region further comprises four framework regions (FWs): HC-FW1, HC-FW2, HC-FW3, and HC-FW4. In some embodiments, the light chain variable region further comprises four framework regions (FWs): LC-FW1, LC-FW2, LC-FW3, and LC-FW4.

5 [0038] In some embodiments, each the heavy chain and light chain FW regions are derived from an IgG antibody. In some embodiments, each the heavy chain and light chain FW regions are derived from an IgA antibody.

[0039] In some embodiments, an antibody described herein or the functional fragment thereof is a chimeric antibody, a heavy chain antibody, a single chain antibody, a humanized antibody, a human antibody, a
10 monoclonal antibody, a deimmunized antibody, a bispecific antibody, a multispecific antibody, a multivalent antibody, or a combination thereof.

[0040] In some embodiments, the antigen-binding fragment of an antibody described herein comprises a Fab, Fab', Fab'-SH, Fv, scFv, F(ab')₂, a diabody, a linear antibody, a single domain antibodies (sdAb), a VHH domain, or a multi-specific antibody formed from antibody fragments. In some embodiments, the variable
15 domain specifically binds to GD2, CD20, CD47, CD38, CD19, EGFR, HER2, PD-L1, or CD25. In some embodiments, the antibody or the functional fragment thereof further comprises an enzyme, a substrate, cofactor, a fluorescent marker, a chemiluminescent marker, a peptide tag, a magnetic particle, a drug, a toxin, a radionuclide, a binding site for secondary antibodies, a metal binding domain, or a combination thereof.

[0041] Provided herein is a pharmaceutical composition of comprising the antibody or the functional
20 fragment thereof of any one of aspects above and a pharmaceutically acceptable carrier.

[0042] Provided herein is a method of treating or a composition for use in treating a subject in need thereof, comprising administering to the subject a therapeutic dose of the antibody or the functional fragment thereof of any one of aspects above or the pharmaceutical composition disclosed above.

[0043] In some embodiments, the antibody or the functional fragment thereof or the pharmaceutical
25 composition is cytolytic to a target cell. In some embodiments, the target cell is a cancer cell. In some embodiments, the antibody or the functional fragment thereof or the pharmaceutical composition inhibits tumor growth. In some embodiments, the antibody or the functional fragment thereof or the pharmaceutical composition is administered subcutaneously, intravenously, intradermally, intraperitoneally, orally, intramuscularly or intracranially. In some embodiments, the antibody or the functional fragment thereof or
30 the pharmaceutical composition is administered to the subject in combination with a second therapeutic agent. In some embodiments, the second therapeutic agent comprises an anti-cancer agent, a chemotherapeutic agent, radiation therapy, a cytotoxic agent, a NSAID, a corticosteroid, a dietary supplement such as an antioxidant, or a combination thereof. In some embodiments, the second therapeutic agent is administered

prior to, concurrently, or after administering the antibody or the functional fragment thereof or the pharmaceutical composition.

[0044] Provided herein is an isolated nucleic acid encoding the antibody or the functional fragment thereof of any one of aspects above.

- 5 **[0045]** Provided herein is an isolated nucleic acid molecule encoding a heavy chain polypeptide, the isolated nucleic acid molecule comprising a first nucleic acid sequence encoding a IgA heavy chain constant region, wherein the first nucleic acid sequence is selected from any one of SEQ ID NOs: 25-32. In some embodiments, the isolated nucleic acid molecule aspect above, further comprises a second nucleic acid sequence encoding a variable heavy chain region, wherein the second nucleic acid sequence is selected from
10 any one of SEQ ID NOs:87-84.

[0046] Provided herein is a vector comprising the isolated nucleic acid molecule of any one of aspects above.

- [0047]** Provided herein is a host cell comprising the isolated nucleic acid molecule of any one of aspects above. In some embodiments, the host cell of any one of aspects above, further comprises an isolated nucleic acid molecule encoding a light chain polypeptide, wherein the isolated nucleic acid molecule encoding the
15 light chain polypeptide comprises a nucleic acid sequence encoding a variable light chain region, wherein the nucleic acid sequence is selected from any one of SEQ ID NOs:101-108. In some embodiments, the isolated nucleic acid molecule encoding the light chain polypeptide further comprises a nucleic acid sequence encoding a kappa light chain constant region, wherein the nucleic acid sequence comprises a sequence of SEQ ID NO:32.

- 20 **[0048]** Provided herein is a host cell expressing the antibody or the functional fragment thereof of any one of aspects above. In some embodiments, the host cell is a bacterial cell or a mammalian cell. In some embodiments, the host cell is a CHO cell, or a HEK293 cell.

- [0049]** Provided herein is a method of producing an antibody or the functional fragment thereof, the method comprising: (a) culturing the host cell of any one of aspects above in a medium under conditions permitting
25 expression of a polypeptide encoded by the isolated nucleic acid molecule and assembling of the antibody or the functional fragment thereof; and (b) purifying the antibody or the functional fragment thereof from the cultured host cell or the medium of the host cell. In some embodiments, the purifying is by size exclusion chromatography.

- 30 **[0050]** Provided herein is an antibody or a functional fragment thereof that selectively binds to a CD20 polypeptide or a variant thereof, comprising: (a) an IgA heavy chain constant region, wherein the IgA heavy chain constant region comprises an amino acid sequence selected from any one of SEQ ID NOs:16-21; and (b) a variable heavy chain region, wherein the variable heavy chain region comprises at least one of a complementarity-determining region heavy chain 1 (HC-CDR1) of SEQ ID NO: 33, HC-CDR2 of SEQ ID NO: 41, and HC-CDR3 of SEQ ID NO: 49; (c) a variable light chain region, wherein the variable light chain

region comprises at least one of a complementarity-determining region light chain 1 (LC-CDR1) of SEQ ID NO:57, LC-CDR2 of SEQ ID NO:65, and LC-CDR3 of SEQ ID NO:73; or (d) the variable heavy chain of (b) and the variable light chain of (c). In some embodiments, the variable heavy chain comprises a polypeptide sequence having at least 95% sequence identity to the amino acid sequence of SEQ ID NO: 81, and the variable light chain comprises a polypeptide sequence having at least 95% sequence identity to the amino acid sequence of SEQ ID NO: 95.

[0051] Provided herein is an antibody or a functional fragment thereof that selectively binds to a GD2 protein or a variant thereof, comprising: (a) an IgA heavy chain constant region, wherein the IgA heavy chain constant region comprises an amino acid sequence selected from any one of SEQ ID NOs: 16-21; and (b) a variable heavy chain region, wherein the variable heavy chain region comprises at least one of a complementarity-determining region heavy chain 1 (HC-CDR1) of SEQ ID NO: 34, HC-CDR2 of SEQ ID NO: 42, and HC-CDR3 of SEQ ID NO: 50; (c) a variable light chain region, wherein the variable light chain region comprises at least one of a complementarity-determining region light chain 1 (LC-CDR1) of SEQ ID NO: 58, LC-CDR2 of SEQ ID NO: 66, and LC-CDR3 of SEQ ID NO: 74; or (d) the variable heavy chain of (b) and the variable light chain of (c). In some embodiments, the variable heavy chain comprises a polypeptide sequence having at least 95% sequence identity to the amino acid sequence of SEQ ID NO: 4, and the variable light chain comprises a polypeptide sequence having at least 95% sequence identity to the amino acid sequence of SEQ ID NO: 5.

[0052] Provided herein is an antibody or a functional fragment thereof that selectively binds to a Her2 protein or a variant thereof, comprising: (a) an IgA heavy chain constant region, wherein the IgA heavy chain constant region comprises an amino acid sequence selected from SEQ ID NOs: 16-21; and (b) a variable heavy chain region, wherein the variable heavy chain region comprises at least one of a complementarity-determining region heavy chain 1 (HC-CDR1) of SEQ ID NO: 35, HC-CDR2 of SEQ ID NO: 43, and HC-CDR3 of SEQ ID NO: 51; (c) a variable light chain region, wherein the variable light chain region comprises at least one of a complementarity-determining region light chain 1 (LC-CDR1) of SEQ ID NO: 59, LC-CDR2 of SEQ ID NO: 67, and LC-CDR3 of SEQ ID NO: 75; or (d) the variable heavy chain of (b) and the variable light chain of (c). In some embodiments, the variable heavy chain comprises a polypeptide sequence having at least 95% sequence identity to the amino acid sequence of SEQ ID NO: 82, and the variable light chain comprises a polypeptide sequence having at least 95% sequence identity to the amino acid sequence of SEQ ID NO: 96.

[0053] Provided herein is an antibody or a functional fragment thereof that selectively binds to a gp75 protein or a variant thereof, comprising: (a) an IgA heavy chain constant region, wherein the IgA heavy chain constant region comprises an amino acid sequence selected from SEQ ID NOs: 16-21; and (b) a variable heavy chain region, wherein the variable heavy chain region comprises at least one of a complementarity-

determining region heavy chain 1 (HC-CDR1) of SEQ ID NO: 36, HC-CDR2 of SEQ ID NO: 44, and HC-CDR3 of SEQ ID NO: 52; (c) a variable light chain region, wherein the variable light chain region comprises at least one of a complementarity-determining region light chain 1 (LC-CDR1) of SEQ ID NO: 60, LC-CDR2 of SEQ ID NO: 68, and LC-CDR3 of SEQ ID NO: 76; or (d) the variable heavy chain of (b) and the variable light chain of (c). In some embodiments, the variable heavy chain comprises a polypeptide sequence having at least 95% sequence identity to the amino acid sequence of SEQ ID NO: 83, and the variable light chain comprises a polypeptide sequence having at least 95% sequence identity to the amino acid sequence of SEQ ID NO: 97.

[0054] Provided herein is an antibody or a functional fragment thereof that selectively binds to a CTLA4 protein or a variant thereof, comprising: (a) an IgA heavy chain constant region, wherein the IgA heavy chain constant region comprises an amino acid sequence selected from SEQ ID NOs: 16-21; and (b) a variable heavy chain region, wherein the variable heavy chain region comprises at least one of a complementarity-determining region heavy chain 1 (HC-CDR1) of SEQ ID NO: 37, HC-CDR2 of SEQ ID NO: 45, and HC-CDR3 of SEQ ID NO: 53; (c) a variable light chain region, wherein the variable light chain region comprises at least one of a complementarity-determining region light chain 1 (LC-CDR1) of SEQ ID NO: 61, LC-CDR2 of SEQ ID NO: 69, and LC-CDR3 of SEQ ID NO: 77; or (d) the variable heavy chain of (b) and the variable light chain of (c). In some embodiments, the variable heavy chain comprises a polypeptide sequence having at least 95% sequence identity to the amino acid sequence of SEQ ID NO: 84, and the variable light chain comprises a polypeptide sequence having at least 95% sequence identity to the amino acid sequence of SEQ ID NO: 98.

[0055] Provided herein is an antibody or a functional fragment thereof that selectively binds to a CD47 protein or a variant thereof, comprising: (a) an IgA heavy chain constant region, wherein the IgA heavy chain constant region comprises an amino acid sequence selected from SEQ ID NOs: 16-21; and (b) a variable heavy chain region, wherein the variable heavy chain region comprises at least one of a complementarity-determining region heavy chain 1 (HC-CDR1) of SEQ ID NO: 38, HC-CDR2 of SEQ ID NO: 46, and HC-CDR3 of SEQ ID NO: 54; (c) a variable light chain region, wherein the variable light chain region comprises at least one of a complementarity-determining region light chain 1 (LC-CDR1) of SEQ ID NO: 62, LC-CDR2 of SEQ ID NO: 70, and LC-CDR3 of SEQ ID NO: 78; or (d) the variable heavy chain of (b) and the variable light chain of (c). In some embodiments, the variable heavy chain comprises a polypeptide sequence having at least 95% sequence identity to the amino acid sequence of SEQ ID NO: 85, and the variable light chain comprises a polypeptide sequence having at least 95% sequence identity to the amino acid sequence of SEQ ID NO: 99.

[0056] Provided herein is an antibody or a functional fragment thereof comprising: an antigen binding domain; and a constant domain, wherein the constant domain comprises an IgA2 heavy chain constant region

comprising an IgA2 CH1 region, an IgA2 CH2 region, and an IgA2 CH3 region, and wherein the IgA2 heavy chain constant region comprises: a N45.2G amino acid substitution, a N135Q amino acid substitution, a deletion of C147, and a deletion of Y148, relative to the WT IgA2 heavy chain constant region comprising an amino acid sequence of SEQ ID NO: 1, numbering according to IMGT scheme.

- 5 [0057] Provided herein is an antibody or a functional fragment thereof comprising: an antigen binding domain; and a constant domain, wherein the constant domain comprises an IgA2 heavy chain constant region comprising an IgA2 CH1 region, an IgA2 CH2 region, and an IgA2 CH3 region, and wherein the IgA2 heavy chain constant region comprises: a N45.2G amino acid substitution, a N114T amino acid substitution, a I115L amino acid substitution, a T116S amino acid substitution, an amino acid substitution selected from the group
10 consisting of N15.2G, N15.2Q, and N15.2T, and a deletion of C-terminal amino acids P131-Y148, relative to the WT IgA heavy chain constant region comprising an amino acid sequence of SEQ ID NO: 1, numbering according to the IMGT scheme.

- [0058] In some embodiments, the antibody or the functional fragment thereof of aspect above, further comprises: a C86S amino acid substitution, a P124R amino acid substitution, a deletion of C147, a deletion
15 of Y148, a L15.3I amino acid substitution, a T16S amino acid substitution or a combination thereof,

- [0059] relative to the WT IgA heavy chain constant region comprising an amino acid sequence of SEQ ID NO: 1, numbering according to IMGT scheme. In some embodiments, the IgA2 heavy chain constant region comprises an amino acid sequence that is at least 95% identical to a sequence selected from any one of SEQ ID NOs: 18-21. In some embodiments the antibody or the functional fragment thereof is aglycosylated.

- 20 [0060] In some embodiments the antibody or the functional fragment thereof has increased circulatory half-life than a corresponding IgA comprising one or more of wild type amino acid residues N45.2, N114, I115, T116, N15.2, or a C-terminal amino acid residues P131-Y148, numbering according to the IMGT scheme.

- [0061] In some embodiments the antibody or the functional fragment thereof induces increased antibody dependent cell mediated cytotoxicity (ADCC), compared to a corresponding antibody comprising an IgG
25 heavy chain constant domain. In some embodiments the antibody or the functional fragment thereof exhibits increased thermostability, compared to a corresponding IgA comprising one or more of wild type amino acid residues N45.2, N114, I115, T116, N15.2, or a C-terminal amino acid residues P131-Y148, numbering according to the IMGT scheme, numbering according to the IMGT scheme. In some embodiments the antibody or the functional fragment thereof exhibits decreased glycosylation, compared to a corresponding
30 IgA comprising one or more of wild type amino acid residues N45.2, N114, I115, T116, N15.2, or a C-terminal amino acid residues P131-Y148, numbering according to the IMGT scheme.

INCORPORATION BY REFERENCE

[0062] All publications, patents, and patent applications mentioned in this specification are herein incorporated by reference to the same extent as if each individual publication, patent, or patent application
5 was specifically and individually indicated to be incorporated by reference.

BRIEF DESCRIPTION OF THE FIGURES

[0063] The features of the present disclosure are set forth with particularity in the appended claims. A better understanding of the features and advantages of the present disclosure will be obtained by reference to the following detailed description that sets forth illustrative embodiments, in which the principles of the
10 disclosure are utilized, and the accompanying drawings of which:

[0064] FIG. 1 is an illustration showing the amino acid sequence of IgA2 heavy chain (UniProt Reference No.: A0A0G2JMB2) (SEQ ID NO: 1). The highlighted amino acids depict the amino acids subject to substitution in some embodiments described herein. The underlined sequence indicates the tail piece, which all or part of can be deleted in some embodiments described herein.

15 [0065] FIG. 2 is an illustration showing the amino acid sequence of IgA2 heavy chain (UniProt Reference No.: P01877) (SEQ ID NO: 2). The highlighted amino acids depict the amino acids subject to substitution in some embodiments described herein. The underlined sequence indicates the tail piece, which all or part of can be deleted in some embodiments described herein.

20 [0066] FIG. 3 is an illustration showing the amino acid sequence of a representative IgA2 heavy chain (UniProt Reference No.: A0A286Y5Y5) (SEQ ID NO: 3). The highlighted amino acids depict the amino acids subject to substitution or deletion in some embodiments described herein. In some further embodiments, a corresponding amino acid can be deleted in a comparable IgA antibody or a comparable chimeric antibody with an IgA2 constant region.

25 [0067] FIGs. 4A-4D show a schematic representation of engineered IgA variants relative to a wild type IgA (IgA2(m1)). FIG. 4A shows representation of wild type (WT) IgA; IgA2m1 antibody. The wild type (WT) IgA; IgA2m1 contains three N-glycosylation sites in its CH domains and 1 glycosylation site in its tailpiece.

30 [0068] FIG. 4B shows representation of an engineered IgA3.0+ (plus) variant. The engineered IgA3.0+ variant is generated by engineering IgA2m1 antibody to contain a stabilized heavy and light chain linkage via a CH1-P124R mutation, removal of two free cysteines of which 1 is mutated to serine (CH2-C86S) and the second (CH3-CHS-C147del) is removed by deletion of two final amino acids of the tailpiece. In addition, three N-linked glycosylation sites have been removed by substituting critical amino acids in three N-glycosylation motifs. Mutation in the three N-glycosylation motifs include CH1-N45.2G; CH2-N114T-I115L-T116S; CH3-CHS-N135Q.

[0069] FIG. 4C shows representation of an engineered IgA3.0- or IgA3.0min variant contains deletion of the entire tailpiece (CH3-CHS-P131-Y148del). The IgA3.0min variant contains a stabilized heavy and light chain linkage (CH1-P124R mutation), deletion of entire tailpiece (CH3-CHS-P131-Y148del), lacks two free cysteines of which 1 is mutated to serine (CH2-C86S) and the second (CH3-CHS-C147del) is deletion of the tailpiece. In addition, three N-linked glycosylation sites have been removed by substituting critical amino acids in 2 N-glycosylation motifs i.e., CH1-N45.2G and CH2-N114T-I115L-T116S and deletion of CH3-CHS-N135Qdel by deletion of tailpiece.

[0070] FIG. 4D shows representation of an engineered IgA4.0 variant that contains all the features of the IgA3.0min and further contains a mutation in the final N-linked glycosylation motif CH2-N15.2. Therefore, IgA4.0 variant contains a stabilized heavy and light chain linkage (CH1-P124R mutation), deletion of entire tailpiece (CH3-CHS-P131-Y148del), lacks two free cysteines of which 1 is mutated to serine (CH2-C86S) and the second (CH3-CHS-C147del) is deletion of the tailpiece. In addition, four N-linked glycosylation sites have been removed by substituting critical amino acids in four N-glycosylation motifs (CH1-N45.2G; CH2-N114T-I115L-T116S; CH3-CHS-N135Q; and one of CH2-N15.2G; CH2-N15.2Q; ;CH2-N15.2T; or CH2-N15.2T-L15.3I-T16S). IgA4.0 variant is an aglycosylated IgA.

[0071] FIGs. 5A and 5B demonstrate yield of IgA3.0min antibody variants.

[0072] FIG. 5A shows concentration of anti-Her2 IgA antibody i.e., IgA2-Her2 as determined by ELISA on supernatants of HEK293F cells transfected with different heavy chain (HC):light chain (LC):pAdvantage ratios.

[0073] FIG. 5B shows concentration of anti-CD47 IgA3.0min antibody i.e., IgA3.0min antibody containing anti-CD47 variable domains (IgA3.0min-C47A8-CQ) as determined by ELISA on supernatants of HEK293F cells transfected with different heavy chain (HC):light chain (LC):pAdvantage ratios. The variable domains are obtained from C47A8-CQ antibody.

[0074] FIGs. 6A-6C show comparison of antibody yield of IgA3.0min antibody variants in HEK293F cells versus ExpiCHO-S cells as determined in ELISA.

[0075] FIG. 6A shows comparison of production rate in HEK293F cells versus ExpiCHO cells of anti-CD20 IgA3.0min-Obi antibody. IgA3.0min-Obi antibody contains IgA3.0min with Obinutuzumab (Obi) variable domains.

[0076] FIG. 6B shows comparison of production rate in HEK293F cells versus ExpiCHO cells of anti-Her2 IgA3.0min-Her2 antibody. IgA3.0min-Her2 antibody contains IgA3.0min with anti-Her2 variable domains.

[0077] FIG. 6C shows comparison of production rate in HEK293F cells versus ExpiCHO cells of anti-mCTLA4 IgA3.0min-mCTLA4 antibody. IgA3.0min-mCTLA4 contains IgA3.0min with anti-mCTLA4 variable domains.

- [0078] FIGs. 7A-7D demonstrate concentration of IgA3.0min antibody variants in ExpiCHO-S cells transfected with different heavy chain (HC):light chain (LC):pAdvantage ratios.
- [0079] FIG. 7A shows concentration of anti-GD2 IgA3.0min-ch14.18 antibody. IgA3.0min-ch14.18 antibody contains IgA3.0min with ch14.18 variable domains.
- 5 [0080] FIG. 7B shows concentration of anti-gp75 IgA3.0min-TA99 antibody. IgA3.0min-TA99 antibody contains IgA3.0min with TA99 variable domains.
- [0081] FIG. 7C shows concentration of anti-Her2 IgA3.0min-Her2 antibody. IgA3.0min-Her2 antibody contains IgA3.0min with anti-Her2 variable domains (anti-Her2 variable domains were derived from anti-Her2 antibody Trastuzumab).
- 10 [0082] FIG. 7D shows concentration of anti-CD20 IgA3.0min-Obi antibody. IgA3.0min-Obi antibody contains IgA3.0min with Obinutuzumab (Obi) variable domains.
- [0083] FIGs. 8A-8D show production rate of IgA4.0 antibody variants in ExpiCHO-S cells transfected with different heavy chain (HC):light chain (LC):pAdvantage ratios.
- [0084] FIG. 8A shows production of anti-CD20 IgA4.0_NG-Obi variant. IgA4.0_NG-Obi variant contains
- 15 IgA4.0 with CH2-N15.2G mutation in glycosylation site and variable domains from Obi.
- [0085] FIG. 8B shows production of anti-CD20 IgA4.0_NT-Obi variant. IgA4.0_NT-Obi variant contains IgA4.0 with CH2-N15.2T mutation in glycosylation site and variable domains from Obi.
- [0086] FIG. 8C shows production of anti-CD20 IgA4.0_NQ-Obi variant. IgA4.0_NQ-Obi variant contains IgA4.0 with CH2-N15.2Q mutation in glycosylation site and variable domains from Obi
- 20 [0087] FIG. 8D shows production of anti-CD20 IgA4.0_NLT-TIS-Obi variant. IgA4.0_NLT-TIS-Obi variant contains IgA4.0 with N15.2T-L15.3I-T16S mutation in glycosylation site and variable domains from Obi.
- [0088] FIG. 9A-9B show elution profiles of IgA3.0min-Obi using a KappaSelect column (GE Healthcare) from supernatants from the producing cell lines ExpiCHO-S (FIG. 9A) and HEK293F (FIG.9B).
- [0089] FIGs. 9C-9D show SEC separation profiles of IgA3.0min-Obi from the producing cell lines
- 25 ExpiCHO-S (FIG. 9C) and HEK293F (FIG. 9D), no aggregates are observed upon SEC separation.
- [0090] FIGs. 10A-10B show elution profiles of IgA3.0min-Her2 using a KappaSelect column (GE Healthcare) from supernatants from the producing cell lines ExpiCHO-S (FIG. 10A) and HEK293F (FIG.10B).
- [0091] FIGs. 10C-10D show SEC elution profiles of IgA3.0min-Her2 from the producing cell lines
- 30 ExpiCHO-S (FIG. 10C) and HEK293F (FIG. 10D), no aggregates are observed upon SEC.
- [0092] FIG. 11A shows typical elution profiles of IgA4.0-Obi using a KappaSelect column (GE Healthcare) from supernatants from the producing cell line ExpiCHO-S.
- [0093] FIG. 11B shows SEC elution profile of IgA4.0-Obi from the producing cell line ExpiCHO-S, no aggregates are observed upon SEC.

- [0094] FIGs. 12A-12E shows binding of IgA3.0+-Obi and IgA3.0min-Obi to CD20 positive Daudi cells by flow cytometry.
- [0095] FIG. 12A shows unstained CD20+Daudi cells control.
- [0096] FIG. 12B shows Daudi cells stained with secondary antibody only negative control.
- 5 [0097] FIG. 12C shows stained with anti-CD20 IgA Obi (5ug/mL) positive control.
- [0098] FIG. 12D shows Daudi cells stained with IgA3.0+-Obi (supernatant). FIG. 12E shows Daudi cells stained with IgA3.0min-Obi (supernatant). FIGs. 12D-12E shows Daudi cells stained with supernatants from HEK293F cells transfected with IgA3.0min-Obi or IgA3.0+-Obi. Both variants IgA3.0+-Obi (FIG. 12D) and IgA3.0min-Obi (FIG. 12E) bind to CD20 positive Daudi to the same extent as IgA Obi (FIG. 12C).
- 10 [0099] FIG. 13 shows the binding of the Fc-part to PMN by IgA3.0min-Obi as assessed by a PMN binding assay. PMNs were added to ELISA plate wells coated with antibody in different concentrations. A series of washing determines the binding strength of the Fc par to PMN of IgA3.0min-Obi and IgA2. After the 6th wash binding is plotted. IgA3.0min-Obi shows equal to better binding to PMNs as compared to wild type IgA2.
- 15 [0100] FIG. 14 shows equal coating concentrations of IgA3.0min compared to wild type IgA2.
- [0101] FIG. 15 shows binding analysis of IgA4.0 variants. Supernatants from ExpiCHO-S cells transfected with IgA4.0 variants; IgA4.0_NT-Obi, IgA4.0_NQ-Obi, IgA4.0_NG-Obi, and IgA4.0_NLT-TIS-Obi, were assessed for binding on CD20 expressing SKBR3 cells. All variants of IgA4.0-Obi bind to SKBR3-CD20 to the same extent as supernatant from ExpiCHO-S cells transfected with IgA3.0min-Obi or purified IgA3.0min-Obi.
- 20 [0102] FIGs. 16A-16D shows IgA variants induce PMN-mediated ADCC against target cells.
- [0103] FIG. 16A shows IgA3.0min-Her2 antibody induces ADCC to a similar or better extent than IgA2-Her2 of SKBR3 cells as determined by chromium release assay.
- [0104] FIGs. 16B-16C show IgA3.0min-Obi antibody produced by HEK293F cells or ExpiCHO-S cells induce similar level of ADCC of Ramos cells (FIG. 16B) and Daudi cells (FIG. 16C) as assessed by chromium release assay.
- 25 [0105] FIG. 16D shows induction of ADCC by IgA variants against Daudi cells. Purified IgA3.0min-Obi, supernatant from cells transfected with IgA3.0min-Obi, IgA4.0_NG-Obi, IgA4.0_NQ-Obi, IgA4.0_NT-Obi, and IgA4.0_NLT-TIS-Obi show increased ADCC compared to purified IgG1-Obi. IgA4.0-obi variants; IgA4.0_NG-Obi, IgA4.0_NQ-Obi, IgA4.0_NT-Obi, and IgA4.0_NLT-TIS-Obi show similar levels of ADCC induction compared to Purified IgA3.0min-Obi, supernatant from cells transfected with IgA-3.0min-Obi.
- 30 [0106] FIGs. 17A-17C show thermostability of IgA variants. Using a SYPRO Orange thermal shift assay the thermostability of IgA3.0min and IgA4.0 has been determined. FIG. 17A demonstrates the engineered IgA3.0min variants; IgA3.0min with Obi variable domains and IgA3.0min with 2.3D11 variable domains

showed increased thermostability compared to wild type IgA2(m1), i.e., IgA2 with ch14.18 variable domains and IgA2 with Her2 variable domains.

[0107] FIG. 17A also demonstrates the engineered IgA4.0min variants; IgA4.0 NQ, IgA4.0 NT, IgA4.0 NLT-TIS, and IgA4.0 NG all with Obi variable domains showed increased thermostability compared to wild type IgA2(m1), i.e., IgA2 with ch14.18 variable domains and IgA2 with Her2 variable domains. The increased thermostability of engineered IgA3.0min variants and engineered IgA4.0 variants relative wild type IgA2(m1) is indicated by the shift to the higher temperatures.

[0108] FIG. 17B shows a plot of average T_m values of analysed engineered IgA3.0min variants and engineered IgA4.0 variants relative to wild type IgA2(m1). The plot shows engineered IgA3.0min variants and engineered IgA4.0 variants are more thermostable compared to wild type IgA2.

[0109] FIG. 17C shows antibodies exposed to increasing temperatures tested for their functionality in a PMN-ADCC against CD20 expressing Raji cells. Wild type IgA2(m1) shows a decremental efficacy from 47°C on, while both IgA3.0min-Obi and IgA4.0_NT-Obi are still effective. At 71°C both IgA3.0min-Obi as well as IgA4.0_NT-Obi give more than 50% killing efficacy while IgA2(m1) is dysfunctional at this temperature.

[0110] FIG. 18 shows effects of deglycosylation of engineered IgA3.0min variants and engineered IgA4.0 variant compared to wild type IgA2 by PNGase F treatment. Wild type IgA2; IgA2(m1)-UMAB10 shows the largest shift, whereas engineered IgA3.0min variant; IgA3.0min-Obi shows a minor shift indicating reduced glycosylation compared to wild type IgA2. Engineered IgA4.0 variant; IgA4.0_NG-Obi does not show a shift indicating reduced glycosylation compared to engineered IgA3.0min variant and wild type IgA2.

[0111] FIGs. 19A-19G show total glycosylation profile of wild type IgA2 and engineered IgA variants as determined with MALDI-TOF-MS.

[0112] FIG. 19A shows total glycosylation profile of HEK293 cells produced wild type IgA2; IgA2-Her2.

[0113] FIG. 19B shows total glycosylation profile of HEK293 cells produced engineered IgA3.0min variant; IgA3.0min-Her2. The profile shows all signals from a single glycosylation site present in engineered IgA3.0min variant.

[0114] FIG. 19C shows glycosylation site specific analysis of engineered IgA3.0min variant; IgA3.0min-Obi produced in HEK293F and ExpiCHO-S. IgA3.0min-Obi produced in ExpiCHO-S shows less free galactoses implying to be less susceptible to ASGPR-dependent clearance in the liver.

[0115] FIGs. 19D-19G show native MS analysis of four IgA4.0 variants; IgA4.0_NG-Obi (FIG. 19D), IgA4.0_NQ-Obi (FIG. 19E), IgA4.0_NT-Obi (FIG. 19F), and IgA4.0_NLT-TIS-Obi (FIG. 19G) Large plot displays broad mass range of antibody, inset zooms in on highest peak. Observed mass is virtually equal to theoretical mass, excluding presence of bulky N-glycans, indicating absence of glycosylation in IgA4.0 variants.

[0116] FIGs. 20A-20B show pharmacokinetics and pharmacodistribution analysis of engineered IgA variants. An amount of 100 µg of wild-type-IgG1-dinutuximab, wild type-IgA2-dinutuximab and IgA3.0min-dinutuximab, IgA3.0min-Obinituzumab, IgA4.0-Obinituzumab were injected into BALB/c mice and blood was analysed at indicated time points by ELISA.

5 [0117] FIG. 20A shows that IgA3.0min variant; IgA3.0min-dinutuximab showed extended half-life as compared to its wild type IgA2 counterpart.

[0118] FIG. 20B shows comparison between IgA3.0min and IgA4.0 in an Obinituzumab format. Engineered IgA4.0 variant; IgA4.0-obinituzumab showed a better half-life profile than engineered IgA3.0min variant; IgA3.0min-Obinituzumab. Obinituzumab IgA4.0 can still be detected after 120 hours.

10 [0119] FIGs. 21A-21B shows biodistribution of IgG1 dinutuximab and engineered IgA3.0min variant, IgA3.0min dinutuximab that contains variable domains from dinutuximab. Indium-111 radiolabeled IgG1 dinutuximab and engineered IgA3.0min variant, IgA3.0min dinutuximab were i.v. injected into mice for biodistribution analysis. Mice were monitored to trace antibody distribution after 24 hours (FIG. 21A) and after 48 hours (FIG. 21B). A clear infiltration of both IgG1 and engineered IgA3.0min variants to the tumor is
15 observed.

[0120] FIG. 22 shows quantification of the distribution of radiolabeled of IgG1 dinutuximab and engineered IgA3.0min variant, IgA3.0min dinutuximab distribution in mice from FIGs. 21A-21B. The ratio of tumor signal to liver signal has been determined to adjust for background signals. Plot shows engineered IgA3.0min variant, IgA3.0min dinutuximab showed increased tumor/liver ratio compared to the IgG1 dinutuximab.

20 [0121] FIG. 23 shows inhibition of tumor growth in an established tumor model upon administration of engineered IgA3.0min variant; IgA3.0min-Her2. A431-Luc2-Her2 cells were injected intraperitoneally on day 0 in hCD89 Transgenic or non-transgenic SCID mice followed by a subcutaneous pegG-CSF injection on day 6. Bioluminescence signal was measured from day 6 onwards and mice were randomized into different treatment groups on this day, and significant signal was measured. Treatment started on day 7 with daily
25 intraperitoneal injections of 10ug IgA3.0min-Her2 for 10 days. Bioluminescence signal was measured on the indicated time points.

DETAILED DESCRIPTION OF THE DISCLOSURE

[0122] The following description and examples illustrate embodiments of the present disclosure in detail. It is to be understood that this disclosure is not limited to the particular embodiments described herein and as
30 such can vary. Those of skill in the art will recognize that there are numerous variations and modifications of this disclosure, which are encompassed within its scope.

[0123] All terms are intended to be understood as they would be understood by a person skilled in the art. Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which the disclosure pertains.

5 [0124] The section headings used herein are for organizational purposes only and are not to be construed as limiting the subject matter described.

10 [0125] Although various features of the present disclosure may be described in the context of a single embodiment, the features can also be provided separately or in any suitable combination. Conversely, although the present disclosure may be described herein in the context of separate embodiments for clarity, the present disclosure can also be implemented in a single embodiment.

Definitions

15 [0126] The following definitions supplement those in the art and are directed to the current application and are not to be imputed to any related or unrelated case, *e.g.*, to any commonly owned patent or application. Although any methods and materials similar or equivalent to those described herein can be used in the practice for testing of the present disclosure, the preferred materials and methods are described herein. Accordingly, the terminology used herein is for the purpose of describing particular embodiments only, and is not intended to be limiting.

20 [0127] In this application, the use of the singular includes the plural unless specifically stated otherwise. It must be noted that, as used in the specification, the singular forms “a,” “an” and “the” include plural referents unless the context clearly dictates otherwise. In this application, the use of “or” means “and/or” unless stated otherwise. Furthermore, use of the term “including” as well as other forms, such as “include,” “includes,” and “included,” is not limiting.

25 [0128] Reference in the specification to “some embodiments,” “an embodiment,” “one embodiment” or “other embodiments” means that a particular feature, structure, or characteristic described in connection with the embodiments is included in at least some embodiments, but not necessarily all embodiments, of the present disclosure.

30 [0129] As used in this specification and claim(s), the words “comprising” (and any form of comprising, such as “comprise” and “comprises”), “having” (and any form of having, such as “have” and “has”), “including” (and any form of including, such as “includes” and “include”) or “containing” (and any form of containing, such as “contains” and “contain”) are inclusive or open-ended and do not exclude additional, un-recited elements or method steps. It is contemplated that any embodiment discussed in this specification can be implemented with respect to any method or composition of the present disclosure, and vice versa.

Furthermore, compositions of the present disclosure can be used to achieve methods of the present disclosure.

[0130] The term “about” or “approximately” means within an acceptable error range for the particular value as determined by one of ordinary skill in the art, which will depend in part on how the value is measured or determined, *i.e.*, the limitations of the measurement system. For example, “about” can mean within 1 or more than 1 standard deviation, per the practice in the art. Alternatively, “about” can mean a range of up to 20%, up to 10%, up to 5%, or up to 1% of a given value. In another example, the amount “about 10” includes 10 and any amounts from 9 to 11.

[0131] In yet another example, the term “about” in relation to a reference numerical value can also include a range of values plus or minus 10%, 9%, 8%, 7%, 6%, 5%, 4%, 3%, 2%, or 1% from that value. Alternatively, particularly with respect to biological systems or processes, the term “about” can mean within an order of magnitude, preferably within 5-fold, and more preferably within 2-fold, of a value. Where particular values are described in the application and claims, unless otherwise stated the term “about” meaning within an acceptable error range for the particular value should be assumed.

[0132] As used herein, the term “antibody” refers to an immunoglobulin (Ig) whether natural or partly or wholly synthetically produced. The term also covers any polypeptide or protein having a binding domain which is, or is homologous to, an antigen-binding domain. The term further includes “antigen-binding fragments” or “functional fragment thereof”, or “fragment of an antibody”, “antibody fragment”, “functional fragment of an antibody” and other interchangeable terms for similar binding fragments such as described below.

[0133] An antibody includes, for example, monoclonal antibodies, chimeric antibodies, humanized antibodies, human antibodies, recombinant antibodies, chemically engineered antibodies, deimmunized antibodies, affinity-matured antibodies, multispecific antibodies (for example, bispecific antibodies and polyreactive antibodies), heteroconjugate antibodies, antibody fragments, and combinations thereof (e.g., a monoclonal antibody that is also deimmunized, a humanized antibody that is also deimmunized, etc.).

[0134] An antibody can be, for example, murine, chimeric, humanized, heteroconjugate, bispecific, diabody, triabody, or tetrabody. The antigen binding fragment can include, for example, Fab’, F(ab’)2, Fab, Fv, rIgG, scFv, hcAbs (heavy chain antibodies), a single domain antibody, VHH, VNAR, sdAbs, or nanobody.

[0135] The term “monoclonal antibodies,” as used herein, refers to antibodies that are produced by a single clone of B-cells and bind to the same epitope. In contrast, “polyclonal antibodies” refer to a population of antibodies that are produced by different B-cells and bind to different epitopes of the same antigen. A whole antibody typically consists of four polypeptides: two identical copies of a heavy (H) chain polypeptide and two identical copies of a light (L) chain polypeptide. Each of the heavy chains contains one N-terminal variable (VH) region and three C-terminal constant (CH1, CH2 and CH3) regions, and each light chain contains one N-terminal variable (VL) region and one C-terminal constant (CL) region. The variable regions of each pair of light and heavy chains form the antigen binding site of an antibody. The VH and VL regions

have a similar general structure, with each region comprising four framework regions, whose sequences are relatively conserved. The framework regions are connected by three complementarity determining regions (CDRs). The three CDRs, known as CDR1, CDR2, and CDR3, form the “hypervariable region” of an antibody, which is responsible for antigen binding.

5 [0136] As used herein a “chimeric antibody” is an antibody that comprises an amino acid sequence derived from two different species or, or two different sources, and includes synthetic molecules. By way of non-limiting example, an antibody that comprises a non-human CDR and a human variable region framework or constant or Fc region, an antibody with binding domains from two different monoclonal antibodies, or an antibody comprising a mutation of one or more amino acid residues to increase or decrease biological activity
10 or binding of a part of the antibody. In certain embodiments, recombinant antibodies are produced from a recombinant DNA molecule or synthesized. In certain embodiments, the antibodies described herein are a polypeptide(s) encoded by one or more polynucleotides.

[0137] As used herein, “recognize” refers to the association or binding between an antigen binding domain and an antigen. As used herein, an “antigen” refers to an antigenic substance that can trigger an immune
15 response in a host. An antigenic substance can be a molecule, such as a costimulatory molecule that can trigger an immune response in a host.

[0138] As used herein, an “antibody construct” refers to a construct that contains an antigen binding domain and an Fc domain.

[0139] As used herein, a “binding domain” refers to an antibody or non-antibody domain.

20 [0140] As used herein, an “antigen binding domain” refers to a binding domain from an antibody or from a non-antibody that can bind to an antigen. An antigen binding domain can be a tumor antigen binding domain or a binding domain that can bind to an antigen (such as a molecule) on an antigen presenting cell. Antigen binding domains can be numbered when there is more than one antigen binding domain in a given conjugate or antibody construct (e.g., first antigen binding domain, second antigen binding domain, third antigen
25 binding domain, etc.). Different antigen binding domains in the same conjugate or construct can target the same antigen or different antigens (e.g., first antigen binding domain that can bind to a tumor antigen, second antigen binding domain that can bind to a molecule on an antigen presenting cell (APC antigen), and third antigen binding domain that can bind to an APC antigen). The term “antigen binding domain” refers to a fragment of an antibody that comprises the area which specifically binds to an epitope, and is complementary
30 to part or all of an antigen. An antigen binding domain may be provided by, for example, one or more antibody variable domains (also called antibody variable regions). Particularly, an antigen binding domain comprises an antibody light chain variable region (VL) and an antibody heavy chain variable region (VH).

[0141] As used herein, the term “antigen” means a molecule or portion of a molecule that can react with a recognition site on an antibody. The term “antigen” also includes a molecule or a portion of a molecule that

can, either by itself or in conjunction with an adjuvant or carrier, elicit an immune response (also called an “immunogen”). The term “antigen” as used herein includes molecules or portions of molecules (epitopes) that can elicit production of antibodies or that can bind to antibodies. The term includes materials that react strongly and with high specificity, and also includes materials that react weakly and/or with low affinity to an antibody.

[0142] The term “epitope,” as used herein, refers to an antigenic determinant that interacts with a specific antigen binding site in the variable region of an antibody molecule known as a paratope. A single antigen may have more than one epitope. Thus, different antibodies may bind to different areas on an antigen and may have different biological effects. Epitopes may be either conformational or linear. A conformational epitope is produced by spatially juxtaposed amino acids from different segments of the linear polypeptide chain. A linear epitope is one produced by adjacent amino acid residues in a polypeptide chain. In certain circumstance, an epitope may include moieties of saccharides, phosphoryl groups, or sulfonyl groups on the antigen. Various techniques known to persons of ordinary skill in the art can be used to determine whether an antigen-binding domain of an antibody “interacts with one or more amino acids” within a polypeptide or protein. Exemplary techniques include, e.g., routine cross-blocking assay such as that described Antibodies, Harlow and Lane (Cold Spring Harbor Press, Cold Spring Harb., NY), alanine scanning mutational analysis, peptide blots analysis (Reineke, 2004, *Methods Mol Biol* 248:443-463), and peptide cleavage analysis. In addition, methods such as epitope excision, epitope extraction and chemical modification of antigens can be employed (Tomer, 2000, *Protein Science* 9:487-496). Another method that can be used to identify the amino acids within a polypeptide with which an antigen-binding domain of an antibody interacts is hydrogen/deuterium exchange detected by mass spectrometry. In general terms, the hydrogen/deuterium exchange method involves deuterium-labeling the protein of interest, followed by binding the antibody to the deuterium-labeled protein. Next, the protein/antibody complex is transferred to water to allow hydrogen-deuterium exchange to occur at all residues except for the residues protected by the antibody (which remain deuterium-labeled). After dissociation of the antibody, the target protein is subjected to protease cleavage and mass spectrometry analysis, thereby revealing the deuterium-labeled residues, which correspond to the specific amino acids with which the antibody interacts. See, e.g., Ehring (1999) *Analytical Biochemistry* 267(2):252-259; Engen and Smith (2001) *Anal. Chem.* 73:256A-265A. X-ray crystallography of the antigen/antibody complex may also be used for epitope mapping purposes.

[0143] As used herein, an “antibody antigen binding domain” refers to a binding domain from an antibody that can bind to an antigen.

[0144] As used herein, an “Fc domain” refers to an Fc domain from an antibody or from a non- antibody that can bind to an Fc receptor. As used herein, an “Fc domain” and an “Fc comprising domain” can be used interchangeably.

[0145] As used herein, a “target binding domain” refers to a construct that contains an antigen binding domain from an antibody or from a non-antibody that can bind to an antigen.

[0146] As used herein, the abbreviations for the natural L- enantiomeric amino acids are conventional and can be as follows: alanine (A, Ala); arginine (R, Arg); asparagine (N, Asn); aspartic acid (D, Asp); cysteine (C, Cys); glutamic acid (E, Glu); glutamine (Q, Gln); glycine (G, Gly); histidine (H, His); isoleucine (I, Ile); leucine (L, Leu); lysine (K, Lys); methionine (M, Met); phenylalanine (F, Phe); proline (P, Pro); serine (S, Ser); threonine (T, Thr); tryptophan (W, Trp); tyrosine (Y, Tyr); valine (V, Val). Unless otherwise specified, X can indicate any amino acid. In some aspects, X can be asparagine (N), glutamine (Q), histidine (H), lysine (K), or arginine (R).

10 [0147] The phrase “pharmaceutically acceptable” is employed herein to refer to those compounds, materials, compositions, and/or dosage forms which are, within the scope of sound medical judgment, suitable for use in contact with the tissues of human beings and animals without excessive toxicity, irritation, allergic response, or other problem or complication, commensurate with a reasonable benefit/risk ratio.

[0148] The phrase “pharmaceutically acceptable excipient” or “pharmaceutically acceptable carrier” as used
15 herein means a pharmaceutically acceptable material, composition or vehicle, such as a liquid or solid filler, diluent, excipient, solvent or encapsulating material. Each carrier must be “acceptable” in the sense of being compatible with the other ingredients of the formulation and not injurious to the patient. Some examples of materials which can serve as pharmaceutically acceptable carriers include: (1) sugars, such as lactose, glucose and sucrose; (2) starches, such as corn starch and potato starch; (3) cellulose, and its derivatives, such as
20 sodium carboxymethyl cellulose, ethyl cellulose and cellulose acetate; (4) powdered tragacanth; (5) malt; (6) gelatin; (7) talc; (8) excipients, such as cocoa butter and suppository waxes; (9) oils, such as peanut oil, cottonseed oil, safflower oil, sesame oil, olive oil, corn oil and soybean oil; (10) glycols, such as propylene glycol; (11) polyols, such as glycerin, sorbitol, mannitol and polyethylene glycol; (12) esters, such as ethyl oleate and ethyl laurate; (13) agar; (14) buffering agents, such as magnesium hydroxide and aluminum
25 hydroxide; (15) alginic acid; (16) pyrogen- free water; (17) isotonic saline; (18) Ringer's solution; (19) ethyl alcohol; (20) phosphate buffer solutions; and (21) other non-toxic compatible substances employed in pharmaceutical formulations.

[0149] The terms “cancer,” “tumor,” “proliferative disease,” “malignancy,” or “malignant disease” relate to
30 diseases in which a group of cells display uncontrolled growth or unwanted growth. Cancer cells can also spread to other locations, which can lead to the formation of metastases. Spreading of cancer cells in the body can, for example, occur via lymph or blood. Uncontrolled growth, intrusion and metastasis formation are also termed malignant properties of cancers. These malignant properties differentiate cancers from benign tumors, which typically do not invade or metastasize.

[0150] “Antigen recognition moiety” or “antibody recognition domain” refers to a molecule or portion of a molecule that specifically binds to an antigen. In one embodiment, the antigen recognition moiety is an antibody, antibody like molecule or fragment thereof and the antigen is a tumor antigen or an infectious disease antigen.

5 [0151] The terms “fragment of an antibody,” “antibody fragment,” “functional fragment of an antibody,” “antigen-binding portion” or their grammatical equivalents are used interchangeably herein to mean one or more fragments or portions of an antibody that retain the ability to specifically bind to an antigen (see, generally, Holliger et al., *Nat. Biotech.*, 23(9):1126-1129 (2005)). The antibody fragment desirably comprises, for example, one or more CDRs, the variable region (or portions thereof), the constant region (or portions thereof), or combinations thereof. Examples of antibody fragments include, but are not limited to, (i) 10 a Fab fragment, which is a monovalent fragment consisting of the VL, VH, CL, and CH1 domains; (ii) a F(ab')₂ fragment, which is a bivalent fragment comprising two Fab fragments linked by a disulfide bridge at the stalk region; (iii) a Fv fragment consisting of the VL and VH domains of a single arm of an antibody; (iv) a single chain Fv (scFv), which is a monovalent molecule consisting of the two domains of the Fv fragment 15 (i.e., VL and VH) joined by a synthetic linker which enables the two domains to be synthesized as a single polypeptide chain (see, e.g., Bird et al., *Science*, 242: 423-426 (1988); Huston et al., *Proc. Natl. Acad. Sci. USA*, 85: 5879-5883 (1988); and Osbourn et al., *Nat. Biotechnol.*, 16: 778 (1998)) and (v) a diabody, which is a dimer of polypeptide chains, wherein each polypeptide chain comprises a VH connected to a VL by a peptide linker that is too short to allow pairing between the VH and VL on the same polypeptide chain, 20 thereby driving the pairing between the complementary domains on different VH-VL polypeptide chains to generate a dimeric molecule having two functional antigen binding sites. Antibody fragments are known in the art and are described in more detail in, e.g., U.S. Pat. No. 8,603,950. Other antibody fragments can include variable fragments of heavy chain antibodies (VHH).

[0152] The term “conservative amino acid substitution” or “conservative mutation” refers to the replacement 25 of one amino acid by another amino acid with a common property. A functional way to define common properties between individual amino acids is to analyze the normalized frequencies of amino acid changes between corresponding proteins of homologous organisms (Schulz, G. E. and Schirmer, R. H., *Principles of Protein Structure*, Springer-Verlag, New York (1979)). According to such analyses, groups of amino acids may be defined where amino acids within a group exchange preferentially with each other, and therefore 30 resemble each other most in their impact on the overall protein structure (Schulz, G. E. and Schirmer, R. H., *supra*). Examples of conservative mutations include amino acid substitutions of amino acids within the sub-groups above, for example, lysine for arginine and vice versa such that a positive charge may be maintained; glutamic acid for aspartic acid and vice versa such that a negative charge may be maintained; serine for threonine such that a free –OH can be maintained; and glutamine for asparagine such that a free –NH₂ can be

maintained. Alternatively or additionally, the therapeutic IgA antibodies can comprise the amino acid sequence of the reference protein with at least one non-conservative amino acid substitution.

[0153] The terms “non-conservative mutation” or “non-conservative amino acid substitution” involve amino acid substitutions between different groups, for example, lysine for tryptophan, or phenylalanine for serine, etc. In this case, it is preferable for the non-conservative amino acid substitution to not interfere with, or inhibit the biological activity of the therapeutic IgA antibody. The non-conservative amino acid substitution may enhance the biological activity of the therapeutic IgA antibody, such that the biological activity of the therapeutic IgA antibody is increased as compared to the wild type therapeutic IgA antibody.

[0154] As used herein, “humanized” antibodies refer to forms of non-human (e.g. murine) antibodies that are specific chimeric immunoglobulins, immunoglobulin chains, or fragments thereof (such as Fv, Fab, Fab', F(ab')₂ or other antigen-binding subsequences of antibodies) that contain minimal sequence derived from non-human immunoglobulin. For the most part, humanized antibodies are human immunoglobulins (recipient antibody) in which residues from a complementary determining region (CDR) of the recipient are replaced by residues from a CDR of a non-human species (donor antibody) such as mouse, rat, or rabbit having the desired specificity, affinity, and capacity. In some instances, Fv framework region (FR) residues of the human immunoglobulin are replaced by corresponding non-human residues. Furthermore, the humanized antibody may comprise residues that are found neither in the recipient antibody nor in the imported CDR or framework sequences but are included to further refine and optimize antibody performance. In general, the humanized antibody will comprise substantially all of at least one, and typically two, variable domains, in which all or substantially all of the CDR regions correspond to those of a non-human immunoglobulin and all or substantially all of the FR regions are those of a human immunoglobulin consensus sequence. The humanized antibody optimally also will comprise at least a portion of an immunoglobulin constant region or domain (Fc), typically that of a human immunoglobulin. Other forms of humanized antibodies have one or more CDRs (one, two, three, four, five, six) which are altered with respect to the original antibody, which are also termed one or more CDRs “derived from” one or more CDRs from the original antibody.

[0155] As used herein, an “isolated antibody” is one that has been separated and/or recovered from a component of its natural environment. Contaminant components of its natural environment are materials that would interfere with diagnostic or therapeutic uses of the antibody, and may include enzymes, hormones, and other proteinaceous or non-proteinaceous components. In preferred embodiments, the antibody is purified: (1) to greater than 95% by weight of antibody as determined by the Lowry method, and most preferably more than 99% by weight; (2) to a degree sufficient to obtain at least 15 residues of N-terminal or internal amino acid sequence by use of a spinning cup sequenator; or (3) to homogeneity as shown by SDS-PAGE under reducing or non-reducing conditions and using Coomassie blue or, preferably, silver staining. Isolated antibody includes the antibody in situ within recombinant cells since at least one component of the antibody's

natural environment will not be present. Ordinarily, however, isolated antibody will be prepared by at least one purification step.

[0156] By "substantially purified" is meant that the antibody, or the functional fragment thereof is substantially free of cellular material or other contaminating proteins from the cell or tissue source from which it is derived, or is substantially free from chemical precursors or other chemicals when chemically synthesized. The language includes preparations of an antibody, which is separated from cellular components of the cells from which it is isolated or recombinantly produced. Thus, an antibody, that is substantially free of cellular material includes preparations having less than about 30%, 20%, 10% or 5% (by dry weight) of contaminating protein and culture medium. In some embodiments, the antibody can be purified by chromatography, for example, size exclusion chromatography or ion exchange chromatography.

[0157] As used herein, the term "Complementarity Determining Regions" (CDRs, i.e., CDR1, CDR2, and CDR3) refers to the amino acid residues of an antibody variable domain the presence of which are necessary for antigen binding. Each variable domain typically has three CDR regions identified as CDR1, CDR2 and CDR3. The CDRs of variable heavy chain can be CDR-H1, CDR-H2 and CDR-H3. The CDRs of variable light chain can be CDR-L1, CDR-L2 and CDR-L3. Exemplary hypervariable loops occur at amino acid residues 26-32 (L1), 50-52 (L2), 91-96 (L3), 26-32 (H1), 53-55 (H2), and 96-101 (H3). (Chothia and Lesk, J. Mol. Biol. 196:901-917 (1987)). Exemplary CDRs (CDR-L1, CDR-L2, CDR-L3, CDR-H1, CDR-H2, and CDR-H3) occur at amino acid residues 24-34 of L1, 50-56 of L2, 89-97 of L3, 31-35B of H1, 50-65 of H2, and 95-102 of H3 (Kabat et al., Sequences of Proteins of Immunological Interest, 5th ed. (1991)). Thus, the HVs may be comprised within the corresponding CDRs and references herein to the "hypervariable loops" of VH and VL domains should be interpreted as also encompassing the corresponding CDRs, and vice versa, unless otherwise indicated. The more highly conserved regions of variable domains are called the framework region (FR), as defined below. The variable domains of native heavy and light chains each comprise four FRs (FR1, FR2, FR3 and FR4, respectively), largely adopting a [beta]-sheet configuration, connected by the three hypervariable loops. The hypervariable loops in each chain are held together in close proximity by the FRs and, with the hypervariable loops from the other chain, contribute to the formation of the antigen-binding site of antibodies. Structural analysis of antibodies revealed the relationship between the sequence and the shape of the binding site formed by the complementarity determining regions (Chothia et al., J. Mol. Biol. 227: 799-817 (1992)); Tramontano et al., J. Mol. Biol, 215: 175-182 (1990)). Despite their high sequence variability, five of the six loops adopt just a small repertoire of main-chain conformations, called "canonical structures". These conformations are first of all determined by the length of the loops and secondly by the presence of key residues at certain positions in the loops and in the framework regions that determine the conformation through their packing, hydrogen bonding or the ability to assume unusual main-chain conformations.

[0158] A “variable region” of an antibody refers to the variable region of the antibody light chain or the variable region of the antibody heavy chain, either alone or in combination. The variable regions of the heavy and light chain each consist of four framework regions (FR) connected by three complementarity determining regions (CDRs) also known as hypervariable regions. The CDRs in each chain are held together in close proximity by the FRs and, with the CDRs from the other chain, contribute to the formation of the antigen-binding site of antibodies. There are at least two techniques for determining CDRs: (1) an approach based on cross-species sequence variability (i.e., Kabat et al. Sequences of Proteins of Immunological Interest, (5th ed., 1991, National Institutes of Health, Bethesda Md.)); and (2) an approach based on crystallographic studies of antigen-antibody complexes (Allazikani et al (1997) J. Molec. Biol. 273:927-948)). A CDR may refer to CDRs defined by either approach or by a combination of both approaches.

[0159] A “constant region” of an antibody refers to the constant region of the antibody light chain, i.e., a light chain constant region or the constant region of the antibody heavy chain, i.e., a heavy chain constant region either alone or in combination. The constant region does not vary with respect to antigen specificity.

[0160] As used herein, the term “heavy chain region” includes amino acid sequences derived from the constant domains of an immunoglobulin heavy chain. A polypeptide comprising a heavy chain region comprises at least one of: a CH1 domain, a hinge (e.g., upper, middle, and/or lower hinge region) domain, a CH2 domain, a CH3 domain, or a variant or fragment thereof. In an embodiment, an antibody or an antigen-binding fragment thereof may comprise the Fc region of an immunoglobulin heavy chain (e.g., a hinge portion, a CH2 domain, and a CH3 domain). In another embodiment, an antibody or an antigen-binding fragment thereof lacks at least a region of a constant domain (e.g., all or part of a CH2 domain). In certain embodiments, at least one, and preferably all, of the constant domains are derived from a human immunoglobulin heavy chain. For example, in one preferred embodiment, the heavy chain region comprises a fully human hinge domain. In other preferred embodiments, the heavy chain region comprising a fully human Fc region (e.g., hinge, CH2 and CH3 domain sequences from a human immunoglobulin). In certain embodiments, the constituent constant domains of the heavy chain region are from different immunoglobulin molecules. For example, a heavy chain region of a polypeptide may comprise a domain derived from an IgA molecule and a hinge region derived from an IgA1 or IgA2 molecule. In other embodiments, the constant domains are chimeric domains comprising regions of different immunoglobulin molecules. For example, a hinge may comprise a first region from an IgA1 molecule and a second region from an IgA2 molecule. As set forth above, it will be understood by one of ordinary skill in the art that the constant domains of the heavy chain region may be modified such that they vary in amino acid sequence from the naturally occurring (wild-type) immunoglobulin molecule. That is, the polypeptides of the invention disclosed herein may comprise alterations or modifications to one or more of the heavy chain constant domains (CH1, hinge, CH2 or CH3) and/or to the light chain constant domain (CL). Exemplary modifications include additions, deletions or

substitutions of one or more amino acids in one or more domains. In some embodiments, the modifications are selected from modifications in Table 2.

5 [0161] The antibodies or antigen-binding fragment thereof of the present disclosure can comprise a CDR3 region that is a length of at least about 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, or 30 amino acids in length. The antibodies or antigen-binding fragment thereof of the present disclosure can comprise a CDR3 region that is at least about 18 amino acids in length.

10 [0162] As used herein, the term "hinge region" includes the region of a heavy chain molecule that joins the CH1 domain to the CH2 domain. The hinge region can comprise approximately 25 residues and is flexible, thus allowing the two N-terminal antigen binding regions to move independently. Hinge regions can be subdivided into three distinct domains: upper, middle, and lower hinge domains (Roux et al. J. Immunol. 1998 161:4083).

[0163] As used herein, the term "Fv" is the minimum antibody fragment that contains a complete antigen-recognition and -binding site. This fragment consists of a dimer of one heavy- and one light-chain variable region domain in tight, non-covalent association.

15 [0164] From the folding of these two domains emanate six hypervariable loops (three loops each from the H and L chain) that contribute the amino acid residues for antigen binding and confer antigen binding specificity to the antibody. However, even a single variable domain (or half of an Fv comprising only three CDRs specific for an antigen) has the ability to recognize and bind antigen, although at a lower affinity than the entire binding site.

20 [0165] "Heavy chain variable region" or "VH" with regard to an antibody refers to the fragment of the heavy chain that contains three CDRs interposed between flanking stretches known as framework regions, these framework regions are generally more highly conserved than the CDRs and form a scaffold to support the CDRs.

25 [0166] "light chain variable region" or "VL" with regard to an antibody refers to the fragment of the light heavy chain that contains three CDRs interposed between flanking stretches known as framework regions, these framework regions are generally more highly conserved than the CDRs and form a scaffold to support the CDRs.

30 [0167] Six hypervariable loops (three loops each from the Heavy and Light chain) contribute the amino acid residues for antigen-binding and confer antigen-binding specificity to the antibody. However, even a single variable domain (or half of an Fv comprising only three CDRs specific for an antigen) has the ability to recognize and bind antigen, although at a lower affinity than the entire binding site.

[0168] "Framework" or FR residues are those variable domain residues other than the hypervariable region residues.

- [0169] It is understood in the art that an antibody is a glycoprotein having at least two heavy (H) chains and two light (L) chains interconnected by disulfide bonds, or an antigen-binding portion thereof. A heavy chain is comprised of a heavy chain variable region (VH) and a heavy chain constant region (CH1, CH2 and CH3). A light chain is comprised of a light chain variable region (VL) and a light chain constant region (CL). The variable regions of both the heavy and light chains comprise framework regions (FRs or FWRs) and hypervariable regions (HVRs). The HVRs are the amino acid residues of an antibody that are responsible for antigen binding. The hypervariable region generally comprises amino acid residues from a complementarity determining region (CDR), which have the highest sequence variability and/or involved in antigen recognition. With the exception of CDR1 in VH, CDRs generally comprise the amino acid residues that form the hypervariable loops. CDRs also comprise “specificity determining residues,” or “SDRs,” which are residues that contact antigen. SDRs are contained within regions of the CDRs called abbreviated-CDRs, or a-CDRs. Exemplary a-CDRs (a-CDR-L1, a-CDR-L2, a-CDR-L3, a-CDR-H1, a-CDR-H2, and a-CDR-H3) occur at amino acid residues 31-34 of L1, 50-55 of L2, 89-96 of L3, 31-35B of H1, 50-58 of H2, and 95-102 of H3. (See, e.g., Fransson, *Front. Biosci.* 13:1619-1633 (2008))
- [0170] Unless otherwise indicated, HVR residues and other residues in the variable domain (e.g., FR residues) are numbered herein according to Kabat et al., *supra*. A variable region is a domain of an antibody heavy or light chain that is involved in binding the antibody to antigen. (See, e.g., Kindt et al. *Kuby Immunology*, 6th ed., W.H. Freeman and Co., p.91 (2007)). A single VH or VL domain may be sufficient to confer antigen-binding specificity. Furthermore, antibodies that bind a particular antigen may be isolated using a VH or VL domain from an antibody that binds the antigen to screen a library of complementary VL or VH domains, respectively. (See, e.g., Portolano et al., *J. Immunol.* 150:880-887 (1993); Clarkson et al., *Nature* 352:624-628 (1991)). The four FWR regions are typically more conserved while CDR regions (CDR1, CDR2 and CDR3) represent hypervariable regions and are arranged from NH2 terminus to the COOH terminus as follows: FWR1, CDR1, FWR2, CDR2, FWR3, CDR3, and FWR4. The variable regions of the heavy and light chains contain a binding domain that interacts with an antigen while, depending of the isotype, the constant region(s) may mediate the binding of the immunoglobulin to host tissues or factors. An antibody also includes chimeric antibodies, humanized antibodies, and recombinant antibodies, human antibodies generated from a transgenic non-human animal, as well as antibodies selected from libraries using enrichment technologies available to the artisan.
- [0171] The term “antibody heavy chain,” refers to the larger of the two types of polypeptide chains present in antibody molecules in their naturally occurring conformations, and which normally determines the class to which the antibody belongs.

[0172] The term “antibody light chain,” refers to the smaller of the two types of polypeptide chains present in antibody molecules in their naturally occurring conformations. Kappa (“κ”) and lambda (“λ”) light chains refer to the two major antibody light chain isotypes.

- 5 [0173] An antibody or antigen-binding fragment thereof “specifically binds” or “preferentially binds” to a target if it binds with greater affinity and/or avidity than it binds to epitopes on unrelated polypeptides. The specificity of an antibody or antigen-binding fragment or portion thereof can be determined based on affinity and/or avidity. Methods to determine such specific binding are also well known in the art. According to certain embodiments of the present disclosure, the antibodies or antigen-binding fragment thereof can bind to a human cancer antigen but not to a cancer antigen from other species. Alternatively, the antibodies or
- 10 antigen-binding fragment thereof, in certain embodiments, bind to human cancer antigen and to cancer antigen from one or more non-human species. For example, the antibodies or antigen-binding fragment thereof can bind to human cancer antigen and can bind or not bind, as the case may be, to one or more of mouse, rat, guinea pig, hamster, gerbil, pig, cat, dog, rabbit, goat, sheep, cow, horse, camel, cynomolgus, marmoset, rhesus or chimpanzee cancer antigen.
- 15 [0174] The affinity, represented by the equilibrium constant for the dissociation (K_D) of an antigen with an antigen-binding protein, is a measure for the binding strength between an antigenic determinant and an antigen-binding site on the antigen-binding protein: the lesser the value of the K_D , the stronger the binding strength between an antigenic determinant and the antigen-binding molecule. Alternatively, the affinity can also be expressed as the affinity constant (K_A), which is $1/K_D$. As will be clear to the skilled person, affinity
- 20 can be determined in a manner known per se, depending on the specific antigen of interest. Accordingly, an antibody or antigen-binding fragment thereof as defined herein is said to be “specific for” a first target or antigen compared to a second target or antigen when it binds to the first antigen with an affinity (as described above, and suitably expressed, for example as a K_D value) that is at least 50 times, such as at least 100 times, and preferably at least 1000 times, and up to 10,000 times or more better than the affinity with which said
- 25 amino acid sequence or polypeptide binds to another target or polypeptide. Preferably, when an antibody or antigen-binding fragment thereof is “specific for” a target or antigen, compared to another target or antigen, it can bind the target or antigen, but does not bind the other target or antigen. However, as understood by one of ordinary skill in the art, in some embodiments, where a binding site on a target is shared or partially shared by multiple, different ligands, an antibody or antigen-binding fragment thereof can specifically bind to a target,
- 30 such as cancer associated antigen, and have the functional effect of, for example, inhibiting/preventing tumor progression.

[0175] In some embodiments, an antibody provided herein has a dissociation constant (K_D) of about 1 μ M, 100 nM, 10 nM, 5 nM, 2 nM, 1 nM, 0.5 nM, 0.1 nM, 0.05 nM, 0.01 nM, or 0.001 nM or less (*e.g.*, 10^{-8} M or less, *e.g.*, from 10^{-8} M to 10^{-13} M, *e.g.*, from 10^{-9} M to 10^{-13} M). Another aspect of the invention provides for

- an antibody or antigen-binding fragment thereof with an increased affinity for its target, for example, an affinity matured antibody. An affinity matured antibody is an antibody with one or more alterations in one or more hypervariable regions (HVRs), compared to a parent antibody which does not possess such alterations, such alterations resulting in an improvement in the affinity of the antibody for antigen. These antibodies can bind to antigen with a K_D of about 5×10^{-9} M, 2×10^{-9} M, 1×10^{-9} M, 5×10^{-10} M, 2×10^{-10} M, 1×10^{-10} M, 5×10^{-11} M, 1×10^{-11} M, 5×10^{-12} M, 1×10^{-12} M, or less. In some embodiments, the present disclosure provides an antibody or antigen-binding fragment thereof which has an increased affinity of at least 1.5 fold, 2 fold, 2.5 fold, 3 fold, 4 fold, 5 fold, 10 fold, 20 fold or greater as compared to a WT IgA antibody or a WT IgG antibody, containing the heavy chain sequence and light chain sequence, or both. In other embodiments, the antibody or a functional fragment thereof provided herein competes for binding to the same epitope as a corresponding antibody from which the variable domains are derived. In some embodiments, the antibody or antigen-binding fragment thereof that binds to the same epitope, and/or competes for binding to the same epitope as an antibody exhibits effector function activities, such as, for example, Fc-mediated cellular cytotoxicity, including ADCC activity.
- [0176]** K_D can be measured by any suitable assay. For example, K_D can be measured by a radiolabeled antigen-binding assay (RIA) (See, e.g., Chen et al., J. Mol. Biol. 293:865-881 (1999); Presta et al., Cancer Res. 57:4593-4599 (1997)). For example, K_D can be measured using a surface plasmon resonance assay (e.g., using a BIACORE®-2000 or a BIACORE®-3000). For example, K_D can be measured using a competitive ELISA.
- [0177]** Avidity is the measure of the strength of binding between an antigen-binding molecule and the pertinent antigen. Avidity is related to both the affinity between an antigenic determinant and its antigen-binding site on the antigen-binding molecule, and the number of pertinent binding sites present on the antigen-binding molecule. Typically, antigen-binding proteins will bind to their cognate or specific antigen with a dissociation constant (K_D of 10^{-5} to 10^{-12} moles/liter or less, and preferably 10^{-7} to 10^{-12} moles/liter or less and more preferably 10^{-8} to 10^{-12} moles/liter (*i.e.* with an association constant (K_A) of 10^5 to 10^{12} liter/moles or more, and preferably 10^7 to 10^{12} liter/moles or more and more preferably 10^8 to 10^{12} liter/moles). Any K_D value greater than 10^{-4} mol/liter (or any K_A value lower than 10^4 M⁻¹) is generally considered to indicate non-specific binding. The K_D for biological interactions which are considered meaningful (*e.g.*, specific) are typically in the range of 10^{-10} M (0.1 nM) to 10^{-5} M (10000 nM). The stronger an interaction is, the lower is its K_D . Preferably, a binding site on an anti-LAP antibody or antigen-binding fragment thereof described herein will bind with an affinity less than 500 nM, preferably less than 200 nM, more preferably less than 10 nM, such as less than 500 pM. Specific binding of an antigen-binding protein to an antigen or antigenic determinant can be determined in any suitable manner known per se, including, for example, Scatchard analysis and/or competitive binding assays, such as radioimmunoassays (RIA), enzyme

immunoassays (EIA) and sandwich competition assays, and the different variants thereof known per se in the art; as well as other techniques as mentioned herein.

[0178] The term “kon”, as used herein, is intended to refer to the rate constant for association of an antibody or antigen-binding fragment thereof to an antigen.

- 5 **[0179]** The term “Koff”, as used herein, is intended to refer to the rate constant for dissociation of an antibody or antigen-binding fragment thereof from the antibody/antigen complex.

- [0180]** The term “recombinant human antibody”, as used herein, includes all human antibodies that are prepared, expressed, created or isolated by recombinant means, such as (a) antibodies isolated from an animal (e.g., a mouse) that is transgenic or transchromosomal for human immunoglobulin genes or a hybridoma
10 prepared therefrom (described further below), (b) antibodies isolated from a host cell transformed to express the human antibody, e.g., from a transfectoma, (c) antibodies isolated from a recombinant, combinatorial human antibody library, and (d) antibodies prepared, expressed, created or isolated by any other means that involve splicing of human immunoglobulin gene sequences to other DNA sequences. Such recombinant human antibodies have variable regions in which the framework and CDR regions are derived from
15 immunoglobulin sequences, disclosed herein. In certain embodiments, however, such recombinant human antibodies can be subjected to in vitro mutagenesis (or, when an animal transgenic for human Ig sequences is used, in vivo somatic mutagenesis) and thus the amino acid sequences of the VH and VL regions of the recombinant antibodies are sequences that, while derived from and related to human immunoglobulin VH and VL sequences, may not naturally exist within the human antibody germline repertoire in vivo.
- 20 **[0181]** In the context of an antibody or antigen-binding fragment thereof, the term “specificity” or “specific for” refers to the number of different types of antigens or antigenic determinants to which a particular antibody or antigen-binding fragment thereof can bind. The specificity of an antibody or antigen-binding fragment or portion thereof can be determined based on affinity and/or avidity. The affinity, represented by the equilibrium constant for the dissociation (K_D) of an antigen with an antigen-binding protein, is a measure
25 for the binding strength between an antigenic determinant and an antigen-binding site on the antigen-binding protein: the lesser the value of the K_D , the stronger the binding strength between an antigenic determinant and the antigen-binding molecule. Alternatively, the affinity can also be expressed as the affinity constant (K_A), which is $1/K_D$. As will be clear to the skilled person, affinity can be determined in a manner known per se, depending on the specific antigen of interest. Accordingly, an antibody or antigen-binding fragment thereof as
30 defined herein is said to be “specific for” a first target or antigen compared to a second target or antigen when it binds to the first antigen with an affinity (as described above, and suitably expressed, for example as a K_D value) that is at least 50 times, such as at least 100 times, and preferably at least 1000 times, and up to 10,000 times or more better than the affinity with which said amino acid sequence or polypeptide binds to another target or polypeptide. Preferably, when an antibody or antigen-binding fragment thereof is “specific for” a

target or antigen, compared to another target or antigen, it can bind the target or antigen, but does not bind the other target or antigen.

[0182] However, as understood by one of ordinary skill in the art, in some embodiments, where a binding site on a target is shared or partially shared by multiple, different ligands, an antibody or antigen binding fragment thereof can specifically bind to a target antigen, and have the functional effect of, for example, inhibiting/preventing tumor progression.

[0183] Avidity is the measure of the strength of binding between an antigen-binding molecule and the pertinent antigen. Avidity is related to both the affinity between an antigenic determinant and its antigen binding site on the antigen-binding molecule, and the number of pertinent binding sites present on the antigen-binding molecule. Typically, antigen-binding proteins will bind to their cognate or specific antigen with a dissociation constant (K_D of 10^{-5} to 10^{-12} moles/liter or less, and preferably 10^{-7} to 10^{-12} moles/liter or less and more preferably 10^{-8} to 10^{-12} moles/liter (*i.e.* with an association constant (K_A) of 10^5 to 10^{12} liter/moles or more, and preferably 10^7 to 10^{12} liter/moles or more and more preferably 10^8 to 10^{12} liter/moles). Any K_D value greater than 10^{-4} mol/liter (or any K_A value lower than 10^4 M⁻¹) is generally considered to indicate non-specific binding. The K_D for biological interactions which are considered meaningful (*e.g.*, specific) are typically in the range of 10^{-10} M (0.1 nM) to 10^{-5} M (10000 nM). The stronger an interaction is, the lower is its K_D . Preferably, a binding site on an anti-LAP antibody or antigen-binding fragment thereof described herein will bind with an affinity less than 500 nM, preferably less than 200 nM, more preferably less than 10 nM, such as less than 500 pM. Specific binding of an antigen-binding protein to an antigen or antigenic determinant can be determined in any suitable manner known per se, including, for example, Scatchard analysis and/or competitive binding assays, such as radioimmunoassays (RIA), enzyme immunoassays (EIA) and sandwich competition assays, and the different variants thereof known per se in the art; as well as other techniques as mentioned herein.

[0184] The term "fusion protein" as used herein refers to a polypeptide that comprises an amino acid sequence of an antibody or fragment thereof and an amino acid sequence of a heterologous polypeptide (*i.e.*, an unrelated polypeptide).

[0185] In some embodiments, the antibody or antigen binding fragment thereof of the present disclosure is a single domain antibody. The expression "single domain antibody" (sdAbs) or "single variable domain (SVD) antibody" generally refers to a single variable region (VH or V) wherein the antibody-antigen binding can be imparted. In other words, single variable domain does not need to recognize the target antigen by interacting with another variable region. A single domain antibody monomers single arm antigen binding by each antibody variable region (VH * VJ composition. Examples of single domain antibodies include those derived from camelids (camels and llamas) and cartilaginous fish (*e.g.* nurse sharks) antibodies and those antibodies (Ward et al from human and mouse antibodies by

recombinant methods, Nature (1989) 341: 544-546; Dooley and Flajnik, Dev Comp Immunol (2006) 30: 43-56; Muyldermans et, TrendBiochem Sci (2001) 26: 230-235; Holt et, Trends Biotechnol (2003): 21: 484-490; W0 2005/035572; TO 03/035694; Davies and Riechmann, Febs Lett (1994) 339: 285-290; W000 / 29 004; W0 02/051870) and a single variable region of an antibody can be other than a single domain antibody
 5 variable regions or variable domains are present in an antigen binding arm (e.g., homo- or hetero-multimer together).

[0186] As used herein the term ‘modification’ refers to an amino acid substitution or an amino acid deletion of one or more amino acid residues in a heavy chain constant region of an antibody, compared to a WT heavy chain constant region of a WT antibody. In some embodiments, the modification is in an amino acid residue
 10 in the IgA CH1 region of the heavy chain constant region. In some embodiments, the modification is in an amino acid residue in the IgA CH2 region of the heavy chain constant region. In some embodiments, the modification is in an amino acid residue in the IgA CH3 region of the heavy chain constant region. In some embodiments, the modification is selected from Table 2. In some embodiments, the one or more modifications disclosed herein result in an improved property of the antibody comprising the one or more
 15 modifications compared to a corresponding WT antibody.

[0187] The term "improved property" means a characteristic associated with an antibody comprising the one or more modifications disclosed herein, that is improved compared to the parent WT antibody not comprising the one or more modifications. Such an improved property includes, but is not limited to, increased thermostability, increased circulating half-life, increased ADCC, decreased aggregation, decreased
 20 aggregation with serum proteins, increased tumor targeting, increased stability, decreased glycosylation, increased binding a Fc α R expressed on an immune cell. In some embodiments, the improved property is an increase in one or more effector functions of the antibody or the functional fragment thereof, as compared to a corresponding WT IgA. Effector functions are biological activities attributable to the Fc region of an antibody, which vary with the antibody isotype. Examples of antibody effector functions include: C1q binding
 25 and complement dependent cytotoxicity (CDC); Fc receptor binding; antibody-dependent cell-mediated cytotoxicity (ADCC); phagocytosis.

[0188] In some embodiments, the antibodies or a functional fragment thereof disclosed herein (e.g., comprising the one or modifications disclosed herein in the IgA heavy chain constant region), have at least 2%, 3%, 4%, 5%, 7%, 8%, 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, or at least 100% of one or
 30 more improved property, compared to a corresponding WT IgA antibody or a corresponding WT IgG antibody.

[0189] As used herein the term “corresponding” unmodified antibody means a wild type antibody of the same sequence as the antibody comprising one or more select modifications disclosed herein, but without the one or more selected modifications described herein, in particular in the heavy chain constant region. In some

embodiments, the corresponding antibody can be a WT IgA antibody comprising a WT IgA heavy chain constant region. In some embodiments, the corresponding antibody can be a WT IgG antibody comprising a WT IgG heavy chain constant region. In some embodiments, a corresponding WT IgA antibody is a WT IgA1 antibody. In some embodiments, the corresponding WT IgA antibody is WT IgA2 antibody. In some
5 embodiments, a corresponding WT IgA antibody comprises a wild type IgA heavy chain constant region comprising an amino acid sequence set forth in SEQ ID NO: 1. In some embodiments, a corresponding WT IgA2 antibody comprises a wild type IgA2 heavy chain constant region comprising an amino acid sequence set forth in SEQ ID NO: 1. In some embodiments, a corresponding WT IgA antibody comprises a wild type
10 IgA heavy chain constant region comprising an amino acid sequence set forth in SEQ ID NO: 2. In some embodiments, a corresponding WT IgA antibody comprises a wild type IgA heavy chain constant region comprising an amino acid sequence set forth in SEQ ID NO: 2. In some embodiments, the WT IgA heavy chain constant region comprises an amino acid sequence set forth in SEQ ID NO: 1. In some embodiments, the WT IgA2 heavy chain constant region comprises an amino acid sequence set forth in SEQ ID NO: 1. In some embodiments, the amino acid modifications disclosed herein are relative to amino acid residues at select
15 positions in the WT IgA heavy chain constant region (e.g., WT IgA2 heavy chain constant region) comprising an amino acid sequence set forth in SEQ ID NO: 1, numbering according to IMGT scheme.

[0190] In some embodiments, the WT IgA heavy chain constant region comprises an amino acid sequence set forth in SEQ ID NO: 2. In some embodiments, the WT IgA2 heavy chain constant region comprises an amino acid sequence set forth in SEQ ID NO: 2. In some embodiments, the amino acid modifications
20 disclosed herein are relative to amino acid residues at select positions in the WT IgA heavy chain constant region (e.g., WT IgA2 heavy chain constant region) comprising an amino acid sequence set forth in SEQ ID NO: 2, numbering according to IMGT scheme.

[0191] In some embodiments, the WT IgA heavy chain constant region comprises an amino acid sequence set forth in SEQ ID NO: 3. In some embodiments, the WT IgA2 heavy chain constant region comprises an amino acid sequence set forth in SEQ ID NO: 3. In some embodiments, the amino acid modifications
25 disclosed herein are relative to amino acid residues at select positions in the WT IgA heavy chain constant region (e.g., WT IgA2 heavy chain constant region) comprising an amino acid sequence set forth in SEQ ID NO: 3, numbering according to IMGT scheme.

[0192] The term "Fab" as used herein is intended to refer to a region of an antibody composed of one
30 constant and one variable domain of each of the heavy and the light chains (monovalent antigen-binding fragment), but wherein the heavy chain is truncated such that it lacks the CH2 and CH3 domain (ie VH, CH1, VL, and CL), and may also lack some or all of the hinge region. It can be produced by digestion of a whole antibody with the enzyme papain. Fab may refer to this region in isolation, or this region in the context of a full length antibody, immunoglobulin construct or Fab fusion protein.

[0193] The term Fab' as used herein can be obtained by treating a whole antibody with pepsin, followed by reduction, to yield a molecule consisting of an intact light chain and a portion of a heavy chain comprising a VH and a single constant domain. Two Fab' fragments are obtained per antibody treated in this manner.

[0194] By "scFv" it is meant an antibody fragment comprising the VFI and VL domains of an antibody, wherein these domains are present in a single polypeptide chain. See, for example, U.S. Pat. Nos. 4,946,778, 5,260,203, 5,455,030, and 5,856,456. Generally, the Fv polypeptide further comprises a polypeptide linker between the VH and VL domains that enables the scFv to form the desired structure for antigen-binding. For a review of scFv see Pluckthun (1994) The Pharmacology of Monoclonal Antibodies vol 113 ed. Rosenberg and Moore (Springer-Verlag, New York) pp 269-315. The VFI and VL domain complex of Fv fragments may also be stabilized by a disulfide bond (US Pat. No. 5,747,654).

[0195] The term "in vivo half-life" or "circulating half-life" as used herein refers to the circulation of an antibody or a functional fragment thereof in a given animal and is represented by a time required for half the quantity administered in the animal to be cleared from the circulation.

[0196] The term "increased circulating half life" as used herein means that the antibody comprising one or more modifications relative to a WT IgA antibody as provided according to the invention has a greater persistence in the serum or plasma and/or takes a greater period of time to reduce to half the maximal measured serum or plasma concentration relative to the same antibody, that does not contain the same modifications, i.e., a WT antibody e.g., WT IgA antibody.

[0197] The term "increased thermostability" means a higher retention of a biological activity e.g., ADCC, binding to an antigen, binding to a Fc α R, of an antibody or a functional fragment disclosed herein after a period of incubation at a temperature relative to a corresponding WT IgA antibody. The increased thermostability of the can be assessed, for example, under conditions of one or more (e.g., several) temperatures. For example, the one or more (e.g., several) temperatures can be any temperature or temperatures in the range of 45°C to 95°C, e.g., 45, 50, 55, 60, 65, 70, 75, 80, 85, or 95°C (or in between, e.g., 62°C, 68°C, 72°C, etc.) at one or more (e.g., several) pHs in the range of 3 to 9, e.g., 3.0, 3.5, 4.0, 4.5, 5.0, 5.5, 6.0, 6.5, 7.0, 7.5, 8.0, 8.5, or 9.0 (or in between) for a suitable period (time) of incubation, e.g., 1 minute, 5 minutes, 10 minutes, 15 minutes, 20 minutes, 25 minutes, 30 minutes, 45 minutes, or 60 minutes (or in between, e.g., 23 minutes, 37 minutes, etc.), such that the variant retains residual activity. However, longer periods of incubation can also be used. The term "increased thermostability" can be used interchangeably with "improved thermostability".

[0198] As used herein, the term "naturally occurring " as it relates to a glycosylation site or an amino acid residue in a IgA heavy chain constant region, refers to the fact that the glycosylation site or the amino acid residue can be found in nature in the IgA heavy chain constant region, for example, can be isolated from a

source in nature and has not been intentionally modified in the laboratory by a human (including a virus). A polypeptide or polynucleotide sequence present in an organism is a sequence naturally occurring.

[0199] The terms “disease”, “disorder”, or “condition” are used interchangeably herein, refer to any alternation in state of the body or of some of the organs, interrupting or disturbing the performance of the functions and/or causing symptoms such as discomfort, dysfunction, distress, or even death to the person afflicted or those in contact with a person. A disease or disorder can also be related to a distemper, ailing, ailment, malady, disorder, sickness, illness, complaint, or affectation.

[0200] The term “in need thereof” when used in the context of a therapeutic or prophylactic treatment, means having a disease, being diagnosed with a disease, or being in need of preventing a disease, e.g., for one at risk of developing the disease. Thus, a subject in need thereof can be a subject in need of treating or preventing a disease.

[0201] As used herein, the term “administering,” refers to the placement of a compound (e.g., an antibody or antigen binding fragment thereof as disclosed herein) into a subject by a method or route that results in at least partial delivery of the agent at a desired site. Pharmaceutical compositions comprising an antibody or antigen binding fragment thereof, disclosed herein can be administered by any appropriate route which results in an effective treatment in the subject, including but not limited to intravenous, intraarterial, injection or infusion directly into a tissue parenchyma, etc. Where necessary or desired, administration can include, for example, intracerebroventricular (“icv”) administration, intranasal administration, intracranial administration, intracelial administration, intracerebellar administration, or intrathecal administration.

[0202] The term “cancer” refers to a disease characterized by the rapid and uncontrolled growth of aberrant cells. Cancer cells can spread locally or through the bloodstream and lymphatic system to other parts of the body.

[0203] The term “anti-tumor effect” refers to a biological effect which can be manifested by various means, including but not limited to, e.g., a decrease in tumor volume, a decrease in the number of tumor cells, a decrease in the number of metastases, an increase in life expectancy, decrease in tumor cell proliferation, decrease in tumor cell survival, or amelioration of various physiological symptoms associated with the cancerous condition. An “anti-tumor effect” can also be manifested by the ability of the peptides, polynucleotides, cells and antibodies of the invention in prevention of the occurrence of tumor in the first place.

[0204] As used herein, a “subject”, “patient”, “individual” and like terms are used interchangeably and refers to a vertebrate, a mammal, a primate, or a human. Mammals include, without limitation, humans, primates, rodents, wild or domesticated animals, including feral animals, farm animals, sport animals, and pets. Primates include, for example, chimpanzees, cynomolgus monkeys, spider monkeys, and macaques, e.g., Rhesus. Rodents include, for example, mice, rats, woodchucks, ferrets, rabbits and hamsters. Domestic and

game animals include, for example, cows, horses, pigs, deer, bison, buffalo, feline species, e.g., domestic cat, and canine species, e.g., dog, fox, wolf, avian species, e.g., chicken, emu, ostrich, and fish, e.g., trout, catfish and salmon. The terms, "individual," "patient" and "subject" are used interchangeably herein. A subject can be male or female.

5 [0205] In some embodiments, the subject is a mammal. The mammal can be a human, non-human primate, mouse, rat, dog, cat, horse, or cow, but is not limited to these examples. Mammals other than humans can be advantageously used as subjects that represent animal models of conditions or disorders associated with uncontrolled cell growth (e.g., a cancer). Non-limiting examples include murine tumor models. In addition, the compositions and methods described herein can be used to treat domesticated animals and/or pets. A
10 subject can be one who has been previously diagnosed with or identified as suffering from a cancer. A subject can be one who is diagnosed and currently being treated for, or seeking treatment, monitoring, adjustment or modification of an existing therapeutic treatment, or is at a risk of developing a given disorder (e.g., cancer).

[0206] A "cytotoxic agent" refers to an agent that has a cytotoxic and/or cytostatic effect on a cell. A "cytotoxic effect" refers to the depletion, elimination and/or the killing of a target cell(s). A "cytostatic effect"
15 refers to the inhibition of cell proliferation.

[0207] As used herein, the terms "protein", "peptide" and "polypeptide" are used interchangeably to designate a series of amino acid residues connected to each other by peptide bonds between the alpha-amino and carboxy groups of adjacent residues. The terms "protein", "peptide" and "polypeptide" refer to a polymer of amino acids, including modified amino acids (e.g., phosphorylated, glycosylated, etc.) and amino
20 acid analogs, regardless of its size or function. "Protein" and "polypeptide" are often used in reference to relatively large polypeptides, whereas the term "peptide" is often used in reference to small polypeptides, but usage of these terms in the art overlaps. The terms "protein", "peptide" and "polypeptide" are used interchangeably herein when referring to a gene product and fragments thereof. These terms encompass, e.g., native and artificial proteins, protein fragments and polypeptide analogs (such as muteins, variants, and fusion
25 proteins) of a protein sequence as well as post-translationally, or otherwise covalently or non-covalently, modified proteins. A peptide, polypeptide, or protein may be monomeric or polymeric. A polypeptide can have the amino acid sequence of naturally occurring polypeptide from any mammal. Such native sequence polypeptide can be isolated from nature or can be produced by recombinant or synthetic means. In some embodiments, the polypeptide is a "variant". "Variant" means a biologically active polypeptide having at least
30 about 80% amino acid sequence identity with the native sequence polypeptide after aligning the sequences and introducing gaps, if necessary, to achieve the maximum percent sequence identity, and not considering any conservative substitutions as part of the sequence identity. Such variants include, for instance, polypeptides wherein one or more amino acid residues are added, or deleted, at the N- or C-terminus of the polypeptide. In some embodiments, a variant will have at least about 80% amino acid sequence identity. In

some embodiments, a variant will have at least about 90% amino acid sequence identity. In some embodiments, a variant will have at least about 95% amino acid sequence identity with the native sequence polypeptide. A “derivative” of a polypeptide is a polypeptide (e.g., an antibody) that has been chemically modified, e.g., via conjugation to another chemical moiety (such as, for example, polyethylene glycol or albumin, e.g., human serum albumin), phosphorylation, and glycosylation

[0208] The terms “increased”, “increase”, or “enhance” are all used herein to generally mean an increase by a statically significant amount; for the avoidance of doubt, the terms “increased”, “increase”, or “enhance”, mean an increase of at least 10% as compared to a reference level, for example an increase of at least about 10%, at least about 20%, or at least about 30%, or at least about 40%, or at least about 50%, or at least about 60%, or at least about 70%, or at least about 80%, or at least about 90% or up to and including a 100% increase or any increase between 10-100% as compared to a reference level, or at least about a 2-fold, or at least about a 3-fold, or at least about a 4-fold, or at least about a 5-fold or at least about a 10-fold increase, or any increase between 2-fold and 10-fold or greater as compared to a reference level.

[0209] The term “fusion protein” as used herein refers to a polypeptide that comprises an amino acid sequence of an antibody or fragment thereof and an amino acid sequence of a heterologous polypeptide (i.e., an unrelated polypeptide).

[0210] The terms, “decrease”, “reduce”, “reduction”, “lower” or “lowering,” or “inhibit” are all used herein generally to mean a decrease by a statistically significant amount. For example, “decrease”, “reduce”, “reduction”, or “inhibit” means a decrease by at least 10% as compared to a reference level, for example a decrease by at least about 20%, or at least about 30%, or at least about 40%, or at least about 50%, or at least about 60%, or at least about 70%, or at least about 80%, or at least about 90% or up to and including a 100% decrease (e.g., tumor size after treatment as compared to a reference level prior to the treatment), or any decrease between 10-100% as compared to a reference level. In the context of a marker or symptom, by these terms is meant a statistically significant decrease in such level. The decrease can be, for example, at least 10%, at least 20%, at least 30%, at least 40% or more, and is preferably down to a level accepted as within the range of normal for an individual without a given disease. Reduce or inhibit can refer to, for example, the symptoms of the disorder being treated, the presence or size of metastases or micrometastases, the size of the primary tumor, the presence or the size of the dormant tumor.

[0211] The terms “synthetic polynucleotide,” “synthetic gene” or “synthetic polypeptide,” as used herein, mean that the corresponding polynucleotide sequence or portion thereof, or amino acid sequence or portion thereof, is derived, from a sequence that has been designed, or synthesized de novo, or modified, compared to an equivalent naturally-occurring sequence. Synthetic polynucleotides (antibodies or antigen-binding fragments) or synthetic genes can be prepared by methods known in the art, including but not limited to, the chemical synthesis of nucleic acid or amino acid sequences. Synthetic genes are typically different from

naturally-occurring genes, either at the amino acid, or polynucleotide level, (or both) and are typically located within the context of synthetic expression control sequences. Synthetic gene polynucleotide sequences, may not necessarily encode proteins with different amino acids, compared to the natural gene; for example, they can also encompass synthetic polynucleotide sequences that incorporate different codons but which encode the same amino acid (i.e., the nucleotide changes represent silent mutations at the amino acid level).

IgA Antibodies

[0212] IgA has two subclasses (IgA1 and IgA2) and can be produced as a monomeric as well as a dimeric form and secretory form. In some embodiments, the IgA antibody can be monomeric. In some embodiments, the IgA antibody can comprise one or more IgA1 amino acid sequences. In some embodiments, the IgA antibody can comprise one or more IgA2 amino acid sequences. In some embodiments, the IgA antibody can comprise one or more IgA1 amino acid sequences and one or more IgA2 amino acid sequences.

[0213] In some embodiments, the IgA antibody is an IgA2 antibody of allotype: IgA2m(1), IgA2(m)2, or IgA2n. In some embodiments, the IgA2m(1) antibody is a Caucasian IgA2m(1) antibody. In some embodiments, the IgA2m(2) antibody is an African IgA2m(2) antibody or an Asian IgA2m(2) antibody.

[0214] In some embodiments, the IgA antibody comprises a heavy chain constant region comprising at least 60%, 70%, 75%, 80%, 85%, 90%, 95%, or 99% IgA amino acids. In some embodiments, the IgA antibody comprises a light chain constant region comprising at least 50, 60%, 70%, 75%, 80%, 85%, 90%, 95%, or 99% IgA amino acids.

[0215] In some embodiments, the IgA antibody comprises a heavy chain constant region comprising one or more of an IgA CH3 region, an IgA CH2, or an IgA CH1 region, or any combination thereof. In some embodiments, the IgA antibody comprises a light chain region comprising an IgA CH1 region. In some embodiments, the IgA antibody comprises a light chain region comprising a kappa light chain constant region. In some embodiments, the IgA antibody comprises a heavy chain constant region comprising one or more amino acid of an IgG CH3 region, an IgG CH2, or an IgG CH1 region, or any combination thereof. In some embodiments, the IgA antibody comprises a heavy chain constant region comprising an IgA CH3 region, an IgA CH2, and an IgA CH1 region, or any combination thereof. In some embodiments, the IgA antibody comprises a heavy chain constant region comprising an IgA CH3 region, an IgA CH2, and an IgG CH1 region, or any combination thereof. In some embodiments, the IgA2 antibody comprises a heavy chain constant region comprising one or more of an IgA2 CH3 region, an IgA2 CH2, or an IgA2 CH1 region, or any combination thereof.

[0216] In some embodiments, the IgA antibody comprises an IgG light chain variable region. In some embodiments, the IgA antibody comprises an IgG heavy chain variable region. In some embodiments, the IgA antibody comprises an IgG light chain variable region and an IgG heavy chain variable region.

[0217] In some embodiments, the IgA antibody can be a humanized antibody. In some embodiments, the IgA antibody can be a chimeric antibody. In some embodiments, the IgA antibody can be a human antibody.

[0218] In some embodiments, the IgA antibody can be a monospecific antibody. In some embodiments, the IgA antibody can be a bi-specific antibody. In some embodiments, the IgA antibody can be a tri-specific antibody. In some embodiments, the IgA antibody can be a multi-specific antibody.

[0219] In some embodiments, the IgA antibody can be a bispecific antibody. In some embodiments, the IgA antibody co-engages two antigens at the cell surface. In some examples, the binding of the IgA antibody to two different antigens is sequential. For example, the binding of the IgA antibody to the first antigen occurs first and thereby restricts the space explored by the second antibody arm. Consequentially, there can be a significant increase in local concentration of the second antigen, which can facilitate the binding of the second antibody arm.

[0220] In some embodiments, the IgA antibody can comprise at least a portion of the Fc domain. In some embodiments, the IgA antibody comprises a heavy chain constant region comprising a CH3, CH2, and CH1 domain. In some embodiments, the IgA antibody comprises a light chain constant region comprising a CH1.

[0221] In some embodiments, the IgA antibody induces complement-dependent cytotoxicity (CDC). In some embodiments, the IgA antibody induces polymorphonuclear neutrophil (PMN)-mediated tumor cell lysis. In some embodiments, the IgA antibody induces programmed cell death (PCD) via a caspase-independent pathway. In some embodiments, the IgA antibody induces antibody-dependent cell-mediated cytotoxicity (ADCC). In some embodiments, the IgA antibody induces antibody-dependent cell-mediated cytotoxicity (ADCC) mediated by neutrophils.

[0222] In some embodiments, the IgA antibody can have a superior ability to recruit neutrophils for antibody-dependent cell-mediated cytotoxicity (ADCC) compared to a corresponding IgG antibody. In some embodiments, the IgA antibody can require lower effector:target (E:T) ratios. In some embodiments, the IgA antibody can require lower tumor-opsonizing antibody concentrations compared to other types of antibodies (e.g., IgG). In some embodiments, the IgA antibody can trigger neutrophil-mediated phagocytosis or trogocytosis of tumor cells following IgA antibody-neutrophil engagement. In some embodiments, the IgA antibody can trigger trogoptosis of tumor cells following IgA antibody-neutrophil engagement. This mechanism of killing tumor cells is mediated mainly by interacting with the Fc receptor for IgA (FcαRI; CD89), which is the best characterized IgA receptor. FcαRI is expressed on monocytes, macrophages, granulocytes, subsets of dendritic cells, and Kupffer cells and binds both monomeric and dimeric IgA isoforms with median affinity. Binding of IgA to FcαRI mediates effector functions such as phagocytosis, oxidative burst, cytokine release, antigen presentation, and ADCC. In humans, two IgA isotypes, IgA1 and IgA2, and three allotypes, IgA2m(1), IgA2m(2) and IgA2n, have been distinguished. In some embodiments, IgA antibodies trigger polymorphonuclear cell (PMN) mediated ADCC more efficiently than IgG antibodies.

[0223] In some embodiments, the IgA does not bind a B cell, a T cell, a platelet, and/or an erythrocyte. For example, in some embodiments, the IgA antibody can have low immunogenicity.

[0224] In some embodiments, the IgA antibody is a therapeutic antibody. In some embodiments, the IgA antibody can be a recombinant antibody. In some embodiments, the IgA antibody is made in a cell line. In some embodiments, the cell line is CHO. In some embodiments, the cell line is SP20. In some embodiments, the cell line is a HEK 239 cell line. In some embodiments, the HEK 293 cell line is HEK 293 F.

[0225] In some embodiments, the antibody or a functional fragment thereof provided herein comprises an IgA heavy chain constant region comprising having at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% sequence identity to the amino acid sequence of any one of SEQ ID NO:16-21. In some embodiments, the antibody or a functional fragment thereof provided herein comprises a variable heavy chain domain from an IgG antibody. In some embodiments, the antibody or a functional fragment thereof provided herein comprises a variable light chain domain from an IgG antibody. The variable domains can be derived from, for example, Dinutuximab, Obinutuzumab, Unituxin, TA99, 2.3D11, C47A8-CQ, UMAB10, and Trastuzumab and Rituxan (Rituximab), other antibodies having therapeutic activity encompassed within the scope of the invention include, but are not limited to, Avastin, Herceptin, 3F8, 8H9, Abagovomab, Abciximab, Actoxumab, Adalimumab, Adecatumumab, Aducanumab, Afelimomab, Afutuzumab, Alacizumab pegol, ALD518, Alemtuzumab, Alirocumab, Altumomab pentetate, Amatuximab, Anatumomab mafenatox, Anifrolumab, Anrukinzumab, Apolizumab, Arcitumomab, Aselizumab, Atinumab, Atlizumab, Atorolimumab, Bapineuzumab, Basiliximab, Bavixumab, Bectumomab, Belimumab, Benralizumab, Bertilimumab, Besilesomab, Bevacizumab, Bezlotoxumab, Biciromab, Bimagrumab, Bivatuzumab mertansine, Blinatumomab, Blosozumab, Brentuximab vedotin, Briakinumab, Brodalumab, Canakinumab, Cantuzumab mertansine, Cantuzumab ravtansine, Caplacizumab, Capromab pendetide, Carlumab, Catumaxomab, cBR96-doxorubicin immunoconjugate, CC49, Cedelizumab, Certolizumab pegol, Cetuximab, Ch.14.18, Citatuzumab bogatox, Cixutumumab, Clazakizumab, Clenoliximab, Clivatuzumab tetraxetan, Conatumumab, Concizumab, CR6261, Crenezumab, Dacetuzumab, Daclizumab, Dalotuzumab, Daratumumab, Demcizumab, Denosumab, Detumomab, Dorlimomab aritox, Drozitumab, Duligotumab, Dupilumab, Dusigitumab, Ecomeximab, Eculizumab, Edobacomab, Edrecolomab, Efalizumab, Efungumab, Eldelumab, Elotuzumab, Elsilimomab, Enavatuzumab, Enlimomab pegol, Enokizumab, Enoticumab, Ensituximab, Eptumomab cituxetan, Epratuzumab, Erlizumab, Ertumaxomab, Etaracizumab, Etrolizumab, Evolocumab, Exbivirumab, Fanolesomab, Faralimumab, Farletuzumab, Fasinumab, FBT A05, Felvizumab, Fezakinumab, Ficlaturumab, Figitumumab, Flanvotumab, Fontolizumab, Foralumab, Foravirumab, Fresolimumab, Fulranumab, Futuximab, Galiximab, Ganitumab, Gantenerumab, Gavilimumab, Gemtuzumab ozogamicin, Gevokizumab, Girentuximab, Glembatumumab vedotin, Golimumab, Gomiliximab, Guselkumab, Ibalizumab, Ibritumomab tiuxetan, Icrucumab, Igovomab, IMAB362, Imciromab,

- Imgatuzumab, Inclacumab, Indatuximab ravtansine, Infliximab, Inolimomab, Inotuzumab ozogamicin, Intetumumab, Ipilimumab, Iratumumab, Itolizumab, Ixekizumab, Keliximab, Labetuzumab, Lambrolizumab, Lampalizumab, Lebrikizumab, Lemalesomab, Lerdelimumab, Lexatumumab, Libivirumab, Ligelizumab, Lintuzumab, Lirilumab, Lodelcizumab, Lorvotuzumab mertansine, Lucatumumab, Lumiliximab,
- 5 Mapatumumab, Margetuximab, Maslimomab, Matuzumab, Mavrilimumab, Mepolizumab, Metelimumab, Milatuzumab, Minretumomab, Mitumomab, Mogamulizumab, Morolimomab, Motavizumab, Moxetumomab pasudotox, Muromonab-CD3, Nacolomab tafenatox, Namilumab, Naptumomab estafenatox, Narnatumab, Natalizumab, Nebacumab, Necitumumab, Nerelimomab, Nesvacumab, Nimotuzumab, Nivolumab, Nofetumomab merpentin, Ocaratuzumab, Ocrelizumab, Odulimumab, Ofatumumab, Olaratumab,
- 10 Olokizumab, Omalizumab, Onartuzumab, Ontuxizumab, Oportuzumab monatox, Oregovomab, Orticumab, Otelixizumab, Otlertuzumab, Oxelumab, Ozanezumab, Ozoralizumab, Pagibaximab, Palivizumab, Panitumumab, Pankomab, Panobacumab, Parsatuzumab, Pascolizumab, Pateclizumab, Patritumab, Pentumomab, Perakizumab, Pertuzumab, Pexelizumab, Pidilizumab, Pinatuzumab vedotin, Pintumomab, Placulumab, Polatuzumab vedotin, Ponezumab, Priliximab, Pritoxaximab, Pritumumab, PRO 140,
- 15 Quilizumab, Racotumomab, Radretumab, Rafivirumab, Ramucirumab, Ranibizumab, Raxibacumab, Regavirumab, Reslizumab, Rilotumumab, Rituximab, Robatumumab, Roledumab, Romosozumab, Rontalizumab, Rovelizumab, Ruplizumab, Samalizumab, Sarilumab, Satumomab pendetide, Secukinumab, Seribantumab, Setoxaximab, Sevirumab, SGN-CD19A, SGN-CD33A, Sibrotuzumab, Sifalimumab, Siltuximab, Simtuzumab, Siplizumab, Sirukumab, Solanezumab, Solitomab, Sonpecizumab, Sontuzumab,
- 20 Stamulumab, Sulesomab, Suvizumab, Tabalumab, Tacatuzumab tetraxetan, Tadocizumab, Talizumab, Tanezumab, Taplitumomab paptox, Tefibazumab, Telimomab aritox, Tenatumomab, Teneliximab, Teplizumab, Teprotumumab, TGN1412, Ticilimumab, Tigatuzumab, Tildrakizumab, TNX-650, Tocilizumab, Toralizumab, Tositumomab, Tovetumab, Tralokinumab, Trastuzumab, TRBS07, Tregalizumab, Tremelimumab, Tucotuzumab celmoleukin, Tuvirumab, Ublituximab, Urelumab, Urtoxazumab,
- 25 Ustekinumab, Vantictumab, Vapaliximab, Vatelizumab, Vedolizumab, Veltuzumab, Vepalimumab, Vesencumab, Visilizumab, Volociximab, Vorsetuzumab mafodotin, Votumumab, Zalutumumab, Zanolimumab, Zatuximab, Ziralimumab, hu14.18K322A, Zolimomab aritox, rituximab (Rituxan®, CD20), trastuzumab (Herceptin®), alemtuzumab (Campath®, CD52), ibritumomab tiuxetan (Zevalin®, CD20) tositumomab-I-131 (Bexxar®: CD20), cetuximab (Erbix®), bevacizumab (VEGF), panitumumab
- 30 (Vectibix®, EGFR), ofatumumab (Arzerra®, CD20), ipilimumab (Ypervoy®, CTLA-4), brentuximab vedotin (Alectris®, CD30), pertuzumab (Perjeta®, HER2), adotrastuzumab, ematansine (Kadcyla®, HER2), obinutuzumab (Gazyva®, CD20), nivolumab and pembrolizumab (anti-PD-1s).

[0226] In one aspect, an antibody or antigen-binding fragment thereof comprises a heavy chain variable region (VH) sequence having at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100%

sequence identity to the amino acid sequence of any one of SEQ ID NOS: 4, 7, 81-86. In some embodiments, a VH sequence having at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% identity contains substitutions (e.g., conservative substitutions), insertions, or deletions relative to the reference sequence, but an antibody comprising that sequence retains the ability to bind to same antigen as of the WT antibody. In some embodiments, a total of 1 to 10 amino acids have been substituted, inserted and/or deleted in the amino acid sequence of any one of SEQ ID NOS: 4, 7, 81-86. In some embodiments, substitutions, insertions, or deletions occur in regions outside the CDRs (e.g., in the FRs). Optionally, the antibody comprises the VH sequence of the amino acid sequence of SEQ ID NOS: 4, 7, 81-86, including one or more post-translational modifications of that sequence. In a particular embodiment, the VH comprises one, two or three CDRs selected from: (a) HC-CDR1 comprising the amino acid sequence of any one of SEQ ID NOS: 33-40, (b) HC-CDR2 comprising the amino acid sequence of any one of SEQ ID NO: 41-48, and (c) HC-CDR3 comprising the amino acid sequence of any one of SEQ ID NO: 49-56.

[0227] In one aspect, an antibody or antigen-binding fragment thereof, is provided, wherein the antibody or antigen-binding fragment thereof comprises a light chain variable region (VL) having at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% sequence identity to the amino acid sequence of any one of SEQ ID NOS: 5, 8, 95-100. In some embodiments, a VL sequence having at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% identity contains substitutions (e.g., conservative substitutions), insertions, or deletions relative to the reference sequence, but an antibody or antigen-binding fragment thereof comprising that sequence retains the ability to bind to same antigen as the WT antibody. In some embodiments, a total of 1 to 10 amino acids have been substituted, inserted and/or deleted in any one of the amino acid sequences of any one of SEQ ID NOS: 5, 8, 95-100. In some embodiments, the substitutions, insertions, or deletions occur in regions outside the CDRs (e.g., in the FRs). Optionally, the antibody or antigen-binding fragment thereof comprises the VL sequence of any one of SEQ ID NO: 5, 8, 95-100., including post-translational modifications of that sequence. In a particular embodiment, the VL comprises one, two or three CDRs selected from (a) LC-CDR1 comprising the amino acid sequence of any one of SEQ ID NOS: 57-64; (b) LC-CDR2 comprising the amino acid sequence of any one of SEQ ID NOS: 65-72; and (c) LC-CDR3 comprising the amino acid sequence of any one of SEQ ID NOS: 73-80.

[0228] In one aspect, an antibody or antigen-binding fragment thereof is provided, wherein the antibody or antigen-binding fragment thereof comprises a VH as in any of the embodiments provided above, and a VL as in any of the embodiments provided above. In some embodiments, the antibody or antigen-binding fragment thereof comprises a VH and a VL, wherein the VH comprises the amino acid sequence of any one of SEQ ID NOS: 4, 7, 81-86, and wherein the VL comprises the amino acid sequence in any one of SEQ ID NOS: 5, 8, 95-100, and optionally including post-translational modifications of those sequences.

[0229] In one aspect, an antibody or antigen-binding fragment thereof is provided, wherein the antibody or antigen-binding fragment thereof comprises a VH selected from any VH in Table 9. In one aspect, an antibody or antigen-binding fragment thereof is provided, wherein the antibody or antigen-binding fragment thereof comprises a VL selected from any VL in Table 9. In one aspect, an antibody or antigen-binding fragment thereof is provided, wherein the antibody or antigen-binding fragment thereof comprises a VH selected from any VH in Table 9 and a VL selected from any VL in Table 9. In one aspect, an antibody or antigen-binding fragment thereof is provided, wherein the antibody or antigen-binding fragment thereof comprises a VH selected from any VH in Table 9 and a VL selected from any VL in Table 9, wherein the selected VH and VL are paired according to Table 9. In one aspect, an antibody or antigen-binding fragment thereof is provided, wherein the antibody or antigen-binding fragment thereof comprises a HC-CDR3 selected from any HC-CDR3 in Table 5 and a LC-CDR3 selected from any LC-CDR3 in Table 6, wherein the selected HC-CDR3 and LC-CDR3 are paired according to Table 9. In one aspect, an antibody or antigen-binding fragment thereof is provided, wherein the antibody or antigen-binding fragment thereof comprises a HC-CDR2 selected from any CDR-H2 in Table 5 and a LC-CDR2 selected from any CDR-L2 in Table 6, wherein the selected HC-CDR2 and LC-CDR2 are paired according to Table 9. In one aspect, an antibody or antigen-binding fragment thereof is provided, wherein the antibody or antigen-binding fragment thereof comprises a HC-CDR1 selected from any HC-CDR1 in Table 5 and a LC-CDR1 selected from any LC-CDR1 in Table 6, wherein the selected HC-CDR1 and LC-CDR1 are paired according to Table 9. In one aspect, an antibody or antigen-binding fragment thereof is provided, wherein the antibody or antigen-binding fragment thereof comprises a HC-CDR1, a HC-CDR2, and a HC-CDR3 selected from any HC-CDR1, a HC-CDR2, and a HC-CDR3 in Table 5 and a LC-CDR1, a LC-CDR2, and a LC-CDR3 selected from any LC-CDR1, a LC-CDR2, and a LC-CDR3 in Table 6, wherein the selected HC-CDR1, a HC-CDR2, a HC-CDR3, LC-CDR1, a LC-CDR2, and a LC-CDR3 are paired according to Table 9. In some embodiments, the antibody or an antigen binding fragment thereof comprises any one of a HC-CDR1, a HC-CDR2, and a HC-CDR3 selected from any HC-CDR1, a HC-CDR2, and a HC-CDR3 in Table 5 and any IgA heavy chain constant region selected from SEQ ID NOs: 16-21. In some embodiments, the antibody or an antigen binding fragment thereof comprises any one of a LC-CDR1, a LC-CDR2, and a LC-CDR3 selected from any LC-CDR1, a LC-CDR2, and a LC-CDR3 in Table 6 and any IgA heavy chain constant region selected from SEQ ID NOs: 16-21.

[0230] In some embodiments, the antibody or an antigen binding fragment thereof comprises any one of a HC-CDR1, a HC-CDR2, and a HC-CDR3 selected from any HC-CDR1, a HC-CDR2, and a HC-CDR3 in Table 5 and any one of a LC-CDR1, a LC-CDR2, and a LC-CDR3 selected from any LC-CDR1, a LC-CDR2, and a LC-CDR3 in Table 6 and any IgA heavy chain constant region selected from SEQ ID NOs: 16-21, wherein the selected HC-CDR1, HC-CDR2, HC-CDR3, selected LC-CDR1, LC-CDR2, LC-CDR3 and selected IgA heavy chain constant region are paired according to Table 9.

IgA Antibody Modifications

[0231] Described herein are IgA antibodies comprising one or more amino acid substitution and/or one or more amino acid deletions.

5 [0232] In some embodiments, the amino acid numbering of an IgA antibody described herein is indicated according to IMGT unique numbering for C-DOMAIN and C-LIKE-DOMAIN (as disclosed in “IMGT unique numbering for immunoglobulin and T cell receptor constant domains and Ig superfamily C-like domains.” Dev Comp Immunol. 2005;29(3):185-203, the entire contents of which are incorporated by reference herein). In some embodiments, the amino acid modifications disclosed herein are relative to amino
10 acid residues at select positions in the WT IgA heavy chain constant region (e.g., WT IgA2 heavy chain constant region) comprising an amino acid sequence set forth in SEQ ID NO: 1, numbering according to IMGT scheme. It is noted that US62/824,864 which is incorporated in its entirety herein also includes antibodies and antibody constructs comprising modified IgA heavy chain constant regions numbered according to the IMGT numbering scheme. Some amino acids were inadvertently mislabeled therein
15 and specific amino acid positions as described in US62/824,864 correspond to the following specific amino acid positions as provided herein. Positions C92, N120, I121, and T122 as referred to in US62/824,864 correspond to amino acid residues C86, N114, I115, and T116 in antibodies described herein which are correctly labeled as per the IMGT numbering scheme (IMGT numbering depicted in table 11, for reference to position numbering only). The reference wild type heavy chain constant
20 region sequence for these antibodies as disclosed in US62/824,864 and the current application is the same, namely SEQ ID NO:1 as shown in FIG. 1. As such the inadvertent mislabeling in US62/824,864 of residues C86, N114, I115, and T116 in the IMGT naming scheme as C92, N120, I121, and T122 would be evident to the skilled artisan.

[0233] In some embodiments, the IgA antibody disclosed herein comprises a deletion of at least four
25 glycosylation sites within the constant region. In some embodiments, the IgA antibody disclosed herein comprises a deletion of at least three N-linked glycosylation sites in the constant region of the antibody. In some embodiments, the IgA antibody comprises a deletion of at least three N-linked glycosylation sites in the constant region of the antibody and at least one O-linked glycosylation site in the constant region of the antibody. In some embodiments, the IgA antibodies, or a functional fragment disclosed herein comprises one
30 or more modifications disclosed herein (e.g., Table 2) in an IgA heavy chain constant region. In some embodiments, the IgA antibodies, or a functional fragment disclosed herein comprises one or more modifications disclosed herein (e.g., Table 2) in an IgA1 heavy chain constant region. In some embodiments, the IgA antibodies disclosed herein, or a functional fragment disclosed herein comprises one or more

modifications disclosed herein (e.g., Table 2) in an IgA2 heavy chain constant region. In some embodiments, the one or more modifications are in an amino acid residue within a CH1 region, CH2 region and/or CH3 region of a IgA1 heavy chain constant region. In some embodiments,

[0234] In some embodiments, the IgA antibodies comprise a deleted tail piece. In some embodiments, the one or more modifications are in an amino acid residue within a CH1 region, a CH2 region and/or a CH3 region of an IgA2 heavy chain constant region. In some embodiments, an IgA CH1 region comprises an amino acid sequence that at least about 80%, 85%, 90%, 95%, 99% or 100% identical to an amino acid sequence set forth in SEQ ID NO: 109. In some embodiments, a IgA CH2 region comprises an amino acid sequence that at least about 80%, 85%, 90%, 95%, 99% or 100% identical to an amino acid sequence set forth in SEQ ID NO: 110. In some embodiments, a IgA CH2 region comprises an amino acid sequence that at least about 80%, 85%, 90%, 95%, 99% or 100% identical to an amino acid sequence set forth in SEQ ID NO: 111. In some embodiments, a IgA antibody, or a functional fragment thereof as disclosed herein comprises a deletion of the 3-20, 3-19, 3-18, 3-17, 3-16, 3-15, 3-14, 3-13, 3-12, 3-11, 3-10, 3-9, 3-8, 3-7, 3-6, 3-5, 3-4 C-terminal amino acids. In some embodiments, the IgA2 antibody comprises a deletion of the 3-20, 3-19, 3-18, 3-17, 3-16, 3-15, 3-14, 3-13, 3-12, 3-11, 3-10, 3-9, 3-8, 3-7, 3-6, 3-5, 3-4 C-terminal amino acids. In some embodiments, the C-terminal amino acids are from amino acids 131-148 of the IgA2 antibody, numbering according to IMGT scheme.

[0235] In some embodiments, the IgA2 antibody comprises a deletion of amino acids 131-148, numbering according to IMGT scheme. In some embodiments, the IgA2 antibody comprises a deletion of amino acids 147-148, numbering according to IMGT scheme. In some embodiments, the IgA2 antibody comprises a deletion of amino acids 146-148, numbering according to IMGT scheme. In some embodiments, the IgA2 antibody comprises a deletion of amino acids 145-148, numbering according to IMGT scheme. In some embodiments, the IgA2 antibody comprises a deletion of amino acids 144-148, numbering according to IMGT scheme. In some embodiments, the IgA2 antibody comprises a deletion of amino acids 143-148, numbering according to IMGT scheme. In some embodiments, the IgA2 antibody comprises a deletion of amino acids 142-148, numbering according to IMGT scheme. In some embodiments, the IgA2 antibody comprises a deletion of amino acids 141-148, numbering according to IMGT scheme. In some embodiments, the IgA2 antibody comprises a deletion of amino acids 140-148, numbering according to IMGT scheme. In some embodiments, the IgA2 antibody comprises a deletion of amino acids 139-148, numbering according to IMGT scheme. In some embodiments, the IgA2 antibody comprises a deletion of amino acids 138-148, numbering according to IMGT scheme. In some embodiments, the IgA2 antibody comprises a deletion of amino acids 137-148, numbering according to IMGT scheme. In some embodiments, the IgA2 antibody comprises a deletion of amino acids 136-148, numbering according to IMGT scheme. In some embodiments, the IgA2 antibody comprises a deletion of amino acids 135-148, numbering according to IMGT scheme. In

some embodiments, the IgA2 antibody comprises a deletion of amino acids 134-148, numbering according to IMGT scheme. In some embodiments, the IgA2 antibody comprises a deletion of amino acids 133-148, numbering according to IMGT scheme. In some embodiments, the IgA2 antibody comprises a deletion of amino acids 132-148, numbering according to IMGT scheme. In some embodiments, the IgA2 antibody comprises a deletion of amino acids 131-148, numbering according to IMGT scheme. In some embodiments, the IgA2 antibody comprises a deletion of amino acids P131-Y148, numbering according to IMGT scheme.

[0236] In some embodiments, the IgA antibody comprises a mutation of the C-terminal asparagine (N) amino acid. In some embodiments, the mutation is a non-conservative amino acid substitution. In some embodiments, the mutation deletes the glycosylation site of the C-terminal asparagine (N) amino acid of the IgA. In some embodiments, the IgA2 antibody comprises a mutation of N135, numbering according to IMGT scheme. In some embodiments, the IgA2 antibody comprises a non-conservative mutation of N135, numbering according to IMGT scheme. In some embodiments, the IgA2 antibody comprises a N135Q mutation, numbering according to IMGT scheme.

[0237] In some embodiments, the IgA antibody or a functional fragment thereof disclosed herein comprises a modification in at least two naturally occurring glycosylation sites in the IgA heavy chain constant region. In some embodiments, the IgA antibody or a functional fragment thereof disclosed herein comprises a modification in at least three naturally occurring glycosylation sites in the IgA heavy chain constant region. In some embodiments, the IgA antibody or a functional fragment thereof disclosed herein comprises a modification in at least four naturally occurring glycosylation site. In some embodiments, the glycosylation site comprises a N-linked glycosylation site. In some embodiments, the glycosylation site comprises a naturally occurring asparagine residue. In some embodiments, the glycosylation site is in a CH2 region, in a CH3 region, and/or a CH1 region. In some embodiments, the IgA heavy chain constant region comprises a modification at N45.2G, N15.2, L15.3, T16, N114, I115, T116, N135, or a combination thereof, numbering according to IMGT scheme. In some embodiments, the IgA heavy chain constant region comprises an amino acid substitution that is N45.2G, N45.2A, N15.2G, N15.2Q, N15.2T, L15.3I, T16S, N114T, I115L, T116S, N135Q, or a combination thereof, numbering according to IMGT scheme. In some embodiments, provided herein is an aglycosylated antibody or a functional fragment thereof. In some embodiments, the aglycosylated antibody comprises a modification at all four naturally occurring glycosylation site in IgA2 heavy chain constant region. In some embodiments, an aglycosylated antibody provided herein comprises a modification at residues; N45.2, N15.2, L15.3, T16, N114, I115, T116, and N135, numbering according to IMGT scheme. In some embodiments, an aglycosylated antibody provided herein comprises a modification at residues; N45.2, N15.2, N114, I115, T116, and N135, numbering according to IMGT scheme.

[0238] In some embodiments, an aglycosylated antibody provided herein comprises a modification at residues; N45.2, N15.2, L15.3, T16, N114, I115, T116, and a deletion of the C-terminal tailpiece residues

P131-Y148, numbering according to IMGT scheme. In some embodiments, an aglycosylated antibody provided herein comprises a modification at residues; N45.2, N15.2, N114, I115, T116, and P131-Y148, numbering according to IMGT scheme.

[0239] In some embodiments, provided herein is an antibody or a functional fragment thereof that comprises a modification at residue N45.2, N15.2, L15.3, T16, N114, I115, T116, and N135, numbering according to IMGT scheme. In some embodiments, provided herein is an antibody or a functional fragment thereof that comprises a modification at residue N45.2, N15.2, N114, I115, T116, and N135, numbering according to IMGT scheme. In some embodiments, the IgA heavy chain constant region comprises an amino acid substitution that is N45.2G, N45.2A, N15.2G, N114T, I115L, T116S, and N135Q, numbering according to IMGT scheme. In some embodiments, the IgA heavy chain constant region comprises an amino acid substitution that is N45.2G, N45.2A, N15.2Q, L15.3I, T16S, N114T, I115L, T116S, and N135Q. In some embodiments, the IgA heavy chain constant region comprises an amino acid substitution that is N45.2G, N45.2A, N15.2G, N15.2T, N114T, I115L, T116S, and N135Q, numbering according to IMGT scheme. In some embodiments, an antibody comprises a combination of acid substitution that is N45.2G, N45.2A, N15.2G, N114T, I115L, T116S, and N135Q in the IgA heavy chain constant region is an aglycosylated antibody, numbering according to IMGT scheme. In some embodiments, the aglycosylated antibody a combination of acid substitution that is N45.2G, N45.2A, N15.2G, T16S, N114T, I115L, T116S, and deletion of the C-terminal tailpiece.

[0240] In some embodiments, an antibody comprises a combination of acid substitution that is N45.2G, N45.2A, N15.2Q, N114T, I115L, T116S, and N135Q in the IgA heavy chain constant region is an aglycosylated antibody, numbering according to IMGT scheme. In some embodiments, the aglycosylated antibody a combination of acid substitution that is N45.2G, N45.2A, N15.2Q, T16S, N114T, I115L, T116S, and deletion of the C-terminal tailpiece. In some embodiments, an antibody comprises a combination of acid substitution that is N45.2G, N45.2A, N15.2T, N114T, I115L, T116S, and N135Q in the IgA heavy chain constant region is an aglycosylated antibody, numbering according to IMGT scheme. In some embodiments, the aglycosylated antibody a combination of acid substitution that is N45.2G, N45.2A, N15.2T, T16S, N114T, I115L, T116S, and deletion of the C-terminal tailpiece. In some embodiments, an antibody comprises a combination of acid substitution that is N45.2G, N45.2A, N15.2T, L15.3I, T16S N114T, I115L, T116S, and N135Q in the IgA heavy chain constant region is an aglycosylated antibody, numbering according to IMGT scheme. In some embodiments, the aglycosylated antibody a combination of acid substitution that is N45.2G, N45.2A, N15.2T, L15.3I, T16S, T16S, N114T, I115L, T116S, and deletion of the C-terminal tailpiece.

[0241] In some embodiments, the aglycosylated antibody exhibits increased circulating half-life relative to a corresponding antibody comprising at least one glycosylation site (e.g., WT residues N45.2, N114, N15.2, and N135, numbering according to IMGT scheme) or a corresponding WT IgA antibody. In some

embodiments, the aglycosylated antibody exhibits decreased aggregation relative to a corresponding antibody comprising at least one glycosylation site (e.g., WT residues N45.2, N114, N15.2. and N135, numbering according to IMGT scheme) or a WT IgA antibody. In some embodiments, the aglycosylated antibody exhibits decreased aggregation with serum proteins relative to a corresponding antibody comprising at least one glycosylation site (e.g., WT residues N45.2, N114, N15.2. and N135, numbering according to IMGT scheme) or a corresponding WT IgA antibody. In some embodiments, the aglycosylated antibody exhibits increased thermostability relative to a corresponding antibody comprising at least one glycosylation site (e.g., WT residues N45.2, N114, N15.2. and N135, numbering according to IMGT scheme) or a corresponding WT IgA antibody. In some embodiments, the aglycosylated antibody exhibits binding to Fc receptor with increased binding affinity relative to a corresponding WT IgA. In some embodiments, the aglycosylated antibody induces ADCC of a target cell. In some embodiments, the aglycosylated antibody specifically binds a target antigen. In some embodiments, the aglycosylated antibody specifically binds a target cell.

[0242] In some embodiments, an antibody comprises a mutation of N45.2, numbering according to IMGT scheme. In some embodiments, the IgA2 antibody comprises a non-conservative mutation of N45.2, numbering according to IMGT scheme. In some embodiments, the IgA2 antibody comprises a N45.2G, numbering according to IMGT scheme. In some embodiments, the IgA2 antibody has an increased circulating half-life compared to an IgA2 antibody that does not have a mutation in the N45.2 amino acid.

[0243] In some embodiments, an antibody comprises a mutation of P124, numbering according to IMGT scheme. In some embodiments, the IgA2 antibody comprises a non-conservative mutation of P124, numbering according to IMGT scheme. In some embodiments, the IgA2 antibody comprises a P124R, numbering according to IMGT scheme. In some embodiments, the IgA2 antibody has an increased circulating half-life compared to an IgA2 antibody that does not have a mutation in the P124 amino acid. In some embodiments, the IgA2 antibody has an increased stability compared to an IgA2 antibody that does not have a mutation in the P124 amino acid.

[0244] In some embodiments, an antibody as described herein comprises a mutation of C86 (inadvertently referred to as C92 in US 62/824,864 which is incorporated in its entirety herein, as described above), numbering according to IMGT scheme. In some embodiments, the antibody comprises a non-conservative mutation of C86, numbering according to IMGT scheme. In some embodiments, the antibody comprises a C86S, numbering according to IMGT scheme. In some embodiments, the antibody has a decreased aggregation compared to an antibody that does not have a mutation in the C86 amino acid. In some embodiments, the antibody has a decreased aggregation with serum proteins compared to an antibody that

does not have a mutation in the C86 amino acid. In some embodiments, the antibody has a decreased aggregation in vitro or in vivo compared to an antibody that does not have a mutation in the C86 amino acid.

[0245] In some embodiments, the IgA2 antibody comprises a mutation of C86 according to IMGT scheme.

In some embodiments, the IgA2 antibody comprises a non-conservative mutation of C86, numbering

5 according to IMGT scheme. In some embodiments, the IgA2 antibody comprises a C86S, numbering

according to IMGT scheme. In some embodiments, the IgA2 antibody has a decreased aggregation compared

to an IgA2 antibody that does not have a mutation in the C86 amino acid. In some embodiments, the IgA2

antibody has a decreased aggregation with serum proteins compared to an IgA2 antibody that does not have a

mutation in the C86 amino acid. In some embodiments, the IgA2 antibody has a decreased aggregation in

10 vitro or in vivo compared to an IgA2 antibody that does not have a mutation in the C86 amino acid.

[0246] In some embodiments, the antibody comprises a mutation of N114 (inadvertently referred to as N120 in US 62/824,864 which is incorporated in its entirety herein, as described above), numbering according to

IMGT scheme. In some embodiments, the antibody comprises a non-conservative mutation of N114,

numbering according to IMGT scheme. In some embodiments, the antibody comprises a N114T, numbering

15 according to IMGT scheme. In some embodiments, the antibody has an increased circulating half-life

compared to an IgA2 antibody that does not have a mutation in the N114 amino acid.

[0247] In some embodiments, the antibody comprises a mutation of I115 (inadvertently referred to as I121 in US 62/824,864 which is incorporated in its entirety herein, as described above), numbering according to

IMGT scheme. In some embodiments, the antibody comprises a non-conservative mutation of I115,

20 numbering according to IMGT scheme. In some embodiments, the antibody comprises an I115L, numbering

according to IMGT scheme. In some embodiments, the antibody has an increased circulating half-life

compared to an IgA2 antibody that does not have a mutation in the I115 amino acid.

[0248] In some embodiments, the antibody comprises a mutation of T116 (inadvertently referred to as T122 in US 62/824,864 which is incorporated in its entirety herein, as described above), numbering according to

25 IMGT scheme. In some embodiments, the antibody comprises a non-conservative mutation of T116,

numbering according to IMGT scheme. In some embodiments, the antibody comprises a T116S, numbering

according to IMGT scheme. In some embodiments, the IgA2 antibody has an increased circulating half-life

compared to an IgA2 antibody that does not have a mutation in the T116 amino acid.

[0249] In some embodiments, the IgA2 antibody comprises a mutation of N114, numbering according to

30 IMGT scheme. In some embodiments, the IgA2 antibody comprises a non-conservative mutation of N114,

numbering according to IMGT scheme. In some embodiments, the IgA2 antibody comprises a N14T,

numbering according to IMGT scheme. In some embodiments, the IgA2 antibody has an increased circulating

half-life compared to an IgA2 antibody that does not have a mutation in the N114 amino acid.

[0250] In some embodiments, the IgA2 antibody comprises a mutation of I115, according to IMGT scheme. In some embodiments, the IgA2 antibody comprises a non-conservative mutation of I115, numbering according to IMGT scheme. In some embodiments, the IgA2 antibody comprises an I115L, numbering according to IMGT scheme. In some embodiments, the IgA2 antibody has an increased circulating half-life compared to an IgA2 antibody that does not have a mutation in the I115 amino acid.

[0251] In some embodiments, the IgA2 antibody comprises a mutation of T116, numbering according to IMGT scheme. In some embodiments, the IgA2 antibody comprises a non-conservative mutation of T116, numbering according to IMGT scheme. In some embodiments, the IgA2 antibody comprises a T116S, numbering according to IMGT scheme. In some embodiments, the IgA2 antibody has an increased circulating half-life compared to an IgA2 antibody that does not have a mutation in the T116 amino acid.

[0252] In some embodiments, the IgA2 antibody comprises a mutation of N15.2, according to IMGT scheme. In some embodiments, the IgA2 antibody comprises a non-conservative mutation of N15.2, numbering according to IMGT scheme. In some embodiments, the IgA2 antibody comprises an N15.2G, N15.2Q, or N15.2T, numbering according to IMGT scheme. In some embodiments, the IgA2 antibody has an increased circulating half-life compared to an IgA2 antibody that does not have a mutation in the N15.2 amino acid.

[0253] In some embodiments, the IgA2 antibody comprises a mutation of L15.3, according to IMGT scheme. In some embodiments, the IgA2 antibody comprises a non-conservative mutation of L15.3, numbering according to IMGT scheme. In some embodiments, the IgA2 antibody comprises an L15.3I, numbering according to IMGT scheme. In some embodiments, the IgA2 antibody has an increased circulating half-life compared to an IgA2 antibody that does not have a mutation in the L15.3 amino acid.

[0254] In some embodiments, the IgA2 antibody comprises a mutation of T16, according to IMGT scheme. In some embodiments, the IgA2 antibody comprises a non-conservative mutation of T16, numbering according to IMGT scheme. In some embodiments, the IgA2 antibody comprises an T16S, numbering according to IMGT scheme. In some embodiments, the IgA2 antibody has an increased circulating half-life compared to an IgA2 antibody that does not have a mutation in the T16S amino acid.

[0255] In some embodiments, the IgA2 antibody comprises a mutation of C147, numbering according to IMGT scheme. In some embodiments, the IgA2 antibody comprises a non-conservative mutation of C147, numbering according to IMGT scheme. In some embodiments, the IgA2 antibody comprises a deletion of amino acid C147, numbering according to IMGT scheme. In some embodiments, the IgA2 antibody has a decreased aggregation compared to an IgA2 antibody that does not have a mutation in the C147 amino acid.

[0256] In some embodiments, the IgA2 antibody comprises a mutation of Y148, numbering according to IMGT scheme. In some embodiments, the IgA2 antibody comprises a non-conservative mutation of Y148,

numbering according to IMGT scheme. In some embodiments, the IgA2 antibody comprises a deletion of amino acid Y148, numbering according to IMGT scheme.

[0257] In some embodiments, the IgA antibody comprises one or more albumin binding domains. In some embodiments, the one or more albumin binding domains are fused to a light chain or heavy chain of a IgA constant region. In some embodiments, the one or more albumin binding domains are fused to a heavy chain of a IgA constant region. In some embodiments, the one or more albumin binding domains are fused to a C-terminal region of a CH3 region of a heavy chain of a IgA constant region. In some embodiments, the IgA2 antibody has an increased circulating half-life compared to an IgA2 antibody that does not comprise one or more albumin binding domains. In some embodiments, the IgA2 antibody comprises one or more albumin binding domain and has circulating half-life within that of 1%, 5%, or 10% of a corresponding IgG antibody. In some embodiments, the IgA2 antibody comprises one or more albumin binding domain and has circulating half-life greater than that of a corresponding IgG antibody.

[0258] In some embodiments, the IgA antibody comprises one or more mutations described in Lohse S. et al. Cancer Res. 2015; 76(2):403-17; Meyer S. et al. mAbs. 2016; 8(1):87-98; or Leusen J. et al. Molecular Immunology. 2015; 68: 35-39.

[0259] In some embodiments, the one or more mutation or deletion results in increased or decreased circulating half-life of the IgA antibody. In some embodiments, the one or more mutation or deletion results in increased circulating half-life of the IgA antibody. For example, the one or more mutations can increase the serum half-life of the IgA antibody to up to 21 days or more in humans. Furthermore, the one or more mutations can increase the serum half-life of the IgA antibody to up to 9 days or more in mice. In some embodiments, the one or more mutations can increase the serum half-life of the IgA antibody to a level comparable to that of an immunoglobulin G (IgG) molecule. In some embodiments, the one or more mutation or deletion results in decreased circulating half-life of the IgA antibody.

[0260] In some embodiments, the one or more mutations can increase the serum half-life of the IgA antibody for at least about 7 days to about 30 days or more. In some embodiments, the one or more mutations can increase the serum half-life of the IgA antibody for at least about 7 days. In some embodiments, the one or more mutations can increase the serum half-life of the IgA antibody for at most about 30 days. In some embodiments, the one or more mutations can increase the serum half-life of the IgA antibody for about 7 days to about 8 days, about 7 days to about 9 days, about 7 days to about 10 days, about 7 days to about 15 days, about 7 days to about 20 days, about 7 days to about 25 days, about 7 days to about 30 days, about 8 days to about 9 days, about 8 days to about 10 days, about 8 days to about 15 days, about 8 days to about 20 days, about 8 days to about 25 days, about 8 days to about 30 days, about 9 days to about 10 days, about 9 days to about 15 days, about 9 days to about 20 days, about 9 days to about 25 days, about 9 days to about 30 days, about 10 days to about 15 days, about 10 days to about 20 days, about 10 days to about 25 days, about 10

- days to about 30 days, about 15 days to about 20 days, about 15 days to about 25 days, about 15 days to about 30 days, about 20 days to about 25 days, about 20 days to about 30 days, or about 25 days to about 30 days. In some embodiments, the one or more mutations can increase the serum half-life of the IgA antibody for about 7 days, about 8 days, about 9 days, about 10 days, about 15 days, about 20 days, about 25 days, or about 30 days. Accordingly, in some embodiments, the antibodies or a functional fragment thereof disclosed herein exhibit a greater circulating half-life compared to a corresponding WT IgA antibody. In some embodiments, the antibodies exhibit a circulating half-life that is greater by at least about 2%, 5%, 10%, 12%, 15%, 20%, at 25%, 50 %, 65%, 70%, 75%, 85%, 90%, 95%, 99%, 100%, 150%, and 200%, relative to a corresponding WT IgA antibody.
- 10 **[0261]** In some embodiments, the IgA antibody exhibits increased stability. In some embodiments, the one or more mutation and/or one or more deletion results in increased stability of the IgA antibody compared to a corresponding IgA antibody which does not comprise the one or mutation and/or one or more deletion.
- [0262]** In some embodiments, the IgA antibody exhibits decreased aggregation. Antibody aggregation is a more common manifestation of physical instability. Protein aggregates generally have reduced activity and more importantly, greater immunogenicity potential because of the multiplicity of epitopes and/or conformational changes. Immunoglobulin aggregates are known to cause serious renal failure and anaphylactoid reactions such as headache, fever, and chills. It is therefore advantageous to decrease aggregation in antibody therapeutics. Additionally, the aggregate level in commercial intravenous immunoglobulin products is limited to less than 5% based on the World Health Organization (WHO)
- 15 standards. In some embodiments, the one or more mutations results in decreased aggregation. In some embodiments, the one or more mutation and/or one or more deletion results in decreased aggregation of the IgA antibody compared to a corresponding IgA antibody which does not comprise the one or mutation and/or one or more deletion. In some embodiments, the antibodies or a functional fragment thereof disclosed herein exhibit a decreased aggregation compared to a corresponding WT IgA antibody. In some embodiments, the antibodies exhibit aggregation that is decreased by at least about 2%, at least 5%, at least 10%, at least 12%, at least 15%, at least 20%, at least 25%, at least 50 %, at least 65%, at least 70%, at least 75%, at least 85%, at least 90%, at least 95%, at least 99%, at least 100%, at least 150%, and at least 200%, relative to a corresponding WT IgA antibody. In some embodiments, the antibodies or a functional fragment thereof disclosed herein exhibit a decreased aggregation with serum protein compared to a corresponding WT IgA antibody. In some embodiments, the antibodies exhibit aggregation that is decreased by at least 2%, at least 5%, at least 10%, at least 12%, at least 15%, at least 20%, at least 25%, at least 50 %, at least 65%, at least 70%, at least 75%, at least 85%, at least 90%, at least 95%, at least 99%, at least 100%, at least 150%, and at least 200%, relative to a corresponding WT IgA antibody.
- 20
- 25
- 30

[0263] In some embodiments, the IgA antibodies provided herein have an aggregate level ranging from at least about 0.1 % to about 5 % at most. In some embodiments, the IgA antibodies provided herein have an aggregate level ranging from at least about 0.1 %. In some embodiments, the IgA antibodies provided herein have an aggregate level ranging from at most about 5 %. In some embodiments, the IgA antibody's provided herein have an aggregate level ranging from about 0.1 % to about 0.5 %, about 0.1 % to about 1 %, about 0.1 % to about 2 %, about 0.1 % to about 3 %, about 0.1 % to about 4 %, about 0.1 % to about 5 %, about 0.5 % to about 1 %, about 0.5 % to about 2 %, about 0.5 % to about 3 %, about 0.5 % to about 4 %, about 0.5 % to about 5 %, about 1 % to about 2 %, about 1 % to about 3 %, about 1 % to about 4 %, about 1 % to about 5 %, about 2 % to about 3 %, about 2 % to about 4 %, about 2 % to about 5 %, about 3 % to about 4 %, about 3 % to about 5 %, or about 4 % to about 5 %. In some embodiments, the IgA antibodies provided herein have an aggregate level ranging from about 0.1 %, about 0.5 %, about 1 %, about 2 %, about 3 %, about 4 %, or about 5 %.

[0264] Therapeutic antibodies disclosed herein may comprise synthetic amino acids in place of one or more naturally-occurring amino acids. Such synthetic amino acids are known in the art, and include, for example, aminocyclohexane carboxylic acid, norleucine, α -amino n-decanoic acid, homoserine, S-acetylaminomethylcysteine, trans-3- and trans-4-hydroxyproline, 4-aminophenylalanine, 4-nitrophenylalanine, 4-chlorophenylalanine, 4-carboxyphenylalanine, β -phenylserine β -hydroxyphenylalanine, phenylglycine, α -naphthylalanine, cyclohexylalanine, cyclohexylglycine, indoline-2-carboxylic acid, 1,2,3,4-tetrahydroisoquinoline-3-carboxylic acid, aminomalonic acid, aminomalonic acid monoamide, N'-benzyl-N'-methyl-lysine, N',N'-dibenzyl-lysine, 6-hydroxylysine, ornithine, α -aminocyclopentane carboxylic acid, α -aminocyclohexane carboxylic acid, α -aminocycloheptane carboxylic acid, α -(2-amino-2-norbornane)-carboxylic acid, α,γ -diaminobutyric acid, α,β -diaminopropionic acid, homophenylalanine, and α -tert-butylglycine.

[0265] Methods of substituting or deleting amino acids are known in the art. For example, amino acid substitutions or deletions can be made by site-directed mutagenesis (for example, Zoller and Smith Nucl. Acids Res. 10:6487 (1982)). Mutagenesis can be performed by synthesizing an oligonucleotide having one or more modifications within the sequence of the constant domain of an antibody to be modified. The antibodies of the present disclosure (e.g., comprising one or more modifications in the IgA heavy chain constant region) can be prepared using any mutagenesis procedure known in the art, such as site-directed mutagenesis, synthetic gene construction, semi-synthetic gene construction, random mutagenesis, shuffling, etc. Site-specific mutagenesis allows the production of mutants through the use of specific oligonucleotide sequences which encode the DNA sequence of the desired mutation, as well as a sufficient number of adjacent oligonucleotides to provide a primer sequence of sufficient size and sequence complexity to form a stable duplex on both sides of the deletion junction being traversed. Typically, a primer of about 17 to about 75

nucleotides or more in length is preferred, with about 10 to about 25 or more residues on both sides of the junction of the sequence being altered. A number of such primers introducing a variety of different mutations at one or more positions may be used to generate a library of mutants.

[0266] The technique of site-specific mutagenesis is well known in the art, (see, e.g., Kunkel et al., *Methods Enzymol.*, 154:367-82, 1987). In general, site-directed mutagenesis is performed by first obtaining a single-stranded vector or melting apart of two strands of a double stranded vector which includes within its sequence a DNA sequence which encodes the desired peptide. An oligonucleotide primer bearing the desired mutated sequence is prepared, generally synthetically. This primer is then annealed with the single-stranded vector, and subjected to DNA polymerizing enzymes such as T7 DNA polymerase, in order to complete the synthesis of the mutation-bearing strand. Thus, a heteroduplex is formed wherein one strand encodes the original non-mutated sequence and the second strand bears the desired mutation. This heteroduplex vector is then used to transform or transfect appropriate cells, such as *E. coli* cells, and clones are selected which include recombinant vectors bearing the mutated sequence arrangement. As will be appreciated, the technique typically employs a phage vector which exists in both a single stranded and double stranded form. Typical vectors useful in site-directed mutagenesis include vectors such as the M13 phage. These phage are readily commercially available and their use is generally well known to those skilled in the art. Double stranded plasmids are also routinely employed in site directed mutagenesis which eliminates the step of transferring the gene of interest from a plasmid to a phage. Site directed mutagenesis has also been used to identify amino acid residues that influence plasma clearance of murine IgG1 hinge-Fc fragments as described in Kim Jin-Kyoo et al., (1994) *Eur. J. Immunol.* 24:542-548).

[0267] Alternatively, the use of PCR with commercially available thermostable enzymes such as Taq DNA polymerase may be used to incorporate a mutagenic oligonucleotide primer into an amplified DNA fragment that can then be cloned into an appropriate cloning or expression vector. See, e.g., Tomic et al., *Nucleic Acids Res.*, 18(6):1656, 1987, and Upender et al., *Biotechniques*, 18(1):29-30, 32, 1995, for PCR-mediated mutagenesis procedures. PCR employing a thermostable ligase in addition to a thermostable polymerase may also be used to incorporate a phosphorylated mutagenic oligonucleotide into an amplified DNA fragment that may then be cloned into an appropriate cloning or expression vector (see e.g., Michael, *Biotechniques*, 16(3):410-2, 1994).

[0268] Other methods known to those of skill in art of producing sequence variants of the Fc region of an antibody or an FcR binding domain thereof can be used. For example, recombinant vectors encoding the amino acid sequence of the constant domain of an antibody or a fragment thereof may be treated with mutagenic agents, such as hydroxylamine, to obtain sequence variants. Mutants that result in desirable properties, example, increased ADCC, decreased aggregation, increased affinity for FcR and/or increased in vivo half-life can be screened using routine assays such as those described later.

[0269] Synthetic gene construction entails in vitro synthesis of a designed polynucleotide molecule to encode a polypeptide of interest. Gene synthesis can be performed utilizing a number of techniques, such as the multiplex microchip-based technology described by Tian et al. (2004, Nature 432: 1050-1054) and similar technologies wherein oligonucleotides are synthesized and assembled upon photo-programmable microfluidic chips. Single or multiple amino acid substitutions, deletions, and/or insertions can be made and tested using known methods of mutagenesis, recombination, and/or shuffling, followed by a relevant screening procedure, such as those disclosed by Reidhaar-Olson and Sauer, 1988, Science 241: 53-57; Bowie and Sauer, 1989, Proc. Natl. Acad. Sci. USA 86: 2152-2156; WO 95/17413 ; or WO 95/22625 . Other methods that can be used include error-prone PCR, phage display (e.g., Lowman et al., 1991, Biochemistry 30: 10832-10837; U.S. Patent No. 5,223,409 ; WO 92/06204) and region-directed mutagenesis (Derbyshire et al., 1986, Gene 46: 145; Ner et al., 1988, DNA 7: 127).

[0270] Mutagenesis/shuffling methods can be combined with high-throughput, automated screening methods to detect activity of cloned, mutagenized polypeptides expressed by host cells (Ness et al., 1999, Nature Biotechnology 17: 893-896). Mutagenized DNA molecules that encode active polypeptides can be recovered from the host cells and rapidly sequenced using standard methods in the art. These methods allow the rapid determination of the importance of individual amino acid residues in a polypeptide.

[0271] Semi-synthetic gene construction is accomplished by combining aspects of synthetic gene construction, and/or site-directed mutagenesis, and/or random mutagenesis, and/or shuffling. Semi-synthetic construction is typified by a process utilizing polynucleotide fragments that are synthesized, in combination with PCR techniques. Defined regions of genes may thus be synthesized de novo, while other regions may be amplified using site-specific mutagenic primers, while yet other regions may be subjected to error-prone PCR or non-error prone PCR amplification. Polynucleotide subsequences may then be shuffled.

Other Covalent Modifications

[0272] Covalent modifications of the antibody are also included within the scope of this invention. They may be made by chemical synthesis or by enzymatic or chemical cleavage of the antibody, if applicable. Other types of covalent modifications of the antibody are introduced into the molecule by reacting targeted amino acid residues of the antibody with an organic derivatizing agent that is capable of reacting with selected side chains or the N- or C-terminal residues.

[0273] Cysteinyll residues most commonly are reacted with a haloacetate (and corresponding amines), such as chloroacetic acid or chloroacetamide, to give carboxymethyl or carboxyamidomethyl derivatives. Cysteinyll residues also are derivatized by reaction with bromotrifluoroacetone, alpha-bromo-(5 imidozoyl)propionic acid, chloroacetyl phosphate, N-alkylmaleimides, 3-nitro-2-pyridyl disulfide, methyl 2-pyridyl disulfide, p-chloromercuribenzoate, 2-chloromercuri-4-nitrophenol, or chloro-7-nitrobenzo-2-oxa-1,3-diazole.

[0274] Histidyl residues are derivatized by reaction with diethylpyrocarbonate at pH 5.5-7.0 because this agent is relatively specific for the histidyl side chain. Para-bromophenacyl bromide also is useful; the reaction is preferably performed in 0.1 M sodium cacodylate at pH 6.0. Lysiny and amino-terminal residues are reacted with succinic or other carboxylic acid anhydrides. Derivatization with these agents has the effect of reversing the charge of the lysiny residues. Other suitable reagents for derivatizing alpha amino-containing residues include imidoesters such as methyl picolinimide, pyridoxal phosphate, pyridoxal, chloroborohydride, trinitrobenzenesulfonic acid, methylisourea, 2,4-pentanedione, and transaminase-catalyzed reaction with glyoxylate.

[0275] Arginy residues are modified by reaction with one or several conventional reagents, among them phenylgly-oxal, 2,3-butanedione, 1,2-cyclohexanedione, and ninhydrin. Derivatization of arginine residues requires that the reaction be performed in alkaline conditions because of the high pKa of the guanidine functional group. Furthermore, these reagents may react with the groups of lysine as well as the arginine epsilon-amino group.

[0276] The specific modification of tyrosyl residues may be made, with particular interest in introducing spectral labels into tyrosyl residues by reaction with aromatic diazonium compounds or tetranitromethane. Most commonly, N-acetylimidazole and tetranitromethane are used to form O-acetyl tyrosyl species and 3-nitro derivatives, respectively. Tyrosyl residues are iodinated using ¹²⁵I or ¹³¹I to prepare labeled proteins for use in radioimmunoassay. Carboxyl side groups (aspartyl or glutamyl) are selectively modified by reaction with carbodiimides (R-N. dbd.C.dbd.N-R'), where R and R' are different alkyl groups, such as 1-cyclohexyl-3-(2-morpholinyl-4-ethyl) carbodiimide or 1-ethyl-3-(4-azonia-4,4-dimethylpentyl)carbodiimide. Furthermore, aspartyl and glutamyl residues are converted to asparaginy and glutaminy residues by reaction with ammonium ions.

[0277] Glutaminy and asparaginy residues are frequently deamidated to the corresponding glutamyl and aspartyl residues, respectively. These residues are deamidated under neutral or basic conditions. The deamidated form of these residues falls within the scope of this invention. Other modifications include hydroxylation of proline and lysine, phosphorylation of hydroxyl groups of seryl or threony residues, methylation of the alpha amino groups of lysine, arginine, and histidine side chains (T. E. Creighton, *Proteins: Structure and Molecular Properties*, W.H. Freeman & Co., San Francisco, pp. 79-86 (1983)), acetylation of the N-terminal amine, and amidation of any C-terminal carboxyl group.

[0278] Another type of covalent modification involves chemically or enzymatically coupling glycosides to the antibody. These procedures are advantageous in that they do not require production of the antibody in a host cell that has glycosylation capabilities for N- or O-linked glycosylation. Depending on the coupling mode used, the sugar(s) may be attached to (a) arginine and histidine, (b) free carboxyl groups, (c) free sulfhydryl groups such as those of cysteine, (d) free hydroxyl groups such as those of serine, threonine, or

hydroxyproline, (e) aromatic residues such as those of phenylalanine, tyrosine, or tryptophan, or (f) the amide group of glutamine. These methods are described in W087I 05330 published 11 Sep. 1987, and in Aplin and Wriston, CRC Crit. Rev. Biochem., pp. 259-306 (1981).

[0279] Removal of any carbohydrate moieties present on the antibody may be accomplished chemically or enzymatically. Chemical deglycosylation requires exposure of the antibody to the compound
5 trifluoromethanesulfonic acid, or an equivalent compound. This treatment results in the cleavage of most or all sugars except the linking sugar (N-acetyl- glucosamine or N-acetylgalactosamine), while leaving the antibody intact. Chemical deglycosylation is described by Hakimuddin, et al. Arch. Biochem. Biophys. 259: 52 (1987) and by Edge et al. Anal. Biochem., 118: 131 (1981). Enzymatic cleavage of carbohydrate moieties
10 on antibodies can be achieved by the use of a variety of endo- and exoglycosidases as described by Thotakura et al. Meth. Enzymol. 138: 350 (1987).

[0280] Another type of covalent modification of the antibody comprises linking the antibody to one of a variety of nonproteinaceous polymers, e.g., polyethylene glycol, poly- propylene glycol, polyoxyethylated polyols, polyoxyethylated sorbitol, polyoxyethylated glucose, polyoxyethylated glycerol, polyoxyalkylenes,
15 or polysaccharide polymers such as dextran. Such methods are known in the art, see, e.g. U.S. Pat. Nos. 4,640,835; 4,496,689; 4,301,144; 4,670,417; 4,791,192, 4,179,337, 4,766,106, 4,179,337, 4,495,285, 4,609,546 or EP 315 456.

[0281]

20 IgA Antibody Targets

[0282] The IgA antibodies described herein can be used to target an antigen expressed on the surface of a cell. In some embodiments, the IgA antibody comprises an antigen binding region that specifically binds to an antigen expressed on the surface of a target cell (e.g., cancer cell). In some embodiments, the antigen is a human antigen. In some embodiments, the target cell is a human cell. In some embodiments, the IgA antibody
25 comprises an antigen binding domain that specifically binds to an antigen of one of the following proteins: CD20, GD2, CD47, CD38, EGFR, HER2, PD-L1, CD25, CD33, BCMA, CD44, CD21, CD64, α -Folate receptor, CAIX, CD30, ROR1, CEA, EGP-2, EGP-40, HER3, Folate-binding Protein, GD3, IL-13R- α 2, KDR, EDB-F, mesothelin, EGFR, MUC-1, MAGE-A1, MUC16, h5T4, PSMA, TAG-72, EGFRvIII, CD123, VEGF-R2, BCMA, CD19, CD22, CD30, CD33, CD123, CD38, CD44, CD70, CD274, CD45, CD123,
30 CD138, CD171, ROR1, EGFR, EphA2, FBP, FAP, CEA, EGP2, EGP40, TAG72, PSMA, PSA, PAP, hsp70-2, M-CSF, LAGE-Ia, p53, NKG2D ligand, B7-H6, IL-13 R α 2, IL-11 R α , MUC1, MUC16, CA9, GD3, HMW-MAA, CD171, Lewis Y, G250/CAIX, HLA-AI MAGE A1, HLA-A2 NY-ESO-1, PSC1, PCTA-1, MAGE, ELF2M, IGF-I, IGF-II, IGF-I receptor, hTERT, WT1, MUC1, LMP2, HPV16, HPV18, RGL4, MelanA, MART, ML-IAP, AFP, BCR, ABL, CYP1B1, PLAC1, BORIS, NY-BR-1, RGS5, SART3, EphA2,

Glypican-3, 5T4, 8H9, $\alpha\beta$ 6 integrin, B7-H3, B7-H6, CAIX, CA9, CSPG4, EGP2, EGP40, EPCAM, ERBB3, ERBB4, ErbB3/4, FAP, FAR, FBP, KDR, MCSP, Mucl, Mucl6, NCAM, PRAME, ROR1, CD44v7/8, 8H9, NCAM, VEGF-R, TAG72, RAGE-1, MN-CA IX, RU1, RU2 (AS), fetal AchR, TEM1, TEM8, PAX5, OY-TES1, LCK, HMWMAA, AKAP-4, SSX2, XAGE 1, tie 2, PDGFR- β , kallikrein 4, PBF, PRAME, HSDL1, CA125, TADG-12, MUC16, mannan-MIC-1, HERV-K-MEL, KK-LC-1, KM-HN-1, LAGE-1, MAGE-A4, SP17, SSX4, TAG1, TAG2, ENAH, mammaglobin-A, NY-BR-1, BAGE-1, HERV-K-MEL, KK-LC-1, KM-KN-1, LAGE1, MAGE1A, MAGEA2, mucink, TRAG3, c-myc, cyclin B1, p62, DKK1, RU2AS, k-ras, ME1, NFYC, STEAP1, FGF5, RU2AS, hsp70-2, ARTC1, B-RAF, beta-catenin, CDC27, CDK4, CDK12, CDKN2A, CLPP, CSNK1A1, FN1, GAS7, GPNMB, HAUS3, LDLR-fucosyltransferase, MART2, MATN, MUM1, MUM2, MUM3, neo-PAP, myosin, PPP1R3B, PRDX5, PTPRK, RBAF600, SIRT2, SNRPD1, triosephosphate isomerase, OA1, RAB38, TRP1, TRP2, melan-A, BAGE1, GAGE1, GAGE2, GAGE8, GAGE3, GAGE4, GAGE5, GAGE6, GAGE7, GNTVF, LY6K, TRAG3, CASP8, SAGE, DEK-CAN, EFTUD2, FLT3-ITD, cyclin A1, FNDC3B, MAGEAG, G250, hepsin, intestinal carboxyl esterase, PBF, CASP5, COA1, OGT, OS9, CALCA, MDM2, alpha actinin4, elongation factor 2, fos-related antigen 1, legumain, sperm protein 17, carbonic anhydrase IX, folate receptor- α , neutrophil elastase, ephrinB2, glioma-associated antigen, β -human chorionic gonadotropin, alphafetoprotein thyroglobulin, telomerase reverse transcriptase, intestinal carboxy esterase, prostatein, or survivin.

Ganglioside G2

[0283] In some embodiments, the IgA antibody or a functional fragment thereof disclosed herein (i.e., antibodies comprising one or more modifications disclosed herein in the IgA heavy chain constant region) specifically binds GD2. In some embodiments, the IgA antibody binds O-acetylated GD2. In some embodiments, the IgA antibody binds GD2 and does not bind O-acetylated GD2. In some embodiments, the IgA antibody binds O-acetylated GD2 and does not bind GD2. In some embodiments, the antibody binds GD2 and binds O-acetylated GD2. In some embodiments, an antibody or a functional fragment disclosed herein specifically binds a human GD2 polypeptide. The polypeptide and coding nucleic acid sequences of GD2 of human origin and those of a number of animals are publicly available, e.g., from the NCBI website.

[0284] In some embodiments, the IgA antibody comprises at least CDR3 of the heavy chain of antibody ch14.18. In some embodiments, the IgA antibody comprises one, two, or three of CDR1, CDR2, or CDR3 of the heavy chain of antibody ch14.18. In some embodiments, the IgA antibody comprises one, two, or three of CDR1, CDR2, or CDR3 of the light chain of antibody ch14.18. In some embodiments, the IgA antibody comprises one, two, or three of CDR1, CDR2, or CDR3 of the heavy chain of antibody ch14.18, and one, two, or three of CDR1, CDR2, or CDR3 of the light chain of antibody ch14.18. In some embodiments, the IgA antibody comprises CDR1, CDR2, or CDR3 of the heavy chain of antibody ch14.18, and CDR1, CDR2, or

CDR3 of the light chain of antibody ch14.18. In some embodiments, the IgA antibody comprises at least CDR3 of the heavy chain of antibody ch14.18 and an IgA hinge. In some embodiments, the IgA antibody comprises at least CDR3 of the heavy chain of antibody ch14.18 an IgA hinge, a CH1 IgA region. In some embodiments, the IgA antibody comprises at least CDR3 of the heavy chain of antibody ch14.18 an IgA hinge, a CH1 IgA region, and a CH2 IgA region. In some embodiments, the IgA antibody comprises at least CDR3 of the heavy chain of antibody ch14.18 an IgA hinge, a CH1 IgA region, a CH2 IgA region, and a CH3 IgA region. In some embodiments, the IgA antibody comprises a heavy chain variable region comprising the amino acid sequence:

EVQLLQSGPELEKPGASVMISCKASGSSFTGYNMNWVRQNIQKSLEWIGAIDPYYGGTSYNQKFKGRATLTVD

10 KSSSTAYMHLKSLTSEDSAVYYCVSGMEYWGQGTSTVTVSS [SEQ ID NO: 4]. In some embodiments, the IgA antibody comprises a light chain variable region comprising the amino acid sequence:

EIVMTQSPATLSVSPGERATLSCRSSQSLVHRNGNTYLHWYLQKPGQSPK

LLIHKVSNRFSGVDPDRFSGSGSGTDFTLKISRVEAEDLGVYFCSQSTHVPPLTFGAGTKLELK [SEQ ID NO:

5]. In some embodiments, the IgA antibody comprises a heavy chain variable region comprising the amino acid sequence:

EVQLLQSGPELEKPGASVMISCKASGSSFTGYNMNWVRQNIQKSLEWIGAIDPYYGGTSYNQKFKGRATLTVD

KSSSTAYMHLKSLTSEDSAVYYCVSGMEYWGQGTSTVTVSS [SEQ ID NO: 4], and a light chain variable region comprising the amino acid sequence:

EIVMTQSPATLSVSPGERATLSCRSSQSLVHRNGNTYLHWYLQKPGQSPKLLIHKVSNRFSGVDPDRFSGSGSG

20 TDFTLKISRVEAEDLGVYFCSQSTHVPPLTFGAGTKLELK [SEQ ID NO: 5].

[0285] In some embodiments, the IgA antibody comprises a variable heavy chain comprising one or more of: CDR1 comprising the amino acid sequence: EFTFTDYY [SEQ ID NO: 10]; CDR2 comprising the amino acid sequence: IRNRANGYTT [SEQ ID NO: 11]; CDR3 comprising the amino acid sequence: ARVSNWAFDY [SEQ ID NO: 12]. In some embodiments, the IgA antibody comprises a variable light chain comprising one or more of: CDR1 comprising the amino acid sequence: QSLKNNNGNTFL [SEQ ID NO: 13]; CDR2 comprising the amino acid sequence: KVS [SEQ ID NO: 14]; CDR3 comprising the amino acid sequence: SQSTHIPPYT [SEQ ID NO: 15].

[0286] In some embodiments, the IgA antibody comprises a variable heavy chain comprising one or more of: CDR1 comprising the amino acid sequence: EFTFTDYY [SEQ ID NO: 10]; CDR2 comprising the amino acid sequence: IRNRANGYTT [SEQ ID NO: 11]; CDR3 comprising the amino acid sequence: ARVSNWAFDY [SEQ ID NO: 12]; and a variable light chain comprising one or more of: CDR1 comprising the amino acid sequence: QSLKNNNGNTFL [SEQ ID NO: 13]; CDR2 comprising the amino acid sequence: KVS [SEQ ID NO: 14]; CDR3 comprising the amino acid sequence: SQSTHIPPYT [SEQ ID NO: 15].

[0287] In some embodiments, the anti-GD2 IgA antibody comprises one or more of (e.g., two, three, four, five, or six) HC-CDR1, HC-CDR2, HC-CDR3, LC-CDR1, LC-CDR2, LC-CDR3 of 3F8 antibody. In some embodiments, the anti-GD2 antibody comprises a variable heavy and/or variable light chain of 3F8 antibody.

[0288] In one aspect, the present disclosure provides an antibody or antigen-binding fragment thereof
5 comprises one or more variable regions selected from the group consisting of (a) VH comprising the amino acid sequence of SEQ ID NO: 4, (b) VL comprising the amino acid sequence of SEQ ID NO: 5, and (c) a combination thereof.

[0289] In one aspect, the present disclosure provides an antibody or antigen-binding fragment thereof
10 comprising at least one, two, three, four, five, or six CDRs selected from (a) HC-CDR1 comprising the amino acid sequence of SEQ ID NO: 34; (b) HC-CDR2 comprising the amino acid sequence of SEQ ID NO: 42; (c) HC-CDR3 comprising the amino acid sequence of SEQ ID NO: 50; (d) LC-CDR1 comprising the amino acid sequence of SEQ ID NO: 58; (e) LC-CDR2 comprising the amino acid sequence of SEQ ID NO: 66; and (f) LC-CDR3 comprising the amino acid sequence of SEQ ID NO: 74.

[0290] In one aspect, the present disclosure provides an antibody or antigen-binding fragment thereof
15 comprising at least one, at least two, or all three VH CDR sequences selected from (a) HC-CDR1 comprising the amino acid sequence of SEQ ID NO: 34; (b) HC-CDR2 comprising the amino acid sequence of SEQ ID NO: 42; and (c) HC-CDR3 comprising the amino acid sequence of SEQ ID NO: 50; and (d) a VL comprising the amino acid sequence of SEQ ID NO: 5.

[0291] In one aspect, the present disclosure provides an antibody or antigen-binding fragment thereof
20 comprising at least one, at least two, or all three VL CDR sequences selected from (a) LC-CDR1 comprising the amino acid sequence of SEQ ID NO: 58; (b) LC-CDR2 comprising the amino acid sequence of SEQ ID NO: 66; and (c) LC-CDR3 comprising the amino acid sequence of SEQ ID NO: 74; and a VH comprising the amino acid sequence of SEQ ID NO: 4.

[0292] In one aspect, the present disclosure provides an antibody or antigen-binding fragment thereof
25 comprising the CDRs: HC-CDR3 comprising the amino acid sequence of SEQ ID NO: 50; and LC-CDR3 comprising the amino acid sequence of SEQ ID NO: 74.

[0293] In one aspect, the disclosure herein provides an antibody or antigen-binding fragment thereof
30 comprising at least one, at least two, or all three VL CDR sequences selected from (a) LC-CDR1 comprising the amino acid sequence of SEQ ID NO: 58; (b) LC-CDR2 comprising the amino acid sequence of SEQ ID NO: 66 and (c) LC-CDR3 comprising the amino acid sequence of SEQ ID NO: 74. In one aspect, the disclosure provides an antibody or antigen-binding fragment thereof comprising at least one, at least two, or all three VH CDR sequences selected from (a) HC-CDR1 comprising the amino acid sequence of SEQ ID NO: 34; (b) HC-CDR2 comprising the amino acid sequence of SEQ ID NO: 42 and (c) HC-CDR3 comprising the amino acid sequence of SEQ ID NO: 50.

[0294] In one aspect, the disclosure provides an antibody or antigen-binding fragment thereof comprising the CDRs: (a) HC-CDR1 comprising the amino acid sequence of SEQ ID NO: 34; (b) HC-CDR2 comprising the amino acid sequence of SEQ ID NO: 42; (c) HC-CDR3 comprising the amino acid sequence of SEQ ID NO: 50; (d) LC-CDR1 comprising the amino acid sequence of SEQ ID NO: 58; (e) LC-CDR2 comprising the amino acid sequence of SEQ ID NO: 66; and (f) LC-CDR3 comprising the amino acid sequence of SEQ ID NO: 74.

[0295] In one aspect, an antibody or antigen-binding fragment thereof comprises a VH sequence having at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% sequence identity to the amino acid sequence of SEQ ID NO: 4. In some embodiments, a VH sequence having at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% identity contains substitutions (e.g., conservative substitutions), insertions, or deletions relative to the reference sequence, but an antibody or antigen-binding fragment thereof comprising that sequence retains the ability to bind to antigen. In some embodiments, a total of 1 to 10 amino acids have been substituted, inserted and/or deleted in the amino acid sequence of SEQ ID NO: 4. In some embodiments, substitutions, insertions, or deletions occur in regions outside the CDRs (e.g., in the FRs).
Optionally, the antibody or antigen-binding fragment thereof comprises the VH sequence of the amino acid sequence of SEQ ID NO: 4, including post-translational modifications of that sequence. In a particular embodiment, the VH comprises one, two or three CDRs selected from: (a) HC-CDR1 comprising the amino acid sequence of SEQ ID NO: 34, (b) HC-CDR2 comprising the amino acid sequence of SEQ ID NO: 42, and (c) HC-CDR3 comprising the amino acid sequence of SEQ ID NO: 50.

[0296] In one aspect, an antibody or antigen-binding fragment thereof is provided, wherein the antibody or antigen-binding fragment thereof comprises a light chain variable domain (VL) having at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% sequence identity to the amino acid sequence of SEQ ID NO: 5. In some embodiments, a VL sequence having at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% identity contains substitutions (e.g., conservative substitutions), insertions, or deletions relative to the reference sequence, but an antibody or antigen-binding fragment thereof comprising that sequence retains the ability to bind to antigen. In some embodiments, a total of 1 to 10 amino acids have been substituted, inserted and/or deleted in any one of the amino acid sequence of SEQ ID NO: 5. In some embodiments, the substitutions, insertions, or deletions occur in regions outside the CDRs (e.g., in the FRs).
Optionally, the antibody or antigen-binding fragment thereof comprises the VL sequence of SEQ ID NO: 5, including post-translational modifications of that sequence. In a particular embodiment, the VL comprises one, two or three CDRs selected from (a) LC-CDR1 comprising the amino acid sequence of SEQ ID NO: 58; (b) LC-CDR2 comprising the amino acid sequence of SEQ ID NO: 66; and (c) LC-CDR3 comprising the amino acid sequence of SEQ ID NO: 74.

[0297] In one aspect, an antibody or antigen-binding fragment thereof is provided, wherein the antibody or antigen-binding fragment thereof comprises a VH as in any of the embodiments provided above, and a VL as in any of the embodiments provided above. In some embodiments, the antibody comprises a VH comprising the amino acid sequence of SEQ ID NO: 4, and a VL sequence in SEQ ID NO: 5, including post-translational modifications of those sequences. In some embodiments, the antibody or antigen binding fragment thereof comprises a IgA heavy chain constant region having at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% sequence identity to the amino acid sequence of any one of SEQ ID NOs: 16-21. In some embodiments, the antibody or antigen binding fragment thereof comprises a kappa light chain constant region of SEQ ID NO: 31.

CD20

[0298] In some embodiments, the IgA antibody specifically binds CD20. In some embodiments, the IgA antibody comprises an antigen binding domain that comprises a Type II or a Type I/II CD20 binding region. In some embodiments, an antibody or a functional fragment disclosed herein specifically binds a human CD20 polypeptide. The polypeptide and coding nucleic acid sequences of CD20 of human origin and those of a number of animals are publicly available, e.g., from the NCBI website. In some embodiments, the IgA antibody comprises an antigen binding domain which specifically binds to a CD20 epitope, wherein said CD20 epitope is within the following amino acid sequence: YNCEPANPSEKNSPSTQYCYS [SEQ ID NO: 6].

[0299] In some embodiments, the variable region of the CD20 antibody is described in PCTNL2017050581, which is incorporated by reference herein in its entirety. In some embodiments, the heavy chain variable region comprises one, two, or three of CDR1, CDR2, OR CDR3 within the following amino acid sequence:

QAYLQQSGAELVRPGASVKMSCKASGYTFT

SYNLHWVKQTPRQGLEWIGAIYPGNGDTSYNQKFKGKATLTVDKSSSTAYMQLSRLTSED

SAVYFCARSNSYGSTYWFYFDVWGTGTTVTVSS [SEQ ID NO: 7]. In some embodiments, the light chain variable region comprises one, two, or three of CDR1, CDR2, OR CDR3 within the following amino acid sequence: QIVLSQSPAVLFASPGEKVTMTCRARSSVSMDWYQQKPRSSPKPWIYATSNLASGVPARFSGSGSGTSYSLTISRVEAEDAATYYCQQWTSNPPTFGSGTKLEIKRADAAPTVSIFPPSS [SEQ ID NO: 8].

[0300] In some embodiments, the heavy chain variable region comprises the following amino acid sequence:

QAYLQQSGAELVRPGASVKMSCKASGYTFTSYNLHWVKQTPRQGLEWIGAIYPGNGDTSYNQKFKGKATLTVDKSSSTAYMQLSRLTSED SAVYFCARSNSYGSTYWFYFDVWGTGTTVTVSS [SEQ ID NO: 7]. In some

embodiments, the light chain variable region comprises the following amino acid sequence:

QIVLSQSPAVLFASPGEKVTMTCRARSSVSMDWYQQKPRSSPKPWIYATSNLASGVPARFSGSG

SGTSYSLTISRVEAEDAATYYCQQWTSNPPTFGSGTKLEIKRADAAPTVSIFPPSS [SEQ ID NO: 8].

- [0301] In some embodiments, the IgA antibody comprises at least CDR3 of the heavy chain of antibody Obinutuzumab. In some embodiments, the IgA antibody comprises one, two, or three of CDR1, CDR2, or CDR3 of the heavy chain of antibody Obinutuzumab. In some embodiments, the IgA antibody comprises one, two, or three of CDR1, CDR2, or CDR3 of the light chain of antibody Obinutuzumab. In some embodiments, the IgA antibody comprises one, two, or three of CDR1, CDR2, or CDR3 of the heavy chain of antibody Obinutuzumab, and one, two, or three of CDR1, CDR2, or CDR3 of the light chain of antibody Obinutuzumab. In some embodiments, the IgA antibody comprises CDR1, CDR2, or CDR3 of the heavy chain of antibody Obinutuzumab, and CDR1, CDR2, or CDR3 of the light chain of antibody Obinutuzumab. In some embodiments, the IgA antibody comprises at least CDR3 of the heavy chain of antibody Obinutuzumab and an IgA hinge. In some embodiments, the IgA antibody comprises at least CDR3 of the heavy chain of antibody Obinutuzumab an IgA hinge, a CH1 IgA region. In some embodiments, the IgA antibody comprises at least CDR3 of the heavy chain of antibody Obinutuzumab an IgA hinge, a CH1 IgA region, and a CH2 IgA region. In some embodiments, the IgA antibody comprises at least CDR3 of the heavy chain of antibody Obinutuzumab an IgA hinge, a CH1 IgA region, a CH2 IgA region, and a CH3 IgA region.
- 15 [0302] In some embodiments, the IgA antibody specifically binds the CD20 epitope EPANPSEK.
- [0303] In some embodiments, the IgA antibody specifically binds CD20 and has increased programmed cell death (PCD) function compared to rituximab with a constant region of the same isotype. In some embodiments, the IgA antibody specifically binds CD20 and has increased antibody dependent cell mediated cytotoxicity (ADCC) functionality compared to rituximab with a constant region of the same isotype. In some
- 20 embodiments, the IgA antibody specifically binds CD20 and has increased complement dependent cytotoxicity (CDC) functionality compared to rituximab with a constant region of the same isotype.
- [0304] In some embodiments, the IgA antibody has a shorter circulating half-life compared to a corresponding IgG antibody. In some embodiments, administration of the anti-CD20 IgA antibody is associated with fewer side effects from B cell depletion compared to a corresponding IgG antibody. In some
- 25 embodiments, administration of the anti-CD20 antibody is associated with faster B-cell replenishment compared to a corresponding IgG antibody.
- [0305] In one aspect, the present disclosure provides an antibody or antigen-binding fragment thereof comprising one or more variable regions selected from the group consisting of (a) VH comprising the amino acid sequence of SEQ ID NO: 81, (b) VL comprising the amino acid sequence of SEQ ID NO: 95, and (c) a
- 30 combination thereof.
- [0306] In one aspect, the present disclosure provides an antibody or antigen-binding fragment thereof comprising at least one, two, three, four, five, or six CDRs selected from (a) HC-CDR1 comprising the amino acid sequence of SEQ ID NO: 33; (b) HC-CDR2 comprising the amino acid sequence of SEQ ID NO: 41; (c) HC-CDR3 comprising the amino acid sequence of SEQ ID NO: 49; (d) LC-CDR1 comprising the amino acid

sequence of SEQ ID NO: 57; (e) LC-CDR2 comprising the amino acid sequence of SEQ ID NO: 65; and (f) LC-CDR3 comprising the amino acid sequence of SEQ ID NO: 73.

5 [0307] In one aspect, the present disclosure provides an antibody or antigen-binding fragment thereof comprising at least one, at least two, or all three VH CDR sequences selected from (a) HC-CDR1 comprising the amino acid sequence of SEQ ID NO: 33; (b) HC-CDR2 comprising the amino acid sequence of SEQ ID NO: 41; and (c) HC-CDR3 comprising the amino acid sequence of SEQ ID NO: 49; and (d) a VL comprising the amino acid sequence of SEQ ID NO: 95.

10 [0308] In one aspect, the present disclosure provides an antibody or antigen-binding fragment thereof comprising at least one, at least two, or all three VL CDR sequences selected from (a) LC-CDR1 comprising the amino acid sequence of SEQ ID NO: 57; (b) LC-CDR2 comprising the amino acid sequence of SEQ ID NO: 65; and (c) LC-CDR3 comprising the amino acid sequence of SEQ ID NO: 73; and a VH comprising the amino acid sequence of SEQ ID NO: 81.

15 [0309] In one aspect, the present disclosure provides an antibody or antigen-binding fragment thereof comprising the CDRs: HC-CDR3 comprising the amino acid sequence of SEQ ID NO: 49; and LC-CDR3 comprising the amino acid sequence of SEQ ID NO: 73.

20 [0310] In one aspect, the disclosure herein provides an antibody or antigen-binding fragment thereof comprising at least one, at least two, or all three VL CDR sequences selected from (a) LC-CDR1 comprising the amino acid sequence of SEQ ID NO: 57; (b) LC-CDR2 comprising the amino acid sequence of SEQ ID NO: 65 and (c) LC-CDR3 comprising the amino acid sequence of SEQ ID NO: 73. In one aspect, the disclosure provides an antibody or antigen-binding fragment thereof comprising at least one, at least two, or all three VH CDR sequences selected from (a) HC-CDR1 comprising the amino acid sequence of SEQ ID NO: 33; (b) HC-CDR2 comprising the amino acid sequence of SEQ ID NO: 41 and (c) HC-CDR3 comprising the amino acid sequence of SEQ ID NO: 49.

25 [0311] In one aspect, the disclosure provides an antibody or antigen-binding fragment thereof comprising the CDRs: (a) HC-CDR1 comprising the amino acid sequence of SEQ ID NO: 33; (b) HC-CDR2 comprising the amino acid sequence of SEQ ID NO: 41; (c) HC-CDR3 comprising the amino acid sequence of SEQ ID NO: 49; (d) LC-CDR1 comprising the amino acid sequence of SEQ ID NO: 57; (e) LC-CDR2 comprising the amino acid sequence of SEQ ID NO: 65; and (f) LC-CDR3 comprising the amino acid sequence of SEQ ID NO: 73.

30 [0312] In one aspect, an antibody or antigen-binding fragment thereof comprises a VH sequence having at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% sequence identity to the amino acid sequence of SEQ ID NO: 81. In some embodiments, a VH sequence having at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% identity contains substitutions (e.g., conservative substitutions), insertions, or deletions relative to the reference sequence, but an antibody or antigen-binding fragment thereof

comprising that sequence retains the ability to bind to antigen. In some embodiments, a total of 1 to 10 amino acids have been substituted, inserted and/or deleted in the amino acid sequence of SEQ ID NO: 81. In some embodiments, substitutions, insertions, or deletions occur in regions outside the CDRs (e.g., in the FRs).

Optionally, the antibody or antigen-binding fragment thereof comprises the VH sequence of the amino acid sequence of SEQ ID NO: 81, including post-translational modifications of that sequence. In a particular embodiment, the VH comprises one, two or three CDRs selected from: (a) HC-CDR1 comprising the amino acid sequence of SEQ ID NO: 33, (b) HC-CDR2 comprising the amino acid sequence of SEQ ID NO: 41, and (c) HC-CDR3 comprising the amino acid sequence of SEQ ID NO: 49.

[0313] In one aspect, an antibody or antigen-binding fragment thereof is provided, wherein the antibody or antigen-binding fragment thereof comprises a light chain variable domain (VL) having at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% sequence identity to the amino acid sequence of SEQ ID NO: 95. In some embodiments, a VL sequence having at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% identity contains substitutions (e.g., conservative substitutions), insertions, or deletions relative to the reference sequence, but an antibody or antigen-binding fragment thereof comprising that sequence retains the ability to bind to antigen. In some embodiments, a total of 1 to 10 amino acids have been substituted, inserted and/or deleted in any one of the amino acid sequence of SEQ ID NO: 95. In some embodiments, the substitutions, insertions, or deletions occur in regions outside the CDRs (e.g., in the FRs). Optionally, the antibody or antigen-binding fragment thereof comprises the VL sequence of SEQ ID NO: 95, including post-translational modifications of that sequence. In a particular embodiment, the VL comprises one, two or three CDRs selected from (a) LC-CDR1 comprising the amino acid sequence of SEQ ID NO: 57; (b) LC-CDR2 comprising the amino acid sequence of SEQ ID NO: 5; and (c) LC-CDR3 comprising the amino acid sequence of SEQ ID NO: 73.

[0314] In one aspect, an antibody or antigen-binding fragment thereof is provided, wherein the antibody or antigen-binding fragment thereof comprises a VH as in any of the embodiments provided above, and a VL as in any of the embodiments provided above. In some embodiments, the antibody comprises a VH comprising the amino acid sequence of SEQ ID NO: 81, and a VL sequence in SEQ ID NO: 95, including post-translational modifications of those sequences. In some embodiments, the antibody or antigen binding fragment thereof comprises a IgA heavy chain constant region having at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% sequence identity to the amino acid sequence of any one of SEQ ID NOs: 16-21. In some embodiments, the antibody or antigen binding fragment thereof comprises a kappa light chain constant region of SEQ ID NO: 31.

[0315] In some embodiments, provided herein is an antibody or a functional fragment thereof that specifically binds CD47. In some embodiments, the IgA antibody comprises at least CDR3 of the heavy chain of antibody UMAB10. In some embodiments, the IgA antibody comprises one, two, or three of CDR1,

CDR2, or CDR3 of the heavy chain of antibody UMAB10. In some embodiments, the IgA antibody comprises one, two, or three of CDR1, CDR2, or CDR3 of the light chain of antibody UMAB10. In some embodiments, the IgA antibody comprises one, two, or three of CDR1, CDR2, or CDR3 of the heavy chain of antibody UMAB10, and one, two, or three of CDR1, CDR2, or CDR3 of the light chain of antibody UMAB10. In some
5 UMAB10. In some embodiments, the IgA antibody comprises CDR1, CDR2, or CDR3 of the heavy chain of antibody UMAB10, and CDR1, CDR2, or CDR3 of the light chain of antibody UMAB10. In some embodiments, the IgA antibody comprises at least CDR3 of the heavy chain of antibody UMAB10 and an IgA hinge. In some embodiments, the IgA antibody comprises at least CDR3 of the heavy chain of antibody UMAB10 an IgA hinge, a CH1 IgA region. In some embodiments, the IgA antibody comprises at least CDR3
10 of the heavy chain of antibody UMAB10 an IgA hinge, a CH1 IgA region, and a CH2 IgA region. In some embodiments, the IgA antibody comprises at least CDR3 of the heavy chain of antibody UMAB10 an IgA hinge, a CH1 IgA region, a CH2 IgA region, and a CH3 IgA region.

[0316] In one aspect, the present disclosure provides an antibody or antigen-binding fragment thereof comprising one or more variable regions selected from the group consisting of (a) VH comprising the amino
15 acid sequence of SEQ ID NO: 86, (b) VL comprising the amino acid sequence of SEQ ID NO: 100, and (c) a combination thereof.

[0317] In one aspect, the present disclosure provides an antibody or antigen-binding fragment thereof comprising at least one, two, three, four, five, or six CDRs selected from (a) HC-CDR1 comprising the amino acid sequence of SEQ ID NO: 40; (b) HC-CDR2 comprising the amino acid sequence of SEQ ID NO: 48; (c)
20 HC-CDR3 comprising the amino acid sequence of SEQ ID NO: 56; (d) LC-CDR1 comprising the amino acid sequence of SEQ ID NO: 64; (e) LC-CDR2 comprising the amino acid sequence of SEQ ID NO: 72; and (f) LC-CDR3 comprising the amino acid sequence of SEQ ID NO: 80.

[0318] In one aspect, the present disclosure provides an antibody or antigen-binding fragment thereof comprising at least one, at least two, or all three VH CDR sequences selected from (a) HC-CDR1 comprising
25 the amino acid sequence of SEQ ID NO: 40; (b) HC-CDR2 comprising the amino acid sequence of SEQ ID NO: 48; and (c) HC-CDR3 comprising the amino acid sequence of SEQ ID NO: 56; and (d) a VL comprising the amino acid sequence of SEQ ID NO: 8.

[0319] In one aspect, the present disclosure provides an antibody or antigen-binding fragment thereof comprising at least one, at least two, or all three VL CDR sequences selected from (a) LC-CDR1 comprising
30 the amino acid sequence of SEQ ID NO: 64; (b) LC-CDR2 comprising the amino acid sequence of SEQ ID NO: 72; and (c) LC-CDR3 comprising the amino acid sequence of SEQ ID NO: 80; and a VH comprising the amino acid sequence of SEQ ID NO: 7.

[0320] In one aspect, the present disclosure provides an antibody or antigen-binding fragment thereof comprising the CDRs: HC-CDR3 comprising the amino acid sequence of SEQ ID NO: 56; and LC-CDR3 comprising the amino acid sequence of SEQ ID NO: 80.

[0321] In one aspect, the disclosure herein provides an antibody or antigen-binding fragment thereof comprising at least one, at least two, or all three VL CDR sequences selected from (a) LC-CDR1 comprising the amino acid sequence of SEQ ID NO: 64; (b) LC-CDR2 comprising the amino acid sequence of SEQ ID NO: 72 and (c) LC-CDR3 comprising the amino acid sequence of SEQ ID NO: 80. In one aspect, the disclosure provides an antibody or antigen-binding fragment thereof comprising at least one, at least two, or all three VH CDR sequences selected from (a) HC-CDR1 comprising the amino acid sequence of SEQ ID NO: 40; (b) HC-CDR2 comprising the amino acid sequence of SEQ ID NO: 48 and (c) HC-CDR3 comprising the amino acid sequence of SEQ ID NO: 56.

[0322] In one aspect, the disclosure provides an antibody or antigen-binding fragment thereof comprising the CDRs: (a) HC-CDR1 comprising the amino acid sequence of SEQ ID NO: 40; (b) HC-CDR2 comprising the amino acid sequence of SEQ ID NO: 48; (c) HC-CDR3 comprising the amino acid sequence of SEQ ID NO: 56; (d) LC-CDR1 comprising the amino acid sequence of SEQ ID NO: 64; (e) LC-CDR2 comprising the amino acid sequence of SEQ ID NO: 72; and (f) LC-CDR3 comprising the amino acid sequence of SEQ ID NO: 80.

[0323] In one aspect, an antibody or antigen-binding fragment thereof comprises a VH sequence having at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% sequence identity to the amino acid sequence of SEQ ID NO: 7. In some embodiments, a VH sequence having at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% identity contains substitutions (e.g., conservative substitutions), insertions, or deletions relative to the reference sequence, but an antibody or antigen-binding fragment thereof comprising that sequence retains the ability to bind to antigen. In some embodiments, a total of 1 to 10 amino acids have been substituted, inserted and/or deleted in the amino acid sequence of SEQ ID NO: 7. In some embodiments, substitutions, insertions, or deletions occur in regions outside the CDRs (e.g., in the FRs). Optionally, the antibody or antigen-binding fragment thereof comprises the VH sequence of the amino acid sequence of SEQ ID NO: 7, including post-translational modifications of that sequence. In a particular embodiment, the VH comprises one, two or three CDRs selected from: (a) HC-CDR1 comprising the amino acid sequence of SEQ ID NO: 40, (b) HC-CDR2 comprising the amino acid sequence of SEQ ID NO: 48, and (c) HC-CDR3 comprising the amino acid sequence of SEQ ID NO: 56.

[0324] In one aspect, an antibody or antigen-binding fragment thereof is provided, wherein the antibody or antigen-binding fragment thereof comprises a light chain variable domain (VL) having at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% sequence identity to the amino acid sequence of SEQ ID NO: 8. In some embodiments, a VL sequence having at least 90%, 91%, 92%, 93%, 94%, 95%, 96%,

97%, 98%, or 99% identity contains substitutions (e.g., conservative substitutions), insertions, or deletions relative to the reference sequence, but an antibody or antigen-binding fragment thereof comprising that sequence retains the ability to bind to antigen. In some embodiments, a total of 1 to 10 amino acids have been substituted, inserted and/or deleted in any one of the amino acid sequence of SEQ ID NO: 8. In some
5 embodiments, the substitutions, insertions, or deletions occur in regions outside the CDRs (e.g., in the FRs). Optionally, the antibody or antigen-binding fragment thereof comprises the VL sequence of SEQ ID NO: 8, including post-translational modifications of that sequence. In a particular embodiment, the VL comprises one, two or three CDRs selected from (a) LC-CDR1 comprising the amino acid sequence of SEQ ID NO: 64; (b) LC-CDR2 comprising the amino acid sequence of SEQ ID NO: 72; and (c) LC-CDR3 comprising the
10 amino acid sequence of SEQ ID NO: 80.

[0325] In one aspect, an antibody or antigen-binding fragment thereof is provided, wherein the antibody or antigen-binding fragment thereof comprises a VH as in any of the embodiments provided above, and a VL as in any of the embodiments provided above. In some embodiments, the antibody comprises a VH comprising the amino acid sequence of SEQ ID NO: 7, and a VL sequence in SEQ ID NO: 8, including post-translational
15 modifications of those sequences. In some embodiments, the antibody or antigen binding fragment thereof comprises an IgA heavy chain constant region having at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% sequence identity to the amino acid sequence of any one of SEQ ID NOS: 16-21. In some embodiments, the antibody or antigen binding fragment thereof comprises a kappa light chain constant region of SEQ ID NO: 31.

20 *Her2*

[0326] In some embodiments, provided herein is an antibody or a functional fragment thereof that specifically binds Her2. In some embodiments, the IgA antibody comprises at least CDR3 of the heavy chain of antibody Trastuzumab. In some embodiments, the IgA antibody comprises one, two, or three of CDR1, CDR2, or CDR3 of the heavy chain of antibody Trastuzumab. In some embodiments, the IgA antibody
25 comprises one, two, or three of CDR1, CDR2, or CDR3 of the light chain of antibody Trastuzumab. In some embodiments, the IgA antibody comprises one, two, or three of CDR1, CDR2, or CDR3 of the heavy chain of antibody Trastuzumab, and one, two, or three of CDR1, CDR2, or CDR3 of the light chain of antibody Trastuzumab. In some embodiments, the IgA antibody comprises CDR1, CDR2, or CDR3 of the heavy chain of antibody Trastuzumab, and CDR1, CDR2, or CDR3 of the light chain of antibody Trastuzumab. In some
30 embodiments, the IgA antibody comprises at least CDR3 of the heavy chain of antibody Trastuzumab and an IgA hinge. In some embodiments, the IgA antibody comprises at least CDR3 of the heavy chain of antibody Trastuzumab an IgA hinge, a CH1 IgA region. In some embodiments, the IgA antibody comprises at least CDR3 of the heavy chain of antibody Trastuzumab an IgA hinge, a CH1 IgA region, and a CH2 IgA region.

In some embodiments, the IgA antibody comprises at least CDR3 of the heavy chain of antibody Trastuzumab, an IgA hinge, a CH1 IgA region, a CH2 IgA region, and a CH3 IgA region.

[0327] In one aspect, the present disclosure provides an antibody or antigen-binding fragment thereof comprising one or more variable regions selected from the group consisting of (a) VH comprising the amino acid sequence of SEQ ID NO: 82, (b) VL comprising the amino acid sequence of SEQ ID NO: 96, and (c) a combination thereof.

[0328] In one aspect, the present disclosure provides an antibody or antigen-binding fragment thereof comprising at least one, two, three, four, five, or six CDRs selected from (a) HC-CDR1 comprising the amino acid sequence of SEQ ID NO: 35; (b) HC-CDR2 comprising the amino acid sequence of SEQ ID NO: 43; (c) HC-CDR3 comprising the amino acid sequence of SEQ ID NO: 51; (d) LC-CDR1 comprising the amino acid sequence of SEQ ID NO: 59; (e) LC-CDR2 comprising the amino acid sequence of SEQ ID NO: 67; and (f) LC-CDR3 comprising the amino acid sequence of SEQ ID NO: 75.

[0329] In one aspect, the present disclosure provides an antibody or antigen-binding fragment thereof comprising at least one, at least two, or all three VH CDR sequences selected from (a) HC-CDR1 comprising the amino acid sequence of SEQ ID NO: 35; (b) HC-CDR2 comprising the amino acid sequence of SEQ ID NO: 43; and (c) HC-CDR3 comprising the amino acid sequence of SEQ ID NO: 51; and (d) a VL comprising the amino acid sequence of SEQ ID NO: 96.

[0330] In one aspect, the present disclosure provides an antibody or antigen-binding fragment thereof comprising at least one, at least two, or all three VL CDR sequences selected from (a) LC-CDR1 comprising the amino acid sequence of SEQ ID NO: 59; (b) LC-CDR2 comprising the amino acid sequence of SEQ ID NO: 67; and (c) LC-CDR3 comprising the amino acid sequence of SEQ ID NO: 75; and a VH comprising the amino acid sequence of SEQ ID NO: 82.

[0331] In one aspect, the present disclosure provides an antibody or antigen-binding fragment thereof comprising the CDRs: HC-CDR3 comprising the amino acid sequence of SEQ ID NO: 51; and LC-CDR3 comprising the amino acid sequence of SEQ ID NO: 75.

[0332] In one aspect, the disclosure herein provides an antibody or antigen-binding fragment thereof comprising at least one, at least two, or all three VL CDR sequences selected from (a) LC-CDR1 comprising the amino acid sequence of SEQ ID NO: 59; (b) LC-CDR2 comprising the amino acid sequence of SEQ ID NO: 67 and (c) LC-CDR3 comprising the amino acid sequence of SEQ ID NO: 75. In one aspect, the disclosure provides an antibody or antigen-binding fragment thereof comprising at least one, at least two, or all three VH CDR sequences selected from (a) HC-CDR1 comprising the amino acid sequence of SEQ ID NO: 35; (b) HC-CDR2 comprising the amino acid sequence of SEQ ID NO: 43 and (c) HC-CDR3 comprising the amino acid sequence of SEQ ID NO: 51.

[0333] In one aspect, the disclosure provides an antibody or antigen-binding fragment thereof comprising the CDRs: (a) HC-CDR1 comprising the amino acid sequence of SEQ ID NO: 35; (b) HC-CDR2 comprising the amino acid sequence of SEQ ID NO: 43; (c) HC-CDR3 comprising the amino acid sequence of SEQ ID NO: 51; (d) LC-CDR1 comprising the amino acid sequence of SEQ ID NO: 59; (e) LC-CDR2 comprising the amino acid sequence of SEQ ID NO: 67; and (f) LC-CDR3 comprising the amino acid sequence of SEQ ID NO: 75.

[0334] In one aspect, an antibody or antigen-binding fragment thereof comprises a VH sequence having at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% sequence identity to the amino acid sequence of SEQ ID NO: 82. In some embodiments, a VH sequence having at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% identity contains substitutions (e.g., conservative substitutions), insertions, or deletions relative to the reference sequence, but an antibody or antigen-binding fragment thereof comprising that sequence retains the ability to bind to antigen. In some embodiments, a total of 1 to 10 amino acids have been substituted, inserted and/or deleted in the amino acid sequence of SEQ ID NO: 82. In some embodiments, substitutions, insertions, or deletions occur in regions outside the CDRs (e.g., in the FRs).
Optionally, the antibody or antigen-binding fragment thereof comprises the VH sequence of the amino acid sequence of SEQ ID NO: 82, including post-translational modifications of that sequence. In a particular embodiment, the VH comprises one, two or three CDRs selected from: (a) HC-CDR1 comprising the amino acid sequence of SEQ ID NO: 35, (b) HC-CDR2 comprising the amino acid sequence of SEQ ID NO: 43, and (c) HC-CDR3 comprising the amino acid sequence of SEQ ID NO: 51.

[0335] In one aspect, an antibody or antigen-binding fragment thereof is provided, wherein the antibody or antigen-binding fragment thereof comprises a light chain variable domain (VL) having at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% sequence identity to the amino acid sequence of SEQ ID NO: 96. In some embodiments, a VL sequence having at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% identity contains substitutions (e.g., conservative substitutions), insertions, or deletions relative to the reference sequence, but an antibody or antigen-binding fragment thereof comprising that sequence retains the ability to bind to antigen. In some embodiments, a total of 1 to 10 amino acids have been substituted, inserted and/or deleted in any one of the amino acid sequence of SEQ ID NO: 96. In some embodiments, the substitutions, insertions, or deletions occur in regions outside the CDRs (e.g., in the FRs).
Optionally, the antibody or antigen-binding fragment thereof comprises the VL sequence of SEQ ID NO: 96, including post-translational modifications of that sequence. In a particular embodiment, the VL comprises one, two or three CDRs selected from (a) LC-CDR1 comprising the amino acid sequence of SEQ ID NO: 59; (b) LC-CDR2 comprising the amino acid sequence of SEQ ID NO: 67; and (c) LC-CDR3 comprising the amino acid sequence of SEQ ID NO: 75.

[0336] In one aspect, an antibody or antigen-binding fragment thereof is provided, wherein the antibody or antigen-binding fragment thereof comprises a VH as in any of the embodiments provided above, and a VL as in any of the embodiments provided above. In some embodiments, the antibody comprises a VH comprising the amino acid sequence of SEQ ID NO: 82, and a VL sequence in SEQ ID NO: 96, including post-translational modifications of those sequences. In some embodiments, the antibody or antigen binding fragment thereof comprises an IgA heavy chain constant region having at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% sequence identity to the amino acid sequence of any one of SEQ ID NOs: 16-21. In some embodiments, the antibody or antigen binding fragment thereof comprises a kappa light chain constant region of SEQ ID NO: 31.

gp75

[0337] In some embodiments, provided herein is an antibody or a functional fragment thereof that specifically binds gp75 or tyrosine related protein 1. In some embodiments, an antibody or a functional fragment disclosed herein specifically binds a human gp75 polypeptide. The polypeptide and coding nucleic acid sequences of gp75 of human origin and those of a number of animals are publicly available, e.g., from the NCBI website. In some embodiments, the IgA antibody comprises at least CDR3 of the heavy chain of antibody TA99. In some embodiments, the IgA antibody comprises one, two, or three of CDR1, CDR2, or CDR3 of the heavy chain of antibody TA99. In some embodiments, the IgA antibody comprises one, two, or three of CDR1, CDR2, or CDR3 of the light chain of antibody TA99. In some embodiments, the IgA antibody comprises one, two, or three of CDR1, CDR2, or CDR3 of the heavy chain of antibody TA99, and one, two, or three of CDR1, CDR2, or CDR3 of the light chain of antibody TA99. In some embodiments, the IgA antibody comprises CDR1, CDR2, or CDR3 of the heavy chain of antibody TA99, and CDR1, CDR2, or CDR3 of the light chain of antibody TA99. In some embodiments, the IgA antibody comprises at least CDR3 of the heavy chain of antibody TA99 and an IgA hinge. In some embodiments, the IgA antibody comprises at least CDR3 of the heavy chain of antibody TA99 an IgA hinge, a CH1 IgA region. In some embodiments, the IgA antibody comprises at least CDR3 of the heavy chain of antibody TA99 an IgA hinge, a CH1 IgA region, and a CH2 IgA region. In some embodiments, the IgA antibody comprises at least CDR3 of the heavy chain of antibody TA99 an IgA hinge, a CH1 IgA region, a CH2 IgA region, and a CH3 IgA region.

[0338] In one aspect, the present disclosure provides an antibody or antigen-binding fragment thereof comprises one or more variable regions selected from the group consisting of (a) VH comprising the amino acid sequence of SEQ ID NO: 83, (b) VL comprising the amino acid sequence of SEQ ID NO: 97, and (c) a combination thereof.

[0339] In one aspect, the present disclosure provides an antibody or antigen-binding fragment thereof comprising at least one, two, three, four, five, or six CDRs selected from (a) HC-CDR1 comprising the amino

acid sequence of SEQ ID NO: 36; (b) HC-CDR2 comprising the amino acid sequence of SEQ ID NO: 44; (c) HC-CDR3 comprising the amino acid sequence of SEQ ID NO: 52; (d) LC-CDR1 comprising the amino acid sequence of SEQ ID NO: 60; (e) LC-CDR2 comprising the amino acid sequence of SEQ ID NO: 68; and (f) LC-CDR3 comprising the amino acid sequence of SEQ ID NO: 76.

- 5 **[0340]** In one aspect, the present disclosure provides an antibody or antigen-binding fragment thereof comprising at least one, at least two, or all three VH CDR sequences selected from (a) HC-CDR1 comprising the amino acid sequence of SEQ ID NO: 36; (b) HC-CDR2 comprising the amino acid sequence of SEQ ID NO: 44; and (c) HC-CDR3 comprising the amino acid sequence of SEQ ID NO: 52; and (d) a VL comprising the amino acid sequence of SEQ ID NO: 97.
- 10 **[0341]** In one aspect, the present disclosure provides an antibody or antigen-binding fragment thereof comprising at least one, at least two, or all three VL CDR sequences selected from (a) LC-CDR1 comprising the amino acid sequence of SEQ ID NO: 60; (b) LC-CDR2 comprising the amino acid sequence of SEQ ID NO: 68; and (c) LC-CDR3 comprising the amino acid sequence of SEQ ID NO: 76; and a VH comprising the amino acid sequence of SEQ ID NO: 83.
- 15 **[0342]** In one aspect, the present disclosure provides an antibody or antigen-binding fragment thereof comprising the CDRs: HC-CDR3 comprising the amino acid sequence of SEQ ID NO: 52; and LC-CDR3 comprising the amino acid sequence of SEQ ID NO: 76.
- 20 **[0343]** In one aspect, the disclosure herein provides an antibody or antigen-binding fragment thereof comprising at least one, at least two, or all three VL CDR sequences selected from (a) LC-CDR1 comprising the amino acid sequence of SEQ ID NO: 60; (b) LC-CDR2 comprising the amino acid sequence of SEQ ID NO: 68 and (c) LC-CDR3 comprising the amino acid sequence of SEQ ID NO: 76. In one aspect, the disclosure provides an antibody or antigen-binding fragment thereof comprising at least one, at least two, or all three VH CDR sequences selected from (a) HC-CDR1 comprising the amino acid sequence of SEQ ID NO: 36; (b) HC-CDR2 comprising the amino acid sequence of SEQ ID NO: 44 and (c) HC-CDR3 comprising the amino acid sequence of SEQ ID NO: 52.
- 25 **[0344]** In one aspect, the disclosure provides an antibody or antigen-binding fragment thereof comprising the CDRs: (a) HC-CDR1 comprising the amino acid sequence of SEQ ID NO: 36; (b) HC-CDR2 comprising the amino acid sequence of SEQ ID NO: 44; (c) HC-CDR3 comprising the amino acid sequence of SEQ ID NO: 52; (d) LC-CDR1 comprising the amino acid sequence of SEQ ID NO: 60; (e) LC-CDR2 comprising the amino acid sequence of SEQ ID NO: 68; and (f) LC-CDR3 comprising the amino acid sequence of SEQ ID NO: 76.
- 30 **[0345]** In one aspect, an antibody or antigen-binding fragment thereof comprises a VH sequence having at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% sequence identity to the amino acid sequence of SEQ ID NO: 83. In some embodiments, a VH sequence having at least 90%, 91%, 92%, 93%,

- 94%, 95%, 96%, 97%, 98%, or 99% identity contains substitutions (e.g., conservative substitutions), insertions, or deletions relative to the reference sequence, but an antibody or antigen-binding fragment thereof comprising that sequence retains the ability to bind to antigen. In some embodiments, a total of 1 to 10 amino acids have been substituted, inserted and/or deleted in the amino acid sequence of SEQ ID NO: 83. In some
- 5 embodiments, substitutions, insertions, or deletions occur in regions outside the CDRs (e.g., in the FRs). Optionally, the antibody or antigen-binding fragment thereof comprises the VH sequence of the amino acid sequence of SEQ ID NO: 83, including post-translational modifications of that sequence. In a particular embodiment, the VH comprises one, two or three CDRs selected from: (a) HC-CDR1 comprising the amino acid sequence of SEQ ID NO: 36, (b) HC-CDR2 comprising the amino acid sequence of SEQ ID NO: 44, and
- 10 (c) HC-CDR3 comprising the amino acid sequence of SEQ ID NO: 52.
- [0346]** In one aspect, an antibody or antigen-binding fragment thereof is provided, wherein the antibody or antigen-binding fragment thereof comprises a light chain variable domain (VL) having at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% sequence identity to the amino acid sequence of SEQ ID NO: 97. In some embodiments, a VL sequence having at least 90%, 91%, 92%, 93%, 94%, 95%, 96%,
- 15 97%, 98%, or 99% identity contains substitutions (e.g., conservative substitutions), insertions, or deletions relative to the reference sequence, but an antibody or antigen-binding fragment thereof comprising that sequence retains the ability to bind to antigen. In some embodiments, a total of 1 to 10 amino acids have been substituted, inserted and/or deleted in any one of the amino acid sequence of SEQ ID NO: 97. In some embodiments, the substitutions, insertions, or deletions occur in regions outside the CDRs (e.g., in the FRs).
- 20 Optionally, the antibody or antigen-binding fragment thereof comprises the VL sequence of SEQ ID NO: 97, including post-translational modifications of that sequence. In a particular embodiment, the VL comprises one, two or three CDRs selected from (a) LC-CDR1 comprising the amino acid sequence of SEQ ID NO: 60; (b) LC-CDR2 comprising the amino acid sequence of SEQ ID NO: 68; and (c) LC-CDR3 comprising the amino acid sequence of SEQ ID NO: 76.
- 25 **[0347]** In one aspect, an antibody or antigen-binding fragment thereof is provided, wherein the antibody or antigen-binding fragment thereof comprises a VH as in any of the embodiments provided above, and a VL as in any of the embodiments provided above. In some embodiments, the antibody comprises a VH comprising the amino acid sequence of SEQ ID NO: 83, and a VL sequence in SEQ ID NO: 97, including post-translational modifications of those sequences. In some embodiments, the antibody or antigen binding
- 30 fragment thereof comprises an IgA heavy chain constant region having at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% sequence identity to the amino acid sequence of any one of SEQ ID NOs: 16-21. In some embodiments, the antibody or antigen binding fragment thereof comprises a kappa light chain constant region of SEQ ID NO: 31.

CTLA4

[0348] In some embodiments, provided herein is an antibody or a functional fragment thereof that specifically binds cytotoxic T-lymphocyte-associated protein 4 (CTLA4). In some embodiments, an antibody or a functional fragment disclosed herein specifically binds a human CTLA4 polypeptide. In some embodiments, an antibody or a functional fragment disclosed herein specifically binds a mouse CTLA4 polypeptide. The polypeptide and coding nucleic acid sequences of CTLA4 of human origin and those of a number of animals are publicly available, e.g., from the NCBI website.

[0349] In one aspect, the present disclosure provides an antibody or antigen-binding fragment thereof comprising one or more variable regions selected from the group consisting of (a) VH comprising the amino acid sequence of SEQ ID NO: 84, (b) VL comprising the amino acid sequence of SEQ ID NO: 98, and (c) a combination thereof.

[0350] In one aspect, the present disclosure provides an antibody or antigen-binding fragment thereof comprising at least one, two, three, four, five, or six CDRs selected from (a) HC-CDR1 comprising the amino acid sequence of SEQ ID NO: 37; (b) HC-CDR2 comprising the amino acid sequence of SEQ ID NO: 45; (c) HC-CDR3 comprising the amino acid sequence of SEQ ID NO: 53; (d) LC-CDR1 comprising the amino acid sequence of SEQ ID NO: 61; (e) LC-CDR2 comprising the amino acid sequence of SEQ ID NO: 69; and (f) LC-CDR3 comprising the amino acid sequence of SEQ ID NO: 77.

[0351] In one aspect, the present disclosure provides an antibody or antigen-binding fragment thereof comprising at least one, at least two, or all three VH CDR sequences selected from (a) HC-CDR1 comprising the amino acid sequence of SEQ ID NO: 37; (b) HC-CDR2 comprising the amino acid sequence of SEQ ID NO: 45; and (c) HC-CDR3 comprising the amino acid sequence of SEQ ID NO: 53; and (d) a VL comprising the amino acid sequence of SEQ ID NO: 98.

[0352] In one aspect, the present disclosure provides an antibody or antigen-binding fragment thereof comprising at least one, at least two, or all three VL CDR sequences selected from (a) LC-CDR1 comprising the amino acid sequence of SEQ ID NO: 61; (b) LC-CDR2 comprising the amino acid sequence of SEQ ID NO: 69; and (c) LC-CDR3 comprising the amino acid sequence of SEQ ID NO: 77; and a VH comprising the amino acid sequence of SEQ ID NO: 84.

[0353] In one aspect, the present disclosure provides an antibody or antigen-binding fragment thereof comprising the CDRs: HC-CDR3 comprising the amino acid sequence of SEQ ID NO: 53; and LC-CDR3 comprising the amino acid sequence of SEQ ID NO: 77.

[0354] In one aspect, the disclosure herein provides an antibody or antigen-binding fragment thereof comprising at least one, at least two, or all three VL CDR sequences selected from (a) LC-CDR1 comprising the amino acid sequence of SEQ ID NO: 61; (b) LC-CDR2 comprising the amino acid sequence of SEQ ID NO: 69 and (c) LC-CDR3 comprising the amino acid sequence of SEQ ID NO: 77. In one aspect, the disclosure provides an antibody or antigen-binding fragment thereof comprising at least one, at least two, or

all three VH CDR sequences selected from (a) HC-CDR1 comprising the amino acid sequence of SEQ ID NO: 37; (b) HC-CDR2 comprising the amino acid sequence of SEQ ID NO: 45 and (c) HC-CDR3 comprising the amino acid sequence of SEQ ID NO: 53.

[0355] In one aspect, the disclosure provides an antibody or antigen-binding fragment thereof comprising the CDRs: (a) HC-CDR1 comprising the amino acid sequence of SEQ ID NO: 37; (b) HC-CDR2 comprising the amino acid sequence of SEQ ID NO: 45; (c) HC-CDR3 comprising the amino acid sequence of SEQ ID NO: 53; (d) LC-CDR1 comprising the amino acid sequence of SEQ ID NO: 61; (e) LC-CDR2 comprising the amino acid sequence of SEQ ID NO: 69; and (f) LC-CDR3 comprising the amino acid sequence of SEQ ID NO: 77.

[0356] In one aspect, an antibody or antigen-binding fragment thereof comprises a VH sequence having at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% sequence identity to the amino acid sequence of SEQ ID NO: 84. In some embodiments, a VH sequence having at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% identity contains substitutions (e.g., conservative substitutions), insertions, or deletions relative to the reference sequence, but an antibody or antigen-binding fragment thereof comprising that sequence retains the ability to bind to antigen. In some embodiments, a total of 1 to 10 amino acids have been substituted, inserted and/or deleted in the amino acid sequence of SEQ ID NO: 84. In some embodiments, substitutions, insertions, or deletions occur in regions outside the CDRs (e.g., in the FRs). Optionally, the antibody or antigen-binding fragment thereof comprises the VH sequence of the amino acid sequence of SEQ ID NO: 84, including post-translational modifications of that sequence. In a particular embodiment, the VH comprises one, two or three CDRs selected from: (a) HC-CDR1 comprising the amino acid sequence of SEQ ID NO: 37, (b) HC-CDR2 comprising the amino acid sequence of SEQ ID NO: 45, and (c) HC-CDR3 comprising the amino acid sequence of SEQ ID NO: 53.

[0357] In one aspect, an antibody or antigen-binding fragment thereof is provided, wherein the antibody or antigen-binding fragment thereof comprises a light chain variable domain (VL) having at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% sequence identity to the amino acid sequence of SEQ ID NO: 98. In some embodiments, a VL sequence having at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% identity contains substitutions (e.g., conservative substitutions), insertions, or deletions relative to the reference sequence, but an antibody or antigen-binding fragment thereof comprising that sequence retains the ability to bind to antigen. In some embodiments, a total of 1 to 10 amino acids have been substituted, inserted and/or deleted in any one of the amino acid sequence of SEQ ID NO: 98. In some embodiments, the substitutions, insertions, or deletions occur in regions outside the CDRs (e.g., in the FRs). Optionally, the antibody or antigen-binding fragment thereof comprises the VL sequence of SEQ ID NO: 98, including post-translational modifications of that sequence. In a particular embodiment, the VL comprises one, two or three CDRs selected from (a) LC-CDR1 comprising the amino acid sequence of SEQ ID NO: 61;

(b) LC-CDR2 comprising the amino acid sequence of SEQ ID NO: 69; and (c) LC-CDR3 comprising the amino acid sequence of SEQ ID NO: 77.

[0358] In one aspect, an antibody or antigen-binding fragment thereof is provided, wherein the antibody or antigen-binding fragment thereof comprises a VH as in any of the embodiments provided above, and a VL as in any of the embodiments provided above. In some embodiments, the antibody comprises a VH comprising the amino acid sequence of SEQ ID NO: 84, and a VL sequence in SEQ ID NO: 98, including post-translational modifications of those sequences. In some embodiments, the antibody or antigen binding fragment thereof comprises an IgA heavy chain constant region having at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% sequence identity to the amino acid sequence of any one of SEQ ID NOs: 16-21. In some embodiments, the antibody or antigen binding fragment thereof comprises a kappa light chain constant region of SEQ ID NO: 31.

CD47

[0359] In some embodiments, the IgA antibody specifically binds CD47. In some embodiments, an antibody or a functional fragment disclosed herein specifically binds a human CD47 polypeptide. The polypeptide and coding nucleic acid sequences of CD47 of human origin and those of a number of animals are publicly available, e.g., from the NCBI website. In some embodiments, the IgA antibody reduces SIRP α binding to CD47 expressed on the surface of a cancer cell. In some embodiments, the IgA antibody binds to CD47 with a binding affinity (Kd) ranging from about 0.5 μ M to about 999 μ M as compared to a corresponding wild type antibody. In some embodiments, the IgA antibody binds to CD47 with a binding affinity (Kd) ranging from about 1 mM to about 1000 mM as compared to a corresponding wild type antibody. In some embodiments, the IgA antibody inhibits a human CD47 interaction with a signal regulatory protein α (SIRP α). In some embodiments, the inhibition of interaction between said human CD47 and said SIRP α increases a potential of said IgA antibody. In some embodiments, the inhibition of the interaction between said human CD47 and said SIRP α increases phagocytosis and clearance of cancer cells at a tumor site. In some embodiments, the cancer cells are IgA-opsonized cancer cells. In some embodiments, the IgA antibody comprises an antigen binding domain that binds CD47 and an antigen binding domain that specifically binds a tumor associated antigen (e.g., one described herein).

[0360] In some embodiments, the IgA antibody binds CD47. In some embodiments, the IgA antibody reduces CD47 binding of a cancer cell. For example, the IgA antibody can inhibit a human CD47 interaction with a signal regulatory protein α (SIRP α). Furthermore, the inhibition of interaction between the human CD47 and the SIRP α can increase a potential of the IgA antibody. The inhibition of interaction between the human CD47 and SIRP α can increase phagocytosis and clearance of cancer cells at a tumor site. For example, the cancer cells can be IgA-opsonized cancer cells. In some embodiments, the IgA

antibodies described herein have a low affinity binding to CD47 that prevents the binding of the IgA antibody to CD47 on a cell other than a cancer cell. In some embodiments, the low affinity CD47 arm of the IgA antibody described herein binds to a tumor cell expressing CD47. In some examples, the low affinity CD47 arm of the IgA antibody described herein does not bind to a cell expressing CD47 that is not a tumor cell. In some embodiments, the IgA antibody binds to CD47 with a binding affinity (K_d) of at least about 0.01 micromolar (μM) to about 999 μM or more. In some embodiments, the IgA antibody binds to CD47 with a binding affinity (K_d) of at least about 0.01 μM . In some embodiments, the IgA antibody binds to CD47 with a binding affinity (K_d) of at most about 999 μM . In some embodiments, the IgA antibody binds to CD47 with a binding affinity (K_d) of about 0.01 μM to about 0.1 μM , about 0.01 μM to about 0.5 μM , about 0.01 μM to about 1 μM , about 0.01 μM to about 5 μM , about 0.01 μM to about 10 μM , about 0.01 μM to about 50 μM , about 0.01 μM to about 100 μM , about 0.01 μM to about 200 μM , about 0.01 μM to about 300 μM , about 0.01 μM to about 500 μM , about 0.01 μM to about 999 μM , about 0.1 μM to about 0.5 μM , about 0.1 μM to about 1 μM , about 0.1 μM to about 5 μM , about 0.1 μM to about 10 μM , about 0.1 μM to about 50 μM , about 0.1 μM to about 100 μM , about 0.1 μM to about 200 μM , about 0.1 μM to about 300 μM , about 0.1 μM to about 500 μM , about 0.1 μM to about 999 μM , about 0.5 μM to about 1 μM , about 0.5 μM to about 5 μM , about 0.5 μM to about 10 μM , about 0.5 μM to about 50 μM , about 0.5 μM to about 100 μM , about 0.5 μM to about 200 μM , about 0.5 μM to about 300 μM , about 0.5 μM to about 500 μM , about 0.5 μM to about 999 μM , about 1 μM to about 5 μM , about 1 μM to about 10 μM , about 1 μM to about 50 μM , about 1 μM to about 100 μM , about 1 μM to about 200 μM , about 1 μM to about 300 μM , about 1 μM to about 500 μM , about 1 μM to about 999 μM , about 5 μM to about 10 μM , about 5 μM to about 50 μM , about 5 μM to about 100 μM , about 5 μM to about 200 μM , about 5 μM to about 300 μM , about 5 μM to about 500 μM , about 5 μM to about 999 μM , about 10 μM to about 50 μM , about 10 μM to about 100 μM , about 10 μM to about 200 μM , about 10 μM to about 300 μM , about 10 μM to about 500 μM , about 10 μM to about 999 μM , about 50 μM to about 100 μM , about 50 μM to about 200 μM , about 50 μM to about 300 μM , about 50 μM to about 500 μM , about 50 μM to about 999 μM , about 100 μM to about 200 μM , about 100 μM to about 300 μM , about 100 μM to about 500 μM , about 100 μM to about 999 μM , about 200 μM to about 300 μM , about 200 μM to about 500 μM , about 200 μM to about 999 μM , about 300 μM to about 500 μM , about 300 μM to about 999 μM , or about 500 μM to about 999 μM . In some embodiments, the IgA antibody binds to CD47 with a binding affinity (K_d) of about 0.01 μM , about 0.1 μM , about 0.5 μM , about 1 μM , about 5 μM , about 10 μM , about 50 μM , about 100 μM , about 200 μM , about 300 μM , about 500 μM , or about 999 μM .

[0361] In some embodiments, the IgA antibody binds to CD47 with a binding affinity (K_d) of about 1 mM to about 1,000 millimolar (mM). In some embodiments, the IgA antibody binds to CD47 with a binding affinity (K_d) of at least about 1 mM. In some embodiments, the IgA antibody binds to CD47 with a binding

affinity (K_d) of at most about 1,000 mM. In some embodiments, the IgA antibody binds to CD47 with a binding affinity (K_d) of about 1 mM to about 5 mM, about 1 mM to about 10 mM, about 1 mM to about 50 mM, about 1 mM to about 100 mM, about 1 mM to about 200 mM, about 1 mM to about 300 mM, about 1 mM to about 400 mM, about 1 mM to about 500 mM, about 1 mM to about 600 mM, about 1 mM to about 800 mM, about 1 mM to about 1,000 mM, about 5 mM to about 10 mM, about 5 mM to about 50 mM, about 5 mM to about 100 mM, about 5 mM to about 200 mM, about 5 mM to about 300 mM, about 5 mM to about 400 mM, about 5 mM to about 500 mM, about 5 mM to about 600 mM, about 5 mM to about 800 mM, about 5 mM to about 1,000 mM, about 10 mM to about 50 mM, about 10 mM to about 100 mM, about 10 mM to about 200 mM, about 10 mM to about 300 mM, about 10 mM to about 400 mM, about 10 mM to about 500 mM, about 10 mM to about 600 mM, about 10 mM to about 800 mM, about 10 mM to about 1,000 mM, about 50 mM to about 100 mM, about 50 mM to about 200 mM, about 50 mM to about 300 mM, about 50 mM to about 400 mM, about 50 mM to about 500 mM, about 50 mM to about 600 mM, about 50 mM to about 800 mM, about 50 mM to about 1,000 mM, about 100 mM to about 200 mM, about 100 mM to about 300 mM, about 100 mM to about 400 mM, about 100 mM to about 500 mM, about 100 mM to about 600 mM, about 100 mM to about 800 mM, about 100 mM to about 1,000 mM, about 200 mM to about 300 mM, about 200 mM to about 400 mM, about 200 mM to about 500 mM, about 200 mM to about 600 mM, about 200 mM to about 800 mM, about 200 mM to about 1,000 mM, about 300 mM to about 400 mM, about 300 mM to about 500 mM, about 300 mM to about 600 mM, about 300 mM to about 800 mM, about 300 mM to about 1,000 mM, about 400 mM to about 500 mM, about 400 mM to about 600 mM, about 400 mM to about 800 mM, about 400 mM to about 1,000 mM, about 500 mM to about 600 mM, about 500 mM to about 800 mM, about 500 mM to about 1,000 mM, about 600 mM to about 800 mM, about 600 mM to about 1,000 mM, or about 800 mM to about 1,000 mM. In some embodiments, the IgA antibody binds to CD47 with a binding affinity (K_d) of about 1 mM, about 5 mM, about 10 mM, about 50 mM, about 100 mM, about 200 mM, about 300 mM, about 400 mM, about 500 mM, about 600 mM, about 800 mM, or about 1,000 mM.

[0362] In some embodiments, provided herein is an antibody or a functional fragment thereof that specifically binds CD47. In some embodiments, the IgA antibody comprises at least CDR3 of the heavy chain of antibody 2.3D11 or of antibody C47A8-CQ. In some embodiments, the IgA antibody comprises one, two, or three of CDR1, CDR2, or CDR3 of the heavy chain of antibody 2.3D11 or of antibody C47A8-CQ. In some embodiments, the IgA antibody comprises one, two, or three of CDR1, CDR2, or CDR3 of the light chain of antibody 2.3D11 or of antibody C47A8-CQ. In some embodiments, the IgA antibody comprises one, two, or three of CDR1, CDR2, or CDR3 of the heavy chain of antibody 2.3D11 or of antibody C47A8-CQ, and one, two, or three of CDR1, CDR2, or CDR3 of the light chain of antibody 2.3D11 or of antibody C47A8-CQ. In some embodiments, the IgA antibody comprises CDR1, CDR2, or CDR3 of the heavy chain

of antibody 2.3D11 or of antibody C47A8-CQ, and CDR1, CDR2, or CDR3 of the light chain of antibody 2.3D11 or of antibody C47A8-CQ. In some embodiments, the IgA antibody comprises at least CDR3 of the heavy chain of antibody 2.3D11 or of antibody C47A8-CQ and an IgA hinge. In some embodiments, the IgA antibody comprises at least CDR3 of the heavy chain of antibody 2.3D11 or of antibody C47A8-CQ an IgA hinge, a CH1 IgA region. In some embodiments, the IgA antibody comprises at least CDR3 of the heavy chain of antibody 2.3D11 or of antibody C47A8-CQ an IgA hinge, a CH1 IgA region, and a CH2 IgA region. In some embodiments, the IgA antibody comprises at least CDR3 of the heavy chain of antibody 2.3D11 or of antibody C47A8-CQ an IgA hinge, a CH1 IgA region, a CH2 IgA region, and a CH3 IgA region.

[0363] In one aspect, the present disclosure provides an antibody or antigen-binding fragment thereof comprising one or more variable regions selected from the group consisting of (a) VH comprising the amino acid sequence of SEQ ID NO: 85, (b) VL comprising the amino acid sequence of SEQ ID NO: 99, and (c) a combination thereof.

[0364] In one aspect, the present disclosure provides an antibody or antigen-binding fragment thereof comprising at least one, two, three, four, five, or six CDRs selected from (a) HC-CDR1 comprising the amino acid sequence of SEQ ID NO: 38; (b) HC-CDR2 comprising the amino acid sequence of SEQ ID NO: 46; (c) HC-CDR3 comprising the amino acid sequence of SEQ ID NO: 54; (d) LC-CDR1 comprising the amino acid sequence of SEQ ID NO: 62; (e) LC-CDR2 comprising the amino acid sequence of SEQ ID NO: 70; and (f) LC-CDR3 comprising the amino acid sequence of SEQ ID NO: 78.

[0365] In one aspect, the present disclosure provides an antibody or antigen-binding fragment thereof comprising at least one, at least two, or all three VH CDR sequences selected from (a) HC-CDR1 comprising the amino acid sequence of SEQ ID NO: 38; (b) HC-CDR2 comprising the amino acid sequence of SEQ ID NO: 46; and (c) HC-CDR3 comprising the amino acid sequence of SEQ ID NO: 54; and (d) a VL comprising the amino acid sequence of SEQ ID NO: 99.

[0366] In one aspect, the present disclosure provides an antibody or antigen-binding fragment thereof comprising at least one, at least two, or all three VL CDR sequences selected from (a) LC-CDR1 comprising the amino acid sequence of SEQ ID NO: 62; (b) LC-CDR2 comprising the amino acid sequence of SEQ ID NO: 70; and (c) LC-CDR3 comprising the amino acid sequence of SEQ ID NO: 78; and a VH comprising the amino acid sequence of SEQ ID NO: 85.

[0367] In one aspect, the present disclosure provides an antibody or antigen-binding fragment thereof comprising the CDRs: HC-CDR3 comprising the amino acid sequence of SEQ ID NO: 54; and LC-CDR3 comprising the amino acid sequence of SEQ ID NO: 78.

[0368] In one aspect, the disclosure herein provides an antibody or antigen-binding fragment thereof comprising at least one, at least two, or all three VL CDR sequences selected from (a) LC-CDR1 comprising the amino acid sequence of SEQ ID NO: 62; (b) LC-CDR2 comprising the amino acid sequence of SEQ ID

NO: 70 and (c) LC-CDR3 comprising the amino acid sequence of SEQ ID NO: 78. In one aspect, the disclosure provides an antibody or antigen-binding fragment thereof comprising at least one, at least two, or all three VH CDR sequences selected from (a) HC-CDR1 comprising the amino acid sequence of SEQ ID NO: 38; (b) HC-CDR2 comprising the amino acid sequence of SEQ ID NO: 46 and (c) HC-CDR3 comprising the amino acid sequence of SEQ ID NO: 54.

[0369] In one aspect, the disclosure provides an antibody or antigen-binding fragment thereof comprising the CDRs: (a) HC-CDR1 comprising the amino acid sequence of SEQ ID NO: 38; (b) HC-CDR2 comprising the amino acid sequence of SEQ ID NO: 46; (c) HC-CDR3 comprising the amino acid sequence of SEQ ID NO: 54; (d) LC-CDR1 comprising the amino acid sequence of SEQ ID NO: 62; (e) LC-CDR2 comprising the amino acid sequence of SEQ ID NO: 70; and (f) LC-CDR3 comprising the amino acid sequence of SEQ ID NO: 78.

[0370] In one aspect, an antibody or antigen-binding fragment thereof comprises a VH sequence having at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% sequence identity to the amino acid sequence of SEQ ID NO: 85. In some embodiments, a VH sequence having at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% identity contains substitutions (e.g., conservative substitutions), insertions, or deletions relative to the reference sequence, but an antibody or antigen-binding fragment thereof comprising that sequence retains the ability to bind to antigen. In some embodiments, a total of 1 to 10 amino acids have been substituted, inserted and/or deleted in the amino acid sequence of SEQ ID NO: 85. In some embodiments, substitutions, insertions, or deletions occur in regions outside the CDRs (e.g., in the FRs).

Optionally, the antibody or antigen-binding fragment thereof comprises the VH sequence of the amino acid sequence of SEQ ID NO: 85, including post-translational modifications of that sequence. In a particular embodiment, the VH comprises one, two or three CDRs selected from: (a) HC-CDR1 comprising the amino acid sequence of SEQ ID NO: 38, (b) HC-CDR2 comprising the amino acid sequence of SEQ ID NO: 46, and (c) HC-CDR3 comprising the amino acid sequence of SEQ ID NO: 54.

[0371] In one aspect, an antibody or antigen-binding fragment thereof is provided, wherein the antibody or antigen-binding fragment thereof comprises a light chain variable domain (VL) having at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% sequence identity to the amino acid sequence of SEQ ID NO: 99. In some embodiments, a VL sequence having at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% identity contains substitutions (e.g., conservative substitutions), insertions, or deletions relative to the reference sequence, but an antibody or antigen-binding fragment thereof comprising that sequence retains the ability to bind to antigen. In some embodiments, a total of 1 to 10 amino acids have been substituted, inserted and/or deleted in any one of the amino acid sequence of SEQ ID NO: 99. In some embodiments, the substitutions, insertions, or deletions occur in regions outside the CDRs (e.g., in the FRs). Optionally, the antibody or antigen-binding fragment thereof comprises the VL sequence of SEQ ID NO: 99,

including post-translational modifications of that sequence. In a particular embodiment, the VL comprises one, two or three CDRs selected from (a) LC-CDR1 comprising the amino acid sequence of SEQ ID NO: 62; (b) LC-CDR2 comprising the amino acid sequence of SEQ ID NO: 70; and (c) LC-CDR3 comprising the amino acid sequence of SEQ ID NO: 78.

- 5 **[0372]** In one aspect, an antibody or antigen-binding fragment thereof is provided, wherein the antibody or antigen-binding fragment thereof comprises a VH as in any of the embodiments provided above, and a VL as in any of the embodiments provided above. In some embodiments, the antibody comprises a VH comprising the amino acid sequence of SEQ ID NO: 85, and a VL sequence in SEQ ID NO: 99, including post-translational modifications of those sequences. In some embodiments, the antibody or antigen binding
- 10 fragment thereof comprises an IgA heavy chain constant region having at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% sequence identity to the amino acid sequence of any one of SEQ ID NOs: 16-21. In some embodiments, the antibody or antigen binding fragment thereof comprises a kappa light chain constant region of SEQ ID NO: 31.

- [0373]** In one aspect, the present disclosure provides an antibody or antigen-binding fragment thereof
- 15 comprises one or more variable regions selected from the group consisting of (a) VH comprising the amino acid sequence of SEQ ID NO: 86, (b) VL comprising the amino acid sequence of SEQ ID NO: 100, and (c) a combination thereof.

- [0374]** In one aspect, the present disclosure provides an antibody or antigen-binding fragment thereof comprising at least one, two, three, four, five, or six CDRs selected from (a) HC-CDR1 comprising the amino
- 20 acid sequence of SEQ ID NO: 39; (b) HC-CDR2 comprising the amino acid sequence of SEQ ID NO: 47; (c) HC-CDR3 comprising the amino acid sequence of SEQ ID NO: 54; (d) LC-CDR1 comprising the amino acid sequence of SEQ ID NO: 63; (e) LC-CDR2 comprising the amino acid sequence of SEQ ID NO: 71; and (f) LC-CDR3 comprising the amino acid sequence of SEQ ID NO: 79.

- [0375]** In one aspect, the present disclosure provides an antibody or antigen-binding fragment thereof
- 25 comprising at least one, at least two, or all three VH CDR sequences selected from (a) HC-CDR1 comprising the amino acid sequence of SEQ ID NO: 39; (b) HC-CDR2 comprising the amino acid sequence of SEQ ID NO: 47; and (c) HC-CDR3 comprising the amino acid sequence of SEQ ID NO: 55; and (d) a VL comprising the amino acid sequence of SEQ ID NO: 100.

- [0376]** In one aspect, the present disclosure provides an antibody or antigen-binding fragment thereof
- 30 comprising at least one, at least two, or all three VL CDR sequences selected from (a) LC-CDR1 comprising the amino acid sequence of SEQ ID NO: 63; (b) LC-CDR2 comprising the amino acid sequence of SEQ ID NO: 71; and (c) LC-CDR3 comprising the amino acid sequence of SEQ ID NO: 79; and a VH comprising the amino acid sequence of SEQ ID NO: 86.

[0377] In one aspect, the present disclosure provides an antibody or antigen-binding fragment thereof comprising the CDRs: HC-CDR3 comprising the amino acid sequence of SEQ ID NO: 55; and LC-CDR3 comprising the amino acid sequence of SEQ ID NO: 79.

[0378] In one aspect, the disclosure herein provides an antibody or antigen-binding fragment thereof comprising at least one, at least two, or all three VL CDR sequences selected from (a) LC-CDR1 comprising the amino acid sequence of SEQ ID NO: 63; (b) LC-CDR2 comprising the amino acid sequence of SEQ ID NO: 71 and (c) LC-CDR3 comprising the amino acid sequence of SEQ ID NO: 79. In one aspect, the disclosure provides an antibody or antigen-binding fragment thereof comprising at least one, at least two, or all three VH CDR sequences selected from (a) HC-CDR1 comprising the amino acid sequence of SEQ ID NO: 39; (b) HC-CDR2 comprising the amino acid sequence of SEQ ID NO: 47 and (c) HC-CDR3 comprising the amino acid sequence of SEQ ID NO: 55.

[0379] In one aspect, the disclosure provides an antibody or antigen-binding fragment thereof comprising the CDRs: (a) HC-CDR1 comprising the amino acid sequence of SEQ ID NO: 39; (b) HC-CDR2 comprising the amino acid sequence of SEQ ID NO: 47; (c) HC-CDR3 comprising the amino acid sequence of SEQ ID NO: 55; (d) LC-CDR1 comprising the amino acid sequence of SEQ ID NO: 63; (e) LC-CDR2 comprising the amino acid sequence of SEQ ID NO: 71; and (f) LC-CDR3 comprising the amino acid sequence of SEQ ID NO: 79.

[0380] In one aspect, an antibody or antigen-binding fragment thereof comprises a VH sequence having at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% sequence identity to the amino acid sequence of SEQ ID NO: 86. In some embodiments, a VH sequence having at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% identity contains substitutions (e.g., conservative substitutions), insertions, or deletions relative to the reference sequence, but an antibody or antigen-binding fragment thereof comprising that sequence retains the ability to bind to antigen. In some embodiments, a total of 1 to 10 amino acids have been substituted, inserted and/or deleted in the amino acid sequence of SEQ ID NO: 86. In some embodiments, substitutions, insertions, or deletions occur in regions outside the CDRs (e.g., in the FRs). Optionally, the antibody or antigen-binding fragment thereof comprises the VH sequence of the amino acid sequence of SEQ ID NO: 86, including post-translational modifications of that sequence. In a particular embodiment, the VH comprises one, two or three CDRs selected from: (a) HC-CDR1 comprising the amino acid sequence of SEQ ID NO: 39, (b) HC-CDR2 comprising the amino acid sequence of SEQ ID NO: 47, and (c) HC-CDR3 comprising the amino acid sequence of SEQ ID NO: 55.

[0381] In one aspect, an antibody or antigen-binding fragment thereof is provided, wherein the antibody or antigen-binding fragment thereof comprises a light chain variable domain (VL) having at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% sequence identity to the amino acid sequence of SEQ ID NO: 100. In some embodiments, a VL sequence having at least 90%, 91%, 92%, 93%, 94%, 95%, 96%,

97%, 98%, or 99% identity contains substitutions (e.g., conservative substitutions), insertions, or deletions relative to the reference sequence, but an antibody or antigen-binding fragment thereof comprising that sequence retains the ability to bind to antigen. In some embodiments, a total of 1 to 10 amino acids have been substituted, inserted and/or deleted in any one of the amino acid sequence of SEQ ID NO: 100. In some
5 embodiments, the substitutions, insertions, or deletions occur in regions outside the CDRs (e.g., in the FRs). Optionally, the antibody or antigen-binding fragment thereof comprises the VL sequence of SEQ ID NO: 100, including post-translational modifications of that sequence. In a particular embodiment, the VL comprises one, two or three CDRs selected from (a) LC-CDR1 comprising the amino acid sequence of SEQ ID NO: 63; (b) LC-CDR2 comprising the amino acid sequence of SEQ ID NO: 71; and (c) LC-CDR3 comprising the
10 amino acid sequence of SEQ ID NO: 79.

[0382] In one aspect, an antibody or antigen-binding fragment thereof is provided, wherein the antibody or antigen-binding fragment thereof comprises a VH as in any of the embodiments provided above, and a VL as in any of the embodiments provided above. In some embodiments, the antibody comprises a VH comprising the amino acid sequence of SEQ ID NO: 86, and a VL sequence in SEQ ID NO: 100, including post-
15 translational modifications of those sequences. In some embodiments, the antibody or antigen binding fragment thereof comprises an IgA heavy chain constant region having at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% sequence identity to the amino acid sequence of any one of SEQ ID NOs: 16-21. In some embodiments, the antibody or antigen binding fragment thereof comprises a kappa light chain constant region of SEQ ID NO: 31.

20 **Improved property and function of IgA antibodies**

[0383] In some embodiments, the antibodies or a functional fragment thereof disclosed herein exhibit an improved stability relative to a corresponding wild type IgA antibody. The term “increased stability” as used herein includes increased thermostability and/or decreased aggregation. Enhanced or improved stability can be determined, for example, by accelerated stability studies. Exemplary accelerated stability studies include,
25 but are not limited to, studies featuring increased storage temperatures. A decrease in the formation of aggregates observed for a antibody as compared to a corresponding WT IgA antibody indicates an increased stability. Stability of antibodies and functional fragment thereof of the present disclosure can be tested by measuring the change in the melting temperature transition of an antibody as compared to the corresponding wild type IgA antibody. immunoglobulin. In such an embodiment, increased stability or increased
30 thermostability would be evident as an increase in the melting temperature transition in the antibody or a functional fragment thereof relative to a corresponding WT IgA or a functional fragment thereof. In some embodiments, the antibodies of a functional fragment thereof have a higher melting temperature than a corresponding WT IgA antibody. In some embodiments, the melting temperature of the antibodies or a functional fragment thereof of the present disclosure is higher than a corresponding WT IgA antibody by at

least about 0.2 fold, 0.3 fold, 0.4 fold, 0.5 fold, 0.6 fold, 0.8 fold, 1.0 fold or higher. In some embodiments, the melting temperature of the antibodies or a functional fragment thereof of the present disclosure is higher than a corresponding WT IgA antibody by at least about 1 ° C, 2 ° C, 3 ° C, 4 ° C, 5 ° C, 6 ° C, 7 ° C, 8 ° C, 9 ° C, 10 ° C, 12 ° C, 15 ° C, or more. In some embodiments, the antibodies, or a functional fragment thereof of the present disclosure have a melting temperature of at least about 60 ° C, 65 ° C, 70 ° C., 75 ° C, 80 ° C, 85 ° C, 90 ° C or more. In some embodiments, this increased stability is in the absence of additional disulfide bonds. Specifically, the increased stability is in the absence of additional disulfide bonds in the IgA heavy chain constant region. In one embodiment, the CH3 domain of the antibody disclosed herein does not contain additional disulfide bonds compared to the wild type CH3 domain. In an alternative embodiment, the CH3 domain of the antibody disclosed herein contains at least one disulfide bond compared to the wild type CH3 domain

[0384] Additional methods for measuring protein aggregation are described in U.S. patent application Ser. No. 10/176,809, and US20030022243A1, the contents of which are incorporated herein by reference in their entities. A variety of analytical techniques for measuring protein stability are available in the art, such as those outlined below: Peptide and Protein Drug Delivery, 247-301, Vincent Lee Ed., Marcel Dekker, Inc., New York, New York, Pubs., 1991; and Jones, A. Adv. Drug Delivery Rev. 10: 29-90, 1993. Stability can be measured at a selected temperature for a selected time. Stability is determined qualitatively and / or quantitatively in a variety of different ways, including determination of aggregate formation (e.g., using size exclusion chromatography or by measuring turbidity and / or visual inspection). it can. The method is as follows: assessment of charge heterogeneity using cation exchange chromatography or capillary zone electrophoresis; analysis of amino-terminal or carboxy-terminal sequences; mass spectrometric analysis; comparing reduced or complete antibodies SDS-PAGE analysis; peptide map (eg, trypsin or LYS-C) analysis; determination of biological activity or antigen-binding function of an antibody. Instability includes one or more of any of the following: aggregation, oxidation (e.g. Met oxidation), isomerization (e.g. Asp isomerization), clipping / hydrolysis / fragmentation (E.g., fragmentation of the hinge region), succinimide formation, unpaired cysteine, N-terminal extension, C-terminal processing, and the like. The term "reduced aggregation" refers to reduction of aggregation of an antibody or a functional fragment thereof of the present disclosure with other antibody molecules and/or with other macromolecule including serum proteins such as albumin, as compared to aggregation exhibited by a corresponding WT IgA antibody.

[0385] The increased thermostability of an antibody relative to a corresponding WT IgA antibody or a variant thereof can be determined by differential scanning calorimetry (DSC) using methods standard in the art (see, for example, Sturtevant, 1987, Annual Review of Physical Chemistry 38: 463-488). The increased thermostability of an antibody relative to a corresponding WT IgA antibody or a variant thereof can also be determined using protein thermal unfolding analysis. Alternatively, the increased thermostability of of

an antibody relative to a corresponding WT IgA antibody or a variant thereof can be determined using any application assay for the antibody, where the performance of the antibody is compared to the WT. For example, ADCC of a target cell, binding to an antigen, or binding to a Fc α R on an immune cell.

Immune effector functions of IgA antibodies

- 5 [0386] Provided herein are engineered IgA variants or antibody comprising one or more modifications within their heavy chain constant region relative to a corresponding WT IgA antibody comprising a WT heavy chain constant region. In some embodiments, the antibodies or the functional fragments thereof disclosed herein (e.g., antibodies comprising one or more modifications disclosed herein) exhibit improved properties.
- 10 [0387] In some embodiments, an antibody or a functional fragment thereof, disclosed herein, exhibits at least one increase in effector function as compared to a corresponding WT IgA antibody or a functional fragment thereof. Effector functions are biological activities attributable to the Fc region of an antibody, which vary with the antibody isotype. Examples of antibody effector functions include: C1q binding and complement dependent cytotoxicity (CDC); Fc receptor binding; antibody-dependent cell-mediated cytotoxicity (ADCC); phagocytosis. For example, an antibody or a functional fragment thereof, disclosed herein exhibits at least one
- 15 increase in effector function as compared to a corresponding WT IgA antibody or a functional fragment thereof or a corresponding WT IgG antibody. Antibody-dependent cellular cytotoxicity (ADCC) is the result of the formation of a complex between the IgA Fab portion of the antibody with an antigen on a cell surface and binding of the Fc portion to the Fc receptors (Fc α Rs), on immune effector cells. For example, an increase in effector function can be increased binding affinity to an Fc receptor (e.g., Fc α Rs), increased ADCC;
- 20 increased cell mediated immunity; increased binding to cytotoxic CD8 T cells; increased binding to NK cells; increased binding to macrophages; increased binding to polymorphonuclear cells; increased binding to monocytes; increased binding to macrophages; increased binding to large granular lymphocytes; increased binding to granulocytes; direct signaling inducing apoptosis; increased dendritic cell maturation; or increased T cell priming, increased opsonization, or increased opsonophagocytosis. In some embodiments, the
- 25 antibodies or functional fragment thereof induce lysis of cancer cells. Lysis can be induced by any mechanism, such as by mediating an effector function, such as C1q binding and complement dependent cytotoxicity (CDC); Fc receptor binding; antibody-dependent cell-mediated cytotoxicity (ADCC); phagocytosis, or direct induction of cell apoptosis.

ADCC

- 30 [0388] "ADCC activity" refers to the ability of an antibody to elicit an ADCC reaction. ADCC is a cell-mediated reaction in which antigen-nonspecific cytotoxic cells that express FcRs (e.g., natural killer (NK) cells, neutrophils, and macrophages) recognize antibody bound to the surface of a target cell and subsequently cause lysis of (i.e., "kill") the target cell (e.g., cancer cell). The primary mediator cells can be natural killer (NK) cells, neutrophils. ADCC activity can be assessed directly using an in vitro assay, e.g., a ⁵¹Cr release

assay using peripheral blood mononuclear cells (PBMC) and/or NK effector cells as described in the Examples and Shields et al. (2001) J. Biol. Chem., 276:6591-6604, or another suitable method known in the art. ADCC activity may be expressed as a concentration of antibody at which the lysis of target cells is half-maximal. Accordingly, in some embodiments, the concentration of an antibody or antigen binding fragment thereof of the disclosure, at which the lysis level is the same as the half-maximal lysis level by the wild-type control, is at least 2-, 3-, 5-, 10-, 20-, 50-, 100-fold lower than the concentration of the wild-type control itself.

[0389] Additionally, in some embodiments, the antibody or a functional fragment thereof of the present disclosure can exhibit a higher maximal target cell lysis as compared to a corresponding wild-type IgA. For example, the maximal target cell lysis of an antibody or a functional fragment thereof of the disclosure can be 2%, 3%, 4%, 5%, 10%, 15%, 20%, 25% or more higher than that of a corresponding WT IgA antibody or a corresponding WT IgG antibody. In some embodiments, the antibodies or a functional fragment thereof disclosed herein induces increased ADCC, compared to a corresponding WT IgG antibody comprising an IgG heavy chain constant domain. In some embodiments, the ADCC is increased by at least 2%, at least 5%, at least 10%, at least 12%, at least 15%, at least 20%, at least 25%, at least 50 %, at least 65%, at least 70%, at least 75%, at least 85%, at least 90%, at least 95%, at least 99%, at least 100%, at least 150%, and at least 200%, relative to a corresponding WT IgG antibody.

CDC

[0390] “Complement dependent cytotoxicity” or “CDC” refer to the ability of a molecule to lyse a target (e.g. cancer cell) in the presence of complement. The complement activation pathway is initiated by the binding of the first component of the complement system (C1q) to a molecule (e.g. an antibody) complexed with a cognate antigen. To assess complement activation, a CDC assay, e.g. as described in Gazzano-Santoro et al., J. Immunol. Methods 202:163 (1996), may be performed.

Fc receptor binding

[0391] In some embodiments, the antibodies or a functional fragment thereof bind a Fc receptor. In some embodiments, the Fc receptor is expressed on an immune effector cell. In some embodiments, the antibodies, or a functional fragment thereof of the present disclosure bind an Fc receptor on an immune effector cell and induce an effector function such as ADCC, CDC, and/or cytolysis of a target cell (e.g., a cancer cell). In some embodiments, the antibodies or a functional fragment thereof bind an IgA receptor. In one embodiment, the IgA receptor is an Fc-alpha receptor (FcαR), such as an FcαR for human IgA. In some embodiments, the FcαR are expressed on an immune effector cell. FcαRs are present on immune effector cells, for example, monocytes, macrophages, neutrophils, and other myeloid cells. FcαRs can also be found on metamyelocytes, myelocytes, promyelocytes and some myeloblasts from, e.g., bone marrow. Such receptors can also be found

on myeloid cell lines, e.g., U937, PLB985, and HL60 cells. It has also been suggested that Fc α Rs are present on lymphocytes. Expression of Fc α Rs can be increased by activation of myeloid cells. For example, stimulation of U937 cells and PLB985 cells with Phorbol Myristic Acetate (PMA) increases the cell surface level of Fc α R several folds (Maliszewski, et al. (1990) J Exp. Med. 172:1665). Other agents which can increase the surface level of Fc α Rs include calcitriol, 1-25 dihydroxy vitamin D3, and interferon- γ (IFN- γ).

[0392] Fc α Rs are capable of interacting with IgA1 and IgA2, in the form of monomers, dimers, and polymers. Binding of antibodies or a functional fragment thereof of the present disclosure to an immune effector cell bearing these receptors (e.g., neutrophils) induces a variety of effector functions, such as phagocytosis, antibody dependent cellular cytotoxicity (ADCC), inflammatory mediator release, lysozyme production, and superoxide anion production (Maliszewski, et al. (1990) J Exp. Med. 172:1665).

[0393] Accordingly, the antibodies or a functional fragment thereof of the present disclosure (e.g., antibodies comprising one or more modifications disclosed herein) are capable of triggering at least one Fc-receptor mediated effector cell function. The term "Fc-receptor mediated effector cell function" is intended to include effector functions, such as those set forth above, which are triggered by binding of immunoglobulin, e.g., IgA, to an Fc receptor on an effector immune cell. An effector immune cell is a cell which is involved in the effector phase of an immune response, as opposed to the cognitive and activation phases of an immune response. Effector immune cells include lymphocytes (e.g., B cells and T cells including cytolytic T cells (CTLs)), killer cells, natural killer cells, macrophages, monocytes, eosinophils, neutrophils, polymorphonuclear cells, granulocytes, mast cells, and basophils. An effector immune cell can phagocytose a target antigen, target cell, or microorganism. An effector immune cell can also lyse a target cell or a microorganism.

[0394] In some embodiments, an effector immune cell is one that can induce ADCC (e.g., That, neutrophils are capable of inducing ADCC) - dependent cells. For example, monocytes, macrophages, which express FcR are to kill target cells specifically, is involved in presenting antigens to other components of the immune system, or binding to antigen-presenting cells. In another embodiment, the effector immune cells can be one that induces phagocytosis for the target antigen, target cell, or microorganism. Effector immune cell can be a macrophage inducing macrophage activity against a target antigen or a target cell or soluble.

[0395] The term "target cell" as used herein refers to a cell that can be targeted by the antibodies or a functional fragment thereof of the present disclosure. In some embodiments, a target cell is a cell expressing or overexpressing an antigen specifically recognized by the antibodies or a functional fragment thereof of the present disclosure. It refers to any cell. In other embodiments, the target cells include tumor cells. The tumor cells can be, for example, of any type, including breast cancer, ovarian cancer, prostate cancer, testicular cancer, lung cancer, colon cancer, rectal cancer, pancreatic cancer, liver cancer, CNS cancer, renal cancer, head cancer, neck cancer, blood cancer and lymphatic cancer of the tumor cells in cancer. In addition to tumor

cells, a target cell can be, for example, IgE- producing lymphocytes for the treatment of autoantibodies targeting or for the production of lymphocytes, allergy or for the treatment of autoimmune diseases. Further, the target may be a microorganism (bacterium or virus) or a soluble antigen (such as rheumatoid factor similar to, or different autoantibodies and toxins). Microorganism includes a pathogen (e.g., virus, bacteria, fungi, protozoa). In some embodiments, a target cell can be a lymphocyte, for example, a CD20 expressing B cell. [0396] In some embodiments, the antibody disclosed herein, exhibit increased binding affinity of to the Fc receptor on an effector immune cell by at least about 2%, at least 5%, at least 10%, at least 12%, at least 15%, at least 20%, at least 25%, at least 50 %, at least 65%, at least 70%, at least 75%, at least 85%, at least 90%, at least 95%, or more relative to a corresponding WT IgA.

Binding to target antigen

[0397] In some embodiments, the antibody and functional fragment thereof of the present disclosure bind a specific target antigen. The term "antigen" refers to any natural or synthetic immunogenic substance immunogenic substance, a fragment or portion of an immunogenic substance, a peptide epitope or a hapten (hapten). Also the term "antigen" is also included in the non-composite material in the form of a non-immunogenic and, when complexed immunogenic. The term "non-complexed" is connected to form a molecular complex of the present invention includes the transfer of material. The term is connected to "composite" includes a substance that forms a molecular complex of the present invention. In some embodiments a target antigen is expressed or overexpressed on a target cell. In some embodiments, binding of the antibodies or a functional fragment thereof to a target antigen induces Fc mediated cytolysis of the target cell expressing the target antigen (e.g., by ADCC). In some embodiments binding of the antibodies or a functional fragment thereof of a present disclosure to a target antigen induces "neutralizing activity" of the target antigen. The term "neutralizing activity" as used herein refers to the ability of the antibodies or a functional fragment thereof to block binding of a cognate ligand to a target antigen.

Reduced glycosylation

[0398] In some embodiments, the antibodies or the functional fragment thereof of the present disclosure exhibit reduced glycosylation compared to a corresponding WT IgA. The antibodies can be completely aglycosylated or partially. The term "glycosylation" refers to the covalent linking of one or more carbohydrates to a polypeptide. Typically, glycosylation is a posttranslational event which can occur within the intracellular milieu of a cell or extract therefrom. The term glycosylation includes, for example, N-linked glycosylation (where one or more sugars are linked to an asparagine residue) and/or O-linked glycosylation (where one or more sugars are linked to an amino acid residue having a hydroxyl group (e.g., serine or threonine). In preferred embodiments, glycosylation refers to N-linked glycosylation. In some

embodiments, reduced glycosylation can be achieved by modifying the asparagine residue in a naturally occurring glycosylation site. In some embodiments, reduced glycosylation is achieved by modifying amino acid residues that are near or within a naturally occurring glycosylation motif or a naturally occurring glycosylation site, for example, an N-linked glycosylation site that contains the amino acid sequence NXT or NXS. In some embodiments, reduced glycosylation is achieved by modifying at least one, at least two, at least three, at least four or more naturally occurring glycosylation site. In some embodiments, the naturally occurring glycosylation site is in a CH2 region. In some embodiments, the naturally occurring glycosylation site is in a CH3 region. In some embodiments, the naturally occurring glycosylation site is in a CH1 region. Consensus motifs, that is, the amino acid sequence recognized by various glycosyl transferases, have been described. For example, the consensus motif for an N-linked glycosylation motif is frequently NXT or NXS, where X can be any amino acid except proline. Accordingly, to identify potential glycosylation sites within an antibody or Fc-containing fragment, the sequence of the antibody is examined, for example, by using publicly available databases such as the website provided by the Center for Biological Sequence Analysis (see www.cbs.dtu.dk/services/NetNGlyc/ for predicting N-linked glycosylation sites) and www.cbs.dtu.dk/services/NetOGlyc/ for predicting O-linked glycosylation sites). Methods to measure glycosylation are known in the art, for example, see Roth et al, International Journal of Carbohydrate Chemistry, Volume 2012, Article ID 640923, and WO2003102018A2, the contents of which are incorporated herein by reference in its entirety.

[0399] In some embodiments, the antibody disclosed herein, exhibit reduced glycosylation by at least about 2%, at least 5%, at least 10%, at least 12%, at least 15%, at least 20%, at least 25%, at least 50 %, at least 65%, at least 70%, at least 75%, at least 85%, at least 90%, at least 95%, or more relative to a corresponding WT IgA. In some embodiments, the antibodies or a functional fragment disclosed herein are partially glycosylated relative to a corresponding WT IgA antibody, for example, less than 100%, 90%, 80%, 70%, 60%, 50%, 40%, 30%, 20%, 10%, 5% or less. In some embodiments, antibodies or a functional fragment thereof disclosed herein are completely glycosylated relative to a corresponding WT IgA antibody.

Biodistribution

In some embodiments, the IgA antibodies or a functional fragment thereof disclosed herein exhibit increased biodistribution relative to a corresponding WT IgA antibody. As used herein, the term “biodistribution” refers to the cellular and/or tissue and/or organ distribution of an IgA antibody or a functional fragment thereof disclosed herein that is administered or delivered to a subject. As used herein, the terms “increased biodistribution” generally refer to the ability of an IgA antibody or a functional fragment thereof disclosed herein that has an increased distribution to a target site, for example, targeted to a tumor or a tumor cell, as compared to the response caused by either vehicle or a corresponding WT IgA antibody. Various methods known to the skilled artisan can be used to assess and increased biodistribution of cells or tissues, including,

but not limited to: nuclear medicine, whole body autoradiography, micro-autoradiography; phosphor imaging, cryo-imaging, nano-secondary ion mass spectroscopy (nanoSIMS), matrix-assisted laser desorption imaging (MALDI-MS), radiography (X-Ray), magnetic resonance imaging (MRI), computed tomography (CT), micro-ultrasound single photon emission CT (SPECT), positron emission tomography (PET) and the like. An increased biodistribution of the antibody to a target site, includes an at least 5%, 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, 100%, 125%, 150%, 175%, 200%, or greater increase compared to a vehicle or a WT IgA antibody. An “increased” or “enhanced” tissue distribution or biodistribution is typically a “statistically significant” increase, and may include an increase that is 1.1, 1.2, 1.5, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 30 or more times (e.g., 500, 1000 times) (including all integers and decimal points in between and above 1, e.g., 1.5, 1.6, 1.7, 1.8, etc.) the distribution of a vehicle or a control composition.

Methods of making an antibody:

[0400] In some embodiments, the antibody or antigen binding fragment thereof is prepared in a host cell. In some embodiments, the antibody or antigen binding fragment thereof is isolated from a host cell. In some embodiments, the antibody or antigen binding fragment thereof is prepared in a cell-free system. An isolated nucleic acid molecule encoding the antibody, portion or polypeptide of the present disclosure can be recombined with vector DNA (e.g., expression vector) in accordance with conventional techniques, including blunt-ended or staggered-ended termini for ligation, restriction enzyme digestion to provide appropriate termini, filling in of cohesive ends as appropriate, alkaline phosphatase treatment to avoid undesirable joining, and ligation with appropriate ligases. Techniques for such manipulations are disclosed, e.g., by Maniatis et al., Molecular Cloning, Lab. Manual (Cold Spring Harbor Lab. Press, NY, 1982 and 1989), and Ausubel, 1987, 1993, and can be used to construct nucleic acid sequences which encode an antibody molecule or antigen binding region thereof. Accordingly, the disclosure provides for a vector or expression vector comprising the isolated nucleic acids set forth herein. In one embodiment, the nucleic acid coding for the light chain and that coding for the heavy chain are isolated separately by the procedures outlined above. In one embodiment, the isolated nucleic acid encoding the light chain and that coding for the heavy chain may be inserted into separate expression plasmids, or together in the same plasmid, so long as each is under suitable promoter and translation control.

[0401] Once the isolated nucleic acid molecule is placed into an expression vector, they are then transfected into host cells such as E. coli cells, simian COS cells, human embryonic kidney 293 cells (e.g., 293E cells), Chinese hamster ovary (CHO) cells, or myeloma cells that do not otherwise produce immunoglobulin protein, to obtain the synthesis of the antibody or an antigen-binding fragment thereof in the recombinant host cells. Recombinant production of antibodies is well known in the art. Many vectors are available. The vector components generally include, but are not limited to, one or more of the following:

a signal sequence, an origin of replication, one or more selective marker genes, an enhancer element, a promoter, and a transcription termination sequence.

5 [0402] The isolated nucleic acid molecules are operably linked to an expression control sequence in the vector DNA. Expression control sequence refers to DNA sequences necessary for the expression of an operably linked coding sequence in a particular host organism. The control sequences that are suitable for prokaryotes, for example, include a promoter, optionally an operator sequence, and a ribosome binding site. Eukaryotic cells are known to utilize promoters, polyadenylation signals, and enhancers. The term “operably linked” or “transcriptional control” refers to functional linkage between a regulatory sequence and a heterologous nucleic acid sequence resulting in expression of the latter. For example, a first nucleic acid sequence is operably linked with a second nucleic acid sequence when the first nucleic acid sequence is placed in a functional relationship with the second nucleic acid sequence. For instance, a promoter is operably linked to a coding sequence if the promoter affects the transcription or expression of the coding sequence. Operably linked DNA sequences can be contiguous with each other and, e.g., where necessary to join two protein coding regions, are in the same reading frame.

15 [0403] Nucleic acid is operably linked when it is placed into a functional relationship with another nucleic acid sequence. For example, a promoter or enhancer is operably linked to a coding sequence if it affects the transcription of the sequence; or a ribosome binding site is operably linked to a coding sequence if it is positioned so as to facilitate translation. Generally, operably linked means that the DNA sequences being linked are contiguous, and/or can be contiguous and in reading phase. However, enhancers do not have to be contiguous. Linking is accomplished by ligation at convenient restriction sites. If such sites do not exist, the synthetic oligonucleotide adaptors or linkers are used in accordance with conventional practice.

20 [0404] Cell, cell line, and cell culture are often used interchangeably and all such designations herein include progeny. Transformants and transformed cells include the primary subject cell and cultures derived therefrom without regard for the number of transfers. It is also understood that all progeny may not be precisely identical in DNA content, due to deliberate or inadvertent mutations. Mutant progeny that have the same function or biological activity as screened for in the originally transformed cell are included. Where distinct designations are intended, it will be clear from the context. In an alternative embodiment, suitable encoding nucleic acid sequences can be designed according to a universal codon table, based on the known amino acid sequence of an immunoglobulin of interest.

30 [0405] Amino acid sequence variants of the desired antibody may be prepared by introducing appropriate nucleotide changes into the encoding DNA, or by peptide synthesis. Such variants include, for example, deletions from, and/or insertions into and/or substitutions of, residues within the amino acid sequences of the antibodies. Any combination of deletion, insertion, and substitution is made to arrive at the final construct, provided that the final construct possesses the desired characteristics. The amino acid changes

also may alter post-translational processes of the monoclonal, human, humanized, or variant antibody, such as changing the number or position of glycosylation sites.

[0406] Nucleic acid molecules encoding amino acid sequence variants of the antibody are prepared by various methods. These methods include, but are not limited to, isolation from a natural source (in the case of naturally occurring amino acid sequence variants) or preparation by oligonucleotide-mediated (or site-directed) mutagenesis, PCR mutagenesis, and cassette mutagenesis of an earlier prepared variant or a non-variant version of the antibody.

[0407] The present disclosure also provides isolated nucleic acid molecules encoding for antibodies or antigen binding fragment thereof, described herein, optionally operably linked to regulatory control sequences recognized by a host cell, vectors and host cells comprising the nucleic acids, and recombinant techniques for the production of the antibodies, which may comprise culturing the host cell so that the nucleic acid is expressed and, optionally, recovering the antibody from the host cell culture or culture medium.

[0408] For recombinant production of an antibody or antigen binding fragment thereof, the nucleic acid molecule encoding it can be isolated and inserted into a replicable vector for further cloning (amplification of the DNA) or for expression. Accordingly provided herein are isolated antibody or antigen binding fragment thereof. In some embodiments, the antibodies or the present disclosure or antigen binding fragment thereof can be recombinant antibody. The term "recombinant antibody" refers to an antibody that is expressed from a cell or cell line transfected with an expression vector (or possibly more than one expression vector, typically two expression vectors) comprising the coding sequence of the antibody, where said coding sequence is not naturally associated with the cell. In one embodiment, a recombinant antibody has a glycosylation pattern that is different than the glycosylation pattern of an antibody having the same sequence if it were to exist in nature. In one embodiment, a recombinant antibody is expressed in a mammalian host cell which is not a human host cell. Notably, individual mammalian host cells have unique glycosylation patterns.

[0409] In some embodiments, the antibodies or antigen binding fragment of the present disclosure is synthetic. The polypeptides of the present disclosure can be purified by isolation/purification methods for proteins generally known in the field of protein chemistry.

[0410] In one aspect, provided herein is a host cell comprising the isolated nucleic acid molecules described herein or a vector comprising said isolated nucleic acid molecules described herein. The vector can be a cloning vector or an expression vector. Suitable host cells for cloning or expressing the DNA in the vectors herein are the prokaryote, yeast, or higher eukaryote cells described above. Suitable prokaryotes for this purpose include eubacteria, such as Gram-negative or Gram-positive organisms, for example, Enterobacteriaceae such as Escherichia, e.g., E. coli, Enterobacter, Erwinia, Klebsiella, Proteus, Salmonella,

e.g., *Salmonella typhimurium*, *Serratia*, e.g., *Serratia marcescens*, and *Shigella*, as well as Bacilli such as *B. subtilis* and *B. licheniformis* (e.g., *B. licheniformis* 41 P disclosed in DD 266,710 published Apr. 12, 1989), *Pseudomonas* such as *P. aeruginosa*, and *Streptomyces*. One preferred *E. coli* cloning host is *E. coli* 294 (ATCC 31,446), although other strains such as *E. coli* B, *E. coli* XI 776 (ATCC 31,537), and *E. coli* W3110 (ATCC 27,325) are suitable. These examples are illustrative rather than limiting.

[0411] In addition to prokaryotes, eukaryotic microbes such as filamentous fungi or yeast are suitable cloning or expression hosts for antibody-encoding vectors. *Saccharomyces cerevisiae*, or common baker's yeast, is the most commonly used among lower eukaryotic host microorganisms. However, a number of other genera, species, and strains are commonly available and useful herein, such as *Schizosaccharomyces pombe*; *Kluyveromyces* hosts such as, e.g., *K. lactis*, *K. fragilis* (ATCC 12,424), *K. bulgaricus* (ATCC 16,045), *K. wickerhamii* (ATCC 24,178), *K. waltii* (ATCC 56,500), *K. drosophilae* (ATCC 36,906), *K. thermotolerans*, and *K. marxianus*; *Yarrowia* (EP 402,226); *Pichia pastoris* (EP 183,070); *Candida*; *Trichoderma reesei* (EP 244,234); *Neurospora crassa*; *Schwanniomyces* such as *Schwanniomyces occidentalis*; and filamentous fungi such as, e.g., *Neurospora*, *Penicillium*, *Tolypocladium*, and *Aspergillus* hosts such as *A. nidulans* and *A. niger*.

[0412] Examples of invertebrate cells include plant and insect cells. Numerous baculoviral strains and variants and corresponding permissive insect host cells from hosts such as *Spodoptera frugiperda* (caterpillar), *Aedes aegypti* (mosquito), *Aedes albopictus* (mosquito), *Drosophila melanogaster* (fruitfly), and *Bombyx mori* have been identified. A variety of viral strains for transfection are publicly available, e.g., the L-1 variant of *Autographa californica* NPV and the Bm-5 strain of *Bombyx mori* NPV, and such viruses may be used as the virus herein according to the present invention, particularly for transfection of *Spodoptera frugiperda* cells.

[0413] Plant cell cultures of cotton, corn, potato, soybean, petunia, tomato, tobacco, *lemna*, and other plant cells can also be utilized as hosts. However, interest has been greatest in vertebrate cells, and propagation of vertebrate cells in culture (tissue culture) has become routine procedure. Examples of useful mammalian host cell lines are Chinese hamster ovary cells, including CHOK1 cells (ATCC CCL61), DXB-11, DG-44, and Chinese hamster ovary cells/DHFR (CHO, Urlaub et al., Proc. Natl. Acad. Sci. USA 77: 4216 (1980)); monkey kidney CV1 line transformed by SV40 (COS-7, ATCC CRL 1651); human embryonic kidney line (293 or 293 cells subcloned for growth in suspension culture, [Graham et al., J. Gen. Virol. 36: 59 (1977)]); baby hamster kidney cells (BHK, ATCC CCL 10); mouse Sertoli cells (TM4, Mather, Biol. Reprod. 23: 243-251 (1980)); monkey kidney cells (CV1 ATCC CCL 70); African green monkey kidney cells (VERO-76, ATCC CRL-1587); human cervical carcinoma cells (HELA, ATCC CCL 2); canine kidney cells (MDCK, ATCC CCL 34); buffalo rat liver cells (BRL 3A, ATCC CRL 1442); human lung cells (W138, ATCC CCL 75); human liver cells (Hep G2, HB 8065); mouse mammary tumor (MMT 060562, ATCC

CCL51); TRI cells (Mather et al., *Annals N.Y Acad. Sci.* 383: 44-68 (1982)); MRC 5 cells; FS4 cells; and a human hepatoma line (Hep G2).

[0414] Host cells are transformed or transfected with the above-described expression or cloning vectors for antibody production and cultured in conventional nutrient media modified as appropriate for inducing promoters, selecting transformants, or amplifying the genes encoding the desired sequences. In addition, novel vectors and transfected cell lines with multiple copies of transcription units separated by a selective marker are particularly useful and preferred for the expression of antibodies, described herein.

[0415] For transfection of the expression vectors and production of the chimeric, humanized, or composite human antibodies described herein, the recipient cell line can be a myeloma cell. Myeloma cells can synthesize, assemble and secrete immunoglobulins encoded by transfected immunoglobulin nucleic acid sequences and possess the mechanism for glycosylation of the immunoglobulin. For example, in some embodiments, the recipient cell is the recombinant Ig-producing myeloma cell SP2/0 (ATCC #CRL 8287). SP2/0 cells produce only immunoglobulin encoded by the transfected genes. Myeloma cells can be grown in culture or in the peritoneal cavity of a mouse, where secreted immunoglobulin can be obtained from ascites fluid. Other suitable recipient cells include lymphoid cells such as B lymphocytes of human or non-human origin, hybridoma cells of human or non-human origin, or interspecies heterohybridoma cells. An expression vector carrying a chimeric, humanized, or composite human antibody construct or antibody polypeptide described herein can be introduced into an appropriate host cell by any of a variety of suitable means, including such biochemical means as transformation, transfection, conjugation, protoplast fusion, calcium phosphate -precipitation, and application with polycations such as diethylaminoethyl (DEAE) dextran, and such mechanical means as electroporation, direct microinjection, and microprojectile bombardment. Johnston et al., 240 *Science* 1538 (1988), as known to one of ordinary skill in the art.

[0416] Yeast provides certain advantages over bacteria for the production of immunoglobulin H and L chains. Yeasts carry out post-translational peptide modifications including glycosylation. A number of recombinant DNA strategies exist that utilize strong promoter sequences and high copy number plasmids which can be used for production of the desired proteins in yeast. Yeast recognizes leader sequences of cloned mammalian gene products and secretes peptides bearing leader sequences (i.e., pre-peptides). Hitzman et al., 11th Intl. Conf. Yeast, Genetics & Molec. Biol. (Montpelier, France, 1982). Yeast gene expression systems can be routinely evaluated for the levels of production, secretion and the stability of antibody polypeptide or antigen binding fragment peptide thereof, and assembled chimeric, humanized, or composite human antibodies, fragments and regions thereof. Any of a series of yeast gene expression systems incorporating promoter and termination elements from the actively expressed genes coding for glycolytic enzymes produced in large quantities when yeasts are grown in media rich in glucose can be utilized. Known glycolytic genes can also provide very efficient transcription control signals. For example,

the promoter and terminator signals of the phosphoglycerate kinase (PGK) gene can be utilized. A number of approaches can be taken for evaluating optimal expression plasmids for the expression of cloned immunoglobulin cDNAs in yeast.

5 [0417] Bacterial strains can also be utilized as hosts for the production of the antibody molecules or fragments thereof described herein, *E. coli* K12 strains such as *E. coli* W3110 (ATCC 27325), *Bacillus* species, enterobacteria such as *Salmonella typhimurium* or *Serratia marcescens*, and various *Pseudomonas* species can be used. Plasmid vectors containing replicon and control sequences which are derived from species compatible with a host cell are used in connection with these bacterial hosts. The vector carries a replication site, as well as specific genes which are capable of providing phenotypic selection in
10 transformed cells. A number of approaches can be taken for evaluating the expression plasmids for the production of chimeric, humanized, or composite humanized antibodies and fragments thereof encoded by the cloned immunoglobulin cDNAs or CDRs in bacteria {see Glover, 1985; Ausubel, 1987, 1993; Sambrook, 1989; Colligan, 1992-1996}.

[0418] Host mammalian cells can be grown in vitro or in vivo. Mammalian cells provide post-translational
15 modifications to immunoglobulin protein molecules including leader peptide removal, folding and assembly of H and L chains, glycosylation of the antibody molecules, and secretion of functional antibody protein. Mammalian cells which can be useful as hosts for the production of antibody proteins, in addition to the cells of lymphoid origin described above, include cells of fibroblast origin, such as Vero (ATCC CRL 81) or CHO-K1 (ATCC CRL 61) cells. Exemplary eukaryotic cells that can be used to express polypeptides
20 include, but are not limited to, COS cells, including COS 7 cells; 293 cells, including 293-6E cells; CHO cells, including CHO-S and DG44 cells; PER.C6.RTM. cells (Crucell); and NSO cells. In some embodiments, a particular eukaryotic host cell is selected based on its ability to make desired post-translational modifications to the variable heavy chains and/or variable light chains. For example, in some
25 embodiments, CHO cells produce polypeptides that have a higher level of sialylation than the same polypeptide produced in 293 cells.

[0419] In some embodiments, polypeptides of the antibodies or antigen binding fragment thereof, disclosed herein can be produced in vivo in an animal that has been engineered or transfected with one or more nucleic acid molecules encoding the polypeptides, according to any suitable method.

[0420] In some embodiments, an antibody or antigen binding fragment thereof is produced in a cell-free
30 system. Nonlimiting exemplary cell-free systems are described, e.g., in Sitaraman et al., *Methods Mol. Biol.* 498: 229-44 (2009); Spirin, *Trends Biotechnol.* 22: 538-45 (2004); Endo et al., *Biotechnol. Adv.* 21: 695-713 (2003).

[0421] Many vector systems are available for the expression of H and L chain nucleic acid sequence in mammalian cells (see Glover, 1985). Different approaches can be followed to obtain complete

H2L2 antibodies. As discussed above, it is possible to co-express H and L chains in the same cells to achieve intracellular association and linkage of H and L chains into complete tetrameric H2L2 antibodies and/or antigen binding fragment peptides. The co-expression can occur by using either the same or different plasmids in the same host. Genes for both H and L chains and/or CDR3 regions peptides can be placed into the same plasmid, which is then transfected into cells, thereby selecting directly for cells that express both chains. Alternatively, cells can be transfected first with a plasmid encoding one chain, for example the L chain, followed by transfection of the resulting cell line with an H chain plasmid containing a second selectable marker. Cell lines producing antigen binding peptide fragments and/or H2L2 molecules via either route could be transfected with plasmids encoding additional copies of peptides, H, L, or H plus L chains in conjunction with additional selectable markers to generate cell lines with enhanced properties, such as higher production of assembled H2L2 antibody molecules or enhanced stability of the transfected cell lines. [0422] Additionally, plants have emerged as a convenient, safe and economical alternative main-stream expression systems for recombinant antibody production, which are based on large scale culture of microbes or animal cells. Antibodies can be expressed in plant cell culture, or plants grown conventionally. The expression in plants may be systemic, limited to sub-cellular plastids, or limited to seeds (endosperms). Several plant-derived antibodies have reached advanced stages of development (see, e.g., Biolex, NC). [0423] In some aspects, provided herein are methods and systems for the production of a humanized antibody, which is prepared by a process which comprises maintaining a host transformed with a first expression vector which encodes the light chain of the humanized antibody and with a second expression vector which encodes the heavy chain of the humanized antibody under such conditions that each chain is expressed and isolating the humanized antibody formed by assembly of the thus-expressed chains. The first and second expression vectors can be the same vector. Also provided herein are DNA sequences encoding the light chain or the heavy chain of the humanized antibody; an expression vector which incorporates a said DNA sequence; and a host transformed with a said expression vector. Generating a humanized antibody from the sequences and information provided herein can be practiced by those of ordinary skill in the art without undue experimentation. In one approach, there are four general steps employed to humanize a monoclonal antibody. These are: (1) determining the nucleotide and predicted amino acid sequence of the starting antibody light and heavy variable domains; (2) designing the humanized antibody, i.e., deciding which antibody framework region to use during the humanizing process; (3) the actual humanizing methodologies/techniques; and (4) the transfection and expression of the humanized antibody.

Purification

[0424] The term “isolated” means altered or removed from the natural state. For example, a nucleic acid or a polypeptide (e.g. antibodies or antigen binding fragment thereof disclosed herein) naturally present in a

living animal is not “isolated,” but the same nucleic acid or polypeptide partially or completely separated from the coexisting materials of its natural state is “isolated.” An isolated nucleic acid molecule or polypeptide can exist in substantially purified form, or can exist in a non-native environment such as, for example, a host cell. An isolated polypeptide is one that has been separated and/or recovered from a component of its natural environment. Contaminant components of its natural environment are materials that would interfere with diagnostic or therapeutic uses of the polypeptide, and may include enzymes, hormones, and other proteinaceous or non-proteinaceous components. In some embodiments, the antibodies or antigen binding fragments thereof of the instant disclosure can be purified by a suitable method. In preferred embodiments, the polypeptide is purified: (1) to greater than 95% by weight of polypeptide as determined by the Lowry method, and most preferably more than 99% by weight; (2) to a degree sufficient to obtain at least 15 residues of N-terminal or internal amino acid sequence by use of a spinning cup sequenator; or (3) to homogeneity as shown by SDS-PAGE under reducing or non-reducing conditions and using Coomassie blue or, preferably, silver staining. Isolated antibody includes the polypeptide in situ within recombinant cells since at least one component of the polypeptide’s natural environment will not be present. Ordinarily, however, isolated polypeptide will be prepared by at least one purification step. In one aspect, disclosed herein is a purified antibody or antigen-binding fragment as provided herein.

[0425] Once expressed, the whole antibodies, their dimers, individual light and heavy chains, or other immunoglobulin forms of the present invention can be recovered and purified by known techniques, e.g., immunoabsorption or immunoaffinity chromatography, chromatographic methods such as HPLC (high performance liquid chromatography), ammonium sulfate precipitation, gel electrophoresis, or any combination of these. See generally, Scopes, *PROTEIN PURIF.* (Springer-Verlag, NY, 1982). Substantially pure immunoglobulins of at least about 90% to 95% homogeneity are advantageous, as are those with 98% to 99% or more homogeneity, particularly for pharmaceutical uses. When using recombinant techniques, the antibody can be produced intracellularly, in the periplasmic space, or directly secreted into the medium, including from microbial cultures. If the antibody is produced intracellularly, as a first step, the particulate debris, either host cells or lysed fragments, is removed, for example, by centrifugation or ultrafiltration. Better et al. *Science* 240: 1041-1043 (1988); ICSU Short Reports 10: 105 (1990); and *Proc. Natl. Acad. Sci. USA* 90: 457-461 (1993) describe a procedure for isolating antibodies which are secreted to the periplasmic space of *E. coli*. (See also, [Carter et al., *Bio/Technology* 10: 163-167 (1992)].

[0426] The antibody composition isolated from microbial or mammalian cells can be purified using, for example, hydroxylapatite chromatography cation or avian exchange chromatography, and affinity chromatography, with affinity chromatography being the preferred purification technique. The suitability of protein A as an affinity ligand depends on the species and isotype of any immunoglobulin Fe domain that is present in the antibody. Protein A can be used to purify antibodies that are based on human $\gamma 1$, $\gamma 2$, or $\gamma 4$

heavy chains (Lindmark et al., J. Immunol. Meth. 62: 1-13 (1983)). Protein G is recommended for all mouse isotypes and for human $\gamma 3$ (Guss et al., EMBO J. 5: 15671575 (1986)). The matrix to which the affinity ligand is attached is most often agarose, but other matrices are available. Mechanically stable matrices such as controlled pore glass or poly(styrenedivinyl)benzene allow for faster flow rates and shorter processing times than can be achieved with agarose. Where the antibody comprises a CH 3 domain, the Bakerbond ABX™ resin (J. T. Baker, Phillipsburg, N.J.) is useful for purification. Other techniques for protein purification such as fractionation on an ion-exchange column, ethanol precipitation, Reverse Phase HPLC, chromatography on silica, chromatography on heparin SEPHAROSE™ chromatography on an anion or cation exchange resin (such as a polyaspartic acid column), chromatofocusing, SDS-PAGE, and ammonium sulfate precipitation are also available depending on the antibody to be recovered. Once purified, partially or to homogeneity as desired, a humanized or composite human antibody can then be used therapeutically or in developing and performing assay procedures, immunofluorescent staining, and the like. See generally, Vols. I & II Immunol. Meth. (Lefkovits & Pernis, eds., Acad. Press, NY, 1979 and 1981).

[0427] Functional activities of an antibody or antigen-binding fragment disclosed herein. Such functional activities include biological activity and ability to bind to a cancer cell antigen. Additionally, a polypeptide having functional activity means the polypeptide exhibits activity similar, but not necessarily identical to, an activity of an antibody described herein, including mature forms, as measured in a particular assay, such as, for example, a biological assay, with or without dose dependency. In the case where dose dependency does exist, it need not be identical to that of the antibodies of the disclosure, but rather substantially similar to the dose-dependence in a given activity as compared to the antibodies set forth herein (i.e., the candidate polypeptide will exhibit greater activity, or not more than about 25-fold less, about 10-fold less, or about 3-fold less activity relative to the antibodies described herein).

25 Nucleic Acid Molecules Encoding Antibodies

[0428] Using the information provided herein, for example, nucleic acid and amino acid sequences of the antibodies; a nucleic acid molecule encoding the antibodies or antigen-binding fragment thereof can be easily obtained by a skilled artisan. Such a nucleic acid molecule can be obtained, for example, using conventional methods disclosed in the art. Nucleic acid molecules of the present disclosure may be in the form of RNA, such as mRNA, hnRNA, tRNA or any other form, or in the form of DNA, including but not limited to, cDNA and genomic DNA obtained by cloning or produced synthetically, or any combinations thereof. The DNA may be triplex, duplex or single-stranded, or any combination thereof. Any portion of at least one strand of the DNA or RNA may be the coding strand, also known as the sense strand, or it can be the antisense strand, also known as the antisense strand.

[0429] “Polynucleotide,” or “nucleic acid molecule,” as used interchangeably herein, refer to polymers of nucleotides of any length, and include DNA and RNA. The nucleotides can be deoxyribonucleotides, ribonucleotides, modified nucleotides or bases, and/or their analogs, or any substrate that can be incorporated into a polymer by DNA or RNA polymerase. A nucleic acid molecule can comprise modified nucleotides, such as methylated nucleotides and their analogs. If present, modification to the nucleotide structure may be imparted before or after assembly of the polymer. The sequence of nucleotides may be interrupted by non-nucleotide components. A polynucleotide may be further modified after polymerization, such as by conjugation with a labeling component. Other types of modifications include, for example, “caps”, substitution of one or more of the naturally occurring nucleotides with an analog, internucleotide modifications such as, for example, those with uncharged linkages (e.g., methyl phosphonates, phosphotriesters, phosphoamidates, cabamates, etc.) and with charged linkages (e.g., phosphorothioates, phosphorodithioates, etc.), those containing pendant moieties, such as, for example, proteins (e.g., nucleases, toxins, antibodies, signal peptides, ply-L-lysine, etc.), those with intercalators (e.g., acridine, psoralen, etc.), those containing chelators (e.g., metals, radioactive metals, boron, oxidative metals, etc.), those containing alkylators, those with modified linkages (e.g., alpha anomeric nucleic acids, etc.), as well as unmodified forms of the polynucleotide(s). Further, any of the hydroxyl groups ordinarily present in the sugars may be replaced, for example, by phosphonate groups, phosphate groups, protected by standard protecting groups, or activated to prepare additional linkages to additional nucleotides, or may be conjugated to solid supports. The 5' and 3' terminal OH can be phosphorylated or substituted with amines or organic capping group moieties of from 1 to 20 carbon atoms. Other hydroxyls may also be derivatized to standard protecting groups. Polynucleotides can also contain analogous forms of ribose or deoxyribose sugars that are generally known in the art, including, for example, 2'-O-methyl-, 2'-O-allyl, 2'-fluoro- or 2'-azido-ribose, carbocyclic sugar analogs, α -anomeric sugars, epimeric sugars such as arabinose, xyloses or lyxoses, pyranose sugars, furanose sugars, sedoheptuloses, acyclic analogs and abasic nucleoside analogs such as methyl riboside. One or more phosphodiester linkages may be replaced by alternative linking groups. These alternative linking groups include, but are not limited to, embodiments wherein phosphate is replaced by P(O)S(“thioate”), P(S)S(“dithioate”), “(O)NR₂ (“amidate”), P(O)R, P(O)OR', CO or CH₂ (“formacetal”), in which each R or R' is independently H or substituted or unsubstituted alkyl (1-20 C) optionally containing an ether (—O—) linkage, aryl, alkenyl, cycloalkyl, cycloalkenyl or araldyl. Not all linkages in a polynucleotide need be identical. The preceding description applies to all polynucleotides referred to herein, including isolated nucleic acid, RNA and DNA.

[0430] In the context of the present invention, the following abbreviations for the commonly occurring nucleic acid bases are used. “A” refers to adenosine, “C” refers to cytosine, “G” refers to guanosine, “T”

refers to thymidine, and “U” refers to uridine. In some embodiments, the nucleic acid molecule comprises an isolated nucleic acid.

[0431] The nucleic acids can be present in whole cells, in a cell lysate, or in a partially purified or substantially pure form. A nucleic acid molecule is “isolated” or “rendered substantially pure” when
5 purified away from other cellular components or other contaminants, e.g., other cellular nucleic acids or proteins, by standard techniques, including, but not limited to alkaline/SDS treatment, CsCl banding, column chromatography, agarose gel electrophoresis and others well known in the art. See, F. Ausubel, et al., ed. (1987) Current Protocols in Molecular Biology, Greene Publishing and Wiley Interscience, New York. A nucleic acid according to at least some embodiments of the disclosure can be, for example, DNA
10 or RNA and may or may not contain intronic sequences. In a preferred embodiment, the nucleic acid is a cDNA molecule.

[0432] Another aspect of the present disclosure pertains to nucleic acid molecules comprising nucleic acid sequences that encode the antibody polypeptide or a functional fragment thereof, described herein or antigen-binding fragment thereof. In some embodiments, the isolated nucleic acid molecule comprises a
15 nucleic acid sequence encoding a modified IgA heavy chain constant region. In some embodiments, the nucleic acid sequence encoding a modified IgA heavy chain constant region comprises a sequence selected from any one of SEQ ID NOs 25-32. In some embodiments, the nucleic acid sequence encoding a variable heavy chain polypeptide is selected from SEQ ID NOs.: 87-84. In some embodiments, the isolated nucleic acid molecule comprises a nucleic acid sequence encoding a light chain variable region polypeptide of an
20 antibody. In some embodiments, the nucleic acid sequence encoding a light chain variable region polypeptide is selected from SEQ ID NOs.: 101-108.

[0433] Once DNA fragments encoding VH and VL segments are obtained, these DNA fragments can be further manipulated by standard recombinant DNA techniques, for example to convert the variable region genes to full-length antibody chain genes, to Fab fragment genes or to a scFv gene. In these manipulations,
25 a VL- or VH-encoding DNA fragment is operatively linked to another DNA fragment encoding another protein, such as an antibody constant region or a flexible linker. The term “operatively linked”, as used in this context, is intended to mean that the two DNA fragments are joined such that the amino acid sequences encoded by the two DNA fragments remain in-frame. The isolated DNA encoding the VH region can be converted to a full-length heavy chain gene by operatively linking the VH-encoding DNA to another DNA
30 molecule encoding heavy chain constant regions (CH1, CH2 and CH3). The sequences of human heavy chain constant region genes are known in the art (see e.g., Kabat, E. A., et al. (1991) Sequences of Proteins of Immunological Interest, Fifth Edition, U.S. Department of Health and Human Services, NIH Publication No. 91-3242) and DNA fragments encompassing these regions can be obtained by standard PCR amplification. The heavy chain constant region can be an IgA1 or a IgA2 constant region. For a Fab

fragment heavy chain gene, the VH-encoding DNA can be operatively linked to another DNA molecule encoding only the heavy chain CH1 constant region.

[0434] The isolated DNA encoding the VL region can be converted to a full-length light chain gene (as well as a Fab light chain gene) by operatively linking the VL-encoding DNA to another DNA molecule encoding the light chain constant region, CL. The sequences of human light chain constant region genes are known in the art (see e.g., Kabat, E. A., et al. (1991) Sequences of Proteins of Immunological Interest, Fifth Edition, U.S. Department of Health and Human Services, NIH Publication No. 91-3242) and DNA fragments encompassing these regions can be obtained by standard PCR amplification. The light chain constant region can be a kappa or lambda constant region, but most preferably is a kappa constant region.

[0435] To create a scFv gene, the VH- and VL-encoding DNA fragments are operatively linked to another fragment encoding a flexible linker, e.g., encoding the amino acid sequence (Gly-4-Ser)₃, such that the VH and VL sequences can be expressed as a contiguous single-chain protein, with the VL and VH regions joined by the flexible linker (see e.g., Bird et al. (1988) Science 242:423-426; Huston et al. (1988) Proc. Natl. Acad. Sci. USA 85:5879-5883; McCafferty et al., (1990) Nature 348:552-554).

[0436] Nucleic acid molecules isolated from the present disclosure can include nucleic acid molecules comprising an open reading frame (ORF), optionally with one or more introns, e.g., but not limited to, at least one specified portion of at least a CDR, as CDR1, CDR2 and / or CDR3 of at least one light chain; nucleic acid molecules comprising the coding sequence of a cancer associated antibody disclosed herein or variable region e.g., variable regions of the light chain; and nucleic acid molecules comprising a nucleotide sequence substantially different from those described above but which, due to the degeneracy of the genetic code, still encode at least antibody or antigen binding fragment thereof as described herein and / or as it is known in the art. Of course, the genetic code is well known in the art. Therefore, it would be routine for one skilled in the art to generate such degenerate nucleic acid variants encoding specific antibodies of the present disclosure. See for example, Ausubel et al., Supra, and such nucleic acid variants are included in the present invention.

[0437] Nucleic acid molecules comprising nucleic acid sequence that encode one or more chains of an antibody are provided herein. In some embodiments, a nucleic acid molecule comprises a nucleic acid sequence that encodes a heavy chain or a light chain of an antibody. In some embodiments, a nucleic acid molecule comprises both a nucleic acid sequence that encodes a heavy chain and a nucleic acid sequence that encodes a light chain, of an antibody. In some embodiments, a first nucleic acid molecule comprises a first nucleic acid sequence that encodes a heavy chain and a second nucleic acid molecule comprises a second nucleic acid sequence that encodes a light chain.

[0438] In some embodiments, the heavy chain and the light chain are expressed from one nucleic acid molecule, or from two separate nucleic acid molecules, as two separate polypeptides. In some

embodiments, such as when an antibody is an scFv, a single nucleic acid sequence encodes a single polypeptide comprising both a heavy chain and a light chain linked together.

[0439] In some embodiments, a nucleic acid sequence encoding a heavy chain or light chain of an antibody disclosed herein comprises a nucleic acid sequence that encodes at least one of the CDRs provided herein.

5 In some embodiments, a nucleic acid sequence encoding a heavy chain or light chain of an antibody disclosed herein comprises a sequence that encodes at least 3 of the CDRs provided herein. In some embodiments, a nucleic acid sequence encoding a heavy chain or light chain of an antibody comprises a sequence that encodes at least 6 of the CDRs provided herein. In some embodiments, a nucleic acid sequence encoding a heavy chain or light chain of an antibody comprises a nucleotide sequence that
10 encodes a leader sequence, which, when translated, is located at the N terminus of the heavy chain or light chain. The leader sequence may be the native heavy or light chain leader sequence, or may be another heterologous leader sequence. The term "leader sequence" refers to a sequence of amino acid residues located at the N-terminus of a polypeptide that facilitates secretion of a polypeptide from a mammalian cell. A leader sequence can be cleaved upon export of the polypeptide from the mammalian cell, forming a
15 mature protein. Leader sequences can be natural or synthetic, and they can be heterologous or homologous to the protein to which they are attached.

[0440] In some embodiments, the nucleic acid molecule is one that encodes for any of the amino acid sequences for the variable light chain and variable heavy chain in the Tables 7-8 herein. In some embodiments, the nucleic acid sequence is one that is at least 80% identical to a nucleic acid encoding any
20 of the amino acid sequences variable light chain and variable heavy chain in the Tables 7-8 herein, for example, at least 80, 85, 90, 91, 92, 93, 94, 95, 96, 97, 98, or 99% identical. In some embodiments, the nucleic acid is one that hybridizes to any one or more of the nucleic acid sequences provided herein. In some of the embodiments, the hybridization is under moderate conditions. In some embodiments, the hybridization is under highly stringent conditions, such as: at least about 6X SSC and 1% SDS at 65°C, with
25 a first wash for 10 minutes at about 42°C with about 20% (v/v) formamide in 0.1X SSC, and with a subsequent wash with 0.2 X SSC and 0.1% SDS at 65°C.

[0441] Nucleic acid molecules can be constructed using recombinant DNA techniques conventional in the art. In some embodiments, a nucleic acid molecule is placed in an expression vector that is suitable for expression in a selected host cell.

30 **[0442]** Vectors comprising nucleic acid molecules that encode the antibodies or antigen binding fragment herein are provided. Vectors comprising nucleic acid molecules that encode a heavy chains and/or a light chains are also provided. Such vectors include, but are not limited to, DNA vectors, phage vectors, viral vectors, retroviral vectors, etc. In one embodiment, the nucleic acid coding for the light chain and that coding for the heavy chain are isolated separately by the procedures outlined above. In one embodiment, the

isolated nucleic acid encoding the light chain and that coding for the heavy chain may be inserted into separate expression plasmids, or together in the same plasmid, so long as each is under suitable promoter and translation control. In some embodiments, the heavy chain and light chain are expressed as part of a single polypeptide, such as, for example, when the antibody is an scFv.

- 5 [0443] In some embodiments, a first vector comprises a nucleic acid molecule that encodes a heavy chain and a second vector comprises a nucleic acid molecule that encodes a light chain. In some embodiments, the first vector and second vector are transfected into host cells in similar amounts (such as similar molar amounts or similar mass amounts). In some embodiments, a mole- or mass-ratio of between 5:1 and 1:5 of the first vector and the second vector is transfected into host cells. In some embodiments, a mass ratio of
10 between 1:1 and 1:5 for the vector encoding the heavy chain and the vector encoding the light chain is used. In some embodiments, a mass ratio of 1:2 for the vector encoding the heavy chain and the vector encoding the light chain is used. In some embodiments, a vector is selected that is optimized for expression of polypeptides in CHO or CHO-derived cells, or in NSO cells. Exemplary such vectors are described, for example, in Running Deer et al., *Biotechnol. Prog.* 20:880-889 (2004).
- 15 [0444] In one aspect, the present disclosure provides methods for treatment or prevention of cancer comprising administering nucleic acid molecules, wherein the nucleic acid molecules encode for a VH, VL, CDR3 region of VH or CDR 3 region of VL or antigen binding fragment thereof, wherein the nucleic acid molecule comprises a sequence disclosed herein (e.g. Table 3 or Table 4) by way of gene therapy. Gene therapy refers to therapy performed by the administration to a subject of an expressed or expressible nucleic
20 acid. In this embodiment of the invention, the nucleic acids produce their encoded protein that mediates a prophylactic or therapeutic effect. Any of the methods for gene therapy available in the art can be used according to the embodiments herein.
- [0445] For general reviews of the methods of gene therapy, see Goldspiel et al., 1993, *Clinical Pharmacy* 12:488-505; Wu and Wu, 1991, *Biotherapy* 3:87-95; Tolstoshev, 1993, *Ann. Rev. Pharmacol. Toxicol.* 32:573-596; Mulligan, 1993, *Science* 260:926-932; and Morgan and Anderson, 1993, *Ann. Rev. Biochem.* 62:191-217; May, 1993, *TIBTECH* 11(5):155-215. Methods commonly known in the art of recombinant DNA technology which can be used are described in Ausubel et al. (eds.), 1993, *Current Protocols in Molecular Biology*, John Wiley & Sons, NY; and Kriegler, 1990, *Gene Transfer and Expression, A Laboratory Manual*, Stockton Press, NY. Delivery of a therapeutic antibody to appropriate cells can be
25 effected via gene therapy ex vivo, in situ, or in vivo by use of any suitable approach known in the art, including by use of physical DNA transfer methods (e.g., liposomes or chemical treatments) or by use of viral vectors (e.g., adenovirus, adeno-associated virus, or a retrovirus). For example, for in vivo therapy, a nucleic acid encoding the desired antibody, either alone or in conjunction with a vector, liposome, or precipitate may be injected directly into the subject, and in some embodiments, may be injected at the site
30

where the expression of the antibody compound is desired. For ex vivo treatment, the subject's cells are removed, the nucleic acid is introduced into these cells, and the modified cells are returned to the subject either directly or, for example, encapsulated within porous membranes which are implanted into the patient. See, e.g. U.S. Pat. Nos. 4,892,538 and 5,283,187. There are a variety of techniques available for introducing

5 nucleic acids into viable cells. The techniques vary depending upon whether the nucleic acid is transferred into cultured cells in vitro, or in vivo in the cells of the intended host. Techniques suitable for the transfer of nucleic acid into mammalian cells in vitro include the use of liposomes, electroporation, microinjection, cell fusion, DEAE-dextran, and calcium phosphate precipitation. A commonly used vector for ex vivo delivery of a nucleic acid is a retrovirus.

- 10 **[0446]** The term “host cell” as used herein refers to the particular subject cell transfected with a nucleic acid molecule and the progeny or potential progeny of such a cell. Progeny of such a cell may not be identical to the parent cell transfected with the nucleic acid molecule due to mutations or environmental influences that may occur in succeeding generations or integration of the nucleic acid molecule into the host cell genome.
- 15 **[0447]** Other in vivo nucleic acid transfer techniques include transfection with viral vectors (such as adenovirus, Herpes simplex I virus, or adeno-associated virus) and lipid-based systems. The nucleic acid and transfection agent are optionally associated with a microparticle. Exemplary transfection agents include calcium phosphate or calcium chloride co-precipitation, DEAE-dextran-mediated transfection, quaternary ammonium amphiphile DOTMA ((dioleoyloxypropyl) trimethylammonium bromide, commercialized as
- 20 Lipofectin by GIBCO-BRL))(Felgner et al., (1987) Proc. Natl. Acad. Sci. USA 84, 7413-7417; Malone et al. (1989) Proc. Natl. Acad. Sci. USA 86 6077-6081); lipophilic glutamate diesters with pendent trimethylammonium heads (Ito et al. (1990) Biochem. Biophys. Acta 1023, 124-132); the metabolizable parent lipids such as the cationic lipid dioctadecylamido glycylspermine (DOGS, Transfectam, Promega) and dipalmitoylphosphatidyl ethanolamylspermine (DPPES)(J. P. Behr (1986) Tetrahedron Lett. 27, 5861-
- 25 5864; J. P. Behr et al. (1989) Proc. Natl. Acad. Sci. USA 86, 6982-6986); metabolizable quaternary ammonium salts (DOTB, N-(1-[2,3-dioleoyloxy]propyl)-N,N,N-trimethylammonium methylsulfate (DOTAP)(Boehringer Mannheim), polyethyleneimine (PEI), dioleoyl esters, ChoTB, ChoSC, DOSC)(Leventis et al. (1990) Biochim. Inter. 22, 235-241); 3beta[N-(N',N'-dimethylaminoethane)-carbamoyl]cholesterol (DC-Chol), dioleoylphosphatidyl ethanolamine (DOPE)/3beta[N-(N',N'
- 30 dimethylaminoethane)-carbamoyl] cholesterolDC-Chol in one to one mixtures (Gao et al., (1991) Biochim. Biophys. Acta 1065, 8-14), spermine, spermidine, lipopolyamines (Behr et al., Bioconjugate Chem, 1994, 5: 382-389), lipophilic polylysines (LPLL) (Zhou et al., (1991) Biochim. Biophys. Acta 939, 8-18), [(1, 1,3,3 tetramethylbutyl)cresoxy]ethoxy]ethyl]dimethylbnzylammonium hydroxide (DEBDA hydroxide) with excess phosphatidylcholine/cholesterol (Ballas et al., (1988) Biochim. Biophys. Acta 939, 8-18),

cetyltrimethylammonium bromide (CTAB)/DOPE mixtures (Pinnaduwa et al, (1989) *Biochim. Biophys. Acta* 985, 33-37), lipophilic diester of glutamic acid (TMAG) with DOPE, CTAB, DEBDA, didodecylammonium bromide (DDAB), and stearylamine in admixture with phosphatidylethanolamine (Rose et al., (1991) *Biotechnology* 10, 520-525), DDAB/DOPE (TransfectACE, GIBCO BRL), and

5 oligogalactose bearing lipids. Exemplary transfection enhancer agents that increase the efficiency of transfer include, for example, DEAE-dextran, polybrene, lysosome-disruptive peptide (Ohmori N I et al, *Biochem Biophys Res Commun* Jun. 27, 1997; 235(3):726-9), chondroitin-based proteoglycans, sulfated proteoglycans, polyethylenimine, polylysine (Pollard H et al. *J Biol Chem*, 1998 273 (13):7507-11), integrin-binding peptide CYGGRGDTP, linear dextran nonasaccharide, glycerol, cholesteryl groups

10 tethered at the 3'-terminal internucleoside link of an oligonucleotide (Letsinger, R. L. 1989 *Proc Natl Acad Sci USA* 86: (17):6553-6), lysophosphatide, lysophosphatidylcholine, lysophosphatidylethanolamine, and 1-oleoyl lysophosphatidylcholine.

[0448] In some situations, it may be desirable to deliver the nucleic acid with an agent that directs the nucleic acid containing vector to target cells. Such "targeting" molecules include antibodies specific for a

15 cell-surface membrane protein on the target cell, or a ligand for a receptor on the target cell. Where liposomes are employed, proteins which bind to a cell-surface membrane protein associated with endocytosis may be used for targeting and/or to facilitate uptake. Examples of such proteins include capsid proteins and fragments thereof tropic for a particular cell type, antibodies for proteins which undergo internalization in cycling, and proteins that target intracellular localization and enhance intracellular half-

20 life. In other embodiments, receptor-mediated endocytosis can be used. Such methods are described, for example, in Wu et al., 1987 or Wagner et al., 1990. For review of the currently known gene marking and gene therapy protocols, see Anderson 1992. See also WO 93/25673 and the references cited therein.

Chimeric antigen receptor

[0449] In one aspect, the disclosure herein, provides a chimeric antigen receptor comprising, an antigen

25 binding fragment, disclosed herein, a transmembrane domain, and an intracellular signaling domain. The term "chimeric Antigen Receptor" (CAR), "artificial T cell receptor", "chimeric T cell receptor", or "chimeric immunoreceptor" as used herein refers to an engineered receptor, which grafts an arbitrary specificity onto an immune effector cell. CARs typically have an extracellular domain (ectodomain), which comprises an antigen-binding domain, a transmembrane domain, and an intracellular (endodomain) domain.

30 The term "signaling domain" refers to the functional portion of a protein which acts by transmitting information within the cell to regulate cellular activity via defined signaling pathways by generating second messengers or functioning as effectors by responding to such messengers.

[0450] An "intracellular signaling domain," as the term is used herein, refers to an intracellular portion of a molecule. The intracellular signaling domain generates a signal that promotes an immune effector function

of the CAR containing cell, e.g., a CART cell. Examples of immune effector function, e.g., in a CART cell, include cytolytic activity and helper activity, including the secretion of cytokines.

[0451] In an embodiment, the intracellular signaling domain can comprise a primary intracellular signaling domain. Exemplary primary intracellular signaling domains include those derived from the molecules responsible for primary stimulation, or antigen dependent stimulation. In an embodiment, the intracellular signaling domain can comprise a costimulatory intracellular domain. Exemplary costimulatory intracellular signaling domains include those derived from molecules responsible for costimulatory signals, or antigen independent stimulation. For example, in the case of a CART, a primary intracellular signaling domain can comprise a cytoplasmic sequence of a T cell receptor, and a costimulatory intracellular signaling domain can comprise cytoplasmic sequence from co-receptor or costimulatory molecule.

[0452] A primary intracellular signaling domain can comprise a signaling motif which is known as an immunoreceptor tyrosine-based activation motif or ITAM. Examples of ITAM containing primary cytoplasmic signaling sequences include, but are not limited to, those derived from CD3 zeta, FcR gamma, FcR beta, CD3 gamma, CD3 delta, CD3 epsilon, CD5, CD22, CD79a, CD79b, and CD66d DAP10 and DAP12.

[0453] The term “zeta” or alternatively “zeta chain”, “CD3-zeta” or “TCR-zeta” is defined as the protein provided as GenBank Acc. No. BAG36664.1, or the equivalent residues from a non-human species, e.g., mouse, rodent, monkey, ape and the like, and a “zeta stimulatory domain” or alternatively a “CD3-zeta stimulatory domain” or a “TCR-zeta stimulatory domain” is defined as the amino acid residues from the cytoplasmic domain of the zeta chain that are sufficient to functionally transmit an initial signal necessary for T cell activation. In one aspect the cytoplasmic domain of zeta comprises residues 52 through 164 of GenBank Acc. No. BAG36664.1 or the equivalent residues from a non-human species, e.g., mouse, rodent, monkey, ape and the like, that are functional orthologs thereof.

[0454] The term “costimulatory molecule” refers to the cognate binding partner on a T cell that specifically binds with a costimulatory ligand, thereby mediating a costimulatory response by the T cell, such as, but not limited to, proliferation. Costimulatory molecules are cell surface molecules other than antigen receptors or their ligands that are required for an efficient immune response. Costimulatory molecules include, but are not limited to, an MHC class I molecule, BTLA and a Toll ligand receptor, as well as OX40, CD2, CD27, CD28, CD5, ICAM-1, LFA-1 (CD11a/CD18) and 4-1BB (CD137).

[0455] A costimulatory intracellular signaling domain can be derived from the intracellular portion of a costimulatory molecule. A costimulatory molecule can be represented in the following protein families: TNF receptor proteins, Immunoglobulin-like proteins, cytokine receptors, integrins, signaling lymphocytic activation molecules (SLAM proteins), and activating NK cell receptors. Examples of such molecules include CD27, CD28, 4-1BB (CD137), OX40, GITR, CD30, CD40, ICOS, BAFFR, HVEM, lymphocyte

function-associated antigen-1 (LFA-1), CD2, CD7, LIGHT, NKG2C, SLAMF7, NKp80, CD160, B7-H3, and a ligand that specifically binds with CD83, and the like.

[0456] The intracellular signaling domain can comprise the entire intracellular portion, or the entire native intracellular signaling domain, of the molecule from which it is derived, or a functional fragment thereof.

- 5 **[0457]** In another aspect, the antigen binding fragment comprises a humanized antibody or antibody fragment. In one embodiment, the antigen binding fragment comprises one or more (e.g., one, two, or all three) light chain complementary determining region 1 (LC-CDR1), light chain complementary determining region 2 (LC-CDR2), and light chain complementary determining region 3 (LC-CDR3) of an antibody described herein, and one or more (e.g., one, two, or all three) heavy chain complementary determining region 1 (HC-CDR1), heavy chain complementary determining region 2 (HC-CDR2), and heavy chain complementary determining region 3 (HC-CDR3) of an antibody described herein.

Transmembrane domain

- [0458]** With respect to the transmembrane domain, in various embodiments, a CAR can be designed to comprise a transmembrane domain that is attached to the extracellular domain of the CAR. A
15 transmembrane domain can include one or more additional amino acids adjacent to the transmembrane region, e.g., one or more amino acid associated with the extracellular region of the protein from which the transmembrane was derived (e.g., 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 up to 15 amino acids of the extracellular region) and/or one or more additional amino acids associated with the intracellular region of the protein from which the transmembrane protein is derived (e.g., 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 up to 15 amino acids of the
20 intracellular region). In one aspect, the transmembrane domain is one that is associated with one of the other domains of the CAR is used. In some instances, the transmembrane domain can be selected or modified by amino acid substitution to avoid binding of such domains to the transmembrane domains of the same or different surface membrane proteins, e.g., to minimize interactions with other members of the receptor complex. In one aspect, the transmembrane domain is capable of homodimerization with another CAR on
25 the CAR T-cell surface. In a different aspect, the amino acid sequence of the transmembrane domain may be modified or substituted so as to minimize interactions with the binding domains of the native binding partner present in the same CAR T-cell.

- [0459]** The transmembrane domain may be derived either from a natural or from a recombinant source. Where the source is natural, the domain may be derived from any membrane-bound or transmembrane
30 protein. In one aspect the transmembrane domain is capable of signaling to the intracellular domain(s) whenever the CAR has bound to a target. A transmembrane domain, for example, can include at least the transmembrane region(s) of e.g., the alpha, beta or zeta chain of the T-cell receptor, CD28, CD3 epsilon, CD45, CD4, CD5, CD8, CD9, CD16, CD22, CD33, CD37, CD64, CD80, CD86, CD134, CD137, CD154.

[0460] In some instances, the transmembrane domain can be attached to the extracellular region of the CAR, e.g., the antigen-binding domain of the CAR, via a hinge, e.g., a hinge from a human protein. For example, in one embodiment, the hinge can be a human Ig (immunoglobulin) hinge, e.g., an IgG4 hinge, or a CD8a hinge. In one aspect, the hinge or spacer comprises an IgG4 hinge.

5

Cytoplasmic Domain

[0461] The cytoplasmic domain or region of the CAR includes an intracellular signaling domain. An intracellular signaling domain is generally responsible for activation of at least one of the normal effector functions of the immune cell in which the CAR has been introduced. The term “effector function” refers to a specialized function of a cell. Effector function of a T cell, for example, may be cytolytic activity or helper activity including the secretion of cytokines. The term “intracellular signaling domain” refers to the portion of a protein which transduces the effector function signal and directs the cell to perform a specialized function. While usually the entire intracellular signaling domain can be employed, in many cases it is not necessary to use the entire chain. To the extent that a truncated portion of the intracellular signaling domain is used, such truncated portion may be used in place of the intact chain as long as it transduces the effector function signal. The term intracellular signaling domain is thus meant to include any truncated portion of the intracellular signaling domain sufficient to transduce the effector function signal.

10

15

20

[0462] Examples of intracellular signaling domains for use in the CAR of the invention include the cytoplasmic sequences of the T cell receptor (TCR) and co-receptors that act in concert to initiate signal transduction following antigen receptor engagement, as well as any derivative or variant of these sequences and any recombinant sequence that has the same functional capability.

25

[0463] It is known that signals generated through the TCR alone are insufficient for full activation of the T cell and that a secondary and/or costimulatory signal is also required. Thus, T cell activation can be said to be mediated by two distinct classes of cytoplasmic signaling sequences: those that initiate antigen-dependent primary activation through the TCR (primary intracellular signaling domains) and those that act in an antigen-independent manner to provide a secondary or costimulatory signal (secondary cytoplasmic signaling domain, e.g., a costimulatory domain).

30

[0464] A primary signaling domain regulates primary activation of the TCR complex either in a stimulatory way, or in an inhibitory way. Primary intracellular signaling domains that act in a stimulatory manner may contain signaling motifs which are known as immunoreceptor tyrosine-based activation motifs or ITAMs.

[0465] Examples of ITAM containing primary intracellular signaling domains that are of particular use in the invention include those of TCR zeta, FcR gamma, FcR beta, CD3 gamma, CD3 delta, CD3 epsilon, CD5, CD22, CD79a, CD79b, and CD66d. In one embodiment, a CAR of the invention, e.g., a CAR

comprises a intracellular signaling domain, e.g., a primary signaling domain, of CD3-zeta. In one embodiment, a primary signaling domain comprises a modified ITAM domain, e.g., a mutated ITAM domain which has altered (e.g., increased or decreased) activity as compared to the native ITAM domain. In one embodiment, a primary signaling domain comprises a modified ITAM-containing primary intracellular signaling domain, e.g., an optimized and/or truncated ITAM-containing primary intracellular signaling domain. In an embodiment, a primary signaling domain comprises one, two, three, four or more ITAM motifs.

[0466] The intracellular signaling domain of the CAR can comprise the CD3-zeta signaling domain by itself or it can be combined with any other desired intracellular signaling domain(s) useful in the context of a CAR of the invention. For example, the intracellular signaling domain of the CAR can comprise a CD3 zeta chain portion and a costimulatory signaling domain. The costimulatory signaling domain refers to a portion of the CAR comprising the intracellular domain of a costimulatory molecule. A costimulatory molecule is a cell surface molecule other than an antigen receptor or its ligands that is required for an efficient response of lymphocytes to an antigen. Examples of such molecules include CD27, CD28, 4-1BB (CD137), OX40, CD30, CD40, PD-1, ICOS, lymphocyte function-associated antigen-1 (LFA-1), CD2, CD7, LIGHT, NKG2C, B7-H3, and a ligand that specifically binds with CD83, and the like. For example, CD27 costimulation has been demonstrated to enhance expansion, effector function, and survival of human CART cells in vitro and augments human T cell persistence and antitumor activity in vivo (Song et al. Blood. 2012; 119(3):696-706).

[0467] The intracellular signaling sequences within the cytoplasmic portion of a CAR of the invention may be linked to each other in a random or specified order. Optionally, a short oligo- or polypeptide linker, for example, between 2 and 10 amino acids (e.g., 2, 3, 4, 5, 6, 7, 8, 9, or 10 amino acids) in length may form the linkage between intracellular signaling sequence. In one embodiment, a glycine-serine doublet can be used as a suitable linker. In one embodiment, a single amino acid, e.g., an alanine, a glycine, can be used as a suitable linker.

[0468] In one aspect, the intracellular signaling domain is designed to comprise two or more, e.g., 2, 3, 4, 5, or more, costimulatory signaling domains. In an embodiment, the two or more, e.g., 2, 3, 4, 5, or more, costimulatory signaling domains, are separated by a linker molecule, e.g., a linker molecule described herein. In one embodiment, the intracellular signaling domain comprises two costimulatory signaling domains. In some embodiments, the linker molecule is a glycine residue. In some embodiments, the linker is an alanine residue.

[0469] In some embodiments, CAR does not actually recognize the entire antigen; instead it binds to only a portion of the antigen's surface, an area called the antigenic determinant or epitope.

[0470] In some embodiments, a CAR described herein include (including functional portions and functional variants thereof) glycosylated, amidated, carboxylated, phosphorylated, esterified, N-acylated, cyclized via, e.g., a disulfide bridge, or converted into an acid addition salt and/or optionally dimerized or polymerized, or conjugated.

5 **Immunoconjugates**

[0471] In one aspect of the present disclosure, the antibodies or antigen binding fragment thereof, disclosed herein, can initiate a potent immune response against the tumor and/or are capable of direct cytotoxicity. In this regard, the antibodies or antigen binding fragment thereof herein may elicit tumor cell lysis by either complement-mediated or antibody-dependent cell cytotoxicity (ADCC) mechanisms, both of which require
10 an intact Fc portion of the immunoglobulin molecule for interaction with effector cell Fc receptor sites or complement proteins. In addition, antibodies that exert a direct biological effect on tumor growth are useful in the practice of the disclosure. Potential mechanisms by which such directly cytotoxic antibodies may act include inhibition of cell growth, modulation of cellular differentiation, modulation of tumor angiogenesis factor profiles, and the induction of apoptosis. The mechanism by which a particular antibody or an antigen
15 binding fragment thereof, disclosed herein, exerts an anti-tumor effect may be evaluated using any number of in vitro assays designed to determine ADCC, ADMMC, complement-mediated cell lysis, and so forth, as is generally known in the art.

[0472] The antibodies or antigen binding fragment thereof, disclosed herein, may be administered in their "naked" or unconjugated form, or may have therapeutic agents conjugated to them. In one embodiment,
20 antibodies or antigen binding fragment thereof are used as a radiosensitizer. In such embodiments, the antibodies or antigen binding fragment are conjugated to a radiosensitizing agent. The term "radiosensitizer," as used herein, is defined as a molecule, preferably a low molecular weight molecule, administered to animals in therapeutically effective amounts to increase the sensitivity of the cells to be radiosensitized to electromagnetic radiation and/or to promote the treatment of diseases that are treatable
25 with electromagnetic radiation. Diseases that are treatable with electromagnetic radiation include neoplastic diseases, benign and malignant tumors, and cancerous cells.

[0473] The terms "electromagnetic radiation" and "radiation" as used herein include, but are not limited to, radiation having the wavelength of 10-20 to 100 meters. Preferred embodiments of the present disclosure can employ for example, the electro-magnetic radiation of: gamma-radiation c10-20 to 10-13 m), X-ray
30 radiation (10-12 to 10-9 m), ultraviolet light (10 nm to 400 nm), visible light (400 nm to 700 nm), infrared radiation (700 nm to 1.0 mm), and microwave radiation (1 mm to 30 cm).

[0474] Radiosensitizers are known to increase the sensitivity of cancerous cells to the toxic effects of electromagnetic radiation. Many cancer treatment protocols currently employ radiosensitizers activated by the electromagnetic radiation of X-rays. Examples of X-ray activated radiosensitizers include, but are not

limited to, the following: metronidazole, misonidazole, desmethylmisonidazole, pimonidazole, etanidazole, nimorazole, mitomycin C, RSU 1069, SR 4233, E09, RB 6145, nicotinamide, 5-bromode-oxyuridine (BUdR), 5-iododeoxyuridine (IUdR), bromode-oxycytidine, fluorodeoxyuridine (FUdR), hydroxyurea, cisplatin, and therapeutically effective analogs and derivatives of the same.

5 **[0475]** Photodynamic therapy (PDT) of cancers employs visible light as the radiation activator of the sensitizing agent. Examples of photodynamic radiosensitizers include the following, but are not limited to: hematoporphyrin derivatives, Photofrin®, benzoporphyrin derivatives, NPe6, tin etioporphyrin (SnET2), pheoborbide-a, bacteriochlorophyll-a, naphthalocyanines, phthalocyanines, zinc phthalocyanine, and therapeutically effective analogs and derivatives of the same.

10 **[0476]** In another embodiment, the antibody may be conjugated to a receptor (such streptavidin) for utilization in tumor pretargeting wherein the antibody-receptor conjugate is administered to the patient, followed by removal of unbound conjugate from the circulation using a clearing agent and then administration of a ligand (e.g., avidin) which is conjugated to a cytotoxic agent (e.g., a radionuclide).

[0477] The present disclosure further provides the above-described antibodies or antigen binding thereof in
15 detectably labeled form. Antibodies can be detectably labeled through the use of radioisotopes, affinity labels (such as biotin, avidin, etc.), enzymatic labels (such as horseradish peroxidase, alkaline phosphatase, etc.) fluorescent or luminescent or bioluminescent labels (such as FITC or rhodamine, etc.), paramagnetic atoms, and the like. Procedures for accomplishing such labeling are well known in the art; for example, see (Sternberger, L. A. et al., J. Histochem. Cytochem. 18:315 (1970); Bayer, E. A. et al., Meth. Enzym. 62:308
20 (1979); Engval, E. et al., Immunol. 109:129 (1972); Goding, J.W. J. Immunol. Meth. 13:215 (1976)).

[0478] "Label" refers to a detectable compound or composition which is conjugated directly or indirectly to the antibody. The label may itself be detectable by itself (e.g., radioisotope labels or fluorescent labels) or, in the case of an enzymatic label, may catalyze chemical alteration of a substrate compound or composition which is detectable. Alternatively, the label may not be detectable on its own but may be an element that is
25 bound by another agent that is detectable (e.g. an epitope tag or one of a binding partner pair such as biotin-avidin, etc.). Thus, the antibody may comprise a label or tag that facilitates its isolation, and methods of the invention to identify antibodies include a step of isolating the antigen/antibody through interaction with the label or tag.

[0479] Exemplary therapeutic immunoconjugates comprise the antibody described herein conjugated to a
30 cytotoxic agent such as a chemotherapeutic agent, toxin (e.g., an enzymatically active toxin of bacterial, fungal, plant or animal origin, or fragments thereof), or a radioactive isotope (i.e., a radioconjugate). Fusion proteins are described in further detail below.

[0480] In some embodiments, antibodies and antigen binding fragments thereof disclosed herein can be conjugated to a therapeutic agent such as a chemotherapeutic cytotoxin, such as a cytostatic or cytotoxic

agent (e.g., paclitaxol, cytochalasin B or diphtheria toxin, taxol, cytochalasin B, gramicidin D, ethidium bromide, emetine, mitomycin, etoposide, teniposide, vincristine, vinblastine, colchicin, doxorubicin, daunorubicin, dihydroxy anthracindione, mitoxantrone, mithramycin, actinomycin D, 1-dehydrotestosterone, glucocorticoids, procaine, tetracaine, lidocaine, propranolol, and puromycin and analogs or homologs thereof.), antimetabolites (e.g., methotrexate, 6-mercaptopurine, 6-thioguanine, cytarabine, fludarabine, 5-fluorouracil decarbazine), alkylating agents (e.g., mechlorethamine, thioepa chlorambucil, melphalan, carmustine (BSNU) and lomustine (CCNU), cyclophosphamide, busulfan, dibromomannitol, streptozotocin, mitomycin C, and cis-dichlorodiamine platinum (II) (DDP) cisplatin), anthracyclines (e.g., daunorubicin (formerly daunomycin) and doxorubicin), antibiotics (e.g., dactinomycin (formerly actinomycin), bleomycin, mithramycin, and anthramycin (AMC), and anti-mitotic agents, a thrombotic or anti-angiogenic agent or a radioactive label. Examples of suitable cytotoxic agents and chemotherapeutic agents for forming immunoconjugates are known in the art, see for example, WO 05/103081). In another embodiment, antibodies and antigen binding fragments thereof disclosed herein are conjugated to a detectable substrate such as, e.g., an enzyme, fluorescent marker, chemiluminescent marker, bioluminescent material, or radioactive material. In some embodiments of the aspects described herein, the antibody and antibody fragments thereof disclosed herein are conjugated to a toxin (e.g., an enzymatically active toxin of bacterial, fungal, plant or animal origin, or fragments thereof), a small molecule, an siRNA, a nanoparticle, a targeting agent (e.g., a microbubble), or a radioactive isotope (i.e., a radioconjugate). Such conjugates are referred to herein as "immunoconjugates". Such immunoconjugates can be used, for example, in diagnostic, theranostic, or targeting methods.

[0481] Enzymatically active toxins and fragments thereof which can be used include diphtheria A chain, nonbinding active fragments of diphtheria toxin, exotoxin A chain (from *Pseudomonas aeruginosa*), ricin A chain, abrin A chain, modeccin A chain, alpha-sarcin, Aleurites fordii proteins, dianthin proteins, Phytolacca americana proteins (PAPI, PAPII, and PAP-S), momordica charantia inhibitor, curcin, croton, saponaria officinalis inhibitor, gelonin, mitogellin, restrictocin, phenomycin, enomycin and the tricothecenes. A variety of radioisotopes are available for the production of radioconjugate antibodies. Examples include, but are not limited to, ²¹²Bi, ¹³¹I, ¹³¹In, ⁹⁰Y and ¹⁸⁶Re.

[0482] Conjugates of the antibodies or antigen binding fragments thereof described herein and a cytotoxic agent can be made using any of a variety of bifunctional protein coupling agents such as N-succinimidyl-3-(2-pyridyldithiol) propionate (SPDP), iminothiolane (IT), bifunctional derivatives of imidoesters (such as dimethyl adipimidate HCL), active esters (such as disuccinimidyl suberate), aldehydes (such as glutaraldehyde), bis-azido compounds (such as bis (p-azidobenzoyl) hexanediamine), bis-diazonium derivatives (such as bis-(p-diazoniumbenzoyl)-ethylenediamine), diisocyanates (such as tolyene 2,6-diisocyanate), and bis-active fluorine compounds (such as 1,5-difluoro-2,4-dinitrobenzene). For example, a

ricin immunotoxin can be prepared as described in Vitetta et al., 238 Science 1098 (1987). Carbon-14-labeled l-isothiocyanatobenzyl-3-methyldiethylene triaminepentaacetic acid (MX-DTPA) is an exemplary chelating agent for conjugation of radionucleotide to the antibody. See WO 94/11026.

[0483] In other embodiments, the antibody or portion thereof can be conjugated to a "receptor" (e.g., streptavidin) for utilization in tumor pretargeting wherein the antibody-receptor conjugate is administered to the subject, followed by removal of unbound conjugate from the circulation using a clearing agent and then administration of a "ligand" (e.g., avidin) which is conjugated to a cytotoxic agent (e.g. a radionucleotide). In some embodiments, the antibody or antibody fragment thereof can be conjugated to biotin, and the biotin conjugated antibody or antibody fragment thereof can be further conjugated or linked to a streptavidin-bound or-coated agent, such as a streptavidin-coated microbubble, for use in, for example, molecular imaging of angiogenesis.

[0484] Techniques for conjugating such therapeutic moiety to antibodies are well known, see, e.g., Amon et al., "Monoclonal Antibodies For Immunotargeting Of Drugs In Cancer Therapy", in Monoclonal Antibodies And Cancer Therapy, Reisfeld et al. (eds.), pp. 243-56 (Alan R. Liss, Inc. 1985); Hellstrom et al., "Antibodies For Drug Delivery", in Controlled Drug Delivery (2nd Ed.), Robinson et al. (eds.), pp. 623-53 (Marcel Dekker, Inc. 1987); Thorpe, "Antibody Carriers Of Cytotoxic Agents In Cancer Therapy: A Review", in Monoclonal Antibodies '84: Biological And Clinical Applications, Pinchera et al. (eds.), pp. 475-506 (1985); "Analysis, Results, And Future Prospective Of The Therapeutic Use Of Radiolabeled Antibody In Cancer Therapy", in Monoclonal Antibodies For Cancer Detection And Therapy, Baldwin et al. (eds.), pp. 303-16 (Academic Press 1985), and Thorpe et al., "The Preparation And Cytotoxic Properties Of Antibody-Toxin Conjugates", Immunol. Rev., 62: 119-58 (1982).

[0485] Production of immunoconjugates is described in U.S. Pat. No. 6,306,393. Immunoconjugates can be prepared by indirectly conjugating a therapeutic agent to an antibody component. General techniques are described in Shih et al., Int. J. Cancer 41 :832-839 (1988); Shih et al., Int.J. Cancer 46:1101-1106 (1990); and Shih et al., U.S. Pat. No. 5,057,313. The general method involves reacting an antibody component having an oxidized carbohydrate portion with a carrier polymer that has at least one free amine function and that is loaded with a plurality of drug, toxin, chelator, boron addends, or other therapeutic agent. This reaction results in an initial Schiff base (imine) linkage, which can be stabilized by reduction to a secondary amine to form the final conjugate.

[0486] The carrier polymer is preferably an aminodextran or polypeptide of at least 50 amino acid residues, although other substantially equivalent polymer carriers can also be used. Preferably, the final immunoconjugate is soluble in an aqueous solution, such as mammalian serum, for ease of administration and effective targeting for use in therapy. Thus, solubilizing functions on the carrier polymer will enhance the serum solubility of the final immunoconjugate. In particular, an aminodextran will be preferred.

[0487] The process for preparing an immunoconjugate with an aminodextran carrier typically begins with a dextran polymer, advantageously a dextran of average molecular weight of about 10,000-100,000. The dextran is reacted with an oxidizing agent to affect a controlled oxidation of a portion of its carbohydrate rings to generate aldehyde groups. The oxidation is conveniently affected with glycolytic chemical reagents such as NaIO₄, according to conventional procedures.

[0488] The oxidized dextran is then reacted with a polyamine, preferably a diamine, and more preferably, a mono- or polyhydroxy diamine. Suitable amines include ethylene diamine, propylene diamine, or other like polymethylene diamines, diethylene triamine or like polyamines, 1,3-diamino-2-hydroxypropane, or other like hydroxylated diamines or polyamines, and the like. An excess of the amine relative to the aldehyde groups of the dextran is used to ensure substantially complete conversion of the aldehyde functions to Schiff base groups.

[0489] A reducing agent, such as NaBH₄, NaBH₃CN or the like, is used to effect reductive stabilization of the resultant Schiff base intermediate. The resultant adduct can be purified by passage through a conventional sizing column to remove cross-linked dextrans. Other conventional methods of derivatizing a dextran to introduce amine functions can also be used, e.g., reaction with cyanogen bromide, followed by reaction with a diamine. The aminodextran is then reacted with a derivative of the particular drug, toxin, chelator, immunomodulator, boron addend, or other therapeutic agent to be loaded, in an activated form, preferably, a carboxyl-activated derivative, prepared by conventional means, e.g., using dicyclohexylcarbodiimide (DCC) or a water soluble variant thereof, to form an intermediate adduct.

[0490] Alternatively, polypeptide toxins such as poke- weed antiviral protein or ricin A-chain, and the like, can be coupled to aminodextran by glutaraldehyde condensation or by reaction of activated carboxyl groups on the protein with amines on the aminodextran. Chelators for radiometals or magnetic resonance enhancers are well-known in the art. Typical are derivatives of ethylenediaminetetraacetic acid (EDTA) and diethylenetriaminepentaacetic acid (DTPA). These chelators typically have groups on the side chain by which the chelator can be attached to a carrier. Such groups include, e.g., benzylisothiocyanate, by which the DTPA or EDTA can be coupled to the amine group of a carrier. Alternatively, carboxyl groups or amine groups on a chelator can be coupled to a carrier by activation or prior derivatization and then coupling, all by well-known means.

[0491] Boron addends, such as carboranes, can be attached to antibody components by conventional methods. For example, carboranes can be prepared with carboxyl functions on pendant side chains, as is well known in the art. Attachment of such carboranes to a carrier, e.g., aminodextran, can be achieved by activation of the carboxyl groups of the carboranes and condensation with amines on the carrier to produce an intermediate conjugate. Such intermediate conjugates are then attached to antibody components to produce therapeutically useful immunoconjugates, as described below.

[0492] A polypeptide carrier can be used instead of aminodextran, but the polypeptide carrier should have at least 50 amino acid residues in the chain, preferably 100-5000 amino acid residues. At least some of the amino acids should be lysine residues or glutamate or aspartate residues. The pendant amines of lysine residues and pendant carboxylates of glutamine and aspartate are convenient for attaching a drug, toxin, immunomodulator, chelator, boron addend or other therapeutic agent. Examples of suitable polypeptide carriers include polylysine, polyglutamic acid, polyaspartic acid, co-polymers thereof, and mixed polymers of these amino acids and others, e.g., serines, to confer desirable solubility properties on the resultant loaded carrier and immunoconjugate.

[0493] Conjugation of the intermediate conjugate with the antibody component is effected by oxidizing the carbohydrate portion of the antibody component and reacting the resulting aldehyde (and ketone) carbonyls with amine groups remaining on the carrier after loading with a drug, toxin, chelator, immunomodulator, boron addend, or other therapeutic agent. Alternatively, an intermediate conjugate can be attached to an oxidized antibody component via amine groups that have been introduced in the intermediate conjugate after loading with the therapeutic agent. Oxidation is conveniently effected either chemically, e.g., with NaIO₄ or other glycolytic reagent, or enzymatically, e.g., with neuraminidase and galactose oxidase. In the case of an aminodextran carrier, not all of the amines of the aminodextran are typically used for loading a therapeutic agent. The remaining amines of aminodextran condense with the oxidized antibody component to form Schiff base adducts, which are then reductively stabilized, normally with a borohydride reducing agent.

[0494] Analogous procedures are used to produce other immunoconjugates according to the invention. Loaded polypeptide carriers preferably have free lysine residues remaining for condensation with the oxidized carbohydrate portion of an antibody component. Carboxyls on the polypeptide carrier can, if necessary, be converted to amines by, e.g., activation with DCC and reaction with an excess of a diamine.

[0495] The final immunoconjugate is purified using conventional techniques, such as sizing chromatography on Sephacryl S-300 or affinity chromatography using one or more CD84Hy epitopes. Alternatively, immunoconjugates can be prepared by directly conjugating an antibody component with a therapeutic agent. The general procedure is analogous to the indirect method of conjugation except that a therapeutic agent is directly attached to an oxidized antibody component. It will be appreciated that other therapeutic agents can be substituted for the chelators described herein. Those of skill in the art will be able to devise conjugation schemes without undue experimentation.

[0496] As a further illustration, a therapeutic agent can be attached at the hinge region of a reduced antibody component via disulfide bond formation. For example, the tetanus toxoid peptides can be constructed with a single cysteine residue that is used to attach the peptide to an antibody component. As an alternative, such peptides can be attached to the antibody component using a heterobifunctional cross-

linker, such as N-succinyl 3-(2-pyridyldithio)propionate (SPDP). Yu et al., *Int. J. Cancer* 56:244 (1994). General techniques for such conjugation are well-known in the art. See, for example, Wong, *Chemistry Of Protein Conjugation and Cross-Linking* (CRC Press 1991); Upeslakis et al., "Modification of Antibodies by Chemical Methods," in *Monoclonal Antibodies: Principles and Applications*, Birch et al. (eds.), pages 187-230 (Wiley-Liss, Inc. 1995); Price, "Production and Characterization of Synthetic Peptide-Derived Antibodies," in *Monoclonal Antibodies: Production, Engineering and Clinical Application*, Ritter et al. (eds.), pages 60-84 (Cambridge University Press 1995).

[0497] Conjugates of the antibody and cytotoxic agent are made using a variety of bifunctional protein coupling agents such as N-succinimidyl-3-(2-pyridyldithiol) propionate (SPDP), iminothiolane (IT), bifunctional derivatives of imidoesters (such as dimethyl adipimidate HCL), active esters (such as disuccinimidyl suberate), aldehydes (such as glutaraldehyde), bis-azido compounds (such as bis (p-azido-benzoyl) hexanediamine), bis-diazonium derivatives (such as bis-(p-diazoniumbenzoyl)-ethylenediamine), diisocyanates (such as tolyene 2,6-diisocyanate), and bis-active fluorine compounds (such as 1,5-difluoro-2,4-dinitrobenzene). For example, a ricin immunotoxin can be prepared as described in Vitetta et al., *Science* 238: 1098 (1987). Carbon-14-labeled 1-isothiocyanatobenzyl-3-methyl diethylene triaminepentaacetic acid (MX-DTPA) is an exemplary chelating agent for conjugation of radionuclide to the antibody (see, e.g., W094/11026).

[0498] As described above, carbohydrate moieties in the Fc region of an antibody can be used to conjugate a therapeutic agent. However, the Fc region may be absent if an antibody fragment is used as the antibody component of the immunoconjugate. Nevertheless, it is possible to introduce a carbohydrate moiety into the light chain variable region of an antibody or antibody fragment. See, for example, Leung et al., *J. Immunol.* 154:5919 (1995); Hansen et al., U.S. Pat. No. 5,443,953. The engineered carbohydrate moiety is then used to attach a therapeutic agent. **[0404]** In addition, those of skill in the art will recognize numerous possible variations of the conjugation methods. For example, the carbohydrate moiety can be used to attach polyethyleneglycol in order to extend the half-life of an intact antibody, or antigen-binding fragment thereof, in blood, lymph, or other extracellular fluids. Moreover, it is possible to construct a "divalent immunoconjugate" by attaching therapeutic agents to a carbohydrate moiety and to a free sulfhydryl group. Such a free sulfhydryl group may be located in the hinge region of the antibody component.

[0499] In some embodiments, provided herein is a method for diagnosing or treating a condition associated with a target antigen. In some embodiments, the condition associated with the antigen results due to an increase in the expression or biological activity of the antigen. In some embodiments, the condition associated with the antigen results due to a decrease in the expression or biological activity of the antigen. In some embodiments, the antibody or antigen-binding fragment thereof disclosed herein binds an antigen

and, thereby, partially or substantially inhibits the expression and/or at least one biological activity of the antigen. An antibody, or specified portion or variant thereof, that partially or preferably substantially inhibits the expression and/or at least one biological activity of an antigen disclosed herein can bind the protein antigen or fragment thereof and thereby inhibit activities mediated through the antigen e.g., binding of antigen to one or more receptor or ligand known in the art to bind a particular antigen. In some 5 embodiments, the antibody can inhibit an antigen activity by about 20-120%, preferably by at least about 10, 20, 30, 40, 50, 55, 60, 65, 70, 75, 80, 85, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100% or more. An antibody, or specified portion or variant thereof, that partially or preferably substantially increases the expression and/or at least one biological activity of an antigen disclosed herein can bind the protein antigen or fragment thereof and thereby increase activities mediated through the antigen. In some embodiments, the 10 antibody can increase an antigen activity by about 20-120%, preferably by at least about 10, 20, 30, 40, 50, 55, 60, 65, 70, 75, 80, 85, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100% or more.

[0500] The capacity of an antibody to increase an antigen-dependent activity is preferably assessed by at least one suitable assay, as described herein and/or as known in the art. In some embodiments, the antigen 15 related condition can be an immune related disease, a cardiovascular disease, an infectious disease, a malignant disease or a neurologic disease.

Methods of Treatment

[0501] Disclosed herein, in some embodiments, are methods of treating a subject in need thereof, comprising administering to the subject, a therapeutic dose of a therapeutic IgA antibody described herein 20 or a pharmaceutical composition comprising said therapeutic IgA antibody described herein. In some embodiments, the subject has a cancer, an infectious disease, or an autoimmune disease.

[0502] In some embodiments, the subject has a cancer. In some embodiments, the subject has an inflammatory disorder. In some embodiments, the cancer is associated with expression of a tumor associated antigen described herein. In some embodiments, the cancer is associated with expression of 25 CD47, CD20, GD2, CD38, CD19, EGFR, HER2, PD-L1, CD25, CD33, BCMA, CD44, α -Folate receptor, CAIX, CD30, ROR1, CEA, EGP-2, EGP-40, HER3, Folate-binding Protein, GD3, IL-13R- α 2, KDR, EDB-F, mesothelin, CD22, EGFR, MUC-1, MAGE-A1, MUC16, h5T4, PSMA, TAG-72, EGFRvIII, CD123 or VEGF-R2.

[0503] In some embodiments, disclosed herein are methods of administering a therapeutic IgA antibody or pharmaceutical composition described herein, to a subject having a cancer associated with an 30 overexpression of CD20. In some embodiments, disclosed herein are methods of administering a therapeutic IgA antibody or pharmaceutical composition described herein to a subject having a cancer associated with an overexpression of GD2. In some embodiments, disclosed herein are methods of administering a therapeutic IgA antibody or pharmaceutical composition described herein to a subject having a cancer

associated with an overexpression of mesothelin. In some embodiments, disclosed herein are methods of administering a modified effector cell to a subject having a cancer associated with an overexpression of CD38, CD19, EGFR, HER2, PD-L1, CD25, CD33, BCMA, CD44, α -Folate receptor, CAIX, CD30, ROR1, CEA, EGP-2, EGP-40, HER3, Folate-binding Protein, GD2, GD3, IL-13R-a2, KDR, EDB-F, mesothelin, CD22, EGFR, MUC-1, MAGE-A1, MUC16, h5T4, PSMA, TAG-72, EGFRvIII, CD123 or VEGF-R2.

[0504] In some embodiments, the cancer is a metastatic cancer. In other embodiments, the cancer is a relapsed or refractory cancer. In some embodiments, a cancer is a solid tumor or a hematologic malignancy. In some embodiments, the cancer is a solid tumor. In other embodiments, the cancer is a hematologic malignancy. In some embodiments, the cancer is a metastatic cancer. In some embodiments, the cancer is a relapsed or refractory cancer.

[0505] In some embodiments, the cancer is a solid tumor. Exemplary solid tumors include, but are not limited to, anal cancer; appendix cancer; bile duct cancer (*i.e.*, cholangiocarcinoma); bladder cancer; brain tumor; breast cancer; cervical cancer; colon cancer; cancer of Unknown Primary (CUP); esophageal cancer; eye cancer; fallopian tube cancer; gastroenterological cancer; kidney cancer; liver cancer; lung cancer; medulloblastoma; melanoma; oral cancer; ovarian cancer; pancreatic cancer; parathyroid disease; penile cancer; pituitary tumor; prostate cancer; rectal cancer; skin cancer; stomach cancer; testicular cancer; throat cancer; thyroid cancer; uterine cancer; vaginal cancer; vulvar cancer; or glioblastoma.

[0506] In some embodiments, the cancer is a hematologic malignancy. In some embodiments, a hematologic malignancy comprises a lymphoma, a leukemia, a myeloma, or a B-cell malignancy. In some embodiments, a hematologic malignancy comprises a lymphoma, a leukemia or a myeloma. In some embodiments, exemplary hematologic malignancies include chronic lymphocytic leukemia (CLL), small lymphocytic lymphoma (SLL), high risk CLL, non-CLL/SLL lymphoma, prolymphocytic leukemia (PLL), follicular lymphoma (FL), diffuse large B-cell lymphoma (DLBCL), mantle cell lymphoma (MCL), Waldenstrom's macroglobulinemia, multiple myeloma, extranodal marginal zone B cell lymphoma, nodal marginal zone B cell lymphoma, Burkitt's lymphoma, non-Burkitt high grade B cell lymphoma, primary mediastinal B-cell lymphoma (PMBL), immunoblastic large cell lymphoma, precursor B-lymphoblastic lymphoma, B cell prolymphocytic leukemia, lymphoplasmacytic lymphoma, splenic marginal zone lymphoma, plasma cell myeloma, plasmacytoma, mediastinal (thymic) large B cell lymphoma, intravascular large B cell lymphoma, primary effusion lymphoma, or lymphomatoid granulomatosis. In some embodiments, the hematologic malignancy comprises a myeloid leukemia. In some embodiments, the hematologic malignancy comprises acute myeloid leukemia (AML) or chronic myeloid leukemia (CML).

[0507] In some embodiments, the antibody or fragment thereof binds one or more of G_{D2}, ALK, hNET, G_{D3}, and CD20. In some embodiments, the G_{D2} positive tumor is neuroblastoma, retinoblastoma, melanoma, small cell lung cancer, glioblastoma, osteosarcoma, rhabdomyosarcoma, Ewing's sarcoma, liposarcoma,

fibrosarcoma, leiomyosarcoma, and any combinations thereof. In an aspect, the antibody or fragment thereof binds a neuroblastoma cell. In some embodiments, the ALK (anaplastic lymphoma kinase) positive tumor is an anaplastic large-cell lymphoma, an adenocarcinoma of the lung, a neuroblastoma, an inflammatory myofibroblastic tumor, a renal cell carcinomas, esophageal squamous cell carcinoma, breast cancer, a colonic adenocarcinoma, a glioblastoma multiforme or an anaplastic thyroid cancer. In some embodiments, the hNET (human norepinephrine transporter) positive tumor is a bladder tumor, breast tumor, prostate tumor, carcinoma, leukemia, liver cancer, lung cancer, lymphoma, Hodgkin's lymphoma, Non-Hodgkin's lymphoma, melanoma, neuroblastoma, ovarian tumor, pancreatic tumor or a retinoblastoma. In some case, the G_{D3} positive tumor is a neuroectodermal tumor of the center nervous system, glioma, neuroblastoma, retinoblastoma, ependymoma, sarcoma, melanoma, breast cancer, ovarian cancer, glioblastoma, Ewing's sarcoma, or small cell lung carcinoma. In some embodiments the CD20 positive tumor is a leukemia, a lymphoma or a neuroblastoma.

[0508] In other embodiments, disclosed herein are methods of administering to a subject having an infection due to an infectious disease. An infectious disease can be a disease resulting from a bacterial, viral or fungi infection. In other embodiments, exemplary viral pathogens include those of the families of Adenoviridae, Epstein-Barr virus (EBV), Cytomegalovirus (CMV), Respiratory Syncytial Virus (RSV), JC virus, BK virus, HSV, HHV family of viruses, Picornaviridae, Herpesviridae, Hepadnaviridae, Flaviviridae, Retroviridae, Orthomyxoviridae, Paramyxoviridae, Papovaviridae, Polyomavirus, Rhabdoviridae, and Togaviridae. Exemplary pathogenic viruses cause smallpox, influenza, mumps, measles, chickenpox, ebola, and rubella. Exemplary pathogenic fungi include Candida, Aspergillus, Cryptococcus, Histoplasma, Pneumocystis, and Stachybotrys. Exemplary pathogenic bacteria include Streptococcus, Pseudomonas, Shigella, Campylobacter, Staphylococcus, Helicobacter, E. coli, Rickettsia, Bacillus, Bordetella, Chlamydia, Spirochetes, and Salmonella.

[0509] In some embodiments, the IgA antibody is administered with one or more additional therapeutic agents. In some embodiments, the IgA antibody and one or more additional therapeutic agents are co-administered. In some embodiments, the IgA antibody and one or more additional therapeutic agents are sequentially administered.

[0510] In some embodiments, the combination therapy can include one or more antibodies of the disclosure co-formulated with, and/or co-administered with, one or more additional therapeutic agents, e.g., chemotherapeutic or anti-neoplastic agents, such as cytokine and growth factor inhibitors, immunosuppressants, anti-inflammatory agents, metabolic inhibitors, enzyme inhibitors, and/or cytotoxic or cytostatic agents. The term "combination" in this context means that the agents are given substantially contemporaneously, either simultaneously or sequentially. Exemplary chemotherapeutic agents include, but are not limited to, aldesleukin, altretamine, amifostine, asparaginase, bleomycin, capecitabine, carboplatin,

carmustine, cladribine, cisapride, cisplatin, cyclophosphamide, cytarabine, dacarbazine (DTIC), dactinomycin, docetaxel, doxorubicin, dronabinol, duocarmycin, etoposide, filgrastim, fludarabine, fluorouracil, gemcitabine, granisetron, hydroxyurea, idarubicin, ifosfamide, interferon alpha, irinotecan, lansoprazole, levamisole, leucovorin, megestrol, mesna, methotrexate, metoclopramide, mitomycin, mitotane, mitoxantrone, omeprazole, ondansetron, paclitaxel (Taxol™), pilocarpine, prochloroperazine, saproin, tamoxifen, taxol, topotecan hydrochloride, vinblastine, vincristine and vinorelbine tartrate.

[0511] In some embodiments, the IgA antibodies described herein can be combined with an effective dose of other antibodies that have been used in treatment of cancer including, without limitation the following FDA approved monoclonal antibodies: rituximab (Rituxan®, CD20: chimeric IgG1), trastuzumab (Herceptin®, HER2: chimeric IgG1), alemtuzumab (Campath®, CD52: humanized IgG1), ibritumomab tiuxetan (Zevalin®, CD20: murine, IgG1, radiolabeled (Yttrium 90)), tositumomab-I-131 (Bexxar®: CD20, murine, IgG2a, radiolabeled (Iodine 131)), cetuximab (Erbix®, EGFR: chimeric, IgG1), bevacizumab (VEGF: humanized, IgG4), panitumumab (Vectibix®, EGFR: human IgG2), ofatumumab (Arzerra®, CD20: human IgG1), ipilimumab (Ypervoy®, CTLA-4: human IgG1), brentuximab vedotin (Adectris®, CD30: chimeric, IgG1, drug-conjugate), pertuzumab (Perjeta®, HER2: humanized IgG1, drug conjugate), adotrastuzumab ematansine (Kadcyla®, HER2: humanized, IgG1, drug-conjugate), obinutuzumab (Gazyva®, CD20: humanized and glycol-engineered), nivolumab and pembrolizumab (anti-PD-1s), etc. Trastuzumab targets the HER-2 antigen.

[0512] In some embodiments, the anti-CD47 IgA antibody is administered in combination with a HER2 inhibitor, an anti-HER2 antibody, an EGFR inhibitor, an anti-EGFR antibody. In some embodiments, the anti-CD47 IgA antibody is administered in combination with trastuzumab. In some embodiments, the anti-CD47 IgA antibody is administered in combination with a c-kit inhibitor. Dosages

[0513] Provided herein are compositions comprising IgA antibodies or antigen binding fragment thereof for treatment (including prevention) of a disease (e.g., cancer). In some embodiments, the compositions are pharmaceutical compositions comprising a pharmaceutically acceptable carrier. The compositions are administered in an amount effective for treatment (including prophylaxis) of cancer. In some embodiments, the compositions (e.g., the antibodies or the antigen binding fragment thereof or the nucleic acid molecules encoding said antibody or antigen binding fragment thereof) are administered in an amount effective for enhancing an immune response and/or increasing T cell activation in a subject. The compositions are to be used for in vivo administration to a subject by any available means, such as parenteral administration. For administration to a subject, a composition or medicament comprising the antibodies or antigen binding fragment thereof described herein can be sterile, which can readily accomplished by filtration through sterile filtration membranes, or other methods known to those of skill in the art. In one embodiment, a composition or medicament has been treated to be free of pyrogens or endotoxins. Testing pharmaceutical compositions

or medicaments for pyrogens or endotoxins and preparing pharmaceutical compositions or medicaments free of pyrogens or endotoxins or preparing pharmaceutical compositions or medicaments that have endotoxins at a clinically-acceptable level, are well understood to one of ordinary skill in the art. Commercial kits are available to test pharmaceutical compositions or medicaments for pyrogens or endotoxins.

- 5 [0514] The compositions to be used for in vivo administration, such as parenteral administration, in the methods described herein can be sterile, which is readily accomplished by filtration through sterile filtration membranes, or other methods known to those of skill in the art.

- [0515] The IgA antibodies or antigen binding fragments thereof, describe herein, are formulated, dosed, and administered in a fashion consistent with good medical practice. Factors for consideration in this context
10 include the particular disorder being treated, the particular subject being treated, the clinical condition of the individual subject, the cause of the disorder, the site of delivery of the agent, the method of administration, the scheduling of administration, and other factors known to medical practitioners. A "therapeutically effective amount" of a substance/molecule, agonist or antagonist may vary according to factors such as the disease state, age, sex, and weight of the individual, and the ability of the substance/molecule, agonist or
15 antagonist to elicit a desired response in the individual. A therapeutically effective amount is also one in which any toxic or detrimental effects of the substance/molecule, agonist or antagonist are outweighed by the therapeutically beneficial effects. A therapeutically effective amount may be delivered in one or more administrations. A therapeutically effective amount refers to an amount effective, at dosages and for periods of time necessary, to achieve the desired therapeutic and/or prophylactic result. The "therapeutically
20 effective amount" to be administered will be governed by such considerations, and refers to the minimum amount necessary to ameliorate, treat, or stabilize, the cancer; to increase the time until progression (duration of progression free survival) or to treat or prevent the occurrence or recurrence of a tumor, a dormant tumor, or a micrometastases. The antibodies or antigen binding fragment thereof, disclosed herein, is optionally formulated with one or more additional therapeutic agents currently used to prevent or treat cancer or a risk
25 of developing a cancer. The effective amount of such other agents depends on the amount of antibody or antigen binding fragment thereof present in the formulation, the type of disorder or treatment, and other factors discussed above. These are generally used in the same dosages and with administration routes as used herein before or about from 1 to 99% of the heretofore employed dosage.

- [0516] The dose of antibody may vary depending upon the age and the size of a subject to be administered,
30 target disease, conditions, route of administration, and the like. The preferred dose is typically calculated according to body weight or body surface area. When an antibody or antigen binding fragment thereof disclosed herein is used for treating a condition or disease in an adult patient, it may be advantageous to intravenously administer the antibody of the present invention normally at a single dose of about 0.01 to about 20 mg/kg body weight, more preferably about 0.02 to about 7, about 0.03 to about 5, or about 0.05 to

about 3 mg/kg, about 5 mg/kg, about 7.5 mg/kg, about 10 mg/kg, or about 15 mg/kg body weight. Depending on the severity of the condition, the frequency and the duration of the treatment can be adjusted. Effective dosages and schedules for administering may be determined empirically; for example, patient progress can be monitored by periodic assessment, and the dose adjusted accordingly. Moreover, interspecies scaling of dosages can be performed using well-known methods in the art (e.g., Mordenti et al., 1991, *Pharmaceut. Res.* 8:1351).

[0517] In some embodiments, the compositions herein can comprise a prophylactically effective amount, e.g., when administering to a subject at a risk of cancer or in earlier stages of a disease. A “prophylactically effective amount” refers to an amount effective, at dosages and for periods of time necessary, to achieve the desired prophylactic result. Typically, since a prophylactic dose is used in subjects prior to or at an earlier stage of disease, the prophylactic dose is lower than the therapeutic dose.

[0518] Advantageously, the pharmaceutical compositions for oral or parenteral use described above are prepared into dosage forms in a unit dose suited to fit a dose of the active ingredients. Such dosage forms in a unit dose include, for example, tablets, pills, capsules, injections (ampoules), suppositories, etc.

[0519] The administration can be, for example, by one or more separate administrations, or by continuous infusion. For repeated administrations over several days or longer, depending on the condition, the treatment is sustained until, for example, the cancer is treated, as measured by the methods known in the art. However, other dosage regimens can be useful. In one non-limiting example, an antibody or antigen binding fragment thereof, disclosed herein is administered once every week, every two weeks, or every three weeks, at a dose range from about 5 mg/kg to about 15 mg/kg, including but not limited to 5 mg/kg, 7.5 mg/kg, 10 mg/kg or 15 mg/kg. The progress of using the methods described herein can be easily monitored by conventional techniques and assays. The duration of a therapy using the methods described herein will continue for as long as medically indicated or until a desired therapeutic effect (e.g., those described herein) is achieved. In certain embodiments, the administration of one or more antibodies or antigen binding fragment thereof, or compositions, described herein, is continued for 1 month, 2 months, 4 months, 6 months, 8 months, 10 months, 1 year, 2 years, 3 years, 4 years, 5 years, 10 years, 20 years, or for a period of years up to the lifetime of the subject.

Efficacy of treatment

[0520] The efficacy of the treatment methods for example, for cancer, comprising administering the IgA antibodies or antigen binding fragment thereof, or pharmaceutical compositions of the present disclosure can be measured by various endpoints commonly used in evaluating cancer treatments, including but not limited to, tumor regression, tumor weight or size shrinkage, time to progression, duration of survival, progression free survival, overall response rate, duration of response, and quality of life. The antibodies or antigen

binding fragments thereof disclosed herein can require unique measures and definitions of clinical responses to drugs. In the case of cancers, the therapeutically effective amount of the antibodies, antigen binding fragments thereof disclosed herein or compositions comprising the same can reduce the number of cancer cells; reduce the tumor size; inhibit (i.e., slow to some extent and preferably stop) cancer cell infiltration into peripheral organs; inhibit (i.e., slow to some extent and preferably stop) tumor metastasis; inhibit, to some extent, tumor growth; and/or relieve to some extent one or more of the symptoms associated with the disorder. To the extent the antibodies or antigen binding fragment thereof, disclosed herein, act to prevent growth and/or kill existing cancer cells; it can be cytostatic and/or cytotoxic. For cancer therapy, efficacy in vivo can, for example, be measured by assessing the duration of survival, duration of progression free survival (PFS), the response rates (RR), duration of response, and/or quality of life. In some embodiments, the IgA antibodies or a functional fragment thereof disclosed herein inhibits tumor growth by at least about 2%, 3%, 5%, 6%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 60%, 70%, 80%, 90% or more relative to an untreated subject. In some embodiments, the IgA antibodies or a functional fragment thereof disclosed herein inhibits tumor engraftment by at least about 2%, 3%, 5%, 6%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 60%, 70%, 80%, 90% or more relative to an untreated subject. In some embodiments, the IgA antibodies or a functional fragment thereof disclosed herein induce cytolysis of tumor cell. In some embodiments, the IgA antibodies or a functional fragment thereof disclosed herein induce increased cytolysis of tumor cell by at least about 2%, 3%, 5%, 6%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 60%, 70%, 80%, 90% or more relative to a corresponding WT IgA.

[0521] In other embodiments, described herein are methods for increasing progression free survival of a human subject susceptible to or diagnosed with a cancer, for example, skin cancer, such as cutaneous melanoma. Time to disease progression is defined as the time from administration of the drug until disease progression or death. In a preferred embodiment, the combination treatment of the invention using an antibody or antigen binding fragment thereof, disclosed herein, and one or more chemotherapeutic agents may significantly increase progression free survival by at least about 1 month, 1.2 months, 2 months, 2.4 months, 2.9 months, 3.5 months, such as by about 1 to about 5 months, when compared to a treatment with chemotherapy alone. In another embodiment, the methods described herein may significantly increase response rates in a group of human subjects susceptible to or diagnosed with a cancer that are treated with various therapeutics. Response rate is defined as the percentage of treated subjects who responded to the treatment. In one embodiment, the combination treatment described herein using an antibody or antigen binding fragment thereof, disclosed herein, such as a recombinant antibody or antigen binding fragment thereof, and one or more chemotherapeutic agents significantly increases response rate in the treated subject group compared to the group treated with chemotherapy alone.

- [0522] As used herein, the terms "treat," "treatment," "treating," or "amelioration" refer to therapeutic treatments, wherein the object is to reverse, alleviate, ameliorate, inhibit, slow down or stop the progression or severity of a condition associated with, a disease or disorder. The term "treating" includes reducing or alleviating at least one adverse effect or symptom of a condition, disease or disorder associated with a chronic immune condition, such as, but not limited to, a chronic infection or a cancer. Treatment is generally "effective" if one or more symptoms or clinical markers are reduced. Alternatively, treatment is "effective" if the progression of a disease is reduced or halted. That is, "treatment" includes not just the improvement of symptoms or markers, but also a cessation of at least slowing of progress or worsening of symptoms that would be expected in absence of treatment. Beneficial or desired clinical results include, but are not limited to, alleviation of one or more symptom(s), diminishment of extent of disease, stabilized (i.e., not worsening) state of disease, delay or slowing of disease progression, amelioration or palliation of the disease state, and remission (whether partial or total), whether detectable or undetectable. The term "treatment" of a disease also includes providing relief from the symptoms or side-effects of the disease (including palliative treatment).
- [0523] For example, in some embodiments, the methods described herein comprise administering an effective amount of the antibodies or antigen binding fragment thereof, described herein, to a subject in order to alleviate a symptom of a disease, for example, cancer. As used herein, "alleviating a symptom of a cancer" is ameliorating or reducing any condition or symptom associated with a disease. As compared with an equivalent untreated control, such reduction or degree of prevention is at least 5%, 10%, 20%, 40%, 50%, 60%, 80%, 90%, 95%, or 100% as measured by any standard technique. Ideally, the cancer is completely cleared as detected by any standard method known in the art, in which case the cancer is considered to have been treated. A patient who is being treated for a cancer is one who a medical practitioner has diagnosed as having such a condition. Diagnosis can be by any suitable means. Diagnosis and monitoring can involve, for example, detecting the level of cancer cells in a biological sample (for example, a tissue or lymph node biopsy, blood test, or urine test), detecting the level of a surrogate marker of the cancer in a biological sample, detecting symptoms associated with the specific cancer, or detecting immune cells involved in the immune response typical of such a cancer.
- [0524] The term "effective amount" as used herein refers to the amount of an antibody or antigen binding fragment thereof or composition comprising the same needed to alleviate at least one or more symptom of the disease or disorder, and relates to a sufficient amount of pharmacological composition to provide the desired effect. The term "therapeutically effective amount" therefore refers to an amount of an antibody or antigen binding fragment thereof disclosed herein, that is sufficient to effect a particular effect when administered to a typical subject. An effective amount as used herein would also include an amount sufficient to delay the development of a symptom of the disease, alter the course of a symptom disease (for

example but not limited to, slow the progression of a symptom of the disease), or reverse a symptom of the disease. Thus, it is not possible to specify the exact "effective amount". For any given case, however, an appropriate "effective amount" can be determined by one of ordinary skill in the art using only routine experimentation.

- 5 [0525] Effective amounts, toxicity, and therapeutic efficacy can be determined by standard pharmaceutical procedures in cell cultures or experimental animals, e.g., for determining the LD50 (the dose lethal to 50% of the population) and the ED50 (the dose therapeutically effective in 50% of the population). The dosage can vary depending upon the dosage form employed and the route of administration utilized. The dose ratio between toxic and therapeutic effects is the therapeutic index and can be expressed as the ratio LD50/ED50-
- 10 Compositions and methods that exhibit large therapeutic indices are preferred. A therapeutically effective dose can be estimated initially from cell culture assays. Also, a dose can be formulated in animal models to achieve a circulating plasma concentration range that includes the IC50 (i.e., the concentration of the antibody or antigen binding fragment thereof), which achieves a half-maximal inhibition of symptoms as determined in cell culture, or in an appropriate animal model. Levels in plasma can be measured, for
- 15 example, by high performance liquid chromatography. The effects of any particular dosage can be monitored by a suitable bioassay. The dosage can be determined by a physician and adjusted, as necessary, to suit observed effects of the treatment.

- [0526] The treatment and/or prevention of cancer includes, but is not limited to, alleviating symptoms associated with cancer, the inhibition of the progression of cancer, the promotion of the regression of cancer,
- 20 the promotion of the immune response, inhibition of tumor growth, inhibition of tumor size, inhibition of metastasis, inhibition of cancer cell growth, inhibition of cancer cell proliferation, or cause cancer cell death.

Modes of administration

- [0527] The IgA antibodies or antigen binding fragment thereof, described herein, can be administered to a subject in need thereof by any appropriate route which results in an effective treatment in the subject. As
- 25 used herein, the terms "administering," and "introducing" are used interchangeably and refer to the placement of an antibody or antibody portion thereof into a subject by a method or route which results in at least partial localization of such agents at a desired site, such as a site of inflammation or cancer, such that a desired effect(s) is produced.

- [0528] In some embodiments, the antibodies or antigen binding fragment thereof, described herein, or
- 30 compositions comprising the same is administered to a subject having a cancer, to be inhibited by any mode of administration that delivers the agent systemically or to a desired surface or target, and can include, but is not limited to, injection, infusion, instillation, and inhalation administration. Oral administration forms are also contemplated herein. "Injection" includes, without limitation, intravenous, intramuscular, intraarterial, intrathecal, intraventricular, intracapsular, intraorbital, intracardiac, intradermal, intraperitoneal,

transtracheal, subcutaneous, subcuticular, intraarticular, sub capsular, subarachnoid, intracranial, intraspinal, intracerebro spinal, and intrasternal injection and infusion.

[0529] The phrases "parenteral administration" and "administered parenterally" as used herein, refer to modes of administration other than enteral and topical administration, usually by injection. The phrases "systemic administration," "administered systemically", "peripheral administration" and "administered peripherally" as used herein refer to the administration of the bispecific or multispecific polypeptide agent other than directly into a target site, tissue, or organ, such as a tumor site, such that it enters the subject's circulatory system and, thus, is subject to metabolism and other like processes.

[0530] In some embodiments, the antibodies, or antigen binding fragment thereof, described herein, or compositions comprising the same can be administered via intravenous administration as a bolus or by continuous infusion over a period of time, by intramuscular, intraperitoneal, intracerebrospinal, subcutaneous, intra-articular, intrasynovial, intrathecal, oral, topical, or inhalation routes. Local administration, for example, to a tumor or cancer site where angiogenesis is occurring, is particularly desired if extensive side effects or toxicity is associated with the use of the antibodies, or antigen binding fragment thereof, described herein, or compositions comprising the same. An ex vivo strategy can also be used for therapeutic applications in some embodiments. Ex vivo strategies involve transfecting or transducing cells obtained from a subject with a nucleic acid sequence, disclosed herein. The transfected or transduced cells are then returned to the subject. The cells can be any of a wide range of types including, without limitation, hematopoietic cells (e.g., bone marrow cells, macrophages, monocytes, dendritic cells, T cells, or B cells), fibroblasts, epithelial cells, endothelial cells, keratinocytes, or muscle cells.

[0531] In some embodiments, an antibody or antigen binding fragment thereof, disclosed herein, or a composition comprising the same is administered by any suitable means, including parenteral, subcutaneous, intraperitoneal, intrapulmonary, and intranasal, and, if desired for local immunosuppressive treatment, intralesional administration.

[0532] Parenteral infusions include intramuscular, intravenous, intraarterial, intraperitoneal, or subcutaneous administration. In some embodiments, the antibody or antigen binding fragment thereof or compositions of the disclosure are suitably administered by pulse infusion, particularly with declining doses of the antibody. Preferably the dosing is given by injections, most preferably intravenous or subcutaneous injections, depending in part on whether the administration is brief or chronic. In some embodiments, the antibody or antigen binding fragment thereof or compositions of the disclosure are administered locally, e.g., by direct injections, when the disorder or location of the tumor permits, and the injections can be repeated periodically. In some embodiments, the antibody or antigen binding fragment thereof or compositions of the disclosure can also be delivered systemically to the subject or directly to the tumor cells, e.g., to a tumor or a

tumor bed following surgical excision of the tumor, in order to prevent or reduce local recurrence or metastasis, for example of a dormant tumor or micrometastases.

[0533] Antibody-targeted sonoporation methods are contemplated for use in some embodiments of the methods for inhibiting tumors described herein, in order to enhance the efficacy and potency of the therapeutic compositions comprising antibodies and antigen binding fragment thereof provided herein. As used herein, "sonoporation" refers to the use of sound, preferably at ultrasonic frequencies, or the interaction of ultrasound with contrast agents (e.g., stabilized microbubbles) for temporarily modifying the permeability of cell plasma membranes, thus allowing uptake of large molecules, such as therapeutic agents. The membrane permeability caused by the sonoporation is transient, leaving the agents trapped inside the cell after the ultrasound exposure. Sonoporation employs acoustic cavitation of microbubbles to enhance delivery of large molecules.

[0534] Accordingly, in some embodiments of the methods, the antibody or antigen binding fragment thereof described herein, mixed with ultrasound contrast agents, such as microbubbles, can be injected locally or systemically into a subject in need of treatment for cancer, and ultrasound can be coupled and even focused into the defined area, e.g., tumor site, to achieve targeted delivery. In some embodiments, the methods use focused ultrasound methods to achieve targeted delivery. As used herein, HIFU or "High Intensity Focused Ultrasound" refers to a non-invasive therapeutic method using high-intensity ultrasound to heat and destroy malignant or pathogenic tissue without causing damage to overlying or surrounding health tissue. As described in Khaibullina et al., 49 J. Nucl. Med. 295 (2008), and WO 2010127369, HIFU can also be used as a means of delivery of therapeutic agents, such as antibodies or antibody fragments thereof.

[0535] Methods using contrast-enhanced ultrasound (CEUS) are also contemplated for use with an antibody or antigen binding fragment thereof, described herein. Contrast-enhanced ultrasound (CEUS) refers to the application of ultrasound contrast medium and ultrasound contrast agents to traditional medical sonography. Ultrasound contrast agents refer to agents that rely on the different ways in which sound waves are reflected from interfaces between substances. A variety of microbubble contrast agents are available for use with the compositions and methods described herein. Microbubbles can differ in their shell makeup, gas core makeup, and whether or not they are targeted. Targeting ligands that bind to receptors characteristic of angiogenic disorders, can be conjugated to microbubbles, enabling the microbubble complex to accumulate selectively in areas of interest, such as diseased or abnormal tissues. This form of molecular imaging, known as targeted contrast-enhanced ultrasound, will only generate a strong ultrasound signal if targeted microbubbles bind in the area of interest. Targeted contrast-enhanced ultrasound has many applications in both medical diagnostics and medical therapeutics. In some embodiments, an antibody or antigen binding fragment thereof, described herein, is administered to a subject in need of treatment for a cancer or a tumor, using a targeted ultrasound delivery.

Pharmaceutical Compositions and Dosage Forms

[0536] Disclosed herein, in certain embodiments, are pharmaceutical compositions comprising an IgA antibody or a functional fragment thereof disclosed herein for administration in a subject.

5 [0537] In some embodiments, pharmaceutical compositions comprising a IgA antibody described herein are formulated in a conventional manner using one or more physiologically acceptable carriers including excipients and auxiliaries which facilitate processing of the active compounds into preparations which can be used pharmaceutically. Proper formulation is dependent upon the route of administration chosen. A summary of pharmaceutical compositions described herein is found, for example, in Remington: The Science and Practice of Pharmacy, Nineteenth Ed (Easton, Pa.: Mack Publishing Company, 1995); Hoover, John E., Remington's Pharmaceutical Sciences, Mack Publishing Co., Easton, Pennsylvania 1975; Liberman, H.A. and Lachman, L., Eds., Pharmaceutical Dosage Forms, Marcel Decker, New York, N.Y., 1980; and Pharmaceutical Dosage Forms and Drug Delivery Systems, Seventh Ed. (Lippincott Williams & Wilkins 1999).

15 [0538] Pharmaceutical compositions are optionally manufactured in a conventional manner, such as, by way of example only, by means of conventional mixing, dissolving, granulating, dragee-making, levigating, emulsifying, encapsulating, entrapping or compression processes.

[0539] In certain embodiments, compositions may also include one or more pH adjusting agents or buffering agents, including acids such as acetic, boric, citric, lactic, phosphoric and hydrochloric acids; bases such as sodium hydroxide, sodium phosphate, sodium borate, sodium citrate, sodium acetate, sodium lactate and tris-hydroxymethylaminomethane; and buffers such as citrate/dextrose, sodium bicarbonate and ammonium chloride. Such acids, bases and buffers are included in an amount required to maintain pH of the composition in an acceptable range.

20 [0540] In other embodiments, compositions may also include one or more salts in an amount required to bring osmolality of the composition into an acceptable range. Such salts include those having sodium, potassium or ammonium cations and chloride, citrate, ascorbate, borate, phosphate, bicarbonate, sulfate, thiosulfate or bisulfite anions; suitable salts include sodium chloride, potassium chloride, sodium thiosulfate, sodium bisulfite and ammonium sulfate.

25 [0541] The pharmaceutical compositions described herein are administered by any suitable administration route, including but not limited to, oral, parenteral (*e.g.*, intravenous, subcutaneous, intramuscular, intracerebral, intracerebroventricular, intra-articular, intraperitoneal, or intracranial), intranasal, buccal, sublingual, or rectal administration routes. In some embodiments, the pharmaceutical composition is formulated for parenteral (*e.g.*, intravenous, subcutaneous, intramuscular, intracerebral, intracerebroventricular, intra-articular, intraperitoneal, or intracranial) administration.

30

[0542] The pharmaceutical compositions described herein are formulated into any suitable dosage form, including but not limited to, aqueous oral dispersions, liquids, gels, syrups, elixirs, slurries, suspensions and the like, for oral ingestion by an individual to be treated, solid oral dosage forms, aerosols, controlled release formulations, fast melt formulations, effervescent formulations, lyophilized formulations, tablets, powders, pills, dragees, capsules, delayed release formulations, extended release formulations, pulsatile release formulations, multiparticulate formulations, and mixed immediate release and controlled release formulations.

[0543] In some embodiments, the pharmaceutical compositions are formulated into capsules. In some embodiments, the pharmaceutical compositions are formulated into solutions (for example, for IV administration). In some embodiments, the pharmaceutical composition is formulated as an infusion. In some embodiments, the pharmaceutical composition is formulated as an injection.

[0544] The pharmaceutical solid dosage forms described herein optionally include a compound described herein and one or more pharmaceutically acceptable additives such as a compatible carrier, binder, filling agent, suspending agent, flavoring agent, sweetening agent, disintegrating agent, dispersing agent, surfactant, lubricant, colorant, diluent, solubilizer, moistening agent, plasticizer, stabilizer, penetration enhancer, wetting agent, anti-foaming agent, antioxidant, preservative, or one or more combination thereof.

[0545] In still other aspects, using standard coating procedures, such as those described in Remington's Pharmaceutical Sciences, 20th Edition (2000), a film coating is provided around the compositions. In some embodiments, the compositions are formulated into particles (for example for administration by capsule) and some or all of the particles are coated. In some embodiments, the compositions are formulated into particles (for example for administration by capsule) and some or all of the particles are microencapsulated. In some embodiments, the compositions are formulated into particles (for example for administration by capsule) and some or all of the particles are not microencapsulated and are uncoated.

[0546] In certain embodiments, compositions provided herein may also include one or more preservatives to inhibit microbial activity. Suitable preservatives include mercury-containing substances such as merfen and thiomersal; stabilized chlorine dioxide; and quaternary ammonium compounds such as benzalkonium chloride, cetyltrimethylammonium bromide and cetylpyridinium chloride.

[0547] "Proliferative disease" as referred to herein means a unifying concept that excessive proliferation of cells and turnover of cellular matrix contribute significantly to the pathogenesis of several diseases, including cancer is presented.

[0548] "Patient" or "subject" as used herein refers to a mammalian subject diagnosed with or suspected of having or developing a physiological condition, for instance a cancer or an autoimmune condition or an infection. In some embodiments, the term "patient" refers to a mammalian subject with a higher than average likelihood of developing cancer. Exemplary patients may be humans, apes, dogs, pigs, cattle, cats,

horses, goats, sheep, rodents and other mammals that can benefit from the therapies disclosed herein. Exemplary human patients can be male and/or female. "Patient in need thereof" or "subject in need thereof" is referred to herein as a patient diagnosed with or suspected of having a disease or disorder, for instance, but not restricted to a proliferative disorder such as cancer. In some embodiments, a cancer is a solid tumor or a hematologic malignancy. In some embodiments, the cancer is a solid tumor. In other embodiments, the cancer is a hematologic malignancy. In some embodiments, the cancer is a metastatic cancer. In some
5 embodiments, the cancer is a relapsed or refractory cancer. In some embodiments, the cancer is a solid tumor. Exemplary solid tumors include, but are not limited to, anal cancer; appendix cancer; bile duct cancer (*i.e.*, cholangiocarcinoma); bladder cancer; brain tumor; breast cancer; cervical cancer; colon cancer; cancer of Unknown Primary (CUP); esophageal cancer; eye cancer; fallopian tube cancer;
10 gastroenterological cancer; kidney cancer; liver cancer; lung cancer; medulloblastoma; melanoma; oral cancer; ovarian cancer; pancreatic cancer; parathyroid disease; penile cancer; pituitary tumor; prostate cancer; rectal cancer; skin cancer; stomach cancer; testicular cancer; throat cancer; thyroid cancer; uterine cancer; vaginal cancer; vulvar cancer; or glioblastoma. In some embodiments leukemia can be, for
15 instance, acute lymphoblastic leukemia (ALL), acute myeloid leukemia (AML), chronic lymphocytic leukemia (CLL) and chronic myeloid leukemia (CML).

[0549] "Administering" is referred to herein as providing the compositions of the present disclosure to a patient. By way of example and not limitation, composition administration, *e.g.*, injection, may be performed by intravenous (*i.v.*) injection, sub-cutaneous (*s.c.*) injection, intradermal (*i.d.*) injection,
20 intraperitoneal (*i.p.*) injection, or intramuscular (*i.m.*) injection. One or more such routes may be employed. Parenteral administration can be, for example, by bolus injection or by gradual perfusion over time. Alternatively, or concurrently, administration may be by the oral route. Additionally, administration may also be by surgical deposition of a bolus or pellet of cells, or positioning of a medical device. In an
25 embodiment, a composition of the present disclosure may comprise engineered cells or host cells expressing nucleic acid sequences described herein, or a vector comprising at least one nucleic acid sequence described herein, in an amount that is effective to treat or prevent proliferative disorders. A pharmaceutical composition may comprise a target cell population as described herein, in combination with one or more pharmaceutically or physiologically acceptable carriers, diluents or excipients. Such
30 compositions may comprise buffers such as neutral buffered saline, phosphate buffered saline and the like; carbohydrates such as glucose, mannose, sucrose or dextrans, mannitol; proteins; polypeptides or amino acids such as glycine; antioxidants; chelating agents such as EDTA or glutathione; adjuvants (*e.g.*, aluminum hydroxide); and preservatives.

[0550] As used herein, the terms "treatment," "treating," and its grammatical equivalents refer to obtaining a desired pharmacologic and/or physiologic effect. In embodiments, the effect is therapeutic, *i.e.*, the effect

partially or completely cures a disease and/or adverse symptom attributable to the disease. To this end, the method described herein comprises administering a “therapeutically effective amount” of the composition comprising the host cells expressing the nucleic acid sequence described herein, or a vector comprising the nucleic acid sequences described herein.

5 [0551] The terms “therapeutically effective amount”, “therapeutic amount”, “immunologically effective amount”, “anti-tumor effective amount”, “tumor inhibiting effective amount” or their grammatical equivalents refers to an amount effective, at dosages and for periods of time necessary, to achieve a desired result. The therapeutically effective amount may vary according to factors such as the disease state, age, sex, and weight of the individual and the ability of a composition described herein to elicit a desired
10 response in the individual. The precise amount of the compositions of the present disclosure to be administered can be determined by a physician with consideration of individual differences in age, weight, tumor size, extent of infection or metastasis, and condition of the patient (subject).

[0552] Alternatively, the pharmacologic and/or physiologic effect of administration of one or more compositions described herein to a patient or a subject may be “prophylactic,” *i.e.*, the effect completely or
15 partially prevents a disease or symptom thereof.

[0553] A “prophylactically effective amount” refers to an amount effective, at dosages and for periods of time necessary, to achieve a desired prophylactic result (*e.g.*, prevention of disease onset).

[0554] “Antifoaming agents” reduce foaming during processing which can result in coagulation of aqueous dispersions, bubbles in the finished film, or generally impair processing. Exemplary anti-foaming agents
20 include silicon emulsions or sorbitan sesquoleate.

[0555] Antioxidants” include, for example, butylated hydroxytoluene (BHT), sodium ascorbate, ascorbic acid, sodium metabisulfite and tocopherol. In certain embodiments, antioxidants enhance chemical stability where required.

[0556] Formulations described herein may benefit from antioxidants, metal chelating agents, thiol
25 containing compounds and other general stabilizing agents. Examples of such stabilizing agents, include, but are not limited to: (a) about 0.5% to about 2% w/v glycerol, (b) about 0.1% to about 1% w/v methionine, (c) about 0.1% to about 2% w/v monothioglycerol, (d) about 1 mM to about 10 mM EDTA, (e) about 0.01% to about 2% w/v ascorbic acid, (f) 0.003% to about 0.02% w/v polysorbate 80, (g) 0.001% to about 0.05% w/v. polysorbate 20, (h) arginine, (i) heparin, (j) dextran sulfate, (k) cyclodextrins, (l) pentosan
30 polysulfate and other heparinoids, (m) divalent cations such as magnesium and zinc; or (n) combinations thereof.

[0557] “Binders” impart cohesive qualities and include, *e.g.*, alginic acid and salts thereof; cellulose derivatives such as carboxymethylcellulose, methylcellulose (*e.g.*, Methocel®), hydroxypropylmethylcellulose, hydroxyethylcellulose, hydroxypropylcellulose (*e.g.*, Klucel®),

ethylcellulose (*e.g.*, Ethocel®), and microcrystalline cellulose (*e.g.*, Avicel®); microcrystalline dextrose; amylose; magnesium aluminum silicate; polysaccharide acids; bentonites; gelatin; polyvinylpyrrolidone/vinyl acetate copolymer; crospovidone; povidone; starch; pregelatinized starch; tragacanth, dextrin, a sugar, such as sucrose (*e.g.*, Dipac®), glucose, dextrose, molasses, mannitol, sorbitol, xylitol (*e.g.*, Xylitab®), and lactose; a natural or synthetic gum such as acacia, tragacanth, ghatti gum, mucilage of isapol husks, polyvinylpyrrolidone (*e.g.*, Polyvidone® CL, Kollidon® CL, Polyplasdone® XL-10), larch arabogalactan, Veegum®, polyethylene glycol, waxes, sodium alginate, and the like.

[0558] A “carrier” or “carrier materials” include any commonly used excipients in pharmaceuticals and should be selected on the basis of compatibility with compounds disclosed herein, such as, compounds of ibrutinib and an anticancer agent, and the release profile properties of the desired dosage form. Exemplary carrier materials include, *e.g.*, binders, suspending agents, disintegration agents, filling agents, surfactants, solubilizers, stabilizers, lubricants, wetting agents, diluents, and the like. “Pharmaceutically compatible carrier materials” may include, but are not limited to, acacia, gelatin, colloidal silicon dioxide, calcium glycerophosphate, calcium lactate, maltodextrin, glycerine, magnesium silicate, polyvinylpyrrolidone (PVP), cholesterol, cholesterol esters, sodium caseinate, soy lecithin, taurocholic acid, phosphatidylcholine, sodium chloride, tricalcium phosphate, dipotassium phosphate, cellulose and cellulose conjugates, sugars sodium stearoyl lactylate, carrageenan, monoglyceride, diglyceride, pregelatinized starch, and the like. See, *e.g.*, Remington: The Science and Practice of Pharmacy, Nineteenth Ed (Easton, Pa.: Mack Publishing Company, 1995); Hoover, John E., Remington’s Pharmaceutical Sciences, Mack Publishing Co., Easton, Pennsylvania 1975; Liberman, H.A. and Lachman, L., Eds., Pharmaceutical Dosage Forms, Marcel Decker, New York, N.Y., 1980; and Pharmaceutical Dosage Forms and Drug Delivery Systems, Seventh Ed. (Lippincott Williams & Wilkins 1999).

[0559] “Dispersing agents,” and/or “viscosity modulating agents” include materials that control the diffusion and homogeneity of a drug through liquid media or a granulation method or blend method. In some embodiments, these agents also facilitate the effectiveness of a coating or eroding matrix. Exemplary diffusion facilitators/dispersing agents include, *e.g.*, hydrophilic polymers, electrolytes, Tween® 60 or 80, PEG, polyvinylpyrrolidone (PVP; commercially known as Plasdone®), and the carbohydrate-based dispersing agents such as, for example, hydroxypropyl celluloses (*e.g.*, HPC, HPC-SL, and HPC-L), hydroxypropyl methylcelluloses (*e.g.*, HPMC K100, HPMC K4M, HPMC K15M, and HPMC K100M), carboxymethylcellulose sodium, methylcellulose, hydroxyethylcellulose, hydroxypropylcellulose, hydroxypropylmethylcellulose phthalate, hydroxypropylmethylcellulose acetate stearate (HPMCAS), noncrystalline cellulose, magnesium aluminum silicate, triethanolamine, polyvinyl alcohol (PVA), vinyl pyrrolidone/vinyl acetate copolymer (S630), 4-(1,1,3,3-tetramethylbutyl)-phenol polymer with ethylene oxide and formaldehyde (also known as tyloxapol), poloxamers (*e.g.*, Pluronic F68®, F88®, and F108®),

which are block copolymers of ethylene oxide and propylene oxide); and poloxamines (*e.g.*, Tetronic 908®, also known as Poloxamine 908®, which is a tetrafunctional block copolymer derived from sequential addition of propylene oxide and ethylene oxide to ethylenediamine (BASF Corporation, Parsippany, N.J.)), polyvinylpyrrolidone K12, polyvinylpyrrolidone K17, polyvinylpyrrolidone K25, or polyvinylpyrrolidone K30, polyvinylpyrrolidone/vinyl acetate copolymer (S-630), polyethylene glycol, *e.g.*, the polyethylene glycol can have a molecular weight of about 300 to about 6000, or about 3350 to about 4000, or about 7000 to about 5400, sodium carboxymethylcellulose, methylcellulose, polysorbate-80, sodium alginate, gums, such as, *e.g.*, gum tragacanth and gum acacia, guar gum, xanthans, including xanthan gum, sugars, cellulose, such as, *e.g.*, sodium carboxymethylcellulose, methylcellulose, sodium carboxymethylcellulose, polysorbate-80, sodium alginate, polyethoxylated sorbitan monolaurate, polyethoxylated sorbitan monolaurate, povidone, carbomers, polyvinyl alcohol (PVA), alginates, chitosans and combinations thereof. Plasticizers such as cellulose or triethyl cellulose can also be used as dispersing agents. Dispersing agents particularly useful in liposomal dispersions and self-emulsifying dispersions are dimyristoyl phosphatidyl choline, natural phosphatidyl choline from eggs, natural phosphatidyl glycerol from eggs, cholesterol and isopropyl myristate.

[0560] Combinations of one or more erosion facilitator with one or more diffusion facilitator can also be used in the present compositions.

[0561] The term “diluent” refers to chemical compounds that are used to dilute the compound of interest prior to delivery. Diluents can also be used to stabilize compounds because they can provide a more stable environment. Salts dissolved in buffered solutions (which also can provide pH control or maintenance) are utilized as diluents in the art, including, but not limited to a phosphate buffered saline solution. In certain embodiments, diluents increase bulk of the composition to facilitate compression or create sufficient bulk for homogenous blend for capsule filling. Such compounds include *e.g.*, lactose, starch, mannitol, sorbitol, dextrose, microcrystalline cellulose such as Avicel®; dibasic calcium phosphate, dicalcium phosphate dihydrate; tricalcium phosphate, calcium phosphate; anhydrous lactose, spray-dried lactose; pregelatinized starch, compressible sugar, such as Di-Pac® (Amstar); mannitol, hydroxypropylmethylcellulose, hydroxypropylmethylcellulose acetate stearate, sucrose-based diluents, confectioner’s sugar; monobasic calcium sulfate monohydrate, calcium sulfate dihydrate; calcium lactate trihydrate, dextrates; hydrolyzed cereal solids, amylose; powdered cellulose, calcium carbonate; glycine, kaolin; mannitol, sodium chloride; inositol, bentonite, and the like.

[0562] “Filling agents” include compounds such as lactose, calcium carbonate, calcium phosphate, dibasic calcium phosphate, calcium sulfate, microcrystalline cellulose, cellulose powder, dextrose, dextrates, dextran, starches, pregelatinized starch, sucrose, xylitol, lactitol, mannitol, sorbitol, sodium chloride, polyethylene glycol, and the like.

- [0563] “Lubricants” and “glidants” are compounds that prevent, reduce or inhibit adhesion or friction of materials. Exemplary lubricants include, *e.g.*, stearic acid, calcium hydroxide, talc, sodium stearyl fumarate, a hydrocarbon such as mineral oil, or hydrogenated vegetable oil such as hydrogenated soybean oil (Sterotex®), higher fatty acids and their alkali-metal and alkaline earth metal salts, such as aluminum, calcium, magnesium, zinc, stearic acid, sodium stearates, glycerol, talc, waxes, Stearowet®, boric acid, sodium benzoate, sodium acetate, sodium chloride, leucine, a polyethylene glycol (*e.g.*, PEG-4000) or a methoxypolyethylene glycol such as Carbowax™, sodium oleate, sodium benzoate, glyceryl behenate, polyethylene glycol, magnesium or sodium lauryl sulfate, colloidal silica such as Syloid™, Cab-O-Sil®, a starch such as corn starch, silicone oil, a surfactant, and the like.
- 5
- [0564] “Plasticizers” are compounds used to soften the microencapsulation material or film coatings to make them less brittle. Suitable plasticizers include, *e.g.*, polyethylene glycols such as PEG 300, PEG 400, PEG 600, PEG 1450, PEG 3350, and PEG 800, stearic acid, propylene glycol, oleic acid, triethyl cellulose and triacetin. In some embodiments, plasticizers can also function as dispersing agents or wetting agents.
- 10
- [0565] “Solubilizers” include compounds such as triacetin, triethylcitrate, ethyl oleate, ethyl caprylate, sodium lauryl sulfate, sodium docusate, vitamin E TPGS, dimethylacetamide, N- methylpyrrolidone, N- hydroxyethylpyrrolidone, polyvinylpyrrolidone, hydroxypropylmethyl cellulose, hydroxypropyl cyclodextrins, ethanol, n-butanol, isopropyl alcohol, cholesterol, bile salts, polyethylene glycol 200-600, glycofurol, transcutool, propylene glycol, and dimethyl isosorbide and the like.
- 15
- [0566] “Stabilizers” include compounds such as any antioxidation agents, buffers, acids, preservatives and the like.
- 20
- [0567] “Suspending agents” include compounds such as polyvinylpyrrolidone, *e.g.*, polyvinylpyrrolidone K12, polyvinylpyrrolidone K17, polyvinylpyrrolidone K25, or polyvinylpyrrolidone K30, vinyl pyrrolidone/vinyl acetate copolymer (S630), polyethylene glycol, *e.g.*, the polyethylene glycol can have a molecular weight of about 300 to about 6000, or about 3350 to about 4000, or about 7000 to about 5400, sodium carboxymethylcellulose, methylcellulose, hydroxypropylmethylcellulose, hydroxymethylcellulose acetate stearate, polysorbate-80, hydroxyethylcellulose, sodium alginate, gums, such as, *e.g.*, gum tragacanth and gum acacia, guar gum, xanthans, including xanthan gum, sugars, cellulose, such as, *e.g.*, sodium carboxymethylcellulose, methylcellulose, sodium carboxymethylcellulose, hydroxypropylmethylcellulose, hydroxyethylcellulose, polysorbate-80, sodium alginate, polyethoxylated sorbitan monolaurate, polyethoxylated sorbitan monolaurate, povidone and the like.
- 25
- 30
- [0568] “Surfactants” include compounds such as sodium lauryl sulfate, sodium docusate, Tween 60 or 80, triacetin, vitamin E TPGS, sorbitan monooleate, polyoxyethylene sorbitan monooleate, polysorbates, polaxomers, bile salts, glyceryl monostearate, copolymers of ethylene oxide and propylene oxide, *e.g.*, Pluronic® (BASF), and the like. Some other surfactants include polyoxyethylene fatty acid glycerides and

vegetable oils, *e.g.*, polyoxyethylene (60) hydrogenated castor oil; and polyoxyethylene alkylethers and alkylphenyl ethers, *e.g.*, octoxynol 10, octoxynol 40. In some embodiments, surfactants may be included to enhance physical stability or for other purposes.

[0569] "Viscosity enhancing agents" include, *e.g.*, methyl cellulose, xanthan gum, carboxymethyl cellulose, hydroxypropyl cellulose, hydroxypropylmethyl cellulose, hydroxypropylmethyl cellulose acetate stearate, hydroxypropylmethyl cellulose phthalate, carbomer, polyvinyl alcohol, alginates, acacia, chitosans and combinations thereof.

[0570] "Wetting agents" include compounds such as oleic acid, glyceryl monostearate, sorbitan monooleate, sorbitan monolaurate, triethanolamine oleate, polyoxyethylene sorbitan monooleate, polyoxyethylene sorbitan monolaurate, sodium docusate, sodium oleate, sodium lauryl sulfate, sodium docusate, triacetin, Tween 80, vitamin E TPGS, ammonium salts and the like.

[0571] Optionally, the formulations comprising the compositions described herein contain a pharmaceutically acceptable salt, typically, *e.g.*, sodium chloride, and preferably at about physiological concentrations. Optionally, the formulations of the invention can contain a pharmaceutically acceptable preservative. In some embodiments the preservative concentration ranges from 0.1 to 2.0%, typically v/v. Suitable preservatives include those known in the pharmaceutical arts. Benzyl alcohol, phenol, m-cresol, methylparaben, and propylparaben are examples of preservatives. Optionally, the formulations of the invention can include a pharmaceutically acceptable surfactant at a concentration of 0.005 to 0.02%.

[0572] The compositions described herein can be specially formulated for administration of the antibody or antigen binding fragment thereof to a subject in solid, liquid or gel form, including those adapted for the following: (1) parenteral administration, for example, by subcutaneous, intramuscular, intravenous or epidural injection as, for example, a sterile solution or suspension, or sustained-release formulation; (2) topical application, for example, as a cream, ointment, or a controlled-release patch or spray applied to the skin; (3) intravaginally or intrarectally, for example, as a pessary, cream or foam; (4) ocularly; (5) transdermally; (6) transmucosally; or (7) nasally. Additionally, an antibody or antigen binding fragment thereof, or compositions of the present disclosure can be implanted into a patient or injected using a drug delivery system. See, *e.g.*, Urquhart et al., 24 Ann. Rev. Pharmacol. Toxicol. 199 (1984); *Controlled Release of Pesticides & Pharmaceuticals* (Lewis, ed., Plenum Press, New York, 1981); U.S. Patents No. 3,773,919, No. 3,270,960.

[0573] The compositions disclosed herein, comprising an antibody or antigen binding fragment, described herein, can also contain more than one active compound as necessary for the particular indication being treated, preferably those with complementary activities that do not adversely affect each other. For example, the composition can further comprise a cytotoxic agent, cytokine, growth inhibitory agent and/or an angiogenesis inhibitor such as a VEGFR antagonist. Such molecules are suitably present in combination in amounts that are effective for the purpose intended. The active ingredients of the compositions comprising an

antibody or antigen binding fragment thereof described herein can also be entrapped in microcapsules prepared, for example, by coacervation techniques or by interfacial polymerization, for example, hydroxymethylcellulose or gelatin-microcapsules and poly-(methylmethacrylate) microcapsules, respectively, in colloidal drug delivery systems (for example, liposomes, albumin microspheres, microparticle, 5 microemulsions, nano-particles and nanocapsules) or in macroemulsions. Such techniques are disclosed in Remington's Pharmaceutical Sciences (16th ed., Osol, ed., 1980). The pharmaceutical composition can be also delivered in a vesicle, in particular a liposome (see Langer 1990 Science 249:1527-1533; Treat et al. (1989) in Liposomes in the Therapy of Infectious Disease and Cancer, Lopez-Berestein and Fidler (eds.), Liss, New York, pp. 353-365; Lopez-Berestein, *ibid.*, pp. 317-327; see generally *ibid.*). Liposomes include emulsions, 10 foams, micelles, insoluble monolayers, phospholipid dispersions, lamellar layers and the like, and can serve as vehicles to target the M-CSF antibodies to a particular tissue as well as to increase the half life of the composition. A variety of methods are available for preparing liposomes, as described in, e.g., U.S. Pat. Nos. 4,837,028 and 5,019,369, which patents are incorporated herein by reference.

[0574] Liposomes containing the antibody are prepared by methods known in the art, such as described in 15 Epstein et al., Proc. Natl. Acad. Sci. USA 82: 3688 (1985); Hwang et al., Proc. Natl Acad. Sci. USA 77: 4030 (1980); and U.S. Pat. Nos. 4,485,045 and 4,544,545. Liposomes with enhanced circulation time are disclosed in U.S. Pat. No. 5,013,556. Particularly useful liposomes can be generated by the reverse phase evaporation method with a lipid composition comprising phosphatidylcholine, cholesterol and PEG-derivatized phosphatidylethanolamine (PEG-PE). Liposomes are extruded through filters of defined pore size to yield 20 liposomes with the desired diameter. Fab' fragments of the antibody of the present invention can be conjugated to the liposomes as described in Martin et al., J. Biol. Chem. 257: 286-288 (1982) via a disulfide interchange reaction. A chemotherapeutic agent (such as Doxorubicin) is optionally contained within the liposome [see, e.g., Gabizon et al., J. National Cancer Inst. 81(19): 1484 (1989)].

[0575] In some embodiments, sustained-release preparations can be used. Suitable examples of sustained- 25 release preparations include semipermeable matrices of solid hydrophobic polymers containing an antibody or antigen binding fragment of the present disclosure, in which the matrices are in the form of shaped articles, e.g., films, or microcapsule. Examples of sustained-release matrices include polyesters, hydrogels (for example, poly(2-hydroxyethyl-methacrylate), or poly(vinylalcohol)), polylactides (U.S. Patent No. 3,773,919), copolymers of L-glutamic acid and γ ethyl-L-glutamate, non-degradable ethylene-vinyl acetate, 30 degradable lactic acid-glycolic acid copolymers such as the LUPRON DEPOT™ (injectable microspheres composed of lactic acid-glycolic acid copolymer and leuprolide acetate), and poly-D-(-)-3-hydroxybutyric acid. While polymers such as ethylene-vinyl acetate and lactic acid-glycolic acid enable release of molecules for over 100 days, certain hydrogels release proteins for shorter time periods. When encapsulated antibodies remain in the body for a long time, they can denature or aggregate as a result of exposure to moisture at 37°C,

resulting in a loss of biological activity and possible changes in immunogenicity. Rational strategies can be devised for stabilization depending on the mechanism involved. For example, if the aggregation mechanism is discovered to be intermolecular S-S bond formation through thiodisulfide interchange, stabilization can be achieved by modifying sulfhydryl residues, lyophilizing from acidic solutions, controlling moisture content, using appropriate additives, and developing specific polymer matrix compositions. In certain situations, the pharmaceutical composition can be delivered in a controlled release system. In one embodiment, a pump may be used (see Langer, *supra*; Sefton 1987 CRC Crit. Ref. Biomed. Eng. 14:201). In another embodiment, polymeric materials can be used. In yet another embodiment, a controlled release system can be placed in proximity of the composition's target, thus requiring only a fraction of the systemic dose (see, e.g., Goodson, 1984, in *Medical Applications of Controlled Release*, *supra*, vol. 2, pp. 115-138).

[0576] A pharmaceutical composition of the present disclosure can be delivered, e.g., subcutaneously or intravenously with a standard needle and syringe. In addition, with respect to subcutaneous delivery, a pen delivery device readily has applications in delivering a pharmaceutical composition of the present invention. Such a pen delivery device can be reusable or disposable. A reusable pen delivery device generally utilizes a replaceable cartridge that contains a pharmaceutical composition. Once all of the pharmaceutical composition within the cartridge has been administered and the cartridge is empty, the empty cartridge can readily be discarded and replaced with a new cartridge that contains the pharmaceutical composition. The pen delivery device can then be reused. In a disposable pen delivery device, there is no replaceable cartridge. Rather, the disposable pen delivery device comes prefilled with the pharmaceutical composition held in a reservoir within the device. Once the reservoir is emptied of the pharmaceutical composition, the entire device is discarded. Numerous reusable pen and autoinjector delivery devices have applications in the subcutaneous delivery of a pharmaceutical composition of the present invention. Examples include, but certainly are not limited to AUTOPEN™ (Owen Mumford, Inc., Woodstock, UK), DISETRONIC™ pen (Disetronic Medical Systems, Burghdorf, Switzerland), HUMALOG MIX 75/25™ pen, HUMALOG™ pen, HUMALIN70130™ pen (Eli Lilly and Co., Indianapolis, Ind.), NOVOPEN™ I, II and III (Novo Nordisk, Copenhagen, Denmark), NOVOPEN JUNIOR™ (Novo Nordisk, Copenhagen, Denmark), BD™ pen (Becton Dickinson, Franklin Lakes, N.J.), OPTIPEN™, OPTIPEN PRO™, OPTIPEN STARLET™, and OPTICLIK™ (Sanofi-Aventis, Frankfurt, Germany), to name only a few. Examples of disposable pen delivery devices having applications in subcutaneous delivery of a pharmaceutical composition include, but certainly are not limited to the SOLOSTAR™ pen (Sanofi-Aventis), the FLEXPEN™ (Novo Nordisk), and the KWIKPEN™ (Eli Lilly).

[0577] The injectable preparations may include dosage forms for intravenous, subcutaneous, intracutaneous and intramuscular injections, drip infusions, etc. These injectable preparations may be prepared by methods publicly known. For example, the injectable preparations may be prepared, e.g., by dissolving, suspending or emulsifying the antibody or its salt described above in a sterile aqueous medium or an oily medium

conventionally used for injections. As the aqueous medium for injections, there are, for example, physiological saline, an isotonic solution containing glucose and other auxiliary agents, etc., which may be used in combination with an appropriate solubilizing agent such as an alcohol (e.g., ethanol), a polyalcohol (e.g., propylene glycol, polyethylene glycol), a nonionic surfactant [e.g., polysorbate 80, HCO-50 (polyoxyethylene (50 mol) adduct of hydrogenated castor oil)], etc. As the oily medium, there are employed, e.g., sesame oil, soybean oil, etc., which may be used in combination with a solubilizing agent such as benzyl benzoate, benzyl alcohol, etc. The injection thus prepared is preferably filled in an appropriate ampoule.

[0578] Compositions of the present disclosure can be in the form of, for example, granules, powders, tablets, capsules, syrup, suppositories, injections, emulsions, elixirs, suspensions or solutions. The amount of the aforesaid antibody contained can be about 5 to about 500 mg per dosage form in a unit dose; especially in the form of injection, it is preferred that the aforesaid antibody is contained in about 5 to about 100 mg and in about 10 to about 250 mg for the other dosage forms.

[0579] For oral, buccal, and sublingual administration, powders, suspensions, granules, tablets, pills, capsules, gelcaps, and caplets are acceptable as solid dosage forms. These can be prepared, for example, by mixing one or more compounds of the instant invention, or pharmaceutically acceptable salts or tautomers thereof, with at least one additive such as a starch or other additive. Suitable additives are sucrose, lactose, cellulose sugar, mannitol, maltitol, dextran, starch, agar, alginates, chitins, chitosans, pectins, tragacanth gum, gum arabic, gelatins, collagens, casein, albumin, synthetic or semi-synthetic polymers or glycerides. Optionally, oral dosage forms can contain other ingredients to aid in administration, such as an inactive diluent, or lubricants such as magnesium stearate, or preservatives such as paraben or sorbic acid, or anti-oxidants such as ascorbic acid, tocopherol or cysteine, a disintegrating agent, binders, thickeners, buffers, sweeteners, flavoring agents or perfuming agents. Tablets and pills may be further treated with suitable coating materials known in the art.

[0580] Liquid dosage forms for oral administration may be in the form of pharmaceutically acceptable emulsions, syrups, elixirs, suspensions, and solutions, which may contain an inactive diluent, such as water. In some embodiments, pharmaceutical formulations and medicaments may be prepared as liquid suspensions or aqueous solutions, for example, using a sterile liquid, such as, but not limited to, an oil, water, an alcohol, and combinations of these. In some embodiments, pharmaceutical compositions can be prepared in a lyophilized form. The lyophilized preparations can comprise a cryoprotectant known in the art. The term “cryoprotectants” as used herein generally includes agents, which provide stability to the protein from freezing-induced stresses. Examples of cryoprotectants include polyols such as, for example, mannitol, and include saccharides such as, for example, sucrose, as well as including surfactants such as, for example, polysorbate, poloxamer or polyethylene glycol, and the like. Cryoprotectants also contribute to the tonicity of

the formulations. Pharmaceutically suitable surfactants, suspending agents, emulsifying agents, may be added for oral or par- enteral administration.

[0581] As noted above, suspensions may include oils. Such oils include, but are not limited to, peanut oil, sesame oil, cottonseed oil, corn oil and olive oil. Suspension preparation may also contain esters of fatty acids such as ethyl oleate, isopropyl myristate, fatty acid glycerides and acetylated fatty acid glycerides. Suspension formulations may include alcohols, such as, but not limited to, ethanol, iso- propyl alcohol, hexadecyl alcohol, glycerol and propylene glycol. Ethers, such as but not limited to, poly(ethyleneglycol), petroleum hydrocarbons such as mineral oil and petrolatum; and water may also be used in suspension formulations.

[0582] For nasal administration, the pharmaceutical formulations and medicaments may be a spray or aerosol containing an appropriate solvent(s) and optionally other compounds such as, but not limited to, stabilizers, antimicrobial agents, antioxidants, pH modifiers, surfactants, bio- availability modifiers and combinations of these. A propellant for an aerosol formulation may include compressed air, nitrogen, carbon dioxide, or a hydrocarbon based low boiling solvent.

[0583] Injectable dosage forms generally include aqueous suspensions or oil suspensions which may be prepared using a suitable dispersant or wetting agent and a suspending agent. Injectable forms may be in solution phase or in the form of a suspension, which is prepared with a solvent or diluent. Acceptable solvents or vehicles include sterilized water, Ringer's solution, or an isotonic aqueous saline solution. Alternatively, sterile oils may be employed as solvents or suspending agents. Preferably, the oil or fatty acid is non-volatile, including natural or synthetic oils, fatty acids, mono-, di- or tri-glycerides.

[0584] For injection, the pharmaceutical formulation and/ or medicament may be a powder suitable for reconstitution with an appropriate solution as described above. Examples of these include, but are not limited to, freeze dried, rotary dried or spray dried powders, amorphous powders, granules, precipitates, or particulates. For injection, the formulations may optionally contain stabilizers, pH modifiers, surfactants, bioavailability modifiers and combinations of these.

[0585] For rectal administration, the pharmaceutical formulations and medicaments may be in the form of a suppository, an ointment, an enema, a tablet or a cream for release of compound in the intestines, sigmoid flexure and/or rectum. Rectal suppositories are prepared by mixing one or more compounds of the instant invention, or pharmaceutically acceptable salts or tautomers of the compound, with acceptable vehicles, for example, cocoa butter or polyethylene glycol, which is present in a solid phase at normal storing temperatures, and present in a liquid phase at those temperatures suitable to release a drug inside the body, such as in the rectum. Oils may also be employed in the preparation of formulations of the soft gelatin type and suppositories. Water, saline, aqueous dextrose and related sugar solutions, and glycerols may be employed in the preparation of suspension formulations which may also contain suspending agents such as

pectins, carbomers, methyl cellulose, hydroxypropyl cellulose or carboxymethyl cellulose, as well as buffers and preservatives.

[0586] The concentration of an antibody or an antigen binding fragment thereof in these compositions can vary widely, i.e., from less than about 10%, usually at least about 25% to as much as 75% or 90% by weight and will be selected primarily by fluid volumes, viscosities, etc., in accordance with the particular mode of administration selected. Actual methods for preparing orally, topically and parenterally administrable compositions will be known or apparent to those skilled in the art and are described in detail in, for example, Remington's Pharmaceutical Science, 19th ed., Mack Publishing Co., Easton, Pa. (1995), which is incorporated herein by reference.

[0587] In another embodiment of the invention, an article of manufacture containing materials useful for the treatment of the diseases, disorders or conditions described above is provided, including for treatment of cancer. The article of manufacture comprises a container and a label. Suitable containers include, for example, bottles, vials, syringes, and test tubes. The containers may be formed from a variety of materials such as glass or plastic. The container holds a composition which is effective for treating the condition and may have a sterile access port (for example the container may be an intravenous solution bag or a vial having a stopper pierceable by a hypodermic injection needle). The active agent in the composition is the antibody of the invention. The label on or associated with, the container indicates that the composition is used for treating the condition of choice. The article of manufacture may further comprise a second container comprising a pharmaceutically- acceptable buffer, such as phosphate-buffered saline, Ringer's solution and dextrose solution. It may further include other materials desirable from a commercial and user stand- point, including other buffers, diluents, filters, needles, syringes, and package inserts with instructions for use. Pharmaceutical compositions and medicaments described herein are useful in treating a cancerous disease.

Diagnostic and other Uses

[0588] Provided herein are methods of using the antibodies for detection, diagnosis and monitoring of a disease, disorder or condition associated with the antigen expression (either increased or decreased relative to a normal sample, and/or inappropriate expression, such as presence of expression in tissues(s) and/or cell(s) that normally lack the epitope expression). Provided herein are methods of determining whether a patient will respond to antibody therapy.

[0589] In some embodiments, the method comprises detecting whether the patient has cells that express target antigen using an antibody disclosed herein. In some embodiments, the method of detection comprises contacting the sample with an antibody or antigen binding fragment thereof of the disclosure, and determining whether the level of binding differs from that of a reference or comparison sample (such as a control). In some embodiments, the method may be useful to determine whether the antibodies or polypeptides described herein are an appropriate treatment for the subject.

[0590] In some embodiments, the cells or cell/tissue lysate are contacted with an antibody and the binding between the antibody and the cell is determined. When the test cells show binding activity as compared to a reference cell of the same tissue type, it may indicate that the subject would benefit from treatment with an antibody. In some embodiments, the test cells are from human tissues. In some embodiments, the test cells are from human blood.

[0591] Various methods known in the art for detecting specific antibody-antigen binding can be used. Exemplary immunoassays which can be conducted include fluorescence polarization immunoassay (FPIA), fluorescence immunoassay (FIA), enzyme immunoassay (EIA), nephelometric inhibition immunoassay (NIA), enzyme linked immunosorbent assay (ELISA), and radioimmunoassay (RIA). An indicator moiety, or label group, can be attached to the subject antibodies and is selected so as to meet the needs of various uses of the method which are often dictated by the availability of assay equipment and compatible immunoassay procedures.

[0592] Appropriate labels include, without limitation, radionuclides (for example ¹²⁵I, ¹³¹I, ³⁵S, ³H, or ³²P), enzymes (for example, alkaline phosphatase, horseradish peroxidase, luciferase, or β -galactosidase), fluorescent moieties or proteins (for example, fluorescein, rhodamine, phycoerythrin, GFP, or BFP), or luminescent moieties (for example, QdotTM nanoparticles supplied by the Quantum Dot Corporation, Palo Alto, Calif.). General techniques to be used in performing the various immunoassays noted above are known to those of ordinary skill in the art.

[0593] For purposes of diagnosis, the antibodies or antigen binding fragment thereof can be labeled with a detectable moiety including but not limited to radioisotopes, fluorescent labels, and various enzyme-substrate labels known in the art. Methods of conjugating labels to an antibody are known in the art.

[0594] In some embodiments, the antibodies need not be labeled, and the presence thereof can be detected using a second labeled antibody which binds to the first antibody. The antibodies or antigen binding fragment thereof of the present invention may be used as affinity purification agents for a cancer associated antigen or in diagnostic assays for a cancer associated antigen protein, e.g., detecting its expression in specific cells, tissues, or serum. The antibodies or antigen binding fragment thereof, disclosed herein, may also be used for in vivo diagnostic assays. Generally, for these purposes the antibody is labeled with a radionuclide (such as ^{111}In , ^{99}Tc , ^{14}C , ^{131}I , ^{125}I , ^3H , ^{32}P or ^{35}S) so that the tumor can be localized using immunoscintigraphy.

[0595] The antibodies of the present invention may be employed in any known assay method, such as competitive binding assays, direct and indirect sandwich assays, such as ELISAs, and immunoprecipitation assays. Zola, *Monoclonal Antibodies: A Manual of Techniques*, pp. 147-158 (CRC Press, Inc. 1987). The antibodies may also be used for immunohistochemistry, to label tumor samples using methods known in the art. As a matter of convenience, the antibody of the present invention can be provided in a kit, i.e., a packaged combination of reagents in predetermined amounts with instructions for performing the diagnostic assay. Where the antibody is labeled with an enzyme, the kit will include substrates and cofactors required by the enzyme (e.g., a substrate precursor which provides the detectable chromophore or fluorophore). In addition, other additives may be included such as stabilizers, buffers (e.g., a block buffer or lysis buffer) and the like. The relative amounts of the various reagents may be varied widely to provide for concentrations in solution of the reagents which substantially optimize the sensitivity of the assay. Particularly, the reagents may be provided as dry powders, usually lyophilized, including excipients which on dissolution will provide a reagent solution having the appropriate concentration.

Kits

[0596] Provided herein are also kits, medicines, compositions, and unit dosage forms for use in any of the methods described herein. Provided herein is a kit comprising a therapeutically effective amount of at least one of the IgA antibody or antigen binding fragment thereof disclosed herein. In some embodiments, the kit further comprises a second therapeutic agent (e.g., a chemotherapeutic agent). In some embodiments, the antibody or antigen binding fragment thereof is an aqueous form or a lyophilized form. The kit further comprises a diluent or a reconstitution solution.

[0597] Kits can include one or more containers comprising an antibody (or unit dosage forms and/or articles of manufacture). In some embodiments, a unit dosage is provided wherein the unit dosage contains a predetermined amount of a composition comprising an antibody (e.g., a therapeutically effective amount), with or without one or more additional agents. In some embodiments, such a unit dosage is supplied in single-use prefilled syringe for injection. In some embodiments, the composition comprising the antibody or

antigen binding fragment thereof can comprise saline, sucrose, or the like; a buffer, such as phosphate, or the like; and/or be formulated within a stable and effective pH range. In some embodiments, the antibody or antigen binding fragment thereof can be provided as a lyophilized powder that may be reconstituted upon addition of an appropriate liquid, for example, sterile water. In some embodiments, the antibody or antigen binding fragment thereof further comprises one or more substances that inhibit protein aggregation, including, but not limited to, sucrose and arginine. In some embodiments, the antibody or antigen binding fragment thereof further comprises heparin and/or a proteoglycan.

[0598] In some embodiments, kits further comprise instructions for use in the treatment of cancer in accordance with any of the methods described herein. The kit may further comprise a description of selection an individual suitable or treatment. Instructions supplied in the kits are typically written instructions on a label or package insert (for example, a paper sheet included in the kit), but machine-readable instructions (for example, instructions carried on a magnetic or optical storage disk) are also acceptable. In some embodiments, the kit further comprises another therapeutic agent (e.g., an anti-cancer antibody or a chemotherapeutic agent)

[0599] The kits are in suitable packaging. Suitable packaging includes, but is not limited to, vials, bottles, jars, flexible packaging (for example, sealed Mylar or plastic bags), and the like. Kits may optionally provide additional components such as buffers and interpretative information. The present application thus also provides articles of manufacture, which include vials (such as sealed vials), bottles, jars, flexible packaging, and the like.

EXAMPLES

Example 1. Engineered Therapeutic IgA Antibodies

[0600] A first wild type IgA2 antibody was engineered to incorporate the following amino acid substitutions and deletions: N45.2G, P124R, C86S, N114T, I115L, T116S, deletion of C147, and deletion of Y148, numbering according to IMGT scheme.

[0601] A second wild type IgA2 antibody was engineered to mutate the asparagine at position 135 to a glutamine. The N135Q mutation deletes an N-glycosylation site of the IgA2 antibody. The N135Q mutation can be combined with additional mutations of N45.2G, P124R, C86S, N114T, I115L, T116S, deletion of C147, and deletion of Y148, numbering according to IMGT scheme.

[0602] A third wild type IgA2 antibody was engineered to delete the entire tail piece, amino acids 131-148 (PTHINVSVMMAEADGTCY) (SEQ ID NO: 9). The N135 N-glycosylation site of the IgA2 antibody. The tail piece deletion can be combined with additional mutations of N45.2G, P124R, C86S, N114T, I115L, and T116S, numbering according to IMGT scheme.

- [0603] Table 1 details a comparison of the percent identity, percent positives, gaps, and total length of wild type IgA2 (IgA2_Valerius_hgnc_id=HGNC:5479); an engineered IgA2 antibody comprising the following amino acid substitutions and deletion: N45.2G, P124R, C86S, N114T, I115L, T116S, deletion of C147, and deletion of Y148 (IgA2 - 2.0) (numbering according to IMGT scheme); an engineered IgA2 antibody described herein comprising a N135Q mutation (IgA3.0 glycosylation mutated hgnc id=HGNC:5470); an engineered IgA2 antibody described herein comprising a deletion of the tail piece amino acids 131-148 (IgA2 - 3.0_tailpiece_del_); IgA2 (tr|A0A0G2JMB2|A0A0G2JMB2 HUMAN); IgA2 (sp|P01877|IGHA2_HUMAN); and IgA2 (tr|A0A286Y5Y5|A0A286Y5Y5 HUMAN).

- 10 [0604] Table 1. Comparison of Engineered Antibody Amino Acid Sequences

Name	Alternative name	Identities (%)	Positives (%)	Gaps	Total length
IgA2_Valerius_hgnc_id=HGNC:5479:		100	100	0	340
IgA2 - 2.0		332/338(98%)	334/338(98%)	0	338
IgA2 - 3.0 glycosylation_mutated_hgnc_id=HGNC:5479	IgA3.0+	331/338(98%)	333/338(98%)	0	338
IgA2 - 3.0_tailpiece_del_	IgA3.0min	316/322(98%)	318/322(98%)	0	322
IgA4.0_NG		315/322(98%)	317/322(98%)	0	322
IgA4.0_NQ		315/322(98%)	317/322(98%)	0	322
IgA4.0_NT		315/322(98%)	317/322(98%)	0	322
IgA4.0_NLT-TIS		313/322(97%)	317/322(98%)	0	322
tr A0A0G2JMB2 A0A0G2JMB2_HUMAN		100	100	0	340
sp P01877 IGHA2_HUMAN		333/340 (98%)	337/340 (99%)	0	340
tr A0A286Y5Y5 A0A286Y5Y5_HUMAN		316/321(98%)	319/321(99%)	0	391

Example 2 Engineered IgA variants

- [0605] Protein glycosylation plays a major role in function and serum half-life of an antibody in circulation. Incomplete glycosylation of IgA leads to clearance by dedicated receptors. The main receptor for the internalization of incompletely glycosylated proteins is the asialoglycoprotein receptor (ASGRP1). This receptor recognizes terminal galactose on N-glycosylated proteins, and is an important mediator of IgA clearance from serum. The IgA2(m1) antibody has a complicated N-linked glycosylation pattern formed by a total of 4 N-linked glycosylation sites (CH1-N45.2, CH2-N15.2, CH2-N114, CH3_CHS-N135). To enhance half-life of IgA2(m1) critical N-glycosylation motifs have been engineered to only contain either a single N-

glycosylation motif or none at all. A single N-linked glycosylation site resulted in a more complete glycosylation profile increasing half-life and subsequently less antibody clearance. Yet, a fully aglycosylated IgA will not be subjected to glycosylation-dependent clearance at all, considerably augmenting half-life. Moreover, this aglycosylated variant will allow production in lower organisms, significantly increasing yields and reducing manufacturing costs. Hence, single- or non-glycosylated antibodies will simplify biopharmaceutical manufacturing and be an ideal candidate for therapeutic purposes.

Design of engineered IgA variants

[0606] The genetic sequence of the IgA2(m1) heavy chain was the backbone for IgA engineering and N-linked glycosylation motifs were silenced, and stabilizing mutations were introduced (Table 2; FIGs. 4A-4D). This resulted in a series of molecules: IgA3.0-(min), IgA3.0+(plus) and IgA4.0.

IgA3.0+variants

[0607] The IgA3.0+ molecule was modified to allow for covalent binding between the heavy- and light-chain by the CH1-P124R mutation. Two Cysteine residues were modified or removed (CH2-C86S; CH3_CHS-C147del_Y148del) to prevent the formation of cysteine bridges with serum proteins, preventing dimeric aggregates and/or complex formation. Furthermore, three N-linked glycosylation motifs were silenced, by substitution of critical amino acids in these motifs (CH1-N45.2G; CH2-N114T; CH3_CHS-N135Q).

IgA3.0- or IgA3.0min variants

[0608] The IgA3.0min molecule contains the same mutations as the CH1 and CH2 in IgA3.0+, but in contrast to IgA3.0+, in IgA3.0min almost the entire tailpiece was deleted (CH3_CHS-P131-Y148del).

IgA4.0 variant or aglycosylated variant

[0609] The IgA4.0 molecules were created by silencing the only remaining N-linked glycosylation motif (CH2-N15.2) by 4 separate amino acid substitutions, thereby creating 4 fully aglycosylated IgA2 molecules.

25

Table 2 lists the mutations in engineered IgA variants; IgA3.0+, IgA3.0min, and IgA4.0 relative to the wild type (WT) IgA2(m1)

Domain	Modification	Description	IgA2 (m1)	IgA3.0 +	IgA3.0 min	IgA4.0
CH1	N45.2G	Remove N-linked glycosylation motif	X	✓	✓	✓
	P124R	Covalent heavy-/light-chain interaction	X	✓	✓	✓
CH2	N15.2G	Remove N-linked glycosylation motif	X	X	X	✓
	N15.2Q	Remove N-linked glycosylation motif	X	X	X	✓
	N15.2T	Remove N-linked glycosylation motif	X	X	X	✓
	L15.3I	Remove N-linked glycosylation motif				
	T16S	Remove N-linked glycosylation motif				
	C86S	Prevent cysteine bridging	X	✓	✓	✓
	N114T	Remove N-linked glycosylation motif	X	✓	✓	✓
	I115L	Remove N-linked glycosylation motif				
	T116S	Remove N-linked glycosylation motif				
CH3-CHS	N135Q	Remove N-linked glycosylation motif	X	✓	X	X

	C147del	Deletion preventing cysteine bridging	X	✓	X	X
	Y148del	Deletion	X	✓	X	X
	P131-Y148del	Deletion tailpiece	X	X	✓	✓

Example 3

Methods

Cloning IgA3.0/IgA4.0

- 5 [0610] To clone the IgA3.0+ and IgA3.0min molecules, synthetic DNA covering the entire constant region of IgA3.0 (CH1-CH2-CH3-CHS) was ordered (Baseclear, Leiden NL) and cloned in the pEE14.4 vector, replacing the IgA2(m1) sequence. In some cases of IgA3.0min molecules, gBlocks (IDT) covering both the variable and constant regions of IgA3.0min (VH-CH1-CH2-CH3-CHS) were cloned into the pcDNA3.4 vector. Based on the IgA3.0min vector, IgA4.0 molecules were cloned by replacing the CH1-CH2-CH3-CHS
- 10 region of IgA3.0min in the pcDNA3.4 vector with gBlocks (IDT) containing the proper mutations for IgA4.0.

Production

- [0611] For the production of the IgA3.0+ and IgA3.0min HEK293F cells (Thermo Fisher) were used
- 15 according to the manufacturer's instructions. For DNA complexing, separate vectors for IgA3.0 + or IgA3.0min heavy chain (HC), along with vectors for kappa light chain (LC) and pAdvantage (pAdv; Promega) were mixed in optimal HC:LC:pAdv ratios and complexed with 293fectin prior to transfection.
- [0612] For the production of the IgA3.0min and IgA4.0 ExpiCHO-S cells (Thermo Fisher) were used according to manufacturer's instructions. For DNA complexing, separate vectors for IgA3.0+ or IgA3.0min
- 20 or IgA4.0 heavy chain, along with vectors for kappa light chain and pAdvantage (Promega) were mixed in optimal HC:LC:pAdv ratios and complexed with Expifectamine prior to transfection.

Purification

- [0613] The procedure for the purification of IgA was identical for all IgA variants, and is performed by
- 25 capturing kappa light chains from clarified and filtered cell culture supernatant using FPCL with a HiTrap KappaSelect column (GE Healthcare), followed by a separation by a Size Exclusion Chromatography procedure using a HiPrep 26/60 Sephacryl S-300 HR column.

Binding assays

[0614] To determine binding of the engineered antibodies, both the binding of the variable portion and the Fc portion was assessed.

5 [0615] Clarified supernatants of IgA3.0min-Obinituzumab (IgA3.0min-Obi) or IgA3.0+- Obinituzumab (IgA3.0+ -Obi) from HEK293F productions were tested on CD20-expressing Daudi cells in a FACS binding experiment. Clarified supernatants of Obinituzumab-IgA4.0 from ExpiCHO-S productions were used undiluted in a FACS binding experiment.

10 [0616] In short, supernatants were incubated with Daudi cells for 1 hour on ice. After washing, cells were incubated with PE-labeled anti-IgA antibody (Southern Biotech) for 45 minutes. After washing cells were fixed with PFA and measured on a Canto II (BD). CD20 recognizing control antibodies were added at concentration of 5-10ug/mL.

15 [0617] To determine whether the antibody Fc-region is able to bind to Fc α R, IgA2(m1) and IgA3.0min-Obi (25 μ g/mL) were coated overnight in an ELISA plate in carbonate buffer pH9.0. Plates were blocked with 1% BSA after which CD89-expressing Calcein-labeled healthy donor polymorphonuclear neutrophils (PMN) were allowed to bind the plate for 45 minutes at 37°C. Subsequently, after every two washing steps, binding was determined by comparing the remaining signal to the input signal (no wash). An additional control to determine if coating concentrations were equal, was performed by staining an ELISA plate overnight with serially diluted IgA2(m1) and IgA3.0min antibody after which their presence was detected with anti-hIgA-HRP (Southern Biotech).

20

ADCC

[0618] Target cells were loaded with ⁵¹Cr (Perkin-Elmer), washed twice and incubated with serially diluted IgA antibodies and healthy donor PMN's for 4 hours. Chromium release was measured in the supernatant and specific lysis was calculated using following the formula: ((experimental cpm – basal cpm) / (maximal cpm – basal cpm)) x 100, with maximal lysis determined by incubating labelled cells with 1.25% triton and minimal lysis in the absence of antibodies and effector cells. In the case of IgA4.0, undiluted supernatants of ExpiCHO-S productions were assessed.

25

Thermostability

30 [0619] Thermal stability was analyzed in thermal shift assay using Sypro Orange (Life Technologies). A total of 12,5 μ g of antibody diluted in 25 μ L PBS and 3x SYPRO Orange (final concentration) was transferred to a white 96-well thin-wall PCR plate (Roche) and sealed with Optical-Quality Sealing Tape (Roche). Plates were heated in a ViiA7 (Roche) from 37°C to 99°C with a heat-rate of 1.6°C/second and 1

minute incubation at each degree. Fluorescence was recorded simultaneously using 490 and 575 nm as excitation and emission wavelengths, respectively.

[0620] To assess functionality of destabilized IgA molecules, antibodies in PBS were incubated at different temperatures (4°C, 23°C-95°C in incremental steps of 12°C) for 5 minutes in a thermal cycler. After incubation complete medium was added and antibodies were used directly in an ADCC at a final concentration of 10 µg/mL.

Glycosylation analysis

[0621] PNGase F treatments were performed according to manufacturer's instructions (NEB). In short, antibodies were first denatured for 10 minutes at 100°C, and subsequently NP-40, Glycobuffer and PNGase F enzyme were added and incubated for 1 hour at 37°C. Samples were taken in Laemmli buffer with 20mM DTT and run on a 10% Mini Protean TGX SDS-PAGE (Bio-Rad). Gels were stained with InstantBlue (Expedeon) for 10 minutes, and rinsed with water.

[0622] Glycan identification and quantification was determined by mass spectrometry. For the Her2-IgA2(m1) a reflectron positive mode MALDI-TOF-MS analysis of the released N-glycosylation of IgA2 antibodies after linkage-specific sialic acid derivatization was performed as described before (Meyer, Nederend et al. 2015). For the IgA3.0min-Her2, IgA3.0min-Obi, and IgA4.0-Obi variants an LC/MS2 approach has been taken.

[0623] Antibodies were denatured, reduced, alkylated and proteolytically digested with GluC (Roche (Indianapolis, IN)) and trypsin (Sigma-Aldrich (Steinheim, Germany)). For this, 10 µg of antibody was brought to 100 mM Tris-HCl (pH 8.5), 5 mM Tris(2-carboxyethyl)-phosphine (TCEP, Sigma-Aldrich (Steinheim, Germany)), 30 mM chloroacetamide (CAA, Sigma-Aldrich (Steinheim, Germany)) and 1% sodium deoxycholate (SDC, Sigma-Aldrich (Steinheim, Germany)), water (MQ) (generated from a Q-POD or Q-Gard 1 system (Millipore), operated at ≥ 18.2 MΩ). This mixture was incubated for 4 h at 37 °C with GluC with an enzyme:protein ratio of 1:75 w/w, followed by an overnight incubation at 37 °C with trypsin (1:100 w/w). Hereafter, the SDC was precipitated by bringing the samples to 0.5% trifluoroic acid (TFA, Sigma-Aldrich (Steinheim, Germany)) and centrifugation at maximum speed for 10 min. The supernatant was collected for solid-phase extraction (SPE).

[0624] For SPE, the use of an Oasis µElution HLB 96-well plate (Waters, Wexford, Ireland) positioned on a vacuum manifold was made. The plate was conditioned with acetonitrile (ACN, BioSolve Valkenswaard, The Netherlands) equilibrated with 0.5% TFA, loaded with the supernatant, washed with 0.5% TFA, and peptides were finally eluted with 50% ACN 0.5% TFA. The recovered eluate was dried by means of rotary evaporation, and reconstituted in 2% formic acid for subsequent LC-MS2 analysis.

- [0625] For each of the digested and desalted samples, 100 ng was analyzed by use of an Agilent 1290 Infinity HPLC system (Agilent Technologies, Waldbronn, Germany), equipped with flow-splitter to achieve nanoflow, hyphenated to an Orbitrap Fusion Tribrid mass spectrometer (Thermo Fisher Scientific, Bremen, Germany). The samples were separated on a 2 cm trap column (100 μ m inner diameter, packed with 3 μ m ReproSil-Pur C18-AQ; Dr. Maisch GmbH, Ammerbuch-Entringen, German) coupled to a 50 cm analytical column (50 μ m inner diameter, packed with 2.7 μ m Poroshell 120 EC-C18; Agilent Technologies, Amstelveen, The Netherlands). Buffer A consisted of 0.1% formic acid, buffer B of 0.1% formic acid in 80% ACN. The LC gradient was as follows: 0-5 min: 100% A (the rest being B), 5-53 min: 87% A to 60% A, 53-58 min 0% A, 58-65 min 100% A.
- 10 [0626] Mass spectrometry was performed in positive ion mode, with electrospray ionization from a coated fused silica emitter at 2 kV spray voltage. Each sample was measured in triplicate, using the same MS1 acquisition methods, but different MS2 methods. For the MS1 scans, the mass range was set from m/z 350 to 2000 with a resolution of 60,000, an AGC target of 400,000 with maximum injection time of 50 ms. Each of the three MS2 methods initiated HCD fragmentation (30% normalized collision energy; NCE) on the highest
- 15 charge state, lowest m/z signals within a 3 s cycle time, using an exclusion time of 30 s. MS2 by HCD was recorded with a resolution of 30,000 from m/z 120 to 4000, with an AGC target of 50,000 and a maximum injection time of 50 ms. For MS2 method 1, only HCD fragmentation was performed. For MS2 method 2, the detection of at least 3 oxonium ions (Hex: 127.0390, 145.0495, 163.0601; HexNAc: 138.0550, 168.0655, 186.0761, 204.0867; PhosphoHex: 243.0264; NeuAc: 274.0921, 292.1027; Complex: 366.1395, 405.0793,
- 20 407.1660, 512.1974, 657.2349) within the HCD spectrum triggered stepping-HCD on the same precursor signal, combining the HCD fragments with NCEs of 10%, 25% and 40%. Stepping-HCD was recorded with a resolution of 30,000 from m/z 120 to 4000, with an AGC target of 200,000 and a maximum injection time of 250 ms For MS2 method 3, detection of the oxonium ions triggered EThcD (30% supplemental activation), which was recorded with a resolution of 30,000 from m/z 120 to 4000, with an AGC target of
- 25 200,000 and a maximum injection time of 250 ms.
- [0627] Bottom-up data was interpreted with Byonic v3.3.11 (Protein Metrics Inc.). Raw data was searched with C-terminal cleavage sites at Arg and Lys (trypsin), and Glu and Asp (GluC). 3 missed cleavages were allowed, using a precursor mass tolerance of 10 ppm and fragment mass tolerance of 20 ppm. Cys carbamidomethylation was included as fixed modification and Met oxidation as variable modification. For
- 30 N-glycosylation 279 compositions were included following the pathways for N-glycan biosynthesis.
- [0628] Skyline (v3.7.0.11317) was used to perform relative quantification. For this, each of the peptides found glycosylated by Byonic were integrated, which included all major miscleavages and oxidation variants. For each of these, the aforementioned 279 glycan compositions were integrated from each LC-MS2 run. Integrations obtained as such were subsequently curated to adhere to the following criteria: 1) ≤ 5 ppm error

to the theoretical mass, 2) having an idotp of ≥ 0.85 with the theoretical isotopic pattern, 3) eluting within ± 2 min of the mean retention time for that peptide, 4) having no apparent overlapping isotopic patterns. The resulting list of glycopeptides was in agreement with the annotation by Byonic, and further used for relative quantification. For each of the curated peptide glycoforms, the MS1 areas were integrated and peptides
5 informing on the same N-glycosylation site combined. As alternative way of quantification, the number of peptide spectrum matches (PSMs) was counted for each glycopeptide combination that informed on a given glycosylation site.

[0629] For visualization of glycan species, the recommendations of the Consortium for Functional Glycomics were followed. Glycan cartoons were constructed using GlycoWorkbench (v2.1 build 146). For
10 native MS analysis antibodies were buffer exchanged to 150 mM ammonium acetate pH 7.5, using a Vivaspin 500 30 kDa molecular weight cut-off filter (Sartorius Stedim Biotech, Germany) by 10x 15 min centrifugation at 15.000 x g. Following buffer exchange, the antibodies were adjusted to an approximate concentration of 3 μ M with 150 mM ammonium acetate pH 7.5.

[0630] Native MS was performed on a modified Exactive Plus Orbitrap instrument with extended mass
15 range (EMR) (Thermo Fisher Scientific, Bremen), calibrated using a 25 mg/mL CsI solution. Voltage settings of the transport multipoles and ion lenses were manually optimized to provide good transmission at the required m/z range. Electrospray ionization was achieved from a gold-coated glass capillary, employing a capillary voltage of 1.2 kV, while the MS was operated with a source fragmentation of 80 V, a source temperature of 250 °C, a collision energy of 80 V, and a resolution of 35000 at m/z 200. Desolvation of
20 molecules was further achieved by addition of nitrogen to the HCD cell to reach a gas pressure of approximately 3.7×10^{-10} bar. Masses were calculated from the charge state distributions by fitting the charges to the lowest standard deviation on the mass (typically resulting in mass errors lower than 1 Da).

Pharmacokinetic/Pharmacodynamic studies

25 [0631] For pharmacokinetics/pharmacodynamics of IgA BALB/cByJ mice (Jackson Laboratory) were injected i.v. with 100 μ g (1mg/mL) of antibody. Blood was collected at the indicated time points, centrifuged to separate and collect serum. ELISAs were performed on diluted serum.

Biodistribution

30 [0632] Antibodies were conjugated to a chelator prior to radio-labeling. Antibody was spun through a 30 kDa centrifugal filter at 12,000 g for 8min. Subsequently, 500 μ L of 0.1 M sodium bicarbonate (pH 8.2) was added to the centrifugal filter and spun again at 12,000 g for 8 min. The centrifugal filter was inverted into a new tube and centrifuged at 1,000 g for 2 min, yielding about 40 μ L to which 60 μ L of sodium bicarbonate buffer was added, resulting in 100 μ L. A 20-fold molar excess of p-SCN-Bn-DTPA (2 mg/mL in dry DMSO

(used 1 mg in 500 μ L)) was added and vortexed for 30 sec, and incubated 1 hour at 37°C. Next, the BnDTPA-IgG1-dinutximab or BnDTPA-IgA3.0min-dinutximab conjugate was run through a G50 column, eluting with 0.5 M MES buffer (pH 5.4) in 100 μ L fractions for 12 fractions. The 5 most concentrated fractions were pooled and spun through a 30 kDa centrifugal filter at 12,000 g for 8mins. A volume of 500 μ L 0.5 M MES buffer (pH 5.4) was added to the centrifugal filter and centrifuged for further 8 min at 12,300 g. The centrifugal filter was inverted into a new microcentrifuge tube and centrifuged at 1,000 g for 2 min. Protein concentration was measured by Nanodrop.

[0633] To radiolabel the antibody 150 μ L Indium solution (55.5 MBq) was added to 150 μ g antibody in MES buffer in a microcentrifuge tube and incubated for 45mins at room temperature. The reaction mixture was run through a G50 column, and eluted with PBS (pH 7.4). Sample is collected in a microcentrifuge tube in 100 μ L fractions (3drops) for 16 fractions, activity is checked and highest fractions were combined.

[0634] To check purity of the radiolabeled antibody by an iTLC strip, a total of 2 μ L of the reaction mixture was pipetted onto the iTLC and dried for 2 minutes. A volume of 0.1 M citrate buffer was added to measurement cylinder covering the bottom. The iTLC strip was placed into measurement cylinder and allowed to draw the citrate buffer up until it is about 1 cm from the top. The strip was removed and scanned using radio-TLC.

In vivo i.p model

[0635] Mice were maintained in the animal facility of the University of Utrecht. Experiments were conducted using female hCD89 Tg or hCD89 NTg mice backcrossed on a CB17/ lcr-Prkdcscid/lcrIcoclrl (Charles river) background. Mice were housed in groups of 5 under a 12:12 light dark cycle, with food and water available ad libitum. All experiments were performed in accordance with international guidelines and approved by the national Central Authority for Scientific Procedures on Animals (CCD) and the local experimental animal welfare body.

[0636] An amount of 1×10^5 A431-Luc2-Her2 cells was injected i.p. on day 0 followed by a s.c. injection with 20 μ g pegG-CSF (Amgen) on day 6. Bioluminescence (BLI) signal was measured on a U-OI system (MiLabs) 10 minutes after i.p. injection of 2.5mg Luciferin (Promega) in PBS. On day 6 mice were randomized into the different treatment groups. Treatment started on day 7 with daily i.p. injections of 10 μ g IgA3.0min-Her2 for 10 days. BLI signal was measured on the indicated time points.

Example 4

Results

[0637] Plasmids encoding IgA2, IgA3.0min, IgA3.0+, and IgA4.0 molecules against various targets were transfected in HEK293F and/or ExpiCHO-S cells. The IgA3.0min-C47A8-CQ antibody was expressed to

similar or better levels as wild type IgA2(m1)-Her2 as shown in FIGs.5A-5B, where different HC:LC:pAdv ratios have been tested in HEK293F cells. This also holds true for IgA3.0+.

[0638] As a comparison between production cell lines, IgA3.0- molecules (e.g, IgA3.0min-Obi, IgA3.0min-Her2, IgA3.0min-mCTLA4, targeting three different antigens CD20, Her2, and mCTLA4 respectively) have been produced in both HEK293F and ExpiCHO-S cells FIGs.6A-6C. Expression levels of IgA3.0min molecules were higher in ExpiCHO-S cells, where they reached almost 1g/L as determined by sandwich ELISA on supernatants. In addition, different IgA3.0min molecules (IgA3.0min-ch14.18, IgA3.0min-TA99, IgA3.0min-Her2, IgA3.0min-Obi, targeting various antigens GD2, gp75, Her2, CD20 respectively have been produced successfully in ExpiCHO-S cells (FIGs. 7A-7D). These data indicate that engineered IgA3.0min and IgA3.0+ variants were expressed properly, independent of production system choice.

[0639] Also, all the four aglycosylated IgA4.0 variants expressed in ExpiCHO-S cells showed similar expression levels as IgA3.0min antibodies (FIGs. 8A-8D). This indicates that none of the introduced mutations prevent expression of engineered IgA4.0 variants, and that a fully aglycosylated IgA antibody was properly expressed.

[0640] Subsequent kappa light chain dependent purification of IgA3.0min variants; IgA3.0min-Obi, (FIGs. 9A-9B)) and IgA3.0min-Her2 (FIGs. 10A-10B) followed by their respective SEC separation (FIGs. 9C-9D for IgA3.0min-Obi, and FIGs. 10C-10D demonstrating IgA3.0min-Her2, revealed no aggregates that are normally seen with wild type IgA2(m1) due to cysteine in the tailpiece. By removing this cysteine (IgA3.0+) or the entire tailpiece (IgA3.0min) no aggregation occurred. Likewise, no aggregate formation was observed in the SEC separation of IgA4.0 variant; IgA4.0-Obi (FIGs. 11A-11B), which like IgA3.0min variants also lacks the tailpiece.

[0641] Next, the binding capacities of the IgA3.0+-Obi and IgA3.0min-Obi molecules was determined by flow cytometry on Daudi cells. As expected, both the IgA3.0+-Obi as well as the IgA3.0min-Obi antibodies were able to bind to CD20 on Daudi cells, indicating their Fab fragments were properly folded (FIGs. 12A-12E). Next, to assess whether the fully aglycosylated IgA4.0 variants were also still able to bind target antigen, a flow cytometric assay was performed using CD20 complemented SKBR3 cells (FIG. 15). These FACS binding data indicated engineered IgA4.0 variants (IgA4.0_NT-Obi, IgA4.0_NQ-Obi, IgA4.0_NG-Obi, IgA4.0_NG-Obi, and IgA4.0_NLT-TIS-Obi) bind CD20 as well as the engineered IgA3.0min molecules, confirming the folding of the Fab regions are unaffected by the engineering.

[0642] To assess whether the Fc portion still retains the ability to bind Fc α R, an IgA-Fc α R binding assay has been performed where neutrophils were allowed to bind IgA coated plates followed by a series of washes. After a number of washes neutrophils were still capable of binding the IgA molecules, where IgA3.0min-Obi show similar or better binding profiles than IgA2(m1) (FIG. 13). To ensure the amount of

IgA molecules presented to the neutrophils was equivalent, an IgA detection step was executed indicating equal concentrations of antibody of IgA2(m1) and IgA3.0min (FIG. 14).

[0643] To demonstrate engineered IgA molecules are functional and are able to induce PMN-mediated ADCC, chromium release assays were performed. On SKBR3-CD20 cells the IgA3.0min-Her2 antibody was able to induce ADCC as good as, or better than IgA2(m1)-Her2 (FIG. 16A). Moreover, IgA3.0min-Obi antibodies produced in ExpiCHO-S cells induced ADCC in both Daudi and Ramos cells as efficient as with HEK293F-produced antibodies (FIGs. 16B-16C). Furthermore, the fully aglycosylated IgA4.0 variants induced ADCC to similar levels as IgA3.0min-Obi (FIG. 16D), showing full functionality with regards to killing. This is in stark contrast to aglycosylated IgG1 which functionality is severely compromised or ablated (Jefferis 2009). These data indicate that, despite the modifications introduced for engineering IgA variants, both IgA3.0min as well as IgA4.0 still retained their function in inducing ADCC.

[0644] In order to see whether the engineered IgA3.0min and IgA4.0 variants were more thermally stable, a SYPRO thermal shift was executed. Antibodies were incubated in the presence of SYPRO Orange in a thermal cycler slowly raising its temperature. When the protein unfolds, SYPRO Orange is able to bind hydrophobic pockets and fluorescent signal can be detected. FIG. 17A shows the thermal stability profiles of two wild type IgA2(m1) antibodies versus two engineered IgA3.0min molecules and four engineered IgA4.0 antibodies. The shift in the curves indicate IgA3.0min variants and IgA4.0 variants were more stable at higher temperatures than wild type IgA2(m1). The T_m values calculated by a non-linear regression analysis have been averaged for the different IgA molecules assessed (FIG. 17B). Both types of engineered IgA variants (IgA3.0min and IgA4.0) show improved stability at higher temperatures compared to wild type IgA2(m1).

[0645] To translate the observed stability into functionality anti-CD20 IgA antibodies; wild type IgA2 and IgA3.0min-Obi were heated to certain temperatures and tested in a chromium release ADCC using Raji cells (FIG. 17C). Signals were normalized to the control which was antibody incubated at 4°C. The killing efficiency of thermally exposed antibodies differed between the wild type IgA2(m1) and engineered IgA3.0min/IgA4.0 molecules. The first showing a decremental profile at exposure at 47°C or higher, while the latter still showed over 50% activity after exposure 71°C. All activity of the three antibodies tested was lost at temperatures higher than 83°C. This indicates that the stability of the antibodies is important in terms of functionality, and the improved thermostability of the engineered IgA variants relative to wild type is advantageous.

[0646] As the engineered IgA3.0min and IgA4.0 molecules have less or no sites for N-linked glycosylation in comparison to wild type IgA2(m1), a PNGase F assay has been performed to visualize remaining N-glycosylation (FIG. 18). For this, IgA molecules were treated with PNGase F which removes all N-linked glycosylation by cleaving between the innermost GlcNac and asparagine. Subsequently, samples were run on

a reducing SDS-PAGE gel. As expected, the wild type IgA2(m1) (HEK293F-produced) shows a major shift in molecular weight, due to the presence of four N-linked glycans, while the shift of IgA3.0min-Obi (HEK293F-produced) is minor due to only as single glycosylation event. Yet, the aglycosylated IgA4.0 variants (ExpiCHO-S-produced) does not show a shift at all, indicating this antibody is not glycosylated.

- 5 **[0647]** To assess the glycosylation profiles in more detail, antibodies were analyzed by mass spectrometry. The glycosylation profiles of Wild type IgA2(m1)-Her2 and IgA3.0min-Her2 (FIGs. 19A-19B) were analyzed. Several N-glycan species were detected by MALDI-TOF-MS analysis in IgA2(m1)-Her2 reflecting the overall glycosylation on the four N-linked glycosylation motifs. Yet, the peaks of the IgA3.0min-Her2 profile obtained from a LC/MS2 analysis only came from the single N-linked glycosylation motif left, thus
- 10 also reflecting the overall glycosylation profile as no other sites have been modified. When comparing antibodies from the HEK293F and ExpiCHO-S antibody production systems by LC/MS2 a substantial difference in glycosylation profiles was observed (FIG. 19C). In ExpiCHO-S the majority (>50%) of the glycosylated antibodies were terminated on bi-antennary GlcNac while the HEK293F showed a more diverse array of glycosylation species of the hybrid or complex type. The unfavorable free galactoses were markedly
- 15 more present in the HEK293F-produced antibodies than in ExpiCHO-S-produced antibodies. These results showed ExpiCHO-S-produced antibodies showed less heterogeneous glycosylation and will have better half-life characteristics with regard to ASGPR-induced clearance. To determine whether glycosylation events occur in IgA4.0 molecules a native MS approach was taken, analyzing full, undigested antibodies. All four IgA4.0 variants displayed a mass very close to the theoretical peptide backbone (FIGs. 19D-19G). There
- 20 is a consistent loss of 36 Da amongst all four variants, which could not be attributed to N-glycan species, as these are typically 1500-2500 Da per glycan. The observed mass difference is likely to be formation of two pyroglutamates accounting for 36 Da. Thus, this analysis showed that the IgA4.0 antibody variants lack N-glycans on their backbone, demonstrating that aglycosylated IgA antibodies are fully functional in in vitro assays.
- 25 **[0648]** To determine whether the loss of glycosylation sites in engineered IgA3.0 variants and IgA4.0 variants lead to a better half-life in serum, a pharmacokinetics study was set-up. BALB/c mice were i.v. injected with 100µg antibody and analysed serum samples at the indicated timepoints by ELISA. Two antibodies in different formats, dinutuximab and obinutuzumab were assessed. Detected serum wild type IgG1-Dinutuximab levels were higher than wild type IgA2-Dinutuximab levels which was rapidly cleared
- 30 (FIG. 20A). The difference in serum levels between IgA2 and IgA3.0min at 24 hours can be attributed to the clearance by the ASGPR within the first hours after injection. This demonstrated that the distribution phase is less pronounced in IgA3.0min, due to less N-glycosylation-dependent clearance by ASGPR. The data showed an increased serum half-life was observed with the IgA3.0min- Dinutuximab antibodies as compared to wild type IgA2- Dinutuximab. When assessing the serum half-life of Ig4.0-Obinutuzumab, a clear increase

in serum half-life is observed compared to its IgA3.0min counterpart, IgA3.0min-Obinutuzumab (FIG. 20B). This demonstrates that engineered IgA variants has extended serum-half life relative to their wild type counterpart IgA and thus can leverage a similar effect on tumor using less antibody.

[0649] To define if the IgA3.0min antibody distributes to tumors, a biodistribution experiment was performed where the antibodies were radiolabeled with ¹¹¹Indium to monitor them in vivo. For this, GD2-expressing IMR-32 neuroblastoma cells were subcutaneously injected in NSG mice. After 42 days of outgrowth, mice were injected intravenously with radiolabeled wild type IgG1-dinutuximab and IgA3.0min-dinutuximab. Mice were screened 24 and 48 hours after injection (FIGs. 21A-21B). Radioactive signal was observed both in the liver and in the tumor, indicating antibodies infiltrate tumors. Liver signals represent accumulated antibodies from the bloodstream that are being catabolized. By determining the tumor/liver ratio the tumor-specific signal would be corrected for background. Initially after 24 hours the signals observed are similar, but at 48 hours the IgA3.0min- Dinutuximab signal is more tumor-specific than its IgG1 counterpart (FIG. 22).

[0650] Next the efficacy of IgA3.0min in vivo was determined. For this an established tumor model with hCD89 transgenic and non-transgenic SCID was used (Brandsma, Ten Broeke et al. 2015). Mice were injected i.p. with A431-luc2-Her2 cells. Once tumors were established, mice were treated by administering either PBS or IgA3.0min-Her2 i.v. (FIG. 23). Tumor growth was monitored by BLI on the indicated time points. Tumors showed exponential outgrowth in the PBS group. However, mice treated with IgA3.0min-Her2 showed no outgrowth of tumor and control the cancer after 58 days, demonstrating the efficacy of engineered IgA variants.

[0651] The results demonstrated that compared to wild type IgA2(m1) all engineered IgA variants; IgA3.0+, IgA3.0min, and IgA4.0 show improved biopharmaceutical properties while still retaining their functionality. Engineered IgA variants are expressed to high levels, and can be purified without aggregates, resulting in a pure and more stable IgA antibody that still displays robust binding to target antigen and efficient killing of target cells both in vitro as well as in vivo.

Table 3 lists exemplary amino acid sequences and nucleic acid sequences of the heavy chain constant region of the engineered IgA variants

IgA constant chain	SEQ ID NO:	Amino acid sequence	SEQ ID NO:	Nucleic acid sequence
IgA2m1	1	ASPTSPKVFPLSLDSTPQDGNVV VACLQGFFPQEPLSVTWSESG QNV TARNFPQS QDASGDLYTTSS QLTLPATQCPDGKSVTCHVKHY TNPSQDVTVPVPPPPPCCHPR LSLHRPALEDLLL GSEANLTCTL	22	gccagccccaccagccccaaggtgttccccctgagc ctggacagcaccccccaggacggcaactgtgtgt ggcctgcctgtgtcagggtcttccccaggagcc cctgagcgtgacctggagcgagagcgccagaac gtgaccgccagggaacttccccccagccaggacgc cagcggcgacctgtacaccaccagcagcagctga

		TGLRDASGATFTWTPSSGKSAV QGPPERDLCGCYSVSSVLP QPWNHGETFTCTAAHPELKTPL TANITKSGNTFRPEVHLLPPPSEE LALNELVTLTCLARGFSPKDV LVRWLQGSQELPREKYLTWASR QEPSQGTTFVAVTSILRVAAEDW KKGDTFSCMVGHEALPLAFTQK TIDRLAGKPTHVNVSVVMAEVD GTCY		ccctgcccgccaccagtgccccgacggcaagagc gtgacctgccacgtgaagcactacaccaacccagc caggacgtgacctgcccgtgccccgtgccccccc cccccctgtgcccacccagcgtgacctgcacag gcccgcctggaggacctgctgctgggcagcgagg ccaacctgacctgcacctgacctgacctgaggac gccagcggcgccaccttcacctgacccccagcag cggcaagagcgccgtgcaggcccccccgaggagg gacctgtgaggctgctacagcgtgagcagcgtgctg cccggctgcgccagccctggaaccacggcgaga ccttcacctgcaccggcgccaccccgagctgaaga cccccctgaccgccaacatcaccaagagcggaac acctcaggccccgaggtgcacctgctgccccccc agcgaggagctggccctgaacgagctggtgacct gacctgcctggccaggggcttcagcccaaggacg tgctggtgaggtggctgcagggcagccagagctg cccagggagaagtacctgacctggccagcaggga ggagcccagccagggcaccaccaccttcgccgtga ccagcalcctgagggtggccgccgaggactggaag aaggcgagaccttcagctgcatgtgtggccacga ggccctgcccctggccttcaccagaagaccatcga caggctggccggcaagccaccacgtgaacgtga gcgtggtgatggccgaggtggacggcacctgctac
IgA2m2	2	ASPTSPKVFPLSLDSTPQDGNVV VACL VQGFFPQEPLSVTWSESG QNVTARNFPPSQDASGDLYTTSS QLTLPATQCPDGKSVTCHVKHY TNSSQDVTVPCRVP PPPPCCHPR LSLHRPALEDLLL GSEANLTCTL TGLRDASGATFTWTPSSGKSAV QGPPERDLCGCYSVSSVLP QPWNHGETFTCTAAHPELKTPL TANITKSGNTFRPEVHLLPPPSEE LALNELVTLTCLARGFSPKDV LVRWLQGSQELPREKYLTWASR QEPSQGTTFVAVTSILRVAAED WKKGETFSCMVGHEALPLAFTQ KTIDRMAGKPTHINVSVMMAEA DGTCY	23	gccagccccaccagccccaggtgttccccctgagc ctggacagcaccccccaggacggcaacgtgtgtgt ggccctgcctgggtgagggcttcttccccaggagcc cctgagcgtgacctggagcagagcgccagaac gtgaccgccaggaaacttccccccagccaggacgc cagcgcgacctgtacaccaccagcagcagctga ccctgcccgccaccagtgccccgacggcaagagc gtgacctgccacgtgaagcactacaccaacagcagc caggacgtgacctgcccgtcagggtgccccccc cccccctgtgcccacccagcgtgacctgcacag gcccgcctggaggacctgctgctgggcagcgagg ccaacctgacctgcacctgacctgacctgaggac gccagcggcgccaccttcacctgacccccagcag cggcaagagcgccgtgcaggcccccccgaggagg gacctgtgaggctgctacagcgtgagcagcgtgctg cccggctgcgccagccctggaaccacggcgaga ccttcacctgcaccggcgccaccccgagctgaaga cccccctgaccgccaacatcaccaagagcggaac acctcaggccccgaggtgcacctgctgccccccc agcgaggagctggccctgaacgagctggtgacct gacctgcctggccaggggcttcagcccaaggacg tgctggtgaggtggctgcagggcagccagagctg cccagggagaagtacctgacctggccagcaggga ggagcccagccagggcaccaccacctacgccgtga ccagcalcctgagggtggccgccgaggactggaag aaggcgagaccttcagctgcatgtgtggccacga ggccctgcccctggccttcaccagaagaccatcga

				caggatggccggcaagcccaccacatcaacgtga gcgtggtgatggccgagccgacggcacctgctac
IgA2_A0 A286YE Y5	3	ASPTSPKVFPLSLDSTPQDGNVV VACLVQGFFPQEPLSVTWSESG QNV TARNFPPSQDASGDLYTTSS QLTLPATQCPDGKSVTCHVKHY TNSSQDVTVPCRVP PPPPCCHPR LSLHRPALEDLLL GSEANLTCTL TGLRDASGATFTWTPSSGKSAV QGPPERDLCGCYSVSSVLPGCA QPWNHGETFTCTAAHPELKTPL TANITKSGNTFRPEVHLLPPPSEE LALNELVTLTCLARGFSPKDVL VRWLQGSQELPREKYLTWASR QEPSQGTTTYAVTSILRVAED WKKGETFSCMV GHEALPLAFTQ KTIDRMAGSCCVADWQMPPPY VVLDPQETLEETPGANLWPT TITFLTFLLSLFYSTALT VTSVR GPSGKREGPQY	24	gccagccccaccagcccc aagggtgtccctgagc ctggacagcacccccaggacggcaacgtggtggt ggcctgcctggtgcagggttcttccccaggagcc cctgagcgtgacctggagcgagagcgccagaac gtgaccgccaggaaacttccccccaggcaggacgc cagcggcgacctgtacaccaccagcagcagctga ccctgcccgccaccagtgccccgacggcaagagc gtgacctgccacgtgaagcactacccaacagcagc caggacgtgacctgtccctgcagggtgccccccc ccccccctgtgccacccccaggctgacctgcacag gccccccctggaggacctgtgtgtggcgagcagg ccaacctgacctgcacctgacctgaggagc gccagcggcgccaccttcacctggacccccagcag cggcaagagcgccgtgcaggcccccccgagagg gacctgtgcgctgtacagcgtgagcagcgtgtgt cccggctgcgccagccctggaaccacggcgaga ccttcacctgcaccgcccacccccagctgaaga ccccctgaccgccaacatcaccaagagcgcaac acctcaggccccgaggtgcacctgtgtccccccc agcgaggagctggccctgaacgagctggtgacct gacctgcctggccagggtgtcagccccaggacg tgctggtgaggtggtgcaggcgagccaggagctg cccagggagaagtacctgacctggccagcaggca ggagcccagccaggcgaccaccacctacgccgtga ccagcatcctgagggtggccgccaggactggaag aaggcgagaccttcagctgcatggtggccacga ggccctgcccctggccttaccagaagaccatcga caggatggccggcagctgtgtgtggccgactggc agatgccccccccctacgtgtgtgtgacctgccc aggagaccttgaggaggagacccccggcgccaa cctgtggcccaccaccatcacttctgacctgttcc tgctgacctgttctacagaccgacctgacctgac cagcgtgagggccccagcggaagaggaggagg ccccagctac
IgA3.0+	16	ASPTSPKVFPLSLDSTPQDGNVV VACLVQGFFPQEPLSVTWSESG QGV TARNFPPSQDASGDLYTTSS QLTLPATQCPDGKSVTCHVKHY TNPSQDVTVPCRVP PPPPCCHPR LSLHRPALEDLLL GSEANLTCTL TGLRDASGATFTWTPSSGKSAV QGPPERDLCGCYSVSSVLPGSAQ PWNHGETFTCTAAHPELKTPLT ATLSKSGNTFRPEVHLLPPPSEEL ALNELVTLTCLARGFSPKDVLV RWLQGSQELPREKYLTWASRQE	25	gctagcccaacctctcctaagggtgtccctctgacct ggacagcaccctcaggatggaaatgtgtgtgtg cctgtgtgtgcagggttcttccacaagaccct gtccgtgactggagcgaatctggacaggcgctgac cgccagaaacttcccaccttctcaggacgccttggc gacctgtacaccaccttctcagctgacctgcctgc cacacagtgcctgatggcaagtctgtgacctgccac gtgaagcactacccaatcctagccaggacgtgacc gtgccttgagagttcctcctcctccacctgtgtcac cctcggctgtctgtcacagaccgctctggaagatc tgctgtgtggctctgaggccaacctgacatgtacct gaccggcctgagagatgcttctggcgccaccttacc

		PSQGTTFVAVTSILRVAEDWK KGDTFSCMVGHEALPLAFTQKT IDRLAGKPTHVQVSVM AEVDG T		tggacacctccagcggaaagtcgctgttcaggga cctcctgagaggacgtgctgctgttactctgtgtc tagtgtgctgctggcagcggccagccttggaatcat ggcgagacattacctgtaccgctgctcaccggag ctgaaaacccctctgaccgccacactgtccaagtccg gcaacaccttccggcctgaagtgcactgtgctcctcc acctagcgaggaaactggccctgaatgagctgggtcac cctgacctgtctggccaggggctttagccctaaggac gtgctcgttagatggctgcagggtcccaagagctg cccagagagaagtatctgacctggcctctcggcaa gagccatctcagggcaccacaacctttgccgtgacc agcatcctgagagtggccgccgaagattggaagaa ggcgacaccttcagctgcatgtcggacatgaagc cctgcctctggcttaccagaaaaccatcgacaga ctggccggcaagcccacccatgtccaagtgtctgtt tcatggcggagggtggacggcacc
IgA3.0mi n	17	ASPTSPKVFPLSLDSTPQDGNVV VACL VQGFFPQEPLSVTWSESG QGV TARNFPPSQDASGDLYTTSS QLTLPATQCPDGKSVTCHVKHY TNPSQDVTVPCRVP PPPPCCHPR LSLHRPALEDLLL GSEANLTCTL TGLRDASGATFTWTPSSGKSAV QGPPERDLCGCYSVSSVLP GSAQ PWNHGETFTCTAAHPELKTPLT ATLSKSGNTFRPEVHLLPPPSEEL ALNELVTLTCLARGFSPKDV LV RWLQGSQELPREKYLTWASRQE PSQGTTFVAVTSILRVAEDWK KGDTFSCMVGHEALPLAFTQKT IDRLAGK	26	gctagcccaacctctcctaaggtgttccctctgagcct ggacagcaccctcaggatggaaatgtgtgtgtg cctgtctggtgcaggatcttcccacaagaccct gtccgtgacttgagcgaatctggacaggcgtgac cgccagaaaactcccaccttctcagcagcctctggc gacctgtacaccaccttctcagctgacctgcctgc cacacagtgcctgatggcaagtctgtgacctgccac gtgaagcactacaccaatcctagccaggacgtgacc gtgccttgagagttcctcctcctccacctgtgtcac cctcggctgtctctgcacagaccgctctggaagatc tgctgtggtgctctgaggccaacctgacatgtaccct gaccggcctgagagatgcttctggcgccacctttacc tggacacctccagcggaaagtcgctgttcaggga cctcctgagaggacgtgctgctgttactctgtgtc tagtgtgctgctggcagcggccagccttggaatcat ggcgagacattacctgtaccgctgctcaccggag ctgaaaacccctctgaccgccacactgtccaagtccg gcaacaccttccggcctgaagtgcactgtgctcctcc acctagcgaggaaactggccctgaatgagctgggtcac cctgacctgtctggccaggggctttagccctaaggac gtgctcgttagatggctgcagggtcccaagagctg cccagagagaagtatctgacctggcctctcggcaa gagccatctcagggcaccacaacctttgccgtgacc agcatcctgagagtggccgccgaagattggaagaa ggcgacaccttcagctgcatgtcggacatgaagc cctgcctctggcttaccagaaaaccatcgacaga ctggccggcaag
IgA4.0_ NG	18	ASPTSPKVFPLSLDSTPQDGNVV VACL VQGFFPQEPLSVTWSESG	27	gctagcccaacctctcctaaggtgttccctctgagcct ggacagcaccctcaggatggaaatgtgtgtgtg cctgtctggtgcaggatcttcccacaagaccct

		QGV TARNFPPSQDASGDLYTTSS QLTLPATQCPDGKSVTCHVKHY TNPSQDVTVPCRVP PPPPCCHPR LSLHRPALEDLLL GSEAGLTCTL TGLRDASGATFTWTPSSGKSAV QGPPERDLCGCY SVSSVLPGSAQ PWNHGETFTCTAAHPELKTPLT ATLSKSGNTFRPEVHLLPPPSEEL ALNELVTLTCLARGFSPKDV LV RWLQGSQELPREKYLTWASRQE PSQGT TTF AVTSILRVAEDWK KGDTFSCMVGHEALPLAFTQKT IDRLAGK		gtccgtgacttggagcgaatctggacagggcgtgac cgccagaaacttcccaccttctcaggacgcctctggc gacctgtacaccaccttctcagctgacctgcctgc cacacagtgcctgatggcaagtctgtgacctgccac gtgaagcactacaccaatcctagccaggacgtgacc gtgccttgacaggttctcctctccaccttgcctgcac cctcggctgtctctgcacagaccgctctggaagatc tgctgctgggctctgagggcggcctgacatgtacct gaccggcctgagagatgcttctggcgccacctttacc tggacaccttccagcggaaagtccgctgttcaggga cctcctgagagggacctgtgcggctgttactctgtgc tagtgtgctgcctggcagcggccagccttggaaatcat ggcgagacattcacctgtaccgctgctcaccggag ctgaaaacccctctgaccgccacctgtccaaagtccg gcaacaccttccggcctgaagtgcactgtgcctcc acctagcgaggaaactggccctgaatgagctgtgcac cctgacctgtctggccaggggctttagccctaaggac gtgctcgttagatggctgcagggtcccaagagctg cccagagagaagtatctgacctgggcctctcgga gagccatctcagggcaccacaacctttgcctgacc agcatcctgagagtgccggccgaagattggaagaa ggggcacaccttcagctgcatgtctggacatgaagc cctgcctctggctttaccagaaaaccatcgacaga ctggccggcaag
IgA4.0_ NQ	19	ASPTSPKVFPLSLDSTPQDGNVV VACL VQGFFPQEPLSVTWSESG QGV TARNFPPSQDASGDLYTTSS QLTLPATQCPDGKSVTCHVKHY TNPSQDVTVPCRVP PPPPCCHPR LSLHRPALEDLLL GSEAQLTCTL TGLRDASGATFTWTPSSGKSAV QGPPERDLCGCY SVSSVLPGSAQ PWNHGETFTCTAAHPELKTPLT ATLSKSGNTFRPEVHLLPPPSEEL ALNELVTLTCLARGFSPKDV LV RWLQGSQELPREKYLTWASRQE PSQGT TTF AVTSILRVAEDWK KGDTFSCMVGHEALPLAFTQKT IDRLAGK	28	gctagcccaacctctctaaggtgttcctctgagcct ggacagcaccctcaggatggaaatgtggtgtgg cctgtctggtgcaggatcttccacaagaccct gtccgtgacttggagcgaatctggacagggcgtgac cgccagaaacttcccaccttctcaggacgcctctggc gacctgtacaccaccttctcagctgacctgcctgc cacacagtgcctgatggcaagtctgtgacctgccac gtgaagcactacaccaatcctagccaggacgtgacc gtgccttgacaggttctcctctccaccttgcctgcac cctcggctgtctctgcacagaccgctctggaagatc tgctgctgggctctgagggccagctgacatgtacct gaccggcctgagagatgcttctggcgccacctttacc tggacaccttccagcggaaagtccgctgttcaggga cctcctgagagggacctgtgcggctgttactctgtgc tagtgtgctgcctggcagcggccagccttggaaatcat ggcgagacattcacctgtaccgctgctcaccggag ctgaaaacccctctgaccgccacctgtccaaagtccg gcaacaccttccggcctgaagtgcactgtgcctcc acctagcgaggaaactggccctgaatgagctgtgcac

				cctgacctgtctggccagggccttagccctaaggac gtgctcgttagatggctgcagggctccaagagctg cccagagagaagtatctgacctgggcctctcgga gagccatctcagggcaccacaacctttgccgtgacc agcatcctgagagtggccgccgaagattggaagaa gggcgacaccttcagctgcatggctggacatgaagc cctgcctctggctttaccagaaaaccatcgacaga ctggccggcaag
IgA4.0_ NT	20	ASPTSPKVFPLSLDSTPQDGNVV VACLVQGFFPQEPLSVTWSESG QGV TARNFPPSQDASGDLYTTSS QLTLPATQCPDGKSVTCHVKHY TNPSQDVTVPCRVP PPPPCCHPR LSLHRPALEDLLLGEATLTCTL TGLRDASGATFTWTPSSGKSAV QGPPERDLCGCYSVSSVLPGSAQ PWNHGETFTCTAAHPELKTPLT ATLSKSGNTFRPEVHLLPPPSEEL ALNELVTLTCLARGFSPKDV LV RWLQGSQELPREKYLTWASRQE PSQGTTF AVTSILRVAEDWK KGDTFSCMVGHEALPLAFTQKT IDRLAGK	29	gctagcccaacctctcctaagggtgtccctctgagcct ggacagcaccctcaggatggaaatgtgtgtgtg cctgtctgtgagggattcttccacaagagccct gtccgtgacttgagcgaatctggacagggcgtgac cgccagaaacttccaccttctcaggacgccttggc gacctgtacaccaccttctcagctgacctgcctgc cacacagtgcctgatggcaagtctgtacctgccac gtgaagcactacaccaatcctagccaggacgtgacc gtgccttgcagagtctcctcctccaccttctgtcac cctcggctgtctctgcacagaccgctctggaagatc tgtgtgtggctctgagggccacctgacatgtacct gaccggcctgagagatgtcttggccacctttacc tggacaccttccagcggaaagtcgctgttcaggga cctcctgagaggacgtgtcggtgttactctgtgtc tagtgtgtgctgctggcagcgccagccttggatcat ggcgagacattcacctgtaccgctgtctaccccgag ctgaaaacccctctgaccgccacatgtccaagtccg gcaacaccttccggcctgaagtgcactgtgtgcctc acctagcgaggaaactggccctgaatgagctgtgtac cctgacctgtctggccagggccttagccctaaggac gtgctcgttagatggctgcagggctccaagagctg cccagagagaagtatctgacctgggcctctcgga gagccatctcagggcaccacaacctttgccgtgacc agcatcctgagagtggccgccgaagattggaagaa gggcgacaccttcagctgcatggctggacatgaagc cctgccttggctttaccagaaaaccatcgacaga ctggccggcaag
IgA4.0_ NLT-TIS	21	ASPTSPKVFPLSLDSTPQDGNVV VACLVQGFFPQEPLSVTWSESG QGV TARNFPPSQDASGDLYTTSS QLTLPATQCPDGKSVTCHVKHY TNPSQDVTVPCRVP PPPPCCHPR LSLHRPALEDLLLGEATISCTLT GLRDASGATFTWTPSSGKSAVQ GPPERDLCGCYSVSSVLPGSAQP	30	gctagcccaacctctcctaagggtgtccctctgagcct ggacagcaccctcaggatggaaatgtgtgtgtg cctgtctgtgagggattcttccacaagagccct gtccgtgacttgagcgaatctggacagggcgtgac cgccagaaacttccaccttctcaggacgccttggc gacctgtacaccaccttctcagctgacctgcctgc cacacagtgcctgatggcaagtctgtacctgccac gtgaagcactacaccaatcctagccaggacgtgacc

		WNHGETFTCTAAHPELKTPLTA TLSKSGNTFRPEVHLLPPPSEELA LNELVTLTCLARGFSPKDLVR WLQGSQELPREKYLTWASRQEP SQGTTTFAVTSILRVAAEDWKK GDTFSCMVGHEALPLAFTQKTI DRLAGK		gtgccttgacagagttcctcctccacctgtgtcac cctcggctgtctgcacagaccgctctggaagac tgcgtctgggctctgaggccaccatcagctgtaccct gaccggcctgagagatgcttctggcgccacctttacc tggacacctccagcggaaagtcgctgttcaggga cctcctgagaggacctgtgcggctgtactctgtgtc tagtgtgtgcctggcagcggccagccttggaatcat ggcgagacattcactgtaccgctgtcaccgag ctgaaaacccctctgaccggcacactgtccaagtccg gcaacacctccggcctgaagtgcactgtgtcctcc acctagcgaggaaactggccctgaatgagctgtgtac cctgacctgtctggcaggggctttagccctaaggac gtgtctgttagatgctgcagggtcccaagagctg ccagagagaagtatctgacctgggctctcggcaa gagcatctcagggcaccacaacctttgccgtgacc agcatcctgagagtgccgcccgaagattggaagaa gggcgacacctcagctgcatggtcgacatgaagc cctgcctctggcttcccccagaaaaccatcgacaga ctggccggcaag
--	--	---	--	---

[0652] Table 4 lists exemplary amino acid sequences and nucleic acid sequences of a kappa light chain constant domain of the engineered IgA variants.

Name	SEQ ID NO	Amino acid sequences	SEQ ID NO	Nucleic acid sequences
kappa light chain constant domain	31	RTVAAPSVFIFPPSDEQLKSGTASV VCLLNNFYPREAKVQWKVDNAL QSGNSQESVTEQDSKSTYLSST LTLKADYEEKHKVYACEVTHQGL SSPVTKSFNRGEC	32	cggacagtggccgctcctccgtgttcattctcc cacttccgacgagcagctgaagtccggcaca gtagcgtggtctgcctgctgaacaactctacc ctccgggaagccaaggtgcagtggaggtgga caatgccctgcagtccggcaactcccaagagt ctgtgaccgagcaggactccaaggacagcac ctacagcctgtctccactgacctgtccaa ggccgactacgagaagcacaaggtgtacgcct gcgaagtgacctcagggcctgtctagccct gtgaccaagtcttcaaccggggcgagtg

5 Table 5 lists exemplary lists amino acid sequences of complementarity determining regions of the variable heavy chain of engineered IgA variants

Antigen	Variable heavy chain of Antigen	SEQ ID NO	HC-CDR1	SEQ ID NO	HC-CDR2	SEQ ID NO	HC-CDR3
---------	---------------------------------	-----------	---------	-----------	---------	-----------	---------

	binding domain						
CD20	Obinutuzumab	33	GYAFSYS WIN	41	RIFPGDG DTDYNG KFKG	49	NVFDGYWLV Y
GD2 Disialoganglosi de expressed on tumors of neuroectoderm al origin	Dinutuximab (also called Ch14.18 or Unituxin)	34	GSSFTGY NMN	42	AIDPYYG GTSYNQ KFKG	50	GMEY
Her2	Trastuzumab	35	GFNIKDT YIH	43	RIYPTNG YTRYADS VKG	51	WGGDGFYA MDY
Tyrosinase- related protein- 1 or gp75 Glycoprotein produced by pigmented melanocytes and melanomas	TA99	36	GFNIKDY FLH	44	WINPDNG NTVYDP KFQG	52	RDYTYEKAA LDY
cytotoxic T- lymphocyte- associated protein 4 (mCTLA-4)		37	GYSITSG YGWN	45	FIYYEGS TYYNPSI KS	53	QTGYFDY
CD47	2.3D11	38	SGVSIRSI NWWN	46	EIYHSGS TNYNPSL KS	54	DGGIAVTDY YYYGLDV
CD47	C47A8-CQ	39	SGYTFTS YYMH	47	IINPSGGS TSYAQKF QG	55	STLWFSEFDY
CD20	UMAB10	40	GFTFPSY NLH	48	AIYPGNG DTSYNQ KFKG	56	SAYYGSNVW FFDV

Table 6 lists exemplary lists amino acid sequences of complementarity determining regions of the variable light chain of engineered IgA variant

Antigen	Variable light chain of Antigen binding domain	SEQ ID NO	LC-CDR1	SEQ ID NO	LC-CDR2	SEQ ID NO	LC-CDR3
CD20	Obinutuzumab	57	RSSKSLHNSNGITYLY	65	QMSNLVS	73	AQNLELPYT
GD2	Dinutuximab (also called Ch14.18 or Unituxin)	58	RSSQSLVHRNGNTYLH	66	KVSNRFS	74	SQSTHVPPLT
Her2	Trastuzumab	59	RASQDVNTAVA	67	SASFLYS	75	QQHYTTPPT
Tyrosinase-related protein-1 or gp75	TA99	60	RASGNIYNYLA	68	DAKTLAD	76	QHFWSLP
cytotoxic T-lymphocyte-associated protein 4 (mCTLA-4)		61	KSSQSLFNSNAKTN YLN	69	YASTRHT	77	QQWYDYPYT
CD47	2.3D11	62	RASESVSSNLA	70	GAFNRAT	78	QQRSDWFT
CD47	C47A8-CQ	63	SGTSSDVGGHNYVS	71	DVTKRPS	79	QSYAGSRVYV
CD20	UMAB10	64	RASSSVSYMD	72	ATSNLAS	80	QQWISNPPT

Table 7 lists exemplary amino acid sequences and nucleic acid sequences of the variable heavy chain of engineered IgA variant. The CDR regions are depicted in bold font

Antigen	Antigen binding domain	SEQ ID NO	Amino acid sequence	SEQ ID NO	Nucleic acid sequence
CD20	Obinutuzumab	81	QVQLVQSGAEVKKPGSS VKVSCKASGYAFSYSWI NWVRQAPGQGLEWMGR IFPGDGD TDYNGKFKGR VTITADKSTSTAYMELSSL RSEDTAVYYCARNVFDG YWL VYWGQGLTVTVSS	87	cagggtgcaattgggtcagctcggcgctg aagttaagaagcctgggagttcagtga ggctcctgcaaggctccggatacgcc ttcagctattcttggatcaattgggtgcg caggcgctggacaaggctcgagt gatgggacggatcttccggcgatggg gatactactacaatggaaattcaagg gcagagtcacaattaccggcgacaatc cactagcacagcctatatggagctgagc agcctgagatctgaggacacggcctg tattactgtcaagaaatgtcttggatggt actggctgttactggggcagggaac cctggtcaccgtctcgaca
GD2	Dinutuximab (also called Ch14.18 or Unituxin)	4	EVQLQSGPELEKPGASV MISCKASGSSFTGYNMN WVRQNIKGSLEWIGAI DP YYGGTSYNQKFKGR ATL TVDKSSSTAYMHLKSLTS EDSAVYYCVSGMEYWG QGTSTVTVSS	88	gaagtgcagctgggtgcagagcggcgc ggaagtggaaaaaccggcgcgagcgc tgaaaattagctgcaaacgcgagcggca gcagcttaccggctataacatgaactgg gtgcgccagaacattggcaaaagcctg gaatggattggcgcgattgatccgtatta tggcggcaccagctataaccagaaattt aaaggccgcgcgaccctgaccgtggat aaaagcaccagcaccgcgtatatgcatc tgaaaagcctgcgcagcgaagataccg cgggtgtattattgcgtgagcggcatgga atattggggccagggcaccagcgtgac cgtgagcagc
Her2	Trastuzumab	82	EVQLVESGGGLVQPGGSL RLSCAASG FN IKD TYI HW VRQAPGKGLEWVARI YP TNGYTRYADSVKGR FTIS ADTSKNTAYLQMNSLRA EDTAVYYCSR WG GD GF YAM DYWGQGLTVTVSS	89	Gaggtgcagctgggtcgagagcggcgg agggctgggtcagccaggcggcagcc tgaggctgtcctgcgccagcggctt caacatcaaggacacctacatccactgg gtcggcagggccaggaaggcct ggagtgggtggcaggtatctacccac caacggctacaccagatacgccgacag cgtgaagggcaggttcacatcagcgc cgacaccagaagaacaccgctacct gcagatgaacagcctgagggccgagg acaccggcgtgtactactgcagcagatg gggcggggacggcttctacgtatgga ctactggggccagggcaccctggtgac

					cgtgagcagc
Tyrosinase-related protein-1 or gp75	TA99	83	EVQLQQSGAELVRPGALV KLSCKTS GFNIKDYFLH WVRQRPDQGLEWIGWIN PDNGNTVYDPKFQGTAS LTADTSSNTVYLQLSGLT SEDTAVYFCTR RDYTYE KAALDYWGQGT TVTVST	90	gaggttcagttgcagcagctctggcgccg aactcgttagacctggcgctctggttaag ctgtcctgcaagacctccggttcaatat caaggactacttctgcactgggtccga cagaggcctgaccaaggactggaatgg atcggtggatcaaccccgacaacggc aacaccgtgtacgacctaaagtccagg gcaccgcttcttgaccgccgacacctc ttccaataccgtgtacctgcagctgtccg gcctgacctctgaggataccgccgtgta cttctgaccagacgggactacacctac gagaaggccgctctggattattggggcc agggcacaaccgtgaccgtgtctaca
cytotoxic T-lymphocyte-associated protein 4 (mCTLA-4)		84	QVKLEESGPGLVNPSQSL SLSCSVT GYSITSGYGWN WIRQFPGQKVEWMGFIY YEGSTYYNPSIKSRISITR DTSKNQFFLQVNSVTTED TATYYCAR QTGYFDYW GQGTMTVTVSS	91	caggtgaaactggaagaagcggccc gggcctggtaacccgagccagacct gagcctgagctgcagcgtgaccggcta tagcattaccagcggctatggctggaac tggattcgccagttccgggccagaaag tggaatggatgggctttatttattatgaag gcagcacctattataacccgagcattaaa agccgcattagcattaccgcgatacca gcaaaaaccagtttttctgcaggtgaac agcgtgaccaccgaagataccgcgacc tattattgcgcgcgaccagcggtatttt gattattggggccagggcaccatggtga ccgtgagcagcgggtgagtg
CD47	2.3D11	85	QVQLQESGPGLVKPSGTL SLTCAV SGVSIRSINWWN WVRQPPGKGLEWIGEIIY HSGSTNYNPSLKS RVTIS VDKSKNQFSLKLNSVTAA DTAVYYCARD GGLAVTD YYYYGLDVWGQGT TVTVSS	92	caggtgcagctgcaggaaagcggccc gggcctggtaaacccgagcggcacct gagcctgacctgcgcggtgagcggcgt gagcattcgcagcattaaactggtggaac tgggtgcgccagccgccggcgaagg cctggaatggattggcgaaattatcata gcggcagcaccactataacccgagcc tgaagccgcgtgaccattagcgtgga taaaagcaaaaccagtttagcctgaaa ctgaacagcgtgaccgcggcgatacc gcgggtgattattgcgcgcgcatggcg gcattgcggtgaccgattattattatg gcctggatgtgtgggccagggcacca ccgtgaccgtgagcagc
CD47	C47A8-CQ	86	QVQLVQSGAEVKKPGAS VKVSKAS GYTFTSYM	93	caggtgcagctggcagagcggcgc ggaagtgaiaaaacggcgcgagcgc

			HWVRQAPGQGLEWMGII NPSGGSTSYAQKFQGRV TMTRDTSTSTVYMELSSL RSEDTAVYYCARSTLWF SEFDYWGQGLTVTVSS		tgaagtgaagctgcaaacgagcgccgt atactttaccagctattatgcattgggt gcgccaggcgccggccaggccctgg aatggatgggcattattaacccgagcgg cggcagcaccagctatgcgcagaaatt cagggccgctgaccatgaccgcgat accagcaccagcaccgtgtatatggaac tgagcagcctgcgcagcgaagataccg cgggtgattattgcgcgcgagcaccct gtggtttagcgaatttgattattggggcca gggcaccctggtgaccgtgagcagc
CD20	UMAB10	7	QAYLQSGADLVRPGAS VKMSCKASGFTFPSYNL HWVKQTPRQGLEWIGAI YPGNGDTSYNQKFKGK ATLTVDKSSSTAYMQLSS LTSEDSAVYFCARSAYYG SNVWFFDVWGTGTTVTV SS	94	caagcctacctgcagcagctctggcgcc gatctctgctggcctgcccctctgtga agatgagctgtaaacccagcggcttcac cttccccagctacaacctgcactgggtc aagcagacccccagacagggcctgga atggatcggagccatctaccccgcaa cggcgacacctctacaaccagaagttc aagggaagggcacctgaccgtgga caagagcagcagcaccgcctacatgca gctgagcagcctgaccagcaggaca gcgccgtgtacttctgtccagaagcgc ctactacggcagcaacgtgtggttcttcg acgtgtggggcaccggcaccaccgtga cagtgcatct

Table 8 lists exemplary amino acid sequences and nucleic acid sequences of the variable light chain of engineered IgA variant. The CDR regions are depicted in bold font

Antigen	Antigen binding domain	SEQ ID NO	Amino acid sequence	SEQ ID NO	Nucleic acid sequence
CD20	Obinutuzumab	95	DIVMTQTPLSLPVT PGEPASISCRSSKS LLHSNGITYLYWY LQKPGQSPQLLIYQ MSNLVSGVPDRFS GSGSGTDFTLKISR VEAEDVGVYYCA QNLELPYTFGGGT KVEIK	101	gatatcgtgatgaccagacacctctgagcctgctg tgacacctggcgagcctgcttccatctctgcccgtc ctctaagtcctgctgcacttaacggcatcacctacc tgtactggatctgcagaagccccggcagctcctca gctgctgatctaccagatgtccaacctggtgtctggc gtgccccagagatttccggctctggctctggcaccg acttcacctgaagatctccagagtgaagccgagg acgtgggcgtgtactattgtgccagaacctggaact gccctacacctttggcggaggcacaaggtggaat caag
GD2	Dinutuximab (also called	5	EIVMTQSPATLSVS PGERATLSCRSSQS	102	gaaattgtgatgaccagagccccggcgaccctgag cgtgagccccggcgaaacgcgcgaccctgagctgc

	Ch14.18 or Unituxin)		LVHRNGNTYLHW YLQKPGQSPKLLIH KVSNRFSGV PDRF SGSGSGTDFTLKIS RVEAEDLG VYFCS QSTHVPPLTFGAG TKLELK		cgcagcagccagagcctggtgcatcgcaacggcaa cacctatctgcatgtgtatctgcagaaacgggccag agcccgaactgctgattcataaagtgcgaaccgct ttagcggcgtgcggatcgctttagcggcagcggca gcggcaccgattttaccctgaaaattagccgctgga agcgggaagatctggcgctgtattttgcagccagagc acccatgtgccgccgtgaccttggcgcgggcacc aaactggaactgaaa
Her2	Trastuzumab	96	DIQMTQSPSSLSAS VGDRVITICRASQ DVNTA VAWYQQK PGKAPKLLIYSASF LYSGVPSRFSGSRS GTDFTLTISSLQPE DFATYYCQQHYT TPPTFGQGTKVEIK	103	gacatccagatgacccagtcctcactctctgtccgc ctctgtggcgacagagtaccatcacctgtagagc cagccagcagtgtaacaccggcgtggcttggtatca gcagaagcctggcaaggccccctaaagctgctgatcta ctcgcctcctctgtactctggcgtgccctccagatt ctcggcagcagatctggcaccgactttaccctgaca atctccagcctgcagcctgaggacttcgccactact actgccagcagcactacaccacacctccaaccttgg ccagggcaccaaggtggaatacaag
Tyrosinas e-related protein-1 or gp75	TA99	97	IQMSQSPASLSASV GETVTITCRASGNI YNYLAWYQQKQG KSPHLLVYDAKTL ADGVPSRFSGSGSG TQYSLKISSLQTED SGNYCYQHFWSL PFTFGSGTKLEIK	104	atccagatgagtcagtcctccggcctccatctgcatc tgtgggagaaactgtcaccatcacatgctgagcaagt ggaaatatttacaattatttagatggtatcagcagaaa cagggaaaaatctcctcactcctggtctatgatgcaa aaaccttagcagatggtgtgccatcaaggttcagtg cagtggtcagggacacaaattctctcaagattagc agccttcagactgaagattctgggaattattactgtca acattttggagcttccattcacgttcggctcggggac caagctggaataaaa
cytotoxic T- lymphocy te- associate d protein 4 (mCTLA -4)		98	DVMMTQSPSSMSV SAGEKATISCKSSQ SLFNSNAKTNYLN WYMQKPGQSPKLL TYYASTRHTGVPD RFRGSGSGTDFTLT ISSVQDEDQAFYYC QQWYDYPYTFGA GTKLEIK	105	gatgtgatgatgacccagagcccgcagcagcatgag cgtgagcgcggcgaaaaagcgaccattagctgca aaagcagccagagcctgttaacagcaacgcgaaa accaactatctgaactggtatatgcagaaaccgggcc agagcccgaactgctgacctattatgcgagcacc gccataccggcgtgccggatcgcttcggcgagcg gcagcggcaccgattttaccctgaccattagcagcgt gcaggatgaagatcaggcgtttattattgccagcagt ggtatgattatccgtataccttggcgcgggcaccaa actggaaataaaa
CD47	2.3D11	99	EIVLTQSPATLSLSP GERATLSCRASES VSSNLAWYQQKPG QAPRLLIYGAFNR ATGIPARFSGSGSG TDFTLTISSLEPEDF	106	gaaattgtgctgacccagagcccggcgaccctgag cctgagccccggcgaaacgcgcgacctgagctgcc gcgcgagcgaagcgtgagcagcaacctggcgtg gtatcagcagaaaccggcgagcgccgcgcctgc tgatttatggcgcgtttaaccgcgcgaccggcattcc

			AVYYC QQRSDWF TFGGGTKVEIK		ggcgcgctttagcggcagcggcagcggcaccgatt ttaccctgaccattagcagcctggaaccggaagatttt gcgggtgtattattgccagcagcgcagcgattggttac ctttggcggcggcaccgaaagtggaaattaaa
CD47	C47A8-CQ	100	QSVLTQPSSVSASP GQSITIS CSGTSSDV GGHNY VSQYQQH PGKAPKLMY DVT KRPSG VPDRFSGS KSGNTASLTVSGL QAEDEADYY CQSY AGSRVY VFGTGTK LTVL	107	cagagcgtgctgaccagccgagcagcgtgagcg cgagcccgggccagagcattaccattagctgcagc ggcaccagcagcgtgtggcgggccataactatgtg agctggtatcagcagcattccgggcaagcgccgaa actgatgattatgatgtgaccaaagcccgagcggc gtgccggtatcgttttagcggcagcaaaagcggcaa caccgcgagcctgaccgtgagcggcctgcagcg gaagatgaagcggattattattgccagagctatgcgg gcagccgcgtgtatgtttggcaccggcaccgaaact gaccgtgctg
CD20	UMAB10	8	QIVLSQSPAILSASP GEKVTMT CRASS VSYMD WYQKPG SSPKWIY ATSNLA SGVPTRFSGSGSGT SYSLTISRVEAEDA ATYYC QQWISNPP TFGAGTKLDLK	108	cagatcgtgctgagccagagcccccgcacccctgagt gctagccctggcgagaaagtgacctgacctgcag agccagcagcagcgtgtcctacatggactggtatca gcagaagcccgagcagcccaagccctggtatct acgccacaagcaatctggccagcggcgtgcccaca agattttccggcagcggcctctggcaccagctacagc ctgacctagccgggtggagccgaagatgccgc cacctactactgccagcagtggtatcagcaaccccc cacctttggagccggcaccgaaagtggatctgaag

Table 9 lists exemplary IgA variant antibodies capable of binding different antigen. The table shows exemplary pairings of sequences of the variable light chain, variable heavy chain, IgA heavy chain constant region and kappa light chain constant region for the IgA variant antibodies.

Exemplary engineered IgA antibody	Antigen	HC-CD R1 of SEQ ID NO:	HC-CD R2 of SEQ ID NO:	HC-CD R3 of SEQ ID NO:	Variable heavy chain of SEQ ID NO:	LC-CD R1 of SEQ ID NO:	LC-CD R2 of SEQ ID NO:	LC-CD R3 of SEQ ID NO:	Variable light chain of SEQ ID NO:	IgA heavy chain constant region of SEQ ID NO:	Kappa light chain constant region of SEQ ID NO:
IgA3.0+-Obi	CD20	33	41	49	81	57	65	73	95	16	31
IgA3.0+-ch14.18	GD2	34	42	50	4	58	66	74	5	16	31

IgA3.0+- Her2	Her2	35	43	51	82	59	67	75	96	16	31
IgA3.0+- TA99	gp75	36	44	52	83	60	68	76	97	16	31
IgA3.0+- mCTLA4	CTLA 4	37	45	53	84	61	69	77	98	16	31
IgA.3.0+- 2.3D11	CD47	38	46	54	85	62	70	78	99	16	31
IgA3.0+- C47A8- CQ	CD47	39	47	55	86	63	71	79	100	16	31
IgA3.0+- UMAB10	CD20	40	48	56	7	64	72	80	8	16	31
IgA3.0mi n-Obi	CD20	33	41	49	81	57	65	73	95	17	31
IgA3.0mi n-ch14.18	GD2	34	42	50	4	58	66	74	5	17	31
IgA3.0mi n-Her2	Her2	35	43	51	82	59	67	75	96	17	31
IgA3.0mi n-TA99	gp75	36	44	52	83	60	68	76	97	17	31
IgA3.0mi n- mCTLA4	CTLA 4	37	45	53	84	61	69	77	98	17	31
IgA.3.0mi n-2.3D11	CD47	38	46	54	85	62	70	78	99	17	31
IgA3.0mi n-C47A8- CQ	CD47	39	47	55	86	63	71	79	100	17	31
IgA3.0mi n- UMAB10	CD20	40	48	56	7	64	72	80	8	17	31
IgA4.0_N G-Obi	CD20	33	41	49	81	57	65	73	95	18	31
IgA4.0_N T-Obi	CD20	33	41	49	81	57	65	73	95	20	31

IgA4.0_N Q-Obi	CD20	33	41	49	81	57	65	73	95	19	31
IgA4.0_N LT-TIS- Obi	CD20	33	41	49	81	57	65	73	95	21	31
IgA4.0_N G- ch14.18	GD2	34	42	50	4	58	66	74	5	18	31
IgA4.0_N T- ch14.18	GD2	34	42	50	4	58	66	74	5	20	31
IgA4.0_N Q- ch14.18	GD2	34	42	50	4	58	66	74	5	19	31
IgA4.0_N LT-TIS- ch14.18	GD2	34	42	50	4	58	66	74	5	21	31
IgA4.0_N G- Her2	Her2	35	43	51	82	59	67	75	96	18	31
IgA4.0_N T- Her2	Her2	35	43	51	82	59	67	75	96	20	31
IgA4.0_N Q- Her2	Her2	35	43	51	82	59	67	75	96	19	31
IgA4.0_N LT-TIS- Her2	Her2	35	43	51	82	59	67	75	96	21	31
IgA4.0_N G- TA99	gp75	36	44	52	83	60	68	76	97	18	31
IgA4.0_N T- TA99	gp75	36	44	52	83	60	68	76	97	20	31
IgA4.0_N Q- TA99	gp75	36	44	52	83	60	68	76	97	19	31
IgA4.0_N LT-TIS- TA99	gp75	36	44	52	83	60	68	76	97	21	31
IgA4.0_N G-	CTLA 4	37	45	53	84	61	69	77	98	18	31

mCTLA4											
IgA4.0_N T- mCTLA4	CTLA 4	37	45	53	84	61	69	77	98	20	31
IgA4.0_N Q- mCTLA4	CTLA 4	37	45	53	84	61	69	77	98	19	31
IgA4.0_N LT-TIS- mCTLA4	CTLA 4	37	45	53	84	61	69	77	98	21	31
IgA4.0_N G- 2.3D11	CD47	38	46	54	85	62	70	78	99	18	31
IgA4.0_N T- 2.3D11	CD47	38	46	54	85	62	70	78	99	20	31
IgA4.0_N Q- 2.3D11	CD47	38	46	54	85	62	70	78	99	19	31
IgA4.0_N LT-TIS- 2.3D11	CD47	38	46	54	85	62	70	78	99	21	31
IgA4.0_N G- C47A8- CQ	CD47	39	47	55	86	63	71	79	100	18	31
IgA4.0_N T- C47A8- CQ	CD47	39	47	55	86	63	71	79	100	20	31
IgA4.0_N Q- C47A8- CQ	CD47	39	47	55	86	63	71	79	100	19	31
IgA4.0_N LT-TIS- C47A8- CQ	CD47	39	47	55	86	63	71	79	100	21	31
IgA4.0_N G- UMAB10	CD20	40	48	56	7	64	72	80	8	18	31

IgA4.0_N T- UMAB10	CD20	40	48	56	7	64	72	80	8	20	31
IgA4.0_N Q- UMAB10	CD20	40	48	56	7	64	72	80	8	19	31
IgA4.0_N LT-TIS- UMAB10	CD20	40	48	56	7	64	72	80	8	21	31

Table 10 lists exemplary sequence of CH1, CH2 and CH3 domains of WT IgA heavy chain constant region

Name	SEQ ID NO:	Amino acid sequence
CH1 domain of WT IgA2 heavy chain constant region	109	ASPTSPKVFPLSLDSTPQDGNVVVACLV QGFFPQEPLSVTWSESGQNVTARNPPPSQ DASGDLYTTSSQLTLPATQCPDGKSVTC HVKHYTNPSQDVTVP
CH2 domain of WT IgA2 heavy chain constant region	110	CCHPRLSLHRPALEDLLLGSEANLTCTL TGLRDASGATFTWTPSSGKSAVQGPPER DLCGCYSVSSVLPGCAQPWNHGETFTC TAAHPELKTPLTANITKS
CH3 domain of WT IgA2 heavy chain constant region including C-terminal Tail-piece	111	GNTFRPEVHLLPPPSEELALNELVTLTCL ARGFSPKDVLRWLQGSQELPREKYLT WASRQEPSQGTTFVAVTSILRVAAEDW KKGDTFSCMVGHEALPLAFTQKTIDRL AGKPTHVNVSVVMAEVDGTCY

Table 11

IMGT numbering table
IMGT accession number J00221
Weblink: <http://www.imm.org/IMGTconserve/TCRseq/aj:aleg/insg.php?accession=Hmch33&accessions&group=TCR&gene=IGHA2>
Consulted on 27MAR2020

J00221	IGHA2*01 F CH1 gDNA	1.8 1.7 1.6 1.5 1.4 1.3 1.2 1.1 1 2 3 4 5 6 7 8 9 10 11 12	A S P T S P K V F P L S L D S T	gca tcc ccg acc agc ccc aag gtc ttc ccg ctg agc ctc gac agc acc
J00221	IGHA2*01 F CH1 gDNA	13 14 15 15.1 15.2 15.3 16 17 18 19 20 21 22 23 24 25 26 27 28 29	P Q gat ggg aac gtc gtc gca tgc ctg gtc cag ggc ttc ttc	ccc caa
J00221	IGHA2*01 F CH1 gDNA	30 31 34 35 36 37 38 39 40 41 42 43 44 45 45.1 45.3 45.5	P Q E P L S V T W S E S G Q N V	ccc cag ... gag cca ctc agt gtc acc tgg agc gaa agc gga cag aac gtc
J00221	IGHA2*01 F CH1 gDNA	45.7 77 78 79 80 81 82 83 84 84.1 84.3 84.5 84.7 85.6 85.4	T A R N F P P S Q D A S G D	... acc gcc aga aac ttc cca cct agc cag gat gcc tcc ggg gac
J00221	IGHA2*01 F CH1 gDNA	85.3 85.2 85.1 85 86 87 88 89 90 91 92 93 94 95 96 96.1 96.2 97 98 99	L Y T T S S Q L T L P A T Q C P D G	ctg tac acc acg agc agc cag ctg acc ctg ccg gcc aca cag tgc cca gac ggc
J00221	IGHA2*01 F CH1 gDNA	100 101 102 103 104 105 106 107 108 109 110 111 112 113 114 115 116 117 118 119	K S V T C H V K H Y T N P S Q D V	aag tcc gtc aca tgc cac gtc aag cac tac acg aat ccc agc cag gat gtc
J00221	IGHA2*01 F CH1 gDNA	120 121 122 123 124	T V P C E	act gtc ccc tgc cca

181

J00221	IGHA2*01 F H gDNA	1 2 3 4 5 6 7 8 9 10 11 12 V P P P P P gtt ccc cca cct ccc cca
J00221	IGHA2*01 F CH2 gDNA	1.8 1.7 1.6 1.5 1.4 1.3 1.2 1.1 1 2 3 4 5 6 7 8 9 10 11 12 C C H P R L S L H R P A L E D L L L tgc tgc cac ccc cga ctg tgc ctg cac cga ccg gcc ctc gag gac ctg ctc tta
J00221	IGHA2*01 F CH2 gDNA	AB 13 14 15 15.1 15.2 15.3 16 17 18 19 20 21 22 23 24 25 26 27 28 29 G S E A N L T C T L T G L R D A S G A ggt tca gaa gcg aac ctc acg tgc aca ctg acc ggc ctg aga gat gcc ... tct ggt gcc
J00221	IGHA2*01 F CH2 gDNA	BC CD 30 31 34 35 36 37 38 39 40 41 42 43 44 45 45.1 45.3 45.5 T F T W T P S S G K S A V Q G acc ttc acc tgg acg ccc tca agt ggg aag agc gct gtt caa gga
J00221	IGHA2*01 F CH2 gDNA	DE 45.7 77 78 79 80 81 82 83 84 84.1 84.3 84.5 84.7 85.6 85.4 P P E R D L C G C Y S V S S cca cct gag cgt gac ctc tgt ggc tgc tac agc gtg tcc agt
J00221	IGHA2*01 F CH2 gDNA	EF 85.2 85.3 85.1 85 86 87 88 89 90 91 92 93 94 95 96 96.1 96.2 97 98 99 V L P G C A Q P W N H G E T F T C T gtc ctg cct gcc tgt gcc cag cca tgg aac cat ggg gag acc ttc acc tgc act
J00221	IGHA2*01 F CH2 gDNA	FG 100 101 102 103 104 105 106 107 108 109 110 111 112 113 114 115 116 117 118 A A H P E L K T P L T A N L T K S gct gcc cac ccc gag ttg aag acc cca cta acc gcc aac atc aca aaa tcc
J00221	IGHA2*01 F CH3-CHS gDNA	1.8 1.7 1.6 1.5 1.4 1.3 1.2 1.1 1 2 3 4 5 6 7 8 9 10 11 12 G N T F R P E V H L L P P P S E gga aac aca ttc cgg ccc gag gtc cac ctg ctg ccg ccg ctg tgc gag
J00221	IGHA2*01 F CH3-CHS gDNA	AB 13 14 15 15.1 15.2 15.3 16 17 18 19 20 21 22 23 24 25 26 27 28 29 E L A L N E L V T L T C L A R G F S gag ctg gcc ctg aac gag ctg gtg acg ctg acg tgc ctg gca cgt gcc ttc agc
J00221	IGHA2*01 F CH3-CHS gDNA	BC CD 30 31 34 35 36 37 38 39 40 41 42 43 44 45 45.1 45.3 45.5 P K D V L V R W L Q G S Q E L P R E ccc aag gat gtg ctg gtt cgc tgg ctg cag ggg tca cag gag ctg ccc cgc gag
J00221	IGHA2*01 F CH3-CHS gDNA	DE 45.7 77 78 79 80 81 82 83 84 84.1 84.3 84.5 84.7 85.6 85.4 K Y L T W A S R Q E P S Q G 85.7 85.5 ... aag tac ctg act tgg gca tcc cgg cag gag ccc agc cag ggc acc acc
J00221	IGHA2*01 F CH3-CHS gDNA	EF 85.2 85.3 85.1 85 86 87 88 89 90 91 92 93 94 95 96 96.1 96.2 97 98 99 T F A V T S I L R V A A E D W K K G acc ttc gct gtg acc agc ata ctg cgc gtg gca gcc gag gac tgg aag aag ggg
J00221	IGHA2*01 F CH3-CHS gDNA	FG 100 101 102 103 104 105 106 107 108 109 110 111 112 113 114 115 116 117 118 119 D T F S C M V G H E A L P L A F T Q K gac acc ttc tcc tgc atg gtg ggc cac gag gcc ... ctg ccg ctg gcc ttc aca cag aag
J00221	IGHA2*01 F CH3-CHS gDNA	CHS 120 121 122 123 124 125 126 127 128 129 130 131 132 133 134 135 136 137 138 139 T I D R L A G K P T H V N V S V V acc atc gac cgc ttg gcg ggt aaa ccc acc cat gtc aat gtg tct gtt gtc
J00221	IGHA2*01 F CH3-CHS gDNA	140 141 142 143 144 145 146 147 148 M A E V D G T C Y atg gcg gag gtg gac ggc acc tgc tac

[0653] While preferred embodiments of the present disclosure have been shown and described herein, it will be obvious to those skilled in the art that such embodiments are provided by way of example only. Numerous variations, changes, and substitutions will now occur to those skilled in the art without departing from the present disclosure. It should be understood that various alternatives to the embodiments described
5 herein, or combinations of one or more of these embodiments or aspects described therein may be employed in practicing the present disclosure. It is intended that the following claims define the scope of the present disclosure and that methods and structures within the scope of these claims and their equivalents be covered thereby.

CLAIMS

What is claimed is:

1. An antibody or a functional fragment thereof that comprises:
 - a. an antigen binding domain; and
 - 5 b. a constant domain, wherein the constant domain comprises an immunoglobulin A (IgA) heavy chain constant region,
wherein the IgA heavy chain constant region comprises an IgA CH2 region and an IgA CH3 region;
wherein the IgA heavy chain constant region comprises a modification of at least two naturally occurring glycosylation sites, as compared to a corresponding WT IgA heavy chain constant region, and
- 10 wherein each of the at least two naturally occurring glycosylation sites is in the IgA CH2 region or in the IgA CH3 region.
2. The antibody or the functional fragment thereof of claim 1, wherein the at least two naturally occurring glycosylation sites are two naturally occurring N-linked glycosylation sites.
3. The antibody or the functional fragment thereof of claim 1 or 2, wherein the at least two naturally occurring
- 15 glycosylation sites comprises a naturally occurring asparagine (N) amino acid residue as compared to a corresponding wild type IgA.
4. The antibody or the functional fragment thereof of any one of claims 1-3, wherein the modification comprises an amino acid substitution, or an amino acid deletion of one, or both of the at least two naturally occurring glycosylation sites.
- 20 5. The antibody or the functional fragment thereof of claim 4, wherein the amino acid substitution is a non-conservative amino acid substitution.
6. The antibody or the functional fragment thereof of any one of claims 1-5, wherein the IgA heavy chain constant region comprises an amino acid substitution at:
 - i. N114 and N135,
 - 25 ii. N114 and N15.2, or
 - iii. N135 and N15.2,relative to the WT IgA heavy chain constant region comprising an amino acid sequence of SEQ ID NO: 1, numbering according to IMGT scheme.
7. The antibody or the functional fragment thereof of any one of claims 1-6, wherein the IgA heavy chain
- 30 constant region comprises:
 - i. a N114T amino acid substitution and a N135Q amino acid substitution,
 - ii. a N114T amino acid substitution and an amino acid substitution selected from the group consisting of N15.2G, N15.2Q, and N15.2T, or

- iii. a N135Q amino acid substitution and an amino acid substitution from the group consisting of N15.2G, N15.2Q, and N15.2T, relative to the WT IgA heavy chain constant region comprising an amino acid sequence of SEQ ID NO: 1, numbering according to IMGT scheme.
- 5 8. The antibody or the functional fragment thereof of any one of claims 1-8, wherein the IgA heavy chain constant region comprises a modification of at least three naturally occurring glycosylation **sites**, as compared to a corresponding wild type IgA.
9. The antibody or the functional fragment thereof of claim 8, wherein the at least three naturally occurring glycosylation sites are three naturally occurring N-linked glycosylation sites.
- 10 10. The antibody or the functional fragment thereof of claim 8 or 9, wherein the at least three naturally occurring glycosylation sites comprises a naturally occurring asparagine (N) amino acid residue as compared to a corresponding wild type IgA.
11. The antibody or the functional fragment thereof of any one of claims 8-10, wherein the modification is an amino acid substitution or an amino acid deletion.
- 15 12. The antibody or the functional fragment thereof of claim 11, wherein the amino acid substitution is a non-conservative substitution.
13. The antibody or the functional fragment thereof of any one of claims 8-12, wherein the IgA heavy chain constant region comprises an amino acid substitution at :
- 20 N114, N135, and N15.2, relative to the WT IgA heavy chain constant region comprising an amino acid sequence of SEQ ID NO: 1, numbering according to IMGT scheme.
14. The antibody or the functional fragment thereof of any one of claims 8-13, wherein the IgA heavy chain constant region comprises:
- i. a N114T amino acid substitution,
- ii. a N135Q amino acid substitution, and
- 25 iii. an amino acid substitution selected from the group consisting of N15.2G, N15.2Q, and N15.2T, relative to the WT IgA heavy chain constant region comprising an amino acid sequence of SEQ ID NO: 1, numbering according to the IMGT scheme.
15. The antibody or the functional fragment thereof of any one of claims 1-14, wherein the IgA heavy chain constant region further comprises an IgA CH1 region.
- 30 16. The antibody or the functional fragment thereof of claim 15, wherein the heavy chain constant region comprises a modification of at least one naturally occurring N-linked glycosylation site in the IgA CH2 region, at least one naturally occurring glycosylation site in the IgA CH3 regions, and at least one naturally occurring glycosylation site within the IgA CH1 region, as compared to a corresponding wild type IgA.

17. The antibody or the functional fragment thereof of claim 15 or 16, wherein the IgA heavy chain constant region comprises an amino acid substitution at:
- i. N45.2, N114, and N135, or
 - ii. N45.2, N15.2, and N135,
- 5 relative to the WT IgA heavy chain constant region comprising an amino acid sequence of SEQ ID NO: 1, numbering according to IMGT scheme.
18. The antibody or the functional fragment thereof of claim 17, wherein the IgA heavy chain constant region comprises:
- i. a N45.2G amino acid substitution, a N114T amino acid substitution, and a N135Q amino acid substitution;
- 10 or
- ii. the N45.2G amino acid substitution, the N135Q amino acid substitution, and an amino acid substitution selected from a group consisting of N15.2G, N15.2Q, and N15.2T,
- relative to the WT IgA heavy chain constant region comprising an amino acid sequence of SEQ ID NO: 1, numbering according to IMGT scheme.
- 15 19. The antibody or the functional fragment thereof of claim 17, wherein the heavy chain constant region comprises a modification of at least two naturally occurring N-linked glycosylation site in the IgA CH2 region, and at least one naturally occurring glycosylation site within the IgA CH1 region, as compared to a corresponding wild type IgA.
20. The antibody or the functional fragment thereof of claim 19, wherein the IgA heavy chain constant region
- 20 comprises an amino acid substitution at:
- i. N45.2, N114, and N15.2G, numbering according to IMGT scheme.
21. The antibody or the functional fragment thereof of claim 20, wherein the IgA heavy chain constant region comprises:
- i. a N45.2G amino acid substitution,
- 25 ii. a N114T amino acid substitution, and
- iii. an amino acid substitution selected from the group consisting of N15.2G, N15.2Q, and N15.2T,
- relative to the WT IgA heavy chain constant region comprising an amino acid sequence of SEQ ID NO: 1, numbering according to IMGT scheme
22. The antibody or the functional fragment thereof of any one of claims 1-21, wherein the antibody or the
- 30 functional fragment thereof exhibits a greater circulating half-life compared to a corresponding WT IgA antibody.
23. The antibody or the functional fragment thereof of any one of claims 1-22, wherein the antibody or the functional fragment thereof exhibits decreased aggregation compared to a corresponding WT IgA antibody.

24. The antibody or the functional fragment thereof of any one of claims 1-23, wherein the antibody or the functional fragment thereof exhibits decreased aggregation with serum proteins, compared to a corresponding WT IgA antibody.
25. The antibody or the functional fragment thereof of any one of claims 1-24, wherein the antibody or the functional fragment thereof induces increased antibody dependent cell mediated cytotoxicity, compared to a corresponding antibody comprising an IgG heavy chain constant region.
26. The antibody or the functional fragment thereof of any one of claims 1-25, wherein the antibody or the functional fragment thereof exhibits increased thermostability, compared to a corresponding WT IgA antibody.
27. The antibody or the functional fragment thereof of any one of claims 1-26, wherein the antibody or the functional fragment thereof exhibits decreased glycosylation, compared to a corresponding WT IgA antibody.
28. The antibody or the functional fragment thereof of any one of claims 1-27, wherein the IgA heavy chain constant region exhibits binding to a Fc α R expressed on an immune effector cell with increased affinity, compared to a corresponding WT IgA antibody.
29. The antibody or the functional fragment thereof of any one of claims 1-28, wherein the IgA heavy chain constant region comprises a modification of at least four naturally occurring glycosylation sites, as compared to a corresponding wild type IgA.
30. The antibody or the functional fragment thereof of claim 29, wherein the at least four naturally occurring glycosylation sites are four naturally occurring N-linked glycosylation sites.
31. The antibody or the functional fragment thereof of claim 29 or claim 30, wherein the at least four naturally occurring glycosylation sites comprises a naturally occurring asparagine (N) amino acid residue.
32. The antibody or the functional fragment thereof of any one of claims 29-31, wherein the heavy chain constant region comprises a modification of at least two naturally occurring N-linked glycosylation sites within the IgA CH2 region, at least one naturally occurring N-linked glycosylation sites within the IgA CH3 region, and at least one naturally occurring N-linked glycosylation site within the IgA CH1 region, as compared to a corresponding wild type IgA.
33. The antibody or the functional fragment thereof of any one of claims 29-32, wherein the IgA heavy chain constant region comprises an amino acid substitution at amino acid residues: N45.2, N114, N135, and N15.2, relative to the WT IgA heavy chain constant region comprising an amino acid sequence of SEQ ID NO: 1, numbering according to IMGT scheme.
34. The antibody or the functional fragment thereof of any one of claims 29-33, wherein the IgA heavy chain constant region comprises a non-conservative amino acid substitution at amino acid residues: N45.2, N114, N135, and N15.2, numbering according to IMGT scheme.

35. The antibody or the functional fragment thereof of any one of claims 29-34, wherein the IgA heavy chain constant region comprises:
- i. a N45.2G amino acid substitution,
 - ii. a N114T amino acid substitution,
 - 5 iii. a N135Q amino acid substitution, and
 - iv. an amino acid substitution selected from the group consisting of N15.2G, N15.2Q, and N15.2T, relative to the WT IgA heavy chain constant region comprising an amino acid sequence of SEQ ID NO: 1, numbering according to IMGT scheme.
- 10 36. The antibody or the functional fragment thereof of any one of claims 29-35, wherein the antibody or the functional fragment thereof exhibits a greater circulating half-life compared to a corresponding IgA antibody comprising at least one naturally occurring N-linked glycosylation site.
37. The antibody or the functional fragment thereof of any one of claims 29-36, wherein the antibody or the functional fragment thereof induces increased antibody dependent cell mediated cytotoxicity (ADCC), compared to a corresponding antibody comprising an IgG CH2 domain and an IgG CH3 domain.
- 15 38. The antibody or the functional fragment thereof of any one of claims 29-37, wherein the antibody or the functional fragment thereof exhibits increased thermostability, compared to a corresponding IgA antibody comprising at least one naturally occurring N-linked glycosylation site.
39. The antibody or the functional fragment thereof of any one of claims 29-38, wherein the antibody or the functional fragment thereof exhibits decreased glycosylation, compared to a corresponding IgA antibody
- 20 comprising at least one naturally occurring N-linked glycosylation site.
40. The antibody or the functional fragment thereof of any one of claims 29-39, wherein the IgA heavy chain constant region exhibits binding to a Fc α R expressed on an immune effector cell with increased affinity, compared to a corresponding IgA antibody comprising at least one naturally occurring N-linked glycosylation site.
- 25 41. The antibody or the functional fragment thereof of any one of claims 1-40, wherein the IgA heavy chain constant region comprises a modification of at least one naturally occurring cysteine (C) amino acid residue, as compared to a corresponding wild type IgA.
42. The antibody or the functional fragment thereof of claim 41, wherein the modification is an amino acid substitution or an amino acid deletion of the at least one naturally occurring cysteine (C) amino acid residue.
- 30 43. The antibody or the functional fragment thereof of claim 42, wherein the amino acid substitution is a non-conservative amino acid substitution of the at least one naturally occurring cysteine (C) amino acid residue.
44. The antibody or the functional fragment thereof of any one of claims 41-43, wherein the at least one naturally occurring cysteine amino acid residue is C147 or C86, relative to the WT IgA heavy chain constant region comprising an amino acid sequence of SEQ ID NO: 1, numbering according to IMGT scheme.

45. The antibody or the functional fragment thereof of claim 44, wherein the IgA heavy chain constant region comprises a C86S amino acid substitution, relative to the WT IgA heavy chain constant region comprising an amino acid sequence of SEQ ID NO: 1, numbering according to IMGT scheme.
46. The antibody or the functional fragment thereof of any one of claims 44-45, wherein the IgA heavy chain
5 constant region comprises a deletion of C147, relative to the WT IgA heavy chain constant region comprising an amino acid sequence of SEQ ID NO: 1, numbering according to the IMGT scheme.
47. The antibody or the functional fragment thereof of any one of claims 44-46, wherein the antibody or the functional fragment thereof exhibits decreased aggregation compared to a corresponding WT IgA antibody or the functional fragment thereof.
- 10 48. The antibody or the functional fragment thereof of any one of claims 44-47, wherein the antibody or the functional fragment thereof exhibits decreased aggregation with serum proteins compared to a corresponding WT IgA construct.
49. The antibody or the functional fragment thereof of any one of claims 44-48, wherein the IgA heavy chain constant region comprises a modification of at least two naturally occurring cysteine (C) amino acid residues,
15 as compared to a corresponding wild type IgA.
50. The antibody or the functional fragment thereof of claim 49, wherein the modification comprises an amino acid substitution of one, or both of the at least two naturally occurring cysteine (C) amino acid residues.
51. The antibody or the functional fragment thereof of claim 49, wherein the modification comprises a
20 deletion of one, or both of the at least two naturally occurring cysteine (C) amino acid residues.
52. The antibody or the functional fragment thereof of claim 49, wherein the IgA heavy chain constant region comprises an amino acid substitution one of the at least two naturally occurring cysteine (C) amino acid residues, and a deletion of one of the at least two naturally occurring cysteine (C) amino acid residues.
53. The antibody or the functional fragment thereof of any one of claims 49-52, wherein the at least two
25 naturally occurring cysteine amino acid residue are C147 and C86, relative to the WT IgA heavy chain constant region comprising an amino acid sequence of SEQ ID NO: 1, numbering according to IMGT scheme.
54. The antibody or the functional fragment thereof of claim 53, wherein the IgA heavy chain constant region comprises a deletion of C147, relative to the WT IgA heavy chain constant region comprising an amino acid
30 sequence of SEQ ID NO: 1, numbering according to IMGT scheme.
55. The antibody or the functional fragment thereof of claim 53, wherein the IgA heavy chain constant region comprises an amino acid substitution of C86, relative to the WT IgA heavy chain constant region comprising an amino acid sequence of SEQ ID NOs: 1, numbering according to IMGT scheme.

56. The antibody or the functional fragment thereof of claim 53, wherein the IgA heavy chain constant region comprises an amino acid substitution of the naturally occurring C86 amino acid residue, and a deletion of the naturally occurring C147 amino acid residue, relative to the WT IgA heavy chain constant region comprising an amino acid sequence of SEQ ID NO: 1, numbering according to IMGT scheme.
- 5 57. The antibody or the functional fragment thereof of claim 56, wherein the IgA heavy chain constant region comprises an C86S amino acid substitution, relative to the WT IgA heavy chain constant region comprising an amino acid sequence of SEQ ID NO: 1, numbering according to IMGT scheme.
58. The antibody or the functional fragment thereof of any one of claims 49-57, wherein the antibody or the functional fragment thereof exhibits decreased aggregation compared to a corresponding WT IgA antibody.
- 10 59. The antibody or the functional fragment thereof of any one of claims 49-58, wherein the antibody or the functional fragment thereof exhibits decreased aggregation with serum proteins compared to a corresponding WT IgA antibody.
60. The antibody or the functional fragment thereof of any one of claims 1-59, wherein the IgA heavy chain constant region comprises a modification of at least one naturally occurring tyrosine (Y) amino acid residue,
- 15 as compared to a corresponding wild type IgA.
61. The antibody or the functional fragment thereof of claim 60, wherein the modification is an amino acid substitution or a deletion of the at least one naturally occurring tyrosine (Y) amino acid residue.
62. The antibody or the functional fragment thereof of claim 60, wherein the amino acid substitution is a non-conservative amino acid mutation of the at least one naturally occurring tyrosine (Y) amino acid residue, as
- 20 compared to a WT IgA antibody.
63. The antibody or the functional fragment thereof of any one of claims 60-62, wherein the at least one tyrosine residue is Y148, relative to the WT IgA heavy chain constant region comprising an amino acid sequence of SEQ ID NO: 1, numbering according to IMGT scheme.
64. The antibody or the functional fragment thereof of claim 63, wherein amino acid Y148 is deleted, relative
- 25 to the WT IgA heavy chain constant region comprising an amino acid sequence of SEQ ID NO: 1, numbering according to IMGT scheme.
65. The antibody or the functional fragment thereof of any one of claims 60-64, wherein the antibody or the functional fragment thereof exhibits decreased aggregation compared to a corresponding WT IgA antibody.
66. The antibody or the functional fragment thereof of any one of claims 60-65, wherein the antibody or the
- 30 functional fragment thereof exhibits decreased aggregation with serum proteins compared to a corresponding WT IgA antibody.
67. The antibody or the functional fragment thereof of any one of claims 1-66, wherein the IgA heavy chain constant region comprises a modification of at least one naturally occurring threonine (T) amino acid residue, as compared to a corresponding wild type IgA antibody.

68. The antibody or the functional fragment thereof of claim 67, wherein the modification comprises an amino acid substitution or deletion of the at least one naturally occurring threonine (T) amino acid residue, as compared to a corresponding wild type IgA antibody.
69. The antibody or the functional fragment thereof of claim 68, wherein the amino acid substitution is a non-conservative amino acid substitution of the at least one naturally occurring threonine (T) amino acid residue.
70. The antibody or the functional fragment thereof of any one of claims 67-69, wherein the at least one naturally occurring threonine amino acid residue is T116 or T16, relative to the WT IgA heavy chain constant region comprising an amino acid sequence of SEQ ID NO: 1, numbering according to IMGT scheme.
71. The antibody or the functional fragment thereof of any one of claims 67-70, wherein the antibody or the functional fragment thereof exhibits decreased aggregation compared to a corresponding WT IgA antibody.
72. The antibody or the functional fragment thereof of any one of claims 67-71, wherein the antibody or the functional fragment thereof exhibits decreased aggregation with serum proteins compared to a corresponding WT IgA antibody.
73. The antibody or the functional fragment thereof of any one of claims 1-72, wherein the IgA heavy chain constant region comprises a modification of at least two naturally occurring threonine (T) amino acid residues, as compared to a corresponding wild type IgA.
74. The antibody or the functional fragment thereof of claim 73, wherein the modification comprises an amino acid substitution or a deletion of one, or both the at least two naturally occurring threonine (T) amino acid residues.
75. The antibody or the functional fragment thereof of claim 74, wherein the amino acid substitution is a non-conservative amino acid substitution of one, or both of the at least two naturally occurring threonine (T) amino acid residues.
76. The antibody or the functional fragment thereof of any one of claims 73-75, wherein the at least two naturally occurring threonine (T) amino acid residue are T116 or T16, relative to the WT IgA heavy chain constant region comprising an amino acid sequence of SEQ ID NO: 1, numbering according to IMGT scheme.
77. The antibody or the functional fragment thereof of claim 76, wherein the IgA heavy chain constant region comprises a T116S amino acid substitution, relative to the WT IgA heavy chain constant region comprising an amino acid sequence of SEQ ID NO: 1, numbering according to IMGT scheme.
78. The antibody or the functional fragment thereof of claim 76, wherein the IgA heavy chain constant region comprises a T16S amino acid substitution, relative to the WT IgA heavy chain constant region comprising an amino acid sequence of SEQ ID NO: 1, numbering according to IMGT scheme.

79. The antibody or the functional fragment thereof of any one of claims 1-78, wherein the IgA heavy chain constant region comprises a modification of at least one naturally occurring isoleucine (I) amino acid residue, as compared to a corresponding wild type IgA.
80. The antibody or the functional fragment thereof of claim 79, wherein the modification comprises an amino acid substitution or a deletion of the at least one naturally occurring isoleucine (I) amino acid residue.
81. The antibody or the functional fragment thereof of claim 80, wherein the amino acid substitution comprises a non-conservative amino acid substitution of the at least one naturally occurring isoleucine (I) amino acid residue.
82. The antibody or the functional fragment thereof of any one of claims 79-81, wherein the at least one naturally occurring isoleucine (I) residue is I115, relative to the WT IgA heavy chain constant region comprising an amino acid sequence of SEQ ID NO: 1, numbering according to IMGT scheme.
83. The antibody or the functional fragment thereof of claim 82, wherein the IgA heavy chain constant region comprises a I115L amino acid substitution, relative to the WT IgA heavy chain constant region comprising an amino acid sequence of SEQ ID NO: 1, numbering according to IMGT scheme.
84. The antibody or the functional fragment thereof of any one of claims 1-83, wherein the IgA heavy chain constant region comprises a modification of at least one naturally occurring leucine (L) amino acid residue, as compared to a corresponding wild type IgA.
85. The antibody or the functional fragment thereof of claim 84, wherein the modification comprises an amino acid substitution or a deletion of the at least one naturally occurring leucine (L) amino acid residue.
86. The antibody or the functional fragment thereof of claim 85, wherein the amino acid substitution is a non-conservative amino acid substitution of the at least one naturally occurring leucine (L) amino acid residue.
87. The antibody or the functional fragment thereof of any one of claims 84-86, wherein the at least one naturally occurring leucine (L) residue is L15.3, relative to the WT IgA heavy chain constant region comprising an amino acid sequence of SEQ ID NO: 1, numbering according to IMGT scheme.
88. The antibody or the functional fragment thereof of claim 87, wherein the IgA heavy chain constant region comprises a L15.3I amino acid substitution, relative to a WT IgA heavy chain constant region comprising an amino acid sequence of SEQ ID NO: 1-3, numbering according to IMGT scheme.
89. The antibody or the functional fragment thereof of any one of claims 1-88, wherein the IgA heavy chain constant region comprises a modification of at least one naturally occurring proline (P) amino acid residue, as compared to a corresponding wild type IgA.
90. The antibody or the functional fragment thereof of claim 89, wherein the IgA heavy chain constant region comprises an amino acid substitution or a deletion of the at least one naturally occurring proline (P) amino acid residue.

91. The antibody or the functional fragment thereof of claim 89, wherein the amino acid substitution is a non-conservative amino acid substitution of the at least one naturally occurring proline (P) amino acid residue.
92. The antibody or the functional fragment thereof of any one of claims 89-91, wherein the at least one naturally occurring proline (P) residue is P124, relative to the WT IgA heavy chain constant region
- 5 comprising an amino acid sequence of SEQ ID NO: 1, numbering according to IMGT scheme.
93. The antibody or the functional fragment thereof of claim 92, wherein the IgA heavy chain constant region comprises a P124R amino acid substitution, relative to the WT IgA heavy chain constant region comprising an amino acid sequence of SEQ ID NO: 1, numbering according to IMGT scheme.
94. The antibody or the functional fragment thereof of any one of claims 89-93, wherein the antibody or the
- 10 functional fragment thereof exhibits greater stability, compared to a corresponding WT IgA antibody.
95. The antibody or the functional fragment thereof of any one of claims 89-94, wherein the antibody or the functional fragment thereof exhibits greater stability between a heavy chain and a light chain, compared to a corresponding WT IgA antibody.
96. The antibody or the functional fragment thereof of any one of claims 89-95, wherein the antibody or the
- 15 functional fragment thereof has a covalent linkage between the heavy chain and the light chain.
97. The antibody or the functional fragment thereof of any one of claims 89-96, wherein the antibody or the functional fragment thereof comprises a disulfide bond between a cysteine (C) amino acid residue of the heavy chain and a cysteine (C) amino acid residue of the light chain.
98. The antibody or the functional fragment thereof of any one of claims 1-97, wherein the IgA heavy chain
- 20 constant region comprises an amino acid sequence selected from any one of SEQ ID NOs: 16-21.
99. The antibody or the functional fragment thereof of any one of claims 1-98, wherein the antibody is an allotype IgA2m(1) antibody or IgA2m(2).
100. The antibody or the functional fragment thereof of any one of claims 1-99, wherein the antibody is an allotype Caucasian IgA2m(1) antibody.
- 25 101. The antibody or the functional fragment thereof of any one of claims 1-100, wherein the IgA heavy chain constant region comprises a modification of a C-terminal IgA tail piece, as compared to a corresponding wild type IgA.
102. The antibody or the functional fragment thereof of any one of claims 1-101, wherein the modification comprises a deletion of the C-terminal 25, 24, 23, 22, 21, 20, 19, 18, 17, 16, 15, 12, 10, 8, 6, 4, or 2 amino
- 30 acids.
103. The antibody or the functional fragment thereof of any one of claims 1-102, wherein the modification comprises a deletion of the C-terminal 18 amino acids.

104. The antibody or the functional fragment thereof of any one of claims 1-103, wherein the modification comprises a deletion of C-terminal amino acid residues 131-148, relative to the WT IgA heavy chain constant region comprising an amino acid sequence of SEQ ID NO: 1, numbering according to IMGT scheme.
105. The antibody or the functional fragment thereof of any one of claims 1-104, wherein the IgA heavy chain constant region comprises a deletion of the C-terminal amino acid residues P131-Y148 relative to the WT IgA heavy chain constant region comprising an amino acid sequence of SEQ ID NO: 1, numbering according to IMGT scheme.
106. The antibody or the functional fragment thereof of any one of claims 1-105, wherein the IgA heavy chain constant region comprises an IgA1 constant region comprising an IgA1 CH2 region and an IgA1 CH3 region.
107. The antibody or the functional fragment thereof of claim 106, wherein the IgA1 constant region further comprises an IgA1 CH1 region.
108. The antibody or the functional fragment thereof of any one of claims 1-106, wherein the IgA heavy chain constant region comprises an IgA2 constant region comprising an IgA2 CH2 region and an IgA2 CH3 region.
109. The antibody or the functional fragment thereof of claim 108, wherein the IgA2 constant region further comprises an IgA2 CH1 region.
110. The antibody or the functional fragment thereof of any one of claims 1-109, wherein the antibody or the functional fragment thereof exhibits a circulating half-life within 1%, 5%, 10%, 20%, or 30% of the circulating half-life of a corresponding antibody comprising the antigen binding domain and an IgA heavy chain constant region comprising an IgG CH2 region and an IgG CH3 region.
111. An antibody or a functional fragment thereof comprising:
- a. an antigen binding domain; and
 - b. a constant domain, wherein the constant domain comprises an IgA2 heavy chain constant region comprising an IgA2 CH1 region, IgA2 CH2 region and an IgA2 CH3 region; wherein the IgA2 heavy chain constant region comprises a N135Q amino acid substitution, numbering according to IMGT scheme, as compared to a corresponding relative to the WT IgA heavy chain constant region comprising an amino acid sequence of SEQ ID NO: 1.
112. The antibody or the functional fragment thereof of claim 111, wherein the heavy chain constant region further comprises:
- i. a N45.2G amino acid substitution,
 - ii. a N114T amino acid substitution,
 - iii. a I115L amino acid substitution,
 - iv. a T116S amino acid substitution, and
 - v. an amino acid substitution selected from the group consisting of N15.2G, N15.2Q, and N15.2T,

relative to the WT IgA2 heavy chain constant region comprising an amino acid sequence of SEQ ID NO: 1,

numbering according to IMGT scheme.

113. The antibody or the functional fragment thereof of claim 111 or claim 112, wherein the heavy chain constant region further comprises:

- i. a C86S amino acid substitution,
- ii. a P124R amino acid substitution,
- iii. a deletion of C147,
- iv. a deletion of Y148,
- v. a L15.3I amino acid substitution,
- vi. a T16S amino acid substitution or
- vii. a combination thereof,

relative to the WT IgA2 heavy chain constant region comprising an amino acid sequence of SEQ ID NO: 1, numbering according to IMGT scheme.

114. An antibody or a functional fragment thereof comprising:

- a. an antigen binding domain; and
- b. a constant domain, wherein the constant domain comprises an IgA2 heavy chain constant region comprising an IgA2 CH2 region and an IgA2 CH3 region, and wherein the IgA2 heavy chain constant region comprises:

- i. a C86S amino acid substitution,
- ii. a N114T amino acid substitution,
- iii. a I115L amino acid substitution,
- iv. a T116S amino acid substitution, and
- v. a N135Q amino acid substitution, relative to the WT IgA2 heavy chain constant region comprising an amino acid sequence of SEQ ID NO: 1, numbering according to IMGT scheme.

115. An antibody or a functional fragment thereof comprising:

- a. an antigen binding domain; and
- b. a constant domain, wherein the constant domain comprises an IgA2 heavy chain constant region comprising an IgA2 CH1 region, an IgA2 CH2 region, and an IgA CH3 region, and wherein the IgA2 heavy chain constant region comprises:

- i. a N45.2G amino acid substitution,
- ii. a C86S amino acid substitution,
- iii. a N114T amino acid substitution,

- iv. a I115L amino acid substitution,
- v. a T116S amino acid substitution,
- vi. a N135Q amino acid substitution,
- vii. a deletion of C147, and
- 5 viii. a deletion of Y148, relative to the WT IgA2 heavy chain constant region comprising an amino acid sequence of SEQ ID NO: 1, numbering according to IMGT scheme.

116. The antibody or the functional fragment thereof of claim 115, wherein the IgA2 heavy chain constant region comprises an amino acid sequence that is at least 95% identical to SEQ ID NO: 16.

117. An antibody or a functional fragment thereof comprising:

- 10 a. an antigen binding domain; and
 - b. a constant domain, wherein the constant domain comprises an IgA2 heavy chain constant region comprising an IgA2 CH1 region, an IgA2 CH2 region, and an IgA2 CH3 region, and wherein the IgA2 heavy chain constant region comprises:
 - i. a N45.2G amino acid substitution,
 - 15 ii. a C86S amino acid substitution,
 - iii. a N114T amino acid substitution,
 - iv. a I115L amino acid substitution,
 - v. a T116S amino acid substitution, and
 - vi. a deletion of C-terminal tail piece, relative to the WT IgA2 heavy chain constant
- 20 region comprising an amino acid sequence of SEQ ID NO: 1, numbering according to IMGT scheme.

118. The antibody or the functional fragment thereof of claim 117, wherein the deletion of C-terminal tail piece comprises a deletion of C-terminal amino acid residues P131-Y148, relative to the WT IgA heavy chain constant region comprising an amino acid sequence of SEQ ID NO: 1, numbering according to the IMGT

25 scheme.

119. The antibody or the functional fragment thereof of claim 117 or claim 118, wherein the IgA2 heavy chain constant region comprises an amino acid sequence that is at least 95% identical to SEQ ID NO: 17.

120. An antibody or a functional fragment thereof comprising:

- 30 a. an antigen binding domain; and
- b. a constant domain, wherein the constant domain comprises an IgA2 heavy chain constant region comprising an IgA2 CH1 region, an IgA2 CH2 region, and an IgA2 CH3 region; wherein the IgA2 heavy chain constant region comprises:
 - i. a N45.2G amino acid substitution,
 - ii. a N114T amino acid substitution,

- 5 iii. a I115L amino acid substitution,
 iv. a T116S amino acid substitution,
 v. a N135Q amino acid substitution,
 vi. an amino acid substitution selected from the group consisting of N15.2G, N15.2Q,
 and N15.2T,
 vii. a L15.3I amino acid substitution, and
 viii. a T16S amino acid substitution,
 relative to the WT IgA2 heavy chain constant region comprising an amino acid
 sequence of SEQ ID NO: 1,
10 numbering according to the IMGT scheme.
121. An antibody or a functional fragment thereof comprising:
- a. an antigen binding domain; and
 b. a constant domain, wherein the constant domain comprises an IgA2 heavy chain constant
 region comprising an IgA2 CH1 region, an IgA2 CH2 region, and an IgA2 CH3 region, and
15 wherein the IgA2 heavy chain constant region comprises:
- i. a N45.2G amino acid substitution,
 ii. a N114T amino acid substitution,
 iii. a I115L amino acid substitution,
 iv. a T116S amino acid substitution,
20 v. an amino acid substitution selected from the group consisting of N15.2G, N15.2Q,
 and N15.2T,
 vi. a L15.3I amino acid substitution,
 vii. a T16S amino acid substitution, and
 viii. a deletion of C-terminal amino acids P131-Y148, relative to the WT IgA2 heavy
25 chain constant region comprising an amino acid sequence of SEQ ID NO: 1,
 numbering according to the IMGT scheme.
122. The antibody of construct of claim 121, further comprising:
- i. a C86S amino acid substitution,
 ii. a P124R amino acid substitution,
30 iii. a deletion of C147,
 iv. a deletion of Y148, or
 v. a combination thereof,
 relative to the WT IgA2 heavy chain constant region comprising an amino acid sequence of SEQ ID
NO: 1, numbering according to IMGT scheme.

123. The antibody or the functional fragment thereof of claim 121 or claim 122, wherein the IgA2 heavy chain constant region comprises an amino acid sequence that is at least 95% identical to a sequence selected from any one of SEQ ID NOs: 18-21.
124. The antibody or the functional fragment thereof of any one of claims 121-123 wherein the antibody or
5 the functional fragment thereof is aglycosylated.
125. The antibody or the functional fragment thereof of any one of claims 121-124, wherein the antibody or the functional fragment thereof has increased circulatory half life than a corresponding IgA comprising one or more of wild type amino acid residues N45.2, N114, I115, T116, N15.2, L15.3, T16, or a C-terminal amino acid residues P131-Y148, numbering according to the IMGT scheme.
- 10 126. The antibody or the functional fragment thereof of any one of claims 121-125, wherein the antibody or the functional fragment thereof induces increased antibody dependent cell mediated cytotoxicity (ADCC), compared to a corresponding antibody comprising an IgG heavy chain constant domain.
127. The antibody or the functional fragment thereof of any one of claims 121-126, wherein the antibody or the functional fragment thereof exhibits increased thermostability, compared to a corresponding IgA
15 comprising one or more of wild type amino acid residues N45.2, N114, I115, T116, N15.2, L15.3, T16, or a C-terminal amino acid residues P131-Y148, numbering according to the IMGT scheme, numbering according to the IMGT scheme.
128. The antibody or the functional fragment thereof of any one of claims 121-127, wherein the antibody or the functional fragment thereof exhibits decreased glycosylation, compared to a corresponding IgA
20 comprising one or more of wild type amino acid residues N45.2, N114, I115, T116, N15.2, L15.3, T16, or a C-terminal amino acid residues P131-Y148, numbering according to the IMGT scheme.
129. The antibody or the functional fragment thereof of any one of claims 121-128, wherein the IgA heavy chain constant region exhibits binding to Fc α R expressed on an immune effector cell with increased affinity, compared to a comprising one or more of wild type amino acid residues N45.2, N114, I115, T116, N15.2,
25 L15.3, T16, or a C-terminal amino acid residues P131-Y148, numbering according to the IMGT scheme.
130. The antibody or the functional fragment thereof of any one of claims 1-129, wherein the IgA heavy chain constant region further comprises a hinge region.
131. The antibody or the functional fragment thereof of claim 130, wherein the hinge region comprises an IgA hinge amino acid sequence or a variant or a fragment thereof.
- 30 132. The antibody or the functional fragment thereof of claim 131, wherein the hinge region comprises a human IgA hinge amino acid sequence or a variant or a fragment thereof.
133. The antibody or the functional fragment thereof of claim 130 or 131, wherein the hinge is an IgA1 hinge or an IgA2 hinge, or a variant or a fragment thereof.

134. The antibody or the functional fragment thereof of any one of claims 1-133, wherein the constant domain further comprises a light chain constant region.
135. The antibody or the functional fragment thereof of claim 134, wherein the light chain constant region is a kappa light chain constant region, wherein the kappa light chain constant region comprises a sequence of SEQ ID NO: 31
136. The antibody or the functional fragment thereof of any one of claims 1-135, wherein the IgA heavy chain constant region further comprises one or more albumin binding regions.
137. The antibody or the functional fragment thereof of claim 136, wherein the one or more albumin binding domains are fused to the C-terminus of the CH3 region.
138. The antibody or the functional fragment thereof of claim 136, wherein the constant region comprises the light chain constant region and the one or more albumin binding domain is fused to the light chain constant region.
139. The antibody or the functional fragment thereof of any one of claims 136-138, wherein the antibody or the functional fragment thereof has a greater circulating half-life compared to a corresponding IgA antibody that does not comprise the one or more albumin binding domains.
140. The antibody or the functional fragment thereof of any one of claims 1-139, wherein the antibody or the functional fragment thereof exhibits decreased complement-dependent cytotoxicity (CDC) compared to a corresponding antibody comprising an IgG heavy chain constant domain when measured in a suitable *in vitro* CDC assay.
141. The antibody or the functional fragment thereof of any one of claims 1-140, wherein the antibody or the functional fragment thereof exhibits increased antibody-dependent cell-mediated cytotoxicity (ADCC) compared to a corresponding WT IgA antibody when measured in a suitable *in vitro* ADCC assay.
142. The antibody or the functional fragment thereof of any one of claims 1-141, wherein the antigen binding domain comprises a heavy chain variable region and a light chain variable region.
143. The antibody or the functional fragment thereof of claim 142, wherein the heavy chain variable region comprises at least one of complementarity determining regions (CDRs):
HC-CDR1 comprising an amino acid sequence selected from any one of SEQ ID NOs: 33-40,
HC-CDR2 comprising an amino acid sequence selected from any one of SEQ ID NOs: 41-48; and
HC-CDR3 comprising an amino acid sequence selected from any one of SEQ ID NOs: 49-56.
144. The antibody or the functional fragment thereof of claim 142 or claim 143, wherein the light chain variable region comprises at least one of complementarity determining regions (CDRs):
LC-CDR1 comprising an amino acid sequence selected from any one of SEQ ID NOs: 57-64;
LC-CDR2 comprising an amino acid sequence selected from any one of SEQ ID NOs: 65-72; and
LC-CDR3 comprising an amino acid sequence selected from any one of SEQ ID NOs: 73-80.

145. The antibody or the functional fragment thereof of any one of claims 142-144, wherein the heavy chain variable region comprises a sequence having at least 95% identity to an amino acid sequence selected from any one of SEQ ID NOs: 4, 7, 81-86.
146. The antibody or the functional fragment thereof of any one of claims 142-145, wherein the light chain variable region comprises a sequence having at least 95% identity to an amino acid sequence selected from any one of SEQ ID NOs: 5, 8, 95-100.
147. The antibody or the functional fragment thereof of any one of claims 143-146, wherein each of the heavy chain and light chain CDRs are derived from an IgG antibody.
148. The antibody or the functional fragment thereof of any one of claims 142-147, wherein the heavy chain variable region further comprises four framework regions (FWs): HC-FW1, HC-FW2, HC-FW3, and HC-FW4.
149. The antibody or the functional fragment thereof of any one of claims 142-148, wherein the light chain variable region further comprises four framework regions (FWs): LC-FW1, LC-FW2, LC-FW3, and LC-FW4.
150. The antibody or the functional fragment thereof of any one of claims 148-149, wherein each the heavy chain and light chain FW regions are derived from an IgG antibody.
151. The antibody or the functional fragment thereof of any one of claims 148-149, wherein each the heavy chain and light chain FW regions are derived from an IgA antibody.
152. The antibody or the functional fragment thereof of any one of claims 1-150, wherein the antibody or the functional fragment thereof is
153. The antibody is a chimeric antibody, a heavy chain antibody, a single chain antibody, a humanized antibody, a human antibody, a monoclonal antibody, a deimmunized antibody, a bispecific antibody, a multispecific antibody, a multivalent antibody, or a combination thereof.
154. The antibody or the functional fragment thereof of any one of claims 1-152, wherein the antigen-binding fragment comprises a Fab, Fab', Fab'-SH, Fv, scFv, F(ab')₂, a diabody, a linear antibody, a single domain antibodies (sdAb), a camelid VHH domain, or a multi-specific antibody formed from antibody fragments,
155. The antibody or the functional fragment thereof of any one of claims 1-153, wherein the variable domain specifically binds to GD2, CD20, CD47, CD38, CD19, EGFR, HER2, PD-L1, or CD25.
156. The antibody or the functional fragment thereof of any one of claims 1-154, wherein the antibody or the functional fragment thereof further comprises an enzyme, a substrate, cofactor, a fluorescent marker, a chemiluminescent marker, a peptide tag, a magnetic particle, a drug, a toxin, a radionuclide, a binding site for secondary antibodies, a metal binding domain, or a combination thereof.
157. A pharmaceutical composition of comprising the antibody or the functional fragment thereof of any one of claims 1-155 and a pharmaceutically acceptable carrier.

158. A method of treating a subject in need thereof, comprising administering to the subject a therapeutic dose of the antibody or the functional fragment thereof of any one of claims 1-155 or the pharmaceutical composition of claim 156.
159. The method of claim 156 or claim 157, wherein the antibody or the functional fragment thereof or the pharmaceutical composition is cytolytic to a target cell.
160. The method of claim 158, wherein the target cell is a cancer cell.
161. The method of any one of claims 157-159, wherein the antibody or the functional fragment thereof or the pharmaceutical composition inhibits tumor growth.
162. The method of any one of claims 157-160, wherein the antibody or the functional fragment thereof or the pharmaceutical composition is administered subcutaneously, intravenously, intradermally, intraperitoneally, orally, intramuscularly or intracranially.
163. The method of any one of claims 157-161, wherein the antibody or the functional fragment thereof or the pharmaceutical composition is administered to the subject in combination with a second therapeutic agent.
164. The method of claim 162, wherein the second therapeutic agent comprises an anti-cancer agent, a chemotherapeutic agent, radiation therapy, a cytotoxic agent, a NSAID, a corticosteroid, a dietary supplement such as an antioxidant, or a combination thereof.
165. The method of any one of claims 162-163, wherein the second therapeutic agent is administered prior to, concurrently, or after administering the antibody or the functional fragment thereof or the pharmaceutical composition.
166. An isolated nucleic acid encoding the antibody or the functional fragment thereof of any one of claims 1-155.
167. An isolated nucleic acid molecule encoding a heavy chain polypeptide, the isolated nucleic acid molecule comprising a first nucleic acid sequence encoding a IgA heavy chain constant region, wherein the first nucleic acid sequence is selected from any one of SEQ ID NOs: 25-32.
168. The isolated nucleic acid molecule of claim 166, further comprising a second nucleic acid sequence encoding a variable heavy chain region, wherein the second nucleic acid sequence is selected from any one of SEQ ID NOs:87-84.
169. A vector comprising the isolated nucleic acid molecule of any one of claims 165-167.
170. A host cell comprising the isolated nucleic acid molecule of any one of claims 165-167.
171. The host cell of claim 169, further comprising an isolated nucleic acid molecule encoding a light chain polypeptide, wherein the isolated nucleic acid molecule encoding the light chain polypeptide comprises a nucleic acid sequence encoding a variable light chain region, wherein the nucleic acid sequence is selected from any one of SEQ ID NOs:101-108.

172. The host cell of claim 170, wherein the isolated nucleic acid molecule encoding the light chain polypeptide further comprises a nucleic acid sequence encoding a kappa light chain constant region, wherein the nucleic acid sequence comprises a sequence of SEQ ID NO:32.
173. A host cell expressing the antibody or the functional fragment thereof of any one of claims 1-155.
- 5 174. The host cell of any one of claims 169-172, wherein the host cell is a bacterial cell or a mammalian cell.
175. The host cell of claim 173, wherein the host cell is a CHO cell, or a HEK293 cell.
176. A method of producing an antibody or the functional fragment thereof, the method comprising:
- (a) culturing the host cell of any one of claims 169-174 in a medium under conditions permitting expression of a polypeptide encoded by the isolated nucleic acid molecule and assembling of the antibody or
- 10 the functional fragment thereof; and
- (b) purifying the antibody or the functional fragment thereof from the cultured host cell or the medium of the host cell.
177. The method of claim 175, wherein the purifying is by size exclusion chromatography.
178. An antibody or a functional fragment thereof that selectively binds to a CD20 polypeptide or a variant
- 15 thereof, comprising:
- (a) an IgA heavy chain constant region, wherein the IgA heavy chain constant region comprises an amino acid sequence selected from any one of SEQ ID NOs:16-21; and
- (b) a variable heavy chain region, wherein the variable heavy chain region comprises at least one of a complementarity-determining region heavy chain 1 (HC-CDR1) of SEQ ID NO: 33, HC-CDR2 of SEQ ID
- 20 NO: 41, and HC-CDR3 of SEQ ID NO: 49;
- (c) a variable light chain region, wherein the variable light chain region comprises at least one of a complementarity-determining region light chain 1 (LC-CDR1) of SEQ ID NO:57, LC-CDR2 of SEQ ID NO:65, and LC-CDR3 of SEQ ID NO:73; or
- (d) the variable heavy chain of (b) and the variable light chain of (c).
- 25 179. The antibody or the functional fragment thereof of claim 177, wherein the variable heavy chain comprises a polypeptide sequence having at least 95% sequence identity to the amino acid sequence of SEQ ID NO: 81, and the variable light chain comprises a polypeptide sequence having at least 95% sequence identity to the amino acid sequence of SEQ ID NO: 95.
180. An antibody or a functional fragment thereof that selectively binds to a GD2 protein or a variant thereof,
- 30 comprising:
- (a) an IgA heavy chain constant region, wherein the IgA heavy chain constant region comprises an amino acid sequence selected from any one of SEQ ID NOs: 16-21; and

(b) a variable heavy chain region, wherein the variable heavy chain region comprises at least one of a complementarity-determining region heavy chain 1 (HC-CDR1) of SEQ ID NO: 34, HC-CDR2 of SEQ ID NO: 42, and HC-CDR3 of SEQ ID NO: 50;

5 (c) a variable light chain region, wherein the variable light chain region comprises at least one of a complementarity-determining region light chain 1 (LC-CDR1) of SEQ ID NO: 58, LC-CDR2 of SEQ ID NO: 66, and LC-CDR3 of SEQ ID NO: 74; or

(d) the variable heavy chain of (b) and the variable light chain of (c).

181. The antibody or the functional fragment thereof of claim 180, wherein the variable heavy chain comprises a polypeptide sequence having at least 95% sequence identity to the amino acid sequence of SEQ ID NO: 4, and the variable light chain comprises a polypeptide sequence having at least 95% sequence identity to the amino acid sequence of SEQ ID NO: 5.

182. An antibody or a functional fragment thereof that selectively binds to a Her2 protein or a variant thereof, comprising:

15 (a) an IgA heavy chain constant region, wherein the IgA heavy chain constant region comprises an amino acid sequence selected from SEQ ID NOs: 16-21; and

(b) a variable heavy chain region, wherein the variable heavy chain region comprises at least one of a complementarity-determining region heavy chain 1 (HC-CDR1) of SEQ ID NO: 35, HC-CDR2 of SEQ ID NO: 43, and HC-CDR3 of SEQ ID NO: 51;

20 (c) a variable light chain region, wherein the variable light chain region comprises at least one of a complementarity-determining region light chain 1 (LC-CDR1) of SEQ ID NO: 59, LC-CDR2 of SEQ ID NO: 67, and LC-CDR3 of SEQ ID NO: 75; or

(d) the variable heavy chain of (b) and the variable light chain of (c).

183. The antibody or the functional fragment thereof of claim 182, wherein the variable heavy chain comprises a polypeptide sequence having at least 95% sequence identity to the amino acid sequence of SEQ ID NO: 82, and the variable light chain comprises a polypeptide sequence having at least 95% sequence identity to the amino acid sequence of SEQ ID NO: 96.

184. An antibody or a functional fragment thereof that selectively binds to a gp75 protein or a variant thereof, comprising:

30 (a) an IgA heavy chain constant region, wherein the IgA heavy chain constant region comprises an amino acid sequence selected from SEQ ID NOs: 16-21; and

(b) a variable heavy chain region, wherein the variable heavy chain region comprises at least one of a complementarity-determining region heavy chain 1 (HC-CDR1) of SEQ ID NO: 36, HC-CDR2 of SEQ ID NO: 44, and HC-CDR3 of SEQ ID NO: 52;

- (c) a variable light chain region, wherein the variable light chain region comprises at least one of a complementarity-determining region light chain 1 (LC-CDR1) of SEQ ID NO: 60, LC-CDR2 of SEQ ID NO: 68, and LC-CDR3 of SEQ ID NO: 76; or
- (d) the variable heavy chain of (b) and the variable light chain of (c).
- 5 185. The antibody or the functional fragment thereof of claim 184, wherein the variable heavy chain comprises a polypeptide sequence having at least 95% sequence identity to the amino acid sequence of SEQ ID NO: 83, and the variable light chain comprises a polypeptide sequence having at least 95% sequence identity to the amino acid sequence of SEQ ID NO: 97.
186. An antibody or a functional fragment thereof that selectively binds to a CTLA4 protein or a variant thereof, comprising:
- 10 (a) an IgA heavy chain constant region, wherein the IgA heavy chain constant region comprises an amino acid sequence selected from SEQ ID NOs: 16-21; and
- (b) a variable heavy chain region, wherein the variable heavy chain region comprises at least one of a complementarity-determining region heavy chain 1 (HC-CDR1) of SEQ ID NO: 37, HC-CDR2 of SEQ ID
- 15 NO: 45, and HC-CDR3 of SEQ ID NO: 53;
- (c) a variable light chain region, wherein the variable light chain region comprises at least one of a complementarity-determining region light chain 1 (LC-CDR1) of SEQ ID NO: 61, LC-CDR2 of SEQ ID NO: 69, and LC-CDR3 of SEQ ID NO: 77; or
- (d) the variable heavy chain of (b) and the variable light chain of (c).
- 20 187. The antibody or the functional fragment thereof of claim 186, wherein the variable heavy chain comprises a polypeptide sequence having at least 95% sequence identity to the amino acid sequence of SEQ ID NO: 84, and the variable light chain comprises a polypeptide sequence having at least 95% sequence identity to the amino acid sequence of SEQ ID NO: 98.
188. An antibody or a functional fragment thereof that selectively binds to a CD47 protein or a variant thereof, comprising:
- 25 (a) an IgA heavy chain constant region, wherein the IgA heavy chain constant region comprises an amino acid sequence selected from SEQ ID NOs: 16-21; and
- (b) a variable heavy chain region, wherein the variable heavy chain region comprises at least one of a complementarity-determining region heavy chain 1 (HC-CDR1) of SEQ ID NO: 38, HC-CDR2 of SEQ ID
- 30 NO: 46, and HC-CDR3 of SEQ ID NO: 54;
- (c) a variable light chain region, wherein the variable light chain region comprises at least one of a complementarity-determining region light chain 1 (LC-CDR1) of SEQ ID NO: 62, LC-CDR2 of SEQ ID NO: 70, and LC-CDR3 of SEQ ID NO: 78; or
- (d) the variable heavy chain of (b) and the variable light chain of (c).

189. The antibody or the functional fragment thereof of claim 188, wherein the variable heavy chain comprises a polypeptide sequence having at least 95% sequence identity to the amino acid sequence of SEQ ID NO: 85, and the variable light chain comprises a polypeptide sequence having at least 95% sequence identity to the amino acid sequence of SEQ ID NO: 99.

5 190. An antibody or a functional fragment thereof comprising:

- a. an antigen binding domain; and
- b. a constant domain, wherein the constant domain comprises an IgA2 heavy chain constant region comprising an IgA2 CH1 region, an IgA2 CH2 region, and an IgA2 CH3 region, and wherein the IgA2 heavy chain constant region comprises:

- 10 i. a N45.2G amino acid substitution,
 - ii. a N135Q amino acid substitution,
 - iii. a deletion of C147, and
 - iv. a deletion of Y148,
- relative to the WT IgA2 heavy chain constant region comprising an amino acid
15 sequence of SEQ ID NO: 1, numbering according to IMGT scheme.

191. An antibody or a functional fragment thereof comprising:

- a. an antigen binding domain; and
- b. a constant domain, wherein the constant domain comprises an IgA2 heavy chain constant region comprising an IgA2 CH1 region, an IgA2 CH2 region, and an IgA2 CH3 region, and
20 wherein the IgA2 heavy chain constant region comprises:

- i. a N45.2G amino acid substitution,
- ii. a N114T amino acid substitution,
- iii. a I115L amino acid substitution,
- iv. a T116S amino acid substitution,
- 25 v. an amino acid substitution selected from the group consisting of N15.2G, N15.2Q, and N15.2T, and
- vi. a deletion of C-terminal amino acids P131-Y148, relative to the WT IgA heavy chain constant region comprising an amino acid sequence of SEQ ID NO: 1, numbering according to the IMGT scheme.

30 192. The antibody of construct of claim 191, further comprising:

- i. a C86S amino acid substitution,
- ii. a P124R amino acid substitution,
- iii. a deletion of C147,
- iv. a deletion of Y148,

- v. a L15.3I amino acid substitution,
- vi. a T16S amino acid substitution or
- vii. a combination thereof,

relative to the WT IgA heavy chain constant region comprising an amino acid sequence of SEQ ID

5 NO: 1, numbering according to IMGT scheme.

193. The antibody or the functional fragment thereof of claim 191 or claim 192, wherein the IgA2 heavy chain constant region comprises an amino acid sequence that is at least 95% identical to a sequence selected from any one of SEQ ID NOs: 18-21.

10 194. The antibody or the functional fragment thereof of any one of claims 191-194 wherein the antibody or the functional fragment thereof is aglycosylated.

195. The antibody or the functional fragment thereof of any one of claims 191-194, wherein the antibody or the functional fragment thereof has increased circulatory half-life than a corresponding IgA comprising one or more of wild type amino acid residues N45.2, N114, I115, T116, N15.2, or a C-terminal amino acid residues P131-Y148, numbering according to the IMGT scheme.

15 196. The antibody or the functional fragment thereof of any one of claims 121-125, wherein the antibody or the functional fragment thereof induces increased antibody dependent cell mediated cytotoxicity (ADCC), compared to a corresponding antibody comprising an IgG heavy chain constant domain.

20 197. The antibody or the functional fragment thereof of any one of claims 121-126, wherein the antibody or the functional fragment thereof exhibits increased thermostability, compared to a corresponding IgA comprising one or more of wild type amino acid residues N45.2, N114, I115, T116, N15.2, or a C-terminal amino acid residues P131-Y148, numbering according to the IMGT scheme, numbering according to the IMGT scheme.

25 198. The antibody or the functional fragment thereof of any one of claims 121-127, wherein the antibody or the functional fragment thereof exhibits decreased glycosylation, compared to a corresponding IgA comprising one or more of wild type amino acid residues N45.2, N114, I115, T116, N15.2, or a C-terminal amino acid residues P131-Y148, numbering according to the IMGT scheme.