

PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION****International Bureau****INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)**

(51) International patent classification⁶: B01J 31/02, 31/24, C07C 45/50	A1	(11) International publication number: WO 99/13982 (43) International publication date: 25 March 1999 (25.03.99)
(21) International application number: PCT/EP98/05491 (22) International filing date: 29 August 1998 (29.08.98) (30) Data relating to the priority: 197 40 672.6 16 September 1997 (16.09.97) DE (71) Applicant (for all designated States except US): AVENTIS RESEARCH & TECHNOLOGIES GMBH & CO. KG [DE/DE]; D-65926 Frankfurt am Main (DE). (72) Inventors; and (75) Inventors/Applicants (US only): BOGDANOVIC, Sandra [DE/DE]; Wolfgangstrasse 35, D-60322 Frankfurt am Main (DE).	(81) Designated states: JP, SG, US, European Patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published With the International Search Report. Before expiry of the period provided for amending the claims, will be republished if such amendments are received.	

As printed

(54) Title: METHOD FOR PREPARING ALDEHYDES BY HYDROFORMYLATION**(54) Bezeichnung:** VERFAHREN ZUR HERSTELLUNG VON ALDEHYDEN DURCH HYDROFORMULIERUNG**(57) Abstract**

The invention concerns a catalyst containing: (a) more than 70 wt. % of a polyethyleneglycol of formula (I): $H-(OCH_2CH_2)_n-OH$, in which: n is a whole number between 3 and 16; and the mean molecular weight in number is less than 650; (b) rhodium in elementary or combined form; (g) between 2 and 25 wt. % of a ligand comprising at least one phosphorus atom; (h) at most, 25 wt. % of water. The invention also concerns the use of said catalyst for preparing aldehydes.

(57) Zusammenfassung

Katalysator, enthaltend: (a) mehr als 70 Gew.-% eines Polyethylenglykols der Formel (I): $H-(OCH_2CH_2)_n-OH$, in der n eine ganze Zahl zwischen 3 und 16 darstellt; und das zahlenmittlere Molgewicht weniger als 650 beträgt; (b) Rhodium in elementarer oder gebundener Form; (g) 2 bis 25 Gew.-% eines Liganden, der mindestens 1 Phosphoratom aufweist; (h) höchstens 25 Gew.-% Wasser; sowie dessen Verwendung zur Herstellung von Aldehyden.