What is claimed is:

1. ~~A method of treating incontinence associated with sexual activity comprising administering a therapeutically effective amount~~ Use of a botulinum toxin in the manufacture of a medicament for treating incontinence associated with sexual activity , wherein the medicament is to be administered to the bladder of a patient in need thereof~~, thereby treating the incontinence associated with sexual activity~~.

2. The ~~method~~ use of claim 1, wherein the botulinum toxin is selected from the group consisting of types A, B, C, D, E, and, G.

3. The ~~method~~ use of claim 2, wherein the botulinum toxin is type A or type B.

4. The ~~method~~ use of claim 1, wherein the incontinence associated with sexual activity occurs following a surgical procedure for a prostate disorder.

5. The ~~method~~ use of claim 4, wherein the surgical procedure is selected from the group consisting of radical prostatectomy, laparoscopic radical prostatectomy, transurethral resection of the prostate, transurethral microwave therapy, transurethral needle ablation, and cryosurgery.

6. ~~A method of preventing post-surgical incontinence associated with sexual activity comprising administering a therapeutically effective amount~~Use of a botulinum toxin in the menufacture of a medicamend for preventing post-surgical incontinence associated with sexual activity , wherein the medicament is to be administered to the bladder of a patient, wherein the botulinum toxin is to be administered prior to the onset of incontinence.

7. The ~~method~~ use of claim 6, wherein the patient is to be administered the botulinum toxin at a time selected from the group consisting of before surgery, during surgery, after surgery, and combinations thereof.

8. The ~~method~~ use of claim 7, wherein the patient is to be administered the botulinum toxin before surgery.

9. The ~~method~~ use of claim 7, wherein the patient is to be administered the botulinum toxin during surgery.

10. The ~~method~~ use of claim 8, wherein the patient is to be further administered botulinum toxin after surgery.

11. ~~A method of treating incontinence associated with sexual activity in a patient previously diagnosed with a urogenic abnormality, comprising administering a therapeutically effective amount~~Use of a botulinum toxin in the manufacture of a medicament for treating incontinence associated with sexual activity in a patient previously diagnosed with a urogenic abnormality to a patient in need thereof~~, thereby treating the incontinence associated with sexual activity~~.

12. The ~~method~~ use of claim 11, wherein the botulinum toxin is selected from the group consisting of types A, B, C, D, E, and, G.

13. A method of treating incontinence associated with sexual activity comprising

a) administering a therapeutically effective amount of a botulinum toxin to the bladder of a patient in need thereof, and;

b) administering a therapeutically effective amount of an anitcholinergic drug to a patient in need thereof;

thereby treating the incontinence associated with sexual activity.

14. The method of claim 13, wherein the botulinum toxin is selected from the group consisting of types A, B, C, D, E, and, G.

15. The method of claim 14, wherein the botulinum toxin is type A or type B.