
Claims

1. Use of a DHODH inhibitor compound for the manufacture of a medicament for use in a local ocular route administration in a method to treat an ocular disease, wherein the compound is  .
2. The use according to claim 1, wherein the ocular disease is uveitis, dry eye, age-related macular degeneration (AMD), optic neuritis, retrobulbar neuritis, ocular inflammation or discomfort or trauma caused by or associated with the use of contact lenses, ocular inflammation, discomfort or trauma caused by or associated with refractive surgery, blepharitis, an optic nerve disease or disorder, or a conjunctivitis condition.
3. The use according to claim 1 or 2, wherein a subject is suffering from uveitis, dry eye, age-related macular degeneration (AMD), conjunctivitis, keratitis, keratoconjunctivitis, vernal keratoconjunctivitis (VKC), and atopic keratoconjunctivitis (AKC), or the subject is suffering from uveitis, dry eye, age-related macular degeneration (AMD), or conjunctivitis, or the subject is suffering from age-related macular degeneration (AMD).
4. The use according to any one of claims 1 to 3, wherein the ocular disease is caused by an adenovirus.
5. The use according to any one of claims 1 to 4, wherein said compound is to be administered to the subject in a therapeutically effective amount.
6. The use according to any one of claims 1 to 5, wherein said compound is to be administered to the subject at least once per day or said compound is administered as a slow release formulation at least once per month, or at least every second month, or at least every third month, preferably at least every sixth month.
7. The use according to any one of claims 1 to 5 in a method for treating an ocular disease through injection of said compound into the vitreous body with a frequency of injections not exceeding one injection per month.
8. A pharmaceutical composition comprising the compound according to claim 1, which is formulated for a local ocular route administration such as intravitreous, topical, periocular injections (subconjunctival, peribulbar, latero-bulbar, retrobulbar, sub-tenon, suprachoroidal), intra- or periocular implants (intrascleral, periscleral,

episcleral), intravitreous implants or suprachoroidal implants or particles or polymeric composition, or any releasing systems such as emulsions, solid non-biodegradable or degradable implants or tablets, mini pumps or any topical formulations.

9. An ophthalmic formulation of a pharmaceutical composition according to claim 8 , optionally in form of a sterile eye drop formulation or as a sterile solution for intravitreal injection.
10. The formulation according to claim 9, wherein the frequency of administration is once daily or in the range from about one to about eight times a day, once per week, once per month, every second month or every third month or every sixth month.
11. The formulation according to claims 9 or 10, wherein a therapeutically effective amount of the compound reaches the posterior segment of the eye.