AZAINDOLES USEFUL AS INHIBITORS OF JAK AND OTHER PROTEIN KINASES

TECHNICAL FIELD OF THE INVENTION

[0001] The present invention relates to inhibitors of protein kinases. The invention also provides pharmaceutical compositions comprising the compounds of the invention and methods of using the compositions in the treatment of various disorders.

BACKGROUND OF THE INVENTION

[0002] The search for new therapeutic agents has been greatly aided in recent years by a better understanding of the structure of enzymes and other biomolecules associated with diseases. One important class of enzymes that has been the subject of extensive study is protein kinases.

[0003] Protein kinases constitute a large family of structurally related enzymes that are responsible for the control of a variety of signal transduction processes within the cell. (See, Hardie, G. and Hanks, S. *The Protein Kinase Facts Book, I and II,* Academic Press, San Diego, CA: 1995). Protein kinases are thought to have evolved from a common ancestral gene due to the conservation of their structure and catalytic function. Almost all kinases contain a similar 250-300 amino acid catalytic domain. The kinases may be categorized into families by the substrates they phosphorylate (e.g., protein-tyrosine, protein-serine/threonine, lipids, etc.). Sequence motifs have been identified that generally correspond to each of these kinase families (See, for example, Hanks, S.K., Hunter, T., *FASEB J.* **1995**, *9*, 576-596; Knighton *et al.*, *Science* **1991**, *253*, 407-414; Hiles *et al.*, *Cell* **1992**, *70*, 419-429; Kunz *et al.*, *Cell* **1993**, *73*, 585-596; Garcia-Bustos *et al.*, *EMBO J.* **1994**, *13*, 2352-2361).

[0004] In general, protein kinases mediate intracellular signaling by effecting a phosphoryl transfer from a nucleoside triphosphate to a protein acceptor that is involved in a signaling pathway. These phosphorylation events act as molecular on/off switches that can modulate or regulate the target protein biological function. These phosphorylation events are ultimately triggered in response to a variety of extracellular and other stimuli. Examples of such stimuli include environmental and chemical stress signals (e.g., osmotic shock, heat shock, ultraviolet radiation, bacterial endotoxin, and H_2O_2), cytokines (e.g., interleukin-1 (IL-1) and tumor necrosis factor α (TNF-α)), and growth factors (e.g., granulocyte macrophage-colony-stimulating factor (GM-CSF), and fibroblast growth factor (FGF)). An extracellular stimulus may affect one or more cellular responses related to cell growth, migration, differentiation, secretion of hormones, activation of transcription factors, muscle contraction, glucose metabolism, control of protein synthesis, and regulation of the cell cycle.

[0005] Many diseases are associated with abnormal cellular responses triggered by protein kinase-mediated events as described above. These diseases include, but are not limited to, autoimmune diseases, inflammatory diseases, bone diseases, metabolic diseases, neurological and neurodegenerative diseases, cancer, cardiovascular diseases, allergies and asthma, Alzheimer's disease, and hormone-related diseases. Accordingly, there has been a substantial effort in medicinal chemistry to find protein kinase inhibitors that are effective as therapeutic agents.

[0006] The Janus kinases (JAK) are a family of tyrosine kinases consisting of JAK1, JAK2, JAK3 and TYK2. The JAKs play a critical role in cytokine signaling. The down-stream substrates of the JAK family of kinases include the signal transducer and activator of transcription (STAT) proteins. JAK/STAT signaling has been implicated in the mediation of many abnormal immune responses such as allergies, asthma, autoimmune diseases such as transplant rejection, rheumatoid arthritis, amyotrophic lateral sclerosis and multiple sclerosis as well as in solid and hematologic malignancies such as leukemias and lymphomas. The pharmaceutical intervention in the JAK/STAT pathway has been reviewed [*Frank Mol. Med.* **5**, 432-456 (1999) & Seidel, et al, *Oncogene* **19**, 2645-2656 (2000)].

[0007] JAK1, JAK2, and TYK2 are ubiquitously expressed, while JAK3 is predominantly expressed in hematopoietic cells. JAK3 binds exclusively to the common cytokine receptor gamma chain (γ_c) and is activated by IL-2, IL-4, IL-7, IL-9, and IL-15. The proliferation and survival of murine mast cells induced by IL-4 and IL-9 have, in fact, been shown to be dependent on JAK3- and γ_c- signaling [Suzuki et al, *Blood* **96**, 2172-2180 (2000)].

[0008] Cross-linking of the high-affinity immunoglobulin (Ig) E receptors of sensitized mast cells leads to a release of proinflammatory mediators, including a number of vasoactive cytokines resulting in acute allergic, or immediate (type I) hypersensitivity reactions [Gordon et al, *Nature* **346**, 274-276 (1990) & Galli, *N. Engl. J. Med.*, **328**, 257-265 (1993)]. A crucial role for JAK3 in IgE receptor-mediated mast cell responses *in vitro* and *in vivo* has been established [Malaviya, et al, *Biochem. Biophys. Res. Commun.* **257**, 807-813 (1999)]. In addition, the prevention of type I hypersensitivity reactions, including anaphylaxis, mediated by mast cell-activation through inhibition of JAK3 has also been reported [Malaviya et al, *J. Biol. Chem.* **274**,27028-27038 (1999)]. Targeting mast cells with JAK3 inhibitors modulated mast cell degranulation *in vitro* and prevented IgE receptor/antigen-mediated anaphylactic reactions *in vivo*.

[0009] A recent study described the successful targeting of JAK3 for immune suppression and allograft acceptance. The study demonstrated a dose-dependent survival of Buffalo heart allograft in Wistar Furth recipients upon administration of inhibitors of JAK3 indicating the possibility of regulating unwanted immune responses in graft versus host disease [Kirken, *Transpl. Proc.* **33**, 3268-3270 (2001)].

[0010] IL-4-mediated STAT-phosphorylation has been implicated as the mechanism involved in early and late stages of rheumatoid arthritis (RA). Up-regulation of proinflammatory cytokines in RA synovium and synovial fluid is a characteristic of the disease. It has been demostrated that IL-4–mediated activation of IL-4/STAT pathway is mediated through the Janus Kinases (JAK 1 & 3) and that IL-4-associated JAK kinases are expressed in the RA synovium [Muller-Ladner, et al, *J. Immunol.* **164**, 3894-3901 (2000)].

[0011] Familial amyotrophic lateral sclerosis (FALS) is a fatal neurodegenerative disorder affecting about 10% of ALS patients. The survival rates of FALS mice were

increased upon treatment with a JAK3 specific inhibitor. This suggested that JAK3 plays a role in FALS [Trieu, et al, *Biochem. Biophys. Res. Commun.* **267**, 22-25 (2000)].

[0012] Signal transducer and activator of transcription (STAT) proteins are activated by, among others, the JAK family kinases. Results form a recent study suggested the possibility of intervention in the JAK/STAT signaling pathway by targeting JAK family kinases with specific inhibitors for the treatment of leukemia [Sudbeck, et al, *Clin. Cancer Res.* **5**, 1569-1582 (1999)]. JAK3 specific compounds were shown to inhibit the clonogenic growth of JAK3-expressing cell lines DAUDI, RAMOS, LC1;19, NALM-6, MOLT-3 and HL-60.

[0013] In animal models, TEL/JAK2 fusion proteins have induced myeloproliferative disorders and in hematopoietic cell lines, introduction of TEL/JAK2 resulted in activation of STAT1, STAT3, STAT5, and cytokine-independent growth [Schwaller, et al, *EMBO J.* **17**, 5321-5333 (1998)].

[0014] Inhibition of JAK 3 and TYK 2 abrogated tyrosine phosphorylation of STAT3, and inhibited cell growth of mycosis fungoides, a form of cutaneous T cell lymphoma. These results implicated JAK family kinases in the constitutively activated JAK/STAT pathway that is present in mycosis fungoides [Nielsen, et al, *Proc. Nat. Acad. Sci. U.S.A.* **94**, 6764-6769 (1997)]. Similarly, STAT3, STAT5, JAK1 and JAK2 were demonstrated to be constitutively activated in mouse T cell lymphoma characterized initially by LCK over-expression, thus further implicating the JAK/STAT pathway in abnormal cell growth [Yu, et al, *J. Immunol.* **159**, 5206-5210 (1997)]. In addition, IL-6 –mediated STAT3 activation was blocked by an inhibitor of JAK, leading to sensitization of myeloma cells to apoptosis [Catlett-Falcone, et al, *Immunity* **10**, 105-115 (1999)].

[0015] One kinase family of interest is Rho-associated coiled-coil forming protein serine/threonine kinase (ROCK), which is believed to be an effector of Ras-related small GTPase Rho. The ROCK family includes p160ROCK (ROCK-1) (Ishizaki *et al., EMBO J.* **1996**, *15*, 1885-1893) and ROKα/Rho-kinase/ROCK-II (Leung *et al., J. Biol. Chem.* **1995**, *270*, 29051-29054; Matsui *et al., EMBO J.* **1996**, *15*, 2208-2216; Nakagawa *et al., FEBS Lett.* **1996**, *392*, 189-193), protein kinase PKN (Amano *et al., Science* **1996**, *271*, 648-650; Watanabe *et al., Science* **1996**, *271*, 645-648), and citron

and citron kinase (Madaule *et al. Nature*, **1998**, *394*, 491-494; Madaule *et al., FEBS Lett.* **1995**, *377*, 243-248). The ROCK family of kinases have been shown to be involved in a variety of functions including Rho-induced formation of actin stress fibers and focal adhesions (Leung *et al., Mol. Cell Biol.* **1996**, *16*, 5313-5327; Amano *et al., Science,* **1997**, *275*, 1308-1311; Ishizaki *et al., FEBS Lett.* **1997**, *404*, 118-124) and in downregulation of myosin phosphatase (Kimura *et al., Science,* **1996**, *273*, 245-248), platelet activation (Klages *et al., J. Cell. Biol.*, **1999**, *144*, 745-754), aortic smooth muscle contraction by various stimuli (Fu *et al., FEBS Lett.*, **1998**, *440*, 183-187), thrombin-induced responses of aortic smooth muscle cells (Seasholtz *et al., Cir. Res.*, **1999**, *84*, 1186-1193), hypertrophy of cardiomyocytes (Kuwahara *et al., FEBS Lett.*, **1999**, *452*, 314-318), bronchial smooth muscle contraction (Yoshii *et al., Am. J. Respir. Cell Mol. Biol.*, **1999**, *20*, 1190-1200), smooth muscle contraction and cytoskeletal reorganization of non-muscle cells (Fukata *et al., Trends in Pharm. Sci* **2001**, *22*, 32-39), activation of volume-regulated anion channels (Nilius *et al., J. Physiol.*, **1999**, *516*, 67-74), neurite retraction (Hirose *et al., J. Cell. Biol.*, **1998**, *141*, 1625-1636), neutrophil chemotaxis (Niggli, *FEBS Lett.*, **1999**, *445*, 69-72), wound healing (Nobes and Hall, *J. Cell. Biol.*, **1999**, *144*, 1235-1244), tumor invasion (Itoh *et al., Nat. Med.*, **1999**, *5*, 221-225) and cell transformation (Sahai *et al., Curr. Biol.*, **1999**, *9*, 136-145).

[0016] More specifically, ROCK has been implicated in various diseases and disorders including hypertension (Satoh *et al., J. Clin. Invest.* **1994**, *94*, 1397-1403; Mukai *et al., FASEB J.* **2001**, *15*, 1062-1064; Uehata *et al., Nature* **1997**, *389*, 990-994; Masumoto *et al., Hypertension*, **2001**, *38*, 1307-1310), cerebral vasospasm (Sato *et al., Circ. Res.* **2000**, *87*, 195-200; Miyagi *et al., J. Neurosurg.* **2000**, *93*, 471-476; Tachibana *et al., Acta Neurochir (Wien)* **1999**, *141*, 13-19), coronary vasospasm (Shimokawa *et al., Jpn. Cir. J.* **2000**, *64*, 1-12; Kandabashi *et al., Circulation* **2000**, *101*, 1319-1323; Katsumata *et al., Circulation* **1997**, *96*, 4357-4363; Shimokawa *et al., Cardiovasc. Res.* **2001**, *51*, 169-177; Utsunomiya *et al., J. Pharmacol.* **2001**, *134*, 1724-1730; Masumoto *et al., Circulation* **2002**, *105*, 1545-1547), bronchial asthma (Chiba *et al., Comp. Biochem. Physiol. C Pharmacol. Toxicol. Endocrinol.* **1995**, *11*, 351-357; Chiba *et al., Br. J. Pharmacol.* **1999**, *127*, 597-600; Chiba *et al., Br. J. Pharmacol.* **2001**, *133*, 886-890; Iizuka *et al., Eur. J. Pharmacol.* **2000**, *406*, 273-

279), preterm labor (Niro *et al., Biochem. Biophys. Res. Commun.* **1997**, *230*, 356-359; Tahara *et al., Endocrinology* **2002**, *143*, 920-929; Kupittayanant *et al., Pflugers Arch.* **2001**, *443*, 112-114), erectile dysfunction (Chitaley *et al., Nat. Med.* **2001**, *7*, 119-122; Mills *et al., J. Appl. Physiol.* **2001**, *91*, 1269-1273), glaucoma (Honjo *et al., Arch. Ophthalmol.* **2001**, 1171-1178; Rao *et al., Invest. Ophthalmol. Vis. Sci.* **2001**, *42*, 1029-1037), vascular smooth muscle cell proliferation (Shimokawa *et al., Cardiovasc. Res.* **2001**, *51*, 169-177; Morishige *et al., Arterioscler. Thromb. Vasc. Biol.* **2001**, *21*, 548-554; Eto *et al., Am. J. Physiol. Heart Circ. Physiol.* **2000**, *278*, H1744-H1750; Sawada *et al., Circulation* **2000**, *101*, 2030-2023; Shibata *et al., Circulation* **2001**, *103*, 284-289), myocardial hypertrophy (Hoshijima *et al., J. Biol. Chem.* **1998**, *273*, 7725-77230; Sah *et al., J. Biol. Chem.* **1996**, *271*, 31185-31190; Kuwahara *et al., FEBS Lett.* **1999**, *452*, 314-318; Yanazume *et al., J. Biol. Chem.* **2002**, *277*, 8618-8625), malignoma (Itoh *et al., Nat. Med.* **1999**, *5*, 221-225; Genda *et al., Hepatology* **1999**, *30*, 1027-1036; Somlyo *et al., Biochem. Biophys. Res. Commun.* **2000**, *269*, 652-659), ischemia/reperfusion-induced injury (Ikeda *et al., J. of Surgical Res.* **2003**, *109*, 155-160; Miznuma *et al. Transplantation* **2003**, *75*, 579-586), endothelial dysfunction (Hernandez-Perera *et al., Circ. Res.* **2000**, *87*, 616-622; Laufs *et al., J. Biol. Chem.* **1998**, *273*, 24266-24271; Eto *et al., Circ. Res.* **2001**, *89*, 583-590), Crohn's Disease and colitis (Segain *et al. Gastroenterology* **2003**, *124(5)*, 1180-1187), neurite outgrowth (Fournier *et al. J. Neurosci.* **2003**, *23*, 1416-1423), Raynaud's Disease (Shimokawa *et al. J. Cardiovasc. Pharmacol.* **2002**, *39*, 319-327), angina (Utsunomiya et al. Br. J. Pharmacol. 2001, 134, 1724-1730; Masumoto et al, Circulation 2002, 105, 1545-1547; Shimokawa et al, J. Cardiovasc. Pharmacol., 2002, 40, 751-761; Satoh et al., Jpn. J. Pharmacol., 2001, 87, 34-40), Alzheimer's disease (Zhou et al., Science 2003, 302, 1215-1218), benign prostatic hyperplasia (Rees et al., J. Urology, 2003, 170, 2517-2522), and atherosclerosis (Retzer *et al. FEBS Lett.* **2000**, *466*, 70-74; Ishibashi *et al. Biochim. Biophys. Acta* **2002**, *1590*, 123-130). Accordingly, the development of inhibitors of ROCK kinase would be useful as therapeutic agents for the treatment of disorders implicated in the ROCK kinase pathway.

[0017] The Aurora proteins are a family of three highly related serine/threonine kinases (termed Aurora-A, -B and –C) that are essential for progression through the

mitotic phase of cell cycle. Specifically Aurora-A plays a crucial role in centrosome maturation and segregation, formation of the mitotic spindle and faithfull segregation of chromosomes. Aurora-B is a chromosomal passenger protein that plays a central role in regulating the alignment of chromosomes on the meta-phase plate, the spindle assembly checkpoint and for the correct completion of cytokinesis.

[0018] Overexpression of Aurora-A, - B or -C has been observed in a range of human cancers including colorectal, ovarian, gastric and invasive duct adenocarcinomas. In addition amplification of the AURKA locus that encodes for Aurora-A correlates with poor prognosis for patients with node-negative breast cancer. Furthermore overexpression of Aurora-A has been shown to transform mammalian fibroblasts, giving rise to aneuploid cells containing multipolar spindles.

[0019] A number of studies have now demonstrated that depletion or inhibition of Aurora-A or –B in human cancer cell lines by siRNA, dominant negative or neutralising antibodies disrupts progression through mitosis with accumulation of cells with 4N DNA, and in some cases this is followed by endoreduplication and cell death.


[0020] Protein kinases are attractive and proven targets for new therapeutic agents to treat a range if human diseases, with examples including Gleevec and Tarceva. The Aurora kinases are especially attractive due to their association with numerous human cancers and the role they play in promoting proliferation of these cancer cells (Harrington et al., *Nature Med.*, 2004, 10: 262-267).

[0021] Accordingly, there is a great need to develop inhibitors of JAK, ROCK and Aurora, preferably JAK-3, ROCK and Aurora A, protein kinases that are useful in treating various diseases or conditions associated with JAK, ROCK and Aurora activation, particularly given the inadequate treatments currently available for the majority of these disorders.

SUMMARY OF THE INVENTION

[0022] It has now been found that compounds of this invention, and pharmaceutically acceptable compositions thereof, are effective as inhibitors of JAK, ROCK and Aurora protein kinases. In certain embodiments, these compounds are

effective as inhibitors of JAK-3, ROCK and Aurora protein kinases. These compounds have the general formula **I:**



(I)


or a pharmaceutically acceptable salt thereof, wherein R^1, R^2, R^3, R^4, X^1, X^2, X^3, R^5 and x are as defined below and in subsets herein.

[0023] These compounds and pharmaceutical compositions thereof are useful for treating or preventing a variety of disorders, including, but not limited to, heart disease, diabetes, Alzheimer's disease, immunodeficiency disorders, inflammatory diseases, hypertension, allergic diseases, autoimmune diseases, destructive bone disorders such as osteoporosis, proliferative disorders, infectious diseases, immunologically-mediated diseases, and viral diseases. The compositions are also useful in methods for preventing cell death and hyperplasia and therefore may be used to treat or prevent reperfusion/ischemia in stroke, heart attacks, and organ hypoxia. The compositions are also useful in methods for preventing thrombin-induced platelet aggregation. The compositions are especially useful for disorders such as chronic myelogenous leukemia (CML), acute myeloid leukemia (AML), acute promyelocytic leukemia (APL), rheumatoid arthritis, asthma, osteoarthritis, ischemia, cancer (including, but not limited to, ovarian cancer, breast cancer and endometrial cancer), liver disease including hepatic ischemia, heart disease such as myocardial infarction and congestive heart failure, pathologic immune conditions involving T cell activation, and neurodegenerative disorders.

DETAILED DESCRIPTION OF THE INVENTION

[0024] *1. General Description of Compounds of the Invention:*

[0025] The present invention relates to compounds of formula **I:**



I

or pharmaceutically acceptable salts thereof, wherein:

R^1 is T-R' or is $-Si(R')_3$;

R^2, R^3, and R^4 are each independently halogen, CN, NO_2, or V-R';

X^1, X^2 and X^3 are each independently N, or CH, wherein the hydrogen atom of CH is optionally replaced by R^5;

x is 1, 2, 3, or 4;

each occurrence of R^5 is independently halogen, CN, NO_2, or U-R';

T, V, and U are each independently a bond or an optionally substituted C_1-C_6 alkylidene chain, wherein up to two methylene units of the chain are optionally and independently replaced by –NR'-, -S-, -O-, -CS-, $-CO_2-$, -OCO-, -CO-, -COCO-, -CONR'-, -NR'CO-, $-NR'CO_2-$, $-SO_2NR'-$, $-NR'SO_2-$, -CONR'NR'-, -NR'CONR'-, -OCONR'-, -NR'NR'-, $-NR'SO_2NR'-$, -SO-, $-SO_2-$, -PO-, $-PO_2-$, or –POR'-; and

each occurrence of R' is independently hydrogen or an optionally substituted group selected from a C_1-C_6 aliphatic group, a 3-8-membered saturated, partially unsaturated, or fully unsaturated monocyclic ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur, or an 8-12 membered saturated, partially unsaturated, or fully unsaturated bicyclic ring system having 0-5 heteroatoms independently selected from nitrogen, oxygen, or sulfur; or two occurrences of R' are taken together with the atom(s) to which they are bound to form an optionally substituted 3-12 membered saturated, partially unsaturated, or fully unsaturated monocyclic or bicyclic ring having 0-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur. In certain embodiments, the two occurrences of R' that form a ring will be on a single substituent (e.g., R^1, R^2, R^3, R^4 or on a single R^5 substituent) and form a monocyclic or bicyclic ring. In other embodiments, the two occurrences of R' are on two substituents (e.g., on two R^5 substituents) and can form a bicyclic fused ring with the ring to which the R^5

substituents are attached. However, the two occurrences of R' do not form a tricyclic ring whether they are a bound to a single substituent or to two separate substituents.

[0026] In certain embodiments, for compounds described directly above:

a. if R^1 is substituted cyclopentyl, x is 1, X^1 and X^3 are CH, then X^2 is not C-R^5, where R^5 is fluoro or OMe;

b. if R^2 and R^3 are simultaneously H and R^1 and R^4 are independently selected from H or Me, x is 1, X^1 and X^3 are CH, then X^2 is not C-R^5, where R^5 is OMe, NO_2, or fluoro;

c. if R^1, R^2, R^3 and R^4 are simultaneously H, x is 1, R^5 is -SMe, NH_2 or an optionally substituted NH-piperidine, and X^1 and X^2 are N, then X^3 is not CH;

d. if R^2, R^3 and R^4 are simultaneously H, X^1, X^2 and X^3 are CH, and two R^5 form a fused optionally substituted bicyclic ring with the ring to which they are attached, then R^1 is not $CH_2CH_2N(Me)_2$.

e. if R^2 and R^3 are simultaneously H, R^4 is NH_2, and X^1, X^2 and X^3 are CH, then R^1 is not substituted phenyl;

f. if R^2, R^3 and R^4 are simultaneously H, then R^1 is not $Si(R')_3$.

g. if R^1, R^2 and R^4 are simultaneously H and (i) X^2 and X^3 are CH or CR^5 or (ii) any one of X^1, X^2 or X^3 are N, then R^3 is not phenyl or phenyl substituted with O-phenyl or $N(Me)_2$.

[0027] *2. Compounds and Definitions:*

[0028] Compounds of this invention include those described generally above, and are further illustrated by the classes, subclasses, and species disclosed herein. As used herein, the following definitions shall apply unless otherwise indicated. For purposes of this invention, the chemical elements are identified in accordance with the Periodic Table of the Elements, CAS version, Handbook of Chemistry and Physics, 75th Ed. Additionally, general principles of organic chemistry are described in "Organic Chemistry", Thomas Sorrell, University Science Books, Sausalito: 1999, and "March's Advanced Organic Chemistry", 5th Ed., Ed.: Smith, M.B. and March, J., John Wiley & Sons, New York: 2001, the entire contents of which are hereby incorporated by reference.

[0029] As described herein, compounds of the invention may optionally be substituted with one or more substituents, such as are illustrated generally above, or as exemplified by particular classes, subclasses, and species of the invention. It will be appreciated that the phrase "optionally substituted" is used interchangeably with the phrase "substituted or unsubstituted." In general, the term "substituted", whether preceded by the term "optionally" or not, refers to the replacement of hydrogen radicals in a given structure with the radical of a specified substituent. Unless otherwise indicated, an optionally substituted group may have a substituent at each substitutable position of the group, and when more than one position in any given structure may be substituted with more than one substituent selected from a specified group, the substituent may be either the same or different at every position. Combinations of substituents envisioned by this invention are preferably those that result in the formation of stable or chemically feasible compounds. The term "stable", as used herein, refers to compounds that are not substantially altered when subjected to conditions to allow for their production, detection, and preferably their recovery, purification, and use for one or more of the purposes disclosed herein. In some embodiments, a stable compound or chemically feasible compound is one that is not substantially altered when kept at a temperature of 40°C or less, in the absence of moisture or other chemically reactive conditions, for at least a week.

[0030] The term "aliphatic" or "aliphatic group", as used herein, means a straight-chain (i.e., unbranched) or branched, substituted or unsubstituted hydrocarbon chain that is completely saturated or that contains one or more units of unsaturation, or a monocyclic hydrocarbon or bicyclic hydrocarbon that is completely saturated or that contains one or more units of unsaturation, but which is not aromatic (also referred to herein as "carbocycle" "cycloaliphatic" or "cycloalkyl"), that has a single point of attachment to the rest of the molecule. Unless otherwise specified, aliphatic groups contain 1-20 aliphatic carbon atoms. In some embodiments, aliphatic groups contain 1-10 aliphatic carbon atoms. In other embodiments, aliphatic groups contain 1-8 aliphatic carbon atoms. In still other embodiments, aliphatic groups contain 1-6 aliphatic carbon atoms, and in yet other embodiments aliphatic groups contain 1-4 aliphatic carbon atoms. In some embodiments, "cycloaliphatic" (or "carbocycle" or "cycloalkyl") refers to a monocyclic C_3-C_8 hydrocarbon or bicyclic C_8-C_{12}

hydrocarbon that is completely saturated or that contains one or more units of unsaturation, but which is not aromatic, that has a single point of attachment to the rest of the molecule wherein any individual ring in said bicyclic ring system has 3-7 members. Suitable aliphatic groups include, but are not limited to, linear or branched, substituted or unsubstituted alkyl, alkenyl, alkynyl groups and hybrids thereof such as (cycloalkyl)alkyl, (cycloalkenyl)alkyl or (cycloalkyl)alkenyl.

[0031] The term "heteroaliphatic", as used herein, means aliphatic groups wherein one or two carbon atoms are independently replaced by one or more of oxygen, sulfur, nitrogen, phosphorus, or silicon. Heteroaliphatic groups may be substituted or unsubstituted, branched or unbranched, cyclic or acyclic, and include "heterocycle", "heterocyclyl", "heterocycloaliphatic", or "heterocyclic" groups.

[0032] The term "heterocycle", "heterocyclyl", "heterocycloaliphatic", or "heterocyclic" as used herein means non-aromatic, monocyclic, bicyclic, or tricyclic ring systems in which one or more ring members are an independently selected heteroatom. In some embodiments, the "heterocycle", "heterocyclyl", "heterocycloaliphatic", or "heterocyclic" group has three to fourteen ring members in which one or more ring members is a heteroatom independently selected from oxygen, sulfur, nitrogen, or phosphorus, and each ring in the system contains 3 to 7 ring members.

[0033] The term "heteroatom" means one or more of oxygen, sulfur, nitrogen, phosphorus, or silicon (including, any oxidized form of nitrogen, sulfur, phosphorus, or silicon; the quaternized form of any basic nitrogen or; a substitutable nitrogen of a heterocyclic ring, for example N (as in 3,4-dihydro-2*H*-pyrrolyl), NH (as in pyrrolidinyl) or NR^+ (as in N-substituted pyrrolidinyl)).

[0034] The term "unsaturated", as used herein, means that a moiety has one or more units of unsaturation.

[0035] The term "alkoxy", or "thioalkyl", as used herein, refers to an alkyl group, as previously defined, attached to the principal carbon chain through an oxygen ("alkoxy") or sulfur ("thioalkyl") atom.

[0036] The terms "haloalkyl", "haloalkenyl" and "haloalkoxy" means alkyl, alkenyl or alkoxy, as the case may be, substituted with one or more halogen atoms. The term "halogen" means F, Cl, Br, or I.

[0037] The term "aryl" used alone or as part of a larger moiety as in "aralkyl", "aralkoxy", or "aryloxyalkyl", refers to monocyclic, bicyclic, and tricyclic ring systems having a total of five to fourteen ring members, wherein at least one ring in the system is aromatic and wherein each ring in the system contains 3 to 7 ring members. The term "aryl" may be used interchangeably with the term "aryl ring". The term "aryl" also refers to heteroaryl ring systems as defined hereinbelow.

[0038] The term "heteroaryl", used alone or as part of a larger moiety as in "heteroaralkyl" or "heteroarylalkoxy", refers to monocyclic, bicyclic, and tricyclic ring systems having a total of five to fourteen ring members, wherein at least one ring in the system is aromatic, at least one ring in the system contains one or more heteroatoms, and wherein each ring in the system contains 3 to 7 ring members. The term "heteroaryl" may be used interchangeably with the term "heteroaryl ring" or the term "heteroaromatic".

[0039] An aryl (including aralkyl, aralkoxy, aryloxyalkyl and the like) or heteroaryl (including heteroaralkyl and heteroarylalkoxy and the like) group may contain one or more substituents and thus may be "optionally substituted". Unless otherwise defined above and herein, suitable substituents on the unsaturated carbon atom of an aryl or heteroaryl group are generally selected from halogen; -R°; -OR°; -SR°; phenyl (Ph) optionally substituted with R°; -O(Ph) optionally substituted with R°; -$(CH_2)_{1-2}$(Ph), optionally substituted with R°; -CH=CH(Ph), optionally substituted with R°; a 5-6 membered heteroaryl or heterocyclic ring optionally substituted with R°; -NO_2; -CN; -$N(R°)_2$; -NR°C(O)R°; -NR°C(S)R°; -NR°C(O)$N(R°)_2$; -NR°C(S)$N(R°)_2$; -NR°CO_2R°; -NR°NR°C(O)R°; -NR°NR°C(O)$N(R°)_2$; -NR°NR°CO_2R°; -C(O)C(O)R°; -C(O)CH_2C(O)R°; -CO_2R°; -C(O)R°; -C(S)R°; -C(O)$N(R°)_2$; -C(S)$N(R°)_2$; -OC(O)$N(R°)_2$; -OC(O)R°; -C(O)N(OR°) R°; -C(NOR°) R°; -$S(O)_2$R°; -$S(O)_3$R°; -$SO_2N(R°)_2$; -S(O)R°; -NR°$SO_2N(R°)_2$; -NR°SO_2R°; -N(OR°)R°; -C(=NH)-$N(R°)_2$; -$P(O)_2$R°; -$PO(R°)_2$; -$OPO(R°)_2$; or -$(CH_2)_{0-2}$NHC(O)R°; wherein each independent occurrence of R° is selected from hydrogen, optionally substituted C_{1-6} aliphatic, an unsubstituted 5-6 membered heteroaryl or heterocyclic ring, phenyl, -O(Ph), or -CH_2(Ph), or, notwithstanding the definition above, two independent occurrences of R°, on the same substituent or

different substituents, taken together with the atom(s) to which each R° group is bound, to form an optionally substituted 3-12 membered saturated, partially unsaturated, or fully unsaturated monocyclic or bicyclic ring having 0-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur.

[0040] Optional substituents on the aliphatic group of R° are selected from NH_2, $NH(C_{1-4}aliphatic)$, $N(C_{1-4}aliphatic)_2$, halogen, $C_{1-4}aliphatic$, OH, $O(C_{1-4}aliphatic)$, NO_2, CN, CO_2H, $CO_2(C_{1-4}aliphatic)$, $O(haloC_{1-4}$ aliphatic), or $haloC_{1-4}aliphatic$, wherein each of the foregoing $C_{1-4}aliphatic$ groups of R° is unsubstituted.

[0041] An aliphatic or heteroaliphatic group, or a non-aromatic heterocyclic ring may contain one or more substituents and thus may be "optionally substituted". Unless otherwise defined above and herein, suitable substituents on the saturated carbon of an aliphatic or heteroaliphatic group, or of a non-aromatic heterocyclic ring are selected from those listed above for the unsaturated carbon of an aryl or heteroaryl group and additionally include the following: =O, =S, $=NNHR^*$, $=NN(R^*)_2$, $=NNHC(O)R^*$, $=NNHCO_2(alkyl)$, $=NNHSO_2(alkyl)$, or $=NR^*$, where each R^* is independently selected from hydrogen or an optionally substituted C_{1-6} aliphatic group.

[0042] Unless otherwise defined above and herein, optional substituents on the nitrogen of a non-aromatic heterocyclic ring are generally selected from $-R^+$, $-N(R^+)_2$, $-C(O)R^+$, $-CO_2R^+$, $-C(O)C(O)R^+$, $-C(O)CH_2C(O)R^+$, $-SO_2R^+$, $-SO_2N(R^+)_2$, $-C(=S)N(R^{+1})_2$, $-C(=NH)-N(R^+)_2$, or $-NR^+SO_2R^+$; wherein R^+ is hydrogen, an optionally substituted C_{1-6} aliphatic, optionally substituted phenyl, optionally substituted -O(Ph), optionally substituted $-CH_2(Ph)$, optionally substituted $-(CH_2)_{1-2}(Ph)$; optionally substituted -CH=CH(Ph); or an unsubstituted 5-6 membered heteroaryl or heterocyclic ring having one to four heteroatoms independently selected from oxygen, nitrogen, or sulfur, or, notwithstanding the definition above, two independent occurrences of R^+, on the same substituent or different substituents, taken together with the atom(s) to which each R^+ group is bound, form an optionally substituted 3-12 membered saturated, partially unsaturated, or fully unsaturated monocyclic or bicyclic ring having 0-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur.



[0043] Optional substituents on the aliphatic group or the phenyl ring of R^+ are selected from $-NH_2$, $-NH(C_{1-4}$ aliphatic), $-N(C_{1-4}$ aliphatic$)_2$, halogen, C_{1-4} aliphatic, -OH, $-O(C_{1-4}$ aliphatic), $-NO_2$, -CN, $-CO_2H$, $-CO_2(C_{1-4}$ aliphatic), -O(halo C_{1-4} aliphatic), or halo(C_{1-4} aliphatic), wherein each of the foregoing C_{1-4}aliphatic groups of R^+ is unsubstituted.

[0044] The term "alkylidene chain" refers to a straight or branched carbon chain that may be fully saturated or have one or more units of unsaturation and has two points of attachment to the rest of the molecule.

[0045] The term "protecting group", as used herein, refers to an agent used to temporarily block one or more desired reactive sites in a multifunctional compound. In certain embodiments, a protecting group has one or more, or preferably all, of the following characteristics: a) reacts selectively in good yield to give a protected substrate that is stable to the reactions occurring at one or more of the other reactive sites; and b) is selectively removable in good yield by reagents that do not attack the regenerated functional group. Exemplary protecting groups are detailed in Greene, T.W., Wuts, P. G in "Protective Groups in Organic Synthesis", Third Edition, John Wiley & Sons, New York: 1999, the entire contents of which are hereby incorporated by reference. The term "nitrogen protecting group", as used herein, refers to an agents used to temporarily block one or more desired nitrogen reactive sites in a multifunctional compound. Preferred nitrogen protecting groups also possess the characteristics exemplified above, and certain exemplary nitrogen protecting groups are also detailed in Chapter 7 in Greene, T.W., Wuts, P. G in "Protective Groups in Organic Synthesis", Third Edition, John Wiley & Sons, New York: 1999, the entire contents of which are hereby incorporated by reference.

[0046] As detailed above, in some embodiments, two independent occurrences of R° (or R^+, R, R' or any other variable similarly defined herein), are taken together with the atom(s) to which they are bound to form an optionally substituted 3-12 membered saturated, partially unsaturated, or fully unsaturated monocyclic or bicyclic ring having 0-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur.

[0047] Exemplary rings that are formed when two independent occurrences of R° (or R^+, R, R' or any other variable similarly defined herein), are taken together with the atom(s) to which each variable is bound include, but are not limited to the

following: a) two independent occurrences of R^o (or R^+, R, R' or any other variable similarly defined herein) that are bound to the same atom and are taken together with that atom to form a ring, for example, $N(R^o)_2$, where both occurrences of R^o are taken together with the nitrogen atom to form a piperidin-1-yl, piperazin-1-yl, or morpholin-4-yl group; and b) two independent occurrences of R^o (or R^+, R, R' or any other variable similarly defined herein) that are bound to different atoms and are taken together with both of those atoms to form a ring, for example where a phenyl group is substituted with two occurrences of OR^o , these two occurrences of R^o are taken together with the oxygen atoms to which they are bound to form a fused 6-membered oxygen containing ring: . It will be appreciated that a variety of other rings can be formed when two independent occurrences of R^o (or R^+, R, R' or any other variable similarly defined herein) are taken together with the atom(s) to which each variable is bound and that the examples detailed above are not intended to be limiting.

[0048] Unless otherwise stated, structures depicted herein are also meant to include all isomeric (e.g., enantiomeric, diastereomeric, and geometric (or conformational)) forms of the structure; for example, the R and S configurations for each asymmetric center, (Z) and (E) double bond isomers, and (Z) and (E) conformational isomers. Therefore, single stereochemical isomers as well as enantiomeric, diastereomeric, and geometric (or conformational) mixtures of the present compounds are within the scope of the invention. Unless otherwise stated, all tautomeric forms of the compounds of the invention are within the scope of the invention. Additionally, unless otherwise stated, structures depicted herein are also meant to include compounds that differ only in the presence of one or more isotopically enriched atoms. For example, compounds having the present structures except for the replacement of hydrogen by deuterium or tritium, or the replacement of a carbon by a ^{13}C- or ^{14}C-enriched carbon are within the scope of this invention. Such compounds are useful, for example, as analytical tools or probes in biological assays.

[0049] *3. Description of Exemplary Compounds:*

[0050] As described generally above, R^1 is T-R', or is $-Si(R')_3$. In certain embodiments, when R^1 is T-R', T is an optionally substituted C_1-C_6alkylidene chain wherein up to two methylene units are optionally and independently replaced with –O-, -S-, -NR'-, -OCO-, -COO-, $-SO_2$-or –CO-, and R' is hydrogen, C_1-C_4-alkyl, or an optionally substituted 5- or 6-membered aryl or heteroaryl group. In other embodiments of R^1, R' may additionally be C_1-C_4-aliphatic. In other embodiments, when R^1 is $-Si(R')_3$, R' is hydrogen, C_1-C_4-alkyl, or an optionally substituted 5- or 6-membered saturated, partially unsaturated, or fully unsaturated monocyclic ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur. In still other embodiments, R^1 is hydrogen, C_1-C_4alkyl, -COR', $-SO_2R'$, or $-Si(R')_3$. In yet other embodiments, R^1 is hydrogen, methyl, ethyl, n-propyl, isopropyl, n-butyl, p-toluenesulfonyl (Ts), t-butyldimethylsilyl (TBS), triisopropylsilyl (TIPS), or triethylsilyl (TES). Exemplary R^1 groups are also depicted in Tables 1 and 2 herein.


[0051] As described generally above, R^2, R^3, and R^4 are each independently halogen, CN, NO_2, or V-R'. In certain embodiments, R^2, R^3, and R^4 are each independently hydrogen, R', halogen, CN, NO_2, $-N(R')_2$, $-CH_2N(R')_2$, -OR', $-CH_2OR'$, -SR', $-CH_2SR'$, -COOR', -NR'COR', $-CON(R')_2$, $-SO_2N(R')_2$, $-CONR'(CH_2)_2N(R')_2$, $-CONR'(CH_2)_3N(R')_2$, $-CONR'(CH_2)_4N(R')_2$, $-O(CH_2)_2OR'$, $O(CH_2)_3OR'$, $O(CH_2)_4OR'$, $-O(CH_2)_2N(R')_2$, $-O(CH_2)_3N(R')_2$, or $-O(CH_2)_4N(R')_2$. In other embodiments, R^2, R^3, and R^4 are each independently Cl, Br, F, -CN, -COOH, -COOMe, $-NH_2$, $-N(CH_3)_2$, $-N(Et)_2$, $-N(iPr)_2$, $-O(CH_2)_2OCH_3$, $-CONH_2$, $-COOCH_3$, -OH, $-CH_2OH$, $-NHCOCH_3$, $-SO_2NH_2$, $-SO_2N(Me)_2$, or an optionally substituted group selected from C_1-C_4alkyl, C_1-C_4alkyloxy, a 3-8-membered saturated, partially unsaturated, or fully unsaturated monocyclic ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur, or an 8-12 membered saturated, partially unsaturated, or fully unsaturated bicyclic ring system having 0-5 heteroatoms independently selected from nitrogen, oxygen, or sulfur. In still other embodiments, R^2, R^3, and R^4 are each hydrogen. In other embodiments, one of R^2, R^3, or R^4 is hydrogen. In yet other embodiments, two of R^2, R^3, or R^4 is hydrogen. In yet other embodiments, R^2 and R^4 are both hydrogen, and R^3 is halogen, CN, NO_2, or V-R'. In still other embodiments, R^2 and R^4 are both hydrogen, and R^3 is an

optionally substituted group selected from a 3-8-membered saturated, partially unsaturated, or fully unsaturated monocyclic ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur, or an 8-12 membered saturated, partially unsaturated, or fully unsaturated bicyclic ring system having 0-5 heteroatoms independently selected from nitrogen, oxygen, or sulfur. In yet other embodiments, R^2 and R^4 are both hydrogen, and R^3 is an optionally substituted 5- or 6-membered saturated, partially unsaturated, or fully unsaturated ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur. In yet other embodiments, R^2 and R^4 are both hydrogen, and R^3 is an optionally substituted ring selected from phenyl, pyridyl, pyrimidinyl, thiazolyl, oxazolyl, thienyl, furyl, pyrrolyl, pyrazolyl, triazolyl, pyrazinyl, thiadiazolyl, or oxadiazolyl. In yet other embodiments, when R^1, R^2 and R^4 are H, then R^3 is not an optionally substituted phenyl. In yet other embodiments, when R^1, R^2 and R^4 are H, then R^3 is not an aryl, heteroaryl, carbocyclyl or heterocyclyl ring. Exemplary R^2, R^3, and R^4 groups also include those shown below in Tables 1 and 2.


[0052] As described above, R^2, R^3, and R^4 are each optionally substituted, and in certain embodiments, R^2, R^3, and R^4 are each optionally and independently substituted with z occurrences of R^6, wherein z is 0-5 and R^6 is =O, =NR", =S, halogen, -CN, -NO_2, or Z-R", wherein Z is a bond or an optionally substituted C_1-C_6 alkylidene chain, wherein up to two methylene units of the chain are optionally and independently replaced by –NR"-, -S-, -O-, -CS-, -CO_2-, -OCO-, –CO-, -COCO-, -CONR"-, -NR"CO-, -NR"CO_2-, -SO_2NR"-, -NR"SO_2-, -CONR"NR"-, -NR"CONR"-, -OCONR"-, -NR"NR"-, -NR"SO_2NR"-, -SO-, -SO_2-, -PO-, -PO_2- or –POR"-, and and each occurrence of R" is independently hydrogen or an optionally substituted C_1-C_6 aliphatic group, a 3-8-membered saturated, partially unsaturated, or fully unsaturated monocyclic ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur, or an 8-12 membered saturated, partially unsaturated, or fully unsaturated bicyclic ring system having 0-5 heteroatoms independently selected from nitrogen, oxygen, or sulfur; or two occurrences of R" are taken together with the atom(s) to which they are bound to form an optionally substituted 3-12 membered saturated, partially unsaturated, or fully unsaturated monocyclic or bicyclic ring having 0-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur. In

other embodiments, z is 0, 1, 2, or 3, and each occurrence of R^6 is independently hydrogen, R", -CH_2R", halogen, CN, NO_2, -N(R")$_2$, -CH_2N(R")$_2$, -OR", -CH_2OR", -SR", -CH_2SR", -COOR", -NR"COR", -NR"COOR", -CON(R")$_2$, -SO_2N(R")$_2$, -CONR"$(CH_2)_2$N(R")$_2$, -CONR$(CH_2)_3$N(R")$_2$, -CONR"$(CH_2)_4$N(R")$_2$, -O$(CH_2)_2$OR", O$(CH_2)_3$OR", O$(CH_2)_4$OR", -O$(CH_2)_2$N(R")$_2$, -O$(CH_2)_3$N(R")$_2$, -O$(CH_2)_4$N(R")$_2$, -NR"CH(CH_2OH)R", -NR"CH(CH_2CH_2OH)R", -NR"(CH_2)R", -NR"$(CH_2)_2$R", -NR"$(CH_2)_3$R", -NR"$(CH_2)_4$R", -NR"(CH_2)N(R")$_2$, -NR"$(CH_2)_2$N(R")$_2$, -NR"$(CH_2)_3$N(R")$_2$, -NR"$(CH_2)_4$N(R")$_2$, -NR"(CH_2)OR", -NR"$(CH_2)_2$OR", -NR"$(CH_2)_3$OR", or –NR"$(CH_2)_4$OR". In other embodiments, R^6 may additionally be -NR"CH(CH_3)R". In still other embodiments, z is 1, 2, or 3 and each occurrence of R^6 is independently F, Cl, Br, CN, OH, NH_2, -CH_2OH, C_1-C_6alkyl, -O(C_1-C_6alkyl), -CH_2O(C_1-C_6alkyl), -CO(C_1-C_6alkyl), -COO(C_1-C_6alkyl), -NHSO_2(C_1-C_6alkyl), -SO_2NH_2, -$CONH_2$, -CON(C_1-C_6alkyl), -SO_2(C_1-C_6alkyl), -SO_2phenyl, phenyl, benzyl, -N(C_1-C_6alkyl)$_2$, or -S(C_1-C_6alkyl), wherein each of the foregoing phenyl, benzyl, and C_1-C_6alkyl groups is independently and optionally substituted, and wherein each of the foregoing C_1-C_6alkyl groups is linear, branched, or cyclic. Additional exemplary R^6 groups are depicted in Table 1.


[0053] As described generally above for compounds of formula **I**, X^1, X^2 and X^3 are each independently N, or CH, wherein the hydrogen atom of CH is optionally replaced by R^5. In certain embodiments, two of X^1, X^2, or X^3 is N, and the remaining one of X^1, X^2, or X^3 is CH, wherein the hydrogen atom of CH is optionally replaced by R^5. In certain other embodiments, one of X^1, X^2, or X^3 is N, and the remaining two of X^1, X^2, or X^3 is CH, wherein the hydrogen atom of CH is optionally replaced by R^5. In yet other embodiments, each of X^1, X^2 and X^3 is CH, wherein the hydrogen atom of CH is optionally replaced by R^5. In certain other exemplary embodiments, compounds have one of formulae **I-A**, **I-B**, **I-C** or **I-D**:




I-A



I-B




I-C



I-D.

[0054] In other embodiments, compounds have formula **I-E**:



I-E

[0055] As described generally for compounds of formula **I** above, x is 1, 2, 3, or 4; and each occurrence of R^5 is independently halogen, CN, NO_2, or U-R', wherein each occurrence of U is independently a bond or an optionally substituted C_1-C_6 alkylidene chain, wherein up to two methylene units of the chain are optionally and independently replaced by –NR'-, -S-, -O-, -CS-, -CO_2-, -OCO-, -CO-, -COCO-, -CONR'-, -NR'CO-, -NR'CO_2-, -SO_2NR'-, -NR'SO_2-, -CONR'NR'-, -NR'CONR'-, -OCONR'-, -NR'NR'-, -NR'SO_2NR'-, -SO-, -SO_2-, -PO-, -PO_2-, or –POR'-; and each occurrence of R' is independently hydrogen or an optionally substituted C_1-C_6 aliphatic group, a 3-8-membered saturated, partially unsaturated, or fully unsaturated monocyclic ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur, or an 8-12 membered saturated, partially unsaturated, or fully unsaturated bicyclic ring system having 0-5 heteroatoms independently selected from nitrogen, oxygen, or sulfur; or two occurrences of R', are taken together with the atom(s) to which they are bound to form an optionally substituted 3-12 membered saturated, partially unsaturated, or fully unsaturated monocyclic or bicyclic ring having 0-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur.

In certain embodiments, each occurrence of R^5 is independently hydrogen, R', -CH_2R', halogen, CN, NO_2, -N(R')$_2$, -CH_2N(R')$_2$, -OR', -CH_2OR', -SR', –CH_2SR', -COOR', -NR'COR', -NR'$COCH_2$R', -NR'$CO(CH_2)_2$R', -NR'COOR', -CON(R')$_2$, -

$SO_2N(R')_2$, -$CONR'(CH_2)_2N(R')_2$, -$CONR(CH_2)_3N(R')_2$, -$CONR'(CH_2)_4N(R')_2$, -$O(CH_2)_2OR'$, $O(CH_2)_3OR'$, $O(CH_2)_4OR'$, -$O(CH_2)_2N(R')_2$, -$O(CH_2)_3N(R')_2$, -$O(CH_2)_4N(R')_2$, -$NR'CH(CH_2OH)R'$, -$NR'CH(CH_2CH_2OH)R'$, -$NR'(CH_2)R'$, -$NR'(CH_2)_2R'$, -$NR'(CH_2)_3R'$, -$NR'(CH_2)_4R'$, -$NR'(CH_2)N(R')_2$, -$NR'(CH_2)_2N(R')_2$, -$NR'(CH_2)_3N(R')_2$, -$NR'(CH_2)_4N(R')_2$, -$NR'(CH_2)OR'$, -$NR'(CH_2)_2OR'$, -$NR'(CH_2)_3OR'$, or -$NR'(CH_2)_4OR'$. In certain embodiments, R^5 may also be -$NR'CH(CH_3)R'$, $NR'CH(CF_3)R'$, -$NR'CH(CH_3)C(O)OR'$, -$NR'CH(CF_3)C(O)OR'$,-$NR'CH(CH_2CH_3)R'$, -$NR'CH_2C(O)N(R')_2$, -$NR'CH(CH_3)C(O)N(R')_2$, $NR'CH(CF_3)C(O)N(R')_2$, -$NR'CH(CH_2CH_3)C(O)N(R')_2$, -$NR'CH(CH(CH_3)_2)C(O)N(R')_2$, -$NR'CH(C(CH_3)_3)C(O)N(R')_2$, -$NR'CH(CH_2CH(CH_3)_2)C(O)N(R')_2$, -$NR'CH(CH_2OR^9)C(O)N(R')_2$ or -$NR'CH(CH_2CH_2N(Me)_2)C(O)N(R')_2$. In certain exemplary embodiments, x is 1, 2, or 3, and at least one occurrence of R^5 is $-N(R')_2$, -$NR'CH(CH_2OH)R'$, -$NR'CH(CH_2CH_2OH)R'$, -$NR'(CH_2)R'$, -$NR'(CH_2)_2R'$, $NR'(CH_2)N(R')_2$, or -$NR'(CH_2)_2N(R')_2$. In other embodiments, x is 1, 2, or 3, and at least one occurrence of R^5 is -OR'. In other embodiments, x is 1, 2, or 3, and at least one occurrence of R^5 is halogen. In yet other embodiments, x is 1, 2, or 3, and at least one occurrence of R^5 is -$NR'COR'$, -$NR'COCH_2R'$, -$NR'CO(CH_2)_2R'$, -$NR'CH(CH_2OH)R'$, -$NR'CH(CH_2OMe)R'$, -$NR'CH(CH_2OEt)R'$, -$NR'CH(CH_2OCF_3)R'$, -$NR'CH(CH_2CH_2OH)R'$, -$NR'CH(CH_2CH_2OMe)R'$, -$NR'CH(CH_2CH_2OEt)R'$, -$NR'CH(CH_2CH_2OCF_3)R'$, -$NR'CH(CH_3)C(O)OR'$, -$NR'CH(CF_3)C(O)OR'$, -$NR'CH(CH_3)C(O)N(R')_2$, -$NR'CH(CF_3)C(O)N(R')_2$, -$NR'CH(CH_2CH_3)C(O)N(R')_2$, -$NR'CH(CH_2OH)C(O)N(R')_2$, -$NR'CH(CH_2OMe)C(O)N(R')_2$, -$NR'CH(CH_2OEt)C(O)N(R')_2$ or -$NR'CH(CH_2OCF_3)C(O)N(R')_2$, wherein R' is an optionally substituted C_1-C_4 aliphatic; $NHCH_2C(O)NHR'$, -$NHCH(CH_3)C(O)NHR'$, -$NHCH(CH_2CH_3)C(O)NHR'$, -$NHCH(CH(CH_3)_2)C(O)NHR'$, -$NHCH(C(CH_3)_3)C(O)NHR'$, -$NHCH(CH_2CH(CH_3)_2)C(O)NHR'$, -$NHCH(CH_2OH)C(O)NHR'$, -$NHCH(CH_2OMe)C(O)NHR'$ or -$NHCH(CH_2CH_2N(Me)_2)C(O)NHR'$, wherein R' is an optionally substituted C_1-C_4 aliphatic; -NHR', -$NH(CH_2)R'$, -$NH(CH_2)_2R'$, -$NHCH(CH_3)R'$, -$NHCH_2C(O)NHR'$, -$NHCH(CH_3)C(O)NHR'$, -$NHCH(CH_2CH_3)C(O)NHR'$, -$NHCH(CH(CH_3)_2)C(O)NHR'$, -$NHCH(C(CH_3)_3)C(O)NHR'$, -

$NHCH(CH_2CH(CH_3)_2)C(O)NHR'$, $-NHCH(CH_2OH)C(O)NHR'$, $-NHCH(CH_2OMe)C(O)NHR'$ or $-NHCH(CH_2CH_2N(Me)_2)C(O)NHR'$, wherein R' is an optionally substituted C_1-C_4 aliphatic; $-NHCH(CH_3)R'$, wherein R' is optionally substituted phenyl; H, halogen, CH_3, CF_3, COOH, COOMe or OR', wherein R' is C_1-C_4 aliphatic.

[0056] In still other embodiments, x is 1, 2, or 3, and at least one occurrence of R^5 is an optionally substituted C_1-C_6 aliphatic group, a 3-8-membered saturated, partially unsaturated, or fully unsaturated monocyclic ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur, or an 8-12 membered saturated, partially unsaturated, or fully unsaturated bicyclic ring system having 0-5 heteroatoms independently selected from nitrogen, oxygen, or sulfur.


[0057] In yet other embodiments, x is 1 or 2, and each occurrence of R^5 is independently halogen, R', CN, $-CH_2CN$, $-(CH_2)_2CN$, NO_2, $-CH_2NO_2$, $-(CH_2)_2NO_2$, $CON(R')_2$, $-CH_2CON(R')_2$, $-(CH_2)_2CON(R')_2$, COOR', $-CH_2COOR'$, $-(CH_2)_2COOR'$, $-SO_2N(R')_2$, $-CH_2SO_2N(R')_2$, $-(CH_2)_2SO_2N(R')_2$, $-NR'SO_2R'$, $-CH_2NR'SO_2R'$, $-(CH_2)_2NR'SO_2R'$, $NR'CON(R')_2$, $-CH_2NR'CON(R')_2$, $-(CH_2)_2NR'CON(R')_2$, $-NR'SO_2N(R')_2$, $-CH_2NR'SO_2N(R')_2$, $-(CH_2)_2NR'SO_2N(R')_2$, -COCOR', $-CH_2COCOR'$, $-(CH_2)_2COCOR'$, $-N(R')_2$, $-CH_2N(R')_2$, $-(CH_2)_2N(R')_2$, -OR', $-CH_2OR'$, $-(CH_2)_2OR'$, $-NR'COR'$, $-NR'COCH_2R'$, $-NR'CO(CH_2)_2R'$, $-CH_2NR'COR'$, or $-(CH_2)_2NR'COR'$. In yet other embodiments, R^5 is CN, $-CH_2CN$, $-(CH_2)_2CN$, $-NO_2$, $-CH_2NO_2$, $-(CH_2)_2NO_2$, OR', $-CH_2OR'$, $-CON(R')_2$, $-SO_2N(R')_2$, $-N(R')_2$, or R'. In still other embodiments, each occurrence of R^5 is independently hydrogen, halogen, CN, $-CH_2CN$, $-(CH_2)_2CN$, NO_2, $-CH_2NO_2$, $-(CH_2)_2NO_2$, $-CONH_2$, $-CON(C_1$-C_4alkyl), $-SO_2NH_2$, $-SO_2N(C_1$-C_4alkyl), NH_2, $-N(C_1$-C_4alkyl), -OH, $-O(C_1$-C_4alkyl), $-CH_2OH$, $-CH_2O(C_1$-C_4alkyl), or an optionally substituted 5- or 6-membered unsaturated ring wherein 0-3 ring carbon atoms is optionally replaced by oxygen, sulfur, or nitrogen.

[0058] As described generally for compounds of formula I above, R^5 is optionally substituted with y occurrences of R^7, wherein y is 0-5 and R^7 is =O, =NR", =S, halogen, -CN, $-NO_2$, or W-R", wherein W is a bond or an optionally substituted C_1-C_6 alkylidene chain, wherein up to two methylene units of the chain are optionally and independently replaced by –NR"-, -S-, -O-, -CS-, $-CO_2$-, -OCO-, -CO-, -COCO-, -


CONR"-, -NR"CO-, -NR"CO_2-, -SO_2NR"-, -NR"SO_2-, -CONR"NR"-, -NR"CONR"-, -OCONR"-, -NR"NR"-, -NR"SO_2NR"-, -SO-, -SO_2-, -PO-, -PO_2-, or –POR"-, and and each occurrence of R" is independently hydrogen or an optionally substituted C_1-C_6 aliphatic group, a 3-8-membered saturated, partially unsaturated, or fully unsaturated monocyclic ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur, or an 8-12 membered saturated, partially unsaturated, or fully unsaturated bicyclic ring system having 0-5 heteroatoms independently selected from nitrogen, oxygen, or sulfur; or two occurrences of R", are taken together with the atom(s) to which they are bound to form an optionally substituted 3-12 membered saturated, partially unsaturated, or fully unsaturated monocyclic or bicyclic ring having 0-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur. In other embodiments, y is 0, 1, 2, or 3, and each occurrence of R^7 is independently hydrogen, R", -CH_2R", halogen, CN, NO_2, -N(R")$_2$, -CH_2N(R")$_2$, -OR", -CH_2OR", -SR", -CH_2SR", -COOR", -NR"COR", -NR"COOR", -CON(R")$_2$, -SO_2N(R")$_2$, -CONR"$(CH_2)_2$N(R")$_2$, -CONR$(CH_2)_3$N(R")$_2$, -CONR"$(CH_2)_4$N(R")$_2$, -O$(CH_2)_2$OR", O$(CH_2)_3$OR", O$(CH_2)_4$OR", -O$(CH_2)_2$N(R")$_2$, -O$(CH_2)_3$N(R")$_2$, -O$(CH_2)_4$N(R")$_2$, -NR"CH(CH_2OH)R", -NR"CH(CH_2CH_2OH)R", -NR"(CH_2)R", –NR"$(CH_2)_2$R", -NR"$(CH_2)_3$R", -NR"$(CH_2)_4$R", -NR"(CH_2)N(R")$_2$, -NR"$(CH_2)_2$N(R")$_2$, -NR"$(CH_2)_3$N(R")$_2$, -NR"$(CH_2)_4$N(R")$_2$, -NR"(CH_2)OR", -NR"$(CH_2)_2$OR", -NR"$(CH_2)_3$OR", or –NR"$(CH_2)_4$OR". In certain embodiments, R^7 may also be -NR'CH(CH_3)R'. In other embodiments, y is 1, 2, or 3 and each occurrence of R^7 is independently F, Cl, Br, CN, OH, NH_2, -CH_2OH, C_1-C_6alkyl, -O(C_1-C_6alkyl), -CH_2O(C_1-C_6alkyl), -CO(C_1-C_6alkyl), -COO(C_1-C_6alkyl), -NHSO_2(C_1-C_6alkyl), -SO_2NH_2, -$CONH_2$, -CON(C_1-C_6alkyl), -SO_2(C_1-C_6alkyl), -SO_2phenyl, phenyl, benzyl, -N(C_1-C_6alkyl)$_2$, or -S(C_1-C_6alkyl), wherein each of the foregoing phenyl, benzyl, and C_1-C_6alkyl groups is independently and optionally substituted, and wherein each of the foregoing C_1-C_6alkyl groups is linear, branched, or cyclic. Additional exemplary R^7 groups are depicted in Table 1.

[0059] In still other embodiments, x is 1, 2, or 3; at least one occurence of R^5 is –N(R')$_2$, -NR'CH(CH_2OH)R', -NR'CH(CH_2CH_2OH)R', -NR'(CH_2)R', -NR'$(CH_2)_2$R', NR'(CH_2)N(R')$_2$,-NR'$(CH_2)_2$N(R')$_2$, -OR', -NR'COR', -NR'$COCH_2$R', or -NR'$CO(CH_2)_2$R'; and R' is a C_1-C_6aliphatic group or a 3-8-membered saturated,


partially unsaturated, or fully unsaturated monocyclic ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur, or an 8-12 membered saturated, partially unsaturated, or fully unsaturated bicyclic ring system having 0-5 heteroatoms independently selected from nitrogen, oxygen, or sulfur; or two occurrences of R' are taken together with the atom(s) to which they are bound to form an optionally substituted 3-12 membered saturated, partially unsaturated, or fully unsaturated monocyclic or bicyclic ring having 0-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur, wherein each occurrence of R' is optionally substituted with y occurrences of R^7. In certain embodiments, R' is hydrogen, C_1-C_6alkyl optionally substituted with 1-3 occurrences of R^7, or is a 5-10-membered monocyclic or bicyclic saturated, partially unsaturated or fully unsaturated ring having 0-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur, wherein the ring is optionally substituted with 1-3 occurrences of R^7. In other embodiments, R' is hydrogen, C_1-C_4alkyl optionally substituted with 1-3 occurrences of R^7, or is a ring selected from:




i




ii




iii




iv




v




vi




vii




viii




ix




x




xi




xii




xiii




xiv




xv




xvi




xvii




xviii




xix




xx




xxi




xxii




xxiii




xxiv




xxv




xxvi




xxvii




xxviii




xxix




xxx




xxxi




xxxii




xxxiii




xxxiv




xxxv




xxxvi




xxxvii




xxxviii




xxxix




xL




xLi




xLii




xLiii



xLiv




xLv




xLvi


wherein y and R^7 are described generally and in subsets above.

In another embodiment, R' may be  **xLvii** , wherein y and R^7 are described generally and in subsets above.


[0060] In yet other embodiments, R^5 is $-N(R')_2$ and the two occurrences of R' are taken together with the nitrogen atom to which they are bound to form an optionally substituted 3-10-membered monocyclic or bicyclic heterocyclic ring. In certain embodiments, the ring is selected from:




a




b




c




d




e




f




g




h




i




j




k




l



m



n




o

wherein y and R^7 are described generally and in subsets above.

[0061] Certain additional subsets of compounds of general formula I include:

[0062] **I. Compounds of formula I-A:**



I-A

wherein R^1, R^2, R^3, R^4, R^5 and x are each described generally above and in subsets described above and herein.

[0063] In some embodiments, for compounds of formula **I-A**:

a. R^1 is:

i. T-R', wherein T is a bond or an optionally substituted C_1-C_6alkylidene chain wherein up to two methylene units are optionally and independently replaced with –O-, -S-, -NR'-, -OCO-, -COO-, $-SO_2$-or –CO-, and R' is hydrogen, C_1-C_4-alkyl, or an optionally substituted 5- or 6-membered aryl or heteroaryl group, or

ii. $-Si(R')_3$, wherein R' is hydrogen, C_1-C_4-alkyl, or an optionally substituted 5- or 6-membered saturated, partially unsaturated, or fully unsaturated monocyclic ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur;

b. R^2, R^3, and R^4 are each independently hydrogen, R', halogen, CN, NO_2, $-N(R')_2$, $-CH_2N(R')_2$, -OR', $-CH_2OR'$, -SR', $-CH_2SR'$, -COOR', -NR'COR', $-NR'COCH_2R'$, $-NR'CO(CH_2)_2R'$, $-CON(R')_2$, $-SO_2N(R')_2$, $-CONR'(CH_2)_2N(R')_2$, $-CONR'(CH_2)_3N(R')_2$, $-CONR'(CH_2)_4N(R')_2$, $-O(CH_2)_2OR'$, $O(CH_2)_3OR'$, $O(CH_2)_4OR'$, $-O(CH_2)_2N(R')_2$, $-O(CH_2)_3N(R')_2$, or $-O(CH_2)_4N(R')_2$; wherein R^2, R^3, and R^4 are each optionally substituted with z occurrences of R^6, wherein z is 0-5 and R^6 is =O, =NR", =S, halogen, -CN, $-NO_2$, or Z-R", wherein Z is a bond or an optionally substituted C_1-C_6 alkylidene chain, wherein up to two methylene units of the chain are optionally and independently replaced by –NR"-, -S-, -O-, -CS-, $-CO_2$-, -OCO-, -CO-, -COCO-, -CONR"-, -NR"CO-, $-NR"CO_2$-, $-SO_2NR"$-, $-NR"SO_2$–, -CONR"NR"-, -NR"CONR"-, -OCONR"-, -NR"NR"-, $-NR"SO_2NR"$-, -SO-, $-SO_2$-, -PO-, $-PO_2$-, or –POR"-, and each occurrence of R" is independently hydrogen or an optionally substituted C_1-C_6 aliphatic group, a 3-8-membered saturated, partially unsaturated, or fully unsaturated monocyclic ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur,

or an 8-12 membered saturated, partially unsaturated, or fully unsaturated bicyclic ring system having 0-5 heteroatoms independently selected from nitrogen, oxygen, or sulfur; or two occurrences of R" are taken together with the atom(s) to which they are bound to form an optionally substituted 3-12 membered saturated, partially unsaturated, or fully unsaturated monocyclic or bicyclic ring having 0-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur; and

c. each occurrence of R^5 is independently hydrogen, R', -CH_2R', halogen, CN, NO_2, -N(R')$_2$, -CH_2N(R')$_2$, -OR', -CH_2OR', -SR', -CH_2SR', -COOR', -NR'COR', -NR'$COCH_2$R', -NR'$CO(CH_2)_2$R', -NR'COOR', -CON(R')$_2$, -SO_2N(R')$_2$, -CONR'$(CH_2)_2$N(R')$_2$, -CONR$(CH_2)_3$N(R')$_2$, -CONR'$(CH_2)_4$N(R')$_2$, -O$(CH_2)_2$OR', O$(CH_2)_3$OR', O$(CH_2)_4$OR', -O$(CH_2)_2$N(R')$_2$, -O$(CH_2)_3$N(R')$_2$, -O$(CH_2)_4$N(R')$_2$, -NR'CH(CH_2OH)R', -NR'CH(CH_2CH_2OH)R', -NR'(CH_2)R', -NR'$(CH_2)_2$R', -NR'$(CH_2)_3$R', -NR'$(CH_2)_4$R', -NR'(CH_2)N(R')$_2$, -NR'$(CH_2)_2$N(R')$_2$, -NR'$(CH_2)_3$N(R')$_2$, -NR'$(CH_2)_4$N(R')$_2$, -NR'(CH_2)OR', -NR'$(CH_2)_2$OR', -NR'$(CH_2)_3$OR', or -NR'$(CH_2)_4$OR', wherein R^5 is optionally substituted with y occurrences of R^7, wherein y is 0-5 and R^7 is =O, =NR", =S, halogen, -CN, -NO_2, or W-R', wherein W is a bond or an optionally substituted C_1-C_6 alkylidene chain, wherein up to two methylene units of the chain are optionally and independently replaced by –NR"-, -S-, -O-, -CS-, -CO_2-, -OCO-, -CO-, -COCO-, -CONR"-, -NR"CO-, -NR"CO_2-, -SO_2NR"-, -NR"SO_2-, -CONR"NR"-, -NR"CONR"-, -OCONR"-, -NR"NR"-, -NR"SO_2NR"-, -SO-, -SO_2-, -PO-, -PO_2-, or –POR"-, and and each occurrence of R" is independently hydrogen or an optionally substituted C_1-C_6 aliphatic group, a 3-8-membered saturated, partially unsaturated, or fully unsaturated monocyclic ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur, or an 8-12 membered saturated, partially unsaturated, or fully unsaturated bicyclic ring system having 0-5 heteroatoms independently

selected from nitrogen, oxygen, or sulfur; or two occurrences of R" are taken together with the atom(s) to which they are bound to form an optionally substituted 3-12 membered saturated, partially unsaturated, or fully unsaturated monocyclic or bicyclic ring having 0-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur.

[0064] In still other embodiments, for compounds of formula **I-A**:

a. R^1 is hydrogen, C_1-C_4alkyl, -COR', $-SO_2R'$, or $-Si(R')_3$;

b. R^2, R^3, and R^4 are each independently Cl, Br, F, -CN, -COOH, -COOMe, $-NH_2$, $-N(CH_3)_2$, $-N(Et)_2$, $-N(iPr)_2$, $-O(CH_2)_2OCH_3$, $-CONH_2$, $-COOCH_3$, -OH, $-CH_2OH$, $-NHCOCH_3$, $-SO_2NH_2$, $-SO_2N(Me)_2$, or an optionally substituted group selected from C_1-C_4alkyl, C_1-C_4alkyloxy, a 3-8-membered saturated, partially unsaturated, or fully unsaturated monocyclic ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur, or an 8-12 membered saturated, partially unsaturated, or fully unsaturated bicyclic ring system having 0-5 heteroatoms independently selected from nitrogen, oxygen, or sulfur; wherein R^2, R^3, and R^4 are each independently and optionally substituted with z occurrences of R^6, wherein z is 0, 1, 2, or 3, and each occurrence of R^6 is independently hydrogen, R", $-CH_2R"$, halogen, CN, NO_2, $-N(R")_2$, $-CH_2N(R")_2$, -OR", $-CH_2OR"$, -SR", $-CH_2SR"$, -COOR", -NR"COR", -NR"COOR", $-CON(R")_2$, $-SO_2N(R")_2$, $-CONR"(CH_2)_2N(R")_2$, $-CONR(CH_2)_3N(R")_2$, $-CONR"(CH_2)_4N(R")_2$, $-O(CH_2)_2OR"$, $O(CH_2)_3OR"$, $O(CH_2)_4OR"$, $-O(CH_2)_2N(R")_2$, $-O(CH_2)_3N(R")_2$, $-O(CH_2)_4N(R")_2$, $-NR"CH(CH_2OH)R"$, $-NR"CH(CH_2CH_2OH)R"$, $-NR"(CH_2)R"$, $-NR"(CH_2)_2R"$, $-NR"(CH_2)_3R"$, $-NR"(CH_2)_4R"$, $-NR"(CH_2)N(R")_2$, $-NR"(CH_2)_2N(R")_2$, $-NR"(CH_2)_3N(R")_2$, $-NR"(CH_2)_4N(R")_2$, $-NR"(CH_2)OR"$, $-NR"(CH_2)_2OR"$, $-NR"(CH_2)_3OR"$, or $-NR'(CH_2)_4OR"$;

c. each occurrence of R^5 is independently hydrogen, R', $-CH_2R'$, halogen, CN, NO_2, $-N(R')_2$, $-CH_2N(R')_2$, -OR', $-CH_2OR'$, -SR', -

CH_2SR', -COOR', -NR'COR', -NR'$COCH_2R'$, -NR'$CO(CH_2)_2R'$, -NR'COOR', -$CON(R')_2$, -$SO_2N(R')_2$, -$CONR'(CH_2)_2N(R')_2$, -$CONR(CH_2)_3N(R')_2$, -$CONR'(CH_2)_4N(R')_2$, -$O(CH_2)_2OR'$, $O(CH_2)_3OR'$, $O(CH_2)_4OR'$, -$O(CH_2)_2N(R')_2$, -$O(CH_2)_3N(R')_2$, -$O(CH_2)_4N(R')_2$, -NR'$CH(CH_2OH)R'$, -NR'$CH(CH_2CH_2OH)R'$, -NR'$(CH_2)R'$, -NR'$(CH_2)_2R'$, -NR'$(CH_2)_3R'$, -NR'$(CH_2)_4R'$, -NR'$(CH_2)N(R')_2$, -NR'$(CH_2)_2N(R')_2$, -NR'$(CH_2)_3N(R')_2$, -NR'$(CH_2)_4N(R')_2$, -NR'$(CH_2)OR'$, -NR'$(CH_2)_2OR'$, -NR'$(CH_2)_3OR'$, or -NR'$(CH_2)_4OR'$, wherein R^5 is optionally substituted with y occurrences of R^7, wherein y is 0, 1, 2, or 3, and each occurrence of R^7 is independently hydrogen, R", -CH_2R", halogen, CN, NO_2, -$N(R")_2$, -$CH_2N(R")_2$, -OR", -CH_2OR", -SR", -CH_2SR", -COOR", -NR"COR", -NR"COOR", -$CON(R")_2$, -$SO_2N(R")_2$, -$CONR"(CH_2)_2N(R")_2$, -$CONR(CH_2)_3N(R")_2$, -$CONR"(CH_2)_4N(R")_2$, -$O(CH_2)_2OR$", $O(CH_2)_3OR$", $O(CH_2)_4OR$", -$O(CH_2)_2N(R")_2$, -$O(CH_2)_3N(R")_2$, -$O(CH_2)_4N(R")_2$, -NR"$CH(CH_2OH)R$", -NR"$CH(CH_2CH_2OH)R$", -NR"$(CH_2)R$", -NR"$(CH_2)_2R$", -NR"$(CH_2)_3R$", -NR"$(CH_2)_4R$", -NR"$(CH_2)N(R")_2$, -NR"$(CH_2)_2N(R")_2$, -NR"$(CH_2)_3N(R")_2$, -NR"$(CH_2)_4N(R")_2$, -NR"$(CH_2)OR$", -NR"$(CH_2)_2OR$", -NR"$(CH_2)_3OR$", or –NR"$(CH_2)_4OR$".

[0065] In other embodiments, for compounds of formula **I-A** and subsets described directly above, R^1 is hydrogen, methyl, ethyl, n-propyl, isopropyl, n-butyl, p-toluenesulfonyl (Ts), t-butyldimethylsilyl (TBS), triisopropylsilyl (TIPS), or triethylsilyl (TES).

[0066] In still other embodiments, for compounds of formula **I-A** and subsets described directly above, R^2, R^3, and R^4 are each hydrogen. In other embodiments, one of R^2, R^3, or R^4 is hydrogen. In yet other embodiments, two of R^2, R^3, or R^4 is hydrogen. In yet other embodiments, R^2 and R^4 are both hydrogen, and R^3 is halogen, CN, NO_2, or V-R'. In still other embodiments, R^2 and R^4 are both hydrogen, and R^3 is an optionally substituted group selected from a 3-8-membered saturated, partially unsaturated, or fully unsaturated monocyclic ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur, or an 8-12 membered

saturated, partially unsaturated, or fully unsaturated bicyclic ring system having 0-5 heteroatoms independently selected from nitrogen, oxygen, or sulfur. In yet other embodiments, R^2 and R^4 are both hydrogen, and R^3 is an optionally substituted 5- or 6-membered saturated, partially unsaturated, or fully unsaturated ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur. In yet other embodiments, R^2 and R^4 are both hydrogen, and R^3 is an optionally substituted ring selected from phenyl, pyridyl, pyrimidinyl, thiazolyl, oxazolyl, thienyl, furyl, pyrrolyl, pyrazolyl, triazolyl, pyrazinyl, thiadiazolyl, or oxadiazolyl. In yet other embodiments, R^3 is selected from H, Cl, Br, F, -CN, -COOH, -COOMe, $-NH_2$, $-N(R')_2$, $-NO_2$, -OR', $-CON(R')_2$, -COOR', -OH, -SR', $-C(R')_2OR'$, -N(R')COR', -N(R')C(O)OR', $-SO_2NH_2$, $-SO_2N(R')_2$, or an optionally substituted group selected from C_1-C_4 aliphatic, C_1-C_4 alkyloxy or $-C\equiv C-C_1$-C_4 aliphatic. In a further embodiment, R^2 and R^4 are both hydrogen and R^3 is selected from the immediately preceding list.

[0067] In other embodiments, for compounds of formula **I-A** and subsets described directly above, R^2, R^3, and R^4 are each independently and optionally substituted with z occurrences of R^6, wherein z is 1, 2, or 3 and each occurrence of R^6 is independently F, Cl, Br, CN, OH, NH_2, $-CH_2OH$, C_1-C_6alkyl, $-O(C_1$-C_6alkyl), $-CH_2O(C_1$-C_6alkyl), $-CO(C_1$-C_6alkyl), $-COO(C_1$-C_6alkyl), $-NHSO_2(C_1$-C_6alkyl), $-SO_2NH_2$, $-CONH_2$, $-CON(C_1$-C_6alkyl), $-SO_2(C_1$-C_6alkyl), $-SO_2$phenyl, phenyl, benzyl, $-N(C_1$-C_6alkyl$)_2$, or $-S(C_1$-C_6alkyl), wherein each of the foregoing phenyl, benzyl, and C_1-C_6alkyl groups is independently and optionally substituted, and wherein each of the foregoing C_1-C_6alkyl groups is linear, branched, or cyclic.


[0068] In certain exemplary embodiments, for compounds of formula **I-A** and subsets described directly above, x is 1, 2, or 3, and at least one occurrence of R^5 is $-N(R')_2$, $-NR'CH(CH_2OH)R'$, $-NR'CH(CH_2CH_2OH)R'$, $-NR'(CH_2)R'$, $-NR'(CH_2)_2R'$, $NR'(CH_2)N(R')_2$, or $-NR'(CH_2)_2N(R')_2$. In other embodiments, x is 1, 2, or 3, and at least one occurrence of R^5 is -OR'. In yet other embodiments, x is 1, 2, or 3, and at least one occurrence of R^5 is -NR'COR', $-NR'COCH_2R'$, or $-NR'CO(CH_2)_2R'$. In still other embodiments, x is 1, 2, or 3, and at least one occurrence of R^5 is an optionally substituted C_1-C_6 aliphatic group, a 3-8-membered saturated, partially unsaturated, or fully unsaturated monocyclic ring having 0-3 heteroatoms

independently selected from nitrogen, oxygen, or sulfur, or an 8-12 membered saturated, partially unsaturated, or fully unsaturated bicyclic ring system having 0-5 heteroatoms independently selected from nitrogen, oxygen, or sulfur.

[0069] In still other embodiments, for compounds of formula **I-A** and subsets described directly above, R^5 is optionally substituted with y occurrences of R^7, wherein y is 1, 2, or 3 and each occurrence of R^7 is independently F, Cl, Br, CN, OH, NH_2, $-CH_2OH$, C_1-C_6alkyl, $-O(C_1$-C_6alkyl), $-CH_2O(C_1$-C_6alkyl), $-CO(C_1$-C_6alkyl), $-COO(C_1$-C_6alkyl), $-NHSO_2(C_1$-C_6alkyl), $-SO_2NH_2$, $-CONH_2$, $-CON(C_1$-C_6alkyl), $-SO_2(C_1$-C_6alkyl), $-SO_2$phenyl, phenyl, benzyl, $-N(C_1$-C_6alkyl$)_2$, or $-S(C_1$-C_6alkyl), wherein each of the foregoing phenyl, benzyl, and C_1-C_6alkyl groups is independently and optionally substituted, and wherein each of the foregoing C_1-C_6alkyl groups is linear, branched, or cyclic. In yet other embodiments, for compounds of formula **I-A** and subsets described directly above, x is 1, 2, or 3; at least one occurence of R^5 is $-N(R')_2$, $-NR'CH(CH_2OH)R'$, $-NR'CH(CH_2CH_2OH)R'$, $-NR'(CH_2)R'$, $-NR'(CH_2)_2R'$, $NR'(CH_2)N(R')_2$, $-NR'(CH_2)_2N(R')_2$, $-OR'$, $-NR'COR'$, $-NR'COCH_2R'$, or $-NR'CO(CH_2)_2R'$; and R' is a C_1-C_6aliphatic group or a 3-8-membered saturated, partially unsaturated, or fully unsaturated monocyclic ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur, or an 8-12 membered saturated, partially unsaturated, or fully unsaturated bicyclic ring system having 0-5 heteroatoms independently selected from nitrogen, oxygen, or sulfur; or two occurrences of R', are taken together with the atom(s) to which they are bound to form an optionally substituted 3-12 membered saturated, partially unsaturated, or fully unsaturated monocyclic or bicyclic ring having 0-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur, wherein each occurrence of R' is optionally substituted with y occurrences of R^7. In certain embodiments, R' is hydrogen, C_1-C_6alkyl optionally substituted with 1-3 occurrences of R^7, or is a 5-10-membered monocyclic or bicyclic saturated, partially unsaturated or fully unsaturated ring having 0-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur, wherein the ring is optionally substituted with 1-3 occurrences of R^7. In other embodiments, R' is hydrogen, C_1-C_4alkyl optionally substituted with 1-3 occurrences of R^7, or is a ring selected from (i)-(xLvi) or (xLvii) described in [0059]. In yet other embodiments, R^5 is $-N(R')_2$ and the two occurrences of R' are taken together with the

nitrogen atom to which they are bound to form an optionally substituted 3-10-membered monocyclic or bicyclic heterocyclic ring. In certain embodiments, the ring is selected from (a)-(o) described in [0060].

[0070] In yet other embodiments, x is 1 and compounds have general formula **I-A-i**:




I-A-i

wherein R^1, R^2, R^3, and R^4 are described generally and in subsets above and herein, and R^5 is $-N(R')_2$, $-NR'CH(CH_2OH)R'$, $-NR'CH(CH_2CH_2OH)R'$, $-NR'(CH_2)R'$, $-NR'(CH_2)_2R'$, $NR'(CH_2)N(R')_2$, $-NR'(CH_2)_2N(R')_2$, $-OR'$, $-NR'COR'$, $-NR'COCH_2R'$, or $-NR'CO(CH_2)_2R'$,; and R' is a C_1-C_6aliphatic group or a 3-8-membered saturated, partially unsaturated, or fully unsaturated monocyclic ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur, or an 8-12 membered saturated, partially unsaturated, or fully unsaturated bicyclic ring system having 0-5 heteroatoms independently selected from nitrogen, oxygen, or sulfur; or two occurrences of R', are taken together with the atom(s) to which they are bound to form an optionally substituted 3-12 membered saturated, partially unsaturated, or fully unsaturated monocyclic or bicyclic ring having 0-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur, wherein each occurrence of R' is optionally substituted with y occurrences of R^7. In some embodiments, R^5 is $N(R')_2$. In certain embodiments, R' is hydrogen, C_1-C_6alkyl optionally substituted with 1-3 occurrences of R^7, or is a 5-10-membered monocyclic or bicyclic saturated, partially unsaturated or fully unsaturated ring having 0-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur, wherein the ring is optionally substituted with 1-3 occurrences of R^7. . In other embodiments, R' is hydrogen, C_1-C_4alkyl optionally substituted with 1-3 occurrences of R^7, or is a ring selected from (i)-(xLvi) or (xLvii) described in [0059]. In yet other embodiments, R^5 is $-N(R')_2$ and the two occurrences of R' are taken together with the nitrogen atom to which they are bound

to form an optionally substituted 3-10-membered monocyclic or bicyclic heterocyclic ring. In certain embodiments, the ring is selected from (a)-(o) described in [0060].

[0071] In still other embodiments, x is 1 and compounds have general formula **I-A-ii**:




I-A-ii

wherein R^1, R^2, R^3, and R^4 are described generally and in subsets above and herein, and R^5 is $-N(R')_2$, $-NR'CH(CH_2OH)R'$, $-NR'CH(CH_2CH_2OH)R'$, $-NR'(CH_2)R'$, $-NR'(CH_2)_2R'$, $NR'(CH_2)N(R')_2$, $-NR'(CH_2)_2N(R')_2$, $-OR'$, $-NR'COR'$, $-NR'COCH_2R'$, or $-NR'CO(CH_2)_2R'$; and R' is a C_1-C_6aliphatic group or a 3-8-membered saturated, partially unsaturated, or fully unsaturated monocyclic ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur, or an 8-12 membered saturated, partially unsaturated, or fully unsaturated bicyclic ring system having 0-5 heteroatoms independently selected from nitrogen, oxygen, or sulfur; or two occurrences of R', are taken together with the atom(s) to which they are bound to form an optionally substituted 3-12 membered saturated, partially unsaturated, or fully unsaturated monocyclic or bicyclic ring having 0-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur, wherein each occurrence of R' is optionally substituted with y occurrences of R^7. In some embodiments, R^5 is $N(R')_2$. In certain embodiments, for each of the subsets described above, R' is hydrogen, C_1-C_6alkyl optionally substituted with 1–3 occurrences of R^7, or is a 5-10-membered monocyclic or bicyclic saturated, partially unsaturated or fully unsaturated ring having 0-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur, wherein the ring is optionally substituted with 1-3 occurrences of R^7. . In other embodiments, R' is hydrogen, C_1-C_4alkyl optionally substituted with 1-3 occurrences of R^7, or is a ring selected from (i)-(xLvi) or (xLvii) described in [0059]. In yet other embodiments, R^5 is $-N(R')_2$ and the two occurrences of R' are taken together with the nitrogen atom to which they are bound to form an optionally substituted 3-10-

membered monocyclic or bicyclic heterocyclic ring. In certain embodiments, the ring is selected from (a)-(o) described in [0060].


[0072] In still other embodiments, R^1, R^2, R^3 and R^4 are each hydrogen and compounds of formula **I-A-iii** are provided:



I-A-iii

wherein x is 1, 2, or 3; and at least one occurence of R^5 is $-N(R')_2$, -$NR'CH(CH_2OH)R'$, -$NR'CH(CH_2CH_2OH)R'$, -$NR'(CH_2)R'$, -$NR'(CH_2)_2R'$, $NR'(CH_2)N(R')_2$,-$NR'(CH_2)_2N(R')_2$, -OR', -NR'COR', -$NR'COCH_2R'$, or -$NR'CO(CH_2)_2R'$; and R' is a C_1-C_6aliphatic group optionally substituted with y occurrences of R^7, or is a ring selected from (i)-(xLvi) or (xLvii) described in [0059]. In yet other embodiments, R^5 is $-N(R')_2$ and the two occurrences of R' are taken together with the nitrogen atom to which they are bound to form an optionally substituted 3-10-membered monocyclic or bicyclic heterocyclic ring. In certain embodiments, the ring is selected from (a)-(o) described in [0060].

[0073] In still other embodiments, R^1, R^2, R^3 and R^4 are each hydrogen, and x is 1, and compounds of formula **I-A-iv** are provided:




I-A-iv

wherein R^5 is $-N(R')_2$, -$NR'CH(CH_2OH)R'$, -$NR'CH(CH_2CH_2OH)R'$, -$NR'(CH_2)R'$, -$NR'(CH_2)_2R'$, $NR'(CH_2)N(R')_2$,-$NR'(CH_2)_2N(R')_2$, -OR', -NR'COR', -$NR'COCH_2R'$, or -$NR'CO(CH_2)_2R'$; and R' is a C_1-C_6aliphatic group optionally substituted with y occurrences of R^7 or is a ring selected from (i)-(xLvi) or (xLvii) described in [0059]. In yet other embodiments, R^5 is $-N(R')_2$ and the two occurrences of R' are taken together with the nitrogen atom to which they are bound to form an

optionally substituted 3-10-membered monocyclic or bicyclic heterocyclic ring. In certain embodiments, the ring is selected from (a)-(o) described in [0060].


[0074] In still other embodiments, R^1, R^2, R^3 and R^4 are each hydrogen, x is 1, and compounds of formula **I-A-v** are provided:



I-A-v

wherein R^5 is $-N(R')_2$, $-NR'CH(CH_2OH)R'$, $-NR'CH(CH_2CH_2OH)R'$, $-NR'(CH_2)R'$, $-NR'(CH_2)_2R'$, $NR'(CH_2)N(R')_2$, $-NR'(CH_2)_2N(R')_2$, $-OR'$, $-NR'COR'$, $-NR'COCH_2R'$, or $-NR'CO(CH_2)_2R'$; and R' is a C_1-C_6aliphatic group optionally substituted with y occurrences of R^7 or is a ring selected from (i)-(xLvi) or (xLvii) described in [0059]. In yet other embodiments, R^5 is $-N(R')_2$ and the two occurrences of R' are taken together with the nitrogen atom to which they are bound to form an optionally substituted 3-10-membered monocyclic or bicyclic heterocyclic ring. In certain embodiments, the ring is selected from (a)-(o) described in [0060].

[0075] In still other embodiments, R^1, R^2, and R^4 are each hydrogen, and compounds of formula **I-A-vi** are provided:



I-A-vi


wherein:

R^3 is an optionally substituted group selected from a 3-8-membered saturated, partially unsaturated, or fully unsaturated monocyclic ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur, or an 8-12 membered saturated, partially unsaturated, or fully unsaturated bicyclic ring system having 0-5 heteroatoms independently selected from nitrogen, oxygen, or sulfur;

x is 1, 2, or 3; and at least one occurence of R^5 is $-N(R')_2$, $-NR'CH(CH_2OH)R'$, $-NR'CH(CH_2CH_2OH)R'$, $-NR'(CH_2)R'$, $-NR'(CH_2)_2R'$, $NR'(CH_2)N(R')_2$, $-NR'(CH_2)_2N(R')_2$, $-OR'$, $-NR'COR'$, $-NR'COCH_2R'$, or $-NR'CO(CH_2)_2R'$; and R' is a C_1-C_6aliphatic group optionally substituted with y occurrences of R^7, or is a ring selected from (i)-(xLvi) or (xLvii) described in [0059]. In yet other embodiments, R^5 is $-N(R')_2$ and the two occurrences of R' are taken together with the nitrogen atom to which they are bound to form an optionally substituted 3-10-membered monocyclic or bicyclic heterocyclic ring. In certain embodiments, the ring is selected from (a)-(o) described in [0060].

[0076] In yet other embodiments for compounds described directly above, R^3 is an optionally substituted 5- or 6-membered saturated, partially unsaturated, or fully unsaturated ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur. In yet other embodiments, R^3 is an optionally substituted ring selected from phenyl, pyridyl, pyrimidinyl, thiazolyl, oxazolyl, thienyl, furyl, pyrrolyl, pyrazolyl, triazolyl, pyrazinyl, thiadiazolyl, or oxadiazolyl. As described generally above, R^3 is optionally substituted with z occurrences of R^6. In certain embodiments, wherein z is 0, 1, 2, or 3, and each occurrence of R^6 is independently hydrogen, R", $-CH_2R"$, halogen, CN, NO_2, $-N(R")_2$, $-CH_2N(R")_2$, -OR", $-CH_2OR"$, -SR", $-CH_2SR"$, -COOR", -NR"COR", -NR"COOR", $-CON(R")_2$, $-SO_2N(R")_2$, $-CONR"(CH_2)_2N(R")_2$, $-CONR(CH_2)_3N(R")_2$, $-CONR"(CH_2)_4N(R")_2$, $-O(CH_2)_2OR"$, $O(CH_2)_3OR"$, $O(CH_2)_4OR"$, $-O(CH_2)_2N(R")_2$, $-O(CH_2)_3N(R")_2$, $-O(CH_2)_4N(R")_2$, $-NR"CH(CH_2OH)R"$, $-NR"CH(CH_2CH_2OH)R"$, $-NR"(CH_2)R"$, $-NR"(CH_2)_2R"$, $-NR"(CH_2)_3R"$, $-NR"(CH_2)_4R"$, $-NR"(CH_2)N(R")_2$, $-NR"(CH_2)_2N(R")_2$, $-NR"(CH_2)_3N(R")_2$, $-NR"(CH_2)_4N(R")_2$, $-NR"(CH_2)OR"$, $-NR"(CH_2)_2OR"$, $-NR"(CH_2)_3OR"$, or $-NR'(CH_2)_4OR"$.

[0077] In still other embodiments, R^1, R^2, R^3 and R^4 are each hydrogen, and x is 1, and compounds of formula **I-A-vii** are provided:




I-A-vii

wherein:

R^3 is an optionally substituted group selected from a 3-8-membered saturated, partially unsaturated, or fully unsaturated monocyclic ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur, or an 8-12 membered saturated, partially unsaturated, or fully unsaturated bicyclic ring system having 0-5 heteroatoms independently selected from nitrogen, oxygen, or sulfur;
R^5 is $-N(R')_2$, $-NR'CH(CH_2OH)R'$, $-NR'CH(CH_2CH_2OH)R'$, $-NR'(CH_2)R'$, $-NR'(CH_2)_2R'$, $NR'(CH_2)N(R')_2$, $-NR'(CH_2)_2N(R')_2$, $-OR'$, $-NR'COR'$, $-NR'COCH_2R'$, or $-NR'CO(CH_2)_2R'$; and R' is a C_1-C_6aliphatic group optionally substituted with y occurrences of R^7, or is a ring selected from (i)-(xLvi) or (xLvii) described in [0059]. In yet other embodiments, R^5 is $-N(R')_2$ and the two occurrences of R' are taken together with the nitrogen atom to which they are bound to form an optionally substituted 3-10-membered monocyclic or bicyclic heterocyclic ring. In certain embodiments, the ring is selected from (a)-(o) described in [0060].

[0078] In yet other embodiments for compounds described directly above, R^3 is an optionally substituted 5- or 6-membered saturated, partially unsaturated, or fully unsaturated ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur. In yet other embodiments, R^3 is an optionally substituted ring selected from phenyl, pyridyl, pyrimidinyl, thiazolyl, oxazolyl, thienyl, furyl, pyrrolyl, pyrazolyl, triazolyl, pyrazinyl, thiadiazolyl, or oxadiazolyl. As described generally above, R^3 is optionally substituted with z occurrences of R^6. In certain embodiments, wherein z is 0, 1, 2, or 3, and each occurrence of R^6 is independently hydrogen, R", $-CH_2R"$, halogen, CN, NO_2, $-N(R")_2$, $-CH_2N(R")_2$, $-OR"$, $-CH_2OR"$, $-SR"$, $-CH_2SR"$, $-COOR"$, $-NR"COR"$, $-NR"COOR"$, $-CON(R")_2$, $-SO_2N(R")_2$, $-CONR"(CH_2)_2N(R")_2$, $-CONR(CH_2)_3N(R")_2$, $-CONR"(CH_2)_4N(R")_2$, $-O(CH_2)_2OR"$, $O(CH_2)_3OR"$, $O(CH_2)_4OR"$, $-O(CH_2)_2N(R")_2$, $-O(CH_2)_3N(R")_2$, $-O(CH_2)_4N(R")_2$, $-NR"CH(CH_2OH)R"$, $-NR"CH(CH_2CH_2OH)R"$, $-NR"(CH_2)R"$, $-NR"(CH_2)_2R"$, $-NR"(CH_2)_3R"$, $-NR"(CH_2)_4R"$, $-NR"(CH_2)N(R")_2$, $-NR"(CH_2)_2N(R")_2$, $-NR"(CH_2)_3N(R")_2$, $-NR"(CH_2)_4N(R")_2$, $-NR"(CH_2)OR"$, $-NR"(CH_2)_2OR"$, $-NR"(CH_2)_3OR"$, or $-NR'(CH_2)_4OR"$.

[0079] In still other embodiments, R^1, R^2, R^3 and R^4 are each hydrogen, x is 1, and compounds of formula **I-A-viii** are provided:



I-A-viii

wherein:


R^3 is an optionally substituted group selected from a 3-8-membered saturated, partially unsaturated, or fully unsaturated monocyclic ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur, or an 8-12 membered saturated, partially unsaturated, or fully unsaturated bicyclic ring system having 0-5 heteroatoms independently selected from nitrogen, oxygen, or sulfur; R^5 is $-N(R')_2$, -$NR'CH(CH_2OH)R'$, -$NR'CH(CH_2CH_2OH)R'$, -$NR'(CH_2)R'$, -$NR'(CH_2)_2R'$, $NR'(CH_2)N(R')_2$, -$NR'(CH_2)_2N(R')_2$, -OR', -NR'COR', -NR'COCH$_2$R', or -$NR'CO(CH_2)_2R'$; and R' is a C_1-C_6aliphatic group optionally substituted with y occurrences of R^7, or is a ring selected from (i)-(xLvi) or (xLvii) described in [0059]. In yet other embodiments, R^5 is -$N(R')_2$ and the two occurrences of R' are taken together with the nitrogen atom to which they are bound to form an optionally substituted 3-10-membered monocyclic or bicyclic heterocyclic ring. In certain embodiments, the ring is selected from (a)-(o) described in [0060].

[0080] In yet other embodiments for compounds described directly above, R^3 is an optionally substituted 5- or 6-membered saturated, partially unsaturated, or fully unsaturated ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur. In yet other embodiments, R^3 is an optionally substituted ring selected from phenyl, pyridyl, pyrimidinyl, thiazolyl, oxazolyl, thienyl, furyl, pyrrolyl, pyrazolyl, triazolyl, pyrazinyl, thiadiazolyl, or oxadiazolyl. As described generally above, R^3 is optionally substituted with z occurrences of R^6. In certain embodiments, wherein z is 0, 1, 2, or 3, and each occurrence of R^6 is independently hydrogen, R", -CH_2R", halogen, CN, NO_2, -$N(R")_2$, -$CH_2N(R")_2$, -OR", -CH_2OR", -SR", -CH_2SR", -COOR", -NR"COR", -NR"COOR", -$CON(R")_2$, -$SO_2N(R")_2$, -

$CONR''(CH_2)_2N(R'')_2$, $-CONR(CH_2)_3N(R'')_2$, $-CONR''(CH_2)_4N(R'')_2$, $-O(CH_2)_2OR''$, $O(CH_2)_3OR''$, $O(CH_2)_4OR''$, $-O(CH_2)_2N(R'')_2$, $-O(CH_2)_3N(R'')_2$, $-O(CH_2)_4N(R'')_2$, $-NR''CH(CH_2OH)R''$, $-NR''CH(CH_2CH_2OH)R''$, $-NR''(CH_2)R''$, $-NR''(CH_2)_2R''$, $-NR''(CH_2)_3R''$, $-NR''(CH_2)_4R''$, $-NR''(CH_2)N(R'')_2$, $-NR''(CH_2)_2N(R'')_2$, $-NR''(CH_2)_3N(R'')_2$, $-NR''(CH_2)_4N(R'')_2$, $-NR''(CH_2)OR''$, $-NR''(CH_2)_2OR''$, $-NR''(CH_2)_3OR''$, or $-NR'(CH_2)_4OR''$.

[0081] Certain additional subsets of compounds of general formula I include:

[0082] **II. Compounds of formula I-C:**



I-C

wherein R^1, R^2, R^3, R^4, R^5 and x are each described generally above and in subsets described above and herein.

[0083] In some embodiments, for compounds of formula I-C:

a. R^1 is:

   i. T-R', wherein T is a bond or an optionally substituted C_1-C_6alkylidene chain wherein up to two methylene units are optionally and independently replaced with –O-, -S-, -NR'-, -OCO-, -COO-, $-SO_2-$ or –CO-, and R' is hydrogen, C_1-C_4-alkyl, or an optionally substituted 5- or 6-membered aryl or heteroaryl group, or

   ii. $-Si(R')_3$, wherein R' is hydrogen, C_1-C_4-alkyl, or an optionally substituted 5- or 6-membered saturated, partially unsaturated, or fully unsaturated monocyclic ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur;

b. R^2, R^3, and R^4 are each independently hydrogen, R', halogen, CN, NO_2, $-N(R')_2$, $-CH_2N(R')_2$, -OR', $-CH_2OR'$, -SR', $-CH_2SR'$, -COOR', -NR'COR', $-CON(R')_2$, $-SO_2N(R')_2$, $-CONR'(CH_2)_2N(R')_2$, $-CONR'(CH_2)_3N(R')_2$, $-CONR'(CH_2)_4N(R')_2$, $-O(CH_2)_2OR'$,

$O(CH_2)_3OR'$, $O(CH_2)_4OR'$, $-O(CH_2)_2N(R')_2$, $-O(CH_2)_3N(R')_2$, or $-O(CH_2)_4N(R')_2$; wherein R^2, R^3, and R^4 are each optionally substituted with z occurrences of R^6, wherein z is 0-5 and R^6 is =O, =NR", =S, halogen, -CN, $-NO_2$, or Z-R", wherein Z is a bond or an optionally substituted C_1-C_6 alkylidene chain, wherein up to two methylene units of the chain are optionally and independently replaced by —NR"-, -S-, -O-, -CS-, $-CO_2-$, -OCO-, -CO-, -COCO-, -CONR"-, -NR"CO-, -NR"CO_2-, $-SO_2NR"-$, -NR"SO_2-, -CONR"NR"-, -NR"CONR"-, -OCONR"-, -NR"NR"-, -NR"SO_2NR"-, -SO-, $-SO_2-$, -PO-, $-PO_2-$, or –POR"-, and each occurrence of R" is independently hydrogen or an optionally substituted C_1-C_6 aliphatic group, a 3-8-membered saturated, partially unsaturated, or fully unsaturated monocyclic ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur, or an 8-12 membered saturated, partially unsaturated, or fully unsaturated bicyclic ring system having 0-5 heteroatoms independently selected from nitrogen, oxygen, or sulfur; or two occurrences of R" are taken together with the atom(s) to which they are bound to form an optionally substituted 3-12 membered saturated, partially unsaturated, or fully unsaturated monocyclic or bicyclic ring having 0-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur; and

c. each occurrence of R^5 is independently hydrogen, halogen, R', CN, $-CH_2CN$, $-(CH_2)_2CN$, NO_2, $-CH_2NO_2$, $-(CH_2)_2NO_2$, $CON(R')_2$, $-CH_2CON(R')_2$, $-(CH_2)_2CON(R')_2$, COOR', $-CH_2COOR'$, $-(CH_2)_2COOR'$, $-SO_2N(R')_2$, $-CH_2SO_2N(R')_2$, $-(CH_2)_2SO_2N(R')_2$, $-NR'SO_2R'$, $-CH_2NR'SO_2R'$, $-(CH_2)_2NR'SO_2R'$, $NR'CON(R')_2$, $-CH_2NR'CON(R')_2$, $-(CH_2)_2NR'CON(R')_2$, $-NR'SO_2N(R')_2$, $-CH_2NR'SO_2N(R')_2$, $-(CH_2)_2NR'SO_2N(R')_2$, -COCOR', $-CH_2COCOR'$, $-(CH_2)_2COCOR'$, $-N(R')_2$, $-CH_2N(R')_2$, $-(CH_2)_2N(R')_2$, -OR', $-CH_2OR'$, $-(CH_2)_2OR'$, -NR'COR', $-NR'COCH_2R'$, $-NR'CO(CH_2)_2R'$,; $-CH_2NR'COR'$, or $-(CH_2)_2NR'COR'$, wherein R^5 is optionally substituted with y occurrences of R^7, wherein y is 0-5 and R^7 is =O,

=NR", =S, halogen, -CN, -NO_2, or W-R', wherein W is a bond or an optionally substituted C_1-C_6 alkylidene chain, wherein up to two methylene units of the chain are optionally and independently replaced by –NR"-, -S-, -O-, -CS-, -CO_2-, -OCO-, -CO-, -COCO-, -CONR"-, -NR"CO-, -NR"CO_2-, -SO_2NR"-, -NR"SO_2-, -CONR"NR"-, -NR"CONR"-, -OCONR"-, -NR"NR"-, -NR"SO_2NR"-, -SO-, -SO_2-, -PO-, -PO_2-, or –POR"-, and and each occurrence of R" is independently hydrogen or an optionally substituted C_1-C_6 aliphatic group, a 3-8-membered saturated, partially unsaturated, or fully unsaturated monocyclic ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur, or an 8-12 membered saturated, partially unsaturated, or fully unsaturated bicyclic ring system having 0-5 heteroatoms independently selected from nitrogen, oxygen, or sulfur; or two occurrences of R" are taken together with the atom(s) to which they are bound to form an optionally substituted 3-12 membered saturated, partially unsaturated, or fully unsaturated monocyclic or bicyclic ring having 0-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur.

[0084] In still other embodiments, for compounds of formula **I-C**:

a. R^1 is hydrogen, C_1-C_4alkyl, -COR', -SO_2R', or –Si(R')$_3$;

b. R^2, R^3, and R^4 are each independently Cl, Br, F, -CN, -COOH, -COOMe, -NH_2, –N(CH_3)$_2$, -N(Et)$_2$, -N(iPr)$_2$, -O(CH_2)$_2$$OCH_3$, -$CONH_2$, -$COOCH_3$, -OH, -$CH_2$OH, -$NHCOCH_3$, -$SO_2NH_2$, -$SO_2$N(Me)$_2$, or an optionally substituted group selected from C_1-C_4alkyl, C_1-C_4alkyloxy, a 3-8-membered saturated, partially unsaturated, or fully unsaturated monocyclic ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur, or an 8-12 membered saturated, partially unsaturated, or fully unsaturated bicyclic ring system having 0-5 heteroatoms independently selected from nitrogen, oxygen, or sulfur; wherein R^2, R^3, and R^4 are each independently and optionally substituted with z occurrences of R^6, wherein z is 0, 1, 2, or 3, and each occurrence of R^6 is

independently hydrogen, R", -CH_2R", halogen, CN, NO_2, -N(R")$_2$, -CH_2N(R")$_2$, -OR", -CH_2OR", -SR", -CH_2SR", -COOR", -NR"COR", -NR"COOR", -CON(R")$_2$, -SO_2N(R")$_2$, -CONR"$(CH_2)_2$N(R")$_2$, -CONR$(CH_2)_3$N(R")$_2$, -CONR"$(CH_2)_4$N(R")$_2$, -O$(CH_2)_2$OR", O$(CH_2)_3$OR", O$(CH_2)_4$OR", -O$(CH_2)_2$N(R")$_2$, -O$(CH_2)_3$N(R")$_2$, -O$(CH_2)_4$N(R")$_2$, -NR"CH(CH_2OH)R", -NR"CH(CH_2CH_2OH)R", -NR"(CH_2)R", -NR"$(CH_2)_2$R", -NR"$(CH_2)_3$R", -NR"$(CH_2)_4$R", -NR"(CH_2)N(R")$_2$, -NR"$(CH_2)_2$N(R")$_2$, -NR"$(CH_2)_3$N(R")$_2$, -NR"$(CH_2)_4$N(R")$_2$, -NR"(CH_2)OR", -NR"$(CH_2)_2$OR", -NR"$(CH_2)_3$OR", or –NR'$(CH_2)_4$OR"; each occurrence of R^5 is independently CN, -CH_2CN, -$(CH_2)_2$CN, -NO_2, -CH_2NO_2, -$(CH_2)_2NO_2$, OR', -CH_2OR', -CON(R')$_2$, -SO_2N(R')$_2$, -N(R')$_2$, or R', wherein R^5 is optionally substituted y occurrences of R^7, wherein y is 0, 1, 2, or 3, and each occurrence of R^7 is independently hydrogen, R", -CH_2R", halogen, CN, NO_2, -N(R")$_2$, -CH_2N(R")$_2$, -OR", -CH_2OR", -SR", -CH_2SR", -COOR", -NR"COR", -NR"COOR", -CON(R")$_2$, -SO_2N(R")$_2$, -CONR"$(CH_2)_2$N(R")$_2$, -CONR$(CH_2)_3$N(R")$_2$, -CONR"$(CH_2)_4$N(R")$_2$, -O$(CH_2)_2$OR", O$(CH_2)_3$OR", O$(CH_2)_4$OR", -O$(CH_2)_2$N(R")$_2$, -O$(CH_2)_3$N(R")$_2$, -O$(CH_2)_4$N(R")$_2$, -NR"CH(CH_2OH)R", -NR"CH(CH_2CH_2OH)R", -NR"(CH_2)R", -NR"$(CH_2)_2$R", -NR"$(CH_2)_3$R", -NR"$(CH_2)_4$R", -NR"(CH_2)N(R")$_2$, -NR"$(CH_2)_2$N(R")$_2$, -NR"$(CH_2)_3$N(R")$_2$, -NR"$(CH_2)_4$N(R")$_2$, -NR"(CH_2)OR", -NR"$(CH_2)_2$OR", -NR"$(CH_2)_3$OR", or –NR"$(CH_2)_4$OR".

[0085] In other embodiments, for compounds of formula **I-C** and subsets described directly above, R^1 is hydrogen, methyl, ethyl, n-propyl, isopropyl, n-butyl, p-toluenesulfonyl (Ts), t-butyldimethylsilyl (TBS), triisopropylsilyl (TIPS), or triethylsilyl (TES).

[0086] In still other embodiments, for compounds of formula **I-C** and subsets described directly above, R^2, R^3, and R^4 are each hydrogen. In other embodiments, one of R^2, R^3, or R^4 is hydrogen. In yet other embodiments, two of R^2, R^3, or R^4 is hydrogen. In yet other embodiments, R^2 and R^4 are both hydrogen, and R^3 is halogen,

CN, NO_2, or V-R'. In still other embodiments, R^2 and R^4 are both hydrogen, and R^3 is an optionally substituted group selected from a 3-8-membered saturated, partially unsaturated, or fully unsaturated monocyclic ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur, or an 8-12 membered saturated, partially unsaturated, or fully unsaturated bicyclic ring system having 0-5 heteroatoms independently selected from nitrogen, oxygen, or sulfur. In yet other embodiments, R^2 and R^4 are both hydrogen, and R^3 is an optionally substituted 5- or 6-membered saturated, partially unsaturated, or fully unsaturated ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur. In yet other embodiments, R^2 and R^4 are both hydrogen, and R^3 is an optionally substituted ring selected from phenyl, pyridyl, pyrimidinyl, thiazolyl, oxazolyl, thienyl, furyl, pyrrolyl, pyrazolyl, triazolyl, pyrazinyl, thiadiazolyl, or oxadiazolyl.


[0087] In other embodiments, for compounds of formula **I-C** and subsets described directly above, R^2, R^3, and R^4 are each independently and optionally substituted with z occurrences of R^6, wherein z is 1, 2, or 3 and each occurrence of R^6 is independently F, Cl, Br, CN, OH, NH_2, $-CH_2OH$, C_1-C_6alkyl, $-O(C_1$-C_6alkyl), $-CH_2O(C_1$-C_6alkyl), $-CO(C_1$-C_6alkyl), $-COO(C_1$-C_6alkyl), $-NHSO_2(C_1$-C_6alkyl), $-SO_2NH_2$, $-CONH_2$, $-CON(C_1$-C_6alkyl), $-SO_2(C_1$-C_6alkyl), $-SO_2$phenyl, phenyl, benzyl, $-N(C_1$-C_6alkyl$)_2$, or $-S(C_1$-C_6alkyl), wherein each of the foregoing phenyl, benzyl, and C_1-C_6alkyl groups is independently and optionally substituted, and wherein each of the foregoing C_1-C_6alkyl groups is linear, branched, or cyclic.

[0088] In certain exemplary embodiments, for compounds of formula **I-C** and subsets described directly above, x is 1, 2, or 3, and at least one occurrence of R^5 is halogen, CN, $-CH_2CN$, $-(CH_2)_2CN$, NO_2, $-CH_2NO_2$, $-(CH_2)_2NO_2$, $-CONH_2$, $-CON(C_1$-C_4alkyl), $-SO_2NH_2$, $-SO_2N(C_1$-C_4alkyl), NH_2, $-N(C_1$-C_4alkyl), -OH, $-O(C_1$-C_4alkyl), $-CH_2OH$, $-CH_2O(C_1$-C_4alkyl), or an optionally substituted 5- or 6-membered unsaturated ring wherein 0-3 ring carbon atoms is optionally replaced by oxygen, sulfur, or nitrogen. In other embodiments, x is 1, 2, or 3, and at least one occurrence of R^5 is -OR'.

[0089] In still other embodiments, for compounds of formula **I-C** and subsets described directly above, R^5 is optionally substituted with y occurrences of R^7, wherein y is 1, 2, or 3 and each occurrence of R^7 is independently F, Cl, Br, CN, OH,

NH_2, $-CH_2OH$, C_1-C_6alkyl, $-O(C_1$-C_6alkyl), $-CH_2O(C_1$-C_6alkyl), $-CO(C_1$-C_6alkyl), $-COO(C_1$-C_6alkyl), $-NHSO_2(C_1$-C_6alkyl), $-SO_2NH_2$, $-CONH_2$, $-CON(C_1$-C_6alkyl), $-SO_2(C_1$-C_6alkyl), $-SO_2$phenyl, phenyl, benzyl, $-N(C_1$-C_6alkyl$)_2$, or $-S(C_1$-C_6alkyl), wherein each of the foregoing phenyl, benzyl, and C_1-C_6alkyl groups is independently and optionally substituted, and wherein each of the foregoing C_1-C_6alkyl groups is linear, branched, or cyclic.

[0090] In yet other embodiments, x is 1 and compounds have general formula **I-C-i**:




I-C-i

wherein R^1, R^2, R^3, and R^4 are described generally and in subsets above and herein, and R^5 is halogen, R', CN, $-CH_2CN$, $-(CH_2)_2CN$, NO_2, $-CH_2NO_2$, $-(CH_2)_2NO_2$, $CON(R')_2$, $-CH_2CON(R')_2$, $-(CH_2)_2CON(R')_2$, COOR', $-CH_2COOR'$, $-(CH_2)_2COOR'$, $-SO_2N(R')_2$, $-CH_2SO_2N(R')_2$, $-(CH_2)_2SO_2N(R')_2$, $-NR'SO_2R'$, $-CH_2NR'SO_2R'$, $-(CH_2)_2NR'SO_2R'$, $NR'CON(R')_2$, $-CH_2NR'CON(R')_2$, $-(CH_2)_2NR'CON(R')_2$, $-NR'SO_2N(R')_2$, $-CH_2NR'SO_2N(R')_2$, $-(CH_2)_2NR'SO_2N(R')_2$, -COCOR', $-CH_2COCOR'$, $-(CH_2)_2COCOR'$,$-N(R')_2$, $-CH_2N(R')_2$, $-(CH_2)_2N(R')_2$, -OR', $-CH_2OR'$, $-(CH_2)_2OR'$, $-NR'COR'$, $-CH_2NR'COR'$, or $-(CH_2)_2NR'COR'$, and R' is a C_1-C_6aliphatic group or a 3-8-membered saturated, partially unsaturated, or fully unsaturated monocyclic ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur, or an 8-12 membered saturated, partially unsaturated, or fully unsaturated bicyclic ring system having 0-5 heteroatoms independently selected from nitrogen, oxygen, or sulfur; or two occurrences of R', are taken together with the atom(s) to which they are bound to form an optionally substituted 3-12 membered saturated, partially unsaturated, or fully unsaturated monocyclic or bicyclic ring having 0-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur, wherein each occurrence of R' is optionally substituted with y occurrences of R^7. In some embodiments, R^5 is CN, $-CH_2CN$, $-(CH_2)_2CN$, -

NO_2, $-CH_2NO_2$, $-(CH_2)_2NO_2$, OR', $-CH_2OR'$, $-CON(R')_2$, $-SO_2N(R')_2$, $-N(R')_2$, or R'. In certain embodiments, R' is hydrogen, C_1-C_6alkyl optionally substituted with 1-3 occurrences of R^7, or is a 5-10-membered monocyclic or bicyclic saturated, partially unsaturated or fully unsaturated ring having 0-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur, wherein the ring is optionally substituted with 1-3 occurrences of R^7. In other embodiments, R' is hydrogen, C_1-C_4alkyl optionally substituted with 1-3 occurrences of R^7, or is a ring selected from (i)-(xLvi) or (xLvii) described in [0059].

[0091] In yet other embodiments, x is 1 and compounds have general formula **I-C-ii:**




I-C-ii

wherein R^1, R^2, R^3, and R^4 are described generally and in subsets above and herein, and R^5 is halogen, R', CN, $-CH_2CN$, $-(CH_2)_2CN$, NO_2, $-CH_2NO_2$, $-(CH_2)_2NO_2$, $CON(R')_2$, $-CH_2CON(R')_2$, $-(CH_2)_2CON(R')_2$, COOR', $-CH_2COOR'$, $-(CH_2)_2COOR'$, $-SO_2N(R')_2$, $-CH_2SO_2N(R')_2$, $-(CH_2)_2SO_2N(R')_2$, $-NR'SO_2R'$, $-CH_2NR'SO_2R'$, $-(CH_2)_2NR'SO_2R'$, $NR'CON(R')_2$, $-CH_2NR'CON(R')_2$, $-(CH_2)_2NR'CON(R')_2$, $-NR'SO_2N(R')_2$, $-CH_2NR'SO_2N(R')_2$, $-(CH_2)_2NR'SO_2N(R')_2$, -COCOR', $-CH_2COCOR'$, $-(CH_2)_2COCOR'$,$-N(R')_2$, $-CH_2N(R')_2$, $-(CH_2)_2N(R')_2$, -OR', $-CH_2OR'$, $-(CH_2)_2OR'$, $-NR'COR'$, $-CH_2NR'COR'$, or $-(CH_2)_2NR'COR'$, and R' is a C_1-C_6aliphatic group or a 3-8-membered saturated, partially unsaturated, or fully unsaturated monocyclic ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur, or an 8-12 membered saturated, partially unsaturated, or fully unsaturated bicyclic ring system having 0-5 heteroatoms independently selected from nitrogen, oxygen, or sulfur; or two occurrences of R', are taken together with the atom(s) to which they are bound to form an optionally substituted 3-12 membered saturated, partially unsaturated, or fully unsaturated

monocyclic or bicyclic ring having 0-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur, wherein each occurrence of R' is optionally substituted with y occurrences of R^7. In some embodiments, R^5 is CN, $-CH_2CN$, $-(CH_2)_2CN$, $-NO_2$, $-CH_2NO_2$, $-(CH_2)_2NO_2$, OR', $-CH_2OR'$, $-CON(R')_2$, $-SO_2N(R')_2$, $-N(R')_2$, or R'. In certain embodiments, R' is hydrogen, C_1-C_6alkyl optionally substituted with 1-3 occurrences of R^7, or is a 5-10-membered monocyclic or bicyclic saturated, partially unsaturated or fully unsaturated ring having 0-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur, wherein the ring is optionally substituted with 1-3 occurrences of R^7. In other embodiments, R' is hydrogen, C_1-C_4alkyl optionally substituted with 1-3 occurrences of R^7, or is a ring selected from (i)-(xLvi) or (xLvii) described in [0059].

[0092] In still other embodiments, each occurrence of R^5 is independently CN, $-CH_2CN$, $-(CH_2)_2CN$, $-NO_2$, $-CH_2NO_2$, $-(CH_2)_2NO_2$, OR', $-CH_2OR'$, $-CON(R')_2$, $-SO_2N(R')_2$, $-N(R')_2$, or R'. In still other embodiments, each occurrence of R^5 is independently hydrogen, halogen, CN, $-CH_2CN$, $-(CH_2)_2CN$, NO_2, $-CH_2NO_2$, $-(CH_2)_2NO_2$, $-CONH_2$, $-CON(C_1$-C_4alkyl), $-SO_2NH_2$, $-SO_2N(C_1$-C_4alkyl), NH_2, $-N(C_1$-C_4alkyl), -OH, $-O(C_1$-C_4alkyl), $-CH_2OH$, $-CH_2O(C_1$-C_4alkyl), or an optionally substituted 5- or 6-membered unsaturated ring wherein 0-3 ring carbon atoms is optionally replaced by oxygen, sulfur, or nitrogen, wherein R^5 is optionally substituted by 0-3 occurrences of R^7.

[0093] In yet other embodiments, x is 2 and compounds have general formula **I-C-iii:**




I-C-iii

wherein R^1, R^2, R^3, and R^4 are described generally and in subsets above and herein, and each occurrence of R^5 is independently halogen, R', CN, $-CH_2CN$, $-(CH_2)_2CN$, NO_2, $-CH_2NO_2$, $-(CH_2)_2NO_2$, $CON(R')_2$, $-CH_2CON(R')_2$, $-(CH_2)_2CON(R')_2$, COOR',

$-CH_2COOR'$, $-(CH_2)_2COOR'$, $-SO_2N(R')_2$, $-CH_2SO_2N(R')_2$, $-(CH_2)_2SO_2N(R')_2$, $-NR'SO_2R'$, $-CH_2NR'SO_2R'$, $-(CH_2)_2NR'SO_2R'$, $NR'CON(R')_2$, $-CH_2NR'CON(R')_2$, $-(CH_2)_2NR'CON(R')_2$, $-NR'SO_2N(R')_2$, $-CH_2NR'SO_2N(R')_2$, $-(CH_2)_2NR'SO_2N(R')_2$, $-COCOR'$, $-CH_2COCOR'$, $-(CH_2)_2COCOR'$, $-N(R')_2$, $-CH_2N(R')_2$, $-(CH_2)_2N(R')_2$, $-OR'$, $-CH_2OR'$, $-(CH_2)_2OR'$, $-NR'COR'$ $-CH_2NR'COR'$, or $-(CH_2)_2NR'COR'$, and R' is a C_1-C_6aliphatic group or a 3-8-membered saturated, partially unsaturated, or fully unsaturated monocyclic ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur, or an 8-12 membered saturated, partially unsaturated, or fully unsaturated bicyclic ring system having 0-5 heteroatoms independently selected from nitrogen, oxygen, or sulfur; or two occurrences of R', are taken together with the atom(s) to which they are bound to form an optionally substituted 3-12 membered saturated, partially unsaturated, or fully unsaturated monocyclic or bicyclic ring having 0-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur, wherein each occurrence of R' is optionally substituted with y occurrences of R^7. In some embodiments, R^5 is CN, $-CH_2CN$, $-(CH_2)_2CN$, $-NO_2$, $-CH_2NO_2$, $-(CH_2)_2NO_2$, OR', $-CH_2OR'$, $-CON(R')_2$, $-SO_2N(R')_2$, $-N(R')_2$, or R'. In certain embodiments, R' is hydrogen, C_1-C_6alkyl optionally substituted with 1-3 occurrences of R^7, or is a 5-10-membered monocyclic or bicyclic saturated, partially unsaturated or fully unsaturated ring having 0-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur, wherein the ring is optionally substituted with 1-3 occurrences of R^7. In other embodiments, R' is hydrogen, C_1-C_4alkyl optionally substituted with 1-3 occurrences of R^7, or is a ring selected from (i)-(xLvi) or (xLvii) described in [0059].

[0094] In still other embodiments, each occurrence of R^5 is independently CN, $-CH_2CN$, $-(CH_2)_2CN$, $-NO_2$, $-CH_2NO_2$, $-(CH_2)_2NO_2$, OR', $-CH_2OR'$, $-CON(R')_2$, $-SO_2N(R')_2$, $-N(R')_2$, or R'. In still other embodiments, each occurrence of R^5 is independently hydrogen, halogen, CN, $-CH_2CN$, $-(CH_2)_2CN$, NO_2, $-CH_2NO_2$, $-(CH_2)_2NO_2$, $-CONH_2$, $-CON(C_1$-C_4alkyl), $-SO_2NH_2$, $-SO_2N(C_1$-C_4alkyl), NH_2, $-N(C_1$-C_4alkyl), -OH, $-O(C_1$-C_4alkyl), $-CH_2OH$, $-CH_2O(C_1$-C_4alkyl), or an optionally substituted 5- or 6-membered unsaturated ring wherein 0-3 ring carbon atoms is optionally replaced by oxygen, sulfur, or nitrogen, wherein R^5 is optionally substituted by 0-3 occurrences of R^7.

[0095] In still other embodiments, R^1, R^2, R^3 and R^4 are each hydrogen, and x is 1, and compounds of formula **I-C-iv** are provided:




I-C-iv

wherein R^5 is halogen, R', CN, $-CH_2CN$, $-(CH_2)_2CN$, NO_2, $-CH_2NO_2$, $-(CH_2)_2NO_2$, $CON(R')_2$, $-CH_2CON(R')_2$, $-(CH_2)_2CON(R')_2$, COOR', $-CH_2COOR'$, $-(CH_2)_2COOR'$, $-SO_2N(R')_2$, $-CH_2SO_2N(R')_2$, $-(CH_2)_2SO_2N(R')_2$, $-NR'SO_2R'$, $-CH_2NR'SO_2R'$, $-(CH_2)_2NR'SO_2R'$, $NR'CON(R')_2$, $-CH_2NR'CON(R')_2$, $-(CH_2)_2NR'CON(R')_2$, $-NR'SO_2N(R')_2$, $-CH_2NR'SO_2N(R')_2$, $-(CH_2)_2NR'SO_2N(R')_2$, -COCOR', $-CH_2COCOR'$, $-(CH_2)_2COCOR'$,–$N(R')_2$, $-CH_2N(R')_2$, $-(CH_2)_2N(R')_2$, -OR', $-CH_2OR'$, $-(CH_2)_2OR'$, $-NR'COR'$, $-CH_2NR'COR'$, or $-(CH_2)_2NR'COR'$, and R' is a C_1-C_6aliphatic group or a 3-8-membered saturated, partially unsaturated, or fully unsaturated monocyclic ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur, or an 8-12 membered saturated, partially unsaturated, or fully unsaturated bicyclic ring system having 0-5 heteroatoms independently selected from nitrogen, oxygen, or sulfur; or two occurrences of R', are taken together with the atom(s) to which they are bound to form an optionally substituted 3-12 membered saturated, partially unsaturated, or fully unsaturated monocyclic or bicyclic ring having 0-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur, wherein each occurrence of R' is optionally substituted with y occurrences of R^7. In some embodiments, R^5 is CN, $-CH_2CN$, $-(CH_2)_2CN$, $-NO_2$, $-CH_2NO_2$, $-(CH_2)_2NO_2$, OR', $-CH_2OR'$, $-CON(R')_2$, $-SO_2N(R')_2$, $-N(R')_2$, or R'. In certain embodiments, R' is hydrogen, C_1-C_6alkyl optionally substituted with 1-3 occurrences of R^7, or is a 5-10-membered monocyclic or bicyclic saturated, partially unsaturated or fully unsaturated ring having 0-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur, wherein the ring is optionally substituted with 1-3 occurrences of R^7. In other embodiments, R' is hydrogen, C_1-C_4alkyl optionally substituted with 1-3 occurrences of R^7, or is a ring selected from (i)-(xLvi) or (xLvii) described in [0059].

[0096] In still other embodiments, each occurrence of R^5 is independently CN, -CH_2CN, -$(CH_2)_2CN$, -NO_2, -CH_2NO_2, -$(CH_2)_2NO_2$, OR', -CH_2OR', -$CON(R')_2$, -$SO_2N(R')_2$, -$N(R')_2$, or R'. In still other embodiments, each occurrence of R^5 is independently hydrogen, halogen, CN, -CH_2CN, -$(CH_2)_2CN$, NO_2, -CH_2NO_2, -$(CH_2)_2NO_2$, -$CONH_2$, -$CON(C_1$-C_4alkyl), -SO_2NH_2, -$SO_2N(C_1$-C_4alkyl), NH_2, -$N(C_1$-C_4alkyl), -OH, -$O(C_1$-C_4alkyl), -CH_2OH, -$CH_2O(C_1$-C_4alkyl), or an optionally substituted 5- or 6-membered unsaturated ring wherein 0-3 ring carbon atoms is optionally replaced by oxygen, sulfur, or nitrogen, wherein R^5 is optionally substituted by 0-3 occurrences of R^7.

[0097] In yet other embodiments, R^1, R^2, R^3 and R^4 are each hydrogen, x is 1 and compounds have general formula **I-C-v**:




I-C-v

wherein R^5 is halogen, R', CN, -CH_2CN, -$(CH_2)_2CN$, NO_2, -CH_2NO_2, -$(CH_2)_2NO_2$, $CON(R')_2$, -$CH_2CON(R')_2$, -$(CH_2)_2CON(R')_2$, COOR', -CH_2COOR', -$(CH_2)_2COOR'$, -$SO_2N(R')_2$, -$CH_2SO_2N(R')_2$, -$(CH_2)_2SO_2N(R')_2$, -NR'SO_2R', -CH_2NR'SO_2R', -$(CH_2)_2$NR'SO_2R', NR'$CON(R')_2$, -CH_2NR'$CON(R')_2$, -$(CH_2)_2$NR'$CON(R')_2$, -NR'$SO_2N(R')_2$, -CH_2NR'$SO_2N(R')_2$, -$(CH_2)_2$NR'$SO_2N(R')_2$, -COCOR', -CH_2COCOR', -$(CH_2)_2$COCOR',–$N(R')_2$, -$CH_2N(R')_2$, -$(CH_2)_2N(R')_2$, -OR', -CH_2OR', -$(CH_2)_2OR'$, -NR'COR', -CH_2NR'COR', or –$(CH_2)_2$NR'COR', and R' is a C_1-C_6aliphatic group or a 3-8-membered saturated, partially unsaturated, or fully unsaturated monocyclic ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur, or an 8-12 membered saturated, partially unsaturated, or fully unsaturated bicyclic ring system having 0-5 heteroatoms independently selected from nitrogen, oxygen, or sulfur; or two occurrences of R', are taken together with the atom(s) to which they are bound to form an optionally substituted 3-12 membered saturated, partially unsaturated, or fully unsaturated monocyclic or bicyclic ring having 0-4 heteroatoms independently selected from

nitrogen, oxygen, or sulfur, wherein each occurrence of R' is optionally substituted with y occurrences of R^7. In some embodiments, R^5 is CN, $-CH_2CN$, $-(CH_2)_2CN$, $-NO_2$, $-CH_2NO_2$, $-(CH_2)_2NO_2$, OR', $-CH_2OR'$, $-CON(R')_2$, $-SO_2N(R')_2$, $-N(R')_2$, or R'. In certain embodiments, R' is hydrogen, C_1-C_6alkyl optionally substituted with 1-3 occurrences of R^7, or is a 5-10-membered monocyclic or bicyclic saturated, partially unsaturated or fully unsaturated ring having 0-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur, wherein the ring is optionally substituted with 1-3 occurrences of R^7. In other embodiments, R' is hydrogen, C_1-C_4alkyl optionally substituted with 1-3 occurrences of R^7, or is a ring selected from (i)-(xLvi) or (xLvii) described in [0059].

[0098] In still other embodiments, each occurrence of R^5 is independently CN, $-CH_2CN$, $-(CH_2)_2CN$, $-NO_2$, $-CH_2NO_2$, $-(CH_2)_2NO_2$, OR', $-CH_2OR'$, $-CON(R')_2$, $-SO_2N(R')_2$, $-N(R')_2$, or R'. In still other embodiments, each occurrence of R^5 is independently hydrogen, halogen, CN, $-CH_2CN$, $-(CH_2)_2CN$, NO_2, $-CH_2NO_2$, $-(CH_2)_2NO_2$, $-CONH_2$, $-CON(C_1$-C_4alkyl), $-SO_2NH_2$, $-SO_2N(C_1$-C_4alkyl), NH_2, $-N(C_1$-C_4alkyl), -OH, $-O(C_1$-C_4alkyl), $-CH_2OH$, $-CH_2O(C_1$-C_4alkyl), or an optionally substituted 5- or 6-membered unsaturated ring wherein 0-3 ring carbon atoms is optionally replaced by oxygen, sulfur, or nitrogen, wherein R^5 is optionally substituted by 0-3 occurrences of R^7.

[0099] In yet other embodiments, R^1, R^2, R^3 and R^4 are each hydrogen, x is 2 and compounds have general formula **I-C-vi**:



I-C-vi


wherein each occurrence of R^5 is independently halogen, R', CN, $-CH_2CN$, $-(CH_2)_2CN$, NO_2, $-CH_2NO_2$, $-(CH_2)_2NO_2$, $CON(R')_2$, $-CH_2CON(R')_2$, $-(CH_2)_2CON(R')_2$, COOR', $-CH_2COOR'$, $-(CH_2)_2COOR'$, $-SO_2N(R')_2$, $-CH_2SO_2N(R')_2$, $-(CH_2)_2SO_2N(R')_2$, $-NR'SO_2R'$, $-CH_2NR'SO_2R'$, $-(CH_2)_2NR'SO_2R'$, $NR'CON(R')_2$, $-CH_2NR'CON(R')_2$, $-(CH_2)_2NR'CON(R')_2$, $-NR'SO_2N(R')_2$, -

$CH_2NR'SO_2N(R')_2$, $-(CH_2)_2NR'SO_2N(R')_2$, -COCOR', $-CH_2COCOR'$, $-(CH_2)_2COCOR'$,–$N(R')_2$, $-CH_2N(R')_2$, $-(CH_2)_2N(R')_2$, -OR', $-CH_2OR'$, $-(CH_2)_2OR'$, -NR'COR', $-CH_2NR'COR'$, or $-(CH_2)_2NR'COR'$, and R' is a C_1-C_6aliphatic group or a 3-8-membered saturated, partially unsaturated, or fully unsaturated monocyclic ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur, or an 8-12 membered saturated, partially unsaturated, or fully unsaturated bicyclic ring system having 0-5 heteroatoms independently selected from nitrogen, oxygen, or sulfur; or two occurrences of R', are taken together with the atom(s) to which they are bound to form an optionally substituted 3-12 membered saturated, partially unsaturated, or fully unsaturated monocyclic or bicyclic ring having 0-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur, wherein each occurrence of R' is optionally substituted with y occurrences of R^7. In some embodiments, R^5 is CN, $-CH_2CN$, $-(CH_2)_2CN$, $-NO_2$, $-CH_2NO_2$, $-(CH_2)_2NO_2$, OR', $-CH_2OR'$, $-CON(R')_2$, $-SO_2N(R')_2$, $-N(R')_2$, or R'. In certain embodiments, R' is hydrogen, C_1-C_6alkyl optionally substituted with 1-3 occurrences of R^7, or is a 5-10-membered monocyclic or bicyclic saturated, partially unsaturated or fully unsaturated ring having 0-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur, wherein the ring is optionally substituted with 1-3 occurrences of R^7. In other embodiments, R' is hydrogen, C_1-C_4alkyl optionally substituted with 1-3 occurrences of R^7, or is a ring selected from (i)-(xLvi) or (xLvii) described in [0059].


[00100] In still other embodiments, each occurrence of R^5 is independently CN, $-CH_2CN$, $-(CH_2)_2CN$, $-NO_2$, $-CH_2NO_2$, $-(CH_2)_2NO_2$, OR', $-CH_2OR'$, $-CON(R')_2$, $-SO_2N(R')_2$, $-N(R')_2$, or R'. In still other embodiments, each occurrence of R^5 is independently hydrogen, halogen, CN, $-CH_2CN$, $-(CH_2)_2CN$, NO_2, $-CH_2NO_2$, $-(CH_2)_2NO_2$, $-CONH_2$, $-CON(C_1$-C_4alkyl), $-SO_2NH_2$, $-SO_2N(C_1$-C_4alkyl), NH_2, $-N(C_1$-C_4alkyl), -OH, $-O(C_1$-C_4alkyl), $-CH_2OH$, $-CH_2O(C_1$-C_4alkyl), or an optionally substituted 5- or 6-membered unsaturated ring wherein 0-3 ring carbon atoms is optionally replaced by oxygen, sulfur, or nitrogen, wherein R^5 is optionally substituted by 0-3 occurrences of R^7.

[00101] Representative examples of compounds of formula I are set forth below in Table 1.


[00102] Table 1. Examples of Compounds of Formula I:




I-1




I-2




I-3




I-4




I-5




I-6




I-7




I-8




I-9




I-10




I-11




I-12




I-13




I-14




I-15




I-16




I-17




I-18




I-19




I-20




I-21




I-22




I-23




I-24




I-25




I-26




I-27




I-28




I-29




I-30




I-31




I-32




I-33




I-34




I-35




I-36




I-37




I-38




I-39




I-40




I-41




I-42




I-43




I-44




I-45




I-46




I-47




I-48




I-49




I-50




I-51




I-52



I-53




I-54



I-55

[00103] Certain additional subsets of compounds of general formula I include:

[00104] **Compounds of formula I-B:**



I-B

wherein R^1, R^2, R^3, R^4, R^5 and x are each described generally above and in subsets described above and herein.

[00105] In some embodiments, for compounds of formula I-B:

a. R^1 is:

i. T-R', wherein T is a bond or an optionally substituted C_1-C_6alkylidene chain wherein up to two methylene units are optionally and independently replaced with –O-, -S-, -NR'-, -OCO-, -COO-, $-SO_2$-or –CO-, and R' is hydrogen, C_1-C_4-alkyl, or an optionally substituted 5- or 6-membered aryl or heteroaryl group, or

ii. $-Si(R')_3$, wherein R' is hydrogen, C_1-C_4-alkyl, or an optionally substituted 5- or 6-membered saturated, partially unsaturated, or fully unsaturated monocyclic ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur;

b. R^2, R^3, and R^4 are each independently hydrogen, R', halogen, CN, NO_2, -N(R')$_2$, -CH_2N(R')$_2$, -OR', -CH_2OR', -SR', -CH_2SR', -COOR', -NR'COR', -NR'C(O)OR', -NR'$COCH_2$R', -NR'CO(CH_2)$_2$R', -CON(R')$_2$, -SO_2N(R')$_2$, -CONR'(CH_2)$_2$N(R')$_2$, -CONR'(CH_2)$_3$N(R')$_2$, -CONR'(CH_2)$_4$N(R')$_2$, -O(CH_2)$_2$OR', O(CH_2)$_3$OR', O(CH_2)$_4$OR', -O(CH_2)$_2$N(R')$_2$, -O(CH_2)$_3$N(R')$_2$, or -O(CH_2)$_4$N(R')$_2$; wherein R^2, R^3, and R^4 are each optionally substituted with z occurrences of R^6, wherein z is 0-5 and R^6 is =O, =NR", =S, halogen, -CN, -NO_2, or Z-R", wherein Z is a bond or an optionally substituted C_1-C_6 alkylidene chain, wherein up to two methylene units of the chain are optionally and independently replaced by –NR"-, -S-, -O-, -CS-, -CO_2-, -OCO-, -CO-, -COCO-, -CONR"-, -NR"CO-, -NR"CO_2-, -SO_2NR"-, -NR"SO_2-, -CONR"NR"-, -NR"CONR"-, -OCONR"-, -NR"NR"-, -NR"SO_2NR"-, -SO-, -SO_2-, -PO-, -PO_2-, or –POR"-, and each occurrence of R" is independently hydrogen or an optionally substituted C_1-C_6 aliphatic group, a 3-8-membered saturated, partially unsaturated, or fully unsaturated monocyclic ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur, or an 8-12 membered saturated, partially unsaturated, or fully unsaturated bicyclic ring system having 0-5 heteroatoms independently selected from nitrogen, oxygen, or sulfur; or two occurrences of R" are taken together with the atom(s) to which they are bound to form an optionally substituted 3-12 membered saturated, partially unsaturated, or fully unsaturated monocyclic or bicyclic ring having 0-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur; and

c. each occurrence of R^5 is independently hydrogen, R', -CH_2R', halogen, CN, NO_2, -N(R')$_2$, -CH_2N(R')$_2$, -OR', -CH_2OR', -SR', -CH_2SR', -COOR', -NR'COR', -NR'$COCH_2$R', -NR'CO(CH_2)$_2$R', -NR'COOR', -CON(R')$_2$, -SO_2N(R')$_2$, -CONR'(CH_2)$_2$N(R')$_2$, -CONR(CH_2)$_3$N(R')$_2$, -CONR'(CH_2)$_4$N(R')$_2$, -O(CH_2)$_2$OR', O(CH_2)$_3$OR', O(CH_2)$_4$OR', -O(CH_2)$_2$N(R')$_2$, -O(CH_2)$_3$N(R')$_2$, -

$O(CH_2)_4N(R')_2$, -NR'CH(CH_2OH)R', -NR'CH(CH_2CH_2OH)R', -NR'(CH_2)R', -NR'$CH_2(CH_3)$R', -NR'(CH_2)$_2$R', -NR'(CH_2)$_3$R', -NR'(CH_2)$_4$R', -NR'(CH_2)N(R')$_2$, -NR'(CH_2)$_2$N(R')$_2$, -NR'(CH_2)$_3$N(R')$_2$, -NR'(CH_2)$_4$N(R')$_2$, -NR'(CH_2)OR', -NR'(CH_2)$_2$OR', -NR'(CH_2)$_3$OR', or -NR'(CH_2)$_4$OR', wherein R^5 is optionally substituted with y occurrences of R^7, wherein y is 0-5 and R^7 is =O, =NR", =S, halogen, -CN, -NO_2, or W-R', wherein W is a bond or an optionally substituted C_1-C_6 alkylidene chain, wherein up to two methylene units of the chain are optionally and independently replaced by –NR"-, -S-, -O-, -CS-, -CO_2-, -OCO-, -CO-, -COCO-, -CONR"-, -NR"CO-, -NR"CO_2-, -SO_2NR"-, -NR"SO_2-, -CONR"NR"-, -NR"CONR"-, -OCONR"-, -NR"NR"-, -NR"SO_2NR"-, -SO-, -SO_2-, -PO-, -PO_2-, or –POR"-, and and each occurrence of R" is independently hydrogen or an optionally substituted C_1-C_6 aliphatic group, a 3-8-membered saturated, partially unsaturated, or fully unsaturated monocyclic ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur, or an 8-12 membered saturated, partially unsaturated, or fully unsaturated bicyclic ring system having 0-5 heteroatoms independently selected from nitrogen, oxygen, or sulfur; or two occurrences of R" are taken together with the atom(s) to which they are bound to form an optionally substituted 3-12 membered saturated, partially unsaturated, or fully unsaturated monocyclic or bicyclic ring having 0-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur.

[00106] In still other embodiments, for compounds of formula **I-B**:

a. R^1 is hydrogen, C_1-C_4alkyl, -COR', -SO_2R', or –Si(R')$_3$;

b. R^2 and R^4 are each independently Cl, Br, F, -CN, -COOH, -COOMe, -NH_2, –N(CH_3)$_2$, -N(Et)$_2$, -N(iPr)$_2$, -O(CH_2)$_2$$OCH_3$, -$CONH_2$, -$COOCH_3$, -OH, -$CH_2$OH, -NHCO$CH_3$, -$SO_2NH_2$, -$SO_2$N(Me)$_2$, or an optionally substituted group selected from C_1-C_4alkyl, C_1-C_4alkyloxy, a 3-8-membered saturated, partially unsaturated, or fully unsaturated monocyclic ring having 0-3 heteroatoms independently selected from

nitrogen, oxygen, or sulfur, or an 8-12 membered saturated, partially unsaturated, or fully unsaturated bicyclic ring system having 0-5 heteroatoms independently selected from nitrogen, oxygen, or sulfur; wherein R^2 and R^4 are each independently and optionally substituted with z occurrences of R^6, wherein z is 0, 1, 2, or 3, and each occurrence of R^6 is independently hydrogen, R", -CH_2R", halogen, CN, NO_2, -N(R")$_2$, -CH_2N(R")$_2$, -OR", -CH_2OR", -SR", -CH_2SR", -COOR", -NR"COR", -NR"COOR", -CON(R")$_2$, -SO_2N(R")$_2$, -CONR"$(CH_2)_2$N(R")$_2$, -CONR$(CH_2)_3$N(R")$_2$, -CONR"$(CH_2)_4$N(R")$_2$, -O$(CH_2)_2$OR", O$(CH_2)_3$OR", O$(CH_2)_4$OR", -O$(CH_2)_2$N(R")$_2$, -O$(CH_2)_3$N(R")$_2$, -O$(CH_2)_4$N(R")$_2$, -NR"CH(CH_2OH)R", -NR"CH(CH_2CH_2OH)R", -NR"(CH_2)R", -NR"$(CH_2)_2$R", -NR"$(CH_2)_3$R", -NR"$(CH_2)_4$R", -NR"(CH_2)N(R")$_2$, -NR"$(CH_2)_2$N(R")$_2$, -NR"$(CH_2)_3$N(R")$_2$, -NR"$(CH_2)_4$N(R")$_2$, -NR"(CH_2)OR", -NR"$(CH_2)_2$OR", -NR"$(CH_2)_3$OR", or –NR'$(CH_2)_4$OR";

c. R^3 is independently Cl, Br, F, -CN, -COOH, -COOMe, -NH_2, –N$(CH_3)_2$, -N$(Et)_2$, -N$(iPr)_2$, -O$(CH_2)_2OCH_3$, -$CONH_2$, -$COOCH_3$, -OH, -CH_2OH, -$NHCOCH_3$, -SO_2NH_2, -SO_2N$(Me)_2$, or an optionally substituted group selected from C_1-C_4alkyl, C_1-C_4alkyloxy, wherein R^3 is independently and optionally substituted with z occurrences of R^6, wherein z is 0 or 1, and each occurrence of R^6 is independently hydrogen, R", -CH_2R", halogen, CN, NO_2, -N(R")$_2$, -CH_2N(R")$_2$, -OR", -CH_2OR", -SR", -CH_2SR", -COOR", -NR"COR", -NR"COOR", -CON(R")$_2$, -SO_2N(R")$_2$, -CONR"$(CH_2)_2$N(R")$_2$, -CONR$(CH_2)_3$N(R")$_2$, -CONR"$(CH_2)_4$N(R")$_2$, -O$(CH_2)_2$OR", O$(CH_2)_3$OR", O$(CH_2)_4$OR", -O$(CH_2)_2$N(R")$_2$, -O$(CH_2)_3$N(R")$_2$, -O$(CH_2)_4$N(R")$_2$, -NR"CH(CH_2OH)R", -NR"CH(CH_2CH_2OH)R", -NR"(CH_2)R", -NR"$(CH_2)_2$R", -NR"$(CH_2)_3$R", -NR"$(CH_2)_4$R", -NR"(CH_2)N(R")$_2$, -NR"$(CH_2)_2$N(R")$_2$, -NR"$(CH_2)_3$N(R")$_2$, -NR"$(CH_2)_4$N(R")$_2$, -NR"(CH_2)OR", -NR"$(CH_2)_2$OR", -NR"$(CH_2)_3$OR", or –NR'$(CH_2)_4$OR";

d. each occurrence of R^5 is independently hydrogen, R', -CH_2R', halogen, CN, NO_2, -N(R')$_2$, -CH_2N(R')$_2$, -OR', -CH_2OR', -SR', -CH_2SR', -COOR', -NR'COR', -NR'COCH_2R', -NR'CO(CH_2)$_2$R', -NR'COOR', -CON(R')$_2$, -SO_2N(R')$_2$, -CONR'(CH_2)$_2$N(R')$_2$, -CONR(CH_2)$_3$N(R')$_2$, -CONR'(CH_2)$_4$N(R')$_2$, -O(CH_2)$_2$OR', O(CH_2)$_3$OR', O(CH_2)$_4$OR', -O(CH_2)$_2$N(R')$_2$, -O(CH_2)$_3$N(R')$_2$, -O(CH_2)$_4$N(R')$_2$, -NR'CH(CH_2OH)R', -NR'CH(CH_2CH_2OH)R', -NR'(CH_2)R', -NR'CH_2(CH_3)R', -NR'(CH_2)$_2$R', -NR'(CH_2)$_3$R', -NR'(CH_2)$_4$R', -NR'(CH_2)N(R')$_2$, -NR'(CH_2)$_2$N(R')$_2$, -NR'(CH_2)$_3$N(R')$_2$, -NR'(CH_2)$_4$N(R')$_2$, -NR'(CH_2)OR', -NR'(CH_2)$_2$OR', -NR'(CH_2)$_3$OR', or -NR'(CH_2)$_4$OR', wherein R^5 is optionally substituted with y occurrences of R^7, wherein y is 0, 1, 2, or 3, and each occurrence of R^7 is independently hydrogen, R", -CH_2R", halogen, CN, NO_2, -N(R")$_2$, -CH_2N(R")$_2$, -OR", -CH_2OR", -SR", -CH_2SR", -COOR", -NR"COR", -NR"COOR", -CON(R")$_2$, -SO_2N(R")$_2$, -CONR"(CH_2)$_2$N(R")$_2$, -CONR(CH_2)$_3$N(R")$_2$, -CONR"(CH_2)$_4$N(R")$_2$, -O(CH_2)$_2$OR", O(CH_2)$_3$OR", O(CH_2)$_4$OR", -O(CH_2)$_2$N(R")$_2$, -O(CH_2)$_3$N(R")$_2$, -O(CH_2)$_4$N(R")$_2$, -NR"CH(CH_2OH)R", -NR"CH(CH_2CH_2OH)R", -NR"(CH_2)R", -NR"(CH_2)$_2$R", -NR"(CH_2)$_3$R", -NR"(CH_2)$_4$R", -NR"(CH_2)N(R")$_2$, -NR"(CH_2)$_2$N(R")$_2$, -NR"(CH_2)$_3$N(R")$_2$, -NR"(CH_2)$_4$N(R")$_2$, -NR"(CH_2)OR", -NR"(CH_2)$_2$OR", -NR"(CH_2)$_3$OR", or –NR"(CH_2)$_4$OR".

[00107] In other embodiments, for compounds of formula **I-B** and subsets described directly above, R^1 is hydrogen, methyl, ethyl, n-propyl, isopropyl, n-butyl, p-toluenesulfonyl (Ts), t-butyldimethylsilyl (TBS), triisopropylsilyl (TIPS), or triethylsilyl (TES).

[00108] In still other embodiments, for compounds of formula **I-B** and subsets described directly above, R^2, R^3, and R^4 are each hydrogen. In other embodiments, one of R^2, R^3, or R^4 is hydrogen. In yet other embodiments, two of R^2, R^3, or R^4 is hydrogen. In yet other embodiments, R^2 and R^4 are both hydrogen, and R^3 is halogen,

CN, NO_2, or V-R'. In still further embodiments, R^2 and R^4 are both hydrogen and R^3 is halogen. In yet further embodiments, R^2 and R^4 are both hydrogen and R^3 is Cl.

[00109] In other embodiments, for compounds of formula **I-B** and subsets described directly above, R^2, R^3, and R^4 are each independently and optionally substituted with z occurrences of R^6, wherein z is 1, 2, or 3 and each occurrence of R^6 is independently F, Cl, Br, CN, OH, NH_2, -CH_2OH, C_1-C_6alkyl, -O(C_1-C_6alkyl), -CH_2O(C_1-C_6alkyl), -CO(C_1-C_6alkyl), -COO(C_1-C_6alkyl), -$NHSO_2$(C_1-C_6alkyl), -SO_2NH_2, -$CONH_2$, -CON(C_1-C_6alkyl), -SO_2(C_1-C_6alkyl), -SO_2phenyl, phenyl, benzyl, –N(C_1-C_6alkyl)$_2$, or -S(C_1-C_6alkyl), wherein each of the foregoing phenyl, benzyl, and C_1-C_6alkyl groups is independently and optionally substituted, and wherein each of the foregoing C_1-C_6alkyl groups is linear, branched, or cyclic with the proviso that R^3 is not phenyl.

[00110] In certain exemplary embodiments, for compounds of formula **I-B** and subsets described directly above, x is 1, 2, or 3, and at least one occurrence of R^5 is –N(R')$_2$, -NR'CH(CH_2OH)R', -NR'CH(CH_2CH_2OH)R', -NR'(CH_2)R', -NR'CH_2(CH_3)R', -NR'(CH_2)$_2$R', NR'(CH_2)N(R')$_2$, or -NR'(CH_2)$_2$N(R')$_2$. In other embodiments, x is 1, 2, or 3, and at least one occurrence of R^5 is -OR'. In yet other embodiments, x is 1, 2, or 3, and at least one occurrence of R^5 is -NR'COR', -NR'$COCH_2$R', or -NR'CO(CH_2)$_2$R'. In still other embodiments, x is 1, 2, or 3, and at least one occurrence of R^5 is an optionally substituted C_1-C_6 aliphatic group, a 3-8-membered saturated, partially unsaturated, or fully unsaturated monocyclic ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur, or an 8-12 membered saturated, partially unsaturated, or fully unsaturated bicyclic ring system having 0-5 heteroatoms independently selected from nitrogen, oxygen, or sulfur.


[00111] In still other embodiments, for compounds of formula **I-B** and subsets described directly above, R^5 is optionally substituted with y occurrences of R^7, wherein y is 1, 2, or 3 and each occurrence of R^7 is independently F, Cl, Br, CN, OH, NH_2, -CH_2OH, C_1-C_6alkyl, -O(C_1-C_6alkyl), -CH_2O(C_1-C_6alkyl), -CO(C_1-C_6alkyl), -COO(C_1-C_6alkyl), -$NHSO_2$(C_1-C_6alkyl), -SO_2NH_2, -$CONH_2$, -CON(C_1-C_6alkyl), -SO_2(C_1-C_6alkyl), -SO_2phenyl, phenyl, benzyl, -N(C_1-C_6alkyl)$_2$, or -S(C_1-C_6alkyl), wherein each of the foregoing phenyl, benzyl, and C_1-C_6alkyl groups is independently

and optionally substituted, and wherein each of the foregoing C_1-C_6alkyl groups is linear, branched, or cyclic.

[00112] In yet other embodiments, for compounds of formula **I-B** and subsets described directly above, x is 1, 2, or 3; at least one occurence of R^5 is $-N(R')_2$, -NR'CH(CH_2OH)R', -NR'CH(CH_2CH_2OH)R', -NR'(CH_2)R', -NR'CH_2(CH_3)R', -NR'(CH_2)$_2$R', NR'(CH_2)N(R')$_2$,-NR'(CH_2)$_2$N(R')$_2$, -OR', -NR'COR', -NR'$COCH_2$R', or -NR'CO(CH_2)$_2$R'; and R' is a C_1-C_6aliphatic group or a 3-8-membered saturated, partially unsaturated, or fully unsaturated monocyclic ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur, or an 8-12 membered saturated, partially unsaturated, or fully unsaturated bicyclic ring system having 0-5 heteroatoms independently selected from nitrogen, oxygen, or sulfur; or two occurrences of R', are taken together with the atom(s) to which they are bound to form an optionally substituted 3-12 membered saturated, partially unsaturated, or fully unsaturated monocyclic or bicyclic ring having 0-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur, wherein each occurrence of R' is optionally substituted with y occurrences of R^7. In certain embodiments, R' is hydrogen, C_1-C_6alkyl optionally substituted with 1-3 occurrences of R^7, or is a 5-10-membered monocyclic or bicyclic saturated, partially unsaturated or fully unsaturated ring having 0-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur, wherein the ring is optionally substituted with 1-3 occurrences of R^7. In other embodiments, R' is hydrogen, C_1-C_4alkyl optionally substituted with 1-3 occurrences of R^7, or is a ring selected from (i)-(xLvi) or (xLvii) described in [0059]. In yet other embodiments, R^5 is -N(R')$_2$ and the two occurrences of R' are taken together with the nitrogen atom to which they are bound to form an optionally substituted 3-10-membered monocyclic or bicyclic heterocyclic ring. In certain embodiments, the ring is selected from (a)-(o) described in [0060].

wherein y and R^7 are described generally and in subsets above.


[00113] In yet other embodiments, x is 1 and compounds have general formula **I-B-i**:



I-B-i

wherein R^1, R^2, R^3, and R^4 are described generally and in subsets above and herein, and R^5 is $-N(R')_2$, $-NR'CH(CH_2OH)R'$, $-NR'CH(CH_2CH_2OH)R'$, $-NR'(CH_2)R'$, $-NR'CH_2(CH_3)R'$ $-NR'(CH_2)_2R'$, $NR'(CH_2)N(R')_2$, $-NR'(CH_2)_2N(R')_2$, $-OR'$, $-NR'COR'$, $-NR'COCH_2R'$, or $-NR'CO(CH_2)_2R'$,; and R' is a C_1-C_6aliphatic group or a 3-8-membered saturated, partially unsaturated, or fully unsaturated monocyclic ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur, or an 8-12 membered saturated, partially unsaturated, or fully unsaturated bicyclic ring system having 0-5 heteroatoms independently selected from nitrogen, oxygen, or sulfur; or two occurrences of R', are taken together with the atom(s) to which they are bound to form an optionally substituted 3-12 membered saturated, partially unsaturated, or fully unsaturated monocyclic or bicyclic ring having 0-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur, wherein each occurrence of R' is optionally substituted with y occurrences of R^7. In some embodiments, R^5 is $N(R')_2$. In certain embodiments, R' is hydrogen, C_1-C_6alkyl optionally substituted with 1-3 occurrences of R^7, or is a 5-10-membered monocyclic or bicyclic saturated, partially unsaturated or fully unsaturated ring having 0-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur, wherein the ring is optionally substituted with 1-3 occurrences of R^7. In other embodiments, R' is hydrogen, C_1-C_4alkyl optionally substituted with 1-3 occurrences of R^7, or is a ring selected from (i)-(xLvi) or (xLvii) described in [0059]. In yet other embodiments, R^5 is $-N(R')_2$ and the two occurrences of R' are taken together with the nitrogen atom to which they are bound to form an optionally substituted 3-10-membered monocyclic or bicyclic heterocyclic ring. In certain embodiments, the ring is selected from (a)-(o) described in [0060].


[00114] In yet other embodiments, x is 0-3 and compounds have general formula **I-B-ii**:



I-B-ii

wherein R^1, R^2, R^3, and R^4 are described generally and in subsets above and herein, and each R^5 is independently selected from halogen, optionally substituted C_1-C_6 alkyl, -SR', -CN, -COOH, -CO_2R', -CON((R')$_2$, –N(R')$_2$, -NR'CH(CH_2OH)R', -NR'CH(CH_2CH_2OH)R', -NR'(CH_2)R', -NR'CH(CH_3)R' -NR'(CH_2)$_2$R', NR'(CH_2)N(R')$_2$,-NR'(CH_2)$_2$N(R')$_2$, -OR', -NR'COR', -NR'COCH$_2$R', or -NR'CO(CH_2)$_2$R',; and R' is a C_1-C_6 aliphatic group or a 3-8-membered saturated, partially unsaturated, or fully unsaturated monocyclic ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur, or an 8-12 membered saturated, partially unsaturated, or fully unsaturated bicyclic ring system having 0-5 heteroatoms independently selected from nitrogen, oxygen, or sulfur; or two occurrences of R', are taken together with the atom(s) to which they are bound to form an optionally substituted 3-12 membered saturated, partially unsaturated, or fully unsaturated monocyclic or bicyclic ring having 0-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur, wherein each occurrence of R' is optionally substituted with y occurrences of R^7. In some embodiments, R^5 is N(R')$_2$. In certain embodiments, R' is hydrogen, C_1-C_6 alkyl optionally substituted with 1-3 occurrences of R^7, or is a 5-10-membered monocyclic or bicyclic saturated, partially unsaturated or fully unsaturated ring having 0-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur, wherein the ring is optionally substituted with 1-3 occurrences of R^7. In other embodiments, R' is hydrogen, C_1-C_4 alkyl optionally substituted with 1-3 occurrences of R^7, or is a ring selected from (i)-(xLvi) or (xLvii) described in [0059]. In yet other embodiments, R^5 is -N(R')$_2$ and the two occurrences of R' are taken together with the nitrogen atom to which they are bound to form an optionally substituted 3-10-membered monocyclic or bicyclic heterocyclic ring. In certain embodiments, the ring is selected from (a)-(o) described in [0060].

[00115] In yet other embodiments, x is 1-4 and compounds have general formula **I-B-iii:**




I-B-iii

wherein:

R^3 is an optionally substituted group selected from halogen, optionally substituted C_{1-6} alkyl, CN, $N(R')_2$, CO_2R', $NR'COR'$, $CON(R')_2$, $CH_2N(R')_2$, OR', SR', CH_2OR';

x is 1, 2, or 3; and at least one occurence of R^5 is selected from halogen, optionally substituted C1-C6 aliphatic group, -SR', -CN, -COOH, $-CO_2R'$, $-CON((R')_2$, $-N(R')_2$, $-NR'CH(CH_2OH)R'$, $-NR'CH(CH_2CH_2OH)R'$, $-NR'(CH_2)R'$, $-NR'CH(CH_3)R'$ - $NR'(CH_2)_2R'$, $NR'(CH_2)N(R')_2$,$-NR'(CH_2)_2N(R')_2$, $-OR'$, $-NR'COR'$, - $NR'COCH_2R'$, or $-NR'CO(CH_2)_2R'$; and R' is selected from hydrogen, a C_1-C_6aliphatic group optionally substituted with y occurrences of R^7, or is a ring selected from (i)-(xLvi) or (xLvii) described in [0059]. In yet other embodiments, R^5 is - $N(R')_2$ and the two occurrences of R' are taken together with the nitrogen atom to which they are bound to form an optionally substituted 3-10-membered monocyclic or bicyclic heterocyclic ring. In certain embodiments, the ring is selected from (a)-(o) described in [0060].

[00116] In still other embodiments, R^1, R^2 and R^4 are each hydrogen, and x is 0-3, and compounds of formula **I-B-iv** are provided:




I-B-iv

wherein:

R^3 is an optionally substituted group selected from halogen, optionally substituted C1-6 aliphatic, CN, $N(R')_2$, CO_2R', $NR'COR'$, $CON(R')_2$, $CH_2N(R')_2$, OR', SR', CH_2OR';

R^5 is selected from halogen, optionally substituted C1-C6 aliphatic group, -SR', -CN, -COOH, $-CO_2R'$, $-CON((R')_2$, $-N(R')_2$, $-NR'CH(CH_2OH)R'$, $-NR'CH(CH_2CH_2OH)R'$, $-NR'(CH_2)R'$, $-NR'CH(CH_3)R'$ $-NR'(CH_2)_2R'$, $NR'(CH_2)N(R')_2$, $-NR'(CH_2)_2N(R')_2$, -OR', -NR'COR', $-NR'COCH_2R'$, or $-NR'CO(CH_2)_2R'$; and R' is selected from hydrogen, a C_1-C_6aliphatic group optionally substituted with y occurrences of R^7, or is a ring selected from (i)-(xLvi) or (xLvii) described in [0059]. In yet other embodiments, R^5 is $-N(R')_2$ and the two occurrences of R' are taken together with the nitrogen atom to which they are bound to form an optionally substituted 3-10-membered monocyclic or bicyclic heterocyclic ring. In certain embodiments, the ring is selected from (a)-(o) described in [0060].

[00117] In still other embodiments, R^1, R^2 and R^4 are each hydrogen, and x is 2, and compounds of formula **I-B-v** are provided:



I-B-v


wherein:

R^3 is an optionally substituted group selected from halogen, optionally substituted C_{1-6} aliphatic, CN, $N(R')_2$, CO_2R', $NR'COR'$, $CON(R')_2$, $CH_2N(R')_2$, OR', SR', CH_2OR';

R^5 is selected from halogen, optionally substituted C_1-C_6 aliphatic group, -SR', -CN, -COOH, $-CO_2R'$, $-CON((R')_2$, $-N(R')_2$, $-NR'CH(CH_2OH)R'$, $-NR'CH(CH_2CH_2OH)R'$, $-NR'(CH_2)R'$, $-NR'CH(CH_3)R'$ $-NR'(CH_2)_2R'$, $NR'(CH_2)N(R')_2$, $-NR'(CH_2)_2N(R')_2$, -OR', -NR'COR', $-NR'COCH_2R'$, or $-NR'CO(CH_2)_2R'$; and R' is selected from hydrogen, a C_1-C_6aliphatic group optionally substituted with y occurrences of R^7, or is a ring selected from (i)-(xLvi) or (xLvii)

described in [0059]. In yet other embodiments, R^5 is -N(R')$_2$ and the two occurrences of R' are taken together with the nitrogen atom to which they are bound to form an optionally substituted 3-10-membered monocyclic or bicyclic heterocyclic ring. In certain embodiments, the ring is selected from (a)-(o) described in [0060].

[00118] In still other embodiments, R^1, R^2 and R^4 are each hydrogen, and x is 2, and compounds of formula **I-B-vi** are provided:



I-B-vi


wherein:

R^3 is an optionally substituted group selected from halogen, optionally substituted C1-6 aliphatic, CN, N(R')$_2$, CO_2R', NR'COR', CON(R')$_2$, CH_2N(R')$_2$, OR', SR', CH_2OR';

R^5 is selected from halogen, optionally substituted C1-C6 aliphatic group, -SR', -CN, -COOH, -CO_2R', -CON((R')$_2$, –N(R')$_2$, -NR'CH(CH_2OH)R', -NR'CH(CH_2CH_2OH)R', -NR'(CH_2)R', -NR'CH(CH_3)R' -NR'(CH_2)$_2$R', NR'(CH_2)N(R')$_2$,-NR'(CH_2)$_2$N(R')$_2$, -OR', -NR'COR', -NR'$COCH_2$R', or -NR'$CO(CH_2)_2$R'; and R' is selected from hydrogen, a C_1-C_6aliphatic group optionally substituted with y occurrences of R^7, or is a ring selected from (i)-(xLvi) or (xLvii) described in [0059]. In yet other embodiments, R^5 is -N(R')$_2$ and the two occurrences of R' are taken together with the nitrogen atom to which they are bound to form an optionally substituted 3-10-membered monocyclic or bicyclic heterocyclic ring. In certain embodiments, the ring is selected from (a)-(o) described in [0060].

[00119] Certain additional subsets of compounds of general formula I include:

[00120] Compounds of formula I-E:



I-E

wherein R^1, R^2, R^3, R^4, R^5 and x are each described generally above and in subsets described above and herein.

[00121] In some embodiments, for compounds of formula **I-E**:

a. R^1 is:

i. T-R', wherein T is a bond or an optionally substituted C_1-C_6alkylidene chain wherein up to two methylene units are optionally and independently replaced with –O–, -S-, -NR'-, -OCO-, -COO-, $-SO_2$-or –CO-, and R' is hydrogen, C_1-C_4-alkyl, or an optionally substituted 5- or 6-membered aryl or heteroaryl group, or

ii. –Si(R')$_3$, wherein R' is hydrogen, C_1-C_4-alkyl, or an optionally substituted 5- or 6-membered saturated, partially unsaturated, or fully unsaturated monocyclic ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur;

b. R^2, R^3, and R^4 are each independently hydrogen, R', halogen, CN, NO_2, -N(R')$_2$, -CH_2N(R')$_2$, -OR', -CH_2OR', -SR', -CH_2SR', -COOR', -NR'COR', -NR'C(O)OR', -NR'$COCH_2$R', -NR'$CO(CH_2)_2$R', -CON(R')$_2$, -SO_2N(R')$_2$, -CONR'$(CH_2)_2$N(R')$_2$, -CONR'$(CH_2)_3$N(R')$_2$, -CONR'$(CH_2)_4$N(R')$_2$, -$O(CH_2)_2$OR', $O(CH_2)_3$OR', $O(CH_2)_4$OR', -$O(CH_2)_2$N(R')$_2$, -$O(CH_2)_3$N(R')$_2$, or -$O(CH_2)_4$N(R')$_2$; wherein R^2, R^3, and R^4 are each optionally substituted with z occurrences of R^6, wherein z is 0-5 and R^6 is =O, =NR", =S, halogen, -CN, -NO_2, or Z-R", wherein Z is a bond or an optionally substituted C_1–C_6 alkylidene chain, wherein up to two methylene units of the chain are optionally and independently replaced by –NR"-, -S-, -O-, -CS-, -CO_2-, -OCO-, -CO-, -COCO-, -CONR"-, -NR"CO-, -NR"CO_2-, -SO_2NR"-, -NR"SO_2-

, -CONR"NR"-, -NR"CONR"-, -OCONR"-, -NR"NR"-, -NR"SO_2NR"-, -SO-, -SO_2-, -PO-, -PO_2-, or –POR"-, and each occurrence of R" is independently hydrogen or an optionally substituted C_1-C_6 aliphatic group, a 3-8-membered saturated, partially unsaturated, or fully unsaturated monocyclic ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur, or an 8-12 membered saturated, partially unsaturated, or fully unsaturated bicyclic ring system having 0-5 heteroatoms independently selected from nitrogen, oxygen, or sulfur; or two occurrences of R" are taken together with the atom(s) to which they are bound to form an optionally substituted 3-12 membered saturated, partially unsaturated, or fully unsaturated monocyclic or bicyclic ring having 0-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur;

c. each occurrence of R^5 is independently hydrogen, R', -CH_2R', halogen, CN, NO_2, -N(R')$_2$, -CH_2N(R')$_2$, -OR', -CH_2OR', -SR', -CH_2SR', -COOR', -NR'COR', -NR'$COCH_2$R', -NR'$CO(CH_2)_2$R', -NR'COOR', -CON(R')$_2$, -SO_2N(R')$_2$, -CONR'$(CH_2)_2$N(R')$_2$, -CONR$(CH_2)_3$N(R')$_2$, -CONR'$(CH_2)_4$N(R')$_2$, -O$(CH_2)_2$OR', O$(CH_2)_3$OR', O$(CH_2)_4$OR', -O$(CH_2)_2$N(R')$_2$, -O$(CH_2)_3$N(R')$_2$, -O$(CH_2)_4$N(R')$_2$, -NR'CH(CH_2OH)R', -NR'CH(CH_2CH_2OH)R', -NR'(CH_2)R', -NR'CH(CH_3)R', -NR'$(CH_2)_2$R', -NR'$(CH_2)_3$R', -NR'$(CH_2)_4$R', -NR'(CH_2)N(R')$_2$, -NR'$(CH_2)_2$N(R')$_2$, -NR'$(CH_2)_3$N(R')$_2$, -NR'$(CH_2)_4$N(R')$_2$, -NR'(CH_2)OR', -NR'$(CH_2)_2$OR', -NR'$(CH_2)_3$OR', or -NR'$(CH_2)_4$OR', wherein R^5 is optionally substituted with y occurrences of R^7, wherein y is 0-5 and R^7 is =O, =NR", =S, halogen, -CN, -NO_2, or W-R', wherein W is a bond or an optionally substituted C_1-C_6 alkylidene chain, wherein up to two methylene units of the chain are optionally and independently replaced by –NR"-, -S-, -O-, -CS-, -CO_2-, -OCO-, -CO-, -COCO-, -CONR"-, -NR"CO-, -NR"CO_2-, -SO_2NR"-, -NR"SO_2-, -CONR"NR"-, -NR"CONR"-, -OCONR"-, -NR"NR"-, -NR"SO_2NR"-, -SO-, -SO_2-, -PO-, -PO_2-, or –POR"-, and and each occurrence of R" is

independently hydrogen or an optionally substituted C_1.C_6 aliphatic group, a 3-8-membered saturated, partially unsaturated, or fully unsaturated monocyclic ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur, or an 8-12 membered saturated, partially unsaturated, or fully unsaturated bicyclic ring system having 0-5 heteroatoms independently selected from nitrogen, oxygen, or sulfur; or two occurrences of R" are taken together with the atom(s) to which they are bound to form an optionally substituted 3-12 membered saturated, partially unsaturated, or fully unsaturated monocyclic or bicyclic ring having 0-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur.

[00122] In still other embodiments, for compounds of formula **I-E**:

a. R^1 is hydrogen, C_1-C_4alkyl, -COR', $-SO_2R'$, or $-Si(R')_3$;

b. R^2 and R^4 are each independently Cl, Br, F, -CN, -COOH, -COOMe, $-NH_2$, $-N(CH_3)_2$, $-N(Et)_2$, $-N(iPr)_2$, $-O(CH_2)_2OCH_3$, $-CONH_2$, $-COOCH_3$, -OH, $-CH_2OH$, $-NHCOCH_3$, $-SO_2NH_2$, $-SO_2N(Me)_2$, or an optionally substituted group selected from C_1-C_4alkyl, C_1-C_4alkyloxy, a 3-8-membered saturated, partially unsaturated, or fully unsaturated monocyclic ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur, or an 8-12 membered saturated, partially unsaturated, or fully unsaturated bicyclic ring system having 0-5 heteroatoms independently selected from nitrogen, oxygen, or sulfur; wherein R^2, R^3, and R^4 are each independently and optionally substituted with z occurrences of R^6, wherein z is 0, 1, 2, or 3, and each occurrence of R^6 is independently hydrogen, R", $-CH_2R$", halogen, CN, NO_2, $-N(R")_2$, $-CH_2N(R")_2$, -OR", $-CH_2OR$", -SR", $-CH_2SR$", -COOR", -NR"COR", -NR"COOR", $-CON(R")_2$, $-SO_2N(R")_2$, $-CONR"(CH_2)_2N(R")_2$, $-CONR(CH_2)_3N(R")_2$, $-CONR"(CH_2)_4N(R")_2$, $-O(CH_2)_2OR$", $O(CH_2)_3OR$", $O(CH_2)_4OR$", $-O(CH_2)_2N(R")_2$, $-O(CH_2)_3N(R")_2$, $-O(CH_2)_4N(R")_2$, -NR"CH(CH_2OH)R", -NR"CH(CH_2CH_2OH)R", -NR"(CH_2)R", $-NR"(CH_2)_2R$", $-NR"(CH_2)_3R$", $-NR"(CH_2)_4R$", $-NR"(CH_2)N(R")_2$, $-NR"(CH_2)_2N(R")_2$,

-NR"$(CH_2)_3N(R")_2$, -NR"$(CH_2)_4N(R")_2$, -NR"$(CH_2)OR$", -NR"$(CH_2)_2OR$", -NR"$(CH_2)_3OR$", or -NR'$(CH_2)_4OR$";

c. R^3 is independently Cl, Br, F, -CN, -COOH, -COOMe, $-NH_2$, $-N(CH_3)_2$, $-N(Et)_2$, $-N(iPr)_2$, $-O(CH_2)_2OCH_3$, $-CONH_2$, $-COOCH_3$, -OH, $-CH_2OH$, $-NHCOCH_3$, $-SO_2NH_2$, $-SO_2N(Me)_2$, or an optionally substituted group selected from C_1-C_4alkyl, C_1-C_4alkyloxy, wherein R^3 is independently and optionally substituted with z occurrences of R^6, wherein z is 0 or 1, and each occurrence of R^6 is independently hydrogen, R", $-CH_2R$", halogen, CN, NO_2, $-N(R")_2$, $-CH_2N(R")_2$, -OR", $-CH_2OR$", -SR", $-CH_2SR$", -COOR", -NR"COR", -NR"COOR", $-CON(R")_2$, $-SO_2N(R")_2$, $-CONR"(CH_2)_2N(R")_2$, $-CONR(CH_2)_3N(R")_2$, $-CONR"(CH_2)_4N(R")_2$, $-O(CH_2)_2OR$", $O(CH_2)_3OR$", $O(CH_2)_4OR$", $-O(CH_2)_2N(R")_2$, $-O(CH_2)_3N(R")_2$, $-O(CH_2)_4N(R")_2$, -NR"$CH(CH_2OH)R$", -NR"$CH(CH_2CH_2OH)R$", -NR"$(CH_2)R$", -NR"$(CH_2)_2R$", -NR"$(CH_2)_3R$", -NR"$(CH_2)_4R$", -NR"$(CH_2)N(R")_2$, -NR"$(CH_2)_2N(R")_2$, -NR"$(CH_2)_3N(R")_2$, -NR"$(CH_2)_4N(R")_2$, -NR"$(CH_2)OR$", -NR"$(CH_2)_2OR$", -NR"$(CH_2)_3OR$", or -NR'$(CH_2)_4OR$";

d. each occurrence of R^5 is independently hydrogen, R', $-CH_2R$', halogen, CN, NO_2, $-N(R')_2$, $-CH_2N(R')_2$, -OR', $-CH_2OR$', -SR', $-CH_2SR$', -COOR', -NR'COR', -NR'$COCH_2R$', -NR'$CO(CH_2)_2R$', -NR'COOR', $-CON(R')_2$, $-SO_2N(R')_2$, $-CONR'(CH_2)_2N(R')_2$, $-CONR(CH_2)_3N(R')_2$, $-CONR'(CH_2)_4N(R')_2$, $-O(CH_2)_2OR$', $O(CH_2)_3OR$', $O(CH_2)_4OR$', $-O(CH_2)_2N(R')_2$, $-O(CH_2)_3N(R')_2$, $-O(CH_2)_4N(R')_2$, -NR'$CH(CH_2OH)R$', -NR'$CH(CH_2CH_2OH)R$', -NR'$(CH_2)R$', -NR'$CH(CH_3)R$', -NR'$(CH_2)_2R$', -NR'$(CH_2)_3R$', -NR'$(CH_2)_4R$', -NR'$(CH_2)N(R')_2$, -NR'$(CH_2)_2N(R')_2$, -NR'$(CH_2)_3N(R')_2$, -NR'$(CH_2)_4N(R')_2$, -NR'$(CH_2)OR$', -NR'$(CH_2)_2OR$', -NR'$(CH_2)_3OR$', or -NR'$(CH_2)_4OR$', wherein R^5 is optionally substituted with y occurrences of R^7, wherein y is 0, 1, 2, or 3, and each occurrence of R^7 is independently hydrogen, R", $-CH_2R$", halogen, CN, NO_2, $-N(R")_2$, $-CH_2N(R")_2$, -OR", $-CH_2OR$", -SR", -

CH_2SR", -COOR", -NR"COR", -NR"COOR", -$CON(R")_2$, -$SO_2N(R")_2$, -CONR"$(CH_2)_2N(R")_2$, -CONR$(CH_2)_3N(R")_2$, -CONR"$(CH_2)_4N(R")_2$, -$O(CH_2)_2$OR", $O(CH_2)_3$OR", $O(CH_2)_4$OR", -$O(CH_2)_2N(R")_2$, -$O(CH_2)_3N(R")_2$, -$O(CH_2)_4N(R")_2$, -NR"$CH(CH_2OH)$R", -NR"$CH(CH_2CH_2OH)$R", -NR"(CH_2)R", -NR"$(CH_2)_2$R", -NR"$(CH_2)_3$R", -NR"$(CH_2)_4$R", -NR"$(CH_2)N(R")_2$, -NR"$(CH_2)_2N(R")_2$, -NR"$(CH_2)_3N(R")_2$, -NR"$(CH_2)_4N(R")_2$, -NR"(CH_2)OR", -NR"$(CH_2)_2$OR", -NR"$(CH_2)_3$OR", or –NR"$(CH_2)_4$OR".

[00123] In other embodiments, for compounds of formula **I-E** and subsets described directly above, R^1 is hydrogen, methyl, ethyl, n-propyl, isopropyl, n-butyl, p-toluenesulfonyl (Ts), t-butyldimethylsilyl (TBS), triisopropylsilyl (TIPS), or triethylsilyl (TES).

[00124] In still other embodiments, for compounds of formula **I-E** and subsets described directly above, R^2, R^3, and R^4 are each hydrogen. In other embodiments, one of R^2, R^3, or R^4 is hydrogen. In yet other embodiments, two of R^2, R^3, or R^4 is hydrogen. In yet other embodiments, R^2 and R^4 are both hydrogen, and R^3 is halogen, CN, NO_2, or V-R'. In still further embodiments R^2 and R^4 are both hydrogen and R^3 is halogen. In yet further embodiments R^2 and R^4 are both hydrogen and R^3 is Cl.

[00125] In other embodiments, for compounds of formula **I-E** and subsets described directly above, R^2, R^3, and R^4 are each independently and optionally substituted with z occurrences of R^6, wherein z is 1, 2, or 3 and each occurrence of R^6 is independently F, Cl, Br, CN, OH, NH_2, $-CH_2OH$, C_1-C_6alkyl, -O(C_1-C_6alkyl), -CH_2O(C_1-C_6alkyl), -CO(C_1-C_6alkyl), -COO(C_1-C_6alkyl), -$NHSO_2$(C_1-C_6alkyl), -SO_2NH_2, -$CONH_2$, -CON(C_1-C_6alkyl), -SO_2(C_1-C_6alkyl), -SO_2phenyl, phenyl, benzyl, -N(C_1-C_6alkyl)$_2$, or -S(C_1-C_6alkyl), wherein each of the foregoing phenyl, benzyl, and C_1-C_6alkyl groups is independently and optionally substituted, and wherein each of the foregoing C_1-C_6alkyl groups is linear, branched, or cyclic.

[00126] In certain exemplary embodiments, for compounds of formula **I-E** and subsets described directly above, x is 1, 2, or 3, and at least one occurrence of R^5 is –N(R')$_2$, -NR'$CH(CH_2OH)$R', -NR'$CH(CH_2CH_2OH)$R', –NR'(CH_2)R', -NR'$CH(CH_3)$R', -NR'$(CH_2)_2$R', NR'$(CH_2)N(R')_2$, or -NR'$(CH_2)_2N(R')_2$. In other

embodiments, x is 1, 2, or 3, and at least one occurrence of R^5 is -OR'. In yet other embodiments, x is 1, 2, or 3, and at least one occurrence of R^5 is -NR'COR', -NR'$COCH_2$R', or -NR'$CO(CH_2)_2$R'. In still other embodiments, x is 1, 2, or 3, and at least one occurrence of R^5 is an optionally substituted C_1-C_6 aliphatic group, a 3-8-membered saturated, partially unsaturated, or fully unsaturated monocyclic ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur, or an 8-12 membered saturated, partially unsaturated, or fully unsaturated bicyclic ring system having 0-5 heteroatoms independently selected from nitrogen, oxygen, or sulfur.


[00127] In still other embodiments, for compounds of formula **I-E** and subsets described directly above, R^5 is optionally substituted with y occurrences of R^7, wherein y is 1, 2, or 3 and each occurrence of R^7 is independently F, Cl, Br, CN, OH, NH_2, -CH_2OH, C_1-C_6alkyl, -O(C_1-C_6alkyl), -CH_2O(C_1-C_6alkyl), -CO(C_1-C_6alkyl), -COO(C_1-C_6alkyl), -$NHSO_2$(C_1-C_6alkyl), -SO_2NH_2, -$CONH_2$, -CON(C_1-C_6alkyl), -SO_2(C_1-C_6alkyl), -SO_2phenyl, phenyl, benzyl, -N(C_1-C_6alkyl)$_2$, or -S(C_1-C_6alkyl), wherein each of the foregoing phenyl, benzyl, and C_1-C_6alkyl groups is independently and optionally substituted, and wherein each of the foregoing C_1-C_6alkyl groups is linear, branched, or cyclic.

[00128] In yet other embodiments, for compounds of formula **I-E** and subsets described directly above, x is 1, 2, or 3; at least one occurence of R^5 is –N(R')$_2$, -NR'CH(CH_2OH)R', -NR'CH(CH_2CH_2OH)R', -NR'(CH_2)R', -NR'CH(CH_3)R', -NR'(CH_2)$_2$R', NR'(CH_2)N(R')$_2$,-NR'(CH_2)$_2$N(R')$_2$, -OR', -NR'COR', -NR'$COCH_2$R', or -NR'$CO(CH_2)_2$R'; and R' is a C_1-C_6aliphatic group or a 3-8-membered saturated, partially unsaturated, or fully unsaturated monocyclic ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur, or an 8-12 membered saturated, partially unsaturated, or fully unsaturated bicyclic ring system having 0-5 heteroatoms independently selected from nitrogen, oxygen, or sulfur; or two occurrences of R', are taken together with the atom(s) to which they are bound to form an optionally substituted 3-12 membered saturated, partially unsaturated, or fully unsaturated monocyclic or bicyclic ring having 0-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur, wherein each occurrence of R' is optionally substituted with y occurrences of R^7. In certain embodiments, R' is

hydrogen, C_1-C_6alkyl optionally substituted with 1-3 occurrences of R^7, or is a 5-10-membered monocyclic or bicyclic saturated, partially unsaturated or fully unsaturated ring having 0-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur, wherein the ring is optionally substituted with 1-3 occurrences of R^7. In other embodiments, R' is hydrogen, C_1-C_4alkyl optionally substituted with 1-3 occurrences of R^7, or is a ring selected from (i)-(xLvi) or (xLvii) described in [0059]. In yet other embodiments, R^5 is $-N(R')_2$ and the two occurrences of R' are taken together with the nitrogen atom to which they are bound to form an optionally substituted 3-10-membered monocyclic or bicyclic heterocyclic ring. In certain embodiments, the ring is selected from (a)-(o) described in [0060].

wherein y and R^7 are described generally and in subsets above.

[00129] In yet other embodiments, x is 1 and compounds have general formula **I-E-i**:




I-E-i

wherein R^1, R^2, R^3, and R^4 are described generally and in subsets above and herein, and R^5 is $-N(R')_2$, $-NR'CH(CH_2OH)R'$, $-NR'CH(CH_2CH_2OH)R'$, $-NR'(CH_2)R'$, $-NR'CH_2(CH_3)R'$ $-NR'(CH_2)_2R'$, $NR'(CH_2)N(R')_2$, $-NR'(CH_2)_2N(R')_2$, $-OR'$, $-NR'COR'$, $-NR'COCH_2R'$, or $-NR'CO(CH_2)_2R'$,; and R' is a C_1-C_6aliphatic group or a 3-8-membered saturated, partially unsaturated, or fully unsaturated monocyclic ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur, or an 8-12 membered saturated, partially unsaturated, or fully unsaturated bicyclic ring system having 0-5 heteroatoms independently selected from nitrogen, oxygen, or sulfur; or two occurrences of R', are taken together with the atom(s) to which they are bound to form an optionally substituted 3-12 membered saturated, partially unsaturated, or fully unsaturated monocyclic or bicyclic ring having 0-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur, wherein each occurrence of R' is optionally substituted with y occurrences of R^7. In some embodiments, R^5 is $N(R')_2$. In certain embodiments, R' is hydrogen, C_1-C_6alkyl

optionally substituted with 1-3 occurrences of R^7, or is a 5-10-membered monocyclic or bicyclic saturated, partially unsaturated or fully unsaturated ring having 0-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur, wherein the ring is optionally substituted with 1-3 occurrences of R^7. In other embodiments, R' is hydrogen, C_1-C_4alkyl optionally substituted with 1-3 occurrences of R^7, or is a ring selected from (i)-(xLvi) or (xLvii) described in [0059]. In yet other embodiments, R^5 is $-N(R')_2$ and the two occurrences of R' are taken together with the nitrogen atom to which they are bound to form an optionally substituted 3-10-membered monocyclic or bicyclic heterocyclic ring. In certain embodiments, the ring is selected from (a)-(o) described in [0060].

[00130] In still other embodiments, R^1, R^2, and R^4 are each hydrogen, and compounds of formula **I-E-ii** are provided:




I-E-ii

wherein:

R^3 is an optionally substituted group selected from halogen, optionally substituted C1-6 alkyl, CN, N(R')2, CO2R', NR'COR', CON(R')2, CH2N(R')2, OR', SR', CH2OR';

x is 1, 2, or 3; and at least one occurence of R^5 is $-N(R')_2$, -NR'CH(CH_2OH)R', -NR'CH(CH_2CH_2OH)R', -NR'(CH_2)R', -NR $CH_2(CH_3)$R', -NR'(CH_2)$_2$R', NR'(CH_2)N(R')$_2$,-NR'(CH_2)$_2$N(R')$_2$, -OR', -NR'COR', -NR'$COCH_2$R', or -NR'CO(CH_2)$_2$R'; and R' is a C_1-C_6aliphatic group optionally substituted with y occurrences of R^7, or is a ring selected from (i)-(xLvi) or (xLvii) described in [0059]. In yet other embodiments, R^5 is $-N(R')_2$ and the two occurrences of R' are taken together with the nitrogen atom to which they are bound to form an optionally substituted 3-10-membered monocyclic or bicyclic heterocyclic ring. In certain embodiments, the ring is selected from (a)-(o) described in [0060].

[00131] In still other embodiments, R^1, R^2 and R^4 are each hydrogen, and x is 0-3, and compounds of formula **I-E-iii** are provided:




I-E-iii

wherein:

R^3 is an optionally substituted group selected from halogen, optionally substituted C1-6 alkyl, CN, $N(R')_2$, CO_2R', NR'COR', $CON(R')_2$, $CH_2N(R')_2$, OR', SR', CH_2OR';

R^5 is $-N(R')_2$, -NR'CH(CH_2OH)R', -NR'CH(CH_2CH_2OH)R', -NR'(CH_2)R', -NR'$CH_2(CH_3)$R', -NR'(CH_2)$_2$R', NR'(CH_2)N(R')$_2$,-NR'(CH_2)$_2$N(R')$_2$, -OR', -NR'COR', -NR'$COCH_2$R', or -NR'CO(CH_2)$_2$R'; and R' is a C_1-C_6aliphatic group optionally substituted with y occurrences of R^7, or is a ring selected from (i)-(xLvi) or (xLvii) described in [0059]. In yet other embodiments, R^5 is -$N(R')_2$ and the two occurrences of R' are taken together with the nitrogen atom to which they are bound to form an optionally substituted 3-10-membered monocyclic or bicyclic heterocyclic ring. In certain embodiments, the ring is selected from (a)-(o) described in [0060].

[00132] In still other embodiments, R^1, R^2 and R^4 are each hydrogen, and x is 1, and compounds of formula **I-E-iv** are provided:



I-E-iv


wherein:



R^3 is an optionally substituted group selected from halogen, optionally substituted C_{1-6} alkyl, CN, $N(R')_2$, CO_2R', NR'COR', $CON(R')_2$, $CH_2N(R')_2$, OR', SR', CH_2OR'; R^5 is $-N(R')_2$, -NR'CH(CH_2OH)R', -NR'CH(CH_2CH_2OH)R', -



$NR'(CH_2)R'$, $-NR'CH_2(CH_3)R'$, $-NR'(CH_2)_2R'$, $NR'(CH_2)N(R')_2$, $-NR'(CH_2)_2N(R')_2$, $-OR'$, $-NR'COR'$, $-NR'COCH_2R'$, or $-NR'CO(CH_2)_2R'$; and R' is a C_1-C_6aliphatic group optionally substituted with y occurrences of R^7, or is a ring selected from (i)-(xLvi) or (xLvii) described in [0059]. In yet other embodiments, R^5 is $-N(R')_2$ and the two occurrences of R' are taken together with the nitrogen atom to which they are bound to form an optionally substituted 3-10-membered monocyclic or bicyclic heterocyclic ring. In certain embodiments, the ring is selected from (a)-(o) described in [0060].



Representative examples of compounds of formula I are set forth below in Table 2. Compounds of Table 2 may also be represented by II-x, where x is the compound number indicated in Table 2.



Table 2. Examples of Compounds of Formula **I**:



Cmpd No.	Compound
1	



Cmpd No.	Compound
2	
3	



Cmpd No.	Compound
4	
5	



Cmpd No.	Compound
6	
7	



Cmpd No.	Compound
8	
9	



Cmpd No.	Compound
10	
11	



Cmpd No.	Compound
12	
13	



Cmpd No.	Compound
14	
15	



Cmpd No.	Compound
16	
17	



Cmpd No.	Compound
18	
19	



Cmpd No.	Compound
20	
21	



Cmpd No.	Compound
22	
23	



Cmpd No.	Compound
24	
25	



Cmpd No.	Compound
26	
27	



Cmpd No.	Compound
28	
29	



Cmpd No.	Compound
30	
31	



Cmpd No.	Compound
32	
33	



Cmpd No.	Compound
34	
35	



Cmpd No.	Compound
36	
37	



Cmpd No.	Compound
38	
39	



Cmpd No.	Compound
40	
41	



Cmpd No.	Compound
42	
43	



Cmpd No.	Compound
44	
45	



Cmpd No.	Compound
46	
47	



Cmpd No.	Compound
48	
49	



Cmpd No.	Compound
50	
51	



Cmpd No.	Compound
52	
53	



Cmpd No.	Compound
54	
55	



Cmpd No.	Compound
56	
57	



Cmpd No.	Compound
58	
59	



Cmpd No.	Compound
60	
61	



Cmpd No.	Compound
62	
63	



Cmpd No.	Compound
64	
65	



Cmpd No.	Compound
66	
67	



Cmpd No.	Compound
68	
69	



Cmpd No.	Compound
70	
71	



Cmpd No.	Compound
72	
73	



Cmpd No.	Compound
74	
75	



Cmpd No.	Compound
76	
77	



Cmpd No.	Compound
78	
79	



Cmpd No.	Compound
80	
81	



Cmpd No.	Compound
82	
83	



Cmpd No.	Compound
84	
85	



Cmpd No.	Compound
86	
87	



Cmpd No.	Compound
88	
89	



Cmpd No.	Compound
90	
91	



Cmpd No.	Compound
92	
93	



Cmpd No.	Compound
94	
95	



Cmpd No.	Compound
96	
97	



Cmpd No.	Compound
98	
99	



Cmpd No.	Compound
100	
101	



Cmpd No.	Compound
102	
103	



Cmpd No.	Compound
104	
105	



Cmpd No.	Compound
106	
107	



Cmpd No.	Compound
108	
109	



Cmpd No.	Compound
110	
111	



Cmpd No.	Compound
112	
113	



Cmpd No.	Compound
114	
115	



Cmpd No.	Compound
116	
117	



Cmpd No.	Compound
118	
119	



Cmpd No.	Compound
120	
121	



Cmpd No.	Compound
122	
123	



Cmpd No.	Compound
124	
125	



Cmpd No.	Compound
126	
127	



Cmpd No.	Compound
128	
129	



Cmpd No.	Compound
130	
131	



Cmpd No.	Compound
132	
133	



Cmpd No.	Compound
134	
135	



Cmpd No.	Compound
136	
137	



Cmpd No.	Compound
138	
139	



Cmpd No.	Compound
140	
141	



Cmpd No.	Compound
142	
143	



Cmpd No.	Compound
144	
145	



Cmpd No.	Compound
146	
147	



Cmpd No.	Compound
148	
149	



Cmpd No.	Compound
150	
151	



Cmpd No.	Compound
152	
153	



Cmpd No.	Compound
154	
155	



Cmpd No.	Compound
156	
157	



Cmpd No.	Compound
158	
159	



Cmpd No.	Compound
160	
161	



Cmpd No.	Compound
162	
163	



Cmpd No.	Compound
164	
165	



Cmpd No.	Compound
166	
167	



Cmpd No.	Compound
168	
169	



Cmpd No.	Compound
170	
171	



Cmpd No.	Compound
172	
173	



Cmpd No.	Compound
174	
175	



Cmpd No.	Compound
176	
177	



Cmpd No.	Compound
178	
179	



Cmpd No.	Compound
180	
181	



Cmpd No.	Compound
182	
183	



Cmpd No.	Compound
184	
185	



Cmpd No.	Compound
186	
187	



Cmpd No.	Compound
188	
189	



Cmpd No.	Compound
190	
191	



Cmpd No.	Compound
192	
193	



Cmpd No.	Compound
194	
195	



Cmpd No.	Compound
196	
197	



Cmpd No.	Compound
198	
199	



Cmpd No.	Compound
200	
201	



Cmpd No.	Compound
202	
203	



Cmpd No.	Compound
204	
205	



Cmpd No.	Compound
206	
207	



Cmpd No.	Compound
208	
209	



Cmpd No.	Compound
210	
211	



Cmpd No.	Compound
212	
213	



Cmpd No.	Compound
214	
215	



Cmpd No.	Compound
216	
217	



Cmpd No.	Compound
218	
219	



Cmpd No.	Compound
220	
221	



Cmpd No.	Compound
222	
223	



Cmpd No.	Compound
224	
225	



Cmpd No.	Compound
226	
227	



Cmpd No.	Compound
228	
229	



Cmpd No.	Compound
230	
231	



Cmpd No.	Compound
232	
233	



Cmpd No.	Compound
234	
235	



Cmpd No.	Compound
236	
237	



Cmpd No.	Compound
238	
239	



Cmpd No.	Compound
240	
241	



Cmpd No.	Compound
242	
243	



Cmpd No.	Compound
244	
245	



Cmpd No.	Compound
246	
247	



Cmpd No.	Compound
248	
249	



Cmpd No.	Compound
250	
251	



Cmpd No.	Compound
252	
253	



Cmpd No.	Compound
254	
255	



Cmpd No.	Compound
256	
257	



Cmpd No.	Compound
258	
259	



Cmpd No.	Compound
260	
261	



Cmpd No.	Compound
262	
263	



Cmpd No.	Compound
264	
265	



Cmpd No.	Compound
266	
267	



Cmpd No.	Compound
268	
269	



Cmpd No.	Compound
270	
271	



Cmpd No.	Compound
272	
273	



Cmpd No.	Compound
274	
275	



Cmpd No.	Compound
276	
277	



Cmpd No.	Compound
278	
279	



Cmpd No.	Compound
280	
281	



Cmpd No.	Compound
282	
283	



Cmpd No.	Compound
284	
285	



Cmpd No.	Compound
286	
287	



Cmpd No.	Compound
288	
289	



Cmpd No.	Compound
290	
291	



Cmpd No.	Compound
292	
293	



Cmpd No.	Compound
294	
295	



Cmpd No.	Compound
296	
297	



Cmpd No.	Compound
298	
299	



Cmpd No.	Compound
300	
301	



Cmpd No.	Compound
302	
303	



Cmpd No.	Compound
304	
305	



Cmpd No.	Compound
306	
307	



Cmpd No.	Compound
308	
309	



Cmpd No.	Compound
310	
311	



Cmpd No.	Compound
312	
313	



Cmpd No.	Compound
314	
315	



Cmpd No.	Compound
316	
317	



Cmpd No.	Compound
318	
319	



Cmpd No.	Compound
320	
321	



Cmpd No.	Compound
322	
323	



Cmpd No.	Compound
324	
325	



Cmpd No.	Compound
326	
327	



Cmpd No.	Compound
328	
329	



Cmpd No.	Compound
330	
331	



Cmpd No.	Compound
332	
333	



Cmpd No.	Compound
334	
335	



Cmpd No.	Compound
336	
337	



Cmpd No.	Compound
338	
339	



Cmpd No.	Compound
340	
341	



Cmpd No.	Compound
342	
343	



Cmpd No.	Compound
344	
345	



Cmpd No.	Compound
346	
347	



Cmpd No.	Compound
348	
349	



Cmpd No.	Compound
350	
351	



Cmpd No.	Compound
352	
353	



Cmpd No.	Compound
354	
355	



Cmpd No.	Compound
356	
357	



Cmpd No.	Compound
358	
359	



Cmpd No.	Compound
360	
361	



Cmpd No.	Compound
362	
363	



Cmpd No.	Compound
364	
365	



Cmpd No.	Compound
366	
367	



Cmpd No.	Compound
368	
369	



Cmpd No.	Compound
370	
371	



Cmpd No.	Compound
372	
373	



Cmpd No.	Compound
374	
375	



Cmpd No.	Compound
376	
377	



Cmpd No.	Compound
378	
379	



Cmpd No.	Compound
380	
381	



Cmpd No.	Compound
382	
383	



Cmpd No.	Compound
384	
385	



Cmpd No.	Compound
386	
387	



Cmpd No.	Compound
388	
389	



Cmpd No.	Compound
390	
391	



Cmpd No.	Compound
392	
393	



Cmpd No.	Compound
394	
395	



Cmpd No.	Compound
396	
397	



Cmpd No.	Compound
398	
399	



Cmpd No.	Compound
400	
401	



Cmpd No.	Compound
402	
403	



Cmpd No.	Compound
404	
405	



Cmpd No.	Compound
406	
407	



Cmpd No.	Compound
408	
409	



Cmpd No.	Compound
410	
411	



Cmpd No.	Compound
412	
413	



Cmpd No.	Compound
414	
415	



Cmpd No.	Compound
416	
417	



Cmpd No.	Compound
418	
419	



Cmpd No.	Compound
420	
421	



Cmpd No.	Compound
422	
423	



Cmpd No.	Compound
424	
425	



Cmpd No.	Compound
426	
427	



Cmpd No.	Compound
428	
429	



Cmpd No.	Compound
430	
431	



Cmpd No.	Compound
432	
433	



Cmpd No.	Compound
434	
435	



Cmpd No.	Compound
436	
437	



Cmpd No.	Compound
438	
439	



Cmpd No.	Compound
440	
441	



Cmpd No.	Compound
442	
443	



Cmpd No.	Compound
444	
445	



Cmpd No.	Compound
446	
447	



Cmpd No.	Compound
448	
449	



Cmpd No.	Compound
450	
451	



Cmpd No.	Compound
452	
453	



Cmpd No.	Compound
454	
455	



Cmpd No.	Compound
456	
457	



Cmpd No.	Compound
458	
459	

Cmpd No.	Compound
460	
461	

Cmpd No.	Compound
462	
463	

Cmpd No.	Compound
464	
465	

Cmpd No.	Compound
466	
467	

Cmpd No.	Compound
468	
469	

[00133] *4. Uses, Formulation and Administration*

[00134] *Pharmaceutically acceptable compositions*

[00135] As discussed above, the present invention provides compounds that are inhibitors of protein kinases, and thus the present compounds are useful for the treatment of diseases, disorders, and conditions including, but not limited to a proliferative disorder, a cardiac disorder, a neurodegenerative disorder, psychotic disorders, an autoimmune disorder, a condition associated with organ transplant, an inflammatory disorder, an immunologically mediated disorder, a viral disease, or a

bone disorder. In certain embodiments, the compounds are useful for the treatment of immune responses such as allergic or type I hypersensitivity reactions or asthma; autoimmune diseases such as transplant rejection, graft versus host disease, rheumatoid arthritis, amyotrophic lateral sclerosis, and multiple sclerosis; neurodegenerative disorders such as Familial amyotrophic lateral sclerosis (FALS); and solid and hematologic malignancies such as leukemias and lymphomas.

[00136] These compounds and pharmaceutical compositions thereof are also useful for treating or preventing a variety of disorders, including, but not limited to, heart disease, diabetes, Alzheimer's disease, immunodeficiency disorders, inflammatory diseases, hypertension, allergic diseases, autoimmune diseases, destructive bone disorders such as osteoporosis, proliferative disorders, infectious diseases, immunologically-mediated diseases, and viral diseases. The compositions are also useful in methods for preventing cell death and hyperplasia and therefore may be used to treat or prevent reperfusion/ischemia in stroke, heart attacks, and organ hypoxia. The compositions are also useful in methods for preventing thrombin-induced platelet aggregation. The compositions are especially useful for disorders such as chronic myelogenous leukemia (CML), acute myeloid leukemia (AML), acute promyelocytic leukemia (APL), rheumatoid arthritis, asthma, osteoarthritis, ischemia, cancer (including, but not limited to, ovarian cancer, breast cancer and endometrial cancer), liver disease including hepatic ischemia, heart disease such as myocardial infarction and congestive heart failure, pathologic immune conditions involving T cell activation, and neurodegenerative disorders.

[00137] Accordingly, in another aspect of the present invention, pharmaceutically acceptable compositions are provided, wherein these compositions comprise any of the compounds as described herein, and optionally comprise a pharmaceutically acceptable carrier, adjuvant or vehicle. In certain embodiments, these compositions optionally further comprise one or more additional therapeutic agents.

[00138] It will also be appreciated that certain of the compounds of present invention can exist in free form for treatment, or where appropriate, as a pharmaceutically acceptable derivative thereof. According to the present invention, a pharmaceutically acceptable derivative includes, but is not limited to, pharmaceutically acceptable prodrugs, salts, esters, salts of such esters, or any other

adduct or derivative which upon administration to a patient in need is capable of providing, directly or indirectly, a compound as otherwise described herein, or a metabolite or residue thereof.

[00139] As used herein, the term "pharmaceutically acceptable salt" refers to those salts which are, within the scope of sound medical judgement, suitable for use in contact with the tissues of humans and lower animals without undue toxicity, irritation, allergic response and the like, and are commensurate with a reasonable benefit/risk ratio. A "pharmaceutically acceptable salt" means any non-toxic salt or salt of an ester of a compound of this invention that, upon administration to a recipient, is capable of providing, either directly or indirectly, a compound of this invention or an inhibitorily active metabolite or residue thereof. As used herein, the term "inhibitorily active metabolite or residue thereof" means that a metabolite or residue thereof is also an inhibitor of a JAK-3, ROCK and Aurora.

[00140] Pharmaceutically acceptable salts are well known in the art. For example, S. M. Berge *et al.*, describe pharmaceutically acceptable salts in detail in *J. Pharmaceutical Sciences*, **1977**, *66*, 1-19, incorporated herein by reference. Pharmaceutically acceptable salts of the compounds of this invention include those derived from suitable inorganic and organic acids and bases. Examples of pharmaceutically acceptable, nontoxic acid addition salts are salts of an amino group formed with inorganic acids such as hydrochloric acid, hydrobromic acid, phosphoric acid, sulfuric acid and perchloric acid or with organic acids such as acetic acid, oxalic acid, maleic acid, tartaric acid, citric acid, succinic acid or malonic acid or by using other methods used in the art such as ion exchange. Other pharmaceutically acceptable salts include adipate, alginate, ascorbate, aspartate, benzenesulfonate, benzoate, bisulfate, borate, butyrate, camphorate, camphorsulfonate, citrate, cyclopentanepropionate, digluconate, dodecylsulfate, ethanesulfonate, formate, fumarate, glucoheptonate, glycerophosphate, gluconate, hemisulfate, heptanoate, hexanoate, hydroiodide, 2-hydroxy-ethanesulfonate, lactobionate, lactate, laurate, lauryl sulfate, malate, maleate, malonate, methanesulfonate, 2-naphthalenesulfonate, nicotinate, nitrate, oleate, oxalate, palmitate, pamoate, pectinate, persulfate, 3-phenylpropionate, phosphate, picrate, pivalate, propionate, stearate, succinate, sulfate, tartrate, thiocyanate, p-toluenesulfonate, undecanoate, valerate salts, and the like.

Salts derived from appropriate bases include alkali metal, alkaline earth metal, ammonium and $N^+(C_{1-4}alkyl)_4$ salts. This invention also envisions the quaternization of any basic nitrogen-containing groups of the compounds disclosed herein. Water or oil-soluble or dispersable products may be obtained by such quaternization. Representative alkali or alkaline earth metal salts include sodium, lithium, potassium, calcium, magnesium, and the like. Further pharmaceutically acceptable salts include, when appropriate, nontoxic ammonium, quaternary ammonium, and amine cations formed using counterions such as halide, hydroxide, carboxylate, sulfate, phosphate, nitrate, loweralkyl sulfonate and aryl sulfonate.

[00141] As described above, the pharmaceutically acceptable compositions of the present invention additionally comprise a pharmaceutically acceptable carrier, adjuvant, or vehicle, which, as used herein, includes any and all solvents, diluents, or other liquid vehicle, dispersion or suspension aids, surface active agents, isotonic agents, thickening or emulsifying agents, preservatives, solid binders, lubricants and the like, as suited to the particular dosage form desired. Remington's Pharmaceutical Sciences, Sixteenth Edition, E. W. Martin (Mack Publishing Co., Easton, Pa., 1980) discloses various carriers used in formulating pharmaceutically acceptable compositions and known techniques for the preparation thereof. Except insofar as any conventional carrier medium is incompatible with the compounds of the invention, such as by producing any undesirable biological effect or otherwise interacting in a deleterious manner with any other component(s) of the pharmaceutically acceptable composition, its use is contemplated to be within the scope of this invention. Some examples of materials which can serve as pharmaceutically acceptable carriers include, but are not limited to, ion exchangers, alumina, aluminum stearate, lecithin, serum proteins, such as human serum albumin, buffer substances such as phosphates, glycine, sorbic acid, or potassium sorbate, partial glyceride mixtures of saturated vegetable fatty acids, water, salts or electrolytes, such as protamine sulfate, disodium hydrogen phosphate, potassium hydrogen phosphate, sodium chloride, zinc salts, colloidal silica, magnesium trisilicate, polyvinyl pyrrolidone, polyacrylates, waxes, polyethylene-polyoxypropylene-block polymers, wool fat, sugars such as lactose, glucose and sucrose; starches such as corn starch and potato starch; cellulose and its derivatives such as sodium carboxymethyl cellulose, ethyl cellulose and cellulose

acetate; powdered tragacanth; malt; gelatin; talc; excipients such as cocoa butter and suppository waxes; oils such as peanut oil, cottonseed oil; safflower oil; sesame oil; olive oil; corn oil and soybean oil; glycols; such a propylene glycol or polyethylene glycol; esters such as ethyl oleate and ethyl laurate; agar; buffering agents such as magnesium hydroxide and aluminum hydroxide; alginic acid; pyrogen-free water; isotonic saline; Ringer's solution; ethyl alcohol, and phosphate buffer solutions, as well as other non-toxic compatible lubricants such as sodium lauryl sulfate and magnesium stearate, as well as coloring agents, releasing agents, coating agents, sweetening, flavoring and perfuming agents, preservatives and antioxidants can also be present in the composition, according to the judgment of the formulator.

[00142] *Uses of Compounds and Pharmaceutically acceptable compositions*

[00143] In yet another aspect, a method for the treatment or lessening the severity of a proliferative disorder, a cardiac disorder, a neurodegenerative disorder, a psychotic disorder, an autoimmune disorder, a condition associated with organ transplant, an inflammatory disorder, an immunologically mediated disorder, a viral disease, or a bone disorder is provided comprising administering an effective amount of a compound, or a pharmaceutically acceptable composition comprising a compound to a subject in need thereof. In certain embodiments of the present invention an "effective amount" of the compound or pharmaceutically acceptable composition is that amount effective for treating or lessening the severity of a proliferative disorder, a cardiac disorder, a neurodegenerative disorder, a psychotic disorder, an autoimmune disorder, a condition associated with organ transplant, an inflammatory disorder, an immunologically mediated disorder, a viral disease, or a bone disorder.

[00144] In other aspects, the invention comprises a method for treating or lessening the severity of a variety of disorders, including, but not limited to, heart disease, diabetes, Alzheimer's disease, immunodeficiency disorders, inflammatory diseases, hypertension, allergic diseases, autoimmune diseases, destructive bone disorders such as osteoporosis, proliferative disorders, infectious diseases, immunologically-mediated diseases, and viral diseases. In other aspects, the invention comprises methods for preventing cell death and hyperplasia and therefore may be used to treat

or prevent reperfusion/ischemia in stroke, heart attacks, and organ hypoxia. In other embodiments, the invention comprises methods for preventing thrombin-induced platelet aggregation. The invention also comprises methods for treating, lessening the severity or preventing disorders such as chronic myelogenous leukemia (CML), acute myeloid leukemia (AML), acute promyelocytic leukemia (APL), rheumatoid arthritis, asthma, osteoarthritis, ischemia, cancer (including, but not limited to, ovarian cancer, breast cancer and endometrial cancer), liver disease including hepatic ischemia, heart disease such as myocardial infarction and congestive heart failure, pathologic immune conditions involving T cell activation, and neurodegenerative disorders.

[00145] The compounds and compositions, according to the method of the present invention, may be administered using any amount and any route of administration effective for treating or lessening the severity of a proliferative disorder, a cardiac disorder, a neurodegenerative disorder, an autoimmune disorder, a condition associated with organ transplant, an inflammatory disorder, an immunologically mediated disorder, a viral disease, or a bone disorder. The exact amount required will vary from subject to subject, depending on the species, age, and general condition of the subject, the severity of the infection, the particular agent, its mode of administration, and the like. The compounds of the invention are preferably formulated in dosage unit form for ease of administration and uniformity of dosage. The expression "dosage unit form" as used herein refers to a physically discrete unit of agent appropriate for the patient to be treated. It will be understood, however, that the total daily usage of the compounds and compositions of the present invention will be decided by the attending physician within the scope of sound medical judgment. The specific effective dose level for any particular patient or organism will depend upon a variety of factors including the disorder being treated and the severity of the disorder; the activity of the specific compound employed; the specific composition employed; the age, body weight, general health, sex and diet of the patient; the time of administration, route of administration, and rate of excretion of the specific compound employed; the duration of the treatment; drugs used in combination or coincidental with the specific compound employed, and like factors well known in the medical arts. The term "patient", as used herein, means an animal, preferably a mammal, and most preferably a human.

[00146] The pharmaceutically acceptable compositions of this invention can be administered to humans and other animals orally, rectally, parenterally, intracisternally, intravaginally, intraperitoneally, topically (as by powders, ointments, or drops), bucally, as an oral or nasal spray, or the like, depending on the severity of the infection being treated. In certain embodiments, the compounds of the invention may be administered orally or parenterally at dosage levels of about 0.01 mg/kg to about 50 mg/kg and preferably from about 1 mg/kg to about 25 mg/kg, of subject body weight per day, one or more times a day, to obtain the desired therapeutic effect.

[00147] Liquid dosage forms for oral administration include, but are not limited to, pharmaceutically acceptable emulsions, microemulsions, solutions, suspensions, syrups and elixirs. In addition to the active compounds, the liquid dosage forms may contain inert diluents commonly used in the art such as, for example, water or other solvents, solubilizing agents and emulsifiers such as ethyl alcohol, isopropyl alcohol, ethyl carbonate, ethyl acetate, benzyl alcohol, benzyl benzoate, propylene glycol, 1,3-butylene glycol, dimethylformamide, oils (in particular, cottonseed, groundnut, corn, germ, olive, castor, and sesame oils), glycerol, tetrahydrofurfuryl alcohol, polyethylene glycols and fatty acid esters of sorbitan, and mixtures thereof. Besides inert diluents, the oral compositions can also include adjuvants such as wetting agents, emulsifying and suspending agents, sweetening, flavoring, and perfuming agents.

[00148] Injectable preparations, for example, sterile injectable aqueous or oleaginous suspensions may be formulated according to the known art using suitable dispersing or wetting agents and suspending agents. The sterile injectable preparation may also be a sterile injectable solution, suspension or emulsion in a nontoxic parenterally acceptable diluent or solvent, for example, as a solution in 1,3-butanediol. Among the acceptable vehicles and solvents that may be employed are water, Ringer's solution, U.S.P. and isotonic sodium chloride solution. In addition, sterile, fixed oils are conventionally employed as a solvent or suspending medium. For this purpose any bland fixed oil can be employed including synthetic mono- or diglycerides. In addition, fatty acids such as oleic acid are used in the preparation of injectables.

[00149] The injectable formulations can be sterilized, for example, by filtration through a bacterial-retaining filter, or by incorporating sterilizing agents in the form

of sterile solid compositions which can be dissolved or dispersed in sterile water or other sterile injectable medium prior to use.

[00150] In order to prolong the effect of a compound of the present invention, it is often desirable to slow the absorption of the compound from subcutaneous or intramuscular injection. This may be accomplished by the use of a liquid suspension of crystalline or amorphous material with poor water solubility. The rate of absorption of the compound then depends upon its rate of dissolution that, in turn, may depend upon crystal size and crystalline form. Alternatively, delayed absorption of a parenterally administered compound form is accomplished by dissolving or suspending the compound in an oil vehicle. Injectable depot forms are made by forming microencapsule matrices of the compound in biodegradable polymers such as polylactide-polyglycolide. Depending upon the ratio of compound to polymer and the nature of the particular polymer employed, the rate of compound release can be controlled. Examples of other biodegradable polymers include poly(orthoesters) and poly(anhydrides). Depot injectable formulations are also prepared by entrapping the compound in liposomes or microemulsions that are compatible with body tissues.

[00151] Compositions for rectal or vaginal administration are preferably suppositories which can be prepared by mixing the compounds of this invention with suitable non-irritating excipients or carriers such as cocoa butter, polyethylene glycol or a suppository wax which are solid at ambient temperature but liquid at body temperature and therefore melt in the rectum or vaginal cavity and release the active compound.

[00152] Solid dosage forms for oral administration include capsules, tablets, pills, powders, and granules. In such solid dosage forms, the active compound is mixed with at least one inert, pharmaceutically acceptable excipient or carrier such as sodium citrate or dicalcium phosphate and/or a) fillers or extenders such as starches, lactose, sucrose, glucose, mannitol, and silicic acid, b) binders such as, for example, carboxymethylcellulose, alginates, gelatin, polyvinylpyrrolidinone, sucrose, and acacia, c) humectants such as glycerol, d) disintegrating agents such as agar--agar, calcium carbonate, potato or tapioca starch, alginic acid, certain silicates, and sodium carbonate, e) solution retarding agents such as paraffin, f) absorption accelerators such as quaternary ammonium compounds, g) wetting agents such as, for example,

cetyl alcohol and glycerol monostearate, h) absorbents such as kaolin and bentonite clay, and i) lubricants such as talc, calcium stearate, magnesium stearate, solid polyethylene glycols, sodium lauryl sulfate, and mixtures thereof. In the case of capsules, tablets and pills, the dosage form may also comprise buffering agents.

[00153] Solid compositions of a similar type may also be employed as fillers in soft and hard-filled gelatin capsules using such excipients as lactose or milk sugar as well as high molecular weight polyethylene glycols and the like. The solid dosage forms of tablets, dragees, capsules, pills, and granules can be prepared with coatings and shells such as enteric coatings and other coatings well known in the pharmaceutical formulating art. They may optionally contain opacifying agents and can also be of a composition that they release the active ingredient(s) only, or preferentially, in a certain part of the intestinal tract, optionally, in a delayed manner. Examples of embedding compositions that can be used include polymeric substances and waxes. Solid compositions of a similar type may also be employed as fillers in soft and hard-filled gelatin capsules using such excipients as lactose or milk sugar as well as high molecular weight polethylene glycols and the like.

[00154] The active compounds can also be in micro-encapsulated form with one or more excipients as noted above. The solid dosage forms of tablets, dragees, capsules, pills, and granules can be prepared with coatings and shells such as enteric coatings, release controlling coatings and other coatings well known in the pharmaceutical formulating art. In such solid dosage forms the active compound may be admixed with at least one inert diluent such as sucrose, lactose or starch. Such dosage forms may also comprise, as is normal practice, additional substances other than inert diluents, e.g., tableting lubricants and other tableting aids such a magnesium stearate and microcrystalline cellulose. In the case of capsules, tablets and pills, the dosage forms may also comprise buffering agents. They may optionally contain opacifying agents and can also be of a composition that they release the active ingredient(s) only, or preferentially, in a certain part of the intestinal tract, optionally, in a delayed manner. Examples of embedding compositions that can be used include polymeric substances and waxes.

[00155] Dosage forms for topical or transdermal administration of a compound of this invention include ointments, pastes, creams, lotions, gels, powders, solutions,

sprays, inhalants or patches. The active component is admixed under sterile conditions with a pharmaceutically acceptable carrier and any needed preservatives or buffers as may be required. Ophthalmic formulation, ear drops, and eye drops are also contemplated as being within the scope of this invention. Additionally, the present invention contemplates the use of transdermal patches, which have the added advantage of providing controlled delivery of a compound to the body. Such dosage forms can be made by dissolving or dispensing the compound in the proper medium. Absorption enhancers can also be used to increase the flux of the compound across the skin. The rate can be controlled by either providing a rate controlling membrane or by dispersing the compound in a polymer matrix or gel.

[00156] As described generally above, the compounds of the invention are useful as inhibitors of protein kinases. In one embodiment, the compounds and compositions of the invention are inhibitors of one or more of JAK-3, ROCK and Aurora isoforms, and thus, without wishing to be bound by any particular theory, the compounds and compositions are particularly useful for treating or lessening the severity of a disease, condition, or disorder where activation of JAK-3, ROCK or Aurora is implicated in the disease, condition, or disorder. When activation of JAK-3, ROCK or Aurora is implicated in a particular disease, condition, or disorder, the disease, condition, or disorder may also be referred to as a "JAK-3-mediated disease", "ROCK-mediated disease", "Aurora-mediated disease" or disease symptom. Accordingly, in another aspect, the present invention provides a method for treating or lessening the severity of a disease, condition, or disorder where activation or one or more of JAK-3, ROCK or Aurora isoforms is implicated in the disease state.

[00157] The activity of a compound utilized in this invention as an inhibitor of JAK-3, ROCK or Aurora may be assayed *in vitro, in vivo* or in a cell line. *In vitro* assays include assays that determine inhibition of either the phosphorylation activity or ATPase activity of activated JAK-3, ROCK or Aurora. Alternate *in vitro* assays quantitate the ability of the inhibitor to bind to JAK-3, ROCK or Aurora. Inhibitor binding may be measured by radiolabelling the inhibitor prior to binding, isolating the inhibitor/JAK-3, inhibitor/ROCK or inhibitor/Aurora complex and determining the amount of radiolabel bound. Alternatively, inhibitor binding may be determined by

running a competition experiment where new inhibitors are incubated with JAK-3, ROCK or Aurora bound to known radioligands.

[00158] The term "measurably inhibit", as used herein means a measurable change in JAK-3, ROCK or Aurora activity between a sample comprising said composition and a JAK-3, ROCK or Aurora kinase and an equivalent sample comprising JAK-3, ROCK or Aurora kinase in the absence of said composition.

[00159] The term "JAK-mediated disease", as used herein means any disease or other deleterious condition in which a JAK family kinase is known to play a role. Such conditions include, without limitation, immune responses such as allergic or type I hypersensitivity reactions, asthma, autoimmune diseases such as transplant rejection, graft versus host disease, rheumatoid arthritis, amyotrophic lateral sclerosis, and multiple sclerosis, neurodegenerative disorders such as Familial amyotrophic lateral sclerosis (FALS), as well as in solid and hematologic malignancies such as leukemias and lymphomas.

[00160] The term "ROCK-mediated condition" or "disease", as used herein, means any disease or other deleterious condition in which ROCK is known to play a role. The term "ROCK-mediated condition" or "disease" also means those diseases or conditions that are alleviated by treatment with a ROCK inhibitor. Such conditions include, without limitation, hypertension, angina pectoris, cerebrovascular contraction, asthma, peripheral circulation disorder, premature birth, cancer, erectile dysfunction, arteriosclerosis, spasm (cerebral vasospasm and coronary vasospasm), retinopathy (e.g., glaucoma), inflammatory disorders, autoimmune disorders, AIDS, osteoporosis, myocardial hypertrophy, ischemia/reperfusion-induced injury, and endothelial dysfunction.

[00161] The term "Aurora-mediated condition" or "disease", as used herein, means any disease or other deleterious condition in which Aurora is known to play a role. The term "Aurora-mediated condition" or "disease" also means those diseases or conditions that are alleviated by treatment with a Aurora inhibitor. Such conditions include, without limitation, immune responses such as allergic or type I hypersensitivity reactions, asthma, autoimmune diseases such as transplant rejection, graft versus host disease, rheumatoid arthritis, amyotrophic lateral sclerosis, and multiple sclerosis, neurodegenerative disorders such as Familial amyotrophic lateral

sclerosis (FALS), as well as in solid and hematologic malignancies such as leukemias and lymphomas.

[00162] It will also be appreciated that the compounds and pharmaceutically acceptable compositions of the present invention can be employed in combination therapies, that is, the compounds and pharmaceutically acceptable compositions can be administered concurrently with, prior to, or subsequent to, one or more other desired therapeutics or medical procedures. The particular combination of therapies (therapeutics or procedures) to employ in a combination regimen will take into account compatibility of the desired therapeutics and/or procedures and the desired therapeutic effect to be achieved. It will also be appreciated that the therapies employed may achieve a desired effect for the same disorder (for example, an inventive compound may be administered concurrently with another agent used to treat the same disorder), or they may achieve different effects (e.g., control of any adverse effects). As used herein, additional therapeutic agents that are normally administered to treat or prevent a particular disease, or condition, are known as "appropriate for the disease, or condition, being treated".

[00163] Examples of agents the inhibitors of this invention may also be combined with include, without limitation: treatments for Alzheimer's Disease such as Aricept® and Excelon®; treatments for Parkinson's Disease such as L-DOPA/carbidopa, entacapone, ropinrole, pramipexole, bromocriptine, pergolide, trihexephendyl, and amantadine; agents for treating Multiple Sclerosis (MS) such as beta interferon (e.g., Avonex® and Rebif®), Copaxone®, and mitoxantrone; treatments for asthma such as albuterol and Singulair®; agents for treating schizophrenia such as zyprexa, risperdal, seroquel, and haloperidol; anti-inflammatory agents such as corticosteroids, TNF blockers, IL-1 RA, azathioprine, cyclophosphamide, and sulfasalazine; immunomodulatory and immunosuppressive agents such as cyclosporin, tacrolimus, rapamycin, mycophenolate mofetil, interferons, corticosteroids, cyclophosphamide, azathioprine, and sulfasalazine; neurotrophic factors such as acetylcholinesterase inhibitors, MAO inhibitors, interferons, anti-convulsants, ion channel blockers, riluzole, and anti-Parkinsonian agents; agents for treating cardiovascular disease such as beta-blockers, ACE inhibitors, diuretics, nitrates, calcium channel blockers, and statins; agents for treating liver disease such as corticosteroids, cholestyramine,

interferons, and anti-viral agents; agents for treating blood disorders such as corticosteroids, anti-leukemic agents, and growth factors; and agents for treating immunodeficiency disorders such as gamma globulin.

[00164] The amount of additional therapeutic agent present in the compositions of this invention will be no more than the amount that would normally be administered in a composition comprising that therapeutic agent as the only active agent. Preferably the amount of additional therapeutic agent in the presently disclosed compositions will range from about 50% to 100% of the amount normally present in a composition comprising that agent as the only therapeutically active agent.

[00165] The compounds of this invention or pharmaceutically acceptable compositions thereof may also be incorporated into compositions for coating implantable medical devices, such as prostheses, artificial valves, vascular grafts, stents and catheters. Accordingly, the present invention, in another aspect, includes a composition for coating an implantable device comprising a compound of the present invention as described generally above, and in classes and subclasses herein, and a carrier suitable for coating said implantable device. In still another aspect, the present invention includes an implantable device coated with a composition comprising a compound of the present invention as described generally above, and in classes and subclasses herein, and a carrier suitable for coating said implantable device.

[00166] Vascular stents, for example, have been used to overcome restenosis (re-narrowing of the vessel wall after injury). However, patients using stents or other implantable devices risk clot formation or platelet activation. These unwanted effects may be prevented or mitigated by pre-coating the device with a pharmaceutically acceptable composition comprising a kinase inhibitor. Suitable coatings and the general preparation of coated implantable devices are described in US Patents 6,099,562; 5,886,026; and 5,304,121. The coatings are typically biocompatible polymeric materials such as a hydrogel polymer, polymethyldisiloxane, polycaprolactone, polyethylene glycol, polylactic acid, ethylene vinyl acetate, and mixtures thereof. The coatings may optionally be further covered by a suitable topcoat of fluorosilicone, polysaccarides, polyethylene glycol, phospholipids or combinations thereof to impart controlled release characteristics in the composition.


[00167] Another aspect of the invention relates to inhibiting JAK-3, ROCK or Aurora activity in a biological sample or a patient, which method comprises administering to the patient, or contacting said biological sample with a compound of formula I or a composition comprising said compound. The term "biological sample", as used herein, includes, without limitation, cell cultures or extracts thereof; biopsied material obtained from a mammal or extracts thereof; and blood, saliva, urine, feces, semen, tears, or other body fluids or extracts thereof.

[00168] Inhibition of JAK-3, ROCK or Aurora kinase activity in a biological sample is useful for a variety of purposes that are known to one of skill in the art. Examples of such purposes include, but are not limited to, blood transfusion, organ-transplantation, biological specimen storage, and biological assays.

EXAMPLES

[00169] Although certain exemplary embodiments are described in detail below, it will be appreciated that additional compounds of general formula I can be prepared according to the methods described generally herein using appropriate starting materials by methods generally available to one of ordinary skill in the art.

[00170] **Scheme 1**



[00171] 3-Bromo-1*H*-pyrrolo[2,3-*b*]pyridine (2).

[00172] Azaindole **1** (4g, 0.025 mol) in 100 mL of chloroform was cooled to 0°C. Bromine in 20 mL of chloroform was added dropwise the resulting mixture was stirred at 0°C for 1 h. The resulting suspension was diluted with 0.5N HCl and the aqueous layer was made basic with 0.5 N NaOH and the solid filtered to provide 4g (82%) of crude product **2** that was used directly for the next step.

[00173] 3-Bromo-1-(toluene-4-sulfonyl)-1*H*-pyrrolo[2,3-*b*]pyridine (**3**).

[00174] 3g (0.015 mol) of **2** in 20 mL of anhydrous THF was cooled to –78°C and n-BuLi 2.5 M in hexanes (6.7 mL, 0.167 mol) was added dropwise. After 15 minutes of stirring, tosyl chloride in 5 mL of THF was added dropwise. The cooling bath was removed and the reaction mixture stirred at rt for 1 h. Extracted with ether and the organic phase washed with brine, dried with magnesium sulfate and concentrated in vacuo to give a white solid that was passed through a pad of silica (70% EtOAc; 30% Hexanes) to give 4.65 g (84%) of **3**. ^{1}H NMR $CDCl_3$ 8.4 (s, 1H), 8.1 (d, 2H), 7.8 (s, 2H), 7.2 (m, 3H), 2.3 (s, 3H).

[00175] 3-(4,4,5,5-Tetramethyl-[1,3,2]dioxaborolan-2-yl)-1-(toluene-4-sulfonyl)-1*H*-pyrrolo[2,3-*b*]pyridine(**4**).

[00176] **3** (850 mg, 0.0024 mol) was dissolved in 20 mL of DME and pinnacol borane (921 mg, 0.0036 mol), Pd_2Cl_2 $(dppf)_2$ (197 mg, 0.24 mmol)and KOAc (713 mg, 0.00726 mol) were added and the mixture was stirred and refluxed at 90°C for 18h. Diluted with ethyl acetate and the organic phase washed with water and brine then dried (Na_2SO_4) and concentrated in vacuo. The residue was subjected to flash chromatography (20% EtOAc/ 80% hexanes) to give 900 mg (99%) of the desired product **4**.

[00177] 3-(2-Methylsulfanyl-pyrimidin-4-yl)-1-(toluene-4-sulfonyl)-1*H*-pyrrolo[2,3*b*] pyridine (**5**).

[00178] A mixture of boronic ester **4** (900 mg, 0.0023 mol), 4-chloro-2-thiomethyl pyrimidine (341 mg, 0.0029 mol), $Pd(Ph_3P)_4$ (260 mg, 0.23 mmol) and 2 M sodium carbonate (3.4 mL, 0.0068 mol) in 20 mL of DME was refluxed under nitrogen for 18 h. Diluted with ethyl acetate and the organic phase washed with water and brine then dried (Na_2SO_4) and concentrated in vacuo. The residue was subjected to flash chromatography (40% EtOAc/ 60% hexanes) to give 460 mg (51%) of the desired

product **5**. ^{1}H NMR $CDCl_3$ 8.8 (d, 1H), 8.7 (m, 2H), 8.4 (s, 1H), 8.1 (d, 2H), 7.2 (m, 4H), 2.6 (s, 3H), 2.3 (s, 3H).

[00179] 3-(2-Methanesulfonyl-pyrimidin-4-yl)-1-(toluene-4-sulfonyl)-1*H*-pyrrolo[2,3-*b*]pyridine (**6**).

[00180] Pyrimidine **5** (460 mg, 0.0012 mol) was dissolved in 20 mL of methanol-water (1:1) then oxone (2.14 g, 0.0035 mol) was added and the reaction was refluxed for 18 h. The methanol was removed in vacuo and the aqueous was extracted with ethyl acetate. The organic phase washed with water and brine then dried (Na_2SO_4) and concentrated in vacuo. The residue was subjected to flash chromatography (40% EtOAc/ 60% hexanes) to give 160 mg (32%) of the desired product **6**. LCMS ES^+ = 428.9.

[00181] Benzyl-[4-(1*H*-pyrrolo[2,3-*b*]pyridin-3-yl)-pyrimidin-2-yl]-amine (**7**).

[00182] A solution of **6** (20 mg, 0.047 mmol) and benzylamine (0.007 mL, 0.061 mmol) in 1 mL of ethanol was heated in a seal tube at 80°C for 18 h. The solvent was evaporated and the crude product was purified by preparative TLC (50% EtOAc/ 50% hexanes) to give 20 mg of product which was deprotected with 2 mL of 3N NaOH in methanol for 4 h. Added 2 mL of 3N HCl and evaporated to dryness. Reverse phase HPLC (20-70% MeCN-water with 0.1% TFA (20 mL/min) gave 10 mg (75%) of **7**. ^{1}H NMR $DMSOD_6$ 8.7 (s, 1H), 8.4 (s, 1H), 8.25 (d, 1H), 8.1 (d, 1H), 7.4 (m, 2H), 7.25 (m, 3H), 7.2 (s, 1H), 7.15 (dd, 1H), 7.1 (s, 1H), 4.8 (s, 2H). LCMS ES^+ = 302.0.

[00183] 3-(6-Chloro-pyrimidin-4-yl)-1-(toluene-4-sulfonyl)-1*H*-pyrrolo[2,3-*b*]pyridine (**8**).

[00184] A solution of boronate **4** (0.11 g, 0.276 mmol), 4,6-dichloropyrimidine (0.049g, 0.331 mmol), catalytic $Pd(PPh_3)_4$ and excess potassium carbonate in DMF was heated in the microwave at 160^{0}C for 5 minutes resulting in conversion to product by TLC (20% EtOAc:hexanes). The reaction was partitioned between EtOAc/H_2O, extracted, stripped down in vacuo and purified by silica column (eluent:5% EtOAc:hexanes) giving **8** (0.035g) as a white solid in 33% yield.

[00185] 1-{4-[6-(1*H*-Pyrrolo[2,3-*b*]pyridin-3-yl)-pyrimidin-4-yl]-[1,4]diazepan-1-yl}-ethanone (**9**).

[00186] A solution of **8** (0.035g, 0.091mmol), N-acetyl-homopiperizine (0.025g, 0.181mmol), excess potassium carbonate in DMF was heated at 80^{0}C resulting in a

color change after 5 minutes. LC/MS and TLC indicated conversion to tosyl protected product after 15 minutes. 1 ml of 6N NaOH and 1ml of methanol were added to the reaction resulting in the immediate removal the tosyl group by LC/MS (M+1=337). The reaction was partitioned between EtOAc/H_2O and extracted. The crude product was purified by preparative HPLC giving **9** (0.022g) as a clear oil in 73% yield.

[00187] NMR: MeOD 2.0 bs(2H), 2.1 s(3H), 3.6 m(2H), 3.8-4.3 bm(6H), 7.1 d(1H), 7.3 m(1H), 8.2 d(1H), 8.35 m(2H), 8.65 d(1H). LC/MS(M+1)=337

[00188] Benzyl-(2-chloro-pyrimidin-4-yl)-amine (**10**).

[00189] To a solution of 2,4-dichloropyrimidine (0.15g, 1.0 mmol), benzylamine (0.109ml, 1.0mmol) in THF was added DIPEA (0.526ml, 3.0 mmol) and the reaction was heated at reflux for 2 hours resulting in formation of a 4:1 mixture of regioisomers (desired versus undesired) by TLC (5%Methanol:methylene chloride). The reaction was stripped down in vacuo and purified by silica column (eluent: 2% methanol:methylene chloride) giving 0.12g (0.548 mmol) of desired product **10** in 54% yield. LC/MS(M+1) = 220

[00190] Benzyl-(6-chloro-pyrimidin-4-yl)-amine (**12**).

[00191] Benzyl amine (0.697ml, 6.76 mmol) was added to 4,6-dichloropyrimidine (1.0g, 6.76 mmol) neat causing a vigorous reaction and color change. The reaction was slowly diluted with methylene chloride resulting in a white precipitate. 1 ml of triethyl amine was added and TLC indicated conversion to product (5% MeOH:methylene chloride). The reaction was loaded directly onto a silica and purified (eluent:2%MeOH:methylene chloride) giving 1.17g (5.32 mmol) of **12** as a yellow wax in 79% yield.

[00192] Benzyl-[2-(1*H*-pyrrolo[2,3-*b*]pyridin-3-yl)-pyrimidin-4-yl]-amine (**11**).


[00193] A solution of **4** (0.36g, 0.09 mmol), **10** (0.028g, 0.108 mmol), 2.0M Na_2CO_3 (0.108ml, 0.271mmol) and catalytic $PdCl_2(PPh_3)_2$ in 1 ml of DMSO was heated in the microwave at 160^{0}C for 5 minutes resulting in conversion to tosyl protected product by LC/MS(M+1)=456. NaOtBu (.026g, 0.271mmol) was added to the reaction and it was heated in the microwave for 5 minutes at 160^{0}C resulting in complete conversion to product **11**. The reaction was filtered and purified by preparative HPLC giving 0.0032g of **11** as a white solid in 11% yield.

[00194] NMR: MeOD 4.95 bs(2H), 6.6 d(1H), 7.2 m(1H), 7.3-7.5 mm(5H), 8.0 d(1H), 8.3 d(1H), 8.4 s(1H), 8.55 d(1H). LC/MS(M+1)= 302

[00195] Final product **13** was formed as described for **11** giving 0.012g (0.039 mmol) of **13** as a white solid. NMR: MeOD 4.8 s(2H), 7.05 s(1H), 7.2-7.6 m (6H), 8.2 s(1H), 8.4 m(2H), 8.55 s(1H). LC/MS(M+1)=302

[00196]

Scheme 2 Synthesis of 5-H-azaindole pyrimidine and pyridine analogues



[00197] 3-(2-Chloro-pyridin-4-yl)-1-(toluene-4-sulfonyl)-1*H*-pyrrolo[2,3-*b*]pyridine (**14**).

[00198] A mixture of azaindole **3** (80 mg, 0.23 mmol), 2-chloropyridine-4-boronic acid (41 mg, 0.27 mmol), $Pd(Ph_3P)_4$ (20 mg, 0.11 mmol) and 2 M sodium carbonate (.34 mL, O.68 mmol) in 2 mL of DME was microwaved on high at 160°C for under nitrogen for 15 minutes. Diluted with ethyl acetate and the organic phase washed with water and brine then dried (Na_2SO_4) and concentrated in vacuo. The residue was subjected to flash chromatography (40% EtOAc/ 60% hexanes) to give 60 mg (68%) of the desired product **14**. ^{1}H NMR $CDCl_3$ 8.6 (d, 2H), 8.1 (m, 4H), 7.25 (m, 5H), 2.5 (s, 3H). The 2-chloropyridine could be displaced, for example, with benzylamine in a microwave reaction at 250°C for 30 min followed by deprotection to give compound **15**.

[00199] 1-(5-Bromo-1*H*-pyrrolo[2,3-*b*]pyridin-3-yl)-ethanone (**16**).

[00200] 7-Azaindole **13** (2.16g, 0.011 mol) was dissolved in 75 mL of dry DCM. Aluminium trichloride (4.36 g, 0.0327 mol) was added to the solution and the reaction mixture was stirred for 1 h at rt. Acetyl chloride (1.16 mL, 0.0164 mol) was adde dropwise to the mixture and stirred at rt for 18 h. 20 mL of methanol was added and the reaction for 1 h. Concentrated in vacuo and suspended in water-EtOAc mixture. Extraction with EtOAc and drying of the organic gave, after concentration in vacuo, 2.37 g (91%) of compound **14**. ^{1}H NMR $CDCl_3$ 9.5 (bs, 1H), 8.8 (s, 1H), 8.5 (s, 1H), 7.9 (s, 1H), 2.4 (s, 3H).

[00201] 1-[5-Bromo-1-(toluene-4-sulfonyl)-1*H*-pyrrolo[2,3-*b*]pyridin-3-yl]-3-dimethylamino-propenone (**17**).

[00202] To a suspension of NaH (263 mg, 0.0104 mol) in dry THF at 0°C was slowly added **16** (2.37g, 0.01 mol) in THF. Stirred for 0°C for 15 min then p-toluensulfonyl chloride (2.27 g, 0.012 mol) in THF was added and the reaction miture stirred for 18 h at rt. Quenched with water and extracted with ethyl acetate. The organic layer was dried and concentrated in vacuo to 3.37g of a solid residue that was used directly for the next step. The above intermediate was mixed with DMF-DMA (5.7 mL, 0.0428 mol) and heated at 100°C for 22 h. Concentrated in vacuo and subjected to flash chromatography (60% EtOAc/ 40% hexanes) to give 3.37 g (76%

from **17**) of the desired product **15**. ^{1}H NMR $CDCl_3$ 8.8 (s, 1H), 8.3 (s, 1H), 8.1 (s, 1H), 8.05 (d, 2H), 7.8 (d, 1H), 7.2 (s, 2H),5.7 (d, 1H), 3.0 (bs, 6H), 2.2 (s, 3H).

[00203] 6-(5-Bromo-1*H*-pyrrolo[2,3-*b*]pyridin-3-yl)-1*H*-pyrimidine-2-thione (**18**).

[00204] To a freshly prepared solution of sodium (175 mg, 0.0076 mol) in ethanol (18mL) were consecutively added compound **17** (1.0 g, 0.0022 mol) and thiourea (187 mg, 0.0025 mol). The mixture was heated to reflux for 4 h. The solvent was removed and the residue was dissolved in 8 mL of water. The solution was neutralized with 1 n HCl and extracted with ethyl acetate. The organic phase was dried and concentrated in vacuo to provide 570 mg (75%) of technically clean **18**. ^{1}H NMR $CDCl_3$ 12.2 (s, 1H), 8.6 (s, 1H), 8.3 (s, 1H), 8.2 (s, 1H), 7.6 (d, 1H), 5.8 (d, 1H).

[00205] 5-Bromo-3-(2-methylsulfanyl-pyrimidin-4-yl)-1-(toluene-4-sulfonyl)-1*H*-pyrrolo[2,3-*b*]pyridine (**19**).


[00206] A suspension of **18** (565 mg, 0.0018 mol), sodium acetate (526 mg, 0.0039 mol), methyl iodide (0.126 mL, 0.0020 mol) in 15 mL of ethanol-THF (9:1) was heated to reflux for 2 h. Cooled to rt and filtered. Chromatography (70% EtOAc-30% hexane) provided 48 mg (9%) of the thiomethyl intermediate. Tosylation provided 49 mg (70%) of the desired compound **19**. ^{1}H NMR $CDCl_3$ 8.8 (d, 1H), 8.5 (m, 2H), 8.35 (s, 1H), 8.1 (d, 2H), 7.25 (m, 3H), 2.7 (s, 3H), 2.3 (s, 3H).

[00207] Benzyl-[4-(5-bromo-1*H*-pyrrolo[2,3-*b*]pyridin-3-yl)-pyrimidin-2-yl]-amine (**20**).

[00208] This compound could be prepared from **19** via oxone oxidation and benzylamine displacement following Scheme 1.

[00209]

Scheme 3 Synthesis of 5-Substituted-azaindole pyrimidine and pyridine analogues



[00210] 5-Bromo-1-(*tert*-butyl-dimethyl-silanyl)–1*H*-pyrrolo[2,3-*b*]pyridine (**19**).

[00211] To a suspension of NaH (200 mg, 0.0078 mol) in 15 mL of dry DMF at 0°C was added azaindole **13** (1.36 g, 0.007 mol) in 5 mL of DMF. Stirred at 0°C for 1o min. and TBDMSCl in 3 mL of DMF was added and the resulting mixture stirred O.N. Taken in EtOAc and washed with water, brine. The organic layer was dried and concentrated in vacuo to an oil that was used directly for the next step without further purification. ^{1}H NMR $CDCl_3$ 8.3 (d, 1H), 7.9 (d, 1H), 7.2 (d, 1H), 6.4 (d, 1H), 0.85 (s, (H), 0.5 (s, 6H).

[00212] 1-(*tert*-Butyl-dimethyl-silanyl)-5-pyridin-3-yl-1*H*-pyrrolo[2,3-*b*]pyridine (**20**).

[00213] To a solution of **19** (1.77 g, 0.0057 mol), 3-diethylpyridyl borane (1,0 g, 0.0068 mol) in 80 mL of DME, was added $PdCl_2(Ph_3P)_2$ (320 mg, 0.455 mmol) and aqueous 2M sodium carbonate (8.5 mL, 0.017 mol) under nitrogen. The reaction mixture was refluxed for 3 h. Cooled to rt and diluted with ethyl acetate and washed with brine. The organic layer was dried and concentrated to an oil that was subjected to flash chromatography (30% EtOAc-70% hexanes) to give 1 g (57%) of the desired material **20**. ^{1}H NMR $CDCl_3$ 8.9 (s, 1H), 8.5 (d, 1H), 8.4 (d, 1H), 8.0 (d, 1H), 7.9 (d, 1H), 7.5 (d, 1H), 7.2 (d, 1H), 6.5 (d, 1H), 0.85 (s, 9H), 0.55 (s, 6H).

[00214] 3-Bromo-1-(*tert*-butyl-dimethyl-silanyl)–5-pyridin-3-yl-1*H*-pyrrolo[2,3-*b*]pyridine (**21**).

[00215] A solution of bromine (0.166 mL, 0.0032 mol) in 1 mL of CCl_4 was added dropwise to a stirred solution of **20** (1.0 g, 0.0032 mol) and pyridine (0.314 mL, 0.0039 mol) in dry $CHCl_3$ (30 mL) at 0°C. The reaction mixture was stirred at 0°C for 1 h and then neutralized with 10 mL of a mixture of sodium bicarbonate-sodium thiosulfate (1:1). The organic layer was separated and further extracted with dichloromethane (3X 10 mL) and the combined organics dried and concentrated in vacuo. The residue was subjected to flash chromatography (30% EtOAc-70% hexanes) to give 0.97 g (78%) of the desired material **21**. ^{1}H NMR $CDCl_3$ 8.9 (s, 1H), 8.55 (d, 1H), 8.4 (d, 1H), 8.0 (d, 1H), 7.9 (d, 1H), 7.4 (dd, 1H), 7.2 (s, 1H), 0.9 (s, 9H), 0.5 (s, 6H).

[00216] 5-Pyridin-3-yl-3-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-1-(toluene-4-sulfonyl)-1*H*-pyrrolo[2,3-*b*]pyridine (**22**).

[00217] Compound **21** could be desilylated with, for example, 10% HCl and tosylated with, for example, NaH and p-toluenesulfonyl chloride. Compound **22** could be prepared using the same protocol as for compound **4** in Scheme 1.

[00218] 3-(2-Methanesulfonyl-pyrimidin-4-yl)-5-pyridin-3-yl-1-(toluene-4-sulfonyl)-1*H*-pyrrolo[2,3-*b*]pyridine (**24**).

[00219] Compound **24** could be prepared using the same protocol as for compound **6** in Scheme 1.

[00220] 3-(6-Chloro-pyrimidin-4-yl)-5-pyridin-3-yl-1-(toluene-4-sulfonyl)-1*H*-pyrrolo[2,3-*b*]pyridine (**25**)


[00221] Compound **25** could be prepared using the same protocol as for compound **8** in Scheme 1.

[00222] 3-(2-Chloro-pyridin-4-yl)-5-pyridin-3-yl-1-(toluene-4-sulfonyl)-1*H*-pyrrolo[2,3-*b*]pyridine (**26**).

[00223] Compound **26** could be prepared using the same protocol as for compound **14** in Scheme 2.

[00224]

Scheme 4



[00225] 2-Methoxy-4-[1-(toluene-4-sulfonyl)-1*H*-pyrrolo[2,3-*b*]pyridin-3-yl]-benzonitrile (**29**).

[00226] In a tube was placed boronic acid **28** (35.4 mg, 200 μmol), and tosyl protected azaindole **3** (65.5mg, 194 μmol), with sodium carbonate (61.8mg, 583μmol) and tetrakis(triphenylphosphine)palladium(0) (8.3mg, 7.2 μmol). Water (323mg) and Ethylene glycol dimethyl ether (848mg) were added and the mixture was deoxygenated. The tube was sealed and heated to 160C for 10 minutes with magnetic stirring, utilizing microwave irradiation. The crude product was extracted with ethyl acetate and water. The organics were washed with brine, dried over sodium sulfate, filtered and concentrated to give crude product (83mg). The crude was purified via flash chromatography and eluted with a gradient from 1:1 ethyl acetate/hexane to 100% ethyl acetate to 4/4/1 ethyl acetate/hexane/7N ammonia in methanol to give **29** (42mg, 54%) and **30** (7.7 mg, 15%).

[00227] 2-Methoxy-4-(1*H*-pyrrolo[2,3-*b*]pyridin-3-yl)-benzonitrile (**30**).


[00228] The tosyl protecting group was removed from **29** by refluxing in dioxane (5ml) with 1N sodium hydroxide (200 μl, 2eq.) for 4.5 hours. The reaction was concentrated and extracted with ethyl acetate and saturated aqueous sodium bicarbonate. The organics were washed with brine, dried over sodium sulfate, filtered and concentrated to give **30** (30mg, 100%). ^{1}H NMR CD_3CN 10.00 (s, 1H), 8.32 (m, 2H), 7.83 (s, 1H), 7.65 (d, 1H), 7.40 (m, 2H), 7.20 (dd, 1H), 4.02 (s, 3H). LC/MS(M+1)=250

[00229] 2-Methoxy-4-(1*H*-pyrrolo[2,3-*b*]pyridin-3-yl)-benzamide (**31**).

[00230] The nitrile **30** (10mg, 40 μmol) was dissolved in dimethyl sulfoxide (0.5 ml). To this was added potassium carbonate (20mg) and 30% hydrogen peroxide in water (100 μl). The reaction was warmed to approximately 60C for 15 minutes. The reaction was concentrated to dryness and triturated in water. The resulting precipitate was filtered and dried in-vacuo overnight to give **31** (10mg, 93%). ^{1}H NMR DMSO-d6 11.99 (s, 1H), 8.32 (d, 1H), 8.24 (m, 1H), 8.01 (m, 1H), 7.88 (d, 1H), 7.59 (s, 1H), 7.40 (s, 1H), 7.37 (m, 2H), 7.15 (dd, 1H), 4.00 (s, 3H). LC/MS(M+1)=268.

[00231]

Scheme 5



[00232] Boronic acids **32B** through **32F** (500μmol) were placed in tubes with azaindole **3** (87 mg, 250 μmol). In each vial was placed sodium carbonate (53 mg, 500 μmol) and tetrakis(triphenylphosphine)palladium(0) (15 mg, 13 μmol). Water (1 ml) and Ethylene glycol dimethyl ether (2 ml) were added and the tubes were deoxygenated and sealed. The tubes were heated to 140 C via microwave, for 10 minutes. The reactions were cooled to 0 C and quenched with 2N hydrochloric acid

in water (1 ml). The reactions were extracted with ethyl acetate. The organics were washed with brine, dried over sodium sulfate, filtered and concentrated. The products were purified on silica (50% ethyl acetate/hexane to 100% ethyl acetate gradient) to give **33B** to **33F.**

[00233] The compounds **33B** to **33F** were dissolved in methanol (20 ml) and treated with 2N sodium hydroxide in water (1 ml) at room temperature, overnight. The reactions were neutralized with 2N hydrochloric acid (1 ml) and concentrated to dryness. The products were purified via chromatography on silica gel (50% ethyl acetate/hexane to 100% ethyl acetate gradient) or mass directed reverse phase chromatography on C18 (15% acetonitrile/water with 0.09% trifluoroacetic acid to 35% acetonitrile/water with 0.09% trifluoroacetic acid over 15 minutes) to give products **34B** to **34F.**

[00234] 3-(3-Benzyloxy-phenyl)-1*H*-pyrrolo[2,3-*b*]pyridine (**34B**).

[00235] ^{1}H NMR DMSO-d6 11.95 (s, 1H), 8.29 (m, 1H), 8.20 (d, 1H), 7.89 (s, 1H), 7.49 (d, 2H), 7.41 (t, 2H), 7.34 (m, 2H), 7.29 (m, 2H), 7.13 (dd, 1H), 6.90 (m, 1H), 5.20 (s, 2H). LC/MS(M+1)=301.

[00236] 3-(4-Benzyloxy-phenyl)-1*H*-pyrrolo[2,3-*b*]pyridine (**34C**).

[00237] ^{1}H NMR DMSO-d6 11.84 (s, 1H), 8.28 (m, 2H), 7.76 (s, 1H), 7.61 (d, 2H), 7.48 (d, 2H), 7.39 (t, 2H), 7.32 (t, 1H), 7.15 (dd, 1H), 7.09 (d, 2H), 5.13 (s, 2H). LC/MS(M+1)=301.

[00238] 3-(3,4-Dimethoxy-phenyl)-1*H*-pyrrolo[2,3-*b*]pyridine (**34D**).

[00239] ^{1}H NMR DMSO-d6 11.90 (s, 1H), 8.31 (d, 1H), 8.28 (d, 1H), 7.80 (s, 1H), 7.21 (m, 2H), 7.19 (dd, 1H), 7.02 (d, 1H), 3.86 (s, 3H), 3.78 (s, 3H). LC/MS(M+1)=255.

[00240] 3-(3,4,5-Trimethoxy-phenyl)-1*H*-pyrrolo[2,3-*b*]pyridine (**34E**).


[00241] ^{1}H NMR DMSO-d6 11.85 (s, 1H), 8.29 (d, 1H), 8.27 (d, 1H), 7.83 (s, 1H), 7.15 (d, 1H), 6.92 (s, 2H), 3.86 (s, 6H), 3.69 (s, 3H). LC/MS(M+1)=285.

[00242] 2-Methoxy-4-(1*H*-pyrrolo[2,3-*b*]pyridin-3-yl)-phenol (**34F**).

[00243] ^{1}H NMR DMSO-d6/D_2O 8.25-8.30 (m, 2H), 7.74 (s, 1H), 7.20 (d, 1H), 7.17 (dd, 1H), 7.10 (dd, 1H), 6.86 (d, 1H), 3.88 (s, 3H). LC/MS(M+1)=241.

[00244]

Scheme 6



[00245] 2-Methoxy-5-[1-(toluene-4-sulfonyl)-1*H*-pyrrolo[2,3-*b*]pyridin-3-yl]-benzaldehyde (**36**).

[00246] In a tube was placed boronate ester **35** (131 mg, 500 μmol), and **3** (175 mg, 500 μmol), with sodium carbonate (114 mg, 1.08 mmol) and tetrakis(triphenylphosphine)palladium(0) (22 mg, 19 μmol). Water (0.9 g) and Ethylene glycol dimethyl ether (2.1 g) were added and the mixture was deoxygenated. The tube was sealed and heated to 110C overnight, with magnetic stirring. The crude product was extracted with ethyl acetate and water. The organics were washed with

brine, dried over sodium sulfate, filtered and concentrated to give crude product. The crude was purified via flash chromatography and eluted with a gradient from 1:1 ethyl acetate/hexane to 100% ethyl acetate to 4/4/1 ethyl acetate/hexane/7N ammonia in methanol to give **36** (98 mg, 48%).

[00247] 2-Methoxy-5-[1-(toluene-4-sulfonyl)-1*H*-pyrrolo[2,3-*b*]pyridin-3-yl]-benzonitrile (**37**).

[00248] The aldehyde **36** (98 mg, 241 μmol) was dissolved in tetrahydrofuran (5 ml). To this was added 30% aqueous ammonium hydroxide (3 ml) and iodine (79 mg, 313 μmol). The reaction was stirred at room temperature overnight. The reaction was diluted with water and extracted with ether and ethyl acetate. The organics were washed with aqueous sodium sulfite (50 mg) in water (10ml). The organics were washed with brine, dried over sodium sulfate, filtered and concentrated to give **37** (95.8 mg, 98%).


[00249] 2-Methoxy-5-(1*H*-pyrrolo[2,3-*b*]pyridin-3-yl)-benzonitrile (**38**).

[00250] The nitrile **37** (95.8mg, 237 μmol) was dissolved in dioxane (5ml) and treated with 2N sodium hydroxide in water (250 μl, 500 μmol). The reaction was sealed and heated to reflux for 2 hours. The reaction was concentrated to dryness and extracted with ethyl acetate and water. The organics were washed with brine, dried over sodium sulfate, filtered and concentrated to give crude product (72.3 mg). The crude was purified via flash chromatography and eluted with a gradient from 1:1 ethyl acetate/hexane to 100% ethyl acetate to 4/4/1 ethyl acetate/hexane/7N ammonia in methanol to give **38** (26.7 mg, 45%).

[00251] 2-Methoxy-5-(1*H*-pyrrolo[2,3-*b*]pyridin-3-yl)-benzamide (**39**).


[00252] The de-tosylated nitrile **38** (26.7 mg, 107μmol) was dissolved in dimethyl sulfoxide (2.49g). To this was added potassium carbonate (32.7 mg, 237 μmol) and 30% hydrogen peroxide in water (85.5 mg) and the reaction was stirred at room temperature overnight. The reaction was diluted with water and the resulting precipitate was washed with water, acetonitrile and ether to give final product **39** (25.3 mg, 88%). ^{1}H NMR DMSO-d6 11.86 (s, 1H), 8.28 (m, 1H), 8.19 (d, 1H), 8.11 (m, 1H), 7.81 (m, 2H), 7.70 (s, 1H), 7.55 (s, 1H), 7.21 (d, 1H), 7.16 (m, 1H), 3.91 (s, 3H). LC/MS(M+1)=268

[00253] 4-Bromo-2-methoxybenzonitrile:



[00254] In a 1L round-bottomed flask, 53.94 g (270 mmol) of 4-bromo-2-fluorobenzonitrile was dissolved in 500 mL of THF. Sodium methoxide (21.99 g, 407 mmol) was added and the mixture was heated to reflux until TLC (SiO_2: CH_2Cl_2) showed complete consumption of starting material. The mixture was poured into 1N HCl and the THF was evaporated in vacuo. The remaining mixture was extracted with diethyl ether. The extract was dried ($MgSO_4$) and filtered over a plug of silica gel. The plug was eluted with CH_2Cl_2 and the filtrate was evaporated in vacuo to afford 45.56 g (80%) of the product as a white solid. ^{1}H NMR (500 MHz, $CDCl_3$) δ 7.42 (d, 1H), 7.17 (dd, 1H), 7.14 (d, 1H), 3.95 (s, 3H).


[00255] 4-Cyano-3-methoxyphenyl boronic acid (**28**).




[00256] A 3-necked 1L round-bottomed flask was equipped with an overhead stirrer, and a nitrogen line. The flask was charged with 45.56 g (215 mmol) of 4-bromo-2-methoxybenzonitrile, 64 mL (277 mmol) of tri-isopropylborate and 500 mL of THF. The solution was cooled to –78°C in a dry-ice/acetone bath. n-Butyllithium (2.5M, 110 mL, 275 mmol) was added dropwise via addition funnel. The mixture was stirred for 30 minutes and 2N HCl was added. The mixture was stirred for an hour and poured into water. The mixture was extracted with Et_2O. The organic solution was backextracted with 1N NaOH. The aqueous layer was washed with Et_2O and acidified with conc. HCl. The mixture was extracted with Et_2O. The organic extract was dried ($MgSO_4$) and evaporated in vacuo to afford 20.77 g (55%) of the product as a white solid. ^{1}H NMR (500 MHz, d6-DMSO) δ 7.68 (d, 1H), 7.59 (s, 1H), 7.47 (d, 1H), 3.95 (s, 3H).

[00257] **Schemes 7 and 8:**

Scheme 7:



Scheme 8:



[00258] 5-Chloro-3-iodo-pyridin-2-ylamine (40).

[00259] Iodine (16.28 g, 64 mmol) was added to a mixture of 1-amino-5-chloropyridine (8.25 g, 64 mmol) and silver sulfate (20 g, 64 mmol) in 400 mL of ethanol and the mixture was stirred at rt for 20 h. The mixture was filtered over celite and the solvent removed in vacuo. The residue was dissolved in DCM (600 mL) and washed with 5% aqueous NaOH (500 mL), water and brine. The organic layer was dried and concentrated in vacuo to solid residue that was subjected to flash chromatography (20% EtOAc-80%hexanes) to give 9.8 g (60%) of **40**. . ^{1}H NMR (500 MHz, $CDCl_3$) 7.9 (s, 1H), 7.7 (s, 1H), 5.0 (bs, 2H).

[00260] 5-Chloro-2-(triethyl-silanyl)-1*H*-pyrrolo[2,3-*b*]pyridine (**41**).

[00261] A mixture of **40** (9.5 g, 37.3 mmol) was dissolved in 320 mL of DMF and triethylsilylacetylene (20 mL, 112 mmol), Pd_2Cl_2 (dppf)$_2$ (1.52 g, 1.9 mmol), lithium chloride (1.58 g, 37.3 mmol)and 2M Na_2CO_3 (7.9 mL, 74.7 mL) were added and the mixture was stirred and refluxed at 90°C for 15h under nitrogen. Diluted with ethyl acetate-ether mixture (1:1) and the organic phase washed with water and brine then dried (Na_2SO_4) and concentrated in vacuo. The residue was subjected to flash chromatography (20% EtOAc/ 80% hexanes) to give 4.37 g (44%) of the desired product **41**. ^{1}H NMR (500 MHz, $CDCl_3$) 9.4 (bs, 1H), 8.2 (s, 1H), 7.9 (s, 1H), 6.6 (s, 1H), 0.9 (t, 9H), 0.75 (q, 6H).

[00262] 5-Chloro-1*H*-pyrrolo[2,3-*b*]pyridine (**42**).

[00263] Compound **41** (4.37 g, 0.0164 mol) was dissolved in THF. Molecular sieve (10g of 3A) was added followed by TBAF (32.75 mL, 0.0328). The reaction mixture was stirred for 5 h at rt. Diluted with ethyl acetate and washed several times with water, brine and the organic layer was dried and concentrated in vacuo to an oil that was subjected to flash chromatography (30% EtOAc-70% hexanes) to give 02.3 g (90%) of the desired material **42**. ^{1}H NMR DMSO d_6 11.8 (bs, 1H), 8.2 (s, 1H), 8.1 (s, 1H), 7.5 (s, 1H), 6.5 (s, 1H).

[00264] 1-(*tert*-Butyl-dimethyl-silanyl)-5-chloro-1*H*-pyrrolo[2,3-*b*]pyridine (**43**).

[00265] In a sealed tube equipped with a septa under nitrogen was added **42** (600 mg, 0.00396 mol) and 20 mL of dry THF and the solution cooled to 0°C. NaH (110 mg, 0.00435 mol) was added portionwise and after 15 min. of stirring at 0°C, TBSCl (656 mg, 0.00435 mol) was added. The septa was replaced by a Teflon screw cap and the sealed tube heated at 80°C for 3h. Cooled and neutralized with ammonium

chloride solution and extracted with hexanes. The organic phase was dried and concentrated in vacuo to an oil that was subjected to a short plug filtration (10% EtOAC-90% hexanes) to give 871 mg (82%) of compound **43**. ^{1}H NMR (500 MHz, $CDCl_3$) 8.2 (s, 1H), 7.8 (s, 1H), 7.25 (s, 1H), 6.4 (s, 1H), 0.9 (s, 9H), 0.6 (s, 6H).

[00266] Compound **43** can be brominated at C-3 as previously described in Scheme 3 for compound **21**.

[00267] Compound **19** can be lithiated with, for example, t-BuLi and quenched with DMF to provide the 5-formyl azaindole **44**. Compound **19** can also be treated with carbon moxoxide and methanol in the presence of a palladium catalyst , for example, $Pd(Ph_3P)_4$ to provide the 5-carbomethoxy azaindole **44**. Functionnal group interconversion to an acid, primary, secondary and tertiary amides such as **45** by standard transformations. Compound **44** can be homologated via , for example, a Wittig reaction to compounds like **49**.


[00268] Compound **19** can be cyanated with, for example, KCN in heating DMF in the presence of a catalyst such as, copper or palladium to provide the 5-cyano azaindole **46**.

[00269] Amination of compound **19** with, for example, and amine in the presence of a palladium catalyst can provide 5-aminated azaindoles such as **48**.

[00270] Compound **19** can be arylated or heterroarylated using, for example, Suzuki or Stille coupling to provide compound like **47**.


[00271] In another aspect, other synthetic schemes and compound syntheses are provided:

[00272] **Scheme 9**



[00273]

Example A




(1)

5-Chloro-3-iodo-pyridin-2-ylamine (1): A round bottom flask equipped with an air condenser was charged with 5-chloro pyridin-2-ylamine (26g, 0.2mol), acetic acid (78ml) and water (18ml). This was followed by dropwise addition of concentrated sulphuric acid (2.6ml), portionwise addition of periodic acid (9.5g, 0.04mol), and iodine (20g, 0.08mol). The reaction mixture was vigorously stirred at 80°C for 6h and then allowed to cool the room temperature.


The reaction mixture was poured onto ice (~700g). The pH of the suspension was adjusted to 8-9 with a 5M aqueous NaOH solution. A brown solid was filtered off and

solubilised in EtOAc (1.2l). The organic was washed with a saturated aqueous solution of $Na_2S_2O_3$, 1M NaOH solution and brine. The organic was dried over $MgSO_4$ and concentrated. The residue was recristallised from cyclohexane to afford an orange solid (42g, 80%). 1H NMR ($CDCl_3$): 4.8-5.1 (2H, brs), 7.9 (1H, s), 8.0 (1H, s)



(2)


5-Chloro-3-(trimethyl-silanylethynyl)-pyridin-2-ylamine (2): A 250ml round bottom flask was charged with 5-chloro-3-iodo-pyridin-2-ylamine (1) (42g, 165mmol), THF (100ml), copper iodide (315mg, 1.65mmol) and $PdCl_2(PPh_3)_2$ (1.15g, 1.65 mmol) under nitrogen. Triethylamine (70ml, 0.5mol), and trimethylsilyl acetylene (30ml, 0.21mol). The reaction mixture was stirred at room temperature for 2h. The reaction mixture was then cooled to 0°C and diethyl ether was added. The suspension was filtered through a celite and thoroughly washed with diethyl ether. The filtrate was concentrated and pre-absorbed onto silica gel and purified by column chromatography using as eluent, pentane/DCM 10% to 100%, to afford an off white solid (36g, 100%). 1H NMR ($CDCl_3$): 0.3 (9H, s), 5.0-5.1 (2H, brs), 7.6 (1H, s), 7.9 (1H, s). MS (ES+): 225, 227.



(3)


5-Chloro-1H-pyrrolo[2,3-b]pyridine (3): A solution of potassium tert-butoxide (36g, 320mmol) in N-methylpyrrolidone (70ml) was heated to 80°C under nitrogen. A solution of 5-chloro-3-(trimethyl-silanylethynyl)-pyridin-2-ylamine (2) (36g, 160mmol) in NMP (200ml) was added dropwise via a dropping funnel. The reaction mixture was stirred at 80°C for a further fifty minutes. The reaction mixture was allowed to cool to room temperature. Brine (500ml) was added to the reaction mixture and extracted with diethyl ether (5 x 200ml). The combined organics were washed with brine, dried over magnesium sulfate and concentrated in vacuo. The residue was

purified by column chromatography, using as eluent pentane/EtOAc 0% to 40%, and further recrystallised from cyclohexane to afford the title compound (10g, 41%). 1H NMR ($CDCl_3$) 6.5 (1H, s), 7.4 (1H, s), 8.0 (1H, s), 8.2 (1H, s), 10.4-10.6 (1H, brs). MS (ES+) 153.




(4)

3-Bromo-5-chloro-1H-pyrrolo[2,3-b]pyridine (4): A solution of bromine (3.5ml) in chloroform (40ml) was added dropwise to an ice-cold solution of the 5-chloro-1*H*-pyrrolo[2,3-*b*]pyridine (3) (10g, 65mM) in chloroform (260ml). The reaction mixture was stirred for 60 minutes at 0°C. The reaction mixture was then hydrolysed with water and the pH of the solution was adjusted to 10. The resulting solid was removed by filtration, and the . aqueous was extracted with dichloromethane. The organic was washed with water, dried over magnesium sulfate and concentrated in vacuo to afford the title compound (10.5g, 69%). 1H NMR (DMSO-d6) 7.8 (1H,s), 7.9 (1H, s), 8.3 (1H, s)




(5)

3-Bromo-5-chloro-1-(toluene-4-sulfonyl)-1H-pyrrolo[2,3-b]pyridine (5): Sodium hydride (2.2g, 54mmol) was added portionwise to an ice-cold solution of 3-bromo-5-chloro-1H-pyrrolo[2,3-b]pyridine (4) (10.5g, 45mmol) in dimethylformamide (70ml) under nitrogen. Thirty minutes later, tosyl chloride (8.7g, 46mmol) was added to the reaction mixture and the reaction mixture was stirred at room temperature for 18h. The reaction mixture was hydrolysed with water (~150ml) and a brown solid was obtained by filtration and dried in vacuo to afford the title compound (14,8g, 85%). 1H NMR ($CDCl_3$) 2.4 (3H, s), 7.3-7.4 (2H, d), 7.8-7.9 (2H. 2s), 8.1-8.2 (2H, d), 8.4 (1H, s); MS(ES+) 387




(6)

5-Chloro-3-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-1-(toluene-4-sulfonyl-1H-[2,3-b]pyridine (6): A 500ml round bottom flask was charged under nitrogen with 3-bromo-5-chloro-1-(toluene-4-sulfonyl)-1H-pyrrolo[2,3-b]pyridine (5) (6.9g, 18mmol), bispinacolato diboron (6.9g, 27mmol), $PdCl_2(dppf)_2$ (1.5g, 1.8mmol), potassium acetate (5.3g, 54mmol) and dimethoxyethane (100ml). The reaction mixture was stirred at 90°C for 18h. The reaction mixture was diluted with ethyl acetate (200ml) and washed with brine. The organic was dried over magnesium sulfate and concentrated in vacuo. The residue was purified by column chromatography, using as eluent pentane/EtOAc 0% to 20% and then triturated with pentane to afford the title compound (4g, 50%). 1H NMR(CDCl3) 1.4 (9H, s), 2.4 (3H, s), 7.2 (2H, d), 8.00-8.05 (2H, d), 8.10 (2H, s), 8.3 (1H, s). MS(ES+) 433



(7)

(6-Bromo-pyridin-2-yl)-(4-chloro-benzyl)-amine (7): A microwave vial was charged with 4-chloro benzylamine (700mg; 5 mmol; 5 equivalents) and 2,6-dibromo pyridine (238mg, 1mmol). The reaction mixture was stirred in the microwave at 150°C for three times ten minutes (200W maximum). The reaction mixture was diluted with diethyl ether (50ml), washed with 10% aqueous citric acid, saturated aqueous sodium bicarbonate and brine. The organic was dried over magnesium sulfate and, after filtration, concentrated in vacuo to afford an oil as the title compound (300 mg, 100% yield). LC/MS: 299 [M+H]



(8)

(4-Chloro-benzyl)-[6-(5-chloro-1H-pyrrolo[2,3-b]pyridin-3yl)-pyridin-2yl]-amine (8): A microwave vial was charged with (6-bromo-pyridin-2-yl)-(4-chloro-benzyl)-amine (300mg, 1mmol), 5-chloro-3-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-1-(toluene-4-sulfonyl-1H-[2,3-b]pyridine (7) (215mg, 0.5 mmol), tetrakis(triphenylphosphine)palladium (60mg, 0.05mmol), 2M sodium hydroxide (0.75ml) and dimethoxyethane (5ml). The suspension was degassed with nitrogen. The reaction mixture was stirred in the microwave at 130°C for ten minutes (200W maximum). It was then diluted with ethyl acetate (60ml), washed with brine twice, dried over magnesium sulfate and, after filtration, concentrated in vacuo. The compound was purified by flash chromatography (eluent: petroleum ether/ ethyl acetate 60/40) to afford 50 mg of the tosyl protected title compound. This residue was taken up in a mixture of methanol and tetrahydrofuran (1/3ml). 1M sodium hydroxide solution (1ml) was added to the reaction mixture, which was then stirred at room temperature for three hours.. The reaction mixture was then concentrated in vacuo and the residue was triturated with methanol. The suspension was filtered to afford the title compound (35mg, 10%). 1H NMR (DMSO-d6): 4.60-4.70 (2H, m), 6.35-6.40 (1H, d), 7.05-7.10 (1H, d), 7.20-7.25 (1H, t), 7.35-7.45 (4H, m), 8.15-8.20 (2H, m), 8.55-8.60 (1H, s), 12.5 (1H, s). LC/MS: 369 [M+H], 367 [M-H]

[00274] Table 3 below depicts data for certain exemplary compounds. Compound numbers correspond to those compounds depicted in Table 1. A "-" indicates that the measurement was not made.

Table 3

Cmpd #	LC_MASS_ PLUS	NMR_RESULT
1	-	8.9 1H, d; 8.35 1H, s; 8.25 1H, d; 8.1 1H,d; 7.2 1H, dd, 7.1 1H, d; 6.5 1H s
2	309.10	CDCl3 9.21 (m, 1H) 9.18 (s, 1H) 8.51 (s, 1H) ,8.16 (dd, J = 5.4, 1.0, 1H) 7.34 (dd, J = 8.0, 5.4, 1H),7.25 (m 1H) 7.19 (m, 1H) 7.03 (m, 1H) 6.85 (s, 1H)
3	324.00	MeOD (500 MHz) 8.76 (s, 1H) 8.45 (s, 1H) 8.43 (s, 1H),8.34 (s, 1H) 7.66 (m, 1H) 7.40 (m, 1H) 7.38 (m, 1H),7.27 (m, 2H)
4	288.00	MeOD,7.2 s(1H), 7.25 t(1H), 7.35 m(1H), 7.5 t(2H), 7.65 d(2H), 8.25 s(1H), 8.4 m(2H), 8.7 s(1H)

Cmpd #	LC_MASS_ PLUS	NMR_RESULT
5	310.00	MeOD,0.6 m(2H), 1.0 m(2H), 1.6 s(3H), 1.8 m(1H), 2.1 m(1H), 2.3 d(1H), 2.6 d(1H),4.3 s(1H), 6.0 m(2H), 6.95 s(1H), 7.05 m(1H), 7.5 m(2H)
6	212.00	MeOD,7.05 s(1H), 7.35 m (1H), 8.2 s(1H), 8.4 m(2H), 8.55 s(1H)
7	302.00	MeOD,4.8 s(2H), 7.05 s(1H), 7.2-7.6 m (6H), 8.2 s(1H), 8.4 m(2H), 8.55 s(1H)
8	240.10	1H 8.94, 1H 8.19, 1H 8.14, 2 H 7.25, 1H 7.02, 2H 3.52,,3H 1.31
9	337.00	MeOD,2.0 bs(2H), 2.1 s(3H), 3.6 m(2H), 3.8-4.3 bm(6H), 7.1 d(1H), 7.3 m(1H), 8.2 d(1H), 8.35 m(2H), 8.65 d(1H)
10	332.00	MeOD,3.8 s(3H), 4.7 bs(2H), 7.0 m(3H), 7.35 m(4H), 8.2 s(1H), 8.4 m(2H), 8.6 bs(1H)
11	308.00	MeOD,1.05 m(2H), 1.3 m(3H), 1.7 m(2H), 1.75-1.9 m(4H), 3.5 d(2H), 7.0 s(1H), 7.3 (1H), 8.1 s(1H), 8.35 d(1H), 8.4 d(1H), 8.6 s(1H)
12	310.00	MeOD1.4 m(2H), 1.7 m(2H), 2.0 m(1H), 2.65 s(1H), 3.45 m(2H), 3.6 m(1H), 4.0 d(2H), 7.0 s(1H), 7.35 m(1H), 8.2 s(1H), 8.35 d(1H), 8.45 m(1H), 8.6 s(1H)
13	240.00	MeOD;1.3 t(3H), 3.65 m(2H), 7.0 s(1H), 7.3 m(1H), 8.2 s(1H), 8.4 d(2H), 8.6 m(1H)
14	256.00	MeOD;3.7-3.9 dm(4H), 7.0 s(1H), 7.3 m(1H), 8.1 s(1H), 8.4 d(2H), 8.6 bs(1H)
15	360.10	^{1}H NMR 500MHz (DMSO-d6) 12.35 ppm, 1H, s;,11.02 ppm, 1H, s; 8.88 ppm, 1H, s; 8.63 ppm, 1H, d;,8.42 ppm, 1H, s; 8.33 ppm, 2H, m; 7.25 ppm, 2H, m;,6.97 ppm, 2H, m; 6.87 ppm, 1H, d; 3.78 ppm, 2H, s;,3.77 ppm, 3H, s.
16	302.00	DMSO d_6 8.7(bs, 1H); 8.4 (bs,1H); 8.25 (d, 1H); 8.1 (d, 1H); 7.4 (m, 2H); 7.25 (m, 3H); 7.2 (bs, 1H); 7.15 (bs, 1H); 4.7 (bs, 2H)
17	313.00	DMSO,7.2 q(2H), 8.3 mm(3H), 8.6 d(2H), 8.85 dd(2H), 9.1 s(1H), 12.35 bs(2H)
18	336.00	MeOD,4.8 s (2H), 7.0 s(1H), 7.3 bs(5H), 8.1 s(1H), 8.3 m(2H), 8.6 s(1H)
19	295.00	-
20	337.00	-
21	347.90	-
22	347.90	-
23	365.90	-
24	422.90	-
25	343.95	-
26	328.00	MeOD,3.1 bm(2H), 4.0-4.4 bd(2H), 5.0-5.3 bs(2H), 7.25 s(1H), 7.3-7.35 m(4H), 7.4 m(1H), 8.3 s(1H), 8.4 d(1H), 8.5 d(1H), 8.7 s(1H)

Cmpd #	LC_MASS_PLUS	NMR_RESULT
27	366.10	^{1}H NMR 500 MHz (DMSO-d6) 12.33 ppm, 1H, s;11.03 ppm, 1H, s; 8.86 ppm, 1H, s; 8.62 ppm, 1H, m;8.43 ppm, 1H, s; 8.29 ppm, 2H, m; 7.45 ppm, 2H, m;7.25 ppm, 1H, m; 7.20 ppm, 2H, m; 3.87 ppm, 2H, s; .
28	348.00	^{1}H NMR 500 MHz (DMSO-d6) 12.42 ppm, 1H, s;11.05 ppm, 1H, s; 9.72 ppm, 1H, s; 8.89 ppm, 1H, s;8.63 ppm, 1H, m; 8.43 ppm, 1H, s; 8.31 ppm, 2H, m;7.25 ppm, 6H, m; 1H, m; 3.87 ppm, 2H, s; 3.00 ppm, 3H, s.
29	423.10	^{1}H NMR 500 MHz (DMSO-d6) 12.33 ppm, 1H, s;11.00 ppm, 1H, s; 8.84 ppm, 1H, s; 8.59 ppm, 1H, m;8.45 ppm, 1H, s; 8.30 ppm, 2H, m; 7.30 ppm, 6H, m;3.83 ppm, 2H, s.
30	330.00	^{1}H NMR 500 MHz (DMSO-d6) 12.35 ppm, 1H, s;11.10 ppm, 1H, s; 8.88 ppm, 1H, s; 8.62 ppm, 1H, m;8.42 ppm, 1H, s; 8.31 ppm, 2H, m; 7.30 ppm, 5H, m;3.90 ppm, 2H, s;
31	348.00	^{1}H NMR 500 MHz (DMSO-d6) 12.35 ppm, 1H, s;11.10 ppm, 1H, s; 8.88 ppm, 1H, s; 8.62 ppm, 1H, m;8.42 ppm, 1H, s; 8.31 ppm, 2H, m; 7.30 ppm, 5H, m;3.90 ppm, 2H, s;
32	309.00	12.25 (s, 1H), 8.67 (d, 1H), 8.40 (s, 1H),,8.30 (d, 1H), 8.27 (d, 1H), 7.23 dd, 1H), ,7.09 (d, 1H), 4.63 (d, 2H), 3.30 (s, 2H), 2.80 (m, 2H), 2.45 (m, 1H), 1.07 (d, 6H),DMSO-d6,
33	351.00	12.28 (s, 1H), 8.72 (d, 1H), 8.42 (s, 1H),,8.31 (d, 1H), 8.28 (d, 1H), 7.25 (dd, 1H), ,7.14 (d, 1H), 4.71 (d, 2H), 4.10-4.70 (m, 2H),3.30 (s, 1H), 3.12 (s, 2H), 2.09 (s, 3H), ,1.21 (m, 6H),DMSO-d6,
34	308.00	8.90 (s, 1H), 8.60 (s, 1H), 8.39 (m, 1H), ,8.01 (m, 1H), 7.33 (m, 2H), 4.80 (s, 10H, solvent), ,3.55 (m, 1H), 3.30 (m, 14H, solvent), 1.92 (m, 2H), 1.81 (m, 3H), 1.72 (m, 1H),,1.25-1.40 (m, 3H), 1.08-1.25 (m, 2H),MeOH-d4,
35	385.00	MeOD,1.8-2.8 bm(2H), 2.35 bm(2H), 3.2 bm(1H), 3.6 (2H), 4.5-4.5 bs (3H), 7.0 bs(1H), 7.3 m(1H), 7.s(5H), 8.2 s(1H), 8.3 m(2H), 8.6 s(1H)
36	367.00	MeOD,1.25 t(3H), 1.55 q(2H), 2.1 d(2H), 3.1 m(2H), 4.15 m(4H), 4,6 m(1H), 7.0 s(1H), 7.3 m(1H), 8.2 s(1H), 8.3 d(1H), 8.4 d(1H), 8.6 s(1H)
37	337.00	12.32 (s, 1H), 9.3-10.5 (s, 4H), 9.01 (d, 1H),,8.51 (s, 1H), 8.48 (d, 1H), 8.19 (d, 1H), ,7.53 (m, 1H), 7.25 (d, 1H), 4.82 (m, 1/2H), ,4.25-4.53 (m, 2 1/2H), 3.81-3.93 (m, 1/2H), 3.58-3.71 (m, 1 1/2H), 3.14-3.58 (m, 1 1/2H), ,2.12 (s, 3H), 1.18-1.40 (m, 3H),CD3CN,
38	337.00	12.4 (s, 1H), 8.96 (d, 1H),,8.48 (s, 1H), 8.40 (d, 1H), 8.12 (d, 1H), ,7.49 (m, 1H), 7.20 (d, 1H), 4.8 (m, 1/2H), ,4.2-4.5 (m, 2 1/2H), 3.79-3.91 (m, 1/2H), 3.58-3.69 (m, 1 1/2H), 3.12-3.58 (m, 1 1/2H), ,2.1 (s, 3H), 1.12-1.35 (m, 3H),CD3CN,,
39	351.00	MeOD,1.4 s(1H), 1.7 t(2H), 2.3 d(2H), 7.0 s(1H), 7.3 m(1H), 8.1 s(1H), 8.3 d(1H), 8.35 d(1H), 8.6 s(1H)

Cmpd #	LC_MASS_PLUS	NMR_RESULT
40	378.90	12.9 (s, 1H); 9.1 (s, 1H); 8.9 (s, 1H); 8.8 (s, 1H); 8.7 (bs, 2H); 8.5 (d, 1H); 8.3 (d, 1H); 7.6 (bs, 1H); 7.4 (m ,6H); 4.7 (s, 2H)
41	335.90	$CDCl_3$ 8.7 (bs, 2H); 8.4 (bs, 2H); 8.3 (s, 1H); 7.4 (m, 5H); 4.8 (s, 2H)
42	362.00	10.94 (m, 1H), 8.96 (d, 1H), 8.71 (s, 1H), 8.55 (s, 1H), ,8.19 (dd, 1H), 7.24 (d, 1H), 4.5-6.0 (6H), 3.7-4.3 (m, 6H),,3.60 (m, 2H), 2.1 (s, 3H)
43	302.00	MeOD,4.95 bs(2H), 6.6 d(1H), 7.2 m(1H), 7.3-7.5 mm(5H), 8.0 d(1H), 8.3 d(1H), 8.4 s(1H), 8.55 d(1H)
44	230.10	1H 8.45 ,3H 8.35, 2H 7.82, 1H 7.25
45	250.00	10.00 (s, 1H), 8.32 (m, 2H), 7.83 (s, 1H), 7.65 (d, 1H),,7.40 (m, 2H), 7.20 (dd, 1H), 4.02 (s, 3H),,CD3CN
46	268.00	11.99 (s, 1H), 8.32 (d, 1H), 8.24 (m, 1H), ,8.01 (m, 1H), 7.88 (d, 1H), 7.59 (s, 1H),,7.40 (s, 1H), 7.37 (m, 2H), 7.15 (dd, 1H), ,4.00 (s, 3H),,DMSO-d6
47	268.00	-
48	238.00	11.98 (s, 1H), 8.32 (d, 1H), 8.26 (m, 1H), 8.15 (s, 1H),,8.00 (s, 1H), 7.91 (d, 1H), 7.82 (d, 1H), 7.71 (d, 1H), ,7.47 (t, 1H), 7.34 (s, 1H), 7.16 (dd, 1H),DMSO-d6
49	238.00	11.97 (s, 1H), 8.29 (d, 1H), 8.21 (d, 1H), 7.92 (d, 1H), ,7.85 (d, 2H), 7.71 (d, 2H), 7.17 (s, 1H), 7.10 (dd, 1H),DMSO-d6
50	292.00	11.97 (s, 1H), 8.48 (s, 1H), 8.29 (m, 2H), 8.00 (d, 1H),,7.88 (d, 1H), 7.71 (dd, 1H), 7.61 (s, 1H), 7.27 (d, 1H), ,7.21 (dd, 1H), 4.01 (s, 3H),DMSO-d6
51	301.00	-
52	301.00	-
53	255.00	-
54	285.00	-
55	241.00	-

[00275] Table 4 below depicts data for certain exemplary compounds. Compound numbers correspond to those compounds depicted in Table 2. A blank indicates that the measurement was not made.

Table 4

Cmpd No.	LC_MASS _PLUS M+1 (obs)	NMR_RESULT (1H NMR)	RT (mins)
1	380.00	12.68 (m, 1H), 9.20 (s, 1H), 8.82 (s, 1H), 8.62 (m , 1H), 8.28 (m, 1H), 8.05 (s, 1H), 7.35 (s, 1H), 7.25 (m, 1H), 3.6-4.2 (m, 6H), 3.45 (m, 2H), 1.7-2.1 (m, 5H) DMSO-d6	1.45
2	379.00		1.70
3	414.00		2.30
4	335.90		2.60
5	371.00		2.10
6		MeOD 5.0 s(2H), 7.1 s(1H), 7 .3 m(1H), 7.6 m(1H), 8.1 d(1H), 8.2 s(1H), 8.3-8.4m(2H), 8.55 d(1H), 8.6 s(1H), 8.7 s(1H)	
7		MeOD 5.1 s(2H), 7.25 m(1H), 7.35 m(1H), 7.9 d(2H), 8.25 s(1H), 8.4 d(2H),8.65 s(1H), 8.7 d(2H)	
8	385.00, 385.00	11.35 (s, 1H), 9.25 (s, 1H), 9.10 (s, 1H), 9.00 (s, 1H), 8.69 (d, 1H), 7.82 (d, 1H), 5.00 (m, 2H), 4.10 (m, 2H), 2.73 (s, 3H), 1.91 (m, 6H)DMSO-d6,12.5 (s, 1H), 8.70 (s, 1H), 8.50 (s, 1H), 8.31 (s, 1H), 8.29 (d, 1H), 7.10 (d, 1H), 4.66 (d, 2H), 4.2-4.6 (m, 2H), 3.15 (m, 2H), 2.09 (s, 3H), 1.20 (m 6H) DMSO-d6	2.02, 4.63
9	328.00	12.73 (s, 1H), 8.71 (s, 1H), 8.67 (s, 1H), 8.32 (s, 1H), 8.25 (d, 1H), 7.21 (m, 1H), 3.65-3.95 (m, 4H), 1.82 (m, 4H), 1.53 (m, 4H)DMSO-d6	2.35
10	346.00		2.23
11	328.00	12.75 (s, 1H), 8.71 (s, 1H), 8.68 (s, 1H), 8.35 (s, 1H), 8.25 (d, 1H), 7.22 (d, 1H), 4.62 (m, 2H), 3.10 (m, 2H), 1.75 (m, 3H), 1.20 (m, 2H), 0.91 (m, 3H)DMSO-d6	2.40
12	332.00	DMDO d6 12.6 (s, 1H); 9,5 (bs, 1H); 8.8 (s, 1H); 8.4 (s, 1H); 8.1 (s, 1H); 7.8 (bs, 1H): 7.4 (m ,5H); 7.1 (bs, 1H); 4.8 (d, 2H), 3.8 (s, 3H)	2.10
13	346.00	DMSO d6 12.8 (s, 1H); 9,4 (bs, 1H); 8.8 (s, 1H); 8.3 (s, 1H); 8.2 (s, 1H); 7.9 (bs, 1H): 7.4 (m ,5H); 7.2 (bs, 1H); 5.2 (bs, 1H), 3.8 (s, 3H); 1.5 (d, 3H)	2.20
14	346.00	DMSO d6 12.8 (s, 1H); 9,4 (bs, 1H); 8.8 (s, 1H); 8.3 (s, 1H); 8.2 (s, 1H); 7.9 (bs, 1H): 7.4 (m ,5H); 7.2 (bs, 1H); 5.2 (bs, 1H), 3.8 (s, 3H); 1.5 (d, 3H)	2.20
15	413.00	MeOD 1.2 s(3H), 1.9 m(1H), 2.2 m(1H), 2.55 m(1H), 2.6 s(3H), 3.2-3.4 m(2H), 3.55 m(1H), 3.8 m(1H), 4.5 s(2H), 5.5 bs(0.7H), 7.1 s(1H), 7 .35 m(1H), 7.5-7.6 mm(5H), 8.3 s(1H), 8.4-8.5 mm(2H), 8.7 s(1H)	1.31

Cmpd No.	LC_MASS _PLUS M+1 (obs)	NMR_RESULT (1H NMR)	RT (mins)
16	320.00	MeOD 4.9 s(2H), 7.0-7.6 mm(7H), 8_2 s(1H), 8.3 d(1H), 8.7 s(1H)	1.98
17		MeOD 1.2 d(3H), 1.9 m(1H), 2.2 m(1H), 2.4 s(1H), 2.6 m(3H), 3.0 s(1H)3.5 m(1H), 3.8 m(1H), 7.1 s(1H), 7.2 d(1H), 7.35 m(1H), 7.7 d(1H), 8.3 s(1H), 8.4 d(1H), 8.5 d(1H), 8.7 s(1H)	
18	380.00, 380.00	11.95 (s, 1H), 10.47 (s, 3H), 8.51 (s, 1H), 8.17 (s, 1H), 8.09 (s, 1H), 7.77 (t, 1H), 7.18 (d, 1H), 6.87 (d, 1H), 4.4-4.6 (s, 2H) 3.99 (s, 3H), 3.32 (m, 2H), 2.13 (s, 3H), 1.35 (m, 6H) DMSO-d6,9.82 (s, 1H), 8.32 (d, 1H), 8.08 (d, 1H), 7.93 (s, 1H), 7.58 (t, 1H), 7.20 (d, 1H), 6.66 (d, 1H), 4.2-4.7 (m, 4H), 3.91 (s, 3H), 3.12 (m, 2H), 2.10 (s, 3H), 1.35 (m, 6H) CD3CN	1.98, 2.02
19	384.00, 384.00, 384.00, 384.00, 384.00, 384.00, 384.00, 384.00, 384.00, 384.00, 384.00	10.75 (s, 1H), 8.46 (s, 1H), 8.39 (s, 1H), 8.10 (s, 1H), 7.91 (t, 1H), 7.19 (d, 1H), 7.02 (d, 1H), 6.25-6.75 (s, 9H), 4.4-4.6 (s, 2H) 4.10 (d, 2H), 3.49 (m, 2H), 2. 18 (s, 3H), 1.38 (d, 6H) DMSO-d6,good,good,good,11-12.2 (s, 2H), 10.6-11.0 (s, 1H), 8.54 (s, 1H), 8.35 (s, 1H), 8.05 (s, 1H), 7.69 (t, 1H), 7.11 (d, 1H), 6.89 (d, 1H), 4.2-4.7 (s, 2H), 4.18 (d, 2H), 3.35 (m, 2H), 2.17 (s, 3H), 1.38 (m, 6H) CD3CN,12.20 (s, 1H), 8.71 (s, 1H), 8.25 (d, 2H), 7.55 (t, 1H), 7.17 (d, 1H), 6.69 (d, 1H), 4.0-4.5 (s, 2H), 4.29 (d, 2H), 3.05 (m, 2H), 2.09 (s, 3H), 1.28 (m, 6H),12.24 (s, 1H), 8.73 (s, 1H), 8.27 (d, 2H), 7.58 (t, 1H), 7.18 (d, 1H), 6.71 (d, 1H), 4.2-4.5 (m, 14H), 3.08 (m, 2H), 2.32 (s, 3H), 2.09 (s, 3H), 1.28 (s, 6H) DMSO-d6,12.20 (s, 1H), 8.72 (s, 1H), 8.25 (d, 2H), 7.57 (t, 1H), 7.19 (d, 1H), 6.71 (d, 1H), 4.3-4.7 (s, 1H), 4.3 (d, 2H), 3.27 (s, 9H), 3.07 (m, 2H), 2.70 (q, 1H), 2.10 (s, 3H), 1.29 (s, 6H) DMSO-d6 DMSO-d6,12.31 (s, 1H), 8.70 (s, 1H), 8.29 (m, 2H), 7.65 (m, 1H), 7.20 (d, 1H0, 6.80 (d, 1H), 6.5-6.8 (s, 4H), 4.28 (d, 2H), 4.0-4.6 (s, 1H), 3.15 (m, 2H), 2.41 (s, 6H),good	2.39, 2.70, 3.26, 2.42, 3.22, 2.39, 3.35, 3.38, 3.32, 3.32, 3.32

Cmpd No.	LC_MASS _PLUS M+1 (obs)	NMR_RESULT (1H NMR)	RT (mins)
20	335.00	10.55 (s, 1H), 8.38 (m, 1H), 8.30 (m, 2H), 7.89 (t, 1H), 7.41 (m, 4H), 7.34 (m, 1H), 7.12 (d, 1H), 6.73 (d, 1H), 4.7-5.7 (s, 4H) 4.60 (s, 2H) CD3CN	2.39
21	336.00	MeOD 2.65 s(0.3H,DMSO), 4.9 s(2H), 7.1 s(1H), 7.3-7.5 m(6H), 8.2 s(1H), 8.4-8.5 m(2H), 8.7 s(1H)	2.20
22	370.00	MeOD 4.9 s(2H), 7.1 s(1H), 7.3 bs(1H), 7.6-7.8 m(5H), 8.2 s(1H), 8.4 m(2H), 8.675 s(1H)	2.60
23	378.00	MeOD 2.65 s(0.9H,DMSO)4.9 s(2H), 7.1 s(1H), 7.3-7.5 m(6H), 7.6-7.7 m(5H), 8.2 s(1H), 8.4 m(2H), 8.65 s(1H)	2.60
24	367.00	DMSO d6 12.5 (bs, 1H); 8.5 (bs, 1H); 8.2 (m, 2H); 8.0 (d, 1H); 7.3 (d, 1H); 4.0 (bs, 2H); 3.9 (s, 3H); 3.8 (bs, 2H); 3.5 (t, 2H); 2.5 (s, 3H); 1.9 (bs, 2H)	1.70
25	381.10, 381.40	DMSO d6 12.5 (bs, 1H); 8.4 (s, 1H); 8.3 (d, 1H); 8.25 (s, 1H); 8.1 (s, 1H); 7.3 (d, 1H); 4.7 (app d, 2H); 3.9 (s, 3H); 3.2 (bd, 2H); 2.1 (s, 3H); 1.2 (bs, 6H),DMSO d6 12.3 (bs, 1H); 8.5 (s, 1H); 8.25 (m, 2H); 8.1 (d, 1H); 7.2 (d, 1H); 4.65 (d, 4H); 3.9 (s, 3H); 3.2 (bd, 2H); 2.1 (s, 3H); 1.2 (bs, 6H)	2.00, 2.00
26	316.00	H1 MeOD: 1.3 m(1H), 3.4 s(3H), 5.1 s(2H), 7.0 s(1H), 7.1-7.4 m(7H), 8.1 s(1H), 8.3 d(1H), 8.6 s(1H)	2.10
27	320.00	H1 DMSO: 4.7 s(2H), 7.2 t(4H), 7.3 m(1H), 7.55 m(2H), 8.3 m(3H), 8.7 s(1H), 9.1-9.3 bs(0.75H)	2.00
28	317.00	H1 MeOD:1.8 bs(2H), 4.7 s(2H), 6.7-6.85 mm(4H)7.0-7.2 m(1H), 7.3 m(1H), 8.15 s(1H), 8.3 m(2H), 8.6 s(1H)	1.60
29	338.00	MeOD 4.9 s(2H), 6.9 m(1H), 7.0 m(2H), 7.2 s(1H), 7.3 m(1H), 8.2 s(1H), 8.4 m(2H), 8.65 s(1H)	2.00
30	403.20	CD3CN, 10.7 (bs, 1H); 8.15 (s, 1H); 8-8.1 (m, 3H); 7 (d, 1H); 3.8-4.05 (bs, 4H); 3.8 (s, 3H); 3.6 (bs, 2H); 3.3 (t, 2H); 2.8 (s, 3H); 2.4 (s, 2H)	1.90
31	417.20	CD3CN, 10.7 (s, 1H); 8.15 (s, 1H); 8 (m, 3H); 7 (d, 1H); 3.8-4 (bs, 4H); 3.85 (s, 3H); 3.6 (bs, 2H); 3.3 (m, 4H); 2.9 (m, 2H); 1.1 (m, 3H)	2.00
32	427.00	12.33 (s, 1H), 8.78 (d, 1H), 8.37 (d, 1H), 8.28 (d, 1H), 7.87 (s, 1H), 7.69 (d, 1H), 7.49 (s, 1H), 7.44 (d, 1H), 4.37 (s, 2H), 3.5-4.2 (m, 14H), 3.10 (m, 2H), 2.08 (s, 3H), 1.32 (m, 6H) DMSO-d6	2.24

Cmpd No.	LC_MASS _PLUS M+1 (obs)	NMR_RESULT (1H NMR)	RT (mins)
33	397.30	DMSO-d6 12.25 (1H, bs); 8.5 (1H, s); 8.3 (1H, d); 8.2 (1H, d); 8.05 (1H, d); 7.15 (1H, bs); 4.0 (4H, m); 3.9 (3H, s); 3.8-4.0 (4H, bm); 3.6-3.8 (2H, bs); 3.4 (2H, bs); 1.1 (3H, bm)	2.10
34	411.30	DMSO-d6 12.25 (1H, bs); 8.5 (1H, s); 8.3 (1H, d); 8.2 (1H, d); 8.05 (1H, d); 7.15 (1H, bs); 3.85 (3H, s); 3.7-4.0 (8H, bm); 3.6-3.8 (2H, bs); 3.4 (2H, bs); 1.5 (2H, bs); 0.8 (3H, bm)	2.30
35	408.30, 408.29	DMSOd6 12.4 (s, 1H); 9.5 (s,1H); 8.8 (s, 1H); 8.3 (s, 1H); 7.5 (dd,1H); 7.2 (d, 1H); 6.7 (d, 1H); 4.6 (bs, 2H); 4.3 (d, 2H); 3.9 (s, 3H); 3.1 (bd, 2H); 2.1 (s, 3H); 1.2 (bs, 6H)	2.90, 2.94
36	428.00	12.41 (s, 1H), 8.78 (s, 1H), 8.41 (s, 1H), 8.29 (s, 1H), 7.95 (d, 1H), 7.45 (d, 1H), 6.3-7.5 (s, 1H), 4.39 (s, 2H), 3.69 (m, 2H), 3.23 (m, 2H), 2.05 (s, 3H), 1.29 (m, 6H) DMSO-d6	2.38
37	465.40, 465.30	DMSOd6 11.9 (s, 1H); 9.4 (s, 1H); 9.2 (s, 1H); 8.2 (s, 1H); 7.5 (dd, 1H); 7.2 (d, 1H); 6.6 (d, 1H); 4.4 (bs, 4H); 3.1 (bs, 2H); 2.1 (s, 3H); 1.5 (s, 9H); 1.2 (bs, 6H)	3.10, 3.10
38	365.20	8.9 (s, 1H); 8.4 (s, 1H); 8.2 (s, 1H); 7.7 (dd, 1H); 7.2 (d, 1H); 6.9 (d, 1H); 4.4 (bs, 4H); 3.3 (bs, 2H); 2.2 (s, 3H); 1.4 (bs, 6H)	2.53
39	315.10	DMSO-d6: 12.3 (1H, bs); 8.3 (2H, m); 8.2 (1H, bs); 7.8 (1H, bs); 7.5 (6 H, m); 6.8 (1H, bs); 4.7 (2H, s); 4-4.5 (1H, bs); 2.3 (3H, s)	2.30
40	350.09	DMSO-d6: 11.9 (bs, 1H); 8.5 (d, 1H); 8.2 (s, 1H); 8.1 (s, 1H); 7.5 (t, 1H); 7.1 (t, 1H); 6.5 (d, 1H); 3.9-4.5 (bs, 1H); 4.0 (t, 1H); 3.7 (m, 4H); 3.4 (dt, 2H); 2.4 (s, 3H); 2.1 (s, 3H); 1.9 (m, 2H)	1.90
41	364.20	DMSO-d6: 11.9 (s, 1H); 8.5 (s, 1H); 8.2 (s, 1H); 8.1 (s, 1H); 7.6 (dd, 1H); 7.2 (d, 1 H); 6.7 (d, 1H); 4.5-5.4 (bs, 1H); 4.3 (d, 4H); 3.1 (bd, 2H); 2.4 (s, 3H); 2.1 (s, 3H); 1.3 (bs, 6H)	2.40
42	379.00	12.6-13.0 (s, 1H), 12.38 (s, 1H), 8.65 (s, 1H), 8.61 (m, 1H), 8.40 (m, 1H), 8.21 (m, 1H), 8.06 (d, 1H), 7.49 (d, 2H), 7.31 (t, 2H), 7.21 (t, 1H), 7.18 (d, 1H), 6.1-6.9 (s, 1H), 4.88 (m, 2H), 3.4-4.7 (s, 2H) DMSO-d6	3.18
43	393.00	12.39 (s, 1H), 8.60 (s, 1H), 8.50 (t, 1H), 8.41 (s, 1H), 8.22 (s, 1H), 8.09 (d, 1H), 7.39 (m, 2H), 7.29 (m, 2H), 7.20 (m, 1H), 6.3-6.8 (m, 3H), 4.90 (d, 2H), 3.85 (s, 3H) DMSO-d6	3.92
44	379.00	12.12 (s, 1H), 8.20 (s, 1H), 8.10 (s, 1H), 7.82 (m, 2H), 7.32 (m, 3H), 7.21 (m, 2H), 6.50 (d, 1H), 4.60 (s, 2H), 3.8-4.5 (s, 8H) DMSO-d6	2.20

Cmpd No.	LC_MASS _PLUS M+1 (obs)	NMR_RESULT (1H NMR)	RT (mins)
45	393.00	9.94 (s, 1H), 8.20 (d, 2H), 7.88 (d, 1H), 7.73 (s, 1H), 7.31-7.42 (m, 4H), 7.26 (t, 1H), 6.44 (d, 1H), 6.27 (m, 1H), 4.68 (d, 2H), 3.70 (s, 3H) CD3CN	2.52
46	378.00	mixture	2.94
47	394.27	DMSOD6 12.3 (s, 1H); 9.4 (s, 1H); 8.8 (s, 1H); 8.2 (s, 1H); 7.5 (dd, 1H); 7.2 (d, 1H); 6.7 (d, 1H); 4.4 (d, 4H); 3.1 (bd, 2H); 2.1 (s, 3H); 1.2 (bs, 6H)	2.47
48	319.20	DMSOd6 12.3 (bs, 1H); 8.4 (bd, 1H); 8.3 (d, 2H); 7.6 (bs, 1H); 7.5 (m, 4H); 7.2 (dd, 1H); 7.1 (bd, 1H); 6.5 (bs, 1H); 4.5 (s, 2H)	2.30
49	354.30	DMSOd6 12.2 (bs, 1H); 8.4 (app.t, 1H); 8.2 (m, 2H); 7.5 (dd, 1H); 7.1 (dd, 1H); 6.5 (bs, 1H); 3.9 (t, 1H); 3.8-3.6 (m, 4H); 3.4 (dt, 2H); 2.1 and 1.8 (s, 3H); 2.0 (t, 1H)	2.20
50	368.40	DMSO D6 12.1 (s, 1H); 8.5 (dd, 1H); 8.2 (s, 2H); 7.6 (dd, 1H); 7.2 (d, 1H); 6.8 (d, 1H); 4.3 (bd, 4H); 3.1 (dd, 2H); 2.1 (s, 3H); 1.2 (bs, 6H)	2.90
51	462.00	11.60 (s, 1H), 8.70 (s, 1H), 8.48 (s, 1H), 8.11 (s, 1H), 7.82 (t, 1H), 7.15 (d, 1H), 7.1-8.1 (s, 4H), 6.95 (d, 1H), 4.55 (s, 2H), 4.14 (m, 2H), 3.68 (m, 6H), 3.39 (m, 2H), 2.15 (s, 3H), 1.38 (m, 6H) CD3CN	1.84
52	490.00	11.29 (s, 1H), 8.90 (s, 1H), 8.50 (s, 1H), 8.61 (m, 2H), 8.11 (s, 1H), 7.68 (t, 1H), 7.12 (d, 1H), 6.78 (d, 1H), 4.0-4.5 (m, 12H), 3.40 (m, 2H), 3.27 (m, 4H), 2.15 (s, 3H), 1.25-1.45 (m, 12H)	1.89
53	436.40	DMSO d6 12.4 (s, 1H); 9.4 (s, 1H); 8.8 (s,1H); 8.3 (s,1H); 7.5 (t,1H); 7.2 (d, 1H); 6.7 (d, 1H); 5.2 (sept, 1H); 4.3 (bs, 4H); 3.1 (bs, 2H); 2.1 (s, 3H);1.3 (d, 6H0; 1.2 (bs, 6H)	3.40
54	461.00		
55	475.00		
56	462.00		
57	476.00		
58	475.00		
59	407.00		
60	421.00		
61	499.30	DMSO-d6: 11.9 ppm (s, 1H), 10.2 (s, 1H), 9.2 (s, 1H), 8.2 (s, 1H), 8.15 (s, 1H), 7.5 (t, 1H), 7.15-7.25 (dd, 4H), 7.15 (s, 1H), 6.7 (d, 1H), 4.2-4.8 (bs, 4H), 3.0 (bd, 2H), 2.5 (s, 6H), 2.3 (s, 3H), 2.0 (s, 3H)	3.40

Cmpd No.	LC_MASS _PLUS M+1 (obs)	NMR_RESULT (1H NMR)	RT (mins)
62	451.30	DMSO-d6: 11.9 ppm (s, 1H), 9.6 (s, 1H), 9.1 (s, 1H), 8.15 (s, 1H), 8.1 (s, 1H), 7.5 (t, 1H), 7.1 (d, 1H), 6.7 (d, 1H), 4.35 (m, 4H), 4.1 (t, 2H), 3.05 (bd, 2H), 2.1 (s, 3H), 1.7 (m, 2H), 1.3 (bs, 6H), 0.95 (t, 3H)	2.90
63	465.40	DMSO-d6: 11.9 ppm (s, 1H), 9.6 (s, 1H), 9.1 (s, 1H), 8.15 (s, 1H), 8.1 (s, 1H), 7.5 (t, 1H), 7.1 (d, 1H), 6.7 (d, 1H), 4.35 (m, 4H), 3.9 (d, 2H), 3.05 (bd, 2H), 2.1 (s, 3H), 1.9 (m, 1H), 1.3 (bs, 6H), 0.95 (d, 6H)	3.20
64	479.40	DMSO-d6: 11.9 ppm (s, 1H), 9.6 (s, 1H), 9.1 (s, 1H), 8.15 (s, 1H), 8.1 (s, 1H), 7.5 (t, 1H), 7.1 (d, 1H), 6.7 (d, 1H), 4.35 (m, 4H), 3.85 (s, 2H), 3.05 (bd, 2H), 2.1 (s, 3H), 1.3 (bs, 6H), 0.95 (s, 9H)	3.40
65	451.40	DMSO-d6: 11.9 ppm (s, 1H), 9.6 (s, 1H), 9.1 (s, 1H), 8.15 (s, 1H), 8.1 (s, 1H), 7.5 (t, 1H), 7.1 (d, 1H), 6.7 (d, 1H), 4.9 (q, 1H), 4.35 (m, 4H), 3.05 (bd, 2H), 2.1 (s, 3H), 1.3 (m, 12H)	2.90
66	423.30	DMSO-d6: 11.9 ppm (s, 1H), 9.6 (s, 1H), 9.1 (s, 1H), 8.15 (s, 1H), 8.1 (s, 1H), 7.5 (t, 1H), 7.1 (d, 1H), 6.7 (d, 1H), 4.35 (m, 4H), 3.7, (s, 3H), 3.05 (bd, 2H), 2.1 (s, 3H), 1.3 (bs, 6H)	2.40
67	428.00	12.31 (s, 1H), 8.71 (d, 1H), 8.41 (d, 1H), 8.29 (d, 1H), 7.56 (s, 1H), 7.07 (s, 1H), 4.0-4.8 (m, 4H), 3.3-3.9 (m, 20H), 3.13 (m, 2H), 2.10 (s, 3H), 1.16 (m, 6H)	3.20
68	427.00	12.54 (s, 1H), 8.50 (d, 1H), 8.35 (d, 1H), 8.31 (d, 1H), 8.09 (s, 1H), 8.07 (s, 1H), 4.0-5.4 (m, 9H), 3.3 (m, 2H), 2.09 (s, 3H), 1.26 (m, 6H)	3.70
69	485.30	DMSO d6 11.9 (s, 1H), 10.3 (s, 1H), 9.2 (s, 1H), 8.25 (s, 1H), 8.15 (s, 1H), 7.5 (t, 1H), 7.45 (t, 2H), 7.25 (d, 3H), 7.15 (d, 1H), 6.7 (d, 1H), 4.0-4.8 (bs, 4H), 3.0 (bd, 2H), 2.0 (s, 3H), 1.2 (bs, 6H)	3.20
70	437.30	DMSO d6 11.8 (s, 1H), 9.6 (s, 1H), 9.1 (s, 1H), 8.2 (s, 1H), 8.1 (s, 1H), 7.5 (t, 1H), 7.1 (d, 1H), 6.7 (d, 1H), 4.4 (m, 4H), 4.15 (q, 2H), 3.0 (bd, 2H), 2.0 (s, 3H), 1.25 (m, 9H)	3.20
71	515.30	DMSO d6 11.9 (s, 1H), 10.2 (s, 1H), 9.2 (s, 1H), 8.2 (s, 1H), 8.15 (s, 1H), 7.5 (t, 1H), 7.1 (m, 3H), 7.0 (d, 2H), 6.7 (d, 1H), 4.0-4.8 (bs, 4H), 3.8 (s, 3H), 3.0 (bd, 2H), 2.0 (s, 3H), 1.2 (m, 6H)	3.20
72	445.00	12.17 (s, 1H), 8.41 (d, 1H), 8.27 (d, 1H), 7.88 (d, 1H), 7.77 (s, 1H), 7.61 (s, 1H), 7.59 (s, 1H), 7.50 (s, 1H), 6.48 (s, 1H), 4.05 (m, 2H), 3.10 (m, 2H), 2.08 (s, 3H), 1.31 (m, 6H) DMSO-d6	2.84
73	422.40	DMSOd6 12.3 (bs, 1H); 9.3 (s, 1H); 8.8 (s, 1H); 8.2 (s, 1H); 7.5 (dd, 1H); 7.2 (d, 1H); 6.8 (d, 1H); 4.4 (q, 2H); 4.3 (bs, 4H); 3.1 (bs, 2H); 2.1 (s, 3H); 1.3 (t, 3H); 1.2 (bs, 6H)	3.20

Cmpd No.	LC_MASS _PLUS M+1 (obs)	NMR_RESULT (1H NMR)	RT (mins)
74	450.40	DMSOd6 12.3 (bs, 1H); 9.4 (s, 1H); 8.9 (s, 1H); 8.2 (s, 1H); 7.7 (dd, 1H); 7.2 (d, 1H); 6.8 (d, 1H); 4.4 (bd, 4H); 4.1 (d, 2H); 3.1 (bs, 2H); 2.1 (s, 3H); 2.05 (m, 1H); 1.3 (bs, 6H); 1.0 (d, 6H)	3.70
75	429.00	good	3.70
76	399.00	good	3.17
77	366.30	MeOD4 8.4 (s, 1H); 8.1 (s, 1H); 7.9 (s, 1H); 7.6 (dd, 1H); 7.2 (d, 1H); 6.8 (d, 1H); 4.3 (bs, 4H); 3.2 (bs, 2H); 2.2 (s, 3H); 1.3 (bs, 6H)	2.10
78	429.00		
79	423.00		
80	423.00		
81	399.00	12.43 (s, 1H), 8.48 (s, 1H), 8.33 (d, 1H), 8.06 (d, 1H), 7.59 (d, 1H), 6.87 (d, 1H),4.7-6.3 (m, 4H), 4.2-4.6 (m, 1H), 4.20 (m, 2H), 3.05 (m, 2H), 2.06 (s, 3H), 1.25 (m, 6H) DMSO-d6	2.21
82	368.50	DMSO d6 12.3 (s, 1H); 8.9 (s, 1H); 8.2 (s, 1H); 7.6 (s, 1H); 7.2 (s, 1H); 6.7 (s, 1H); 4.4 (d, 1H); 3.7-4.1 (m, 5H); 2.9 (s, 3H); 1.9-2.4 (m, 5H); 1.6 (s, 1H)	2.10
83	386.40	DMSO d6 12.2 ppm (s, 1H), 8.7 (s, 1H), 8.26 (s, 1H), 8.23 (s, 1H), 7.5 (t, 1H), 7.1 (d, 1H), 6.5 (d, 1H), 3.8 (m, 2H), 3.75 (t, 2H), 3.6 (m, 2H), 3.55 (s, 1H), 3.45 (s, 1H), 3.35 (m, 2H), 1.9 (m, 3H)	3.10
84	400.40	DMSO d6 12.2 ppm (s, 1H), 8.7 (s, 1H), 8.26 (s, 1H), 8.23 (s, 1H), 7.5 (t, 1H), 7.1 (d, 1H), 6.5 (d, 1H), 4.0 (m, 1H), 3.9 (m, 1H), 3.8 (m, 2H), 3.75 (m, 2H), 3.65 (m, 2H), 3.35 (m, 2H), 1.9 (m, 2H), 1.0 (dt, 3H)	3.30
85	369.90	DMSO-D6: 12.2 (1H, s); 8.7 (1H, s); 8.25 (2H, m); 7.6 (1H, dd); 7.25 (1H, d); 6.7 (1H, d); 3.8-3.5 (8H, m); 2.4 (2H, q); 1.05 (3H, t).	2.62
86	384.00	DMSO-D6: 12.2 (1H, s); 8.7 (1H, s); 8.25 (2H, m); 7.6 (1H, dd); 7.25 (1H, d); 6.7 (1H, d); 3.8-3.5 (8H, m); 2.4 (2H, q); 1.55 (2H, sextet); 0.95 (3H, t).	2.86
87	302.90	DMSO d6: 12.2 (s, 1H), 10.3 (s, 1H), 9.3 (d, 1H), 8.3 (d, 1H), 8.2 (d, 1H), 7.7 (t, 1H), 7.6 (m, 2H), 3.7 (s, 3H)	3.20
88	330.90	DMSO d6: 12.2 (s, 1H), 10.3 (s, 1H), 9.3 (s, 1H), 8.35 (s, 1H), 8.25 (s, 1H), 7.7 (t, 1H), 7.6 (d, 1H), 7.55 (d, 1H), 4.1 (t, 2H), 1.7 (m, 2H), 2 0 (t, 3H)	3.90
89	398.00	DMSO-D6: 12.2 (1H, s); 8.7 (1H, s); 8.25 (2H, m); 7.6 (1H, dd); 7.2 (1H, d); 6.7 (1H, d); 3.6-3.3 (8H, m); 2.3 (2H, d); 2.0 (1H, m); 0.9 (6H, d).	3.08

Cmpd No.	LC_MASS _PLUS M+1 (obs)	NMR_RESULT (1H NMR)	RT (mins)
90	344.80	DMSO d6: 12.2 ppm (s, 1H), 10.3 (s, 1H), 9.3 (s, 1H), 8.3 (s, 1H), 8.25 (s, 1H), 7.7 (t, 1H), 7.6 (d, 1H), 7.55 (d, 1H), 3.95 (d, 2H), 2.0 (m, 1H), 1.0 (d, 6H)	4.20
91	378.80	DMSO d6: 12.2 ppm (s, 1H), 10.4 (s, 1H), 9.3 (s, 1H), 8.3 (s, 1H), 8.2 (s, 1H), 7.7 (t, 1H), 7.62 (d, 1H), 7.57 (d, 1H), 7.5 (m, 2H), 7.4 (m, 2H), 5.2 (s, 2H)	4.20
92	409.00	12.56 (s, 1H), 8.73 (d, 1H), 8.52 (d, 1H), 8.32 (d, 1H), 8.05 (d, 1H), 7.52 (d, 1H), 4.0-4.7 (m, 4H), 3.24 (m, 2H), 2.09 (s, 3H), 1.36 (m, 6H) DMSO-d6	2.95
93	409.00	12.54 (s, 1H), 8.57 (d, 1H), 8.36 (d, 1H), 8.33 (d, 1H), 7.93 (d, 1H), 6.92 (d, 1H), 4.0-4.6 (m, 4H), 3.2 (m, 2H), 2.05 (s, 3H), 1.24 (m, 6H) DMSO-d6	2.82
94	437.00	DMSO d6 8.2 (s, 1H); 8.1 (s, 1H); 8.0 (s, 1H); 7.6 (dd, 1H); 7.1 (d, 1H); 6.8 (d, 1H); 4.5 (t, 2H); 4.4 (bs, 2H); 4.3 (d, 2H); 3.7 (bs, 2H); 3.1 (bs, 2H); 3.0 (s, 6H); 2.0 (s, 3H); 1.2 (bs, 6H)	1.59
95	479.00	DMSO d6 8.2 (s, 1H); 8.1 (s, 1H); 7.9 (s, 1H); 7.5 (dd, 1H); 7.0 (d, 1H); 6.6 (d, 1H); 4.7 (t, 2H); 4.4 (d, 2H); 4.0-3.8 (bs, 6H); 3.8 (t, 2H); 3.7-3.5 (bs, 4H); 3.1 (bs, 2H); 3.0 (s, 6H); 2.0 (s, 3H); 1.2 (bs, 6H)	1.63
96	436.00	DMSO d6 8.2 (s, 1H); 8.1 (s, 1H); 7.9 (s, 1H); 7.5 (dd, 1H); 7.0 (d, 1H); 6.6 (d, 1H); 4.7 (t, 2H); 4.4 (d, 2H); 4.0-3.8 (bs, 6H); 3.8 (t, 2H); 3.7-3.5 (bs, 4H); 3.1 (bs, 2H); 3.0 (s, 6H); 2.0 (s, 3H); 1.2 (bs, 6H)	2.25
97	427.00	12.29 (s, 1H), 8.73 (d, 1H), 8.30 (m, 2H), 8.09 (s, 1H), 7.59 (s, 1H), 7.52 (s, 1H), 7.06 (s, 1H), 4.1-4.8 (m, 4H), 3.1 (m, 2H), 2.09 (s, 3H), 1.25 (m, 6H) DMSO-d6	2.35
98	409.00	12.39 (s, 1H), 8.70 (d, 1H), 8.43 (d, 1H), 8.31 (s, 1H), 7.52 (s, 1H), 7.17 (s, 1H), 4.39 (m, 4H), 3.15 (m, 2H), 2.09 (s, 3H), 1.25 (m, 6H) DMSO-d6	3.04
99	300.90	CDCl3: 10.3 (s, 1H); 9.7 (s, 1H); 8.3 (s, 1H); 8.1 (s, 1H); 7.9 (s, 1H); 7.7 (dd, 1H); 6.8 (d, 1H); 6.5 (d, 1H); 3.05 (d, 2H); 2.0 (m, 1H); 1.0 (d 6H).	2.01
100	364.90	DMSO-D6: 12.3 (br s, 1H); 8.6 (s, 1H); 8.3 (s, 1H); 8.2 (s, 1H); 7.5 (m, 3H); 7.3 (m, 2H); 7.2 (m 1H); 7.1 (m, 1H); 6.5 (br s, 1H); 5.05 (br s 1H); 3.75 (m, 1H); 3.7 (m, 1H).	1.99
101	364.90	CDCl3: 10.6 (br s, 1H); 10.25 (br s, 1H); 8.3 (s, 1H); 8.1 (s, 1H); 7.9 (s, 1H); 7.6 (dd, 1H); 7.3 (m, 6H); 6.8 (d, 1H); 6.25 (d, 1H); 4.6 (m, 1H); 3.95 (m, 1H); 3.9 (m, 1H).	1.98

Cmpd No.	LC_MASS _PLUS M+1 (obs)	NMR_RESULT (1H NMR)	RT (mins)
102	302.90	DMSO-D6 (~2:1 rotational mixture): 12.6 (br s, 1H); 12.2 (br s, 1H); 8.7 (s, .33H); 8.6 (s, .67H); 8.3 (s, .67H); 8.25 (s, 1H); 8.2 (s, .33H); 7.8 (s, .67H); 7.5 (s, .33H); 7.15 (d, .67H); 7.05 (d, .33H); 6.8 (s, .67H); 6.45 (s, .33H); 4.5 (m, 1H); 4.1 (m, 1H); 3.6 (m, 2H); 1.3 (d, .6 H); 1.2 (d, 2.4H).	1.62
103	302.90	DMSO-D6 (~2:1 rotational mixture): 12.65 (br s, 1H); 12.25 (br s, 1H); 8.7 (s, .4H); 8.6 (s, .6H); 8.35 (s, .67H); 8.3 (s, 1H); 8.25 (s, .33H); 7.8 (s, .67H); 7.5 (s, .33H); 7.2 (d, .67H); 7.1 (d, .33H); 6.8 (s, .67H); 6.5 (s, .33H); 4.5 (m, 1H); 4.05 (m, 1H); 3.5 (m, 2H); 1.3 (d, .6 H); 1.2 (d, 2.4H).	1.63
104	272.80	DMSO d6 12.3 (s, 1H); 8.4 (s, 1H); 8.3 (s, 1H); 8.1 (s, 1H); 8.0 (d,1H); 7.8 (d, 1H).	2.61
105	271.90	DMSO d6 12.2 (s, 1H); 8.4 (s, 1H); 8.3 (s, 1H); 8.1 (s, 1H); 7.9 (d,1H); 7.8 (d, 1H).	2.26
106	273.80	DMSO d6 12.4 (s, 1H); 9.2 (s, 1H); 8.9 (s, 1H); 8.5 (s, 1H); 8.3 (s, 1H); 8.2 (d,1H); 8.0 (d, 1H).	2.34
107	272.90	DMSO d6 12.3 (s, 1H); 9.1 (s, 1H); 8.9 (s, 1H); 8.5 (s, 1H); 8.3 (s, 1H); 8.1 (d,1H); 8.0 (bs, 1H); 7.9 (d, 1H); 7.5 (bs, 1H).	2.01
108	506.00	DMSO-d6: 12.2 (s, 1H); 9.6 (br s, 1H); 9.25 (s, 1H); 8.8 (s, 1H); 8.25 (s, 1H); 7.6 (dd, 1H); 7.2 (d, 1H); 6.7 (d, 1H); 4.4 (br s, 4H); 4.0 (m, 2H); 3.8-3.6 (m, 6H); 3.35 (m, 2H); 3.2 (br s, 2H); 3.1 (br s, 2 H); 2.1 (s, 3H); 1.3 (s, 6H).	1.59
109	533.00	DMSO-d6: 12.2 (s, 1H); 9.2 (s, 1H); 8.75 (s, 1H); 8.6 (br s, 1H); 8.2 (s, 1H); 7.6 (dd, 1H); 7.2 (d, 1H); 6.75 (d, 1H); 4.4 (br s, 4H); 3.6-2.7 (m, 17H); 2.1 (s, 3H); 1.9 (m, 2H); 1.3 (s, 6H).	1.46
110	452.00	12.09 (s, 1H), 8.28 (d, 1H), 8.20 (d, 1H), 7.77 (d, 1H), 7.11 (d, 1H), 6.97 (d, 1H), 4.0-4.7 (m, 4H), 3.30 (s, 8H), 3.18 (m, 2H), 2.10 (s, 3H), 1.28 (m, 6H) DMSO-d6	2.31
111	452.00	12.32 (s, 1H), 8.76 (d, 1H), 8.37 (d, 1H), 8.31 (d, 1H), 7.71 (s, 1H), 7.27 (s, 1H), 4.0-4.7 (m, 4H), 3.30 (s, 6H), 3.20 (m, 2H), 2.09 (s, 3H), 1.29 (m, 6H) DMSO-d6	2.59
112	366.00	DMSOd6 12.2 (bs, 1H); 8.2 (s, 1H); 8.1 (s, 1H); 8.0 (s,1H); 7.6 (dd, 1H); 7.1 (dd, 1H); 6.5 (dd, 1H); 4.0 (m, 1H); 3.9 (s, 3H); 3.8 (m, 2H); 3.75 (m, 1H); 3.7 (m, 2H); 3.5 (m, 2H); 2.0 (s, 3H); 1.9 (m, 2H)	1.90
113	407.00	DMSO d6: 11.8 ppm (s, 1H), 10 (s, 1H), 9.2 (s, 1H), 8.2 (s, 1H), 8.1 (s, 1H), 7.5 (t, 1H), 7.1 (d, 1H), 6.7 (d, 1H), 4.4 (m, 4H), 3.05 (m, 2H), 2.1 (s, 3H), 2.08 (s, 3H), 1.3 (bs, 6H)	2.10

Cmpd No.	LC_MASS _PLUS M+1 (obs)	NMR_RESULT (1H NMR)	RT (mins)
114	421.00	DMSO d6: 11.8 ppm (s, 1H), 9.9 (s, 1H), 9.3 (s, 1H), 8.2 (s, 1H), 8.1 (s, 1H), 7.5 (t, 1H), 7.1 (d, 1H), 6.7 (d, 1H), 4.4 (m, 4H), 3.0 (m, 2H), 2.4 (m, 2H), 2.1 (s, 3H), 1.3 (bs, 6H), 1.1 (t, 3H)	2.20
115	491.00	DMSO d6: 11.9 ppm (s, 1H), 10 (s, 1H), 9.3 (s, 1H), 8.6 (bs, 1H), 8.2 (s, 1H), 8.15 (s, 1H), 7.5 (t, 1H), 7.1 (d, 1H), 6.7 (d, 1H), 4.4 (m, 4H), 3.4 (bs, 2H), 3.2 (bs, 4H), 3.0 (m, 2H), 2.9 (bs, 4H), 2.1 (s, 3H), 1.3 (bs, 6H)	1.50
116	492.00	DMSO d6: 12 ppm (s, 1H), 10.7 (bs, 1H), 9.3 (s, 1H), 8.22 (s, 1H), 8.19 (s, 1H), 7.5 (t, 1H), 7.2 (d, 1H), 6.7 (d, 1H), 4.4 (m, 4H), 4.2 (bs, 2H), 3.8 (t, 4H), 3.1 (m, 4H), 3.0 (m, 2H), 2.1 (s, 3H), 1.3 (bs, 6H)	1.60
117	435.00	DMSO d6: 11.8 ppm (s, 1H), 9.9 (s, 1H), 9.3 (s, 1H), 8.2 (s, 1H), 8.1 (s, 1H), 7.5 (t, 1H), 7.1 (d, 1H), 6.7 (d, 1H), 4.4 (m, 4H), 3.0 (m, 2H), 2.6 (m, 1H), 2.1 (s, 3H), 1.3 (bs, 6H), 1.1 (d, 6H)	2.40
118	419.00	DMSO d6: 11.9 ppm (s, 1H), 10.2 (s, 1H), 9.4 (s, 1H), 8.2 (s, 1H), 8.1 (s, 1H), 7.5 (t, 1H), 7.1 (d, 1H), 6.7 (d, 1H), 6.5 (m, 1H), 6.3 (d, 1H), 5.8 (d, 1H), 4.4 (m, 4H), 3.0 (m, 2H), 2.1 (s, 3H), 1.3 (bs, 6H)	2.30
119	493.00	DMSO d6: 11.8 ppm (s, 1H), 10.0 (s, 1H), 9.3 (s, 1H), 8.2 (s, 1H), 8.1 (s, 1H), 7.5 (t, 1H), 7.1 (d, 1H), 6.7 (d, 1H), 4.4 (m, 4H), 4.0 (m, 4H), 3.0 (m, 2H), 2.6 (m, 2H), 2.1 (s, 3H), 1.3 (bs, 6H), 1.2 (t, 3H)	2.50
120	452.00	8.79 (d, 1H), 8.46 (s, 1H), 8.26 (d, 1H), 8.07 (s, 1H), 7.69 (d, 1H), 7.40 (d, 1H), 5.71 (s, 2H), 4.91 (s, 2H), 4.20 (s, 2H), 3.70 (s, 4H), 3.40 (m, 2H), 3.25 (s, 1H), 3.12 (s, 1H), 3.03 (s, 1H), 2.95 (m, 2H), 1.99 (s, 3H), 1.09 (m, 6H) CD3CN with 4 drops of DMSO-d6	2.43
121	506.30	DMSO d6: 11.9 ppm (s, 1H), 10.3 (S, 1H), 9.3 (S, 1H), 8.2 (s, 1H), 8.1 (s, 1H), 7.5 (t, 1H), 7.1 (d, 1H), 6.7 (d, 1H), 4.4 (m, 4H), 4.0 (d, 2H), 3.7 (t, 2H), 3.5 (m, 4H), 3.1 (m, 4H), 3.0 (t, 2H), 2.1 (s, 3H), 1.3 (bs, 6H)	1.70
122	272.90	DMSO-d6: 12.2 (br s, 1H); 8.8 (s, 1H); 8.25 (m, 2H); 7.55 (dd, 1H); 7.1 (d, 1H); 6.5 (d, 1H); 3.1 (s, 6H).	2.05
123	315.00	DMSO-d6: 12.2 (br s, 1H); 8.8 (s, 1H); 8.25 (m, 2H); 7.5 (dd, 1H); 7.1 (d, 1H); 6.45 (m, 1H); 3.5 (d, 2H); 3.1 (s, 3H); 2.15 (m, 1H); 0.95 (d, 6H).	3.55
124	492.00	DMSO-d6: 12.2 (br s, 1H); 9.7 (br s, 1H); 9.2 (m, 1H); 8.8 (s, 1H); 8.2 (s, 1H); 7.5 (m, 1H); 7.1 (m, 1H); 6.5 (m, 1H); 4.1-3.5 (m, 14H); 3.4 (m, 4H); 3.2 (br s, 2H); 2.0-1.8 (m, 5H).	2.24

Cmpd No.	LC_MASS _PLUS M+1 (obs)	NMR_RESULT (1H NMR)	RT (mins)
125	519.10	DMSO-d6: 12.2 (br s, 1H); 9.2 (d, 1H); 8.8 (s, 1H); 8.65 (br s, 1H); 8.2 (s, 1H); 7.5 (m, 1H); 7.1 (m, 1H); 6.5 (dd, 1H); 4.1-3.1 (m, 18H); 3.0 (br s, 4H); 2.8 (s, 3H); 2.0-1.8 (m, 5H).	1.75
126	379.90	DMSO-d6: 11.8 (br s, 1H); 9.2 (s, 1H); 8.8 (s, 1H); 8.0 (s, 1H); 7.5 (dd, 1H); 7.1 (d, 1H); 6.5 (dd, 1H); 4.0 (m, 1H); 3.8-3.65 (m, 5H); 3.4 (m, 2H); 2.0-1.8 (m, 5H).	1.83
127	477.90	DMSO d6: 11.8 ppm (s, 1H), 9.0 (s, 1H), 8.6 (s, 1H), 8.2 (s, 1H), 8.1 (s, 1H), 7.5 (t, 1H), 7.1 (d, 1H), 6.7 (d, 1H), 4.3 (m, 4H), 3.6 (t, 4H), 3.5 (t, 4H), 3.0 (m, 2H), 2.1 (s, 3H), 1.2 (bs, 6H)	2.10
128	505.00	DMSO d6: 11.9 ppm (s, 1H), 10.3 (s, 1H), 9.3 (s, 1H), 8.2 (s, 1H), 8.1 (s, 1H), 7.5 (t, 1H), 7.1 (d, 1H), 6.7 (d, 1H), 4.4 (m, 4H), 3.3 (m, 12H), 3.0 (m, 2H), 2.8 (m, 2H), 2.1 (3H), 1.3 (bs, 6H)	1.50
129	355.10	MeOD-D4: 8.4 (s, 1H); 8.35 (s, 1H); 8.1 (s, 1H); 8.0 (dd, 1H); 7.2 (d, 1H); 7.0 (d, 1H); 3.75 (m, 1H); 1.9-1.8 (m, 3H); 1.7 (m, 1H); 1.55 (m, 1H); 1.3-1.1 (m, 9H).	2.40
130	315.10	MeOD-D4: 8.4 (s, 1H); 8.35 (s, 1H); 8.15 (s, 1H); 8.0 (dd, 1H); 7.2 (d, 1H); 7.0 (d, 1H); 3.8 (m, 1H); 1.9 (m, 1H); 1.3 (d, 3H); 1.0 (m, 6H).	2.00
131	367.00	MeOD-D4: 8.3 (m, 2H); 8.1 (s, 1H); 7.9 (dd, 1H); 7.5 (m, 2H); 7.2 (d, 1H); 7.1 (m, 2H); 6.75 (d, 1H); 5.0 (m, 1H); 1.65 (d, 3H).	2.40
132	342.80	DMSO-D6: 12.5 (br s, 1H); 8.7 (br s, 1H); 8.35-8.15 (m, 2H); 7.7 (br s, 1H); 7.1 (m, 1H); 6.8 (br s, 1H); 3.8 (br s, 1H); 3.7-3.5 (m, 2H); 2.0 (m, 1H); 1.25-0.8 (m, 8H).	2.00
133	360.90	DMSO-D6: 12.3 (br s, 1H); 8.8 (br s, 1H); 8.25 (m, 2H); 7.6 (br s, 1H); 7.4-7.1 (m, 5H); 6.5 (br s, 1H); 5.6 (br s, 1H); 3.0 (m, 1H); 2.9 (m, 1H); 2.6 (m, 1H); 1.95 (m, 1H).	3.00
134	330.80	DMSO-D6: 12.5 (br s, 1H); 8.7 (br s, 1H); 8.3-8.25 (m, 2H); 7.6 (br s, 1H); 7.2 (br s, 1H); 6.7 (br s, 1H); 3.8 (br s, 1H); 3.65-3.5 (m, 3H); 2.0 (m, 1H); 1.0 (m, 7H).	2.00
135	330.80	DMSO-D6: 12.5 (br s, 1H); 8.6 (br s, 1H); 8.3-8.2 (m, 2H); 7.7 (br s, 1H); 7.1 (br s, 1H); 6.8 (br s, 1H); 3.8 (br s, 1H); 3.7-3.5 (m, 3H); 2.0 (m, 1H); 1.0 (m, 7H).	2.00
136	376.90	DMSO-D6: 12.5 (br s, 1H); 8.7 (br s, 1H); 8.3 (m, 2H); 7.4-7.1 (m, 5H); 5.5 (br s, 1H); 4.7 (m, 1H); 3.2 (m, 1H); 2.9 (m, 1H).	2.40
137	376.90	DMSO-D6: 12.1 (br s, 1H); 8.8 (br s, 1H); 8.2 (m, 2H); 7.45-7.05 (m, 5H); 6.5-6.4 (m, 2H); 5.5 (br s, 1H); 4.7 (m, 1H); 3.15 (m, 1H); 2.9 (m, 1H).	2.40

Cmpd No.	LC_MASS _PLUS M+1 (obs)	NMR_RESULT (1H NMR)	RT (mins)
138	430.00	DMSO-d6 11.9 ppm (s, 1H), 10.2 (s, 1H), 9.3 (s, 1H), 8.3 (s, 1H), 8.1 (s, 1H), 7.5 (m, 1H), 7.1 (m, 1H), 6.5 (m, 2H), 6.3 (d, 1H), 5.8 (d, 1H), 3.5-4.0 (m, 10H), 1.9 (m, 2H)	2.30
139	405.10	DMSO-d6 11.9 ppm (s, 1H), 10.2 (s, 1H), 9.3 (s, 1H), 8.3 (s, 1H), 8.1 (s, 1H), 7.5 (m, 1H), 7.1 (m, 1H), 6.5 (m, 2H), 6.3 (s, 1H), 5.8 (s, 1H), 4.0 (t, 1H), 3.8 (dt, 2H), 3.7 (m, 3H), 3.4 (m, 2H), 2.0 (s, 1H), 1.9 (t, 1H), 1.8 (s, 3H)	2.00
140	393.10	DMSO-d6 11.9 ppm (s, 1H), 10.0 (s, 1H), 9.1 (s, 1H), 8.2 (s, 1H), 8.1 (s, 1H), 7.5 (m, 1H), 7.1 (m, 1H), 6.5 (m, 2H), 4.0 (t, 1H), 3.8 (dt, 2H), 3.7 (m, 3H), 3.4 (m, 2H), 2.1 (s, 3H), 2.0 (s, 1H), 1.9 (t, 1H), 1.8 (s, 3H)	1.70
141	407.00	DMSO-d6 11.8 ppm (s, 1H), 9.9 (s, 1H), 9.1 (s, 1H), 8.2 (s, 1H), 8.1 (s, 1H), 7.5 (m, 1H), 7.1 (m, 1H), 6.5 (m, 2H), 4.0 (t, 1H), 3.8 (dt, 2H), 3.7 (m, 3H), 3.4 (m, 2H), 2.4 (q, 2H), 2.0 (s, 1H), 1.9 (t, 1H), 1.8 (s, 3H), 1.1 (t, 3H)	1.90
142	334.80	DMSOD6 12.2 (bs, 1H); 8.3 (s, 1H); 8.15 (s, 1H); 8.0 (s, 1H); 7.6 (d, 2H); 7.5 (dd, 2H); 7.4 (dd, 2H); 7.3 (d, 2H); 6.95 (d, 1H); 5.1 (s, 2H)	4.60
143	287.80	DMSO D6 12.1 (bs, 1H); 8.3 (s, 2H); 7.9 (s, 1H); 7.55 (d, 1H); 7.4 (dd, 2H); 7.2 (d, 1H); 2.95 (hept, 1H); 1.25 (d, 6H)	4.60
144	271.90	DMSO D6 12.1 (bs, 1H); 8.3 (s, 2H); 7.9 (s, 1H); 7.3 (d, 1H); 7.2 (bs, 2H); 6.8 (app d, 1H); 2.9 (s, 6H)	2.60
145	333.80		4.10
146	382.90	DMSO d6 12.1 (bs, 1H); 8.3 (s, 1H); 8.25 (s, 1H); 8.0 (s, 1H); 7.3 (dd, 1H); 7.2 (m, 2H); 6.9 (s, 1H); 4.2 (bs, 2H); 3.6 (d, 2H); 2.8 (bs, 2H); 2.0 (s, 3H); 1.3 (bs, 6H)	3.70
147	378.10		3.00
148	288.10		2.00
149	395.10	DMSO-D6: 12.4 (s, 1H); 8.7 (d, 1H); 8.4 (s, 1H); 8.25 (s, 1H), 7.5 (s, 1H); 6.9 (d, 1H); 4.0-3.65 (m, 6H); 3.4 (m, 2H); 2.0-1.8 (m, 5H).	3.30
150	326.00	DMSO-D6: 12.3 (s, 1H); 8.9 (s, 1H); 8.4 (s, 1H); 8.25 (s, 1H), 7.35 (s, 1H); 7.25 (m, 1H); 6.6 (s, 1H); 3.2 (m, 2H); 2.0 (m, 1H); 1.0 (d, 6H).	3.70
151	353.00	DMSO-D6: 12.4 (s, 1H); 8.75 (br s, 1H)8.7 (s, 1H); 8.45 (s, 1H); 8.3 (s, 1H), 7.6 (s, 1H); 7.0 (s, 1H); 4.0 (m, 2H); 3.8 (m, 2H); 3.35 (m, 2H); 3.2 (m, 2H); 2.1 (m, 2H).	1.80
152	519.20	DMSO d6: 11.9 ppm (s, 1H), 10.1 (s, 1H), 9.3 (s, 1H), 9.0 (bs, 1H), 8.2 (s, 1H), 8.1 (s, 1H), 7.5 (t, 1H), 7.15 (d, 1H), 6.7 (d, 1H), 4.3 (m, 4H), 3.0-3.8 (m, 14H), 3.1 (m, 2H), 2.1 (s, 3H), 2.0 (q, 2H), 1.3 (bs, 6H)	1.40

Cmpd No.	LC_MASS _PLUS M+1 (obs)	NMR_RESULT (1H NMR)	RT (mins)
153	520.20	DMSO d6: 11.9 ppm (s, 1H), 10.1 (s, 1H), 9.3 (s, 1H), 8.2 (s, 1H), 8.1 (s, 1H), 7.5 (t, 1H), 7.1 (d, 1H), 6.7 (d, 1H), 4.3 (m, 4H), 4.0 (d, 2H), 3.7 (t, 4H), 3.5 (d, 2H), 3.2 (m, 2H), 3.1 (m, 4H), 2.1 (s, 3H), 2.0 (q, 2H), 1.3 (bs, 6H)	1.60
154	433.10	DMSO d6: 11.9 ppm (s, 1H), 9.9 (s, 1H), 9.3 (s, 1H), 8.3 (s, 1H), 8.1 (s, 1H), 7.5 (t, 1H), 7.1 (d, 1H), 6.7 (d, 1H), 5.9 (s, 1H), 5.5 (s, 1H), 4.4 (m, 4H), 3.0 (m, 2H), 2.1 (s, 3H), 2.0 (s, 3H), 1.3 (bs, 6H)	2.20
155	473.10	DMSO d6: 11.8 ppm (s, 1H), 9.6 (s, 1H), 9.3 (s, 1H), 8.3 (s, 1H), 8.1 (s, 1H), 7.5 (t, 1H), 7.1 (d, 1H), 6.75 (t, 1H), 6.7 (d, 1H), 4.4 (m, 4H), 3.0 (m, 2H), 2.3 (m, 2H), 2.2 (m, 2H), 2.1 (s, 3H), 1.7 (m, 2H), 1.6 (m, 2H), 1.3 (bs, 6H)	2.60
156	469.10	DMSO d6: 11.9 ppm (s, 1H), 10.3 (s, 1H), 9.4 (s, 1H), 8.4 (s, 1H), 8.1 (s, 1H), 8.0 (d, 2H), 7.6 (m, 4H), 7.2 (d, 1H), 6.7 (d, 1H), 4.4 (m, 4H), 3.1 (m, 2H), 2.1 (s, 3H), 1.3 (bs, 6H)	2.50
157	459.10	DMSO d6: 11.9 ppm (s, 1H), 9.7 (s, 1H), 9.3 (s, 1H), 8.3 (s, 1H), 8.1 (s, 1H), 7.5 (t, 1H), 7.1 (d, 1H), 6.75 (s, 1H), 6.7 (d, 1H), 4.4 (m, 4H), 3.1 (m, 2H), 2.6 (dt, 4H), 2.1 (s, 3H), 1.9 (q, 2H), 1.3 (bs, 6H)	2.40
158	475.20	DMSO d6: 11.8 ppm (s, 1H), 9.8 (s, 1H), 9.3 (s, 1H), 8.15 (s, 1H), 8.1 (s, 1H), 7.5 (t, 1H), 7.1 (d, 1H), 6.7 (d, 1H), 4.4 (m, 4H), 3.1 (m, 2H), 2.4 (m, 1H), 2.1 (s, 3H), 1.8 (dd, 4H), 1.4 (q, 2H), 1.3 (bs, 10H)	2.60
159	379.00	12.03 (s, 1H), 8.7-9.8 (s, 5H), 8.40 (s, 1H), 8.08 (s, 1H), 7.96 (s, 1H), 7.77 (t, 1H), 7.15 (d, 1H), 6.90 (d, 1H), 4.15-4.7 (m, 4H), 3.31 (m, 2H), 2.95 (s, 3H), 2.16 (s, 3H), 1.37 (m, 6H) CD3CN	1.70
160	440.90	DMSO d6 12.3 (bs, 1H); 10.0 (bs, 1H); 8.6 (s, 1H); 8.2 (s, 1H); 8.1 (s, 1H); 7.3 (s, 1H); 7.1 (s, 1H); 4.4 (bs, 2H); 4.2 (d, 2H); 3.1 (bd, 2H); 2.05 (s, 3H); 2.01 (s, 3H); 1.3 (bs, 6H)	2.40
161	447.00	DMSO d6: 11.8 ppm (s, 1H), 9.7 (s, 1H), 9.3 (s, 1H), 8.3 (s, 1H), 8.1 (s, 1H), 7.5 (t, 1H), 7.1 (d, 1H), 6.7 (d, 1H), 6.5 (q, 1H), 4.4 (m, 4H), 3.1 (m, 2H), 2.1 (s, 3H), 1.9 (s, 3H), 1.8 (d, 3H), 1.2 (bs, 6H)	2.40
162	475.90	DMSO d6: 11.9 ppm (s, 1H), 10.5 (s, 1H), 9.4 (s, 1H), 9.3 (s, 1H), 8.5 (s, 1H), 8.45 (s, 1H), 8.1 (s, 1H), 7.6 (t, 1H), 7.1 (d, 1H), 6.7 (d, 1H), 4.3 (m, 4H), 3.1 (m, 2H), 2.1 (s, 3H), 1.3 (bs, 6H)	2.30

Cmpd No.	LC_MASS _PLUS M+1 (obs)	NMR_RESULT (1H NMR)	RT (mins)
163	470.00	DMSO d6: 11.9 ppm (s, 1H), 10.6 (s, 1H), 9.4 (s, 1H), 9.3 (s, 1H), 9.2 (s, 1H), 8.8 (d, 1H), 8.4 (s, 1H), 8.4 (d, 1H), 8.2 (s, 1H), 7.7 (q, 1H), 7.6 (t, 1H), 7.2 (d, 1H), 6.7 (d, 1H), 4.4 (m, 4H), 3.1 (m, 2H), 2.1 (s, 3H), 1.3 (bs, 6H)	2.10
164	471.00	DMSO d6: 11.9 ppm (s, 1H), 10.9 (s, 1H), 9.5 (s, 1H), 9.3 (s, 1H), 9.0 (s, 1H), 8.8 (s, 1H), 8.5 (s, 1H), 8.2 (s, 1H), 7.6 (t, 1H), 7.1 (d, 1H), 6.7 (d, 1H), 4.4 (m, 4H), 3.0 (m, 2H), 2.1 (s, 3H), 1.3 (bs, 6H)	2.30
165	433.00	DMSO d6: 11.8 ppm (s, 1H), 10.2 (s, 1H), 9.3 (s, 1H), 8.2 (s, 1H), 8.1 (s, 1H), 7.5 (t, 1H), 7.1 (d, 1H), 6.7 (d, 1H), 4.3 (bd, 4H), 3.0 (bd, 2H), 2.1 (s, 1H), 1.8 (m, 1H), 1.3 (bs, 6H), 0.8 (m, 4H)	2.30
166	461.10	DMSO d6: 11.8 ppm (s, 1H), 9.9 (s, 1H), 9.3 (s, 1H), 8.2 (s, 1H), 8.1 (s, 1H), 7.5 (t, 1H), 7.1 (d, 1H), 6.7 (d, 1H), 4.4 (bd, 4H), 3.0 (bd, 2H), 2.8 (q, 1H), 2.1 (s, 1H), 1.9 (m, 1H), 1.8 (m, 1H), 1.7 (m, 1H), 1.6 (m, 1H), 1.3 (bs, 6H)	2.60
167	460.00	DMSO d6: 12.0 ppm (s, 1H), 10.9 (s, 1H), 9.3 (s, 1H), 8.8 (s, 1H), 8.4 (s, 1H), 8.2 (s, 1H), 7.6 (t, 1H), 7.2 (s, 1H), 7.15 (d, 1H), 6.7 (d, 1H), 4.4 (bd, 4H), 3.1 (bd, 2H), 2.1 (s, 3H), 1.3 (bs, 6H)	2.30
168	483.00	DMSO d6: 11.9 ppm (s, 1H), 10.3 (s, 1H), 9.4 (s, 1H), 8.4 (s, 1H), 8.2 (s, 1H), 7.85 (s, 1H), 7.8 (d, 1H), 7.6 (t, 1H), 7.4 (s, 2H), 7.4 (q, 1H), 7.2 (d, 1H), 6.7 (d, 1H), 4.4 (bd, 4H), 3.1 (bd, 2H), 2.4 (s, 3H), 2.1 (s, 3H), 1.3 (bs, 6H)	2.70
169	487.00	DMSO d6: 11.9 ppm (s, 1H), 10.4 (s, 1H), 9.4 (s, 1H), 8.4 (s, 1H), 8.2 (s, 1H), 7.9 (d, 1H), 7.8 (d, 1H), 7.6 (q, 1H), 7.55 (t, 1H), 7.45 (t, 1H), 7.2 (d, 1H), 6.7 (d, 1H), 4.4 (bd, 4H), 3.1 (bd, 2H), 2.1 (s, 3H), 1.3 (bs, 6H)	2.70
170	437.00	DMSO d6: 11.9 ppm (s, 1H), 9.8 (s, 1H), 9.3 (s, 1H), 8.3 (s, 1H), 8.1 (s, 1H), 7.5 (t, 1H), 7.1 (d, 1H), 6.7 (d, 1H), 4.4 (bd, 4H), 4.1 (s, 2H), 3.4 (s, 1H), 3.0 (bd, 2H), 2.1 (s, 3H), 1.3 (bs, 6H)	2.10
171	476.00	DMSO d6: 12.0 ppm (s, 1H), 10.5 (s, 1H), 9.0 (s, 1H), 8.3 (s, 1H), 8.2 (s, 1H), 7.6 (t, 1H), 7.1 (d, 1H), 6.7 (d, 1H), 4.4 (bd, 4H), 3.3 (m, 2H), 3.0 (bd, 2H), 2.3 (m, 2H), 2.1 (s, 3H), 1.5-1.9 (m, 4H), 1.3 (bs, 6H)	1.70
172	487.00	DMSO d6: 11.9 ppm (s, 1H), 10.4 (s, 1H), 9.4 (s, 1H), 8.3 (s, 1H), 8.2 (s, 1H), 7.7 (t, 1H), 7.6 (q, 1H), 7.5 (t, 1H), 7.4 (q, 2H), 7.2 (d, 1H), 6 .7 (d, 1H), 4.4 (bd, 4H), 3.1 (bd, 2H), 2.1 (s, 3H), 1.3 (bs, 6H)	2.60
173	502.90	DMSO d6: 11.9 ppm (s, 1H), 10.5 (s, 1H), 9.4 (s, 1H), 8.3 (s, 1H), 8.2 (s, 1H), 7.4-7.7 (m, 5H), 7.2 (d, 1H), 6 .7 (d, 1H), 4.4 (bd, 4H), 3.1 (bd, 2H), 2.0 (s, 3H), 1.3 (bs, 6H)	2.60

Cmpd No.	LC_MASS _PLUS M+1 (obs)	NMR_RESULT (1H NMR)	RT (mins)
174	502.90	DMSO d6: 11.9 ppm (s, 1H), 10.5 (s, 1H), 9.4 (s, 1H), 8.4 (s, 1H), 8.2 (s, 1H), 8.0 (d, 1H), 7.7 (s, 1H), 7.6 (t, 1H), 7.5 (t, 1H), 7.2 (d, 1H), 6 .7 (d, 1H), 4.4 (bd, 4H), 3.1 (bd, 2H), 2.1 (s, 3H), 1.3 (bs, 6H)	2.90
175	307.00	DMSO d6 12.4 (bs, 1H); 8.4 (s, 1H); 8.3 (s, 1H); 8.2 (s, 1H); 7.95 (d, 2H); 7.9 (s, 1H)	2.80
176	253.70	DMSO d6 12.4 (bs, 1H); 8.5 (s, 1H); 8.3 (s, 1H); 8.2 (s, 1H); 8.0 (d, 2H); 7.9 (d, 2H)	3.10
177	350.95	DMSO d6 12.4 (bs, 1H); 8.35 (s, 1H); 8.3 (s, 1H); 8.2 (m, 2H); 8.1 (s, 1H); 7.9 (s, 1H)	3.16
178	287.01	DMSO d6 12.2 (bs, 1H); 8.4 (s, 1H); 8.3 (s, 1H); 8.0 (s, 1H); 7.65 (s, 1H); 7.6 (dd, 1H); 7.4 (dd, 1H); 7.2 (dd, 1H); 3.8 (S, 2H)	2.69
179	494.00	DMSO d6: 11.9 ppm (s, 1H), 10.5 (s, 1H), 9.4 (s, 1H), 8.45 (s, 1H), 8.4 (s, 1H), 8.3 (d, 1H), 8.2 (s, 1H), (8.1 (d, 1H), 7.8 (t, 1H), 7.6 (t, 1H), 7.2 (d, 1H), 6.7 (d, 1H), 4.4 (bd, 4H), 3.1 (bd, 2H), 2.1 (s, 3H), 1.3 (bs, 6H)	2.60
180	449.10	DMSO d6: 11.8 ppm (s, 1H), 9.9 (s, 1H), 9.3 (s, 1H), 8.2 (s, 1H), 8.1 (s, 1H), 7.6 (t, 1H), 7.1 (d, 1H), 6.7 (d, 1H), 4.4 (bd, 4H), 3.1 (bd, 2H), 2.4 (m, 1H), 2.1 (s, 3H), 1.6 (m, 1H), 1.4 (m, 1H), 1.3 (bs, 6H), 1.1 (d, 3H), 0.9 (t, 3H)	2.50
181	433.00	DMSO d6: 11.9 ppm (s, 1H), 10.0 (s, 1H), 9.3 (s, 1H), 8.2 (s, 1H), 8.1 (s, 1H), 7.5 (t, 1H), 7.1 (d, 1H), 6.8 (m, 1H), 6.7 (d, 1H), 6.2 (d, 1H), 4.4 (bd, 4H), 3.1 (bd, 2H), 2.1 (s, 3H), 1.9 (d, 3H), 1.3 (bs, 6H)	2.30
182	474.90	DMSO d6: 11.9 ppm (s, 1H), 10.3 (s, 1H), 9.3 (s, 1H), 8.4 (s, 1H), 8.2 (s, 1H), 8.0 (d, 1H), 7.9 (d, 1H), 7.6 (t, 1H), 7.25 (t, 1H), 7.2 (d, 1H), 6.7 (d, 1H), 4.4 (bd, 4H), 3.1 (bd, 2H), 2.1 (s, 3H), 1.3 (bs, 6H)	2.50
183	228.80	DMSO d6 12.2 (bs, 1H); 8.3 (s, 1H); 8.25 (s, 1H); 8.0 (s, 1H); 7.7 (d, 2H); 7.4 (dd); 7.2 (dd, 1H)	3.40
184	230.10	DMSO d6 12.2 (bs, 1H); 8.3 (s, 1H); 8.25 (s, 1H); 8.0 (s, 1H); 7.7 (d, 2H); 7.4 (dd); 7.2 (dd, 1H)	1.60
185	243.10	DMSO d6 12.2 (bs, 1H); 8.3 (s, 1H); 8.25 (s, 1H); 7.9 (s, 1H); 7.5 (dd, 2H); 7.3 (dd, 1H); 7.1 (d, 1H); 2.3 (s, 3H)	3.40
186	334.10	DMSO d6 12.2 (bs, 1H); 8.25 (s, 1H); 8.2 (s, 1H); 8.0 (s, 1H); 7.5 (d, 1H); 7.4 (dd, 2H); 7.3 (d, 1H); 7.05 (dd, 2H); 6.7 (d, 2H); 6.55 (dd, 1H); 4.3 (s, 2H)	2.60
187	401.80	DMSO d6 12.2 (bs, 1H); 8.3 (s, 1H); 8.05 (s, 1H); 8.0 (s, 1H); 7.5 (m, 5H); 7.3 (dd, 1H); 7.2 (s, 1H); 7.1 (s, 1H); 6.8 (s, 1H); 4.4 (s, 2H)	3.90
188	291.02		2.30

Cmpd No.	LC_MASS _PLUS M+1 (obs)	NMR_RESULT (1H NMR)	RT (mins)
189	487.00	DMSO d6: 11.9 ppm (s, 1H), 10.3 (s, 1H), 9.3 (s, 1H), 8.4 (s, 1H), 8.15 (s, 1H), 8.1 (m, 2H), 7.5 (t, 1H), 7.4 (t, 2H), 7.15 (d, 1H), 6.7 (d, 1H), 4.4 (bd, 4H), 3.1 (bd, 2H), 2.1 (s, 3H), 1.3 (bs, 6H)	2.60
190	460.00	DMSO d6: 11.9 ppm (s, 1H), 10.6 (bs, 1H), 9.3 (s, 1H), 8.5 (s, 1H), 8.2 (s, 1H), 7.5 (t, 1H), 7.1 (d. 1H), 6.7 (d, 1H), 4.4 (bd, 4H), 3.1 (bd, 2H), 2.1 (s, 3H), 1.2 (bs, 6H)	2.00
191	445.00	DMSO d6: 11.9 ppm (s, 1H), 10.2 (bs, 1H), 9.4 (s, 1H), 8.3 (s, 1H), 8.1 (s, 1H), 7.5 (t, 1H), 7.2 (m, 1H), 7.1 (d. 1H), 6.7 (d, 1H), 6.6 (m, 1H), 6.3 (d, 1H), 5.7 (d, 1H), 5.5 (d, 1H), 4.4 (bd, 4H), 3.1 (bd, 2H), 2.1 (s, 3H), 1.2 (bs, 6H)	2.40
192	452.90	DMSO d6: 11.9 ppm (s, 1H), 10.3 (s, 1H), 9.3 (s, 1H), 8.2 (s, 1H), 8.1 (s, 1H), 7.5 (t, 1H), 7.4 (d, 1H), 7.1 (d. 1H), 6.7 (d, 1H), 6.7 (s, 1H), 4.4 (bd, 4H), 3.1 (bd, 2H), 2.1 (s, 3H), 1.2 (bs, 6H)	2.50
193	456.10	DMSO-d6: 11.9 (bs, 1H); 8.3 (s, 1H); 8.1 (m, 2H); 7.55 (dd, 1H); 7.45 (d, 2H); 7.4 (dd, 2H); 7.3 (m, 1H); 7.15 (d, 1H); 6.7 (d, 1H); 5.2 (s, 2H); 4.2 (d, 4H); 3.1 (m, 2H); 2.05 (s, 3H); 1.25 (s, 6H).	3.00
194	451.20	DMSO-d6: 11.9 (bs, 1H); 8.25 (s, 1H); 8.1 (s, 1H); 8.05 (s, 1H); 7.55 (dd, 1H); 7.15 (d, 1H); 6.7 (d, 1H); 4.3 (d, 4H); 4.15 (m, 2H); 3.25 (m, 2H); 3.05 (m, 2H); 2.8 (s, 6H); 2.15 (m, 2H); 2.05 (s, 3H); 1.25 (s, 6H).	1.60
195	279.80	(d4-methanol, free base) 8.87 (s, 1H), 8.70 (s, 1H), 8.50 (s, 1H), 8.38 (s, 1H), 8.20 (d, 1H), 8.18 (d, 1H), 8.10 (m, 1H), 7.96 (m, 1H)	2.86
196	311.80	(d4-methanol, salt) 8.88 (s, 1H), 8.25 (m, 2H), 7.86 (s, 1H), 7.34 (m, 1H), 2.72 (s, 3H)	3.95
197	293.80	(d4-methanol, salt) 8.71-7.45 (m, 8H), 2.90 (s, 3H)	2.33
198	293.90	(d4-methanol, salt) 8.53-7.82 (m, 7 H), 7.41 (s, 1H), 4.34 (s, 3H)	2.33
199	278.90	(d4-methanol, salt) 8.39 (dd, 1H), 8.25 (s, 1H), 8.11 (s, 1H), 7.94-7.79 (m, 5H), 7.51-7.44 (m, 2H)	3.87
200	279.90	(d4-methanol, salt) 9.16-7.99 (m, 9H)	3.87
201	259.90	(d4-methanol, salt) 8.86 (d, 1H), 8.23 (s, 1H), 8.09 (s, 1H), 7.99 (m, 1H), 7.70 (d, 1H), 7.39 (dd, 1H), 6.65 (d, 1H), 4.17 (s, 3H)	3.46
202	279.80	(d4-methanol, salt) 9.01 (d, 1H), 8.98 (d, 1H), 8.32 (s, 1H), 8.31 (s, 1H), 8.28 (d, 1H), 8.17 (d, 1H), 8.01-7.97 (m, 2H), 7.94 (s, 1H)	2.50
203	287.90	(CDCl3) 10.00 (br s, 1H), 8.49 (s, 1H), 8.27 (s, 1H), 8.00 s, 1H), 7.71 (m, 1H), 7.48 (m, 1H), 7.20 (m, 1H), 7.19 (d, 1H), 1.97 (s, OH, 1H), 1.57 (s, 6H)	2.56

Cmpd No.	LC_MASS _PLUS M+1 (obs)	NMR_RESULT (1H NMR)	RT (mins)
204	412.90	DMSO d6 12.2 (bs, 1H); 8.7 (t, 1H); 8.4 (s, 1H); 8.2 (s, 1H); 8.1 (s, 1H); 8.0 (s, 1H); 4,8 (q, 1H); 3.9 (s, 3H); 3.85 (m, 2H); 1.15 (d, 3H)	1.90
205	394.20		2.10
206	408.10		2.20
207	420.20		2.30
208	493.20		1.70
209	452.20		2.20
210	343.00	1H NMR (CDCl3) 0.30 (9H, s), 7.33-7.54 (8H, m), 7.59-7.68 (2H, m), 8.05-8.08 (1H, m), 8.61-8.65 (1H, m), 9.70 (1H, brs).	
211	349.10	(400 MHz, DMSO-d6) 7.30-7.43(8H, br m), 7.48(2H, m), 7.98(1H,d), 8.34(1H, d), 12.36(1H, br m).	
212	364.40	(400 MHz, DMSO-d6) 7.00(2H, d), 7.23(2H, d), 7.34(3H, m), 7.49(2H, dd), 7.97(1H, s), 8.33(1H, d) and 12.38(1H, s).	
213	271.00	1H NMR (CDCl3) 7.30-7.57 (6H, m), 7.60-7.77 (5H, m), 8.44-8.48 (1H, m), 8.65-8.70 (1H, m), 11.21 (1H, brs).	
214	301.00	1H NMR (DMSO) 3.84 (3H, s), 6.80-6.88 91H, m), 7.25-7.30 (1H, m), 7.32-7.41 (3H, m), 7.45-7.54 (2H, m), 7.72-7.81 (2H, m), 7.90-7.98 (1H, m), 8.39-8.43 91H, m), 8.54-8.60 (1H, m), 12.04 (1H, brs).	
215	315.00	1H NMR (DMSO) 1.36 (3H, t, J = 6.9Hz), 4.10 (2H, q, J = 6.9Hz), 6.80-6.86 (1H, m), 7.22-7.29 (1H, m), 7.30-7.41 (3H, m), 7.45-7.55 (2H, m), 7.73-7.80 (2H, m), 7.91-7.98 (1H, m), 8.38-8.41 (1H, m), 8.53-8.59 (1H, m), 12.02 91H, brs).	
216	331.00	1H NMR (DMSO) 3.70-3.80 (6H, m), 6.81-6.90 (1H, m), 7.02-7.15 (2H, m), 7.32-7.40 (1H, m), 7.42-7.51 (2H, m), 7.66-7.79 (3H, m), 8.11-8.16 (1H, m), 8.50-8.56 (1H, m), 11.95 (1H, brs).	
217	436.70	(400 MHz, DMSO-d6) 12.31(1H, s), 8.32(1H, d), 7.94(1H, d), 7.52(2H, d), 7.33(3H, m), 7.10(2H, d), 6.98(2H, d), 3.75(4H, dd) and 3.15(4H, dd).	
218	349.00	1H NMR (DMSO) 3.30 (3H, s), 7.34-7.45 (1H, m), 7.49-7.83 (6H, m), 8.16 (1H, s), 8.18-8.28 (2H, m), 8.45-8.49 91H, m), 8.58-8.62 (1H, m), 12.21 (1H, brs).	
219	342.00	1H NMR (DMSO) 3.00 (6H, s), 7.35-7.42 (1H, m), 7.45-7.68 (4H, m), 7.75-7.90 (4H, m), 8.03 (1H, s), 8.45-8.50 (1H, m), 8.56-8.60 (1H, m), 12.11 (1H, brs).	

Cmpd No.	LC_MASS _PLUS M+1 (obs)	NMR_RESULT (1H NMR)	RT (mins)
220	370.00	1H NMR (DMSO) 0.91 (3H, t, J = 7.4Hz), 1.30-1.42 (2H, m), 1.49-1.59 (2H, m), 3.25-3.34 (2H, m), 7.35-7.42 (1H, m), 7.46-7.58 (3H, m), 7.68-7.81 (3H, m), 7.92-8.03 (2H, m), 8.19-8.22 (1H, m), 8.44-8.49 (1H, m), 8.50-8.61 (2H, m), 12.10 (1H, brs).	
221	367.00	1H NMR (DMSO) 3.36 (3H, s), 7.28-7.39 (2H, m), 7.50-7.88 (5H, m), 8.10-8.26 (3H, m), 8.43-8.46 (1H, m), 8.57-8.59 (1H, m), 12.25 (1H, brs).	
222	349.00	1H NMR (DMSO) 3.79 (3H, s), 3.86 (3H, s), 7.00-7.05 (1H, m), 7.26-7.37 (4H, m), 7.78-7.89 (2H, m), 8.35-8.40 (1H, m), 8.50-8.56 (1H, m), 11.93 (1H, brs).	
223	307.00	1H NMR (DMSO) 7.22-7.37 (4H, m), 7.78-7.88 (4H, m), 7.92 (1H, brs), 8.36-8.41 91H, m), 8.51-8.58 (1H, m), 12.02 (1H, brs).	
224	319.00	1H NMR (DMSO) 3.80 (3H, s), 6.99-7.06 (2H, m), 7.26-7.36 (2H, m), 7.68-7.84 (5H, m), 8.33-8.39 (1H, m), 8.50-8.57 (1H, m), 11.91 (1H, brs).	
225	452.00	1H NMR (DMSO) 3.30-3.80 (8H, m), 7.25-7.32 (1H, m), 7.50-7.63 (2H, m), 7.80 (1H, s), 7.88-7.99 (2H, m), 8.05 (1H, s), 8.49-8.54 (1H, m), 8.60-8.68 (1H, m), 12.20 (1H, brs).	
226	410.00	1H NMR (DMSO) 1.16 (3H, t), 3.20-3.42 (2H, m), 7.50-7.76 (3H, m), 7.88-7.95 (2H, m), 7.98-8.06 (2H, m), 8.20 (1H, s), 8.51-8.66 (3H, m), 12.18 (1H, brs).	
227	332.00	1H NMR (DMSO) 7.28-7.39 (3H, m), 7.79-8.03 (7H, m), 8.09 (1H, brs), 8.48-8.51 (1H, m), 8.53-8.59 (1H, m), 12.14 (1H, brs).	
228	332.00	1H NMR (DMSO) 7.30-7.48 (3H, m), 7.42 (1H, brs), 7.50-7.58 (1H, m), 7.70-7.86 (3H, m), 7.92-8.11 (3H, m), 8.20 (1H, brs), 8.41-8.49 (1H, m), 8.54-8.59 (1H, m), 12.09 (1H, brs).	
229	299.00	1H NMR (DMSO) 7.38-7.43 91H, m), 7.48-7.55 (2H, m), 7.78-7.83 (2H, m), 7.97 92H, d, J = 8.2Hz), 8.09 (2H, d, J = 8.2Hz), 8.21 (1H, s), 8.54-8.62 92H, m), 10.00 91H, s), 12.30 (1H, brs).	
230	342.00	1H NMR (DMSO) 1.90 93H, s), 4.20-4.30 92H, m), 7.27-7.40 (3H, m), 7.45-7.51 92H, m), 7.70-7.80 (4H, m), 7.85-7.91 (1H, m), 8.30-8.41 (2H, m), 8.51-8.60 (1H, m), 12.00 (1H, brs).	
231	313.00	1H NMR (DMSO) 2.60 (3H, s), 7.36-7.42 (1H, m), 7.47-7.55 (2H, m), 7.76-7.83 (2H, m), 7.95-8.08 (4H, m), 8.15 (1H, s), 8.50-8.56 (1H, m), 8.58-8.63 91H, m), 12.24 (1H, brs).	

Cmpd No.	LC_MASS _PLUS M+1 (obs)	NMR_RESULT (1H NMR)	RT (mins)
232	286.00	1H NMR (DMSO) 5.08 (2H, brs), 6.63-6.70 (2H, m), 7.30-7.52 95H, m), 7.60-7.66 (1H, m), 7.70-7.80 (2H, m), 8.30-8.35 (1H, m), 8.50-8.55 (1H, m), 11.75 (1H, brs).	
233		1H NMR (DMSO) 2.40-2.48 (3H, m), 7.36-7.55 (4H, m), 7.78–7.85 (4H, m), 8.02-8.09 (2H, m), 8.15 (1H, s), 8.52-8.56 (1H, m), 8.59-8.62 (1H, m), 12.25 (1H, brs).	
234	364.00	1H NMR (DMSO) 3.00 (3H, s), 7.25-7.54 (5H, m), 7.70-7.90 (5H, m), 8.40-8.48 (1H, m), 8.52-8.60 (1H, m), 9.73 (1H, brs), 12.00 (1H, brs).	8.93
235	304.00	1H NMR (DMSO) 7.15(1H, s), 7.35 (1H, brs), 7.78 (1H, s), 7.85–8.08 (7H, m), 8.31 (1H, s), 8.49 91H, brs), 8.61 (1H, brs), 12.08 (1H, brs).	7.68
236	320.00	1H NMR (DMSO) 7.30 (1H, s), 7.60-7.76 (2H, m), 7.82-8.10 (7H, m), 8.55 (1H, brs), 8.70 (1H, s), 12.10 (1H, brs).	8.14
237	370.00	(400 MHz, MeOH-d4): 4.70-4.85 (2H, brs), 6.35-6.45 (1H, s), 7.30-7.50 (5H, m), 8.05-8.15 (1H, d), 8.20-8.25 (1H, s), 8.60–8.70 (1H, brs)	4.87
238	388.00	1H NMR (DMSO-d6): 3.33 (1H, s), 4.75-4.80 (2H, m), 7.35–7.45 (4H, m), 8.15 (1H, s), 8.20-8.25 (2H, m), 8.28-8.35 1H, m), 8.50 (1H, s)	4.87
239	344.00	1H NMR (DMSO) 3.85 (3H, s), 6.91-7.00 (1H, m), 7.26-7.49 (4H, m), 7.85-8.10 (6H, m), 8.50 (1H, brs), 8.60 (1H, brs), 12.15 (1H, brs).	8.38
240	308.24	1H NMR (DMSO-d6): 7.22 (1H, m), 7.47 (2H, m), 7.85 (1H, s), 8.21 (1H, s), 8.64 (1H, s), 11.19 (1H, br s)	4.93
241	299.38	1H NMR (CDCl3): 2.04 (4H, s), 3.53 (4H, s), 6.22 (1H, d), 6.86 (1H, d), 7.44 (1H, t), 7.81 (1H, s), 8.17 (1H, s), 8.87 (1H, s)	5.15
242	369.00	1H NMR (DMSO-d6): 4.60-4.70 (2H, m), 6.35-6.40 (1H, d), 7.05-7.10 (1H, d), 7.20-7.25 (1H, t), 7.35-7.45 (4H, m), 8.15-8.20 (2H, m), 8.55-8.60 (1H, s), 12.5 (1H, s)	4.99
243	321.00	1H NMR (DMSO-d6): 6.70-6.75 (1H, d), 6.90-6.95 (1H, t), 7.30-7.35 (3H, qd), 7.50-7.65 (3H, m), 8.25-8.30 (2H, m), 8.80 (1H, s), 9.05 (1H, s), 12.5 (1H, s)	4.84
244	401.49	1H NMR (DMSO-d6): 4.63 (2H, d), 5.20 (1H, br s), 6.45 (1H, d), 7.08 (1H, d), 7.27 (1H, t), 7.37 (2H, s), 7.43 (1H, t), 7.64 (1H, s), 8.16 (2H, d), 8.39 (1H, s), 12.48 (1H, br s)	5.20
245	513.00	(d6-DMSO, 400MHz) 3.16 (3H, s), 5.30 (2H, s), 7.52 (1H, d), 7.60 (2H, d), 7.70 (1H, d), 7.81 (1H, t), 7.91 (2H, d), 8.17 (1H, d), 8.28 (1H, d), 8.36 (1H, s), 12.31 (1H, s)	4.89
246	245.00	(d6-DMSO, 400MHz) 5.98 (2H, brs), 6.25 (1H, d), 7.01 (1H, d), 7.37 (1H, t), 8.18 (1H, s), 8.24 (1H, d), 8.96 (1H, d), 12.12 (1H, s)	3.80

Cmpd No.	LC_MASS _PLUS M+1 (obs)	NMR_RESULT (1H NMR)	RT (mins)
247	413.00	(d6-DMSO, 400MHz) 3.15 (3H, s), 4.71 (2H, s), 6.38 (1H, d), 7.06 (1H, d), 7.42 (1H, t), 7.65 (2H, d), 7.88 (2H, d), 8.18 (2H, s), 8.53 (1H, s), 12.12 (1H, s)	4.00
248	372.00	1H NMR (DMSO-d6): 7.05-7.10 (1H, t), 7.35-7.40 (1H, m), 7.60-7.65 (1H, t), 7.70–7.80 (3H, m), 7.82-7.88 (1H, d), 7.95 (1H, s), 8.25 (1H, s), 8.40 (1H, s), 12.5 (1H, s)	5.22
249	349.00	(d6-DMSO, 400MHz) 7.56 - 7.60 (2H, m), 7.63 (1H, d), 7.69 (1H, d), 7.82 (1H, t), 7.97 (1H, s), 8.00 (2H, d), 8.27 (1H, s), 8.42 (1H, s), 9.28 (1H, s), 10.81 (1H, s), 12.29 (1H, s)	4.59
250	345.00	(d6-DMSO, 400MHz) 1.53 (9H, s), 7.56 (2H, dd), 7.69 (1H, t), 8.25 (1H, d), 8.36 (1H, s), 9.30 (1H, s), 10.06 (1H, s), 12.22 (1H, s)	4.90
251	405.00	NMR (DMSO) 4.6 (2H, d, CH2), 6.4 (H, d, ar), 7.05 (H, d, ar), 7.3 (H, m, ar), 7.4 (H, d, ar), 7.45 (H, t, ar), 7.6 (H, d, ar), 7.65 (H, s, ar), 8.15 (2H, s, ar), 8.5 (H, s, NH) and 12.15 (H, s, NH).	5.15
252	349.00	1H NMR (DMSO-d6): 1.50-1.55 (3H, d), 5.05-5.15 (2H, m), 6.30-3.35 (1H, d), 6.95-7.05 (1H, d), 7.10-7.15 (1H, d), 7.15-7.20 (1H, t), 7.25-7.40 (3H, m), 7.45-7.50 (2H, d), 8.15 (1H, s), 8.20 (1H, s), 8.75 (1H, s), 12.5 (1H, s)	4.93
253	315.70	CDCl3 3.23 (6H, m), 6.45 (1H, d), 7.03 (1H, d), 7.43 (1H, m), 7.5-7.57 (3H, m), 7.70-7.73 (2H, m), 7.88 (1H, s), 8.65 (1H, s), 9.10 (1H, s), 9.25 (1H, s).	10.37
254	349.00	NMR (DMSO) 2.2 (3H, s, CH3), 4.5 (2H, s, CH2), 6.3 (H, s, ar), 6.8-7.4 (7H, m, ar), 8.1-8.3 (2H, m, ar) and 8.7 (H, s, NH).	5.03
255	407.00	1H NMR (DMSO) 3.01 (3H, s), 7.28-7.39 (3H, m), 7.70-8.10 (8H, m), 8.45 (1H, brs), 8.55 (1H, brs), 9.80 (1H, brs), 12.14 (1H, brs).	
256	407.00	1H NMR (DMSO) 2.45 (3H, brs), 7.31(1H, brs), 7.51 (1H, brs), 7.80-8.15 (10H, m), 8.60 (1H, brs), 8.67 (1H, brs), 12.21 (1H, brs).	
257	414.53	1H NMR (DMSO-d6): 1.44 (9H, s), 3.41 (4H, s), 3.57 (4H, d), 6.69 (1H, d), 7.22 (1H, d), 7.57 (1H, t), 8.27 (2H, s), 8.67 (1H, s), 12.25 (1H, s	2.50
258	365.40	1H NMR (DMSO-d6): 3.70 (3H, s), 6.35 (1H, d), 6.88 (2H, d), 7.03 (1H, d), 7.11 (1H, t), 7.34 (2H, d), 7.37 (1H, t), 8.18 (1H, s), 8.21 (1H, d), 8.76 (1H, s), 12.13 (1H, s)	4.78

Cmpd No.	LC_MASS _PLUS M+1 (obs)	NMR_RESULT (1H NMR)	RT (mins)
259	329.70	CDCl3 1.05 (3H, t), 1.7-1.8 (2H, m), 3.4-3.5 (2H, m), 4.6 (1H, s), 6.32 (1H, d), 7.03 (1H, d), 7.45 (1H, m), 7.5-7.6 (3H, m), 7.75 (2H, d), 7.78 (1H, s), 8.65 (1H, s), 8.97 (1H, s), 9.12 (1H, s)	4.99
260	349.00	(400 MHz, MeOH-d4): 1.55-1.60 (3H, d), 5.05-5.15 (1H, m), 6.30-6.35 (1H, d), 6.95-7.05 (1H, d), 7.10-7.15 (1H, m), 7.15-7.20 (1H, t), 7.25-7.40 (3H, m), 7.45-7.50 (2H, d), 7.95 (1H, s) 8.15 (1H, s), 8.75 (1H, s)	4.93
261	445.00	NMR (DMSO) 4.7 (2H, d, CH2), 6.35 (H, d, ar), 7.15 (H, d, ar), 7.2 (H, t, ar), 7.3-7.55 (7H, m, ar), 7.6 (s, H, ar), 8.2 (H, s, ar), 8.5 (H, s, NH) and 8.9 (H, s, NH).	5.30
262	391.00	NMR (DMSO) 2.2 (3H, s, Me), 4.6 (2H, d, CH2), 6.3 (H, d, ar), 7.05-7.1 (4H, m, ar), 7.25 (2H,, d, ar), 7.3-7.45 (4H, m, ar), 7.6 (2H, d, ar), 8.15 (H, s, ar), 8.5 (H, s, NH) and 9.05 (H, s, NH).	5.14
263	355.00	1H NMR (DMSO-d6): 6.70-6.80 (1H, d), 7.10-7.20 (1H, t), 7.30-7.40 (2H, m), 7.50-7.60 (2H, m), 7.80-7.90 (1H, d), 8.15 (1H, s), 8.25 (1H, s), 8.50 (2H, s), 12.2 (1H, s)	4.92
264	349.00	1H NMR (DMSO-d6): 2.95-3.00 (2H, t), 3.55-3.60 (2H, m), 6.30-6.35 (1H, d), 6.65-6.70 (1H, t), 7.05-7.10 (1H, d), 7.15-7.20 (1H, t), 7.30 (4H, m), 7.32-7.40 (1H, t), 8.17 (1H, s), 8.20 (1H, s), 8.85 (1H, s), 12.2 (1H, s)	4.67
265	363.00	NMR (DMSO) 6.7 (H, d, ar), 6.9 (H, t, ar), 7.35-7.4 (2H, m, ar), 7.4-7.5 (4H, m, ar), 7.5-7.75 (5H, m, ar), 8.25 (H, s, ar), 8.6 (H, s, ar), 8.9 (H, s, ar), 9.3 (H, s, NH) and 12.1 (1H, s, NH).	4.64
266	397.00	NMR (DMSO) 6.8 (H, d, ar), 7.0 (H, m, ar), 7.2 (H, m, ar), 7.2-7.6 (7H, m, ar), 7.65 (H, m, ar), 8.0 (H, m, ar), 8.2 (H, s, ar), 8.4 (H, m, NH), 8.5 (H, s, ar), 8.7 (H, s, ar) and 12.1 (H, brs, NH).	5.20
267	314.43,314.00	1H NMR (DMSO-d6): 3.11 (4H, s), 3.88 (4H, d), 6.78 (1H, d), 7.30 (1H, d), 7.65 (1H, t), 8.30 (2H, d), 8.63 (1H, s), 8.77 (2H, br s), 12.30 (1H, s),DMSO D6 2.82-2.93 (4H, m), 3.48-3.55 (4H, m), 6.65 (1H, d), 7.15 (1H, d), 7.55 (1H, t), 8.25-8.30 (2H, m), 8.69 (1H, s)	3.42,2.75
268	287.60	CDCl3 4.48 (2H, s), 6.45 (1H, d), 7.12 (1H, d), 7.42 (1H, t), 7.5-7.6 (3H, m), 7.75 (2H, t), 7.88 (1H, s), 8.7 (1H, s), 8.88 (1H, s), 9.52 (1H, s)	2.79

Cmpd No.	LC_MASS _PLUS M+1 (obs)	NMR_RESULT (1H NMR)	RT (mins)
269	377.60	CDCl3 4.7 (2H, d), 5.0 (1H, s), 6.35 (1H, d), 7.08 (1H, d), 7.3-7.4 (9H, t), 7.5-7.6 (3H, m), 7.68 (2H, d), 7.88 (1H, s), 8.65 (1H, s), 8.95 (1H, s), 9.43 (1H, s)	5.00
270	289.00	1H NMR (DMSO-d6): 3.45-3.50 (2H, m), 3.60-3.65 (2H, m), 4.70-4.77 (1H, t), 6.25-6.30 (1H, d), 6.50-6.55 (1H, t), 7.00-7.05 (1H, d), 7.30-7.40 (1H, t), 8.20 (1H, s), 8.25 (1H, s), 8.85 (1H, s), 12.2 (1H, s)	3.79
271	330.00	1H NMR (DMSO-d6): 2.00-2.10 (2H, m), 2.80-2.85 (6H, s), 3.20-3.30 (2H, m), 3.40-3.50 (2H, m), 6.25-6.30 (1H, m), 6.60-6.80 (1H, m), 7.00-7.05 (1H, d), 7.30-7.40 (1H, t), 8.20-8.30 (2H, m), 8.85 (1H, s), 12.2 (1H, s)	3.34
272	225.00	(d6-DMSO) 2.90 (3H, d), 6.24 (1H, d), 6.42 - 6.44 (1H, m), 7.00 (1H, d), 7.14 (1H, dd), 7.37 (1H, t), 8.08 (1H, s), 8.23 (1H, d), 8.85 (1H, d), 11.87 (1H, s)	3.32
273	355.00	1H NMR (DMSO-d6): 6.60-6.65 (1H, m), 6.70-6.80 (1H, d), 7.00-7.10 (1H, m), 7.15-7.30 (1H, m), 7.20-7.30 (1H, m), 7.55-7.65 (2H, m), 7.90 (1H, s), 8.15 (1H, s), 8.30 (1H, s), 8.60 (1H, s), 8.90-8.95 (1H, m), 12.2 (1H, s)	
274	349.48	1H NMR (DMSO-d6): 3.14 (3H, s), 4.92 (4H, s), 6.42 (1H, d), 7.09 (1H, d), 7.23 (1H, t), 7.31 (4H, m), 7.48 (1H, t), 8.11 (1H, s), 8.21 (1H, s), 8.55 (1H, s), 12.20 (1H, br s)	5.25
275	328.46	1H NMR (DMSO-d6): 2.24 (3H, s), 3.31 (4H, s), 3.56 (4H, s), 6.67 (1H, d), 7.19 (1H, d), 7.54 (1H, t), 8.27 (2H, s), 8.69 (1H, s), 12.23 (1H, s)	4.37
276	354.64	1H NMR (DMSO-d6): 2.07 (3H, s), 3.57 (8H, m), 6.70 (1H, d), 7.23 (1H, d), 7.58 (1H, t), 8.28 (2H, s), 8.68 (1H, s), 12.25 (1H, s)	4.25
277	331.70	DMSO 3.55 (1H,m), 3.65 (1H,m), 4.82 (1H,m), 6.35 (1H,d), 6.52 (1H,m), 7.05 (1H,d), 7.4 (1H,t), 7.55 (1H,t), 7.75 (1H,d), 8.15 (1H,s), 8.58 (1H,s), 9.08 (1H,s)	4.24
278	301.70	DMSO 2.92 (3H,s), 6.30 (1H,d), 6.51 (1H,m), 7.08 (1H,d), 7.35-7.42 (2H,m), 7.5-7.6 (2H,m), 7.72 (2H,d), 8.13 (1H,s), 8.65 (1H,s), 9.18 (1H,s)	4.62
279	315.00	NMR(DMSO) 1.1 (6H, d, iPr), 2.85 (H, m, CH), 3.3 (H, brs, NH), 7.55 (H, d, ar), 7.65 (H, t, ar), 7.85 (H, t, ar), 8.2 (H, s, ar), 8.3 (H, s, ar), 9.1 (H, s, ar) and 10.3 (H, s, NH).	4.47
280	363.00	NMR (DMSO) 3.85 (2H, s, CH2), 7.15 (5H, m, ar), 7.6 (H, d, ar), 7.7 (H, t, ar), 7.9 (h, d, ar), 8.3 (H, s, ar), 84 (H, s, ar), 9.1 (H, s, ar), 10.6 (H, s, NH) and 12.3 (H, brs, NH).	4.72

Cmpd No.	LC_MASS _PLUS M+1 (obs)	NMR_RESULT (1H NMR)	RT (mins)
281	383.00	(DMSO) 7.45-7.7 (3H, m, ar), 7.7-7.8 (2H, m, ar), 7.8 (H, t, ar), 8.0 (H, m, ar), 8.25 (H, s, ar), 8.4 (H, s, ar), 9.2 (H, s, ar), 11.1 (H, s, NH), and 12.3 (H, s, NH).	4.68
282	383.00	DMSO) 6.6 (H, m, ar), 7.7 (2H, d ar), 7.85 (H, t, ar), 8.0 (H, m, ar), 8.1 (H, s, ar), 8.25 (H, s, ar), 8.45 (H, s, ar), 9.3 (H, s, ar), 10.8 (0.5H, s, NH) and 12.3 (0.5H, s, NH).	4.95
283	287.00	(DMSO) 2.2 (3H, s, Me), 7.55 (H, m, ar), 7.7 (H, m, ar), 7.85 (H, m, ar), 8.2 (H, m,ar), 8.4 (H, m, ar), 10.4 (H, s, NH) and 12.3 (H, s, NH)	4.05
284	335.00	1H NMR (DMSO-d6): 3.50-3.55 (3H, m), 6.40-6.45 (1H, m), 7.25-7.30 (2H, m), 7.32-7.37 (2H, m), 7.45-7.55 (3H, m), 8.15 (1H, s), 8.30 (1H, s), 8.45 (1H, s), 12.2 (1H, s)	5.27
285	386.46	1H NMR (CDCl3): 1.69 (3H, d, J = 6.9 Hz), 5.29 (1H, m), 5.42 (1H, 6.8 Hz), 7.07 (2H, m), 7.44 (2H, m), 8.04 (1H, m), 8.09 (1H, m), 8.27 (1H, m), 8.65 (1H, m), 8.87 (1H, br s)	4.97
286	368.46	1H NMR (CDCl3): 1.69 (3H, d, J = 6.8), 5.33 (1H, m), 5.45 (1H, m), 7.38 (2H, m), 7.47 (2H, m), 8.04 (1H, d), 8.08 (1H, d), 8.27 (1H, d), 8.70 (2H, m)	4.90
287	369.80	DMSO 1.1-1.3 (6H,m), 1.6-1.7 (2H,m), 2.0-2.08 (2H,m), 4.0 (1H,m), 6.26 (1H,d), 6.35 (1H,d), 7.0 (1H,d), 7.3-7.4 (2H,m), 7.52 (2H,t), 7.7 (2H,d), 8.15 (1H,s), 8.55 (1H,s), 9.1 (1H,s)	5.27
288	355.00	(DMSO) 1.1-1.5 (5H, m, cyhex), 1.6-1.9 (5H, m, cyhex), 2.65 (1H, m, cyhex), 7.6 (H, d, ar), 7.75 (H, t, ar), 7.95 (H, d, ar), 8.3 (H, s, ar), 8.4 (H, s, ar), 9.15 (H, s, ar), 10.35 (H, s, NH) and 12.2 (H, s, NH).	4.95
289	313.45	1H NMR (DMSO-d6): 1.55 (6H, m), 2.01 (2H, s), 4.20 (1H, m), 6.27 (1H, d), 6.54 (1H, d), 7.01 (1H, d), 7.34 (1H, t), 8.23 (1H, s), 8.30 (1H, s), 8.96 (1H, s), 12.13 (1H, s)	5.09
290	327.45	1H NMR (DMSO-d6): 1.27 (3H, m), 1.45 (2H, m), 1.69 (1H, d), 1.81 (2H, d), 2.09 (2H, d), 3.81 (1H, m), 6.24 (1H, d), 6.39 (1H, d), 6.97 (1H, d), 7.31 (1H, t), 8.20 (2H, s), 8.88 (1H, s), 12.20 (1H, br s)	5.25
291	341.49	1H NMR (DMSO-d6): 1.51 (6H, m), 1.67 (4H, m), 2.01 (2H, 2), 4.02 (1H, s), 6.26 (1H, d), 6.44 (1H, d), 6.98 (1H, d), 7.33 (1H, t), 8.19 (1H, s), 8.24 (1H, s), 8.93 (1H, s), 12.13 (1H, br s)	5.37
292	273.38	1H NMR (DMSO-d6): 2.94 (3H, s), 3.53 (3H, s), 6.56 (1H, s), 7.05 (1H, s), 7.58 (1H, s), 8.31 (1H, s), 8.36 (1H, s), 8.78 (1H, s)	4.59

Cmpd No.	LC_MASS _PLUS M+1 (obs)	NMR_RESULT (1H NMR)	RT (mins)
293	341.55	1H NMR (CDCl3): 2.04 (4H, m), 3.61 (4H, m), 6.24 (1H, d, J = 8.3), 6.98 (1H, d, J = 7.4), 7.39 (1H, m), 7.48-7.52 (3H, m), 7.70 (2H, m), 7.88 (1H, m), 8.61 (1H, s), 9.16 (1H, s), 9.33 (1H, br s)	5.32
294	350.00	DMSO) 7.7-7.8 (2H, m, ar), 7.9 (H, t, ar), 8.05 (H, d, ar), 8.15 (H, t, ar), 8.25 (H, d, ar), 8.3 (H, s, ar), 8.45 (H, s, ar), 8.8 (H, d, ar), 9.0 (H, s, ar), 10.6 (H, s, NH) and 12.4 (H, s, NH).	4.85
295	383.00	(400 MHz, MeOH-d4): 7.85-7.95 (2H, d), 8.05-8.10 (2H, d), 8.25-8.30 (2H, s), 8.35-8.37 (1H, s), 8.70 (1H, s)	3.98
296	350.00	(DMSO) 7.6-7.7 (H, m, ar), 7.75 (H, m, ar), 7.85 (H, m, ar), 8.0 (H, m, ar), 8.3 (H, m, ar), 8.4 (H, m, ar), 8.8 (H, s, ar), 9.15 (h, s, ar), 9.3 (H, s, ar), 11.0 (H, s, ar) and 12.35 (H, s, NH)	4.22
297	395.55	1H NMR (DMSO-d6): 3.69 (6H, d), 4.41 (2H, br s), 6.34 (1H, s), 6.88 (2H, m), 6.94 (2H, s), 7.02 (1H, m), 7.36 (1H, t), 8.19 (2H, m), 8.76 (1H, s), 12.13 (1H, s)	4.64
298	257.59	1H NMR (DMSO-d6): 2.89 (3H, s), 6.26 (1H, d), 6.54 (1H, d), 7.03 (1H, d), 7.38 (1H, t), 8.21 (1H, s), 8.24 (1H, s), 8.94 (1H, s), 12.14 (1H, s)	4.40
299	354.00	(400 MHz, MeOH-d4): 2.35-2.40 (3H, s), 7.30-7.45 (4H, m), 8.05-8.15 (3H, m), 8.20-8.25 (1H, s)	4.87
300	368.00	(400 MHz, DMSO-d6): 2.20 (6H, s), 7.25 (3H, m), 7.70-7.80 (1H, s), 8.05 (1H, m), 8.15 (1H, m), 8.30 (1H, m), 9.10 (1H, s), 12.2 (1H, s)	4.93
301	406.00	(400 MHz, DMSO-d6): 7.80-7.85 (1H, m), 7.90-7.95 (2H, d), 8.05-8.10 (2H, d), 8.15-8.35 (3H, m), 8.50 (1H, s), 8.60 (1H, s), 9.50-9.55 (1H, s), 9.85-9.90 (1H, s), 12.2 (1H, s)	4.53
302		(400 MHz, MeOH-d4): 2.10-2.0 (1H, m), 2.70-2.80 (1H, m), 2.95-3.05 (1H, m), 3.10-3.20 (1H, m), 5.90-6.00 (1H, t), 7.20-7.40 (4H, m), 8.05-8.10 (1H, s), 8.15-8.20 (2H, d), 8.85 (1H, s)	
303	342.60	(DMSO) 2.9-3.1(4H, m), 7.0-7.5(6H, m), 7.7-7.8(2H, m), 7.8-8.2(6H, m), 11.9(0.7H, s)	4.30
304	346.50	(DMSO) 7.1-7.2(3H, m), 7.2-7.4(3H, m), 7.8-7.9(2H, m), 7.9-8.1(3H, m), 8.1-8.2(1H, s), 8.4(1H, s), 8.5(1H, s), 12.3-12.4(0.7H, s)	4.34
305	398.00	(400 MHz, CDCl3): 1.65-1.70 (3H, d), 5.25-5.30 (1H, m), 5.40-5.45 (1H, m), 6.80-6.85 (1H, d), 7.0 (1H, s), 7.05-7.10 (1H, d), 7.30-7.35 (1H, t), 8.05-8.10 (2H, m), 8.30 (1H, s), 8.70 (1H, s), 9.20-9.50 (1H, brs)	4.87
306	384.00	(400 MHz, MeOH-d4): 1.65-1.70 (3H, d), 5.40-5.50 (1H, m), 6.65-6.80 (1H, d), 6.95 (1H, s), 7.00-7.05 (1H, d), 7.15-7.20 (1H, t), 8.00-8.10 (2H, m), 8.20 (1H, s), 8.70 (1H, s)	4.45

Cmpd No.	LC_MASS _PLUS M+1 (obs)	NMR_RESULT (1H NMR)	RT (mins)
307	410.80	CDCl3 1.7 (3H,d), 3.97 (3H,s), 5.42 (1H, br s), 5.57 (1H,m), 7.03 (2H,t), 7.5-7.6 (2H,m), 8.12 (1H,s), 8.2 (1H,s), 9.1 (1H,s) , 9.52 (1H,s), 10.15 (1H,br s)	4.64
308	405.00	(d6-DMSO, 400MHz) 4.73 (2H, s), 7.38 (2H, d), 7.54 (2H, d), 8.30 (1H, d), 8.50 (1H, s), 8.62 (2H, d)	5.50
309	409.80	DMSO 1.62 (3H,d), 2.88 (3H,m), 5.58 (1H,d), 7.1 (2H,m), 7.62 (2H,m), 8.1-8.25 (3H,m), 8.55 (1H,m), 8.72 (1H,s), 9.15 (1H,s), 12.3 (1H,s)	3.95
310	413.51	1H NMR (DMSO-d6): 1.58 (3H, d), 5.50 (1H, t), 7.67 (2H, d), 7.79 (2H, d), 8.16 (3H, m), 8.21 (2H, d), 8.37 (1H, s)	4.82
311	402.48	1H NMR (DMSO-d6): 1.71 (3H, d), 5.43 (1H, m), 7.28 (1H, s), 7.54 (2H, d), 7.61 (2H, d), 7.66 (1H, s), 7.94 (1H, s), 8.26 (1H, s), 8.47 (1H, s), 12.49 (1H, s)	4.89
312	388.42	1H NMR (DMSO-d6): 4.74 (2H, d), 7.28 (1H, t), 7.38 (2H, m), 7.48 (1H, s), 8.19 (1H, s), 8.23 (2H, m), 8.33 (1H, t), 8.52 (1H, s), 12.33 (1H, s)	4.95
313	383.50	1H NMR (DMSO-d6): 1.66 (3H, d), 5.45 (1H, q), 6.85 (2H, d), 7.31 (1H, s), 8.20 (1H, d), 8.23 (1H, s), 8.54 (2H, d)	4.64
314	389.00	(d6-DMSO, 400MHz) 5.69 (2H, s), 7.49 (2H, d), 7.59 (2H, d), 8.32 (1H, s), 8.42 (1H, s), 8.65 - 8.67 (2H, m), 12.56 (1H, s)	5.32
315	306.46	1H NMR (DMSO-d6): 4.48 (1H, m), 8.23 (1H, s), 8.35 (2H, m), 8.68 (1H, s), 12.59 (1H, s)	4.80
316	355.46	1H NMR (DMSO-d6): 4.87 (2H, d), 7.36 (1H, m), 7.51 (1H, d), 7.87 (1H, m), 8.12 (1H, s), 8.24 (1H, s), 8.29 (1H, d), 8.43 (1H, s), 8.62 (2H, d), 12.39 (1H, s)	4.10
317	355.46	1H NMR (DMSO-d6): 4.86 (2H, d), 7.73 (1H, m), 8.26 (2H, s), 8.30 (1H, d), 8.44 (1H, s), 8.54 (1H, s), 8.65 (1H, d), 8.86 (1H, s) 12.43 (1H, s)	4.10
318	355.46	1H NMR (DMSO-d6): 4.93 (2H, d), 7.90 (2H, d), 8.16 (1H, s), 8.23 (2H, d), 8.31 (1H, s), 8.55 (1H, s), 8.78 (2H, d), 12.37 (1H, s)	4.00
319	231.38	1H NMR (DMSO-d6): 7.27 (1H, t), 8.32 (1H, s), 8.41 (1H, s), 8.80 (3H, m)	4.20
320	455.00	1H NMR (DMSO-d6): 1.55-1.60 (3H, d), 2.15-2.20 (6H, s), 2.55-2.60 (2H, m), 3.95-4.00 (2H, m), 5.35-5.40 (1H, m), 6.60-6.40 (1H, m), 7.00-7.10 (2H, m), 7.15-7.20 (1H, t), 8.05-8.10 (1H, m), 8.15-8.20 (2H, m), 8.25 (1H, s), 8.60 (1H, s), 12.2 (1H, s)	4.49

Cmpd No.	LC_MASS _PLUS M+1 (obs)	NMR_RESULT (1H NMR)	RT (mins)
321	453.00	1H NMR (DMSO-d6): 1.60-1.70 (3H, d), 2.20-2.30 (2H, m), 3.4-3.5 (4H, m), 5.15-5.20 (1H, m), 5.55-5.62 (1H, m), 6.85-6.90 (1H, d), 7.05-7.10 (1H, s), 7.18-7.23 (1H, d), 7.35-7.40 (1H, t), 8.30-8.35 (3H, m), 8.50 (1H, s)	4.25
322	382.00	NMR (DMSO) 1.65 (3H, d, CH3), 2.15 (3H, s, CH3), 5.55 (H, m, alpha), 7.1 (H, d, ar), 7.1-7.2 (2H, m, ar), 7.6 (2H, m, ar), 8.05 (1H, s, ar), 8.2 (1H, s, ar), 8.3 (1H, s,ar), 8.63 (1H, s, ar) and 12.3 (1H, s, ar)	4.99
323	412.00	NMR (DMSO) 1.5 (3H, d, CH3), 5.4 (H, m alpha), 7.0 (H, s, ar), 7.1-7.25 (2H,, m, ar), 7.5-7.6 (2H, m, ar), 8.3 (H, s, ar), 8.4 (2H, brs, ar), 8.65 (H, s, ar) and 12.4 (H, s, ar).	4.10
324	382.00	NMR (DMSO) 1.5 (3H, s, CH3), 2.25 (3H, s, CH3), 5.3 (H, m, alpha), 6.2 (H, m, ar),7.3 (2H, m, ar), 7.5 (2H, m, ar), 7.75 (H, m, ar), 8.2 (H, m, ar), 8.3 (H, m, ar), 8.65 (H, m, ar) and 12.3 (H, m, ar)	4.92
325	361.44	1H NMR (DMSO-d6): 1.53 (3H, m), 1.81 (2H, m), 1.98 (1H, m), 2.90 (1H, m), 3.31 (1H, m), 3.43 (1H, m), 3.58 (2H, m), 7.85 (1H, s), 8.30 (2H, d), 8.36 (1H, s), 8.68 (1H, d), 8.84 (1H, s), 12.52 (1H, s)	3.70
326	361.44	1H NMR (DMSO-d6): 1.34 (1H, m), 1.60 (1H, m), 1.92 (2H, m), 2.20 (1H, m), 2.78 (2H, m), 3.24 (3H, m), 3.54 (1H, m), 7.99 (1H, s), 8.23 (1H, m), 8.30 (2H, d), 8.60 (1H, d), 8.70 (1H, s), 12.45 (1H, s)	3.59
327	361.44, 361.00	1H NMR (DMSO-d6): 1.41 (2H, m), 1.92 (2H, m), 2.08 (1H, m), 2.83 (2H, m), 3.29 (2H, m), 3.44 (2H, m), 7.97 (1H, s), 8.20 (1H, s), 8.25 (1H, s), 8.30 (1H, s), 8.48 (1H, d), 8.71 (1H, s), 12.43 (1H, s),1H NMR (DMSO-d6): 1.41 (2H, m), 1.92 (2H, m), 2.08 (1H, m), 2.83 (2H, m), 3.29 (2H, m), 3.44 (2H, m), 8.20 (1H, s), 8.25 (1H, s), 8.30 (1H, s), 8.48 (1H, d), 8.71 (1H, s), 12.43 (1H, s)	3.57, 3.55
328	347.40	1H NMR (DMSO-d6): 1.94 (1H, m), 1.96 (2H, m), 2.14 (1H, m), 3.20 (2H, m), 3.71 (2H, m), 3.83 (1H, m), 7.89 (1H, s), 8.30 (4H, m), 8.69 (1H, s), 8.91 (1H, s), 12.58 (1H, s)	3.67
329	347.40	1H NMR (DMSO-d6): 1.77 (1H, m), 2.12 (1H, m), 2.77 (1H, m), 3.00 (1H, m), 3.17 (1H, m), 3.34 (2H, m), 3.60 (2H, m), 7.98 (1H, s), 8.23 (1H, d), 8.31 (2H, m), 8.71 (3H, m), 12.53 (1H, s)	3.54
330	394.45	1H NMR (DMSO-d6): 2.04 (4H, m), 2.84 (2H, m), 3.33 (1H, s), 5.61 (1H, m), 7.18 (3H, m), 7.27 (1H, d), 8.00 (1H, d), 8.21 (1H, t), 8.23 (1H, s), 8.26 (1H, s), 8.68 (1H, s), 12.34 (1H, s)	5.28

Cmpd No.	LC_MASS _PLUS M+1 (obs)	NMR_RESULT (1H NMR)	RT (mins)
331	483.00	(d6-DMSO, 400 MHz) 1.44 (9H, s), 1.54 (3H, d), 5.34 - 5.37 (1H, m), 7.11 (1H, brs), 7.19 (2H, d), 7.63 (1H, s), 8.10 - 8.12 (2H, m), 8.18 (1H, d), 8.23 (1H, d), 8.54 (1H, d), 9.27 (1H, brs), 12.29 (1H, brs)	4.95
332	264.00	NMR (DMSO) 7.3 (2H, brs, NH2), 8.1 (1H, s, ar), 8.2 (1H, s, ar), 8.3 (1H, s, ar), 8.95 (1H, s, ar) and 12.35 (1H, s, NH).	3.87
333	383.00	(d6-DMSO, 400MHz) 1.55 (3H, d), 5.36 - 5.40 (1H, m), 6.76 (1H, brs), 6.98 (1H, brs), 7.21 (2H, brs), 8.16 (1H, d), 8.23 (1H, d), 8.26 (2H, d), 8.58 (1H, s), 12.36 (1H, s)	4.42
334	382.80	DMSO 1.58 (3H,d), 4.65 (2H,d), 5.28 (1H,t), 5.5-5.6 (1H,m), 7.15 (2H,m), 7.6 (2H,m), 8.05 (1H,m), 8.2 (1H,s), 8.26 (1H,m), 8.68 (1H,s), 11.95 (1H,s)	3.90
335	456.00	1H NMR (MeOD-d4): 1.60-1.70 (3H, d), 3.60 (3H, s), 4.65 (2H, s), 5.40-5.45 (1H, m), 6.70-6.75 (1H, d), 7.03 (1H, s), 7.12-7.18 (1H, d), 7.25-7.30 (1H, t), 8.00-8.07 (2H, m), 8.15 (1H, s), 8.60 (1H, s)	4.70
336	442.00	1H NMR (DMSO-d6): 1.35-1.40 (3H, d), 4.00-4.05 (2H, s), 5.25-5.35 (1H, m), 6.50-6.55 (1H, d), 6.95-7.00 (2H, m), 7.05-7.15 (1H, m), 8.05-8.13 (3H, m), 8.18 (1H, s), 8.55 (1H, s), 12.5 (1H, s)	3.82
337	363.00	1H NMR (CD3OD): 1.60-1.70 (3H, d),3.10-3.20 (3H, s), 3.35-3.45 (3H, s), 5.30-5.35 (1H, m), 8.15-8.20 (2H, m), 8.30-8.35 (1H, s), 8.75-8.80 (1H, s)	4.07
338	375.00	1H NMR (DMSO-d6): 0.30-0.40 (1H, m), 0.42-0.47 (1H, m), 1.35-1.40 (3H, d), 2.55-2.60 (1H, m), 4.40-4.50 (1H, m), 7.58-7.63 (1H, d), 8.15-8.25 (3H, m), 8.28 (1H, s), 8.65 (1H, s), 12.0 (1H, s)	3.84
339	398.47	1H NMR (CDCl3): 1.63 (3H, m), 3.44 (1H, s), 4.57 (2H, m), 5.31 (1H, br s), 6.02 (1H, br s), 7.06 (2H, m), 7.39 (2H, m), 8.19 (1H, m), 8.30 (1H, m), 8.76 (1H, br s), 8.93 (1H, br s)	4.59
340	349.00, 349.00	1H NMR (DMSO-d6): 1.35-1.40 (3H, d), 2.55-2.60 (3H, m), 4.40-4.50 (1H, m), 7.58-7.63 (1H, d), 8.00-8.05 (1H, m), 8.15-8.25 (2H, m), 8.28 (1H, s), 8.65 (1H, s), 12.0 (1H, s),1H NMR (DMSO-d6): 1.35-1.40 (3H, d), 2.55-2.60 (3H, m), 4.40-4.50 (1H, m), 7.58-7.63 (1H, d), 8.00-8.05 (1H, m), 8.15-8.25 (2H, m), 8.28 (1H, s), 8.65 (1H, s), 12.0 (1H, s)	3.59, 2.84

Cmpd No.	LC_MASS _PLUS M+1 (obs)	NMR_RESULT (1H NMR)	RT (mins)
341	428.00	1H NMR (DMSO-d6): 1.35-1.40 (3H, d), 3.6-3.7 (2H, m), 3.95-4.00 (2H, m), 4.80-4.90 (1H, m), 5.4-5.5 (1H. m), 6.7-6.8 (1H, m), 7.00-7.10 (2H, m), 7.20-7.25 (1H, m), 8.10-8.25 (3H, m), 8.65 (1H, s), 12.0 (1H, s)	4.47
342		(400 MHz, DMSO) 1.57 (3H, d), 5.50-5.57 (1H, m), 7.11-7.15 (2H, m), 7.57-7.60 (2H, m), 8.20 (1H, d), 8.25 (1H, d), 8.36 (1H, s), 9.15 (1H, d), 9.45 (1H, d), 12.90 (1H, brs).	
343		(400 MHz, DMSO) 1.55 (3H, d), 4.85 (2H, s), 5.45-5.53 (1H, m), 7.09-7.14 (2H, m), 7.55-7.58 (2H, m), 7.62 (1H, s), 7.72 (1H, d), 7.87 (1H, s), 7.95 (1H, d), 8.10 (1H, d), 11.55 (1H, s).	
344	333.00	1H NMR (CD3OD): 2.40-2.50 (2H, m), 2.6-2.7 (2H, m), 3.50-3.60 (3H, m), 3.7-3.8 (1H, m), 5.2-5.3 (1H, m), 8.40 (1H, s), 8.50 (1H, m), 8.55 (1H, s), 8.75 (1H, s)	3.54
345	347.00, 347.40	1H NMR (DMSO-d6): 1.80-1.90 (2H, m), 2.1-2.2 (2H, m), 3.05-3.15 (2H, m), 3.4-3.5 (2H, m), 4.30-4.40 (1H, m), 7.65-7.70 (1H, m), 8.30-8.35 (2H, m), 8.40-8.50 (1H, s), 8.60-8.70 (1H, s), 8.75-8.80 (1H, m), 12.0 (1H, s),1H NMR (DMSO-d6): 1.84 (2H, m), 2.33 (2H, m), 3.14 (2H, m), 3.44 (2H, m), 4.36 (1H, m), 7.88 (1H, s), 8.25 (1H, s), 8.30 (2H, d), 8.37 (1H, m), 8.60 (1H, m), 8.69 (1H, s), 12.44 (1H, s)	3.50, 3.50
346	423.56	1.73 (3 H, d, J 6.8, Me), 3.06 (3 H, br s, Me), 3.24 (3 H, br s, Me), 5.53-5.47 (1 H, m, CH), 5.75 (1 H, br s, NH), 7.07-7.03 (2 H, m, 2 x ArH), 7.46-7.43 (2 H, m, 2x ArH), 8.14 (1 H, d, J 3.7, ArH), 8.46 (1 H, d, J 1.8, ArH), 8.77 (2 H, s, 2 x ArH).	4.02
347	347.47	1H NMR (DMSO-d6): 1.66 (1H, m), 1.85 (1H, m), 1.98 (1H, m), 2.11 (1H, m), 2.90 (2H, m), 3.32 (1H, m), 3.47 (1H, m), 4.46 (1H, m), 7.57 (1H, d), 8.31 (3H, m), 8.66 (3H, m), 12.47 (1H, s)	3.68
348	348.44	1H NMR (DMSO-d6): 1.67 (2H, m), 1.96 (2H, d), 3.49 (2H, t), 3.96 (2H, d), 4.30 (1H, m), 7.62 (1H, d), 8.18 (1H, s), 8.22 (1H, s), 8.29 (1H, s), 8.72 (1H, s),12.36 (1H, s)	4.42
349	360.46	1H NMR (DMSO-d6): 1.17 (5H, m), 1.70 (6H, m), 2.67 (2H, d), 8.27 (1H, s), 8.42 (2H, s), 8.72 (1H, s), 12.57 (1H, s)	5.39
350	396.46	(DMSO, D6); 1.55 (3 H, d, J 6.8, Me), 5.53 (1 H, t, J 6.8, CH), 7.13-7.09 (2 H, m, 2 x ArH), 7.63-7.60 (2 H, m, 2 x ArH), 8.15-8.14 (1 H, m, NH), 8.22-8.19 (2 H, m, 2 x ArH), 8.83 (1 H, s, ArH), 9.30 (1 H, s, ArH).	3.27
351	392.00	1H NMR (DMSO-d6): 1.35-1.40 (9H, s), 1.50-1.55 (3H, d), 4.10-4.15 (1H, m) 4.40-4.50 (1H, m), 7.95-8.00 (1H, d), 8.20 (1H, s), 8.30 (1H, m), 8.35 (1H, m), 8.75 (1H, s)	4.87

Cmpd No.	LC_MASS _PLUS M+1 (obs)	NMR_RESULT (1H NMR)	RT (mins)
352	347.00	(d6-DMSO, 400MHz) 1.86 - 1.94 (1H, m), 2.28 (3H, s), 2.33 (1H, brs), 2.53 - 2.57 (1H, m), 2.60 - 2.67 (2H, m), 2.84 - 2.88 (1H, m), 4.62 (1H, brs), 7.67 (1H, d), 8.17 (1H, d), 8.21 (1H, s), 8.28 (1H, d), 12.35 (1H, s)	4.00
353	255.00	DMSO D6 7.60 (1H, d), 8.32 (1H, s), 8.43 (1H, s), 8.56 (1H, s), 8.82-8.86 (2H, m) 12.55 (1H, s).	4.52
354	382.46	1H NMR (DMSO-d6): 0.95 (3H, t), 1.85 (1H, m), 1.97 (1H, m), 7.19 (1H, t), 7.34 (2H, t), 7.53 (2H, d), 8.08 (1H, d), 8.16 (2H, s), 8.26 (1H, s), 8.66 (1H, s),12.31 (1H, s)	4.92
355	374.50	1H NMR (DMSO-d6): 1.15 (8H, m), 1.28 (2H, m), 1.72 (2H, m), 1.84 (2H, m), 4.23 (1H, m), 7.43 (1H, d), 8.12 (1H, s), 8.18 (1H, s), 8.27 (1H, s), 8.73 (1H, s),12.32 (1H, s)	5.47
356	346.43	1H NMR (DMSO-d6): 0.85 (1H, m), 1.48 (4H, m), 1.68 (1H, d), 1.81 (2H, m), 2.04 (2H, m), 4.03 (1H, m), 7.49 (1H, d), 8.13 (1H, s), 8.19 (1H, s), 8.29 (1H, s), 8.73 (1H, s), 12.21 (1H, s)	5.25
357	432.70	CDCl3 0.8 (3H,d), 5.45 (1H,m), 5.6 (1H, br s), 7.1 (2H,m), 7.45 (2H,m), 8.15 (1H,s), 8.4 (1H,s), 8.5-8.6 (1H, br s), 8.78 (1H,s), 9.65 (1H, br s)	5.00
358	439.00	(DMSO) 1.27 (3 H, t, J 7.1), 1.56 (3 H, d, J 7.0), 4.17 (2 H, q, J 7.1), 5.64 (1 H, t, J 7.0), 7.10 (2 H, t, J 8.9), 7.60-7.56 (2 H, m), 8.04-8.00 (2 H, m), 8.15 (1 H, d, J 3.8), 8.22 (1 H, d, J 2.1), 8.96 (1 H, br s), 9.65 (1 H, br s), 11.96 (1 H, br s).	4.39
359	425.51	1H NMR (DMSO): 1.52 (3H, m), 2.85 (3H, m), 5.33 (1H, m), 7.00 (1H, s), 7.15 (2H, m), 7.48 (2H, m), 8.24 (2H, m), 8.49 (1H, s), 8.66 (1H, s), 8.87 (1H, m), 12.4 (1H, br s)	4.82
360	336.00	1H NMR (DMSO-d6): 1.45-1.50 (3H, d), 4.50-4.60 (1H, m), 8.05-8.25 (5H, m), 8.60 (1H, m), 12.3-12.4 (1H, s), 12.50-1.270 (1H, brs)	3.20
361	363.00	1H NMR (CD3OD): 1.10-1.20 (3H, t), 1.90-2.20 (2H, m), 2.85 (3H, s), 4.65-4.70 (1H, m), 8.25-8.35 (3H, m), 8.65 (1H, s)	3.92
362	377.00	1H NMR (CD3OD): 1.10-1.20 (6H, m), 2.3-2.4 (1H, m), 2.80 (3H, s), 4.70-4.75 (1H, d), 8.25-8.35 (3H, m), 8.80 (1H, s)	4.27
363	377.00	1H NMR (CD3OD): 1.05-1.15 (6H, m), 1.50-1.60 (3H, m), 3.95-4.05 (1H, m), 4.50-4.65 (1H, m), 8.10-8.25 (3H, m), 8.80 (1H, s)	4.07
364	391.00	1H NMR (CD3OD): 0.70-0.80 (6H, m), 1.55-1.60 (3H, d), 1.70-1.80 (1H, m), 2.85-2.95 (1H, m), 3.10-3.15 (1H, m), 4.50-4.65 (1H, m), 8.10-8.25 (3H, m), 8.80 (1H, s)	4.22

Cmpd No.	LC_MASS _PLUS M+1 (obs)	NMR_RESULT (1H NMR)	RT (mins)
365	381.00	1H NMR (CD3OD): 1.55-1.60 (3H, d), 3.45-3.55 (2H, m), 4.25-4.35 (1H, m), 4.40-4.45 (1H, m), 4.55-4.65 (1H, m), 8.10-8.25 (3H, m), 8.80 (1H, s)	3.79
366	362.00	(d6-DMSO, 400MHz) 1.23 - 1.33 (2H, m), 1.72 (2H, d), 1.99 - 2.04 (1H, m), 3.27 (2H, t), 3.41 (2H, t), 3.85 - 3.89 (2H, m), 7.80 (1H, t), 8.14 (1H, d), 8.21 (1H, s), 8.28 (1H, d), 8.74 (1H, d), 12.35 (1H, brs)	4.50
367	363.00	1H NMR (DMSO-d6): 0.95-1.05 (3H, t), 1.35-1.40 (3H, d), 3.00-3.10 (2H, m), 4.45-4.55 (1H, m), 7.55-7.60 (1H, d), 8.05-8.10 (1H, m), 8.18-8.22 (2H, m), 8.28 (1H, s), 8.70 (1H, s),	3.92
368	456.00, 456.07	1H NMR (DMSO-d6): 1.40-1.50 (3H, d), 2.80 (3H, s), 2.85-3.05 (2H, m), 3.15-3.25 (2H, m), 4.50-4.60 (1H, m), 7.00-7.05 (1H, m), 7.60-7.65 (1H, d), 8.12-8.17 (1H, m), 8.19 (1H, m), 8.22 (1H, m), 8.28 (1H, s), 8.70 (1H, s),,1H NMR (DMSO): 1.45 (3H, d), 2.89 (2H, m), 3.15 (2H, m), 3.31 (3H, s), 4.55 (1H, m), 7.04 (1H, s), 7.69 (1H, d), 8.19 (2H, m), 8.26 (1H, s), 8.67 (1H, s), 12.35 (1H, s)	3.57, 2.84
369	417.00, 416.90	1H NMR (CD3OD): 1.55-1.60 (3H, d), 3.85-4.00 (2H, m), 4.70-4.80 (1H, m), 8.10-8.25 (3H, m), 8.80 (1H, s),DMSO d6 12.4 (bs, 1H); 8.8 (t, 1H); 8.7 (s, 1H); 8.3 (s, 2H); 8.2 (s, 1H); 7.9 (bs, 1H); 4.7 (q, 1H); 4.0 (m, 2H); 1.3 (d, 3H)	4.09, 2.30
370	412.44	1H NMR (DMSO): 1.60 (3H, m), 5.51 (1H, br s), 7.08-7.21 (2H, m), 7.49-7.53 (2H, m), 8.29 (1H, m), 8.39 (1H, m), 8.53 (1H, m), 8.79 (2H, m), 12.6 (1H, s)	4.22
371	436.48	1H NMR (CDCl3): 1.64 (3H, m), 5.31 (1H, m), 5.77 (1H, m), 7.05 (2H, m), 7.40 (2H, m), 7.95 (1H, m), 8.32 (1H, br s), 8.58 (1H, m), 8.94 (1H, m)	5.17
372	375.00	(d6-DMSO, 400MHz) 1.14 - 1.24 (2H, m), 1.75 - 1.80 (5H, m), 2.11 (3H, s), 2.75 (2H, d), 3.40 (2H, t), 7.80 (1H, t), 8.13 (1H, d), 8.20 (1H, s), 8.28 (1H, d), 8.73 (1H, d), 12.35 (1H, s)	3.95
373	348.00	(d6-DMSO, 400MHz) 1.66 - 1.74 (1H, m), 1.99 - 2.08 (1H, m), 2.67 - 2.74 (1H, m), 3.49 - 3.51 (2H, m), 3.58 - 3.67 (2H, m), 3.73 (1H, t), 3.79 - 3.84 (1H, m), 7.88 (1H, t), 8.16 (1H, d), 8.22 (1H, s), 8.29 (1H, d), 8.74 (1H, d), 12.36 (1H, brs)	3.27
374	376.70	CDCl3 1.75 (3H,d), 3.23 (1H,s), 5.30-5.35 (1H,m), 5.5-5.56 (1H,m), 7.08-7.18 (2H,m), 7.5-7.58 (2H,m), 8.22 (2H,m), 8.55 (1H, br s), 8.9 (1H,s), 10.65 (1H, br s)	3.54

Cmpd No.	LC_MASS _PLUS M+1 (obs)	NMR_RESULT (1H NMR)	RT (mins)
375		1H NMR (CD3OD): 1.68-1.73 (3H, d), 4.50 (2H, s), 5.40-5.50 (1H, m), 6.7-6.8 (1H, m), 7.05-7.10 (1H, m), 7.15-7.20 (1H, d), 7.30-7.40 (1H, t), 8.30-8.40 (2H, m), 8.45 (1H, s), 8.50 (1H, s)	3.25
376	455.00	1H NMR (CD3OD): 1.68-1.73 (3H, d), 2.75-2.80 (3H, s), 4.50 (2H, s), 5.40-5.50 (1H, m), 6.8-6.9 (1H, m), 7.05-7.10 (1H, m), 7.15-7.20 (1H, d), 7.30-7.40 (1H, t), 8.05-8.10 (2H, m), 8.20 (1H, s), 8.60 (1H, s)	3.38
377	469.00	1H NMR (CD3OD): 1.68-1.73 (3H, d), 2.15-2.20 (2H, m), 2.90-2.95 (6H, s), 3.10-3.20 (2H, m), 4.05-4.10 (2H, m), 5.50-5.60 (1H, m), 6.8-6.9 (1H, m), 7.05-7.10 (1H, m), 7.15-7.20 (1H, d), 7.30-7.40 (1H, t), 8.20-8.24 (1H, d), 8.25-8.30 (2H, m), 8.60 (1H, s)	3.29
378	406.75	CDCl3 1.75 (3H,d), 3.55 (1H,s), 4.65 (2H,s), 5.30-5.35 (1H,m), 5.42-5.48 (1Hm), 7.1-7.18 (2H,m), 7.45-7.52 (2H,m), 8.1-8.18 (2H,m), 8.45 (1H, br s), 8.78 (1H,s), 9.4 (1H, br s)	3.15
379	347.47	1H NMR (CDCl3 / MeOD): 0.83 (2H, m), 1.94 (1H, m), 2.32 (1H, m), 3.00 (1H, m), 3.30 (1H, m), 3.36 (2H, m), 3.46 (1H, m), 3.60 (1H, m), 3.87 (1H, m), 8.15 (1H, s), 8.24 (1H, s), 8.29 (1H, s), 8.68 (1H, s)	3.54
380	333.51	1H NMR (DMSO-d6): 2.15 (1H, m), 2.30 (2H, m), 3.35 (2H, m), 3.58 (1H, m), 4.77 (1H, m), 7.87 (1H, s), 8.29 (3H, m), 8.81 (1H, s), 8.94 (2H, br s), 12.45 (1H, s)	5.00
381	333.40	1H NMR (DMSO-d6): 2.17 (1H, m), 2.34 (1H, m), 3.34 (3H, m), 3.58 (1H, m), 4.79 (1H, m), 7.87 (1H, d), 8.27 (3H, m), 8.68 (1H, s), 8.81 (2H, br s), 12.45 (1H, s)	5.00
382	302.37	1H NMR (DMSO-d6): 3.16 (1H, s), 4.29 (2H, m), 8.29 (4H, m), 8.88 (1H, s), 12.40 (1H, s)	3.22
383	403.48	1H NMR (DMSO-d6): 0.82 (2H, m), 1.11 (1H, m), 1.22 (1H, m), 1.85 (2H, t), 1.98 (3H, s), 2.07 (1H, br s), 3.00 (1H, t), 3.51 (1H, s), 3.83 (1H, m), 4.40 (1H, d), 8.40 (1H, s), 8.44 (1H, d), 8.65 (1H, s), 8.88 (1H, s), 9.20 (1H, br s), 12.91 (1H, s)	3.18
384	420.80	CDCl3 1.72 (3H,d), 3.55 (3H,s), 4.45 (2H,s), 5.32-5.36 (1H,m), 5.47-5.53 (1H,m), 7.05-7.1 (2H,m), 7.45-7.5 (2H,m), 8.15 (2H,m), 8.5 (1H, br s), 8.85 (1H,s), 10.05 (1H, br s)	3.59
385	391.00	1H NMR (CD3OD): 1.15-1.20 (9H, s), 2.80 (3H, s), 4.70-4.75 (1H, s), 8.05-8.10 (1H, m), 8.12-8.15 (2H, m), 8.85 (1H, s)	3.50

Cmpd No.	LC_MASS _PLUS M+1 (obs)	NMR_RESULT (1H NMR)	RT (mins)
386	396.53	1H NMR (DMSO): 1.18 (3H, t), 1.55 (3H, d), 2.56 (2H, q), 5.51 (1H, m), 7.07-7.15 (3H, m), 7.52 (2H, m), 7.98 (1H, s), 8.14 (1H, s), 8.23 (1H, d), 8.60 (1H, d), 12.22 (1H, s)	3.88
387	365.00	(d6-DMSO, 400MHz) 2.61 (3H, d), 3.82 - 3.83 (2H, m), 4.60 (1H, dd), 5.02 (1H, t), 7.32 (1H, d), 8.05 (1H, d), 8.20 (1H, s), 8.24 (1H, d), 8.27 (1H, d), 8.66 (1H, d), 12.36 (1H, brs)	2.55
388	379.00	(d6-DMSO, 400MHz) 2.61 (3H, d), 3.30 (3H, s), 3.74 (2H, brs), 4.82 (1H, dd), 7.54 (1H, d), 8.13 (1H, d), 8.23 (2H, brs), 8.27 (1H, s), 8.66 (1H, s), 12.36 (1H, brs)	2.82
389	441.00	1H NMR (CD3OD): 1.68-1.73 (3H, d), 2.80-2.85 (3H, s), 3.10-3.25 (2H, m), 3.35-3.55 (2H, m), 4.55-4.60 (1H, m), 8.10 (1H, m), 8.15 (1H, s), 8.20 (1H, s), 8.75 (1H, s)	2.68
390	453.00	1H NMR (CD3OD): 1.68-1.73 (3H, d), 2.10-2.50 (2H, m), 2.90-3.20 (4H, m), 4.50-4.60 (2H, m), 8.10-8.15 (2H, m), 8.20 (1H, s), 8.80 (1H, s)	1.74
391	379.00	1H NMR (CD3OD): 1.68-1.73 (3H, d), 3.35-3.40 (2H, m), 3.50-3.60 (2H, m), 4.60-4.70 (1H, m), 8.10 (1H, m), 8.15 (1H, s), 8.20 (1H, s), 8.80 (1H, s)	2.63
392	393.00	1H NMR (CD3OD): 1.68-1.73 (3H, d), 3.15-3.20 (3H, s), 3.30-3.60 (4H, m), 4.60-4.70 (1H, m), 8.10 (1H, m), 8.15 (1H, s), 8.20 (1H, s), 8.80 (1H, s)	2.80
393	391.00	(d6-DMSO, 400MHz) 0.89 (3H, d), 0.94 (3H, d), 1.58 - 1.81 (3H, m), 2.60 (3H, d), 4.61 - 4.67 (1H, m), 7.59 (1H, d), 8.03 (1H, d), 8.21 (1H, d), 8.22 (1H, s), 8.27 (1H, d) 8.68 (1H, d), 12.36 (1H, s)	3.33
394		(400 MHz, DMSO) 1.56 (3H, d), 5.41-5.48 (1H, m), 7.15 (2H, t), 7.56 (2H, dd), 8.17 (1H, d), 8.21 (1H, d), 8.28 (1H, s), 8.64 (1H, d), 8.84 (1H, d), 12.75 (1H, brs).	
395	367.00	(400 MHz, DMSO) 1.50 (3H, d), 5.10-5.14 (1H, m), 6.32 (1H, d), 6.99 (1H, d), 7.08-7.14 (3H, m), 7.34 (1H, t), 7.46-7.49 (2H, m), 8.14 (1H, s), 8.20 (1H, d), 8.64 (1H, s), 12.11 (1H, brs).,MeOD-D4: 8.35 (m, 2H); 8.1 (s, 1H); 7.95 (dd, 1H); 7.5 (m, 2H); 7.2 (d, 1H); 7.15 (m, 2H); 6.8 (d, 1H); 5.0 (m, 1H); 1.65 (d, 3H).	2.50
396		(400 MHz, DMSO) 1.60 (3H, d), 3.88 (3H, s), 7.15 (2H, t), 7.46-7.49 (2H, m), 7.98 (1H, d), 8.25-8.30 (2H, m), 8.34 (1H, d), 8.56 (1H, s), 12.65 (1H, brs).	
397		(400 MHz, DMSO) 1.51 (3H, s), 5.31-5.36 (1H, m), 7.12 (2H, t), 7.41-7.46 (3H, m), 7.80-8.08 (2H, m), 8.17 (1H, d), 11.98 (1H, brs).	

Cmpd No.	LC_MASS _PLUS M+1 (obs)	NMR_RESULT (1H NMR)	RT (mins)
398		(400 MHz, DMSO)1.51 (3H, d), 5.33-5.38 (1H, m), 7.14 (2H, t), 7.42-7.45 (2H, m), 7.70-7.76 (3H, m), 7.88 (1H, d), 8.21 (1H, s), 12.20 (1H, brs).	
399	339.00	DMSO D6 2.78-2.92 (4H, m), 3.50-3.62 (4H, m), 7.05 (1H, s), 7.54 (1H, s), 8.27 (1H, s), 8.45 (1H, s), 8.68 (1H, s)	1.24
400	403.00	(d6-DMSO, 400MHz) 3.90 - 3.95 (2H, m), 4.12 (2H, d), 7.97 (1H, t), 8.15 (1H, s), 8.24 (1H, d), 8.27 (1H, d), 8.63 (1H, d), 8.75 (1H, t), 12.34 (1H, brs)	2.93
401	350.00	NMR (DMSO) 1.3 (3H, d, CH3), 2.6-2.8 (2H, m, CH2), 4.65 (H, m, alpha), 7.15 (H, m, ar), 8.15 (H, m, ar), 8.25 (2H, s, ar), 8.7 (H, s, NH), 12.1-12.5 (2H, NH and COOH, brs x 2)	2.59
402	363.00	NMR (DMSO) 1.25 (3H, d, CH3), 1.75 (3H, m, CH3), 3.0 (2H, CH2, m), 4.65 (H, m, alpha), 7.5 (H, d, ar), 8.0 (H, m, NH), 8.1 (H, d, ar), 8.15 (H, s, ar), 8.25 (H, s, ar), 8.7 (H, s, NH) and 12.45 (H, s, NH).	2.90
403	379.38	1H NMR (DMSO-d6): 1.73 (2H, m), 1.99 (2H, m), 2.84 (2H, m), 4.55 (1H, m), 7.74 (2H, br s), 7.95 (1H, d), 8.19 (1H, s), 8.28 (2H, m), 8.73 (1H, s), 12.38 (1H, s), 12.99 (1H, br s)	2.21
404	418.49	1H NMR (DMSO-d6): 1.74 (3H, m), 5.82 (1H, m), 7.20 (2H, m), 7.61 (2H, m), 7.70 (1H, s), 7.84 (1H, s), 7.99 (1H, s), 8.39 (1H, s), 8.54 (1H, s), 8.66 (1H, s), 8.79 (1H, br s), 9.82 (1H, br s), 13.08 (1H, br s)	3.85
405	344.00	2.82-2.88 (4H, m), 3.47-3.52 (4H, m), 4.50 (2H, s), 5.31 (1H, br s), 6.61 (1H, s), 7.15 (1H, s), 8.20 (1H, s), 8.25 (1H, s), 8.70 (1H, s), 12.25 (1H, br s)	2.34
406	417.00	1H NMR (CD3OD): 1.1-1.4 (6H, m), 1.6-2.0 (7H, m), 3.0-3.15 (1H, m), 3.65-3.80 (1H, m), 5.6-5.7 (1H, m), 8.15 (1H, m), 8.20-8.25 (2H, m), 8.80 (1H, s)	3.37
407	429.00	1H NMR (DMSO-d6): 1.45-1.50 (3H, d), 3.80-4.00 (5H, m), 4.60-4.65 (1H, m), 6.80-6.85 (1H, m), 7.95 (1H, s), 8.10 (1H, s), 8.25 (1H, s), 8.70 (2H, m), 12.2 (1H, s)	3.13
408	443.00	1H NMR (DMSO-d6): 1.37-1.47 (6H, m), 3.85-4.05 (2H, m), 4.10-4.20 (2H, qd), 4.60-4.65 (1H, m), 6.65-6.70 (1H, d), 7.95 (1H, s), 8.10 (1H, s), 8.25 (1H, s), 8.70-8.80 (2H, m), 12.2 (1H, s)	3.33
409	342.80	DMSO D6 3.22-3.30 (4H, m), 3.86-3.91 (4H, m), 7.22 (1H, s), 7.69 (1H, s), 8.33 (1H, s), 8.46 (1H, s), 8.68 (1H, s), 10.07 (1H, s), 12.48 (1H, s)	2.80
410	457.00	1H NMR (DMSO-d6): 1.35-1.40 (6H, m), 1.45-1.50 (3H, d), 3.80-4.10 (2H, m), 4.60-4.65 (2H, m), 6.50-6.60 (1H, m), 7.95 (1H, s), 8.10 (1H, s), 8.25 (1H, s), 8.70-8.80 (2H, m), 12.5 (1H, s)	3.46

Cmpd No.	LC_MASS _PLUS M+1 (obs)	NMR_RESULT (1H NMR)	RT (mins)
411	423.00	1H NMR (DMSO-d6): 1.45-1.50 (3H, d), 3.80-4.00 (2H, m), 4.50-4.60 (1H, m), 4.75 (1H,s), 8.05 (1H, m), 8.30-8.40 (2H, m), 8.50-8.60 (1H, m), 8.95-9.000 (1H, m), 9.20 (1H, m), 12.5 (1H, s)	3.22
412	338.70	DMSO D6 2.80-2.90 (4H, m), 3.48-3.61 (4H, m), 4.43 (1H, s), 6.66 (1H, s), 7.25 (1H, s), 8.27 (1H, s), 8.35 (1H, s), 8.66 (1H, s), 12.28 (1H, br s).	3.16
413	399.00	1H NMR (CD3OD): 1.5-1.6 (3H, d), 3.85-4.10 (2H, m), 4.60-4.75 (1H, m), 6.40-6.50 (1H, s), 8.15 (1H, d), 8.20-8.25 (2H, m), 8.90 (1H, s)	2.93
414	497.00	1H NMR (CD3OD): 1.5-1.6 (3H, d), 3.00-3.10 (3H, s), 3.30-3.60 (6H, brm), 3.90-4.05 (2H, m), 4.45-4.55 (1H, m), 6.50-6.60 (1H, s), 8.20 (1H, s), 8.30 (1H, s), 8.50 (1H, s), 8.70-8.80 (1H, m)	3.08
415	583.00	1H NMR (DMSO-d6): 1.30-1.35 (3H, d), 1.40 (9H, s), 3.35-3.40 (4H, m), 3.65-3.72 (4H, m), 3.9-4.0 (2H, m), 4.40-4.50 (1H, m), 6.35-6.40 (1H,s), 7.10-7.20 (1H, m), 8.15 (1H, s), 8.25 (1H, s), 8.50 (1H, s), 8.60-8.70 (1H, m), 12.5 (1H, s)	3.69
416	483.00	1H NMR (CD3OD): 1.50-1.60 (3H, d), 3.40-3.45 (4H, m), 3.95-4.05 (2H, m), 4.10-4.20 (2H, m), 4.6-4.7 (1H, m), 6.55 (1H, s), 8.35 (1H, s), 8.40 (2H, m), 8.90 (1H, s)	2.73
417	458.00	1H NMR (CD3OD): 1.40-1.50 (3H, d), 3.50-3.55 (2H, m), 3.75-3.85 (2H, m), 3.95-4.05 (2H, m), 4.10-4.20 (2H, m), 4.45-4.55 (1H, m), 6.30-6.35 (1H, s), 8.00-8.05 (1H, s), 8.25 (2H, m), 8.75 (1H, s)	2.88
418	467.74	1H NMR (DMSO-d6): 1.43 (3H, d), 3.93 (2H, m), 4.56 (1H, m), 8.04 (1H, s), 8.35 (1H, d), 8.39 (1H, d), 8.57 (1H, m), 8.66 (1H, m), 8.99 (1H, s), 12.67 (1H, s)	3.35
419	485.00	1H NMR (CDCl3): 1.50-1.55 (3H, d), 2.25-2.30 (6H, s), 2.5-2.7 (2H, m), 3.50-3.60 (2H, m), 3.80-4.10 (2H, m), 4.50-4.60 (1H, m), 5.55-5.60 (1H, m), 5.90 (1H, s), 7.60-7.90 (2H, m), 8.20 (1H, s), 8.45-8.55 (1H, brs), 10.5-10.8 (1H, brs)	2.82
420	424.80	1H NMR (DMSO): 1.44 (3H, d), 3.91 (2H, m), 4.58 (1H, m), 8.38 (1H, m), 8.66-8.72 (3H, m), 9.20 (1H, d)	3.12
421	453.77	1H NMR (DMSO-d6): 3.92 (2H, m), 4.06 (2H, s), 8.06 (1H, s), 8.33 (1H, d), 8.44 (1H, s), 8.59 (2H, m), 8.95 (1H, s), 12.76 (1H, s)	3.22
422	413.02	1H NMR (DMSO): 1.68 (3H, d), 5.67 (1H, m), 7.19 (2H, m), 7.61 (2H, m), 8.32 (1H, s), 8.47 (2H, m), 8.85 (1H, m), 9.18 (1H, s), 12.83 (1H, br s)	3.90

Cmpd No.	LC_MASS _PLUS M+1 (obs)	NMR_RESULT (1H NMR)	RT (mins)
423	383.76	1H NMR (DMSO): 1.65 (2H, m), 5.56 (1H, m), 5.75 (1H, br s), 7.22 (2H, m), 7.50(2H, m), 7.56 (1H, s), 8.35 (2H, s), 8.39 (1H, m), 12.75 (1H, br s), 13.5 (1H, s)	3.45
424	481.00	1H NMR (DMSO-d6): 1.45-1.50 (3H, d), 3.05-3.20 (3H, m), 3.80-4.00 (2H, m), 5.30-5.50 (1H, m), 8.00-8.10 (1H, m), 8.30-8.40 (1H, m), 8.50-8.80 (2H, m), 12.5 (1H, s)	3.67
425	439.81	1H NMR (DMSO): 1.52 (3H, d), 2.99 (6H, m), 5.36 (1H, m), 6.43 (1H, br s), 7.16 (2H, m), 7.49 (2H, m), 8.11 (1H, m), 8.22 (2H, m), 8.51 (1H, br s), 12.36 (1H, s)	3.56
426	478.00	1H NMR (DMSO-d6): 1.28 (9H, s), 1.40-1.50 (3H, d), 2.90-3.15 (4H, m), 4.50-4.60 (1H, m), 6.70-6.80 (1H, m), 7.60-7.65 (1H, d), 8.08-8.13 (1H, m), 8.15-8.20 (2H, m), 8.25 (1H, m), 8.70 (1H, s), 12 (1H, s)	3.23
427	435.82	1H NMR (DMSO-d6): 3.92 (2H, m), 4.24 (2H, s), 7.43 (1H, m), 7.59 (1H, m), 7.77 (2H, m), 8.22 (2H, s), 8.78 (1H, m), 8.81 (1H, m), 8.92 (1H, s), 12.35 (1H, s)	3.28
428	293.63	1H NMR (DMSO-d6): 1.25 (3H, t), 2.93 (2H, q), 8.34 (1H, s), 8.37 (1H, s), 8.57 (1H, s), 8.07 (1H, s), 12.84 (1H, br s)	3.28
429	441.00	1H NMR (CD3OD): 1.30-1.50 (6H, dd), 1.60-1.70 (3H, d), 3.05-3.15 (1H, m), 3.85-4.05 (2H, m), 4.80-4.90 (1H, m), 8.10 (1H, s), 8.20-8.25 (2H, m), 8.80 (1H, s)	3.45
430	359.00	1H NMR (CD3OD): 1.30-1.40 (6H, d), 2.88 (3H, s), 2.95-3.05 (1H, m), 4.20-4.25 (2H, m), 8.05 (1H, s), 8.20-8.25 (2H, m), 8.80 (1H, s)	3.00
431	378.73	1H NMR (CD3OD): 1.70-1.75 (3H, d), 2.90-3.15 (2H, m), 3.5-3.6 (2H, m); 3.80-3.85 (1H, m), 8.35-8.45 (3H, m), 8.60-8.70 (1H, m)	2.54
432	470.19	1H NMR (CD3OD): 1.30-1.40 (6H, d), 1.60 (3H, d), 2.95 (3H, s), 3.85-4.20 (2H, m), 4.80-4.90 (1H, qd), 7.60 (1H, s), 8.05 (1H, s), 8.20 (1H, s)	3.43
433	439.00	1H NMR (CD3OD): 1.45-1.55 (3H, d), 1.80-1.90 (2H, m), 2.65-2.80 (2H, m); 4.50-4.60 (1H, m), 7.00-7.10 (5H, m), 8.10-8.20 (3H, m), 8.80 (1H, s)	3.33
434	443.00	1H NMR (CD3OD): 1.65-1.75 (3H, d), 2.90-2.95 (2H, m), 3.50-3.60 (2H, m); 3.70-3.80 (3H, m), 4.90-5.00 (1H, m), 7.30 (1H, s), 8.40-8.45 (2H, m), 8.50 (1H, s), 8.60 (1H, s), 8.80 (1H, s)	2.77
435	448.00	1H NMR (CD3OD): 1.75-1.85 (3H, d), 3.05-3.15 (2H, m), 3.25-3.30 (2H, m); 3.45-3.55 (2H, m), 3.60-3.80 (4H, m), 3.90-4.00 (2H, m), 4.95-5.00 (1H, m), 8.40-8.45 (2H, m), 8.50 (1H, s), 8.60 (1H, s)	2.81

Cmpd No.	LC_MASS _PLUS M+1 (obs)	NMR_RESULT (1H NMR)	RT (mins)
436	432.00	1H NMR (CD3OD): 1.75-1.85 (3H, d), 2.90-2.10 (4H, m), 3.00-3.10 (2H, m); 3.20-3.30 (2H, m), 3.55-3.75 (4H, m), 4.95-5.00 (1H, m), 8.40-8.45 (2H, m), 8.50 (1H, s), 8.60 (2H, s)	2.64
437	478.19	1H NMR (CD3OD): 1.65-1.70 (3H, d), 2.90-3.00 (2H, m), 3.50-3.70 (2H, m), 4.80-4.90 (1H, m), 6.80-6.90 (1H, t), 6.95-7.05 (3H, m), 7.20-7.25 (1H, d), 7.35-7.40 (1H, d), 8.35-8.40 (3H, m), 8.50 (1H, m)	3.18
438	440.00	1H NMR (CD3OD): 1.65-1.70 (3H, d), 3.70-3.90 (4H, m), 4.80-4.90 (1H, m), 7.80-7.85 (1H, t), 7.90-7.95 (1H, t), 8.35-8.45 (3H, m), 8.50 (2H, s), 8.60 (1H, m)	2.93
439	446.00	1H NMR (CD3OD): 1.75-1.85 (3H, d), 1.90-2.10 (6H, m), 2.80-2.90 (2H, m); 3.00-3.10 (2H, m), 3.40-3.75 (4H, m), 4.95-5.00 (1H, m), 8.40-8.45 (2H, m), 8.50 (1H, s), 8.60 (2H, s)	2.59
440	435.14	1H NMR (CD3OD): 1.00-1.05 (6H, d), 1.75-1.85 (3H, d), 3.20-3.40 (5H, m), 4.60-4.70 (1H, m), 8.10-8.30 (3H, m), 8.80 (1H, s)	3.15
441	447.00	1H NMR (CD3OD): 1.75-1.85 (3H, d), 3.10-3.80 (8H, m), 4.95-5.00 (1H, m), 8.40-8.45 (2H, m), 8.50 (1H, s), 8.60 (1H, s)	2.66
442	443.00	1H NMR (CD3OD): 1.75-1.85 (3H, d), 2.05-2.15 (2H, m), 3.55-3.60 (1H, m); 3.75-3.80 (1H, m), 4.10-4.20 (2H, m), 4.95-5.00 (1H, m), 7.50 (2H, s), 8.35 (1H, s), 8.40-8.60 (3H, m), 8.63 (1H, s), 8.87 (1H, s)	2.75
443	428.00	1H NMR (CD3OD): 1.60-1.70 (3H, d), 3.40-3.50 (1H, m), 3.70-3.80 (1H, m); 4.00-4.05 (2H, m), 4.80-4.90 (1H, m), 5.90 (1H, s), 6.60 (1H, s), 8.35-8.55 (7H, m), 8.6 (2H, s),	3.16
444	420.09	1H NMR (DMSO-d6): 1.26 (3H, d), 1.68 (3H, s), 3.05 (4H, m), 4.54 (1H, m), 7.78 (2H, m), 8.13 (1H, s), 8.27 (3H, m), 8.66 (1H, s), 12.40 (1H, s)	2.68
445	470.12	1H NMR (CDCl3 + CD3OD drops): 1.19 (5H, m), 1.66 (3H, d), 2.76 (2H, m), 3.06 (2H, m), 4.69 (1H, m), 8.10 (1H, s), 8.18 (1H, s), 8.30 (1H, s), 8.64 (1H, s)	2.88
446	518.19	1H NMR (CDCl3 + CD3OD drops): 2.83 (2H, t), 3.33 (1H, s), 3.40 (1H, m), 3.58 (1H, m), 4.05 (2H, m), 4.59 (1H, m), 6.94 (1H, br s), 7.06 (2H, d), 7.56 (2H, d), 7.92 (1H, s), 8.03 (1H, s), 8.12 (1H, s), 8.12 (1H, s), 8.71 (1H, s)	2.88
447	423.00	1H NMR (CD3OD): 1.05-1.10 (3H, t), 1.60-1.70 (3H, d), 2.35-2.40 (2H, m), 2.45-2.55 (2H, m); 3.40-3.50 (2H, m), 4.55-4.60 (1H, m), 8.10 (1H, s), 8.15 (1H, s), 8.20 (1H, s), 8.80 (1H, s)	3.23

Cmpd No.	LC_MASS _PLUS M+1 (obs)	NMR_RESULT (1H NMR)	RT (mins)
448	446.00	1H NMR (CD3OD): 1.60-1.70 (4H, m), 1.90-2.05 (2H, m), 2.10-2.20 (1H, m), 2.25-2.40 (1H, m); 2.70-2.80 (3H, m), 2.90-3.00 (1H, m), 3.05-3.10 (1H, m), 3.35-3.45 (2H, m), 3.50-3.60 (2H, m), 3.65-3.75 (2H, m), 4.90-5.00 (1H, m), 8.35-8.40 (2H, m), 8.50 (1H, s), 8.60-8.65 (1H, s)	2.62
449	440.00	1H NMR (CD3OD): 1.65-1.70 (3H, d), 3.70-3.90 (4H, m), 4.80-4.90 (1H, m), 7.80-7.85 (1H, t), 8.35-8.45 (3H, m), 8.45-8.55 (2H, s), 8.60 (1H, s), 8.75 (1H, s)	2.90
450	452.00	(d6-DMSO, 400MHz) 1.36 (3H, d), 2.28 (3H, s), 2.78 (3H, s), 2.79 - 2.99 (2H, m), 3.15 - 3.25 (2H, m), 4.50 (1H, brs), 6.22 (1H, brs), 7.05 (1H, brs), 7.38 (1H, d), 8.13 (1H, brs), 8.22 (1H, s), 8.26 (1H, d), 8.77 (1H, brs), 12.30 (1H, s)	2.76
451	492.00	1H NMR (CD3OD): 1.40-1.45 (9H, s), 1.65-1.70 (3H, d), 2.80-2.85 (3H, m), 2.95-3.00 (1H, m), 3.05-3.10 (1H, m), 3.40-3.50 (2H, m), 4.60-4.70 (1H, m), 8.15 (1H, s), 8.25 (1H, s), 8.30 (1H, s), 8.80 (1H, s)	3.42
452	392.00	1H NMR (CD3OD): 1.65-1.70 (3H, d), 3.20-3.40 (2H, m), 3.70 (3H, s), 3.75-3.80 (2H, m), 4.50-4.60 (1H, m), 6.25-6.30 (1H, d), 6.35-6.40 (1H, t), 6.55-6.65 (2H, m), 7.90 (1H, s), 8.05 (1H, s), 8.15 (1H, s), 8.55 (1H, s)	2.59
453	485.00	1H NMR (CD3OD): 1.65-1.70 (3H, d), 3.20-3.40 (2H, m), 3.70 (3H, s), 3.75-3.80 (2H, m), 4.50-4.60 (1H, m), 6.25-6.30 (1H, d), 6.35-6.40 (1H, t), 6.55-6.65 (2H, m), 7.90 (1H, s), 8.05 (1H, s), 8.15 (1H, s), 8.55 (1H, s)	3.13
454	484.00	1H NMR (CD3OD): 1.08-1.13 (6H, m), 1.55-1.60 (3H, d), 2.85-3.05 (2H, m), 3.10-3.25 (2H, m), 3.35-3.40 (1H, m), 4.60-4.70 (1H, m), 8.10 (1H, s), 8.15 (1H, s), 8.20 (1H, s), 8.80 (1H, s)	2.94
455	468.00	(d6-DMSO, 400MHz) 1.39 (3H, brs), 2.70 - 2.96 (5H, m), 3.16 - 3.26 (2H, m), 3.84 (3H, s), 4.31 - 4.36 (1H, m), 6.60 (1H, s), 7.03 (1.4H, brs), 7.43 (0.6H, brs), 8.01 (1H, t), 8.26 (1H, s), 8.41 (1H, s), 8.70 (0.4H, brs), 9.09 (0.6H, brs), 12.39 (1H, s)	2.96
456	420.00	(d6-DMSO, 400MHz) 1.81 - 1.88 (1H, m), 1.94 - 1.98 (1H, m), 2.15 (6H, s), 2.24 - 2.28 (1H, m), 2.38 - 2.44 (1H, m), 2.86 (3H, s), 3.23 (3H, s), 5.38 - 5.40 (1H, m), 7.83 (1H, d), 8.15 (1H, s), 8.20 (1H, d), 8.28 (1H, d), 8.72 (1H, d), 12.35 (1H, s)	2.90
457	455.13	1H NMR (DMSO): 1.47 (3H, d), 3.36 (1H, m), 3.51 (1H, m), 3.80 (2H, m), 6.64 (2H, d), 6.82 (1H, t), 7.09 (2H, t), 8.07 (1H, br s), 8.28 (4H, m), 8.66 (1H, s), 12.50 (1H, s)	3.29
458	512.22	1H NMR (DMSO): 1.45 (3H, d), 3.07 (4H, m), 4.55 (1H, m), 4.93 (2H, m), 7.21 (1H, m), 7.29 (5H, m), 8.01 (1H, br s), 8.17 (1H, m), 8.28 (3H, m), 8.65 (1H, s), 12.44 (1H, s)	3.34

Cmpd No.	LC_MASS _PLUS M+1 (obs)	NMR_RESULT (1H NMR)	RT (mins)
459	479.16	1H NMR (DMSO + CD3OD drops): 1.41 (3H, d), 3.27 (2H, m), 3.58 (1H, m), 3.60 (1H, m), 4.53 (1H, m), 7.42 (2H, m), 7.60 (2H, m), 8.13 (1H, s), 8.24 (2H, m), 8.43 (1H, s), 8.51 (1H, s)	3.09
460	460.17	1H NMR (DMSO): 1.48 (5H, m), 1.75 (2H, m), 2.10 (2H, t), 2.92 (2H, m), 3.05 (4H, m), 4.53 (1H, m), 7.90 (1H, br s), 8.04 (1H, m), 8.28 (3H, m), 8.66 (1H, s), 12.41 (1H, s)	2.86
461	388.04	1H NMR (DMSO): 1.46 (3H, d), 2.59 (2H, m), 3.35 (1H, m), 4.57 (1H, m), 7.95 (1H, br s), 8.23 (1H, s), 8.28 (2H, s), 8.47 (1H, t), 8.65 (1H, s)	2.84
462	555.18	1H NMR (DMSO): 1.45 (3H, d), 2.60 (2H, m), 3.26 (2H, m), 3.85 (2H, d), 4.56 (1H, m), 7.22 (1H, t), 7.37 (2H, d), 7.69 (1H, d), 8.21 (4H, m), 8.66 (1H, s), 12.31 (1H, s)	3.59
463	419.00	1H NMR (CD3OD): 0.70-0.75 (3H, t), 1.05-1.15 (7H, m), 1.35- 1.45 (2H, m), 1.55-1.60 (3H, d), 3.90-4.00 (1H, m), 4.60-4.65 (1H, m), 8.10 (1H, s), 8.15 (1H, s), 8.20 (1H, s), 8.80 (1H, s)	3.50
464	405.00	1H NMR (CD3OD): 0.70-0.75 (6H, t), 1.30-1.40 (2H, m), 1.40-1.50 (1H, m), 1.55-1.60 (3H, d), 2.90-3.00 (1H, m), 3.15-3.20 (1H. m), 4.70-4.75 (1H, m), 8.10 (1H, s), 8.15 (1H, s), 8.20 (1H, s), 8.80 (1H, s)	3.40
465	391.00	1H NMR (CD3OD): 0.70-0.75 (3H, t), 1.05-1.10 (3H, d), 1.35- 1.45 (2H, m), 1.55-1.60 (3H, d), 3.80-3.90 (1H, m), 4.70-4.75 (1H, m), 8.10 (1H, s), 8.15 (1H, s), 8.20 (1H, s), 8.80 (1H, s)	3.30
466	407.00	1H NMR (CD3OD): 1.55-1.60 (3H, d), 2.40-2.50 (2H, m), 3.35-3.55 (2H. m), 4.65-4.70 (1H, m), 8.25 (1H, s), 8.30-8.40 (2H, m), 8.70 (1H, s)	2.50
467	420.00	1H NMR (CD3OD): 1.55-1.60 (3H, d), 2.30-2.40 (2H, m), 3.35-3.60 (5H, m), 4.60-4.70 (1H, m), 8.15 (1H, s), 8.20 (1H, s), 8.25 (1H, s), 8.80 (1H, s)	2.80
468	448.00	1H NMR (CD3OD): 1.00-1.10 (6H, m), 1.55-1.60 (3H, d), 2.30-2.40 (2H, m), 3.35-3.50 (3H, m), 3.80-3.90 (2H, m), 4.60-4.70 (1H, m), 8.15 (1H, s), 8.20 (1H, s), 8.25 (1H, s), 8.80 (1H, s)	3.00
469	484.00	1H NMR (CD3OD): 0.90-1.00 (6H, m), 1.55-1.60 (3H, d), 2.95-3.05 (1H, m), 3.10-3.25 (2H. m). 3.50-3.60 (1H. m), 3.70-3.80 (1H, m), 4.65-4.70 (1H, m), 8.10 (1H, s), 8.30-8.40 (2H, m), 8.70 (1H, s)	3.00

[00276] B) Biological Data:

[00277] Example 1: JAK3 Inhibition Assay

[00278] Compounds were screened for their ability to inhibit JAK3 using a standard radioactive enzyme assay. 1.5μL/well of a DMSO stock containing serial dilutions of a compound of the present invention (concentrations ranging from 667 μM to 46 nM) was placed in a 96 well polycarbonate plate. 50 μL per well of kinase buffer (100 mM HEPES (pH 7.4), 10 mM $MgCl_2$, 25 mM NaCl, 1 mM DTT, and 0.01% bovine serum albumin (BSA)) containing 2 μM poly(Glu)$_4$Tyr and 10 μM ATP was also added to the plate. To initiate the reaction, 50 μL kinase buffer containing 2 nM JAK3 enzyme was added. Final ATP concentration was 5 μM [γ-^{33}P] ATP (200 μCi ^{33}P ATP/μmol ATP (Perkin Elmer, Cambridge, MA). After 20 minutes at room temperature (25°C), the reaction was stopped by adding 50 μL 20% trichloroacetic acid (TCA)/0.4 mM ATP to each well. The entire content of each well was then transferred to a 96 well glass fiber filter plate using a TomTek Cell Harvester. After washing, 60 μL scintillation fluid was added and ^{33}P incorporation detected using a Perkin Elmer TopCount. After removing mean background values for all of the data points the data was fit using Prism software to obtain a K_i(app). Inhibition of JAK2 was measured as above except that final poly(Glu)$_4$Tyr concentration was 15 μM and final ATP concentration was 12 μM.

[00279] Example 2: ROCK Inhibition Assays

[00280] Compounds were screened for their ability to inhibit ROCK I (AA 6-553) activity using a standard coupled enzyme system (Fox et al. (1998) *Protein Sci.* 7, 2249). Reactions were carried out in a solution containing 100 mM HEPES (pH 7.5), 10 mM $MgCl_2$, 25 mM NaCl, 2 mM DTT and 1.5% DMSO. Final substrate concentrations in the assay were 45 μM ATP (Sigma Chemicals, St Louis, MO) and 200 μM peptide (American Peptide, Sunnyvale, CA). Reactions were carried out at 30 °C and 45 nM ROCK I. Final concentrations of the components of the coupled enzyme system were 2.5 mM phosphoenolpyruvate, 350 μM NADH, 30 μg/ml pyruvate kinase and 10 μg/ml lactate dehydrogenase.

[00281] Some compounds were screened for their ability to inhibit ROCK using a standard radioactive enzyme assay. Assays were carried out in a solution containing

100 mM HEPES (pH 7.5), 10 mM $MgCl_2$, 25 mM NaCl, 2 mM DTT, and 1.5% DMSO. Final substrate concentrations in the assay were 13 μM [γ-^{33}P] ATP (25mCi ^{33}P ATP/mmol ATP, Perkin Elmer, Cambridge, MA /Sigma Chemicals, St Louis, MO) and 27 μM Myelin Basic Protein (MBP). Final enzyme concentration in the assay was 5 nM ROCK. Assays were carried out at room temperature. 1.5μl of DMSO stock containing serial dilutions of the compound of the present invention (concentrations ranging from 10 μM to 2.6nM) was placed in a 96 well plate. 50 μl of Solution 1 (100 mM HEPES (pH 7.5), 10 mM $MgCl_2$, 26 mM [γ-^{33}P] ATP) was added to the plate. The reaction was initiated by addition of 50 μl of Solution 2 (100 mM HEPES (pH 7.5), 10 mM $MgCl_2$, 4 mM DTT, 54 mM MBP and 10 nM ROCK). After 2 hours the reaction was quenched with 50 μL of 30% trichloroacetic acid (TCA, Fisher) containing 9mM ATP. Transfer of 140 μL of the quenched reaction to a glass fiber filter plate (Corning, Cat. No.3511) was followed by washing 3 times with 5% TCA. 50 μL of Optima Gold scintillation fluid (Perkin Elmer) was added and the plates were counted on a Top Count (Perkin Elmer). After removing mean background values for all of the data points the data was fit using Prism software to obtain a K_i(app).

[00282] Example 3: Aurora Inhibition Assay

[00283] Compounds were screened for their ability to inhibit full length Aurora-A (AA 1-403) activity using a standard coupled enzyme system (Fox *et al.*, Protein Sci., 7, pp. 2249 (1998)). Reactions were carried out in a solution containing 100 mM HEPES (pH 7.5), 10 mM $MgCl_2$, 25 mM NaCl, 300 μM NADH, 1 mM DTT and 3% DMSO. Final substrate concentrations in the assay were 200 μM ATP (Sigma Chemicals, St Louis, MO) and 800 μM peptide (LRRASLG, American Peptide, Sunnyvale, CA). Reactions were carried out at 30 °C and 35 nM Aurora-A. Final concentrations of the components of the coupled enzyme system were 2.5 mM phosphoenolpyruvate, 200 μM NADH, 60 μg/ml pyruvate kinase and 20 μg/ml lactate dehydrogenase.

[00284] An assay stock buffer solution was prepared containing all of the reagents listed above with the exception of ATP and the test compound of interest. The assay stock buffer solution (60 μl) was incubated in a 96 well plate with 2 μl of the test

compound of interest at final concentrations spanning 0.002 μM to 30 μM at 30 °C for 10 min. Typically, a 12 point titration was conducted by preparing serial dilutions (from 1 mM compound stocks) with DMSO of the test compounds in daughter plates. The reaction was initiated by the addition of 5 μl of ATP (final concentration 200 μM). Rates of reaction were obtained using a Molecular Devices Spectramax plate reader (Sunnyvale, CA) over 10 min at 30 °C. The Ki values were determined from the rate data as a function of inhibitor concentration using computerized nonlinear regression (Prism 3.0, Graphpad Software, San Diego, CA). Aurora-C activity was screened in a similar manner using Aurora-C protein. Aurora-B activity was screened using a radioactive assay, such as described in Examples 1 and 2, but using Aurora-B protein.

[00285] Table 5 below depicts enzyme inhibition data (K_i) for certain exemplary compounds. Compound numbers correspond to those compounds depicted in Table 1.

[00286] Table 6 below depicts enzyme inhibition data (K_i) for certain exemplary compounds. Compound numbers correspond to those compounds depicted in Table 2.

[00287] In Tables 5 and 6, "A" represents a K_i of less than 0.5 μM, "B" represents a K_i of between 0.5 and 5.0μM, and "C" represents a K_i greater than 5.0 μM for the indicated enzyme. If more than one value of K_i has been determined, the average K_i is indicated. If no value is indicated, then the K_i was not determined. For ROCK, the term "Enzyme" indicates that an enzyme-linked assay was used; the term "^{33}P" indicates that a radioactive assay was used.

Table 5. Enzyme Inhibition Data for Compounds of Table 1

Compound Number	Aurora A	JAK2	JAK3	ROCK	
				Enzyme	^{33}P
1				A	A
2	B				A
3	A	A	A		B
4		B	A		A
5		B	C		A
6		B	B		A
7	B	A	A		A

Compound Number	Aurora A	JAK2	JAK3	ROCK	
				Enzyme	^{33}P
8	B	A	A		A
9	B	B	B		B
10		A	A		A
11		A	A		A
12		A	B		B
13		B	B		A
14		A	A		A
15					A
16	A	A	A		A
17	B	A	B		B
18	B	A	A		A
19		A	A		A
20		A	A		A
21	A				A
22					A
23					A
24					A
25	B				A
26		B	B		B
27					A
28					A
29					A
30	A				A
31	A				A
32		B	A		A
33	B	A	A		A
34		A	A		A
35		B	C		B
36		B	B		B
37		A	A		A
38	B	A	A		A
39		C	C		B
40		A	A		A
41		A	A		B
42		A	A		B
43	B	A	A		A
44	B	B	B		A
45		B	B		B
46		B	A		A
47		B	B		B
48		B	B		B
49		B	A		B
50		B	B		B
51	B	B	A		A

Compound Number	Aurora A	JAK2	JAK3	ROCK	
				Enzyme	^{33}P
52		C	C		B
53		C	B		B
54		C	C		B
55		B	B		A

Table 6. Enzyme Inhibition Data for Compounds of Table 2

Compound No.	Aurora A	Aurora B	Aurora C	JAK2	JAK3	ROCK	
						Enzyme	^{33}P
1				A	A		B
2				A	A		B
3				A	A		A
4	A			A	A		A
5				A	A		B
6				A	A		A
7				B	B		B
8	A			A	A		A
9				A	A		B
10	A			A	A		A
11				A	A		B
12				A	A		A
13				A	A		B
14	B			A	A		A
15				B	B		B
16				A	A		A
17				B	B		A
18	B			A	A	B	B
19	A			A	A	A	A
20	A			A	A		B
21				A	A		A
22				A	A		A
23				B	B		A
24	B			A	A		A
25	A			A	A		A
26				A	A		A
27				A	A		A
28				A	A		A
29				A	A		A
30	A			A	A		A
31				A	A		A
32	A			A	A		B
33	A			A	A	A	A
34				A	A	A	B
35	A			A	A	A	A

Compound No.	Aurora A	Aurora B	Aurora C	JAK2	JAK3	ROCK	
						Enzyme	^{33}P
36	A			A	A	B	B
37				A	A	B	A
38	B			A	A	B	B
39	A			A	A	B	
40				A	A	B	
41	A			A	A	B	
42	A			A	A	B	
43	A			B	B	B	
44	A			B	B	B	
45	A			A	A	B	
46	A			A	A	B	
47	B			A	A	B	
48	A			A	A	B	
49	B			A	A	B	
50	B			A	A	A	
51				B	B	B	
52	B			C	C	B	
53	A			A	A	A	
54	A			B	B	B	
55				B	B		
56	B			C	B		
57	B			C	B	B	
58	B			A	A	B	
59	B			A	A	B	
60	B			B	B	B	
61				A	A	B	
62				A	A	A	
63				A	A	A	
64				A	A	B	
65				A	A	B	
66	B			A	A	B	
67	A			A	A	B	
68	A			A	A	B	
69	B			A	A	B	
70	B			A	A	B	
71				A	A	B	
72	A			B	A	B	
73				A	A	A	
74				A	A	A	
75	A			A	A	A	
76	A			A	A	B	
77				A	A	B	
78	A			A	A	B	
79	A			A	A	B	

Compound No.	Aurora A	Aurora B	Aurora C	JAK2	JAK3	ROCK	
						Enzyme	^{33}P
80	A			A	A	B	
81	A			B	B	B	
82				A	A	A	
83				A	A	B	
84				A	A	B	
85				A	A	B	
86				A	A	B	
87				A	A	B	
88				A	A	B	
89				A	A	B	
90				A	A	B	
91				B	B	B	
92	A			A	A	A	
93	A			A	A	B	
94				B	B	B	
95	B			B	B	B	
96	B			A	A	B	
97	A			A	A	B	
98	A			A	A	B	
99	A			A	A	B	
100	A			A	A	B	
101	A			A	A	B	
102	B			A	A	B	
103	B			A	A	B	
104				A	A	B	
105				A	A	B	
106				A	A	B	
107				A	A	B	
108				A	A	B	
109				A	A	B	
110	B			B	B	B	
111				A	A	B	
112				A	A	B	
113	B			A	A	B	
114	B			A	A	B	
115				A	A	B	
116				A	A	B	
117	B			A	A	B	
118	B			A	A	B	
119	B			A	A	B	
120	A			A	A	B	
121	B			A	A	B	
122	B			A	A	B	
123	B			A	A	B	

Compound No.	Aurora A	Aurora B	Aurora C	JAK2	JAK3	ROCK	
						Enzyme	^{33}P
124	B			A	A	B	
125	B			A	A	B	
126				C	B	B	
127				A	A	B	
128	B			A	A	B	
129				B	A	B	
130	A			A	A	B	
131	A			A	A	B	
132	A			A	A	B	
133				A	A	B	
134				A	A	B	
135				A	A	A	
136				A	A	A	
137				A	A	B	
138				A	A	B	
139				A	A	B	
140				A	A	B	
141				A	A	B	
142				B	A	B	
143				B	A	B	
144				A	A	B	
145				A	A	B	
146				A	A	B	
147				A	A	B	
148				B	A	A	
149				A	A	B	
150	B			A	A	B	
151	A			A	A	A	
152	B			A	A	B	
153	B			A	A	B	
154	B			A	A	B	
155	B			A	A	B	
156	B			A	A	A	
157	B			A	A	B	
158	B			A	A	B	
159	B			A	A	B	
160				A	A	B	
161	B			A	A	B	
162	B			A	A	B	
163	B			A	A	B	
164	B			A	A	B	
165	B			A	A	B	
166	B			A	A	B	
167	B			A	A	B	

Compound No.	Aurora A	Aurora B	Aurora C	JAK2	JAK3	ROCK	
						Enzyme	^{33}P
168	B			A	A	B	
169	B			A	A	B	
170	B			A	A	B	
171				A	A	B	
172	B			A	A	A	
173				A	A	A	
174				A	A		
175				A	A		
176				A	A		
177	A			A	A	B	
178				B	B	B	
179	B			A	A	B	
180				A	A	B	
181	B			A	A	B	
182	B			A	A	B	
183				A	A	B	
184				B	A	B	
185				B	A		
186	A			A	A		
187				B	A		
188				C	B		
189				A	A	B	
190	B			A	A	B	
191	B			A	A	A	
192	B			A	A	A	
193				A	A	A	
194				A	A	B	
195				B	A	B	
196				C	B	B	
197				B	A	B	
198				C	B	B	
199				B	A	B	
200				B	A	B	
201				A	A	B	
202				B	B	B	
203				A	A	B	
204					A	B	
205					A	B	
206					A	B	
207					A	B	
208					A	B	
209					A	B	
210	B						
211	A						B

Compound No.	Aurora A	Aurora B	Aurora C	JAK2	JAK3	ROCK	
						Enzyme	^{33}P
212	A	B					B
213	A						B
214	B						B
215	B						B
216	A						B
217	A					B	B
218	A					A	A
219	A					B	B
220	B					B	
221	A					A	
222	A					A	
223	B					B	
224	A					A	
225	B					B	
226	B						
227	A					A	
228	A					A	
229	A					B	
230	B					B	
231	A					A	
232	A					B	
233						B	
234	A						
235	A					A	
236	A						
237	A	A	A			B	
238	A	A	A			B	
239	A					A	
240	A					B	
241	B						
242	A	B	A			B	
243	A					B	
244	B					B	
245	A						
246	B					B	
247	A					B	
248	A					B	
249	A					B	
250	A					B	
251	A					B	
252	A		A			B	
253	A					A	
254	A					B	
255	A					A	

Compound No.	Aurora A	Aurora B	Aurora C	JAK2	JAK3	ROCK	
						Enzyme	^{33}P
256	A					A	
257	B					B	
258	A					B	
259	A					A	
260	A		A			B	
261	B					B	
262	B					B	
263	A					B	
264	A					B	
265	A					B	
266	A					B	
267	A					A	
268	B						
269	A						
270	B					B	
271	A					B	
272	B						
273	A					B	
274	A					B	
275	B					B	
276	A					B	
277	A					A	
278	A					A	
279	B						
280	A						
281	A						
282	A						
283	B						
284	B					B	
285	A	A	A			B	
286	A					B	
287	A						
288	A						
289	A					B	
290	A	B	B			B	
291	A	B				B	
292	C					B	
293	B					B	
294	A					B	
295	A					B	
296	A					B	
297	A					B	
298	B					B	
299	A	A				A	

Compound No.	Aurora A	Aurora B	Aurora C	JAK2	JAK3	ROCK	
						Enzyme	^{33}P
300	A	A				A	
301	B					B	
302	A	A				B	
303						A	
304						B	
305	A	A				B	
306	A	A	A			B	
307	A					A	
308	B					B	
309	B					B	
310	A	A				B	
311	A					B	
312	A	A	A			B	
313	A					B	
314	B						
315	A	B				B	
316	A	A	A			B	
317	A	A	A			B	
318	A					B	
319	B						
320	A						
321	A	B	A			B	
322	A	A				B	
323	A	A	A			B	
324	A	A	A			B	
325	A	A	A			B	
326	A	A	A			A	
327	A	A	A			A	
328	A	A	A			B	
329	A	A	A			A	
330	A	A				B	
331	A		A			B	
332	A						
333	A	A	A			B	
334	A						
335	A	A					
336	A					B	
337	A					B	
338	A					B	
339	A	A	A			B	
340	A	A	A			B	
341	A	A				B	
342	A					B	
343	B						


Compound No.	Aurora A	Aurora B	Aurora C	JAK2	JAK3	ROCK	
						Enzyme	^{33}P
344	A					A	
345	A	B	B			A	
346	B						
347	A	B	B			A	
348	A	A	A			B	
349	A	A	A			B	
350	B						
351	A					B	
352	A						
353						B	
354	A	A				B	
355	A	A	A			B	
356	A	A	A			B	
357	A					B	
358	B						
359	A	A	A			B	
360	B					B	
361	A	A	A			B	
362	A	A	A			B	
363	A					B	
364	A					B	
365	A					B	
366	A					B	
367	A					B	
368	A	A	A			B	
369	A	A	A		A	B	
370	A	A	A			B	
371	A	A	A	A		B	
372	A					B	
373	A					B	
374	A	A	A			B	
375	A	A				B	
376	A					B	
377	A					B	
378	A					B	
379	A					A	
380	A	A				B	
381	A					A	
382	A	B					
383	A					B	
384	A					B	
385	A	A				B	
386	A	A	A	A		B	
387	A	A	A			B	

Compound No.	Aurora A	Aurora B	Aurora C	JAK2	JAK3	ROCK	
						Enzyme	^{33}P
388	A	A				B	
389	A					B	
390	B					B	
391	A		A		A	B	
392	A					B	
393	A					B	
394	A					B	
395	A	A		A	A	B	
396	A					B	
397	A	A				B	
398	A					B	
399	A					A	
400	A		A			B	
401	A					B	
402	B					B	
403	B						
404	A	B				B	
405	A						
406	A	B				B	
407	A		A			B	
408	A					B	
409	A					A	
410	A					B	
411	A		A			B	
412	A						
413	A	A				B	
414	B	B				B	
415	A	A				B	
416	A	B				B	
417	A	A				B	
418	A	A				B	
419	B	B					
420	A	A	A			B	
421	A	B				B	
422	A	A				B	
423	A	A				B	
424	A	A				B	
425	A	A				B	
426	A	A				B	
427	A	B	B			B	
428	A	B				B	
429	A	B	B			B	
430	A	B	B			B	
431	A	A	A			B	

Compound No.	Aurora A	Aurora B	Aurora C	JAK2	JAK3	ROCK	
						Enzyme	^{33}P
432	B	B				B	
433	B	B				B	
434	B	B				B	
435	B	B				B	
436	B	A				B	
437	A	B				B	
438	B	B				B	
439	A	A				B	
440	B	B				B	
441	B	B		B		B	
442	A	B		B		B	
443	A	A	A	A		B	
444	A	A	A	B		B	
445	A	B	A				
446	A	A	A	B		B	
447	A	B	B			B	
448	B	B				B	
449	A	B				B	
450	A	A	A	B		B	
451	B	B				B	
452	A	A	A			B	
453	B	B				B	
454	A	B	B			B	
455	A	B	B			B	
456	A	B				B	
457	B	B					
458	A	A	A				
459	B	B					
460	B	B					
461	A	A	A				
462	B	B					
463	A	B					
464	B						
465	A						
466	B						
467	A						
468	A						
469	A						

CLAIMS

1. A compound of formula **(IB):**



IB

or pharmaceutically acceptable salts thereof, wherein:

R^1 is hydrogen, T-R' or $-Si(R')_3$;

R^2, R^3, and R^4 are each independently hydrogen, halogen, CN, NO_2, or V-R';

x is 1, 2, 3, or 4;

each occurrence of R^5 is independently halogen, CN, NO_2, or U-R', wherein at least one R^5 is other than H;

T, V, and U are each independently a bond or an optionally substituted C_1-C_6 alkylidene chain, wherein up to two methylene units of the chain are optionally and independently replaced by –NR'-, -S-, -O-, -CS-, -CO_2-, -OCO-, -CO-, -COCO-, -CONR'-, -NR'CO-, -NR'CO_2-, -SO_2NR'-, -NR'SO_2-, -CONR'NR'-, -NR'CONR'-, -OCONR'-, -NR'NR'-, -NR'SO_2NR'-, -SO-, or -SO_2-, -PO-, -PO_2-, or –POR'-; and

each occurrence of R' is independently, hydrogen or an optionally substituted group selected from a C_1-C_6 aliphatic group, a 3-8-membered saturated, partially unsaturated, or fully unsaturated monocyclic ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur, or an 8-12 membered saturated, partially unsaturated, or fully unsaturated bicyclic ring system having 0-5 heteroatoms independently selected from nitrogen, oxygen, or sulfur; or two occurrences of R', are taken together with the atom(s) to which they are bound to form an optionally substituted 3-12 membered saturated, partially unsaturated, or fully

unsaturated monocyclic or bicyclic ring having 0-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur;

provided that:

a) if R^2, R^3 and R^4 are simultaneously H, then R^1 is not $Si(R')_3$; and

b) if R^1, R^2 and R^4 are simultaneously H and any one of X^1, X^2 or X^3 are N, then R^3 is not phenyl or phenyl substituted with O-phenyl or $N(Me)_2$.

2. The compound of claim 1, wherein:

a) R^1 is T-R', wherein T is a bond or is an optionally substituted C_1-C_6alkylidene chain wherein up to two methylene units are optionally and independently replaced with –O-, -S-, -NR'-, -OCO-, -COO-, $-SO_2$-or –CO-, and R' is hydrogen, C_1-C_4-alkyl, or an optionally substituted 5- or 6-membered aryl or heteroaryl group; or

R^1 is $-Si(R')_3$, R' is hydrogen, C_1-C_4-alkyl, or an optionally substituted 5- or 6-membered saturated, partially unsaturated, or fully unsaturated monocyclic ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur.

3. The compound of claim 1, wherein R^1 is hydrogen, C_1-C_4alkyl, -COR', $-SO_2R'$, or $-Si(R')_3$.

4. The compound of claim 1, wherein R^1 is hydrogen, methyl, ethyl, n-propyl, isopropyl, n-butyl, p-toluenesulfonyl (Ts), t-butyldimethylsilyl (TBS), triisopropylsilyl (TIPS), or triethylsilyl (TES).

5. The compound of claim 1, wherein R^2, R^3, and R^4 are each independently hydrogen, R', halogen, CN, NO_2, $-N(R')_2$, $-CH_2N(R')_2$, -OR', $-CH_2OR'$, -SR', $-CH_2SR'$, -COOR', -NR'COR', -NR'COCH2R', or -NR'CO(CH2)2R', $-CON(R')_2$, $-SO_2N(R')_2$, $-CONR'(CH_2)_2N(R')_2$, $-CONR'(CH_2)_3N(R')_2$, $-CONR'(CH_2)_4N(R')_2$, $-O(CH_2)_2OR'$, $O(CH_2)_3OR'$, $O(CH_2)_4OR'$, $-O(CH_2)_2N(R')_2$, $-O(CH_2)_3N(R')_2$, or $-O(CH_2)_4N(R')_2$.

6. The compound of claim 1, wherein R^2, R^3, and R^4 are each independently Cl, Br, F, -CN, -COOH, -COOMe, $-NH_2$, $-N(CH_3)_2$, $-N(Et)_2$, $-N(iPr)_2$, $-O(CH_2)_2OCH_3$, $-CONH_2$, $-COOCH_3$, -OH, $-CH_2OH$, $-NHCOCH_3$, $-SO_2NH_2$, $-SO_2N(Me)_2$, or an optionally substituted group selected from C_1-C_4alkyl, C_1-C_4alkyloxy, a 3-8-membered saturated, partially unsaturated, or fully unsaturated monocyclic ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur, or an 8-12 membered saturated, partially unsaturated, or fully unsaturated bicyclic ring system having 0-5 heteroatoms independently selected from nitrogen, oxygen, or sulfur.

7. The compound of claim 1 wherein:

a) R^2, R^3, and R^4 are each is hydrogen;

b) one of R^2, R^3, or R^4 is hydrogen;

c) two of R^2, R^3, or R^4 is hydrogen;

d) R^2 and R^4 are both hydrogen, and R^3 is halogen, CN, NO_2, or V-R';

e) R^2 and R^4 are both hydrogen, and R^3 is an optionally substituted group selected from a 3-8-membered saturated, partially unsaturated, or fully unsaturated monocyclic ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur, or an 8-12 membered saturated, partially unsaturated, or fully unsaturated bicyclic ring system having 0-5 heteroatoms independently selected from nitrogen, oxygen, or sulfur;

f) R^2 and R^4 are both hydrogen, and R^3 is an optionally substituted 5- or 6-membered saturated, partially unsaturated, or fully unsaturated ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur; or

g) R^2 and R^4 are both hydrogen, and R^3 is an optionally substituted ring selected from phenyl, pyridyl, pyrimidinyl, thiazolyl, oxazolyl, thienyl, furyl, pyrrolyl, pyrazolyl, triazolyl, pyrazinyl, thiadiazolyl, or oxadiazolyl.

8. The compound of claim 1, wherein R^2, R^3, and R^4 are each optionally substituted with z occurrences of R^6, wherein z is 0-5 and R^6 is =O, =NR", =S, halogen, -CN, $-NO_2$, or Z-R", wherein Z is a bond or an optionally substituted

C_1-C_6 alkylidene chain, wherein up to two methylene units of the chain are optionally and independently replaced by –NR"-, -S-, -O-, -CS-, -CO_2-, -OCO-, -CO-, -COCO-, -CONR"-, -NR"CO-, -NR"CO_2-, -SO_2NR"-, -NR"SO_2-, -CONR"NR"-, -NR"CONR"-, -OCONR"-, -NR"NR"-, -NR"SO_2NR"-, -SO-, -SO_2-, -PO-, -PO_2-, or –POR"-, and and each occurrence of R" is independently hydrogen or an optionally substituted C_1.C_6 aliphatic group, a 3-8-membered saturated, partially unsaturated, or fully unsaturated monocyclic ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur, or an 8-12 membered saturated, partially unsaturated, or fully unsaturated bicyclic ring system having 0-5 heteroatoms independently selected from nitrogen, oxygen, or sulfur; or two occurrences of R" are taken together with the atom(s) to which they are bound to form an optionally substituted 3-12 membered saturated, partially unsaturated, or fully unsaturated monocyclic or bicyclic ring having 0-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur.

9. The compound of claim 8, wherein z is 0, 1, 2, or 3, and each occurrence of R^6 is independently hydrogen, R", -CH_2R", halogen, CN, NO_2, -N(R")$_2$, -CH_2N(R")$_2$, -OR", -CH_2OR", -SR", -CH_2SR", -COOR", -NR"COR", -NR"COOR", -CON(R")$_2$, -SO_2N(R")$_2$, -CONR"$(CH_2)_2$N(R")$_2$, -CONR$(CH_2)_3$N(R")$_2$, -CONR"$(CH_2)_4$N(R")$_2$, -O$(CH_2)_2$OR", O$(CH_2)_3$OR", O$(CH_2)_4$OR", -O$(CH_2)_2$N(R")$_2$, -O$(CH_2)_3$N(R")$_2$, -O$(CH_2)_4$N(R")$_2$, -NR"CH(CH_2OH)R", -NR"CH(CH_2CH_2OH)R", -NR"(CH_2)R", -NR"$(CH_2)_2$R", -NR"$(CH_2)_3$R", -NR"$(CH_2)_4$R", -NR"(CH_2)N(R")$_2$, -NR"$(CH_2)_2$N(R")$_2$, -NR"$(CH_2)_3$N(R")$_2$, -NR"$(CH_2)_4$N(R")$_2$, -NR"(CH_2)OR", -NR"$(CH_2)_2$OR", -NR"$(CH_2)_3$OR", or –NR"$(CH_2)_4$OR".

10. The compound of claim 8, wherein z is 1, 2, or 3 and each occurrence of R^6 is independently F, Cl, Br, CN, OH, NH_2, -CH_2OH, C_1-C_6alkyl, -O(C_1-C_6alkyl), -CH_2O(C_1-C_6alkyl), -CO(C_1-C_6alkyl), -COO(C_1-C_6alkyl), -NHSO_2(C_1-C_6alkyl), -SO_2NH_2, -$CONH_2$, -CON(C_1-C_6alkyl), -SO_2(C_1-C_6alkyl), -SO_2phenyl, phenyl, benzyl, -N(C_1-C_6alkyl)$_2$, or -S(C_1-C_6alkyl), wherein each of the foregoing

phenyl, benzyl, and C_1-C_6alkyl groups is independently and optionally substituted, and wherein each of the foregoing C_1-C_6alkyl groups is linear, branched, or cyclic.

11. The compound of claim 1, wherein each occurrence of R^5 is independently hydrogen, R', -CH_2R', halogen, CN, NO_2, -N(R')$_2$, -CH_2N(R')$_2$, -OR', -CH_2OR', -SR', -CH_2SR', -COOR', -NR'COR', -NR'$COCH_2$R', -NR'$CO(CH_2)_2$R', -NR'COOR', -CON(R')$_2$, -SO_2N(R')$_2$, -CONR'$(CH_2)_2$N(R')$_2$, -CONR$(CH_2)_3$N(R')$_2$, -CONR'$(CH_2)_4$N(R')$_2$, -$O(CH_2)_2$OR', $O(CH_2)_3$OR', $O(CH_2)_4$OR', -$O(CH_2)_2$N(R')$_2$, -$O(CH_2)_3$N(R')$_2$, -$O(CH_2)_4$N(R')$_2$, -NR'CH(CH_2OH)R', -NR'CH(CH_2CH_2OH)R', -NR'(CH_2)R', -NR'$(CH_2)_2$R', -NR'$(CH_2)_3$R', -NR'$(CH_2)_4$R', -NR'(CH_2)N(R')$_2$, -NR'$(CH_2)_2$N(R')$_2$, -NR'$(CH_2)_3$N(R')$_2$, -NR'$(CH_2)_4$N(R')$_2$, -NR'(CH_2)OR', -NR'$(CH_2)_2$OR', -NR'$(CH_2)_3$OR', or -NR'$(CH_2)_4$OR'.

12. The compound of claim 1, wherein x is 1, 2, or 3, and at least one occurrence of R^5 is –N(R')$_2$, -NR'CH(CH_2OH)R', -NR'CH(CH_2CH_2OH)R', -NR'(CH_2)R', -NR'$(CH_2)_2$R', NR'(CH_2)N(R')$_2$, or -NR'$(CH_2)_2$N(R')$_2$.

13. The compound of claim 1, wherein x is 1, 2, or 3, and at least one occurrence of R^5 is -OR'.

14. The compound of claim 1, wherein x is 1, 2, or 3, and at least one occurrence of R^5 is -NR'COR', -NR'$COCH_2$R', or -NR'$CO(CH_2)_2$R'.

15. The compound of claim 1, wherein x is 1, 2, or 3, and at least one occurrence of R^5 is an optionally substituted C_1.C_6 aliphatic group, a 3-8-membered saturated, partially unsaturated, or fully unsaturated monocyclic ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur, or an 8-12 membered saturated, partially unsaturated, or fully unsaturated

bicyclic ring system having 0-5 heteroatoms independently selected from nitrogen, oxygen, or sulfur.

16. The compound of claim 1, wherein x is 1 or 2, and each occurrence of R^5 is independently halogen, R', CN, $-CH_2CN$, $-(CH_2)_2CN$, NO_2, $-CH_2NO_2$, $-(CH_2)_2NO_2$, $CON(R')_2$, $-CH_2CON(R')_2$, $-(CH_2)_2CON(R')_2$, COOR', $-CH_2COOR'$, $-(CH_2)_2COOR'$, $-SO_2N(R')_2$, $-CH_2SO_2N(R')_2$, $-(CH_2)_2SO_2N(R')_2$, $-NR'SO_2R'$, $-CH_2NR'SO_2R'$, $-(CH_2)_2NR'SO_2R'$, $NR'CON(R')_2$, $-CH_2NR'CON(R')_2$, $-(CH_2)_2NR'CON(R')_2$, $-NR'SO_2N(R')_2$, $-CH_2NR'SO_2N(R')_2$, $-(CH_2)_2NR'SO_2N(R')_2$, -COCOR', $-CH_2COCOR'$, $-(CH_2)_2COCOR'$,–$N(R')_2$, $-CH_2N(R')_2$, $-(CH_2)_2N(R')_2$, -OR', $-CH_2OR'$, $-(CH_2)_2OR'$, $-NR'COR'$, $-NR'COCH_2R'$, $-NR'CO(CH_2)_2R'$, $-CH_2NR'COR'$, or $-(CH_2)_2NR'COR'$.

17. The compound of claim 16, wherein R^5 is CN, $-CH_2CN$, $-(CH_2)_2CN$, $-NO_2$, $-CH_2NO_2$, $-(CH_2)_2NO_2$, OR', $-CH_2OR'$, $-CON(R')_2$, $-SO_2N(R')_2$, $-N(R')_2$, or R'.

18. The compound of claim 16, wherein each occurrence of R^5 is independently hydrogen, halogen, CN, $-CH_2CN$, $-(CH_2)_2CN$, NO_2, $-CH_2NO_2$, $-(CH_2)_2NO_2$, $-CONH_2$, $-CON(C_1\text{-}C_4alkyl)$, $-SO_2NH_2$, $-SO_2N(C_1\text{-}C_4alkyl)$, NH_2, $-N(C_1\text{-}C_4alkyl)$, -OH, $-O(C_1\text{-}C_4alkyl)$, $-CH_2OH$, $-CH_2O(C_1\text{-}C_4alkyl)$, or an optionally substituted 5- or 6-membered unsaturated ring wherein 0-3 ring carbon atoms is optionally replaced by oxygen, sulfur, or nitrogen.

19. The compound of claim 1, wherein R^5 is optionally substituted with y occurrences of R^7, wherein y is 0-5 and R^7 is =O, =NR", =S, halogen, -CN, $-NO_2$, or W-R", wherein W is a bond or an optionally substituted $C_1\text{-}C_6$ alkylidene chain, wherein up to two methylene units of the chain are optionally and independently replaced by –NR"-, -S-, -O-, -CS-, $-CO_2$-, -OCO-, -CO-, -COCO-, -CONR"-, -NR"CO-, $-NR"CO_2$-, $-SO_2NR"$-, -

NR"SO_2-, -CONR"NR"-, -NR"CONR"-, -OCONR"-, -NR"NR"-, -NR"SO_2NR"-, -SO-, -SO_2-, -PO-, -PO_2-, or –POR"-, and and each occurrence of R" is independently hydrogen or an optionally substituted C_1-C_6 aliphatic group, a 3-8-membered saturated, partially unsaturated, or fully unsaturated monocyclic ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur, or an 8-12 membered saturated, partially unsaturated, or fully unsaturated bicyclic ring system having 0-5 heteroatoms independently selected from nitrogen, oxygen, or sulfur; or two occurrences of R", are taken together with the atom(s) to which they are bound to form an optionally substituted 3-12 membered saturated, partially unsaturated, or fully unsaturated monocyclic or bicyclic ring having 0-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur.

20. The compound of claim 19, wherein y is 0, 1, 2, or 3, and each occurrence of R^7 is independently hydrogen, R", -CH_2R", halogen, CN, NO_2, -N(R")$_2$, -CH_2N(R")$_2$, -OR", -CH_2OR", -SR", -CH_2SR", -COOR", -NR"COR", -NR"COOR", -CON(R")$_2$, -SO_2N(R")$_2$, -CONR"$(CH_2)_2$N(R")$_2$, -CONR$(CH_2)_3$N(R")$_2$, -CONR"$(CH_2)_4$N(R")$_2$, -O$(CH_2)_2$OR", O$(CH_2)_3$OR", O$(CH_2)_4$OR", -O$(CH_2)_2$N(R")$_2$, -O$(CH_2)_3$N(R")$_2$, -O$(CH_2)_4$N(R")$_2$, -NR"CH(CH_2OH)R", -NR"CH(CH_2CH_2OH)R", -NR"(CH_2)R", -NR"$(CH_2)_2$R", -NR"$(CH_2)_3$R", -NR"$(CH_2)_4$R", -NR"(CH_2)N(R")$_2$, -NR"$(CH_2)_2$N(R")$_2$, -NR"$(CH_2)_3$N(R")$_2$, -NR"$(CH_2)_4$N(R")$_2$, -NR"(CH_2)OR", -NR"$(CH_2)_2$OR", -NR"$(CH_2)_3$OR", or –NR"$(CH_2)_4$OR".


21. The compound of claim 19, wherein y is 1, 2, or 3 and each occurrence of R^7 is independently F, Cl, Br, CN, OH, NH_2, -CH_2OH, C_1-C_6alkyl, -O(C_1-C_6alkyl), -CH_2O(C_1-C_6alkyl), -CO(C_1-C_6alkyl), -COO(C_1-C_6alkyl), -$NHSO_2$(C_1-C_6alkyl), -SO_2NH_2, -$CONH_2$, -CON(C_1-C_6alkyl), -SO_2(C_1-C_6alkyl), -SO_2phenyl, phenyl, benzyl, -N(C_1-C_6alkyl)$_2$, or -S(C_1-C_6alkyl), wherein each of the foregoing phenyl, benzyl, and C_1-C_6alkyl groups is

independently and optionally substituted, and wherein each of the foregoing C_1-C_6alkyl groups is linear, branched, or cyclic.


22. The compound of claim 1, wherein x is 1, 2, or 3; at least one occurence of R^5 is $-N(R')_2$, $-NR'CH(CH_2OH)R'$, $-NR'CH(CH_2CH_2OH)R'$, $-NR'(CH_2)R'$, $-NR'(CH_2)_2R'$, $NR'(CH_2)N(R')_2$, $-NR'(CH_2)_2N(R')_2$, $-OR'$, $-NR'COR'$, $-NR'COCH_2R'$, or $-NR'CO(CH_2)_2R'$; and R' is a C_1-C_6aliphatic group or a 3-8-membered saturated, partially unsaturated, or fully unsaturated monocyclic ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur, or an 8-12 membered saturated, partially unsaturated, or fully unsaturated bicyclic ring system having 0-5 heteroatoms independently selected from nitrogen, oxygen, or sulfur; or two occurrences of R' are taken together with the atom(s) to which they are bound to form an optionally substituted 3-12 membered saturated, partially unsaturated, or fully unsaturated monocyclic or bicyclic ring having 0-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur, wherein each occurrence of R' is optionally substituted with y occurrences of R^7.

23. The compound of claim 22, wherein R' is hydrogen, C_1-C_6alkyl optionally substituted with 1-3 occurrences of R^7, or is a 5-10-membered monocyclic or bicyclic saturated, partially unsaturated or fully unsaturated ring having 0-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur, wherein the ring is optionally substituted with 1-3 occurrences of R^7.


24. The compound of claim 22, wherein R' is hydrogen, C_1-C_4alkyl optionally substituted with 1-3 occurrences of R^7, or is a ring selected from:




i




ii




iii




iv




v




vi




vii




viii




ix




x




xi




xii




xiii




xiv




xv




xvi




xvii




xviii




xix




xx




xxi




xxii




xxiii




xxiv




xxv




xxvi




xxvii




xxviii




xxix




xxx




xxxi




xxxii




xxxiii




xxxiv




xxxv




xxxvi




xxxvii




xxxviii




xxxix




xL




xLi




xLii




xLiii



xLiv




xLv




xLvi

wherein y is 0-5 and R^7 is =O, =NR", =S, halogen, -CN, -NO_2, or W-R", wherein W is a bond or an optionally substituted C_1-C_6 alkylidene chain, wherein up to two methylene units of the chain are optionally and independently replaced by –NR"-, -S-, -O-, -CS-, -CO_2-, -OCO-, -CO-, -COCO-, -CONR"-, -NR"CO-, -NR"CO_2-, -SO_2NR"-, -NR"SO_2-, -CONR"NR"-, -NR"CONR"-, -OCONR"-, -NR"NR"-, -NR"SO_2NR"-, -SO-, -SO_2-, -PO-, -PO_2-, or –POR"-, and and each occurrence of R" is independently hydrogen or an optionally substituted C_1-C_6 aliphatic group, a 3-8-membered saturated, partially unsaturated, or fully unsaturated monocyclic ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur, or an 8-12 membered saturated, partially unsaturated, or fully unsaturated bicyclic ring system having 0-5 heteroatoms independently selected from nitrogen, oxygen, or sulfur; or two occurrences of R", are taken together with the atom(s) to which they are bound to form an optionally substituted 3-12 membered saturated, partially unsaturated, or fully unsaturated monocyclic or bicyclic ring having 0-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur.


25. The compound of claim 1, wherein R^5 is $-N(R')_2$ and the two occurrences of R' are taken together with the nitrogen atom to which they are bound to form an optionally substituted 3-10-membered monocyclic or bicyclic heterocyclic ring. In certain embodiments, the ring is selected from:




a




b




c




d




e




f




g




h




i




j




k




l



m



n




o

wherein y is 0-5 and R^7 is =O, =NR", =S, halogen, -CN, $-NO_2$, or W-R", wherein W is a bond or an optionally substituted C_1-C_6 alkylidene chain, wherein up to two methylene units of the chain are optionally and independently replaced by –NR"-, -S-, -O-, -CS-, $-CO_2$-, -OCO-, -CO-, -COCO-, -CONR"-, -NR"CO-, $-NR"CO_2$-, $-SO_2NR"$-, $-NR"SO_2$-, -CONR"NR"-, -NR"CONR"-, -OCONR"-, -NR"NR"-, $-NR"SO_2NR"$-, -SO-, $-SO_2$-, -PO-, $-PO_2$-, or –POR"-, and and each occurrence of R" is independently hydrogen or an optionally substituted C_1-C_6 aliphatic group, a 3-8-membered saturated, partially unsaturated, or fully unsaturated monocyclic ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur, or an 8-12 membered saturated, partially unsaturated, or fully unsaturated bicyclic ring system having 0-5 heteroatoms independently selected from nitrogen, oxygen, or

sulfur; or two occurrences of R", are taken together with the atom(s) to which they are bound to form an optionally substituted 3-12 membered saturated, partially unsaturated, or fully unsaturated monocyclic or bicyclic ring having 0-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur.

26. A compound which is:



I-43

27. The compound according to claim 1, wherein R^2 and R^4 are each independently selected from H, Cl, Br, F, -CN, -COOH, -COOR', $-NH_2$, $-N(CH_3)_2$, $-N(Et)_2$, $-N(iPr)_2$, -NHR', -OR', -SR', $-O(CH_2)_2OCH_3$, $-CONH_2$, -CONHR', $CON(R')_2$, -OH, $-CH_2OH$, CH_2NH_2, $-CH_2NHR'$, -CH2N(R')$_2$, $-C(R')_2N(R')_2$, $-NHCOCH_3$, -N(R')COR', $-SO_2NH_2$, $-SO_2N(Me)_2$, or an optionally substituted group selected from C_1-C_6 aliphatic, C_1-C_6 alkyloxy, a 3-8-membered saturated, partially unsaturated, or fully unsaturated monocyclic ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur, or an 8-12 membered saturated, partially unsaturated, or fully unsaturated bicyclic ring system having 0-5 heteroatoms independently selected from nitrogen, oxygen, or sulfur.

28. The compound according to claim 27, wherein R^2 and R^4 are each hydrogen.

29. The compound according to claim 1, wherein R^1 is selected from hydrogen, C_1-C_4alkyl, -COR', -SO_2R', or –Si(R')$_3$.

30. The compound according to claim 29, wherein R^1 is hydrogen.

31. The compound according to claim 1 wherein R^3 is selected from H, Cl, Br, F , -CN, -COOH, -COOMe, -NH_2, –N(R')$_2$, -NO_2, -OR', -CON(R')$_2$, -COOR', -OH, -SR', -C(R')$_2$OR', -N(R')COR', -N(R')C(O)OR', -SO_2NH_2, -SO_2N(R')$_2$, or an optionally substituted group selected from C_1-C_4 aliphatic, C_1-C_4 alkyloxy or -C≡C-C_1-C_4 aliphatic.

32. The compound according to claim 31, wherein R^3 is selected from H, Cl, -Br, -CN, -COOH, -COOMe, -CONHR', -CON(Me)$_2$, -CH_2OH, -NO_2, -NH_2 or an optionally substituted C_1-C_4 aliphatic.

33. The compound according to claim 32, wherein R^3 is selected from Cl, -Br, -CN or an optionally substituted C_1-C_4 aliphatic.

34. The compound according to claim 32, wherein R^3 is Cl.

35. The compound according to claim 1, wherein R^1, R^2 and R^4 are hydrogen and R^3 is selected from Cl, -Br, -CN or an optionally substituted C_1-C_4 aliphatic.

36. The compound according to claim 37, wherein R^3 is H, OR', NR'C(O)R', NR'C(O)OR', CON(R')$_2$ or COOMe.

37. The compound according to claim 32, wherein R^3 is OR', NR'C(O)R' or NR'C(O)OR'.

38. The compound according claim 1, wherein R^1, R^2 and R^4 are hydrogen and R^3 is selected from H, OR', NR'C(O)R', NR'C(O)OR', CON(R')$_2$ or COOMe.

39. The compound according to claim 1, wherein x is 1, 2, or 3; wherein each occurrence of R^5 is independently selected from R', -CH_2R', halogen, CN, NO_2, -N(R')$_2$, -CH_2N(R')$_2$, -OR', -CH_2OR', -SR', -CH_2SR', -COOR', -NR'COR', -NR'COR^{8}R', -NR'COOR', -CON(R')$_2$, -SO_2N(R')$_2$, -CONR'R^{8}N(R')$_2$, -OR^{8}OR', -OR^{8}N(R')$_2$, -NR'CH(R^9)R', -NR'CH(R^9)C(O)OR', -N(R')R^{8}R', -N(R')R^{8}R', -N(R')R^{8}N(R')$_2$, -N(R')R^{8}OR', -NR'CH(R^9)R', -NR'CH_2C(O)N(R')$_2$, or -NR'CH(R^9)C(O)N(R')$_2$, wherein R^8 is an optionally substituted C_1-C_4 alkyl and R^9 is an optionally substituted C_1-C_6 aliphatic.

40. The compound according to claim 39, wherein each occurrence of R^5 is independently selected from R', $-CH_2R'$, halogen, $-CN$, $-NO_2$, $-N(R')_2$, $-CH_2N(R')_2$, $-OR'$, $-CH_2OR'$, $-SR'$, $-CH_2SR'$, $-COOR'$, $-NR'COR'$, $-NR'COCH_2R'$, $-NR'CO(CH_2)_2R'$, $-NR'COOR'$, $-CON(R')_2$, $-SO_2N(R')_2$, $-CONR'(CH_2)_2N(R')_2$, $-CONR(CH_2)_3N(R')_2$, $-CONR'(CH_2)_4N(R')_2$, $-O(CH_2)_2OR'$, $O(CH_2)_3OR'$, $O(CH_2)_4OR'$, $-O(CH_2)_2N(R')_2$, $-O(CH_2)_3N(R')_2$, $-O(CH_2)_4N(R')_2$, $-NR'CH(CH_2OR^9)R'$, $-NR'CH(CH_2CH_2OR^9)R'$, $-NR'CH(CH_3)R'$, $NR'CH(CF_3)R'$, $-NR'CH(CH_3)C(O)OR'$, $-NR'CH(CF_3)C(O)OR'$,$-NR'(CH_2)R'$, $-NR'(CH_2)_2R'$, $-NR'(CH_2)_3R'$, $-NR'(CH_2)_4R'$, $-NR'(CH_2)N(R')_2$, $-NR'(CH_2)_2N(R')_2$, $-NR'(CH_2)_3N(R')_2$, $-NR'(CH_2)_4N(R')_2$, $-NR'(CH_2)OR'$, $-NR'(CH_2)_2OR'$, $-NR'(CH_2)_3OR'$, $-NR'(CH_2)_4OR'$, $-NR'CH(CH_2CH_3)R'$, $-NR'CH_2C(O)N(R')_2$, $-NR'CH(CH_3)C(O)N(R')_2$, $NR'CH(CF_3)C(O)N(R')_2$, $-NR'CH(CH_2CH_3)C(O)N(R')_2$, $-NR'CH(CH(CH_3)_2)C(O)N(R')_2$, $-NR'CH(C(CH_3)_3)C(O)N(R')_2$, $-NR'CH(CH_2CH(CH_3)_2)C(O)N(R')_2$, $-NR'CH(CH_2OR^9)C(O)N(R')_2$ or $-NR'CH(CH_2CH_2N(Me)_2)C(O)N(R')_2$.

41. The compound according to claim 39, wherein at least one occurrence of R^5 is independently selected from $-NR'CH(CH_2OH)R'$, $-NR'CH(CH_2OMe)R'$, $-NR'CH(CH_2OEt)R'$, $-NR'CH(CH_2OCF_3)R'$, $-NR'CH(CH_2CH_2OH)R'$, $-NR'CH(CH_2CH_2OMe)R'$, $-NR'CH(CH_2CH_2OEt)R'$, $-NR'CH(CH_2CH_2OCF_3)R'$, $-NR'CH(CH_3)C(O)OR'$, $-NR'CH(CF_3)C(O)OR'$, $-NR'CH(CH_3)C(O)N(R')_2$, $-NR'CH(CF_3)C(O)N(R')_2$, $-NR'CH(CH_2CH_3)C(O)N(R')_2$, $-NR'CH(CH_2OH)C(O)N(R')_2$, $-NR'CH(CH_2OMe)C(O)N(R')_2$, $-NR'CH(CH_2OEt)C(O)N(R')_2$ or $-NR'CH(CH_2OCF_3)C(O)N(R')_2$, wherein R' is an optionally substituted C_1-C_4 aliphatic.

42. The compound according to claim 39, wherein said at least one occurrence of R^5 is selected from $NHCH_2C(O)NHR'$, $-NHCH(CH_3)C(O)NHR'$, $-NHCH(CH_2CH_3)C(O)NHR'$, $-NHCH(CH(CH_3)_2)C(O)NHR'$, -

$NHCH(C(CH_3)_3)C(O)NHR'$, $-NHCH(CH_2CH(CH_3)_2)C(O)NHR'$, $-NHCH(CH_2OH)C(O)NHR'$, $-NHCH(CH_2OMe)C(O)NHR'$ or $-NHCH(CH_2CH_2N(Me)_2)C(O)NHR'$, wherein R' is an optionally substituted C_1-C_4 aliphatic.

43. The compound according to claim 39, wherein at least one occurrence of R^5 is independently selected from $-NHR'$, $-NH(CH_2)R'$, $-NH(CH_2)_2R'$, $-NHCH(CH_3)R'$, $-NHCH_2C(O)NHR'$, $-NHCH(CH_3)C(O)NHR'$, $-NHCH(CH_2CH_3)C(O)NHR'$, $-NHCH(CH(CH_3)_2)C(O)NHR'$, $-NHCH(C(CH_3)_3)C(O)NHR'$, $-NHCH(CH_2CH(CH_3)_2)C(O)NHR'$, $-NHCH(CH_2OH)C(O)NHR'$, $-NHCH(CH_2OMe)C(O)NHR'$ or $-NHCH(CH_2CH_2N(Me)_2)C(O)NHR'$, wherein R' is an optionally substituted phenyl.

44. The compound according to claim 39, wherein said at least one occurrence of R^5 is $-NHCH(CH_3)R'$, wherein R' is optionally substituted phenyl.


45. The compound according to claim 39, wherein at least one occurrence of R^5 is H, halogen, CH_3, CF_3, COOH, COOMe or OR', wherein R' is C_1-C_4 aliphatic.

46. The compound according to claim 39, wherein x is 2 or 3 and at least one occurrence of R^5 is F.


47. The compound according to claim 39, wherein R^1, R^2 and R^4 are hydrogen and R^5 is selected from -NHR', -NH(CH_2)R', -NH(CH_2)$_2$R', -NHCH(CH_3)R', -NHCH$_2$C(O)NHR', -NHCH(CH_3)C(O)NHR', -NHCH(CH_2CH_3)C(O)NHR', -NHCH(CH(CH_3)$_2$)C(O)NHR', -NHCH(C(CH_3)$_3$)C(O)NHR', -NHCH(CH_2CH(CH_3)$_2$)C(O)NHR', -NHCH(CH_2OH)C(O)NHR', -NHCH(CH_2OMe)C(O)NHR' or -NHCH(CH_2CH_2N(Me)$_2$)C(O)NHR', wherein R' is an optionally substituted phenyl.

48. The compound according to claim 39, wherein R^1, R^2 and R^4 are hydrogen and R^5 is selected from -NR'CH(CH_2OH)R', -NR'CH(CH_2OMe)R', -NR'CH(CH_2OEt)R', -NR'CH(CH_2OCF_3)R', -NR'CH(CH_2CH_2OH)R', -NR'CH(CH_2CH_2OMe)R', -NR'CH(CH_2CH_2OEt)R', -NR'CH($CH_2CH_2OCF_3$)R', -NR'CH(CH_3)C(O)OR', -NR'CH(CF_3)C(O)OR', -NR'CH(CH_3)C(O)N(R')$_2$, -NR'CH(CF_3)C(O)N(R')$_2$, -NR'CH(CH_2CH_3)C(O)N(R')$_2$, -NR'CH(CH_2OH)C(O)N(R')$_2$, -NR'CH(CH_2OMe)C(O)N(R')$_2$, -NR'CH(CH_2OEt)C(O)N(R')$_2$ or -NR'CH(CH_2OCF_3)C(O)N(R')$_2$, wherein R' is an optionally substituted C_1-C_4 aliphatic.


49. The compound according to claim 1 , wherein R' is selected from hydrogen, a C_1-C_6aliphatic group optionally substituted with y occurrences of R^7, or R' is a ring selected from:




i




ii




iii




iv




v




vi




vii




viii




ix




x




xi




xii




xiii




xiv




xv




xvi




xvii




xviii




xix




xx




xxi




xxii




xxiii




xxiv




xxv




xxvi




xxvii




xxviii




xxix




xxx




xxxi




xxxii




xxxiii




xxxiv




xxxv




xxxvi




xxxvii




xxxviii




xxxix




xL




xLi




xLii




xLiii




xLiv



xLv




xLvi



xLvii


or two occurrences of R' are taken together with the nitrogen atom to which they are bound to form an optionally substituted 3-10-membered monocyclic or bicyclic heterocyclic ring selected from:




a




b




c




d




e




f




g




h




i




j




k




l



m



n




o

wherein y is 0, 1, 2, or 3, and each occurrence of R^7 is independently hydrogen, R", -CH_2R", halogen, CN, NO_2, -N(R")$_2$, -CH_2N(R")$_2$, -OR", -CH_2OR", -SR", -CH_2SR", -COOR", -NR"COR", -NR"COOR", -CON(R")$_2$, -SO_2N(R")$_2$, -CONR"$(CH_2)_2$N(R")$_2$, -CONR$(CH_2)_3$N(R")$_2$, -CONR"$(CH_2)_4$N(R")$_2$, -O$(CH_2)_2$OR", O$(CH_2)_3$OR", O$(CH_2)_4$OR", -O$(CH_2)_2$N(R")$_2$, -O$(CH_2)_3$N(R")$_2$, -O$(CH_2)_4$N(R")$_2$, -NR"CH(CH_2OH)R", -NR"CH(CH_2CH_2OH)R", -NR"(CH_2)R", -NR"$(CH_2)_2$R", -NR"$(CH_2)_3$R", -NR"$(CH_2)_4$R", -NR"(CH_2)N(R")$_2$, -NR"$(CH_2)_2$N(R")$_2$, -NR"$(CH_2)_3$N(R")$_2$, -NR"$(CH_2)_4$N(R")$_2$, -NR"(CH_2)OR", -NR"$(CH_2)_2$OR", -NR"$(CH_2)_3$OR", or –NR"$(CH_2)_4$OR", and each occurrence of R" is independently hydrogen or an optionally substituted C_1-C_6 aliphatic group, a 3-8-membered saturated, partially unsaturated, or fully unsaturated monocyclic ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur, or an 8-12 membered saturated, partially unsaturated, or fully unsaturated bicyclic ring system


having 0-5 heteroatoms independently selected from nitrogen, oxygen, or sulfur; or two occurrences of R", are taken together with the atom(s) to which they are bound to form an optionally substituted 3-12 membered saturated, partially unsaturated, or fully unsaturated monocyclic or bicyclic ring having 0-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur.

50. The compound according to claim 1, wherein x is 2, 3 or 4 and at least two R^5 on adjacent ring members are R', and wherein said two R^5 are taken together with the atoms to which they are bound to form an optionally substituted 3-12 membered saturated, partially saturated or fully unsaturated monocyclic or bicyclic ring having 0-4 heteroatoms independently selected from nitrogen, oxygen or sulfur.

51. The compound according to claim 50, wherein said compound has formula I-B-vii or I-B-viii:



I-B-vii




I-B-viii



wherein y is 0, 1, 2, or 3, and each occurrence of R^7 is independently hydrogen, R", -CH_2R", halogen, CN, NO_2, -N(R")$_2$, -CH_2N(R")$_2$, -OR", -CH_2OR", -SR", -CH_2SR", -COOR", -NR"COR", -NR"COOR", -CON(R")$_2$, -SO_2N(R")$_2$, -CONR"$(CH_2)_2$N(R")$_2$, -CONR$(CH_2)_3$N(R")$_2$, -CONR"$(CH_2)_4$N(R")$_2$, -O$(CH_2)_2$OR", O$(CH_2)_3$OR", O$(CH_2)_4$OR", -O$(CH_2)_2$N(R")$_2$, -O$(CH_2)_3$N(R")$_2$, -O$(CH_2)_4$N(R")$_2$, -



NR"CH(CH_2OH)R", -NR"CH(CH_2CH_2OH)R", -NR"(CH_2)R", -NR"(CH_2)$_2$R", -NR"(CH_2)$_3$R", -NR"(CH_2)$_4$R", -NR"(CH_2)N(R")$_2$, -NR"(CH_2)$_2$N(R")$_2$, -NR"(CH_2)$_3$N(R")$_2$, -NR"(CH_2)$_4$N(R")$_2$, -NR"(CH_2)OR", -NR"(CH_2)$_2$OR", -NR"(CH_2)$_3$OR", or –NR"(CH_2)$_4$OR", and each occurrence of R" is independently hydrogen or an optionally substituted C_1-C_6 aliphatic group, a 3-8-membered saturated, partially unsaturated, or fully unsaturated monocyclic ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur, or an 8-12 membered saturated, partially unsaturated, or fully unsaturated bicyclic ring system having 0-5 heteroatoms independently selected from nitrogen, oxygen, or sulfur; or two occurrences of R", are taken together with the atom(s) to which they are bound to form an optionally substituted 3-12 membered saturated, partially unsaturated, or fully unsaturated monocyclic or bicyclic ring having 0-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur.



52. A compound selected from: Table 2.



Table 2. Examples of Compounds of Formula **I**:



Cmpd No.	Compound
1	



Cmpd No.	Compound
2	
3	



Cmpd No.	Compound
4	
5	



Cmpd No.	Compound
6	
7	



Cmpd No.	Compound
8	
9	



Cmpd No.	Compound
10	
11	



Cmpd No.	Compound
12	
13	



Cmpd No.	Compound
14	
15	



Cmpd No.	Compound
16	
17	



Cmpd No.	Compound
18	
19	



Cmpd No.	Compound
20	
21	



Cmpd No.	Compound
22	
23	



Cmpd No.	Compound
24	
25	



Cmpd No.	Compound
26	
27	



Cmpd No.	Compound
28	
29	



Cmpd No.	Compound
30	
31	



Cmpd No.	Compound
32	
33	



Cmpd No.	Compound
34	
35	



Cmpd No.	Compound
36	
37	



Cmpd No.	Compound
38	
39	



Cmpd No.	Compound
40	
41	



Cmpd No.	Compound
42	
43	



Cmpd No.	Compound
44	
45	



Cmpd No.	Compound
46	
47	



Cmpd No.	Compound
48	
49	



Cmpd No.	Compound
50	
51	



Cmpd No.	Compound
52	
53	



Cmpd No.	Compound
54	
55	



Cmpd No.	Compound
56	
57	



Cmpd No.	Compound
58	
59	



Cmpd No.	Compound
60	
61	



Cmpd No.	Compound
62	
63	



Cmpd No.	Compound
64	
65	



Cmpd No.	Compound
66	
67	



Cmpd No.	Compound
68	
69	



Cmpd No.	Compound
70	
71	



Cmpd No.	Compound
72	
73	



Cmpd No.	Compound
74	
75	



Cmpd No.	Compound
76	
77	



Cmpd No.	Compound
78	
79	



Cmpd No.	Compound
80	
81	



Cmpd No.	Compound
82	
83	



Cmpd No.	Compound
84	
85	



Cmpd No.	Compound
86	
87	



Cmpd No.	Compound
88	
89	



Cmpd No.	Compound
90	
91	



Cmpd No.	Compound
92	
93	



Cmpd No.	Compound
94	
95	



Cmpd No.	Compound
96	
97	



Cmpd No.	Compound
98	
99	



Cmpd No.	Compound
100	
101	



Cmpd No.	Compound
102	
103	



Cmpd No.	Compound
104	
105	



Cmpd No.	Compound
106	
107	



Cmpd No.	Compound
108	
109	



Cmpd No.	Compound
110	
111	



Cmpd No.	Compound
112	
113	



Cmpd No.	Compound
114	
115	



Cmpd No.	Compound
116	
117	



Cmpd No.	Compound
118	
119	



Cmpd No.	Compound
120	
121	



Cmpd No.	Compound
122	
123	



Cmpd No.	Compound
124	
125	



Cmpd No.	Compound
126	
127	



Cmpd No.	Compound
128	
129	



Cmpd No.	Compound
130	
131	



Cmpd No.	Compound
132	
133	



Cmpd No.	Compound
134	
135	



Cmpd No.	Compound
136	
137	



Cmpd No.	Compound
138	
139	



Cmpd No.	Compound
140	
141	



Cmpd No.	Compound
142	
143	



Cmpd No.	Compound
144	
145	



Cmpd No.	Compound
146	
147	



Cmpd No.	Compound
148	
149	



Cmpd No.	Compound
150	
151	



Cmpd No.	Compound
152	
153	



Cmpd No.	Compound
154	
155	



Cmpd No.	Compound
156	
157	



Cmpd No.	Compound
158	
159	



Cmpd No.	Compound
160	
161	



Cmpd No.	Compound
162	
163	



Cmpd No.	Compound
164	
165	



Cmpd No.	Compound
166	
167	



Cmpd No.	Compound
168	
169	



Cmpd No.	Compound
170	
171	



Cmpd No.	Compound
172	
173	



Cmpd No.	Compound
174	
175	



Cmpd No.	Compound
176	
177	



Cmpd No.	Compound
178	
179	



Cmpd No.	Compound
180	
181	



Cmpd No.	Compound
182	
183	



Cmpd No.	Compound
184	
185	



Cmpd No.	Compound
186	
187	



Cmpd No.	Compound
188	
189	



Cmpd No.	Compound
190	
191	



Cmpd No.	Compound
192	
193	



Cmpd No.	Compound
194	
195	



Cmpd No.	Compound
196	
197	



Cmpd No.	Compound
198	
199	



Cmpd No.	Compound
200	
201	



Cmpd No.	Compound
202	
203	



Cmpd No.	Compound
204	
205	



Cmpd No.	Compound
206	
207	



Cmpd No.	Compound
208	
209	



Cmpd No.	Compound
210	
211	



Cmpd No.	Compound
212	
213	



Cmpd No.	Compound
214	
215	



Cmpd No.	Compound
216	
217	



Cmpd No.	Compound
218	
219	



Cmpd No.	Compound
220	
221	



Cmpd No.	Compound
222	
223	



Cmpd No.	Compound
224	
225	



Cmpd No.	Compound
226	
227	



Cmpd No.	Compound
228	
229	



Cmpd No.	Compound
230	
231	



Cmpd No.	Compound
232	
233	



Cmpd No.	Compound
234	
235	



Cmpd No.	Compound
236	
237	



Cmpd No.	Compound
238	
239	



Cmpd No.	Compound
240	
241	



Cmpd No.	Compound
242	
243	



Cmpd No.	Compound
244	
245	



Cmpd No.	Compound
246	
247	



Cmpd No.	Compound
248	
249	



Cmpd No.	Compound
250	
251	



Cmpd No.	Compound
252	
253	



Cmpd No.	Compound
254	
255	



Cmpd No.	Compound
256	
257	



Cmpd No.	Compound
258	
259	



Cmpd No.	Compound
260	
261	



Cmpd No.	Compound
262	
263	



Cmpd No.	Compound
264	
265	



Cmpd No.	Compound
266	
267	



Cmpd No.	Compound
268	
269	



Cmpd No.	Compound
27O	
271	



Cmpd No.	Compound
272	
273	



Cmpd No.	Compound
274	
275	



Cmpd No.	Compound
276	
277	



Cmpd No.	Compound
278	
279	



Cmpd No.	Compound
280	
281	



Cmpd No.	Compound
282	
283	



Cmpd No.	Compound
284	
285	



Cmpd No.	Compound
286	
287	



Cmpd No.	Compound
288	
289	



Cmpd No.	Compound
290	
291	



Cmpd No.	Compound
292	
293	



Cmpd No.	Compound
294	
295	



Cmpd No.	Compound
296	
297	



Cmpd No.	Compound
298	
299	



Cmpd No.	Compound
300	
301	



Cmpd No.	Compound
302	
303	



Cmpd No.	Compound
304	
305	



Cmpd No.	Compound
306	
307	



Cmpd No.	Compound
308	
309	



Cmpd No.	Compound
310	
311	



Cmpd No.	Compound
312	
313	



Cmpd No.	Compound
314	
315	



Cmpd No.	Compound
316	
317	



Cmpd No.	Compound
318	
319	



Cmpd No.	Compound
320	
321	



Cmpd No.	Compound
322	
323	



Cmpd No.	Compound
324	
325	



Cmpd No.	Compound
326	
327	



Cmpd No.	Compound
328	
329	



Cmpd No.	Compound
330	
331	



Cmpd No.	Compound
332	
333	



Cmpd No.	Compound
334	
335	



Cmpd No.	Compound
336	
337	



Cmpd No.	Compound
338	
339	



Cmpd No.	Compound
340	
341	



Cmpd No.	Compound
342	
343	



Cmpd No.	Compound
344	
345	



Cmpd No.	Compound
346	
347	



Cmpd No.	Compound
348	
349	



Cmpd No.	Compound
350	
351	



Cmpd No.	Compound
352	
353	



Cmpd No.	Compound
354	
355	



Cmpd No.	Compound
356	
357	



Cmpd No.	Compound
358	
359	



Cmpd No.	Compound
360	
361	



Cmpd No.	Compound
362	
363	



Cmpd No.	Compound
364	
365	



Cmpd No.	Compound
366	
367	



Cmpd No.	Compound
368	
369	



Cmpd No.	Compound
370	
371	



Cmpd No.	Compound
372	
373	



Cmpd No.	Compound
374	
375	



Cmpd No.	Compound
376	
377	



Cmpd No.	Compound
378	
379	



Cmpd No.	Compound
380	
381	



Cmpd No.	Compound
382	
383	



Cmpd No.	Compound
384	
385	



Cmpd No.	Compound
386	
387	



Cmpd No.	Compound
388	
389	



Cmpd No.	Compound
390	
391	



Cmpd No.	Compound
392	
393	



Cmpd No.	Compound
394	
395	



Cmpd No.	Compound
396	
397	



Cmpd No.	Compound
398	
399	



Cmpd No.	Compound
400	
401	



Cmpd No.	Compound
402	
403	



Cmpd No.	Compound
404	
405	



Cmpd No.	Compound
406	
407	



Cmpd No.	Compound
408	
409	



Cmpd No.	Compound
410	
411	



Cmpd No.	Compound
412	
413	



Cmpd No.	Compound
414	
415	



Cmpd No.	Compound
416	
417	



Cmpd No.	Compound
418	
419	



Cmpd No.	Compound
420	
421	



Cmpd No.	Compound
422	
423	



Cmpd No.	Compound
424	
425	



Cmpd No.	Compound
426	
427	



Cmpd No.	Compound
428	
429	



Cmpd No.	Compound
430	
431	



Cmpd No.	Compound
432	
433	



Cmpd No.	Compound
434	
435	



Cmpd No.	Compound
436	
437	



Cmpd No.	Compound
438	
439	



Cmpd No.	Compound
440	
441	



Cmpd No.	Compound
442	
443	



Cmpd No.	Compound
444	
445	



Cmpd No.	Compound
446	
447	



Cmpd No.	Compound
448	
449	



Cmpd No.	Compound
450	
451	



Cmpd No.	Compound
452	
453	



Cmpd No.	Compound
454	
455	



Cmpd No.	Compound
456	
457	



Cmpd No.	Compound
458	
459	

Cmpd No.	Compound
460	
461	

Cmpd No.	Compound
462	
463	

Cmpd No.	Compound
464	
465	

Cmpd No.	Compound
466	
467	

Cmpd No.	Compound
468	
469	

53. A pharmaceutical composition comprising a compound according to any one of claims 1, 26 or 52 and a pharmaceutically acceptable carrier, adjuvant, or vehicle.

54. The composition of claim 53, further comprising an additional therapeutic agent selected from a chemotherapeutic or anti-proliferative agent, a treatment for Alzheimer's Disease, a treatment for Parkinson's Disease, an agent for treating Multiple Sclerosis (MS), a treatment for asthma, an agent for treating schizophrenia, an anti-inflammatory agent, an immunomodulatory or immunosuppressive agent, a neurotrophic factor, an agent for treating cardiovascular disease, an agent for treating destructive bone disorders, an agent for treating liver disease, an agent for treating a blood disorder, or an agent for treating an immunodeficiency disorder.

55. A use of

a) a composition according to claim 53; or
b) a compound according to claim 1

in the manufacture of a medicament for inhibiting JAK-3, ROCK or Aurora kinase activity in a patient.

56. A method of inhibiting JAK 3, ROCK or Aurora kinase activity in a biological sample comprising the step of contacting said biological sample with:

a) a composition according to claim 53; or
b) a compound according to claim 1.

57. A use of a composition according to claim 53 or a compound according to claim 1 in the manufacture of a medicament for treating or alleviating a disease or disorder selected from immune responses such as allergic or type I hypersensitivity reactions or asthma; autoimmune diseases such as transplant rejection, graft versus host disease, rheumatoid arthritis, amyotrophic lateral sclerosis, or multiple sclerosis; neurodegenerative disorders; or solid and hematologic malignancies.

58. The use according to claim 57, wherein the medicament comprises an additional therapeutic agent.

59. The use according to claim 57, wherein the disease is an autoimmune disease and is transplant rejection, graft versus host disease, rheumatoid arthritis, amyotrophic lateral sclerosis, or multiple sclerosis.

60. The use according to claim 57, wherein the disease is hypertension, angina pectoris, cerebrovascular contraction, asthma, peripheral circulation disorder, premature birth, cancer, erectile dysfunction, arteriosclerosis, spasm (cerebral vasospasm, and coronary vasospasm), retinopathy (e.g., glaucoma), inflammatory disorders, autoimmune disorders, AIDS, osteoporosis, myocardial hypertrophy, ischemia/reperfusion-induced injury, or endothelial dysfunction.

61. A use of a composition according to claim 53 or a compound according to claim 1 in the manufacture of a medicament for treating, preventing or lessening the severity of a disease or disorder selected from heart disease, diabetes, Alzheimer's disease, immunodeficiency disorders, inflammatory diseases, hypertension, allergic diseases, autoimmune diseases, destructive bone disorders, osteoporosis, proliferative disorders, infectious diseases, immunologically-mediated diseases, viral diseases, hyperplasia, reperfusion/ischemia in stroke, heart attacks, organ hypoxia., thrombin-induced platelet aggregation, chronic myelogenous leukemia (CML), acute myeloid leukemia (AML), acute promyelocyte leukemia (APL), rheumatoid artiuitis, asthma, osteoarthritis, ischemia, cancer, hepatic ischemia, myocardi al infarction, congestive heart failure, pathologic immune conditions involving T cell activation, and neurodegenerative disorders.

62. The use according to claim 61, wherein said cancer is selected from melanoma, lymphoma, neuroblastoma, leukemia, or a cancer selected from colon, breast, lung, kidney, ovary, endometrial, pancreatic, renal, central nervous system, cervical, prostate, or cancer of the gastric tract, CML, AML, APL, acute lymphocytic leukemia (ALL), mastocytosis or gastrointestinal stromal tumor (GIST).