


CLAIMS

1. A compound represented by general formula (Ib):



wherein

ring A is an optionally substituted 5-membered or 6-membered hydrocarbon cyclic ring, or an optionally substituted 5-membered or 6-membered heterocyclic ring having up to three heteroatoms independently selected from the group consisting of O, N, S and NH;
 10 Y1 and Y2 are each independently selected from the group consisting of:

a single bond,

15

—Z—O—Z—,

—Z—NR^b—Z—,

—Z—O—C(=O)—Z—, —Z—C(=O)—O—Z—,

—Z—NR^b—C(=O)—Z—, —Z—C(=O)—NR^b—Z—,

—Z—NR^b—C(=O)—O—Z—, —Z—O—C(=O)—NR^b—Z—;

20

where each Z is independently selected from the group consisting of a single bond, optionally substituted saturated aliphatic chain and optionally substituted unsaturated aliphatic chain;

where R^b is H or C₁-C₆ alkyl, and

with the proviso that when ring A is 1,4-phenylene, Y1 is not





25

—(CH₂)—O—C(=O)— and Y2 is not —C(=O)—O—(CH₂)—.

2. The compound according to claim 1, wherein ring A is selected from the group consisting of benzene, furan, thiophene, pyrrole, oxazole, isoxazole, isothiazole, tetrahydrofuran, tetrahydrothiophene,

pyrrolidine, pyridine, pyrone, pyridazine, pyrimidine, pyrazine, triazine, piperidine and piperazine.

3. The compound according to any one of claims 1 to 2, wherein ring A is
5 selected from any one of the general formulae (II) to (V):




wherein

R^{3a}, R^{3b} and R^{3c} are each independently selected from the group
consisting of a hydrogen, hydroxy, halogen, cyano, amino, nitro,
carboxyl, alkyl, alkenyl, alkynyl, alkoxy, haloalkyl, haloalkoxy,
15 cycloalkyl, cycloalkenyl, alkylcarbonyl and alkoxy carbonyl.


4. The compound according to claim 1, wherein
Y¹ is selected from the group consisting of –Z–O–Z– and
–Z–O–C(=O)–Z–;
20 Y² is selected from the group consisting of –Z–O–Z– and
–Z–C(=O)–O–Z–;
each Z is independently selected from the group consisting of a single
bond and C₁–C₆ alkyl.

5. The compound according to any one of claims 1 to 4 selected from the following:




bis((2-oxo-1,3-dioxolan-4-yl)methyl)furan-2,5-dicarboxylate (**FBC-1**);


5



4,4'-(((furan-2,5-diylbis(methylene))bis(oxy))bis(methylene))bis(1,3-dioxolan-2-one) (**FBC-2**);




- 10 bis((2-oxo-1,3-dioxolan-4-yl)methyl)pyridine-2,5-dicarboxylate (**PBC**);



bis((2-oxo-1,3-dioxolan-4-yl)methyl)pyridine-2,6-dicarboxylate (**PBC-2**); and


15



4,4'-(((tetrahydrofuran-3,4-diyl)bis(oxy))bis(methylene))bis(1,3-dioxolan-2-one) (**HFBC**).

6. A method of preparing the compound according to any one of claims 1 to 5, the method comprising:

converting a precursor compound represented by general formula (VI) to the compound of any one of claims 1-5 through one or more chemical reactions:




wherein

- ring A is an optionally substituted 5-membered or 6-membered hydrocarbon cyclic ring, or an optionally substituted 5-membered or 6-membered heterocyclic ring having up to three heteroatoms independently selected from the group consisting of O, N, S and NH;
 R^4 and R^5 are each independently selected from the group consisting of $-OH$, $-C(=O)H$, $-C(=O)-OH$, $-NR^cR^d$, $-C(=O)-NR^cR^d$, $-(C_1-C_6 \text{ alkyl})-OH$, $-(C_1-C_6 \text{ alkyl})-C(=O)H$, $-(C_1-C_6 \text{ alkyl})-C(=O)-OH$, $-(C_1-C_6 \text{ alkyl})-NR^cR^d$ and $-(C_1-C_6 \text{ alkyl})-C(=O)-NR^cR^d$,
where R^c and R^d are independently selected from the group consisting of H or C_1-C_6 alkyl,
wherein at least one of the one or more chemical reactions is carried out in the presence of a halogenated compound.

7. The method according to claim 6, wherein ring A is selected from the group consisting of disubstituted benzene, disubstituted furan, disubstituted thiophene, disubstituted pyrrole, disubstituted oxazole, disubstituted isoxazole, disubstituted isothiazole, disubstituted tetrahydrofuran, disubstituted tetrahydrothiophene, disubstituted pyrrolidine, disubstituted pyridine, disubstituted pyrone, disubstituted pyridazine, disubstituted pyrimidine, disubstituted pyrazine,

disubstituted triazine, disubstituted piperidine and disubstituted piperazine.

8. The method according to any one of claims 6 to 7, wherein the precursor compound is selected from the group consisting of 5-hydroxymethylfurfural (HMF), furan-2,5-diyl dimethanol (FDM), furan-2,5-dicarboxylic acid (FDCA), furan-2,5-diyl dimethanamine (FBA), pyridine-2,5-dicarboxylic acid, pyridine-2,6-dicarboxylic acid and 1,4-anhydroerythritol.
9. A reaction product of the reaction between one or more compounds according to any one of claims 1 to 5 and one or more amine containing compounds, the reaction product having hydroxyl groups and urethane/carbamate linkages.
10. The reaction product according to claim 9, wherein the reaction product is a polymer having a repeating unit represented by general formula (VIIb) or a derivative thereof:



(VIIb)

wherein

ring A is an optionally substituted 5-membered or 6-membered hydrocarbon cyclic ring, or an optionally substituted 5-membered or 6-membered heterocyclic ring having up to three heteroatoms independently selected from the group consisting of O, N, S and NH; Y¹ and Y² are each independently selected from the group consisting of:

a single bond,

–Z–O–Z–,

–Z–NR^b–Z–,

–Z–O–C(=O)–Z–, –Z–C(=O)–O–Z–,

5 –Z–NR^b–C(=O)–Z–, –Z–C(=O)–NR^b–Z–,

–Z–NR^b–C(=O)–O–Z–, –Z–O–C(=O)–NR^b–Z–;

where each Z is independently selected from the group consisting of a single bond, optionally substituted saturated aliphatic chain and optionally substituted unsaturated aliphatic chain;

10 where R^b is H or C₁–C₆ alkyl;





R⁶ is selected from the group consisting of optionally substituted alkyl, optionally substituted alkenyl, optionally substituted alkynyl, optionally substituted polyether, optionally substituted 5-membered or 6-membered hydrocarbon cyclic ring and an optionally substituted 5-membered or 6-membered heterocyclic ring having up to three heteroatoms independently selected from the group consisting of O, and S;

X³ and X⁴ are each independently selected from the group consisting of a single bond and –(C₁–C₆ alkyl); and

20 with the proviso that when ring A is 1,4-phenylene, Y¹ is not –(CH₂)–O–C(=O)– and Y² is not –C(=O)–O–(CH₂)–.

11. The reaction product according to any one of claims 9 to 10, wherein ring A is selected from the group consisting of benzene, furan, thiophene, pyrrole, oxazole, isoxazole, isothiazole, tetrahydrofuran, tetrahydrothiophene, pyrrolidine, pyridine, pyrone, pyridazine, pyrimidine, pyrazine, triazine, piperidine and piperazine.

12. The reaction product according to any one of claims 9 to 11, wherein ring A is selected from any one of the general formulae (II) to (V):




wherein


- 10 R^{3a} , R^{3b} and R^{3c} are each independently selected from the group consisting of a hydrogen, hydroxy, halogen, cyano, amino, nitro, carboxyl, alkyl, alkenyl, alkynyl, alkoxy, haloalkyl, haloalkoxy, cycloalkyl, cycloalkenyl, alkylcarbonyl and alkoxy carbonyl.

- 15 13. The reaction product according to any one of claims 10 to 12, wherein Y^1 is selected from the group consisting of $-Z-O-Z-$ and $-Z-O-C(=O)-Z-$;
 Y^2 is selected from the group consisting of $-Z-O-Z-$ and $-Z-C(=O)-O-Z-$;
 20 each Z is independently selected from the group consisting of a single bond and C₁-C₆ alkyl.


14. The reaction product according to any one of claims 9 to 13 selected from the following:




(1)




(2)



(3)




(4)




(5)


98




(6)




(7)




(8)



(9)




(10)




(11)


99




(12)




(13)




(14)



(15)




(16)




(17)


100



(18)



(19)



(20)

or a derivative thereof.

15. The reaction product according to any one of claims 9 to 14, wherein
 10 the product has one or more of the following properties: number
 average molecular weight (M_n) in the range of 2,000 g/mol to 50,000
 g/mol, peak molecular weight (M_p) in the range of 1,500 g/mol to
 60,000 g/mol and the polydispersity index (PDI) is in the range of 1.0
 to 5.0, wherein the number average molecular weight, peak molecular

weight and polydispersity index are determined by gel permeation chromatography using polymethyl methacrylate (PMMA) calibration.

16. A method of preparing the reaction product according to any one of
5 claims 9 to 15, the method comprising:
 reacting one or more compounds according to any one of claims
 1 to 5 with one or more amine containing compounds to obtain the
 reaction product.
- 10 17. The method according to claim 16, wherein the amine containing
 compound comprises at least two amine functional groups.
18. The method according to claim 16, wherein the amine containing
 compound is selected from the group consisting of furan-2,5-
15 diyldimethanamine (FBA), xylene diamine (XDA), diaminopentane
 (DAP), hexamethylenediamine (HDA), ethylenediamine,
 diaminopropane, diaminobutane, ether diamine, polyether diamine,
 dimer diamine, lysine, isophorone diamine and phenylenediamine.
- 20 19. The method according to any one of claims 6 to 8 and 16 to 18, wherein
 the method is devoid of a step containing the use of isocyanates as a
 reactant.
20. The method according to any one of claims 16 to 18, further comprising
25 a step of functionalising one or more hydroxyl groups present in the
 reaction product.
21. The method according to any one of claims 16 to 18, further comprising
 a step of grafting a polymer to one or more hydroxyl groups present in
30 the reaction product.

22. The method according to any one of claims 16 to 18, further comprising a step of grafting one or more molecular entities or polymers to one or more furan rings present in the reaction product.
- 5 23. A functionalised or grafted product obtained according to any one of claims 20 to 22, wherein the functionalised or grafted polymer has one or more of the following properties: solubility or dispersibility in water, solubility or dispersibility in oil, photo or thermo or redox or pH response and crosslinking ability under air, photo, thermal, redox or ionic
10 conditions.