



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : G05B 1/03	A1	(11) International Publication Number: WO 99/59037 (43) International Publication Date: 18 November 1999 (18.11.99)
(21) International Application Number: PCT/US99/10027 (22) International Filing Date: 7 May 1999 (07.05.99) (30) Priority Data: 60/084,742 8 May 1998 (08.05.98) US 60/090,046 19 June 1998 (19.06.98) US (71) Applicant (for all designated States except US): ROSETTA INPHARMATICS, INC. [US/US]; 12040 115th Avenue, N.E., Kirkland, WA 98034 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): FRIEND, Stephen, H. [US/US]; 2653 Cascade Avenue S., Seattle, WA 98144 (US). STOUGHTON, Roland [US/US]; 425 West Spruce Street, San Diego, CA 92103 (US). (74) Agents: ANTLER, Adriane, M. et al.; Pennie & Edmonds LLP, 1155 Avenue of the Americas, New York, NY 10036 (US).	(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>	
(54) Title: METHODS OF DETERMINING PROTEIN ACTIVITY LEVELS USING GENE EXPRESSION PROFILES		
(57) Abstract		
The present invention provides methods for determining the level of protein activity in a cell by: (i) measuring abundances of cellular constituents in a cell in which the activity of a specific protein is to be determined so that a diagnostic profile is thus obtained; (ii) measuring abundances of cellular constituents that occur in a cell in response to perturbations in the activity of said protein to obtain response profiles and interpolating said response profiles to generate response curves; and (iii) determining a protein activity level at which the response profile extracted from the response curves best fits the measured diagnostic profile, according to some objective measure. In alternative embodiments, the present invention also provides methods for identifying individuals having genetic mutations or polymorphisms that disrupt protein activity, and methods for identifying drug activity <i>in vivo</i> by determining the activity levels of proteins which interact with said drugs.		