


**CHIRAL SYNTHESIS OF N-{3,4-DIFLUORO-2-[(2-FLUORO-4-
IODOPHENYL)AMINO]-6-METHOXYPHENYL}-1-[2,3-DIHYDROXY-
PROPYL]CYCLOPROPANESULFONAMIDES.**

5 FIELD OF THE INVENTION

The present invention relates to a novel enantioselective method of preparing (S)- and (R)-enantiomers of N-{3,4-difluoro-2-[(2-fluoro-4-iodophenyl)amino]-6-methoxyphenyl}-1-[2,3-dihydroxy-propyl]cyclopropanesulfonamide, to novel
10 intermediate compounds, and to the use of said novel intermediate compounds for the preparation of said (S)- and (R)-enantiomers of N-{3,4-difluoro-2-[(2-fluoro-4-iodophenyl)amino]-6-methoxyphenyl}-1-[2,3-dihydroxy-propyl]cyclopropanesulfonamide :



15

BACKGROUND TO THE INVENTION


N-{3,4-difluoro-2-[(2-fluoro-4-iodophenyl)amino]-6-methoxyphenyl}-1-[(2S)-
2,3-dihydroxy-propyl]cyclopropanesulfonamide (hereinafter referred to as
20 “(S)-14”, “BAY 86-9766” or “RDEA 119”) is a highly potent and selective MEK1/2 inhibitor currently under development in clinical trials for treatment

of late stage cancer patients refractory or intolerant to other anticancer therapies [ref. 1].

The initial synthesis of (S)-14, shown in Scheme A, *infra*, published in US 2008/0058340 [ref. 2] comprises an osmium catalyzed dihydroxylation of the allyl sulfonamide substituted core followed by chromatographic separation of the enantiomers using a chiral stationary phase : the initial synthesis of (S)-14 provides the target compound as a racemic mixture that needs to be separated by chiral chromatography [ref. 2].

10

Scheme A: Racemic synthesis of (S)-14 according to US 2008/0058340 :



NMO: 4-methylmorpholine 4-oxide monohydrate

In Scheme A, *supra*, a racemic mixture of the (S)- and (R)-enantiomers of N-{3,4-difluoro-2-[(2-fluoro-4-iodophenyl)amino]-6-methoxyphenyl}-1-[2,3-dihydroxy-propyl]cyclopropanesulfonamide is produced which, as seen, must be separated by chiral chromatography, in order to provide the single enantiomers. This separation of the enantiomers by chiral chromatography after the last step of the synthesis is a significant drawback in that in addition to the chiral separation step, more than double amounts of all intermediates have to be produced to obtain the same quantity of (S)-14 (RDEA 119).

As mentioned *supra*, another drawback of prior art synthesis of Scheme A is the use of very toxic osmium tetroxide which requires additional effort to remove the content of osmium to acceptable levels.

5


Another synthesis of (S)-14, shown in Scheme B, *infra*, published in PCT patent application under WO 2011/009541 A1 [ref. 7], describes a chiral preparation of (R)- and (S)-N-{3,4-difluoro-2-[(2-fluoro-4-iodophenyl)amino]-6-methoxyphenyl}-1-[2,3-dihydroxy-propyl]cyclopropanesulfonamide and

10

Scheme B, *infra*, illustrates the synthesis of (R)-N-(3,4-difluoro-2-(2-fluoro-4-iodophenylamino)-6-methoxyphenyl)-1-(2,3-dihydroxypropyl)cyclopropane-1-sulfonamide according to WO 2011/009541 A1 [ref. 7] :

15

SCHEME B



20

The synthesis as described in WO 2011/009541 A1 starts from commercially available glycidol which is converted via protection of the alcohol and coupling (Step A) with a protected acetylene followed by deprotection (Step B) and iodination with 9-I-BBN (Step C) to provide 4-iodopent-4-ene-1,2-diol, both HO
5 groups of which are protected (Step D). A cyclopropyl group is introduced across the alkene of diol-protected 4-iodopent-4-ene-1,2-diol (Step E) to form the protected 3-(1-iodocyclopropyl)propane-1,2-diol derivative, which is then deprotected (Step F) and protected again (Step G), before the iodo group of which is transformed into a sulfonyl chloride group in Step H.

10


It has been discovered, and this provides the basis of the present invention, that (S)-14 can be synthesised *via* a chiral synthesis of sodium 1-{[(4S)-2,2-dimethyl-1,3-dioxolan-4-yl]methyl}cyclopropanesulfonate (hereinafter referred to also as compound “(S)-7”) starting from (S)-epichlorohydrin (hereinafter
15 also referenced as compound “(S)-1”) and alternatively from enantiomerically pure glycidol derivatives, as illustrated in Schemes 1 and 2, *infra*.

20

The *R*-enantiomers have been prepared in the same manner.

Scheme 1, *infra*, represents a general illustration of the steps used in the chiral synthesis of sodium 1-{[(4S)-2,2-dimethyl-1,3-dioxolan-4-yl]methyl}cyclopropanesulfonate ((S)-7) according to the present invention :

SCHEME 1




It is seen that the steps in Scheme 1 relate to the enantioselective synthesis of (S)-7, using the (S)-enantiomers of all the intermediates involved in said synthesis : as is understood by the person skilled in the art, the enantioselective synthesis of (R)-7 is identical to the synthesis of (S)-7 as illustrated in Scheme 1, *supra*, except that it uses the (R)-enantiomers of all the intermediates instead of the (S)-enantiomers.

Starting from economically priced chiral epoxyp propane derivatives, chiral sodium 1-[[*(4S)*-2,2-dimethyl-1,3-dioxolan-4-yl]methyl]cyclopropanesulfonate ((*S*)-7) was prepared (Scheme 1, *supra*), which was converted to chiral 1-[[*(4S)*-2,2-dimethyl-1,3-dioxolan-4-yl]methyl]cyclopropane sulfonyl chloride
5 ((*S*)-10), which, in turn, was converted to the final product, N-{3,4-difluoro-2-[(2-fluoro-4-iodophenyl)amino]-6-methoxyphenyl}-1-[(2*S*)-2,3-dihydroxy-propyl]cyclopropanesulfonamide ((*S*)-14), according to Scheme 2, *infra*.

Scheme 2, *infra*, represents a general illustration of the steps used in the
10 chiral synthesis of N-{3,4-difluoro-2-[(2-fluoro-4-iodophenyl)amino]-6-methoxyphenyl}-1-[(2*S*)-2,3-dihydroxy-propyl]cyclopropanesulfonamide ((*S*)-14) from sodium 1-[[*(4S)*-2,2-dimethyl-1,3-dioxolan-4-yl]methyl]cyclopropanesulfonate ((*S*)-7) according to the present invention :

SCHEME 2



- 5 It is seen that the steps in Scheme 2 relate to the enantioselective synthesis of (S)-14, using the (S)-enantiomers of all the intermediates involved in said

synthesis : as is understood by the person skilled in the art, the enantioselective synthesis of (R)-14 is identical to the synthesis of (S)-14, except that it uses the (R)-enantiomers of all the intermediates instead of the (S)-enantiomers.

5

The synthesis of RDEA 119 as depicted in Schemes 1 and 2 of the present invention delivers compound 14 containing 79-85% of the desired enantiomer (S)-14 (via steps A, B, C), hence, only 15-21% of the undesired isomer needs removing via chromatography. The amounts of intermediates and waste as well as the production costs of (S)-14 are thus reduced significantly using the synthesis of the present invention. Further, the use of toxic osmium tetroxide is avoided.

10

Further, the synthesis of RDEA 119 as depicted in Schemes 1 and 2 of the present invention delivers compound 14 as the desired enantiomer (S)-14 (via steps E, F, G), alleviating the technical problems of the separation of the enantiomers by chiral chromatography, and of the use of toxic osmium tetroxide. The amounts of intermediates and waste as well as the production costs of (S)-14 are thus reduced significantly using the synthesis of the present invention.

20

The synthesis of the sulfonyl chloride (S-10) depicted in Schemes 1 and 2 according to the present invention involves 5 steps (Steps A, B, C, D and H, or Steps E, F, G, D and H), including one protection step (Step C, starting from (S)-1, or Step G, starting from (S)-8).

25

Further, the synthesis of the sulfonyl chloride (S-10) depicted in Schemes 1 and 2 according to the present invention proceeds with an overall yield of >60% from (S)-8 (Steps, E, F, G, D, H).

30

All reagents used in the synthesis of the sulfonyl chloride (S-10) depicted in Schemes 1 and 2 according to the present invention (phosphoryl chloride, dimethoxypropane, sodium methoxide) are cheap and available in bulk.

5 The synthesis of the sulfonyl chloride (S-10) depicted in Schemes 1 and 2 according to the present invention provides the chiral sodium 1-{[(4S)-2,2-dimethyl-1,3-dioxolan-4-yl]methyl}cyclopropanesulfonate (S)-7, as an intermediate, which is a solid which can be purified by crystallization.

10 As seen in the Experimental section, Example 9b, it was found that the reaction of 1-{[(4R)-2,2-dimethyl-1,3-dioxolan-4-yl]methyl}cyclopropanesulfonyl chloride (compound (R)-10) with 5,6-difluoro-N1-(4-fluoro-2-iodophenyl)-3-methoxybenzene-1,2-diamine (compound 12), in the presence of 4-dimethylaminopyridine in pyridine, provided, under the conditions
15 described, only a percentage yield of 38.9 % of N-{3,4-difluoro-2-[(2-fluoro-4-iodophenyl)amino]-6-methoxyphenyl}-1-{[(4R)-2,2-dimethyl-1,3-dioxolan-4-yl]methyl}cyclopropanesulfonamide (compound (R)-13). However, surprisingly, the bromide-promoted method of the present invention as depicted in Scheme 2, Steps J and K) provides the pure (S)-14 product in very good yield : as seen
20 specifically in the Experimental section, Example 9a, it was surprisingly found that the reaction of 1-{[(4S)-2,2-dimethyl-1,3-dioxolan-4-yl]methyl}cyclopropanesulfonyl chloride (compound (S)-10) with 5,6-difluoro-N1-(4-fluoro-2-iodophenyl)-3-methoxybenzene-1,2-diamine (compound 12), in the presence of tetrabutylammonium bromide in pyridine and sulfolane,
25 provided, under the conditions described, a percentage yield of 92.2 % of N-{3,4-difluoro-2-[(2-fluoro-4-iodophenyl)amino]-6-methoxyphenyl}-1-{[(4S)-2,2-dimethyl-1,3-dioxolan-4-yl]methyl}cyclopropanesulfonamide (compound (S)-13), thus representing an increase in percentage yield of product of over 50% when directly comparing the method in Example 9a with the method in
30 Example 9b. This is clearly advantageous from a chemical developmental point

of view, in view of the productivity of the method and the decrease in the amount of impurities.

Further, more importantly, as seen in the Experimental Section, Example 10a,
5 it was surprisingly found that :

- the reaction of 1-{[(4S)-2,2-dimethyl-1,3-dioxolan-4-yl]methyl}cyclopropanesulfonyl chloride (compound (S)-10) with 5,6-difluoro-N1-(4-fluoro-2-iodophenyl)-3-methoxybenzene-1,2-diamine
10 (compound 12), in the presence of tetrabutylammonium bromide in pyridine and sulfolane, then, after complete conversion to N-{3,4-difluoro-2-[(2-fluoro-4-iodophenyl)amino]-6-methoxyphenyl}-1-{[(4S)-2,2-dimethyl-1,3-dioxolan-4-yl]methyl}cyclopropanesulfonamide (compound (S)-13),
15 - followed by stirring the resulting reaction mixture with hydrochloric acid,

provided, under the conditions described, a percentage yield of 91.6 % of N-{3,4-difluoro-2-[(2-fluoro-4-iodophenyl)amino]-6-methoxyphenyl}-1-[(2S)-2,3-dihydroxy-propyl]cyclopropanesulfonamide (compound (S)-14) : this “one-pot”
20 method of Example 10a is clearly advantageous from a chemical developmental point of view, in terms of process efficiency.

Further, it was surprisingly found that the preparation of (S)-13 from (S)-10-Br and the one pot conversion to R-14 proceeds much cleaner and at a
25 temperature of 23°C, as compared to the preparation of (S)-13 from (S)-10 and the one pot conversion to R-14, which proceeds at a temperature of 70°C.

The method of preparation of (S)-14 from (S)-10 of the present invention, particularly the one-pot method of preparing (S)-14 from (S)-10, as described
30 and defined herein, is thus clearly surprisingly technically advantageous.

The following description provides further general details of each step shown in Schemes 1 and 2, *supra*, together with the technical advantages thereof.

I. In re. Scheme 1, Steps A, B, C, E, F, G, and D, *supra* :


5

Synthesis of Butyl 1-[(4S)-2,2-dimethyl-1,3-dioxolan-4-yl]methyl}cyclopropanesulfonates 6

Deprotonation of butyl cyclopropanesulfonate 2 with n-butyllithium and reaction with (S)-epichlorohydrin (S)-1 provided the chloroalcohol 3 in good yield (*vide* Scheme 1, Step A, *supra*, and Scheme 3, *infra*). Chloroalcohol 3 was converted quantitatively to the epoxide 4 with aqueous sodium hydroxide in tetrahydrofuran (*vide* Scheme 1, Step B, and Scheme 3, *infra*).

Scheme 3: Stereoselective synthesis of epoxides 4 starting from (S)-epichlorohydrin (S)-1 :


20



25

Performing the first step of the sequence without additives resulted in up to 65% by-product formation caused by competitive self alkylation (*vide* Scheme 4, *infra*). Several additives such as TMEDA (N,N,N',N'-Tetramethylethylenediamine), DMI (1,3-Dimethyl-2-imidazolidinone), DMPU (1,3-Dimethyl-3,4,5,6-tetrahydro-2(1H)-pyrimidinone), CuI, Lil were screened but showed no significant improvement : it was surprisingly found that 1 equivalent of boron trifluoride etherate led to a significant decrease in by-product formation :

Scheme 4: By-product formation




- 5 From reactions performed with magnesium bromide, 1-bromo-3-chloropropan-2-ol was isolated as the only product, no conversion was observed when triethylborane was used as additive.

Of all additives tested, boron trifluoride etherate was identified as the most
 10 promising. Good results have been obtained with 1.0 equivalent of boron trifluoride etherate whereas more equivalents resulted in formation of polymers.

The reaction of isopropyl cyclopropanesulfonate with (S)-2-(benzyloxymethyl)oxirane in the presence of 0.83 equivalents of hexamethyl
 15 phosphoric triamide (HMPT) was published in international patent application WO 2010/145197 A1 [ref. 3]. As HMPT exhibits carcinogenic and mutagenic properties, it was preferred to use boron trifluoride etherate instead of HMPT as additive for industrial preparation (*vide* Scheme 5, *infra*).

20

Scheme 5: Direct conversion of epoxides 4 to dioxolanes 6 (*vide* Scheme 1, Step C, *supra*) :




- a) BF_3 etherate, acetone: 58.2% ee
 b) phosphomolybdic acid hydrate, acetone: 69.8% ee

As shown with (**R**)-**4** direct conversion of the epoxide **4** with acetone catalyzed by boron trifluoride etherate or phosphomolybdic acid hydrate yielded dioxolane **6** as a mixture of enantiomers with 58.2 and 69.8% ee respectively
 5 (scheme 5).

Alternatively, chiral diol **5** was obtained, without racemization, from TBDMS- and THP-protected glycidols (**9a** and **9b**, respectively) and butyl cyclopropanesulfonate (**2**) under the same conditions optimized for the
 10 preparation of chloroalcohol **3** in good yield (*vide* Scheme 1, Step E, *supra*). Fortunately, TBDMS- and THP-protecting groups could be easily removed by acidic treatment of protected intermediates **9a** and **9b** providing enantiomerically pure diol **5** (*vide* Scheme 1, Step F, *supra*, and Scheme 6, *infra*).

15


Scheme 6: Chiral synthesis starting from protected glycidols



PPTS = pyridinium p-toluenesulfonate

Reactions performed without additives resulted in up to 65% by-product formation caused by competitive self alkylation (*vide* Scheme 7, *infra*). Therefore several additives like TMEDA, DMI, DMPU, CuI, Lil were screened but showed no significant improvement (*vide* Scheme 7, *infra*).

5

Scheme 7 : By-product formation

10 From reactions performed with magnesium bromide 1-bromo-3-chloro-propan-2-ol was isolated as sole product, no conversion was observed when triethylborane was used as additive. Of all additives tested boron trifluoride etherate was identified as most promising. Good results have been obtained with 1.0 equivalent of boron trifluoride etherate whereas more equivalents
15 resulted in formation of polymers.

The reaction of isopropyl cyclopropanesulfonate with (S)-2-(benzyloxymethyl)oxirane in the presence of 0.83 equivalents of hexamethyl phosphoric triamide (HMPT) was published [ref. 3]. As HMPT exhibits carcinogenic and mutagenic properties we preferred boron trifluoride etherate
20 to HMPT as additive for industrial preparation.

Chiral diol 5 was obtained without racemization from TBDMS- as well as THP-protected glycidols (8a and 8b, respectively) and butyl cyclopropanesulfonate 2.

25 Fortunately TBDMS- and THP-protecting groups could be easily removed by acidic treatment providing enantiomerically pure diol 5 (*vide* scheme 1, step F, *supra*).


Chiral diol **5** was converted subsequently to the desired dioxolane **6** with 2,2-dimethoxypropane catalyzed by pyridinium p-toluenesulfonate in good yield (*vide* Scheme 1, Step G, *supra*).

5 II. *In re*. Scheme 2, Steps H, J (and I), and K, *supra* :

Synthesis of (S)-**14** via sulfonyl chloride (S)-**10**

As seen *supra*, formation of sodium sulfonate **7** was achieved by using sodium methoxide instead of the considerably more expensive potassium thiocyanate
10 [refs. 1,2] (*vide* Scheme 1, Step D, *supra*, and Scheme 8, *infra*). Pure sodium sulfonate **7** was obtained from crystallization with ethanol or isopropanol. Reaction of **7** in pyridine with phosphoryl chloride [ref. 4] proved to be a more economical route to sulfonyl chloride **10** compared to cyanuric chloride [ref.
15 5] as chlorination reagent (*vide* Scheme 2, Step H, *supra*, and Scheme 8, *infra*). After aqueous work up, **10** was obtained in good yield without deterioration of the dioxolane protecting group. Reactions with oxalyl chloride or triphenylphosphine/sulfurylchloride [ref. 6] were less effective. Enantiomeric purity of **10** was assessed by chiral HPLC of the corresponding
20 anilides **11** (*vide* Scheme 2, Step I, *supra*, and Scheme 8, *infra*). The reaction of sulfonyl chloride **10** with aniline was accelerated by addition of lithium bromide.

Scheme 8 : Synthesis of chiral amides




- 5 Starting from sulfonyl chloride (S)-10 enantiomerically pure (S)-14 was prepared with tetrabutyl ammonium bromide as additive at 70°C in sulfolane/pyridine 2:1 followed by acidic deprotection of (S)-13 during aqueous work up (*vide* Scheme 2, Step K, *supra*, and Scheme 9, *infra*). The amide formation was accelerated by addition of tetrabutyl ammonium bromide
- 10 (*vide* Scheme 9, *infra*).


Sulfonyl chloride (R)-10 furnished the enantiomer (R)-14 accordingly (*vide* Scheme 9, *infra*).

The enantiomeric purity of 14 was determined by chiral HPLC.

15 Scheme 9 : Stereoselective synthesis of (S)-14



Hence, in a first aspect, the present invention relates to a method of preparing N-{3,4-difluoro-2-[(2-fluoro-4-iodophenyl)amino]-6-methoxyphenyl}-1-[(2S)-2,3-dihydroxy-propyl]cyclopropanesulfonamide of formula (S)-14 :




5


(S)-14,

via the following steps shown in Schemes 1 and 2, *infra* :


SCHEME 1



SCHEME 2



In an embodiment of the first aspect, the present invention relates to a method of preparing a compound of formula (S)-14 :



5


(S)-14 ;

comprising :

i) the following step J :


wherein a compound of formula (S)-10 :

10



(S)-10

is allowed to react with a compound of formula (12) :




15

(12)

in the presence of a bromide, such as tetra-n-butylammonium bromide or lithium bromide for example, optionally in the presence of a base, such as pyridine for example, optionally in a solvent, such as sulfolane for example,

thereby providing a reaction mixture containing a compound of formula (S)-13 :



5

and, then,

ii) the following step K :

10 wherein an aqueous mineral acid, such as hydrochloric acid for example, is added to said reaction mixture containing a compound of formula (S)-13,


thereby providing a compound of formula (S)-14.

15

Alternatively, in the above-mentioned step J, a catalyst, such as 4-dimethylaminopyridine for example, can be used instead of said bromide.

Alternatively, in scheme 2, step J, the compound of formula (S)-10-Br :


20



may be used instead of the compound (S)-10.

Hence, in an embodiment of the first aspect, the present invention relates to a method of preparing a compound of formula (S)-14 :

5




(S)-14 ;

comprising :


10 i) the following step J :

wherein a compound of formula (S)-10-Br :



(S)-10-Br

15 is allowed to react with a compound of formula (12) :




(12)

optionally in the presence of a base, such as pyridine for example, optionally in a solvent, such as sulfolane for example,

thereby providing a reaction mixture containing a compound of formula (S)-

5 13 :



and, then,


10 ii) the following step K :

wherein an aqueous mineral acid, such as hydrochloric acid for example, is added to said reaction mixture containing a compound of formula (S)-13,

15 thereby providing a compound of formula (S)-14.

In another embodiment of the first aspect, the present invention relates to a method of preparing a compound of formula (S)-14 :

24




(S)-14 ;

comprising the following step K :

wherein an aqueous mineral acid, such as hydrochloric acid for example, is

5 added to a compound of formula (S)-13 :




((S)-13)

;

thereby providing a compound of formula (S)-14.

10 In a further embodiment of the first aspect of the present invention, the above-mentioned compound of formula (S)-13 :


25




is prepared by the following step J :

wherein a compound of formula (S)-10 :

5



is allowed to react with a compound of formula (12) :




10

in the presence of a bromide, such as tetra-n-butylammonium bromide or lithium bromide for example, optionally in the presence of a base, such as pyridine for example, optionally in a solvent, such as sulfolane for example,

15 thereby providing a reaction mixture containing a compound of formula (S)-13 :


26



Alternatively, in the above-mentioned step J, a catalyst, such as 4-dimethylaminopyridine for example, can be used instead of said bromide.

5


In a further embodiment of the first aspect of the present invention, the above-mentioned compound of formula (S)-13 :



10


is prepared by the following step J :

wherein a compound of formula (S)-10-Br :




15

is allowed to react with a compound of formula (12) :




optionally in the presence of a base, such as pyridine for example, optionally
5 in a solvent, such as sulfolane for example,

thereby providing a reaction mixture containing a compound of formula (S)-
13 :



10


In a further embodiment of the first aspect of the present invention, the
above-mentioned compound of formula (S)-10 :



15

is prepared by allowing a compound of formula (S)-7 :

28




(S)-7

to react with a chlorinating agent, such as $\text{P}(\text{O})\text{Cl}_3$, PCl_3 or SOCl_2 for example, optionally in a solvent, such as pyridine for example, thereby providing a compound of formula (S)-10.

5


In a further embodiment of the first aspect of the present invention, the above-mentioned compound of formula (S)-10-Br :



(S)-10-Br

10

is prepared by allowing a compound of formula (S)-7 :




(S)-7

to react with a brominating agent, such as $\text{P}(\text{O})\text{Br}_3$ or PBr_3 for example, optionally in a solvent, such as pyridine for example, thereby providing a compound of formula (S)-10-Br.

15


In a further embodiment of the first aspect of the present invention, the above-mentioned compound of formula (S)-7 :

29



(S)-7


is prepared by allowing a compound of formula (S)-6 :



(S)-6


- 5 to react with a sodium alkoxide, such as sodium methoxide for example, optionally in a solvent, such as methanol for example, thereby providing a compound of formula (S)-7.

In a further embodiment of the first aspect, the present invention relates to a
10 method of preparing a compound of formula (S)-10 :



(S)-10


wherein a compound of formula (S)-7' :




(S)-7'

- 15 is allowed to react with a chlorinating agent, such as P(O)Cl₃, PCl₃ or SOCl₂ for example, optionally in a solvent, such as pyridine for example, thereby providing a compound of formula (S)-10.

In a further embodiment of the first aspect of the present invention, the above-mentioned compound of formula (S)-7' :



5 is prepared by allowing a compound of formula (S)-6 :




to react with a potassium alkoxide, such as potassium methoxide for example, optionally in a solvent, such as methanol for example, thereby providing a compound of formula (S)-7'.

10


A further embodiment of the first aspect of the present invention relates to a method of preparing N-{3,4-difluoro-2-[(2-fluoro-4-iodophenyl)amino]-6-methoxyphenyl}-1-[(2S)-2,3-dihydroxy-propyl]cyclopropanesulfonamide of formula (S)-14 :

15



comprising the following step D :


wherein a compound of formula (S)-6 :



(S)-6


is allowed to react with a sodium alkoxide, such as sodium methoxide for example, optionally in a solvent, such as methanol for example,

5 thereby providing a compound of formula (S)-7 :



(S)-7


A further embodiment of the first aspect of the present invention relates to a
 10 method of preparing N-{3,4-difluoro-2-[(2-fluoro-4-iodophenyl)amino]-6-methoxyphenyl}-1-[(2S)-2,3-dihydroxy-propyl]cyclopropanesulfonamide of formula (S)-14 :



(S)-14 ;

15 comprising the following step D :
 wherein a compound of formula (S)-6 :


32



(S)-6

is allowed to react with a potassium alkoxide, such as potassium methoxide for example, optionally in a solvent, such as methanol for example, thereby providing a compound of formula (S)-7' :


5



(S)-7'


In a further embodiment of the first aspect of the present invention, the above-mentioned compound of formula (S)-6 :

10



(S)-6

is prepared by allowing a compound of formula (S)-4 :



(S)-4

15

to react with either :


a) boron trifluoride, optionally in the form of a complex, such as a boron trifluoride etherate complex for example, such as a boron trifluoride

tetrahydrofuran complex, or a boron trifluoride diethylether complex for example, optionally in a solvent such as acetone for example, or

b) phosphomolybdic acid hydrate, optionally in a solvent such as acetone for example,

5 thereby providing a compound of formula (S)-6.


In a further embodiment of the first aspect of the present invention, the above-mentioned compound of formula (S)-4 :



(S)-4

10

is prepared by allowing a compound of formula (S)-3 :




(S)-3 ;

15

to react with a base, such as sodium hydroxide for example, optionally in a solvent, such as water for example, thereby providing a compound of formula (S)-4.


20 In a further embodiment of the first aspect of the present invention, the above-mentioned compound of formula (S)-3 :



(S)-3 ;

25 is prepared by :

a) allowing a compound of formula 2 :



to react with a base, such as n-butyllithium for example, optionally in a solvent, such as tetrahydrofuran for example,

5

b) adding a compound of formula (S)-1 and boron trifluoride, optionally in the form of a complex, such as a boron trifluoride etherate complex for example, such as a boron trifluoride tetrahydrofuran complex, or a boron trifluoride diethylether complex for example ; and

10

c) allowing the thus-formed deprotonated compound of formula 2 to react with a compound of formula (S)-1 :




15

(S)-1 ;

thereby providing a compound of formula (S)-3.

In a further embodiment of the first aspect of the present invention, the above-mentioned compound of formula (S)-6 :


20



(S)-6

is prepared by allowing a compound of formula (S)-5 :

35




(S)-5

to react with 2,2-dimethoxypropane, optionally in a solvent, such as tetrahydrofuran for example and optionally in the presence of a catalyst, such as pyridinium p-toluenesulfonate for example,
 5 thereby providing a compound of formula (S)-6.

In a further embodiment of the first aspect of the present invention, the above-mentioned compound of formula (S)-5 :

10



(S)-5

is prepared by allowing a compound of formula (S)-9 :



15


(S)-9 ;

wherein PG represents a protecting group, such as a t-butyldimethylsilyl-group (compound (S)-9a) or a tetrahydropyranyl group (compound (S)-9b), for example,
 20 to react with an acid, such as hydrochloric acid or p-toluene sulfonic acid for example, optionally in a solvent, such as methanol for example, thereby providing a compound of formula (S)-5.

In a further embodiment of the first aspect of the present invention, the above-mentioned compound of formula (S)-9 :

25

36




(S)-9 ;

wherein PG represents a t-butyldimethylsilyl- group (compound (S)-9a) or a tetrahydropyranyl group (compound (S)-9b),

5 is prepared by :

a) allowing a compound of formula 2 :



(2)


to react with a base, such as n-butyllithium for example, optionally in a solvent, such as tetrahydrofuran for example,

10

b) adding a compound of formula (S)-8 and boron trifluoride, optionally in the form of a complex, such as a boron trifluoride etherate complex for example, such as a boron trifluoride tetrahydrofuran complex, or a boron trifluoride diethylether complex for example ; and

15

c) allowing the thus-formed deprotonated compound of formula 2 to react with a compound of formula (S)-8 :



(S)-8,

20 wherein PG represents a t-butyldimethylsilyl- group (compound (S)-8a) or a tetrahydropyranyl group (compound (S)-8b), respectively, thereby providing a compound of formula (S)-9.

In a further embodiment of the first aspect, the present invention relates to a
25 method of preparing the compound of formula (S)-14, wherein each of said


steps A to K as shown in Schemes 1 and 2, *supra*, are described in relation to Schemes 3 to 9, *supra*.

In accordance with a second aspect, the present invention relates to intermediate compounds which are useful in the preparation of the above-mentioned compound of formula (S)-14.

In an embodiment of said second aspect, the present invention relates to a compound :

10

1-{[(4S)-2,2-dimethyl-1,3-dioxolan-4-yl]methyl}cyclopropanesulfonyl bromide (S)-10-Br :




15

(S)-10-Br.

In an embodiment of said second aspect, the present invention relates to a compound :

20 potassium 1-{[(4S)-2,2-dimethyl-1,3-dioxolan-4-yl]methyl}cyclopropane-sulfonate (S)-7' :




(S)-7' .

In an embodiment of said second aspect, the present invention relates to a compound :

25

Butyl 1-[[*(4S)*-2,2-dimethyl-1,3-dioxolan-4-yl]methyl]cyclo propane sulfonate
(*S*)-6 :




5

(*S*)-6 .


In an embodiment of said second aspect, the present invention relates to a compound :

10 Butyl 1-[(*2S*)-2,3-dihydroxypropyl]cyclopropanesulfonate (*S*)-5 :

(*S*)-5 .

15 In an embodiment of said second aspect, the present invention relates to a compound :

Butyl 1-[(*2S*)-oxiran-2-ylmethyl]cyclopropanesulfonate (*S*)-4 :


(*S*)-4 .

20

In an embodiment of said second aspect, the present invention relates to a compound :

Butyl 1-[(*2S*)-3-chloro-2-hydroxypropyl]cyclopropanesulfonate (*S*)-3 :


39



(S)-3 .

In an embodiment of said second aspect, the present invention relates to a
5 compound :

Butyl 1-[(2S)-3-[[tert-butyl(dimethyl)silyl]oxy}-2-hydroxypropyl] cyclopropane-
sulfonate (S)-9a :




10

(S)-9a .

In an embodiment of said second aspect, the present invention relates to a
compound :

15 Butyl 1-[(2S)-2-hydroxy-3-(tetrahydro-2H-pyran-2-yloxy)propyl]cyclopropane
sulfonate (S)-9b :




(S)-9b.

20 In a third aspect, the present invention relates to the use of the intermediate
compounds according to the second aspect, *supra*, for preparing the compound
of formula (S)-14, *supra*.

In an embodiment of said third aspect, the present invention relates to the use of :

1-{[(4S)-2,2-dimethyl-1,3-dioxolan-4-yl]methyl}cyclopropanesulfonyl bromide

5 (S)-10-Br :




(S)-10-Br

10 for preparing the compound of formula (S)-14, *supra*.

In an embodiment of said third aspect, the present invention relates to the use of :

15 potassium 1-{[(4S)-2,2-dimethyl-1,3-dioxolan-4-yl]methyl}cyclopropane-sulfonate (S)-7' :



(S)-7' ;


for preparing the compound of formula (S)-14, *supra*.

20

In an embodiment of said third aspect, the present invention relates to the use of :

Butyl 1-{[(4S)-2,2-dimethyl-1,3-dioxolan-4-yl]methyl}cyclo propane sulfonate

25 (S)-6 :




(S)-6 ;

for preparing the compound of formula (S)-14, *supra*.

5 In an embodiment of said third aspect, the present invention relates to the use of :

Butyl 1-[(2S)-2,3-dihydroxypropyl]cyclopropanesulfonate (S)-5 :




10

(S)-5 ;

for preparing the compound of formula (S)-14, *supra*.

15 In an embodiment of said third aspect, the present invention relates to the use of :

Butyl 1-[(2S)-oxiran-2-ylmethyl]cyclopropanesulfonate (S)-4 :



20


(S)-4 ;

for preparing the compound of formula (S)-14, *supra*.

In an embodiment of said third aspect, the present invention relates to the use of :

25

Butyl 1-[(2S)-3-chloro-2-hydroxypropyl]cyclopropanesulfonate (S)-3 :




(S)-3 ;

- 5 for preparing the compound of formula (S)-14, *supra*.

In an embodiment of said third aspect, the present invention relates to the use of :

- 10 Butyl 1-[(2S)-3-{[tert-butyl(dimethyl)silyl]oxy}-2-hydroxypropyl] cyclopropane-sulfonate (S)-9a :




(S)-9a ;

- 15 for preparing the compound of formula (S)-14, *supra*.

In an embodiment of said third aspect, the present invention relates to the use of :


- 20 Butyl 1-[(2S)-2-hydroxy-3-(tetrahydro-2H-pyran-2-yloxy)propyl]cyclopropane sulfonate (S)-9b :



(S)-9b ;

for preparing the compound of formula (S)-14, *supra*.


In a fourth aspect, the present invention relates to a method of preparing N-
{3,4-difluoro-2-[(2-fluoro-4-iodophenyl)amino]-6-methoxyphenyl}-1-[(2R)-2,3-
5 dihydroxy-propyl]cyclopropanesulfonamide of formula (R)-14 :




(R)-14,

via the following steps shown in Schemes C and D, *infra* :

SCHEME C




SCHEME D



In a further embodiment of the fourth aspect, the present invention relates to a method of preparing the compound of formula (R)-14, wherein each of said steps A to K as shown in Schemes C and D, *supra*, are described in analogy to the preparation of (S)-14 in Schemes A and B, and Schemes 3 to 9, *supra*.

5

In an embodiment of the fourth aspect, the present invention relates to a method of preparing a compound of formula (R)-14 :



10


(R)-14 ;

comprising :

i) the following step J :

wherein a compound of formula (R)-10 :


15



(R)-10

is allowed to react with a compound of formula (12) :


47



in the presence of a bromide, such as tetra-n-butylammonium bromide or lithium bromide for example, optionally in the presence of a base, such as pyridine for example, optionally in a solvent, such as sulfolane for example,

5

thereby providing a reaction mixture containing a compound of formula (R)-13 :



10 and, then,

ii) the following step K :


wherein an aqueous mineral acid, such as hydrochloric acid for example, is
 15 added to said reaction mixture containing a compound of formula (R)-13,

thereby providing a compound of formula (R)-14.

Alternatively, in the above-mentioned step J, a catalyst, such as 4-dimethylaminopyridine for example, can be used instead of said bromide.

Alternatively, in scheme D, step J, the compound of formula (R)-10-Br :


5



(R)-10-Br

may be used instead of the compound (R)-10.

- 10 Hence, in an embodiment of the fourth aspect, the present invention relates to a method of preparing a compound of formula (R)-14 :




(R)-14 ;

- 15 comprising :

i) the following step J :


wherein a compound of formula (R)-10-Br :

49



(R)-10-Br

is allowed to react with a compound of formula (12) :




(12)

5

optionally in the presence of a base, such as pyridine for example, optionally in a solvent, such as sulfolane for example,

thereby providing a reaction mixture containing a compound of formula (R)-

10 13 :



((R)-13)

;

and, then,


15 ii) the following step K :

wherein an aqueous mineral acid, such as hydrochloric acid for example, is added to said reaction mixture containing a compound of formula (R)-13,

thereby providing a compound of formula (R)-14.

5

In another embodiment of the fourth aspect, the present invention relates to a method of preparing a compound of formula (R)-14 :




10

(R)-14 ;

comprising the following step K :

wherein an aqueous mineral acid, such as hydrochloric acid for example, is added to a compound of formula (R)-13 :




((R)-13)

;

15

thereby providing a compound of formula (R)-14.


In a further embodiment of the fourth aspect of the present invention, the above-mentioned compound of formula (R)-13 :



5


is prepared by the following step J :

wherein a compound of formula (R)-10 :



10


is allowed to react with a compound of formula (12) :



in the presence of a bromide, such as tetra-n-butylammonium bromide or lithium bromide for example, optionally in the presence of a base, such as pyridine for example, optionally in a solvent, such as sulfolane for example,

15


thereby providing a reaction mixture containing a compound of formula (R)-13 :



- 5 Alternatively, in the above-mentioned step J, a catalyst, such as 4-dimethylaminopyridine for example, can be used instead of said bromide.

In a further embodiment of the fourth aspect of the present invention, the above-mentioned compound of formula (R)-13 :

10




is prepared by the following step J :

wherein a compound of formula (R)-10-Br :


15

53



(R)-10-Br

is allowed to react with a compound of formula (12) :




(12)

5

optionally in the presence of a base, such as pyridine for example, optionally in a solvent, such as sulfolane for example,

thereby providing a reaction mixture containing a compound of formula (R)-

10 13 :




((R)-13)

In a further embodiment of the fourth aspect of the present invention, the above-mentioned compound of formula (R)-10 :


15

54



(R)-10

is prepared by allowing a compound of formula (R)-7 :




(R)-7

to react with a chlorinating agent, such as $\text{P}(\text{O})\text{Cl}_3$, PCl_3 or SOCl_2 for example,
 5 optionally in a solvent, such as pyridine for example, thereby providing a compound of formula (R)-10.


In a further embodiment of the fourth aspect of the present invention, the above-mentioned compound of formula (R)-10-Br :

10



(R)-10-Br

is prepared by allowing a compound of formula (R)-7 :




(R)-7

15


to react with a brominating agent, such as $\text{P}(\text{O})\text{Br}_3$ or PBr_3 for example,
 optionally in a solvent, such as pyridine for example, thereby providing a compound of formula (R)-10-Br.

In a further embodiment of the fourth aspect of the present invention, the above-mentioned compound of formula (R)-7 :



5


is prepared by allowing a compound of formula (R)-6 :



to react with a sodium alkoxide, such as sodium methoxide for example, optionally in a solvent, such as methanol for example, thereby providing a compound of formula (R)-7.


10

In a further embodiment of the fourth aspect, the present invention relates to a method of preparing a compound of formula (R)-10 :




15

wherein a compound of formula (R)-7' :




is allowed to react with a chlorinating agent, such as $P(O)Cl_3$, PCl_3 or $SOCl_2$ for example, optionally in a solvent, such as pyridine for example, thereby providing a compound of formula (R)-10.

- 5 In a further embodiment of the fourth aspect of the present invention, the above-mentioned compound of formula (R)-7' :



(R)-7'

is prepared by allowing a compound of formula (R)-6 :




(R)-6

10

to react with a potassium alkoxide, such as potassium methoxide for example, optionally in a solvent, such as methanol for example, thereby providing a compound of formula (R)-7'.

- 15 A further embodiment of the fourth aspect of the present invention relates to a method of preparing N-{3,4-difluoro-2-[(2-fluoro-4-iodophenyl)amino]-6-methoxyphenyl}-1-[(2R)-2,3-dihydroxy-propyl]cyclopropanesulfonamide of formula (R)-14 :

57




(R)-14 ;

comprising the following step D :

wherein a compound of formula (R)-6 :


5



(R)-6

is allowed to react with a sodium alkoxide, such as sodium methoxide for example, optionally in a solvent, such as methanol for example, thereby providing a compound of formula (R)-7 :

10




(R)-7

A further embodiment of the fourth aspect of the present invention relates to a method of preparing N-{3,4-difluoro-2-[(2-fluoro-4-iodophenyl)amino]-6-methoxyphenyl}-1-[(2R)-2,3-dihydroxy-propyl]cyclopropanesulfonamide of formula (R)-14 :

15

58




(R)-14 ;

comprising the following step D :

wherein a compound of formula (R)-6 :


5



(R)-6

is allowed to react with a potassium alkoxide, such as potassium methoxide for example, optionally in a solvent, such as methanol for example, thereby providing a compound of formula (R)-7' :

10




(R)-7'


In a further embodiment of the fourth aspect of the present invention, the above-mentioned compound of formula (R)-6 :

15

59



is prepared by allowing a compound of formula (R)-4 :




5

to react with either :


- a) boron trifluoride, optionally in the form of a complex, such as a boron trifluoride etherate complex for example, such as a boron trifluoride tetrahydrofuran complex, or a boron trifluoride diethylether complex for example, optionally in a solvent such as acetone for example, or
 - b) phosphomolybdic acid hydrate, optionally in a solvent such as acetone for example,
- thereby providing a compound of formula (R)-6.

- 15 In a further embodiment of the fourth aspect of the present invention, the above-mentioned compound of formula (R)-4 :



is prepared by allowing a compound of formula (R)-3 :


20



to react with a base, such as sodium hydroxide for example, optionally in a solvent, such as water for example, thereby providing a compound of formula (R)-4.

5

In a further embodiment of the fourth aspect of the present invention, the above-mentioned compound of formula (R)-3 :




10

(R)-3 ;

is prepared by :

a) allowing a compound of formula 2 :



(2)

15 to react with a base, such as n-butyllithium for example, optionally in a solvent, such as tetrahydrofuran for example,

b) adding a compound of formula (R)-1 and boron trifluoride, optionally in the form of a complex, such as a boron trifluoride etherate complex for example, 20 such as a boron trifluoride tetrahydrofuran complex, or a boron trifluoride diethylether complex for example ; and

c) allowing the thus-formed deprotonated compound of formula 2 to react with a compound of formula (R)-1 :

25




61

(R)-1 ;


thereby providing a compound of formula (R)-3.

In a further embodiment of the fourth aspect of the present invention, the
 5 above-mentioned compound of formula (R)-6 :



(R)-6

10 is prepared by allowing a compound of formula (R)-5 :




(R)-5

to react with 2,2-dimethoxypropane, optionally in a solvent, such as tetrahydrofuran for example and optionally in the presence of a catalyst, such
 15 as pyridinium p-toluenesulfonate for example,
 thereby providing a compound of formula (R)-6.

In a further embodiment of the fourth aspect of the present invention, the
 above-mentioned compound of formula (R)-5 :


20



(R)-5

is prepared by allowing a compound of formula (R)-9 :

62




(R)-9 ;

wherein PG represents a protecting group, such as a t-butyldimethylsilyl-group (compound (R)-9a) or a tetrahydropyranyl group (compound (R)-9b), for example,

to react with an acid, such as hydrochloric acid or p-toluene sulfonic acid for example, optionally in a solvent, such as methanol for example, thereby providing a compound of formula (R)-5.


In a further embodiment of the fourth aspect of the present invention, the above-mentioned compound of formula (R)-9 :



(R)-9 ;

wherein PG represents a t-butyldimethylsilyl- group (compound (R)-9a) or a tetrahydropyranyl group (compound (R)-9b), is prepared by :

a) allowing a compound of formula 2 :




(2)

to react with a base, such as n-butyllithium for example, optionally in a solvent, such as tetrahydrofuran for example,

b) adding a compound of formula (R)-8 and boron trifluoride, optionally in the form of a complex, such as a boron trifluoride etherate complex for example, such as a boron trifluoride tetrahydrofuran complex, or a boron trifluoride diethylether complex for example ; and

c) allowing the thus-formed deprotonated compound of formula 2 to react with a compound of formula (R)-8 :



5

(R)-8,


wherein PG represents a t-butyldimethylsilyl- group (compound (R)-8a) or a tetrahydropyranyl group (compound (R)-8b), respectively, thereby providing a compound of formula (R)-9.

10 In a further embodiment of the fourth aspect, the present invention relates to a method of preparing the compound of formula (R)-14, wherein each of said steps A to K as shown in Schemes C and D, *supra*, are described in relation to Schemes 3 to 9, *supra*.

15 In accordance with a fifth aspect, the present invention relates to intermediate compounds which are useful in the preparation of the above-mentioned compound of formula (R)-14.

In an embodiment of said fifth aspect, the present invention relates to a
20 compound :

1-{[(4*R*)-2,2-dimethyl-1,3-dioxolan-4-yl]methyl}cyclopropanesulfonyl bromide
(R)-10-Br :




25

(R)-10-Br .

In an embodiment of said fifth aspect, the present invention relates to a compound :


potassium 1-{[(4R)-2,2-dimethyl-1,3-dioxolan-4-yl]methyl}cyclopropane-
5 sulfonate (R)-7' :



In an embodiment of said fifth aspect, the present invention relates to a compound :

10


Butyl 1-{[(4R)-2,2-dimethyl-1,3-dioxolan-4-yl]methyl}cyclo propane sulfonate
(R)-6 :



15

In an embodiment of said fifth aspect, the present invention relates to a compound :


Butyl 1-[(2R)-2,3-dihydroxypropyl]cyclopropanesulfonate (R)-5 :



20

In an embodiment of said fifth aspect, the present invention relates to a compound :


Butyl 1-[(2R)-oxiran-2-ylmethyl]cyclopropanesulfonate (R)-4 :



5

In an embodiment of said fifth aspect, the present invention relates to a compound :

Butyl 1-[(2R)-3-chloro-2-hydroxypropyl]cyclopropanesulfonate (R)-3 :




10

In an embodiment of said fifth aspect, the present invention relates to a compound :

15

Butyl 1-[(2R)-3-{[tert-butyl(dimethyl)silyl]oxy}-2-hydroxypropyl] cyclopropane-sulfonate (R)-9a :




20

In an embodiment of said fifth aspect, the present invention relates to a compound :

Butyl 1-[(2R)-2-hydroxy-3-(tetrahydro-2H-pyran-2-yloxy)propyl]cyclopropane sulfonate (R)-9b :

25

66




(R)-9b.

In a sixth aspect, the present invention relates to the use of the intermediate
 5 compounds according to the fifth aspect, *supra*, for preparing the compound
 of formula (R)-14, *supra*.

In an embodiment of said sixth aspect, the present invention relates to the use
 of :

10

1-{[(4R)-2,2-dimethyl-1,3-dioxolan-4-yl]methyl}cyclopropanesulfonyl bromide
 (R)-10-Br :



15


(R)-10-Br

for preparing the compound of formula (R)-14, *supra*.

In an embodiment of said sixth aspect, the present invention relates to the use
 20 of :

potassium 1-{[(4R)-2,2-dimethyl-1,3-dioxolan-4-yl]methyl}cyclopropane-
 sulfonate (R)-7' :


67



for preparing the compound of formula (R)-14, *supra*.

- 5 In an embodiment of said sixth aspect, the present invention relates to the use of :

Butyl 1-[(4R)-2,2-dimethyl-1,3-dioxolan-4-yl]methyl}cyclo propane sulfonate (R)-6 :




10

for preparing the compound of formula (R)-14, *supra*.

- 15 In an embodiment of said sixth aspect, the present invention relates to the use of :

Butyl 1-[(2R)-2,3-dihydroxypropyl]cyclopropanesulfonate (R)-5 :




20

for preparing the compound of formula (R)-14, *supra*.

- In an embodiment of said sixth aspect, the present invention relates to the use of :

Butyl 1-[(2R)-oxiran-2-ylmethyl]cyclopropanesulfonate (R)-4 :



(R)-4 ;


5

for preparing the compound of formula (R)-14, *supra*.

In an embodiment of said sixth aspect, the present invention relates to the use of :

10

Butyl 1-[(2R)-3-chloro-2-hydroxypropyl]cyclopropanesulfonate (R)-3 :



(R)-3 ;


15

for preparing the compound of formula (R)-14, *supra*.

In an embodiment of said sixth aspect, the present invention relates to the use of :

20

Butyl 1-[(2R)-3-{[tert-butyl(dimethyl)silyl]oxy}-2-hydroxypropyl] cyclopropane-sulfonate (R)-9a :



(R)-9a ;


25

for preparing the compound of formula (R)-14, *supra*.

In an embodiment of said sixth aspect, the present invention relates to the use of :

5

Butyl 1-[(2R)-2-hydroxy-3-(tetrahydro-2H-pyran-2-yloxy)propyl]cyclopropane sulfonate (R)-9b :



(R)-9b ;

10

for preparing the compound of formula (R)-14, *supra*.

Within the context of the present invention the term “solvent”, as optionally present in any reaction step of the method of the invention, is understood, as is by the person skilled in the art, as meaning any substance in which other materials dissolve to form a solution, such as, without being limited to : a polar solvent, such as a polar protic solvent, such as water, n-butanol, isopropanol, n-propanol, ethanol, methanol, or formic acid or acetic acid, etc., for example ; a polar aprotic solvent, such as 1,4-dioxane, tetrahydrofuran, 1,2-dimethoxyethane, acetone, acetonitrile, dimethylformamide, sulfolane, pyridine or dimethylsulphoxide, etc., for example ; or a non-polar solvents, such as pentane, hexane, benzene, toluene, diethyl ether, methyl ethyl ketone, dichloromethane, chloroform, tetrachloromethane, ethyl acetate, etc., for example ; or any mixture of the solvents listed above.

25

It is understood that any combination of the definitions given in the above-mentioned embodiments is possible within the context of the present invention.

The invention will be better understood upon reading the Examples below, which are provided as an illustration of the present invention. The Examples below in no way whatsoever constitute a limitation of the present invention as
5 described in the present text and as defined in the claims appended hereto.

EXPERIMENTAL SECTION

EXAMPLES

10 Abbreviations used :

The following abbreviations used in the Examples have the following meanings :

	TMEDA	N,N,N',N'-Tetramethylethylenediamine
15	DMI	1,3-Dimethyl-2-imidazolidinone
	DMPU	1,3-Dimethyl-3,4,5,6-tetrahydro-2(1H)-pyrimidinone
	conc.	concentrated
	h	hour / hours
	THF	tetrahydrofuran
20	¹ H-NMR	proton nuclear magnetic resonance spectroscopy : chemical shifts (δ) are given in ppm.
	s	singlet
	bs	broad singlet
	d	doublet
25	dd	doublet of doublets
	t	triplet

General

30 Butyl cyclopropanesulfonate 2 and 5,6-difluoro-N1-(4-fluoro-2-iodophenyl)-3-methoxybenzene-1,2-diamine can be prepared as described in [ref. 2, Example

12, Step B, and Example 50, Step B, respectively]. All other reagents are commercially available.

Example 1 : General Procedure for the synthesis of substituted Butyl 1-[2-hydroxypropyl]cyclopropanesulfonates (GP1) :

A solution of butyl cyclopropanesulfonate 2 (1.2 g, 6.7 mmol) in THF (24 mL) was cooled to -78°C and n-butyllithium (1.6M in n-hexane, 5.0 mL, 8.1 mmol) was added within 5 min. The mixture was stirred for 30 min and epoxypropane derivative 1 or 8 (8.1 mmol) was added followed by addition of boron trifluoride diethyl etherate (0.84 mL, 6.7 mmol). The solution was stirred for 1 h at -78°C, the cooling bath was removed and stirring was continued for 2 additional hours. For aqueous work-up diluted hydrochloric acid (10%, 0.8 mL), water (10 mL) and ethyl acetate (20 mL) were added. The organic layer was concentrated at 40°C to afford the corresponding butyl 1-[2-hydroxypropyl]cyclopropanesulfonate 3.

According to **GP1** the following compounds were prepared :

Example 1a : Butyl 1-[(2S)-3-chloro-2-hydroxypropyl]cyclopropanesulfonate (S)-3




(S)-3 was obtained from (S)-epichlorohydrin (S)-1 in 81.3% yield.

R_f: 0.19 (silica gel, n-hexane/ethyl acetate 3:1)

MS (DCI): m/z = 288.0 [M+NH₄]⁺(100), 271 [M+H]⁺(14)


¹H NMR (400 MHz, Chloroform-*d*) δ = 0.90 - 1.04 (m, 4 H) 1.15 - 1.23 (m, 1 H) 1.39 - 1.57 (m, 4 H) 1.67 - 1.80 (m, 2 H) 1.90 (dd, J=15.28, 8.68 Hz, 1 H) 2.15 (dd, J=15.41, 3.67 Hz, 1 H) 2.5 - 3.0 (br. s, 2H) 3.51 - 3.73 (m, 2 H) 4.13 - 4.34 (m, 3 H) ppm

Example 1b : Butyl 1-[(2*R*)-3-chloro-2-hydroxypropyl]cyclopropanesulfonate (*R*)-3



5 (*R*)-3 was obtained from (*R*)-epichlorohydrin (*R*)-1 in 98.7% yield. Analytical data see (*S*)-3.

Example 1c : Butyl 1-[3-chloro-2-hydroxypropyl]cyclopropanesulfonate *rac*-3




10

rac-3 was obtained from epichlorohydrin in 72.5% yield. Analytical data see (*S*)-3

Example 1d : Butyl 1-[(2*S*)-3-{[*tert*-butyl(dimethyl)silyl]oxy}-2-hydroxypropyl]cyclopropanesulfonate (*S*)-9a

15



(*S*)-9a was prepared from *tert*-butyl(dimethyl)[(2*S*)-oxiran-2-ylmethoxy]silane in 100% yield (crude product).


20 R_f : 0.43 (silica gel, n-hexane/ethyl acetate 3:1)

GC-MS: m/z = 384.1 [$M+NH_4$]⁺(100), 367.0 [$M+H$]⁺(8)

The crude product was converted without further purification to (*S*)-5

Example 1e : Butyl 1-[(2*R*)-3-{[*tert*-butyl(dimethyl)silyl]oxy}-2-hydroxypropyl]cyclopropanesulfonate (*R*)-9a

25



(*R*)-**9a** was prepared from tert-butyl(dimethyl)[(2*R*)-oxiran-2-ylmethoxy]silane in 100% yield (crude product).


5 GC-MS: 70% (*R*)-**9a**, 7% (*R*)-**5**

R_f : 0.43 (silica gel, n-hexane/ethyl acetate 3:1)

MS (DCI): m/z = 384.0 $[M+NH_4]^+$ (34), 367.1 $[M+H]^+$ (38), 270.1 $[M_{(R)-2}+NH_4]^+$ (100)

The crude product was converted without further purification to (*R*)-**5**

10 **Example 1f : Butyl 1-[(2*R*)-2-hydroxy-3-(tetrahydro-2H-pyran-2-yloxy)propyl]cyclopropanesulfonate (*R*)-**9b****



15 (*R*)-**9b** was prepared from 2-[(2*R*)-oxiran-2-ylmethoxy]tetrahydro-2H-pyran in 100% yield (crude product) and converted without purification to (*R*)-**5**

R_f : 0.29 (silica gel, n-hexane/ethyl acetate 3:1)

MS (DCI): m/z = 354.0 $[M+NH_4]^+$ (5), 337.1 $[M+H]^+$ (4), 270.1 $[M_{(R)-2}+NH_4]^+$ (100), 253.1 $[M_{(R)-2}+H]^+$ (15)


20

Example 2 : General Procedure for the Synthesis of Butyl 1-[oxiran-2-ylmethyl]cyclopropane-sulfonates 4 (GP2) :

To a solution of **3** (22.1 mmol) in tetrahydrofuran (29 mL) was added a solution
 25 of sodium hydroxide (33.2 mmol) in water (22 mL) and the mixture was stirred overnight at room temperature to complete conversion (TLC silica gel, toluene/ethyl acetate 5:1). The reaction mixture was washed with ethyl

acetate (30 mL), the organic layer was concentrated and the crude product purified by chromatography (silica gel, toluene/ethyl acetate 5:1). According to GP2 the following compounds were prepared:

5 **Example 2a : Butyl 1-[(2S)-oxiran-2-ylmethyl]cyclopropanesulfonate (S)-4**




(S)-4 was obtained from (S)-3 in 85.0% yield

R_f: 0.40 (silica gel, toluene/ethyl acetate 5:1)

MS (DCI): m/z = 252.1 [M+NH₄]⁺(100), 235.0 [M+H]⁺(10),


10 ¹H NMR (400 MHz, Chloroform-*d*) δ = 0.89 - 1.06 (m, 4 H) 1.14 - 1.23 (m, 1 H)
1.37 - 1.55 (m, 4 H) 1.61 - 1.79 (m, 3 H) 2.40 (m, *J*=15.28, 4.52 Hz, 1 H) 2.48
(dd, *J*=4.89, 2.69 Hz, 1 H) 2.76 - 2.85 (m, 1 H) 3.14 - 3.23 (m, 1 H) 4.25 (td,
J=6.48, 0.98 Hz, 2 H) ppm.

15 **Example 2b : Butyl 1-[(2R)-oxiran-2-ylmethyl]cyclopropanesulfonate (R)-4**



(R)-4 was obtained from (R)-3 in 86.3% yield. Analytical data see (S)-4

Example 2c : Butyl 1-[oxiran-2-ylmethyl]cyclopropanesulfonate rac-4



20

rac-4 was obtained from *rac*-3 in 92.5% yield. Analytical data see (S)-4


25

Example 3 : General Procedure for the Synthesis of Butyl 1-[2,3-dihydroxypropyl]cyclopropane-sulfonates 5 by Acidic Deprotection of 9a (GP3) :

- 5 To **9a** (6.7 mmol) was added hydrochloric acid (10%, 4 ml) and methanol (24.5 mL) and it was stirred until TLC (silica gel, hexanes/ethyl acetate 3:1) indicated complete conversion (1h). For aqueous work-up ethyl acetate (50 mL) was added and it was washed with a saturated aq. solution of sodium hydrogen carbonate (2 x 20 mL). The organic layer was concentrated and the
10 residual oil was purified by chromatography (silica gel, 1. hexanes/ ethyl acetate 3:1, 2. ethyl acetate).

According to **GP3** the following compounds were prepared:


- 15 **Example 3a : Butyl 1-[(2S)-2,3-dihydroxypropyl]cyclopropanesulfonate (S)-5**



(S)-5 was obtained from **(S)-9a** in 97.1% yield

- 20 R_f : 0.36 (silica gel, ethyl acetate);
MS (DCI): m/z = 270.1 $[M+NH_4]^+$ (100), 253.1 $[M+H]^+$ (17)
 1H NMR (400 MHz, Chloroform- d) δ = 0.90 - 1.03 (m, 4 H) 1.12 - 1.19 (m, 1 H)
1.38 - 1.54 (m, 4 H) 1.68 - 1.79 (m, 2 H) 1.84 (dd, J =15.41, 9.29 Hz, 1 H) 2.00
(dd, J =15.41, 3.42 Hz, 1 H) 2.55 (m, 1 H) 3.23 (d, J =3.91 Hz, 1 H) 3.47 (dt,
25 J =11.37, 5.81 Hz, 1 H) 3.60 - 3.72 (m, 1 H) 4.11 (dt, J =6.11, 2.81 Hz, 1 H) 4.25
(td, J =6.60, 1.47 Hz, 2 H) ppm

Example 3b : Butyl 1-[(2*R*)-2,3-dihydroxypropyl]cyclopropanesulfonate (*R*)-5




5 (*R*)-5 was obtained from (*R*)-9a in 88.2 yield.

R_f : 0.36 (silica gel, ethyl acetate);

MS (DCI): m/z = 270.1 [$M+NH_4$]⁺(100), 253.1 [$M+H$]⁺(17)

¹H NMR (400 MHz, Chloroform-*d*) δ = 0.90 - 1.03 (m, 4 H) 1.12 - 1.19 (m, 1 H)
 1.38 - 1.54 (m, 4 H) 1.68 - 1.79 (m, 2 H) 1.84 (dd, J =15.41, 9.29 Hz, 1 H) 2.00
 10 (dd, J =15.41, 3.42 Hz, 1 H) 2.55 (m, 1 H) 3.23 (d, J =3.91 Hz, 1 H) 3.47 (dt,
 J =11.37, 5.81 Hz, 1 H) 3.60 - 3.72 (m, 1 H) 4.11 (dt, J =6.11, 2.81 Hz, 1 H) 4.25
 (td, J =6.60, 1.47 Hz, 2 H) ppm

Example 3c : Preparation of Butyl 1-[(2*R*)-2,3-dihydroxypropyl]cyclopropanesulfonate (*R*)-5 by Hydrolysis of (*R*)-9b



Crude (*R*)-9b (10 mmol) was dissolved in methanol (38 mL), *p*-toluene sulfonic acid (30 mg) was added and stirring was continued for 5.5 h until the reaction was completed (TLC, silica gel, ethyl acetate). The mixture was concentrated and the residue purified by chromatography (silica gel, ethyl acetate) to yield
 20 a colorless oil.

Yield: 1.1 g, 40.4 %

R_f : 0.36 (silica gel, ethyl acetate);

MS (DCI): m/z = 270.1 [$M+NH_4$]⁺(100), 253.1 [$M+H$]⁺(17)

25 ¹H NMR (400 MHz, Chloroform-*d*) δ = 0.90 - 1.03 (m, 4 H) 1.12 - 1.19 (m, 1 H)
 1.38 - 1.54 (m, 4 H) 1.68 - 1.79 (m, 2 H) 1.84 (dd, J =15.41, 9.29 Hz, 1 H) 2.00
 (dd, J =15.41, 3.42 Hz, 1 H) 2.55 (m, 1 H) 3.23 (d, J =3.91 Hz, 1 H) 3.47 (dt,


$J=11.37$, 5.81 Hz, 1 H) 3.60 - 3.72 (m, 1 H) 4.11 (dt, $J=6.11$, 2.81 Hz, 1 H) 4.25 (td, $J=6.60$, 1.47 Hz, 2 H) ppm

Example 4 : General Procedure for the Synthesis of Butyl 1-[[2,2-dimethyl-1,3-dioxolan-4-yl]methyl]-cyclopropanesulfonates 6 from diols 5

To a solution of **5** (0.6 g, 4.0 mmol) in THF (2.4 mL) was added 2,2-dimethoxypropane (1.5 mL, 11.9 mmol) and pyridinium *p*-toluenesulfonate (29.9 mg, 0.1 mmol). It was stirred until complete conversion overnight (TLC, silica gel, ethyl acetate). Ethyl acetate (20 mL) and a saturated aq. solution of sodium hydrogen carbonate (10 mL) were added; the organic layer was washed with brine (10 mL), dried over magnesium sulfate and concentrated at 40°C. The residue was purified by chromatography (silica gel, hexane/ethyl acetate 3:1).

15

Example 4a : Butyl 1-[[*(4S)*-2,2-dimethyl-1,3-dioxolan-4-yl]methyl]cyclopropanesulfonate (*S*)-6



20

(S)-6 was obtained from **(S)-5** in 84.1% yield.

R_f : 0.40 (silica gel, n-hexane/ethyl acetate 3:1)


MS (DCI): m/z = 310.1 [$M+NH_4$]⁺(27), 293.0 [$M+H$]⁺(100), 277.0 (6), 235.0 (14), 196.0 (3)

¹H NMR (500 MHz, Chloroform-*d*) δ = 0.83 - 0.96 (m, 4 H) 1.12 - 1.20 (m, 1 H) 1.27 (s, 3 H) 1.31 (s, 3 H) 1.33 - 1.44 (m, 4 H) 1.62 - 1.70 (m, 2 H) 1.75 (dd, $J=15.11$, 8.32 Hz, 1 H) 2.27 (dd, $J=15.15$, 4.31 Hz, 1 H) 3.46 (dd, $J=8.05$, 6.98

25

Hz, 1 H) 4.05 (dd, $J=8.16$, 6.10 Hz, 1 H) 4.12 - 4.22 (m, 2 H) 4.28 - 4.39 (m, 1 H) ppm

Example 4b : Butyl 1-[[*(4R)*-2,2-dimethyl-1,3-dioxolan-4-yl]methyl]cyclopropanesulfonate (*R*)-6



(*R*)-6 was obtained from (*R*)-5 in 74.8% yield.

10 R_f : 0.40 (silica gel, n-hexane/ethyl acetate 3:1)


MS (DCI): m/z = 310.1 [$M+NH_4$]⁺(27), 293.0 [$M+H$]⁺(100), 277.0 (6), 235.0 (14), 196.0 (3)

¹H NMR (500 MHz, Chloroform-*d*) δ = 0.83 - 0.96 (m, 4 H) 1.12 - 1.20 (m, 1 H) 1.27 (s, 3 H) 1.31 (s, 3 H) 1.33 - 1.44 (m, 4 H) 1.62 - 1.70 (m, 2 H) 1.75 (dd, $J=15.11$, 8.32 Hz, 1 H) 2.27 (dd, $J=15.15$, 4.31 Hz, 1 H) 3.46 (dd, $J=8.05$, 6.98 Hz, 1 H) 4.05 (dd, $J=8.16$, 6.10 Hz, 1 H) 4.12 - 4.22 (m, 2 H) 4.28 - 4.39 (m, 1 H) ppm

Example 5 : General Procedure for the Synthesis of Butyl 1-[[2,2-dimethyl-1,3-dioxolan-4-yl]methyl]-cyclopropanesulfonates 6 directly from epoxides 4 (GP4)


According to GP4 the following compounds were prepared:

Example 5a i) : Synthesis of Butyl 1-[[*(4R)*-2,2-dimethyl-1,3-dioxolan-4-yl]methyl]cyclopropanesulfonate (*(R)*-6 from Epoxide (*(R)*-4



To a solution of (*(R)*-4 (12.5 g, 53.3 mmol) in 115 mL acetone was added boron trifluoride diethyl etherate (1.8 g, 12.8 mmol). The solution was stirred for 2 h
5 at room temperature. Ethyl acetate (235 mL) and sat. aq. sodium carbonate (235 mL) were added, the organic layer was separated and concentrated in vacuum. The residue was dissolved in ethyl acetate (250 mL) and washed twice with water (50 mL each). The organic layer was concentrated in vacuum.
10 Yield: 15.3 g, 98.1%
Optical purity: 58.2% ee (determined by chiral HPLC of the corresponding anilide (*(R)*-11)

Example 5a ii) : Synthesis of Butyl 1-[[*(4R)*-2,2-dimethyl-1,3-dioxolan-4-yl]methyl]cyclopropanesulfonate (*(R)*-6 from (*(R)*-4



To a solution of (*(R)*-4 (0.6 g g 2.5 mmol) in 5 mL acetone was added phosphomolybdic acid hydrate (50 mg, 0.03 mmol). It was stirred for 2 h at
20 room temperature until TLC (silica gel, ethyl acetate) indicated complete conversion. The crude product was purified by chromatography (silica gel, ethyl acetate) to furnish an oil.
Yield: 0.65 g, 86.8%
Optical purity: 69.8% ee (determined by chiral HPLC of the corresponding anilide (*(R)*-11)


Example 6 : General Procedure for the Synthesis of Sodium 1-[[2,2-dimethyl-1,3-dioxolan-4-yl]methyl]-cyclopropanesulfonates 7 (GP5)

To a solution of **6** (0.5 g, 1.7 mmol) in 1,2-dimethoxyethane (3.3 mL) was
5 added a solution of sodium methoxide (30% in methanol, 0.49 mL, 2.6 mmol).
It was stirred under reflux temperature until complete conversion (TLC, silica
gel, ethyl acetate) (1.5 h) and concentrated at 45°C to yield a solid. The solid
was stirred with hexanes (5 mL), isolated by filtration, crystallized from
ethanol (3.5 mL per g) or isopropanol (8.0 mL per g) under reflux temperature
10 and dried in vacuo at 45°C.

According to **GP5** the following compounds were prepared :

Example 6a : Sodium 1-[(4S)-2,2-dimethyl-1,3-dioxolan-4-yl]methyl}cyclopropanesulfonate (S)-7

15



(S)-7 was obtained from **(S)-6** in quantitative yield.

20 MS: m/z = 281.0 [M+Na]⁺


¹H NMR (400 MHz, DMSO-*d*₆) δ = 0.36 - 0.50 (m, 2 H) 0.78 - 0.90 (m, 2 H) 1.23
(s, 3 H) 1.26 (s, 3 H) 1.69 (dd, *J*=14.18, 7.58 Hz, 1 H) 1.91 - 2.01 (m, 1 H) 3.48
(t, *J*=7.70 Hz, 1 H) 4.02 (dd, *J*=7.82, 5.87 Hz, 1 H) 4.44 (s, 1 H) ppm

Optical purity: >99.9 % ee

25 (obtained from **(S)-8** via steps E, F, G, D (cf. Scheme 1 *supra*)

(determined by chiral HPLC of the corresponding anilide **(S)-11**)

Example 6b : Sodium 1-[(4*R*)-2,2-dimethyl-1,3-dioxolan-4-yl]methyl}cyclopropanesulfonate (*R*)-7



5

(*R*)-7 was obtained from (*R*)-6 in quantitative yield.

MS: $m/z = 281.0 [M+Na]^+$

1H NMR (400 MHz, DMSO- d_6) $\delta = 0.36 - 0.50$ (m, 2 H) $0.78 - 0.90$ (m, 2 H) 1.23 (s, 3 H) 1.26 (s, 3 H) 1.69 (dd, $J=14.18, 7.58$ Hz, 1 H) $1.91 - 2.01$ (m, 1 H) 3.48 (t, $J=7.70$ Hz, 1 H) 4.02 (dd, $J=7.82, 5.87$ Hz, 1 H) 4.44 (s, 1 H) ppm

10

Optical purity: 99.4 % ee (*R*)-7 obtained from (*R*)-8a

99.7 % ee (*R*)-7 obtained from (*R*)-8b

(determined by chiral HPLC of the corresponding anilide (*R*)-11)

15


Example 7 : General Procedures for the Synthesis of 1-[[2,2-dimethyl-1,3-dioxolan-4-yl]methyl}cyclopropanesulfonyl chlorides 10 (GP6a and GP6b)

GP6a: To a suspension of **7** (0.50 g, 1.9 mmol) in 5 mL pyridine was added under cooling phosphoric trichloride ($P(O)Cl_3$) (0.30 g, 1.9 mmol) and the mixture was stirred overnight at room temperature. The suspension was added to water (10 mL) and toluene (15 mL). The aqueous layer was separated and washed with toluene (10 mL). The combined organic layers were concentrated at $45^\circ C$. Two portions of toluene (5 mL each) were added and stripped off to remove any traces of hydrochloric acid.

25

According to **GP6a** the following compounds were prepared :

Example 7a : 1-[[*(4S)*-2,2-dimethyl-1,3-dioxolan-4-yl]methyl]cyclopropanesulfonyl chloride (*S*)-10



5 (*S*)-10 was obtained from (*S*)-7 in 96.0% yield.


R_f: 0.53 (silica gel, n-hexane/ethyl acetate 1:1)

MS (DCI): *m/z* = 255.1 [*M*+H]⁺(100), 223.1 (40), 196.2 (10), 148.2 (4)

¹H NMR (400 MHz, Chloroform-*d*) δ = 1.29 - 1.37 (m, 4 H) 1.39 (s, 3 H) 1.56 - 1.65 (m, 1 H) 1.70 - 1.87 (m, 2 H) 2.06 (dd, *J*=15.41, 9.05 Hz, 1 H) 2.56 - 2.67 (m, 1 H) 3.59 (dd, *J*=7.83, 7.09 Hz, 1 H) 4.15 (dd, *J*=8.07, 6.11 Hz, 1 H) 4.40 (d, *J*=6.11 Hz, 1 H) ppm

Example 7b : 1-[[*(4R)*-2,2-dimethyl-1,3-dioxolan-4-yl]methyl]cyclopropanesulfonyl chloride (*R*)-10

15



(*R*)-10 was obtained from (*R*)-7 in 86.2% yield.

R_f: 0.53 (silica gel, n-hexane/ethyl acetate 1:1)

20 MS (DCI): *m/z* = 255.1 [*M*+H]⁺(100), 223.1 (40), 196.2 (10), 148.2 (4)

¹H NMR (400 MHz, Chloroform-*d*) δ = 1.29 - 1.37 (m, 4 H) 1.39 (s, 3 H) 1.56 - 1.65 (m, 1 H) 1.70 - 1.87 (m, 2 H) 2.06 (dd, *J*=15.41, 9.05 Hz, 1 H) 2.56 - 2.67 (m, 1 H) 3.59 (dd, *J*=7.83, 7.09 Hz, 1 H) 4.15 (dd, *J*=8.07, 6.11 Hz, 1 H) 4.40 (d, *J*=6.11 Hz, 1 H) ppm


25

GP6b: To a suspension of **7** (5.0 g, 19.0 mmol) in 36 mL pyridine was added at 60°C a solution of thionyl chloride (SOCl₂) (13.8 g, 116.0 mmol) in toluene (38 mL) over 10 min. The mixture was stirred for 30 min at 60°C. After cooling to room temperature, toluene (75 mL) was added and the reaction suspension
5 was added to cold water (60 mL). The organic layer was separated and washed twice with cold water (60 mL each). The combined organic layers were concentrated at 50°C.

According to **GP6b** the following compound was prepared :

10

Example 7c : 1-{[(4S)-2,2-dimethyl-1,3-dioxolan-4-yl]methyl}cyclopropanesulfonyl chloride (S)-10



15 **(S)-10** was obtained from **(S)-7** in 95.1% yield.


Example 7-Br : General Procedure for the Synthesis of 1-{[2,2-dimethyl-1,3-dioxolan-4-yl]methyl}cyclopropanesulfonyl bromides 10-Br (GP6-Br)

20

To a suspension of **7** (5.00 g, 19.4 mmol) in 50 mL pyridine was added under cooling phosphoric tribromide (P(O)Br₃) (8.33 g, 29.0 mmol) and the mixture was stirred for 1h at room temperature. The suspension was added to water (100 mL) and toluene (150 mL). The aqueous layer was separated and washed
25 with toluene (100 mL). The combined organic layers were concentrated at 45°C. Two portions of toluene (50 mL each) were added and stripped off to remove any traces of hydrochloric acid.

According to **GP6-Br** the following compounds were prepared :

Example 7d.1 : 1-{[(4*S*)-2,2-dimethyl-1,3-dioxolan-4-yl]methyl}cyclopropanesulfonyl bromide (*S*)-10-Br



5

(*S*)-10-Br was obtained from (*S*)-7 in 89.8% yield.


R_f: 0.53 (silica gel, n-hexane/ethyl acetate 2:1)

MS (DCI): m/z = 317.9, 315.9 [M+NH₄]⁺(20), 301.0, 299.0 [M+H]⁺(100), 196.2
10 (18), 136.1 (46), 130.2 (16), 102.2 (70)

¹H NMR (400 MHz, Chloroform-*d*) δ = 1.31 - 1.43 (m, 1 H) 1.34 (s, 3 H) 1.40 (s, 3 H) 1.61 - 1.69 (m, 1 H) 1.69 - 1.77 (m, 1 H) 1.84 (m, 1 H) 2.04 (dd, *J*=15.41, 9.05 Hz, 1 H) 2.57 - 2.64 (m, 1 H) 3.60 (dd, *J*=8.19, 6.72 Hz, 1 H) 4.16 (dd, *J*=8.31, 6.11 Hz, 1 H) 4.39 (dd, *J*=6.24, 3.06 Hz, 1 H) ppm

15

Example 7e : 1-{[(4*R*)-2,2-dimethyl-1,3-dioxolan-4-yl]methyl}cyclopropanesulfonyl bromide (*R*)-10-Br



20

(*R*)-10-Br was obtained according to **GP6-Br** from (*R*)-7 in 86.9% yield.

R_f: 0.55 (silica gel, n-hexane/ethyl acetate 2:1)

MS (DCI): m/z = 318.0, 316.0 [M+NH₄]⁺(25), 301.0, 299.0 [M+H]⁺(100), 238.2
25 (20), 221.1 (34), 196.1 (70), 180.1 (20)

¹H NMR (400 MHz, Chloroform-*d*) δ = 1.31 - 1.43 (m, 1 H) 1.34 (s, 3 H) 1.40 (s, 3 H) 1.61 - 1.69 (m, 1 H) 1.69 - 1.77 (m, 1 H) 1.84 (m, 1 H) 2.04 (dd, *J*=15.41, 9.05 Hz, 1 H) 2.57 - 2.64 (m, 1 H) 3.60 (dd, *J*=8.19, 6.72 Hz, 1 H) 4.16 (dd, *J*=8.31, 6.11 Hz, 1 H) 4.39 (dd, *J*=6.24, 3.06 Hz, 1 H) ppm

Example 8 : General Procedure for the Synthesis of 1-[[2,2-dimethyl-1,3-dioxolan-4-yl]methyl]cyclopropanesulfonamides 11 (GP7)


5 To **10** (0.67 mmol) in dichloromethane (0.35 mL) were added at room temperature lithium bromide (57.9 mg, 0.67 mmol), pyridine (270 μ L, 3.3 mmol) and aniline (54.7 μ L, 0.6 mmol). The suspension was stirred overnight, dichloromethane (5 mL) was added, the organic layer was washed three times with hydrochloric acid (2.5%, 17 mL each), dried over magnesium sulfate and
10 concentrated to an orange oil.

The enantiomeric purity of **6** was determined by chiral HPLC (CHIRALPAK IB, 5 μ m column length 25 cm, internal diameter 4.6 mm; temperature of the column oven 35°C; eluent n-heptane/ethanol 95:5 v/v; flow rate 1.0 mL/min; detection wavelength 228 nm; retention time (*R*)-**11** approx. 10.6 min, (*S*)-**11**
15 approx. 13.4 min.

According to **GP7** the following compounds were prepared:

Example 8a : 1-[[2,2-dimethyl-1,3-dioxolan-4-yl]methyl]-N-phenylcyclopropanesulfonamide (*S*)-11

20



(*S*)-**11** was obtained from (*S*)-**10** in 70.0% yield.


R_f: 0.48 (silica gel, n-hexane/ethyl acetate 1:1)

25 MS (DCI): m/z = 329.0 [M+NH₄]⁺(67), 312.2 [M+H]⁺(100), 254.1 (35), 223.1 (5).

¹H NMR (400 MHz, DMSO-*d*₆) δ = 0.80 - 0.92 (m, 1 H) 0.92 - 1.01 (m, 1 H) 1.07 (s, 2 H) 1.16 - 1.24 (m, 3 H) 1.26 (s, 3 H) 1.89 - 2.02 (m, 1 H) 2.02 - 2.18 (m, 1

H) 3.41 (s, 1 H) 3.96 (s, 1 H) 4.15 - 4.31 (m, 1 H) 7.09 (s, 1 H) 7.18 - 7.27 (m, 1 H) 7.29 (d, $J=7.34$ Hz, 2 H) 9.80 (br. s, 1 H) ppm

Example 8b : 1-[[*(4R)*-2,2-dimethyl-1,3-dioxolan-4-yl]methyl]-*N*-phenylcyclopropanesulfonamide (*R*)-11



(*R*)-11 was obtained from (*R*)-10 in 81.9% yield.


10 R_f: 0.48 (silica gel, n-hexane/ethyl acetate 1:1)

MS (DCI): m/z = 329.0 [$M+NH_4$]⁺(67), 312.2 [$M+H$]⁺(100), 254.1 (35), 223.1 (5).

¹H NMR (400 MHz, DMSO-*d*₆) δ = 0.80 - 0.92 (m, 1 H) 0.92 - 1.01 (m, 1 H) 1.07 (s, 2 H) 1.16 - 1.24 (m, 3 H) 1.26 (s, 3 H) 1.89 - 2.02 (m, 1 H) 2.02 - 2.18 (m, 1 H) 3.41 (s, 1 H) 3.96 (s, 1 H) 4.15 - 4.31 (m, 1 H) 7.09 (s, 1 H) 7.18 - 7.27 (m, 1 H) 7.29 (d, $J=7.34$ Hz, 2 H) 9.80 (br. s, 1 H) ppm

Example 9a : Synthesis of *N*-{3,4-difluoro-2-[(2-fluoro-4-iodophenyl)amino]-6-methoxyphenyl}-1-[[*(4S)*-2,2-dimethyl-1,3-dioxolan-4-yl]methyl]cyclopropanesulfonamide (*S*)-13

20



To a mixture of 5,6-difluoro-N1-(4-fluoro-2-iodophenyl)-3-methoxybenzene-1,2-diamine [ref. 2, Example 50, Step B] (0.9 g, 2.3 mmol) and tetrabutylammonium bromide (0.74 g, 2.3 mmol) in pyridine (0.76 mL) and sulfolane (1.5 mL) was added (**S**)-**10** (0.8 g, 3.0 mmol). It was stirred at 70°C
5 for 18h. It was diluted with ethyl acetate (20 mL) and hydrochloric acid (10%, 5 mL). The organic layer was separated, washed with sat. aq. sodium hydrogen carbonate (20 mL), and with water (10 mL). The organic layer was dried over magnesium sulfate and concentrated to a solid. The solid was purified by chromatography (silica gel, hexane/ethyl acetate 1:1).

10 Yield: 1.3 g, 92.2%

R_f = 0.46 (silica gel hexanes/ethyl acetate 1:1)

MS (DCI): m/z = 613.3 [$M+H$]⁺(100), 555.2 (30), 459.1 (20), 393.0 (45) 267.1 (10), 219.0 (5)

¹H NMR (400 MHz, DMSO-*d*₆) δ = 0.82 - 1.02 (m, 4 H) 1.24 (s, 3 H) 1.29 (s, 3 H)
15 2.07 (dd, J =15.28, 7.70 Hz, 1 H) 2.35 (dd, J =15.28, 4.52 Hz, 1 H) 3.46 (t, J =7.70 Hz, 1 H) 3.83 (s, 3 H) 4.03 (m, J =7.34 Hz, 2 H) 4.19 - 4.29 (m, 1 H) 6.48 (td, J =8.80, 4.65 Hz, 1 H) 7.07 (dd, J =12.59, 7.21 Hz, 1 H) 7.33 (d, J =8.56 Hz, 1 H) 7.36 - 7.41 (br. s, 1 H) 7.56 (dd, J =10.76, 1.71 Hz, 1 H) 9.14 (s, 1 H) ppm


20 (**S**)-**13** was obtained from (**S**)-**10**-Br in 100.0% yield

To a mixture of 5,6-difluoro-N1-(4-fluoro-2-iodophenyl)-3-methoxybenzene-1,2-diamine [ref. 2, Example 50, Step B] (0.9 g, 2.3 mmol) in pyridine (0.76 mL) and sulfolane (1.5 mL) was added (**S**)-**10**-Br (0.94 g, 3.0 mmol). It was
25 stirred at 23°C for 18h. It was diluted with ethyl acetate (20 mL) and hydrochloric acid (10%, 5 mL). The organic layer was separated, washed with sat. aq. sodium hydrogen carbonate (20 mL), and with water (10 mL). The organic layer was dried over magnesium sulfate and concentrated to a solid. The solid was purified by chromatography (silica gel, hexane/ethyl acetate
30 1:1).

Yield: 1.4 g, 100.0%

Example 9b : Synthesis of N-{3,4-difluoro-2-[(2-fluoro-4-iodophenyl)amino]-6-methoxyphenyl}-1-[[(4*R*)-2,2-dimethyl-1,3-dioxolan-4-yl]methyl}cyclopropanesulfonamide (*R*)-13

5



A mixture of (*R*)-10 (400 mg, 1.57 mmol), 5,6-difluoro-N1-(4-fluoro-2-iodophenyl)-3-methoxybenzene-1,2-diamine (618 mg, 1.57 mmol) and
 10 4-dimethyl aminopyridine (38 mg, 0.31 mmol) in pyridine (5 mL) was stirred at 50°C for 45h. The solvent was stripped off in vacuum and the residue purified by chromatography (silica gel 1. dichloromethane, 2. methanol) to give 820 mg of crude product.

R_f = 0.28 (silica gel, dichloromethane)

15 The crude product was purified by a second chromatography (silica gel, hexanes/ethyl acetate 1:1).


Yield: 350 mg 38.9%

R_f = 0.46 (silica gel, hexanes/ethyl acetate 1:1)

MS (DCI): m/z = 613.3 [$M+H$]⁺(100), 555.2 (30), 459.1 (20), 393.0 (45) 267.1
 20 (10), 219.0 (5)

¹H NMR (400 MHz, DMSO-*d*₆) δ = 0.82 - 1.02 (m, 4 H) 1.24 (s, 3 H) 1.29 (s, 3 H) 2.07 (dd, J =15.28, 7.70 Hz, 1 H) 2.35 (dd, J =15.28, 4.52 Hz, 1 H) 3.46 (t, J =7.70 Hz, 1 H) 3.83 (s, 3 H) 4.03 (m, J =7.34 Hz, 2 H) 4.19 - 4.29 (m, 1 H) 6.48 (td, J =8.80, 4.65 Hz, 1 H) 7.07 (dd, J =12.59, 7.21 Hz, 1 H) 7.33 (d, J =8.56 Hz, 1 H) 7.36 - 7.41 (br. s, 1 H) 7.56 (dd, J =10.76, 1.71 Hz, 1 H) 9.14 (s, 1 H) ppm
 25

Example 10a.1 : Synthesis of N-{3,4-difluoro-2-[(2-fluoro-4-iodophenyl)amino]-6-methoxyphenyl}-1-[(2S)-2,3-dihydroxypropyl]cyclopanesulfonamide (S)-14



5

To a mixture of 5,6-difluoro-N1-(4-fluoro-2-iodophenyl)-3-methoxybenzene-1,2-diamine [ref. 2, Example 50, Step B] (0.9 g, 2.3 mmol) and tetrabutylammonium bromide (TBAB) (0.74 g, 2.3 mmol) in pyridine (0.76 mL) and sulfolane (1.5 mL) was added (S)-10 (0.8 g, 3.0 mmol). It was stirred at 70 °C for 18h to achieve complete conversion to (S)-13. The reaction mixture was stirred with hydrochloric acid (10%, 4 mL) for 1h. It was diluted with ethyl acetate (20 mL), the organic layer was washed two times with water (30 mL, 2x 15 mL), dried over magnesium sulfate and concentrated to a solid. The solid was purified by chromatography (silica gel, ethyl acetate).

15

Yield: 1.2 g, 91.6%

R_f: 0.12 (silica gel, ethyl acetate)

MS (DCI): m/z 590.1 [M+NH₄]⁺(40), 573.0 [M+H]⁺(28), 395.0 (76), 372.9 (18), 269.0 (100), 247.0 (25), 130.0 (20), 114.0 (16)

¹H NMR (400 MHz, DMSO-*d*₆) δ = 0.88-1.04 (m, 4 H) 1.79 (dd, *J*=14.98, 9.41 Hz, 1 H) 2.32 (dd, *J*=14.92, 2.38 Hz, 1 H) 3.21 - 3.37 (m, 2 H) 3.56 - 3.66 (m, 1 H) 3.83 (s, 3 H) 4.50 (d, *J*=5.32 Hz, 1 H) 4.57 (t, *J*=5.53 Hz, 1 H) 6.49 (td, *J*=8.85, 4.55 Hz, 1 H) 7.03 (dd, *J*=12.62, 7.24 Hz, 1 H) 7.34 (d, *J*=9.29 Hz, 1 H) 7.45 (br. s, 1 H) 7.56 (dd, *J*=10.91, 1.86 Hz, 1 H) 9.04 (br. s, 1 H) ppm

Optical purity: >99.9% ee

25

(S)-14 was obtained from (S)-13 which was prepared from (S)-10-Br

A mixture of (S)-13 (1.3 g, 2.1 mmol) in pyridine (0.76 mL), sulfolane (1.5 mL) and hydrochloric acid (10%, 4 mL) was stirred for 1.5h. It was diluted with ethyl acetate (20 mL) and water (30 mL), the organic layer was washed two times with water (30 mL, 2x 15 mL), dried over magnesium sulfate and concentrated to a solid. The solid was purified by chromatography (silica gel, ethyl acetate).

Yield: 1.1 g, 91.6%


R_f: 0.12 (silica gel, ethyl acetate)

MS (DCI): m/z 590.1 [M+NH₄]⁺(40), 573.0 [M+H]⁺(28), 395.0 (76), 372.9 (18), 269.0 (100), 247.0 (25), 130.0 (20), 114.0 (16)

¹H NMR (400 MHz, DMSO-*d*₆) δ = 0.88-1.04 (m, 4 H) 1.79 (dd, *J*=14.98, 9.41 Hz, 1 H) 2.32 (dd, *J*=14.92, 2.38 Hz, 1 H) 3.21 - 3.37 (m, 2 H) 3.56 - 3.66 (m, 1 H) 3.83 (s, 3 H) 4.50 (d, *J*=5.32 Hz, 1 H) 4.57 (t, *J*=5.53 Hz, 1 H) 6.49 (td, *J*=8.85, 4.55 Hz, 1 H) 7.03 (dd, *J*=12.62, 7.24 Hz, 1 H) 7.34 (d, *J*=9.29 Hz, 1 H) 7.45 (br. s, 1 H) 7.56 (dd, *J*=10.91, 1.86 Hz, 1 H) 9.04 (br. s, 1 H) ppm

Optical purity: >99.5% ee

Example 10b : N-{3,4-difluoro-2-[(2-fluoro-4-iodophenyl)amino]-6-methoxyphenyl}-1-[(2*R*)-2,3-dihydroxy-propyl]cyclopropanesulfonamide (R)-14



25

A solution of (R)-13 (120 mg, 0.6 mmol) in tetrahydrofuran (2.5 mL) and hydrochloric acid (10%, 2 mL) was stirred at room temperature until TLC (silica

gel, ethyl acetate) indicated complete conversion (2h). Ethyl acetate (2.5 mL) and a saturated aq. solution of sodium hydrogen carbonate (2.5 mL) were added, the aqueous layer was extracted with ethyl acetate (2.0 mL), the combined organic layer was washed with water (2 x 2 mL), dried over magnesium sulfate and concentrated to a solid. The solid was purified by chromatography (silica gel 1. ethyl acetate, 2. ethyl acetate/methanol 10:1).

Yield: 61 mg 54.4%

R_f: 0.12 (silica gel, ethyl acetate)

MS (DCI): m/z 590.1 [M+NH₄]⁺(40), 573.0 [M+H]⁺(28), 395.0 (76), 372.9 (18), 269.0 (100), 247.0 (25), 130.0 (20), 114.0 (16)

¹H NMR (400 MHz, DMSO-*d*₆) δ = 0.88-1.04 (m, 4 H) 1.79 (dd, *J*=14.98, 9.41 Hz, 1 H) 2.32 (dd, *J*=14.92, 2.38 Hz, 1 H) 3.21 - 3.37 (m, 2 H) 3.56 - 3.66 (m, 1 H) 3.83 (s, 3 H) 4.50 (d, *J*=5.32 Hz, 1 H) 4.57 (t, *J*=5.53 Hz, 1 H) 6.49 (td, *J*=8.85, 4.55 Hz, 1 H) 7.03 (dd, *J*=12.62, 7.24 Hz, 1 H) 7.34 (d, *J*=9.29 Hz, 1 H) 7.45 (br. s, 1 H) 7.56 (dd, *J*=10.91, 1.86 Hz, 1 H) 9.04 (br. s, 1 H) ppm

Optical purity: >99.9% ee

(R)-14 was obtained from **(R)-10-Br**

To a mixture of 5,6-difluoro-N1-(4-fluoro-2-iodophenyl)-3-methoxybenzene-1,2-diamine [ref. 2, Example 50, Step B] (0.8 g, 2.0 mmol) in pyridine (0.68 mL) and sulfolane (1.3 mL) was added **(R)-10-Br** (0.7 g, 2.4 mmol). It was stirred at 23 °C for 18h to achieve complete conversion to **(R)-13**. The reaction mixture was stirred with hydrochloric acid (10%, 6 mL) for 1.5h. It was diluted with ethyl acetate (20 mL) and water (30 mL), the organic layer was washed two times with water (30 mL, 2x 15 mL), dried over magnesium sulfate and concentrated to a solid. The solid was purified by chromatography (silica gel, ethyl acetate).

Yield: 1.1 g, 90.5%

R_f: 0.12 (silica gel, ethyl acetate)

MS (DCI): m/z 590.1 [M+NH₄]⁺(40), 573.0 [M+H]⁺(28), 395.0 (76), 372.9 (18), 269.0 (100), 247.0 (25), 130.0 (20), 114.0 (16)

¹H NMR (400 MHz, DMSO-*d*₆) δ = 0.88-1.04 (m, 4 H) 1.79 (dd, *J*=14.98, 9.41 Hz, 1 H) 2.32 (dd, *J*=14.92, 2.38 Hz, 1 H) 3.21 - 3.37 (m, 2 H) 3.56 - 3.66 (m, 1 H)
5 3.83 (s, 3 H) 4.50 (d, *J*=5.32 Hz, 1 H) 4.57 (t, *J*=5.53 Hz, 1 H) 6.49 (td, *J*=8.85, 4.55 Hz, 1 H) 7.03 (dd, *J*=12.62, 7.24 Hz, 1 H) 7.34 (d, *J*=9.29 Hz, 1 H) 7.45 (br. s, 1 H) 7.56 (dd, *J*=10.91, 1.86 Hz, 1 H) 9.04 (br. s, 1 H) ppm

Optical purity: >99.6% ee


- 10 The enantiomeric purity of **14** was determined by chiral HPLC (CHIRALPAK AD-H, 5 μm column length 25 cm, internal diameter 4.6 mm; temperature of the column oven 35°C; eluent n-heptane/isopropanol 8:2 v/v; flow rate 1.5 mL/min; detection wavelength 300nm; retention time (*S*)-**14** approx. 10 min, (*R*)-**14** approx. 16 min.

References :

- [1] C. Iverson, G. Larson, C. Lai, L.-T. Yeh, C. Dadson, P. Weingarten, T.
5 Appleby, T. Vo, A. Maderna, J.-M. Vernier; R. Hamatake, J.N. Miner, B.
Quart, Cancer Res. **2009**, 69, 6839-6847
- [2] U.S. Patent application published under 2008/0058340
- [3] PCT patent application published under WO 2010/145197 A1
- [4] S. Fujita, Synthesis **1982**, 423
- 10 [5] G. Blotny, Tetrahedron Lett. **2003**, 44, 1499-1501
- [6] J. Huang, T.S. Widlanski, Tetrahedron Lett. **1992**, 33, 2657-2660
- [7] PCT patent application published under WO 2011/009541 A1

CLAIMS

1. A method of preparing a compound of formula (S)-14 :




(S)-14 ;

comprising :


i) the following step J :

wherein a compound of formula (S)-10 :



(S)-10


is allowed to react with a compound of formula (12) :



(12)

in the presence of a bromide, such as tetra-n-butylammonium bromide or lithium bromide for example, optionally in the presence of a base, such as pyridine for example, optionally in a solvent, such as sulfolane for example,

thereby providing a reaction mixture containing a compound of formula (S)-13 :




5 and, then,

ii) the following step K :

wherein an aqueous mineral acid, such as hydrochloric acid for example, is
10 added to said reaction mixture containing a compound of formula (S)-13,


thereby providing a compound of formula (S)-14 ;

said compound of formula (S)-10 :




15

being prepared by allowing a compound of formula (S)-7 :




to react with a chlorinating agent, such as $P(O)Cl_3$, PCl_3 or $SOCl_2$ for example, optionally in a solvent, such as pyridine for example, thereby providing a compound of formula (S)-10 ;

said compound of formula (S)-7 :



(S)-7


being prepared by allowing a compound of formula (S)-6 :



(S)-6

to react with a sodium alkoxide, such as sodium methoxide for example, optionally in a solvent, such as methanol for example, thereby providing a compound of formula (S)-7.

2. A method of preparing a compound of formula (S)-14 :




(S)-14 ;

comprising :


i) the following step J :

wherein a compound of formula (S)-10-Br :



(S)-10-Br

is allowed to react with a compound of formula (12) :




(12)

5

optionally in the presence of a base, such as pyridine for example, optionally in a solvent, such as sulfolane for example,

thereby providing a reaction mixture containing a compound of formula (S)-

10 13 :



((S)-13)

;


and, then,

15 ii) the following step K :

wherein an aqueous mineral acid, such as hydrochloric acid for example, is added to said reaction mixture containing a compound of formula (S)-13,


thereby providing a compound of formula (S)-14 ;

said compound of formula (S)-10-Br :



(S)-10-Br


being prepared by allowing a compound of formula (S)-7 :



(S)-7


to react with a brominating agent, such as $P(O)Br_3$, PBr_3 or $SOBr_2$ for example, optionally in a solvent, such as pyridine for example, thereby providing a compound of formula (S)-10-Br;

said compound of formula (S)-7 :



(S)-7


being prepared by allowing a compound of formula (S)-6 :



(S)-6


to react with a sodium alkoxide, such as sodium methoxide for example, optionally in a solvent, such as methanol for example, thereby providing a compound of formula (S)-7.

3. The method of claim 1 or 2, wherein said compound of formula (S)-6 :



is prepared by allowing a compound of formula (S)-4 :

5




to react with either :

- a) boron trifluoride, optionally in the form of a complex, such as a boron trifluoride etherate complex for example, such as a boron trifluoride tetrahydrofuran complex, or a boron trifluoride diethylether complex for example, optionally in a solvent such as acetone for example, or
 - b) phosphomolybdic acid hydrate, optionally in a solvent such as acetone for example,
- thereby providing a compound of formula (S)-6.


15

4. The method of claim 3, wherein said compound of formula (S)-4 :




is prepared by allowing a compound of formula (S)-3 :

20



to react with a base, such as sodium hydroxide for example, optionally in a solvent, such as water for example, thereby providing a compound of formula (S)-4.


- 5 5. The method of claim 4, wherein said compound of formula (S)-3 :



is prepared by :

10

- a) allowing a compound of formula 2 :




to react with a base, such as n-butyllithium for example, optionally in a solvent, such as tetrahydrofuran for example,

15

- b) adding a compound of formula (S)-1 and boron trifluoride, optionally in the form of a complex, such as a boron trifluoride etherate complex for example, such as a boron trifluoride tetrahydrofuran complex, or a boron trifluoride diethylether complex for example ; and

20


- c) allowing the thus-formed deprotonated compound of formula 2 to react with a compound of formula (S)-1 :



25


thereby providing a compound of formula (S)-3.

6. The method of claim 1 or 2, wherein said compound of formula (S)-6 :



(S)-6

is prepared by allowing a compound of formula (S)-5 :




(S)-5

to react with 2,2-dimethoxypropane, optionally in a solvent, such as tetrahydrofuran for example and optionally in the presence of a catalyst, such as pyridinium p-toluenesulfonate for example,


thereby providing a compound of formula (S)-6.

7. The method of claim 6, wherein said compound of formula (S)-5 :



(S)-5

is prepared by allowing a compound of formula (S)-9 :




(S)-9 ;

wherein PG represents a protecting group, such as a t-butyldimethylsilyl-group (compound (S)-9a) or a tetrahydropyranyl group (compound (S)-9b), for example,

to react with an acid, such as hydrochloric acid or p-toluene sulfonic acid for example, optionally in a solvent, such as methanol for example,

thereby providing a compound of formula (S)-5.


8. The method of claim 7, wherein said compound of formula (S)-9 :



5 wherein PG represents a t-butyldimethylsilyl- group (compound (S)-9a) or a tetrahydropyranyl group (compound (S)-9b),

is prepared by :

a) allowing a compound of formula 2 :




10 to react with a base, such as n-butyllithium for example, optionally in a solvent, such as tetrahydrofuran for example,

b) adding a compound of formula (S)-8 and boron trifluoride, optionally in the form of a complex, such as a boron trifluoride etherate complex for example,

15 such as a boron trifluoride tetrahydrofuran complex, or a boron trifluoride diethylether complex for example ; and

c) allowing the thus-formed deprotonated compound of formula 2 to react with a compound of formula (S)-8 :




20 wherein PG represents a t-butyldimethylsilyl- group (compound (S)-8a) or a tetrahydropyranyl group (compound (S)-8b), respectively, thereby providing a compound of formula (S)-9.

25

9. One or more relevant compounds selected from :


1-[[*(4S)*-2,2-dimethyl-1,3-dioxolan-4-yl]methyl]cyclopropanesulfonyl bromide
(*S*)-10-Br :



5

(*S*)-10-Br ;


1-[[*(4R)*-2,2-dimethyl-1,3-dioxolan-4-yl]methyl]cyclopropanesulfonyl bromide
(*R*)-10-Br :



10

(*R*)-10-Br ;


15 potassium 1-[[*(4S)*-2,2-dimethyl-1,3-dioxolan-4-yl]methyl]cyclopropane-
sulfonate (*S*)-7' :



(*S*)-7'


;

20 potassium 1-[[*(4R)*-2,2-dimethyl-1,3-dioxolan-4-yl]methyl]cyclopropane-
sulfonate (*R*)-7' :




(R)-7' ;

Butyl 1-[[[(4S)-2,2-dimethyl-1,3-dioxolan-4-yl]methyl]cyclopropanesulfonate
5 (S)-6 :




(S)-6 ;

Butyl 1-[[[(4R)-2,2-dimethyl-1,3-dioxolan-4-yl]methyl]cyclopropanesulfonate (R
10)-6 :




(R)-6 ;

Butyl 1-[(2S)-2,3-dihydroxypropyl]cyclopropanesulfonate (S)-5 :




(S)-5 ;

Butyl 1-[(2R)-2,3-dihydroxypropyl]cyclopropanesulfonate (R)-5 :




(R)-5 ;

Butyl 1-[(2S)-oxiran-2-ylmethyl]cyclopropanesulfonate (S)-4 :




(S)-4 ;

Butyl 1-[(2R)-oxiran-2-ylmethyl]cyclopropanesulfonate (R)-4 :




(R)-4 ;

Butyl 1-[(2S)-3-chloro-2-hydroxypropyl]cyclopropanesulfonate (S)-3 :




(S)-3 ;

Butyl 1-[(2R)-3-chloro-2-hydroxypropyl]cyclopropanesulfonate (R)-3 :



(R)-3 ;


Butyl 1-[(2S)-3-[[tert-butyl(dimethyl)silyl]oxy]-2-hydroxypropyl]cyclopropane-sulfonate (S)-9a :



(S)-9a ;


Butyl 1-[(2R)-3-[[tert-butyl(dimethyl)silyl]oxy]-2-hydroxypropyl]cyclopropane-sulfonate (R)-9a :

106



(R)-9a ;


Butyl 1-[(2S)-2-hydroxy-3-(tetrahydro-2H-pyran-2-yloxy)propyl]cyclopropane sulfonate (S)-9b :



5

(S)-9b ;

Butyl 1-[(2R)-2-hydroxy-3-(tetrahydro-2H-pyran-2-yloxy)propyl]cyclopropane sulfonate (R)-9b :



10

(R)-9b;

used in any one of the method steps of claims 1-8.