


CLAIMS

1. A compound of Formula (I), or a pharmaceutically acceptable salt or solvate thereof, as shown below:



Formula (I)

wherein:

X_1 is selected from N or CR^a , wherein R^a is selected from hydrogen, (1-2C)alkyl, halogen, hydroxy, (1-2C)alkoxy, (1-2C)haloalkyl, (1-2C)haloalkoxy, (2-4C)alkenyl, (2-4C)alkynyl, nitro, cyano or NR^bR^c , wherein R^b and R^c are independently selected from hydrogen or (1-2C)alkyl;

X_2 is selected from N or CR^d , wherein R^d is selected from hydrogen, (1-2C)alkyl, fluoro, chloro, bromo, hydroxy, (1-2C)alkoxy, (1-2C)haloalkyl or (1-2C)haloalkoxy;

R^1 is selected from hydrogen or a group of the formula:



wherein:

L is absent or (1-5C)alkylene optionally substituted by one or more substituents selected from (1-2C)alkyl or oxo;


Y is absent or O, S, SO, SO_2 , $\text{N(R}^e\text{)}$, C(O) , C(O)O , OC(O) , $\text{C(O)N(R}^e\text{)}$, $\text{N(R}^e\text{)C(O)}$, $\text{N(R}^e\text{)C(O)N(R}^f\text{)}$, $\text{N(R}^e\text{)C(O)O}$, $\text{OC(O)N(R}^e\text{)}$, $\text{S(O)}_2\text{N(R}^e\text{)}$, or $\text{N(R}^e\text{)SO}_2$, wherein R^e and R^f are each independently selected from hydrogen or (1-4C)alkyl; and

Z is hydrogen, (1-6C)alkyl, (2-6C)alkenyl, (2-6C)alkynyl, aryl, (3-10C)cycloalkyl, (3-10C)cycloalkenyl, heteroaryl or heterocyclyl; wherein Z is optionally further substituted by one or more substituent groups independently selected from (1-4C)alkyl, halo, (1-4C)haloalkyl, (1-4C)haloalkoxy, amino, (1-4C)aminoalkyl, cyano, hydroxy, carboxy,

carbamoyl, sulphamoyl, mercapto, ureido, NR^gR^h , OR^g , $\text{C(O)}\text{R}^g$, $\text{C(O)}\text{OR}^g$, $\text{OC(O)}\text{R}^g$, $\text{C(O)}\text{N(R}^g\text{)R}^h$, $\text{N(R}^g\text{)C(O)R}^h$, $\text{S(O)}_y\text{R}^g$ (where y is 0, 1 or 2), $\text{SO}_2\text{N(R}^g\text{)R}^h$, $\text{N(R}^g\text{)SO}_2\text{R}^g$, $\text{Si(R}^g\text{)(R}^h\text{)R}^i$ or $(\text{CH}_2)_z\text{NR}^g\text{R}^h$ (where z is 1, 2 or 3); wherein R^g , R^h and R^i are each independently selected from hydrogen, (1-6C)alkyl or (3-6C)cycloalkyl; or R^g and R^h can be linked such that, together with the nitrogen atom to which they are attached, they form a 4-9 membered heterocyclic ring which is optionally substituted by one or more substituents selected from (1-4C)alkyl, halo, (1-4C)haloalkyl, (1-4C)haloalkoxy, (1-4C)alkoxy, (1-4C)alkylamino, amino, cyano or hydroxy;

R^2 is selected from:

- (i) hydrogen or methyl;
- (ii) a group of the formula:



wherein:

 denotes the point of attachment;

W_1 is selected from CR^4R^5 or C(O) , wherein R^4 and R^5 are independently selected from hydrogen, (1-2C)alkyl, fluoro, hydroxy, cyano, nitro, (1-2C)alkoxy, (1-2C)haloalkyl, (1-2C)haloalkoxy or NR^iR^k , wherein R^i and R^k are independently selected from hydrogen or (1-2C)alkyl; or


R^4 and R^5 can be linked such that, together with the carbon atom to which they are attached, they form a 3-6-membered carbocyclic ring or a 3-6-membered heterocyclic ring, which is optionally substituted by one or more substituents selected from (1-2C)alkyl, halo, (1-2C)haloalkyl, (1-2C)haloalkoxy, (1-2C)alkoxy, (1-2C)alkylamino, amino, cyano or hydroxy;

W_2 is selected from cyano, a 5- or 6-membered heteroaryl, phenyl, $\text{C(O)}\text{R}^l$, SO_2R^l , $\text{C(O)}\text{OCH}_3$, $\text{C(O)}\text{N(H)CH}_3$, $\text{CR}^6\text{R}^7\text{R}^8$ or NR^lR^m , wherein

R^l and R^m are independently selected from hydrogen or (1-4C)alkyl, and wherein:

R^6 is selected from hydrogen, (1-2C)alkyl, fluoro, chloro, bromo, hydroxy, amino, cyano, nitro, (1-2C)alkoxy, (1-2C)haloalkyl or (1-2C)haloalkoxy;

R^7 is selected from hydrogen, (1-2C)alkyl, fluoro, chloro, bromo, hydroxy, cyano, nitro, (1-2C)alkoxy, (1-2C)haloalkyl, (1-2C)haloalkoxy or a group of the formula:



wherein:

Y_2 is absent or selected from O, $N(R^n)$, S, SO, SO_2 , C(O), C(O)O, OC(O), C(O) $N(R^n)$, $N(R^n)$ C(O), $S(O)_2N(R^n)$, or $N(R^n)SO_2$, wherein R^n is selected from hydrogen or (1-2C)alkyl;


L_2 is absent or (1-2C)alkylene; and

Z_2 is hydrogen, (1-6C)alkyl, (2-4C)alkenyl, (2-4C)alkynyl, phenyl, (3-6C)cycloalkyl, 5-6 membered heteroaryl or a 4-6-membered heterocyclyl; wherein Z_2 is optionally substituted by one or more substituents selected from (1-4C)alkyl, halo, (1-4C)haloalkyl, (1-4C)haloalkoxy, (1-4C)alkoxy, (1-4C)alkylamino, amino, cyano, nitro, hydroxy, $C(O)R^o$, $C(O)OR^o$, $OC(O)R^o$, $C(O)N(R^o)R^p$, $NR^oC(O)R^p$, wherein R^o and R^p are independently selected from hydrogen or (1-4C)alkyl; and

R^8 is selected from (1-2C)alkyl, $-C(O)OR^q$, OR^q , $-C(O)NR^q$, NR^qR^r , phenyl or a 5-membered heteroaryl, wherein R^q and R^r are independently selected from hydrogen or (1-2C)alkyl;

or R^6 and R^7 can be linked such that, together with the carbon atom to which they are attached, they form a 3-6-membered carbocyclic ring or a 3-6-membered heterocyclic ring, which is optionally substituted by one or more substituents selected from (1-2C)alkyl, halo, (1-2C)haloalkyl, (1-2C)haloalkoxy, (1-2C)alkoxy, (1-2C)alkylamino, amino, cyano or hydroxyl; or

(iii) a group of the formula:



wherein:


denotes the point of attachment;

ring A is a 4 to 6 membered cycloalkyl or heterocyclyl ring, optionally substituted with one or more substituent groups selected from (1-2C)alkyl, halo, hydroxy, cyano or (1-2C)alkoxy;

W_3 is selected from NR^{100} or $CR^{101}R^{102}$, wherein R^{100} is selected from hydrogen, (1-2C)alkyl, (1-4C)haloalkyl, (1-4C)hydroxyalkyl, $-C(O)-CH_3$ or $-C(O)OR^{ab}$, wherein R^{ab} is (1-4C)alkyl, R^{101} and R^{102} are each independently selected from hydrogen, (1-2C)alkyl, cyclopropyl, fluoro, chloro, bromo, hydroxy, amino, cyano, nitro, (1-2C)alkoxy, (1-2C)haloalkyl, (1-2C)haloalkoxy, $-C(O)OR^{ac}$, $-NR^{ac}R^{ad}$, phenyl or a 5-membered heteroaryl, wherein R^{ac} and R^{ad} are independently selected from hydrogen or (1-2C)alkyl; and

R^3 is selected from:

i) a group of Formula A shown below:



Formula A

wherein:

denotes the point of attachment;

X_a and X_b are independently selected from N or CR^{x1} , wherein R^{x1} is selected from hydrogen, fluoro, chloro, bromo, (1-

2C)alkyl, (1-2C)alkoxy, cyano, nitro, acetylenyl, CH₂F, CF₂H or CF₃;

R⁹ is selected from hydrogen, fluoro, chloro, bromo, (1-2C)alkyl, (1-2C)alkoxy, cyano, nitro, acetylenyl, CH₂F, CF₂H or CF₃;


R¹⁰ is selected from hydrogen, halo, (1-4C)alkyl, (1-4C)alkoxy, (1-4C)haloalkyl, (1-4C)haloalkoxy, cyano, nitro, (2-4C)alkenyl, (2-4C)alkynyl or a group of the formula:



wherein:

Y₃ is absent or O, N(R^s)(CR^sR^t)_{q₁} (where q₁ is 0, 1 or 2), S, SO, SO₂, C(O), C(O)O, OC(O), C(O)N(R^s), N(R^s)C(O), N(R^s)C(O)N(R^t), N(R^s)C(O)O, OC(O)N(R^s), S(O)₂N(R^s), N(R^s)SO₂, wherein R^s and R^t are each independently selected from hydrogen or (1-4C)alkyl; and

Z₃ is hydrogen, (1-6C)alkyl, aryl, (3-6C)cycloalkyl, (2-4C)alkenyl, (2-4C)alkynyl, (3-6C)cycloalkenyl, heteroaryl or 4 to 11-membered heterocyclyl; wherein Z₃ is optionally further substituted by one or more substituent groups independently selected from oxo, (1-4C)alkyl, (3-6C)cycloalkyl, halo, (1-4C)haloalkyl, (1-4C)haloalkoxy, (1-4C)hydroxyalkyl, amino, cyano, hydroxy, amido, carboxy, carbamoyl, sulphamoyl, mercapto, C(O)NR^uR^v, NR^uR^v or OR^u, wherein R^u and R^v are each independently selected from hydrogen, (1-4C)alkyl or (3-6C)cycloalkyl; and/or Z₃ is optionally further substituted by a group of the formula:




wherein:

L_Z is absent or a (1-5C)alkylene optionally substituted by one or more substituents selected from (1-2C)alkyl or oxo; and

W_z is aryl, heteroaryl, 4- to 7-membered heterocyclyl, 3- to 6-membered carbocyclyl, halo, (1-4C)haloalkyl, (1-4C)haloalkoxy, cyano, hydroxy, (1-4C)alkoxy, $C(O)R^{xa}$, $COOR^{xa}$, $C(O)NR^{xa}R^{xb}$ or $NR^{xa}R^{xb}$, wherein R^{xa} and R^{xb} are each independently selected from hydrogen or (1-4C)alkyl; and wherein each aryl, heteroaryl, 4- to 7-membered heterocyclyl or 3- to 6-membered carbocyclyl is optionally further substituted by one or more substituent groups independently selected from (1-4C)alkyl, halo, (1-4C)haloalkyl, amino, cyano or hydroxy;

ii) a group of Formula B shown below:



Formula B

wherein:

 denotes the point of attachment;

X_c , X_d and X_e are independently selected from N, CH, CF, CCl, C-CN or CCH_3 ;

R^{11} is selected from hydrogen, halo, (1-4C)alkyl, (1-4C)alkoxy, (1-4C)haloalkyl, (1-4C)haloalkoxy, cyano, nitro, (2-4C)alkenyl, (2-4C)alkynyl or a group of the formula:




wherein:

Y_5 is absent or O, $N(R^w)$, $C(O)$, $C(O)O$, $OC(O)$, $C(O)N(R^w)$, $N(R^w)C(O)$, $N(R^w)C(O)N(R^x)$, $N(R^w)C(O)O$, $OC(O)N(R^w)$, $S(O)_2N(R^w)$, $N(R^w)SO_2$, wherein R^w and R^x are each independently selected from hydrogen or (1-4C)alkyl; and

Z_5 is hydrogen, (1-6C)alkyl, aryl, (3-8C)cycloalkyl, (3-8C)cycloalkenyl, heteroaryl or heterocyclyl; wherein Z_5 is optionally further substituted by one or more substituent groups independently selected from (1-4C)alkyl, halo, (1-4C)haloalkyl, (1-4C)haloalkoxy, (1-4C)hydroxyalkyl, amino, cyano, hydroxy, amido, carboxy, carbamoyl, sulphamoyl, mercapto, NR^yR^z , OR^y , wherein R^y and R^z are each independently selected from hydrogen, (1-4C)alkyl or (3-6C)cycloalkyl;

iii) a group of Formula C shown below:



Formula C

wherein:

 denotes the point of attachment;

R^{12} is selected from fluoro, chloro, bromo, (1-2C)alkyl, (1-2C)alkoxy, cyano, nitro, (2-4C)alkenyl, (2-4C)alkynyl, CH_2F , CF_2H or CF_3 ;


X_f and X_g are independently selected from N or CR^{13} , wherein R^{13} is selected from hydrogen, fluoro, chloro, (1-2C)alkyl, (1-2C)haloalkyl or (1-2C)haloalkoxy;

X_h , X_i and X_j are independently selected from N or CR^{14} , wherein R^{14} is selected from hydrogen, halo, (1-2C)alkyl, (1-2C)alkoxy, (1-2C)haloalkyl or (1-2C)haloalkoxy;

with the proviso that:

- (i) when R^3 is a group of Formula B, no more than two of X_c , X_d and X_e are nitrogen;

- (ii) when R^3 is a group of Formula C, no more than three of X_f , X_g , X_h , X_i and X_j are nitrogen;
- (iii) when X_1 and X_2 are CH, R^1 and R^2 are hydrogen or methyl, R^3 is a group of Formula A, X_a is N, X_b is CH and R^9 is methyl or fluoro, R^{10} is not a methylsulfonylaminophenyl or an aminosulfonylphenyl; and
- (iv) when Y_3 is NH, each of R^1 and R^2 are not hydrogen or methyl; and
- (v) the compound is not one of the following:



2. A compound of Formula (I), or a pharmaceutically acceptable salt or solvate thereof, according to claim 1, wherein X_1 is selected from N or CR^a , wherein R^a is selected from hydrogen, methyl, fluoro, chloro, hydroxy, OCH_3 , CH_2F , CHF_2 , CF_3 , OCF_3 , acetylenyl, cyano or NH_2 .
3. A compound of Formula (I), or a pharmaceutically acceptable salt or solvate thereof, according to claims 1 or 2, wherein X_2 is selected from N or CH.
4. A compound of Formula (I), or a pharmaceutically acceptable salt or solvate thereof, according to any one of claims 1 to 3, wherein X_1 is CH and X_2 is CH.
5. A compound of Formula (I), or a pharmaceutically acceptable salt or solvate thereof, according to any one of claims 1 to 4, wherein R^1 is selected from hydrogen or a group of the formula:



wherein:

L is absent or (1-5C)alkylene optionally substituted by one or more substituents selected from (1-2C)alkyl;

Y is absent or O, S, SO, SO₂, N(R^e), C(O), C(O)O, OC(O), C(O)N(R^e), N(R^e)C(O), S(O)₂N(R^e), or N(R^e)SO₂, wherein R^e is selected from hydrogen or (1-4C)alkyl; and

Z is hydrogen, (1-6C)alkyl, (2-6C)alkenyl, (2-6C)alkynyl, aryl, (3-10C)cycloalkyl, (3-10C)cycloalkenyl, heteroaryl or heterocyclyl; wherein Z is optionally further substituted by one or more substituent groups independently selected from (1-4C)alkyl, halo, (1-4C)haloalkyl, (1-4C)haloalkoxy, amino, (1-4C)aminoalkyl, cyano, hydroxy, carboxy, carbamoyl, sulphamoyl, mercapto, ureido, NR^gR^h, OR^g, C(O)R^g, C(O)OR^g, OC(O)R^g, C(O)N(R^g)R^h, N(R^g)C(O)R^h, S(O)_yR^g (where y is 0, 1 or 2), SO₂N(R^g)R^h, N(R^g)SO₂R^g, Si(R^g)(R^h)Rⁱ or (CH₂)_zNR^gR^h (where z is 1, 2 or 3); wherein R^g, R^h and Rⁱ are each independently selected from hydrogen, (1-6C)alkyl or (3-6C)cycloalkyl; or R^g and R^h can be linked such that, together with the nitrogen atom to which they are attached, they form a 4-9 membered heterocyclic ring.

6. A compound of Formula (I), or a pharmaceutically acceptable salt or solvate thereof, according to any one of claims 1 to 5, wherein R¹ is selected from hydrogen or a group of the formula:



wherein:

L is absent or (1-5C)alkylene;


Y is absent or O, SO₂, C(O), C(O)O, C(O)N(R^e) or S(O)₂N(R^e), wherein R^e is selected from hydrogen or (1-4C)alkyl; and

Z is hydrogen, (1-6C)alkyl, (2-6C)alkenyl, (2-6C)alkynyl, aryl, (3-10C)cycloalkyl, (3-10C)cycloalkenyl, heteroaryl or heterocyclyl; wherein Z is optionally further substituted by one or more substituent groups independently selected from (1-2C)alkyl, halo, (1-2C)haloalkyl, (1-2C)haloalkoxy, amino, (1-2C)aminoalkyl, cyano, hydroxy, NR^gR^h or

OR^g; wherein R^g, R^h and Rⁱ are each independently selected from hydrogen or (1-4C)alkyl.

7. A compound of Formula (I), or a pharmaceutically acceptable salt or solvate thereof, according to any one of claims 1 to 6, wherein R² is selected from:

- (i) hydrogen or methyl;
- (ii) a group of the formula:



wherein:

 denotes the point of attachment;


W₁ is selected from CR⁴R⁵ or C(O), wherein R⁴ and R⁵ are independently selected from hydrogen, (1-2C)alkyl, fluoro, hydroxy, cyano, (1-2C)alkoxy, CH₂F, CF₂H or amino; or

R⁴ and R⁵ can be linked such that, together with the carbon atom to which they are attached, they form a 3-6-membered carbocyclic ring or a 3-6-membered heterocyclic ring, which is optionally substituted by one or more substituents selected from methyl, fluoro, chloro, OCH₃, amino, cyano or hydroxy;

W₂ is selected from cyano, a 5- or 6-membered heteroaryl, phenyl, C(O)OCH₃, C(O)N(H)CH₃, CR⁶R⁷R⁸ or NR^lR^m, wherein R^l and R^m are independently selected from hydrogen or (1-2C)alkyl, and wherein:

R⁶ is selected from hydrogen, (1-2C)alkyl, fluoro, chloro, hydroxy, amino, cyano or (1-2C)alkoxy;

R⁷ is selected from hydrogen, (1-2C)alkyl, fluoro, chloro, hydroxy, cyano, (1-2C)alkoxy, (1-2C)haloalkyl, (1-2C)haloalkoxy or a group of the formula:



wherein:


Y_2 is absent or selected from O, C(O), C(O)O, OC(O), C(O)N(Rⁿ) or N(Rⁿ)C(O), wherein Rⁿ is selected from hydrogen or (1-2C)alkyl;

L_2 is absent or (1-2C)alkylene; and

Z_2 is hydrogen, (1-6C)alkyl, (2-4C)alkenyl, (2-4C)alkynyl, phenyl, (3-6C)cycloalkyl, 5-6 membered heteroaryl or a 4-6-membered heterocyclyl; wherein Z_2 is optionally substituted by one or more substituents selected from (1-2C)alkyl, halo, (1-2C)haloalkyl, (1-2C)haloalkoxy, (1-2C)alkoxy, (1-2C)alkylamino, amino, cyano, nitro or hydroxy; and

R^8 is selected from (1-2C)alkyl, -C(O)OR^q, OR^q, NR^qR^r, phenyl or a 5-membered heteroaryl, wherein R^q and R^r are independently selected from hydrogen or (1-2C)alkyl; or

(iii) a group of the formula:



wherein:


 denotes the point of attachment;

ring A is a 5-membered cycloalkyl or heterocyclyl ring;

W_3 is selected from NR¹⁰⁰ or CR¹⁰¹R¹⁰², wherein R¹⁰⁰ is selected from hydrogen, (1-2C)alkyl or -C(O)OR^{ab}, wherein R^{ab} is (1-2C)alkyl, R¹⁰¹ and R¹⁰² are each independently selected from hydrogen, (1-2C)alkyl, fluoro, chloro, hydroxy, (1-2C)alkoxy, CH₂F, CF₂H, CF₃, -C(O)OR^{ac} or -NR^{ac}R^{ad}, and wherein R^{ac} and R^{ad} are independently selected from hydrogen or (1-2C)alkyl.

8. A compound of Formula (I), or a pharmaceutically acceptable salt or solvate thereof, according to any one of claims 1 to 7, wherein R² is selected from:

- (i) hydrogen or methyl;
- (ii) a group of the formula:



wherein:


 denotes the point of attachment;

W₁ is selected from CHR⁴ or C(O), wherein R⁴ is selected from hydrogen, (1-2C)alkyl, fluoro, hydroxy, cyano, (1-2C)alkoxy, CH₂F, CF₂H or amino;

W₂ is selected from cyano, a 5- or 6-membered heteroaryl, phenyl, C(O)OCH₃, C(O)N(H)CH₃, CR⁶R⁷R⁸ or NR^lR^m, wherein R^l and R^m are independently selected from hydrogen or (1-2C)alkyl, and wherein:

R⁶ is selected from hydrogen, (1-2C)alkyl, fluoro, chloro, hydroxy, amino, cyano, (1-2C)alkoxy, CH₂F or CF₂H;

R⁷ is selected from hydrogen, (1-2C)alkyl, fluoro, chloro, hydroxy, cyano, (1-2C)alkoxy, (1-2C)haloalkyl, (1-2C)haloalkoxy or a group of the formula:



wherein:

Y₂ is absent or selected from O, C(O)O, C(O)N(Rⁿ) or N(Rⁿ)C(O), wherein Rⁿ is selected from hydrogen or (1-2C)alkyl;


L₂ is absent or (1-2C)alkylene; and

Z₂ is hydrogen, (1-6C)alkyl, (2-4C)alkenyl, (2-4C)alkynyl, phenyl, (3-6C)cycloalkyl, 5-6 membered heteroaryl or a 4-6-membered heterocyclyl; wherein Z₂ is optionally substituted by one or more substituents selected from (1-2C)alkyl, halo, (1-2C)haloalkyl, (1-2C)haloalkoxy, (1-2C)alkoxy, (1-2C)alkylamino, amino, cyano or hydroxy; and

R⁸ is selected from (1-2C)alkyl, -C(O)OR^q, OR^q, NR^qR^r, phenyl or a 5-membered heteroaryl, wherein R^q and R^r are independently selected from hydrogen or (1-2C)alkyl; or

R^4 and R^7 can be linked such that, together with the carbon atoms to which they are attached, they form a 4-6 membered carbocyclic ring or a 4-6 membered heterocyclic ring; or

(iii) a group of the formula:



wherein:


denotes the point of attachment;

ring A is a 5-membered cycloalkyl or heterocyclyl ring;

W_3 is selected from NR^{100} or $CR^{101}R^{102}$, wherein R^{100} is selected from hydrogen, (1-2C)alkyl or $-C(O)OR^{ab}$, wherein R^{ab} is (1-2C)alkyl, R^{101} is selected from hydrogen or methyl and R^{102} is selected from (1-2C)alkyl, hydroxy, (1-2C)alkoxy, $C(O)OR^{ac}$ or $-NR^{ac}R^{ad}$, and wherein R^{ac} and R^{ad} are independently selected from hydrogen or (1-2C)alkyl.

9. A compound of Formula (I), or a pharmaceutically acceptable salt or solvate thereof, according to any one of claims 1 to 8, wherein R^3 is selected from:

i) a group of Formula A shown below:



Formula A

wherein:

denotes the point of attachment;

X_a and X_b are independently selected from N or CR^{x1} , wherein R^{x1} is selected from hydrogen, fluoro, chloro,

bromo, (1-2C)alkyl, (1-2C)alkoxy, cyano, nitro, acetylenyl, CH₂F, CF₂H or CF₃;

R⁹ is selected from hydrogen, fluoro, chloro, bromo, (1-2C)alkyl, (1-2C)alkoxy, cyano, nitro, acetylenyl, CH₂F, CF₂H or CF₃;


R¹⁰ is selected from hydrogen, halo, (1-4C)alkyl, (1-4C)alkoxy, (1-4C)haloalkyl, (1-4C)haloalkoxy, cyano, nitro, (2-4C)alkenyl, (2-4C)alkynyl or a group of the formula:



wherein:

Y₃ is absent or N(R^s)(CR^sR^t)_{q₁} (where q₁ is 0, 1 or 2), S, C(O), C(O)O, C(O)N(R^s), N(R^s)C(O), S(O)₂N(R^s) or N(R^s)SO₂, wherein R^s is selected from hydrogen or (1-4C)alkyl; and

Z₃ is hydrogen, (1-6C)alkyl, aryl, (3-6C)cycloalkyl, (2-4C)alkenyl, (2-4C)alkynyl, (3-6C)cycloalkenyl, 5- or 6-membered heteroaryl or 4 to 11-membered heterocyclyl; wherein Z₃ is optionally further substituted by one or more substituent groups independently selected from oxo, (1-4C)alkyl, (3-6C)cycloalkyl, halo, (1-4C)haloalkyl, (1-4C)haloalkoxy, amino, cyano, hydroxy, amido, carboxy, carbamoyl, sulphamoyl, mercapto, C(O)NR^uR^v, NR^uR^v or OR^u, wherein R^u and R^v are each independently selected from hydrogen, (1-4C)alkyl or (3-6C)cycloalkyl; and/or Z³ is optionally further substituted by a group of the formula:




wherein:

L_Z is absent or a (1-3C)alkylene; and

W_z is phenyl, 5- or 6-membered heteroaryl, 6-membered heterocyclyl, halo, (1-4C)haloalkyl, (1-4C)haloalkoxy, cyano, hydroxy, (1-4C)alkoxy, $C(O)R^{xa}$, $COOR^{xb}$, $C(O)NR^{xa}R^{xb}$ or $NR^{xa}R^{xb}$, wherein R^{xa} and R^{xb} are each independently selected from hydrogen or (1-4C)alkyl;

ii) a group of Formula B shown below:



Formula B

wherein:

 denotes the point of attachment;

X_c , X_d and X_e are independently selected from N, CH, CF, CCl, C-CN or CCH_3 ;

R^{11} is selected from hydrogen, halo, (1-4C)alkyl, (1-4C)alkoxy, (1-4C)haloalkyl, (1-4C)haloalkoxy, cyano, nitro, (2-4C)alkenyl, (2-4C)alkynyl or a group of the formula:




wherein:

Y_5 is absent or O, $N(R^w)$, $C(O)$, $C(O)O$, $C(O)N(R^w)$ or $S(O)_2N(R^w)$, wherein R^w is selected from hydrogen or (1-4C)alkyl; and

Z_5 is hydrogen, (1-6C)alkyl, aryl, (3-8C)cycloalkyl, (3-8C)cycloalkenyl, heteroaryl or heterocyclyl; wherein Z_5 is optionally further substituted by one or more substituent groups independently selected from (1-4C)alkyl, halo,

(1-4C)haloalkyl, (1-4C)haloalkoxy, amino, cyano, hydroxy, amido, carboxy, carbamoyl, sulphamoyl, mercapto, NR^yR^z , OR^y , wherein R^y and R^z are each independently selected from hydrogen, (1-4C)alkyl or (3-6C)cycloalkyl;

iii) a group of Formula C shown below:



Formula C

wherein:

 denotes the point of attachment;


R^{12} is selected from fluoro, chloro, bromo, (1-2C)alkyl, (1-2C)alkoxy, cyano, nitro, (2-4C)alkynyl, CH_2F , CF_2H or CF_3 ;

X_f and X_g are independently selected from N or CR^{13} , wherein R^{13} is selected from hydrogen, fluoro, chloro, methyl, CH_2F , CF_2H or CF_3 ;

X_h , X_i and X_j are independently selected from N or CR^{14} , wherein R^{14} is selected from hydrogen, halo, (1-2C)alkyl, (1-2C)alkoxy, (1-2C)haloalkyl or (1-2C)haloalkoxy.

10. A compound of Formula (I), or a pharmaceutically acceptable salt or solvate thereof, according to any one of claims 1 to 9, wherein R^3 is selected from:

i) a group of Formula A shown below:



Formula A

wherein:

 denotes the point of attachment;

X_a and X_b are independently selected from N or CR^{x1} , wherein R^{x1} is selected from hydrogen, fluoro, chloro, bromo, (1-2C)alkyl, (1-2C)alkoxy, cyano, nitro, acetylenyl, CH_2F , CF_2H or CF_3 ;

R^9 is selected from hydrogen, fluoro, chloro, bromo, (1-2C)alkyl, (1-2C)alkoxy, cyano, nitro, acetylenyl, CH_2F , CF_2H or CF_3 ;


R^{10} is selected from hydrogen, halo, (1-4C)alkyl, (1-4C)alkoxy, (1-4C)haloalkyl, (1-4C)haloalkoxy, cyano, nitro, (2-4C)alkenyl, (2-4C)alkynyl or a group of the formula:



wherein:

Y_3 is absent or $N(R^s)(CR^sR^t)_{q_1}$ (where q_1 is 0, 1 or 2), S, C(O), C(O)O, C(O)N(R^s), N(R^s)C(O) or S(O)₂N(R^s), wherein R^s is selected from hydrogen or (1-4C)alkyl; and

Z_3 is hydrogen, (1-6C)alkyl, aryl, (3-6C)cycloalkyl, (2-4C)alkenyl, (2-4C)alkynyl, (3-6C)cycloalkenyl, heteroaryl or 4 to 11-membered heterocyclyl; wherein Z_3 is optionally further substituted by one or more substituent groups independently selected from oxo, (1-4C)alkyl, (3-6C)cycloalkyl, halo, (1-4C)haloalkyl, (1-4C)haloalkoxy, amino, cyano, hydroxy, amido, carboxy, carbamoyl, sulphamoyl, mercapto, C(O)NR^uR^v, NR^uR^v or OR^u, wherein R^u and R^v are each independently selected from hydrogen, (1-4C)alkyl or (3-6C)cycloalkyl; and/or Z^3 is optionally further substituted by a group of the formula:




wherein:

L_z is absent or a (1-5C)alkylene optionally substituted by one or more substituents selected from (1-2C)alkyl or oxo; and

W_z is phenyl, 5- or 6-membered heteroaryl, 6-membered heterocyclyl, halo, (1-4C)haloalkyl, (1-4C)haloalkoxy, cyano, hydroxy, (1-4C)alkoxy, $C(O)R^{xa}$, $COOR^{xa}$, $C(O)NR^{xa}R^{xb}$ or $NR^{xa}R^{xb}$, wherein R^{xa} and R^{xb} are each independently selected from hydrogen or (1-4C)alkyl;

ii) a group of Formula B shown below:



Formula B

wherein:

 denotes the point of attachment;

X_c , X_d and X_e are independently selected from N, CH, CF, CCl, C-CN or CCH_3 ;

R^{11} is selected from hydrogen, halo, (1-4C)alkyl, (1-4C)alkoxy, (1-4C)haloalkyl, (1-4C)haloalkoxy, cyano, nitro, (2-4C)alkenyl, (2-4C)alkynyl or a group of the formula:




wherein:

Y_5 is absent or O, $N(R^w)$, $C(O)$, $C(O)O$, $C(O)N(R^w)$ or $S(O)_2N(R^w)$, wherein R^w is selected from hydrogen or (1-4C)alkyl; and

Z_5 is hydrogen, (1-6C)alkyl, aryl, (3-8C)cycloalkyl, (3-8C)cycloalkenyl, heteroaryl or heterocyclyl; wherein Z_5 is optionally further substituted by one or more substituent groups independently selected from (1-4C)alkyl, halo, (1-4C)haloalkyl, (1-4C)haloalkoxy, amino, cyano, hydroxy, amido, carboxy, carbamoyl, sulphamoyl, mercapto, NR^yR^z , OR^y , wherein R^y and R^z are each independently selected from hydrogen, (1-4C)alkyl or cyclopropyl.

11. A compound of Formula (I), or a pharmaceutically acceptable salt or solvate thereof, according to any one of claims 1 to 10, wherein R^3 is a group of Formula A shown below:



Formula A

wherein:

 denotes the point of attachment;

X_a is CH or N;

X_b is selected from N or CR^{x1} , wherein R^{x1} is selected from hydrogen, fluoro, chloro, bromo, methyl, OCH_3 , cyano or acetylenyl;

R^9 is selected from chloro or cyano;


R^{10} is selected from hydrogen, halo, (1-4C)alkyl, (1-4C)alkoxy, (1-4C)haloalkyl, (1-4C)haloalkoxy, cyano or a group of the formula:



wherein:

Y_3 is absent or $N(R^s)(CH_2)_{q_1}$ (where q_1 is 0 or 1), $C(O)$, $C(O)O$ or $C(O)N(R^s)$, wherein R^s is selected from hydrogen or methyl; and

Z_3 is hydrogen, (1-6C)alkyl, aryl, (3-6C)cycloalkyl, 5- or 6-membered heteroaryl or 4 to 9-membered heterocyclyl; wherein Z_3 is optionally further substituted by one or more substituent groups independently selected from oxo, (1-3C)alkyl, cyclopropyl, halo, (1-2C)haloalkyl, (1-2C)haloalkoxy, amino, cyano, hydroxy, amido, carboxy, $C(O)NR^uR^v$, NR^uR^v or OR^u , wherein R^u and R^v are each independently selected from hydrogen or methyl; or Z^3 is optionally further substituted by a group of the formula:



wherein:

L_z is absent or a (1-3C)alkylene; and

W_z is phenyl, 5- or 6-membered heteroaryl, 6-membered heterocyclyl, halo, (1-2C)haloalkyl, (1-2C)haloalkoxy, cyano, hydroxy, (1-2C)alkoxy, $C(O)R^{xa}$, $COOR^{xa}$, $C(O)NR^{xa}R^{xb}$ or $NR^{xa}R^{xb}$, wherein R^{xa} and R^{xb} are each independently selected from hydrogen or methyl.

12. A compound, or a pharmaceutically acceptable salt or solvate thereof, selected from one of the following:

6-Chloro-5-cyano-4-[[3-(3-hydroxy-3-methyl-butyl)-1-methyl-2-oxo-benzimidazol-5-yl]amino]-*N*-methyl-pyridine-2-carboxamide;

2-chloro-4-[[3-(3-hydroxy-3-methyl-butyl)-1-methyl-2-oxo-benzimidazol-5-yl]amino]-6-methyl-pyridine-3-carbonitrile;

6-chloro-5-cyano-4-[(1,3-dimethyl-2-oxo-benzimidazol-5-yl)amino]pyridine-2-carboxylic acid;

6-chloro-5-cyano-*N*-methyl-4-[[1-methyl-2-oxo-3-[(3*S*)-3-pyrazol-1-ylbutyl]benzimidazol-5-yl]amino]pyridine-2-carboxamide;

6-chloro-5-cyano-4-[[3-[(3*R*)-3-hydroxybutyl]-1-methyl-2-oxo-benzimidazol-5-yl]amino]pyridine-2-carboxylic acid;

6-chloro-5-cyano-*N*-methyl-4-[(1-methyl-2-oxo-3*H*-benzimidazol-5-yl)amino]pyridine-2-carboxamide;

6-chloro-5-cyano-4-[[3-(3-hydroxy-3-methyl-butyl)-2-oxo-1-(tetrahydropyran-4-ylmethyl)benzimidazol-5-yl]amino]-*N*-methyl-pyridine-2-carboxamide;

Ethyl 7-[[3-(3-hydroxy-3-methyl-butyl)-1-methyl-2-oxo-benzimidazol-5-yl]amino]pyrazolo[1,5-*a*]pyrimidine-5-carboxylate;

2-chloro-4-[[3-(3-hydroxy-3-methyl-butyl)-1-methyl-2-oxo-benzimidazol-5-yl]amino]pyridine-3-carbonitrile;

Methyl 4-[6-[(2-chloro-3-cyano-4-pyridyl)amino]-3-methyl-2-oxo-benzimidazol-1-yl]-2-methyl-butanoate;

2-bromo-4-[[3-(3-hydroxy-3-methyl-butyl)-1-methyl-2-oxo-benzimidazol-5-yl]amino]pyridine-3-carbonitrile;

2-chloro-4-[[3-[(3*R*)-3-hydroxybutyl]-1-methyl-2-oxo-benzimidazol-5-yl]amino]-6-methyl-pyridine-3-carbonitrile;

5-[(2,5-dichloro-4-pyridyl)amino]-3-[(3*R*)-3-hydroxybutyl]-1-methyl-benzimidazol-2-one;

5-[(2,5-dichloropyrimidin-4-yl)amino]-3-(3-hydroxy-3-methyl-butyl)-1-methyl-benzimidazol-2-one;

Ethyl 7-[[3-[(3*R*)-3-hydroxybutyl]-1-methyl-2-oxo-benzimidazol-5-yl]amino]pyrazolo[1,5-*a*]pyrimidine-5-carboxylate;

4-chloro-6-[[3-(3-hydroxy-3-methyl-butyl)-1-methyl-2-oxo-benzimidazol-5-yl]amino]pyrimidine-5-carbonitrile;

5-[(2,3-dichloro-4-pyridyl)amino]-3-[(3*R*)-3-hydroxybutyl]-1-methyl-benzimidazol-2-one;

Ethyl 3-fluoro-7-[(3-(2-hydroxybutyl)-1-methyl-2-oxo-2,3-dihydro-1*H*-benzo[*d*]imidazol-5-yl)amino]pyrazolo[1,5-*a*]pyrimidine-5-carboxylate;

Methyl 6-chloro-5-cyano-4-[[3-(4-methoxy-3-methyl-4-oxo-butyl)-1-methyl-2-oxo-benzimidazol-5-yl]amino]pyridine-2-carboxylate;

Ethyl 6-chloro-5-cyano-4-[[3-(4-methoxy-3-methyl-4-oxo-butyl)-1-methyl-2-oxo-benzimidazol-5-yl]amino]pyridine-2-carboxylate;

Isopropyl 6-chloro-5-cyano-4-[[3-(4-methoxy-3-methyl-4-oxo-butyl)-1-methyl-2-oxo-benzimidazol-5-yl]amino]pyridine-2-carboxylate;

Ethyl 6-chloro-5-cyano-4-[[3-[(3*R*)-3-hydroxybutyl]-1-methyl-2-oxo-benzimidazol-5-yl]amino]pyridine-2-carboxylate;

6-Chloro-5-cyano-4-[[3-(4-methoxy-3-methyl-4-oxo-butyl)-1-methyl-2-oxo-benzimidazol-5-yl]amino]pyridine-2-carboxylic acid;

Methyl 3-[6-[(2-chloro-3-cyano-4-pyridyl)amino]-3-methyl-2-oxo-benzimidazol-1-yl]-2-methyl-propanoate;

Methyl 4-[6-[(5-chloro-2-methyl-pyrimidin-4-yl)amino]-3-methyl-2-oxo-benzimidazol-1-yl]-2-methyl-butanoate;

6-Chloro-5-cyano-4-[[3-[(3*R*)-3-hydroxybutyl]-1-methyl-2-oxo-benzimidazol-5-yl]amino]-*N*-methyl-pyridine-2-carboxamide;

Methyl 4-[6-[[2-chloro-3-cyano-6-(3-methyl-1,2,4-oxadiazol-5-yl)-4-pyridyl]amino]-3-methyl-2-oxo-benzimidazol-1-yl]-2-methyl-butanoate;

Methyl (2*S*)-2-amino-4-[6-[(2-chloro-3-cyano-4-pyridyl)amino]-3-methyl-2-oxo-benzimidazol-1-yl]butanoate;

Methyl 4-[6-[[2-chloro-3-cyano-6-(methylcarbamoyl)-4-pyridyl]amino]-3-methyl-2-oxo-benzimidazol-1-yl]-2-methyl-butanoate;

Methyl 4-[6-[[6-(but-3-ynylcarbamoyl)-2-chloro-3-cyano-4-pyridyl]amino]-3-methyl-2-oxo-benzimidazol-1-yl]-2-methyl-butanoate;

Methyl 4-[6-[[2-chloro-3-cyano-6-(dimethylcarbamoyl)-4-pyridyl]amino]-3-methyl-2-oxo-benzimidazol-1-yl]-2-methyl-butanoate;

6-Chloro-5-cyano-*N*-[2-(dimethylamino)ethyl]-4-[(1,3-dimethyl-2-oxo-benzimidazol-5-yl)amino]pyridine-2-carboxamide;

Ethyl 7-[[3-(4-methoxy-3-methyl-4-oxo-butyl)-1-methyl-2-oxo-benzimidazol-5-yl]amino]pyrazolo[1,5-*a*]pyrimidine-5-carboxylate;

Methyl 4-[6-[(2-chloro-3-cyano-4-pyridyl)amino]-3-methyl-2-oxo-benzimidazol-1-yl]-2-hydroxy-butanoate;

2-Chloro-4-[[3-(2,3-dihydroxy-3-methyl-butyl)-1-methyl-2-oxo-benzimidazol-5-yl]amino]pyridine-3-carbonitrile;

Methyl 4-[6-[(2-chloro-3-cyano-4-pyridyl)amino]-3-methyl-2-oxo-benzimidazol-1-yl]-2-methoxy-butanoate;

Methyl 4-[6-[(2-chloro-3-cyano-4-pyridyl)amino]-3-methyl-2-oxo-benzimidazol-1-yl]-2-ethoxy-butanoate;

Methyl 4-[6-[(2-chloro-3-cyano-4-pyridyl)amino]-3-methyl-2-oxo-benzimidazol-1-yl]butanoate;

methyl 4-[6-[(2-chloro-3-cyano-4-pyridyl)amino]-3-methyl-2-oxo-benzimidazol-1-yl]-2-(cyclopropylmethoxy)butanoate;

2-chloro-4-[[3-(2-hydroxy-3-pyrazol-1-yl-propyl)-1-methyl-2-oxo-benzimidazol-5-yl]amino]pyridine-3-carbonitrile;

2-chloro-4-[[3-(2-cyanobutyl)-1-methyl-2-oxo-benzimidazol-5-yl]amino]pyridine-3-carbonitrile;

2-chloro-4-[[3-[(3*S*)-3-hydroxybutyl]-1-methyl-2-oxo-benzimidazol-5-yl]amino]pyridine-3-carbonitrile;

Methyl 2-[[6-[(2-chloro-3-cyano-4-pyridyl)amino]-3-methyl-2-oxo-benzimidazol-1-yl]methyl]cyclopentanecarboxylate;

Methyl (2*R*)-2-amino-4-[6-[(2-chloro-3-cyano-4-pyridyl)amino]-3-methyl-2-oxo-benzimidazol-1-yl]butanoate;

N-[3-[6-[(2-Chloro-3-cyano-4-pyridyl)amino]-3-methyl-2-oxo-benzimidazol-1-yl]-1-methyl-propyl]acetamide;

5-Chloro-*N*-ethyl-4-[[3-(3-hydroxy-3-methyl-butyl)-1-methyl-2-oxo-benzimidazol-5-yl]amino]pyridine-2-carboxamide;

5-[[5-Chloro-2-(3,5-dimethylpyrazol-1-yl)pyrimidin-4-yl]amino]-3-(3-hydroxy-3-methyl-butyl)-1-methyl-benzimidazol-2-one;

5-((5-chloro-2-((3*R*,5*S*)-4,4-difluoro-3,5-dimethylpiperidin-1-yl)pyrimidin-4-yl)amino)-3-(3-hydroxy-3-methylbutyl)-1-methyl-1,3-dihydro-2*H*-benzo[d]imidazol-2-one;

ethyl 1-(5-chloro-4-((3-(3-hydroxy-3-methylbutyl)-1-methyl-2-oxo-2,3-dihydro-1*H*-benzo[d]imidazol-5-yl)amino)pyrimidin-2-yl)-1*H*-pyrazole-4-carboxylate;

ethyl 1-(5-chloro-4-((3-(3-hydroxy-3-methylbutyl)-1-methyl-2-oxo-2,3-dihydro-1*H*-benzo[d]imidazol-5-yl)amino)pyrimidin-2-yl)-3,5-dimethyl-1*H*-pyrazole-4-carboxylate;

5-((5-chloro-2-(3-(trifluoromethyl)-1*H*-pyrazol-1-yl)pyrimidin-4-yl)amino)-3-(3-hydroxy-3-methylbutyl)-1-methyl-1,3-dihydro-2*H*-benzo[d]imidazol-2-one;

5-((5-chloro-2-(3,5-dimethyl-1*H*-pyrazol-1-yl)pyrimidin-4-yl)amino)-3-(3,5-dihydroxy-3-methylpentyl)-1-methyl-1,3-dihydro-2*H*-benzo[d]imidazol-2-one;

5-((5-chloro-2-(3,5-dimethyl-1*H*-pyrazol-1-yl)pyrimidin-4-yl)amino)-3-(3-hydroxy-3-methylpentyl)-1-methyl-1,3-dihydro-2*H*-benzo[d]imidazol-2-one;

5-((5-chloro-2-(dimethylamino)pyrimidin-4-yl)amino)-3-(3-hydroxy-3-methylbutyl)-1-methyl-1,3-dihydro-2*H*-benzo[d]imidazol-2-one;

5-((5-chloro-2-(3,5-dimethyl-1*H*-pyrazol-1-yl)pyrimidin-4-yl)amino)-1-methyl-1,3-dihydro-2*H*-benzo[d]imidazol-2-one;

1-(5-chloro-4-((3-(3-hydroxy-4-methoxy-3-methylbutyl)-1-methyl-2-oxo-2,3-dihydro-1H-benzo[d]imidazol-5-yl)amino)pyrimidin-2-yl)-N,N-dimethylpiperidine-4-carboxamide;

5-((5-chloro-2-(3,5-dimethyl-1H-pyrazol-1-yl)pyrimidin-4-yl)amino)-3-(3-hydroxy-4-methoxy-3-methylbutyl)-1-methyl-1,3-dihydro-2H-benzo[d]imidazol-2-one;

1-(5-chloro-4-((3-(3,5-dihydroxy-3-methylpentyl)-1-methyl-2-oxo-2,3-dihydro-1H-benzo[d]imidazol-5-yl)amino)pyrimidin-2-yl)-N,N-dimethylpiperidine-4-carboxamide;

1-(5-chloro-4-((3-(3-hydroxy-3-methylpentyl)-1-methyl-2-oxo-2,3-dihydro-1H-benzo[d]imidazol-5-yl)amino)pyrimidin-2-yl)-N,N-dimethylpiperidine-4-carboxamide;

5-((5-chloro-2-(1H-pyrazol-1-yl)pyrimidin-4-yl)amino)-3-(3-hydroxy-3-methylbutyl)-1-methyl-1,3-dihydro-2H-benzo[d]imidazol-2-one;

5-((5-chloro-2-(4-chloro-3,5-dimethyl-1H-pyrazol-1-yl)pyrimidin-4-yl)amino)-3-(3-hydroxy-3-methylbutyl)-1-methyl-1,3-dihydro-2H-benzo[d]imidazol-2-one;

6-(3,5-dimethyl-1H-pyrazol-1-yl)-4-((3-(3-hydroxy-3-methylbutyl)-1-methyl-2-oxo-2,3-dihydro-1H-benzo[d]imidazol-5-yl)amino)nicotinonitrile;

5-((5-chloro-2-(3,5-dimethyl-1H-pyrazol-1-yl)-6-methylpyrimidin-4-yl)amino)-3-(3-hydroxy-3-methylbutyl)-1-methyl-1,3-dihydro-2H-benzo[d]imidazol-2-one;

5-((5-bromo-2-(3,5-dimethyl-1H-pyrazol-1-yl)pyrimidin-4-yl)amino)-3-(3-hydroxy-3-methylbutyl)-1-methyl-1,3-dihydro-2H-benzo[d]imidazol-2-one;

5-((5-chloro-2-(3-methyl-1H-pyrazol-1-yl)pyrimidin-4-yl)amino)-3-(3-hydroxy-3-methylbutyl)-1-methyl-1,3-dihydro-2H-benzo[d]imidazol-2-one;

5-((5-chloro-2-(5-methyl-1H-pyrazol-1-yl)pyrimidin-4-yl)amino)-3-(3-hydroxy-3-methylbutyl)-1-methyl-1,3-dihydro-2H-benzo[d]imidazol-2-one;

5-((5-chloro-2-(1H-indazol-3-yl)pyrimidin-4-yl)amino)-3-(3-hydroxy-3-methylbutyl)-1-methyl-1,3-dihydro-2H-benzo[d]imidazol-2-one;

5-((5-chloro-2-(1H-indazol-1-yl)pyrimidin-4-yl)amino)-3-(3-hydroxy-3-methylbutyl)-1-methyl-1,3-dihydro-2H-benzo[d]imidazol-2-one;

2-chloro-4-((3-(2-(1-hydroxycyclobutyl)ethyl)-1-methyl-2-oxo-2,3-dihydro-1H-benzo[d]imidazol-5-yl)amino)nicotinonitrile;

2-chloro-4-((1-methyl-3-(2-(methylsulfonyl)ethyl)-2-oxo-2,3-dihydro-1H-benzo[d]imidazol-5-yl)amino)nicotinonitrile;

2-chloro-4-((1-methyl-2-oxo-3-(3-oxopentyl)-2,3-dihydro-1H-benzo[d]imidazol-5-yl)amino)nicotinonitrile;

2-chloro-4-((1-methyl-3-((2-methyltetrahydrofuran-3-yl)methyl)-2-oxo-2,3-dihydro-1H-benzo[d]imidazol-5-yl)amino)nicotinonitrile;

2-chloro-4-((1-methyl-3-(2-(2-methyl-1,3-dioxolan-2-yl)ethyl)-2-oxo-2,3-dihydro-1H-benzo[d]imidazol-5-yl)amino)nicotinonitrile;

tert-butyl 2-((6-((2-chloro-3-cyanopyridin-4-yl)amino)-3-methyl-2-oxo-2,3-dihydro-1H-benzo[d]imidazol-1-yl)methyl)azetidine-1-carboxylate;

5-((6-((5-chloro-2-(2,2,6,6-tetramethylmorpholino)pyrimidin-4-yl)amino)-3-methyl-2-oxo-2,3-dihydro-1H-benzo[d]imidazol-1-yl)methyl)-5-(2-hydroxypropan-2-yl)-3-methyloxazolidin-2-one;

1-(5-chloro-4-((3-(3-hydroxy-3-methylbutyl)-1-methyl-2-oxo-2,3-dihydro-1H-benzo[d]imidazol-5-yl)amino)pyrimidin-2-yl)-N,N-dimethylpiperidine-4-carboxamide;

5-((5-chloro-2-(piperidin-1-yl)pyrimidin-4-yl)amino)-3-(3-hydroxy-3-methylbutyl)-1-methyl-1,3-dihydro-2H-benzo[d]imidazol-2-one;

5-((5-chloro-2-((3S,5R)-3,5-dimethylpiperidin-1-yl)pyrimidin-4-yl)amino)-3-(3-hydroxy-3-methylbutyl)-1-methyl-1,3-dihydro-2H-benzo[d]imidazol-2-one;

5-((5-chloro-2-(isopropylamino)pyrimidin-4-yl)amino)-3-(3-hydroxy-3-methylbutyl)-1-methyl-1,3-dihydro-2H-benzo[d]imidazol-2-one;

5-((5-chloro-2-(3-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-3-(3-hydroxy-3-methylbutyl)-1-methyl-1,3-dihydro-2H-benzo[d]imidazol-2-one;

5-((5-chloro-2-(3-(trifluoromethyl)piperidin-1-yl)pyrimidin-4-yl)amino)-3-(3-hydroxy-3-methylbutyl)-1-methyl-1,3-dihydro-2H-benzo[d]imidazol-2-one;

5-((5-chloro-2-(ethyl(methyl)amino)pyrimidin-4-yl)amino)-3-(3-hydroxy-3-methylbutyl)-1-methyl-1,3-dihydro-2H-benzo[d]imidazol-2-one;

5-((5-chloro-2-((2R,6S)-2,6-dimethylmorpholino)pyrimidin-4-yl)amino)-3-(3-hydroxy-3-methylbutyl)-1-methyl-1,3-dihydro-2H-benzo[d]imidazol-2-one;

5-((5-chloro-2-(2,2-dimethyl-6-(trifluoromethyl)morpholino)pyrimidin-4-yl)amino)-3-(3-hydroxy-3-methylbutyl)-1-methyl-1,3-dihydro-2H-benzo[d]imidazol-2-one;

5-((6-((5-chloro-2-((2R,6S)-2,6-dimethylmorpholino)pyrimidin-4-yl)amino)-3-methyl-2-oxo-2,3-dihydro-1H-benzo[d]imidazol-1-yl)methyl)-5-ethyl-3-methyloxazolidin-2-one;

5-((6-((5-chloro-2-(3-(trifluoromethyl)-1H-pyrazol-1-yl)pyrimidin-4-yl)amino)-3-methyl-2-oxo-2,3-dihydro-1H-benzo[d]imidazol-1-yl)methyl)-5-ethyl-3-methyloxazolidin-2-one;

5-((6-((5-chloro-2-((2R,6S)-2,6-dimethylmorpholino)pyrimidin-4-yl)amino)-3-methyl-2-oxo-2,3-dihydro-1H-benzo[d]imidazol-1-yl)methyl)-5-ethyloxazolidin-2-one;

5-((6-((5-chloro-2-(3-(trifluoromethyl)-1H-pyrazol-1-yl)pyrimidin-4-yl)amino)-3-methyl-2-oxo-2,3-dihydro-1H-benzo[d]imidazol-1-yl)methyl)-5-ethyloxazolidin-2-one;

5-((5-chloro-2-((2R,6S)-2,6-dimethylmorpholino)pyrimidin-4-yl)amino)-6-fluoro-3-(3-hydroxy-3-methylbutyl)-1-methyl-1,3-dihydro-2H-benzo[d]imidazol-2-one;

5-((5-chloro-2-((3S,5R)-3,5-dimethylpiperidin-1-yl)pyrimidin-4-yl)amino)-6-fluoro-3-(3-hydroxy-3-methylbutyl)-1-methyl-1,3-dihydro-2H-benzo[d]imidazol-2-one;

5-((5-chloro-2-(3-(trifluoromethyl)-1H-pyrazol-1-yl)pyrimidin-4-yl)amino)-6-fluoro-3-(3-hydroxy-3-methylbutyl)-1-methyl-1,3-dihydro-2H-benzo[d]imidazol-2-one;

5-((5-chloro-2-(4-(trifluoromethyl)-1H-pyrazol-1-yl)pyrimidin-4-yl)amino)-3-(3-hydroxy-3-methylbutyl)-1-methyl-1,3-dihydro-2H-benzo[d]imidazol-2-one;

5-((5-chloro-2-morpholinopyrimidin-4-yl)amino)-3-(3-hydroxy-3-methylbutyl)-1-methyl-1,3-dihydro-2H-benzo[d]imidazol-2-one;

5-((5-chloro-2-((2S,6R)-2-cyclopropyl-6-methylmorpholino)pyrimidin-4-yl)amino)-3-(3-hydroxy-3-methylbutyl)-1-methyl-1,3-dihydro-2H-benzo[d]imidazol-2-one;

5-((5-chloro-2-((2R,6R)-2-cyclopropyl-6-methylmorpholino)pyrimidin-4-yl)amino)-3-(3-hydroxy-3-methylbutyl)-1-methyl-1,3-dihydro-2H-benzo[d]imidazol-2-one;

5-((5-chloro-2-(methylthio)pyrimidin-4-yl)amino)-3-(3-hydroxy-3-methylbutyl)-1-methyl-1,3-dihydro-2H-benzo[d]imidazol-2-one;

5-((2-bromo-5-chloropyridin-4-yl)amino)-3-(3-hydroxy-3-methylbutyl)-1-methyl-1,3-dihydro-2H-benzo[d]imidazol-2-one;

2-chloro-4-((3-((5-ethyl-2-oxooxazolidin-5-yl)methyl)-1-methyl-2-oxo-2,3-dihydro-1H-benzo[d]imidazol-5-yl)amino)nicotinonitrile;

4-chloro-6-((6-fluoro-3-(3-hydroxy-3-methylbutyl)-1-methyl-2-oxo-2,3-dihydro-1H-benzo[d]imidazol-5-yl)amino)pyrimidine-5-carbonitrile;

2-chloro-4-((3-((5-ethyl-3-methyl-2-oxooxazolidin-5-yl)methyl)-1-methyl-2-oxo-2,3-dihydro-1H-benzo[d]imidazol-5-yl)amino)nicotinonitrile;

2-chloro-4-((6-fluoro-3-(3-hydroxy-3-methylbutyl)-1-methyl-2-oxo-2,3-dihydro-1H-benzo[d]imidazol-5-yl)amino)nicotinonitrile;

5-((2,5-dichloropyrimidin-4-yl)amino)-3-(3-hydroxy-3-methylpentyl)-1-methyl-1,3-dihydro-2H-benzo[d]imidazol-2-one;

5-((5-chloropyrazolo[1,5-a]pyrimidin-7-yl)amino)-3-(3-hydroxy-3-methylbutyl)-1-methyl-1,3-dihydro-2H-benzo[d]imidazol-2-one;

3-(3-hydroxy-3-methylbutyl)-1-methyl-5-((2,5,6-trichloropyrimidin-4-yl)amino)-1,3-dihydro-2H-benzo[d]imidazol-2-one;

(R)-6-chloro-5-cyano-4-((3-(3-methoxybutyl)-1-methyl-2-oxo-2,3-dihydro-1H-benzo[d]imidazol-5-yl)amino)-N-methylpicolinamide;

4-((3-(3-acetamido-3-methylbutyl)-1-methyl-2-oxo-2,3-dihydro-1H-benzo[d]imidazol-5-yl)amino)-6-chloro-5-cyano-N-methylpicolinamide;

5-((5,6-dichloropyrimidin-4-yl)amino)-3-(3-hydroxy-3-methylbutyl)-1-methyl-1,3-dihydro-2H-benzo[d]imidazol-2-one;

2-chloro-4-((3-(((1S,2S)-2-ethyl-2-hydroxycyclopentyl)methyl)-1-methyl-2-oxo-2,3-dihydro-1H-benzo[d]imidazol-5-yl)amino)nicotinonitrile;

2-chloro-4-((3-(((1S,2S)-2-hydroxy-2-methylcyclopentyl)methyl)-1-methyl-2-oxo-2,3-dihydro-1H-benzo[d]imidazol-5-yl)amino)nicotinonitrile;

5-((5-chloro-2-(1-methyl-1H-pyrazol-3-yl)pyrimidin-4-yl)amino)-3-(3-hydroxy-3-methylbutyl)-1-methyl-1,3-dihydro-2H-benzo[d]imidazol-2-one;

5-((5-chloro-2-(1,3-dimethyl-1H-pyrazol-5-yl)pyrimidin-4-yl)amino)-3-(3-hydroxy-3-methylbutyl)-1-methyl-1,3-dihydro-2H-benzo[d]imidazol-2-one;

5-((5-chloro-2-(2,4-dimethylthiazol-5-yl)pyrimidin-4-yl)amino)-3-(3-hydroxy-3-methylbutyl)-1-methyl-1,3-dihydro-2H-benzo[d]imidazol-2-one;

5-((5-chloro-2-(thiophen-2-yl)pyrimidin-4-yl)amino)-3-(3-hydroxy-3-methylbutyl)-1-methyl-1,3-dihydro-2H-benzo[d]imidazol-2-one;

5-((5-chloro-2-(1-methyl-1H-imidazol-2-yl)pyrimidin-4-yl)amino)-3-(3-hydroxy-3-methylbutyl)-1-methyl-1,3-dihydro-2H-benzo[d]imidazol-2-one;

2-chloro-4-((1-methyl-2-oxo-3-((4-(2,2,2-trifluoroethyl)morpholin-3-yl)methyl)-2,3-dihydro-1H-benzo[d]imidazol-5-yl)amino)nicotinonitrile;

2-chloro-4-((3-(2-(dimethylamino)butyl)-1-methyl-2-oxo-2,3-dihydro-1H-benzo[d]imidazol-5-yl)amino)nicotinonitrile;

(S)-2-chloro-4-((3-((1-ethylpyrrolidin-2-yl)methyl)-1-methyl-2-oxo-2,3-dihydro-1H-benzo[d]imidazol-5-yl)amino)nicotinonitrile;

2-chloro-4-((1-methyl-2-oxo-3-((1-(2,2,2-trifluoroethyl)piperidin-2-yl)methyl)-2,3-dihydro-1H-benzo[d]imidazol-5-yl)amino)nicotinonitrile;

(S)-2-chloro-4-((3-((1-(2-fluoroethyl)pyrrolidin-2-yl)methyl)-1-methyl-2-oxo-2,3-dihydro-1H-benzo[d]imidazol-5-yl)amino)nicotinonitrile;

(S)-2-chloro-4-((3-((1-(2-hydroxyethyl)pyrrolidin-2-yl)methyl)-1-methyl-2-oxo-2,3-dihydro-1H-benzo[d]imidazol-5-yl)amino)nicotinonitrile;

2-chloro-4-((3-(((2R,4S)-4-fluoro-1-(2,2,2-trifluoroethyl)pyrrolidin-2-yl)methyl)-1-methyl-2-oxo-2,3-dihydro-1H-benzo[d]imidazol-5-yl)amino)nicotinonitrile;

2-chloro-4-((3-(3-(ethylamino)butyl)-1-methyl-2-oxo-2,3-dihydro-1H-benzo[d]imidazol-5-yl)amino)nicotinonitrile;

2-chloro-4-((3-(3-hydroxy-3-methylhex-5-yn-1-yl)-1-methyl-2-oxo-2,3-dihydro-1H-benzo[d]imidazol-5-yl)amino)nicotinonitrile;

2-chloro-4-((3-(3-hydroxy-4-methoxy-3-methylbutyl)-1-methyl-2-oxo-2,3-dihydro-1H-benzo[d]imidazol-5-yl)amino)nicotinonitrile;

2-chloro-4-((3-(3-hydroxy-3-methylpentyl)-1-methyl-2-oxo-2,3-dihydro-1H-benzo[d]imidazol-5-yl)amino)nicotinonitrile;

2-chloro-4-((1-methyl-2-oxo-3-(4,4,4-trifluoro-3-hydroxy-3-methylbutyl)-2,3-dihydro-1H-benzo[d]imidazol-5-yl)amino)nicotinonitrile;

2-chloro-4-((3-(3-hydroxy-2,3-dimethylbutyl)-1-methyl-2-oxo-2,3-dihydro-1H-benzo[d]imidazol-5-yl)amino)nicotinonitrile;

2-chloro-4-((3-(3-hydroxy-3,4-dimethylpentyl)-1-methyl-2-oxo-2,3-dihydro-1H-benzo[d]imidazol-5-yl)amino)nicotinonitrile;

5-((5-chloro-2-(2,2,6,6-tetramethylmorpholino)pyrimidin-4-yl)amino)-3-(3-hydroxy-3-methylbutyl)-1-methyl-1,3-dihydro-2H-benzo[d]imidazol-2-one;

5-((5-chloro-2-(2-(trifluoromethyl)morpholino)pyrimidin-4-yl)amino)-3-(3-hydroxy-3-methylbutyl)-1-methyl-1,3-dihydro-2H-benzo[d]imidazol-2-one;

5-((3-chloro-2-fluoropyridin-4-yl)amino)-3-(3-hydroxy-3-methylbutyl)-1-methyl-1,3-dihydro-2H-benzo[d]imidazol-2-one;

5-((2,3-dichloropyridin-4-yl)amino)-3-(3-hydroxy-3-methylbutyl)-1-methyl-1,3-dihydro-2H-benzo[d]imidazol-2-one;

5-((3-bromopyridin-4-yl)amino)-3-(3-hydroxy-3-methylbutyl)-1-methyl-1,3-dihydro-2H-benzo[d]imidazol-2-one;

5-((5-chloro-2-(trifluoromethyl)pyridin-4-yl)amino)-3-(3-hydroxy-3-methylbutyl)-1-methyl-1,3-dihydro-2H-benzo[d]imidazol-2-one;

5-((3-chloropyridin-4-yl)amino)-3-(3-hydroxy-3-methylbutyl)-1-methyl-1,3-dihydro-2H-benzo[d]imidazol-2-one;

5-((5-chloro-2-(3,5-dimethyl-1H-pyrazol-1-yl)pyridin-4-yl)amino)-3-(3-hydroxy-3-methylbutyl)-1-methyl-1,3-dihydro-2H-benzo[d]imidazol-2-one;

5-((5-chloro-2-(2-oxopyrrolidin-1-yl)pyrimidin-4-yl)amino)-3-(3-hydroxy-3-methylbutyl)-1-methyl-1,3-dihydro-2H-benzo[d]imidazol-2-one;

(S)-5-((5-chloro-2-(2-(hydroxymethyl)pyrrolidin-1-yl)pyrimidin-4-yl)amino)-3-(3-hydroxy-3-methylbutyl)-1-methyl-1,3-dihydro-2H-benzo[d]imidazol-2-one;

(S)-7-((3-(3-hydroxy-3-methylbutyl)-1-methyl-2-oxo-2,3-dihydro-1H-benzo[d]imidazol-5-yl)amino)-5-(2-(hydroxymethyl)pyrrolidin-1-yl)pyrazolo[1,5-a]pyrimidine-3-carbonitrile;

2-chloro-4-((1-methyl-3-(2-(2-methyloxiran-2-yl)ethyl)-2-oxo-2,3-dihydro-1H-benzo[d]imidazol-5-yl)amino)nicotinonitrile;

2-chloro-4-((3-(2-(3,5-dimethyl-2-oxooxazolidin-5-yl)ethyl)-1-methyl-2-oxo-2,3-dihydro-1H-benzo[d]imidazol-5-yl)amino)nicotinonitrile;

2-chloro-4-((1-methyl-3-((5-methyl-2-oxooxazolidin-4-yl)methyl)-2-oxo-2,3-dihydro-1H-benzo[d]imidazol-5-yl)amino)nicotinonitrile;

(S)-5-((5-chloro-2-(2,2,6,6-tetramethylmorpholino)pyrimidin-4-yl)amino)-1-methyl-3-((1-(2,2,2-trifluoroethyl)pyrrolidin-2-yl)methyl)-1,3-dihydro-2H-benzo[d]imidazol-2-one;

5-((5-chloro-2-((3R,5S)-4,4-difluoro-3,5-dimethylpiperidin-1-yl)pyrimidin-4-yl)amino)-1,3-bis(3-hydroxy-3-methylbutyl)-1,3-dihydro-2H-benzo[d]imidazol-2-one;

5-((5-chloro-2-(2,2,6,6-tetramethylmorpholino)pyrimidin-4-yl)amino)-1,3-bis(3-hydroxy-3-methylbutyl)-1,3-dihydro-2H-benzo[d]imidazol-2-one;

5-((5-chloro-2-(4,4-difluoropiperidin-1-yl)pyrimidin-4-yl)amino)-3-(3-hydroxy-3-methylbutyl)-1-methyl-1,3-dihydro-2H-benzo[d]imidazol-2-one;

5-((5-chloro-2-(3,3-difluoro-8-azabicyclo[3.2.1]octan-8-yl)pyrimidin-4-yl)amino)-3-(3-hydroxy-3-methylbutyl)-1-methyl-1,3-dihydro-2H-benzo[d]imidazol-2-one;

5-((5-chloro-2-(2,2-difluoro-7-azaspiro[3.5]nonan-7-yl)pyrimidin-4-yl)amino)-3-(3-hydroxy-3-methylbutyl)-1-methyl-1,3-dihydro-2H-benzo[d]imidazol-2-one;

5-((5-chloro-2-(4-(trifluoromethyl)piperidin-1-yl)pyrimidin-4-yl)amino)-3-(3-hydroxy-3-methylbutyl)-1-methyl-1,3-dihydro-2H-benzo[d]imidazol-2-one;

5-((5-chloro-2-(4,4-difluoropiperidin-1-yl)pyridin-4-yl)amino)-3-(3-hydroxy-3-methylbutyl)-1-methyl-1,3-dihydro-2H-benzo[d]imidazol-2-one;

5-((5-chloro-2-((2R,6S)-2,6-dimethylmorpholino)pyridin-4-yl)amino)-3-(3-hydroxy-3-methylbutyl)-1-methyl-1,3-dihydro-2H-benzo[d]imidazol-2-one;

1-(5-chloro-4-((3-(3-hydroxy-3-methylbutyl)-1-methyl-2-oxo-2,3-dihydro-1H-benzo[d]imidazol-5-yl)amino)pyridin-2-yl)-N,N-dimethylpiperidine-4-carboxamide;

(R)-2-chloro-4-((3-(3-hydroxybutyl)-1-methyl-2-oxo-2,3-dihydro-1H-benzo[d]imidazol-5-yl)amino)nicotinonitrile;

(S)-2-chloro-4-((3-((1-(2,2-difluoroethyl)pyrrolidin-2-yl)methyl)-1-methyl-2-oxo-2,3-dihydro-1H-benzo[d]imidazol-5-yl)amino)nicotinonitrile;

5-((5-chloro-2-(4,4-difluoro-3-(hydroxymethyl)piperidin-1-yl)pyrimidin-4-yl)amino)-3-(3-hydroxy-3-methylbutyl)-1-methyl-1,3-dihydro-2H-benzo[d]imidazol-2-one;

5-((5-chloro-2-(4,4-difluoro-3-(methoxymethyl)piperidin-1-yl)pyrimidin-4-yl)amino)-3-(3-hydroxy-3-methylbutyl)-1-methyl-1,3-dihydro-2H-benzo[d]imidazol-2-one;

5-((5-chloro-2-(4,4-difluoro-3-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-3-(3-hydroxy-3-methylbutyl)-1-methyl-1,3-dihydro-2H-benzo[d]imidazol-2-one;

5-((5-chloro-2-(4-(trifluoromethyl)piperidin-1-yl)pyrimidin-4-yl)amino)-3-(3-hydroxy-3-methylbutyl)-1-methyl-1,3-dihydro-2H-benzo[d]imidazol-2-one;

5-((5-chloro-2-((3R,4S)-3,4-difluoropyrrolidin-1-yl)pyrimidin-4-yl)amino)-3-(3-hydroxy-3-methylbutyl)-1-methyl-1,3-dihydro-2H-benzo[d]imidazol-2-one;

5-((5-chloro-2-((3S,5R)-3,4,5-trimethylpiperazin-1-yl)pyrimidin-4-yl)amino)-3-(3-hydroxy-3-methylbutyl)-1-methyl-1,3-dihydro-2H-benzo[d]imidazol-2-one;

5-((2-((1R,5S)-8-azabicyclo[3.2.1]octan-8-yl)-5-chloropyrimidin-4-yl)amino)-3-(3-hydroxy-3-methylbutyl)-1-methyl-1,3-dihydro-2H-benzo[d]imidazol-2-one;

5-((2-((1R,5S)-3-azabicyclo[3.2.1]octan-3-yl)-5-chloropyrimidin-4-yl)amino)-3-(3-hydroxy-3-methylbutyl)-1-methyl-1,3-dihydro-2H-benzo[d]imidazol-2-one;

5-((5-chloro-2-((3aR,7aS)-octahydro-2H-isoindol-2-yl)pyrimidin-4-yl)amino)-3-(3-hydroxy-3-methylbutyl)-1-methyl-1,3-dihydro-2H-benzo[d]imidazol-2-one;

5-((5-chloro-2-((3R,4S)-3,4-dimethylpyrrolidin-1-yl)pyrimidin-4-yl)amino)-3-(3-hydroxy-3-methylbutyl)-1-methyl-1,3-dihydro-2H-benzo[d]imidazol-2-one;

6-((5-chloro-2-(2,2,6,6-tetramethylmorpholino)pyrimidin-4-yl)amino)-1,3-bis(3-hydroxy-3-methylbutyl)-1,3-dihydro-2H-imidazo[4,5-b]pyridin-2-one;

6-((5-chloro-2-((3R,5S)-4,4-difluoro-3,5-dimethylpiperidin-1-yl)pyrimidin-4-yl)amino)-1,3-bis(3-hydroxy-3-methylbutyl)-1,3-dihydro-2H-imidazo[4,5-b]pyridin-2-one;

5-((5-chloro-2-(8,8-difluoro-3-azabicyclo[3.2.1]octan-3-yl)pyrimidin-4-yl)amino)-3-(3-hydroxy-3-methylbutyl)-1-methyl-1,3-dihydro-2H-benzo[d]imidazol-2-one;

5-((2-((1r,3r,5r,7r)-2-azaadamantan-2-yl)-5-chloropyrimidin-4-yl)amino)-3-(3-hydroxy-3-methylbutyl)-1-methyl-1,3-dihydro-2H-benzo[d]imidazol-2-one;

4-chloro-6-((5-chloro-2-(2,2,6,6-tetramethylmorpholino)pyrimidin-4-yl)amino)-1,3-bis(3-hydroxy-3-methylbutyl)-1,3-dihydro-2H-benzo[d]imidazol-2-one;

5-((5-chloro-2-(3-(methoxymethyl)piperidin-1-yl)pyrimidin-4-yl)amino)-3-(3-hydroxy-3-methylbutyl)-1-methyl-1,3-dihydro-2H-benzo[d]imidazol-2-one;

5-((5-chloro-2-(6,6-difluoro-3-azabicyclo[3.1.1]heptan-3-yl)pyrimidin-4-yl)amino)-3-(3-hydroxy-3-methylbutyl)-1-methyl-1,3-dihydro-2H-benzo[d]imidazol-2-one;

5-((5-chloro-2-((3S,5R)-3-hydroxy-5-methylpiperidin-1-yl)pyrimidin-4-yl)amino)-3-(3-hydroxy-3-methylbutyl)-1-methyl-1,3-dihydro-2H-benzo[d]imidazol-2-one;

5-((5-chloro-2-((3R,5S)-3,5-dimethylazepan-1-yl)pyrimidin-4-yl)amino)-3-(3-hydroxy-3-methylbutyl)-1-methyl-1,3-dihydro-2H-benzo[d]imidazol-2-one;

5-((5-chloro-2-(3-phenylpiperidin-1-yl)pyrimidin-4-yl)amino)-3-(3-hydroxy-3-methylbutyl)-1-methyl-1,3-dihydro-2H-benzo[d]imidazol-2-one;

5-((2-(4-((1H-pyrazol-1-yl)methyl)piperidin-1-yl)-5-chloropyrimidin-4-yl)amino)-3-(3-hydroxy-3-methylbutyl)-1-methyl-1,3-dihydro-2H-benzo[d]imidazol-2-one;

5-((5-chloro-2-((3S,5R)-3,5-dimethylpiperidin-1-yl)pyridin-4-yl)amino)-3-(3-hydroxy-3-methylbutyl)-1-methyl-1,3-dihydro-2H-benzo[d]imidazol-2-one;

5-((5-chloro-2-((3R,5S)-3,5-dimethylpiperidine-1-carbonyl)pyridin-4-yl)amino)-3-(3-hydroxy-3-methylbutyl)-1-methyl-1,3-dihydro-2H-benzo[d]imidazol-2-one;

5-((5-chloro-2-((3S,5R)-4,4-difluoro-3,5-dimethylpiperidine-1-carbonyl)pyridin-4-yl)amino)-3-(3-hydroxy-3-methylbutyl)-1-methyl-1,3-dihydro-2H-benzo[d]imidazol-2-one;

5-((5-chloro-2-((3S,5R)-4,4-difluoro-3,5-dimethylpiperidin-1-yl)pyrimidin-4-yl)amino)-1-(2-(dimethylamino)ethyl)-3-(3-hydroxy-3-methylbutyl)-1,3-dihydro-2H-benzo[d]imidazol-2-one;

5-((5-chloro-2-((3S,5R)-4,4-difluoro-3,5-dimethylpiperidin-1-yl)pyrimidin-4-yl)amino)-3-(3-hydroxy-3-methylbutyl)-1-(2-morpholinoethyl)-1,3-dihydro-2H-benzo[d]imidazol-2-one;

ethyl (E)-4-(5-((5-chloro-2-((3S,5R)-4,4-difluoro-3,5-dimethylpiperidin-1-yl)pyrimidin-4-yl)amino)-3-(3-hydroxy-3-methylbutyl)-2-oxo-2,3-dihydro-1H-benzo[d]imidazol-1-yl)but-2-enoate;

5-((5-Chloro-2-(2,2,2-trifluoroethoxy)pyrimidin-4-yl)amino)-3-(3-hydroxy-3-methylbutyl)-1-methyl-1,3-dihydro-2H-benzo[d]imidazol-2-one;

5-[[5-chloro-2-[(3S,5R)-4,4-difluoro-3,5-dimethyl-1-piperidyl]pyrimidin-4-yl]amino]-1-(2-hydroxyethyl)-3-(3-hydroxy-3-methyl-butyl)benzimidazol-2-one;

5-((5-chloro-2-((3S,5R)-4,4-difluoro-3,5-dimethylpiperidin-1-yl)pyrimidin-4-yl)amino)-1-(2,2-dimethoxyethyl)-3-(3-hydroxy-3-methylbutyl)-1,3-dihydro-2H-benzo[d]imidazol-2-one;

1-(5-chloro-4-((3-(3-hydroxy-3-methylbutyl)-1-methyl-2-oxo-2,3-dihydro-1H-benzo[d]imidazol-5-yl)amino)pyrimidin-2-yl)piperidine-4-carbonitrile;

1-(5-chloro-4-((3-(3-hydroxy-3-methylbutyl)-1-methyl-2-oxo-2,3-dihydro-1H-benzo[d]imidazol-5-yl)amino)pyrimidin-2-yl)piperidine-3-carbonitrile;

5-((5-chloro-2-(4-(morpholinomethyl)piperidin-1-yl)pyrimidin-4-yl)amino)-3-(3-hydroxy-3-methylbutyl)-1-methyl-1,3-dihydro-2H-benzo[d]imidazol-2-one;

5-((5-chloro-2-(3-(morpholinomethyl)piperidin-1-yl)pyrimidin-4-yl)amino)-3-(3-hydroxy-3-methylbutyl)-1-methyl-1,3-dihydro-2H-benzo[d]imidazol-2-one;

5-((5-chloro-2-(2-methyl-2,4,5,7-tetrahydro-6H-pyrazolo[3,4-c]pyridin-6-yl)pyrimidin-4-yl)amino)-3-(3-hydroxy-3-methylbutyl)-1-methyl-1,3-dihydro-2H-benzo[d]imidazol-2-one;

5-((5-chloro-2-(1-methyl-1,4,5,7-tetrahydro-6H-pyrazolo[3,4-c]pyridin-6-yl)pyrimidin-4-yl)amino)-3-(3-hydroxy-3-methylbutyl)-1-methyl-1,3-dihydro-2H-benzo[d]imidazol-2-one;

5-((5-chloro-2-(3-(hydroxymethyl)piperidin-1-yl)pyrimidin-4-yl)amino)-3-(3-hydroxy-3-methylbutyl)-1-methyl-1,3-dihydro-2H-benzo[d]imidazol-2-one;

5-((5-chloro-2-(4-morpholinopiperidin-1-yl)pyrimidin-4-yl)amino)-3-(3-hydroxy-3-methylbutyl)-1-methyl-1,3-dihydro-2H-benzo[d]imidazol-2-one;

5-((2-(4-(1H-pyrazol-1-yl)piperidin-1-yl)-5-chloropyrimidin-4-yl)amino)-3-(3-hydroxy-3-methylbutyl)-1-methyl-1,3-dihydro-2H-benzo[d]imidazol-2-one; or

5-((5-chloro-2-(2-(hydroxymethyl)morpholino)pyrimidin-4-yl)amino)-3-(3-hydroxy-3-methylbutyl)-1-methyl-1,3-dihydro-2H-benzo[d]imidazol-2-one.

13. A compound according to any one of the preceding claims, or a pharmaceutically acceptable salt or hydrate thereof, for use in therapy.
14. A pharmaceutical composition comprising a compound according to any one of claims 1 to 12, or a pharmaceutically acceptable salt or hydrate thereof and a pharmaceutically acceptable carrier or excipient.
15. A compound according to any one of claims 1 to 12, or a pharmaceutically acceptable salt, hydrate or solvate thereof, or a pharmaceutical composition according to claim 13, for use in the treatment of cancer.
16. The compound or the pharmaceutical composition for use according to claim 15, wherein said cancer is diffuse large B-cell lymphoma (DLBCL), follicular lymphoma (FL), Burkitt lymphoma (BL), angioimmunoblastic T-cell lymphoma (AITL), acute lymphoblastic leukaemia (ALL), chronic myeloid leukaemia (CML), multiple myeloma, breast cancer, non-small cell lung cancer (NSCLC) or squamous cell carcinomas (SCC) of the head and neck, oesophagus, lung or ovary
17. Use of a compound according to any of claims 1 to 12 or a pharmaceutically acceptable salt or hydrate thereof, or a pharmaceutical composition according to claim 14 in the manufacture of a medicament for the treatment of cancer in a subject in need of such treatment, wherein a therapeutically effective amount of the compound or the pharmaceutical composition is to be administered.
18. The use according to claim 17, wherein said cancer is diffuse large B-cell lymphoma (DLBCL), follicular lymphoma (FL), Burkitt lymphoma (BL), angioimmunoblastic T-cell lymphoma (AITL), acute lymphoblastic leukaemia (ALL), chronic myeloid leukaemia

(CML), multiple myeloma, breast cancer, non-small cell lung cancer (NSCLC) or squamous cell carcinomas (SCC) of the head and neck, oesophagus, lung or ovary.